



23/ 964

65(1)









James Lee

James Lee

Engraved by Thomas G. Sharp

AN
INTRODUCTION
TO THE
SCIENCE OF BOTANY,



CHIEFLY EXTRACTED FROM THE
WORKS OF LINNÆUS;
TO WHICH ARE ADDED,
SEVERAL NEW TABLES AND NOTES,
AND
A LIFE OF THE AUTHOR.

BY THE LATE JAMES LEE,
PURSEYMAN, AND FLORIST, AT THE VINEYARD, HAMMERSMITH.

FOURTH EDITION, CORRECTED AND ENLARGED.
BY JAMES LEE,
SON AND SUCCESSOR TO THE AUTHOR.

LONDON:

PRINTED FOR F. C. AND J. RIVINGTON; WILKIE AND ROBIN-
SON; J. WALKER; WHITE AND CO.; LONGMAN, HURST, REES,
AND ORME; CADELL AND DAVIES; LACKINGTON, ALLEN, AND
CO.; J. BOOKER; J. MAWMAN; B. CROSBY AND CO.; W.
BAYNES; SHERWOOD, NEELY, AND JONES; J. JOHNSON AND
CO.; AND C. AND R. BALDWIN.

1810.

647828

THE UNIVERSITY OF TORONTO

LIBRARY



Printed by S. Hamilton, Weybridge.

DEDICATION.



TO JAMES EDWARD SMITH, M. D.

PRESIDENT OF THE LINNÆAN SOCIETY

&c. &c.

Vineyard, Hammersmith, May 1, 1810.

SIR,

THERE is no gentleman in this country better able to appreciate the merit of my father's work on Botany, than yourself; and when it is considered at what an early period of the establishment of the true science his "*Introduction*" came forth, and how much it has done to introduce the SEXUAL SYSTEM to the notice of the public, it cannot fail to be pleasing to YOU, who are daily advancing the Science of Botany, to see the same work continued in its publication, with all the modern improvements, which have poured in like a torrent upon the botanist, and made the science assume, as it were, a new face. Although many elementary

works have since appeared, still I believe I may with confidence assert, that none has been contrived on a more useful plan than the present one; nor can any work boast of so many useful TABLES, by all acknowledged to be extremely necessary for constant reference, and which makes the present work an invaluable companion to all persons who are anxious to acquire the true knowledge of plants. Unless for these TABLES, first contrived by my father, your admirable "Introduction to Systematic Botany" would have superseded the necessity of the republication of the present edition; but these TABLES, peculiar to our publication, renders our work of great public utility: and you will find, Sir, that we have here paid very considerable attention to the TABLES, and even ventured to add some additional ones. I have the honour to remain,


Dear Sir,

With the greatest respect and devotion,

Your obliged and faithful servant,

JAMES LEE.

SKETCH
OF THE
LIFE AND WRITINGS
OF
THE LATE JAMES LEE,
BY ROBERT JOHN THORNTON, M. D.



THE lives of Botanists seldom present any very remarkable features to interest the general reader. The destroyers of kingdoms, like the history of highwaymen in the Newgate Calendar, engross the attention of the majority of mankind. ADAM, in his state of primitive virtue, cultivating Paradise with EVE, has, with some, less of attraction than when he viewed the dismal fruits of the fall of man, seen in the murder of ABEL by CAIN. Still, however, there are to be found persons, who rather delight in the short account of *British Worthies*, whose trophies are bloodless, to reading of the feats of men whose actions are a disgrace to humanity.

Mr. JAMES LEE was born at Selkirk, in Scotland, of respectable parents ; and being a youth of very

promising talents, he was sent to England in the year 1715, to be under the immediate protection of the EARL OF ILLA, who continued his education, and gave him the free use of his library. He had an early love for plants, and studied Botany at a period when few persons in England had any knowledge of the science. In his time a meteor appeared in the North, which was the great LINNÆUS, who was born in the year 1707. As the foundation of the reputation of LEE depended upon the Reform that this transcendent genius wrought in Botany, and since his "*Introduction to Botany*," as he expresses in the title-page, is but a transcript of the mind of that most distinguished naturalist, it may not be improper in a work like the present, to say a few words respecting the progress of the Science of Botany. Previous to the time of LINNÆUS, NEHEMIAH GREW, an Englishman, flourished a very eminent physiologist, who consulted not books, but Nature, and wrote his "*Vegetable Anatomy*," in 1682. In this work he mentions the *Sexes of Plants*, relating a conversation he held on the subject, with Sir THOMAS MILLINGTON, Savilian Professor of Natural History at Oxford, and President of the Royal London College of Physicians. SEBASTIAN VAILLANT also wrote "*a Discourse on the Structure of Flowers*," confirming the doctrine of the *Sexes of Plants*, which LINNÆUS acknowledges to have read, and which might have laid the foundation of his building up a

System on this important discovery. TOURNEFORT also flourished before the period of LINNÆUS, and his fame in 1683, procured him the appointment of Botanic Professor in the King's Garden. At the expense of the King of France, in pursuit of plants, he travelled over all the countries of Europe, and spent three years in the Levant. His glory is, to have formed a *System*, beautiful in itself, but suited to a limited knowledge of plants, which could then be accommodated to such a system; and to have invented the method of forming plants into their respective *Genera*, since perfected by LINNÆUS. His "*Elements of Botany*" evince a vast knowledge of the genera and the species of plants, and this botanical work is one of which the French are, even to the present day, passionately fond. He rose to be President of the head of the faculty at Paris. This illustrious botanist was born in 1656, and died in 1707, the same year that LINNÆUS came into the world.

RAY was the contemporary of TOURNEFORT, somewhat prior, being born in 1628, and from his studies at Cambridge, his health declined, and he was obliged, for its recovery, to go much in the fields. In these excursions, plants naturally presented themselves, and he hence became enamoured of the science of Botany. He first published a "*Catalogue of the Plants growing about Cambridge.*" Travelling abroad, his vast mind collected a knowledge of various

plants, and, like TOURNEFORT, he wished to dispose these into a method, and invented a celebrated "*System*," more perfect than that of TOURNEFORT, but less simple and practical than that of LINNÆUS; and arranged under his own System, the "*Historia Plantarum*," "a History of Plants," in three large folio volumes, being a description of all the species of plants known at that period.

The botanical world found the chaotic mass somewhat removed by these illustrious men: but still the science of Botany was of difficult attainment, and many new Plants could not be reduced to the Systems of either TOURNEFORT or RAY, when LINNÆUS turned his attention to this science. He soon became the pride and wonder of the age! Like the sun, when he flourished, all preceding botanists hid their diminished heads, and are now only read to know the state of natural science before the period of LINNÆUS! He ranged throughout every path of Nature, and left nothing unattempted or unaccomplished! He may be truly said to have lived, if life is to be computed by acquisitions, for he saw and described more than others had seen and done in a thousand years, and each day with him appears, from his gigantic achievements, an age! He introduced truth, order, precision, and perfection, into Natural History! He borrowed from none, his labours are all original! Attacked by numerous and ran-

corous adversaries, who cowardly and morosely addressed the prejudices of the vulgar against him, he only retorted by embellishing his portrait with a monkey teasing a bear, in allusion to their characters, and sensible of his own. His "*Sexual System*" first proved his uncommon genius. His "*Philosophia Botanica*," "*Botanical Philosophy*," cleared away all the obscurities in Botany, and formed it into a science. His "*Genera Plantarum*," containing a full description of the minutest parts of each genus of plants, showed the most consummate patience, the nicest observation, and the greatest skill. His "*Species Plantarum*," and his "*Systema Naturæ*," "*Species of Plants*," and "*System of Nature*," evince not only such an acquaintance with all plants, as is truly astonishing, but also with all the wonderful works of God throughout nature. His "*travels*," and works on "*medicine*," are only little considered from the superlative excellence of his other labours. He was honoured and encouraged by the patronage of the King and Queen of Sweden. His only opponent in this country was Sir HANS SLOANE, President of the Royal Society, who was envious of his fame, and who treated him, when in this country, somewhat rudely. Adored, beloved, honoured, LINNÆUS saw his darling pursuits advance with rapid steps, not only in Sweden, but throughout every country of Europe, and himself looked up to as the FATHER of NATURAL HISTORY. Pupils of his own choice, supported by his government, traversed

the globe; and sent him its produce to arrange. When he died, in the year 1778, the KING OF SWEDEN, in his annual address, mentioned him as a public loss, and the whole university attended his funeral; and there was also, on this occasion, a general mourning.

LEE, who was passionately fond of Botany, in all probability saw LINNÆUS when he visited this country; and it is well known, that he afterwards corresponded with that naturalist, and sent him specimens of such rare plants as were in his possession.

The EARL OF ILLA, observing the bent of the mind of LEE, promoted his entering into partnership with Mr. KENNEDY, who was a nurseryman and florist of some eminence at that period, at Hammersmith.

It was there he conceived and executed the plan of transfusing into our language, the learned improvements of the great LINNÆUS, and his System; and he preferred the form, he then gave it, to a literal translation of the "*Philosophia Botanica*" of that great genius. It commences with the flower, as being the part most attractive and interesting to the young botanist; and the first ten chapters give a clear exposition of the seven component parts of fructification: in the next ten chapters the reader is advanced

into the difficulties of the science ; and the twenty-first chapter treats of the *Sexes of Plants*. In LINNÆUS'S "*Philosophia Botanica*" the learner, on the contrary, is made to begin from the root in the ground, and may thus get disgusted with the science at the very onset. In laying down his principles, LINNÆUS gave few or no examples ; this he probably reserved for his lectures, and we may here remark, that his fame had attracted to Upsal three thousand pupils, some of whom were nobles ; and that persons from all countries flocked to him, even from our own. The distinguishing merit of LEE'S work is, that it abounds with examples. There is scarce a single axiom laid down, but four or five illustrations are given ; and this arose from his being practically acquainted not only with native plants, of which he had formed a large and valuable *Hortus Siccus*, but also with exotic botany.

He next explains, in what he calls Part II., the *Sexual System* of LINNÆUS, and illustrates the *Classes* and *Orders* by an enumeration of all the *Genera* which arrange themselves under this system. By some, this crowding in of names may be objected ; but it may be right to observe, that the chief merit of this Introduction is, that he does not go slovenly to work, and an idle person can never expect to become a botanist.

He treats next of the *Genera of Plants*, entering upon the discoveries of former botanists, and closes this part with TABULAR ILLUSTRATIONS, the merit of which mode of instruction must be acknowledged by every person.

The science of Botany may be compared to a ladder, being only an artificial aid by which we mount up to a knowledge of plants. Parts I. and II. may be called the FIRST STEPS IN BOTANY. It may be objected by some, that he has no where given us the derivations of the terms used in this science; but it should be observed, that he wrote principally for the unlearned, and calls his book only an "Introduction."

The next great advance in Botany is, the knowledge of the *species of plants*; and to obtain this, the student must make more progress into the science of Botany. The *genera* are founded upon the fructification alone, the *species* upon all the parts of plants. In Part III. he lays down the general plan, then treats of roots, trunks, leaves, &c. which he does in a very able manner, often giving many examples, illustrative of the terms; and he closes with some more useful TABLES, and a short exposition of all the terms of Botany, being a direct translation of the "*Termini Botanici*" of LINNÆUS, finishing with Plates, copied from LINNÆUS, which, in this

new edition, are considerably improved. We need not here enter widely into the merits of this work, which has gone through several editions, and is generally the first book that the botanist purchases; and has laid the foundation of the knowledge of Botany, which principally exists at this day.

Other introductions possess also considerable merit. The learned President of the Linnæan Society has favoured the world with one that surpasses, from its clearness and elegance, all power of praise, but still they want TABLES; and, I am persuaded, this will ever hold its rank, as a popular Introduction, and even attract more purchasers, from this very consideration of the many useful TABLES it contains.

The “*Elements of Botany*,” by ROSE, which is a more direct translation of the “*Philosophia Botanica*” of LINNÆUS, possessing the same order, though deservedly recommended, has fallen into general disuse from this very cause, and so we may predict of all the other elementary books on the science of Botany.

But, to return to the subject of our memoir. The great LINNÆUS felt no jealousy at the manner LEE had adopted to diffuse Botany amongst his countrymen; but, on the contrary, in testimony to his knowledge, named a new plant after him, LEEA.

But the knowledge of Mr. LEE was not confined to Botany, he was also an adept in ~~et~~ymology, conchology, and natural history in general, of which he made a most superb collection, which is still in the possession of his son; and this cabinet, possessing many unique specimens of insects and shells, is often quoted by Fabricius, and other eminent authors.

He sent out persons to different quarters of the globe, to collect new plants; and his extensive stoves, green-house, and nursery, was the emporium of all that was curious and interesting in Botany. He discovered what islands had belonged to Europe, and what to Asia, by the *heath* (ERICA), which is abundantly dispersed over Europe, Africa, and America; but is not to be found in Asia, or any of its islands, which once formed a part of that continent.

Although the great exertions made to extend the Royal Garden, at Kew, and large sums expended, made that the chief repository of new and rare plants, still Mr. LEE's Nursery, at Hammersmith, took, at any rate, the second lead; and the two together has gradually, and, imperceptibly as it were, greatly enriched our gardens, and extended the Science of Botany. Prints of new plants are for ever acknowledging the favour of Mr. LEE.

As might be expected from an author, LEE'S Garden was always open to the curious; nor was he ever backward in communicating knowledge; whereas Mr. MILLER concealed the names of his valuable collection in the Chelsea Gardens; and the papers, which contained his foreign seeds, were industriously thrown into the Thames; and such is the ardour of Botany, although the acquisition was often to be swam for, these were fished for up again, and the names of the new plants, then introduced, was thus known to Mr. LEE, and others, in a way which greatly surprised the author of the Gardener's Dictionary.

LEE might have died rich, but he was notoriously generous; and cared not what expenses he was at for the attainment of rare plants; and when he possessed such as might have procured him a golden harvest, he chose rather to give duplicates away to lovers of Botany, before the selling them to the rich but careless collectors of flowers, rather led to them through ostentation, than from a laudable enthusiasm in the pursuit of knowledge. He never concealed his methods of propagating plants; and he generally observed, that, for want of insects to further the nuptials of plants, or a proper degree of ventilation, or rather favouring breezes, or from some defect in the escape of the pollen from the anthers, that the seeds in stove plants are in general unproductive; and for a series of years *artificial impregnation* has been performed at Hammersmith.

which always secured an increase, and proves the practical value of science.

He had the felicity of having his company courted by all the illustrious botanists of the day, as the Rev. Dr. HALES, the celebrated author of "Vegetable Statics;" the Rev. Dr. COLIN MILNE, author of a "Philosophical and Systematic Dictionary," a work that is in the hands of every botanist, which has gone through four editions; the illustrious Dr. FOTHERGILL, the great patron of Botany; MILLER, author of a "Botanical Dictionary," being the very first work of its kind; Professor MARTYN, author of "Letters on Botany," and editor of a new edition of Miller's "Dictionary," which, from the additions made, may be almost styled his own; Dr. WITHERING, author of a "New Arrangement of British Plants," a work which has considerably advanced the Science of Botany; and the celebrated JOHN HUNTER, a great lover of natural history.

The Marchioness of Rockingham was exceedingly fond of plants, and Mr. Lee used to dine once every week at Hillingdon, with this amiable lady; and the Marquis always gave him a hearty welcome, with a hospitality becoming a nobleman. In short, he was esteemed and courted by a numerous circle of the first people; and he lived to the very advanced age of eighty. With patient resignation he met the expected summons, July 1795, and was universally regretted by great and poor.

To sum up the character of Lee, he was an ardent enthusiast in the pursuit of natural science;

although he had, what is proverbial, plenty of bowing, still he was courted by the great, rather for his attainments in Botany, and clear vigorous understanding, than for a politeness which appeared in him natural: he was very conscientious in all his dealings; he was generous to a fault; his garden was the resort of science, nor was his house, or purse, ever shut against persons of that description: having received a better education than gardeners usually get, he passed with the vulgar, and mankind in general, for a prodigy in knowledge: he was temperate in his way of living, hence he attained a green old age: he had a wife, who was kind and most affectionate, by whom he had one son and three daughters, the eldest of whom, Ann, was so eminent as an artist, that her botanical drawings are esteemed as *chef d'œuvres*: he had the good fortune to live to see them all well-married; I mean, as relates to both happiness and competency; and his son, the present Mr. Lee, who inherits the wisdom, liberality, and virtues of his father, is blessed with a daughter, who has drawn all the numerous heaths (*Ericas*), so as even to rival her aunt.

I shall conclude my memoirs with wishing the surviving family all prosperity and happiness, and my readers many particles of that sacred flame, which animated old Lee in his love for plants, to the very latest period of his life.



C O N T E N T S.



PART I.

Chap.	Page
I. <i>Of the Seven Parts of Fructification</i>	1
II. <i>Of the Calyx</i>	2
III. <i>Of the Corolla</i>	4
IV. <i>Of the Stamina</i>	7
V. <i>Of the Pistillum</i>	8
VI. <i>Of the Pericarpium</i>	10
VII. <i>Of the Seeds</i>	11
VIII. <i>Of the Receptacle</i>	13
IX. <i>Of the distinct Characters of the Parts of Fructification</i>	14
X. <i>Of the most natural Structure of the Parts of Fructification</i>	16
XI. <i>Of the different Structures of the Calyx</i>	18
XII. <i>Of the different Structures of the Corolla</i>	21
XIII. <i>Of the different Structures of the Stamina</i>	24
XIV. <i>Of the different Structures of the Pistillum</i>	26
XV. <i>Of the different Structures of the Pericarpium</i>	29
XVI. <i>Of the different Structures of the Seeds</i>	31
XVII. <i>Of the different Structures of the Receptacle</i>	32
XVIII. <i>Of the Singularities in the Structure of the Parts of Fructification</i>	33
XIX. <i>Of Aggregate Flowers</i>	36
XX. <i>Of Luxuriant Flowers, commonly called double</i>	39
XXI. <i>Of the Sexes in Plants</i>	46

PART II.

Chap.	Page
I. <i>Of the Sexual System, and its Origin</i>	49
II. <i>Explanation of the Titles of the Twenty-four Classes</i>	58
III. <i>Explanation of the Titles of the Orders</i>	62
IV. <i>Of the first Class, Monandria</i>	65
V. <i>Of the second Class, Diandria</i>	66
VI. <i>Of the third Class, Triandria</i>	67
VII. <i>Of the fourth Class, Tetrandria</i>	68
VIII. <i>Of the fifth Class, Pentandria</i>	70
IX. <i>Of the sixth Class, Hexandria</i>	74
X. <i>Of the seventh Class, Heptandria</i>	76
XI. <i>Of the eighth Class, Octandria</i>	77
XII. <i>Of the ninth Class, Enneandria</i>	78
XIII. <i>Of the tenth Class, Decandria</i>	79
XIV. <i>Of the eleventh Class, Dodecandria</i>	81
XV. <i>Of the twelfth Class, Icosandria</i>	82
XVI. <i>Of the thirteenth Class, Polyandria</i>	84
XVII. <i>Of the fourteenth Class, Didynamia</i>	86
XVIII. <i>Of the fifteenth Class, Tetradynamia</i>	89
XIX. <i>Of the sixteenth Class, Monadelphia</i>	92
XX. <i>Of the seventeenth Class, Diadelphia</i>	95
XXI. <i>Of the eighteenth Class, Polyadelphia</i>	100
XXII. <i>Of the nineteenth Class, Syngenesia</i>	101
XXIII. <i>Of the twentieth Class, Gynandria</i>	106
XXIV. <i>Of the twenty-first Class, Monœcia</i>	109
XXV. <i>Of the twenty-second Class, Diœcia</i>	111
XXVI. <i>Of the twenty-third Class, Polygamia</i>	114
XXVII. <i>Of the twenty-fourth Class, Cryptogamia</i>	116
XXVIII. <i>Of the Appendix</i>	119
XXIX. <i>Of Generic Distinctions</i>	120
XXX. <i>By what Parts of Fructification the Genus may with the most certainty be determined</i>	122
XXXI. <i>Of the Genera rejected by the Sexual System, as not established on the Fructification</i>	125

C O N T E N T S.

xxiii

Chap.	Page
XXXII. <i>Of the Genera rejected by the System, as grounded on the Variations of some Parts only of the Fructification.....</i>	129
XXXIII. <i>Of the Genera rejected by the System, as grounded on a Difference in the Fruit only.....</i>	132
TABLE I. <i>Changes in the Names of Genera, arranged in Alphabetical Order.....</i>	136
TABLE II. <i>The Linnæan Genera, with References and Synonymes</i>	167
TABLE III. <i>The Linnæan Genera, alphabetically arranged, with the Classical and English Names, and accented; with a Reference also to their Classes and Orders</i>	299
TABLE IV. <i>An Alphabetical Catalogue of English and Scotch Names of Plants, from the most approved Authors, referred to their respective Genera.....</i>	353
TABLE V. <i>The Names of Plants, in Latin and French.....</i>	390

PART III.

Chap.	
I. <i>Of Vegetables, and their Kinds.....</i>	413
II. <i>Of Roots.....</i>	415
III. <i>Of the Herb.....</i>	416
IV. <i>Of the Trunk.....</i>	417
V. <i>Of simple Leaves.....</i>	423
VI. <i>Of compound Leaves.....</i>	432
VII. <i>Of determinate Leaves.....</i>	434
VIII. <i>Of the Fulcra of Plants.....</i>	438
IX. <i>Of the Hybernacula of Plants.....</i>	439
X. <i>Of the Habit of Plants.....</i>	440
XI. <i>Of Placentation.....</i>	441
XII. <i>Of Radication.....</i>	443
XIII. <i>Of Ramification.....</i>	444
XIV. <i>Of Intorsion.....</i>	445
XV. <i>Of Gemmation.....</i>	447
XVI. <i>Of Foliation.....</i>	448
XVII. <i>Of Stipulation.....</i>	450
XVIII. <i>Of Pubescence.....</i>	452

Chap.	Page
XIX. <i>Of Glandulation</i>	455
XX. <i>Of Lactescence</i>	456
XXI. <i>Of Inflorescence</i>	457
XXII. <i>Of Specific Distinctions</i>	458
XXIII. <i>Of Varieties</i>	466
TABLE VI. <i>Definitions of the Botanic Terms, in Latin and English, from the Termini Botanici of Linnæus</i> ...	474
TABLE VII. <i>A Botanical Dictionary</i>	505
TABLE VIII. <i>Derivations of the Botanic Terms, alphabetically arranged</i>	514

INTRODUCTION
TO THE
SCIENCE OF BOTANY.

PART THE FIRST.

CHAPTER I.

OF THE SEVEN PARTS OF FRUCTIFICATION.

BY fructification we are to understand both the *flower* and *fruit* of plants, which cannot well be separated; for though the fruit does not swell and ripen till after the flower is fallen, its rudiment, or first beginning, is in the flower, of which it properly makes a part. *Linnaeus* defines “the fructification to be a temporary part of vegetables, allotted to re-production, terminating the old vegetable, and beginning the new.” It consists of seven principal parts, viz.

1. CALYX, vulgarly called *empalement*, or *flower-cup*.
2. COROLLA, *foliation*, vulgarly called *the flower*.
3. STAMINA, vulgarly called *the chives*.
4. PISTILLUM, vulgarly styled *the pointal*.

5. PERICARPIUM, *seed-vessel*.
6. SEMINA, *the seeds themselves*.
7. RECEPTACLE*, *base*, on which the fructification is seated.

All these parts, and their several uses, will be particularly explained in the following chapters; and it is sufficient to observe here, that the four first, viz. Calyx, Corolla, Stamina, and Pistillum, are properly parts of the *flower*; and the three last, Pericarpium, Semina, and Receptacle, parts of the *fruit*; and that it is from the number, proportion, positions, and other circumstances attending these parts of fructification, that the classes of vegetables, and the genera they contain, are to be characterized according to the Sexual System.

CHAP. II.

OF THE CALYX.

THE CALYX is, according to Linnæus, “the termination of the *cortex*, or *outer bark* of the plant; which, after accompanying the trunk or stem through all its branches, breaks out with the flower, and is present in the fructification in this new form.” Its chief use is to enclose and protect the other parts†. It has received different appellations, according to the circumstances with which it is attended, viz.

1. PERIANTHIUM, a *flower-cup*, when its station is close to the fructification. - If it includes the stamina, and not the germen,

* For the derivations of these terms, vide Doctor Thornton’s “Grammar of Botany.”

EDITOR.

† It sometimes serves the office of pericarpium, as in the LAMIUM, *nettle*, and frequently accompanies the fruit. In the PATAGONULA and *egg plant* it is observed to grow to a larger size in the fruit than it had in the flower. EDITOR.

it is the *perianthium of the flower*;...if the germen, but not the stamina, the *perianthium of the fruit*;...but if it includes both, it is the *perianthium of the fructification*.

2. INVOLUCRUM, a *cover*, when stationed at the foot of an umbel, at a distance from the flower*; it is an universal involucre, if it is under the universal umbel;...or a partial one, if under a partial†.

3. AMENTUM, *catkin*, when it proceeds from one common receptacle, resembling the chaff of an ear of corn.

4. SPATHA, *sheath*, when it bursts lengthways, and puts forth a *spadix*‡.

5. GLUME, *husk*, in grasses, which it folds over with its valves; and the sharp point or beard issuing from the glume is called an *arista*.

6. CALYPTRA, a *veil*, in mosses, where it is placed over the *antheræ*, tops of the *stamina*, and is hooded like a monk's cowl.

7. VOLVA, from its *involving*, or *infolding*, in the *fungi*, or *mushroom tribe*, where it is membranaceous, and rent on all sides.

It is sometimes difficult to distinguish a *calyx* from the *bractea*, *floral leaf*§, such as is found to accompany the fructification

* Sometimes this part does not attend an umbel, as in the *anemony*. EDITOR.

† For the definition of an umbel, vide p. 13. EDITOR.

‡ *Spadix* properly signifies the receptacle of a palm: see p. 14. But *spatha* is not confined only to such plants as have a *spadix* in this sense of the term, but is applied to NARCISUS, GALANTHUS, PANCRATIUM, and many others, whose flower-stalks come out of a sheath. *Spadix* therefore is here to be understood in a more general sense: agreeable to such latitude we shall find it used in CAILA, DRACONTIUM, POTHOS, ARUM, and ZOSTERA, as well as in the PALMS. AUTHOR.

§ In many plants there are found green leaves amongst the flowers, that differ in shape from the original leaves of the plant. These are the *Bractea*, or *floral leaves*,

of the *TILIA*, *LAVENDULA*, *MELAMPYRUM*, and others. They may be distinguished by this certain rule, that a *calyx* always withers when the fruit is ripe, if not before; but the *bractea* will remain longer. Without attending to this, mistakes might easily be made in *HELLEBORUS*, *NIGELLA*, *PASSIFLORA*, *HEPATICA*, *PEGANUM*, and others, in which the calyx is wanting. The distinction between a *calyx* and *corolla* in doubtful cases will be treated of in the next chapter. In many flowers the calyx is deciduous, dropping off the instant the flower begins to expand; this is the case with *EPIMEDIUM* and *PAPAVER*.

CHAP. III.

OF THE COROLLA.

THE COROLLA is said by Linnæus "to be the termination of the *liber*, or *inner bark*, continued to, and accompanying the fructification in this new form of painted leaves."

Its *use* is the same as that of the calyx, serving as an inner work of defence, for the parts it encloses, as the calyx, which is usually of stronger texture, does for an outer one.

The leaves of which the corolla consists are called *petals*; by which appellation they are conveniently distinguished from the green leaves of the plant, with which they might else be con-

here spoken of. They are commonly situated on the flower-stalks, and sometimes so near to the flower, as to be mistaken for its calyx. AUTHOR.

founded*. The petal is defined by *Linnaeus* "as a corollaceous covering to the flower," meaning that it encloses and protects in the manner of a *corolla*, or *wreath*. If the corolla be

MONOPETALOUS, of one petal; it consists of two parts, viz. The *tube*, or lower part, which is usually tube-shaped; and the *limb*, or upper part, which usually spreads wider. And the limb again, according to its figure, is either *campanulate* (*bell-shaped*), that is, bellying out, and without a tube;...*infundibuliform* (*funnel-shaped*), that is, of the figure of a cone, and standing on a tube;...*hypocrateriform* (*salver-shaped*), that is, plain or flat, and standing on a tube;...*rotato-plane* (*wheel-shaped and flat*), without a tube;...or *ringent* (*gaping*), that is, irregular and personated with two lips. But if the corolla be

POLYPETALOUS, of many petals; each petal consists of *unguis*, a *claw*, which is the lower part fastened to the base; and *lamina*, a *thin plate*, which is the upper part, and usually spreading. A polypetalous corolla is *cruciform* (*cross-shaped*) when it consists of four petals that are equal and spreading;...and *papilionaceous* (*butterfly-shaped*) when it is irregular, consisting of four petals,

* *Petal* (in the Greek *πεταλον*) signifies leaves in general: but there being another *Greek* word (*φυλλον*) nearly of the same signification, the modern botanists have borrowed this to express the leaves of the flower. The ancients seem to have had no distinct term in use to express this part of the fructification. Thus *Virgil*, in describing his *amellus*, which is a species of *aster*, the flower of which has a yellow middle, and purple rays, calls it a golden flower, surrounded with purple leaves.

*Aureus ipse Flos, sed in foliis, quæ plurima circum
Funduntur, violæ subluet purpura nigræ.*

GEORG. IV.

This loose expression, which is chargeable rather on the language than the poet, has misled all his translators; as is rightly observed by *Martin*, in his note on this passage. Thus *Addison* makes the real leaves of the plant purple:

The flower itself is of a golden hue,
THE LEAVES inclining to a darker blue.
THE LEAVES shoot thick about the root, and grow
Into a bush; and shade the turf below.

ADDISON.

of which the under one resembles the keel of a ship, the upper one rises, and the two side ones stand single*.

There belongs also to the corolla a part called the *nectarium*, which has been but newly distinguished, having been by former botanists confounded with the *petals*. It is by *Linnaeus* defined to be “the part which bears the honey, and belongs to the flower only.” This part affords a wonderful variety in the manner of its appearance. In some plants it is very large, as in the *NARCISSUS* and *AQUILEGIA*; in the former of which the *cup*, and in the latter the *horns*, are *nectaria*: in others it is scarce discoverable, even with glasses. In some plants it is united with, and makes part of the petals: in others it is detached from them. Its shape and situation are also as various. Its use is not known, unless the supposition of its secreting the honey may be depended upon†.

Between the *calyx* and *corolla* nature has put no absolute limits; as is plain from the *DAPHNIS*, in which plant they grow together, and are united in the margin, like a leaf of the *BUXUS*; but they may be commonly distinguished by their position in respect of the *stamina*, the *petal* and *stamina* being ranged alternately; whereas the segments of the *calyx* and the *stamina* answer to each other. That this is their natural situation, appears from the complete flowers in the classes *tetrandria*‡ and *pentandria*§: And the use of applying this rule will be found in the instances of *CHENOPodium*, *URTICA*, and *PARIETARIA*; where it decides, that the single cover in those *genera* is a *perianthium*, and that it is the *co.olla* that is wanting. Should we infer, where only one of the two covers appears, that it is a *corolla*, because that is a more principal part, there would be no certainty from such an infer-

* The under petal is called the *CARINA*, *keel*; the two side petals, the *ALÆ*, *wings*; and the upper petal, *VEXILLUM*, *banner*. EDITOR.


† There seems much confusion in this part: in fact, whatever is not calyx, corolla, stamina, and pistillum, is *nectary* with botanists, whether it secretes honey, or not. EDITOR.

‡ This is explained in Part II. Chap. VII.

§ This is explained in Part I. Chap. VIII.

ence: as is evident from the AMMANIA, ISNARDA, PEPLIS, RUELLIA and CAMPANULA, in all which the *corolla* is often found wanting, but not the *calyx*.

That the calyx, as proceeding from the *cortex* of the plant, is coarser and thicker than the corolla, which is produced by the soft, pliant, coloured *liber*, is obvious to every one. But there are no limits determinable from any such circumstances, unless it be from the colour; and even this is not sufficient; for the perianthium of the BARTSIA is crimson-coloured; and there are also many flowers whose *corollas* are coloured, when in the state of flowering, but which afterwards harden and turn green, and remain on the plant like a calyx; as for instance, the HELLEBORUS and ORNITHOGALUM. The EUPHORBIA has likewise deceived many, who have described it as monopetalous, taking the *calyx* for the *corolla*.



CHAP. IV.

OF THE STAMINA.

THE STAMINA are the male part of the flower. *Linnaeus* defines them as a “viscus of the plant, designed for the preparation of the *pollen* ;” of which we shall speak presently.

Each single *stamen* consists of two parts*, *viz.*

1. FILAMENTUM, *the filament or thread*; which serves to elevate the *anthera*, or *summit*, and at the same time connects it with the flower.

2. ANTHERA, the *summit* itself; which contains within it the pollen, and when come to maturity discharges the same.

* This is not always the case, as some stamens are complete, having nothing but the anther, as in the CANNA INDICA, *Indian canna*, &c. EDITOR.

The POLLEN, *meal*, contained within the antheræ, is a fine dust secreted therein, and destined for the impregnation of the *germen*; of which part we shall speak in the next chapter.

The stamina being, as I have said, the male part of the flower, the construction and distribution of the *Sexual System* is principally founded upon, and regulated by it; as will appear in the explanation of the System. It is sufficient to observe here, that such flowers as want this part are called *female*; such as have it, but want the female part, described in the next chapter, *male*; such as have them both, *hermaphrodite**; and such as have neither, *neuter*†.

CHAP. V.

OF THE PISTILLUM.

THE PISTILLUM is the female part of the flower: it is defined by *Linnaeus* “as a viscus of the plant, designed for the reception of the pollen.” It consists of three parts‡.

1. The GERMEN; which is the rudiment of the fruit accompanying the flower, but not yet arrived at maturity.

* This odious term should be expelled the lovely science of botany, and the term *bisexual* substituted in its place; for the analogy does not hold, nor are there any truly *hermaphrodite* flowers, as with animals. The two sexes are contiguous, and marry; but with snails, and earth worms, which have the two sexes in the same person, these are really *hermaphrodites*; but are not competent to reproduce of themselves, but have relationships with others of the same species; therefore, the analogy here does not hold. EDITOR.

† Neuter, or *barren*.

‡ This is not always the case, as the style in many instances is wanting, as in *PAFAVER*, *poppy*; *TULIPA*, *tulip*, &c. EDITOR.

2. The *STYLE*, which is the part that serves to elevate the stigma from the germen.

3. The *STIGMA*, which is the summit of the pistillum, and covered with a moisture for the breaking of the pollen.

It has been said in the last chapter, that the *pollen* was destined for the impregnation of the germen: this is performed in the following manner. The *antheræ*, which at the first opening of the flower are whole, burst open soon after, and discharge the pollen, which dispersing itself about the flower, part of it lodges on the surface of the stigma, where it is detained by the moisture with which that part is covered*; and each single grain or atom of the *pollen* bursting and dissolving in this liquor, as it has been observed to do by the microscope, is supposed to discharge something still more subtle, that impregnates the germen below. What the substance is that is so discharged, and whether it actually passes through the style into the germen, seems yet undetermined†, it being difficult to observe such minute parts: but whatever be the operation by which Nature produces the effect in question, the cause as far as it has been here explained, is scarce^{ly} disputable; and accordingly we see, that after this impregnation[^], when the parts of the flower that have done their office are fallen away, the germen swells to a fruit big with seeds, by which the species is propagated. The *pistillum* being, as I have said, the *female part* of the flower, is of great consequence in the *Sexual System*, as well as the *male part*, as will appear when the System comes to be explained.

* This is beautifully seen in the *AMARYLLIS FORMOSISSIMA*, on whose stigma may be observed a large limpid globule of an adhesive nature, to catch the fertilising pollen. Vide Dr. Thornton's "New Illustration of the Sexual System, with a Dissertation on the Sexes of Plants." EDITOR.

† This dispute is now settled. The *pollen*, Linnæus, in his Dissertation on the Sexes of Plants, has proved, does not pass the style, as in the *MIRABILIS*, *marvel of Peru*, where each globule of pollen is larger than the style, but only the most subtle exhalation. EDITOR.

CHAP. VI.

OF THE PERICARPIUM.

THE PERICARPIUM, *seed-vessel*, is the germen described in the last chapter, grown to maturity. It is defined by *Linnaeus* "as a viscus of the plant filled with seeds, which it discharges when ripe."

It is distinguished, according to the circumstances that attend it, by the following appellations.

1. CAPSULA, a *capsule*, is a hollow pericarpium, which cleaves or parts in some determinate manner... The enclosure of the capsule, which surrounds and covers the fruit externally, is called a *valvule*; the partitions which divide the capsule into sundry compartments or cells, *dissepiments*; the substance which passes through the capsule, and connects the several partitions and seeds, *columella*; and the cells, or hollow compartments of the capsule in which the seeds are lodged, *loculaments*.

2. SILIQUA, a *pod*, is a pericarpium of two *valves**, wherein the seeds are fastened along both the sutures or joinings of the valves.

3. LEGUMEN, a *pod* also, is a pericarpium of two valves, wherein the seeds are fastened along one suture only.

4. CONCEPTACULUM, a *conceptacle*, is a pericarpium of a single valve, which opens on one side lengthways, and has not the seeds fastened to it.

* The author has called the separate pieces which constitute the pericarp, *valvules*, and those of the other kinds, *valves*. This distinction of names is seldom or never observed. EDITOR.

5. DRUPA, a *drupe*, is a fleshy or pulpy pericarpium, without valve, containing a stone.

6. POMUM, a *pome*, is a fleshy or pulpy pericarpium, without valve, containing a capsule.

7. BACCA, a *berry*, is a fleshy or pulpy pericarpium, without valve, the seeds within which have no other covering.

8. STROBILUS, a *strobile*, is a pericarpium formed of an *amentum**.

CHAP. VII.

OF THE SEEDS.

THE SEED, according to the definition of *Linnaeus*, "is a deciduous part of the vegetable, the rudiment of a new one, quickened for vegetation by the sprinkling of the pollen." Its distinctions are,

A SEED, *properly so called*, which is a rudiment of a new vegetable, furnished with sap, and covered with a bladdery coat or tunic. It consists of,

1. CORCULUM, the first principle of the new plant within the seed.

2. PLUMULA, a scaly part of the corculum, which ascends.

3. ROSTELLUM, a plain part of the corculum, which descends.

4. COTYLEDON, a side lobe of the seed, of a porous substance, and perishable.

* See Chap. II.

5. HILUM, an external mark or scar on the seed, where it was fastened within the fruit.

6. ARILLUS, the proper exterior coat, or tunic of the seed, which comes off of itself.

7. CORONULA, *the little crown* of a seed, which is either CALY-
CULUS, the *calyx* of a floret, adhering to the seed, and assisting
it to fly, or PAPPUS, *a down*, which is a feathery, or hairy crown,
answering the same end, and connected with the seed by STIPES,
*a trunk**, which here signifies a thread on which the down is
raised and supported.

8. ALA, *wing*, a membrane affixed to the seed, and which by
its flying helps to disperse it.

9. NUX, *a nut*, which is a seed enclosed with an *osseous epider-*
mis, a bony or hard outer skin, commonly called the *shell*.

10. PROPAGO, which is the seed of a moss, first discovered by
Linnæus, who peeled off the bark, and detected it in the year
1750. These seeds have neither tunic nor cotyledon, but con-
sist only of the plumula of a naked corculum, where the rostel-
lum is inserted into the calyx of the plant.

* Sometimes, however, this part, the *stipes*, is wanting, and the *pappus* is imme-
diately connected with the seed, when, like the *anther* and *stigma*, it is termed *sessile*.

CHAP. VIII.

OF THE RECEPTACLE.

THE RECEPTACLE is the base, which connects the other six parts of fructification. Its various appellations are as follow.

I. A PROPER RECEPTACLE is that which belongs only to the parts of a single fructification: and this is called...1. *A receptacle of the fructification*, when it is common to both flower and fruit; ...2. *A receptacle of the flower*, when it is a base to which the parts of the flower only are fastened, without the germen;...3. *A receptacle of the fruit*, when it is a base for the fruit only, remote from the receptacle of the flower;...4. *A receptacle of the seeds*, when it is a base that fastens the seeds within the pericarpium.

II. A COMMON RECEPTACLE is that which connects many florets in such a manner, as that the taking away any of them would cause an irregularity. *Palea, a chaff*, is a thin substance, springing from the receptacle to part the florets.

III. UMBELLA, *an umbel*, is a receptacle which, from a common centre, runs out into thread-shaped foot-stalks, of proportionate lengths...It is called *a simple umbel*, when it has no subdivisions; *a compound umbel*, when each foot-stalk is terminated by an *umbellula*, or *little umbel*; and in this case the umbel that bears the umbellula on its foot-stalks, is called a *universal umbel*; and the umbellula which proceeds from the universal umbel, a *partial umbel*.

IV. CYMA, a *cyme*, is a receptacle that runs into long fastigi-

ate peduncles*, proceeding from the same universal centre, but with irregular partial ones.

V. SPADIX is the receptacle of a palm†, produced within a *spatha*, or *sheath*, on the branches that bear fruit.

CHAP. IX.

OF THE DISTINCT CHARACTERS OF THE PARTS OF
FRUCTIFICATION.

THE parts of fructification, with their subdivisions, having been explained separately in the preceding chapters, we shall here give a view of them all together, with the proper distinguishing character assigned to each by *Linnaeus*, beginning with the vegetable itself.

The essence of the *vegetable* consists in its *fructification*;...the essence of the *fructification* consists in the *flower* and *fruit*;...the essence of the *flower* consists in the *anthera* and *stigma*;...the essence of the *fruit* consists in the *seeds*. We shall give now a short *definition* of THESE PARTS.

POLLEN is the fine powder of vegetables, designed to burst in

* *Peduncles*, *flower-stalks*, are called *fastigate*, when their lengths are so proportioned, that the flowers which they support form an even surface. AUTHOR.

† This is the proper sense of the term, as employed by the ancients: but *spadix* is now used in a more general sense, viz. to express all flower-stalks that come out of a *spatha*. See the note on this subject in Chap. II. This definition, by *Linnaeus*, therefore, appears to be too strict. AUTHOR.

a liquor appropriated to that purpose*, and discharge thereon, by its elastic force, a substance not distinguishable by the naked eye.

A SEED is a deciduous part of a plant, fraught with the rudiment of a new plant, and quickened by the pollen.

ANTHERA is a vessel that produces and discharges the pollen.

PERICARPIUM is a vessel that produces and discharges the seeds.

FILAMENTUM is the foot-stalk that supports† the anthera; and fastens it to the vegetable‡.

GERMEN is the rudiment of the pericarpium, not yet arrived at maturity.

STIGMA is the moistened summit of the germen: its existence is chiefly at the time when the anthera is discharging its pollen.

STYLUS is the foot-stalk of the stigma, that connects it with the germen.

COROLLA and CALYX are the *teguments* or *covers* of the stamina and pistillum; the calyx arising from the *cortical epidermis*, or *outer bark*, and the corolla from the *liber*, or *inner bark*.

RECEPTACULUM is that part which connects the parts before mentioned§.

From these characters the following principles may be deduced.

1. That every *vegetable* is furnished with *flower* and *fruit*; there being no species where these are wanting.

* If the pollen be placed on a damp plate, all its particles, which have determinate shapes, will explode. The moisture on the stigma of plants effects the same purpose. EDITOR.

† Elevates. EDITOR.

‡ And attaches it to the flower. EDITOR.

§ Is the connecting medium betwixt the PEDUNCLE, *flower-stalk*, and *flower*.

2. That there is no *fructification* without *anthera*, *stigma*, and *seed*.

3. That the *anthera* and *stigma* constitute a *flower*, whether the covers are present or wanting.

4. That the *seed* constitutes a *fruit*, whether there be a *pericarpium* or not.

In respect to the *seed*; its essence consists in the *corculum*, which is fastened to the cotyledon, and involved therein, and closely covered with its proper tunic.

The essence of the *corculum* consists in the *plumula*, which is the vital speck of the plant itself, extremely small in its dimensions, but increasing like a bud in growth. The *rostellum*, however, must be included, being the base of the *plumula*, which descends, and strikes root, being the part originally contiguous to the mother plant.

That the *propagines*, or seeds of mosses, consist only of the *plumula* and *rostellum*, has been already shown*.

CHAP. X.

OF THE MOST NATURAL STRUCTURE OF THE PARTS OF FRUCTIFICATION.

IN considering the structure of the parts of fructification, the principal objections to be attended to are, 1. The *number* of each part. 2. Its *figure*. 3. Its *proportion*; by which is to be understood its height in respect to the rest: and 4. Its *situation*; which will include also its *insertion* and *connexions*. As to any other differences, such as a difference in the size, colour, smell, or taste, it is not safe to allow any weight to them,

* See Chap. VII.

as they might lead us to make distinctions not justifiable by the true principles of the science.

As the number, figure, proportion, and situation of the parts are variable, we shall consider, 1. THE MOST NATURAL STRUCTURE, or that which most frequently occurs; and this we shall make the subject of the present chapter. 2. THE DIFFERENCES in structure, arising from the variation of the parts in different plants, which will take up a few of the succeeding chapters; and 3. THE SINGULAR STRUCTURES, or such as are observed in a few genera only; for which we shall allot a chapter by itself.

The MOST NATURAL STRUCTURE of the parts, in respect to NUMBER is, to have the calyx divided into as many segments as the corolla;...the filaments equal in number to the segments of the corolla and calyx;...a single anthera on each filament;...the divisions of the pistillum equal in number to the cells of the pericarpium, or the receptacles of the seeds; the most common number *five* (whence the extent of the classes *Pentandria** and *Syngenesia*†);...and the corolla and calyx also *quinquifid*, *cut into five segments*.

In respect to FIGURE, to have the calyx less spreading than the corolla;...the corolla widening gradually;...the stamina and pistillum upright and tapering;...the pericarpium big with seeds, swelling and extending after the rest of the parts (the calyx excepted) are fallen off.

In respect to PROPORTION, to have the calyx less than the corolla;...the pistillum of equal length with the stamina in an upright flower, but longer in an inverted one;...if the flower slope downward, the stamina and pistillum inclining towards the under side; but if it slope upwards, placed close under the upper side.

In respect to SITUATION, to have the perianthium surrounding the receptacle;...the corolla placed on the receptacle, and alternate with the perianthium;...the filaments placed within the corolla, but corresponding with the perianthium;...the *antheræ*

* See Part II. Chap. VIII.

† See Part II. Chap. XXII.

seated on the tops of the filaments;...The germen possessing the centre of the receptacle;...the style standing on the top of the germen;...the stigma seated on the top of the style. When the stigma and style are fallen, the germen grows to a pericarpium, supported by the calyx, and including the seeds, which are affixed to the receptacle of the fruit. The receptacle of the flower is generally under the pericarpium, being not so often found to grow either round it, or over it.

CHAP. XI.

OF THE DIFFERENT STRUCTURES OF THE CALYX.

HAVING shown *the most natural Structure* of the parts of the fructification in the last chapter, we come now to their DIFFERENCES, or variations (which are the foundation of the *genera*), and their characters; and of these we shall treat in their order, beginning with the *calyx*.

The variations of the calyx, in respect to NUMBER, will take in the terms also that respect its *composition, parts, and segments*.

In respect to *number*, it is either *single*, as in PRIMULA, and most flowers;...*double*, as in MALVA, HIBISCUS, and BIXA;...or *wanting*, as in TULIPA, FRITILLARIA, and many of the liliaceous flowers*.

In respect to *composition*, it is either *imbricate*, that is, composed of various scales, lying over each other, as in HIERACIUM,

* It is to be hoped, that the student will not be deterred by these nice observations and distinctions in the science of botany. Such as may find their memories too much fatigued may pass on at once to part second, the Sexual System, p. 72.

SONCHIUS, and CAMELLIA;...*squarrose*, that is, composed of scales, divaricated on all sides, and spreading widely open, as in CARDUUS, ONOPORDUM, and CONYZA;...*auctus*, *augmented*; that is, having a series of distinct leaves, shorter than its own, that surround its base externally, as in COREOPSIS, BIDENS, CREPIS, and DIANTHUS;...or *multiflorous*, *many flowered*, that is, common to many florets, as in SCABIOSA, and in the plants of the class *Syngenesia**.

In respect to its *parts*, it is either *monophyllous*, of one leaf, as in DATURA and PRIMULA;...*diphyllous*, of two, as in FUMARIA, and FUMARIA BULBOSA;...*triphyllous*, of three, as in TRADESCANTIA;...*tetraphyllous*, of four, as in SAGINA, EPIMEDIUM, and in the plants of the class *Tetradynamia*†;...*pentaphyllous*, of five, as in CISTUS, ADONIS, and CERBERA;...*hexaphyllous*, of six, as in BERBERIS;...or *decaphyllous*, of ten, as in HIBISCUS.

In respect to its *segments* (which chiefly concern the monophyllous calyx) it is either *integer*, *whole*, as in GENIPA;...*bifid*, *divided in two segments*, as in UTRICULARIA;...*trifid*, *in three*, as in ALISMA, and CLIFFORTIA;...*quadrifid*, *in four*, as in RHINANTHUS;...*quinquesid*, *in five*, as in NICOTIANA;...*sexsid*, *in six*, as in PAVIA;...*octosid*, *in eight*, as in TORMENTILLA;...*decemsid*, *in ten*, as in POTENTILLA, and FRAGARIA;...or *duodecemsid*, *in twelve*, as in LYTHRUM.

The variations of the *calyx*, in respect to *FIGURE*, will also include the terms respecting its *equality*, *margin*, and *apex*, or *top*.

In respect to *figure*, it is either *globose*, *globe-shaped*, as in CUCUBALUS;...*clavate*, *club-shaped*, as in SILENE;...*reflex*, *bent back*, as in ASCLEPIAS;...or *erect*, *upright*, as in PRIMULA, and NICOTIANA.

In respect to *equality*, it is either *equal* as in LYCHNIS;...*unequal*, as in HELIANTHEMUM; or with the segments *alternately shorter*, as in TORMENTILLA, and POTENTILLA.

In respect to its *margin*, it is either *integerrimus*, *very entire*, as

* See Part II. Chap. XXII.

† See Part II. Chap. XVIII.

in most plants;...*serrate, sawed*, as in some species of *HYPERICUM*;...or *ciliate, fringed with hairs, like an eye-lash*, as in some species of *CENTAUREA*.

In respect to its *apex, or top*, it is either *acute, sharp*, as in *PRIMULA*, and *ANDROSACE*;...*acuminate, pointed*, as in *HYOSCYAMUS*;...*obtuse, blunt*, as in *NYMPHÆA*, and *GARCINIA*;...or with one of its indents *lopped off*, as in *VERBENA*.

In respect to *PROPORTION*, it is either *longer* than the corolla, as in *AGROSTEMMA*, *SAGINA*, and some species of *ANTIRRHINUM*;...*equal* to it, as in some species of *CERASTIUM*;...or *shorter*, as in *SILENE*.

In respect to *SITUATION*, it is either a calyx of the *flower*, as in *LINNÆA* and *MORINA*;...of the *fruit*, as in *LINNÆA* and *MORINA**;...or of the *fructification*, as in *PÆONIA*.

The *DURATION* of the calyx may also be considered. In respect to which it is either *caducous, falling off at the first opening of the flower*, as in *PAPAVER* and *EPIMEDIUM*;...*deciduous* with the corolla, as in *BERBERIS*, and in the plants of the class *Tetradynamia*†;...or *persisting*, till the fruit is come to maturity, as in the plants of the class *Didynamia*‡.

VARIATIONS OF AN INVOLUCRUM.

The preceding varieties of the calyx chiefly respect a perianthium. An *involutrum* is either *monophyllous*, as in *BUPLEURUM*;...*diphyllous*, as in *EUPHORBIA*;...*triphyllous*, as in *BUTOMUS* and *ALISMA*;...*tetraphyllous*, as in *CORNUS*;...*pentaphyllous*, as in *DAUCUS*;...or *hexaphyllous*, as in *HÆMANTHUS*.

* The *Linnaea* and *Morina* have each of them two *calyxes*, one of the flower, the other of the fruit; which is the reason of their being given as instances of both cases.

† See Part II. Chap. XVIII.

‡ See Part II. Chap. XVII.

VARIATIONS OF A SPATHA.

A *spatha* is either *monophyllous*, as in NARCISSEUS;...*diphyllous*, as in STRATIOTES;...or *imbricate*, as in MUSA.

 CHAP. XII.

OF THE DIFFERENT STRUCTURES OF THE COROLLA.

THE variations of the corolla, in respect to NUMBER, concern either *petals*, or *laciniæ*, *segments*: the variations of the nectarium shall be given separate.

The corolla, in respect to its *petals*, is either *monopetalous*, or *consisting of one petal*, as in CONVULVUS and PRIMULA;...*dipetalous*, of two, as in CIRCEA and COMMELINA;...*tripetalous*, of three, as in ALISMA and SAGITTARIA;...*tetrapetalous*, of four, as in the class *Tetradynamia**;...*pentapetalous*, of five, as in umbelliferous plants†;...*hexapetalous*, of six, as in TULIPA, LILIUM, PODOPHYLLUM;...*enneapetalous*, of nine, as in THEA, MAGNOLIA, and LIRIODENDRON;...or *polypetalous*, of many, as in NYMPHEA.

In respect to its *laciniæ* (which concern rather the monopetalous than the polypetalous, being but rarely observed in the latter) it has either *two*, as in ALSINE and CIRCEA;...*three*, as in

* See Part II. Chap. XVIII.

† The umbelliferous plants are in the order *Digynia* of the class *Pentandria*; see Part II. Chap. VIII.

HOLOSTEUM and HYPECOUM ;...*four*, as in LYCHNIS ;...or *five*, as in RESEDA.

The variations of the corolla, in respect to FIGURE, will include what also concerns its *Equality*, and its *Margin*.

In respect to *Figure*, it is either *undulate, waved*, as in GLORIOSA ;...*plicate, folded*, as in CONVULVULUS ;...*revolute, rolled back*, as in ASPARAGUS and MEDEOLA ;...or *tort, twisted*, as in NERIUM, ASCLEPIAS, and VINCA. Its more considerable variations, in respect to figure, have been already shown in Chap. III.

In respect to *Equality*, it is either *equal*, as in PRIMULA ;...*unequal*, as in BUTOMUS ;...*regular*, as in AQUILEGIA ;...or *irregular*, as in ACONITUM and LAMIUM.

In respect to its *Margin*, it is either *crenate, notched*, as in LINUM ;...*serrate, sawed*, as in TILIA and ALISMA ;...*ciliate, fringed*, as in RUTA, MENYANTHES, and TROPÆOLUM ;...*denticulate* between the segments ; that is, having a *denticulus, or little jag*, at the bottom of the divisions, as in SAMOLUS and SIDEROXYLUM ;...or with a *hairy surface*, as in MENYANTHES and LASIANTHUS, a species of HYPERIUM.

In respect to PROPORTION, it may be very *long*, as in CATESBÆA, SIPHONANTHUS, BRUNSFELSIA, and CRANIOLARIA ;...or very *short*, as in SAGINA, CENTUNCULUS, and RIBES.

In respect to SITUATION, the base of the corolla is usually close to the perianthium, if there be one. It is, indeed, separated from it by the germen, in ADOXA, SANGUISORBA and MIRABILIS ; but these instances are very rare.

In respect to DURATION, it is either *persisting, lasting till the fruit is ripe*, as in NYMPHÆA ;...*caducous, dropping as soon as the flower is blown*, as in ACTÆA and THALICTRUM ;...*deciduous, dropping off with the flower*, which is the most common ;...or *marcescent, withering, but not falling*, as in CAMPANULA, ORCHIS, CUCUMIS, CUCURBITA, and BRYONIA.

VARIATIONS OF THE NECTARIUM.

It has been already said, Chap. III. that the *nectarium*, by the former botanists, had been confounded with the petals; but though it commonly attends upon, and makes part of the corolla, it is often found distinct from it, as in the instances of ACONITUM, AQUILEGIA, HELLEBORUS, ISOPYRUM, NIGELLA, GARIDELLA, EPI-MEDIUM, PARNASSIA, THEOBROMA, CHERLERIA, and SAUVAGESIA; which sufficiently proves that it should be distinguished from the petals. The *nectarium* affords very singular varieties, especially if it grows distinct from the petals. It admits of the following principal distinctions.

CALCARIATE *nectaria*, such as resemble a *calcar*, or *spur*; and these are either in *monopetalous corolla*, as in ANTIRRHINUM, VALERIANA, PINGUICULA, and UTRICULARIA;...or in POLYPETALOUS, as in ORCHIS, DELPHINIUM, VIOLA, IMPATIENS, and FUMARIA.

Nectaria that lie within the SUBSTANCE of the *petals*, as in FRITILLARIA, LILIUM, SWERTIA, IRIS, HERMANNIA, UVULARIA, HYDROPHYLLUM, MYOSURUS, RANUNCULUS, BROMELIA, ERYTHRONIUM, BERBERIS, and VALISNERIA.

Nectaria that CROWN the *corolla*, as in PASSIFLORA, NARCISSUS, PANCRATIUM, OLAX, LYCHNIS, SILENE, CORONARIA, STAPELIA, ASCLEPIAS, CYNANCHIUM, NEPENTHES, CHERLERIA, CLUSIA, HAMAMELIS, and DIOSMA.

Nectaria of SINGULAR construction, as in RESEDA, CARDIOSPERMUM, AMOMUM, COSTUS, CURCUMA, GREWIA, URTICA, ANDRACHINE, EPIDENDRUM, HELICTERES, and SALIX.

CALYCINE *nectaria*, such as are found upon the *calyx*, as in TROPEOLUM, MONOTROPA, BISCUTELLA, and MALPIGHIA.

STAMINEOUS *nectaria*, such as attend the *stamina*; and these are either upon the *antheræ*, as in ADENANTHERA;...or upon the filaments, as in LAURUS, DICTAMNUS, ZYGOPHYLLUM, COMMELINA, MIRABILIS, PLUMBAGO, CAMPANULA, and ROELLA.

PISTILLACEOUS *nectaria*, such as accompany the *pistillum*. These

are upon the germen, as in HYACINTHUS, IRIS, BUTOMUS, CHIERANTHUS, HESPERIS, &c.

RECEPTACULACEOUS *nectaria*, such as join to the *receptacle*, as in LATHRÆA, HELXINE, COLLINSONIA, SEDUM, COTYLEDON, SEMPERVIVUM, &c. MERCURIALIS, KIGGELLARIA, CLUTIA, PHYLLANTHUS, MELIANTHUS, and DIOSMA.



·CHAP. XIII.

OF THE DIFFERENT STRUCTURES OF THE STAMINA.

THE stamina consisting each of a filament and an anthera, (see Chap. IV.), we shall speak first of the variations of the filaments.

As the terms respecting the NUMBER of the stamina will be explained in the chapters that treat of the sexual system, we shall omit here what concerns the number of the filaments themselves, to avoid repetition; but they are sometimes found to have *laciniæ*, *segments*; and these are either *two*, as in SALVIA;...*three*, as in FUMARIA;...or *nine*, as in the class *Diadelphia**.

The FIGURE of the filaments is either *capillary*, *like hairs*, as in PLANTAGO;...*plane*, *flat*, as in ORNITHOGALUM;...*cuneiform*, *wedge-shaped*, as in THALICTRUM;...*spiral*, *screw-shaped*, as in HIRTELLA;...*subulate*, *awl-shaped*, as in TULIPA;...*emarginate*, *nicked*, or *notched*, as in PORRUM;...*reflex*, *bent back*, as in GLORIOSA;...or *hirsute*, *hairy*, as in TRADESCANTIA, and ANTHERICUM.

The PROPORTION of the filaments is either *unequal*, as in DAPHNE, LYCHNIS, and SAXIFRAGA;...*irregular*, as in LONICERA, and the class *Didynamia*†;...*very long*, as in TRICHOSTEMA, PLANTAGO, and HIRTELLA;...or *very short*, as in TRIGLOCHIN.

* See Part II. Chap. XX.

† See Part II. Chap. XVII.

The SITUATION of the filaments, is either *opposite* to the leaves or segments of the calyx, as in URTICA;...or *alternate*, with them, as in ELÆAGNUS. In *monopetalous* flowers they are inserted into the corolla; but scarce ever in *polypetalous*. In the class *Icosandria** they are always inserted in the calyx, as they are also in EPILOBIUM, OENOTHERA, JUSSIÆA, LUDWIGIA, OLDENLANDIA, ISNARDA, AMMANIA, PEPLIS, LYTHRUM, GLAUX, and RHEXIA; and in some APETALOUS† flowers, as in ELÆAGNUS; but it is more common for them to be inserted into the *receptacle*, like the calyx and corolla.

VARIATIONS OF THE ANTHERÆ.

The NUMBER of the antheræ is either a *single* one to each filament, as in the generality of plants;...*one* common to *three*, as in CUCURBITA;...*one* to *five*, as in the whole class *Syngenesia*‡; *two* to each filament, as in MERCURIALIS;...*three* to each, as in FUMARIA;...*five* to *three* filaments, as in BRYONIA;...or *five* to each, as in THEOBROMA.

In some plants that have single antheræ to the filaments, some of the antheræ are *wanting*; thus *one* is wanting in CLEONIA and MARTYNIA;...*two* in PINGUICULA and VERBENA;...*three* in GRATIOLA, and in some BIGNONIAS and GERANIUMS;...*four* in CURCUMA;...and *five* in PENTAPETES and some GERANIUMS.

The number of cells that contain the pollen, is either *one*, as in MERCURIALIS;...*two*, as in HELLEBORUS;...*three*, as in ORCHIS;...or *four*, as in FRITILLARIA.

The FIGURE of the antheræ is either *oblong*, as in LILIUM;...*globose*, as in MERCURIALIS;...*sagittate*, arrow-shaped, as in CROCUS;...*angulate*, cornered, as in TULIPA;...or *cornute*, horned, as in HAMAMELIS, ERICA, VACCINIUM, and PYROLA.

They BURST either on the *side*, as in LEUCOIMUM, and most flowers;...on the *apex*, as in GALANTHUS and KIGGELLARIA;...or

* See Part II. Chap. XV.

† Without petals.

‡ See Part II. Chap. XXII.

from the *apex*, to the *base* through the whole length, as in *EPIMEDIUM* and *LEONTICE*.

They are *FASTENED* either by their *base*, as in most plants;... their *tops*, as in *COLCHICUM*;...their *sides*, as in *CANNA*;...or grow to the *nectarium*, as in *COSTUS*.

Their *SITUATION* is either on the *tops* of the filaments, as in most plants;...on the *sides* of the filaments, as in *PARIS* and *ASARUM*;...on the *pistillum*, as in *ARISTOLOCHIA*;...or on the *receptacle*, as in *ARUM*.

The *FIGURE* of the particles of the pollen appears, by glasses, to be either *globus echinatus*, a *prickly ball*, as in *HELIANTHUS*;... *perforate*, as in *GERANIUM*;...*double*, as in *SYMPHYTUM*;...*rotatodentate*, *wheel-shaped*, as in *MALVA*;...*angulate*, *cornered*, as in *VIOLA*;...*reniform*, *kidney-shaped*, as in *NARCISSUS*;... or *folia sonvoluta*, a *leaf rolled up*, as in *BORAGO*.

CHAP. XIV.

OF THE DIFFERENT STRUCTURES OF THE PISTILLUM.

THE Pistillum consists* of three parts, *Germen*, *Stylus*, and *Stigma*. Of these the germen being no other than the rudiment of the pericarpium, its variations will be considered under that head in the next chapter; nor need we speak here of the number of the styles, as that will be treated of in the explanation of the *Sexual System*†; but as the style is often divided, we must consider its laciniae.

* The author should have said *usually consists of*. EDITOR.

† See Part II. Chap. III. in which the titles of the orders, which are governed chiefly by the number of styles, are explained.

STYLE—The style, in respect to its LACINIÆ, is either *bifid*, as in PERSICARIA and CORNUTIA;...*trifid*, as in CLETHRA and FRANKENIA;...*quadrifid*, as in RHAMNUS;...*quinquefid*, as in GERANIUM;...or *dichotomous, halved, and each lacinia halved again*, as in CORDIA.

The FIGURE of the style is either *cylindric, like a rolling stone*, as in MONOTROPA;...*angulate, cornered*, as in CANNA;...*subulate, awl-shaped*, as in GERANIUM;...*capillary, like hairs*, as in CERATOCARPUS;...or *thicker towards the top*, as in LEUCOIMUM.

In respect to LENGTH, it is either *very long*, as in TAMARINDUS, CASSIA, CAMPANULA, SCORZONERA and ZEA;...*very short*, as in PAPAVER;...or of the length of the *stamina*, as in NICOTIANA, and most flowers.

In respect to THICKNESS, it is either *thicker* than the *stamina*, as in LEUCOIMUM;...*thinner*, as in CERATOCARPUS;...or of *equal* thickness with them, as in LAMIUM.

Its SITUATION is either on the *apex* of the germen, as is too common to need example;...both *above* and *below* the germen, as in CAPPARIS and EUPHORBIA (unless the lower part in these be considered as the extension of the receptacle);...or on the *side* of the germen, as in ROSA, RUBUS, and the rest of the plants of the order *Polygynia*, in the class of *Icosandria**, and also in HIRTELLA and SURIANA.

As to its DURATION, it is sometimes *persisting*, as in the class *Tetradynamia*†.

STIGMA—The NUMBER of the stigmata is either a *single* one, as in most flowers;...*two*, as in SYRINGA;...*three*, as in CAMPANULA;...*four*, as in EPILOBIUM and PARNASSIA;...or *five*, as in PYROLA.

The LACINIÆ of the stigma are either *convolute, rolled together*, as in CROCUS;...*capillary*, as in RUMEX;...*revolute, rolled back*, as in DIANTHUS, CAMPANULA, and in the class *Syngenesia*‡;...or *bent to the left*, as in SILENE;...and in respect to their *number*, the stigma may be *sexpartite, divided into six parts*, as in ASARUM;...or *multifid, with many divisions*, as in TURNERA.

* See Part II. Chap. XV.

† See Part II. Chap. XVIII.

‡ See Part II. Chap. XXII.

The FIGURE of the stigma is either *capitate, headed*, as in TRIBELLUS, HUGONIA, VINCA, IPOMŒA, and CLUSIA;...*globose, globe-shaped*, as in PRIMULA, HOTTONIA, LINNŒA, and LIMOSELLA;...*ovate, egg-shaped*, as in GENIPA;...*obtuse, blunt*, as in ANDROMEDA;...*truncate, lopped*, as in MARANTA;...*pressed down obliquely*, as in ACTŒA and DAPHNE;...*emarginate, notched*, as in MELICA;...*orbiculate, rounded*, as in LYTHRUM;...*peltate, like a pelta, or little shield*, as in SARRACENA, NYMPHŒA, CLUSIA, and PAPAVER;...*coroniform, crown-shaped*, as in PYROLA;...*cruciform, cross-shaped*, as in PENŒA;...*uncinate, hooked*, as in VIOLA and LANTANA;...*canaliculate, grooved, or channelled*, as in COLCHICUM;...*concave, hollow*, as in VIOLA;...*angulate, cornered*, as in MUNTINGIA;...*striate, streaked*, as in PAPAVER;...*plumose, feathery*, as in RHEUM, TRIGLOCHIN, TAMARIX, and in grasses;...or *pubescent, downy*, as in CUCUBALUS and LATHYRUS.

In respect to LENGTH, it may be *filiiform, thread-like*, as in ZEA;...or *as long as the style*, as in GENIPA.

In respect to THICKNESS, it may be *foliaceous, resembling a thin leaf*, as in IRIS.

In respect to DURATION, it is either *marcescent, withering*, as in most plants;...or *persisting*, as in SARRACENA, HYDRANGŒA, NYMPHŒA, and PAPAVER.

CHAP. XV.

OF THE DIFFERENT STRUCTURES OF THE PERI-
CARPIUM.

THE variations of the pericarpium itself, in respect to NUMBER, arise properly from the number of its capsules; that is, the number of parts into which the fruit is *externally* divided, the internal divisions respecting the loculaments.

In respect to external division, the pericarpium is either *absent*, as in the order *Gymnospermia* of the class *Didynamia**;...*Unicapsular*, consisting of one capsule, as in LYCHNIS;...*bicapsular*, of two, as in PÆONIA and ASCLEPIAS;...*tricapsular*, of three, as in VERATRUM and DELPHINIUM;...*quadricepsular*, of four, as in RHODIOLA;...*quinquecapsular*, of five, as in AQUILEGIA;...or *multicapsular*, of many, as in CALTHA, TROLLIUS, and HELLEBORUS.

The fruit in respect to the *loculaments*, or internal divisions of the pericarpium, is either *unilocular*, of one cell, as in TRIENTALIS and PRIMULA;...*bilocular*, of two, as in HYOSCYAMUS, SINAPIS, and NICOTIANA;...*trilocular*, of three, as in LILIUM;...*quadrilocular*, of four, as in EUONYMUS;...*quinquelocular*, of five, as in PYROLA;...*sexlocular*, of six, as in ASARUM and ARISTOLOCHIA;...*octolocular*, of eight, as in the species of LINUM, called RADIOLA;...*decemlocular*, of ten, as in LINUM;...or *multilocular*, of many, as in NYMPHÆA.

The pericarpium, in respect to the number of its *valvules*, or outer inclosures, is either *bivalve*, of two valves, as in CHELIDONIUM and BRASSICA;...*trivalve*, of three, as in VIOLA, POLEMONIUM, and HELIANTHEMUM;...*quadrivalve*, of four, as in LUDWIGIA and OENOTHERA;...or *quinquevalve*, of five, as in HOTTONIA.

* See Part II. Chap. XVIII.

30 DIFFERENT STRUCTURES OF THE PERICARP.

The *dissepiments* are either *parallel* to the valvules, as in LUNARIA and DRABA;...or placed the *contrary way*, as in BISCUTELLA and THLASPI.

The most considerable differences in the *FIGURE* of the pericarpium, with the names assigned for each, have been explained in Chap. VI. It varies farther in being *turbinate*, *narrowing like a child's top*, as in PYRUS;...*inflated, puffed*, as in CARDIOSPERMUM and STAPHYLÆA;...*membranaceous, composed of thin membranes*, as in ULMUS;...*triquetrous, tetragonous, pentagonous, of three, four, or five sides*, as in AVERRHOA, ZYGOPHYLLUM, &c....or *articulate, jointed*, as in ORNITHOPYS, HEDYSARUM, and RAPHINUS.

The *OPENING* of the pericarpium for discharging the seeds when the fruit is ripe, is either at the *apex*, which may be *quadridentate, split into four segments*, as in DIANTHUS;...*quinquedentate, into five*, as in ALSINE;...or *decudentate, into ten*, as in CERASTIUM;...opening at the *base into three parts*, as in TRIGLOCHIN and CAMPANULA;...or *into five parts*, as in LEDUM;...at the *angles, corners, longitudinally, lengthways*, as in OXALIS and ORCHIS;...through a *pore, hole*, as in CAMPANULA;...or *horizontally across the middle*, as in ANAGALLIS, PLANTAGO, AMARANTHUS, PORTULACA, and HYOSCYAMUS.

All fruit that is *articulate, jointed*, opens at every one of the joints, each of which is *monospermous, single seeded*.

The *CONFINEMENT* of the seeds is sometimes *elastic, bursting like a spring*, as in OXALIS, ELATERIUM, MOMORDICA, IMPATIENS, CARDAMIME, PHYLLANTHUS, EUPHORBIA, JUSTICIA, RUELLIA, DICTAMNUS, HURA, RICINUS, TRAGIA, JATROPHA, CROTON, CLUSIA, ACALYPHA.

The *SITUATION* of the pericarpium is at the receptacle of the flower, either placed *under* it, as in VACCINIUM and EPILOBIUM;...*over* it, as in AREUTUS and TULIPA;...or both *above and below* it, as in SAXIFRAGA and LOBELIA.

CHAP. XVI.

OF THE DIFFERENT STRUCTURES OF THE SEEDS.

IN respect to the NUMBER of seeds contained within the fruit, plants are either *monospermous*, having one seed, as in POLYGONUM and COLLINSONIA;...*dispermous*, two, as in DAUCUS;...*trispermous*, three, as in EUPHORBIA;...or *tetraspermous*, four, as in TOURNEFORTIA.

In respect to the number of loculements of the seed itself, it has but one in most plants;...but is *bilocular*, with two cells, in CORNUS, XANTHIUM, LOCUSTA, VALERIANA, and CORDIA.

In respect to its FIGURE, it is either *cinct*, girt, as in ARENARIA and BRYONIA;...*cordiform*, heart-shaped, as in MEDEOLA;...*reniform*, kidney-shaped, as in ANACARDIUM and PHASEOLUS;...*ovate**, egg-shaped, as in POLYGALA and ISATIS;...or *echinate*, prickly, like an echinus, or hedge-hog, as in LAPPULA, a species of MYOSOTIS.

In respect to their SUBSTANCE, they are *osseous*, bony, as in CORYLUS, LITHOSPERMUM, and *nuts* of all kinds;...or *callous*, tough, as in CITRUS.

The CORONULA, *little crown*, that attends many seeds, is either *calyculus*, a *small calyx* formed of the perianthium of the flower, as in SCABIOSA, KNAUTIA, AGERATUM, and ARCTOTIS;...or *pappus*, a *down*; and this *pappus* is either *capillary*, like a hair, that is *simple* and *filiform*;...*thread-shaped*, as in HIERACIUM and SONCHUS;...*plumose*, *feathery*, that is, *shaggy* and *compound*, as in CREPIS, SCORZONERA, and TRAGOPOGON;...*paleaceous*, *chaffy*, as in BIDENS, SILPHIUM, TAGETES, and COREOPSIS;...or *wanting*, as in TANACETUM.

* The term *ovate* is used to express an elliptical figure when it is broader at one end than the other; and the term *oval* for the same figure, when the ends are alike.

32 DIFFERENT STRUCTURES OF THE RECEPTACLE.

The seed has an ARILLUS*, in COFFEA, JASMINUM, CYNOGLOSSUM, CUCUMIS, DICTAMNUS, DIOSMA, CELASTRUS, and EUONYMUS.

The seeds in respect to SIZE may be *very small*, as in CAMPANULA, LOBELIA, TRACHELIUM, and AMMANIA;...or *very large*, as in COCCUS.

In respect to SITUATION, they are either *nidulantia, nesting*, that is, dispersed about the pulp, as in NYMPHÆA;...fastened to the *suture*, as in plants that are *siliquose, podded*;...fastened to the *columella*, as in MALVA;...or placed on *receptacles*, as in NICOTIANA and DATURA.

The HILUM of the seed is evident in CARDIOSPERMUM and STAPHYLÆA.

The CORCULUM is close to the hilum.

CHAP. XVII.

OF THE DIFFERENT STRUCTURES OF THE RECEPTACLE.

IT is in the class SYNGENESIA†, which contains the compound flowers, that the varieties of the receptacle are principally to be considered.

In respect to its FIGURE, it is either *plane, flat*, as in ACHILLEA;...*convex, rounding*, as in MATRICARIA;...or *conic, shaped like a cone*, as in ANTHEMIS and MELAMPODIUM.

In respect to its SURFACE, it is either *naked*, as in MATRICARIA;...*punctate, dotted*, as in TRAGOPOGON;...*villose, shaggy*, as in

* See Chap. VII.

† See Part II. Chap. XXII.

ANDRYALA;...*setose, bristly*, as in CENTAUREA;...of *paleaceous, chaffy*, as in HYPOCHERIS and ANTHEMIS.

In some simple flowers the fruit has *separate* receptacles, as in MAGNOLIA, UVARIA, and MICHELIA.

CHAP. XVIII.

OF THE SINGULARITIES IN THE STRUCTURE OF THE PARTS OF FRUCTIFICATION.

BY a singular structure of the parts of fructification is to be understood such a one as is observed but in very few genera; it is directly opposed to the natural structure explained in Chap. X. For instances of this we may mention the ARUM, whose stamina are within the pistilla;...the ADOXA, whose germen separates the corolla from the calyx;...the SALVIA, whose filaments are *articulate, jointed*;...the ERIOCAULON, whose stamina are placed on the germen, and whose corolla and calyx are below the germen;...and the MAGNOLIA, the receptacle of whose fruit is *capitate, headed*, the seeds, which are like berries, hanging by a thread out of the capsule; but to take the parts in their order.

The CALYX is usually less coloured than the COROLLA; but in the AMERICAN BARTSIA the *perianthium* is red;...in the HERBACEOUS CORNUS the *petals* are black, but the *involucrum* white;...and in the AMERICAN CORNUS the *involucrum* is red, and *cordate*, that is, *heart-shaped*. In ASTRANTIA the *involucrum* is coloured; and in PALMS the *spathæ* are red; where the corolla is wanting, the *perianthium* is apt to be more coloured, especially when the flowers are blowing, as in ORNITHOGALUM, PERSICARIA, and POLYGONUM:...where either the calyx or the corolla is found to be less

coloured, the leaves often take a colour, as in *AMARANTHUS TRICOLOR*.

In most plants the *STAMINA* and *PETALS* are inserted into the *receptacle*, in the bottom of the flower; but the plants of the class *Icosandria** have a monophyllous calyx, the inner side of which is girt with a line, into which the stamina and petals are fastened; and the calyx is also observed to support the flowers in some other plants, as in *LYTHRUM*, *EPILOBIUM*, *CENOTHERA*, *AMMANIA*; *ISNARDA*, *PEPLIS*, and *ELÆAGNUS*. In some plants the receptacle is lined on all sides with the perianthium, and the corolla adheres to the perianthium as though it were glued to it; this is found in the *cucurbitaceous*† plants, such as *CUCURBITA*, *PASSIFLORA*, *FEVILLEA*, *MOMORDICA*, *TRICHOSANTHES*, *CUCUMIS*, *BRYONIA*, *SICYOS*, *MELOTHRIA*, and *GRONOVIA*; the same is also observed in *CACTUS*. In some others there is a receptacle that elevates the pericarpium, as in *PASSIFLORA*, *CAPPARIS*, *BREYNIA*, *ARUM*, *CALLA*, *DRACONTIUM*, *POTHOS*, *ZOSTERA*, *NEPENTHES*, *CLUTIA*, *HELICTERES*, and *SISYRINCHIUM*.

In monopetalous flowers the stamina are usually inserted into the petal, but they are separate from it in the *planta bicornes*‡, viz. in *LEDUM*, *AZALEA*, *ANDROMEDA*, *CLETHRA*, *ERICA*, *MYRSINE*, *MEMECYLUM*, *SANTALUM*, *VACCINIUM*, *ARBUTUS*, *ROYENA*, *DIOSPYROS*, *MELASTOMA*, and *PYROLA*; they are separate also in *CISSUS* and *ALOE*. In polypetalous flowers the stamina are usually separate from the petals. But this also has a few exceptions; for in the *STATICE*, which is pentapetalous, the filaments are inserted in the claws of the petals; in *MELANTHIUM*, which is hexapetalous, they are inserted in the petals; and in the *LYCHNIS*, which is pentapetalous, as also in *SAPONARIA*, *CUCUBALUS*, *SILENE*, and *AGROSTEMMA*, which were formerly ranged with the *LYCHNIS*, every other stamen is fastened to the claws of the petals.

The *ANTHERS* are commonly placed on the tops of the filaments: but they stick close to the sides of the filaments in *PA*.

* See Part II. Chap. XV.

† So called from their affinity to the cucurbita.

‡ Having two horns; these plants have been so called from their bifid *Antheræ*.

RIS and ASARUM, and adhere to the stigma without filaments in ARISTOLÓCHIA.

The singularities of the NECTARIUM have been already mentioned in Chap. XII.

The PISTIL is commonly placed within the *anthers*: but in the ARUM there is this singularity, that the receptacle runs out into a club, the base of which is occupied by the pistilla, and the upper part by the stanina; so that here the pistilla stand on the outside of and surround the stamina; and in the ETHIOPIAN CALLA these parts are disposed in the same manner. The RUMEX is singular in the insertion of its stamina.

The STYLE is commonly placed on the top of the germen. Some exceptions to this have been given in Chap. XIV. to these may be added PASSERINA, GNIDIA, STRUTHIA, and STELLARIA.

The PERICARP is generally shut; but in RESEDA and DATISCA it is always open: in PARNASSIA it gapes at the time of flowering, and closes afterwards.

That the pericarpia are ever found one within another, the greater containing the smaller ones, *Linnaeus* refuses to admit; for although there is the appearance of such a singularity in MAGNOLIA, UVARIA, and MICHELIA, he thinks the outer pericarp is in such cases to be looked upon only as a common receptacle.

Where the pericarp is a berry, it is distinguishable into *proper* berries, those which are formed of the pericarp;...and *improper* or *singular*, such as are formed of any of the other parts.

The berry is improper or singular in the following instances, viz. When it is a *calyx*, as in BLITUM, MORUS, BASELLA, EPHEDRA, COIX, ROSA and CORIARIA;...a *receptacle*, as in TAXUS, RHIZOPHORA, ANACARDIUM, OCHNA, LAURUS, FICUS, DORSTENIA, and FRAGARIA;...a *seed*, as in RUBUS, MAGNOLIA, UVARIA, MICHELIA, PRASIMUM, UVULARIA, PANAX, ADONIS, CRAMBE, and OSTEOSPERMUM;...an *Arillus*, as in EUONYMUS and CELASTRUS;...a *nectarium*, as in MIRABILIS;...a *corolla*, as in ADOXA, POTERIUM, and CORIARIA;...a *capsule*, as in EUONYMUS, ANDROSEMUM, CUCUBALUS, and EPIDENDRUM;...a *dry berry*, as in LINNEA, GALIUM, &c. TETRAGONIA, MYRICA, TRIENTALIS, TROPEOLUM, XAN-

THIUM, JUGLANS, PTELEA, ULMUS, COMARUM, AMYGDALUS, and MIRABILIS; ... a *capsule* externally, as in DILLENIA, CLUSIA, NYMPHLEA, CAPPARIS, BREYNIA, MORISONIA, STRATIOTES, CYCLAMEN, and STRYCHINUS; ... a *hollow berry*, as in STAPHYLEA, CARDIOSPERMUM, and CAPSICUM; ... a *conceptacle*, as in ACTÆA; ... a *legumen*, as in HYMENÆA, CASSIA, INGA, and CERATONIA; ... of a *strobilus*, as in ANNONA and JUNIPERUS.

The berry does not naturally burst, being soft, and the dispersion of the seeds being designed to be by means of animals.

The berries in the CAPE ADONIS are evidently *aggregate*, many united in one.

CHAP. XIX.

OF AGGREGATE FLOWERS.

COMPLETE flowers are either simple or aggregate. Simple flowers differ from aggregate in this, that they have not any part of fructification common to many flowers, as is the case with aggregate. Flowers are called aggregate when many *flosculi*, *florets*, are by the mediation of some part of the fructification common to them all, so united, that no one of them could be taken out without destroying the form of the whole, of which it was a part. The common part in aggregate flowers is either the receptacle or the calyx. A partial flower of the aggregate one is called *flosculus*, a *floret*. Aggregate flowers are primarily divisible into seven kinds, viz. 1. The *aggregate*, properly so called. 2. The *compound*. 3. The *umbellate*. 4. The *cymose*. 5. The *mentaceous*. 6. The *glumose*. 7. The *spadiceous*: all which we shall explain in their turn.

1. An AGGREGATE flower, properly so called, has a receptacle that is *dilate*, *extended in breadth*, the florets standing on

*peduncles, foot-stalks**, as in SCABIOSA, KNAUTIA, DIPSACUS, CEPHALANTHUS, GLOBULARIA, LEUCADENDRON, PROTEA, BRUNIA, BARRERIA, and STATICE.

2. A COMPOUND flower† is an aggregate one, comprehending many florets that are *sessile*, without *peduncles*, on a common receptacle that is entire, and having also a common perianthium, but furnished with antheræ that grow together in the form of a cylinder.

The properties of a compound flower are, 1. A common receptacle enlarged and undivided. 2. A common perianthium, surrounding all the florets. 3. The florets monopetalous and sessile. 4. The antheræ of each floret five in number, and growing together in a cylinder. 5. A monospermous germen under each of the florets. Of these properties, the two last are essential to a compound flower; but observe, that there are some whose calyx contains only a single floret, as ECHINOPS, STÆBE, CORYMBIUM, and ARTEMISIA.

Compound flowers are of three kinds: 1. *Ligulate*, when all the *corollulæ, little corollæ* of the florets, are *plane, flat*, shaped like *ligula, a narrow tongue, or fillet*, and expanded towards the outer side. 2. *Tubulose*, when all the *corollulæ* of the florets are tubulose, and nearly equal. 3. *Radiate, having rays*, when the *corollulæ* of the *disk, middle parts*, are tubulose, and those of the *circumference, margin*, of another form: which variation affords three cases, viz. when the *corollulæ* of the circumference are either *ligulate*, as in ACHILLEA;...*tubulose*, but unlike the tubulous florets of the disk, as in CENTAUREA;...or *naked*, as in ARTEMISIA and GNAPHALIUM. A compound flower usually consists of many florets, but rarely of a determinate number of them.

3. An UMBELLATE flower is an aggregate one, consisting of

* *Peduncle* is the foot-stalk of a flower only; the foot-stalk of a leaf is called a *petiole*.

† These are the flowers of the class *Syngenesia*, see Part II. Chap. XXII.

many florets placed on a receptacle, on fastigate peduncles* that are all produced from the same point. A *simple* umbel is when the receptacle is but once divided into peduncles;...a *compound* umbel is when all the common peduncles are subdivided into *umbellulæ*, *little umbels*;...an *umbellula* therefore is a *partial* umbel.

Umbellate flowers, properly so called†, have the following properties: 1. A common receptacle divided into peduncles in the manner above-mentioned, whether the umbel produced be *plane*, *flat*; *convex*, *rounding*; or *concave*, *hollow*. 2. A germen under the corollula. 3. Five distinct stamina that are deciduous. 4. A bifid pistillum. 5. Two seeds joined at their summits.

A *radiate* umbel is when the marginal petals are larger than those of the disk, as in TORDYLIUM, CAUCALIS, CORIANDRUM, AMMI, and some species of HERACLEUM; an umbel may vary also in having the flowers of the margin differing in sex from those of the disk, as in ASTRANTIA, CAUCALIS, ARTEDIA, (ENANTHE, and SCANDIX. The *involucrum* varies, in being either *tetraphyllous*, of four leaves, as in HYDROCOTYLE, SISON, and CUMINUM;...*pentaphyllous*, of five, as in BUPLEURUM, SCANDIX, and BUBON;...*heptaphyllous*, of seven, as in LIGUSTICUM;...*decaphyllous*, of ten, as in ARTEDIA:...with the *partial involucrum dimidiate*, *halved*, going but half round, as in ÆTHUSA, CORIANDRUM, and SANICULA;... or *caducous*, *falling off*, as in FERULA and HERACLEUM.

4. A CYMOSE flower is an aggregate one, of many florets, placed on a receptacle upon fastigate‡ peduncles, the primary ones of which issue from the same centre, as in an umbel; but the secondary, or partial ones, lie dispersed without order; which circumstance distinguishes the cyma from the umbel, as in OPULUS, OPHIORRHIZA, and the species of cornus, called VIRGA-SANGUINEA, or *bloody-rod*.

* See the first note in Chap. VIII.

† The umbellate flowers, properly so called, belong to the order *Digynia*, of the class *Pentandria*. See Part II. Chap. VIII.

‡ See the first note on Chap. VIII.

5. An AMENTACEOUS aggregate flower has a *filiform, thread-shaped* receptacle, along which are disposed *amentaceous squamæ, scales* that form an *amentum, or catkin*, as in XANTHIUM, AMBROSIA, PARTHENIUM, IVA, ALNUS, BETULA, SALIX, POPULUS, CORYLUS, CARPINUS, JUGLANS, FAGUS, QUERCUS, LIQUIDAMBAR, CYNOMORION, FICUS, DORSTENIA, PARIETARIA, URTICA, PINUS, ABIES, CUPRESSUS, THUYA, JUNIPERUS, TAXUS, and ÉPHEDRA.

6. A GLUMOSE aggregate flower has a filiform receptacle, the base of which is furnished with a common *glume, husk*, as in BROMUS, FESTUCA, AVENA, ARUNDO, BRIZA, POA, AIRA, UNIOLA, CYNOSURUS, MELICA, ELYMUS, LOLIUM, TRITICUM, SECALE, HORDEUM, SCIRPUS, CYPERUS, and CAREX.

7. A SPADICEOUS aggregate flower is, when there is a receptacle common to many florets, placed within a *spatha* or sheath; such a receptacle is called a *spadix*, and is either *branched*, as in palms, or *simple*. In this last case the florets may be disposed either all *round* it, as in CALLA, DRACONTIUM, and POTHOS;... on the *lower* part of it, as in ARUM;...or on *one side* of it, as in ZOSTERA.



CHAP. XX.

OF LUXURIANT FLOWERS, COMMONLY CALLED DOUBLE.

A FLOWER is said to be luxuriant, when some of the parts of fructification are augmented in number, and others thereby excluded. The luxuriance is commonly owing to the luxuriance of its nourishment; the part multiplied is usually the corolla, but sometimes the calyx also; and by this increase of the covers,

the essential parts of fructification are destroyed. Luxuriant flowers are divisible into, 1. *Multiplicate, multiplied*. 2. *Pleni, full*. And 3. *Proliferous, producing young*; to which may be added, 4. *Mutilate, maimed*; such as are deficient in some part, which stand opposed to the luxuriant ones: all these shall be explained in their order.

1. Flowers are said to be MULTIPLICATE, when by the increase of the corolla only a part of the stamina are excluded; and this distinguishes them from the *flores pleni, full flowers*, in which the multiplication of the corolla is so great as to exclude them all. Multiply flowers are distinguished into *duplicate, triplicate, quadruplicate, &c.* that is, having a *double, treble, or quadruple* series, or row, according to the number of the repetitions of the corolla. The *polypetalous* flowers are the most subject to multiplication; the *monopetalous* are multiplied likewise, but it is very uncommon to meet with them full. A *coloured perianthium*, though it may have the appearance of a repetition of the corolla, ought not to be considered as such; for though this appearance is in some degree *monstrous, unnatural*, it is no multiplication.

2. A flower is said to be PLENUS, *full*, when the corolla is so far multiplied as to exclude all the stamina, as was before observed. The *plenitude, fullness*, is occasioned by the stamina running into petals, with which the flower is so crowded as frequently to choak the pistillum also. The parts essential to generation being thus destroyed in full flower it is evident they must be barren; wherefore no good seed is to be expected from them*. And for the same reason of their imperfection, we should be cautious also of constituting a genus from them; for the characters of a genus should be drawn from the parts when in their natural state, and not when in a state of luxuriance.

Plenitude is chiefly incidental to polypetalous flowers, as in

* Some few, as the *Pionia, Papaver, and Nigella*, perfect their seed: but these are rather multiply flowers than full ones.

MALUS, PYRUS, PESICA, CERASUS, AMYGDALUS, MYRTUS, ROSA, FRAGARIA, RANUNCULUS, CALTHA, HEPATICA, ANEMONE, AQUILEGIA, NIGELLA, PAPAVER, PÆONIA, DIANTHUS, SILENE, LYCHNIS, CORONARIA, LILIUM, FRITILLARIA, TULIPA, NARCISSUS, COLCHICUM, CROCUS, CHEIRANTHUS, HESPERIS, MALVA, ALCEA, and HIBISCUS.

Plenitude of monopetalous flowers is by some authors held a contradiction ; but this cannot be granted ; for there are instances of it in COLCHICUM, CROCUS, HYACINTHUS, and POLIANTHES : however, it is rare that their luxuriancy passes duplicity. When they are filled, it is by the multiplication of the *laciniæ, segments* ; whereas the polypetalous are usually filled by the multiplication of the petals ; but the manner in which the *impletion, filling*, is brought about, must be more particularly considered.

The impletion is either in simple or compound flowers ; we shall begin with the simple.

The impletion of SIMPLE flowers is by the increase either of the petals, or of the nectarium. The impletion of the AQUILEGIA is observed to be after three different manners, viz. either, 1. By multiplying its petals, and excluding the nectaria ; 2. By multiplying its nectaria, and excluding its petals ; or, 3. By multiplying its nectaria, and retaining its petals ; in which last case the five petals remain, and the spaces between them are each of them filled up with a triple case of nectaria ; that is, three nectaria buried one within another.

The impletion of the NIGELLA is by multiplying the nectaria only ; that of the NARCISSUS two ways, by multiplying either the nectarium only, or both nectarium and petals ; that of DELPHINIUM, for the most part, by multiplying the petals, and excluding the nectarium. The change wrought in the SAPONARIA ANG-LICANA is remarkable, the flower from pentapetalous becoming truly monopetalous ; and the alteration in the PELORIA is also very singular* : but the most extraordinary instance of plenitude

* The *Peloria* is a plant which has been found in some parts of *Sweden*, growing amongst the species of *Antirrhinum* called *Linaria*. It resembles the *Linaria* so

is that of the *OPULUS FLORE GLOBOSO*, commonly called the *Gelder rose*. In the common simple *OPULUS*, the flowers are produced on a cyma, which consists of a great number of *campanulate, bell-shaped*, hermaphrodite flowers in the disk, and of others in the circumference, whose corollæ are larger, flat, and wheel-shaped; and that are *barren, wanting the pistillum*. But in the *OPULUS FLORE GLOBOSO*, all the flowers of the disk are barren also, and shaped like those of the circumference; so that the impletion here arises only from the additional number of barren flowers, the corollæ of which are of a larger size; and in this it resembles the impletion of the compound flowers, of which we shall presently speak.

Before we leave the simple flowers, it will be of use to remark, that a simple flower, in a state of luxuriancy, may in all cases be distinguished from a compound one in its natural state, by this rule; that in *simple* flowers, how much soever multiplied, there is but one pistillum in the centre of the flower, common to the whole multiplication; whereas in *compound* flowers, each of the florets is furnished with its own pistillum and stamina.

We come now to the impletion of *COMPOUND* flowers; that these are of three kinds, *ligulate, tubulose**, and *radiate*, has been shown and explained in Chap. XIX. where it has also been seen, that there is not either in the ligulate or tubulose any distinction of disk or radius, all the florets in these being alike; but that the contrary is the very characteristic of the radiate; now this being attended to, the manner of the impletion will be easily understood. Compound flowers gain their impletion two ways, either by the radius, or the disk. We shall begin with the first.

Impletion by the *radius* is when, by the multiplication of the

nearly, in every thing but the flower, that they are not to be known one from the other, till their flowers appear; and even in the flowers they agree in the calyx, pericarpium, and seeds, and also in colour; which has given rise to a supposition, that the *Peloria* is only a *Linaria* in a monstrous state; see the Dissertation of *Daniel Rüdberg* on the *Peloria*, in the *Amanitates Academicæ* vol. I. p. 280. This is now known to be the fact, as the *Peloria* has been known to return back to the *Linaria*; and flowers of both kinds have been found on some plants. EDITOR.

* Tubulose, *tubular*. EDITOR.

radius, the disk of the flower is filled up; as in *HELIANTHUS*, *CALENDULA*, *CHRYSANTHEMUM*, *ANTHEMIS*, *MATRICARIA*, *PTARMICA*, *TAGETES*, and the species of *CENTAUREA*, called *CYANUS*. In this sort of impletion, which belongs only to radiate flowers, it is observable, that all the florets which fill up the disk follow the conditions of those of the radius; so that if the florets of the radius in the natural flower have a pistillum, all those of the full flower will have one also, as in *MATRICARIA*, *BELLIS*, *CHRYSANTHEMUM*, and *TAGETES*; or if they have no pistillum, then it will also be wanting in the full one, as in *HELIANTHUS*, *CALENDULA*, and *CENTAUREA*; and the same holds true of the male part also; for as the florets of the radius in the natural flower are never furnished with antheræ, so these are wanting also in all those of the full ones. This last remark is of great use to distinguish a radiate full flower, from a ligulate natural one; which might be confounded in many cases, were we not apprised that there are antheræ in the latter, but none in the former; by this rule in *CHRYSANTHEMUM*, *HELIANTHUS*, *CALENDULA*, and *TAGETES*, when the disk is destroyed by the multiplication of the radius, we know by the defect of antheræ, that it is only the luxuriancy of a radiate flower, as in *HIERACIUM*, *LEONTODON*, and *SONCHUS*; by the presence of the antheræ we know the flowers to be ligulate and natural.

Impletion by the *disk* is, when there is no multiplication of the radius; but the corollulæ of the disk run out into length, and have their brims less divided: this manner of impletion seems to concern only the *radiate* and the *tubulose**. In the *radiate*, it will so far affect the radius, as to change its flowers from ligulate to tubulose: instances of this manner of impletion may be had in *BELLIS*, *MATRICARIA*, and *TAGETES*. In the *CARDUUS* of the oats, which is a species of *SERRATULA*, the corollulæ are both lengthened and enlarged. In respect to the *ligulate* flowers, if

* This is not expressly asserted, as the distinction is omitted in the *Philosophia Botanica* of *Linnaeus*; but it appears to be his meaning, by his speaking of the impletion of ligulate flowers separately afterwards.

we confine ourselves to the two-fold manner of impletion, after the author, whose divisions we have adopted, we shall be obliged to call their impletion also, an impletion by the disk; though the manner of it differs from that last explained, and the expression does not so well answer to flowers, that in the botanical sense of the term have properly no disk at all. But not to stop at too great niceties, their impletion is by the lengthening of their stigmata, and the enlarging and diverging of their germina; by which augmentations, the full flowers are to be distinguished from the natural ones, as in SCORZONERA and LAPSANA VULGARIS; which last, *Linnaeus* tells us, is frequently found with a full flower at *Upsal*.

3. Flowers are said to be PROLIFEROUS, when one flower grows out of another: this generally happens in full flowers, the fullness being the cause of their becoming prolific. Prolification is after two manners; 1. From the centre; 2. From the side.

Prolification from the *centre*, which happens in simple flowers, is when the pistillum shoots up into another flower, standing on a single peduncle; of which there are instances in DIANTHUS, RANUNCULUS, ANEMONE, GEUM, and ROSA.

Prolification from the *side*, which happens in aggregate flowers, properly so called (see Chap. XIX.), is when many pedunculate flowers are produced out of one common calyx; of which there are instances in BELLIS, CALENDULA, HIERACIUM, and SCABIOSA.

In *umbellate* flowers, the prolification is by the increase of the umbellulæ, one simple umbellula producing another, as in CORNUS and PERICLYMENUM; and in this manner compound umbels will become *supradecomposed*, more than compounded a second time, as in SELINUM and THYSSELINUM.

A prolific flower is called *frondose**, leafy, when it pro-

* *Frons*, with the ancients (though frequently used, in respect to trees, in the same sense with *folium*, a leaf) implied, in its proper signification, a part of the wood of the tree with the leaf; or as we should express it, a *twig with leaves*; and for this reason they never applied the term to the leaves of herbs (which were always

duces leaves ; this rarely happens ; but instances of it have been found in ROSA, ANEMONE, and others. The other kinds of proliferation are frequent enough.

4. MUTILATE flowers are the reverse of luxuriant. *Linnaeus* confines the term to those flowers only that want the corollæ, though they ought to be furnished with it ; which often happens in IPOMÆA, CAMPANULA, RUELLIA, VIOLA, TUSSILAGO, and CUCUBALUS. The cause of this defect he ascribes chiefly to the want of sufficient heat.

The luxuriancy of the *calyx*, mentioned in the beginning of this chapter, is very unfrequent, but not without instances ; in DIANTHUS CARYOPHYLLUS there is a variety, in which the *squamæ*, *scales*, of the calyx, are so multiplied as to constitute a perfect spike, in a manner most singular. The GRAMINA, *grasses*, of the Alps, become full by their *glumæ*, *husks*, shooting out into leaves, as in a species of the FESTUCA ; and in SALIX ROSEA, and PLANTAGE ROSEA, the *squamæ* of the amentum of the former, and the bractæ* of the spike in the latter, will shoot into leaves also.

Linnaeus has enumerated some tribes of plants, which are not found subject to luxuriancy ; but as the heads, under which he has ranged them, are taken from the systems of preceding writers, and not from the sexual, it would perplex the reader to explain them ; and we shall therefore omit them. The curious may have recourse to them in the *Philosophia Botanica*, p. 81.

called *folia*), but only to those of trees. *Linnaeus* has availed himself of this old distinction to make it a botanical term ; which he applies to express the circumstances of *palms* and *flices*, *ferns* ; in the former of which the branches, and in the latter even the stem itself is an actual leaf : and here again he applies it to the leafy proliferation in question, calling it *frondose*, rather than *foliaceus*, for the like reason. AUTHOR.

* Floral leaves.

CHAP. XXI.

OF THE SEXES IN PLANTS.

THE distinction of flowers into male, female, bisexual, and neuter, has been already explained in Chap. IV. To which we must add, that bisexual flowers are sometimes distinguishable into *male* and *female* bisexual: this is, when, although the flower contains the parts belonging to each sex, one of them proves abortive or ineffectual; if the defect be in the *stamina*, it is a *female* bisexual; if in the *pistillum*, a *male* one. The case wherein this distinction becomes necessary, happens very rarely. It will be shown in the course of this chapter.

Plants, in respect to sex, take their denominations from the sex of their flowers, in the manner following:

1. **BISEXUAL** plants are such as upon the same root bear flowers, that are all composed of the two sexes in the same corolla, as in most genera.

2. **ANDROGYNOUS**, *male* and *female*, such as upon the same root bear both male and female flowers, as in the class *Monœcia**.

3. **MALE**, such as upon the same root bear male flowers only, as in the class *Diœcia*†.

4. **FEMALE**, such as upon the same root bear female flowers only, as in the class *Diœcia*.

* See Part II. Chap. XXIV.

† See Part II. Chap. XXV.

5. POLYGAMOUS*, such as either on the *same*, or on *different* roots, bear bisexual flowers, and flowers of either or of both sexes, as in the class *Polygamia*†.

Of plants that are polygamous on the *same* root, there are three cases: 1st. *Male bisexual* and *Female bisexual* flowers; which is a very rare case; but is observed in MUSA. 2. *Bisexual* and *male* flowers, as in VERATRUM, CELTIS, ÆGILOPS, and VALANTIA. 3. *Bisexual* and *female* flowers, as in PARIETARIA and ATRIPLEX.

Of such as are polygamous on *two* distinct roots, the cases are four; 1. *Bisexual* flowers and *male*, as in PANAX, NYSSA, and DIOSPYROS. 2. *Bisexual* flowers and *female*, as in FRAXINUS. 3. *Bisexual* flowers and both *male* and *female*, as in GLEDITSIA‡. 4. *Androgynous*|| and *male*, as in ARCTOPUS. Of plants that are polygamous on *three* distinct roots, there is but one case, viz. *Androgynous*, *male*, and *female*, as in FICUS§.

* See the signification of this term explained in the account of the title of the class *Polygamia*, in Part II. Chap. XXVI.

† See Part II. Chap. XXVI.

‡ In the *Gleditsia*, which is the only known instance of this case, the male flowers and the bisexual are produced upon the same plant, and the females on a distinct one.

|| This case and the next, having no bisexual flowers, seem to be exceptions to the definition of polygamous plants.

§ The instance of this case given in the *Philosophia Botanica* is the *Empetrum*; but that *genus* is removed to the class *Diœcia*, in the last edition of the *Genera Plantarum*; where a note informs us, that the bisexual flowers, which the author had once seen on a plant of this *genus*, could not afterwards be ever found again. We have therefore changed this instance for the *Ficus*, the only other instance left of this singular case. Some have asserted, that the *Ficus* is only male and female; and this age hath refuted the opinion of Camerarius, who maintained, that the seeds of figs produce never any plants. Linnæus asserts that trees have been raised in Holland from the seed of fruit imported from Italy. But if the fruit be produced in France, England, Germany, or Sweden, where there are no wild figs, the seeds produce nothing; on the other hand, if those seeds are sown, which grew in Italy or the Greek islands, where the male fig abounds, the plants spring up with ease, putting forth leaves which at first are like those of the Mallow.

The best proof of the sexes in plants is drawn from the production of *hybrids*, or *bastards*, which is well observed in CABBAGES. One Richard Baal, a gardener at Brentford, sold a great quantity of *cauliflower* seeds, which he raised in his own garden, to several gardeners in the suburbs of London, who carefully sowed the seeds in good ground, but they produced mostly the *common long-leafed cabbage*, for which reason they complained they were imposed upon, and commenced a suit against Baal in Westminster Hall; the judge's opinion was, that Baal must return the gardeners the money he had received, and also make good their loss of time and crops, being wholly unacquainted with the sexes of plants. *Vide Ray's History, vol. 1. p. 42.* This apparent fraud we ought not to ascribe to the poor gardener, for it depended wholly on the impregnation by the common sorts; wherefore, if any one doth possess an excellent sort of cabbage, he ought not to let it flower in the same bed with any other of an inferior sort, lest the good sort should be impregnated with the dust of the other, and produce a degenerate race*.

* Those who wish for further information upon this curious point will do well to consult Doctor Thornton's superb new Illustration of the Sexual System of Carolus Von Linnæus. EDITOR.

PART SECOND.

CHAPTER I.

OF THE SEXUAL SYSTEM, AND ITS ORIGIN.

THE SEXUAL SYSTEM, as its title imports, is founded on a discovery, that there is in vegetables, as well as in animals, a *distinction of the sexes*. This was not wholly unknown to the ancients; but their knowledge of it was very imperfect.

It has been seen in the course of this work, that the flowers of the generality of vegetables are *bisexual*, containing within them the characters of both sexes; and we shall see in the classes *Monæcia* and *Diaecia*, the sexes are parted, and allotted to different flowers; and that in the class *Diaecia* in particular, the sexes are even on different plants, the male flowers growing all upon one plant, and the female upon another. Now this last circumstance the ancients had observed: indeed it could hardly escape their notice; for the Palm-tree, whose fruit was in esteem, being of the class *Diaecia*, a very little observation was requisite to teach them, that in these trees the flowers of the male were necessary to ripen the fruit of the female. Accordingly we find, in

the account given by *Herodotus** of the country about *Babylon*, where these trees are in plenty, that it was a custom with the natives, in their culture of this plant, to assist the operations of nature, by gathering the flowers of the male trees, and carrying them to the female. By this means they secured the ripening of the fruit, which might else, from unfavourable seasons, or the want of a proper intermixture of the trees of each sex, have been precarious, or at least not to have been expected in equal quantities.

It seems pretty extraordinary, that this discovery should not have led the ancients to detect the whole process of Nature in the propagation of the various species of vegetables; and yet it does not appear, by any of their writings that are come down to us, that they went farther than this obvious remark upon the palm-tree, and some similar notions concerning the fig. They had indeed, from what they saw in these plants, formed a notion, that all others were male and female likewise†; but this notion was false, the far greater part having bisexual flowers, and serves to convince us, that what they discovered of the palm and fig, was only a right guess, and not founded on any knowledge of the anatomy of flowers, either in those trees, or any others.

In this dark state the doctrine of the sexes of vegetables remained, not only through all the ages of antiquity, but almost to the end of the last century, the moderns seeing no more of this doctrine than the ancients had done before them; and hence we have to this very hour in use, the false distinctions of male and female species of *cornus*, *pæony*, *cistus*, and many others, which have all bisexual flowers, the distinction in these cases being

* Book the first.

† Thus *Theophrastus* says, in his *History of Plants* :

“ In trees, considered universally, and taking in each several kind, there are, as has been said, many differences. One of these is common to them all, namely, that by which they are distinguished into female and male, of which the one bears fruit, the other not, in some kinds; in those in which both bear fruit, that of the female is the best, unless these are to be called males, for so they are called by some.

Hist. Pl. Book iii. Chap. IX.

grounded on nothing more than some difference in the habit of the two species with which the sexes are no ways concerned.

The honour of having first suggested the true sexual distinctions in plants appears to be due to our countryman, Sir *Thomas Millington*, from whose hints Dr. *Grew*, as the doctor himself acknowledges, was led to the observations he has given on this subject, in his *Anatomy of Plants**. After this, *Camerarius*, *Moreland*, *Geoffroy*, *Vaillant*, *Blair*, *Jussieu*, and *Bradley*, pursued their enquiries and experiments so far as to remove all doubt concerning these discoveries; and lastly, *Linnaeus* added his observations, and founded thereon the system of botany, which we are going to explain in this work.

The sexual hypothesis, on its first appearance, was received with all that caution that becomes an enlightened age; and nature was traced experimentally through all her variations, before it was universally assented to. *Tournefort* refused to give it any place in his system; and *Pontedera*, though he had examined it, treated it as chimerical; but the proofs which *Linnaeus* has stated amongst the aphorisms of his *Fundamenta Botanica*†, and farther explained and illustrated in his *Philosophia Botanica*‡, are so clear, that the birth of animals is not more evidently the consequence of an intercourse between the sexes, than that of vegetables; and it would be now as ridiculous for any one, who has looked at the arguments, to doubt of the one as of the other.

We shall not attempt to lay all these proofs before the reader; our business is to explain, not demonstrate; but as it may be satisfactory to see some one fact established, that carries conviction with it, we shall here give an extract of a letter from *Berlin*,

* Published in the year 1682. The doctor expresses himself thus: "In discourse hereof with our learned *Savilian* professor, Sir *Thomas Millington*, he told me, he conceived that the attire doth serve as the male, for the generation of the seed. I immediately replied, that I was of the same opinion, and gave him some reasons for it, and answered some objections which might oppose them, &c." *Anat. of Plants*, p. 171.

† Aphorism 132 to 150.

‡ Page 86 to 96.

inserted in the *Philosophical Transactions**, concerning a remarkable experiment made on the palm-tree.

*Professor Mylius's Letter to Doctor Watson, dated at Berlin,
February 20, 1750—51.*

“The sex of plants is very well confirmed, by an experiment that has been made here on the *palma major foliis flabelliformibus*. There is a great tree of this kind in the garden of the Royal Academy. It has flowered and bore fruit these thirty years, but the fruit never ripened, and when planted, it did not vegetate. The palm-tree, as you know, is a *planta diœcia*; that is, one of those in which the male and female parts of generation are upon different plants. We having therefore no male plants, the flowers of our female were never impregnated with the farina of the male. There is a male plant of this kind in a garden at *Leipsic*, twenty *German* miles from *Berlin*. We procured from thence, in *April*, 1749, a branch of male flowers, and suspended it over our female ones; and our experiment succeeded so well, that our palm-tree produced more than an hundred perfectly ripe fruit; from which we have already eleven young palm-trees. This experiment was repeated last year, and our palm-tree bore above two thousand ripe fruit. As I do not remember a like experiment, I thought it convenient to mention it to you; and, if you think proper, be pleased to communicate it to the Royal Society.”

This letter, which was read to the Society the 2d of *May*, 1751, with some ingenious observations on the same subject, by *Dr. Watson*, F. R. S. to whom it was addressed†, has established the fact, attested by the ancients, concerning the palm-tree, which some may, perhaps, have looked upon as fabulous; and,

* Vol. XLVII. Page 169.

† Printed also in the *Philosophical Transactions* with the letter.

as the fructification in other vegetables, though it may differ in particuar circumstances, has yet, in general, a manifest conformity with that of the palm-tree, in respect to the parts supposed to be the organs of generation, which are discoverable either on the same, or on a separate flower, in all but the class *Cryptogamia*, where they are too minute for observation; so from this single experiment we may fairly draw an argument, by analogy, for the confirmation of the whole sexual hypothesis: but there are, as has been said, other, and stronger proofs. We have already directed the reader to those stated by *Linnaeus*; whoever desires farther satisfaction concerning this point, may see the several demonstrations collected, and methodically connected in the *Sponsalia Plantarum* of *J. Gustavus Walkbloom*, published in the *Amœnitates Academicae* at *Leyden*, in 1749.

The SEXUAL SYSTEM was invented by *Linnaeus*, professor of physie and botany, at *Upsal*. It is founded on the parts of fructification described in the former part of this work: these having been observed with more accuracy, since the discovery of the uses for which Nature has assigned them, a new set of principles have been derived from them; by means of which, the distribution of plants has been brought to a greater precision, and rendered more conformable to true philosophy in this system, than in any one of those which preceded it. The author of it does not pretend to call it a *natural* one; he gives it as *artificial* only, and modestly owns his inability to detect the order pursued by Nature in her vegetable productions: but of this he seems confident, that no natural system can ever be framed, without taking in the materials, out of which he has raised his own; and urges the necessity of admitting *artificial systems* for convenience, till one *truly natural* shall appear*.

* *Linnaeus* has given *Fragmenta Methodi naturalis*, *Fragments of the natural Method*, in which he has made a distribution of plants under various *orders*, putting together in each, such as appear to have a *natural affinity* to each other. This appear, after a long and fruitless search after the *natural method*, he gives as the result of his

By the *Sexual System* plants are disposed according to the *number*, *proportion*, and *situation* of the *stamina* and *pistilla*, &c. The manner of their distribution will appear in the following chapters. We shall here only speak in general of the divisions of the system.

The first general division of the whole body of vegetables is into twenty-four *classes*; these are again subdivided into *orders*, the orders into *genera*, the genera into *species*, and the species into *varieties*, where there are any worthy of note. Of these divisions, we shall treat of the three first only in this second part. These more immediately respect the theory of the science than the other two, which, though systematic divisions likewise, have, as our author observes, a nearer relation to the practice; and it is in these also that the principal improvements in the management of the science are more particularly included.

As the classes and orders of the system will be separately treated of in the following chapters, we shall conclude this introductory one with a table, exhibiting their titles at one view, in the order in which they stand in the system; that the reader may have recourse thereto, as he finds occasion*.

own speculation, for the assistance of such as may engage in the same pursuit. See his *Classes Plantarum*, p. 485, and *Phil. Bot.* p. 27.

* Should any difficulty occur to the young student, in comprehending the SEXUAL SYSTEM of *Carolus Von Linnæus*, the reader is referred to Doctor Thornton's very easy explanation of that admirable system, in his "GRAMMAR OF BOTANY," where symbolical characters have been used, so that no capacity can fail of at once perfectly comprehending the Sexual System, or to his more expensive and elaborate work the "NEW ILLUSTRATION OF THE SEXUAL SYSTEM OF LINNÆUS," *Folio*. This last work was honoured by a diamond ring presented to the Doctor, by ALEXANDER, the present emperor of Russia.

CLASSES AND ORDERS OF THE SEXUAL SYSTEM.

CLASSES.	ORDERS.
I. MONANDRIA	1. <i>Monogynia.</i> 2. <i>Digynia.</i>
II. DIANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Digynia.</i> 3. <i>Trigynia.</i>
III. TRIANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Digynia.</i> 3. <i>Trigynia.</i>
IV. TETRANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Digynia.</i> 3. <i>Tetragynia.</i>
V. PENTANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Digynia.</i> 3. <i>Trigynia.</i> 4. <i>Tetragynia.</i> 5. <i>Pentagynia.</i> 6. <i>Polygynia.</i>
VI. HEXANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Digynia.</i> 3. <i>Trigynia.</i> 4. <i>Tetragynia.</i> 5. <i>Polygynia.</i>
VII. HEPTANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Digynia.</i> 3. <i>Tetragynia.</i> 4. <i>Heptagynia.</i>
VIII. OCTANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Digynia.</i> 3. <i>Trigynia.</i> 4. <i>Tetragynia.</i>
IX. ENNEANDRIA	{ 1. <i>Monogynia.</i> 2. <i>Trigynia.</i> 3. <i>Hexagynia.</i>

CLASSES.	ORDERS.
X. DECANDRIA	{ 1. <i>Monogynia</i> . 2. <i>Digynia</i> . 3. <i>Tri- gynia</i> . 4. <i>Pentagynia</i> . 5. <i>De- cagynia</i> .
XI. DODECANDRIA	{ 1. <i>Monogynia</i> . 2. <i>Digynia</i> . 3. <i>Tri- gynia</i> . 4. <i>Pentagynia</i> . 5. <i>Do- decagynia</i> .
XII. ICOSANDRIA	{ 1. <i>Monogynia</i> . 2. <i>Digynia</i> . 3. <i>Tri- gynia</i> . 4. <i>Pentagynia</i> . 5. <i>Po- lygynia</i> .
XIII. POLYANDRIA	{ 1. <i>Monogynia</i> . 2. <i>Digynia</i> . 3. <i>Tri- gynia</i> . 4. <i>Tetragynia</i> . 5. <i>Pen- tagynia</i> . 6. <i>Hexagynia</i> . 7. <i>Po- lygynia</i> .
XIV. DIDYNAMIA	1. <i>Gymnospermia</i> . 2. <i>Angiospermia</i> .
XV. TETRADYNAMIA	1. <i>Siliculosa</i> . 2. <i>Siliquosa</i> .
XVI. MONADELPHIA	{ 1. <i>Triandria</i> . 2. <i>Pentandria</i> . 3. <i>Oc- tandria</i> . 4. <i>Emneandria</i> . 5. <i>De- candria</i> . 6. <i>Endecandria</i> . 7. <i>Do- decandria</i> . 8. <i>Polyandria</i> .
XVII. DIADELPHIA	{ 1. <i>Pentandria</i> . 2. <i>Hexandria</i> . 3. <i>Oc- tandria</i> . 4. <i>Decandria</i> .
XVIII. POLYADELPHIA	{ 1. <i>Pentandria</i> . 2. <i>Icosandria</i> . 3. <i>Po- lyandria</i> .
XIX. SYNGENESIA	{ 1. <i>Polygamia æqualis</i> . 2. <i>Polygamia superflua</i> . 3. <i>Polygamia frustra- nea</i> . 4. <i>Polygamia necessaria</i> . 5. <i>Polygamia segregata</i> . 6. <i>Mo- nogamia</i> .

CLASSES.	ORDERS.
XX. GYNANDRIA	{ 1. <i>Diandria</i> . 2. <i>Triandria</i> . 3. <i>Tetrandria</i> . 4. <i>Pentandria</i> . 5. <i>Hexandria</i> . 6. <i>Decandria</i> . 7. <i>Dodecandria</i> . 8. <i>Polyandria</i> .
XXI. MONŒCIA	{ 1. <i>Monandria</i> . 2. <i>Diandria</i> . 3. <i>Triandria</i> . 4. <i>Tetrandria</i> . 5. <i>Pentandria</i> . 6. <i>Hexandria</i> . 7. <i>Heptrandria</i> . 8. <i>Polyandria</i> . 9. <i>Monadelphia</i> . 10. <i>Syngenesia</i> . 11. <i>Gynandria</i> .
XXII. DIŒCIA	{ 1. <i>Monandria</i> . 2. <i>Diandria</i> . 3. <i>Triandria</i> . 4. <i>Tetrandria</i> . 5. <i>Pentandria</i> . 6. <i>Hexandria</i> . 7. <i>Octandria</i> . 8. <i>Enneandria</i> . 9. <i>Decandria</i> . 10. <i>Dodecandria</i> . 11. <i>Polyandria</i> . 12. <i>Monadelphia</i> . 13. <i>Syngenesia</i> . 14. <i>Gynandria</i> .
XXIII. POLYGAMIA	1. <i>Monœcia</i> . 2. <i>Diœcia</i> . 3. <i>Triœcia</i> .
XXIV. CRYPTOGAMIA	{ 1. <i>Filices</i> . 2. <i>Musci</i> . 3. <i>Algæ</i> . 4. <i>Fungi</i> .
APPENDIX	1. <i>Palma</i> *.

* Vide Plate of the Classes, at the end of this work.

CHAP. II.

EXPLANATION OF THE TITLES OF THE TWENTY-FOUR CLASSES.

HAVING, in the preceding chapter, given the divisions of the system, we shall in this explain the meaning of the terms used for the titles of the classes. As these terms in the *Greek* language, from whence they are taken, are all expressive of the principal circumstance that obtains in the class to which they are applied, the explanation of them will itself give us a good insight into the proper characters of the several classes, and the sexual distinctions on which they are founded: however it will be necessary to say something more particular concerning many of them afterwards in the chapters we shall allot for each of them separately.

Class 1. MONANDRIA. 2. DIANDRIA. 3. TRIANDRIA. 4. TETRANDRIA. 5. PENTANDRIA. 6. HEXANDRIA. 7. HEPTANDRIA. 8. OCTANDRIA. 9. ENNEANDRIA. 10. DECANDRIA.—These ten classes, which consist of bisexual flowers, take their denominations from the number of stamina, or male parts of the flower. The word here compounded with the numerical terms, signifies a *male*; so that the title *Monandria* expresses that the flowers of this class have but *one male*, that is, one stamen; *Diandria*, *two* stamina; *Triandria*, *three*; *Tetrandria*, *four*; *Pentandria*, *five*; *Hexandria*, *six*; *Heptandria*, *seven*; *Octandria*, *eight*; *Enneandria*, *nine*; and *Decandria*, *ten*. It must be observed, however, that the flowers being *bisexual*, as above mentioned, is in all these classes a necessary condition; for should the female part be wanting, the plant would belong to some other class, notwithstanding the number of stamina may be such as would otherwise

refer it to one of these : and this caution we give once for all to avoid repetitions, that when we use the term *bisexual*, we mean that it is a condition not to be dispensed with.

Class 11. DODECANDRIA.—This term, in the *Greek*, imports that the flowers have *twelve males*, or stamina. However, the class is not confined to this number, but includes all such *bisexual* flowers as are furnished with any number of stamina, from *twelve* to *nineteen* inclusive : no flowers have yet been found to have eleven stamina, which is the reason no class has been allotted to that number.

Class 12. ICOSANDRIA.—This term imports, that the flowers have *twenty* males, or stamina ; but here again the title is to be understood with great latitude ; for though the plants that belong to this class are rarely found with less than twenty stamina, yet they frequently have a greater number : and they are therefore not to be known with certainty from those of the next class, without having recourse to their classic character ; which, not being expressed in the title, we forbear the explanation of here, as we shall give it in the chapter allotted for this class.

Class 13. POLYANDRIA.—This term imports, that the flowers have *many* stamina.

Class 14. DIDYNAMIA.—This term signifies the *power*, or *superiority* of *two*, and is applied to this class, because its flowers have four stamina, of which there are two longer than the rest. This circumstance alone is sufficient to distinguish this class from the fourth, where the four stamina are equal ; but the flowers of this class have also their particular character, besides what the title expresses, their corollæ being mostly *ringent*, as will be shown in its place*.

Class 15. TETRADYNAMIA.—This term expresses the *power*, or *superiority* of *four* ; and accordingly there are in the flowers of

* See Chap. XVII. See also Part I. Chap. III. where the term *ringent* is explained.

this class six stamina, four of which are longer than the rest; which circumstance distinguishes them from those of the sixth class, where the six stamina are equal: but these flowers have their particular character also, their corollæ being *cruciform**,

Class 16. *MONADELPHIA*.—The word here, compounded with the numerical term, signifies a *brother*. This relation is employed to express the union of the filaments of the stamina, which in this class do not stand separate, but join at the base, and form one substance, out of which they proceed as from a common mother; and the title of the class expresses a *single* brotherhood, meaning that there is but *one* set of stamina so united, which distinguishes the class from the two following ones. The number of stamina in this class is not limited: the flowers have their particular character†.

Class 17. *DIADELPHIA*.—This term expresses a *double* brotherhood, or *two* sets of stamina, united in the manner explained in the preceding class. The number of the stamina is not limited: the flowers of this class have a very particular character, their corolla being *papilionaceous*, as will be shown in its place‡.

Class 18. *POLYADELPHIA*.—This term expresses *many* brotherhoods, or sets of stamina; the flowers have no classic character, farther than is expressed in the title.

Class 19. *SYNGENESIA*.—This class contains the compound flowers described in Part I. Chap. 19. The title signifies *congeneration*, alluding to the circumstance of the stamina; in which, though the filaments stand separate, yet the antheræ, subservient to generation, are united in a cylinder, and perform their office *together*. The classic character will be explained in its place||.

* See Chap. XVIII. See also Part I. Chap. III. where the term *cruciform* is explained.

† See Chap. XIX.

‡ See Chap. XX. See also Part I. Chap. III. for the explanation of the term *papilionaceous*.

|| See Chap. XXII.

Class 20. GYNANDRIA.—The term is compounded of two words, that signify *wife* and *husband*; and alludes to the singular circumstance of this class, in the flowers of which the stamina grow out of the pistillum.

Class 21. MONŒCIA.—The word here, compounded with the numerical term, signifies a *house* or *habitation*. To understand the application of this title, we must know, that the plants of this class are not *bisexual*, but *androgynous**, the flowers that have the stamina wanting the pistillum, and those that have the pistillum wanting the stamina. Now the term *monœcia*, which signifies a *single* house, alludes to this circumstance, that in this class the male and female flowers are both found on the *same* plant, whereas in the next they have *distinct* habitations.

Class 22. DICŒCIA.—This term, which signifies *two* houses, is applied to this class (the plants of which are *male* and *female*), to express the circumstance of the *male* flowers being on one plant, and the *female* on another; the contrary of which is the case of the androgynous class *Monœcia* last explained.

Class 23. POLYGAMIA.—The term signifies *plurality* of *marriages*. This class produces, either upon the same or different plants, *bisexual* flowers, and also flowers of *one* sex only, be it male or female.

Class 24. CRYPTOGAMIA†.—The term signifies *concealment* of *marriages*; this class consisting of such plants as either bear their flowers concealed within the fruit‡, or have them so small, as to be imperceptible.

* See Part I. Chap. XXI.

† Perhaps the Greek words should have been expressed; but the editor was fearful of adding them, as Mr. Lee has knowingly omitted them. These may, however, be seen in Doctor Thornton's GRAMMAR OF BOTANY.

‡ The Ficus, whose flowers are within the fruit, used to be put in this class; but is since removed to the twenty-third class, *Polygamia*.

CHAP. III.

EXPLANATION OF THE TITLES OF THE ORDERS.

THE titles of the orders have been given in Chap. I. It remains to explain them.

Class 1 to 13, inclusive.—The orders of the first thirteen classes take their denominations from the number of the *pistilla*, or female part of the plant, which is usually reckoned from the *base* of the *style*, if there be any; but if the style be wanting, the number is fixed from the *stigmata*. The *Greek* word, compounded with the numerical terms in the titles of these orders, signifies a *wife*: MONOGYNIA implies *one wife*, or one style; DIGYNIA, *two styles*; TRIGYNIA, *three*; TETRAGYNIA, *four*; PENTAGYNIA, *five*; HEXAGYNIA, *six*; DECAGYNIA, *ten*; and POLYGYNIA, *many*. These are the titles that occur in the orders of these thirteen classes; and this general explanation of them will be thought sufficient, as from the table given in the first chapter it appears how they are employed in the classes.

Class 14. DIDYNAMIA.—Of the three orders of this class the two first are founded on a distinction in the fruit. The title of the first order, GYMNOSPERMIA, is expressive of such plants as have *naked* seeds; and that of the second, ANGIOSPERMIA, of such as have their seeds in a *vessel*, or *pericarpium*. A third order, POLYPETALA, is expressive of such plants as have *many petals*: this order seems to have been established in favour of one genus of plants only, the *melianthus*, the flowers of which are *polypetalous*, though those of all the rest of this class are *monopetalous**.

* This order is properly omitted in the *Systema Naturæ*, published in 1756. See the note on this order, in Chap. XVII.

Class 15. TETRADYNAMIA.—The two orders of this class are founded on a distinction in the *pericarpium*. In the first order, SILICULOSA, the *pericarpium* is a SILICULA, *little siliqua*; which differs from the SILIQUA in being round, and having the apex of the dissepiment, which had been the style, prominent beyond the valves, often so far as to be equal in length to the siliqua. In the second order, SILIQUOSA, the *pericarpium* is a SILIQUA, which is long, and without any remarkable extension of the style.

Class 16. MONADELPHIA. 17. DIADELPHIA. 18. POLYADELPHIA. The orders of these three classes are founded on the number of the stamina in each brotherhood, or distinct set of stamina. The titles of the orders being the same that are used for the titles of the early classes of the system, the explanation need not be repeated here.

Class 19. SYNGENESIA.—To understand the orders of this class, we must explain what is meant by *polygamy* in flowers. We have already treated of polygamous *plants*, and shown that the term *polygamous*, as there applied, alluded to the intercommunication of the male or female flowers with the bisexual ones, either upon the same, or a distinct plant; but in respect to flowers, the term is applied to a single flower only; for the flowers of this class being compound, a *polygamy* arises from the intercommunication of the several *florets* in one and the same flower. Now the *polygamy* of *flowers*, in this sense of the word, affords four cases, which are the foundations of the four first orders of this class. First order, POLYGAMIA ÆQUALIS, *equal polygamy*, is when *all* the florets are bisexual. Second order, POLYGAMIA SUPERFLUA, *superfluous polygamy*, when *some* of the florets are *bisexual*, and *others female* only; for in this case, as the fructification is perfected in the *bisexual*, the addition of the females is a superfluity. Third order, POLYGAMIA FRUSTRANEA, *frustraneous* or *ineffectual polygamy*, when *some* of the florets are *bisexual*, and *others neuter*; for in this case the addition of the neuters is of no assistance to the fructification. Fourth order, POLYGAMIA NECESSARIA, *ne-*

cessary polygamy, when some of the florets are *male*, and the rest *female*; for in this case, there being no bisexual, the polygamy arising from the composition of the florets of different sexes, is *necessary* to perfect the fructification. Fifth order, POLYGAMIA SEGREGATA. The title signifies to be separated, the plants of this order having partial cups growing out of the common calyx which surround and divide the flosculi or florets. Sixth order, MONOGAMIA: the title signifies a *single* marriage, and is opposed to the *polygamia* of the four other orders; for in this, though the antheræ are united, which is the essential character of the flowers of this class, the flower is *simple*, and not compounded of *many* florets, as in the other orders.

Class 20. GYNANDRIA.—The orders of this class are founded on the number of stamina. The titles have been already explained.

Class 21. MONÆCIA. 22. DIÆCIA. These two classes, whose flowers have no fixed character, but that of not being bisexual, take in the characters of almost every other class; and the orders have accordingly been disposed under the titles of those classes to which their respective flowers would have belonged if the stamina and pistillum had been under the same covers. As the explanation of all these titles has been given in the last chapter in the explanation of the classes, it need not be repeated here.

Class 23. POLYGAMIA.—In this class the titles of the two first orders are the same with the titles of the twenty-first and twenty-second classes, and are to be understood in the same manner; that is, 1. MONÆCIA, when the polygamy is on the *same* plant; and 2. DIÆCIA, when it is on *distinct* plants. The order TRIÆCIA has been established in favour of a single genus, the *ficus*; in which the polygamy is on *three* distinct plants, one producing *male* flowers, another *female*, and a third *bisexual*, or *androgynous*.

Class 24. CRYPTOGAMIA.—The orders of this class are, 1. FILICES, *ferns*. 2. MUSCI, *mosses*. 3. ALGÆ, *flags*; and 4. FUNGI,

mushrooms. As the explanation of the character of these orders will come more properly into the chapters that treat particularly of each class, we shall content ourselves here with having interpreted the titles as above.

CHAP. IV.

OF THE FIRST CLASS, MONANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with but *one* stamen. The orders are *two*, viz.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. This order contains twenty *genera*, distinguished into 1. *Scitamina*, with an inferior fruit, one-celled or three-celled, viz. CANNA (*Indian reed*)... AMOMUM... COSTUS ... ALPINIA... MARANTA ... CURCUMA (*turmeric*) ... KEMPTFERIA... THALIA ... MYROSMA... RENEALMIA..... HELLENIA..... HEDYCHIUM... HORNSTEDTIA... and PHRYMUM. 2. *Fruit inferior, four-celled*; LOPEZIA. 3. *Fruit superior*, PHYLIDRUM... CUCULLARIA... QUALEA... USTERIA. 4. *One-seeded*... BOERHAAVIA... SALICORHIA (*saltwort*)... HIPPURIS (*mare's tail*)... POLLICHIA... MITHRIDATEA. 5. *Naked seeds*, CHARA... ZOSTERA (*sea-wrack*).

Order 2. DIGYNIA, comprehending such plants as have two styles. This order contains five *genera*, viz. CORISPERMUM... CALITRICHE... BLITUM (*strawberry blite*)... CINNA... MINIARUM... and LACISTEMA.

CHAP. V.

OF THE SECOND CLASS, DIANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *two* stamina. The orders are three, viz.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. This order contains forty-one *genera*, distinguished into, 1. Such as have *regular corolla, one-petalled, flowers inferior*, of which there are eleven, viz. NYCTANTHES...JASMINUM (*jasmine*)...LIGUSTRUM (*privet*)...PHILLYREA...OLEA (*olive*)...CHIONANTHUS (*snow-drop tree*)...SYRINGA...ERANTHEMUM...WULFENIA...PIMELEA...and GALIPEA. 2. Such as have *irregular corolla, and the fruit angiospermous*; of which there are *thirteen*, viz. VERONICA (*speedwell*)...PÆDEROTA...JUSTICIA...DIANTHERA...GRATIOLA...SCHWENKIA...PINGUICULA (*butter-wort*)...UTRICULARIA (*bladder-wort*)...CALCEOLARIA (*ladies' slipper*)...CYRTANDRA...BAEA...GIINIA...and SCIURIS. 3. Such as have an *irregular corolla, and the fruit gymnospermous*; of which there are *nine*, viz. VERBENA (*vervain*)...LYCOPUS (*water horehound*)...AMETHYSTEA...CUNILA...ZIZIPHORA...MONARDA...ROSMARINUS (*rosemary*)...SALVIA (*sage*)...and COLLINSONIA. 4. *Flowers inferior, polypetalous*; of which there are four, viz. FONTANESIA...LITHOPHYLA...LINOCIERA...and DIALIUM. 5. *Flowers superior*, MONNA...CIRCEA (*enchanter's nightshade*)...and GLOBBA. 6. *Flowers apetalous*, ANCISTRUM...and ARUNA.

Order 2. DIGYNIA, comprehending such plants that have *two* styles. This order contains but two genera, viz. ANTHOXANTHUM (*vernal grass*)...and CRYPISIS.

Order 3. TRIGYNIA, comprehending such plants that have *three* styles. There is but one genus of this order, viz. *Piper*.

CHAP. VI.

OF THE THIRD CLASS, TRIANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with three stamina. The orders are three.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. This order contains forty-five genera, distinguished into, 1. *Those whose flowers are superior*, of which there are ten, viz. VALERIANA (*valerian*)... MELOTHRIA... DILATRIS... MELOTRIA ...CROCUS... ANTHOLYZA... GLADIOLUS... IRIS... IXIA... ARISTEA... MORÆA. 2. *Flowers inferior, not glumaceous*, of which there are twenty-three, viz. WACHENDORFIA ... COMMELINA... CALLISIA... XYRIS... WITSENIA... MARICA... ZIPHIDIUM... GOMMELINA... OXYBAPHUS... MACROLOBIUM... ROHRIA... HYPOCRATEA... TONSELLA... LÆFLINGIA... WILLICHIA... SYENA... RUMPHIA... FISSILIA... CNEORUM... COMOCLADIA... OLAX... ROTALA... ORTEGIA... POLYCHEMUM. 3. Such as have an *imbricated amentum*, and are *gymnospermous*; of which there are twelve, viz. SCHÆNUS... CYPERUS... SCIRPUS ... ERIOPHORUM... LYCEUM... NARDUS... KYLLINGIA... FUIRENA... MASSANIA... MIEGIA... CENCHRUS... and POMMERCULLIA.

Order 2. DIGYNIA, comprehending such plants as have *two* styles. This order contains thirty-three genera, viz. CORNUCOPIA... SACCHARUM... PANICUM... PHILEUM (*cat's-tail grass*)... ALOPECURUS (*fox-tail grass*)... MILIUM (*millet-grass*)... AGROSTIS (*bent-grass*)... AIRA... MELICA... POA (*meadow-grass*)... BRIZA (*quaking-grass*) ... UNIOLA ... DACTYLIS (*cock's-foot grass*) ... CYNOSURUS (*dog's-tail grass*)... FESTUCA... BROMUS..... STIPA (*feather-grass*)... AVENA (*oat*)... LAGURUS (*hare's-tail grass*) ... ARUNDO (*reed*)... ARISTIDA... LOLIUM (*darnel*)... FLYMUS... SECALE (*rye*)... HORDEUM

(barley) ... TRITICUM (*wheat*) ... PHALARIS (*canary-grass*) ... PASPALUM ... ROTTBOELLIA ... PEROTIS ... LEERSIA ... PAPPOPHORUM ... and LAPPAGO.

Order 3. TRIGYNIA, comprehending such plants as have *three* styles. This order contains twelve genera, viz. ERIOCAULON ... MONTIA ... PROSERPINACA ... TRIPLARIS ... HOLOSTEUM ... POLYCARPON ... MOLLUGO ... MINUARTIA ... QUERIA ... LECHEA ... KÆNIGIA ... and DONATIA.

CHAP. VII.

OF THE FOURTH CLASS, TETRANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *four* stamina. The flowers of this class may be known from those of the fourteenth by this distinction, that the stamina are of an equal length; whereas in those of the fourteenth, which have four stamina likewise, there are two long and two short. The orders of this class are *three*; viz.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. This order contains ninety-four genera, distinguished into, 1. *Flowers monopetalous, one-seeded, inferior*, viz. GLOBULARIA. 2. *Flowers monopetalous, one-seeded, superior, aggregate*, as DIPSACUS (*teasel*) ... KNAUTIA ... SCABIOSA (*scabious*) ... and ALLIONIA. 3. *Flowers monopetalous, four-seeded*, as MATTUŞHKEA. 4. *Flowers monopetalous, one-fruited, inferior*, as PYROSTRIA ... MYONIMA ... PETITIA ... AQUARTIA ... ROUSSEA ... CALLICARPA ... WALLO니아 ... WITHERINGIA ... ÆGIPHILA ... CEFHALANTHUS ... LASIOSTOMA

...SCOPARIA... CENTUNCULUS... PLANTAGO... POLYPRENUM... BUD-
LEIA... EXACUM... MYRMECIA... LABATIA... PENÆA... and BLÆRIA.
5. *Flowers monopetalous, one-fruited, superior, as* CIOMELIA...
CUNNINGHAMIA... SCOLOSANTHUS... PAVETTA... IKORA... PETESIA...
CATESBÆA... FRÆLICHIA... HOFFMANNIA... ERNODEA... SIDERODEN-
DRUM... COCCOCYPSILUM... MITCHELLA... HEDYOTIS... OLDENLANDIA
... HYDROPHYLAX... MANETTIA... CARPHALEA... BELLARDIA... SAN-
GUISORBA (*great burnet*). 6. *Flowers monopetalous, dicoccous, in-*
ferior; HOUSTONIA. 7. *Flowers monopetalous, dicoccous, superior,*
stellate; RUBIA (madder)... GALIUM (bed-straw)... ASPERULA... SHE-
RARDIA... SPERMACOCE... KNOXIA... DIODIA... CRUCIANELLA. 8.
Flowers monopetalous, tetracoccous, inferior; SIPHONANTHUS. 9.
Flowers four-petalled, viz. EPIMEDIUM ... CORNUS ... FAGARA'...
AMANNIA... PTELEA... LUDWIGIA... SANTALUM... TRAPA... SAMARA
BLACKBURNIA... SKIMMIA... MONETIA... HARTOGIA... CURTISIA...
OTHERA... ORIXA... CISSUS... and GLOSSOMA. 10. *Flowers incom-*
plete, viz. DORSTENIA... ELÆAGNUS... KRAMERIA... RIVINA... SAL-
VADORA... CAMPHOROSMA... ALCHEMILLA... STRUTHIOLA... COMETES
... OPERCULARIA... PROTEA... RUPALA... BANKSIA... EMBOTRIUM...
POTHOS, GONATOCARPUS, ACCENA, ISNARDIA.

Order 2. DIGYNIA, comprehending such plants as have *two* styles. This order contains *seven* genera, viz. CRUZITA, BUF-
FONIA, HAMAMELIS, CUSCUTA, HYPECOUM, GALOPINA, and
NERTERIA.

Order 3. TRIGYNIA, has *one* genus only; BOSCIA.

Order 4. TETRAGYNIA, comprehending such plants as have
four styles. This order contains *seven* genera, viz. ILEX (*holly*)
... COLDENIA... POTAMOGETON... RUPPIA... SAGINA... MYGINDA...
and TILLÆA.

CHAP. VIII.

OF THE FIFTH CLASS, PENTANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *five stamina*. The orders are seven.

Order 1. MONOGYNIA, *one pistillum*. This order contains 208 genera, distinguished into, 1. *Flowers monopetalous, inferior, one-seeded*, of which there are *six* genera; MIRABILIS (*Marvel of Peru*) ... TRICATUS... PLUMBAGO... WEIGELIA... QUINCHAMALA... CORYMBIUM. 2. *Flowers monopetalous, inferior, two-seeded*; ASPERIFOLIÆ, of which there are *two* species; CERINTHE (*honey-wort*)... MESSERSCHMIDIA. 3. *Flowers monopetalous, inferior, four-seeded*; ASPERIFOLIÆ, of which there are *twelve* species... ECHIUM (*viper's bugloss*)... HELIOTROPIUM (*turnsole*) ... PULMONARIA ... LITHOSPERMUM (*gromwell*) ... ONOSMA ... SYMPHYTUM (*comfrey*)... BORAGO (*borage*)... LYCOPSIS... BUGLOSS... ASPERUGO... CYNOGLOSSUM (*hound's-tongue*) ... ANCHUSA (*alkanet*) ... MYOSOTIS (*mouse-ear, scorpion's-grass*). 4. *Flowers monopetalous, inferior, five-seeded*, of which there is *one* species; NOLANA. 5. *Flowers monopetalous, inferior, with the seeds enclosed in a pericarp*, of which there are *eighty-nine* genera; CORIS... HYDROPHYLLUM (*water-leaf*)... GALAX... BARRERIA ... CORTUSA ... ANAGALLIS (*pimpernel*)... LYSIMACHIA (*loostrike*)... DORENA ... CYCLAMEN (*sow-bread*)... DODECATHÉON (*Meadia, or Virginia cowslip*)... SOLDANELLA... LITA... PRIMULA (*primrose*)... ANDROSACE... ARETIA... BACOPA... HOTTONIA (*water-violet*)... SHEFFIELDIA... MENYANTHES (*buck-bean*)... ALLAMANDA... THEOPHRASTA... GENIOSTOMA... SPIGELIA (*worm-grass*)... SPHENOCLEA... OPHIORHIZA... RETZIA... CONVULVULUS (*bind-weed*)... LISIANTHUS... DATURA... HYOSCYAMUS (*henbane*)... NICOTIANA (*tobacco*) ... VERBASCUM (*mullein*) ... CHEIRONIA ...

PORANA... DIAPENSIA... PHLOX... POLEMONIUM (*Jacob's ladder*)...
 CANTUA... IPOMŒA... BROSSEA... AZALEA... EPACRIS... NERIUM (*ole-
 andar or rose-bay*)... ECHITES... PLUMIERIA... CAMERARIA... TA-
 BERNÆMONTANA... VINCA (*periwinkle*)... CERBERA... THOUINIA...
 TECTONA (*teak-tree*)... ARDISIA... BRUMELIA... GYNOPOGON... LAU-
 GERIA... VARRONIA... CORDIA... IGNATIA... EHRETIA... STYPHELIA...
 WILLOUGHBEIA... CARISSA... JACQUINIA... MYRSINE... BLADHIA...
 PÆDERIA... RAUWOLFIA... ARDUINA... CESTRUM... FAGRÆA...
 TOURNEFORTIA... STRYCHNOS (*poison-nut*)... CAPSICUM... SOLANUM
 (*night-shade*)... PHYSALIS (*winter cherry*)... JABOROSA... ATROPA...
 ELLISIA... LYCIUM... CRYPTOSTOMUM... CUMAX... TRIGUERA... SO-
 LANDRA... MENAIS... LEEA... SIDEROXYLUM (*iron-wood*)... CHRYSO-
 PHYLLUM (*star-apple*)... BASSOVIA... BRËBOTRYS. 6. *Flowers no-
 nopetalous, superior, of which there are thirty-eight genera*; SAMO-
 LUS... VIRECTA... BELLONIA... MACROCENENUM... DENTELLA... CHI-
 MARIIS... RONDELETIA... CINCHONA (*bark*)... PORTLANDIA... ROEL-
 LA... GOODENIA... PHYTEUMA... TRACHELIUM (*throat-wort*)... CAM-
 PANULA (*bell-flower*)... LOBELIA... SCHÆVOLA... SCHŒPSIA... MAT-
 THIOLA... MORINDA... PSYCOTRIA... COFFEA... CHIOCOCCA... SERISSA...
 CEPHIAELIS... VANGUENA... SOLENA... WEBERA... GARDENIA... UCRI-
 ANA... CANEPIORA... BERTIERA... LONICERA (*honey-suckle*)... TRI-
 OSTEUM... PLOCAMA... MUSSENDA... SCHWENKFELDIA... HAMELLIA
 ... ERITHALIS. 7. *Flowers tetrapetalous, one species*; STRŒMIA. 8.
Flowers pentapetalous, inferior, contains thirty-seven species; HIR-
 TELLA... RHAMNUS (*buck-thorn*)... CLEONOTHUS... CELASTRUS (*staff-
 tree*)... EUONYMUS (*spindle-tree*)... STAAVIA... EUPAREA... BILLARDI-
 ERA... RUYSCHIA... VITIS (*wine*)... ESCALLONIA... MANGIFERA (*man-
 go-tree*)... ZIZYPHUS... SCHREBERA... ELÆODENDRUM... WALKERA...
 CORYNOCARPUS... HUMBOLDTIA... PILOCARPUS... CEDRELA... CALO-
 DENDRUM... SCOPOLIA... POLYCARDIA... PITTOSPORUM... BUTTENRIA
 ... AYENIA... GLUTA... DIOSMA... SPRENGELIA... HOVENIA... NAUCLEA
 ... IMPATIENS (*balsam*)... CLAYTONIA... RORIDULA... ITEA... ÆGI-
 CERAS... SAUVAGESIA... VENTILAGO... BRUNIA. 9. *Flowers pentape-
 talous, superior, contains thirteen genera*; RIBES (*currant*)... HE-
 DERA (*ivy*)... PLECTRONIA... STRUMPFIA... PHYLICA... CARPODETUS...
 GRONOVIA... JASIONE... CYPHIA... ARGOPHYLLUM... LIGHTFOOTIA...

LAGÆCIA...CONOCARPUS. 10. *Flowers incomplete, inferior, contains six genera*; ACHYRANTHES...CHENOLIA...CELOSIA (*cock's-comb*)...ILLECEBRUM...GLAUX...COLLETIA. 11. *Flowers incomplete, superior*; THESIUM...HELICONIA...STRELITZIA.

Order 2. DIGYNIA, *two pistilla*, contains *eighty* genera, distinguished into, 1. *Flowers monopetalous, inferior, which contains sixteen genera*; STAPELIA...CYNANCHUM...PERIPLOCA...HOSTEA...APOCYNUM (*dog's-bane*)...PERGULARIA...ASCLEPIAS (*swallow-wort*)...CEROPEGIA...MELODINUS...SWERTIA...GENTIANA...CRESSA...NAMA...HYDROLEA...ROCHEFORTRA...DICHONDRA. 2. *Flowers pentapetalous, inferior, contains five genera*; VELESIA...LINCONIA...BUMALDA...HEUCHERA...ANABASIS. 3. *Flowers incomplete, contains eight genera*; SALSOLA (*saltwort*)...CHENOPODIUM (*goose-foot*)...BETA (*beet*)...HERNIARIA (*rupture-wort*)...GOMPHRENA...BOSEA...ULMUS (*elm*)...MICROTEA. 4. *Flowers pentapetalous, superior, capsuled, contains one genus*; VASSLIA. 5. *Flowers pentapetalous, superior, two-seeded, contains fifty genera*; UMBELLATE*, *with both genera and partial umbels*; PHYLIS...ERYNGIUM (*eringo*)...HYDROCOTYLE...AZORELLA...CUSSONIA...SANICULA (*sanicle*)...ASTRANTIA (*masterwort*)...HERACLEUM (*cow-parsnip*)...ÆNANTHE (*water-dropwort*)...ECHINOPHORA (*prickly samphire*)...CAUCALIS...ARTEDIA...DAUCUS...TORDYLIUM (*hartwort*)...CORIANDRUM (*coriander*)...LASERPITIUM (*laserwort*)...PEUCEDANUM (*sulphurwort*)...AMMI (*bishop's-wced*)...HASSELQUISTIA...CONIUM (*hemlock*)...EXOCANTHA...BUNIUM (*earthnut*)...ATHAMANTA (*stone-parsley*)...BUPLEURUM...SIUM (*water-parsnip*)...SELINUM...CUMINUM (*cumin*)...FERULA (*giant-fennel*)...CRITHMUM (*samphirc*)...BUBON...CACHRYS...LIGUSTICUM (*lovage*)...MEUM...ANGELICA...SISON (*honestwort*). 6. *With partial involucre only*; ÆTHUSA (*fool's parsley*)...SCANDIX (*chervil*)...CHÆROPHYLLUM (*cow's parsley*)...PHELLANDRIUM (*water-hemlock*)

* These plants, and those of the two distinctions next following, which are *gynodispermous* also, are the *umbellate* plants of *Tournefort's* seventh class. See his *Institution*, R. H. In dry soils they are aromatic, warm, resolvent, and carminative; but in moist places poisonous. The virtue is in the roots and seeds. AUTHOR.

...IMPERATORIA.....SESELI....CICUTA (*water-cowbane*). 7. *Without any involucre, or scarcely any general involucre, and never any partial one; SMYRNIUM (Alexanders)... CARUM (caraway)... THAPSIA... PASTINACA (parsnip)... ANETHUM (dill)... ÆGOPIDIUM (goat-weed)... APIUM (smallage and parsley)... PIMPINELLA (burnet saxifrage).*

Order 3. TRIGYNIA, *three pistilla*, contains *twenty* genera; VIBURNUM (*wayfaring tree*) ... SAMBUCUS (*elder*)... SEMECARPUS... RHUS... CRASSINE... REICHELIA... SPATHELIA... STAPHYLEA (*bladder-nut tree*)... TAMARIX (*tamarisk*)... DRYPIS... TURNERA... SALMASIA... SAROTHTA ... ALSINE (*chickweed*), ... TELEPHEUM ... CORRIGIOLA... PORTULACARIA... PHARMACEUM... XYLOPHYLLA... BASELLA.

Order 4. TETRAGYNIA, *four pistilla*, contains *two* orders; PARNASSIA... EVOLVULUS.

Order 5. PENTAGYNIA, *five pistilla*, contains *eleven* genera which are disposed under the following arrangement. 1. *Flowers superior; ARALIA... GLOSSOPETALUM.* 2. *Flowers inferior; URSULA... GISECKIA... LINUM (flax)... ALDROVANDA... DROSELA (sundew)... MAHERNIA... COMMERSONIA... SIBBALDIA... STATICE (thrift and sea lavender).*

Order 6. DECAGYNIA, *ten pistilla*, contains *one* genus only; SCHEFFLERA.

Order 7. POLYGYNIA, contains *two* genera; MYOSURUS (*mouse-tail*)... ZANTHORHIZA.

CHAP. IX.

OF THE SIXTH CLASS, HEXANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *six* stamina. The flowers of this class may be known from those of the fifteenth, by this distinction, that the stamina are of *equal* length; whereas in those of the fifteenth, which have six stamina likewise, there are *four long*, and *two short*. The orders of this class are *five*, containing *one hundred and twenty-one* genera.

Order 1. MONOGYNIA, *one pistillum*, contains 93 genera, which fall under the following subdivisions: 1. *Flowers furnished both with calyx and corolla, but without spathes*, which contains 24 genera; BROMELIA (*ananas, or pine-apple*)...PITCAIRNIA...TILLANDSIA...BURMANNIA...TRADESCANTIA (*spiderwort*)...STEPHANIA...FRANKENIA (*sea-heath*)...COSSIGNEA...LORANTHUS...HILLIA...SCHRADERA...DUROIA...RICHARDIA...TACCA...BARBACENIA...BERBERIS (*barberry*)...LEONTICE...NANDINA...PRINOS (*winter-berry*)...PSATHURIA...ISERTIA...CANARINA...ACHRAS...CAPURA. 2. *Flowers furnished with calyx, corolla, and spathes*, which contains three genera; CORYPHA...LICUALA...MNASIMUM. 3. *Flowers spathaceous or glumaceous*, which contains twenty genera; URANIA...HEMANTHUS (*blood-flower*)...LEUCOIUM (*snow-flake*)...STRUMARIA...GALANTHUS (*snowdrop*)...NARCISSUS...PANCRATIUM...AMARYLLIS...CRINUM...CYRTANTHUS...EUSTEPHIA...AGAPANTHUS...PONTERA...BULBOCODIUM...TULBAGIA...ALLIUM (*garlic, onion, &c.*)...CURCULIGO...APHYLLANTHES...MASSONIA...HYPOXIS. 4. *Flowers naked*, which contains thirty-eight genera; XEROPHYTA...ALSTRÆMERIA...LANARIA...HEMEROCALLIS (*day-lily*)...AGAVE...GETHYLLIS...ALOE...ALETRIS...VELTHEIMIA...POLYANTHES...CON-

VALLARIA (*lily of the valley, and Solomon's seal*)... SANSEVIERA...
 HYACINTHUS (*hyacinth*)... DRIMIA... MILLEA... ASPHODELUS (*asphodel*)... EUCOMIS... ANTHERICUM... ENARGEA... PHORMIUM... LACHENALIA... ORNITHOGALUM (*star of Bethlehem*)... ERIOSPERMUM...
 ... SCILLA (*squill*)... CYANELLA... PHILEZIA... LINDERA... DRACÆNA (*dragon-tree*)... ASPARAGUS (*sparrow-grass*)... POLLIA... GLORIOSA (*superb lily*)... ERYTHRONIUM (*dog-tooth violet*)... UVULARIA... FRITILLARIA (*fritillary, crown imperial*)... LILIUM (*lily*)... TULIPA (*tulip*)... YUCCA (*Adam's needle*)... ALBUCA. 5. *Flowers incomplete, which contains six genera*; ORONTIUM... ACORUS (*sweet flag*)... CALAMUS (*rattan*)... JUNCUS (*rush*)... THRINAX... PEPLIS (*water purslane*). 6. *Grasses, containing three genera*; BAMBUSIA (*bambu cane*)... GAHNIA... EBRIHARTA.

Order 2. DIGYNIA, *two pistilla*, contains *four* genera; FALKIA... ATRAPHAXIS... NEETRIS... ORYZA (*rice*).

Order 3. TRIGYNIA, *three pistilla*, contains *ten* genera, thus subdivided: 1. *Flowers inferior, containing nine* genera; WURMBEA... COLCHICUM (*meadow saffron*)... MELANTHIUM... MEDEOLA... HELONIAS... TRILLIUM... TRIGLOCHIN (*arrow-grass*)... RUMEX... SCHEUCHZERIA. 2. *Flowers superior, containing one* genus; FLAGELLARIA.

Order 4. HEXAGYNIA, *six pistilla*, contains *two* genera; DAMASONIUM... WENDLANDIA.

Order 5. POLYGYNIA, *many pistilla*, has but *one* genus; ALISMA.

CHAP. X.

OF THE SEVENTH CLASS, HEPTANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *seven* stamina. The orders of this class are four, viz.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. This order contains *eleven* genera, falling under these divisions: 1. *Flowers complete*, which contains *six* genera, viz. TRIENTALIS (*chickweed, winter-green*) DISANDRA ... ÆSCULUS (*horse-chestnut*)...PETROCARYA...PANCOVIA...JONESIA. 2. *Flowers incomplete*, which contains *five* genera; PISONIA...PETIVERIA... DRACONTIUM...CALLA...HOULTUYNIA.

Order 2. DIGYNIA, comprehending such plants as have *two* styles. This order contains but one genus, viz. LIMEUM.

Order 3. TETRAGYNIA, comprehending such plants as have *four* styles. Of this order there are but two genera, viz. SAURURUS (*lizard's tail*)...and ASTRANTHUS.

Order 4. HEPTAGYNIA, containing such plants as have *seven* styles. Of this order there is but one genus, viz. SEPTAS.

CHAP. XI.

OF THE EIGHTH CLASS, OCTANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *eight* stamina. The orders are four, viz.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. Of this order there are *fifty-three* genera, arranging under two heads: 1. *Flowers complete*, which contains *forty-two* genera, viz. MIMUSOPS...CUPANIA...DIMOCARPUS...TROPÆOLUM (*Indian cress*)...BÆCKIA...EPHIELIS...MOLINEA...HONCKENYA...HAGENIA...MEMECYLON...COMBRETUM...ROXBURGHIA...EPILOBIUM (*willow-herb*)...GAURA...ÆNOTHERA (*tree primrose*)...VITMANIA...RHEXIA...OSBECKIA...TETRATHECA...GRISLEA...KOELREUTERIA...PERSOONIA...GUAREA...CORREA...ANTICHORUS...ALLOPHYLUS...ORNITHROPHE...JAMBOLIFERA...XYLOCARPUS...ZIMENIA...LAWSONIA...MELICOCCA...AMYRIS...MELICOPE...GNIDIA...FUCHSIA...HEDWIGIA...MICHAXIA...CHLORA (*yellow centaury*)...VACCINIUM (*bilberry, whortleberry, cranberry*)...MENZIESIA...ERICA (*heath*). 2. *Flowers incomplete*, which contains *eleven* genera, as, OPHIRA...GRUBBIA...BUGINVILLEA...LACHNÆA...DIRCA...DAPHNE (*meze-reon, spurge-laurel, &c.*)...PASSERINA...STELLERA...DODONÆA...VALENTINIA...CEDROTA.

Order 2. DIGYNIA, comprehending such plants as have *two* styles. This order contains *five* genera, viz. GALENIA...WEINMANNIA...MÆHRINGIA...SCHMEIDELIA...and CODIA.

Order 3. TRIGYNIA, comprehending such plants as have *three* styles. This order contains *seven* genera, viz. POLYGONUM...

COCCOLOBA...PAULLINIA...CARDIOSPERMUM...SAPINDUS...SERIANA
...and PONÆA.

Order 4. TETRAGYNIA, comprehending such plants as have *four* styles. This order contains *six* genera, viz. PARIS...ADOXA (*tuberous moscatell*)...ELATINE...HALORAGIS...VEREA ...and FORSKOLEA.



CHAP. XII.

OF THE NINTH CLASS, ENNEANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *nine* stamina. The orders are *three*, containing *seven* genera, viz.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. This order contains *five* genera, viz. LAURUS (*bay*, &c.)...ANACARDIUM (*cashew-nut*)...CASSYTA...PANKE...and PLEGORHIZA.

Order 2. TRIGYNIA, comprehending such plants as have *three* styles. This order contains but *one* genus, viz. RHEUM (*rhubarb*).

Order 3. HEXAGYNIA, comprehending such plants as have *six* styles. Of this order there is but *one* genus, viz. BUTOMUS (*flowering rush*).

CHAP. XIII.

OF THE TENTH CLASS, DECANDRIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *ten* stamina. The orders are five, containing *one hundred and twenty* genera, viz.

Order 1. MONOGYNIA, comprehends such plants as have *one* style. This order contains fifty-six genera, distinguished into, 1. Such as have *flowers polypetalous, irregular*, of which there are *nineteen* genera, viz. SOPHORA... ANAGYRIS... CERCIS (*Judas' tree*)... BAUHINIA (*mountain ebony*)... PARKINSONIA... CASSIA... CÆSALPINIA... BASILETTO... GUILANDINA (*bonduc or neckar tree*)... DICTAMNUS (*fraxinella*)... PODALYRIA... PULTENÆA... HYMENÆA (*locust-tree*)... MYROXYLON... TOLUIFERA... CUBÆA... HYPERANTHERA... GÆRTNERA... GOMPHIA... RHODORA. 2. *Flowers polypetalous, equal*, of which there are *thirty-eight* genera, viz. RUTA (*rue*)... HEMATOXYLON (*log-wood*)... ADENANTHERA... MELIA (*bread-tree*)... TRICHILIA... ZYGOPHYLLUM (*bean-caper*)... QUASSIA... FAGONIA... TRIBULUS (*caltrops*)... THRYALLIS... MURRAYA... MONOTROPA (*yellow bird's-nest*)... JUSSIEUA... LIMONIA... MELASTOMA... LEDUM... QUISQUALIS... BERGERA... BUCIDA... CLETHRA... PYROLA (*winter-green*)... PROSOPIS... HEISTERIA... TURRÆA... DIONÆA (*Venus's fly-trap*)... EKEBERGIA... CYNOMETRA... SCHOTIA... CADIA... GILBERTIA... SANDORICUM... SWEITENIA (*mahogany*)... GUAIACUM (*lignum-vitæ*)... ZWINGERA... CERATOPETALUM... SCHOUSBÆA... PETALOMA... COOKIA... MERIANIA. 3. *Flowers monopetalous, equal*, containing *twelve* genera, viz. PANZERA... NICANDRA... CODON... INOCARPUS... STRIGILIA... ANDROMEDA... RHODODENDRON... KALMIA... EPIGÆA... GUALTERIA... ARBUTUS... and STYRAX.

Order 2. DIGYNIA, comprehending such plants as have *two* styles. Of this order there are *twelve* genera, viz. ROYENA... HYDRANGEA... CUNONIA... CHRYSOSPLENIUM (*golden saxifrage*)... SAXIFRAGA... TIARELLA... MITELLA... SCLERANTHUS (*knawell*)... TRIANTHEMA... SAPONARIA (*soapwort*)... DIANTHUS (*pink*)...and SILENE (*catch-fly*).

Order 3. TRIGYNIA, comprehending such plants as have *three* styles. Of this order there are *thirteen* genera, viz. CUCUBALUS (*campion*)... STELLARIA (*stickwort*)... ARENARIA (*sandwort*)... CHERLERIA... GARIDELLA... MALPIGHIA... BANISTERIA... TRIOPTERIS... ERYTHROXYLON... HIRÆA... DEUTZIA... BRUNNICHIA... GYPSOPHILA.

Order 4. PENTAGYNIA, comprehending such plants as have *five* styles. Of this order there are *sixteen* genera, viz. AVERHOA... SPONDIAS (*hog-plum*)... COTYLEDON (*navelwort*)... SEDUM (*stonecrop*)... PENTHORUM... OXALIS (*sorrel*)... SURIANA... LYCHNIS... AGROSTEMMA (*campion cockle*)... CERASTIUM (*mouse-ear chickweed*)... SPERGULA (*spurrey*)... GRIELUM... BERGIA... CNESTIS... JONQUETIA... ROBERGIA.

Order 5. DECAGYNIA, comprehending such plants as have *ten* styles. This order contains only *two* genera, viz. NEURADA...and PHYTOLACCA.

CHAP. XIV.

OF THE ELEVENTH CLASS, DODECANDRIA.

THIS class, notwithstanding its title, which is expressive of *twelve* stamina, consists of such plants as bear *bisexual* flowers, furnished with any number of stamina, from *twelve* to *nineteen* inclusive*. The orders are *six*, including *forty-one* genera, viz.

Order 1. MONOGYNIA, comprehends such plants as have but *one* style. This order contains *thirty-one* genera, which fall under the following sections: 1. *Corolla* none, of which there are *three* genera, viz. ASARUM (*asarabacca*)...BOCCONIA...STERCULARIA. 2. *Corolla* cut in four divisions, of which there are *seven* genera, viz. RHIZOPHORA...GARCINIA (*mangostan*) ...CRATÆVA...HALESIA ...APACTIS...DODECAS...and CRENÆA. 3. *Corolla* *five-petalled*, of which there are *thirteen* genera...TOMEX...EURYA...TRIUMFETTA...PEGANUM...KLEINHOFIA...NITRARIA...ARISTOTELIA...GRANGERIA...VATICA...HUDSONIA...CANELLA...PORTULACCA (*purslane*) ...TALINUM. 4. *Corolla* *six-petalled*, of which there are *five* genera...LYTRUM (*loosetrife*) ...CUPHEA...GINORIA...BLAKEA...AGATHOPHYLUM. 5. *Corolla* *seven-petalled*, of which there is *one* genus...BEFARIA. 6. *Corolla* *eight-cleft*, of which there is *one* genus...BASSIA. 7. *Corolla* *ten-petalled*, of which there is *one* genus...DECUMARIA.

Order 2. DIGYNIA, comprehends such plants as have *two*

* *Tormentilla* is an exception, belonging to the next class, though it has but *sixteen* stamina. The characters of the fructification in the next class, over-rule the number of the *stamina* expressed in its title. AUTHOR.

styles. Of this order there are *two* genera, viz. HELIOCARPUS ...and AGRIMONIA (*agrimony*).

Order 3. TRIGYNIA, comprehends such plants as have *three* styles. This order contains *three* genera, viz. RESEDA (*dyer's-weed*)...EUPHORBIA (*spurge*)...and VISMEA.

Order 4. TETRAGYNIA, contains such plants as have *four* styles, comprehending *two* genera, APONOGETON...CALLIGONUM.

Order 5. PENTAGYNIA, comprehends such plants as have *five* styles. This order contains *two* genera, viz. GLINUS...BLACKWELLIA.

Order 6. DODECAGYNIA, comprehends such plants as have *twelve* styles. This order contains but one genus, viz. SEMPERVIVUM (*houseleek*).



CHAP. XV.

OF THE TWELFTH CLASS, ICOSANDRIA*.

THIS class consists of such plants as bear *bisexual* flowers, of the following characters, viz. 1. A calyx monophyllous, and concave. 2. The corolla fastened by its claws to the inner side of the calyx. 3. The stamina twenty or more. As the number of stamina in this class, notwithstanding its title, is not limited, an attention must be had to the two first characters, to di-

* This class furnishes the fruits most in esteem.

stinguish the flowers from those of the next class, with which they might otherwise be confounded. The orders are five, viz.

Order 1. MONOGYNIA, comprehends such plants as have but *one* style. This order contains *twenty-one* genera, which fall under two sections. 1. *Calyx superior*, containing *thirteen* genera, viz. CACTUS (*melon thistle*) ... EUGENIA...PHILADELPHUS (*syringa*) ...PSIDIUM (*guava*) ... MYRTUS (*myrtle*) ...PUNICA (*pomegranate*) ...LEPTOSPERMUM... FABRICIA... METROSIDEROS... ROBINSONIA... CALYPTRANTHES...EUCALYPTUS...and FÆTIDIA. 2. *Calyx inferior*, including *eight* genera, viz. SONNERATIA...AMYGDALUS (*almond, peach, nectarine*)...PRUNUS (*plum, cherry, apricot, laurel*)...CHRYSOBALANUS (*cocoa plum*)...PLINIA...BANARA...ANTHERYLIUM...and SCOLOPIA.

Order 2. DIGYNIA, comprehending such plants as have *two* styles. Of this order there are *two* genera, viz. CRATÆGUS (*hawthorn*)...WALDSTEINIA.

Order 3. TRIGYNIA, comprehending such plants as have *three* styles. This order contains *two* genera, viz. SORBUS (*service*) ...and SESUVIUM.

Order 4. PENTAGYNIA, comprehending such plants as have *five* styles. This order contains *six* genera, viz. MESPILUS (*medlar*) ...PYRUS (*pear, apple, quince*)...TETRAGONIA...MESEMBRYANTHEMUM (*fig marygold*)...AIZOON...and SPIRÆA.

Order 5. POLYGYNIA, comprehending such plants as have *many* styles. This order contains *nine* genera, viz. ROSA (*rose*)...RUBUS (*bramble raspberry*) ...FRAGARIA (*strawberry*) ...POTENTILLA (*cinquefoil*) ...TORMENTILLA (*septfoil*) ... GEUM (*avens*) ... DRYAS (*mountain avens*) ...COMARUM (*marsh cinquefoil*)...and CALYCAN-THUS (*allspice*).

CHAP. XVI.

OF THE THIRTEENTH CLASS, POLYANDRIA*.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *many* stamina. The distinction between this class and the twelfth, may be known by having recourse to the characters of the twelfth class in the preceding chapter. The orders are six, containing *eighty-six* genera, viz.

Order 1. MONOGYNIA, comprehending such plants as have but *one* style. This order contains *forty-nine* genera, distinguished into, 1. *Such as have one petal*, of which there are *three* genera, viz. SWARTIA... MARCGRAVIA... and TERNSTRÆMIA. 2. *Three-petalled*, of which there is *one* genus, TRILIX. 3. *Four-petalled*, of which there are *nine* genera, viz. CAPPARIS (*capers*)... ACTÆA (*herb Christopher*)... CHELIDONIUM (*celandine*)... PAPAVER (*poppy*)... SARRACENA... MAMMEA... SPARMANNIA... CALOPHYLLUM... and GRIAS. 4. *Such as have five petals*, of which there are *twenty-one* genera, viz. OCHNA... TILIA (*lime tree*)... ELÆOCARPUS... CISTUS... CORCHORUS... LOASA... VALLEA... STERBECKIA... BONNETIA... LIGNOTIS... FREZIERA... MARILA... CISTUS... LEMNISCIA... MYRODENDRUM... SARRACENIA... AUBLETIA... OCHNA... ASCIUM... GREWIA... MUNTINGEA... and MICROCOS. 5. *Such as have six petals*, which contain *five* genera... ARGEMONE (*prickly poppy*)... LAGERSTRÆMIA... ALANGIUM... THEA (*tea tree*)... LECYTHIS. 6. *Such as have eight petals*, containing *one* genus only, SANGUINARIA (*puccoon*). 7.

* The fruits of this class are often poisonous; which makes it necessary to distinguish them from those of the last, which abounds with eatable fruits.

Such as have nine petals, *PODOPHYLLUM* (*duck's foot, or May-apple*). 8. Ten petals, one genus, *BIXA* (*anotta*). 9. Many petals, one genus, *NYMPHÆA* (*water lily*). 10. Without petals, seven genera, viz. *PROCKIA...MÆRUA...LUDIA...SLOANEA...RYANIA...LÆTIA...and SEGUIERIA.*

Order 2. *DIGYNIA*, comprehends such plants as have two styles. This order contains five genera, viz. *PÆONIA* (*peony*)...*CURATELLA...FOTHERGILLA...TRICHOCARPUS...LACIS.*

Order 3. *TRIGYNIA*, comprehends such plants as have three styles. This order contains three genera, viz. *DELPHINIUM* (*larkspur*)...*ACONITUM* (*wolf's-bane*)...and *HOMALIUM.*

Order 4. *TETRAGYNIA*, comprehends such plants as have four styles. This order contains five genera, viz. *TETRACERA...CARYOCAR...CIMICIFUGA...WINTERA...and WAHLBOMIA.*

Order 5. *PENTAGYNIA*, comprehends such plants as have five styles. This order contains four genera, viz. *AQUILEGIA* (*columbine*)...*NIGELLA* (*fennel flower*)...and *REAUMURIA.*

Order 6. *POLYGYNIA*, comprehends such plants as have many styles. This order contains twenty-one genera, viz. *DILLENIA...LIRIODENDRON* (*tulip tree*)...*MAGNOLIA...MICHELIA...UVARIA...ANNONA* (*custard apple*)...*ANEMONE...ATRAGENE...CLEMATIS* (*virgin's bower*)...*THALICTRUM* (*meadow rue*)...*ADONIS...ILLICIUM* (*uniseed tree*)...*RANUNCULUS* (*crowfoot*)...*TROLLIUS* (*globe flower*)...*ISOPYRUM...HELLEBORUS* (*hellebore*)...*CALTHA* (*marsh marygold*)...*HYDRASTIS* (*yellow root*)...*UNONA...XYLOPIA...NELUMBIVM.*

CHAP. XVII.

OF THE FOURTEENTH CLASS, DIDYNAMIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *four* stamina; *two* of which are longer than the rest. This circumstance would suffice to distinguish it from the fourth class, in which the four stamina are *equal*; however, as the flowers of this class have a particular structure, there are general characters which will nearly serve for the whole class; and these we will give at length.

Characters of the Class Didynamia.

CALYX—A perianthium, monophyllous, erect, tubulate, quinquefid, with segments for the most part unequal, and persisting.

COROLLA—Monopetalous and erect, the base of which contains the honey, and does the office of a nectarium. The upper lip straight: the lower spreading and trifid. The middle lacinia the broadest.

STAMINA—Four filaments, subulate, inserted in the tube of the corolla, and inclined towards the back thereof. The two inner and nearest the shortest. All of them parallel, and rarely exceeding the length of the corolla. The antheræ lodged under the upper lip of the corolla in pairs; in each of which respectively the two antheræ approach each other.

PISTILLUM—The germen commonly above the receptacle. The style single, filiform, bent in the same form as the fila-

ments, usually placed within them, a little exceeding them in length, and slightly curved towards the summit. The stigma for the most part emarginate.

PERICARPIUM—Either wanting (see the first order), or, if present, usually bilocular (see the second order).

SEEDS—If no pericarpium, four seeds, lodged within the hollow of the calyx, as in a capsule; but if there be a pericarpium, more numerous, and fastened to a receptacle placed in the middle of the pericarpium.

The flowers of this class are for the most part almost upright, but inclining a little at an acute angle from the stem, that the corolla may more easily cover the antheræ, and that the pollen may fall on the stigma, and not be injured with the rain. The essential character is in the four stamina; of which the two nearest are shorter, and all four close to each other, and transmitted with the single style of the pistillum, through a corolla that is unequal.

The orders of this class are two, comprehending *one hundred and twenty-five* genera, viz.

Order 1. GYMNOSPERMIA*, includes such plants as have *naked* seeds. This order has these farther characters, viz. the seeds *four* (excepting PHRYMA, which is *monospermous*); and the stigma *bipartite*, and *acute*, with the lower lacinia *reflexed*. It contains *thirty-nine* genera, distinguished into, 1. *Such as have the calyx quinquesid, and nearly equal*, of which there are *twenty-three* genera, viz. AJUGA (*bugle*)... TEUCRIUM (*germander*)... SATUREIA (*savory*) ... HYSSOPUS (*hyssop*)... NEPETA (*cat mint*)... LAVANDULA (*lavender*)... BETONICA (*betony*)... SIDERITIS (*ironwort*)... MENTHA (*mint*)... GLECHOMA (*ground ivy*)... PERILLA... LAMIUM (*archangel*)

* The plants of this order are scented, and are accounted cephalic and resolvent. The virtue is in the leaves. They are the labiati (lipped plants) of Tournefort, and verticillati (plants that flower at the joints, in whirls of Ray's Hist. Plant. 508.

... GALEOPSIS (*hemp nettle*) ... STACHYS (*wound-wort*) ... BALLOTA (*black horehound*) ... MARRUBIUM (*white horehound*) ... LEONURUS (*mother-wort*) ... PHLOMIS... MOLUCCELLA (*Molucca balm*)... ELZ-HOLTZIA... BYSTROPOGON... and HYPTIS. 2. *Such as have the calyx bilabiate, divided into two lips*; of which there are sixteen genera, viz. CLINOPODIUM (*basil*)... ORIGANUM (*marjoram*)... THYMUS (*thyme*)... MELISSA (*balm*)... DRACOCEPHALON (*aragon's head*)... MELITTIS (*bastard balm*)... OCYMUUM (*basil*)... SCUTELLARIA (*scull-cap*) ... PRUNELLA (*self-heal*)... CLEONIA... PRASIMUM... PHRYMA... PLECTRANTHUS... THYMBRA, and SELAGO.

Order 2. ANGIOSPERMIA*, comprehends such plants as have the seeds in a *pericarpium*, which circumstance is constant, and distinguishes this order from the last in every form. To this character may be added that of a stigma, commonly *obtusè*. This order contains *eighty-six* genera, distinguished into, 1. *Such as have a calyx undivided*, which contains *two* genera, ÆGINETIA... TANÆCIUM. 2. *Calyxes bifid*, which contains *eight* genera, OBO-LARIA... OROBANCHE (*broom-rape*)... HEBENSTREITIA... TORENIA ... CASTILLEIA... ACANTHUS... PREMNA... and CRÉSCENTIA (*calabash tree*). 3. *Calyxes quadrifid*, which contains *eleven* genera... LIPPIA... LATHRÆA (*tooth-wort*)... BARTSIA... EUPHRASIA (*eye-bright*) ... RHINANTHUS (*yellow rattle*) ... MELAMPYRUM (*cow wheat*) ... SCHWALBEA... BARLERIA... LÆSELIA... GNELINA... and LANTANA. 4. *Calyxes five-cleft*, which contains *sixty-three* genera, which subdivides into, 1. *Capsule one-celled*, which contains *twelve* genera, AVECENNIA... TOZZIA... PHAYLOPSIS... LIMOSELLA (*mudwort*)... BROWÆLIA... BRUNFELSIA... HOLMSKIOLDIA... LINDERNIA... CONOBEA... COLUMNÆA... VANDELLIA... RUSSELLIA. 2. *Capsule two-celled*, which contains *twenty-eight* genera, ALECTRA... GESNERIA... CYRILLA... SCROPHULARIA (*fig-wort*) ... STEMODIA ... ACHIMENES... CELSIA... HEMIMERIS... SIETHORPIA... CAPRARIA... DIGITALIS (*fox-glove*)... BIGNONIA... INCARVILLEA... RUELIIA... BUCHNERA... ERIUS ... PETREA... MANULEA... ANTIRRHINUM (*snap-dragon*)... ANARRHI-

* These are the *personati* (*personate flowers*) of Tournefort.

NUM...GERARDIA...PEDICULARIS (*louse-wort*)...MIMULUS (*monkey flower*)...DODARTIA...CHELONE...PENTSTEMON...SESAMUM (*oily grain*)...GLOXINIA. 3. *Capsule four-celled*, which contains two genera, TOURETIA...MARTYNIA. 4. *Capsules two*, one genus, MAURANDIA. 5. *A silique*; one genus, MILLINGTONIA. 6. *A nut*; two genera, TORTULA...PEDALUM. 7. *A berry*; five genera, LINNEA...CORNUTIA...OVIEDA...AMASONIA...BESLERIA. 8. *A drupe*; eight genera, BONTIA...SPIELMANNIA...VITEX...MYOPORUM...CYTHAREXYLON...VOLKAMERIA...CLERODENDRON...DURANTA. 5. *Calyx many-cleft*, which comprehends four genera, HYOBANCHE...LEPIDAGATHIS...CYMBARIA...THUNBERGIA. 6. *Many-petalled*, which has only one genus, MELIANTHUS (*honey flower*).

CHAP. -XVIII.

OF THE FIFTEENTH CLASS, TETRADYNAMIA*.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *six* stamina, *two* of which are shorter than the rest, by which last circumstance it may be distinguished from the sixth class, whose flowers have *six equal* stamina. The flowers

* These are the *cruciformes* (*cross-shaped flowers*) of *Tournefort*, and the *siliculosæ*, and the *siliquosæ* (plants that have a *silicula* and *siliqua*) of *Ray's Hist. Plant.* 777. This class is truly *natural*, and has been assumed as such by all systematists. LINNÆUS thinks he has given no other, unless it be CLEOME. The distinction into *siliculose*, and *siliquose*, is admitted by all, as a good distinction. The plants are held to be antiscorbutic and diuretic. The taste in most is watery; mixed with a sharpness. They commonly lose their quality when dried.

of this class are of a particular structure, answering to the following characters :

Characters of the Class Tetradynamia.

CALYX—A *perianthium*, tetraphyllous and oblong ; the leaves of which are ovato-oblong, concave, obtuse, conniving, gibbous downwards at the base, the opposite ones equal and deciduous. The calyx in these flowers is a nectarium*, which is the reason of the base being gibbous.

COROLLA—called cruciform, that has four equal and opposite petals. The *claws* plano-subulate, erect, and somewhat longer than the calyx. The *limb* plane. The *laminae* widening outwards, obtuse, the sides hardly touching one another. The insertion of the petals is in the same circle with the stamina.

STAMINA—The *filaments* six, and subulate ; of which two that are opposite, are of the length of the calyx ; the other four somewhat longer, but not so long as the corolla. The *antheræ* oblong, acuminate, thicker at the base, erect, and with their tops leaning outwards. There is a *nectariferous glandule*, which in the different genera has various appearances : it is seated close to the stamina, and particularly to the two shorter ones, to whose base it is fastened ; and these have a light curvature to prevent their pressing upon it, whereby those filaments become shorter than the rest.

PISTILLUM—The *germen* above the receptacle increasing daily in height. The *style* either of the length of the longer stamina, or wanting. The *stigma* obtuse.

PERICARPIUM—A *siliqua* of two valves, often bilocular, opening from the base to the top. The *dissepiment* projecting at the

* It should be, contains the nectarium, which is explained afterwards to be a gland. EDITOR.

top, beyond the valves, the prominent part thereof having before served as a style.

SEEDS—Roundish, inclining downwards, alternately plunged lengthwise into the dissepiment. The *receptacle* linear, surrounding the dissepiment, and immersed in the sutures of the pericarpium. The orders are two, containing *thirty-four* genera, viz.

Order 1. SILICULOSA, comprehending those plants whose pericarpium is a silicula*. This order contains nineteen genera, subdivided into, 1. *Silicle entire*; that is, *not emarginate at the top*, which contains *nine* genera, viz. MYAGRUM (*gold of pleasure*)... VELLA (*cross-rocket*)...SUBULARIA (*awl-wort*)...DRABA (*whitlow grass*) ...LUNARIA (*honesty*)...CAKILE...PUGIONIUM...BUNIAS (*sea rocket*)...and CRAMBE (*colewort, or sea-kale*). 2. *Silicle emarginate at the end*, which contains *ten* genera...IBERIS (*candy-tuft*)...ALYSSUM (*madwort*)...CLYPEOLA (*treacle mustard*)...PELTARIA...COCHLIARIA (*scurvy-grass*)...LEPIDIUM (*pepper-wort*)...THLASPI (*mithridate mustard*)...ISATIS (*woad*)...BUSCUTELLA...and ANASTATICA (*rose of Jericho*).

Order 2. SILIQUOSA, comprehends those plants whose pericarpium is a siliqua†. This order contains fifteen genera, falling under two divisions, 1. *Calyx closed, with the leaves converging longitudinally*, which contains *ten* genera, viz. DENTARIA (*tooth-wort, or coral-wort*)...ERYSIMUM (*hedge mustard*)...CHEIRANTHUS (*wall-flower, and stock gilliflower*)...HESPERIS (*rocket*)...ARABIS (*wall-cress, and rock-cress*)...TURRITIS (*tower mustard*)...BRASSICA (*cabbage, rape, or cole-seed, turnip*)...RAPHANUS (*radish*)...RICOTIA...and CORDYLOCARPUS. 2. *Calyx gaping, with the leaves distant above*, contains five genera, CLEOME...CARDAMINE (*ladies' smock*)...SINAPIS (*mustard*)...SISYMERIUM (*water-cress, water-rocket*)...and HELIOPHILA.

* See the account of this order in Chap. III.

† See Chap. III.

CHAP. XIX.

OF THE SIXTEENTH CLASS, MONADELPHIA*.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *one* set of *united* stamina. This class consists of eight orders. The characters of the flowers are as follow:

Characters of the Class Monadelphia.

CALYX—A *perianthium* always present, persisting, and in most genera double.

COROLLA—Pentapetalous, the *petals* heart-shaped; the sides of which lap each one over the next, contrary to the motion of the sun.

STAMINA—The *filaments* united below, but distinct upwards if there be more than one†. The exterior ones shorter than the interior. The *antheræ* incumbent.

PISTILLUM—The *receptacle* of the fructification prominent in the centre of the flower. The *germen* erect, surrounding the top of the receptacle in a jointed ring. The *styles* are all united below in one substance with the receptacle, but divided above into as many threads as there are germens. The *stigma* spreading and thin.

* In this class the calyx is of great moment for distinguishing the genera, and fixes the limits with certainty. They were formerly distinguished by the fruit; which not being found sufficient, recourse was had to the leaves of the plant. The plants of this class are esteemed to be emollient and mucilaginous. AUTHOR.

† The *melochia* has five antheræ, but it does not appear that there are any distinct filaments. See its character in the Genera Plantarum. AUTHOR.

PERICARPIUM—A *capsule* divided into as many loculements as there are pistilla. Its figure various in the different genera.

SEEDS—Kidney-shaped.

The *corolla* in this class has been called *monopetalous*; but as the petals are all distinct at the base, it is to be styled more properly *pentapetalous*, notwithstanding the petals cohere by the union of the stamina. The orders are nine, containing *sixty* genera, viz.

Order 1. TRIANDRIA, comprehending such plants as have *three* stamina. This order contains five genera, viz. APHYTEIA... GALAXIA... SISYRINCHIUM... FERRARIA... TAMARINDUS.

Order 2. PENTANDRIA, comprehending such plants as have *five* stamina. This order contains *nine* genera, viz. WALTHERIA... LERCHEA... HERMANNIA... MELOCHIA... SYMPHONIA... ERODIUM (*stork's-bill*)... OZOPHYLLUM... OCHROMA... PASSIFLORA... HERMANNIA... and MELOCHIA*.

Order 3. HEPTANDRIA, comprehends such plants as have *seven* stamina, and includes *one* genus, PELARGONIUM.

Order 4. OCTANDRIA, comprehends such plants as have *eight* stamina. Of this order there are *two* genera, viz. AITONIA... PISTTIA.

Order 5. ENNEANDRIA, comprehends such plants as have *nine* stamina. Of this order there is but *one* genus, viz. DRYANDRA.

Order 6. DECANDRIA, comprehending such plants as have *ten* stamina. This order contains *three* genera, viz. CONARUS... GERANIUM†... HUGONIA... SENBÆA... and CRINODENDRUM.

* The reader will observe, that several of these genera were, by Linnæus, considered to be of the class Gynandria.

† The species of this genus varies singularly in the number of stamina and other circumstances, viz. from 1 to 22 they have seven fertile stamina, the leaves alternate, and many flowers on a peduncle; (these now constitute a new genus, called Pelargo-

Order 7. ENDECANDRIA, comprehending such plants as have *eleven* stamina. Of this order there is only *one* genus, viz. BROWNEA.

Order 8. DODECANDRIA, comprehending such plants as have *twelve* stamina. This order contains *nine* genera, viz. PENTAPETES...MONSONIA...HELICTERES (*screw-tree*)...PLAGIANTHUS...ACTA...PTEROSPERMUM...CIENFUEGIA...DOMBEYA, and ASSONIA.

Order 9. POLYANDRIA, comprehending such plants have *many* stamina. This order contains *twenty-eight* genera, viz. BOMBAX (*silk-cotton*) ... SIDA... ADANSONIA ... ALTHÆA (*marsh-mallow*) ... MALVA (*mallow*) ... LAVATERA... MALOPE... URENA... GOSSYPIUM (*cotton*)...HIBISCUS...STUARTIA...CAMELLIA (*Japan rose*)... MORISONIA... MESUA... MALACHRA...GORDONIA...GUSTAVIA...CAROLINEA...BARRINGTONIA...CROSSOSTYLIS...MYRODIA...POURRETIA...PALAVIA ... LAGUNÆA ... RUIZIA... PAVONIA ...ACHANIA...and KITAIBELIA.

nium): from 23 to 35 they have seven fertile stamina, and the leaves growing opposite; from 36 to 45 five fertile stamina, the calyx five leaves, and the fruit declined; from 46 to 58 ten fertile stamina, and two flowers on a peduncle; from 59 to 68 ten fertile stamina, two flowers on a peduncle, and the plants annual; from 69 to 82 ten fertile stamina, and one flower on a peduncle.

CHAP. XX.

OF THE SEVENTEENTH CLASS, DIADELPHIA*.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *two* sets of *united* stamina†. The characters of the fructification are as follow :

Characters of the Class Diadelphia.

CALYX—A *perianthium* monophyllous, campanulate, and withering. The base gibbous, the lower part thereof fastened to the peduncle, the upper obtuse and melliferous. The brim quinque-dentate, acute, erect, oblique, unequal. The lowest odd denticle longer than the rest; the upper pair shorter and farther asunder. The bottom of the cavity moist with a melleous liquor, including the receptacle.

* The plants of the class *Diadelphia*, are the *papilionaceous*, *butterfly-shaped* plants, of *Tournefort*; *irregular tetrapetalous* of *Rivinus*; and *leguminous* of *Ray's* *Hist. Plant.* 883. Of all the classes, this is the most natural, and has its flowers of the most singular structure. The calyx, though hitherto little attended to, is of great moment for fixing the genera. The legumen was held of consequence by other systematists; but by *Linnaeus* it is made of less account. The leaves of these plants are food for cattle, and the seeds also for quadrupeds of the same kind; the latter are accounted flatulent.

† This circumstance, implied in the title, does not hold through the class, the plants given under the first distinction of the third order having *monadelphious* stamina; the class is therefore not so properly to be fixed from its title, as by the *papilionaceous corolla*, and other characters of the fructification. It may be observed likewise, that in the diadelphious flowers of this class, one of the two stamina is not a set of *united* filaments, as in the other, but only a *single* stamen, detached from the united set. See the characters of the fructification.

COROLLA—Termed papilionaceous, unequal; the petals expressed by distinct names, viz.

Vexillum, the *standard*; a petal covering the rest, incumbent, greater, plano-horizontal, inserted by its claw in the upper margin of the receptacle, approaching to a circular figure when it leaves the calyx, and nearly entire; along it, and especially towards its extremity, runs a line or ridge, that rises up, as if the lower part of the petal had been compressed; the part of the petal next to the base approaching to a semicylindric figure, embraces the parts that lie under it. The disk of the petal is depressed on each side, but the sides of it nearest the margin are reflexed upwards. Where the halved tube ends, and the halved limb begins to unfold itself, are two concave impressions prominent underneath, and compressing the wings that lie under them.

Alæ, the *wings*, two equal petals, one at each side of the flower, placed under the vexillum; incumbent with their margins, parallel, roundish, or oblong, broader upwards, the upper margin straighter, the lower spreading more into a roundness; the base of each wing bifid, the lower division stretching out into a claw, inserted in the side of the receptacle, and about the length of the calyx; the upper shorter and inflexed.

Carina, the *keel*, the lowest petal often bipartite, placed under the vexillum and between the alæ, boat-shaped, concave, compressed on the sides, set like a vessel afloat, mutilate at the base, the lower part of which runs into a claw, of the length of the calyx, and inserted in the receptacle, but the upper and side laciniae are interwoven with that part of the alæ that is of the same shape. The form of the sides of the carina, is much like that of the alæ; and so also is their situation, except that they are lower, and stand within them. The line that forms the *carina*, or *keel*, in this petal, runs straight as far as the middle, and then rises gradually in the segment of a circle, but the mar-

ginal line runs straight to the extremity, where meeting the carinal, they terminate obtusely.

STAMINA—are what is called *diadelphia*. The filaments two, of different forms, viz. a lower one that involves the pistillum, and an upper one incumbent on it. The former of these, from the middle downwards, is cylindraceous, membranaceous, and split lengthwise on its upper side; but the upper half terminates in nine subulate* parts, that are of the same length with, and follow the flexure of, the carina of the corolla, and of which the intermediate or lower radii† are longer by alternate pairs. The upper filament is subulato-setose‡, covering the splitting of the former cylindraceous filament, incumbent on it, answering to it in situation, simple and gradually shorter; its base is detached from the rest, and prepares an outlet for the honey on each side. The antheræ, reckoned all together, are ten, one on the upper filament, and nine on the lower, each of the radii being furnished with a single one; they are small, all of one size, and terminate the radii.

PISTILLUM—Single, growing out of the receptacle, within the calyx. The *germen* oblong, roundish, lightly compressed, straight, of the length of the cylinder of the lower filament which involves it. The *style* subulate, filiform, ascending, having the same length and position as the radii of the filament among which it is placed, and withering. The *stigma* downy, of the length of the style from the part turned upwards, and placed immediately under the antheræ.

PERICARPIUM—A *legumen*, oblong, compressed, obtuse, bivalved, with a longitudinal suture both above and below; each suture straight, though the upper one falls near the base, and the lower one rises near the top. The legumen opens at the upper suture.

* Awl-shaped. AUTHOR.

† Rays, meaning the divisions of the filaments. AUTHOR.

‡ Awl-shaped, and like a bristle. AUTHOR.

SEEDS—A few, roundish, smooth, fleshy, pendulous, marked with an embryo that is a little prominent towards the point of insertion. When the ova* are hatched, the cotyledons† preserve the form of the halved seed.

RECEPTACLE—The proper receptacles of the seeds are very small; very short, thinner towards the base, obtuse at the disk that fastens them, oblong, inserted longitudinally in the upper suture of the legumen only, but placed alternate; so that when the valvulæ have been parted, the seeds adhere alternately to each of the valves.

The ordinary situation of the flowers is obliquely pendulous; that is, at an acute angle from the perpendicular. The orders are *four*, containing *seventy-eight* genera, viz.

Order 1. PENTANDRIA, comprehending such plants as have *five* stamina. Of this order there is only one genus, viz. MONNERIA.

Order 2. HEXANDRIA, comprehending such plants as have *six* stamina. This order contains *two* genera, viz. FUMARIA (*fumetory*)...and SARACA.

Order 3. OCTANDRIA, comprehending such plants as have *eight* stamina. This order contains *three* genera, viz. POLYGALA (*milkwort*)...SECURIDACA...and BREDEMEYERA.

Order 4. DECANDRIA, comprehending such plants as have *ten* stamina. This order contains *fifty* genera, distinguished into, 1. *Such as have monadelphous‡ filaments*; of which there are *twen-*

* Eggs, meaning the seeds themselves, which answer to the eggs of animals, and are as it were hatched when the corculum, or first principle of the new plant begins to strike root and vegetate. See Part I. Chap. VII. AUTHOR.

† Side leaves of the seed. See Part I. Chap. VII. The two seed-leaves, which first appear above ground, are these very cotyledons, which are brought up with the plant, after the corculum has struck; and it is these seed-leaves that are here spoken of. AUTHOR.

‡ One set, or brotherhood. AUTHOR.

ty-nine genera, viz. NISSOLIA...DIPTERIX...PTEROCARPUS...AMERIMNUM...AMORPHA (*bastard indigo*)...TRIGONIA...ERYTHRINA (*coral-tree*)...RUDOLPHIA...BUTEA...ABRUS...LEBECKIA...SPARTIUM (*broom*)...GENISTA (*broom*)...RAFANIA...LUPINUS (*lupine*)...TERAMNUS...ANTHYLLIS (*kidney-vetch*)...PISCIDIA (*Jamaica dog-wood*)...WIBORGIA...SARCOPHYLLUM...BORBONIA...ÆDMANNIA...ULEX (*furze, whins, or gorse*)...ARACHIS (*earth-nut*)...ASPALATHUS...ONONIS (*rest-harrow*)...BOSSIEA...CROTALARIA...and PLATYLOBIUM.

2. *Stigma pubescent, stamens diadelphous*, of which there are seven genera; COLUTEA (*bladder-senna*)...PHASEOLUS (*kidney-bean*)...DOLICHOS...OROBUS (*bitter vetch*)...PISUM (*pea*)...LATHYRUS (*everlasting pea*)...and VICIA (*vetch, or tare*).

3. *Legume subbilocular, stamens diadelphous*, of which there are three genera; ASTRAGALUS (*milk-vetch*)...BISERRULA...and PHACA (*bastard vetch*).

4. *Legumes one or two-seeded, stamens diadelphous*, of which there are ten genera; DALBERGIA...DALEA...PSORALEA...TRIFOLIUM (*trefoil*)...DORYCNIUM...HALLIA...STYLOSANTHES...CYLISTA...GLYCYRRHIZA (*liquorice*)...and DIMORPHA.

5. *Legume subarticulate, stamens diadelphous*, of which there are eight genera; ÆSCHYNOMENE...MULLERA...HEDYSARUM (*sainfoin*)...SMITHIA...CORONILLA...ORNITHOPUS (*bird's-foot*)...SCORPIURUS (*caterpillar*)...and HIPPOCREPIS (*horse-shoe vetch*).

6. *Legume one-celled, many-seeded, diadelphous*, of which there are fifteen genera; TRIGONELLA (*fenugreek*)...GLYCINE...CLITORIA...ROBINIA...INDIGOFERA (*indigo*)...CICER (*chick pea*)...ERVUM (*lentil*)...LIPARIA...ACHYRONIA...CYTISUS...DIPHYSA...GALEGA (*goat's rue*)...LOTUS (*bird's-foot trefoil*)...MEDICAGO (*medick lucern*)...and GEOFFROYA.

CHAP. XXI.

OF THE EIGHTEENTH CLASS, POLYADELPHIA.

THIS class consists of such plants as bear *bisexual* flowers, furnished with *many* sets of *united* stamina; the flowers have no particular character farther than is expressed in the title. The orders are four, including *eleven* genera, viz.

Order 1. DECANDRIA, comprehending such plants as have *ten* stamina in each set. Of this order there is only *one* genus, viz. THEOBROMA (*chocolate*).

Order 2. DODECANDRIA, comprehending such plants as have *twelve* stamina in each set. Of this order there are *two* genera, viz. BUBROMA...ABROMA.

Order 3. ICOSANDRIA, comprehending such plants as have *twenty* stamina in each set. Of this order there are *two* genera, viz. CITRUS (*orange and lemon*)...and MELALEUCA.

Order 4. POLYANDRIA, comprehending such plants as have *many* stamina in each set. This order contains *six* genera, viz. HYPERICUM...ASCYRUM...SYMPLOCOS...DURIO...GLABRARIA...and LUHEA.

CHAP. XXII.

OF THE NINETEENTH CLASS, SYNGENESIA*.

THIS class consists of such plants as bear *compound* flowers. We have already paved the way for understanding this class, by the account given of *compound* flowers, in Part I. Chap. XIX. and the explanation of the titles of the class, and its orders, in Chap. II. and III. What is farther necessary here, is to give the characters of the flowers. *Compound* flowers admit of a double description, viz. 1. of the whole flower in its aggregate state, which is termed the *flosculose flower*; and, 2. of the *flosculi, florets*, of which it is composed. We shall begin with the first, which concerns only the *calyx* and *receptacle*, those being the only parts that are in common.

Characters of the Flosculous Flower.

CALYX—The *common calyx* is a perianthium, which contains the florets and the receptacle. It is either *simple, augmented, or imbricated* †. It contracts when the flowers are fallen, but expands and turns back when the seeds are ripe.

RECEPTACLE—The *common receptacle* of the fructification receives many sessile florets on its disk, which is either *concave... plane...convex...pyramidal...or globosc.* The surface of the disk

* This class of *compound* flowers is a natural one, if we except the last order; which, upon the systematic principles assumed, could not be refused an admission into it. Its plants are commonly bitter and stomachic. AUTHOR.

It has, however, been abolished by Dr. Smith, president of the Linnæan Society, with the approbation of Professor Martyn, and this is admitted by the generality of botanists. EDITOR.

† See these terms explained in Part I. Chap. XI.

is either *naked*, without any other inequality than that of being lightly dotted;...*villose*, covered with upright hairs;...or *paleaceous*, covered with *paleæ*, *chaffs*, or *straws*, that are linear, subulate, compressed, and erect, and serve to part the florets.

Characters of the Florets.*

CALYX—A small perianthium, often quinquepartite, seated on the germen, persisting, and becoming the crown of the seed.

COROLLA—Monopetalous, with a long and very narrow tube. It is seated on the germen; and is either *tubulate*, with the limb campanulate and quinquefid, and the laciniaë spreading and turning back;...*ligulate*, with the limb linear, plane, turned outwards, and the top whole;...*tridentate*, or *quinquedentate*, or *wanting*, having no limb, and often no tube.

STAMINA—The *filaments* five, capillary very short, inserted in the neck of the corollulaë. The *antheræ* five, linear erect; and by the union of their sides forming a cylinder, that is tubulate, quinquedentate, and of the length of the limb.

PISTILLUM—The *germen* oblong, placed under the receptacle of the flower; the *style* filiform, erect, of the length of the stamina, and perforating the cylinder of the antheræ; the *stigma* bipartite, the laciniaë revolute, and spreading asunder.

PERICARPIUM—No true one, though in some there is a coriaceous†

* The character here given is of a *bisexual* floret; but the florets may also be either *male*, *female*, or *neutral*, as the orders show; it may not be improper, therefore, to observe, in general, upon these classic characters, which our author has drawn with such minute exactness, that they should be understood as collected only from the circumstances that most frequently occur in the class; and liable to variation, not in particular genera only, but even through the whole orders of the class in some cases. AUTHOR.

† Leathery. EDITOR.

SEED—A single one, oblong, often tetragonous, but commonly narrower at the base. It is either crowned,...or with the *crown wanting*. The crown is of two kinds, either a *pappus*,...or a *perianthium*:...if a *pappus*, it is either sessile,...or placed on a *stipes*; and consists of many *radii*, that are placed in a *round*, and are either *simple*,...*radiate*,...or *ramose*: when the crown is a *perianthium*, it is such as is described above under that head.

The essence of a *flosculose* flower consists in having the antheræ united in a cylinder, and a single seed below the receptacle of the floret*. The orders of this class are six, containing *seventy-three* genera, viz.

Order 1. POLYGAMIA ÆQUALIS, comprehends such plants as have *compound* flowers, of which the florets are all bisexual. This order contains *thirty* genera, distinguished into 1. SEMIFLOSCULOSA, with all the corollets *ligulate*, which contains *thirty* genera; SCOLYMUS (*golden thistle*)...CICHORIUM (*succory, endive*)...CATANACHE...SERIOLA...HYPOCHÆRIS (*cat's-ear*)...GEROPOGON (*old man's beard*)...ROTHIA...ANDRYALA...TRIPTILION...TRAGOPOGON (*gout's-beard*)...ARNOPOGON...HELMINTIA...PICRIS (*ox-tongue*)...ASPARGIA...SCORZONERA... (*viper's grass*)...LEONTODON (*dandelion*)...CREPIS (*hawk's beard*)...CHONDRILLA (*gum-succory*)...PRENANTHES...LACTUCA (*lettuce*)...HIERACIUM (*hawk-weed*)...SONCHUS (*sow-thistle*)...ZACINTHA...LAPSANA (*nipple-wort*)...RHAGADIOLUS...KRIGIA...HYOSERIS (*swine's succory*)...HEDYPTIS (*hawk-bit*)...THRINZIA...TOLPIS. 2. CAPITATI, flowers in a head, all the corollets *tubular*; spreading at the tip, contains *nineteen* genera; ATRACTYLIS...ACARNA...SERRATULA (*saw-wort*)...CARTHAMUS (*bastard saffron*)...CARLINA (*carline thistle*)...ARCTIUM (*burdock*)...

* That the essence of a *flosculose*, or *compound* flower, does not consist either in the common calyx or receptacle, *Linnaeus* argues from hence; that the common calyx is wanting in *echinops*, and the common receptacle in *milleria*, though both those genera belong to this class; and that, on the other hand, the common calyx is found in *scabiosa*, and the common receptacle in *dipsacus*, both which plants belong to the class *Tetrandria*, though they have, with the *gomphrena* and others, been falsely ranged with the compound flowers. AUTHOR.

PTERONIA... STOBÆA... LACHNOSPERMUM... BARNADESIA... CYNARA
(artichoke)... JOHANNIA... CNICUS... CARDUUS *(thistle)*... ONOSERIS...
 STOKESIA... LIATRIS... VERNONIA... ONOPORDON *(cotton-thistle)*. 3.
 DISCOIDEI, *all the corollets tubular, erect-parallel, flattish at the tip,*
dense, which contains twenty-four genera; STEHELINA... HAYNEA
 ... CALEA *(halbert-weed)*... BIDENS *(bur-marygold)*... SPILANTHES...
 ANTHANASIA... SANTOLINA *(lavender-cotton)*... CÆSULIA... TARCHO-
 NANTHUS *(African fleabane)*... KUHNIA... EUPATORIUM *(hemp agri-*
mony)... CHRYSOCOMA *(golden locks)*... MILKANIA... KLEINIA... CA-
 CALIA... LAVENIA... AGERATUM... STEVIA... HYMENOPAPPUS... CE-
 PHALOPHORA... PENTZIA... ETHULIA... PIQUERIA... BALSAMITA.

Order 2. POLYGAMIA SUPERFLUA, comprehends such plants as
 have the florets of the *disk* bisexual, and those of the *radius* female.
 This order contains *sixty-one* genera, distinguished by, 1. DISCOIDEI,
corollets of the ray obscure or none, which contains ten genera; ARTE-
 MISIA *(south-ernwood, wormwood, mugwort)*... CARPESIMUM... TANA-
 CETUM *(tansy)*... COTULA... BACCHARIS... CONYZA *(fleabane)*... GNA-
 PHALIUM *(cudweed)*... ELICHRYSUM... XERANTHEMUM... ANACYCLUS.
 2. SEMIFLOSCULI, SUBBILABIATI, *subbilabiate*, which includes *two*
 genera; DENEKIA... PERDICIUM. 3. RADIATI, *corollets of the disk*
floscular, of the ray ligulate, which contains *forty-nine* genera;
 MADIA... BELLIS *(daisy)*... MATRICARIA... LIDBECKIA... CHRYSAN-
 THEMUM *(ox-eye daisy, corn-marygold)*... PYRETHRUM *(feverfew)*...
 COLUMELLIA... DORONICUM *(leopard's bane)*... ARNICA... INULA *(ele-*
campane, fleabane)... ERIGERON *(fleabane)*... SOLIDAGO *(golden rod)*
 ... CINERARIA *(fleurwort)*... SENECIO *(groundel, ragwort)*... TUSSILAGO
 ... ASTER *(starwort)*... BOEBERA... MUTISIA... BELLIUM... ACTINEA
 ... TAGETES... HELENIUM ... PECTIS... SCHKUHRIA... HETEROSPER-
 MUM... BOLTONIA ... LEYSERA... SEIGESBECKIA... ECLIPTRA... AN-
 THEMIS *(chamomile)*... ACHILLEA *(milfoil)*... TETRAGONOTHECA...
 XIMENESIA... PHAETHUSA... GEORGINA... RELHANIA... PASCALIA...
 BUPHTHALMUM *(ox-eye)*... RHANTERIUM... SANVITALIA... AMELLUS
 ... TRIDAX... ROSENIA... VERBESINA... SCHLECHTENDALIA... GALIN-
 SOGEA... ZINNIA... BALBISIA... and STARKEA.

Order 3. POLYGAMIA FRUSTRANEA, comprehends such plants as have the florets of the *disk bisexual*, and those of the *radius neuter*. This order contains *sixteen* genera, all *radiate*, viz. HELIANTHUS (*sun-flower*)...RUDBECKIA...COREOPSIS...GORTERIA...OSMITES...ZÆGEA...CENTAUREA*...SCLEROCARPUS...DIDELTA...MUSSINIA...LAPEYROUSIA...BEREKHEYA...TITHONIA...GALARDIA...COSMEA...and PALLUSIA.

Order 4. POLYGAMIA NECESSARIA, comprehends such plants as have flowers of the *disk male*, and those of the *radius female*. This order contains *twenty-two* genera, most of which are *radiate*, viz. MILLERIA...SILPHIUM...CHRYSOGONUM...MELAMPIDIUM...CALENDULA (*marygold*)...ARCTOTIS...OSTEOSPERMUM...OTHONNA (*African ragwort*)...POLYMNIA...ERIOCEPHALUS...FILAGO (*cudweed*)...MICROPUS...BALTIMORA...HIPPIA...PSIADIA...UNXIA...IVA...WEDELIA...ACICARPHA...PARTHENIUM...ARCTOTHECA...TRIXIS.

Order 5. POLYGAMIA SEGREGATA. This order comprehends such plants as have *many* partial florets contained in the common calyx, which separate and surround the floscula. This order contains *sixteen* genera; TETRANTHUS...ROLANDRA...NAUENBURGIA...CALYCERA...NOCCÆA...BOOPIS...STÆBEA...ÆDERA...BROTERA...ECHINOPS (*globe-thistle*)...ELEPHANTOPUS (*elephant's foot*)...NASSAUVIA...JUNGLIA...GUNDELIA...SPHÆRANTHUS...and CRASPEDIA.

Order 6. MONOGAMIA, comprehends such plants as have *simple* flowers. This order contains seven genera, viz. STRUMPFIA...SERIPHIMUM...CORYMBIUM...JASIONE...LOBELIA (*cardinal's flower*)...VIOLA (*violet*)...and IMPATIENS (*balsam*)†.

* The corollulæ of the *centaurea* are all *tubulose*, but those of the *radius* differ from those of the *disk*, which brings it within the definition of a *radiate* flower; however, *Linnæus*, in his description of the *centaurea*, in the *Genera Plantarum*, has not called the corolla *radiate*, but *tubulosa difformis*, *tubulose of different forms*.
AUTHOR.

† The plants of this class are removed, by general consent, into the class *PENTANDRIA*, to which they properly belong. Here they disfigure a class that has the strong recommendation of being altogether natural. We have preserved them, however, here, to illustrate the *Sexual System of Linnæus*. EDITOR.

CHAP. XXIII.

OF THE TWENTIETH CLASS, GYNANDRIA*.

THIS class consists of such plants as have the *stamina* growing either upon the *style* itself, or upon a *receptacle* that stretches out into the form of a style, and supports both the *stamina* and the *pistillum*. The orders are nine, viz.

Order 1. DIANDRIA†, comprehending such plants as have *two* *stamina*. The flowers of this order have a most singular structure, answering to the following description.

Characters of the Order Diandria, of the Class Gynandria.

The *germen* is always contort‡; the *petals* are five; of which the two inner ones usually approach, and form a *galea*||; the lower lip of which becomes a *nectarium*, and serves also for a *pistillum* and sixth petal. The *style* grows to the inner margin of the *nectarium*, in such a manner as to be, with its stigma, scarce either of them distinguishable. The *filaments* are always two, supporting as many *antheræ*; which are narrower downwards; naked, or without tunic, and divisible, like the pulp of a *citrus*. These last are covered by little cells, that are open underneath, and grow to the inner margin itself of the *nectarium*. The *fruit*

* All the flowers of this class have a monstrous appearance, owing to the uncommon situation of the parts of fructification. AUTHOR.

† This order is a *natural* one, the genera differing only in respect of the *nectarium*. This part *Linneus* considers as a mark of distinction for these genera, far preferable to the root, though not received as such by former botanists. AUTHOR.

‡ Twisted like a screw. EDITOR.

|| Helmet. EDITOR.

is a *capsule*, that is unilocular, trivalved, and splits in the angles under the carinate* ribs. The seeds are scobiform†, numerous, affixed to a linear receptacle in each valvule‡.

Order 1. DIANDRIA, comprehending such plants as have two stamina. This order contains *eleven* genera, viz. ORCHIS...SATYRIUM...OPHRYS...SERAPIAS...LIMODORUM...ARETHUSA...CYPRI-PEDIUM...EPIDENDRUM...GUNNERA...FORSTERA...and DISA.

Order 2. TRIANDRIA, comprehending such plants as have *three* stamina. This order contains *four* genera, viz. SISYRINCHIUM...FERRARIA...STILAGO...and SALACIA.

Order 3. TETRANDRIA, comprehending such plants as have *four* stamina. Of this order there is but *one* genus, viz. NE-PENTHES.

Order 4. PENTANDRIA, comprehending such plants as have *five* stamina. This order contains *three* genera, viz. PASSIFLORA...GLUTA...and AYENIA.

Order 5. HEXANDRIA, comprehending such plants as have *six* stamina. This order contains *two* genera, viz. ARISTOLOCHIA...and PISTIA.

Order 6. OCTANDRIA, comprehending such plants as have *eight* stamina. Of this order there is only one genus, viz. SCOPOLIA.

Order 7. DECANDRIA, comprehending such plants as have *ten* stamina. Of this order there are but *two* genera, viz. HELIC-TERES...and KLEINHOVIA.

Order 8. DODECANDRIA, comprehending such plants as have *twelve* stamina. This order contains but one genus, viz. CYTINUS.

* Keel-shaped. EDITOR.

† Like filings or saw-dust; *i. e.* very small. EDITOR.

‡ For figures illustrative of these plants, vide Doctor Thornton's Practical Botany, vol. I.

Order 9. POLYANDRIA, comprehending such plants as have *many* stamina. This order contains *eight* genera, viz. GREWIA... XYLOPIA... ARUM... DRACONTIUM... CALLA... POTHOS... AMBROSINI...and ZOSTERA.

Or, in another view of the Sexual System, as improved by Wildenow, this class contains *four* orders, which embrace *thirty-three* genera.

Order 1. MONANDRIA, comprehends such plants of this class sh have only *one* stamen, which contains *twenty-six* genera, which branch out into two divisions: 1. ORCHIDEÆ, *with spurs*, containing *six* genera; ORCHIS... HABENARIA... BONATEA... LIMODORUM... DISA...and SATYRIUM. 2. ORCHIDEÆ, *without spurs*, containing *twenty* genera; PTERYGODIUM... DISPERIS ... CORYCIUM... OPHRYS... SERAPIAS (*helleborine*)... NEOTTIA... CRANICHIS... THELYMITRA... DIURIS... ARETHUSA... EPIPACTIS... MALAXIS... CYMBIDIUM ... ONCIDIUM... EPIDENDRUM... VANILLA... AERIDES... DENDROBIUM ... STELIS...and LEPANTHES.

Order 2. DIANDRIA, comprehends plants with *two* stamina, including *four* orders; CYPRIPEDIUM (*ladies' slipper*)... STYLIDIUM... FORSTERA...and GUNNERA.

Order 3. TRIANDRIA, comprehends plants with *three* stamens, including *two* genera; SALACIA...and RHOPIMUM.

Order 4. HEXANDRIA, comprehends plants having *six* stamens, and has only *one* genus, ARISTOLOCHIA (*birthwort*).

CHAP. XXIV.

OF THE TWENTY-FIRST CLASS, MONŒCIA.

THIS class consists of such plants as have no *bisexual* flowers, but bear both *male* and *female* flowers on the same plant*. The orders of this class are *eleven*, containing *a hundred and twenty-six* genera, viz.

Order 1. MONANDRIA, comprehends such plants as have their *male* flowers furnished with *one* stamen. This order contains *eleven* genera, viz. ZANICHELLIA... CERATOCARPUS... CYNOMORIUM ... ELATERIUM... CHARA... ÆGOPHICON... ARTOCARPUS (*bread-fruit*) ... NIPA... CASUARINA... PHYLLACHNE... CAULINIA.

Order 2. DIANDRIA, comprehends such plants as have their *male* flowers furnished with *two* stamina. This order contains *three* genera, viz. LEMNA (*duck-weed*)... ANGURIA... PODOSTEMUM.

Order 3. TRIANDRIA, comprehends such plants as have their *male* flowers furnished with *three* stamina. This order contains *sixteen* genera, viz. TYPHA (*cat's tail, or reed mace*)... SPARGANGLUM (*bur-reed*)... ZEA (*mays, or Indian corn*)... COIX (*Job's tears*)... TRIPSACUM... OLYRA... CAREX (*sedge*)... AXYRIS... TRAGIA... HERNANDIA... ZEUGITES... KOBRESIA... SCLERIA... ERIOCAULON... COMPTONIA... and ACHARIA.

Order 4. TETRANDRIA, comprehends such plants as have their *male* flowers furnished with *four* stamina. This order contains *sixteen* genera, viz. BUXUS (*box*)... URTICA (*nettle*)... MORUS (*mulberry*)... CICCICA... SERPICULA... LITTORELLA... AUCUBA... DIOTIS...

* These are the androgynous plants. See Part I. Chap. XXI.

BŒMERIA... PROCRI... TRICERA... PACHYSANDRA... EMPLEURUM...
ALNUS (*alder*)... NAJAS... and ARGYTHAMNIA.

Order 5. PENTANDRIA, comprehends such plants as have the *male* flowers furnished with *five* stamina. This order contains *ten* genera, viz. XANTHIUM... AMBROSIA... AMARANTHUS... NEPHELIUM... CLIBADIUM... CROTONOPSIS... POLYCHROA... LUFFA... FRANSERIA... and SCHISANDRA.

Order 6. HEXANDRIA, comprehends such plants as have their *male* flowers furnished with *six* stamina. Of this order there are *nine* genera, viz. ZIZANIA... PHARUS... SAGUS (*sago*)... COCOS (*cocoa*)... ELATE... BACTRIS... GUETTARDA... EPIBATERIUM... and POMETIA.

Order 7. POLYANDRIA, comprehends such plants as have their *male* flowers furnished with *many* stamina. This order contains *twenty-six* genera, viz. CERATOPHYLLUM (*hornwort*)... MYRIOPHYLLUM (*water-milfoil*)... SAGITTARIA (*arrow-head*)... BEGONIA... THELIGONUM... POTERIUM (*burnet*)... QUERCUS (*oak*)... JUGLANS (*walnut*)... FAGUS (*beech*)... CUSTANEA (*chestnut*)... CORYLUS (*hazel*)... CARPINUS (*hornbeam*)... BETULA (*birch*)... PLATANUS (*plane-tree*)... ARUM*... CALADIUM... THOA... SALISBURIA (*maiden-hair tree*)... HEDYOSMUM... ACIDOTON... MABEA... PARIANA... GARCIA... MANICARIA... CARYOTA... and LIQUIDAMBAR.

Order 8. MONADELPHIA, comprehends such plants as have their *male* flowers furnished with *one* set of *united* stamina. This order contains *twenty-seven* genera, viz. HURA (*sānd-box tree*)... PINUS (*pine, fir, larch*)... CUPRESSUS (*cypress*)... THUJA (*arbor vitæ*)... ACALYPHA... DALECHAMPIA... PLUKENETIA... CUPANIA... CROTON (*tallow-tree*)... RICINUS (*palma Christi*)... JATROPIA (*physic-nut*)... STERCULIA... HIPPOMANE (*manchineel-tree*)... STILLINGIA... GNETUM... CYTINUS... BRADLEYA... NISSA... ARECA (*cabbage-tree*)... GLEONO-

* This was contained formerly in the class Gynandria, but was altered by Willdenow, and others, into this class. EDITOR.

MA...PHYLLANTHUS...AGYNEIA...EPISTYLUM...SIPHONIA...SAPIUM...OMPHALEA...HECATEA...ALEURITIS...and MYRANTHUS.

Order 9. SYNGENESIA, comprehends such plants as have their *male* flowers furnished with stamina, of which the antheræ are *united*. This order contains *six* genera, viz. TRICHOLANTHES (*snake-gourd*)...MOMORDICA (*squirting cucumber*)...CUCUMIS (*cucumber, melon*)...CUCURBITA (*gourd, pompion*)...CICYOS...and BRYONIA (*bryony*)*.

Order 10. GYNANDRIA, comprehending such plants as have their *male* flowers furnished with stamina that grow out of a kind of *style*, or imperfect *pistillum*, the perfect one being in the female flower. This order contains two genera, viz. ANDRACHINE and HYPHYDRA.

CHAP. XXV.

OF THE TWENTY-SECOND CLASS, DICECIA.

THIS class consists of such plants as have no *bisexual* flowers, but bear *male* and *female* flowers on *distinct* plants†. The orders of this class are fourteen, containing *one hundred and eleven* genera, viz.

* These genera all melt into the order Monadelphia, according to Willdenow, the Syngenesia being by him confined to compound flowers. They are here preserved distinct, as an illustration of the Sexual System. EDITOR.

† There are many plants which have male and female flowers on *distinct* plants; but which are not admitted to this class, because this circumstance happens to one species only, and not to the whole genus. Instances of this are met with in *Morus*, *Urtica*, *Laurus*, *Croton*, *Rumex*, *Silene*, *Carex*, *Rhus*, *Valeriana*, *Rhamnus*, and *Cucubulus*. But it is observable, that in the plants that stand under the first di-

Order 1. MONANDRIA, comprehends such plants as have their *male* flowers furnished with *one* stamen. This order contains *seven* genera, viz. PANDANUS...PHUCAGROSTIS...MONIMIA...ASCARINA...DIDYMELES...DAHLIA...PHELYPÆA.

Order 2. DIANDRIA, comprehends such plants as have their *male* flowers furnished with *two* stamina. This order contains *five* genera, viz. VALLISNERIA...SALIX (*willow, sallow, osier*)...CECROPIA...CERATIOLA...and BORYA.

Order 3. TRIANDRIA, comprehends such plants as have their *male* flowers furnished with *three* stamina. This order contains *ten* genera, viz. EMPETRUM...OSYRIS...CATURUS...RESTIO...MABA...PHŒNIX (*date palm*)...HELWINGIA...STILAGO...WILLDENOVIA, and ELEGIA.

Order 4. TETRANDRIA, comprehending such plants as have their *male* flowers furnished with *four* stamina. This order contains *thirteen* genera, viz. VISCUM (*misseltoe*)...HIPPOPHAE (*sea buckthorn*)...MYRICA...TROPHIS...BATIS...MONTINIA...BRUCEA...SCHEFFERIA...CAVANILLA...NAGEIA...ANTHROSPERMUM...KŒLE-RA...and BROUSSONETIA.

Order 5. PENTANDRIA, comprehends such plants as have their *male* flowers furnished with *five* stamina. This order contains *seventeen* genera, viz. PISTACIA...ZANTHOXYLUM...ASTRONIUM...IRESINE...ANTIDESMA...SPINACIA (*spinach*)...ACNIDA...CANNABIS (*hemp*)...HUMULUS (*hop*)...ZANONIA...FEUILLEA...CANARIUM...PICRAMNIA...SECURINEGA...ACNIDA...FLUGGEA, and MELICYTUS.

Order 6. HEXANDRIA, comprehends such plants as have their *male* flowers furnished with *six* stamina. This order contains *ten*

stinction, in the order *Monogynia*, of the class *Pentandria*, which are the *Asperifolia* (*rough-leaved plants*) of Ray, and also in the plants of the classes *Didynamia*, *Tetradynamia*, and *Diadelphica*, there have not been found any species where the sexes are on distinct plants: this may be accounted for from the structure of the flowers in those classes. AUTHOR.

genera, viz. TAMUS (*black bryony root*) ... SMILAX ... RAJANIA... DIOSCOREA... BRAUNIA ... FERREOLA... CHAMÆDOREA... MAURITIA ... BORASSUS...and ELAIS.

Order 7. OCTANDRIA, comprehends such plants as have their *male* flowers furnished with *eight* stamina. This order contains five genera, viz. POPULUS (*poplar*)... RHODIOLA (*rose-root*)... MARGARITARIA... COMMIPHORA... HERMESIA.

Order 8. ENNEANDRIA, comprehends such plants as have their *male* flowers furnished with *nine* stamina. This order contains three genera, viz. MERCURIALIS (*mercury*)... HYDROCHARIS ... TRIPLARIS.

Order 9. DECANDRIA, comprehends such plants as have their *male* flowers furnished with *ten* stamina. This order contains five genera, viz. CARICA (*pawpaw tree*) ... KIGGELARIA... CORIARIA... SCHINUS... GYMNOCLADUS.

Order 10. DODECANDRIA, comprehends such plants as have their *male* flowers furnished with *twelve* stamina. This order contains five genera, viz. MENISPERMUM... DATISCA... EUCLEA... STRATIOTES...and TOXICODENDRUM.

Order 11. ICOSANDRIA, comprehends such plants as have their *male* flowers furnished with *many* stamina inserted into the calyx. Of this order there are four genera, viz. FLACOURTIA... ROTTLEA... GELONIUM...and HEDICARYA.

Order 12. POLYANDRIA, comprehends such plants as have their *male* flowers furnished with *many* stamina. Of this order there are nine genera, viz. CLIFFORTIA ... PERULA... TREWIA... XYLOSMA... HISINGERA... EMBRYOPTERIS... HAMADRYAS... CYCAS... ZAMIA.

Order 13. MONADELPHIA, comprehending such plants as have their *male* flowers furnished with *one* set of *united* stamina. This order contains *sixteen* genera, viz. TAXUS (*yew tree*)... JUNIPERUS

(*juniper, cedar, savin*)...EPHEDRA (*shrubby horse-tail*)...CISSAMPELOS... NAPÆA...ADELIA...ARAUCARIA... EXCÆCARIA ... HORSFIELDIA...MYRISTICA (*nutmeg-tree*)...DRYANDRA... BATSIA...LATANIA...LOUREIRA...ZANTHE...ALCHORNEA...and NEPENTES.

Order 14. SYNGENESIA, comprehends such plants as have their *male* flowers furnished with stamina, of which the *antheræ* are united. Of this order there is but one genus, viz. RUSCUS (*butcher's-broom*)*.

Order 15. GYNANDRIA, comprehends such plants as have their *male* flowers furnished with stamina that grow out of a kind of style, or *imperfect pistillum*, the *perfect* one being in the female flower. Of this order there is but one genus, viz. CLUYTIA.

CHAP. XXVI.

OF THE TWENTY-THIRD CLASS, POLYGAMIA.

THIS class consists of such plants as bear *bisexual* flowers; and also either *male* or *female* flowers, or *both*. The orders of this class are *three*, containing *fifty-four* genera, viz.

Order 1. MONÆCIA, comprehends such plants as have the polygamy on the *same* plant. This order contains *forty-five* genera, viz. MUSA (*plantain tree, and banana tree*)...HOLCUS (*soft grass, Indian millet*)...TETRAPOGON...ANTHEROPOGON...ÆGOPOGON...ELYONURUS...ISCHÆMUM...ÆGILOPS...CHLORIS...ANDROPOGON...COLLA-

* This, by Willdenow, is inserted in the class Monadelphia. ERITREA

DOA...APLUDA...ANTHISTIRIA...VALANTIA...PLANERA...DIDY-
 MANDRA...CASTELA...OPHIOXYLON...COPROSMA...CELTIS (*nettle-*
tree)...KERNERA...VERATRUM (*white hellebore*)...MARTINEZIA...
 CEROXYLON...TRATTINNICKIA...STALAGMITIS...GYROCARPUS...
 AILANTHUS...GIMBERNATIA...GOUANIA (*chaw-stick*)...BRIEDELIA
 ...SCHRANKIA...DESMANTHUS...ACACIA...INGA...MIMOSA (*sensitive*
and humble plant)...BRABEIUM...HERITIERA...HYPELATE...TER-
 MINALIA...CLUSIA...FERONIA...HERMAS...PARIETARIA (*pellitory*)
 ...and ATRIPLEX (*orache*).

Order 2. DIÆCIA, comprehends such plants as have the poly-
 gamy on *two distinct* plants. This order contains *twenty-eight*
 genera, viz. PANAX (*ginseng*)...DIOSPYROS (*date plum*)...CHRY-
 SITRIX...SPINIFEX...ELEPHANTUSA...NUNNEZIA...CHAMÆROPS (*fan*
palm)...BREYNIA...PENNANTIA...STILBE...NYSSA...HAMILTONIA...
 LAUROPHYLLUS...FRAXINUS (*ash tree*)...RICHERIA...ISQUIERDA...
 BURSERA...GRISELINIA...HYDNOCARPUS...ARCTOPUS...GLEDIT-
 SCHIA (*three-thorned acacia*)...SCHLEICHERA...BROSIMUM...CABAL-
 LERIA...LARDIZABALA...SMEGMARIA...KAGENECKIA...and CERA-
 TONIA (*carob tree*).

Order 3. TRIÆCIA, comprehending such plants as have the
 polygamy on *three distinct* plants. This order contains *one* genus
 viz. FICUS* (*the fig-tree*).

* To understand this order, the singular manner of the fructification must be ex-
 plained. The fruit of the *Ficus* is not a *pericarpium*, but a *receptacle*, the interior
 sides of which support the flowers, which by this means are enclosed within it.
 These flowers in the cultivated fig-trees are *female* only, but there is a sort known
 by the name of *Caprificus*, that has *male* flowers, and another again called *Erinosyce*,
 which is *androgynous*, having both *male* and *female* flowers distinct, though lodged
 within the same receptacle. Here then we have the *Triæcious* polygamy explained;
 and if the descriptions of *de la Hire* may be trusted, there are figs that contain
bisexual flowers; which give us even a fourth habitation for the sexes. Thus
 much suffices to explain the order; but there is an objection naturally arising from
 hence to the doctrine of the sexes; the obviating which, will furnish the opportunity
 of a necessary remark. It will be asked, how it happens that the fruit of our fig-
 trees ripen, if the plants are of one sex only, and have no assistance from the
 male? The answer is this: the fruit is in all cases to be distinguished from the
 seed contained within it: if the male be wanting, the seed will not vegetate when

CHAP. XXVII.

OF THE TWENTY-FOURTH CLASS, CRYPTOGAMIA*.

THIS class consists of such plants as *conceal* their fructification, having their flowers either *within* the fruit, or so *small*, as not to be perceptible to the naked eye. The fructification in these is also of an uncommon structure. The orders are four, containing seventy-eight genera, viz.

Order 1. FILICES, *ferns*, comprehending such plants as are dorsiferous†. What is known of the fructification of these plants, amounts only to the few characters following :

Characters of the Filices.

CALYX—A *squama* growing out of the leaf, opening on one of its sides ; and under which there are pedunculate *globules* ; each

sown, but the *fruit* may nevertheless swell, and come to an appearance of perfection ; and so it is observed to do in the instance in question, and in many others, especially where the fruit is formed of one of the parts less connected with the seed ; as calyx, receptacle, &c. though it is more common for it to drop off before it ripens, if not fecundated by the male. AUTHOR.

* The plants of this class are often of a dangerous quality.

† Bearing the fruit on the back of the leaf. These have been called also *epiphyllouspermous*, a Greek compound expressive of the same circumstance ; *capillary*, as being esteemed good for the hair ; and *acaules*, *without stems* ; for in these plants, what rises out of the ground is plainly a leaf only ; one of the characters of a stem or trunk is, to be alike on every side ; but in the stalks of ferns, there is manifestly a front and back, the former being flat and channelled, and the latter convex ; which shows them to be leaves. AUTHOR.

globule is girt with an elastic ring, which breaks elastically, and sheds a dust, which are the seeds.

This order contains thirty-one genera, arranging under two heads: 1. *Annular*; i. e. *having the capsules girt with an elastic ring, contrary to the valves*, contains nineteen genera, which are as follows, viz. ACROSTICUM...PTERIS (*brake*)...BLECHNUM...HEMIONITIS...LONCHITIS...ASPENIUM (*spleenwort*)...POLYPODIUM...ADIANTUM... TRICHOMANES.....DAREA...SCOLOPENDRUM (*hart's tongue*)... WOODWARDIA... LINDSÆA ... VITTARIA ... DAVALLIA... DICKSONIA ... CYATHEA ... HYMENOPHYLLUM...and SCHIZÆA. 2. *Exannular, capsules without rings*, GLEICHENIA...MARATTIA...DANÆA...EQUISETUM (*horse-tail*)...OPHIOGLOSSUM (*adder's tongue*)...OSMUNDA (*moon-wort*)...LYCOPIDIUM (*club-moss*)...PORELLA...SALVINIA...MARSILEA...PILULARIA...and ISOETES (*quill-wort*).

Order 2. MUSCI, *mosses*. The character of the plants comprehended under this title are, antheræ without filaments; the female flowers distinct, and without any pistillum; and the seeds, consisting only of a naked corculum, without cotyledon or tunic. The genera of this order have been distinguished by *Linnaeus*, according to the following circumstance, viz. The antheræ, *with* or *without* a *calyptra**, placed on the same plant as the female floret, or on a *distinct* one; and the female *aggregate*, or *single*. The order contains *twenty-two* genera; 1. *With no peristome*, three genera...PHASCUM (*earth-moss*)...SPHAGNUM (*bog-moss*)...GYMNOSTOMUM (*beardless moss*). 2. *With a simple peristome*, which contains nine genera...TETRAPHIS (*four-toothed moss*)...ANDRÆA...SPLACHNUM (*gland moss*)...ENCALYPTA (*extinguisher moss*)...PTEROGONIUM (*wing moss*)...GRIMMIA...DICRANUM (*fork moss*)...TRICHOSTOMUM (*fringe moss*)...TORTULA (*screw moss*). 3. *With a double peristome*, which contains ten genera...ORTHOTRICHUM (*bristle moss*)...NECKERA...FUNARIA (*cord moss*)...BUXBAUMIA...BARTRAMIA...MNIUM (*spring moss*)...BRYUM (*thread moss*)

* A veil.

...HYPNUM (*feather moss*)...FONTINALIS (*water moss*)...and POLYTRICHUM (*hair moss*).

Order 3. HEPATICÆ*, which comprehends *sic* genera...MARCHANTIA...JUNGERMANNIA...TARGIONIA...ANTHROCEROS...BLASIA...and RICCIA.

Order 4. ALGÆ, *flags*. The plants comprehended under this order have their root, stem, and leaf, all in one. The characters of the fructification of this order are not yet known, excepting the few descriptions given by *Michelius*. The genera are six, viz. LICHEN (*liver-wort*)...TREMELLA...FUCUS...ULVA...CONFERVA...and BYSSUS.

Order 5. FUNGI, *mushrooms*. The genera of this order are given by *Linnaeus*, after the method of *Dillenius*. The fructification being imperfectly known, no character can be assigned for this order, farther than the title, which is familiar to every one. The genera are *twelve*; 1. *Capped*, four genera, viz. AGARICUS...BOLETUS...HYDNUM...PHALLUS. 2. *Without a cap*, eight genera...CLATHRUS...HELVELLA...PEZIZA...CLAVARIA...LYCOPERDON...MUCOR...OCTOSPORA...and SPERIA.

* These were formerly blended with the algæ, but have since been separated by the learned president of the Linnæan Society, Dr. Smith. EDITOR.

† *Linnaeus* tells us, he preferred the method of *Dillenius* for the fungi to that of *Michelius*; because it was plain to every one; whereas that of *Michelius*, though that author has thrown great light upon this tribe, required too nice an inspection.

AUTHOR.

CHAP. XXVIII.

OF THE APPENDIX.

BESIDES the twenty-four classes explained in the preceding chapters, *Linnaeus* has in his *Genera Plantarum* given an *Appendix*, which in the *Ordo Generum*, prefixed to that work, he calls the twenty-fifth class*. It contains only one order, viz.

PALMÆ, comprehending such plants as have a *spudix* and *spatha*. This order contains nine genera, viz. CHAMÆROPS...BO-RASSUS...CORYPHA...COCOS...PHŒNIX...ELAIS...ARECA....ELATE ...and CARYOTA†.

* *Linnaeus*, in the first edition of his *Genera Plantarum*, made two orders in his *Appendix*, which, in the last edition of the *Systema Naturæ*, he has reduced to one, finding, after more mature examination, all the plants in his second order fell naturally under the other classes and orders, to which they properly belonged.

† These have become now better understood, and have been incorporated by *Willdenow*, *Dr. Smith*, and other able botanists, into their respective classes. EDITOR.

CHAP. XXIX.

OF GENERIC DISTINCTIONS.

HAVING now gone through the explanation of the CLASSES and ORDERS of the system, we come to the distinctions of the GENERA. These, by the theory of the Sexual System, are to be regulated by the *fructification* only. The parts of fructification known to the earlier botanists were few, and might be well thought insufficient for distinguishing the vegetable productions of nature: they therefore had recourse to the habit of plants, and other circumstances; and by this means a great number of genera were established, which the new system is obliged to reject. Of these we shall give the reader an ample list of instances in Chap. XXXI.

The *fructification* being admitted as the only foundation of the generic distinctions, all vegetables that *agree* in their parts of fructification are to be put together under one genus; and all such as *differ* in those parts, are to be divided. The characteristic mark of each genus is to be fixed from the number, figure, proportion, and situation, of all the parts: but as there are few genera wherein all the parts are constant in every one of the species, we ought, wherever it is possible, to fix upon some one single circumstance that is constant, and make it the *essential* character. This in most genera may be had: thus the essence of PRUNELLA, TORENIA, EUPHRASIA, ALYSSUM, and CRAMBE, lies in the *denticles of the stamina*;...that of CURCUMA, CHELONE, BIGNONIA, and MARTYNIA, in a *mutilated stamen*;...the RANUNCULUS is distinguished by its *nectarium*, which is a pore in the claws of its petals;...HYDROPHYLLUM by the same part, which in that genus is a *closed chink* in the laciniaë of the corolla;...and HELLE-

BORUS and NIGELLA also, by their *tubulose nectaria*;...in PANCRATIUM the *stamina* are *inserted* in the *nectarium*, which distinguishes it from NARCISSUS;...in HYOSCYAMUS, there is a *covering to the capsules*, by which it is known from PHYSALIS;...the RESEDA has always a *lateral nectarium*, but varies in its corolla and pistillum;...the CAMPANULA has a *quinquevalved nectarium*, but is inconstant in the corolla and capsule;...and lastly, the IRIS has a *stigma* of singular construction, but varies in the beard of its corolla.

There is, however, no one part of fructification that can be relied on as a constant characteristic mark for all genera; it being found, that the part which is constant in some genera, will be inconstant in others: thus in CARICA the flowers of the male plant are *monopetalous*, and those of the female *pentapetalous*;...in MYRICA some species have *naked seeds*, others *berries*;...in FRAXINUS some have a *naked flower*, and others a *corolla*;...in GERANIUM some have *regular corolla*, and others *irregular*;...in LINUM some are *pentapetalous*, others *tetrapetalous*;...in ACONITUM some are *tricapular*, and others *quinquecapsular*;...and in TRIFOLIUM some are *monopetalous*, others *polypetalous*; some *monospermous*, and others *polyspermous*.

This inconstancy of particular parts in many genera has been another source of error amongst the earlier botanists, who have parted many plants from their congeners on this account. Of these mistakes we shall give an ample list in Chap. XXXII.

When the characteristic mark of any genus is wanting in any particular species, we should proceed with caution, lest we confound genera that should be distinguished: for want of this caution the ERICA and ANDROMEDA had been joined, but were parted afterwards on account of the *two horns in the anthera* of the ERICA;...the ADONIS had been joined to the RANUNCULUS, but was parted from it again, on observing that it *wanted the nectariferous pore*;...and the ALOE and AGAVE had been blended, till it was observed that *in the latter* the *stamina* were inserted in the corolla, and not in the receptacle.

When the characteristic mark of any genus is observed in

some species of another genus near of kin to it, a like caution is again necessary on the other hand, lest we should multiply the genera, by parting species that should stand together: thus we find, that in *SEDUM*, *SEMPERVIVUM*, *RHODIOLA*, *CRASSULA*, *TILLÆA*, and *COTYLEDON*, the nectaria adhere to the base of the pistillum;...in *EPILOBIUM* and *CENOTHERA* the calyx is tubulose;... in *MESPILUS*, *CRATÆGUS*, and *SORBUS*, the structure of the flower is alike;...and in both *ALNUS* and *BETULA*, there are three florets on the foliole of the amentum*.

CHAP. XXX.

BY WHAT PARTS OF FRUCTIFICATION THE GENUS MAY WITH THE MOST CERTAINTY BE DETERMINED.

THE more constant any part of the fructification is found through the several species of any genus, the more it may be relied on with certainty, as a characteristic mark for that genus. Thus in *HYPECOUM* the *nectarium* is constant, but not the *siliqua*; ...the *CONVALLARIA* is constant in its *spotted berry*, but not in its *corolla*;...the *LOBELIA* in its *corolla*, but not in its *fruit*;...the *CASSIA* in its *corolla*, but not in its *siliqua*;...and the *VERBENA* in its *calyx* and *corolla*, but not in its *stamina* and *seeds*.

In some genera one part of the fructification is found to be the most constant, and in others another; but there is no part that

* The *alnus* and *betula* are joined by *Linnæus* under the title of *Betula*. The rest of these instances he has kept separate, notwithstanding the doubt raised here concerning the propriety of distinguishing them. AUTHOR.

is not liable sometimes to a variation. Thus we find the *pericarpium* variable in IMPATIENS, CAMPANULA, PRIMULA, PAPAVER, CISTUS, FUMARIA, and ARBUTUS;...the *calyx* in NYMPHÆA, and CORNUS; the *corolla* in VACCINIUM, CONVALLARIA, ANDRŌMEDA, GENTIANA, and LINUM;...and the *seeds* in RANUNCULUS, and ALISMA.

If the *flowers* agree, but the fruits differ, the genus ought not to be parted. Thus in those extensive genera, the CASSIA, HEDYSARUM, SOPHORA, LAVATERA, HIBISCUS, and MIMOSA, so great a number of species have been ranged under the same genus, on account of the conformity in the *flowers*, though there is a variation in the *fruit*.

That the *figure* of the *flowers* is more certain than that of the *fruit*, appears from many examples; as from CAMPANULA, PRIMULA, ANTIRRHINUM, ALISMA, HIBISCUS, &c.; but the *proportion* of the parts is subject to very great variation.

The *number* of the parts is more liable to variation than their *figure*, and is found sometimes to vary even upon the same plant; as in RUTA, CHRYSOSPLENIUM, MONOTROPA, TETRAGONIA, EUONYMUS, PHILADELPHUS, and ADOXA, in the flowers of all which the number of the parts varies from five to four. In these doubtful cases, the natural number must be collected from the primary flower; but in the variations of the *number* of the parts, there is a proportional affinity worth remarking. In *flowers* the stamina usually vary from ten to eight, and from five to four; the corolla and calyx from five to four, and the whole flower from four to three; and the *fruit* also usually varies from five to three, and from five to four.

The *situation* of the parts is the most constant, very rarely varying in the same genus.

The *regularity* of the *petals* is not so much to be depended on as some former botanists* have thought; for we see in *geranium* the *European* species have regular corollæ, but the *African* ones irregular.

* *Rivinus* in particular. AUTHOR.

The *nectarium* nature has made of the greatest consequence. This part, which had not even a name, till *Linnaeus* had distinguished it, is a decisive mark in all the following genera, viz. in ORCHIS...SATYRIUM...MONOTROPA...FUMARIA...VIOLA...MALPIGHIA...BANNISTERIA...ADENANTHERA...COMMELINA...LAURUS...HELXINE...DICTAMNUS...ZYGOPHYLLUM...SWERTIA...LILIUM...FRITILLARIA...HYDROPHYLLUM...RANUNCULUS...HERMANNIA...BERBERIS...STAPHYLEA...PASSIFLORA...NARCISSUS...PANCRACTIUM...MIRABILIS...NERIUM...STAPELIA...ASCLEPIAS...DIOSMA...CAMPANULA...PLUMBAGO...HYACINTHUS...RHODODENDRUM...CHEIRANTHUS...SINAPIS...KIGGELARIA...CLUTIA...AQUILEGIA...NIGELLA...ACONITUM...PARNASSIA...EPIMEDIUM...THEOBROMA...RESEDA...GREWIA...HELLEBORUS...ISOPYRUM...TROPÆOLUM...and IMPATIENS.

The *stamina* and *calyx*, being less subject to luxuriancy, are far more certain than the *petals*.

The *corolla* varies as to its figure in many genera; as in VACCINIUM...PYROLA...ANDROMEDA...NICOTIANA...MENYANTHES...PRIMULA...VERONICA...GENTIANA...HYACINTHUS...SCABIOSA...and NARCISSUS. It varies also as to number, being in RANUNCULUS *pentapetalous* in some species, and *polypetalous* in others; ... in HELLEBORUS also, *pentapetalous* and *polypetalous*; ...in STATICE, *pentapetalous* and *monopetalous*; ...and in FUMARIA, *dipetalous* and *tetrapetalous*; ...and the number is also sometimes variable in the same species; as is observed in CARICA, and JATROPIA.

The structure of the *pericarpium* was formerly thought to be of great consequence in determining the genera; but there are examples without number that demonstrate the contrary. There are a great many genera that have been established on distinctions in the pericarpium, and that are now rejected; of these we shall give an ample list in Chap. XXXIII.

The characters of *luxuriant* flowers, whether *barren** or *mutilate*, cannot be allowed any place in determining the genera;

* Barren flowers are such as have lost the *stamina*, which is the case of full flowers. *Mutilate* are those that are incomplete, wanting the *corolla* or *perianthium*.

for in full flowers no number of petals can be assigned, and the stamina are generally wanting, the number of which makes a part of the generic character; and in mutilate flowers, as in some species of *CAMPANULA*, *IPOMÆA*, and *RUELLIA*, the corolla would be excluded from the description, contrary to the nature of the other species of the genus. But as the calyx*, in full flowers, is scarce ever altered, it may detect the genus; and the lowest series of petals in polypetalous corollæ remaining the same in respect to number, the genus may also be often known by that character; as in *PAPAVER*, *NIGELLA*, and *ROSA*.

CHAP. XXXI.

OF THE GENERA REJECTED BY THE SEXUAL SYSTEM, AS NOT ESTABLISHED ON THE FRUCTIFICATION.

WE have observed, in Chap. XXIX. that the earlier botanists had admitted many genera, on distinctions that were not grounded on the *parts of fructification*, but on the *habit of plants*, and on *other circumstances*, which are now considered as *specific distinctions* only: of these we shall here give an ample list. The

* Some systematists have distributed the whole body of vegetables by the differences of the *calyx*; and in such systems the full flowers, as our author observes, are more easily referred to their proper genus than in his own, the calyx not being subject to luxuriancy. Instances of this are in *hepatica*, *ranunculus*, and *alcea*.

AUTHOR.

We frequently find the calyx also luxuriant, and the lower leaves of the corolla increased in number. Luxuriant plants are easily known, when the single are well understood, *primâ facie*, and the young student would do well at first to pass these over. EDITOR.

reader will here take notice, that under the first column are ranged the genera that are abolished; and over-against them, in the second, the genus to which they are severally to be referred*, with the specific difference that had given occasion to the false distinction.

OLD GENERA.

NEW GENERA.

<i>Limodorum</i>	ORCHIS, with a fibrose root.
<i>Bistorta</i>	POLYGONUM, with a fleshy root.
<i>Rapa</i>	BRASSICA, with a gibbose root.
<i>Sisarum</i>	SIMUM, with a tuberosc root.
<i>Hermodactylus</i>	IRIS, with a tuberosc root.
<i>Sisyrinchium</i>	IRIS, with a double bulb, one over the other.
<i>Xiphium</i>	IRIS, with a tunicated bulb.
<i>Lilio Fritillaria</i>	FRITILLARIA, with a squamose bulb.
<i>Mesomora</i>	CORNUS, with an herbaceous stem.
<i>Anacamseros</i>	SEDUM, with an erect stem.
<i>Psyllum</i>	PLANTAGO, with a branching stem.
<i>Bellis Leucanthemum</i> ...	BELLIS, with a leafy stem.
<i>Pilosella</i>	HIERACIUM, with a naked stem.
<i>Suber</i>	QUERCUS, with a fungous bark.
<i>Larix</i>	ABIEST†, with fasciculate leaves.
<i>Genistella</i>	GENISTA, with jointed leaves.
<i>Potamopithys</i>	ALSINASTRUM‡, with leaves not starry.
<i>Lupinaster</i>	TRIFOLIUM, with digitate leaves.
<i>Dracunculus</i>	ARUM, with pedate leaves.
<i>Trichomanes</i>	ASPENIUM, with pinnate leaves.

* The names and the generic arrangement of vegetables having undergone many alterations during the progress of the improvements made in the science, the new genera, to which these false ones are referred in this and the following lists, do not all stand under the titles given to them in the later editions of the works of *Linnaeus*. Where this happens, we shall explain it by a note; choosing that method rather than to alter the lists themselves, which we have taken from the *Philosophia Botanica*. AUTHOR.

† Now *Pinus*.

‡ Now *Elatine*. AUTHOR.

OLD GENERA.

NEW GENERA.

<i>Clymenum</i>	LATHYRUS, with pinnate leaves.
<i>Muscoides</i>	{ JUNGERMANNIA, with leaves many times imbricate.
<i>Lentiscus</i>	{ TEREBINTHUS*, with no odd foliole to the leaves.
<i>Faba</i>	VICIA, with leaves that have no cirrhus.
<i>Cytisogenista</i>	SPARTIUM, with leaves simple and triple.
<i>Colocasia</i>	ARUM, with leaves not ear-shaped.
<i>Cirsium</i>	CARDUUS, with leaves without thorns.
<i>Coronopus</i>	COCHLEARIA, with a pinnatifid leaf.
<i>Coronopus</i>	PLANTAGO, with dentate leaves.
<i>Ilex</i>	QUERCUS, with denticulate leaves.
<i>Scorzoneroides</i>	SCORZONERA, with dentate leaves.
<i>Anguria</i>	CUCURBITA, with multifid leaves.
<i>Alcea</i> †.....	MALVA, with multifid leaves.
<i>Millefolium</i>	PTARMICA, with leaves minutely divided.
<i>Cicutaria</i>	LIGUSTICUM, with a cicuta leaf.
<i>Cedrus</i>	JUNIPERUS, with a cypress leaf.
<i>Ranunculoides</i>	RANUNCULUS, with capillary leaves.
<i>Alhagi</i>	HEDYSARUM, with simple leaves.
<i>Nissolia</i>	LATHYRUS, with simple leaves.
<i>Marsilea</i>	JUNGERMANNIA, with simple leaves.
<i>Balsamita</i>	TANACETUM, with undivided leaves.
<i>Cepa</i>	ALLIUM, with fistulous leaves.
<i>Aphaca</i>	LATHYRUS, with no leaves, but stipulæ only.
<i>Mimosa</i>	ACACIA‡, with sensitive leaves.
<i>Oxyoides</i>	OXALIS, with sensitive winged leaves.
<i>Aurantium</i>	CITRUS, with cordate petioles§.
<i>Calamintha</i>	MELISSA, with branching peduncles .

* Now *Pistacia*. AUTHOR.

† *Alcea* is still the title of a genus, though of a different one; being applied to the *Malva Rosea*, or *Hollyhock*. AUTHOR.

‡ *Mimosa* is now the title of the whole genus, including the *Acacias*. AUTHOR.

§ Footstalks of the leaves. EDITOR.

|| Footstalks of the flowers. EDITOR.

OLD GENERA.	NEW GENERA.
<i>Cotinus</i>	RHUS, with woolly peduncles.
<i>Virga Sanguinea</i>	CORNUS, with a naked cyme.
<i>Corona Imperialis</i> ...	{ FRITILLARIA, with a head of leaves on the racemus.
<i>Stæchas</i>	LAVANDULA, with bracteæ on the spike.
<i>Carex</i>	CYPEROIDES*, with androgynous spikes.
<i>Chamæpithys</i>	TEUCRIUM, with sparsed leaves.
<i>Acinos</i>	THYMUS, with sparsed leaves.
<i>Limonium</i>	STATICE, with sparsed leaves.
<i>Chomædryes</i>	TEUCRIUM, with verticillate leaves.
<i>Thymbra</i>	SATUREIA, with sparsed leaves.
<i>Volubilis</i>	IPOMOEA, with flowers in heads.
<i>Polium</i>	TEUCRIUM, with cymose flowers.
<i>Castanea</i>	FAGUS, with flowers in spikes.
<i>Fagopyrum</i>	{ POLYGONUM, with spiked flowers, and a fibrose root.
<i>Majorana</i>	ORIGANUM, with rounder spikes of flowers.
<i>Malus</i>	PYRUS, with a distinct face.
<i>Cydonia</i>	PYRUS, with a distinct face.
<i>Armeniaca</i>	PRUNUS, with a distinct face.
<i>Cerasus</i>	PRUNUS, with a distinct face.
<i>Lauro-Cerasus</i>	PRUNUS, with a distinct face.
<i>Limon</i>	CITRUS, with a distinct face.
<i>Napus</i>	BRASSICA, with a distinct face.
<i>Absinthium</i>	ARTEMISIA, with the outward face distinct.
<i>Abrotanum</i>	ARTEMISIA, with the outward face distinct.
<i>Bellidiastrum</i>	DORONICUM, with a distinct habit.
<i>Euphorbia</i>	TITHYMALUS†, with the habit not branching.
<i>Usnea</i>	LICHEN, with the habit capillary.
<i>Coralloides</i>	LICHEN, with the habit caulescent.
<i>Clavaria</i>	CORALLOIDES‡, with the habit not branching.

* *Carex* is now the title of the genus. AUTHOR.

† *Euphorbia* is now the title of the genus.

‡ Now *Clavaria*.

OLD GENERA.

NEW GENERA.

<i>Tuber</i>	LYCOPERDON, with a more solid substance.
<i>Fungoides</i>	{ ELVELA, with a substance smooth on both sides.
<i>Lycoperdoides</i>	LYCOPERDON, with a cellular substance.
<i>Amanita</i>	AGARICUS, with the pileus on a stipes.
<i>Phallus</i>	{ BOLETUS, with a volva at the base of the stipes.
<i>Phalloboletus</i>	{ BOLETUS, with a pileus not closed in the sides.
<i>Polyporus</i>	{ BOLETUS, with pores not to be distinguished.
<i>Erinaceus</i>	ULEX, thick-set with spines.
<i>Thysselinum</i>	SELINUM, with a milky juice.
<i>Moly</i>	ALLIUM, with a sweet scent.
<i>Acetosa</i>	LAPATHUM*, with an acid taste.
<i>Colocynthis</i>	ANGURIA†, with a bitter fruit.

 CHAP. XXXII.

OF THE GENERA REJECTED BY THE SYSTEM, AS GROUNDED ON THE VARIATIONS OF SOME PARTS ONLY OF THE FRUCTIFICATION.

IT has been observed, in Chap. XXIX., that there are few genera, wherein all the parts of fructification are constant in every species; and that this inconstancy of particular parts had been another source of error in former botanists. We shall here

* Now *Rumex*.† Now *Cucumis*.

give a list of these mistakes, referring the old genera to the new titles, in the same manner as we did those in the list given in the preceding chapter.

OLD GENERA.	NEW GENERA.
<i>Arisarum</i>	ARUM, with a hooded spatha.
<i>Asteriscus</i>	BUPHTHALMUM, with a starry leafy calyx.
<i>Silybum</i>	CARDUUS, with a thorny calyx.
<i>Moldavica</i>	{ DRACOCEPHALUM, with the calyx gibbous and bilabiate.
<i>Tithymaloides</i>	{ EUPHORBIA, with the calyx gibbous and irregular.
<i>Trionum</i>	HIBISCUS, with an inflated calyx.
<i>Ficaria</i>	{ RANUNCULUS, with a triphyllous calyx, and polypetalous.
<i>Iva</i>	TEUCRIUM, with a gibbous calyx.
<i>Lunularia</i>	{ MARCHANTIA, with the common calyx. quadrifid.
<i>Leucanthemum</i>	{ CHRYSANTHEMUM, with the squamæ of the calyx narrow.
<i>Cardiaca</i>	LEONURUS*, with a quinquedentate calyx.
<i>Paronychia</i>	{ HERNIARIA, with the leaves of the calyx hooded.
<i>Pseudo-Dictamnus</i>	MERRUBIUM, with a funnel-shaped calyx.
<i>Anemone-Ranunculus</i>	{ ANEMONOIDES†, with a pentapetalous corolla.
<i>Linaria</i>	ANTIRRHINUM, with a tailed corolla.
<i>Valerianoides</i>	VALERIANA, with a tailed corolla.
<i>Bromelia</i>	ANANAS‡, with a tripetalous corolla.
<i>Opuntia</i>	MELOCACTUS , with a polypetalous corolla.
<i>Glacium</i>	CHELIDONIUM, with a rosaceous corolla.

* The scarlet *leonurus* of the Cape is removed to the genus *phlomis*, on account of its wanting the shining points on the antheræ; but the title *leonurus* is nevertheless applied to the *cardiaca*.

† Now *Anemone*.

‡ *Bromelia* is now the title of the genus.

|| Now *Cactus*.

OLD GENERA.

NEW GENERA.

<i>Polygonatum</i>	LIL. CONVALLIUM*, with a tubulose corolla.
<i>Centaurium minus</i>	GENTIANA, with a funnel-shaped corolla.
<i>Liliastrum</i>	HEMEROCALLIS, with a hexapetalous corolla.
<i>Borbonia</i>	LAURUS, with a pentaphylloideous calyx.
<i>Benjoe</i>	LAURUS, with an octofid corolla.
<i>Auricula Ursi</i>	PRIMULA, with an hypocrateriform corolla.
<i>Triphylloides</i>	TRIFOLIUM, with a monopetalous corolla.
<i>Oxycooccus</i>	VACCINIUM, with a tetrapetalous corolla.
<i>Bonarota</i>	VERONICA, with a tubulose corolla.
<i>Zannonia</i>	COMMELINA, with a tripetalous corolla.
<i>Borraginoides</i>	BORRAGO, with an infundibuliform corolla.
<i>Horminum</i>	{ SALVIA, with a galeate galea, and a concave beard.
<i>Sclarea</i>	{ SALVIA, with a falcate galea, and a concave beard.
<i>Phelypæa</i>	{ CLANDESTINA†, with a galea of the corolla bifid.
<i>Murucuja</i>	PASSIFLORA, with an undivided nectarium.
<i>Sherardia</i> ‡.....	VERBENA, with two stamina.
<i>Stellaris</i>	{ ORNITHOGALUM, with stamina that are not flat.
<i>Porrum</i>	ALLIUM, with trifid stamina.
<i>Dodonæa</i>	ILEX, with a trifid flower.
<i>Hypocistis</i>	ASARUM, with a quadrifid flower.
<i>Radiola</i>	LINUM, with a quadrifid flower.
<i>Unifolium</i>	CONVALLARIA, with a quadrifid flower.
<i>Bernhardia</i>	CROTON, with dioecious flowers.
<i>Petasites</i>	TUSSILAGO, with fasciculate flowers.
<i>Ananthocyclus</i>	COTULA, with flosculose flowers.
<i>Ceratocephalus</i>	BIDENS, with radiate flowers.
<i>Doria</i>	SOLIDAGO, with few florets in the radius.
<i>Medium</i>	CAMPANULA, with fruit quinquelocular.

* Now *Convallaria*.† Now *Lathræa*.‡ The title *Sherardia* is still in use, but is applied to another genus;

OLD GENERA:

NEW GENERA.

<i>Speculum Veneris</i>	CAMPANULA, with siliquose fruit.
<i>Cornucopioides</i>	VALERIANA, with an irregular flower.
<i>Limonioides</i>	STATICE, with a monopetalous flower.
<i>Viscaria</i>	SILENE, with a quinquelocular fruit.
<i>Tetragonolobus</i>	LOTUS, with an angular fruit.

 CHAP. XXXIII.

 OF THE GENERA REJECTED BY THE SYSTEM, AS GROUNDED ON
 A DIFFERENCE IN THE FRUIT ONLY.

IT has been observed, in Chap. XXX., that a great many genera had been established on account of differences in the pericarpium, but that they have since been abolished. Of these the following is a list; in which, as in the preceding lists, it will appear where they are now ranged.

OLD GENERA.

NEW GENERA.

<i>Clandestina</i>	ANBLATUM*, with an elastic fruit.
<i>Trollius</i> †.....	HELLEBORUS, with a multicapsular fruit.
<i>Sesamoides</i>	RESEDA, with a multicapsular fruit.
<i>Lycopersicon</i>	SOLANUM, with a multicapsular fruit.
<i>Ascyrum</i> ‡.....	HYPERICUM, with a quinquecapsular fruit.
<i>Dortmanna</i>	RAPUNTIIUM§, with a bilocular fruit.

* Now *Lathraea*.† *Trollius* and *Helleborus* are parted again.‡ The title *Ascyrum* is still in use for another genus.§ Now *Labellia*.

OLD GENERA.

NEW GENERA.

<i>Helianthemum</i> ...	CISTUS, with an unilocular fruit.
<i>Androsænum</i>	HYPERICUM, with an unilocular fruit.
<i>Pavia</i>	ESCULUS, with an unilocular fruit.
<i>Asarina</i>	ANTIRRHINUM, with multivalvular fruit.
<i>Elatine</i>	{ ANTIRRHINUM, with the fruit bursting on the side.
<i>Nelumbo</i>	{ NYMPHÆA, with the fruit perforate at the top.
<i>Raphanistrum</i>	RAPHANUS, with articulate fruit.
<i>Cakile</i>	BUNIAS, with articulate fruit.
<i>Ulmaria</i>	FILIPENDULA*, with twisted fruit.
<i>Persica</i>	AMYGDALUS, with a succulent fruit.
<i>Cassia</i>	SENNA†, with a succulent fruit.
<i>Inga</i>	ACACIA‡, with a succulent fruit.
<i>Malvaviscus</i>	HIBISCUS, with a succulent fruit.
<i>Lobelia</i>	RAPUNTIUM§, with a drupaceous fruit.
<i>Pereskia</i>	CACTUS, with a leafy fruit.
<i>Sabina</i>	JUNIPERUS, with a warted fruit.
<i>Bihai</i>	MUSA, with a trispermous fruit.
<i>Alaternus</i>	RHAMNUS, with a trispermous fruit.
<i>Frangula</i>	RHAMNUS, with a dispermous fruit.
<i>Dracunculus</i>	HÆMANTHUS, with monospermous fruit.
<i>Onobrychis</i>	HEDYSARUM, with monospermous fruit.
<i>Malvinda</i>	ABUTILON , with a fruit not inflatè.
<i>Cysticapnos</i>	FUMARIA, with an inflatè fruit.
<i>Impatiens</i>	BALSAMINA¶, with an attenuate fruit.

* Now *Spiræa*.

† *Cassia* is now the title of the genus, which includes the *cassia fistula*, and many other species; but the *cassia lignæa* of *Sumatra*, whose bark so nearly resembles that of the *cinnamomum*, is a *laurus*, as is the *cinnamomum* also; and the two plants are by some supposed to be the same.

‡ Now *Mimosa*.

§ *Lobelia* is now the title of the genus.

|| Now *Sida*.

¶ *Impatiens* is now the title of the genus.

OLD GENERA.	NEW GENERA.
<i>Guazuma</i>	CACAO*, with a reticulate fruit.
<i>Paliurus</i>	RHAMNUS, with a shield-shaped fruit.
<i>Alisma</i>	DAMASONIUM†, with a fruit not corniculate.
<i>Securidaca</i> ‡.....	CORONILLA, with faulchion-shaped fruit.
<i>Melo</i>	CUCUMIS, with an ovate fruit.
<i>Melopepo</i>	CUCURBITA, with a sulcate fruit.
<i>Rapistrum</i>	CRAMBE, with a fruit that does not open.
<i>Radicala</i>	SISYMBRIUM, with a siliculose fruit.
<i>Bluttaria</i>	VERBASCUM, with a rounder fruit.
<i>Persea</i>	{ LAURUS, with a fruit that is berried on every side.
<i>Cururi</i>	{ SERIANA§, with a fruit that bears seeds at the top.
<i>Bursa Pastoris</i>	THLAPSI, with a fruit that has no margin.
<i>Nasturtium</i>	LEPIDIUM, with a margin to the fruit.
<i>Valerianella</i>	VALERIANA, with a fruit not pappose.
<i>Anemonoides</i>	ANEMONE, with naked seeds.
<i>Eupatoriophalacrum</i> ...	VERBESINA, with naked seeds.
<i>Lcontodontoides</i>	HYOSERIS, with seeds almost naked.
<i>Atractylis</i> 	{ CARTHAMUS, with an obsolete crown to the seeds.
<i>Carthamoides</i>	CARTHAMUS, with pappose seeds.
<i>Zazinth</i>	LAPSANA, with pappose seeds.
<i>Alypum</i>	GLOBULARIA, with pappose seeds.
<i>Xeranthemoides</i>	XERANTHEMUM, with a feathered pappus.
<i>Astercropterus</i>	ASTER, with a feathered pappus.
<i>Acarna</i>	CNICUS, with a feathered pappus.
<i>Achyrophorus</i>	HYPOCHÆRIS, with a feathered pappus.
<i>Carlinoides</i>	CARLINA, with an obsolete pappus.

* Now *Theobroma*.† *Alisma* is now the title of the genus.‡ *Securidaca* is still a title, but of a different genus.§ Now *Paullinia*.|| *Atractylis* is still a title, but applied to another genus.

OLD GENERA.

NEW GENERA.

<i>Viticella</i>	CLEMATIS, with tailed seeds.
<i>Nymphoides</i>	MENYANTHES, with an arillus to the seed.
<i>Karatas</i>	BROMELIA, with no arillus to the seed.
<i>Tragopogonoides</i>	TRAGOPOGON, with bent seeds.
<i>Tinus</i>	VIBURNUM, with pear-shaped seeds.
<i>Opulus</i>	VIBURNUM, with heart-shaped seeds.
<i>Persicaria</i>	POLYGONUM, with triangular seeds.
<i>Emerus</i>	CORONILLA, with cylindrical seeds.
<i>Feniculum</i>	ANETHUM, with thick seeds.
<i>Lens</i>	CICER, with lens-shaped seeds.
<i>Pepo</i>	CUCURBITA, with seeds not emarginate.
<i>Falcaria</i>	SIUM, with slender seeds.
<i>Cerinthoides</i>	CERINTHE, with four distinct seeds.
<i>Blæria</i>	SHERARDIA, with echinate seeds*.

These changes will be better seen from the annexed Tables.

* For origin of the names of the genera, both classical and English, and the essential and natural generic characters, the reader is referred to Doctor Thornton's Practical Botany, being a New Illustration of the Genera of Plants; with a Description, and Plates of Dissections of each Genus. This book will be found a useful introduction to that work.

The reader is also referred to another work, entitled The Botanists' Vade Mecum; where all the essential generic characters are given by themselves: a most useful pocket-companion to the botanist.

TABLE I.
 CHANGES IN THE NAMES OF GENERA,
 ARRANGED IN
 ALPHABETICAL ORDER.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
A		
<i>Abies, Tourn</i>	Fir.....	PINUS.
<i>Abrotanum, Tourn</i>	Southernwood.....	ARTEMISIA.
<i>Absinthium, Tourn</i> <i>and Vaill. A. G.</i>	Wormwood.....	ARTEMISIA.
<i>Abutilon, Dill. Elth.</i> <i>and Tourn.</i>	Indian Mallow.....	SIDA.
<i>Abutilon, Dill. Elth</i>	Carolina Mallow.....	MALVA.
<i>Acacia, Tourn</i>		MIMOSA.
<i>Acajou, Tourn</i>	Cashew Nut.....	ANACARDIUM.
<i>Acarna, Vaill. A. G</i>	Blessed Thistle.....	CNICUS.
<i>Acetosa, Tourn</i>	Sorrel.....	RUMEX.
<i>Achyraantha, Dill. Elth</i>		ACHYRANTHES.
<i>Achyronia, Royen</i>	African Broom.....	ASPALATHUS.
<i>Achyrophorus, Vaill. A. G</i>		HYPOCHÆRIS.
<i>Acinodendron, Lin</i> <i>gen. pl. ed. prim.</i>	American Gooseberry.....	MELASTOMÆ.
<i>Acinos, Dill. gen</i>	Wild, or Stone Basil.....	THYMUS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
<i>Acnide, Mitch</i>		ACNIDA.
<i>Adhatoda, Tourn</i>	Malabar Nut.....	JUSTICIA.
<i>Ægilops, Dill. gen</i>	Oat Grass.....	BROMUS.
<i>Ageratum, Tourn</i>		ERINUS.
<i>Agnanthus, Vaill. A. G.</i>		CORNUTIA.
<i>Agrimonoides, Tourn</i> ...	Bastard Agrimony.....	AGRIMONIA.
<i>Ahouai, Tourn</i>		CERBERA.
<i>Alaternus, Tourn</i>	False Phyllyrea.....	RHAMNUS.
<i>Alcea, Tourn</i>	Vervain Mallow.....	MALVA.
<i>Alchimilla, Tourn</i>	Ladies' Mantle.....	ALCHEMILLA.
<i>Alga, Raj. Ang</i>	Grass-wrack.....	ZOSTERA.
<i>Algoïdes, Vaill. A. G.</i>		ZANNICHELLIA
<i>Alhagi, Tourn</i>	French Honeysuckle.....	HEDYSARUM.
<i>Alkekengi, Tourn</i>	Winter Cherry.....	PHYSALIS.
<i>Alnus, Tourn</i>	Alder.....	BETULA.
<i>Aloïdes, Boer. Lugd</i>	Water Soldier.....	STRATIOTES.
<i>Alpina, Plum</i>		ALPINIA.
<i>Alsinastrum, Vaill. B. P.</i>		ELATINE.
<i>Alsine, Tourn</i>	Great Chickweed.....	STELLARIA.
<i>Alsinella, Dill. gen</i>		SAGINA.
<i>Alsinoides, Raj</i>		BUFONIA.
<i>Alsinoides, Vaill. B. P.</i>		MONTIA.
<i>Alypum, Niss. A. G.</i> ...	Blue Daisy.....	GLOBULARIA.
<i>Alyssoides, Tourn</i>	Madwort.....	ALYSSUM.
<i>Amanita, Dill</i>	Agaric.....	AGARICUS.
<i>Amaranthi species, Tourn</i>		AMARANTHUS.
<i>Amaranthoides, Tourn</i> ...	Globe Amaranth.....	GOMPHRENA.
<i>Amberboi, Vaill</i>	Sweet Oriental Cyanus... called Sweet Sultan.	CENTAUREA.
<i>Amethystina, Amman</i>		AMETHYSTEA.
<i>and Hall.</i>		
<i>Ammoides, Boerh</i>	Bishop's Weed.....	AMMI.
<i>AMPANA, Hort. Mal</i>	Malabar Palm (male).....	BORASSUS.
<i>Anacampseros, Tourn</i> ...	Orpine.....	SEDUM.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
<i>Anacampseros, Lin.....</i> <i>gen. pl. ed. prim.</i>	Evergreen African Purslane	PORTULACA.
<i>Anagallidastrum, Mich.....</i>		CENTUNCULUS.
<i>Ananas, Tourn.....</i>	Pine Apple.....	BROMELIA.
<i>Ananthocyclos, Vaill.....</i> <i>A.G. and Dill. Elth.</i>		COTULA.
<i>Anapodophyllum,.....</i> <i>Tourn.</i>	Duck's-foot, or May Apple	PODOPHYLLUM
<i>Androsæmum, Tourn....</i>	Tutsan, or Park Leaves....	HYPERICUM.
<i>Anemone ranunculus,..</i> <i>Dill. gen.</i>	Wind Flower.....	ANEMONE.
<i>Anemonoides, Dill.gen...</i> <i>and Vaill. A. G.</i>	Wood Anemone.....	ANEMONE.
<i>Anemoniospermos, Com.....</i> <i>Hort. Amst.</i>		ARCTOTIS.
<i>Angiopteris, Mitch.....</i>		ONOCLEA.
<i>Anguina, Trew.....</i>	Water Dragons.....	CALLA.
<i>Anguina, Mich.....</i>	Serpent Cucumber.....	TRICHOSAN- THES.
<i>Anguria, Tourn.....</i>	Water Melon.....	CUCURBITA.
<i>Anonis, Tourn.....</i>	Rest-harrow.....	ONONIS.
<i>Anonymos, Gron. virg.....</i>		CHELONE.
<i>Antanisophyllum, Vaill.</i> <i>A. G.</i>	Hog-weed.....	BOERHAAVIA.
<i>Anthyllis, Magn. char.....</i>		CRESSA.
<i>Aparine, Tourn.....</i>	Clivers, or Goose Grass...	GALIUM.
<i>Aphaca, Tourn.....</i>	Yellow Vetchling.....	LATHYRUS.
<i>Aphyllon, Mich.....</i>	Single-flowered Broom... Rape.	OROBANCHE.
<i>Apios, Boerh.....</i>	Knobbed-rooted Liquor- ice Vetch.	GLYCINE.
<i>Apocynum, Tourn.....</i>	Dog's Bane.....	ASCLEPIAS.
<i>Aponogeton, Pont. Anth.</i>	Triple-headed Pond-weed	ZANICHELLIA.
<i>Aquifolium, Tourn.....</i>	Holly.....	ILEX.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Arachidna, <i>Plumb</i>	Ground Nut.....	ARACHIS.
Arachidnoides, <i>Niss</i>	Ground Nut.....	ARACHIS.
A. G.		
Araliastrum, <i>Vaill</i>	Ginseng.....	PANAX.
Arapabaca, <i>Plumb</i>	Worm Grass.....	SPIGELIA.
Arctotheca, <i>Vaill.</i> A. G.....		ARCTOTIS.
Arisarum, <i>Tourn</i>	Friar's Cowl.....	ARUM.
Armeniaca, <i>Tourn</i>	Apricot.....	PRUNUS.
Aronia, <i>Mitch</i>	Floating Arum.....	ORONTIUM.
Aruncus, <i>Lin. gen. pl.</i> ... <i>ed. prim.</i>	Greater Meadow-sweet....	SPIRÆA.
Asarina, <i>Tourn</i>	Snap-dragon, with Ivy Leaves.	ANTIRRHINUM
Ascyrum, <i>Tourn</i>	St. Peter's Wort, with great Flowers.	HYPERICUM.
Aspergillus, <i>Mich</i>		BYSSUS.
Asteriscus, <i>Dill. Elth</i> ...	Bastard Chrysanthemum...	SILPHIUM.
Asteriscus, <i>Tourn. Vaill</i> ...	Ox Eye.....	BUPHTHALMUM
A. G. and <i>Dill. Elth.</i>		
Asterocephalus, <i>Vaill</i> ...	Scabious.....	SCABIOSA.
A. G.		
Asteroides, <i>Tourn.</i>	Ox Eye.....	BUPHTHALMUM
<i>and Vaill. A. G.</i>		
Asteropterus, <i>Vaill.</i> A. G.	Star-wort.....	ASTER.
Astragaloides, <i>Tourn</i>	Bastard Milk-vetch.....	PHACA.
Atractylis, <i>Vaill.</i> A. G.	Distaff Thistle.....	CARTHAMUS.
Aurantium, <i>Tourn</i>	Orange.....	CITRUS.
Aureliana, <i>Laft</i>	Ginseng.....	PANAX.
Auricula Ursi, <i>Tourn</i> ...	Auricula, or Bear's Ear....	PRIMULA.
Azederach, <i>Tourn</i>	Bead Tree.....	MELIA.
B		
Baccharis, <i>Vaill.</i> A. G.	Lavender Cotton.....	SANTOLINA.
Badiaga, <i>Buxb</i>	River Sponge.....	SPONGIA.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
Ballote, <i>Tourn</i>	Black Horehound.....	BALLOTA.
Balsamina, <i>Tourn</i>	Balsam.....	IMPATIENS.
Balsamita, <i>Vaill.</i> A. G.	Costmary.....	TANACETUM.
Barba capræ, <i>Tourn</i> ...	Greater Meadow-sweet...	SPIRÆA.
Belladona, <i>Tourn</i>	Deadly Nightshade.....	ATROPA.
Bellidiastrum, <i>Mich</i>	Middle Daisy.....	DORONICUM.
Bellidioides, <i>Vaill.</i> A.G.	Greater, or Ox-eye Daisy	CHRYSANTHE- MUM.
Bellis-Leucanthemum,...	Annual Daisy.....	BELLIS.
<i>Mich. gen.</i>		
Benzoë, <i>Boerh</i>	Benjamin Tree.....	LAURUS.
Bermudiana, <i>Tourn. and</i>		SISYRINCHIUM.
<i>Dill. Elth.</i>		
Bernhardia, <i>Houst.</i> A.A.	Bastard Ricinus.....	CROTON.
Bidentis species, <i>Dill</i> ...	Tick-seeded Sun-flower...	COREOPSIS.
<i>Elth.</i>		
Bihai, <i>Plum</i>	Banana.....	MUSA.
Bistorta, <i>Tourn</i>	Bistort, or Snake-weed....	POLYGONUM.
Blairia, <i>Houst.</i> A. A....	Vervain.....	VERBENA.
Blattaria, <i>Tourn</i>	Moth Mullein.....	VERBASCUM.
Boletus, <i>Mich</i>		PHALLUS.
Bonarota, <i>Mich</i>	Rock Germander.....	VERONISA.
Bonduc, <i>Plum</i>	Nickar Tree.....	GUILANDINA.
Boraginoides, <i>Boerh</i>	Indian Borage.....	BORRAGO.
Borbonia, <i>Plum</i>	Red Bay of Carolina.....	LAURUS.
Botrytis, <i>Mich</i>		BYSSUS.
Bovista, <i>Dill</i>		LYCOPERDON.
Bryonioides, <i>Dill. Elth.</i>	Single-seeded Cucumber...	SICYOS.
Bucca-ferrea, <i>Mich</i>		RUPPIA.
Buglossum, <i>Tourn</i>	Bugloss.....	ANCHUSA.
Bugula, <i>Tourn</i>	Bugle.....	AJUGA.
Bulbine, <i>Lin. gen. pl</i> ...	Cape Spiderwort.....	ANTHERICUM.
<i>ed. prim.</i>		
Bulbocastanum, <i>Tourn.</i>	Pig-nut, or Earth-nut....	BUNIUM.
Buphthalmum, <i>Tourn</i> ...	Ox-eye, of old authors....	ANTHEMIS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Bupleuroides, <i>Boerh.</i>	Bastard Hare's-ear.....	PHYLLUS.
Bursa Pastoris, <i>Tourn.</i> ...	Shepherd's Pouch.....	THLASPI.
C		
Caapeba, <i>Plum.</i>		CISSAMPELOS.
Cacalanthemum, <i>Dill.</i>		CACALIA.
<i>Elth.</i>		
Cacao, <i>Tourn.</i>	Chocolate Nut.....	THEOBROMA.
Cainito, <i>Plum.</i>	Star Apple.....	CHRYSOPHYLLUM.
Calaba, <i>Plum.</i>		CALOPHYLLUM.
Calamintha, <i>Tourn.</i>	Calamint.....	MELISSA.
Calamus aromaticus, ...	Sweet Rush.....	ACORUS.
<i>Pet. gen. and Mich.</i>		
Calceolus, <i>Tourn.</i>	Ladies' Slipper.....	CYPRIPEDIUM.
Calcitrapa, <i>Vaill.</i>	Star Thistle.....	CENTAUREA.
Calcitrapoides, <i>Vaill.</i> ...	Thorny Knapweed.....	CENTAUREA.
Caltha, <i>Tourn. and Vaill.</i>	Marigold.....	CALENDULA.
A. G.		
Camara, <i>Plum. and Dill.</i>	American Viburnum.....	LANTANA.
<i>Elth.</i>		
Cameraria, <i>Dill. gen.</i>	Small Water Chickweed, ...	MONTIA.
or Blinks.		
Camphora, <i>Gronov. diss.</i>	Camphor Tree.....	LAURUS.
Camphorata, <i>Tourn.</i>	Stinking Ground-Pine.....	CAMPHOROSMA.
Cannabina, <i>Tourn. cor.</i> ...	Bastard Hemp.....	DATISCA.
Cannacorus, <i>Tourn.</i>	Indian Flowering Reed...	CANNA.
Capnoides, <i>Tourn.</i>	Fumatory.....	FUMARIA.
Caprifolium, <i>Tourn.</i>	Honeysuckle.....	LONICERA.
Caprificus, <i>Pont. Anth.</i> ...	Wild Fig-tree.....	FICUS.
Caraguata, <i>Plum.</i>		TILLANDSIA.
Caraxeron, <i>Vaill. A. G.</i>	Globe Amaranth.....	GOMPHRENA.
Cardamindum, <i>Tourn.</i> ...	Indian Cress.....	TROPÆOLUM.
Cardiaca, <i>Tourn.</i>	Motherwort.....	LEONURUS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
Cardispermum, <i>Trant</i> ... A. G.	Marigold.....	CALENDULA.
Cardui species, <i>Tourn</i> ...	Woolly Thistle.....	ONOPORDUM.
Carelia, <i>Pont. diss</i>	Bastard Hemp-Agrimony..	AGERATUM.
Carimpana, <i>Hort. Mal.</i>	Malabar Palm (female)...	BORASSUS.
Carlinoides, <i>Vaill.</i> A. G.	Carline Thistle.....	CARLINA.
Carpobolus, <i>Mich</i>		LYCOPERDON.
Carthamoides, <i>Vail.</i> A. G.	Bastard Saffron.....	CARTHAMUS.
Carui, <i>Tourn</i>	Caraway.....	CARUM.
Caryophyllata, <i>Tourn</i> ...	Avens, or Herb Bennet....	GEUM.
Caryophyllodendron, ... <i>Vaill.</i> A. G.	Clove-tree.....	CARYOPHYLLUS
Caryophyllus, <i>Tourn</i>	Pink, Clove July-Flower, .. Sweet William, &c.	DIANTHUS.
Caryophyllus aromati- cus, <i>Tourn.</i>	Clove-tree.....	CARYOPHYLLUS
Casia, <i>Tourn</i>	Poet's Cassia.....	OSYRIS.
Cassida, <i>Tourn</i>	Skull Cap.....	SCUTELLARIA.
Castanea, <i>Tourn</i>	Chestnut.....	FAGUS.
Castorea, <i>Plum</i>		DURANTA.
Catanance, <i>Tourn</i>	Candy Lion's Foot.....	CATANANCHE.
Cataria, <i>Tourn</i>	Cat-mint.....	NEPETA.
Cedrus, <i>Tourn</i>	Cedar.....	JUNIPERUS.
Ceiba, <i>Plum</i>	Silk Cotton-Tree.....	BOMBAX.
Centaureum majus, <i>Tour.</i>	Centaury.....	CENTAUREA.
Centaureum minus, <i>Tour.</i>	Lesser Centaury.....	GENTIANA.
Cepa, <i>Tourn</i>	Onion.....	ALLIUM.
Cerasus, <i>Tourn</i>	Cherry.....	PRUNUS.
Ceratocephaloides, <i>Vail.</i> A. G.		VERBESINA.
Ceratocephalus, <i>Vaill.</i> A. G.		BIDENS.
Ceratoides, <i>Tourn. Cor</i>		AXYRIS.
Cereus, <i>Juss.</i> A. G. ...	Torch Thistle.....	CACTUS.
Cerinthoides, <i>Boerh</i>	Honeywort	CERINTHE.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
<i>Cervispina, Dill. gen.</i>	Buckthorn.....	RHAMNUS.
<i>Chærophylli species, Tourn.</i>	Wild Chervil.....	CHÆROPHYL- LUM.
<i>Chamæbuxus, Tourn.</i>	Low Box.....	POLYGALA.
<i>Chamæcerasus, Tourn.</i>	Dwarf Cherry, or Upright Honeysuckle.	LONICERA.
<i>Chamædaphne, Buxh.</i>		ANDROMEDA.
A. R.		
<i>Chamædaphne, Mitch.</i>		MITCHELLA.
<i>Chamædrys, Tourn.</i>	Germander.....	TEUCRIUM..
<i>Chamæjasme, Amm.</i>		STELLERA.
<i>Chamælea, Tourn.</i>	Widow Wail.....	CNEORUM.
<i>Chamælinum, Vaill.</i>	Least Rupture-wort, or	LINUM.
B. P. All-seed.		
<i>Chamæmelum, Tourn.</i>	Chamomile.....	ANTHEMIS.
<i>and Vaill.</i> A. G.		
<i>Chamænerion, Tourn.</i>	Rosebay, or Willow Herb...	EPILOBIUM.
<i>Chamæpitys, Tourn.</i>	Ground Pine.....	TEUCRIUM.
<i>Chamærhododendros, Tourn.</i>	Dwarf Rosebay.....	RHODODEN- DRON.
<i>Chamæriphe, Pont.</i>	Dwarf Palm.....	CHAMÆROPS.
<i>Chenopodio-morus, Boer.</i>	Strawberry Spinach, or	BLITUM.
Blite.		
<i>Christophoriana, Tourn.</i>	Herb Christopher.....	ACTEA.
<i>Chrysanthemoides, A. G. Dill. gen. & Elth.</i>	Hard-seeded Chrysanthe- mum.	OSTEOSPER- MUM.
<i>Chrysocome, Dill. gen.</i>	Golden Locks.....	CHRYSOCOMA.
<i>Cicuta, Tourn.</i>	Hemlock.....	CONIUM.
<i>Cicutaria, Tourn.</i>	Great broad-leaved Ba- stard Hemlock.	LIGUSTICUM.
<i>Cinara, Tourn.</i>	Artichoke.....	CYNARA.
<i>Cinnamomum, L. B. and Burm. Zeyl.</i>	Herm. H. Cinnamon Tree:	LAURUS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
<i>Cirsium, Tourn. and Vaill. A. G.</i>	Soft, or Gentle Thistle	CARDUUS.
<i>Citreum, Tourn.</i>	Citrøn	CITRUS.
<i>Clandestina, Tourn.</i>	Broom Rape, with great purple flowers; or, Great purple Herb- bane.	LATHRÆA.
<i>Clematitis, Tourn.</i>	Virgin's Bower	CLEMATIS.
<i>Clitorius, Dill. Elth.</i>		CLITÓRIA.
<i>Clymenum, Tourn.</i>	Chichling Vetch	LATHYRUS.
<i>Coa, Plum.</i>		HIPPOCRATEA.
<i>Codda Panna, Hort. Mal.</i>		CORYPHA.
<i>Coffè, Juss. A. G.</i>	Coffee Tree	COFFEA.
<i>Colocasia, Boerh.</i>	Great Egyptian Arum	ARUM.
<i>Colocynthis, Tourn.</i>	Coloquintida, or Bitter Gourd.	CUCUMIS.
<i>Coma aurea, Boerh.</i>	Golden Locks	CHRYSOCOMA.
<i>Conocarpodendron, Boerh.</i>	Silver Tree	PROTEA.
<i>Convolvulo Tithymalus, Boerh.</i>		DALECHAMPIA.
<i>Conyzella, Dill. gen.</i>		ERIGERON.
<i>Conyzoides, Dill. gen.</i>		ERIGERON.
<i>Conyzoides, Tourn. A. G.</i>		CARPESIUM.
<i>Coral, Dill. Elth.</i>	Coral Tree	ERYTHRINA.
<i>Corallo fungus, Vaill. B. P.</i>		CLAVARIA.
<i>Corallodendron, Tourn.</i>	Coral Tree	ERYTHRINA.
<i>Coralloides, Tourn. and Mich.</i>		CLAVARIA.
<i>Coralloides, Dill. Musc.</i>	Liverwort	LICHEN.
<i>Cordyline, Roy. Lugd.</i>	Adam's Needle	YUCCA.
<i>Corindum, Tourn.</i>	Heart-seed, or Heart-pea	CARDIOSPER- MUM.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Cornucopiodes, <i>Scheuch</i>		CORNUCOPIE.
Corona imperialis, <i>Tour.</i>	Crown Imperial.....	FRITILLARIA.
Corona solis, <i>Vaill.</i> A. G...	Sunflower.....	HELIANTHUS.
<i>Tourn. & Dill. Elth.</i>		
Coronopus, <i>Tourn</i>	Buck's-horn Plantain.....	PLANTAGO.
Corrigiola, <i>Dill. gen.</i> ...	Verticillate Knot-grass... <i>and Mehr.</i>	ILLECEBRUM.
Cortusa, <i>Plum</i>		THALIA.
Corydalis, <i>Dill. gen</i>	Bladder Fumatory.....	FUMARIA.
Cotinus, <i>Tourn</i>	Venice Sumach.....	RHUS.
Cotula, <i>Tourn</i>		ANACYCLUS.
Courbaril, <i>Plum</i>	Locust Tree.....	HYMENEÆ.
Crepis, <i>Vaill.</i> A. G.....	Tangier Sow-Thistle.....	SCORZONERA.
Crocodilium, <i>Vaill.</i>	Centaury without stems...	CENTAUREA.
Crocoddilodes, <i>Vaill.</i>	Distaff Thistle.....	ATRACYLIS.
Cruciata, <i>Tourn</i>	Crosswort.....	VALANTIA.
Cucularia, <i>Juss.</i> A. G...	Fumatory with a naked... stalk.	FUMARIA.
Cujete, <i>Plum</i>	Calabash Tree.....	CRESCENTIA.
Cuminoides, <i>Tourn</i>	Wild or Bastard Cumir ...	LAGOECIA.
Cururu, <i>Plum</i>		PAULLINIA.
Cyanus, <i>Tourn. and</i> ...	Bluebottle.....	CENTAUREA.
<i>Vaill.</i> A. G.		
Cyathoides, <i>Mich</i>	Cup Mushroom.....	PEZIZA.
Cydonia, <i>Tourn</i>	Quince Tree.....	PYRUS.
Cynocrambe, <i>Tourn</i>	Dog's Cabbage.....	THELIGONUM.
Cynoglossoides, <i>Isnard</i> ...	Borragé.....	BORRAGO.
A. G.		
Cynomorium, <i>Garc</i>		CYNOMETRA.
Cynorrhinchium, <i>Mitch</i>		MIMULUS.
Cyperella, <i>Mich</i>		SCHÆNUS.
Cyperoides, <i>Tour. Scheu</i>		CAREX.
<i>and Mich.</i>		
Cysticapnos, <i>Boerh</i>	Bladder Fumatory.....	FUMARIA.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
D		
<i>Dalea, Lin. gen. pl. ed. prim.</i>		PSORALEA.
<i>Damasonium, Tourn. and Vaill. A. G.</i>	Star-headed Water Plantain.	ALISMA.
<i>Dantia, Petit. gen.</i>		ISNARDIA.
<i>Dens Canis, Tourn.</i>	Dog's-Tooth Violet.	ERYTHRONIUM.
<i>Dens Leonis, Tourn.</i>	Dandelion.	LEONTODON.
<i>Dichotophyllum, Dill. gen.</i>		CERATOPHYLLUM.
<i>Diconangia, Mich.</i>		ITEA.
<i>Dimorphotheca, Vaill. A. G.</i>	Marigold.	CALENDULA.
<i>Diototheca, Vaill. A. G.</i>		MORINA.
<i>Dodonæa, Plum.</i>	Holly, with winged leaves	ILEX.
<i>Doria, Dill. gen. & Elth.</i>	Golden Rod.	SOLIDAGO.
<i>Dortmanna, Rudb. A. S.</i>	Water Gladiole.	LOVELIA.
<i>Dracunculoides, Boerh.</i>	Blood-Flower.	HÆMANTHUS.
<i>Dracunculus, Tourn.</i>	Dragons.	ARUM.
<i>Duglassia, Houst. A. A.</i>		VOLKAMERIA.
E		
<i>Echinopus, Tourn. and Vaill. A. G.</i>	Globe Thistle.	ECHINOPS.
<i>Echinoides, Dill. gen.</i>		LYCOPSIS.
<i>Elate, Mus. Cliff.</i>	Common Palm, or Date Tree	PHENIX.
<i>Elaterium, Boerh.</i>	Wild, Spirting, or Ass's Cucumber.	MOMORDICA.
<i>Elatine, Dill. gen.</i>	Fluellin, or Female Speedwell.	ANTIRRHINUM.
<i>Elephas, Tourn.</i>	Elephant's Head.	RHINANTHUS.
<i>Elichrysum, Tourn. and Dill. Elth.</i>	Cassidony, Golden-locks, or Eternal Flower.	GNAPHALIUM.
<i>Elymus, Mich.</i>		ZIZANIA.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
<i>Emerus, Tourn</i>	Scorpion Senna.....	CORONILLA.
<i>Enula, Cesalp. and Magnol.</i>	Elecampane.....	INULA.
<i>Ephemerum, Tourn</i>	Virginian Spiderwort.....	TRADESCANTIA
<i>Erebinthus, Mitch</i>		VICIA.
<i>Eresia, Plum</i>		THEOPHRASTA.
<i>Ericæ species, Tourn</i>		ANDROMEDA.
<i>Erinacea, Tourn</i>	Spanish Hedgehog Thorn	ANTHYLLIS.
<i>Erinaceus, Dill. & Mich</i>		HYDNUM.
<i>Eriocephalus, Vaill. A.G.</i>	Spear Thistle.....	CARDUUS.
<i>Eriophorus, Vaill. A.G.</i>	Downy Sow-Thistle, or... Woolly Hawk-weed.	ANDRYALA.
<i>Erucago, Tourn</i>	Square-codded Rocket..... of Montpellier.	BUNIAS.
<i>Euonymoides, Isnar. A.G.</i>	Staff Tree.....	CELASTRUS.
<i>Eupatoriophalacron, Dill. Elth. and Vaill. A. G.</i>		VERBESINA.
<i>Euphorbium, Isnar. A.G.</i>	Burning Thorny Plant.....	EUPHORBIA.
F		
<i>Faba, Tourn</i>	Bean.....	VICIA.
<i>Fabago, Tourn</i>	Bean Caper.....	ZYGOPHYLLUM
<i>Fagopyrum, Tourn</i>	Buck Wheat, or Brank.....	POLYGONUM.
<i>Ferrum equinum, Tour.</i>	Horseshoe Vetch.....	HIPPOCREPIS.
<i>Ficaria, Dill. gen.</i>	Pilewort, or Lesser Ce- landine.	RANUNCULUS.
<i>Ficoida, Niss. A. G. Dill. gen. and Elth.</i>		AIZOON.
<i>Ficoides, Tourn. A. G.</i>	Fig Marigold.....	MESEMBRYAN- THEMUM.
<i>Filago, Vaill. A. G. &... Tourn.</i>	Cudweed.....	GNAPHALIUM.
<i>Filipendula, Tourn</i>	Dropwort.....	SPIRÆA.
<i>Fluvialis, Vaill. A.G. and Mich.</i>		NAIAS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
<i>Fœniculum, Tourn.</i>	Fennel.....	ANETHUM.
<i>Fœnum Græcum, Tourn.</i>	Fenugreek.....	TRIGONELLA.
<i>Franca, Mich.</i>		FRANKENIA.
<i>Frangula, Tourn.</i>	Black or Berry-bearing... Alder.	RHAMNUS.
<i>Fungoidaster, Mich.</i>		ELVELA.
<i>Fungoides, Mich.</i>		ELVELA.
<i>Fungoides, Dill.</i>		CLAVARIA.
<i>Fungoidis species, Vaill. B. P.</i>	Cup Mushroom.....	PAZIZA.
<i>Fungoidis species, Vail. B. P.</i>		ELVELA.
G		
<i>Gale, Tourn. A. G. & Dill. gen.</i>	Sweet Willow, Gale, or... Dutch Myrtle.	MYRICA.
<i>Galeobdolon, Dill. gen.</i>	Yellow Archangel, or... Dead Nettle.	GALEOPSIS.
<i>Galeopsis, Tourn.</i>	Base Horehound.....	STACHYS.
<i>Gallium, Tourn.</i>	Ladies' Bed-straw, or... Cheese Rennet.	GALIUM.
<i>Geaster, Mich.</i>		LYCOPERDON.
<i>Genista, Tourn.</i>	Broom.....	SPARTIUM.
<i>Genista-spartium, Tour.</i>	Furze, Whins, or Gorse...	ULEX.
<i>Genistella, Tourn.</i>	Dwarf Broom.....	GENISTA.
<i>Gerbera, Lin. gen. pl. ed. prim.</i>		ARNICA.
<i>Gesnera, Plum.</i>		GESNERIA.
<i>Geum, Tourn.</i>	Kidneywort.....	SAXIFRAGA.
<i>Glaucium, Tourn.</i>	Horned Poppy.....	CHELIDONIUM.
<i>Glaucoides, Mich.</i>	Water Purslane.....	PEPLIS.
<i>Gnaphaloides, Tourn.</i>	Bastard Cudweed.....	MICROPUS.
<i>Graminifolia, Dill. gen.</i>	Triple-headed Pond-weed...	ZANNICHELLIA.
<i>Granadilla, Tourn. & Dill. Elth.</i>	Passion Flower.....	PASSIFLORA.
<i>Grossularia, Tourn.</i>	Gooseberry.....	RIBES.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
Guaicana, <i>Tourn</i>	Indian Date-Plum.....	DIOSPYROS.
Guaiava, <i>Tourn</i>	Bay Plum.....	PSIDIUM.
Guanabanus, <i>Plum</i>	Custard-Apple.....	ANNONA.
Guazuma, <i>Plum</i>	Bastard Cedar of Jamaica	THEOBROMA.
Guidonia, <i>Plum</i>		SAMYDA.
H		
Hacub, <i>Vaill. A. G.</i>		GUNDELIA.
Harmala, <i>Tourn</i>	Wild Syrian Rue.....	PEGANUM.
Hedypnois, <i>Tourn</i>		HYOSERIS.
Heisteria, <i>Lin. gen. pl.</i>		POLYGALA.
	<i>ed. prim.</i>	
Heleniastrum, <i>Vail. A. G.</i>	Bastard Sunflower.....	HELENIA.
Helenium, <i>Vail. A. G.</i>	Starwort.....	ASTER.
Helenium, <i>Moris. Raj.</i>	Elecampane.....	INULA.
	<i>Herm Rivin. Rupp.</i>	
	<i>Knaut. and Vaill.</i>	
Helianthemum, <i>Tourn</i> ...	Dwarf Cistus, or Little... Sunflower.	CISTUS.
Helichrysoides, <i>Vaill</i>		SERIPHIMUM.
	A. P.	
Helichrysoides, <i>Vaill</i>		GNAPHALIUM.
	A. G.	
Helichrysum, <i>Vail. A. G.</i>	Cassidony, Golden-locks,... or Eternal Flower.	GNAPHALIUM.
Helleborine, <i>Tourn</i>	Bastard Hellebore.....	SERAPIAS.
Helminthotheca, <i>Vaill</i>		PICRIS.
	A. G.	
Helxine, <i>Lin. gen. pl.</i> ...	Buck-wheat, or Brank....	POLYGONUM.
	<i>ed. prim.</i>	
Henna, <i>Ludw</i>		LAWSONIA.
Hepatica, <i>Dill. gen</i>	Noble Liverwort, or He- patica.	ANEMONE.
Hepatica, <i>Mich</i>		MARCHANTIA.
Herba Paris, <i>Tourn</i>	True-love, or One-berry	PARIS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
<i>Hermodactylus</i> , <i>Tourn.</i>	Tuberose Iris.....	IRIS.
<i>Hieracioides</i> , <i>Vaill.</i>	A.G. Bastard Hawkweed.....	CREPIS.
<i>Hippocastanum</i> , <i>Tourn.</i>	Horse Chestnut.....	ÆSCULUS.
<i>Hippuris</i> , <i>Dill. gen. &</i>	CHARA.
<i>Pont. Anth.</i>		
<i>Horminum</i> , <i>Tourn.</i>Clary.....	SALVIA.
<i>Hyacinthus stellaris</i> ,.....	Star Hyacinth.....	SCILLA.
<i>Raj. Meth.</i>		
<i>Hydroceratophyllon</i> ,.....		CERATOPHYLL- LUM.
<i>Vaill. A. G.</i>		
<i>Hydrophace</i> , <i>Buxb. cent.</i>	Duck-meat.....	LEMNA.
<i>Hypericoides</i> , <i>Plum.</i>St. Peter's Wort.....	ASCYRUM.
<i>Hypocistis</i> , <i>Tourn.</i>Rape of Cistus.....	ASARUM.
<i>Hypophyllocarpoden-</i>	PROTEA.
<i>dron</i> , <i>Boerh.</i>		
<i>Hypopitys</i> , <i>Dill. gen.</i>	MONOTROPA.
<i>Hysterophorus</i> , <i>Vaill.</i>Bastard Feverfew.....	PARTHENIUM.
A. G.		

I

<i>Jabotapita</i> , <i>Plum.</i>	OCHNA.
<i>Jacea</i> , <i>Tourn. Dill. gen.</i>Knapweed.....	CENTAUREA,
<i>and Vaill.</i>		
<i>Jacobææ species</i> , <i>Tour.</i>Ragworts (sundry, of old... authors).	SOLIDAGO.
<i>Vaill. A. G.</i>		
<i>Jacobææ species</i> , <i>Tour.</i>Ragworts (sundry, of old... authors).	SENECIO.
<i>Jacobæastrum</i> , <i>Vaill.</i>African Ragwort.....	OTHONNA.
A. G.		
<i>Jacobæoides</i> , <i>Vail.</i>	A.G. African Ragwort.....	OTHONNA.
<i>Jalapa</i> , <i>Tourn.</i>Marvel of Peru.....	MIRABILIS.
<i>Jan-raja</i> , <i>Plum.</i>	RAJANIA.
<i>Jasminoides</i> , <i>Niss.</i>	A. G. Bastard Jasmine.....	LYCIUM.
<i>Ïcaco</i> , <i>Plum.</i>Cocoa Plum.....	CHRYSOBALA- NUS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
<i>Ilex, Tourn.</i>	Evergreen Oak.....	QUERCUS.
<i>Indigo, Isnard, A. G.</i> ...	Goat's Rue.....	GALEGA.
<i>Inga, Plum.</i>	MIMOSA.
<i>Jonthlaspi, Tourn.</i>	Treacle Mustard.....	CLYPEOLA.
<i>Isora, Plum.</i>	Screw Tree.....	HELICTERES.
<i>Juncago, Tourn. & Mich.</i>	Arrow-headed Grass.....	TRIGLOCHIA.
<i>Jussievia, Houst. A. A.</i>	JATROPHA.

K

<i>Kali, Tourn.</i>	Glasswort.....	SALSOLA.
<i>Karatas, Plum.</i>	Pine-apple.....	BROMELIA.
<i>Katovindel, Hort. Mal.</i>	Palm, or Date Tree.....	PHENIX.
<i>Kæmpfera, Houst. A. A.</i>	Vervain.....	VERBENA.
<i>Keratophyton, Boerh.</i>	LITHOXYLUM.
<i>Ketmia, Tourn.</i>	Althæa Frutex, or Syrian... Mallow.	HIBISCUS.

Kleinia, Lín. gen. pl.... Foreign Colt's-foot.....CACALIA.
ed. prim.

Knawel, Dill. gen.... ..German Knot-grass.....SCLERANTHUS.
Kodda-pail, Plum......Water Houseleek of Egypt PISTIA.

L

<i>Lacryma Job, Tourn.</i>	Job's Tears.....	COIX.
<i>Lampsana, Vaill. A. G.</i>	Nipplewort.....	LAPSANA.
<i>LANCISIA, Pont. diss.</i>	COTULA.
<i>Lapathum, Tourn.</i>	Dock.....	RUMEX.
<i>Lappa, Tourn. & Vail.</i> ...	Burdock.....	ARCTIUM.

A. G.

<i>Larix, Tourn.</i>	Larch Tree.....	PINUS.
<i>Laurentia, Mich.</i>	LOBELIA.
<i>Laurocerasus, Tourn.</i> ...	Laurel.....	PRUNUS.
<i>Ledum, Mich.</i>	ANDROMEDA.
<i>Lens, Tourn.</i>	Lentils.....	ERVUM.
<i>Lentibularia, Vaill. A.</i> ...	Water Milfoil.....	UTRICULARIA.

G. and Dill. gen.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
Lenticula, <i>Mich. and...</i> <i>Dill. gen.</i>	Duck-meat.....	LEMNA.
Leontodontoides, <i>Mich.</i> <i>gen.</i>	HYOSERIS.
Leontopetalon, <i>Tourn.</i>	Lion's Leaf.....	LEONTICE.
Lepidocarpodendron, <i>Boerh.</i>	PROTEA.
Leptostachia, <i>Mich.</i>	PHRYMA.
Leucanthemum, <i>Tourn.</i>	Chrysanthemum with... white rays, or Ox- eye Daisy.	CHRYSANTHE- MUM.
Leucojum, <i>Tourn.</i>	Stock July-Flower, and... Wall Flower.	CHEIRANTHUS,
Lichen, <i>Dill. Musc.</i>	MARCHANTIA,
Lichenastrum, <i>Dill.</i> <i>Musc.</i>	JUNGERMAN- NIA.
Lichenoides, <i>Dill. Musc.</i>	LICHEN.
Lilac, <i>Tourn.</i>	Lilac, or Pipe Tree.....	SYRINGA.
Liliastrum, <i>Tourn.</i>	White Day Lily, St. Bru- no's Lily, or Great Sa- voy Spiderwort.	HEMEROCAL- LIS.
Lilio-asphodelus, <i>Tourn.</i>	Day Lily, or Lily Aspho- del.	HEMEROCAL- LIS.
Lilio-hyacinthus, <i>Tourn.</i>	Lily-Hyacinth.....	SCILLA.
Lilionarcissus, <i>Tourn.</i>	Lily-Daffodil.....	AMARYLLIS.
Lilium convallium, <i>Tourn.</i>	Lily of the Valley.....	CONVALLARIA.
Limnopoce, <i>Vaill. A.G.</i>	HIPPURIS.
Limodorum, <i>Tourn.</i>	Purple Bird's-Nest.....	ORCHIS.
Limon, <i>Tourn.</i>	Lemon.....	CITRUS.
Limonium, <i>Tourn.</i>	Sea Lavender.....	STATICE.
Linagrostis, <i>Mich. &</i> <i>Tourn.</i>	Cotton Grass.....	ERIOPIHORUM.
Linaria, <i>Tourn.</i>	Toad Flax.....	ANTIRRHINUM.
Lingua cervina, <i>Tourn.</i>	Hart's Tongue.....	ASPENIUM.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Linocarpon, <i>Mich</i>	Least Rupturewort, or... All-seed.	LINUM.
Lirium, <i>Roy</i>	Lily.....	LILIUM.
Lithophyton, <i>Tourn</i>		LITHOXYLON.
Lonchitis, <i>Tourn</i>	Rough Spleenwort.....	POLYPODIUM.
Luffa, <i>Tourn. A.G. Dill</i> ...	Egyptian Cucumber..... <i>gen. and Elth.</i>	MOMORDICA.
Lunularia, <i>Mich</i>		MARCHANTIA.
Lupinaster, <i>Buxb</i>		TRIFOLIUM.
Lupulus, <i>Tourn</i>	Hop.....	HUMULUS.
Luteola, <i>Tourn</i>	Wild Woad, or Dyer's... Weed.	RESEDA.
Lychnidea, <i>Dill. Elth</i> ...	Bastard Lychnis.....	PHLOX.
Lychni scabiosa, <i>Boerh</i>		KNAUTIA.
Lycogala, <i>Mich</i>		MUCOR.
Lycoperdastrum, <i>Mich</i>		LYCOPERDON.
Lycoperdoides, <i>Mich</i>		LYCOPERDON.
Lycopersicon, <i>Tourn</i> ...	Wolf's Peach, or Love... Apple.	SOLANUM.
Lycopodioides, <i>Dill. Musc</i>		LYCOPODIUM.

M

Malachodendron, <i>Mitch</i>		STEWARTIA.
Malacoides, <i>Tourn</i>	Bastard Mallow.....	MALOPE.
Malva, <i>Tourn</i>	Rose Mallow, or Holly-... hock.	ALCEA.
Malvaviscus, <i>Dill. Elth.</i>	Berry-bearing Hibiscus...	HIBISCUS.
Malvinda, <i>Dill. Elth</i> ...	Indian Mallow, with sin-... gle Seeds.	SIDA.
Malus, <i>Tourn</i>	Apple.....	PYRUS.
Mamei, <i>Plum</i>	Mammee.....	MAMMEA.
Mancanilla, <i>Plum</i>	Manchineel.....	HIPPOMANE.
Mangles, <i>Plum</i>	Pee-kaudel of the Indians	RHIZOPHORA.
Mangostans, <i>Garc. A.A.</i>	Mangostan.....	GARCINIA.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Manihot, <i>Tourn. and...</i> <i>Dill. Elth.</i>	Cassava.....	JATROPHA.
Maurocena, <i>Lin. gen...</i> <i>pl. ed. prim.</i>	Hottentot Cherry.....	CASSINE.
Mays, <i>Tourn.....</i>	Indian or Turkey Wheat.	ZEA.
Medica, <i>Tourn.....</i>	Snail Trefoil, and Medic... or Lucern Grass.	MEDICAGO.
Melanoschœnus, <i>Mich...</i> <i>gen.</i>	Round Black-headed..... Marsh-Rush, or Bog Rush.	SCHŒNUS.
Melilobus, <i>Mitch.....</i>	Three-thorned Acacia.....	GLEDITSIA.
Melilotus, <i>Tourn.....</i>	Melilot.....	TRIFOLIUM.
Melo, <i>Tourn.....</i>	Melon.....	CUCUMIS.
Melocactus, <i>Tourn.....</i>	Melon Thistle.....	CACTUS.
Melongena, <i>Tourn.....</i>	Mad Apple, or Egg Plant	SOLANUM.
Melopepo, <i>Tourn.....</i>	Buckler Gourd.....	CUCURBITA.
Memecylum, <i>Mich.....</i>	Trailing Arbutus.....	EPIGEA.
Methonica, <i>Tourn.....</i>	Superb Lily.....	GLORIOSA.
Meum, <i>Tourn.....</i>	Spignel.....	ATHAMANTA.
Michelia, <i>Houst. A. A.....</i>		PONTERIA.
Michelia, <i>Amm. Act. Pet.....</i>		GMELINA.
Microleuconymphaea, ... <i>Boerh.</i>	Frog's Bit.....	HYDROCHARIS.
Millefolium, <i>Tourn.....</i>	Yarrow, or Milfoil.....	ACHILLEA.
Mitra, <i>Houst.....</i>		OPHIORRHIZA.
Mitreola, <i>Lin. gen. pl.....</i> <i>ed. prim.</i>		OPHIORRHIZA.
Moldavica, <i>Tourn.....</i>	Turkey or Moldavian..... Baum.	DRACOCEPHALUM.
Molle, <i>Tourn.....</i>	Peruvian Mastich.....	SCHINUS.
Molucca, <i>Tourn.....</i>	Molucca Baum.....	MOLUCELLA.
Moly, <i>Boerh.....</i>	Moly with Lily Flowers, ... or Homer's Moly.	ALLIUM.
Monbin, <i>Plum.....</i>	Brasilian Plum.....	SPONDIAS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
Monilifera, <i>Vaill.</i>	A. G. -Hard-seeded Chrysan- themum.	OSTEOSPER- MUM.
Monospermalthæa, <i>Isnar</i>		WALTHERIA.
A. G.		
Montia, <i>Houst.</i>	A. A.....	HELIOCARPUS.
Morocarpus, <i>Rupp</i>	Blite, or Strawberry Spinach	BLITUM.
Morsu ranæ, <i>Tour.</i>	A. G. Frog's Bit.....	HYDROCHARIS.
Moschatellina, <i>Tourn</i> ...	Tuberose Moschatel, or... Hollow Root.	ADOXA.
Mucilago, <i>Mich</i>		MUCOR.
Murucuja, <i>Tourn</i>	Passion Flower.....	PASSIFLORA.
Muscari, <i>Tourn</i>	Grape Hyacinth.....	HYACINTHUS.
Muscoides, <i>Mich</i>		JUNGERMANNIA
Myosotis, <i>Tourn</i>	Mouse-ear Chickweed.....	CERASTIUM.
Myosuros, <i>Dill. gen</i>	Mouse-tail.....	MYOSURUS.
Myrobatindum, <i>Vaill</i> ...	American Viburnum.....	LANTANA.
A. G.		

N

Narcisso-Leucojum,.....	Greater Snow-drop.....	LEUCOJUM.
<i>Tourn.</i>		
Nasturtium, <i>Tourn</i>	Cress.....	LEPIDIUM.
Nelumbo, <i>Tourn</i>	Indian Water-Lily.....	NYPHÆA.
Nhandiroba, <i>Plum</i>		FEVILLEA.
Ninsi, <i>Breyn. diss</i>	Ginseng.....	PANAX.
Nummularia, <i>Nov. gen</i>		HOLOSTEUM.
Nux, <i>Tourn. and Boerh.</i>	Walnut.....	JUGLANS.
Nymphoides, <i>Tourn</i>	Lesser Yellow Water Lily, ... with fringed flowers.	MENYANTHES.

O

Obeliscotheca, <i>Vaill</i>	Dwarf Sunflower.....	RUDBECKIA.
A. G. and <i>Dill. Elth.</i>		
Ochrus, <i>Tourn</i>	Wildwinged Pea.....	PISUM.
Odontitis, <i>Dill. gen</i>	Red Meadow-Eyebright...	EUPHRASIA.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
Omphalodes, <i>Tourn.</i>	Venus's Navelwort.....	CYNOGLOSSUM.
Onagra, <i>Tourn.</i>	Tree Primrose.....	CENOTHERA.
Onobrychis, <i>Tourn.</i>	Cock's Head, or Sainfoin...	HEPYSARUM.
Ophris, <i>Tourn.</i>	Twyblade.....	OPHRYS.
Opulus, <i>Tourn. & Vaill.</i> ...	Marsh Elder, or Gelder...	VIBURNUM.
A. G.	Rose.	
Opuntia, <i>Tourn.</i>	Indian Fig, or Prickly Pear	CACTUS.
Orchidion, <i>Mitch.</i>		ARETHUSA.
Oreoselinum, <i>Tourn.</i>	Mountain Parsley.....	ATHAMANTA.
Ornithopodium, <i>Tourn.</i>	Bird's Foot.....	ORNITHOPUS.
Ornus, <i>Mich.</i>	Ash.....	FRAXINUS.
Orobanchoides, <i>Tourn.</i>		MONOTROPA,
A. G.		
Ostrya, <i>Mich.</i>	Hornbeam.....	CARPINUS.
Oxycoccus, <i>Tourn.</i>	Marsh Whortleberries, ...	VACCINIUM.
	Moss Berries, or Moor	
	Berries.	
Oxyoides, <i>Garc. A. A.</i>	Sensitive Wood-Sorrel.....	OXALIS.
Oxys, <i>Tourn.</i>	Wood Sorrel.....	OXALIS.
P		
Padus, <i>Lin. gen. pl.</i>	Bird Cherry.....	PRUNUS.
<i>ed. prim.</i>		
Paliurus, <i>Tourn.</i>	Christ's Thorn.....	RHAMNUS,
Panacea, <i>Mitch.</i>	Ginseng.....	PANAX.
Panicastrella, <i>Mich.</i>		CENCHRUS.
Papaya, <i>Tourn.</i>	Papaw.....	CARICA.
Papia, <i>Mich.</i>		ORVALA.
Paronychia, <i>Tourn.</i>	Mountain Knot-grass.....	ILLECEBRUM.
Partheniastrum, <i>Niss.</i> ...	Bastard Feverfew.....	PARTHENIUM.
A. G. <i>Dill. gen. & Elth.</i>		
Patagonica, <i>Dill. Elth.</i>		PATAGONULA.
Pavia, <i>Boerh.</i>	Scarlet Horse-chestnut.....	ÆSCULUS.
Pedicularis species, <i>Tour.</i>	Yellow Rattle, Cock's-...	RHINANTHUS.
	comb, or Lousewort.	

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
<i>Pelecinus, Tourn</i>	Clusius's Foreign Mat- chet-Vetch.	<i>DESMODIUM</i> .
<i>Persea, Plum</i>	Tree Milk-tree, with rough tree-bark.	<i>LAURUS</i> .
<i>Pentagonoloba, Vaill.</i> A. G.	<i>DESMODIUM</i> .
<i>Pentaplyllum, Tourn</i>	Cinquefoil, whose leaves are not all to quill.	<i>POTENTILLA</i> .
<i>Pentastachyllum, Dill. gen.</i>	Water Milk-tree.....	<i>DIODAPHYL- LUM</i> .
<i>Pepo, Tourn</i>	Pumpkin.....	<i>CUCURBITA</i> .
<i>Percepiet, Dill. gen.</i>	Parsley Piert.....	<i>APHANIS</i> .
<i>Perekin, Plum. Lin.</i> <i>gen. pl. et prin.</i>	Gooseberry of the Ame- ricans, or Blad Apple.	<i>CACTUS</i> .
<i>Pericyanenum, Tourn</i>	Trumpet Honeysuckle.....	<i>LONICERA</i> .
<i>Persea, Plum</i>	Avocado or Avogato Pear	<i>LAURUS</i> .
<i>Persica, Tourn</i>	Peach.....	<i>AMYGDALUS</i> .
<i>Persicaria, Tourn</i>	Arse-smart, or Persicaria	<i>POLYGONUM</i> .
<i>Pervinca, Tour</i>	Periwinkle.....	<i>VINCA</i> .
<i>Petasis, Tourn. and</i> <i>Vaill. A. G.</i>	Butterburr, or Pestilent- wort.	<i>TUSSILAGO</i> .
<i>Petilium, Lin. gen. pl.</i> <i>ed. prim.</i>	Crown Imperial.....	<i>FRITILLARIA</i> .
<i>Phalangium, Tourn</i>	Spiderwort.....	<i>ANTHERICUM</i> .
<i>Phalloboletus, Mich</i>	<i>PHALLUS</i> .
<i>Phillyreastrum, Vaill.</i> A. G.	<i>MORINDA</i> .
<i>Pilosella, Vaill. A. G.</i>	Creeping Mouse-ear.....	<i>HIERACIUM</i> .
<i>Pimpinella, Tourn</i>	Burnet.....	<i>POTERIUM</i> .
<i>Pinastella, Dill. gen.</i>	<i>HIPPURIS</i> .
<i>Pinguin, Dill. Eth.</i>	Wild Ananas.....	<i>BROMELIA</i> .
<i>Pittonia, Plum</i>	<i>TOURNEFORTIA</i> .
<i>Plantaginella, Dill. gen.</i>	Least Water-Plantain.....	<i>LIMOSELLA</i> .
<i>Plantanocephalus, Vaill.</i> A. G.	Button-wood.....	<i>CEPHALANTHUS</i> .

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
<i>Poliifolia</i> , <i>Burb.</i> A. R.	Marsh Cistus, or Rosemary	ANDROMEDA.
<i>Polium</i> , <i>Tourn.</i>	Poley Mountain.....	TEUCRIUM.
<i>Polyacantha</i> , <i>Vaill.</i>	Casaubon's Thistle, sup- posed the true Fish Thistle or Acarna of Theophrastus.	CARDUUS.
<i>Polygaloides</i> , <i>Dill. gen.</i>	Milkwort.....	POLYGALA.
<i>Polygonatum</i> , <i>Tourn.</i>	Solomon's Seal.....	CONVALLARIA.
<i>Polygonifolia</i> , <i>Dill. gen.</i>		CORRIGIOLA.
<i>Polygonoides</i> , <i>Tourn.</i>		CALLIGONUM.
<i>Polyporus</i> , <i>Mitch.</i>		BOLETUS.
<i>Populago</i> , <i>Tourn.</i>	Marsh Marigold.....	CALTHA.
<i>Porophyllum</i> , <i>Vaill.</i>	Cacalin, with perforate leaves.	CACALIA.
<i>Porrum</i> , <i>Tourn.</i>	Leek.....	ALLIUM.
<i>Portula</i> , <i>Dill. gen.</i>	Water Purslane.....	PEPLIS.
<i>Portulacastrum</i> , <i>B. Jus.</i>	Horse Purslane.....	TRIANTHEMA.
<i>Potamopithys</i> , <i>Burb.</i> A. R.		ELATINE.
<i>Primula veris</i> , <i>Tourn.</i> ...	Primrose.....	PRIMULA.
<i>Provenzalia</i> , <i>Petit. Gen.</i>	Water Dragons.....	CALLA.
<i>Pseudoacacia</i> , <i>Tourn.</i> ...	False Acacia.....	ROBINIA.
<i>Pseudocyperus</i> , <i>Mich.</i>		SCHÆNUS.
<i>Pseudodictamnus</i> , <i>Tour.</i>	Bastard Dittany.....	MARRUBIUM.
<i>Pseudoruta</i> , <i>Mich.</i>	Three-leaved Rue.....	RUTA.
<i>Psyllium</i> , <i>Tourn.</i>	Fleawort.....	PLANTAGO.
<i>Ptarmica</i> , <i>Tourn.</i>	Sneezewort, Bastard Pel- litory, or Goose-tongue.	ACHILLEA.
<i>Pterocephalus</i> , <i>Vaill.</i>	Scabious.....	SCABIOSA.
A. G.		
<i>Pterospermadendron</i> , <i>Am.</i>		PENTAPETES.
<i>Pulsatilla</i> , <i>Tourn.</i>	Pasque Flower.....	ANEMONE.
Q		
<i>Quamoclit</i> , <i>Tourn.</i>		IPOMŒA.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Quinquefolium, <i>Tourn</i> ...	Cinquefoil.....	POTENTILLA.
Quinquina, <i>Condam</i> A. G.	True Jesuits'-Bark Tree....	CINCHONA.
R		
Radicula, <i>Dill. gen</i>	Water Radish.....	SISYMBRIUM.
Radiola, <i>Dill. gen</i>	Least Rupturewort, or..... All-seed.	LINUM.
Ranunculoides, <i>Va. A.G.</i>	Water Crowfoot.....	RANUNCULUS.
Rapa, <i>Tourn</i>	Turnep.....	BRASSICA.
Raphanistrum, <i>Tourn</i> ...	White-flowered Char-..... lock, with jointed pods.	RAPHANUS.
Rapistrum, <i>Tourn</i>	Sea Cabbage.....	CRAMBE.
Rapunculus, <i>Tourn</i>	Rampions.....	PHYTEUMA.
Rapuntium, <i>Tourn. &</i> ... <i>Dill. Elth.</i>	Cardinal Flower.....	LOBELIA.
Rhabarbarum, <i>Tourn</i> ...	Rhubarb.....	RHEUM.
Rhagadioloides, <i>Va. A.G.</i>		HYOSERIS.
Rhagadiolus, <i>Vail. A. G.</i> <i>and Tourn.</i>		LAPSANA.
Rhamnoides, <i>Tourn</i>	Bastard Rhamnus, or Sea.. Buckthorn.	HIPPOPHAE.
Rhaponticoides, <i>Vaill</i> ...	Centaury.....	CENTAUREA.
Rhapontium, <i>Vaill</i>	Centaury.....	CENTAUREA.
Ribesium, <i>Dill. Elth</i> ...	Currant Tree.....	RIBES.
Ricinocarpus, <i>Boer. & Bur</i>		ACALYPHA.
Ricinoides, <i>Tourn</i>	Bastard Ricinus.....	CROTON.
Rivina, <i>Plum</i>		RIVINIA.
Royenia, <i>Houst. A. A.</i>		LÆSELIA.
Rojoc, <i>Plum</i>		MORINDA.
Ros solis, <i>Tourn</i>	Sun-dew.....	DROSERA.
Rubeola, <i>Tourn</i>	Petty Madder.....	CRUCIANELLA.
Rudbeckia, <i>Houst. A. A.</i>	Button Tree.....	CONOCARPUS.
Ruppia, <i>Act. Ang</i>	Grass Wrack.....	ZOSTERA.
Ruta muraria, <i>Tourn</i>	Wall-rue, or Tentwort....	ASPLENIUM.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
S		
Sabina, <i>Boerh</i>	Savine.....	JUNIPERUS.
Sagitta, <i>Dill. gen. and</i> ...	Arrow-head.....	SAGITTARIA.
<i>Vaill. A. G.</i>		
Salicaria, <i>Tourn</i>	Willow-herb, or Purple... Loosestrife.	LYTHRUM.
Salvinia, <i>Mich</i>		MARSILEA.
Santolinoides, <i>Vaill. A. G</i>		ANACYCLUS.
<i>and Mich. gen.</i>		
Sapota, <i>Plum</i>	Sapota.....	ACHRAS.
Sassafras, <i>Off</i>	Sassafras Tree.....	LAURUS.
Saururus, <i>Plum</i>	Lizard's Tail.....	PIPER.
Schunda Pana, <i>Hort. Mal</i>		CARYOTA.
Scirpocyperus, <i>Mitch</i> ...	Rush Grass.....	SCIRPUS.
Scirpoides, <i>Mont</i>		CAREX.
Sclarea, <i>Tourn</i>	CLARY.....	SALVIA.
Scorodoprasum, <i>Mich</i> ...	Great round-headed or.... Turkey Garlick.	ALLIUM.
Scorpioides, <i>Tourn</i>	Caterpillars.....	SCORPIURUS.
Scorzoneroides, <i>Vaill</i> ...	Viper's Grass.....	SCORZONERA.
<i>A. G.</i>		
Sebestena, <i>Dill. Elth</i> ...	Sebesten.....	CORDIA.
Securidaca, <i>Tourn</i>	The True Hatchet-Vetch... or Sicklewort.	CORONILLA.
Sedi species, <i>Tourn</i>	Houseleek.....	SEMPERVIVUM.
Selaginoides, <i>Dill. Musc</i>		LYCOPODIUM.
Selago, <i>Dill. Musc</i>	Upright Fir-Moss.....	LYCOPODIUM.
Senecionis species, <i>D. Elt</i>		ERIGERON.
Senna, <i>Tourn</i>	Senna of the shops.....	CASSIA.
Seriana, <i>Plum</i>		PAULLINIA.
Sesamoides, <i>Tourn</i>	Bastard Rocket.....	RESEDA.
Sherardia, <i>Vaill</i>	Vervain.....	VERBENA.
Sherardia, <i>Pont. Epist</i>		GALENIA.
Sicyoides, <i>Tourn</i>	Single-seeded Cucumber..	SICYOS.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
Siliqua, <i>Tourn</i>	Carob Tree, or St. John's... Bread.	CERATONIA.
Siliquastrum, <i>Tourn</i>	Judas Tree.....	CERCIS.
Silybium, <i>Vaill.</i> A. G.	Milk Thistle, or Lady's... Thistle.	CARDUUS.
Sinapi, <i>Tourn</i>	Mustard.....	SINAPIS.
Sinapistrum, <i>Tourn</i>	Bastard Mustard.....	CLAOME.
Siphonanthemum, <i>Amm</i>		SIPHONANTHUS
<i>Act. Petrop.</i>		
Sisarum, <i>Tourn</i>	Skirret.....	SIMUM.
Sisyrrinchium, <i>Tourn</i>	Iris with a double bulb,... called Spanish Nut.	IRIS.
Sloana, <i>Plum</i>	Apeiba of the Brasilians	SLOANEA.
Solanoides, <i>Tourn.</i> A. G.	American Nightshade....	RIVINA.
Sorgum, <i>Mich</i>	Indian Millet.....	HOLCUS.
Spartium, <i>Tourn</i>	Single-seeded Broom.....	GENISTA.
Sphondylium, <i>Tourn</i>	Cow Parsnep.....	HERACLEUM.
Sphondylococcus, <i>Mitc.</i>	Johnsonia.....	CALLICARPA.
Stachyarpagophora,.....	Cock's-comb.....	CELOSIA.
<i>Vaill.</i> A. G.		
Staphylodendron, <i>Tour.</i>	Bladder Nut.....	STAPHYLÆA.
Stellaria, <i>Dill. gen</i>		CALLITRICHE.
Stellaris, <i>Dill. gen</i>	Yellow Star of Bethlehem	ORNITHOGALUM
Stœchas, <i>Tourn</i>	French Lavender.....	LAVANDULA.
Stramonium, <i>Tou. & Pont.</i>	Thorn Apple.....	DATURA.
Stratiotes, <i>Vaill.</i> A. G.	Water Milfoil, or Water... Violet.	HOTTONIA.
Stratiotes, <i>Dill. gen</i> ,.....	Frog's-bit.....	HYDROCHARIS.
Struthia, <i>Royen</i>		GNIDIA.
Suber, <i>Tourn</i>	Cork Tree.....	QUERCUS.
Succisa, <i>Vaill.</i> A. G.....	Devil's-bit.....	SCABIOSA.
Suillus, <i>Mich</i>		BOLETUS.
Symphoricarpos, <i>Di. Elt.</i>	Shrubby St. Peter's-wort...	LONICERA.
Syringa, <i>Tourn</i>	Mock Orange, or Syringa	PHILADELPHUS

GÉNÉRIC NAMES REJECTED.	ENGLISH NAMES.	LINNÆAN GENERA.
T		
Tamariscus, <i>Tourn</i>	Tamarisk.....	TAMARIX.
Tamius, <i>Tourn</i>	Black Bryony.....	TAMUS.
Tapia, <i>Plum</i>	Garlick Pear.....	CRATEVA.
Taraxaconastrum, <i>Vaill</i>		HYOSERIS.
A. G.		
Taraxaconoides, <i>Vaill</i> ...	Dandelion.....	LEONTODON.
A. G.		
Tarchonanthus, <i>Va. Act.</i>	Jesuit's Bark Tree, false- ly so called.	IVA.
Telephiastrum, <i>Dill. Elt.</i>	African Purslane.....	PORTULACA.
Telephioides, <i>Tour. &</i>	Bastard Orpine.....	ANDRACHNE.
<i>Dill. Elth.</i>		
Tenga, <i>Hort. Mal</i>	Cocoa Nut.....	COCOS.
Terebinthus, <i>Tourn</i>	Turpentine Tree.....	PISTACIA.
Ternatea, <i>Tourn. A. G.</i>		CLITORIA.
Tetrahit, <i>Dill. gen</i>	Bastard Hemp.....	GALEOPSIS.
Thlaspidium, <i>Tourn</i>	Buckler Mustard.....	BISCUTELLA.
Thymbra, <i>Tourn</i>	Savory, with verticillate flowers.	SATUREJA.
Thymelæa, <i>Tourn</i>	Mezereon, or Spurge- Laurel.	DAPHNE.
Thysselinum, <i>Tourn</i>	Milky Parsley.....	SELINUM.
Tinus, <i>Tour. & Vaill. A. G.</i>	Laurustinus.....	VIBURNUM.
Titanokeratophyton, <i>Bo</i>		LITHOXYLON.
Tithymaloides, <i>Tourn</i> ...	Bastard Spurge.....	EUPHORBIA.
Tithymaloides (an) <i>Klein</i>	Cabbage Tree, or Car- <i>Monagr</i> nation Tree.	CACALIA.
Tithymalus, <i>Tourn</i>	Spurge.....	EUPHORBIA.
Tournefortia, <i>Pont. Epis.</i>	Amber Tree.....	ANTHOSPER- MUM.
Toxicodendron, <i>Tourn.</i>	Poison Tree	RHUS.
Tragacantha, <i>Tourn</i>	Goat's-horn.....	ASTRAGALUS.
Tragopogonoides, <i>Vaill</i> ...	Goat's-beard with crook- ed seeds.	TRAGOPOGON.
A. G.		

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Tragoselinum, <i>Tourn</i> ...	Burnet Saxifrage.....	PIMPINELLA.
Tribuloides, <i>Tourn</i>	Water Caltrops.....	TRAPA.
Trichomanes, <i>Tourn</i>	English Black Maiden- hair.	ASPLENUM.
Trifolium, <i>Mich</i>	White-flowered Meadow- Trefoil, Honeysuckle Grass, or Dutch Clover.	TRIFOLIUM.
Trilopus, <i>Mitch</i>	Witch Hazel.....	HAMAMELIS.
Triosteospermum, <i>Dill</i> ...	Fever-root, Doctor Tin- <i>Elth.</i> ker's Weed, or False Ipecacuana.	TRIOSTEUM.
Trixis, <i>Mitch</i>		PROSERPINACA.
Tulipifera, <i>Catesb</i>	Tulip Tree.....	LIRIODENDRON.
Tuna, <i>Dill. Elth</i>	Indian Fig, or Prickly Pear	CACTUS.
Tunica, <i>Dill. Elth</i>	Pink.....	DIANTHUS.
V		
Valdia, <i>Plum</i>		OVIEDA.
Valerianella, <i>Tour. and</i> ...	Lamb's Lettuce, or Corn- <i>Vaill.</i> Sallad.	VALERIANA.
Vallisneroides, <i>Mich</i>		VALISNERIA.
Vanilla, <i>Plum</i>	Vanilla.....	EPIDENDRUM.
Vanrheedia, <i>Plum</i>		RHEEDIA.
Vesicaria, <i>Rivinus</i>	Heart-seed, or Heart Pea	CARDIOSPER- MUM.
Vesicaria, <i>Tourn</i>	Madwort with bladdery pods.	ALYSSUM.
Virgaurea, <i>Tour. and</i> ...	Golden Rod.....	SOLIDAGO.
<i>Vaill. A. G.</i>		
Virga sanguinea, <i>Dill</i> ...	Female Dog-wood, Dog- berry, or Gatter Tree.	CORNUS.
Viscago, <i>Dill. Elth</i>	Viscous Campion, or Catch-fly.	SILENE.
Viticella, <i>Mitch</i>		GALAX.

GENERIC NAMES REJECTED.	ENGLISH NAMES.	LINNEAN GENERA.
Viticella, <i>Dill. gen.</i>	Virgin's Bower, or Lady's Bower.	CLEMATIS.
Vitis Idea, <i>Tourn.</i>	Whortleberry.....	VACCINIUM.
Ulmaria, <i>Tourn.</i>	Meadow-sweet, or Queen of the Meadows.	SPIRÆA.
Unifolium, <i>Dill. gen.</i>	One-blade.....	CONVALLARIA.
Volubilis, <i>Dill. Elth.</i>		IPOMŒA.
Usnea, <i>Dill. Musc.</i>	Tree Moss.....	LICHEN.
Uva ursi, <i>Tourn.</i>	Spanish Redwhorts, or Bearberries.	ARBUTUS.
Vulneraria, <i>Tourn.</i>	Kidney Vetch, or Lady's Finger.	ANTHYLLIS.

X

Xeranthemoides, <i>Dill.</i>		XERANTHE- MUM.
<i>Elth.</i>		
Xiphium, <i>Tourn.</i>	Bulbous Iris.....	IRIS.
Xylon, <i>Lin. gen. pl. ed. pr.</i>	Silk Cotton Tree.....	BOMBAX.
Xylon, <i>Tourn.</i>	Cotton.....	GOSSYPIUM.
Xylosteum, <i>Tourn.</i>	Fly Honeysuckle.....	LONICERA.

Z

Zacintha, <i>Vaill. A. G. and T.</i>	Wart Succory.....	LAPSANA.
Zanonia, <i>Plum.</i>		COMMELINA.
Ziziphus, <i>Tourn.</i>	Jujuba Tree.....	RHAMNUS.


I N D E X
OF
AUTHORS REFERRED TO
IN TABLE II.



<i>Amm.</i>	Ammannus.	<i>Knaut.</i>	Knautius.
<i>Battar.</i>	Battarra.	<i>Kram.</i>	Kramerus.
<i>Blackw.</i>	Blackwell. Edit. Nor.	<i>Linn.</i>	Linnæus.
<i>Boerh.</i>	Boerhavius.	— <i>E. N.</i> —	Editio novissima, 1754.
<i>Burm.</i>	Burmannus.	— <i>Gen.</i> —	Genera Planta- rum, Holm. 1754.
<i>Buxb.</i>	Buxbaumius.	— <i>Spec.</i> —	Species Planta- rum, Holm. 1753.
<i>Bauh.</i>	Bauhinus.	— <i>Syst.</i> —	Systema Naturæ, Holm. 1759.
<i>Cord.</i>	Cordus.	<i>Lob.</i>	Lobelius.
<i>Dill.</i>	Dillenius.	<i>Ludw.</i>	Ludwigius.
<i>Dod.</i>	Dodonæus.	<i>Mag.</i>	Magnolius.
<i>Fev.</i>	Feuillæus.	<i>Malp.</i>	Malpighius.
<i>Garc.</i>	Garcinus.	<i>March.</i>	Marchantius.
<i>Gled.</i>	Gleditschius.	<i>Mars.</i>	Marsilius.
<i>H. M.</i>	Hortus Malabaricus.	<i>Mich.</i>	Michelius.
<i>Hall.</i>	Hallerus.	<i>Niss.</i>	Nissolius.
<i>Heb.</i>	Hebenstreitius.	<i>Off.</i>	Officinarum.
<i>Heist.</i>	Heisterus.	<i>P. Alp.</i>	Prosper Alpinus.
<i>Herm.</i>	Hermannus.	<i>Pet.</i>	Petitus.
<i>Houst.</i>	Houstonus.	<i>Petiv.</i>	Petiverius.
<i>Imp.</i>	Imperatus.	<i>Pis.</i>	Piso.
<i>Johr.</i>	Johrenius.		
<i>Isn.</i>	Isnardus.		
<i>Juss.</i>	Jussieus.		
<i>Kæmpf.</i>	Kæmpferus.		

<i>Pluck.</i>	Pluckenetus.	<i>Schaff. G.</i>	— Beschreibung des
<i>Plum.</i>	Plumierus.		Gichtschwammes,
<i>Pont.</i>	Pontedera.		1760.
<i>Rai.</i>	Raius.	<i>Schaw.</i>	Schawius.
<i>Riv.</i>	Rivinus.	<i>Scheuchz.</i>	Scheuchzerus.
<i>Roy.</i>	Royenius.	<i>Sig.</i>	Sigesbeckius.
<i>Rudb.</i>	Rudbeckius.	<i>Sloan.</i>	Sloanus.
<i>Rupp.</i>	Ruppius.	<i>Tab.</i>	Tabernæmontanus.
<i>Schaff. A.</i>	Schæfferi erleichterte	<i>Tourn.</i>	Tournefortius.
	Arzneykräuterwis- <i>Trag.</i>		Tragus.
	senschaft, 1759.	<i>Trew.</i>	Trewius.
— <i>B.</i>	— Beobachtuntun- <i>Vaill.</i>		Vaillantius.
	gen der Schwämme	<i>Weinm.</i>	Weinmannius.
	um Regensb, 1759.		

TABLE II.
THE LINNÆAN GENERA,
WITH
REFERENCES AND SYNONYMES.



1. ACALYPHA.

Linn. *Gen.* 959. *Spec.* 1003. *Syst.* 959. Ludw. 897.
RICINOCARPOS. Boerh.

2. ACANTHUS.

Linn. *Gen.* 711. *Spec.* 639. *Syst.* 711. Mill. i. 14. Tourn. *tab.*
80, 81. Weinm. *tab.* 13. Ludw. 239.

3. ACER.

Linn. *Gen.* 1023. *Spec.* 1054. *Syst.* 1023. Hall. 421. Ludw.
551. Mill. i. 14. Tourn. *tab.* 386. Weinm. *tab.* 14-17.

4. ACHILLEA.

Linn. *Gen.* 871. *Spec.* 896. *Syst.* 871. Hall. 712. Ludw. 358.
MILLEFOLIUM. Tourn. *tab.* 283. Blackw. *tab.* 18. Mill. ii. 47.
Schæff. A. 122. Weinm. *tab.* 729, 730.
PTARMICA. Tourn. *tab.* 283. Blackw. *tab.* 276. Mill. ii. 165.
Schæff. A. 123. Weinm. *tab.* 837.

5. ACHRAS.

Linn. *Gen.* 1093. *Spec.* 1190. *Syst. No.* 1093, p. 1381.

SAPOTA. Plum. Ludw. 1046.

6. ACHYRANTHES.

Linn. *Gen.* 254. *Spec.* 204. *Syst.* 254. Ludw. 772.

ACHYRACANTHA. Dill.

7. ACNIDA.

Linn. *Gen.* 987. *Spec.* 1027. *Syst.* 987.

ACNIDE. Mitch.

8. ACONITUM.

Linn. *Gen.* 603. *Spec.* 532. *Syst.* 603. Hall. 312. Ludw. 653.

Mill. i. 17. Tourn. *tab.* 239, 240. Weinm. *tab.* 22-24.

NAPELLUS. Riv. ANTHORA. Riv.

9. ACORUS.

Linn. *Gen.* 392. *Spec.* 324. *Syst.* 392. Hall. 259. Ludw. 784.

Mill. iii. 8. Schæff. A. 245. Weinm. *tab.* 25.

CALAMUS AROMATICUS. Mich.

10. ACROSTICHUM.

Linn. *Gen.* 1037. *Spec.* 1067. *Syst.* 1037.

RUTA MURARIA. Tourn. *tab.* 317. Blackw. *tab.* 219. ADIANTHUM

ALBUM. Off. Schæff. A. 304. Weinm. *tab.* 26.

ACROSTICHUM. Hall. 134. Ludw. 942. ASPLENIUM. Hall. 134.

Ludw. 943.

11. ACTÆA.

Linn. *Gen.* 568. *Spec.* 504. *Syst.* 568.

CHRISTOPHORIANA. Tourn. *tab.* 154. Hall. 305. Ludw. 457.

Mill. i. 205. Weinm. *tab.* 384.

12. ADANSONIA.

Linn. *Gen.* 1094. *Spec.* 1190. *Syst. No.* 1094, p. 1382-1144.

13. ADELIA.

Linn. *Syst.* 1298.

14. ADENANTHERA.

Linn. *Gen.* 472. *Spec.* 384. *Syst.* 472. Ludw. 556.

15. ADIANTUM.

Linn. *Gen.* 1044. *Spec.* 1094. *Syst.* 1044. Blackw. *tab.* 367.
Ludw. 945. Mill. i. 19. Weinm. *tab.* 26, 27.

16. ADONIS.

Linn. *Gen.* 618. *Spec.* 547. *Syst.* 618. Hall. 319. Ludw. 753.
Mill. i. 20. iii. 9. Weinm. *tab.* 28.

17. ADOXA.

Linn. *Gen.* 450. *Spec.* 367. *Syst.* 450.
MOSCHATELLINA. Tourn. *tab.* 68. Hall. 412. Ludw. 137. Mill.
ii. 59. Weinm. 737.

18. ÆGILOPS.

Linn. *Gen.* 1018. *Spec.* 1050. *Syst.* 1018. Ludw. 847.

19. ÆGINETIA.

Linn. *Gen.* 695. *Spec.* 632, *Syst.* 695. Ludw. 1036.

20. ÆGOPODIUM.

Linn. *Gen.* 330. *Spec.* 265. *Syst.* 330.
PODAGRARIA. Riv. Hall. 427. Ludw. 658.

21. ÆSCHYNOMENE.

Linn. *Gen.* 769. *Spec.* 713. *Syst.* 769. Ludw. 499.

22. ÆSCULUS.

Linn. *Gen.* 420. *Spec.* 344. *Syst.* 420.
HIPPOCASTANUM. Tourn. *tab.* 382. Ludw. 630. Mill. i. 407.
Weinm. *tab.* 342. CASTANEA EQUINA. Rai.
PAVIA. Boerh. Ludw. 632. Mill. ii. 110.

23. ÆTHUSA.

Linn. *Gen.* 317. *Spec.* 256. *Syst.* 317. Hall. 433. Ludw. 692.
CYNAPIUM. Riv.

24. AGARICUS.

Linn. *Gen.* 1074. *Spec.* 1171. *Syst.* 1074. Schæff. B. § 70.
 AMANITA. Dill. Ludw. 963.
 FUNGUS. Mich. Battar. Gled. Hall. 24. Tourn. *tab.* 327.
 AGARICO-FUNGUS. Hall. 57.

25. AGAVE.

Linn. *Gen.* 390. *Spec.* 323. *Syst.* 390.

26. AGERATUM.

Linn. *Gen.* 843. *Spec.* 839. *Syst.* 843. Mill. i. 20. Weinm. *tab.* 29.
 CARELIA. Pont. Ludw. 299.

27. AGRIMONIA.

Linn. *Gen.* 534. *Spec.* 448. *Syst.* 534.
 AGRIMONIA. Tourn. *tab.* 155. Blackw. *tab.* 21. Hall. 407. Ludw.
 606. Mill. i. 21. Schæff. A. 195. Weinm. *tab.* 29.
 AGRIMONOIDES. Tourn. *tab.* 155. Ludw. 549. Mill. i. 23. iii. 9.

28. AGROSTEMMA.

Linn. *Gen.* 516. *Spec.* 435. *Syst.* 516.
 LYCHNIS. Hall. 376. Ludw. 573.

29. AGROSTIS.

Linn. *Gen.* 74. *Spec.* 61. *Syst.* 74. Hall. 218. 229. Ludw. 821.

30. AIRA.

Linn. *Gen.* 75. *Spec.* 63. *Syst.* 75. Ludw. 825.

31. AJUGA.

Linn. *Gen.* 624. *Spec.* 561. *Syst.* 624.
 BUGULA. Tourn. *tab.* 98. Hall. 633. Ludw. 191. Mill. i. 147.
 CONSOLIDA MEDIA. Off. Weinm. *tab.* 407.

32. AIZOON.

Linn. *Gen.* 553. *Spec.* 488. *Syst.* 553. Ludw. 808. Mill. iii. 9.
 FICOIDEA. Niss. Mill. i. 316.

33. ALCEA.

Linn. *Gen.* 750. *Spec.* 687. *Syst.* 750. Ludw. 145.

MALVA. Tourn. *tab.* 24. Ludw. 144. Mill. ii. 4. Schæff. A. 48.
50. Weinm. *tab.* 693-697.

34. ALCHEMILLA.

Linn. *Gen.* 153. *Spec.* 123. *Syst.* 153.

ALCHIMILLA. Tourn. *tab.* 289. Blackw. *tab.* 72. Hall. 184. Ludw.
764. Mill. i. 25. Schæff. A. 284. Weinm. *tab.* 36, 37.

35. ALDROVANDA.

Linn. *Gen.* 350. *Spec.* 281. *Syst.* 350.

36. ALETRIS.

Linn. *Gen.* 387. *Spec.* 319. *Syst.* 387.

37. ALISMA.

Linn. *Gen.* 418. *Spec.* 342. *Syst.* 418.

PLANTAGO AQUATICA. Boerh. Ludw. 384.

DAMASONIUM. Tourn. *tab.* 132. Hall. 300. Ludw. 385. Mill. i.
265.

38. ALLIONIA.

Linn. *Syst.* No. 1112. *p.* 1361. 890.

39. ALLIUM.

Linn. *Gen.* 370. *Spec.* 294. *Syst.* 370.

ALLIUM. Tourn. *tab.* 206. Hall. 296, 297. Ludw. 724. Mill. i.
26. iii. 11. Schæff. A. 249. Weinm. *tab.* 38, 39.

CEPA. Tourn. *tab.* 205. Hall. 295. Ludw. 724. Mill. i. 192.
Schæff. A. 250. Weinm. *tab.* 349.

PORRUM. Tourn. *tab.* 204. Hall. 294. Ludw. 724. Mill. i. 158.
Schæff. A. 251. Weinm. *tab.* 828.

SCORODOPRASUM. Mich. Ludw. 724.

MOLY. Boerh. Ludw. 427. Mill. ii. 56. Weinm. *tab.* 734.

40. ALLOPHYLUS.

Linn. *Gen.* 428. *Spec.* 348. *Syst.* 428.

41. ALOE.

Linn. *Gen.* 389. *Spec.* 319. *Syst.* 389. Blackw. *tab.* 229. Ludw.
116. Mill. i. 27. iii. 12. Tourn. *tab.* 191. Weinm. *tab.* 42-75.

42. ALOPECURUS.

Linn. *Gen.* 72. *Spec.* 60. *Syst.* 72. Ludw. 818. Hall. 205.

43. ALPINIA.

Linn. *Gen.* 4. *Spec.* 2. *Syst.* 4. Ludw. 173. Mill. iii. 12.

ALPINA. Plum.

44. AISINE.

Linn. *Gen.* 342. *Spec.* 272. *Syst.* 342. Blackw. *tab.* 164. Hall. 385. Ludw. 569. Tourn. *tab.* 126. Weinm. *tab.* 76-78.

45. ALTHÆA.

Linn. *Gen.* 749. *Spec.* 686. *Syst.* 749. Blackw. *tab.* 90. Hall. 364. Ludw. 146. Mill. i. 30. iii. 12. Schæff. A. 49. Weinm. *tab.* 79-83.

46. ALYSSUM.

Linn. *Gen.* 722. *Spec.* 650. *Syst.* 722. Ludw. 429.

ALYSSON. Tourn. *tab.* 104. Hall. 537. Mill. i. 31. Weinm. *tab.* 973.

ALYSSOIDES. Tourn. *tab.* 104. Mill. i. 31.

VESICARIA. Tourn.

47. AMARANTHUS.

Linn. *Gen.* 941. *Spec.* 989. *Syst.* 941. Blackw. *tab.* 317. Ludw. 882. Mill. i. 33. Tourn. *tab.* 118. Hall. 176. Weinm. *tab.* 84-99.

48. AMARYLLIS.

Linn. *Gen.* 367. *Spec.* 292. *Syst.* 367. Mill. iii. 13. Ludw. 723. LILIO-NARCISSUS. Tourn. *tab.* 207. Mill. i. 509. Weinm. *tab.* 672.

49. AMBROSIA.

Linn. *Gen.* 938. *Spec.* 987. *Syst.* 938. Ludw. 858. Mill. i. 34. Tourn. *tab.* 252.

50. AMELLUS.

Linn. *Syst.* No. 1162. p. 1377. 1225.

51. AMETHYSTEA.

Linn. *Gen.* 32. *Spec.* 21. *Syst.* 32.

AMETHYSTINA. Amm.

52. AMMANNIA.

Linn. *Gen.* 144. *Spec.* 119. *Syst.* 144. Ludw. 393.

53. AMMI.

Linn. *Gen.* 297. *Spec.* 243. *Syst.* 297. Ludw. 697. Mill. i. 35.
Weinm. *tab.* 99, 100. Tourn. *tab.* 159.

AMMOIDES. Boerh.

54. AMOMUM.

Linn. *Gen.* 2. *Spec.* 1. *Syst.* 2. Ludw. 170. Weinm. *tab.* 101.
ZINGIBER. Boerh.

55. AMORPHA.

Linn. *Gen.* 768. *Spec.* 713. *Syst.* 768. Ludw. 286.

56. AMYGDALUS.

Linn. *Gen.* 545. *Spec.* 472. *Syst.* 545.

AMYGDALUS. Tourn. *tab.* 402. Blackw. *tab.* 105. Ludw. 596.

Mill. i. 35. Schæff. A. 185. Weinm. *tab.* 101, 102.

PERSICA. Tourn. *tab.* 400. Blackw. *tab.* 101. Ludw. 597. Mill.

ii. 115. Schæff. A. 186. Weinm. *tab.* 707.

57. AMYRIS.

Linn. *Syst. No.* 1130. *p.* 1367. 100.

58. ANABASIS.

Linn. *Gen.* 276. *Spec.* 223. *Syst.* 276.

59. ANACARDIUM.

Linn. *Gen.* 467. *Spec.* 383. *Syst.* 467. Blackw. *tab.* 369. Ludw.
1021

ACAJOU. Tourn. *tab.* 435. Mill. i. 13. iii. 8. Weinm. *tab.* 104.

60. ANACYCLUS.

Linn. *Gen.* 869. *Spec.* 892. *Syst.* 869. Ludw. 312.

SANTALINOIDES. Vaill.

COTULA. Tourn. *tab.* 282.

61. ANAGALLIS.

Linn. *Gen.* 189. *Spec.* 148. *Syst.* 189. Blackw. *tab.* 43. 274.

Hall. 481. Ludw. 41. Mill. i. 36. iii. 17. Schæff. A. 327.

Tourn. *tab.* 59. Weinm. *tab.* 106, 107.

62. ANAGYRIS.

Linn. *Gen.* 457. *Spec.* 374. *Syst.* 457. Ludw. 635. Mill. i. 37.
Weinm. *tab.* 108, 109. Tourn. *tab.* 415.

63. ANASTATICA.

Linn. *Gen.* 715. *Spec.* 641. *Syst.* 715. Ludw. 426.

64. ANCHUSA.

Linn. *Gen.* 167. *Spec.* 133. *Syst.* 167. Blackw. *tab.* 112. Mill. i.
47. Weinm. *tab.* 117.

BUGLOSSUM. Tourn. *tab.* 53. Hall. 523. Ludw. 32. Mill. i. 146.
Schæff. A. 37. Weinm. *tab.* 271.

ALCANNA. Off. Schæff. A. 42.

65. ANDRACHNE.

Linn. *Gen.* 973. *Spec.* 1014. *Syst.* 973. Ludw. 862. Mill. iii. 19.
TELEPHIOIDES. Tourn. Mill. ii. 314.

66. ANDROMEDA.

Linn. *Gen.* 485. *Spec.* 393. *Syst.* 485.

LEDUM. Mich. Hall. 217.

CHAMÆDAPHNE. Buxb.

POLIIFOLIA. Buxb.

ERICÆ SPECIES. Tourn. *tab.* 373. B. Ludw. 139.

67. ANDROPOGON.

Linn. *Gen.* 1014. *Spec.* 1045. *Syst.* 1014.

68. ANDROSACE.

Linn. *Gen.* 179. *Spec.* 141. *Syst.* 179. Ludw. 44. Mill. iii. 20.
Tourn. *tab.* 46.

69. ANDRYALÀ.

Linn. *Gen.* 820. *Spec.* 808. *Syst.* 820. Ludw. 342.

ERIOPHORUS. Vaill.

70. ANEMONE.

Linn. *Gen.* 614. *Spec.* 538. *Syst.* 614. Ludw. 756.

ANEMONE. Tourn. *tab.* 147. Hall. 321. Mill. i. 47. Weinm. *tab.*
118-128.

ANEMONOIDES. Dill. Mill. iii. 20.

ANEMONE-RANUNCULUS. Dill.

TRINITAS. Hall. 320. HEPATICA. Dill. Blackw. *tab.* 207. Mill. i. 401. Schæff. A. 260. Weinm. *tab.* 570.

PULSATILLA. Tourn. *tab.* 148. Mill. ii. 168. Weinm. *tab.* 838-840.

71. ANETHUM.

Linn. *Gen.* 326. *Spec.* 263. *Syst.* 326.

ANETHUM. Tourn. *tab.* 169. Ludw. 657. Mill. i. 50. Schæff. A. 236. Weinm. *tab.* 129.

FÆNICULUM. Tourn. *tab.* 164. Blackw. *tab.* 288. Hall. 425. Ludw. 669. Mill. i. 326. Schæff. A. 237. Weinm. *tab.* 513.

72. ANGELICA.

Linn. *Gen.* 309. *Spec.* 250. *Syst.* 309. Hall. 445. Ludw. 661. Mill. i. 50. iii. 21. Schæff. A. 239. Tourn. *tab.* 167. Weinm. *tab.* 130-136.

73. ANNONA.

Linn. *Gen.* 613. *Spec.* 536. *Syst.* 613.

GUANABANUS. Plum. Mill. i. 384. iii. 125.

ANONA. Ludw. 744.

74. ANTHEMIS.

Linn. *Gen.* 870. *Spec.* 893. *Syst.* 870.

CHAMEMELUM. Tourn. *tab.* 281. Blackw. *tab.* 67. Hall. 716. Ludw. 357. Mill. i. 200. CHAMOMILLA. Off. Schæff. A. 127. Weinm. *tab.* 362-364.

BUPHTHALMUM. Tourn. *tab.* 282. Ludw. 362. Mill. i. 149.

ANTHEMIS. Mich.

75. ANTHERICUM.

Linn. *Gen.* 380. *Spec.* 310. *Syst.* 380. Hall. 291.

PHALANGIUM. Tourn. *tab.* 193. Mill. ii. 134. Ludw. 713. Weinm. *tab.* 807.

BULBINE. Linn. *edit. prior.* ASPHODELITIS. Boerh.

76. ANTHOCEROS.

Linn. *Gen.* 1064. *Spec.* 1139. *Syst.* 1064. Hall. 127. Ludw. 981.

77. ANTHOLYZA.

Linn. *Gen.* 56. *Spec.* 37. *Syst.* 56.

78. ANTHOSPERMUM.

Linn. *Gen.* 1029. *Spec.* 1058. *Syst.* 1029. Ludw. 1035. Mill.
iii. 22.

TOURNEFORTIA. Pont.

79. ANTHOXANTHUM.

Linn. *Gen.* 40. *Spec.* 28. *Syst.* 40. Hall. 230. Ludw. 812.

80. ANTHYLLIS.

Linn. *Gen.* 773. *Spec.* 719. *Syst.* 773. Ludw. 475. Weinm. *tab.*
142.

VULNERARIA. Tourn. *tab.* 211. Hall. 569. Mill. ii. 466.

ERINACEA. Tourn.

BARBA JOVIS. Boerh.

81. ANTIDESMA.

Linn. *Gen.* 985. *Spec.* 1027. *Syst.* 985.

82. ANTIRRHINUM.

Linn. *Gen.* 668. *Spec.* 612. *Syst.* 668. Hall. 613. Ludw. 247.

ANTIRRHINUM. Tourn. *tab.* 75. Mill. i. 60. Weinm. *tab.* 144.

LINARIA. Tourn. *tab.* 76. Blackw. *tab.* 115. Hall. 613. Mill. i.
518. Schæff. A. 78. Weinm. *tab.* 664, 665.

ASARINA. Tourn. *tab.* 76.

ELATINE. Riv. Blackw. *tab.* 170. Weinm. *tab.* 476.

83. APHANES.

Linn. *Gen.* 154. *Spec.* 123. *Syst.* 154. Ludw. 770.

PERCEPIER. Dill. Hall. 184.

84. APHYLLANTHES.

Linn. *Gen.* 369. *Spec.* 294. *Syst.* 369. Ludw. 725. Tourn. *tab.*
430.

85. APIUM.

Linn. *Gen.* 329. *Spec.* 264. *Syst.* 329. Blackw. *tab.* 172. Hall.
427. Ludw. 695. Mill. i. 65. Tourn. *tab.* 160. Schæff. A. 226.
Weinm. *tab.* 150.

86. APLUDA.

Linn. *Gen.* 89. *Spec.* 82. *Syst.* No. 89. p. 1306. 1383.

87. APOCYNUM.

Linn. *Gen.* 269. *Spec.* 213. *Syst.* 269. Ludw. 98. Mill. i. 67.
iii. 23. Tourn. *tab.* 20. Weinm. *tab.* 151.

88. AQUILEGIA.

Linn. *Gen.* 605. *Spec.* 533. *Syst.* 605. Hall. 310. Ludw. 752.
Mill. i. 71. iii. 25. Schæff. A. 259. Tourn. *tab.* 242. Weinm.
tab. 160-164.

89. ARABIS.

Linn. *Gen.* 732. *Spec.* 664. *Syst.* 732. Ludw. 414. Hall. 561.

90. ARACHIS.

Linn. *Gen.* 787. *Spec.* 741. *Syst.* 787. Ludw. 483. Mill. iii. 26.
ARACHIDNA. Plum. Weinm.
ARACHIDNOIDES. Niss.

91. ARALIA.

Linn. *Gen.* 346. *Spec.* 273. *Syst.* 346. Ludw. 548. Mill. i. 72.
Tourn. *tab.* 154.

92. ARBUTUS.

Linn. *Gen.* 488. *Spec.* 395. *Syst.* 488.
ARBUTUS. Tourn. *tab.* 368. Hall. 415. Ludw. 140. Mill. i. 73.
Weinm. *tab.* 166.
UVA URSI. Tourn. *tab.* 370. Hall. 415. Mill. ii. 466. Schæff.
A. 43.

93. ARCTIUM.

Linn. *Gen.* 830. *Spec.* 816. *Syst.* 830. Hall. 675.
LAPPA. Tourn. *tab.* 256. Ludw. 324.
BARDANA. Dod. Blackw. *tab.* 117. Schæff. A. 99. Weinm. *tab.*
231.

94. ARCTOPUS.

Linn. *Gen.* 1030. *Spec.* 1058. *Syst.* 1030.

95. ARCTOTIS.

Linn. *Gen.* 886. *Spec.* 922. *Syst.* 886. Mill. iii. 26.

ANTHOTHECA. Vaill.

ANEMONOSPERMOS. Boerh. Ludw. 363. Mill. i. 50.

96. ARECA.

Linn. *Gen.* 1090. *Spec.* 1189. *Syst.* 1090.

97. ARENARIA.

Linn. *Gen.* 505. *Spec.* 423. *Syst.* 505.

ALSINE. Hall. 385-387. Ludw. 569.

98. ARETHUSA.

Linn. *Gen.* 905. *Spec.* 950. *Syst.* 905.

ORCHIDION. Mitch.

99. ARETIA.

Linn. *Gen.* 178. *Spec.* 141. *Syst.* 178. Hall. 485. Ludw. 44.

100. ARGEMONE.

Linn. *Gen.* 574. *Spec.* 508. *Syst.* 574. Ludw. 448. Mill. i. 73.

Tourn. *tab.* 121.

101. ARISTIDA.

Linn. *Gen.* 88. *Spec.* 82. *Syst.* 88.

102. ARISTOLOCHIA.

Linn. *Gen.* 911. *Spec.* 960. *Syst.* 911. Blackw. *tab.* 255-257.

Hall. 196. Ludw. 283. Mill. i. 74. Tourn. *tab.* 71. Schæff. A.

52. Weinm. *tab.* 167, 168.

103. ARNICA.

Linn. *Gen.* 861. *Spec.* 884. *Syst.* 861.

GERBERA. Linn. *edit. prior.* Ludw. 356. Weinm. *tab.* 469.

104. ARTEDIA.

Linn. *Gen.* 295. *Spec.* 242. *Syst.* 295. Ludw. 667.

105. ARTEMISIA.

Linn. *Gen.* 849. *Spec.* 845. *Syst.* 849. Ludw. 296. Hall. 694.

ARTEMISIA. Tourn. *tab.* 260. Mill. i. 75. Schæff. A. 103. Weinm. *tab.* 170, 171.

ABROTANUM. Tourn. Mill. i. 6. iii. 4. Schæff. A. 104. Weinm. tab. 4-6.

ABSINTHIUM. Tourn. tab. 260. Blackw. tab. 17. Mill. i. 8. Schæff. A. 105. Weinm. tab. 7-9.

DRACO. Boerh. DRACUNCULUS. Bauh. Blackw. tab. 116.

106. ARUM.

Linn. *Gen.* 915. *Spec.* 964. *Syst.* 915. Hall. 260. Ludw. 810.
ARUM. Tourn. tab. 69. Blackw. tab. 228. Hall. 261. Mill. i. 77.
Schæff. A. 53. Weinm. tab. 172-177.

ARISARUM. Tourn. tab. 70. Mill. i. 74. iii. 27. Weinm. tab. 169.

COLOCASIA. Boerh.

DRACUNCULUS. Tourn. tab. 70. Blackw. tab. 269. Mill. i. 277.
Weinm. tab. 472.

107. ARUNDO.

Linn. *Gen.* 87. *Spec.* 81. *Syst.* 87. Hall. 221. Ludw. 829. Mill.
i. 78. iii. 28. Weinm. tab. 178-180.

108. ASARUM.

Linn. *Gen.* 522. *Spec.* 442. *Syst.* 522.

ASARUM. Tourn. tab. 286. Blackw. tab. 359. Hall. 195. Ludw.
801. Mill. i. 79. iii. 29. Schæff. A. 268. Weinm. tab. 181.

HUPOCISTUS. Tourn. tab. 477. Ludw. 804. Mill. i. 421.

109. ASCLEPIAS.

Linn. *Gen.* 270. *Spec.* 214. *Syst.* 270.

ASCLEPIAS. Tourn. tab. 22. Blackw. tab. 96. Hall. 525. Ludw.
99. Mill. i. 80. iii. 29.

APOCYNUM. Tourn. tab. 21. Weinm. tab. 152, seq.

VINCETOXICUM. Off. Schæff. A. 35. Weinm. tab. 1011. a, b.

110. ASCYRUM.

Linn. *Gen.* 809. *Spec.* 787. *Syst.* 809. Mill. i. 80.

HYPERICOIDES. Plum. Ludw. 453.

111. ASPALATHUS.

Linn. *Gen.* 767. *Spec.* 711. *Syst.* 767.

ACHYRONIA. Royen. Ludw. 476.

112. ASPARAGUS.

Linn. *Gen.* 382. *Spec.* 313. *Syst.* 382. Blackw. *tab.* 332. Ludw. 730. Mill. i. 81. iii. 29. Schæff. A. 246. Tourn. *tab.* 154. Weinm. *tab.* 182.

113. ASPERUGO.

Linn. *Gen.* 173. *Spec.* 138. *Syst.* 173. Hall. 522. Ludw. 40. Mill. i. 84. Tourn. *tab.* 54.

APARINE MAJOR. Weinm. *tab.* 148. a.

114. ASPERULA.

Linn. *Gen.* 113. *Spec.* 103. *Syst.* 113.

RUBEOLA. Hall. 457.

115. ASPHODELUS.

Linn. *Gen.* 379. *Spec.* 309. *Syst.* 379. Blackw. *tab.* 233-238. Ludw. 119. Mill. i. 85. Schæff. A. 12. Tourn. *tab.* 178. Weinm. *tab.* 184.

116. ASPLENIUM.

Linn. *Gen.* 1042. *Spec.* 1078. *Syst.* 1042. Hall. 134. Ludw. 943. ASPLENIUM. Tourn. *tab.* 318. Mill. i. 85.

LINGUA CERVINA. Tourn. *tab.* 319. Blackw. *tab.* 138. Weinm. *tab.* 667, *seq.*

TRICHOMANES. Tourn. *tab.* 315. a, b. Blackw. *tab.* 370. Weinm. *tab.* 26. d.

117. ASTER.

Linn. *Gen.* 858. *Spec.* 872. *Syst.* 858. Hall. 725. Ludw. 352. Mill. i. 86. iii. 30. Tourn. *tab.* 274. Weinm. *tab.* 187-196.

ASTER. Vaill.

HELERIUM. Vaill.

ASTEROPTERUS. Vaill.

118. ASTRAGALUS.

Linn. *Gen.* 799. *Spec.* 755. *Syst.* 799. Hall. 565.

ASTRAGALUS. Tourn. *tab.* 233. Ludw. 508. Mill. i. 90. iii. 31. Weinm. *tab.* 196.

TRAGACANTHA. Tourn. *tab.* 234. Blackw. *tab.* 264. Ludw. 639. Mill. ii. 346. Weinm. *tab.* 977. d.

119. ASTRANTIA.

Linn. *Gen.* 290. *Spec.* 235. *Syst.* 290. Hall. 439. Ludw. 655.
Mill. i. 90. Tourn. *tab.* 166.

120. ATHAMANTA.

Linn. *Gen.* 301. *Spec.* 244. *Syst.* 301.

MEUM. Tourn. *tab.* 165. Hall. 426. Ludw. 677. Mill. ii. 47.
Schæff. A. 234. Weinm. *tab.* 729. a.

OREOSELINUM. Tourn. *tab.* 169. Ludw. 665. Mill. ii. 93. SELI-
NUM. Hall. 143.

121. ATRACTYLIS.

Linn. *Gen.* 837. *Spec.* 829. *Syst.* 837. Ludw. 365.

CROCODILODES. Vaill.

122. ATRAGENE.

Linn. *Gen.* 615. *Spec.* 542. *Syst.* 615.

123. ATRAPHAXIS.

Linn. *Gen.* 405. *Spec.* 333. *Syst.* 405. Ludw. 371. Mill. iii. 27.

124. ATRIPLEX.

Linn. *Gen.* 1021. *Spec.* 1052. *Syst.* 1021. Blackw. *tab.* 99, 100,
Tourn. *tab.* 286. Hall. 173. Ludw. 775. Mill. i. 91. iii. 31.
Schæff. A. 275. Weinm. *tab.* 200-204.

125. ATROPA.

Linn. *Gen.* 222. *Spec.* 181. *Spec.* 222.

BELLADONA. Tourn. *tab.* 13. Hall. 508. Ludw. 90. Mill. i. 117.
iii. 38. Schæff. A. 38. Weinm. *tab.* 235.

126. AVENA.

Linn. *Gen.* 85. *Spec.* 79. *Syst.* 85. Hall. 222, 223. Ludw. 830.
Mill. i. 93. iii. 31. Schæff. A. 302. Tourn. *tab.* 297. Weinm.
tab. 205.

127. AVERRHOA.

Linn. *Gen.* 511. *Spec.* 428. *Syst.* 511. Ludw. 578.

128. AVICENNIA.

Linn. *Gen.* 125. *Spec.* 110. *Syst.* 125. Ludw. 16.

129. ANYRIS.

Linn. *Gen.* 929. *Spec.* 979. *Syst.* 929.

130. AYENIA.

Linn. *Syst. No.* 1164. *p.* 1378. 1247.

131. AZALEA.

Linn. *Gen.* 195. *Spec.* 150. *Syst.* 195. Hall. 416. Ludw. 71.
LEDUM. Hall. 417.
CHAMÆRHODODENDROS. Tourn.

132. BACCHARIS.

Linn. *Gen.* 853. *Spec.* 860. *Syst.* 853. Mill. iii. 34.
CONYZA. Ludw. 306.

133. BÆCKEA.

Linn. *Gen.* 442. *Spec.* 358. *Syst.* 442.

134. BALLOTA.

Linn. *Gen.* 639. *Spec.* 582. *Syst.* 639.
BALLOTE. Tourn. *tab.* 85. Blackw. *tab.* 136. Hall. 648. Ludw.
204. Mill. i. 100. Weinm. *tab.* 711. b.
MARRUBIASTRUM. Riv.

135. BANISTERIA.

Linn. *Gen.* 509. *Spec.* 427. *Syst.* 509. Ludw. 568. Mill. i. 101.

136. BARLERIA.

Linn. *Gen.* 703. *Spec.* 636. *Syst.* 703. Ludw. 230. Mill. i. 109.
iii. 35.

137. BARRERIA.

Linn. *Gen.* 347. *Spec.* 274. *Syst.* 347. Ludw. 1038.

138. BARTRAMIA.

Lin. *Gen.* 480. *Spec.* 398. *Syst.* ...

139. BARTSIA.

Linn. *Gen.* 657. *Spec.* 602. *Syst.* 657. Ludw. 243.

STÆHELINA. Hall. 624.

140. BASELLA.

Linn. *Gen.* 343. *Spec.* 272. *Syst.* 343. Ludw. 27. Mill. i. 110.

141. BATIS.

Linn. *Syst. No.* 1152. p. 1380. 1289.

142. BAUHINIA.

Linn. *Gen.* 459. *Spec.* 374. *Syst.* 459. Ludw. 645. Mill. i. 112.
iii. 36.

143. BEGONIA.

Linn. *Gen.* 1024. *Spec.* 1056. *Syst.* 1024. Ludw. 1044. Tourn.
tab. 442.

144. BELLIS.

Linn. *Gen.* 864. *Spec.* 886. *Syst.* 864. Blackw. *tab.* 200. Hall
722. Ludw. 347. Mill. i. 118. Schæff. A. 117. Tourn. *tab.*
280. Weinm. *tab.* 236, 237.

145. BELLONIA.

Linn. *Gen.* 207. *Spec.* 172. *Syst.* 207. Ludw. 1013. Mill. i. 118.

146. BERBERIS.

Linn. *Gen.* 399. *Spec.* 330. *Syst.* 399. Blackw. *tab.* 165. Hall.
424. Ludw. 729. Mill. i. 119. Schæff. A. 258. Tourn. *tab.*
385. Weinm. *tab.* 240.

147. BESLERIA.

Linn. *Gen.* 673. *Spec.* 619. *Syst.* 673. Ludw. 263. Mill. i. 122.

148. BETA.

Linn. *Gen.* 274. *Spec.* 222. *Syst.* 274. Blackw. *tab.* 235. Ludw.
777. Mill. i. 123. Schæff. A. 277. Tourn. 286. Weinm. *tab.*
241, 242.

TABLE II.

149. BETONICA.

Linn. *Gen.* 631. *Spec.* 573. *Syst.* 631. Blackw. *tab.* 46. Hall.
645. Ludw. 203. Mill. i. 123. Schæff. A. 67. Tourn. *tab.*
96. Weinm. *tab.* 243.

150. BETULA.

Linn. *Gen.* 933. *Spec.* 982. *Syst.* 933.

BETULA. Tourn. *tab.* 360. Blackw. *tab.* 240. Hall. 158. Ludw.
879. Mill. i. 124. Schæff. A. 292. Weinm. *tab.* 244.

ALNUS. Tourn. *tab.* 359. Hall. 157. Ludw. 878. Mill. i. 27.
iii. 11. Weinm. *tab.* 40, 41.

151. BIDENS.

Linn. *Gen.* 840. *Spec.* 831. *Syst.* 840. Hall. 709. Ludw. 313.
Mill. i. 124. Tourn. *tab.* 262.

CERATOCEPHALUS. Vaill.

152. BIGNONIA.

Linn. *Gen.* 677. *Spec.* 622. *Syst.* 677. Ludw. 1025. Mill. i. 125.
Tourn. *tab.* 72.

GELSEMINUM. Weinm. *tab.* 530. c.

153. BISCUTELLA.

Linn. *Gen.* 724. *Spec.* 652. *Syst.* 724. Hall. 541. Ludw. 420.
THLASPIDIUM. Tourn. *tab.* 101. Mill. ii. 331.

PERSPICILLUM. Heist.

154. BISERRULA.

Linn. *Gen.* 800. *Spec.* 762. *Syst.* 800.

PELECINUS. Tourn. *tab.* 234. Ludw. 509. Mill. ii. 111.

155. BIXA.

Linn. *Gen.* 581. *Spec.* 512. *Syst.* 581. Ludw. 749.

156. BLÆRIA.

Linn. *Gen.* 130. *Spec.* 112. *Syst.* 130.

157. BLAKEA.

Linn. No. 1141. p. 1370, 1044.

158. BLASIA.

Linn. *Gen.* 1062. *Spec.* 1138. *Syst.* 1062. Ludw. 984.

159. BLECHNUM.

Linn. *Gen.* 1039. *Spec.* 1077. *Syst.* 1039.

160. BLITUM.

Linn. *Gen.* 14. *Spec.* 4. *Syst.* 14. Hall. 774.

CHENOPODIOMORUS. Boerh. Mill. i. 205.

MOROCARPUS. Rup. Ludw. 761.

161. BOBARTIA.

Lin. *Gen.* 66. *Spec.* 54. *Syst.* 66.

162. BOCCONIA.

Linn. *Gen.* 569. *Spec.* 505. *Syst.* 569. Ludw. 459. Mill. i. 132.

163. BOERHAVIA.

Linn. *Gen.* 9. *Spec.* 3. *Syst.* 9. Ludw. 6. Mill. iii. 41.

ANTANISOPHYLLUM. Vaill.

164. BOLETUS.

Linn. *Gen.* 1075. *Spec.* 1176. *Syst.* 1075. Gled, *tab.* iii. Schæff.
B. 870.

SUILLUS. Mich. Hall. 29. Ludw. 964. FUNGUS. Tourn. *tab.*
328.

POLYPORUS. Mich. Hall. 25. Ludw. 965.

AGARICO-POLYPORUS. Hall. 26.

CERIOMYCES. AGARICUS. Battarr. *tab.* iv. xxxviii.

165. BOMBAX.

Linn. *Gen.* 580. *Spec.* 511. *Syst.* No. 580. p. 1141.

XYLON. Linn. *edit. prior.* Ludw. 523.

CEIBA. Plum. Mill. i. 188. iii. 54.

166. BONTIA.

Linn. *Gen.* 709. *Spec.* 638. *Syst.* 709. Ludw. 1026. Mill. i. 133.

167. BORAGO.

Linn. *Gen.* 172. *Spec.* 137. *Syst.* 172.

BORRAGO. Tourn. *tab.* 53. Blackw. *tab.* 36. Hall. 524. Ludw.
31. Mill. i. 134. iii. 42. Schæff. A. 39. Weinm. *tab.* 253,
254.

BORRAGINOIDES. Boerh. CYNOGLOSSOIDES. Isnard.

168. BORASSUS.

Lin. *Gen.* 1085. *Spec.* 1187. *Syst.* 1085.

AMPANA. H. M. CARIMPANA. H. M.

169. BORBONIA.

Linn. *Gen.* 764. *Spec.* 707. *Syst.* 764. Ludw. 638.

170. BOSEA.

Linn. *Gen.* 280. *Spec.* 225. *Syst.* 280. Ludw. 781. Mill. iii. 42.

171. BRABEUM.

Linn. *Gen.* 149. *Spec.* 121. *Syst.* 149. Ludw. 391. Mill. iii. 43.

172. BRASSICA.

Linn. *Gen.* 734. *Spec.* 666. *Syst.* 734.

BRASSICA. Tourn. *tab.* 106. Ludw. 405. Mill. i. 137. Schæff. A.
151. Weinm. *tab.* 256.

RAPA. Tourn. *tab.* 113. Blackw. *tab.* 226. Ludw. 406. Mill. ii.
189. Schæff. A. 152. Weinm. *tab.* 859.

NAPUS. Tourn. Blackw. *tab.* 224. Mill. ii. 66. Weinm. *tab.*
746. a.

173. BREYNIA.

Linn. *Gen.* 566. *Spec.* 503. *Syst.* ...Ludw. 451. Mill. i. 144.

174. BRIZA.

Linn. *Gen.* 78. *Spec.* 70. *Syst.* 78. Ludw. 837.

175. BROMELIA.

Linn. *Gen.* 356. *Spec.* 285. *Syst.* 356.

ANANAS. Tourn. *tab.* 426, 427, 428. Ludw. 381. Mill. i. 37. iii.
17. Weinm. *tab.* 110-116.

KARATAS. Plum. Mill. i. 471.

PINGVIN. Dill.

176. BROMUS.

Linn. *Gen.* 83. *Spec.* 76. *Syst.* 83. Hall. 227. Ludw. 834.
ÆGILOPS. Dill.

177. BROSSÆA.

Linn. *Gen.* 1095. *Spec.* 1190. *Syst.* 1095. Ludw. 1047.

178. BROWALLIA.

Linn. *Gen.* 691. *Spec.* 631. *Syst.* 691. Ludw. 237.

179. BRUNIA.

Linn. *Gen.* 242. *Spec.* 199. *Syst.* 242. Ludw. 514.

180. BRUNSFELSIA.

Linn. *Gen.* 230. *Spec.* 191. *Syst.* 230. Mill. i. 145.
BRUNFELSIA. Plum. Ludw. 93.

181. BRYONIA.

Linn. *Gen.* 970. *Spec.* 1012. *Syst.* 970. Blackw. *tab.* 37. Hall.
505. Ludw. 856. Mill. i. 145. Schæff. A. 22. Tourn. *tab.* 28.
Weinm. *tab.* 269:

182. BRYUM.

Linn. *Gen.* 1057. *Spec.* 1115. *Syst.* 1057. Ludw. 957.

183. BUBON.

Linn. *Gen.* 312. *Spec.* 253. *Syst.* 312. Ludw. 687.

184. BUCEPHALON.

Linn. *Gen.* 1096. *Spec.* 1190. *Syst.* 1096. Ludw. 1048.

185. BUCHNERA.

Linn. *Gen.* 690. *Spec.* 630. *Syst. No.* 580. p. 690. Ludw. 20.

186. BUCIDA.

Linn. *Gen. No.* 1135. p. 1368. 1025.

TABLE II.

187. BUDDLEIA.

Linn. *Gen.* 131. *Spec.* 112. *Syst.* 131. Ludw. 21.

188. BUFONIA.

Linn. *Gen.* 41. *Spec.* 123. *Syst.* 42.

ALSINOIDES. Rai.

189. BULBOCODIUM.

Linn. *Gen.* 368. *Spec.* 294. *Syst.* 368. Ludw. 727. Mill. i. 149.

190. BUNIAS.

Linn. *Gen.* 737. *Spec.* 669. *Syst.* 737.

ERUCAGO. Tourn. *tab.* 103. Ludw. 430. Mill. i. 301.

191. BUNIUM.

Linn. *Gen.* 298. *Spec.* 243. *Syst.* 298. Ludw. 686.

BULBOCASTANUM. Tourn. *tab.* 161. Hall. 782. Mill. i. 148.

Weinm. *tab.* 273.

192. BUPHTHALMUM.

Lin. *Gen.* 876. *Spec.* 903. *Syst.* 876. Hall. 710. Ludw. 362.

BUPHTHALMUM. Tourn. *tab.* 282. Mill. i. 149.

ASTERISCUS. Tourn. *tab.* 283. Blackw. *tab.* 272. Mill. i. 88.

ASTEROIDES. Tourn. *tab.* 487. Mill. i. 88.

193. BUPLEURUM.

Linn. *Gen.* 291. *Spec.* 236. *Syst.* 291. Tourn. *tab.* 163. Hall. 436.

Ludw. 685. Mill. i. 150. iii. 44. Weinm. *tab.* 273. 1.

PERFOLIATA. Riv. Blackw. *tab.* 95. Schæff. A. 230. Weinm.

tab. 801. b, c, d.

194. BURMANNIA.

Linn. *Gen.* 359. *Spec.* 287. *Syst.* 359. Ludw. 380.

195. BUTOMUS.

Linn. *Gen.* 455. *Spec.* 372. *Syst.* 455. Hall. 299. Ludw. 387.

Mill. i. 151. Tourn. *tab.* 143.

196. BUXBAUMIA.

Linn. *Syst.* 1332.

197. BUXUS.

Linn. *Gen.* 934. *Spec.* 983. *Syst.* 934. Blackw. *tab.* 196. Hall. 163. Ludw. 881. Mill. i. 151. Schæff. A. 285. Tourn. *tab.* 345. Weinm. *tab.* 275.

198. BYSSUS.

Linn. *Gen.* 1071. *Spec.* 1168. *Syst.* 1071.

ASPERGILLUS. Mich. Hall. 6.

BOFRYTIS. Mich. Hall. 7.

EMBOLUS. Hall. 8.

199. BYTTNERIA.

Linn. *Syst.* No. 1125. p. 1365. 939.

200. CACALIA.

Linn. *Gen.* 841. *Spec.* 834. *Syst.* 841.

CACALIA. Tourn. *tab.* 258. Mill. iii. 44.

CACALIANthemum. Dill. Mill. i. 152. KLEINIA. Linn. *edit. prior.* Mill. iii. 151.

POROPHYLLUM. Vaill.

201. CACHRYS.

Linn. *Gen.* 304. *Spec.* 246. *Syst.* 304. Ludw. 688. Mill. i. 153. Tourn. *tab.* 172.

202. CACTUS.

Linn. *Gen.* 539. *Spec.* 466. *Syst.* 539. Mill. iii. 45.

CEREUS. Juss. Ludw. 162. Mill. i. 196. iii. 59. Weinm. *tab.* 354. 358.

OPUNTIA. Tourn. *tab.* 122. Ludw. 163. Mill. ii. 90. Weinm. *tab.* 766. TUNA. Dill.

MELOCACTUS. Tourn. *tab.* 425. Ludw. 161. Mill. ii. 38. Weinm. *tab.* 474.

PERESKIA. Plum. Ludw. 164. Mill. ii. 112.

203. CÆSALPINA.

Linn. *Gen.* 463. *Spec.* 380. *Syst.* 463. Ludw. 1049. Mill. i. 158.

204. CALAMUS.

Linn. *Gen.* 395. *Spec.* 325. *Syst.* 395.

205. CALENDULA.

Linn. *Gen.* 885. *Spec.* 921. *Syst.* 885. Blackw. *tab.* 106. Weinm.
tab. 282, *seq.*

CALTHA. Tourn. *tab.* 284. Hall. 722. Ludw. 348. Mill. i. 161.
iii. 48. Schæff. A. 132.

DIMORPHOTHECA. Vaill.

206. CALLA.

Linn. *Gen.* 917. *Spec.* 968. *Syst.* 917. Ludw. 811.

PROVENZALIA. Petit.

ANGUINA. Trew.

207. CALLICARPA.

Linn. *Gen.* 127. *Spec.* 111. *Syst.* 127.

SPONDYLOCOCCOS. Mitch.

208. CALLIGONUM.

Linn. *Gen.* 601. *Spec.* 530. *Syst.* 601. Ludw. 805.

POLYGONOIDES. Tourn. *tab.* 478.

209. CALLITRICHE.

Linn. *Gen.* 13. *Spec.* 969. *Syst.* 13.

STELLARIA. Dill.

210. CALOPHYLLUM.

Linn. *Gen.* 586. *Spec.* 513. *Syst.* 586.

CALABA. Plum. Mill. i. 160. Ludw. 455.

211. CALTHA.

Linn. *Gen.* 623. *Spec.* 558. *Syst.* 623. Weinm. *tab.* 184.

POPULAGO. Tourn. *tab.* 145. Hall. 319. Ludw. 624. Mill. ii.
157.

212. CALYCANTHUS.

Linn. *Syst. No.* 1144. *p.* 1371. 1066.

213. CAMBOGIA.

Linn. *Gen.* 576. *Spec.* ... *Syst.* 576. Blackw. *tab.* 392.

214. CAMELLIA.

Linn. *Gen.* 759. *Spec.* 698. *Syst.* 759. Ludw. 154.

215. CAMERARIA.

Linn. *Gen.* 264. *Spec.* 210. *Syst.* 264. Ludw. 105. Mill. iii. 48.

216. CAMOCLADIA.

Linn. *Syst. No.* 1109. *p.* 1360. 861.

217. CAMPANULA.

Linn. *Gen.* 201. *Spec.* 163. *Syst.* 201. Hall. 490. Ludw. 63.
Mill. i. 162. iii. 48. Tourn. *tab.* 37. Weinm. *tab.* 286-294.

218. CAMPHOROSMA.

Linn. *Gen.* 152. *Spec.* 122. *Syst.* 152.

CAMPHORATA. Tourn. Ludw. 765. Mill. i. 165.

219. CANNA.

Linn. *Gen.* 1. *Spec.* 1. *Syst.* 1. Ludw. 168. Weinm. *tab.* 296-
298.

CANNACORUS. Tourn. *tab.* 192. Mill. i. 166.

220. CANNABIS.

Linn. *Gen.* 988. *Spec.* 1027. *Syst.* 988. Blackw. *tab.* 322. Ludw.
925. Mill. i. 165. Schæff. A. 278. Tourn. *tab.* 309. Weinm.
tab. 299.

221. CAPPARIS.

Linn. *Gen.* 567. *Spec.* 503. *Syst.* 567. Ludw. 458. Mill. i. 167.
Schæff. A. 156. Tourn. *tab.* 139. Weinm. *tab.* 303.

222. CAPRARIA.

Linn. *Gen.* 686. *Spec.* 628. *Syst.* 686. Ludw. 17.

SAMOLOIDES. Boerh. Mill. ii. 231.

223. CAPSICUM.

Linn. *Gen.* 225. *Spec.* 188. *Syst.* 225. Blackw. *tab.* 129. Ludw.
88. Mill. i. 168. Tourn. *tab.* 66.

224. CARDAMINE.

Linn. *Gen.* 727. *Spec.* 654. *Syst.* 727. Blackw. *tab.* 223. Hall.
557. Ludw. 415. Mill. i. 170. Tourn. *tab.* 109. Weinm. *tab.*
751. c.

225. CARDIOSPERMUM.

Linn. *Gen.* 447. *Spec.* 366. *Syst.* 447. Ludw. 442.

CORINDUM. Tourn. *tab.* 246. Mill. i. 229.

VESICARIA. Riv.

226. CARDUUS.

Linn. *Gen.* 832. *Spec.* 820. *Syst.* 832. Ludw. 321. Schæff. A.

97, 98. Weinm. *tab.* 308, *seq.*

CARDUUS. Vaill. Blackw. *tab.* 79. Hall. 676.

POLYACANTHA. Vaill.

SILYBUM. Vaill. Hall. 684.

CIRSIUM. Tourn. *tab.* 255. Mill. i. 210. iii. 64.

ERIOCEPHALUS. Vaill.

227. CAREX.

Linn. *Gen.* 928. *Spec.* 972. *Syst.* 928. Ludw. 869.

CYPEROIDES. Tourn. *tab.* 300. Hall. 234.

CAREX. Dill. Mich.

ULVA. Hall. 242.

228. CARICA.

Linn. *Gen.* 1000. *Spec.* 1036. *Syst.* 1000.

PAPAYA. Tourn. *tab.* 441. Ludw. 906. Mill. ii. 106.

229. CARLINA.

Linn. *Gen.* 836. *Spec.* 828. *Syst.* 836. Hall. 685. Ludw. 317.

Mill. i. 171, Schæff. A. 100. Tourn. *tab.* 285. Weinm. *tab.* 319.

CARLINA. Vaill.

CARLINOIDES. Vaill.

230. CARPESIUM.

Linn. *Gen.* 852. *Spec.* 859. *Syst.* 852.

CONYZOIDES. Tourn.

231. CARPINUS.

Linn. *Gen.* 952. *Spec.* 998. *Syst.* 952. Hall. 158. Ludw. 889.

Mill. i. 172. Tourn. *tab.* 348. Weinm. *tab.* 319. c.

OSTRYA. Mich.

232. CARTHAMUS.

Linn. *Gen.* 838. *Spec.* 830. *Syst.* 838. Ludw. 325. Mill. i. 173.

Schæff. A. 96. Tourn. *tab.* 258. Weinm. *tab.* 320.

CARTHAMUS. Vaill.

ATRACTYLIS. Vaill. Hall. 685.

CARTHAMOIDES. Vaill.

233. CARUM.

Linn. *Gen.* 327. *Spec.* 263. *Syst.* 327. Hall. 428. Ludw. 671.

CARUI. Tourn. *tab.* 160. Mill. i. 173. Schæff. A. 231. Weinm. *tab.* 321.

234. CARYOPHYLLUS.

Linn. *Gen.* 594. *Spec.* 515. *Syst.* 594. Ludw. 449. Weinm. *tab.* 324.

CARYOPHYLLUS AROMATICUS. Tourn. *tab.* 432. Blackw. *tab.* 338.

CARYOPHYLLODENDRON. Vaill.

235. CARYOTA.

Linn. *Gen.* 1092. *Spec.* 1189. *Syst.* 1092.

SCHUNDA-PANA. H. M.

236. CASSIA.

Linn. *Gen.* 461. *Spec.* 376. *Syst.* 461. Ludw. 641.

CASSIA. Tourn. *tab.* 392. Blackw. *tab.* 381. Mill. i. 179. iii. 51.
Weinm. *tab.* 340.

SENNA. Tourn. *tab.* 390. Mill. ii. 252. Weinm. *tab.* 915.

237. CASSINE.

Linn. *Gen.* 333. *Spec.* 268. *Syst.* 333. Ludw. 113.

MAUROCENIA. Linn. *edit. prior.* Mill. iii. 181.

238. CASSYTHA.

Linn. *Gen.* 52. *Spec.* 35. *Syst.* 52.

239. CATANANCHE.

Linn. *Gen.* 824. *Spec.* 812. *Syst.* 824. Ludw. 344.

CATANANCE. Tourn. *tab.* 271. Mill. i. 184.

240. CATESBÆA.

Linn. *Gen.* 121. *Spec.* 109. *Syst.* 121. Ludw. 1010. Mill. iii. 53.

241. CAUCALIS.

Linn. *Gen.* 294. *Spec.* 240. *Syst.* 294. Hall. 448. Ludw. 681.

Mill. i. 185. Tourn. *tab.* 171. Weinm. *tab.* 344.

TABLE II.

242. CEANOTHUS.

Linn. *Gen.* 237. *Spec.* 195. *Syst.* 237.

243. CECROPIA.

Linn. *Gen.* No. 1171. p. 1380. 1286.

244. CEDRELA.

Linn. *Syst.* No. 1124. p. 1365. 940.

CEDRO. Loeffl.

245. CELASTRUS.

Linn. *Gen.* 239. *Spec.* 196. *Syst.* 239. Ludw. 524. Mill. i. 189.
iii. 54.

EUONYMOIDES. Isn.

246. CELOSIA.

Linn. *Gen.* 255. *Spec.* 205. *Syst.* 255. Ludw. 515.

STACHYARPAGOPHORA. Vaill.

247. CELSIA.

Linn. *Gen.* 675. *Spec.* 621. *Syst.* 675. Ludw. 255. Mill. iii. 55.
THRYALLIS. Sig.

248. CELTIS.

Linn. *Gen.* 1012. *Spec.* 1043. *Syst.* 1012. Ludw. 782. Mill. i.
190. Tourn. *tab.* 383.

249. CENCHRUS.

Linn. *Gen.* 1017. *Spec.* 1049. *Syst.* 1017. Ludw. 845.

PANICASTRELLA. Mich.

250. CENTAUREA.

Linn. *Gen.* 880. *Spec.* 909. *Syst.* 880. Ludw. 366.

CENTAURIUM MAJUS. Tourn. *tab.* 256. Blackw. *tab.* 93. Mill. i.
191. iii. 56. Weinm. *tab.* 347.

JACEA. Tourn. *tab.* 254. Mill. i. 442. iii. 143.

CYANUS. Tourn. *tab.* 254. Blackw. *tab.* 66. 270. Mill. i. 155.

Schæff. A. 131. Weinm. *tab.* 451.

CALCITRAPA. Vaill. Hall. 689.

CALCITRAPOIDES. Vaill.

RHAPONTICUM. Vaill. Blackw. *tab.* 93. Hall. 687.

RHAPONTICOIDES. Vaill.

AMBERBOI. Vaill.

CROCODILUM. Vaill.

251. CENTUNCULUS.

Linn. *Gen.* 135. *Spec.* 116. *Syst.* 135. Ludw. 18.

ANAGALLIDASTRUM. Mich.

252. CEPHALANTHUS.

Linn. *Gen.* 105. *Spec.* 95. *Syst.* 105. Ludw. 293. Mill. iii. 56.

PLATANOCEPHALUS. Vaill.

253. CERASTIUM.

Linn. *Gen.* 518. *Spec.* 437. *Syst.* 518. Ludw. 570.

MYOSOTIS. Tourn. *tab.* 126. Hall. 383. Mill. ii. 63. Weinm.
tab. 740. a.

254. CERATOCARPUS.

Linn. *Gen.* 921. *Spec.* 969. *Syst.* 921.

255. CERATONIA.

Linn. *Gen.* 983. *Spec.* 1026. *Syst.* 983. Ludw. 923.

SILIQUA. Tourn. *tab.* 344. Blackw. *tab.* 209. Mill. ii. 260. iii.
58. Weinm. *tab.* 922. a.

256. CERATOPHYLLUM.

Linn. *Gen.* 944. *Spec.* 992. *Syst.* 944. Hall. 202. Ludw. 886.

HYDROCERATOPHYLLUM. Vaill.

DYCHOTOPHYLLUM. Dill.

257. CERBERA.

Linn. *Gen.* 260. *Spec.* 208. *Syst.* 260. Ludw. 79.

AHOVAL. Tourn. *tab.* 434. Mill. i. 23. iii. 9.

258. CERCIS.

Linn. *Gen.* 458. *Spec.* 374. *Syst.* 458.

SILQUASTRUM. Tourn. *tab.* 414. Ludw. 646. Mill. ii. 261. Weinm.
tab. 922. b.

259. CERINTHE.

Linn. *Gen.* 171. *Spec.* 136. *Syst.* 171. Ludw. 34. Hall. 515.

CERINTHE. Tourn. *tab.* 56. Mill. i. 198. Weinm. *tab.* 359, 360.
CERINTHOIDES. Boerh.

260. CEROPEGIA.

Linn. *Gen.* 266. *Spec.* 211. *Syst.* 266. Ludw. 1018.

261. CESTRUM.

Linn. *Gen.* 231. *Spec.* 191. *Syst.* 231. Ludw. 92. Mill. iii. 59.

262. CHÆROPHYLLUM.

Linn. *Gen.* 320. *Spec.* 258. *Syst.* 320. Hall. 452. Ludw. 675.
Mill. i. 198. Tourn. *tab.* 166.

CEREFOLIUM. Schæff. A. 224. Weinm. *tab.* 353.

263. CHAMÆROPS.

Linn. *Gen.* 1084. *Spec.* 1187. *Syst.* 1084. Ludw. 382. Weinm.
tab. 784.

CHAMÆRIPHES. Pont.

264. CHARA.

Linn. *Gen.* 1066. *Spec.* 1156. *Syst.* 1066. Hall. 196. Ludw. 954.
HIPPURIS. Dill.

265. CHEIRANTHUS.

Linn. *Gen.* 730. *Spec.* 661. *Syst.* 730. Ludw. 412.

GHEIRI. Schæff. A. 154. KEIRI. Rupp. Blackw. *tab.* 179.

LEUCOIUM. Tourn. *tab.* 107. Mill. i. 502. Weinm. *tab.* 646.

266. CHELIDONIUM.

Linn. *Gen.* 572. *Spec.* 505. *Syst.* 572.

CHELIDONIUM. Tourn. *tab.* 116. Blackw. *tab.* 91. Hall. 305.

Ludw. 447. Mill. i. 203. Schæff. A. 139. Weinm. *tab.* 366. a.

GLAUCIUM. Tourn. *tab.* 130. Hall. 304. Ludw. 452. Mill. i.
370.

267. CHELONE.

Linn. *Gen.* 666. *Spec.* 611. *Syst.* 666. Ludw. 240. Mill. i. 203.

ANONYMOS. Gron.

268. CHENOPODIUM.

Linn. *Gen.* 273. *Spec.* 218. *Syst.* 273. Tourn. *tab.* 288. Hall. 174.

Ludw. 776. Mill. i. 204. Blackw. *tab.* 311-314.

BONUS HENRICUS. Off. Schæff. A. 276.

269. CHERLERIA.

Linn. *Gen.* 506. *Spec.* 425. *Syst.* 506. Hall. 391. Ludw. 571.

270. CHIOCOCCA.

Linn. *Syst. No.* 1120. *p.* 1363. 917.

271. CHIONANTHUS.

Linn. *Gen.* 21. *Spec.* 8. *Syst.* 21. Ludw. 1009. Mill. iii. 61.

272. CHIRONIA.

Linn. *Gen.* 227. *Spec.* 189. *Syst.* 227. Ludw. 61.

273. CHONDRILLA.

Linn. *Gen.* 815. *Spec.* 796. *Syst.* 815. Hall. 755. Ludw. 334.
Mill. i. 205. iii. 61. Tourn. *tab.* 268. Weinm. *tab.* 308.

274. CHRYSANTHEMUM.

Linn. *Gen.* 866. *Spec.* 887. *Syst.* 866. Ludw. 349.

CHRYSANTHEMUM. Tourn. *tab.* 280. Mill. i. 206. Weinm. *tab.*
371, *seq.*

LEUCANTHEMUM. Tourn. *tab.* 492. Mill. i. 500. Weinm. *tab.* 238.

BELLIS MAJOR. Blackw. *tab.* 42. BELLIS PRATENSIS. Off.
Schæff. A. 126.

BELLIDOIDES. Vaill.

MATRICARIA. Vaill. Tourn. Hall. 718. Ludw. 349.

PYRETHRUM. Hall. 720.

275. CHRYSOBALANUS.

Linn. *Gen.* 585. *Spec.* 513. *Syst.* 585. Ludw. 598. Mill. iii. 62.

ICACO. Plum. Mill. i. 455.

276. CHRYSOCOMA.

Linn. *Gen.* 845. *Spec.* 840. *Syst.* 845. Hall. 703. Ludw. 307.
Mill. iii. 63.

CHRYSOCOME. Dill. COMA AUREA. Boerh. Mill. i. 219. Weinm.
tab. 406.

277. CHRYSOGONUM.

Linn. *Gen.* 883. *Spec.* 920. *Syst.* 883. Ludw. 364.

278. CHRYSOPHYLLUM.

Linn. *Gen.* 233. *Spec.* 192. *Syst.* 233. Ludw. 83. Mill. iii. 63.
CAINITO. Plum. Mill. i. 159.

279. CHRYSOSPLENIUM.

Linn. *Gen.* 403. *Spec.* 398. *Syst.* 493. Hall. 189. Ludw. 791.
Mill. i. 207. Tourn. *tab.* 60. Weinm. *tab.* 380.

280. CICER.

Linn. *Gen.* 783. *Spec.* 738. *Syst.* 783.

CICER. Tourn. *tab.* 210. Ludw. 480. Mill. i. 207. Schæff. A.
160. Weinm. *tab.* 380, c. d.

LENS. Tourn. *tab.* 210. Hall. 601. Ludw. 479. Mill. i. 497.
Schæff. A. 159. Weinm. *tab.* 637. a, b.

281. CICHORIUM.

Linn. *Gen.* 825. *Spec.* 813. *Syst.* 825. Blackw. *tab.* 177. 183.
Hall. 761. Ludw. 345. Mill. i. 207. Schæff. A. 110. Tourn.
tab. 272. Weinm. *tab.* 381.

282. CICUTA.

Linn. *Gen.* 316. *Spec.* 255. *Syst.* 316.

SIUM. Hall. 435. Ludw. 693. Mill. i. 209.

283. CINCHONA.

Linn. *Gen.* 208. *Spec.* 172. *Syst.* 208. Ludw. 1014.

QUINQUINA. Off. Weinm. *tab.* 367. a.

284. CINNA.

Linn. *Gen.* 15. *Spec.* 5. *Syst.* 15.

285. CIRCÆA.

Linn. *Gen.* 24. *Spec.* 9. *Syst.* 24. Hall. 456. Ludw. 369. Mill. i.
210. Tourn. *tab.* 155. Weinm. *tab.* 389.

286. CISSAMPELOS.

Linn. *Gen.* 993. *Spec.* 1031. *Syst. No.* 993. p. 1298.

CAAPEBA. Plum. Mill. i. 152.

287. CISSUS.

Linn. *Gen.* 137. *Spec.* 117. *Syst.* 137.

288. CISTUS.

Linn. *Gen.* 598. *Spec.* 523. *Syst.* 598.

CISTUS. Tourn. *tab.* 136. Blackw. *tab.* 197. Ludw. 590. Mill. i.
210. iii. 65. Weinm. *tab.* 390.

HELIANTHEMUM. Tourn. *tab.* 128. Hall. 358. Ludw. 583. Mill.
i. 395. Weinm. *tab.* 390. e.

289. CITHAREXYLUM.

Linn. *Gen.* 678. *Spec.* 625. *Syst.* 678.

290. CITRUS.

Linn. *Gen.* 807. *Spec.* 782. *Syst.* 807. Ludw. 605.

CITREUM. Tourn. *tab.* 395, 396. Blackw. *tab.* 361. Mill. i. 212.
Schæff. A. 192. Weinm. *tab.* 702. a.

AURANTIUM. Tourn. *tab.* 393, 394. Blackw. *tab.* 349. Mill. i. 94.
iii. 807. Schæff. A. 193. Weinm. *tab.* 701.

LIMON. Tourn. *tab.* 397. Blackw. *tab.* 362. Mill. i. 516. iii. 163.
Schæff. A. 194. Weinm. *tab.* 702. b.

291. CLATHRUS.

Linn. *Gen.* 1078. *Spec.* 1179. *Syst.* 1078. Battarr. *tab.* ii. Gled.
tab. iv. Ludw. 969. Schæff. B. § 70.

CLATHRUS. Mich.

CLATHROIDES. Mich. Hall. 9.

CLATHROIDASTRUM. Mich. Hall. 8.

SPHÆROCEPHALUS. Hall. 9.

BUXBAUMIA. Hall. 10.

292. CLAVARIA.

Linn. *Gen.* 1081. *Spec.* 1182. *Syst.* 1081. Ludw. 974. Schæff.
B. § 70.

CLAVARIA. Mich. Hall. 14. Battarr. *tab.* iii.

CORALLOIDES. Tourn. *tab.* 332. Hall. 14. Battarr. *tab.* i.

CORALLO-FUNGUS. Vaill.

FUNGOIDES. Dill.

293. CLAYTONIA.

Linn. *Gen.* 253. *Spec.* 204. *Syst.* 253. Ludw. 521.

294. CLEMATIS.

Linn. *Gen.* 616. *Spec.* 543. *Syst.* 616. Hall. 333. Ludw. 466.

Weinm. *tab.* 399, *seq.*

CLEMATITIS. Tourn. *tab.* 150. Mill. i. 213.

VITICELLA. Dill.

FLAMMULA. Rupp.

295. CLEOME.

Linn. *Gen.* 740. *Spec.* 671. *Syst.* 740. Ludw. 470.

SINAPISTRUM. Tourn. *tab.* 116. Mill. ii. 262.

296. CLERODENDRUM.

Linn. *Gen.* 707. *Spec.* 637. *Syst.* 707. Ludw. 264.

297. CLETHRA.

Linn. *Gen.* 489. *Spec.* 396. *Syst.* 489. Ludw. 648. Mill. iii. 66.

298. CLIFFORTIA.

Linn. *Gen.* 1004. *Spec.* 1038. *Syst.* 1004. Ludw. 934. Mill. iii.
66.

299. CLINPODIUM.

Linn. *Gen.* 644. *Spec.* 587. *Syst.* 644. Hall. 653. Ludw. 218.
Mill. i. 215. iii. 67. Tourn. *tab.* 92. Weinm. *tab.* 399.

300. CLITORIA.

Linn. *Gen.* 796. *Spec.* 753. *Syst.* 796. Ludw. 493.

TERNATEA. Tourn. Mill. ii. 318.

CLITORIUS. Dill.

301. CLUSIA.

Linn. *Gen.* 577. *Spec.* 509. *Syst. No.* 577. p. 1309. Ludw. 591.
Mill. iii. 67.

302. CLUTIA.

Linn. *Gen.* 1009. *Spec.* 1042. *Syst.* 1009. Ludw. 912. Mill. i.
215. iii. 68.

303. CLYPEOLA.

Linn. *Gen.* 723. *Spec.* 652. *Syst.* 723. Ludw. 400. Mill. iii. 69.
CLYPEOLA. Hall. 539. JONTHLASPI. Tourn. *tab.* 99. Mill. i. 459.

304. CNEORUM.

Linn. *Gen.* 47. *Spec.* 34. *Syst.* 47. Mill. iii. 69.
CHAMÆLEA. Tourn. *tab.* 421. Ludw. 375. Mill. i. 200.

305. CNICUS.

Linn. *Gen.* 833. *Spec.* 826. *Syst.* 883. Ludw. 323. Mill. i. 216.
iii. 69. Tourn. *tab.* 257.

ARCANA. Vaill.

306. COCCOLOBA.

Linn. *Syst No.* 1132. p. 1367. 1007.

307. COCHLEARIA.

Linn. *Gen.* 720. *Spec.* 647. *Syst.* 720. Blackw. *tab.* 218. 227.
Ludw. 425. Mill. i. 217. iii. 71. Schæff. A. 142. Tourn. *tab.*
101. Weinm. *tab.* 401. a.

308. COCOS.

Linn. *Gen.* 1088. *Spec.* 1188. *Syst.* 1088.

TENGA. H. M.

309. COFFEA.

Linn. *Gen.* 209. *Spec.* 172. *Syst.* 209. Blackw. *tab.* 337. Ludw.
75.

COFFE. Juss.

310. COIX.

Linn. *Gen.* 927. *Spec.* 972. *Syst.* 927. Mill. iii. 71. Ludw. 871.
LACRHYMA JOB. Tourn. *tab.* 306. Mill. i. 479. Weinm. *tab.*
618. c.

311. COLCHICUM.

Linn. *Gen.* 415. *Spec.* 341. *Syst.* 415. Hall. 282. Ludw. 129.
Mill. i. 218. Tourn. *tab.* 181, 182. Weinm. *tab.* 403, 404.

312. COLDENIA.

Linn. *Gen.* 159. *Spec.* 125. *Syst.* 159.

313. COLLINSONIA.

Linn. *Gen.* 38. *Spec.* 28. *Syst.* 38. Ludw. 175. Mill. iii. 71.

314. COLUMNEA.

Linn. *Gen.* 710. *Spec.* 638. *Syst.* 710. Ludw. 268. Mill. iii. 72.

315. COLUTEA.

Linn. *Gen.* 776. *Spec.* 723. *Syst.* 776. Hall. 575. Ludw. 506.
Mill. i. 218. iii. 72. Tourn. *tab.* 418. Weinm. *tab.* 406.

316. COMARUM.

Linn. *Gen.* 563. *Spec.* 502. *Syst.* 563. Hall. 337. Mill. iii. 73.
QUINQUEFOLIUM. Weinm. *tab.* 847. d.
POTENTILLA. Ludw. 621.

317. COMBRETUM.

Linn. *Syst. No.* 1129. p. 1366. 999.

318. COMMELINA.

Linn. *Gen.* 58. *Spec.* 40. *Syst.* 58. Mill. i. 220. iii. 73.
ZANONIA. Plum.

319. CONFERVA.

Linn. *Gen.* 1070. *Spec.* 1164. *Syst.* 1070. Hall. 1. Ludw. 995.

320. CONIUM.

Linn. *Gen.* 299. *Spec.* 243. *Syst.* 299.
CICUTA. Tourn. *tab.* 160. Hall. 433. Ludw. 691. Weinm. *tab.*
382, *seq.*

321. CONNARUS.

Linn. *Gen.* 744. *Spec.* 675. *Syst.* 744.

322. CONOCARPUS.

Linn. *Gen.* 213. *Spec.* 176. *Syst.* 213. Ludw. 774. Mill. iii. 74.
RUDBECKIA. Houst.

323. CONVALLARIA.

Linn. *Gen.* 383. *Spec.* 314. *Syst.* 383. Mill. iii. 75.
LILIUM CONVALLIUM. Tourn. *tab.* 14. Blackw. *tab.* 70. Hall.
286. Ludw. 127. Mill. i. 516. Schæff. A. 10. Weinm. *tab.*
653. b, c.

POLYGONATUM. Tourn. *tab.* 14. Blackw. *tab.* 251. Hall. 286.
Ludw. 126. Mill. ii. 155. SIGILLUM SALOMONIS. Off.
Schæff. A. 11. Weinm. *tab.* 920.

UNIFOLIUM. Dill. Hall. 280. Ludw. 394. Weinm. *tab.* 653. l.

324. CONVULVULUS.

Linn. *Gen.* 198. *Spec.* 153. *Syst.* 198. Blackw. *tab.* 38. Hall. 488.
Ludw. 66. Mill. i. 222. Tourn. *tab.* 17. Weinm. *tab.* 413, *seq.*

325. CONYZA.

Linn. *Gen.* 854. *Spec.* 861. *Syst.* 854. Blackw. *tab.* 103. Ludw.
306. Hall. 704. Mill. i. 225. Tourn. *tab.* 259. Weinm. *tab.*
422, *seq.*

326. CORCHORUS.

Linn. *Gen.* 599. *Spec.* 529. *Syst.* 599. Ludw. 588. Mill. i. 228.
Tourn. *tab.* 135.

327. CORDIA.

Linn. *Gen.* 228. *Spec.* 190. *Syst.* 228. Ludw. 78. Mill. iii. 76.
MYXA. Boerh. Mill. ii. 65. SEBESTENA. Dill. Blackw. *tab.*
398. Weinm. *tab.* 910. a.

328. COREOPSIS.

Linn. *Gen.* 879. *Spec.* 907. *Syst.* 879. Mill. iii. 77.

329. CORIANDRUM.

Linn. *Gen.* 318. *Spec.* 256. *Syst.* 318. Blackw. *tab.* 176. Ludw.
690. Mill. i. 229. Schæff. A. 223. Tourn. *tab.* 168. Weinm.
tab. 429. a.

330. CORIARIA.

Linn. *Gen.* 1002. *Spec.* 1037. *Syst.* 1002. Ludw. 576. Mill. i.
229. iii. 78.

331. CORIS.

Linn. *Gen.* 216. *Spec.* 177. *Syst.* 216. Ludw. 271. Mill. i. 230.
Tourn. *tab.* 423. Weinm. *tab.* 429. b.

332. CORISPERMUM.

Linn. *Gen.* 12. *Spec.* 4. *Syst.* 12. Ludw. 367. Mill. iii. 78.

333. CORNUCOPIÆ.

Linn. *Gen.* 67. *Spec.* 54. *Syst.* 67. Ludw. 813.

CORNUCOPIOIDES. Scheuchz.

334. CORNUS.

Linn. *Gen.* 139. *Spec.* 117. *Syst.* 139. Blackw. *tab.* 121. Hall.
463. Ludw. 395. Mill. i. 230. Schæff. A. 133. Tourn. *tab.*
410. Weinm. *tab.* 429. d.

VIRGA SANGUINEA. Dill.

335. CORNUTIA.

Linn. *Gen.* 684. *Spec.* 628. *Syst.* 684. Ludw. 266. Mill. i. 232.
GNANTHUS. Vaill.

336. CORONILLA.

Linn. *Gen.* 789. *Spec.* 742. *Syst.* 789. Ludw. 500.

CORONILLA. Tourn. *tab.* 419. Hall. 573. Mill. i. 235. Weinm.
tab. 430.

SECURIDACA. Tourn. *tab.* 224. Hall. 576.

EMERUS. Tourn. *tab.* 418. Hall. 573. Mill. i. 296.

337. CORRIGIOLA.

Linn. *Gen.* 340. *Spec.* 271. *Syst.* 340. Ludw. 536. Hall. 391.

POLYGONIFOLIA. Dill.

338. CORTUSA.

Linn. *Gen.* 181. *Spec.* 144. *Syst.* 181. Ludw. 47. Mill. i. 237.
iii. 79. Weinm. *tab.* 430. f.

339. CORYLUS.

Linn. *Gen.* 953. *Spec.* 998. *Syst.* 953. Blackw. *tab.* 293. Hall.
159. Ludw. 891. Mill. i. 237. Schæff. A. 290. Tourn. *tab.*
347. Weinm. *tab.* 431.

340. CORYMBIUM.

Linn. *Gen.* 895. *Spec.* 928. *Syst.* 895. Ludw. 1012.

341. CORYPHA.

Linn. *Gen.* 1086. *Spec.* 1187. *Syst.* 1086.

CODDA-PANNA. H. M.

342. COSTUS.

Linn. *Gen.* 3. *Spec.* 2. *Syst.* 3. Blackw. *tab.* 394. Ludw. 171.

Mill. iii. 80. Weinm. *tab.* 432. a.

343. COTULA.

Linn. *Gen.* 868. *Spec.* 891. *Syst.* 868. Ludw. 294.

ANANTHOCYLLUS. Vaill. LANCISIA. Pont.

344. COTYLEDON.

Linn. *Gen.* 512. *Spec.* 429. *Syst.* 512. Blackw. *tab.* 263. Ludw.

142. Mill. i. 238. Tourn. *tab.* 19. Weinm. *tab.* 433, *seq.*

345. CRAMBE.

Linn. *Gen.* 739. *Spec.* 671. *Syst.* 739. Ludw. 398.

CRAMBE. Tourn. *tab.* 100. Mill. i. 240.

RAPISTRUM. Tourn. *tab.* 99. Mill. ii. 191. Weinm. *tab.* 862. a, b.

346. CRANIOLARIA.

Linn. *Gen.* 670. *Spec.* 618. *Syst.* 670. Ludw. 189.

347. CRASSULA.

Linn. *Gen.* 352. *Spec.* 282. *Syst.* 352. Mill. i. 240. iii. 81.

COTYLEDON. Ludw. 142.

TELEPHIUM. Weinm. *tab.* 967. d.

348. CRATÆGUS.

Linn. *Gen.* 547. *Spec.* 475. *Syst.* 547. Blackw. *tab.* 149. Hall.

353. Ludw. 608. Mill. i. 241.

MESPILUS. Weinm. *tab.* 727.

349. CRATÆVA.

Linn. *Gen.* 528. *Spec.* 444. *Syst.* 528.

TAPIA. Plum. Ludw. 511. Mill. ii. 313.

350. CREPIS.

Linn. *Gen.* 819. *Spec.* 805. *Syst.* 819.

HIERACIOIDES. Vaill. Hall. 750. HIERACIUM. Ludw. 337.

351. CRESCENTIA.

Linn. *Gen.* 680. *Spec.* 626. *Syst.* 680. Ludw. 267.

CUIETE. Plum. Mill. i. 253.

352. CRESSA.

Linn. *Gen.* 277. *Spec.* 223. *Syst.* 277.

ANTHYLLIS. Magn.

353. CRINUM.

Linn. *Gen.* 366. *Spec.* 291. *Syst.* 366. Ludw. 1020. Mill. iii. 82.

354. CRITHMUM.

Linn. *Gen.* 303. *Spec.* 246. *Syst.* 303. Ludw. 666. Mill. i. 241.
Tourn. 169.

355. CROCUS.

Linn. *Gen.* 53. *Spec.* 36. *Syst.* 53. Blackw. *tab.* 137. Hall. 281.
Ludw. 7. Mill. i. 241. Schæff. A. 9. Tourn. *tab.* 183, 184.
Weinm. *tab.* 438.

356. CROTALARIA.

Linn. *Gen.* 771. *Spec.* 714. *Syst.* 771. Ludw. 484. Mill. i. 247.
Weinm. *tab.* 439.

357. CROTON.

Linn. *Gen.* 960. *Spec.* 1004. *Syst.* 960. Ludw. 863.RICINOIDES. Tourn. *tab.* 423. Mill. ii. 202.

BERNHARDIA. Houst. Mill. i. 120.

358. CRUCIANELLA.

Linn. *Gen.* 118. *Spec.* 108. *Syst.* 118.RUBEOLA. Tourn. *tab.* 50. Ludw. 12. Mill. ii. 209. Weinm. *tab.*
872.

359. CRUZETA.

Linn. *Syst. No.* 1117. *p.* 1362. 900.

360. CUCUBALUS.

Linn. *Gen.* 502. *Spec.* 414. *Syst.* 502. Blackw. *tab.* 268. Mill. i.
248. Tourn. *tab.* 176.

LYCHNIS. Ludw. 573.

361. CUCUMIS.

Linn. *Gen.* 969. *Spec.* 1011. *Syst.* 969. Ludw. 854.

CUCUMIS. Tourn. *tab.* 31. Blackw. *tab.* 4. Mill. i. 248. Schæff.
A. 16. Weinm. *tab.* 441.

COLOCYNTHIS. Tourn. Mill. i. 218. Schæff. A. 18. Weinm. *tab.*
405.

ANGURIA. Tourn. *tab.* 35. Blackw. *tab.* 157. Mill. i. 51. Ci-
TRULLUS. Rai. Schæff. A. 19.

MELO. Tourn. *tab.* 32. Blackw. *tab.* 329. Mill. ii. 32. iii. 185.
Schæff. A. 17. Weinm. *tab.* 721.

362. CUCURBITA.

Linn. *Gen.* 968. *Spec.* 1010. *Syst.* 968. Ludw. 855.

CUCURBITA. Tourn. *tab.* 36. Mill. i. 252. Schæff. A. 21. Weinm.
tab. 442.

PEPO. Tourn. *tab.* 33. Mill. ii. 112.

MELOPEPO. Tourn. *tab.* 34. Mill. ii. 40. Weinm. *tab.* 723.

363. CUMINUM.

Linn. *Gen.* 313. *Spec.* 254. *Syst.* 313. Ludw. 670. Mill. i. 254.
Schæff. A. 228. Weinm. *tab.* 447.

364. CUNILA.

Linn. *Syst. No.* 1106. *p.* 1359.

365. CUNONIA.

Linn. *Syst. No.* 1136. *p.* 1368. 1025.

366. CUPANIA.

Linn. *Gen.* 246. *Spec.* 200. *Syst.* 246. Ludw. 1033.

367. CUPRESSUS.

Linn. *Gen.* 958. *Spec.* 1002. *Syst.* 958. Blackw. *tab.* 127. Ludw.
875. Schæff. A. 295. Tourn. *tab.* 358. Weinm. *tab.* 448. a, b.

CYPRESSUS. Mill. i. 258.

TABLE II.

368. CURATELLA.

Linn. *Syst. No.* 1151. *p.* 1373. 1079.

369. CURCUMA.

Linn. *Gen.* 6. *Spec.* 2. *Syst.* 6. Ludw. 169. Weinm. *tab.* 448. e.

370. CUSCUTA.

Linn. *Gen.* 156. *Spec.* 124. *Syst.* 156. Hall. 468. Ludw. 26.
Schæff. A. 322. Tourn. *tab.* 422. Weinm. *tab.* 449. a.

371. CYANELLA.

Linn. *Gen.* 386. *Spec.* ... *Syst.* 386.

372. CYCAS.

Lin. *Gen.* 1087. *Spec.* 1188. *Syst.* 1087.

373. CYCLAMEN.

Linn. *Gen.* 184. *Spec.* 145. *Syst.* 184. Blackw. *tab.* 147. Hall.
499. Ludw. 48. Mill. i. 256. Schæff. A. 328. Tourn. *tab.*
68. Weinm. *tab.* 453.

374. CYMBARIA.

Linn. *Gen.* 669. *Spec.* 618. *Syst.* 669. Ludw. 234.

375. CYNANCHUM.

Linn. *Gen.* 268. *Spec.* 212. *Syst.* 268. Ludw. 101.

376. CYNARA.

Linn. *Gen.* 835. *Spec.* 827. *Syst.* 835. Ludw. 322.
CINARA. Tourn. *tab.* 253. Mill. i. 75. 209. Weinm. *tab.* 388.

377. CYNOGLOSSUM.

Linn. *Gen.* 168. *Spec.* 134. *Syst.* 168. Hall. 521.
CYNOGLOSSUM. Tourn. *tab.* 57. Blackw. *tab.* 249. Ludw. 33.
Mill. i. 257. Schæff. A. 40. Weinm. *tab.* 454.
OMPHALODES. Tourn. *tab.* 58. Ludw. 38. Mill. ii. 86.

378. CYNOMETRA.

Linn. *Gen.* 466. *Spec.* 382. *Syst.* 466.

CYNOMORIUM. Garc.

379. CYNOMORIUM.

Linn. *Gen.* 922. *Spec.* 970. *Syst.* 922. Ludw. 865.

380. CYNOSURUS.

Linn. *Gen.* 81. *Spec.* 72, 73. *Syst.* 81. Hall. 232. Ludw. 833.

381. CYFERUS.

Linn. *Gen.* 61. *Spec.* 44. *Syst.* 61. Blackw. *tab.* 316. Hall. 246.
Ludw. 840. Tourn. *tab.* 299. Weinm. *tab.* 455.

382. CYPRIPIEDUM.

Linn. *Gen.* 906. *Spec.* 951. *Syst.* 906.

CALCEOLUS. Tourn. *tab.* 249. Hall. 276. Ludw. 705. Mill. i. 160.

HELLEBORINE. Weinm. *tab.* 567. a.

383. CYTISUS.

Linn. *Gen.* 785. *Spec.* 739. *Syst.* 785. Hall. 591. Ludw. 478.

Mill. i. 261. iii. 85. Tourn. *tab.* 416. Weinm. *tab.* 456, *seq.*

384. DACTYLIS.

Linn. *Gen.* 80. *Spec.* 71. *Syst.* 80. Ludw. 824.

385. DALECHAMPIA.

Linn. *Gen.* 1022. *Spec.* 1054. *Syst.* 1022. Ludw. 888. Mill. i.
265.

CONVOLVULO-TITHYMALUS. Boerh.

386. DALIBARDA.

Linn. *Gen.* 555. *Spec.* 491. *Syst.* 555.

387. DAPHNE.

Linn. *Gen.* 436. *Spec.* 356. *Syst.* 436.

THYMELEA. Tourn. *tab.* 366. Blackw. *tab.* 62. Hall. 137. Ludw.

132. Mill. ii. 333. LAUREOLA. Weinm. *tab.* 633.

COCCOGNIDIUM. Off. Schæff. A. 1.

388. DATISCA.

Linn. *Gen.* 1003. *Spec.* 1037. *Syst.* 1003.

CANNABINA. Tourn. *tab.* 488. Mill. i. 166.

389. DATURA.

Linn. *Gen.* 218. *Spec.* 179. *Syst.* 218. Schæff. A. 329.

STRAMONIUM. Tourn. *tab.* 43, 44. Blackw. *tab.* 313. Hall. 489.

Ludw. 57. Mill. ii. 304. iii. 271. Weinm. *tab.* 951, *seq.*

390. DAUCUS.

Linn. *Gen.* 296. *Spec.* 242. *Syst.* 296. Hall. 450.

DAUCUS. Tourn. *tab.* 161. Ludw. 680. Mill. i. 265. Schæff. A. 242. Weinm. *tab.* 458.

VISNAGA. Riv. Ludw. 696. Mill. ii. 400.

LIBANOTIS. Riv. Hall. 450. Ludw. 682.

391. DELIMA.

Linn. *Gen.* 590. *Spec.* ... *Syst.* 590.

392. DELPHINIUM.

Linn. *Gen.* 602. *Spec.* 530. *Syst.* 602. Blackw. *tab.* 265. Hall. 314. Ludw. 740. Mill. i. 267. Tourn. *tab.* 241.

CONSOLIDA REGALIS. Off. Blackw. *tab.* 26. Schæff. A. 254. Weinm. *tab.* 408, 409. CALCATRIPPA. Heuch.

393. DENTARIA.

Linn. *Gen.* 726. *Spec.* 653. *Syst.* 726. Hall. 556. Ludw. 416. Mill. i. 268. Tourn. *tab.* 110. Weinm. *tab.* 460. c, d.

394. DIANTHERA.

Linn. *Gen.* 37. *Spec.* 27. *Syst. No.* 37. p. 850. Ludw. 1039.

395. DIANTHUS.

Linn. *Gen.* 500. *Spec.* 409. *Syst.* 500.

CARYOPHYLLUS. Tourn. *tab.* 174. Blackw. *tab.* 85. Mill. i. 174. Weinm. *tab.* 326, *seq.*

TUNICA. Dill. Hall. 381. Ludw. 563. Schæff. A. 181.

396. DIAPENSIA.

Linn. *Gen.* 177. *Spec.* 141. *Syst.* 177. Ludw. 1015.

397. DICTAMNUS.

Linn. *Gen.* 468. *Spec.* 383. *Syst.* 468. Blackw. *tab.* 75. Ludw. 649. Mill. i. 269. Schæff. A. 220. Weinm. *tab.* 461.
FRAXINELLA. Tourn. *tab.* 243. Hall. 311. Mill. i. 330.

398. DIGITALIS.

Linn. *Gen.* 676. *Spec.* 621. *Syst.* 676. Blackw. *tab.* 16. Hall. 616. Ludw. 249. Mill. i. 269. iii. 88. Tourn. *tab.* 73. Weinm. *tab.* 463.

399. DILLENIA.

Linn. *Gen.* 608. *Spec.* 535. *Syst.* 608. Ludw. 626.

400. DIODIA.

Lin. *Gen.* 114. *Spec.* 104. *Syst.* 114. Ludw. 250.

401. DIOSCOREA.

Linn. *Gen.* 995. *Spec.* 1032. *Syst.* 995. Ludw. 928. Mill. i. 270. iii. 88.

402. DIOSMA.

Linn. *Gen.* 241. *Spec.* 198. *Syst.* 241. Ludw. 546. Mill. iii. 89.

403. DIOSPYROS.

Linn. *Gen.* 1027. *Spec.* 1057. *Syst.* 1027. Mill. iii. 89.
GUAIIACANA. Tourn. *tab.* 371. Ludw. 135. Mill. i. 382. Weinm. *tab.* 559.

404. DIPSACUS.

Linn. *Gen.* 107. *Spec.* 97. *Syst.* 107. Blackw. *tab.* 50. Hall. 672. Ludw. 287. Mill. i. 271. Tourn. *tab.* 265. Weinm. *tab.* 465. *seq.*

405. DIRAC.

Linn. *Gen.* 437. *Spec.* 358. *Syst.* 437.

406. DODARTIA.

Linn. *Gen.* 698. *Spec.* 633. *Syst.* 698. Ludw. 251. Mill. i. 271. iii. 90. Tourn. *tab.* 478.

407. DODECATHEON.

Linn. *Gen.* 183. *Spec.* 144. *Syst.* 183.

MEADIA. Catesb.

408. DOLICHOS.

Linn. *Gen.* 778. *Spec.* 725. *Syst.* 778. Ludw. 492.

409. DORONICUM.

Linn. *Gen.* 862. *Spec.* 885. *Syst.* 862.DORONICUM. Tourn. *tab.* 277. Blackw. *tab.* 239. Ludw. 354.Mill. i. 275. Schæff. A. 118. Weinm. *tab.* 469.

BELLIDIASTRUM. Mich. Hall. 723.

410. DORSTENIA.

Linn. *Gen.* 147. *Spec.* 121. *Syst.* 147. Ludw. 940. Mill. i. 276.

411. DRABA.

Linn. *Gen.* 717. *Spec.* 642. *Syst.* 717. Hall. 538. Ludw. 428.

412. DRACOCEPHALUM.

Linn. *Gen.* 648. *Spec.* 594. *Syst.* 648.DRACOCEPHALON. Tourn. *tab.* 83. Ludw. 200. Mill. i. 277. iii. 91.Weinm. *tab.* 470.MOLDAVICA. Tourn. *tab.* 85. Ludw. 224. Mill. ii. 54. Weinm. *tab.* 719.

413. DRACONTIUM.

Linn. *Gen.* 916. *Spec.* 967. *Syst.* 916. Ludw. 550.

414. DROSERA.

Linn. *Gen.* 351. *Spec.* 281. *Syst.* 351.ROS SOLIS. Tourn. *tab.* 127. RORELLA. Rupp. Hall. 372. Ludw.544. Schæff. A. 179. Weinm. *tab.* 872. c, d.

415. DRYAS.

Linn. *Gen.* 562. *Spec.* 501. *Syst.* 562. Hall. 335. Ludw. 757.CARYOPHYLLATÆ SPECIES. Tourn. *tab.* 151. B. E.

416. DRYPIS.

Linn. *Gen.* 501. *Spec.* 413. *Syst.* 501. Ludw. 574.

417. DURANTA.

Linn. *Gen.* 704. *Spec.* 637. *Syst.* 704.

CASTOREA. Plum. Ludw. 1050. Mill. i. 184.

418. EBENUS.

Linn. *Spec.* 764. *Syst. No.* 1159. *p.* 1376. 1176.

419. ECHINOPHORA.

Linn. *Gen.* 292. *Spec.* 239. *Syst.* 292. Ludw. 1051. Mill. i.288. Tourn. *tab.* 423.

420. ECHINOPS.

Linn. *Gen.* 829. *Spec.* 814. *Syst.* 829.ECHINOPUS. Tourn. *tab.* 262. Ludw. 300. Mich. i. 288.

421. ECHIUM.

Linn. *Gen.* 175. *Spec.* 139. *Syst.* 175. Blackw. *tab.* 299 Hall.513. Ludw. 270. Mill. i. 289. Tourn. *tab.* 54. Weinm. *tab.*

474, 475.

422. EHRETIA.

Linn. *Syst. No.* 1119. *p.* 1363. 936.

423. ELÆAGNUS.

Linn. *Gen.* 148. *Spec.* 121. *Syst.* 148. Ludw. 768. Mill. i. 291.Tourn. *tab.* 489.

424. ELEOCARPUS.

Linn. *Gen.* 589. *Spec.* 515. *Syst.* 589.

425. ELATE.

Linn. *Gen.* 1091. *Spec.* 1189. *Syst.* 1091.

426. ELATINE.

Linn. *Gen.* 451. *Spec.* 367. *Syst.* 451. Ludw. 444.

POTAMOPITHYS. Buxb. ALSINASTRUM. Vaill.

427. ELEPHANTOPUS.

Linn. *Gen.* 827. *Spec.* 814. *Syst.* 827. Ludw. 309. Mill. i. 292.

428. ELLISIA.

Linn. *Syst. No.* 1156. *p.* 1375. 1121.

429. ELVELA.

Linn. *Gen.* 1079. *Spec.* 1180. *Syst.* 1079. *Gled. tab.* ii. Schæff.
B. §. 70.

FUNGOIDASTER. Mich. 82. Hall. 58. Ludw. 970.

FUNGOIDES. Mich. 86. Ludw. 971.

430. ELYMUS.

Linn. *Gen.* 91. *Spec.* 83. *Syst.* 91.

431. EMPETRUM.

Linn. *Gen.* 977. *Spec.* 1022. *Syst.* 977. Hall. 162. Ludw. 860.
Mill. i: 296. Tourn. *tab.* 421.

432. EPHEDRA.

Linn. *Gen.* 1007. *Spec.* 1040. *Syst.* 1007. Hall. 145. Ludw. 931.
Mill. i. 297. Tourn. *tab.* 477.

433. EPIDENDRUM.

Linn. *Gen.* 907. *Spec.* 952. *Syst.* 907.

VANILLA. Plum. Ludw. 707. Mill. ii. 366. Weinm. *tab.* 1002. c.

434. EPIGÆA.

Linn. *Gen.* 486. *Spec.* 395. *Syst.* 486.

MEMECYLUM. Mitch.

435. EPILOBIUM.

Linn. *Gen.* 426. *Spec.* 347. *Syst.* 426. Hall. 408. Mill. iii. 95.
CHAMÆNERION. Tourn. *tab.* 157. Ludw. 435. Mill. i. 201.
Weinm. *tab.* 688. a.

436. EPIMEDIUM.

Linn. *Gen.* 138. *Spec.* 117. *Syst.* 138. Ludw. 389. Mill. i. 297.
Tourn. *tab.* 117. Weinm. *tab.* 478. b.

437. EQUISETUM.

Linn. *Gen.* 1033. *Spec.* 1061. *Syst.* 1033. Blackw. *tab.* 217. Hall.
142. Ludw. 953. Schæff. A. 303. Tourn. *tab.* 307. Weinm.
tab. 479, 480.

438. ERANTHEMUM.

Linn. *Gen.* 23. *Spec.* 9. *Syst.* 23.

439. ERICA.

Linn. *Gen.* 435. *Spec.* 352. *Syst.* 435. Hall. 418. Mill. i. 300.

Tourn. *tab.* 373. a. Weinm. *tab.* 481.

ERICOIDES. Ludw. 131.

440. ERIGERON.

Linn. *Gen.* 855. *Spec.* 863. *Syst.* 855. Hall. 724.

SENECIONIS SPECIES. Dill. CONYZA. Ludw. 306. CONYZOIDES.

Dill. CONYZELLA. Dill.

441. ERINUS.

Linn. *Gen.* 689. *Spec.* 630. *Syst.* 689. Ludw. 258.

AGERATUM. Tourn. *tab.* 422. Hall. 629. Mill. i. 20. iii. 9. Weinm. *tab.* 29. a.

442. ERIOCAULON.

Linn. *Gen.* 95. *Spec.* 87. *Syst.* 95. Ludw. 838.

443. ERIOCEPHALUS.

Linn. *Gen.* 890. *Spec.* 926. *Syst.* 890.

444. ERIOPHORUM.

Linn. *Gen.* 63. *Spec.* 52. *Syst.* 63.

LINAGROSTIS. Mich. Tourn. Hall. 250. Ludw. 844.

445. ERITHALIS.

Linn. *Syst.* No. 1123. p. 1364. 930.

446. ERVUM.

Linn. *Gen.* 784. *Spec.* 738. *Syst.* 784. Hall. 602. Ludw. 482.

Mill. i. 301. Schæff. A. 161. Tourn. *tab.* 221. Weinm. *tab.* 484. a.

447. ERYNGIUM.

Linn. *Gen.* 287. *Spec.* 232. *Syst.* 287. Blackw. *tab.* 297. Hall.

455. Ludw. 528. Mill. i. 302. iii. 97. Schæff. A. 243. Tourn. *tab.* 173. Weinm. *tab.* 484, 485.

448. ERYSIMUM.

Linn. *Gen.* 729. *Spec.* 660. *Syst.* 729. Blackw. *tab.* 28. Ludw. 409. Mill. i. 302. Schæff. A. 148. Tourn. *tab.* 111. Weinm. *tab.* 487. a.

SISYMBRIUM. Hall. 547.

ALIARIA. Rai. Blackw. *tab.* 372. Weinm. *tab.* 37.

449. ERYTHRINA.

Linn. *Gen.* 762. *Spec.* 706. *Syst.* 762. Ludw. 498.

CORALLODELLON. Tourn. *tab.* 446. Mill. i. 227.

CORAL. Dill.

450. ERYTHRONIUM.

Linn. *Gen.* 375. *Spec.* 305. *Syst.* 375. Hall. 290. Ludw. 720. DENS CANIS. Tourn. *tab.* 202. Mill. i. 268. Weinm. *tab.* 460. a, b.

451. ERYTHROXYLON.

Linn. *Syst. No.* 1137. p. 1369. 1035.

452. EUGENIA.

Linn. *Gen.* 542. *Spec.* 470. *Syst.* 542. Ludw. 454.

453. EUONYMUS.

Linn. *Gen.* 240. *Spec.* 197. *Syst.* 240. Hall. 423. Ludw. 527. Mill. i. 303. Tourn. *tab.* 388. Weinm. *tab.* 494.

454. EUPATORIUM.

Linn. *Gen.* 842. *Spec.* 836. *Syst.* 842. Blackw. *tab.* 110. Hall. 704. Ludw. 308. Mill. i. 304. Schæff. A. 108. Tourn. *tab.* 259. Weinm. *tab.* 495, 496.

455. EUPHORBIA.

Linn. *Gen.* 536. *Spec.* 450. *Syst.* 536. Ludw. 166. Mill. iii. 98. EUPHORBIVM. Isn. Blackw. *tab.* 339, 340. Mill. i. 305. Weinm. *tab.* 497, 498.

ESULA. Riv. Schæff. A. 47. Weinm. *tab.* 487, *seq.* TITHYMALUS. Tourn. *tab.* 18. Hall. 189. Mill. ii. 338. iii. 278.

TITHYMALOIDES. Tourn. Mill. ii. 338.

456. EUPHRASIA.

Linn. *Gen.* 659. *Spec.* 604. *Syst.* 659.

EUPHRASIA. Tourn. *tab.* 78. Hall. 628. Ludw. 248. Mill. i. 306.

Schæff. A. 60. Weinm. *tab.* 499.

ODONTITIS. Dill. Hall. 627.

457. EXACUM.

Linn. *Syst.* 132. *Spec.* 112. *Syst.* 132.

458. EXCÆCARIA.

Linn. *Gen. post No.* 978. p. 1288.

459. FAGARA.

Linn. *Syst. No.* 1115. p. 1362. 897.

460. FAGONIA.

Linn. *Gen.* 475. *Spec.* 386. *Syst.* 475. Ludw. 559. Mill. i. 308.

Tourn. *tab.* 141.

461. FAGUS.

Linn. *Gen.* 951. *Spec.* 997. *Syst.* 951.

FAGUS. Tourn. *tab.* 351. Hall. 160. Ludw. 894. Mill. i. 309.

Weinm. *tab.* 501. a.

CASTANEA. Tourn. *tab.* 352. Blackw. *tab.* 330. Hall. 160. Ludw.

895. Mill. i. 182. iii. 52. Weinm. *tab.* 343.

462. FERULA.

Linn. *Gen.* 305. *Spec.* 246. *Syst.* 305. Ludw. 662. Mill. i. 310.

Tourn. *tab.* 170. Weinm. *tab.* 504.

463. FESTUCA.

Linn. *Gen.* 82. *Spec.* 73. *Syst.* 82. Hall. 209. Ludw. 835.

464. FEVILLEA.

Linn. *Gen.* 972. *Spec.* 1013. *Syst. No.* 972. p. 1292. Ludw. 850.

NHANDIROBA. Plum.

465. FICUS.

Linn. *Gen.* 1032. *Spec.* 1059. *Syst.* 1032. Blackw. *tab.* 125.

Ludw. 866. Mill. i. 319. iii. 105. Schæff. A. 264. Tourn. *tab.*
420. Weinm. *tab.* 508.

CAPRIFICUS. Pont.

466. FILAGO.

Linn. *Gen.* 891. *Spec.* 927. *Syst.* 891.

467. FLAGELLARIA.

Linn. *Gen.* 406. *Spec.* 333. *Syst.* 406.

468. FONTINALIS.

Linn. *Gen.* 1053. *Spec.* 1107. *Syst.* 1053. Hall. 95. Ludw. 961.

469. FRAGARIA.

Linn. *Gen.* 558. *Spec.* 494. *Syst.* 558. Blackw. *tab.* 77. Hall. 343.
Mill. i. 328. Schæff. A. 208. Tourn. *tab.* 152. Weinm. *tab.*
514.

POTENTILLA. Ludw. 621.

470. FRANKENIA.

Linn. *Gen.* 401. *Spec.* 331. *Syst.* 401.

FRANCA. Mich. Ludw. 554.

471. FRAXINUS.

Linn. *Gen.* 1026. *Spec.* 1057. *Syst.* 1026. Blackw. *tab.* 328. Hall.
167. Ludw. 910. Mill. i. 330. Schæff. A. 291. Tourn. *tab.*
343. Weinm. *tab.* 515.

ORNUS. Mich.

472. FRITILLARIA.

Linn. *Gen.* 372. *Spec.* 303. *Syst.* 372.

FRITILLARIA. Tourn. *tab.* 201. Hall. 290. Ludw. 718. Mill. i.
331. Weinm. *tab.* 515.

CORONA IMPERIALIS. Tourn. *tab.* 197, 198. Ludw. 719. Mill. i.
232. Weinm. *tab.* 661.

473. FUCHSIA.

Linn. *Gen.* 1097. *Spec.* 1191. *Syst. No.* 1097. p. 893. Ludw.
1052. Mill. i. 336.

474. FUCUS.

Linn. *Gen.* 1068. *Spec.* 1158. *Syst.* 1068.

475. FUMARIA.

Linn. *Gen.* 760. *Spec.* 699. *Syst.* 760. Hall. 604. Ludw. 471.
 FUMARIA. Tourn. *tab.* 237. Blackw. *tab.* 237. Mill. i. 337.
 Schæff. A. 157. Weinm. *tab.* 430.
 CAPNOIDES. Tourn. *tab.* 237. Mill. i. 166.
 CORYDALIS. Dill. CYSTICAPNOS. Boerh. Mill. i. 261.
 PSEUDOFUMARIA. Riv.
 CAPNORCHIS. Boerh.

476. GALANTHUS.

Linn. *Gen.* 362. *Spec.* 288. *Syst.* 362. Ludw. 722. Mill. iii. 114.
 LEUCOIUM. Weinm. *tab.* 642. b.

477. GALAX.

Linn. *Gen.* 244. *Spec.* 200. *Syst.* 244.

VITICELLA. Mitch.

478. GALEGA.

Linn. *Gen.* 770. *Spec.* 714. *Syst.* 770. Blackw. *tab.* 92. Hall.
 576. Ludw. 497. Mill. i. 341. Schæff. A. 162. Tourn. *tab.*
 222. Weinm. *tab.* 528.

INDIGO. Isn.

CRACCA. Linn. *Syst.* 1172. *Spec.* 752.

479. GALENIA.

Linn. *Gen.* 443. *Spec.* 359. *Syst.* 443. Ludw. 792. Mill. iii. 115.
 SHERARDIA. Pont.

480. GALEOPSIS.

Linn. *Gen.* 637. *Spec.* 579. *Syst.* 637. Hall. 644. Ludw. 193.
 Mill. i. 342.

TETRAHIT. Dill. GALEORDOLON. Dill.

481. GALIUM.

Linn. *Gen.* 117. *Spec.* 105. *Syst.* 117. Hall. 458, *seq.*
 GALLIUM. Tourn. *tab.* 39. Blackw. *tab.* 168. Ludw. 11. Mill. i.
 343. Weinm. *tab.* 530.

APARINE. Blackw. *tab.* 39. Hall. 458. Ludw. 10. Mill. iii. 22.
Weinm. *tab.* 146, 147.

482. GARCINIA.

Linn. *Gen.* 526. *Spec.* 443. *Syst.* 526. Ludw. 460.

MANGOSTANS. Garc.

483. GARIDELLA.

Linn. *Gen.* 507. *Spec.* 425. *Syst.* 507. Ludw. 612. Mill. i. 343.
Tourn. *tab.* 430.

484. GAULTHERIA.

Linn. *Gen.* 487. *Spec.* 395. *Syst.* 487.

485. GAURA.

Linn. *Gen.* 425. *Spec.* 347. *Syst.* 425.

486. GENIPA.

Linn. *Gen.* 229. *Spec.* ... *Syst.* 229. 981. Ludw. 94. Tourn. *tab.*
436, 437.

487. GENISTA.

Linn. *Gen.* 766. *Spec.* 709. *Syst.* 766. Hall. 592. Ludw. 477.
SPARTIUM. Tourn. *tab.* 412. Mill. ii. 284. Weinm. *tab.* 533.
GENISTELLA. Tourn. *tab.* 413.

488. GENTIANA.

Linn. *Gen.* 285. *Spec.* 227. *Syst.* 285. Hall. 473.

GENTIANA. Tourn. *tab.* 40. Ludw. 97. Mill. i. 350. Weinm. *tab.*
534, 535.

CENTAURIUM MINUS. Tourn. *tab.* 48. Ludw. 106. Mill. i. 192.
Schæff. A. 326. Weinm. *tab.* 348.

489. GERANIUM.

Linn. *Gen.* 476. *Spec.* 676. *Syst.* 746. Blackw. *tab.* 58. 150. Hall.
365. Ludw. 640. Mill. i. 351. Schæff. A. 221. Tourn. *tab.*
142. Weinm. *tab.* 535, *seq.*

PELARGONIUM. Burm.

GRUINALIS. Rupp. Riv.

490. GERARDIA.

Linn. *Gen.* 665. *Spec.* 610. *Syst.* 665. Ludw. 253.

491. GESNERIA.

Linn. *Gen.* 667. *Spec.* 612. *Syst.* 667.

GESNERA. Plum. Ludw. 257. Mill. i. 355. iii. 117.

492. GETHYLLIS.

Linn. *Gen.* 523. *Spec.* 442. *Syst.* 523. Ludw. 158.

MANGLES. Plum.

493. GEUM.

Linn. *Gen.* 561. *Spec.* 500. *Syst.* 561.

CARYOPHYLLATA. Tourn. *tab.* 151. *A. D. F. G.* Blackw. *tab.* 253.

Hall. 335. Ludw. 622. Mill. i. 173. Schæff. *A.* 209. Weinm. *tab.* 323.

494. GLADIOLUS.

Linn. *Gen.* 55. *Spec.* 36. *Syst.* 55. Hall. 279. Ludw. 187. Mill. i. 359. iii. 118. Tourn. *tab.* 190. Weinm. *tab.* 546.

495. GLAUX.

Linn. *Gen.* 257. *Spec.* 207. *Syst.* 257. Ludw. 46. Mill. i. 371. Tourn. *tab.* 60.

496. GLECOMA.

Linn. *Gen.* 634. *Spec.* 578. *Syst.* 634.

HEDERA TERRESTRIS. Off. Blackw. *tab.* 225. Schæff. *A.* 81. Weinm. *tab.* 564. b, c, d.

CHAMÆCLEMA. Boerh. Hall. 652. Ludw. 219. Mill. i. 199.

497. GLEDITSIA.

Linn. *Gen.* 1027. *Spec.* 1056. *Syst.* 1025. Ludw. 911.

MELILOBUS. Mitch.

498. GLINUS.

Linn. *Gen.* 537. *Spec.* 463. *Syst.* 537.

499. GLOBULARIA.

Linn. *Gen.* 106. *Spec.* 95. *Syst.* 106. Hall. 667. Ludw. 288.

Weinm. *tab.* 238. a. Mill. i. 371. Tourn. *tab.* 265.

ALYPUM. Niss.

TABLE II.

500. GLORIOSA.

Linn. *Gen.* 374. *Spec.* 305. *Syst.* 374. Mill. iii. 120.
METHONICA. Tourn. *A. G.* Ludw. 716. Mill. ii. 47.

501. GLYCINE.

Linn. *Gen.* 797. *Spec.* 753. *Syst.* 797. Mill. iii. 121.
APIOS. Boerh. Ludw. 510. Mill. i. 65. Weinm. *tab.* 149.

502. GLYCYRRHIZA.

Linn. *Gen.* 788. *Spec.* 741. *Syst.* 788. Ludw. 633. Mill. i. 372.
Tourn. *tab.* 210. Weinm. *tab.* 547.
LIQUIRITIA. Off. Schæff. A. 222.

503. GMELINA.

Linn. *Gen.* 681. *Spec.* 626. *Syst.* 681. Ludw. 1024.
MICHELIA. Amm.

504. GNAPHALIUM.

Linn. *Gen.* 850. *Spec.* 850. *Syst.* 850. Hall. 698. Ludw. 305.
Schæff. A. 101. Weinm. *tab.* 549, *seq.*
ELICHRYSUM. Tourn. *tab.* 259. Mill. i. 293. iii. 95. HELICHRY-
SUM. Vaill.
FILAGO. Tourn. Vaill. *tab.* 259.
HELICHRYSOIDES. Vaill.

505. GNIDIA.

Linn. *Gen.* 438. *Spec.* 358. *Syst.* 438.
STRUTHIA. Roy.

506. GOMPHRENA.

Linn. *Gen.* 279. *Spec.* 224. *Syst.* 279.
AMARANTHOIDES. Tourn. *tab.* 429. Mill. i. 32. iii. 13.
CARAXERON. Vaill.

507. GORTERIA.

Linn. *Syst. No.* 1163. *Spec.* 1377. 1229.

508. GOSSYPIU.

Linn. *Gen.* 755. *Spec.* 693. *Syst.* 755. Blackw. *tab.* 357. Ludw.
150. Mill. iii. 121.
XYLON. Tourn. *tab.* 27. Mill. ii. 529. Weinm. *tab.* 551. e.

509. GRATIOLOA.

Linn. *Gen.* 27. *Spec.* 17. *Syst.* 27. Hall. 617. Ludw. 182. Schæff.
A. 61. Weinm. *tab.* 558. a.

510. GREWIA.

Linn. *Gen.* 914. *Spec.* 964. *Syst.* 914. Ludw. 601. Mill. iii. 124.

511. GRIAS.

Linn. *Syst.* No. 1145. p. 1371. 1075.

512. GRISLEA.

Linn. *Gen.* 427. *Spec.* 348. *Syst.* 427. Ludw. 439.

513. GRONOVIA.

Linn. *Gen.* 248. *Spec.* 202. *Syst.* 248. Ludw. 520. Mill. iii. 124.

514. GUAIAACUM.

Linn. *Gen.* 465. *Spec.* 381. *Syst.* 465. Blackw. *tab.* 350. Ludw.
599. Mill. i. 383. Weinm. *tab.* 560.

LIGNUM VITÆ, SANCTUM. Off.

515. GUETTARDA.

Linn. *Gen.* 943. *Spec.* 991. *Syst.* 943.

516. GUILANDINA.

Linn. *Gen.* 464. *Spec.* 381. *Syst.* 464.

BONDUC. Plum. Ludw. 914. Mill. i. 133. iii. 41.

517. GUNDELIA.

Linn. *Gen.* 828. *Spec.* 814. *Syst.* 828. Mill. i. 387. iii. 127.
Tourn. *tab.* 486.

HACUB. Vaill. Ludw. 297.

518. GYPSOPHILA.

Linn. *Gen.* 498. *Spec.* 406. *Syst.* 498.

519. HEMANTHUS.

Linn. *Gen.* 394. *Spec.* 325. *Syst.* 394. Ludw. 124. Mill. i. 387.
iii. 127. Tourn. *tab.* 433. Weinm. *tab.* 562.

DRANCUNCULOIDES. Boerh.

TABLE II.

520. HÆMATOXYLUM.

Linn. *Gen.* 471. *Spec.* 384. *Syst.* 471. Ludw. 555. Mill. iii. 128.
CAMPECIA. Sloan.

521. HALESIA.

Linn. *Syst. Gen. No.* 1138. *p.* 1369. 1044.

522. HALLERIA.

Linn. *Gen.* 679. *Spec.* 625. *Syst.* 679. Ludw. 269. Mill. iii. 129.

523. HAMAMELIS.

Linn. *Gen.* 155. *Spec.* 124. *Syst.* 155. Ludw. 396. Mill. iii. 129.
TRILOPUS. Mitch.

524. HARTOGIA.

Linn. *Syst. No.* 1126. *p.* 1365. 939.

525. HASSELQUISTIA.

Linn. *Syst. p.* 957.

526. HEBENSTRETIA.

Linn. *Gen.* 688. *Spec.* 629. *Syst.* 688. Ludw. 236.

527. HEDERA.

Linn. *Gen.* 249. *Spec.* 202. *Syst.* 249. Blackw. *tab.* 188. Hall.
165. Ludw. 526. Mill. i. 390. Schæff. A. 176. Tourn. *tab.*
384. Weinm. *tab.* 563. d.

528. HEDYOTIS.

Linn. *Gen.* 110. *Spec.* 101. *Syst.* 110.

529. HEDYSARUM.

Linn. *Gen.* 793. *Spec.* 745. *Syst.* 793.

HEDYSARUM. Tourn. *tab.* 225. Hall. 577. Ludw. 504. Mill. i.
391. Weinm. *tab.* 565.

ONOBRYCHIS. Tourn. *tab.* 211. Hall. 577. Ludw. 472. Mill. ii.
87. Weinm. *tab.*

ALHAGI. Tourn. *tab.* 489. Ludw. 504.

530. HELENIUM

Linn. *Gen.* 863. *Spec.* 886. *Syst.* 863. Mill. i. 393. Weinm. *tab.*
565. d.

HELENIASTRUM. Vaill. Mill. i. 392. iii. 135.

531. HELIANTHUS.

Linn. *Gen.* 877. *Spec.* 904. *Syst.* 877. Ludw. 360. Mill. iii. 135.

CORONA SOLIS. Tourn. *tab.* 279. Mill. i. 233.

532. HELICTERES.

Linn. *Gen.* 913. *Spec.* 963. *Syst.* 913. Ludw. 651.

ISORA. Plum. Mill. i. 463.

533. HELIOCARPUS.

Linn. *Gen.* 533. *Spec.* 448. *Syst.* 533. Ludw. 462.

MONTIA. Houst. Mill. ii. 57. iii. 196.

534. HELIOTROPIUM.

Linn. *Gen.* 164. *Spec.* 130. *Syst.* 164. Hall. 520. Mill. i. 398.

Tourn. *tab.* 57. Weinm. *tab.* 566.

535. HELLEBORUS.

Lin. *Gen.* 622. *Spec.* 557. *Syst.* 622. Blackw. *tab.* 57. Hall. 317.

Ludw. 625. Mill. i. 400. iii. 136. Schæff. A. 172. Tourn.

tab. 144. Weinm. *tab.* 569.

536. HELONIAS.

Linn. *Gen.* 416. *Spec.* 342. *Syst.* 416.

537. HEMEROCALLIS.

Linn. *Gen.* 391. *Spec.* 324. *Syst.* 391. Hall. 290. Mill. iii. 137.

LILIO-ASPHODELUS. Tourn. *tab.* 179. Ludw. 120. Mill. i. 508.

Weinm. *tab.* 651.

LILIASTRUM. Tourn. *tab.* 194. Ludw. 714.

538. HEMIONITIS.

Linn. *Gen.* 1040. *Spec.* 1077. *Syst.* 1040. Ludw. 948. Mill. i.

401.

539. HERACLEUM.

Linn. *Gen.* 307. *Spec.* 249. *Syst.* 307.

SPHONDYLIVM. Tourn. *tab.* 170. Hall. 447. Ludw. 664. Mill.

ii. 286. Weinm. *tab.* 944.

540. HERNANNIA.

Linn. *Gen.* 742. *Spec.* 673. *Syst.* 742. Ludw. 522. Mill. i. 402.
iii. 138. Tourn. *tab.* 432.

541. HERMANDIA.

Linn. *Gen.* 931. *Spec.* 981. *Syst.* 931. Ludw. 1053. Mill. i. 403.

542. HERNIARIA.

Linn. *Gen.* 272. *Spec.* 218. *Syst.* 272. Blackw. *tab.* 320. Hall.
182. Ludw. 779. Mill. i. 404. Schæff. A. 274. Tourn. *tab.*
288. Weinm. *tab.* 570. f, g.

543. HESPERIS.

Linn. *Gen.* 731. *Spec.* 663. *Syst.* 731. Hall. 563. Ludw. 411.
Mill. i. 404. iii. 139. Tourn. *tab.* 108. Weinm. *tab.* 571.

544. HEUCHERA.

Linn. *Gen.* 283. *Spec.* 226. *Syst.* 283. Ludw. 530.

545. HIBISCUS.

Linn. *Gen.* 756. *Spec.* 693. *Syst.* 756. Mill. iii. 139.
KETMIA. Tourn. *tab.* 26. Ludw. 151. Mill. i. 472. Weinm. *tab.*
617.

MALVAVISCUS. Dill.

TRIONUM. Linn. *edit. prior.*

546. HIERACIUM.

Linn. *Gen.* 818. *Spec.* 799. *Syst.* 818. Hall. 742. Ludw. 337.
HIERACIUM. Tourn. *tab.* 267. Mill. i. 405. Weinm. *tab.* 573, *seq.*
AURICULA MURIS. Rai. PILOSELLA. Vaill. Mill. i. 97. Schæff.
A. 114. Weinm. *tab.* 206.

547. HIPPOCRATEA.

Linn. *Gen.* 1098. *Spec.* 1191. *Syst. No.* 1098. p. 988.
COA. Plum. Ludw. 1054. Mill. i. 216.

548. HIPPOCREPIS.

Linn. *Gen.* 791. *Spec.* 744. *Syst.* 791. Ludw. 503.

FERRUM EQUINUM. Tourn. *tab.* 225. Hall. 573. Mill. i. 310.
Weinm. *tab.* 504. d.

549. HIPPOMANE.

Linn. *Gen.* 1099. *Spec.* 1191. *Syst. No.* 1099. p. 1383. 1259
MANCANILLA. Plum. Ludw. 1055. Mill. ii. 7.

550. HIPPOPHAE.

Linn. *Gen.* 980. *Spec.* 1023. *Syst.* 980. Ludw. 921.
RHAMNOIDES. Tourn. *tab.* 481. Hall. 161. Mill. ii. 198.

551. HIPPURIS.

Linn. *Gen.* 11. *Spec.* 4. *Syst.* 11.
LIMNOPEUCE. Vaill. Hall. 197. Ludw. 937.
PINASTELLA. Dill.

552. HIRTELLA.

Linn. *Gen.* 44. *Spec.* 34. *Syst.* 44. Ludw. 512.

553. HOLCUS.

Linn. *Gen.* 1015. *Spec.* 1047. *Syst.* 1015. Ludw. 846.
SORGUM. Mich.

554. HOLOSTEUM.

Linn. *Gen.* 98. *Spec.* 88. *Syst.* 98.
NUMMULARIA. Nov. Gen. ALSINE. Hall. 385. Ludw. 569.

555. HORDEUM.

Linn. *Gen.* 93. *Spec.* 84. *Syst.* 93. Hall. 204. Ludw. 828. Mill.
i. 407. Schæff. A. 300. Tourn. *tab.* 295. Weinm. *tab.* 577.

556. HORMINUM.

Linn. *Gen.* 649. *Spec.* 596. *Syst.* 649. Ludw. 223.

557. HOTTONIA.

Linn. *Gen.* 186. *Spec.* 145. *Syst.* 186. Hall. 487. Ludw. 45.
Mill. i. 409.

STRATIOTES. Vaill. MYRIOPHYLLUM. Riv.

558. HOUSTONIA.

Linn. *Gen.* 116. *Spec.* 105. *Syst.* 116.

RUBEOLA. Ludw. 12.

559. HUGONIA.

Linn. *Gen.* 745. *Spec.* 675. *Syst.* 745. Ludw. 577.

560. HUMULUS.

Linn. *Gen.* 989. *Spec.* 1028. *Syst.* 989.LUPULUS. Tourn. *tab.* 309. Hall. 166. Ludw. 924. Mill. i. 535.
Schæff. A. 273. Weinm. *tab.* 675.

561. HURA.

Linn. *Gen.* 965. *Spec.* 1008. *Syst. No.* 965. p. 1383. Ludw. 1045.
Mill. i. 410. iii. 141.

562. HYACINTHUS.

Linn. *Gen.* 385. *Spec.* 316. *Syst.* 385.HYACINTHUS. Tourn. *tab.* 180. Blackw. *tab.* 61. Ludw. 121.
Mill. i. 411. iii. 142. Weinm. *tab.* 583, *seq.*MUSCARI. Tourn. *tab.* 180. Hall. 285. Ludw. 122. Mill. ii. 61.
iii. 198. Weinm. *tab.* 583. c, d, e, f.

563. HYDNUM.

Linn. *Gen.* 1076. *Spec.* 1178. *Syst.* 1076. Gled. *tab.* iii.
Schæff. B. § 70.

ERINACEUS. Dill. Hall. 31. Ludw. 966.

ECHIN-AGARICUS. Hall. 32.

564. HYDRANGEA.

Linn. *Gen.* 492. *Spec.* 397. *Syst.* 492. Ludw. 567.

565. HYDRASTIS.

Linn. *Syst. No.* 1153. p. 1374. 1088.

566. HYDROCHARIS.

Linn. *Gen.* 999. *Spec.* 1036. *Syst.* 999. Hall. 301. Ludw. 909.
MICROLEUCONYMPHÆA. Boerh. STRATIOTES. Dill. MORSUS
RANÆ. Tourn.

567. HYDROCOTYLE.

Linn. *Gen.* 288. *Spec.* 234. *Syst.* 288. Hall. 425. Ludw. 654.
Mill. i. 416. Tourn. *tab.* 173.

568. HYDROPHYLLUM.

Linn. *Gen.* 187. *Spec.* 146. *Syst.* 187. Ludw. 49. Mill. i. 416.
Tourn. *tab.* 16. Weinm. *tab.* 588. a.

569. HYMENÆA.

Linn. *Gen.* 1100. *Spec.* 1192. *Syst. No.* 1100. p. 1016.
COURBARIL. Plum. Ludw. 1056. Mill. i. 239.

570. HYOSCYAMUS.

Linn. *Gen.* 219. *Spec.* 179. *Syst.* 219. Blackw. *tab.* 111. Hall.
512. Ludw. 275. Mill. i. 418. Schæff. A. 89. Tourn. *tab.* 42.
Weinm. *tab.* 588.

571. HYOSERIS.

Linn. *Gen.* 821. *Spec.* 808. *Syst.* 821. Ludw. 331.

TARAXOCONASTRUM. Vaill.

LEONTODONTOIDES. Mich.

HEDYPNOIS. Tourn. *tab.* 271. Ludw. 330. Mill. i. 390. RIIA-
GADIOLOIDES. Vaill.

572. HYPECOUM.

Linn. *Gen.* 157. *Spec.* 124. *Syst.* 157. Ludw. 467.

HYPECOON. Tourn. *tab.* 115. Mill. i. 419.

573. HYPERICUM.

Linn. *Gen.* 808. *Spec.* 783. *Syst.* 808. Hall. 360.

HYPERICUM. Tourn. *tab.* 131. Blackw. *tab.* 15. Ludw. 610. Mill.
i. 419. Schæff. A. 198. Weinm. *tab.* 589, *seq.*

ASCYRUM. Tourn. *tab.* 131. Ludw. 615. Mill. i. 80. Weinm. *tab.*
589. d.

ANDROÆMUM. Tourn. *tab.* 128. Blackw. *tab.* 94. Ludw. 611.
Mill. i. 47. Weinm. *tab.* 589. e, f.

574. HYPNUM.

Linn. *Gen.* 1058. *Spec.* 1122. *Syst.* 1058. Hall. 97. Ludw. 958.

575. HYPOCHÆRIS.

Linn. *Gen.* 822. *Spec.* 810. *Syst.* 822.

HYPOCHÆRIS. Hall. 759. Ludw. 346. ACHYROPHORUS. Vaill.

576. HYPOXIS.

Linn. *Syst. No.* 1127. *p.* 1366. 986.

577. HYSSOPUS.

Linn. *Gen.* 628. *Spec.* 569. *Syst.* 628. Blackw. *tab.* 296. Ludw. 225. Mill. i. 441. Schæff. A. 70. Tourn. *tab.* 95. Weinm. *tab.* 591.

578. JAMBOLIFERA.

Linn. *Gen. No.* 430. *p.* 349. *Syst.* 430.

579. JASIONE.

Linn. *Gen.* 896. *Spec.* 928. *Syst.* 896. Hall. 496.

RAPUNCULUS. Ludw. 276.

580. JASMINUM.

Linn. *Gen.* 17. *Spec.* 7. *Syst.* 17. Blackw. *tab.* 13. Ludw. 4. Mill. i. 449. Schæff. A. 324. Tourn. *tab.* 308. Weinm. *tab.* 602.

581. JATROPHA.

Linn. *Gen.* 961. *Spec.* 1006. *Syst.* 961.

MANIHOT. Tourn. *tab.* 438. Ludw. 864. Mill. ii. 8.

JUSSIEUIA. Houst.

582. IBERIS.

Linn. *Gen.* 721. *Spec.* 648. *Syst.* 721. Hall. 542. Ludw. 421. Mill. iii. 145.

THLASPIDIUM. Riv.

583. ILEX.

Linn. *Gen.* 158. *Spec.* 125. *Syst.* 158.

AQUIFOLIUM. Tourn. *tab.* 371. Blackw. *tab.* 205. Hall. 467. Ludw. 28. Mill. i. 69. Weinm. *tab.* 158.

DODONÆA. Plum. Ludw. 796. Mill. iii. 91.

584. ILLECEBRUM.

Linn. *Gen.* 256. *Spec.* 206. *Syst.* 256.

CORRIGIOLA. Dill. Ludw. 536.

PARONYCHIA. Tourn. *tab.* 281. Mill. i. 108.

585. ILLICIAM.

Linn. *Syst. No.* 1142. *p.* 1370. 1050.

586. IMPATIENS.

Linn. *Gen.* 899. *Spec.* 937. *Syst.* 899. Hall. 505.BALSAMINA. Tourn. *tab.* 235. Ludw. 469. Mill. i. 101. iii. 35.Weinm. *tab.* 218. 222.IMPATIENS. Weinm. *tab.* 603.

587. IMPERATORIA.

Linn. *Gen.* 321. *Spec.* 259. *Syst.* 321. Blackw. *tab.* 279. Hall.445. Ludw. 660. Mill. i. 458. Schæff. A. 238. Weinm. *tab.*

604.

588. INDIGOFERA.

Linn. *Gen.* 794. *Spec.* 751. *Syst.* 794. Weinm. *tab.* 605.

589. INULA.

Linn. *Gen.* 860. *Spec.* 881. *Syst.* 860.

ENULA. Cæsalp. Schæff. A. 119. HELENIUM. Riv. Mill. i. 393.

Weinm. *tab.* 565.

590. IPOMŒA.

Linn. *Gen.* 199. *Spec.* 159. *Syst.* 199.QUAMOCLIT. Tourn. *tab.* 39. Ludw. 67. Mill. ii. 180.

VOLUBILIS. Dill.

591. IRESINE.

Linn. *Syst. No.* 1174. *p.* 1381. 1291.

592. IRIS.

Linn. *Gen.* 57. *Spec.* 38. *Syst.* 57. Hall. 279.IRIS. Tourn. *tab.* 186. 188. Blackw. *tab.* 69. 261. Ludw. 8. Mill.i. 459. Schæff. A. 6, 7. Weinm. *tab.* 607.

XIPHION. Tourn. *tab.* 189. Mill. *ii.* 527. Weinm. *tab.* 609.

SISYRINCHIUM. Tourn. Mill. *ii.* 264. Weinm. *tab.* 925.

HERMODACTYLUS. Tourn. Mill. *i.* 403.

593. ISATIS.

Linn. *Gen.* 738. *Spec.* 670. *Syst.* 738. Blackw. *tab.* 246. Hall.
535. Ludw. 401. Mill. *i.* 462. Tourn. *tab.* 100. Weinm. *tab.*
614.

594. ISCHÆMUM.

Linn. *Gen.* 1016. *Spec.* 1049. *Syst.* 1016.

SCHÆNANTHUS. Scheuchz. Ludw. 1060.

595. ISNARDIA.

Linn. *Gen.* 145. *Spec.* 120. *Syst.* 145. Ludw. 766.

DANTIA. Petit.

596. ISCETES.

Linn. *Gen.* 1048. *Spec.* 1100. *Syst.* 1048.

597. ISOPYRUM.

Linn. *Gen.* 621. *Spec.* 557. *Syst.* 621. Ludw. 607.

598. ITEA.

Linn. *Gen.* 243. *Spec.* 199. *Syst.* 243. Ludw. 55.

DICONANGIA. Mich.

599. IVA.

Linn. *Gen.* 940. *Spec.* 988. *Syst.* 940.

TARCHONANTHUS. Vaill.

600. JUGLANS.

Linn. *Gen.* 950. *Spec.* 997. *Syst.* 950. Blackw. *tab.* 247. Ludw.
892. Mill. *iii.* 147. Schæff. A. 294.

NUX JUGLANS. Tourn. *tab.* 346. Mill. *ii.* 79.

601. JUNCUS.

Linn. *Gen.* 396. *Spec.* 325. *Syst.* 396. Hall. 252. Ludw. 787.
Mill. *i.* 464. Tourn. *tab.* 127. Weinm. *tab.* 614.

602. JUNGERMANNIA.

Linn. *Gen.* 1059. *Spec.* 1131. *Syst.* 1059. Ludw. 962.

JUNGERMANNIA. Mich. Hall. 120.

MUSCOIDES. Mich. Hall. 123.

LICHENASTRUM. Dill.

603. JUNIPERUS.

Linn. *Gen.* 1005. *Spec.* 1038. *Syst.* 1005. Ludw. 919.

JUNIPERUS. Tourn. *tab.* 361. Blackw. *tab.* 187. Hall. 146. Mill.
i. 464. iii. 147. Schæff. A. 265. Weinm. *tab.* 615.

SABINA. Boerh. Blackw. *tab.* 214. Mill. ii. 220. Schæff. A. 266.
Weinm. *tab.* 876.

CEDRUS. Tourn. *tab.* 361. Mill. i. 186, 187. Weinm. *tab.* 346.

604. JUSSIEA.

Linn. *Gen.* 478. *Spec.* 388. *Syst.* 478. Ludw. 553.

605. JUSTICIA.

Linn. *Gen.* 26. *Spec.* 15. *Syst.* 26. Mill. i. 468.

ADHATODA. Tourn. *tab.* 79. Ludw. 184. Mill. i. 19. ECBULIUM.
Weinm. *tab.* 473.

606. IXIA.

Linn. *Gen.* 54. *Spec.* 36. *Syst.* 54. Ludw. 843.

607. IXORA.

Linn. *Gen.* 122. *Spec.* 110. *Syst.* 122. Ludw. 24.

608. KÆMPFERIA.

Linn. *Gen.* 7. *Spec.* 2. *Syst.* 7. Ludw. 372.

KÆMPFERA. Mill. i. 472. iii. 149.

609. KALMIA.

Linn. *Gen.* 482. *Spec.* 391. *Syst.* 482.

610. KIGGELARIA.

Linn. *Gen.* 1001. *Spec.* 1037. *Syst.* 1001. Ludw. 915. Mill. iii.
151.

TABLE II.

611. KNAUTIA.

Linn. *Gen.* 109. *Spec.* 101. *Syst.* 109. Ludw. 290. Mill. iii. 153.
 LYCHNISCABIOSA. Boerh. LIMNESIUM. Sigb.

612. KNOXIA.

Linn. *Gen.* 115. *Spec.* 104. *Syst.* 115.

613. KRAMERIA.

Linn. *Syst. No.* 1116. p. 1362. 899.

614. LACHNÆA.

Linn. *Gen.* 441. *Spec.* 500. *Syst.* 441. Ludw. 789.

615. LACTUCA.

Linn. *Gen.* 814. *Spec.* 795. *Syst.* 814. Blackw. *tab.* 88. Hall. 755.
 Ludw. 340. Mill. i. 479. Schæff. A. 113. Tourn. *tab.* 267.
 Weinm. *tab.* 619, *seq.*

616. LÆTIA.

Linn. *Syst. No.* 1149. p. 1373. 1074.

617. LAGERSTRÆMIA.

Linn. *Syst. No.* 1146. p. 1372. 1076.

618. LAGÆCIA.

Linn. *Gen.* 251. *Spec.* 203. *Syst.* 251. Ludw. 513. Mill. iii. 154.
 CUMINOIDES. Tourn. *tab.* 155. Mill. i. 254.

619. LAGURUS.

Linn. *Gen.* 86. *Spec.* 81. *Syst.* 86. Ludw. 822.

620. LAMIUM.

Linn. *Gen.* 636. *Spec.* 579. *Syst.* 636. Blackw. *tab.* 182. Hall.
 640. Ludw. 192. Mill. i. 481. Tourn. *tab.* 85.
 GALEOPSIS. Riv.

621. LANTANA.

Linn. *Gen.* 683. *Spec.* 626. *Syst.* 683. Ludw. 265. Mill. iii. 156.
 CAMARA. Plum. Mill. i. 161.

MYROBATINDUM. Vaill.

VIBURNUM. Weinm. *tab.* 1007.

622. LAPSANA.

Linn. *Gen.* 823. *Spec.* 811. *Syst.* 823.

LAPSANA. Tourn. *tab.* 272. Hall. 759. Ludw. 328. Mill. i. 482.
Weinm. *tab.* 673.

RHAGADIOLUS. Tourn. *tab.* 272. Ludw. 329.

ZACINTHA. Tourn. *tab.* 269. Ludw. 332. Mill. ii. 531.

623. LASERPITIUM.

Linn. *Gen.* 306. *Spec.* 248. *Syst.* 306. Hall. 440. Ludw. 684.
Mill. ii. 260. iii. 158. Tourn. *tab.* 172. Weinm. *tab.* 647. a.

624. LATHRÆA.

Linn. *Gen.* 661. *Spec.* 605. *Syst.* 661.

CLANDESTINA. Tourn. *tab.* 424. Ludw. 229.

ANBLATUM. Tourn. *tab.* 481. SQUAMARIA. Riv. Hall. 611.
Ludw. 227.

PHELYPÆA. Linn. *edit. prior.*

625. LATHYRUS.

Linn. *Gen.* 781. *Spec.* 729. *Syst.* 781. Hall. 594. Ludw. 488.

LATHYRUS. Tourn. *tab.* 216, 217. Mill. i. 490. Weinm. *tab.* 628,
seq.

CLYMENUM. Tourn. *tab.* 218. Mill. i. 215. Weinm. *tab.* 295.

APHACA. Tourn. *tab.* 223. Mill. i. 65.

626. LAVANDULA.

Linn. *Gen.* 630. *Spec.* 572. *Syst.* 630. Ludw. 220.

LAVENDULA. Tourn. *tab.* 93. Blackw. *tab.* 294, 295. Mill. i. 492.
Schæff. A. 80.

STÆCHAS. Tourn. *tab.* 95. Blackw. *tab.* 241. Mill. ii. 303.

627. LAVATERA.

Linn. *Gen.* 752. *Spec.* 690. *Syst.* 752. Ludw. 147. Mill. i. 492.
Tourn. *A. G.*

628. LAURUS.

Linn. *Gen.* 452. *Spec.* 369. *Syst.* 452. Ludw. 133.

LAURUS. Tourn. *tab.* 367. Black. *tab.* 175. Mill. i. 495. iii. 160.

Schæff. A. 3. Weinm. *tab.* 634.

CINNAMOMUM. Burm. Blackw. *tab.* 354.

CAMPHORA. Boerh. Blackw. *tab.* 347. Mill. i. 165.

PERSEA. Plum. Mill. ii. 114.

BORBONIA. Plum. Mill. iii. 41.

BENZOE. Boerh. Mill. i. 119.

SASSAFRAS. Off. Black. *tab.* 267.

629. LAWSONIA.

Linn. *Gen.* 433. *Spec.* 349. *Syst.* 433.

HENNA. Ludw. 432.

630. LECHEA.

Linn. *Gen.* 102. *Spec.* 90. *Syst.* 102.

631. LECYTHIS.

Linn. *Syst. No.* 1147. p. 1372. 1071.

632. LEDUM.

Linn. *Gen.* 483. *Spec.* 391. *Syst.* 483. Ludw. 560.

633. LEMNA.

Linn. *Gen.* 923. *Spec.* 970. *Syst.* 923.

LENTICULA. Mich. Blackw. *tab.* 380. Hall. 128. Ludw. 977.

Weinm. *tab.* 637.

HYDROPHACE. Buxb.

634. LEONTICE.

Linn. *Gen.* 381. *Spec.* 312. *Syst.* 381.

LEONTOPETALON. Tourn. *tab.* 484. Ludw. 709. Mill. i. 499.

Weinm. *tab.* 630.

635. LEONTODON.

Linn. *Gen.* 817. *Spec.* 798. *Syst.* 817.

DENS LEONIS. Tourn. *tab.* 266. Mill. i. 268. TARAXACUM. Off.
Hall. 739. Blackw. *tab.* 1. Ludw. 339. Schæff. A. 115. Weinm.
tab. 966.

TARAXACONOIDES. Vaill.

636. LEONURUS.

Linn. *Gen.* 641. *Spec.* 584. *Syst.* 641.

LEONURUS. Tourn. *tab.* 87. Ludw. 198. Mill. i. 499. Weinm.
tab. 640.

CARDIACA. Tourn. *tab.* 87. Blackw. *tab.* 171. Hall. 639. Ludw.
199. Mill. i. 170. Schæff. A. 76. Weinm. *tab.* 307.

637. LEPIDIUM.

Linn. *Gen.* 718. *Spec.* 643. *Syst.* 718.

LEPIDIUM. Tourn. *tab.* 103. Ludw. 423. Mill. i. 500. Weinm.
tab. 641. a, b.

NASTURTIUM. Tourn. *tab.* 102. Blackw. *tab.* 23. Hall. 543. Ludw.
422. Mill. ii. 71. Schæff. A. 144. Weinm. *tab.* 751, 752.

638. LEUCADENDRON.

Linn. *Gen.* 102. *Spec.* 91. *Syst.* 102. Ludw. 291.

LEPIDOCARPODENDRON. Boerh. Mill. i. 500.

HYPOPHYLLOCARPODENDRON. Boerh.

CONOCARPODENDRON. Boerh. Mill. i. 221.

639. LEUCOIUM.

Linn. *Gen.* 363. *Spec.* 289. *Syst.* 363. Hall. 284. Ludw. 721.

NARCISSO-LEUCOIUM. Tourn. *tab.* 208. A. F. Mill. ii. 66. Weinm.
tab. 642. a.

640. LICHEN.

Linn. *Gen.* 1065. *Spec.* 1140. *Syst.* 1065. Blackw. *tab.* 335, 336.

Hall. 63. Mill. i. 504. Tourn. *tab.* 325. Weinm. *tab.* 648.

LICHEN. Mich. Ludw. 985.

LICHENOIDES. Dill. Ludw. 986.

CORALLOIDES. Dill.

ULNEA. Dill.

641. LIGUSTICUM.

Linn. *Gen.* 308. *Spec.* 250. *Syst.* 308. Hall. 434. Ludw. 688.
 LIGUSTICUM. Tourn. *tab.* 171. Blackw. *tab.* 275. Mill. i. 506. iii.
 162. Schæff. A. 240. Weinm. *tab.* 647. a.
 CICUTARIA. Tourn. *tab.* 171. Mill. i. 209.

642. LIGUSTRUM.

Linn. *Gen.* 18. *Spec.* 7. *Syst.* 18. Blackw. *tab.* 140. Hall. 527.
 Ludw. 5. Mill. i. 506. iii. 162. Schæff. A. 320. Tourn. *tab.* 367.
 Weinm. *tab.* 650. b.

643. LILIUM.

Linn. *Gen.* 371. *Spec.* 302. *Syst.* 371. Blackw. *tab.* 11. Hall.
 289. Ludw. 715. Mill. i. 513. Schæff. A. 247. Tourn. *tab.*
 195. Weinm. *tab.* 651.

644. LINEUM.

Linn. *Syst. No.* 1128. *p.* 1366. 995.

645. LIMODORUM.

Linn. *Gen.* 904. *Spec.* 950. *Syst.* 904. Hall. 278. Ludw. 706.
 Tourn. *tab.* 250?

646. LIMOSELLA.

Linn. *Gen.* 694. *Spec.* 631. *Syst.* 694. Hall. 609. Ludw. 238.
 PLANTAGINELLA. Dill.

647. LINNÆA.

Linn. *Gen.* 692. *Spec.* 631. *Syst.* 692. Ludw. 23.
 SERPILLIFOLIA. Buxb.

648. LINUM.

Linn. *Gen.* 349. *Spec.* 277. *Syst.* 349. Blackw. *tab.* 160. 368.
 Hall. 370. Ludw. 547. Mill. i. 520. Schæff. A. 210, 211.
 Tourn. *tab.* 176. Weinm. *tab.* 669.
 RADIOLA. Dill.
 CHAMÆLINUM. Mich. LINOCARPON. Mich.

649. LIPPIA.

Linn. *Gen.* 699. *Spec.* 633. *Syst.* 699. Ludw. 228. Mill. i. 521.

650. LIQUIDAMBAR.

Linn. *Gen.* 955. *Spec.* 999. *Syst.* 955. Ludw. 893. Mill. iii. 164.
Weinm. *tab.* 953.

651. LIRIODENDRON.

Linn. *Gen.* 609. *Spec.* 535. *Syst.* 609. Ludw. 759.

TULIPIFERA. Catesb. Mill. ii. 360. Weinm. *tab.* 997, 998.

652. LITHOSPERMUM.

Linn. *Gen.* 166. *Spec.* 132. *Syst.* 166. Hall. 517. Ludw. 36.
Mill. i. 521. Schæff. A. 41. Tourn. *tab.* 55. Weinm. *tab.* 671.

653. LITHOXYLON.

Linn. *Gen.* 1073. *Spec.* ... *Syst.* ...

LITHOPHYTON. Tourn. *tab.* 341.

KERATOPHYTON. Boerh. Ludw. 1003.

TITANOKERATOPHYTON. Boerh.

654. LOBELIA.

Linn. *Gen.* 897. *Spec.* 929. *Syst.* 897. Ludw. 272.

LOBELIA. Mill. i. 522.

RAPUNTIUM. Tourn. *tab.* 51. Mill. ii. 192.

DORTMANNA. Rudb.

LAURENTIA. Mich.

655. LÆFLINGIA.

Linn. *Gen.* 50. *Spec.* 35. *Syst.* 50.

656. LÆSELIA.

Linn. *Gen.* 685. *Spec.* 628. *Syst.* 685. Ludw. 259.

ROYENI. Houst.

657. LOLIUM.

Linn. *Gen.* 90. *Spec.* 83. *Syst.* 90. Hall. 206. Ludw. 836.

658. LONCHITIS.

Linn. *Gen.* 1041. *Spec.* 1078. *Syst.* 1041. Ludw. 947. Mill. i. 525.

659. LONICERA.

Linn. *Gen.* 210. *Spec.* 173. *Syst.* 210. Mill. iii. 166.

CAPRIFOLIUM. Tourn. *tab.* 378. Blackw. *tab.* 25. Hall. 464. Ludw. 278. Mill. i. 108. Weinm. *tab.* 802.

PERICLYMENUM. Tourn. *tab.* 378. Ludw. 280. Mill. ii. 113. Weinm. *tab.* 801.

CHAMÆCERASUS. Tourn. *tab.* 379. Hall. 464. Ludw. 279. Mill. i. 109.

XYLOSTEUM. Tourn. *tab.* 379. Ludw. 281. Mill. ii. 530. Weinm. *tab.* 1022.

SYMPHORICARPOS. Dill. Ludw. 70.

660. LORANTHUS.

Linn. *Gen.* 400. *Spec.* 331. *Syst.* 400.

661. LOTUS.

Linn. *Gen.* 803. *Spec.* 773. *Syst.* 803. Hall. 571. Ludw. 490. Mill. i. 525. iii. 186. Tourn. *tab.* 227. Weinm. *tab.* 672.

662. LUDWIGIA.

Linn. *Gen.* 142. *Spec.* 118. *Syst.* 142. Ludw. 392. Mill. iii. 168.

663. LUNARIA.

Linn. *Gen.* 725. *Spec.* 653. *Syst.* 725. Hall. 540. Ludw. 417. Mill. i. 533. Tourn. *tab.* 105.

664. LUPINUS.

Linn. *Gen.* 774. *Spec.* 721. *Syst.* 774. Blackw. *tab.* 282. Ludw. 494. Mill. i. 534. Schæff. A. 158. Tourn. *tab.* 213. Weinm. *tab.* 674.

665. LYCHNIS.

Linn. *Gen.* 517. *Spec.* 436. *Syst.* 517. Ludw. 573. Mill. i. 543. Tourn. *tab.* 175. A. B. Weinm. *tab.* 678, *seq.*

666. LYCIUM.

Linn. *Gen.* 232. *Spec.* 191. *Syst.* 232.

JASMINOIDES. Dill. Mill. i. 447.

667. LYCOPERDON.

Linn. *Gen.* 1082. *Spec.* 1183. *Syst.* 1082. Battarr. *tab.* 39. Gled.
tab. 5. Ludw. 973. Schæff. A. 318. B. §. 70.

LYCOPERDON. Mich. Tourn. *tab.* 331. Hall. 11. BOVISTA. Dill.

LYCOPERDOIDES. Mich. Hall. 13.

LYCOPERDASTRUM. Mich. Hall. 13.

GEASTER. Mich. Hall. 13.

CARPOBOLUS. Mich. Hall. 13.

TUBER. Mich. Hall. 14. TARTUFI. Imp.

668. LYCOPODIUM.

Linn. *Gen.* 1049. *Spec.* 1100. *Syst.* 1049. Ludw. 955.

LYCOPODIUM. Dill. Hall. 92. Weinm. *tab.* 737. c.

LYCOPODIOIDES. Dill. Hall. 93.

SELAGO. Dill. Hall. 94.

SELAGINOIDES. Dill. Hall. 94.

669. LYCOPSIS.

Linn. *Gen.* 174. *Spec.* 138. *Syst.* 174. Hall. 523. Ludw. 32.

ECHIOIDES. Dill.

670. LYCOPUS.

Linn. *Gen.* 31. *Spec.* 21. *Syst.* 31. Hall. 660. Ludw. 178. Tourn.
tab. 89.

671. LYGEUM.

Linn. *Gen.* 64. *Spec.* ... *Syst.* 64.

672. LYSIMACHIA.

Linn. *Gen.* 188. *Spec.* 146. *Syst.* 188. Hall. 480. Ludw. 42.

LYSIMACHIA. Tourn. *tab.* 59. Blackw. *tab.* 278. Mill. i. 547.

Weinm. *tab.* 688, *seq.*

NUMMULARIA. Bauh. Mill. ii. 79. Schæff. A. 26. Weinm. *tab.*
758. e, f.

673. LYTHRUM.

Linn. *Gen.* 532. *Spec.* 446. *Syst.* 532.

SALICARIA. Tourn. *tab.* 129. Hall. 405. Ludw. 734. Mill. ii. 226. Weinm. *tab.* 688. i.

674. MAGNOLIA.

Linn. *Gen.* 610. *Spec.* 535. *Syst.* 610. Ludw. 760. Mill. ii. 1. iii. 170.

675. MALOPE.

Linn. *Gen.* 753. *Spec.* 692. *Syst.* 753. Mill. iii. 171.

MALACOIDES. Tourn. *tab.* 25. Ludw. 149. Mill. ii. 3.

676. MALPIGHIA.

Linn. *Gen.* 508. *Spec.* 425. *Syst.* 508. Ludw. 575. Mill. ii. 13. iii. 172.

677. MALVA.

Linn. *Gen.* 751. *Spec.* 687. *Syst.* 751. Hall. 362.

MALVA. Tourn. *tab.* 23, 24. Blackw. *tab.* 22. Mill. i. 4. iii. 173.

Schæff. A. 50. Weinm. *tab.* 693, *seq.*

ALCEA. Tourn. *tab.* 25. Blackw. *tab.* 309. Mill. i. 24. iii. 11.

Schæff. A. 330. Weinm. *tab.* 33.

ABUTILON. Dill. Mill. i. 9.

678. MAMMEA.

Linn. *Gen.* 583. *Spec.* 512. *Syst.* 583. Ludw. 456.

MAMEI. Plum. Mill. ii. 6.

679. MANDRAGORA.

Linn. *Gen.* 221. *Spec.* 181. *Syst.* 221. Blackw. *tab.* 364. Ludw. 89. Mill. ii. 7. Schæff. A. 27. Tourn. *tab.* 12. Weinm. *tab.* 709.

680. MANGIFERA.

Linn. *Gen.* 245. *Spec.* 200. *Syst.* 245.

681. MARANTA.

Linn. *Gen.* 5. *Spec.* 2. *Syst.* 5. Ludw. 172. Mill. ii. 9.

682. MARCGRAVIA.

Linn. *Gen.* 564. *Spec.* 503. *Syst.* 564. Ludw. 165.

683. MARCHANTIA.

Linn. *Gen.* 1061. *Spec.* 1137. *Syst.* 1061.

MARCHANTIA. Mich. Hall. 126.

HEPATICA. Mich. Hall. 126.

MARSILEA. Mich. Hall. 124.

LUNULARIA. Mich. Hall. 125.

LICHEN. Dill. Hall. 126.

684. MARRUBIUM.

Linn. *Gen.* 640. *Spec.* 582. *Syst.* 640.

MARRUBIUM. Tourn. *tab.* 91. Hall 648. Ludw. 212. Mill. ii. 10.

Schæff. A. 92. Weinm. *tab.* 709.

PSEUDODICTAMNUS. Tourn. *tab.* 89. Ludw. 213. Mill. ii. 165.

685. MARSILEA.

Linn. *Gen.* 1046. *Spec.* 1099. *Syst.* 1046.

SALVINIA. Mich. Ludw. 979.

686. MARTYNIA.

Linn. *Gen.* 671. *Spec.* 618. *Syst.* 671. Ludw. 260. Mill. ii. 11.

687. MATRICARIA.

Linn. *Gen.* 687. *Spec.* 890. *Syst.* 867. Blackw. *tab.* 192. Mill. ii.

12. Schæff. A. 128. Tourn. *tab.* 281. Weinm. *tab.* 713.

688. MATTHIOLA.

Linn. *Gen.* 1101. *Spec.* 1192. *Syst.* 1101. Ludw. 1016.

689. MEDEOLA.

Linn. *Gen.* 411. *Spec.* 339. *Syst.* 411. Ludw. 732.

690. MEDICAGO.

Linn. *Gen.* 805. *Spec.* 778. *Syst.* 805. Ludw. 505.

MEDICAGO. Tourn. *tab.* 231. Mill. ii. 27.

MEDICA. Tourn. *tab.* 231. Hall. 578. Mill. ii. 23. MEDICA
COCHLEATA. Mill. ii. 25. Weinm. *tab.* 401.
FALCATA. Riv. Weinm. *tab.* 501.

691. MELAMPODIUM.

Linn. *Gen.* 884. *Spec.* 921. *Syst.* 884. Ludw. 315.

692. MELAMPYRUM.

Linn. *Gen.* 660. *Spec.* 605. *Syst.* 660. Hall. 625. Ludw. 244.
Mill. ii. 29. Tourn. *tab.* 78. Weinm. *tab.* 716, 717.

693. MELANTHIUM.

Linn. *Gen.* 410. *Spec.* 339. *Syst.* 410. Ludw. 1061.

694. MELASTOMA.

Linn. *Gen.* 481. *Spec.* 389. *Syst.* 481. Ludw. 562. Mill. iii. 183.
ACINODENDRON. Linn. *edit. prior.*

695. MELIA.

Linn. *Gen.* 473. *Spec.* 384. *Syst.* 473. Mill. iii. 34.
AZEDARACH. Tourn. *tab.* 387. Ludw. 561. Mill. i. 99. Weinm.
tab. 217.

696. MELIANTHUS.

Linn. *Gen.* 712. *Spec.* 639. *Syst.* 712. Ludw. 468. Mill. ii. 30.
Tourn. *tab.* 245. Weinm. *tab.* 717. c.

697. MELICA.

Linn. *Gen.* 76. *Spec.* 66. *Syst.* 76. Hall. 215. Ludw. 823. Mill.
i. 823.

698. MELISSA.

Linn. *Gen.* 647. *Spec.* 592. *Syst.* 647.
MELISSA. Tourn. *tab.* 91. Blackw. *tab.* 27. Hall. 651. Ludw.
214. Mill. ii. 32. iii. 47. Schæff. A. 69. Weinm. *tab.* 719.
CALAMINTHA. Tourn. *tab.* 92. Blackw. *tab.* 166. Hall. 650. Ludw.
215. Mill. i. 160. Schæff. A. 68. Weinm. *tab.* 282, 283.

699. MELITTIS.

Linn. *Gen.* 650. *Spec.* 597. *Syst.* 650. Ludw. 216.

MELISSA. Hall. 650. MELISSOPHYLLON. Riv.

700. MELOCHIA.

Linn. *Gen.* 743. *Spec.* 674. *Syst.* 743. Ludw. 545.

701. MELOTHRIA.

Linn. *Gen.* 48. *Spec.* 35. *Syst.* 48. Ludw. 9. Mill. iii. 192.

702. MEMECYLON.

Linn. *Gen.* 432. *Spec.* 349. *Syst.* 432.

703. MENISPERMUM.

Linn. *Gen.* 413. *Spec.* 340. *Syst.* 413. Ludw. 733. Mill. ii. 41.

704. MENTHA.

Linn. *Gen.* 633. *Spec.* 576. *Syst.* 633. Blackw. *tab.* 22. 290. 292.

Hall. 657. Ludw. 217. Mill. ii. 42. Schæff. A. 65. Tourn. *tab.* 89. Weinm. *tab.* 724.

705. MENTZELIA.

Linn. *Gen.* 595. *Spec.* 516. *Syst.* 595. Ludw. 585. Mill. ii. 42.

706. MENYANTHES.

Linn. *Gen.* 185. *Spec.* 145. *Syst.* 185. Ludw. 51.

MENOANTHES. Hall. 487. MENYANTHES. Tourn. *tab.* 15. Mill.

ii. 43. TRIFOLIUM FIBRINUM. Off. Schæff. A. 315. Weinm. *tab.* 979. d.

NYMPHOIDES. Tourn. *tab.* 67. Weinm. *tab.* 761. f.

707. MERCURIALIS.

Linn. *Gen.* 998. *Spec.* 1035. *Syst.* 998. Blackw. *tab.* 162. Hall.

194. Ludw. 933. Mill. ii. 43. Schæff. A. 267. Tourn. *tab.* 308. Weinm. *tab.* 726.

708. MESEMBRYANTHEMUM.

Linn. *Gen. No.* 552. *p.* 480. *Syst.* 552.

FICOIDES. Tourn. Ludw. 167. Mill. i. 316. iii. 105. Weinm. *tab.*

709. MESPILUS.

Linn. *Gen.* 549. *Spec.* 478. *Syst.* 549. Blackw. *tab.* 154. Hall.
352. Ludw. 617. Mill. ii. 44. iii. 192. Schæff. A. 199. Tourn.
tab. 410. Weinm. *tab.* 728.

710. MESUA.

Linn. *Gen.* 591. *Spec.* 515. *Syst.* 591. Ludw. 450.

711. MICHELIA.

Linn. *Gen.* 611. *Spec.* 536. *Syst.* 611. Ludw. 745.

712. MICROCOS.

Linn. *Gen.* No. 588. *p.* 514. *Syst.* 588.

713. MICROPUS.

Linn. *Gen.* No. 892. *p.* 927. *Syst.* 892. Ludw. 316. Mill. i. 373.
GNAPHALODES. Tourn. *tab.* 439.

714. MILIUM.

Linn. *Gen.* 73. *Spec.* 61. *Syst.* 73. Hall. 219. Ludw. 820. Mill.
ii. 47. Tourn. *tab.* 298. Weinm. *tab.* 729.

715. MILLERIA.

Linn. *Gen.* 881. *Spec.* 919. *Syst.* 881. Ludw. 1027. Mill. ii. 48.

716. MIMOSA.

Linn. *Gen.* 597. *Spec.* 516. *Syst.* No. 597. *p.* 1310.
MIMOSA. Tourn. *tab.* 375. Ludw. 157. Mill. ii. 48. Weinm.
tab. 731.
ACACIA. Tourn. *tab.* 375. Blackw. *tab.* 345. Ludw. 156. Mill. i.
10. Weinm. *tab.* 10.
INGA. Plum. Mill. i. 458.

717. MIMULUS.

Linn. *Gen.* 701. *Spec.* 634. *Syst.* 701.

CYNORRHYNCHIUM. Mitch.

718. MIMUSOPS.

Linn. *Gen.* 429. *Spec.* 349. *Syst.* 429.

719. MINUARTIA.

Linn. *Gen.* 100. *Spec.* 89. *Syst.* 100.

720. MIRABILIS.

Linn. *Gen.* 215. *Spec.* 177. *Syst.* 215. Weinm. *tab.* 732.

JALAPA. Tourn. *tab.* 50. Ludw. 29. Mill. i. 446.

721. MITCHELLA.

Linn. *Gen.* 126. *Spec.* 111. *Syst.* 126.

CHAMÆDAPHNE. Mitch.

722. MITELLA.

Linn. *Gen.* 496. *Spec.* 406. *Syst.* 496. Ludw. 565. Mill. ii. 53.

Tourn. *tab.* 126.

723. MNIMUM.

Linn. *Gen.* 1056. *Spec.* 1109. *Syst.* 1056. Hall. 119. Ludw. 959.

724. MÖHRINGIA.

Linn. *Gen.* 444. *Spec.* 359. *Syst.* 444. Ludw. 440.

725. MOLLUGO.

Linn. *Gen.* 99. *Spec.* 89. *Syst.* 99. Ludw. 762. Mill. iii. 195.

726. MOLUCCELLA.

Linn. *Gen.* 643. *Spec.* 587. *Syst.* 643.

MOLUCCA. Tourn. *tab.* 88. Ludw. 211. Mill. ii. 55. Weinm. *tab.*

728. c.

727. MOMORDICA.

Linn. *Gen.* 967. *Spec.* 1009. *Syst.* 96

MOMORDICA. Tourn. *tab.* 29, 30. Ludw. 852. Mill. ii. 56.

Schæff. A. 325. Weinm. *tab.* 735.

CUCUMIS ASININUS. Rai. Blackw. *tab.* 108. ELATERIUM. Boerh.
Ludw. 853. Mill. i. 527. Schæff. A. 20.
LUFFA. Tourn. Dill. Mill. i. 527.

728. MONARDA.

Linn. *Gen.* 34. *Spec.* 22. *Syst.* 34. Ludw. 179. Mill. iii. 195.

729. MONNIERIA.

Linn. *Syst.* No. 1157. p. 1375. 1153.

730. MONOTROPA.

Linn. *Gen.* 477. *Spec.* 387. *Syst.* 477.

HYPOPHYTES. Dill. Hall. 411. Ludw. 437. Weinm. *tab.* 774. a.
OROBANCHOIDES. Tourn.

731. MONTIA.

Linn. *Gen.* 96. *Spec.* 87. *Syst.* 96. Hall. 608. Mill. ii. 57.

CAMERARIA. Dill.

ALSINOIDES. Vaill.

732. MORINA.

Linn. *Gen.* 39. *Spec.* 28. *Syst.* 39. Ludw. 174. Mill. ii. 58.
Tourn. *tab.* 480.

DIOTOTHECA. Vaill.

733. MORINDA.

Linn. *Gen.* 212. *Spec.* 176. *Syst.* 212. Ludw. 1017.

ROIOC. Plum.

PHILLYREASTRUM. Vaill.

734. MORISONIA.

Linn. *Gen.* 565. *Spec.* 503. *Syst.* 565. Ludw. 461.

735. MORUS.

Linn. *Gen.* 936. *Spec.* 986. *Syst.* 936. Blackw. *tab.* 126. Ludw.
880. Mill. ii. 58. iii. 196. Schæff. A. 271. Tourn. *tab.* 362.
Weinm. *tab.* 736.

736. MUCOR.

Linn. *Gen.* 1083. *Spec.* 1185. *Syst.* 1083. Battarr. *tab.* 10. Gled.
tab. 6. Ludw. 991. Schæff. B. § 70.

MUCOR. Mich. Hall. 7.

MUCILAGO. Mich. Hall. 5.

LYCOGALA. Mich. Hall. 7.

737. MUNTINGIA.

Linn. *Gen.* 575. *Spec.* 509. *Syst.* 575. Ludw. 600. Mill. ii. 60.
iii. 197.

738. MUSA.

Linn. *Gen.* 1010. *Spec.* 1013. *Syst.* 1010. Ludw. 370. Mill. ii. 60.
BIBAL. Plum. Mill. i. 126.

739. MUSSÆNDA.

Linn. *Gen.* 214. *Spec.* 177. *Syst.* 214.

740. MYAGRUM.

Linn. *Gen.* 713. *Spec.* 640. *Syst.* 713. Ludw. 399. Mill. ii. 62.

741. MYOSOTIS.

Linn. *Gen.* 165. *Spec.* 131. *Syst.* 165.

SCORPIURUS. Knaut. Hall. 519.

LITHOSPERMI SPECIES. Tourn.

742. MYOSURUS.

Linn. *Gen.* 355. *Spec.* 284. *Syst.* 355. Ludw. 619. Mill. iii. 198.
MYOSUROS. Dill. CAUDA MURIS. Weinm. *tab.* 346. a.

743. MYRICA.

Linn. *Gen.* 981. *Spec.* 1024. *Syst.* 981. Ludw. 922. Mill. i. 340.
iii. 198.

GALE. Tourn. Mill. i. 340.

744. MYRIOPHYLLUM.

Linn. *Gen.* 945. *Spec.* 992. *Syst.* 945.

PENTAPTEROPHYLLUM. Dill. Ludw. 797.

PENTAPTERIS. Hall. 201.

745. MYRSINE.

Linn. *Gen.* 238. *Spec.* 196. *Syst.* 238. Ludw. 95.

746. MYRTUS.

Linn. *Gen.* 543. *Spec.* 471. *Syst.* 543. Blackw. *tab.* 114. Ludw. 603. Mill. ii. 63. Schæff. A. 190. Weinm. *tab.* 745.

747. NAIAS.

Linn. *Gen.* 974. *Spec.* 1015. *Syst.* 974. Ludw. 848.
FLUVIALIS. Vaill.

748. NAMA.

Linn. *Gen.* 282. *Spec.* 226. *Syst.* 282.

749. NAPÆA.

Linn. *Gen.* 748. *Spec.* 686. *Syst.* 748.

750. NARCISSUS.

Lin. *Gen.* 364. *Spec.* 289. *Syst.* 364. Hall. 284. Ludw. 118.
Mill. ii. 67. iii. 199. Tourn. *tab.* 185. Weinm. *tab.* 747, *seq.*

751. NARDUS.

Linn. *Gen.* 65. *Spec.* 53. *Syst.* 65. Hall. 203. Ludw. 814.

752. NEPENTHES.

Linn. *Gen.* 909. *Spec.* 955. *Syst.* 909. Ludw. 767.

753. NEPETA.

Linn. *Gen.* 629. *Spec.* 570. *Syst.* 629. Schæff. A. 72. Weinm.
tab. 725. c.

CATARIA. Hall. 649. Ludw. 205. Mill. i. 185. iii. 53.

754. NERIUM.

Linn. *Gen.* 262. *Spec.* 209. *Syst.* 262. Ludw. 103. Mill. ii. 75.
Tourn. *tab.* 374. Weinm. *tab.* 754, *seq.*

755. NEURADA.

Linn. *Gen.* 520. *Spec.* 441. *Syst.* 520. Ludw. 579.

756. NICOTIANA.

Linn. *Gen.* 220. *Spec.* 180. *Syst.* 220. Blackw. *tab.* 146. Ludw.
58. Mill. ii. 277. Tourn. *tab.* 41. Weinm. *tab.* 757. a.

TABACUM. Rai. Schæff. A. 31.

757. NIGELLA.

Linn. *Gen.* 606. *Spec.* 534. *Syst.* 606. Hall. 316. Ludw. 614.
 Mill. ii. 78. Schæff. A. 171. Tourn. *tab.* 134. Weinm. *tab.*
 757. b, c.

758. NITRARIA.

Linn. *Syst. No.* 1139. *p.* 1369. 1044. ♀

759. NYCTANTHES.

Linn. *Gen.* 16. *Spec.* 6. *Syst.* 16. Tourn. *tab.* 368.

760. NYMPHÆA.

Linn. *Gen.* 579. *Spec.* 510. *Syst.* 579. Ludw. 750.
 NYMPHÆA. Tourn. *tab.* 137, 138. Hall. 302. Mill. ii. 81. Schæff.
 A. 262. Weinm. *tab.* 761.
 NELUMBO. Tourn.
 LEUCONYMPHÆA. Boerh.

761. NYSSA.

Linn. *Gen.* 1028. *Spec.* 1058. *Syst.* 1028. Ludw. 913.

762. OBOLARIA.

Linn. *Gen.* 696. *Spec.* 632. *Syst.* 696. Ludw. 256.

763. OCHNA.

Linn. *Gen.* 584. *Spec.* 513. *Syst.* 584.
 JABOTAFITA. Plum. Ludw. 1063.

764. OCYMUM.

Linn. *Gen.* 651. *Spec.* 597. *Syst.* 651. Ludw. 226. Mill. ii. 83.
 iii. 200. Tourn. *tab.* 96. Weinm. *tab.* 762.
 BASILICUM. Rai. Blackw. *tab.* 104. Schæff. A. 91.

765. CENANTHE.

Linn. *Gen.* 314. *Spec.* 254. *Syst.* 314. Hall. 431. Ludw. 678.
 Mill. ii. 84. Tourn. *tab.* 166.

766. CENOTHERA.

Linn. *Gen.* 424. *Spec.* 316. *Syst.* 424.

ONAGRA. Tourn. *tab.* 156. Hall. 410. Ludw. 434. Mill. ii. 86.
Weinm. *tab.* 688. b.

767. OLAX.

Linn. *Gen.* 45. *Spec.* 34. *Syst.* 45.

768. OLDENLANDIA.

Linn. *Gen.* 143. *Spec.* 119. *Syst.* 143. Ludw. 1032. Mill. ii. 84.

769. OLEA.

Linn. *Gen.* 20. *Spec.* 8. *Syst.* 20. Blackw. *tab.* 199. 213. Ludw.
2. Mill. ii. 85. Schæff. A. 319. Tourn. *tab.* 370. Weinm. *tab.*
764.

770. OLYRA.

Linn. *Syst. No.* 1168. p. 1379. 1261.

771. OMPHALEA.

Linn. *Syst. No.* 1166. p. 1378. 1264.

772. ONOCLEA.

Linn. *Gen.* 1034. *Spec.* 1062. *Syst.* 1034.

ANGIOPTERIS. Mich.

773. ONONIS.

Linn. *Gen.* 772. *Spec.* 716. *Syst.* 772. Schæff. A. 165.

ANONIS. Tourn. *tab.* 229. Blackw. *tab.* 301. Hall. 588. Ludw.
485. Mill. i. 59. Schæff. A. 165. Weinm. *tab.* 139. 141.

774. ONOPORDUM.

Linn. *Gen.* 834. *Spec.* 827. *Syst.* 834. Hall. 674. Ludw. 310.
CARDUI SPECIES. Tourn. *tab.* 253.

775. OPHIOGLOSSUM.

Linn. *Gen.* 1035. *Spec.* 1062. *Syst.* 1035. Hall. 131. Ludw.
952. Mill. ii. 89. Tourn. *tab.* 325. Weinm. *tab.* 765. c, f.

776. OPHIORRHIZA.

Linn. *Gen.* 193. *Spec.* 150. *Syst.* 193.

777. OPHIOXYLON.

Linn. *Gen.* 1011. *Spec.* 1043. *Syst.* 1011.

778. OPHIRYS.

Linn. *Gen.* 902. *Spec.* 945. *Syst.* 902.

OPHIRIS. Tourn. *tab.* 250. Hall. 277. Ludw. 702. Mill. ii. 125.

Weinm. *tab.* 769. a. BIFOLIUM. Mill. i. 125.

779. ORCHIS.

Linn. *Gen.* 900. *Spec.* 939. *Syst.* 900. Hall. 262. Ludw. 698.

Mill. ii. 92. Tourn. *tab.* 247. Weinm. *tab.* 767, *seq.*

780. ORIGANUM.

Linn. *Gen.* 645. *Spec.* 588. *Syst.* 645. Hall. 656. Ludw. 221.

ORIGANUM. Tourn. *tab.* 94. Blackw. *tab.* 280. Mill. ii. 93.

Schæff. A. 64. Weinm. *tab.* 772.

MAJORANA. Tourn. Blackw. *tab.* 319. Mill. ii. 2. Schæff. A.

56. Weinm. *tab.* 691, 692.

781. ORNITHOGALUM.

Linn. *Gen.* 377. *Spec.* 306. *Syst.* 377. Hall. 294. Ludw. 713.

Mill. ii. 94. iii. 200. Tourn. *tab.* 203. *A, B, H, I, K.* Weinm. *tab.* 773.

STELLARIS. Dill.

782. ORNITHOPUS.

Linn. *Gen.* 790. *Spec.* 743. *Syst.* 790.

ORNITHOPODIUM. Tourn. *tab.* 224. Hall. 572. Ludw. 502. Mill.

ii. 96. Weinm. *tab.* 773.

783. OROBANCHE.

Linn. *Gen.* 697. *Spec.* 632. *Syst.* 697. Hall. 610. Ludw. 231.

Tourn. *tab.* 81. Weinm. *tab.* 774.

APHYLLON. Mich.

784. OROBUS.

Linn. *Gen.* 780. *Spec.* 728. *Syst.* 780. Blackw. *tab.* 208. Hall.

602. Ludw. 489. Mill. ii. 96. Tourn. *tab.* 214. Weinm. *tab.*

775.

785. ORONTIUM.

Linn. *Gen.* 393. *Spec.* 324. *Syst.* 393.

ARONIA. Mitch.

786. ORTEGIA.

Linn. *Gen.* 49. *Spec.* 560. *Syst.* 49.

787. ORVALA.

Linn. *Gen.* 635. *Spec.* 578. *Syst.* 635. Ludw. 197.

PAPIA. Mich.

788. ORYZA.

Linn. *Gen.* 404. *Spec.* 333. *Syst.* 404. Ludw. 839. Mill. ii. 97.

Tourn. *tab.* 296. Weinm. *tab.* 775. f.

789. OSBECKIA.

Linn. *Gen.* 422. *Spec.* 345. *Syst.* 422.

790. OSMUNDA.

Linn. *Gen.* 1036. *Spec.* 1063. *Syst.* 1036. Blackw. *tab.* 324.

Hall. 130. Ludw. 951. Mill. ii. 98. Tourn. *tab.* 324.

791. OSTEOSPERMUM.

Linn. *Gen.* 887. *Spec.* 923. *Syst.* 887. Ludw. 350. Mill. iii. 202.

MONILIFERA. Vaill. CHRYSANTHEMOIDES. Tourn. Mill. i. 205.

Weinm. *tab.* 369.

792. OSYRIS.

Linn. *Gen.* 978. *Spec.* 1022. *Syst.* 978. Ludw. 920. Mill. iii. 203.

CASIA. Tourn. *tab.* 488. Mill. i. 179.

793. OTHONNA.

Linn. *Gen.* 888. *Spec.* 924. *Syst.* 888. Ludw. 355. Mill. iii. 203.

JACOBÆASTRUM. Vaill.

794. OVIEDA.

Linn. *Gen.* 705. *Spec.* 637. *Syst.* 705.

VALDIA. Plum. Ludw. 81.

795. OXALIS.

Linn. *Gen.* 515. *Spec.* 433. *Syst.* 515.

OXYS. Tourn. *tab.* 19. Blackw. *tab.* 308. Hall. 364. Ludw. 143.
Mill. ii. 98.

ACETOSELLA. Off. Schæff. A. 44. Weinm. *tab.* 20.

OXYOIDES. Garc. LUIULA. Sig.

796. PÆONIA.

Linn. *Gen.* 600. *Spec.* 530. *Syst.* 600. Blackw. *tab.* 65. 245. Hall.
310. Mill. ii. 99. Schæff. A. 212. Weinm. *tab.* 776. *seq.*

PÆONIA. Tourn. *tab.* 146. Ludw. 751.

797. PANAX.

Linn. *Gen.* 1031. *Spec.* 1058. *Syst.* No. 1031. p. 1314.

ARALIASTRUM. Vaill. Ludw. 532.

AURELIANA. Lafit.

NINSI. Breyn.

PANACEA. Mitch.

798. PANCRATIUM.

Linn. *Gen.* 365. *Spec.* 290. *Syst.* 365. Ludw. 728. Mill. ii. 103.

799. PANICUM.

Linn. *Gen.* 70. *Spec.* 55. *Syst.* 70. Hall. 233. Ludw. 819. Mill.
ii. 104. Weinm. *tab.* 786.

800. PAPAVER.

Linn. *Gen.* 573. *Spec.* 506. *Syst.* 573. Blackw. *tab.* 2. Hall. 303.
Ludw. 446. Mill. ii. 105. Schæff. A. 137, 138. Tourn. *tab.* 119.
Weinm. *tab.* 788. *seq.*

801. PARIETARIA.

Linn. *Gen.* 1020. *Spec.* 1052. *Syst.* 1020. Blackw. *tab.* 156. Hall.
177. Ludw. 763. Mill. ii. 106. Schæff. A. 272. Tourn. *tab.*
289. Weinm. *tab.* 798.

802. PARIS.

Linn. *Gen.* 449. *Spec.* 367. *Syst.* 449. Hall. 412. Ludw. 445.

Mill. iii. 206. Schæff. A. 155. Weinm. *tab.* 799. a, b, c.

HERBA PARIS. Tourn. *tab.* 117. Blackw. *tab.* 286. Mill. i. 402.

803. PARKINSONIA.

Linn. *Gen.* 460. *Spec.* 375. *Syst.* 460. Ludw. 643. Mill. ii. 107.

804. PARNASSIA.

Linn. *Gen.* 345. *Spec.* 273. *Syst.* 345. Hall. 316. Ludw. 516.
Mill. ii. 107. Tourn. *tab.* 127.

805. PARTHENIUM.

Linn. *Gen.* 939. *Spec.* 988. *Syst.* 939. Ludw. 311.

PARTHENIASTRUM. Niss. Mill. ii. 109. iii. 207.

HYSTEROPHORUS. Vaill.

806. PASPALUM.

Linn. *Syst. No.* 1107. *p.* 1359. 855.

807. PASSERINA.

Linn. *Gen.* 440. *Spec.* 559. *Syst.* 440. Hall. 187. Ludw. 790.
Mill. iii. 207.

SANAMUNDA. Magnol.

808. PASSIFLORA.

Linn. *Gen.* 910. *Spec.* 955. *Syst.* 910.

GRANADILLA. Tourn. *tab.* 124. Ludw. 539. Mill. i. 376. ii. 60.

MURUCUIA. Tourn. *tab.* 125. Mill. ii. 60.

809. PASTINACA.

Linn. *Gen.* 324. *Spec.* 262. *Syst.* 324. Blackw. *tab.* 379. Hall.
446. Ludw. 656. Mill. ii. 109. Schæff. *A.* 235. Tourn. *tab.*
170. Weinm. *tab.* 799.

810. PATAGONULA.

Linn. *Gen.* 191. *Spec.* 149. *Syst.* 191. Ludw. 72.

PATAGONICA. Dill.

811. PAVETTA.

Linn. *Gen.* 124. *Spec.* 110. *Syst.* 124.

812. PAULLINIA.

Linn. *Gen.* 446. *Spec.* 365. *Syst.* 446. Ludw. 441.

SERIANA. Plum. Mill. ii. 254.

CURURU. Plum. Mill. i. 255.

813. PECTIS.

Linn. *Syst.* No. 1160. p. 1376. 1221.

814. PEDALIUM.

Linn. *Syst.* No. 1155. p. 1375. 1123.

815. PEDICULARIS.

Linn. *Gen.* 664. *Spec.* 607. *Syst.* 664. Hall. 620. Ludw. 241.
Mill. ii. 111. Tourn. *tab.* 77. Weinm. *tab.* 800. A, D, H, I, K, L.

816. PEGANUM.

Linn. *Gen.* 530. *Spec.* 444. *Syst.* 530.

HARMALA. Tourn. *tab.* 133. Blackw. *tab.* 310. Ludw. 586. Mill.
i. 388.

817. PENÆA.

Linn. *Gen.* 129. *Spec.* 111. *Syst.* 129.

818. PENTAPETES.

Linn. *Gen.* 757. *Spec.* 698. *Syst.* 757.

PTEROSPERMADENDRON. Amm. Ludw. 1059.

819. PENTHORUM.

Linn. *Gen.* 514. *Spec.* 432. *Syst.* 514. Ludw. 800.

820. PEPLIS.

Linn. *Gen.* 402. *Spec.* 332. *Syst.* 402. Hall. 406. Ludw. 710.
ANDRACHNOIDES. Sigesb. PORTULA. Dill. GLAUCOIDES. Mich.

821. PERIPLOCA.

Linn. *Gen.* 267. *Spec.* 211. *Syst.* 267. Ludw. 102. Mill. ii. 113.
iii. 203. Tourn. *tab.* 22.

822. PETESIA.

Linn. *Syst.* No. 1113. p. 1361. 894.

TABLE II.

823. PETIVERIA.

Linn. *Gen.* 417. *Spec.* 342. *Syst.* 417. Ludw. 788. Mill. ii. 127.

824. PETREA.

Linn. *Gen.* 682. *Spec.* 626. *Syst.* 682. Ludw. 1040.

825. PEUCEDANUM.

Linn. *Gen.* 302. *Spec.* 245. *Syst.* 302. Hall. 442. Ludw. 668.
Mill. ii. 127. Schæff. A. 241. Tourn. *tab.* 169. Weinm. *tab.*
806.

826. PEZIZA.

Linn. *Gen.* 1080. *Spec.* 1180. *Syst.* 1080. Gled. *tab.* 4. Hall. 18.
Schæff. B. §. 70.

CYATHOIDES. Mich. Hall. 17. Ludw. 972.

FUNGOIDIS SPECIES. Vaill.

FUNGUS MEMBRANACEUS. Battafr. *tab.* 3.

827. PHACA.

Linn. *Gen.* 798. *Spec.* 755. *Syst.* 798. Mill. iii. 218.

ASTRAGALOIDES. Tourn. *tab.* 223. Hall. 569. Ludw. 481. Mill. i.
89.

828. PHALARIS.

Linn. *Gen.* 69. *Spec.* 54. *Syst.* 59. Ludw. 816.

829. PHALLUS.

Linn. *Gen.* 1077. *Spec.* 1178. *Syst.* 1077. Gled. *tab.* 1. Ludw.
968. Schæff. B. §. 70. G.

PHALLUS. Mich. Hall. 24.

BOLETUS. Mich. Hall. 23. MORCHELA. Dill. Battarr. *tab.* 2.

PHALLOBOLETUS. Mich. Battarr. *tab.* 40.

830. PHARMALEUM.

Linn. *Gen.* 341. *Spec.* 272. *Syst.* 341. Ludw. 783.

831. PHARUS.

Linn. *Syst.* 1169. *Spec.* p. 1379. 1269.

832. PHASGUM.

Linn. *Gen.* 1052. *Spec.* 1106. *Syst.* 1052.

833. PHASEOLUS.

Linn. *Gen.* 777. *Spec.* 723. *Syst.* 777. Ludw. 491. Mill. ii. 135.
Schæff. A. 169. *Tourn. tab.* 232. *Weinm. tab.* 807, *seq.*

834. PHELLANDRIUM.

Linn. *Gen.* 315. *Spec.* 255. *Syst.* 315. Hall. 432. Ludw. 694.
Tourn. tab. 161.

835. PHILADELPHUS.

Linn. *Gen.* 540. *Spec.* 470. *Syst.* 540.

SYRINGA. *Tourn. tab.* 389. Ludw. 463.

JASMINUM. *Weinm. tab.* 602. g.

836. PHILLYREA.

Linn. *Gen.* 19. *Spec.* 7. *Syst.* 19. Ludw. 3. Mill. ii. 137. iii. 219.
Tourn. tab. 367. *Weinm. tab.* 809. f.

837. PHLEUM.

Linn. *Gen.* 71. *Spec.* 59. *Syst.* 71. Hall. 231. Ludw. 817.

838. PHILOMIS.

Linn. *Gen.* 642. *Spec.* 584. *Syst.* 642. Ludw. 196. Mill. ii. 139.
Tourn. tab. 82.

839. PHLOX.

Linn. *Gen.* 197. *Spec.* 151. *Syst.* 197.

LYCHNOIDEA. Dill. Ludw. 69.

840. PHŒNIX.

Linn. *Gen.* 1089. *Spec.* 1188. *Syst.* 1089. Ludw. 908.

ELATE. Mus. Cliff.

KATOVINDEL. H. M.

841. PHRYMA.

Linn. *Gen.* 656. p. 601. *Syst.* 656.

LEPTOSTACHIA. Mitch.

842. PHYLICA.

Linn. *Gen.* 236. *Spec.* 195. *Syst.* 236. Mill. iii. 221.

PHILYCA. Ludw. 68.

843. PHYLLANTHUS.

Linn. *Gen.* 932. *Spec.* 981. *Syst.* 932. Blackw. *tab.* 400. Ludw. 868. Mill. iii. 222.

844. PHYLLIS.

Linn. *Gen.* 286. *Spec.* 232. *Syst.* 286. Ludw. 529. Mill. iii. 223.
BUPLEUROIDES. Mill. i. 149.

845. PHYSALIS.

Linn. *Gen.* 223. *Spec.* 182. *Syst.* 223.
ALKEKENGI. Tourn. *tab.* 64. Blackw. *tab.* 161. Hall. 508. Ludw. 86. Mill. i. 25. Weinm. *tab.* 931. a.

846. PHYTEUMA.

Linn. *Gen.* 203. *Spec.* 170. *Syst.* 203.
RAPUNCULUS. Tourn. *tab.* 38. Hall. 497. Ludw. 276. Mill. ii. 191. Weinm. *tab.* 862. d, e, f.

847. PHYTOLACCA.

Linn. *Gen.* 521. *Spec.* 444. *Syst.* 521. Ludw. 580. Mill. ii. 140.
Tourn. *tab.* 154.

848. PICRIS.

Linn. *Gen.* 812. *Spec.* 792. *Syst.* 812. Hall. 751. Ludw. 335.
HELMINTHOTHECA. Vaill.

849. PILULARIA.

Linn. *Gen.* 1347. *Spec.* 1100. *Syst.* 1047. Hall. 129. Ludw. 978.

850. PIMPINELLA.

Linn. *Gen.* 328. *Spec.* 263. *Syst.* 328. Ludw. 672. Schæff. A. 232.

TRAGOSELINUM. Tourn. *tab.* 163. Hall. 428. Mill. ii. 349. Weinm. *tab.* 811, 812.

ANISUM. Rai. Blackw. *tab.* 374. Ludw. 695. Weinm. *tab.* 138.

851. PINGUICULA.

Linn. *Gen.* 28. *Spec.* 17. *Syst.* 28. Hall. 611. Ludw. 1022. Mill. iii. 224. Tourn. *tab.* 74. Weinm. *tab.* 812. h.

852. PINUS.

Linn. *Gen.* 956. *Spec.* 1000. *Syst.* 956.

PINUS. Tourn. *tab.* 355. Blackw. *tab.* 189, 190. Hall. 149. Ludw. 885. Mill. ii. 142. Schæff. A. 297. Weinm. *tab.* 813.

ABIES. Tourn. *tab.* 353, 354. Blackw. *tab.* 203. Hall. 147. Ludw. 883. Mill. i. 1. iii. 1. Schæff. A. 298. Weinm. *tab.* 1. 3.

LARIX. Tourn. *tab.* 357. Hall. 148. Ludw. 884. Mill. i. 488. iii. 158. Schæff. A. 299. Weinm. *tab.* 627.

CEDRUS LIBANI. Edw. Mill. i. 186.

853. PIPER.

Linn. *Gen.* 42. *Spec.* 28. *Syst.* 42. Blackw. *tab.* 355, 356. Ludw. 938. Weinm. *tab.* 814, *seq.*

854. PISCIDIA.

Linn. *Gen.* No. 1158. *p.* 1376. 1155.

PISCIPULA. Loeffl. ICHTHYOMETRA. Brown.

855. PISONIA.

Linn. *Gen.* 984. *Spec.* 1026. *Syst.* 984. *p.* 1384. Blackw. *tab.* 348. Ludw. 904. Mill. ii. 143.

PENTAGONOTHECA. Vaill.

856. PISTACIA.

Linn. *Gen.* 982. *Spec.* 1025. *Syst.* 982.

TEREBINTHUS. Tourn. *tab.* 345. Ludw. 926. Mill. ii. 316. Weinm. *tab.* 816.

LENTISCUS. Tourn. Blackw. *tab.* 195. Ludw. 927. Mill. i. 498. Weinm. *tab.* 638. a.

857. PISTIA.

Linn. *Gen.* 912. *Spec.* 963. *Syst.* 912.

KODDA-PAIL. Plum.

ARISTOLOCHIA. Ludw. 283.

858. PISUM.

Linn. *Gen.* 779. *Spec.* 727. *Syst.* 779. Ludw. 487.

PISUM. Tourn. *tab.* 215. Blackw. *tab.* 83. Mill. ii. 144. Schæff. A. 163. Weimm. *tab.* 817, *seq.*

OCHRUS. Tourn. *tab.* 219, 220. Mill. ii. 81.

859. PLANTAGO.

Linn. *Gen.* 133. *Spec.* 112. *Syst.* 133. Hall. 470. Ludw. 22.

PLANTAGO. Tourn. *tab.* 48. Blackw. *tab.* 14. 35. Schæff. A. 13. Weimm. *tab.* 820, *seq.*

CORONOPUS. Tourn. *tab.* 49. Blackw. *tab.* 120. Mill. i. 236. Weimm. *tab.* 430. g.

PSYLLIUM. Tourn. *tab.* 49. Mill. ii. 165. Schæff. A. 14. Weimm. *tab.* 837.

860. PLATANUS.

Linn. *Gen.* 954. *Spec.* 999. *Syst.* 954. Ludw. 898. Mill. ii. 148. Tourn. *tab.* 363.

861. PLINIA.

Linn. *Gen.* 596. *Spec.* 516. *Syst.* 596. Ludw. 160. Mill. ii. 149.

862. PLUKENETIA.

Linn. *Gen.* 964. *Spec.* 1192. *Syst.* 964. Ludw. 1064.

863. PLUMBAGO.

Linn. *Gen.* 196. *Spec.* 151. *Syst.* 196. Ludw. 30. Mill. ii. 150. Tourn. *tab.* 58. Weimm. *tab.* 460. f.

864. PLUMERIA.

Linn. *Gen.* 263. *Spec.* 209. *Syst.* 263. Ludw. 104. Mill. ii. 150. Tourn. *tab.* 439.

865. POA.

Linn. *Gen.* 77. *Spec.* 67. *Syst.* 77. Hall. 211. Ludw. 831.

866. PODOPHYLLUM.

Linn. *Gen.* 571. *Spec.* 505. *Syst.* 571.

ANAPODOPHYLLON. Tourn. *tab.* 122. Ludw. 746. Mill. i. 42. iii. 19.

867. POINCIANA.

Linn. *Gen.* 462. *Spec.* 380. *Syst.* 462. Ludw. 642. Mill. ii. 151.
Tourn. *tab.* 391.

868. POLEMONIUM.

Linn. *Gen.* 200. *Spec.* 162. *Syst.* 200. Hall. 490. Ludw. 65.
Mill. ii. 153. Tourn. *tab.* 61.

VALERIANA GRÆCA. Weimn. *tab.* 1002. g.

869. POLIANTHES.

Linn. *Gen.* 384. *Spec.* 316. *Syst.* 384. Ludw. 123. Mill. iii. 229.
TUBEROSA. Heist. HYACINTHUS TUBEROSUS. Boerh. Mill. i. 415.

870. POLYCARPON.

Linn. *Syst.* 1110. *Spec. p.* 1360. 881.

871. POLYCNEMUM.

Linn. *Gen.* 51. *Spec.* 35. *Syst.* 51.

CAMPHORATA. Ludw. 765.

872. POLYGALA.

Linn. *Gen.* 761. *Spec.* 701. *Syst.* 761. Ludw. 386.

POLYGALA. Tourn. *tab.* 79. Hall. 606. Mill. ii. 154. Weimn. *tab.*
823.

CHAMÆBUXUS. Tourn. POLYGALOIDES. Dill. Hall. 607. Weimn.
tab. 301. a.

PENÆA. Plum.

HEISTERIA. Linn. *edit. prior.*

873. POLYGONUM.

Linn. *Gen.* 445. *Spec.* 359. *Syst.* 445.

POLYGONUM. Tourn. *tab.* 290. Blackw. *tab.* 315. Hall. 182.
Ludw. 793. Schæff. A. 282. Weimn. *tab.* 824, *seq.* CENTINO-
DIUM. Mill. i. 192.

BISTORTA. Tourn. *tab.* 291. Blackw. *tab.* 254. Hall. 178. Ludw.
795. Mill. i. 127. Schæff. A. 280. Weimn. *tab.* 244, 245.

PERSICARIA. Tourn. *tab.* 290. Blackw. *tab.* 118, 119. Hall. 179.
Ludw. 778. Mill. ii. 125. Schæff. A. 279. Weimn. *tab.* 803,
seq.

FAGOPYRUM. Tourn. *tab.* 290. Hall. 172. Ludw. 794. Mill. i.
309. Schæff. A. 281. Weinm. *tab.* 501. d.

HELXINE. Linn. *edit. prior.*

HYDROPIPER. Dod. Mill. i. 417.

874. POLYMNIA.

Linn. *Gen.* 889. *Spec.* 926. *Syst.* 889.

875. POLYPODIUM.

Linn. *Gen.* 1043. *Spec.* 1082. *Syst.* 1043. Hall. 137. Ludw. 944.

POLYPODIUM. Tourn. *tab.* 316. Blackw. *tab.* 215. Mill. ii. 156.
Schæff. A. 310. Weinm. *tab.* 825. h.

LONCHITIS. Tourn. *tab.* 314. Mill. i. 324.

FILIX. T. Blackw. *tab.* 323. Schæff. A. 311. Weinm. *tab.* 510,
seq.

876. POLYPREMUM.

Linn. *Gen.* 128. *Spec.* 111. *Syst.* 128.

877. POLYTRICHUM.

Linn. *Gen.* 1055. *Spec.* 1109. *Syst.* 1055. Ludw. 956. Hall. 106.

ADIANTHUM AUREUM. Rai. Blackw. *tab.* 371.

878. PONTEDERIA.

Linn. *Gen.* 361. *Spec.* 288. *Syst.* 361. Ludw. 284.

MICHELIA. Houst.

879. POPULUS.

Linn. *Gen.* 996. *Spec.* 1034. *Syst.* 996. Blackw. *tab.* 248. Hall.
156. Ludw. 932. Mill. ii. 157. Schæff. A. 296. Tourn. *tab.*
365. Weinm. *tab.* 826, *seq.*

880. PORELLA.

Linn. *Gen.* 1050. *Spec.* 1106. *Syst.* 1050.

881. PORTLANDIA.

Linn. *Syst.* No. 1121. *p.* 1364. 928.

882. PORTULACA.

Linn. *Gen.* 531. *Spec.* 445. *Syst.* 531. Blackw. *tab.* 287. Hall. 392. Ludw. 582. Mill. ii. 159. iii. 16. Schæff. A. 173. Tourn. *tab.* 118. Weinm. *tab.* 828.

ANACAMPSEROS. Linn. *edit. prior.*

TELEPHIASTUM. Dill.

883. POTAMOGETON.

Linn. *Gen.* 160. *Spec.* 126. *Syst.* 160. Hall. 199. Ludw. 397. Tourn. *tab.* 103. Weinm. *tab.* 829, *seq.*

884. POTENTILLA.

Linn. *Gen.* 559. *Spec.* 495. *Syst.* 559. Ludw. 621. Mill. iii. 232.

QUINQUEFOLIUM. Tourn. *tab.* 153. Hall. 340. Mill. ii. 184.

Schæff. A. 206. Weinm. *tab.* 847. PENTAPHYLLUM. Rai.

PENTAPHYLLOIDES. Tourn. Mill. ii. 111.

ANSERINA. Blackw. *tab.* 6. Schæff. A. 207. Weinm. *tab.* 142.

885. POTERIUM.

Linn. *Gen.* 948. *Spec.* 994. *Syst.* 948. Ludw. 19.

PIMPINELLA. Tourn. *tab.* 68. Hall. 469. Mill. ii. 141.

886. POTHOS.

Linn. *Gen.* 918. *Spec.* 968. *Syst.* 918.

887. PRASIUM.

Linn. *Gen.* 655. *Spec.* 601. *Syst.* 655. Ludw. 195. Mill. iii. 233.

888. PRENANTHES.

Linn. *Gen.* 816. *Spec.* 797. *Syst.* 816. Hall. 754. Ludw. 336. Mill. iii. 233.

889. PRIMULA.

Linn. *Gen.* 180. *Spec.* 142. *Syst.* 180. Hall. 482. Ludw. 50.

PRIMULA VERIS. Tourn. *tab.* 47. Blackw. *tab.* 52. 226. Mill. ii.

159. Schæff. A. 25. Weinm. *tab.* 831, *seq.*

AURICULA URSI. Tourn. *tab.* 46. Mill. i. 98. iii. 34. Weinm. *tab.* 207. 216.

890. PRINOS.

Linn. *Gen.* 398. *Spec.* 330. *Syst.* 398. Ludw. 128. Mill. iii. 234.

891. PROCKIA.

Linn. *Syst. No.* 1148. p. 1372. 1074.

892. PROSERPINACA.

Linn. *Gen.* 97. *Spec.* 88. *Syst.* 97.

TRIXIS. Mich.

893. PROTEA.

Linn. *Gen.* 104. *Spec.* 94. *Syst.* 104. Ludw. 292. Mill. iii. 234.

CONOCARPODENDRON. Boerh.

894. PRUNELLA.

Linn. *Gen.* 654. *Spec.* 600. *Syst.* 654.

BRUNELLA. Tourn. *tab.* 84. Blackw. *tab.* 24. Hall. 636. Ludw.

201. Mill. i. 144. Schæff. A. 73. Weinm. *tab.* 268.

CONSOLIDA MINOR. Offic.

895. PRUNUS.

Linn. *Gen.* 546. *Spec.* 473. *Syst.* 546.

PRUNUS. Tourn. *tab.* 398. Blackw. *tab.* 305. Hall. 355. Ludw.

594. Mill. ii. 161. Schæff. A. 187. Weinm. *tab.* 835.

ARMENIACA. Tourn. *tab.* 399. Blackw. *tab.* 281. Ludw. 595.

Mill. i. 68. Weinm. *tab.* 698.

CERASUS. Tourn. *tab.* 401. Hall. 356. Ludw. 593. Mill. i. 194.

Schæff. A. 188. Weinm. *tab.* 350, *seq.*

PADUS. Linn. *edit. prior.* Hall. 357. Ludw. 592. Mill. iii.

203. LAUKOCERASUS. Tourn. *tab.* 403. Mill. i. 493.

896. PSIDIUM.

Linn. *Gen.* 541. *Spec.* 470. *Syst.* 541.

GUAIAVA. Tourn. *tab.* 443. Ludw. 602. Mill. i. 384. Weinm.

tab. 561. h.

897. PSORALEA.

Linn. *Gen.* 801. *Spec.* 762. *Syst.* 801. Ludw. 637.

DALEA. Linn. *edit. prior.* Mill. iii. 87.

898. PSYCHOTRIA.

Linn. *Syst.* No. 1122. *p.* 1364. 929.

PSYCHOTROPHUM. Broun.

899. PTELEA.

Linn. *Gen.* 141. *Spec.* 118. *Syst.* 141. Ludw. 1031. Mill. iii. 235.

900. PTERIS.

Linn. *Gen.* 1038. *Spec.* 1073. *Syst.* 1038. Ludw. 946.

FILIX. Hall. 132.

901. PULMONARIA.

Linn. *Gen.* 169. *Spec.* 135. *Syst.* 169. Blackw. *tab.* 376. Hall.
516. Ludw. 37. Mill. ii. 167. Schæff. A. 36. Tourn. *tab.* 55.
Weinm. *tab.* 956. b.

902. PUNICA.

Linn. *Gen.* 544. *Spec.* 472. *Syst.* 544. Blackw. *tab.* 97. 145.
Ludw. 736. Mill. ii. 169. Schæff. A. 257. Tourn. *tab.* 407.

903. PYROLA.

Linn. *Gen.* 490. *Spec.* 396. *Syst.* 490. Ludw. 650. Mill. ii. 170.
Schæff. A. 218. Tourn. *tab.* 134. Weinm. *tab.* 841.

PIROLA. Hall. 420.

904. PYRUS.

Linn. *Gen.* 550. *Spec.* 479. *Syst.* 550. Hall. 351. Ludw. 618.

PYRUS. Tourn. *tab.* 404. Mill. ii. 171. iii. 173. Weinm. *tab.* 842.
MALUS. Tourn. *tab.* 406. Black. *tab.* 144. Mill. ii. 5. Weinm.
tab. 704.

CYDONIA. Tourn. *tab.* 405. Blackw. *tab.* 137. Mill. i. 257. Schæff.
A. 200. Weinm. *tab.* 703.

905. QUERCUS.

Linn. *Gen.* 949. *Spec.* 994. *Syst.* 949. Ludw. 890.

QUERCUS. Tourn. *tab.* 349. Hall. 159. Mill. ii. 181. Schæff. A.
293. Weinm. *tab.* 845.

ILEX. Tourn. *tab.* 350. Blackw. *tab.* 186. Weinm. *tab.* 603.

SUBER. Tourn. Blackw. *tab.* 193. Mill. ii. 306. Weinm. *tab.* 954.

906. QUERIA.

Linn. *Gen.* 101. *Spec.* 90. *Syst.* 101.

907. RAIANIA.

Linn. *Gen.* 994. *Spec.* 1032. *Syst.* 994. Ludw. 930.

JAN-RAIA. Plum.

908. RANDIA.

Linn. *Gen.* 194. *Spec.* 1192. *Syst.* 194. Ludw. 1065. Mill. ii. 185.

909. RANUNCULUS.

Linn. *Gen.* 619. *Spec.* 548. *Syst.* 619.

RANUNCULUS. Tourn. *tab.* 149. A. C. Blackw. *tab.* 31. Hall. 321.

Ludw. 623. Mill. ii. 186. iii. 244. Schæff. A. 213. Weinm. *tab.* 849, *seq.*

FICARIA. Dill. Hall. 321. Ludw. 755. CHELIDONIUM MINUS.

Boerh. Blackw. *tab.* 51. Mill. i. 203. Schæff. A. 261. Weinm. *tab.* 366. b. MARISCA. Sigb.

RANUNCULOIDES. Vaill.

910. RAPHANUS.

Linn. *Gen.* 736. *Spec.* 669. *Syst.* 736.

RAPHANUS. Tourn. *tab.* 114. Blackw. *tab.* 81. Hall. 555. Ludw.

404. Mill. ii. 190. Schæff. A. 145. Weinm. *tab.* 860, *seq.*

RAPHANISTRUM. Tourn. *tab.* 115. Ludw. 402. Mill. ii. 190.

911. RAUWOLFIA.

Linn. *Gen.* 259. *Spec.* 208. *Syst.* 259. Ludw. 74. Mill. ii. 194.

912. REAUMURIA.

Linn. *Syst. No.* 1152. *p.* 1374. 1081.

913. RENEALMIA.

Linn. *Gen.* 358. *Spec.* 286. *Syst.* 358. Ludw. 125.

914. RESEDA.

Linn. *Gen.* 535. *Spec.* 448. *Syst.* 535.

RESEDA. Tourn. *tab.* 238. Hall. 315. Ludw. 737. Mill. ii. 199.
Weinm. *tab.* 863.

LUTEOLA. Tourn. *tab.* 238. Blackw. *tab.* 283. Hall. 315. Ludw.
738. Mill. i. 542. Weinm. *tab.* 676.

SESAMOIDES. Tourn. *tab.* 238. Ludw. 739.

915. RHACOMA.

Linn. *Syst. No.* 1114. *p.* 1361. 896.

CROSSOPETALUM. Brown.

916. RHAMNUS.

Linn. *Gen.* 235. *Spec.* 193. *Syst.* 235.

RHAMNUS. Tourn. *tab.* 366. Hall. 163. Ludw. 84. Mill. ii. 199.
Weinm. *tab.* 864. a, b.

FRANGULA. Tourn. *tab.* 383. Blackw. *tab.* 152. Hall. 164.
Ludw. 76. Mill. i. 329. Schæff. A. 4. Weinm. *tab.* 514. a.

CERVISPINA. Dill. Blackw. *tab.* 135. Ludw. 903. Schæff. A. 2.
Weinm. *tab.* 945. c.

PALIURUS. Tourn. *tab.* 381. Ludw. 112. Mill. ii. 100.

ALATERNUS. Tourn. *tab.* 366. Ludw. 82. Mill. i. 23. iii. 10.
Weinm. *tab.* 31.

ZIZIPHUS. Tourn. *tab.* 403. Ludw. 77. JUJUBE. C. B. Mill. ii.
545. Schæff. A. 5.

917. RHEEDIA.

Linn. *Gen.* 1102. *Spec.* 1193. *Syst.* 1102.

VANRHEEDIA. Plum.

918. RHEUM.

Linn. *Gen.* 454. *Spec.* 371. *Syst.* 454. Ludw. 138.

RHABARBARUM. Tourn. *tab.* 18.

919. RHEXIA.

Linn. *Gen.* 423. *Spec.* 346. *Syst.* 423. Ludw. 433.

920. RHINANTHUS.

Linn. *Gen.* 658. *Spec.* 603. *Syst.* 658.

PEDICULARIS SPECIES. Tourn. *tab.* 77. *B, F, M, N, O, P.* Weinm. *tab.* 800.

ELEPHAS. Tourn. *tab.* 482. Ludw. 246. Mill. i. 292.

CRISTA GALLI. Riv. ALECTOROLOPHUS. Hall. 623. Ludw. 242.

921. RHIZOPHORA.

Linn. *Gen.* 524. *Spec.* 443. *Syst.* 524.

MANGLES. Plum. Ludw. 1041.

922. RHODIOLA.

Linn. *Gen.* 997. *Spec.* 1035. *Syst.* 997.

SEDUM. Hall. 394. Ludw. 613.

RHODIA RADIX. Off. Schæff. A. 134.

923. RHODODENDRON.

Linn. *Gen.* 481. *Spec.* 392. *Syst.* 484.

CHAMERHODODENDROS. Tourn. *tab.* 373. Mill. i. 202. iii. 60.

924. RHUS.

Linn. *Gen.* 331. *Spec.* 265. *Syst.* 331.

RHUS. Tourn. *tab.* 381. Ludw. 535. Mill. ii. 200. Weinm. *tab.* 864.

TOXICODENDRON. Tourn. *tab.* 381. Ludw. 535. Mill. ii. 344.

COTINUS. Tourn. *tab.* 380. Ludw. 534. Mill. i. 238. iii. 80. Weinm. *tab.* 432.

VERNIX. Kæmpf. Ludw. 535.

925. RIBES.

Linn. *Gen.* 247. *Spec.* 200. *Syst.* 247. Hall. 345. Ludw. 533.

RIBES. Tourn. Blackw. *tab.* 285. Mill. ii. 201. Schæff. A. 177. Weinm. *tab.* 865. RIBESIUM. Dill.

GROSSULARIA. Tourn. *tab.* 409. Blackw. *tab.* 277. Mill. i. 381. Weinm. *tab.* 558.

926. RICCIA.

Linn. *Gen. No.* 1063. *p.* 1138. *Syst.* 1063. Hall. 128. Ludw. 980.

927. RICHARDIA.

Linn. *Gen.* 397. *Spec.* 330. *Syst.* 397. Ludw. 114.

928. RICINUS.

Linn. *Gen.* 962. *Spec.* 1007. *Syst.* 962. Ludw. 890. Mill. ii. 203.

Tourn. *tab.* 397. Weinm. *tab.* 866.

PALMA CHRISTI. Mill. ii. 203.

CATAPUTIA MAJOR. Ruel. Blackw. *tab.* 148. Schæff. A. 283.

929. RIVINA.

Linn. *Gen.* 150. *Spec.* 121. *Syst.* 150. Mill. iii. 250.

RIVINIA. Plum. SOLANOIDES. Tourn. Ludw. 769. Mill. ii. 268.

930. ROBINIA.

Linn. *Gen.* 775. *Spec.* 722. *Syst.* 775. Ludw. 495. Mill. iii. 251.

PSEUDOACACIA. Tourn. *tab.* 417. Mill. ii. 163. LABURNUM. Sigesb.

931. ROELLA.

Linn. *Gen.* 202. *Spec.* 170. *Syst.* 202. Ludw. 62.

932. RONDELETIA.

Linn. *Gen.* 206. *Spec.* 172. *Syst.* 206. Ludw. 60. Mill. ii. 205.

933. ROSA.

Linn. *Gen.* 556. *Spec.* 491. *Syst.* 556. Blackw. *tab.* 8. 78. 82.

Hall. 347. Ludw. 623. Mill. ii. 205. iii. 252. Schæff. A. 201,
202. Tourn. *tab.* 408. Weinm. *tab.* 868.

934. ROSMARINUS.

Linn. *Gen.* 35. *Spec.* 23. *Syst.* 35. Blackw. *tab.* 159. Ludw. 177.

Mill. ii. 209. Schæff. A. 57. Tourn. *tab.* 92. Weinm. *tab.* 872.
a, b.

935. ROYENA.

Linn. *Gen.* 491. *Spec.* 397. *Syst.* 491. Ludw. 141. Mill. ii. 253.

936. RUBIA.

Linn. *Gen.* 119. *Spec.* 109. *Syst.* 119. Blackw. *tab.* 326. Hall. 462. Ludw. 25. Mill. ii. 209. Schæff. A. 15. Tourn. *tab.* 38. Weinm. *tab.* 873.

937. RUBUS.

Linn. *Gen.* 557. *Spec.* 492. *Syst.* 557. Blackw. *tab.* 45. 279. Hall. 343. Ludw. 627. Mill. iii. 211. Schæff. A. 205. Tourn. *tab.* 385. Weinm. *tab.* 874.

938. RUDBECKIA.

Linn. *Gen.* 878. *Spec.* 906. *Syst.* 878. Ludw. 361. Mill. iii. 254. OBELISCOTHECA. Vaill. CALCANTHEMUM. Mill. ii. 81.

939. RUELLIA.

Linn. *Gen.* 702. *Spec.* 634. *Syst.* 702. Ludw. 235. Mill. ii. 212.

940. RUMEX.

Linn. *Gen.* 407. *Spec.* 333. *Syst.* 407. Ludw. 786. ACETOSA. Tourn. *tab.* 287. Blackw. *tab.* 130. 262. 306, 307. Hall. 169. Mill. i. 16. Schæff. A. 287. Weinm. *tab.* 28. LAPATHUM. Tourn. Hall. 168. 172. Mill. i. 487. Schæff. A. 288. Weinm. *tab.* 624, *seq.*

941. RUMPHIA.

Linn. *Gen.* 1103. *Spec.* 1193. *Syst.* 1103. Ludw. 377.

942. RUPPIA.

Linn. *Gen.* 161. *Spec.* 127. *Syst.* 161. Ludw. 941. BUCCA FERREA. Mich.

943. RUSCUS.

Linn. *Gen.* 1008. *Spec.* 1041. *Syst.* 1008. Blackw. *tab.* 155. Ludw. 902. Mill. ii. 212. Schæff. A. 286. Tourn. *tab.* 15. Weinm. *tab.* 875. a, b.

944. RUTA.

Linn. *Gen.* 469. *Spec.* 383. *Syst.* 469. Blackw. *tab.* 7. Hall. 411.

Ludw. 436. Mill. ii. 213. Schæff. A. 133. Tourn. *tab.* 133.

Weinm. *tab.* 875. c, d, e.

PSEUDORUTA. Mich.

945. SACCHARUM.

Linn. *Gen.* 68. *Spec.* 54. *Syst.* 68. Ludw. 815.

946. SAGINA.

Linn. *Gen.* 162. *Spec.* 128. *Syst.* 162.

ALSINELLA. Dill. ALSINE. Hall. 390. Ludw. 569.

947. SAGITTARIA.

Linn. *Gen.* 946. *Spec.* 993. *Syst.* 946. Hall. 300. Ludw. 861.

SAGITTA. Dill. Weinm. *tab.* 876. b, c.

948. SALICORNIA.

Linn. *Gen.* 10. *Spec.* 3. *Syst.* 10. Ludw. 936. Mill. ii. 227.

Tourn. *tab.* 485.

949. SALIX.

Linn. *Gen.* 976. *Spec.* 1015. *Syst.* 976. Blackw. *tab.* 327. Ludw.

918. Hall. 151. Mill. ii. 227. iii. 258. Tourn. *tab.* 364. Weinm.

tab. 877, *seq.*

950. SALSOLA.

Linn. *Gen.* 275. *Spec.* 222. *Syst.* 275.

KALI. Tourn. *tab.* 128. Ludw. 773. Mill. i. 470. Weinm. *tab.*

616.

951. SALVADORA.

Linn. *Gen.* 151. *Spec.* 122. *Syst.* 151.

952. SALVIA.

Linn. *Gen.* 36. *Spec.* 23. *Syst.* 36. Hall. 638. Ludw. 176.

SALVIA. Tourn. *tab.* 83. Blackw. *tab.* 10. Mill. ii. 229. Schæff.

A. 62. Weinm. *tab.* 879, *seq.*

HORMINUM. Tourn. *tab.* 82. Blackw. *tab.* 258. Mill. i. 408.
Weinm. *tab.* 578.

SCLAREA. Tourn. *tab.* 82. Blackw. *tab.* 122. Mill. ii. 244.
Schæff. A. 63. Weinm. *tab.* 579.

953. SAMBUCUS.

Linn. *Gen.* 334. *Spec.* 269. *Syst.* 334. Blackw. *tab.* 151. Hall.
465. Ludw. 111. Mill. ii. 230. Schæff. A. 23. Tourn. *tab.* 376.
Weinm. *tab.* 881.

954. SAMOLUS.

Linn. *Gen.* 205. *Spec.* 171. *Syst.* 205. Ludw. 43. Mill. ii. 231.
Tourn. *tab.* 60.

955. SAMYDA.

Linn. *Gen.* 525. *Spec.* 443. *Syst. No.* 525. p. 1024. 1382. Ludw.
1068.

GUIDONIA. Plum. Mill. i. 385. iii. 126.

956. SANGUINARIA.

Linn. *Gen.* 570. *Spec.* 505. *Syst.* 570. Ludw. 748. Mill. iii. 259.

957. SANGUISORBA.

Linn. *Gen.* 136. *Spec.* 116. *Syst.* 136. Ludw. 19. Mill. iii. 260.
PIMPINELLA. Tourn. *tab.* 68. Hall. 469. Mill. ii. 141. Weinm.
tab. 810.

958. SANICULA.

Linn. *Gen.* 289. *Spec.* 235. *Syst.* 289. Blackw. *tab.* 63. Hall.
449. Ludw. 679. Mill. ii. 232. Schæff. A. 229. Tourn. *tab.*
173. Weinm. *tab.* 835. a.

959. SANTALUM.

Linn. *Gen.* 431. *Spec.* 349. *Syst.* 431. Ludw. 136. Weinm. *tab.*
883.

960. SANTOLINA.

Linn. *Gen.* 847. *Spec.* 842. *Syst.* 847. Blackw. *tab.* 346. Ludw.
312. Mill. ii. 232. Tourn. *tab.* 260.

BACCHARIS. Vaill.

961. SAPINDUS.

Linn. *Gen.* 448. *Spec.* 367. *Syst.* 448. Ludw. 443. Mill. ii. 233.
Tourn. *tab.* 440.

962. SAPONARIA.

Linn. *Gen.* 499. *Spec.* 408. *Syst.* 499. Blackw. *tab.* 113. Hall.
378. Ludw. 564. Schæff. A. 180. Weinm. *tab.* 686. c.

963. SAROTHRA.

Linn. *Gen.* 344. *Spec.* 272. *Syst.* 344.

964. SARRACENIA.

Linn. *Gen.* 578. *Spec.* 510. *Syst.* 578. Ludw. 589. Tourn. *tab.*
476.

965. SATUREIA.

Linn. *Gen.* 626. *Spec.* 567. *Syst.* 626. Ludw. 209.

SATUREIA. Tourn. Blackw. *tab.* 318. Mill. ii. 235. Schæff. A. 71.
Weinm. *tab.* 885. c.

THYMBRA. Tourn. Weinm. *tab.* 975. c.

966. SATYRIUM.

Linn. *Gen.* 901. *Spec.* 944. *Syst.* 901. Blackw. *tab.* 53. Schæff.
A. 244.

ORCHIS. Hall. 262. Ludw. 698.

967. SAURURUS.

Linn. *Gen.* 414. *Spec.* 341. *Syst.* 414. Ludw. 785. Mill. ii. 235.

968. SAUVAGESIA.

Linn. *Gen.* 252. *Spec.* 203. *Syst.* 252.

SAUVAGEA. Ludw. 747.

969. SAXIFRAGA.

Linn. *Gen.* 494. *Spec.* 398. *Syst.* 494. Hall. 399. Ludw. 566.

SAXIFRAGA. Tourn. *tab.* 129. Blackw. *tab.* 56. Mill. ii. 236.
Schæff. A. 182. Weinm. *tab.* 886.

GEUM. Tourn. *tab.* 129. Mill. i. 355.

970. SCABIOSA.

Linn. *Gen.* 108. *Spec.* 98. *Syst.* 108. Ludw. 289.

SCABIOSA. Tourn. *tab.* 263, 264. Blackw. *tab.* 185. Hall. 669.

Mill. ii. 258. Schæff. A. 94. Weinm. *tab.* 886, *seq.*

SUCCISA. Vaill. Blackw. *tab.* 142. Hall. 671. MORSUS DIABOLI.

Schæff. A. 95. Weinm. *tab.* 889. d, e, f.

ASTEROCEPHALUS. Vaill. Hall. 668.

PTEROCEPHALUS. Vaill.

971. SCANDIX.

Linn. *Gen.* 319. *Spec.* 256. *Syst.* 319.

SCANDIX. Tourn. *tab.* 173. Hall. 454. Ludw. 673. Mill. ii. 241.

Weinm. *tab.* 290.

MYRRHIS. Tourn. Hall. 453. Ludw. 674. Mill. i. 63. Weinm.

tab. 749. b.

CEREFOLIUM. Riv. CHÆROPHYLLUM. Blackw. Tourn. *tab.* 166.

236. Hall. 452. Ludw. 674. Schæff. A. 83.

972. SCHEUCHZERIA.

Linn. *Gen.* 408. *Spec.* 338. *Syst.* 408. Ludw. 787. Hall. 258.

973. SCHINUS.

Linn. *Gen.* 479. *Spec.* 388. *Syst. No.* 479. p. 1034.

MOLLE. Tourn. Ludw. 1042. Mill. ii. 55.

974. SCHÆNUS.

Linn. *Gen.* 60. *Spec.* 42. *Syst.* 60. Ludw. 842.

CYPERELLA. Mich.

PSEUDOCYPERUS. Mich.

MELANOSCHÆNUS. Mich.

975. SCHWALBEA.

Linn. *Gen.* 662. *Spec.* 606. *Syst.* 662. Ludw. 188.

976. SCILLA.

Linn. *Gen.* 378. *Spec.* 308. *Syst.* 378.

SCILLA. Ludw. 712. Mill. ii. 243. Schæff. A. 248. Weinm. *tab.* 890. d.

LILIOHYACINTHUS. Tourn. *tab.* 196. B, F, G. Ludw. 713. Mill. i. 509.

HYACINTHUS STELLARIS. Rai. Weinm. *tab.* 587.

977. SCIRPUS.

Linn. *Gen.* 62. *Spec.* 47. *Syst.* 62. Hall. 247. Ludw. 841.
Tourn. *tab.* 300.

978. SCLERANTHUS.

Linn. *Gen.* 497. *Spec.* 406. *Syst.* 497. Ludw. 793.
KNAWEL. Rai. Hall. 186.

979. SCOLYMUS.

Linn. *Gen.* 826. *Spec.* 813. *Syst.* 826. Ludw. 343. Mill. ii. 244.
Tourn. *tab.* 273. Weinm. *tab.* 906.

980. SCOPARIA.

Linn. *Gen.* 134. *Spec.* 116. *Syst.* 134.

981. SCORPIURUS.

Linn. *Gen.* 792. *Spec.* 744. *Syst.* 792. Ludw. 501.
SCORPIOIDES. Tourn. *tab.* 226. Mill. ii. 245.

982. SCORZONERA.

Linn. *Gen.* 811. *Spec.* 790. *Syst.* 811. Ludw. 341.
SCORZONERA. Tourn. *tab.* 269. Hall. 757. Mill. ii. 246. Schæff.
A. 112.
SCORZONEROIDES. Vaill.

983. SCROPHULARIA.

Linn. *Gen.* 674. *Spec.* 619. *Syst.* 674. Blackw. *tab.* 86, 87.
Hall. 618. Ludw. 252. Mill. ii. 246. Schæff. A. 79. Tourn.
tab. 74, Weinm. *tab.* 908, *seq.*

984. SCURRULA.

Linn. *Gen.* 123. *Spec.* 110. *Syst.* 123.

985. SCUTELLARIA.

Linn. *Gen.* 653. *Spec.* 598. *Syst.* 653. Mill. iii. 261. Weinm.
tab. 909. d.
CASSIDA. Tourn. *tab.* 84, Hall. 635. Ludw. 202. Mill. i. 180.

TABLE II.

986. SECALE.

Linn. *Gen.* 92. *Spec.* 84. *Syst.* 92. Ludw. 826. Mill. ii. 248.
Weinm. *tab.* 910.

987. SECURIDACA.

Linn. *Gen.* 763. *Spec.* 707. *Syst.* 763. Ludw. 507. Mill. ii. 248,

988. SEDUM.

Linn. *Gen.* 513. *Spec.* 430. *Syst.* 513. Ludw. 613.

SEDUM. Tourn. *tab.* 140. *A, B, G, H, K, M.* Blackw. *tab.* 366.

Hall. 392. Mill. ii. 248. Schæff. *A.* 184. Weinm. *tab.* 911,
seq.

ANACAMPSEROS. Tourn. Mill. i. 36. iii. 16.

989. SEGUIERIA.

Linn. *Syst. No.* 1150. *p.* 1373. 1074.

990. SELAGO.

Linn. *Gen.* 687. *Spec.* 629. *Syst.* 687.

CAMPHORATA. Ludw. 765.

991. SELINUM.

Linn. *Gen.* 300. *Spec.* 244. *Syst.* 300. Hall. 443. Ludw. 665.

THYSSELINUM. Tourn.

992. SEMPERVIVUM.

Linn. *Gen.* 538. *Spec.* 464. *Syst.* 538. Schæff. *A.* 263.

SEDI SPECIES. Tourn. *tab.* 140. *C, E, I.* Hall. 392. Ludw. 613.

993. SENECIO.

Linn. *Gen.* 857. *Spec.* 866. *Syst.* 857.

SENECIO. Tourn. *tab.* 260. Blackw. *tab.* 132. Hall. 730. Ludw.

303. Mill. ii. 252. iii. 263. Schæff. *A.* 120. Weinm. *tab.* 915. *d.*

JACOBÆE SPECIES. Tourn. Mill. i. 445.

994. SERAPIAS.

Linn. *Gen.* 903. *Spec.* 949. *Syst.* 903.

HELLEBORINE. Tourn. *tab.* 249. Hall. 274. Ludw. 699. Mill. i.

398. Weinm. *tab.* 567.

995. SERIPHIMUM.

Linn. *Gen.* 894. *Spec.* 928. *Syst.* 894.

HELICHRYSOIDES. Vaill. Ludw. 305.

996. SERRATULA.

Linn. *Gen.* 831. *Spec.* 816. *Syst.* 831. Ludw. 318. Mill. ii. 255.
Weinm. *tab.* 917.

997. SESAMUM.

Linn. *Gen.* 700. *Spec.* 634. *Syst.* 700. Ludw. 261. Mill. ii. 256.
Weinm. *tab.* 918.

998. SESELI.

Linn. *Gen.* 322. *Spec.* 259. *Syst.* 322. Hall. 430. Ludw. 676.
Mill. ii. 256. Weinm. *tab.* 918. d.

999. SESUVIUM.

Linn. *Syst. No.* 1143. *p.* 1371. 1058.

HALIMUM. Loeffl.

1000. SHERARDIA.

Linn. *Gen.* 112. *Spec.* 102. *Syst.* 112. Hall. 457. Ludw. 13.
Mill. ii. 257.

DILLENIA. Heist.

1001. SIBBALDIA.

Linn. *Gen.* 354. *Spec.* 284. *Syst.* 354. Hall. 342.

SIBBALDIA. Ludw. 543.

1002. SIBTHORPIA.

Linn. *Gen.* 693. *Spec.* 631. *Syst.* 693.

1003. SICYOS.

Linn. *Gen.* 971. *Spec.* 1013. *Syst.* 971. Ludw. 857.

SICYOIDES. Tourn. *tab.* 28. Mill. ii. 258.

BRYONIOIDES. Dill.

1004. SIDA.

Linn. *Gen.* 747. *Spec.* 683. *Syst.* 747.

MALVINDA. Dill. Ludw. 148.

ABUTILON. Tourn. *tab.* 25. Ludw. 153. Mill. iii. 6.

1005. SIDERITIS.

Linn. *Gen.* 632. *Spec.* 574. *Syst.* 632. Hall. 647. Ludw. 203.

Mill. ii. 258. Schæff. A. 86. Tourn. *tab.* 90. Weinm. *tab.* 919.

1006. SIDEROXYLON.

Linn. *Gen.* 234. *Spec.* 192. *Syst.* 234. Hall. 647. Mill. ii. 258.

iii. 265.

SIDEROXYLUM. Ludw. 85.

1007. SIGESBECKIA.

Linn. *Gen.* 873. *Spec.* 900. *Syst.* 873.

VERBESINA. Ludw. 314.

1008. SILENE.

Lin. *Gen.* 503. *Spec.* 416. *Syst.* 503. Ludw. 573.

VISCAGO. Dill. Hall. 373. 375.

1009. SILPHIUM.

Linn. *Gen.* 882. *Spec.* 919. *Syst.* 882. Ludw. 362. Mill. iii. 265.

ASTERISCUS. Dill. Mill. iii. 30.

1010. SINAPIS.

Linn. *Gen.* 735. *Spec.* 668. *Syst.* 735.

SINAPI. Tourn. *tab.* 112. Blackw. *tab.* 29. Hall. 553. Ludw. 408.

Mill. ii. 262. iii. 266. Schæff. A. 150. Weinm. *tab.* 923.

1011. SIPHONANTHUS.

Linn. *Gen.* 120. *Spec.* 109. *Syst.* 120. Ludw. 1011.

SIPHONANTHEMUM. Annm.

1012. SISON.

Linn. *Gen.* 311. *Spec.* 252. *Syst.* 311. Ludw. 1034.

1013. SISYMEIUM.

Linn. *Gen.* 728. *Spec.* 657. *Syst.* 728. Hall. 547. Ludw. 410.

Mill. ii. 263. Tourn. *tab.* 109.

RADICULA. Dill.

1014. SISYRINCHIUM.

Linn. *Gen.* 908. *Spec.* 954. *Syst.* 908. Mill. ii. 264.

BERMUDIANA. Tourn. *tab.* 208. Ludw. 708. Mill. i. 119. iii. 38.

1015. SIUM.

Linn. *Gen.* 310. *Spec.* 251. *Syst.* 310.

SIUM. Tourn. *tab.* 162. Hall. 435. Ludw. 693. Mill. ii. 265.

SISARUM. Tourn. *tab.* 163. Mill. ii. 263. Weinm. *tab.* 924. d.

1016. SLOANEA.

Linn. *Gen.* 582. *Spec.* 512. *Syst.* 582. Ludw. 802.

SLOANA. Plum.

1017. SMILAX.

Linn. *Gen.* 992. *Spec.* 1028. *Syst.* 992. Blackw. *tab.* 393. Ludw.

917. Mill. ii. 265. Tourn. *tab.* 421.

1018. SMYRNIUM.

Linn. *Gen.* 325. *Spec.* 262. *Syst.* 325. Ludw. 689. Mill. ii. 267.

Tourn. *tab.* 168. Weinm. *tab.* 926.

1019. SOLANDRA.

Linn. *Syst. No.* 1170. *p.* 1380. 1269.

1020. SOLANUM.

Linn. *Gen.* 224. *Spec.* 184. *Syst.* 224. Ludw. 87.

SOLANUM. Tourn. *tab.* 62. Blackw. *tab.* 34. 107. Hall. 506.

Mill. ii. 268. Schæff. A. 28. Weinm. *tab.* 927, *seq.*

LYCOPERSICON. Tourn. *tab.* 63. Blackw. *tab.* 133. Mill. i. 547.
iii. 168.

MELONGENA. Tourn. *tab.* 65. Mill. ii. 40. iii. 192. Weinm. *tab.*
934.

1021. SOLDANELLA.

Linn. *Gen.* 182. *Spec.* 144. *Syst.* 182. Hall. 488. Ludw. 52.

Mill. ii. 272. Schæff. A. 45. Tourn. *tab.* 16. Weinm. *tab.* 420.

1022. SOLIDAGO.

Linn. *Gen.* 859. *Spec.* 878. *Syst.* 859. Hall. 729. Ludw. 353.

Mill. iii. 298.

JACOBÆÆ SPECIES. Tourn. Mill. i. 444.

VIRGA AUREA. Tourn. *tab.* 275. Blackw. *tab.* 169. Mill. ii. 398.
iii. 289.

DORIA. Dill. Mill. i. 275.

1023. SONCHUS.

Linn. *Gen.* 813. *Spec.* 793. *Syst.* 813. Blackw. *tab.* 30. 130. Hall.
752. Ludw. 338. Mill. ii. 272. Tourn. *tab.* 268. Weinm. *tab.*
938.

CREPIS. Vaill.

1024. SOPHORA.

Linn. *Gen.* 456. *Spec.* 373. *Syst.* 456. Ludw. 644.

1025. SORBUS.

Linn. *Gen.* 548. *Spec.* 477. *Syst.* 548. Blackw. *tab.* 173, 174.
Hall. 350. Ludw. 609. Mill. i. 281. Schæff. A. 197. Weinm.
tab. 941.

1026. SPARGANIUM.

Linn. *Gen.* 925. *Spec.* 971. *Syst.* 925. Hall. 259. Ludw. 872.
Tourn. *tab.* 302. Weinm. *tab.* 942.

1027. SPARTIUM.

Linn. *Gen.* 765. *Spec.* 708. *Syst.* 765.

GENISTA. Tourn. *tab.* 411. Blackw. *tab.* 244. Ludw. 634. Mill. i.
347. Weinm. *tab.* 532, *seq.*

1028. SPERGULA.

Linn. *Gen.* 519. *Spec.* 440. *Syst.* 519. Mill. ii. 285.

ALSINE. Hall. 387. Ludw. 569.

1029. SPERMACOCE.

Linn. *Gen.* 111. *Spec.* 102. *Syst.* 111. Ludw. 14.

1030. SPHERANTHUS.

Linn. *Gen.* 893. *Spec.* 927. *Syst.* 893. Ludw. 1030.

1031. SPHAGNUM.

Linn. *Gen.* 1051. *Spec.* 1106. *Syst.* 1051. Hall. 95. Ludw. 960.

1032. SPIGELIA.

Linn. *Gen.* 192. *Spec.* 149. *Syst.* 192. Ludw. 59.

ARAPABACA. Plum.*

1033. SPINACIA.

Linn. *Gen.* 986. *Spec.* 1027. *Syst.* 986. Blackw. *tab.* 49. Ludw. 929. Mill. ii. 286. Tourn. *tab.* 308. Weinm. *tab.* 946.

1034. SPIRÆA.

Linn. *Gen.* 554. *Spec.* 489. *Syst.* 554.

SPIRÆA. Tourn. *tab.* 389. Ludw. 616. Mill. ii. 288. iii. 270. Weinm. *tab.* 947.

FILIPENDULA. Tourn. *tab.* 150. Hall. 306. Ludw. 741. Mill. i. 324. Weinm. *tab.* 509. c.

ULMARIA. Tourn. *tab.* 141. Ludw. 620. Mill. ii. 462. Weinm. *tab.* 918.

ARUNCUS. Linn. *edit. prior.* BARBA CAPRÆ. Tourn. *tab.* 141. Ludw. 916. Weinm. *tab.* 229.

1035. SPLACHNUM.

Linn. *Gen.* 1054. *Spec.* 1108. *Syst.* 1054.

1036. SPONDIAS.

Linn. *Gen.* 453. *Spec.* 371. *Syst. No.* 453. p. 1036. 1382. Ludw. 552.

MONBIN. Plum. Mill. ii. 56.

1037. SPONGIA.

Linn. *Gen.* 1072. *Spec.* 1169. *Syst.* 1072. Ludw. 1002. Tourn. *tab.* 342.

BADIAGA. Buxb.

1038. STACHYS.

Linn. *Gen.* 638. *Spec.* 580. *Syst.* 638. Hall. 642. Ludw. 194.

STACHYS. Tourn. *tab.* 86. Mill. ii. 289. Weinm. *tab.* 948.

GALEOPSIS. Tourn. *tab.* 86. Blackw. *tab.* 84. Mill. i. 342. Weinm. *tab.* 529.

1039. STEHELINA.

Linn. *Gen.* 844. *Spec.* 840. *Syst.* 844. Ludw. 319.

1040. STAPELIA.

Linn. *Gen.* 271. *Spec.* 217. *Syst.* 271. Ludw. 100.

FRITILLARIACRASSA. Hortulanorum.

1041. STAPHYLEA.

Linn. *Gen.* 336. *Spec.* 270. *Syst.* 336.

STAPHYLODENDRON. Tourn. *tab.* 386. Hall. 423. Ludw. 531.

Mill. i. 290. Weinm. *tab.* 816. b.

1042. STATICE.

Linn. *Gen.* 348. *Spec.* 274. *Syst.* 348.

STATICE. Tourn. *tab.* 177. Ludw. 540. Mill. ii. 302. iii. 271.

LIMONIUM. Tourn. *tab.* 177. Ludw. 541. Mill. i. 518. Weinm. *tab.* 663.

1043. STELLARIA.

Linn. *Gen.* 504. *Spec.* 421. *Syst.* 504.

ALSINE. Tourn. *tab.* 126. Ludw. 569.

1044. STELLERA.

Linn. *Gen.* 439. *Spec.* 559. *Syst.* 439.

CHAMÆIASME. Amm.

1045. STEMODIA.

Linn. *Syst. No.* 1154. *p.* 1374. 1118.

STEMODIACRA. Brown.

1046. STERCULA.

Linn. *Gen.* 963. *Spec.* 1007. *Syst.* 963.

1047. STEWARTIA.

Linn. *Gen.* 758. *Spec.* 698. *Syst.* 758.

MELACHODENDRON. Mich.

1048. STIPA.

Linn. *Gen.* 84. *Spec.* 78. *Syst.* 84.

1049. STÆBE.

Linn. *Gen.* 839. *Spec.* 831. *Syst.* 839. Ludw. 304.

1050. STRATIOTES.

Linn. *Gen.* 607. *Spec.* 535. *Syst.* 607. Ludw. 388. Mill. iii. 272.
 ALOIDES. Boerh. Mill. i. 30. iii. 13.

1051. STRYCHNOS.

Linn. *Gen.* 226. *Spec.* 189. *Syst.* 226. Ludw. 91.
 NUX VOMICA. Offic. Blackw. *tab.* 395.

1052. STYRAX.

Linn. *Gen.* 527. *Spec.* 444. *Syst.* 527. Ludw. 159. Mill. ii. 305.
 Tourn. *tab.* 369. Weinm. *tab.* 953.

1053. SUBULARIA.

Linn. *Gen.* 716. *Spec.* 642. *Syst.* 716.

1054. SURIANA.

Linn. *Gen.* 353. *Spec.* 284. *Syst.* 353. Ludw. 542. Mill. ii. 307.

1055. SWERTIA.

Linn. *Gen.* 284. *Spec.* 226. *Syst.* 284.
 GENTIANA. Ludw. 97.

1056. SYMPHYTUM.

Linn. *Gen.* 170. *Spec.* 136. *Syst.* 170. Blackw. *tab.* 252. Hall.
 514. Ludw. 35. Mill. ii. 307. Schæff. A. 38. Tourn. *tab.* 56.
 Weinm. *tab.* 958.
 CONSOLIDA MAJOR. Off.

1057. SYRINGA.

Linn. *Gen.* 22. *Spec.* 9. *Syst.* 22. Mill. ii. 308. Weinm. *tab.* 958,
 959.
 LILAC. Tourn. *tab.* 372. Ludw. 1. Mill. i. 507.

1058. TABERNÆMONTANA.

Linn. *Gen.* 265. *Spec.* 210. *Syst.* 265. Ludw. 1019. Mill. ii. 309.

1059. TAGETES.

Linn. *Gen.* 865. *Spec.* 887. *Syst.* 865. Ludw. 351. Mill. ii. 310.
 Tourn. *tab.* 278. Weinm. *tab.* 960, *seq.*

1060. TAMARINDUS.

Linn. *Gen.* 46. *Spec.* 34. *Syst.* 46. Blackw. *tab.* 201. 221. Ludw. 373. Mill. ii. 310. Tourn. *tab.* 445. Weinm. *tab.* 964.
TAMARINTHUS. Mill. ii. 310.

1061. TAMARIX.

Linn. *Gen.* 337. *Spec.* 270. *Syst.* 337.

TAMARISCUS. Tourn. Blackw. *tab.* 331. Hall. 419. Ludw. 518.
Mill. ii. 311. Schæff. A. 175. Weinm. *tab.* 965.

1062. TAMUS.

Linn. *Gen.* 991. *Spec.* 1028. *Syst.* 991.

TAMNUS. Tourn. *tab.* 28. Hall. 165. Ludw. 907. Mill. ii. 311.

1063. TANACETUM.

Linn. *Gen.* 848. *Spec.* 843. *Syst.* 848. Hall. 693.

TANACETUM. Tourn. *tab.* 261. Mill. ii. 312. Schæff. A. 106.
Weinm. *tab.* 965.

BALSAMITA. Vaill. Blackw. *tab.* 98. Mill. i. 101. iii. 35. Schæff.
A. 107. Weinm. *tab.* 966.

1064. TARCHONANTHUS.

Linn. *Gen.* 846. *Spec.* 842. *Syst.* 846. Ludw. 320.

1065. TARGIONIA.

Linn. *Gen.* 1060. *Spec.* 1136. *Syst.* 1060. Ludw. 982.

1066. TANUS.

Linn. *Gen.* 1006. *Spec.* 1040. *Syst.* 1006. Hall. 146. Ludw. 935.
Mill. ii. 313. Tourn. *tab.* 362. Weinm. *tab.* 961. a.

1067. TELEPHIUM.

Linn. *Gen.* 339. *Spec.* 271. *Syst.* 339. Ludw. 517. Mill. ii. 316.
Schæff. A. 184. Tourn. *tab.* 128. Weinm. *tab.* 967, *seq.*

1068. TETRACERA.

Linn. *Gen.* 604. *Spec.* 533. *Syst.* 604. Ludw. 807.

1069. TETRAGONIA.

Linn. *Gen.* 551. *Spec.* 480. *Syst.* 551. Ludw. 806.

TETRAGONOCARPOS. Boerh. Mill. ii. 319.

1070. TETRAGONOTHECA.

Linn. *Gen.* 875. *Spec.* 903. *Syst.* 875. Ludw. 1029. Mill. iii. 274.

1071. TEUCRIUM.

Linn. *Gen.* 625. *Spec.* 562. *Syst.* 625. Ludw. 190.

TEUCRIUM. Tourn. *tab.* 90. Mill. ii. 319. Weinm. *tab.* 969.

POLIUM. Tourn. *tab.* 97. Mill. ii. 153. Weinm. *tab.* 822. e, f.

MARUM. Boerh. Blackw. *tab.* 47. Mill. ii. 12. Weinm. *tab.* 712.

CHAMÆDRYS. Tourn. *tab.* 97. Blackw. *tab.* 180. Hall. 630. Mill. i. 199. Schæff. A. 84. Weinm. *tab.* 361.

CHAMÆFITYS. Tourn. *tab.* 98. Mill. i. 202. Schæff. A. 85. Weinm. *tab.* 365.

SCORDIUM. Rai. Mill. ii. 245. Schæff. A. 82. SALVIA AGRESTIS. Blackw. *tab.* 9. Weinm. *tab.* 907. a.

IVA. Dill.

1072. THALIA.

Linn. *Gen.* 8. *Spec.* 1193. *Syst.* 8. Ludw. 1066.

CORTUSA. Plum.

1073. THALICTRUM.

Linn. *Gen.* 617. *Spec.* 545. *Syst.* 617. Hall. 307. Ludw. 465.

Mill. ii. 321. Tourn. *tab.* 270. Weinm. *tab.* 971.

1074. THAPSIA.

Linn. *Gen.* 323. *Spec.* 261. *Syst.* 323. Ludw. 659. Mill. ii. 322.

Tourn. *tab.* 171. Weinm. *tab.* 972. a, b.

1075. THEA.

Linn. *Gen.* 593. *Spec.* 515. *Syst.* 593. Blackw. *tab.* 351. Ludw.

735. Weinm. *tab.* 972. d.

1076. THELIGONUM.

Linn. *Gen.* 947. *Spec.* 993. *Syst.* 947.

CYNOCRAMBE. Tourn. *tab.* 485. Ludw. 887.

1077. THEOBROMA.

Linn. *Gen.* 806. *Spec.* 782. *Syst.* 806.

CACAO. Tourn. *tab.* 444. Blackw. *tab.* 378. Ludw. 604. Mill. i.
154. Weinm. *tab.* 277.

GUAZUMA. Plum. Ludw. 604. Mill. i. 385.

1078. THEOPHRASTA.

Linn. *Gen.* 190. *Spec.* 149. *Syst.* 190. Ludw. 53.

ERESIA. Plum.

1079. THESIUM.

Linn. *Gen.* 258. *Spec.* 207. *Syst.* 258. Hall. 183. Ludw. 771.

LINOPHYLLUM. Pont. LINOSYRIS. Rupp.

1080. THLASPI.

Linn. *Gen.* 719. *Spec.* 645. *Syst.* 719.

THLASPI. Tourn. *tab.* 101. *F, G, H, I, K.* Blackw. *tab.* 68. Ludw.
418. Mill. ii. 329. Schæff. A. 140. Weinm. *tab.* 973, 974.

BURSA PASTORIS. Tourn. *tab.* 103. Blackw. *tab.* 5. Ludw. 419.
Mill. i. 151. Schæff. A. 141. Weinm. *tab.* 274.

1081. THUJA.

Linn. *Gen.* 957. *Spec.* 1002. *Syst.* 957. Ludw. 876. Mill. ii. 332.
iii. 276. Tourn. *tab.* 358.

ARBOR VITÆ. Blackw. *tab.* 210.

1082. THYMERA.

Linn. *Gen.* 627. *Spec.* 569. *Syst.* 627. Mill. ii. 333.

1083. THYMUS.

Linn. *Gen.* 646. *Spec.* 590. *Syst.* 646. Hall. 654. Ludw. 210.

THYMUS. Tourn. *tab.* 93. Mill. ii. 336. Schæff. A. 74. Weinm.
tab. 975.

SERPILLUM. Tourn. *tab.* 93. Mill. ii. 255. Schæff. A. 75. Weinm.
tab. 916.

ACINOS. Dill. Mill. i. 17.

MASTICHINA. Boerh.

1084. TIARELLA.

Linn. *Gen.* 495. *Spec.* 405. *Syst.* 495.

1085. TILIA.

Linn. *Gen.* 587. *Spec.* 514. *Syst.* 587. Hall. 357. Ludw. 581.
Mill. ii. 336. Schæff. A. 191. Weinm. *tab.* 976.

1086. TILLÆA.

Linn. *Gen.* 163. *Spec.* 128. *Syst.* 163. Ludw. 378.

1087. TILLANDSIA.

Linn. *Gen.* 357. *Spec.* 286. *Syst.* 357. Ludw. 115.
CARAGUATA. Plum.

1088. TINUS.

Linn. *Syst.* No. 1133. p. 1367. 1010.
VOLKAMERIA. Brown.

1089. TOLUIFERA.

Linn. *Gen.* 470. *Spec.* 384. *Syst.* 470. Ludw. 652.

1090. TOMEX.

Linn. *Gen.* 140. *Spec.* 118. *Syst.* 140.

1091. TORDYLIUM.

Linn. *Gen.* 293. *Spec.* 293. *Syst.* 239. Hall. 448. Ludw. 663.
Mill. ii. 343. Tourn. *tab.* 170. Weinm. *tab.* 977.

1092. TORENIA.

Linn. *Gen.* 672. *Spec.* 619. *Syst.* 672.

1093. TORMENTILLA.

Linn. *Gen.* 560. *Spec.* 500. *Syst.* 560. Ludw. 464. Mill. ii. 344.
Schæff. A. 136. Tourn. *tab.* 153. Weinm. *tab.* 977. o.
POTENTILLA. Hall. 341.

1094. TOURNEFORTIA.

Linn. *Gen.* 176. *Spec.* 140. *Syst.* 176. Ludw. 80.
PITTONIA. Plum. Mill. ii. 146.

TABLE II.

1095. TOZZIA.

Linn. *Gen.* 663. *Spec.* 607. *Syst.* 663. Hall. 609. Ludw. 233.

1096. TRACHELIUM.

Linn. *Gen.* 204. *Spec.* 171. *Syst.* 204. Ludw. 64. Mill. ii. 345.
Tourn. *tab.* 50.

1097. TRADESCANTIA.

Linn. *Gen.* 360. *Spec.* 288. *Syst.* 360.
EPHEMERUM. Tourn. *tab.* 193. Ludw. 379. Mill. i. 279.

1098. TRAGIA.

Linn. *Gen.* 930. *Spec.* 980. *Syst.* 930. Ludw. 849. Mill. ii. 348.

1099. TRAGOPOGON.

Linn. *Gen.* 810. *Spec.* 789. *Syst.* 810. Hall. 758. Ludw. 333.
Mill. ii. 348. iii. 279. Weinm. *tab.* 978.

TRAGOPOGON. Tourn. *tab.* 270. BARBA HIRCI. Off. Schæff. A.
109.

TRAGOPOGONOIDES. Vaill.

1100. TRAPA.

Linn. *Gen.* 146. *Spec.* 120. *Syst.* 146. Ludw. 390.

TRIBULOIDES. Tourn. *tab.* 431. Hall. 468. TRIBULUS AQUATICUS. Rai.

1101. TREMELLA.

Linn. *Gen.* 1067. *Spec.* 1157. *Syst.* 1067.

1002. TREWIA.

Linn. *Gen.* 1104. *Spec.* 1193. *Syst.* 1104. Ludw. 803.

1103. TRIANTHEMA.

Linn. *Gen.* 273. *Spec.* 223. *Syst.* 273.

PORTULACASTRUM. Juss.

1104. TRIBULUS.

Linn. *Gen.* 476. *Spec.* 386. *Syst.* 476. Ludw. 557. Mill. ii. 350.
Tourn. *tab.* 141. Weinm. *tab.* 979. a.

1105. TRICHILIA.

Linn. *Syst. No.* 1134. *p.* 1368. 1020.

1106. TRICHOMANES.

Linn. *Gen.* 1045. *Spec.* 1097. *Syst.* 1045. Ludw. 949. Mill. ii.
350. Schæff. A. 308.

1107. TRICHOSANTHES.

Linn. *Gen.* 966. *Spec.* 1008. *Syst.* 966.

ANGUINA. Mich.

1108. TRICHOSTEMA.

Linn. *Gen.* 652. *Spec.* 598. *Syst.* 652. Ludw. 207.

1109. TRIDAX.

Linn. *Gen.* 872. *Spec.* 900. *Syst.* 872. Ludw. 1028.

1110. TRIENTALIS.

Linn. *Gen.* 419. *Spec.* 344. *Syst.* 419. Ludw. 130.

1111. TRIFOLIUM.

Linn. *Gen.* 802. *Spec.* 764. *Syst.* 802. Ludw. 473.

TRIFOLIUM. Tourn. *tab.* 228. Blackw. *tab.* 20. Hall. 580. Mill.
ii. 351. Schæff. A. 316. Weinm. *tab.* 979, *seq.* TRIPHYLLUM.
Sigb.

TRIFOLIASTRUM. Mich.

MELILOTUS. Tourn. *tab.* 229. Blackw. *tab.* 80. 284. Hall. 587.
Mill. ii. 30. Schæff. A. 167. Weinm. *tab.* 718.

LUPINASTER. Buxb.

TRIFOLIOIDES. Knaut. TRIPHYLLOIDES. Pont.

1112. TRIGLOCHIN.

Linn. *Gen.* 409. *Spec.* 333. *Syst.* 409. Hall. 253. Ludw. 383.

JUNCAGO. Tourn. *tab.* 142.

1113. TRIGONELLA.

Linn. *Gen.* 804. *Spec.* 776. *Syst.* 804. Ludw. 496.

FÆNUM GRÆCUM. Tourn. *tab.* 270. Mill. i. 327. Schæff. A. 140.
Weinm. *tab.* 514. a, b.

TABLE II.

1114. TRILLIUM.

Linn. *Gen.* 412. *Spec.* 339. *Syst.* 412.

1115. TRIOPTERIS.

Linn. *Gen.* 510. *Spec.* 428. *Syst.* 510. Ludw. 799.

1116. TRIOSTEUM.

Linn. *Gen.* 211. *Spec.* 176. *Syst.* 211.

TRIOSTEOSPERMUM. Dill. Ludw. 282. Mill. ii. 353.

1117. TRIPLARIS.

Linn. *Syst. No.* 1111. p. 1360. 881.

1118. TRIPSACUM.

Linn. *Syst. No.* 1167. p. 1379. 1261.

1119. TRITICUM.

Linn. *Gen.* 94. *Spec.* 85. *Syst.* 94. Blackw. *tab.* 40. Hall. 207.

Ludw. 827. Mill. ii. 354. Tourn. *tab.* 292, 293. Weinm. *tab.* 981.

1120. TRIUMFETTA.

Linn. *Gen.* 529. *Spec.* 444. *Syst.* 529. Ludw. 587. Mill. ii. 355.

1121. TROLLIUS.

Linn. *Gen.* 620. *Spec.* 556. *Syst.* 620.

HELLEBORO-RANUNCULUS. Boerh. Ludw. 758. Weinm. *tab.* 569. d. RANUNCULUS GLOBOSUS. Rai.

1122. TROPÆOLUM.

Linn. *Gen.* 421. *Spec.* 345. *Syst.* 421. Schæff. A. 217.

CORDAMINDUM. Tourn. *tab.* 244.

ACRIVIOLA. Boerh. Ludw. 631, Mill. i. 18. iii. 9. Weinm. *tab.* 753.

1123. TROPHIS.

Linn. *Syst. No.* 1173. p. 1381. 1289.

1024. TULIPA.

Linn. *Gen.* 376. *Spec.* 305. *Syst.* 376. Ludw. 717. Mill. ii. 355.

Tourn. *tab.* 199, 200. Weinm. *tab.* 982, *seq.*

1125. TURNERA.

Linn. *Gen.* 338. *Spec.* 271. *Syst.* 338. Ludw. 537. Mill. ii. 362.

1126. TURRITIS.

Linn. *Gen.* 733. *Spec.* 666. *Syst.* 733. Hall. 560. Ludw. 413.
Mill. ii. 362.

1127. TUSSILAGO.

Linn. *Gen.* 856. *Spec.* 865. *Syst.* 856. Ludw. 301.

TUSSILAGO. Tourn. *tab.* 276. Blackw. *tab.* 204. Mill. ii. 362.

Weinm. *tab.* 999. FARFARA. Schæff. A. 129.

PETASITES. Tourn. *tab.* 258. Blackw. *tab.* 222. Hall. 706. Mill.
ii. 126. Schæff. A. 130. Weinm. *tab.* 805. b.

1128. TYPHA.

Linn. *Gen.* 924. *Spec.* 971. *Syst.* 924. Hall. 260. Ludw. 873.
Tourn. *tab.* 301.

1129. VACCINIUM.

Linn. *Gen.* 434. *Spec.* 349. *Syst.* 434. Ludw. 134. Mill. iii. 284.
Weinm. *tab.* 1000.

VITIS IDÆA. Tourn. *tab.* 377. Hall. 413. Mill. ii. 461.

OXYCOCCUS. Tourn. *tab.* 431. Hall. 413.

1130. VALANTIA.

Linn. *Gen.* 1019. *Spec.* 1051. *Syst.* 1019.

VALANTIA. Tourn.

CRUCIATA. Tourn. *tab.* 39. Ludw. 11. Mill. i. 247. Weinm:
tab. 439. d, e.

1131. VALERIANA.

Linn. *Gen.* 43. *Spec.* 31. *Syst.* 43. Weinm. *tab.* 1001, 1002.

VALERIANA. Tourn. *tab.* 52. Blackw. *tab.* 250. 271. Hall. 662.
Ludw. 184. Mill. ii. 365. Schæff. A. 54.

VALERIANELLA. Tourn. *tab.* 52. Hall. 666. Ludw. 185. Mill. ii.
365.

1132. VALLISNERIA.

Linn. *Gen.* 975. *Spec.* 1015. *Syst.* 975. Ludw. 901.

VALLISNEROIDES. Mich.

TABLE II.

1133. VARRONIA.

Linn. *Syst.* 1118. *Spec.* 1363. 916.

1134. VATERIA.

Linn. *Gen.* 592. *Spec.* 515. *Syst.* 592. Ludw. 584.

1135. VELEZIA.

Linn. *Gen.* 403. *Spec.* 332. *Syst.* 403.

1136. VELLA.

Linn. *Gen.* 714. *Spec.* 641. *Syst.* 714. Ludw. 427. Mill. iii. 285.

1137. VERATRUM.

Linn. *Gen.* 1013. *Spec.* 1044. *Syst.* 1013. Hall. 298. Ludw. 731.
Mill. ii. 384. Tourn. *tab.* 145.

HELLEBORUS ALBUS. Rai. Blackw. *tab.* 74. Schæff. A. 253.
Weinm. *tab.* 568.

1138. VERBASCUM.

Linn. *Gen.* 217. *Spec.* 177. *Syst.* 217. Hall. 509. Ludw. 274.
VERBASCUM. Tourn. *tab.* 61. Blackw. *tab.* 3. Mill. ii. 385. Schæff.
A. 88. Weinm. *tab.* 1003.

BLATTARIA. Tourn. Mill. i. 131. iii. 40. Weinm. *tab.* 245. 249.

1139. VERBENA.

Linn. *Gen.* 30. *Spec.* 18. *Syst.* 30. Hall. 662. Ludw. 222.
VERBENA. Tourn. *tab.* 94. Blackw. *tab.* 41. Ludw. 222. Schæff.
A. 77. Weinm. *tab.* 1004.

SHERARDIA. Vaill.

BLAIRIA. Houst.

KEMPFERA. Houst.

1140. VERBESINA.

Linn. *Gen.* 874. *Spec.* 901. *Syst.* 874. Ludw. 314.
EUPATORIOPHALACRON. Dill. Mill. i. 303.
CERATOCEPHALOIDES. Vaill.

1141. VERONICA.

Linn. *Gen.* 25. *Spec.* 9. *Syst.* 25. Hall. 527. Ludw. 183.

VERONICA. Tourn. *tab.* 60. Blackw. *tab.* 134. Mill. ii. 387. Schæff. A. 58. Weinm. *tab.* 1004, *seq.* BARNAROTA. Mich.

BECCABUNGA. Tourn. Blackw. *tab.* 48. Mill. i. 117. Schæff. A. 59. Weinm. *tab.* 233.

1142. VIBURNUM.

Linn. *Gen.* 332. *Spec.* 267. *Syst.* 332.

VIBURNUM. Tourn. *tab.* 377. Hall. 467. Ludw. 110. Mill. ii. 391. iii. 287. Weinm. *tab.* 1007, 1008.

TINUS. Tourn. *tab.* 377. Ludw. 108. Mill. ii. 337.

OPULUS. Tourn. *tab.* 376. Hall. 463. Ludw. 109. Mill. ii. 89.

1143. VICIA.

Linn. *Gen.* 782. *Spec.* 734. *Syst.* 782. Ludw. 486.

VICIA. Tourn. *tab.* 221. Hall. 597. Mill. ii. 393. Weinm. *tab.* 1009.

FABA. Tourn. *tab.* 212. Blackw. *tab.* 19. Mill. i. *p.* 307. *p.* 101. Schæff. A. 168. Weinm. *tab.* 500.

1144. VINCA.

Linn. *Gen.* 261. *Spec.* 209. *Syst.* 261.

PERVINCA. Tourn. *tab.* 45. Blackw. *tab.* 59. Hall. 526. Ludw. 56. Mill. ii. 125. Schæff. A. 34. Weinm. *tab.* 1010.

1145. VIOLA.

Linn. *Gen.* 898. *Spec.* 933. *Syst.* 898. Blackw. *tab.* 44, 45. Hall. 500. Ludw. 629. Mill. ii. 396. iii. 288. Schæff. A. 215. Tourn. *tab.* 236. Weinm. *tab.* 1011.

1146. VISCUM.

Linn. *Gen.* 979. *Spec.* 1023. *Syst.* 979. Blackw. *tab.* 184. Hall. 162. Ludw. 877. Mill. ii. 400. Schæff. A. 270. Weinm. *tab.* 1013. b.

1147. VITEX.

Linn. *Gen.* 708. *Spec.* 938. *Syst.* 708. Black. *tab.* 139. Ludw. 262. Mill. ii. 401. Tourn. *tab.* 373.

AGNUS CASTUS. Off. Schæff. A. 90. Weinm. *tab.* 30.

1148. VITIS.

Linn. *Gen.* 250. *Spec.* 202. *Syst.* 250. Blackw. *tab.* 153. Hall. 166. Ludw. 525. Mill. ii. 401. iii. 290. Schæff. A. 174. Tourn. *tab.* 384. Weinm. *tab.* 1014, *seq.*

1149. ULEX.

Linn. *Gen.* 786. *Spec.* 741. *Syst.* 786. Ludw. 636. Mill. iii. 295. GENISTA-SPARTIUM. Tourn. *tab.* 412.

1150. ULMUS.

Linn. *Gen.* 281. *Spec.* 225. *Syst.* 281. Hall. 167. Ludw. 780. Mill. ii. 462. Tourn. *tab.* 372. Weinm. *tab.* 1018. b.

1151. ULVA.

Linn. *Gen.* 1069. *Spec.* 1163. *Syst.* 1069. Ludw. 869.

1152. UNIOLA.

Linn. *Gen.* 79. *Spec.* 71. *Syst.* 79. Ludw. 832.

1153. VOLKAMERIA.

Linn. *Gen.* 706. *Spec.* 637. *Syst.* 706. Ludw. 254. DOUGLASSIA. Heist. Mill. ii. 276.

1154. URENA.

Linn. *Gen.* 754. *Spec.* 692. *Syst.* 754. Ludw. 152. Mill. iii. 296.

1155. URTICA.

Linn. *Gen.* 935. *Spec.* 983. *Syst.* 935. Blackw. *tab.* 12. 321. Hall. 177. Ludw. 874. Mill. ii. 465. Schæff. A. 269. Weinm. *tab.* 1019, *seq.*

1156. UTRICULARIA.

Linn. *Gen.* 29. *Spec.* 18. *Syst.* 29. LATIBULARIA. Vaill. Hall. 612. Ludw. 181.

1157. UVARIA.

Linn. *Gen.* 612. *Spec.* 536. *Syst.* 612.

1158. UVULARIA.

Linn. *Gen.* 373. *Spec.* 304. *Syst.* 373. Ludw. 726. Schæff. A. 252.

1159. WACHENDORFIA.

Linn. *Syst.* 1108. *Spec.* 1359. 864.

1160. WALTHERIA.

Linn. *Gen.* 741. *Spec.* 673. *Syst.* 741. Ludw. 519. Mill. iii. 300.

MONOSPERMALTHEA. Isii.

1161. WEINMANNIANA.

Linn. *Syst. No.* 1131. p. 1367. 1005.

1162. WINTERANA.

Linn. *Gen. No.* 1140. *No.* 1370. 1045.

1163. XANTHIUM.

Linn. *Gen.* 937. *Spec.* 987. *Syst.* 937. Hall. 161. Ludw. 859.

Mill. ii. 525. Tourn. *tab.* 252. Weinm. *tab.* 1021.

1164. XERANTHEMUM.

Linn. *Gen.* 851. *Spec.* 857. *Syst.* 851. Hall. 709. Weinm. *tab.*
1021.

XERANTHEMUM. Tourn. *tab.* 284. Ludw. 326.

XERANTHEMOIDES. Dill.

1165. XIMENIA.

Linn. *Gen.* 1105. *Spec.* 1193. *Syst.* 1105. Ludw. 1067.

1166. XYLOPIA.

Linn. *Syst. No.* 1165. p. 1378. 1250.

1167. XYRRIS.

Linn. *Gen.* 59. *Spec.* 42. *Syst.* 59.

1168. YUCCA.

Linn. *Gen.* 388. *Spec.* 319. *Syst.* 388. Ludw. 117. Mill. ii. 531.

Weinm. *tab.* 1023.

CORDYLINE. Roy.

1169. ZANNICHELLIA.

Linn. *Gen.* 920. *Spec.* 969. *Syst.* 920.

ALGOIDES. Vaill.

APONOGETON. Pont.

GRAMINIFOLIA. Dill.

1170. ZANONIA.

Linn. *Gen.* 990. *Spec.* 1028. *Syst.* 990. Ludw. 905.

1171. ZANTHOXYLUM.

Linn. *Gen.* 335. *Spec.* 270. *Syst. No.* 335. p. 1290. Mill. iii. 309.

1172. ZEA.

Linn. *Gen.* 926. *Spec.* 971. *Syst.* 926.

MAYS. Tourn. *tab.* 303, 304, 305. Ludw. 870. Mill. ii. 22. iii. 182.

1173. ZINNIA.

Linn. *Syst. No.* 1161. p. 1377. 1221.

1174. ZIZANIA.

Linn. *Gen.* 942. *Spec.* 991. *Syst.* 942. Ludw. 899.

ELYMUS. Mich.

1175. ZIZIPHORA.

Linn. *Gen.* 33. *Spec.* 21. *Syst.* 33. Ludw. 180. Mill. iii. 311.

1176. ZOSTERA.

Linn. *Gen.* 919. *Spec.* 968. *Syst.* 919.

AIGA. Rai.

RUFFIA. Act. Angl.

1177. ZYGOPHYLLUM.

Linn. *Gen.* 474. *Spec.* 385. *Syst.* 474.

FABAGO. Tourn. *tab.* 135. Ludw. 558. Mill. i. 308.

TABLE III.

THE

LINNÆAN GENERA,

ALPHABETICALLY ARRANGED,

WITH

THE CLASSICAL AND ENGLISH NAMES;

AND ACCENTED,

With a Reference also to their Classes and Orders;

- Abróma, *Class xviii. Polyadelphia, Order i. Pentandria*
Abrus, *Class xvii. Diadelphia, Order iv. Decandria*
Ακαΐφφα, *Class xxi. Monœcia, Order ix. Monadelphia*
Acánthus (Bear's Breech), *Class xiv. Didynamia, Order ii. Angiosperma*
Acæna, *Class iv. Tetrandria, Order i. Monogynia*
Acer (Maple), *Class xxiii. Polygamia, Order i. Monœcia*
Achilléa (Málfoil), *Class xix. Syngenesia, Order ii. Polyg. sup.*
Achras (Sapota), *Class vi. Hexandria, Order i. Monogynia*
Achyránthes, *Class v. Pentandria, Order i. Monogynia*
Acnída, *Class xxii. Diœcia, Order v. Pentandria*
Aconítum (Wolfsbane), *Class xiii. Polyandria, Order iii. Trigynia*
Acorus (Sweet Rush), *Class vi. Hexandria, Order i. Monogynia*
Acróstichum (Forked Fern), *Class xxiv. Cryptogamia, Order i. Filices*
Actæa (Herb Christopher), *Class xviii. Polyandria, Order i. Monogynia*

- Adansónia (Æthiopian Sourgourd), *Class* xvi. Monadelphia, *Order* vii. Polyandria
- Adélia, *Class* xxii. Diœcia, *Order* xii. Monadelphia
- Adenanthéra (Bastard Flower-fence), *Class* x. Decandria, *Order* i. Monogynia
- Adiántum (Maiden Hair), *Class* xxiv. Cryptogamia, *Order* i. Filices
- Adónis (Bird's-eye), *Class* xiii. Polyandria, *Order* vii. Polygynia
- Adóxa (Tuberous Moscatel, or Hollow Root), *Class* viii. Octandria, *Order* iv. Tetragynia
- Ægilops, *Class* xxiii. Polygamia, *Order* i. Monœcia
- Ægiphila, *Class* iv. Tetrandria, *order* i. Monogynia
- Ægopódium (Herb Gerard, Gout-wort, or wild Angelica), *Class* v. Pentandria, *Order* ii. Digynia
- Ægópricon, *Class* xxi. Monœcia, *Order* i. Monandria
- Æschynómene (Bastard sensitive Plant), *Class* vii. Diadelphia, *Order* iv. Decandria
- Æsculus (Horse Chestnut), *Class* vii. Heptandria, *Order* i. Monogynia
- Æthúsa (Lesser Hemlock), or Fool's Parsley, *Class* v. Pentandria, *Order* ii. Digynia
- Agáricus, Agaric, *Class* xxiv. Cryptogamia, *Order* iv. Fungi
- Agáve (American Aloë), *Class* vi. Hexandria, *Order* i. Monogynia
- Agératum (Bastard Hemp Agrimony), *Class* xix. Syngenesia, *Order* i. Polyg. æqu.
- Agrimónia (Agrimony), *Class* xi. Dodecandria, *Order* ii. Digynia
- Agrostémma (Campion, or wild Lichnis), *Class* x. Decandria, *Order* v. Pentagynia
- Agróstis (Bent Grass), *Class* iii. Triandria, *Order* ii. Digynia
- Agynéja, *Class* xxi. Monœcia, *Order* ii. Gynandria
- Aira (Hair Grass), *Class* xi. Triandria, *Order* ii. Digynia
- Aitoma, *Class* xvi. Monadelphia, *Order* viii. Octandria
- Ajuga (Bugle), *Class* xiv. Didynamia, *Order* ii. Gymnosperma
- Aizoon, *Class* xii. Icosandria, *Order* v. Pentagynia
- Albúca, *Class* vi. Hexandria, *Order* i. Monogynia
- Alicea (Hollyhock, or Rose Mallow), *Class* xvi. Monadelphia, *Order* vii. Polyandria

- Alchemilla (Ladies' Mantle), *Class* iv. Tetrandria, *Order* i. Monogynia
- Aldrovanda, *Class* v. Pentandria, *Order* i. Monogynia
- Alétris (Bastard Aloë), *Class* vi. Hexandria, *Order* i. Monogynia
- Alisma (Water Plantain), *Class* vi. Hexandria, *Order* v. Polygynia
- Allamánda, *Class* v. Pentandria, *Order* i. Monogynia
- Alliónia, *Class* iv. Tetrandria, *Order* i. Monogynia
- Allium (Garlic), *Class* vi. Hexandria, *Order* i. Monogynia
- Allophýllus, *Class* viii. Octandria, *Order* i. Monogynia
- Aloë, *Class* vi. Hexandria, *Order* i. Monogynia
- Alopecúrus (Foxtail Grass), *Class* iii. Triandria, *Order* ii. Digynia
- Alpínia, *Class* i. Monandria, *Order* i. Monogynia
- Alsíne (Chickweed), *Class* v. Pentandria, *Order* i. Monogynia
- Alstónia, *Class* xiii. Polyandria, *Order* i. Monogynia
- Alstrœméria, *Class* vi. Hexandria, *Order* i. Monogynia
- Althæa (Marshmallow), *Class* xvi. Monadelphia, *Order* vii. Polyandria
- Alýssum (Madwort), *Class* xv. Tetradyndamia, *Order* ii. Silicul.
- Amaránthus (Amaranth, or Flower-gentle), *Class* xxi. Monœcia, *Order* v. Pentandria
- Amarýllis (Lily Daffodil), *Class* vi. Hexandria, *Order* i. Monogy.
- Ambrósia, *Class* xxi. Monœcia, *Order* v. Pentandria
- Ambrosína, *Class* xx. Gynandria, *Order* ix. Polyandria
- Aniéllus, *Class* xix. Syngenesia, *Order* ii. Polyg. super.
- Amethystéa, *Class* ii. Diandria, *Order* i. Monogynia
- Ammánia, *Class* iv. Tetrandria, *Order* i. Monogynia
- Ammi (Bishop's Weed), *Class* v. Pentandria, *Order* i. Monogynia
- Amúnum (Ginger), *Class* i. Monandria, *Order* i. Monogynia
- Amórpha (Bastard Indigo), *Class* xvii. Diadelphia, *Order* iv. Decandria
- Amýgdalus (Almond, or Peach), *Class* xii. Icosandria, *Order* i. Monogynia
- Amýris, *Class* viii. Octandria, *Order* i. Monogynia
- Anábasis (Berry-bearing Glasswort), *Class* v. Pentandria, *Order* ii. Digynia
- Anacárdium (Cashew Nut), *Class* ix. Enneandria, *Order* i. Monogynia

- Anacýlus*, *Class* xix. Syngenesia, *Order* ii. Polyg. super.
Anagáltis (Pimpernel), *Class* v. Pentandria, *Order* i. Monogynia
Anágyris (Stinking Bean Trefoil), *Class* x. Decandria, *Order* i.
 Monogynia
Anastática (Rose of Jericho), *Class* xv. Tetradyndamia, *Order* ii.
 Siliculosa
Anchúsa (Bugloss), *Class* v. Pentandria, *Order* i. Monogynia
Ancístum, *Class* ii. Diandria, *Order* i. Monogynia
Andráchne (Bastard Orpine), *Class* xxi. Monœcia, *Order* ii. Gy-
 nandria
Andrómeda (Marsh Cistus), *Class* x. Decandria, *Order* i. Mono-
 gynia
Andropógon, *Class* xxiii. Polygamia, *Order* i. Monœcia
Andrósace, *Class* v. Pentandria, *Order* i. Monogynia
Andrýala (Downy Sow-thistle), *Class* xix. Syngenesia, *Order* i.
 Polyg. æqu.
Anemóne (Wind Flower), *Class* xiii. Polyandria, *Order* vii. Poly-
 gynia
Anéthum (Dill), *Class* v. Pentandria, *Order* ii. Digynia
Angélica *Class* v. Pentandria, *Order* ii. Digynia
Angúria, *Class* xxi. Monœcia, *Order* ii. Diandria
Amóna (Custard Apple), *Class* xiii. Polyandria, *Order* vii. Po-
 lygynia
Anthemis (Chamomile), *Class* xix. Syngenesia, *Order* ii. Polyg.
 super.
Anthéricum (Spider-wort), *Class* vi. Hexandria, *Order* i. Mono-
 gynia
Anthistíria, *Class* iii. Triandria, *Order* ii. Digynia
Anthóceros, *Class* xxiv. Cryptogamia, *Order* iii. Algæ
Anthospérmum (Amber Tree), *Class* xxiii. Polygamia, *Order* ii.
 Dicœcia
Anthoxánthum (Vernal Grass), *Class* ii. Diandria, *Order* ii. Digynia
Antholiza, *Class* iii. Triandria, *Order* i. Monogynia
Anthýllis (Kidney Vetch, or Lady's Finger), *Class* xvii. Diadel-
 phia, *Order* iv. Decandria
Antichórus, *Class* viii. Octandria, *Order* i. Monogynia
Audidésma, *Class* xxii. Dicœcia, *Order* v. Pentandria

- Antirrhinum* (Snap-dragon, or Calf's-snout), *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
Apáctis, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*
Aphanes (Parsley-piert), *Class* iv. *Tetrandria*, *Order* ii. *Digynia*
Aphyllánthes, *Class* vi. *Hexandria*, *Order* i. *Monogynia*
Aphytéia, *Class* xvi. *Monadelphia*, *Order* i. *Triandria*
Apium (Parsley), *Class* v. *Pentandria*, *Order* ii. *Digynia*
Aplúda, *Class* xxiii. *Polygamia*, *Order* i. *Monœcia*.
Apócynum (Dog's-bane), *Class* v. *Pentandria*, *Order* ii. *Digynia*
Aponogéton, *Class* vii. *Heptandria*, *Order* iv. *Tetragynia*
Aquártia, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
Aquilégia (Columbine,) *Class* xiii. *Polyandria*, *Order* v. *Pentagyn.*
Aquilícia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Arábis (Bastard Tower Mustard), *Class* xv. *Tetradynamia*, *Order* i. *Siliquosa*
Aráchis (Ground Nut), *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
Arália (Berry-bearing Angelica), *Class* v. *Pentandria*, *Order* ii. *Digynia*
Arbutus (Strawberry-tree), *Class* x. *Decandria*, *Order* i. *Monogyn.*
Arctium (Burdock), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqu.*
Arctopus, *Class* xxiii. *Polygamia*, *Order* ii. *Diœcia*
Arctótis, *Class* xix. *Syngenesia*, *Order* iv. *Polyg. necessaria*
Arduína (Bastard Lycium), *Class* v. *Pentandria*, *Order* i. *Monogynia*
Aréca (Areca Nut), Appendix, *Palmæ*
Arenária (Sea Chickweed), *Class* x. *Decandria*, *Order* iii. *Trigynia*
Arethúsa, *Class* xx. *Gynandria*, *Order* i. *Diandria*
Arétia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Argenóné (Prickly Poppy), *Class* xiii. *Polyandria*, *Order* i. *Monogynia*
Argophýllum, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Aristída, *Class* iii. *Triandria*, *Order* ii. *Digynia*
Aristótelia, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*
Aristolóchia (Birthwort), *Class* xx. *Gynandria*, *Order* v. *Hexand.*
Arnica, *Class* xix. *Syngenesia*, *Order* ii. *Polyg. super.*
Artédia, *Class* v. *Pentandria*, *Order* ii. *Digynia*
Artemísia (Mugwort), *Class* xix. *Syngenesia*, *Order* ii. *Polyg. sup.*

- Artocárpus, *Class* xxi. Monœcia, *Order* i. Monandria
 Arum (Wake Robin, or Cuckoo Pint), *Class* xx. Gynandria, *Order* ix. Polyandria
 Arúnde (Reed), *Class* iii. Triandria, *Order* ii. Digynia
 Asárum (Asarabacca), *Class* xi. Dodecandria, *Order* i. Monogy.
 Asclépias (Swallow-wort), *Class* v. Pentandria, *Order* ii. Digynia
 Ascyrum (St. Peter's-wort), *Class* xviii. Polyadelphia, *Order* iii.
 Polyandria
 Aspálathus (Áfrican Broom), *Class* xvii. Diadelphia, *Order* iv.
 Decandria
 Aspáragus (Asparagus), or Sperge, *Class* v. Hexandria, *Order* i.
 Monogynia
 Asperúgo (Wild Buglos, or Goose Grass), *Class* v. Pentandria,
Order i. Monogynia
 Aspérula (Woodroof), *Class* iv. Tetrandia, *Order* i. Monogynia
 Asphódelus (Asphodel, or King's Spear), *Class* vi. Hexandria,
Order i. Monogynia
 Asplénium (Spleen-wort, or Milt-waste), *Class* xxiv. Cryptogamia,
Order i. Filices
 Aster (Star-wort), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
 Astrágalus (Liquorice Vetch, or Milk Vetch), *Class* xvii. Diadelphia,
Order iv. Decandria
 Astrántia (Black Masterwort), *Class* v. Pentandria, *Order* ii. Digynia
 Astrónium, *Class* xxii. Diœcia, *Order* v. Pentandria
 Athamánta (Spignel), *Class* v. Pentandria, *Order* ii. Digynia
 Athanásia, *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
 Atráctylis (Distaff Thistle), *Class* xix. Syngenesia, *Order* i. Polyg.
 æqua.
 Atrágene, *Class* xiii. Polyandria, *Order* vii. Polygynia
 Atrapháxis, *Class* vi. Hexandria, *Order* ii. Digynia
 Atriplex (Orach), *Class* xxiii. Polygamia, *Order* ii. Diœcia
 Atropa (Deadly Nightshade), *Class* v. Pentandria, *Order* i. Monogynia
 Aucúba, *Class* xxii. Monœcia, *Order* iv. Tetrandria
 Avéna (Oats), *Class* iii. Triandria, *Order* ii. Digynia
 Avérrhoa, *Class* x. Decandria, *Order* v. Pentagynia

- Avicénna, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
 Axýris, *Class* xxii. *Monœcia*, *Order* iii. *Triandria*
 Ayénia, *Class* xx. *Gynandria*, *Order* iv. *Pentandria*
 Azáleá (American upright Honeysuckle), *Class* v. *Pentandria*,
Order i. *Monogynia*

B

- Báccharis (Plowman's Spikenard), *Class* xix. *Syngenesia*, *Order*
 ii. *Polyg. super.*
 Bæckéa, *Class* viii. *Octandria*, *Order* i. *Monogynia*
 Ballóta (Black Horehound), *Class* xiv. *Didynamia*, *Order* i. *Gym-*
nosperma
 Baltimóra, *Class* xix. *Syngenesia*, *Order* iv. *Polyg. necess.*
 Banistéria, *Class* x. *Decandria*, *Order* iii. *Trigynia*
 Bánksia, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
 Barléria *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
 Barnadésia, *Class* xix. *Syngenesia*, *Order* i. *Pol. æqualis*
 Barringtónia, *Class* xvi. *Monadelphia*, *Order* vii. *Polyandria*
 Bártsia, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
 Basélla (Malabar Nightshade), *Class* v. *Pentandria*, *Order* iii.
Trigynia
 Bássia, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*
 Bátis, *Class* xxii. *Dicecia*, *Order* iv. *Tetrandria*
 Bauhínia (Mountain Ebony), *Class* x. *Decandria*, *Order* i. *Mono-*
gynia
 Befária, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*
 Begónia, *Class* xxi. *Monœcia*, *Order* vii. *Polyandria*
 Béllium, *Class* xix. *Syngenesia*, *Order* ii. *Polygamia super.*
 Béllis (Daisy), *Class* xix. *Syngenesia*, *Order* ii. *Polygamia super.*
 Bellónia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Bérberis (Berberry, or Piperidge Bush), *Class* vi. *Hexandria*,
Order i. *Monogynia*
 Bérgia, *Class* x. *Decandria*, *Order* v. *Pentagynia*
 Besléria, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
 Béta (Beet), *Class* v. *Pentandria*, *Order* ii. *Digynia*
 Betónica (Betony), *Class* xiv. *Didynamia*, *Order* i. *Gymnosper.*

- Bétula (Birch), *Class* xxi. Monœcia, *Order* iii. Triandria
- Bídens (Water Hemp Agrim.), *Class* xix. Syngenesia, *Order* i.
Polyg. æqual.
- Bignónia (Trumpet Flower), *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Biscutélla (Buckler Mustard), *Class* xv. Tetradynamia, *Order* i.
Siliculosa
- Bisérula, *Class* xvii. Diadelphia, *Order* iv. Decandria
- Bíxa (Anotta), *Class* xiii. Polyandria, *Order* i. Monogynia
- Bládhia, *Class* v. Pentandria, *Order* i. Monogynia
- Blakéa, *Class* xi. Dodecandria, *Order* i. Monogynia
- Blásia, *Class* xxiv. Cryptogamia, *Order* iii. Algæ
- Blæria, *Class* iv. Tetrandria, *Order* i. Monogynia
- Bléchnum, *Class* xxiv. Cryptogamia, *Order* i. Filices
- Blítum (Strawberry Spinach, or Blite), *Class* i. Monandria, *Order* ii. Digynia
- Bobártia, *Class* iii. Triandria, *Order* ii. Digynia
- Boccónia, *Class* xi. Dodecandria, *Order* i. Monogynia
- Boerháavia (American Hog-weed), *Class* i. Monandria, *Order* i.
Monogynia
- Bolétus, *Class* xxiv. Cryptogamia, *Order* iv. Fungi
- Bómbax (Silk Cotton Tree), *Class* xvi. Monadelphia, *Order* vii.
Polyandria
- Bóntia, *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Borássus, Appendix, Palma
- Borbónia, *Class* xvii. Diadelphia, *Order* iv. Decandria
- Borágo (Borrage), *Class* v. Pentandria, *Order* i. Monogynia
- Boséa (Yerva-mora, or Golden-rod Tree), *Class* v. Pentandria,
Order ii. Digynia
- Brabéium (African Almond), *Class* xxiii. Polygamia, *Order* i.
Monœcia
- Brássica (Cabbage), *Class* xv. Tetradynamia, *Order* i. Siliquosa
- Bráthys, *Class* xiii. Polyandria, *Order* v. Pentagynia
- Bríza (Quaking Grass), *Class* iii. Triandria, *Order* ii. Digynia
- Bromélia (Ananas, or Pine Apple), *Class* vi. Hexandria, *Order* i:
Monogynia

- Brómus (Brome Grass), *Class* iii. Triandria, *Order* ii. Digynia
 Brossæa, Appendix, Palmæ
 Browállia, *Class* xiv. Didynamia, *Order* ii. Angiosperma
 Brownéa, *Class* xvi. Monadelphia, *Order* v. Euneandria
 Brunélla,
 Brúnia, *Class* v. Pentandria, *Order* i. Monogynia
 Brunsfélsia, *Class* v. Pentandria, *Order* i. Monogynia
 Brucea, *Class* xxii. Diœcia, *Order* iv. Tetrandria
 Bryónia (Bryony), *Class* xxi. Monœcia, *Order* x. Syngenesia
 Bryúm, *Class* xxiv. Cryptogamia, *Order* ii. Musci
 Búbon (Macedonian Parsley), *Class* v. Pentandria, *Order* ii. Digynia
 Bucída, *Class* x. Decandria, *Order* i. Monogynia
 Buchnéra, *Class* xiv. Didynamia, *Order* ii. Angiosperma
 Buddléia, *Class* xv. Tetrandria, *Order* i. Monogynia
 Bufónia, *Class* iv. Tetrandria, *Order* ii. Digynia
 Bulbocódium, *Class* vi. Hexandria, *Order* i. Monogynia
 Bumálda, *Class* v. Pentandria, *Order* ii. Digynia
 Búnias, *Class* iv. Tetrodynamia, *Order* i. Siliquosa
 Búnium (Pig-nut, or Earth-nut), *Class* v. Pentandria, *Order* ii. Digynia
 Buphthálmuni (Ox-eye), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
 Bupleúrum (Hare's Ear), *Class* v. Pentandria, *Order* ii. Digynia
 Burmánnia, *Class* vi. Hexandria, *Order* i. Monogynia
 Burséra, *Class* vi. Hexandria, *Order* i. Monogynia
 Bútonus (Flowering Rush, or Water Gladiolus), *Class* ix. Enneandria, *Order* vi. Hexagynia
 Buxbáumia, *Class* xxiv. Cryptogamia, *Order* ii. Musci
 Búxus (Box Tree), *Class* xxi. Monœcia, *Order* iv. Tetrandria
 Býssus, *Class* xxiv. Cryptogamia, *Order* iii. Algæ
 Buttnéria, *Class* v. Pentandria, *Order* i. Monogynia

C

- Cacália (Alpine Colt's-foot), *Class* xix. Syngenesia, *Order* i. Polyg. æqu.

- Cactus (Melon Thistle), *Class* xii. Icosandria, *Order* i. Monogynia
- Cáchrys, *Class* v. Pentandria, *Order* ii. Digynia
- Cæsalpínia (Brasiletto), *Class* x. Decandria, *Order* i. Monogynia
- Caléa, *Class* xix. Syngenesia, *Order* i. Polyg. æqual.
- Caléndula (Marygold), *Class* xix. Syngenesia, *Order* iv. Polyg. neces.
- Cálamus, *Class* vi. Hexandria, *Order* i. Monogynia
- Calceolária, *Class* ii. Diandria, *Order* i. Monogynia
- Calycánthus (Virginian All-spice), *Class* xii. Icosandria, *Order* v. Polygynia
- Cála (African Arum), *Class* xx. Gynandria, *Order* ix. Polyandria
- Callicárpa (Johnsonia), *Class* iv. Tetrandria, *Order* i. Monogynia
- Calligonum, *Class* xiii. Polyandria, *Order* ii. Bigynia
- Callisia, *Class* iv. Triandria, *Order* i. Monogynia
- Callitriche (Star-headed Water Chickweed), *Class* i. Monandria, *Order* ii. Digynia
- Calódendrum, *Class* v. Pentandria, *Order* i. Monogynia
- Calophýllum, *Class* xiii. Polyandria, *Order* i. Monogynia
- Cáltha (Marsh Marygold), *Class* xiii. Polyandria, *Order* vii. Polygynia
- Cambógia, *Class* xiii. Polyandria, *Order* i. Monogynia
- Caméllia, *Class* xvi. Monadelphía, *Order* vii. Polyandria
- Camerária, *Class* v. Pentandria, *Order* i. Monogynia
- Campánula (Bell-flower), *Class* v. Pentandria, *Order* i. Monogynia
- Camocládia, *Class* iv. Triandria, *Order* i. Monogynia
- Camphorósma, *Class* iv. Tetrandria, *Order* i. Monogynia
- Canarína, *Class* vi. Hexandria, *Order* i. Monogynia
- Canárium, *Class* xxii. Dicœcia, *Order* v. Pentandria
- Canélla, *Class* xi. Dodecandria, *Order* i. Monogynia
- Cámma (Indian Flowering Reed), *Class* i. Monandria, *Order* i. Monogynia
- Cánnabis (Hemp), *Class* xxii. Dicœcia, *Order* v. Pentandria
- Cápparis (Caper Bush), *Class* xiii. Polyandria, *Order* i. Monogynia

- Caprária, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
- Cápsicum (Guinea Pepper), *Class* v. *Pentandria*, *Order* i. *Monogy.*
- Capúra, *Class* vi. *Hexandria*, *Order* i. *Monogynia*
- Cardamíne (Lady's Smock), *Class* xv. *Tetradynamia*, *Order* i. *Siliquosa*
- Cardiospérmum (Heart Pea), *Class* viii. *Octandria*, *Order* iii. *Trigynia*
- Cárduus (Thistle), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqual.*
- Cárex, *Class* xxi. *Monœcia*, *Order* iii. *Triandria*
- Caríca (Papaw), *Class* xxii. *Dioœcia*, *Order* ix. *Decandria*
- Carissa, *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Carlína (Carline Thistle), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
- Carolinéa, *Class* xvi. *Monadelphia*, *Order* vii. *Polyandria*
- Caróxylon, *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Carpésium, *Class* xix. *Syngenesia*, *Order* ii. *Polyg. super.*
- Cárpinus (Hornbeam), *Class* xxi. *Monœcia*, *Order* 8. *Polyandria*
- Cárthamus (Bastard Saffron), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
- Cárum (Carui, or Carraway), *Class* v. *Pentandria*, *Order* ii. *Digynia*
- Carýocar, *Class* xiii. *Polyandria*, *Order* iv. *Tetragynia*
- Caryophýllus (Clove Tree), *Class* xiii. *Polyandria*. *Order* i. *Monogynia*
- Carýota, *Appendix*, *Palmæ*
- Cassia (Wild Senna), *Class* x. *Decandria*, *Order* i. *Monogynia*
- Cássine (Hottentot Cherry), *Class* v. *Pentandria*, *Order* iii. *Trigynia*
- Cassýt, *Class* ix. *Enneandria*, *Order* i. *Monogynia*
- Castilléia, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
- Casuarína, *Class* xxi. *Monœcia*, *Order* i. *Monandria*
- Catananche (Candy Lion's Foot), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
- Catesbæa (Lily Thorn), *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
- Cáturus, *Class* xxii. *Dioœcia*, *Order* v. *Pentandria*
- Caúcalis (Bastard Parsley), *Class* v. *Pentandria*, *Order* ii. *Digyn.*

- Ceanóthus (New Jersey Tea), *Class v. Pentandria, Order i. Monogynia*
- Cecrópia, *Class xxii. Diœcia, Order ii. Diandria*
- Cedréla, *Class v. Pentandria, Order i. Monogynia*
- Celéstrus (Staif Tree), *Class v. Pentandria, Order i. Monogynia*
- Celósia (Cock's-comb), *Class v. Pentandria, Order i. Monogynia*
- Célsia, *Class xiv. Didynamia, Order ii. Angiosperma*
- Céltis (Nettle Tree), *Class xxiii. Polygamia, Order i. Monœcia*
- Cénchrus, *Class xxiii. Polygamia, Order i. Monœcia*
- Centaureá (Centaury), *Class xix. Syngenesia, Order iii. Polyg. frustr.*
- Centélla, *Class xxi. Monœcia, Order iv. Tetrandria*
- Centúnculus, *Class iv. Tetrandria, Order i. Monogynia*
- Cephaláanthus (Button Wood), *Class iv. Tetrandria, Order i. Monogynia*
- Cerástium (Mouse-ear Chickweed), *Class x. Decandria, Order iv. Pentagynia*
- Ceratocárpus, *Class xxi. Monœcia, Order i. Monandria*
- Cerafónia (Carob Tree, or St. John's Bread), *Class xxiii. Polygamia, Polyœcia*
- Ceratophýllum (Horned Pond Weed), *Class xxi. Monœcia, Order viii. Polyandria*
- Cérbera, *Class v. Pentandria, Order i. Monogynia*
- Cércis (Judas Tree), *Class x. Decandria, Order i. Monogynia*
- Cerínthe (Honey-wort), *Class v. Pentandria, Order i. Monogyn.*
- Ceropégia, *Class v. Pentandria, Order i. Monogynia*
- Cestrum (Bastard Jasmine), *Class v. Pentandria, Order i. Monogyn.*
- Chærophýllum (Wild Chervil), *Class v. Pentandria, Order ii. Digynia*
- Chalcás, *Class x. Decandria, Order i. Monogynia*
- Chamærops (Dwarf-palm, or Palmeto), *Appendix, Palmæ*
- Chamira, *Class xv. Tetradynamia, Order i. Siliquosa*
- Chára, *Class xxi. Monœcia, Order i. Monandria*
- Cheiránthus (Stock July Flower), *Class xv. Tetradynamia, Order i. Siliquosa*
- Chelidónium (Celendine), *Class xiii. Polyandria, Order i. Monogynia*

- Chelóne, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
- Chenólea, *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Chenopódium (Goose-foot, or Wild Orach), *Class* v. *Pentandria*,
Order ii. *Digynia*
- Cherléria, *Class* x. *Decandria*, *Order* iii. *Tiigynia*
- Chiocócca, *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Chionánthus (Snow-drop Tree, or Fringe Tree), *Class* ii. *Diandria*,
Order i. *Monogynia*
- Chirónia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Chlóra, *Class* viii. *Octandria*, *Order* i. *Monogynia*
- Chondrilla (Gum Succory), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
- Chrysánthemum (Corn Marygold), *Class* xix. *Syngenesia*, *Order* ii. *Polyg. super.*
- Chrýsitrix, *Class* xxiii. *Polygamia*, *Order* ii. *Dicœcia*
- Chrysobálanus (Cocoa Plumb), *Class* xii. *Icosandria*, *Order* i. *Monogynia*
- Chrysócoma (Golden Locks), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
- Chrysógonum, *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
- Chrysophýllum (Star Apple), *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Chrysosplénium (Golden Saxifrage), *Class* x. *Decandria*, *Order* ii. *Digynia*
- Cicca, *Class* xxi. *Monœcia*, *Order* iv. *Tetrandria*
- Cícer (Chich Peas), *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
- Chichórium (Succory, or Endiye), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
- Cicúta (Water Hemlock), *Class* v. *Pentandria*, *Order* ii. *Digynia*
- Cimicífuga, *Class* xiii. *Polyandria*, *Order* iv. *Tetragynia*
- Chinchóna, *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Cinna, *Class* i. *Monandria*, *Order* ii. *Digynia*
- Cinerária (Sky-flower), *Class* xix. *Synge*, *Order* ii. *Polyg. sup.*
- Circæa (Enchanter's Nightshade), *Class* ii. *Diandria*, *Order* i. *Monogynia*
- Cissámpelos, *Class* xxii. *Dicœcia*, *Order* xii. *Monadelpkia*

- Cissus*, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
Cistus (Rock Rose), *Class* xiii. *Polyandria*, *Order* i. *Monogynia*
Citharóxylon (Fiddle Wood), *Class* xiv. *Didynamia*, *Order* ii.
 Angiosperma
Citrus (Citron), *Class* xviii. *Polyadelphia*, *Order* ii. *Icosandria*
Cláthrus, *Class* xxiv. *Cryptogamia*, *Order* iv. *Fungi*
Clavária, *Class* xxiv. *Cryptogamia*, *Order* iv. *Fungi*
Claytónia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Clématis (Virgin's Bower), *Class* xiii. *Polyandria*, *Order* vii. *Polygynia*
Cléome (Bastard Mustard), *Class* xv. *Tetradynamia*, *Order* i.
 Siliquosa
Cleónia, *Class* xiv. *Didynamia*, *Order* i. *Gymnosperma*
Clerodéndrum, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
Clibádium, *Class* xxi. *Monœcia*, *Order* v. *Pentandria*
Cléthra, *Class* x. *Decandria*, *Order* i. *Monogynia*
Cleyera, *Class* xiii. *Polyandria*, *Order* i. *Monogynia*
Cliffortia, *Class* xxii. *Diœcia*, *Order* xi. *Polyandria*
Clinopódium (Field Basil), *Class* xiv. *Didynamia*, *Order* i. *Gymnosperma*
Clitória, *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
Clúsia (Balsam Tree), *Class* xxiii. *Polygamia*, *Order* i. *Monœcia*
Clútia, *Class* xxii. *Diœcia*, *Order* xiv. *Gynandria*
Clypéola (Treacle Mustard), *Class* xv. *Tetradynamia*, *Order* ii.
 Siliculosa
Cneórum (Widow Wail), *Class* iii. *Triandria*, *Order* i. *Monogynia*
Cnicus (Blessed Thistle), *Class* x. *Syngenesia*, *Order* i. *Polygæqualis*
Cochleária (Scurvy-grass, or Spoon-wort), *Class* xv. *Tetradynamia*, *Order* ii. *Siliculosa*
Cócos (Cocoa-Nut), *Palmæ*
Códia, *Class* viii. *Octandria*, *Order* ii. *Digynia*
Coccóloba, *Class* viii. *Octandria*, *Order* iii. *Trigynia*
Códon, *Class* x. *Decandria*, *Order* i. *Monogynia*
Cofféa (Coffee-Tree), *Class* v. *Pentandria*, *Order* i. *Monogynia*
Coix (Job's Tears), *Class* xxi. *Monœcia*, *Order* iii. *Triandria*

- Cólchicum* (Meadow Saffron), *Class vi. Hexandria, Order iii.*
Trigynia
- Coldénia*, *Class iv. Tetrandria, Order iii. Tetragynia*
- Collinsonia*, *Class ii. Diandria, Order i. Monogynia*
- Columnéa*, *Class xiv. Didynamia, Order ii. Angiosperma*
- Colútea* (Bladder Senna), *Class xvii. Diadelphia, Order iv. Decandria*
- Cómarum* (Marsh Cinquefoil), *Class xii. Icosandria, Order v. Polygynia*
- Combrétum*, *Class viii. Octandria, Order i. Monogynia*
- Cométes*, *Class iv. Tetrandria, Order i. Monogynia*
- Commelína*, *Class iii. Triandria, Order i. Monogynia*
- Commersónia*, *Class v. Pentandria, Order v. Pentagynia*
- Comocládia*, *Class iii. Triandria, Order i. Monogynia*
- Conférva*, *Class xxiv. Cryptogamia, Order iii. Algæ*
- Coníum* (Hemlock), *Class v. Pentandria, Order ii. Digynia*
- Cónnarus*, *Class xvi. Monadelphia, Order iv. Decandria*
- Conocárpus* (Button-Tree), *Class v. Pentandria, Order i. Monogynia*
- Convallária* (Lily of the Valley), *Class vi. Hexandria, Order i. Monogynia*
- Convólulus* (Bind Weed), *Class v. Pentandria, Order i. Monogynia*
- Conýza* (Flea-bane), *Class xix. Syngenesia, Order iii. Polyg. frustr.*
- Copáifera*, *Class x. Decandria, Order i. Monogynia*
- Coprósmia*, *Class v. Pentandria, Order ii. Digynia*
- Córchorus* (Jew's Mallow), *Class xiii. Polyandria, Order i. Monogynia*
- Córdia* (Sibestan), *Class v. Pentandria, Order i. Monogynia*
- Coreópsis* (Tick-seeded Sun-flower), *Class xix. Syngenesia, Order iii. Polyg. frustr.*
- Coriándrum* (Coriander), *Class v. Pentandria, Order ii. Digynia*
- Coriária* (Myrtle-leaved Sumach), *Class xxii. Diœcia, Order ix. Decandria*
- Córis* (Heathlow Pine), *Class v. Pentandria, Order i. Monogynia*

- Corispérmum (Tick-seed), *Class i. Monandria, Order ii. Digynia*
 Cornucópiæ, *Class iii. Triandria, Order ii. Digynia*
 Córnuſ (Dog-wood, or Cornelian Cherry), *Class iv. Tetrandria,*
Order i. Monogynia
 Cornútia, *Class xiv. Didynamia, Order ii. Angiosperma*
 Coronílla (Jointed-podded Coluta), *Class xvii. Diadelphia, Order*
iv. Decandria
 Corrigiôla, *Class v. Pentandria, Order iii. Trigynia*
 Cortúſa (Bear's-ear Sanicle), *Class v. Pentandria, Order i. Mo-*
nogynia
 Córylus (Hazel, or Nut-tree), *Class xxi. Monœcia, Order viii.*
Polyandria
 Corýmbium, *Class xix. Syngenesia, Order vi. Monogynia*
 Corynocárpus, *Class v. Pentandria, Order i. Monogamia*
 Corýpha, Palmæ
 Cóstus, *Class i. Monandria, Order i. Monogynia*
 Cótula, *Class xix. Syngenesia, Order ii. Polyg. superfl.*
 Cotylédon (Navel-wort), *Class x. Decandria, Order iv. Penta-*
gynia
 Crámbe (Sea Cabbage), *Class xv. Tetradynamia, Order i. Sili-*
quosa
 Craméria, *Class iv. Tetrandria, Order i. Monogynia*
 Craneolária, *Class xiv. Didynamia, Order ii. Angiosperma*
 Crássula (Lesser Orpine), *Class v. Pentandria, Order v. Pentagyn.*
 Cratægus (Wild Service), *Class xii. Icosandria, Order ii. Di-*
gynia
 Cratæva (Garlic Pear), *Class xi. Dodecandria, Order i. Monogyn.*
 Crépis (Bastard Hawk-weed), *Class xix. Syngenesia, Order i. Po-*
lyg. æqualis
 Crescéntia (Calabash Tree), *Class xiv. Didynamia, Order ii. An-*
giosperma
 Créſſa, *Class v. Pentandria, Order ii. Digynia*
 Crínium (Asphodel Lily), *Class vi. Hexandria, Order i. Monogyn.*
 Críthmum (Samphire), *Class v. Pentandria, Order ii. Digynia*
 Crócus (Saffron), *Class iii. Triandria, Order i. Monogynia*
 Crotalária, *Class xvii. Diadelphia, Order iv. Decandria*

- Cróton* (Tallow-Tree, or Bastard Ricinus), *Class* xxi. Monœcia,
Order ix. Monadelphia
- Crucianélla* (Petty Madder), *Class* iv. Tetrandria, *Order* i. Monogynia
- Cruzíta*, *Class* iv. Tetrandria, *Order* ii. Digynia
- Cucúbalus* (Berry-bearing Chickweed), *Class* x. Decandria, *Order* iii. Trigynia
- Cúcumis* (Cucumber), *Class* xxi. Monœcia, *Order* x. Syngenesia
- Cucúrbita* (Gourd), *Class* xxi. Monœcia, *Order* x. Syngenesia
- Cumínium* (Cumin), *Class* v. Pentandria, *Order* ii. Digynia
- Cuníla*, *Class* ii. Diandria, *Order* i. Monogynia
- Cunónia*, *Class* x. Decandria, *Order* ii. Digynia
- Cupánia*, *Class* xxi. Monœcia, *Order* ix. Monadelphia
- Cupréssus* (Cypress), *Class* xxi. Monœcia, *Order* ix. Monadelph.
- Curatélla*, *Class* xiii. Polyandria, *Order* ii. Digynia
- Curcúma* (Turmerick), *Class* i. Monandria, *Order* i. Monogynia
- Cuscúta* (Dodder), *Class* iv. Tetrandria, *Order* ii. Digynia
- Cussónia*, *Class* v. Pentandria, *Order* ii. Digynia
- Cyanélla*, *Class* vi. Hexandria, *Order* i. Monogynia
- Cýcas* (Sego Palm), *Class* xxiv. Cryptogamia, *Order* i. Filices
- Cýclamen* (Sow-bread), *Class* v. Pentandria, *Order* i. Monogyn.
- Cymbária*, *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Cynánchum*, *Class* v. Pentandria, *Order* ii. Digynia
- Cýnara* (Artichoke), *Class* xix. Syngenesia, *Order* i. Polyg. æqu.
- Cynoglóssum* (Hound's Tongue), *Class* v. Pentandria, *Order* i. Monogynia
- Cynométra*, *Class* x. Decandria, *Order* i. Monogynia
- Cynomórium*, *Class* xxi. Monœcia, *Order* i. Monandria
- Cynosúrus* (Dog's-tail Grass), *Class* v. Triandria, *Order* ii. Digynia
- Cyperus* (English Galingale), *Class* iii. Triandria, *Order* i. Monogynia
- Cypripédium* (Lady's Slipper), *Class* xx. Gynandria, *Order* ii. Diandria
- Cyrilla*, *Class* v. Pentandria, *Order* i. Monogynia
- Cýtinus*, *Class* xx. Gynandria, *Order* viii. Dodecandria

Cýtísus (Base-Tree Trefoil), *Class xvii. Diadelphia, Order iv. Decandria*

D

Dáctylis (Cock's-foot Grass), *Class iii. Triandria, Order ii. Digynia*

Dáis, *Class x. Decandria, Order i. Monogynia*

Dalbérgia, *Class xvii. Diadelphia, Order iii. Octandria*

Dalechâmpia, *Class xxi. Monœcia, Order ix. Monadelphia*

Dáphne (Mézereon, or Spurge Laurel), *Class viii. Octandria, Order i. Monogynia*

Dátisca (Bastard Hemp), *Class xxii. Dioœcia, Order x. Dodecand.*

Dátúra (Thorn Apple), *Class v. Pentandria, Order i. Monogynia*

Dáucus (Carrot), *Class v. Pentandria, Order ii. Digynia*

Decumária, *Class xi. Dodecandria, Order i. Monogynia*

Delína, *Class xiii. Polyandria, Order i. Monogynia*

Delphínium (Larkspur), *Class xiii. Polyandria, Order iii. Trigyn.*

Dentária (Tooth-wort), *Class xv. Tetradymania, Order i. Siliquosa*

Deutzia, *Class x. Decandria, Order iii. Trigynia*

Diálium, *Class ii. Diandria, Order i. Monogynia*

Dianthéra, *Class ii. Diandria, Order i. Monogynia*

Diánthus (Pink, or Carnation), *Class x. Decandria, Order ii. Digynia*

Diapénsia, *Class v. Pentandria, Order i. Monogynia*

Dictámnus (Fraxinella, or White Dittany), *Class x. Decandria, Order i. Monogynia*

Didelta, *Class xix. Syngenesia, Order iii. Polyg. frustr.*

Digitális (Fox-glove), *Class xiv. Didynamia, Order ii. Angiosperma*

Dílatris, *Class iii. Triandria, Order i. Monogynia*

Dillénia, *Class xiii. Polyandria, Order vii. Polygynia*

Diódia, *Class iv. Tetrandria, Order i. Monogynia*

Dionæa (Venus's Fly-trap), *Class x. Decandria, Order i. Monogynia*

Dioscoréa, *Class xxii. Dioœcia, Order vi. Hexandria*

- Diósma* (African Spirea), *Class* v. Pentandria, *Order* i. Monogynia
Diospýrus (Indian Date Plum), *Class* xxiii. Polygamia, *Order* ii. Dicoecia
Dirca (Leather-wood), *Class* viii. Octandria, *Order* i. Monogyn.
Dipsacus (Teazel), *Class* iv. Tetrandria, *Order* i. Monogynia
Dísa, *Class* xx. Gynandria, *Order* ii. Diandria
Disándra, *Class* vii. Heptandria, *Order* i. Monogynia
Dodártia, *Class* xiv. Didynamia, *Order* ii. Angiosperma
Dódecas, *Class* xi. Dodecandria, *Order* i. Monogynia
Dodecáthcon Meadia, *Class* v. Pentandria, *Order* i. Monogynia
Dodonæa, *Class* viii. Octandria, *Order* i. Monogynia
Dólichos, *Class* xvii. Diadelphia, *Order* iv. Decandria
Doræna, *Class* v. Pentandria, *Order* i. Monogynia
Dombeya, *Class* xiv. Didynamia, *Order* ii. Angiosperma
Dorónicum (Leopard's Bane), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
Dorsténia (Contrayerva), *Class* xv. Tetrandria, *Order* i. Monogyn.
Drába (Whitlow Grass), *Class* xvi. Tetradynamia, *Order* ii. Siliquosa
Dracæna, *Class* vi. Hexandria, *Order* i. Monogynia
Dracocéphalum (Dragon's Head), *Class* xiv. Didynamia, *Order* ii. Angiosperma
Dracóntium (Dragons), *Class* xx. Gynandria, *Order* ix. Polyandria
Drósera (Sun-dew), *Class* v. Pentandria, *Order* v. Pentagynia
Dryandra, *Class* xvii. Monadelphia, *Order* v. Enneandria
Drýas, *Class* xii. Icosandria, *Order* v. Polygynia
Drýpis, *Class* v. Pentandria, *Order* iii. Trigynia
Duránta, *Class* xiv. Didynamia, *Order* ii. Angiosperma
Dúrio, *Class* xviii. Polyadelphia, *Order* iii. Polyandria
Duróia, *Class* vi. Hexandria, *Order* i. Monogynia

E

- Ebenus* (Ebony of Crete), *Class* xvii. Diadelphia, *Order* iv. Decandria

- Echinóphora (Prickly Parsnip), *Class v. Pentandria, Order ii.*
Digynia
- Echinops (Globe Thistle), *Class xix. Syngenesia, Order v. Poly-*
gamia segregata
- Echítes, *Class v. Pentandria, Order i. Monogynia*
- Echium (Viper's Bugloss), *Class v. Pentandria, Order i. Monogy.*
- Eclípta, *Class xix. Syngenesia, Order ii. Polyg. super.*
- Ehrháta, *Class vi. Hexandria, Order i. Monogynia*
- Ehrétia, *Class v. Pentandria, Order i. Monogynia*
- Ekebergia, *Class x. Decandria, Order i. Monogynia*
- Elæagnus (Wild Olive), *Class iv. Tetrandria, Order i. Monogyn.*
- Elæocarpus, *Class xiii. Polyandria, Order i. Monogynia*
- Eláís, Palmæ
- Elaiodendron, *Class v. Pentandria, Order i. Monogynia*
- Elate, Palmæ
- Elatérium, *Class xxi. Monœcia, Order i. Monandria*
- Elatine (Water-wort), *Class viii. Octandria, Order iii. Trigynia*
- Elephántopus (Elephant's Foot), *Class xix. Syngenesia, Order*
v. Polygamia segregata
- Ellisia, *Class v. Pentandria, Order i. Monogynia*
- Elymus, *Class iii. Triandria, Order ii. Digynia*
- Embothrium, *Class iv. Tetrandria, Order i. Monogynia*
- Empetrum (Black-berried Heath, or Crow-berries), *Class xxii.*
Diœcia, Order iii. Triandria
- Epácris, *Class v. Pentandria, Order i. Monogynia*
- Ephedra (Shrubby Horse-tail), *Class xxii. Diœcia, Order xii.*
Monadelphia
- Epidéndrum (Vanilla, or Vanelloe), *Class xx. Gynandria, Or-*
der iv. Diandria
- Epigæa (Trailing Arbutus), *Class x. Decandria, Order i. Mono-*
gynia
- Epilóbium (Willow Herb, or French Willow), *Class viii. Oc-*
tandria, Order i. Monogynia
- Epimédium (Barren-wort), *Class iv. Tetrandria, Order i. Mono-*
gynia
- Equisétum (Horse-tail), *Class xxiv. Cryptogamia, Order i. Filices.*
- Eránthemum, *Class ii. Diandria, Order i. Monogynia*

- Erica* (Heath), *Class* viii. *Octandria*, *Order* i. *Monogynia*
Erigeron, *Class* xix. *Syngenesia*, *Order* ii. *Polyg. super.*
Eriónus, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
Eriocáulon, *Class* iii. *Triandria*, *Order* iii. *Trigynia*
Eriocéphalus, *Class* xix. *Syngenesia*, *Order* iv. *Polyg. neces.*
Erióphorum, *Class* iii. *Triandria*, *Order* i. *Monogynia*
Eriðhalis, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Ervum (Bitter Vetch), *Class* xvii. *Diadelphia*, *Order* iv. *Decandr.*
Eryngium (Eryngo, or Sea Holly), *Class* v. *Pentandria*, *Order* ii.
Digynia
Erysimum (Hedge Mustard), *Class* xv. *Tetradynamia*, *Order* i.
Siliquosa
Erythrina (Coral-tree), *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
Erythronium (Dog's-tooth Violet), *Class* vi. *Hexandria*, *Order* i.
Monogynia
Erythroxylon, *Class* x. *Decandria*, *Order* iii. *Trigynia*
Escallonia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Ethulia, *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
Eucléa, *Class* xxii. *Diœcia*, *Order* x. *Dodecandria*
Eugénia, *Class* xii. *Icosandria*, *Order* i. *Monogynia*
Evólulus, *Class* v. *Pentandria*, *Order* iv. *Tetragynia*
Euonymus (Spindle-tree), *Class* v. *Pentandria*, *Order* i. *Monogynia*
Eupatorium (Hemp Agrimony), *Class* xix. *Syngenesia*, *Order* i.
Polyg. æqualis
Euphorbia (Burning Thorny Plant, or Spurge), *Class* xi. *Dodecandria*, *Order* iii. *Trigynia*
Euphrasia (Eyebright), *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
Eurya, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*
Exacum, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
Excœcaria, *Class* xxii. *Dicœcia*, *Order* iii. *Triandria*

F

- Fagára*, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
Fagónia, *Class* x. *Decandria*, *Order* i. *Monogynia*

- Fágus (Beech), *Class* xxii. Monœcia, *Order* viii. Polyandria
 Falkía, *Class* vi. Hexandria, *Order* ii. Digynia
 Férula (Fennel Giant), *Class* v. Pentandria, *Order* ii. Digynia
 Ferrária, *Class* xx. Gynandria, *Order* iii. Triandria
 Festúca (Fescue Grass), *Class* iii. Triandria, *Order* ii. Digynia
 Fevíllea, *Class* xxii. Diœcia, *Order* v. Pentandria
 Ficus (Fig), *Class* xxiii. Polygamia, *Order* v. Polyœcia
 Filágo (CottonWeed), *Class* xix. Syngenesia, *Order* iv. Polygamia
 necessaria
 Flacourtía, *Class* xxii. Diœciœ, *Order* xii. Icosandria
 Flagellária, *Class* vi. Hexandria, *Order* iii. Trigynia
 Fontínalis (Water Moss), *Class* xxiv. Cryptogamia, *Order* ii.
 Musci
 Forskohléa, *Class* x. Decandria, *Order* iv. Pentagynia
 Fórstera, *Class* xx. Gynandria, *Order* ii. Diandria
 Fothergílla, *Class* xiii. Polyandria, *Order* ii. Digynia
 Fragária (Strawberry), *Class* xii. Icosandria, *Order* v. Polygyn.
 Frankénia, *Class* vi. Hexandria, *Order* i. Monogynia
 Fráxinus (Ash), *Class* xxiii. Polygamia, *Order* ii. Diœcia
 Fritillária (Fritillary), *Class* vi. Hexandria, *Order* i. Monogynia
 Fúchsia, *Class* viii. Octandria, *Order* i. Monogynia
 Fúcus (Wrack, or Sea Weed), *Class* xxiv. Cryptogamia, *Order* iii.
 Algæ
 Fuiréna, *Class* iii. Triandria, *Order* i. Monogynia
 Fumária (Fumitory), *Class* xvii. Diadelphia, *Order* ii. Hexandria
 Fúsanus, *Class* xxiii. Polygamia, *Order* i. Monœcia

G

- Ghínia, *Class* vi. Hexandria, *Order* ii. Digynia
 Galánthus (Snow-drop), *Class* vi. Hexandria, *Order* i. Monogyn.
 Gálax, *Class* v. Pentandria, *Order* i. Monogynia
 Galaxia, *Class* xvi. Monadelphia, *Order* i. Triandria
 Galéga (Goat's Rue), *Class* xvii. Diadelphia, *Order* iv. Decandr.
 Galénia, *Class* viii. Octandria, *Order* ii. Digynia
 Galeópsis (Hedge Nettle), *Class* xiv. Didynamia, *Order* i. Gymnosperma

- Gárium (Lady's Bed-straw), *Class* iv. Tetrandria, *Order* i. Monogynia
- Galopina, *Class* iv. Tetrandria, *Order* ii. Digynia
- Garcinia, *Class* xi. Dodecandria, *Order* i. Monogynia
- Gardénia (Cape Jasmine), *Class* v. Pentandria, *Order* i. Monogynia
- Garidélla (Fennel-Flower of Crete), *Class* x. Decandria, *Order* iii. Trigynia
- Gaulthéria, *Class* x. Decandria, *Order* i. Monogynia
- Gáura (Virginian Loosestrife), *Class* viii. Octandria, *Order* i. Monogynia
- Génipa, *Class* v. Pentandria, *Order* i. Monogynia
- Genísta (Single-seeded Broom), *Class* xvii. Diadelphia, *Order* iv. Decandria
- Gentiána (Gentian, or Fell-wort), *Class* v. Pentandria, *Order* ii. Digynia
- Geoffróya, *Class* xvii. Diadelphia, *Order* iv. Decandria
- Geránium (Crane's Bill), *Class* xvi. Monadelphia, *Order* iv. Decandria
- Gerárdia, *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Geropógon, *Class* xix. Syngenesia, *Order* i. Polyg. æqualis
- Gesnéria, *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Gethýllis, *Class* xi. Dodecandria, *Order* i. Monogynia
- Géum (Aven's, or Herb Bennet), *Class* xii. Icosandria, *Order* v. Polygynia
- Ginóra, *Class* xi. Dodecandria, *Order* i. Monogynia
- Ginkgo, *Planta Obscura.*
- Gisékia, *Class* v. Pentandria, *Order* v. Pentagynia
- Glabrária, *Class* xiii. Polyadelphia, *Order* iii. Polyandria
- Gladíolus (Corn Flag), *Class* iii. Triandria, *Order* i. Monogynia
- Glaúx (Sea Milk-wort, or Black Salt-wort), *Class* v. Pentandria, *Order* i. Monogynia
- Glecóma (Ground Ivy, or Gill), *Class* xiv. Didynamia, *Order* i. Gymnosperma
- Gledítsia (Three-thorned Acacia), *Class* xxiii. Polygamia, *Order* ii. Diœcia
- Glinus, *Class* xi. Dodecandria, *Order* v. Pentagynia

- Glóbbá, *Class* ii. Diandria, *Order* i. Monogynia.
- Globulária (Blue Daisy), *Class* iv. Tetrandria, *Order* i. Monogyn.
- Gloriósa (Superb Lily), *Class* vi. Hexandria, *Order* i. Monogyn.
- Glúta, *Class* xx. Gynandria, *Order* v. Pentandria
- Glýcine (Carolina Kidney-bean Tree), *Class* xvii. Diadelphia, *Order* iv. Decandria
- Glycyrrhíza (Liquorice), *Class* xvii. Diadelphia, *Order* iv. Decandria
- Gmelína, *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Gnaphálium (Cudweed), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
- Gnétum, *Class* xxii. Monœcia, *Order* ix. Monadelphia
- Gnidia, *Class* viii. Octandria, *Order* i. Monogynia
- Gomózia, *Class* iv. Tetrandria, *Order* ii. Digynia
- Gomphréna (Globe Amaranth), *Class* v. Pentandria, *Order* ii. Digynia
- Gonocárpus, *Class* iv. Tetrandria, *Order* i. Monogynia
- Gordónia, *Class* xvi. Monadelphia, *Order* vii. Polyandria
- Gortéria, *Class* xix. Syngenesia, *Order* iii. Polyg. frustr.
- Gossýpium (Cotton), *Class* xvi. Monadelphia, *Order* vii. Polyandria
- Gouánia, *Class* xxiii. Polygamia, *Order* i. Monœcia
- Gratióla (Hedge Hyssop), *Class* ii. Diandria, *Order* i. Monogyn.
- Gréwia, *Class* xx. Gynandria, *Order* ix. Polyandria
- Grías, *Class* xiii. Polyandriá, *Class* i. Monogynia
- Griélum, *Class* x. Decandria, *Order* v. Pentagynia
- Grisléa, *Class* viii. Octandria, *Order* i. Monogynia
- Gronóvia, *Class* v. Pentandria, *Order* i. Monogynia
- Gúajacum (Lignum Vitæ), *Class* x. Decandria, *Order* i. Monogyn.
- Guaréa, *Class* viii. Octandria, *Order* i. Monogynia
- Guettárda, *Class* xxi. Monœcia, *Order* vii. Heptandria
- Guilandína (Bonduc, or Nickar-tree), *Class* x. Decandria, *Order* i. Monogynia
- Gundélia, *Class* xix. Syngenesia, *Order* v. Polygamia segregata
- Gúnnera, *Class* xx. Gynandria, *Order* i. Diandria
- Gustávia, *Class* xvi. Monadelphia, *Order* vii. Polyandria
- Gypsóphila, *Class* x. Decandria, *Order* ii. Digynia

H

- Hæmánthus* (Blood Flower), *Class* vi. *Hexandria*, *Order* i. *Monogynia*
Hæmatóxyllum (Logwood), *Class* x. *Decandria*, *Order* i. *Monogynia*
Halésia, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*
Haléria (African Fly-honeysuckle), *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
Halóragis, *Class* viii. *Octandria*, *Order* iv. *Tetragynia*
Hamamélis (Witch Hazel), *Class* iv. *Tetrandria*, *Order* ii. *Digyn.*
Hainéllia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Hartógia, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
Hasselquístia, *Class* v. *Pentandria*, *Order* ii. *Digynia*
Hebénstrétia, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
Hédera (Ivy), *Class* v. *Pentandria*, *Order* i. *Monogynia*
Hedycárya, *Class* xxii. *Diœcia*, *Order* xi. *Polyandria*
Hedyótis, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
Hedýsarum (French Honeysuckle), *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
Heistéria, *Class* x. *Decandria*, *Order* i. *Monogynia*
Helénium (Bastard Sunflower), *Class* xix. *Syngenesia*, *Order* ii. *Polyg. super.*
Heliánthus (Sunflower), *Class* xix. *Syngenesia*, *Order* iii. *Polyg. frustr.*
Helicónia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
Helictéres (Skrew Tree), *Class* xx. *Gynandria*, *Order* vii. *Decandria*
Heliocárpus, *Class* xi. *Dodecandria*, *Order* ii. *Digynia*
Helióphila, *Class* xv. *Tetradynamia*, *Order* i. *Siliquosa*
Heliotrópium (Turn-sole), *Class* v. *Pentandria*, *Order* i. *Monogynia*
Helónias, *Class* vi. *Hexandria*, *Order* iii. *Trigynia*
Hellebórus (Black Hellebore), *Class* xiii. *Polyandria*, *Order* vii. *Polygynia*
Helvélla, *Class* xxiv. *Cryptogamia*, *Order* iv. *Fungi*
Hemerocállis (Day Lily, or Lily Asphodel), *Class* vi. *Hexandria*, *Order* i. *Monogynia*

- Hemionítis (Mule's Fern), *Class xxiv. Cryptogamia, Order i. Filices*
- Heniméris, *Class xiv. Didynamia, Order ii. Angiosperma*
- Heracléum (Cow Parsnep), *Class v. Pentandria, Order ii. Digynia*
- Hermánia, *Class xvi. Monadelphia, Order ii. Pentandria*
- Hérmas, *Class xxiii. Polygamia, Order i. Monœcia*
- Hernándia (Jack in a Box), *Class xxi. Monœcia, Order iii. Triandria*
- Herniária (Rupture Wort), *Class v. Pentandria, Order ii. Digynia*
- Hésperis (Dame's Violet, Rocket, or Queen's July Flower), *Class xvi. Tetradyndamia, Order i. Siliquosa*
- Heúchéra, *Class v. Pentandria, Order ii. Digynia*
- Hibíscus (Althea Frutex, or Syrian Mallow), *Class xvi. Monadelphia, Order vii. Polyandria*
- Hierácium (Hawk-weed), *Class xix. Syngenesia, Order i. Polyg. æqualis*
- Híllia, *Class vi. Hexandria, Order i. Monogynia*
- Híppia, *Class xix. Syngenesia, Order iv. Polygamia necessaria*
- Hippocratéa, *Class iii. Triandria, Order i. Monogynia*
- Hippocrépis (Horseshoe Vetch), *Class xvii. Diadelphía, Order iv. Decandria*
- Hippómane (Manchineel), *Class xxi. Monœcia, Order ix. Monadelphia*
- Hippóphaë (Sea Buckthorn), *Class xxii. Diœcia, Order iv. Tetrandia*
- Hippúris, *Class i. Monandria, Order i. Monogynia*
- Hiræa, *Class x. Decandria, Order iii. Trigynia*
- Hirtélla, *Class v. Pentandria, Order i. Monogynia*
- Hólcus (Indian Millet), *Class xxiii. Polygamia, Order i. Monœcia*
- Holósteum, *Class iii. Triandria, Order iii. Trigynia*
- Hopéa, *Class xviii. Polyadelphia, Order iii. Polyandria*
- Hórdeum (Barley), *Class iii. Triandria, Order ii. Digynia*
- Hormínium (Pyrenæan Clary), *Class xiv. Didynamia, Order i. Gymnosperma*
- Hottónia (Water Milfoil, or Water Violet), *Class v. Pentandria, Order i. Monogynia*
- Hovénia, *Class v. Pentandria, Order i. Monogynia*

- Houstónia, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
 Houtuynia, *Class* xiii. *Polyandria*, *Order* vii. *Polygynia*
 Hudsónia, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*
 Hugónia, *Class* xvi. *Monadelphia*, *Order* iv. *Decandria*
 Húmulus (Hop), *Class* xxii. *Diœcia*, *Order* v. *Pentandria*
 Húra (Sand-box Tree), *Class* xxi. *Monœcia*, *Order* ix. *Monadelphia*
 Hyacínthus (Hyacinth), *Class* vi. *Hexandria*, *Order* i. *Monogyn.*
 Hýdnum, *Class* xxiv. *Cryptogamia*, *Order* iv. *Fungi*
 Hydránga, *Class* x. *Decandria*, *Order* i. *Monogynia*
 Hydrástis (Yellow Root), *Class* xiii. *Polyandria*, *Order* vii. *Polygynia*
 Hydrócharis (Frog's-bit), *Class* xxii. *Diœcia*, *Order* viii. *Enneandria*
 Hydrocótyle (Water Navel-wort), *Class* v. *Pentandria*, *Order* ii. *Digynia*
 Hydrólea, *Class* v. *Pentandria*, *Order* ii. *Digynia*
 Hydrophýlax, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
 Hydrophýllum (Water Leaf), *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Hymenæa (Locust-tree, or Courbaril), *Class* x. *Decandria*, *Order* i. *Monogynia*
 Hyobánche, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
 Hyoscýamus (Henbane), *Class* v. *Pentandria*, *Order* i. *Monogyn.*
 Hyóseris, *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
 Hypécoum, *Class* iv. *Tetrandria*, *Order* ii. *Digynia*
 Hypericum (St. John's Wort), *Class* xviii. *Polyadelphia*, *Order* iii. *Polyandria*
 Hýpnum, *Class* xxi. *Cryptogamia*, *Order* ii. *Musci*
 Hypochæris, *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqualis*
 Hypóxis, *Class* vi. *Hexandria*, *Order* i. *Monogynia*
 Hyssópus (Hyssop), *Class* xiv. *Didynamia*, *Order* i. *Gymnosper.*

I

- Jacquínia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Jambolífera, *Class* viii. *Octandria*, *Order* i. *Monogynia*

- Jasione (Sheep Scabious), *Class* xix. Syngenesia, *Order* vi. Monogynia
- Jasmínium (Jasmine), *Class* ii. Diandria, *Order* i. Monogynia
- Játropha (Cassava), *Class* xxi. Monœcia, *Order* ix. Monadelph.
- Iberis (Candy Tuft, or Sciatic Cress), *Class* iv. Tetrodynamia, *Order* ii. Siliculosa
- Ignátia, *Class* v. Pentandria, *Order* i. Monogynia
- Ilex (Holly), *Class* iv. Tetrandria, *Order* iii. Tetragynia
- Illecébrum (Mountain Knot Grass), *Class* v. Pentandria, *Order* i. Monogynia
- Illicium, *Class* xiii. Polyandria, *Order* vii. Polygynia
- Impátiens (Balsam, or Female Balsamine), *Class* xix. Syngenesia, *Order* vi. Monogamia
- Imperatória (Master-wort), *Class* v. Pentandria, *Order* ii. Digynia
- Indigófera (Indigo), *Class* xvii. Diadelphia, *Order* iv. Decandria
- Inocárpus, *Class* x. Decandria, *Order* i. Monogynia
- Inula (Elacampane), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
- Ipomœa (Quamoclit), *Class* v. Pentandria, *Order* i. Monogynia
- Irésine, *Class* xxii. Diœcia, *Order* v. Pentandria
- Iris (Flower de Luce), *Class* iii. Triandria, *Order* i. Monogynia
- Isátis (Woad), *Class* xv. Tetrodynamia, *Order* i. Siliquosa
- Ischæmum, *Class* xxiii. Polygamia, *Order* i. Monœcia
- Isnárdia, *Class* xiv. Tetrandria, *Order* i. Monogynia
- Isœëtis, *Class* xxiv. Cryptogamia, *Order* i. Filices
- Isofýrum, *Class* xiii. Polyandria, *Order* vii. Polygynia
- Itea, *Class* v. Pentandria, *Order* i. Monogynia
- Iva (Jesuits' Bark Tree), *Class* xxi. Monœcia, *Order* v. Pentand.
- Júglans (Wallnut), *Class* xxi. Monœcia, *Order* viii. Polyandria
- Júncus (Rush), *Class* vi. Hexandria, *Order* i. Monogynia
- Jungermánua, *Class* xxiv. Cryptogamia, *Order* iii. Algæ
- Júngia, *Class* xix. Syngenesia, *Order* v. Polyg. segreg.
- Juníperus (Juniper), *Class* xxii. Diœcia, *Order* xii. Monadelph.
- Jussiaea, *Class* x. Decandria, *Order* i. Monogynia
- Justícia (Malabar Nut), *Class* ii. Diandria, *Order* i. Monogynia
- Ixia, *Class* iii. Triandria, *Order* i. Monogynia
- Ixora, *Class* iv. Tetrandria, *Order* i. Monogynia

K

- Kálmnia (Dwarf American Laurel), *Class* x. Decandria, *Order* i.
 Monogynia
- Kæmpféria, *Class* i. Monandria, *Order* i. Monogynia
- Kiggelária, *Class* xxii. Diœcia, *Order* ix. Decandria
- Kleinhóvia, *Class* xx. Gynandria, *Order* vii. Decandria
- Knáutia, *Class* iv. Tetrandria, *Order* i. Monogynia
- Knóxia, *Class* iv. Tetrandria, *Order* i. Monogynia
- Kœnigia, *Class* iii. Triandria, *Order* iii. Trigynia
- Krameria, *Class* iv. Tetrandria, *Order* i. Monogynia
- Kúhnia, *Class* v. Pentandria, *Order* i. Monogynia
- Kyllinga, *Class* iii. Triandria, *Order* i. Monogynia

L

- Lachenália, *Class* vi. Hexandria, *Order* i. Monogynia
- Lachnæa, *Class* viii. Octandria, *Order* i. Monogynia
- Lactúca (Lettuce), *Class* xix. Syngenesia, *Order* i. Polyg. æqu.
- Lœtia, *Class* xiii. Polyandria, *Order* i. Monogynia
- Lagerstrœmia, *Class* xiii. Polyandria, *Order* i. Monogynia
- Lagœcia (Bastard Cumin), *Class* v. Pentandria, *Order* i. Monogyn.
- Lagúrus (Hare's-tail Grass), *Class* iii. Triandria, *Order* ii. Digyn.
- Lámium (Dead Nettle, or Archangel), *Class* xiv. Didynamia,
Order i. Gynnosperma
- Lántana (American Viburnum), *Class* xiv. Didynamia, *Order* ii.
 Angiosperma
- Lápsana (Nipple-wort), *Class* xix. Syngenesia, *Order* i. Polyg.
 æqualis
- Laserpítium (Laser-wort), *Class* v. Pentandria, *Order* ii. Digyn.
- Lathræa, *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Lathýrus (Chichling Vetch), *Class* xvii. Diadelphia, *Order* iv.
 Decandria
- Lavéndula (Lavender), *Class* xiv. Didynamia, *Order* ii. Angiosp.
- Lavatéra; *Class* xvi. Monadelphia, *Order* vii. Polyandria
- Laugéria, *Class* v. Pentandria, *Order* i. Monogynia
- Láurus (Bay), *Class* ix. Enneandria, *Order* i. Monogynia
- Lawsónia, *Class* viii. Octandria, *Order* i. Monogynia

- Leéa, *Class* xxi. Monœcia, *Order* v. Pentandria
 Lechéa, *Class* iii. Triandria, *Order* iii. Trigynia
 Lécythis, *Class* xiii. Polyandria, *Order* i. Monogynia
 Léдум (Marsh Cistus, or Wild Rosemary), *Class* x. Decandria,
Order i. Monogynia
 Lémna (Duck Meat), *Class* xxi. Monœcia, *Order* ii. Diandria
 Leóntice (Lion's Leaf), *Class* vi. Hexandria, *Order* i. Monogynia
 Leóntodon (Dandelion), *Class* xix. Syngenesia, *Order* i. Polyg.
 æqualis
 Leonúrus (Lion's Tail), *Class* xiv. Didynamia, *Order* i. Gymnosp.
 Lepídium (Dittander, or Pepper-wort), *Class* xv. Tetradynamia,
Order ii. Siliculosa
 Lerchéa, *Class* xvi. Monadelphía, *Order* ii. Pentandria
 Leucójum (Greater Snow-drop), *Class* vi. Hexandria, *Order* i.
 Monogynia
 Leyséra, *Class* xix. Syngenesia, *Order* ii. Polyg. super.
 Lichen (Liver-wort), *Class* xxiv. Cryptogamia, *Order* iii. Algæ.
 Licuala, *Class* vi. Hexandria, *Order* i. Monogynia
 Ligústicum (Lovage), *Class* v. Pentandria, *Order* ii. Digynia
 Ligústrum (Privet), *Class* ii. Diandria, *Order* i. Monogynia
 Lílium (Lily), *Class* vi. Hexandria, *Order* i. Monogynia
 Liméum, *Class* vii. Heptandria, *Order* ii. Digynia
 Limodórum, *Class* xx. Gynandria, *Order* i. Diandria
 Limónia, *Class* x. Decandria, *Order* i. Monogynia
 Limosélla (Least Water Plantain), *Class* xiv. Didynamia, *Order* ii.
 Angiosperma
 Lindérnia, *Class* xiv. Didynamia, *Order* ii. Angiosperma
 Lincónia, *Class* v. Pentandria, *Order* ii. Digynia
 Lindera, *Class* vi. Hexandria, *Order* i. Monogynia
 Linnæa, *Class* xiv. Didynamia, *Order* ii. Angiosperma
 Línium (Flax), *Class* v. Pentandria, *Order* v. Pentagynia
 Lipária, *Class* xvii. Diadelphia, *Order* iv. Decandria
 Lippia, *Class* xiv. Didynamia, *Order* ii. Angiosperma
 lygynia
 Liquidámber (Sweet Gum), *Class* xxi. Monœcia, *Order* viii.
 Polyandria
 Liriodéndrum (Tulip Tree), *Class* xiii. Polyandria, *Order* vii. Po-
 lisiánthus, *Class* v. Pentandria, *Order* i. Monogynia

- Lithospérmum (Gromwell), *Class v. Pentandria, Order i. Monogyn.*
 Littorélla, *Class xxi. Monœcia, Order iv. Tetrandia*
 Lobélia (Cardinal Flower), *Class xix. Syngenesia, Order vi. Monogamia*
 Lœflingia, *Class iii. Triandria, Order i. Monogynia*
 Lœsélia, *Class xiv. Didynamia, Order ii. Angiosperma*
 Lolium (Darnel, or Rye-grass), *Class iii. Triandria, Order ii. Digynia*
 Lonchítis (Rough Spleen-wort), *Class xxiv. Cryptogamia, Order i. Filices*
 Lonicéra (Honeysuckle), *Class v. Pentandria, Order i. Monogyn.*
 Loósa, *Class xiii. Polyandria, Order i. Monogynia*
 Loránthus, *Class vi. Hexandria, Order i. Monogynia*
 Lótus (Bird's-foot Trefoil), *Class xvii. Diadelphia, Order iv. Decandria*
 Ludwígia, *Class iv. Tetrandia, Order i. Monogynia*
 Lunária (Moon-wort, Satin Flower, or Honesty), *Class xv. Tetradynamia, Order ii. Siliculosa*
 Lupínus (Lupine), *Class xvii. Diadelphia, Order iv. Decandria*
 Lýchnis (Campion), *Class xviii. Decandria, Order v. Pentagynia*
 Lýcium (Box-thorn), *Class v. Pentandria, Order i. Monogynia*
 Lycopérdon, *Class xxiv. Cryptogamia, Order iv. Fungi*
 Lycopódium (Wolf's-claw Moss), *Class xxiv. Cryptogamia, Order ii. Musci*
 Lycopsis, *Class v. Pentandria, Order i. Monogynia*
 Lýcopus (Water Horehound), *Class x. Decandria, Order i. Monogynia*
 Lygéum (Hooded Matweed), *Class iii. Triandria, Order i. Monogynia*
 Lysimáchia (Loosestrife), *Class v. Pentandria, Order i. Monogyn.*
 Lýthrum (Willow Herb), *Class xi. Dodecandria, Order i. Monogynia*

M

- Mába, *Class xxii. Diœcia, Order iii. Triandria*
 Macrocnémum, *Class v. Pentandria, Order i. Monogynia*
 Magnólia (Laurel-leaved Tulip-tree), *Class xiii. Polyandria, Order vii. Polygynia*

- Mahérnia, *Class v. Pentandria, Order v. Pentagynia*
 Málachra, *Class xvi. Monadelphia, Order vii. Polyandria*
 Málope (Bastard Mallow), *Class xvi. Monadelphia, Order vii. Polyandria*
 Malpíghia (Barbadoes Cherry), *Class x. Decandria, Order iii. Trigynia*
 Málva (Mallow), *Class xvi. Monadelphia, Order vii. Polyandria*
 Mamméa (Mammee), *Class xiii. Polyandria, Order i. Monogyn.*
 Manéttia, *Class iv. Tetrandria, Order i. Monogynia*
 Mangífera (Mango Tree), *Class v. Pentandria, Order i. Monogyn.*
 Manisúris, *Class xxiii. Polygamia, Order i. Monœcia*
 Manúlea, *Class xiv. Didynamia, Order ii. Angiosperma*
 Maránta (Indian Arrow Root), *Class i. Monandria, Order i. Monogynia*
 Marcgrávia, *Class xiii. Polyandria, Order vii. Polygynia*
 Marchántia, *Class xxiv. Cryptogamia, Order iii. Algæ*
 Margaritária, *Class xxii. Dioœcia, Order viii. Enneandria*
 Marrúbium (Horehound), *Class xiv. Didynamia, Order i. Gymnosperma*
 Marsiléa, *Class xxiv. Cryptogamia, Order i. Filices*
 Martýnia, *Class xiv. Didynamia, Order ii. Angiosperma*
 Massónia, *Class vi. Hexandria, Order i. Monogynia*
 Matricária (Feverfew), *Class xix. Syngenesia, Order ii. Polyg. super.*
 Matthiôla, *Appendix*
 Maurítia, *Appendix*
 Medéola (Climbing African Asparagus), *Class vi. Hexandria, Order iii. Trigynia*
 Medicágo (Snail and Moon Trefoil), *Class xvii. Diadelphia, Order iv. Decandria*
 Melaléuca, *Class xviii. Polyadelphia, Order iii. Polyandria*
 Melampódium, *Class xix. Syngenesia, Order iv. Polyg. necess.*
 Melampýrum (Cow-wheat), *Class xiv. Didynamia, Order ii. Angiosperma*
 Melánthium, *Class vi. Hexandria, Order iii. Trigynia*
 Melástoma (American Gooseberry), *Class x. Decandria, Order i. Monogynia*

- Mélia (Bead Tree), *Class* x. Decandria, *Order* i. Monogynia
- Meliánthus (Honey-flower), *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Mélica, *Class* iii. Triandria, *Order* ii. Digynia
- Melicócca, *Class* viii. Octandria, *Order* i. Monogynia
- Melíssa (Baum), *Class* xiv. Didynamia, *Order* i. Gymnosperma
- Melíttis (Baum-leaved Archangel, or Bastard Baum), *Class* xiv. Didynamia, *Order* i. Gymnosperma
- Melóchia, *Class* xvi. Monadelphía, *Order* ii. Pentandria
- Melódinus, *Class* v. Pentandria, *Order* ii. Digynia
- Melótheria (Small Creeping Cucumber), *Class* ix. Triandria, *Order* i. Monogynia
- Memécylon, *Class* viii. Octandria, *Order* i. Monogynia
- Menáís, *Class* v. Pentandria, *Order* i. Monogynia
- Menispérmum (Moon Seed), *Class* xxii. Diœcia, *Order* x. Dodecandria
- Méntha (Mint), *Class* xiv. Didynamia, *Order* i. Gymnosperma
- Mentzélia, *Class* xiii. Polyandria, *Order* i. Monogynia
- Menyánthes (Bog-bean, or Marsh Trefoil), *Class* v. Pentandria, *Order* i. Monogynia
- Mercuriális (Mercury), *Class* xxii. Diœcia, *Order* viii. Enneand.
- Mesembryánthemum (Fig Marygold), *Class* xii. Icosandria, *Order* iv. Pentagynia
- Messerschmídia, *Class* v. Pentandria, *Order* i. Monogynia
- Méspilus (Medlar), *Class* xii. Icosandria, *Order* iv. Pentagynia
- Mésua (Indian Rose Chesnut), *Class* xvi. Monadelphía, *Order* viii. Polyandria
- Michélia, *Class* xiii. Polyandria, *Order* vii. Polygynia
- Mícropus (Bastard Cudweed), *Class* xix. Syngenesia, *Order* iv. Polyg. neces.
- Mílium (Millet), *Class* iii. Triandria, *Order* ii. Digynia
- Milléria, *Class* xix. Syngenesia, *Order* iv. Polyg. neces.
- Millingtónia, *Class* xiv. Didynamia, *Order* ii. Angiosperma
- Mimósa (Sensitive Plant), *Class* xxiii. Polygamia, *Order* i. Monœcia
- Mímulus (Monkey Flower), *Class* xiv. Didynamia, *Order* ii. Angiosperma

- Mimúsops, *Class* viii. Octandria, *Order* ii. Digynia
 Minuártia, *Class* iii. Triandria, *Order* iii. Trigynia
 Mirábilis (Marvel of Peru), *Class* v. Pentandria, *Order* i. Monogynia
 Mitchélla, *Class* iv. Tetrandria, *Order* i. Monogynia
 Mitélla (Bastard American Sanicle), *Class* x. Decandria, *Order* ii. Digynia
 Mníarum, *Class* i. Monandria, *Order* ii. Digynia
 Mníum, *Class* xxiv. Cryptogamia, *Order* ii. Musci
 Mœhríngia (Mountain Chickweed), *Class* viii. Octandria, *Order* ii. Digynia
 Mollúgo, *Class* iii. Triandria, *Order* iii. Trigynia
 Moluccélla (Molucca Baum), *Class* xiv. Didynamia, *Order* i. Gymnosperma
 Mómórica (Male Balsam Apple), *Class* xxi. Monœcia, *Order* x. Syngenesia
 Monárda (Oswego Tea), *Class* ii. Diandria, *Order* i. Monogynia
 Monetia, *Class* iv. Tetrandria, *Order* i. Monogynia
 Monniéria, *Class* xvii. Diadelphia, *Order* i. Pentandria
 Monotrópa, *Class* x. Decandria, *Order* i. Monogynia
 Monsónia, *Class* xviii. Polyadelphia, *Order* 0. Dodecandria
 Móntia (Blinks), *Class* iii. Triandria, *Order* iii. Trigynia
 Montínia, *Class* xxii. Diœcia, *Order* iv. Tetrandria
 Moræa, *Class* iii. Triandria, *Order* i. Monogynia
 Morína, *Class* ii. Diandria, *Order* i. Monogynia
 Morinda, *Class* v. Pentandria, *Order* i. Monogynia
 Morisónia, *Class* xiii. Polyandria, *Order* i. Monogynia
 Mórús (Mulberry Tree), *Class* xxi. Monœcia, *Order* iv. Tetrand.
 Múcor, *Class* xxiv. Cryptogamia, *Order* iv. Fungi
 Mulléra, *Class* xvii. Diadelphia, *Order* iv. Decandria
 Muncháusia, *Class* xviii. Polyadelphia, *Order* 0. Polyandria
 Muntíngia, *Class* xiii. Polyandria, *Order* i. Monogynia
 Murráya, *Class* x. Decandria, *Order* i. Monogynia
 Músa (Plantain Tree), *Class* xiii. Polyandria, *Order* i. Monœcia
 Mussænda, *Class* v. Pentandria, *Order* i. Monogynia
 Mutísia, *Class* xix. Syngenesia, *Order* ii. Polyg. super.
 Myágrum (Gold of Pleasure), *Class* xv. Tetradynamia, *Order* ii. Siliculosa

- Mygínda, *Class* iv. *Tetrandria*, *Order* iii. *Tetragynia*
 Myosótis (Mouse-ear Scorpion Grass), *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Myosúrus (Mouse-tail), *Class* v. *Pentandria*, *Order* i. *Monogyn.*
 Myrica (Candleberry Myrtle-gale, or Sweet Willow), *Class* xxii. *Diœcia*, *Order* iv. *Tetrandria*
 Myriophýllum (Water Milfoil), *Class* xxi. *Monœcia*, *Order* viii. *Polyandria*
 Myrósmá, *Class* i. *Monandria*, *Order* i. *Monogynia*
 Mýrsine (African Box Tree), *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Myróxylon, *Class* x. *Decandria*, *Order* i. *Monogynia*
 Mýrtus (Myrtle), *Class* xii. *Icosandria*, *Order* i. *Monogynia*
 Myristica, *Class* xiii. *Polyandria*, *Order* i. *Monogynia*

N

- Nájas, *Class* xxii. *Diœcia*, *Order* i. *Monandria*
 Náma, *Class* v. *Pentandria*, *Order* ii. *Digynia*
 Nandina, *Class* vi. *Hexandria*, *Order* i. *Monogynia*
 Napæa, *Class* xxii. *Diœcia*, *Order* xii. *Monadelphia*
 Narcíssus (Daffodil), *Class* vi. *Hexandria*, *Order* i. *Monogynia*
 Nárdus, *Class* iii. *Triandria*, *Order* i. *Monogynia*
 Naucléa, *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Nepénthes, *Class* xx. *Gynandria*, *Order* iii. *Tetrandria*
 Népetá (Cat-mint, or Nep), *Class* xiv. *Didynamia*, *Order* i. *Gymnosperma*
 Nephélium, *Class* xxi. *Monœcia*, *Order* v. *Pentandria*
 Nérium (Oleander, or Rose Bay), *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Neuráda, *Class* x. *Decandria*, *Order* v. *Decagynia*
 Nicotiána (Tobacco), *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Nigélla (Fennel Flower, or Devil in a Bush), *Class* xiii. *Polyandria*, *Order* v. *Pentagynia*
 Nigrína, *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Nipa, *Class* xxi. *Monœcia*, *Order* i. *Monandria*
 Nissólia, *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
 Nitrária, *Class* xi. *Dodecandria*, *Order* i. *Monogynia*

- Nolána, *Class v. Pentandria, Order i. Monogynia*
 Nyctánthes (Arabian Jasmine), *Class ii. Diandria, Order i. Monogynia*
 Nymphæa (Water Lily), *Class xiii. Polyandria, Order i. Monogynia*
 Nýssa (Tupelo Tree), *Class xxiii. Polygamia, Order ii. Diœcia*

O

- Obolária, *Class xiv. Didynamia, Order ii. Angiosperma*
 Ochna, *Class xiii. Polyandria, Order i. Monogynia*
 Ocymum (Basil), *Class xiv. Didynamia, Order i. Gymnosperma*
 Œdéra, *Class xix. Syngenesia, Order v. Polygamia segregata*
 Œnánthe (Water Drop-wort), *Class v. Pentandria, Order ii. Digynia*
 Œnothéra (Tree Primrose), *Class viii. Octandria, Order i. Monogynia*
 Olax, *Class iii. Triandria, Order i. Monogynia*
 Oldenlándia, *Class iv. Tetrandria, Order i. Monogynia*
 Oléa (Olive), *Class ii. Diandria, Order i. Monogynia*
 Olýra, *Class xxi. Monœcia, Order iii. Triandria*
 Omphaléa, *Class xxi. Monœcia, Order iii. Triandria*
 Onocléa (Sensible Polypody), *Class xxiv. Cryptogamia, Order i. Filices*
 Onónis (Rest Harrow), *Class xvii. Diadelphia, Order iv. Decandria*
 Onopórdum (Woolly Thistle), *Class xix. Syngenesia, Order i. Polyg. æqual.*
 Onósma, *Class v. Pentandria, Order i. Monogynia*
 Ophioglóssum (Adder's Tongue), *Class xxii. Cryptogamia, Order i. Filices*
 Ophiorrhíza (Serpent's Tongue), *Class v. Pentandria, Order i. Monogynia*
 Ophióxylon, *Class xxiii. Polygamia, Order i. Monœcia*
 Ophíra, *Class viii. Octandria, Order i. Monogynia*
 Ophrys (Twyblade), *Class xx. Gynandria, Order i. Diandria*
 Orchis, *Class xx. Gynandria, Order i. Diandria*

- Origanum (Wild Marjorum), *Class* xiv. *Didynamia*, *Order* vii.
 Gymnosperma
 Oríxa, *Class* xv. *Tetrandria*, *Order* i. *Monogynia*
 Ornithógalum (Star of Bethlehem), *Class* vi. *Hexandria*, *Order* i.
 Monogynia
 Ornithopus (Bird's Foot), *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
 Orobánche (Broom Rape), *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
 Orobus (Bitter Vetch), *Class* xvii. *Diadelphia*, *Order* iv. *Decandr.*
 Oróntium (Floating Arum), *Order* vi. *Hexandria*, *Order* i. *Monogynia*
 Ortégia, *Class* iii. *Triandria*, *Order* i. *Monogynia*
 Orýza (Rice), *Class* vi. *Hexandria*, *Order* ii. *Digynia*
 Osbéckia, *Class* viii. *Octandria*, *Order* i. *Monogynia*
 Osmítes, *Class* xix. *Syngenesia*, *Order* iii. *Polyg. frustr.*
 Osmúnda (Osmund Royal, or Flowering Fern), *Class* xxiv.
 Cryptogamia, *Order* i. *Filices*
 Osteospermum (Hard-seeded Chrysanthemum), *Class* xix. *Syngenesia*, *Order* iv. *Polygamia necessaria*
 Osýris (Poet's Cassia), *Class* xxii. *Dicæcia*, *Order* iii. *Triandria*
 Óthera, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
 Orthónna (African Ragwort), *Class* xix. *Syngenesia*, *Order* iv.
 Polygamia necessaria
 Oviéda, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
 Oxalis (Wood Sorrel), *Class* x. *Decandria*, *Order* iv. *Pentagynia*

P

- Pæderóta, *Class* ii. *Diandria*, *Order* i. *Monogynia*
 Pædéria, *Class* v. *Pentandria*, *Order* i. *Monogynia*
 Pæónia (Pæony), *Class* xiii. *Polyandria*, *Order* ii. *Digynia*
 Pallásia, *Class* xi. *Dodecandria*, *Order* iii. *Trigynia*
 Pánax (Ginseng), *Class* xxiii. *Polygamia*, *Order* ii. *Dicæcia*
 Pancrátium (Sea Daffodil), *Class* vi. *Hexandria*, *Order* i. *Monogynia*
 Pándanus, *Class* xxii. *Dicæcia*, *Order* i. *Monandria*

- Pánicum (Panic Grass), *Class* iii. Triandria, *Order* ii. Digynia
 Papáver (Poppy), *Class* xiii. Polyandria, *Order* i. Monogynia
 Parietária (Pellitory), *Class* xxiii. Polygamia, *Order* i. Monœcia
 Páris (Herb True-love, or One-berry), *Class* viii. Octandria,
Order iv. Tetragynia
 Parkinsónia, *Class* x. Decandria, *Order* i. Monogynia
 Parnássia (Glass of Parnassus), *Class* v. Pentandria, *Order* iv.
 Tetragynia
 Parthénium (Bastard Feverfew), *Class* xxi. Monœcia, *Order* v.
 Pentandria
 Páspalum, *Class* iii. Triandria, *Order* ii. Digynia
 Passerina (Sparrow-wort), *Class* viii. Octandria, *Order* i. Mo-
 nogynia
 Passiflóra (Passion Flower), *Class* xx. Gynandria, *Order* iv. Pen-
 tandria
 Pastináca (Parsnip), *Class* v. Pentandria, *Order* ii. Digynia
 Patagónula, *Class* v. Pentandria, *Order* i. Monogynia
 Pavétta, *Class* iv. Tetrandria, *Order* i. Monogynia
 Paulínia, *Class* viii. Octandria, *Order* ii. Trigynia
 Péctis, *Class* xix. Syngenesia, *Order* ii. Polyg. super.
 Pedálium, *Class* xiv. Didynamia, *Order* ii. Angiosperma
 Pediculáris (Rattle Coxcomb, or Louse-wort), *Class* xiv. Didy-
 namia, *Order* ii. Angiosperma
 Pegánum (Wild Syrian Rue), *Class* xi. Dodecandria, *Order* i. Mo-
 nogynia
 Peltária, *Class* xv. Tetradynamia, *Order* ii. Siliculosa
 Penæa, *Class* iv. Tetrandria, *Order* i. Monogynia
 Pentápedes, *Class* xvi. Monadelphia, *Order* vi. Dodecandria
 Pénthorum, *Class* x. Decandria, *Order* iv. Pentagynia
 Péplis (Water Purslane), *Class* vi. Hexandria, *Order* i. Monogyn.
 Perdícium, *Class* xix. Syngenesia, *Order* ii. Polyg. super.
 Perílla, *Class* xiv. Didynamia, *Order* i. Gymnosperma
 Períploca (Virginian Silk), *Class* v. Pentandria, *Order* ii. Digynia
 Pergulária, *Class* v. Pentandria, *Order* i. Monogynia
 Petésia, *Class* iv. Tetrandria, *Order* i. Monogynia
 Petivéria (Guinea-hen Weed), *Class* vi. Hexandria, *Order* iv.
 Tetragynia

- Petréa, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
- Peucedanum (Hog's Fennel, or Sulphur-wort), *Class* v. *Pentandria*, *Order* ii. *Digynia*
- Pezíza (Cup Mushroom), *Class* xxiv: *Cryptogamia*, *Order* iv. *Fungi*
- Pháca (Bastard Milk Vetch), *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
- Phálaris (Canary Grass), *Class* iii. *Triandria*, *Order* iii. *Trigynia*
- Phállus (Stink-horns), *Class* xxiv. *Cryptogamia*, *Order* iv. *Fungi*
- Pharnacéum, *Class* v. *Pentandria*, *Order* iii. *Trigynia*
- Phárus, *Class* xxi. *Monœcia*, *Order* vi. *Hexandria*
- Pháscum, *Class* xxiv. *Cryptogamia*, *Order* ii. *Musci*
- Phaséolus (Kidney Bean), *Class* xvii. *Diadelphia*, *Order* iv. *Decandria*
- Phellándrium, *Class* v. *Pentandria*, *Order* ii. *Digynia*
- Philadélphus (Mock Orange), *Class* xii. *Icosandria*, *Order* i. *Monogynia*
- Phyllýrea (Mock Privet), *Class* ii. *Diandria*, *Order* i. *Monogyn.*
- Phléum (Cat's-tail Grass), *Class* iii. *Triandria*, *Order* ii. *Digynia*
- Phlómis (Jerusalem Sage), *Class* xiv. *Didynamia*, *Order* i. *Gymnosperma*
- Phlóx (Lychnidea, or Bastard Lychnis), *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Phœnix (Common Palm, or Date Palmæ Tree)
- Phórmium, *Class* vi. *Hexandria*, *Order* i. *Monogynia*
- Phrýma, *Class* xiv. *Didynamia*, *Order* i. *Gymnosperma*
- Phýlica (Bastard Alaternus), *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Philiánthus (Sea-side Laurel), *Class* xxi. *Monœcia*, *Order* iii. *Triandria*
- Phylláchne, *Class* xxi. *Monœcia*, *Order* i. *Monandria*
- Phýllis (Bastard Hare's-ear), *Class* v. *Pentandria*, *Order* ii. *Digyn.*
- Phýsalis (Alkekengi, or Winter Cherry), *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Phytéuma (Rampions), *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Phytolácca (American Nightshade), *Class* x. *Decandria*, *Order* v. *Decagynia*

- Pícris, *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
 Pilulária (Pepper Grass), *Class* xxiv. Cryptogamia, *Order* i.
 Filices
 Pimpinélla (Burnet Saxifrage), *Class* v. Pentandria, *Order* ii. Di-
 gynia
 Píngúicula (Butter-wort), *Class* ii. Diandria, *Order* i. Monogyn.
 Pínus (Pine Tree), *Class* xxi. Monœcia, *Order* ix. Monadelphia
 Píper (Pepper), *Class* ii. Diandria, *Order* iii. Trigynia
 Piscídia, *Class* xvii. Diadelphia, *Order* iv. Decandria
 Pistácia (Pistacia Nut), *Class* xxii. Diœcia, *Order* v. Pentandria
 Pisónia (Fingrigo), *Class* xxiii. Polygamia, *Order* ii. Diœcia
 Pístia, *Class* xx. Gynandria, *Order* v. Hexandria
 Písum (Pea), *Class* xvii. Diadelphia, *Order* iv. Decandria
 Plantágo (Plantain), *Class* iv. Tetrandria, *Order* i. Monogynia
 Plátanus (Plane Tree), *Class* xxi. Monœcia, *Order* viii. Polyandria
 Plectrónia, *Class* v. Pentandria, *Order* i. Monogynia
 Plínia, *Class* xiii. Polyandria, *Order* i. Monogynia
 Plukenétia, *Class* xxi. Monœcia, *Order* ix. Monadelphia
 Plumbágo (Lead-wort), *Class* v. Pentandria, *Order* i. Monogynia
 Pluméria (Red Jasmine), *Class* v. Pentandria, *Order* i. Monogyn.
 Póa, *Class* iii. Triandria, *Order* ii. Digynia
 Podophýllum (Duck's-Foot, or May Apple), *Class* xiii. Polyandria,
 Order i. Monogynia
 Poinciána (Barbadoes Flower Fence), *Class* x. Decandria, *Or-*
 der i. Monogynia
 Polemónium (Greek Valerian), *Class* v. Pentandria, *Order* i. Mo-
 nogynia
 Polyánthes (Tuberose), *Class* vi. Hexandria, *Order* i. Monogynia
 Pollia, *Class* vi. Hexandria, *Order* i. Monogynia
 Polycárpon, *Class* iii. Triandria, *Order* iii. Trigynia
 Polycnémum, *Class* iii. Triandria, *Order* i. Monogynia
 Polýgala (Milk-wort), *Class* xvii. Diadelphia, *Order* iii. Octandria
 Polýgonum (Knot-grass), *Class* viii. Octandria, *Order* iii. Trigynia
 Polymnia, *Class* xix. Syngenesia, *Order* 0. Polyg. neces.
 Polypódium (Polypody), *Class* xxiv. Cryptogamia, *Order* i. Filices
 Polyprémum (Carolina Flax), *Class* iv. Tetrandria, *Order* i. Mo-
 nogynia

- Polytrichum* (Golden Maiden Hair), Cryptogamia, Class xxiv.
 Order ii. Musci
- Pommeréulla*, Class iii. Triandria, Order i. Monogynia
- Pontedéria*, Class vi. Hexandria, Order i. Monogynia
- Pópulus* (Poplar), Class xxii. Diœcia, Class vii. Octandria
- Porána*, Class v. Pentandria, Order i. Monogynia
- Porélla*, Class xxiv. Cryptogamia, Order ii. Musci
- Portlândia*, Class v. Pentandria, Order i. Monogynia
- Portuláca* (Purslane), Class xi. Dodecandria, Order i. Monogyn.
- Potamogéton* (Pond-weed), Class iv. Tetrandria, Order iii. Te-
 tragynia
- Potentilla* (Cinquefoil), Class xii. Icosandria, Order v. Polygynia
- Potérium* (Burnet), Class xxi. Monœcia, Order viii. Polyandria
- Póthos*, Class xx. Gynandria, Order ix. Polyandria
- Prásium* (Shrubby Hedge-Nettle), Class xiv. Didynamia, Order i.
 Gymnosperma
- Prenánthes* (Wild Lettuce), Class xix. Syngenesia, Order i. Polyg. æqua.
- Prémna*, Class xiv. Didynamia, Order ii. Angiosperma
- Prímula* (Primrose), Class v. Pentandria, Order i. Monogynia
- Prínos* (Winter Berry), Class vi. Hexandria, Order i. Monogyn.
- Próckia*, Class xviii. Polyandria, Order i. Monogynia
- Proserpináca*, Class iii. Triandria, Order iii. Trigynia
- Prosópis*, Class x. Decandria, Order i. Monogynia
- Protéa* (Silver Tree), Class iv. Tetrandria, Order i. Monogynia
- Prunélla* (Self-heal), Class xiv. Didynamia, Order i. Gymnosper.
- Prúnus* (Plum-tree), Class xii. Icosandria, Order i. Monogynia
- Psídium* (Guayava, or Bay Plum), Class xii. Icosandria, Order i.
 Monogynia
- Psorálea*, Class xvii. Diadelphia, Order ii. Decandria
- Psychótria*, Class v. Pentandria, Order i. Monogynia
- Ptélea* (Shrub Trefoil), Class iv. Tetrandria, Order i. Monogynia
- Ptéris* (Brakes, or Female Fern), Class xxiv. Cryptogamia, Or-
 der i. Filices
- Pterocárpus*, Class xvii. Diadelphia, Order iv. Decandria
- Pterónia*, Order xix. Syngenesia, Order i. Polyg. æqua.
- Pulmonária* (Lung-wort), Class v. Pentandria, Order i. Monogyn.

Púnica (Pomegranate), *Class xii. Icosandria, Order i. Monogyn.*
 Pýrola (Winter Green), *Class x. Decandria, Order i. Monogynia*
 Pýrus (Pear), *Class xii. Icosandria, Order i. Pentagynia*

Q

Qúassia, *Class x. Decandria, Order i. Monogynia*
 Qúercus (Oak), *Class xxi. Monœcia, Order viii. Polyandria*
 Quéria, *Class iv. Tetrandria, Order iii. Trigynia*
 Quisqúalis, *Class x. Decandria, Order i. Monogynia*

R

Rajánia, *Class xxii. Diœcia, Order vi. Hexandria*
 Rándia, *Class v. Pentandria, Order i. Monogynia*
 Ranúnculus (Crowfoot), *Class xiii. Polyandria, Order vii. Polygynia*
 Ráphanus (Radish), *Class xv. Tetradyndamia, Order i. Siliquosa*
 Rauvólfia, *Class v. Pentandria, Order i. Monogynia*
 Reaumúria, *Class xiii. Polyandria, Order v. Pentagynia*
 Reneálmia, *Class i. Monandria, Order i. Monogynia*
 Reséda (Bastard Rocket), *Class xi. Dodecandria, Order iii. Trigynia*
 Réstio, *Class xxii. Diœcia, Order iii. Triandria*
 Rétzia, *Class v. Pentandria, Order i. Monogynia*
 Rhacóma, *Class iv. Tetrandria, Order i. Monogýnia*
 Rhámnnus (Buckthorn), *Class v. Pentandria, Order i. Monogyn.*
 Rhéedia, *Class xiii. Polyandria, Order i. Monogynia*
 Rhéum (Rhubarb), *Class ix. Enneandria, Order ii. Trigynia*
 Rhéxia, *Class viii. Octandria, Order i. Monogynia*
 Rhinánthus (Elephant's Head), *Class xiv. Didynamia, Order ii. Angiosperma*
 Rhizóphora (Candle of the Indians), *Class xi. Dodecandria, Order i. Monogynia*
 Rhodiola (Rose Root), *Class xxii. Diœcia, Order vii. Octandria*
 Rhododéndron (Dwarf Rose-bay), *Class x. Decandria, Order i. Monogynia*
 Rhús (Sumach), *Class v. Pentandria, Order iii. Trigynia*

- Ríbes (Currant Tree), *Class v. Pentandria, Order i. Monogynia*
 Ríccia (Marsh Liver-wort), *Class xxiv. Cryptogamia, Order iii.*
 Algæ
- Richárdia, *Class vi. Hexandria, Order i. Monogynia*
 Ricinus (Palma Christi), *Class xxi. Monœcia, Order ix. Monadelphía*
- Ricótia, *Class xv. Tetradynamia, Order i. Siliquosa*
 Rivina, *Class iv. Tetrandria, Order i. Monogynia*
 Robínia (False Acacia), *Class xvii. Diadelphia, Order iv. Decandria*
- Roélla, *Class v. Pentandria, Order i. Monogynia*
 Rondelétia, *Class v. Pentandria, Order i. Monogynia*
 Rorídula, *Class v. Pentandria, Order i. Monogynia*
 Rósa (Rose), *Class xii. Icosandria, Order v. Polygynia*
 Rosmarínus (Rosemary), *Class ii. Diandria, Order i. Monogynia*
 Rótala, *Class iii. Triandria, Order i. Monogynia*
 Rottboélla, *Class iii. Triandria, Order ii. Digynia*
 Royéna (African Bladder Nut), *Class x. Decandria, Order ii. Digynia*
- Rúbia (Madder), *Class iv. Tetrandria, Order i. Monogynia*
 Rúbus (Raspberry), *Class xii. Icosandria, Order vi. Polygynia*
 Rudbeckia (Dwarf Sunflower), *Class xix. Syngenesia, Order iii. Polyg. frustr.*
 Ruéllia, *Class xiv. Didynamia, Order ii. Angiosperma*
 Rúmex (Dock), *Class vi. Hexandria, Order iii. Trigynia*
 Rúmphia, *Class iii. Triandria, Order i. Monogynia*
 Rúppia, *Class iv. Tetrandria, Order iii. Tetragynia*
 Rúscus (Knee Holly, or Butchers' Broom), *Class xxii. Diœcia, Order xiii. Syngenesia*
- Russélia, *Class v. Pentandria, Order ii. Digynia*
 Rúta (Rue), *Class x. Decandria, Order i. Monogynia*

S

- Sáccharum (Sugar Cane), *Class iii. Triandria, Order iii. Digynia*
 Sagina (Péarl-wort), *Class iv. Tetrandria, Order iii. Tetragynia*
 Sagittária (Arrow-head), *Class xxi. Monœcia, Order viii. Polyandria*
 Salácia, *Class xx. Gynandria, Order iii. Triandria*

- Salicórnía (Jointed Glass-wort), *Class* i. Monandria, *Order* i.
 Monogynia
- Sálix (Willow), *Class* xxii. Diœcia, *Order* ii. Diandria
- Salsóla (Glass-wort), *Class* v. Pentandria, *Order* ii. Digynia
- Salvadóra, *Class* iv. Tetrandria, *Order* iii. Tetragynia
- Sálvia (Sage), *Class* ii. Diandria, *Order* i. Monogynia
- Sámara, *Class* iv. Tetrandria, *Order* i. Monogynia
- Sambúcus (Elder), *Class* v. Pentandria, *Order* iii. Trigynia
- Sámolus (Round-leaved Water Pimpernel), *Class* v. Pentandria,
Order i. Monogynia
- Samýða, *Class* x. Decandria, *Order* i. Monogynia
- Sanguinária (Puccoon), *Class* xiii. Polyandria, *Order* i. Mono-
 gynia
- Sanguisórba (Greater Wild Burnet), *Class* iv. Tetrandria, *Order* i.
 Monogynia
- Sanícula (Sanicle), *Class* v. Pentandria, *Order* ii. Digynia
- Sántalum (Saunders), *Class* iv. Tetrandria, *Order* i. Monogynia
- Santolína (Lavender Cotton), *Class* xix. Syngenesia, *Order* i.
 Polyg. æqua.
- Sapíndus (Soap-berry), *Class* viii. Octandria, *Order* iii. Trigynia
- Saponária (Soap-wort), *Class* x. Decandria, *Order* ii. Digynia
- Saráca, *Class* xvii. Diadelphia, *Order* ii. Hexandria
- Sarracénia (Side-saddle Flower), *Class* xiii. Polyandria, *Order* i.
 Monogynia
- Saróthra (Bastard Gentian), *Class* v. Pentandria, *Order* iii. Trigyn.
- Saturéja (Savory), *Class* xiv. Didynamia, *Order* i. Gymnosperma
- Saurúrus (Lizard's Tail) *Class* vii. Heptandria, *Order* iii. Trigynia
- Satýrium (Lizard Flower), *Class* xx. Gynandria, *Order* ii. Diandr.
- Sauvagésia, *Class* v. Pentandria, *Order* i. Monogynia
- Saxífraga (Saxifrage), *Class* x. Decandria, *Order* ii. Digynia
- Scabiósa (Scabious), *Class* iv. Tetrandria, *Order* i. Monogynia
- Scabrita, *Class* iv. Tetrandria, *Order* i. Monogynia
- Scándix (Shepherd's Needle, or Venus's Comb), *Class* v. Pentan-
 dria, *Order* ii. Digynia
- Scævola, *Class* v. Pentandria, *Order* i. Monogynia
- Scheuchzéria (Lesser Flowering Rush), *Class* vi. Hexandria, *Or-*
der iii. Trigynia

- Scheffieldia, *Class* v. Pentandria, *Order* i. Monogynia
 Schínus (Indian Mastick), *Class* xxii. Diœcia, *Order* ix. Decandr.
 Schmedélia, *Class* viii. Octandria, *Order* ii. Digynia
 Schœnus (Bastard Cypress), *Class* iii. Triandria, *Order* i. Monogynia
 Schrebéra, *Class* v. Pentandria, *Order* ii. Digynia
 Schwálbea, *Class* xiv. Didynamia, *Order* ii. Angiosperma
 Schwénkia, *Class* ii. Diandria, *Order* i. Monogynia
 Scílla (Squill), *Class* vi. Hexandria, *Order* i. Monogynia
 Scírpus (Rush Grass), *Class* iii. Triandria, *Order* i. Monogynia
 Scleránthus (German Knot-grass, or Knawel), *Class* x. Decandria,
Order ii. Digynia
 Scólymus (Golden Thistle), *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
 Scopária, *Class* iv. Tetrandria, *Order* i. Monogynia
 Scopólia, *Class* xx. Gynandria, *Order* vi. Octandria
 Scorpiúrus (Caterpillars), *Class* xvii. Diadelphia, *Order* iv. Decandria
 Scorzonéra (Viper Grass), *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
 Scrophulária (Fig-wort), *Class* xiv. Didynamia, *Order* ii. Angiosperma
 Scutellária (Skull-cap), *Class* xiv. Didynamia, *Order* i. Gymnosperma
 Secále (Rye), *Class* iii. Triandria, *Order* ii. Digynia
 Securidáca, *Class* xvii. Diadelphia, *Order* iii. Octandria
 Sédum (Lesser Houseleek), *Class* x. Decandria, *Order* iv. Pentagynia
 Seguíéria, *Class* xiii. Polyandria, *Order* i. Monogynia
 Selágo, *Class* xiv. Didynamia, *Order* iii. Angiosperma
 Selínium (Milk Parsley), *Class* v. Pentandria, *Order* ii. Digynia
 Semecárpus, *Class* v. Pentandria, *Order* ii. Trigynia
 Sempervívum (Houseleek), *Class* xi. Dodecandria, *Order* v. Dodecagy.
 Senécio (Groundsel), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
 Séptas, *Class* vi. Heptandria, *Order* iv. Heptagynia

- Serápias (Helleborine), *Class xx. Gynandria, Order ii. Diandria*
 Seríola, *Class xix. Syngenesia, Order i. Polyg. æqua.*
 Seríphium, *Class xix. Syngenesia, Order i. Monogamia*
 Serpícula, *Class xxi. Monœcia, Order iv. Tetrandria*
 Serrátula (Saw-wort), *Class xix. Syngenesia, Order i. Polyg. æqua.*
 Sesámum (Oily Purging Grain), *Class xiv. Didynamia, Order ii. Angiosperma*
 Séseli (Hartwort of Marseilles), *Class v. Pentandria, Order ii. Digynia*
 Sesúvium, *Class xii. Icosandria, Order iii. Trigynia*
 Sheffiéldia
 Sherardia (Little Field Madder), *Class iv. Tetrandia, Order i. Monogynia*
 Sibbáldia, *Class v. Pentandria, Order v. Pentagynia*
 Sibthórpa, *Class xiv. Didynamia, Order ii. Angiosperma*
 Sícyos (Single-seeded Cucumber), *Class xxi. Monœcia, Order x. Syngenesia*
 Sída (Indian Mallow), *Class xvi. Monadelphia, Order vii. Polyandria*
 Sideritis (Iron-wort), *Class xiv. Didynamia, Order i. Gymnosp.*
 Sideróxylon (Iron-wood), *Class v. Pentandria, Order i. Monogyn.*
 Sigesbéckia, *Class xix. Syngenesia, Order i. Polyg. super.*
 Siléne (Viscous Champion), *Class x. Decandria, Order iii. Trigyn.*
 Sílphium (Bastard Chrysanthemum), *Class xix. Syngènesia, Order iv. Polyg. necess.*
 Sinápis (Mustard), *Class xv. Tetradynamia, Order i. Siliquosa*
 Siphonánthus, *Class iv. Tetrandria, Order i. Monogynia*
 Sírium, *Class iv. Tetrandria, Order i. Monogynia*
 Síson (Bastard Stone Parsley), *Class v. Pentandria, Order ii. Digynia*
 Sisýmbrium (Water Cresses), *Class xv. Tetradynamia, Order i. Siliquosa*
 Sisýrinchium (Bermudiana), *Class xx. Gynandria, Order ii. Trigynia*
 Sium (Water Parsnep), *Class v. Pentandria, Order ii. Digynia*
 Škimmia, *Class iv. Tetrandria, Order i. Monogynia*

- Sloanéa (Apeiba of Brasilians), *Class* xiii. Polyandria, *Order* i. Monogynia
- Smílax (Rough Bindweed), *Class* xxii. Diœcia, *Order* vi. Hexandria
- Smýrnum (Alexanders), *Class* v. Pentandria, *Order* ii. Digynia
- Solándra, *Class* xxiii. Polygamia, *Order* i. Monœcia
- Solánum (Nightshade), *Class* v. Pentandria, *Order* i. Monogynia
- Soldanélla (Soldanel), *Class* v. Pentandria, *Order* i. Monogynia
- Solidágo (Golden Rod), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
- Sónchús (Sow Thistle), *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
- Sonnerátia, *Class* xii. Icosandria, *Order* i. Monogynia
- Sophóra, *Class* x. Decandria, *Order* i. Monogynia
- Sórbus (Service Tree), *Class* xii. Icosandria, *Order* iii. Trigynia
- Spárgánium (Burr Reed), *Class* xxi. Monœcia, *Order* iii. Triandr.
- Sparrmánia, *Class* xiii. Polyandria, *Order* i. Monogynia
- Spártium (Broom), *Class* xiv. Diadelphia, *Order* iv. Decandria
- Spathélia, *Class* v. Pentandria, *Order* iii. Trigynia
- Spérgula (Spurrey), *Class* x. Decandria, *Order* iv. Pentagynia
- Spermacóce (Button Weed), *Class* iv. Tetrandria, *Order* i. Monogynia
- Spæránthus (Globe Flower), *Class* xix. Syngenesia, *Order* v. Polyg. segreg.
- Sphágnum (Bog-moss), *Class* xxiv. Cryptogamia, *Order* ii. Musci
- Spigélia (Worm-grass), *Class* v. Pentandria, *Order* i. Monogynia
- Spilánthus, *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
- Spinácia (Spinach), *Class* xxii. Diœcia, *Order* v. Pentandria
- Spínifex, *Class* xxiii. Polygamia, *Order* i. Monœcia
- Spiræa (Spiræa Frutex), *Class* xii. Icosandria, *Order* iv. Pentagyn.
- Spláchnum, *Class* xxiv. Cryptogamia, *Order* ii. Musci
- Spóndias (Brazilian Plum), *Class* x. Decandria, *Order* iv. Pentagynia
- Stáchys (Base Horehound), *Class* xiv. Didynamia, *Order* i. Gymnosperma
- Stæhelína, *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
- Stapélia, *Class* v. Pentandria, *Order* ii. Digynia

- Staphyléa (Bladder Nut), *Class v. Pentandria, Order iii. Trigynia*
 Státice (Thrift, or Sea Pink), *Class v. Pentandria, Order v. Pentagynia*
 Stellária (Great Chickweed), *Class x. Decandria, Order iii. Trigin.*
 Stelléra (German Groundsel), *Class viii. Octandria, Order i. Monogynia.*
 Stemódia, *Class xiv. Didynamia, Order ii. Angiosperma*
 Stercúlia, *Class xxi. Monœcia, Order ix. Monadelphia*
 Stéris, *Class v. Pentandria, Order ii. Digynia*
 Stéwartia, *Class xvi. Monadelphia, Order viii. Polyandria*
 Stipa (Feather-grass), *Class iii. Triandria, Order ii. Digynia*
 Stilágo, *Class xx. Gynandria, Order ii. Triandria*
 Stílbe, *Class xxiii. Polygamia, Order ii. Diœcia*
 Stillíngia, *Class xxi. Monœcia, Order ix. Monadelphia*
 Stœbe (Bastard Æthiopian Elichrysum), *Class xix. Syngenesia, Order v. Polyg. segreg.*
 Stratiótes (Water Soldier), *Class xiii. Polyandria, Order vi. Hexagynia*
 Struthíola, *Class iv. Tetrandria, Order i. Monogynia*
 Strúmpfia, *Class xix. Syngenesia, Order vi. Monogynia*
 Strýchnos, *Class v. Pentandria, Order i. Monogynia*
 Stýrax (Storax Tree), *Class xi. Dodecandria, Order i. Monogyn.*
 Subulária (Rough-leaved Alysson), *Class xv. Tetradynamia, Order ii. Siliculosa*
 Suriána, *Class x. Decandria, Order iv. Pentagynia*
 Swértia (Marsh Gentian), *Class v. Pentandria, Order ii. Digyn.*
 Symphónia, *Class xvi. Monadelphia, Order ii. Pentandria*
 Sýmphytum (Comphrey), *Class v. Pentandria, Order i. Monogynia*
 Sýmplocas, *Class xviii. Polyadelphia, Order iii. Polyandria*
 Syrínga (Lilac), *Class ii. Diandria, Order i. Monogynia*
 Swieténia (Mahogany Tree), *Class x. Decandria, Order i. Monogynia*

T

- Tabernæmontána, *Class x. Decandria, Order i. Monogynia*
 Tácca, *Class xi. Dodecandria, Order iii. Trigynia*
 Tagétes (African Marygold), *Class xix. Syngenesia, Order ii. Polyg. super.*

- Tamaríndus (Tamarind Tree), *Class* iii. Triandria, *Order* i. Monogynia
- Támarix (Tamarisk), *Class* v. Pentandria, *Order* iii. Trigynia
- Támus (Black Bryony), *Class* xxii. Diœcia, *Order* vi. Hexandria
- Tanacétum (Tansey), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
- Tarchonánthus (Shrubby African Fleabane), *Class* xix. Syngenesia, *Order* i. Polyg. æqua.
- Targiónia, *Class* xxiv. Cryptogamia, *Order* iii. Algæ
- Táxus (Yew Tree), *Class* xxii. Diœcia, *Order* xii. Monadelphia
- Téctona, *Class* v. Pentandria, *Order* i. Monogynia
- Teléphiúm (True Orpine), *Class* v. Pentandria, *Order* iii. Trigynia
- Terminália, *Class* xxiii. Polygamia, *Order* i. Monœcia
- Ternstrómia, *Class* xiii. Polyandria, *Order* i. Monogynia
- Tetráçera, *Class* xiii. Polyandria, *Order* iii. Trigynia
- Tetragónia, *Class* xii. Icosandria, *Order* iv. Penægynia
- Teúcrium (Germander), *Class* xiv. Didynamia, *Order* i. Gymnosperma
- Thalía, *Class* i. Monandria, *Order* i. Monogynia
- Thalíctruni (Meadow Rue), *Class* xiii. Polyandria, *Order* vii. Polygynia
- Thápsia (Deadly Carrot, or Scorching Fennel), *Class* v. Pentandria, *Order* ii. Digynia.
- Théa (Tea Tree), *Class* xiii. Polyandria, *Order* i. Monogynia
- Theligonum (Dog's Cabbage), *Class* xxi. Monœcia, *Order* viii. Polyandria
- Theobróma (Chocolate Nut), *Class* xvii. Polyadelphia, *Order* i. Pentandria
- Theophrásta, *Class* v. Pentandria, *Order* i. Monogynia
- Thésium (Bastard Toad Flax), *Class* v. Pentandria, *Order* i. Monogynia
- Thláspi (Mithridate Mustard, or Treacle Mustard), *Class* xv. Tetradynamia, *Order* ii. Siliculosa
- Thouínia, *Class* ii. Diandria, *Order* i. Monogynia
- Thyrállis, *Class* x. Decandria, *Order* i. Monogynia
- Thúja (Arbor Vitæ), *Class* xxi. Monœcia, *Order* ix. Monadelph.
- Thunbérgia, *Class* xiv. Didynamia, *Order* ii. Angiosperma

- Thýmbra (Mountain Hyssop), *Class* xiv. *Didynamia*, *Order* i.
Gymnosperma
- Thýnfus (Thyme), *Class* xiv. *Didynamia*, *Order* i. *Gymnosperma*
- Tiarélla, *Class* x. *Decandria*, *Order* ii. *Digynia*
- Tília (Lime Tree), *Class* xiii. *Polyandria*, *Order* i. *Monogynia*.
- Tillæa (Small Annual Houseleek), *Class* xiv. *Tetrandria*, *Order* i.
Monogynia
- Tillándsia, *Class* vi. *Hexandria*, *Order* i. *Monogynia*
- Tínus, *Class* ix. *Enneandria*, *Order* i. *Monogynia*
- Toluífera (Balsam of Tolu Tree), *Class* x. *Decandria*, *Order* i.
Monogynia
- Tomex, *Class* iv. *Tetrandria*, *Order* i. *Monogynia*
- Tordýlium (Hartwort of Crete), *Class* v. *Pentandria*, *Order* ii.
Digynia
- Torénia, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
- Tormentílla (Tormentil), *Class* xii. *Icosandria*, *Order* v. *Polygynia*
- Tournefórtia, *Class* v. *Pentandria*, *Order* i. *Monogynia*
- Tózzia, *Class* xiv. *Didynamia*, *Order* ii. *Angiosperma*
- Trachélium (Umbelliferous Throat-wort), *Class* v. *Pentandria*,
Order i. *Monogynia*
- Tradescántia (Virginian Spider-wort), *Class* vi. *Hexandria*, *Order* i. *Monogynia*
- Trágia, *Class* xxi. *Monœcia*, *Order* iii. *Triandria*
- Tragopógon (Goat's Beard), *Class* xix. *Syngenesia*, *Order* i. *Polyg. æqua*.
- Trápa (Water Caltrops), *Class* iv. *Tetrandria*, *Order* i. *Monogyn.*
- Tremélla, *Class* xxiv. *Cryptogamia*, *Order* iii. *Algæ*
- Tréwia, *Class* xiii. *Polyandria*, *Order* i. *Monogynia*
- Triánthema (Horse Purslane), *Class* x. *Decandria*, *Order* i. *Monogynia*
- Tríbulus (Caltrops), *Class* x. *Decandria*, *Order* i. *Monogynia*
- Tríchilia, *Class* x. *Decandria*, *Order* i. *Monogynia*
- Trichománes, *Class* xxiv. *Cryptogamia*, *Order* i. *Filices*
- Trichosánthes (Serpent Cucumber), *Class* xxi. *Monœcia*, *Order* x.
Syngenesia
- Trichostéma, *Class* xiv. *Didynamia*, *Order* i. *Gymnosperma*

- Trídax (Trailing Starwort of Vera Cruz), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
- Trientális (Winter-green, with Chickweed Flowers), *Class* vii. Heptandria, *Order* i. Monogynia
- Trifólium (Trefoil), *Class* xvii. Diadelphia, *Order* iv. Decandria
- Triglóchin (Arrow-headed Grass), *Class* vi. Hexandria, *Order* iii. Trigynia
- Trigonélla (Fenugreek), *Class* xvii. Diadelphia, *Order* iv. Decandria
- Tríllium (Herb Truelove of Canada), *Class* vi. Hexandria, *Order* iii. Trigynia
- Trílix, *Class* xiii. Polyandria, *Order* i. Monogynia
- Triópteris, *Class* x. Decandria, *Order* iii. Trigynia
- Triósteum (Fever-root, or False Ipecacuana), *Class* v. Pentandria, *Order* i. Monogynia
- Tripláris, *Class* iii. Triandria, *Order* iii. Trigynia
- Trípsacum, *Class* xxi. Monœcia, *Order* iii. Triandria
- Tríticum (Wheat), *Class* iii. Triandria, *Order* iii. Digynia
- Triumfétta, *Class* xi. Dodecandria, *Order* i. Monogynia
- Tróllius (Globe Ranunculus), *Class* xiii. Polyandria, *Order* vii. Polygynia
- Tropæolum (Indian Cress), *Class* viii. Octandria, *Order* i. Monogynia
- Tróphis, *Class* xxii. Diœcia, *Order* iv. Tetrandria
- Tulbágia, *Class* vi. Hexandria, *Order* i. Monogynia
- Túlipa (Tulip), *Class* vi. Hexandria, *Order* i. Monogynia
- Turnéra, *Class* v. Pentandria, *Order* iii. Trigynia
- Turræa, *Class* x. Decandria, *Order* i. Monogynia
- Turrítis (Tower Mustard), *Class* xv. Tetradinamia, *Order* i. Siliquosa
- Tussilágo (Colt's Foot), *Class* xix. Syngenesia, *Order* ii. Polyg. super.
- Týpha (Cat's-tail, or Reedmace), *Class* xxi. Monœcia, *Order* iii. Triandria
- V
- Vaccínium (Whortle Berry), *Class* viii. Octandria. *Order* i. Monogynia

- Vahlia*, *Class v.* Pentandria, *Order ii.* Digynia
Valániu (Crosswort), *Class xxiii.* Polygamia, *Order i.* Monœcia
Valeriána (Vaterian), *Class iii.* Triandria, *Order i.* Monogynia
Válbea, *Class xiii.* Polyandria, *Order i.* Monogynia
Vallisnéria, *Class xxii.* Diœcia, *Order ii.* Diandria
Vandéllia, *Class xiv.* Didynamia, *Order ii.* Angiosperma
Varrónia, *Class v.* Pentandria, *Order i.* Monogynia
Vatéria, *Class xiii.* Polyandria, *Order i.* Monogynia
Vática, *Class xi.* Dodecandria, *Order i.* Monogynia
Valézia, *Class vi.* Hexandria, *Order ii.* Digynia
Vélla (Spanish Cress), *Class xv.* Tetradynamia, *Order ii.* Sili-
 culosa
Verátrum (White Hellebore), *Class xxiii.* Polygamia, *Order i.*
 Monœcia
Verbáscum (Mullein), *Class v.* Pentandria, *Order i.* Monogynia
Verbéna (Vervain), *Class ii.* Diandria, *Order i.* Monogynia
Verbesúna, *Class xix.* Syngenesia, *Order ii.* Polyg. super.
Verónica (Speedwell), *Class ii.* Diandria, *Order i.* Monogynia
Vibúrnum (Pliant Mealy Tree, or Wayfaring Tree), *Class v.*
 Pentandria, *Order iii.* Trigynia
Vícia (Vetch), *Class xvii.* Diadelphia, *Order iv.* Decandria
Víncia (Periwinkle), *Class v.* Pentandria, *Order i.* Monogynia
Viola (Violet), *Class xix.* Syngenesia, *Order i.* Monogynia
Vírécta, *Class v.* Pentandria, *Order i.* Monogynia
Viscum (Mistletoe), *Class xxii.* Diœcia, *Order iv.* Tetrandria
Visnéa, *Class xi.* Dodecandria, *Order iii.* Trigynia
Vítex (Agnus Castus, or Chaste Tree), *Class xiv.* Didynamia,
Order ii. Angiosperma .
Vítis (Vine), *Class v.* Pentandria, *Order i.* Monogynia
Volkaméria, *Class xiv.* Didynamia, *Order ii.* Angiosperma
Ulex (Furze, Whins, or Gorfs), *Class vii.* Diadelphia, *Order iv.*
 Decandria
Ulmus (Elm Tree), *Class v.* Pentandria, *Order ii.* Digynia
Ulva (Laver), *Class xxiv.* Cryptogamia, *Order iii.* Algæ
Uníola (Sea-side Oats of Carolina), *Class iii.* Tirandira, *Order ii.*
 Digynia
Unóna, *Class xiii.* Polyandria, *Order vii.* Polygynia

Uréna (Indian Mallow), *Class* xvi. Monadelphia, *Order* vii. Polyandria.

Unxia, *Class* xix. Syngenes. *Order* ii. Polyg. superfl.

Urtica (Nettle), *Class* xxi. Monœcia, *Order* iv Tetrandria

Utriculária (Water Milfoil), *Class* x. Decandria, *Order* i. Monogynia

Uvária, *Class* xiii. Polyandria, *Order* vii. Polygynia

Uvulária, *Class* vi. Hexandria, *Order* i. Monogynia

W

Wachendórfia *Class* iii. Triandria, *Order* i. Monogynia

Walthéria, *Class* xvi. Monadelphia. *Order* ii. Pentandria

Weigéla, *Class* v. Pentandria, *Order* i. Monogynia

Weinmánnia, *Class* viii. Octandria, *Order* ii. Digynia

Willíchia, *Class* iii. Triandria, *Order* i. Monogynia

Winteránia

Wintéra, *Class* xiii. Pólyandria, *Order* vii. Polygynia

Witsénia, *Class* iii. Triandria, *Order* i. Monogynia

Wulfénia, *Class* ii. Diandria, *Order* i. Monogynia

Wurbéa, *Class* vi. Hexandria, *Order* iii. Trigynia

X

Xánthium (Lesser Burdock), *Class* xxi. Monœcia, *Order* v. Pentandria

Xeránthemum (Austrian sneezewort, or Eternal Flower), *Class* xix. Syngenesia, *Order* ii. Polygamia superflua

Ximénia, *Class* viii. Octandria, *Order* i. Monogynia.

Xylophýlla, *Class* v. Pentandria, *Order* iii. Trigynia

Xylóphia, *Class* xx. Gynandria, *Order* ix. Polyandria

Xýris, *Class* iv. Tetrandria, *Order* i. Monogynia.

Y

Yúcca (Adam's Needle), *Class* vi. Hexandria, *Order* i. Monogynia

Z

Zámia, *Class* xxiv. Cryptogamia, *Order* i. Filices.

- Zanichéllia (Triple-headed Pond-weed), *Class* xxi. Monœcia,
Order i. Monandria
- Zanónia, *Class* xxii. Diœcia, *Order* v. Pentandria
- Zanthóxylum (Tooth-ach Tree), *Class* xxii. Diœcia, *Order* v.
Pentandria
- Zea (Indian, or Turkey-wheat), *Class* xxi. Monœcia, *Order* iii.
Triandria
- Zinnia, *Class* xix. Syngenes. *Order* ii. Polyg. super.
- Zizânia, *Class* xxi. Monœcia, *Order* vi. Hexandria
- Zizíphora (Syrian Field Basil), *Class* ii. Diandria, *Order* i. Mo-
nogynia
- Zœgea, *Class* xix. Syngenesia, *Order* iii. Polyg. frustr.
- Zostéra (Grass-wrack), *Class* xx. Gynandria, *Order* ix. Polyand.
- Zygophýllum (Bean Caper), *Class* x. Decandria, *Order* i. Mono-
gynia

TABLE IV.



AN

ALPHABETICAL CATALOGUE

OF

ENGLISH AND SCOTCH NAMES

OF

PLANTS,

FROM THE MOST APPROVED AUTHORS,

Referred to their respective Genera.



A

Abele, <i>Populus</i>	Agnus castus, <i>Vitex</i>
Abelmosk, <i>Hibiscus</i>	Agrimony, <i>Agrimonia</i>
Acacia, <i>Mimosa</i>	Agrimony, Hemp, <i>Eupatorium</i>
Acacia, false, <i>Robinia</i>	Agrimony, Bastard Hemp, <i>Age-</i>
Acacia, German, <i>Prunus</i>	<i>ratum</i>
Acacia, three-thorned, <i>Gleditsia</i>	Agrimony, Naked-headed
Acajou, <i>Anacardium</i>	Hemp, <i>Verbesina</i>
Aconite, <i>Aconitum</i>	Agrimony, Water Hemp, <i>Bidens</i>
Aconite, Winter, <i>Helleborus</i>	Ague Tree, <i>Laurus</i>
Adam's Apple, <i>Citrus</i>	Aikraw, <i>Lichen scrob.</i>
Adam's Needle, <i>Yucca</i>	Alaternus, Bastard, <i>Phylica</i>
Adder's Wort, <i>Polygonum</i>	Alder, <i>Betula</i>
Adder's Tongue, <i>Ophioglossum</i>	Alder, Black, or Berry-bearing
Adragant, Gum, see <i>Tragacanth</i>	<i>Rhamnus</i>
Agaric, <i>Agaricus</i>	Ale-cost, <i>Tanacetum</i>

Ale-hoof, <i>Glechoma</i>	Apple, Custard, <i>Annona</i>
Alexanders, <i>Smyrniun</i>	Apple, Love, <i>Solanum</i>
Alkali, <i>Salicornia</i>	Apple, Mad, <i>Solanum</i>
Alkanet, <i>Lithospermum</i>	Apple, Male Balsam, <i>Momordica</i>
Alkekengi, <i>Physalis</i>	Apple, May, <i>Podophyllum</i>
All-good, <i>Chenopodium</i>	Apple, Pine, <i>Bromelia</i>
All-heal, Clowns, <i>Stachys</i>	Apple, Purple, <i>Annona</i>
All-heal, Hercules's, <i>Pastinaca</i>	Apple, Soap, <i>Sapindus</i>
All-heal, Hercules's, <i>Heracleum</i>	Apple, Sour, <i>Annona</i>
All-seed, <i>Linum</i>	Apple, Star, <i>Chrysophyllum</i>
All-spice, <i>Myrtus</i>	Apple, Sugar, <i>Annona</i>
Alligator Pear, <i>Laurus</i>	Apple, Sweet, <i>Annona</i>
Almond, <i>Amygdalus</i>	Apple, Thorn, <i>Datura</i>
Almond, African, <i>Brabejum</i>	Apple, Water, <i>Annona</i>
Almond, Ethiopian, <i>Brabejum</i>	Apricot, <i>Prunus</i>
Aloe, American, <i>Agave</i>	Arbor Vitæ, <i>Thuya</i>
Aloe, Water, <i>Stratiotes</i>	Arbutus Trailing, <i>Epigæa</i>
Althæa frutex, <i>Hibiscus</i>	Arcel, <i>Lichen omph.</i>
Alysson, Rough-leaved, <i>Subularia</i>	Ar-nuts, <i>Avena elet</i>
Amaranth, <i>Amaranthus</i>	Archangel, <i>Lamium</i>
Amaranth, Globe, <i>Gomphrena</i>	Archangel, Baulm-leav'd, <i>Melittis</i>
Amber Tree, <i>Anthospermum</i>	Archangel, Yellow, <i>Galeopsis</i>
Amellus of Virgil, <i>Aster</i>	Arrowhead, <i>Sagittaria</i>
Amomum Plinii, <i>Solanum</i>	Arrow-headed Grass, <i>Triglochin</i>
Amomum, German, <i>Sison</i>	Arrow-root, Indian, <i>Maranta</i>
Ananas, <i>Bromelia</i>	Arse-smart, <i>Polygonum</i>
Angelica, Berry-bearing, <i>Aralia</i>	Artichoke, <i>Cynara</i>
Angelica, Wild, <i>Ægopodium</i>	Artichoke, Jerusalem, <i>Helianthus</i>
Angelica Tree, <i>Aralia</i>	Arum, African, <i>Calla</i>
Anife, <i>Pimpinella</i>	Arum, Floating, <i>Orontium</i>
Anotta, <i>Bixa</i>	Asarabacca, <i>Asarum</i>
Apeiba of the Brasilians, <i>Sloanea</i>	Ash, <i>Fraxinus</i>
Apple, <i>Pyrus</i>	Ash, Mountain, <i>Sorbus</i>
Apple, Adam's, <i>Citrus</i>	Ash, Poison, <i>Rhus</i>
Apple, Blad, <i>Cactus</i>	Ash-weed, <i>Ægopodium</i>

Asparagus, Climbing, African, Bark, <i>Ilathera</i> , <i>Clusia</i>	Bark, Winter's, <i>Laurus</i>
<i>Medeola</i>	Barley, <i>Hordeum</i>
Asp, or Aspen Tree, <i>Populus</i>	Barren-wort, <i>Epimedium</i>
Asphodel, <i>Asphodelus</i>	Base-tree Trefoil, <i>Cytisus</i>
Asphodel, African, <i>Anthericum</i>	Basil, <i>Ocimum</i>
Asphodel, Lily, <i>Hemerocallis</i>	Basil, Field, <i>Clinopodium</i>
Asphodel, Lily, <i>Crinum</i>	Basil, American Field, <i>Monarda</i>
Asses Cucumber, <i>Momordica</i>	Basil, Syrian Field, <i>Ziziphora</i>
Atamasco Lily, <i>Amaryllis</i>	Basil, Stone, <i>Thymus</i>
Avens, <i>Geum</i>	Basil, Wild, <i>Thymus</i>
Avocado Pear, <i>Laurus</i>	Batchelor's Buttons, <i>Lychnis</i>
Auricula, <i>Primula</i>	Batchelor's Pear, <i>Solanum</i>
Auricula, Borrage-leaved, <i>Verbascum</i>	Baulm, <i>Melissa</i>
Ax-vetch. See Hatchet-vetch	Baulm, Bastard, <i>Melittis</i>
Azarole, <i>Crataegus</i>	Baulm, Moldavian, <i>Dracocephalum</i>
Azerira, <i>Prunus</i>	Baulm, Molucca, <i>Moluccella</i>
	Baulm, Turkey, <i>Dracocephalum</i>
B	Bay, <i>Laurus</i>
Balaustine, <i>Punica</i>	Bay, Loblolly, <i>Gordonia</i>
Balm, see Baulm	Bay, Rose, <i>Nerium</i>
Balm of Gilead, false, <i>Dracocephalon</i>	Bay, Dwarf Rose, <i>Rhododendrum</i>
Balsam, <i>Impatiens</i>	Bay, Mountain Rose, <i>Rhododendrum</i>
Balsam of Tolu, <i>Toluifera</i>	Bay, Sweet-flowering, <i>Magnolia</i>
Balsam Apple, Male, <i>Momordica</i>	Bay Plumb, <i>Psidium</i>
Balsam Tree, <i>Clusia</i>	Bead Tree, <i>Melia</i>
Balsam Tree, <i>Pistacia</i>	Bean, <i>Vicia</i>
Balsam Tree, <i>Copaifera</i>	Bean, Bog, <i>Menyanthes</i>
Balsamine, Female, <i>Impatiens</i>	Bean, white, <i>Crataegus</i>
Bamboo Cane, <i>Arundo</i>	Bean, Kidney, <i>Phaseolus</i>
Banana, <i>Musa</i>	Bean Tree, Kidney, <i>Glycine</i>
Bane-berries, <i>Actæa</i>	Bean Tree of America, <i>Erythrina</i>
Baniam Tree, <i>Ficus</i>	Bean Tree, Binding, <i>Mimosa</i>
Bark, True Jesuit's, <i>Cinchona</i>	Bean Caper, <i>Zygophyllum</i>
Bark, False Jesuit's, <i>Iva</i>	

Bean Trefoil, <i>Cytisus</i>	Bird's Eye, <i>Adonis</i>
Bean Trefoil, Stinking, <i>Anagyris</i>	Bird's Foot, <i>Ornithopus</i>
Bear-berries, <i>Arbutus</i>	Bird's Foot Trefoil, <i>Lotus</i>
Bear-bind, <i>Convolvulus</i>	Bird's Nest, <i>Ophrys</i>
Bear's-breach, <i>Acanthus</i>	Bird's Nest, Purple, <i>Orchis</i>
Bear's-ear, <i>Primula</i>	Birch, <i>Betula</i>
Bear's-ear Sanicle, <i>Cortusa</i>	Birth-wort, <i>Aristolochia</i>
Bear's-foot, <i>Helleborus</i>	Bishop's-weed, <i>Ammi</i>
Beard, Old Man's, <i>Clematis</i>	Bistort, <i>Polygonum</i>
Beech, <i>Fagus</i>	Bitter-gourd, <i>Cucumis</i>
Beet, <i>Beta</i>	Bitter-sweet, <i>Solanum</i>
Bee-flower, <i>Ophrys</i>	Bitter-vetch, <i>Ervum</i>
Behen, White, <i>Cucubalus</i>	Bitter-vetch, <i>Orobus</i>
Bell-flower, <i>Campanula</i>	Bitter-vetch, Jointed podded,
Bells Canterbury, <i>Campanula</i>	<i>Ervum</i>
Bed-straw, <i>Galium</i>	Bitter-wort, <i>Gentiana</i>
Bell-Pepper, <i>Capsicum</i>	Blackberry, <i>Rubus</i>
Belladonna Lily, <i>Amaryllis</i>	Blad Apple, <i>Cactus</i>
Belvidere, <i>Chenopodium</i>	Bladder Nut, <i>Staphylæa</i>
Bellyach-weed, <i>Jatropha</i>	Bladder Nut, African, <i>Royena</i>
Benjamin Tree, <i>Laurus</i>	Bladder Nut, Laurel-leaved, <i>Ilex</i>
Bennet, Herb, <i>Geum</i>	Bladder Senna, <i>Colutea</i>
Berberry, <i>Berberis</i>	Bladder Senna, Jointed podded,
Bermudiana, <i>Sisyrinchium</i>	<i>Coronilla</i>
Betony, <i>Betonica</i>	Blessed Thistle, <i>Cnicus</i>
Betony, Paul's, <i>Veronica</i>	Blindman's Ball, <i>Lycoperdon bor.</i>
Betony, Water, <i>Scrophularia</i>	Blinks, <i>Montia</i>
Big, <i>Hordeum</i>	Blite, <i>Blitum</i>
Bilberry, <i>Vaccinium</i>	Blite, <i>Amaranthus</i>
Bindweed, <i>Convolvulus</i>	Blood-flower, <i>Hæmanthus</i>
Bindweed, Black, <i>Tamus</i>	Blood-wood, <i>Hæmantoxylon</i>
Bindweed, Rough, <i>Smilax</i>	Blood-wort, <i>Rumex</i>
Birch, <i>Betula</i>	Blue-bonnets, <i>Centaurea cyan</i>
Birch of Jamaica, <i>Pistacia</i>	Blue-bottle, <i>Centaurea</i>
Bird-cherry, <i>Prunus</i>	Bogbean, <i>Menyanthes</i>
Bird Pepper, <i>Capsicum</i>	Bogberries, <i>Vaccinium</i>

Bogwhorts, <i>Vaccinium</i>	Buckler, Mustard, <i>Biscutiella</i>
Bonduc, <i>Guilandina</i>	Buck's-horn Plantain, <i>Plantago</i>
Bonnet Pepper, <i>Capsicum</i>	Buck's-horn, Warded, <i>Cochlearia</i>
Bore-cole, <i>Brassica</i>	Buck-thorn, <i>Rhamnus</i>
Borage, <i>Borrage</i>	Buck-thorn, Sea, <i>Hippophaë</i>
Bottle-flower, <i>Centaurea</i>	Buck-wheat, <i>Polygonum</i>
Box, <i>Buxus</i>	Bugbane. See Bogbean
Box, African, <i>Myrsine</i>	Bugle, <i>Ajuga</i>
Box, Low, <i>Polygala</i>	Bugloss, <i>Achusa</i>
Boxthorn, <i>Lycium</i>	Bugloss, Small wild, <i>Asperugo</i>
Brackens, <i>Pteris</i>	Bugloss, Viper's, <i>Echium</i>
Brakes, <i>Pteris</i>	Bullace Tree, <i>Chrysophyllum</i>
Bramble, <i>Rubus</i>	Bullace Tree, <i>Prunus</i>
Brank, <i>Polygonum</i>	Burdock, <i>Arctium</i>
Brank Ursine, <i>Acanthus</i>	Burdock, Lesser, <i>Xanthium</i>
Brasletto, <i>Cesalpina</i>	Bur-Marygold, <i>Bidens</i>
Break-stone, <i>Saxifraga</i>	Burnet, Garden, <i>Poterium</i>
Break-stone, Parsley, <i>Aphanès</i>	Burnet, Greater wild, <i>Sanguisorba</i>
Briar, Sweet, <i>Rosa</i>	Burnet Saxifrage, <i>Pimpinella</i>
Briar, Wild, <i>Rosa</i>	Burning Thorny Plant, <i>Euphorbia</i>
Brimstone-wort, <i>Peucedanum</i>	Bur Reed, <i>Sparganium</i>
Bristol, Flower of, <i>Lychnis</i>	Butcher's Broom, <i>Ruscus</i>
Broccoli, <i>Brassica</i>	Butter Burr, <i>Tussilago</i>
Brooklime, <i>Veronica</i>	Butter-cups, <i>Ranunculus</i>
Broom, <i>Spartium</i>	Butter-wort, <i>Pinguicula</i>
Broom, African, <i>Aspalathus</i>	Button Tree, <i>Conocarpus</i>
Broom, Dyer's, <i>Genista</i>	Button Weed, <i>Spermacoce</i>
Broom, Dwarf, <i>Genista</i>	Button Wood, <i>Cephalanthus</i>
Broom, Single-seeded, <i>Genista</i>	
Broom, Rape, <i>Orobauche</i>	C
Broom, Rape, with great Purple Flowers, <i>Lathræa</i>	Cabbage, <i>Brassica</i>
Brown-wort, <i>Scrophularia</i>	Cabbage, Dog's, <i>Theligonum</i>
Brown-wort, <i>Prunella</i>	Cabbage, Sea, <i>Crambe</i>
Bryony, <i>Bryonia</i>	Cabbage Tree, <i>Cacalia</i>
Bryony, Black, <i>Tamus</i>	Calabash, <i>Cucurbita</i>
	Calabash Tree, <i>Crescentia</i>

Calamint, <i>Melissa</i>	Cashew-nut, <i>Anacardium</i>
Calamint, Water, <i>Mentha</i>	Cassava, <i>Jatropha</i>
Cale, <i>Brassica</i>	Cassia, Poet's, <i>Osyris</i>
Cale, Sea, <i>Crambe</i>	Cassidony, <i>Gnaphalium</i>
Caltrops, <i>Tribulus</i>	Cassiobury Bush, <i>Cassine</i>
Caltrops, Water, <i>Trapa</i>	Catchfly, <i>Silene</i>
Calve's Snout, <i>Antirrhinum</i>	Catmint, <i>Nepeta</i>
Cammock, <i>Ononis</i>	Cat's-foot, <i>Glechoma</i>
Campeachy Wood, <i>Hæmatoxylon</i>	Cat's-foot Mountain, <i>Gnaphalium</i>
Camphire Tree, <i>Laurus</i>	Cat's-tail, <i>Typha</i>
Campion, <i>Angrostemma</i>	Caterpillars, <i>Scorpiurus</i>
Campion, <i>Lychnis</i>	Cauliflower, <i>Brassica</i>
Campion, Viscous, <i>Silene</i>	Cedar, <i>Juniperus</i>
Canary-grass, <i>Phalaris</i>	Cedar of Jamaica, Bastard, <i>Theo-</i>
Candle of the Indians. See Kandel	<i>broma</i>
Candleberry Myrtle, <i>Myrica</i>	Cedar, White, <i>Cupressus</i>
Candy Carrot, <i>Athamanta</i>	Cedar of Busaco, <i>Cupressus</i>
Candy Lion's Foot, <i>Catananche</i>	Cedar of Libanus, <i>Pinus</i>
Candy Tuft, <i>Iberis</i>	Celandine, <i>Chelidonium</i>
Candy Tuft Tree, <i>Iberis</i>	Celandine, Lesser, <i>Ranunculus</i>
Cane or Reed, <i>Arundo</i>	Celandine Tree, <i>Bocconia</i>
Cane, Sugar, <i>Saccharum</i>	Celeriac, <i>Apium</i>
Canterbury Bells, <i>Campanula</i>	Celery, <i>Apium</i>
Caper-Bush, <i>Capparis</i>	Céntaury, <i>Centaurea</i>
Caper, Bean, <i>Zygophyllum</i>	Centaury, Lesser, <i>Gentiana</i>
Caraway, <i>Carum</i>	Ceterach, <i>Asplenium</i>
Cardinal-flower, <i>Lobelia</i>	Chamomile, <i>Anthemis</i>
Carline Thistle, <i>Carlina</i>	Champignon, <i>Agaricuscam</i>
Carnation, <i>Dianthus</i>	Chardon, <i>Cynara</i>
Carnation, Spanish, <i>Poinciana</i>	Charlock, <i>Sinapi</i>
Carnation Tree, <i>Cacalia</i>	Charlock, White-flowered, with
Carob Tree, <i>Ceratonia</i>	jointed Pods, <i>Raphanus</i>
Carrot, <i>Daucus</i>	Chaste Tree, <i>Vitex</i>
Carrot, Candy, <i>Athamanta</i>	Cheese Rennet, <i>Galium</i>
Carrot, Deadly, <i>Thapsia</i>	Cherry, <i>Prunus</i>
Carui, <i>Carum</i>	Cherry, Barbadoes, <i>Malpighia</i>

Cherry, Bird, <i>Prunus</i>	Ciboules, <i>Allium</i>
Cherry, Cornelian, <i>Cornus</i>	Cicely, Sweet, <i>Scandix</i>
Cherry, Dwarf, <i>Lonicera</i>	Cinnamon Tree, <i>Laurus</i>
Cherry, Hottentot, <i>Cassine</i>	Cinnamon, White, <i>Laurus</i>
Cherry, Winter, <i>Physalis</i>	Cinquefoil, <i>Potentilla</i>
Cherry, Winter, <i>Solanum</i>	Cinquefoil, Marsh, <i>Comarum</i>
Cherry of the Alps, <i>Lonicera</i>	Cistus, Marsh, <i>Ledum</i>
Cherry Laurel, <i>Prunus</i>	Cistus, Lesser Marsh, <i>Andromeda</i>
Chervil, Garden, <i>Scandix</i>	Cistus, Nettle-leaved, <i>Turnera</i>
Chervil, Wild, <i>Cherophyllum</i>	Cistus, Rape of, <i>Asarum</i>
Chesnut, <i>Fagus</i>	Citron, <i>Citrus</i>
Chesnut, Horse, <i>Æsculus</i>	Citrus, <i>Cucurbita</i>
Chesnut, Indian Rose, <i>Mesua</i>	Cives, <i>Allium</i>
Chich Peas, <i>Cicer</i>	Clary, <i>Salvia</i>
Chiches, <i>Cicer</i>	Clary, Pyrænean, <i>Horminum</i>
Chichling Vetch, <i>Lathyrus</i>	Clivers, <i>Galium</i>
Chickweed, <i>Alsine</i>	Cloud-berry, <i>Rubuscam</i>
Chickweed, African, <i>Mollugo</i>	Clove July Flower, <i>Dianthus</i>
Chickweed, Berry-bearing, <i>Cucubalis</i>	Clove Tree, <i>Caryophyllum</i>
Chickweed, Great, <i>Stellariu</i>	Clover, <i>Trifolium</i>
Chickweed, Mountain, <i>Machringia</i>	Clover, Dutch, <i>Trifolium</i>
Chickweed, Mouse-ear, <i>Cerastium</i>	Clown's, Allheal, <i>Stachys</i>
Chickweed, Small-water, <i>Montia</i>	Clown's Wound-wort, <i>Stachys</i>
China Root, <i>Smilax</i>	Cob-nut, <i>Corylus</i>
China Rose, <i>Hibiscus</i>	Cock's-comb, <i>Celosia</i>
Chinquapin, <i>Fagus</i>	Cock's-comb, <i>Pedicularis</i>
Chocolate-nut, <i>Theobroma</i>	Cock's-comb, Yellow, <i>Rhinanthus</i>
Christmas Rose, <i>Helleborus</i>	Cock's-head, <i>Hedysarum</i>
Christopher, Herb, <i>Actæa</i>	Cocoa-nut, <i>Cocos</i>
Christ's-thorn, <i>Rhamnus</i>	Cocoa-plumb, <i>Chrysobalanus</i>
Chrysanthemum, Bastard, <i>Silphium</i>	Codlin Tree, <i>Pyrus</i>
Chrysanthemum, Hard-seeded, <i>Osteospermum</i>	Codlins and Cream, <i>Epilobium</i>
	Coffee Tree, <i>Coffea</i>
	Cole-seed, <i>Brassica</i>
	Cole-rape, <i>Brassica</i>
	Cole-wort, <i>Brassica</i>

Cole-wort, Sea, <i>Crambe</i>	Costmary, <i>Tanacetum</i>
Cole-wort, Sea, <i>Convolvulus</i>	Cotton, <i>Gossypium</i>
Coloquintida, <i>Cucumis</i>	Cotton, Lavender, <i>Santolina</i>
Colt's-foot, <i>Tussilago</i>	Cotton Tree, Silk, <i>Bombax</i>
Colt's-foot, Alpine, <i>Cacalia</i>	Cotton Grass, <i>Eriophorum</i>
Colt's-foot, Foreign, <i>Cacalia</i>	Cotton Weed, <i>Filago</i>
Columbine, <i>Aquilegia</i>	Coventry Bells, <i>Campanula</i>
Columbine Feathered, <i>Thalic-</i> <i>trum</i>	Courbaril, <i>Hymenæa</i>
Colutea, Jointed podded, <i>Coro-</i> <i>nilla</i>	Cow-quakes, <i>Briza</i>
Comphry, <i>Symphytum</i>	Cowslip, <i>Primula</i>
Consound, Greater, <i>Symphytum</i>	Cowslip, American, <i>Dodecatheon</i>
Consound, Lesser, <i>Bellis</i>	Cowslip, Jerusalem, <i>Pulmonaria</i>
Consound, Middle, <i>Ajuga</i>	Cowslip, Mountain, <i>Pulmonaria</i>
Consound, Royal, <i>Delphinium</i>	Cow's Lungwort, <i>Verbascum</i>
Consound, Saracen's, <i>Solidago</i>	Cow Parsnep, <i>Heracleum</i>
Consound, the True Saracen's, <i>Senecio</i>	Cow Weed, <i>Charophyllum</i>
Contrayerva, <i>Dorstenia</i>	Cow Wheat, <i>Melampyrum</i>
Contrayerva of Hernandez, <i>Pas-</i> <i>siflora</i>	Coxcomb. See Cock's-comb
Convall, Lily, <i>Convallaria</i>	Crab Tree, <i>Pyrus</i>
Coral Tree, <i>Erythrina</i>	Crake-berries, <i>Empetrum</i>
Coral-wort, <i>Dentaria</i>	Cranberries, <i>Vaccinium</i>
Coriander, <i>Coriandrum</i>	Crane's Bill, <i>Geranium</i>
Cork Tree, <i>Quercus suber</i>	Creeper, Virginian, <i>Hedera</i>
Corn, Indian, <i>Zea</i>	Cress, <i>Lepidium</i>
Corn Flag, <i>Gladiolus</i>	Cress, Indian, <i>Tropaeolum</i>
Corn Marigold, <i>Chrysanthemum</i>	Cress, Sciatica, <i>Iberis</i>
Corn Parsley, <i>Sison</i>	Cress, Spanish, <i>Vella</i>
Corn Rocket, <i>Bunias</i>	Cress, Swines, <i>Cochlearia</i>
Corn Rose, <i>Papaver</i>	Cress, Wall, <i>Turritis</i>
Corn Salad, <i>Valeriana</i>	Cress, Warded, <i>Cochlearia</i>
Cornel Tree, <i>Cornus</i>	Cress, Water, <i>Sisymbrium</i>
Cornelian Cherry, <i>Cornus</i>	Cross, Winter, <i>Erisimum</i>
	Cross, Jerusalem, <i>Lychnis</i>
	Cross, Knights, <i>Lychnis</i>
	Cross, Scarlet, <i>Lychnis</i>
	Cross-wort, <i>Valantia</i>

Crow-berries, <i>Empetrum</i>	Daffodil, Sea, <i>Pancreatum</i>
Crow-foot, <i>Ranunculus</i>	Daisy, <i>Bellis</i>
Crow-sick, <i>Conferva riv.</i>	Daisy, Blue, <i>Globularia</i>
Crown Imperial, <i>Fritillaria</i>	Daisy, Globe, <i>Globularia</i>
Cuckoo Flower, <i>Cardamine</i>	Daisy, Greater, <i>Chrysanthemum</i>
Cuckoo-pint, <i>Arum</i>	Daisy, Middle, <i>Doronicum</i>
Cucumber, <i>Cucumis</i>	Daisy, Ox-eye, <i>Chrysanthemum</i>
Cucumber, Asses, <i>Momordica</i>	Dame's Violet, <i>Hesperis</i>
Cucumber, Egyptian, <i>Momordica</i>	Damson Tree, <i>Prunus</i>
	Damson Tree, <i>Chrysophyllum</i>
Cucumber, Serpent, <i>Trichosanthes</i>	Dandelion, <i>Leontodon</i>
	Dane-wort, <i>Sambucus</i>
Cucumber, Single-seeded, <i>Sicyos</i>	Darnel, <i>Lolium</i>
Cucumber, Small creeping, <i>Melothria</i>	Date Plumb, Indian, <i>Diospyros</i>
	Date Tree, <i>Phœnix</i>
Cucumber, Spirting, <i>Momordica</i>	Day Lily, <i>Hemerocallis</i>
Cucumber, Wild, <i>Momordica</i>	Dead Nettle, <i>Lamium</i>
Cudweed, <i>Gnaphalium</i>	Dead Nettle, Yellow, <i>Galeopsis</i>
Cudweed, Bastard, <i>Micropus</i>	Deadly Carrot, <i>Thapsia</i>
Cullions, <i>Orchis</i>	Deadly Nightshade, <i>Atropa</i>
Cullions, Soldier's, <i>Orchis</i>	Devil in a Bush, <i>Nigella</i>
Cumin, <i>Cuminum</i>	Devil's Bit, <i>Scabiosa</i>
Cumin, Bastard, <i>Lagœcia</i>	Devil's Bit, Yellow, <i>Leontodon</i>
Cumin, Wild, <i>Lagœcia</i>	Dewberry Bush, <i>Rubus</i>
Cup Mushroom, <i>Peziza</i>	Dier's Broom, <i>Genista</i>
Currant Tree, <i>Ribes</i>	Dier's Weed, <i>Reseda</i>
Cushion Lady's, <i>Saxifraga</i>	Dier's Weed, <i>Genista</i>
Cushion, Sea, <i>Statice</i>	Dill, <i>Anethum</i>
Custard, Apple, <i>Annona</i>	Distaff Thistle, <i>Atractylis</i>
Cypress, <i>Cupressus</i>	Distaff Thistle, <i>Carthamus</i>
Cypress, Summer, <i>Chenopodium</i>	Dittander, <i>Lepidium</i>
	Dittany, <i>Origanum</i>
	Dittany, Bastard, <i>Marrubium</i>
	Dittany, White, <i>Dictamnus</i>
Daffodil, <i>Narcissus</i>	Dock, <i>Rumex</i>
Daffodil, Lily, <i>Amaryllis</i>	Doctor Tinker's Weed, <i>Triosticum</i>
Daffodil, Lily, <i>Pancreatum</i>	

Dodder, <i>Cuscuta</i>	Egg Plant, <i>Solanum</i>
Dodder of Thyme, <i>Cuscuta</i>	Eglantine, <i>Rosa</i>
Dog's Bane, <i>Apocynum</i>	Elder Tree, <i>Sambucus</i>
Dog's Bane, <i>Asclepias</i>	Elder, Marsh, <i>Viburnum</i>
Dog-berry, <i>Cornus</i>	Elecampane, <i>Inula</i>
Dog's Cabbage, <i>Theligomum</i>	Elecampane, Bastard, <i>Helenia</i>
Dog's Rue, <i>Scrophularia</i>	Elemi Tree, Gum, <i>Pistacia</i>
Dog's Stones, <i>Orchis</i>	Elephant's Foot, <i>Elephantopus</i>
Dog's Tooth, or Dog's Tooth Violet, <i>Erythronium</i>	Elephant's Head, <i>Rhinanthus</i>
Dogwood, <i>Cornus</i>	Elichrysum, Bastard Ethiopian, <i>Stabe</i>
Dogwood of Jamaica, <i>Erythrina</i>	Eller, <i>Bettula alb.</i>
Double Tongue, <i>Ruscus</i>	Elm, <i>Ulmus</i>
Dove's Foot, <i>Geranium</i>	Enchanter's Nightshade, <i>Circea</i>
Dragons, <i>Dracontium</i>	Endive, <i>Chichorium</i>
Dragons, <i>Arum</i>	Eryngo, <i>Eryngium</i>
Dragon's Head, <i>Dracocephalum</i>	Eschalot, <i>Allium</i>
Dragon's Water, <i>Calla</i>	Eternal Flower, <i>Xeranthema</i>
Dragon's Wort, <i>Artemisia</i>	Eternal Flower, <i>Gnaphalium</i>
Dragon, Gum, see Tragacanth	Eternal Flower, <i>Gomphrena</i>
Dragon, Wild, <i>Artemisia</i>	Evergreen, <i>Aizoon</i>
Drop-wort, <i>Spiræa</i>	Evergreen, <i>Semprevivum</i>
Drop-wort, Hemlock, <i>Ænanthe</i>	Everlasting, <i>Xeranthemum</i>
Drop-water, <i>Ænanthe</i>	Everlasting, <i>Gomphrena</i>
Duck's-meat, <i>Lemna</i>	Everlasting, <i>Gnaphalium</i>
Duck's-meat, Starry, <i>Callitriche</i>	Euonymus, Climbing, <i>Celastrus</i>
Duck's-foot, <i>Podophyllum</i>	Euonymus, Bastard, <i>Kiggellaria</i>
Dulse, <i>Fucus palm</i>	Euonymus, Bastard, <i>Celastrus</i>
Dwale, <i>Atropa</i>	Eye-bright, <i>Euphrasia</i>

E

Ebony, Cretan, *Ebenus*
 Ebony, False, *Poinciana*
 Ebony of the Alps, *Cytisus*
 Ebony, Mountain, *Bauhinia*
 Edders, *Arum*

F

Fairy Mushroom, *Agaricus cor.*
 Farting Tree, *Hura*
 Faufel Nut, *Areca*
 Felwort, *Gentiana*
 Felon-wort, *Solanum*

Fennel, <i>Anethum</i>	Finochia, <i>Anethum</i>
Fennel Hog's, <i>Peucedanum</i>	Fir, <i>Pinus</i>
Fennel, Scorching, <i>Thapsia</i>	Fir Moss, Upright, <i>Lycopodium</i>
Fennel, Sea, <i>Crithmum</i>	Fish Thistle, <i>Carduus</i>
Fennel Flower, <i>Nigella</i>	Flag, or Flag-flower, <i>Iris</i>
Fennel Flower of Crete, <i>Gari-</i>	Flag, Corn, <i>Gladiolus</i>
<i>della</i>	Flag, Sweet-scented, <i>Acorus</i>
Fennel Giant, <i>Ferula</i>	Flax, <i>Linum</i>
Fenugreek, <i>Trigonella</i>	Flax, Carolina, <i>Polypremum</i>
Fern, Common Male, <i>Polypodium</i>	Flax, Toad, <i>Antirrhinum</i>
Fern, Common Female, <i>Polypo-</i>	Fleabane, <i>Coryza</i>
<i>dium</i>	Fleabane, Marsh, <i>Inula</i>
Fern, Flowering, <i>Osmunda</i>	Fleabane, Middle, <i>Inula</i>
Fern, Common, or True Mules,	Fleabane, Shrubby African, <i>Tar-</i>
<i>Asplenium</i>	<i>chonanthus</i>
Fern, Mules, <i>Hemionitis</i>	Fleabane Tree, <i>Tarchonanthus</i>
Fern, Sweet, <i>Scandix</i>	Flea-wort, <i>Plantago</i>
Feverfew, <i>Matricaria</i>	Flix-weed, <i>Sisymbrium</i>
Feverfew, Bastard, <i>Parthenium</i>	Flower of Bristol, <i>Lychnis</i>
Fever-root, <i>Triosteum</i> ,	Flower of Constantinople, <i>Lychnis</i>
Fever-weed, <i>Eryngium</i>	Flower Gentle, <i>Amaranthus</i>
Fiddle-wood, <i>Citharexylum</i>	Flower of an Hour, <i>Hibiscus</i>
Field Basil, <i>Clenopodium</i>	Flower de Luce, <i>Iris</i>
Field Basil, American, <i>Monarda</i>	Flower-fence of Barbadoes, <i>Po-</i>
Field Basil, Syrian, <i>Ziziphora</i>	<i>inciana</i>
Fig, <i>Ficus</i>	Flower-fence, Bastard, <i>Adenan-</i>
Fig, Indian, <i>Cactus</i>	<i>thera</i>
Fig, Infernal, <i>Argemone</i>	Fluellin, <i>Antirrhinum</i>
Fig, Pharoah's, <i>Ficus</i>	Fly Honeysuckle, <i>Lonicera</i>
Fig, Pharoah's, <i>Musa</i>	Fly Honeysuckle, African, <i>Hal-</i>
Fig, Marigold, <i>Mesembryanthe-</i>	<i>leria</i>
<i>mum</i>	Fly Bane, <i>Silene</i>
Fig Tree, Cochineal, <i>Cactus</i>	Fly-wort, <i>Silene</i>
Fig-wort, <i>Scrophularia</i>	Fool's Parsley, <i>Æthusa</i>
Filberd, <i>Corylus</i>	Fool's Stones, <i>Orchis</i>
Fingrigo, <i>Pisonia</i>	Four o'clock Flower, <i>Mirabilis</i>

Fox Glove, <i>Digitalis</i>	Germander, Water, <i>Teucrium</i>
Fox-tail Grass, <i>Alopecurus</i>	Gilead, False Baulm of, <i>Draco-</i>
Frankincense, Jews, <i>Styrax</i>	<i>cephalum</i>
Frankincense Tree, <i>Pinus</i>	Gill, <i>Glechoma</i>
Fraxinella, <i>Dictamnus</i>	Gilly-flower, see July-flower
French Bean, <i>Phaseolus</i>	Ginger, <i>Amomum</i>
French Honeysuckle, <i>Hedysarum</i>	Ginseng, <i>Panax</i>
Fresh-water Soldier, <i>Stratiotes</i>	Gladiole, Water, <i>Butomus</i>
Friar's Cowl, <i>Arum</i>	Gladiole, Water, <i>Lobelia</i>
Fringe Tree, <i>Chionanthus</i>	Gladwin, Stinking, <i>Iris</i>
Fritillary, <i>Fritillaria</i>	Glass-wort, <i>Salsola</i>
Fritillary Coxcomb, <i>Stapelia</i>	Glass-wort, Berry-bearing, <i>Ana-</i>
Frog's Bit, <i>Hydrocharis</i>	<i>basis</i>
Fuller's Thistle, <i>Dipsacus</i>	Glass-wort, Jointed, <i>Salicornia</i>
Fumatory, <i>Fumaria</i>	Globe Amaranth, <i>Gomphrena</i>
Furze, <i>Ulex</i>	Globe Daisy, <i>Globularia</i>
Fustic Tree, <i>Morus</i>	Globe Flower, <i>Spharanthus</i>
	Globe Ranunculus, <i>Trollius</i>
	Globe Thistle, <i>Echinops</i>
G	Goat's Beard, <i>Tragopogon</i>
Gale, or Sweet Gale, <i>Myrica</i>	Goat's-Rue, <i>Galega</i>
Galingale, <i>Cyperus</i>	Goat's Stones, greater, <i>Satyrium</i>
Garavanees, <i>Cicer</i>	Goat's Stones, lesser, <i>Orchis</i>
Garlick, <i>Allium</i>	Goat's Thorn, <i>Astragalus</i>
Garlick Pear, <i>Crateva</i>	Gold of Pleasure, <i>Myagrum</i>
Gatter Tree, <i>Cornus</i>	Golden Cups, <i>Ranunculus</i>
Gelder Rose. <i>Viburnum</i>	Golden Lung-wort, <i>Hieracium</i>
Gelder Rose, Currant-leaved,	Golden Maiden-hair, <i>Polytrichum</i>
<i>Spiræa</i>	Golden Mouse-ear, <i>Hieracium</i>
Gelder Rose, Virginian, <i>Spiræa</i>	Golden Rod, <i>Solidago</i>
Gentian, <i>Gentiana</i>	Golden Rod Tree, <i>Bosea</i>
Gentian, Bastard, <i>Sarothra</i>	Golden Samphire, <i>Inula</i>
Gentianella, <i>Gentiana</i>	Golden Saxifrage, <i>Chrysopenium</i>
Gentle, Flower, <i>Amaranthus</i>	Golden Thistle, <i>Scolymus</i>
Gerard, Herb, <i>Ægopodium</i>	Golden Locks, <i>Chrysocoma</i>
Germander, <i>Teucrium</i>	Golden Locks, <i>Gnaphalium</i>
Germander, Rock, <i>Veronica</i>	

Good Henry, <i>Chenopodium</i>	Gromwell, German, <i>Stellera</i>
Gooseberry, <i>Ribes</i>	Ground Ivy, <i>Glechoma</i>
Gooseberry, American, <i>Melastoma</i>	Ground Nut, <i>Arachis</i>
	Ground Pine, <i>Teucrium</i>
Gooseberry of the Americans, <i>Cactus</i>	Ground Pine, Stinking, <i>Camphorosma</i>
Gooseberry of Barbadoes, <i>Cactus</i>	Groundsel, <i>Senecio</i>
Goose Foot, <i>Chenopodium</i>	Groundsel Tree, <i>Baccharis</i>
Goose Grass, <i>Galium</i>	Groundsel Tree, with a Ficoides Leaf, <i>Cacalia</i>
Goose Grass, great, <i>Asperugo</i>	Guava. See Guayava
Goose Tongue, <i>Achillea</i>	Guava, French, <i>Cassia</i>
Go to bed at Noon, <i>Tragopogon</i>	Guayava, <i>Psidium</i>
Gorss, <i>Ulex</i>	Guills, <i>Chrysanthemum</i> , <i>seg.</i>
Gourd, <i>Cucurbita</i>	Gum Elemi Tree, <i>Pistachia</i>
Gourd, Bitter, <i>Cucumis</i>	Gum Succory, <i>Chondrilla</i>
Gourd, Ethiopian, Sour, <i>Adansonia</i>	Gum Tragacanth, <i>Astragalus</i>
Gourd Tree, Indian, <i>Crescentia</i>	Gum, Sweet, <i>Liquidambar</i>
Gout-wort, <i>Ægopodium</i>	
Gowan, <i>Bellis</i>	H
Grace, Herb of, <i>Ruta</i>	Hag-berries, <i>Prunus pad.</i>
Grain, Oily Purging, <i>Sesamum</i>	Hag-taper, <i>Verbascum thap.</i>
Grain, Scarlet, <i>Quercus</i>	Hair-bells, <i>Hyacinthus</i>
Grain, Scarlet, <i>Cactus</i>	Hare's-ear, <i>Beuphrium</i>
Grape, <i>Vitis</i>	Hare's-ear, Bastard, <i>Phyllis</i>
Grape, Mangrove, <i>Polygonum</i>	Hare's Lettuce, <i>Sonchus</i>
Grape, Sea-side, <i>Polygonum</i>	Hart's-horn Plantain, <i>Plantago</i>
Grape Hyacinth, <i>Hyacinthus</i>	Hart's-tongue, <i>Asplenium</i>
Grass of Parnassus, <i>Parnassia</i>	Hart-wort, <i>Seseli</i>
Grass Vetch, Crimson, <i>Lathyrus</i>	Hart-wort of Crete, <i>Tordylium</i>
Grass Wrack, <i>Zostera</i>	Hart-wort, Shrubby, of Ethiopia, <i>Bupleurum</i>
Gravel-bind, <i>Convolvulus</i>	
Greek Valerian, <i>Polemonium</i>	Hart-wort of Marseilles, <i>Seseli</i>
Green-weed, <i>Genista</i>	Hatchet Vetch, Tree, <i>Coronilla</i>
Grim the Collier, <i>Hieracium</i>	Hatchet Vetch, Clusius's, Foreign, <i>Biserrula</i>
Gromwell, or Gromil, <i>Lithospermum</i>	

Hawk-weed, <i>Hieracium</i>	Helmet-flower, <i>Aconitum</i>
Hawk-weed, Bastard, <i>Crepis</i>	Hemlock, <i>Conium</i>
Hawk-weed, Trailing crooked-seeded, <i>Hyoseris</i>	Hemlock, Great broad-leaved Bastard, <i>Ligusticum</i>
Hawk-weed, Woolly, <i>Andryala</i>	Hemlock, Lesser, <i>Æthusa</i>
Hawthorn, or Haw, <i>Cratægus</i>	Hemlock, Water, <i>Cicuta</i>
Hawthorn, Black American, <i>Viburnum</i>	Hemlock Drop-wort, <i>Cœnanthe</i>
Hay, Burgundian, <i>Medicago</i>	Hemp, <i>Cannabis</i>
Hazel, or Hazel Nut, <i>Corylus</i>	Hemp, Bastard, <i>Datisca</i>
Hazel, Witch, <i>Hamamelis</i>	Hemp, Bastard, <i>Galeopsis</i>
Hazel, Witch, <i>Ulmus</i>	Hemp Agrimony, <i>Eupatorium</i>
Hart Pea, <i>Cardiospermum</i>	Hemp Agrimony, Bastard, <i>Ageratum</i>
Heart Seed, <i>Cardiospermum</i>	Hemp Agrimony, Naked-headed, <i>Verbesina</i>
Heart's Ease, <i>Viola</i>	Hemp Agrimony, Water, <i>Bidens</i>
Heath, <i>Erica</i>	Henbane, <i>Hyoscyamus</i>
Heath, Berry-bearing, <i>Empetrum</i>	Henbane, Yellow, <i>Nicotiana</i>
Heath, Black-berried, <i>Empetrum</i>	Henweed, Guinea, <i>Petiveria</i>
Heath, Mountain, <i>Saxifraga</i>	Hepatica, <i>Anemone</i>
Heath, Low Pine, <i>Coris</i>	Hep Tree, <i>Rosa</i>
Heath Peas, <i>Orobus</i>	Herb-bane, <i>Orobanche</i>
Hedge-hog, <i>Medicago</i>	Herb-bane, Great Purple, <i>Lathræa</i>
Hedge-hog Thistle, <i>Cactus</i>	Herb Bennet, <i>Geum</i>
Hedge-hog Thorn, Spanish, <i>Anthyllis</i>	Herb Christopher, <i>Actæa</i>
Hedge Hyssop, <i>Gratiola</i>	Herb Gerard, <i>Ægopodium</i>
Hedge Mustard, <i>Erysium</i>	Herb of Grace, <i>Ruta</i>
Hedge Nettle, <i>Galeopsis</i>	Herb Mastick, <i>Satureia</i>
Hedge Nettle, Shrubby, <i>Prasium</i>	Herb Paris, <i>Paris</i>
Hellebore, <i>Helleborus</i>	Herb Paris of Canada, <i>Trillium</i>
Hellebore, Bastard, <i>Serapias</i>	Herb Robert, <i>Geranium</i>
Hellebore, Black, <i>Helleborus</i>	Herb Trinity, <i>Viola</i>
Hellebore, Fennel-leaved Black, <i>Adonis</i>	Herb Truelove, <i>Paris</i>
Hellebore, White, <i>Veratrum</i>	Herb Truelove of Canada, <i>Trillium</i>
Helleborine, <i>Serapias</i>	

Herb Two-pence, <i>Lysimachia</i>	Hop-tree, <i>Ilex</i>
Herb, Blessed, <i>Geum</i>	Horehound, <i>Macrabium</i>
Herb, St. Bartholomew's, <i>Ilex</i>	Horehound, Base, <i>Stachys</i>
Herb, Willow, <i>Epilobium</i>	Horehound, Bastard, <i>Sideritis</i>
Herb, Willow, <i>Lythcum</i>	Horehound, Black, <i>Ballota</i>
Herb, Willow, <i>Lisymachia</i>	Horehound, Stinking Marsh, Bastard, <i>Glechoma</i>
Hercules's Allheal, <i>Pastiuaca</i>	Horehound, Water, <i>Lycopus</i>
Hercules's Allheal, <i>Heracleum</i>	Hornbeam, <i>Carpinus</i>
Hercules's Club, <i>Zanthoxylon</i>	Horns, <i>Medicago</i>
Hicory Nut, <i>Jugians</i>	Horse Chesnut, <i>Æsculus</i>
High Taper, <i>Verbascum</i>	Horse Purslane, <i>Trianthema</i>
Hind-berry, <i>Rubus</i>	Horse-radish, <i>Cochleavia</i>
Hog Plumb-tree, <i>Spodias</i>	Horse-shoe Vetch, <i>Hippocrepis</i>
Hog's Fennel, <i>Peucedanum</i>	Horse-tail, <i>Equisetum</i>
Hog-weed of the Americans, <i>Boerhaavia</i>	Horse-tail, Shrubby, <i>Ephedra</i>
Hollow Root, <i>Adoxa</i>	Horse-tongue, <i>Ruscus</i>
Holly, <i>Ilex</i>	Hottentot Cherry, <i>Cassine</i>
Holly, Knee, <i>Ruscus</i>	Hound's-tongue, <i>Cynoglossum</i>
Holly, Sea, <i>Eryngium</i>	Houseleek, <i>Sempervivum</i>
Hollyhock, <i>Alcea</i>	Houseleek, Lesser, <i>Sedum</i>
Holy Thistle, <i>Cnicus</i>	Houseleek, Small, annual, <i>Tillæa</i>
Honesty, <i>Lunaria</i>	Houseleek, Water of Egypt, <i>Pistia</i>
Hone-wort, <i>Sison</i>	Hyacinth, <i>Hyacinthus</i>
Honey-flower, <i>Melianthus</i>	Hyacinth, African Blue, umbel- lated, <i>Crinum</i>
Honey Locust, <i>Gleditsia</i>	Hyacinth, Lily, <i>Scilla</i>
Honeysuckle, <i>Lonicera</i>	Hyacinth, Peruvian, <i>Scilla</i>
Honeysuckle, African Fly, <i>Hal- leria</i>	Hyacinth, Starry, <i>Scilla</i>
Honeysuckle, American Upright, <i>Azalea</i>	Hyssop, <i>Hyssopus</i>
Honeysuckle, French, <i>Hedysa- rum</i>	Hyssop, Hedge, <i>Gratiola</i>
Honeysuckle Grass, <i>Trifolium</i>	Hyssop, Mountain, <i>Thymbra</i>
Honey-wort, <i>Cerithe</i>	I
Hop, <i>Humulus</i>	Jacinth, <i>Hyacinthus</i>

Jack in a Box, <i>Hernandia</i>	Indigo, Bastard, <i>Amorpha</i>
Jack by the Hedge, <i>Erysimum</i>	Infernal Fig, <i>Argemone</i>
Jacob's Ladder, <i>Polemonium</i>	Job's Tears, <i>Coin</i>
Jacobæa Lily, <i>Amaryllis</i>	Johnsonia, <i>Callicarpa</i>
Jalap, <i>Mirabilis</i>	Johnquill, <i>Narcissus</i>
Jasmine, <i>Jasminum</i>	Ipecacuana, Bastard, <i>Asclepias</i>
Jasmine, Arabian, <i>Nyctanthes</i>	Ipecacuana, False, <i>Triosteum</i>
Jasmine, Bastard, <i>Cestrum</i>	Iris, Uvaria, <i>Aletris</i>
Jasmine, Bastard, <i>Lycium</i>	Iron-wood, <i>Sideroxylum</i>
Jasmine, Ilex-leaved, <i>Lantana</i>	Iron-wort, <i>Sideritis</i>
Jasmine, Fennel-leaved, <i>Ipomœa</i>	Judas-tree, <i>Circis</i>
Jasmine, Persian, <i>Syringa</i>	Jujube-tree, <i>Rhamnus</i>
Jasmine, Red, <i>Plumeria</i>	July-flower, Clove, <i>Dianthus</i>
Jasmine, Scarlet, <i>Bignonia</i>	July-flower, Queen's, <i>Hesperis</i>
Jasmine, Yellow, <i>Bignonia</i>	July-flower, Stock, <i>Cheiranthus</i>
Jericho, Rose of, <i>Anastatica</i>	Juniper, <i>Juniperus</i>
Jersey, Thea, New, <i>Ceanothus</i>	Jupiter's Beard, <i>Anthyllis</i>
Jerusalem Artichoke, <i>Helianthus</i>	Jupiter's Beard, American,
Jerusalem Cowslip, <i>Pulmonaria</i>	<i>Amorpha</i>
Jerusalem Cross, <i>Lychnis</i>	Jupiter's Distaff, <i>Salvia</i>
Jerusalem Oak, <i>Chenopodium</i>	Ivy, <i>Hedera</i>
Jerusalem, Sage, <i>Phlomis</i>	Ivy, Bindweed-leaved, <i>Meni-</i>
Jerusalem, Sage of, <i>Pulmonaria</i>	<i>spermum</i>
Jessamine, see Jasmine	Ivy, Ground, <i>Glechoma</i>
Jesuit's Bark-tree, True, <i>Chin-</i>	Ivy-tree of America, <i>Kalmia</i>
<i>chona</i>	
Jesuit's Bark-tree, False, <i>Iva</i>	K
Jew's Frankincense, <i>Styrax</i>	Kale, Sea, <i>Crambe</i>
Jew's Mallow, <i>Corchorus</i>	Kali, <i>Salsola</i>
Ilathera Bark, <i>Clusia</i>	Kali, Egyptian, <i>Mesembryanthe-</i>
Immortal Eagle Flower, <i>Impa-</i>	<i>mum</i>
<i>tiens</i>	Kali, Sal, <i>Salicornia</i>
Immortal Flower, <i>Gomphrena</i>	Kandel of the Indians, <i>Rhizo-</i>
Indian God Tree, <i>Ficus</i>	<i>phora</i>
Indian Shot, <i>Canna</i>	Kelp, <i>Salicornia</i>
Indigo, <i>Indigofera</i>	Kermes, <i>Quercus</i>

Kidney Bean, <i>Phaseolus</i>	Lark's Heel, <i>Delphinium</i>
Kidney Bean-tree of Carolina, <i>Glycine</i>	Lark's Spur, <i>Delphinium</i>
Kidney Vetch, <i>Anthyllis</i>	Laserwort, <i>Laserpitium</i>
Kidney-wort, <i>Saxifraga</i>	Lavender, <i>Lavandula</i>
King's Spear, <i>Asphodelus</i>	Lavender, Sea, <i>Statice</i>
Knapweed, <i>Centaurea</i>	Lavender Cotton, <i>Santolina</i>
Knapweed, Thorny, <i>Centaurea</i>	Laver, <i>Ulva</i>
Knawel, <i>Scleranthus</i>	Laurel, <i>Prunus</i>
Knee Holly, <i>Ruscus</i>	Laurel, Alexandrian, <i>Ruscus</i>
Knee Holm, <i>Ruscus</i>	Laurel, Dwarf, of America, <i>Kalmia</i>
Knight's Cross, <i>Lychnis</i>	Laurel, Flax-leaved, <i>Daphne</i>
Knot Berries, <i>Rubus</i>	Laurel, Sea-side, <i>Phyllanthus</i>
Knot Grass, <i>Polygonum</i>	Laurel, Spurge, <i>Daphne</i>
Knot Grass, German, <i>Scleranthus</i>	Laurustinus, <i>Viburnum</i>
Knot Grass, Mountain, <i>Illecebrum</i>	Lauskraut, <i>Delphinium</i>
Knot Grass, Verticillate, <i>Illecebrum</i>	Lead-wort, <i>Plumbago</i>
	Leather-wood, <i>Dirca</i>
	Leek, <i>Allium</i>
	Lemon, <i>Citrus</i>
	Lemon, Water, <i>Passiflora</i>
	Lentils, <i>Ervum</i>
	Lentisk, <i>Pistacia</i>
	Lentisk, African, <i>Schinus</i>
	Lentisk, Peruvian, <i>Schinus</i>
	Leopard's Bane, <i>Doronicum</i>
	Lettuce, <i>Lactuca</i>
	Lettuce, Hare's, <i>Sonchus</i>
	Lettuce, Lamb's, <i>Valeriana</i>
	Lettuce, Wild, <i>Prenanthes</i>
	Life, Tree of, <i>Thuja</i>
	Life, Wood of, <i>Guaiacum</i>
	Life Everlasting, <i>Gnaphalium</i>
	Lignum Aloes, <i>Cordica</i>
	Lignum Vitæ, <i>Guaiacum</i>
	Lilac, <i>Syringa</i>

Lily, <i>Lilium</i>	Liquorice, Wild, <i>Capraria</i>
Lily, African Scarlet, <i>Amaryllis</i>	Liquorice, Wild, <i>Glycine</i>
Lily, Asphodel, <i>Crimum</i>	Liquorice Vetch, <i>Astragalus</i>
Lily, Atamasco, <i>Amaryllis</i>	Liquorice Vetch, Knobbed-root- ed, <i>Glycine</i>
Lily, Belladonna, <i>Amaryllis</i>	Live-ever, <i>Sedum</i>
Lily, St. Bruno's, <i>Heimerocallis</i>	Live-long, <i>Sedum</i>
Lily, Convall, <i>Convallaria</i>	Liver-wort, <i>Lichen</i>
Lily, Day, <i>Heimerocallis</i>	Liver-wort, Marsh, <i>Riccia</i>
Lily, Guernsey, <i>Amaryllis</i>	Liver-wort, Noble, <i>Anemone</i>
Lily, Jacobæa, <i>Amaryllis</i>	Lizard's-tail, <i>Saururus</i>
Lily, Japan, <i>Amaryllis</i>	Lizard's-tail, <i>Piper</i>
Lily, May, <i>Convallaria</i>	Loblolly Bay, <i>Gordonia</i>
Lily, Mexican, <i>Amaryllis</i>	Locker Gowlands, <i>Trollius</i>
Lily, Persian, <i>Fritillaria</i>	Locust, <i>Melanthus</i>
Lily, Superb, <i>Gloriosa</i>	Locust, <i>Ceratonia</i>
Lily, Water, <i>Nymphaea</i>	Locust, Bastard, <i>Hymenææ</i>
Lily, Lesser Yellow Water, with fringed Flowers, <i>Menyanthes</i>	Locust-tree, <i>Hymenææ</i>
Lily, Zeylon, <i>Amaryllis</i>	Locust-tree, <i>Robinia</i>
Lily, Asphodel, <i>Heimerocallis</i>	Locust-tree, Honey, <i>Gleditsia</i>
Lily, Daffodil, <i>Amaryllis</i>	Logwood, <i>Hamatoxylon</i>
Lily, Daffodil, <i>Pancreatium</i>	London Pride, <i>Saxifraga</i>
Lily, Hyacinth, <i>Scilla</i>	Loose-strife, <i>Lysimachia</i>
Lily, Thorn, <i>Catesbaa</i>	Loose-strife, Podded, <i>Ipilobium</i>
Lily of the Valley, <i>Convallaria</i>	Loose-strife, Purple, <i>Lythrum</i>
Lime, <i>Citrus</i>	Loose-strife, Spiked, <i>Lythrum</i>
Lime, Brook, <i>Veronica</i>	Loose-strife, Yellow Virginian, <i>Gaura</i>
Lime-tree, <i>Tilia</i>	
Ling, <i>Erica</i>	Lords and Ladies, <i>Arum</i>
Linden-tree, <i>Tilia</i>	Lotus, or Lote-tree, <i>Celtis</i>
Lion's-foot, Candy, <i>Catananche</i>	Lotus, supposed of Homer, <i>Dios- pyros</i>
Lion's leaf, <i>Leontice</i>	
Lion's-tail, <i>Leonurus</i>	Lotus, Honey, <i>Trifolium</i>
Lipplehout, <i>Cassine</i>	Lovage, <i>Ligusticum</i>
Liquorice, <i>Glycyrrhiza</i>	Love, Tree of, <i>Cercis</i>
Liquorice, Wild, <i>Astragalus</i>	Love Apple, <i>Solanum</i>

Love in a Mist, <i>Passiflora</i>	Mallow, Jew's, <i>Corchorus</i>
Love lies a bleeding, <i>Antarantus</i>	Mallow, Indian, <i>Sida</i>
Louse-wort, <i>Pedicularis</i>	Mallow, Indian, <i>Urena</i>
Louse-wort, Yellow, <i>Rhinanthus</i>	Mallow, Marsh, <i>Althwa</i>
Lucern Grass, <i>Medicago</i>	Mallow, Rose, <i>Alcea</i>
Lucken-Gowan, <i>Trollius</i>	Mallow, Syrian, <i>Hibiscus</i>
Lung-wort, <i>Pulmonaria</i>	Mallow, Tree, <i>Lavatera</i>
Lung-wort, Cow's, <i>Verbascum</i>	Mallow, Varied leav'd, <i>Lavatera</i>
Lung-wort, Golden, <i>Hieracium</i>	Mallow, Venetian, <i>Lavatera</i>
Lupine, <i>Lupinus</i>	Mallow, Vervain, <i>Malva</i>
Lust-wort, <i>Drosera</i>	Mallow, Yellow, <i>Sida</i>
Lychnidea, <i>Phlox</i>	Mammee, <i>Mammea</i>
Lychnis, Bastard, <i>Phlox</i>	Mammee, Sapota, <i>Achras</i>
Lychnis, Wild, <i>Agrostema</i>	Manchineel-tree, <i>Hippomane</i>
	Mandrake, <i>Mandragora</i>
	Mango-tree, <i>Mangifera</i>
	Mangostan, or Mangosteen, <i>Garciana</i>
	Mangrove Grape, <i>Polygonum</i>
	Mangrove-tree of America, <i>Rhizophora</i>
	Manihot, <i>Jatropha</i>
	Maple, <i>Acer</i>
	Maracœck, <i>Passiflora</i>
	Marigold, <i>Calendula</i>
	Marigold, African, <i>Tagetes</i>
	Marigold, Corn, <i>Chrysanthemum</i>
	Marigold, Fig, <i>Mesembryanthemum</i>
	Marigold, French, <i>Tagetes</i>
	Marigold, Marsh, <i>Caltha</i>
	Marjoram, Common or Sweet, <i>Origanum</i>
	Marjoram, Bastard, <i>Origanum</i>
	Marjoram, Pot, <i>Origanum</i>
	Marjoram, Spanish, <i>Urtica</i>

Marjoram, Wild, <i>Origanum</i>	Medic, <i>Medicago</i>
Marjoram, Winter Sweet, <i>Origanum</i>	Medic, Bastard, <i>Medicago</i>
	Medic, Sea, <i>Medicago</i>
Marsh-mallow. See Mallow	Medic, Vetch, <i>Hedysarum</i>
Martagon, <i>Lilium</i>	Medic, Vetchling, <i>Hedysarum</i>
Marvel of Peru, <i>Mirabilis</i>	Medlar, <i>Mespilus</i>
Marum, Common, <i>Satureia</i>	Medusa's Head, <i>Euphorbia</i>
Marum, Pennyroyal-scented, <i>Melissa</i>	Melancholy Thistle, <i>Carduus</i>
	Melancholy-tree, <i>Nyctanthes</i>
Marum, Syrian or Cretan, <i>Origanum</i>	Melilot, <i>Trifolium</i>
	Melon, <i>Cucumis</i>
Master-wort, <i>Imperatoria</i>	Melon, Water, <i>Cucurbita</i>
Master-wort, Black, <i>Astrantia</i>	Melon-thistle, <i>Cactus</i>
Mastich, Herb, <i>Satureia</i>	Mercury, <i>Mercurialis</i>
Mastich, Indian, <i>Schinus</i>	Mercury, English, <i>Chenopodium</i>
Mastich, Peruvian, <i>Schinus</i>	Mezereon, <i>Daphne</i>
Mastich-tree, <i>Pistachia</i>	Meu, <i>Athamanta</i>
Mastich-tree, Indian, <i>Schinus</i>	Mignonette, <i>Reseda</i>
Mastich Thyme, <i>Satureia</i>	Milfoil, <i>Achillea</i>
Mastich Thyme, <i>Thymus</i>	Milfoil, Water, <i>Hottonia</i>
Martfellon, <i>Centaurea</i>	Milfoil, Water, <i>Myriophyllum</i>
Mat-weed, Hooded, <i>Lygeum</i>	Milfoil, Water, <i>Utricularia</i>
Maudlin, <i>Achillea</i>	Milk Vetch, <i>Astragalus</i>
May Apple, <i>Podophyllum</i>	Milk Vetch, Bastard, <i>Phaca</i>
May Bush, <i>Cratægus</i>	Milk Wood, <i>Bignonia</i>
May Lily, <i>Convallaria</i>	Milk-wort, <i>Polygala</i>
May Weed, <i>Anthemis</i>	Milk-wort, <i>Euphorbia</i>
Mays, <i>Zea</i>	Milk-wort, Sea, <i>Glaur</i>
Meadia, <i>Dòdecathenon</i>	Millett, <i>Panicum</i>
Meadow Rue, <i>Thalictrum</i>	Millet-grass, <i>Milium</i>
Meadow Saffron, <i>Colchicum</i>	Millet, Indian, <i>Holcus</i>
Meadow Saxifrage, <i>Peucedanum</i>	Milt-waste, <i>Asplenium</i>
Meadow-sweet, <i>Spiræa</i>	Mint, <i>Mentha</i>
Meadow-sweet, Greater, <i>Spiræa</i>	Mint, Cat, <i>Nepeta</i>
Meadow, Queen of the, <i>Spiræa</i>	Mistletoe, <i>Viscum</i>
Mealy-tree, Pliant, <i>Viburnum</i>	Mithridate Mustard, <i>Thlaspi</i>

Mithridate Mustard, Bastard, Mulberry-tree, <i>Morus</i>	
<i>Iberis</i>	Mulberry Blite, <i>Blitum</i>
Mock Orange, <i>Philadelphus</i>	Mule Fairchild's, <i>Dianthus</i>
Mock Privet, <i>Phillyrea</i>	Mule-wort, <i>Hemionitis</i>
Moldavian Baulm, <i>Dracocephalum</i>	Mule's Fern, <i>Hemionitis</i>
	Mullein, <i>Verbascum</i>
Molucca Baulm, <i>Moluccella</i>	Mullein, Moth, <i>Verbascum</i>
Moly with Lily-flowers, or Ho-	Mushrooms, <i>Agaricus</i>
mer's, <i>Allium</i>	Mushrooms, Cup, <i>Peziza</i>
Money-wort, <i>Lysimachia</i>	Musk Seed, <i>Hibiscus</i>
Monk's-head, <i>Leontodon</i>	Mustard, <i>Sinapis</i>
Monk's-hood, <i>Aconitum</i>	Mustard, Bastard, <i>Cleome</i>
Monk's Rhubarb, <i>Rumex</i>	Mustard Buckler, <i>Biscutella</i>
Monster, <i>Fritillaria</i>	Mustard, Hedge, <i>Erysimum</i>
Moon Seed, <i>Menispermum</i>	Mustard, Mithridate, <i>Thlaspi</i>
Moon Trefoil, <i>Medicago</i>	Mustard, Bastard Mithridate,
Moon-wort, <i>Lunaria</i>	<i>Iberis</i>
Moor Berries, <i>Vaccinium</i>	Mustard, Tower, <i>Turritis</i>
Moschatel, Tuberose, <i>Adoxa</i>	Mustard, Bastard Tower, <i>Arabis</i>
Moss-tree, <i>Lichen</i>	Mustard, Treacle, <i>Clypeola</i>
Moss, Upright Fir, <i>Lycopodium</i>	Mustard, Treacle, <i>Thlaspi</i>
Moss, Water, <i>Fontinalis</i>	Myrtle, <i>Myrtus</i>
Moss-berries, <i>Vaccinium</i>	Myrtle, Candleberry, <i>Myrica</i>
Moth Mullein, <i>Verbascum</i>	Myrtle, Dutch, <i>Myrica</i>
Mother of Thyme, <i>Thymus</i>	
Mother-wort, <i>Leonurus</i>	N
Mouse-ear, <i>Hieracium</i>	Naked Ladies, <i>Colchicum</i>
Mouse-ear, Creeping, <i>Hieracium</i>	Naples, Star of, <i>Ornithogalum</i>
	Narcissus, Third, of Matthiolum,
Mouse-ear, Golden, <i>Hieracium</i>	<i>Pancratium</i>
Mouse-ear Chickweed, <i>Cerastium</i>	Naseberry-tree, <i>Sloanea</i>
Mouse-ear Scorpion-grass, <i>Myosotis</i>	Nasturtion, <i>Tropaeolum</i>
	Navel-wort, <i>Cotyledon</i>
Mouse-tail, <i>Myosorus</i>	Navel-wort, Bastard, <i>Crassula</i>
Mugweed, <i>Valantia cru.</i>	Navel-wort, False, <i>Crassula</i>
Mug-wort, <i>Artemisia</i>	Navel-wort, Venus's, <i>Cynoglossum</i>

Navel-wort, Water, <i>Hydrocotyle</i>	Nut, Hazel, <i>Corylus</i>
Navew, <i>Brassica</i>	Nut, Malabar, <i>Justicia</i>
Nectarine, <i>Amygdalus</i>	Nut, Pease Earth, <i>Lathyrus</i>
Nep, <i>Nepeta</i>	Nut, Physic, <i>Jatropha</i>
Nettle, <i>Urtica</i>	Nut, Physic, <i>Croton</i>
Nettle, Dead, <i>Lamium</i>	Nut, Pig, <i>Bunium</i>
Nettle, Hedge, <i>Galeopsis</i>	Nut, Pistacia, <i>Pistacia</i>
Nettle, Shrubby Hedge, <i>Prasium</i>	Nut, Purging, <i>Croton</i>
Nettle-tree, <i>Celtis</i>	Nut, Purging, <i>Jatropha</i>
Network, <i>Eriocaulon dec.</i>	Nut, Spanish, <i>Iris</i>
Nickar-tree, <i>Guilandina</i>	Nut, Walnut, <i>Juglans</i>
Nightshade, <i>Solanum</i>	
Nightshade, American, <i>Phyto-</i>	O
<i>lacca</i>	Oak, <i>Quercus</i>
Nightshade, American, <i>Rivina</i>	Oak, Dwarf, <i>Teucrium</i>
Nightshade, Bastard, <i>Rivina</i>	Oak of Cappadocia, <i>Ambrosia</i>
Nightshade, Deadly, <i>Atropa</i>	Oak of Jerusalem, <i>Chenopodium</i>
Nightshade, Enchanter's, <i>Circæa</i>	Oak, Poison; <i>Rhus</i>
Nightshade, Malabar, <i>Basella</i>	Oats, <i>Avena</i>
Nightshade, Three-leaved, <i>Tril-</i>	Oats, Seaside, of Carolina, <i>Uniola</i>
<i>lium</i>	Oats, Wild-bearded, <i>Bromus</i>
Nipple-wort, <i>Lapsana</i>	Oat-grass, <i>Bromus</i>
Noli me tangere, <i>Impatiens</i>	Oil Nut, <i>Ricinus</i>
Noli me tangere, <i>Momordica</i>	Oil Seed, <i>Ricinus</i>
None so pretty, <i>Saxifraga</i>	Oil-tree, <i>Ricinus</i>
Nonsuch, <i>Lychnis</i>	Oily Purging Grain, <i>Sesamum</i>
Nose-bleed, <i>Achillea</i>	Okra, <i>Hibiscus</i>
Nut-tree, <i>Corylus</i>	Old Man's Beard, <i>Clematis</i>
Nut, Bladder, <i>Staphylæa</i>	Old Man's head, <i>Dianthus</i>
Nut, Cashew, <i>Anacardium</i>	Oleander, <i>Nerium</i>
Nut, Chocolate, <i>Theobroma</i>	Oleaster, <i>Elæagnus</i>
Nut, Cob, <i>Corylus</i>	Olive, <i>Olea</i>
Nut, Cocoa, <i>Cocos</i>	Olive, Spurge, <i>Daphne</i>
Nut, Earth, <i>Bunium</i>	Olive, Wild, <i>Elæagnus</i>
Nut, Fausel, <i>Areca</i>	Olive, Wild, of Barbadoes,
Nut, Ground, <i>Arachis</i>	<i>Bontia</i>

One Berry, <i>Paris</i>	Palm, Common or Greater, or
One Blade, <i>Convallaria</i>	Date-tree, <i>Phoenix</i>
Onion, <i>Allium</i>	Palm, Lesser or Dwarf, <i>Chamæ-</i>
Onion, Sea, <i>Scilla</i>	<i>rops</i>
Orange, <i>Citrus</i>	Palm, The Cocoa Nut, <i>Cocos</i>
Orange, Mock, <i>Philadelphus</i>	Palm, the Fausel Nut, <i>Areca</i>
Origany, <i>Origanum</i>	Palm, Malabar, called Ampana
Oroonoka, <i>Nicotiana</i>	and Carimpana, <i>Borassus</i>
Orpine, <i>Sedum</i>	Palm, Wild Malabar, called Ka-
Orpine, Bastard, <i>Abrachne</i>	<i>ton Indel, Elate</i>
Orpine, Lesser, <i>Crassula</i>	Palm, Mountain, with largest
Orpine, True of Imperatus, <i>Te-</i>	Leaves, called Codda Panna,
<i>lephium</i>	<i>Corypha</i>
Orrach, <i>Atriplex</i>	Palm with ringed Stems, called
Orrach, Berry-bearing, <i>Blitum</i>	<i>Todda Panna, Cycas</i>
Orrach, Creeping shrubby, <i>Atra-</i>	Palm with bipennate Leaves,
<i>phaxis</i>	called Schunda Panna, <i>Cary-</i>
Orrach, Wild, <i>Chenopodium</i>	<i>ota</i>
Osier, <i>Salix</i>	Palma Christi, <i>Ricinus</i>
Osmund Royal, <i>Osmunda</i>	Palmetto, <i>Chamærops</i>
Ox-eye, <i>Bupthalmum</i>	Panic, <i>Panicum</i>
Ox-eye of old Authors, <i>Anthemis</i>	Panic-grass, <i>Panicum</i>
Ox-eye Daisy, <i>Chrysanthemum</i>	Pansies, <i>Viola</i>
Ox-lips, <i>Primula</i>	Papaw-tree, <i>Carica</i>
Ox-tongue, <i>Picris</i>	Papaw-tree of North America,
Oswego Tea, <i>Monarda</i>	<i>Annona</i>
Oyster-green, <i>Ulva luc</i>	Paraguay Tea, <i>Ilex</i>
	Park-leaves, <i>Hypericum</i>
	Parsley, <i>Apium</i>
	Parsley, Bastard, <i>Caucilis</i>
	Parsley, Corn, <i>Sison</i>
	Parsley, Fool's, <i>Æthusa</i>
	Parsley, Macedonian, <i>Bubon</i>
	Parsley, Milky, <i>Selinum</i>
	Parsley, Mountain, <i>Athamanta</i>

P

Parsley, Stone, <i>Bubon</i>	Pear, Garlick, <i>Cratæva</i>
Parsley, Bastard Stone, <i>Sison</i>	Pear, Prickly, <i>Cactus</i>
Parsley, Wild, <i>Sison</i>	Pearl-wort, <i>Sagina</i>
Parsley, Wild of America, <i>Cardiospermum</i>	Pellitory, <i>Parietaria</i>
Parsley, Break-stone, <i>Aphanes</i>	Pellitory, Bastard, <i>Achillea</i>
Parsley Piert, <i>Aphanes</i>	Pellitory, Double, <i>Achillea</i>
Parsnep, <i>Pastinaca</i>	Pellitory of Spain, <i>Anthemis</i>
Parsnep, Cow's, <i>Heracleum</i>	Pellitory of Spain, False, <i>Chrysanthemum</i>
Parsnep, Prickly, <i>Echinophora</i>	Pellitory-tree, <i>Zanthoxylum</i>
Parsnep, Water, <i>Sium</i>	Pellitory of the Wall, <i>Parietaria</i>
Parnassus, Grass of, <i>Parnassia</i>	Pennyroyal, <i>Mentha</i>
Pasque-flower, <i>Anemone</i>	Pennyroyal, Virginian, <i>Satureia</i>
Passion-flower, <i>Passiflora</i>	Pennywort, Marsh, <i>Hydrocotyle</i>
Patience, <i>Rumex</i>	Pennywort, Wall, <i>Cotyledon</i>
Paul's Betony, <i>Veronica</i>	Pennywort, Water, <i>Hydrocotyle</i>
Pea, <i>Pisum</i>	Penguin, <i>Bromelia</i>
Pea, Chich, <i>Cicer</i>	Penstemon, <i>Chelone</i>
Pea, Chickling, <i>Lathyrus</i>	Peony. See Pæony
Pea, Earth-nut, <i>Lathyrus</i>	Pepper, <i>Piper</i>
Pea, Everlasting, <i>Lathyrus</i>	Pepper, Barbary, <i>Capsicum</i>
Pea, Heart, <i>Cardiospermum</i>	Pepper, Bell, <i>Capsicum</i>
Pea, Heath, <i>Orobus</i>	Pepper, Bird, <i>Capsicum</i>
Pea, Painted Lady, <i>Lathyrus</i>	Pepper, Bonnet, <i>Capsicum</i>
Pea, Pigeon, <i>Cytisus</i>	Pepper, Guinea, <i>Capsicum</i>
Pea, Sweet-scented, <i>Lathyrus</i>	Pepper, Jamaica, <i>Myrtus</i>
Pea, Tangier, <i>Lathyrus</i>	Pepper, Indian, <i>Capsicum</i>
Pea, Winged, <i>Lotus</i>	Pepper, Long, <i>Piper</i>
Pea, Wood, <i>Orobus</i>	Pepper, Poor Man's, <i>Lepidium</i>
Peach, <i>Amygdalus</i>	Pepper, Wall, <i>Sedum</i>
Peach, Wolf's, <i>Solanum</i>	Pepper, Water, <i>Polygonum</i>
Pear, <i>Pyrus</i>	Pepper-grass, <i>Pilularia</i>
Pear, Avocado, Avocado, or Alligator, <i>Laurus</i>	Pepper-pot, <i>Capsicum</i>
Pear, Bachelor's, <i>Solanum</i>	Pepper-tree, <i>Vitis</i>
	Pepper-wort, <i>Lepidium</i>

Percepier, <i>Aphanes</i>	Pink, <i>Dianthus</i>
Periwinkle, <i>Vinca</i>	Pink, Indian, <i>Ipomœa</i>
Persicaria, <i>Polygonum</i>	Pink, Indian, <i>Lonicera</i>
Pestilent-wort, <i>Tussilago</i>	Pink, Sea, <i>Statice</i>
Petroseline Wortle, <i>Apium</i>	Pinpillow. See Pimpillo
Petty Madder, <i>Crucianella</i>	Pipe-tree, <i>Syringa</i>
Petty Whin, <i>Ononis</i>	Pipe-tree, Pudding, <i>Cassia</i>
Pharoah's Fig, <i>Musa</i>	Piperidge Bush, <i>Berberis</i>
Pharoah's Fig, <i>Ficus</i>	Pippen, <i>Pyrus</i>
Pheasant's Eye, <i>Adonis</i>	Piquets, <i>Dianthus</i>
Phyllyrea, False, <i>Rhamnus</i>	Pishamin Plum, <i>Diospyros</i>
Phu, <i>Valeriana</i>	Pistacia Nut, <i>Pistacia</i>
Physic, Nut, <i>Jatropha</i>	Pistacia-tree, Black Virginian,
Physic, Nut, <i>Croton</i>	<i>Hamamelis</i>
Physic, Pork, <i>Phytolacca</i>	Pistacia, Hazel-leaved, <i>Hama-</i>
Pick-tooth, <i>Daucus</i>	<i>melis</i>
Pigeon Pea, <i>Cytisus</i>	Pitch-tree, <i>Pinus</i>
Pig Nut, <i>Bunium</i>	Plane-tree, <i>Platanus</i>
Pig Nut, <i>Juglans</i>	Plane-tree, False, <i>Acer</i>
Pilewort, <i>Ranunculus</i>	Plant, Burning Thorny, <i>Euphor-</i>
Pimento, <i>Myrtus</i>	<i>bia</i>
Pimpernel, <i>Anagallis</i>	Plant, Egg, <i>Solanum</i>
Pimpernel, Water, <i>Veronica</i>	Plant, Humble, <i>Mimosa</i>
Pimpernel, Round-leaved Wa-	Plant, Sensitive, <i>Mimosa</i>
ter, <i>Samolus</i>	Plant, Bastard Sensitive, <i>Æschy-</i>
Pimpernel, Yellow, of the	<i>nomene</i>
Woods, <i>Lysimachia</i>	Plantain, <i>Plantago</i>
Pimpillo, <i>Cactus</i>	Plantain, Water, <i>Alisma</i>
Pinaster, <i>Pinus</i>	Plantain, Least Water, <i>Limosella</i>
Pine-tree, <i>Pinus</i>	Plantain, Star-headed Water,
Pine, Ground, <i>Teucrium</i>	<i>Alisma</i>
Pine, Stinking Ground, <i>Campho-</i>	Plantain Shot, <i>Canna</i>
<i>rosma</i>	Plantain-tree, <i>Musa</i>
Pine, Heath-low, <i>Coris</i>	Pliant Mealy-tree, <i>Viburnum</i>
Pine-apple, <i>Bromelia</i>	Plowman's Spikenard, <i>Baccharis</i>
Pine-apple, Wild, <i>Renealmia</i>	Plowman's Spikenard, <i>Conyza</i>

Plum-tree, <i>Prunus</i>	Potatoe, Indian, <i>Dioscorea</i>
Plum, American Black, <i>Chryso-</i> <i>balanus</i>	Potatoe, Spanish, <i>Convolvulus</i>
Plum, Bay, <i>Psidium</i>	Prick Wood, <i>Euonymus</i>
Plum, Brazilian, <i>Spondias</i>	Primrose, <i>Primula</i>
Plum, Cocoa, <i>Chrysobalanus</i>	Primrose, Night, <i>Cenothera</i>
Plum, Hog, <i>Spondias</i>	Primrose, Peerless, <i>Narcissus</i>
Plum, Indian Date, <i>Diospyros</i>	Primrose-tree, <i>Cenothera</i>
Plum, Maiden, <i>Chrysobalanus</i>	Prince's Feather, <i>Amaranthus</i>
Plum, Pishamin, Persimon, or Pitchumon, <i>Diospyros</i>	Privet, <i>Ligustrum</i>
Poccoon. See Fuccoon	Privet, Evergreen, <i>Rhamnus</i>
Pockwood, <i>Guaiacum</i>	Privet, Mock, <i>Phillyrea</i>
Poet's Cassia, <i>Osyris</i>	Privy-saugh, <i>Ligustrum</i>
Poet's Rosemary, <i>Osyris</i>	Puccoon, <i>Sanguinaria</i>
Poison Ash, <i>Rhus</i>	Pudding-grass, <i>Meutha</i>
Poison Berry, <i>Cestrum</i>	Pudding Pipe-tree, <i>Cassia</i>
Poison Bush, <i>Euphorbia</i>	Puff-balls, <i>Lycoperdon bov.</i>
Poison Oak, <i>Rhus</i>	Pumpion. See Pompion
Poison Tree, <i>Rhus</i>	Pumpkin. See Pompion
Poke, Virginian, <i>Phytolacca</i>	Purging Grain, Oily, <i>Sesamum</i>
Poley, Mountain, <i>Teucrium</i>	Purging Nut, <i>Croton</i>
Poley, Grass, <i>Lythrum</i>	Purging Nut, <i>Jatropha</i>
Polypody, <i>Polypodium</i>	Purging Thorn, <i>Rhamnus</i>
Pomegranate, <i>Punica</i>	Purple Apple, <i>Amoma</i>
Pompion, <i>Cucurbita</i>	Purslane, <i>Portulaca</i>
Pond-weed, <i>Potamogeton</i>	Purslane, Horse, <i>Trianthema</i>
Pond-weed, Triple-headed, <i>San-</i> <i>nichellia</i>	Purslane, Sea, <i>Atriplex</i>
Poplar, <i>Populus</i>	Purslane, Water, <i>Peplis</i>
Poppy, <i>Papaver</i>	Purslane, Tree Sea, <i>Atriplex</i>
Poppy, Horned, <i>Chelidonium</i>	
Poppy, Prickly, <i>Argemone</i>	Q
Poppy, Spatling, <i>Cucubalus</i>	Quamoclit, <i>Ipomœa</i>
Pork, Physic, <i>Phytolacca</i>	Queen of the Meadows, <i>Spirœa</i>
Potatoe, <i>Solanum</i>	Queen's July-flower, <i>Hesperis</i>
	Queen's Violet, <i>Hesperis</i>
	Quick, <i>Cratagus</i>
	Quicken, <i>Sorbus</i>

Quickbean-tree, <i>Sorbus</i>	Red Whorts, Spanish, <i>Arbutus</i>
Quince-tree, <i>Pyrus</i>	Reddish. See Radish
Quill-wort, <i>Isotes lac.</i>	Reed, <i>Arundo</i>
	Reed, Burr, <i>Sparganium</i>
	Reed, Indian Flowering, <i>Canna</i>
	Reed Mace, <i>Typha</i>
	Rennet, Cheese, <i>Galium</i>
	Rest Harrow, <i>Ononis</i>
	Rhamnus, Bastard, <i>Hippophaë</i>
	Rhendecr, <i>Lichen ran.</i>
	Rhubarb, <i>Rheum</i>
	Rhubarb, Monk's, <i>Rumex</i>
	Ribwort, <i>Plantago</i>
	Rice, <i>Oryza</i>
	Ricinus, Bastard, <i>Croton</i>
	Roane-tree, <i>Sorbus</i>
	Robert, Herb, <i>Geranium</i>
	Rocamboles, <i>Allium</i>
	Rock Germander, <i>Veronica</i>
	Rock Rose, <i>Cistus</i>
	Rocket, <i>Brassica</i>
	Rocket, Bastard, <i>Reseda</i>
	Rocket, Corn, <i>Bunias</i>
	Rocket, Marsh, <i>Sisymbrium</i>
	Rocket, Sea, <i>Bunias</i>
	Rocket, Square-codded, of Montpelier, <i>Bunias</i>
	Rocket, Water, <i>Sisymbrium</i>
	Rocket, Winter, <i>Sisymbrium</i>
	Rocket, or Dame's Violet, <i>Hesperis</i>
	Rod, Aaron's, <i>Solidago</i>
	Rod, Golden, <i>Solidago</i>
	Rod-tree, Golden, <i>Bosea</i>
	Rod, Shepherd's, <i>Dipsacus</i>
	Roe-buck Berries, <i>Rubus sax.</i>
R	
Radish, <i>Raphanus</i>	
Radish, Horse, <i>Cochlearia</i>	
Radish, Water, <i>Sisymbrium</i>	
Ragged Robin, <i>Lychnis</i>	
Ragwort, Common, <i>Senecio</i>	
Ragwort, African, <i>Othonna</i>	
Ragworts, Sundry, of old Authors, <i>Senecio</i>	
Ragworts, Sundry, of old Authors, <i>Solidago</i>	
Rampions, Horned, <i>Phyteuma</i>	
Rampions, Crested, <i>Lobelia</i>	
Rampions, Common Esculent, <i>Campanula</i>	
Rampions with scabious Heads, <i>Jasione</i>	
Ramsons, <i>Allium</i>	
Ranunculus, Globe, <i>Trollius</i>	
Rape, <i>Brassica</i>	
Rape, Broom, <i>Orobanchè</i>	
Rape, Cole, <i>Brassica</i>	
Rape of Cistus, <i>Asarum</i>	
Raspberry, <i>Rubus</i>	
Rattle, <i>Pedicularis</i>	
Rattle, Yellow, <i>Rhinanthus</i>	
Rattlesnake-Root, Senegaw, <i>Polygala</i>	
Rattlesnake-Root, Dr. Witts's, <i>Prenanthes</i>	
Rattlesnake-Weed, <i>Eryngium</i>	
Redbud, <i>Cercis</i>	

Root, Indian Arrow, <i>Maranta</i>	Rue, <i>Ruta</i>
Root, China, <i>Smilax</i>	Rue, Dog's, <i>Scrophularia</i>
Root, False China, <i>Senecio</i>	Rue, Goat's, <i>Galega</i>
Root, Fever, <i>Triosteum</i>	Rue, Meadow, <i>Thalictrum</i>
Root, Hollow, <i>Adoxa</i>	Rue, Wall, <i>Asplenium</i>
Root, Rose, <i>Rhodiola</i>	Rue, Wild Syrian, <i>Peganum</i>
Root, Snake, <i>Aristolochia</i>	Rupture-wort, <i>Herniaria</i>
Root, Snake, black or wild, of America, <i>Actæa</i>	Rupture-wort, Least, <i>Linum</i>
Root, Dr. Witts's Rattlesnake, <i>Prenanthes</i>	Rush, <i>Juncus</i>
Root, Senegaw Rattlesnake, <i>Po- lygala</i>	Rush, Flowering, <i>Butomus</i>
Root, Sweet, <i>Glycyrrhiza</i>	Rush, Lesser flowering, <i>Scheuch- zeria</i>
Rose, <i>Rosa</i>	Rush, Round, black-headed, Marsh or Bog, <i>Schænus</i>
Rose, China, <i>Hibiscus</i>	Rush, Sweet, <i>Acorus</i>
Rose, Christmas, <i>Helleborus</i>	Rush-grass, <i>Scirpus</i>
Rose, Corn, <i>Papaver</i>	Ruyschiana, <i>Dracocephalon</i>
Rose, Gelder, <i>Viburnum</i>	Rye, <i>Secale</i>
Rose, Gelderland, <i>Viburnum</i>	Rye, Wild, <i>Hordeum</i>
Rose, Virginian Gelder, <i>Spiræa</i>	Rye-grass, <i>Hordeum</i>
Rose, Martinico, <i>Hibiscus</i>	S
Rose, Rock, <i>Cistus</i>	Saffron, <i>Crocus</i>
Rose of Jericho, <i>Anastatica</i>	Saffron, Bastard, <i>Carthamus</i>
Rose Bay, <i>Nerium</i>	Saffron, Meadow, <i>Colchicum</i>
Rose Bay Dwarf, <i>Rhododendrum</i>	Sage, <i>Salvia</i>
Rose Bay, Mountain, <i>Rhodo- dendrum</i>	Sage, Wild, <i>Teucrium</i>
Rose Bay Willow-herb, <i>Epilobium</i>	Sage, Indian Wild, <i>Lantana</i>
Rose Mallow, <i>Alcea</i>	Sage, Wood, <i>Teucrium</i>
Rose Root, <i>Rhodiola</i>	Sage of Jerusalem, <i>Pulmonaria</i>
Rosemary, <i>Rosemarinus</i>	Sage of Jerusalem, <i>Phlomis</i>
Rosemary, Poet's, <i>Osyris</i>	Sage-tree, <i>Phlomis</i>
Rosemary, Wild, <i>Ledum</i>	Saint Bartholomew's Herb, <i>Ilex</i>
Rosemary, Lesser Wild, <i>Andro- meda</i>	Saint Bruno's Lily, <i>Hemerocallis</i>
	Saint John's Bread, <i>Ceratonia</i>
	Saint John's-wort, <i>Hypericum</i>

Saint Peter's-wort, <i>Ascyrum</i>	Saunders, <i>Santalum</i>
Saint Peter's-wort, <i>Hypericum</i>	Savory, <i>Satureia</i>
Saint Peter's-wort, Shrubby, <i>Lonicera</i>	Savoys, <i>Brassica</i>
Saintfoin, <i>Hedysarum</i>	Saw-wort, <i>Serratula</i>
Sallad, Corn, <i>Valeriana</i>	Saxifrage, <i>Saxifraga</i>
Sal-kali, <i>Salicornia</i>	Saxifrage, Burnet, <i>Pimpinella</i>
Sallow, <i>Salix</i>	Saxifrage, Golden, <i>Chrysosplenium</i>
Salsafy, <i>Tragopogon</i>	Saxifrage, Meadow, <i>Peucedanum</i>
Salt-wort, <i>Salicornia</i>	Scabious, <i>Scabiosa</i>
Salt-wort, Black, <i>Glaur</i>	Scabious, Sheep's, <i>Jasione</i>
Samphire, <i>Crithmum</i>	Scallion, <i>Allium</i>
Samphire, Golden, <i>Inula</i>	Scammony, Syrian, <i>Convolvulus</i>
Sand-box Tree, <i>Hura</i>	Scammony of Montpellier, <i>Cynanchum</i>
Sanders. See Saunders	Sciatica Cress, the True, <i>Lepidium</i>
Sanicle, <i>Sanicula</i>	Sciatica Cress, <i>Iberis</i>
Sanicle, <i>Saxifraga</i>	Scorching Fennel, <i>Thapsia</i>
Sanicle, American Bastard, <i>Mitella</i>	Scorpion-grass, <i>Scorpiurus</i>
Sanicle, Bear's-ear, <i>Cortusa</i>	Scorpion-grass, Mouse-ear, <i>Myosotis</i>
Sappadillo-tree, <i>Sloanea</i>	Scorpion Senna, <i>Coronilla</i>
Sapota, <i>Achras</i>	Scorpion's Thorn, <i>Ulex</i>
Sapota Mammee, <i>Achras</i>	Screw-tree. See Skrew-tree
Saracen's Consound, <i>Solidago</i>	Scull-cap. See Skull-cap
Saracen's Consound, the True, <i>Senecio</i>	Scurvy-grass, <i>Cochlearia</i>
Saracen's Wound-wort, <i>Solidago</i>	Sea-beard, <i>Conferva rup.</i>
Saracen's Wound-wort, the True, <i>Senecio</i>	Sea-Weed, <i>Fucus</i>
Sassafras-tree, <i>Laurus</i>	Sebesten, <i>Cordia</i>
Sassafy. See Salsafy	Sedum Pyramidal, <i>Saxifraga</i>
Satin-flower, <i>Lunaria</i>	Seed, Heart, <i>Cardiospermum</i>
Satin, White, <i>Lunaria</i>	Segs, <i>Iris pseu.</i>
Sauce alone, <i>Erysimum</i>	Self-heal, <i>Brunella</i>
Savin, <i>Juniperus</i>	Self-heal, <i>Sanicula</i>
Savin-tree, Indian, <i>Bauhinia</i>	Senna of the Shops, <i>Cassia</i>
	Senna, Bastard, <i>Cassia</i>

Senna, Bastard, <i>Colutea</i>	Silk, Virginian, <i>Periploca</i>
Senna, Jointed-podded Bladder, <i>Coronilla</i>	Silver Bush, <i>Anthyllis</i>
Senna, Scorpion, <i>Coronilla</i>	Silver-tree, <i>Prosea</i>
Senna, Wild, <i>Cassia</i>	Silver-weed, <i>Potentilla</i>
Senegaw Rattlesnake Root, <i>Polygala</i>	Simpla Nobla, <i>Phyllis</i>
Sengreen, <i>Semprevivum</i>	Simpler's Joy, <i>Verbena</i>
Sensitive Plant, <i>Mimosa</i>	Skirret, <i>Sium</i>
Sensitive Plant, Bastard, <i>Eschynomene</i>	Skull-cap, <i>Scutellaria</i>
Septfoil, <i>Tormentilla</i>	Skrew-tree, <i>Helicteres</i>
Sermountain, <i>Laserpitium</i>	Sloe-tree, <i>Prunus</i>
Serpent Cucumber, <i>Trichosanthes</i>	Sloke, <i>Uta</i>
Serpent's Tongue, <i>Ophioglossum</i>	Smallage, <i>Apium</i>
Service-tree, <i>Sorbus</i>	Snails, <i>Medicago</i>
Service, Maple-leaved, <i>Cratægus</i>	Snail Clover, <i>Medicago</i>
Service, Wild, <i>Cratægus</i>	Snail Trefoil, <i>Medicago</i>
Setfoil. See Septfoil	Snakeweed, <i>Polygonum</i>
Setwall. See Zedoary	Snake-root, <i>Aristolochia</i>
Setwall, Garden, <i>Valeriana</i>	Snake-root, Black or Wild, of America, <i>Actæa</i>
Setter-wort, <i>Helleborus</i>	Snap-tree, <i>Justicia</i>
Shaddock, <i>Citrus</i>	Snap-dragon, <i>Antirrhinum</i>
Shallot. See Eschalot	Snap-dragon of America, <i>Ruellia</i>
Shavegrass, <i>Equisetum</i>	Sneeze-wort, <i>Achillea</i>
Sheep Scabious, <i>Jasione</i>	Sneeze-wort, Austrian, <i>Xeranthemum</i>
Shepherd's Needle, <i>Scandix</i>	Snowball-tree, <i>Viburnum</i>
Shepherd's Pouch, <i>Thlaspi</i>	Snowberry-bush, <i>Lonicera</i>
Shepherd's Rod, <i>Dipsacus</i>	Snowdrop, <i>Galanthus</i>
Shepherd's Staff, <i>Dipsacus</i>	Snowdrop, Greater, <i>Lencojunz</i>
Shot, Indian, <i>Canna</i>	Snowdrop-tree, <i>Chionanthus</i>
Shot, Plantain, <i>Canna</i>	Soap Apple, <i>Sapindus</i>
Sickle-wort, <i>Coronilla</i>	Soap Berry, <i>Sapindus</i>
Side-saddle Flower, <i>Sarracena</i>	Soap-wort, <i>Saponaria</i>
Silk Cotten-tree, <i>Bombax</i>	Soldanel, <i>Soldanella</i>
	Soldanel of the Shops, <i>Convolvulus</i>

Soldier, Water, <i>Stratiotes</i>	Spider-wort, Great Savoy, <i>Hemerocallis</i>
Soldier, Fresh Water, <i>Stratiotes</i>	
Soldier's Cullions, <i>Orchis</i>	Spider-wort, Virginian, <i>Tradescantia</i>
Solomon's Seal, <i>Cowallaria</i>	
Solomon's Seal, Pennsylvanian, <i>Uvularia</i>	Spignel, <i>Athamanta</i>
	Spignel, Wild, <i>Seseli</i>
Sorgo, <i>Holcus</i>	Spike-grass, Winged, <i>Stipa</i>
Sorrel, <i>Rumex</i>	Spikenard, Indian, or True*
Sorrel, Indian Red, <i>Hibiscus</i>	Spikenard, Bastard French, <i>Nardus</i>
Sorrel, Indian White, <i>Hibiscus</i>	
Sorrel, Wood, <i>Oxalis</i>	Spikenard, Celtic, <i>Valeriana</i>
Sorrel-tree, <i>Andromeda</i>	Spikenard, False, <i>Lavandula</i>
Sorrowful-tree, <i>Nyctanthes</i>	Spikenard, Plowman's, <i>Baccharis</i>
Sour Gourd, Æthiopian, <i>Adansonia</i>	
	Spikenard, Plowman's, <i>Coryza</i>
Sour Soap, <i>Annona</i>	Spikenard, Wild, <i>Asarum</i>
Southernwood, <i>Artemisia</i>	Spinach, <i>Spinacia</i>
South-sea Tea, <i>Ilex</i>	Spinach, Strawberry, <i>Blitum</i>
Sow-bread, <i>Cyclamen</i>	Spindle-tree, <i>Euonymus</i>
Sow-ruck, <i>Rumex acet.</i>	Spindle-tree, Climbing, <i>Celastrus</i>
Sow Thistle, <i>Sonchus</i>	Spindle-tree, Bastard, <i>Kiggellaria</i>
Sow Thistle, <i>Prenanthes</i>	Spindle-tree, Bastard, <i>Celastrus</i>
Sow Thistle, Downy, <i>Andryala</i>	Spiræa Frutex, <i>Spiræa</i>
Sow Thistle, Tangier, <i>Scorzonera</i>	Spiræa, African, <i>Diosma</i>
Sparrow-grass. See Asparagus	Spirting Cucumber, <i>Momordica</i>
Sparrow-wort, <i>Passerina</i>	Spleen-wort, <i>Asplenium</i>
Sparrow-wort, Tragus's, <i>Stellera</i>	Spleen-wort, Rough, <i>Lonchitis</i>
Spatling Poppy, <i>Cucubalus</i>	Spleen-wort, Rough, <i>Polypodium</i>
Spear-wort, <i>Ranunculus</i>	Spoon-wort, <i>Cochlearia</i>
Speerage. See Asparagus	Sponge, <i>Spongia</i>
Speedwell, <i>Veronica</i>	Sponge-tree, <i>Biemosa</i>
Speedwell, Female, <i>Antirrhinum</i>	Spurge, <i>Euphorbia</i>
Spice Wood, <i>Laurus</i>	Spurge, Bastard, <i>Euphorbia</i>
Spice, All, <i>Myrtus</i>	Spurge Laurel, <i>Daphne</i>
Spider-wort, <i>Anthericum</i>	Spurge Olive, <i>Daphne</i>

*Unknown.

Spurrey, <i>Spergula</i>	Strawberry, Barren, <i>Potentilla</i>
Squash, <i>Cucurbita</i>	Strawberry, Barren, <i>Fragaria</i>
Squill, <i>Scilla</i>	Strawberry Blite, <i>Blitum</i>
Squill, Lesser White, <i>Pancreatum</i>	Strawberry Spinach, <i>Blitum</i>
Staff-tree, <i>Celastrus</i>	Strawberry-tree, <i>Arbutus</i>
Staff, Shepherd's, <i>Dipsacus</i>	Succory, <i>Cichorium</i>
Stag's-horn-tree, <i>Rhus</i>	Succory, Gum, <i>Chondrilla</i>
Star of Alexandria, <i>Ornithogalum</i>	Succory, Wart, <i>Lapsana</i>
	Sugar Cane, <i>Saccharum</i>
Star Apple, <i>Chrysophyllum</i>	Sulphur-wort, <i>Peucedanum</i>
Star of Bethlehem, <i>Ornithogalum</i>	Sultan-flower, <i>Centaurea</i>
Star of Constantinople, <i>Ornithogalum</i>	Sumach, <i>Rhus</i>
	Sumach, Myrtle-leaved, <i>Coriaria</i>
Star Hyacinth, <i>Scilla</i>	Sumach, Tanner's or Currier's, <i>Coriaria</i>
Star of Naples, <i>Ornithogalum</i>	
Star Thistle, <i>Centaurea</i>	Sundew, <i>Drosera</i>
Star-wort, <i>Aster</i>	Sun-flower, <i>Helianthus</i>
Star-wort, Bastard, <i>Bupthalmum</i>	Sun-flower, Bastard, <i>Helenia</i>
Star-wort, Trailing, of Vera-Cruz, <i>Tridax</i>	Sun-flower, Dwarf, <i>Rudbeckia</i>
Star-wort, Yellow, <i>Inula</i>	Sun-flower, Dwarf, <i>Tetragonotheca</i>
Star-wort, Yellow, <i>Bupthalmum</i>	Sun-flower, Little, <i>Cistus</i>
Staves Acre, <i>Delphinium</i>	Sun-flower, Tick-seeded, <i>Coreopsis</i>
Stich-wort, <i>Stellaria</i>	Sun-flower, Willow-leaved, <i>Helenia</i>
Stink-horns, <i>Phallus</i>	
Stock, <i>Cheiranthus</i>	Superb Lily, <i>Gloriosa</i>
Stock July-flower, <i>Cheiranthus</i>	Swallow-wort, <i>Asclepias</i>
Stock, Dwarf Annual, <i>Hesperis</i>	Sweet Briar, <i>Rosa</i>
Stock, Virginian, <i>Hesperis</i>	Sweet Cicely, <i>Scandix</i>
Stone-crop, <i>Sedum</i>	Sweet Gum, <i>Liquidambar</i>
Stone-crop-tree, <i>Chenopodium</i>	Sweet John, <i>Dianthus</i>
Stone Parsley, <i>Bubon</i>	Sweet Root, <i>Glycyrrhiza</i>
Stone Parsley, Bastard, <i>Sison</i>	Sweet Sop, <i>Annona</i>
Storax-tree, <i>Styrax</i>	Sweet Sultan, <i>Centaurea</i>
Storax, Liquid, <i>Liquidambar</i>	Sweet Weed, <i>Capraria</i>
Strawberry, <i>Fragaria</i>	

Sweet William, <i>Dianthus</i>	Thistle, Golden, <i>Scolymus</i>
Sweet William of Barbadoes, <i>Ipomœa</i>	Thistle, Hedge-hog, <i>Cactus</i>
Swine's Cress, <i>Cochlearia</i>	Thistle, Holy, <i>Cnicus</i>
Sycamore, <i>Ficus</i>	Thistle, Ladies', <i>Carduus</i>
Sycamore, False, <i>Acer</i>	Thistle, Melancholy, <i>Carduus</i>
Syringa, commonly called, <i>Philadelphus</i>	Thistle, Melon, <i>Cactus</i>
	Thistle, Milk, <i>Carduus</i>
	Thistle, Soft, <i>Carduus</i>
	Thistle, Sow, <i>Sonchus</i>
	Thistle, Sow, <i>Prenanthes</i>
	Thistle, Downy Sow, <i>Andryala</i>
	Thistle, Star, <i>Centaurea</i>
	Thistle, Torch, <i>Cactus</i>
	Thistle, Woolly, <i>Onopordon</i>
	Thongs, <i>Fucus</i>
	Thorn, Black, <i>Prunus</i>
	Thorn, Box, <i>Lycium</i>
	Thorn, Christ's, <i>Rhamnus</i>
	Thorn, Egyptian, <i>Mimosa</i>
	Thorn, Evergreen, <i>Mespilus</i>
	Thorn, Goat's, <i>Astragalus</i>
	Thorn, Lily, <i>Catesbœa</i>
	Thorn, Purging, <i>Rhamnus</i>
	Thorn, Scorpion's, <i>Ulex</i>
	Thorn, Spanish Hedge-hog, <i>Anthyllis</i>
	Thorn, White, <i>Cratagus</i>
	Thorn, Apple, <i>Datura</i>
	Thorny Plant, Burning, <i>Euphorbia</i>
	Thorough Wax, <i>Bupleurum</i>
	Three Faces under a Hood, <i>Viola</i>
	Three-leaved Grass, <i>Trifolium</i>
	Thrift, <i>Statice</i>
	Throat-wort, Blue umbelliferous, <i>Trachelium</i>

Throat-wort, <i>Campanula</i>	Trefoil, Shrub, <i>Ptelea</i>
Thyme, <i>Thymus</i>	Trefoil, Snail, <i>Medicago</i>
Thyme, Dodder of, <i>Cuscuta</i>	Trefoil, Thorny, of Candia, <i>Fa-</i> <i>gonia</i>
Thyme, Mastick, <i>Satureia</i>	Trefoil Tree, <i>Cytisus</i>
Tickseed, <i>Corispermum</i>	Trefoil. Base-tree, <i>Cytisus</i>
Tills, <i>Ervum</i>	Trinity Herb, <i>Viola</i>
Timothy-grass, <i>Phleum</i>	Triple Ladies' Traces, <i>Ophrys</i>
Tinker's Weed, <i>Triosteum</i>	True-love, <i>Paris</i>
Toad Flax, <i>Antirrhinum</i>	True-love of Canada, <i>Trillium</i>
Tobacco, <i>Nicotiana</i>	Truffles, <i>Lycoperdon tub.</i>
Tolu-tree, Balsam of, <i>Toluifera</i>	Trumpet-flower, <i>Bignonia</i>
Tomatoes, <i>Solanum</i>	Tuberose, <i>Polyanthes</i>
Tooth-ach-tree, <i>Zanthoxylum</i>	Tulip, <i>Tulipa</i>
Tooth-pick, <i>Daucus</i>	Tulip, African, <i>Hæmanthes</i>
Tooth-wort, <i>Dentaria</i>	Tulip, Chequered, <i>Fritillaria</i>
Tooth-wort, <i>Plumbago</i>	Tulip-flower, <i>Bignonia</i>
Torch Thistle, <i>Cactus</i>	Tulip-tree, <i>Liriodendrum</i>
Tormentil, <i>Tormentilla</i>	Tulip-tree, Laurel-leaved, <i>Mag-</i> <i>nolia</i>
Touch me not, <i>Impatiens</i>	Tun-hoof, <i>Glechoma</i>
Touch me not, <i>Momordica</i>	Tapelo-tree, <i>Nyssa</i>
Tower Mustard, <i>Turrîtis</i>	Turbith Indian, or of the Shops, <i>Convolvulus</i>
Tower Mustard, Bastard, <i>Arabis</i>	Turbith, Garganic, <i>Thapsia</i>
Tragacanth, Gum, <i>Astragalus</i>	Turkey-feather, <i>Ulva par.</i>
Tragus's Sparrow-wort, <i>Stellera</i>	Turk's Cap, <i>Lilium</i>
Traveller's Joy, <i>Clematis</i>	Turk's Head, <i>Cactus</i>
Treacle Mustard, <i>Clypeola</i>	Turk's Turban, <i>Ranunculus</i>
Treacle Mustard, <i>Thlaspi</i>	Turnep, <i>Brassica</i>
Tree Moss, <i>Lichen</i>	Turnep, French, <i>Brassica</i>
Trefoil, <i>Trefolium</i>	Turmeric, <i>Curcuma</i>
Trefoil, Bean, <i>Cytisus</i>	Turnsole, <i>Heliotropium</i>
Trefoil, Stinking Bean, <i>Anagyris</i>	Turpentine-tree, <i>Pistacia</i>
Trefoil, Bird's-foot, <i>Lotus</i>	Tutsan, <i>Hypericum</i>
Trefoil, Marsh, <i>Menyanthes</i>	Two-pence, Herb, <i>Lysimachia</i>
Trefoil, Moon, <i>Medicago</i>	
Trefoil of Montpellier, Shrub, <i>Lotus</i>	

Tway Blade, <i>Ophrys</i>	Vetchling, Yellow, <i>Lathyrus</i>
Twy Blade, <i>Ophrys</i>	Viburnum, American, <i>Lantana</i>
	Vine, <i>Vitis</i>
V	Vine, Black, <i>Tamus</i>
Valerian, <i>Valeriana</i>	Vine, Climbing Five-leaved, of
Valerian, Greek, <i>Polemonium</i>	Canada, <i>Hedera</i>
Vanilla, or Vaneloe, <i>Epidendrum</i>	Vine, Spanish Arbor, <i>Ipomœa</i>
Vernal-grass, <i>Anthoxanthum</i>	Vine, White, <i>Bryonia</i>
Venus's Comb, <i>Scandix</i>	Violet, <i>Viola</i>
Venus's Looking-glass, <i>Camp-</i>	Violet, Bulbous, <i>Galanthus</i>
<i>nula</i>	Violet, Calathian, <i>Gentiana</i>
Venus's Navel-wort, <i>Cynoglossum</i>	Violet, Dame's, <i>Hesperis</i>
Vervain, <i>Verbena</i>	Violet, Dog's Tooth, <i>Erythronium</i>
Vervain Mallow, <i>Malva</i>	Violet, Queen's, <i>Hesperis</i>
Vetch, <i>Vicia</i>	Violet, Water, <i>Hottonia</i>
Vetch, Ax. See Hatchet Vetch	Viper's Buglos, <i>Echium</i>
Vetch, Bitter, <i>Ervum</i>	Viper's Grass, <i>Scorzonera</i>
Vetch, Bitter, <i>Orobus</i>	Virgin's Bower, <i>Clematis</i>
Vetch, Jointed-podded Bitter,	Vitæ, Arbor, <i>Thuya</i>
<i>Ervum</i>	Vitæ, Lignum, <i>Guaiacum</i>
Vetch, Chichling, <i>Lathyrus</i>	Umbrella-tree, <i>Magnolia</i>
Vetch, Crimson Grass, <i>Lathyrus</i>	
Vetch, Hatchet, <i>Coronilla</i>	W
Vetch, Clusius's Foreign Hatchet,	Wake Robin, <i>Arum</i>
<i>Biserrula</i>	Wall-flower, <i>Cheiranthus</i>
Vetch, Horse-shoe, <i>Hippocrepis</i>	Walnut, <i>Juglans</i>
Vetch, Kidney, <i>Anthyllis</i>	Walnut, Jamaica, <i>Hura</i>
Vetch Liquorice, <i>Astragalus</i>	Wall-wort, <i>Sambucus</i>
Vetch, Knobbed-rooted Liquor-	Wanhom, <i>Kæmpferia</i>
ice, <i>Glycine</i>	Ware-sea, <i>Fucus ves.</i>
Vetch, Milk, <i>Astragalus</i>	Wart Succory, <i>Lapsana</i>
Vetch, Bastard Milk, <i>Phaca</i>	Wart-wort, <i>Euphorbia</i>
Vetch, Venetian, <i>Orobus</i>	Wart-wort, <i>Heliotropium</i>
Vetch, Medic, <i>Hedysarum</i>	Wart-wort, <i>Lapsana</i>
Vetchling, <i>Hedysarum</i>	Water-leaf, <i>Hydrophyllum</i>
Vetchling, Medic., <i>Heilysarum</i>	Water Soldier, <i>Stratiotes</i>

Wayfaring-tree, <i>Viburnum</i>	Wind-flower, <i>Anemone</i>
Weld, <i>Reseda</i>	Wind-seed, <i>Arctotis</i>
Wheat, <i>Triticum</i>	Winged Spiked Grass, <i>Stipa</i>
Wheat, Buck, <i>Polygonum</i>	Winter Berry, <i>Prinos</i>
Wheat, Cow, <i>Myiampyrum</i>	Winter Bloom, <i>Azalea</i>
Wheat, French, <i>Polygonum</i>	Winter Cherry, <i>Physalis</i>
Wheat, Indian, <i>Zea</i>	Winter Cherry, <i>Solanum</i>
Wheat, Turkey, <i>Zea</i>	Winter Green, <i>Pyrola</i>
Whin, <i>Ulex</i>	Winter Green, Ivy-flowering, <i>Kalmia</i>
Whin, Petty, <i>Ononis</i>	Winter Green, with Chickweed Flowers, <i>Trientalis</i>
Whistles, Sea, <i>Fucus nod.</i>	Winter's Bark, <i>Laurus</i>
White Beam-tree, <i>Cratagus</i>	Witch Hazel, <i>Hamamelis</i>
White, Leaf-tree, <i>Cratagus</i>	Witch Hazel, <i>Ulmus</i>
White Satin, <i>Lunaria</i>	Woad, <i>Isatis</i>
White Wood, <i>Bignonia</i>	Woad, Wild, <i>Reseda</i>
Whitlow Grass, <i>Draba</i>	Wolf's Bane, <i>Aconitum</i>
Whitlow Grass, Rue-leaved, <i>Saxifraga</i>	Wolf's Bane, Winter, <i>Helleborus</i>
Whortle Berry, <i>Vaccinium</i>	Wolf's Peach, <i>Solanum</i>
Whortle Berry, African, <i>Royena</i>	Woodbind, <i>Lonicera</i>
Whorts, Black, <i>Vaccinium</i>	Woodbind, Spanish, <i>Ipomæa</i>
Whorts, Bog, <i>Vaccinium</i>	Wood of Life, <i>Guaiacum</i>
Whorts, Red, <i>Vaccinium</i>	Wood Anemone, <i>Anemone</i>
Whorts, Spanish Red, <i>Arbutus</i>	Wood Sorrel, <i>Oxalis</i>
Wicken-tree, <i>Sorbus</i>	Woodroof, <i>Asperula</i>
Widow Wail, <i>Cneorum</i>	Woodwaxen, <i>Genista</i>
Willow, <i>Salix</i>	Worm-grass, <i>Spigelia</i>
Willow, French, <i>Epilobium</i>	Worm-seed, <i>Chenopodium</i>
Willow, Spiked, of Theophras- tus, <i>Spiræa</i>	Wormwood, <i>Artemisia</i>
Willow, Sweet, <i>Myrica</i>	Wormwood, Wild, <i>Parthenium</i>
Willow, Herb, <i>Epilobium</i>	Wortle, Petroseline, <i>Apium</i>
Willow, Herb, <i>Lythrum</i>	Would, <i>Reseda</i>
Willow, Herb, <i>Lysimachia</i>	Wound-wort of Achilles, <i>Achillea</i>
Willow Herb, Rosebay, <i>Epilo- bium</i>	Wound-wort, Clown's, <i>Stachys</i>
	Wound-wort, Saracen's, <i>Solidago</i>

Wound-wort, Saracen's, the true, Yellow Weed, *Reseda*

Senecio

Yerva Mora, *Bosea*

Wrack, *Fucus*

Yew Tree, *Taxus*

Wrack, Grass, *Zostera*

Z

Y

Zedoary, Round, *Kempferia*

Yams, *Dioscorea*

Zedoary, Long, *Amomum*

Yapon, *Ilex*

Zerumbith, *Amomum*

Yarrow, *Achillea*

TABLE V.

THE NAMES OF PLANTS

IN LATIN AND FRENCH.

A

<i>Abies</i> , Sapin	<i>Actea</i> , Actée
<i>Ablania</i> , Ablania	<i>Adansonia</i> , Adansonier, Baobab
<i>Abroma</i> , Ambrôme	<i>Adenanthera</i> , Condori
<i>Abrus</i> , Abrus	<i>Adenia</i> , Adénia
<i>Acalypha</i> , Ricinelle	<i>Adiantum</i> , Adiante, Capillaire
<i>Acanthi</i> , J. Acanthes	<i>Adonis</i> , Adonis, Adonide
<i>Acanthus</i> , Acanthe	<i>Adoxa</i> , Moschatelle
<i>Acer</i> , Erable	<i>Ægilops</i> , Egilope
<i>Acera</i> , Erables	<i>Ægopodium</i> , Podagraire
<i>Achillea</i> , Achillée	<i>Æschinomene</i> , Nélitte
<i>Achras</i> , Sapotillier	<i>Æsculus</i> , Marronnier
<i>Achyranthes</i> , Cadélari	<i>Æthusa</i> , Æthuse
<i>Acnida</i> , Acnide	<i>Agallochum</i> , La M. Agalloche
<i>Aconitum</i> , Aconit	<i>Alisma</i> , Fluteau
<i>Acorus</i> , Acore	<i>Allium</i> , Ail
<i>Acotyledones</i> , Acotylédons	<i>Aloë</i> , Aloës
<i>Acrostichum</i> , Acrostique	<i>Alopecurus</i> , Vulpin

<i>Alpinia</i> , Alpinia	<i>Amethystea</i> , Amethystée
<i>Alsine</i> , Morgeline	<i>Ammannia</i> , Ammane
<i>Alstonia</i> , Alstonia	<i>Ammi</i> , Ammi
<i>Astroëmeria</i> , Pélégrine	<i>Anomum</i> , Anôme
<i>Althæa</i> , Guimauve	<i>Amorpha</i> , Amorpha
<i>Alyssum</i> , Alysson, Alysse	<i>Amygdalæa</i> , Sous-ordre des Rosacées de Amandiers
<i>Amanita</i> , Amanite	<i>Amygdalus</i> , Amandier
<i>Agaricus</i> , Agaric	<i>Amyris</i> , Balsamier
<i>Agathophyllum</i> , Raven-tsara	<i>Anacardium</i> , Anacarde
<i>Agave</i> , Agavé	<i>Anacyclus</i> , Anacycle
<i>Ageratum</i> , Agérate	<i>Anagallis</i> , Mouron
<i>Aggregata</i> , Aggrégées	<i>Anagyris</i> , Anagyris, Bois puant
<i>Agrimonia</i> , Aigremoine	<i>Anastatica</i> , Jérose
<i>Agrostemma</i> , Agrostemma	<i>Anavinga</i> , Anavinga
<i>Agrostis</i> , Agrostis	<i>Anchusa</i> , Buglose
<i>Agyneja</i> , Agynei	<i>Ancistrum</i> , Ancistrum
<i>Aira</i> , Canche	<i>Andrachne</i> , Andrachné
<i>Ajuga</i> , Bugle	<i>Andromeda</i> , Andromède
<i>Aizoon</i> , Aizoon, ou Lanquette	<i>Andropogon</i> , Barbon
<i>Albuca</i> , Albuca	<i>Androsace</i> , Androsace
<i>Alcea</i> , Alcée	<i>Androsæmum</i> , Voyez. <i>Hypéri-</i> <i>cum</i>
<i>Alchimilla</i> , Alchimille ou Pied- de-Lion	<i>Andryala</i> , Andryale
<i>Aldrovanda</i> , Aldrovande	<i>Anemone</i> , Anémone
<i>Aletris</i> , Alétris	<i>Anethum</i> , Aneth
<i>Algæ</i> , Algues	<i>Angelica</i> , Angélique
<i>Amaranthi</i> , Amarantes	<i>Anguillaria</i> , Gærtm. <i>Badula</i> . J. Anguillaire. V. <i>Badula</i>
<i>Amaranthus</i> , Amaranthe	<i>Anguria</i> , Angourie
<i>Amaryllis</i> , Amaryllis	<i>Aniba</i> , Aniba
<i>Amasonia</i> , Amasone, Ama- sonie	<i>Anomalæa</i> , T. Anomales
<i>Ambora</i> , Tamboul	<i>Anona</i> , Anone, Corossol
<i>Ambrosia</i> , Ambrosie	<i>Anonæa</i> , Anones
<i>Amellus</i> , Amelle	<i>Anthemis</i> , Camomille
<i>Amentaceæ</i> , Amentacées	<i>Anthericum</i> , Anthéric
<i>Amentaceæ (Arbores)</i> , Arbres, à Chaton, ou Amentacés	<i>Anthoceros</i> , Ancthocère

<i>Antholyza</i> , Antholyze	<i>Aroidæ</i> , J. Aroides
<i>Anthoxanthum</i> , Flouve	<i>Artedia</i> , Artédie
<i>Anthyllis</i> , Anthyllide	<i>Artemisia</i> , Armoise
<i>Antidesma</i> , Antidesma	<i>Artocarpus</i> , Jaquier
<i>Antirrhinum</i> , Mufflier	<i>Arum</i> , Arum, Gouet
<i>Apactis</i> , Apactis	<i>Arundo</i> , Roseau
<i>Apetalæ (Arbores)</i> , Arbres	<i>Asarum</i> , Asaret, Cabaret
Apétales	<i>Ascarina</i> , Ascarine
<i>Aphanes</i> , Aphanès, Percepier	<i>Asclepias</i> , Asclépiade
<i>Aphytëia</i> , Aphytée	<i>Ascyrum</i> , Ascyre
<i>Apium</i> , Persil	<i>Aspalathus</i> , Aspalat
<i>Apluda</i> , Aplude	<i>Asparagi</i> J. Asperges
<i>Apocynæ</i> , Apocinées	<i>Asparagus</i> , Asperge
<i>Apocinum</i> , Apocin, ou Apocyn	<i>Asperifoliæ</i> , Apresfeuilles
<i>Aponogeton</i> , Aponoget	<i>Asperugo</i> , Rapette
<i>Aquilaria</i> , Aquilaria, Garo	<i>Asperula</i> , Aspérule
<i>Aquilegia</i> , Ancolie	<i>Asphodeli</i> J. Asphodèles
<i>Aquilicia</i> , Aquilice	<i>Asphodelus</i> , Asphodèle
<i>Arabis</i> , Arabette	<i>Asplenium</i> , Doradille
<i>Arachis</i> , Arachide	<i>Assonia</i> , Assonia
<i>Aralia</i> , Aralie	<i>Aster</i> , Astère
<i>Araliæ</i> , Aralies	<i>Astragalus</i> , Astragale
<i>Araucaria</i> , Araucaria, Pin du Chili	<i>Astrantia</i> , Astrance
<i>Arbustiflæ</i> , Arbustives	<i>Astronium</i> , Astronium
<i>Arbutus</i> , Arbousier	<i>Athamantha</i> , Athamanthe
<i>Arctium</i> , Bardane	<i>Athanasia</i> , Athanasie
<i>Arctotis</i> , Arctotide	<i>Atractylis</i> , Atractylide
<i>Areca</i> , Arec ou Arèque	<i>Atragene</i> , Atragène
<i>Arenaria</i> , Sabline	<i>Atriplices</i> J. Arroches
<i>Argemone</i> , Argémone	<i>Atriplex</i> , Arroche
<i>Aristida</i> , Aristide	<i>Atropa</i> , Belladone
<i>Aristolochia</i> , Aristoloche	<i>Avena</i> , Avoine
<i>Aristolochiæ</i> , Aristoloches	<i>Azerrhou</i> , Carambolier
<i>Aristotelia</i> , Mâqui du Chili	<i>Aurantia</i> J. Orangers
<i>Armeniaca</i> , Abricotier	<i>Auricularia</i> , Auriculaire
<i>Arnica</i> , Arnica	<i>Axyris</i> , Axyris

<i>Ayenia</i> , Ayénia	<i>Bignonia</i> , Bignones
<i>Aylantus</i> , Langit	<i>Biscutella</i> , Lunetière
<i>Aytonia</i> , Aïton	<i>Bisserula</i> , Double-scie
<i>Azalea</i> , Azalée	<i>Bixa</i> , Rocou
<i>Azima</i> , Azima	<i>Blahdia</i> , Blahdia
	<i>Blasia</i> , Blasie
B	<i>Blechnum</i> , Bleigne
<i>Baccharis</i> , Bacchante	<i>Blittum</i> , Blète ou Blite
<i>Badula</i> , Bois de pintade	<i>Bobartia</i> , Bobarte
<i>Balanophora</i> , Balanophore	<i>Bocconia</i> , Boccône
<i>Ballota</i> , Ballote	<i>Boehmeria</i> , Boehmer
<i>Balsimina</i> , Balsamine	<i>Boerhaavia</i> , Boerhavie
<i>Baltimora</i> , Baltimore	<i>Boletus</i> , Bolet
<i>Banisteria</i> , Banisteria	<i>Bombar</i> , Fromager
<i>Barbula</i> (Hedw.)	<i>Borbonia</i> , Borbonia
<i>Barbylus</i> , Barbyl	<i>Borragineæ</i> , Borraginées
<i>Barleria</i> , Barrélière	<i>Borrago</i> , Bourrache
<i>Barnadesia</i> , Barnadez	<i>Brabeium</i> , Brabei
<i>Bartramia</i> (Hedw.)	<i>Brassica</i> , Chou
<i>Basella</i> , Baselle	<i>Briza</i> , Amourette
<i>Basilæa</i> , Basilée	<i>Bromelia</i> , Ananas
<i>Bassia</i> , Illipé	<i>Bromeliæ</i> , Ananas
<i>Bassovia</i> , Bassove	<i>Bromus</i> , Brôme
<i>Batis</i> , Bâtis	<i>Browallia</i> , Broualle
<i>Bauhinia</i> , Bauhinia ou Bauhin	<i>Brownæa</i> , Brounéa
<i>Begonia</i> , Bégône	<i>Brunella</i> , Brunelle
<i>Bellis</i> , Pâquerette	<i>Brunia</i> , Brunia
<i>Bellium</i> , Bellium	<i>Bryonia</i> , Bryône
<i>Berberides</i> , Vinettiers	<i>Bryum</i> , Bry
<i>Berberis</i> , Vinettier	<i>Bubon</i> , Bubone
<i>Beta</i> , Bette	<i>Bucida</i> , Grignon
<i>Betonica</i> , Bétoine	<i>Budleïa</i> , Budlèje
<i>Betula</i> , Bouleau	<i>Bufonia</i> , Buffone
<i>Bicornes</i> , Bicornées	<i>Bugula</i> , Bugle
<i>Bidens</i> , Bident	<i>Bulbocodium</i> , Bulbocode
<i>Bignonia</i> , Bignone	<i>Bunias</i> . Voyez Caméline

<i>Bunium</i> , Terre-Noix	<i>Canarium</i> , Canari
<i>Buphtalmum</i> , Buphtalme	<i>Candelares</i> , L.....
<i>Buplevrum</i> , Buplèvre	<i>Canna</i> , Balisier
<i>Burmannia</i> , Burmanne	<i>Cannabis</i> , Chanvre
<i>Butomus</i> , Butôme	<i>Cannæ</i> , Balisiers
<i>Butonica</i> , Butonic	<i>Cantharellus</i> , Chanterelle
<i>Buxbaumia</i> , Buxbaume	<i>Cantua</i> , Cantu
<i>Buxus</i> , Buis	<i>Capitatae</i> , Capitées
<i>Byssus</i> , Byssus, Byssa	<i>Capparides</i> , Capriers
	<i>Capparis</i> , Câprier
C	<i>Capraria</i> , Capraire
<i>Cacalia</i> , Cacalie	<i>Cuprifolia</i> , Chevrefeuille
<i>Cachrys</i> , Armarinthe	<i>Caprifolium</i> , Chèvrefeuille
<i>Cacti</i> , Cactiers	<i>Capsicum</i> , Piment
<i>Cactus</i> , Cactier	<i>Capura</i> , Capura
<i>Cæsalpina</i> , Bresillet	<i>Caragana</i> , Caragan
<i>Calamus</i> . Voyez <i>Acorus</i>	<i>Caraipa</i> , Caraïpa
<i>Calceolaria</i> , Calcéolaire	<i>Cardamine</i> , Cresson
<i>Calcitrapa</i> , Chausse-Trape	<i>Cardiospermum</i> , Corinde
<i>Calea</i> , Caléa	<i>Carduus</i> , Chardon
<i>Calendula</i> , Souci	<i>Carex</i> , Carex, Caret, Laiche
<i>Calinea</i> , Calinéa	<i>Carlina</i> , Carline
<i>Calla</i> , Calle, Chou-calle	<i>Carpesium</i> , Carpèse
<i>Callitriche</i> , Callitrique	<i>Carpinus</i> , Charme
<i>Calodendrum</i> , Calodendrum	<i>Carthamus</i> , Carthame
<i>Calophyllum</i> , Calaba	<i>Carum</i> , Carvi
<i>Caltha</i> , Populage	<i>Caryocar</i> , Caryocar
<i>Calycanthemæ</i> , Calycanthèmes	<i>Caryophylleæ</i> , Caryophyllées
<i>Calycanthus</i> , Calycanth	<i>Caryophyllus</i> , Girofflier
<i>Cambogia</i> , Cambogier, Guttier	<i>Cassine</i> , Cassine
<i>Cameruria</i> , Camérier	<i>Cassuvium</i> , Acajou
<i>Campanaceæ</i> , Campanacées	<i>Cassytha</i> , Cassythe
<i>Campaniformes</i> , Campani- formes	<i>Casuarina</i> , Filao
<i>Campanula</i> , Campanule	<i>Catalpa</i> . Voyez Bignonæ
<i>Campanulæ</i> , Campanules	<i>Catanance</i> , Cupidone
	<i>Catka</i> , Catha

<i>Catimbium</i> , Catimban	<i>Chrysanthemum</i> , Chrysanthème
<i>Catonia</i> , Catonia	<i>Crysocoma</i> , Chrysocôme
<i>Caucalis</i> , Caucalide	<i>Chrysophyllum</i> , Caïmitier
<i>Ceanothus</i> , Céanothus	<i>Chrysozplenium</i> , Dorine
<i>Cecropia</i> , Coulekin	<i>Ciathea</i> , Smith
<i>Cedrela</i> , Cédrel	<i>Cicer</i> , Ciche, Pois-ciche ou Pois-chiche
<i>Celastrus</i> , Célastre	<i>Cichoraceæ</i> , Chicoracées
<i>Celosia</i> , Passe-velours	<i>Chichorium</i> , Chicorée
<i>Celtis</i> , Micocoulier	<i>Cicuta</i> , Ciguë
<i>Cenchrus</i> , Râcle	<i>Cicutaria</i> , Cicutaire
<i>Centaurea</i> , Centaurée	<i>Cimicifuga</i> , Cimicaire
<i>Centunculus</i> , Centenille	<i>Chinchona</i> , Quinquina
<i>Ceodes</i> , Céodès	<i>Cinara</i> , Artichaut
<i>Cephalanthus</i> , Céphalant	<i>Cinarocephalæ</i> , Cinarocéphales
<i>Cerastium</i> , Céraiste	<i>Cineraria</i> , Cinéraire
<i>Cerasus</i> , Cerisier	<i>Cinna</i> , Cinna
<i>Ceratonia</i> , Caroubier	<i>Circæa</i> , Circée
<i>Ceratophyllum</i> , Cornifle	<i>Cissampelos</i> , Cissampelos
<i>Cerbera</i> , Ahouï	<i>Cissus</i> , Cissus, Achèt
<i>Cercis</i> , Gainier	<i>Cisti</i> , Cistes
<i>Cercodea</i> , Cercodéa	<i>Cistus</i> , Ciste
<i>Cerinthé</i> , Melinet	<i>Citras</i> , Citronier
<i>Cestrum</i> , Cestreau	<i>Clathrus</i> , Clathre
<i>Chærophyllum</i> , Cerfeuil	<i>Clavaria</i> , Clavaire
<i>Chamærops</i> , Palmier-éven-tail	<i>Clausena</i> , Clausèna
<i>Chara</i> , Charagne	<i>Clématis</i> , Clématite
<i>Cheiranthus</i> , Giroflée	<i>Cleome</i> , Mozambé
<i>Chelidonium</i> , Chélidoine	<i>Clethra</i> , Clethra
<i>Chelone</i> , Galane	<i>Cleyera</i> , Cleyèra
<i>Chenopodium</i> , Chénopode, An- sérine	<i>Clibadium</i> , Clibade
<i>Chionanthus</i> —	<i>Cliffortia</i> , Cliffort
<i>Chironia</i> , Chérone	<i>Clinopodium</i> , Clinopode
<i>Chloranthus</i> , Chloranthus ou Ni- grine	<i>Clitoria</i> , Clitorie
<i>Chondrilla</i> , Chondrille	<i>Clusia</i> , Clusia
	<i>Clutia</i> , Clutia

<i>Clypeola</i> , Clypéole	<i>Coreopsis</i> , Coréope
<i>Cneorum</i> , Camelée	<i>Coriandrum</i> , Coriandre
<i>Cnestis</i> , Cnestis	<i>Coriaria</i> , Corroyer
<i>Cnicus</i> , Cnique	<i>Cornus</i> , Cornouiller
<i>Coadunata</i> , Connées	<i>Coronaria</i> , Coronaires
<i>Colloloba</i> , Raisinier	<i>Coronilla</i> , Coronille
<i>Cochlearia</i> , Cochléaria, Vansone	<i>Corrigiola</i> , Corrigiole
<i>Cocos</i> , Coco	<i>Corydalis</i> , Corydalis
<i>Codon</i> , Codon	<i>Corylus</i> , Coudrier
<i>Cænopteris</i> , Berg.....	<i>Corymbifera</i> , Corymbifères
<i>Coffea</i> , Cafféyer	<i>Corymbium</i> , Corymbiole
<i>Coix</i> , Larme de Job	<i>Corypha</i> , Corypha
<i>Colchicum</i> , Colchique	<i>Coryspermum</i> , Corysperme
<i>Coldenia</i> , Coldene	<i>Costus</i> , Costus
<i>Columnifera</i> , Columnifères	<i>Cotula</i> , Cotule
<i>Colutea</i> , Baguenaudier	<i>Cotyledon</i> , Cotylédone, Coty- lette
<i>Comarum</i> , Comaret	<i>Coutarea</i> , Coutaréa
<i>Combretum</i> , Combretum	<i>Crambe</i> , Crambé
<i>Cometes</i> , Comete	<i>Crassula</i> , Crassule
<i>Commelina</i> , Commeline	<i>Cratægus</i> , Alisier
<i>Commersonia</i> , Commerson	<i>Crepis</i> , Crévide
<i>Compositæ (flores)</i> , Composées	<i>Crescentia</i> , Calebassier
<i>Conferva</i> , Conferve	<i>Crinodendrum</i> , Crinodendron, Patagna
<i>Conifera</i> , Conifères	<i>Crinum</i> , Crinole
<i>Conium</i> , Conium	<i>Crithmum</i> , Criste, Bacille
<i>Comarus</i> , Connas	<i>Crocus</i> , Crocuse
<i>Conocarpus</i> , Manglier	<i>Crossostylis</i> , Crostyle
<i>Contorta</i> , Contournées	<i>Crotalaria</i> , Crotalaire
<i>Convallaria</i> , Muguet	<i>Croton</i> , Croton
<i>Convolvuli</i> , Liserons	<i>Crucianella</i> , Crucianelle
<i>Convolvulus</i> , Liseron	<i>Crucifera</i> , Crucifères
<i>Conyza</i> , Conyse	<i>Crupina</i> , Adans. <i>Serratula</i> (il y aussi une <i>Centaurea crupina</i>).
<i>Copaïfera</i> , Copaïer	Voyez <i>Serratula</i>
<i>Corchorus</i> , Corète	
<i>Cordia</i> , Sébestier	

<i>Cucubalus</i> , Cucubale	<i>Damasonium</i> , J. <i>Alisma Damasonium</i> , L. V. <i>Alisma</i>
<i>Cucumis</i> , Concombre	<i>Daphne</i> , Lauréole. (Lauréol.)
<i>Cucurbita</i> , Courge	<i>Darea</i> , Darée
<i>Cucurbitaceæ</i> , Cucurbitacées	<i>Datisca</i> , Cannabine
<i>Culmineæ</i> , Culminées	<i>Datura</i> , Datura
<i>Cuminum</i> , Cumin	<i>Daucus</i> , Carotte
<i>Cupania</i> , Cupani	<i>Davallia</i> , Smith...
<i>Cupressus</i> , Cyprès	<i>Delphinium</i> , Dauphinelle, Pied d'Alouette
<i>Curcuma</i> , Curcuma	<i>Dentaria</i> , Dentaire
<i>Cuscuta</i> , Cascute	<i>Denudataæ</i> , Nues ou Dépouillées
<i>Cussonia</i> , Cussonia	<i>Deutzia</i> , Deutz
<i>Cyanilla</i> , Cyanelle	<i>Dialium</i> , Diali
<i>Cyanus</i> , Ambrette	<i>Dianella</i> , Dianelle
<i>Cyathus</i> ...Réuni aux Pezize par Bulliard	<i>Dianthera</i> , Dianthèra
<i>Cycas</i> , Cycas	<i>Dianthus</i> , Œillet
<i>Cyclamen</i> , Cyclame	<i>Diapensia</i> , Diapenze
<i>Cydonia</i> , Coignassier	<i>Dichondra</i> , Dichondre
<i>Cymosæ</i> , Cimoides	<i>Dicksonia</i> , L'Hérit, Smith...
<i>Cynanchum</i> , Cynanch	<i>Dicranum</i> , Hew, Bridel...
<i>Cynoglossum</i> , Cynoglosse	<i>Dicotyledones</i> , Dicotyledons
<i>Cynometra</i> , Cynometra	<i>Dictamnus</i> , Dictamne
<i>Cynomorium</i> , Cynomoire	<i>Didelta</i> , Didelta (Dideltoïde)
<i>Cynosurus</i> , Crételle	Didymodon, Hedw. Brid....
<i>Cyperoïdeæ</i> , Cyperoïdes ou Souchets	<i>Diervilla</i> , Diervilla
<i>Cyperus</i> , Souchet, Souchette	<i>Digitalis</i> , Digitale
<i>Cypripedium</i> , Sabot, Sabotine	<i>Dillenia</i> , Dillen
<i>Cytinus</i> , Cytinel	<i>Dionæa</i> , Dionée
<i>Cytisus</i> , Cytise	<i>Dioscorea</i> , Dioscorée, Ignose
D	<i>Diosma</i> , Diosma
<i>Dactylis</i> , Dactyle	<i>Diospyros</i> , Plaqueminier
<i>Dalbergia</i> , Dalberg	<i>Dipsaceæ</i> . Dipsacées
<i>Dalea</i> , Dalée	<i>Dipsacus</i> , Cardère

<i>Dirca</i> , Dirca	<i>Embelia</i> , Embelia
<i>Dobera</i> , Dobèra	<i>Embothrium</i> , Embothrium
<i>Dodartia</i> , Dodarte	<i>Empetrum</i> , Camarine (Empe- trum).
<i>Dodecatheon</i> , Dodécathéone	<i>Encelia</i> , Encélie
Giroselle	<i>Ensata</i> , Gladiées
<i>Dodonæa</i> , Dodonéa	<i>Ephedra</i> , Ephédra
<i>Dolichos</i> , Dolique	<i>Epidendrum</i> , Epidendrone
<i>Doliocarpus</i> , Doliocarpus	<i>Epilobium</i> , Epilobe
<i>Dombeya</i> , Dombey	<i>Epimedium</i> , Epimède
<i>Doræna</i> , Dorèna	<i>Equisetum</i> , Prêle
<i>Doronicum</i> , Doronic (Doronique).	<i>Erharta</i> , Erharte
<i>Dorstenia</i> , Dorstène	<i>Erica</i> , Bruyère (Erica).
<i>Draba</i> , Drâve	<i>Ericæ</i> , Bruyères
<i>Dracæna</i> , Sang-Dragon	<i>Erigeron</i> , Vergerolle
<i>Dracocephalum</i> , Dracocephale,	<i>Eriocaulon</i> , Joncinelle
Moldavique	<i>Eriocephalus</i> , Eriocéphale, (Eri- océphal)
<i>Dracontium</i> , Draconte	<i>Eriophorum</i> , Linaigrette
<i>Drosera</i> , Rossolis (Drosère).	<i>Erum</i> , Ers, (Érse)
<i>Drupacæa</i> , Drupacées	<i>Eryngium</i> , Panicaut, (Panicau- de)
<i>Dryandra</i> , Dryandra	<i>Erysimum</i> , Vêlar, (Vêlare)
<i>Dryas</i> , Dryade	<i>Erysimum</i> —
<i>Dumosæ</i> , Buissonées	<i>Erythrina</i> , Erythrina
E	<i>Erythronium</i> , Erythronium, Ery- thron
<i>Echinops</i> , Echinops	<i>Erythroxyllum</i> , Erythroxyllon
<i>Echites</i> , Echites	<i>Euclea</i> , Eucléa
<i>Echium</i> , Vipérine	<i>Eugenia</i> , Eugénia, Sambosier
<i>Ehretia</i> , Cabrillet	<i>Eupatorium</i> , Eupatoire
<i>Elæagni</i> , Chalefs	<i>Euphorbia</i> , Euphorbe
<i>Elæagnus</i> , Chalef	<i>Euphorbia</i> , Euphorbes
<i>Elæocarpus</i> , Eléocarpus	<i>Euphrasia</i> , Euphraise
<i>Elatarium</i> , J. <i>Momordica</i> <i>Elate-</i> <i>rium</i> , L. Voy. <i>Momordica</i>	<i>Eurya</i> , Eurya
<i>Elatine</i> , Elatine	<i>Evea</i> , Evéa
<i>Elymus</i> , Elyme	

Evolvulus, Liseret*Evonymus*, Fusian

F

Faba J. *Vicia*, *Faba* L. Voy. *Vicia**Fagara*, Fagara*Fagonia*, Fagone*Fagus*, Hêtre*Ferraria*, Ferrarée*Ferula*, Férule*Festuca*, Féstouque*Fevillea*, Févillée, Nandirobe*Ficaria*, *Ranunculus ficaria*,Voy. *Ranunculus**Ficoidea*, J. Ficoïdes*Ficus*, Figuier*Filago*, Cotonière*Filices*, Fougères*Flagellaria*, Flagellaire

..... Fleurs à étamines

(plantes à)

..... Fleurs ni fruits.

(plantes sans)

Flosculosi (*flores*), Flosculeuses*Fontinalis*, Fontinale*Forskalea*, Forskale*Forstera*, Forstère*Fothergilla*, Fothergilla*Fragaria*, Fraisier*Frankenia*, Franckène*Fraxinus*, Fresne*Fritillaria*, Fritillaire

..... Fruits sans fleurs.

(Plantes à)

Fuchsia, Fuchsia*Fuci*, *Fucus* (les), Sous-ord*Fucus*, Fucus*Fugosia*, Fugosie*Fumaria*, Fumeterre.*Fungi*, Champignons

G

Galanthus, Galantine*Galax*, Galax*Galega*, Galéga Lavanèse*Galenia*, Galiène*Galeopsis*, Galéope*Galipwa*, Galipier*Gallium*, Galliet*Garcinia*, Mangoustan*Garidella*, Garidelle*Geniostoma*, Geniostome*Genipa*, Génipayer*Genista*, Genest*Gentiana*, Gentiane*Gentianæ*, Gentianes*Geoffræa*, Geoffréa*Gerania*, Geraines*Geranium*, Géranium, Geraine*Geropogon*, Géropogone*Gethyllis*, Gethyllide*Geum*, Benoite*Gevuina*, Geouin*Glabraria*, Glabraria*Gladiolus*, Glayeul*Glaucium*, Glaucienne*Glaux*, Glauce*Glecoma*, Glécome*Gleditsia*, Fêvier*Glinus*, Glinole*Globba*, Globba*Globularia*, Globulaire

<i>Glochidion</i> , Glochidion	<i>Helicteres</i> , Helictères
<i>Gluta</i> , Gluta	<i>Heliocarpus</i> , Héliocarpe
<i>Glycine</i> , Glycine	<i>Heliotropium</i> , Héliotrope
<i>Glycyrrhiza</i> , Réglisse	<i>Helleborus</i> , Hellébore
<i>Gnaphalium</i> , Gnaphale	<i>Helonias</i> , Hélonias
<i>Gnidia</i> , Guidienne	<i>Helvella</i> , Helvèle
<i>Gomphrena</i> , Amaranthine	<i>Hemerocallis</i> , Hémérocalles
<i>Gonocarpus</i> , Gonocarpe	<i>Hemionitis</i> , Hémionite
<i>Gossypium</i> , Cotonier	<i>Hepaticæ</i> , Hépatiques
<i>Graminæ</i> , Graminées	<i>Hericius</i> , Urchin
<i>Gratiola</i> , Gratiolle	<i>Hermannia</i> , Hermannia
<i>Grewia</i> , Grewia, Greuvier	<i>Hernias</i> , Hernias
<i>Gronovia</i> , Gronove	<i>Hernandia</i> , Hernandia
<i>Guaiacum</i> , Plaqueminiens	<i>Herniaria</i> , Herniole
<i>Guaiacum</i> , Gayac	<i>Hesperidæ</i> , Hespéridées
<i>Guettarda</i> , Guettard	<i>Hesperis</i> , Julienne
<i>Guilandina</i> , Bonduc	<i>Hibiscus</i> , Ketmie
<i>Gundelia</i> , Gondèle	<i>Hieracium</i> , Épervière
<i>Gunnera</i> , Gunnère	<i>Hippia</i> , Hippia
<i>Guttiferæ</i> , Guttiers	<i>Hippocratea</i> , Béjuco
<i>Gypsophyla</i> , Gypsophyle	<i>Hippocrepis</i> , Hippocrépide
	<i>Hippomane</i> , Mancenillier
H	<i>Hippomanica</i> , Hippomanique
<i>Hæmanthus</i> , Hæmanthe	<i>Hippophaë</i> , Argousier
<i>Hæmatoxylum</i> , Campêche	<i>Hippuris</i> , Pesse
<i>Hamamelis</i> , Hamamelis	<i>Hirtella</i> , Hirtelle
<i>Hamelia</i> , Hamelia	<i>Holcus</i> , Houque
<i>Hedera</i> , Lierre	<i>Holeraceæ</i> Oléracées ou Potagères
<i>Hedycaria</i> , Hedycaria	<i>Holosteum</i> , Holostée
<i>Hedyotis</i> , Hedyotis	<i>Homalium</i> , Homali, Acomat
<i>Hedypnois</i> , Hedyppoïde	<i>Hordeum</i> , Orge
<i>Hedysarum</i> , Sain-Foin	<i>Hottonia</i> , Hottone, Plumeau
<i>Helenium</i> , Hélène	<i>Houttuynia</i> , Houttuynie
<i>Helianthemum</i> , Helianthème	<i>Humulus</i> , Honblou
<i>Helianthus</i> , Hélianthe	<i>Hura</i> , Sâblier
<i>Heliconia</i> , Bihai	

<i>Hyacinthus</i> , Jacinthe	<i>Irides</i> . J. Iris
<i>Hydnum</i> , Hydne	<i>Iris</i> , Iris
<i>Hydrastis</i> , Hydrastis	<i>Isatis</i> , Pastel
<i>Hydrocharides</i> . Morrènes	<i>Isnardia</i> , Isnarde
<i>Hydrocharis</i> , Morrène	<i>Isoëtes</i> , Isote
<i>Hydrocotyle</i> , Hydrocotyle	<i>Itea</i> , Itéa
<i>Hydrophylax</i> , Hydrophylax	<i>Iva</i> , Iva
<i>Hymenea</i> , Courbaril	<i>Ixia</i> , Ixie
<i>Hyobanche</i> , Hyobanche	<i>Ixora</i> , Ixora
<i>Hyosciamus</i> , Jusquiame	
<i>Hyoseris</i> , Hyoséride	J
<i>Hypecoum</i> , Hypécoon	<i>Jacaranda</i> , Jacaranda
<i>Hyperica</i> . Millepertuis	<i>Jacea</i> , Jacée
<i>Hypericum</i> , Millepertuis	<i>Jusione</i> , Jusione
<i>Hypnum</i> , Hypne	<i>Jusmineæ</i> . Jasminées
<i>Hypochæris</i> , Hypochæride	<i>Jasminum</i> , Jasmin
<i>Hypoxis</i> , Hypoxis	<i>Jatropha</i> , Jatropha, Médecinier
<i>Hypoxylum</i> , Hypoxylon	<i>Juglans</i> , Noyer
<i>Hyssopus</i> , Hyssope	<i>Juncago</i> . Voy. Triglochine
	<i>Junci</i> . Jones
I	<i>Juncus</i> , Jonc
<i>Iberis</i> , Ibéride	<i>Jungermannia</i> , Jongermanne
<i>Icica</i> , Iciquier	<i>Juniperus</i> , Genévrier
<i>Ilex</i> , Houx	<i>Jussiaea</i> , Jussienne
<i>Illecebrum</i> , Illécébrum	<i>Justitia</i> , Carmantine
<i>Illicium</i> , Badiane	
<i>Imbricaria</i> , Imbricaria, Bois de Natte	K
<i>Imperatoria</i> , Impératoire	<i>Kæmpferia</i> , Zédoaire
<i>Imperialis</i> , Impériale	<i>Kalmia</i> , Kalmia
<i>Indigofera</i> , Indigotier	<i>Kiggellaria</i> , Kiggellaria
<i>Infundibuliformes</i> . Infundibu- liformes	<i>Kleinhovia</i> , Kleinilovia
<i>Inula</i> , Inule, Année	<i>Kænigia</i> , Kænige
<i>Inundata</i> , Inondées	<i>Koëltreutera</i> , Hedw....
<i>Ipomæa</i> , Ipoméé, Quamoclit	<i>Krameria</i> , Kramer
	<i>Kuhnia</i> , Kuhnie

L	
<i>Labiata</i> , Labiées	<i>Ligusticum</i> , Livêche
<i>Lactuca</i> , Laitue	<i>Ligustricum</i> , Troène
<i>Lagerstromia</i> , Lagerstromia	<i>Lilac</i> , <i>Syringa</i> . Voyez <i>Syringa</i>
<i>Lagetta</i> , Lagetto	<i>Lilia</i> , Lis
<i>Lagoëcia</i> , Lagœcie	<i>Liliaceæ</i> , Liliacées
<i>Lamium</i> , Lamier	<i>Lilium</i> , Lis
<i>Lampsana</i> , Lampsane	<i>Limcum</i> , Liméole
<i>Lantana</i> , Lantana, Camara	<i>Limonia</i> , Limonellier
<i>Laserpitium</i> , Laser	<i>Limosella</i> , Limoselle
<i>Latania</i> , Latanier	<i>Linaria</i> , Linaire
<i>Lathræa</i> , Clandestine	<i>Linnæa</i> , Linnée
<i>Lathyrus</i> , Gesse	<i>Linum</i> , Lin
<i>Lavandula</i> , Lavande	<i>Liparia</i> , Lipari
<i>Lavatera</i> , Lavatère	<i>Liquidambar</i> , Liquidambar
<i>Lauri</i> , Lauriers	<i>Liriodendrum</i> , Tuñpier
<i>Laurus</i> , Laurier	<i>Lithospermum</i> , Grémil
<i>Lausonia</i> , Lausonia, Henné	<i>Littorella</i> , Litorelle
<i>Lechea</i> , Léchéa	<i>Lobelia</i> , Lobélie
<i>Lecythis</i> , Lecythis, Quatèle	<i>Lolium</i> , Ivroie
<i>Ledum</i> , Lède	<i>Lomentaceæ</i> , Lomentacées
<i>Leca</i> , Léca	<i>Lonchitis</i> , Lonchite
<i>Leersia</i> , Hedw....	<i>Lonicera</i> , Chèvrefeuille
<i>Leguminosæ</i> , Légumineuses	<i>Lontarus</i> , Lontar
<i>Lemna</i> . Voyez <i>Marsilea</i>	<i>Lophanthus</i> , Lophanthe
<i>Lenticula</i> , Lenticule, Canillée	<i>Loranthus</i> , Loranthe
<i>Leontice</i> , Léontice	<i>Lotus</i> , Lotier
<i>Leontodon</i> , Leontodon Liondent	<i>Lunaria</i> , Luneaire
<i>Leonurus</i> , Agripaume	<i>Lupinus</i> , Lupin
<i>Lepidium</i> , Pâsse-rage	<i>Lurida</i> , Livides
<i>Lepra</i> , Lèpre	<i>Lychnis</i> , Lychnide
<i>Lerchea</i> , Lerchéa	<i>Lycium</i> , Lyciet
<i>Leskia</i> , Hedw....	<i>Lycoperdon</i> , Lycoperdon, Vesse-
<i>Leucodium</i> , Nivéole	Loup
<i>Leysera</i> , Leysera ou Leyser	<i>Lycopodium</i> , Lycopode
<i>Lichen</i> , Lichen	<i>Licopsis</i> , Lycopsite
	<i>Lycopus</i> , Lycope

<i>Lygeum</i> , Alvarde	<i>Melastomæ</i> , Mélastômes
<i>Lysimachia</i> , Lysimaque	<i>Melia</i> , Azédarach
<i>Lysimachia</i> , Lysimachies	<i>Melia</i> , Azédarachs
<i>Lythrum</i> , Salicaire	<i>Melianthus</i> , Mélianthé
	<i>Melica</i> , Mélique
M	<i>Melicope</i> , Mélicope
<i>Maba</i> , Maba	<i>Melicytus</i> , Mélicyte
<i>Marua</i> , Mœrua	<i>Melilotus</i> , <i>Trifolium</i> , <i>Melilotus</i> ,
<i>Magnolia</i> , Magnolier	Mélilot
<i>Magnolia</i> , Magnoliers	<i>Melissa</i> , Melisse
<i>Mahurea</i> , Mahuré	<i>Melittis</i> , Mélitte, Melissot
<i>Malachra</i> , Malacre	<i>Melochia</i> , Mélochias
<i>Malope</i> , Malope	<i>Melothria</i> , Mélothrie
<i>Malpighia</i> , Malpighi	<i>Menisperma</i> , Menispermes
<i>Malpighia</i> , Malpighies	<i>Menispermum</i> , Ménisperme
<i>Malva</i> , Mauve	<i>Mentha</i> , Menthe
<i>Malvaceæ</i> , Malvacées	<i>Mentzélia</i> , Mentzélie
<i>Malus</i> , Pommier	<i>Menyanthes</i> , Ményanthe
<i>Mandragora</i> , <i>Atropa</i> <i>Mandrago-</i>	<i>Mercurialis</i> , Mercuriale
<i>ra</i> , Mandragore. Voy. <i>Atropa</i>	<i>Merulius</i> , Mérule (Champ.)
<i>Maranta</i> , Galanga	<i>Meryta</i> , Méryta
<i>Marattia</i> , Smith, Maratte	<i>Mespilus</i> , Néflier
<i>Marchantia</i> , Hépatique	<i>Mesembryanthemum</i> , Ficoïde
<i>Margaritaria</i> , Margaritaire	<i>Messerschmidia</i> , Arguze
<i>Marrubium</i> , Marrube	<i>Methonica</i> , Méthonique
<i>Marsilea</i> , Marsile	<i>Micropus</i> , Micrope
<i>Martynia</i> , Cornaret, Bicorne	<i>Milium</i> , Mil
<i>Massonia</i> , Massione	<i>Milleria</i> , Millérie
<i>Matricaria</i> , Matricaire	<i>Mimosa</i> , Mimosa, Acacie
<i>Meborea</i> , Méboré	<i>Mimusops</i> , Mimusops
<i>Medeola</i> , Médéole	<i>Mirabilis</i> , Nictage
<i>Medicago</i> , Luserne	<i>Mitchella</i> , Mitchelle
<i>Meesia</i> , Hedw.	<i>Mitella</i> , Mitelle
<i>Melaleuca</i> , Mélaleuca	<i>Mniarum</i> , Mniarum
<i>Melampyrum</i> , Mélampyre	<i>Mnium</i> , Mnie
<i>Melastoma</i> , Mélastôme	<i>Molucella</i> , Molucelle

<i>Mollugo</i> , Mollugine	N
<i>Momordica</i> , Momordique	<i>Naiides</i> , Naïades
<i>Monopetalæ</i> (<i>Arbores</i>), Arbres	<i>Naias</i> , Naïade
Monopétales	<i>Nandina</i> , Nandina
<i>Monniera</i> , Monnière	<i>Napæa</i> , Napée
<i>Monocotyledones</i> , Monocotyle- dons	<i>Narcissi</i> , Narcisses
<i>Monotropa</i> , Monotrope	<i>Narcissus</i> , Narcisse
<i>Monsonia</i> , Monsone	<i>Nardus</i> , Nard
<i>Montia</i> , Montie	<i>Nastus</i> , Nastus
<i>Morinda</i> , Royoc	<i>Neckera</i> , Hedw.
<i>Moringa</i> , Moringa, Ben.	<i>Nelumbium</i> , <i>Nymphæa</i> , <i>Nelumbo</i> , Nélumbo
<i>Morus</i> , Murier	<i>Nepenthes</i> , Népente
<i>Moscharia</i> , Moscaire	<i>Nepeta</i> , Cataire
<i>Mourera</i> , Mourère	<i>Nephelium</i> , Néphélie
<i>Moutabea</i> , Moutabéa	<i>Nerium</i> , Nerion, Laur-Rose
<i>Mucor</i> , Mucor, Moisissure	<i>Nevrada</i> , Névrade
<i>Multisiliquosæ</i> , Multisiliqueuses	<i>Nicotiana</i> , Nicotiane, Tabac
<i>Munchausia</i> , Munchausia	<i>Nidularia</i> , Nidulaire, Bull
<i>Muricata</i> , Muriquées	<i>Nigella</i> , Nigelle
<i>Musa</i> , Bananier	<i>Nipa</i> , Nipa
<i>Musæ</i> , Bananiers	<i>Nitraria</i> , Nitraire
<i>Musci</i> , Mousses	<i>Nolana</i> , Nolane
<i>Mutisia</i> , Mutis	<i>Nucamentacæ</i> , Nucamentacées
<i>Myagrum</i> , Caméline	<i>Nyctagines</i> , Nyctages
<i>Myosotis</i> , Scorpionne	<i>Nyctago</i> , Nyctage
<i>Myosurus</i> , Myosure	<i>Nyctanthes</i> , Nyctanthe
<i>Myrica</i> , Gâlé	<i>Nymphæa</i> , Nénuphar
<i>Myriophyllum</i> , Myriofle	
<i>Myriothecca</i> , Myriothèque	O
<i>Myristica</i> , Muscadier	<i>Obolaria</i> , Obolaire
<i>Myrosma</i> , Myrosme	<i>Ochrosia</i> , Ochrosia, Bois jaune
<i>Myroxylum</i> , Myroxylon	<i>Ocimum</i> , Basilic
<i>Myrsine</i> , Myrsiné	<i>Octoblepharum</i> , Hedw....
<i>Myrti</i> , Myrthes	<i>Octospora</i> , Hedw....
<i>Myrtus</i> , Myrthe	<i>Oedera</i> , Oëder

<i>Ænanthe</i> , Ænanthe	<i>Paliurus</i> , Paliure
<i>Ænothera</i> , Onagre	<i>Palmæ</i> , Palmiers
<i>Olax</i> , Olax	<i>Panax</i> , Gin-seng
<i>Olea</i> , Olivier	<i>Pancreatium</i> , Pancraïis
<i>Olyra</i> , Olyre	<i>Pandanus</i> , Baquois
<i>Omphalea</i> , Omphaléa	<i>Panicum</i> , Panic
<i>Onagrea</i> , Onagres	<i>Papaver</i> , Pavot
<i>Onoclea</i> , Onoclée	<i>Papaveracea</i> , Papaveracée
<i>Ononis</i> , Bugrane	<i>Papaya</i> , Papayer
<i>Onopordum</i> , Onoporde	<i>Papilionacea</i> , Papilionacées
<i>Ophioglossum</i> , Ophioglosse	<i>Papilionacea</i> (<i>Arbores</i>) Arbres
<i>Ophryse</i> , Ophryse	Papilionacés
<i>Oppositi-folia</i> , Composées a	<i>Pariana</i> , Pariane
Feuilles opposées	<i>Parietaria</i> , Pariétaire
<i>Orchidea</i> , Orchidées	<i>Paris</i> , Parisette
<i>Orchis</i> , Orchis	<i>Parnassia</i> , Parnassie
<i>Origanum</i> , Origan	<i>Parthenium</i> , Parthène
<i>Orixa</i> , Orixa	<i>Paspalum</i> , Paspale
<i>Ornithogalum</i> , Ornithogale	<i>Passerina</i> , Passerine
<i>Ornithopus</i> , Ornithope, Pied-	<i>Passiflora</i> , Grenadille
d'Oiseau	<i>Pastinaca</i> , Panais
<i>Orobanche</i> , Orobanche	<i>Paullinia</i> , Paullinia
<i>Orobus</i> , Orobe	<i>Pavonia</i> , Pavonia
<i>Orontium</i> , Oronce	<i>Payrola</i> , Payrola
<i>Orthotricum</i> , Hedw....	<i>Pediculares</i> , Pédiculaires
<i>Oryza</i> , Riz	<i>Pedicularis</i> , Pédiculaire
<i>Osbeckia</i> , Osbeckie	<i>Peganum</i> , Harmale
<i>Osmunda</i> , Osmonde	<i>Pekca</i> , Pékéa
<i>Osyris</i> , Rouvet	<i>Peltigera</i>
<i>Ouratea</i> , Ouratéa	<i>Penæa</i> , Pénéa
<i>Ourisia</i> , Ourisie	<i>Pennantia</i> , Pennantia
<i>Oxalis</i> , Oxalide	<i>Pentapctes</i> , Pentapètes
	<i>Penthorum</i> , Penthôte
P	<i>Peplis</i> , Péplide
<i>Pachira</i> , Pachira	<i>Perebea</i> , Pérébéa
<i>Pæonia</i> , Pivoine	<i>Perforata</i> , Perforées

<i>Perilla</i> , Pérille	<i>Pisonia</i> , Pisonia
<i>Periploca</i> , Périploca	<i>Pistacia</i> , Pistachier
<i>Personata</i> , Personées	<i>Pistia</i> , Pistie
<i>Petiveria</i> , Pétivérie	<i>Pisum</i> , Pois
<i>Peucedanum</i> , Peucedanum	<i>Plantagines</i> , Plantains
<i>Peziza</i> , Pezize	<i>Plantago</i> , Plantain
<i>Phaca</i> , Phâce	<i>Platanus</i> , Platâne
<i>Phalaris</i> , Alpiste	<i>Plegorhiza</i> , Guaïcura
<i>Phalangium</i> , Phalangère	<i>Plumbagines</i> , Dentelaires
<i>Phallus</i> , Morille	<i>Plumbago</i> , Dentelaire
<i>Pharnaceum</i> , Pharnace	<i>Plumeria</i> , Frangipanier
<i>Pharus</i> , Pharelle	<i>Poa</i> , Pâturin
<i>Phascom</i> , Phasque	<i>Podophyllum</i> , Podophylle
<i>Phaseolus</i> , Haricot	<i>Pohlia</i> , Hedw.
<i>Phellandrium</i> , Phellandre	<i>Poinciana</i> , Poincillade
<i>Philadelphus</i> , Syringa	<i>Polemonia</i> , Polémoines
<i>Phleum</i> , Fléole	<i>Polemonium</i> , Polémoine
<i>Phlomis</i> , Phlomide	<i>Polianthes</i> , Tubéreuse
<i>Phlox</i> , Phloxe	<i>Polycnemum</i> , Polycnème
<i>Phœnix</i> , Dattier	<i>Polygala</i> , Polygale
<i>Phormium</i> , Phormion	<i>Polygoneæ</i> , Polygonées
<i>Phylica</i> , Phylica	<i>Polygonum</i> , Renouée
<i>Phyllachne</i> , Phyllachné	<i>Polymnia</i> , Polymnie
<i>Phyllanthus</i> , Phyllanthe	<i>Polypodium</i> , Polypode
<i>Phyllirea</i> , Phylliréa ou Filaria	<i>Polytrichum</i> , Polytric
<i>Physalis</i> , Coqueret	<i>Pomaceæ</i> , Pomacées
<i>Phytolacca</i> , Phytolacca	Id. Sous-ordre des Rosacées de
<i>Picris</i> , Picride	J. Pommiers
<i>Pibularia</i> , Pilulaire	<i>Pommereulla</i> , Pommereulle
<i>Pimpinella</i> , Pimprenelle (Ombellif), Boucage	<i>Pongatium</i> , Pongati
<i>Pinguicula</i> , Grassette	<i>Pontederia</i> , Pontédérie
<i>Pinus</i> , Pin	<i>Populus</i> , Peuplier
<i>Piper</i> , Poivre	<i>Porana</i> , Porana
<i>Piperita</i> , Poivrées	<i>Porella</i> ...
<i>Piscidia</i> , Piscidia	<i>Portulaca</i> , Pourpier
	<i>Portulacææ</i> , Portulacées

R

<i>Potalia</i> , Potalie	<i>Radiati</i> (fiores), Radiées
<i>Potamogeton</i> , Potamot	<i>Rajania</i> , Rajania
<i>Potentilla</i> , Sous-ordre de Rosa- cées de J. Potentilles	<i>Ranunculacea</i> , Renonculacées
<i>Potentilla</i> , Potentille	<i>Ranunculus</i> , Renoncule
<i>Poterium</i> , Pimprenelle	<i>Raphanus</i> , Radis ou Raifort
<i>Pothos</i> , Pothos	<i>Raputia</i> , Raputier
<i>Precia</i> , Précoces	<i>Ravenala</i> , Ravenal
<i>Primula</i> , Primule	<i>Reseda</i> , Réséda
<i>Prockia</i> , Sous-ordre des Rosa- cées de J. Prockies	<i>Restio</i> , Restion
<i>Prockia</i> , Prockia	<i>Reticularia</i> , Réticulaire
<i>Proserpinaca</i> , Proserpine	<i>Rhæades</i> , Rhéades
<i>Protea</i> , Protée	<i>Rhagadiolus</i> , Rhagadiole
<i>Protea</i> , Protées	<i>Rhamni</i> , Nepruns
<i>Prunus</i> , Prunier	<i>Rhamnus</i> , Nerprun
<i>Psidium</i> , Goyavier	<i>Rhaponticum</i> , Rhapontic
<i>Psoralea</i> , Psorale	<i>Rheum</i> , Rhubarbe
<i>Psyllium</i> , Pulicaire	<i>Rhexia</i> , Rhéxie
<i>Ptelea</i> , Ptéléa	<i>Rhinanthus</i> , Cocrète
<i>Pterigynandrum</i> , Hedw....	<i>Rhizobolus</i> , Gærtn....
<i>Pteris</i> , Ptéride	<i>Rhizophora</i> , Palétuvier
<i>Pterocarpus</i> , Ptérocarpe	<i>Rhododendra</i> , Rosages
<i>Pulmonaria</i> , Pulmonaire	<i>Rhododendrum</i> , Rosage
<i>Punica</i> , Grenadier	<i>Rhus</i> , Sumac
<i>Putamineæ</i> .	<i>Ribes</i> , Groseiller
<i>Pyrola</i> , Pyrole	<i>Riccia</i> , Riccie
<i>Pyrus</i> , Poirier	<i>Ricinus</i> , Ricin
	<i>Ricotia</i> , Ricotie
	<i>Robinia</i> , Robinia
	<i>Ropourea</i> , Ropouréa
	<i>Roridula</i> , Roridula
	<i>Rosæ</i> , Sous-ordre des Rosacées de Rosiers
	<i>Rosa</i> , Rosier
	<i>Rosaceæ</i> , Rosacées

Q

Qualea, Qualéa
Quassia, Quassia
Quercus, Chêne
Quillaja, Quillaï

<i>Rosaceæ</i> (<i>Arbores</i>), Arbres Ro- sacés	<i>Sanguisorba</i> , Sous-ordre des Ro- sacées de J. Les Pimprenelles
<i>Rosmarinus</i> , Romarin	ou Sanguisorbes
<i>Rotatæ</i> , Plantes à fleur en Roue	<i>Sanguisorba</i> , Sanguisorbe
<i>Rottbollia</i> , Rottbolle	<i>Sanicula</i> , Sanicle
<i>Rubia</i> , Garance	<i>Santalum</i> , Santal
<i>Rubiaceæ</i> , Rubiacées	<i>Santolina</i> , Santoline
<i>Rubus</i> , Ronce	<i>Sapindi</i> , Savoniers
<i>Rudbeckia</i> , Rudbecke	<i>Sapindus</i> , Savonier
<i>Ruellia</i> , Crustolle	<i>Saponaria</i> , Saponaire
<i>Rumex</i> , Patience	<i>Sapota</i> , Sapotilliers
<i>Ruppia</i> , Ruppie	<i>Saraca</i> , Saraca
<i>Ruscus</i> , Fragon	<i>Sarmentaceæ</i> , Sarmentacées
<i>Ruta</i> , Rue	<i>Sarracenia</i> , Sarracène
<i>Rutaceæ</i> , Rutacées	<i>Sassia</i> , Sassia
<i>Ruyschia</i> , Ruysch	<i>Saturcïa</i> , Sariette
	<i>Satyrium</i> , Satyrion
	<i>Saururus</i> ...
S	
<i>Saccharum</i> , Cannamelle, Canne à Sucre	<i>Sauvagesia</i> , Sauvagèse
<i>Sagina</i> , Sagine	<i>Saxifrage</i> , Saxifrage
<i>Sagittaria</i> , Sagittaire	<i>Saxifragæ</i> , Saxifrages
<i>Sagus</i> , Sagouyer	<i>Scabiosa</i> , Scabieuse
<i>Salacia</i> , Salacia	<i>Scabridæ</i> , Scabrides
<i>Salicaria</i> , Salicaïres	<i>Scandix</i> ...
<i>Salicornia</i> , Salicorne	<i>Schafferia</i> , Schæffer
<i>Salix</i> , Saule	<i>Schefflera</i> , Schefflère
<i>Salsola</i> , Soude	<i>Scheuchzeria</i> , Scheuchzère
<i>Salvia</i> , Sauge	<i>Schinus</i> (Molle)
<i>Salvinia</i> , Salvinie	<i>Schizæa</i> , Smith.
<i>Sambucus</i> , Sureau	<i>Schmidelia</i> , Schmidel
<i>Samolus</i> , Samole ou Mouron d'eau	<i>Schænus</i> , Choin
<i>Samyda</i> , Samyda	<i>Scilla</i> , Scille
<i>Sanguinaria</i> , Sanguinaire	<i>Scirpus</i> , Scirpe
	<i>Scitamineæ</i> , Scitaminées ou Epicées

<i>Sclerocarpus</i> , Sclérocarme	<i>Sideroxylum</i> , Argan
<i>Scelopendrium</i> , Scolopendre	<i>Sigesbeckia</i> , Sigesbeckie
<i>Scolymus</i> , Scolyme	<i>Silene</i> , Siléné
<i>Scopolia</i> , Scopoli	<i>Siliquosa</i> , Siliqueuses
<i>Scorpiurus</i> , Chenillette	<i>Silphium</i> , Silphium
<i>Scorzonera</i> , Scorsonère	<i>Simbuleta</i> , Simbulêta
<i>Scrophularia</i> , Scrophulaire	<i>Sinapis</i> , Moutarde
<i>Scrophulariæ</i> , Scrophulaires	<i>Siparuna</i> , Siparuna
<i>Scutellaria</i> , Toque	<i>Sison</i> , Sison
<i>Secale</i> , Seigle	<i>Sisymbrium</i> , Sysimbre
<i>Securidaca</i> , Sécuridaca	<i>Sisyrrinchium</i> , Bermudienne
<i>Sedum</i> , Orpin	<i>Smithia</i> , Smithe
<i>Seguiera</i> , Séguier	<i>Siun</i> , Berle
<i>Selago</i> , Selago	<i>Skimmia</i> , Skimmia
<i>Selinum</i> , Sélinum	<i>Smilax</i> , Smilax
<i>Semi-Flosculosi (flores)</i> , Demi-Flosculeuses	<i>Smyrniun</i> , Mâceron
<i>Sempervivæ</i> , Joubarbes	<i>Solaneæ</i> , Solanées
<i>Sempervivum</i> , Joubarbe	<i>Solanum</i> , Morelle
<i>Senecio</i> , Sénéçon	<i>Soldanella</i> , Soldanelle
<i>Senticosæ</i> , Sentiqueuses ou Touffues	<i>Solidago</i> , Verge d'Or
<i>Sepiariæ</i> , Sépiaires ou de Haie	<i>Sonchus</i> , Laitron
<i>Septas</i> , Septas	<i>Sonneratia</i> , Pagapate
<i>Serapias</i> , Sérapias	<i>Sophora</i> , Sophora
<i>Seriola</i> , Sérieole	<i>Soramia</i> , Soramia
<i>Seriphium</i> , Armoselle	<i>Sorbus</i> , Sorbier
<i>Serratula</i> , Sarrête	<i>Soulamea</i> , Soulaméa
<i>Sesamum</i> , Sésame	<i>Souroubea</i> , Souroubéa
<i>Seseli</i> , Séséli	<i>Sparganium</i> , Ruban d'eau
<i>Sesuvium</i> , Sésuvium	<i>Spathuceæ</i> , Spathacées
<i>Sherardia</i> , Shérarde	<i>Spergula</i> , Spargoute
<i>Sicyos</i> , Sicyos	<i>Spermacoce</i> , Spermacocée
<i>Sida</i> , Abutilon	<i>Sphæranthus</i> , Sphæranthe
<i>Sideritis</i> , Crapaudine	<i>Sphæria</i> , Variolaria Variolaire
	<i>Sphagnum</i> , Sphaigne
	<i>Spigelia</i> , Spigèle

<i>Spinacia</i> , Epinars	<i>Tamarindus</i> , Tamarinier
<i>Spireæ</i> , Sous-ordre des Rosa- cées de J. Spirées	<i>Tamarix</i> , Tamaris
<i>Spiræa</i> , Spirée	<i>Tamnus</i> , Tâme ou Taminier
<i>Splachnum</i> , Splaigne	<i>Tanacetum</i> , Tanaïs
<i>Stachys</i> , Stachyde	<i>Tapura</i> , Tapura
<i>Stapelia</i> , Stapélie	<i>Taraxacum</i> , Pissenlit
<i>Staphylea</i> , Staphyléa	<i>Tarchonanthus</i> , Tarconanthe
<i>Statice</i> , Staticée	<i>Targionia</i> , Targiône
<i>Statuminatæ</i> , Statuminées ou Appuis de Vigne	<i>Taxus</i> , If.
<i>Stellaria</i> , Stellaire	<i>Telephium</i> , Télèphe
<i>Stellatæ</i> , Etoilées	<i>Temus</i> , Témô
<i>Stelleria</i> , Stèllère	<i>Terminalia</i> , Badomier
<i>Sterculia</i> , Sterculia	<i>Tetracera</i> , Tétracera
<i>Stilbe</i> , Stilbé	<i>Tetragonia</i> , Tétragône
<i>Stipa</i> , Stipe	<i>Teucrium</i> , Germandrée
<i>Stæbe</i> , Stæbé	<i>Thalia</i> , Thalia
<i>Stratiotes</i> , Stratiote	<i>Thalictrum</i> , Pigamon
<i>Strumpfia</i> , Strumpfia	<i>Thapsia</i> , Thapsie
<i>Struthiola</i> , Struthiola	<i>Thea</i> , Thé
<i>Styrchnos</i> , Vomique	<i>Theligonum</i> , Theligône
<i>Styrax</i> , Aliboufier	<i>Theobroma</i> , Cacaoyer
<i>Succulentæ</i> , Succulentes ou Plantes Grassés	<i>Therebintaceæ</i> , Thérébintacées
<i>Suillus</i> . J...	<i>Therebinthus</i> , Thérébinthe
<i>Suriana</i> , Suriana	<i>Thesium</i> , Thesium
<i>Swartzia</i> , Hedw....	<i>Thlaspi</i> , Thlaspi
<i>Swietenia</i> , Mahogon	<i>Thuya</i> , Thuya
<i>Symphytum</i> , Consonde	<i>Thymeleæ</i> , Thymélées
<i>Symplocos</i> , Symplocos	<i>Thymus</i> , Thym
	<i>Tiarella</i> , Tiarelle
T	<i>Tilia</i> , Tilleul
<i>Tabernæmontana</i> , Taberné	<i>Tilliaceæ</i> , Tiliacées
<i>Tacca</i> (Herbe), Tacca	<i>Tillæa</i> , Tillée
<i>Tagetes</i> , Œillet d'Inde	<i>Tillandsia</i> , Tillandsia
	<i>Timmia</i> , Hedw....
	<i>Tolwifera</i> , Tolut
	<i>Tomex</i> , Tomex

<i>Tonina</i> , Tonine	<i>Trollius</i> , Trolle
<i>Tontelea</i> , Tontéléa	<i>Tropaeolum</i> , Capucine
<i>Tordylium</i> , Tordylium	<i>Trophis</i> , Trophis
<i>Tormentilla</i> , Tormentille	<i>Tuber</i> , Truffe
<i>Tortula</i> , Hedw....	<i>Tulbagia</i> , Tulbagie
<i>Tounatea</i> , Tounatéa	<i>Tulipa</i> , Tulipe
<i>Tournefortia</i> , Tournefort	<i>Turnera</i> , Turnéra
<i>Touroulia</i> , Tourouli	<i>Turræa</i> , Turræa
<i>Trachelium</i> , Gantelé	<i>Turritis</i> , Turrète ou Totrette
<i>Tradescantia</i> , Ephémère	<i>Tussilago</i> , Tussilage
<i>Tragia</i> , Tragia	<i>Typha</i> , Massette
<i>Tragopogon</i> , Cersifis	<i>Typhæ</i> , Massettes
<i>Trapa</i> , Mâcre	
<i>Tremella</i> , Trémelle	U
<i>Trewia</i> , Tréwia	<i>Ulex</i> , Ajonc
<i>Trianthema</i> , Trianthème	<i>Ulmus</i> , Orme
<i>Trichia</i> , Bull. Capilline	<i>Ulva</i> , Ulve
<i>Tribulus</i> , Herse	<i>Ulvæ</i> , Ulves, Sous-ordre
<i>Trichilia</i> , Trichilia	<i>Umbelliferæ</i> , Ombellifères
<i>Trichomanes</i> , Trichomâne	<i>Uniola</i> , Uniolle
<i>Trichosanthes</i> , Anguine	<i>Urena</i> , Uréna
<i>Tricocœ</i> , Arbres portant un fruit à trois Coques	<i>Urtica</i> , Ortie
<i>Trifolium</i> , Trèfle	<i>Urticæ</i> , Orties
<i>Triglochin</i> , Triglochine	<i>Utricularia</i> , Utriculaire
<i>Trigonella</i> , Trigonelle, Fenu- grec	<i>Uvaria</i> , Uvaria
<i>Trihilatæ</i> , Triumbliquées	<i>Uvularia</i> , Uvulaire
<i>Trilix</i> , Trilix	V
<i>Trillium</i> , Trillie	<i>Vaccinium</i> , Airelle
<i>Triopteris</i> , Triopteris	<i>Vaginales</i> , Vaginales ou Plantes à gânes
<i>Triosteum</i> , Triosté	<i>Valantia</i> , Croisette
<i>Tripetaloidæ</i> , Tripétaloïdes	<i>Valeriana</i> , Valériane
<i>Tripsacum</i> , Tripsaque	<i>Vallea</i> , Valléa
<i>Triticum</i> , Froment	<i>Vallisneria</i> , Vallisnérie
<i>Trionfetta</i> , Lapullier	<i>Vanilla</i> , Vanille

<i>Vantanea</i> , Vantanéa	<i>Weinmannia</i> , Weinmannia
<i>Variolaria</i> , Bull. Variolaire	<i>Willichia</i> , Willichia
<i>Vepreculæ</i> (Ce sont les Tymé- lées de J.)	<i>Wittaria</i> , Smith....
<i>Veratrum</i> , Varaire	<i>Woedwardia</i> , Id....
<i>Verbascum</i> , Molène	X
<i>Verbena</i> , Verveine	<i>Xanthium</i> , Lampourde
<i>Verbesina</i> , Verbésine	<i>Xeranthemum</i> , Immortelle
<i>Verticillatæ</i> , Verticillées	<i>Xerophyta</i> , Xérophyta
<i>Veronica</i> , Véronique	<i>Xilopia</i> , Xilopia
<i>Verrucaria</i> , Hoffm....	<i>Ximenia</i> , Ximédia
<i>Viburnum</i> , Viorne	<i>Xylophylla</i> , Xylophylla
<i>Vicia</i> , Vesce	
<i>Vinca</i> , Pervenche	Y
<i>Viola</i> , Violette	<i>Yucca</i> , Yucca
<i>Viscum</i> , Gui	
<i>Vites</i> , Vignes	Z
<i>Vitex</i> , Gattilier	<i>Zamia</i> , Zamia
<i>Vitices</i> , Gattiliers	<i>Zanichellia</i> , Zanichelle
<i>Vitis</i> , Vigne	<i>Zanthoxylum</i> , Clavaliér
<i>Vochisia</i> , Vochy	<i>Zea</i> , Maïs
	<i>Zinnia</i> , Zinnie
W	<i>Zizania</i> , Zizane ou Zizanie
<i>Wachendorfa</i> , Wachendorfe	<i>Ziziphus</i> , Jujubier
<i>Webera</i> , Hedw....	<i>Zoëgea</i> , Zoégée
<i>Weissia</i> , Hedw....	<i>Zostera</i> , Zostère
<i>Weigela</i> , Weigéla	<i>Zygophyllum</i> , Fabagelle

PART THIRD.

CHAPTER I.

OF VEGETABLES, AND THEIR KINDS.

VEGETABLES are divisible into the seven *families*, or *tribes*, as follows :

1. FUNGI, *mushrooms*.
2. ALGÆ, *algæ*; whose root, leaf, and stem are all one.
3. MUSCI, *mosses*; whose antheræ have no filaments, and are placed at a distance from the female flower; and whose seeds also want their proper tunic and cotyledons.
4. FILICES, *ferns*; whose fructification is on the back of the *frondes**.
5. GRAMINA, *grasses*†; which have simple leaves, a jointed culm or stem, a glumose calyx, and a single seed.

* *Leaves* of the ferns and palms so called; see the explanation of the term *frons*, in Chap. IV. AUTHOR.

† This tribe includes the various sorts of corn as well as the *grasses*. AUTHOR.

6. PALMÆ, *palms*; which have simple stems, that are *frondose** at the summit, and have their fructifications on a spadix issuing from a spatha.

7. PLANTS, which include all that do not enter into any of the other divisions. These are,

1. *Herbaceous*, when they die down to the root every year; for in the perennial kinds, the buds are all produced on the root below the surface of the ground.

2. *Shrubs*, when their stems come up *without* buds†.

3. *Trees*, when their stems come up *with* buds.

Vegetables are each primarily divisible into, 1. The *root*...2. The *herb* or plant itself...3. The *fructification*. Of these the *last* has been already treated of in the first book: the two others, upon which the specific differences of vegetables more immediately depend, come now under consideration, and will be the subject matter of the ensuing chapters‡.

* See the term *frons*, explained in Chap. IV. AUTHOR.

† Nature has put no limits between a tree and a shrub, which is only a vulgar distinction. This *Linnæus* acknowledges; and argues, that his own distinction, though he thinks it the best, is nevertheless exceptionable; inasmuch as there are seldom any buds upon the large trees in India; all which must, therefore, by this definition, notwithstanding their great height, be ranked with shrubs. AUTHOR.

‡ It may not be improper here to obviate an objection that may be made to the method pursued in this work. It may be asked, if the matter of this third part would not have stood more properly in the first. In answer to this it is admitted, that the *order of nature* would thereby have been more directly followed: but the design of this work was not so much to follow the order of nature, as to explain the *System of Linnæus*; and as the Classes, Orders, and Genera, which come first in the system, are grounded on the fructification, the beginning with that part of the vegetable was indispensably necessary. AUTHOR.

CHAP. II.

OF ROOTS.

THE root, whose office is to draw up nourishment, and which also produces the herb, with its fructification, consists of two parts, viz. CAUDEX, the *stock* or body of the root; and RADICULA, the *radicle* or little root.

CAUDEX, the *body* of the root both ascends and descends.

The *ascending caudex* raises itself gradually above ground, serving often as a trunk, and produces the herb or plant*.

The *descending caudex* strikes gradually downward into the ground, and puts forth radicles. It has been distinguished, according to its various structure, into

1. PERPENDICULAR, when it runs directly downwards.
2. HORIZONTAL, when it extends itself transversely under the earth.
3. SIMPLE, when it has no subdivisions.
4. RAMOSE, *branching*; when it is divided into lateral branches.
5. FUSIFORM, *spindle-shaped*; when it is oblong, thick, and tapering, as in DAUCUS and PASTINACA.
6. TUBEROSE, *knobbed*; when it consists of roundish bodies collected into a fascicle or bunch, as in PEONEA,...HEMEROCALLIS,...HELIANTHUS,...SOLANUM,...and FILIPENDULA.

* *Linnaeus* infers from hence, that all trees and shrubs are to be considered as roots above ground; and that this is the reason that trees, when inverted, put forth leaves from the descending stem, and roots from the ascending. AUTHOR.

7. REPENT, *creeping*; when it runs out to a distance, and puts forth radicles from space to space.

8. FIBROSE, when it consists only of fibrose radicles.

9. PRÆMORSE, *bitten off*; when the lower part is truncate, and the termination not tapering, as in SCABIOSA,...PLANTAGO,...and VALERIANA*.

RADICULA, the *radicle*, is the fibrose part of the root, which terminates the descending caudex, and enables the root to draw nourishment for the support of the vegetable.

CHAP. III.

OF THE HERB.

THE herb is a part of the vegetable arising from the root, and terminated by the fructification. It comprehends,

1. The TRUNK, which serves to multiply the herb, and leads immediately from the root to the fructification. It is clothed with the leaves, and terminated by the fructification. See Chap. IV.

2. The LEAVES, whose office is to transpire and attract, like the lungs in animals, and to afford shade. See Chap. V. VI. VII.

3. The FULCRA, *props*; which serve as stāys to strengthen the plant; but may, however, be taken off without destroying it. See Chap. VIII.

* For figures of these, *vide* our Plates, attached at the end of this work.

4. The HYBERNACULA, *winterings**; each of which is a compendium of the herb upon its root before it begins to grow. See Chap. IX.

CHAP. IV.

OF THE TRUNK.

TRUNCUS, the *trunk*, is that which produces the leaves and fructification: it is of seven kinds, viz. *Caulis*,...*culmis*,...*scapus*,...*pedunculus*,...*petiolus*,...*frons*,...and *stipes*.

I. CAULIS, a *stem*, is the proper trunk of the herb, and serves to elevate the leaves and fructification: it is either *simple*...or *compound*.

SIMPLE stems are such as proceed in a continued series towards their summits; and these may be,

1. INTEGRI, *entire*; when they are most simple, having scarce any branches.

2. NUDI, *naked*; when they are destitute of leaves, as in EU-PHOREIA,...CACTUS,...STAPELIA,...EPHEDRA,...and CUSCUTA.

3. FOLIATE, *leafy*; when they are furnished with leaves.

4. FLEXUOSE, *bending* different ways, when the direction of the stem changes at every joint, as in PTELIA.

5. VOLUBILES, *twining*; when they ascend spirally by the branch of some other plant†: these wind either to the *left*, according to the motion of the sun (as it is commonly phrased),

* These are the bulbs and buds. EDITOR.

† Vide Plate V. Figure 4, of this work.

as in HUMULUS,...HELIXINE,...LONICERA,.....and TAMUS; or to the *right*, contrary to the sun's motion, as in CONVULVUS,... BASELLA,...PHASEOLUS,...CYNANCHE,...EUPHORBIA,... and EUPATORIUM.

6. RECLINATE, *reclined*; when they bend in an arch towards the earth.

7. PROCUMBENT, *lying upon the ground*; when their direction is horizontal.

8. REPENT, *creeping*; when, by lying upon the ground, they put forth roots at certain intervals, as in HEDERA and BIGNONIA*.

9. SARMENTOSE†; when they are repent and subnude‡.

10. PARASITIC§; when they grow not out of the ground, but on some other plant.

11. TERETES, *round*; when they are cylindric.

12. ANCIPITES, *double-edged*; when they have two opposite angles; and also DIGONUS, TRIGONUS, TETRAGONUS, PENTAGONUS, POLYGONUS, having *two, three, four, five, or many angles*, which are all species of ancipites: also,

13. TRIQUETROUS, *three-square*; when they have three plane sides; and,

14. TRIANGULAR, QUADRANGULAR, QUINQUANGULAR, MULTANGULAR; when they have *three, four, five, or many sides or angles*.

15. SULCATE, *furrowed*; when they are cut in with broad and deep grooves or channels.

16. STRIATE, *streaked*; when they are marked with very thin hollow lines.

* Vide Plate V. Figure 2, of this work.

† From SARMENTUM, a long shoot, such as those of a vine. EDITOR.

‡ Almost naked or bare of leaves. EDITOR.

§ Supporting themselves on others, like parasites. EDITOR.

17. GLABRI, *smooth*; when they have a smooth surface.
18. VILLOSE, *hairy* or *shaggy*; when there is a down of soft hairs upon them.
19. SCABROUS, *rough*; when they are covered with little projecting points.
20. HISPID*; when they are covered with stiff bristles.
21. RAMOSE, *branchy*; when they are furnished with lateral branches; and these are,
22. ASCENDING; when the branches incline upwards.
23. DIFFUSE; when the branches are spreading.
24. DISTICH, in *two rows*; when the branches are produced in a horizontal situation.
25. BRACHIATE, having *arms*; when the branches are opposite, and each pair is crossed by the pair next above or below it†.
26. RAMOSISSIMI, *very branchy*; when the branches are many, and without order.
27. FULCRATE, *propt*; when the branches descend to the root, as in FICUS.
28. PROLIFEROUS; when they send forth branches only from the centre of the apex, as in PINUS.
- The rest as in *entire* stems.

COMPOUND stems, are such as are subdivided into RAMULI, *small branches*, and diminish as they ascend. These are either,

1. DICHOTOMUS, *forked*; when the division is always in two parts ‡.

* The word expresses a greater degree of roughness. EDITOR.

† Vide Plate V. Figure 8, of this work.

‡ Vide Plate V. Figure 7, of this work.

2. SUBDIVIDED; when they are divided into branches irregularly or without order: or,

3. ARTICULATE, *jointed*; when they are distinguished from space to space, by knots or joints, as in PIPER*.

II. CULMUS, a *straw*, is the proper stem or trunk of a grass, and serves to elevate and support both the leaves and the fructification: it admits of most of the distinctions already given for a caulis or stem; besides which, it may be either.

1. ENODIS, *without knots*; when it is continuous, and not intercepted by joints.

2. ARTICULATE, *jointed*; when it is connected by various joints.

3. SQAMOSE, *scaly*; when it is covered with imbricate scales†.

III. SCAPUS, a *stalk*, is an *universal* trunk, raising the fructification, but not the leaves, as in NARCISSUS,...PYROLA,...CONVAL-LARIA,...and HYACINTHUS‡.

IV. A PEDUNCLE, or *foot-stalk* of a *flower*, is a *partial* trunk, raising the fructification, but not the leaves.

PEDICELLUS, is a *partial peduncle*.

The determination of *peduncles* respects *place* and *manner*.

Determination in respect to *place*, shows where the base of the peduncle is inserted into the plant: and in this respect peduncles are,

1. RADICAL, belonging to the *root*; when they come out immediately from the root.

2. CAULINE, belonging to the *stem*; when they are placed on the stem.

3. RAMEOUS, belonging to the *branches*; when they come out upon the branches.

* Vide Plate V. Figure 5, of this work.

† Vide Plate V. Figure 1, of this work.

‡ Vide Plate V. Figure 6, of this work.

4. AXILLARY*, coming out from the wings; that is, either between the leaf and the stem, or between the branch and the stem.

5. TERMINAL, when they *terminate* the branches or stem.

6. SOLITARY, when there comes out but *one* from the same place.

7. SPARSE, *scattered*; when they are numerous, and come out without order.

Determination in respect to *manner*, shows how the flowers are placed and connected on the summits of the *peduncles*: and in this respect *peduncles* have the following variations:

1. UNIFLOROUS, BIFLOROUS, TRIFLOROUS, or MULTIFLOROUS peduncles, are such as bear *one, two, three, or many flowers*, according to the number of the fructifications on a single peduncle.

2. FASCICULUS, a *bunch*, is a collection of flowers that are erect, parallel, forming a flat or even surface, and close to one another; as in DIANTHIUS BARBATUS†.

3. CAPITULUM, a *little head*, is composed of a number of flowers, collected almost into a globular form, as in GOMPHRENA.

4. SPICA, a *spike*, has sessile flowers that are alternate and dispersed about a common peduncle that is simple. It is called SPICA SECUNDA, a *single-rowed spike*, when the flowers are all turned *one way*: and SPICA DISTICHA, a *double-rowed spike*, when the flowers stand *two ways*.

5. A CORYMBUS‡, is a kind of spike, the flowers of which have

* From AXILLA, an *arm-pit*. EDITOR.

† Sweet William. EDITOR.

‡ *Corymbus*, in its ancient and proper signification, meant a bunch of ivy berries: but is now used as a botanical term, for all fructifications that are produced in this manner. EDITOR.

each its proper PEDICELLUS*, or *partial foot-stalk*, raised to a proportionable height, as in SPIRÆA OPULIFOLIA,...and LEDUM.

6. A PANICLE, is a fructification dispersed on peduncles variously subdivided. It is a DIFFUSE panicle, when the pedicelli are *divaricate, spreading asunder*; and a COARCTATE or *confined* one, when they stand close to each other.

7. A THRYBUS, is a panicle contracted into an ovate form, as in SYRINGA and PETASITES.

8. A RACEMUS† consists of a peduncle that has short lateral branches, as in VITIS...and RIBES.

9. VERTICILLUS, a *whorl*, expresses a number of flowers that are subsessile‡, and are produced in rings round the stems.

V. A PETIOLE, or *foot-stalk of a leaf*, is a species of trunk that fastens the leaves, but not the fructification; which circumstance distinguishes it from a peduncle, which is the foot-stalk of a flower, as has been explained above. There are some cases where the fructification and leaves are born on the same foot-stalks, as in TURNERA...and HIBISCUS; but these instances are very rare.

VI. FRONS§, is a species of trunk, composed of a branch and leaf blended together; and is frequently united with the fructification: it belongs properly to the PALMS...and FILICES||.

* In the *Philosophia Botanica*, it is not *Pedicellus*, but *Petiolus*; which seems to be a mistake, this term being applied to leaves only. It may be translated *Peduncle*. EDITOR.

† *Racemus*, anciently signified a bunch of grapes. EDITOR.

‡ With no foot-stalks, or with very short ones. EDITOR.

§ There is no expression answerable to this term in our language. See the note at page 67. AUTHOR.

|| Vide Plate V. Figure 3, of this work.

VII. STIPES*, is used to express the base or trunk of a *frons*, and is applied only to the PALMS...FILICES...and FUNGI:

CHAP V.

OF SIMPLE LEAVES.

LEAVES are to be considered in three respects, viz. as SIMPLE...2. COMPOUND...3. DETERMINATE. We shall in this chapter treat only of the simple.

SIMPLE leaves are such as have only a single leaf on a petiole. They differ in respect to *circumscription...angles...sinus...apices...margin...superficies...and substance*:

I. CIRCUMSCRIPTION considers the form of the circumference of leaves where there are no angles or sinuations: in which respect leaves are,

1. ORBICULATE, *round*; when the longitudinal and transverse diameters are equal, and the circumference circular.
2. SUBROTUND, *roundish*; when the figure is nearly orbiculate.
3. OVATE, *egg-shaped*; when the longitudinal diameter exceeds the transverse, and the base is circumscribed with the segment of a circle, but the apex is narrower.
4. OVAL, or *elliptic*; when the longitudinal diameter exceeds

* The word in its proper signification means a trunk or stock of any plant: but the sense in which the term is received in botany is as here explained: it is used also to express the thread or fine trunk that supports the pappus in downy seeds. See Part I., Chap. VII. AUTHOR.

the transverse, and the circumscription of both upper and lower extremity is narrower than the segment of a circle.

5. PARABOLIC, in the form of a *parabola**; when the longitudinal diameter exceeds the transverse, and the figure contracting from the base upwards becomes SEMIOVATE, *half-egg-shaped*.

6. SPATULATE, resembling a *spatula*†; when the figure is roundish, but lengthened out by the addition of a linear base that is narrower.

7. CUNEIFORM, *wedge-shaped*; when the longitudinal diameter exceeds the transverse, and the figure gradually contracts downwards.

8. OBLONG, when the longitudinal diameter is twice, thrice, &c. the length of the transverse, and the circumscription of each of the extremities is narrower than the segment of a circle.

II. ANGLES are the prominent parts of a horizontal leaf. In respect to these, a leaf is,

1. LANCEOLATE, *spear-shaped*; when the figure is oblong, narrowing gradually at each end towards the extremity.

2. LINEAR; when it is every where of the same breadth, though sometimes narrowing at the extremities only.

3. ACEROSE, *chaffy*; when it is linear and persisting as in PINUS, ... ABIES, ... JUNIPERUS, ... and TAXUS.

4. SUBULATE, *awl-shaped*; when it is linear below, but gradually contracting towards the top.

5. TRIANGULAR, *three-cornered*; when the disk is surrounded by three prominent angles.

* A geometric curve so called. EDITOR.

† A surgeon's instrument so called. EDITOR.

6. QUADRANGULAR, *quincunangular*, &c. *four-cornered*, *five-cornered*, &c. when four or five prominent angles lie round the disk.

7. DELTOID, shaped like a *delta**; when the figure is a rhombus; that is, having four angles, of which the two lateral ones are less distant from the centre than those at the extremities.

8. ROTUND, round; when it has no angles.

III. SINUS, a *hollow*, is a term used to express those openings or cavities in leaves, which distinguish them into parts: in respect to these, leaves are said to be,

1. RENIFORM, *kidney-shaped*; when they are roundish, and hollowed at the base, without any angles.

2. CORDIFORM, *heart-shaped*; when they are ovate, and hollowed at the base, and the hinder or lower part has no angles.

3. LUNULATE, *moon-shaped*; when they are round, and hollowed at the base, and the lower part has no angles.

4. SAGITTATE, *arrow-shaped*; when they are triangular, hollowed at the base, and are furnished with angles at the lower part.

5. HASTATE, *javelin-shaped*; when they are triangular, the base and sides hollowed, and the angles spreading.

6. PANDURÆFORM, *pandure-shaped*†; when they are oblong, broader above than below, and contracted in the sides.

7. FISSA, *cloven*; when they are divided by linear sinusses,

* A Greek letter so called. The figure of the *delta* is a triangle, which does not exactly answer to the character here given of a *deltoid* leaf.

† A musical instrument of the lute kind, but now disused: the shape of it, as given by *Marsemus, Harm. Instr.* l. 1. does not answer to that of the leaves here explained; the figure of which comes nearer to that of the body of a violoncello or violin. AUTHOR.

and have their margins straight ; and from the number of such divisions they are called BIFID, TRIFID, QUADRIFID, MULTIFID, &c. cut into *two, three, four, five, or many segments.*

8. LOBATE, *lobed* ; when they are divided to the middle into parts that stand wide from each other, and have their margins convex ; and from the number of these they are called BILOBE, TRILOBE, QUADRILLOBE, or QUINQUELOBE ; consisting of *two, three, four, or five lobes.*

9. PALMATE, *handed* ; when they are cut longitudinally into many parts, nearly equal ; the divisions extending themselves downward, almost to the base, where the segments cohere.

10. PINNATIFID, cut into *wings* ; when they are divided transversely into laciniaë that are oblong and horizontal.

11. LYRATE, *lyre-shaped* ; when they are divided transversely into laciniaë, of which the upper ones are larger, and the lower ones farther asunder.

12. LACINIATE, *jagged* ; when they are variously divided into parts, and those parts in like manner indeterminately subdivided.

13. SINUATE, *hollowed* ; when they have broad and spreading openings in the sides.

14. PARTITE, *divided* ; when they are separated down to the base ; and from the number of the divisions they are BIPARTITE, TRIPARTITE, QUADRIPARTITE, QUINQUEPARTITE, or MULTIPARTITE ; divided into *two, three, four, five, or many parts.*

15. INTEGRA, *entire* ; when they are without divisions, and have no sinus or opening. This stands opposed to all the kinds of divided leaves before described.

IV. APEX, *tip*, is the extremity in which the leaf terminates. Leaves, in respect to their apices, are called,

1. TRUNCATE, *lopped* ; when they end in a transverse line,

2. *PRÆMORSE*, *bitten in the fore-part*; when they are very obtuse, and are terminated by unequal notches or incisions.

3. *RETUSE*, *blunted*; when they terminate in an obtuse sinus.

4. *EMARGINATE*, *nicked*; when they terminate in a notch.

5. *OPTUSE*, *blunt*; when they terminate, as it were, within a segment of a circle.

6. *ACUTE*, *sharp*; when they terminate in an acute angle.

7. *ACUMINATE*, *pointed*; when they terminate in a subulate apex.

8. *CIRRHOSE*, *tendriled*; when they terminate in a clasper or tendril, as in *GLORIOSA*,...*FLAGELLARIA*,...and *NISSOLIA*.

V. The *MARGIN* of a leaf is the outermost boundary of its sides, exclusive of its disk. Leaves, in respect to their margin, are,

1. *SPINOSE*, *thorny*, or prickly; when the margin of the leaf runs into points that are hard, stiff, and pungent.

2. *INERM*, *unarmed* or smooth: which is opposed to spinose.

3. *DENTATE*, *toothed* or indented; when the margin ends in horizontal points, that are of the consistence of the leaf, and are separated by intermediate spaces.

4. *SERRATE*, *sawed*; when the margin is cut into sharp imbricate angles, that point towards the extremity of the leaf: if they point towards the base, the leaf is said to be *RETRORSUM SERRATE*, *sawed backwards*.

5. *DUPLICATO-SERRATE*, *doubly sawed*; when there is a two-fold serrature, the less upon the greater.

6. *CRENATE*, *notched*; when the margin is cut into angles, that point towards neither of the extremities; and these are obtusely

crenate, when the angles are rounded; or acutely crenate, when the angles are pointed.

7. *DUPPLICATO-CRENATE*, *doubly notched*; when the notches are two-fold, the less upon the greater.

8. *REPAND*, *bending back again*; when the margin is terminated with angles, and interjacent sinusses, that are both inscribed with the segments of circles*.

9. *CARTILAGINEOUS*, *bristly*; when the edge of the leaf is strengthened by a tough border, the substance of which differs from that of the leaf.

10. *CILIATE*, *lashed or fringed*; when the margin is surrounded on all sides with parallel bristles.

11. *LACERA*, *rent or ragged*; when they are variously cut on the margin into unlike segments.

12. *EROSE*, *gnarwed*; when the leaf is sinuate, and has other very small obtuse sinusses or hollows on its margin.

13. *INTEGERRIMA*, *very entire*; when the outermost margin is entire and quite free from notches.

VI. *SUPERFICIES*, *surface*, is the outside, or what covers the disk of the leaf, and respects both the *supine*† disk or face of the leaf, and *prone* disk or back of it. Leaves, in respect to their surface, are,

1. *VISCID*, *clammy*; when they are smeared over with a juice that is not fluid, but tenacious, sticky.

2. *TOMENTOSE*, *downy*; when they are covered with a nap of interwoven hairs, scarce perceptible, that gives them a whiteness.

* A serpentine edge. EDITOR.

† *Supine* is what lies on its back, or face upwards; and *prone*, the contrary: these terms are, therefore, well applied to the upper and under disk or face of a leaf. EDITOR.

3. LANATE, *woolly*; when they are covered, as it were, with a spider's web, as in SALVIA...and SIDERITIS.
4. PILOSE, *hairy*; when their surface is covered with distinct hairs, that rise to some length.
5. HIRSUITE, *rough with hair*; when they are hairy in a greater degree.
6. VILLOSE, *sluggish*; when they are covered with a coarser hair or shag.
7. HISPID, *rough*; when the disk is covered with a stiffish sort of bristles, that are frangible.
8. SCAEROUS, *rugged*; when the disk is covered with tubercles, little knobs.
9. ACULEATE, *prickly*; when the disk is beset with points that are sharp and stiff.
10. STRIATE, *streaked*; when the surface is cut in, or scored longitudinally with parallel lines.
11. PAPPILLOSE, *nipply*; when it is covered with *vesicles*, or *little bladders*.
12. PUNCTATE, *dotted*; when it is besprinkled with hollow points or dots.
13. NITID, *bright*; when the smoothness of the leaves causes them to shine.
14. PPLICATE, *plaited*; when the disk of the leaf rises and falls in angles towards the margin, as in ALCHEMILLA.
15. UNDULATE, *waved*; when the disk of the leaf rises and falls in convexities towards the margin.
16. CRISP, *curled*; when the circumference of the leaf becomes larger than the disk admits of, and is hereby forced to undulate. All curled leaves are monsters.

17. *RUGOSE, wrinkled*; when the veins of the leaves contract into a narrower compass than the disk, so that the substance between them is obliged to rise, as in *SALVIA*.

18. *CONCAVE, hollow*; when the margin of the leaf contracts, and becomes less than the circumscription of the disk, by which means the disk is depressed.

19. *VENOSE, veiny*; when the vessels are branched all over the leaves, and their anastomose* or joinings are plain to the naked eye.

20. *NERVOSE*; when they have simple unbranched vessels, that extend themselves from the base to the apex.

21. *COLOURED*; when they change their green for some other colour, as in *AMARANTHUS TRICOLOR*†.

22. *GLABRA, smooth*; when the surface is void of all inequality.

VII. The *SUBSTANCE* of a leaf respects the conditions of its sides: in this respect leaves are,

1. *TERETES* ‡, *round*, like a pillar; when they are for the most part cylindric.

2. *SEMICYLINDRIC*, like a *halved cylinder*; when they are round on one side, and flat on the other.

3. *TUBULOSE*, like a *tube* or pipe; when upon cutting them they appear to be hollow within.

4. *CARNOSE, fleshy* or succulent; when they are filled with a pulp.

* A term in anatomy, expressing the union of veins and arteries; or where they pass from one branch to the other in smaller channels. EDITOR.

† Three-coloured.

‡ Round one way and long the other: our language has no distinct term to express roundness in this sense; the figure is, by mathematicians, called a cylinder, from a Greek word, signifying to roll; a body of this figure being the best adapted to that sort of motion.

5. COMPRESSED, *flatted*; when they are so compressed by their opposite marginal sides, that the substance of the leaf becomes greater than the disk.

6 PLANE, *level*; when they have both surfaces every where parallel.

7. GIBBOUS, *bunched*; when, by the plenty of the pulp, both the surfaces are rendered convex.

8. CONVEX, *rounding*; when the disk rises higher than the sides.

9. DEPRESSED, *pressed down*; when the sides rise higher than the disk.

10. CANALICULATE, *channelled*; when a deep furrow runs along it, and sinks it almost to a half cylinder.

11. ANCIPITES, *double-faced*; when the disk is convex, and there are two prominent longitudinal angles.

12. ENSIFORM, *sword-shaped*; when they are ancipites, and grow narrower from the base to the apex.

13. ACINACIFORM, *fulchion* or *scimitar-shaped*; when they are fleshy and compressed, with one edge convex and narrow, and the other straighter and broader.

14. DOLABRIFORM, *hatchet-shaped*; when their figure is roundish, compressed, and obtuse; gibbous outwardly, with a sharp edge, and taper towards the lower part.

15. LINGUEFORM, *tongue-shaped*; when they are linear, fleshy, obtuse, convex underneath, and often with a cartilaginous margin.

16. TRIQUETROUS, *three-corned*; when they are subulate, and have three flat longitudinal sides.

17. SULCATE, *furrowed*; when they are scored longitudinally

with numerous angles or ridges, and as many hollows or channels betwixt them.

18. *CARINATE*, *keeled*; when the prone part of the disk is prominent longitudinally.

19. *MEMBRANACEOUS*; when they have no perceptible pulp between the two surfaces*.

CHAP. VI.

OF COMPOUND LEAVES.

A *LEAF* is said to be compound, when there are more than one upon a common petiole or foot-stalk.

COMPOUND leaves are to be considered in respect to *structure* and *degree*.

I. By the *STRUCTURE* of a compound leaf is to be understood the insertion of the folioles or lesser leaves, of which it is compounded; and in this respect leaves are called,

1. *COMPOUND*; when a single petiole furnishes more than one leaf.

2. *ARTICULATE*, *jointed*; when one leaf grows out at the top of another.

3. *DIGITATE*, *fingered*; when the apex of a single petiole connects many folioles; and they are termed *BINATE*, *TERNATE*, or

* For the figures of these leaves, *vide* our Plates at the end of this work.

QUINATE, growing *two, three, or five* together, according to the number of folioles, of which the digitate leaf consists.

4. PINNATE, *winged*; when the sides of a single petiole connect many folioles.

5. PINNATE WITH AN ODD ONE; when it is terminated by an odd foliole.

6. A CIRRHOSE PINNATE LEAF; when it terminates in a cirrhus or clasper.

7. AN ABRUPT PINNATE LEAF; when it is terminated neither by a foliole nor cirrhus.

8. OPPOSITELY PINNATE; when the folioles stand opposite to each other.

9. ALTERNATELY PINNATE; when the folioles are produced alternately.

10. INTERRUPTEDLY PINNATE; when the folioles are alternately less.

11. ARTICULATELY PINNATE; when the petiole common to all the folioles is articulate, jointed.

12. DECURSIVELY PINNATE; when the folioles are decurrent, running down; that is, extend themselves downwards along the petiole.

13. CONJUGATE; when the pinnate leaf consists of two folioles only.

II. DEGREE, in a compound leaf, respects the subdivision of the common petiole. In respect to which leaves are,

1. DECOMPOUND; when a petiole once divided connects many folioles.

2. BIGEMINATE; when a dichotomus* petiole connects four folioles on its apices.

* Forked or halved, and each division forked again: EDITOR.

3. BITERNATE, or DUPLICATO-TERNATE; when there are three folioles on a petiole, and each foliole is ternate, as in EPIMEDIUM.

4. BIPINNATE, or DUPLICATO-PINNATE; when the folioles of a pinnate leaf are pinnate.

5. PEDATE, *foot-shaped* or branching; when a bifid petiole connects many folioles on its inside only, as in PASSIFLORA and ARUM.

6. SUPRA-DECOMPOUND; when many folioles are born on a petiole, that has been any number of times subdivided.

7. TRITERNATE, or TRIPPLICATO-TERNATE; when a petiole bears three folioles that are each of them ternate.

8. TRIPINNATE, or TRIPPLICATO-PINNATE; when a petiole bears many folioles, each of which are bipinnate*.

CHAP. VII.

OF DETERMINATE LEAVES.

BY the DETERMINATION of leaves is to be understood their character, expressed from some circumstance foreign to their own particular structure or configuration; as from their *place, situation, insertion, or direction*.

I. By the PLACE of a leaf is meant the part where it is fastened to the plant. In respect to which leaves are called,

* Vide Plates of Leaves at the end of this work.

1. SEMINAL, *seed* leaves; which before were the cotyledons, and are the first which appear.

2. RADICAL, *root* leaves; such as proceed from the root.

3. CAULINE, *stem* leaves; such as grow on the stem.

4. RAMEOUS, *branch* leaves; such as grow on the branches.

5. AXILLARY*, such as are placed at the coming out of the branches.

6. FLORAL, *flower* leaves; such as are placed at the coming out of the flower.

II. By SITUATION is meant the disposition of the leaves on the stem of the plant. In respect to which, leaves are called,

1. STELLATE, *starry*; or VERTICILLATE, *whorled*; when the stalk is surrounded in whorls by more than two leaves; and these again receive the denomination of *tern*, *quatern*, *quine*, *sene*, &c. according to the number of leaves of which the star or whorl is composed, as in NERIUM,...BRABEJUM,...and HIPPURIS.

2. OPPOSITE; when the cauline leaves come out in pairs facing each other, and each pair is crossed by the next, so that they point four different ways.

3. ALTERNATE; when they come out singly, and follow in a gradual order.

4. SPARSE, *scattered*; when they come out in plenty about the plant without order.

5. CONFERT, *crowded*; when they come out in quantities, so as almost to cover the branches, and leave hardly any space between them.

6. IMBRICATE; when they are confert and erect, so as to lie over one another, each covering a part of the following one.

* From *axilla*, an *armpit*. EDITOR.

7. FASCICULATE, *bundled*; when many come out from the same point, as in LARIX.

8. DISTICH, *in two rows*; when the leaves all respect two sides of the branches only, as in ABIES and DIERVILLA.

III. In respect to their INSERTION (which is usually at the base), leaves are called,

1. PELTATE, *shield-fashioned*; when the petiole is inserted into the disk of the leaf, and not into its *base* or margin, as in NYMPHŒA, ...HERNANDRIA, ...and COLOCASIA.

2. PETIOLATE; when there is a petiole fastened to the leaf at the margin of the base.

3. SESSILE; when the leaf has no petiole, but is fastened immediately to the stem.

4. DECURRENT, *running down*; when the base of a sessile leaf extends itself downwards along the stem beyond the proper base or termination of the leaf, as in VERBESINA, ...CARDUUS, ...and SPHÆRANTHUS.

5. AMPLEXICAUL, *embracing the stalk*; when the base of the leaf embraces the sides of the stem crosswise on both sides; or SEMIAMPLEXICAUL, *half embracing the stalk*; which only differs from *Amplexicaul*, in that it is in a less degree.

6. PERFOLIATE; when the base of the leaf is continued across the stem till it meets again, so as to embrace it all around, as in BUPLEURUM.

7. CONNATE, *growing together*; when two opposite leaves join, and are united in one, as in LONICERA and EUPATORIUM.

8. VAGINANT, forming a *vagina* or sheath; when the base of the leaf forms a cylindric tube that invests the branch.

IV. In respect to their DIRECTION, leaves are called,

1. ADVERSE; when their sides are not turned towards heaven but towards the south, as in AMOMUM.
2. OBLIQUE; when the base of the leaf looks towards heaven, and the apex or tip towards the horizon, as in PROTEA and FRITILLARIA.
3. INFLEX, *bending inwards*; when the leaf is bowed upwards towards the stem.
4. ADPREST; when the disk of the leaf lies close to the stem.
5. ERECT, *upright*; when the angle they form with the stem is extremely small.
6. PATENT, *spreading*; when they make an acute angle with the stem.
7. HORIZONTAL; when they stand at right angles with the stem.
8. RECLINED, or, as some term it, REFLEX; when they are bowed downwards, so that the apex or tip is lower than the base.
9. REVOLUTE, *rolled back*; when they are rolled downwards.
10. DEPENDENT, *hanging down*; when they point directly to the ground.
11. RADICANT, *rooting*; when the leaves strike root.
12. NATANT, *floating*; when they lie on the surface of the water, as in NYMPHŒA and POTAMOGITON.
13. DEMERSE, *sunk*; when they are hid beneath the surface of the water*.

* Vide Plate 9, at the end of this work.

CHAP. VIII.

OF THE FULCRA OF PLANTS.

FULCRUM, a *prop*, is a term used to express those small parts of plants, of which the chief use is to strengthen and support them.

FULCRA are of seven kinds, viz. STIPULA, ... BRACTEA ... SPINA, ... ACULEUS, ... CIRRHUS, ... GLANDULA, ... and PILUS; all which we shall explain in their order.

1. STIPULA, is a scale or small leaf, stationed on each side the base of the petioles or peduncles, when they are first appearing, as in *papilionaceous* flowers; and also in TAMARINDUS, ... CASSIA, ... ROSA, ... MELIANTHUS, ... LIRIODENDRON, ... ARMENIACA, ... PERSICA, ... PADUS, and others.

2. BRACTEA, a *floral leaf*, is so called, when it differs in shape and colour from the rest, as in TILIA, ... FUMARIA BULBOSA, ... STÆCHAS, ... and HORMINUM.

3. SPINA, a *thorn*; is a kind of sharp weapon or armature, protruded from the wood of the plant, as in PRUNUS, ... RHAMNUS, ... HIPPOPHAE, ... CELASTRUS, ... and LYCIUM: it will often disappear by culture, as in PYRUS.

4. ACULEUS, a *prickle*, is the same sort of armature, proceeding from the cortex of the plant only, as in ROSA, ... RUBUS, ... RIBES, ... and BERBERIS.

5. CIRRHUS, a *clasper* or *tendrill*, is a filiform spiral band, by which a plant fastens itself to any other body, as in VITIS, ... BANISTERIA, ... CARDIOSPERMUM, ... PISUM, ... and BIGNONIA.

6. GLANDULA, a *little gland*; is a kind of pap or teat, serving for the excretion of some humour: its situation is commonly on the petioles, the serratures of the leaves, or the tender stipulæ.

7. PILUS, a *hair*, is a sort of bristle, serving as an excretory duct to the plants.

CHAP. IX.

OF THE HYBERNACULA OF PLANTS.

THE HYBERNACULUM, *winter-lodge*, is that part of a plant which encloses and protects the embryo, or future shoot, from external injuries: it is of two kinds, viz. BULBUS, a *bulb*; and GEMMA, a *bud*.

1. A BULB, is an *hybernacle*, placed on the descending caudex: it is of various kinds, viz. a *squamose* bulb, when it consists of *imbricate lamellæ**, as in LILIUM;...a *solid* bulb, when it consists of a *solid* substance, as in TULIPA;...a *tunicate* bulb, when it consists of many *tunics* or coats, as in CÉPA;...and an *articulate* or jointed bulb, when it consists of *lamellæ* that are linked together, as in LATHRÆA,...MARTINIA,...and ADOXA.

2. GEMMA, a *bud*, is an *hybernacle* placed on the ascending caudex: it consists either of *stipulæ*, of *petioles*, of the *rudiments* of leaves, or of *cortical squamæ*†.

BUDS are of various kinds. In the generality of plants they are *floriferous*; that is, producing both leaves and flowers;

* Thin plates or scales. EDITOR.

† Scales of the bark. EDITOR.

but in *ALNUS* they bear leaves only;...in *POPULUS*, *FRAXINUS*, and some species of *SALIX*, they bear *leaves* and *flowers* distinctly;...in *CORYLUS* and *CARPINUS*, leaves and *female* flowers;...in *PINUS* and *ABIES*, leaves and *male* flowers;...and in *DAPHNE*, *ULMUS*, *CORNUS*, and *AMYGDALUS*, leaves and *bisexual* flowers:...in *DENTARIA*, *ORNITHOGALUM*, *LILIUM*, and *SAXIFRAGA*, the buds are *de-idiuous*.

In several plants there are *no* buds, as in *PHILADELPHUS*,...*FRANGULA*,...*ALATERNUS*,...*PALIURUS*,...*JATROPHA*,...*HIBISCUS*,...*BAHOBAB*,...*JUSTICIA*,...*CASSIA*,...*MIMOSA*,...*GLEDITSIA*,...*ERYTHRINA*,...*ANAGYRIS*,...*MEDICAGO*,...*NERIUM*,...*VIBURNUM*,...*RHUS*,...*TAMARIX*,...*HEDERA*,...*ERICA*,...*MALPIGHIA*,...*LAVATERA*,...*SOLANUM*,...*ASCLEPIAS*,...*RUTA*,...*GERANIUM*,...*PETIVERIA*,...*PERESKIA*,...*CUPRESSUS*,...*THUYA*,...and *SABINA*.

In *cold* countries there are but few plants without buds, and in *hot* countries but few that have any.

CHAP. X.

OF THE HABIT OF PLANTS.

BY the *HABIT*, or *external face* of plants, is to be understood a certain conformity between vegetables that belong to the same genus, or are *near of kin* to each other*. This conformity may

* This definition of the habit of plants, which we have taken from the *Philosophia Botanica*, seems to agree better with the old state of botany, when plants were actually ranged according to their external face, than with the modern system that ranges them by the fructification: for plants that, by the system, are neither of the

be in respect to various circumstances, as *placentation*, *radication*, *ramification*, *intorsion*, *gemmaion*, *foliation*, *stipulation*, *pubescence*, *glandulation*, *lutescence*, *inflorescence*, &c. As each of the terms here enumerated will furnish us with a separate chapter, we shall forbear the explanation of them here.

CHAP. XI.

OF PLACENTATION.

BY PLACENTATION* is meant the disposition of the *cotyledons* at the time when the seed is beginning to grow. Plants, in respect to *placentation*, are termed,

I. ACOTYLEDONES, *without cotyledons*, when this part is wanting, as in *MOSSSES*.

same genus, nor have any systematic affinity, will often have a great conformity in their habit; whilst those of the same genus will have their habits distinct. The habits of plants was the invention of the earlier botanists, who knew no better rule for the distribution of vegetables: and, indeed, *Linnaeus* himself is induced to admit, that it is often a good guide; and that *Casper Bauhine*, and others, had in many cases discovered the affinity of plants by the habit, when systematists had failed in attempting the same by their artificial rules; nor does he think even the fructification, which is the invention of the moderns, sufficient for detecting all the classes of vegetables, though he considers it as the primary guide to the natural method so much sought after by those who have cultivated this science.

AUTHOR.

* The *cotyledons* of the seed in vegetables answer the purpose of the *placenta* in the animal œconomy; and hence the disposition of the *cotyledons* is called *placentation*. AUTHOR.

II. MONOCOTYLEDONES, with a *single cotyledon**; and these are either,

1. PERFORATE, as in GRASSES.
2. UNILATERAL, as in PALMS; or,
3. REDUCED, as in CEPA.

III. DICOTYLEDONES, having *two cotyledons*; and these are either,

1. IMMUTATE, *unchanged*, as in the class *Didynamia*; and in plants whose pericarpium is a *legumen*, *pomum*, or *drupa*†.

2. PPLICATE, *folded*, as in GOSSYPIUM.

3. DUPLICATE, *doubled*, as in MALVA; and in the class *Tetradynamia*.

4. OBOVATE, *rolled up*, as in HELXINE.

5. SPIRAL, *turning like a screw*, as in SALSOLA, ...SALICORNIA, ... CERATOCARPUS, ...BASELLA, ...and all *oleraceous* plants‡; or,

6. REDUCED, as in *umbellate* plants.

IV. POLYCOTYLEDONES, with *many cotyledons*, as in PINUS, ... CUPRESSUS, and LINUM.

* *Linnaeus* observes, that the *Monocotyledones* are properly *Acotyledones*; the cotyledons remaining within the seed. AUTHOR.

† See these terms explained in Part I. Chap. VI. EDITOR.

‡ Pot herbs. The *oleraceous* plants make an order in the *Fragmenta Methodi Naturalis* of *Linnaeus*; consisting of SPINACIA—BLITUM—BETA—GALENIA—ATRIPLEX—CHENOPODIUM—RIVINIA—PETIVERIA—HERNIARIA—ILLECEBRUM—POLYCNEMUM—AXYRIS—ACHYRANTHES—AMARANTHUS—GOMPHRENA—CELOSIA—CERATOCARPUS—CORISPERNUM—CALLITRICHE—SALSOLA—SALICORNIA, and ANABASIS. AUTHOR.

CHAP. XII.

OF RADICATION.

BY RADICATION is meant the disposition of the *root* of the plant, which is to be considered in respect to the ascending caudex and the radicles, as has been shown in Chap. II. where the principal characters of roots have been explained. Roots are farther distinguished into,

I. BULBOSE, consisting of a *bulb*; and these are either,

1. SQUAMOSE, *scaly*, as in LILIUM.

2. TUNICATE, *coated*, as in CEPA.

3. DUPLICATE, *double*, as in FERTILLARIA; or,

4. SOLID, as in TULIPA.

II. TUBEROSE, *knobbed*; and these are either,

1. PALMATE, *handed*, as in ORCHIS.

2. FASCICULATE, *bundled*, as in PÆONIA; or

3. PENDULOUS, *hanging*, as in FILIPENDULA and ELÆAGNUS.

III. ARTICULATE, *jointed*, as in LATHRÆA,...OXALIS,...MARTY-
NIA,...and DENTARIA.

IV. FUSIFORM, *spindle-shaped*, as in PASTINACA,...DAUCUS,...
and RAPHANUS.

V. GLOBOSE, *globe-shaped*, as in BUNIUM, and in some species
of RANUNCULUS and CHEROPHYLLUM.

CHAP. XIII.

OF RAMIFICATION.

RAMIFICATION is the manner in which a tree produces its *branches*, with the situation of which that of the *leaves* is also connected*.

Some plants have no *branches*, though they have *leaves* which are placed on the stem. This is the case with DICTAMNUS,... PEONIA,... EPIMEDIUM,... and PODOPHYLLUM.

Leaves *opposite* or *alternate* are generally a mark of great difference in plants: a few genera, however, must be excepted, which have some species with opposite leaves, and others with alternate, as in EUPHORBIA,... CISTUS,... LANTANA,... ANTIRRHINUM, ... LILIUM,... and EPILOBIUM.

In ANTIRRHINUM,... JASMINUM,... VERONICA,... and BORAGO,... the lower leaves at the branches are *opposite*, and the upper ones at the flowers *alternate*.

In POTENTILLA SUPINA, and in POTAMOGITON, the lower leaves are *alternate*, and the upper ones on the branches *opposite*.

In NERIUM the lower leaves are *opposite*, and the upper ones *ternate*.

In RUSCUS the lower leaves are *ternate* and the upper ones *alternate*.

* The doctrines delivered here under the head of *Ramification* do not answer to the title, the greater part respecting rather the situation of the *leaves* than that of the *branches*: they might, with more propriety, have been collected under a head of *foliation*; but as the term *foliation* is used to express the habit of plants, in respect to the position of leaves in the bud, before they disclose themselves, as will be shown in Chap. XVI. these doctrines could not have stood under the same head, without a confusion in the use of the term; and this seems to be the reason why *Linnaeus*, whom we follow, has given them in this place. AUTHOR.

In *COREOPSIS ALTERNIFOLIA*, and in *ANTIRRHINUM CHALEPENSE*, the lower leaves are *quatern*, and the upper ones *alternate*.

The *natural* situation of the leaves in plants that are much branched is best concluded from the *radical* leaves.

CHAP. XIV.

OF INTORSION.

INTORSION, *winding*, is the flexion or bending of any part of a plant towards one side.

CAULES VOLUBILES, *twining stems*, wind either,

1. SINISTRORSUM, to the *left*, as in TAMUS, ... DIOSCOREA, ... RAJANIA, ... MENISPERMUM, ... CISSAMPELOS, ... HIPPOCRATEA, ... LONICERA, ... HUMULUS, ... and HELXINE; or,

2. DEXTRORSUM, to the *right*, as in PHASEOLUS, ... DOLICHOS, ... CLITORIA, ... GLYCINE, ... SECURIDACA, ... CONVULVULUS, ... IPOMŒA, ... CYNANCHE, ... PERIPLOCA, ... CEROPEGIA, ... EUPHORBIA, ... TRAGIA, ... BASELLA, ... EUPATORIUM, ... and TOURNEFORTIA.

CIRRI VOLUBILES, *twining clasps*, wind to the *right*, and *back* again. Most *leguminous* plants have cirri of this kind: in SMILAX, and in most species of PIPER, the *petioles* are cirrhiferous.

COROLLÆ bend to the *left**, in ASCLEPIAS, ... NERIUM, ... VINCA,

* Supposing yourself placed in the centre, and looking towards the south.

...RAUWOLFIA,...PERIPLOCA,...and STAPELIA ;...and to the *right* in PEDICULARIS.

In TRIENTALIS there is this singularity, that the petals are all *imbricate*, one side of each folding over the next towards the *right*.

In GENTIANA, the imbrication of the petals before they are unfolded is contrary to the sun.

The PISTILLA incline to the *left* in CUCUBALUS and SILENE.

The GERMINA are twisted to the *left* in HELICTERES and ULMARIA.

FLOWERS, in respect to *intorsion*, have,

A *resupination**, which is, when the upper lip of the corolla look towards the ground, and the upper lip towards heaven, as in the EUROPEAN VIOLE, ...AJUGA ORIENTALIS, ...OCYUM, ...and some species of SATYRIUM ; or

An *obliquity*, as in the species of HYSSOPUS, called LOPANTHUS, ...NEPETA SIBIRICA, ...and some species of PEDICULARIS.

SPICÆ, *spikes*, are,

Spiral, as in CLAYTONIA, and in some *asperifolious*† plants ; or, *incurvate, crooked*, as in SAURURUS, ...MIMOSA, ...PETIVERIA, ...PAPAYER, ...SEDUM RUBRUM, ...and LILIUM MARTAGON.

In several plants there is found a *contorsion* of the fibres, which answers the end of an *hygrometer*‡. Thus in AVENA, there is an arista or beard, that is twisted like a rope : in some GERANIUMS, the arillus of the seed has a spiral tail ; and in MNIUM, the peduncles are twisted contrary ways above and below.

* *Resupination* is, when any thing is thrown on its back, or lies face upwards.

EDITOR.

† The *asperifoliæ* belong to the class *Pentandria*. See Part II. Chap. VIII.

EDITOR.

‡ An instrument for measuring the degree of dryness or moisture of the air. The fibres of the plant, here instanced being affected by the quality of the air, the spiral part twists or untwists, as the weather varies ; and by observing this, the temperature of the air may be discovered. EDITOR.

CHAP. XV.

OF GEMMATION.

GEMMATION is the construction of the *gem* or *bud*, which is formed either of *leaves*, *stipulæ*, *petioles*, or *squamæ*. Those that are formed of the leaves will be considered in the next Chapter, under the head of *foliation*; the rest are distinguishable into,

PETIOLAR *buds*, which are either,

1. *Opposite*, as in LIGUSTRUM, ... PHILLYREA, ... NYCTANTHES, ... SYRINGA, ... HYPERICUM, ... CORIARIA, ... BUXUS, ... JASMINUM, ... VACCINIUM, ... ARBUTUS, ... ANDROMEDA, ... LEDUM, ... DAPHNE, ... LAURUS, ... MYRICA, ... LINNÆA, ... DIERVILLA, ... LONICERA, ... EUONYMUS, ... FRAXINUS, ... ACER, ... ESCULUS, ... BIGNONIA, ... OPULUS, ... SAMBUCUS, ... and PSIDIUM; or,

2. *Alternate*, as in SALIX, ... SPIRÆA, ... GENISTA, ... SOLANUM, ... HIPPOPHAE, ... BERBERIS, ... ILEX, ... RIBES, ... JUGLANS, ... PISTACHIA, ... and PLUMBAGO.

STIPULACEOUS *buds*, which are either,

1. *Opposite*, as in CEPHALANTHUS and RHAMNUS CATHARTICUS; or,

2. *Alternate*, as in POPULUS, ... TILIA, ... ULMUS, ... QUERCUS, ... FAGUS, ... CARPINUS, ... CORYLUS, ... BETULA, ... ALNUS, ... FICUS, ... and MORUS.

STIPULACEO-PETIOLAR *buds*, which are,

1. *Alternate*, as in SORBUS, ... CRATÆGUS, ... PRUNUS, ... MESPILUS, ... PYRUS, ... MALUS, ... COTONEASTER, ... AMYGDALUS, ... CERASUS, ... PADUS, ... MELIANTHUS, ... ROSA, ... RUBUS, ... VITIS, ... ROBINIA, ... CYTISUS, ... POTENTILLA FRUTICOSA, ... and STAPHYLEA.

2. *Anomalous, or irregular buds*, as in ABIES,... PINUS,...and TAXUS.

In many plants the buds are *wanting*, as has been shown in Chap. IX.

CHAP. XVI.

OF FOLIATION.

BY FOLIATION is to be understood the *complicate*, or folded state the leaves are in, whilst they remain concealed within the *buds* of the plant*. Leaves, in respect to the manner of their *complication*, are either,

1. INVOLUTE *rolled in*; when their lateral margins are rolled spirally inwards on both sides, as in LONICERA,...DIERVILLA,... EUONYMUS,... RHAMNUS CATHARTICUS,...PYRUS,... MALUS,... POPULUS,... PLUMBAGO,... VIOLA,... COMMELINA ANNUA,... PLANTAGO, ...ALISMA,... POTAMOGITON NATANS,... NYMPHŒA,... SAURURUS,... ASTER ANNUUS,... HUMULUS,... URTICA,... HEPATICA,... SAMBUCUS EBULUS,...and STAPHYLEA.

2. REVOLUTE, *rolled back*; when their lateral margins are rolled spirally backward on both sides, as in ROSMARINUS,... TEUCRIUM MARUM,... DRACOCEPHALON,... DIGITALIS,... NERIUM,... ANDROMEDA,... LEDUM,... EPILOBIUM ANGUSTUM,... RUMEX,... PERSICARIA,... POLYGONUM,... PARIETARIA,... PRIMULA,... CARDUUS,... CNICUS,... TUSSILAGO,... SENECIO,... OTHONNA,... POTENTILLA FRUTICOSA,... PTELEA,...and some species of SALIX.

* *Linnaeus* claims the invention of the distinctions given in this Chapter, preceding botanists not having (as he says) attended to the *foliation in buds*. AUTHOR.

3. *OBVOLUTE*, *rolled against* each other; when their respective margins alternately embrace the straight margin of the opposite leaf, as in *DIANTHUS*,...*LYCHNIS*,...*SAPONARIA*,...*EPILOBIUM OPPOSITIFOL.*,...*DIPSACUS*,...*SCABIOSA*,...*VALERIANA*,... *MARRUBIUM*,...*PHLOMIS*,...*SALVIA*,...and *PRASIMUM*.

4. *CONVOLUTE*, *rolled together*; when the margin of one side surrounds the other margin of the same leaf, in the manner of a cowl or hood, as in *CANNA*,...*AMOMUM*,...*CALLA*,...*ARUM*,...*PIPER*,...*HYDROCHARIS*, ... *COMMELINA LUTEA*, ...*PRUNUS ARMENIACA*,... *DODECATHEON*,...*CREPIS*,...*LACTUCA*,...*HIERACIUM*,...*SONCHUS SIBIR.* ...*TRAGOPOGON*,... *OROBUS*,...*VICIA*,... *LATHYRUS*,... *SOLIDAGO*,... *ASTER*, ... *PINGUICULA*, ... *VACCINIUM*,... *PYROLA*,... *BERBERIS*,... *BRASSICA*,...*ARMORACIA*,...*SYMPHYTUM*,...*CYNOGLOSSUM*,...*POTAMOGITON PERFOL.*,...*ERYNGIUM*,... *MENYANTHES*,... *SAXIFRAGA*,... *ARALIA*,...*DICTAMNUS*,...*EPIMEDIUM*,...and many *GRASSES*.

5. *IMBRICATE*; when they are *parallel*, with a straight surface, and lie one over the other, as in *SYRINGA*,...*LIGUSTRUM*,...*PHILLYREA*,...*NYCTANTHES*,...*LINNÆA*,...*CEPHALANTHUS*,...*CORIARIA*,...*HYPERICUM*,...*VALANTIA*,... *JUSTICIA*,... *PORTULACA*,... *LAURUS*,... *DAPHNE*,...*HIPPOPHAE*,...*RUSCUS*,...*CYANUS PERENNIS*,...*MESPILUS GERM.*,...*CAMPANULA*,...*POLEMONIUM*,...and *Sium*.

6. *EQUITANT*, *riding*; when the sides of the leaves lie parallel, and approach in such manner, as the outer embrace the inner (which is not the case with the *conduplicate* explained in the next head), as in *HEMEROCALLIS*,... *IRIS*,... *ACORUS*,...*CAREX*,...*POA*,...and some *grasses*.

7. *CONDUPLICÆ*, *doubled together*; when the sides of the leaf are parallel, and approach each other, as in *QUERCUS*,...*FAGUS*, ...*CORYLUS*,...*CARPINUS*,...*TILIA*,...*PADUS*,...*CERASUS*,...*AMYGDALUS*,...*COTONEASTER*,...*FRANGULA*,...*ALATERNUS*,...*PALIURUS*,...*JUGLANS*,...*PISTACIA*,...*RHUS*,...*FRAXINUS*,...*SORBUS*,...*RUBUS*,...*POTENTILLA VULG.*,...*COMARUM*,...*BIGNONIA*,... *CYTISUS*,... *ROBINIA*,

...PISUM,...MELIANTHUS,...PASTINACA,...HERACLEUM,...LASERPITUM,...POTERIUM,...and most *diadelphous* plants.

PPLICATE, *plaited*; when their complication is in plaits lengthways, like the plicate leaves explained in Chap. V. as in CRATÆGUS, ... BETULA, ... ALNUS, ... FAGUS, ... VITIS, ... ACER, ... OPULUS VIBURNUM, ... RIBES, ... ALTHÆA, ... MALVA, ... HUMULUS, ... URTICA, ... PASSIFLORA, ...and ALCHEMILA.

8. RECLINATE, *reclined*; when the leaves are reflexed downwards towards the petiole, as in PODOPHYLLUM, ... ACONITUM, ... HEPATICA, ... PULSATILLA, ... ANEMONE, ...and ADOXA.

9. CIRCINAL, *compassed**; when the leaves are rolled in spirally downwards, as in FILICES, and some PALMS†.

CHAP. XVII.

OF STIPULATION.

BY STIPULATION is meant the situation and structure of the *stipula*‡, at the base of the leaves.

The *stipula* in different plants are found to be as various as the leaves. They are,

1. WANTING in the *asperifolia*§, the class *Didynamia*, the

* In rings.

† Vide Plate 11, at the end.

‡ See Chap. VIII. AUTHOR.

§ *Pentandria Monogynia*, Distinction 1. See Part II. Chap. VIII. AUTHOR.

*tellate**, *siliquosæ*†, *liliciæ* ‡, *orchideæ*§, and in most compound flowers.

2. PRESENT in the *Papilionaciæ* ||, *Lomentaceæ*¶, and in the class *Icosandria*.

3. GEMINÆ, two together, or with a single one on each side in most plants.

4. SOLITARY, in *MELIANTHUS*, in which the stipula is on the inside; and *RUSCUS*, in which it is on the outside.

5. DECIDUOUS, in *PADUS*,...*CERASUS*,...*AMYGDALUS*; and also** in *POPULUS*,...*TILIA*,...*ULMUS*,...*QUERCUS*,...*FAGUS*,...*CARPINUS*,...*CORYLUS*,...*BETULA*,...*ALNUS*,...*FICUS*,...and *MORUS*.

6. PERSISTING, in the class *Diadelphia*, and in *Icosandria*, *Polygynia*.

7. ADNATE, growing close to the plant, in *ROSA*,...*RUBUS*,...*POTENTILLA*,...*COMARUM*,...and *MELIANTHUS*.

8. SOLUTE, free or loose, in most plants.

9. INTRAFOLIACEOUS, on the inside of the leaves, in *FICUS* and *MORUS*.

10. EXTRAFOLIACEOUS, on the outside of the leaves, in *ALNUS*,...*BETULA*,...*TILIA*,...and the class *Diadelphia*.

* *Tetrandria Monogynia*, Distinction 2. See Part II. Chap. VII. AUTHOR.

† *Tetradynamia Siliquosa*. See Part II. Chap. XVIII. AUTHOR.

‡ *LILIUM*—*FRITILLARIA*—*TULIPA*—and *ERYTHRONIUM* are the *lillicæ* plants; which make an order in the *Methodi Naturalis Fragmenta*. See *Phil. Bot.* page 28. AUTHOR.

§ *ORCHIS*—*SATYRIUM*—*SERAPIS*—*HERMINIUM*—*NEOTTIA*—*OPHRYE*—*CYPRIPEDIUM*—*EPIDENDRUM*—*LIMODORUM*—and *ARETHUSA*, are the *Orchideæ*; which are another order in the *Method Nat. Frag.* See *Phil. Bot.* p. 27. AUTHOR.

|| Class *Diadelphia*. See Part II. Chap. XX. AUTHOR.

¶ *SOPHORA*—*CERCIS*—*BAUHINIA*—*PARKINSONIA*—*CASSIA*—*POINCIANA*—*TAMARINDUS*—*GUILANDINA*—*ADENANTHERA*—*HÆMATOXYLON*—*CÆSALPINIA*—and *MIMOSA*. These are an order in *Meth. Nat. Frag.* See *Phil. Bot.* p. 34. They are called *lomentaceous* from *Lomentum*, which signifies *Bear Meal*. AUTHOR.

** The genera here instanced are the same with those enumerated in the 15th Chapter, as having stipulaceous buds that are alternate, which are those referred to by *Linnaeus* in this place. AUTHOR.

CHAP. XVIII.

OF PUBESCENCE.

PUBESCENCE, *downiness**, is an armature, by which plants are defended from external injuries. *Pubescence* is of the following kinds, viz.

SCABRITIES, *roughness*; which is composed of particles scarce visible to the naked eye†, that are scattered over the surface of the plant. This is distinguishable into,

I. *Scabrities* GLANDULOSA, a *glandulose roughness*; when it consists of little glands, which are either,

1. MILIARY, like grains of *millet*.
2. VESICULAR, composed of *bladders*.
3. LENTICULAR, resembling *lentils*.
4. GLOBULAR, *globe-shaped*; as in ATRIPLEX,...and CHENOPODIUM.
5. SECRETORY, serving for *secretion*.
6. CATENULATE, consisting of little *chains*; or,
7. UTRICULAR, like little *bottles*.

II. *Scabrities* SETACEA, a *bristly roughness*; when it consists of *bristles*, which are either,

1. CYLINDRIC, like a *cylinder*.

* The term *downiness* is not to be taken here in too strict a sense, as the following explanations show. AUTHOR.

† *Guettard* was the first who carefully examined this kind of pubescence.

2. CONIC, like a *cone*.
3. HAMOSE, *hooked*.
4. GLANDULIFEROUS, *bearing glands*.
5. FURCATE, *forked*.
6. SECURIFORM, *hatchet-shaped*, as in HUMULUS.
7. AGGREGATE and *starry*, as in ALYSSUM...and HELICTERES; or
8. AGGREGATE and *simple*, as in HIPPOPHAE.

III. *Scabrities* ARTICULATA, a *jointed roughness*; when it is in *joints*, which are either,

1. SIMPLICES, *simple*.
2. NODOSE, *knotty*.
3. CAUDATE, *tailed*.
4. RAMOSE, *branching*, as in VERBASCUM; or,
5. PLUMOSE, *feathery*.

IV. LANA, *wool*, is a protection to many plants against the scorching heat, as in SIDERITIS CANARIENSIS, ... SALVIA CANARIENSIS, ... the SALVIA called ÆTHIOPIS, ... MARRUBIUM, ... VERBASCUM, ... STACHYS, ... the CARDUUS called ERIOCEPHALUS*, ... and ONOPORDUM.

V. TOMENTUM, *down*, is a defence for plants against winds; it has commonly a whitish, or hoary appearance, as in TOMEX, ... MEDICAGO, ... and HALIMUS.

VI. STRIGÆ†, with their stiff bristles, are of use to prevent

* There is a genus entitled ERIOCEPHALUS, but the plant here meant is the CARDUUS ERIOPHORUS of *Lin. Species Plant.* page 823, which is the CARDUUS CAPITĒ ROTUNDO TOMENTOSO of *Casp. Bauhine*. It was formerly called CORONA FRATRUM. AUTHOR.

† *Linæus* has omitted the definition of this term. It signifies properly a row, or ordinate disposition of things of any sort; and appears, by the instances here

plants from being bruised or destroyed by vermin, as in CACTUS,...MALPIGHIA,...HIBISCUS,...and RUBUS.

VII. HAMI, *hooks*, fasten themselves to animals as they pass by; these are either,

1. TRIGLOCHID, *three-pointed*, as in LAPPULA; or,

2. INCURVATE, *crooked*, as in ARCTIUM,...MARRUBIUM,...XANTHIUM,...and PETIVERIA.

VIII. STIMULI, *stings*, keep off naked animals by their venomous punctures, as in URTICA,... JATROPHA, ...ACALYPHIA, ... and TRAGIA.

IX. ACULEI, *prickles*, keep off particular animals, as in VOLKAMERA, ...PISONIA,...CÆSALPINIA,... MIMOSA,... PARKINSONIA, ... CAPPARIS,...ERYTHRYNA,...ROBINIA,...SOLANUM,...CLEOME,...SMILAX,...CONVOLVULUS,...ARALIA,...DURANTA,...XYLON,...DRYPIS,... EUPHORBIA,... TRAGACANTHA,... and TRAGOPOGON. In HUGONIA the ACULEI are *spiral* or *cirrhose**.

X. FURCÆ, *forks*, are a defence against animals in general, as in BERBERIS, ... RIBES, ... GLEDITSIA, ... MESEMBRYANTHEMUM,... OSTEOSPERMUM,...BALLOTA,... BARLERIA,... FAGONIA,...and POTE RIUM.

XI. SPINÆ, *thorns*, serve to keep off cattle: these are either,

On the *branches*, as in PYRUS,...PRUNUS,...CITRUS,...HIPPOPHÆE,...GMELINA,... RHAMNUS,... LYCIUM,...CATESBEA,... CELASTRUS,... ULEX,...ASPARAGUS,... SPARTIUM,...ACHYRONIA,... XIMENIA,...ONONIS,...STACHYS,...ALYSSUM, and CICHORIUM.

On the *leaves*, as in ALOE,...AGAVE,...YUCCA,... ILEX,... HIPPOMANE,...THEOPHRASTA,...CARLINA,... CYNARA,... ONOPORDUM,...

given, to be applied to thorns or prickles that come out in rows, or in some regular order. No *English* word occurs that is exactly expressive of the term in this sense. AUTHOR.

* From CIRRHUS, a *clasper* or *tendril*. EDITOR.

MORINA, ... ACANTHUS, ... GUNDELIA, ... JUNIPERUS, ... SALSOLA, ... POLYGALA, ... RUSCUS, ... BORBONIA, ... STATICE, ... OVIEDA, ... and CLIFFORTIA.

On the *calyx*, as in CARDUUS, ... CNICUS, ... CENTAUREA, ... MOLUCELLA, ... and GALEOPSIS; or,

On the *fruit*, as in TRAPA, ... TRIBULUS, ... MUREX, ... SPINACHIA, ... AGRIMONIA, ... and DATURA.

CHAP. XIX.

OF GLANDULATION.

GLANDULATION respects the secretory vessels; which are either GLANDULES, ... FOLLICLES, ... or UTRICLES.

I. GLANDULES* are either,

1. *Petiolar*, when they are on the *petioles*, as in RICINUS, ... JATROPHA, ... PASSIFLORA, ... CASSIA, ... and MIMOSA.

2. *Foliaceous*, when they are produced from the *leaves*: and these are either from the *serratures*, as in SALIX; ... from the *base*, as in AMYGDALUS, ... CUCURBITA, ... ELEOCARPUS, ... IMPATIENS, ... PADUS, ... and OPULUS; ... from the *back*, as in URENA, ... TAMARIX, ... and CROTON; ... or from the *surface*, as in PINGUICULA, ... and DROSERA.

3. *Stipular*, when they are produced from the *stipulæ*, as in BAUHINIA, ... and ARMENIACA.

* See Chap. VIII.

4. *Capillary*, like *hairs*, as in RIBES,...ANTIRRHINUM QUADRI-FOLIUM,...SCROPHULARIA,...CERASTIUM,...and SILENE; or,

5. *Pores* only, as in TAMARIX,...and SILENE VISCARIA.

II. FOLLICLES*, are vessels distended with air, as in UTRICULARIA, at the root of which there are roundish vessels that are inflatè, and have two horns;...and in ALDROVANDA also, at the leaves of which there are pot-shaped follicles that are semicircular.

III. UTRICLES†, are vessels filled with a secreted liquor. Thus in NEPENTHES, the extremity of the leaves terminate in a thread, and this thread terminates in a cylinder, the top of which is closed with a lid that opens on the edge;...in SARRACENA also, the leaves are hooded almost like those of NEPENTHES, but sessile at the root;...and in MARGRAVIA, from the centre of the umbel there are vessels produced, which resemble the ringent corolla of the GALEOPSIS, but without the under lip.

CHAP. XX.

OF LACTESCENCE.

LACTESCENCE, *milkiness*, is when a copious juice flows out on any injury done to the plant. The colour of the liquor is either,

1. WHITE, as in EUPHORBIA,... PAPAVER,... ASCLEPIAS,... APO-

* The word signifies a little bladder filled with wind. EDITOR.

† The word signifies a bottle. EDITOR.

CYNUM, ... CYNANCHUM, ... CAMPANULA, ... LOBELIA, ... JASIONE, ...
ACER, ... SELINUM, ... RHUS, ... CACTUS MAMILLARIS, ... and the *semi-*
flosculose flowers of *Tournefort**.

2. YELLOW, as in CHELIDONIUM, ... BOCCONIA, ... SANGUINARIA, ...
CAMBOGIA; or,

3. RED, as in RUMEX SANGUINEA.

CHAP. XXI.

OF INFLORESCENCE.

INFLORESCENCE, is the manner in which the flowers are fastened to the plant by the *peduncle*. Plants, in respect to *Inflorescence*, are distinguished into,

1. VERTICILLATE, with the flowers in *whorls*, as in MARRUBIUM.
2. CORYMBIFEROUS, bearing the flowers in *corymbi*, as in *siliquose* plants†.
3. SPICATE, with the flowers in *spikes*, as in PHYTOLACCA, ... ARUM, ... PHŒNIX, ... PIPER, &c.
4. PANICULATE, with the flowers in *panicles*, as in sundry of the *grasses*.

* SONCHUS—LACTUCA, &c. These make one of the classes of *Tournefort's* Inst. R. H. AUTHOR.

† MYAGRUM—ANASTATICA, &c. The *siliquose* plants make an order in the *Meth. Nat. Frag.* See *Phil. Bot.* page 34, where the plants here meant are enumerated. AUTHOR.

5. AXILLARY flowers are such as come out from the *wings* of the leaves or branches, which is the most common case.

6. OPPOSITIFOLIUS, such as come out *opposite* to the leaves, as in PIPER,... SAURURUS,... PHYTOLACCA,... DULCAMARA, ... VITIS,... CISSUS,... CORCHORUS,... GERANIUM,... RANUNCULUS AQUATILIS,... and the annual species of CISTUS.

7. INTERFOLIACEOUS, such as come out *between* the opposite leaves, but are placed alternately, as in ASCLEPIAS.

8. LATERIFOLIUS, such as come out at the *side* of the base of the leaf, as in CLAYTONIA,... SOLANUM,... and the ASPERIFOLIÆ*.

9. PETIOLAR, when the peduncle is inserted in the *petiole*, as in HIBISCUS,... and TURNERA.

10. CIRRHIFEROUS, such as bear *cirrho*, as in CADIOSPERMUM, ... and VITIS.

11. SUPRA-AXILLARY, such as come out *above* the wings, as in the ASPERIFOLIÆ,... and in POTENTILLA MONSPELIENSIS.

CHAP. XXII.

OF SPECIFIC DISTINCTIONS.

WE have treated of *generic* differences in the last five Chapters of the Second Part of this work, we come now to treat of the *specific* ones. For this a foundation has been lain in the

* *Pentandria Monogynia*, Distinction 1. AUTHOR.

preceding Chapters of this Third Part, by the explanation of those parts of the vegetable on which the difference of the *species* most commonly depends; but it is necessary to observe, that the fructification, which we treated of in the First Part, as preparatory to the distinctions of the classes and genera, has its influence likewise in many cases upon the *species*, as will appear in the course of this Chapter.

Generic differences we have shown to depend on the form of the fructification, and to be confined to that alone. *Specific differences* take their rise from any circumstance, wherein plants of the same genus are found to disagree; provided such circumstance is constant, and not liable to alteration by culture or other accidents. Hence *Linnaeus* asserts, the *species* to be as many as there were different forms of vegetables produced at the creation; and considers all casual differences, as varieties of the same species.

Towards the end of the last century, the desire of increasing the number of plants had so seized the botanists of that time, that *new species* were established on too slight differences, to the great detriment of the science; and the same eagerness led them also to set down as *new genera* what should have been species only. This evil was in some measure unavoidable, whilst there were no fixed principles for the regulation of the science in this respect. A remedy to it was first attempted by *Vaillant*; afterwards by *Jussieu*, *Haller*, *Royen*, *Gronovius*, and others; and lastly by *Linnaeus*, whose aphorisms have brought this work much nearer to perfection. Something indeed seems still wanting to complete these doctrines; but perhaps more is not to be expected, till this branch of natural philosophy receives farther assistance from experiment.

We shall treat in this Chapter of those circumstances by which *species* are distinguished with certainty, reserving the *varieties* for the Chapter following.

The Root often affords a real *specific difference**, and is some-

* In *FUMARIA BULEGIA*, the greater and less sorts with a hollow root, and the

times the chief distinction, as in *SCILLA*, where the species are scarce to be distinguished, but by the bulbs being *tunicate*,...*solid*,...or *squamose*;...and in *ORCHIS*, where the species are known by the roots being *fibrose*,...*round*,...or *testiculate*; but as access cannot always be conveniently had to this part of the plant, it is better to fix the *specific* distinction on some other circumstance, if the case will admit of it.

The *TRUNK* often furnishes a sure mark of distinction. Thus in *HYPERICUM**,...*CONVALLARIA*†,...and *HEDYSARUM*‡, there are many species distinguishable by the *angles* of the stem;...and in *LUPINUS*, the species are not easy to be known, except by the same part being *simple* or *compound*. In *ERIOCAULON*, the most remarkable difference is in the *CULMUS*, which is *quinquangular*,...*hexangular*,...*decangular*, &c. In *PYROLA*, some species are distinguished by a *triquetrous scapus*. In *CITRUS*, the *aurantium* is distinguished from its congeners by its *petioles*, which are winged, or increased by a membrane on each side;...and in *GOMPHRENA*, there is a species§ distinguished by its *peduncles*, which are *diphyllous*, being furnished with two opposite folioles that are placed under the head of the flowers.

The *LEAVES* exhibit most natural and also most elegant specific differences. These have been so amply treated of already, that it would be only repetition to particularise or exemplify the numerous cases that occur of such distinctions.

FULCRA are generally a good mark of distinction, and must be carefully attended to by the botanist, for the determination of the species; as we shall show by many examples, where the difference consists principally in those parts of the plant. Thus,

greater and less sorts with a root *not* hollow, appear by the whole habit of the plants to be varieties only, as will be observed in the next Chapter. AUTHOR.

* *HYPERICUM HIRSUTUM* (*Lin. Spec. Plant.* 786.) *caule tereti*—*HYPERICUM PERFORATUM* (*Lin. Spec. Plant.* 785.) *caule ancipiti*—*HYPERICUM QUADRANGULUM* (*Lin. Spec. Plant.* 785.) *caule quadrangulo*. EDITOR.

† *CONVALLARIA POLYGONATUM* (*Lin. Spec. Plant.* 315.) *caule ancipiti*—*CONVALLARIA MULTIFLORA* (*Lin. Spec. Plant.* 315.) *caule tereti*. EDITOR.

‡ *HEDYSARUM TRIQUETRUM* (*Lin. Spec. Plant.* 746.) *caule triquetro*. EDITOR

§ *GOMPHRENA GLOBOSA* (*Lin. Spec. Plant.* 224.) AUTHOR.

Aculei are remarkable in RUBUS.

Spines in PRUNUS.

Bractea in FUMARIA,...DRACOCEPHALON,...and the *Indian* species of HEDYSARUM ; to which must be added the COMA, which is a bushy head, composed of *bractea*, that are of a large size, and terminate the stem in CORONA IMPERIALIS,...LAVANDULA,...and SALVIA.

Glandules furnish the essential mark in PADUS,...URENA,...MIMOSA,...CASSIA, and many other genera, which it would be impossible to distinguish without being acquainted with this part. They are found on the *serratures*, at the *base* of the leaves, in HELIOCARPUS,...SALIX,...and AMYGDALUS ;...on the *back* of the leaves in PADUS,...URENA,...and PASSIFLORA ;...and on the *aculei* in BAUHINIA ACULEATA, where by the apex of the *aculei* a liquor is secreted. The AMYGDALUS is distinguished from PERSICA only by the glandules of the *serratures* ;...nor could the species of URENA be ever fixed without examining the glandules of the *leaves*. The CONVULVULUS with a tuberculate calyx, is so variable in the shape of its leaves, that it seems divisible into many species, yet it is kept together by the *glandules* : and there is a species of MONARDA, distinguishable from its congeners, by the glandules, that are sprinkled over the corolla.

STIPULÆ are of great consequence in many extensive genera, where the species are liable to confusion. Thus in one species of MELIANTHUS the *stipulae* are *solitary* ;...in the other they are in *pairs* ;...and the CASSIA AURICULATA is rendered distinct from all its congeners, by the shape of its *stipulae*, which are *reniform* and *barbate*.

HYBERNACLES afford likewise a certain specific difference.

That *gems* or buds often differ greatly in the same genus, is proved by RHAMNUS ; in which the various species, viz. CERVISINA,...ALATERNUS,...PALIURUS,... and FRANGULA, have all a difference in their *buds* ; and in that extensive and intricate genus, the SALIX, the species are, by the structure and foliation of the *buds*, distinguished with great certainty.

Bulbs also distinguish the species, as is proved by SCILLA, where

they afford a real, and almost the only distinction; and by their situation in the *axilla* of the leaves, they determine DENTARIUM, ...LILIUM, ...ORNITHOGALUM, ...SAXIFRAGA, ...and BISTORTA.

INFLORESCENCE affords the truest, and in most genera the most elegant distinction. Thus in *spiraea*, the flowers are in some species *duplicato-racemose*; ... in others *corymbose*; ... and in others again, *umbellate*; ... without which characters there would be no certainty of the species.

The *peduncle*, or flower-stalk, which is the foundation of the characters of *inflorescence*, varies as to the manner of its supporting the flowers; and is said to be,

1. FLACCID, *wanting firmness*; when it is so weak as to be bowed down by the weight of the flower itself.

2. CERNUUS, *nodding*; when it is incurvate at the *apex*, so that the flower inclines to one side, or towards the ground, and cannot preserve an erect posture, by reason of the strict curvature of the peduncle, as in CARPESIUM, ... BIDENS RADIATA, ... CARDUUS NUTANS, ... SCABIOSA ALPINA, ... HELIANTHUS ANNUUS, ... and CNICUS SIBIRICUS.

3. Bearing *fastigate* flowers; when the *pedicelli**, or partial foot-stalks elevate the fructification into a *fascicle*, so that they are of an equal height at the top, as if they had been shorn off horizontally, as in DIANTHUS and SILENE.

4. PATULUS, *spreading*; when it is branched out every way, so that the flowers stand remote from each other. This stands opposite to COARCTATE, *close*.

5. Bearing CONGLOMERATE flowers; when it is branched, and bears the flowers in close compact heaps, and is therefore opposed to a *diffuse pannicle*.

* In this, and some other places, the *Philosophia Botanica* has *petiole* for *pedicellus*; but the latter is the proper term for the partial foot-stalk of a flower. See Chap. IV. AUTHOR.

6. ARTICULATE, *jointed*; when it is *furnished* with a *joint*, as in OXALIS, ...SIDA, ...and HIBISCUS.

7. Coming out in *pairs*, as in CAPRARIA, and OLDENLANDIA BIFLORA.

8. TERN, or *three*, from the same axilla, as in IMPATIENS TRIFLORA.

9. FLEXUOSE, *bending* divers ways, or UNDULATE, *waved*, as in AIRA FLEXUOSA.

10. REMAINING on the plant after the fructification is fallen, as in JAMBOLIFERA, ...OCHNA, ...and JUSTICIA.

11. INCRASSATE, *thickened* towards the flower, as in COTULA, ...TRAGOPOGON, and most *cernuous* flowers.

The parts of FRUCTIFICATION often furnish most certain and constant specific differences. *Linnaeus* tells us he was once of a contrary opinion; and held, that as the flower was of short duration, and its parts commonly very minute, recourse should not be had to the fructification for specific differences, till all other ways had been tried and found ineffectual; but as the fructification contains more distinct parts than all the rest of the plant taken together, and certitude is found throughout nature to depend mostly on her minuter parts, he has since readily admitted this distinction.

In GENTIANA, the species cannot any way be distinguished, if the flower is not admitted as a specific character; but they are easily distinguished by their *corollæ*, which vary in being *campaniform*, ...*rotate*, ...*infundibuliform*, ...*quinquefid*, ...*quadrid*, ...*octofid*, &c.

In HYPERICUM, the species are distinguished by the flowers being *trigynous**, or *pentagynous*†.

In GERANIUM, the *African* species are distinguishable from their *European* congeners, by the corolla being irregular, and also by the connexion of their stamina.

* With three styles. EDITOR.

+ With five styles. EDITOR.

IN LICHEN, the fructification is distinguishable into TUBERCULUM, a *little knob*, which is a fructification, consisting of rough points collected like à heap of dust ;... SCUTELLUM, a *small buckler*, which is a concave orbiculate fructification, the margin of which is elevated on every side ;...or PELTA, a *little shield*, which is a plane fructification fastened for the most part to the margin of the leaf*.

IN MOSSES, the CAPITULUM, or *little head*, is an ANTHERA.

IN GRASSES, SPICULA, a *little spike*, is a *partial one* ; the ARISTA is *tortile, twisted*, when it has a twisted joint in the middle. ARTICULUS, a *joint*, is the part of the *culmus* that lies between two *geniculi*, or *knots*.

A *radiate* compound flower consists of *disk* and *radius*. The radius is composed of irregular corollulæ in the circumference ; and the *disk* of smaller corollulæ, that are for the most part regular.

A *decompound* flower contains within the same calyx lesser calyces, that are each of them common to many flowers, as in SPHÆRANTHUS.

The COROLLA is said to be *equal*, when its parts are equal in figure, magnitude, and proportion ;...*unequal*, when the parts answer in proportion, though not in magnitude, so that the flower comes out to be regular ;...*regular*, when it is equal in respect to the figure, magnitude, and proportion of the parts ;...*irregular*, when the parts of the limb differ in figure, magnitude, or proportion. RICTUS, a *gaping*, or *grinning*, is the gap or opening between the two lips of the corolla. FAUX, the *gorge*, or *gullet*, is the opening of the tube of the corolla. PALATUM, the *palate*, is a gibbosity, or bunching out in the *faux* of the *corolla*. CALCAR, a *spur*, is a nectarium extending in a cone in the hinder part of the *corolla*. The *corolla* is URCEOLATE, *pitcher-shaped*,

* The terms explained here, and in the following paragraphs, respect such circumstances of the parts of fructification as concern rather the *specific* differences than the classic, or generic ones ; and we have therefore followed *Linnaeus* in subjoining them to this head, notwithstanding that some few of them have been already mentioned and explained in the First Part of this work. AUTHOR.

when it is inflate and gibbous on all sides, after the manner of that vessel ;...CYATHIFORM, shaped like a *drinking-glass*, when it is cylindric, but widening a little towards the upper part ;...CON-
NIVING, when there is a convergency of the points of the several lobes of the limb ; or, LACERA, *rent*, when the limb is finely cut.

The ANTHERA is *versatile**,...and *incumbent*†, when it is fastened on at its side ;...and *erect*, when it is fastened on at its base.

The PERICARPIUM is INFLATE, *puffed*, when it is hollow, like a bladder, and not filled up with seeds ;...PRISMATIC, *prism-shaped*, when it is a linear *polyedron*, with plane sides ;...TURBINATE, *top-shaped*, when it tapers towards the base, as in PYRUS ;...CONTORT, *twisted*, when it turns spirally, as in ULMARIA,...HELIC-
TERES, ... and THALICTRUM ; ... ACINACIFORM, *falchion-shaped*, when the fruit is compressed, like a blade, one of the longitudinal angles being obtuse, and the other acute ;...ECHINATE, *prickly*, like an *echinus*‡, when it is beset on all sides with spines or *aculei* ;...TOROSE§, *brawny*, when it is here and there gibbous, with brawny swellings or prominences, as in LYCOPERSICON,... and PHYTOLACCA.

* Easily turned about. EDITOR.

† Resting on. EDITOR.

‡ Hedge-hog. EDITOR.

§ *Torus* signifies properly the rise or swelling out of the strong muscles of an arm. EDITOR.

CHAP. XXIII.

OF VARIETIES.

THE collecting of VARIETIES under their proper species, is a work no less necessary than that of collecting the several species under their proper genus. We have observed in the last Chapter, that such differences as are only incidental to vegetables, and are not found constant and unchangeable in them, are to be considered as *varieties* only. These *varieties* are grounded chiefly on the following circumstances, viz. *sex*,...*magnitude*,...*time of flowering*,...*colour*,...*scent*,...*taste*,...*virtues and uses*;...*duration*,...*multitude*,...*pubescence*,...*leaves*,...and *monstrous flowers*. Of all which we shall treat in their order.

The SEX of plants in the class *Diœcia* affords a variety of all others the most natural; for the male and female flowers in this class being upon different plants, these last are distinguished by the fructification, though the species is the same in both. But it must be observed, that this kind of *variety* holds only in the class *Diœcia*; for in the genera that belong to any of the bisexual classes, the same circumstance, whenever it happens, becomes a specific distinction: thus in RUMEX, which belongs to the class *Hexandria*, the ACETOSA and ACETOSELLA, being *diœcius* plants; that is, having their male and female flowers on distinct roots, these species are thereby distinguished from the rest of the genus.

MAGNITUDE is no specific difference, but a variety, being liable to alteration from the soil or climate.

The TIME of flowering is a treacherous mark of a distinct species; and unless supported by other distinctions, can only be considered as a variety.

COLOUR is found so changeable in the same species, that it must be considered as a variety only.

In FLOWERS the colour is most variable, as in TULIPA,...HEPATICÆ,...CYANUS,... CAMPANULA,... AQUILEGIA,... VIOLA,... GALEGA, ... FUMARIA, and others, which it would be tedious to enumerate: ...the most usual change is from *blue* or *red* to *white*. The trifling distinctions which have been made by *anthophili* (*florists*), in some of the genera we have here instanced, from the colours of the corollæ, and to which they have given such pompous names*, are most difficult to attain, and to be accomplished only by a long attention to the subject. Much fashion reigns here.

FRUITS are observed to change their colour as they ripen; the pericarpium, when it is a berry, changing from *green* to *red*, and from *red* to *white*; and in ripe fruits, the colour, whether *white*, *red*, or *blue*, admits of variation, as in PYRUS,... PRUNUS,... CERASUS, and others†.

SEEDS rarely vary in their colour, though there are instances of it in PAPAVER,... AVENA,... PHASEOLUS,... PISUM,... and FABÆ‡.

* <i>Phœbus,</i>	<i>Triumphus Floræ,</i>	
<i>Apollo,</i>	<i>Pompa Floræ,</i>	
<i>Astræa,</i>	<i>Splendor Asiæ,</i>	
<i>Dædalus,</i>	<i>Corona Europæ,</i>	
<i>Cupido,</i>	<i>Gemma Hollandiæ.</i>	AUTHOR.

- † SOLANUM GUINEENSE *fructu nigerrimo* (B.)
- SOLANUM ANNUM baccis luteis (Dillen.)
- SOLANUM JUDAICUM baccis aurantibus (Dillen.)
- RUBUS VULGARIS *major fructu albo* (Raj.)
- RIEBS VULGARE *acidum albus baccas ferens* (J. B.) AUTHOR.
- ‡ PAPAVER HORTENSE *nigro semine* (C. B.)
- PAPAVER HORTENSE *semine albo* (C. B.)
- AVENA VULGARIS *et alba* (C. B.)
- AVENA NIGRA (C. B.)
- PHASEOLUS VULGARIS *fructu violaceo* (Tourn.)
- PHASEOLUS VULGARIS *fructu ex rubro et nigro variegato* (Tourn.)
- PHASEOLUS *fructu albo venis nigris et lituris distincto* (Tourn.)
- PISUM MAXIMUM *fructu nigra linea maculato* (H. R. P.)
- PISUM HORTENSE *flore fructuque variegato* (C. B.)
- FABÆ *ex rubicundo colore purpurascete.* AUTHOR.

Roots are also little subject to alteration in colour; yet a variation is observed in the roots of DAUCUS,...and RAPHANUS*.

LEAVES are rarely found to quit their green, but they are coloured in AMARANTHUS,...and frequently become spotted, as in PERSICARIA,... RANUNCULUS,... ORCHIS,... HIERACIUM, and LACTUCA†.

The *whole* plant is often found to vary in its colour, as in ERYNGIUM,...ABROTANUM,... ARTEMISIA,...ATRIPLEX,...AMARANTHUS,...PORTULACCA,...and LACTUCA‡.

SCENT in plants is, of all other circumstances, the least to be depended on; and therefore all species grounded on a distinction in the *scent* only, are to be rejected, and referred to varieties.

TASTE in plants is a circumstance variable from soil or culture, and not to be depended on as a real difference. The distinctions of gardeners in fruit of the same species, is considered by *Linnaeus* as a *variety* too minute even to enter the province of botany; and therefore the *various* names§, which have been

* DAUCUS SATIVUS *radice alba* (Tourn.)

DAUCUS SATIVUS *radice lutea* (Tourn.)

DAUCUS SATIVUS *radice aurantii coloris* (Tourn.)

DAUCUS SATIVUS *radice atro-rubescete* (Tourn.)

RAPHANUS NIGER (C. B.) AUTHOR.

† PERSICARIA *cum maculis ferrum equinum referentibus* (Tourn.)

RANUNCULUS HEDERACEUS *atra macula notatus.*

ORCHIS PALMATA *palustris maculata* (C. B.)

HIERACIUM *Alpinum maculatum* (Tourn.)

LACTUCA *maculosa* (C. B.) AUTHOR.

‡ ERYNGIUM *latifolium planum caule ex viridi pallescente flore albo* (Tourn.)

ABROTANUM *caulicidis albicantibus* (Tourn.)

ARTEMISIA *vulgaris major caule ex viridi albicante* (Tourn.)

ATRIPLEX *hortensis rubra* (C. B.)

AMARANTHUS *sylvestris maximus Novæ Angliæ spicis purpureis* (Tourn.)

PORTULACCA *sativa foliis flavis* (Moris.)

LACTUCA *capitata rubra* B. AUTHOR.

§ Poma Paradisiacæ,

Prasomila,

Rubelliana,

Borstorphiæna,

Appianæ,

Melmela,

Pyra Falerna,

Favonia,

Boni Christiana,

Crustamina,

Picena,

Libraria.

AUTHOR.

given to these distinctions, cannot be taught in the science of botany, though, for the purposes of horticulture, they have their use.

The VIRTUES and USES of plants furnish no specific difference ; and the distinctions, therefore, of physical writers are not always to be depended on.

The DURATION of plants is no sure mark of distinct species, being often owing rather to the place, than to the nature of the plant. In warm regions, plants that are *annual* with us will become *perennial*,...or *arborescent*, as is found in TROPÆOLUM,...BETA,...MAJORANA,...MALVA ARBOREA, &c.; and on the contrary, cold regions will occasion *perennial* plants to become *annual*, as is observed in RICINUS,...MIRABILIS*, &c.

MULTITUDE, or quantity, is an accidental circumstance in plants, and cannot conclude any thing, whether the increase be of the *plant* itself, or of its *roots, stems, leaves, or fructification*.

PUBESCENCE is an uncertain mark, as by culture and change of soil, plants are subject to lose as well their *spines* as their *hair* or *down*.

LEAVES, though they for the most part furnish most elegant specific differences, as has been observed in the last Chapter, are yet subject to luxuriation in the same species, which must be carefully distinguished. This may respect their *opposition* and *composition*, and also their being *crisp (curled)*,...or *bullate (bladdery)*.

In respect to OPPOSITION, opposite leaves will sometimes become *ternate*,...*quaternate*,...or *quinate*, growing by *threes, fours, or fives* ; and then the stem also from QUADRANGULAR, *square*, will become *polygonous*, of *many sides*†.

* RICINUS and MIRABILIS, are naturally *perennial* plants, and are only killed by frost in cold countries. AUTHOR.

† LYSIMACHIA *lutea major foliis ternis* (Tourn.)

LYSIMACHIA *lutea major foliis quaternis* (Tourn.)

In respect to COMPOSITION, *digitate* leaves will frequently gain an addition of one or more folioles*.

CRISP, *curled* leaves are a very frequent variety. In TANACETUM,... MENTHA,... OCYMUM,... and MATRICARIA, which are scented plants, there is this singularity observable, that when the leaves are curled, the scent is heightened by the crispature†.

BULLATE, *bladdery* leaves are generally produced from such as are RUGOSE, *wrinkled*; and this is owing to the increase of the substance of the leaf within its vessels, which occasions it to swell and rise. In the SAPONARIA CONCAVA ANGLICANA, a *bullate* leaf is produced in a singular manner from the defect of wrinkles; for here the margin of the leaf contracting itself, the leaves become hollow, like a spoon‡.

Plants are sometimes found to vary from *broad-leaved* to *narrow-leaved*; but this variation is less frequent§.

MONSTROUS flowers, such as the *multiplicate*, *full*, or *proliferous*,

LYSIMACHIA lutea major foliis quinis (Tourn.)

ANAGALLIS cœrulea foliis binis ternisque ex adverso nascentibus (Raj.)

ANAGALLIS Phœnicea foliis amplioribus ex adverso quaternis (Tourn.)

SALICARIA trifolia caule Hexagono (Tourn.) AUTHOR.

* TRIFOLIUM quadrifolium hortense album (C. B.) AUTHOR.

† MALVA crispa (J. B.)

MENTHA crispa Danica (Park.)

TANACETUM foliis crispis (C. B.)

MATRICARIA Crispa.

OCYMUM latifolium maculatum vel crispum (C. B.)

‡ OCYMUM foliis bullatis (C. B.) -

BRASSICA undulata (Rem. alim.)

LACTUCA capitata foliis magis rugosis (B.)

LACTUCA capitata major foliis rugosis et contortis (B.)

LACTUCA capitata omnium maxima verrucosa (B.) AUTHOR.

§ HERACLĒUM hirsutum foliis angustioribus (C. B.)

LYCOPUS foliis in profundas lacinas incisis (Tourn.)

BRASSICA angusta apii folio (C. B.)

VERONICA Austriaca foliis tenuissime laciniatis (Tourn.)

SAMBUCUS laciniato folio (C. B.)

SONCHUS asper laciniatus (C. B.)

VALERIANA Sylvestris foliis tenuissime divisis (C. B.) AUTHOR.

derive their origin from natural ones, therefore are to be considered only as a variety from *luxuriance*.

Upon the whole, the *change of soil* is found to have a great effect on the nature of plants; and to this many of the varieties above mentioned must be imputed; as in BUXUS,...XANTHIUM,... ACANTHUS,...CINARA,...PRUNELLA,...MYOSOTIS,...CRISTA GALLI,... and CERINTHE*; which would all return to their old conditions if the soil were changed again: and in like manner the improvements which are made by culture in the plants cultivated for sale, as in VITIS,...MALUM,...PYRUS,...AMYGDALUS,...PERSICA,... ASPARAGUS,...CERASUS; and in *grain, pulse, and fruit* of all kinds, are not to be esteemed as lasting: for all these, if left to themselves in a poor soil, would run off again, and resume the qualities they had when they grew wild.

The SOIL has some effect also upon *leaves*; for though it is less common for the leaves to differ on the same plant, as they do in some species of LEPIDIUM, ...TITHYMALUS, ... RUDBECKIA, ... and HIBISCUS†; yet it is observed, that *watry* soils are apt to produce a division in the lower leaves of the plant, and even to render capillary such as are produced under the water, as in some species of RANUNCULUS and SISYMBRIUM‡; and also in CICUTA,...SIUM,... PHELLANDRIUM....ŒNANTHE, &c.: and on the contrary, that

* BUXUS *arborescens* (C. B.) BUXUS *humilis* (Dod.)
 XANTHIUM (Dod.) XANTHIUM *Canadense majus* (Tourn.)
 ACANTHUS *mollis* (C. B.) ACANTHUS *aculeatus* (C. B.)
 CINARA *aculeata* (C. B.) CINARA *non aculeata* (C. B.)
 BRUNELLA (Dod.) BRUNELLA *caerulea serena flore* (C. B.)
 MYOSOTIS *foliis hirsutis* (H. C.) et *foliis glabris* (H. C.)
 CRISTA GALLI *fœmina* (J. B.) et *mas.* (J. B.)
 CERINTHE *flore ex rubro purpurascens* (C. B.) et *flavo flore asperior* (C. B.)

AUTHOR.

† TITHYMALUS *heterophyllus* (Plum. Pluk. Alm. 112. f. 6.)
 RUDBECKIA *foliis inferioribus trilobis, superioribus indivisis* (Hort. Upsal.)
 HIBISCUS *foliis inferioribus integris, superioribus trilobis* (Hort. Cliff.)
 LEPIDIUM *foliis caulinis pinnato-multifidis, ramis cordatis amplexicaulibus integris* (H. C.) AUTHOR.

‡ RANUNCULUS *aquaticus folio rundo et capillaceo* (C. B.)
 SISYMBRIUM *foliis simplicibus dentatis serratis* (H. C.) AUTHOR.

mountainous plants usually have their upper leaves more divided, and their lower ones more entire, as in PIMPINELLA, ... PETROSELINUM, ... ANISUM, ... and CORIANDRUM.

VARIETIES may generally be explained and reduced under their proper species with ease, by conferring the variable marks of the variety with the natural plant: but there are some few which are attended with difficulty, and require judgement and experience; as in some species of HELLEBORUS*, ... GENTIANA†, ... FUMARIA‡, ... VALERIANA§, ... SCORPIOIDES||, ... and MEDICAGO¶. In respect to the FUMARIA in question, it is known to be one species only, by the minuteness of its perianthium, the scale of its bud, the structure of its leaves, the situation of the branch, the place of the bractea, the corolla, siliqua, seeds, and stigma; but it varies in the division of its bractea, and in the root being more or less hollow: and that the VALERIANS here spoken of are all of the same species, though they differ so greatly in the fruit, and often in having their leaves more cut, is also proved from their dichotomous stems and annual roots, and from the structure of their

* HELLEBORUS *aconiti folio, flore globoso creceo* (Amm. ruth. 101.) TROLLIUS *humilis flore patula* (Barb. cent. 1. p. 15. l. 22.) *Varietas Hellebori Trollii* (Fl. Succ. 475) *Nectarius longitudine corollæ.* AUTHOR.

† GENTIANA *corolla hypoglottici formi Tubo villis clauso, calycis foliis alternis majoribus* (Fl. Leo. 94.) *Varietas gentianæ fauce barbata* (Fl. Succ. 203.) *flore quadrifido et calycinis laciniis alternis duplo latioribus.* AUTHOR.

‡ FUMARIA *bulbosa radice cava et non cava major et minor.* AUTHOR.

§ VALERIANA *arvensis præcox humilis, semine compresso* (T.)

VALERIANA *arvensis præcox humilis, foliis serratis* (T.)

VALERIANA *arvensis serotina albior, semine turgidiore* (Mor.)

VALERIANA *semine umbilicato nudo rotundo* (Mor.)

VALERIANA *semine umbilicato nudo longo* (Mor.)

VALERIANELLA *semine umbilicato hirsuto majore* (Mor.)

VALERIANELLA *semine umbilicato hirtuso minore* (Mor.)

VALERIANELLA *Cretica, fructu vesicario* (Tourn. Cor.)

VALERIANELLA *semine stellato* (C. B.) AUTHOR.

|| SCORPIOIDES *siliqua campoide hispida* (J. B.)

SCORPIOIDES *siliqua cochleata et striata Ulissiponensis* (T.)

SCORPIOIDES *Bupleuri folio siliquis levibus* (Park.)

SCORPIOIDES *siliqua crassa (Boëlli Cr.)* AUTHOR.

¶ MEDICAGO *leguminibus cochleatis, stipulis, dentatis, caule diffuso* (H. C.)

AUTHOR.

leaves, corollæ, and seeds. Nor should the species of SCORPIURUS and MEDICAGO here instanced be either of them parted, although there is so remarkable a diversity in the fruit of the individuals. In the MEDICAGO* in particular, the forms of the real snails, which nature has imitated in these plants, are scarce more diversified than is the fruit of this mimic species; so that the botanist, who is studious of varieties, would hardly find any end to his labour, of pursuing nature through the various shapes which she has so sportively adopted.

The whole order of the FUNGI, to the scandal of the science, is still a chaos, the botanists not being yet able, in these, to decide with certainty what is a *species*, and what a *variety*†.

* *Medicago scutellata,*
orbiculata,
echinata,
turbinata,
coronata,
doliata,
ciliaris,
tornata,

Medicago hirsuta, .
lupulina,
spinosa,
rugosa,
polycarpus,
dicarpus,
Arabiaca,
Cretica.

AUTHOR.

† Much, however, is expected in this department, from the ingenious labours of the President of the Linnæan Society, Dr. Smith. EDITOR.

TABLE VI.

DEFINITIONS

OF

THE BOTANIC TERMS,

IN LATIN AND ENGLISH,

FROM THE TERMINI BOTANICI OF LINNÆUS.

Consisting of the various Kinds of ROOTS, TRUNKS, BRANCHES, LEAVES, and FRUCTIFICATION, &c. in their natural Arrangement*.

RADIX, THE ROOT,

An Organ by which a Plant receives its Nourishment.

I. DURATION.

- 1 ANNUA, *annual*, that dies in one year.
- 2 BIENNIS, *biennial*, that dies in the space of two years.
- 3 PERENNIS, *perennial*, that regenerates several years successively.

II. FIGURE.

- 4 FIBROSA, *fibrous*, consisting entirely of filaments.
- 5 RAMOSA, *ramous*, subdivided into branchy fibres.
- 6 FUSIFORMIS, *spindle-shaped*, simple, and gradually lessening downward.

* The reader will find in Doctor Thornton's "Elementary Botanical Plates," a number of very excellent plates, explanatory of the definitions of the science, which may be consulted together with our plates.

- 7 PRÆMORSA, *bitten*, or gnawed.
- 8 REPENS, *creeping horizontally*, and putting forth radicles downward, and shooting upwards.
- 9 ARTICULATA, *jointed*, divided into joints.
- 10 DENTATA, *toothed*, having rows of knobs, like teeth.
- 11 GLOBOSA, *round* (158), roots springing from the sides of others.
- 12 TUBEROSA, *tuberous*, consisting of fleshy bodies connected by slender fibres.
- 13 FASCICULARIS, *bunched*, fleshy roots sessile, connected at the base (150).
- 14 PALMATA, *handed*, fleshy lobate roots, like fingers (184).
- 15 BULBOSA, furnished with a bulb (655).
- 16 GRANULATA, *granulated*, round fleshy roots, like seeds.

TRUNCUS, THE TRUNK OR STEM,

The Organ which supports the Branches, Leaves, and Fructification:

I. KINDS.

- 17 CAULIS, a *stem*, which elevates the fructification and leaves.
- 18 CULMUS, a *straw*, properly the trunk of grasses.
- 19 SCAPUS, a *stalk*, elevating the fructification and not the leaves.
- 20 STIPES, a *trunk*, that expands itself into a leaf.

II. DURATION.

- 21 HERBACEOUS, *herb-like*, that perishes every year; an annual stem, not woody.
- 22 SUFFRUTICOSUS, *suffruticous*, *half shrubby*, the root permanent, and the branches sometimes withering.
- 23 FRUTICOSUS, *shrubby*, with perennial stalks arising from the root, that are woody.
- 24 ARBOREUS, *tree-like*, with a single woody stem from the same root.

- 25 SOLIDUS, *solid*, without internal pores.
- 26 INANIS, *pithy*, filled with a spongy substance.
- 27 FISTULOSUS, *fistulous*, hollow like a pipe.

III. DIRECTION.

- 28 ERECTUS, *erect*, rising nearly to a perpendicular direction.
- 29 STRICTUS, *straight*, perpendicular without flexure.

- 30 RIGIDUS, *hard*, not easily bent.
- 31 LAXUS, *loose*, easily bent.
- 32 OBLIQUUS, *awry*, in a direction neither perpendicular nor horizontal.
- 33 ADSCENDENS, *rising upwards*, with a curve like an arch.
- 34 DECLINATUS, *declined*, bending downwards archways.
- 35 INCURVATUS, *incurvate*, bending inwards.
- 36 NUTANS, *nodding*, the top or head bent downwards.
- 37 DIFFUSUS, *diffuse*, with spreading branches.
- 38 PROCUMBENS, *procumbent*, lying on the ground.
- 39 STOLONIFERUS, *producing shoots*, or runners from the root.
- 40 SARMENTOSUS, *thread-like*, producing roots from the joints.
- 41 REPENS, *creeping*, trailing on the ground, and here and there producing roots.
- 42 RADICANS, *rooting*, striking root laterally, and fixing to other bodies.
- 43 GENICULATUS, *jointed*, divided by knots or round swellings.
- 44 FLEXUOSUS, *waved*, bent backwards and forwards from bud to bud.
- 45 SCANDENS, *climbing*, generally by the support of some other body.
- 46 VOLUBILIS, *twining*, growing round some other body in a spiral ascending direction.
- a. DEXTRORSUM, twining from the right to the left.
- b. SINISTRORSUM, twining from the left to the right.

IV. FIGURE.

- 47 TERES, *round*, cylinder-shaped without angles.
- 48 SEMITERES, *half round*, semicylindrical.
- 49 COMPRESSUS, *flattened*, with two opposite sides flat.
- 50 ANCEPS, *two-edged*, flattened, with two opposite sides sharp.
- 51 ANGULATUS, *angulated*, having three or more angles formed by as many intermediate longitudinal cavities.
- a. ACUTANGULUS, *sharp-angled*.
- b. OBTUSANGULUS, *obtusely-angled*.
- 52 TRIQUETER, *three-sided*, having three sides that are quite flat

- 53 TRIGONUS, TETRAGONUS, &c. three-cornered, four-cornered &c. having three, four, or more prominent angles lengthways.
- 54 NUDUS, *naked*, without leaves or other covering.
- 55 APHYLLUS, *without leaves*.
- 56 FOLIATUS, *leafy*, furnished with leaves.
- 57 VAGINATUS, *sheathed*, surrounded with a sheath, formed by the base of the leaf.
- 58 SQUAMOSUS, *squamous*, covered with scales.
- 59 IMBRICATUS, *imbricate*, covered with leaves or scales, placed like tiles, or the scales of fishes.

V. SURFACE.

- 60 SUBEROSUS, *suberous*, the outward bark soft, but elastic, like cork.
- 61 RIMOSUS, *rimous*, the outward bark full of cracks and fissures.
- 62 TUNICATUS, *tunicated*, coated with skins or membranes.
- 63 LEVIS, *smooth*, free from protuberances or inequalities.
- 64 STRIATUS, *striate*, marked with small lines.
- 65 SULCATUS, *sulcate*, furrowed with deep hollow lines.
- 66 GLABER, *slippery*, smooth and glossy, like glass.
- 67 SCABER, *scabrous*, covered with rough prominences.
- 68 MURICATUS, *muricated*, covered with sharp points or prickles.
- 69 TOMENTOSUS, *tomentose*, covered with down.
- 70 LANATUS, *woolly*.
- 71 VILLOSUS, *villous*, covered with soft hair.
- 72 PILOSUS, *pilose*, covered with long hairs that are thinly placed.
- 73 HISPIDUS, *hispid*, covered with stiff hairs or bristles.
- 74 ACULEATUS, *aculeate*, armed with prickles, 378.
- 75 SPINOSUS, *spinous*, armed with thorns, 384.
- 76 URENS, *stinging*, armed with stings, 391.
- 77 STIPULATUS, *stipulate*, having stipula, 291.
- 78 MEMBRANATUS, *membranated*, flat, like a thin pellucid leaf.
- 79 BULBIFERUS, *bearing bulbs*, 655.

VI. COMPOSITION.

- 80 ENODIS, *without knots or joints*, the thickness uniform.
- 81 SIMPLICISSIMUS, *very simple*, with few or no branches.

- 82 SIMPLEX, *simple*, that rises uniform and regular to the top.
 83 INTEGER, *entire*, undivided.
 84 ARTICULATUS, *jointed*.
 85 PROLIFER, *proliferous*, sending forth branches only from the apex.
 86 DICHOTOMUS, *branched always by two*, forked.
 87 BRACHIATUS, *brachiate*, branching opposite, the upper pair crossing the next below.
 88 SUBRAMOSUS, *subramous*, having few lateral branches.
 89 RAMOSUS, *ramous*, having many lateral branches.
 90 RAMOSISSIMUS, *many branches*, subdivided without order, in all directions.
 91 VIRGATUS, *virgated*, with many slender twigs.
 92 PANICULATUS, *paniculated*, whose branches are variously subdivided.
 93 FASTIGIATUS, *fastigiate*, branches arising from a centre to an equal height.
 94 PATENS, *spreading*, 134.
 95 DIVARICATUS, *divaricate*, branches forming an obtuse angle from the trunk, 105.

RAMI PARTES CAULIS,

The Branches Parts of the Stem.

- 96 ALTERNI, *alternate*, when they come out single, and follow in gradual order, 115.
 97 DISTICHI, *distichous*, in two rows.
 98 SPARSI, *sparsed*, scattered without order, 118.
 99 CONFERTI, *crowded*, 119.
 100 OPPOSITI, *opposite*, 126.
 101 VERTICILLATI, *verticillate*, branches surrounding the stem, or at the joints, like the rays of a vessel.
 102 ERECTI, *erect*, upright, perpendicular.
 103 COARCTATI, *close together*, almost touching towards the top.
 104 DIVERGENTES, *divergent*, branches growing from the trunk at right angles, like rays from a centre.
 105 DIVARICATI, *divaricate*, branches shooting from the trunk, so as to make an obtuse angle.
 106 DEFLEXI, *deflex*, bending downwards archwise.

- 107 REFLEXI, *reflex*, bending back towards the trunk.
 108 RETROFLEXI, *retroflex*, bending backward and forward towards the trunk.
 109 FULCRATI, *fulcrate*, having props or supports.

THE LEAVES,

The Organs by which Plants are put in Motion.

I. THEIR PLACE.

- 110 RADICALE, *radical*, springing from the root.
 111 CAULINUM, *cauline*, springing from the stem.
 112 RAMEUM, *rameous*, growing on the branches.
 113 AXILLARE, *axillary*, placed at the insertion of the branch.
 114 FLORALE, *floral*, placed near the flower, and are commonly smaller.

II. SITUATION.

- 115 ALTERNA, *alternate*, when they come out single, and follow in a gradual order.
 116 DISTICHA, *distichous*, disposed in two opposite rows, though inserted on all sides.
 117 BIFARIA, *bifarious*, inserted only on two opposite sides of a branch or middle rib.
 118 SPARSA, *sparsed*, scattered in no certain order.
 119 CONFERTA, *confert*, crowded together.
 120 IMBRICATA, *imbricate*, lying over one another like scales of fishes.
 121 FASCICULATA, *fasciculate*, growing in bunches from one point.
 122 GEMINA, TRINA, &c. *two, three*, or more together from the same point.
 123 CONFLUENTIA, *confluent*, growing together or running into one another at the base.
 124 APPROXIMATA, *approximate*, mutually approaching each other.
 125 REMOTA, *remote*, placed at some distance from each other.
 126 OPPOSITA, *opposite*, growing opposite, but in such a manner that each pair crosses the other above and below.

- 127 DECUSSATA, *decussated*, where the pairs cross each other in a regular manner.
- 128 VERTICILLATA, *verticillate*, *whorled*, where three or more leaves surround the stem.
- 129 TERNATA, QUATERNA, &c. *three or four together*, &c. according to the number of leaves surrounding each joint.

III. DIRECTION.

- 130 ERECTUM, *erect*, upright, perpendicular.
- 131 STRICTUM, *straight*, quite perpendicular, without flexure or bending.
- 132 RIGIDUM, *rigid*, stiff, not easily bent.
- 133 ADPRESSUM, *adprest*, the disk of the leaf pressed towards the stem.
- 134 PATENS, *patent*, spreading, making an acute angle with the stem.
- 135 HORIZONTALIS, *horizontal*, growing from the stem at right angles.
- 136 ASSURGENS, *assurgent*, bending upwards, 33.
- 137 INFLEXUM, *inflex*, bending inwards towards the stem.
- 138 RECLINATUM, *reclinate*, bending downwards archwise, the apex ascending.
- 139 RECURVATUM, *recurvate*, bent backwards, in the form of an arch, the convex side upwards.
- 140 REVOLUTUM, *revolute*, rolled back, in form of a scroll.
- 141 PENDENS, *pendent*, hanging with the point downwards.
- 142 OBLIQUUM, *oblique*, the base looking upwards, the apex to the horizon.
- 143 VERTICALE, *vertical*, leaves so situated that the base is perpendicular to the apex.
- 144 RESUPINATUM, *resupinate*, when the lower disk of the leaf looks upwards.
- 145 SUBMERSUM, *submersed*, sunk under the surface of the water.
- 146 NATANS, *natant*, floating on the surface of the water.
- 147 RADICANS, *radicant*, striking root.

IV. INSERTION.

- 148 PETIOLATUM, *petiolate*, having a petiole or footstalk, 290.
- 149 PELTATUM, *peltate*, having the foot-stalk inserted into the disk of the leaf.
- 150 SESSILE, *sessile*, sitting immediately on the stem, without a foot-stalk.
- 151 ADNATUM, *adnate*, the upper disk of the leaf adhering to the stem by an attachment of its base.
- 152 COADUNATA, *condunate*, several growing together at their base.
- 153 DECURRENS, *decurrent*, where the base of a sessile leaf is elongated, and runs down the stem.
- 154 AMPLEXICAULE, *amplexicaul*, embracing the stem with its base.
- 155 PERFOLIATUM, *perfoliate*, where the base of the leaf entirely surrounds the stem, or when the stalk grows through the centre of the leaf.
- 156 CONNATA, *connate*, where two opposite leaves grow together at their bases.
- 157 VAGINANS, *vaginant*, where the base of the leaf forms a tubular sheath that surrounds the stem.

V. FIGURE.

- 158 SUBROTUNDUM, *subrotund*, almost round, nearly circular.
- 159 ORBICULATUM, *orbiculate*, of a circular figure.
- 160 OVATUM, *ovate*, egg-shaped.
- 161 OVALE, *oval*, the shape of an egg, when both ends are equal.
- 162 OBLONGUM, *oblong*, twice the length of its breadth.
- 163 PARABOLICUM, *parabolic*, like the smaller end of an egg.
- 164 CUNEIFORME, *cuneiform*, wedge-shaped, tapering from the apex to the base.
- 165 SPATULATUM, *spatulate*, rounded at the apex, and narrower and linear at the base.
- 166 ROTUNDATUM, *rotundate*, rounded, or with angles in a circle.
- 167 LANCEOLATUM, *lanceolate*, oblong, and tapering towards both extremities.
- 168 ELLIPTICUM, *elliptical*, an oval whose ends are equal.
- 169 LINEARE, *linear*, every where of the same breadth.

170 ACEROSUM, *acerose*, linear, and permanent, like chaff, or the leaves of pines.

VI. ANGLES.

171 INTEGRUM, *entire*, undivided, without divisions.

172 TRIANGULARE, *triangular*, &c. three-angled, &c.

173 DELTOIDEUM, *deltoid*, a leaf whose angles are formed like the Greek delta.

174 RHOMBEUM, *rombus-shaped*, an irregular four-sided figure, resembling the ace of diamonds.

VII. SINUSES.

175 TRAPEZIFORME, *trapeziform*, a figure of four unequal sides.

176 CORDATUM, *cordate*, heart-shaped.

177 RENIFORME, *reniform*, kidney-shaped.

178 LUNATUM, *lunate*, shaped like a half moon.

179 SAGITTATUM, *sagittate*, arrow-shaped.

180 HASTATUM, *hastate*, spear-shaped.

181 RUNCINATUM, *runcinate*, like the teeth of a great saw, whose serratures are bent downwards

182 PANDURIFORME, *panduriform*, fiddle-shaped.

183 FISSUM, *slit*, divided into linear partitions.

184 LOBATUM, *lobate*, divided into lobes.

185 BILOBUM, TRILOBUM, &c. *two and three-lobed*, &c. according to the number of lobes.

186 PARTITUM, *partite*, divided almost to the base; the number of divisions are expressed by the terms *Bipartite*, *Tripartite*, &c.

187 PALMATUM, *palmate*, divided like a hand.

188 LYRATUM, *lyrate*, *lyre-shaped*, with transverse divisions broadest at the apex, the lower ones gradually less and more distant.

189 PINNATIFIDUM, *pinnatifid*, deeply divided into transverse, lateral, oblong segments.

190 SINUATUM, *sinuate*, divided into lateral hollows.

191 LACINIATUM, *laciniate*, divided into segments.

192 SQUARROSUM, *squarrose*, divided into elevated segments, not plane or parallel, as in the calyx of some syngenesious plants.

VIII. MARGIN.

- 193 INTEGERRIMUM, *very entire*, without any incision.
- 194 CREMATUM, *crenate*, where the margin is notched at right angles to the centre without inclining to either extremity.
- 195 SERRATUM, *serrate*, sawed, notches like the teeth of a saw, inclining all the same way, either towards the point or base.
- 196 CILIATUM, *ciliate*, where bristles are arranged in a parallel order on the margin of the leaf, like eye-lashes.
- 197 DENTATUM, *dentate, toothed*, points like teeth protruding from the margin of the leaf, at some distance from each other.
- 198 SPINOSUM, *spinose*, where the margin is armed with sharp spines.
- 199 CARTILAGINEUM, *cartilaginous*, where the margin is hard and tough.
- 200 REPANDUM, *repand*, where the margin is waved.
- 201 LACERUM, *lacerate*, where the margin is variously divided, as if torn.
- 202 EROSUM, *erose*, where the margin is sinuate, as if gnawed with teeth.
- 203 MEMBRANACEUM, *membranaceous*, where the margin is thin and pellucid.
- 204 DÆDALEUM, *dedalous*, where the margin has many various windings and turnings.

IX. APEX.

- 205 OBTUSUM, *obtuse*, where the point is rounded.
- 206 EMARGINATUM, *emarginate*, where the apex is notched.
- 207 RETUSUM, *retuse*, terminating in an obtuse hollow.
- 208 PRÆMORSUM, *premorse*, where the termination appears as if bitten off.
- 209 TRUNCATUM, *truncate*, terminating in a line, as if cut off.
- 210 ACUTUM, *acute*, terminating in a sharp angle.
- 211 ACUMINATUM, *acuminate*, terminating in a sharp point.
- 212 CUSPIDATUM, *cuspidate*, terminating in a point, like a spear.

- 213 MUCRONATUM, *mucronate*, terminating in a small prickle.
- 214 CIRRHOSUM, *cirrhose*, terminating in a clasper or tendril, 292.
- X. SURFACE.
- 215 NUDUM, *naked*, without hairs or excrescences.
- 216 GLABRUM, *smooth*, slippery.
- 217 NITIDUM, *glossy*, smooth and shining.
- 218 LUCIDUM, *lucid*, bright, reflecting light.
- 219 COLORATUM, *coloured*, of a colour different from green.
- 220 NERVOSUM, *nervous*, with nerves extended from the base to the apex.
- 221 TRINERVE, where three nerves join at the base and apex.
- 222 TRIPLINERVE, where three nerves are each divided into three more above the base.
- 223 TRINERVATUM, where three nerves run into each other at the base.
- 224 ENERVE, *without nerves*, opposite to nervous.
- 225 LINEATUM, *lined*, with depressed nerves or hollow lines.
- 226 SULCATUM, *furrowed*, with deep lines.
- 227 VENOSUM, *veined*, with veins many ways.
- 228 RUGOSUM, *rugose*, wrinkled, shrivelled, rough.
- 229 BULLATUM, *studded*, bladdery, alternately convex and concave.
- 230 LACUNOSUM, where the disk of the leaf is depressed into deep cavities between the veins that run parallel from the disk to the margin.
- 231 AVENE, *without veins*.
- 232 PUNCTATUM, *punctate*, with hollow scattered punctures.
- 233 PAPILLOSUM, *papillose*, covered with fleshy punctures.
- 234 PAPULOSUM, *papulose*, covered with vascular punctures.
- 235 VISCIDUM, *viscid*, covered with a viscid humour.
- 236 VILLOSUM, *villous*, covered with soft hairs.
- 237 TOMENTOSUM, *downy*, covered with downy hairs.
- 238 SERICEUM, *silky*, covered with soft silky hairs.
- 239 LANATUM, *woolly*, covered with woolly hairs.
- 240 BARBATUM, *bearded*, hairs growing in tufts.
- 241 PILOSUM, *pilous*, covered with long hairs that appear distinctly.

242 SCABRUM, *rough*, covered with rigid punctures raised above the surface.

243 HISPIDUM, *hispid*, covered with hard bristles.

244 ACULEATUM, *prickly*, covered with sharp prickles, 378.

245 STRIGOSUM, *strigous*, armed with lance-shaped prickles, 167.

XI. EXPANSION.

246 PLANUM, *plane*, with a flat equal surface.

247 CANALICULATUM, *channelled*, a deep channel or furrow, running lengthways.

248 CONCAVUM, *concave*, when the disk is arched from the margin, and forms a hollow.

249 CONVEXUM, *convex*, opposite to concave: these two terms arise from the same cause, the margin being too tight for the expansion of the disk; therefore if a leaf is concave on one side, it is convex on the other.

250 CUCULLATUM, *hollowed*, when the sides of a leaf press together at the base, and expand towards the apex.

251 PLICATUM, *plaited*, folded in sharp flexures from the disk to the margin.

252 UNDATUM, *waved*, the flexures or folds being obtuse from the disk to the margin.

253 CRISPUM, *curled*, where the margin is plaited, but the folds do not reach to the middle rib of the disk.

XII. SUBSTANCE.

254 MEMBRANACEUM, *skinny*, pellucid, without any fleshy substance.

255 SCARIOSUM, of a *dry parched substance*, that sounds when touched.

256 GIBBUM, *gouty*, when both sides of a leaf is bunched out by a copious quantity of pulp.

257 TERES, *cylindrical*, or pillar-shaped.

258 DEPRESSUM, more pulpy in the disk, and flatted towards the sides.

259 COMPRESSUM, more flatted in the disk, and pulpy towards the sides.

- 260 CARINATUM, *carinate*, the lower part of the disk prominent lengthwise.
- 261 COMPACTUM, *compact*, of a solid substance.
- 262 TUBULOSUM, *tubulous*, the inside hollow, without pith.
- 263 PULPOSUM, *pulpous*, of a fleshy pulpy substance.
- 264 CARNOSUM, *fleshy*, the inside of a solid pulp.
- 265 TRIQUETRUM, *triquetrous*, three-cornered lengthwise.
- 266 ANCEPS, *two-angled*, or edged lengthwise.
- 267 LINGULATUM, *tongue-shaped*, linear, fleshy, the lower side convex.
- 268 ENSIFORME, *sword-shaped*, doubled-edged, gradually lessening from the base to the point.
- 269 SUBULATUM, *subulate*, linear at the base, and smaller towards the point.
- 270 ACINACIFORME, *scymitar-shaped*, fleshy, and compressed, one side convex sharp, the other straight and thicker.
- 271 DOLABRIFORME, *hatchet-shaped*, compressed and half-round, gibbous outward, the edge sharp, the lower part rounded.

XIII. DURATION.

- 272 DECIDUUM, *deciduous*, finished, and falling off in one summer.
- 273 CADUCUM, *cadent*, falling off, short duration, not abiding through the summer.
- 274 PERSISTENS, *persisting*, abiding, lasting or remaining more than one summer.
- 275 PERENNE, *perennial*, continuing green many years.
- 276 SEMPERVIRENS, *evergreen*, green at all times of the year.

XIV. COMPOSITION.

- 277 ARTICULATUM, *articulate*, a leaf having a little leaf growing out of its point.
- 278 CONJUGATUM, *conjugate*, winged, the little leaves or wings coming by pairs.
- 279 DIGITATUM, *digitate*, a single foot-stalk connecting the little leaves at its top.

- 280 BINATUM, TERNATUM, QUINATUM, &c. terminating by two, three, or five little leaves or folioles.
- 281 PEDATUM, *pedate*, like the toes of the feet, the foot-stalk dividing sideways obliquely, and connecting many folioles.
- 282 PINNATUM, *pinnate*, winged, a simple foot-stalk, connecting many little leaves sidewise.
- 283 BIJUGUM (thus TRIJUGA, QUADRIJUGA, QUINQUEJUGA, SEJUGA, &c.) winged, but the little leaves coming by pairs, and are four, six, eight, ten, twelve, &c.
- CUM IMPARI, winged, not terminating in pairs, but with an odd foliole.
- AERUPTÉ PINNATUM, *abruptly winged*, terminating without a tendril, or an odd foliole.
- CIRROSUM, *cirrhous*, terminating in a tendril or clasper, 292.
- FOLIOLIS OPPOSITIS (126), the little leaves growing opposite.
- FOLIOLIS ALTERNIS (115), the little leaves growing alternate.
- RUPTIS, the little leaves alternately smaller, broken.
- DECURSIVIS, the foot-stalks of the little leaves running down the middle rib, or rachi, 153.

XV. DECOMPOSITION.

- 284 BIGEMINUM, *the foot-stalk forked by twos* (86), connecting many little leaves.
- 285 BITERNATUM, *doubled by threes*, 280.
- 286 BIPINNATUM, *double winged*, 282.

XVI. TRIPLE COMPOSITION.

- 287 TERGEMINUM, *triple budded*.
- 288 TRITERNATUM, *three times three*.
- 289 TRIPINNATUM, *three ways winged*:

FULCRA, PROPS,

Supports for the better sustaining the different Parts of Plants.

- 290 PETIOLUS, a *foot-stalk* that sustains the leaf.
- 291 STIPULA, a *scale* at the base of the foot-stalk which it supports.

- 292 CIRRHUS, *clasper or tendril*, growing like threads, in a spiral form, which takes hold of plants, or any other body near it.
- 293 PUBES, *downy hairs* in all plants.
- 294 ARMA, *armed with points*, to keep off animals from hurting them.
- 295 BRACTEA, *floral leaves*, the face and texture different from other leaves.
- 296 PEDUNCULUS, the *foot-stalk*, or prop that sustains the fructification.

PETIOLUS, FOOT-STALK OF THE LEAF.

I. FIGURE.

- 297 LINEARIS (169), *linear*, every where the same breadth.
- 298 ALATUS, *winged*, spread out at the sides.
- 299 CLAVATUS, *clubbed*, thickened towards the point.
- 300 MEMBRANACEUS, *flat*, thin, and generally pellucid.
- 301 TERES (257), *rounded like a cylinder*, pillar-shaped.
- 302 SEMITERES (48), *half-rounded*, like a split column.
- 303 TRIQUETER (52), *three-sided*.

II. MAGNITUDE.

- 304 BREVISSIMUS, *very short*, when the length of the foot-stalk is not equal to the length of the leaf.
- 305 BREVIS, *short*, not quite so long as the leaf.
- 306 MEDIOCRIS, *of the length of the leaf*.
- 307 LONGUS, *longer than the leaf*.
- 308 LONGISSIMUS, *something longer than the leaf*.

III. INSERTION.

- 309 INSERTUS, *inserted*, joined.
- 310 ADNATUS (151), *adhering to*.
- 311 DECURRENS (153), *running down the branch*.
- 312 AMPLEXICAULIS (154), *embracing the stalk with its base*.
- 313 APPENDICULATUS, a *leafy appendage* adhering to the base of a leaf.

IV. DIRECTION.

- 314 ERECTUS (130), *upright*.
 315 PATENS (134), *spreading*.
 316 ASSURGENS (136), *bending upwards in a kind of arch*.
 317 RECURVATUS (139), *bent backwards*.

V. SURFACE.

- 318 GLABER (216), *smooth*.
 319 ACULEATUS (244), *prickly*.
 320 NUDUS (215), *naked*.
 321 ARTICULATUS (34), *jointed*.
 322 SPINESCENS, *hard, and sharp*.

STIPULÆ, APPENDAGES TO THE LEAF.

- 323 GEMINÆ, *two and two, by pairs*.
 324 SOLITARÆ, *single scattered*.
 325 LATERALES, *inserted in the sides*.
 326 EXTRAFOLIACÆ, *on the outside, below the base of the petiole*.
 327 INTRAFOLIACÆ, *on the inside, above the base of the petiole*.
 328 OPPOSITIFOLIACÆ, *opposite, placed on the sides at the base of the leaf*.
 329 CADUCÆ (273), *falling off, withering before the leaf*.
 330 DECIDUÆ (272), *falling annually*.
 331 PERSISTENTES, *abiding after the leaf falls off*.
 332 SPINESCENTES (322), *hard and sharp, like a spine or prickle*.
 333 SESSILES (150), *squat, having no foot-stalk*.
 334 ADNATÆ (151), *adhering to the branch by an attachment of its upper surface*.
 335 DECURRENTES (153), *running down the branch*.
 336 VAGINANTES (157), *surrounding the stem like a sheath*.
 337 SUBULATE (269), *awl-shaped*.
 338 LANCEOLATÆ (167), *lance-shaped*.
 339 SAGITTATÆ (179), *arrow-shaped*.
 340 LUNATÆ (178), *moon-shaped*.
 341 ERECTÆ (130), *upright*.

- 342 PATENTIS (134), *spreading*.
 343 INTEGRÆ (193), *entire*.
 344 SERRATÆ (195), *sawed*.
 345 CILIATÆ (196), *lashed, like the eye*.
 346 DENTATÆ (197), *toothed*.
 347 FISSÆ (183), *split*.

CIRRHUS, A TENDRIL OR CLASPER.

- 348 AXILLARIS (113), *at the insertion of the branch*.
 349 FOLIARIS, *sitting on a leaf*.
 350 PETIOLARIS, *growing on the foot-stalk of the leaf, 290*.
 351 PEDUNCULARIS (296), *growing on the foot-stalk of the flower*.
 352 SIMPLEX, *undivided*.
 353 TRIFIDUS, *divided in three parts*.
 354 MULTIFIDUS, *divided in many parts*.
 355 CONVOLUTUS, *twisting in the same direction as the sun, in rings*.
 356 REVOLUTUS, *revolute, rolled back in half spiral rings*.

PUBES, DOWN OR PUBESCENCE.

- 357 PILI, *excretory ducts, long distinct hairs*.
 358 LANA, *wool, curled hairs and thick*.
 359 BARBA, *bearded tufts of parallel hairs*.
 360 TOMENTUM, *down, hairs scarcely conspicuous*.
 361 STRIGÆ, *strong hard flat hairs*.
 362 SETÆ, *bristles, rigid round hairs*.
 363 SIMPLICES, *single, not divided*.
 364 HAMOSÆ, *hooked, by which they easily adhere to animals*.
 365 RAMOSÆ, s. FURCATÆ, *subdivided into little branches, or forked*.
 366 PLUMOSÆ, *feathery, composed of fine down, or hairs*.
 367 STELLATÆ, *starry, disposed crosswise*.
 368 HAMI, *hooks, prickles with recurved points*.
 369 GLOCHIDES, *prickles, with the points turned back, having many teeth*.

- 370 GLANDULA, *glands*, little glands for throwing out the excrementitious humour of plants; these are either *sessiles*, squat; *stipulata*, having a foot-stalk; or, *porus*, having a pore, often perforating a leaf.
- 371 UTRICULUS, *little vessels*, replete with secretory liquor.
- 372 FOLIACEI, inserted in the leaves.
- 373 PETIOLARES (350), inserted in the foot-stalk of the leaf.
- 374 PEDUNCULARES (351), inserted in the foot-stalk of the flower.
- 375 STIPULARES (291), inserted in the stipula.
- 376 VISCOSITAS, a humour of a clammy quality.
- 377 GLUTINOSITAS, a humour whose quality is of a lubricating slippery nature.

ARMA, ARMS.

- 378 ACULEI, *sharp prickles* fixed in the bark of plants.
- 379 RECTI, *straight*, without bending.
- 380 INCURVI, *bent inwards*.
- 381 RECURVI, *bent outwards*.
- 382 FURCÆ, *prickles divided into many forks*.
- 383 BIFIDÆ and TRIFIDÆ, *by two and three*, or according to the number of divisions.
- 384 SPINA, a *spine*, a prickle fixed in the wood of the trunk or branch.
- 385 TERMINALIS, *terminating the branch*.
- 386 AXILLARIS (113), growing from the insertion of the branch.
- 387 CALYCINA, growing on the cup.
- 388 FOLIARIS (349), growing on the leaf.
- 389 SIMPLEX (363), single.
- 390 DIVISA, divided at the point.
- 391 STIMULI, *stings*, that make inflammatory punctures, which go off with an itching.

BRACTÆ, FLORAL LEAVES.

- 392 COLORATÆ (219), coloured.
- 393 CADUCÆ (273), falling off with the flower.

- 394 DECIDUE (272), falling off.
 395 PERSISTENTES (274), abiding.
 396 COMA, a *bractæ*, terminating the stalk above the flower, distinguished by its magnitude or colour.

PEDUNCULUS, FOOT-STALK OF A FLOWER.

- 397 PARTIALIS, in some flowers growing from the common foot-stalk.
 398 COMMUNIS, a *foot-stalk* common to many flowers.
 399 PEDICELLUS, a *little foot-stalk*, proper to flowers that have a common foot-stalk, 398.
 400 SCAPUS, a *peduncle*, rising from the root, resembling a stalk.

I. PLACE.

- 401 RADICALIS (110), springing from the root.
 402 CAULINUS (111), springing from the stem.
 403 RAMEUS (112), growing from the branch.
 404 PETIOLARIS (350), growing from the petiole.
 405 CIRRHIFERUS (292), growing from the tendril or clasper.
 406 TERMINALIS (385), terminating the branch.
 407 AXILLARIS (113), at the insertion of the branch or leaf.
 408 OPPOSITIFOLIUS (328), having opposite leaves.
 409 LATERIFLORUS (325), flowering at the sides.
 410 INTRAFOLIACEUS (327), within the leaves.
 411 EXTRAFOLIACEUS (326), on the outside of the leaves.

II. SITUATION.

- 412 ALTERNI (115), alternate.
 413 SPARSI (118), scattered.
 414 OPPOSITI (126), opposite.
 415 VERTICILLATI (128), in circles round the stem.

III. NUMBER.

- 416 SOLITARIUS (324), single.
 417 GEMINATUS (323), by twos.
 418 UMBELLULA SESSILIS, many peduncles from the same centre, produced of the same height.

IV. DIRECTION.

- 419 ADPRESSUS (133), pressed towards the stem.
 420 ERECTUS (130), upright.
 421 PATENS (134), spreading.
 422 CERNUUS, the point looking downwards.
 423 RESUPINATUS (144), looking upwards.
 424 DECLINATUS (34), bent downwards archwise.
 425 NUTANS (36), *nodding*, hanging downward.
 426 FLACCIDUS, *slender*, weak, when the weight of a proper flower makes it hang downwards.
 427 ASCENDENS (33), rising upwards archwise.
 428 PENDULUS, hanging loose.
 429 STRICTUS (29), straight.
 430 FLEXUOSUS, bending from one flower to another.
 431 RETROFRACTUS, bent backward and forward, as if broken.
 432 UNIFLORUS, BIFLORUS, TRIFLORUS, &c. MULTIFLORUS, one flower, two flowers, three flowers, &c. many flowers, according to the number of flowers growing on the foot-stalk.

V. STRUCTURE.

- 433 TERES (47), *round*, like a cylinder.
 434 TRIQUETER (52), *three-sided*.
 435 TETRAGONUS (53), *four-angled*.
 436 FILIFORMIS, *thread-shaped*, every where of equal thickness.
 437 ATTENUATUS, lessening gradually in thickness towards the point.
 438 CLAVATUS, *clubbed*, thick towards the point, 209.
 439 INCRASSATUS, *gradually thickening upwards*.
 440 NUDUS (215), *naked*.
 441 SQUAMOSUS (58), *scaly*.
 442 FOLIATUS (56), *leafy*.
 443 BRACTEATUS (295), *furnished with floral leaves*.
 444 GENICULATUS (43), *jointed*.
 445 ARTICULATUS (84), *knotted*.

INFLORESCENTIA, INFLORESCENCE,

Is the manner by which Flowers are joined to the Plant by the Peduncle or Foot-stalk.

- 446 VERTICILLUS, *whorled*, many flowers growing round the stalk in a circle.
- 447 SESSILES, *squat*, without any manifest foot-stalk.
- 448 PEDUNCULATUS, a *peduncle*, elevating the flowers.
- 449 NUDUS (450, 451), opposite to the following.
- 450 INVOLUCRATUS (520), furnished with an involucre.
- 451 BRACTEATUS (443), having floral leaves.
- 452 CONFERTUS, foot-stalks crowded together.
- 453 DISTANS, the foot-stalks distant
- 454 CAPITULUM, a *head*, flowers collected into a globe or head.
- 455 SUBROTUNDUM (456), *nearly of a globular figure*, almost round.
- 456 GLOBOSUM, *globular*, perfectly round.
- 457 DIMIDIATUM, *halved*, like a globe cut into two parts.
- 458 FOLIOSUM, *leafy*, leaves intermixed with the flowers.
- 459 NUDUM, *naked*, without leaves or bristles.
- 460 FASCICULUS, *bunched*, a flower growing in bunches.
- 461 SPICA, *sessile flowers*, growing alternate on a common peduncle.
- 462 SIMPLEX, a *single spike*, undivided.
- 463 COMPOSITA, many little spikes growing from the common peduncle.
- 464 GLOMERATA, many little spikes crowded together.
- 465 OVATA (160), *egg-shaped*.
- 466 VENTRICOSA (256), *swollen*, gouty.
- 467 CYLINDRICA, *pillar-shaped*.
- 468 INTERRUPTA, *spikes alternately smaller*.
- 469 IMERICATA (120), *scaled*.
- 470 ARTICULATA (84), *knotted*, jointed.
- 471 RAMOSA, *branching variously*.
- 472 LINEARIS (169), *linear*, of equal width, lengthwise.

- 473 CILIATA (196), *lashed*.
- 474 FOLIACEA, *leafy*.
- 475 COMOSA, *terminating in little leaves*.
- 476 CORYMBUS (461), *a kind of spike, whose flowers are furnished with foot-stalks, so proportioned to their situation, as to elevate all the flowers of the spike to the same height*.
- 477 THYRSUS (489), *a kind of crowded panicle, of an ovate form*.
- 478 RACEMUS, *a bunch of flowers, the peduncles coming at the sides*.
- 479 SIMPLEX, *undivided*.
- 480 COMPOSITUS, *divided into many*.
- 481 UNILATERALIS, *all the flowers growing on one side*.
- 482 SECUNDUS, *the flowers all bending to one side*.
- 483 PEDATUS (281), *the foot-stalk coming on one side, like the toes of the feet*.
- 484 CONJUGATUS (278), *joined by twos*.
- 485 ERECTUS (130), *upright*.
- 486 LAXUS (31), *loose, not closely connected*.
- 487 NUDUS (459), *naked*.
- 488 FOLIATUS (56), *leafy*.
- 489 PANICULA, *flowers scattered on peduncles that are divided in different forms*.
- 490 SIMPLEX, *always few flowers*.
- 491 COMPOSITA, *many florets coming together*.

FRUCTIFICATIO, FRUCTIFICATION.

Temporary Parts of Vegetables, called the Generation.

- 492 CALYX, *a flower cup*, is the termination of the outer bark of the plant, present in the fructification.
- 493 PERIANTHIUM, *a flower cup*, whose station is close to the fructification.
- 494 FRUCTIFICATIONIS, when it includes the stamina and germen.
- 495 FLORIS, containing the stamina without the germen.
- 496 FRUCTUS, containing the germen without the stamina.

- 497 PROPRIUM, without respect to the flower.
- 498 MONOPHYLLUM, consisting of one leaf.
- 499 POLYPHYLLUM, consisting of many leaves.
- 500 2—5 FIDUM (183), divided into two, three, four, or five divisions.
- 501 2—5 PARTITUM (186), divided almost to the base, from two to five.
- 502 INTEGRUM, *entire* (171), undivided.
- 503 TUBULOSUM (262), *tube-shaped*.
- 504 PATENS (134), *spreading*.
- 505 REFLEXUM, the parts bent backwards.
- 506 INFLATUM, puffed out like a bladder.
- 507 ABBREVIATUM, shorter than the tube of the corolla.
- 508 OBTUSUM (205), the divisions rounded.
- 509 ACUTUM (210), the divisions sharp.
- 510 SPINOSUM (75), bearing spines.
- 511 ACULEATUM (244), bearing prickles.
- 512 SUPERUM, when the germen is below the receptacle.
- 513 INFERUM, when the germen is above the receptacle.
- 514 COMMUNE, a *common calyx*, containing many florets, as in compound flowers.
- 515 IMBRICATUM, *scaled*, various scales lying over one another.
- 516 SQUARROSUM, with scales pointing many ways.
- 517 SCARIOSUM, *having scales*; their margins are membranaceous, hard, dry, and sounding when touched.
- 518 TURBINATUM, *top-shaped*, like an obverse cone.
- 519 CALYCVLATUM, *when a lesser calyx is added*, and encircles the base of the larger one.
- 520 INVOLUCRUM, a kind of calyx, standing remote from the flower.
- 521 UNIVERSALE, in umbelliferous plants, standing under the universal umbel.
- 522 PARTIALE, an involucre, standing under the partial umbel.
- 523 PROPRIUM, always under the flower.
- 524 GLUMA, a *husk*, a cup belonging to the grasses, whose flowers it embraces, with the valves folded over.

- 525 UNIFLORA, *one-flowered*, when it embraces one flower.
- 526 MULTIFLORA, *many-flowered*, when it includes many flowers.
- 527 UNIVALVIS, *one-valved*, when there is constantly but one scale.
- 528 BIVALVIS, *two-valved*, when there are two valves.
- 529 MULTIVALVIS, *many-valved*, when there are many scales, or more than two.
- 530 COLORATA (219), *coloured*.
- 531 GLABRA (216), *smooth*.
- 532 HISPIDA (243), *hispid, covered with hard hairs*.
- 533 MUTICA, *blunt, without point*, or arista.
- 534 ARISTA, *a beard*, growing on the husk.
- 535 TERMINALIS, *terminal*, fixed to the top of the husk.
- 536 DORSALIS, *dorsal*, fixed on the outside of the husk.
- 537 RECTA, *straight*, growing perpendicular:
- 538 TORTILIS, *twisted*.
- 539 GENICULATA (43), *jointed*.
- 540 RECURVATA (139), *recurved*.
- 541 AMENTUM (635), *a catkin, or ament*, proceeding from a common receptacle, resembling the chaff of corn.
- 542 SPATHA, *a sheath*, a kind of cup bursting out lengthwise.
- 543 UNIVALVIS, *of one valve*, opening on one side.
- 544 DIMIDIATA, *dimidiate, halved*, the inner one covering the fructification on one side, and the outer one on the other.
- 545 CALYPTRA, *a veil, or hood*, covering the antheræ, in mosses.
- 546 RECTA, *straight*, every where equal.
- 547 OBLIQUA, *oblique*, bent on one side.
- 548 VOLVA, *a membranaceous calyx* belonging to the fungi.
- 549 APPROXIMATA, *approximate*, close to the head.
- 550 REMOTA, *remote*, at some distance from the head.
- 551 COROLLA, the termination of the inner bark, present in the flower.
- 552 PETALUM, *a petal*, a part of the corolla when divided into many.
- 553 TUBUS, *a tube*, the lower part of a flower, with one petal.
- 554 UNGUIS, *a claw*, the lower part of a polypetalous flower, by which it is fixed to the receptacle.
- 555 LIMBUS, *limb*, the upper part of a monopetalous corolla.

- 556 LAMINA, *the upper spreading part* of a polypetalous flower.
 MONOPETALA, VEL POLYPETALA, &c. *monopetalous* or *polypetalous*, from one to many petals, or according to number.
- 557 REGULARIS, *regular, of an equal figure*, the size of all the parts proportioned to one another.
- 558 IRREGULARIS, *irregular*, when the limb and other parts are disproportionate.
- 559 INÆQUALIS, *unequal*, when the different sizes of the parts do not correspond in proportion to one another.
- 560 GLOBOSA, *globose, globe-shaped*.
- 561 CAMPANULATA, *campanulate, bell-shaped*.
- 562 INFUNDIBULIFORMIS, *funnel-shaped*.
- 563 ROTATA, *rotate, wheel-shaped*.
- 564 HYPOCRATERIFORMIS, *salver-shaped*.
- 565 RINGENS, *ringent, gaping*, irregular, with two lips.
 GALEA, *helmet*, the upper lip gaping.
 LABIUM, *lip*.
- 566 FAUX, *the jaws gaping between the divisions of the corollæ*, where the tube terminates.
- 567 PERSONATA (565), *personate, gaping*, but shut between the lips, with a palate.
- 568 CRUCIATA, *cruciform, cross-shaped*, having four equal spreading petals.
- 569 CONCAVA (248), *hollow*.
- 570 PATENS (134), *patent, spreading*.
- 571 PAPILIONACEA, *papilionaceous, butterfly-shaped*, irregular.
Carina, the keel, the lower petal often in form of a boat.
Vexillum, the standard, or upper petal ascending. *Alæ*, the wings, standing single on each side.
- 572 COMPOSITA, *compound flowers*, having many florets in a common perianthium, above the common receptacle.
- 573 LIGULATA, *ligulate, tongue-shaped*, florets whose limb is plane, and expanded outward.
- 574 TUBULOSA, *tubular*, florets that are all tubular and equal.
- 575 RADIATA, *radiate*, when the florets are tubular in the disk, and radiate and ligulate in the margin.
- 576 NECTARIUM, *nectary, honey-cell*, that part of the flower bearing honey.

- 577 PROPRIUM, *proper*, so called, as a distinct part from the petal.
- 578 PETALINUM, when inserted into the petal.
- 579 STAMEN, the male organ of generation furnished with a viscus, designed for the preparation of the pollen.
- 580 FILAMENTUM, *filament, threads*, the part that elevates, and is connected to the antheræ.
- 581 ÆQUALIA, *equal*, when they are all of an equal length.
- 582 INEQUALIA, *unequal*, when some are long, and others short.
- 583 CONNATA, *connate, when joined in one body*, but their number, figure, and insertion is expressed.
- 584 ANTHERA, *anther*, that part of the flower big with the pollen, which it throws forth when come to maturity.
- 585 DISTINCTÆ, *not cohering*.
- 586 CONNATÆ, jointed by the sides into one body.
- 587 POLLEN, *powder of the antheræ*, destined for the impregnation of the germen, and bursting in a viscous humour, into fine atoms, is, by a prolific breeze, scattered on the stigma.
- 588 PISTILLUM, a viscous humour adhering to the fruit for the reception of the pollen, and the female organ of generation.
- 589 GERMEN, the immature rudiment of the fruit within the flower.
- 590 SUPERUM, *above*, when included in the corollæ.
- 591 INFERUM, *beneath*, when below the corollæ.
- 592 STYLUS, *style*, that part of the pistillum which elevates the stigma from the germen.
- 593 STYGMA, the top of the pistil, furnished with a moist humour.
- 594 PERICARPIUM, *pericarp*, the germen of the plant big with the seeds, which it emits when mature.
- 595 CAPSULA, *capsule, a hollow pericarpium*, which cleaves or opens in some determinate manner.
- 596 VALVULA, *valve, an opening*, a part of a capsule or outer cover to the fruit.

- 597 LOCULAMENTUM, *a kind of arched cell*, for the lodgement of the seeds.
- 598 DISSEPIENTUM, *partitions of the fruit*, which divide the pericarpium into cells.
- 599 BICAPSULARIS, *two capsules*; TRICAPSULARIS, &c. *three capsules*, or according to the number.
- 600 BILOCULARIS, &c. *two cells*, &c. according to the number.
- 601 TRICOCCA, a capsule with three protuberant knobs, which divide into three cells.
- 602 DIDYMA, a capsule with two gibbous knobs, which divide into two cells.
- 603 SILIQUA, *a pericarpium of two valves*, in which the seeds are fixed alternately to the opposite sutures.
- 604 COMPRESSA, *flatted*, the opposite sides coming nearly together.
- 605 TORULOSA, *brawny protuberances*, when the pericarpium is bunched out by the seeds.
- 606 ARTICULATA, interrupted by arched joints.
- 607 PARALLELUM DISSEPIENTUM, the same width or diameter of the dissepiment to which the valves adhere.
- 608 TRANSVERSUM DISSEPIENTUM, dissepiments running cross-wise.
- 609 LEGUMEN, *a pericarpium of two valves*, the seeds fixed to one suture only.
- 610 ISTHMUS INTERCEPTUM, pods with various cross-divisions, forming distinct cells.
- 611 FOLLICULUS, *a pericarpium of one valve*, gaping lengthwise on one side, without the seeds being fixed to the suture.
- 612 DRUPA, *a pulpy pericarpium*, without valves, containing a stone or nut, 633.
- 613 SUCCULENTA, containing a pulpy humour.
- 614 SICCA, opposite the foregoing, dry.
- 615 POMUM, *an apple*, a fleshy pericarpium without valves, containing a capsule.
- 616 BACCA, *a berry*, a pulpy pericarpium without valves, containing naked seeds.
- 617 NIDULANTIA, seeds nestling in the pulp of a berry.

- 618 STROBILUS, *a pericarpium formed from an amentum*, with hard scales lying over each other, as in the pine tree.
- 619 SEMEN, *seed*, the rudiment of a new plant; are known according to the number, figure, superficies, and consistence.
- 620 HILUM, *the eye*, an external scar of the seed, where it has been fixed to the fruit or receptacle.
- 621 CORCULUM, the essence of a new plant within the seed.
- 622 PLUMULA, *part of the corculum*, the ascending scaly part of the plant.
- 623 ROSTELLUM, the descending part of the corculum that forms the root.
- 624 COTYLEDON, *the side lobes* of the seed of a porous substance, and perishing.
- 625 CORONA, *a crown*, a little cup adhering to the top of the seed, by which it flies.
- 626 PAPPUS, *a downy feathered cup*, adhering to the top of the seed, by which it flies.
- 627 STIPITATUS, *a kind of thread-like trunk*, elevating the down, and connecting it with the seeds.
- 628 CAPILLARIS, hairs undivided.
- 629 PLUMOSUS, having feathery hairs.
- 630 CAUDA, a thread terminating the seed:
- 631 HAMUS, a hooked seed adhering to animals.
- 632 ALA, *a membranaceous wing*, fixed to the seed.
- 633 NUX, *a nut*, a seed covered with a bony epidermis, having one, two, or more cells.
- 634 ARILLUS, the proper exterior coat of a seed that falls off spontaneously, and is either cartilaginous or succulent.
- 635 RECEPTACULUM, *the base*, by which the parts of fructification are connected.
- 636 COMMUNE, containing many flowers and fruit.
- 637 PUNCTATUM, a receptacle marked with hollow punctures.
- 638 PILOSUM (241), hairy.
- 639 PALEACEUM, chaffy scales which distinguish the florets.
- 640 PLANUM (246), *plain*, a flat surface.

- 641 CONVEXUM (249), the disk elevated.
- 642 CONICUM, *cone-shaped*, rounded and lessening towards the point.
- 643 SUBULATUM (269), awl-shaped.
- 644 COMPOSITUS-FLOS, *a compound flower*, with the receptacle spread out and entire, the florets sessile.
- 645 AGGREGATUS-FLOS, *an aggregate flower*, the receptacle enlarged, and the florets on little peduncles.
- 646 UMBELLA, *an umbel*, a receptacle which, from a common centre, runs out into thread-shaped foot-stalks of proportionate lengths.
- 647 SIMPLEX, when the foot-stalks proceed from one and the same centre of the receptacle.
- 648 COMPOSITA, when every foot-stalk of the general umbel produces a partial umbel.
- 649 UNIVERSALIS, composed of many simple umbels.
- 650 PARTIALIS, *a little umbel*, a part supported by the universal umbel.
- 651 PROLIFERA, an umbel more than decompound.
- 652 CYMA, a receptacle producing many foot-stalks from the same centre, that are of unequal lengths, the partial ones irregular on long fastigate peduncles.
- 653 RACHIS, *a thread-shaped receptacle*, the flowers adhering to it lengthwise, and forming a spike.
- 654 SPADIX, *a receptacle of a palm*, produced within a spatha or sheath, divided into branches that bear the fruit.
- 655 BULBUS, is an hybernacle placed on the descending caudex, and contains the rudiment of the plant and leaf that perishes.
- 656 SOLIDUS, *a solid fleshy bulb*, without any internal divisions.
- 657 TUNICATUS, bulbs having coats lying over each other, like the onion.
- 658 SQUAMATUS, bulbs consisting of imbricated scales, as in the lily.
- 659 CAULINUS, bulbs growing on the stalk of the plant.
- 660 GEMMA, *a bud*, is a hybernacle of the future plant with its leaves.

- 661 PETIOLARIS, enclosing the rudiment of the leaves.
- 662 STIPULARIS, enclosing the stipula.
- 663 CORTICALIS, consisting of cortical squamæ.
- 664 FOLIARIS, containing the leaf, and not the flowers.
- 665 FLORALIS, containing the flowers, and not the leaf.
- 666 COMMUNIS, containing both the leaf and the flowers.
- 667 VERNATIO, the position of the leaf within the bud.
- 668 CONDUPLICATA, when the parallel sides of a leaf approach.
- 669 CONVOLUTA, rolled together in a spiral form.
- 670 INVOLUTA, rolled inwards spirally from the lateral margins.
- 671 REVOLUTA, rolled spirally backwards from the lateral margins.
- 672 OEVOLUTA, rolled together, one margin embracing the other alternately.
- 673 EQUITANTIA, when the sides of the leaves lie parallel, the outward one embracing the inner one.
- 674 IMBRICATA, a parallel straight surface, lying over each other.
- 675 PLICATA, *plaited*, when their complication is in plaits lengthwise.
- 676 RECLINATA, *reclined*, reflexed downwards towards the petiole.
- 677 SPIRALIA, *spiral*, twisted in transverse plaits, so that the apex becomes the centre.
- 678 ÆSTIVATIO, *the complication of the corollæ*, before the unfolding of the flower.
- 679 CONVOLUTA, *rolled together*, 669.
- 680 IMBRICATA (674), *imbricate*.
- 681 CONDUPLICATA (668), when the parallel sides of the leaf approach.
- 682 VALVATA, having valves.
- 683 INÆQUIVALVIS, with unequal valves.
- 684 SOMNUS, *sleep*, the change that leaves of plants undergo in the night.
- 685 CONNIVENS, when the upper disk of two opposite leaves or folioles are pressed together so as to appear one leaf.

- 686 INCLUDENS, when the leaves are alternate, and in the night press against the stalk, so as to include it.
- 687 CIRCUMSEPIENS, when leaves growing in a horizontal position, erect themselves in the night, by clasping together in the form of a funnel.
- 688 MUNIENS, when the leaves have foot-stalks spreading horizontally, become dependent, in form of a hollow arch.
- 689 CONDUPLICANS, *doubling*, when the folioles lightly approach each other with their upper disk, so that both are covered.
- 690 INVOLVENS, when the points of the upright folioles are pressed together, and form a cavity between.
- 691 DIVERGENS, when the base of the folioles approach, and the points are spreading.
- 692 DEPENDENS, when the folioles hang downwards.
- 693 INVERTENS, when the folioles hang down, and are at the same time inverted.
- 694 IMBRICANS, the folioles imbricated, 120.

MENSURA, THEIR MEASURE.

- 695 LINEARIS, *linear*, the twelfth part of an inch.
- 696 UNGUICULARIS, the length of a nail.
- 697 POLICARIS, the length of the outward joint of the thumb.
- 698 PALMARIS, the width of the hand.
- 699 SPITHAMÆUS, *a span*, the length between the point of the thumb and fore finger.
- 700 DODRANTALIS, nine inches, the space between the point of the thumb and little finger, when extended.
- 701 PEDALIS, *a foot*, the space from the bending of the elbow to the base of the thumb.
- 702 ORGYALIS, *a fathom*, or six feet, the height of a man, or the space between the extreme points of the fingers, when the arms are extended.

TABLE VII.



A

BOTANICAL DICTIONARY*.



A

ABBREVIATUM PERIANTHIUM, *shortened*, when the cup is shorter than the tube of the flower.

ABORTIENS FLOS, *barren flowers*, such as produce no fruit.

ABRUPTUM FOLIUM PINNATUM, *winged leaves*, ending without either foliole or cirrus.

ACAULIS, *without stalk or stem*.

ACEROSUM FOLIUM, *chaffy leaves*, when they are linear and abiding, as in PINUS, ABIES, and JUNIPERUS.

ACICULARIS, *needle-shaped*, as in SCIRPUS ACICULARIS.

ACINACIFORME, *fulchion or scimitar-shaped*, as in MESEMBRYANTHEMUM ACINACIFORME.

ACINI, the *small berries* which compose the fruit of a mulberry or bramble.

ACOTYLEDONES, plants whose seeds have *no cotyledons*, or seminal leaves.

ACULEI, *prickles* fixed in the rind or surface of the bark.

* The reader who may wish for fuller information on this part of the science, should consult Dr. Colin Milne's Botanical Dictionary, third edition, lately published.

- ACULEATUS CAULIS, a stalk or stem furnished with prickles.
- ACUMINATUM FOLIUM, a leaf ending in a point.
- ACUTUM FOLIUM, leaves terminating in an acute angle.
- ADNATUM FOLIUM, the disk of the leaf pressing close to the stem of the plant.
- ADPRESSA FOLIA, the disk of the leaf pressed towards the stem.
- ADSCENDENS CAULIS, a stalk or branch inclining upwards.
- ADVERSUM FOLIUM, when the sides of the leaf are turned towards the south.
- AGGREGATUS FLOS, an assemblage of flowers coming in clusters.
- AGGREGATÆ, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- ALA, a wing, the side petals of a papilionaceous blossom, or a membrane added to a seed, stalk, &c.
- ALATUS PETIOLUS, when the foot-stalk of a leaf is winged with membranes.
- ALBURNUM, the white substance that lies between the inner bark and the wood of trees.
- ALGÆ, flags, one of the seven families of plants.
- ALTERNI RAMI FOLIA, when they come out singly, and follow in gradual order.
- AMENTACEÆ, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*, bearing catkins.
- AMENTUM, a catkin.
- AMPLEXICAULE FOLIUM, embracing the stalk when the base of the leaf embraces the stem sideways.
- ANCEPS CAULIS, double-edged, when a stalk is compressed, and forms two opposite acute angles.
- ANDROGYNA, plants bearing male and female flowers on the same root.
- ANGULATUS CAULIS, angulated stalks.
- ANGUSTIFOLIA, narrow-leaved.
- ANGIOSPERMIA, the second order in the class *Didynamia* of *Linnaeus*; containing plants whose seeds are covered with a capsule.
- ANNUA RADIX, an annual root; that which lives but one year.

- ANTHERA, the summit of the stamina bearing the pollen, and is a part of the principal male organ of generation.
- APERTURA, *an aperture*, opening in some species of anthera.
- APETALUS FLOS, having no petals or corolla.
- APEX, *the top*, or summit.
- APHYLLUS CAULIS, *destitute of leaves*.
- APOPHYSIS, an excrescence from the receptacle of the musci.
- APPENDICULATUS PETIOLUS, a little appendage hanging from the extremity of the foot-stalk.
- APROXIMATA FOLIA, leaves growing near each other.
- ARBOR, *a tree*.
- ARBUSTIVA, *a copse of shrubs or trees*, an order of plants in the *Fragmenta Methodi Naturalis of Linnæus*.
- ARCUATUM LEGUMEN, *arched*, a pod that is curved or bent.
- ARILLUS, the proper exterior coat of a seed that falls off spontaneously.
- ARISTA, the beard of corn or grasses.
- ARMA, *arms, weapons*, one of the seven kinds of fulcra of plants.
- ARTICULATUS CAULIS, *culmus*, having knots or joints.
- ARTICULUS CULMI, the straight part of the stalk between the two joints.
- ASPERIFOLIA, *rough-leaved plants*, an order of plants in the *Fragmenta Methodi Naturalis of Linnæus*.
- ASSURGENTIA FOLIA, first bent down, but rising erect towards the apex.
- ATTENUATUS PEDUNCULUS, when the foot-stalk grows smaller towards the flower.
- AUCTUS CALYX, *augmented*, having a series of distinct leaves, shorter than its own, that surround its base.
- AVENIA FOLIA, leaves which have no visible veins.
- AURICULATUM FOLIUM, *an ear-shaped leaf*, when the leaf towards the base has a lobe on each side.
- AXILLARIA FOLIA, growing out of the angles formed by the branches and the stem.

B

- BACCA, *a berry*, or a pulpy pericarpium without valves, in which the seeds are naked.
- BARBA, *a beard*, a species of pubescence, sometimes on the leaves of plants, as on the MESEMERYANTHEMUM BARBATUM.
- BARBATUM FOLIUM, when a bunch of strong hairs terminate the leaves.
- BICORNES, plants whose antheræ have the appearance of two horns. Likewise an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- BIENNIS RADIX, a root which continues to vegetate two years.
- BIFARIA FOLIA, a leaf pointing two ways.
- BIFERÆ PLANTÆ, flowering twice a year.
- BIFIDUM FOLIUM, divided or cloven into two parts.
- BIFLORUS PEDUNCULUS, bearing two flowers on a foot-stalk.
- BIGEMINUM FOLIUM, *a forked foot-stalk*, with two little leaves on the apex of each division.
- BIJUGUM FOLIUM, *a winged leaf*, bearing two pair of foliola.
- BILABIATA COROLLA, a corolla with two lips.
- BILOBUM FOLIUM, a leaf consisting of two lobes.
- BINATA FOLIA, *a digitate leaf*, consisting of two foliola.
- BIPARTITUM FOLIUM, a leaf divided into two segments.
- BIPINNATUM FOLIUM, *doubly winged*, when the folioles of a pinnate leaf are pinnate.
- BITERNATUM FOLIUM, when there are three folioles on a petiole, and each foliole is ternate, as in EPIMEDIUM.
- BIVALVE PERICARPIUM, *consisting of two valves*, as in the SILIQUA and LEGUMEN.
- BRACHIATUS CAULIS, *branching in pairs*; each pair standing at right angles with those above and below.
- BRACHIUM, *the arm*, tenth degree in the *Linnaean Scale* for measuring plants, being twenty-four Parisian inches.
- BRACTEA, *a floral leaf*, these are generally of a different shape and colour from the other leaves of the plant, and are always seated near the fructification.
- BRACTEATUS, having a bractea growing out of it.

- BULBIFERUS CAULIS, *a stalk-bearing bulb*, as in a species called LILIUM BULBIFERUM.
- BULBOSA RADIX, *a bulbous root*, and is either *squamosa*, scaly, as in LILIUM; *tunicata*, coated, as in CEPÆ; *duplicata*, double, as in FRITILLARIA; or *solida*, as in TULIPA.
- BULLATUM FOLIUM, when the surface of the leaf rises above veins, so as to appear like blisters.

C

- CADUCUS CALYX, *to fall off*; a term signifying the shortest time of duration, falling off at the first opening of the flower.
- CALAMARIÆ, *a reed*, an order of plants in the Fragmenta Methodi Naturalis of *Linnæus*.
- CALCARIATUM NECTARIUM, a kind of nectarium resembling a spur, as in the DELPHINIUM.
- CALICULATUS CALYX, a little calyx added to a larger one, as in the COREOPSIS, LEONTICE, &c.
- CALYCANTHEMI, *a calyx*, an order of plants in the Fragmenta Methodi Naturalis of *Linnæus*.
- CALYPTRA, *a veil*, in mosses, where it is placed over the antheræ.
- CALYX, a flower cup, of which there are the following kinds, viz. PERIANTHIUM, ... INVOLUCRUM, ... AMENTUM, ... SPATHA, ... GLUMA, ... CALYPTRA, ... and VOLVA.
- CAMPANACEI, an order of plants in the Fragmenta Methodi Naturalis of *Linnæus*.
- CAMPANULATA COROLLA, bell-shaped flowers.
- CANALICULATUM FOLIUM, leaves having a deep channel running from the base to the apex.
- CANDELARES, an order of plants in the Fragmenta Methodi Naturalis of *Linnæus*.
- CAPILLACEUM FOLIUM, *capillary*, exemplified in the RANUNCULUS AQUATILIS.
- CAPILLARIS PAPPUS, *hairy down*, as in HIERACIUM and SONCHUS.
- CAPILLUS, *hair*, the first degree of the *Linnæan* Scale for measuring plants, the diameter of a hair, and the twelfth part of a line.

- CAPITATI FLORES, *flowers collected into heads*, as in MENTHA AQUATICA and THYMUS SERPYLLUM.
- CAPITULUM, *a little head*, a species of Inflorescentia, in which the flowers are connected into close heads on the tops of the peduncles, as in GOMPHRENA.
- CAPREOLUS, *a tendril*, see CIRRHUS.
- CAPSULA, *a capsule*, a hollow pericarpium, which cleaves or parts in some determinate manner, and consists of VALVULA,...DISSEPIENTUM, ...COLUMELLA,... and LOCULAMENTUM.
- CARINA, *the keel of a boat or ship*, the lower petal of the papilionaceous corolla.
- CARINATUM FOLIUM, when the back of a leaf resembles the keel of a ship.
- CARIOPHYLLÆUS FLOS, *clove-tree*, or flowers growing in the manner of carnations.
- CARNOSUM FOLIUM, *a fleshy leaf*, as in SEDUM DASYPHILLUM.
- CARTILAGINEUM FOLIUM, a leaf, whose brim is furnished with a margin of different substance from the disk.
- CARYOPHYLLÆI, *carnations or pinks*, an order of plants in the Fragmenta Methodi Naturalis of Linnæus.
- CATENULATA SCABRITIES, *species of glandular roughness*, hardly visible to the naked eye, resembling little chains on the surface of some plants.
- CAUDEX, the stem of a tree.
- CAULESCENS, having a stalk or stem.
- CAULINA FOLIA, leaves growing immediately on the stem.
- CAULIS, a stem, a species of Truncus.
- CERNUUS, nodding, or hanging down its head.
- CESPITOSA, plants which produce many stems from one root, and form a surface of turf or sod.
- CILIATUM, whose margin is guarded by parallel bristles, formed like the eye-lash.
- CIRCINALEA FOLIA, *a hoop or ring*, a term of foliation, expressive of the leaves within the gemma, being rolled spirally downward.

- CIRCUMSCISSA CAPSULA, *cut transversely*, as in ANAGALLIS.
- CIRRHIFERUS PEDUNCULUS, a peduncle bearing a tendril, as in VITIS.
- CIRRHOSUM FOLIUM, a leaf that terminates in a tendril, as in GLORIOSA.
- CIRRHUS, *a clasper*, or tendril, one of the fulcra of plants.
- CLASSIS, *a class*, is defined by *Linnæus*, to be an agreement of several genera in the parts of fructification, according to the principles of nature distinguished by art.
- CLAVATUS PETIOLUS, PEDUNCULUS, when the foot-stalk of the leaf or flower is club-shaped, tapering from the base to its apex.
- CLAVICULA, *a little key*, a tendril.
- CLAUSA COROLLA, when the neck of the corolla is close shut in with valves.
- COADUNATÆ, *to gather together*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnæus*.
- COARCTATI RAMI, *close together*, opposed to DIVARICATUS.
- COCHLEATUM LEGUMEN, a pod like the shell of a snail, as in MEDICAGO.
- COLORATUM FOLIUM, *coloured*, when leaves which are generally green, are of a different colour.
- COLUMELLA, *a little column*, the substance that passes through the capsule, and connects the several partitions and seeds.
- COLUMNIFERI, *pillar-shaped*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnæus*.
- COMA, *a bush*, or head of hair, a species of Fulcra, composed of large bractea, which terminates the stalk, as in LAVANDULA, SALVIA, &c.
- COMMUNIS GEMMA, regards the contents of the gemma, containing both flower and fruit.
- COMMUNIS CALYX, when a cup contains both receptacle and flower.
- COMOSÆ, *a head of hair*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnæus*.
- COMOSA RADIX, the fibres which put forth at the base of a bulbous root, resembling a head of hair.

- COMPACTUM FOLIUM, when the leaf is of a compact and solid substance.
- COMPLETUS FLOS, having a perianthium and corolla.
- COMPOSITUS CAULIS, a *compound stem*, diminishing as they ascend.
- COMPOSITUM FOLIUM, when the petiole bears more than one leaf, of which are the following species, viz. ARTICULATUM, ... DIGITATUM, ... CONJUGATUM, ... PEDATUM, ... PINNATUM, ... DECOMPOSITUM, ... SUPRA-DECOMPOSITUM.
- COMPOSITI, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- COMPRESSUS CAULIS, FOLIUM, a leaf resembling a cylinder compressed on the opposite sides.
- CONCAVUM FOLIUM, *hollowed*, the margin forms an arch with the disk.
- CONCEPTACULUM, *conceptacle or receiver*, a pericarpium of a single valve, which opens on the side lengthways, and has not the seeds fastened to it.
- CONDUBLICATUM FOLIUM, *doubled together*, when the sides of the leaf are parallel, and approach each other.
- CONFERTI RAMI, branches crowded together.
- CONFERTUS VERTICILLUS, FLOS, ET FOLIA, when flowers and leaves are formed into whorls round the stalk, and crowded together.
- CONFLUENTIA FOLIA, *to flow together*, as in the pinnated leaf, when the pinnæ run into one another.
- CONGLOBATUS FLOS, when flowers are collected into globular heads.
- CONGLOMERATUS FLOS, flowers irregularly crowded together.
- CONGESTA UMBELLA, flowers collected into a spherical shape, as in the ALLIUM.
- CONICA SCABRITIES, a species of setaceous scabrities, scarce visible to the naked eye, on the surface of plants, formed like cones.
- CONIFERÆ, *plants bearing cones*, such as PINUS, ... CUPRESSUS, &c. an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- CONJUGATUM, *to join or couple together*, a species of pinnate leaf, where the folioles come by pairs.

- CONNATUM, *to grow together*, when two opposite leaves unite at their base, so as to have the appearance of one leaf.
- CONNIVENS COROLLA, when the apices of the petals converge, so as to close the flower, as in TROLLIUS EUROPEUS.
- CONNIVENTES ANTHERÆ, approaching or inclining together.
- CONTINUATUM FOLIUM, *continued*, when the leaf appears to be a continuation of the substance of the stalk.
- CONTORTI, *to twist*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- CONTRARIÆ VALVULÆ, valves are termed contraria, when the dissepimentum is placed transversely between them.
- CONVEXUM FOLIUM, a leaf rising from the margin to the centre of the leaf.
- CONVOLUTUS CIRRHUS, a tendril twining with the same direction with the sun's motion.
- CONVOLUTUM FOLIUM, *a term in foliation*, when the leaf is rolled up like a scroll of paper.
- CONUS. See STROBILUS.
- CORCULUM, the heart and essence of the seed.
- CORDATUM FOLIUM, the heart-shaped leaf.
- CORDIFORMUS, shaped like a heart.
- COROLLA, *a wreath or crown*, one of the seven parts of fructification.
- COROLLULA, *a little corolla*.
- CORONA SEMINIS, a crown adhering to many kinds of seeds serving them as wings, which enables them to disperse.
- CORONARIÆ, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*,
- CORONULA, *a little crown*.
- CORTEX, the outer rind or bark of vegetables.
- CORYDALES, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- CORYMBUS is a kind of spike, the flowers of which have each its proper pedicellus, or partial foot-stalk, raised to a proportional height, as in SPIREA OPULIFOLIA.

- COTYLEDON, *a side-lobe of the seed*, of a porous substance, and perishable, or seminal leaves.
- CRENATUM FOLIUM, *a notched leaf*, when the margin is cut into angles that point towards neither of the extremities, obtusely crenate, when the angles are rounded, or acutely crenate, when the angles are pointed.
- CRISPUM FOLIUM, *a curved leaf*, when the circumference becomes larger than the disk admits of.
- CRISTATUS FLOS, when the flower has a tufted crest, as in POLYGALA.
- CRUCIFORMES FLORES, *cross-shaped flowers*, consisting of four petals, disposed in the form of a cross, as in the class Tetradynamia of *Linnaeus*.
- CRYPTOGAMIA, *hidden marriages*, the twenty-fourth class of the *Linnaean* System.
- CUBITUS, *a cubit*, the ninth degree of the *Linnaean* Scale for measuring plants, from the elbow to the extremity of the middle finger.
- CUCULLATUM FOLIUM, *leaves rolled up lengthways*, in form of a cone, as in GERANIUM CUCULLATUM, &c.
- CUCURBITACEÆ, *gourds*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- CULMINÆ, *the top or crown of any thing*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- CULMUS, *a reed or straw*, the proper stem or trunk of a grass.
- CUSPIDATUM FOLIUM, a leaf, whose apex resembles the point of a spear or lance.
- CUNEIFORME FOLIUM, a wedge-shaped leaf.
- CYATHIFORMIS COROLLA, flowers of the form of a cup.
- CYLINDRACEA SPICA, a spike of flowers in form of a cylinder.
- CYMA, that runs into long fastigate peduncles, proceeding from the same universal centre, but with irregular partial ones.
- CYMOSES FLOS. See CYMA.
- CYMOSEÆ, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.

D

- DÆDALEUM FOLIUM**, a leaf whose texture is remarkably beautiful, and exquisitely wrought.
- DEBILIS CAULIS**, a weak, feeble stalk.
- DECAGYNIA**, *ten females*, the fifth order in the tenth class: flowers that have ten styli.
- DECANDRIA**, *ten males*, the tenth class of *Linnæus*.
- DECAPHYLLUS CALYX**, a calyx consisting of ten leaves.
- DECIDUUM FOLIUM**, leaves that fall off in winter.
- DECLINATUS CAULIS**, a stalk bending towards the earth.
- DECOMPOSITA FOLIA**, when a petiole, once divided, connects many folioles.
- DECUMBENS**, to lie down.
- DECURRENS FOLIUM**, running down, when the base of a sessile leaf extends itself downwards along the stem, beyond the proper base or termination of the leaf.
- DECURSIVE, FOLIUM PINNATUM**, when the bases of the foliole are continued along the sides of the petiolus.
- DECUSSATA FOLIA**, *to divide*, when leaves grow in pairs, and opposite, each pair being opposite alternately.
- DEFLEXUS RAMUS**, a branch bent a little downwards.
- DEFLORATA STAMINA**, having shed or discharged the farina fecundans.
- DEFOLIATIO**, the time in Autumn when plants shed their leaves.
- DELTOIDES FOLIUM**, a leaf formed like the Greek delta, as in **MESEMERYANTHEMUM DELTOIDES**.
- DEMERSUM FOLIUM**, in aquatic plants, leaves sunk below the surface of the water.
- DENTROIDES SURCULUS**, shrub-like, a subdivision of the surculus in the genus Hypnum.
- DENTATUM FOLIUM**, leaves having horizontal points of the same consistence of the leaf, and standing at a little distance from each other.
- DENUDATE**, *stripped naked*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnæus*.

- DEPENDENS FOLIUM, *hanging down*, leaves pointing towards the ground.
- DEPRESSUM FOLIUM, *pressed down*, when the sides rise higher than the disk.
- DIADELPHIA, *two brotherhoods*, the seventeenth class in the Sexual System.
- DIANDRIA, *two males*, the second class in the Sexual System.
- DICHOTOMUS CAULIS, *forked stalks*, when the divisions come by two and two.
- DICOTYLEDONES, when the seeds have two cotyledons that are the placenta of the embryo plant, and afterwards the seed leaves.
- DIDYMA ANTHERA, *twins*, when anthera come by twos on each filament.
- DIDYNAMIA, *the superiority of two*, the fourteenth class in the Sexual System.
- DIFFORMIA FOLIA, *different forms*, when leaves on the same plant come of different forms.
- DIFFUSUS CAULIS, when the branches of the stalk spread different ways.
- DIGITATUM FOLIUM, *fingered*, when the apex of a petiole connects many folioles.
- DIGYNIA, *two females*, the second order in each of the first thirteen classes, except the ninth.
- DIMIDIATUM, *halved*.
- DIÆCIA, the twenty-second class in the Sexual System.
- DIPETALA COROLLA, flowers consisting of two petals, as in CIRCÆA, and COMMELINA.
- DIPHYLLUS CALYX, a calyx consisting of two leaves, as in the PAPAVER and FUMARIA.
- DISCUS, *a disk*, the middle part of a radiate compound flower.
- DISPERMA, plants producing their seeds by twos, as in the umbellatæ.
- DISSECTUM FOLIUM, leaves cut into lacinia, or divisions.
- DISSEPIMENTUM, *partitions of the fruit*, which divide the pericarpium into cells.

- DISSILIENS SILIQUA, pods that burst with elasticity.
- DISTANS VERTICILLUS, when the whorls of flowers, in verticillate plants, stand at a great distance from one another.
- DISTICHA FOLIA, *in two rows*, when leaves all respect two sides of the branches only.
- DIVARICATI RAMI, branches standing wide from each other, in different directions.
- DIVERGENTES RAMI, widening gradually.
- DODECANDRIA, *twelve males*, the eleventh class in the Sexual System.
- DODRANS, *the seventh degree* in the *Linnæan* Scale for measuring the parts of plants, or nine inches.
- DODRANTALIS, *nine inches*.
- DOLABRIFORME FOLIUM, *a leaf resembling an axe*, as in MESENBRY-ANTHEMUM DOLABRIFORME.
- DORSALIS ARISTA, *an awne or beard*, fixed to the back or external part of the gluma.
- DRUPA, *a pulpy pericarpium*, without valves, containing a stone, as in the plum and peach.
- DRUPACEÆ, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- DUMOSÆ, *a bush*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- DUPLICATA RADIX, *a double root*, a species of bulbous root, consisting of two solid bulbs, as in some species of ORCHIS.
- DUPLICATO-SERRATUM FOLIUM, *sawed double*, with lesser teeth within the greater.

E

- EBRACTEATUS RACEMUS, *without a bractea or floral leaf*.
- ECAUDATA COROLLA, *without a tail or spur*, as in ANTIRRHINUM, ...CYMBALARIA.
- ECHINATUM PERICARPIUM, *Pods beset with prickles*, like a hedgehog.
- EFFLORESCENTIA, the precise time when a plant shows its first flowers.

- EMARGINATUM FOLIUM, when the apex of a leaf terminates in a notch: the same may be applied to PETALA and STIGMA.
- ENERVIUM FOLIUM, leaves having no apparent nerves.
- ENNEANDRIA, *nine males*, the ninth class in the Sexual System.
- ENNEAPETALA COROLLA, a flower consisting of nine petals.
- ENODIS CAULIS, CULMUS, stalks and straws, having no knots or joints.
- ENSATÆ, plants having sword-shaped leaves, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- ENSIFORME FOLIUM, leaves shaped like a two-edged sword, tapering towards the point.
- EQUITANTIA FOLIA, *riding*, when the sides of the leaves approach in such a manner as the outer embrace the inner.
- ERECTUS CAULIS, RAMUS, FOLIUM, upright, perpendicular.
- EROSUM FOLIUM, *gnawed*, when the leaf is sinuate, and the margin appears as if it were gnawed or bitten.
- EXSERTA STAMINA, *standing forth*, when the stamina appear above the corolla.
- EXSTIPULATUS, *without stipulæ*.
- EXSUCCUM FOLIUM, when the substance of the leaf is dry.
- EXTRAFOLIACÆ STIPULÆ, stipula growing on the outside of the leaves.

F

- FARCTUM FOLIUM, *stuffed*, opposed to TUBULOSUM.
- FASCICULATA, *bundled*, leaves growing in bunches.
- FASCICULARIS RADIX, *bundled*, tuberous roots growing in bundles.
- FASCIATA PLANTA, when many stalks grow together, like a faggot or bundle.
- FASTIGIATI PEDUNCULI, pedunculi pointed at the apex.
- FAUCES, the jaws or chops.
- FEMINA PLANTA, a plant bearing female flowers on the same root only.
- FIBROSA RADIX, a fibrous root.
- FILAMENTUM, *a thread*, applied to the thread-like part of the stamina.

- FILICES, *ferns*, one of the seven divisions of the vegetable kingdom, and an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- FILIFORM FILAMENTUM, thread-shaped stamina.
- FIMBRIATA PETALA, a fringed petal, as in *MENYANTHUS*.
- FISSUM FOLIUM, a leaf split or cloven half way down.
- FISTULOSUS CAULIS, a piped or hollow stem.
- FLABELLATUM FOLIUM, a fan-shaped leaf.
- FLACCIDUS PEDUNCULUS, the foot-stalk of a flower that is feeble and slender.
- FLAGELLUM, a *twig* or shoot like a whip or thong.
- FLEXUOSUS CAULIS, a stalk, having many turnings or bendings, taking a different direction at every joint.
- FLORALIA FOLIA, floral leaves that immediately attend the flower.
- FLORALIS GEMMA, flower-buds.
- FLOS, a flower.
- FLOSCULUS, a little flower.
- FOLIACEÆ GLANDULÆ, glands growing on the leaves.
- FOLIARIS CIRRHUS, a tendril growing from a leaf.
- FOLIARIS GEMMATIO, leaf-buds.
- FOLIATIO PLANTÆ, the complication of the leaves, whilst folded within the gemma, or bud.
- FOLIATUS CAULIS, a leafy stalk.
- FOLIFERA GEMMA, a bud producing leaves.
- FOLIOLUM, a *little leaf*, one of the single leaves, which together constitute a compound leaf.
- FOLIOSUM CAPITULUM, covered with leaves amongst the flowers or tops of the plant.
- FOLIUM, a leaf.
- FORNICATUM PETALUM, *vaulted or arched*, as in the upper lip of the flowers in the class *Didynamia*.
- FREQUENS PLANTA, plants growing frequently, or commonly, every where.
- FRONDESCENTIA, the season of the year when the leaves of plants are unfolded.
- FRONDOSUS CORDEX, a species of trunk composed of a branch and a leaf blended together, as is frequently united with the fructification.

- FRUCTESCENTIA, the time of the year when a plant scatters its ripe seeds.
- FRUCTIFICATIO, the temporary part of a vegetable appropriated to generation, terminating the old vegetable, and beginning the new.
- FRUSTRANEA POLYGAMIA, *to no purpose*, the third order of the class Syngnesia.
- FRUTEX, a shrub.
- FRUTICOSUS CAULIS, a shrubby stalk.
- FUGACISSIMA PETALA, petals that are fleeting, and of short duration.
- FULCRATUS CAULIS, branches having props. See FULCRUM.
- FULCRUM, a prop, or support.
- FUNGI, *a kind of mushroom*, one of the seven families of plants, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnæus*.
- FURCATA, *forked*.
- FUSIFORM RADIX, a spindle-shaped root.

G

- GALEA, *a helmet*, applied to the corolla of the class Gynandria, as in ORCHIS.
- GALEATUM LABIUM, the lip of a flower shaped like a helmet.
- GEMINÆ STIPULÆ, stipula growing in pairs.
- GEMINATUS PEDUNCULUS, double foot-stalks growing from the same point.
- GEMMA, *a bud*, an hybernaculum on the ascending caudex.
- GEMNATIO, a young bud.
- GEMNIPARUS, bearing buds.
- GENERA PLANTARUM, *genera of plants*, the second subdivision in the *Linnaean System*; it comprehends an assemblage of species, similar in their parts of fructification, under the same class and order.
- GENICULATUS CAULIS, CULMUS, PEDUNCULUS, *a jointed stalk*, straw, or foot-stalk of a flower.
- GENICULA, *little joints*.

- GERMEN**, a sprout or bud, the base of the pistillum, the rudiment of the fruit yet in embryo.
- GIBBUM FOLIUM**, *bunching-out*, or gouty.
- GLABER**, *smooth*, having an even surface.
- GLADIATA SILIQUA**, a sword-shaped pod.
- GLANDULÆ**, a gland, or secretory vessel.
- GLANDULIFERA SCABRITIES**, a kind of bristly roughness on the surface of some plants, on which there are minute glands at the extremity of each bristle.
- GLAREOSIS LOCIS**, *gravelly places*, where plants delight in gravel.
- GLAUCOPHYLLUS**, a blueish or azure-coloured leaf.
- GLOBOSA RADIX**, a round root.
- GLOBULARIS SCABRITIES**, a species of glandular roughness, scarce visible to the naked eye, the small grains of which are exactly globular.
- GLOCHOIDES**, the small points of the pubes of plants. *Linneus* applies this term only to the **HAMI TRIGLOCHOIDES**, with three hooked points.
- GLOMERATA SPICA**, flowers crowded together in a globular form.
- GLUMA**, a husk or chaff, a species of calyx peculiar to corn and grasses.
- GLUTINOSITAS**, like glue or paste.
- GRAMINA**, *grasses*, one of the seven families of the vegetable kingdom.
- GRANULATA RADIX**, roots consisting of many little knobs, like seeds of grain, attached to one another by small strings, as in **SAXIFRAGA GRANULATA**.
- GYMNOSPERMA**, *naked seeded*, the first order of the class *Didynamia*.
- GYNANDRIA**, when the male and female parts are joined together; the twentieth class in the *Linnean* System.

H

- HABITUALIS CHARACTER**, the character or description of a plant, taken from its habit, which consists in the *Placentatio*, *Radicatio*, *Ramificatio*, *Foliatio*, *Stipulatio*, *Pubescentia*, *Inflorescentia*.

- HABITUS**, *the external appearance*: *Linnaeus* defines it, the conformity or affinity that the congeners of vegetables have to one another, in placentation, radification, &c.
- HAMOSIA SETA**, *hooked bristles*.
- HASTATUM FOLIUM**, leaves resembling the head of a spear or halbert.
- HEMISPHERICUS CALYX**, *half round*, or half a sphere.
- HEPTANDRIA**, *seven males*, the seventh class of the Sexual System.
- HERBA**, *an herb*: according to *Linnaeus*, it is the part of the vegetable which arises from the root; it is terminated by the fructification, and comprehends the stem, leaf, props, and hybernacula.
- HERBACEÆ PLANTÆ**, are perennial plants, which annually perish down to the root.
- HERBACEUS CAULIS**, stalks that dry annually.
- HERMAPHRODITUS FLOS**, flowers that contain both sexes, as anthera and stigma.
- HESPERIDÆ**, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- HEXAGONUS CAULIS**, a stalk with six angles.
- HEXANDRIA**, *the sixth class*, in the Sexual System, which produce hermaphrodite flowers, with six stamina of equal length.
- HEXAGYNIA**, an order of plants that produce six styles.
- HEXAPETALA COROLLA**, flowers consisting of six petals.
- HEXAPHYLLIS CALYX**, a flower-cup, consisting of six leaves.
- HIANS COROLLA**, a monopetalous flower that is gaping.
- HIRSUTUS**, *rough*, hairy.
- HISPIDUS CAULIS**, a stalk covered with strong fragile bristles.
- HOLERACEÆ**, *pot herbs*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- HORIZONTALIS FLOS**, flowers growing with their disk parallel to the horizon.
- HYBERNACULUM**, *winter-lodge*, the part of a plant that encloses and secures the embryo from external injuries.
- HYBRIDA**, *a bastard*, a monstrous production of two plants of different species, like the mule in the animal creation.

HYPOCRATERIFORMIS COROLLA, a monopetalous flower, shaped like a cup or salver.

I

ICOSANDRIA, *the twelfth class* in the Sexual System.

IMBERBIS COROLLA, a flower without a beard.

IMBRICATES, *tiled*, when the scales of a stalk, or flower-cup, lie over one another in the manner of tiles upon a house.

IMMUTATÆ, *unaltered*.

IMPAR, *odd*, applied to a pinnated leaf terminating in an odd lobe.

INEQUALIS COROLLA, an unequal flower.

INANIS CAULIS, hollow or empty stalks.

INCANUM FOLIUM, leaves covered with whitish down.

INCISUM FOLIUM, leaves cut into irregular segments.

INCOMPLETUS FLOS, imperfect flowers without petals.

INCRASSATUS PEDUNCULUS, foot-stalks of flowers that increase in thickness as they approach the flowers.

INCUMBENS ANTHERA, anthera which are affixed to the filament sideways.

INCURVATUS CAULIS, a stalk bowed towards the earth.

INDIVISUM FOLIUM, an entire undivided leaf.

INERME FOLIUM, *unarmed*, a leaf without bristles or prickles.

INFERUS FLOS, flowers whose receptacle are situated below the germen.

INFLATUM PERIANTHIUM, a calyx puffed out like a bladder.

INFLEXA FOLIA, to bend inwards towards the stem.

INFLORESCENTIA, *inflorescence*, signifies the various modes in which flowers are joined to the plant by the pedunculus.

INFUNDIBULIFORMIS COROLLA, a monopetalous flower, shaped like a funnel.

INSERTUS PETIOLUS, a foot-stalk inserted into the stem.

INTEGRUM FOLIUM, an entire or undivided leaf.

INTEGERRIMUM FOLIUM, *an entire leaf*, whose margin is destitute of incisions or serratures.

INTERFOLIACEUS PEDUNCULUS, flower-stalks arising from between opposite leaves.

- INTERRUPTUM FOLIUM PINNATUM, when the large folioles of a winged leaf are interrupted alternately by pairs of smaller ones.
- INTERRUPTA SPICA, a spike of flowers, interrupted or broken by small clusters of flowers between the larger ones.
- INTORSIO, writhing or twisting.
- INTRAFOLIACEÆ STIPULÆ, stipulæ growing on the inside of the leaves of the plant.
- INUNDATA LOCA, this term is applied by *Linnaeus* to such places that are overflowed only in winter.
- INVOLUCELLUM, a partial involucre.
- INVOLUCRUM, a cover, the calyx of the umbelliferous plants standing at a distance from the flower.
- INVOLUTA FOLIA, rolled in leaves, when their lateral margins are rolled spirally inwards on both sides.
- IRREGULARIS FLOS, irregular flowers of deformed shapes.
- JUBA, a crest of feathers.
- JULUS, a catkin.

L

- LABIATUS FLOS, a lipped flower.
- LACERUM FOLIUM, a cleft or fissure; leaves whose margin is cut into segments, as if rent or torn.
- LACINIÆ, segments or incisions.
- LACINIATUM FOLIUM, a leaf cut into irregular incisions.
- LACTESCENTIA, milky; those plants are called milky, whose juices are white, yellow, or red.
- LACUNOSUM FOLIUM, leaves that are deeply furrowed, by the veins being sunk below the surface.
- LACUSTRIS PLANTA, plants which grow in lakes of water.
- LAMINA, a thin plate, the upper expanded part of a polypetalous flower.
- LANA, wool, a species of pubescence, which covers the surface of plants.
- LANATUM FOLIUM, a woolly leaf.
- LANCULATUM FOLIUM, a lance-shaped leaf.
- LATERALES FLORES, flowers coming from the sides.
- LAXUS CAULIS, loose, weak, slender.

- LEGUMEN, *pulse*, a pericarpium of two valves, in which the seeds are fixed along one suture only.
- LENTICULARIS SCABRITIES, a species of glandular scabrities, in the form of lentils.
- LEPROSUS, spotted as a leper, exemplified in LICHEN.
- LÆVIS CAULIS, *smooth*, having an even surface.
- LIBER, *the inner rind* or bark of a plant.
- LIGNOSUS CAULIS, *a woody stem*.
- LIGNUM, *wood*.
- LIGULATUS FLOS, when the petals, tubulated at the base, are plane linear towards the middle, and widest at the extremity, in form of a bandage.
- LILIACEÆ, *like a lily*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- LIMBUS, *a border*, the upper expanded part of a monopetalous flower.
- LINEA, *a line*, the second degree in the Linnæan Scale for measuring plants, the twelfth part of an inch.
- LINEARE FOLIUM, *a narrow leaf*, whose opposite margins are almost parallel, as in PINUS.
- LINEATUM FOLIUM, leaves whose superficies are marked with parallel lines, running lengthways.
- LINGULATUM FOLIUM, a leaf shaped like a tongue.
- LOBATUM FOLIUM, when leaves are divided to the middle into parts that stand wide from each other, and have their margins convex.
- LOCULAMENTUM, *a cell*, the divisions of that species or pericarpium, called a CAPSULA.
- LOCUS FOLIORUM, the particular part of the plant to which the leaf is affixed.
- LOMENTACEÆ, *bean meal*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- LONGIUSCULUS, *longish*.
- LONGUM PERIANTHIUM, when the tube of the calyx is equal in length to that of the corollæ.
- LUCIDUM FOLIUM, *clear*, shining.
- LUNATUM FOLIUM, *moon-shaped leaves*, when they are round and hollowed at the base like a half moon.

LUNULATE, shaped like a crescent.

LURIDÆ, pale, wan, an order of plants in the *Fragmenta Methodi Naturalis of Linnæus*.

LUXURIANS FLOS, a luxuriant flower.

LYRATUM FOLIUM, leaves shaped like a harp or lyre.

M

MARCESCENS COROLLA, flowers withering on the plant.

MARGO FOLII, the margin or edge of the leaf.

MAS, *male*. See class *Dicœcia*.

MASCULUS FLOS, *male flowers*, containing antheræ, but no stigma.

MEDULLA, *marrow*, the pith or heart of a plant.

MEMBRANACEUM FOLIUM, when leaves have no distinguishable pulp between their surfaces.

MEMBRANATUS CAULIS, a stalk covered with thick membranes.

MONADELPHIA, *one brother*, the sixteenth class in the *Sexual System*.

MONANDRIA, *one male*, the first class in the *Sexual System*.

MONOCOTYLEDONES, a term in placentation, applied to plants whose seed have a single cotyledon.

MONŒCIA, *one house*, the twenty-first class in the *Sexual System*.

MONOGYNIA, *one female*, the first order of the first thirteen classes in the *Linnæan System*.

MONOPETALA COROLLA, a flower having one petal.

MONOPHYLLUM INVOLUCRUM, consisting of one leaf.

MONOSPERMA, having one seed.

MILIARIS SCABRITIES, a species of glandular roughness appearing on the surface of some plants like grains of millet.

MUCRONATUM FOLIUM, a leaf terminating in a sharp point.

MULTIFIDUM FOLIUM, a leaf divided into many linear segments or divisions.

MULTIFLORUS PEDUNCULUS, a foot-stalk bearing many flowers.

MULTIPARTITUM FOLIUM, a leaf divided into many parts.

MULTIPLICATUS FLOS, a luxuriant flower whose corolla is multiplied so as to exclude some of the stamina.

MULTISILIQÆ, *many pods*, an order of plants in the *Fragmenta Methodi Naturalis of Linnæus*.

- MURICATUS CAULIS, a stalk, whose surface is covered with sharp points, like the murex shell.
- MURICATE, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- MUSCI, *mosses*, one of the seven families in the vegetable kingdom, and an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- MUTICA GLUMA, when the arista is wanting.
- MUTILATUS FLOS, a mutilated flower.

N

- NATANS FOLIUM, a leaf which swims on the surface of water.
- NAVICULARIS VALVULA, when the valve of a seed vessel resembles a ship.
- NECESSARIA POLYGAMIÆ, *necessary marriages*, the fourth order of the nineteenth class in the Sexual System.
- NECTARIUM, that part of the corolla that contains the honey juice.
- NERVOSUM FOLIUM, leaves whose surface is full of nerves or strings.
- NIDULANTIA SEMINA BACCARUM, seeds nestling in the pulp of a berry.
- NITIDUM FOLIUM, a bright shining glossy leaf.
- NUCAMENTACEÆ, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- NUCLEUS, *a kernel*.
- NUDUS CAULIS, a naked stalk.
- NUTANS CAULIS, a nodding stalk.
- NUX, *a nut*.

O

- OBCORDATUM PETALUM, *a heart-shaped petal*, with its apex downwards.
- OBLIQUUM FOLIUM, when the apex of the leaf points obliquely towards the horizon.
- OBLONGUM FOLIUM, an oblong leaf.
- OBSOLETE LOBATUM FOLIUM, leaves having lobes scarce discernible.
- OBTUSUM FOLIUM, leaves blunt or rounded at the apex.

- OBVOLUTUM FOLIUM, rolled against each other, when their respective margins alternately embrace the straight margin of the opposite leaf.
- OCTANDRIA, *eight males*, the eighth class in the Sexual System.
- OFFICINALIS, plants used in medicine, and kept in the apothecaries' shops.
- OPERCULUM, *a cover*, as in the mosses.
- OPPOSITI RAMI FOLIA, branches and leaves that grow by pairs opposite each other.
- ORBICULATUM FOLIUM, round leaves.
- ORCHIDÆ ORCHIS, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- ORDO, *order*.
- ORGYA, a fathom, or six Parisian feet.
- OVALE FOLIUM, an oval leaf.
- OVALIUM, the germen.
- O VATUM FOLIUM, an oval, or egg-shaped leaf.

P

- PAGINA FOLII, the surface of a leaf.
- PALEA, *chaff*, a thin membrane rising from a common receptacle, which separates the flosculi.
- PALEACEUS PAPPUS, chaffy down.
- PALMÆ, *palms*, one of the seven families of the vegetable kingdom.
- PALMATA RADIX, *a handed root*, as in ORCHIS.
- PALMATUM FOLIUM, a leaf shaped like an open hand.
- PALUSTRIS, marshy or fenny.
- PANDURIFORME FOLIUM, *shaped like a guitar*, a musical instrument so called.
- PANICULA, *a panicle*, or loose spike of grass.
- PAPILIONACEUS, *butterfly-shaped flower*, as in the class Diadelphia of *Linnaeus*.
- PAPILIONACEÆ, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.

- PARILOSUM FOLIUM**, a *nipple*, a leaf covered with dots or points, like nipples.
- PAPPUS**, *down*.
- PAPULOSUM FOLIUM**, a leaf whose surface is covered with pimples.
- PARABOLICUM FOLIUM**, a leaf in form of a parabola.
- PARALLELUM DISSEPIMENTUM**, when the dissepiments are parallel to the sides of the pericarpium.
- PARASITICA PLANTA**, plants that grow only out of other plants, as the **VISNUM**.
- PARTIALIS UMBELLA**, a partial umbel.
- PARTIALE INVOLUCRUM**, when at the base of the partial umbel.
- PARTITUM FOLIUM**, a divided leaf.
- PARVUM PERIANTHIUM**, a *little flower-cup*, or comparatively small, opposed to **MAGNUM**.
- PATENS CAULIS, RAMUS, &c.** spreading stalks and branches.
- PATULUS CALYX**, a spreading cup.
- PAUCIFLORIS**, having few flowers.
- PEDALIS CAULIS**, a stalk a foot in height.
- PEDATUM FOLIUM**, a species of compound leaf, whose divisions resemble the toes of a foot, as in **HELLEBORUS FÆTIDA**.
- PEDICELLUS**, a little foot-stalk.
- PEDUNCULARIS CIRRHUS**, a tendril proceeding from the foot-stalk of a flower.
- PEDUNCULATI FLORES**, flowers growing on foot-stalks.
- PEDUNCULUS**, the foot-stalk of a flower.
- PELTATUM FOLIUM**, when the foot-stalk is inserted into the disk of the leaf, and not into its base.
- PENICILLIFORMIA STIGMATA**, a stigma in form of a painter's pencil.
- PENTAGONUS CAULIS**, a five-angled stalk.
- PENTAGYNIA**, *five females*, the fifth order of a class.
- PENTANDRIA**, *five males*, the fifth class in the *Sex. Syst.* of *Linnaeus*.
- PENTAPETALA COROLLA**, a flower consisting of five petals.
- PENTAPHYLLUS CALYX**, a calyx consisting of five leaves.
- PERENNIS RADIX**, a *perennial root*, continuing for many years.
- PERFECTUS FLOS**, *flowers having petals*, the perfect flowers of *Ray*, *Tournefort*, and other botanists.

- PERFOLIATUM FOLIUM, when the base of the leaf entirely surrounds the stem, or when the stalk grows through the centre of the leaf, as in CRASSULA PERFOLIATA.
- PERFORATI COTYLEDONES, *pierced through*, a species of the MONOCOTYLEDONES, exemplified in the GERMINA; also an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- PERIANTHIUM, a kind of calyx so called when contiguous to the fructification.
- PERICARPIUM, a species of pod that contains the seed.
- PERICHÆTIUM, a modification in the receptaculum in the MUSCI and ALGÆ.
- PERPENDICULARIS RADIX, a perpendicular, or downright root.
- PERSONATÆ, *masked*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- PES, a foot.
- PETALIFORMIA STIGMATA, a stigma resembling the shape of a petal.
- PETALODES FLOS, a flower having petals.
- PETALUM, the corollaceous teguments of a flower.
- PETIOLARIS CIRRHUS, a tendril proceeding from the foot-stalk of a leaf.
- PETIOLATUM FOLIUM, a leaf growing on a foot-stalk:
- PETIOLUS, a little foot-stalk.
- PILEUS, *a hat or bonnet*, the orbicular expansion of a mushroom, which covers the fructification.
- PILI, hairs.
- PILOSUM FOLIUM, leaves whose surface is covered with long distinct hairs.
- PINNATIFIDUM FOLIUM (a winged leaf), applied to simple leaves whose laciniaë are transverse to the rachiaë.
- PINNATUM FOLIUM, a winged leaf.
- PIPERITÆ, *pepper*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- PISTILLUM, or female organ of generation, whose office is to receive the farina fecundans.
- PIXIDATUM FOLIUM, a kind of foliage, where one leaf is let into another by a joint, as in EQUISETUM.

- PLACENTATIO, *Cotyledons*, of the seed.
- PLANIPETALUS FLOS, a flower with plain flat petals.
- PLANTÆ, *plants*, one of the seven families of vegetables, comprehending all which are not included in the other six tribes.
- PLANUM FOLIUM, plain flat leaves.
- PLENUS FLOS, a full or double flower.
- PPLICATUM FOLIUM, a plaited leaf.
- PLUMATA SETA, a feathered hair or bristle.
- PLUMOSUS PAPPUS, a kind of soft down.
- PLUMULA, the ascending scaly part of the corculum.
- POLLEN, *meal*, the prolific powder contained in the anthera.
- POLLEX, *a thumb*, the length of the first joint of the thumb, or a Parisian inch.
- POLYADELPHIA, *many brotherhoods*, the eighteenth class in the Sexual System.
- POLYANDRIA, *many males*, the thirteenth class in the Sexual System of *Linnaeus*.
- POLYCOTYLEDONES, *many cotyledons*.
- POLYGAMIA, *many marriages*, the twenty-third class in the Sexual System.
- POLYGYNIA, *many females*, an order of some of the classes in the Sexual System.
- POLYPETALA COROLLA, a flower consisting of many petals.
- POLYPHYLLUM INVOLUCRUM, an involucre of many leaves.
- POLYSTACHIUS CULMUS, a stalk of grass having many spikes.
- POMACEÆ, POMUM, *an apple*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- POMUM, *an apple*.
- PORI, *pores*.
- PRÆMORSA RADIX, *a bitten root*, when it ends abruptly, as in SCABIOSA.
- PRECIÆ, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- PRISMATICUS CALYX, a triangular flower-cup.
- PROCUMBENS CAULIS, lying on the ground.

- PROLIFER FLOS, flowers growing through, or out of one another, either from the centre or side.
- PROMINULUM DISSEPIENTUM, jetting out beyond the valves.
- PRONUM DISCUM FOLII, leaves having their face downwards.
- PROPAGO, *a shoot*, the seed of mosses.
- PROPRIUM INVOLUCRUM, an involucre when at the base of an umbellated flower.
- PSEUDO, *a bastard*.
- PUBES, *down*, or *hair*, one of the seven kinds of fulcra.
- PULPOSUM FOLIUM, a leaf having a pulpy or fleshy substance.
- PULVERATUM FOLIUM, a leaf powdered with a kind of dust like meal, as in PRIMULA FARINOSA.
- PUNCTATUM FOLIUM, a leaf sprinkled with hollow dots or points.
- PUTAMINEÆ, *like a shell*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.

Q

- QUADRANGULARE FOLIUM, *a quadrangular leaf*, having four prominent angles in the circumscription of its disk.
- QUADRIFIDUM FOLIUM, a leaf divided into four parts.
- QUADRIJUGUM FOLIUM, a leaf having four pair of folioles.
- QUADRILOBUM FOLIUM, a leaf consisting of four lobes.
- QUADRIPARTITUM FOLIUM, a leaf consisting of four divisions down to the base.
- QUINA FOLIA, verticillate leaves, coming by fives.
- QUATERNA FOLIA, when verticillate leaves come by fours, having four in each whorl.
- QUINATUM FOLIUM, when a digitate leaf has five folioles.
- QUINQUANGULARE FOLIUM, a leaf having five prominent angles in the circumscription of the disk.
- QUINQUEJUGUM FOLIUM, when a pinnated leaf has five pair of folioles.
- QUINQUELOBUM FOLIUM, a leaf having five lobes.
- QUINQUEFIDUM FOLIUM, a leaf consisting of five divisions, with linear sinuses, and straight margins.
- QUINQUEPARTITUM FOLIUM, consisting of five divisions down to the base.

R

RACEMUS, a bunch of grapes or currants, or any other bunch of berries that bears that resemblance.

RACHIS, the back bone, a species of receptaculum, as in the **PANICUM**.

RACHIS FOLII PINNATI, the middle rib of a winged leaf, to which the folioles are affixed.

RADIATUS FLOS, a species of compound flowers, in which the florets of the disk are tubular, and those of the radius ligulate, as in the class Syngenesia.

RADICALIA FOLIA, leaves proceeding immediately from the root.

RADICANS CAULIS, a stalk bending to the ground, and taking root where it touches the earth.

RADICATUM FOLIUM, leaves shooting out roots.

RADICULA, a little root.

RADIUS, a ray, the ligulate margin of the disk of a compound flower.

RADIX, a root:

RAMEA FOLIA, regards leaves that grow only on the branches, and not on the trunk.

RAMOSISSIMUS CAULIS, stalks abounding with branches irregularly disposed.

RAMUS, a branch of a tree.

RAMOSUS CAULIS, a stalk having many branches.

RECEPTACULUM, a receptacle, the basis on which the parts of fructification are connected.

RECLINATUM FOLIUM, a leaf reclined or bending downward.

RECURVATUM FOLIUM, a leaf bent backwards.

REFLEXUS RAMUS, a branch bent back towards the trunk.

REGULARIS COROLLA, a flower whose parts are regular in its figure and magnitude.

REMOTUS VERTICILLUS, when the whorls of flowers and leaves stand at a distance from one another.

RENIFORME FOLIUM, a kidney shaped leaf.

REFANDUM FOLIUM, a leaf having a bending or waved margin, without any angles.

- REPENS RADIX, a creeping root extending horizontally.
- REPENS CAULIS, a creeping stalk either running along the ground, on trees, or rocks, and striking roots at certain distances.
- REPTANS FLAGELLUM, creeping along the ground, as in FRAGARIA.
- RESTANTES PEDUNCULI, foot-stalks remaining on, after the fructification has fallen off.
- RESUPINATIO FLORUM, when the upper lip of the flower faces the ground, and the lower lip is turned upwards.
- RESUPINATUM FOLIUM, when the lower disk of the leaf looks upward.
- RETROFLEXUS RAMUS, a branch bent in different directions.
- RETROFRACTUS PEDUNCULUS, bent backwards towards its insertion, as if it were broken.
- RETUSUM FOLIUM, when the apex of the leaf is blunt.
- REVOLUTUM FOLIUM, a leaf rolled back.
- RHEADES, *the red poppy*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- RHOMBEUM FOLIUM, a leaf whose shape nearly resembles a rhombus.
- RHOMBOIDEUM FOLIUM, a leaf of a geometrical figure, whose sides and angles are unequal.
- RIGIDUS CAULIS FOLIA, *stiff*, hard, rigid.
- RIMOSUS CAULIS, abounding with clefts and chinks.
- RINGENS, grinning and gaping.
- ROSACEUS FLOS, a flower whose petals are placed in a circle, in form like those of a rose.
- ROSTELLUM, *a little beak*, the descending plain part of the coraculum of the seed.
- ROTACEÆ, *a wheel*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- ROTATUS LIMBUS, COROLLA, a wheel-shaped flower, expanded horizontally, having a tubular basis.
- ROTUNDATUM FOLIUM, a roundish leaf.
- RUBRA LACTESCENTIA, red milkiness in plants.
- RUDERATA LOCA, rubbishy places.
- RUGOSUM FOLIUM, a rough or wrinkled leaf.

S

- SAGITTATUM FOLIUM, an arrow-shaped leaf.
- SARMENTACEÆ, *a twig or shoot of a vine*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- SARMENTOSUS CAULIS, *the shoot of a vine*, naked between each joint, and producing leaves at the joints.
- SCABER CAULIS, ET FOLIUM, *scabby and rough*, having tubercles.
- SCABRIDÆ, *rough*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- SCABRITIS, *a species of pubescens*, composed of particles scarce visible to the naked eye, sprinkled on the surface of plants.
- SCANDENS CAULIS, a climbing stalk.
- SCAPUS, a species of stalk which elevates the fructification, and not the leaves, as in *NARCISsus*.
- SCARIOSUM FOLIUM, leaves dry on the margin that sound when touched.
- SCITAMINIA, *fair, beautiful*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- SCORPIOIDES FLOS, a flower resembling the tail of a scorpion.
- SCUTELLUM, a species of fructification which is orbicular, concave, and elevated in the margin, as in some species of *LICHEN*.
- SCYPHUR, *cup-bearing*, a subdivision of the genus *LICHEN*.
- SECRETORIA SCABRITIES, a species of glandular roughness on the surface of some plants.
- SECUNDA SPICA, a spike of grass with the flowers turned all towards one side.
- SECURIFORMIS PUBESCENTIA, a species of pubes on the surface of some plants, the bristles resembling an axe or hatchet.
- SEMEN, seed.
- SEMINALE FOLIUM, seed leaves.
- SEMPERVIRENS FOLIUM, an ever-green leaf.
- SEMITERES CAULIS, *half a cylinder*, flat on one side, and round on the other.
- SENA FOLIA, *leaves growing in sixes*, as in *GALIUM SPURIUM*.
- SENTICOSÆ, *a briar or bramble*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.

- SEPIARIÆ, *a hedge*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- SERICIUM FOLIUM, a leaf whose surface is of a soft silky texture.
- SERRATUM FOLIUM, a sawed leaf.
- SESSILE FOLIUM, a leaf growing immediately to the stem, without any foot-stalk.
- SETÆ, a bristle, a species of pubescens, covering the surface of some plants.
- SETACEUM FOLIUM, leaves shaped like bristles.
- SEXUS PLANTARUM, plants are distinguished by the sex of their flowers, which are either male, female, or bisexual.
- SILICULA, *a little pod*, a bivalve pericarpium. See class *Tetradynamia*.
- SILIQUA, *a pod*, a pericarpium consisting of two valves, in which the seeds are fixed alternately to each suture.
- SILIQUOSA, the second order in the class *Tetradynamia*.
- SILIQUOSÆ, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- SIMPLEX CAULIS, a simple or single stem.
- SIMPLICISSIMUS CAULIS, the most simple stalk.
- SINUATUM FOLIUM, a leaf whose sides are hollowed or scalloped.
- SITUS FOLIORUM, the disposition of leaves on the stem and branches, which are either starry, by threes, opposite, alternate, scattered, or crowded.
- SOLIDUS CAULIS, a solid stalk or stem.
- SOLITARIUS PEDUNCULUS, when only one flower-stalk proceeds from the same part.
- SOLUTÆ STIPULÆ, *loose*, opposed to *adnatæ*.
- SPADIX, *the receptaculum of a palm*, a pedunculus which proceeds from a spatha.
- SPARSI RAMI, PEDUNCULI FOLIA, scattered without order.
- SPATHA, a species of calyx resembling a sheath.
- SPATHACEÆ, *like a sheath*, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- SPATULATUM FOLIUM, *a leaf in form of a spatula*, an instrument used to spread salve.

- SPECIES PLANTARUM, the third subdivision in the Linnæan System.
- SPICA, *a spike*, a species of inflorescence resembling an ear of corn.
- SPICA SECUNDA, when the flowers all turn towards one side.
- SPICA DISTICHA, when the flowers are in two rows, and look two ways.
- SPICULA, a little spike.
- SPINÆ, thorns or rigid prickles.
- SPINOSUS CAULIS, *strong prickles*, whose roots proceed from the wood of the stem, and not from the surface of the bark.
- SPIRALES COTYLEDONES, seminal leaves twisted spirally.
- SPITHAMA, *a span*, or seven Parisian inches.
- SPLENDENTIA FOLIA, a shining leaf.
- SQUAMOSA RADIX, a scaly root.
- SQUARROSUM, rough, scaly, or scurfy.
- STAMEN, the filaments that sustain the anthera.
- STAMINEUS FLOS, flowers having stamina, and no corolla.
- STATUMINATÆ, *a prop*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- STELLATA FOLIA, *leaves surrounding the stem*, like the rays of a circle.
- STELLATÆ SETA, a species of pubescens called bristles, when they arise from a centre in form of a star, as in the MESEMBRYANTHEMUM BARBATUM.
- STELLATA PLANTA, one of Mr. *Ray's* classes, the Tetrandria Monogynia of *Linnaeus*.
- STELLATÆ, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- STERILIS FLOS, *a barren flower*, masculus of *Linnaeus*.
- STIGMA, apex of the pistillum.
- STIMULI, stings.
- STIPITATUS PAPPUS, a kind of trunk that elevates the down, and connects it with the seed.
- STIPULA, one of the kinds of fulcra of plants, generally growing on each side of the base of the foot-stalks of leaves or flowers, and are either by twos, single, deciduous, abiding, adhering, loose, on the inside of the foot-stalks or on the outside.

STIPULARES GLANDULÆ, glands produced from stipulæ.

STOLO, a shoot, which running on the surface of the ground strikes root at every joint, as in FRAGARIA and others.

STRIATUS CAULIS, CULMUS, &c. channelled streaks, running lengthwise in parallel lines.

STRICTUS CAULIS, straight stiff shoots.

STRIGÆ, ridges, rows.

STROBILUS, a species of pericarpium, formed from an amentum, as the cone of the pine-tree.

STYLUS, that part of the pistillum which elevates the stigma from the germen.

SUBMERSUM FOLIUM, when aquatic plants have their leaves sunk under the surface of the water.

SUBRAMOSUS CAULIS, a stalk having few branches.

SUBROTUNDUM FOLIUM, a leaf almost round.

SUBULATUM FOLIUM, an awl-shaped leaf.

SUCCULENTÆ, juicy, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.

SUFFRUTEX, an under shrub.

SULCATUS CAULIS, CULMUS, a stalk deeply furrowed lengthways.

SUPERFLUA POLYGAMIA, superfluous, the second order in the class Syngenesia.

SUPERUS FLOS, when the receptacle of the flower stands above the germen.

SUPRA-AXILLARIS PEDUNCULUS, the foot-stalk of a flower, whose insertion is above the angle formed by the branch.

SUPRA-DECOMPOSITA FOLIA, are composite leaves which have little leaves growing on a subdivided foot-stalk.

SUPRA-FOLIACEUS, PEDUNCULUS, the foot-stalk of a flower inserted into the stem immediately above the leaf.

SURCULUS, a twig, the stalks or branches of mosses.

SYNGENESIA, to generate together, the nineteenth class in the Sexual System.

T

TEGUMENTUM, a cover, the perianthium and corolla.

TERES CAULIS FOLIUM, a cylindrical stalk or leaf.

- TERGEMINUM FOLIUM COMPOSITUM, *a leaf three times double*, when a dichotomus petiolus is subdivided, having two foliola on the extremity of each division.
- TERMINALIS FLOS, flowers terminating a branch.
- TERNA FOLIA, leaves in whorls by threes.
- TERNATUM FOLIUM, a chequered leaf, whose squares are of different colours.
- TESSELLATUM FOLIUM, a chequered leaf, whose squares are of different colours.
- TETRADYNAMIA, the superiority or power of four, the fifteenth class in the Sexual System.
- TETRAGONUS CAULIS, a four-cornered or square stalk.
- TETRAGYNIA, *four females*, the fourth order of some of the classes in the Sexual System.
- TETRANDRIA, *four males*, the fourth class in the Sexual System.
- TETRAPETALA COROLLA, a flower consisting of four petals.
- TETRAPHYLLUS CALYX, a flower cup consisting of four leaves.
- TETRASPERMA PLANTA, producing four seeds.
- THALAMUS, *a bed*, the receptacle.
- THECA, a sheath.
- THYRSUS, a spike like a pine-cone.
- TOMENTOSUS CAULIS FOLIA, a stalk and leaf covered with a whitish down like wool.
- TOMENTUM, a species of pubescence, covering the surface of some plants of woolly or downy substance.
- TOROSUM PERICARPIMUM, brawny protuberances, like the swelling of the veins when a pericarpium is bunched out by the enclosed seeds.
- TORTA COROLLA, when the petals of a flower are twisted, as in NERIUM.
- TORTILIS ARISTA, awns or beards of corn twisted like a screw.
- TRANSVERSUM DISSEPIMENTUM, when the dissepiiments are at right angles with the sides of the pericarpium.
- TRAPEZIFORME FOLIUM, a leaf having four prominent angles, whose sides are neither equal nor opposite.
- TRIANDRIA, *three males*, the third class in the Sexual System.

- TRIANGULARE FOLIUM, a triangular leaf.
- TRICOCCA CAPSULA, a capsule with three cells, and a single seed in each cell.
- TRICOCCEÆ, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- TRICUSPIDATA, three-pointed:
- TRIFIDUM FOLIUM, a leaf divided into three linear segments, having straight margins:
- TRIFLORUS PEDUNCULUS, a foot-stalk bearing three flowers.
- TRIGONUS CAULIS, a three-sided stalk.
- TRIGYNIA, *three females*, the third order in some of the classes.
- TRIHILLATÆ, a seed having three eyes.
- TRIJUGUM FOLIUM, a winged leaf, with three pair of foliola.
- TRILOBUM FOLIUM, a leaf having three lobes.
- TRINERVUM FOLIUM, a leaf having three strong nerves running from the base to the apex.
- TRICECIA, *three houses*, the third order in the class Polygamia in the Sexual System.
- TRIPARTITUM FOLIUM, a leaf divided into three parts down to the base.
- TRIPETALA COROLLA, a flower consisting of three petals.
- TRIPETALOIDEÆ, *three-petaled*, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- TRIPHYLLUS CALYX, a cup consisting of three leaves.
- TRIPINNATUM FOLIUM, COMPOSITUM, a leaf having a triple series of pinna, or wings.
- TRIPLINERVE FOLIUM, a leaf having three nerves running from the base to the apex.
- TRIQUETRUM FOLIUM, CAULIS, leaves and stalks having three plain sides.
- TRISPERMA, *three-seeded*, as in EUPHORBIA.
- TRITERNATUM FOLIUM, COMPOSITUM, a compound leaf when the divisions of a triple petiolus are subdivided into threes.
- TRIVALVE PERICARPIUM, a pod consisting of three valves.
- TRUNCATUM FOLIUM, a leaf having its apex as it were cut off.
- TRUNCUS, the body or stem of a tree.

- TUBERCULATUS, having pimples or tubercles.
- TUBERCULUM, a little pimple.
- TUBEROSA RADIX, a tuberous or knobbed root.
- TUBULATUM PERIANTHIUM, tubular flowers, as in the class Didynamia.
- TUBULOSI FLOSCULI, tubular florets nearly equal, one of the three divisions of compound flowers.
- TUBUS, a tube, the lower and narrow part of a monopetalous flower.
- TUNICATUS RADIX, a species of bulbous root, having coats lying one over another from the centre to the surface, as in the ONION, TULIP, &c.
- TURBINATUM PERICARPIUM, a kind of pod shaped like a top, narrow at the base and broad at the apex.
- TURGIDUM LEGUMEN, swollen, puffed out, as in ONONIS.
- TURIO, the young buds or shoots of pines.

V

- VAGINALES, sheathed, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- VAGINANS FOLIUM, a leaf like a sheath, whose base infolds the stem.
- VALVULA, a valve, a partition of the external cover of that sort of pericarpium called capsula.
- VEGETABILIA, one of the three kingdoms of nature.
- VENOSUM FOLIUM, the veins which run over the whole surface of a leaf.
- VENTRICOSA SPICA, a spike narrowing at each extremity, and bellying out in the middle.
- VENTRICULOSUS CALYX, a flower-cup bellying out in the middle, but not in so great a degree as *Ventricosus*.
- VEPRECULÆ, a briar or bramble, an order of plants in the Fragmenta Methodi Naturalis of *Linnaeus*.
- VERRUCOSA CAPSULA, a capsule having little knobs or warts on its surface.

- VERSATILIS ANTHERA, when the anthera is fixed by the middle on the point of the filament, and so poised as to turn like the needle of a compass.
- VERTICALIA FOLIA. Leaves so situated that their base is perpendicular above the apex.
- VERTICILLATI RAMI, FLORES, FOLIA, branches, flowers, or leaves surrounding the stem like the rays of a wheel.
- VERTICILLATÆ, an order of plants in the *Fragmenta Methodi Naturalis* of *Linnaeus*.
- VERTICILLUS, a species of inflorescence, in which the flowers grow in whorls, as in *MENTHA*.
- VESICULA, a little bladder.
- VESICULARIS SCABRITIES, a kind of glandular roughness, resembling *VESICULÆ*.
- VEXILLUM, *a standard*, the upright petal of a papilionaceous flower.
- VILLOSUS, CAULIS, FOLIUM, a stalk or leaf covered with soft hairs.
- VIRGATUS CAULIS, *stalks shooting out*; slender, straight branches or rods.
- VISCIDUM FOLIUM, *a leaf whose surface is clammy*.
- VISCOSITAS, *gluey*, clammy.
- ULIGINOSA LOCA, *boggy places*.
- UMBELLA, *an umbel or umbrella*.
- UMBELLATUS FLOS, *an umbellated flower*, as in *PENTANDRIA DIGYNIA*.
- UMBELLULA, *a little umbel*.
- UMBILICATUM FOLIUM, a peltate leaf, shaped like a navel, at the insertion of the foot-stalk.
- UNCINATUM STIGMA, *a hooked stigma*.
- UNDATUM FOLIUM, *a waved leaf*, whose surface rises and falls in waves towards the margin.
- UNDULATA COROLLA, a flower whose petals are waved.
- UNGUIS, *a nail or claw*, that part of a petal that is joined to the receptacle.
- UNICUS FLOS, *one flower*.
- UNICUS RADIX, *a single root*.

- UNIFLORUS PEDUNCULUS, one flower on a foot-stalk.
- UNILATERALIS RACEMUS, a bunch of flowers growing on one side.
- UNIVERSALIS UMBELLA, *an universal umbel.*
- VOLVA, the membranaceous calyx of the fungi.
- VOLUBILIS CAULIS, *a twining stalk.*
- URCEOLATA, COROLLA, *a pitcher-shaped flower.*
- URENS CAULIS, FOLIUM, a leaf or stalk, burning, stinging, as
NETTLES.
- UTRICULI, a species of glandular secretory vessels, on the surface
of various plants.
- VULGARIS, *common*, the trivial name of many plants in the books
of old botanists.

TABLE VIII.

DERIVATIONS

OF

THE BOTANIC TERMS,

ALPHABETICALLY ARRANGED.

A

- ABRUPTUM** *Folium pinnatum*; from *abrumpor*, to be broken.
- ACAULIS** *Herba*; from *à priv.* and *caulis*.
- ACEROSUM** *Folium*; from *acus*, chaff.
- ACICULARIS**; from *acicula*, a pin, or small needle.
- ACINACIFORME**; from *acinaces*, a Persian scymitar.
- ACOTYLEDONES**; from *à priv.* and *Cotyledon*.
- ACULEI**; from *ἄκτις*, *cuspis*, a point.
- ACULEATUS** *Caulis, Folium*; from *aculeus*, a sting.
- ACUMINATUM** *Folium*; from *acuo*, to sharpen.
- ACUTE SERRATUM** *Folium*; from *acuo*, to sharpen, and *serra* a saw.
- ACUTUM** *Folium*; from *acuo*, to whet.
- ADNATUM** *Folium*; from *ad*, to, and *nascor*, to be born, to grow,
growing close to the stem.
- ADPRESSA** *Folia*; from *ad*, to, and *pressus*, pressed.
- ÆSTIVATIO**; from *æstas*, summer.
- AGGREGATUS** *Flos*; from *aggrego*, to assemble.

- ALARIS *Pedunculus*; from *ala*.
- ALATUS *Petiolus*; from *ala*, a wing.
- ALBURNUM; from *albus*, white.
- AMENTACEÆ; from *amentum*, a thong.
- AMENTUM; from *αμμα*, *vinculum*, a bond or thong.
- AMPLEXICAULE *Folium*; from *amplector*, to embrace, and *caulis*, a stem.
- ANDROGYNA *Planta*; from *ανηρ*, *vir*, a man, and *γυνη*, *mulier*, a woman.
- ANGUSTIFOLIA; from *angustus*, narrow, and *folium*, a leaf.
- ANGYOSPERMIA; from *αγγος*, *vas*, a vessel.
- ANNUA *Radix*; from *annus*, a year.
- ANNULATUS *Stipes*; from *annulus*, a small ring.
- ANOMALÆ *Gemmæ*; from *à* priv. and *ουαλος*, *æqualis*.
- ANTHERA; from *ανθος*, *flos*, a flower.
- APETALUS *Flos*; from *à* priv. and *petalum*.
- APEX *Folii*; from *apiendo*, i. e. *ligando*.
- APHYLLUS *Caulis*; from *à*, and *φυλλον*, *folium*, a leaf.
- APOPHYSIS; from *απο*, and *φωο*, *nascor*, to grow from.
- APPENDICULATUS *Petiolus*; from *appendicula*, dim. from *appendix*, a little appendage.
- ARBORESCENS; from *arbor*, a tree.
- ARBOREUS *Caulis*; from *arbor*, a tree.
- ARBUSTIVA; from *arbutum*, a copse of shrubs, or trees; an orchard, a vineyard.
- ARCUATUM *Legumen*; from *arcus*, the curvature of an arch, or of a bow-stick.
- ARILLATA *Semina*; from *arillus*.
- ARISTA; from *areo*, to be dry or parched.
- ARISTATA *Gluma*; from *arista*.
- ARTICULUS *Culmi*; from *artus*, a joint or limb.
- ASCYROYDEÆ; from *Ασκυρον*, Pliny's name for the *Hypericum*.
- ASPERIFOLIÆ; from *asper*, rough, and *folium*, a leaf.
- ASSURGENTIA *Folia*; from *assurgo*, to rise up.
- ATTENUATUS *Pedunculus*; from *attenuor*, to be wasted, worn.
- AUCTUS *Calyx*; from *augeor*, to be increased.

AVENIA *Folia*; from *à*, not, and *vena*, a vein.

AURICULATUM *Foliolum*; from *auricula*, a little ear, dim. from *auris*, the ear.

AURIFORMIS; from *auris*, an ear.

AXILLARIA *Folia*; from *axilla*, the arm-pit.

B

BACCATUM *Receptaculum seminum*; from *bacca*.

BARBATUM *Folium*; from *barba*, a beard.

BICORNES; from *bis*, and *cornu*, a horn.

BIENNIS *Radix*; from *bis*, twice, and *annus*, a year.

BIFARIA *Folia*; from *bis*, and *fari*, to speak.

BIFERÆ *Plantæ*; from *bis*, and *fero*, to bear.

BIFIDUM *Folium*; from *bis*, twice, and *fissum*, cloven.

BIFLORUS *Pedunculus*; from *bis*, and *flos*, a flower.

BIGEMINUM *Folium compositum*; from *bis*, twice, and *geminus*, double.

BIJUGUM *Folium*; from *bis*, and *jugo*, to yoke.

BILABIATUS *Corolla*; from *bis*, and *labium*, a lip.

BILOBUM *Folium*; from *bis*, twice, and *lobus*, the tip of the ear.

BILAMELLATUM *Stigma*; from *bis*, and *lamella*, a thin plate.

BILOCULARIS *Capsula*; from *bis*, and *loculus*, a small place.

BINATA *Folia*; from *binus*, two and two.

BIPARTITUM *Folium*; from *bis*, and *partitus*, divided.

BIPINNATUM *Folium compositum*; from *bis*, and *pinnatum*, winged.

BITERNATUM *Folium compositum*; from *bis*, twice, and *ternus*, threefold.

BIVALVE *Pericarpium*; from *bis*, and *valvæ*, doors or valves.

BLATTARIÆ; from *blatta*, a moth, or little worm.

BRACHIATUS *Caulis*; from *brachium*, an arm.

BRACTEATUS *Pedunculus*; from *bractea*, a floral leaf.

BULBIFERUS *Caulis*; from *bulbus*, a round root.

BULBOSA *Radix*; from *bulbus*, a species of onion.

BULLATUM *Folium*; from *bulla*, a bubble.

C

- CADUCUM *Folium*; from *cado*, to fall.
- CALAMARIÆ; from *calamus*, a reed.
- CALCARATUM *Nectarium*; from *calcar*, a spur.
- CALICULATUS *Calyx*; from *calicula*, dim. from *calyx*.
- CALYCANTHEMI; from *calyx*.
- CALYCIFIBRÆ; from *calyx*, and *fibra*, a fibre.
- CALYCIFLORÆ; from *calyx*, and *flos*.
- CALYPTRA; from *καλυπτω*, *tego*, to cover.
- CALYX; from *καλυπτω*, *tego*, to cover.
- CAMPANACEI; from *campana*, a bell.
- CAMPANIFORMIS *Corolla*; from *campana*, a bell.
- CAMPANULATA *Corolla*; from *campanula*, a little bell.
- CANALICULATUM *Folium*; from *canalicula*, dim. from *canalis*, a channel.
- CANCELLATUS *Pilus*; from *cancelli*, cross bars or trellis.
- CANDELARIS; from *candela*, a candle.
- CAPILLARIS *Pappus*; from *capillus*, hair.
- CAPILLUS (quasi *capitis pilus*), hair.
- CAPITULUM; dim. from *caput*, a head.
- CAPREOLUS; dim. from *caprea*, a branch that produces tendrils.
- CARINATUM *Folium*; from *carina*, the keel or bottom of a ship.
- CARIOPHYLLÆUS *Flos*; from *caryophyllus*, the clove-tree.
- CARNOSUM *Folium*; from *caro*, flesh.
- CARTILAGINEUM *Folium*; from *cartilago*, a cartilage.
- CARYOPHYLLEI; from *caryophyllus*, a pink or gillyflower.
- CATENULATA *Scabrities*; from *catena*, a chain.
- CAUDEX; from *cauda*, a tail.
- CAULESCENS *Planta*; from *caulis*.
- CAULINA *Folia*; from *caulis*, a stem.
- CAULIS; from *καυλος*, a stalk.
- CERNUUS *Pedunculus*, *Flos*; from *cerno*, to discern, *quod terram cernat*.
- CESPITOSA *Planta*; from *cespes*, turf or sod.
- CILIATUM *Folium*; from *cilium*, the eye-lash.
- CIRCINALIA *Folia*; from *circes*, a hoop or ring.

- CIRCUMSCISSA *Capsula*; from *circum*, about, and *cædo*, to cut.
- CIRRHIFERUS *Pedunculus*; from *cirrhus*, and *fero*.
- CIRRHOSUM *Folium*; from *cirrhus*, a tuft or lock of hair.
- CIRRIUS, rather *cirrus*; from *κερας*, *cornu*, a horn, *quod cirri cornuum figuram referant*.
- CLAVÆFORMIS; from *clava*, a club.
- CLAVATUS *Petiolus, Pedunculus*; from *clavis*, a nail, or *clava*, a club.
- CLAVICULA; dim. from *clavis*, a key,
- COADUNATE; from *coaduno*, to join or gather together.
- COARCTATI *Rami*; from *coarcto*, to straiten or press together.
- COCHLEATUM *Legumen*; from *cochlea*, the shell of a snail.
- COLORATUM *Folium*; from *color*, colour.
- COLUMNIFERI; from *columna*, a pillar, and *fero*, to bear.
- COMA; from *κομη*, a bush or head of hair.
- COMOSÆ; from *coma*, a head of hair.
- COMPACTUM *Folium*; from *compingo*, to put together.
- CONDPLICATUM *Folium*; from *con*, and *duplicor*, to be doubled.
- CONFERTI *Rami*; from *confercio*, to fill, to stuff.
- CONFLUENTIA *Folia*; from *confluo*, to flow together.
- CONGLOBATUS *Flos*; from *con*, and *globus*, a ball.
- CLONGLOMORATI *Flores*; from *con*, and *glomus*, a clew.
- CONGESTA *umbella*; from *congeror*, to be heaped.
- CONICA *Scabrities*; from *κωνος*, *conus*, a cone.
- CONIFERÆ; from *κωνος*, a cone, and *fero*, to bear.
- CONJUGATUM *Folium*; from *con*, together, and *jugo*, to couple.
- CONNATUM *Folium*; from *con*, and *nascor*, to be born, to grow together.
- CONNIVENS *Corolla*; from *connivo*, to wink.
- CONTORTI; from *contorqueo*, to twist.
- CONVOLUTUS *cirrus*; from *convolto*, to wrap round.
- CORCULUM; dim. from *cor*, the heart.
- CORDATUM *Folium*; from *cor*, the heart:
- CORIACEOUS *Calyx*; from *corium*, leather.
- COROLLA; dim. from *corona*, a crown.
- COROLLULA; dim. from *corolla*.
- CORONARIÆ; from *corona*, a crown.

- CORONULA; dim. from *corona*.
- CORTEX; from *corium*, a hide, and *tego*, to cover.
- CORTICALIS *Gemmatio*; from *cortex*, rind or bark.
- CORTICATUM *Semen*; from *cortex*.
- CORYDALES; from *κorys*, *galea*, *cassis*, *galerita*, a helmet.
- COTYLEDON; from *κοίτη*, *cavitas*, a cavity.
- CRENATUM *Folium*; from *crena*, a notch.
- CRINITUS; from *crinis*, hair.
- CRISTATUS *Flos*; from *crista*, a tuft or crest.
- CRUCIFORMES *Flores*; from *crux*, a cross, and *forma*, form.
- CRYPTANTHERE; from *κρυπῶ*, *occulto*, to hide, and *ανθος*, *flos*, a flower.
- CRYPTOGAMIA; from *κρυπτός*, *occultus*, concealed, and *γάμος*, *nuptia*, nuptials.
- CUBITUS; from *cubando*, lying down, *quod ad sumendos cibos in ipso cubamus*.
- CUCULLATUM *Folium*; from *cucullus*, a coronet of paper in which grocers put their spices.
- CUCURBITACEÆ; from *cucurbita*, a gourd.
- CULMINALE; from *culmen*, the top or crown of any thing.
- CULMUS; from *καλαμῶς*, *calamus*, a reed or straw.
- CUNEIFORME *Folium*; from *cuneus*, a wedge.
- CUSPIDATUM *Folium*; from *cuspis*, the point of a spear.
- CYATHIFORMIS *Corolla*, *Calyx*; from *cyathus*, a cup.
- CYLINDRACEA *Spica*; from *cylindrus*, a roller, a cylinder.
- CYLINDRICA *Scabrities*; from *cylindrus*.
- CYMA; from *κύμα*, *fætus*.
- CYMOSUS *Flos*; from *cyma*, a sprout.
- CYTINIFORMIS *Calyx*; from *cytinus*, the flower of the pomegranate.

D

- DEDALEUM *Folium*; from *δαίδαλος*, *dædalus*, ingenious.
- DEBILIS, *Caulis*; from *de* and *habilis*.
- DECAGYNIĀ; from *δεκα*, *decem*, ten, and *γυνή*, *mulier*, a woman.
- DECANDRIA; from *δεκα*, *decem*, ten, and *ανηρ*, *maritus*, a husband.
- DECAPHYLLUS *Calyx*; from *δεκα*, *decem*, ten, and *φυλλον*, *folium*, a leaf.
- DECIDUUM *Folium*; from *decido*, to fall down, to die.

- DECUMBENS *Flos*; from *decumbo*, to lie down.
- DECURRENS *Folium*; from *decurro*, to run along.
- DECURSIVE *Folium pinnatum*; from *decurro*, to run along.
- DECUSSATA *Folia*; from *decusso*, to divide.
- DEFLEXUS *Ramus*; from *deflecto*, to bow or bend.
- DEFLORATA; from *de*, and *flos*.
- DEHISCENS *Siliqua*; from *dehisco*, to open, to gape.
- DELTOIDES *Folium*; from Δ , *delta*, the Greek D.
- DEMERSUM *Folium*; from *demergo*, to dive.
- DENDROIDIS *Surculus*; from *δενδρον*, *arbustum*, a shrub.
- DENTATUM *Folium*; from *dens*, a tooth.
- DENTICULATA *Semina*; from *denticulus*, a little tooth.
- DENUDATÆ; from *denudo*, to be stripped naked.
- DEPENDENS *Folium*; from *dependeo*, to hang down.
- DIADELPHIA; from *δισ*, *bis*, two, and *ἀδελφος*, *frater*, a brother.
- DIANDRIA; from *δισ*, *bis*, two, and *ανηρ*, *maritus*, a husband.
- DIANGLE; from *δισ*, and *αγλος*, *vas*, a vessel, or *locumentum*.
- DICHOTOMUS *Caulis*; from *διχοτομος*, *dissectus*, divided.
- DICOTYLEDONES; from *δισ*, and *cotyledon*.
- DICOCCUM *Pericarpium*; from *δισ*, and *κοκος*, *granum*, a grain.
- DIDYMA *Anthera*; from *διδυμος*, *geminus*, twins.
- DIDYNAMIA; from *δισ*, *bis*, two, and *δυναμις*, *potentia*, power.
- DIFFORMIA *Folia*; from *δισ*, and *forma*, form, shape.
- DIGITATUM *Folium*; from *digitus*, a finger.
- DIGYNIA; from *δισ*, and *γυνη*, *mulier*, a woman.
- DIMIDIATUM *Capitulum*; from *dimidius*, half.
- DIOECIA, *δισ*, *bis*, and *οικος*, *domus*, a house.
- DIPETALA *Corolla*; *δισ*, and *πεταλον*, *petalum*.
- DIPHYLLUS *Calyx*; from *δισ*, and *φυλλον*, *folium*, a leaf.
- DIPLOSANTHERÆ; from *διπλος*, *duplex*, double, and *anthera*.
- DISPERMA; from *δισ*, and *sperma*, a seed.
- DISSILIENS *Siliqua*; from *dissilio*, to break, to shiver.
- DISTICHA *Folia, Rami*; from *δισ*, and *στιχος*, *ordo*, rank.
- DIVARICATI *Rami*; from *divarico*, to stride.
- DODECANDRIA; from *δωδεκα*, *duodecim*, twelve, and *ανηρ*, *maritus*, a husband.
- DODRANTALIS *Caulis*; from *dodrans*, nine inches.

- DOLABRIFORME *Folium*; from *dolabra*, an axe.
 DORSALIS *Arista*, probably for *dorsualis*; from *dorsum*, the back.
 DRUPA; from *δρυς*, *arbor*, and *πίπτω*, *cado*, to fall; ripe fruit.
 DRUPACEÆ; from *drupa*.
 DUMOSÆ; from *dumus*, a bush.
 DUPLICATA *Radix*; from *duplex*, double.

E

- EBRACTEATUS *Racemus*; from *è* priv. and *bractea*, a bracteal or floral leaf.
 ECALCARATA *Corolla*; from *è* priv. and *calcar*, a spur.
 ECAUDATA *Corolla*; from *è* priv. and *cauda*, a tail.
 ECHINATUM *Pericarpium*; from *εχινος*, *crinaceus*, a hedgehog.
 EFFLORESCENTIA; from *effloresco*, to blow, to bloom.
 EMARGINATUM *Folium*; from *è*, and *margo*, the margin.
 ENERVIVM, or *enerve Folium*; from *è*, and *nervus*, a nerve, or string.
 ENNEANDRIA; from *εννεα*, *novem*, nine, and *ανηρ*, *maritus*, a husband.
 ENNEAPETALA *Corolla*; from *εννεα*, *novem*, nine, and *πεταλον*, *petalum*.
 ENODIS *Caulis*, *Culmus*; from *è*, and *nodus*.
 ENSATÆ; from *ensis*, a sword.
 ENSIFORME *Folium*; from *ensis*, a sword.
 EQUITANTIA *Folia*; from *equitans*, riding.
 EROSUM *Folium*; from *erodor*, to be gnawed.
 EXCEDENS; from *excedo*.
 ESSERTA *Stamina*; from *exsero*, to put forth.
 EXSTIPULATUS; from *ex*, and *stipula*, stubble or straw.
 EXSUCCUM *Folium*; from *ex*, and *succus*, juice.
 EXTRAFOLIACEÆ *Stipulæ*; from *extra*, and *folium*.

F

- FARCTUM *Folium*; from *farcio*, to stuff, to cram.
 FASCICULATA *Folia*; from *fasciculus*, a little bundle.
 FASCICULARIS *Radix*; from *fascis*, a bundle.
 FASCICULUS; dim. from *fascis*, a bundle.
 FASCIATA *Planta*; from *fascis*, a bundle.

- FASTIGIATI *Pedunculi*; from *fastigium*, the *apex*, or top of a pyramid.
- FIBROSA *Radix*; from *fibra*, a fibre.
- FILAMENTOSA *Radix*; from *filum*, a thread.
- FILAMENTUM; from *filum*, a thread.
- FILICES; from *filum*, a thread.
- FILIFORMIS *Filamentum*, *Stylus*, *Receptaculum*; from *filum*, a thread, and *forma*, form or shape.
- FIMBRICATA *Petala*; from *fimbria*, a border or fringe.
- FISSUM *Folium*; from *findor*, to be cloven.
- FISTULOSUS *Caulis*; from *fistula*, a pipe.
- FLABELLIFOLIÆ; from *flabellum*, a fan.
- FLAGELLUM; from *flagrum*, a whip or thong.
- FLORALIA *Folia*; from *flos*, a flower.
- FLORALIS *Gemma*; from *flos*.
- FLORIFERÆ *Gemma*; from *flos*, and *fero*, to bear.
- FLUVIATILIS; from *fluvius*, a river.
- FOLIARIS *Cirrus*; from *folium*, a leaf.
- FOLIATIO *Plantæ*; from *folium*.
- FOLIFERÆ *Gemma*; from *folium*, and *fero* to bear.
- FOLIOLUM; dim. of *folium*, a green leaf.
- FOLIOSUM *Capitulum*; from *folium*.
- FOLLICULUS; dim. from *follis*, a bag.
- FORNICATUM *Petalum*; from *fornix*, an arch or vault.
- FRONDESCENTIA; from *frons*, a leaf.
- FRONDOSUS *Caudex*; from *frons*.
- FRUCTESCENTIA; from *fructus*, fruit.
- FRUCTIFICATIO; from *fructus*, fruit.
- FRUCTIFLORÆ; from *fructus*, fruit, and *flos*, a flower.
- FRUSTRANLA *Polygamia*; from *frustra*, to no purpose.
- FRUTESCENS *Caulis*; from *frutex*, a shrub.
- FRUTICOSUS *Caulis*; from *frutex*, a shrub.
- FUGACISSIMA *Petala*; from *fugax*, fleeting.
- FULCRATUS *Caulis*, *Ramus*; from *fulcio*, to prop.
- FUNGI; from *σφογγος*, *fungus*.
- FURCATA *Frons*; from *furca*, a fork.
- FUSIFORMIS *Radix*; from *fusus*, a spindle.

G

- GALEA; from γαλή.
- GALEATUM *Labium*; from *galea*, a helmet.
- GEMINATUS *Pedunculus*; from *geminus*, double.
- GEMMA; from *geno*, i. e. *gigno*, or from γεμω, *plenus sum*.
- GEMATIO; from *gemma*, a young bud.
- GEMMIPARUS; from *gemma*, a bud, and *pario*, to bear.
- GENICULATUS *Caulis, Culmus, Pedunculus*; from *genu*, the knee.
- GENICULUM; from *genu*, the knee.
- GIBBUM *Folium*; from *gibba*, a hump on the back.
- GLADIATA *Siliqua*; from *gladius*, a sword or knife.
- GLANDULATIO; from *glans*, an acorn, a gland.
- GLAREOSIS, *locis* understood; from *glareo*, gravel.
- GLAUCOPHYLLUS; from γλαυκος, *glaucus*, blue, and φυλλον, *folium*, a leaf.
- GLOBOSA *Radix*; from *globus*, a globe.
- GLOBULARIS *Scabrities*; dim. from *globus*, a round ball or globe.
- GLOCHIDES; from γλωχίς, *cuspis*, a point.
- GLOMERATA *Spica*; from *glomus*, a clue of yarn or thread.
- GLUMA; from *glubo*, to strip the bark from a tree.
- GLUMOSUS; from *gluma*.
- GLUTINOSITAS; from *gluten*, glue, paste.
- GRANULATA *Radix*; from *granum*, a grain.
- GYMNOSPERMIA; from γυμνος, *nudus*, naked, and σπέρμα, *seed*.
- GYNANDRIA; from γυνή, *mulier*, a woman, and ανήρ, *vir*, a man.

H

- HABITUALIS *Character*; from *habitus*.
- HAMOSA *Seta*; from άμη, *falx*, a hook, asking Mr. Ainsworth's pardon.
- HASTATUM *Folium*; from *hasta*, a spear.
- HEDERIFOLIA; from *hedera*, the ivy.
- HEMISPHERICUS *Calyx*; from ήμισ, *semis*, half, and σφαίρα, *sphæra*, a sphere.
- HEPATICA; from *hepar*, the liver.
- HEPTANDRIA; from έπτα, *septem*, seven, and ανήρ, *maritus*, a husband.

HERBA ; *de etym. parum constat.*

HERBACEÆ *Planta* ; from *herba*, an herb.

HERMAPHRODITUS *Flos* ; from Ερμης *Mercury*, and Αφροδιτη, *Venus*.

HESPERIDÆ ; from *Hesperides*, whose orchards produced golden fruit.

HEXANDRIA ; from ἕξ, *sex*, six, and ανηρ, *vir*, a man.

HEXAGYNIA ; from ἕξ, *sex*, six, and γυνη, *mulier*, a woman.

HEXAPETALA *Corolla* ; from ἕξ, *sex*, and πεταλον, *petalum*.

HEXAPHYLLUS *Calyx* ; from φυλλον, *folium*, a leaf.

HOLERACEÆ ; from *olus*, pot-herbs, or herbs for food.

HORIZONTALIS *Flos* ; from *horizon*.

HYBRIDA *Planta* ; from ὑβρις, *injuria*, injury, dishonour.

HYPOCRATERIFORMIS *Corolla* ; from ὑπο, *ab*, and κρατηρ, a cup.

I

ICOSANDRIA ; from εικοσι, *viginti*, and ανηρ, *maritus*, a husband.

IMBRICATUS, *Caulis, Culmus, Calyx* ; from *imbrex*, a tile.

INANIS *Caulis* ; from *inania*, cobwebs.

INCLUDENS *Calyx* ; from *includo*, to include, or shut up.

INCLUSA *Stamina* ; from *in*, and *claudo*, to shut in.

INCRASSATUS *Pedunculus* ; from *incrasso*, to make thick, to fatten.

INCUMBENS *Anthera* ; from *incumbo*, to lean against.

INERME *Folium* ; from *in* priv. and *arma*.

INFERUS *Flos* ; from *infra* beneath.

INFLATUM *Perianthium* ; from *in*, and *flatus*, a puff, a blast.

INFLEXA *Folia* ; from *inflecto*, to bend inward.

INFUNDIBULIFORMIS *Corolla, nectarium* ; from *infundibulum*, a funnel.

INSERTUS *Petiolus* ; from *inseror*, to be put in.

INSIDENS ; from *insido*, to rest or sit upon.

INTEGERRIMUM *Folium* ; from *integer*, entire.

INTERFOLIACEUS *Pedunculus* ; from *inter*, between, and *folium*, a leaf.

INTORSIO ; from *in*, and *torsio*, writhing.

INTRAFOLIACEÆ *Stipulae* ; from *intra*, within, and *folium*, a leaf.

INUNDATA *loca* ; from *in*, and *unda*, a wave, or water.

- INVOLUCELLUM ; from *involverum*.
 INVOLUCRATUS *Verticillus* ; from *involverum*.
 INVOLUCRUM ; from *in*, and *volvo*, to roll or wrap.
 INVOLUTA *Folia* ; from *in*, and *volvo*, to roll.
 JUNCIFOLIUS ; from *juncus*, a rush, and *folium*.

L

- LABIATUS *Flos* ; from *labium*, a lip.
 LACERUM *Folium* ; from *λακος*, *fissura*, a cleft or fissure.
 LACINIÆ ; from *lacino*, to make holes.
 LACINIATUM *Folium* ; from *lacinia*, a fringe or jag.
 LACTESCENTIA ; from *lac*, milk.
 LACUNOSUM *Folium* ; from *lacuna*, a ditch, a trench.
 LACUSTRIS *Plantæ* ; from *lacus*, a lake.
 LAMELLÆ ; from *lamella*, a small thin plate.
 LANATUM *Folium* ; from *lana*, wool.
 LANCEOLATUM *Folium* ; from *lanceola*, a little lance.
 LATERALES *Flores* ; from *latus*, a side.
 LATERIFOLIUS *Pedunculus* ; from *latus*, a side, and *folium*, a leaf.
 LENTICULARIS *Scabrities* ; dim. from *lens*, a lentil.
 LEPROSUS ; from *lepra*, leprosy.
 LEVIS *Caulis* ; rather *lævis*, smooth.
 LIGNOSUS *Caulis* ; from *lignum*, wood.
 LIGULATUS *Flos* ; from *ligula*, a strap.
 LILIACEÆ ; from *lilium*, the lily.
 LINEA ; *propriè est funiculus ex lino*, a line.
 LINEARE *Folium*, *Pitiohus* ; from *linea*.
 LINEATUM *Folium* ; from *linea*.
 LINGULATUM *Folium* ; from *lingua*, a tongue.
 LITHOPHYTA ; from *λιθος*, *lapis*, a stone, and *φυτον*, *planta*, a
 plant
 LOBATUM *Folium* ; from *λοβος*, *lobus*, the lobe or tip of the ear.
 LOCULUS ; dim. from *locus*, a place.
 LOMENTACEÆ ; from *lomentum*, bean-meal.
 LONGIUSCULUS ; dim. from comp. *longior*.

- LUCIDUM *Folium* ; from *lux*, light.
 LUMBRICIFORMIS ; from *lumbricus*, an earth-worm.
 LUNATUM *Folium* ; from *luna*, the moon.
 LUNULATA *Carina* ; from *lunula* dim. a half-moon.
 LURIDÆ ; from *luridus*, pale, wan.
 LUTEA *Lactescentia* ; from *luteum*, the yolk of an egg.
 LYRATUM *Folium* ; from *lyra*, a harp or lyre.

M

- MARCESCENS *Corolla* ; from *marceo*, to wither.
 MARGO *Folii* ; from *margo*, margin.
 MAS *Planta* ; *etym. incertum*.
 MASCULUS *Flos* ; from *mas*.
 MEDULLA ; from *μυελος*, marrow.
 MEMBRANACEUM *Folium* ; from *membrana*, a membrane.
 METEORICI *Flores solares* ; from *μετεωρος*.
 MINIATUS ; from *minium*, red lead.
 MONADELPHIA ; from *μονος*, *unicus*, one only, and *αδελφος*, *frater* brother.
 MONANDRIA ; from *μονος*, *unicus*, one, and *ανηρ*, *maritus*, a husband.
 MONANGIÆ ; from *μονος*, *unicus*, and *αγος*, *vas*, a vessel, or *loculamentum*.
 MONOCOTYLEDONES ; from *μονος*, *unicus*, one, and *cotyledon*.
 MONŒCIA ; from *μονος*, *unicus*, one, and *οικος*, *domus*, a house.
 MONOGAMIA ; from *μονος*, *unicus*, and *γαμος*, *nuptiæ*, nuptials.
 MONOGYNIA ; from *μονος*, *unicus*, and *γυνη*, *mulier*, a woman.
 MONOPETALA *Corolla* ; from *μονος*, and *πεταλον*, *petalum*.
 MONOPHYLLUM *Intolucrum* ; from *μονος*, *unicus*, one, and *φυλλον*, *folium*, a leaf.
 MONOSPERMA ; from *μονος*, and *sperma*, seed.
 MILIARIS *Scabrities* ; from *milium*, a small grain called *millet*.
 MUCRONATUM *Folium* ; from *μακρος*, *longus*, long.
 MULTIFIDUM *Folium* ; from *multus*, many, and *findo*, to cleave, or divide.

- MULTIFLORUS *Pedunculus* ; from *multus*, many, and *flos*, a flower.
 MULTIPARTITUM *Folium* ; from *multus*, many, and *partitus*, divided.
 MULTIPLEX *Corolla* ; from *multus*, many, and *plicare*, to fold.
 MULTILOULARIS *Drupa* ; from *multus*, and *loculus*, a little cell.
 MULTISILIQUÆ ; from *multus*, many, and *siliqua*, a pod.
 MURICATUS *Caulis* ; from *murex*, a fish, whose shell is covered with sharp points, or prickles.
 MUSCARIOSUS ; from *musca*, a fly.
 MUSCI ; from *μωσχος*, *vitulus* ; properly any thing young, new, or fresh.
 MUTICA *Gluma* ; from *mutilus*, broken off.

N

- NATANS *Folium* ; from *nato*, to swim.
 NAVICULARIS *Vulvula* ; dim. from *navis*, a ship.
 NECTARIUM ; from *nectar*, honey.
 NERVOSUM *Folium* ; from *nervus*, a nerve, or string.
 NIDULANTIA *semina*, *Bacca* ; from *nidus*, a nest.
 NUCAMENTACEÆ ; from *nucamentum*, a cat's tail, or long excrescence hanging down from the pine, fir, &c.

O

- OBCONICUM *Nectarium* ; from *ob*, and *conus*, a cone, a geometrical figure, like a sugar-loaf.
 OBCORDATUM *Petalum* ; from *ob*, and *cordatum*, heart-shaped.
 OBLIQUUM *Folium* ; from *ob* *liquus*, transverse.
 OBOVATUM *Folium* ; from *ob*, and *ovum*, an egg.
 OBTUSUM *Folium* ; from *obtundor*, to be blunted at the point.
 OBVOLUTUM *Folium* ; from *ob*, and *volvo*, to roll.
 OCTANDRIA ; from *ὀκτώ*, *octo*, eight, and *ἀνῆρ* *maritus*, a husband.
 OFFICINALIS ; from *officina*, a shop.
 OLIGANTHERE ; from *ὀλιγός*, *exiguus*, small, few, and *anthera*.
 OPERCULATA *Anthera* ; from *operculum*, a cover.

OPPOSITIFOLIUS *Pedunculus*; from *oppositum*, opposite, and *folium*, a leaf.

ORBICULATUM *Folium*; from *orbis*, an orb, or circle.

ORCHIDEÆ; from *orchis*, the first *genus* in the class *Gynandria*.

ORGYA; from *οργυια*, *orgyia*, six foot.

ORGYALIS *Caulis*; from *οργυια*, *idem*.

OVARIVM; from *ovum*.

OVATUM *Folium*; from *ovum*, an egg.

P

PAGINA *Folii*; from *pagina*, the page of a book.

PALEACEUS *Pappus*; from *palea*, short straw, or chaff.

PALMÆ; from *παλαμη*, the palm of the hand.

PALMARIS *Caulis*; from *palmus*.

PALMATA *Radix*; from *palma*, a hand.

PALMUS; from *palma*, the palm of the hand.

PALUSTRIS; from *palus*, a fen or marsh.

PANDURIFORME *Folium*; from *pandura*, a musical instrument.

PANICULA; from *panus*, a woof about the quill in the shuttle.

PAPILIONACEUS; from *papilio*, a butterfly.

PAPILLOSUM *Folium*; from *papilla*, the nipple.

PAPULOSUM *Folium*; from *papula*, a pimple.

PARASITICUS *Caulis*; from *parasitus*, a parasite.

PAUCIFLORIS; from *pauci*, few, and *flos*, a flower.

PEDATUM *Folium*; from *pes*, a foot.

PEDICELLUS; from *pediculus*, a little foot.

PEDICULUS; dim. from *pes*, a foot.

PEDUNCULARIS *Cirrus*; from *pedunculus*.

PEDUNCULATI *Flores*; from *pedunculus*.

PEDUNCULUS; from *pedo*, one who is splay-footed.

PELTATUM *Folium*; from *pelta*, a target.

PENNATIFOLIÆ; from *penna*, a large feather, and *folium*, a leaf.

PENDULA *Radix*; from *pendeo*, to hang.

PENICILLATUM *Stigma*; from *penicillus*, a pencil.

PENTAGYNIA; from *πεντε*, *quinque*, five, and *γυνη*, *mulier*, a woman.

- PENTANDRIA; from *πεντε*, *quinque*, five, and *ανηρ*, *maritus*, a husband.
- PENTANGIÆ; from *πεντε*, *five*, and *αγχος*, *vas*, a vessel, or *loculamentum*.
- PENTAPETALA *Corolla*; from *πεντε*, *quinque*, and *πεταλον*, *petalum*.
- PENTAPHYLLUS *Calyx*; from *πεντε*, *quinque*, and *φυλλον*, *folium*, a leaf.
- PERENNIS *Radix, folium*; from *per*, *by*, and *annus*, a year.
- PERFOLIATUM *Folium*; from *per* and *folium*.
- PERFORATÆ *Cotyledones*; from *perforor*, to be pierced through.
- PERIANTHIUM; from *περι*, *circum*, about, and *ανθος*, *flos*, a flower.
- PERICARPIUM; from *περι*, *circum*, and *καρπος*, *semen*, seed.
- PERICHÆTIUM; from *περι*, and *χαιτη*, *juba*.
- PERSISTENS *Folium*; from *persisto*, to abide.
- PERSONATÆ; i. e. *personam gerens*, masked.
- PETALIFORMIA *Stigmata*; from *petalum*.
- PETALODES *Flos*; from *petalum*.
- PETALUM; from *πεταω*, *pando*, to expand.
- PETIOLARIS *Cirrus*; from *petiolus*.
- PETIOLATUM *Folium*; from *petiolus*, a foot-stalk.
- PETIOLUS; dim. from *pede*, *quasi pediolus*, a little foot, or from *pettilus*, slender.
- PILEUS *Fungi*; from *πιλος*, *lana coacta*.
- PILOSUM, *Folium*; from *πιλος*, *pilus*, a hair.
- PINNATIFIDUM *Folium*; from *πιννα*, a wing.
- PINNATUM *Folium*; from *pinna*, the large feathers of a wing.
- PIPERATUS; from *piper*, pepper.
- PIPERITÆ; from *piper*, pepper.
- PIXIDATUM *Folium*; from *pixis*, a box.
- PLACENTATIO; from *placenta*.
- PLANIPETALUS *Flos*; from *planus*, plane, flat, and *petalum*, a petal.
- PLANUM *Folium*; from *απλανης*, *planus*.
- PLICATUM *Folium*; from *plico*, to fold.
- PLUMATA *Seta*; from *pluma*, a soft feather:
- PLUMOSUS *Pappus*; from *pluma*, a small soft feather.
- POLLEN; from *παλη*, fine meal, or flour.

- POLLICARIS *Caulis*; from *poller*, a thumb.
- POLYADELPHIA; from *πολυς*, *multus*, many, and *αδελφος*, *frater*, a brother.
- POLYANDRIA; from *πολυς*, *multus*, many, and *ανηρ*, *maritus*, a husband.
- POLYANGIÆ; from *πολυς* many, and *αγχος*, *vas*, a vessel, or *locumentum*.
- POLYCOTYLEDONES; from *πολυς*, and *cotyledon*.
- POLYGAMIA; from *πολυς*, *multus*, many, and *γαμος*, *nuptiæ*, nuptials.
- POLYGYNIA; from *πολυς*, *multus*, many, and *γυνη*, *mulier*, a woman.
- POLYMORPHA; from *πολυς*, *multus*, and *μορφη*, *forma*, shape.
- POLYPETALA *Corolla*; from *πολυς*, *multus*, and *πεταλον*, *petalum*.
- POLYPHYLLUM *Involucrum*; from *πολυς*, *multus*, many, and *φυλλον*, *folium*, a leaf.
- POLYSTACHIUS *Culmus*; from *πολυς*, and *σταχυς*, *spica*.
- POMACEÆ; from *pomum*, an apple, pear, &c.
- PORI; from *πειρω*, *transadigo*, to pierce through.
- POSTICUS *Angulus*; from *post*, ut *anticus* ab *ante*.
- PRATENSIS; from *pratium*, a meadow.
- PRÆCOX; from *præ* et *coquo*, to cook.
- PRÆMORSA *Radix*; from *præmordeo*, to bite.
- PRECIÆ, from *precius*, early.
- PRISMATICUS *Calyx*; from *prisma*, a prism.
- PROLIFER *Flos*; from *proles*, offspring.
- PROMINULUM *Dissepimentum*; from *promineo*, to jet or stand out.
- PRONUM *discum folii*; from *προνος*, *antiq.* having the face downwards.
- PSEUDO; from *ψευδω*, *fallo*, to deceive.
- PULPOSUM *Folium*, from *pulpa*, the pulp, or fleshy part of meat.
- PULVERATUM, or *pulverulentum, folium*; from *pulvis*, powder, dust.
- PULVINATUS *Pileus*; from *pulvinar*, a pillow or cushion.
- PUMILA; from *pumilus*, or rather *pumilio*, a dwarf.
- PUNCTATUM *Folium*; from *punctum*, a point.
- PUTAMINEA; from *putamen*, a shell.

Q

- QUADRIDENTATUS *Pappus*; from *quatuor*, four, and *dens*, a tooth.
- QUADRIFIDUM *Folium*; in *quatuor partes fissus*, four-cleft.
- QUADRIJUGUM *Folium*; from *quatuor*, and *jugo*, to yoke.
- QUADRILOBUM *Folium*; from *quatuor*, and *λοβος*, the tip of the ear.
- QUADRILOCULARIS *Bacca*; from *quatuor*, and *loculus*, a little place.
- QUADRIPARTITUM *Folium*; from *quatuor*, and *partitus*, divided.
- QUINATUM *Folium*; from *quinus*, five.
- QUINQUECOCCUS *fructus*; from *quinque*, and *κοκκος*, *granum*, a grain.
- QUINQUEJUGUM *Folium*; from *quinque*, and *jugo*, to yoke.
- QUINQUELOBUM *Folium*; from *quinque*, five, and *λοβος*, the tip of the ear.
- QUINQUEPARTITUM *Folium*; from *quinque*, and *partitus*, divided.
- QUINQUEFIDUM *Folium*; in *quinque partes fissum*, five-cleft.

R

- RACEMUS; from *ramus*, or from *radendo*.
- RACHIS; from *ραχις*, *dorsum*, the back; or rather, *spina dorsi*, the back-bone.
- RADIATUS *Flos*; from *radius*.
- RADICALIA *Folia*; from *radix*, a root.
- RADICANS *Caulis*; from *radicor*, to take root.
- RADICATUM *Folium*; from *radix*.
- RADICULA; dim. from *radix*.
- RAMEA *Folia*; from *ramus*, a branch.
- RAMOSISSIMUS *Caulis*; from *ramus*, a branch.
- RAMOSUS *Caulis*; from *ramus*, a branch.
- RAMUS; from *ῥαμνος*, a small branch.
- RECLINATUM *Folium*; from *reclino*, to bend.
- RECURVATUM *Folium Petiolus*; from *recurvo*, to bend back.
- REFLEXUS *Ramus*; from *reflecto*, to bend back.

- REMOTUS *Verticillus*; from *removeo*, to remove.
 RENIFORME *Folium*; from *ren*, a kidney.
 REPANDUM *Folium*; from *re*, and *pando*, to bend.
 REPENS *Rudix*; from *repo*, to creep.
 REPTANS *Flagellum*; from *repto*, to creep.
 RESUPINATUM *Folium*; from *resupino*, to turn upwards.
 RETICULATA *Petala*; from *rete*, a net.
 RETROFLEXUS *Ramus*; from *retro*, backward, and *flexus*, bent.
 RETROFRACTUS *Pedunculus*; from *retro*, backwards, and *frangor*,
 to be broken.
 RETUSUM *Folium*; from *retundor*, to be blunted.
 REVOLUTUM *Folium*; from *revolvo*, to roll back.
 RHÆADES; from *rhæus*, the red poppy.
 RHOMBEUM *Folium*; from *rhombus*, a geometrical figure of four
 equal sides, but not right-angled.
 RHOMBOIDEUM *Folium*; from *rhomboides*, a geometrical figure
 whose sides and angles are unequal.
 RINGENS; from *ῥίη*, *nasus*, a nose.
 ROSACEUS *Flos*; from *rosa*, a rose.
 ROSTRATUS *fructus*; from *rostrum*, the beak of a bird.
 ROTACEÆ; from *rota*, a wheel.
 ROTATUS *Limbus Corollæ*; from *rota*, a wheel.
 RUDERATIS, *locis* understood; from *rudis*, rubbish.
 RUGOSUM *Folium*; from *ruga*, a wrinkle.
 RUNCINATUM *Folium*; from *runcina*, a large saw.

S

- SAGITTATUM *Folium*; from *sagitta*, an arrow.
 SANGUINEA; from *sanguis*, blood.
 SARMENTACEÆ; from *sarmentum*, a twig or spray of a vine.
 SARMENTOSUS *Caulis*; from *sarmentum*, the twig of a vine.
 SCABRIDÆ; from *scaber*, rough, rugged.
 SCABRITIES; from *scaber*, rough.
 SCAPUS; from *σκηπῶ*, *innitor*, to lean upon.
 SCARROSUM *Folium*; from *scarreo*, to be rough.

- SCITAMINA ; from *situs*, fair, beautiful, or from *scitamentum*, meat of a pleasant taste.
- SCORPIOIDES *Flos* ; from *scorpio*, a scorpion.
- SCUTELLUM ; from *scutum*, a target.
- SCYPHIFER ; from *σκυφος*, *scyphus*, a cup, and *fero*, to bear.
- SECRETORIA *Scabrities* ; from *secerno*, to separate.
- SECUNDA *Spica* ; from *sequendo*, to follow.
- SECURIFORMIS *Pubescentia* ; from *securis*, an axe or hatchet.
- SEGREGATA *Polygamia* ; *segrego*, to separate.
- SEMINALE *Folium* ; from *semen*, seed.
- SEMPERVIRENS *Folium* ; from *semper*, and *virens*, green.
- SENA *Folia* ; from *sex*, six.
- SENTICOSÆ ; from *sentis*, a brier or bramble.
- SEPIARIÆ ; from *sepes*, a hedge.
- SEPTENIS *foliolis*, from *septem*, seven.
- SEPTICUS ; from *σηπω* *putrefacio*, to rot.
- SEPTUM ; from *sepio*, to enclose.
- SERICEUM *Folium* ; from *sericum*, silk.
- SERPYLLIFOLIA ; from *serpyllum*, thyme, and *folium*.
- SERRATUM *Folium* : from *serra*, a saw.
- SESSILE *Folium* ; from *sedeo*, to sit.
- SETÆ ; from *χαιτη*, *juba*, a horse's mane.
- SETACEUM *Folium* ; from *seta*, a bristle.
- SEXFIDUM *Nectarium* ; from *sex*, six, and *fissum*, cloven.
- SILICULA ; dim. from *siliqua*, a pod.
- SILICULOSA ; from *silicula*, a little pod.
- SILIQUOSA ; from *siliqua*, a pod.
- SINUATUM *Folium* ; from *sinus*, a hollow.
- SOLITARIUS *Pedunculus* ; from *solus*, alone.
- SOLUTÆ *Stipulæ* ; from *solvor*, to be loosed.
- SPADICEUS ; from *spadix*.
- SPATHA ; from *σπατος*, *corium*, skin.
- SPATHACEÆ ; from *spatha*, a sheath, in the language of botany.
- SPATULATUM *Folium* ; from *spathula*, an instrument used to spread salve.
- SPICA ; from *σταχυς*, *Æolice* *σταχυς*, an ear of corn.

- SPICILLA ; dim. from *spica*.
 SPICULA *Graminibus* ; dim. from *spica*.
 SPINESCENS *Petiolus, Stipula* ; from *spina*, a thorn.
 SPINOSUS *Caulis Folium* ; from *spina*, a thorn.
 SPIRALES *Cotyledones* ; from *spira*, a circle, the coil of a cable, &c.
 SPITHAMEUS *Caulis* ; from *spithama*, a span.
 SQUAMOSA *Radix* ; from *squama*, a scale.
 SQUARROSUM *Folium* ; from *ισχαρα, scarra*, or from *squarra*, scurf.
 STAMINÆUS *Flos* ; from *stamen*.
 STATUMINATÆ ; from *statumen*, a prop, a support.
 STELLATA *Folia* ; from *stella*, a star.
 STIGMA ; from *στιζω, signum quod inuritur*, a brand.
 STIMULI : from *στιγμος, stimulus*, per sync. *stimulus*.
 STIPATUS, or *stipitatus, Pappus* ; from *stipes*.
 STIPES ; from *στυπος*, a stump.
 STIPULA ; from *stipa*, tow.
 STIPULARIS *Gemma* ; from *stipula*.
 STIPULATIO ; from *stipula*.
 STIPULATUS *Caulis* ; from *stipula*.
 STOLONIFERUS *Truncus Caulis* ; from *stolo*, a shoot, or scion.
 STRIATUS *Caulis, Culmus, Folia* ; from *stria*, a slight groove.
 STRICTUS *Caulis, Culmus, Folia* ; from *stringo*, to tie fast.
 STRIGÆ ; from *strigo*, pro *stringo*, to grasp, to tighten.
 STRIGOSUM *Folium* ; from *strigando*, standing still, *quippe bos præmacie*.
 STYLUS ; from *στυλος, columna*, a pillar.
 SUBCAULIS ; from *sub* and *acaulis*.
 SUBALARIA *Folia* ; from *sub*, under, and *ala*, a wing.
 SUBEROSUS *Caulis folia* ; from *sub*, and *erodor*, to be eaten into.
 SUBEXCEDENS *Calyx* ; from *sub*, and *excedo*, to surpass.
 SUBMERSUM *Folium* ; from *submergo*, to sink under water.
 SUBRAMOSUS *Caulis* ; from *sub*, and *ramus*, a branch.
 SUBRENIFORMUM *Folium* ; from *sub*, pro *fere*, and *ren*, a kidney.
 SUBROTUNDUM *Folium* ; from *sub*, near to, and *rotundum*, round.
 SUBULATUM *Folium* ; from *subula*, an awl.
 SUCCULENTÆ ; from *succus*, juice.
 SUFFRUTEX ; from *sub*, and *frutex*, a shrub.

- SUFFRUTICOSUS *Caulis*; from *sub*, and *frutex*, a shrub.
 SULCATUS *Caulis, Culmus, Folia*; from *sulcus*, a furrow.
 SUPERFICIES *Folii*; from *super*, i. e. *supra*, and *facies*, a face.
 SUPERUS *Flos*; from *super*, above.
 SUPRA-AXILLARIS *Pedunculus*; from *supra*, above, and *axilla*, the arm-pit.
 SUPRAFOLIACEUS *Pedunculus*; from *supra*, above, and *folium*, a leaf.
 SYLVESTRIS; from *sylva*, a wood or forest.
 SYNGENESIA; from *συν*, *cum*, or *simul*, together, and *γενεσις*, *generatio*.

T

- TERGEMINUM *Folium compositum*; from *ter*, thrice, and *geminus*, double.
 TERNATUM *Folium*; from *ternus*, three.
 TESSELLATUM *Folium*; from *tessella*, the square pieces of wood or stone used in making checkered work.
 TETRADYNAMIA; from *τεσσαρες*, *quatuor*, and *δυναμις*, *potentia*, power.
 TETRAGYNIA; from *τεσσαρες*, *quatuor*, and *γυνη*, *mulier*, a woman.
 TETRANDRIA; from *τεσσαρες*, *quatuor*, four, and *ανηρ*, *maritus*, a husband.
 TETRANGLÆ; from *τεσσαρες*, four, and *αγγος*, *vas*, a vessel, or *loculamentum*.
 TETRAPETALA *Corolla*; from *τεσσαρες*, *quatuor*, and *πεταλον*, *petalum*.
 TETRAPHYLLUS *Calyx*; from *τεσσαρες*, and *φυλλον*, *folium*, a leaf.
 TETRASPERMA *Planta*; from *τεσσαρες*, and *sperma*, seed.
 THALAMUS; from *θαλαμος*, a bed, a chamber.
 TOMENTOSUS *Caulis Folia*; from *tomentum*, short wool, shorn off.
 TOMENTUM; from *τομιον*, *frustum*, a fragment, *seu quod sectione est ablatum*.
 TOROSUM *Pericarpium*; from *torus*, protuberance or swelling, as of the veins.

- TORTA *Covolla* ; from *torqueo*, to writhe, torture.
- TORTILIS *Arista* ; from *torqueo*, to writhe, or twist.
- TORULOSA *Siliqua* ; from *torulus*, dim. from *torus*.
- TRAPEZIFORMIUM *Folium* ; from *trapezium*, a geometrical quadrangle, whose sides are neither equal nor opposite.
- TRIANDRIA ; from *τρεις*, *tres*, three, and *ανηρ*, *maritus*, a husband.
- TRIANGLÆ ; from *τρεις*, three, and *αγγος*, *vas*, a vessel.
- TRICOCCA *Capsula* ; from *κοκκος*, *granum*, grain.
- TRIFIDUM *Folium* ; *in tres partes fissum*.
- TRIFLORUS *Pedunculus* ; from *tres*, and *flos*, a flower.
- TRIGLOCHIDES *Hami* ; from *tres*, and *γλωχίς*, *cuspis*, a point.
- TRIGONUS *Caulis* ; from *τρεις*, *ter*, and *γωνία*, *angulus*, an angle.
- TRIGYNIA ; from *τρεις*, *tres*, three, and *γυνή*, *mulier*, a woman.
- TRIHILATÆ ; from *hilum*, the black of a bean.
- TRIJUGUM *Folium* ; from *tres*, and *jugo*, to yoke.
- TRILOBUM *Folium* ; from *tres*, three, and *λοβος*, the tip of the ear.
- TRINERVIUM *Folium* ; from *tres*, three, and *nervus*, a nerve or string.
- TRIŒCIA ; from *τρεις*, *tres*, three, and *οικος*, *domus*, a house.
- TRIPARTITUM *Folium* ; from *tres*, and *partitus*, divided.
- TRIPETALA *Corolla* ; from *τρεις*, *tres*, and *πεταλον*, *petalum*.
- TRIPETALOIDEÆ ; from *tres*, and *petalum*.
- TRIPHYLLUS *Calyx* ; from *τρεις*, and *φυλλον*, a leaf.
- TRIPINNATUM *Folium compositum* ; from *tres*, three, and *pinnatus*, winged.
- TRIPLINERVIUM *Folium* ; from *triplex*, triple, and *nervus*, a nerve or string.
- TRIQUETRUM *Folium Caulis* ; quasi *triquadrus*, i. e. *quadratus in tres angulos*.
- TRISPERMA ; from *tres* and *sperma*, seed.
- TRITERNATUM *Folium compositum* ; from *tres*, three, and *ternus*, threefold.
- TRIVALVE *Pericarpium* ; from *tres*, and *valvæ*, doors or valves.
- TRUNCATUM *Folium* ; from *truncus*, a stump.
- TUBERCULATUS ; from *tuberculum*, a little pimple or tubercle.
- TUBEROSA *Radix* ; from *tuber*, a knob.

- TUBULOSI *Flosculi* ; from *tuba*, a tube.
 TUNICATUS *Caulis, Radix* ; from *tunica*, a coat.
 TURBINATUM *Pericarpium* ; from *turbo*, a top.
 TURIO ; from *tyro*, a novice.

V

- VAGÆ ; from *vagor*, to wander.
 VAGINALES ; from *vagina*, a sheath.
 VAGINANS *Folium* ; from *vagina*, a sheath.
 VAGINATUS *Caulis Culmus* ; from *vagina*, a sheath.
 VASA ; from *vescendo*, to be eaten, *quod in ea vesce ponantur*.
 VEGETABILIA ; from *vegeto*, to quicken.
 VENOSUM *Folium* ; from *vena*, a vein.
 VENTRICOSA *Spica* ; from *venter*, the belly.
 VENTRICULOSUS *Calyx* ; dim. from *venter*, the belly.
 VEPREULÆ ; dim. from *tepres*, a brier or bramble.
 VERNATIO ; from *ver*, the spring.
 VERRUCOSA *Capsula* ; from *verruca*, a wart.
 VERSATILIS *Anthera* ; from *verto*, to turn.
 VERTICALIA *Folia* ; from *vertex*, the top of any thing.
 VERTICILLATI *Rami, Flores, Folia* ; from *verticillum*, an axis or spindle.
 VERTICILLUS ; from *vertex*, a whirlpool.
 VESICULARIS *Scabrities* ; from *vesica*, a bladder.
 VILLOSUS *Caulis, Folium* ; from *villus*, wool.
 VIRGATUS *Caulis* ; from *virga*, a rod.
 VISCIDUM *Folium* ; from *viscus*, glue.
 VISCOSITAS ; from *viscus*, glue.
 ULIGINOSA *Loca* ; from *uligo*, the natural moisture of the earth.
 UMBELLA ; dim. from *umbra*, a shadow.
 UMBELLULA ; dim. from *umbella*.
 UMBILICATUM *Folium* ; from *umbilicus*.
 UNANGULATUS *Caulis* ; from *unus*, and *angulus*.
 UNCINATUM *Stigma* ; from *uncipus*, an instrument hooked at the end.
 UNDATUM *Folium* ; from *unda*, a wave.

- UNDULATA *Corolla* ; from *undula*, dim. from *unda*, a wave.
UNGUICULARIS *Caulis* ; from *unguis*, a nail of the hand, &c.
UNGUIS ; from *οὐξ*, *idem*.
UNIFLORUS *Pedunculus* ; from *unus*, one, and *flos*, a flower.
UNIFOLIUM ; from *unus*, one, and *folium*, a leaf.
UNILATERALIS *Racemus* ; from *unus*, one, and *latus*, a side:
UNILOCULARIS, *Capsula* ; from *unus*, one, and *loculus*, a little cell.
UNISPERMA *Bacca* ; from *unus*, and *sperma*, seed.
VOLUBILIS *Caulis* ; from *volvo*, to roll.
URCEOLATA *Corolla* ; from *urceolus*, from *urceus*, a pitcher.
URENS *Caulis* ; from *uro*, to burn.
UTRICULI ; from *uter*, a bag, or bottle.
VULGARIS ; from *vulgus*, the common people.



PLATE I.

PARTS OF THE FLOWER.

VIDE PART I. CHAP. I. p. 1.

Fig.

1. A FLOWER, with its *Corolla*, *Pistillum*, and *Stamina*: A, the *Petals* of the *Corolla*; b, the *Germen*; c, the *Style*; d, the *Stigma*; e, the *Filaments*; f, the *Antheræ*.
2. The *Pistillum* and *Stamina*, separate from the *Corolla*: b, the *Germen*; c, the *Style*; d, the *Stigma*; c, c, the *Filaments*, with the *Antheræ* bursting and discharging the *Pollen*.
3. A Flower, whose *Corolla* is *Monopetalous*: A, the *Corolla*; B, the *Perianthium*.
4. A *Polypetalous Corolla*: A, the *Unguis*; B, the *Lamina*.
5. A *Narcissus*, issuing from its *Spatha*: A, the *Flower*; B, the *Spatha*.
6. An *Amentum*.
7. The *Fructification* of a *Moss*: A, the *Calyptra*.
8. A *Fungus*: A, the *Volva*.
9. A *Grass*: A, the *Ghuma*; B, the *Arista*.
10. A *Compound Umbel*: A, the *Universal Umbel*; B, the *Umbellulæ*, or *Partial Umbels*; C, the *Universal Involucrum*; d, the *Partial Involucra*.
11. A *Bractea*, accompanying the flowers of the *Tilia*: A, the *Bractea*.
12. A, the *Pollen*, seen with a microscope; B, an elastic vapour discharged from it.








PLATE II.

PARTS OF THE FRUIT.

VIDE PART I. CHAP. XV. p. 29.

Fig.

1. A *Capsule*: A, the *Valvules*.
2. A, A, a *Receptacle* of the seeds.
3. A *Strobilus*.
4. A *Winged Seed*: A, the *Seed*; b, the *Wing*.
5. A *Legumen*: A, the *Under Sutures*, along which are affixed the seeds.
6. A *Siliqua*: A, B, the two sutures, to which the seeds are fastened alternately.
7. A *Seed*, crowned with a *Pappus*: A, the *Seed*; B, the *Stipes* of the *Pappus*; c, d, a *Feathery Pappus*.
8. The *Seed* of a *Bean*, split in two: A, A, the *Cotyledons*; b, the *Corculum*; c, the *Rostellum*; d, the *Plumula*.
9. A *Drupa*: A, the *Nucleus*, or *Stone*; B, the *Pulp*.
10. A *Pomum*: A, the *Capsule*; B, the *Pulp*.
11. A *Berry*: A, the *Seeds*; B, B, the *Pulp*.
12. A *Seed*, crowned with a *Calyculus*: A, the *Seed*; B, the *Calyculus*.







Drawn by Henderson. Engraved by Weston.

Published Dec. 20-1869 by Sherwood, Nedy & Jones

PLATE III.

CLASSES.

VIDE PART II. CHAP. I. p. 48.

Class.

1. *Monandria.*
2. *Diandria.*
3. *Triandria.*
4. *Tetrandria.*
5. *Pentandria.*
6. *Hexandria.*
7. *Heptandria.*
8. *Octandria.*
9. *Enneandria.*
10. *Decandria.*
11. *Dodecandria.*
12. *Icosandria.*
13. *Polyandria.*
14. *Diynamia.*
15. *Tetradynamia.*
16. *Monadelpkia.*
17. *Diadelphia.*
18. *Polyadelphia.*
19. *Syngenesia.*
20. *Gynandria.*
21. *Monœcia.*
22. *Diœcia.*
23. *Polygamia.*
24. *Cryptogamia.*

PLATE IV.

ROOTS.

VIDE PART III. CHAP. II. p. 415.


Fig.

1. A *Scaly Bulb*, as in the White Lily.
2. A *Solid Bulb*, as in the Crocus.
3. Transverse section of a *Coated Bulb*.
4. A *Tuberous and Pendulous Root*, as in Piony.
5. A *Branched Root*.
6. A *Simple tapering Root*, as in the Carrot.
7. A *Creeping Root*.

1



2



3



4



5



6



7







PLATE V.

TRUNK.

VIDE PART II. CHAP. IV. p. 41.

Fig.

1. A *Squamose Stem.*
2. A *Voluble Stem*, vide also Fig. 4.
3. A *Frons.*
4. A *Scapus.*
5. An *Articulate Stem.*
7. A *Dichotomous Stem.*
8. A *Brachiate Stem.*

PLATE VI.

SIMPLE LEAVES.

VIDE PART III. CHAP. V. p. 423.

Fig.

1. *Orbiculate.*
2. *Subrotund.*
3. *Ovate.*
4. *Oval.*
5. *Oblong.*
6. *Lanceolate.*
7. *Linear.*
8. *Subulate.*
9. *Reniform.*
10. *Cordate.*
11. *Lumulate.*
12. *Triangular.*
13. *Sagittate.*
14. *Cordato-sagittate.*
15. *Hastate.*
16. *Cloven.*
17. *Trilobe.*
18. *Præmorse.*
19. *Lobate.*
20. *Quinquangular.*
21. *Erose.*
22. *Palmate.*
23. *Pinnatifid.*
24. *Laciniate.*
25. *Sinuate.*
26. *Dentato-sinuate.*
27. *Retrorsum-sinuate.*
28. *Partite.*
29. *Repand.*
30. *Dentate.*



Drawn by H. Schimper. Engraved by W. Smith.

Published Dec. 10 1844 by Sherwood, Naylor, & Co.





PLATE VII.

SIMPLE LEAVES CONTINUED.

VIDE PART III. CHAP. V. p. 427.

Fig.

1. *Serrate.*
2. *Duplicato-serrate.*
3. *Duplicato-crenate.*
4. *Cartilagineous.*
5. *Acutely crenate.*
6. *Obtusely crenate.*
7. *Plicate.*
8. *Crenate.*
9. *Crisp.*
10. *Obtuse.*
11. *Acute.*
12. *Acuminate.*
13. *Obtuse, with an Acumen.*
14. *Acutely-emarginate.*
15. *Cuneiform-emarginate.*
16. *Retuse.*
17. *Pilose.*
18. *Tomentose.*
19. *Hispid.*
20. *Ciliate.*
21. *Rugose.*
22. *Venose.*
23. *Nervose.*
24. *Pappillose.*
25. *Linguiform.*
26. *Acinaciform.*
27. *Dolabriform.*
28. *Deltoid.*
29. *Triquetrous.*
30. *Canaliculate.*

PLATE VIII.

SIMPLE LEAVES CONTINUED.

Fig.

1. *Sulcate.*
2. *Teretes.*

COMPOUND LEAVES.

VIDE PART III. CHAP. VI. p. 432:

3. *Binate.*
4. *Ternate, with the folioles sessile.*
5. *Ternate, with the folioles petiolate.*
6. *Digitate.*
7. *Pedate.*
8. *Pinnate, with an odd one.*
9. *abrupt.*
10. *alternately.*
11. *interruptedly.*
12. *cirrhone.*
13. *conjugate.*
14. *decursively.*
15. *articulately.*
16. *Lyrate.*
17. *Biternate.*
18. *Bipinnate.*
19. *Triternate.*












PLATE IX.

COMPOUND LEAVES CONTINUED.

Fig.

1. *Tripinnate abrupt.*
2. with an *odd one.*

DETERMINATE LEAVES.

3. A, *inflex*; B, *erect*; C, *patent*; D, *horizontal*; E, *reclined*; F, *revolute.*
4. A, *seminal*; B, *cauline*; C, *ramous*; D, *floral.*
5. A, *peltate*; B, *petiolate*; C, *sessile*; D, *decurrent*; E, *amplexicaul*; F, *perfoliate*; G, *connate*; H, *vaginant.*
6. A, *articulate*; B, *stellate*; C, *quatern*; D, *opposite*; E, *alternate*; F, *acerose*; G, *imbricate*; H, *fasciculate.*
7. *Parabolic.*
8. *Spatulate.*


PLATE X.

FULCRA.

VIDE PART III. CHAP. III. p. 438.

Fig.

1. A, a *Cirrhus*; C, *Stipula*, with *Concave Glandules* on the leaf.
2. *Pedicellate Glandules*.
3. A, a *Bractea*, differing from the *Leaves*; B, the *Leaves*.
4. A, *Simple Spines*; B, a *Triple Spine*.
5. A, *Simple Aculei*; B, *Triple Aculei*.
6. A, A, *Opposite Leaves*; B, the *Axilla*.



Drawn by Henderson, Engraved by Weston.
Published Dec. 20. 1819 by Sherwood Nedy & Jones.







PLATE XI.

FOLIATION.

VIDE PART III. CHAP. XVI. p. 148.

Fig.

1. *Convolute.*
2. *Involute.*
3. *Revolute.*
4. *Conduplicate.*
5. *Equitant.*
6. *Imbricate.*
7. *Obvolute.*
8. *Plicate.*
9. *Convolute.*
10. *Involute opposite.*
11. *alternate.*
12. *Revolute opposite.*
13. *Equitant ancipital.*
14. *triquetrous.*

PLATE XII.

MISCELLANEOUS.

Fig.


1. A *Corymbus*.
2. Fruit of the *Lily*: A, the *Valvules* of the *Capsule*; B, a *Seed*; C, the *Arillus* opened to discover the *Seed*.
3. A *Verticillus*.
4. A, A, the *Horned Nectaria* in the *Aconitum*; B, B, two *Peduncles* that support them.
5. A *Paleaceous Receptacle* of a *Compound Flower*, shown in *Rudbeckia*: A, A, the *Paleæ* that parts the *Florets* of the *Disk*; B, the *Tubulose Florets* of the *Disk*; C, the *Ligulate Corollulæ* of the *Radius*; D, a *Ligulate Corollula* fallen off.
6. A, A, a *Spatha*; B, a *Spadix*.
7. A *Racemus*.
8. A *Tubulose Floret* of a *Compound Flower*.
9. A *Monopetalous Hypocrateriform Corolla*.
10. A *Nectarium* that crowns the *Corolla*, shown in the *cup* of a *Narcissus*.
11. A *Spike*.
12. A *Calycine Nectarium*, shown in the *flower* of a *Tropæolum*: A, the *Nectarium*.
13. A *Nectarium* of singular construction, shown in a *flower* of the *Parnassia*: A, five heart-shaped *Nectaria*, terminated by threads, each of which is crowned with a little ball.
14. A *Cyma* of the *Laurustinus*.
15. A *Panicle*.

THE END.

1




2



3



4



5



11



9



10



8




15




14



12



13



Drawn by A. G. ... Engraved by W. ...

Published Dec. 20 - 1843, by Sherman & Co., No. 115 S. 3rd St.




KING'S
College
LONDON

GUYPO QK92 LEE

Library
LEE, JAMES
AN ENTRODUCTION TO THE
SCIENCE OF ROTARY
1810

201001183 9



KING'S COLLEGE LONDON



