

**AD-A276 728**



2

**PL-TR-93-2155**

**UNDERGROUND NUCLEAR EXPLOSIONS AT  
AZGIR, KAZAKHSTAN, AND IMPLICATIONS  
FOR IDENTIFYING DECOUPLED NUCLEAR  
TESTING IN SALT**

**Lynn R. Sykes**

**Columbia University in the City of New York  
Box 20  
Low Memorial Library  
New York, NY 10027**

**DTIC  
ELECTE  
FEB 24 1994  
S B D**

**28 June 1993**

**DTIC QUALITY INSPECTED 2**

**Scientific Report No. 2**

**Approved for public release; distribution unlimited.**



**PHILLIPS LABORATORY  
Directorate of Geophysics  
AIR FORCE MATERIEL COMMAND  
HANSCOM AIR FORCE BASE, MA 01731-3010**

94-06027


1299 94-06027




**Best  
Available  
Copy**

The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the Air Force or the U.S. Government.

This technical report has been reviewed and is approved for publication.

  
\_\_\_\_\_  
JAMES F. LEWKOWICZ  
Contract Manager  
Solid Earth Geophysics Branch  
Earth Sciences Division

  
\_\_\_\_\_  
JAMES F. LEWKOWICZ  
Branch Chief  
Solid Earth Geophysics Branch  
Earth Sciences Division

  
\_\_\_\_\_  
DONALD H. ECKHARDT, Director  
Earth Sciences Division

This document has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify PL/IMA, 29 Randolph Road, Hanscom AFB MA 01731-3010. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document require that it be returned.

**DISCLAIMER**

The reproduction quality of the section "Catalog of Reports From Standard Stations of USSR for Ten Tamped Nuclear Explosions at Azgir" is the best available at the time of publication, and is beyond the control of the publisher. This section is provided as a service to the reader.

## NOTICE


We are furnishing you this document even though it does not meet our quality standards as indicated below:

- a.  Missing pages
- b.  Illegible printing
- c.  Illegible graphics and math equations
- d.  Blurred printing
- e.  Small size type
- f.  Broken letters
- g.  Positive image
- h.  Variable contrast
- i.  Half-tones have not reproduced well
- j.  Almost wholly illegible
- k.  Others:

If we can obtain a better document from the source, we will send it to you automatically without charge to replace this one.

If, however, you cannot use this document, please call Autovon 284-7633 or 202-274-7633 for a credit to your account.

# DISCLAIMER NOTICE



THIS DOCUMENT IS BEST QUALITY AVAILABLE. THE COPY FURNISHED TO DTIC CONTAINED A SIGNIFICANT NUMBER OF PAGES WHICH DO NOT REPRODUCE LEGIBLY.

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> 28 June 1993	<b>3. REPORT TYPE AND DATES COVERED</b> Scientific No. 2	
<b>4. TITLE AND SUBTITLE</b> Underground Nuclear Explosions at Azgir, Kazakhstan, and Implications for Identifying Decoupled Nuclear Testing in Salt			<b>5. FUNDING NUMBERS</b> PE 62101F PR 7600 TA 09 WU BK	
<b>6. AUTHOR(S)</b> Lynn R. Sykes			Contract F19628-90-K-0059	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Columbia University in the City of New York Box 20, Low Memorial Library New York, N.Y. 10027			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> Phillips Laboratory 29 Randolph Road Hanscom AFB, Ma. 01731-3010  Contract Manager: James Lewkowicz/GPEH			<b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>  PL-TR-93-2155	
<b>11. SUPPLEMENTARY NOTES</b>				
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b>  APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED			<b>12b. DISTRIBUTION CODE</b>	
<b>13. ABSTRACT (Maximum 200 words)</b>  <p>Bodywave magnitudes, <math>m_b</math> are recomputed for 17 nuclear explosions with yields of about 0.01 to 100 kilotons (kt) at Azgir in western Kazakhstan. Station corrections were developed for Azgir using larger events and then applied in recomputing magnitude of other explosions. Revised values of <math>m_b</math> for three tamped (fully coupled) explosions in salt at Azgir and one at Orenburg of announced yield, <math>Y</math>, were used to obtain the relationship, <math>m_b = 4.425 + 0.832 \log Y</math>. Salt is one of the best coupling geological media for generating seismic waves from underground nuclear explosions. In a special study made of the Azgir explosion of 1.1 kt of 1966 <math>m_b</math> was determined for 16 stations as <math>4.52 \pm .06</math>. For purposes of appreciating the detection capability of a given seismic network, it is important to recognize that a fully-coupled explosion of 1 kt in salt in high-Q (low attenuation) areas of the Former Soviet Union (FSU), like Azgir, has an <math>m_b</math> of 4.4; fully decoupled events of 1 and 10 kt have <math>m_b</math>'s of about 2.6 and 3.4. Most areas of thick salt deposits in the C.I.S. are typified by high Q for P waves and low natural seismic activity.</p> <p>Yields of all known nuclear explosions at Azgir and in other areas of thick salt deposits in the C.I.S. through May 1993 are recalculated. The yields of fully decoupled nuclear explosions of <math>Y \geq 0.5</math> kt that possibly could be detonated in the cavities produced by those events are calculated.</p>				
<b>14. SUBJECT TERMS</b>  yields of Soviet nuclear explosions, nuclear tests in salt, decoupled, evasion			<b>15. NUMBER OF PAGES</b> 130	
			<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified	<b>20. LIMITATION OF ABSTRACT</b> SAR	

Nearly all of those cavities are situated in the area of extensive salt domes to the north of the Caspian Sea in the Pre-Caspian depression. The largest fully decoupled tests, up to a maximum of 4 kt, that could be conducted in cavities crated by past nuclear explosions are largely confined to Azgir itself. An important consideration in identifying cavities in salt created by past nuclear explosions is that the yield of a fully decoupled explosion in them cannot be larger than 5% of that of the explosion that created the cavity. The monitoring of fully decoupled explosions of  $Y \geq 0.5$  kt set off in cavities created by past nuclear explosions would be a relatively easy task since the number of sites at which suitable cavities were created is quite limited. The FSU also conducted six very small nuclear explosions in a water-filled cavity at Azgir created by the 1968 explosion of 25 kt. Those six events, however, are all examples of enhanced seismic coupling at certain frequencies rather than of decoupled testing.

Ten tamped nuclear explosions at Azgir were relocated using seismic data and the locations of shot points on a SPOT satellite image taken in 1988. Most of the shot points are easily recognized on the image even though testing occurred there 9 to 22 years earlier. Since the Azgir area, like much of the Pre-Caspian depression, is arid, it would not be a suitable place for constructing large cavities in salt by solution mining and then using them for clandestine decoupled nuclear testing. A catalog (in Russian) of seismic readings from standard stations of the FSU for the 10 tamped explosions at Azgir was provided to us by Russian scientists and is reproduced in this report.

## CONTENTS

Introduction.....	1
Bodywave Magnitudes.....	4
Magnitude-Yield Relationship for Tamped Nuclear Explosions at Azgir.....	5
Magnitudes of Tamped Explosions in Salt in High Q Areas.....	6
Magnitude-Yield Relationships for Coupled and Decoupled Explosions in Salt in High Q Areas.....	9
Capabilities of Seismic Networks for Identifying Decoupled Nuclear Explosions in C.I.S.....	11
Cavities Formed by Nuclear Explosions in Thick Salt Deposits of C.I.S. That Might Be Usable for Clandestine Testing.....	13
Nuclear Explosions Detonated in a Water-Filled Cavity at Azgir.....	21
Conclusions.....	23
References.....	27
Tables.....	30
Figures.....	36
Appendix A - Accurate Relocation of Nuclear Explosions at Azgir, Kazakhstan, from Satellite Images and Seismic Data: Implications for Monitoring Decoupled Explosions.....	39
Appendix B - Catalog Reports From Standard Stations of USSR for Ten Tamped Nuclear Explosions at Azgir.....	53

<b>Accession For</b>	
NTIS SNA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/ _____	
<b>Availability Codes</b>	
Dist	Avail and/or Special
A-1	



## INTRODUCTION

This paper examines the yields of past nuclear explosions at the Azgir testing area in western Kazakhstan and the yields of other nuclear explosions detonated by the Former Soviet Union (FSU) in or near other thick deposits of salt. Azgir, an area of low population density, which is underlain by salt domes of the Pre-Caspian depression, appears to have been a major testing ground for the technology and effects of peaceful nuclear explosions (PNEs) from 1966 through 1979. Nuclear explosions were also conducted at that site in the cavities created by larger nuclear explosions (Adushkin et al., 1993).

For the Azgir site where nuclear explosions of a variety of sizes from about 1 to 100 kt were conducted in salt (Adushkin et al., 1993), major emphasis is given to developing an  $m_b$ -yield relationship for salt for yields of 1 to 100 kt, calculating yields of past nuclear explosions in salt both at that site and elsewhere in the FSU, and assessing the potential for using large cavities that may remain from those explosions for future clandestine nuclear testing. A future paper will examine past nuclear explosions detonated in large cavities at Azgir in more detail than is done here, the numbers of chemical explosions in that area of comparable size to that of decoupled nuclear explosions, and the potential for conducting and identifying decoupled nuclear explosions of various sizes in other geological media.

In previous work my colleagues and I determined yields for Soviet explosions by using calibration curves for P waves that were based on underground explosions of announced yield for which their P-wave magnitudes were corrected for  $m_b$  bias between the actual test site of the explosion and the Soviet site in question. Sykes (1992) estimated the yields of underground explosions at the Shagan River (Balapan) testing area in eastern Kazakhstan using refined magnitude-yield relationships for both body and Lg waves. This paper seeks to further improve magnitude-yield relationships for nuclear explosions detonated in salt.

The detonation of nuclear explosions in large underground cavities under either a Low-Yield Threshold Test Ban Treaty (LYTTBT) or a Comprehensive Test Ban Treaty (CTBT) would constitute the greatest challenge to verification efforts. That evasion scenario sets the limit on how low a yield can be verified effectively. The OTA Report, *Seismic Verification of Nuclear Testing Treaties* (Office of Technology Assessment, 1988), focused upon the Soviet Union and concluded that between

1 to 2 and 10 kt the only plausible method of evasion is that of nuclear testing in large cavities in salt domes. It also concluded that no method of evading a good monitoring network is credible above 10 kt but that several evasion scenarios, including testing in cavities in bedded salt and in hard rocks, are possible below 1 to 2 kt. Much of my present efforts in nuclear verification are focused upon what I believe is the critical yield regime from 1 to 10 kt, where scientific research over the near term (a few years) seems most likely to have a major impact on the verifiability of either a LYTTBT or a CTBT and on what threshold can be verified effectively, e.g. 1, 3, 10 kt.

The idea that the seismic waves from underground nuclear explosions detonated in large underground cavities could be considerably reduced in size, i. e. decoupled, compared to the those from a tamped explosion was first proposed in 1959 (Latter et al., 1961). A tamped explosion is one where there are insignificant void spaces between the nuclear device and the surrounding rock medium, and a fully decoupled event is one where there is a large enough void space between the device and the surrounding material that no damage is done to the surrounding material (Denny and Goodman, 1990). The general concept of decoupling was confirmed in the Cowboy experiments where small (less than one metric ton) chemical explosions were set off in cavities excavated in a salt dome in Louisiana (Herbst et al., 1961). The United States has detonated only three nuclear explosions in salt. The first two, Gnome in New Mexico in 1962 with a yield of 3.4 kt and Salmon in Mississippi in 1964 of 5.3 kt (Rawson et al., 1966) were tamped events. Both of those explosions produced cavities that remained standing for many years, a situation that is almost unique to salt as a geologic testing medium. Cavities produced by tamped nuclear explosions in the common testing medium at the Nevada Test Site (NTS), tuff (Stevens et al., 1991), and in hard rock nearly always collapse in a relatively short time. Such cavities, of course, are not then available for use in conducting decoupled explosions.

The Sterling nuclear explosion, a decoupled event of 0.38 kt, was detonated in the Salmon cavity at a depth of 828 m in 1966 to test the decoupling concept (Springer et al., 1968; Healey et al., 1971; Denny and Goodman, 1990). It is the only decoupled U.S. nuclear explosion detonated in domed or bedded salt. Its small size, however, has resulted in a long controversy about the feasibility of conducting larger-yield, clandestine nuclear explosions in much larger cavities in salt and

whether such events would be identified or not. Since Gnome and Salmon were both detonated prior to the installation of many high-gain short-period seismic stations and seismic arrays at teleseismic distances, a long debate has ensued about their bodywave magnitudes,  $m_b$ , and about  $m_b$ -yield relationships for tamped and decoupled nuclear explosions in salt.

Unlike the United States, the Former Soviet Union detonated a large number of nuclear explosions in and near thick deposits of salt, including a series of large underground explosions near the town of Azgir in western Kazakhstan and a series of nuclear explosions with yields of about 3 to 15 kt near Astrakhan in the adjacent part of the Russian Republic (Fig. 1). Azgir is located to the north of the Caspian Sea within the Pre-Caspian depression, which contains the world's largest concentrations of salt domes. Several other nuclear explosions were detonated by the U.S.S.R. from 1965 to 1988 in or near salt deposits (Sykes and Ruggi, 1989). Recently released information by the Russians on their program of nuclear explosions conducted in cavities at Azgir (Adushkin et al., 1993) indicates that a decoupled test of 8 kt was detonated in March 1976 within the air-filled cavity created in a salt dome by a tamped nuclear explosion eight times larger. The 1976 event was 20 times larger than the Sterling event and was recorded not only at local distances, like Sterling, but also at regional and teleseismic distances.

Clearly, there is much to be gained in understanding magnitude-yield relationships for nuclear explosions in salt and the feasibility of conducting and identifying clandestine decoupled explosions with yields from 1 to 10 kt by examining data from the much greater number of past Soviet explosions in salt and in urging the release of additional data on events of that type by the Russian Republic. There is no indication, however, that either the United States, the Soviet Union or the C.I.S. has created and evacuated a large solution-mined cavity and used it to test decoupled nuclear explosions.

Several important aspects of decoupling have received little study for more than 20 to 25 years. Since then a great deal of experience has been obtained by the U.S. and several European countries on the rheological properties of salt in conjunction with research on radioactive waste disposal in salt deposits and by industry on the construction and stability of large cavities in salt for petroleum storage and waste disposal. Experience with very large cavities in salt domes is now available from the U.S. Strategic Petroleum Reserve and from Europe, the C.I.S. and other parts of

the U.S. on cavities created in salt for gas storage and on their stability (or lack thereof). In addition yields, depths and cavity dimensions of explosions in salt at Azgir in 1966, 1968 and 1971 and at Orenburg in 1971 have been published (Kedrovskiy, 1970; Izrael' and Grechushkina, 1978; Adushkin et al., 1993) as have the yields, depths and rock types for a number of underground nuclear explosions at the eastern Kazakhstan test site (Bocharov et al., 1989). Thus, the time appears to be ripe for a reassessment of the feasibility of conducting decoupled nuclear tests of various sizes in large cavities in salt domes and ascertaining whether or not such events would be identified by various monitoring techniques.

### BODYWAVE MAGNITUDES

Revised  $m_b$  values were recomputed for all known Soviet underground nuclear explosions in the vicinity of Azgir for which data were available from the International Seismological Center (ISC), the U. S. Geological Survey (USGS) and the Norsar and Hagfors seismic arrays in Scandinavia for the period January 1961 through May 1993. Station corrections for Azgir events were derived for seven large explosions at that site and then applied to all known underground events for that testing area. Similar previous work on magnitude and yield determination are described in Sykes and Ruggi (1986, 1989), Sykes and Ekstrom (1989) and Sykes (1992). Stations used in the recalculations were confined to the distance range  $25^\circ$  to  $95^\circ$ . A major object in this study was to reduce the standard error of the mean (SEM) for  $m_b$  to values as small as 0.015 to 0.03 by both using large numbers of stations (40 to 70 for larger events) and applying station corrections, i.e. making a correction for systematic differences in magnitude at individual stations for a given testing area. Since individual stations typically record explosions from a given test site with amplitudes that are consistently either higher or lower than the mean for each explosion, the application of station corrections considerably reduces the standard deviation of individual readings and avoids biases related to the inclusion or exclusion of individual readings from one event to another. It is particularly important to use station corrections since measurements of  $m_b$  from the stations of countries like Canada or France that operate large networks were not available for some explosions but were available for others.

Recomputed magnitudes for Azgir explosions, their SEM, and other pertinent data are listed in Table 1. As described later, the recalculations of  $m_b$  for the Azgir

site do not represent maximum-likelihood determinations in which allowance is made for non-detection at additional stations (Ringdal, 1976). Instead, I emphasize the application of station corrections obtained from relatively large events to the determination of magnitudes of smaller events within the same testing area.

#### *Small Azgir Explosion of 1966 and 25 kt Event of 1968*

A special study was made of the Azgir explosion of 1.1 kt of 1966 wherein all available WWSSN and Canadian records were searched for the P wave from the event.  $m_b$  was measured at 16 stations, giving an average  $m_b = 4.524 \pm .056$ . (The uncertainty in magnitude as used here and throughout this paper is plus or minus one SEM.) Although small in amplitude, P waves could be readily identified at many of the better WWSSN stations of higher gain and good signal-to-noise ratio. The ability to detect such a small event 27 years ago using analog records from mainly simple (non-array) stations reflects the high coupling of a tamped underground explosion in salt and the efficient propagation (high Q) for P waves from the Azgir area to stations worldwide. A similar study for the 1968 Azgir PNE event of 25 kt gave  $m_b = 5.529 \pm .027$ . For both of the two Azgir events, readings were used from only from those stations for which a station correction was available based on the seven large events at Azgir. This use of station corrections avoids the largest contributor to biased determinations of  $m_b$  for small events, i.e. the inclusion of raw  $m_b$  values that are mostly from stations that systematically report larger than average magnitudes, such as those in Scandinavia. For example, for the seven large Azgir events used in calculating station corrections,  $m_b$  values for Norsar are on the average 0.48 units larger than the event means. Correction for that systematic effect results in subtracting a station correction of  $0.48 \pm 0.08$  from Norsar's raw  $m_b$ 's.

### MAGNITUDE-YIELD RELATIONSHIP FOR TAMPED NUCLEAR

#### EXPLOSIONS AT AZGIR

The revised values of  $m_b$  and the published yields for the Azgir explosions of 1966, 1968 and 1971 and the Orenburg event of 1971, all of which are reported as having been detonated in salt, were used to obtain the  $m_b$ -yield relationship

$$m_b = 4.4250 + 0.832 \log Y. \quad (1)$$

The yields of the explosions at Azgir in 1966 and 1968 were made public more than 23 years ago (Kedrovskiy, 1970) as part of an exchange of data on peaceful uses of

nuclear explosions. The 64 kt yield of the Azgir explosion of 1971 is from Adushkin et al. (1993); the 15 kt value for the 1971 Orenburg event is from Nordyke (1975), Izrael' and Grechushkina (1978) and Borg (1984). The three tamped explosions in salt at Azgir have magnitudes similar to those of explosions of similar yield in hard rock in the southwestern portion of the Shagan River testing area in Central Asia (Fig. 2) and, once corrected for test site bias, to those of the three U.S. explosions of announced yield in granite in Nevada (Sykes, 1992). Thus, the Azgir events in salt exhibit a coupling of explosion energy into seismic wave energy that is about as efficient as that of explosions in granite and other hard rock.

## MAGNITUDES OF TAMPED EXPLOSIONS IN SALT IN HIGH Q AREAS

### *Magnitudes of Salmon, Gnome and Small Events at Azgir*

Even though the United States carried out its few (two tamped and one decoupled) nuclear explosions in salt more than 27 years ago, there has still not developed a consensus (at least in the U.S.) about the magnitudes of those explosions and about  $m_b$ -yield relationships for coupled and decoupled nuclear explosions for areas of thick salt deposits of the C.I.S. Those relationships are, of course, very important to considerations of the yields of decoupled nuclear explosions that might be tested clandestinely under either a LYTTBT or CTBT. Most of the early magnitudes published for those U.S. explosions rely mainly on measurements at distances less than  $20^\circ$  to  $25^\circ$  where amplitudes of P waves vary considerably from one region to another. For that reason most work on magnitude-yield relations since about 1968 has used data only from teleseismic distances, i.e.  $\Delta > 20^\circ$  to  $25^\circ$ . The SIPRI Seismic Study Group (1968, p. 101) pointed out that truly teleseismic data published for the Salmon event give a magnitude of 4.8 to 4.9 whereas the inclusion of data from closer distances leads to lower estimates.

Some workers have argued that the use of teleseismic arrivals from Salmon leads to overestimates of its  $m_b$  since those data come only from stations with relatively high signal levels. Since larger explosions were not conducted at the Gnome and Salmon sites, they cannot be used to develop station corrections that could be applied to those relatively small explosions as I have done for the much greater number and size range of nuclear explosions in salt at Azgir or has been done for major test sites like NTS and Shagan River. The SIPRI Seismic Study Group (1968, p. 102), referring to a presentation by Peter Marshall, points out that stations with a gain of at least

100 K would be needed to detect an event the size of Salmon at teleseismic distances. That study states: "When Jordan et al.'s [1966] stations are examined, it is found that of 10 stations with gains greater than or equal to 100 K, 9 reported an observation." "Values from these averaged to 4.9." The SIPRI study goes on to conclude (p. 102) "There seems therefore a good case for SALMON having a magnitude of between 4.75 and 5.0." Marshall et al. (1979) calculated a magnitude and SEM for Salmon of  $4.87 \pm 0.08$  using data from 11 stations at  $\Delta > 20^\circ$ . When that  $m_b$  value is corrected for the relative attenuation between the Salmon and Azgir testing areas using the data and formulas of Der et al. (1985),  $m_b = 4.90$  is obtained. Using that  $m_b$ , a yield of 3.7 kt is derived from equation (1), which is only somewhat smaller than the announced yield of 5.3 kt. Thus, the calibration data from Azgir explosions are consistent with an  $m_b$  for Salmon near 4.9, i.e. in the range quoted by the SIPRI Seismic Study Group (1968).

Using a maximum likelihood program developed at Teledyne-Geotech (Blandford et al., 1984; R. R. Blandford, written communication to P. G. Richards, 1989) and Jih and Wagner (1991) obtained the following much smaller  $m_b$  values for Salmon: 4.31, 4.45 and 4.20 respectively. In his written communication Blandford states that he used 11 observations of  $m_b$  of Jordan et al. (1966) for  $30^\circ < \Delta < 90^\circ$  as well as assumptions about the noise levels at many other stations in that distance range for which P waves were not observed by Jordan et al. It seems clear that Blandford's assumptions about noise levels in data read by others are crucial in his determination of  $m_b$  using the maximum likelihood procedure that seeks to deal with stations that do not observe an event as well as with those that do. Jih and Wagner (1991) used  $m_b$  values from 6 stations and measurement of noise at 33 others in the range  $20^\circ < \Delta < 90^\circ$  in their determination.

The veracity of the Teledyne-Geotech procedures and estimates of  $m_b$  for Salmon now can be checked with information from Azgir. Jih and Wagner (1991) report  $m_b$ 's (which they call Geotech's maximum-likelihood network  $m_b$ ) of  $4.225 \pm 0.083$ ,  $5.542 \pm 0.040$  and  $3.986 \pm 0.073$  for the Azgir events of 1966, 1968 and April 25, 1975 for which I estimated their  $m_b$ 's to be 4.524, 5.529 (Table 1) and  $4.45 \pm 0.13$  (Table 2). For the 1968 event of 25 kt they used 44  $m_b$  values and 10 noise estimates whereas I used 20 values for which station corrections to  $m_b$  were available from seven large events at Azgir. My estimate, theirs, and that of Marshall et al. (1991) of  $m_b$  5.57, which is also a maximum-likelihood determination, are statistically

indistinguishable for the 1968 explosion. This is not surprising since each used relatively large numbers of  $m_b$  values in their determinations. Also, Ringdal found little or no difference between magnitude estimation techniques for  $m_b \geq 5.0$ .

For the 1966 event of 1.1 kt, however, Jih and Wagner used only three  $m_b$  values but included 10 noise observations whereas I used 16 readings, each of which incorporated a station correction. The situation is even more extreme for the 1975 explosion where Jih and Wagner used merely a *single*  $m_b$  value and 16 noise estimates whereas I used seven values of  $m_b$ , each of which incorporated a station correction. For the 1975 event their  $m_b$  estimate is 0.46 units smaller than mine. Marshall et al. (1991) obtained a maximum-likelihood  $m_b$  of 4.59 for the 1975 event, which is statistically indistinguishable from my value but is 0.60  $m_b$  units larger than that of Jih and Wagner (1991).

Thus, for two of small Azgir events Jih and Wagner (1991) report considerably smaller magnitudes than those obtained by either Marshall et al. (1991) or me. Procedurally it seems that the Teledyne-Geotech  $m_b$  values for small events suffer from a reliance on a very few actual readings, lacking many data from standard sources like ISC, USGS, Norsar, Lasa, the Hagfors array in Sweden and WWSSN stations, as I used in obtaining my estimates. When the number of  $m_b$  measurements in the Geotech database is very small (3 and 1 for the 1966 and 1975 Azgir events) and the number of noise estimates is much larger, it seems clear that their calculated magnitudes and SEM's are systematically too small. Note that this should not be taken as a criticism of the maximum-likelihood technique itself since Marshall et al. (1991) obtained a similar result to mine for the 1975 explosion.

Returning to the Salmon event, it is clear that using just six  $m_b$  values and 33 estimates of noise as Jih and Wagner (1991) did, when at least 11  $m_b$  values were available at distances greater than  $20^\circ$ , probably also led to their very small estimates of  $m_b$  and its SEM for Salmon. Likewise, similar reasoning indicates that the  $m_b$  values of Blandford et al. (1984) and Blandford (written communication, 1989) are also too small.

Not much can be done in utilizing the data from the Gnome explosion since Marshall et al. (1979) report only two  $m_b$  readings for  $\Delta > 20^\circ$ , whose average is 4.13. If the poorly known site bias for the Gnome area with respect to Azgir is like that of NTS with respect to Azgir, 0.323  $m_b$  units (Der et al., 1985),  $m_b = 4.45$  is obtained for Gnome as normalized to Azgir. In view of the large uncertainty in the  $m_b$  deter-



mination for Gnome, however, the  $m_b$  values for explosions at Azgir appear to be more precise since they are not biased by either a lack of station corrections, an uncertain estimate of attenuation, or a very small number of P waves observed at teleseismic distances.

## MAGNITUDE-YIELD RELATIONSHIPS FOR COUPLED AND DECOUPLED EXPLOSIONS IN SALT IN HIGH Q AREAS

### *Is Salt a High or Low Coupling Testing Medium?*

Most workers who have studied the seismic waves from underground nuclear explosions in thick deposits of salt (e. g. SIPRI Seismic Study Group, 1968; Marshall et al., 1979; Rodean, 1981) have concluded that salt is one of the best-coupling, common geologic media, i. e. that tamped explosions of a given yield in salt have among the largest bodywave magnitudes compared to those for other testing media. The large values of  $m_b$  in Fig. 2 for the three Azgir explosions of published yield supports that contention. This is reasonable since salt has a low porosity, and relatively little of the energy of the explosion must be expended in closing pore space, especially air-filled pore space, as is the case with explosions conducted above the water table in less competent sedimentary rocks and alluvium.

Based on their determination of the  $m_b$  of Salmon, Blandford et al. (1984), however, conclude that its magnitude, when corrected for bias between testing areas, was 0.4  $m_b$  units below the magnitude-yield curve for shots in volcanic and granitic rocks in Nevada, Amchitka and Algeria. They conclude from that single data point for Salmon that explosions in salt couple less well than those in the above hard rocks. In the light of my finding that their  $m_b$  is likely to be seriously underestimated for Salmon, however, there is no remaining rationale for concluding that tamped explosions in salt couple less well than those in granite and other hard rocks.

### *Magnitudes of 1 and 10 kt Nuclear Explosions in Salt in High-Q Areas*

Eqn. (1), the magnitude-yield relationship derived earlier for tamped nuclear explosions in salt at Azgir and Orenburg, implies that a fully-coupled (tamped) nuclear explosions of 1 and 10 kt in salt in those and similar high-Q areas of the FSU have  $m_b$ 's of 4.42 and 5.26 respectively. The OTA Report (Office of Technology Assessment, 1988) and a number of other investigators have concluded that the aver-

age decoupling factor (DF) at low frequencies for the Sterling nuclear explosion was about 70 times, i. e. the amount by which its seismic amplitude at low frequencies was reduced compared to that of a tamped explosion of the same yield at that location. Since the yield of Sterling, 0.38 kt, was somewhat larger than that calculated theoretically for full decoupling, 0.21 kt, some workers have maintained that somewhat higher decoupling factors would be expected for a mined cavity in salt of a size such that the salt surrounding the cavity remains wholly in the elastic regime. The OTA Report concluded, however, that the experience from Sterling and the re-analysis of the Cowboy data from chemical explosions in salt, which also gives a maximum DF of about 70 (Murphy, 1980), indicates that larger values of DF are unlikely to be achieved in practice. For purposes of calculating magnitudes of fully decoupled explosions I assume a DF of 70. Subtracting  $\log 70$  from (1), the following expression is obtained for fully decoupled explosions in salt at Azgir and other high-Q areas of the FSU:

$$m_b = 2.58 + 0.832 \log Y. \quad (2)$$

Assuming that attenuation of P waves leaving the high Q (efficient transmission) Azgir site is the same as that for other high-Q areas of the C.I.S., fully decoupled events of 1 and 10 kt would have  $m_b$ 's of 2.58 and 3.41 respectively. Most areas of thick salt deposits in the former U.S.S.R. are typified by high Q for P waves and low natural seismic activity. These magnitudes are higher than has generally been thought previously for fully decoupled nuclear explosions. For example, Murphy et al. (1988) state "Cavity decoupled underground nuclear explosions in the yield range from 1 to 10 kt can be expected to generate seismic signals corresponding to  $m_b$  values in the 2.0 to 3.0 range . . ." One of the smallest magnitudes estimated is that of Werth and Randolph (1966) who concluded that a 5-kt, fully decoupled explosion would be down in amplitude from that of Salmon by a DF of 170 or a magnitude of 2.1. Eqn. (2) gives  $m_b = 3.16$  for 5 kt fully decoupled, a full magnitude larger than their estimate. Their low estimate arises at least in part from using an  $m_b$  (4.35) for Salmon determined from stations as close as  $16^\circ$ , data from a single pair of tamped and decoupled explosions, and too large a decoupling factor in the light of the re-evaluation of the data from the Cowboy set of chemical explosions.

## CAPABILITIES OF SEISMIC NETWORKS FOR IDENTIFYING DECOUPLED NUCLEAR EXPLOSIONS IN C.I.S.

Figure 2 indicates the expected magnitudes of contained underground nuclear explosions of various yields in the territory of the former Soviet Union. The range of  $m_b$ 's for tamped (fully coupled) events in hard rocks and water-saturated rocks is meant to cover the range of experience reported in the literature for differences in coupling near the shot point for those media as well as differences in the attenuation of P waves transmitted to teleseismic distances. Note that the line labelled SW Shagan River on Figure 2, which pertains to a hard rock site with low attenuation for P waves, falls near the top of that regime as do the data points for the three Azgir explosions in salt. Dry porous media (Fig. 2) do not appear to be present in sufficient thicknesses anywhere in the C.I.S. such that explosions could be detonated in them clandestinely with yields larger than 1 to 2 kt (SIPRI Seismic Study Group, 1968; Office of Technology Assessment, 1988). The upper limits of possible underground testing in Figure 2 are taken to be the present 150 kt limit of the TTBT for hard rocks and water saturated rocks and the upper limit of 10 kt for conducting clandestine testing in large cavities in salt domes as described in Office of Technology Assessment (1988).

The OTA Report described the *identification* threshold as of 1988 for the C.I.S. as  $m_b$  4.0 (which is, of course higher than the *detection* threshold for seismic events). That assessment did not include data from stations within the FSU, such as that from the newer IRIS stations. At that capability, tamped explosions at Shagan River would be identified down to several tenths of a kiloton; some events in hard rock and water saturated rock in areas of low P-wave attenuation of the C.I.S. might go unidentified for yields as large as nearly 3 kt. The OTA Report concluded that at an identification threshold of 4.0 that fully decoupled explosions in salt domes might go unidentified for yields as large as 10 kt, a level largely set by the size of cavities that could be constructed and used clandestinely and the low levels of natural seismic activity for most salt domes areas of the Soviet Union.

Members of the panel that advised OTA on their 1988 study, of which I was a part, did not reach a consensus on what the identification threshold for the C.I.S. would be if data from a good internal seismic network were available in addition to data from external stations. There was agreement (p. 92) that "identification can be

accomplished in the U.S.S.R. down to *at least* as low as  $m_b$  3.5." The Report goes on to state "Many experts claim that this identification threshold is too cautious and that with an internal network, identification could be done with high confidence down to  $m_b$  3.0." Figure 2 indicates that an identification capability of  $m_b$  3.5 would include many events in the FSU down to about 0.07 kt, and events in all hard rocks and those below the water table down to 0.75 kt but might miss fully decoupled events as large as 10 kt. A capability to discriminate at  $m_b$  3.0 would lead to the identification of fully decoupled events in salt domes down to yields of 2 to 4 kt.

The regime labelled "salt, fully decoupled" in Figure 2 pertains to high Q areas of the C.I.S, which includes most but not all salt domes. Salt domes in more seismically attenuating areas are concentrated in the Republic of Tadjakistan, which is now a separate country from the Russian Republic. A better identification capability for possible decoupled explosions in that area, if deemed necessary, could be furnished by a local seismic network. The major civil war that has been in progress in Tadjikistan for the last few years, however, would undoubtedly make the conduct of nuclear tests of any kind in that area very difficult, if not impossible so long as the war continues. Likewise, it would likely limit or prohibit the operation of seismic stations in Tadjikistan.

Stevens et al. (1991a) and others imply that the C.I.S. could conduct clandestine nuclear explosions in other dry materials such as dry tuff, which is present in thick quantities at NTS. Dry tuff of significant thicknesses for even small nuclear tests appears to be present in the C.I.S. only in the Caucasus region. Again, a monitoring network in the vicinity of those deposits could provide better identification for that region if it were deemed necessary under a LYTTBT or CTBT. Conducting clandestine nuclear explosions in dry porous alluvium would be very risky for a potential evader since that material is very weak and undergoes significant compaction when nuclear explosions are fired in it. The collapse of the cavity formed by such an explosion, which is almost certain to occur, may well produce significant disturbance at the surface, even if it is overburied. This could readily be detected by satellite photography or air surveillance.

## CAVITIES FORMED BY NUCLEAR EXPLOSIONS IN THICK SALT DEPOSITS OF C.I.S. THAT MIGHT BE USABLE FOR CLANDESTINE TESTING

### *Inventory of Past Nuclear Explosions of FSU in and near Thick Salt Deposits*

The U.S.S.R. carried out a number of nuclear explosions in salt near Azgir and conducted several other PNEs in regions known to contain salt. Cavities produced by nuclear explosions in salt, such as Salmon and Gnome in the United States, have remained standing for many years whereas cavities produced in other rock types usually collapse within short periods of time. Many investigators evaluating the possibility of decoupled nuclear testing that might be conducted under either a CTBT or a LYTTBT have paid considerable attention to the potential use of cavities produced by nuclear explosions in salt. I will show, however, that monitoring of the relatively few areas of the C.I.S. in which cavities of that type could exist and could be used in the future for the full decoupling of explosions with yields larger than 0.5 kt is tractable, given both a problem-solving approach and the inclusion of reasonable verification measures in a treaty to further limit nuclear testing. Instead, more attention needs to be devoted to the feasibility of decoupled testing in the yield range from 1 to 10 kt in large cavities produced by solution mining. More potential sites are available in the C.I.S. for creating large cavities by solution mining than are available from past nuclear explosions in salt.

Table 1 lists underground nuclear explosions that were detonated by the FSU through June 1993 either in or near thick salt deposits. The latest events on that list were conducted in 1988. The judgment as to whether or not an explosion occurred in or near a thick salt deposits was largely based on its location with respect to the maps of salt deposits of Elias et al. (1966) and Rachlin (1985). Subsequent to writing the first draft of this paper Sultanov et al. (1993) published information on Soviet PNEs that includes the rock type at the hypocenter of the explosion. All of the events that they list as occurring in salt are shown in Table 1 without a designation as to rock type; all others that they report in other rock types are so designated in Table 1.

Nuclear explosions detonated in cavities at Azgir are not included in Table 1 but are listed separately in Table 2. As mentioned earlier, yields of tamped explosions at Azgir were determined using recalculated  $m_b$  values that included station corrections derived for that site. The explosions at Azgir listed in Table 1 have calculated yields between 1.3 and 93 kt. That range is in good agreement with the statement by

Adushkin et al. (1993) that tamped nuclear explosions at Azgir were in the yield range from 1 to 100 kt and the reported yield of 1.1 kt for the 1966 explosion. Those events occurred between April 1966 and October 1979 in the area of numerous salt domes to the north of the Caspian Sea. Applying station corrections determined from explosions at Azgir to events more than 100 km away, however, did not reduce the SEM of those  $m_b$  values. Hence, station corrections were not used in recomputing  $m_b$  for the other events in Table 1.

The yields,  $Y(mb)$ , of all of the events in Table 1 were calculated from equation (1). It is assumed in those calculations that all of the events in Table 1 were tamped explosions and that all occurred in salt in areas of low attenuation for P waves transmitted to teleseismic distances. Announced yields are listed in a separate column. In Table 1 origin times and locations for events at Azgir are from Sykes et al. (1993); those of explosions at Astrakhan, Lake Aralsor, Orenburg and Karachaganak are from Marshall et al. (1991); the others are from the ISC bulletins.

#### *Conservative Assumptions about Use of Cavities for Possible Clandestine Testing*

I now estimate the maximum yields of fully decoupled nuclear explosions that could be detonated in cavities created by past Soviet nuclear explosions detonated either in or near thick salt deposits of the C.I.S. Two conservative assumptions are made in ascertaining potential sites where decoupled tests might be performed in the future. One is that all of the events in Table 1 occurred in thick deposits of salt. Sultanov et. al (1993) state that several of the events in Table 1, however, were detonated in either anhydrite, clay, or dolomite and not in salt. Those events are so annotated in Table 1. Without precise knowledge of the depths of explosions and details of the stratigraphy for regions of bedded salt, it is not possible to ascertain whether explosions, in fact, occurred in salt or in some other rock type. An example of this is the series of nuclear explosions (Table 1) that have been detonated south of the town of Mirnyy near  $61.5^{\circ}N$ ,  $112.8^{\circ}E$  within the large region of bedded salt to the northwest of Lake Baikal. Sultanov et. al (1993) indicated that only one of those explosions occurred in salt; the others were conducted in dolomite in conjunction with oil recovery.

Other conservative assumptions are that cavities have remained standing for all of the events in Table 1 and that water present in any of them can be removed so that decoupled testing would be possible. Kedrovskiy (1970) mentions that the small

cavity created by the 1966 explosion at Azgir filled with water. Russian scientists have stated that the larger cavity created by the nearby 1968 explosion filled with water and that six of the very small explosions in Table 2 were detonated in it. As discussed later, they are examples of enhanced coupling at frequencies of about 7 to 9 Hz. rather than decoupled tests.

### *Cavity Volume as a Function of Yield and Depth of Tamped Nuclear Explosions in Salt*

Information about the depths and dimensions of the cavities created in salt by the U.S. explosions Salmon and Gnome, three explosions at Azgir and one near Orenburg are used to calculate yields of fully decoupled nuclear explosions that could be conducted in the cavities assumed to remain standing from the events in Table 1.

Salmon, a fully-tamped explosion of 5.3 kt, was detonated in a salt dome in the state of Mississippi at a depth of 828 m in 1964. The Sterling nuclear explosion, a decoupled event of 0.38 kt, was detonated in the Salmon cavity at the same depth in 1966 (Denny and Goodman, 1990). The Salmon cavity, like that produced by the Gnome explosion in bedded salt in New Mexico and those produced by the tamped Azgir explosions of 1966, 1968 and 1971 in salt domes, was not perfectly spherical in shape. In each case a significant amount of rubble, radioactive products and re-solidified salt accumulated at the bottom of what was initially a more nearly spherically shaped cavity. In the following descriptions and calculations the cavity volume,  $V_C$ , referred to is the remaining volume; it does not include the rubble zone since it is the remaining volume that is pertinent to the conduct of possible decoupled nuclear tests in those cavities.

$V_C$  for the Sterling event was  $19,400 \text{ m}^3$ , giving a mean radius of 16.7 m (Denny and Goodman, 1990). The Gnome explosion of 1962 was conducted in bedded salt at a depth of 361 m and produced a  $V_C$   $27,400 \text{ m}^3$  (Rawson et al, 1966). The 1968 Azgir event of 25 kt was detonated at a depth of 590 m and produced a cavity with a volume of  $140,000 \text{ m}^3$ , giving a mean radius of 32.2 m (Kedrovskiy, 1970). The Soviet explosion at Orenburg of October 22, 1971 of 15 kt in bedded salt was used to produce a cavity for storing gas condensates at depth of 1140 m. Its volume was  $50,000 \text{ m}^3$ , giving a mean radius of 22.9 m (Nordyke, 1975; Izrael' and Grechushkina, 1978; Borg, 1984). The 1971 Azgir explosion of 64 kt was detonated at a depth of 987 m and produced an air-filled cavity with a volume of about  $214,000 \text{ m}^3$  (Adushkin et al, 1993) and a mean radius of 37 m. That cavity was used to conduct

a partially decoupled nuclear explosion of 8 kt in 1976 (Adushkin et al, 1993). The 1966 Azgir explosion of 1.1 kt was detonated at a depth of about 165 m and generated a cavity with a volume of 10,000 m<sup>3</sup> (mean radius of 13.4 m).

### *Containment Criteria for Fully Decoupled Nuclear Explosions*

The minimum depth for a clandestine test in an underground cavity in salt is determined by the requirement that the explosion not produce a crater or other disturbance at the surface and that it be fully contained so as not to leak radioactive products to the surface. For a very weak material like salt, Latter et al. (1961) conclude that the amplitude of the long-term step of pressure on the cavity wall for full decoupling must be less than or equal to one half of the overburden pressure (i.e. half of the vertical stress,  $\rho gh$ ) so as to prevent failure in tension of the surrounding salt material and, hence, to prevent leakage of radioactive gases from the cavity. Since salt, like other geological materials is very weak in tension, the presence of a compressive overburden stress is need to make sure that salt near the cavity wall is not subjected to tensional stress by the pressure step from the explosion. The relationship between a step in cavity pressure,  $P$ , produced by a decoupled explosion of yield,  $Y_D$ , in a cavity of volume,  $V_C$ , and the requirement for containment that  $P$  be less than some constant,  $k$ , times the vertical stress can be written

$$P = (\gamma-1)Y_D/V_C \leq k \rho gh \quad (3)$$

where  $\gamma$  is the ratio of heat capacity at constant pressure to that at constant volume, which is taken to be 1.2 for an air-filled cavity (Latter et al., 1961),  $\rho$  is the average density of the material from the surface to the depth,  $h$ , of the cavity and  $g$  is the gravitational acceleration at the earth's surface. The average density in the following applications is taken to be that of salt at the Salmon site, 2200 kg / m<sup>3</sup> (Stevens et al., 1991), similar to that reported for Azgir (Kedrovskiy, 1970; Adushkin et al., 1993). For the Latter criterion mentioned above,  $k = 0.5$ .

### *Maximum Fully Decoupled Yields Possible for Various Cavities*

Taking the relevant parameters for the cavities created by the Salmon, Orenburg and the 1966, 1968 and 1971 Azgir explosions, the maximum yields of fully decoupled explosions according to the Latter criterion that could be detonated in those cavities (after converting the energy in Joules,  $J$ , to kilotons, where 1 kt = 4.184 x 10<sup>12</sup> J) are 0.21, 0.75, 0.02, 1.1 and 2.7 kt respectively (Table 1). The ratio,  $Y_{FD}/Y$ , of the



yield of the largest fully decoupled event that could be detonated in each of those cavities to the yield of the tamped nuclear explosion that was used to create those cavities is  $1/26$ ,  $1/20.4$ ,  $1/53$ ,  $1/24$  and  $1/24$  for the above five events respectively (Figure 3). For the other events in Table 1 for which information on yield, cavity dimensions and depth have not been published, the maximum yield of a fully decoupled explosion,  $Y_{FD}$ , that could be detonated in the cavities created by those events was obtained by multiplying the announced or calculated yield,  $Y$ , in Table 1 by  $1/20$ , the largest ratio obtained for the above five events. Values of  $Y_{FD}$  are listed in Table 1 if they are greater than or equal to 0.5 kt.

It should be appreciated that fully decoupled and partially decoupled nuclear explosions larger than 1 kt would have to be conducted within a fairly narrow range of depths. An air-filled cavity in salt is likely to deform significantly at depths greater than about 1000 to 1200 m. In fact, it is questionable whether the cavity produced by the Orenburg explosion of 1971 at a depth of 1140 m could be safely evacuated for clandestine testing without it deforming significantly. Since it was intended for storage of gas condensates, the presence of that material in the cavity provides a significant amount of the support for the cavity, allowing it to remain relatively undeformed to a greater depth than would be the case of an air-filled cavity at atmospheric or lower pressures. Russian scientists have stated that several of the cavities created by nuclear explosions in salt at a depth of about 1100 m in the Pre-Caspian depression have either collapsed or suffered enough of a reduction in volume that they could not be used for their intended industrial use, storage of gas condensates. Filmmakers for a National Geographical Society film on the Volga River interviewed health professionals in Astrakhan. They were informed that 13 of the 15 cavities created by nuclear explosions (Table 1) near Astrakhan (Fig. 1) experienced those problems.

Not much purpose is served, however, by debating whether the maximum depth of a stable air-filled cavity in salt is say 1000 or 1200 m. That depth varies from place to place and is dependent upon the temperature gradient, amounts of other evaporite minerals present in addition to NaCl, grain size, and the presence or absence of inclusions of salt brine (Jenyon, 1986). The main point is that cavity stability cannot be insured except at quite shallow depths in the crust, depths shallower than those of many salt bodies and depths much shallower than those of many oil and gas wells. For example, a salt deposit at a depth of say 5 km cannot be used to

create a large cavity. Likewise, an evader determined not to be caught testing at such yields, would use only cavities that are at least several hundred meters deep to insure containment and to prevent bomb-produced products from escaping from the cavity.

Figure 3 shows the scaled cavity volume,  $V_C/Y$ , as a function of depth for tamped explosions in salt of published yield.  $V_C/Y$  clearly decreases with depth. There is a tradeoff, however, between the fact that a shallower tamped explosion produces a larger cavity than a deeper one in the same material and the fact that a larger cavity at a shallower depth is needed to satisfy inequality (3) for a decoupled event of yield,  $Y_D$ . As shown at the top of Figure 3, larger values of  $Y_{FD}/Y$  are obtained at deeper depths. That ratio, however, does not increase much between 360 and 1140 m.

The slow increase of  $Y_{FD}/Y$  with depth,  $h$ , at the top of Figure 3 can be understood as follows. The data in the lower half of Figure 3, not including that for the 1971 explosion, were best fit with regressions of form  $V_C/Y \sim h^{-n}$ . For the solid line, which includes the data point for the shallow small event of 1966,  $n = 0.57$ . When that data point is excluded (dashed line),  $n = 0.83$ . Given the yield of a tamped explosion,  $Y$ , and  $h$  eqn. (3) can be rewritten

$$Y_{FD} \leq k\rho ghV_c / (\gamma - 1) \sim Y h^{1-n} \quad (4)$$

where  $1-n$  is 0.43 with and 0.17 without using data from the small, shallow event of 1966. Thus, for a given yield,  $Y$ , of a tamped explosion in salt, the yield,  $Y_{FD}$  of a fully decoupled explosion that can be detonated in its cavity, increases slowly with the depth of the tamped explosion.

Since the depths of most of the explosions in Table 1 are not known, assuming that they occurred at the depth of the deepest known nuclear explosion in salt in Figure 3 and taking the ratio of  $Y_{FD}/Y = 1/20$  should lead to conservative estimates of  $Y_{FD}$ . Uncertainties in calculating the yields of the events in Table 1 can lead to uncertainties of a factor of 1.5 to 2.0 in  $Y_{FD}$ . My main concern here, however, is what are the possibilities that may be available to test weapons clandestinely of certain approximate sizes.

Sterling was almost, but not fully decoupled;  $P$  was a factor of  $1.8 = 0.38 / 0.21$  times larger than that calculated for full decoupling by the Latter criterion. If Sterling conditions, i. e.  $k = 0.90$ , apply rather than the Latter criterion, the decoupled yields in the last column of Table 1 should be multiplied by 1.8. The value  $k = 0.90$  calculated for that event using equation (3) indicates, however, that  $P$  still did not

exceed the vertical stress ( $k = 1$ ). Experience with cavities in salt used for high-pressure gas storage indicates that leakage may occur when the internal pressure in the cavity exceeds the vertical stress (Berest and Minh, 1981). In fact, Latter (1960) was well aware of this "general rule of thumb" during the initial work on decoupling 33 years ago.

Thus, much larger (20 to 55 times larger) tamped explosions are needed to create cavities than the maximum sizes of fully decoupled explosions that can be detonated in them according to the Latter criterion ( $k=0.5$ ). For Sterling conditions ( $k=0.9$ ) that ratio is 12 to 31. This ratio is important since unclassified data alone are sufficient to identify down to a small yield all past Soviet and U.S. underground nuclear explosions that were detonated either in or near areas of thick salt deposits or in other areas that conceivably could be the sites of such deposits. The yields of fully decoupled explosions that could be detonated in cavities that may remain standing from those events according to the Latter criterion is at least a factor of 20 smaller than the yields of the explosions that generated those cavities. For Sterling conditions, the yield must be at least 12 times smaller.

#### *Inventory of Cavities Suitable for Full Decoupling by Region and Yield*

Table 3 summarizes the yields of fully decoupled nuclear explosions,  $Y_{FD} \geq 0.5$  kt, that could be detonated in the cavities of events in Table 1 both by area and size. Most of the possibilities for such testing are concentrated in the area to the north of the Caspian Sea in the Pre-Caspian depression (Fig. 1). The possibilities for conducting larger tests, up to a maximum of 4.2 kt fully decoupled, are mostly confined to Azgir itself. Possibilities for decoupled testing in cavities created by past nuclear explosions within the area of bedded salt to the northwest of Lake Baikal are few and the yields very small. Sultanov et al. (1993) list only a single small nuclear explosion in salt in that area.

The Bukhara II event of May 1968 (Table 1) was used to put out a fire in a gas well, the drilling of which had encountered an unanticipated fault that had provided pathways for escaping petroleum (Kedrovskiy, 1970; Nordyke, 1975). It was detonated at a depth of 2440 m near the boundary between anhydrite and salt (Kedrovskiy, 1970; Nordyke, 1975). The great depth of the Bukhara II explosion insures that any cavity that may have been created undoubtedly closed soon after detonation, as, in fact, was the intention in putting out the gas fire. The presence of a

fault known to have leaked in the past would make the conduct of a decoupled nuclear test at that site a risky proposition even if a cavity did remain standing.

The two explosions of 1972 in Table 1 to the northeast of Elista and at Lake Aralsor were situated along an 800-km long deep seismic sounding profile for which a cross section is reproduced in Scheimer and Borg (1984). The cross section indicates that salt is only present at a depth exceeding several kilometers for the first explosion and is absent altogether near the second. This is in accord with the statement of Sultanov et al. (1993) that the Lake Aralsor and Elista explosions were conducted in clay and with that by Bogacheva et al. (1965) that the 6800 m deep Aralsor borehole (which was located at or near the explosion) penetrated a complete Triassic sequence of rocks that are undisturbed by salt tectonics, i.e. by the presence of salt diapirism.

If the statements by Russian scientists about the events in Table 1 that were detonated in clay are correct, all of the cavities that may remain standing from past nuclear explosions in salt that could be used for full decoupling of explosions of  $Y \geq 1$  kt are situated at Azgir. If that is the case, the monitoring of such cavities by a combination of a local seismic network, satellite and air photography and on site inspections would be very easy if provisions to that effect are included in any test ban treaty. In any case, it would not be much more difficult in a monitoring program agreed to by treaty to include the sites of the other past large nuclear explosions in Table 3 for which the events are reported to have been detonated in clay.

Thus, the number of cavities produced by past nuclear explosions in the C.I.S. that potentially could be used for clandestine testing of fully or nearly fully decoupled nuclear explosions under either a CTBT or a LYTTBT is very limited. Those sites are confined to a few areas of the former U.S.S.R. Most, and perhaps all, of the larger cavities produced by nuclear explosions that may remain standing are situated in the Republic of Kazakhstan and not in the Russian Republic. One small nuclear explosion in salt with a yield of about 3 kt (Table 1) was detonated in the Ukraine (Sultanov et al., 1993).

The question of using cavities created by past nuclear explosions for partially decoupled events as described by Stevens et al. (1991) will be dealt with in a separate paper along with the feasibility of constructing and using cavities in hard rock for either the full or the partial decoupling of explosions with yields from 1 to 10 kt.

## NUCLEAR EXPLOSIONS DETONATED IN A WATER-FILLED CAVITY AT AZGIR

The Norsar and Hagfors arrays and the ISC Bulletin have each located a number of small events in the general vicinity of Azgir, including the partially decoupled explosion of 1976. Sykes and Lyubomirskiy (1992) made a special study of small events reported by the ISC, Norsar and Lasa in the  $5^{\circ}$  by  $5^{\circ}$  box outlined in Figure 2 that includes Azgir and Astrakhan for the 23-year period 1969 through 1991. They compiled a catalog of chemical and nuclear explosions in that region that they show is complete down to  $m_b$  3.1 since 1969. That  $m_b$  corresponds to a tamped nuclear explosion of about 0.025 kt and to yields of about 1 and 4 kt for decoupling factors of 20 and 70. This capability was made possible by recognizing that the Norsar, Hagfors and Lasa arrays recorded seismic waves from large events at Azgir that have  $m_b$ 's about 0.5 units larger than the average. Thus, those arrays have (or for Lasa had) a detection capability for Azgir that extends down to a very small  $m_b$  and yield. The location capability of those arrays by themselves, however, is poor compared with that of either a local seismic network or data from several arrays well distributed in azimuth. Either of those can be obtained by the installation of appropriate seismic monitoring equipment.

Sykes and Lyubomirskiy (1992) reported that seven small events of  $m_b$  3.02 to 4.45 from their catalog fulfill an origin-time criterion (being detonated exactly on the hour within the uncertainty in estimating origin time) for being either very small tamped or small decoupled nuclear explosions, one of which is the 8-kt partially decoupled event of 1976. Of the 126 other small events in the area, most or all of which are taken to be chemical explosions from their concentration during the work day, the largest two in 23 years were of  $m_b$  4.0. Chemical explosions of  $m_b \geq 3.5$  and those of  $m_b \geq 3.0$  occurred about 1.3 and 4.5 times per year in the entire study area. Thus, the number of chemical explosions per year in that area that would have to be discriminated as such from small decoupled nuclear events under a future test ban is small even at the  $m_b$  3.0 level. A major uncertainty pointed out by the OTA Report is, in fact, how many chemical explosions must be contended with per year

equal in  $m_b$  to that of decoupled explosions of various yields?

Russian scientists stated to me that besides the partially decoupled event of 1976 that four of the other events on the list of Sykes and Lyubomirskiy (1992) were very small nuclear explosions detonated in a water-filled cavity of radius 32m that was created in salt by a previous nuclear explosion. The earliest date of those small explosions, 1975, and the description of a cavity of that radius created by a 25 kt explosion in salt by Kedrovskiy (1970) indicate that the small explosions must have been detonated in the cavity created by the explosion of 1968 at Azgir. Those scientists indicated that two of the events on the list were not nuclear explosions but stated that two yet smaller nuclear explosions that were not on the list had been detonated in the same water-filled cavity on October 30, 1977 and November 30, 1978.

Since nearly all known or inferred Soviet nuclear explosions at Azgir and in the rest of the Pre-Caspian depression were detonated on the hour in a narrow range of local times from 0600 to 1100, I asked Dr. Frode Ringdal to search for possible small signals on those dates in 1977 and 1978 on the Norsar recordings that would have been detonated exactly on the hour during that five-hour time window. He reported that the Hagfors array in Sweden did report P arrivals that were consistent to within a few seconds (Table 2) of events having occurred at Azgir on those two dates at 0700 and 0800 GMT respectively. Norsar, however, was not operational at either of the two expected arrival times. Hagfors recorded all seven of the events in Table 2 that occurred in cavities at Azgir; Norsar recorded five of the events and undoubtedly would have recorded and located the other two if that array had been in operation.

Station corrections for Norsar and Hagfors (and other stations that recorded the explosions in Table 2 of 1975, 1976 and 1979) were used in deriving magnitudes of the seven events. Equation (1) was used to derive approximate yields of the six events fired in the water-filled cavity. It can be seen that the calculated yields of the three smallest events in Table 2 are 0.01 to 0.02 kt. Apparently the small explosions were tested to see if the fundamental frequency of a water-filled cavity surrounded by salt could be excited so as to produce larger than normal seismic waves near that frequency for use in deep seismic sounding. A simple calculation gives a fundamental resonance of about 11.7 Hz; Russian scientists report amplitudes up to

five times those of tamped explosions in salt at frequencies of 7 to 9 Hz. Thus, the six events in the water-filled cavity were not tests of decoupling but of enhanced coupling at certain frequencies commonly recorded in deep seismic sounding.

## CONCLUSIONS

For purposes of appreciating the detection capability of a given seismic network, it is important to recognize, using data from Azgir, that a fully-coupled (tamped) explosion of 1 kt in salt in high-Q areas of the Commonwealth of Independent States (C.I.S.) has an  $m_b$  of 4.4; fully decoupled events of 1 and 10 kt have  $m_b$ 's of about 2.6 and 3.4 respectively (assuming a decoupling factor of 70). These magnitudes are higher than was generally thought during earlier debates on decoupling. Hence, chemical explosions of  $m_b \leq 2.5$  in high Q areas containing salt need not be considered in monitoring a 1 kt threshold treaty or down to that level under a CTBT. Most areas of thick salt deposits in the former U.S.S.R. are typified by high Q (efficient transmission) for P waves and low natural seismic activity. Many of the thick salt deposits of the C.I.S. suitable for construction of large cavities at depth suitable for decoupling, including those few typified by either known natural seismic activity or high attenuation for P waves, are located outside the Russian Republic itself.

Much of the long debate in the United States about the feasibility of conducting and identifying decoupled nuclear explosions in thick salt deposits with yields of say either 1, 10 or 30 kt comes from the very small number--three--of U.S. nuclear explosions in salt, the fact that those events were conducted more than 27 years ago when coverage by sensitive seismic stations and arrays was very limited, and by the very small size, 0.38 kt, of the only U.S. decoupled nuclear explosion in salt, Sterling. Sterling was so small that it was not recorded at teleseismic distances. Environmental considerations, cost and possible continuation of the present testing moratorium will probably prevent the United States in the foreseeable future from conducting a decoupled nuclear explosion in salt in the range 1 to 10 kt (and perhaps from conducting on its territory nuclear events of any size in salt).

The former Soviet Union (FSU), on the other hand, detonated a number of nuclear explosions in salt including tamped events of 1 to 100 kt, a partially decoupled nuclear explosion of 8 kt in the cavity created by a tamped explosion 8 times larger (Adushkin et al., 1993) and six very small nuclear explosions in the water-

filled cavity created by the 25 kt explosion of 1968. Clearly, the decoupling experiment of 1976, with a yield 21 times that of Sterling, is crucial to ascertaining decoupling factors for overdriven (partially decoupled) nuclear weapons tests in the yield range from 1 to 10 kt. The release of additional data on that and other events in salt by the Russian Republic would go far in answering longstanding questions in the United States about decoupling, evasion and the ability to detect decoupled events under either a CTBT or a LYTTBT.

Past nuclear explosions conducted in salt by the FSU for which cavities may remain standing that are large enough for the full decoupling of explosions with yields equal to or larger than 0.5 kt are concentrated in only a few areas. The existence of all cavities of that size or larger that were created by past nuclear explosions in the C.I.S. is known (Tables 1 and 3) since the yields of explosions that created those cavities must be at least 20 times larger in yield than the size of a fully decoupled event that can be detonated in them and at least 12 times larger than that of a nearly-fully decoupled explosion (assuming Salmon/Sterling conditions for the latter). Hence, the monitoring of cavities of that type that may remain standing that were created by past nuclear explosions should be relatively easy at the one kiloton level, providing U.S. stations are allowed to operate under the treaty in the vicinity of the epicenters of those past explosions.

Probably the greatest difficulty in monitoring either a LYTTBT or a CTBT involves cavities created, not by past large nuclear explosions in salt, but by solution mining in other areas of thick salt deposits of either the C.I.S. or the U.S. From an analysis of satellite images Sykes et al. (1993) point out, however, that much of the Pre-Caspian region, including all of the area near Azgir, is arid. Hence, Azgir would not be a suitable place for constructing large cavities in salt by solution mining and then using them for clandestine nuclear testing. An understandable question is then why and how, in such an arid environment did the cavities created by the 1966 and 1968 explosions fill with water? Russian scientists have stated that the water that filled the cavity was of underground origin. I do not have information on how long the cavity took to fill except that it must have filled by the time the first explosion was conducted in it in 1975, seven years later.

The 1966 and 1968 explosions were detonated in the Azgir west salt dome whereas the 1971 explosion, for which the cavity remained dry (Adushkin et al., 1993), and other tamped explosions from 1976 to 1979 were conducted in the Azgir



east dome about 15 km to the east (Sykes et al., 1993). It is not unusual for considerably higher water pressures to exist at depth in the more permeable sediments surrounding salt domes than in the domes themselves. Fractures created by the 1966 and 1968 events may have permitted water to enter the cavities created by those explosions. Would it be possible to exploit such sub-surface waters for solution mining of large cavities? It should be borne in mind, however, that volumes of water many times those the size of the cavity being created are needed in the solution mining process. Whether such large volumes of water are realistically obtainable at Azgir from nearby sub-surface sources is not known. Large volumes of water, if they are in fact available, must then be removed from a large cavity in a clandestine manner if it is then to be used for secret decoupled testing. A prime scenario for constructing such a clandestine cavity would be to use an oil or gas field as a cover operation. Drilling associated with the construction of the cavity might be mistaken for that for oil or gas; nearby existing well might be used for disposal of the brine created by solution mining and the emptying of the cavity. Since satellite images reveal that Azgir is not the site of oil or gas exploitation, however, that scenario is not applicable to that and many other areas within the Pre-Caspian depression. Possible clandestine testing in large cavities created by either solution or conventional mining will be dealt with more fully in a separate paper.

#### ACKNOWLEDGMENTS

I thank Dan Davis and Paul Richards for acting as critical reviewers and V. V. Adushkin, D. D. Sultanov and I. O. Kitov for information on the Soviet program of peaceful nuclear explosions as part of a joint agreement for work on decoupling and nuclear testing between the Institute for Dynamics of the Geospheres, Russian Academy of Sciences, and Lamont-Doherty. Frode Ringdal kindly furnished listings of Norsar detections and magnitudes for the area near Azgir and provided information from Norsar and Hagfors on small events at Azgir. Hans Israelson provided additional information on those events for the Hagfors array. Paul Lyubomirskiy translated a number of papers from Russian on the Pre-Caspian depression. This work was supported by the Dept. of the Air Force, Phillips Laboratory (AFSC), Hanscom Air Force Base, MA under contract F19628-90-K-0059.



## REFERENCES

- Adushkin, V. V., I. O. Kitov, O. P. Kuznetsov and D. D. Sultanov (1993). Seismic efficiency of decoupled nuclear explosions, *Geophys. Res. Lett.* **20**, 1695-1698.
- Adushkin, V. V., I. O. Kitov, O. P. Kuznetsov and D. D. Sultanov (1992b). Experimental results of seismic efficiency of decoupled explosions measurements, in *Abstracts of Papers to be Presented at 14th Annual PL/DARPA Seismic Research Symposium*, Phillips Laboratory, Hanscom Air Force Base, MA., PL-TR-92-2210, ADA256711.
- Berest, P. and D. N. Minh, Stability of cavities in rocksalt (1981). *Proc. Intern. Symp. on Weak Rock, Tokyo*, 473-478.
- Blandford, R. R., R. H. Shumway, R. Wagner and K. L. McLaughlin (1984). Magnitude yield for nuclear explosions at several test sites with allowance for effects of truncated data, amplitude correlation between events within test sites, absorption, and pP, Teledyne Geotech, Alexandria VA, TGAL-TR-83-06, 1-48.
- Bocharov, V. S., S. A. Zelentsov and V. N. Mikhailov (1989). The characteristics of 96 underground nuclear detonations at the Semipalatinsk test range, *Atomic Energy* **67**, 210-214.
- Bogacheva, M. L., Y. M. Vasil'yev, B. K. Proshlyakov, M. M. Charygin and A. G. Shleyfer (1965). A unique sequence of Triassic rocks in the extra deep Aralsor hole (Caspian depression), *Akad. Nauk SSSR, Doklady, Earth Sci.* **165**, (English translation), 33-35.
- Borg, I. (1984) Nuclear explosives--the peaceful side, *New Scientist*, 8 March, pp. 10-13.
- Denny, M. D. and D. M. Goodman (1990). A case study of the seismic source function: Salmon and Sterling reevaluated, *J. Geophys. Res.* **95**, 19,705-19,723.
- Der, Z., T. McElfresh, R. Wagner and J. Burnett (1985). Spectral characteristics of P waves from nuclear explosions and yield estimation, *Bull. Seismol. Soc. Amer.* **75**, 379-390, 1222.
- Elias, M. M., K. Y. Lee and R. J. Sun (1966). *Atlas of Asia and Eastern Europe to Support Detection of Underground Nuclear Testing, 4: Features Affecting Underground Nuclear Testing*, prepared by the U. S. Geological Survey for the Advanced Research Projects Agency, 7 map sheets.
- Healy, J. H., C. Y. King and M. E. O'Neill (1971) Source parameters of the Salmon and Sterling nuclear explosions from seismic measurements, *J. Geophys. Res.* **76**, 3344-3355..
- Herbst, R. F., G. C. Werth and D. L. Springer (1961). Use of large cavities to reduce seismic waves from underground explosions, *J. Geophys. Res.* **66**, 959-978.
- Izrael', Yu. A. and M. P. Grechushkina (1978). The use of peaceful underground nuclear explosions with minimum radioactive contamination of the environment, *Peaceful Nuclear Explosions V*, International Atomic Energy Agency, Vienna, document IAEA-TC-81-5/7, pp. 167-176.
- Jih, R. S. and R. A. Wagner (1991). Recent methodological developments in magnitude determination and yield estimation with applications to Semipalatinsk explosions, Teledyne Geotech, Alexandria VA, final report to Phillips Laboratory, pp. 1-90, PL-TR-91-2212, ADA244503.

- Jenyon, M. K. (1986). *Salt Tectonics*, Elsevier, New York.
- Jordan, J. N., W. V. Mickey, W. Helterbran and D. M. Clark (1966). Travel times and amplitudes from the Salmon explosion, *J. Geophys. Res.* **71**, 3469-3482.
- Kedrovshiy, O. L. (1970). Prospective applications of underground nuclear explosions in the national economy of the USSR, *UCRL-Trans-10477*, (Translation from Russian), Lawrence Radiation Laboratory, University of California, Livermore, CA, 1-47.
- Latter, A. (1960). presentation, in *Technical Working Group 2, Verbatim Record of Seventh Meeting*, Conference on the Discontinuance of Nuclear Weapons Tests, held in Geneva 2 December 1959, GEN/DNT/TWG.2/PV.7 (15 January 1960) Sir William Penny Chairman, pp. 91-110.
- Latter, A. L., R. E. LeLevier, E. A., Martinelli and W. G. McMillan (1961). A method of concealing underground nuclear explosions, *J. Geophys. Res.* **66**, 943-946.
- Marshall, P. D., D. L. Springer and H. C. Rodean (1979). Magnitude corrections for attenuation in the upper mantle, *Geophys. J. R. astr. Soc.* **57**, 609-638.
- Marshall, P. D., R. C. Lilwal, R. C. Stewart and I. Marsden (1991). Seismometer array recordings of P waves from explosions in the North Caspian USSR area, *Atomic Weapons Research Establishment Report O 4/91*, 1-192.
- Murphy, J. R. (1980). An evaluation of the factors influencing the seismic detection of decoupled explosions at regional distances, S-Cubed, Final Report to U.S. Arms Control and Disarmament Agency, SSS-R-80-4579, La Jolla, CA, 1-62.
- Murphy, J. R., J. L. Stevens and N. Rimer (1984). High frequency seismic source characteristics of cavity decoupled underground nuclear explosions, S-Cubed, Maxwell Laboratories, Scientific Report No. 1, SSS-R-88-9595, La Jolla, CA, to Air Force Geophysics Laboratory, Hanscom Air Force Base, Mass., pp. 1-51, AFGL-TR-88-0130, ADA198121.
- Nordyke, M. D. (1975). A review of Soviet data on the peaceful uses of nuclear explosions, *Annals Nuclear Energy* **2**, 657-673.
- Office of Technology Assessment, Congress of the United States (1988). *Seismic Verification of Nuclear Testing Treaties*, OTA-ISC-361, U. S. Government Printing Office, Washington D. C., 1-139.
- Rachlin, J. (1985). Cavity construction opportunities in the Soviet Union, in *Proceedings of the Department of Energy Sponsored Cavity Decoupling Workshop, Pajaro Dunes, California*, D.B. Larson (Editor), Lawrence Livermore National Laboratory, Livermore, CA, Conference 850779, 53-66.
- Rawson, D., P. Randolph, C. Boardman and V. Wheeler (1966). Post-explosion environment resulting from the Salmon event, *J. Geophys. Res.* **71**, 3507-3521.
- Ringdal, F. (1976). Maximum-likelihood estimation of seismic magnitude, *Bull. Seismol. Soc. Amer.* **66**, 789-802.
- Rodean, H. C. (1981). Inelastic processes in seismic wave generation by underground explosions,

- in *Identification of Seismic Sources--Earthquake or Underground Nuclear Explosion*, E. S. Husebye and S. Mykkeltveit, (Editors), Reidel, Dordrecht, 97-189.
- Scheimer, J. F. and I. Y. Borg (1984). Deep seismic sounding with nuclear explosives in the Soviet Union, *Science* **226**, 787-792.
- SIPRI Seismic Study Group (1968). *Seismic Methods for Monitoring Underground Explosions*, D. Davies, Rapporteur, International Institute for Peace and Conflict Research (SIPRI), Stockholm, 1-130.
- Springer, D., M. Denny, J. Healey and W. Mickey (1968). The Sterling experiment: decoupling of seismic waves by a shot-generated cavity, *J. Geophys. Res.* **73**, 5995-6001.
- Stevens, J. L., J. R. Murphy and N. Rimer (1991). Seismic source characteristics of cavity decoupled explosions in salt and tuff, *Bull. Seismol. Soc. Amer.* **81**, 1272-1291.
- Sultanov, D. D. et al. (1993). Investigation of seismic efficiency of Soviet peaceful nuclear explosions conducted in various geological conditions, Part 1. Institute for Dynamics of Geospheres, Moscow, 220 p.
- Sykes, L.R. (1992). Yields of Underground Nuclear Explosions at Azgir and Shagan River, USSR and Implications for Identifying Decoupled Nuclear Testing in Salt, Sci. Rpt. 1, PL-TR-92-2002, Phillips Laboratory, Hanscom Air Force Base, MA, 34 pp, ADA250971.
- Sykes, L. R., J. Deng and P. Lyubomirskiy (1993). Accurate location of nuclear explosions at Azgir, Kazakhstan, from satellite images and seismic data: implications for monitoring decoupled explosions, *Geophys. Res. Lett.* **20**, 1919-1922.
- Sykes, L. R. and G. Ekstrom (1989). Comparison of seismic and hydrodynamic yield determinations for the Soviet joint verification experiment of 1988, *Proc. Natl. Acad. Sci. USA*, **86**, 3456-3460.
- Sykes, L. R. and P. Lyubomirskiy (1992). Analysis of small seismic events near Azgir, Kazakhstan: implications for identifying chemical and decoupled nuclear explosions in a major salt dome province, in *Papers Presented at 14th Annual PL/DARPA Seismic Research Symposium*, Phillips Laboratory, Hanscom Air Force Base, MA., PL-TR-92-2210, ADA256711.
- Sykes, L. R. and S. Ruggi (1986) Soviet underground nuclear testing: inferences from seismic observations and historical perspective, *Working Paper NWD 86-4*, Natural Resources Defense Council, Washington, D. C., 1-88.
- Sykes, L. R. and S. Ruggi (1989). Soviet nuclear testing, in *Soviet Nuclear Weapons, Nuclear Weapons Databook, IV*, T. B. Cochran et al. (editors), Ballanger, New York, 332-382.
- Werth, G. and P. Randolph (1966). The Salmon seismic experiment, *J. Geophys. Res.* **71**, 3405-3413.

Table 1. Tamped Underground Nuclear Explosions in or Near Thick Salt Deposits of C.I.S.

Area	Date		Origin Time Hr. Mn. Sec.	Lat. (°N)	Long. (°E)	mb	n	Y (mb) in kt*	Y (announced) in kt	Y <sub>FD</sub> in kt (if ≥ 0.5 kt)
	Day	Mon. Year								
Azgir	22 Apr	1966	02 58 02.1	47.88	47.89	4.524 ± .056	16	1.3	1.1	-
Bukhara II	21 May	1968	03 59 10.0	38.89	65.10	5.4	135		40.	2.0
Azgir	01 July	1968	04 01 59.9	47.91	47.91	5.529 ± .027	20	21.	25.	1.1
Orenburg	25 June	1970	04 59 52.4	52.2	55.7	4.9		3.7		-
Orenburg	22 Oct	1971	05 00 02.1	51.57	54.52	5.260 ± .043	23	10.1	15.	0.7
Azgir	22 Dec	1971	07 00 00.1	47.90	48.13	6.064 ± .020	22	93.	64.	2.7
Kharkov area	09 July	1972	06 59 57.9	49.78	35.45	4.8	62	2.8		-
Lake Aralsor (clay)	20 Aug	1972	03 00 01.2	49.41	48.13	5.750 ± .037	22	39.		2.0
W. of Orenburg	21 Sept.	1972	09 00 01.4	52.19	51.94	5.0	83	4.9		-
NE of Elista (clay)	03 Oct	1972	08 59 57.8	46.86	44.87	5.864 ± .050	21	54.		2.7
W. of Orenburg	24 Nov	1972	09 00 02.9	52.14	51.83	4.5	40	1.2		-
Orenburg	30 Sept	1973	05 00 00.3	51.60	54.51	5.213 ± .047	20	8.9		-
Azgir	29 Jul	1976	05 00 01.4	47.87	48.14	5.884 ± .015	30	57.		2.8
S. of Mirnyy (dolomite)	05 Nov	1976	03 59 56.9	61.52	112.73	5.100 ± .052	16	6.5		-
Azgir	30 Sept	1977	06 59 59.4	47.89	48.15	4.994 ± .029	25	4.8		-
S. of Mirnyy (dolomite)	08 Oct	1978	00 00 00.	61.53	112.87	5.249 ± .034	39	9.8		-
Azgir	17 Oct	1978	05 00 00.0	47.86	48.11	5.851 ± .014	63	52.		2.6

Area	Date		Origin Time Hr. Mn. Sec.	Lat. (°N)	Long. (°E)	mb	n	Y (mb) in kt*	Y (announced) in kt	Y <sub>FD</sub> in kt (if ≥ 0.5 kt)
	Day	Mon. Year								
Azgir (clay)	18 Dec	1978	08 00 00.0	47.85	48.14	5.977 ± .012	65	73.		3.6
Azgir	17 Jan	1979	07 59 59.1	47.92	48.12	6.027 ± .013	58	84.		4.2
Azgir	14 July	1979	04 59 58.8	47.88	48.12	5.620 ± .012	59	27.		1.4
S. of Mirnyy (dolomite)	07 Oct	1979	20 59 56.9	61.85	113.12	5.0	110	4.9		-
Azgir	24 Oct	1979	06 00 00.3	47.85	48.12	5.762 ± .015	69	41.		2.0
Astrakhan	08 Oct	1980	06 00 00.2	46.75	48.24	5.184 ± .038	44	8.2		-
Kuyumba (dolomite)	01 Nov	1980	12 59 58.0	60.79	97.57	5.208 ± .034	40	8.7		-
Astrakhan	26 Sept	1981	05 00 00.2	46.80	48.25	5.146 ± .035	48	7.4		-
Astrakhan	26 Sept	1981	05 04 00.0	46.81	48.26	5.183 ± .034	36	8.2		-
W. Tura	25 Sept	1982	17 59 57.4	64.33	91.80	5.211 ± .034	41	8.8		-
S. of Mirnyy (dolomite)	10 Oct	1982	04 59 56.9	61.53	112.86	5.323 ± .028	40	12.		0.6
Astrakhan	16 Oct	1982	06 00 00.1	46.77	48.20	5.230 ± .033	33	9.3		-
Astrakhan	16 Oct	1982	06 05 00.1	46.75	48.20	5.272 ± .031	36	10.4		0.5
Astrakhan	16 Oct	1982	06 10 00.1	46.78	48.23	5.255 ± .035	35	10.0		0.5
Astrakhan	16 Oct	1982	06 15 00.1	46.75	48.24	5.381 ± .034	42	14.		0.7
Karachaganak	10 July	1983	04 00 00.2	51.35	53.23	5.313 ± .029	48	12.		0.6
Karachaganak	10 July	1983	04 05 00.1	51.35	53.24	5.350 ± .031	48	13.		0.6

Area	Date		Origin Time Hr. Mn. Sec.	Lat. (°N)	Long. (°E)	mb	n	Y (mb) in kt*	Y (announced) in kt	Y <sub>IP</sub> in kt (if ≥ 0.5 kt)
	Day	Mon. Year								
Karachaganak	10 July	1983	04 10 00.1	51.36	53.25	5.235 ± .027	44	9.4	-	-
Astrakhan	24 Sept	1983	05 00 00.0	46.79	48.26	5.159 ± .070	32	7.6	-	-
Astrakhan	24 Sept	1983	05 05 00.1	46.80	48.23	5.100 ± .046	31	6.5	-	-
Astrakhan	24 Sept	1983	05 10 00.0	46.78	48.27	4.996 ± .046	26	4.9	-	-
Astrakhan	24 Sept	1983	05 15 00.2	46.78	48.24	5.175 ± .040	24	8.0	-	-
Astrakhan	24 Sept	1983	05 20 00.1	46.79	48.22	5.342 ± .033	31	13.	0.6	0.6
Astrakhan	24 Sept	1983	05 25 00.0	46.79	48.24	5.267 ± .044	29	10.3	0.5	0.5
Karachaganak	21 July	1984	03 00 00.0	51.34	53.24	5.331 ± .028	52	12.3	0.6	0.6
Karachaganak	21 July	1984	03 04 59.9	51.38	53.26	5.264 ± .026	52	10.2	0.5	0.5
Karachaganak	21 July	1984	03 10 00.0	51.35	53.25	5.323 ± .026	49	12.	0.6	0.6
Astrakhan	27 Oct	1984	06 00 00.2	46.88	48.11	5.018 ± .042	38	5.2	-	-
Astrakhan	27 Oct	1984	06 05 00.1	46.86	48.03	5.082 ± .046	38	6.2	-	-
Perm Region (limestone)	19 Apr	1987	03 59 57.1	60.62	57.2	4.5	17	1.2	-	-
Perm Region (limestone)	19 Apr	1987	04 04 55.9	60.8	57.5	4.5	19	1.2	-	-
S. of Mirnyy (dolomite)	06 July	1987	23 59 56.9	61.50	112.83	5.1	70	6.5	-	-
S. of Mirnyy (dolomite)	24 July	1987	01 59 56.9	61.46	112.78	5.1	65	6.5	-	-
S. of Mirnyy	12 Aug	1987	01 29 57.1	61.46	112.79	5.0	60	4.9	-	-



Area	Date Day Mon. Year	Origin Time Hr. Mn. Sec.	Lat. (°N)	Long. (°E)	$m_b$	$n$	$Y$ (mb) in $kt^*$	$Y$ (announced) in $kt$	$Y_{FD}$ in $kt$ (if $\geq 0.5$ $kt$ )
SE side, Pre-Caspian depression	03 Oct 1987	15 14 57.7	47.62	56.20	5.3	65	11.		0.6
E. of Kotlas (anhydrite, dolomite)	06 Sept 1988	16 19 58.7	61.33	47.98	4.8	57	2.8		-

\* calculated from  $m_b = 4.4250 + 0.832 \log Y$

**Table 2. Nuclear Explosions in Cavities in Salt at Azgir**

Date	Hr.Min.	$m_b \pm SEM$	$n(m_b)$	Yield (kt)
25 Apr. 1975	05 00	4.45 $\pm$ .13	7	1.1
*29 Mar. 1976	07 00	4.06 $\pm$ .04	7	8**
14 Oct. 1977	07 00	3.42	1	0.06
30 Oct. 1977	07 00	2.77	1	0.01
12 Sept 1978	05 00	3.02	1	0.02
30 Nov. 1978	08 00	3.07	1	0.02
10 Jan. 1979	08 00	4.36 $\pm$ .14	2	0.8

-----  
 \*In air-filled cavity created by 64 kt explosion of 1971; all other events in water-filled cavity created by 25 kt explosion of 1968.

\*\*Yield of 1976 event from Adushkin et al.(1993); other yields, Y, from  $m_b = 4.425 + 0.832 \log Y$

**Table 3. Inventory of Large Cavities Produced by Past Nuclear Explosions in or near Thick Salt Deposits of C.I.S. that may Remain Standing that might be used to Conduct Fully Decoupled Nuclear Tests of Yield,  $Y_{FD} \geq 0.5$  kt.**

REGION	$Y_{FD}(kt)$				
	0.5-0.9	1.0-1.9	2.0-2.9	3.0-3.9	4.0-4.2
<b>Pre-Caspian Depression</b>					
Azgir		2	4	1*	1
Astrakhan	5				
Karachaganak	5				
Lake Aralsor			1*		
Other	2		1*		
<b>Bedded salt to NW of Lake Baikal</b>	1				
<b>Central Asia - Bukhara</b>			1		
<b>Totals</b>	13	2	7	1*	1

-----  
 \* Reported by Russian workers as detonated in clay




Fig. 1. North Caspian Region showing sites of nuclear explosions in and near thick salt deposits (squares) in Pre-Caspian depression and area of special study of small seismic events (boxed region).




Fig. 2. Seismic magnitude as a function of yield for underground nuclear explosions conducted under various testing conditions (hatched areas) in Commonwealth of Independent States (C.I.S.). SW Shagan River denotes regression line for fully-coupled nuclear explosions in southwestern part of that testing area in eastern Kazakhstan (Sykes, 1992). Three upper dots denote data points for fully-coupled explosions in salt at Azgir; dot labelled 1976 denotes partially decoupled explosion at Azgir of March 1976; 150 kt denotes yield limitation of Threshold Test Ban Treaty. Present identification threshold using seismic stations solely external to C.I.S. and range of thresholds with seismic network in C.I.S. from Office of Technology Assessment (1988).




Fig. 3. Top: Calculated maximum fully-decoupled yield,  $Y_{FD}$ , divided by yield,  $Y$ , of tamped explosion creating a cavity of usable volume  $V_C$  as a function of depth,  $h$ , for tamped nuclear explosions in salt for which information has been released on  $Y$ ,  $h$  and  $V_C$ . Data from 1971 explosion were not included in either two regression lines shown. Solid line is regression based on data from four other explosions; dashed, that based on those four data points minus that for shallow event of 1966.

# ACCURATE RELOCATION OF NUCLEAR EXPLOSIONS AT AZGIR, KAZAKHSTAN, FROM SATELLITE

## IMAGES AND SEISMIC DATA : IMPLICATIONS FOR MONITORING DECOUPLED EXPLOSIONS

Lynn Sykes<sup>1</sup>, Jishu Deng<sup>1</sup>, and Paul Lyubomirskiy

Lamont-Doherty Earth Observatory, Palisades, NY 10964

<sup>1</sup>Also at Department of Geological Sciences of Columbia University, NY, NY 10027

*Abstract.* The 10 largest tamped nuclear explosions detonated by the Former Soviet Union in and near two salt domes near Azgir were relocated using seismic data and the locations of shot points on a SPOT satellite image taken in 1988. Many of the shot points are clearly recognized on the satellite image and can be located with an accuracy of 60 m even though testing was carried out at those points many years earlier, i. e. between 1966 and 1979. Onsite inspections and a local seismic monitoring network combined with our accurate locations of previous explosions would insure that any cavities that remain standing from those events could not be used for undetected decoupled nuclear testing down to a very small yield. Since the Azgir area, like much of the Pre-Caspian depression, is arid, it would not be a suitable place for constructing large cavities in salt by solution mining and then using them for clandestine nuclear testing.

### Introduction

The former Soviet Union ( FSU ) conducted a large number of nuclear explosions in thick salt deposits, especially in the Pre-Caspian depression to the north of the Caspian Sea, the world's largest salt dome province. Most of the larger explosions were conducted near the small town of Azgir in westernmost Kazakhstan, a remote site which appears to have been a Soviet testing area for nuclear explosions for peaceful purposes [Sykes, 1991]. A major issue in the monitoring of treaties to further limit or ban nuclear testing has been the possibility that large cavities in salt domes created by either past large nuclear explosions or solution mining could be used to reduce the amplitudes of seismic waves of small nuclear explosions detonated in those cavities [Office of Technology Assessment, 1988]. An event of that type is a so-called decoupled explosion. The FSU detonated several nuclear explosions in large cavities created by tamped (full coupled) events

at Azgir, including a partially decoupled explosion of eight kilotons (kt) in 1976 in an air-filled cavity created by a 64 kt explosion in 1971 [Sykes and Lyubomirskiy, 1992; Adushkin et al., 1993].

To better identify attempts to use cavities of that kind for future decoupled nuclear testing, it is helpful to know accurately the locations of past large nuclear explosions in salt domes and other thick salt deposits. In this paper we use images from the SPOT satellite and seismic data to relocate the 10 largest tamped nuclear explosions at Azgir. From the SPOT data taken in 1988 we obtain absolute locations of likely shot points of these events, which occurred from 1966 to 1979. We use P-wave arrival time data from large numbers of seismic stations for each explosion to obtain relative locations with respect to a master event so as to identify the shot locations on SPOT images that correspond to explosions detected seismically on specific dates. Thurber et al. [1993] used a similar technique to associate shot positions at the Balapan (Shagan River) portion of the eastern Kazakhstan nuclear weapons test site with dates of explosions detected seismically. Leith and Simpson [1990] discuss the use of satellite images in locating seismic events. Marshall et al. [1991] used the Joint Hypocentral Determination (JHD) method to obtain relative locations of 9 of the 10 Azgir events we studied. We include newly released arrival times for the 10 events from standard seismograph stations in the FSU. In addition we comment on monitoring the Azgir region under possible future treaties to ban or limit the testing of nuclear weapons and on the unsuitability of the arid Azgir region for the construction of large cavities by solution mining.

## Data and Methods

We purchased the digital data for three spectral bands of the SPOT imagery taken on September 14, 1988. It covers a region 60 by 60 km centered on 47°59.9'N, 48°01.5'E with a pixel size of 20 m. In our work we processed a number of images of that area and portions thereof. Fig. 1 is a 30 by 20 km part of the original image. The town of Azgir is located at the convergence of roads at the lower left. Clear indicators of past nuclear testing, including disturbed areas and convergence of roads near shot points, can be seen in the area 14 to 18 km to the east and northeast of Azgir. Seven of the eight shot points in that area are clearly visible on the satellite images. Two other areas of ground disturbance and convergence of roads about 6 to 8 km NNW of Azgir, as we will see from the relative seismic locations, appear to have been the shot points of the 1.1 kt explosion



of 1966 and the 25 kt event of 1968 [Kedrovskiy, 1970; Sykes, 1991]. The southernmost shot point of the two, which we associate with the 1966 explosion, is less clear on Fig. 1. We used locations of about 30 road intersections from available maps to provide absolute locations for points on the SPOT image. Individual shot points can be measured with an accuracy of about 60 m (one standard deviation).

In obtaining our seismic locations we used the P wave arrival times from seismic stations within 100° of Azgir from the bulletins of the International Seismological Center (ISC) and standard stations of the national network of the FSU. Fig. 2A shows the 359 stations in that distance range that reported one of the largest Azgir events, that of October 17, 1978, and Fig. 2B shows the stations that reported both that event and the small event of 1.1 kt in 1966. The master-event technique [Bullen and Bolt, 1985] was used to calculate the relative locations and origin time corrections of the nine other events in Table 1 with respect to the master event of October 17, 1978. The depths were held fixed in the calculations. The IASPEI91 earth model was used to calculate ray-parameters. We assumed that the data for a given event with respect to the master event satisfy a normal distribution, and used a Gaussian weighting function to improve the results and minimize uncertainties. We calculated 95% confidence ellipses for the relative epicentral location of each event and show them in Figs. 3 and 4 and Table 1. From Table 1 it is clear that at least 124 stations were used in the relocations except for the smallest event, that of 1966, for which 42 relative P times were available.

## Results

Table 1 lists the events studied, their revised locations and origin times. The origin time of the master event of October 17, 1978 was set at 05 00 00.0 UCT. The origin times of the other 9 explosions were calculated using the calculated hours and minutes in the ISC bulletins and then adding to zero seconds the time corrections calculated with respect to the assumed origin time of the master event. The uncertainty in the calculated time corrections is smaller than 0.1 s. The fact that two of the 9 events have calculated origin times within 0.1 s of an exact hour argues strongly that those events and the master event, in fact, occurred exactly on an hour to within 0.1 s. Likewise, the 1968 event occurred within 0.1 s of an exact minute. All of the events from 1971 to

1979 occurred within 1.4 s of an exact hour. It is a common procedure for nuclear explosions of many countries to be scheduled and then detonated either on an exact hour or minute since the activities of many people and much equipment at many locations involved in monitoring nuclear tests must be coordinated.

The relative locations also show two distinctive groupings of explosions (Fig. 3). The first group consists of the two early explosions of 1966 and 1968, which were detonated north of the town of Azgir in the west Azgir salt dome [Kedrovskiy, 1970; Ministry of Geology, 1983]. The other explosions from 1971 to 1979 were detonated about 18 km east of the first group (upper-right Fig. 1) in the east Azgir salt dome [Ministry of Geology, 1983; Adushkin et al. 1993]. Fig. 4, which is an expanded view of the northeastern part of Fig. 3, indicates our preferred match between SPOT image measurements and seismic locations. Only one of the shot points on the SPOT image in Fig. 4 is somewhat uncertain; the others are very clear picks. The locations from the satellite image in Fig. 4 are more tightly clustered, especially in the east-west direction, than the seismic locations. Since the salt structures in the Azgir area have an upper relief of several kilometers, the use of a standard, spherically symmetrical velocity model in computing the relative seismic locations probably contributes to the blurring of the seismic image.

When the seismic location of the master event is co-located with the shot point we associate with it on the SPOT image, the seismic locations of the other events exhibit a systematic bias in latitude with respect to the SPOT locations. To reduce that bias we allowed the seismic and satellite locations of the master event to differ. We obtained the match in Fig. 4 by minimizing the sum of the squares of the lengths of the arrows connecting the inferred locations by the two methods. That procedure also resulted in the centroid of the seismic locations exactly coinciding with that from the satellite locations. In that case six of the seven 95% confidence ellipses in Fig. 4 overlap at least one possible shot point measured from the SPOT image. The confidence ellipses are generally elongated in the northeast-southwest directions, indicative of the smaller number of stations in those quadrants (Fig. 2).

Russian scientists have stated in writing to us that the explosion on Dec. 18, 1978 was detonated in clay while all of the other events at Azgir were detonated in salt. That event is correlated in Fig. 4 with the only shot point in Fig. 1 for which the false color of the disturbed area near it is red and not white. It is also the only event for which the 95% confidence ellipse does not overlap the corresponding shot point in Fig. 1. An alternate solution is to associate the master event with the southernmost shot point on the satellite image. In that case, however, a large misfit in seismic and

SPOT locations cannot be avoided for the explosion of Dec. 18, 1978. In all of these solutions, however, the association of shot points with seismic locations remains the same for the four northernmost events of Fig. 4.

Our relative seismic locations are very similar to those obtained by Marshall et al. [1991]. Their epicenters differ from the satellite locations on the average by only 2.9 km. The ISC locations show more scatter and differ from the satellite locations on the average by 8.4 km.

Shot depths of 165, 590 and 987 m are published for the explosions of 1966, 1968 and 1971 [Kedrovskiy, 1970; Adushkin et al., 1993]. Since the calculated origin time of the 1971 event in Table 1 is within 0.1 s of an exact hour, we can consider the depths of all of the events, including the master, to be normalized to that of the 1971 event, 987 m. Correcting the 1966 and 1968 events for difference in depth of focus with respect to the 1971 explosion using a salt velocity of 4.2 km/s [Adushkin et al., 1993] would make their origin times earlier than those in Table 1 by 0.2 and 0.1 s. Hence, differences in origin time larger than a few tenths of a second are unlikely to be associated with different depths of detonation. Thus, most of the departures of the calculated origin times from an exact minute in Table 1 probably reflect that some of them were detonated as much as 1 or 2 s off an exact minute or that large differences in structure exist at depth beneath the various shot points.

Some of the shot points to the northeast of Azgir in Fig. 1 are characterized by at least two nearby disturbed areas. Some of these may be additional holes drilled for instrument placement. What we identify as the location of the 1971 explosion in Fig. 4 is characterized by two disturbed areas about 200 m apart of nearly equal intensity in Fig. 1. One of these could be the re-entry hole into the 1971 cavity that was used for the placement of the device for the partially decoupled explosion of 1976 [Adushkin et al., 1993].

## Discussion

Eight of the ten shot points of nuclear explosions conducted near Azgir from 1966 to 1979 are clearly visible on the satellite images taken in 1988. The seismic and satellite data taken together result in quite accurate locations for all 10 events. A combination of onsite inspections, repeated satellite imagery of high resolution and a local seismic monitoring network would insure that any

cavities that remain standing from those events could not be used for undetected decoupled nuclear explosions down to a very small yield. The NORSAR and Hagfors arrays can detect small events of body wave magnitude as low as 2.5 from the Azgir region [Ringdal, 1981; Ringdal and Husebye, 1982; Sykes and Lyubomirskiy, 1992]. Since they are located more than 2000 km from Azgir, the location capability of those arrays by themselves is poor compared with that of a local seismic network. Sykes and Lyubomirskiy [1992] found that the rate of occurrence of chemical explosions in a large area surrounding Azgir is low and that of small earthquakes is even lower. Several small events at Azgir that occurred almost exactly on the hour from 1975 to 1979 are taken to be small nuclear explosions and will be the subject of another paper.

An examination of Fig. 1 and of the entire 60 by 60 km SPOT image that we processed indicates that the region is arid and sparsely populated. The many salt flats and small playa lakes in Fig. 1 and in the larger image indicate a lack of fresh water in almost all of the area. Several small rivers of the Pre-Caspian depression drain into ephemeral lakes [Ministry of Geology, 1983] and not into the two major rivers of the region, the Ural and Volga, that drain into the Caspian Sea. On most of the 60 by 60 km SPOT image the percentage of cultivated land is even smaller than that in Fig. 1. The area near Azgir, like much of the Pre-Caspian depression, would not be one in which large cavities in salt domes could be constructed by solution mining since that process requires great quantities of fresh water. Therefore, it would be practical to excavate such cavities for clandestine decoupled nuclear testing in the Pre-Caspian depression only in a few areas near major rivers and along the coast of the Caspian Sea.

Acknowledgements. We thank D. D. Sultanov for providing a catalog of data for the 10 Azgir explosions for standard stations of the national network of the FSU. We thank D. Davis, W. Y. Kim and P. Richards for critically reading the manuscript and for helpful suggestions. This research was supported by contract F19628-90-K-0059 from the Phillips Laboratory, Hanscom Air Force Base, MA. Lamont-Doherty Earth Observatory Contribution 00000.

## REFERENCES

- Adushkin, V. V., I. O. Kitov, O. P. Kuznetsov and D. D. Sultanov, Seismic efficiency of decoupled nuclear explosions, in press, *Geophys. Res. Lett.*, 1993.
- Bullen, K. E. and B. A. Bolt, *An Introduction to the Theory of Seismology*, 4th Ed., p.273, Cambridge University Press, 1985.
- Kedrovskiy, O. L., Prospective applications of underground nuclear explosions in the national economy of the USSR, UCRL-Trans-10477, 47 pp., Lawrence Radiation Laboratory, University of California, 1970.
- Leith, W. and D. W. Simpson, Monitoring underground nuclear tests, in *Commercial Observation Satellites and International Security*, St. Martin's Press, N. Y., pp. 114-129, 1990.
- Marshall, P. D., R. C. Lilwall, R. C. Stewart and I. Marsden, Seismometer array recordings of P waves from explosions in the north Caspian USSR area, Atomic Weapons Establishment (U. K.), AWE Report No. O 4/91, 91 pp., 1991.
- Ministry of Geology, U.S.S.R. and Kazakh SSR Ministry of Geology, Structural chart of the surface of the salt-bearing complex in the Pre-Caspian depression, *Aerogeologia and Kazneftegazgeologia*, 4 sheets, scale 1:1, 000, 000, ed. by Volchegurckiy, L. F., O. S. Turkov and A. E. Shlezenger, 1983.
- Office of Technology Assessment, Congress of the United States, *Seismic Verification of Nuclear Testing Treaties*, OTA-ISC-361, U. S. Government Printing Office, Washington, DC, 139 pp., 1988.
- Ringdal, F. and E. S. Husebye, Application of arrays in the detection, location, and identification of seismic events, *Bull. Seism. Soc. Am.*, **72**, S201-S224, 1982.
- Ringdal, F., Location of regional events using travel time differentials between P arrival branches,

NORSAR Scientific Report No. 2-80/81, pp.60-69, 1981.

Sykes, L. R, Yields of underground nuclear explosions at Azgir and Shagan River, USSR and implications for identifying decoupled nuclear testing in salt, PL-TR-92-2002, 34 pp., Phillips Laboratory, Hanscom Air Force Base, MA, ADA250971, 1991.

Sykes, L. R. and P. Lyubomirskiy, Analysis of small seismic events near Azgir, Kazakhstan: implications for identifying chemical and decoupled nuclear explosions in a major salt dome province, in Papers Presented at 14th Annual PL/DARPA Seismic Research Symposium, Phillips Laboratory, Hanscom Air Force Base, MA, pp. 415-421, PL-TR-92-2210, ADA256711, 1992.

Thurber, C. H, H. R. Quin and P. G. Richards, Accurate locations of nuclear explosions in Balapan, Kazakhstan, 1987 to 1989, *Geophys. Res. Lett.*, 20-5, 399-402, 1993.

TABLE 1. Origin Times and Locations for the 10 Largest Nuclear Explosions Near Azgir.

Date Day Mon. Year	Origin Time			SPOT Location		95% Confidence Ellipses <sup>1</sup>			Number of Stations
	Hr.	Mn.	Sec.	Lat.(N)	Lon.(E)	Maj.(km)	Min.(km)	Angle	
22 Apr. 1966	02	58	02.1	47°53.02'	47°53.32'	6.6	6.3	-28.5	42
01 July 1968	04	01	59.9	47°54.54'	47°54.84'	4.2	3.0	64.4	124
22 Dec. 1971	07	00	00.1	47°53.80' 47°53.79'	48°07.82' 48°07.66'	2.5	1.8	35.9	170
29 July 1976	05	00	01.4	47°52.25'	48°08.32'	2.4	1.6	45.1	216
30 Sept 1977	06	59	59.4	47°53.27'	48°09.13'	3.2	2.3	47.9	129
17 Oct. 1978	05	00	00.0	47°51.82'	48°06.81'				359
18 Dec. 1978	08	00	00.0	47°51.11'	48°08.65'	2.0	1.3	41.8	281
17 Jan. 1979	07	59	59.1	47°55.14'	48°07.30'	2.2	1.3	45.3	252
14 July 1979	04	59	58.8	47°52.89'	48°07.20'	2.1	1.4	40.2	248
24 Oct. 1979	06	00	00.3	47°50.78'	48°07.36'	2.2	1.4	35.2	266

1. Maj. and Min. are dimensions of semi-major and semi-minor axes of confidence ellipse. Angle in degrees from north to one of semi-major axis; clockwise is positive.

Fig. 1. Digitally processed image of SPOT photography taken on September 14, 1988 of nuclear testing areas near Azgir, west Kazakhstan. The latitudes and longitudes of the four corners clockwise from the northwestern are as follows: 47°58.19'N, 47°52.23'E; 47°54.36'N, 48°15.95'E; 47°43.77'N, 48°12.11'E; 47°47.60'N, 47°48.43'E. Town of Azgir is located at convergence of roads at lower left. Sites of eight explosions from 1971 to 1979 can be seen at upper right of center where shot points and roads are still clearly visible. Two explosions in 1966 and 1968 occurred to the north of Azgir in a separate area. (Image made from base photo, © 1988, CNES/SPOT Image Corporation.)






10 km


5

0





A



B

Fig. 2. ( A ). Station distribution centered on master event of Oct. 17, 1978. Stations farther than 100° were not used in this study. ( B ). Stations that reported both the master event and the small event of 1.1 kt on April 22, 1966.




Fig. 3. Calculated relative locations (pluses) of Azgir explosions with respect to master event with 95% confidence ellipses. Three solid circles are possible sites of nuclear explosions from the SPOT image of Fig. 1. On right hand side, only the inferred SPOT location of the master event is plotted. Town of Azgir is at lower left corner.




Fig. 4. Enlarged view of testing area in Fig. 3 to northeast of town of Azgir. Arrows connect relative locations from seismic data (pluses) with inferred shot points from SPOT image (solid circles). Confidence ellipses indicate uncertainties in seismic locations for other explosions with respect to master event. Event of Dec. 18, 1978 was detonated in clay; all others were in salt.

**CATALOG OF REPORTS FROM STANDARD STATIONS OF USSR FOR  
TEN TAMPED NUCLEAR EXPLOSIONS AT AZGIR**

**(Provided by D. D. Sultanov)**

22.04.1966г. (Северное Каспийского моря - АЗГИР-I)

(02-58)

УСС: Н=02-58-04,0

$\varphi = 47^{\circ}93'N$

$\lambda = 47^{\circ}69'E$   $h = 0км$

МВ = 4.7

ЕССН: 0-02-58-02,27

$\varphi = 47^{\circ}796'N$

$\lambda = 47^{\circ}917'E$   $h = 0км$

расчет -  $h = 33км$

Северное Каспийского моря

№ пп	Станция	Az	$\Delta^{\circ}$	Приб.	Фазы	Время	T сек	M			Mag	L				
								NS	EW	Z						
1.	Грозный (GRO)		4,55 <sup>X</sup>	СК	eeP	02-59-15	1,0			0,4						
						03-00-48	1,8	0,62								
					L <sub>R</sub>	02-04	9,0			0,6	3,20ML					
2.	Махацкала (MAK)		4,7 <sup>X</sup>	СК	ePcy	02-59-17,8	0,6	0,53								
					L <sub>R</sub>	03-06-02	7,0			0,32	3,09ML					
3.	Кр. поляна (KPR)		6,4 <sup>X</sup>	СКМ	iP <sub>1</sub>	02-59-41,8	0,5			0,038						
					P <sub>2</sub>	03-00-16,8	0,6			0,053						
4.	Бакуриани (BKR)		6,5 <sup>X</sup>	СКМ	iP	02-59-42,8	0,9			0,485						
				СК		1,0				0,1						
				СКМ	iX	03-00-31,0	1,0			0,003						
5.	Абастумани (ABS)		6,6 <sup>X</sup>	BCX	eP	02-59-45										
6.	Гори (GOR)		6,7 <sup>X</sup>	СКМ-3	eeP	02-59-45,5	0,8			0,071						
					S <sub>1B3</sub>		1,1	0,128								
					S <sub>2cy</sub>	03-01-57,5	2,6	0,159								
7.	Кировабад		6,8 <sup>X</sup>	СК	P	02-59-47,0										
	Зочи		6,8 <sup>X</sup>	СКМ	iP <sub>1</sub>	02-59-47,0	0,5			0,12						
					P <sub>2</sub>	49,0	0,6			0,08						
9.	Степанован		6,9 <sup>X</sup>	СХ	eP	02-59-48,4										
10.	Герис 188		8,3	СК	eP	03-00-04						-0,				
11.	Алушта		9,3 <sup>X</sup>	СХ	eP	03-00-22,3										
12.	Ялта		9,5 <sup>X</sup>	СХ	iP <sub>1</sub>	03-00-25,1	0,45			0,033						
					P <sub>2</sub>	37	0,6			0,023						
13.	Обнинск		9,5 <sup>X</sup>	Бен-Z	eP	03-00-24,6										
14.	Ст. №1 (Михнево)		9,5 <sup>X</sup>	СКМ	iP	03-00-26	0,45			0,123						
				УСФ	UP		0,4			0,142						
				СКМ	S	02-00,5	0,65			0,072						
				УСФ			0,8			0,129						
				СК			0,8			0,167						

22.04.66

№ пп	Станция	Az°	Δ°	Приб.	Фаза	Время t	T сек	Am			Mag	Res
								NS	EW	Z		
15.	Свердловск		11,1 <sup>X</sup>	СКМ-3	eP	03-00-47,0						
16.	Симферополь		11,25 <sup>X</sup>	СХ <sub>23</sub>	eeP	03-00-49,0	1,0			0,057		
17.	Ашхабад		11,9 <sup>X</sup>	СК	eP	03-01-03						
18.	Боровое		14,6 <sup>X</sup>	СКМ	eP	03-01-32,9	0,8				0,006	
				(S)			1,0			0,008		
19.	Рахов		15,1 <sup>X</sup>	СКМ	eP	03-01-99,5						
20.	Мехгорье		15,55 <sup>X</sup>	СКМ	eX <sub>2</sub>	02-05,2						
21.	Куляб		16,0 <sup>X</sup>	СКМ	eP	03-01-51,8						
22.	Ужгород		16,4 <sup>X</sup>	СКМ	eP	03-01-57,3	0,6				0,016	
23.	Гарм	110	18,4	СКМ	eP	03-02-16,8					0,03	
24.	Андижан		18,6 <sup>X</sup>	СКМ	P	03-02-23,4	0,6				0,07	5,07 MPV
25.	Джержиталь		18,85 <sup>X</sup>	ВЭГИК	P	03-02-27,1	0,1					
26.	Или		20,1 <sup>X</sup>	СК	eP	03-02-41,4						
27.	Апатиты		20,3 <sup>X</sup>	СКМ	<sup>р</sup> MP	<sup>03-02-44</sup> 26,2	18	1,4	1,1			
28.	Талгар	91	20,9	СК	iP <sub>z</sub>	03-02-45,4						-5,5
29.	Курменты		21,3 <sup>X</sup>	СКМ	eP	03-02-53,2						
30.	Усть-Каме- ногорск		21,9 <sup>X</sup>	УСФ-3М	iP	03-02-59,6	0,8				0,017	4,52 MPV
31.	Ельцовка		24,2	СКМ	eP	03-03-22,5	0,8				0,02	4,75 MPV
32.	Чаган-Узун		25,9 <sup>X</sup>	СКМ	eP	03-03-39,2	1,2				0,02	4,67 MPV
33.	Бодайбо	49	39,7	СКМ	iP	03-05-34	0,6				0,026	+0,8
34.	Тикси	27	42,7	СКМ-3	iP	03-05-58	0,9				-0,005	<sup>MPV</sup> 4,45 -0,3
35.	Иультин	18	59,4	СКМ-3	eP	03-08-03,5	0,8				-0,012	<sup>MPV</sup> 5,08 +0,7
36.	Мирный	161	119,0	СКМ	ePKP e	03-15-00 18-15						

(01 July 1968)

- 24 -

I. УП. 1968г. (Севернее Каспийского моря - АЗГИР-2) - Зон. Казахстан  
(04-02)

JSC: H=04-02-00,9

$\varphi = 47^{\circ} 85' N$

$\lambda = 47^{\circ} 72' E$   $h = 0 \text{ км}$

MB = 5,5

ССН: 0=04-01-57,6

$\varphi = 48^{\circ} 005' N$

$\lambda = 47^{\circ} 850' E$   $h = 0 \text{ км}$

MPV

Севернее Каспийского моря

№ п/п	Станция	Az <sup>c</sup>	Δ <sup>c</sup>	Приб.	Фаза	Время	Am			T <sup>т</sup> сек	Mag	Res	
							X.S	EW	Z				
I	2	3	4	5	6	7	8	9	10	11	12	13	
1. Грозный		4,5 <sup>X</sup>	СК-З	P	P	04-03-13,0							
							E	i	41,5				
							E	(S <sub>2</sub> )	04-50,5				
							S <sub>max</sub>			5,6	5,2	2,0	1,0
				M <sub>L</sub>					4,2	2,4	3,4	8,0	4,7 ML
2. Махачкала	36	4,7	СК-З	P	P	04-03-15,6					6		
							e <sub>Lg</sub>		05-40				
							e <sub>L2</sub>		07-18,6	4,3	4,0	4,0	8,0
3. Пятигорск	221	5,2	СК	P	P	04-03-17,0						+0,9	
							e		24				
							e <sub>Pg</sub>		38				
4. Тбилиси	200	6,7	СК	P <sub>2</sub>	P <sub>2</sub>	04-03-37,0						-0,3	
							e <sub>NE</sub>		04-12				
							e <sub>S<sub>2</sub>NE</sub>		05-50				
							e <sub>Lg,NE</sub>		07-00				
5. Гори		6,7 <sup>X</sup>	СВЮМ-3	P	P	04-04-08							
6. Абастумани		6,7 <sup>X</sup>	ВСХ	P	P	04-03-44							
							ГСХ <sub>E</sub>		04-18				
7. Степанован		6,85 <sup>X</sup>	СХ	P	P	04-03-45,2							
							i		04-42,2				
							i		05-03,2				
							i <sub>S<sub>1</sub></sub>						
8. Анапа		7,4 <sup>X</sup>	СКМ	P	P	04-03-53,5							
							e		56				
9. Бакуриани	207	7,0	СКД-З	P	P	04-03-41,5							
							"	i	55,5				
							СК-З	e	57,5				
							"	i	04-14,5				
							"	i	18,1				
							" B	i	21,5				



I	2	3	4	5	6	7	8	9	10	11	12	13
9. Бакурчалли (црэг)					i	04-04-29,0		0,9		1,0		
					i	38,6	1,0			1,0		
					iS <sub>1</sub>	51		1,7		1,2		
					i	06-24,1						
					4x	08,2	0,6			9,0	3,49	
10. Ереван	7,6 <sup>x</sup>			СК	eP <sub>2</sub>	04-03-56,6						
11. Горис 187	8,6			СК	-iP	04-04-02			-0,2	1,0		+1,
					eP <sub>2</sub>	14						
					e	06-59						
					eS <sub>1</sub>	07-04						
					eS <sub>1</sub> (lg)	08-01						
12. Феодосия	8,6 <sup>x</sup>			СК	eP	04-04-11						
					eS <sub>1</sub>	05-53						
					eX	06-03						
					eX	19						
13. Алушта	9,3 <sup>x</sup>			СХ	-iP	04-04-20,1						
					P <sub>max</sub>	21			0,5	0,7		
					eX	36,4						
					eS <sub>1</sub>	05-52,7						
14. Ялта	9,6 <sup>x</sup>			СХ-2	eP	04-04-23,5						
					P <sub>max</sub>	24,0	0,2	0,3	-0,5	0,5		
15. Симферополь	257	10,0		СК-2	eP	04-04-20						+2,
				" В	eX	31						
					iX	55						
				" С	e(S)	05-37						
16. Москва	324	10,0		СВХ	eP	04-04-22						+0.
				"	i	24						
				СВХ <sub>W</sub>	e	34						
				"		46						
				"	e	05-12						
				СГК	e	06-12						
					e	07-06						
					e	10						
					eS <sub>2</sub>	31						
				СК	M <sub>L</sub>		0,4		0,6	8,0	3,88	M <sub>L</sub>
				СКД	M <sub>L</sub> g <sub>2</sub>	09,5	0,4		0,5	8,0		
				СВКД	M <sub>L</sub>				0,3	9,0		

I	2	3	4	5	6	7	8	9	10	11	12	13	
17. Кизил-Арват	I42	10,7	СК	$eP_2$ $eN$		04-04-33 05-01						-0,3	
18. Свердловск	36	11,8	СКМ-3	P		04-04-44 II	0,5	0,5	0,9	11,0	3,94	+3,3	
19. Ванновская	I39	12,5	ВЭГИК	$eP$		04-04-59						+1,4	
20. Ашхабад	I38	12,6	СК	$eP$ $M_{g_1}$		04-04-55,5 10-21,5			1,1	3,5		+3,4	
21. Кишенев	272	12,9	СК- $\Sigma$	$iP$ B CB B $\Delta M$		04-05-01,2 13 07-23 52 04-11,5					8,0	3,91	
22. Боровое	6I	15,1	СКМ	$-P$ "загр Pmax СКМ " СКД	$eS_E$ $eLg_N$ $M\Delta$	04-05-28,1 08-10,7 11-28,3 13	-0,002 0,11	-0,06 0,23	0,07 0,29	0,9 0,9		+3,1	
23. Пулково	325	15,6	СК	$iP$ $i$ $M_1$ $M_2$ $M_3$		04-05-36 44				0,3 0,5 0,45	18 8 6	4,04	+1,8
24. Мекгорье		15,65	СК- $\Sigma$	$eP$ $E\Sigma$ $eX$		04-05-44,1 52,9							
25. Чимкент		15,6	СК	$-iP_2$		04-05-45							
26. Львов	285	15,8	ВЭГ- $\Sigma$	$iP$ СК- $\Sigma$ " E " E	$iP$ $iX$ $eLg_1$ $eLg_2$	04-05-37 47 11-50 12-51							+2,7
27. Самарканд	114	16,1	СК	$eP$ $i$ $M_{L_2}$		04-05-43,0 50 14			0,7	0,4	12	3,78	+1,3
28. Ташкент	105	16,6	СК	$-eP$		04-05-48,5 54,5							12,3
29. Ужгород	282	17,0	СКМ-3	$eP$		04-05-53							+2,4

I	2	3	4	5	6	7	8	9	10	11	12	13
29. Ушггород (прос)					i	04-05-56			+0,067	I,0		
					i	06-08			0,14	10		
30. Душанбе			17,4 <sup>x</sup>	СК	-iP	04-06-07,3						
					edg	12-44,8						
31. Наманган			17,75 <sup>x</sup>	СК	-iP	04-06-11,8						
32. Гарм	II0		18,6	СКМ	eP	04-06-15						-4,3
33. Андижан			18,55 <sup>x</sup>	СК, СКМ	eP <sub>1</sub>	04-06-19						
				"	z	iP <sub>2</sub>			0,9	1,5	5,70 MPV	
				"	E	M <sub>Δ</sub>	16-16		1,0		11,0	
				"	E	M <sub>Δ</sub>			1,5		9,0	4,63 ML
34. Фрунзе			18,8 <sup>x</sup>	СКМ	eP	04-06-25						
				"	P <sub>max</sub>	29			0,1	1,0	5,0 MPV	
				СК	M <sub>Δ</sub>	18	0,6				9,0	4,23 ML
35. Красногорка			19,9 <sup>x</sup>	СК	iP <sub>z</sub>	04-06-38			-0,6	1,0	5,98 MP	
36. Или			19,95 <sup>x</sup>	СК	iP	04-06-38,2						
37. Хорог	II2		20,2	СК-z	P	04-06-36,0	+0,3	-0,4	-0,5	1,0	5,70 MPV	
38. Нарын			20,5 <sup>x</sup>	СКМ-3 <sub>E</sub>	iP	04-06-43,4						
39. Апатиты	344		20,9	СК	-iP	04-06-41,3						-6,6
					e <sub>δ</sub>	10-33						
40. Мургаб			20,9 <sup>x</sup>	СК	eP	04-06-47,6						
					e <sub>δ</sub>	10-43,7						
. Рыбачье			20,9 <sup>x</sup>	СКМ-z	iP	04-06-48						
42. Талгар	92		21,0	СК	+iP	04-06-44,0	-0,05	+0,09	+0,19	11,2	5,25 MPV	+1,
					LM	18	0,36	0,3	0,42	2,5		
43. Курменты			21,3 <sup>x</sup>	СК	iP	04-06-52	0,6	4,8	4,2			
44. Пржевальск			21,7 <sup>x</sup>	СК	iP	04-06-55,9						
45. Новосибирск			22,3 <sup>x</sup>	СК	P	04-07-01,3			0,3	1,8	5,42 MPV	
					δ	11-37,4						
46. Ельцовка	63		24,7	СКМ	iP	04-07-20,0			0,21	0,8	5,76 MPV	+1,5
47. Чаган-Узун			25,85 <sup>x</sup>	СКМ	-iP	04-07-37,4			0,1	1,0	5,45 MPV	
48. Монды	63		33,8	СКМ-3	IP	04-08-11,0						
49. Закаменск			35,1 <sup>x</sup>	СВКМ-3	+P	04-08-58,5			0,014	1,2	4,77 MPV	

I	2	3	4	5	6	7	8	9	10	11	12	13
50. Водайбо	50	39,6	СКМ-3	+iP	04-09-30,6				0,13	0,9	5,86 <sup>нрв</sup>	+I,3
51. Тикси		41,95 <sup>x</sup>	СКМ 3 СК, ВМЗ	+iP	04-09-56,0				+0,046	1,0	5,36 <sup>нрв</sup>	
52. Тупик	53	43,8	СКМ	iP	04-10-06,5	0,028			0,051	0,9	5,46 <sup>нрв</sup>	+2,8
53. Якутск	40	45,7	СКМ-3	+iP	04-10-20,0				0,092	0,8	5,86 <sup>нрв</sup>	II
54. Влади- восток	61	56,2	СКМ	iP	04-10-39,0							
				P <sub>max</sub>			0,5		0,04			
55. Ст. №2	55	52,7	СК	eP	04-11-14,0							+I,
56. Иультин	18	59,2	СКМ-3	iP	04-12-01,0	-0,043	+0,031	+0,09	0,9	5,9	5,9	+0,
57. Ю.Сахалинск	68,2 <sup>x</sup>		СКМ	iP	04-13-06,3				0,08	1,5	5,67 <sup>нрв</sup>	

22.12.1971г. (Северное Каспийского моря - АЗГИР) - Западный Казахстан  
(07-00)

YSC: H=06-59-56,6  
 $\varphi = 47,90N$   
 $\lambda = 48,07E$   $h = 0км$   
MB = 6,0

ECCH: 0=06-59-55,56  
 $\varphi = 48,09N$   
 $\lambda = 48,18E$   $h = 0км$   
MPV = 4,3

Северное Каспийского моря

I	2	3	4	5	6	7	8	9	10	11	12	13
№ шт	Станция	Az°	Δ°	Приб.	Фаза	Время	T сек	Am			Mag	Res
								NS	EW	Z		
1.	Махачкала		4,8 <sup>x</sup>	СК-ВЗ	iP	07-01-15,7						
					(iS)	57						
					Smax		12	11	18	3		
					LM		6	12	12	3,5	4,2ML	
2.	Пятигорск 223		5,4	СКМ-З	iP	07-01-18,0						+1,3
				"	Pmax	21	1,5			3,67		
				СК-З	Pmax	22	2,2			2,81		
				"NS	iS	46						
				SW	i	44	1,4			5,1		
				"Z	Smax	46	1,6			11,8	4,06NS	
				"EW	i	03-00						
					i max	02	1,6	6,7	10,3	12,1		
				"Z	i	04-08						
				"	max	26	10			4,04		
				"EW	i	05-30						
					i max	07-06,6	8	0,9	1,45	1,7		
3.	Анапа		6,5 <sup>x</sup>	СКМ-З	iP	07-01-39,0		+1,2	-3,5	+9		
				"	i	41,5						
				"E	iS	02-30,5						
				СД	ML	07-05,5	8,0	2,0	2,3	2	4,14ML	
4.	Тбилиси 201		6,8	СК-З	eP	07-01 39						+2,2
				"E	e	02-36						
				"NZ	ML	07-50	8,0	1,5		2,3	4,08ML	
5.	Зугдиди		6,8 <sup>x</sup>	СВХ	-iP	07-01-44						
					Pmax		0,5			3,0		
6.	Абастумани		7,0 <sup>x</sup>	СКМЗ-З	+iP	07-01-45,8	0,6			1,36		
				"	e	02-17,8	1,0	0,9				
				"	Smax	25						

22.12.71

- 95 -

I!	2	3	4	5	6	7	8	9	10	11	12	13
7. Шемаха		7,0 <sup>x</sup>	СХ-Л	iP	07-01-47							
			"	iX	02-02		1,8	1,5				
8. Бакуриа- ни	209	7,2	СХМЗ-З	+iP	07-01-42,5	1,4				1,0		+1,2
			СКД	P		0,8				0,04		
			"	i	02-19,5	1,0				0,88		
			"	iS	03-19,6	1,0				2,35		
			"	i	04-07,5	1,2				1,5		
			"~	h	05-3	14	4,14					
			"z	Mh	05.7	12				7,82	4,14 ML	
9. Кирова- бад		7,1 <sup>x</sup>	СХМ-З	iP	07-01-48							
			"c	iX	02-24							
			"	Pm		0,6	0,51	0,67	0,7			
			"	Sm		1,0	0,86	1,1	0,45			
10. Сочи	235	7,4	СК-В	iP	07-01-47							+0,9
			СХМ-В	iP	47,0	0,5	-1,2	-1,2	+3,0			
			"	Pmax	50	0,5			2,8			
			"E	i	02-26,5							
			СД-В	iP	01-47							
			"	Pmax	50	0,6			2,6	3,7		
			c-y	i	02-21							
			СД-Е	i	25							
			СД	iSmax	37	2,0	2,5	4,0			3,7	
			СХМ-Е	i	27							
			СД	ML	05-22	14	1,5	1,3	2,0	3,0 ML		
11. Баку		7,4 <sup>x</sup>	СК	iP	07-01-52							
				i	02-36							
				iz	39,0	4,0						
				iE	05-47							
				ML	08-47	10	5,2	2,0				
12. Ереван		7,8 <sup>x</sup>	СК-З	eP	07-01-58							
			"EЛ	i	04-24							
				ML	09	7,0	1,25	1,13			4,5	+3,5
13. Горис	189	8,7	СК	-iP	07-02-02							
14. Ленкорань		8,9 <sup>x</sup>	СХ	ePM	07-01-13,5	6,0	0,8	0,75	1,5	4,26		

	1	2	3	4	5	6	7	8	9	10	11	12	13
15. Алушта				9,6 <sup>x</sup>	СХ-2	-iP	07-02-22						
					"	Pmax	24	0,7			1,4		
					"	e	33						
					" <sub>2</sub> , c	e S	04-06						
— 16. Ст. AI				9,6									
17. Ялта				9,8 <sup>x</sup>	СХ-2	iP	07-02-24,8						
					, c, в	Pm	25,8	0,6	0,8	0,6	1,5		
					"c, в	S	04-13,6						
18. Москва	323	10,0			СВХ	eP	07-02-24						
					СВК	e	30						
					СДХ	e	36						
					СДК	e	46						
					СДЖ	e	52						
					СВК	e	03-18						
					СГК	e S	04-40						
					"	e	41						
					"	e Lg	05-16						
					"	e	40						
					"	e	48						
					СВКД	M L	07-07,3	14			2,3	4,15 ML	
19. Обнинск	318	10,1			БД-2	+iP	07-02-23,6	1,0		1,8			+0,9
					СХ-вз	i(S)	04-40						
					"св	i(S'S')	55						
					" <sub>2</sub>	L max	07-40	10	4,2	3,5	3,6	4,48 ML	
					ПО-2			10	8,0	3,6	3,0	4,41 ML	
					СКД-2			10	3,4	3,7	3,5	4,48 ML	
20. Симферополь	257	10,2			СХ-св	eP	07-02-24						+2,2
					c	e(S)	04-19						
						S m	20,0	0,9	0,73			4,0 MS	
						S M		1,0		0,42			
21. Кизил-Арват	143	10,6			СХ-2	eP <sub>I</sub>	07-02-30						+2,3
					"	e <sub>2</sub>	34						
					" N	e <sub>3</sub>	38						
					"	i	53						
					"	e <sub>4</sub>	03-35						

	I	2	3	4	5	6	7	8	9	10	11	12	13
— 22. Свердловск	36	11,6	СКМЗ	iP	07-02-43,0								+1,9
			"	Pmax	46,0		1,0				1,2		
			СГ	LM	07-09		1,2	1,5	4				
23. Ашхабад		12,0 <sup>x</sup>	СКМЗ-2	eP	07-02-55								
24. Ванновская	140	12,5	СКМ-3	eP	07-02-52								+4,9
25. Черновицы		14,3 <sup>x</sup>	СКД, СХ	iP	07-03-26,0								
			СХ										
			СКД				1,2	1,8					
			СХ				1,0		6,4				
			СКД, СХ	i	03-34								
			СКД	SP	59								
			СХ	SP	04-01								
			СКД	i	15								
			"	δ	05-29				+2,0				
			СХ										
			СКД				1,4	2,5					
			СХ				1,2		2,7				
			СХ	SS	06-05								
26. Киченев	272	13,1	СК-В	eP	07-03-03,0		1,5				1,1		+2,9
			"	i	15								
			"с	i	08-18								
			"в	i	27								
			СД-2	ML	07-09,3		1,3				1,7	4,25	
← 27. Боровое	62	14,9	СКМ	+iP	07-03-26,2		0,9	+0,01	0,05	+0,04			+0,4
				Pmax	30		0,9	0,20	0,31	0,706			
				eS	05-28,5								
				e	06-06,2								
28. Пулковое		15,2 <sup>x</sup>	СГ	iP	07-03-38								
				i max	40							1,5	
				i	45								
				i max	47		1,8					2,9	
				e(S)	06-32								
			СГ	LM	11,5		8,0	1,1	1,1			4,5 ML	
			СКМ	LM	11,5		8,0	1,1	1,1			4,5 ML	



I	!	2	!	3	!	4	!	5	!	6	!	7	!	8	!	9	!	10	!	11	!	12	!	13
29.	Львов			15,3 <sup>x</sup>		СКМ-з		iP		07-03-39														
						"		e		47														
						"		i		06-21														
						"		e		29														
						"		ML		07-11,6														
30.	Самар-канд	115		15,9		СК		eP		07-03-40														+2,6
31.	Межгорье			15,9 <sup>x</sup>		СКМЗ-з		eP		07-03-46,7														
						з		e		53,7														
						~		i		06-55,7														
32.	Ужгород	281		17,2		СКМ		iP		07-03-52,7		1,0								+0,012				+3,6
						"		iX		58,2														
						"		Pmax		04-00		1,1	0,03	0,07	0,15									
						"		i		03														
						"		i max		15		1,1	0,07	0,16	0,34									
						"		i		07-36														
						"		i		09-41														
33.	Душанбе	114		17,7		СК		+iP		07-04-05										1,5				4,6
								eP		07-30		9,0												
34.	Гарм			18,0 <sup>x</sup>		СКМЗ-з		+iP		07-04-13,1		0,9												0,47
						"		i		23,3		1,0												0,625
						E		i		05-42		1,3												0,225
						СК-з		iP		07-04-13,1		1,8												0,8
						~		i		05-59		3,2												0,4
						~		iL		07-44,3		6,4												0,88 4,51 ML
35.	Фергана			18,1 <sup>x</sup>		СКМЗ, СК-з		+eP		07-04-14,3														
						СК-з		MP		15,3		1,0												0,6
						"E		e		09-40,3		8			1,5									
						"		ML		07-13		11			4,5									5,5 5,09 ML
36.	Андижан			18,3 <sup>x</sup>		СКМЗ-з		-iP		07-04-17,0		0,6												0,1
								MP		20		1												0,9
						СК-з		MP		20		1,5												2,1
						"E		eS		08-31														
						" ~		e Lg		10-07														
						" "		M		07-15		8,0			4,0									3,0 4,98 ML

	I	2	3	4	5	6	7	8	9	10	11	12	13
37. Фрунзе	96	19,2	СКМ	iP	07-04-23,5								-0,1
				i	27,5								
			"	MP	29,0	0,8			0,22				
			СКД	ML	14-20	10		1,5				4,58	ML
38. Хорог	113	20,1	СК-з	P	07-04-33,7								+0,6
39. Талгар	92	20,8	СКМЗ	iP	07-04-42,7	1,1	0,11	0,17	+0,62				+1,0
40. Мургаб		20,8 <sup>X</sup>	СК	iP	07-04-45,1	0,6	-0,5	+0,8	+0,8				
				eS	08-41,1								
41. Апатиты	344	20,9	СКМ, СХ	iP	07-04-42,6								+0,5
			"	i	43,3								
			СКМ-з	Pmax	45,0	1,0				1,3			
			СХ	iS	08-27,8								
42. Прже- вальск	93	21,9	СКМ	eP	07-04-54								+1,9
43. Новосибирск		22,6 <sup>X</sup>	СКМ-з	+iP	07-05-03,0	1,4			0,05				
				Pmax	05,5								
				i	06-01								
44. Ельцов- ка	63	24,5	СКМЗ-з	iP	07-05-19,1	1,0	+0,21	-0,07	+0,65				+1,9
45. Усть- Элегест		29,2 <sup>X</sup>	СКМЗ-з	iP	07-06-05,2								
				Pmax	07,0	1,2			0,2				
46. Хейс	3	32,8	СК	+iP	07-06-36								+3,1
				iPmax	37	1,0			0,12				
47. Монды	63	33,6	СКМ	+P	07-06-43								+3,3
48. Иркутск	61	35,4	СКМЗ	+eP	07 06-57								+1,7
				Pm	58								
				eL2	12,5								
49. Зака- менск	65	35,4	СКМ	iP	07-06-58,4	1,2	+0,026	+0,034	+0,021	+3,			
			"	PmI	07-00	1,2		0,107					
			СКД	L	22-36,6							5,0	ML
			"	LM	23-36,2	16			0,673				
50. Бодайбо	50	39,4	СКМ-3	+iP	07-07-29,5								+0,9
				Pmax	30,0	1,0	0,58	0,47	0,65				

I	2	3	4	5	6	7	8	9	10	11	12	13
51. Тикси	27	42,4	СК, СКМ	iP	07-07-56							+2,4
			СКМ-2	P <sub>max</sub>		58,0	1,1			0,17		
			СК, СКМ	i		09-39						
			ГК	e		46						
			СГК	MZ		31,8	14		0,3			
							10	0,1				
52. Тупик		43,4 <sup>X</sup>	СКМ	+iP	07-08-05,7		0,8	0,05	0,06	0,25		
53. Якутск		45,1 <sup>X</sup>	СК	iP	07-08-20,0		1,0			+0,15		
				P <sub>max</sub>		22	1,0			0,32		
				i		10-14						
5. Ст №2 (Кульдур)		52,1 <sup>^</sup>	УСФ	i	07-10-12							
55. Сеймчан		54,0 <sup>X</sup>	СКМ	P	07-09-27,6							
			"	P <sub>max</sub>		28,6	1,1			0,26		
			"H	MP			1,2	0,06				
							1,3		0,09			
56. Владивос- ток	61	56,0	СКМ, СК	eP	07-09-39,5		1,0			0,12		+1,;
			СК				5,0			0,3		
57. Иультин	18	59,1	СКМ-2	P	07-10-01,1			-1,2	+0,6	+3,5		+0,;
			"2	P <sub>max</sub>		02,0	1,0	0,069	0,05	0,23		
			"E	i		06						
			"N	i		16						
58. Ю. Саха- линск	52	59,7	СКМ3-2	eP	07-10-06,4							+2,;
			"2	P <sub>max</sub>		08						
			"E	e		11						
			"2	e		20,4						
			"2	e		53						
			"2	e		12-18,8						
59. Мирный		119 <sup>K</sup>	СКМ	ePKP	07-18-47							
60. Новола- заревская		127 <sup>K</sup>	СКМ	iPKP	07-19-03		1,0			0,02		
				i		09						
				e		23						

(Зн. 1/2)  
 29 мая 1976г. (АЗГР) Севернее Каспийского моря  
 (05-00) БУСН-04-59-56

УСC : H=04-59-58,0

$\varphi = 47^{\circ} 81' N$

$\lambda = 48^{\circ} 10' E$   $l_1 = 0 \text{ км}$

MB=5,9

$\varphi = 48,1 N$   $\lambda = 48,2 E$   $l_1 = 0 \text{ км}$

$M_p^{сдл} = 5,9 (11с.)$   $M = 4,5 (3с.)$

Севернее Каспийского моря

№	Станция	$\Delta^{\circ}$ по Об- визу	Прис.	Фаза	Время	T сек	$\Delta \mu$			$M_{сдл}$	
							NS	E W	Z	MP	ML
1	2	3	4	5	6	7	8	9	10	11	12
1. Махочкала		5.1	СКМ	$+iP$	05-01-16.3						
				$R_{max}$	22,0	0.7	11.0	5.5	5.5		
				$iS$	45.0						
				$S_{max}$	48.0	0.6	17.0	8.7	4.6		
2. Пятигорск		5.4	СК	$eP$	05-01-19						
				$R_{max}$	20.0	2.0			4.5		
				$eS$	01-43						
				$S_{max}$	47	2.0	6.5				
				$i$	51						
3. Тбилиси		6.8	СКМ	$P$	05-01-39						
				$R_{max}$	01-40						
4. Грозный		4.8 <sup>X</sup>	СКМ	$-iP$	05-01-15.8		+0.7	+0.2	-3.4		
				$R_{max}$	17.0				4.3		
				$iS$	42						
				$S_{max}$	44	1.0	7.1	3.0			
				$ML$	02-07		7.8	4.4		4.4	
5. Бакуриани		7.2	СКМ	$iP$	05-01-49.6						
				$R_{max}$	45.6	1.0			0.68		
				$ML$	05-05,5	12		4.6			
			"	$ML$	05,8	11	3.2				
6. Сочи		7.4	СКМ	$iP$	05-01-49					1.6	
				$i(S)$	02-25			2.5			

1	2	3	4	5	6	7	8	9	10	11	12
7.	Кировабад	7.6	СКМ	-P	05-01-47						
				P <sub>max</sub>		1.0			0.50		
					02-14						
					02-54						
8.	Баку	7.8	СК	P	05-01-54						
					01-55						
					02-00						
					02-12						
9.	Ленинаган	7.9	СКД	eP	05-01-57						
10.	Ереван	8.3	СК	P	05-01-59.0						
			"		02-40						
			"		05-01						
11.	Горис	8.7	ВЭГ	+P	05-02-04						
					02-17.6						
					26.2						
				e	33.2						
				eS	46.2	1.2	1.05				
12.	Симферополь	10.2	СК	-eP	05-02-25.0						
			СХ	e	26						
			СХ	P <sub>max</sub>	27.0	0.4			0.5		
			СК	e	43						
			"СХ	eS	04-20						
			СК	S <sub>max</sub>	04-	1.0	1.0	0.6			
			СХ	S <sub>max</sub>	04-23	1.0	0.4	0.6			
			СК	e	04-43						
			"	eL	06,3						
			СКД	Mh	08-09	14			1.0		3.8
			СК	Mh	08-34	10	0.3	0.7			4.0
13.	Обнинск	10.1	ЭК-1	eP	05-02-28						
			"	P <sub>max</sub>	32.0	1.0					
			(ГИСС)	eS	04-24						
			ГИП-1								
			ЧИСС	M	07-50						
14.	Арги	10.5	СКМ-3	eP	05-02-30						
			"	e	02-50						
			СКД	eS	04-26						
			СКД	M	08-11	12	1.0	1.5	2.0		4.1

I	2	3	4	5	6	7	8	9	10	11	12		
15.	Кзыл-Арват	10.8	СК	eP	05-02-32								
16.	Ашхабад	12.6	СКМ-3	eP	05-02-55.2								
				i	02-57,2								
			СК	e(S)	05-05.0	1.0		1.0					
				M	13-39.1	9.0			1.2				
17.	Киченев	13.1	СК	-P	05-03-04								
				P <sub>max</sub>	05.0	1.0		0.9	1.1				
				i	18								
				i	03-36								
				i	04-16								
				i	05-18								
				LS	38								
				iS <sub>max</sub>	44								
			CD-1	M	09-42	11			1.9	4.5			
			СК	M	09-46	10		1.5	1.8				
18.	Минск	13.7*	СК	eP	05-03-18								
				CD-1	eL	09,4							
				"	M	10-59.0	12		2.0				
19.	Пулково	15.7	ВЭГ	+P	05-03-38.5								
				P <sub>max</sub>	03-48.0	1.3			1.2				
				e	47								
				СГ	e(S)	06-27							
				CD-1	M	11-16	9.0	0.9				4.3	
				СГ	M	11-18	10	0.8	0.5	1.5		4.2	
					10		0.6	1.2					
16. Садарманг	15.9	СК	iP	05-03-40.5									
20.	Львов	15.9	СКД	P	05-03-40								
						49							
						04-18							
						e	05-50						
						e	07-06						
						M	11-47.0	9.0			1.0		
21.	Ташкент	16.4	СКМ	-P	05-03-48								
				"	P <sub>max</sub>	49,5	1.0		1.1	6.0			
				СК	P <sub>max</sub>	49,5	1.5		1.2	5.9			
				СК	i	05-53.5							
				"	i <sub>max</sub>	55.5	2.0		3.0				
				СКМ	i <sub>max</sub>		1.2		2.8				
				СК	M	05-12-34	10	1.5	3.0	1.7	5.4		

1	2	3	4	5	6	7	8	9	10	11	12
22.	Ужгород	17.2	СКОМ	-P	05-03-55.5						
				P <sub>max</sub>	56.0	0.9			0.350		
				+L	03-58.5						
				P <sub>max</sub>	04-03.0	1.0			0.5		
				i	04-08						
				i	15						
				i	30						
				i	37						
				i	44						
				i	55						
				•	05-53						
23.	Душанбе	17.7	СХ	+P	05-04-06						
24.	Андижан	18.7	СКОМ	eP	05-04-18						
			СХ	P <sub>max</sub>	22.0	1.4			1.6		
			СКОМ	P <sub>max</sub>	23.2	1.3			1.43		
			СХ	i	05-32	1.2		1.0			
			"	e	09-07	4.0	1.0				
			"	ML	14-06	10	2.0	2.2			
			"	ML	16-30	9.0	2.2	3.5	2.2		
25.	Куляб	18.7	СХ	+P	05-04-18.4						
				P <sub>max</sub>	21.4	1.0			1.3	6.8	
				eS	10-45.4						
26.	Фрунзе	19.2	СКОМ	eP	05-04-24						
				i	04-28						
				P <sub>max</sub>	30.0	1.0			0.26	6.4	
				L	42						
				i	48						
			СХД	e	05-08						
			СХД	e	05-22						
				e	06-28						
				e	07-08						
				e	07-50						
				e	08-10						
				e	08-40						
				e	09-14						
				e	10-14						
				e(S)	11-18						
				ML	17-00	8.0	1.8				5.5
					71						

1	2	3	4	5	6	7	8	9	10	11	12
27.	Хорог	20.1	СК	P	05-04-35						
28.	Нарын	20.7	СКМ-3	+eP	05-04-42						
				e	05-31.5						
29.	Талгар	20.8	СКМ	+iP	05-04-43.0	1.2	0.11	0.2	0.26	5.5	
			"	iP <sub>max</sub>	45	1.1			0.52	5.8	
			"	iP <sub>max</sub> <sub>a</sub>	48	1.2			0.77	6.0	
			СК	eP	43	2.2			1.0	5.8	
			"	S	08-49.0	8.0			3.0		4.7
			"	iL <sub>e</sub>	13-39	17	0.5		0.7		
			"	M <sub>L</sub>	05-15	11	0.5	0.1	0.8		4.5
30.	Пржевальск	21.9	СКМ	+iP	05-04-55						
			"	P <sub>max</sub>	59.0	1.0			0.20	5.5	
31.	Новосибирск	22.7 <sup>X</sup>	СКМ-3	+iP	05-05-03.5						
			"	P <sub>max</sub>	04.7	1.4			0.454	5.7	
			"	iS	09-10.3						
			СД-1	e	10-28.6						
			"	e	58-45.4						
32.	Норильск	28.7 <sup>X</sup>	СКМ	+iP	05-06-01						
				P <sub>max</sub>	02.0	1.0			0.104		
				i	06-21.5						
				i	06-38						
				i	07-19.5						
				i	07-56.5						
				e	11-58						
				M	20.5	16.0		0.3			
33.	Усть-Элегест	29.3 <sup>X</sup>	СКМ-3	-iP	05-06-05.5	1.0	+0.04	-0.03	-0.09	5.8	
				iS	13-10.5						
34.	Иркутск	35.4	СКМ	+P	05-06-58.2						
			"	P <sub>max</sub>	59.2	1.0	0.066	0.154	0.185	6.0	
			"	h <sub>g</sub>	10.5						
			СКМ	Max	10-45.7	1.9		0.142			
			СКД	eL <sub>e</sub>	11.9						
			СКД	M <sub>L</sub>	12-15.0	9.0	0.38	0.34	0.56		4.6
				eL	22.5						



1	2	3	4	5	6	7	8	9	10	11	12
35. Завьяненск		35.4	СНМ	+P	05-07-00.4	0.8	-0.4	+1.0	+1.7		
			"	P <sub>max</sub>	01.4	1.0		0.07			
			СНМ	e	08-12.4						
			"	e	10-05.2						
			"	L	11-12.4						
36. Тихая	42.3 <sup>2</sup>		СНМ	+P	05-07-57.0	8.0	0.49	0.62	0.69		4.2
			"	P <sub>max</sub>	58.0	1.0		0.09	5.6		
			СН	P <sub>max</sub>	58.0	1.5		0.2	5.8		
			СНМ-3	i	08-02.5						
			"	i	30						
			"	i	35.5						
			"	i	40.5						
			"	i	51.5						
			"	e	09-39.5						
			"	e(S)	14-00						
			"	e	19-31						
			"	e	59						
"	L	30-20.0			14		0.2			4.2	
					12		0.2				
38. Гупши		43.5	СНМ	-P	05-08-05.9	2.0	0.072	0.113	0.220		
39. Млутца		45.5	СНМ	eP	05-08-13.6						
40. Сельчан		54.3	СНМ	+P	05-09-28.5						
			"	P <sub>max</sub>	29.8	1.0		0.17	6.1		
			"	i	10-32.1						
41. Монды		55.3	СНМ	+P	05-09-36						
			"	P <sub>max</sub>		1.0		0.07	5.7		
			"	e	09-58						
			"	e	14-00						
42. Магадан	35.1 <sup>2</sup>		СНМ	L	37-16	18			0.5		4.6
			СНМ	+P	05-09-36						
			"	P <sub>max</sub>	38.0	1.0		0.07			
			"	e	58.5						
			СНМ	e	14-00						
			"	e	15-16						
"	L	37-16			18			0.5			

1	2	3	4	5	6	7	8	9	10	11	12
43.	Нурьган	59.1	СКМ	+P P <sub>max</sub>	05-10-02.5						
			"		10-13	1.0	0.054	0.046	0.17	6.1	
					25						
					54						
					11-14						
				•	11-48						
				•	12-10						
					31						
			СКД	•S	18-08						
			"	•	21-24						
			"	•	26-08						
44.	С.Сакалинск	59.7	СКМ	+P P <sub>max</sub>	05-10-07.3						
					08.1	1.0			0.19	6.2	
				•	10-16.1						
				•	10-29						
				•	10-49.7						
				•	12-19						
45.	Новоза- ревская	121.4	СКМ	ePK, KP	05-18-51.5						
46.	Ст. № I (Михнево)	9.4 <sup>X</sup>	СКМ	eP	05-02-16						
47.	Анапа	8.1 <sup>X</sup>	СКД	P eS ML	05-01-58 02-II 07.8	0.5	1.0	1.0			
48.	Ельцовка	24.5 <sup>X</sup>	СКМ	IP P <sub>max</sub>	05-05-19.8 29.3	9.0	1.5	1.3	2.0		4.0
49.	Аптиты	15.8 <sup>X</sup>	СКМ СКД	P eS ML	05-03-44 07-33 05-13.2	1.2	0.100	0.400	0.376		
50.	Шеки	6.7 <sup>X</sup>	СКМ	+P L	05-01-40.5	12.0	1.4	1.0	1.8		ML4,5 ML4,7
52.	Степанован	7.3 <sup>X</sup>	СКМ	IP	05-01-48.2						
53.	Наманган	18.8 <sup>X</sup>	СК	-IP	05-04-21						
51.	Алушта	9.8 <sup>X</sup>	СХ	IP P <sub>max</sub> СК	05-02-23.6 24.6 05-08.4	0.6 0.6 16.0			0.16 0.78 0.7		

30 <sup>sept.</sup> сентября 1977 (АЗГР) Севернее Каспийского моря -  
(07-00) Западный Казахстан

JSC: H=06-59-55,9

$\varphi = 47^{\circ} 85' N$

$\lambda = 48^{\circ} 13' E$   $h = 0 \text{ км}$

MB=5,0

ВСС: H=0-06-59-59,8

$\varphi = 47,92' N$

$\lambda = 48,31' E$

$R = 0$

MPV (A) = 5,0 (9 ст)

Севернее Каспийского моря

№	Станция	$\Delta^{\circ}$	Az $^{\circ}$	Крб	Фаза	Время	T сек	AN			Mag	Res
								NS	EW	Z		
1	2	3	4	5	6	7	8	9	10	11	12	13
1.	Махочкала	5.04	186	СКМЗ	+P	07-01-15.0						
					eS	02-24						
					ML	04-10.0	14.0	1.0	0.9	0.6	3.1 3.5	
2.	Пятигорск	5.36	224	СКМ	eP	07-01-18.8						
					IS	01-43.6						
3.	Белый уголь	5.1 <sup>X</sup>		ВБГ	+P <sub>Z</sub>	07-01-18.9	0.2	0.02	0.04	0.10		
					Rmax	20.5	0.2	0.58	0.53	0.68		
					+L	01-24.2						
					+IS	01-39.1						
					Smax	01-46.5	1.0	0.5	0.18	0.10		
4.	Зугдиди	6.0 <sup>X</sup>		СК-Σ	eP	07-01-32						
.	Тбилиси	6.3 <sup>X</sup>		СК	eP	07-01-37						
					ePPP	02-09						
					eS	03-15						
					eSS	03-59						
6.	Абастумани	7.0 <sup>X</sup>		СКЛ-Σ	eP	07-01-45.6						
					B	02-23						
7.	Соли	7.42	236	СКМ	+P	07-01-46.1						
						01-58.9						
						04-21.8						
					e	04-59.0						
8.	Степанован	7.0 <sup>X</sup>		СКМ-Σ	P	07-01-47						
						02-18.4						
						30.6						
						44						
9.	Ленинакан	7.87	205	СКД	eP	07-02-23						

30.IX.77

1	2	3	4	5	6	7	8	9	10	11	12	13
10. Горно	8.57	190	ВЭГ	-P	07-02-02.0	0.8				-0.2		
			"	∩	02-15							
			СК	e	04-06							
			"	eS	04-40							
11. Москва	10.19	323	СК	eP	07-02-12							
			"	e	02-35							
			"	e	03-05							
12. Ст. В 1 (Мухомово)	9.0 <sup>X</sup>		СКМ	-P	07-02-15.2	0.4				0.315		
13. Адушта	9.6 <sup>X</sup>		СК-	+eP	07-02-22							
			"	P <sub>max</sub>	26.0	0.5				0.07		
			"	e	02-38							
			Св	e	02-55							
			?	e(S)	04-09							
14. Обвинск	10.21	318	СКМ	+P	07-02-26.1							
			"	P <sub>max</sub>	30.0	0.8				0.15		
			"	e	03-05							
			"	e	04-20							
			"	e	05-12							
			СКД	eL	08-02-30.0	24.0						
15. Арты	10.55	32	СКМ	eP	07-02-28.7							
			"	e	02-31.5							
			"	eS	04-27.7							
			СКД	e	07,6							
16. Свердловск	11.63	35	СКМ-3	+P	07-02-43.0							
			"	P <sub>max</sub>	45.3	1.0				0.12	5.5	
			"	e	02-58							
			"	e	03-05							
			"	e	03-20							
			"	e	04-52							
17. Ашхабад-2	12-48	140	СК	P	07-02-47.3							
			СКМ	-i <sub>max</sub>	03-06.6	0.6				0.09		
			"	+i <sub>Smax</sub>	05-14.3	1.3	0.19					
			СК	+∩	06-22.6							

30.1X.77

I	2	3	4	5	6	7	8	9	10	11	12	13
18. Боровое	14.86	61		СКМ	Р	07-03-26.4	0.7			0.015	4.8	
19. Самарканд	15.1 <sup>X</sup>			СК	еР	07-03-36.0						
					е	40.6						
					-i	46.8	0.6			0.4		
					е(S)	06-49						
					ML	12,59	6.0		0.4		4.2 ML	
							8.0			0.4		
20. Ташкент-2	16.34	105		СКМЗ	еР	07-03-44.0						
				"	Р <sub>max</sub>	53.5	1.2	0.1	0.2	0.3	5.9	
				СК	е	09-13.5						
					е	10-09.5						
					е	12-00						
					еL	08-17-50						
21. Узгород	17.28	281		СКК	+Р	07-03-58.5						
					Р <sub>max</sub>	04-03.5	0.8			0.015	4.8	
					i	04-08						
					i	04-13						
					i PP	06-40						
					i	07-10						
					е	08-10						
					е	12-25						
					е	20-20						
					ML		18.0			1.1	5.6 ML	
22. Душанбе	17.59	114		СК	еР	07-04-04						
23. Гарм	18.28	11		СКМ	Р	07-04-12.2						
					Р <sub>max</sub>	12.2	1.2			0.08	5.3	
24. Фергана	18.0 <sup>X</sup>			СКМ	еР	07-04-13.7						
				СК	L	12-						
25. Рудаб	18.59	114		СК	еР	07-04-16						
					еS	10-47						
26. Анджеван	18.62	103		СКМ	еР	07-04-16.4						
				"	Р <sub>max</sub>	19.8	1.0			0.15	5.6	
				СК	Р <sub>max</sub>	20.0	1.0			0.2	5.7	
					L	05-07						
					е	08-16						
					еL	13-30						
					ML	14-00	13		0.7	0.8		

30.1X.77

1	2	3	4	5	6	7	8	9	10	11	12	13
27.	Бруно	18.7 <sup>x</sup>		СМЗ	еР	07-04-22						
					е	04-26						
					еS	04-50						
					См. еР	04-54.0	0.8	0.04				
28.	Хорог	19.97	113	СМ	еР	07-04-33						
29.	Галгар	20.73	92	СМ	+iP	07-04-41.0						
					R <sub>max</sub>		1.2	0.009	0.009	0.033		
					R <sub>max1</sub>	43.0	1.5	0.031	0.023	0.076	4.9	
					R <sub>max2</sub>	42.0	1.5	0.053	0.029	0.083	5.0	
30.	Новосибирск	22.4 <sup>x</sup>		СМЗ	+iP	07-05-01.3						
					R <sub>max</sub>	08.3	1.4	0.02	0.026	0.056		
					e	08-52.3						
					i	09-08						
				СД-1	M <sub>к</sub>	16-35.0	12.0			0.6		
31.	Апатиты	21.06	344	СМ	+iP	07-05-46.0						
					R <sub>max</sub>	48.0	1.0	0.13	0.1	0.16	5.4	
				СД-1	L	08-01.5	28			1.5		
32.	Ельцовка	24.47	63	СМ	+iP	07-05-18.6						
					R <sub>max</sub>	18.6	1.0			0.100	5.3	
					eS	09-43						
33.	Норильск	28.93	27	СМ	еР	07-05-59.						
					R <sub>max</sub>		1.4			0.017	4.7	
					e	06-35						
					e	06-38						
					e	06-47						
					e	06-59						
					e	07-17						
					e	07-28						
					e	08-03.0	2.1			0.5		

30.1X.77

1	2	3	4	5	6	7	8	9	10	11	12	13
34.	Монди	33.60	63	СНМ	+eP	07-06-41.8						
					P <sub>max</sub>	4279	1.2			0.035	5.2	
35.	Закаменск	35.35	65	СНМ	+P	07-06-57.1						
					P <sub>max</sub>		1.3			0.013	4.7	
					eP	09-25.4						
					P <sub>max</sub>		1.2			0.01	4.4	
36.	Бодайбо	39.36	50	СНМ	+P	07-07-29.0						
					P <sub>max</sub>	29.0	0.6	0.02	0.033	0.08	5.8	
37.	Мульгин	59.23	18	СНМ	eP	07-10-00.5						
					e	10-39						

17 Октября 1970 (А.М.Р.) Севернее, Каспийского моря  
 Западный Казахстан  
 КСХ: 0-04-59-55,0 12,0<sup>5</sup>

$\gamma_{SC} : H=04-59-56,6$   
 $\varphi = 47^{\circ}81'N$   
 $\lambda = 48^{\circ}09' E$   $h=0$  км  
MB=5,8

$\varphi = 47^{\circ}94' N$   
 $\lambda = 48^{\circ}04' E$   $h=0$  км  
 MPV(A)  $\varphi = 6,0$  (по 22 ст)  
 MPV(B)  $\varphi = 6,5$  (по 22 ст)  
 MLH (B)  $\varphi = 4,7$  (по 8 ст)  
 MLV(B)  $\varphi = 4,5$  (по 8 ст)  
 MLC (C)  $\varphi = 4,6$  (по 4 ст)

Севернее Каспийского  
 моря.

№ п/п	Станция	Вр.об.	Вид	Время	T <sub>ср</sub>	Ам			Mag	Δ°	Rz°	R <sub>ср</sub>		
						NS	EW	Z						
1	2	3	4	5	6	7	8	9	10	11	12	13		
1.	Махачкала	СК	+iP	05-01-15.0	0.4	-3.0	+0.8				5.00	184	1.8	
			R <sub>max</sub>	24.0	0.6	22.0	9.2	9.0	7.4					
			MLH	02-12	1.0	35.0	12.0	5.0						-0.7
			LM	06-00	8.0	4.9	4.3	4.5	4.3	MLH				
									4.2	MLV				
2.	Пятигорск	СК	iP	05-01-17.5							5.23	223	1.1	
			R <sub>max</sub>	18.5	2.0			4.0						
			i	01-30										
			i	01-44										
			S <sub>max</sub>	01-46.0	2.0		6.0							
			i	04-24	9.0		5.0							
3.	Зугдиди	СК	+iP	05-01-43							6.8 <sup>x</sup>			
			R <sub>max</sub>		1.0			5.0						
4.	Абастумани	СКМ	+iP	05-01-45.6							7.0 <sup>x</sup>			
			US	02-19.0	0.5	0.67								
5.	Степанован	СММЗ СКД В	+iP	05-01-46.8								7.0 <sup>x</sup>		
			i	02-37.2										
			i	03-45.8										
			i	03-40.6										
			i	04-07.8										
			i	04-15.4										
			i	05-00.2										
6.	Баку	СК	+iP	05-01-53							7.5 <sup>x</sup>			
			i	01-54	0.5			1.5						
			e	02-36	2.0		9.0							



1	2	3	4	5	6	7	8	9	10	11	12	13
7. Ленинград	СНД	+iP	05-01-53.0	2.0				1.0		7.78	204	0.7
		•	02-84									
	ВСТ	•	02-49.0	2.0	8.0							
	СНД	ML	05-00	3.0	8.0	6.0			5.0			
8. Брест	СК	+iP	05-01-58.0							8.16	193	0.3
		i	02-10.5									
		i	04-85.5									
		i	05-38									
9. Горьк	СК	+iP	05-02-02.6							8.52	188	-0.1
	ВСТ	Pmax	03-8	0.6				0.66	6.3			
	СК	•	04-83.8									
	СК	•	05-05									
10. Ст. В 1 (Михнево)	СНМ	+iP	05-02-15.7	0.6				1.080		9.1 <sup>X</sup>		
11. Сандаро- поль	СНД	•P	05-02-21							10.06	257	-3.0
		•S	04-17									-2.0
	СНД	LM	07-21.0	17	0.5	0.6	0.6		3.6 ML 3.5 ML			
	СН-1	LM	07-25.0	16	0.3	0.6	0.7		3.6 ML			
12. Алушта	СХ	•P	05-02-21.7							9.5 <sup>X</sup>		
		-iP	02-22.7	1.0				0.35				
		Pmax	23.2	0.6				0.57				
		i	02-38.6									
		•	03-38									
		•(S)	04-11									
	СК	LM	05-07.7	9.0				0.5	3.7 ML			
13. Москва	СХ	•P	05-02-23.0							10.11	324	-1.6
		•	02-81									
		•	01-46									
		•	03-00									
		•	03-16									
		•	04-17									-3.1
		•	04-46									
		P	05-01									3.7
14. Ялта	СХ	-P	05-02-24.2	0.7	0.3	0.3	0.8			9.7 <sup>X</sup>		
		•	04-05									
		•	04-14									
15. Обнинск	СНМЗ	+P	05-02-25.0							10.12	319	0.3
		Pmax	28.0	1.0	1.1	1.1	1.3	6.5				
	СК	•	03-22									

I	2	3	4	5	6	7	8	9	10	11	12	13
продолж.												
15.Обнинск	СК	•	05-04-06.0									
	СК	LM	07-50.0	8.0	1.5	2.5	2.1	4.5				
								4.3				
	НО	LM	50.0	10	1.3	2.0	1.9	5.5				
								5.6				
16.Арм	СНС	•P	05-02-28.8							10.66	38	-3.2
	Рmax		31.9	0.9			0.35	5.9				
	↓		03-33.5									
	•		03-37.3									
	•		03-39.8									
	LM		03-05.0	11	2.0	2.5	2.5	4.4				
	LM			12			2.0	4.2				
								4.1				
	LM			7.0	1.0		1.3	4.3				
	LM			8.0		1.0		4.1				
17.Клен-Арбат	СК	•P	05-02-24.0							10.73	143	-9.1
	•		03-32									
	•		03-33									
	•		03-38.5									
18.Свердловск	СНС	•P	05-02-44.2							11.75	36	-2.7
	Рmax		47.0	1.0			1.0	6.4				
	•		03-18									
	•		03-32									
	СНД	•	04-50									
	СНД	•	06-04									
	•	LM	03-00.0	12	1.1	1.5	3.5	4.2				
								4.5				
19.Ашгабад	СК	P	05-02-53.5							12.57	139	-4.6
	↓		03-14									
	•		03-34									
	•		03-55									
	•		04-14									
	↓		05-40.0	1.0			0.6					
	↓		06-26									
	↓		06-31.0	5.0			1.0					
	↓		07-17	2.0			1.2					
	•		07-27.2	1.0			0.7					
	•		08-02	3.0			3.5					
	LM		09-55	8.0			2.5	4.6				
	LM		09-56	11			3.0	4.5				
	LM		10-04	9.0			3.0	4.6				

	1	2	3	4	5	6	7	8	9	10	11	12	13
20. Кисловодск	СКД	-P	05-08-02.0								13.04	272	-2.3
		Рmax	03.0	0.8				0.7	6.4				
	СКД	v	03-03.5										
		v	03-10										
		e	05-08										
		e	05-20										
		v	05-22										
		vS	05-28.0										-3.5
		vS	05-34										2.5
	vS	P	06-18										2.6
		v	06-46										
		v	07-15										
		v	07-22										
		ML	09-12.0	10				1.4	4.3				
		ML	09-45	10	2.0	1.1			4.5				
	СД-1	ML	09-50	14				1.6	4.2				
21. Минеральные Воды	СК	e(P)	05-08-16								13.5 <sup>x</sup>		
		eS	07-45										
		eLc	08.9										
	СД-1	ML	09-58	12	1.7	2.1	2.8	5.0	ML				
22. Боровое	СМ	+P	05-03-28.5	0.5	0.019	0.077	0.352	6.2	15.01	61-3.7			
		Рmax	30.4	0.5				0.352	6.2				
		vS	06-03.1										
22. Пуджово	ВАГ	P	05-03-37.5								15.73	325	-2.0
		Рmax	40.0	1.6				0.9	6.2				
		e	06-30										
24. Львов	СКД	eP	05-03-38								15.89	285	-3.6
		v	03-45										
		e	03-52										
		e	03-54										
		e	06-26										
		e	06-52										
		LM	11-16	10	1.6				4.4	MLH			
		LM		13				1.9	4.4	MLV			
		LM		12		2.0			4.5	MLH			

I	2	3	4	5	6	7	8	9	10	11	12	13
27. Самарканд	CK	ср	05-03-89.6							15.4 <sup>x</sup>		
		i	03-41									
		l	03-47.2									
		e	06-29.8									
		ML	12-55.0	9.0	0.8	1.7	1.7		4.6 MLH			
									4.5 MLV			
28. Ташкент	СТМ	+iP	05-03-46.0							16.49	106	-3.8
		P	03-53.5									
		P <sub>max</sub>	54.0	1.5				1.0	6.3			
	CK	P <sub>max</sub>	54.0	2.0	1.0	1.5	2.5	6.6				
	СРМБ	e	06-06									
	"	e	07-07									
	"	e	09-02									
	CK	ML	12-12.0	12			3.6	2.4	4.7 MLH			
									4.6 MLV			
29. Ужгород	СРМБ	+iP	05-03-54.0							17.12	281	-3.1
		P <sub>max</sub>	04-00	1.1				0.1	5.4			
				(0.8)				(0.3)				
		i	04-09									
		i	04-18									
		i	04-30									
		i	04-43									
		i	05-04									
	СРД	i	06-32									
30. Душанбе	CK	+iP	05-04-05							17.73	114	0.1
31. Наманган	CK	ср	05-04-10.2							17.8 <sup>x</sup>		
2. Гари	СРМБ	ср	05-04-13.2							18.42	110	-0.3
		P <sub>max</sub>	13.2	0.9				0.05	5.1			
		P <sub>max</sub>	18	1.4				0.65	6.0			
		e	04-25.8									
		e	10-52.2									
		e	12-51									
		e	14-15									
33. Андриан	СРМБ	+ср	05-04-16.1							18.78	103	-1.8
	"	P <sub>max</sub>	21.5	1.1				1.1	6.3			
	CK	P <sub>max</sub>	21.5	1.6				1.5	6.3			
		e	05-12									
		e	08-54									
		ML	16-10.0	8.0	3.0	3.0	2.0		5.1 MLH			
									4.8 MLV			

I	2	3	4	5	6	7	8	9	10	11	12	13
34. Бруно	СНС	+eP	05-04-22.2							19.31	95	-1.9
		e	04-26.4									
		P <sub>max</sub>	28.0	1.0				0.33	5.8			
		i	04-40									
		e	04-56.2									
		e	08-12.4									
		e	08-24.6									
		e	12-24.5									
35. Талгар	СНС	LM	16-00.6	9.0	2.0				4.7			
	СНС	iP	05-04-41.7	1.0	0.35	0.34	+0.75					
		i	11-10.5	7.2	0.46							
		LM	12	10	0.7	0.84	0.9					
34. Ангити	СНС	+iP	05-04-42.8							21.04	344	0.1
		P <sub>max</sub>	45.6	1.0				0.85	6.1			
	СНС	P <sub>max</sub>	45.6	1.2	1.3	1.2	2.5	6.5				
	СНС	i	04-47									
		iS	08-29.4									-4.1
		iS	08-35									
		i	08-27									
		e	08-52.7									
	СНС	LM	14-30	10	1.3			1.6	4.6 <sup>ML</sup>			
									4.7 <sup>ML</sup>			
	СНС-1	LM		9.0				1.9	4.8 <sup>MLV</sup>			
35. Сентинель-Тинке	СНС	+iP	05-04-43.7							21.12	71	0.1
36. Нарын	СНС	+iP	05-04-49.2							20.73	98	9.5
		e	05-51.4									
22. Талгар	СНС	P	05-04-41.7	1.0	0.35	0.34	+0.75			20.5 <sup>Z</sup>		
		e i	11-10.5	7.2	0.46							
		LM	12	10	0.7	0.84	0.9					
37. Пржевальск	СНС	eP	05-04-54.0							22.00	93	1.4
		P <sub>max</sub>	56.0	1.0	0.08	0.1	0.32	5.7				
	СНС	LM	17-40	10	1.5	2.0	2.0	4.8 <sup>ML</sup>				
38. Новосибирск	СНС	+iP	05-05-02.5							22.85	59	1.4
		P <sub>max</sub>	04.3	1.1	0.12	0.236	0.376	5.8				
		i	06-06.3									
		e	09-13									
		S <sub>max</sub>	20.0	1.7	0.068	0.116		4.5				
	СНС-1	LM	16-24	12	0.8	1.0	1.3	4.5 <sup>ML</sup>				

I	2	3	4	5	6	7	8	9	10	II	I2	I3
39. Вильдовка	СХМЗ	+iP	05-05-19.0							24.63	63	0.9
		P <sub>max</sub>		19.0	0.7	+0.112	+0.290	0.484	6.2			
		e S	09-39									1.1
40. Норильск	СХМЗ	+iP	05-06-00.5							29.03	27	1.9
		P <sub>max</sub>			1.0			0.102	5.6			
		i	06-14									
		i	06-40									
		i	07-14									
		i	07-32									
		i	08-03									
		i	08-51									
		i	09-14									
		i	09-27									
41. Хейс	СХМЗ	+iP	05-06-34.8							32.98	3	1.5
		P <sub>max</sub>		35.8	1.3			0.1	5.6			
		i	06-52									
		i	07-37									
		i	07-50									
		i	08-18									
		i	09-18									
		i	12-07.8									
		e	13-44									
42. Москва	СХМЗ	+iP	05-06-42.7							33.76	63	2.6
		P <sub>max</sub>		43.8	1.0			0.21				
		e S	15-51									
43. Западная	СХМЗ	+iP	05-06-57.7							35.51	64	2.3
		P <sub>max</sub>			1.1			0.06	5.4			
		e	08-12.1									
	СХД	ML	23-55		11.5			0.64	4.6 HLV			
44. Иркутск	СХМЗ	+iP	05-06-57.4							35.56	61	1.1
		P <sub>max</sub>		56.7	1.0			0.192	5.9			
					8.0	0.097						
	СХД	e Le	23-30									
	СХД	LM	24-28		9.0			0.43	4.5 HLV			
45. Бодайбо	СХМЗ	+P	05-07-30							39.50	50	1.3
		P <sub>max</sub>		31.0	0.8	0.142	0.34	0.57	6.5			

??												
1	2	3	4	5	6	7	8	9	10	11	12	13
46. Tumor	CRMS	+P	05-07-56.0							42.61	27	1.9
		P <sub>max</sub>	57.5	1.0				0.084	5.6			
		•	08-05.5									
		•	09-38.5									
47. Sarcos	CRMS	+P	05-08-19.9							45.68	40	0.9
		P <sub>max</sub>	20.4	0.8				0.027	5.3			
			10-16.8									
48. Cellulose	CRMS	+P	05-09-27.4							54.48	38	1.6
		P <sub>max</sub>	28.4	1.0				0.161	6.1			
		•	10-08									
		•	10-30									
	CKI	•	16-15.4									
49. Maragon	CRMS	-P	05-09-35							55.47	36	1.8
		P <sub>max</sub>	36.0	0.6				0.04	5.7			
		•	09-57									
		•	13-56									
50. Int. Cellulose Maragon	CRMS	+P	05-10-06.3							59.91	52	1.4
		P <sub>max</sub>	07.2	0.9			0.08	0.16	6.1			
		•	10-09									
		•	10-51.8									
	IRG	LX	40-06	16			0.9	0.7	5.0 MLH			
									4.8 MLV			
	CKI	LM	40-09	16	0.5	1.0	1.0	1.0	5.1 MLH			
									5.0 MLV			

18 (дек.) декабря 1978 (Август IX) Западно-Казахстан  
(08-00) ЕССН: 0-07-59-56,0

УСС: Н=07-59-56,0  
 $\varphi = 47^{\circ} 78' N$   
 $\lambda = 48^{\circ} 14' E$   $h = 0 \text{ км}$   
МВ=5,9 МС=5,2

$\varphi = 47^{\circ} 92' N$   
 $\lambda = 48^{\circ} 10' E$   
 $h = 0 \text{ км}$   
 МРV(A) = -6.1 (22 от) МРV(B) = -6.1 (4от)

МЛН(B) = -5.1 (8от)  
 МЛН(C) = 5.1  
 МЛV(C) = -4.9 (4от)  
 Запад. Казах. (28-336) - N per 2

№	Станция	Прибор	Фаза	Время	T сек	Δμ			Мад	Δ°	Az°	Res
						NS	EW	Z				
	2	3	4	5	6	7	8	9	10	11	12	13
	Магачнала	СК	+P	08-01-15.0						4.98	185	1.3
			P <sub>max</sub>	19.0	1.4			1.0				
			i	02-37								
			o	02-55								
			i	03-34								
			ML	06-00.0	8.0	17.0	7.0	15.0	4.8			
									4.7			
2.	Пятигорск	СКМЗ	P	08-01-18.2						5.24	223	0.8
			i	01-38								
			i	01-45								
			i	03-01								
			i	04-10								
3.	Бакуриани	СКД	+P	08-01-41.0						6.99	209	1.1
			P <sub>max</sub>		2.0			0.94				
			i	02-24								
			i	02-33								
			i	02-39								
			i	03-03								
			ML		11	18.4			4.9			
4.	Вировабад	СКМЗ	iP	08-01-46.0						7.29	190	-0.3
			i	02-21								
			ML		6.0	2.0	3.0		4.5			
5.	Зочи	СКМЗ	+P	08-01-47.5						7.29	236	1.2



I	2	3	4	5	6	7	8	9	10	II	I2	I3
6.	Баку	СК	∪P	08-01-53.0						7.64	169	1.7
			P <sub>max</sub>		0.1			5.4				
			∪	01-59								
			∪	02-28								
			LM		2.0	15.0	128.0		6.5			
7.	Ленинград		+∪P	08-01-54.0						7.77	204	0.9
			e	04-18								
8.	Бреван		P	08-01-58.2						8.16	199	-0.2
			e	03-47.4								
			∪	04-22.8								
9.	Горис	ВАГ	+∪P	08-02-01.8						8.50	189	-1.5
			P <sub>max</sub>		1.2			0.6				
			∪	02-16.2								
			∪	02-32.4								
			∪	02-44.4								
			∪	04-00.6								
			e	05-07.6								
			∪	05-46								
			LM		6.0	3.8	3.7		4.7			
10.	Ст. № 1 (Михнево)	СНМЗ	∪P	08-02-15.8	0.8			1.5		9.1 <sup>x</sup>		
		СНМ	P <sub>max</sub>	16.0	0.8	1.45	1.2	1.74				
		СК	∪S	16	0.8			2.7				
		СНМ	∪S	04-02	1.6	6.6	7.8					
		СК	LM	07-34	8.0	12.5	15.5	13.0				
11.	Симферополь	СКД	-∪P	08-02-25						10.10	258	-0.3
			P <sub>max</sub>		2.0			1.2				
			∪S	04-19								-1.6
			LM	09-00	10	3.9		2.5	4.5			
									4.3			
		СД-1	LM		10			4.8	4.6			
12.	Москва		+∪P	08-02-26						10.16	324	0.0
			P <sub>max</sub>	29.0	1.5			0.15				
			e	02-36								
			e	03-08								
			LM	08-07.9	9.0			10.1	5.0			

1	2	3	4	5	6	7	8	9	10	11	12	13
13.	Обнинск	СКМЗ	-iP	08-02-24.3						10.17	319	-1.9
			P <sub>max</sub>	27.0	0.8			0.23				
			e	05-20								
		ГЕО	LM	07-40.0	10	6.6	8.1	8.3	4.9	MLH		
									4.8	MLV		
		СК	LM		8.0		7.4	10.0	4.9	MLH		
									5.0	MLV		
		СКД	LM		8.0			1070	5.0			
14.	Арты	СКМЗ	+eP	08-02-28.7						10.65	33	-4.1
			P <sub>max</sub>		1.0			0.05				
			e	02-31.2								
			e	02-33.2								
			e	02-47.2								
			e	02-59								
			e	03-18								
			e	04-20.2								
			e	04-26								
			e	04-42								
			LM	08-04	12	7.0	11.0		5.0	MLH		
			LM		11		38.8		5.5	MLH		
15.	Свердловск	СКМЗ	+iP	08-02-43.5						11.74	35	-4.:
			P <sub>max</sub>	47.5	1.5		1.	1.35				
			e	04-48								
16.	Кишинеу	СК	-eP	08-03-02						13.08	273	-3.:
			l	03-03								
		СКД	l	03-04								
			l	04-37								
			l	04-44								
			l	05-23								
			l	05-28								
		СКД	LM	09-40								
17.	Боровое	СКМЗ	+iP	08-03-26.5	0.5				0.550		14.3 <sup>x</sup>	
18.	Пулково	СКД	eP	08-03-37.0						15.78	325	-3.9
			P <sub>max</sub>	38.0	2.0			3.0				
			l	03-46								
			e	04-07								
			e	04-18								
			eS	06-27								-3.8
			e	06-51								
			e	07-35								
			e	08-24								
		СКД	LM	11-23.0	10	5.3	3.0	7.0	5.0			
									5.1			

I	2	3	4	5	6	7	8	9	10	11	12	13		
19. Самарканд	СК	+	иP	08-03-38.2							15.91	114	-4.5	
			и	03-42.2										
			и	03-46.8										
			е	06-58.4										
			СК	LM	12-52.0	9.5			5.5	5.0	MLV			
" "	"	"	LM		9.0	3.0	6.5		5.1	MLH				
			СКД	+	иP	08-03-39.7						15.94	285	-3
20. Львов	СКД	+	и	03-49										
			и	04-01										
			еS	06-40										-0.7
			и	06-55										
			и	06-58										
			и	07-42										
			LM	11-51	9.0		6.4	6.4	5.1	ML				
			СКД	еP	08-03-56.0							17.17	282	-2.7
21. Ужгород	СКД	е	иP	04-01										
			R <sub>max</sub>		0.8			0.26						
			R <sub>max</sub>		1.1			0.65						
			и	04-07										
			и	04-16										
			и	04-26										
			и	04-32										
			и	04-50										
			и	04-56										
			и	05-05										
			и	05-50										
			и	05-58										
			и	06-30										
22. Душанбе	СК	+	иS	06-55										
			иS	07-03									-6.1	
			иP	08-04-04.0								17.68	114	-1.1
			иS	07-22										1.2
			LM		8.0	6.0		5.0	5.0	MLH				
								5.1	MLV					

I	2	3	4	5	6	7	8	9	10	11	12	13
23. Гари	СКМЗ	+iP	08-04-12.9							18.37	110	-0.8
		P <sub>max</sub>			1.6			1.45				
		e	07-45.3									
			11-05									
24. Куляб	СК	+iP	08-04-16.8							18.72	114	-1.1
	"	P <sub>max</sub>			1.2			2.8				
	"	i	05-05.6									
	"	iS	07-49.3									5.0
	СК	LM	08-15.0	9.0				4.3	5.0	MIV		
25. Андижан	СКМ	+iP	08-04-16.5							18.73	103	-1.6
	"	P <sub>max</sub>			1.6			3.9				
	СК	P <sub>max</sub>		21.5	1.6			4.0				
		e	05-23									
		e	07-25									
		LM	16-15.0	8.0	10.5	12.6	12.3	5.7	MIV			
								5.6	MIV			
26. Фрунзе	СКМЗ	+eP	08-04-22.8							19.26	95	-1.6
		P <sub>max</sub>		30.7	1.4			0.47				
		P <sub>max</sub>			1.5			0.8				
		e	04-47.4									
		e	04-56.2									
		i	05-09.6									
		i	05-11.2									
		e	08-35.2									
		e	09-27									
	СКД	LM	16-13.0	9.0	7.6			3.7	5.3	MIV		
									5.0	MIV		
27. Хорог	СК	iP	08-04-36.0							20.06	112	2.5
		P <sub>max</sub>			3.0			2.0	6.1			
		i	08-23.0									8.9
		i	08-59									
		i	09-07									
		i	11-34									
		i	12-18									
	LM		14-41.0	11.0	1.0	0.4	2.0	4.4	MIV			
								4.7	MIV			

I	2	3	4	5	6	7	8	9	10	II	I2	I3		
28. Нарын	СКМЗ	eP	08-04-35.9							20.68	98	-4.1		
					Рmax	1.6		1.74	6.2					
				eS	08-34.6								7.8	
29. Талгар	СКМЗ	+iP	08-04-42.0							20.85	92	0.3		
					Рmax	43.0	1.3	0.82	6.0					
				СКМ	Рmax	6.0		1.1	5.4					
						04-57								
						08-35								
30. Апатиты	СКМЗ	+iP	08-04-42.8							21.08	344	-1.1		
					Рmax	46.0	1.2	1.9	6.4					
				СКД	Рmax		1.8	3.9	6.5					
						04-53.6								
						05-03								
						05-30								
					e	08-35								0.7
31. Семипалатинск	СКМЗ	+iP	08-04-43.5							21.08	71	-0.5		
					eS	08-30.7								
32. Пржевальск	СКМЗ	+iP	08-04-54.0							21.96	93	1.0		
					Рmax		1.0	0.64	6.0					
				СК	Рmax		1.6	1.5	6.1					
					e	05-01								
					e	09-16								
33. Новосибирск	СКМЗ	+iP	08-05-02.0							22.83	59	0.5		
					Рmax		1.6	1.4	6.2					
					eS	09-04								
				СКД-1	LM	16-26.0	12.8	1.4	3.2	3.6	4.9	MLH	5.0	MLV

I	2	3	4	5	6	7	8	9	10	11	12	13
24. Ельцовна	СКМЗ	+iP	08-05-19.0							34.60	63	0
		P <sub>max</sub>			2.0			1.5	6.2			
		e <sub>S</sub>	09-41									
25. Хельс	СКМЗ	+iP	08-06-35.0							33.00	2	0.7
		P <sub>max</sub>	36.0	1.4				0.2	5.8			
		i	06-51									
		i	07-35									
		e	07-50									
		i	08-13									
		i	09-18									
		i	09-57									
		i	10-46									
	СКД	LM	28-04.0	19	2.3	9.6	4.4	5.2 <sup>MAX</sup>	5.4 <sup>MLH</sup>			
26. Морды	СКМ	+iP	08-06-42.0							33.73	63	1.1
		P <sub>max</sub>		1.6				0.65	6.3			
27. Закамненск	СКМЗ	+iP	08-06-57.3							35.48	64	1.4
		P <sub>max</sub>	56.6	1.4				0.17	5.8			
		e	08-11.9									
	СКД	LM	23-38	14				1.6	4.9 <sup>MLV</sup>			
28. Иркутск	СКМЗ	+iP	08-06-56.5							35.53	61	0.2
	"	P <sub>max</sub>		1.3				0.19	5.8			
	СКД	P <sub>max</sub>		2.0				0.43	6.0			
		e	07-01.8									
		e	07-13.8									
		e	07-20									
		e	07-48									
		e	08-21.8									
	СКД	LM	23-40	13				1.71	4.9 <sup>MLV</sup>			
	"	LM		6.0	0.64				4.9 <sup>MLH</sup>			
	"	LM		14		1.46			4.8 <sup>MLH</sup>			
29. Зоданбо	СКМЗ	+iP	08-07-30							39.55	50	0.1
		P <sub>max</sub>		1.0				0.41	6.3			

1	2	3	4	5	6	7	8	9	10	11	12	13
40. Тикси	СКМ	+iP	08-07-55.5							42.61	27	0.6
	СКД	Pmax	56.5	2.0				0.3	5.8			
	СКМ	Pmax	56.5	1.1				0.18	5.9			
		e	09-39.5									
	СКД	LM	29-03	16				1.3	4.9	M/V		
		LM		16	0.6				4.5	M/H		
		LM		18			1.8		5.0	M/H		
41. Тушик	СКМЗ	+iP	08-08-05.1							43.67	53	1.4
42. Якутск	СКМЗ	-iP	08-08-18.5							45.67	40	-1.2
		i	10-13.5									
		Pmax	10-16.0	1.8				0.5	6.2			
43. Ст. 15 2	СКМ	+iP	08-09-13.0	2.0	0.136	0.256	0.53			52.0 <sup>x</sup>		
		Pmax						0.715				
		i	10-21	1.4				0.126				
		e	14-03									
		e	32-15									
		e	34-18									
44. Сейчан	СКМЗ	-iP	08-09-27.6							54.42	33	1.1
		Pmax	28.4	1.7				0.46	6.3			
		e	09-35.8									
45. Магадан 1	СКМЗ	-iP	08-09-34.5							55.46	36	0.4
		PP	35.3									0.8
		Pmax	36.5	1.0				0.08	5.8			
		e	09-56									
46. Дульгин	СКМЗ	-iP	08-09-54							59.31	18	-0.2
		Pmax	57.0	1.6				0.06	5.4			
		Pmax	10-02	1.0				0.29	6.3			
			10-09									
			10-29									
			10-33									
			10-41									
		e	18-42									
		e	19-00									
	СКД	ML	43-00	14	1.0			1.3	5.1	M/V		
47. Дж. Саха-ЛИНСК	СКМЗ	+ P	08-10-06							59.89	52	0.5
		Pmax	07.1	1.1				0.27	6.2			
		ePP	10-10.7									
		e	10-17									
		e	10-33									
		e	10-52									
		e	21-30									
		e	33-36									

18. XII. 78 Азгир-IX - 235-  
(08-00)

Дополнение из с/бюл. с/карточек

I	2	3	4	5	6	7	8	9	10	11	12	13
48.	Зугдиди	СК	+iP	08-01-44.0								6.8 <sup>x</sup>
49.	Абастумани	СКМЗ	+iP	08-01-45.0								6.9 <sup>x</sup>
			e <sup>3</sup>	02-20.0	0.5	1.0						
50.	Алушта	СХ	iP	08-02-22.8								
			P <sub>max</sub>	23.8	0.6				0.10			
			iPP	02-33								
			e	04-11								
		СК	e	04-44								
			e <sup>L</sup>	06-56								
			LM	08.6	1.0				1.0	4.2MLV		
51.	Ялта	СХ	P	08-02-26.0								9.8 <sup>x</sup>
			P <sub>max</sub>	27.0	0.5				1.4			
			P <sub>max</sub>	27.0	0.5	0.5	0.5					
			S	04-20.2								
			S <sub>max</sub>	23.0	0.9	0.4						
			S <sub>max</sub>	24.4	0.75		0.3					
52.	Минск	СК	eP	08-03-16								13.5 <sup>x</sup>
			P <sub>max</sub>	20.0	2.0	1.4I	5.6	2.4				
			e <sup>S</sup>	05-49								
			e <sup>h<sub>e</sub></sup>	08.4								
		СД-I	LM	09-51	12	4.3			6.3			
		СК	LM	10-08	10		8.8					
		СК	LM	10-10	8.0	5.0			5.0			
53.	Ташкент	СКМЗ	P	08-03-45.3								15.3 <sup>x</sup>
			P <sub>max</sub>	48.3	1.2	0.5	1.0	1.2				
		СК	P <sub>max</sub>		2.0	0.6	1.0	2.5				
			l	53.3								
			iM	54.7	2.0	2.2	3.5	8.5				
			e	07-05								
			e	07-30								
			e <sup>L</sup>	10-11								
		СД-I	LM	12-10	13				11.3	5.3	MLV	
		СКД	LM		12		12			5.4	MLH	
		СК	LM		11		11		8.3	5.2	MLH	
					9	5.5				5.2	MLH	
54.	Наманган	СК	+iP	08-04-09.2	1.4			2.6				17.7 <sup>x</sup>



18.XII.78 Азгир-IX  
(08-00)

I	2	3	4	5	6	7	8	9	10	11	12	13
55.	Фергана	СК	+eP eS L LM	08-04-13.9 07-47.7 10 13-57.2	2.3			2.0		18.1 <sup>x</sup>		
56.	Мургаб	СК	iP eS	08-04-46.0 08-42.0	1.0	0.7	1.0	1.0		20.9 <sup>x</sup>		
57.	Усть Элегест	СКМЗ	+iP LS	08-06-05.2 17-40.2	1.3	0.02	0.08	0.2	6.6	29.2 <sup>x</sup>		

- 238 -  
 (Jan 1979)  
 17 января 1979 (Август УН) - Западный Казахстан  
 Астраханская обл.  
 (08-00)

УСC: H=07-59-55,8  
 $\varphi = 47,96 N$   
 $\lambda = 48,14 E$   $h = 0 км$   
MB=6,0 MS=4,6

УСC: H=07-59-54,6  
 $\varphi = 47,93 N$   
 $\lambda = 48,06 E$   
 $h = 0 км$   
 МК<sub>кн</sub> (A)-5,9 (по 20 ст)  
 МЛ Н(B)-4,8 (по 7 ст)  
 МЛ Н(C)-4,8 (по 9 ст)  
 Западный Казахстан  
 (регистр № 29-336)

№ шт	Станция	Приб. Аппа	Время	T сек	M			Mag	$\Delta^\circ$	Az <sup>o</sup>	RoS		
					NS	EW	Z						
1	2	3	4	5	6	7	8	9	10	11	12	13	
1.	Пятигорск	СНМЗ	iP	08-01-17.6	1,0			2,9		5,23	283	1,7	
			P <sub>max</sub>	01-28									
			i	01-40									
			i	01-46									
			M	02-02									
2.	Баурнани	СНМЗ	ep	08-01-42.0						6,99	209	1,3	
			i	02-22									
			i	02-30									
3.	Зугдиди	СВК	+iP	08-01-43						6,8 <sup>x</sup>			
4.	Абастумани	СВК	+iP	08-01-45						6,9 <sup>x</sup>			
5.	Сочи	СНМЗ	+iP	08-01-46.0						7,28	236	1,3	
			i	01-48									
			i	01-52									
			i	02-18									
			i	02-23									
			i	02-27									
			Mh	05-15.0	14,0					3,8	4,1		
6.	Кировабад	СНМЗ	iP	08-01-46.0						7,30	190	0,9	
			i	01-57									
			i	02-12									
			i	02-28									
			i	03-05									
7.	Степанован	СКД	+iP	08-01-47.0	1,5					1,5		7,0 <sup>x</sup>	
			is	59.8									
			i	02-06.2									
			i	26.6									
			i	45.8	2,0					2,5			
			LM	08-05	12					3,0			

1	2	3	4	5	6	7	8	9	10	11	12	13
8. Ленинград	СКД	+iP	08-01-54.0							7.77	204	2.3
		i	02-82									
		• S	03-12									-0.5
		•	03-52									
		hM	08-16.2	12					2.5	4.1		
9. Брест	СК	+iP	08-01-57.0							8.16	199	-0.1
	СКМЗ	i	02-19									
	СК	(S)	02-44									
		eML	04-41									
10. Горис	ВБГ	+iP	08-02-02		8,0	1,0	1,0	1,0	4,3	8.51	189	0.0
		Pmax		1.6						1.26		
		i	02-15									
	СК	i	02-23.6									
		•	05-18									
		ML	08-48,6	10.0	1.0	0.8	0.8		3.9 ML			
									3.7 ML			
11. Ст. № 1 (Мичурин)	СКМ	+iP	08-02-13.8	0.5				0.07		8.9 <sup>X</sup>		
12. Алушта	СК	iP	08-02-21.5							9.5 <sup>X</sup>		
		Pmax		22.7	0.5			0.8				
		i	02-82									
		i	02-41.5									
		• S	04-08									
		Smax		09.5	1.0	0.72						
13. Симферополь	СКД	• P	08-02-22.1							10.08	257	-1.5
	СХ	• P	22,8									
		Pmax	25,0	0.4				0.5				
	СКД	• S	04-18									
	СКД	ML	07-31	17				1.0	3.7			
	СД-1	ML	07-34	16				1.0	3.7			
14. Москва	СХ	+ P	08-02-23							10.13	324	-1.3
		•	03-11									
15. Ялта	СХ	• P	08-02-24.5							9.6		
		Pmax		25	0.5	0.4	0.4	1.2				
		i	02-44.5									
		S	04-12.5									
		Smax		18.5	0.6	0.3	0.3					
16. Обнинск	СКМЗ	+ P	08-02-24.5							10.14	319	0.1
		Pmax			1.0					1.2		
		•	05-14									
	СК	ML	08-08.0	8.0	7.8	9.7	14.0		5.1 ML			
									5.2 ML			

17. Арты	СКОЗ СКД СКМ	eP	08-02-28.0					10.66	85	-3.5
		P <sub>max</sub>	29.7	0.9			0.2			
		e	02-30							
		e	02-39							
		e	02-44.5							
		e	03-01.7							
		e	04-22.7							-10.0
		e	04-40.7							
		e	05-22.7							
		e	06-27							
	ML	08-14.0	10	1.0	2.0		4.4 <sup>MLH</sup>			
			11			3.2	4.4 <sup>MLV</sup>			
18. Кызыл- Арбат	СК	eP	08-02-31					10.71	143	-1.3
		eS	03-39							
19. Свердловск	СКОЗ	+ P	08-02-42.5					11.74	86	-3.8
		e	04-52							
	ML <sub>2</sub>	08,7	28	1,5	3,0	2,5				
20. Анжыбад	СКОЗ	+i P	08-02-54.8					12.49	139	-1.7
		P <sub>max</sub>	54,8	1.0			0.45			
21. Кашгар	СКД	i P	08-03-02	1,0	1,0	1,0		13.05	273	-1.9
		P <sub>max</sub>	03.0	1.2			1.4			
		i	03-10							
		i	04-29							
		i	04-50							
		i	05-29							
		i	05-49							
	ML	09-37.0	11			2,5	4.5 <sup>MLV</sup>			
22. Боровое	СКМ	-i P	08-03-26.4	0.4			0.777	14.3 <sup>x</sup>		
23. Пулково	ВЭГ	i P	08-03-36.0					15.75	323	-3.1
		P <sub>max</sub>	40.0	1.3			0.7			
	СКД	P <sub>max</sub>	38.0	1.3			1.0			
		i	03-44	1,8	1,9	1,7				
		eS	06-26						-8.7	
	СД-I	ML <sub>2</sub>	10-40	15			1.6	4.3 <sup>MLV</sup>		
		IM <sub>ML<sub>2</sub></sub>	II-38	8,0	1,3	1,4	1,5	4, IM <sup>MLV</sup>		
24. Львов	СКД	eP	08-03-36					15.90	285	-5.3
		i	03-46							
		e	03-57							
		e	06-04							
		ML	11-41.0	10			2.5	4.6 <sup>MLV</sup>		
25. Самар- Канд	СК	- P	08-03-39.2					15.3 <sup>x</sup>		
		i	46.8							
		P <sub>max</sub>	47.4	0.4			2.3			
	ML	13-21.4	9.0	1.0	1.5	1.5				

1	2	3	4	5	6	7	8	9	10	11	12	13
26.	Ужгород	СКС	-iP	08-08-55.5						17.14	281	-1.4
			Р <sub>max</sub>		1.0			0.5				
			i	04-05								
			i	04-10								
			i	04-14								
			i	04-18								
			i	04-23								
			i	04-28								
			i	04-33								
			i	04-53								
			i	05-03								
			i	05-35								
			i	06-08								
			i	07-00								
			i	09-23								
		СКД	ML	12-00.0	12		2.5	3.5	4.6 ML 4.8 ML			
27.	Душанбе	СК	eP	08-04-04.2						17.71	114	0.1
28.	Бергама	СКМ	iP	08-04-04.1						17.3 <sup>X</sup>		
		СК	LM	13-								
29.	Гарм	СКС	eP	08-04-12.4						18.40	110	-0.3
		"	Р <sub>max</sub>	19.3	1.1			0.68				
		СКД	Р <sub>max</sub>		2.0			0.44				
30.	Куляб	СК	+iP	08-04-16.5						18.75	114	-0.5
			Р <sub>max</sub>	08-28.0	1.2			1.3				
			i	05-21								
			e	10-29.7								
31.	Анди- жан	СКС	-eP	08-04-16.2						18.76	103	-0.9
		"	Р <sub>max</sub>	20.0	1.6			2.6				
		СК	Р <sub>max</sub>		1.6			2.8				
			e	08-31								
		СК	LM	16-17	8.0	1.5	4.5	3.6	5.1 ML 5.0 ML			
32.	Фрунзе	СКС	-iP	08-04-21.1						19.29	95	-2.3
			Р <sub>max</sub>	26.6	1.0			0.38				
		СКД	e (PP)	04-41.1								
		"	e	04-53.1								
		СКМ	LS	07-46.5								-9.3
			i	07-48.1								
			i	07-51.1								
			e	07-55.1								
		СК	ML	14-23	10			1.5	4.6 ML			
		СКД	ML	14-25	12			2.0	4.6 ML			

1	2	3	4	5	6	7	8	9	10	11	12	13
32. Тамгар	СМЗ	+ P	08-04-41.0							20.88	92	0.3
		$R_{max}$	41.3	1.2	0.13	0.2	0.5					
		$R_{max}$	44.0	1.4			1.5					
33. Амгунь	СМЗ	+ P	08-04-41.0							21.06	344	-1.3
		$R_{max}$	43.8									-0.7
	СКД	LM	14-24.0	11	2.0	0.5	2.5	4.7 ML				
								4.8 ML				
	СД-1	LM		10			2.6					
34. Семин- ЛЕНСКОЕ	СМЗ	+ P	08-04-42.6							21.11	71	-0.3
		$R_{max}$	43.8	0.8			0.81					
35. Приваловск	СМЗ	+iP	06-04-53.3							21.99	93	1.4
		$R_{max}$		1.2			0.4	5.7				
Новосел- бирск	СМЗ	+iP	08-05-00.3							22.84	59	0.0
		$R_{max}$		1.3			0.25	5.5				
37. Бельцовка	СМЗ	+iP	08-05-09.4							24.62	63	-8.1
		$R_{max}$		1.4			0.95	6.1				
38. Хейс	СМЗ	+iP	08-06-33.4							32.99	2	0.6
		$R_{max}$	34.6	1.4			0.12	5.6				
		i PP	06-38									4.6
		i	07-16									
		i	07-48									
		i	08-41									
		i	09-18									
39. Монах	СМЗ	+iP	08-06-41.0							33.75	63	1.3
		$R_{max}$		1.1			0.35	6.2				
40. Закамское	СМЗ	+ P	08-06-56.5							35.50	64	1.8
		$R_{max}$	57.5	1.2	0,040	0,074	0.12	5.7				
		e	09-24.6	I, I			0,064	5,4				
		e	11-46.5									
		e	16-43.5									
	СКД	ML	23-37.5	10,0	0,42		0.74	4.6 MLV				
41. Иркутск-2	СМЗ	+eP	08-06-56.3							35.55	61	1.2
		$R_{max}$	57,3	1.0	I, II		0.33	6.2				
		e	07-12.8	I, 2			0,205					
		e	07-38.5									
	СКД	LM	23-44	12			0.68	4.6 MLV				
	СКД	LM	23-44	13		0.5		4.4 MLV				

17.1.79

-243-

1	2	3	4	5	6	7	8	9	10	11	12	13
42.	Борисово	СМЗ	+P	08-07-28.6						39.56	50	-0.1
			R <sub>max</sub>		1.0			0.9	6.6			
43.	Тихон	СМЗ	oP	08-09-26.5						42.61	27	33.0
			R <sub>max</sub>	39.5	1.2			0.11	5.7			
44.	Икуток	СМЗ	-P	08-08-18.7						45.68	40	0.3
			R <sub>max</sub>	20.0	1.0			0.53				
			i	10-14								
45.	Ст. Б 2	СМЗ	+P	08-09-12.0	1.2	0.051	0.084	0.19		51.8 <sup>2</sup>		
			Pop	10-22	1.2			0.008				
46.	Сельчан	СМЗ	+P	08-09-25.9						54.43	33	0.7
			R <sub>max</sub>	27.4	1.0			0.02	5.3			
			R <sub>max</sub>		1.1			0.08	5.7			
			R <sub>max</sub>		1.2			0.11	5.8			
			e	08-17.9								
47.	Магадан-1	СМЗ	-iP	08-09-33.8						55.47	36	1.0
			R <sub>max</sub>	36	1.0			0.11	5.9			
			i	09-39								
			i	09-56								
			e	10-08								
			e	10-34								
			e	12-07								
48.	Владивосток	СМЗ	-i	08-10-37.4						56.16	61	59.4
			R <sub>max</sub>		1.1			0.16	6.0			
49.	Куальтан	СМЗ	iP	08-10-00.5						59.31	18	0.7
			R <sub>max</sub>		1.0			0.18	6.1			
			iP	10-05								
			i	10-19								
			i	10-43								
49.	Р. Селен- Линское	СМЗ	+iP	08-10-05						59.90	52	0.8
			R <sub>max</sub>	05.8	1.0			0.28	6.3			
			oP	10-10								5.0
			e	10-25								
			e	10-44								
			e	12-12.5								
			e	12-17								
			oK	12-18.6	1.1			0.06				
			e	18-39								
			e	21-04								
			e	23-57								
			e	26-56								
			e	28-24								
		СМЗ/М		40-08.0	15	0.7	1.5	1.1	5.2 ML			
									8.1 ML			
50.	Новокава- рельская	УСБ	PK EP	08-18-50								
			R <sub>max</sub>	50.5	0.8			0.03				
			e	20-38								

17.1.79

51. Углегорск	СКД	e	08-09-53.3						57.5 <sup>x</sup>
52. Махачкала	СК	+iP	08-01-15.5	0.8	-1.4	-1.1	+4.0		4.8 <sup>x</sup>
		P <sub>max</sub>	18.0	0.8			9.0		
		v	01-22.8						
		e S	02-44						
		e S	02-54						
		S <sub>max</sub>	02-58.0	1.4	11	7.0			
		M <sub>кр</sub>	06-18	8.0	4.5	3.2	4.3	ML	
53. Баку	СК	+iP	08-01-53						7.5 <sup>x</sup>
		e	01-59						
		e	02-40	1.0	5.1	5.0	6.4		
54. Минск	СК	+iP	08-03-16						13.5 <sup>x</sup>
		e <sub>кр</sub>	09,5						
		e <sub>кр</sub>	09,7						
	СД-1	M <sub>кр</sub>	09-50	12			2.4		
55. Мургаб	СК	iP	08-04-45.0	0.8	-0.3	+0.5	+0.6		20.8 <sup>x</sup>
		e S	08-41						
56. Тикси	СВКМЗ	+ P	08-07-54						41.9 <sup>x</sup>
	СКД	+iP	54.5						
	"	P <sub>max</sub>	55.5	1.4			0.2	5.9	
	СВКМЗ	P <sub>max</sub>	55.5	1.1			0.21	6.1	
	"	e	08-03						
	"	e	17.5						
	СКД	M <sub>кр</sub>	28-53	18			0.6	4.5	
				17		0.3	4.3		
57. Петропав- ловск	СКМЗ	e	08-10-23.5						61.8 <sup>x</sup>
		e	12-25						
58. Ст. № 2 (Кульдур)	СКМ	+ P	08-09-12,0	I,2	0,05I	0,084	0,19		51,8 <sup>x</sup>



14 июля 1979 (Аэгур Х1) - Зональный Казахстан  
(05-00)

91

УС: Н=04-59-58,2  
 $\varphi = 47^{\circ}31'N$   
 $\lambda = 48^{\circ}07'E$   $h = 0 \text{ км}$   
 МВ=5,6

СССР-04-59-54,2  
 $\varphi = 49,99'N$   
 $\lambda = 48,10'E$   
 $h = 0 \text{ км}$

МР(А)=5,6 (13ст) М(В)=4,1(4ст)  
 Зап.Казахстан СССР

№ ПП	Станция	Приб.	Фаза	Время	Т СВК	АМ			Мг	$\Delta^{\circ}$	Az $^{\circ}$	ReS
						NS	EW	Z				
1	2	3	4	5	6	7	8	9	10	11	12	13
1.	Грозный		i P	05-01-13.0				2,0		4.96	200	1.3
			e	01-49		2,3	3,3					
2.	Махацкала	СН	+ P	05-01-14.6		2,9	0,9	2,1		5.04	184	1.8
			e S	02-09	1,2			3,1				-3.7
			L M	07-03	8,0	0,9	1,3	1,5	3,7			
3.	Лятигорск	СНМЗ	e P	05-01-16						5.29	223	-0.3
			i	01-25.5								
			i	01-37.5								
			e	01-41.5								
			i	01-43.5								
4.	Бакуриана	СНМЗ	+i P	05-01-40.0						7.05	209	-1.1
			P max		1.0			0.43				
			e	01-41								
			i	02-26								
			i	02-25								
			i	02-32								
			i	02-41								
			M L		12		2.7	4.0				
5.	Зурдиди	СВК	+i P	05-01-43						6.8 <sup>x</sup>		
			P max		0.5			1.0				
			e S	02-21								
			S max		1.0		2.2					
6.	Абастумани	СНМЗ	+i P	05-01-44.0		0.6		0.44		6.8 <sup>x</sup>		
			i S	02-17	0.8			0.4				

1	2	3	4	5	6	7	8	9	10	11	12	13
7.Сочи		СНМЗ	-iP	05-01-45.9						7.33	236	0.9
			i	02-25								
			MH	05-17.0	12			0.5	3.3			
8.Кировабад		СНМЗ	eP	05-01-45.0						7.36	190	-0.4
			i	01-46								
			i	02-53								
			i	02-11								
9.Шемаха			-iP	05-01-48.5						7.36	176	3.0
			i	04-45								
			MH	06-50.0	11	2.0			4.0			
10.Ленинакан		СНМЗ	eP	05-01-52						7.83	204	-0.1
			Pmax		2.0			0.6				
			i	02-32								
			i	03-08								
11.Ереван			-iP	05-01-57.8						8.22	199	0.3
			i	02-10.8								
			i	02-37.2								
12.Горис		ВЭГ	-eP	05-02-01						8.57	189	-61.3
			i	01-08.8								
			i	01-14.8								
			e	04-30.4								
			i	05-04.4								
			e	05-51.8								
13.Ст. № 1 (Чухнево)		СНМ	-iP	05-02-14.6	0.4			0.860		9.0 <sup>x</sup>		
14.Адушта		СХ	-iP	05-02-21.0						9.5 <sup>x</sup>		
			Pmax		22.0	0.6		0.27				
			i	02-28								
			i	02-37.6								
			e	03-32								
			i	03-46.5								
			eS	04-14								
			iS	04-18								
			Sm ax	04-19.5	0.7	0.19						
15.Москва		СХ	+ P	05-02-23.0						10.10	324	-0.4
			Pmax		27.0	1.0		1.1				
			MH	08-00	8.0			0.9	4.0/MH			

1	2	3	4	5	6	7	8	9	10	11	12	13
16.Синфоро- поль	СК	еР	05-02-23.0							10.12	257-0.7	
		Р <sub>нак</sub>		0.4				0.4				
		;	02-37									
		•	04-17									-2.1
		LM	07-28.0	18				0.3	3.1			
	СД	LM	08-08	13				0.4	3.4			
17.Ялта	СК	•	05-02-23.8							9.7 <sup>x</sup>		
		+ P	24.1	0.8	0.2	0.2	0.7					
18.Арчи	СЮМЗ	еР	05-02-27.7							10.60	33	-2.5
		•	02-30.7									
		•	02-42									
		•	02-46.7									
		•	04-18.7									
		•	04-22.7									
		•	04-28.7									
		•	04-46.7									
		•	05-02									
		LM	08-19.0	10	0.6	1.0			4.0	MLH		
		LM		12				1.5	4.1	MLV		
19.Кизил- Арват	СК	еР	05-02-30							10.73	143-2.1	
		•	02-35									
		•	03-25.5									
		•	03-37.5									
20.Свердловск	СЮМЗ	+ P	05-02-43							11.69	36-2.0	
		Р <sub>нак</sub>	46.2	1.2				0.48				
		;	02-48									
		•	02-58									
		•	03-10									
		•	03-26									
		•	03-52									
		•	04-46									
		•	04-57									
		•	05-05									
		•	05-14									
		LM	08-48.0	12	0.5	0.6	1.0		3.8	MLH		
									3.9	MLV		

14.УП.79

1	2	3	4	5	6	7	8	9	10	11	12	13
21.	Ашхабад	СКМЗ	P	05-02-54.6						12.51	139	-1.7
			P <sub>max</sub>	54.6	0.5			0.11				
			e	03-07								
			e	03-12.8								
			e	07-45.1								
			e	08-08								
			e	08-44.6								
			LM	14-10.6	8.5			0.9	4.1			
22.	Кашаев	СКД	-iP	05-03-02.0						13.08	272	-1.8
			P <sub>max</sub>	02.5	1.2			0.5				
			i	05-21								
			i	06-58								
		СКД	LM	09-36.0	11			1.1	4.1			
23.	Боровое	СКМ	iP	05-03-25.5						14.95	61	-2.9
			P <sub>max</sub>		0.5			0.110				
24.	Пудново	ВЗГ	-iP	05-03-36.0						15.72	325	-2.3
			P <sub>max</sub>	38.0	1.6			0.42				
			e	03-44								
			i	03-54								
			e <sup>S</sup>	06-29								
			e	06-52								
		СД-1	LM	11-22.0	9.0	0.6	0.4	0.6	4.1 <sub>MLH</sub>			
									4.1 <sub>MLV</sub>			
25.	Львов	СКД	eP	05-03-37.0						15.93	285	-3.9
			i	03-37								
			i	03-45								
			e	07-32								
			e	08-27								
			LM	11-45.0	10			0.9	4.2 <sub>MLV</sub>			
26.	Самарканд	СК	-iP	05-03-38.5						15.93	114	-2.7
			i	03-41.3								
			i	03-46.3								
		СК	P <sub>max</sub>	46.7	0.5			1.5				
		СК	LM	13-25.3	8.5		0.5	0.7	4.0 <sub>MLH</sub>			
									4.2 <sub>MLV</sub>			

1	2	3	4	5	6	7	8	9	10	11	12	13
27.	Ужгород	СКСЗ	еР	05-03-54.5						17.16	281	-2.2
				03-57								
				04-02								
				04-05								
				04-11								
				04-17								
				04-27								
				04-33								
				04-39.5								
				04-53								
				06-23								
				06-33								
				08-00								
				08-08								
				09-12								
				09-24								
				10-00								
28.	Душанбе	СК	еР	05-04-02.3						17.71	114	-1.3
				13-56.1								
29.	Наманган	СК	еР	05-04-09.2						17.7 <sup>к</sup>		
30.	Гурн	СКСЗ	еР	05-04-12.1						18.40	111	0.0
			Р. ак	17.0	2.0			0.35				
31.	Фергана	СКСЗ	еР	05-04-13.5	0.8			0.24		18.0 <sup>к</sup>		
				04-27.5	1.0			0.3				
		СК	е <sup>с</sup>	07-43.5								
		"	ЛМ	13-49.5	10			1.4				
32.	Кудяб	СК	еР	05-04-15.5						18.74	114	-0.9
			е	04-45.5								
			е	10-29.3								
33.	Анджкан	СК	еР	05-04-15.2						18.74	103	-1.2
		"	Р. ак		1.5			0.8				
		СКСЗ	Р. ак		1.3			0.62				
			е	09-14.2								
			е	10-49.2								
			ЛМ	14-10.0	12	0.7	1.4	1.1	4.5 <sup>MLH</sup>			
									4.3 <sup>MLV</sup>			
			ЛМ	16-16	8.0	0.7	1.6	1.1	4.7 <sup>MLH</sup>			
									4.5 <sup>MLV</sup>			

1	2	3	4	5	6	7	8	9	10	11	12	13
34.	Брунзе	СНМЗ	-eP	05-04-22.0						19.27	95	-0.6
			P <sub>max</sub>	26.5	0.9		0.16					
		СНД	LM	14-27.0	14		0.6	0.6	4.0 <sub>MLH</sub> 4.0 <sub>MLV</sub>			
35.	Хорог	СК	eP	05-04-33.3						20.08	112	1.3
36.	Нарын	СНМ	+iP	05-04-40						20.69	98	1.7
			i	05-39.2								
37.	Талгар	СНМЗ	+iP	05-04-41.0	1.0	-0.028	+0.025	+0.07		20.85	92	1.1
			P <sub>max</sub>	44	0.9	0.19	0.16	0.38	5.8			
			e	10-25.1			0.133					
			e	12-08.1	6.0	0.2						
			e	15-16.1	10		0.268					
38.	Апатиты	СНМЗ	eP	05-04-41.2						21.01	344	-0.2
		СНД	ML	14-10	13	0.58	0.57	0.96	4.3 <sub>MLH</sub> 4.3 <sub>MLV</sub>			
39.	Семипалатинск	СНМЗ	+iP	05-04-42.4						21.06	71	0.4
			e	04-54.2								
			e	05-30.1								
			e	07-44.6								
			e <sub>S</sub>	08-30.7								-1.7
40.	Новосибирск	СНМЗ	+iP	05-05-01						22.79	59	1.7
			P <sub>max</sub>		1.2			0.2	5.5			
				09-09								4.6
		СД-1	LM	16-35.0	12	0.3	0.4		4.1 <sub>ML</sub>			
41.	Ельцовка	СНМЗ	+iP	05-05-17.9						24.57	63	1.4
			P <sub>max</sub>		1.0			0.23	5.7			
			e <sub>S</sub>	09-41								5.7
42.	Хейс	СНМЗ	+eP	05-06-34						32.94	2	2.1
			P <sub>max</sub>	35.0	1.4			0.09	5.5			
			i	07-46								
			i	07-48								
43.	Заваяменск	СНМЗ	+iP	05-06-55.8						35.45	64	2.0
			P <sub>max</sub>		1.0			0.04	5.3			
44.	Бодайбо	СНМЗ	+iP	05-07-28.5						39.50	50	0.8
			P <sub>max</sub>		0.3			0.26	6.6			
45.	Тизес	СНМЗ	+iP	05-07-54.0						42.55	27	1.5
			P <sub>max</sub>	55.5	1.1			0.05	5.4			
			e	09-29.5								

1	2	3	4	5	6	7	8	9	10	11	12	13
46. Куров	СРБЗ	еР	05-09-12.2							45.62	40	-3.2
		R <sub>max</sub>	20.2	1.0			0.08	5.7				
		✓	10-14.2									
		✓	12-46.2									
47. Севидан	СРБЗ	+iP	05-09-25.9							54.37	33	1.6
		R <sub>max</sub>	26.9	1.0			0.09	5.8				
48. Магадані	СРБЗ	- еР	05-09-34.8							55.91	36	2.9
		R <sub>max</sub>	35.0	1.0			0.04	5.5				
		е	10-17									
		е	11-13									
49. Дульгин	СРБЗ	+ P	05-09-59.6							59.25	18	0.7
		е	10-39									
50. Л. Сажо-	СРБЗ	+ uP	05-07-01.4							59.65	52	-182.0
ЛАНСК		е	07-2.6									
		е	07-46									
		е	07-48									
51. Ваку	СК	еР	05-01-51.0							7.3 <sup>x</sup>		
		е	01-57									
		е	02-26									
		S	02-35									
		S <sub>max</sub>	03-03.0	1.6	2.5	1.9						
52. Минск	СК	еР	05-03-15							13.5 <sup>x</sup>		
		е	09-34									
		е	05-53									
		е	12-32									
53. Мургаб	СК	2SS iP	05-04-44.0							20.7 <sup>x</sup>		
		еS	08-41									
54. Новоси-	СКБЗ	+iP	05-05-00.8							22.4 <sup>x</sup>		
бирск		R <sub>max</sub>	01.5	1.2	0.066	0.132	0.206	5.6				
		i	08.5									
		pP	22.2									
		iS	09-08.9									
		S <sub>max</sub>	39.2	1.4	0.015	0.044		4.7				
		i(SS)	13-37.2									
	СД-I	Miq	16-35.2	12	0.3	0.4		4.2				

(October) 268 -  
 24 октября 1979 (АЗГВР) Западное Казakhstan  
 (06-00)

СССР-05-59-55,5

$\psi = 47,95^{\circ}N$

$\lambda = 48,10E$

0

MPV(A)-5,7 (по 15 ст) MLH(D)=4.3

CRM

сд (5ст)

MLV(B)-4,4 (по 9 ст)

CR

Зап. Казахстан ССРС

УРС: H=05-59-56,7

$\psi = 47,79^{\circ}N$

$\lambda = 48,11E$  h=0км

MB=5,8

№ ст	Станция	Приб.	База	Время	T <sub>обс</sub>	Am			Mag	$\Delta^{\circ}$	Az	Pos
						NS	EW	Z				
1	Грозный		JP	06-01-14.4						4.93	200	1.7
				01-40.5								
2	Макашала	СМЗ	+P	06-01-15.5						5.01	185	1.8
			Pmax	19.0	0.5			4.4				
				01-26								
			CS	03-13								
			LM	04-50	9.0	0.4	3.0	1.0	3.9 3.5			
3	Пятигорск	СМЗ	+P	06-01-17.7						5.26	223	0.4
			Pmax	19.2	1.0			2.3				
				01-21.2								
				01-27.2								
				01-28.2								
				01-44								
				01-57								
				03-12.2								
				04-23.2								
4	Степановая	СМЗ	+P	06-01-37.4	1.0			0.5		6.4 <sup>K</sup>		
5	Бакуриани	СМЗ	+P	06-01-42						7.02	209	-0.1
			Pmax		0.8			0.51				
		СМД	Pmax		0.8			0.66				
				02-18								
			LM		10	3.2			4.2			
6	Зугдиди	СВК	+P	06-01-44						6.8 <sup>K</sup>		
			Pmax		0.5			3.3				
7	Абастумани	СМЗ	+P	06-01-45	0.5			0.8				
			CS	02-16	0.5	0.5						

4. Степановая

06-23-07  
 05-41  
 04-08  
 04-20  
 Pmax 06-05-5



1	2	3	4	5	6	7	8	9	10	11	12	13
8. Сочи	СМБ	+P	06-01-47.2							7.31	236	1.1
		Рмак	52.0	0.6				1.3				
			01-56									
			02-37									
			02-52.5									
			03-02									
9. Кировабд	СКД	LM	05-18	13				1.3	3.7			
	СМБ	P	06-01-46.0							7.32	190	-0.3
			03-58									
		e	06-09									
10. Тельма		-P	06-01-46							7.33	176	-0.4
			02-25									
		LM	06-03	8.0				4.5	4.4			
11. Баку	СК	+P	06-01-53							7.68	169	1.8
			01-56									
			02-37									
12. Ленинскан	ВЭГ	eP	06-01-53							7.80	204	0.0
		e	02-34									
13. Гревон	СК	+P	06-01-59.5							8.19	199	1.1
		e	05-32.5									
14. Горис	ВЭГ	+P	06-02-01.4							8.54	189	-1.8
	СМБ		02-16									
			02-31.2									
		e	02-57									
		e	03-40.4									
		e	04-00.2									
15. Нахичевань	СК	-P	06-02-10	1.0	2.8			1.1		8.7 <sup>X</sup>		
		S	02-54	1.5	3.4							
16. Ст. № 1	СМБ	+P	06-02-16	0.7				0.800		9.1 <sup>X</sup>		
17. Алушта	СК	-eP	06-02-21.6							9.6 <sup>X</sup>		
		+P	02-23.2									
		Рмак	23.7	0.5				0.6				
			02-29.1									
			02-47.7									
			03-01									
		e(S)	04-14.7									
	СК	e	05-05									
		e	06-11									
	LM		06-08,9	(9)				0.5	3.7			

1	2	3	4	5	6	7	8	9	10	11	12	13	
18. Самаро- Поль	СХ	СХ	СР	06-02-24.0						10.11	257	-0.9	
			С	04-20									-0.2
19. Москва	СХ	СХ	ЛМ	08-20.0	12			0.5	3.5				
			СР	06-02-25.0							10.13	324	-0.1
			С	02-35									
			С	02-55									
20. Ялта	СХ	СХ	СР	06-02-25.2						9.8 <sup>к</sup>			
			СР	25.9	0.5	-0.4	-0.4	+1.2					
			С	04-19.8									
			С	04-22.7									
21. Обнинск	СХ	СХ	СР	06-02-25						10.14	319	-0.3	
			Р <sub>max</sub>	28.0	1.0			0.9					
			С	05-14									
			ЛМ	07-40.0	10	1.3	1.4	1.7	4.2				
22. Арз	СХ	СХ	СР	06-02-29.6						10.62	33	-2.3	
			С	02-42									
			С	02-59.6									
			С	04-26.6									-6.2
23. Кавказ- Арз	СХ	СХ	ЛМ	08-10	11		2.0		4.2	MLH			
			ЛМ		13			2.4	4.2	MLV			
			СР	06-02-31.5						10.71	143	-1.6	
			С	02-36.5									
24. Свердловск	СХ	СХ	СР	06-02-45						11.71	36	-1.8	
			Р <sub>max</sub>	47.5	1.0			0.88					
			С	02-57									
			С	03-04									
25. Ашгабад	СХ	СХ	СР	06-02-54.4						12.49	139	-2.9	
			С	04-05									
			Р <sub>max</sub>		2.0			0.98					
			ЛМ	05-43									
	СХ	СХ	ЛМ	13-53.5	3.5			1.45	4.7				



1	2	3	4	5	6	7	8	9	10	11	12	13
33.	Ужгород	СМЗ	+ eP	06-08-57.0						17.16	281	-1.1
				08-59								
				04-02								
			P <sub>max</sub>	08.0	0.9			0.25				
				04-04								
				04-07								
				04-14								
				04-20								
				04-29								
				04-34								
				04-53								
				05-09								
				11-41								
				11-56								
34.	Душанбе	СК	+ P	06-04-04.6						17.69	114	-0.2
			LM	14-37.1	9.5		1.5		4.5			
35.	Наманган	СК	-eP	06-04-09.6						17.7 <sup>X</sup>		
36.	Бергана	СМЗ	-P	06-04-14.7	1.0			0.29		18.1 <sup>X</sup>		
		СК	eS	08-52.2								
			ML	14-30.2	9.0	2.0			5.0			
37.	Куляб	СМЗ	+ P	06-04-17.3						18.73	114	-0.3
			P <sub>max</sub>	21.1	1.3			1.0				
				04-30								
			e	10-05.6								
38.	Анджкан	СМ	-P	06-04-16.9						18.74	103	-0.8
			P <sub>max</sub>		1.3			1.2				
		СК	P <sub>max</sub>		1.8			1.5				
			e	08-49								
		СК	LM	16-10	8.0	2.0	3.1	2.3	5.0			
									4.8			
39.	Фрунзе	СМЗ	-eP	06-04-23.4						19.27	95	-0.6
			P <sub>max</sub>	28.0	0.8			0.13				
				04-39.3								
				05-11.8								
			eS	07-52.1								-4.0
			e	08-25								
		СКД	LM	16-09	10	1.4			4.5			
40.	Хорог	СК	eP	06-04-33						20.07	112	-0.2

1	2	3	4	5	6	7	8	9	10	11	12	13
49.	Забайкальск	СНС	+P	06-06-57.8						35.47	64	2.5
		Р.ак			1.4			0.05	5.3			
		е		09-26								
		СД	ЛМ	23-52.4								
50.	Иркутск-2	СНС	еP	06-06-57.8						35.51	61	2.1
		Р.ак			1.2			0.1	5.6			
		СД	ЛМ	23-40	14			0.26	4.1			
51.	Бодайбо	СНС	+P	06-07-30.1						39.52	50	0.9
		Р.ак			0.8			0.49	6.4			
52.	Тирас	СНС	+P	06-07-55.8						42.58	27	1.7
		Р.ак		56.5	1.0			0.06	5.5			
53.	Сейчан	СНС	-eP	06-09-27.9						54.40	33	2.1
54.	Магадан-1	СНС	еP	06-09-35.6						55.44	36	2.1
55.	Мульте	СНС	+P	06-10-01.6						59.28	18	1.1
		Р.ак			0.8			0.13	6.1			
		еP		10-06								4.4
		е		10-18								
56.	Кр. Саха- линск	СНС	+P	06-10-07						59.87	52	2.1
		Р.ак		07.3	0.9			0.05	0.13			
		е		10-12.2								
57.	Новоа- заревская	УОБ	PR, NP	06-18-51						121.26	193	1.8
		Р.ак		52.0	0.8			0.01				

1	2	3	4	5	6	7	8	9	10	11	12	13
41. Нарын	СРМЗ	+	P	06-04-40.5						20.69)	98	0.9
			P <sub>max</sub>		1.4			0.18	5.3			
	СКД		LM	15-14	10			0.97	4.4			
42. Талгар	СРМЗ	+	P	06-04-42.0	1.1	-0.035	+0.043	+0.098		20.85	92	0.8
			P <sub>max</sub>	45.0	1.4	0.23	0.29	0.062	4.0			
			•	05-26								
			•	08-49.4	9.0		0.2					
			•	09-32								
			LM	15-00	12			0.6	4.1			
43. Апатиты	СРМЗ	-	P	06-04-43.3						21.04	344	0.2
			SP	04-44.2								0.9
			•S	05-36								2.8
	СКД		LM	14-26.0	12	1.9	0.7	1.9	4.7			
									4.7			
	СД-1		LM	14-16.0	10			2.4	4.8			
44. Семипалатинск	СРМЗ	+	P	06-04-43.8						21.07	71	0.3
			•	05-15.6								
			•	08-23.1								
45. Пржевальск	СРМЗ	+	P	06-04-55						21.96	93	2.5
			P <sub>max</sub>		1.3			0.32	5.5			
	СК		P <sub>max</sub>		1.6			0.49	5.6			
			•	05-11.5								
			LM	17-30	10			1.2	4.6			
46. Новосибирск	СРМЗ	+	P	06-05-02.2						22.81	59	1.4
			P <sub>max</sub>		1.6			0.42	5.7			
			•S	09-05								-1.0
	СД-1		LM	19-27	12.	0.35	0.86	1.03	4.4			
									4.4			
47. Ельцовка	СРМЗ	+	P	06-05-18.9						24.58	63	0.9
			P <sub>max</sub>		1.0			0.35	5.9			
			•S	09-45								8.1
48. Монды	СРМЗ	+	P	06-06-42						33.72	63	1.7
			P <sub>max</sub>		1.0			0.15	5.8			

Prof. Thomas Ahrens  
Seismological Lab, 252-21  
Division of Geological & Planetary Sciences  
California Institute of Technology  
Pasadena, CA 91125

Prof. Keiiti Aki  
Center for Earth Sciences  
University of Southern California  
University Park  
Los Angeles, CA 90089-0741

Prof. Shelton Alexander  
Geosciences Department  
403 Deike Building  
The Pennsylvania State University  
University Park, PA 16802

Prof. Charles B. Archambeau  
CIRES  
University of Colorado  
Boulder, CO 80309

Dr. Thomas C. Bache, Jr.  
Science Applications Int'l Corp.  
10260 Campus Point Drive  
San Diego, CA 92121 (2 copies)

Prof. Muawia Barazangi  
Institute for the Study of the Continent  
Cornell University  
Ithaca, NY 14853

Dr. Jeff Barker  
Department of Geological Sciences  
State University of New York  
at Binghamton  
Vestal, NY 13901

Dr. Douglas R. Baumgardt  
ENSCO, Inc  
5400 Port Royal Road  
Springfield, VA 22151-2388

Dr. Susan Beck  
Department of Geosciences  
Building #77  
University of Arizona  
Tucson, AZ 85721

Dr. T.J. Bennett  
S-CUBED  
A Division of Maxwell Laboratories  
11800 Sunrise Valley Drive, Suite 1212  
Reston, VA 22091

Dr. Robert Blandford  
AFTAC/TT, Center for Seismic Studies  
1300 North 17th Street  
Suite 1450  
Arlington, VA 22209-2308

Dr. Stephen Bratt  
ARPA/NMRO  
3701 North Fairfax Drive  
Arlington, VA 22203-1714

Dr. Lawrence Burdick  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

Dr. Robert Burrige  
Schlumberger-Doll Research Center  
Old Quarry Road  
Ridgefield, CT 06877

Dr. Jerry Carter  
Center for Seismic Studies  
1300 North 17th Street  
Suite 1450  
Arlington, VA 22209-2308

Dr. Eric Chael  
Division 9241  
Sandia Laboratory  
Albuquerque, NM 87185

Dr. Martin Chapman  
Department of Geological Sciences  
Virginia Polytechnical Institute  
21044 Derring Hall  
Blacksburg, VA 24061

Mr Robert Cockerham  
Arms Control & Disarmament Agency  
320 21st Street North West  
Room 5741  
Washington, DC 20451,

Prof. Vernon F. Cormier  
Department of Geology & Geophysics  
U-45, Room 207  
University of Connecticut  
Storrs, CT 06268

Prof. Steven Day  
Department of Geological Sciences  
San Diego State University  
San Diego, CA 92182

Marvin Denny  
U.S. Department of Energy  
Office of Arms Control  
Washington, DC 20585

Dr. Art Frankel  
U.S. Geological Survey  
922 National Center  
Reston, VA 22092

Dr. Zoltan Der  
ENSCO, Inc.  
5400 Port Royal Road  
Springfield, VA 22151-2388

Dr. Cliff Frolich  
Institute of Geophysics  
8701 North Mopac  
Austin, TX 78759

Prof. Adam Dziewonski  
Hoffman Laboratory, Harvard University  
Dept. of Earth Atmos. & Planetary Sciences  
20 Oxford Street  
Cambridge, MA 02138

Dr. Holly Given  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

Prof. John Ebel  
Department of Geology & Geophysics  
Boston College  
Chestnut Hill, MA 02167

Dr. Jeffrey W. Given  
SAIC  
10260 Campus Point Drive  
San Diego, CA 92121

Eric Fielding  
SNEE Hall  
INSTOC  
Cornell University  
Ithaca, NY 14853

Dr. Dale Glover  
Defense Intelligence Agency  
ATTN: ODT-1B  
Washington, DC 20301

Dr. Petr Firbas  
Institute of Physics of the Earth  
Masaryk University Brno  
Jecna 29a  
612 46 Brno, Czech Republic

Dan N. Hagedon  
Pacific Northwest Laboratories  
Battelle Boulevard  
Richland, WA 99352

Dr. Mark D. Fisk  
Mission Research Corporation  
735 State Street  
P.O. Drawer 719  
Santa Barbara, CA 93102

Dr. James Hannon  
Lawrence Livermore National Laboratory  
P.O. Box 808  
L-205  
Livermore, CA 94550

Prof Stanley Flatte  
Applied Sciences Building  
University of California, Santa Cruz  
Santa Cruz, CA 95064

Prof. David G. Harkrider  
Seismological Laboratory  
Division of Geological & Planetary Sciences  
California Institute of Technology  
Pasadena, CA 91125

Dr. John Foley  
NER-Geo Sciences  
1100 Crown Colony Drive  
Quincy, MA 02169

Prof. Danny Harvey  
CIRES  
University of Colorado  
Boulder, CO 80309

Prof. Donald Forsyth  
Department of Geological Sciences  
Brown University  
Providence, RI 02912

Prof. Donald V. Helmberger  
Seismological Laboratory  
Division of Geological & Planetary Sciences  
California Institute of Technology  
Pasadena, CA 91125



Prof. Eugene Herrin  
Institute for the Study of Earth and Man  
Geophysical Laboratory  
Southern Methodist University  
Dallas, TX 75275

Prof. Robert B. Herrmann  
Department of Earth & Atmospheric Sciences  
St. Louis University  
St. Louis, MO 63156

Prof. Lane R. Johnson  
Seismographic Station  
University of California  
Berkeley, CA 94720

Prof. Thomas H. Jordan  
Department of Earth, Atmospheric &  
Planetary Sciences  
Massachusetts Institute of Technology  
Cambridge, MA 02139

Prof. Alan Kafka  
Department of Geology & Geophysics  
Boston College  
Chestnut Hill, MA 02167

Robert C. Kemerait  
ENSCO, Inc.  
445 Pineda Court  
Melbourne, FL 32940

Dr. Karl Koch  
Institute for the Study of Earth and Man  
Geophysical Laboratory  
Southern Methodist University  
Dallas, Tx 75275

Dr. Max Koontz  
U.S. Dept. of Energy/DP 5  
Forrestal Building  
1000 Independence Avenue  
Washington, DC 20585

Dr. Richard LaCoss  
MIT Lincoln Laboratory, M-200B  
P.O. Box 73  
Lexington, MA 02173-0073

Dr. Fred K. Lamb  
University of Illinois at Urbana-Champaign  
Department of Physics  
1110 West Green Street  
Urbana, IL 61801

Prof. Charles A. Langston  
Geosciences Department  
403 Deike Building  
The Pennsylvania State University  
University Park, PA 16802

Jim Lawson, Chief Geophysicist  
Oklahoma Geological Survey  
Oklahoma Geophysical Observatory  
P.O. Box 8  
Leonard, OK 74043-0008

Prof. Thorne Lay  
Institute of Tectonics  
Earth Science Board  
University of California, Santa Cruz  
Santa Cruz, CA 95064

Dr. William Leith  
U.S. Geological Survey  
Mail Stop 928  
Reston, VA 22092

Mr. James F. Lewkowicz  
Phillips Laboratory/GPEH  
29 Randolph Road  
Hanscom AFB, MA 01731-3010( 2 copies)

Mr. Alfred Lieberman  
ACDA/VI-OA State Department Building  
Room 5726  
320-21st Street, NW  
Washington, DC 20451

Prof. L. Timothy Long  
School of Geophysical Sciences  
Georgia Institute of Technology  
Atlanta, GA 30332

Dr. Randolph Martin, III  
New England Research, Inc.  
76 Olcott Drive  
White River Junction, VT 05001

Dr. Robert Masse  
Denver Federal Building  
Box 25046, Mail Stop 967  
Denver, CO 80225

Dr. Gary McCartor  
Department of Physics  
Southern Methodist University  
Dallas, TX 75275

**Prof. Thomas V. McEvilly**  
Seismographic Station  
University of California  
Berkeley, CA 94720

**Dr. Art McGarr**  
U.S. Geological Survey  
Mail Stop 977  
U.S. Geological Survey  
Menlo Park, CA 94025

**Dr. Keith L. McLaughlin**  
S-CUBED  
A Division of Maxwell Laboratory  
P.O. Box 1620  
La Jolla, CA 92038-1620

**Stephen Miller & Dr. Alexander Florence**  
SRI International  
333 Ravenswood Avenue  
Box AF 116  
Menlo Park, CA 94025-3493

**Prof. Bernard Minster**  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

**Prof. Brian J. Mitchell**  
Department of Earth & Atmospheric Sciences  
St. Louis University  
St. Louis, MO 63156

**Mr. Jack Murphy**  
S-CUBED  
A Division of Maxwell Laboratory  
11800 Sunrise Valley Drive, Suite 1212  
Reston, VA 22091 (2 Copies)

**Dr. Keith K. Nakanishi**  
Lawrence Livermore National Laboratory  
L-025  
P.O. Box 808  
Livermore, CA 94550

**Prof. John A. Orcutt**  
IGPP, A-025  
Scripps Institute of Oceanography  
University of California, San Diego  
La Jolla, CA 92093

**Prof. Jeffrey Park**  
Kline Geology Laboratory  
P.O. Box 6666  
New Haven, CT 06511-8130

**Dr. Howard Patton**  
Lawrence Livermore National Laboratory  
L-025  
P.O. Box 808  
Livermore, CA 94550

**Dr. Frank Pilotte**  
HQ AFTAC/TT  
1030 South Highway A1A  
Patrick AFB, FL 32925-3002

**Dr. Jay J. Pulli**  
Radix Systems, Inc.  
201 Perry Parkway  
Gaithersburg, MD 20877

**Dr. Robert Reinke**  
ATTN: FCTVTD  
Field Command  
Defense Nuclear Agency  
Kirtland AFB, NM 87115

**Prof. Paul G. Richards**  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

**Mr. Wilmer Rivers**  
Teledyne Geotech  
314 Montgomery Street  
Alexandria, VA 22314

**Dr. Alan S. Ryall, Jr.**  
ARPA/NMRO  
3701 North Fairfax Drive  
Arlington, VA 22203-1714

**Dr. Richard Sailor**  
TASC, Inc.  
55 Walkers Brook Drive  
Reading, MA 01867

**Prof. Charles G. Sammis**  
Center for Earth Sciences  
University of Southern California  
University Park  
Los Angeles, CA 90089-0741

**Prof. Christopher H. Scholz**  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Dr. Susan Schwartz  
Institute of Tectonics  
1156 High Street  
Santa Cruz, CA 95064

Secretary of the Air Force  
(SAFRD)  
Washington, DC 20330

Office of the Secretary of Defense  
DDR&E  
Washington, DC 20330

Thomas J. Sereno, Jr.  
Science Application Int'l Corp.  
10260 Campus Point Drive  
San Diego, CA 92121

Dr. Michael Shore  
Defense Nuclear Agency/SPSS  
6801 Telegraph Road  
Alexandria, VA 22310

Dr. Robert Shumway  
University of California Davis  
Division of Statistics  
Davis, CA 95616

Dr. Matthew Sibol  
Virginia Tech  
Seismological Observatory  
4044 Derring Hall  
Blacksburg, VA 24061-0420

Prof. David G. Simpson  
IRIS, Inc.  
1616 North Fort Myer Drive  
Suite 1050  
Arlington, VA 22209

Donald L. Springer  
Lawrence Livermore National Laboratory  
L-025  
P.O. Box 808  
Livermore, CA 94550

Dr. Jeffrey Stevens  
S-CUBED  
A Division of Maxwell Laboratory  
P.O. Box 1620  
La Jolla, CA 92038-1620

Lt. Col. Jim Stobie  
ATTN: AFOSR/NL  
110 Duncan Avenue  
Bolling AFB  
Washington, DC 20332-0001

Prof. Brian Stump  
Institute for the Study of Earth & Man  
Geophysical Laboratory  
Southern Methodist University  
Dallas, TX 75275

Prof. Jeremiah Sullivan  
University of Illinois at Urbana-Champaign  
Department of Physics  
1110 West Green Street  
Urbana, IL 61801

Prof. L. Sykes  
Lamont-Doherty Geological Observatory  
of Columbia University  
Palisades, NY 10964

Dr. David Taylor  
ENSCO, Inc.  
445 Pineda Court  
Melbourne, FL 32940

Dr. Steven R. Taylor  
Los Alamos National Laboratory  
P.O. Box 1663  
Mail Stop C335  
Los Alamos, NM 87545

Prof. Clifford Thurber  
University of Wisconsin-Madison  
Department of Geology & Geophysics  
1215 West Dayton Street  
Madison, WI 53706

Prof. M. Nafi Toksoz  
Earth Resources Lab  
Massachusetts Institute of Technology  
42 Carleton Street  
Cambridge, MA 02142

Dr. Larry Turnbull  
CIA-OSWR/NED  
Washington, DC 20505

Dr. Gregory van der Vink  
IRIS, Inc.  
1616 North Fort Myer Drive  
Suite 1050  
Arlington, VA 22209

Dr. Karl Veith  
EG&G  
5211 Auth Road  
Suite 240  
Suitland, MD 20746

Prof. Terry C. Wallace  
Department of Geosciences  
Building #77  
University of Arizona  
Tuscon, AZ 85721

Dr. Thomas Weaver  
Los Alamos National Laboratory  
P.O. Box 1663  
Mail Stop C335  
Los Alamos, NM 87545

Dr. William Wortman  
Mission Research Corporation  
8560 Cinderbed Road  
Suite 700  
Newington, VA 22122

Prof. Francis T. Wu  
Department of Geological Sciences  
State University of New York  
at Binghamton  
Vestal, NY 13901

ARPA, OASB/Library  
3701 North Fairfax Drive  
Arlington, VA 22203-1714

HQ DNA  
ATTN: Technical Library  
Washington, DC 20305

Defense Intelligence Agency  
Directorate for Scientific & Technical Intelligence  
ATTN: DTIB  
Washington, DC 20340-6158

Defense Technical Information Center  
Cameron Station  
Alexandria, VA 22314 (2 Copies)

TACTEC  
Battelle Memorial Institute  
505 King Avenue  
Columbus, OH 43201 (Final Report)

Phillips Laboratory  
ATTN: XPG  
29 Randolph Road  
Hanscom AFB, MA 01731-3010

Phillips Laboratory  
ATTN: GPE  
29 Randolph Road  
Hanscom AFB, MA 01731-3010

Phillips Laboratory  
ATTN: TSML  
5 Wright Street  
Hanscom AFB, MA 01731-3004

Phillips Laboratory  
ATTN: PL/SUL  
3550 Aberdeen Ave SE  
Kirtland, NM 87117-5776 (2 copies)

Dr. Michel Bouchon  
I.R.I.G.M.-B.P. 68  
38402 St. Martin D'Herès  
Cedex, FRANCE

Dr. Michel Campillo  
Observatoire de Grenoble  
I.R.I.G.M.-B.P. 53  
38041 Grenoble, FRANCE

Dr. Kin Yip Chun  
Geophysics Division  
Physics Department  
University of Toronto  
Ontario, CANADA

Prof. Hans-Peter Harjes  
Institute for Geophysics  
Ruhr University/Bochum  
P.O. Box 102148  
4630 Bochum 1, GERMANY

Prof. Eystein Husebye  
NTNF/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY

David Jepsen  
Acting Head, Nuclear Monitoring Section  
Bureau of Mineral Resources  
Geology and Geophysics  
G.P.O. Box 378, Canberra, AUSTRALIA

Ms. Eva Johannisson  
Senior Research Officer  
FOA  
S-172 90 Sundbyberg, SWEDEN

Dr. Peter Marshall  
Procurement Executive  
Ministry of Defense  
Blacknest, Brimpton  
Reading FG7-FRS, UNITED KINGDOM

Dr. Bernard Massinon, Dr. Pierre Mechler  
Societe Radiomana  
27 rue Claude Bernard  
75005 Paris, FRANCE (2 Copies)

Dr. Svein Mykkeltveit  
NTNT/NORSAR  
P.O. Box 51  
N-2007 Kjeller, NORWAY (3 Copies)

Prof. Keith Priestley  
University of Cambridge  
Bullard Labs, Dept. of Earth Sciences  
Madingley Rise, Madingley Road  
Cambridge CB3 0EZ, ENGLAND

Dr. Jorg Schlittenhardt  
Federal Institute for Geosciences & Nat'l Res.  
Postfach 510153  
D-30631 Hannover , GERMANY

Dr. Johannes Schweitzer  
Institute of Geophysics  
Ruhr University/Bochum  
P.O. Box 1102148  
4360 Bochum 1, GERMANY

Trust & Verify  
VERTIC  
Carrara House  
20 Embankment Place  
London WC2N 6NN, ENGLAND