

The Euro Med treatment of Festuca (Gramineae) — new names and new combinations in Festuca and allied genera

Authors: Foggi, Bruno, Scholz, Hildemar, and Valdés, Benito

Source: Willdenowia, 35(2) : 241-244

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.35.35202>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

BRUNO FOGGI, HILDEMAR SCHOLZ & BENITO VALDÉS

The Euro+Med treatment of *Festuca* (*Gramineae*) – new names and new combinations in *Festuca* and allied genera**Abstract**

Foggi, B., Scholz, H. & Valdés, B.: The Euro+Med treatment of *Festuca* (*Gramineae*) – new names and new combinations in *Festuca* and allied genera. – Willdenowia 35: 241-244. – ISSN 0511-9618; © 2005 BGBM Berlin-Dahlem.
doi:10.3372/wi.35.35202 (available via <http://dx.doi.org/>)

As a consequence of the recognition of *Drymochloa*, *Leucopoa*, *Schedonorus* and *Parafestuca*, and of the reassessment of accepted specific and subspecific taxa, some new names and combinations are required in *Drymochloa*, *Festuca*, *Leucopoa* and *Schedonorus*, which are validated here.

The genus *Festuca* L., considered in its wide sense (but excluding *Vulpia* C. C. Gmel.), includes a series of taxonomic groups that are but rather remotely related among each other and to its core. Some are extra-European, such as *Austrofestuca* (Tzvelev) E. B. Alexeev of Australia and palaeotropical areas, *Helleria* Fourn. of Central and South America, *Pseudobromus* K. Schum. of SE Africa and Madagascar and *Tzvelevia* E. B. Alexeev of the Kerguelen Island; whereas others are well represented in the Euro+Med area.

In her treatment for Flora Europaea (5: 125-153. 1980), Markgraf-Dannenberg recognised *Festuca* s.l. as including 170 species, and she followed the same conservative concept in Davis's Flora of Turkey (9: 400-442. 1985). Since then, a large number of contributions on *Festuca* s.l. were published that have considerably increased the number of recognised taxa, both for Europe and the extra-European territories covered by the Euro+Med project. At the same time, some monophyletic groups were segregated generically from *Festuca* s.str. Whereas Clayton & Renvoize (Gen. Gram. 1986) still recognised *Festuca* in its wide sense, Watson & Dallwitz (Grass Gen. World 1992) treated *Leucopoa* Griseb. as a distinct genus on the basis of its peculiar morphological and biological characters (see Scholz & Strid in Willdenowia 22: 87. 1992); Holub (in Folia Geobot. Phytotax. 19: 95-99. 1984) separated *F.* sect. *Montanae* Hack. as *Drymochloa* Holub, a group of nemoral species with a series of relatively primitive features. *Schedonorus* P. Beauv., characterised by broad leaf blades with basal auricles and a flat or subconcave caryopsis free from the palea, is becoming widely accepted as a separate genus rather than a mere subgenus

of *Festuca*. Alexeev (in Bjull. Moskovsk. Obšč. Isp. Prir., Otd. Biol. 90: 102-109. 1985) separated *F. albida* Lowe, an endemic of the Madeira Islands, as *Parafestuca* E. B. Alexeev.

Festuca L. is being considered for the Euro+Med treatment in a restricted sense that excludes *Drymochloa*, *Leucopoa*, *Schedonorus* and *Parafestuca*. The recognition of these segregate genera, together with the taxonomic and nomenclatural reassessment of several taxa in *Festuca* s.str. make necessary the establishment of some new combinations and new names, as follows:

Drymochloa donax (Lowe) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca donax* Lowe in Trans. Cambridge Philos. Soc. 4: 9. 1831.

Drymochloa drymeja subsp. *exaltata* (C. Presl) Foggi & Signorini, **comb. nov.** ≡ *Festuca exaltata* C. Presl, Fl. Sicul.: 45. 1826 ≡ *Festuca montana* subsp. *exaltata* (C. Presl) K. Richt., Pl. Eur. 1: 106. 1890 ≡ *Schedonorus exaltatus* (C. Presl) Schult. & Schult. f., Mantissa 3: 644. 1817 [“*Festuca sylvatica*” sensu C. Presl, Cyper. Gramin. Sicul.: 37. 1820 (non Huds., Fl. Angl. 38. 1762)].

Festuca gautieri subsp. *lutea* (Hack.) Foggi & Signorini, **comb. nov.** ≡ *Festuca varia* var. *lutea* Hack., Monogr. Festuc. Eur.: 181. 1882 ≡ *Festuca lutea* (Hack.) K. Richt., Pl. Eur. 1: 105. 1890 ≡ *Festuca scoparia* subsp. *lutea* (Hack.) Beldie in Sävulescu, Fl. Rep. Soc. Rom. 12: 496. 1972.

Festuca kozanensis Foggi & J. Müller, **nom. nov.** ≡ *Festuca rechingeri* Markgr.-Dann. in Veröff. Geobot. Inst. ETH Stiftung Rübel Zürich 56: 166. 1976 [non E. B. Alexeev in Bjull. Moskovsk. Obšč. Isp. Prir., Otd. Biol. 80(2): 122. 1975].

Festuca nitida subsp. *macrathera* (Hack.) Foggi & Signorini, **comb. nov.** ≡ *Festuca violacea* var. *macrathera* Hack. in Ann. K. K. Naturhist. Hofmus. 2: 45. 1887 ≡ *Festuca violacea* subsp. *macrathera* (Hack.) Markgr.-Dann. in Bot. J. Linn. Soc. 76: 326. 1978.

Festuca ovina subsp. *molineri* (Litard.) Foggi & J. Müller, **comb. nov.** ≡ *Festuca ovina* var. *molineri* Litard. in Bull. Soc. Bot. France 8(a): 101. 1937.

Festuca plonkae Foggi & Signorini, **nom. nov.** ≡ *Festuca halleri* subsp. *yvesii* Kerguélen & Plonka in Bull. Soc. Bot. Centre-Ouest, ser. 2, 19: 18. 1988 [non *Festuca yvesii* Sennen & Pau in Treb. Inst. Catalana Hist. Nat. 3: 196. 1917].

Festuca polita subsp. *cretica* (Markgr.-Dann.) Foggi & H. Scholz, **comb. nov.** ≡ *Festuca polita* var. *cretica* Markgr.-Dann. in Veröff. Geobot. Inst. ETH Stiftung Rübel Zürich 56: 106. 1976.

Festuca pseudovaria subsp. *winnebachensis* (Wallosek) J. Müller, **comb. nov.** ≡ *Festuca varia* var. *winnebachensis* Wallosek in Folia Geobot. Phytotax. 34: 73. 1999.

Festuca romoi Foggi & Valdés, **nom. nov.** ≡ *Festuca trabutii* Romo in Collect. Bot. (Barcelona) 16: 339. 1986 [non E. B. Alexeev in Novosti Sist. Vysš. Rast. 14: 39. 1977].

Festuca stojanovii (Acht.) [Kožuharov in Phytol. Balcan. 3: 71. 1997, **comb. inval.**, ex] Foggi & Petrova, **comb. nov.** ≡ *Festuca dalmatica* subsp. *stojanovii* Acht. in Izv. Bot. Inst. (Sofija) 3: 17. 1953.

Festuca stricta subsp. *saxatilis* (Schur) Foggi & Signorini, **comb. nov.** ≡ *Festuca saxatilis* Schur, Enum. Pl. Transsilv.: 791. 1866 ≡ *Festuca rupicola* subsp. *saxatilis* (Schur) Rauschert in Feddes Repert. 63: 276. 1960 ≡ *Festuca valesiaca* subsp. *saxatilis* (Schur) E. B. Alexeev in Vestn. Moskovsk. Univ., Ser. 6, Biol. 27(5): 49. 1972.

Leucopoa calabrica (Hack.) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca calabrica* Huter & al. ex Hack., Monogr. Festuc. Eur.: 190. 1882 ≡ *Festuca spectabilis* subsp. *calabrica* (Hack.) Arcang., Comp. Fl. Ital., ed. 2: 61. 1894.

Leucopoa carpatica (F. Dietr.) H. Scholz, **comb. nov.** ≡ *Festuca carpatica* F. Dietr., Nachtr. Vollst. Lex. Gärtn. 3: 333. 1817.

Leucopoa dimorpha (Guss.) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca dimorpha* Guss., Pl. Rar. 34. 1826 ≡ *Festuca laxa* subsp. *dimorpha* (Guss.) St.-Yves in Bull. Soc. Bot. France 71: 132. 1924.

Leucopoa laxa (Host) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca laxa* Host, Icon. Descr. Gram. Austriac. 2: 58. 1802 [= *Festuca canescens* Host, Fl. Austriac. 1: 155. 1827].

Leucopoa pulchella (Schrad.) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca pulchella* Schrad., Fl. Germ.: 336. 1806 ≡ *Schedonorus pulchellus* (Schrad.) P. Beauv., Ess. Agrostogr.: 177. 1812.

Leucopoa pulchella subsp. *jurana* (Gren.) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca pulchella* var. *jurana* Gren. in Mém. Soc. Emul. Doubs, ser. 3, 10: 925. 1864.

Leucopoa spectabilis (Bertol.) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca spectabilis* Bertol., Fl. Ital. 1: 612. 1834 [= *Festuca nemorosa* Dalla Torre & Sarnth., Fl. Tirol 6(2): 272. 1906 = *F. sieberi* Tausch in Flora 20: 127. 1837].

Leucopoa spectabilis subsp. *affinis* (Hack.) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca affinis* Hack. in Heldreich Fl. Graec. Exsicc.: [in schedis] No. 2776. 1852 ≡ *Festuca spectabilis* subsp. *affinis* (Hack.) Hack. in Bot. Centralbl. 8: 408. 1881 [= *Festuca croatica* Pichl. in Kerner, Sched. Fl. Exs. Austro-Hung. 1: 105. 1881].

Leucopoa spectabilis subsp. *carniolica* (Hack.) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca spectabilis* subsp. *carniolica* Hack., Monogr. Festuc. Eur.: 189. 1882 ≡ *Festuca carniolica* (Hack.) K. Richt., Pl. Eur. 1: 106. 1890 ≡ *Festuca siebieri* subsp. *carniolica* (Hack.) Hegi, Ill. Fl. Mitt.-Eur. 1: 351. 1908.

Leucopoa stygia (H. Scholz & Strid) H. Scholz & Foggi, **comb. nov.** ≡ *Festuca stygia* H. Scholz & Strid in Willdenowia 22: 85. 1992.

Schedonorus arundinaceus subsp. *cirtensis* (St.-Yves) H. Scholz & Valdés, **comb. nov.** ≡ *Festuca arundinacea* var. *cirtensis* St.-Yves in Candollea 1: 36. 1922 ≡ *Festuca arundinacea* subsp. *cirtensis* (St.-Yves) Gamisans, Cat. Pl. Vasc. Corse: 63. 1985.

Schedonorus arundinaceus subsp. *corsicus* Foggi & Signorini, **comb. nov.** ≡ *Festuca elatior* subvar. *corsica* Hack., Monogr. Festuc. Eur.: 157. 1882 ≡ *Festuca arundinacea* subsp. *corsica* (Hack.) Kerguélen in Bull. Soc. Bot. France 123: 320. 1976 ≡ *Festuca fenas* subsp. *corsica* (Hack.) K. Richt., Pl. Eur. 1: 102. 1890.

Schedonorus arundinaceus subsp. *mediterraneus* (Hack.) H. Scholz & Valdés, **comb. nov.** ≡ *Festuca elatior* subvar. *mediterranea* Hack., Monogr. Festuc. Eur.: 154. 1882 ≡ *Festuca arundinacea* subsp. *mediterranea* (Hack.) K. Richt., Pl. Eur. 1: 102. 1890 ≡ *Lolium arundinaceum* subsp. *mediterraneum* (Hack.) Banfi & al. in Mem. Soc. Ital. Sci. Nat. Mus. Civico Storia Nat. Milano 33(2): 8. 2005.

Schedonorus arundinaceus subsp. *orientalis* (Hack.) H. Scholz & B. Valdés, **comb. nov.** ≡ *Festuca elatior* subvar. *orientalis* Hack., Monogr. Festuc. Eur.: 154. 1882 ≡ *Festuca arundinacea* subsp. *orientalis* (Hack.) Tzvelev in Spisok Rast. Gerb. Fl. SSSR Bot. Inst. Vsesojuzn. Akad. Nauk 18: 17. 1970.

Schedonorus fontqueri (St.-Yves) H. Scholz & Valdés, **comb. nov.** ≡ *Festuca fontqueri* St.-Yves in Cavanillesia 3: 5. 1930 = *Festuca yvesii* Litard. in Bull. Soc. Hist. Nat. Afrique N. 14: 337. 1923 [non Sennen & Pau in Treb. Inst. Catalana Hist. Nat. 3: 196. 1917].

Schedonorus pauneroi (Cebolla & al.) H. Scholz, **comb. nov.** ≡ *Festuca pauneroi* Cebolla & al. in Candollea 58: 195. 2003.

Schedonorus pratensis subsp. *apenninus* (De Not.) H. Scholz & Valdés, **comb. nov.** ≡ *Festuca pratensis* var. *apennina* (De Not.)

Hegi, Ill. Fl. Mitt.-Eur. 1: 343.1908 ≡ *Schedonorus apenninus* (De Not.) Tzvelev in Novosti Sist. Vysš. Rast. 31: 259. 1998 ≡ *Lolium pratense* subsp. *apenninum* (De Not.) Banfi & al. in Mem. Soc. Ital. Sci. Nat. Mus. Civico Storia Nat. Milano 33(2): 8. 2005.

Acknowledgements

The authors are indebted to Prof. C. Stace, Dr S. Darbyshire and Prof. W. Greuter for their corrections and useful comments.

Addresses of the authors:

Dr Bruno Foggi, Dipartimento di Biologia Vegetale, Università degli Studi di Firenze, via G. La Pira, 4, I-50121 Firenze, Italy; e-mail: bruno.foggi@unifi.it

Prof. Dr Hildemar Scholz, Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin, Königin-Luise-Str. 6-8, D-14191 Berlin, Germany; e-mail: hischo@zedat.fu-berlin.de

Prof. Dr B. Valdés, Departamento de Biología Vegetal y Ecología, Facultad de Biología, Universidad de Sevilla, Avda. Reina Mercedes s/n, 41012 Sevilla, Spain; e-mail: bvaldes@us.es