

Independent Review Report, Reviewer 1

We appreciate very much for the valuable comments of our manuscript. These opinions help to improve academic rigor of our article. Based on your suggestions, we have responded to the comments one by one as listed below, which is also revised in our revised manuscript.

Minor:

1. Line 91 : ‘scientists’ to be replaced with ‘studies’

We appreciate it very much for the suggestion, and we have replaced ‘scientists’ with ‘studies’.

2. Line 93: Citations needed for definition

Thank you for the reminder, and the citations have been added. (Rao and Sivakumar, 1999; Kumar et al., 2009; Rao et al., 2015)

3. Line 183-195 : Quite repetitive writing. Rephrasing needed.

We have revised the text to address your concerns and it is much clearer now.

The revised text reads as follows: “The contribution of SHF to $\frac{\partial T}{\partial t}$ ranged from 0.11°C/5 day to 0.34°C/5 day (Figure 4), always playing a heating role. From April to early May, the contribution of SHF to temperature was around 0.3°C/5 day, indicating that SHF provided substantial thermal support for the increased temperature. By early June, the contribution of SHF to temperature dropped below 0.2°C/5 day. The effect of SHF on warm pool temperature can be divided into two parts that are caused by net surface heat flux (SHF_{Q_{net}}) and the penetrating shortwave radiation at the bottom of the mixed layer (SHF_{Q_{loss}}). In the upper layer of the mixed layer, the temperature change is related to the heat exchange between the surface seawater and the atmosphere, so it is related to net heat flux at the sea surface (SHF_{Q_{net}}). In the lower layer of the mixed layer, the heat flux that can penetrate the bottom of the mixed layer is mainly shortwave radiation. Therefore, the heat loss at the bottom of the mixed layer is mainly related to penetrative shortwave radiation (SHF_{Q_{loss}}). The contribution of $\frac{\partial T}{\partial t}$ gradually increased from 0.41°C/5 day to 0.50°C/5 day in April. It increased slowly but remained at a relatively high level. Then it gradually decreased to less than 0.2°C/5 day. Similar to $\frac{\partial T}{\partial t}$, $\frac{\partial T}{\partial t}$ also gradually increased to -0.17°C/5 day in April and then decreased.”

4. The paper is loosely written. At many places, it seems authors meant something else but have written something else.

We have now worked on both language and readability. We really hope that the language level have been substantially improved. Thanks so much for your useful comments.

5. Line 292 : based on a 0.5 sigma criterion .

Thank you for your suggestions, and we have changed the above as suggested.

6. Figure 13, and its caption : N should be capital In Niño and in all other places.

We are very sorry for our incorrect writing, and we have corrected the errors in the whole paper.

7. Line 328 : Please rewrite the description of IOD if needed in correct tense. This is grammatically incorret.

Thank you for pointing out the error, and we have revised it as: “The IOD is an inherent interannual climate mode in the TIO. During a positive phase, warm waters are pushed to the Western part of the Indian Ocean, while cold deep waters are brought up to the surface in the Eastern Indian Ocean. This pattern is reversed during the negative phase of the IOD (Saji et al., 1999; Kumar et al., 2020)”.

8. Rewrite section 5 with more conciseness and clarity.

Thanks for your comments. According to your suggestions, we have revised the summary and discussion of section 5 of the paper to make them more concise and clear.

Major :

1. Section 2.3 : Are the units consistent in this formulation of Heat budget? The units of ENT1 and ENT2 is degC/s. The units of ENT3 is deg C-m/s. Kindly check and clarify.

After careful checking, we confirmed that the units in this formulation of the heat budget were consistent ($^{\circ}\text{C}/\text{s}$). In our study, since the time interval of the SODA data set used in this paper is 5 days, in order to ensure the accuracy of calculation and the units are consistent, we converted the unit into $^{\circ}\text{C}/5$ days when calculating and drawing.

2. Line 139 : What is advection of mixed layer depth and how is it different from advection : ADV? Please explain. This is very confusing. What is tendency of MLD? Is it the tendency of how depth changes?

“Advection of mixed layer depth” refers to the horizontal variation of the mixed layer depth caused by the horizontal flow of seawater, with an emphasis on the distribution of the mixed layer depth in the horizontal direction. However, the advection term “ADV” refers to the horizontal variation of seawater temperature caused by the horizontal flow of seawater, with an emphasis on the distribution of seawater temperature in the horizontal direction. “tendency of MLD” refers to the trend of changes in mixed layer depth over time, which corresponds to variable

$ENT2 = -H \frac{\partial h}{\partial t} \frac{T - T_h}{h}$ in the formula (line 128), indicating the change in mixed layer temperature caused by local changes in mixed layer depth.

3. Section 2.3 Line154 ; Wh is the bottom of mixed layer? Is it the depth? I understand it is the vertical velocity at the base of the mixed layer from the calculation. However, please write it carefully. It is important for the readers to understand this.

Thank you for pointing this out. The reviewer is correct, and we have corrected it as “ W_h is the vertical velocity at the base of the mixed layer”.

4. Why do you need a dimensionless number H in the heat budget equation?

H is the dimensionless number in the vertical entrainment, which involves the temperature change caused by the local change of mixing depth ($\frac{\partial h}{\partial t}$). When $\frac{\partial h}{\partial t} > 0$, H is 1; When $\frac{\partial h}{\partial t} \leq 0$, H is 0. The calculation formula used here is consistent with that of Kurian and Vinayachandran (2007).

5. Figure 1 & Figure 2 : You define warm pool by 30deg C, but in figure 1, there is no cold pool after 26 May. How do you plot the maximum temperatures in figure 2, it is not clear what area did you choose for it? In Line 151, it is mentioned that area of warm pool has been chosen, but there is no warm pool on those dates.

Thank you for pointing out this problem in our manuscript. According to the revised content, we have redrawn Figure 1 and 2, and added a clear description of the definition: “This article defined a rectangle range of 55.25°E-77.25°E and 5.25°N-20.25°N based on the regional distribution of sea surface temperature in the Arabian Sea (As shown in Figure 1, marked with a dashed rectangle). ASWP was defined as the sea area within the rectangular range with a SST greater than 30°C” . Therefore, it can be seen that there are still small areas of sea temperature above 30°C along the western coast of the Indian Peninsula after May 26th (i.e., May 31st, June 5th). When calculating the highest temperature, we calculate the highest temperature of the sea area with SST greater than 30°C within the rectangular sea area. Corresponding modifications have also been made to Figure 1 and 2.




Figure 1: Climatological SST (1980–2016) in April–June, where the black contour is the 30°C contour. The dashed rectangle represents the selected warm pool range.




Figure 2: Climatological time series of the area and maximum temperature of ASWP, where the blue bars represent the warm pool area (km²) and the solid line represent the maximum warm pool temperature (°C).

6. The ASWP is defined historically in the SEAS, from 31 May onwards the pool seems to be shifting to extreme North of coast. Did you use a latitudinal boundary to define the warm pool, or have the authors taken June 5 contour of 30deg C to calculate area and temp of warm pool?

We apologize for the vague expression of the definition. The definition of ASWP in this article is the sea area with sea surface temperature greater than 30°C within the rectangular area of 55.25°E–77.25°E and 5.25°N–20.25°N. The delineation of this rectangular area is based on the spatial extent of the warm pool when it reaches its maximum area (May 11th). Following your suggestion, a box defining the area has

been added to Figure 1, and a detailed description of the ASWP definition has been added to the text.

7. Line 195: This is anyway clear from blue line in Figure 3.

Thanks for pointing it out. Since this conclusion has already been illustrated in Figure 3, it will not be repeated here at your suggestion.

8. Line 188 : How do authors say that Q_{net} is due to air sea interaction and Q_{loss} is due to SWR penetration. Q_{loss} is due to all factors of Heat flux – Sensible, latent, longwave etc and not only because of reduced SWR penetration. If the authors are too confident about SWR reduction, it should be explained more. Always : $Q = Q_{net} + Q_{loss}$, so saying the SWR penetrated less is not always correct.

We are very sorry for some confusing descriptions. As mentioned by the reviewer, strictly speaking, both Q_{net} and Q_{loss} are indeed related to all factors of heat flux (such as latent heat, sensible heat, longwave and shortwave radiation). In the upper layer of the mixed layer, the temperature change is related to the heat exchange between the surface seawater and the atmosphere, so it is related to net shortwave radiation, longwave radiation, sensible heat, and latent heat at the sea surface. In the lower layer of the mixed layer, the heat flux that can penetrate the bottom of the mixed layer is mainly shortwave radiation (Liu Zhiliang, 2007). Therefore, the heat loss at the bottom of the mixed layer is mainly related to penetrative shortwave radiation.

Therefore, in the calculation formula, $Q_{net} = \frac{Q_{sw} - Q_{lw} \pm Q_{sh} \pm Q_{lh}}{\rho_0 ch}$, $Q_{loss} = \frac{Q_{sw} [re^{-\frac{h}{\beta_R}} + (1-r)e^{-\frac{h}{\beta_B}}]}{\rho_0 ch}$. According to your suggestions, we have revised the paper to make

it more clear.

9. Line 200 onwards : Here SWR penetration is explained. It is not clear at what level have the authors taken SWR in their calculation. Is it at the TOA or Surface of the ocean?

SWR can affect the temperature change of the mixed layer through two aspects, one is the absorption of shortwave radiation in the ocean surface layer, and the other is the loss of shortwave radiation at the bottom of the mixed layer (Li et al., 2016). Therefore, this article reflects the calculation of SWR in two aspects: one is in Q_{net} , which calculates the net shortwave radiation absorbed by the ocean surface layer; the other is in Q_{loss} , which calculates the shortwave radiation that penetrates the bottom of the mixed layer. We corrected it clearly in the revised paper.

10. Is the MLD response to reduced SWR immediate? Or is there a difference in the rate of decay of MLD temperature due to SWR reduction?

Thank you for pointing this out. The revised it as follows: “In addition, although the variation of SST was roughly in line with the trend of SWR, the variation of SST had a lag time of about one month compared with SWR. This trend occurred because

seawater has a large specific heat capacity and there is a lag of one month in the variation of SST.” The results also show a one-month lag (Figure 3).




Figure 3: Monthly Variation of the net shortwave radiation at the sea surface (blue bars and lines; W/m²) and SST (red line; ° C) of the Arabian Sea.

11. Line 211 : what is certain lag?

Thank you for your comment. We’ve changed “ a certain lag ” to “ a lag of one month ”.

12. Line 220 : how does MLD inhibit penetration of SWR. I am not clear with this mechanism. The authors may explain please.

According to the formula $Q_{loss} = \frac{Q_{sw}[re^{\frac{h}{\beta_R}} + (1-r)e^{-\frac{h}{\beta_B}}]}{\rho_0 c h}$ and the spatial distribution

of Q_{loss} and MLD, it can be concluded that Q_{loss} is negatively correlated with MLD. Regarding the mechanism between SWR and MLD, the statement in this paper "the mixing layer greatly inhibited the penetration of shortwave radiation flux into the mixed layer" is inappropriate. The correct statement is that MLD is forced by penetrative shortwave radiation, and the impact of penetrative shortwave radiation on the depth of the mixed layer is controlled by wind speed and net shortwave radiation at the sea surface (Liu Zhiliang, 2007). Thank you for pointing this out. The correction has been made in revised paper.

13. Line 228 onwards : Comment no 1,2,3. I am not clear what each of those terms mean, it is not clearly defined as well. Kindly explain. I only understand ENT1.

The entrainment process in the ocean mixed layer is the product of the entrainment velocity (W_h) and the difference of temperature between the mean temperature in the mixed layer (T) with the temperature just below the mixed layer base (T_h).The entrainment is composed of three terms(Nogueira Neto et al., 2018):

1. the entrainment due to the vertical velocity ($W_h \frac{T-T_h}{h}$).

2. the entrainment due to the local tendency of the mixed layer depth ($H \frac{\partial h}{\partial t} \frac{T-T_h}{h}$). It refers to the trend of changes in mixed layer depth over time.

3. the entrainment due to "advection of the mixed layer depth" $((U \frac{\partial h}{\partial x} + V \frac{\partial h}{\partial y}) \frac{T-T_h}{h})$.

It refers to the horizontal variation of the mixed layer depth caused by the horizontal flow of seawater, with an emphasis on the distribution of the mixed layer depth in the horizontal direction.

14. Line 236 -238 : Do the authors mean that -Low wind speed made MLD more sensitive to heat flux? What is the meaning of this line? I understand MLD is always sensitive to heat flux, atmospheric conditions determine heat flux, the sensitivity is always there because MLD included the Sea surface which is always in contact with the atmosphere. I think the authors mean to write something else, but it is not clear. At the same time, how shallow MLD makes the upper sea water increase rapidly?

Thank you for pointing out this problem in our manuscript. I apologize for my ambiguous expression. Actually I mean that the low wind speeds result in shallower mixed layer, which in turn makes the seawater within the mixed layer more easily heated and sensitive to heat flux. Following your suggestion, the description has been revised as "In the development stage of the ASWP, the low wind speed made the mixed layer shallower, which made the seawater within the mixed layer more easily heated and thus more sensitive to heat flux". Therefore, during the development stage of ASWP, strong net heat flux and shallow mixed layer resulted in rapid warming of the upper sea water.

15. Line 240 onwards : Thermocline and Isothermal layer are not the same. They cannot be used interchangeably.

Thank you for pointing out this problem. We apologize for our error, and we've changed "thermocline" to "isothermal".

16. After reading through, it is not possible for me to point out each and every concern in the vertical entrainment term of heat budget. There is a diverse explanation ranging from penetration of SWR to Rossby waves. The timescales, space scales need a mention. This section needs a major overhaul and rewriting, it is very confusing currently. At the same time, the terms and mechanisms have to be explained carefully and clearly. This section has too many issues to point out and may be worked again.

Thank you for pointing it out. According to your suggestions, we have revised the contents as follows: "The vertical entrainment is closely related to the vertical thermohaline structure of seawater and the depth of the mixed layer. When the mixing of the upper seawater in terms of temperature and salinity is inconsistent, the isothermal layer and mixed layer will differ, and a barrier layer will be formed between the bottom of the mixed layer and the top of the thermocline (Sprintall and Tomczak, 1992). The barrier layer has the characteristics of a strong salinity gradient

and high gravitational stability, which makes it difficult to transport heat from top to bottom by mixing. On the one hand, a strong and stable salinity stratification can effectively inhibit the transfer of non-solar radiation flux to the interior of the ocean, which leads to the warming of the upper mixed layer. On the other hand, the existence of the barrier layer can inhibit the entry of the cold water of the thermocline into the mixed layer, which is not conducive to the exchange of heat, momentum, mass, and nutrients between the mixed layer and the thermocline (Pang Shanshan, 2021). Therefore, In the development stage of the ASWP, the low wind speed made the mixed layer shallower, which made the seawater within the mixed layer more easily heated and thus more sensitive to heat flux. Then the weak wind speed and the existence of anticyclonic circulation (Shankar et al., 2002) associated with the Lakshadweep High trapped the fresh water, contribute to the formation of the barrier layer, and maintain the ASWP by effectively inhibiting the vertical mixing of the upper ocean and causing a warming of the upper mixed layer. In the late stage of ASWP evolution, with the imminent outbreak of the summer monsoon, the stirring effect of wind started to strengthen, the mixing layer became deeper rapidly (Figure 6), and the cooling effect of entrainment was enhanced, which accelerated the decay of the ASWP (Liu Yanliang, 2013).”

17. Section 3.2.3 : The theory of EICC and NEC is explained. But how is it connected with what happens in April- June ASWP is not mentioned. Also, have the authors looked at the current systems during the months of analysis, because putting everything on NEC for the months of April-June may not be okay. The current changes its direction again in these months itself.

Thank you for pointing out this issue in our manuscript. Although the EICC and NEC input low-salinity water into the Arabian Sea from November to March, the weak wind speed and the anticyclonic circulation (Shankar et al., 2002) associated with the Lakshadweep High (Bruce et al., 1994), restrict the low-salinity water within the Arabian Sea and maintain it until May. By June, the summer monsoon is about to break out, causing changes in the surface wind field and surface flow field, gradually spreading the low-salinity water and weakening the salinity stratification.

18. The authors write at Lines 158-159 that they will explain the mechanism of slow decay, however, I failed to find a clear explanation of the same. The authors may end the section with a small paragraph explaining exactly the mechanism of slow decay.

Thank you for the above suggestion. Following your suggestion, we have added a new paragraph at the end of the section 3.2 to explain and summarize the mechanism of ASWP's decay. The new paragraph reads as follows: “In the decline stage of ASWP, the rapid decrease in shortwave radiation flux weakens the heating effect of the sea surface heat flux forcing, while the strengthening of mixing leads to an increase in vertical cooling. The combined effect of these two factors results in a sharp decrease in temperature, leading to the rapid decline of the warm pool.”

19. Section 4.1 : Line 282 : how do authors know that? Citations?

The years of weak warm pools correspond precisely to the years when the Indian Ocean Basin Mode (IOB) is in a negative phase (Guo et al., 2018), during which sea temperatures across the entire Indian Ocean are anomalously cold. Relevant citations have been added to the paper.

Guo, F., Liu, Q., Yang, J., and Fan, L.: Three types of Indian Ocean Basin modes, *Clim. Dyn.*, 51, 4357-4370, 10.1007/s00382-017-3676-z, 2018.

20. Why in strong ASWP years is the average maximum temperature also high?

From Figure 4, it can be seen that the warm pool area and maximum temperature are highest during years of strong ASWP. As analyzed in Section 4.2, this is because years of strong ASWP correspond to a positive phase of IOB, during which the entire Indian Ocean is warming. As a part of the Indian Ocean, the sea temperature in the Arabian Sea also increases, resulting in a higher maximum temperature.




Figure 4: Variation of the area and maximum temperature of different types of ASWP, where the bars represent the area of the warm pool (km²), the solid lines represent the maximum temperature of the warm pool (°C), red represents weak ASWP, blue represents normal ASWP, and yellow represents strong ASWP.

21. What is Climate state averaging?

This refers to the diagnostic results of the climatology from 1980 to 2016. Thank you for pointing out and we have replaced 'climate state averaging' with 'the climatology'.

22. How do you define warming phase and cooling phase? Do you do it separately for each year? Or is there any other technique being used.

The paper analyzes the warming and cooling phases of the climate state (averaging the diagnostic results for different years on the same day) based on the positive and negative values of $\frac{\partial T}{\partial t}$. When $\frac{\partial T}{\partial t} > 0$, it represents a warming phase, while

$\frac{\partial T}{\partial t} < 0$ represents a cooling phase. Since the evolution time of ASWP for different years is different, only the climatically diagnostic results are analyzed here.

23. Line 300 : No clear meaning. I cannot understand what authors mean.

Thank you for pointing this out. The revised text reads as follows: “However, regardless of the year, the contribution of SHF to the temperature change was the largest (78.5% to 81.5%), and the contributions of ENT and ADV were relatively small. Among them, the weaker warm pool in weak ASWP years may be related to the greater cooling effect of ENT at this time (19.7%). In the cooling phase, the contribution of SHF decreased (31.9% to 38.3%), the cooling effect of ENT increased (52.2% to 54.2%), and the contribution of ADV increased slightly (9.5% to 14.6%). SHF and ENT together dominated the temperature change of the ASWP. The larger (smaller) warm pool in strong (weak) ASWP years may be related to the larger (smaller) heating effect of SHF, and the smaller (larger) cooling effect of ENT and ADV.”

24. Section 4.2: Which data has been used for this? How much is the variance explained by each mode can be mentioned in the figure 12 itself. Please mention. The text seems vague for variance explained.

The temperature data used in this paper is from the SODA3.7.2 dataset, and EOF analysis was performed on the temperature in the Arabian Sea from April to June respectively. Thank you for your suggestion, the variance contribution of each mode has been added to the figure 5.




Figure 5: EOF1 (the first column), EOF2 (the second column), and PC1 and PC2 (the third column) of SST anomalies in the Arabian Sea for April–June.

25. Line 313-314 : What is IOB consistent mode? Citation please.. Line 313 is not clear, rewrite.

This refers to the Indian Ocean Basin mode (IOB) (Guo et al., 2018). We have corrected it in the paper.

26. What is B in IOB in Line 318?

Indian Ocean Basin mode (IOB).

27. Probably this is the place (after Line 316/317) where Line 282 should be mentioned rather than there.

Thanks for pointing it out, we have corrected it in the paper.

28. Why do you show PC2 if variance explained is less than 10%?

Thank you for pointing it out. The variance contribution of PC2 from April to June were 12.71%, 12.51% and 11.52%, respectively. We chose to show PC2 in this paper because the variance contribution of PC2 from April to June has passed the significance test, indicating that the spatial and temporal functions of the second mode are valuable.

29. Line 327-328 : Are these the years in which your PC also peaks (Line315) or in which ASWP is found to be stronger (Line 279).

These years are the years when the area of the ASWP was larger. We have corrected it to clearer in the paper.

30. Line 333: is the correlation significant?

Correlation coefficient (r) = 0.57, $P=4.6*10^{-4}$, indicating that the correlation is significant.

31. Line 345: What is IOWP?

Indian Ocean warm pool.

32. Any citation for Line 345-346.

Thank you for the reminder, the citation has been added(Lau and Nath, 2003).

Lau, N.-C. and Nath, M. J.: Atmosphere–Ocean Variations in the Indo-Pacific Sector during ENSO Episodes, *J. Clim.*, 16, 3-20, [https://doi.org/10.1175/1520-0442\(2003\)016<0003:AOVITI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0003:AOVITI>2.0.CO;2), 2003.

33. What is the correlation value of PC1 of ASWP with Nino? Why do you choose to analyse ENSO rather than IOD in lines 348 onwards?

The correlation value between PC1 of ASWP and ENSO ranges from 0.61 to 0.72 ($P<0.01$), when PC1 lags ENSO for 5-7 months. As a part of the Indian Ocean, the Arabian Sea is not only affected by IOB and IOD in the Indian Ocean, but also by ENSO in the Pacific Ocean. Previous studies also show that ENSO has a significant regulatory effect on SST in the tropical Indian Ocean. Therefore, the relationship between ASWP, IOB and IOD is analyzed in Section 4.2. The relationship between ASWP and ENSO is also analyzed later.

34. Line 364 :Vertical entrainment is completely different from what has been used here. Please use a different word. (Again, please note comment no 16.)

Thank you for pointing it out. We have corrected it in the paper.

Thanks again for the suggestive comments of the reviewer!

Independent Review Report, Reviewer 2

We appreciate very much for the valuable comments of our manuscript. These opinions help to improve academic rigor of our article. Based on your suggestions, we have responded the comments one by one as listed below, which is also revised in our revised manuscript.

Review comments:

1. Ln 12: “Indian Ocean summer monsoon.” - Please change it to "Indian summer monsoon".

We appreciate it very much for the suggestion, and we have replaced ‘Indian Ocean summer monsoon’ with ‘Indian summer monsoon’.

2. Ln 26: “The ASWP was more significant (insignificant) in the following year before the summer monsoon after an El Niño (La Niña) event that peaked in the previous winter.” - Please rewrite the sentence for better clarity.

Thank you for pointing this out. The revised text reads as follows: “When the El Niño (La Niña) event peaked in the winter of the previous year, the ASWP that occurred before the summer monsoon was more significant (insignificant) in the following year.” (line 26-28)

3. Ln 33-38: Please provide appropriate references for example, <https://rmets.onlinelibrary.wiley.com/doi/full/10.1002/asl.596> <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49712555503>

Thank you for the reminder, and the citations have been added. (Rao and Sivakumar, 1999; Kurian and Vinayachandran, 2007; Nagamani et al., 2016). The added references can be found at the bottom of this reply letter.

4. Ln 41: “(SST >30.5°C) in the southeastern AS” - please specify the aswp area averaged to get this value. Because of the large variability in the ASWP area, the SST values would change based on the area considered.

Thank you for your reminding, the value here (30.5°C) is wrong, after careful check, it has been corrected to 30.8°C. And the area of the warm pool is was located between the latitudes 10°N to 13°N and longitudes 67°E to 72°E. According to your suggestion, the area range has been added to the revised paper.

5. Ln 42: “Krishnamurt et al. (1988)” - please check the author name carefully.

Thank you for pointing this out, and we have corrected ‘Krishnamurt’ with ‘Krishnamurti’.

6. Ln 43: “Joseph (1990) also suggested that SST maxima occurred one week before the summer monsoon outbreak in the southeastern AS during 1961–1972.” - It is a regular phenomena to have higher SSTs during the summer time due to high insolation and transport of warm coastal waters towards southeastern AS. The question here is whether and how these higher SST values are associated with phase of the ASWP and are consistant with the size and duration of the ASWP.

Thank you for pointing out this problem in our manuscript. I apologize for my

ambiguous expression. Actually I mean that before the onset of the summer monsoon, the warmest area of the warm pool of the tropical oceans is centred over southeast Arabian Sea. It is emphasized that the highest sea surface temperature occurs in the southeastern Arabian Sea. Following your suggestion, the description has been revised in the paper as “ Joseph (1990) also suggested that before the onset of the summer monsoon, the warmest area of the warm pool of the tropical oceans is centred over southeast AS.” (line 44-46)

7. Ln 48: “Arabian Sea” - Please be consistent with using the short and full form of the words. The author may consider using AS here as it was already specified in the above paragraph.

Thank you for pointing this out. After defining the short form of the word, we have replaced 'Arabian Sea' with 'AS' in the subsequent sections of the paper.

8. Ln 49: “In the second phase,” - Please describe the phase. The sentence is inconsistent.

The second phase refers to the cooling phase, which has been changed to “cooling phase” in order to correspond to the “warming stage” in the previous sentence.

9. Ln 51: “He found that” - it is a good practice to avoid gender classification, instead, one can use "the author".

Thank you for your suggestions, and we have changed the above as suggested.

10. Ln 57: “They found that the Western Ghats' orographic impact decreased wind speed in the Arabian Sea's southeast and, consequently, latent heat loss, resulting in a positive heat flux into the ocean.” - Please rephrase the sentence.

Thank you for pointing it out. According to your suggestions, we have revised the contents as follows: “They found that the orographic impact of the West Ghats reduced wind speed in the southeastern AS, thereby reducing latent heat loss and leading to positive heat flux entering the ocean.” (line 59-60)

11. Ln 60: “Sabu and Revichandran (2011)” - Please check the spellings of author names carefully.

After careful checking, we confirmed that the spellings of author names are correct.

12. Ln 62: “intermonsoon “ - a new term?

Intermonsoon, also known as inter-monsoon, refers to the interval between two monsoons. In India, there are two monsoons: the southwest monsoon and the northeast monsoon. Inbetween are two periods referred to as the intermonsoons: spring intermonsoon (March/April) and autumn intermonsoon (October/November).

13. Ln 66: “in Peninsular India” - Please change it to "tropical Indian Ocean".

Thank you for pointing this out, and we have replaced ‘Peninsular India’ with ‘tropical Indian Ocean’.

14. Ln 69: “According to Lau (2000) and Chowdary et al. (2007)” - How does the mechanism explained by Lau (2000) and Chowdary (2007) affect the ASWP as this is mostly confined to eastern TIO, whereas ASWP centered at the central TIO?

According to Lau and Chowdary et al, the ENSO not only affects precipitation

and wind speed in the eastern tropical Indian Ocean, but also heats the entire tropical Indian Ocean by affecting shortwave radiation and latent heat flux, thereby affecting the ASWP.

15. Ln 73: “The response of the warm pool intensity to ENSO does not reach its peak until about 5 months after ENSO peaks.” - Please provide a reference.

Thank you for the reminder, the citation has been added (Lau and Nath, 2003).

Lau, N.-C. and Nath, M. J.: Atmosphere–Ocean Variations in the Indo-Pacific Sector during ENSO Episodes, *J. Clim.*, 16, 3-20, [https://doi.org/10.1175/1520-0442\(2003\)016<0003:AOVITI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0003:AOVITI>2.0.CO;2), 2003.

16. Ln 82: “IOD was more significant and persistent during the years of cooccurrence, and it was characterized by both eastern cooling and western warming.” - Please provide a reference.

Thank you for the reminder, the citation has been added (Chowdary and Gnanaseelan, 2007).

Chowdary, J. S. and Gnanaseelan, C.: Basin-wide warming of the Indian Ocean during El Niño and Indian Ocean dipole years, *Int J Climatol.*, 27, 1421-1438, <https://doi.org/10.1002/joc.1482>, 2007.

17. Ln 90-92: - Please rewrite the sentence for better clarity.

Thank you for pointing it out. According to your suggestions, we have revised the contents as follows: “Although the effects between the large-scale modes (such as ENSO, IOD) and the Indo-Pacific warm pool have been studied, few studies have explored the relationship between various large-scale modes and smaller-scale seas (such as the AS).” (line 93-95)

18. Ln 109: “ JRA-55, the forcing field used in SODA 3.7.2,” - Please provide appropriate citation.

Thank you for the reminder, the citation has been added (Carton et al., 2018).

Carton, J.A., G.A. Chepurin, and L. Chen (2018), SODA3: a new ocean climate reanalysis, *J. Climate*, 31, 6967-6983, <https://doi.org/10.1175/JCLI-D-18-0149.1>

19. Ln 114: “ The data time period is from 1958 to the present.” - Please provide available spatial resolution information also.

Thank you for pointing it out. According to your suggestions, we have revised the contents as follows: “The data time period is from 1958 to the present, with a spatial resolution of $0.5625^\circ \times 0.5625^\circ$.” (line 117-118)

20. Ln 114-115: “day-by-day” - Please change it to "daily".

Thank you for pointing this out, and we have replaced ‘day-by-day’ with ‘daily’.

21. Ln 131: “ $\partial T / \partial x$ and $\partial T / \partial y$ represent the latitudinal and longitudinal spatial variation of mixed layer temperature, respectively.” - isn't it opposite? Please change the wording to zonal and meridional variation, which makes it unambiguous.

Thank you for pointing this out, and we have replaced ‘the latitudinal and longitudinal spatial variation’ with ‘the zonal and meridional variation’.

22. Ln 135: “ In this paper, the mixed layer depth has been defined as the depth at which the seawater is 0.03 kg/m³ higher than the surface density.” - Please provide reference for the MLD criteria considered and justify the same. Please

check <https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2004JC002378>.

Sorry, the definition of the mixed layer was written incorrectly here. In the revised manuscript, we select the criterion to be a density difference of 0.03 kg/m^3 between the reference depth (10 m) and the bottom of the mixed layer. This definition is consistent with De Boyer Montégut et al (De Boyer Montégut et al., 2004; Jofia et al., 2023). The reference depth is therefore chosen to avoid the diurnal cycle of the mixing layer. According to Brainerd et al.(1995), a value of 0.1 kg/m^3 sometimes yielded the depth of the main thermocline, in the case of fossil layers for example, and a value of 0.05 kg/m^3 often falls within the seasonal thermocline rather than at its top. Therefore a threshold of 0.03 kg/m^3 emerged as the appropriate value for the density criterion (De Boyer Montégut et al., 2004).

23. Ln 136: “ T_h is the temperature at the bottom 10 m of the mixed layer;” - Not clear. please elaborate.

Thank you for pointing this out. We have corrected it as “ T_h is the temperature at a depth of 10 m below the bottom of the mixed layer”. (line 142)

24. Ln 141: “This study also used empirical orthogonal function (EOF) analysis to separate the spatial and temporal characteristics of the ASWP and study the relationship between the ASWP and IOD/ENSO using lead lag correlation.” - Please cite any earlier studies done using the similar approach.

Li (2017) studied the mechanism of cycle process between the western Pacific warm pool and ENSO by using empirical orthogonal function analysis. Kim (2012) explore the cause-and-effect relations between ENSO and warm pool variations by calculating the lead-lag correlations. The analysis methods used in this paper is similar to that of Li et al.(Kim et al., 2012; Li Xiaohui, 2017), but we focus on analyzing the results every 5 days from April to June in order to gain some new insights. The relevant references have been added to the revised paper.

25. Figure 1: Please modify the figure 1 and 2 titles by including the details of the sub-plot titles/labels and data used. I don't see the need of including June month's climatology because it is very well known than onset of southwest monsoon in the 1st week of June eventually leads to ASWP decay. You may cite a relevant work on the same in the text. Instead authors can show the March month's conditions.

Thank you for pointing this out. We have corrected it as “ Figure 1: Every 5-days evolution of SST (unit:°C) in April–June, where the black contour is the 30°C contour. The dashed rectangle represents the selected range of Arabian sea warm pool (55.25°E - 77.25°E , 5.25°N - 20.25°N). The data is based on gridded dataset at the period of 1980-2016 from SODA v3.7.2. (<https://www2.atmos.umd.edu/~ocean/>)”.

26. Figure 2: Please mention in the figure title or in the text, how the area was calculated.

Thank you for pointing this out. We have corrected it as “ Figure 2: Time series of the area and maximum temperature of climatological ASWP, where the blue bars represent the warm pool area (km^2) and the solid line represent the maximum warm pool temperature ($^\circ\text{C}$). The area of ASWP was the sea area within the rectangular range with a SST greater than 30°C . The maximum temperature was the highest

temperature of the sea area with SST greater than 30°C within the rectangular sea area”.

27. Ln 151: “ its area and maximum temperature were calculated” - Please specify how the area and maximum temperature calculation are made.

Thank you for pointing out this problem in our manuscript. According to the revised content, we have redrawn Figure 1 and 2, and added a clear description of the definition: “This paper defined a rectangle range of 55.25°E-77.25°E and 5.25°N-20.25°N based on the distribution of sea surface temperature in the AS (As shown in Figure 1, marked with a dashed rectangle). ASWP was defined as the sea area within the rectangular range with SST greater than 30°C”. (line 164-166) When calculating the maximum temperature, we calculate the highest temperature of the sea area with SST greater than 30°C within the rectangular sea area. Corresponding modifications have also been made to Figure 1 and 2.




Figure 1: Every 5-days evolution of SST (unit:°C) in April–June, where the black contour is the 30°C contour. The dashed rectangle represents the selected range of Arabian sea warm pool (55.25°E-77.25°E, 5.25°N-20.25°N). The data is based on gridded dataset at the period of 1980-2016 from SODA v3.7.2. (<https://www2.atmos.umd.edu/~ocean/>)




Figure 2: Time series of the area and maximum temperature of climatological ASWP, where the blue bars represent the warm pool area (km²) and the solid line represent the maximum warm pool temperature (°C). The area of ASWP was the sea area within the rectangular range with a SST greater than 30°C. The maximum temperature was the highest temperature of the sea area with SST greater than 30°C within the rectangular sea area.

28. Ln 154-155: “when the warm pools were in a strong stage.” - Please replace "strong" with "matured" or "developed".

Thank you for pointing this out, and we have replaced ‘strong’ with ‘matured’.

29. Ln 155: “Thereafter, the warm pools decayed rapidly and disappeared almost completely in early June.” - Please provide a citation.

Thank you for the reminder, the citation has been added (Rao et al., 2015).

Rao, R. R., Jitendra, V., GirishKumar, M. S., Ravichandran, M., and Ramakrishna, S. S. V. S.: Interannual variability of the Arabian Sea Warm Pool: observations and governing mechanisms, *Clim. Dyn.*, 44, 2119-2136, 10.1007/s00382-014-2243-0, 2015.

30. Ln 155-158: - Please rewrite/rephrase the sentence for clarity.

Thank you for pointing this out. We have corrected it as “ From Figure 2, it can be seen that the development and decay processes of ASWP are not symmetrical. From the formation to the mature stage of the ASWP, it takes approximately 1.5 months (6 weeks). While from the mature to disappear completely, it only takes 3 weeks. The rate of decline is twice than that of the development. The mechanism influencing this asymmetry in development and decay is discussed below.” (line 172-175)

31. Ln 167: “Throughout the evolution of ASWP, the sums of SHF, ADV, and ENT were basically consistent with the trend of mixed layer temperature.” - This is a strong statement. Authors can notice that, temperature change due to advection is almost constant and only SHF and ENT terms cause temperature to

decrease. Authors can draw another line with only SHF and ENT to check their impact. Notice that their trends are not similar as stated by authors. Please give their trend values in the figure.

Based on your suggestion, we have added a new line with only SHF and ENT in Figure 3. The results indicate that the variation in temperature are mainly related to SHF and ENT, while the impact of ADV is almost constant and minimal. Due to the fact that the calculation in the figure already shows the temperature variation and the contributions of different processes to temperature variation, there is no obvious physical significance in recalculating the trend values of the lines. Moreover, there are many lines in the graph, and adding trend may make the image chaotic and complex. The relevant figure and statements have been revised in the paper.




Figure 3: Contribution of different processes to the temperature variation of the warm pool mixed layer in the Arabian Sea ($^{\circ}\text{C}/5 \text{ day}$), where $\frac{\partial T}{\partial t}$ (solid red line) is the temperature variation of the warm pool mixed layer with time, SHF (solid blue line) represents the sea surface heat flux forcing, ADV (solid green line) represents the horizontal advection, ENT (solid black line) represents the vertical entrainment, and R (black dashed line) represents the residual. The red dashde line represents the sum of SHF, ADV and ENT. The red dashde line represents the sum of SHF and ENT.

32. Ln 172: “mixed layer SST” - Can't be seen from the figure. I am afraid that it is not mixed layer SST, but mixed layer temperature. Please do the necessary corrections elsewhere.

Thank you for pointing this out, and we have replaced ‘mixed layer SST’ with ‘mixed layer temperature’.

33. Ln 173-157: - Not clear whether authors are discussing about SST or mixed layer temperature. Please take care of wording.

Thank you for pointing out. What we want to discuss here is the mixed layer temperature, and the relevant expressions have been corrected.

34. Ln 181: “net sea air heat flux” - Please check.

Thank you for pointing this out, and we have replaced ‘the net sea air heat flux’ with ‘the net surface heat flux’.

35. Ln 194-196: - This will be true most of the time in the tropical oceans with barotropic weather being a dominated phenomenon. There are exceptions like tropical cyclones during which the loss will be higher than the SHF_net.

We agree with the reviewer's opinion. However, in our article, we are discussing the analysis results of climatology, and the results of climatology do differ from those under special weather conditions. Thank you for pointing out, and we have made revisions in the paper based on the above comments.

36. Ln 200-203: - Not true always. A persistent cloud cover may increase the longwave radiation flux. Please check and cite the following articles. <https://spj.science.org/doi/full/10.34133/olar.0003> <https://link.springer.com/article/10.1007/s42452-019-1172-2> Please check for contribution of downwelling longwave radiation also.

Thank you for pointing out. In our article, we discuss the analysis results of climatology. Based on your suggestion, we have plotted the average long wave radiation, short wave radiation, sensible heat, and latent heat fluxes of ASWP (Figure 4). It can be seen that in the Arabian Sea, although the downward long wave radiation flux is stronger than the downward short wave radiation flux, the net long wave radiation is upward and smaller due to the upward long wave radiation flux being greater than the downward long wave radiation flux. Due to the small amount of upward shortwave radiation, the net shortwave radiation is downward and larger, indicating that the ocean is more affected by shortwave radiation flux at this time.




Fig. 4. The average heat fluxes of ASWP. The blue solid line represents the net long wave radiation flux, the orange solid line represents the net short wave radiation flux,

the yellow solid line represents the latent heat flux, and the purple solid line represents the sensible heat flux. The dashed line represents the downward radiation flux, while the dotted line represents the upward radiation flux.

37. Ln 209: “variation of SST was roughly” - SST or mixed layer temperature? Please check and modify elsewhere in the following text.

Thank you for pointing out, and here refers to the mixed layer temperature. The relevant statements have been revised in the paper.

38. Ln 217-221: Authors can discuss the impact of cloud cover and the entrainment and vertical velocity increasing cool subsurface water to change the mixed layer temperature. Here comes the impact of the MLD criteria one would choose.

Thank you for your suggestion. The impact of cloud cover on the mixed layer temperature has been analyzed in the second paragraph of 3.2.1, and the cooling effects of entrainment and subsurface water have been analyzed in section 3.3.2.

39. Ln 231: “SST” - mixed layer temperature?

Thank you for pointing this out, and we have replaced ‘SST’ with ‘mixed layer temperature’.

40. Ln 244: “non-solar radiation flux” - Authors may elaborate what do they mean by "non-solar radiation flux".

Thank you for pointing out. The expression of 'non solar radiation flux' is inappropriate, and has been changed to 'non-shortwave radiation' in the revised paper.

41. Ln 264-267: - Can't understand what do author's want to convey. Please rewrite and provide reason in this section to support using more appropriate citations.

Thank you for pointing out this problem in our manuscript. I apologize for my ambiguous expression. We have revised the contents as follows: “During November and December, there is an equatorward-flowing East India Coastal Current (EICC) off the east coast of India. This current flows around Sri Lanka and continues as a poleward-flowing West India Coastal Current off the west coast of India (Shetye et al., 1996; Rao and Sivakumar, 1999; Mukhopadhyay et al., 2020).”. (line 275-277)

42. Figure 9: Not clear whether area calculated is monthly or pentade? Please provide details.

The area calculated in figure 9 is daily mean every 5 days. Thank you for your suggestions, and we have added detailed instructions in Figure 9.

43. Figure 10: Not clear if this is a pentade mean? Please give more details on the figure to make it clear.

The area calculated in figure 10 is daily mean every 5 days. Thank you for your suggestions, and we have added detailed instructions in Figure 10.

44. Ln 311: “The first mode variance contribution for April, May, and June was all greater than 50%.” - Where is it shown?

Thank you for your suggestion, the variance contribution of each mode has been added to the figure 5.




Figure 5: EOF1 (the first column), EOF2 (the second column), and PC1 and PC2 (the third column) of SST anomalies in the Arabian Sea for April–June.

45. Ln 315: “1987, 1991, 1998, 2010, and 2016” - How do you explain the change in peaks for June month? Similarly explain the negative peaks in PC.

The occurrence of tropical cyclones can affect changes in mixed layer temperature significantly. There are few tropical cyclones in April and May, so the temperature is mainly affected by IOB at this time. June is the peak period for the occurrence of tropical cyclones in the Arabian Sea (Zihan et al., 2023), so the temperature in June is more affected by tropical cyclones. Due to the complex interannual variations of tropical cyclones, which are not always consistent with the changes in IOB, the peaks of PC1 in June differs from the peaks in April and May.

46. Ln 320-322: - Not clear what do the author's mean here?

Thank you for pointing out, and I apologize for my ambiguous expression. Following your suggestion, the description has been revised as: “The variance contribution of the second mode (10.71%, 12.51% and 11.52%) was smaller than the first mode. The spatial distribution patterns vary each month, but there are at least two antiphase extreme centers. The amplitude of PC2 change every month is smaller than the amplitude of PC1.” (line 348-350)

47. Ln 325: “The IOB is the first mode of the Indian Ocean and characterized as a consistent warming or cooling at the Indian Ocean basin scale (Xie et al., 2009).”- Please rewrite. Can't understand what do author's want to convey?

Thank you for pointing it out. We have revised the contents as follows: “A basin-scale warming/cooling is the leading mode of tropical Indian Ocean temperature variability on interannual time scales. It peaks in late winter and persists into the

following spring and summer (Klein et al., 1999; Li et al., 2008). ” (line 304-305)

48. Ln 331: “The left panel of Figure 13 shows that the PC-1 of ASWP was positively correlated with the IOD most of the time.” - It is a strong statement. Can't be seen in the figure. There are many undulation in the correlations which contradicts this statement.

Thank you for pointing out. Based on and in conjunction with your suggestion No. 50, the content of this section has been revised as follow: “The left panel of Figure 13 shows that the highest correlation coefficient between the PC1 and IOD reached 0.56 ($p < 0.01$), indicating that the ASWP had a significant correlation with the IOD. There are three peaks in the lead-lag correlation between PC1 and IOD. The first peak appears at -5, -7, and -8 months ($r = 0.48-0.56$), indicating that the mixed layer temperature lags behind the IOD for 5-8 months. It also means that the temperature from April to June has a higher correlation with the previous winter's IOD. This is consistent with that the IOD peaks in winter (Li et al., 2008). The second peak appears at -2, -1, and 0 months ($r = 0.36-0.49$), indicating that the IOD can also regulate the temperature in near real-time. The third peak appears at 7 months ($r = 0.14-0.34$), indicating that the temperature is not only regulated by IOD, but can also affect the IOD mode, although the effect is relatively small. This is consistent with the research results of Li et al (2008). ” (line 359-366)

49. Ln 336-338: “ You can cite the most recent research on this topic. Please refer following articles. <https://www.nature.com/articles/s41598-023-32840-w> <https://www.nature.com/articles/s41612-020-0127-z> <https://www.nature.com/articles/s41598-018-30552-0>

Thank you for your suggestion. The latest research articles have been cited in the revised paper.

50. Ln 342: “It can be seen that the ASWP was most correlated with a lag of 5–7 months in the Niño3.4 index, indicating that it was modulated by the ENSO.” - From figure 12 and 13, the ENSO impact is stronger than IOD and leading the correlations. How can authors explain this. Also, please describe 3 peaks in the IOD correlation in relation to a smooth decay in ENSO correlation.

From Figure 13, it can be seen that there are three peaks in the lead-lag correlation between PC1 and IOD. The first peak appears at -5, -7, and -8 months ($r = 0.48-0.56$), indicating that the mixed layer temperature lags behind the IOD for 5-8 months. It also means that the temperature from April to June has a higher correlation with the previous winter's IOD. This is consistent with that the IOD peaks in winter (Li et al., 2008). The second peak appears at -2, -1, and 0 months ($r = 0.36-0.49$), indicating that the IOD can also regulate the temperature in near real-time. The third peak appears at 7 months ($r = 0.14-0.34$), indicating that the temperature is not only regulated by IOD, but can also affect the IOD mode, although the effect is relatively small. This is consistent with the research results of Li et al (2008). The correlation coefficient between PC1 and the Niño3.4 index is highest at a lag of 5-7 months ($r = 0.61-0.72$), and sharply decreases after -2 months. This indicates that the temperature in the Arabian Sea is more regulated by ENSO. On the one hand, ENSO can directly affect the temperature changes in the Arabian Sea through atmospheric

bridges (Lau and Nath, 2000; Chowdary and Gnanaseelan, 2007); On the other hand, ENSO can indirectly regulate the temperature changes by influencing the IOD (Behera et al., 2006). So, the correlation coefficient between ENSO index and PC1 is higher than that of IOD, and the impact is stronger. According to your suggestion, the relevant content has been added to the revised paper.

51. Ln 349: “Can changes in the ENSO affect the role of different processes in the evolution of the ASWP? “ - Interesting. can you also include similar analysis with positive and negative IOD years?

Thank you for your suggestion. In fact, we have conducted similar analysis with positive and negative IOD years, but the calculation results indicate that there is not much difference, so we did not include it in the paper for discussion.

52. Ln 354: “SST warmed” - still not clear. Is it SST or mixed layer temperature?

Thank you for pointing out, and here refers to the mixed layer temperature. The relevant statements have been revised in the paper.

53. Ln 354-356: “still not clear. Is it SST or mixed layer temperature?” - Please explain this in detail with appropriate citations?

Thank you for pointing out, and here refers to the mixed layer temperature. The effect of vertical entrainment was enhanced may be related that the convective organization becomes more frequent throughout the tropics, and especially over warmer ocean waters, during El Niño (Sullivan et al., 2019).

54. Ln 366: “SHFQnet was related to SWR,” - Not entirely true. Please check and modify.

Thank you for your suggestion. This has been modified to ' SHF_{Qnet} was mainly related to shortwave radiation'.

55. Ln 368: “SHFQloss was related to the depth of the mixed layer and increases with the onset of the summer monsoon.” - Also on the stratification/properties of the subsurface water including halocline and thermocline variability.

Thank you for your suggestion. Due to the word limit of the paper, the conclusion here is relatively brief, and a more detailed analysis is provided in Chapter 3.2.

56. Ln 382: “which is consistent with the strength of the IOWP peaking about five months after the ENSO peaked.” - Please give proper citations.

Thank you for the reminder, the citation has been added. (Lau and Nath, 2003)

Thanks again for the suggestive comments of the reviewer!

Reference, Reviewer 1:

- Bruce, J. G., Johnson, D. R., and Kindle, J. C.: Evidence for eddy formation in the eastern Arabian Sea during the northeast monsoon, *J. Geophys. Res.: Oceans.*, 99, 7651-7664, <https://doi.org/10.1029/94JC00035>, 1994.
- Guo, F., Liu, Q., Yang, J., and Fan, L.: Three types of Indian Ocean Basin modes, *Clim. Dyn.*, 51, 4357-4370, 10.1007/s00382-017-3676-z, 2018.
- Kumar, P., Hamlington, B., Cheon, S.-H., Han, W., and Thompson, P.: 20th Century Multivariate Indian Ocean Regional Sea Level Reconstruction, *J. Geophys. Res.: Oceans.*, 125, e2020JC016270, <https://doi.org/10.1029/2020JC016270>, 2020.
- Kumar, P. V. H., Joshi, M., Sanilkumar, K. V., Rao, A. D., Anand, P., Kumar, K. M. A., and Rao, C. V. K. P.: Growth and decay of the Arabian Sea mini warm pool during May 2000: Observations and simulations, 528-540, <https://doi.org/10.1016/j.dsr.2008.12.004>,
- Kurian, J. and Vinayachandran, P. N.: Mechanisms of formation of the Arabian Sea mini warm pool in a high-resolution Ocean General Circulation Model, *J. Geophys. Res.: Oceans.*, 112, <https://doi.org/10.1029/2006JC003631>, 2007.
- Lau, N.-C. and Nath, M. J.: Atmosphere–Ocean Variations in the Indo-Pacific Sector during ENSO Episodes, *J. Clim.*, 16, 3-20, [https://doi.org/10.1175/1520-0442\(2003\)016<0003:AOVITI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0003:AOVITI>2.0.CO;2), 2003.
- Li, Y., Han, W., Wang, W., and Ravichandran, M.: Intraseasonal Variability of SST and Precipitation in the Arabian Sea during the Indian Summer Monsoon: Impact of Ocean Mixed Layer Depth, *J. Clim.*, 29, 7889-7910, <https://doi.org/10.1175/JCLI-D-16-0238.1>, 2016.
- Liu Yanliang, Y. W., Li Kuiping: Mixed layer heat budget in Bay of Bengal : Mechanism of the generation and decay of spring warm pool, *Acta Oceanol. Sin.*, 35, 1-8, 2013.
- Liu Zhiliang, C. Z.: the effects of penetration radiation and salinity on mixed layer depth, *Transactions of Oceanology and Limnology*, 2, 19-25, 2007.
- Nogueira Neto, A. V., Giordani, H., Caniaux, G., and Araujo, M.: Seasonal and Interannual Mixed-Layer Heat Budget Variability in the Western Tropical Atlantic From Argo Floats (2007–2012), *J. Geophys. Res.: Oceans.*, 123, 5298-5322, <https://doi.org/10.1029/2017JC013436>, 2018.
- PANG Shanshan, W. X., LIU Hailong, SHAO Caixia: Multi-Scale Variations of Barrier Layer in the Tropical Ocean and Its Impacts on Air-Sea Interaction : A Review, *Adv. Earth Sci.*, 36, 139-153, 10.11867/j.issn.1001-8166.2021.022, 2021.
- Rao, R. and Sivakumar, R.: On the possible mechanisms of the evolution of a mini-warm pool during the pre-summer monsoon season and the genesis of onset vortex in the South-Eastern Arabian Sea, *Q. J. R. Meteorol. Soc.*, 125, <https://doi.org/10.1002/qj.49712555503>, 1999.
- Rao, R. R., Jitendra, V., GirishKumar, M. S., Ravichandran, M., and Ramakrishna, S. S. V. S.: Interannual variability of the Arabian Sea Warm Pool: observations and governing mechanisms, *Clim. Dyn.*, 44, 2119-2136, 10.1007/s00382-014-2243-0, 2015.

Saji, N. H., Goswami, B. N., Vinayachandran, P. N., and Yamagata, T.: A dipole mode in the tropical Indian Ocean, *Nature*, 401, 360-363, [10.1038/43854](https://doi.org/10.1038/43854), 1999.

Shankar, D., Vinayachandran, P. N., and Unnikrishnan, A. S.: The monsoon currents in the north Indian Ocean, *Prog. Oceanogr.*, 52, 63-120, [https://doi.org/10.1016/S0079-6611\(02\)00024-1](https://doi.org/10.1016/S0079-6611(02)00024-1), 2002.

Sprintall, J. and Tomczak, M.: Evidence of the barrier layer in the surface layer of the tropics, *J. Geophys. Res.: Oceans.*, 97, 7305-7316, <https://doi.org/10.1029/92JC00407>, 1992.

Reference, Reviewer 2:

Behera, S. K., Luo, J. J., Masson, S., Rao, S. A., Sakuma, H., and Yamagata, T.: A CGCM Study on the Interaction between IOD and ENSO, *J. Clim.*, 19, 1688-1705, <https://doi.org/10.1175/JCLI3797.1>, 2006.

Brainerd, K. E. and Gregg, M. C.: Surface mixed and mixing layer depths, *J. Geophys. Res.*, 113, C05305, <https://doi.org/10.1029/2002JC001935>, 2008.

Carton, J. A., Chepurin, G. A., and Chen, L.: SODA3: A New Ocean Climate Reanalysis, *J. Clim.*, 31, 6967-6983, <https://doi.org/10.1175/JCLI-D-18-0149.1>, 2018.

Chowdary, J. S. and Gnanaseelan, C.: Basin-wide warming of the Indian Ocean during El Niño and Indian Ocean dipole years, *Int J Climatol.*, 27, 1421-1438, <https://doi.org/10.1002/joc.1482>, 2007.

de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *J. Geophys. Res.: Oceans.*, 109, <https://doi.org/10.1029/2004JC002378>, 2004.

Jofia, J., Girishkumar, M. S., Ashin, K., Sureshkumar, N., Shivaprasad, S., and Pattabhi Ram Rao, E.: Mixed Layer Temperature Budget in the Arabian Sea During Winter 2019 and Spring 2019: The Role of Diapycnal Heat Flux, *J. Geophys. Res.: Oceans.*, 128, e2022JC019088, <https://doi.org/10.1029/2022JC019088>, 2023.

Joseph, P. V.: Warm pool in the Indian Ocean and monsoon onset, *Tropical Ocean-Atmos. News Let.*, 53,1-55, 1990.

Kim, S. T., Yu, J. Y., and Lu, M.-M.: The distinct behaviors of Pacific and Indian Ocean warm pool properties on seasonal and interannual time scales, *J. Geophys. Res.*, 117, <https://doi.org/10.1029/2011JD016557>, 2012.

Klein, S. A., Soden, B. J., and Lau, N.-C.: Remote Sea Surface Temperature Variations during ENSO: Evidence for a Tropical Atmospheric Bridge, *J. Clim.*, 12, 917-932, [https://doi.org/10.1175/1520-0442\(1999\)012<0917:RSSTVD>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<0917:RSSTVD>2.0.CO;2), 1999.

Kurian, J. and Vinayachandran, P. N.: Mechanisms of formation of the Arabian Sea mini warm pool in a high-resolution Ocean General Circulation Model, *J. Geophys. Res.: Oceans.*, 112, <https://doi.org/10.1029/2006JC003631>, 2007.

Lau, N.-C. and Nath, M. J.: Impact of ENSO on the Variability of the Asian–Australian Monsoons as Simulated in GCM Experiments, *J. Clim.*, 13, 4287-4309, [https://doi.org/10.1175/1520-0442\(2000\)013<4287:IOEOTV>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<4287:IOEOTV>2.0.CO;2), 2000.

Lau, N.-C. and Nath, M. J.: Atmosphere–Ocean Variations in the Indo-Pacific Sector during ENSO Episodes, *J. Clim.*, 16, 3-20, [https://doi.org/10.1175/1520-0442\(2003\)016<0003:AOVITI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<0003:AOVITI>2.0.CO;2), 2003.

Li, S., Lu, J., Huang, G., and Hu, K.: Tropical Indian Ocean Basin Warming and East Asian Summer Monsoon: A Multiple AGCM Study, *J. Clim.*, 21, 6080-6088, <https://doi.org/10.1175/2008JCLI2433.1>, 2008.

Li Xiaohui, X. F., Chen Hongying, et al.: Correlation analysis of the cycle process between the western Pacific warm pool and ENSO during 1980-2016, *Journal of Marine Meteorology*, 37(3), 85-94, 1019513/i.cnki.issn2096-3599.2017.03.010, 2017.

Mukhopadhyay, S., Shankar, D., Aparna, S. G., Fernando, V., and Kankonkar, A.: Observed variability of the East India Coastal Current on the continental shelf during 2010–2018, *Journal of Earth System Science*, 129, 106, 10.1007/s12040-020-1353-9, 2020.

Nagamani, P. V., Ali, M. M., Goni, G. J., Udaya Bhaskar, T. V. S., McCreary, J. P., Weller, R. A., Rajeevan, M., Gopala Krishna, V. V., and Pezzullo, J. C.: Heat content of the Arabian Sea Mini Warm Pool is increasing, *Atmospheric Sci. Lett.*, 17, 39-42, <https://doi.org/10.1002/asl.596>, 2016.

Rao, R. and Sivakumar, R.: On the possible mechanisms of the evolution of a mini-warm pool during the pre-summer monsoon season and the genesis of onset vortex in the South-Eastern Arabian Sea, *Q. J. R. Meteorol. Soc.*, 125, <https://doi.org/10.1002/qj.49712555503>, 1999.

Rao, R. R., Jitendra, V., GirishKumar, M. S., Ravichandran, M., and Ramakrishna, S. S. V. S.: Interannual variability of the Arabian Sea Warm Pool: observations and governing mechanisms, *Clim. Dyn.*, 44, 2119-2136, 10.1007/s00382-014-2243-0, 2015.

Shetye, S. R., Gouveia, A. D., Shankar, D., Shenoi, S. S. C., Vinayachandran, P. N., Sundar, D., Michael, G. S., and Nampoothiri, G.: Hydrography and circulation in the western Bay of Bengal during the northeast monsoon, *J. Geophys. Res.: Oceans.*, 101, 14011-14025, <https://doi.org/10.1029/95JC03307>, 1996.

Sullivan, S. C., Schiro, K. A., Stubenrauch, C., and Gentine, P.: The Response of Tropical Organized Convection to El Niño Warming, *Journal of Geophysical Research: Atmospheres*, 124, 8481-8500, <https://doi.org/10.1029/2019JD031026>, 2019.

Zihan, L., Xiaojing, L., Youmin, T., and Yi, L.: Statistical characteristics and the interdecadal variation of tropical cyclones in the North Indian Ocean, *CHINESE JOURNAL OF ATMOSPHERIC SCIENCES*, -, -, - 10.3878/j.issn.1006-9895.2303.22075, 2023.