

DO NOT USE PLANT MATERIAL TREATED WITH
VIEWPOINT® HERBICIDE FOR MULCH OR COMPOST



Viewpoint®

HERBICIDE

Dry Flowable
For Non-Crop Use

Active Ingredients	By Weight
Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)	31.6%
Aminocyclopyrachlor 6-amino-5-chloro-2-cyclopropyl-4-pyrimidinecarboxylic acid	22.8%
Metsulfuron methyl Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-yl)amino]-carbonyl]amino]sulfonyl]benzoate	7.3%
Other Ingredients	38.3%
Total	100.0%

EPA Reg. No. 432-1580 EPA Est. No. 352-IL-001

**KEEP OUT OF REACH
OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside leaflet for complete First Aid Instructions, Precautionary Statements, Directions for Use and Storage and Disposal Instructions

Nonrefillable Container
Net Weight

5 Pounds

84084972

A01821611 151216AV1

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If Swallowed: Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything to an unconscious person. Call a poison control center or doctor for further treatment advice.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Viewpoint[®] Herbicide must be used only in accordance with directions on this label or in separately published BAYER CROPSCIENCE LP labeling.

BAYER CROPSCIENCE LP will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by BAYER CROPSCIENCE LP. User assumes all risks associated with such non-labeled use.

PRODUCT INFORMATION

Viewpoint[®] Herbicide is a dispersible granule that is mixed in water and applied as a spray. Viewpoint[®] Herbicide may be applied by aerial (helicopter) or ground equipment for control of broadleaf weeds and woody species, including many terrestrial and riparian invasive and noxious weeds. Viewpoint[®] Herbicide is registered for general weed and brush control on private, public, and military lands as follows: uncultivated non-agricultural areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas, etc.); uncultivated agricultural areas - non-crop producing (such as farmyards, fuel storage areas, fence rows, non-irrigation ditchbanks, barrier strips, etc.); industrial sites - outdoor (such as lumberyards, pipeline and tank farms, etc.) and natural areas (such as wildlife management areas, wildlife openings, wildlife habitats).

This product may be applied to terrestrial non-crop sites that contain areas of temporary surface water caused by collection of water in equipment ruts or in other depressions created by management activities. It is permissible to treat intermittently flooded low lying sites, seasonally dry flood plains, and transitional areas between upland and lowland sites when no water is present. It is also permissible to treat marshes, swamps, and bogs after water has receded, as well as seasonally dry flood deltas. Viewpoint[®] Herbicide may be applied up to the water's edge. Do not apply directly to water.

Viewpoint[®] Herbicide provides preemergence and/or postemergence control of the broadleaf weeds, vines, and brush species listed in the weeds controlled section of the label. For perennial species on the label, a postemergence application must be used. For best postemergence performance, include an MSO type adjuvant to the spray solution. Excessive wetting of the target plant is not necessary but good spray coverage of the target plant is needed for best results.

Do not apply more than 20 ounces of product per acre per year.

BIOLOGICAL ACTIVITY

Viewpoint[®] Herbicide is quickly taken up by the leaves, stems and roots of plants. The effects of Viewpoint[®] Herbicide may be seen on plants from within a few hours to a few days. The most noticeable symptom is a bending and twisting of stems and leaves. Other advanced symptoms include severe chlorosis, necrosis, stem thickening, growth stunting, leaf crinkling, calloused stems and leaf veins, leaf-cupping, and enlarged roots. Death of treated broadleaf plants may require several more weeks and up to several months for some woody plant species.

Viewpoint[®] Herbicide is rain-fast at 4 hours after application.

IMPORTANT RESTRICTIONS

- Do not apply this product in areas where the roots of desirable trees and/or shrubs may extend unless injury or loss can be tolerated. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend well beyond the tree canopy.
- Do not apply this product if site-specific characteristics and conditions exist that could contribute to movement and unintended root zone exposure to desirable trees or vegetation unless injury or loss can be tolerated.

or plant canopy and coarse or coarser droplet size (ASABE S572.1), or, if specifically using a spinning atomizer nozzle, applicators are required to use a volume mean diameter (VMD) of 385 microns or greater.

- Applications with wind speeds greater than 10 mph are prohibited.
- Applications into temperature inversions are prohibited.
- Apply with the spray boom or nozzle height as low as possible.

See Spray Drift Management Section of this label for additional information.

IMPORTANT PRECAUTIONS

- Certain species may, in particular, be sensitive to low levels of Viewpoint® Herbicide including, but not limited to, conifers (such as Douglas fir, Norway spruce, ponderosa pine, and white pine), deciduous trees (such as aspen, Chinese tallow, cottonwood, honey locust, magnolia, poplar species, redbud, silver maple, and willow species), and ornamental shrubs (such as arborvitae, burning bush, crape myrtle, forsythia, hydrangea, ice plant, magnolia, purple plum, and yew).
- Injury or loss of desirable trees or vegetation may result if Viewpoint® Herbicide is applied on or near desirable trees or vegetation, on areas where their roots extend, or in locations where the treated soil may be washed or moved into contact with their roots. Consider site-specific characteristics and conditions that could contribute to unintended root zone exposure to desirable trees or vegetation. Root zone areas of desirable trees or vegetation are affected by local conditions and can extend beyond the tree canopy. If further information is needed regarding root zone area, consult appropriate state extension service, professional consultant, or other qualified authority.
- Injury to or loss of desirable trees or vegetation may result if equipment is drained or flushed on or near these trees or vegetation, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
- In non-crop areas adjacent to desirable vegetation, avoid overlapping spray applications and shut off spray to the spray boom while starting, turning, slowing, or stopping to avoid injury to desirable vegetation.
- Applications made where runoff water flows onto agricultural land may injure or kill crops, such as but not limited to sugar beets, potatoes, tomatoes, tobacco, soybeans, field beans, alfalfa, grapes, peaches, almonds, and vegetables.
- Applications should be made only when there is little or no hazard from spray drift. Very small quantities of spray, which may not be visible, may seriously injure susceptible plants.
- Exposure to Viewpoint® Herbicide may injure or kill most crops and may injure or kill desirable vegetation. Injury may be more severe when the crops or desirable vegetation are irrigated.
- Caution is advised when using this product in areas where loss of desirable conifer or deciduous trees and/or shrubs as well as other broadleaf plants, including, but not limited to, legumes and wild flowers, cannot be tolerated. Without prior experience, it is necessary that small areas containing these plants be tested for tolerance to Viewpoint® Herbicide and its soil residues before any large scale spraying occurs.
- Low rates of Viewpoint® Herbicide can kill or severely injure most crops. Following a Viewpoint® Herbicide application, the use of spray equipment to apply other pesticides to crops on which Viewpoint® Herbicide is not registered may result in their damage. The most effective way to reduce this crop damage potential is to use dedicated mixing and application equipment.
- Leave treated soil undisturbed to reduce the potential for Viewpoint® Herbicide movement by soil erosion due to wind or water.
- In the case of suspected off-site movement of Viewpoint® Herbicide to cropland, soil samples should be quantitatively analyzed for Viewpoint® Herbicide or any other herbicide which could be having an adverse effect on the crop, in addition to conducting the field bioassay.

to keep accurate records of pesticides applied to individual sites to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural reseller, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants, or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest systems in your area.

PREPARING FOR USE - SITE SPECIFIC CONSIDERATIONS

Understanding the risks associated with the application of Viewpoint® Herbicide is essential to aid in preventing off-site injury to desirable vegetation and agricultural crops. The risk of off-site movement both during and after application may be affected by a number of site specific factors such as the nature, texture and stability of the soil, the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, drainage patterns, and other local physical and environmental conditions. A careful evaluation of the potential for off-site movement from the intended application site, including movement of treated soil by wind or water erosion, must be made prior to using Viewpoint® Herbicide. This evaluation is particularly critical where desirable vegetation or crops are grown on neighboring land for which the use of Viewpoint® Herbicide is not labeled. If prevailing local conditions may be expected to result in off-site movement and cause damage to neighboring desirable vegetation or agricultural crops, do not apply Viewpoint® Herbicide.

Before applying Viewpoint® Herbicide the user must read and understand all label directions, precautions, and restrictions completely, including these requirements for a site specific evaluation. If you do not understand any of the instructions or precautions on the label, or are unable to make a site specific evaluation yourself, consult with your local BAYER CROPS SCIENCE LP Crop Protection representative, local agricultural dealer, university cooperative extension service, land manager, professional applicator, agricultural consultant, or other qualified authorities familiar with the area to be treated. If you still have questions regarding the need for site specific considerations please call 1-800-331-2867.

NON-AGRICULTURAL USES

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Terrestrial non-crop weed control is not within the scope of the Worker Protection Standard. See the Product Information section of this label for a description of noncrop sites.

Do not enter or allow worker entry into treated areas until sprays have dried.

APPLICATION INFORMATION

NON-CROP SITES

Viewpoint® Herbicide is a dispersible granule that is mixed in water and applied as a spray. Viewpoint® Herbicide may be applied by helicopter or ground equipment for control of broadleaf weeds and woody species, including many terrestrial and riparian invasive and noxious weeds.

Table 1. Viewpoint® Herbicide Spray Volume and Use Rate Mixing Instructions

Total Spray Volume gallons/acre	Viewpoint® Herbicide 13 ounces/acre ounces/ 100 gallons of spray	Viewpoint® Herbicide 16 ounces/acre ounces/ 100 gallons of spray	Viewpoint® Herbicide 18 ounces/acre ounces/ 100 gallons of spray	Viewpoint® Herbicide * 20 ounces/acre ounces/ 100 gallons of spray
400	3.25	4	4.5	5
300	4.3	5.3	6	6.7
200	6.5	8	9	10
100	13	16	18	20
75	17.3	21.3	24	26.7
50	26	32	36	40
40	32.5	40	45	50
30	43.3	53.3	60	66.7
25	52	64	72	80
20	65	80	90	100
15	86.7	106.7	120	133.3
10	130	160	180	200

* Do not exceed the maximum use rate of 20 ounces per acre per year.

BAREGROUND WEEDS	Rate 13 to 20 Ounces per Acre
Bittercress	<i>Cardamine sp.</i>
Blackberry	<i>Rubus sp.</i>
Blackeyed-susan	<i>Rudbeckia hirta</i>
Buttercup, bur	<i>Ceratocephala testiculata</i>
Caraway	<i>Carum carvi</i>
Carrot, wild	<i>Daucus carota</i>
Catchfly, conical	<i>Silene conica</i>
Chamomile, false	<i>Matricaria maritima</i>
Chickweed	<i>Stellaria media</i>
Chicory	<i>Cichorium intybus</i>
Cinquefoil, sulfur	<i>Potentilla recta</i>
Clover	<i>Trifolium sp.</i>
Clover, bush	<i>Lespedeza sp.</i>
Clover, sweet	<i>Melilotus officinalis</i>
Cocklebur	<i>Xanthium sp.</i>
Coreopsis, plains	<i>Coreopsis tinctoria</i>
Corncockle	<i>Agrostemma githago</i>
Cowcockle	<i>Vaccaria pyramidata</i>
Crazyweed, silky (locoweed)	<i>Oxytropis sericea</i>
Croton, woolly	<i>Croton capitatus</i>
Crupina	<i>Crupina vulgaris</i>
Daisy, oxeye	<i>Chrysanthemum leucanthemum</i>
Dandelion	<i>Taraxacum officinale</i>
Deadberr	<i>Rubus trivialis</i>
Dock, curly	<i>Rumex crispus</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Dyer's woad	<i>Isatis tinctoria</i>
Falseflax, smallseed	<i>Camelina microcarpa</i>
Fern, old world climbing	<i>Lygodium microphyllum</i>
Fiddleneck, tarweed	<i>Amsinckia lycopsoides</i>
Filaree, redstem	<i>Erodium cicutarium</i>
Fleabane, rough	<i>Conyza asper</i>
Flaxweed	<i>Dasycarpha sophia</i>
Gaillardia, rosering	<i>Gaillardia pulchella</i>
Garlic, wild	<i>Allium vineale</i>
Goldenrod, Canada	<i>Solidago canadensis</i>
Gorse	<i>Ulex europaeus</i>
Greasewood	<i>Sarcobatus vermiculatus</i>
Groundsel, common	<i>Senecio vulgaris</i>
Gumweed	<i>Grindelia sp.</i>
Halogeton	<i>Halogeton glomeratus</i>
Hawkweed, orange	<i>Hieracium aurantiacum</i>
Hemlock, poison	<i>Conium maculatum</i>
Henbane, black	<i>Hyoscyamus niger</i>
Henbit	<i>Lamium amplexicaule</i>
Honeysuckle, Japanese	<i>Lonicera japonica</i>
Houndstongue	<i>Cynoglossum officinale</i>
Iris, wild	<i>Dietes grandiflora</i>
Ironweed, tall	<i>Vernonia angustifolia</i>
Knapweed, diffuse	<i>Centaurea diffusa</i>
Knapweed, Russian	<i>Centaurea repens</i>
Knapweed, spotted	<i>Centaurea biebersteinii</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Kochia (Up to 6 inches) ¹	<i>Kochis scoparia</i>
Lambsquarters	<i>Chenopodium album</i>
Larkspur, duncecap	<i>Delphinium glaucum</i>
Larkspur, tall	<i>Delphinium exaltatum</i>
Lespedeza, sericea	<i>Lespedeza cuneata</i>
Lettuce, miners	<i>Montia perfoliata</i>
Lettuce, prickly	<i>Lactuca scariola</i>
Lettuce, wild	<i>Lactuca virosa</i>
Loosestrife, purple	<i>Lythrum salicaria</i>

Viewpoint® Herbicide

		High Volume Foliar ounces/ 100 gallon	Broadcast Foliar ounces/ acre
BRUSH			
Ash	<i>Fraxinus sp.</i>	13 - 20	16 - 20
Aspen	<i>Populus sp.</i>	13 - 20	16 - 20
Baccharis, eastern	<i>Baccharis halimifolia</i>	13 - 20	16 - 20
Black locust	<i>Robinia pseudoacacia</i>	13 - 20	16 - 20
Black tupelo/gum	<i>Nyssa sylvatica</i>	13 - 20	16 - 20
Boxelder	<i>Acer negundo</i>	13 - 20	16 - 20
Camellthorn	<i>Acacia erioloba</i>	13 - 20	16 - 20
Cherry	<i>Prunus sp.</i>	13 - 20	16 - 20
Cottonwood	<i>Populus sp.</i>	13 - 20	16 - 20
Elderberry	<i>Sambucus sp.</i>	13 - 20	16 - 20
Elm	<i>Ulmus sp.</i>	13 - 20	16 - 20
Hackberry, common	<i>Celtis occidentalis</i>	13 - 20	16 - 20
Hawthorn	<i>Crataegus sp.</i>	13 - 20	16 - 20
Honeysuckle	<i>Lonicera sp.</i>	13 - 20	16 - 20
Hophornbeam	<i>Ostrya sp.</i>	13 - 20	16 - 20
Locust, honey	<i>Gleditsia triacanthos</i>	13 - 20	16 - 20
Mesquite	<i>Prosopis juliflora</i>	13 - 20	16 - 20
Mulberry	<i>Morus sp.</i>	13 - 20	16 - 20
Myrtle dahoon	<i>Ilex myrtifolia</i>	13 - 20	16 - 20
Persimmon, common	<i>Diospyros virginiana</i>	13 - 20	16 - 20
Ocean spray	<i>Holodiscus sp.</i>	13 - 20	16 - 20
Osage orange	<i>Maclura pomifera</i>	13 - 20	16 - 20
Red maple	<i>Acer rubrum</i>	13 - 20	16 - 20
Sassafras	<i>Sassafras albidum</i>	13 - 20	16 - 20
Silver maple	<i>Acer sacharinum</i>	13 - 20	16 - 20
Snowberry	<i>Symphoricarpos sp.</i>	13 - 20	16 - 20
Sugarberry	<i>Celtis laevigata</i>	13 - 20	16 - 20
Sumac	<i>Rhus sp.</i>	13 - 20	16 - 20
Sweetgum	<i>Liquidambar styraciflua</i>	13 - 20	16 - 20
Sycamore	<i>Acer pseudoplatanus</i>	13 - 20	16 - 20
Tallow, Chinese	<i>Sapium sebiferum</i>	13 - 20	16 - 20
Tree of heaven	<i>Ailanthus altissima</i>	13 - 20	16 - 20
Willow	<i>Salix sp.</i>	13 - 20	16 - 20
Yellow poplar	<i>Liriodendron tulipifera</i>	13 - 20	16 - 20
Vaccinium	<i>Vaccinium sp.</i>	13 - 20	16 - 20
Cedar, eastern red ²	<i>Juniperus virginiana</i>	16 - 20	20
Dogwood	<i>Cornus sp.</i>	16 - 20	20
Fir	<i>Abies sp.</i>	16 - 20	20
Hickory	<i>Carya sp.</i>	16 - 20	20
Huisache	<i>Acacia farnesiana</i>	16 - 20	20
Oak	<i>Quercus sp.</i>	16 - 20	20
Pine, loblolly	<i>Pinus taeda</i>	16 - 20	20
Pine, Virginia ^{3,4}	<i>Pinus virginia</i>	16 - 20	20
Privet, Chinese	<i>Ligustrum sinense</i>	16 - 20	20
Rabbitbrush, rubber	<i>Ericameria nauseosa</i>	16 - 20	20
Spruce, black	<i>Picea mariana</i>	16 - 20	20
Spruce, white	<i>Picea glauca</i>	16 - 20	20
Yaupon ³	<i>Ilex vomitoria</i>	16 - 20	20

(continued)

The application volume required will vary with the height and density of the brush and the type of application equipment. In general, aerial application spray volumes range from 15 to 25 gallons per acre.

MIXING INSTRUCTIONS

1. Fill the tank 1/3 to 1/2 full of water.
2. While agitating, add the required amount of Viewpoint® Herbicide.
3. Continue agitation until the Viewpoint® Herbicide is fully dispersed, at least 5 minutes.
4. Once the Viewpoint® Herbicide is fully dispersed, maintain agitation and continue filling tank with water. Viewpoint® Herbicide must be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) and then add the necessary volume of spray adjuvants. Always add spray adjuvants last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply Viewpoint® Herbicide spray mixture within 24 hours of mixing to avoid product degradation.
8. If Viewpoint® Herbicide and a tank mix partner are to be applied in multiple loads, pre-slurry Viewpoint® Herbicide in clean water prior to adding it to the tank. This will prevent the tank mix partner from interfering with the dissolution of the Viewpoint® Herbicide.

SPRAYER CLEANUP

Thoroughly clean all mixing and spray equipment following applications of Viewpoint® Herbicide as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water.
2. Fill the tank with clean water and 1 gallon of household ammonia (contains 3% active) for every 100 gallons of water.
Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank.
Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
Equivalent amounts of an alternate-strength ammonia solution or a commercial cleaner can be used in the cleanup procedure. If a commercial cleaner is used, carefully read and follow the individual cleaner instructions.
3. Remove the nozzles and screens and clean separately in bucket containing cleaning agent and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. Dispose of the rinsate on a labeled site or at an approved waste disposal facility. If a commercial cleaner is used follow the directions for rinsate disposal on the label.
Caution: Do not use chlorine bleach with ammonia as dangerous gases will form. Do not clean equipment in an enclosed area.

Notes:

1. Always start with a clean spray tank.
2. Steam-clean aerial spray tanks to facilitate the removal of any caked deposits.
3. When Viewpoint® Herbicide is tank mixed with other pesticides, all cleanout procedures for each product must be examined and the most rigorous procedure must be followed.
4. In addition to this cleanout procedure, all pre-cleanout guidelines on subsequently applied products must be followed as per the individual labels.

WIND

Drift potential is lowest when applications are made in light to gentle sustained winds (2-10 mph) which are blowing in a constant direction. Many factors, including droplet size and equipment type, also determine drift potential at any given wind speed. **AVOID GUSTY OR WINDLESS CONDITIONS.**

Local terrain can also influence wind patterns. Every applicator is expected to be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

Setting up equipment to produce larger droplets to compensate for droplet evaporation can reduce spray drift potential. Droplet evaporation is most severe when conditions are both hot and dry.

SURFACE TEMPERATURE INVERSIONS

Drift potential is high during a surface temperature inversion. Surface inversions restrict vertical air mixing, which may cause small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Surface inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Mist or fog may indicate the presence of an inversion in humid areas. Inversions may also be identified by producing smoke and observing its behavior. Smoke that remains close to the ground, or moves laterally in a concentrated cloud under low wind conditions indicates a surface inversion. Smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are minimizing drift potential, and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for the intended application, that it is configured properly, and that drift potential has been minimized.

Note: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Read the specific crop use and application equipment instructions to determine if an air assisted field crop sprayer can be used.

SENSITIVE AREAS

Making applications when there is a sustained wind moving away from adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is an effective way to minimize the effect of spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Chemical Producers and Distributors Association (CPDA).

WIND EROSION

Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface must first be settled by rainfall or irrigation.

STORAGE AND DISPOSAL *(continued)*

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums With Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack, or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack, or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums With Liners: Refillable container (fiber drum only). **Refilling Fiber Drum:** Refill this fiber drum with Viewpoint® Herbicide containing aminocyclopyrachlor, metsulfuron methyl, and imazapyr only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. **Disposing of Fiber Drum and/or Liner:** Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. **Refilling Container:** Refill this container with Viewpoint® Herbicide containing aminocyclopyrachlor, metsulfuron methyl, and imazapyr only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact BAYER CROPSCIENCE LP at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact BAYER CROPSCIENCE LP at the number below for instructions. **Disposing of Container:** Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour, or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

(continued)

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury, other property damage, as well as other unintended consequences may result because of factors beyond the control of Bayer CropScience LP. Those factors include, but are not limited to, weather conditions, presence of other materials or the manner of use or application. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

For product information call: 1-800-331-2867

Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

Bayer

OPEN HERE

DO NOT USE PLANT MATERIAL TREATED WITH VIEWPOINT® HERBICIDE FOR MULCH OR COMPOST



Viewpoint®

HERBICIDE

Dry Flowable	
For Non-Crop Use	
Active Ingredients	By Weight
Imazapyr	
(2-[4,5-dihydro-4-methyl-4-(1-methyl-5-oxo-1H-imidazol-2-yl)-3-pyridinecarboxylic acid])	31.6%
Aminocyclopyrachlor	
6-amino-5-chloro-2-cyclopropyl-4-pyrimidinecarboxylic acid	22.8%
Metsulfuron methyl	
Methyl 2-[[[(4-methoxy-6-methyl-1,3,5-triazin-2-ylamino)-carbonyl]amino]sulfonyl]benzoate	7.3%
Other Ingredients	38.3%
Total	100.0%
EPA Reg. No. 432-1580	EPA Est. No. 352-IL-001

KEEP OUT OF REACH OF CHILDREN CAUTION

See attached leaflet for complete First Aid Instructions, Precautionary Statements, Directions for Use and Storage and Disposal Instructions.

Nonrefillable Container Net Weight 5 Pounds 84084972 A01821611 151216A1

ENVIRONMENTAL HAZARDS
This product is toxic to plants. Ditch and run-off may be hazardous to plants in water adjacent to treated areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Surface Water Advisory
This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of aminocyclopyrachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Ground Water Advisory
Aminocyclopyrachlor has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

PHYSICAL AND CHEMICAL HAZARDS
Spray solutions of Viewpoint® herbicide must be mixed, stored, and applied only in stainless steel, fiberglass, plastic, and plastic-lined steel containers. Do not mix, store, or apply Viewpoint Herbicide or spray solutions of Viewpoint Herbicide in unlined steel (except stainless steel) containers or spray tanks.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage and disposal.
Pesticide Storage: Store product in original container only. Store in a cool, dry place.
Pesticide Disposal: Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.
CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Refillable Container" or "Nonrefillable Container" designation.

Bayer (reg'd), the Bayer Cross (reg'd) and Viewpoint® are registered trademarks of Bayer.



(01) 00785740387139

FIRST AID
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF SWALLOWED: Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control center or doctor. DO NOT give anything to an unconscious person. Call a poison control center or doctor for further treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

USER SAFETY RECOMMENDATIONS
USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION
Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.
Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.
PERSONAL PROTECTIVE EQUIPMENT (PPE)
Some materials that are chemical-resistant to this product are polyethylene and polyvinylchloride. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.
Mixers/loaders, applicators, and other handlers must wear:
Long-sleeved shirt and long pants
Shoes plus socks
Chemical resistant gloves for all mixers and loaders, plus applicators using handheld equipment.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.
Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.
Engineering Control Statement: Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(c)(6)]. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Produced for: Bayer Environmental Science A Division of Bayer CropScience LP 2 T. W. Alexander Drive Research Triangle Park, NC 27709 Made in U.S.A. Bayer