

LIGHT UNFLAVORED MESONS

($S = C = B = 0$)

For $l = 1$ (π, b, ρ, a): $u\bar{d}, (u\bar{u}-d\bar{d})/\sqrt{2}, d\bar{u}$;
for $l = 0$ ($\eta, \eta', h, h', \omega, \phi, f, f'$): $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

NODE=MXXX005

NODE=MXXX005

NODE=M014

$f_0(500)$ or σ
was $f_0(600)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=M014

$f_0(500)$ T-MATRIX POLE \sqrt{s}

NODE=M014PP

Note that $\Gamma \approx 2 \text{Im}(\sqrt{s_{\text{pole}}})$.

NODE=M014PP

NODE=M014PP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400–550)–i(200–350) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
(440 ± 10)– i (238 ± 10)	1	ALBALADEJO 12	RVUE Compilation
(445 ± 25)– i (278 ⁺²² _{–18})	2,3	GARCIA-MAR..11	RVUE Compilation
(457 ⁺¹⁴ _{–13})– i (279 ⁺¹¹ _{–7})	2,4	GARCIA-MAR..11	RVUE Compilation
(442 ⁺⁵ _{–8})– i (274 ⁺⁶ _{–5})	5	MOUSSALLAM11	RVUE Compilation
(452 ± 13)– i (259 ± 16)	6	MENNESSIER 10	RVUE Compilation
(448 ± 43)– i (266 ± 43)	7	MENNESSIER 10	RVUE Compilation
(455 ± 6 ⁺³¹ _{–13})– i (278 ± 6 ⁺³⁴ _{–43})	8	CAPRINI 08	RVUE Compilation
(463 ± 6 ⁺³¹ _{–17})– i (259 ± 6 ⁺³³ _{–34})	9	CAPRINI 08	RVUE Compilation
(552 ⁺⁸⁴ _{–106})– i (232 ⁺⁸¹ _{–72})	10	ABLIKIM 07A	BES2 $\psi(2S) \rightarrow \pi^+\pi^-\ J/\psi$
(466 ± 18)– i (223 ± 28)	11	BONVICINI 07	CLEO $D^+ \rightarrow \pi^-\pi^+\pi^+$
(472 ± 30)– i (271 ± 30)	12	BUGG 07A	RVUE Compilation
(484 ± 17)– i (255 ± 10)		GARCIA-MAR..07	RVUE Compilation
(430)– i (325)	13	ANISOVICH 06	RVUE Compilation
(441 ⁺¹⁶ _{–8})– i (272 ⁺⁹ _{–12.5})	14	CAPRINI 06	RVUE $\pi\pi \rightarrow \pi\pi$
(470 ± 50)– i (285 ± 25)	15	ZHOU 05	RVUE
(541 ± 39)– i (252 ± 42)	16	ABLIKIM 04A	BES2 $J/\psi \rightarrow \omega\pi^+\pi^-$
(528 ± 32)– i (207 ± 23)	17	GALLEGOS 04	RVUE Compilation
(440 ± 8)– i (212 ± 15)	18	PELAEZ 04A	RVUE $\pi\pi \rightarrow \pi\pi$
(533 ± 25)– i (249 ± 25)	19	BUGG 03	RVUE
517 – i 240		BLACK 01	RVUE $\pi^0\pi^0 \rightarrow \pi^0\pi^0$
(470 ± 30)– i (295 ± 20)	14	COLANGELO 01	RVUE $\pi\pi \rightarrow \pi\pi$
(535 ⁺⁴⁸ _{–36})– i (155 ⁺⁷⁶ _{–53})	20	ISHIDA 01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
610 ± 14 – i 620 ± 26	21	SUROVTSEV 01	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
(540 ⁺³⁶ _{–29})– i (193 ⁺³² _{–40})		ISHIDA 00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
445 – i 235		HANNAH 99	RVUE π scalar form factor
(523 ± 12)– i (259 ± 7)		KAMINSKI 99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
442 – i 227		OLLER 99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
469 – i 203		OLLER 99B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
445 – i 221		OLLER 99C	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$
(1530 ⁺⁹⁰ _{–250})– i (560 ± 40)		ANISOVICH 98B	RVUE Compilation
420 – i 212		LOCHER 98	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
440 – i 245	22	DOBADO 97	RVUE Compilation
(602 ± 26)– i (196 ± 27)	23	ISHIDA 97	$\pi\pi \rightarrow \pi\pi$
(537 ± 20)– i (250 ± 17)	24	KAMINSKI 97B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, 4\pi$
470 – i 250	25,26	TORNQVIST 96	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
387 – i 305	26,27	JANSSEN 95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
420 – i 370	28	ACHASOV 94	RVUE $\pi\pi \rightarrow \pi\pi$
(506 ± 10)– i (247 ± 3)		KAMINSKI 94	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
370 – i 356	29	ZOU 94B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
408 – i 342	26,29	ZOU 93	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$

470 – $i208$	³⁰ VANBEVEREN 86	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta,$	
(750 ± 50) – $i(450 ± 50)$	³¹ ESTABROOKS 79	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
(660 ± 100) – $i(320 ± 70)$	PROTOPOP... 73	HBC	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
650 – $i370$	³² BASDEVANT 72	RVUE	$\pi\pi \rightarrow \pi\pi$	
¹ Applying the chiral unitary approach at NLO to the K_{e4} data of BATLEY 10 and $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.				NODE=M014PP;LINKAGE=AD
² Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.				NODE=M014PP;LINKAGE=GM
³ Analytic continuation using Roy equations.				NODE=M014PP;LINKAGE=GR
⁴ Analytic continuation using GKPY equations.				NODE=M014PP;LINKAGE=GC
⁵ Using Roy equations.				NODE=M014PP;LINKAGE=MO
⁶ Average of three variants of the analytic K-matrix model. Uses the K_{e4} data of BATLEY 08A and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73 and GRAYER 74.				NODE=M014PP;LINKAGE=ME
⁷ Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.				NODE=M014PP;LINKAGE=MN
⁸ From the K_{e4} data of BATLEY 08A and $\pi N \rightarrow \pi\pi N$ data of HYAMS 73.				NODE=M014PP;LINKAGE=CA
⁹ From the K_{e4} data of BATLEY 08A and $\pi N \rightarrow \pi\pi N$ data of PROTOPOPESCU 73, GRAYER 74, and ESTABROOKS 74.				NODE=M014PP;LINKAGE=AP
¹⁰ From a mean of three different $f_0(500)$ parametrizations. Uses 40k events.				NODE=M014PP;LINKAGE=AL
¹¹ From an isobar model using 2.6k events.				NODE=M014PP;LINKAGE=BO
¹² Reanalysis of ABLIKIM 04A, PISLAK 01, and HYAMS 73 data.				NODE=M014PP;LINKAGE=BU
¹³ Using the N/D method.				NODE=M014PP;LINKAGE=AN
¹⁴ From the solution of the Roy equation (ROY 71) for the isoscalar S-wave and using a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.				NODE=M014PP;LINKAGE=CL
¹⁵ Reanalysis of the data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, ROSSELET 77, PISLAK 03, and AKHMETSHIN 04.				NODE=M014PP;LINKAGE=ZH
¹⁶ From a mean of six different analyses and $f_0(500)$ parameterizations.				NODE=M014PP;LINKAGE=AB
¹⁷ Using data on $\psi(2S) \rightarrow J/\psi\pi\pi$ from BAI 00E and on $\mathcal{T}(nS) \rightarrow \mathcal{T}(mS)\pi\pi$ from BUTLER 94B and ALEXANDER 98.				NODE=M014PP;LINKAGE=GA
¹⁸ Reanalysis of data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, and COHEN 80 in the unitarized ChPT model.				NODE=M014PP;LINKAGE=PE
¹⁹ From a combined analysis of HYAMS 73, AUGUSTIN 89, AITALA 01B, and PISLAK 01.				NODE=M014PP;LINKAGE=PS
²⁰ A similar analysis (KOMADA 01) finds $(580^{+79}_{-30}) - i(190^{+107}_{-49})$ MeV.				NODE=M014PP;LINKAGE=KI
²¹ Coupled channel reanalysis of BATON 70, BENSINGER 71, BAILLON 72, HYAMS 73, HYAMS 75, ROSSELET 77, COHEN 80, and ETKIN 82B using the uniformizing variable.				NODE=M014PP;LINKAGE=SU
²² Using the inverse amplitude method and data of ESTABROOKS 73, GRAYER 74, and PROTOPOPESCU 73.				NODE=M014PP;LINKAGE=DO
²³ Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.				NODE=M014PP;LINKAGE=AA
²⁴ Average and spread of 4 variants (“up” and “down”) of KAMINSKI 97B 3-channel model.				NODE=M014PP;LINKAGE=E
²⁵ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.				NODE=M014PP;LINKAGE=B
²⁶ Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.				NODE=M014PP;LINKAGE=G
²⁷ Analysis of data from FALVARD 88.				NODE=M014PP;LINKAGE=C
²⁸ Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.				NODE=M014PP;LINKAGE=D
²⁹ Analysis of data from OCHS 73, GRAYER 74, and ROSSELET 77.				NODE=M014PP;LINKAGE=F
³⁰ Coupled-channel analysis using data from PROTOPOPESCU 73, HYAMS 73, HYAMS 75, GRAYER 74, ESTABROOKS 74, ESTABROOKS 75, FROGGATT 77, CORDEN 79, BISWAS 81.				NODE=M014PP;LINKAGE=BV
³¹ Analysis of data from APEL 73, GRAYER 74, CASON 76, PAWLICKI 77. Includes spread and errors of 4 solutions.				NODE=M014PP;LINKAGE=A
³² Analysis of data from BATON 70, BENSINGER 71, COLTON 71, BAILLON 72, PROTOPOPESCU 73, and WALKER 67.				NODE=M014PP;LINKAGE=J

$f_0(500)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETERS

NODE=M014M

VALUE (MeV) DOCUMENT ID TECN COMMENT

(400–550) OUR ESTIMATE

NODE=M014M

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

513 ± 32	³³ MURAMATSU 02	CLEO	$e^+e^- \approx 10$ GeV
$478^{+24}_{-23} \pm 17$	AITALA	01B	E791 $D^+ \rightarrow \pi^- \pi^+ \pi^+$
563^{+58}_{-29}	³⁴ ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon \pi\pi$
555	³⁵ ASNER	00	CLE2 $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
540 ± 36	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$
750 ± 4	ALEKSEEV	99	SPEC 1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^- \pi^+ n$
744 ± 5	ALEKSEEV	98	SPEC 1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^- \pi^+ n$
759 ± 5	³⁶ TROYAN	98	5.2 $np \rightarrow np \pi^+ \pi^-$

780±30	ALDE	97	GAM2	450	$pp \rightarrow pp\pi^0\pi^0$
585±20	37 ISHIDA	97			$\pi\pi \rightarrow \pi\pi$
761±12	38 SVEC	96	RVUE	6-17	$\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 860	39,40 TORNQVIST	96	RVUE		$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1165±50	41,42 ANISOVICH	95	RVUE		$\pi^-p \rightarrow \pi^0\pi^0n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
~ 1000	43 ACHASOV	94	RVUE		$\pi\pi \rightarrow \pi\pi$
414±20	38 AUGUSTIN	89	DM2		

33 Statistical uncertainty only.

34 A similar analysis (KOMADA 01) finds 526^{+48}_{-37} MeV.

35 From the best fit of the Dalitz plot.

36 6σ effect, no PWA.

37 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

38 Breit-Wigner fit to S-wave intensity measured in $\pi N \rightarrow \pi^-\pi^+N$ on polarized targets. The fit does not include $f_0(980)$.

39 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

40 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays.

41 Uses $\pi^0\pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+\pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.

42 The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

43 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.

NODE=M014M;LINKAGE=UT

NODE=M014M;LINKAGE=KI

NODE=M014M;LINKAGE=KK

NODE=M014M;LINKAGE=TN

NODE=M014M;LINKAGE=AA

NODE=M014M;LINKAGE=E

NODE=M014M;LINKAGE=B

NODE=M014M;LINKAGE=GG

NODE=M014M;LINKAGE=F

NODE=M014M;LINKAGE=G

NODE=M014M;LINKAGE=D

$f_0(500)$ BREIT-WIGNER WIDTH

NODE=M014W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400-700) OUR ESTIMATE			

NODE=M014W

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

335± 67	44 MURAMATSU	02	CLEO	$e^+e^- \approx 10$ GeV
324 $^{+42}_{-40} \pm 21$	AITALA	01B	E791	$D^+ \rightarrow \pi^-\pi^+\pi^+$
372 $^{+229}_{-95}$	45 ISHIDA	01		$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
540	46 ASNER	00	CLE2	$\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
372± 80	ISHIDA	00B		$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
119± 13	ALEKSEEV	99	SPEC	$1.78 \pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
77± 22	ALEKSEEV	98	SPEC	$1.78 \pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
35± 12	47 TROYAN	98		$5.2 np \rightarrow np\pi^+\pi^-$
780± 60	ALDE	97	GAM2	450 $pp \rightarrow pp\pi^0\pi^0$
385± 70	48 ISHIDA	97		$\pi\pi \rightarrow \pi\pi$
290± 54	49 SVEC	96	RVUE	6-17 $\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 880	50,51 TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
460± 40	52,53 ANISOVICH	95	RVUE	$\pi^-p \rightarrow \pi^0\pi^0n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
~ 3200	54 ACHASOV	94	RVUE	$\pi\pi \rightarrow \pi\pi$
494± 58	49 AUGUSTIN	89	DM2	

44 Statistical uncertainty only.

45 A similar analysis (KOMADA 01) finds 301^{+145}_{-100} MeV.

46 From the best fit of the Dalitz plot.

47 6σ effect, no PWA.

48 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

49 Breit-Wigner fit to S-wave intensity measured in $\pi N \rightarrow \pi^-\pi^+N$ on polarized targets. The fit does not include $f_0(980)$.

50 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

51 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays.

52 Uses $\pi^0\pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+\pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.

53 The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

54 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.

NODE=M014W;LINKAGE=UT

NODE=M014W;LINKAGE=KI

NODE=M014W;LINKAGE=KK

NODE=M014W;LINKAGE=TN

NODE=M014W;LINKAGE=AA

NODE=M014W;LINKAGE=E

NODE=M014W;LINKAGE=B

NODE=M014W;LINKAGE=GG

NODE=M014W;LINKAGE=F

NODE=M014W;LINKAGE=G

NODE=M014W;LINKAGE=D1

$f_0(500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	dominant
Γ_2 $\gamma\gamma$	seen

NODE=M014215;NODE=M014

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

 $f_0(500)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_2
------------------------	-------------	-------------	------	---------	------------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2.05±0.21	55	DAI	14A	RVUE	Compilation	
1.7 ±0.4	56	HOFERICHTER11		RVUE	Compilation	
3.08±0.82	57	MENNESSIER	11	RVUE	Compilation	
2.08±0.2	58	MOUSSALLAM11		RVUE	Compilation	
2.08	59	MAO	09	RVUE	Compilation	
1.2 ±0.4	60	BERNABEU	08	RVUE		
3.9 ±0.6	57	MENNESSIER	08	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
1.8 ±0.4	61	OLLER	08	RVUE	Compilation	
1.68±0.15	61,62	OLLER	08A	RVUE	Compilation	
3.1 ±0.5	63,64	PENNINGTON	08	RVUE	Compilation	
2.4 ±0.4	64,65	PENNINGTON	08	RVUE	Compilation	
4.1 ±0.3	66	PENNINGTON	06	RVUE	$\gamma\gamma \rightarrow \pi^0\pi^0$	
3.8 ±1.5	67,68	BOGLIONE	99	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
5.4 ±2.3	67	MORGAN	90	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
10 ±6		COURAU	86	DM1	$e^+e^- \rightarrow \pi^+\pi^-e^+e^-$	
55 Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.						
56 Using Roy-Steiner equations with $\pi\pi$ phase shifts from an update of COLANGELO 01 and from GARCIA-MARTIN 11A.						
57 Using an analytic K-matrix model.						
58 Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.						
59 Used dispersion theory. The value quoted used the $f_0(500)$ pole position of 457 - i276 MeV.						
60 Using p , n polarizabilities from PDG 06 and fitting to $\pi\pi$ phase motion from GARCIA-MARTIN 07 and σ -poles from GARCIA-MARTIN 07 and CAPRINI 06.						
61 Using twice-subtracted dispersion integrals.						
62 Supersedes OLLER 08.						
63 Solution A (preferred solution based on χ^2 -analysis).						
64 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.						
65 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).						
66 Using unitarity and the σ pole position from CAPRINI 06.						
67 This width could equally well be assigned to the $f_0(1370)$. The authors analyse data from BOYER 90 and MARSISKE 90 and report strong correlation with $\gamma\gamma$ width of $f_2(1270)$.						
68 Supersedes MORGAN 90.						

NODE=M014220

NODE=M014W2
NODE=M014W2

OCCUR=2

NODE=M014W2;LINKAGE=B

NODE=M014W2;LINKAGE=HO

NODE=M014W2;LINKAGE=ME
NODE=M014W2;LINKAGE=MO

NODE=M014W2;LINKAGE=MA

NODE=M014W2;LINKAGE=BE

NODE=M014W2;LINKAGE=OL

NODE=M014W2;LINKAGE=LL

NODE=M014W2;LINKAGE=P1

NODE=M014W2;LINKAGE=P3

NODE=M014W2;LINKAGE=P2

NODE=M014W2;LINKAGE=PE

NODE=M014W2;LINKAGE=A

NODE=M014W2;LINKAGE=BL

 $f_0(500)$ REFERENCES

DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
ALBALADEJO	12	PR D86 034003	M. Albaladejo, J.A. Oller	(MURC)
GARCIA-MAR...	11	PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
GARCIA-MAR...	11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
HOFERICHTER	11	EPJ C71 1743	M. Hoferichter, D.R. Phillips, C. Schat	(BONN+)
MENNESSIER	11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang	
MOUSSALLAM	11	EPJ C71 1814	B. Moussallam	
BATLEY	10	PL B686 101	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY	10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
MENNESSIER	10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang	
MAO	09	PR D79 116008	Y. Mao <i>et al.</i>	
BATLEY	08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BERNABEU	08	PRL 100 241804	J. Bernabeu, J. Prades	(IFIC, GRAN)
CAPRINI	08	PR D77 114019	I. Caprini	
MENNESSIER	08	PL B665 205	G. Mennessier, S. Narison, W. Ochs	
OLLER	08	PL B659 201	J.A. Oller, L. Roca, C. Schat	(MURC, UBA)
OLLER	08A	EPJ A37 15	J.A. Oller, L. Roca	(MURC)
PENNINGTON	08	EPJ C56 1	M.R. Pennington <i>et al.</i>	
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07A	PL B645 19	M. Ablikim <i>et al.</i>	(BES Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>	
GARCIA-MAR...	07	PR D76 074034	R. Garcia-Martin, J.R. Pelaez, F.J. Yndurain	
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)
ANISOVICH	06	IJMP A21 3615	V.V. Anisovich	

NODE=M014

REFID=55923

REFID=54277

REFID=16761

REFID=54121

REFID=53966

REFID=53637

REFID=53975

REFID=53285

REFID=53567

REFID=53657

REFID=52992

REFID=52487

REFID=52271

REFID=52273

REFID=52299

REFID=53976

REFID=53977

REFID=52303

REFID=52309

REFID=51614

REFID=51721

REFID=53252

REFID=51949

REFID=51652

REFID=51137

CAPRINI	06	PRL 96 132001	I. Caprini, G. Colangelo, H. Leutwyler	(BCIP+)	REFID=51076
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
PENNINGTON	06	PRL 97 011601	M.R. Pennington		REFID=51184
ZHOU	05	JHEP 0502 043	Z.Y. Zhou <i>et al.</i>		REFID=50823
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49740
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
BUETTIKER	04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam		REFID=56428; ERROR==1
GALLEGOS	04	PR D69 074033	A. Gallegos <i>et al.</i>		REFID=49769
PELAEZ	04A	MPL A19 2879	J.R. Pelaez		REFID=50347
BUGG	03	PL B572 1	D.V. Bugg		REFID=49586
PISLAK	03	PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=49344
Also		PR D81 119903E	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=53337
MURAMATSU	02	PRL 89 251802	H. Muramatsu <i>et al.</i>	(CLEO Collab.)	REFID=49081
Also		PRL 90 059901 (erratum)	H. Muramatsu <i>et al.</i>	(CLEO Collab.)	REFID=49385
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
BLACK	01	PR D64 014031	D. Black <i>et al.</i>		REFID=48314
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leutwyler		REFID=49180
ISHIDA	01	PL B518 47	M. Ishida <i>et al.</i>		REFID=48354
KOMADA	01	PL B508 31	T. Komada <i>et al.</i>		REFID=48541
PISLAK	01	PRL 87 221801	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=48433
Also		PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=49344
Also		PRL 105 019901E	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=53338
SUROVTSEV	01	PR D63 054024	Y.S. Surovtsev, D. Krupa, M. Nagy		REFID=48310
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
BAI	00E	PR D62 032002	J. Bai <i>et al.</i>	(BES Collab.)	REFID=47955
ISHIDA	00B	PTP 104 203	M. Ishida <i>et al.</i>		REFID=48358
ALEKSEEV	99	NP B541 3	I.G. Alekseev <i>et al.</i>		REFID=46614
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
HANNAH	99	PR D60 017502	T. Hannah		REFID=46935
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
ALEKSEEV	98	PAN 61 174	I.G. Alekseev <i>et al.</i>		REFID=46328
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)	REFID=46372
TROYAN	98	JINRRC 5-91 33	Yu. Troyan <i>et al.</i>		REFID=46615
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
DOBADO	97	PR D56 3057	A. Dobado, J.R. Pelaez		REFID=53964
ISHIDA	97	PTP 98 1005	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45998
KAMINSKI	97B	PL B413 130	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, IPN)	REFID=45778
Also		PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45770
SVEC	96	PR D53 2343	M. Svec	(MCGI)	REFID=44509
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=44375
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)	REFID=44442
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
ACHASOV	94	PR D49 5779	N.N. Achasov, G.N. Shestakov	(NOVM)	REFID=44087
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43799
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)	REFID=44072
ZOU	93	PR D48 R3948	B.S. Zou, D.V. Bugg	(LOQM)	REFID=43672
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
COURAU	86	NP B271 1	A. Courau <i>et al.</i>	(CLER, LALO)	REFID=44510
VANBEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i>	(NIJM, BIEL)	REFID=45769
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BISWAS	81	PRL 47 1378	N.N. Biswas <i>et al.</i>	(NDAM, ANL)	REFID=21106
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP	REFID=20381
MUKHIN	80	JETPL 32 601	K.N. Mukhin <i>et al.</i>	(KIAE)	REFID=44528
		Translated from ZETFP 32 616.			
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
ESTABROOKS	79	PR D19 2678	P. Estabrooks	(CARL)	REFID=20375
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJ	REFID=20367
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)	REFID=11004
CASON	76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL) IJ	REFID=21064
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20642
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)	REFID=21062
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
APEL	73	PL 41B 542	W.D. Apel <i>et al.</i>	(KARL, PISA)	REFID=44532
ESTABROOKS	73	Tallahassee	P.G. Estabrooks <i>et al.</i>	(CERN, MPIM)	REFID=20345
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)	REFID=20349
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108
BAILLON	72	PL 38B 555	P.H. Baillon <i>et al.</i>	(SLAC)	REFID=20093
BASDEVANT	72	PL 41B 178	J.L. Basdevant, C.D. Froggatt, J.L. Petersen	(CERN)	REFID=20095
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)	REFID=44530
BENSINGER	71	PL 36B 134	J.R. Bensinger <i>et al.</i>	(WISC)	REFID=21006
COLTON	71	PR D3 2028	E.P. Colton <i>et al.</i>	(LBL, FNAL, UCLA+)	REFID=44533
ROY	71	PL 36B 353	S.M. Roy		REFID=51107
BATON	70	PL 33B 528	J.P. Baton, G. Laurens, J. Reigner	(SACL)	REFID=20086
WALKER	67	RMP 39 695	W.D. Walker	(WISC)	REFID=20960

$\rho(770)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=M009

NODE=M009

NODE=M009205

NODE=M009205

 $\rho(770)$ MASS

We no longer list S-wave Breit-Wigner fits, or data with high combinatorial background.

NODE=M009M0

NODE=M009M0

NEUTRAL ONLY, e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
775.26±0.25 OUR AVERAGE				
775.02±0.35		1 LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
775.97±0.46±0.70	900k	2 AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
774.6 ±0.4 ±0.5	800k	3,4 ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
775.65±0.64±0.50	114k	5,6 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
775.9 ±0.5 ±0.5	1.98M	7 ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.8 ±0.9 ±2.0	500k	7 ACHASOV 02	SND	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ±1.1		8 BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
775.8 ±0.5 ±0.3	1.98M	9 ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ±0.6 ±0.5	1.98M	10 ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.0 ±0.6 ±1.1	500k	11 ACHASOV 02	SND	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ±0.7 ±5.3		12 BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$, $\mu^+\mu^-$
770.5 ±1.9 ±5.1		13 GARDNER 98	RVUE	$0.28-0.92 e^+e^- \rightarrow \pi^+\pi^-$
764.1 ±0.7		14 O'CONNELL 97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
757.5 ±1.5		15 BERNICHA 94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
768 ±1		16 GESHKEN... 89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$

OCCUR=2

OCCUR=3

OCCUR=3

CHARGED ONLY, τ DECAYS and e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
775.11±0.34 OUR AVERAGE					
774.6 ±0.2 ±0.5	5.4M	17,18 FUJIKAWA 08	BELL	±	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
775.5 ±0.7		18,19 SCHAEEL 05C	ALEP		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
775.5 ±0.5 ±0.4	1.98M	7 ALOISIO 03	KLOE		$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ±1.1 ±0.5	87k	20,21 ANDERSON 00A	CLE2		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
774.8 ±0.6 ±0.4	1.98M	10 ALOISIO 03	KLOE	-	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
776.3 ±0.6 ±0.7	1.98M	10 ALOISIO 03	KLOE	+	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
773.9 ±2.0 $\begin{smallmatrix} +0.3 \\ -1.0 \end{smallmatrix}$		22 SANZ-CILLERO03	RVUE		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
774.5 ±0.7 ±1.5	500k	7 ACHASOV 02	SND	±	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ±0.5		23 PICH 01	RVUE		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

NODE=M009M5

NODE=M009M5

OCCUR=2

OCCUR=3

OCCUR=2

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
763.0±0.3±1.2	600k	24 ABELE 99E	CBAR	0±	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$

NODE=M009M7

NODE=M009M7

NODE=M009M3

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
766.5±1.1 OUR AVERAGE					
763.7±3.2		ABELE 97	CBAR		$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
768 ±9		AGUILAR...	91 EHS		$400 pp$
767 ±3	2935	25 CAPRARO 87	SPEC	-	$200 \pi^-\text{Cu} \rightarrow \pi^-\pi^0\text{Cu}$
761 ±5	967	25 CAPRARO 87	SPEC	-	$200 \pi^-\text{Pb} \rightarrow \pi^-\pi^0\text{Pb}$
771 ±4		HUSTON 86	SPEC	+	$202 \pi^+\text{A} \rightarrow \pi^+\pi^0\text{A}$
766 ±7	6500	26 BYERLY 73	OSPK	-	$5 \pi^- p$
766.8±1.5	9650	27 PISUT 68	RVUE	-	$1.7-3.2 \pi^- p, t < 10$
767 ±6	900	25 EISNER 67	HBC	-	$4.2 \pi^- p, t < 10$

NODE=M009M2

NODE=M009M2

OCCUR=2

NEUTRAL ONLY, PHOTOPRODUCED

NODE=M009M0P
NODE=M009M0P

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
769.0± 1.0 OUR AVERAGE				
771 ± 2 ⁺² / ₋₁	63.5k	28 ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+\pi^-p$
770 ± 2 ±1	79k	29 BREITWEG	98B ZEUS	50-100 γp
767.6± 2.7		BARTALUCCI	78 CNTR	$\gamma p \rightarrow e^+e^-p$
775 ± 5		GLADDING	73 CNTR	2.9-4.7 γp
767 ± 4	1930	BALLAM	72 HBC	2.8 γp
770 ± 4	2430	BALLAM	72 HBC	4.7 γp
765 ±10		ALVENSLEB...	70 CNTR	$\gamma A, t < 0.01$
767.7± 1.9	140k	BIGGS	70 CNTR	$< 4.1 \gamma C \rightarrow \pi^+\pi^-C$
765 ± 5	4000	ASBURY	67B CNTR	$\gamma + Pb$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
771 ± 2	79k	30 BREITWEG	98B ZEUS	50-100 γp

OCCUR=2


OCCUR=2

NEUTRAL ONLY, OTHER REACTIONS

NODE=M009M0R
NODE=M009M0R

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
769.0±0.9 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.					
765 ±6		BERTIN	97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
773 ±1.6		WEIDENAUER	93	ASTE	$\bar{p}p \rightarrow \pi^+\pi^-\omega$
762.6±2.6		AGUILAR-...	91	EHS	400 pp
770 ±2		31 HEYN	81	RVUE	Pion form factor
768 ±4		32,33 BOHACIK	80	RVUE	0
769 ±3		26 WICKLUND	78	ASPK	0 3,4,6 $\pi^\pm N$
768 ±1	76000	DEUTSCH...	76	HBC	0 16 π^+p
767 ±4	4100	ENGLER	74	DBC	0 6 $\pi^+n \rightarrow \pi^+\pi^-p$
775 ±4	32000	32 PROTOPOP...	73	HBC	0 7.1 $\pi^+p, t < 0.4$
764 ±3	6800	RATCLIFF	72	ASPK	0 15 $\pi^-p, t < 0.3$
774 ±3	1700	REYNOLDS	69	HBC	0 2.26 π^-p
769.2±1.5	13300	34 PISUT	68	RVUE	0 1.7-3.2 $\pi^-p, t < 10$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
773.5±2.5		35 COLANGELO	01	RVUE	$\pi\pi \rightarrow \pi\pi$
762.3±0.5±1.2	600k	36 ABELE	99E	CBAR	0 0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
777 ±2	4943	37 ADAMS	97	E665	470 $\mu p \rightarrow \mu XB$
770 ±2		38 BOGOLYUB...	97	MIRA	32 $\bar{p}p \rightarrow \pi^+\pi^-X$
768 ±8		38 BOGOLYUB...	97	MIRA	32 $pp \rightarrow \pi^+\pi^-X$
761.1±2.9		DUBNICKA	89	RVUE	π form factor
777.4±2.0		39 CHABAUD	83	ASPK	0 17 π^-p polarized
769.5±0.7		32,33 LANG	79	RVUE	0
770 ±9		33 ESTABROOKS	74	RVUE	0 17 $\pi^-p \rightarrow \pi^+\pi^-n$
773.5±1.7	11200	25 JACOBS	72	HBC	0 2.8 π^-p
775 ±3	2250	HYAMS	68	OSPK	0 11.2 π^-p

OCCUR=2



- 1 Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
- 2 A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.
- 3 Supersedes ACHASOV 05A.
- 4 A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.
- 5 Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference.
- 6 Update of AKHMETSHIN 02.
- 7 Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.
- 8 From the GOUNARIS 68 parametrization of the pion form factor.
- 9 Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.
- 10 Without limitations on masses and widths.
- 11 Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0\pi\pi} = g_{\rho^\pm\pi\pi}$.
- 12 Using the data of BARKOV 85 in the hidden local symmetry model.
- 13 From the fit to $e^+e^- \rightarrow \pi^+\pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.
- 14 A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.
- 15 Applying the S-matrix formalism to the BARKOV 85 data.
- 16 Includes BARKOV 85 data. Model-dependent width definition.
- 17 $|F_\pi(0)|^2$ fixed to 1.
- 18 From the GOUNARIS 68 parametrization of the pion form factor.
- 19 The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.
- 20 $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
- 21 From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.
- 22 Using the data of BARATE 97M and the effective chiral Lagrangian.
- 23 From a fit of the model-independent parameterization of the pion form factor to the data of BARATE 97M.
- 24 Assuming the equality of ρ^+ and ρ^- masses and widths.
- 25 Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.
- 26 Phase shift analysis. Systematic errors added corresponding to spread of different fits.
- 27 From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.
- 28 Using the KUHN 90 parametrization of the pion form factor, neglecting ρ - ω interference.
- 29 From the parametrization according to SOEDING 66.
- 30 From the parametrization according to ROSS 66.
- 31 HEYN 81 includes all spacelike and timelike F_π values until 1978.
- 32 From pole extrapolation.
- 33 From phase shift analysis of GRAYER 74 data.
- 34 Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.
- 35 Breit-Wigner mass from a phase-shift analysis of HYAMS 73 and PROTOPODESCU 73 data.
- 36 Using relativistic Breit-Wigner and taking into account ρ - ω interference.
- 37 Systematic errors not evaluated.
- 38 Systematic effects not studied.
- 39 From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P -wave intensity. CHABAUD 83 includes data of GRAYER 74.

NODE=M009M0;LINKAGE=LE

NODE=M009M0;LINKAGE=AK
 NODE=M009M0;LINKAGE=AC
 NODE=M009M0;LINKAGE=SN

NODE=M009M5;LINKAGE=GS
 NODE=M009M5;LINKAGE=PT

NODE=M009M0;LINKAGE=CH
 NODE=M009M;LINKAGE=K

NODE=M009M;LINKAGE=DF
 NODE=M009M;LINKAGE=WO

NODE=M009M;LINKAGE=HC
 NODE=M009M;LINKAGE=K2
 NODE=M009M;LINKAGE=G8

NODE=M009M;LINKAGE=AB
 NODE=M009M;LINKAGE=AA
 NODE=M009M;LINKAGE=F
 NODE=M009M5;LINKAGE=FU
 NODE=M009M5;LINKAGE=GO
 NODE=M009M5;LINKAGE=SC
 NODE=M009M;LINKAGE=A6
 NODE=M009M;LINKAGE=K1

NODE=M009M5;LINKAGE=Z
 NODE=M009M;LINKAGE=PC

NODE=M009M;LINKAGE=LB
 NODE=M009M;LINKAGE=Z
 NODE=M009M;LINKAGE=X
 NODE=M009M;LINKAGE=A

NODE=M009M0P;LINKAGE=AB
 NODE=M009M;LINKAGE=B5
 NODE=M009M;LINKAGE=B6
 NODE=M009M;LINKAGE=B
 NODE=M009M;LINKAGE=C
 NODE=M009M;LINKAGE=H
 NODE=M009M;LINKAGE=R

NODE=M009M;LINKAGE=CL

NODE=M009M;LINKAGE=BL
 NODE=M009M;LINKAGE=A1
 NODE=M009M;LINKAGE=QQ
 NODE=M009M;LINKAGE=G

$m_{\rho(770)^0} - m_{\rho(770)^\pm}$

NODE=M009D


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.7±0.8 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
-2.4±0.8		40 SCHAEL	05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
0.4±0.7±0.6	1.98M	41 ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.3±1.1±2.0	500k	41 ACHASOV	02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.6±0.6±1.7	600k	ABELE	99E	CBAR	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
-4 ±4	3000	42 REYNOLDS	69	HBC	-0 $2.26 \pi^- p$
-5 ±5	3600	42 FOSTER	68	HBC	$\pm 0.0 \bar{p} p$
2.4±2.1	22950	43 PISUT	68	RVUE	$\pi N \rightarrow \rho N$

NODE=M009D

- 40 From the combined fit of the τ^- data from ANDERSON 00A and SCHAELE 05C and e^+e^- data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.
- 41 Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.
- 42 From quoted masses of charged and neutral modes.
- 43 Includes MALAMUD 69, ARMENISE 68, BATON 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65, CARMONY 64, GOLDBERGER 64, ABOLINS 63.

NODE=M009D;LINKAGE=SC

NODE=M009D;LINKAGE=CH
 NODE=M009D;LINKAGE=A
 NODE=M009D;LINKAGE=R



$m_{\rho(770)^+} - m_{\rho(770)^-}$

NODE=M009D1

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M009D1

••• We do not use the following data for averages, fits, limits, etc. •••

$1.5 \pm 0.8 \pm 0.7$ 1.98M 44 ALOISIO 03 KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M009D;LINKAGE=WO

44 Without limitations on masses and widths.

$\rho(770)$ RANGE PARAMETER

NODE=M009R

The range parameter R enters an energy-dependent correction to the width, of the form $(1 + q_r^2 R^2) / (1 + q^2 R^2)$, where q is the momentum of one of the pions in the $\pi\pi$ rest system. At resonance, $q = q_r$.

NODE=M009R

VALUE (GeV^{-1}) DOCUMENT ID TECN CHG COMMENT

NODE=M009R

$5.3^{+0.9}_{-0.7}$ CHABAUD 83 ASPK 0 17 $\pi^- p$ polarized

$\rho(770)$ WIDTH

NODE=M009220

We no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

NODE=M009220

NEUTRAL ONLY, e^+e^-

NODE=M009W0
 NODE=M009W0

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

147.8 ± 0.9 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.

149.59 ± 0.67 45 LEES 12G BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$

$145.98 \pm 0.75 \pm 0.50$ 900k 46 AKHMETSHIN 07 $e^+e^- \rightarrow \pi^+\pi^-$

$146.1 \pm 0.8 \pm 1.5$ 800k 47,48 ACHASOV 06 SND $e^+e^- \rightarrow \pi^+\pi^-$

$143.85 \pm 1.33 \pm 0.80$ 114k 49,50 AKHMETSHIN 04 CMD2 $e^+e^- \rightarrow \pi^+\pi^-$

$147.3 \pm 1.5 \pm 0.7$ 1.98M 51 ALOISIO 03 KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$


$151.1 \pm 2.6 \pm 3.0$ 500k 51 ACHASOV 02 SND 0 $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

150.5 ± 3.0 52 BARKOV 85 OLYA 0 $e^+e^- \rightarrow \pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

143.9 ±1.3 ±1.1	1.98M	53	ALOISIO	03	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	OCCUR=2
147.4 ±1.5 ±0.7	1.98M	54	ALOISIO	03	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	OCCUR=3
149.8 ±2.2 ±2.0	500k	55	ACHASOV	02	SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	OCCUR=3
147.9 ±1.5 ±7.5		56	BENAYOUN	98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$, $\mu^+\mu^-$	
153.5 ±1.3 ±4.6		57	GARDNER	98	RVUE	0.28-0.92 $e^+e^- \rightarrow \pi^+\pi^-$	
145.0 ±1.7		58	O'CONNELL	97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
142.5 ±3.5		59	BERNICA	94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
138 ±1		60	GESHKEN...	89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	

WEIGHTED AVERAGE
147.8±0.9 (Error scaled by 2.0)



CHARGED ONLY, τ DECAYS and e^+e^-

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

149.1±0.8 OUR FIT

149.1±0.8 OUR AVERAGE

148.1±0.4±1.7	5.4M	61,62	FUJIKAWA	08	BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$	
149.0±1.2		62,63	SCHAEEL	05c	ALEP	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$	
149.9±2.3±2.0	500k	51	ACHASOV	02	SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	OCCUR=2
150.4±1.4±1.4	87k	64,65	ANDERSON	00A	CLE2	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

143.7±1.3±1.2	1.98M	51	ALOISIO	03	KLOE	\pm 1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
142.9±1.3±1.4	1.98M	54	ALOISIO	03	KLOE	- 1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	OCCUR=2
144.7±1.4±1.2	1.98M	54	ALOISIO	03	KLOE	+ 1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	OCCUR=3
150.2±2.0 ^{+0.7} _{-1.6}		66	SANZ-CILLERO	03	RVUE	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$	
150.9±2.2±2.0	500k	55	ACHASOV	02	SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$	OCCUR=4

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

149.5±1.3	600k	67	ABELE	99E	CBAR	0± 0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
-----------	------	----	-------	-----	------	---	--

NODE=M009W5
NODE=M009W5

NODE=M009W7
NODE=M009W7

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
150.2± 2.4 OUR FIT					
150.2± 2.4 OUR AVERAGE					
152.8± 4.3		ABELE 97	CBAR		$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
155 ±11	2935	68 CAPRARO 87	SPEC	-	$200 \pi^- \text{Cu} \rightarrow \pi^- \pi^0 \text{Cu}$
154 ±20	967	68 CAPRARO 87	SPEC	-	$200 \pi^- \text{Pb} \rightarrow \pi^- \pi^0 \text{Pb}$
150 ± 5		HUSTON 86	SPEC	+	$202 \pi^+ \text{A} \rightarrow \pi^+ \pi^0 \text{A}$
146 ±12	6500	69 BYERLY 73	OSPK	-	$5 \pi^- p$
148.2± 4.1	9650	70 PISUT 68	RVUE	-	$1.7-3.2 \pi^- p, t < 10$
146 ±13	900	EISNER 67	HBC	-	$4.2 \pi^- p, t < 10$

NODE=M009W2
 NODE=M009W2

OCCUR=2

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
151.7± 2.6 OUR AVERAGE					
155 ± 5 ± 2	63.5k	71 ABRAMOWICZ12	ZEUS		$e p \rightarrow e \pi^+ \pi^- p$
146 ± 3 ± 13	79k	72 BREITWEG 98B	ZEUS		$50-100 \gamma p$
150.9± 3.0		BARTALUCCI 78	CNTR		$\gamma p \rightarrow e^+ e^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
138 ± 3	79k	73 BREITWEG 98B	ZEUS		$50-100 \gamma p$
147 ±11		GLADDING 73	CNTR		$2.9-4.7 \gamma p$
155 ±12	2430	BALLAM 72	HBC		$4.7 \gamma p$
145 ±13	1930	BALLAM 72	HBC		$2.8 \gamma p$
140 ± 5		ALVENSLEB... 70	CNTR		$\gamma A, t < 0.01$
146.1± 2.9	140k	BIGGS 70	CNTR		$< 4.1 \gamma C \rightarrow \pi^+ \pi^- C$
160 ±10		LANZEROTTI 68	CNTR		γp
130 ± 5	4000	ASBURY 67B	CNTR		$\gamma + \text{Pb}$

NODE=M009W0P
 NODE=M009W0P

OCCUR=2

OCCUR=2

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
150.9± 1.7 OUR AVERAGE Error includes scale factor of 1.1.					
122 ±20		BERTIN 97C	OBLX		$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
145.7± 5.3		WEIDENAUER 93	ASTE		$\bar{p} p \rightarrow \pi^+ \pi^- \omega$
144.9± 3.7		DUBNICKA 89	RVUE		π form factor
148 ± 6		74,75 BOHACIK 80	RVUE	0	
152 ± 9		69 WICKLUND 78	ASPK	0	$3,4,6 \pi^\pm p N$
154 ± 2	76000	DEUTSCH... 76	HBC	0	$16 \pi^+ p$
157 ± 8	6800	RATCLIFF 72	ASPK	0	$15 \pi^- p, t < 0.3$
143 ± 8	1700	REYNOLDS 69	HBC	0	$2.26 \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
147.0± 2.5	600k	76 ABELE 99E	CBAR	0	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
146 ± 3	4943	77 ADAMS 97	E665		$470 \mu p \rightarrow \mu X B$
160.0 ⁺ 4.1 - 4.0		78 CHABAUD 83	ASPK	0	$17 \pi^- p$ polarized
155 ± 1		79 HEYN 81	RVUE	0	π form factor
148.0± 1.3		74,75 LANG 79	RVUE	0	
146 ±14	4100	ENGLER 74	DBC	0	$6 \pi^+ n \rightarrow \pi^+ \pi^- p$
143 ±13		75 ESTABROOKS 74	RVUE	0	$17 \pi^- p \rightarrow \pi^+ \pi^- n$
160 ±10	32000	74 PROTOPOP... 73	HBC	0	$7.1 \pi^+ p, t < 0.4$
145 ±12	2250	68 HYAMS 68	OSPK	0	$11.2 \pi^- p$
163 ±15	13300	80 PISUT 68	RVUE	0	$1.7-3.2 \pi^- p, t < 10$

NODE=M009W0R
 NODE=M009W0R

⁴⁵ Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

NODE=M009W0;LINKAGE=LE

⁴⁶ A combined fit of AKHMETSIN 07, AULCHENKO 06, and AULCHENKO 05.

NODE=M009W;LINKAGE=AK

⁴⁷ Supersedes ACHASOV 05A.

NODE=M009W0;LINKAGE=AC

⁴⁸ A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

NODE=M009W0;LINKAGE=SN

⁴⁹ Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference.

NODE=M009W5;LINKAGE=GS

⁵⁰ From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSIN 02.

NODE=M009W5;LINKAGE=P2

⁵¹ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

NODE=M009W;LINKAGE=CH

⁵² From the GOUNARIS 68 parametrization of the pion form factor.

NODE=M009W;LINKAGE=K

⁵³ Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.

NODE=M009W;LINKAGE=DF

⁵⁴ Without limitations on masses and widths.

NODE=M009W;LINKAGE=WO

⁵⁵ Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0 \pi \pi} = g_{\rho^\pm \pi \pi}$.

NODE=M009W;LINKAGE=HC

⁵⁶ Using the data of BARKOV 85 in the hidden local symmetry model.

NODE=M009W;LINKAGE=K2

- 57 From the fit to $e^+e^- \rightarrow \pi^+\pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.
- 58 A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.
- 59 Applying the S-matrix formalism to the BARKOV 85 data.
- 60 Includes BARKOV 85 data. Model-dependent width definition.
- 61 $|F_\pi(0)|^2$ fixed to 1.
- 62 From the GOUNARIS 68 parametrization of the pion form factor.
- 63 The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.
- 64 $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
- 65 From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.
- 66 Using the data of BARATE 97M and the effective chiral Lagrangian.
- 67 Assuming the equality of ρ^+ and ρ^- masses and widths.
- 68 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- 69 Phase shift analysis. Systematic errors added corresponding to spread of different fits.
- 70 From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.
- 71 Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
- 72 From the parametrization according to SOEDING 66.
- 73 From the parametrization according to ROSS 66.
- 74 From pole extrapolation.
- 75 From phase shift analysis of GRAYER 74 data.
- 76 Using relativistic Breit-Wigner and taking into account $\rho-\omega$ interference.
- 77 Systematic errors not evaluated.
- 78 From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P -wave intensity. CHABAUD 83 includes data of GRAYER 74.
- 79 HEYN 81 includes all spacelike and timelike F_π values until 1978.
- 80 Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.

NODE=M009W;LINKAGE=G8

NODE=M009W;LINKAGE=AB

NODE=M009W;LINKAGE=AA

NODE=M009W;LINKAGE=F

NODE=M009W5;LINKAGE=FU

NODE=M009W5;LINKAGE=GO

NODE=M009W5;LINKAGE=SC

NODE=M009W;LINKAGE=A6

NODE=M009W;LINKAGE=K1

NODE=M009W5;LINKAGE=Z

NODE=M009W;LINKAGE=LB

NODE=M009W;LINKAGE=Z

NODE=M009W;LINKAGE=X

NODE=M009W;LINKAGE=A

NODE=M009W0P;LINKAGE=AB

NODE=M009W;LINKAGE=B5

NODE=M009W;LINKAGE=B6

NODE=M009W;LINKAGE=C

NODE=M009W;LINKAGE=H

NODE=M009W;LINKAGE=BL

NODE=M009W;LINKAGE=A1

NODE=M009W;LINKAGE=G

NODE=M009W;LINKAGE=B

NODE=M009W;LINKAGE=R

 $\Gamma_{\rho(770)^0} - \Gamma_{\rho(770)^\pm}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.3±1.3 OUR AVERAGE				Error includes scale factor of 1.4.
-0.2±1.0		81 SCHAEEL	05C ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
3.6±1.8±1.7	1.98M	82 ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M009W6

NODE=M009W6

 $\Gamma_{\rho(770)^+} - \Gamma_{\rho(770)^-}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.8±2.0±0.5	1.98M	83 ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M009W16

NODE=M009W16

- 81 From the combined fit of the τ^- data from ANDERSON 00A and SCHAEEL 05C and e^+e^- data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

NODE=M009W6;LINKAGE=SC

- 82 Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

NODE=M009W6;LINKAGE=CH

NODE=M009W16;LINKAGE=WO

- 83 Without limitations on masses and widths.

 $\rho(770)$ DECAY MODES

NODE=M009225;NODE=M009

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\pi\pi$	~ 100	%
$\rho(770)^\pm$ decays		
Γ_2 $\pi^\pm\pi^0$	~ 100	%
Γ_3 $\pi^\pm\gamma$	(4.5 ± 0.5)	$\times 10^{-4}$ S=2.2
Γ_4 $\pi^\pm\eta$	< 6	$\times 10^{-3}$ CL=84%
Γ_5 $\pi^\pm\pi^+\pi^-\pi^0$	< 2.0	$\times 10^{-3}$ CL=84%

DESIG=1;OUR EVAL;→ UNCHECKED ←

NODE=M009;CLUMP=A

DESIG=11;OUR EVAL;→ UNCHECKED ←

DESIG=3

DESIG=5

DESIG=21

$\rho(770)^0$ decays

Γ_6	$\pi^+ \pi^-$	~ 100	%
Γ_7	$\pi^+ \pi^- \gamma$	(9.9 ± 1.6)	$\times 10^{-3}$
Γ_8	$\pi^0 \gamma$	(6.0 ± 0.8)	$\times 10^{-4}$
Γ_9	$\eta \gamma$	(3.00 ± 0.20)	$\times 10^{-4}$
Γ_{10}	$\pi^0 \pi^0 \gamma$	(4.5 ± 0.8)	$\times 10^{-5}$
Γ_{11}	$\mu^+ \mu^-$	[a] (4.55 ± 0.28)	$\times 10^{-5}$
Γ_{12}	$e^+ e^-$	[a] (4.72 ± 0.05)	$\times 10^{-5}$
Γ_{13}	$\pi^+ \pi^- \pi^0$	$(1.01^{+0.54}_{-0.36} \pm 0.34)$	$\times 10^{-4}$
Γ_{14}	$\pi^+ \pi^- \pi^+ \pi^-$	(1.8 ± 0.9)	$\times 10^{-5}$
Γ_{15}	$\pi^+ \pi^- \pi^0 \pi^0$	(1.6 ± 0.8)	$\times 10^{-5}$
Γ_{16}	$\pi^0 e^+ e^-$	< 1.2	$\times 10^{-5}$ CL=90%
Γ_{17}	$\eta e^+ e^-$		

NODE=M009;CLUMP=B
DESIG=12;OUR EVAL;→ UNCHECKED ←
DESIG=60
DESIG=40
DESIG=8
DESIG=80
DESIG=6
DESIG=4
DESIG=7;OUR EVAL;→ UNCHECKED ←
DESIG=22
DESIG=30
DESIG=9
DESIG=10

[a] The $\omega\rho$ interference is then due to $\omega\rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+ \mu^-) = \Gamma(\rho^0 \rightarrow e^+ e^-) \times 0.99785$.

LINKAGE=MD2

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 10 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 10.7$ for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc}
 x_3 & -100 & \\
 \Gamma & 15 & -15 \\
 & x_2 & x_3
 \end{array}$$

	Mode	Rate (MeV)	Scale factor
Γ_2	$\pi^\pm \pi^0$	150.2 \pm 2.4	
Γ_3	$\pi^\pm \gamma$	0.068 \pm 0.007	2.3

DESIG=11
DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 7 branching ratios uses 21 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 6.0$ for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	-100								
x_8	-5	0							
x_9	-1	0	1						
x_{10}	-1	0	0	0					
x_{11}	2	-3	0	0	0				
x_{12}	0	0	-8	-9	0	0			
x_{14}	-1	0	0	0	0	0	0		
Γ	0	0	4	5	0	0	-54	0	
	x_6	x_7	x_8	x_9	x_{10}	x_{11}	x_{12}	x_{14}	

Mode	Rate (MeV)	
Γ_6 $\pi^+ \pi^-$	147.5 \pm 0.9	DESIG=12
Γ_7 $\pi^+ \pi^- \gamma$	1.48 \pm 0.24	DESIG=60
Γ_8 $\pi^0 \gamma$	0.089 \pm 0.012	DESIG=40
Γ_9 $\eta \gamma$	0.0447 \pm 0.0031	DESIG=8
Γ_{10} $\pi^0 \pi^0 \gamma$	0.0066 \pm 0.0012	DESIG=80
Γ_{11} $\mu^+ \mu^-$	[a] 0.0068 \pm 0.0004	DESIG=6
Γ_{12} $e^+ e^-$	[a] 0.00704 \pm 0.00006	DESIG=4
Γ_{14} $\pi^+ \pi^- \pi^+ \pi^-$	0.0027 \pm 0.0014	DESIG=22

$\rho(770)$ PARTIAL WIDTHS

NODE=M009230


$\Gamma(\pi^\pm \gamma)$

Γ_3

NODE=M009W3
NODE=M009W3

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
68 \pm 7 OUR FIT				Error includes scale factor of 2.3.
68 \pm 7 OUR AVERAGE				Error includes scale factor of 2.2. See the ideogram below.
81 \pm 4 \pm 4	CAPRARO	87	SPEC	- 200 $\pi^- A \rightarrow \pi^- \pi^0 A$
59.8 \pm 4.0	HUSTON	86	SPEC	+ 202 $\pi^+ A \rightarrow \pi^+ \pi^0 A$
71 \pm 7	JENSEN	83	SPEC	- 156-260 $\pi^- A \rightarrow \pi^- \pi^0 A$

WEIGHTED AVERAGE
68 \pm 7 (Error scaled by 2.2)



$\Gamma(e^+e^-)$ Γ_{12}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.04 ± 0.06 OUR FIT				
7.04 ± 0.06 OUR AVERAGE				
7.048 ± 0.057 ± 0.050	900k	⁸⁴ AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
7.06 ± 0.11 ± 0.05	114k	^{85,86} AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
6.77 ± 0.10 ± 0.30		BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.12 ± 0.02 ± 0.11	800k	⁸⁷ ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
6.3 ± 0.1		⁸⁸ BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$, $\mu^+\mu^-$

NODE=M009W4
NODE=M009W4

OCCUR=2

 $\Gamma(\pi^0\gamma)$ Γ_8

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
77 ± 17 ± 11	36500	⁸⁹ ACHASOV 03	SND	0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
121 ± 31		DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

NODE=M009W31
NODE=M009W31 $\Gamma(\eta\gamma)$ Γ_9

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
62 ± 17		⁹⁰ DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$

NODE=M009W32
NODE=M009W32 $\Gamma(\pi^+\pi^-\pi^+\pi^-)$ Γ_{14}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.8 ± 1.4 ± 0.5	153	AKHMETSHIN 00	CMD2	0.6–0.97 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M009W33
NODE=M009W33

- ⁸⁴ A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.
⁸⁵ Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference.
⁸⁶ From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02.
⁸⁷ Supersedes ACHASOV 05A.
⁸⁸ Using the data of BARKOV 85 in the hidden local symmetry model.
⁸⁹ Using $\Gamma_{\text{total}} = 147.9 \pm 1.3$ MeV and $B(\rho \rightarrow \pi^0\gamma)$ from ACHASOV 03.
⁹⁰ Solution corresponding to constructive ω - ρ interference.

NODE=M009W4;LINKAGE=AK
NODE=M009W4;LINKAGE=GS
NODE=M009W4;LINKAGE=P2
NODE=M009W4;LINKAGE=AC
NODE=M009W4;LINKAGE=K2
NODE=M009W31;LINKAGE=AV
NODE=M009W32;LINKAGE=L $\rho(770) \Gamma(e^+e^-)\Gamma(i)/\Gamma^2(\text{total})$

NODE=M009233

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_6/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.876 ± 0.023 ± 0.064	800k	^{91,92} ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.72 ± 0.02		⁹³ BENAYOUN 10	RVUE	0.4–1.05 e^+e^-

NODE=M009G4
NODE=M009G4

- ⁹¹ Supersedes ACHASOV 05A.
⁹² A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.
⁹³ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M009G4;LINKAGE=AC
NODE=M009G4;LINKAGE=SN

NODE=M009G4;LINKAGE=BE

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
1.42 ± 0.10 OUR FIT				
1.45 ± 0.12 OUR AVERAGE				
1.32 ± 0.14 ± 0.08	33k	⁹⁴ ACHASOV 07B	SND	0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
1.50 ± 0.65 ± 0.09	17.4k	⁹⁵ AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
1.61 ± 0.20 ± 0.11	23k	^{96,97} AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1.85 ± 0.49		⁹⁸ DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.05 ± 0.02		⁹⁹ BENAYOUN 10	RVUE	0.4–1.05 e^+e^-

NODE=M009G1
NODE=M009G1

- ⁹⁴ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
⁹⁵ From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
⁹⁶ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
⁹⁷ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
⁹⁸ Recalculated by us from the cross section in the peak.
⁹⁹ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M009G1;LINKAGE=AH

NODE=M009G;LINKAGE=AH
NODE=M009G;LINKAGE=AK
NODE=M009G;LINKAGE=BQNODE=M009G;LINKAGE=LP
NODE=M009G1;LINKAGE=BE

$$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \qquad \Gamma_{12}/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8 ±0.4 OUR FIT				
2.8 ±0.4 OUR AVERAGE				
2.90 $^{+0.60}_{-0.55}$ ±0.18	18680	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$
2.37 ±0.53 ±0.33	36500	¹⁰⁰ ACHASOV 03	SND	0.60-0.97 $e^+e^- \rightarrow \pi^0\gamma$
3.61 ±0.74 ±0.49	10625	¹⁰¹ DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$
1.875 ±0.026		¹⁰² BENAYOUN 10	RVUE	0.4-1.05 e^+e^-

NODE=M009G2
NODE=M009G2

¹⁰⁰ Using $\sigma_{\phi \rightarrow \pi^0\gamma}$ from ACHASOV 00 and $m_\rho = 775.97$ MeV in the model with the energy-independent phase of ρ - ω interference equal to $(-10.2 \pm 7.0)^\circ$.

NODE=M009G;LINKAGE=SH

¹⁰¹ Recalculated by us from the cross section in the peak.

NODE=M009G2;LINKAGE=LP

¹⁰² A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M009G2;LINKAGE=BE

$$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \qquad \Gamma_{12}/\Gamma \times \Gamma_{13}/\Gamma$$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
0.903 ±0.076		¹⁰³ BENAYOUN 10	RVUE	0.4-1.05 e^+e^-
4.58 $^{+2.46}_{-1.64}$ ±1.56	1.2M	¹⁰⁴ ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M009G3
NODE=M009G3

¹⁰³ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M009G3;LINKAGE=BE

¹⁰⁴ Statistical significance is less than 3σ .

NODE=M009G3;LINKAGE=AC

$\rho(770)$ BRANCHING RATIOS

NODE=M009235

$$\Gamma(\pi^\pm\eta)/\Gamma(\pi\pi) \qquad \Gamma_4/\Gamma_1$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<60	84	FERBEL 66	HBC	±	$\pi^\pm p$ above 2.5

NODE=M009R4
NODE=M009R4

$$\Gamma(\pi^\pm\pi^+\pi^-\pi^0)/\Gamma(\pi\pi) \qquad \Gamma_5/\Gamma_1$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<20	84	FERBEL 66	HBC	±	$\pi^\pm p$ above 2.5

NODE=M009R1
NODE=M009R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

35 ± 40 JAMES 66 HBC + 2.1 $\pi^+ p$

$$\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-) \qquad \Gamma_{11}/\Gamma_6$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.60 ±0.28 OUR FIT			
4.6 ±0.2 ±0.2	ANTIPOV 89	SIGM	$\pi^- \text{Cu} \rightarrow \mu^+\mu^-\pi^- \text{Cu}$

NODE=M009R5
NODE=M009R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.2 $^{+1.6}_{-3.6}$ ¹⁰⁵ ROTHWELL 69 CNTR Photoproduction

5.6 ±1.5 ¹⁰⁶ WEHMANN 69 OSPK 12 $\pi^- \text{C, Fe}$

9.7 $^{+3.1}_{-3.3}$ ¹⁰⁷ HYAMS 67 OSPK 11 $\pi^- \text{Li, H}$

$$\Gamma(e^+e^-)/\Gamma(\pi\pi) \qquad \Gamma_{12}/\Gamma_1$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.40 ±0.05	¹⁰⁸ BENAKSAS 72	OSPK	$e^+e^- \rightarrow \pi^+\pi^-$

NODE=M009R3
NODE=M009R3

$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \qquad \Gamma_9/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3.00 ±0.21 OUR FIT					
2.90 ±0.32 OUR AVERAGE					

NODE=M009R7
NODE=M009R7

2.79 ±0.34 ±0.03 33k ¹⁰⁹ACHASOV 07B SND 0.6-1.38 $e^+e^- \rightarrow \eta\gamma$

3.6 ±0.9 ¹¹⁰ANDREWS 77 CNTR 0 6.7-10 γCu

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.21 ±1.39 ±0.20 17.4k ^{111,112}AKHMETSHIN 05 CMD2 0.60-1.38 $e^+e^- \rightarrow \eta\gamma$

3.39 ±0.42 ±0.23 ^{110,113,114}AKHMETSHIN 01B CMD2 $e^+e^- \rightarrow \eta\gamma$

1.9 $^{+0.6}_{-0.8}$ ¹¹⁵BENAYOUN 96 RVUE 0.54-1.04 $e^+e^- \rightarrow \eta\gamma$

4.0 ±1.1 ^{110,112}DOLINSKY 89 ND $e^+e^- \rightarrow \eta\gamma$

$$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.8±0.9 OUR FIT					
1.8±0.9±0.3		153	AKHMETSHIN 00	CMD2	0.6-0.97 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••					
<20		90	KURDADZE 88	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M009R13
NODE=M009R13

$$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma(\pi\pi) \quad \Gamma_{14}/\Gamma_1$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••					
<15	90	ERBE 69	HBC	0	2.5-5.8 γp
<20		CHUNG 68	HBC	0	3.2,4.2 $\pi^- p$
<20	90	HUSON 68	HLBC	0	16.0 $\pi^- p$
<80		JAMES 66	HBC	0	2.1 $\pi^+ p$

NODE=M009R11
NODE=M009R11

$$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••					
1.01 ^{+0.54} _{-0.36} ±0.34		1.2M 116	ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<1.2	90		VASSERMAN 88B	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M009R10
NODE=M009R10

$$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi\pi) \quad \Gamma_{13}/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••					
~ 0.01		BRAMON 86	RVUE	0	$J/\psi \rightarrow \omega\pi^0$
<0.01	84	117 ABRAMS 71	HBC	0	3.7 $\pi^+ p$

NODE=M009R6
NODE=M009R6

$$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
1.60±0.74±0.18		118 ACHASOV 09A	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
< 4	90	AULCHENKO 87C	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
<20	90	KURDADZE 86	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

NODE=M009R8
NODE=M009R8

$$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
0.0099±0.0016 OUR FIT				
0.0099±0.0016		119 DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
••• We do not use the following data for averages, fits, limits, etc. •••				
0.0111±0.0014		120 VASSERMAN 88	ND	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<0.005	90	121 VASSERMAN 88	ND	$e^+e^- \rightarrow \pi^+\pi^-\gamma$

NODE=M009R12
NODE=M009R12

OCCUR=2

$$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
6.21 ^{+1.28} _{-1.18} ±0.39	18680	122,123 AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$
5.22±1.17±0.75	36500	123,124 ACHASOV 03	SND	0.60-0.97 $e^+e^- \rightarrow \pi^0\gamma$
6.8 ±1.7		125 BENAYOUN 96	RVUE	0.54-1.04 $e^+e^- \rightarrow \pi^0\gamma$
7.9 ±2.0		123 DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

NODE=M009R9
NODE=M009R9

$$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.2	90	ACHASOV 08	SND	0.36-0.97 $e^+e^- \rightarrow \pi^0 e^+ e^-$
<1.6		AKHMETSHIN 05A	CMD2	0.72-0.84 e^+e^-

NODE=M009R15
NODE=M009R15

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ VALUE (units 10^{-5})

DOCUMENT ID

TECN

COMMENT

NODE=M009R16
NODE=M009R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7

AKHMETSHIN 05A CMD2 0.72-0.84 $e^+ e^-$ $\Gamma(\pi^0 \pi^0 \gamma)/\Gamma_{\text{total}}$ Γ_{10}/Γ VALUE (units 10^{-5})

EVTS

DOCUMENT ID

TECN

COMMENT

NODE=M009R14
NODE=M009R14**4.5 ± 0.8 OUR FIT****4.5^{+0.9}_{-0.8} OUR AVERAGE**5.2^{+1.5}_{-1.3} ± 0.6 190 126 AKHMETSHIN 04B CMD2 0.6^{-0.97}₀ $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

OCCUR=2

4.1^{+1.0}_{-0.9} ± 0.3 295 127 ACHASOV 02F SND 0.36^{-0.97}₀ $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.8^{+3.4}_{-1.8} ± 0.5 63 128 ACHASOV 00G SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$ 105 Possibly large ρ - ω interference leads us to increase the minus error.

NODE=M009R5;LINKAGE=R

106 Result contains $11 \pm 11\%$ correction using SU(3) for central value. The error on the correction takes account of possible ρ - ω interference and the upper limit agrees with the upper limit of $\omega \rightarrow \mu^+ \mu^-$ from this experiment.

NODE=M009R5;LINKAGE=W

107 HYAMS 67's mass resolution is 20 MeV. The ω region was excluded.

NODE=M009R5;LINKAGE=H

108 The ρ' contribution is not taken into account.

NODE=M009R;LINKAGE=KS

109 ACHASOV 07B reports $[\Gamma(\rho(770) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\rho(770) \rightarrow e^+ e^-)] = (1.32 \pm 0.14 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\rho(770) \rightarrow e^+ e^-) = (4.72 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

NODE=M009R7;LINKAGE=AO

110 Solution corresponding to constructive ω - ρ interference.

NODE=M009R7;LINKAGE=A

111 Using $B(\rho \rightarrow e^+ e^-) = (4.67 \pm 0.09) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

NODE=M009R;LINKAGE=AK

112 Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

NODE=M009R7;LINKAGE=AZ

113 The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

NODE=M009R;LINKAGE=BQ

114 Using $B(\rho \rightarrow e^+ e^-) = (4.75 \pm 0.10) \times 10^{-5}$ from AKHMETSHIN 02 and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

NODE=M009R;LINKAGE=BX

115 Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution. Constructive ρ - ω interference solution.

NODE=M009R7;LINKAGE=C

116 Statistical significance is less than 3σ .

NODE=M009R;LINKAGE=NS

117 Model dependent, assumes $l = 1, 2, \text{ or } 3$ for the 3π system.

NODE=M009R6;LINKAGE=G

118 Assuming no interference between the ρ and ω contributions.

NODE=M009R8;LINKAGE=AC

119 Bremsstrahlung from a decay pion and for photon energy above 50 MeV.

NODE=M009R12;LINKAGE=J

120 Superseded by DOLINSKY 91.

NODE=M009R12;LINKAGE=I

121 Structure radiation due to quark rearrangement in the decay.

NODE=M009R12;LINKAGE=N

122 Using $B(\rho \rightarrow e^+ e^-) = (4.67 \pm 0.09) \times 10^{-5}$.

NODE=M009R9;LINKAGE=AK

123 Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}^2$.

NODE=M009R9;LINKAGE=BZ

124 Using $B(\rho \rightarrow e^+ e^-) = (4.54 \pm 0.10) \times 10^{-5}$.

NODE=M009R9;LINKAGE=AS

125 Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

NODE=M009R9;LINKAGE=A

126 This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega\pi^0$, $\omega \rightarrow \pi^0\gamma$, and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0\pi^0$ with a branching ratio $(2.0^{+1.1}_{-0.9} \pm 0.3) \times 10^{-5}$ differing from zero by 2.0 standard deviations.

NODE=M009R14;LINKAGE=AH

127 This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega\pi^0$, $\omega \rightarrow \pi^0\gamma$ and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0\pi^0$ with a branching ratio $(1.9^{+0.9}_{-0.8} \pm 0.4) \times 10^{-5}$ differing from zero by 2.4 standard deviations. Supersedes ACHASOV 00G.

NODE=M009R;LINKAGE=FF

128 Superseded by ACHASOV 02F.

NODE=M009R;LINKAGE=GF

$\rho(770)$ REFERENCES

NODE=M009

ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53101
ACHASOV	08	Translated from ZETF 136 442.			
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=52258
FUJIKAWA	08	Translated from ZETF 134 80.			
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AKHMETSHIN	07	PL B648 28	R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
ACHASOV	06A	Translated from ZETF 130 437.			
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51513
ACHASOV	05A	Translated from ZETFP 84 491.			
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51045
AKHMETSHIN	05	Translated from ZETF 128 1201.			
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50508
ALOISIO	05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=51112
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51060
SCHAEEL	05C	Translated from ZETFP 82 841.			
SCHAEEL	05C	PRPL 421 191	S. Schaeel <i>et al.</i>	(ALEPH Collab.)	REFID=50845
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49610
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49187
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
SANZ-CILLERO	03	EPJ C27 587	J.J. Sanz-Cillero, A. Pich		REFID=49399
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48549
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48816
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48565
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leytwyler		REFID=49180
PICH	01	PR D63 093005	A. Pich, J. Portoles		REFID=48313
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
ACHASOV	00G	Translated from ZETFP 72 411.			
ACHASOV	00G	JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47929
AKHMETSHIN	00	Translated from ZETFP 71 519.			
AKHMETSHIN	00	PL B475 190	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47421
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)	REFID=47468
ABELE	99E	PL B469 270	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47414
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)	REFID=45859
BREITWEG	98B	EPJ C2 247	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=46354
GARDNER	98	PR D57 2716	S. Gardner, H.B. O'Connell		REFID=46366
Also		PR D62 019903 (err.)	S. Gardner, H.B. O'Connell		REFID=47981
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ADAMS	97	ZPHY C74 237	M.R. Adams <i>et al.</i>	(E665 Collab.)	REFID=45533
BARATE	97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45622
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
BOGOLYUB...	97	PAN 60 46	M.Y. Bogolyubsky <i>et al.</i>	(MOSU, SERP)	REFID=45393
O'CONNELL	97	Translated from YAF 60 53.			
O'CONNELL	97	NP A623 559	H.B. O'Connell <i>et al.</i>	(ADLD)	REFID=45860
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
BERNICA	94	PR D50 4454	A. Bernica, G. Lopez Castro, J. Pestieau	(LOUV+)	REFID=44097
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ANTIPOV	89	ZPHY C42 185	Y.M. Antipov <i>et al.</i>	(SERP, JINR, BGNA+)	REFID=40739
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
GESHKEN...	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)	REFID=41017
KURDADZE	88	JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=41121
VASSERMAN	88	Translated from ZETFP 47 432.			
VASSERMAN	88	SJNP 47 1035	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=41019
VASSERMAN	88B	Translated from YAF 47 1635.			
VASSERMAN	88B	SJNP 48 480	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=41020
AULCHENKO	87C	Translated from YAF 48 753.			
AULCHENKO	87C	IYF 87-90 Preprint	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=41370
CAPRARO	87	NP B288 659	L. Capraro <i>et al.</i>	(CLER, FRAS, MILA+)	REFID=40003
BRAMON	86	PL B173 97	A. Bramon, J. Casulleras	(BARC)	REFID=22102
HUSTON	86	PR D33 3199	J. Huston <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=20137
KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=40287
BARKOV	85	Translated from ZETFP 43 497.			
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
JENSEN	83	PR D27 26	T. Jensen <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=20132
HEYN	81	ZPHY C7 169	M.F. Heyn, C.B. Lang	(GRAZ)	REFID=20129
BOHACIK	80	PR D21 1342	J. Bohacik, H. Kuhnelt	(SLOV, WIEN)	REFID=20128
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)	REFID=20126
BARTALUCCI	78	NC 44A 587	S. Bartalucci <i>et al.</i>	(DESY, FRAS)	REFID=20122
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
DEUTSCH...	76	NP B103 426	M. Deuschmann <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20119
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYR	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
BYERLY	73	PR D7 637	W.L. Byerly <i>et al.</i>	(MICH)	REFID=20104
GLADDING	73	PR D8 3721	G.E. Gladding <i>et al.</i>	(HARV)	REFID=20106

HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108
BALLAM	72	PR D5 545	J. Ballam <i>et al.</i>	(SLAC, LBL, TUFTS)	REFID=20094
BENAKSAS	72	PL 39B 289	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20096
JACOBS	72	PR D6 1291	L.D. Jacobs	(SACL)	REFID=20101
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)	REFID=20102
ABRAMS	71	PR D4 653	G.S. Abrams <i>et al.</i>	(LBL)	REFID=20090
ALVENSLEB...	70	PRL 24 786	H. Alvensleben <i>et al.</i>	(DESY)	REFID=20085
BIGGS	70	PRL 24 1197	P.J. Biggs <i>et al.</i>	(DARE)	REFID=20087
ERBE	69	PR 188 2060	R. Erbe <i>et al.</i>	(German Bubble Chamber Collab.)	REFID=20074
MALAMUD	69	Argonne Conf. 93	E.I. Malamud, P.E. Schlein	(UCLA)	REFID=20077
REYNOLDS	69	PR 184 1424	B.G. Reynolds <i>et al.</i>	(FSU)	REFID=20080
ROTHWELL	69	PRL 23 1521	P.L. Rothwell <i>et al.</i>	(NEAS)	REFID=20082
WEHMANN	69	PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)	REFID=20084
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
BATON	68	PR 176 1574	J.P. Baton, G. Laurens	(SACL)	REFID=20056
CHUNG	68	PR 165 1491	S.U. Chung <i>et al.</i>	(LRL)	REFID=20059
FOSTER	68	NP B6 107	M. Foster <i>et al.</i>	(CERN, CDEF)	REFID=20061
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054
HUSON	68	PL 28B 208	R. Huson <i>et al.</i>	(ORSAY, MILA, UCLA)	REFID=20062
HYAMS	68	NP B7 1	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20063
LANZEROTTI	68	PR 166 1365	L.J. Lanzerotti <i>et al.</i>	(HARV)	REFID=20068
PISUT	68	NP B6 325	J. Pisut, M. Roos	(CERN)	REFID=20070
ASBURY	67B	PRL 19 865	J.G. Asbury <i>et al.</i>	(DESY, COLU)	REFID=20038
BACON	67	PR 157 1263	T.C. Bacon <i>et al.</i>	(BNL)	REFID=20039
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
HUWE	67	PL 24B 252	D.O. Huwe <i>et al.</i>	(COLU)	REFID=20049
HYAMS	67	PL 24B 634	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20050
MILLER	67B	PR 153 1423	D.H. Miller <i>et al.</i>	(PURD)	REFID=20051
ALFF....	66	PR 145 1072	C. Alff-Steinberger <i>et al.</i>	(COLU, RUTG)	REFID=10762
FERBEL	66	PL 21 111	T. Ferbel	(ROCH)	REFID=20028
HAGOPIAN	66	PR 145 1128	V. Hagopian <i>et al.</i>	(PENN, SACL)	REFID=20030
HAGOPIAN	66B	PR 152 1183	V. Hagopian, Y.L. Pan	(PENN, LRL)	REFID=20031
JACOBS	66B	UCRL 16877	L.D. Jacobs	(LRL)	REFID=20033
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=10770
ROSS	66	PR 149 1172	M. Ross, L. Stodolsky		REFID=46380
SOEDING	66	PL B19 702	P. Soeding		REFID=46385
WEST	66	PR 149 1089	E. West <i>et al.</i>	(WISC)	REFID=20035
BLIEDEN	65	PL 19 444	H.R. Blieden <i>et al.</i>	(CERN MMS Collab.)	REFID=20016
CARMONY	64	PRL 12 254	D.D. Carmony <i>et al.</i>	(UCB)	REFID=20578
GOLDHABER	64	PRL 12 336	G. Goldhaber <i>et al.</i>	(LRL, UCB)	REFID=20013
ABOLINS	63	PRL 11 381	M.A. Abolins <i>et al.</i>	(UCSD)	REFID=20006

NODE=M001

 $\omega(782)$

$$I^G(J^{PC}) = 0^-(1^--)$$

 $\omega(782)$ MASS

NODE=M001M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
782.65±0.12 OUR AVERAGE		Error includes scale factor of 1.9. See the ideogram below.		
783.20±0.13±0.16	18680	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$
782.68±0.09±0.04	11200	¹ AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
782.79±0.08±0.09	1.2M	² ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
782.7 ±0.1 ±1.5	19500	WURZINGER 95	SPEC	1.33 $p\bar{d} \rightarrow {}^3\text{He}\omega$
781.96±0.17±0.80	11k	³ AMSLER 94C	CBAR	0.0 $\bar{p}p \rightarrow \omega\eta\pi^0$
782.08±0.36±0.82	3463	⁴ AMSLER 94C	CBAR	0.0 $\bar{p}p \rightarrow \omega\eta\pi^0$
781.96±0.13±0.17	15k	AMSLER 93B	CBAR	0.0 $\bar{p}p \rightarrow \omega\pi^0\pi^0$
782.4 ±0.2	270k	WEIDENAUER 93	ASTE	$\bar{p}p \rightarrow 2\pi^+2\pi^-\pi^0$
782.2 ±0.4	1488	KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
782.4 ±0.5	7000	⁵ KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
781.91±0.24		⁶ LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
781.78±0.10		⁷ BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
783.3 ±0.4	433	CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
782.5 ±0.8	33260	ROOS 80	RVUE	0.0-3.6 $\bar{p}p$
782.6 ±0.8	3000	BENKHEIRI 79	OMEG	9-12 $\pi^\pm p$
781.8 ±0.6	1430	COOPER 78B	HBC	0.7-0.8 $\bar{p}p \rightarrow 5\pi$
782.7 ±0.9	535	VANAPEL... 78	HBC	7.2 $\bar{p}p \rightarrow \bar{p}p\omega$
783.5 ±0.8	2100	GESSAROLI 77	HBC	11 $\pi^-p \rightarrow \omega n$
782.5 ±0.8	418	AGUILAR... 72B	HBC	3.9,4.6 K^-p
783.4 ±1.0	248	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K^+K^-\omega$
781.0 ±0.6	510	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K_1^+K_1^-\omega$
783.7 ±1.0	3583	⁸ COYNE 71	HBC	3.7 $\pi^+p \rightarrow \rho\pi^+\pi^+\pi^-\pi^0$
784.1 ±1.2	750	ABRAMOVI... 70	HBC	3.9 $\pi^-\rho$
783.2 ±1.6		⁹ BIGGS 70B	CNTR	<4.1 $\gamma C \rightarrow \pi^+\pi^-\omega$
782.4 ±0.5	2400	BIZZARRI 69	HBC	0.0 $\bar{p}p$

NODE=M001M

OCCUR=2

OCCUR=2


OCCUR=2

- 1 Update of AKHMETSHIN 00C.
- 2 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- 3 From the $\eta \rightarrow \gamma\gamma$ decay.
- 4 From the $\eta \rightarrow 3\pi^0$ decay.
- 5 Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.
- 6 From the $\rho-\omega$ interference in the $\pi^+\pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.
- 7 Systematic uncertainties underestimated.
- 8 From best-resolution sample of COYNE 71.
- 9 From ω - ρ interference in the $\pi^+\pi^-$ mass spectrum assuming ω width 12.6 MeV.

NODE=M001M;LINKAGE=PT
 NODE=M001M;LINKAGE=VH

NODE=M001M;LINKAGE=S1
 NODE=M001M;LINKAGE=S2
 NODE=M001M;LINKAGE=B
 NODE=M001M;LINKAGE=LE

NODE=M001M;LINKAGE=KB
 NODE=M001M;LINKAGE=D
 NODE=M001M;LINKAGE=F



$\omega(782)$ WIDTH

NODE=M001W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.49±0.08 OUR AVERAGE				
8.68±0.23±0.10	11200	1 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.68±0.04±0.15	1.2M	2 ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.2 ±0.3	19500	WURZINGER 95	SPEC	1.33 $pd \rightarrow {}^3\text{He}\omega$
8.4 ±0.1		3 AULCHENKO 87	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.30±0.40		BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.8 ±0.9	1488	KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0 ±0.8	433	CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.1 ±0.8	451	BENAKSAS 72B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
8.13±0.45		4 LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
12 ±2	1430	COOPER 78B	HBC	0.7-0.8 $\bar{p}p \rightarrow 5\pi$
9.4 ±2.5	2100	GESSAROLI 77	HBC	11 $\pi^-p \rightarrow \omega n$
10.22±0.43	20000	5 KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
13.3 ±2	418	AGUILAR-... 72B	HBC	3.9,4.6 K^-p
10.5 ±1.5		BORENSTEIN 72	HBC	2.18 K^-p
7.70±0.9 ±1.15	940	BROWN 72	MMS	2.5 $\pi^-p \rightarrow n\text{MM}$
10.3 ±1.4	510	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K_1^+K_1^-\omega$
12.8 ±3.0	248	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K^+K^-\omega$
9.5 ±1.0	3583	COYNE 71	HBC	3.7 $\pi^+p \rightarrow \rho\pi^+\pi^+\pi^-\pi^0$

NODE=M001W

OCCUR=2

¹ Update of AKHMETSHIN 00C.

² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

³ Relativistic Breit-Wigner includes radiative corrections.

⁴ From the $\rho-\omega$ interference in the $\pi^+\pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.

⁵ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.

NODE=M001W;LINKAGE=PT
NODE=M001W;LINKAGE=VH

NODE=M001W;LINKAGE=D
NODE=M001W;LINKAGE=LE

NODE=M001W;LINKAGE=B

NODE=M001215;NODE=M001

$\omega(782)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 $\pi^+\pi^-\pi^0$	$(89.2 \pm 0.7) \%$		DESIG=1
Γ_2 $\pi^0\gamma$	$(8.28 \pm 0.28) \%$	S=2.1	DESIG=3
Γ_3 $\pi^+\pi^-$	$(1.53^{+0.11}_{-0.13}) \%$	S=1.2	DESIG=2
Γ_4 neutrals (excluding $\pi^0\gamma$)	$(8 \pm \frac{8}{5}) \times 10^{-3}$	S=1.1	DESIG=13
Γ_5 $\eta\gamma$	$(4.6 \pm 0.4) \times 10^{-4}$	S=1.1	DESIG=6
Γ_6 $\pi^0 e^+ e^-$	$(7.7 \pm 0.6) \times 10^{-4}$		DESIG=14
Γ_7 $\pi^0 \mu^+ \mu^-$	$(1.3 \pm 0.4) \times 10^{-4}$	S=2.1	DESIG=11
Γ_8 $\eta e^+ e^-$			DESIG=18
Γ_9 $e^+ e^-$	$(7.28 \pm 0.14) \times 10^{-5}$	S=1.3	DESIG=7
Γ_{10} $\pi^+\pi^-\pi^0\pi^0$	$< 2 \times 10^{-4}$	CL=90%	DESIG=12
Γ_{11} $\pi^+\pi^-\gamma$	$< 3.6 \times 10^{-3}$	CL=95%	DESIG=4
Γ_{12} $\pi^+\pi^-\pi^+\pi^-$	$< 1 \times 10^{-3}$	CL=90%	DESIG=15
Γ_{13} $\pi^0\pi^0\gamma$	$(6.6 \pm 1.1) \times 10^{-5}$		DESIG=5
Γ_{14} $\eta\pi^0\gamma$	$< 3.3 \times 10^{-5}$	CL=90%	DESIG=17
Γ_{15} $\mu^+\mu^-$	$(9.0 \pm 3.1) \times 10^{-5}$		DESIG=8
Γ_{16} 3γ	$< 1.9 \times 10^{-4}$	CL=95%	DESIG=10
Charge conjugation (C) violating modes			
Γ_{17} $\eta\pi^0$	C $< 2.1 \times 10^{-4}$	CL=90%	NODE=M001;CLUMP=A DESIG=9
Γ_{18} $2\pi^0$	C $< 2.1 \times 10^{-4}$	CL=90%	DESIG=193
Γ_{19} $3\pi^0$	C $< 2.3 \times 10^{-4}$	CL=90%	DESIG=16

CONSTRAINED FIT INFORMATION

An overall fit to 15 branching ratios uses 51 measurements and one constraint to determine 10 parameters. The overall fit has a $\chi^2 = 51.8$ for 42 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	22								
x_3	-18	-4							
x_4	-92	-56	1						
x_5	7	7	-1	-9					
x_6	-1	0	0	0	0				
x_7	-1	0	0	0	0	0			
x_9	-38	-33	7	44	-21	0	0		
x_{13}	1	4	0	-2	0	0	0	-1	
x_{15}	0	0	0	0	0	0	0	0	0
	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_9	x_{13}

$\omega(782)$ PARTIAL WIDTHS

NODE=M001218

 $\Gamma(\pi^0\gamma)$ Γ_2

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
880±50	7815	¹ ACHASOV	13	SND 1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
788±12±27	36500	² ACHASOV	03	SND 0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
764±51	10625	DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\gamma$

¹ Systematic uncertainty not estimated.² Using $\Gamma_\omega = 8.44 \pm 0.09$ MeV and $B(\omega \rightarrow \pi^0\gamma)$ from ACHASOV 03.NODE=M001W1
NODE=M001W1NODE=M001W1;LINKAGE=AC
NODE=M001W1;LINKAGE=AD $\Gamma(\eta\gamma)$ Γ_5

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
6.1±2.5	¹ DOLINSKY	89	ND $e^+e^- \rightarrow \eta\gamma$

¹ Using $\Gamma_\omega = 8.4 \pm 0.1$ MeV and $B(\omega \rightarrow \eta\gamma)$ from DOLINSKY 89.NODE=M001W2
NODE=M001W2

NODE=M001W2;LINKAGE=DA

 $\Gamma(e^+e^-)$ Γ_9

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.60 ± 0.02 OUR EVALUATION				
0.591±0.015	11200	^{1,2} AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.653±0.003±0.021	1.2M	³ ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.600±0.031	10625	DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\gamma$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.891 \pm 0.007$ and $\Gamma_{\text{total}} = 8.44 \pm 0.09$ MeV.² Update of AKHMETSHIN 00C.³ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.NODE=M001W7
NODE=M001W7
→ UNCHECKED ←NODE=M001W7;LINKAGE=3P
NODE=M001W7;LINKAGE=PT
NODE=M001W;LINKAGE=VF $\omega(782) \Gamma(e^+e^-)\Gamma(i)/\Gamma^2(\text{total})$

NODE=M001225

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.49±0.11 OUR FIT Error includes scale factor of 1.3.				
6.38±0.10 OUR AVERAGE Error includes scale factor of 1.1.				
6.24±0.11±0.08	11.2k	¹ AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
6.70±0.06±0.27		AUBERT,B	04N	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
6.74±0.04±0.24	1.2M	^{2,3} ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
6.37±0.35		² DOLINSKY	89	ND $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
6.45±0.24		² BARKOV	87	CMD $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.79±0.42	1488	² KURDADZE	83B	OLYA $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.89±0.54	433	² CORDIER	80	DM1 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
7.54±0.84	451	² BENAKSAS	72B	OSPK $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

••• We do not use the following data for averages, fits, limits, etc. •••

6.20±0.13 ⁴BENAYOUN 10 RVUE 0.4–1.05 e^+e^- ¹ Update of AKHMETSHIN 00C.² Recalculated by us from the cross section in the peak.³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.⁴ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.NODE=M001G2
NODE=M001G2NODE=M001G;LINKAGE=PT
NODE=M001G;LINKAGE=LP
NODE=M001G;LINKAGE=VH

NODE=M001G2;LINKAGE=BE

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_2/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
6.02±0.20 OUR FIT Error includes scale factor of 1.9.				
6.45±0.17 OUR AVERAGE				
6.47±0.14±0.39	18680	AKHMETSHIN	05	CMD2 0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
6.50±0.11±0.20	36500	¹ ACHASOV	03	SND 0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
6.34±0.21±0.21	10625	² DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

6.80±0.13 ³BENAYOUN 10 RVUE 0.4–1.05 e^+e^- ¹ Using $\sigma_\phi \rightarrow \pi^0\gamma$ from ACHASOV 00 and $m_\omega = 782.57$ MeV in the model with the energy-independent phase of ρ - ω interference equal to $(-10.2 \pm 7.0)^\circ$.² Recalculated by us from the cross section in the peak.³ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.NODE=M001G4
NODE=M001G4

NODE=M001G;LINKAGE=SH

NODE=M001G4;LINKAGE=LP
NODE=M001G4;LINKAGE=BE

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_3/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.225±0.058±0.041	800k	1 ACHASOV	06 SND	$e^+e^- \rightarrow \pi^+\pi^-$
1.166±0.036		2 BENAYOUN	13 RVUE	0.4–1.05 e^+e^-
1.05 ±0.08		3 DAVIER	13 RVUE	$e^+e^- \rightarrow \pi^+\pi^-(\gamma)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Supersedes ACHASOV 05A.

² A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data. Supersedes BENAYOUN 10.

³ From $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ data of LEES 12G.

NODE=M001G5
NODE=M001G5

NODE=M001G5;LINKAGE=AC
NODE=M001G5;LINKAGE=B

NODE=M001G5;LINKAGE=A

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
3.32±0.28 OUR FIT	Error includes scale factor of 1.1.			
3.18±0.28 OUR AVERAGE				
3.10±0.31±0.11	33k	1 ACHASOV	07B SND	0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
3.17 ^{+1.85} _{-1.31} ±0.21	17.4k	2 AKHMETSHIN	05 CMD2	0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
3.41±0.52±0.21	23k	3,4 AKHMETSHIN	01B CMD2	$e^+e^- \rightarrow \eta\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.50±0.10 ⁵ BENAYOUN 10 RVUE 0.4–1.05 e^+e^-

¹ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.

² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770), \omega(782), \phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

⁵ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M001G3
NODE=M001G3

NODE=M001G3;LINKAGE=AH

NODE=M001G;LINKAGE=AH
NODE=M001G;LINKAGE=AK
NODE=M001G;LINKAGE=BQ

NODE=M001G3;LINKAGE=BE

 $\omega(782)$ BRANCHING RATIOS $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_1/Γ

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.9024±0.0019		1 AMBROSINO	08G KLOE	1.0–1.03 $e^+e^- \rightarrow \pi^+\pi^-\pi^0, 2\pi^0\gamma$
0.8965±0.0016±0.0048	1.2M	2,3 ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.880 ±0.020 ±0.032	11200	3,4 AKHMETSHIN	00C CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.8942±0.0062		3 DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.

² Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

³ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$.

⁴ Using $\Gamma(e^+e^-) = 0.60 \pm 0.02$ keV.

NODE=M001220

NODE=M001R21

NODE=M001R21

NODE=M001R21

NODE=M001R21;LINKAGE=AM

NODE=M001R;LINKAGE=VF

NODE=M001R;LINKAGE=ZL

NODE=M001R;LINKAGE=KH

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.09±0.14		1 AMBROSINO	08G KLOE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0, 2\pi^0\gamma$
9.06±0.20±0.57	18680	2,3 AKHMETSHIN	05 CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
9.34±0.15±0.31	36500	3 ACHASOV	03 SND	0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
8.65±0.16±0.42	1.2M	4,5 ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.39±0.24	9975	6 BENAYOUN	96 RVUE	$e^+e^- \rightarrow \pi^0\gamma$
8.88±0.62	10625	3 DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.

² Using $B(\omega \rightarrow e^+e^-) = (7.14 \pm 0.13) \times 10^{-5}$.

³ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

⁴ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

⁵ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$.

⁶ Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

NODE=M001R28
NODE=M001R28

NODE=M001R28;LINKAGE=AM

NODE=M001R;LINKAGE=AH

NODE=M001R;LINKAGE=VL

NODE=M001R28;LINKAGE=VF

NODE=M001R28;LINKAGE=ZL

NODE=M001R28;LINKAGE=A1

$\Gamma(\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_2/Γ_1

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
9.28±0.31 OUR FIT	Error includes scale factor of 2.3.		
9.05±0.27 OUR AVERAGE	Error includes scale factor of 1.8.		
8.97±0.16	AMBROSINO	08G KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
9.94±0.36±0.38	¹ AULCHENKO	00A SND	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
8.4 ±1.3	KEYNE	76 CNTR	$\pi^-p \rightarrow \omega n$
10.9 ±2.5	BENAKSAS	72C OSPK	$e^+e^- \rightarrow \pi^0\gamma$
8.1 ±2.0	BALDIN	71 HLBC	$2.9\pi^+p$
13 ±4	JACQUET	69B HLBC	$2.05\pi^+p \rightarrow \pi^+p\omega$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9.7 ±0.2 ±0.5	^{2,3} ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.9 ±0.7	² DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\gamma$

NODE=M001R3
 NODE=M001R3

¹ From $\sigma_0^{\omega\pi^0} \rightarrow \pi^0\pi^0\gamma(m_\phi)/\sigma_0^{\omega\pi^0} \rightarrow \pi^+\pi^-\pi^0\pi^0(m_\phi)$ with a phase-space correction factor of 1/1.023.

² Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

³ Using ACHASOV 03. Based on 1.2M events.

NODE=M001R3;LINKAGE=AL

NODE=M001R3;LINKAGE=VL

NODE=M001R3;LINKAGE=VW

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_3/Γ

See also $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.53^{+0.11}_{-0.13} OUR FIT	Error includes scale factor of 1.2.			
1.49±0.13 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
1.46±0.12±0.02	900k	¹ AKHMETSHIN	07	$e^+e^- \rightarrow \pi^+\pi^-$
1.30±0.24±0.05	11.2k	² AKHMETSHIN	04 CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
2.38 ^{+1.77} _{-0.90} ±0.18	5.4k	³ ACHASOV	02E SND	$1.1-1.38 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
2.3 ±0.5		BARKOV	85 OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
1.6 ^{+0.9} _{-0.7}		QUENZER	78 DM1	$e^+e^- \rightarrow \pi^+\pi^-$
3.6 ±1.9		BENAKSAS	72 OSPK	$e^+e^- \rightarrow \pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.75±0.11	4.5M	⁴ ACHASOV	05A SND	$e^+e^- \rightarrow \pi^+\pi^-$
2.01±0.29		⁵ BENAYOUN	03 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1.9 ±0.3		⁶ GARDNER	99 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
2.3 ±0.4		⁷ BENAYOUN	98 RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$
1.0 ±0.11		⁸ WICKLUND	78 ASPK	$3,4,6\pi^\pm N$
1.22±0.30		ALVENSLEB...	71C CNTR	Photoproduction
1.3 ^{+1.2} _{-0.9}		MOFFEIT	71 HBC	$2.8,4.7\gamma p$
0.80 ^{+0.28} _{-0.20}		⁹ BIGGS	70B CNTR	$4.2\gamma C \rightarrow \pi^+\pi^-C$

NODE=M001R15

NODE=M001R15
 NODE=M001R15

¹ A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

² Update of AKHMETSHIN 02.

³ From the $m_{\pi^+\pi^-}$ spectrum taking into account the interference of the $\rho\pi$ and $\omega\pi$ amplitudes.

⁴ Using $\Gamma(\omega \rightarrow e^+e^-)$ from the 2004 Edition of this Review (PDG 04).

⁵ Using the data of AKHMETSHIN 02 in the hidden local symmetry model.

⁶ Using the data of BARKOV 85.

⁷ Using the data of BARKOV 85 in the hidden local symmetry model.

⁸ From a model-dependent analysis assuming complete coherence.

⁹ Re-evaluated under $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ by BEHREND 71 using more accurate $\omega \rightarrow \rho$ photoproduction cross-section ratio.

NODE=M001R15;LINKAGE=AK

NODE=M001R15;LINKAGE=PT

NODE=M001R;LINKAGE=VE

NODE=M001R;LINKAGE=SN


NODE=M001R;LINKAGE=BY

NODE=M001R15;LINKAGE=H4

NODE=M001R15;LINKAGE=Q

NODE=M001R15;LINKAGE=F

NODE=M001R15;LINKAGE=B



$$\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$$

See also $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0172±0.0014 OUR FIT			Error includes scale factor of 1.2.
0.026 ±0.005 OUR AVERAGE			

0.021 $^{+0.028}_{-0.009}$	1,2	RATCLIFF 72	ASPK	$15 \pi^- p \rightarrow n2\pi$
0.028 ±0.006	1	BEHREND 71	ASPK	Photoproduction
0.022 $^{+0.009}_{-0.01}$	3	ROOS 70	RVUE	

¹ The fitted width of these data is 160 MeV in agreement with present average, thus the ω contribution is overestimated. Assuming ρ width 145 MeV.

² Significant interference effect observed. NB of $\omega \rightarrow 3\pi$ comes from an extrapolation.

³ ROOS 70 combines ABRAMOVICH 70 and BIZZARRI 70.

$$\Gamma(\pi^+\pi^-)/\Gamma(\pi^0\gamma)$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.20±0.04	1.98M	1 ALOISIO 03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ Using the data of ALOISIO 02D.

$$\Gamma(\text{neutrals})/\Gamma_{\text{total}}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.091±0.006 OUR FIT				
0.081±0.011 OUR AVERAGE				

0.075±0.025		BIZZARRI 71	HBC	0.0 $p\bar{p}$
0.079±0.019		DEINET 69B	OSPK	1.5 $\pi^- p$
0.084±0.015		BOLLINI 68C	CNTR	2.1 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.073±0.018	42	BASILE 72B	CNTR	1.67 $\pi^- p$

$$\Gamma(\text{neutrals})/\Gamma(\pi^+\pi^-\pi^0)$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.102±0.008 OUR FIT				
0.103 $^{+0.011}_{-0.010}$ OUR AVERAGE				

0.15 ±0.04	46	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
0.10 ±0.03	19	BARASH 67B	HBC	0.0 $\bar{p}p$	
0.134±0.026	850	DIGIUGNO 66B	CNTR	1.4 $\pi^- p$	
0.097±0.016	348	FLATTE 66	HBC	1.4 - 1.7 $K^- p \rightarrow \Lambda MM$	
0.06 $^{+0.05}_{-0.02}$		JAMES 66	HBC	2.1 $\pi^+ p$	
0.08 ±0.03	35	KRAEMER 64	DBC	1.2 $\pi^+ d$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.11 ±0.02	20	BUSCHBECK 63	HBC	1.5 $K^- p$	

 Γ_3/Γ_1

NODE=M001R2
NODE=M001R2
NODE=M001R2

 Γ_3/Γ_2

NODE=M001R33
NODE=M001R33

 $(\Gamma_2+\Gamma_4)/\Gamma$

NODE=M001R14
NODE=M001R14

 $(\Gamma_2+\Gamma_4)/\Gamma_1$

NODE=M001R1
NODE=M001R1

NODE=M001R;LINKAGE=KL

NODE=M001R2;LINKAGE=A

NODE=M001R2;LINKAGE=S
NODE=M001R2;LINKAGE=R

$\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$ $\Gamma_2/(\Gamma_2+\Gamma_4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M001R18
 NODE=M001R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.78 ± 0.07		¹ DAKIN	72	OSPK	$1.4 \pi^- p \rightarrow nMM$
>0.81	90	DEINET	69B	OSPK	

¹ Error statistical only. Authors obtain good fit also assuming $\pi^0\gamma$ as the only neutral decay.

NODE=M001R18;LINKAGE=D

 $\Gamma(\text{neutrals})/\Gamma(\text{charged particles})$ $(\Gamma_2+\Gamma_4)/(\Gamma_1+\Gamma_3)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M001R9
 NODE=M001R9

0.100 ± 0.008 OUR FIT**0.124 ± 0.021**

FELDMAN	67C	OSPK	$1.2 \pi^- p$
---------	-----	------	---------------

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M001R19
 NODE=M001R19

4.6 ± 0.4 OUR FIT Error includes scale factor of 1.1.**6.3 ± 1.3 OUR AVERAGE** Error includes scale factor of 1.2.

6.6 ± 1.7		¹ ABELE	97E	CBAR	$0.0 \bar{p} p \rightarrow 5\gamma$
8.3 ± 2.1		ALDE	93	GAM2	$38\pi^- p \rightarrow \omega n$
$3.0 \begin{smallmatrix} +2.5 \\ -1.8 \end{smallmatrix}$		² ANDREWS	77	CNTR	$6.7-10 \gamma Cu$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.3 \pm 0.5 \pm 0.1$	33k	³ ACHASOV	07B	SND	$0.6-1.38 e^+ e^- \rightarrow \eta\gamma$
$4.44 \begin{smallmatrix} +2.59 \\ -1.83 \end{smallmatrix} \pm 0.28$	17.4k	^{4,5} AKHMETSHIN	05	CMD2	$0.60-1.38 e^+ e^- \rightarrow \eta\gamma$
$5.10 \pm 0.72 \pm 0.34$	23k	⁶ AKHMETSHIN	01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
$0.7 \text{ to } 5.5$		⁷ CASE	00	CBAR	$0.0 p\bar{p} \rightarrow \eta\eta\gamma$
$6.56 \begin{smallmatrix} +2.41 \\ -2.55 \end{smallmatrix}$	3525	^{2,8} BENAYOUN	96	RVUE	$e^+ e^- \rightarrow \eta\gamma$
7.3 ± 2.9		^{2,4} DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$

¹ No flat $\eta\eta\gamma$ background assumed.

² Solution corresponding to constructive ω - ρ interference.

³ ACHASOV 07B reports $[\Gamma(\omega(782) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow e^+ e^-)] = (3.10 \pm 0.31 \pm 0.11) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow e^+ e^-) = (7.28 \pm 0.14) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

⁴ Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁵ Using $B(\omega \rightarrow e^+ e^-) = (7.14 \pm 0.13) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

⁶ Using $B(\omega \rightarrow e^+ e^-) = (7.07 \pm 0.19) \times 10^{-5}$ and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$. Solution corresponding to constructive ω - ρ interference. The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively). Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁷ Depending on the degree of coherence with the flat $\eta\eta\gamma$ background and using $B(\omega \rightarrow \pi^0\gamma) = (8.5 \pm 0.5) \times 10^{-2}$.

⁸ Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

NODE=M001R;LINKAGE=EA
 NODE=M001R19;LINKAGE=A
 NODE=M001R19;LINKAGE=AO

NODE=M001R13;LINKAGE=WL
 NODE=M001R19;LINKAGE=AK
 NODE=M001R19;LINKAGE=TS

NODE=M001R;LINKAGE=CS

NODE=M001R19;LINKAGE=A1

 $\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$ Γ_5/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M001R11
 NODE=M001R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0098 ± 0.0024		¹ ALDE	93	GAM2	$38\pi^- p \rightarrow \omega n$
0.0082 ± 0.0033		² DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$
0.010 ± 0.045		APEL	72B	OSPK	$4-8 \pi^- p \rightarrow n3\gamma$

¹ Model independent determination.

² Solution corresponding to constructive ω - ρ interference.

NODE=M001R11;LINKAGE=A
 NODE=M001R11;LINKAGE=K

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M001R23
 NODE=M001R23

7.7 ± 0.6 OUR FIT**7.7 ± 0.6 OUR AVERAGE**

$7.61 \pm 0.53 \pm 0.64$		ACHASOV	08	SND	$0.36-0.97 e^+ e^- \rightarrow \pi^0 e^+ e^-$
$8.19 \pm 0.71 \pm 0.62$		AKHMETSHIN	05A	CMD2	$0.72-0.84 e^+ e^-$
5.9 ± 1.9	43	DOLINSKY	88	ND	$e^+ e^- \rightarrow \pi^0 e^+ e^-$

$\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.3 ± 0.4 OUR FIT				Error includes scale factor of 2.1.
1.3 ± 0.4 OUR AVERAGE				Error includes scale factor of 2.1.
1.72 ± 0.25 ± 0.14	3k	ARNALDI	09 NA60	158A In-In collisions
0.96 ± 0.23		DZHELADIN	81B CNTR	25-33 $\pi^- p \rightarrow \omega n$

NODE=M001R12
 NODE=M001R12

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<1.1	AKHMETSHIN 05A	CMD2	0.72-0.84 $e^+ e^-$

NODE=M001R34
 NODE=M001R34

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.728 ± 0.014 OUR FIT				Error includes scale factor of 1.3.
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.700 ± 0.016	11200	^{1,2} AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.752 ± 0.004 ± 0.024	1.2M	^{2,3} ACHASOV 03D	RVUE	0.44-2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.714 ± 0.036		² DOLINSKY 89	ND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.72 ± 0.03		² BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.64 ± 0.04	1488	² KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.675 ± 0.069	433	² CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.83 ± 0.10	451	² BENAJSAS 72B	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.77 ± 0.06		⁴ AUGUSTIN 69D	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.65 ± 0.13	33	⁵ ASTVACAT... 68	OSPK	Assume SU(3)+mixing

NODE=M001R13
 NODE=M001R13

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = 0.891 \pm 0.007$. Update of AKHMETSHIN 00C.

² Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}^2$.

³ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+ \pi^-) = (1.70 \pm 0.28)\%$.

⁴ Rescaled by us to correspond to ω width 8.4 MeV. Systematic errors underestimated.

⁵ Not resolved from ρ decay. Error statistical only.

NODE=M001R13;LINKAGE=4P
 NODE=M001R13;LINKAGE=ZL
 NODE=M001R13;LINKAGE=VF
 NODE=M001R13;LINKAGE=E
 NODE=M001R13;LINKAGE=A

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2	90	ACHASOV 09A	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<200	90	KURDADZE 86	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

NODE=M001R5
 NODE=M001R5

 $\Gamma(\pi^+ \pi^- \gamma)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0036	95	WEIDENAUER 90	ASTE	$p\bar{p} \rightarrow \pi^+ \pi^- \pi^+ \pi^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.004	95	BITYUKOV 88B	SPEC	$32 \pi^- p \rightarrow \pi^+ \pi^- \gamma X$

NODE=M001R22
 NODE=M001R22

 $\Gamma(\pi^+ \pi^- \gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_{11}/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.066	90	KALBFLEISCH 75	HBC	2.18 $K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$
<0.05	90	FLATTE 66	HBC	1.2 - 1.7 $K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$

NODE=M001R4
 NODE=M001R4

 $\Gamma(\pi^+ \pi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1 × 10 ⁻³	90	KURDADZE 88	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$

NODE=M001R24
 NODE=M001R24

 $\Gamma(\pi^0 \pi^0 \gamma)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.6 ± 1.1 OUR FIT				
6.5 ± 1.2 OUR AVERAGE				
6.4 ^{+2.4} _{-2.0} ± 0.8	190	¹ AKHMETSHIN 04B	CMD2	0.6-0.97 $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
6.6 ^{+1.4} _{-1.3} ± 0.6	295	ACHASOV 02F	SND	0.36-0.97 $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

NODE=M001R29
 NODE=M001R29

• • • We do not use the following data for averages, fits, limits, etc. • • •

$11.8^{+2.1}_{-1.9} \pm 1.4$	190	² AKHMETSHIN 04B	CMD2	0.6–0.97	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
$7.8 \pm 2.7 \pm 2.0$	63	^{1,3} ACHASOV	00G SND		$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
$12.7 \pm 2.3 \pm 2.5$	63	^{2,3} ACHASOV	00G SND		$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2

¹ In the model assuming the $\rho \rightarrow \pi^0 \pi^0 \gamma$ decay via the $\omega \pi$ and $f_0(500) \gamma$ mechanisms.

² In the model assuming the $\rho \rightarrow \pi^0 \pi^0 \gamma$ decay via the $\omega \pi$ mechanism only.

³ Superseded by ACHASOV 02F.

NODE=M001R29;LINKAGE=A
NODE=M001R29;LINKAGE=B
NODE=M001R;LINKAGE=GF

$\Gamma(\pi^0 \pi^0 \gamma) / \Gamma(\pi^+ \pi^- \pi^0)$

Γ_{13} / Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00045	90	DOLINSKY	89 ND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

NODE=M001R10
NODE=M001R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.08	95	JACQUET	69B HLBC	2.05	$\pi^+ p \rightarrow \pi^+ p \omega$
-------	----	---------	----------	------	--------------------------------------

$\Gamma(\pi^0 \pi^0 \gamma) / \Gamma(\pi^0 \gamma)$

Γ_{13} / Γ_2

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
8.0 ± 1.3 OUR FIT					
8.5 ± 2.9	40 ± 14		ALDE	94B GAM2	$38 \pi^- p \rightarrow \pi^0 \pi^0 \gamma n$

NODE=M001R7
NODE=M001R7

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 50	90	DOLINSKY	89 ND		$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
<1800	95	KEYNE	76 CNTR		$\pi^- p \rightarrow \omega n$
<1500	90	BENAKSAS	72C OSPK		$e^+ e^-$
<1400		BALDIN	71 HLBC	2.9	$\pi^+ p$
<1000	90	BARMIN	64 HLBC	1.3–2.8	$\pi^- p$

$\Gamma(\pi^0 \pi^0 \gamma) / \Gamma(\text{neutrals})$

$\Gamma_{13} / (\Gamma_2 + \Gamma_4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M001R17
NODE=M001R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22 ± 0.07		¹ DAKIN	72 OSPK	1.4	$\pi^- p \rightarrow n \text{MM}$
<0.19	90	DEINET	69B OSPK		

¹ See $\Gamma(\pi^0 \gamma) / \Gamma(\text{neutrals})$.

NODE=M001R17;LINKAGE=D

$\Gamma(\eta \pi^0 \gamma) / \Gamma_{\text{total}}$

Γ_{14} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3.3	90	AKHMETSHIN 04B	CMD2	0.6–0.97 $e^+ e^- \rightarrow \eta \pi^0 \gamma$

NODE=M001R32
NODE=M001R32

$\Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}}$

Γ_{15} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
9.0 ± 3.1 OUR FIT				
9.0 ± 2.9 ± 1.1	18	HEISTER	02C ALEP	$Z \rightarrow \mu^+ \mu^- + X$

NODE=M001R30
NODE=M001R30

$\Gamma(\mu^+ \mu^-) / \Gamma(\pi^+ \pi^- \pi^0)$

Γ_{15} / Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.2	90	WILSON	69 OSPK	12 $\pi^- C \rightarrow \text{Fe}$

NODE=M001R6
NODE=M001R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7	74	FLATTE	66 HBC	1.2 – 1.7	$K^- p \rightarrow \Lambda \mu^+ \mu^-$
<1.2		BARBARO-...	65 HBC	2.7	$K^- p$

$\Gamma(\pi^0 \mu^+ \mu^-) / \Gamma(\mu^+ \mu^-)$

Γ_7 / Γ_{15}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M001R20
NODE=M001R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 ± 0.6	30	¹ DZHELYADIN	79 CNTR	25–33	$\pi^- p$
-----------	----	-------------------------	---------	-------	-----------

¹ Superseded by DZHELYADIN 81B result above.

NODE=M001R20;LINKAGE=S

$\Gamma(3\gamma) / \Gamma_{\text{total}}$

Γ_{16} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	95	¹ ABELE	97E CBAR	0.0 $\bar{p} p \rightarrow 5\gamma$

NODE=M001R27
NODE=M001R27

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2	90	¹ PROKOSHKIN	95 GAM2	38	$\pi^- p \rightarrow 3\gamma n$
----	----	-------------------------	---------	----	---------------------------------

¹ From direct 3γ decay search.

NODE=M001R27;LINKAGE=A

$\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$

Violates C conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.001	90	ALDE	94B	GAM2 $38\pi^- p \rightarrow \eta\pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Γ_{17}/Γ

NODE=M001R25
 NODE=M001R25
 NODE=M001R25

 $[\Gamma(\eta\gamma) + \Gamma(\eta\pi^0)]/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.016	90	¹ FLATTE	66	HBC $1.2 - 1.7 K^- p \rightarrow \Lambda\pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.045	95	JACQUET	69B	HLBC $2.05 \pi^+ p \rightarrow \pi^+ p\omega$
--------	----	---------	-----	---

 $(\Gamma_5 + \Gamma_{17})/\Gamma_1$

NODE=M001R8
 NODE=M001R8

¹ Restated by us using $B(\eta \rightarrow \text{charged modes}) = 29.2\%$.

NODE=M001R8;LINKAGE=A

 $\Gamma(\eta\pi^0)/\Gamma(\pi^0\gamma)$

Violates C conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	¹ STAROSTIN	09	CRYM $\gamma p \rightarrow \eta\pi^0 p$

¹ STAROSTIN 09 reports $[\Gamma(\omega(782) \rightarrow \eta\pi^0)/\Gamma(\omega(782) \rightarrow \pi^0\gamma)] \times [B(\eta \rightarrow 2\gamma)] < 1.01 \times 10^{-3}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

 Γ_{17}/Γ_2

NODE=M001R35
 NODE=M001R35
 NODE=M001R35

NODE=M001R35;LINKAGE=ST

 $\Gamma(2\pi^0)/\Gamma(\pi^0\gamma)$

Violates C conservation and Bose-Einstein statistics.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.59	90	STAROSTIN	09	CRYM $\gamma p \rightarrow 2\pi^0 p$

 Γ_{18}/Γ_2

NODE=M001R36
 NODE=M001R36
 NODE=M001R36

 $\Gamma(3\pi^0)/\Gamma_{\text{total}}$

Violates C conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3×10^{-4}	90	PROKOSHKIN	95	GAM2 $38 \pi^- p \rightarrow 3\pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Γ_{19}/Γ

NODE=M001R26
 NODE=M001R26
 NODE=M001R26

 $\Gamma(3\pi^0)/\Gamma(\pi^0\gamma)$

Violates C conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.72	90	STAROSTIN	09	CRYM $\gamma p \rightarrow 3\pi^0 p$

 Γ_{19}/Γ_2

NODE=M001R37
 NODE=M001R37
 NODE=M001R37

 $\Gamma(3\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

Violates C conservation.

VALUE	CL%	DOCUMENT ID	COMMENT
<0.009	90	BARBERIS	01 450 $p p \rightarrow p_f 3\pi^0 p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Γ_{19}/Γ_1

NODE=M001R31
 NODE=M001R31
 NODE=M001R31

PARAMETER Λ IN $\omega \rightarrow \pi^0\mu^+\mu^-$ DECAY

In the pole approximation the electromagnetic transition form factor for a resonance of mass M is given by the expression:

$$|F|^2 = (1 - M^2/\Lambda^2)^{-2},$$

where for the parameter Λ vector dominance predicts $\Lambda = M_p \approx 0.770$ GeV. The ARNALDI 09 measurement is in obvious conflict with this expectation. Note that for $\eta \rightarrow \mu^+\mu^-\gamma$ decay ARNALDI 09 and DZHELYADIN 80 obtain the value of Λ consistent with vector dominance.

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.668 ± 0.009 ± 0.003	3k	ARNALDI	09	NA60 158A In-In collisions
0.65 ± 0.03		DZHELYADIN	81B	CNTR 25-33 $\pi^- p \rightarrow \omega n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M001LAM
 NODE=M001LAM

NODE=M001LAM

ω (782) REFERENCES

NODE=M001

ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono (PARIN, BERLIN+)		REFID=55357
DAVIER	13	EPJ C73 2597	M. Davier <i>et al.</i>		REFID=55499
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)	REFID=54305
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53101
ARNALDI	09	PL B677 260	R. Arnaldi <i>et al.</i>	(NA60 Collab.)	REFID=52720
STAROSTIN	09	PR C79 065201	A. Starostin <i>et al.</i>	(Crystal Ball Collab. at MAMI)	REFID=53001
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=52258
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52573
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AKHMETSHIN	07	PL B648 28	R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
ACHASOV	06A	Translated from ZETF 130 437.			
AULCHENKO	06	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51513
ACHASOV	05A	Translated from ZETFP 84 491.			
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51045
AKHMETSHIN	05	Translated from ZETF 128 1201.			
AKHMETSHIN	05A	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AULCHENKO	05	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50508
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51060
AKHMETSHIN	04	Translated from ZETFP 82 841.			
AKHMETSHIN	04B	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AUBERT,B	04N	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49610
PDG	04	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ACHASOV	03	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
ACHASOV	03D	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49187
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
BENAYOUN	03	EPJ C29 397	M. Benayoun <i>et al.</i>		REFID=49477
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48816
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48565
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
HEISTER	02C	PL B528 19	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48564
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>		REFID=48324
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
ACHASOV	00G	Translated from ZETFP 72 411.			
ACHASOV	00G	JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47929
AKHMETSHIN	00C	Translated from ZETFP 71 519.			
AULCHENKO	00A	PL B476 33	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47423
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
CASE	00	Translated from ZETF 117 1067.			
ACHASOV	99E	PR D61 032002	T. Case <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47409
GARDNER	99	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
BENAYOUN	98	PR D59 076002	S. Gardner, H.B. O'Connell		REFID=46919
ABELE	97E	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)	REFID=45859
BENAYOUN	96	PL B411 361	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45755
PROKOSHKIN	95	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
WURZINGER	95	SPD 40 273	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=44616
ALDE	94B	Translated from DANS 342 610.			
AMSLER	94C	PR C51 443	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=45209
ALDE	93	PL B340 122	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=44100
ALDE	93	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
ALDE	93	PAN 56 1229	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=43603
AMSLER	93B	Translated from YAF 56 137.			
WEIDENAUER	93	ZPHY C61 35	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=43790
ANTONELLI	92	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43602
DOLINSKY	91	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
WEIDENAUER	90	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
DOLINSKY	89	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BITYUKOV	88B	WEIDENAUER 90 ZPHY C47 353	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=41368
DOLINSKY	88	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
DOLINSKY	88	SJNP 47 800	S.I. Bitjukov <i>et al.</i>	(SERP)	REFID=41021
KURDADZE	88	Translated from YAF 47 1258.			
KURDADZE	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41022
AULCHENKO	87	Translated from YAF 48 442.			
BARKOV	87	JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=41121
BARKOV	87	Translated from ZETFP 47 432.			
BARKOV	87	PL B186 432	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=40007
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
KURDADZE	86	Translated from ZETFP 46 132.			
KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=40287
BARKOV	85	Translated from ZETFP 43 497.			
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
KURDADZE	83B	JETPL 36 274	A.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20244
KURDADZE	83B	Translated from ZETFP 36 221.			

DZHELADIN	81B	PL 102B 296	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20242
CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)	REFID=20240
DZHELADIN	80	PL 94B 548	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10831
ROOS	80	LCN 27 321	M. Roos, A. Pellinen	(HELS)	REFID=20241
BENKHEIRI	79	NP B150 268	P. Benkheiri <i>et al.</i>	(EPOL, CERN, CDEF+)	REFID=20238
DZHELADIN	79	PL 84B 143	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20239
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=20235
QUENZER	78	PL 76B 512	A. Quenzer <i>et al.</i>	(LALO)	REFID=20123
VANAPEL...	78	NP B133 245	G.W. van Apeldoorn <i>et al.</i>	(ZEEM)	REFID=20234
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)	REFID=20230
KEYNE	76	PR D14 28	J. Keyne <i>et al.</i>	(LOIC, SHMP)	REFID=20226
Also		PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20216
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72B	PL 41B 234	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20206
BASILE	72B	Phil. Conf. 153	M. Basile <i>et al.</i>	(CERN)	REFID=20207
BENAKSAS	72	PL 39B 289	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20096
BENAKSAS	72B	PL 42B 507	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20209
BENAKSAS	72C	PL 42B 511	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20517
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)	REFID=20215
BROWN	72	PL 42B 117	R.M. Brown <i>et al.</i>	(ILL, ILLC)	REFID=20211
DAKIN	72	PR D6 2321	J.T. Dakin <i>et al.</i>	(PRIN)	REFID=20212
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)	REFID=20102
ALVENSLEB...	71C	PRL 27 888	H. Alvensleben <i>et al.</i>	(DESY)	REFID=20193
BALDIN	71	SJNP 13 758	A.B. Baldin <i>et al.</i>	(ITEP)	REFID=20195
		Translated from YAF 13	1318.		
BEHREND	71	PRL 27 61	H.J. Behrend <i>et al.</i>	(ROCH, CORN, FNAL)	REFID=20197
BIZZARRI	71	NP B27 140	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20198
COYNE	71	NP B32 333	D.G. Coyne <i>et al.</i>	(LRL)	REFID=20201
MOFFEIT	71	NP B29 349	K.C. Moffeit <i>et al.</i>	(LRL, UCB, SLAC+)	REFID=20204
ABRAMOVI...	70	NP B20 209	M. Abramovich <i>et al.</i>	(CERN)	REFID=20180
BIGGS	70B	PRL 24 1201	P.J. Biggs <i>et al.</i>	(DARE)	REFID=20184
BIZZARRI	70	PRL 25 1385	R. Bizzarri <i>et al.</i>	(ROMA, SYRA)	REFID=20181
ROOS	70	DNPL/R7 173	M. Roos	(CERN)	REFID=20191
		Proc. Daresbury Study Weekend No. 1.			
AUGUSTIN	69D	PL 28B 513	J.E. Augustin <i>et al.</i>	(ORSAY)	REFID=20169
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20171
DEINET	69B	PL 30B 426	W. Deinet <i>et al.</i>	(KARL, CERN)	REFID=20173
JACQUET	69B	NC 63A 743	F. Jacquet <i>et al.</i>	(EPOL, BERG)	REFID=20176
WILSON	69	Private Comm.	R. Wilson	(HARV)	REFID=20179
Also		PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)	REFID=20084
ASTVACAT...	68	PL 27B 45	R.G. Astvatsaturov <i>et al.</i>	(JINR, MOSU)	REFID=20055
BOLLINI	68C	NC 56A 531	D. Bollini <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20164
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)	REFID=20160
FELDMAN	67C	PR 159 1219	M. Feldman <i>et al.</i>	(PENN)	REFID=20161
DIGIUGNO	66B	NC 44A 1272	G. Di Giugno <i>et al.</i>	(NAPL, FRAS, TRST)	REFID=20156
FLATTE	66	PR 145 1050	S.M. Flatte <i>et al.</i>	(LRL)	REFID=20157
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=10770
BARBARO...	65	PRL 14 279	A. Barbaro-Galtieri, R.D. Tripp	(LRL)	REFID=20152
BARMIN	64	JETP 18 1289	V.V. Barmin <i>et al.</i>	(ITEP)	REFID=20149
		Translated from ZETF 45	1879.		
KRAEMER	64	PR 136 B496	R.W. Kraemer <i>et al.</i>	(JHU, NWES, WOOD)	REFID=10755
BUSCHBECK	63	Siena Conf. 1 166	B. Buschbeck <i>et al.</i>	(VIEN, CERN, ANIK)	REFID=20146

$\eta'(958)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

NODE=M002

 $\eta'(958)$ MASS

NODE=M002M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
957.78 ±0.06 OUR AVERAGE				
957.793 ±0.054 ±0.036	3.9k	LIBBY	08	CLEO $J/\psi \rightarrow \gamma\eta'$
957.9 ±0.2 ±0.6	4800	WURZINGER	96	SPEC 1.68 $p d \rightarrow {}^3\text{He}\eta'$
957.46 ±0.33		DUANE	74	MMS $\pi^- p \rightarrow n\text{MM}$
958.2 ±0.5	1414	DANBURG	73	HBC 2.2 $K^- p \rightarrow \Lambda\eta'$
958 ±1	400	JACOBS	73	HBC 2.9 $K^- p \rightarrow \Lambda\eta'$
956.1 ±1.1	3415	¹ BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
957.5 ±0.2		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
959 ±1	630	² BELADIDZE	92C	VES 36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
958 ±1	340	² ARMSTRONG	91B	OMEG 300 $pp \rightarrow pp\eta\pi^+\pi^-$
958.2 ±0.4	622	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
957.8 ±0.2	2420	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
956.3 ±1.0	143	² GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
957.4 ±1.4	535	³ BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
957 ±1		RITTENBERG	69	HBC 1.7-2.7 $K^- p$

NODE=M002M

OCCUR=2

OCCUR=2

¹ Using all η' decays.² Systematic uncertainty not estimated.³ Using η' decays into neutrals. Not independent of the other listed BASILE 71 η' mass measurement.

NODE=M002M;LINKAGE=BS

NODE=M002M;LINKAGE=NS

NODE=M002M;LINKAGE=BA

 $\eta'(958)$ WIDTH

NODE=M002W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.197 ±0.009 OUR FIT					
[0.198 ± 0.009 MeV OUR 2015 FIT]					
0.230 ±0.021 OUR AVERAGE					
0.226 ±0.017 ±0.014	2300	CZERWINSKI	10	MMS	$pp \rightarrow pp\eta'$
0.40 ±0.22	4800	WURZINGER	96	SPEC	1.68 $p d \rightarrow {}^3\text{He}\eta'$
0.28 ±0.10	1000	BINNIE	79	MMS	0 $\pi^- p \rightarrow n\text{MM}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.20 ±0.04		BAI	04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

NODE=M002W

NEW

 $\eta'(958)$ DECAY MODES

NODE=M002215;NODE=M002

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\pi^+\pi^-\eta$	(42.9 ±0.7) %	DESIG=1
Γ_2 $\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$)	(29.1 ±0.5) %	DESIG=9
Γ_3 $\pi^0\pi^0\eta$	(22.3 ±0.8) %	DESIG=2
Γ_4 $\omega\gamma$	(2.62±0.13) %	DESIG=7
Γ_5 ωe^+e^-	(2.0 ±0.4) × 10 ⁻⁴	DESIG=205
Γ_6 $\gamma\gamma$	(2.21±0.08) %	DESIG=6
Γ_7 $3\pi^0$	(2.20±0.20) × 10 ⁻³	DESIG=8
Γ_8 $\mu^+\mu^-\gamma$	(1.08±0.27) × 10 ⁻⁴	DESIG=20
Γ_9 $\pi^+\pi^-\mu^+\mu^-$	< 2.9 × 10 ⁻⁵	90% DESIG=201
Γ_{10} $\pi^+\pi^-\pi^0$	(3.82±0.35) × 10 ⁻³	DESIG=121
Γ_{11} $\pi^0\rho^0$	< 4 %	90% DESIG=18
Γ_{12} $2(\pi^+\pi^-)$	(8.5 ±0.9) × 10 ⁻⁵	DESIG=131
Γ_{13} $\pi^+\pi^-2\pi^0$	(1.8 ±0.4) × 10 ⁻⁴	DESIG=202
Γ_{14} $2(\pi^+\pi^-)$ neutrals	< 1 %	95% DESIG=132
Γ_{15} $2(\pi^+\pi^-)\pi^0$	< 1.9 × 10 ⁻³	90% DESIG=141

Γ_{16}	$2(\pi^+\pi^-)2\pi^0$	< 1	%	95%	DESIG=15
Γ_{17}	$3(\pi^+\pi^-)$	< 3.1	$\times 10^{-5}$	90%	DESIG=203
Γ_{18}	$\pi^+\pi^-e^+e^-$	$(2.4^{+1.3}_{-1.0})$	$\times 10^{-3}$		DESIG=10
Γ_{19}	$\pi^+e^-\nu_e + \text{c.c.}$	< 2.1	$\times 10^{-4}$	90%	DESIG=204
Γ_{20}	γe^+e^-	(4.70 ± 0.30)	$\times 10^{-4}$		DESIG=28
Γ_{21}	$\pi^0\gamma\gamma$	< 8	$\times 10^{-4}$	90%	DESIG=24
Γ_{22}	$4\pi^0$	< 3.2	$\times 10^{-4}$	90%	DESIG=26
Γ_{23}	e^+e^-	< 5.6	$\times 10^{-9}$	90%	DESIG=150
Γ_{24}	invisible	< 5	$\times 10^{-4}$	90%	DESIG=200

**Charge conjugation (C), Parity (P),
Lepton family number (LF) violating modes**

NODE=M002;CLUMP=B

Γ_{25}	$\pi^+\pi^-$	P,CP	< 6	$\times 10^{-5}$	90%	DESIG=111
Γ_{26}	$\pi^0\pi^0$	P,CP	< 4	$\times 10^{-4}$	90%	DESIG=25
Γ_{27}	$\pi^0e^+e^-$	C	[a] < 1.4	$\times 10^{-3}$	90%	DESIG=16
Γ_{28}	ηe^+e^-	C	[a] < 2.4	$\times 10^{-3}$	90%	DESIG=17
Γ_{29}	3γ	C	< 1.0	$\times 10^{-4}$	90%	DESIG=23
Γ_{30}	$\mu^+\mu^-\pi^0$	C	[a] < 6.0	$\times 10^{-5}$	90%	DESIG=22
Γ_{31}	$\mu^+\mu^-\eta$	C	[a] < 1.5	$\times 10^{-5}$	90%	DESIG=21
Γ_{32}	$e\mu$	LF	< 4.7	$\times 10^{-4}$	90%	DESIG=27

[a] C parity forbids this to occur as a single-photon process.

LINKAGE=CS

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 16 branching ratios uses 46 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 52.8$ for 38 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-2							
x_3	-77	-58						
x_4	-11	-13	2					
x_6	-29	-25	32	-1				
x_7	-24	-19	30	0	9			
x_{10}	0	-2	-2	0	-1	-1		
x_{18}	-4	-6	-5	-1	-3	-2	0	
Γ	25	5	-19	3	-71	-5	1	3
	x_1	x_2	x_3	x_4	x_6	x_7	x_{10}	x_{18}

Mode	Rate (MeV)		
Γ_1	$\pi^+\pi^-\eta$	0.085 \pm 0.004	DESIG=1
Γ_2	$\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$)	0.0574 \pm 0.0028	DESIG=9
Γ_3	$\pi^0\pi^0\eta$	0.0440 \pm 0.0023	DESIG=2
Γ_4	$\omega\gamma$	0.00517 \pm 0.00035	DESIG=7
Γ_6	$\gamma\gamma$	0.00435 \pm 0.00013	DESIG=6
Γ_7	$3\pi^0$	(4.3 \pm 0.4) $\times 10^{-4}$	DESIG=8
Γ_{10}	$\pi^+\pi^-\pi^0$	(7.5 \pm 0.8) $\times 10^{-4}$	DESIG=121
Γ_{18}	$\pi^+\pi^-e^+e^-$	(4.7 $^{+2.6}_{-1.9}$) $\times 10^{-4}$	DESIG=10

$\eta'(958)$ PARTIAL WIDTHS

NODE=M002220

 $\Gamma(\gamma\gamma)$ Γ_6

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.35±0.14 OUR FIT				
4.28±0.19 OUR AVERAGE				
4.17±0.10±0.27	2000	1 ACCIARRI	98Q L3	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
4.53±0.29±0.51	266	KARCH	92 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
3.61±0.13±0.48		2 BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\eta'(958)$
4.6 ±1.1 ±0.6	23	BARU	90 MD1	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
4.57±0.25±0.44		BUTLER	90 MRK2	$e^+e^- \rightarrow e^+e^-\eta'(958)$
5.08±0.24±0.71	547	3 ROE	90 ASP	$e^+e^- \rightarrow e^+e^-2\gamma$
3.8 ±0.7 ±0.6	34	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.9 ±0.5 ±0.5	136	4 WILLIAMS	88 CBAL	$e^+e^- \rightarrow e^+e^-2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.7 ±0.6 ±0.9	143	5 GIDAL	87 MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.0 ±0.9		6 BARTEL	85E JADE	$e^+e^- \rightarrow e^+e^-2\gamma$

NODE=M002W4
NODE=M002W4¹No non-resonant $\pi^+\pi^-$ contribution found.²Reevaluated by us using $B(\eta' \rightarrow \rho(770)\gamma) = (30.2 \pm 1.3)\%$.³Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.⁴Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.⁵Superseded by BUTLER 90.⁶Systematic error not evaluated.NODE=M002W4;LINKAGE=AC
NODE=M002W4;LINKAGE=K1
NODE=M002W4;LINKAGE=K2
NODE=M002W4;LINKAGE=K3
NODE=M002W4;LINKAGE=C
NODE=M002W4;LINKAGE=A $\Gamma(e^+e^-)$ Γ_{23}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 × 10⁻³	90	1,2 ACHASOV	15 SND	0.958 $e^+e^- \rightarrow \pi\pi\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<2.0 × 10 ⁻³	90	2 ACHASOV	15 SND	0.958 $e^+e^- \rightarrow \pi\pi\eta$
<2.4 × 10 ⁻³	90	2 AKHMETSHIN	15 CMD3	0.958 $e^+e^- \rightarrow \pi^+\pi^-\eta$
¹ Combining data of ACHASOV 15 and AKHMETSHIN 15.				
² Using η and η' branching fractions from PDG 14.				

NODE=M002W1
NODE=M002W1

OCCUR=2

NODE=M002W1;LINKAGE=A
NODE=M002W1;LINKAGE=B $\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M002223

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

NODE=M002223

 $\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}}$ $\Gamma_6\Gamma_2/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.27±0.04 OUR FIT				
1.26±0.07 OUR AVERAGE Error includes scale factor of 1.2.				
1.09±0.04±0.13		BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\rho(770)^0\gamma$
1.35±0.09±0.21		AIHARA	87 TPC	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.13±0.04±0.13	867	ALBRECHT	87B ARG	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.53±0.09±0.21		ALTHOFF	84E TASS	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.14±0.08±0.11	243	BERGER	84B PLUT	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.73±0.34±0.35	95	JENNI	83 MRK2	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.49±0.13±0.027	213	BARTEL	82B JADE	$e^+e^- \rightarrow e^+e^-\rho\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.85±0.31±0.24	43	BEHREND	83B CELL	$e^+e^- \rightarrow e^+e^-\rho\gamma$

NODE=M002G1
NODE=M002G1 $\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$ $\Gamma_6\Gamma_3/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.97±0.05 OUR FIT			
0.92±0.06±0.11			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.95±0.05±0.08	2 KARCH	90 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
1.00±0.08±0.10	2,3 ANTREASYAN	87 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
¹ Reevaluated by us using $B(\eta \rightarrow \gamma\gamma) = (39.21 \pm 0.34)\%$. Supersedes ANTREASYAN 87 and KARCH 90.			
² Superseded by KARCH 92.			
³ Using $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$.			

NODE=M002G2
NODE=M002G2

NODE=M002G2;LINKAGE=K4

NODE=M002G2;LINKAGE=A
NODE=M002G2;LINKAGE=D

$\eta'(958) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M002224

 $\Gamma(\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_{23}/\Gamma$ NODE=M002G01
NODE=M002G01

VALUE (10^{-3} eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	¹ AKHMETSHIN 15	CMD3	$0.958 e^+e^- \rightarrow \pi^+\pi^-\eta$
¹ AKHMETSHIN 15 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta) \times \Gamma(\eta'(958) \rightarrow e^+e^-)]/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 4.1 \times 10^{-4}$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.				

NODE=M002G01;LINKAGE=A

 $\eta'(958) \text{ BRANCHING RATIOS}$

NODE=M002230

 $\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$ NODE=M002R47
NODE=M002R47

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.429±0.007 OUR FIT				

••• We do not use the following data for averages, fits, limits, etc. •••

0.424±0.011±0.004 1.2k ¹ PEDLAR 09 CLEO $J/\psi \rightarrow \gamma\eta'$ ¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

NODE=M002R47;LINKAGE=PE

 $\Gamma(\pi^+\pi^-\eta(\text{charged decay}))/\Gamma_{\text{total}} \quad 0.286\Gamma_1/\Gamma$ NODE=M002R3
NODE=M002R3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1228±0.0020 OUR FIT				

••• We do not use the following data for averages, fits, limits, etc. •••

0.123 ±0.014 107 RITTENBERG 69 HBC $1.7-2.7 K^-p$ 0.10 ±0.04 10 LONDON 66 HBC $2.24 K^-p \rightarrow \Lambda 2\pi^+ 2\pi^-\pi^0$ 0.07 ±0.04 7 BADIER 65B HBC $3 K^-p$ $\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))/\Gamma_{\text{total}} \quad 0.714\Gamma_1/\Gamma$ NODE=M002R1
NODE=M002R1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.307±0.005 OUR FIT				

••• We do not use the following data for averages, fits, limits, etc. •••

0.314±0.026 281 RITTENBERG 69 HBC $1.7-2.7 K^-p$ $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$ NODE=M002R6
NODE=M002R6
NEW

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.291±0.005 OUR FIT				

[0.291 ± 0.006 OUR 2015 FIT]

••• We do not use the following data for averages, fits, limits, etc. •••

0.287±0.007±0.004 0.2k ¹ PEDLAR 09 CLEO $J/\psi \rightarrow \gamma\eta'$ 0.329±0.033 298 RITTENBERG 69 HBC $1.7-2.7 K^-p$ 0.2 ±0.1 20 LONDON 66 HBC $2.24 K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$ 0.34 ±0.09 35 BADIER 65B HBC $3 K^-p$ ¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

NODE=M002R6;LINKAGE=PE

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta) \quad \Gamma_2/\Gamma_1$ NODE=M002R43
NODE=M002R43
NEW

VALUE	DOCUMENT ID	TECN	COMMENT
0.678±0.017 OUR FIT			

[0.677 ± 0.017 OUR 2015 FIT]

0.683±0.020 OUR AVERAGE0.677±0.024±0.011 PEDLAR 09 CLE3 $J/\psi \rightarrow \eta'\gamma$ 0.69 ±0.03 ABLIKIM 06E BES2 $J/\psi \rightarrow \eta'\gamma$ $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta(\text{neutral decay})) \quad \Gamma_2/0.714\Gamma_1$ NODE=M002R27
NODE=M002R27
NEW

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.950±0.024 OUR FIT				

[0.949 ± 0.024 OUR 2015 FIT]

0.97 ±0.09 OUR AVERAGE0.70 ±0.22 AMSLER 04B CBAR $0 \bar{p}p \rightarrow \pi^+\pi^-\eta$ 1.07 ±0.17 BELADIDZE 92C VES $36 \pi^-\text{Be} \rightarrow \pi^-\eta'\eta\text{Be}$ 0.92 ±0.14 473 DANBURG 73 HBC $2.2 K^-p \rightarrow \Lambda X^0$ 1.11 ±0.18 192 JACOBS 73 HBC $2.9 K^-p \rightarrow \Lambda X^0$

$\Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.223±0.008 OUR FIT

[0.222 ± 0.008 OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.235±0.013±0.004 3.2k ¹ PEDLAR 09 CLEO $J/\psi \rightarrow \gamma\eta'$ ¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.NODE=M002R48
NODE=M002R48
NEW

NODE=M002R48;LINKAGE=PE

 $\Gamma(\pi^0\pi^0\eta(3\pi^0\text{ decay}))/\Gamma_{\text{total}}$ $0.321\Gamma_3/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0716±0.0026 OUR FIT

[0.0712 ± 0.0026 OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.11 ± 0.06 4 BENSINGER 70 DBC $2.2\pi^+d$ NODE=M002R26
NODE=M002R26
NEW $\Gamma(\pi^0\pi^0\eta)/\Gamma(\pi^+\pi^-\eta)$ Γ_3/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.519±0.026 OUR FIT

[0.517 ± 0.026 OUR 2015 FIT]

0.555±0.043±0.013PEDLAR 09 CLE3 $J/\psi \rightarrow \eta'\gamma$ NODE=M002R45
NODE=M002R45
NEW $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi\pi\eta)$ $\Gamma_2/(\Gamma_1+\Gamma_3)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.446±0.012 OUR FIT

[0.447 ± 0.012 OUR 2015 FIT]

0.43 ± 0.02 ± 0.02BARBERIS 98C OMEG 450 $p\bar{p} \rightarrow p_f\eta'p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.15 DAVIS 68 HBC $5.5K^-p$ NODE=M002R7
NODE=M002R7
NEW $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.62±0.13 OUR FIT

[0.0275 ± 0.0023 OUR 2015 FIT]

2.55±0.03±0.1633.2k ¹ ABLIKIM 15AD BES3 $J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.34±0.30±0.04 70 ² PEDLAR 09 CLEO $J/\psi \rightarrow \gamma\eta'$ ¹ Using $B(J/\psi \rightarrow \eta'\gamma) = (5.15 \pm 0.16) \times 10^{-3}$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.² Not independent of other η' branching fractions and ratios in PEDLAR 09.NODE=M002R49
NODE=M002R49
NEWNODE=M002R49;LINKAGE=A
NODE=M002R49;LINKAGE=PE $\Gamma(\omega\gamma)/\Gamma(\pi^+\pi^-\eta)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0610±0.0033 OUR FIT

[0.064 ± 0.006 OUR 2015 FIT]

0.055 ± 0.007 ± 0.001PEDLAR 09 CLE3 $J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.068 ± 0.013 68 ZANFINO 77 ASPK $8.4\pi^-p$ NODE=M002R17
NODE=M002R17
NEW $\Gamma(\omega\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.117±0.007 OUR FIT

[0.124 ± 0.011 OUR 2015 FIT]

0.147±0.016ALDE 87B GAM2 $38\pi^-p \rightarrow n4\gamma$ NODE=M002R33
NODE=M002R33
NEW $\Gamma(\omega e^+e^-)/\Gamma(\omega\gamma)$ Γ_5/Γ_4

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.71±1.34±0.54 ¹ ABLIKIM 15AD BES3 $J/\psi \rightarrow \eta'\gamma$ ¹ Obtained from other ABLIKIM 15AD measurements with common systematics taken into account.NODE=M002R60
NODE=M002R60

NODE=M002R60;LINKAGE=A

 $\Gamma(\omega e^+e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.97±0.34±0.1766 ¹ ABLIKIM 15AD BES3 $J/\psi \rightarrow \eta'\gamma$ ¹ Using $B(J/\psi \rightarrow \eta'\gamma) = (5.15 \pm 0.16) \times 10^{-3}$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.NODE=M002R59
NODE=M002R59

NODE=M002R59;LINKAGE=A

$$\frac{\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / [\Gamma(\pi^+ \pi^- \eta) + \Gamma(\pi^0 \pi^0 \eta) + \Gamma(\omega \gamma)]}{\Gamma_2 / (\Gamma_1 + \Gamma_3 + \Gamma_4)}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.429 ± 0.011 OUR FIT

[0.428 ± 0.011 OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25 ± 0.14 DAUBER 64 HBC 1.95 $K^- p$

NODE=M002R18
 NODE=M002R18
 NEW

$$\frac{[\Gamma(\pi^0 \pi^0 \eta (\text{charged decay})) + \Gamma(\omega (\text{charged decay}) \gamma)] / \Gamma_{\text{total}}}{(0.286\Gamma_3 + 0.89\Gamma_4) / \Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0871 ± 0.0026 OUR FIT

[0.0880 ± 0.0031 OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.045 ± 0.029 42 RITTENBERG 69 HBC 1.7-2.7 $K^- p$

NODE=M002R4
 NODE=M002R4
 NEW

$$\frac{\Gamma(\pi^+ \pi^- \text{ neutrals}) / \Gamma_{\text{total}}}{(0.714\Gamma_1 + 0.286\Gamma_3 + 0.89\Gamma_4) / \Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.394 ± 0.004 OUR FIT

[0.395 ± 0.004 OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 ± 0.1 39 LONDON 66 HBC 2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^- \text{ neutrals}$ 0.35 ± 0.06 33 BADIER 65B HBC 3 $K^- p$

NODE=M002R2
 NODE=M002R2
 NEW

$$\frac{\Gamma(\gamma \gamma) / \Gamma_{\text{total}}}{\Gamma_6 / \Gamma}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.21 ± 0.08 OUR FIT[(2.20 ± 0.08) × 10⁻² OUR 2015 FIT]**2.00 ± 0.15 OUR AVERAGE**1.98^{+0.31}_{-0.27} ± 0.07 114 1 WICHT 08 BELL $B^\pm \rightarrow K^\pm \gamma \gamma$ 2.00 ± 0.18 2 STANTON 80 SPEC 8.45 $\pi^- p \rightarrow n \pi^+ \pi^- 2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.25 ± 0.16 ± 0.03 0.3k 3 PEDLAR 09 CLEO $J/\psi \rightarrow \gamma \eta'$ 1.8 ± 0.2 6000 4 APEL 79 NICE 15-40 $\pi^- p \rightarrow n 2\gamma$ 2.5 ± 0.7 DUANE 74 MMS $\pi^- p \rightarrow n \text{MM}$ 1.71 ± 0.33 68 DALPIAZ 72 CNTR 1.6 $\pi^- p \rightarrow n X^0$ 2.0^{+0.8}_{-0.6} 31 HARVEY 71 OSPK 3.65 $\pi^- p \rightarrow n X^0$

NODE=M002R19
 NODE=M002R19
 NEW

¹ WICHT 08 reports $[\Gamma(\eta'(958) \rightarrow \gamma \gamma) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta' K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta' K^+) = (7.06 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Includes APEL 79 result.³ Not independent of other η' branching fractions and ratios in PEDLAR 09.⁴ Data is included in STANTON 80 evaluation.

NODE=M002R19;LINKAGE=WI

NODE=M002R19;LINKAGE=S
 NODE=M002R19;LINKAGE=PE
 NODE=M002R19;LINKAGE=A

$$\frac{\Gamma(\gamma \gamma) / \Gamma(\pi^+ \pi^- \eta)}{\Gamma_6 / \Gamma_1}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0514 ± 0.0022 OUR FIT

[0.0513 ± 0.0022 OUR 2015 FIT]

0.053 ± 0.004 ± 0.001PEDLAR 09 CLE3 $J/\psi \rightarrow \eta' \gamma$

NODE=M002R46
 NODE=M002R46
 NEW

$$\frac{\Gamma(\gamma \gamma) / \Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma))}{\Gamma_6 / \Gamma_2}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0758 ± 0.0033 OUR FIT

[0.0757 ± 0.0033 OUR 2015 FIT]

0.080 ± 0.008ABLIKIM 06E BES2 $J/\psi \rightarrow \eta' \gamma$

NODE=M002R42
 NODE=M002R42
 NEW

$$\frac{\Gamma(\gamma \gamma) / \Gamma(\pi^0 \pi^0 \eta)}{\Gamma_6 / \Gamma_3}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.099 ± 0.004 OUR FIT**0.105 ± 0.010 OUR AVERAGE** Error includes scale factor of 1.9.0.091 ± 0.009 AMSLER 93 CBAR 0.0 $\bar{p} p$ 0.112 ± 0.002 ± 0.006 ALDE 87B GAM2 38 $\pi^- p \rightarrow n 2\gamma$

NODE=M002R38
 NODE=M002R38

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ (neutral decay) $\Gamma_6/0.714\Gamma_3$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M002R28
 NODE=M002R28

0.139±0.006 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.188±0.058	16	APEL	72	OSPK	$3.8 \pi^- p \rightarrow n X^0$
-------------	----	------	----	------	---------------------------------

 $\Gamma(\text{neutrals})/\Gamma_{\text{total}}$ $(0.714\Gamma_3+0.09\Gamma_4+\Gamma_6)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M002R5
 NODE=M002R5
 NEW

0.184±0.006 OUR FIT

[0.183 ± 0.006 OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.185±0.022	535	BASILE	71	CNTR	$1.6 \pi^- p \rightarrow n X^0$
-------------	-----	--------	----	------	---------------------------------

0.189±0.026	123	RITTENBERG	69	HBC	$1.7-2.7 K^- p$
-------------	-----	------------	----	-----	-----------------

 $\Gamma(3\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R55
 NODE=M002R55

2.20±0.20 OUR FIT[(2.14 ± 0.20) × 10⁻³ OUR 2015 FIT]**3.7 ± 0.4 OUR AVERAGE**[(3.6 ± 0.4) × 10⁻³ OUR 2015 AVERAGE]

4.79±0.59±1.14	183	¹ ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
----------------	-----	----------------------	-----	------	-----------------------------------

3.56±0.22±0.34	309	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
----------------	-----	---------	-----	------	-------------------------------------

¹We have added all systematic uncertainties in quadrature to a single value.

NEW

NEW

NODE=M002R55;LINKAGE=A

 $\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_7/Γ_3

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R32
 NODE=M002R32

99± 9 OUR FIT[(96 ± 9) × 10⁻⁴ OUR 2015 FIT]**78±10 OUR AVERAGE**

86±19	235	BLIK	08	GAMS	$32 \pi^- p \rightarrow \eta' n$
-------	-----	------	----	------	----------------------------------

74±15		ALDE	87B	GAM2	$38 \pi^- p \rightarrow n 6\gamma$
-------	--	------	-----	------	------------------------------------

75±18		BINON	84	GAM2	$30-40 \pi^- p \rightarrow n 6\gamma$
-------	--	-------	----	------	---------------------------------------

NEW

 $\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$ Γ_8/Γ_6

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R29
 NODE=M002R29

4.9±1.2	33	VIKTOROV	80	CNTR	$25,33 \pi^- p \rightarrow 2\mu\gamma$
----------------	----	----------	----	------	--

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R50
 NODE=M002R50

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.29	90	¹ ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
-------	----	----------------------	-----	------	----------------------------------

<2.4	90	² NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
------	----	-------------------	----	------	----------------------------------

¹Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.²Not independent of measured value of Γ_9/Γ_1 from NAIK 09.

NODE=M002R50;LINKAGE=A
 NODE=M002R50;LINKAGE=NA

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_9/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R03
 NODE=M002R03

<0.5	90	¹ NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
----------------	----	-------------------	----	------	----------------------------------

¹NAIK 09 reports [$\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)$] / [$B(\eta \rightarrow 2\gamma)$] < 1.3 × 10⁻³ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

NODE=M002R03;LINKAGE=NA

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\rho^0\gamma)$ (including non-resonant $\pi^+\pi^-\gamma$) Γ_9/Γ_2

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R57
 NODE=M002R57

<1.0	90	ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
----------------	----	---------	-----	------	----------------------------------

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R21
 NODE=M002R21

3.82±0.35 OUR FIT[(3.8 ± 0.4) × 10⁻³ OUR 2015 FIT]**3.9 ± 0.4 OUR AVERAGE**[(3.8 ± 0.4) × 10⁻³ OUR 2015 AVERAGE]

4.28±0.49±1.11	78	¹ ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
----------------	----	----------------------	-----	------	-----------------------------------

3.83±0.15±0.39	1014	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
----------------	------	---------	-----	------	--

3.7 ^{+1.1} _{-0.9} ± 0.4		² NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
---	--	-------------------	----	------	----------------------------------

¹We have added all systematic uncertainties in quadrature to a single value.²Not independent of measured value of Γ_{10}/Γ_1 from NAIK 09.

NEW

NEW

NODE=M002R21;LINKAGE=A
 NODE=M002R21;LINKAGE=NA

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{10}/Γ_1 VALUE (units 10^{-3}) CL% EVTS

DOCUMENT ID TECN COMMENT

8.9 ± 0.8 OUR FIT[(8.8 ± 0.9) × 10^{-3} OUR 2015 FIT]NODE=M002R01
NODE=M002R01
NEW**8.28^{+2.49}_{-2.12} ± 0.04**

20

¹ NAIK

09

CLEO

 $J/\psi \rightarrow \gamma\eta'$

¹ NAIK 09 reports [$\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)$] / [$B(\eta \rightarrow 2\gamma)$] = $(21^{+6}_{-5} \pm 2) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R01;LINKAGE=NA

 $\Gamma(\pi^0\rho^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE CL% EVTS

DOCUMENT ID TECN COMMENT

<0.04

90

RITTENBERG 65

HBC

 $2.7 K^- p$ NODE=M002R10
NODE=M002R10 $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{12}/Γ VALUE (units 10^{-5}) CL% EVTS

DOCUMENT ID TECN COMMENT

8.5 ± 0.9 ± 0.3

199

¹ ABLIKIM

14M

BES3

 $J/\psi \rightarrow \gamma\eta'$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 24

90

² NAIK

09

CLEO

 $J/\psi \rightarrow \gamma\eta'$

<1000

90

RITTENBERG 69

HBC

 $1.7-2.7 K^- p$

¹ ABLIKIM 14M reports [$\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}$] × [$B(J/\psi(1S) \rightarrow \gamma\eta'(958))$] = $(4.40 \pm 0.35 \pm 0.30) \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.15 \pm 0.16) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R24
NODE=M002R24

² Not independent of measured value of Γ_{12}/Γ_1 from NAIK 09.

NODE=M002R24;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{12}/Γ_1 VALUE (units 10^{-3}) CL% EVTS

DOCUMENT ID TECN COMMENT

<0.6

90

¹ NAIK

09

CLEO

 $J/\psi \rightarrow \gamma\eta'$

¹ NAIK 09 reports [$\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)$] / [$B(\eta \rightarrow 2\gamma)$] < 1.4×10^{-3} which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

NODE=M002R04
NODE=M002R04

NODE=M002R04;LINKAGE=NA

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ VALUE (units 10^{-4}) CL% EVTS

DOCUMENT ID TECN COMMENT

1.8 ± 0.4 ± 0.1

84

¹ ABLIKIM

14M

BES3

 $J/\psi \rightarrow \gamma\eta'$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<27

90

² NAIK

09

CLEO

 $J/\psi \rightarrow \gamma\eta'$

¹ ABLIKIM 14M reports [$\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$] × [$B(J/\psi(1S) \rightarrow \gamma\eta'(958))$] = $(9.38 \pm 1.79 \pm 0.89) \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.15 \pm 0.16) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R51;LINKAGE=A

² Not independent of measured value of Γ_{13}/Γ_1 from NAIK 09.

NODE=M002R51;LINKAGE=NA

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{13}/Γ_1 VALUE (units 10^{-3}) CL% EVTS

DOCUMENT ID TECN COMMENT

<6

90

¹ NAIK

09

CLEO

 $J/\psi \rightarrow \gamma\eta'$

¹ NAIK 09 reports [$\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)$] / [$B(\eta \rightarrow 2\gamma)$] < 15×10^{-3} which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

NODE=M002R05
NODE=M002R05

NODE=M002R05;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-)\text{ neutrals})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE CL% EVTS

DOCUMENT ID TECN COMMENT

<0.01

95

DANBURG 73

HBC

 $2.2 K^- p \rightarrow \Lambda X^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<0.01

90

RITTENBERG 69

HBC

 $1.7-2.7 K^- p$ NODE=M002R22
NODE=M002R22 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE CL% EVTS

DOCUMENT ID TECN COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<0.002

90

¹ NAIK

09

CLEO

 $J/\psi \rightarrow \gamma\eta'$

<0.01

90

RITTENBERG 69

HBC

 $1.7-2.7 K^- p$

¹ Not independent of measured value of Γ_{15}/Γ_1 from NAIK 09.

NODE=M002R23
NODE=M002R23

NODE=M002R23;LINKAGE=NA

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{15}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	¹ NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

NODE=M002R06
NODE=M002R06

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 11 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

NODE=M002R06;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	KALBFLEISCH 64B	HBC	$K^-p \rightarrow \Lambda 2(\pi^+\pi^-)+MM$

NODE=M002R16
NODE=M002R16

••• We do not use the following data for averages, fits, limits, etc. •••

<0.01	90	LONDON	66 HBC	Compilation
-------	----	--------	--------	-------------

 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.1	90	¹ ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

NODE=M002R07
NODE=M002R07

••• We do not use the following data for averages, fits, limits, etc. •••

< 53	90	² NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<500	95	KALBFLEISCH 64B	HBC	$K^-p \rightarrow \Lambda 2(\pi^+\pi^-)$

¹ Using $B(J/\psi \rightarrow \gamma\eta'(958)) = (5.16 \pm 0.15) \times 10^{-3}$.

² Not independent of measured value of Γ_{17}/Γ_1 from NAIK 09.

NODE=M002R07;LINKAGE=A
NODE=M002R07;LINKAGE=NA $\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{17}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

NODE=M002R08
NODE=M002R08

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 3(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 3.0 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

NODE=M002R08;LINKAGE=NA

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M002R12
NODE=M002R12**2.4 $\begin{smallmatrix} +1.3 \\ -1.0 \end{smallmatrix}$ OUR FIT**

••• We do not use the following data for averages, fits, limits, etc. •••

$2.11 \pm 0.12 \pm 0.14$	429	¹ ABLIKIM	130 BES3	$J/\psi \rightarrow \gamma\eta'$
--------------------------	-----	----------------------	----------	----------------------------------

$2.5 \begin{smallmatrix} +1.2 \\ -0.9 \end{smallmatrix} \pm 0.5$		² NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
--	--	-------------------	---------	----------------------------------

<6	90	RITTENBERG 65	HBC	2.7 K^-p
----	----	---------------	-----	------------

¹ Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.

² Not independent of measured value of Γ_{18}/Γ_1 from NAIK 09.

NODE=M002R12;LINKAGE=A
NODE=M002R12;LINKAGE=NA $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_{18}/Γ_1

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R02
NODE=M002R02**5.6 $\begin{smallmatrix} +3.0 \\ -2.2 \end{smallmatrix}$ OUR FIT**

$5.52 \begin{smallmatrix} +3.00 \\ -2.30 \end{smallmatrix} \pm 0.03$	8	¹ NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
--	---	-------------------	---------	----------------------------------

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-e^+e^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (14 \begin{smallmatrix} +7 \\ -5 \end{smallmatrix} \pm 3) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R02;LINKAGE=NA

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_{18}/Γ_2

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2\pm0.4\pm0.5	429	ABLIKIM	130 BES3	$J/\psi \rightarrow \gamma\eta'$

NODE=M002R56
NODE=M002R56 $\Gamma(\pi^+e^-\nu_e + \text{c.c.})/\Gamma(\pi^+\pi^-\eta)$ Γ_{19}/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	ABLIKIM	13G BES3	$J/\psi \rightarrow \phi\eta'$

NODE=M002R54
NODE=M002R54 $\Gamma(\gamma e^+e^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R40
NODE=M002R40

••• We do not use the following data for averages, fits, limits, etc. •••

<0.9	90	BRIERE	00 CLEO	10.6 e^+e^-
------	----	--------	---------	---------------

$\Gamma(\gamma e^+ e^-)/\Gamma(\gamma\gamma)$ Γ_{20}/Γ_6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.13 \pm 0.09 \pm 0.07$	864	ABLIKIM	150	BES3 $J/\psi \rightarrow \gamma e^+ e^-$

NODE=M002R00
NODE=M002R00 $\Gamma(\pi^0 \gamma\gamma)/\Gamma(\pi^0 \pi^0 \eta)$ Γ_{21}/Γ_3

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 37	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n 4\gamma$

NODE=M002R35
NODE=M002R35 $\Gamma(4\pi^0)/\Gamma_{total}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.2 \times 10^{-4}$	90	DONSKOV	14	GAM4 $32.5 \pi^- p \rightarrow \eta' n$

NODE=M002R58
NODE=M002R58 $\Gamma(4\pi^0)/\Gamma(\pi^0 \pi^0 \eta)$ Γ_{22}/Γ_3

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R37
NODE=M002R37

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n 8\gamma$
--------	----	------	-----	---

 $\Gamma(e^+ e^-)/\Gamma_{total}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M002R39
NODE=M002R39 $< 5.6 \times 10^{-9}$ (CL = 90%) [$< 2.1 \times 10^{-7}$ (CL = 90%) OUR 2014 BEST LIMIT]

$< 5.6 \times 10^{-9}$	90	¹ ACHASOV	15	SND $0.958 e^+ e^- \rightarrow \pi\pi\eta$
------------------------	----	----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 12 \times 10^{-9}$	90	² AKHMETSHIN	15	CMD3 $0.958 e^+ e^- \rightarrow \pi^+ \pi^- \eta$
-----------------------	----	-------------------------	----	---

$< 2.1 \times 10^{-7}$	90	VOROBYEV	88	ND $e^+ e^- \rightarrow \pi^+ \pi^- \eta$
------------------------	----	----------	----	---

¹ Combining data of ACHASOV 15 and AKHMETSHIN 15 and using $\Gamma(\eta') = 0.198 \pm 0.009$ MeV.² Using $\Gamma_{\eta'(958)} = 198 \pm 9$ keV, $B(\eta'(958) \rightarrow \pi^+ \pi^- \eta) = (42.9 \pm 0.7)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.20)\%$.

NODE=M002R39;LINKAGE=B

NODE=M002R39;LINKAGE=A

 $\Gamma(\text{invisible})/\Gamma_{total}$ Γ_{24}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R52
NODE=M002R52

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9.5	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
---------	----	-------------------	----	---------------------------------------

¹ Not independent of measured value of Γ_{24}/Γ_1 from NAIK 09.

NODE=M002R52;LINKAGE=NA

 $\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$ Γ_{24}/Γ_6

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R44
NODE=M002R44

< 2.4	90	ABLIKIM	13	BES3 $J/\psi \rightarrow \phi\eta'$
---------	----	---------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.69	90	ABLIKIM	06Q	BES $J/\psi \rightarrow \phi\eta'$
----------	----	---------	-----	------------------------------------

 $\Gamma(\text{invisible})/\Gamma(\pi^+ \pi^- \eta)$ Γ_{24}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R09
NODE=M002R09

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.1	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
---------	----	-------------------	----	---------------------------------------

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \text{invisible})/\Gamma(\eta'(958) \rightarrow \pi^+ \pi^- \eta)] / [B(\eta \rightarrow 2\gamma)]$ $< 5.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

NODE=M002R09;LINKAGE=NA

 $\Gamma(\pi^+ \pi^-)/\Gamma_{total}$ Γ_{25}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R20
NODE=M002R20

< 0.6	90	¹ ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^+ \pi^-$
---------	----	----------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 29	90	² MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+ \pi^-$
--------	----	-------------------	-----	---

< 3.3	90	³ MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+ \pi^-$
---------	----	-------------------	-----	---

OCCUR=2

< 800	95	DANBURG	73	HBC $2.2 K^- p \rightarrow \Lambda X^0$
---------	----	---------	----	---

< 200	90	RITTENBERG	69	HBC $1.7-2.7 K^- p$
---------	----	------------	----	---------------------

¹ ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+ \pi^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.15 \times 10^{-3}$.² Taking into account interference with the $\gamma\gamma \rightarrow \pi^+ \pi^-$ continuum.³ Without interference with the $\gamma\gamma \rightarrow \pi^+ \pi^-$ continuum.

NODE=M002R20;LINKAGE=AL

NODE=M002R20;LINKAGE=MO

NODE=M002R20;LINKAGE=MR

$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-4}$	90	¹ ABLIKIM	11G BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$
¹ ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.15 \times 10^{-3}$.				

 Γ_{26}/Γ NODE=M002R53
NODE=M002R53

NODE=M002R53;LINKAGE=AL

 $\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<45	90	ALDE	87B GAM2	$38 \pi^- p \rightarrow n4\gamma$

 Γ_{26}/Γ_3 NODE=M002R36
NODE=M002R36 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	90	BRIERE	00 CLEO	$10.6 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<13	90	RITTENBERG	65 HBC	$2.7 K^- p$

 Γ_{27}/Γ NODE=M002R8
NODE=M002R8 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.4	90	BRIERE	00 CLEO	$10.6 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<11	90	RITTENBERG	65 HBC	$2.7 K^- p$

 Γ_{28}/Γ NODE=M002R9
NODE=M002R9 $\Gamma(3\gamma)/\Gamma(\pi^0\pi^0\eta)$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.6	90	ALDE	87B GAM2	$38 \pi^- p \rightarrow n3\gamma$

 Γ_{29}/Γ_3 NODE=M002R34
NODE=M002R34 $\Gamma(\mu^+ \mu^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	90	DZHELADIN	81 CNTR	$30 \pi^- p \rightarrow \eta' n$

 Γ_{30}/Γ NODE=M002R31
NODE=M002R31 $\Gamma(\mu^+ \mu^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	DZHELADIN	81 CNTR	$30 \pi^- p \rightarrow \eta' n$

 Γ_{31}/Γ NODE=M002R30
NODE=M002R30 $\Gamma(e\mu)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.7	90	BRIERE	00 CLEO	$10.6 e^+ e^-$

 Γ_{32}/Γ NODE=M002R41
NODE=M002R41 $\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

NODE=M002225

$$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha Y|^2 + CX + DX^2$$

X and Y are Dalitz variables; α is complex and C , and D are real-valued. Parameters C and D are not necessarily equal to c and d , respectively, in the generalized parameterization following this one. May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays. Because of different initial assumptions and strong correlations of the parameters we do not average the parameters in the section below.

NODE=M002225

 $Re(\alpha)$ decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.033 \pm 0.005 \pm 0.003$	44k	¹ ABLIKIM	11 BES3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$-0.072 \pm 0.012 \pm 0.006$	7k	² AMELIN	05A VES	$28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
$-0.021 \pm 0.018 \pm 0.017$	6.7k	³ BRIERE	00 CLEO	$10.6 e^+ e^- \rightarrow \eta\pi^+\pi^- X$
$-0.058 \pm 0.013 \pm 0.003$	5.4k	⁴ ALDE	86 GAM2	$38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
-0.08 ± 0.03		^{4,5} KALBFLEISCH	74 RVUE	$\eta' \rightarrow \eta\pi^+\pi^-$

NODE=M002A0
NODE=M002A0¹ See ABLIKIM 11 for the full correlation matrix.² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.³ Assuming $\text{Im}(\alpha) = 0$, $C = 0$, and $D = 0$.⁴ Assuming $C = 0$.⁵ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.NODE=M002A0;LINKAGE=AB
NODE=M002A0;LINKAGE=AMNODE=M002A0;LINKAGE=BR
NODE=M002A0;LINKAGE=A
NODE=M002A0;LINKAGE=KA

$Im(\alpha)$ decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.000±0.049±0.001	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.0 ±0.1 ±0.0	7k	² AMELIN 05A	VES	$28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$
-0.00 ±0.13 ±0.00	5.4k	³ ALDE 86	GAM2	$38 \pi^- p \rightarrow n \eta \pi^0 \pi^0$
0.0 ±0.3		^{3,4} KALBFLEISCH 74	RVUE	$\eta' \rightarrow \eta \pi^+ \pi^-$

¹ See ABLIKIM 11 for the full correlation matrix.

² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

³ Assuming $C = 0$.

⁴ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002IA0
NODE=M002IA0

NODE=M002IA0;LINKAGE=AB
NODE=M002IA0;LINKAGE=AM

NODE=M002IA0;LINKAGE=A
NODE=M002IA0;LINKAGE=KA

C decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
+0.018±0.009±0.003	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.020±0.018±0.004	7k	² AMELIN 05A	VES	$28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.

² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

NODE=M002C0
NODE=M002C0

NODE=M002C0;LINKAGE=AB
NODE=M002C0;LINKAGE=AM

D decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.059±0.012±0.004	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
-0.066±0.030±0.015	7k	² AMELIN 05A	VES	$28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$
0.00 ±0.03 ±0.00	5.4k	³ ALDE 86	GAM2	$38 \pi^- p \rightarrow n \eta \pi^0 \pi^0$
0		^{3,4} KALBFLEISCH 74	RVUE	$\eta' \rightarrow \eta \pi^+ \pi^-$

¹ See ABLIKIM 11 for the full correlation matrix.

² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

³ Assuming $C = 0$.

⁴ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002D0
NODE=M002D0

NODE=M002D0;LINKAGE=AB
NODE=M002D0;LINKAGE=AM

NODE=M002D0;LINKAGE=AL
NODE=M002D0;LINKAGE=KA

 $\eta'(958) \rightarrow \eta \pi \pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 \propto 1 + a Y + b Y^2 + c X + d X^2$$

X and Y are Dalitz variables and a , b , c , and d are real-valued parameters. May be different for $\eta'(958) \rightarrow \eta \pi^+ \pi^-$ and $\eta'(958) \rightarrow \eta \pi^0 \pi^0$ decays. We do not average measurements in the section below because parameter values from each experiment are strongly correlated.

NODE=M002227

NODE=M002227

a decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.047±0.011±0.003	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
-0.066±0.016±0.003	15k	² BLIK 09	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
-0.127±0.016±0.008	20k	³ DOROFEEV 07	VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.

² From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

³ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

NODE=M002DPA
NODE=M002DPA

NODE=M002DPA;LINKAGE=AB
NODE=M002DPA;LINKAGE=BL
NODE=M002DPA;LINKAGE=DO

b decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.069±0.019±0.009	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
-0.063±0.028±0.004	15k	² BLIK 09	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
-0.106±0.028±0.014	20k	³ DOROFEEV 07	VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.

² From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

³ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

NODE=M002DPB
NODE=M002DPB

NODE=M002DPB;LINKAGE=AB
NODE=M002DPB;LINKAGE=BL
NODE=M002DPB;LINKAGE=DO

c decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$+0.019 \pm 0.011 \pm 0.003$	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.107 \pm 0.096 \pm 0.003$	15k	² BLIK 09	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$0.015 \pm 0.011 \pm 0.014$	20k	³ DOROFEEV 07	VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.

² From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

³ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

NODE=M002DPC
NODE=M002DPC

NODE=M002DPC;LINKAGE=AB
NODE=M002DPC;LINKAGE=BL
NODE=M002DPC;LINKAGE=DO

d decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.073 \pm 0.012 \pm 0.003$	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.018 \pm 0.078 \pm 0.006$	15k	² BLIK 09	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$-0.082 \pm 0.017 \pm 0.008$	20k	³ DOROFEEV 07	VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.

² From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay. If $c \equiv 0$ from Bose-Einstein symmetry, $d = -0.067 \pm 0.020 \pm 0.003$.

³ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

NODE=M002DPD
NODE=M002DPD

NODE=M002DPD;LINKAGE=AB
NODE=M002DPD;LINKAGE=BL

NODE=M002DPD;LINKAGE=DO

$\eta'(958) \beta$ PARAMETER

|MATRIX ELEMENT|² = (1 + 2βZ)

See the "Note on η Decay Parameters" in our 1994 edition Physical Review D50 1173 (1994), p. 1454.

NODE=M002226

NODE=M002226

β decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.61 ± 0.08 OUR AVERAGE Error includes scale factor of 1.2. [-0.46 ± 0.22 OUR 2015 AVERAGE Scale factor = 1.4]				
$-0.640 \pm 0.046 \pm 0.047$	1.8k	ABLIKIM	15G BES3	$J/\psi \rightarrow \gamma (\pi^0 \pi^0 \pi^0)$
-0.59 ± 0.18	235	BLIK	08 GAMS	$32 \pi^- p \rightarrow \eta' n$
-0.1 ± 0.3		ALDE	87B GAM2	$38 \pi^- p \rightarrow n 3 \pi^0$

NODE=M002B0
NODE=M002B0
NEW

 $\eta'(958) C$ -NONCONSERVING DECAY PARAMETER

See the note on η decay parameters in the Stable Particle Particle Listings for definition of this parameter.

NODE=M002235

NODE=M002235

DECAY ASYMMETRY PARAMETER FOR $\pi^+ \pi^- \gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.03 ± 0.04 OUR AVERAGE				
-0.019 ± 0.056		AIHARA 87	TPC	$2\gamma \rightarrow \pi^+ \pi^- \gamma$
-0.069 ± 0.078	295	GRIGORIAN 75	STRC	$2.1 \pi^- p$
0.00 ± 0.10	103	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.07 ± 0.08	152	RITTENBERG 65	HBC	$2.1-2.7 K^- p$

NODE=M002A
NODE=M002A

 $\eta'(958) \Rightarrow \gamma \ell^+ \ell^-$ TRANSITION FORM FACTOR SLOPE

Related to the effective virtual meson mass Λ , via slope $\approx \Lambda^{-2}$. See e.g. LANDSBERG 85, eq. (3.8), for a detailed definition.

VALUE (GeV ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
1.62 ± 0.17 OUR AVERAGE				
$1.60 \pm 0.17 \pm 0.08$	864	¹ ABLIKIM 150	BES3	$J/\psi \rightarrow \gamma e^+ e^-$
1.7 ± 0.4	33	¹ VIKTOROV 80		$25,33 \pi^- p \rightarrow 2\mu\gamma$

¹ In the single-pole Ansatz where slope = $1/(\Lambda^2 + \gamma^2)$ with Λ , γ being a Breit-Wigner mass, width for the effective contributing vector meson.

NODE=M002FFL
NODE=M002FFL

NODE=M002FFL

NODE=M002FFL;LINKAGE=A

η' (958) REFERENCES

NODE=M002

ABLIKIM	15AD	PR D92 051101	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56983
ABLIKIM	15G	PR D92 012014	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56602
ABLIKIM	15O	PR D92 012001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56780
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56781
ACHASOV	15	PR D91 092010	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=56788
AKHMETSHIN	15	PL B740 273	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=56386
ABLIKIM	14M	PRL 112 251801	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55904
DONSKOV	14	MPL A29 1450213	S. Donskov <i>et al.</i>	(GAMS-4 π Collab.)	REFID=56321
PDG	14	CPC 38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ABLIKIM	13	PR D87 012009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54829
ABLIKIM	13G	PR D87 032006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54952
ABLIKIM	13O	PR D87 092011	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55388
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55582
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54270
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53646
ABLIKIM	11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53711
CZERWINSKI	10	PRL 105 122001	E. Czerwinski <i>et al.</i>	(COSY-11 Collab.)	REFID=53364
BLIK	09	PAN 72 231	A.M. Blik <i>et al.</i>	(IHEP (Protvino))	REFID=52727
NAIK	09	PRL 102 061801	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52678
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
BLIK	08	PAN 71 2124	A. Blik <i>et al.</i>	(GAMS-4 π Collab.)	REFID=52663
LIBBY	08	PRL 101 182002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=52591
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
DOROFEEV	07	PL B651 22	V. Dorofeev <i>et al.</i>	(VES Collab.)	REFID=51711
MORI	07A	JPSJ 76 074102	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51691
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM	06Q	PRL 97 202002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51487
AMELIN	05A	PAN 68 372	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=50766
AMSLER	04B	EPJ C33 23	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51079
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
BRIERE	00	PRL 84 26	R. Briere <i>et al.</i>	(CLEO Collab.)	REFID=47410
ACCIARRI	98Q	PL B418 399	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46316
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
WURZINGER	96	PL B374 283	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=44992
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)	REFID=43653
AMSLER	93	ZPHY C58 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43311
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)	REFID=43175
KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=42170
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BEHREND	91	ZPHY C49 401	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41497
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
KARCH	90	PL B249 353	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=41377
ROE	90	PR D41 17	N.A. Roe <i>et al.</i>	(ASP Collab.)	REFID=41014
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40564
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
WILLIAMS	88	PR D38 1365	D.A. Williams <i>et al.</i>	(Crystal Ball Collab.)	REFID=40567
AIHARA	87	PR D35 2650	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.) JP	REFID=40009
ALBRECHT	87B	PL B199 457	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40265
ALDE	87B	ZPHY C36 603	D.M. Alde <i>et al.</i>	(LANL, BELG, SERP, LAPP)	REFID=40236
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=40008
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
ALDE	86	PL B177 115	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=20310
BARTEL	85E	PL 160B 421	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=10843
LANDSBERG	85	PRPL 128 301	L.G. Landsberg	(SERP)	REFID=10844
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=20305
BERGER	84B	PL 142B 125	C. Berger	(PLUTO Collab.)	REFID=20306
BINON	84	PL 140B 264	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP+)	REFID=20307
BEHREND	83B	PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
Also		PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20303
JENNI	83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)	REFID=20304
BARTEL	82B	PL 113B 190	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=20300
DZHEL'YADIN	81	PL 105B 239	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10836
STANTON	80	PL B92 353	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGL+)	REFID=40294
VIKTOROV	80	SJNP 32 520	V.A. Viktorov <i>et al.</i>	(SERP)	REFID=20298
APEL	79	PL 83B 131	W.D. Apel, K.H. Augenstein, E. Bertolucci	(KARLK+)	REFID=20295
BINNIE	79	PL 83B 141	D.M. Binnie <i>et al.</i>	(LOIC)	REFID=20296
ZANFINO	77	PRL 38 930	C. Zanfino <i>et al.</i>	(CARL, MCGL, OHIO+)	REFID=20293
GRIGORIAN	75	NP B91 232	A. Grigorian <i>et al.</i>	(+)	REFID=20287
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
DUANE	74	PRL 32 425	A. Duane <i>et al.</i>	(LOIC, SHMP)	REFID=20284
KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)	REFID=20286
DANBURG	73	PR D8 3744	J.S. Danburg <i>et al.</i>	(BNL, MICH) JP	REFID=20280
JACOBS	73	PR D8 18	S.M. Jacobs <i>et al.</i>	(BRAN, UMD, SYRA+)	REFID=20281
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72	PL 40B 680	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20275
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz <i>et al.</i>	(CERN)	REFID=20278
BASILE	71	NC 3A 371	M. Basile <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20270
HARVEY	71	PRL 27 885	E.H. Harvey <i>et al.</i>	(MINN, MICH)	REFID=20272
BENSINGER	70	PL 33B 505	J.R. Bensinger <i>et al.</i>	(WISC)	REFID=20268
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL) I	REFID=20266
DAVIS	68	PL 27B 532	R. Davis <i>et al.</i>	(NWES, ANL)	REFID=20263
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) JP	REFID=11774
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SAFL, AMST)	REFID=20253
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)	REFID=10761
DAUBER	64	PRL 13 449	P.M. Dauber <i>et al.</i>	(UCLA) JP	REFID=20247
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP	REFID=20252

$f_0(980)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See also the minireview on scalar mesons under $f_0(500)$. (See the index for the page number.)

NODE=M003

NODE=M003

 $f_0(980)$ MASS

NODE=M003M1

NODE=M003M1

→ UNCHECKED ←

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
990 ± 20				OUR ESTIMATE
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
989.4 ± 1.3	424	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
989.9 ± 0.4	706	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma 3\pi$
1003 $\begin{smallmatrix} +5 \\ -27 \end{smallmatrix}$		1,2 GARCIA-MAR..11	RVUE	Compilation
996 ± 7		1,3 GARCIA-MAR..11	RVUE	Compilation
996 $\begin{smallmatrix} +4 \\ -14 \end{smallmatrix}$		4 MOUSSALLAM11	RVUE	Compilation
981 ± 43		5 MENNESSIER	10 RVUE	Compilation
1030 $\begin{smallmatrix} +30 \\ -10 \end{smallmatrix}$		6 ANISOVICH	09 RVUE	0.0 $\bar{p}p, \pi N$
977 $\begin{smallmatrix} +11 \\ -9 \end{smallmatrix}$ ± 1	44	7 ECKLUND	09 CLEO	4.17 $e^+e^- \rightarrow D_s^- D_s^{*+} + \text{c.c.}$
982.2 ± 1.0 $\begin{smallmatrix} +8.1 \\ -8.0 \end{smallmatrix}$		8 UEHARA	08A BELL	10.6 $e^+e^- \rightarrow e^+e^- \pi^0 \pi^0$
976.8 ± 0.3 $\begin{smallmatrix} +10.1 \\ -0.6 \end{smallmatrix}$	64k	9 AMBROSINO	07 KLOE	1.02 $e^+e^- \rightarrow \pi^0 \pi^0 \gamma$
984.7 ± 0.4 $\begin{smallmatrix} +2.4 \\ -3.7 \end{smallmatrix}$	64k	10 AMBROSINO	07 KLOE	1.02 $e^+e^- \rightarrow \pi^0 \pi^0 \gamma$
973 ± 3	262 ± 30	11 AUBERT	07AKBABR	10.6 $e^+e^- \rightarrow \phi \pi^+ \pi^- \gamma$
970 ± 7	54 ± 9	11 AUBERT	07AKBABR	10.6 $e^+e^- \rightarrow \phi \pi^0 \pi^0 \gamma$
953 ± 20	2.6k	12 BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
985.6 $\begin{smallmatrix} +1.2+1.1 \\ -1.5-1.6 \end{smallmatrix}$		13 MORI	07 BELL	10.6 $e^+e^- \rightarrow e^+e^- \pi^+ \pi^-$
983.0 ± 0.6 $\begin{smallmatrix} +4.0 \\ -3.0 \end{smallmatrix}$		14 AMBROSINO	06B KLOE	1.02 $e^+e^- \rightarrow \pi^+ \pi^- \gamma$
977.3 ± 0.9 $\begin{smallmatrix} +3.7 \\ -4.3 \end{smallmatrix}$		15 AMBROSINO	06B KLOE	1.02 $e^+e^- \rightarrow \pi^+ \pi^- \gamma$
950 ± 9	4286	16 GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
965 ± 10		17 ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$, $\phi K^+ K^-$
1031 ± 8		18 ANISOVICH	03 RVUE	
1037 ± 31		TIKHOMIROV	03 SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
973 ± 1	2438	19 ALOISIO	02D KLOE	$e^+e^- \rightarrow \pi^0 \pi^0 \gamma$
977 ± 3 ± 2	848	20 AITALA	01A E791	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
969.8 ± 4.5	419	21 ACHASOV	00H SND	$e^+e^- \rightarrow \pi^0 \pi^0 \gamma$
985 $\begin{smallmatrix} +16 \\ -12 \end{smallmatrix}$	419	22,23 ACHASOV	00H SND	$e^+e^- \rightarrow \pi^0 \pi^0 \gamma$
976 ± 5 ± 6		24 AKHMETSHIN	99B CMD2	$e^+e^- \rightarrow \pi^+ \pi^- \gamma$
977 ± 3 ± 6	268	24 AKHMETSHIN	99C CMD2	$e^+e^- \rightarrow \pi^0 \pi^0 \gamma$
975 ± 4 ± 6		25 AKHMETSHIN	99C CMD2	$e^+e^- \rightarrow \pi^0 \pi^0 \gamma$
975 ± 4 ± 6		26 AKHMETSHIN	99C CMD2	$e^+e^- \rightarrow \pi^+ \pi^- \gamma$, $\pi^0 \pi^0 \gamma$
985 ± 10		BARBERIS	99 OMEG	450 $pp \rightarrow p_S p_f K^+ K^-$
982 ± 3		BARBERIS	99B OMEG	450 $pp \rightarrow p_S p_f \pi^+ \pi^-$
982 ± 3		BARBERIS	99C OMEG	450 $pp \rightarrow p_S p_f \pi^0 \pi^0$
987 ± 6 ± 6		27 BARBERIS	99D OMEG	450 $pp \rightarrow K^+ K^-$, $\pi^+ \pi^-$
989 ± 15		BELLAZZINI	99 GAM4	450 $pp \rightarrow pp \pi^0 \pi^0$
991 ± 3		28 KAMINSKI	99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
~ 980		28 OLLER	99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
~ 993.5		OLLER	99B RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
~ 987		28 OLLER	99C RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$
957 ± 6		29 ACKERSTAFF	98Q OPAL	$Z \rightarrow f_0 X$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

960 ± 10		ALDE	98	GAM4		
1015 ± 15		28 ANISOVICH	98B	RVUE	Compilation	
1008		30 LOCHER	98	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
955 ± 10		29 ALDE	97	GAM2	450 $p\bar{p} \rightarrow p\bar{p}\pi^0\pi^0$	
994 ± 9		31 BERTIN	97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
993.2 ± 6.5 ± 6.9		32 ISHIDA	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
1006		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$	
997 ± 5	3k	33 ALDE	95B	GAM2	38 $\pi^-p \rightarrow \pi^0\pi^0n$	
960 ± 10	10k	34 ALDE	95B	GAM2	38 $\pi^-p \rightarrow \pi^0\pi^0n$	OCCUR=2
994 ± 5		AMSLER	95B	CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$	
~ 996		35 AMSLER	95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\pi^0,$ $\pi^0\eta\eta, \pi^0\pi^0\eta$	
987 ± 6		36 ANISOVICH	95	RVUE		
1015		JANSSEN	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
983		37 BUGG	94	RVUE	$\bar{p}p \rightarrow \eta 2\pi^0$	
973 ± 2		38 KAMINSKI	94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
988		39 ZOU	94B	RVUE		
988 ± 10		40 MORGAN	93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}),$ $J/\psi \rightarrow \phi\pi\pi(K\bar{K}),$ $D_s \rightarrow \pi(\pi\pi)$	
971.1 ± 4.0		29 AGUILAR-...	91	EHS	400 $p\bar{p}$	
979 ± 4		41 ARMSTRONG	91	OMEG	300 $p\bar{p} \rightarrow p\bar{p}\pi\pi,$ $p\bar{p}K\bar{K}$	
956 ± 12		BREAKSTONE	90	SFM	$p\bar{p} \rightarrow p\bar{p}\pi^+\pi^-$	
959.4 ± 6.5		29 AUGUSTIN	89	DM2	$J/\psi \rightarrow \omega\pi^+\pi^-$	
978 ± 9		29 ABACHI	86B	HRS	$e^+e^- \rightarrow \pi^+\pi^-X$	
985.0 ^{+9.0} _{-39.0}		ETKIN	82B	MPS	23 $\pi^-p \rightarrow n 2K_S^0$	
974 ± 4		41 GIDAL	81	MRK2	$J/\psi \rightarrow \pi^+\pi^-X$	
975		42 ACHASOV	80	RVUE		
986 ± 10		41 AGUILAR-...	78	HBC	0.7 $\bar{p}p \rightarrow K_S^0 K_S^0$	
969 ± 5		41 LEEPER	77	ASPK	2-2.4 $\pi^-p \rightarrow$ $\pi^+\pi^-n, K^+K^-n$	
987 ± 7		41 BINNIE	73	CNTR	$\pi^-p \rightarrow nMM$	
1012 ± 6		43 GRAYER	73	ASPK	17 $\pi^-p \rightarrow \pi^+\pi^-n$	
1007 ± 20		43 HYAMS	73	ASPK	17 $\pi^-p \rightarrow \pi^+\pi^-n$	
997 ± 6		43 PROTOPOP...	73	HBC	7 $\pi^+p \rightarrow \pi^+p\pi^+\pi^-$	
		1 Quoted number refers to real part of pole position.				NODE=M003M1;LINKAGE=GC
		2 Analytic continuation using Roy equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.				NODE=M003M1;LINKAGE=GM
		3 Analytic continuation using GKPY equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.				NODE=M003M1;LINKAGE=GI
		4 Pole position. Used Roy equations.				NODE=M003M1;LINKAGE=MU
		5 Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.				NODE=M003M1;LINKAGE=ME
		6 On sheet II in a 2-pole solution. The other pole is found on sheet III at (850-100i) MeV				NODE=M003M1;LINKAGE=AO
		7 Using a relativistic Breit-Wigner function and taking into account the finite D_s mass.				NODE=M003M1;LINKAGE=EC
		8 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0} \pi\pi = 0$.				NODE=M003M1;LINKAGE=UE
		9 In the kaon-loop fit.				NODE=M003M1;LINKAGE=AK
		10 In the no-structure fit.				NODE=M003M1;LINKAGE=AS
		11 Systematic errors not estimated.				NODE=M003M1;LINKAGE=NS
		12 FLATTE 76 parameterization. $g_{f_0} \pi\pi = 329 \pm 96$ MeV/c ² assuming $g_{f_0} K\bar{K} / g_{f_0} \pi\pi = 2$.				NODE=M003M1;LINKAGE=BO
		13 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0} \pi\pi = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.				NODE=M003M1;LINKAGE=MO
		14 In the kaon-loop fit following formalism of ACHASOV 89.				NODE=M003M1;LINKAGE=AB
		15 In the no-structure fit assuming a direct coupling of ϕ to $f_0\gamma$.				NODE=M003M1;LINKAGE=AM
		16 FLATTE 76 parameterization. Supersedes GARMASH 05.				NODE=M003M1;LINKAGE=GR
		17 FLATTE 76 parameterization, $g_{f_0} K\bar{K} / g_{f_0} \pi\pi = 4.21 \pm 0.25 \pm 0.21$.				NODE=M003M1;LINKAGE=AL
		18 K-matrix pole from combined analysis of $\pi^-p \rightarrow \pi^0\pi^0n, \pi^-p \rightarrow K\bar{K}n,$ $\pi^+\pi^- \rightarrow \pi^+\pi^-, \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta, \pi^+\pi^-\pi^0, K^+K^-\pi^0, K_S^0 K_S^0\pi^0,$ $K^+K_S^0\pi^-$ at rest, $\bar{p}n \rightarrow \pi^-\pi^-\pi^+, K_S^0 K^- \pi^0, K_S^0 K_S^0\pi^-$ at rest.				NODE=M003M;LINKAGE=KM
		19 From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution.				NODE=M003M1;LINKAGE=KD
		20 Coupled-channel Breit-Wigner, couplings $g_\pi = 0.09 \pm 0.01 \pm 0.01, g_K = 0.02 \pm 0.04 \pm 0.03$.				NODE=M003M;LINKAGE=TL
		21 Supersedes ACHASOV 98I. Using the model of ACHASOV 89.				NODE=M003M;LINKAGE=V9

- 22 Supersedes ACHASOV 98i.
- 23 In the "narrow resonance" approximation.
- 24 Assuming $\Gamma(f_0) = 40$ MeV.
- 25 From a narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.
- 26 From the combined fit of the photon spectra in the reactions $e^+e^- \rightarrow \pi^+\pi^-\gamma$, $\pi^0\pi^0\gamma$.
- 27 Supersedes BARBERIS 99 and BARBERIS 99B
- 28 T-matrix pole.
- 29 From invariant mass fit.
- 30 On sheet II in a 2 pole solution. The other pole is found on sheet III at (1039-93i) MeV.
- 31 On sheet II in a 2 pole solution. The other pole is found on sheet III at (963-29i) MeV.
- 32 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 33 At high $|t|$.
- 34 At low $|t|$.
- 35 On sheet II in a 4-pole solution, the other poles are found on sheet III at (953-55i) MeV and on sheet IV at (938-35i) MeV.
- 36 Combined fit of ALDE 95B, ANISOVICH 94, AMSLER 94D.
- 37 On sheet II in a 2 pole solution. The other pole is found on sheet III at (996-103i) MeV.
- 38 From sheet II pole position.
- 39 On sheet II in a 2 pole solution. The other pole is found on sheet III at (797-185i) MeV and can be interpreted as a shadow pole.
- 40 On sheet II in a 2 pole solution. The other pole is found on sheet III at (978-28i) MeV.
- 41 From coupled channel analysis.
- 42 Coupled channel analysis with finite width corrections.
- 43 Included in AGUILAR-BENITEZ 78 fit.

NODE=M003M;LINKAGE=V8
 NODE=M003M1;LINKAGE=AI
 NODE=M003M;LINKAGE=SM
 NODE=M003M;LINKAGE=ST
 NODE=M003M;LINKAGE=SL
 NODE=M003M1;LINKAGE=BD
 NODE=M003M1;LINKAGE=AN
 NODE=M003M1;LINKAGE=A
 NODE=M003M1;LINKAGE=LC
 NODE=M003M1;LINKAGE=X
 NODE=M003M1;LINKAGE=AA
 NODE=M003M1;LINKAGE=LA
 NODE=M003M1;LINKAGE=LB
 NODE=M003M1;LINKAGE=KL
 NODE=M003M1;LINKAGE=CF
 NODE=M003M1;LINKAGE=C2
 NODE=M003M1;LINKAGE=KM
 NODE=M003M1;LINKAGE=L
 NODE=M003M1;LINKAGE=K
 NODE=M003M1;LINKAGE=B
 NODE=M003M;LINKAGE=B
 NODE=M003M;LINKAGE=R

$f_0(980)$ WIDTH

Width determination very model dependent. Peak width in $\pi\pi$ is about 50 MeV, but decay width can be much larger.

NODE=M003W1
 NODE=M003W1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

10 to 100 OUR ESTIMATE
 [40 to 100 MeV OUR 2015 ESTIMATE]

NODE=M003W1
 NEW;→ UNCHECKED ←

••• We do not use the following data for averages, fits, limits, etc. •••

15.3 ± 4.7	424	ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
9.5 ± 1.1	706	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma 3\pi$
42 +20 -16		1,2 GARCIA-MAR..11		RVUE	Compilation
50 +20 -12		2,3 GARCIA-MAR..11		RVUE	Compilation
48 +22 -6		4 MOUSSALLAM11		RVUE	Compilation
36 ± 22		5 MENNESSIER 10		RVUE	Compilation
70 +20 -32		6 ANISOVICH 09		RVUE	0.0 $\bar{p}p$, πN
91 +30 ± 3 -22	44	7 ECKLUND	09	CLEO	4.17 $e^+e^- \rightarrow D_s^- D_s^{*+} + c.c.$
66.9 ± 2.2 +17.6 -12.5		8 UEHARA	08A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
65 ± 13	262 ± 30	9 AUBERT	07AK	BABR	10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
81 ± 21	54 ± 9	9 AUBERT	07AK	BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\pi^0\gamma$
51.3 +20.8+13.2 -17.7-3.8		10 MORI	07	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
61 ± 9 +14 -8	2584	11 GARMASH	05	BELL	$B^+ \rightarrow K^+\pi^+\pi^-$
64 ± 16		12 ANISOVICH	03	RVUE	
121 ± 23		TIKHOMIROV	03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
~ 70		13 BRAMON	02	RVUE	1.02 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
44 ± 2 ± 2	848	14 AITALA	01A	E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$
201 ± 28	419	15 ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

OCCUR=2

OCCUR=2

122 ± 13	419	16,17	ACHASOV	00H	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
56 ± 20		18	AKHMETSHIN	99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
65 ± 20			BARBERIS	99	OMEG	450 $p p \rightarrow p_s p_f K^+ K^-$	
80 ± 10			BARBERIS	99B	OMEG	450 $p p \rightarrow p_s p_f \pi^+ \pi^-$	
80 ± 10			BARBERIS	99C	OMEG	450 $p p \rightarrow p_s p_f \pi^0 \pi^0$	
48 ± 12 ± 8		19	BARBERIS	99D	OMEG	450 $p p \rightarrow K^+ K^-, \pi^+ \pi^-$	
65 ± 25			BELLAZZINI	99	GAM4	450 $p p \rightarrow p p \pi^0 \pi^0$	
71 ± 14		20	KAMINSKI	99	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, \sigma \sigma$	
~ 28		20	OLLER	99	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$	
~ 25			OLLER	99B	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$	
~ 14		20	OLLER	99C	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, \eta \eta$	
70 ± 20			ALDE	98	GAM4		
86 ± 16		20	ANISOVICH	98B	RVUE	Compilation	
54		21	LOCHER	98	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$	
69 ± 15		22	ALDE	97	GAM2	450 $p p \rightarrow p p \pi^0 \pi^0$	
38 ± 20		23	BERTIN	97C	OBLX	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$	
~ 100		24	ISHIDA	96	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$	
34			TORNQVIST	96	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, K \pi, \eta \pi$	
48 ± 10	3k	25	ALDE	95B	GAM2	38 $\pi^- p \rightarrow \pi^0 \pi^0 n$	
95 ± 20	10k	26	ALDE	95B	GAM2	38 $\pi^- p \rightarrow \pi^0 \pi^0 n$	OCCUR=2
26 ± 10			AMSLER	95B	CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$	
~ 112		27	AMSLER	95D	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$	
80 ± 12		28	ANISOVICH	95	RVUE		
30			JANSSEN	95	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$	
74		29	BUGG	94	RVUE	$\bar{p} p \rightarrow \eta 2\pi^0$	
29 ± 2		30	KAMINSKI	94	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}$	
46		31	ZOU	94B	RVUE		
48 ± 12		32	MORGAN	93	RVUE	$\pi \pi (K \bar{K}) \rightarrow \pi \pi (K \bar{K}), J/\psi \rightarrow \phi \pi \pi (K \bar{K}), D_s \rightarrow \pi (\pi \pi)$	
37.4 ± 10.6		22	AGUILAR-...	91	EHS	400 $p p$	
72 ± 8		33	ARMSTRONG	91	OMEG	300 $p p \rightarrow p p \pi \pi, p p K \bar{K}$	
110 ± 30			BREAKSTONE	90	SFM	$p p \rightarrow p p \pi^+ \pi^-$	
29 ± 13		22	ABACHI	86B	HRS	$e^+ e^- \rightarrow \pi^+ \pi^- X$	
120 ± 281 ± 20			ETKIN	82B	MPS	23 $\pi^- p \rightarrow n 2K_S^0$	
28 ± 10		33	GIDAL	81	MRK2	$J/\psi \rightarrow \pi^+ \pi^- X$	
70 to 300		34	ACHASOV	80	RVUE		
100 ± 80		35	AGUILAR-...	78	HBC	0.7 $\bar{p} p \rightarrow K_S^0 K_S^0$	
30 ± 8		33	LEEPER	77	ASPK	2-2.4 $\pi^- p \rightarrow \pi^+ \pi^- n, K^+ K^- n$	
48 ± 14		33	BINNIE	73	CNTR	$\pi^- p \rightarrow n M M$	
32 ± 10		36	GRAYER	73	ASPK	17 $\pi^- p \rightarrow \pi^+ \pi^- n$	
30 ± 10		36	HYAMS	73	ASPK	17 $\pi^- p \rightarrow \pi^+ \pi^- n$	
54 ± 16		36	PROTOPOP...	73	HBC	7 $\pi^+ p \rightarrow \pi^+ p \pi^+ \pi^-$	

¹ Analytic continuation using Roy equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi \pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.

² Quoted number refers to twice imaginary part of pole position.

³ Analytic continuation using GKPY equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi \pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.

⁴ Pole position. Used Roy equations.

⁵ Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.

⁶ On sheet II in a 2-pole solution. The other pole is found on sheet III at (850-100i) MeV

⁷ Using a relativistic Breit-Wigner function and taking into account the finite D_s mass.

⁸ Breit-Wigner $\pi \pi$ width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0} \pi \pi = 0$.

⁹ Systematic errors not estimated.

¹⁰ Breit-Wigner $\pi \pi$ width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0} \pi \pi = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.

NODE=M003W1;LINKAGE=GC

NODE=M003W1;LINKAGE=GI

NODE=M003W1;LINKAGE=GM

NODE=M003W1;LINKAGE=MU

NODE=M003W1;LINKAGE=ME

NODE=M003W1;LINKAGE=AO

NODE=M003W1;LINKAGE=EC

NODE=M003W1;LINKAGE=UE

NODE=M003W1;LINKAGE=NS

NODE=M003W1;LINKAGE=MO

- 11 Breit-Wigner, solution 1, PWA ambiguous.
- 12 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.
- 13 Using the data of AKHMETSHIN 99C, ACHASOV 00H, and ALOISIO 02D.
- 14 Breit-Wigner width.
- 15 Supersedes ACHASOV 98I. Using the model of ACHASOV 89.
- 16 Supersedes ACHASOV 98I.
- 17 In the "narrow resonance" approximation.
- 18 From the combined fit of the photon spectra in the reactions $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$, $\pi^0 \pi^0 \gamma$.
- 19 Supersedes BARBERIS 99 and BARBERIS 99B
- 20 T-matrix pole.
- 21 On sheet II in a 2 pole solution. The other pole is found on sheet III at (1039–93i) MeV.
- 22 From invariant mass fit.
- 23 On sheet II in a 2 pole solution. The other pole is found on sheet III at (963–29i) MeV.
- 24 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 25 At high $|t|$.
- 26 At low $|t|$.
- 27 On sheet II in a 4-pole solution, the other poles are found on sheet III at (953–55i) MeV and on sheet IV at (938–35i) MeV.
- 28 Combined fit of ALDE 95B, ANISOVICH 94,
- 29 On sheet II in a 2 pole solution. The other pole is found on sheet III at (996–103i) MeV.
- 30 From sheet II pole position.
- 31 On sheet II in a 2 pole solution. The other pole is found on sheet III at (797–185i) MeV and can be interpreted as a shadow pole.
- 32 On sheet II in a 2 pole solution. The other pole is found on sheet III at (978–28i) MeV.
- 33 From coupled channel analysis.
- 34 Coupled channel analysis with finite width corrections.
- 35 From coupled channel fit to the HYAMS 73 and PROTOPODESCU 73 data. With a simultaneous fit to the $\pi\pi$ phase-shifts, inelasticity and to the $K_S^0 K_S^0$ invariant mass.
- 36 Included in AGUILAR-BENITEZ 78 fit.

NODE=M003W1;LINKAGE=GA
 NODE=M003W;LINKAGE=KM

NODE=M003W;LINKAGE=BR
 NODE=M003W;LINKAGE=TL
 NODE=M003W;LINKAGE=V9
 NODE=M003W;LINKAGE=V8
 NODE=M003W1;LINKAGE=AI
 NODE=M003W;LINKAGE=SL

NODE=M003W1;LINKAGE=BD
 NODE=M003W1;LINKAGE=AN
 NODE=M003W1;LINKAGE=LO
 NODE=M003W1;LINKAGE=A
 NODE=M003W1;LINKAGE=X
 NODE=M003W1;LINKAGE=AA

NODE=M003W1;LINKAGE=LA
 NODE=M003W1;LINKAGE=LB
 NODE=M003W1;LINKAGE=KL

NODE=M003W1;LINKAGE=CF
 NODE=M003W1;LINKAGE=C2
 NODE=M003W1;LINKAGE=KM
 NODE=M003W1;LINKAGE=L

NODE=M003W1;LINKAGE=K
 NODE=M003W1;LINKAGE=B
 NODE=M003W;LINKAGE=B
 NODE=M003W;LINKAGE=C

NODE=M003W;LINKAGE=R

$f_0(980)$ DECAY MODES

NODE=M003215;NODE=M003

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	dominant
Γ_2 $K\bar{K}$	seen
Γ_3 $\gamma\gamma$	seen
Γ_4 $e^+ e^-$	

DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=5;OUR EVAL;→ UNCHECKED ←
 DESIG=4

$f_0(980)$ PARTIAL WIDTHS

NODE=M003220

$\Gamma(\gamma\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_3
------------------------	-------------	-------------	------	---------	------------

NODE=M003W4
 NODE=M003W4

0.31 $\begin{smallmatrix} +0.05 \\ -0.04 \end{smallmatrix}$ OUR AVERAGE

0.32 ± 0.05	1	DAI	14A	RVUE	Compilation
0.286 ± 0.017 $\begin{smallmatrix} +0.211 \\ -0.070 \end{smallmatrix}$	2	UEHARA	08A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
0.205 $\begin{smallmatrix} +0.095 +0.147 \\ -0.083 -0.117 \end{smallmatrix}$	3	MORI	07	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
0.42 ± 0.06 ± 0.18	4	OEST	90	JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.16 ± 0.01	5	MENNESSIER	11	RVUE	
0.29 ± 0.21 $\begin{smallmatrix} +0.02 \\ -0.07 \end{smallmatrix}$	6	MOUSSALLAM	11	RVUE	Compilation
0.42	7,8	PENNINGTON	08	RVUE	Compilation
0.10	8,9	PENNINGTON	08	RVUE	Compilation
0.28 $\begin{smallmatrix} +0.09 \\ -0.13 \end{smallmatrix}$	10	BOGLIONE	99	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
0.29 ± 0.07 ± 0.12	11,12	BOYER	90	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
0.31 ± 0.14 ± 0.09	11,12	MARSISKE	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
0.63 ± 0.14	13	MORGAN	90	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$

OCCUR=2

- ¹ Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTAKER 04 as input.
- ² Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $\mathcal{E}f_0 K K / \mathcal{E}f_0 \pi \pi = 0$.
- ³ Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $\mathcal{E}f_0 K K / \mathcal{E}f_0 \pi \pi = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.
- ⁴ OEST 90 quote systematic errors $\begin{matrix} +0.08 \\ -0.18 \end{matrix}$. We use ± 0.18 . Observed 60 events.
- ⁵ Uses an analytic K-matrix model. Compilation.
- ⁶ Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.
- ⁷ Solution A (preferred solution based on χ^2 -analysis).
- ⁸ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.
- ⁹ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).
- ¹⁰ Supersedes MORGAN 90.
- ¹¹ From analysis allowing arbitrary background unconstrained by unitarity.
- ¹² Data included in MORGAN 90, BOGLIONE 99 analyses.
- ¹³ From amplitude analysis of BOYER 90 and MARSISKE 90, data corresponds to resonance parameters $m = 989$ MeV, $\Gamma = 61$ MeV.

NODE=M003W4;LINKAGE=D
 NODE=M003W4;LINKAGE=UE
 NODE=M003W4;LINKAGE=MO
 NODE=M003W4;LINKAGE=H
 NODE=M003W4;LINKAGE=ME
 NODE=M003W4;LINKAGE=MU
 NODE=M003W4;LINKAGE=P1
 NODE=M003W4;LINKAGE=P3
 NODE=M003W4;LINKAGE=P2
 NODE=M003W4;LINKAGE=BL
 NODE=M003W4;LINKAGE=B
 NODE=M003W4;LINKAGE=C
 NODE=M003W4;LINKAGE=A

 $\Gamma(e^+e^-)$ Γ_4

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<8.4	90	VOROBYEV	88 ND	$e^+e^- \rightarrow \pi^0\pi^0$

NODE=M003W3
 NODE=M003W3

 $f_0(980)$ BRANCHING RATIOS

$\Gamma(\pi\pi) / [\Gamma(\pi\pi) + \Gamma(K\bar{K})]$	$\Gamma_1 / (\Gamma_1 + \Gamma_2)$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.52 ± 0.12	9.9k	¹ AUBERT	06O BABR	$B^\pm \rightarrow K^\pm \pi^\pm \pi^\mp$
$0.75^{+0.11}_{-0.13}$		² ABLIKIM	05Q BES2	$\chi_{c0} \rightarrow 2\pi^+ 2\pi^-$, $\pi^+ \pi^- K^+ K^-$
0.84 ± 0.02		³ ANISOVICH	02D SPEC	Combined fit
~ 0.68		OLLER	99B RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
0.67 ± 0.09		⁴ LOVERRE	80 HBC	$4\pi^- p \rightarrow n2K_S^0$
$0.81^{+0.09}_{-0.04}$		⁴ CASON	78 STRC	$7\pi^- p \rightarrow n2K_S^0$
0.78 ± 0.03		⁴ WETZEL	76 OSPK	$8.9\pi^- p \rightarrow n2K_S^0$

¹ Recalculated by us using $\Gamma(K^+K^-) / \Gamma(\pi^+\pi^-) = 0.69 \pm 0.32$ from AUBERT 06O and isospin relations.

² Using data from ABLIKIM 04G.

³ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data.

⁴ Measure $\pi\pi$ elasticity assuming two resonances coupled to the $\pi\pi$ and $K\bar{K}$ channels only.

NODE=M003225

NODE=M003R1
 NODE=M003R1

NODE=M003R1;LINKAGE=AU

NODE=M003R1;LINKAGE=AB
 NODE=M003R;LINKAGE=CH

NODE=M003R1;LINKAGE=B

 $f_0(980)$ REFERENCES

ABLIKIM 15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56781
DAI 14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)	REFID=55923
ABLIKIM 12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54270
GARCIA-MAR... 11	PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=16761
GARCIA-MAR... 11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=54121
MENNESSIER 11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang		REFID=53637
MOUSSALLAM 11	EPJ C71 1814	B. Moussallam		REFID=53975
BATLEY 10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	REFID=53567
MENNESSIER 10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang		REFID=53657
ANISOVICH 09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(CLEO Collab.)	REFID=52719
ECKLUND 09	PR D80 052009	K.M. Ecklund <i>et al.</i>		REFID=53041
BATLEY 08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	REFID=52487
PENNINGTON 08	EPJ C56 1	M.R. Pennington <i>et al.</i>		REFID=52303
UEHARA 08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
AMBROSINO 07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51616
AUBERT 07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
BONVICINI 07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=51721
MORI 07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51652
AMBROSINO 06B	PL B634 148	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51043
AUBERT 06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51141
GARMASH 06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51162
ABLIKIM 05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM 05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ACHASOV 05	PR D72 013006	N.N. Achasov, G.N. Shestakov		REFID=50762
GARMASH 05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=50641
ABLIKIM 04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50187
BUETTAKER 04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam		REFID=56428; ERROR=2
ANISOVICH 03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
TIKHOMIROV 03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423

Translated from YAF 66 860.

ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
ANISOVICH	02D	PAN 65 1545 Translated from YAF 65 1583.	V.V. Anisovich <i>et al.</i>		REFID=48831
BRAMON	02	EPJ C26 253	A. Bramon <i>et al.</i>		REFID=49178
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)	REFID=48312
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47393
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99C	PL B453 325	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46923
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>		REFID=46600
ACKERSTAFF	98Q	EPJ C4 19	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46145
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
ANISOVICH	98B	SPU 41 419 Translated from YAF 62 446. Translated from UFN 168 481.	V.V. Anisovich <i>et al.</i>		REFID=46331
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)	REFID=46372
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ISHIDA	96	PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45770
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=44375
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)	REFID=44442
JANSSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)	REFID=44072
MORGAN	93	PR D48 1185	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=43614
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273 Translated from YAF 48 436.	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)	REFID=20394
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ACHASOV	80	SJNP 32 566 Translated from YAF 32 1098.	N.N. Achasov, S.A. Devyanin, G.N. Shestakov	(NOVM)	REFID=20458
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP	REFID=20381
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+) IJP	REFID=20382
AGUILAR-...	78	NP B140 73	M. Aguilar-Benitez <i>et al.</i>	(MADR, BOMB+)	REFID=20368
CASON	78	PRL 41 271	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20370
LEEPER	77	PR D16 2054	R.J. Leeper <i>et al.</i>	(ISU)	REFID=20365
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)	REFID=11004
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)	REFID=20446
WETZEL	76	NP B115 208	W. Wetzel <i>et al.</i>	(ETH, CERN, LOIC)	REFID=20362
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)	REFID=21062
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
BINNIE	73	PRL 31 1534	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20343
GRAYER	73	Tallahassee	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20347
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108

$a_0(980)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

See our minireview on scalar mesons under $f_0(500)$. (See the index for the page number.)

NODE=M036

NODE=M036

 $a_0(980)$ MASS

NODE=M036205

VALUE (MeV)

DOCUMENT ID

980±20 OUR ESTIMATE Mass determination very model dependent

NODE=M036MX

→ UNCHECKED ←

 $\eta\pi$ FINAL STATE ONLY

NODE=M036M1

NODE=M036M1

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

CHG

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

982.5 ± 1.6 ± 1.1	16.9k	¹ AMBROSINO	09F	KLOE	1.02 $e^+e^- \rightarrow \eta\pi^0\gamma$
986 ± 4		ANISOVICH	09	RVUE	0.0 $\bar{p}p, \pi N$
982.3 + 0.6 + 3.1 - 0.7 - 4.7		² UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
987.4 ± 1.0 ± 3.0		^{3,4} BUGG	08A	RVUE 0	$\bar{p}p \rightarrow \pi^0\pi^0\eta$
989.1 ± 1.0 ± 3.0		^{4,5} BUGG	08A	RVUE 0	$\bar{p}p \rightarrow \pi^0\pi^0\eta$
985 ± 4 ± 6	318	ACHARD	02B	L3	183-209 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
995 + 52 - 10	36	⁶ ACHASOV	00F	SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
994 + 33 - 8	36	⁷ ACHASOV	00F	SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
975 ± 7		BARBERIS	00H		450 $p\rho \rightarrow p_f\eta\pi^0\rho_s$
988 ± 8		BARBERIS	00H		450 $p\rho \rightarrow \Delta_f^{++}\eta\pi^-\rho_s$
~ 1055		⁸ OLLER	99	RVUE	$\eta\pi, K\bar{K}$
~ 1009.2		⁸ OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
993.1 ± 2.1		⁹ TEIGE	99	B852	18.3 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
988 ± 6		⁸ ANISOVICH	98B	RVUE	Compilation
987		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
991		JANSSEN	95	RVUE	$\eta\pi \rightarrow \eta\pi, K\bar{K}, K\pi, \eta\pi$
984.45 ± 1.23 ± 0.34		AMSLER	94C	CBAR	0.0 $\bar{p}p \rightarrow \omega\eta\pi^0$
982 ± 2		¹⁰ AMSLER	92	CBAR	0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
984 ± 4	1040	¹⁰ ARMSTRONG	91B	OMEG ±	300 $p\rho \rightarrow p\rho\eta\pi^+\pi^-$
976 ± 6		ATKINSON	84E	OMEG ±	25-55 $\gamma p \rightarrow \eta\pi n$
986 ± 3	500	¹¹ EVANGELIS...	81	OMEG ±	12 $\pi^-p \rightarrow \eta\pi^+\pi^-\pi^-p$
990 ± 7	145	¹¹ GURTU	79	HBC ±	4.2 $K^-p \rightarrow \Lambda\eta 2\pi$
980 ± 11	47	CONFORTO	78	OSPK -	4.5 $\pi^-p \rightarrow pX^-$
978 ± 16	50	CORDEN	78	OMEG ±	12-15 $\pi^-p \rightarrow n\eta 2\pi$
977 ± 7		GRASSLER	77	HBC -	16 $\pi^\mp p \rightarrow p\eta 3\pi$
989 ± 4	70	WELLS	75	HBC -	3.1-6 $K^-p \rightarrow \Lambda\eta 2\pi$
972 ± 10	150	DEFOIX	72	HBC ±	0.7 $\bar{p}p \rightarrow 7\pi$
970 ± 15	20	BARNES	69C	HBC -	4-5 $K^-p \rightarrow \Lambda\eta 2\pi$
980 ± 10		CAMPBELL	69	DBC ±	2.7 π^+d
980 ± 10	15	MILLER	69B	HBC -	4.5 $K^-N \rightarrow \eta\pi\Lambda$
980 ± 10	30	AMMAR	68	HBC ±	5.5 $K^-p \rightarrow \Lambda\eta 2\pi$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

¹ Using the model of ACHASOV 89 and ACHASOV 03B.² From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.³ Parameterizes couplings to $\bar{K}K, \pi\eta,$ and $\pi\eta'$.⁴ Using AMSLER 94D and ABELE 98.⁵ From the T-matrix pole on sheet II.⁶ Using the model of ACHASOV 89. Supersedes ACHASOV 98B.⁷ Using the model of JAFFE 77. Supersedes ACHASOV 98B.⁸ T-matrix pole.⁹ Breit-Wigner fit, average between a_0^\pm and a_0^0 . The fit favors a slightly heavier a_0^\pm .¹⁰ From a single Breit-Wigner fit.¹¹ From $f_1(1285)$ decay.

NODE=M036M1;LINKAGE=AM

NODE=M036M1;LINKAGE=UE

NODE=M036M1;LINKAGE=BP

NODE=M036M1;LINKAGE=BU

NODE=M036M1;LINKAGE=BT

NODE=M036M1;LINKAGE=V1

NODE=M036M1;LINKAGE=M2

NODE=M036M1;LINKAGE=AN

NODE=M036M1;LINKAGE=BF

NODE=M036M1;LINKAGE=A

NODE=M036M1;LINKAGE=R

K \bar{K} ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
~ 1053		¹² OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
982 ± 3		¹³ ABELE	98	CBAR	$0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
975 ± 15		BERTIN	98B	OBLX ±	$0.0 \bar{p}p \rightarrow K^\pm K_S \pi^\mp$
976 ± 6	316	DEBILLY	80	HBC ±	$1.2-2 \bar{p}p \rightarrow f_1(1285)\omega$
1016 ± 10	100	¹⁴ ASTIER	67	HBC ±	$0.0 \bar{p}p$
1003.3 ± 7.0	143	¹⁵ ROSENFELD	65	RVUE ±	

¹² T-matrix pole.

¹³ T-matrix pole on sheet II, the pole on sheet III is at 1006-i49 MeV.

¹⁴ ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.

¹⁵ Plus systematic errors.

NODE=M036M2
NODE=M036M2

NODE=M036M2;LINKAGE=AN
NODE=M036M2;LINKAGE=Q
NODE=M036M2;LINKAGE=A
NODE=M036M2;LINKAGE=S

 $a_0(980)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
50 to 100 OUR ESTIMATE		Width determination very model dependent. Peak width in $\eta\pi$ is about 60 MeV, but decay width can be much larger.			

• • • We do not use the following data for averages, fits, limits, etc. • • •

75.6 ± 1.6	^{+17.4} ^{-10.0}	¹⁶ UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
80.2 ± 3.8	± 5.4	¹⁷ BUGG	08A	RVUE 0	$\bar{p}p \rightarrow \pi^0\pi^0\eta$
50 ± 13	± 4	318 ACHARD	02B	L3	$183-209 e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
72 ± 16		BARBERIS	00H		$450 pp \rightarrow p_f\eta\pi^0 p_s$
61 ± 19		BARBERIS	00H		$450 pp \rightarrow \Delta_f^{++}\eta\pi^- p_s$
~ 42		¹⁸ OLLER	99	RVUE	$\eta\pi, K\bar{K}$
~ 112		¹⁸ OLLER	99B	RVUE	$\pi\pi \rightarrow \eta\pi, K\bar{K}$
71 ± 7		TEIGE	99	B852	$18.3 \pi^- p \rightarrow \eta\pi^+\pi^- n$
92 ± 20		¹⁸ ANISOVICH	98B	RVUE	Compilation
65 ± 10		¹⁹ BERTIN	98B	OBLX ±	$0.0 \bar{p}p \rightarrow K^\pm K_S \pi^\mp$
~ 100		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
202		JANSSEN	95	RVUE	$\eta\pi \rightarrow \eta\pi, K\bar{K}, K\pi,$ $\eta\pi$
54.12 ± 0.34 ± 0.12		AMSLER	94C	CBAR	$0.0 \bar{p}p \rightarrow \omega\eta\pi^0$
54 ± 10		²⁰ AMSLER	92	CBAR	$0.0 \bar{p}p \rightarrow \eta\eta\pi^0$
95 ± 14	1040	²⁰ ARMSTRONG	91B	OMEG ±	$300 pp \rightarrow p\rho\eta\pi^+\pi^-$
62 ± 15	500	²¹ EVANGELIS...	81	OMEG ±	$12 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
60 ± 20	145	²¹ GURTU	79	HBC ±	$4.2 K^- p \rightarrow \Lambda\eta 2\pi$
60 ⁺⁵⁰ ⁻³⁰	47	CONFORTO	78	OSPK -	$4.5 \pi^- p \rightarrow pX^-$
86.0 ^{+60.0} ^{-50.0}	50	CORDEN	78	OMEG ±	$12-15 \pi^- p \rightarrow n\eta 2\pi$
44 ± 22		GRASSLER	77	HBC -	$16 \pi^\mp p \rightarrow p\eta 3\pi$
80 to 300		²² FLATTE	76	RVUE -	$4.2 K^- p \rightarrow \Lambda\eta 2\pi$
16.0 ^{+25.0} ^{-16.0}	70	WELLS	75	HBC -	$3.1-6 K^- p \rightarrow \Lambda\eta 2\pi$
30 ± 5	150	DEFOIX	72	HBC ±	$0.7 \bar{p}p \rightarrow 7\pi$
40 ± 15		CAMPBELL	69	DBC ±	$2.7 \pi^+ d$
60 ± 30	15	MILLER	69B	HBC -	$4.5 K^- N \rightarrow \eta\pi\Lambda$
80 ± 30	30	AMMAR	68	HBC ±	$5.5 K^- p \rightarrow \Lambda\eta 2\pi$

¹⁶ From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

¹⁷ From the T-matrix pole on sheet II, using AMSLER 94D and ABELE 98.

¹⁸ T-matrix pole.

¹⁹ The $\eta\pi$ width.

²⁰ From a single Breit-Wigner fit.

²¹ From $f_1(1285)$ decay.

²² Using a two-channel resonance parametrization of GAY 76B data.

NODE=M036210

NODE=M036W1

→ UNCHECKED ←

OCCUR=2

NODE=M036W1;LINKAGE=UE

NODE=M036W1;LINKAGE=BU

NODE=M036W1;LINKAGE=AN

NODE=M036W1;LINKAGE=BE

NODE=M036W1;LINKAGE=A

NODE=M036W1;LINKAGE=R

NODE=M036W1;LINKAGE=F

$K\bar{K}$ ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
92 ± 8		²³ ABELE	98	CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 24		²⁴ OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
~ 25	100	²⁵ ASTIER	67	HBC ±	
57 ± 13	143	²⁶ ROSENFELD	65	RVUE ±	

²³ T-matrix pole on sheet II, the pole on sheet III is at 1006-i49 MeV.

²⁴ T-matrix pole.

²⁵ ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.

²⁶ Plus systematic errors.

NODE=M036W2
NODE=M036W2

NODE=M036W2;LINKAGE=Q
NODE=M036W2;LINKAGE=AN
NODE=M036W2;LINKAGE=A
NODE=M036W2;LINKAGE=S

 $a_0(980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi$	dominant
Γ_2 $K\bar{K}$	seen
Γ_3 $\rho\pi$	
Γ_4 $\gamma\gamma$	seen
Γ_5 e^+e^-	

NODE=M036215;NODE=M036

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=2
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6

 $a_0(980)$ PARTIAL WIDTHS **$\Gamma(\gamma\gamma)$**

VALUE (keV)	DOCUMENT ID	TECN
-------------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.30 ± 0.10 ²⁷ AMSLER 98 RVUE

²⁷ Using $\Gamma_{\gamma\gamma} B(a_0(980) \rightarrow \eta\pi) = 0.24 \pm 0.08$ keV.

 Γ_4

NODE=M036217

NODE=M036W4
NODE=M036W4

NODE=M036W4;LINKAGE=A

 $a_0(980)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ **$\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_4/\Gamma$
-------------	------	-------------	------	---------	---------------------------

0.21 $\begin{smallmatrix} +0.08 \\ -0.04 \end{smallmatrix}$ OUR AVERAGE

0.128 $\begin{smallmatrix} +0.003 +0.502 \\ -0.002 -0.043 \end{smallmatrix}$		²⁸ UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$	
0.28 ± 0.04 ± 0.10	44	OEST	90	JADE	$e^+e^- \rightarrow e^+e^-\pi^0\eta$	
0.19 ± 0.07 $\begin{smallmatrix} +0.10 \\ -0.07 \end{smallmatrix}$		ANTREASYAN	86	CBAL	$e^+e^- \rightarrow e^+e^-\pi^0\eta$	

²⁸ From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

NODE=M036220

NODE=M036G1
NODE=M036G1

NODE=M036G1;LINKAGE=UE

 $\Gamma(\eta\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_5/\Gamma$
------------	-----	-------------	------	---------	---------------------------

<1.5 90 VOROBYEV 88 ND $e^+e^- \rightarrow \pi^0\eta$

NODE=M036G2
NODE=M036G2

 $a_0(980)$ BRANCHING RATIOS **$\Gamma(K\bar{K})/\Gamma(\eta\pi)$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
-------	-------------	------	-----	---------	---------------------

0.183 ± 0.024 OUR AVERAGE Error includes scale factor of 1.2.

0.57 ± 0.16	²⁹ BARGIOTTI	03	OBLX	$\bar{p}p$	
0.23 ± 0.05	³⁰ ABELE	98	CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$	
0.166 ± 0.01 ± 0.02	³¹ BARBERIS	98C	OMEG	450 $pp \rightarrow p_f f_1(1285) p_s$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.20 ± 0.15	³² ANISOVICH	09	RVUE	0.0 $\bar{p}p, \pi N$	
1.05 ± 0.07 ± 0.05	³³ BUGG	08A	RVUE	0 $\bar{p}p \rightarrow \pi^0\pi^0\eta$	
~ 0.60	OLLER	99B	RVUE	$\pi\pi \rightarrow \eta\pi, K\bar{K}$	
0.7 ± 0.3	³¹ CORDEN	78	OMEG	12-15 $\pi^- p \rightarrow n\eta 2\pi$	
0.25 ± 0.08	³¹ DEFOIX	72	HBC ±	0.7 $\bar{p} \rightarrow 7\pi$	

NODE=M036225

NODE=M036R2
NODE=M036R2

$\Gamma(\rho\pi)/\Gamma(\eta\pi)$ Γ_3/Γ_1 $\rho\pi$ forbidden.

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.25	70	AMMAR	70	HBC	\pm	4.1,5.5 $K^- p \rightarrow \Lambda \eta 2\pi$
-------	----	-------	----	-----	-------	---

29 Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

30 Using $\pi^0 \pi^0 \eta$ from AMSLER 94D.

31 From the decay of $f_1(1285)$.

32 This is a ratio of couplings.

33 A ratio of couplings, using AMSLER 94D and ABELE 98. Supersedes BUGG 94.

NODE=M036R1

NODE=M036R1

NODE=M036R1

NODE=M036R;LINKAGE=BG

NODE=M036R2;LINKAGE=Q

NODE=M036R2;LINKAGE=L

NODE=M036R2;LINKAGE=AN

NODE=M036R2;LINKAGE=BU

 $a_0(980)$ REFERENCES

NODE=M036

AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=53105
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
BUGG	08A	PR D78 074023	D.V. Bugg	(LOQM)	REFID=52578
ACHASOV	03B	PR D68 014006	N.N. Achasov, A.V. Kiselev		REFID=49476
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48574
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
TEIGE	99	PR D59 012001	S. Teige <i>et al.</i>	(BNL E852 Collab.)	REFID=46613
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45863
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46317
AMSLER	98	RMP 70 1293	C. Amsler		REFID=46601
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20624
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
DEBILLY	80	NP B176 1	L. de Billy <i>et al.</i>	(CURIN, LAUS, NEUC+)	REFID=20461
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)	REFID=20456
CONFORTO	78	LNC 23 419	B. Conforto <i>et al.</i>	(RHEL, TINTO, CHIC+)	REFID=20451
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20452
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20447
JAFFE	77	PR D15 267,281	R. Jaffe	(MIT)	REFID=43673
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)	REFID=20446
GAY	76B	PL 63B 220	J.B. Gay <i>et al.</i>	(CERN, AMST, NIJM) JP	REFID=20445
WELLS	75	NP B101 333	J. Wells <i>et al.</i>	(OXF)	REFID=20444
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
AMMAR	70	PR D2 430	R. Ammar <i>et al.</i>	(KANS, NWES, ANL, WISC)	REFID=20428
BARNES	69C	PRL 23 610	V.E. Barnes <i>et al.</i>	(BNL, SYRA)	REFID=20418
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)	REFID=20419
MILLER	69B	PL 29B 255	D.H. Miller <i>et al.</i>	(PURD)	REFID=20424
		Also	W.L. Yen <i>et al.</i>	(PURD)	REFID=20425
AMMAR	68	PRL 21 1832	R. Ammar <i>et al.</i>	(NWES, ANL)	REFID=20412
ASTIER	67	PL 25B 294	A. Astier <i>et al.</i>	(CDEF, CERN, IRAD)	REFID=20405
		Includes data of BARLOW 67, CONFORTO 67, and ARMENTEROS 65.			
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
CONFORTO	67	NP B3 469	G. Conforto <i>et al.</i>	(CERN, CDEF, IPNP+)	REFID=20411
ARMENTEROS	65	PL 17 344	R. Armenteros <i>et al.</i>	(CERN, CDEF)	REFID=20396
ROSENFELD	65	Oxford Conf. 58	A.H. Rosenfeld	(LRL)	REFID=20399

$\phi(1020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M004

 $\phi(1020)$ MASS

NODE=M004M

NODE=M004M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1019.461±0.019 OUR AVERAGE				
Error includes scale factor of 1.1.				
1019.51 ±0.02 ±0.05		¹ LEES 13Q	BABR	$e^+e^- \rightarrow K^+K^-\gamma$
1019.30 ±0.02 ±0.10	105k	AKHMETSHIN 06	CMD2	$0.98-1.06 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.52 ±0.05 ±0.05	17.4k	AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta\gamma$
1019.483±0.011±0.025	272k	² AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
1019.42 ±0.05	1900k	³ ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$
1019.40 ±0.04 ±0.05	23k	AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1019.36 ±0.12		⁴ ACHASOV 00B	SND	$e^+e^- \rightarrow \eta\gamma$
1019.38 ±0.07 ±0.08	2200	⁵ AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\geq 2\gamma$
1019.51 ±0.07 ±0.10	11169	AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.5 ±0.4		BARBERIS 98	OMEG	$450 pp \rightarrow pp2K^+2K^-$
1019.42 ±0.06	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \text{hadrons}$
1019.7 ±0.3	2012	DAVENPORT 86	MPSF	$400 pA \rightarrow 4KX$
1019.7 ±0.1 ±0.1	5079	ALBRECHT 85D	ARG	$10 e^+e^- \rightarrow K^+K^-X$
1019.3 ±0.1	1500	ARENTON 82	AEMS	$11.8 \text{ polar. } pp \rightarrow KK$
1019.67 ±0.17	25080	⁶ PELLINEN 82	RVUE	
1019.52 ±0.13	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1019.48 ±0.01		LEES 13F	BABR	$D^+ \rightarrow K^+K^-\pi^+$
1019.441±0.008±0.080	542k	⁷ AKHMETSHIN 08	CMD2	$1.02 e^+e^- \rightarrow K^+K^-$
1019.63 ±0.07	12540	⁸ AUBERT,B 05J	BABR	$D^0 \rightarrow \bar{K}^0 K^+K^-$
1019.8 ±0.7		ARMSTRONG 86	OMEG	$85 \pi^+/pp \rightarrow \pi^+/p4Kp$
1020.1 ±0.11	5526	⁸ ATKINSON 86	OMEG	$20-70 \gamma p$
1019.7 ±1.0		BEBEK 86	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
1019.411±0.008	642k	⁹ DIJKSTRA 86	SPEC	$100-200 \pi^\pm, \bar{p}, p, K^\pm, \text{ on Be}$
1020.9 ±0.2		⁸ FRAME 86	OMEG	$13 K^+p \rightarrow \phi K^+p$
1021.0 ±0.2		⁸ ARMSTRONG 83B	OMEG	$18.5 K^-p \rightarrow K^-K^+\Lambda$
1020.0 ±0.5		⁸ ARMSTRONG 83B	OMEG	$18.5 K^-p \rightarrow K^-K^+\Lambda$
1019.7 ±0.3		⁸ BARATE 83	GOLI	$190 \pi^-\text{Be} \rightarrow 2\mu X$
1019.8 ±0.2 ±0.5	766	IVANOV 81	OLYA	$1-1.4 e^+e^- \rightarrow K^+K^-$
1019.4 ±0.5	337	COOPER 78B	HBC	$0.7-0.8 \bar{p}p \rightarrow K_S^0 K_L^0 \pi^+\pi^-$
1020 ±1	383	⁸ BALDI 77	CNTR	$10 \pi^-p \rightarrow \pi^- \phi p$
1018.9 ±0.6	800	COHEN 77	ASPK	$6 \pi^\pm N \rightarrow K^+K^-N$
1019.7 ±0.5	454	KALBFLEISCH 76	HBC	$2.18 K^-p \rightarrow \Lambda K \bar{K}$
1019.4 ±0.8	984	BESCH 74	CNTR	$2 \gamma p \rightarrow pK^+K^-$
1020.3 ±0.4	100	BALLAM 73	HBC	$2.8-9.3 \gamma p$
1019.4 ±0.7		BINNIE 73B	CNTR	$\pi^-p \rightarrow \phi n$
1019.6 ±0.5	120	¹⁰ AGUILAR-... 72B	HBC	$3.9,4.6 K^-p \rightarrow \Lambda K^+K^-$
1019.9 ±0.5	100	¹⁰ AGUILAR-... 72B	HBC	$3.9,4.6 K^-p \rightarrow K^-pK^+K^-$
1020.4 ±0.5	131	COLLEY 72	HBC	$10 K^+p \rightarrow K^+p\phi$
1019.9 ±0.3	410	STOTTLE... 71	HBC	$2.9 K^-p \rightarrow \Sigma/\Lambda K \bar{K}$

OCCUR=2

OCCUR=2

OCCUR=2

- ¹ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.
- ² Update of AKHMETSHIN 99D
- ³ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.
- ⁴ Using a total width of 4.43 ± 0.05 MeV. Systematic uncertainty included.
- ⁵ Using a total width of 4.43 ± 0.05 MeV.
- ⁶ PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DE-GROOT 74.
- ⁷ Strongly correlated with AKHMETSHIN 04.
- ⁸ Systematic errors not evaluated.
- ⁹ Weighted and scaled average of 12 measurements of DIJKSTRA 86.
- ¹⁰ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

NODE=M004M;LINKAGE=C

NODE=M004M;LINKAGE=GS
NODE=M004M;LINKAGE=AENODE=M004M;LINKAGE=G2
NODE=M004M;LINKAGE=F2
NODE=M004M;LINKAGE=RNODE=M004M;LINKAGE=AH
NODE=M004M;LINKAGE=A
NODE=M004M;LINKAGE=B
NODE=M004M;LINKAGE=D **$\phi(1020)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.266 ± 0.031 OUR AVERAGE		Error includes scale factor of 1.2.		
4.29 ± 0.04 ± 0.07		¹ LEES	13Q BABR	$e^+e^- \rightarrow K^+K^-\gamma$
4.30 ± 0.06 ± 0.17	105k	AKHMETSHIN 06	CMD2	$0.98-1.06 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.280 ± 0.033 ± 0.025	272k	² AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
4.21 ± 0.04	1900k	³ ACHASOV	01E SND	$e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$
4.44 ± 0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow$ hadrons
4.5 ± 0.7	1500	ARENTON 82	AEMS	11.8 polar. $pp \rightarrow KK$
4.2 ± 0.6	766	⁴ IVANOV	81 OLYA	$1-1.4 e^+e^- \rightarrow K^+K^-$
4.3 ± 0.6		⁴ CORDIER	80 DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.36 ± 0.29	3681	⁴ BUKIN	78C OLYA	$e^+e^- \rightarrow$ hadrons
4.4 ± 0.6	984	⁴ BESCH	74 CNTR	$2\gamma p \rightarrow \rho K^+K^-$
4.67 ± 0.72	681	⁴ BALAKIN	71 OSPK	$e^+e^- \rightarrow$ hadrons
4.09 ± 0.29		BIZOT	70 OSPK	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.37 ± 0.02		LEES	13F BABR	$D^+ \rightarrow K^+K^-\pi^+$
4.24 ± 0.02 ± 0.03	542k	⁵ AKHMETSHIN 08	CMD2	$1.02 e^+e^- \rightarrow K^+K^-$
4.28 ± 0.13	12540	⁶ AUBERT,B	05J BABR	$D^0 \rightarrow \bar{K}^0 K^+ K^-$
4.45 ± 0.06	271k	DIJKSTRA	86 SPEC	100 π^- Be
3.6 ± 0.8	337	⁴ COOPER	78B HBC	$0.7-0.8 \bar{p}p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$
4.5 ± 0.50	1300	^{4,6} AKERLOF	77 SPEC	400 $pA \rightarrow K^+K^-X$
4.5 ± 0.8	500	^{4,6} AYRES	74 ASPK	$3-6 \pi^- p \rightarrow K^+K^-n, K^-p \rightarrow K^+K^-\Lambda/\Sigma^0$
3.81 ± 0.37		COSME	74B OSPK	$e^+e^- \rightarrow K_L^0 K_S^0$
3.8 ± 0.7	454	⁴ BORENSTEIN	72 HBC	$2.18 K^- p \rightarrow K \bar{K} n$

NODE=M004W

NODE=M004W

NODE=M004W;LINKAGE=C

NODE=M004W;LINKAGE=GS
NODE=M004W;LINKAGE=AENODE=M004W;LINKAGE=D
NODE=M004W;LINKAGE=AH
NODE=M004W;LINKAGE=A **$\phi(1020)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 K^+K^-	(48.9 ± 0.5) %	S=1.1
Γ_2 $K_L^0 K_S^0$	(34.2 ± 0.4) %	S=1.1
Γ_3 $\rho\pi + \pi^+\pi^-\pi^0$	(15.32 ± 0.32) %	S=1.1
Γ_4 $\rho\pi$		
Γ_5 $\pi^+\pi^-\pi^0$		
Γ_6 $\eta\gamma$	(1.309 ± 0.024) %	S=1.2
Γ_7 $\pi^0\gamma$	(1.27 ± 0.06) × 10 ⁻³	

NODE=M004215;NODE=M004

DESIG=1

DESIG=2

DESIG=24

DESIG=16

DESIG=3

DESIG=4

DESIG=7

Γ_8	$\ell^+ \ell^-$	—			DESIG=256;OUR EVAL;→ UNCHECKED ←
Γ_9	$e^+ e^-$	$(2.954 \pm 0.030) \times 10^{-4}$	S=1.1		DESIG=5
Γ_{10}	$\mu^+ \mu^-$	$(2.87 \pm 0.19) \times 10^{-4}$			DESIG=6
Γ_{11}	$\eta e^+ e^-$	$(1.08 \pm 0.04) \times 10^{-4}$			DESIG=17
Γ_{12}	$\pi^+ \pi^-$	$(7.4 \pm 1.3) \times 10^{-5}$			DESIG=8
Γ_{13}	$\omega \pi^0$	$(4.7 \pm 0.5) \times 10^{-5}$			DESIG=25
Γ_{14}	$\omega \gamma$	< 5	%	CL=84%	DESIG=10
Γ_{15}	$\rho \gamma$	< 1.2	$\times 10^{-5}$	CL=90%	DESIG=12
Γ_{16}	$\pi^+ \pi^- \gamma$	$(4.1 \pm 1.3) \times 10^{-5}$			DESIG=9
Γ_{17}	$f_0(980) \gamma$	$(3.22 \pm 0.19) \times 10^{-4}$	S=1.1		DESIG=20
Γ_{18}	$\pi^0 \pi^0 \gamma$	$(1.13 \pm 0.06) \times 10^{-4}$			DESIG=19
Γ_{19}	$\pi^+ \pi^- \pi^+ \pi^-$	$(4.0 \pm_{-2.2}^{2.8}) \times 10^{-6}$			DESIG=15
Γ_{20}	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	< 4.6	$\times 10^{-6}$	CL=90%	DESIG=14
Γ_{21}	$\pi^0 e^+ e^-$	$(1.12 \pm 0.28) \times 10^{-5}$			DESIG=21
Γ_{22}	$\pi^0 \eta \gamma$	$(7.27 \pm 0.30) \times 10^{-5}$	S=1.5		DESIG=22
Γ_{23}	$a_0(980) \gamma$	$(7.6 \pm 0.6) \times 10^{-5}$			DESIG=23
Γ_{24}	$K^0 \bar{K}^0 \gamma$	< 1.9	$\times 10^{-8}$	CL=90%	DESIG=257
Γ_{25}	$\eta'(958) \gamma$	$(6.25 \pm 0.21) \times 10^{-5}$			DESIG=194
Γ_{26}	$\eta \pi^0 \pi^0 \gamma$	< 2	$\times 10^{-5}$	CL=90%	DESIG=195
Γ_{27}	$\mu^+ \mu^- \gamma$	$(1.4 \pm 0.5) \times 10^{-5}$			DESIG=196
Γ_{28}	$\rho \gamma \gamma$	< 1.2	$\times 10^{-4}$	CL=90%	DESIG=250
Γ_{29}	$\eta \pi^+ \pi^-$	< 1.8	$\times 10^{-5}$	CL=90%	DESIG=255
Γ_{30}	$\eta \mu^+ \mu^-$	< 9.4	$\times 10^{-6}$	CL=90%	DESIG=26
Γ_{31}	$\eta U \rightarrow \eta e^+ e^-$	< 1	$\times 10^{-6}$	CL=90%	DESIG=259
Lepton Family number (LF) violating modes					
Γ_{32}	$e^\pm \mu^\mp$	LF	< 2	$\times 10^{-6}$	CL=90%
NODE=M004;CLUMP=A DESIG=258					

CONSTRAINED FIT INFORMATION

An overall fit to 30 branching ratios uses 79 measurements and one constraint to determine 14 parameters. The overall fit has a $\chi^2 = 57.4$ for 66 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-72									
x_3	-53	-21								
x_6	-13	7	2							
x_7	-5	3	1	5						
x_9	30	-25	-10	-32	-15					
x_{10}	-4	3	1	3	2	-11				
x_{12}	-2	1	0	2	1	-5	1			
x_{13}	-2	2	1	2	1	-7	1	0		
x_{17}	0	0	0	0	0	0	0	0	0	
x_{18}	-6	4	2	17	3	-17	2	1	1	0
x_{19}	0	0	0	0	0	-1	0	0	0	0
x_{23}	0	0	0	0	0	0	0	0	0	0
x_{25}	-4	2	1	32	2	-10	1	1	1	0
	x_1	x_2	x_3	x_6	x_7	x_9	x_{10}	x_{12}	x_{13}	x_{17}
x_{19}	0									
x_{23}	0	0								
x_{25}	5	0	0							
	x_{18}	x_{19}	x_{23}							

$\phi(1020)$ PARTIAL WIDTHS

NODE=M004218

 $\Gamma(\eta\gamma)$ Γ_6

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

58.9 ± 0.5 ± 2.4	ACHASOV	00	SND $e^+e^- \rightarrow \eta\gamma$
------------------	---------	----	-------------------------------------

NODE=M004W6
NODE=M004W6 $\Gamma(\pi^0\gamma)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.40 ± 0.16 ^{+0.43} _{-0.40}	ACHASOV	00	SND $e^+e^- \rightarrow \pi^0\gamma$
---	---------	----	--------------------------------------

NODE=M004W7
NODE=M004W7 $\Gamma(\ell^+\ell^-)$ Γ_8

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.320 ± 0.017 ± 0.015	¹ AMBROSINO	05	KLOE 1.02 $e^+e^- \rightarrow \mu^+\mu^-$
-----------------------	------------------------	----	---

NODE=M004W5
NODE=M004W5 $\Gamma(e^+e^-)$ Γ_9

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1.27 ± 0.04 OUR EVALUATION**1.251 ± 0.021 OUR AVERAGE** Error includes scale factor of 1.1.

1.235 ± 0.006 ± 0.022	² AKHMETSHIN	11	CMD2 1.02 $e^+e^- \rightarrow \phi$
1.32 ± 0.05 ± 0.03	³ AMBROSINO	05	KLOE 1.02 $e^+e^- \rightarrow e^+e^-$
1.28 ± 0.05	AKHMETSHIN	95	CMD2 1.02 $e^+e^- \rightarrow \phi$

NODE=M004W8
NODE=M004W8
→ UNCHECKED ← $(\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-))^{\frac{1}{2}}$ $(\Gamma_9\Gamma_{10})^{\frac{1}{2}}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1.320 ± 0.018 ± 0.017	AMBROSINO	05	KLOE 1.02 $e^+e^- \rightarrow \mu^+\mu^-$
------------------------------	-----------	----	---

¹Weighted average of Γ_{ee} and $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ from AMBROSINO 05 assuming lepton universality.

²Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007.

³From forward-backward asymmetry and using $\Gamma_{\text{total}} = 4.26 \pm 0.05$ MeV from the 2004 edition of this Review.

NODE=M004W9
NODE=M004W9

NODE=M004W5;LINKAGE=AM

NODE=M004W8;LINKAGE=AK

NODE=M004W8;LINKAGE=AM

 $\phi(1020) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M004223

 $\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_9/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

0.6340 ± 0.0070 ± 0.0039	¹ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$
---------------------------------	-------------------	-----	--

¹Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770), \omega(782), \phi(1020)$ and their higher mass excitations. The first error combines statistical and systematic uncertainties. The second one is due to the parametrization of the charged kaon form factor and mass calibration.

NODE=M004G01
NODE=M004G01

NODE=M004G01;LINKAGE=A

 $\phi(1020) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M004224

 $\Gamma(K^+K^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

14.46 ± 0.23 OUR FIT Error includes scale factor of 1.1.**14.24 ± 0.30 OUR AVERAGE**

14.27 ± 0.05 ± 0.31	542k	AKHMETSHIN	08	CMD2 1.02 $e^+e^- \rightarrow K^+K^-$
13.93 ± 0.14 ± 0.99	1000k	¹ ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0$

NODE=M004G10
NODE=M004G10 $\Gamma(K_L^0K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

10.10 ± 0.13 OUR FIT**10.06 ± 0.16 OUR AVERAGE**

10.01 ± 0.04 ± 0.17	272k	² AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow K_L^0K_S^0$
10.27 ± 0.07 ± 0.34	500k	¹ ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0$

NODE=M004G6
NODE=M004G6

$$\frac{\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)}{\Gamma_{\text{total}}} \times \frac{\Gamma(e^+e^-)}{\Gamma_{\text{total}}} \quad \Gamma_3/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
4.53 ±0.10 OUR FIT				Error includes scale factor of 1.1.
4.46 ±0.12 OUR AVERAGE				
4.51 ±0.16 ±0.11	105k	AKHMETSHIN 06	CMD2	0.98-1.06 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰
4.30 ±0.08 ±0.21		AUBERT,B	04N BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ γ
4.665 ±0.042 ±0.261	400k	¹ ACHASOV	01E SND	e ⁺ e ⁻ → K ⁺ K ⁻ , K _S K _L , π ⁺ π ⁻ π ⁰
4.35 ±0.27 ±0.08	11169	³ AKHMETSHIN 98	CMD2	e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.38 ±0.12		BENAYOUN 10	RVUE	0.4-1.05 e ⁺ e ⁻


NODE=M004G7
NODE=M004G7

$$\frac{\Gamma(\eta\gamma)}{\Gamma_{\text{total}}} \times \frac{\Gamma(e^+e^-)}{\Gamma_{\text{total}}} \quad \Gamma_6/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
3.87 ±0.07 OUR FIT				Error includes scale factor of 1.2.
3.93 ±0.09 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
4.050 ±0.067 ±0.118	33k	⁴ ACHASOV	07B SND	0.6-1.38 e ⁺ e ⁻ → ηγ
4.093 ^{+0.040} _{-0.043} ±0.247	17.4k	⁵ AKHMETSHIN 05	CMD2	0.60-1.38 e ⁺ e ⁻ → ηγ
3.850 ±0.041 ±0.159	23k	^{6,7} AKHMETSHIN 01B	CMD2	e ⁺ e ⁻ → ηγ
4.00 ±0.04 ±0.11		⁸ ACHASOV	00 SND	e ⁺ e ⁻ → ηγ
3.53 ±0.08 ±0.17	2200	^{9,10} AKHMETSHIN 99F	CMD2	e ⁺ e ⁻ → ηγ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.19 ±0.06		¹¹ BENAYOUN 10	RVUE	0.4-1.05 e ⁺ e ⁻

NODE=M004G2
NODE=M004G2

WEIGHTED AVERAGE
3.93±0.09 (Error scaled by 1.3)



$$\frac{\Gamma(\eta\gamma)}{\Gamma_{\text{total}}} \times \frac{\Gamma(e^+e^-)}{\Gamma_{\text{total}}} \quad \Gamma_6/\Gamma \times \Gamma_9/\Gamma$$

$$\frac{\Gamma(\pi^0\gamma)}{\Gamma_{\text{total}}} \times \frac{\Gamma(e^+e^-)}{\Gamma_{\text{total}}} \quad \Gamma_7/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10 ⁻⁷)	EVTS	DOCUMENT ID	TECN	COMMENT
3.74 ±0.18 OUR FIT				
3.71 ±0.21 OUR AVERAGE				
3.75 ±0.11 ±0.29	18680	AKHMETSHIN 05	CMD2	0.60-1.38 e ⁺ e ⁻ → π ⁰ γ
3.67 ±0.10 ^{+0.27} _{-0.25}		¹² ACHASOV	00 SND	e ⁺ e ⁻ → π ⁰ γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.29 ±0.11		¹¹ BENAYOUN 10	RVUE	0.4-1.05 e ⁺ e ⁻

NODE=M004G3
NODE=M004G3

$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{10}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-8})

DOCUMENT ID

TECN

COMMENT


NODE=M004G5
NODE=M004G5

8.5 $^{+0.5}_{-0.6}$ OUR FIT

8.8 ± 0.9 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

8.36 $\pm 0.59 \pm 0.37$	ACHASOV	01G	SND	$e^+e^- \rightarrow \mu^+\mu^-$
9.9 $\pm 1.4 \pm 0.9$	⁹ ACHASOV	99C	SND	$e^+e^- \rightarrow \mu^+\mu^-$
14.4 ± 3.0	³ VASSERMAN	81	OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
8.6 ± 5.9	³ AUGUSTIN	73	OSPK	$e^+e^- \rightarrow \mu^+\mu^-$

WEIGHTED AVERAGE
8.8 ± 0.9 (Error scaled by 1.5)



$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{10}/\Gamma \times \Gamma_9/\Gamma$$

$$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{12}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-8})

DOCUMENT ID

TECN

COMMENT

NODE=M004G4
NODE=M004G4

2.2 ± 0.4 OUR FIT

2.2 ± 0.4 OUR AVERAGE

2.1 $\pm 0.3 \pm 0.3$	⁹ ACHASOV	00C	SND	$e^+e^- \rightarrow \pi^+\pi^-$
1.95 $^{+1.15}_{-0.87}$	³ GOLUBEV	86	ND	$e^+e^- \rightarrow \pi^+\pi^-$
6.01 $^{+3.19}_{-2.51}$	³ VASSERMAN	81	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

3.31 ± 0.99 ¹³BENAYOUN 13 RVUE 0.4-1.05 e^+e^-

$$\Gamma(\omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{13}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-8})

DOCUMENT ID

TECN

COMMENT

NODE=M004G11
NODE=M004G11

1.40 ± 0.15 OUR FIT

1.37 $\pm 0.17 \pm 0.01$

^{14,15}AMBROSINO 08G KLOE $e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$

$$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{18}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-8})

DOCUMENT ID

TECN

COMMENT

NODE=M004G9
NODE=M004G9

3.34 ± 0.17 OUR FIT

3.33 $^{+0.04+0.19}_{-0.09-0.20}$

¹⁶AMBROSINO 07 KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$

$$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{19}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-9})

EVTS

DOCUMENT ID

TECN

COMMENT

NODE=M004G8
NODE=M004G8

1.2 $^{+0.8}_{-0.7}$ OUR FIT

1.17 $\pm 0.52 \pm 0.64$

3285

⁹AKHMETSHIN 00E CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

- ¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S^0K_L^0$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.
- ² Update of AKHMETSHIN 99D
- ³ Recalculated by us from the cross section in the peak.
- ⁴ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
- ⁵ From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
- ⁶ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
- ⁷ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
- ⁸ From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow 2\gamma) = (39.21 \pm 0.34) \times 10^{-2}$.
- ⁹ Recalculated by the authors from the cross section in the peak.
- ¹⁰ From the $\eta \rightarrow \pi^+\pi^-\pi^0$ decay and using $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.1 \pm 0.5) \times 10^{-2}$.
- ¹¹ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.
- ¹² From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\pi^0 \rightarrow 2\gamma) = (98.798 \pm 0.032) \times 10^{-2}$.
- ¹³ A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data.
- ¹⁴ Recalculated by the authors from the cross section at the peak.
- ¹⁵ AMBROSINO 08G reports $[\Gamma(\phi(1020) \rightarrow \omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(\phi(1020) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (1.22 \pm 0.13 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ¹⁶ Calculated by the authors from the cross section at the peak.

NODE=M004G;LINKAGE=AE

NODE=M004G;LINKAGE=GS
NODE=M004G;LINKAGE=B
NODE=M004G2;LINKAGE=AHNODE=M004G2;LINKAGE=AK
NODE=M004G;LINKAGE=AK
NODE=M004G;LINKAGE=BQNODE=M004G2;LINKAGE=A
NODE=M004G;LINKAGE=A
NODE=M004G2;LINKAGE=C
NODE=M004G7;LINKAGE=BE
NODE=M004G3;LINKAGE=A
NODE=M004G4;LINKAGE=ANODE=M004G11;LINKAGE=AB
NODE=M004G11;LINKAGE=AM

NODE=M004G9;LINKAGE=AM

 $\phi(1020)$ BRANCHING RATIOS

NODE=M004220

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.489±0.005 OUR FIT				Error includes scale factor of 1.1.
0.493±0.010 OUR AVERAGE				
0.492±0.012	2913	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K^+K^-$
0.44 ±0.05	321	KALBFLEISCH 76	HBC	$2.18 K^-p \rightarrow \Lambda K^+K^-$
0.49 ±0.06	270	DEGROOT 74	HBC	$4.2 K^-p \rightarrow \Lambda\phi$
0.540±0.034	565	BALAKIN 71	OSPK	$e^+e^- \rightarrow K^+K^-$
0.48 ±0.04	252	LINDSEY 66	HBC	$2.1-2.7 K^-p \rightarrow \Lambda K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.493±0.003±0.007		¹ AKHMETSHIN 11	CMD2	$1.02 e^+e^- \rightarrow K^+K^-$
0.476±0.017	1000k	² ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0$

NODE=M004R1
NODE=M004R1 **$\Gamma(K_L^0K_S^0)/\Gamma_{\text{total}}$ Γ_2/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.342±0.004 OUR FIT				Error includes scale factor of 1.1.
0.331±0.009 OUR AVERAGE				
0.335±0.010	40644	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K_L^0K_S^0$
0.326±0.035		DOLINSKY 91	ND	$e^+e^- \rightarrow K_L^0K_S^0$
0.310±0.024		DRUZHININ 84	ND	$e^+e^- \rightarrow K_L^0K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.336±0.002±0.006		¹ AKHMETSHIN 11	CMD2	$1.02 e^+e^- \rightarrow K_S^0K_L^0$
0.351±0.013	500k	² ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0$
0.27 ±0.03	133	KALBFLEISCH 76	HBC	$2.18 K^-p \rightarrow \Lambda K_L^0K_S^0$
0.257±0.030	95	BALAKIN 71	OSPK	$e^+e^- \rightarrow K_L^0K_S^0$
0.40 ±0.04	167	LINDSEY 66	HBC	$2.1-2.7 K^-p \rightarrow \Lambda K_L^0K_S^0$

NODE=M004R2
NODE=M004R2 **$\Gamma(K_L^0K_S^0)/\Gamma(K^+K^-)$ Γ_2/Γ_1**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.698±0.014 OUR FIT				Error includes scale factor of 1.1.
0.740±0.031 OUR AVERAGE				
0.70 ±0.06	2732	BUKIN 78C	OLYA	$e^+e^- \rightarrow K_L^0K_S^0$
0.82 ±0.08		LOSTY 78	HBC	$4.2 K^-p \rightarrow \phi$ hyperon
0.71 ±0.05		LAVEN 77	HBC	$10 K^-p \rightarrow K^+K^-\Lambda$
0.71 ±0.08		LYONS 77	HBC	$3-4 K^-p \rightarrow \Lambda\phi$
0.89 ±0.10	144	AGUILAR-...	72B	$3.9,4.6 K^-p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.68 ±0.03		³ AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K_L^0K_S^0, K^+K^-$

NODE=M004R19
NODE=M004R19

$$\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$$

$$\Gamma_2/(\Gamma_1+\Gamma_2)$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.411±0.005 OUR FIT	Error includes scale factor of 1.1.			
0.45 ±0.04 OUR AVERAGE				
0.44 ±0.07		LONDON	66 HBC	2.24 $K^- p \rightarrow \Lambda K\bar{K}$
0.48 ±0.07	52	BADIER	65B HBC	3 $K^- p$
0.40 ±0.10	34	SCHLEIN	63 HBC	1.95 $K^- p \rightarrow \Lambda K\bar{K}$

NODE=M004R5
NODE=M004R5

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1532±0.0032 OUR FIT	Error includes scale factor of 1.1.			
0.151 ±0.009 OUR AVERAGE	Error includes scale factor of 1.7.			
0.161 ±0.008	11761	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.143 ±0.007		DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.155 ±0.002 ±0.005		¹ AKHMETSHIN 11	CMD2	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.159 ±0.008	400k	² ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$
0.145 ±0.009 ±0.003	11169	⁴ AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.139 ±0.007		⁵ PARROUR 76B	OSPK	e^+e^-

NODE=M004R3
NODE=M004R3

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K^+K^-)$$

$$\Gamma_3/\Gamma_1$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.313±0.009 OUR FIT	Error includes scale factor of 1.1.			
0.28 ±0.09	34	AGUILAR-...	72B HBC	3.9,4.6 $K^- p$

NODE=M004R20
NODE=M004R20

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K\bar{K})$$

$$\Gamma_3/(\Gamma_1+\Gamma_2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.184±0.005 OUR FIT	Error includes scale factor of 1.1.		
0.24 ±0.04 OUR AVERAGE			
0.237±0.039	CERRADA 77B	HBC	4.2 $K^- p \rightarrow \Lambda 3\pi$
0.30 ±0.15	LONDON 66	HBC	2.24 $K^- p \rightarrow \Lambda \pi^+\pi^-\pi^0$

NODE=M004R6
NODE=M004R6

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K_L^0 K_S^0)$$

$$\Gamma_3/\Gamma_2$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.448±0.012 OUR FIT	Error includes scale factor of 1.1.			
0.51 ±0.05 OUR AVERAGE				
0.56 ±0.07	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow K_L^0 K_S^0, \pi^+\pi^-\pi^0$
0.47 ±0.06	516	COSME 74	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M004R7
NODE=M004R7

$$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
≈ 0.0087		1.98M	^{6,7} ALOISIO 03	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.0006	90		⁸ ACHASOV 02	SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.23	90		⁸ CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.20	90		⁸ PARROUR 76B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M004R46
NODE=M004R46

$$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma$$

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
1.309±0.024 OUR FIT	Error includes scale factor of 1.2.			
1.26 ±0.04 OUR AVERAGE				
1.246±0.025±0.057	10k	⁹ ACHASOV 98F	SND	$e^+e^- \rightarrow 7\gamma$
1.18 ±0.11	279	¹⁰ AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0 3\gamma$
1.30 ±0.06		¹¹ DRUZHININ 84	ND	$e^+e^- \rightarrow 3\gamma$
1.4 ±0.2		¹² DRUZHININ 84	ND	$e^+e^- \rightarrow 6\gamma$
0.88 ±0.20	290	KURDADZE 83C	OLYA	$e^+e^- \rightarrow 3\gamma$
1.35 ±0.29		ANDREWS 77	CNTR	6.7-10 γCu
1.5 ±0.4	54	¹¹ COSME 76	OSPK	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.38 ±0.02 ±0.02		¹ AKHMETSHIN 11	CMD2	1.02 $e^+e^- \rightarrow \eta\gamma$
1.37 ±0.05 ±0.01	33k	¹³ ACHASOV 07B	SND	0.6-1.38 $e^+e^- \rightarrow \eta\gamma$
1.373±0.014±0.085	17.4k	^{14,15} AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
1.287±0.013±0.063		^{16,17} AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1.338±0.012±0.052		¹⁸ ACHASOV 00	SND	$e^+e^- \rightarrow \eta\gamma$
1.18 ±0.03 ±0.06	2200	¹⁹ AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \eta\gamma$
1.21 ±0.07		²⁰ BENAYOUN 96	RVUE	0.54-1.04 $e^+e^- \rightarrow \eta\gamma$

NODE=M004R11
NODE=M004R11

OCCUR=2

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_7/Γ VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

1.27 ±0.06 OUR FIT**1.31 ±0.13 OUR AVERAGE**

1.30 ±0.13

DRUZHININ 84 ND $e^+e^- \rightarrow 3\gamma$

1.4 ±0.5

32

COSME 76 OSPK e^+e^-

••• We do not use the following data for averages, fits, limits, etc. •••

1.258±0.037±0.077 18680 ^{21,22} AKHMETSHIN 05 CMD2 0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$ 1.226±0.036^{+0.096}_{-0.089} ²³ ACHASOV 00 SND $e^+e^- \rightarrow \pi^0\gamma$ 1.26 ±0.17 ²⁰ BENAYOUN 96 RVUE 0.54-1.04 $e^+e^- \rightarrow \pi^0\gamma$ NODE=M004R17
NODE=M004R17 $\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$ Γ_6/Γ_7

VALUE

DOCUMENT ID TECN COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

10.9±0.3^{+0.7}_{-0.8} ACHASOV 00 SND $e^+e^- \rightarrow \eta\gamma, \pi^0\gamma$ NODE=M004R42
NODE=M004R42 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_9/Γ VALUE (units 10^{-4}) EVTS

DOCUMENT ID TECN COMMENT

2.954±0.030 OUR FIT Error includes scale factor of 1.1.**2.98 ±0.07 OUR AVERAGE** Error includes scale factor of 1.1.2.93 ±0.14 1900k ²⁴ ACHASOV 01E SND $e^+e^- \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0$ 2.88 ±0.09 55600 AKHMETSHIN 95 CMD2 $e^+e^- \rightarrow \text{hadrons}$ 3.00 ±0.21 3681 BUKIN 78C OLYA $e^+e^- \rightarrow \text{hadrons}$ 3.10 ±0.14 ²⁵ PARROUR 76 OSPK e^+e^- 3.3 ±0.3 COSME 74 OSPK $e^+e^- \rightarrow \text{hadrons}$ 2.81 ±0.25 681 BALAKIN 71 OSPK $e^+e^- \rightarrow \text{hadrons}$ 3.50 ±0.27 CHATELUS 71 OSPK e^+e^- NODE=M004R16
NODE=M004R16 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ VALUE (units 10^{-4})

DOCUMENT ID TECN COMMENT

2.87±0.19 OUR FIT**2.5 ±0.4 OUR AVERAGE**2.69±0.46 ²⁶ HAYES 71 CNTR 8.3,9.8 $\gamma C \rightarrow \mu^+\mu^- X$ 2.17±0.60 ²⁶ EARLES 70 CNTR 6.0 $\gamma C \rightarrow \mu^+\mu^- X$

••• We do not use the following data for averages, fits, limits, etc. •••

2.87±0.20±0.14 ²⁷ ACHASOV 01G SND $e^+e^- \rightarrow \mu^+\mu^-$ 3.30±0.45±0.32 ⁴ ACHASOV 99C SND $e^+e^- \rightarrow \mu^+\mu^-$ 4.83±1.02 ²⁸ VASSERMAN 81 OLYA $e^+e^- \rightarrow \mu^+\mu^-$ 2.87±1.98 ²⁸ AUGUSTIN 73 OSPK $e^+e^- \rightarrow \mu^+\mu^-$ NODE=M004R10
NODE=M004R10 $\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ VALUE (units 10^{-4}) EVTS

DOCUMENT ID TECN COMMENT

1.08 ±0.04 OUR AVERAGE[(1.15 ±0.10) × 10^{-4} OUR 2015 AVERAGE]1.075±0.007±0.038 30k ²⁹ BABUSCI 15 KLOE 1.02 $e^+e^- \rightarrow \eta e^+e^-$ |1.19 ±0.19 ±0.12 213 ³⁰ ACHASOV 01B SND $e^+e^- \rightarrow \eta e^+e^-$ 1.14 ±0.10 ±0.06 355 ³¹ AKHMETSHIN 01 CMD2 $e^+e^- \rightarrow \eta e^+e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

1.13 ±0.14 ±0.07 183 ³² AKHMETSHIN 01 CMD2 $e^+e^- \rightarrow \eta e^+e^-$ 1.21 ±0.14 ±0.09 130 ³³ AKHMETSHIN 01 CMD2 $e^+e^- \rightarrow \eta e^+e^-$ 1.04 ±0.20 ±0.08 42 ³⁴ AKHMETSHIN 01 CMD2 $e^+e^- \rightarrow \eta e^+e^-$ 1.3 ^{+0.8}_{-0.6} 7 GOLUBEV 85 ND $e^+e^- \rightarrow \eta e^+e^-$ NODE=M004R24
NODE=M004R24

NEW

OCCUR=2

OCCUR=3

OCCUR=4

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ VALUE (units 10^{-4}) CL%

DOCUMENT ID TECN COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

0.71±0.11±0.09 ⁴ ACHASOV 00C SND $e^+e^- \rightarrow \pi^+\pi^-$ 0.65^{+0.38}_{-0.29} ⁴ GOLUBEV 86 ND $e^+e^- \rightarrow \pi^+\pi^-$ 2.01^{+1.07}_{-0.84} ⁴ VASSERMAN 81 OLYA $e^+e^- \rightarrow \pi^+\pi^-$ <6.6 95 BUKIN 78B OLYA $e^+e^- \rightarrow \pi^+\pi^-$ <2.7 95 ALVENSLEB... 72 CNTR 6.7 $\gamma C \rightarrow C\pi^+\pi^-$ NODE=M004R18
NODE=M004R18

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

DOCUMENT ID TECN COMMENT

 Γ_{13}/Γ NODE=M004R28
NODE=M004R28**4.7±0.5 OUR FIT****5.2^{+1.3}_{-1.1}**35,36 AULCHENKO 00A SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.4±0.6

37 AMBROSINO 08G KLOE $e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$

~ 5.4

38 ACHASOV 00E SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$ 5.5^{+1.6}_{-1.4}±0.336,39 AULCHENKO 00A SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ 4.8^{+1.9}_{-1.7}±0.838 ACHASOV 99 SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$

VALUE CL%

DOCUMENT ID TECN COMMENT

 Γ_{14}/Γ NODE=M004R14
NODE=M004R14

<0.05

84

LINDSEY 66 HBC 2.1-2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-$ neutrals $\Gamma(\rho\gamma)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) CL%

DOCUMENT ID TECN COMMENT

 Γ_{15}/Γ NODE=M004R15
NODE=M004R15

< 0.12

90

40 AKHMETSHIN 99B CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 7

90

AKHMETSHIN 97C CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

<200

84

LINDSEY 66 HBC 2.1-2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-$ neutrals $\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) CL% EVTS

DOCUMENT ID TECN COMMENT

 Γ_{16}/Γ NODE=M004R12
NODE=M004R12**0.41±0.12±0.04**

30175

41 AKHMETSHIN 99B CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.3

90

42 AKHMETSHIN 97C CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

<600

90

KALBFLEISCH 75 HBC 2.18 $K^-p \rightarrow$ $\Lambda\pi^+\pi^-\gamma$

< 70

90

COSME 74 OSPK $e^+e^- \rightarrow \pi^+\pi^-\gamma$

<400

90

LINDSEY 65 HBC 2.1-2.7 $K^-p \rightarrow$ $\Lambda\pi^+\pi^-$ neutrals $\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) CL% EVTS

DOCUMENT ID TECN COMMENT

 Γ_{17}/Γ NODE=M004R30
NODE=M004R30**3.22±0.19 OUR FIT** Error includes scale factor of 1.1.**3.21±0.19 OUR AVERAGE**3.21^{+0.03}_{-0.09}±0.1843 AMBROSINO 07 KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$

2.90±0.21±1.54

44 AKHMETSHIN 99C CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma,$
 $\pi^0\pi^0\gamma$

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.47±0.21

2438

45 ALOISIO 02D KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$ 3.5 ±0.3 ^{+1.3}_{-0.5}

419

46,47 ACHASOV 00H SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$

1.93±0.46±0.50

27188

48 AKHMETSHIN 99B CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

3.05±0.25±0.72

268

49 AKHMETSHIN 99C CMD2 $e^+e^- \rightarrow \pi^0\pi^0\gamma$

1.5 ±0.5

268

50 AKHMETSHIN 99C CMD2 $e^+e^- \rightarrow \pi^0\pi^0\gamma$

OCCUR=2

3.42±0.30±0.36

164

46 ACHASOV 98I SND $e^+e^- \rightarrow 5\gamma$

< 1

90

51 AKHMETSHIN 97C CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

< 7

90

52 AKHMETSHIN 97C CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

OCCUR=2

< 20

90

DRUZHININ 87 ND $e^+e^- \rightarrow \pi^0\pi^0\gamma$ $\Gamma(f_0(980)\gamma)/\Gamma(\eta\gamma)$ VALUE (units 10^{-2})

EVTS DOCUMENT ID TECN COMMENT

 Γ_{17}/Γ_6 NODE=M004R44
NODE=M004R44**2.46±0.15 OUR FIT** Error includes scale factor of 1.1.**2.6 ±0.2 ^{+0.8}_{-0.3}**

419

46 ACHASOV 00H SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ±0.06 OUR AVERAGE					
1.07 ^{+0.01} _{-0.03} ^{+0.06} _{-0.06}			53 AMBROSINO	07 KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.08 ±0.17 ±0.09		268	AKHMETSHIN	99C CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.09 ±0.03 ±0.05		2438	ALOISIO	02D KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.158 ±0.093 ±0.052		419	47,54 ACHASOV	00H SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
<10		90	DRUZHININ	87 ND	$e^+e^- \rightarrow 5\gamma$

NODE=M004R26
 NODE=M004R26

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\eta\gamma)$ Γ_{18}/Γ_6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ±0.04 OUR FIT				
0.865 ±0.070 ±0.017	419	54 ACHASOV	00H SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.90 ±0.08 ±0.07	164	ACHASOV	98I SND	$e^+e^- \rightarrow 5\gamma$

NODE=M004R39
 NODE=M004R39

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
3.93 ±1.74 ±2.14		3285	AKHMETSHIN	00E CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
< 870		90	CORDIER	79 WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M004R22
 NODE=M004R22

 $\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	AKHMETSHIN	00E CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<150	95	BARKOV	88 CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

NODE=M004R27
 NODE=M004R27

 $\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.12 ±0.28 OUR AVERAGE					
1.01 ±0.28 ±0.29		52	55 ACHASOV	02D SND	$e^+e^- \rightarrow \pi^0e^+e^-$
1.22 ±0.34 ±0.21		46	56 AKHMETSHIN	01C CMD2	$e^+e^- \rightarrow \pi^0e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<12	90		DOLINSKY	88 ND	$e^+e^- \rightarrow \pi^0e^+e^-$

NODE=M004R31
 NODE=M004R31

 $\Gamma(\pi^0\eta\gamma)/\Gamma_{\text{total}}$ Γ_{22}/Γ


VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.27 ±0.30 OUR AVERAGE					Error includes scale factor of 1.5. See the ideogram below.
7.06 ±0.22		16.9k	57 AMBROSINO	09F KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
8.51 ±0.51 ±0.57		607	58 ALOISIO	02C KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
7.96 ±0.60 ±0.40		197	59 ALOISIO	02C KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
8.8 ±1.4 ±0.9		36	60 ACHASOV	00F SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
9.0 ±2.4 ±1.0		80	AKHMETSHIN	99C CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
7.01 ±0.10 ±0.20		13.3k	58,61 AMBROSINO	09F KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
7.12 ±0.13 ±0.22		3.6k	59,62 AMBROSINO	09F KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
8.3 ±2.3 ±1.2		20	ACHASOV	98B SND	$e^+e^- \rightarrow 5\gamma$
<250	90		DOLINSKY	91 ND	$e^+e^- \rightarrow \pi^0\eta\gamma$

NODE=M004R32
 NODE=M004R32

OCCUR=2

OCCUR=2

OCCUR=3

 **$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$**

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ
7.6±0.6 OUR FIT						
7.6±0.6 OUR AVERAGE						
7.4±0.7			63	ALOISIO 02C	KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$	
8.8±1.7	36		64	ACHASOV 00F	SND $e^+e^- \rightarrow \eta\pi^0\gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
11 ±2			65	GOKALP 02	RVUE $e^+e^- \rightarrow \eta\pi^0\gamma$	
<500	90			DOLINSKY 91	ND $e^+e^- \rightarrow \pi^0\eta\gamma$	

NODE=M004R33
NODE=M004R33

 $\Gamma(f_0(980)\gamma)/\Gamma(a_0(980)\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ_{23}
6.1±0.6	66	ALOISIO 02C	KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$	

NODE=M004R47
NODE=M004R47

 $\Gamma(K^0\bar{K}^0\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{24}/Γ
<1.9 × 10⁻⁸	90	AMBROSINO 09C	KLOE	$e^+e^- \rightarrow K_S^0 K_S^0 \gamma$	

NODE=M004R48
NODE=M004R48

 $\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{25}/Γ
6.25±0.21 OUR FIT						
6.25±0.30 OUR AVERAGE						
6.25±0.28±0.11	3407		67	AMBROSINO 07A	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\gamma$	
6.7 ^{+2.8} _{-2.4} ±0.8	12		68	AULCHENKO 03B	SND $e^+e^- \rightarrow \eta'\gamma$	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
6.7 ^{+5.0} _{-4.2} ±1.5	7			AULCHENKO 03B	SND $e^+e^- \rightarrow 7\gamma$	
6.10±0.61±0.43	120		69	ALOISIO 02E	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-3\gamma$	
8.2 ^{+2.1} _{-1.9} ±1.1	21		70	AKHMETSHIN 00B	CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$	
4.9 ^{+2.2} _{-1.8} ±0.6	9		71	AKHMETSHIN 00F	CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^- \geq 2\gamma$	
6.4 ±1.6	30		72	AKHMETSHIN 00F	CMD2 $e^+e^- \rightarrow \eta'(958)\gamma$	OCCUR=2
6.7 ^{+3.4} _{-2.9} ±1.0	5		73	AULCHENKO 99	SND $e^+e^- \rightarrow \pi^+\pi^-3\gamma$	
<11	90			AULCHENKO 98	SND $e^+e^- \rightarrow 7\gamma$	
12 ⁺⁷ ₋₅ ±2	6		70	AKHMETSHIN 97B	CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$	
<41	90			DRUZHININ 87	ND $e^+e^- \rightarrow \gamma\eta\pi^+\pi^-$	

NODE=M004R25
NODE=M004R25

$\Gamma(\eta'(958)\gamma)/\Gamma(K_L^0 K_S^0)$ Γ_{25}/Γ_2 VALUE (units 10^{-4}) EVTS

DOCUMENT ID TECN COMMENT

1.83±0.06 OUR FIT1.46^{+0.64}_{-0.54}±0.18

9

74

AKHMETSHIN 00F

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^- \geq 2\gamma$ NODE=M004R43
NODE=M004R43 $\Gamma(\eta'(958)\gamma)/\Gamma(\eta\gamma)$ Γ_{25}/Γ_6 VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

4.77±0.15 OUR FIT**4.78±0.20 OUR AVERAGE**

4.77±0.09±0.19

3407

AMBROSINO 07A

KLOE

1.02 $e^+e^- \rightarrow \pi^+\pi^-7\gamma$

4.70±0.47±0.31

120

75

ALOISIO 02E

KLOE

1.02 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 6.5^{+1.7}_{-1.5}±0.8

21

AKHMETSHIN 00B

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5^{+5.2}_{-4.0}±1.4

6

76

AKHMETSHIN 97B

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ NODE=M004R34
NODE=M004R34 $\Gamma(\eta\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ VALUE (units 10^{-5}) CL%

DOCUMENT ID TECN COMMENT

<2

90

AULCHENKO 98

SND

 $e^+e^- \rightarrow 7\gamma$ NODE=M004R36
NODE=M004R36 $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ VALUE (units 10^{-5}) EVTS

DOCUMENT ID TECN COMMENT

1.43±0.45±0.14

27188

48

AKHMETSHIN 99B

CMD2

 $e^+e^- \rightarrow \mu^+\mu^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ±1.0

824 ± 33

77

AKHMETSHIN 97C

CMD2

 $e^+e^- \rightarrow \mu^+\mu^-\gamma$ NODE=M004R35
NODE=M004R35 $\Gamma(\rho\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{28}/Γ VALUE (units 10^{-4}) CL%

DOCUMENT ID TECN COMMENT

<1.2

90

AULCHENKO 08

CMD2

 $\phi \rightarrow \pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5

90

AKHMETSHIN 98

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ NODE=M004R37
NODE=M004R37 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ VALUE (units 10^{-5}) CL%

DOCUMENT ID TECN COMMENT

< 1.8

90

AKHMETSHIN 00E

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.1

90

AULCHENKO 08

CMD2

 $\phi \rightarrow \eta\pi^+\pi^-$

<30

90

AKHMETSHIN 98

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ NODE=M004R38
NODE=M004R38 $\Gamma(\eta\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ VALUE (units 10^{-6}) CL%

DOCUMENT ID TECN COMMENT

<9.4

90

AKHMETSHIN 01

CMD2

 $e^+e^- \rightarrow \eta e^+e^-$ NODE=M004R45
NODE=M004R45 $\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE CL%

DOCUMENT ID TECN COMMENT

<1 × 10⁻⁶

90

78

BABUSCI 13B

KLOE

1.02 $e^+e^- \rightarrow \eta e^+e^-$ NODE=M004R01
NODE=M004R01

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

³ Theoretical analysis of BRAMON 00 taking into account phase-space difference, electromagnetic radiative corrections, as well as isospin breaking, predicts 0.62. FLOREZ-BAEZ 08 predicts 0.63 considering also structure-dependent radiative corrections. FIS-CHBACH 02 calculates additional corrections caused by the close threshold and predicts 0.68. See also BENAYOUN 01 and DUBYNSKIY 07. BENAYOUN 12 obtains 0.71 ± 0.01 in the HLS model.

⁴ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁵ Using $\Gamma(\phi) = 4.1$ MeV. If interference between the $\rho\pi$ and 3π modes is neglected, the fraction of the $\rho\pi$ is more than 80% at the 90% confidence level.

⁶ From a fit without limitations on charged and neutral ρ masses and widths.

⁷ Adding the direct and $\omega\pi$ contributions and considering the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

⁸ Neglecting the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

⁹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.2 \pm 0.4) \times 10^{-2}$.

¹⁰ From $\pi^+\pi^-\pi^0$ decay mode of η .

NODE=M004R1;LINKAGE=AK

NODE=M004R;LINKAGE=B2
NODE=M004R19;LINKAGE=KHNODE=M004R;LINKAGE=8D
NODE=M004R3;LINKAGE=ENODE=M004R;LINKAGE=L1
NODE=M004R;LINKAGE=L2NODE=M004R;LINKAGE=46
NODE=M004R11;LINKAGE=AC
NODE=M004R11;LINKAGE=Z3

- 11 From 2γ decay mode of η .
 12 From $3\pi^0$ decay mode of η .
 13 ACHASOV 07B reports $[\Gamma(\phi(1020) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow e^+e^-)] = (4.050 \pm 0.067 \pm 0.118) \times 10^{-6}$ which we divide by our best value $B(\phi(1020) \rightarrow e^+e^-) = (2.954 \pm 0.030) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.
 14 Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
 15 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.
 16 Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
 17 The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
 18 From the $\eta \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.
 19 From $\pi^+\pi^-\pi^0$ decay mode of η and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.
 20 Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.
 21 Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$.
 22 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.
 23 From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.
 24 From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.
 25 Using total width 4.2 MeV. They detect 3π mode and observe significant interference with ω tail. This is accounted for in the result quoted above.
 26 Neglecting interference between resonance and continuum.
 27 Using $B(\phi \rightarrow e^+e^-) = (2.91 \pm 0.07) \times 10^{-4}$.
 28 Recalculated by us using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.
 29 Using $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.23)\%$ from PDG 12.
 30 Using $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.32)\%$, $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06)\%$, and $B(\phi \rightarrow e^+e^-) = (3.00 \pm 0.06) \times 10^{-4}$.
 31 The average of the branching ratios separately obtained from the $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$ decays.
 32 From $\eta \rightarrow \gamma\gamma$ decays and using $B(\eta \rightarrow \gamma\gamma) = (39.33 \pm 0.25) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.
 33 From $\eta \rightarrow 3\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.
 34 From $\eta \rightarrow \pi^+\pi^-\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\pi^0 \rightarrow e^+e^-\gamma) = (1.198 \pm 0.032) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.0 \pm 0.4) \times 10^{-2}$, $B(\phi \rightarrow \pi^+\pi^-\pi^0) = (15.5 \pm 0.6) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.
 35 Using the 1996 and 1998 data.
 36 $(2.3 \pm 0.3)\%$ correction for other decay modes of the $\omega(782)$ applied.
 37 Not independent of the corresponding $\Gamma(\omega\pi^0) \times \Gamma(e^+e^-) / \Gamma^2(\text{total})$.
 38 Using the 1996 data.
 39 Using the 1998 data.
 40 Supersedes AKHMETSHIN 97C.
 41 For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible. Supersedes AKHMETSHIN 97C.
 42 For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible.
 43 Obtained by the authors taking into account the $\pi^+\pi^-$ decay mode. Includes a component due to $\pi\pi$ production via the $f_0(500)$ meson. Supersedes ALOISIO 02D.
 44 From the combined fit of the photon spectra in the reactions $e^+e^- \rightarrow \pi^+\pi^-\gamma$, $\pi^0\pi^0\gamma$.
 45 From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution. Superseded by AMBROSINO 07.
 46 Assuming that the $\pi^0\pi^0\gamma$ final state is completely determined by the $f_0\gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$.
 47 Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.
 48 For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.
 49 Neglecting other intermediate mechanisms ($\rho\pi$, $\sigma\gamma$).
 50 A narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.
 51 For destructive interference with the Bremsstrahlung process
 52 For constructive interference with the Bremsstrahlung process
 53 Supersedes ALOISIO 02D.
 54 Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.
 55 Using various branching ratios from the 2000 Edition of this Review (PDG 00).
- NODE=M004R11;LINKAGE=A
 NODE=M004R11;LINKAGE=C
 NODE=M004R11;LINKAGE=AO
 NODE=M004R11;LINKAGE=AH
 NODE=M004R11;LINKAGE=AK
 NODE=M004R;LINKAGE=AK
 NODE=M004R;LINKAGE=BQ
 NODE=M004R;LINKAGE=GA
 NODE=M004R;LINKAGE=FF
 NODE=M004R;LINKAGE=TS
 NODE=M004R17;LINKAGE=AH
 NODE=M004R17;LINKAGE=AK
 NODE=M004R;LINKAGE=3G
 NODE=M004R;LINKAGE=AE
 NODE=M004R16;LINKAGE=E
 NODE=M004R10;LINKAGE=A
 NODE=M004R;LINKAGE=GZ
 NODE=M004R;LINKAGE=VA
 NODE=M004R24;LINKAGE=A
 NODE=M004R;LINKAGE=VM
 NODE=M004R;LINKAGE=H1
 NODE=M004R;LINKAGE=H2
 NODE=M004R;LINKAGE=H3
 NODE=M004R;LINKAGE=H4
 NODE=M004R28;LINKAGE=K2
 NODE=M004R28;LINKAGE=K3
 NODE=M004R28;LINKAGE=AM
 NODE=M004R28;LINKAGE=VS
 NODE=M004R28;LINKAGE=K1
 NODE=M004R;LINKAGE=1N
 NODE=M004R;LINKAGE=2N
 NODE=M004R12;LINKAGE=Z3
 NODE=M004R30;LINKAGE=MB
 NODE=M004R;LINKAGE=SL
 NODE=M004R;LINKAGE=KD
 NODE=M004R;LINKAGE=AI
 NODE=M004R;LINKAGE=U8
 NODE=M004R;LINKAGE=3N
 NODE=M004R;LINKAGE=SM
 NODE=M004R;LINKAGE=ST
 NODE=M004R30;LINKAGE=A
 NODE=M004R30;LINKAGE=B
 NODE=M004R26;LINKAGE=MB
 NODE=M004R;LINKAGE=V8
 NODE=M004R;LINKAGE=DS

- 56 Using $B(\pi^0 \rightarrow \gamma\gamma) = 0.98798 \pm 0.00032$, $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$, and $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$.
- 57 Combined results of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decay modes measurements.
- 58 From the decay mode $\eta \rightarrow \gamma\gamma$.
- 59 From the decay mode $\eta \rightarrow \pi^+\pi^-\pi^0$.
- 60 Supersedes ACHASOV 98B.
- 61 Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.31 \pm 0.20)\%$.
- 62 Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.73 \pm 0.28)\%$.
- 63 Using $M_{a_0(980)} = 984.8$ MeV and assuming $a_0(980)\gamma$ dominance.
- 64 Assuming $a_0(980)\gamma$ dominance in the $\eta\pi^0\gamma$ final state.
- 65 Using data of ACHASOV 00F.
- 66 Using results of ALOISIO 02D and assuming that $f_0(980)$ decays into $\pi\pi$ only and $a_0(980)$ into $\eta\pi$ only.
- 67 AMBROSINO 07A reports $[\Gamma(\phi(1020) \rightarrow \eta'(958)\gamma)/\Gamma_{\text{total}}] / [B(\phi(1020) \rightarrow \eta\gamma)] = (4.77 \pm 0.09 \pm 0.19) \times 10^{-3}$ which we multiply by our best value $B(\phi(1020) \rightarrow \eta\gamma) = (1.309 \pm 0.024) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- 68 Averaging AULCHENKO 03B with AULCHENKO 99.
- 69 Using $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033)\%$.
- 70 Using the value $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06) \times 10^{-2}$.
- 71 Using $B(\phi \rightarrow K_L^0 K_S^0) = (33.8 \pm 0.6)\%$.
- 72 Averaging AKHMETSHIN 00B with AKHMETSHIN 00F.
- 73 Using the value $B(\eta' \rightarrow \eta\pi^+\pi^-) = (43.7 \pm 1.5) \times 10^{-2}$ and $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.31) \times 10^{-2}$.
- 74 Using various branching ratios of K_S^0 , K_L^0 , η , η' from the 2000 edition (The European Physical Journal **C15** 1 (2000)) of this Review.
- 75 From the decay mode $\eta' \rightarrow \eta\pi^+\pi^-$, $\eta \rightarrow \gamma\gamma$.
- 76 Superseded by AKHMETSHIN 00B.
- 77 For $E_\gamma > 20$ MeV.
- 78 For a narrow vector U with mass between 5 and 470 MeV, from the combined analysis of $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow \pi^0\pi^0\pi^0$ from ARCHILLI 12. Measured 90% CL limits as a function of m_U range from 2.2×10^{-8} to 10^{-6} .

NODE=M004R;LINKAGE=5H

NODE=M004R32;LINKAGE=AM

NODE=M004R;LINKAGE=C1

NODE=M004R;LINKAGE=C2

NODE=M004R32;LINKAGE=AF

NODE=M004R32;LINKAGE=AB

NODE=M004R32;LINKAGE=AR

NODE=M004R;LINKAGE=C3

NODE=M004R33;LINKAGE=AF

NODE=M004R;LINKAGE=GK

NODE=M004R;LINKAGE=C4

NODE=M004R25;LINKAGE=AM

NODE=M004R25;LINKAGE=BK

NODE=M004R;LINKAGE=E2

NODE=M004R25;LINKAGE=Q

NODE=M004R;LINKAGE=T2

NODE=M004R;LINKAGE=T3

NODE=M004R25;LINKAGE=AU

NODE=M004R;LINKAGE=T1

NODE=M004R;LINKAGE=E1

NODE=M004R;LINKAGE=KS

NODE=M004R35;LINKAGE=A

NODE=M004R01;LINKAGE=A

Lepton Family number (LF) violating modes

NODE=M00422A

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$					Γ_{32}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 2 \times 10^{-6}$	90	ACHASOV	10A SND	$e^+e^- \rightarrow e^\pm \mu^\mp$	

NODE=M004R29

NODE=M004R29

$\pi^+\pi^-\pi^0 / \rho\pi$ AMPLITUDE RATIO a_1 IN DECAY OF $\phi \rightarrow \pi^+\pi^-\pi^0$

NODE=M004D1

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

NODE=M004D1

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±1.2 OUR AVERAGE					
10.1±4.4±1.7		80k	¹ AKHMETSHIN 06	CMD2	1.017–1.021 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0±1.1±0.6		1.98M	^{2,3} ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M004D1

••• We do not use the following data for averages, fits, limits, etc. •••

$-6 < a_1 < 6$		500k	³ ACHASOV	02 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$-16 < a_1 < 11$	90	9.8k	^{1,4} AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$

¹ Dalitz plot analysis taking into account interference between the contact and $\rho\pi$ amplitudes.

NODE=M004D1;LINKAGE=AK

² From a fit without limitations on charged and neutral ρ masses and widths.

NODE=M004D;LINKAGE=L1

³ Recalculated by us to match the notations of AKHMETSHIN 98.

NODE=M004D;LINKAGE=L3

⁴ Assuming zero phase for the contact term.

NODE=M004D1;LINKAGE=KL

PARAMETER β IN $\phi \rightarrow \eta e^+ e^-$ DECAY

NODE=M004BFP

In the one-pole approximation the electromagnetic transition form factor for $\phi \rightarrow \eta e^+ e^-$ is given as a function of the $e^+ e^-$ invariant mass squared, q^2 , by the expression:

NODE=M004BFP

$$|F(q^2)|^2 = (1 - q^2/\Lambda^2)^{-2},$$

where vector meson dominance predicts parameter $\Lambda \approx 0.770 \text{ GeV}$ ($\Lambda^{-2} \approx 1.687 \text{ GeV}^{-2}$). The slope of this form factor, $\beta = dF/dq^2(q^2=0)$, equals Λ^{-2} in this approximation.

The measurements below obtain β in the one-pole approximation.

VALUE (GeV ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29±0.13 OUR AVERAGE				
$1.28 \pm 0.10^{+0.09}_{-0.08}$	30k	¹ BABUSCI	15 KLOE	$1.02 e^+ e^- \rightarrow \eta e^+ e^-$
3.8 ± 1.8	213	ACHASOV	01B SND	$1.02 e^+ e^- \rightarrow \eta e^+ e^-$

¹ The uncertainty is statistical only with negligible systematic one.

NODE=M004BFP

NODE=M004BFP;LINKAGE=A

$\phi(1020)$ REFERENCES

NODE=M004

BABUSCI	15	PL B742 1	D. Babusi <i>et al.</i>	(KLOE Collab.)	REFID=56374
BABUSCI	13B	PL B720 111	D. Babusi <i>et al.</i>	(KLOE-2 Collab.)	REFID=55068
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono (PARIN, BERLIN+)		REFID=55357
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55127
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
ARCHILLI	12	PL B706 251	F. Archilli <i>et al.</i>	(KLOE-2 Collab.)	REFID=53951
BENAYOUN	12	EPJ C72 1848	M. Benayoun <i>et al.</i>		REFID=54281
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)	REFID=54305
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
AKHMETSHIN	11	PL B695 412	R. Akhmetshin <i>et al.</i>	(CMD2 Collab.)	REFID=53645
ACHASOV	10A	PR D81 057102	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=53352
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
AMBROSINO	09C	PL B679 10	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52969
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=53105
AKHMETSHIN	08	PL B669 217	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=52572
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52573
AULCHENKO	08	JETPL 88 85	V. Aulchenko <i>et al.</i>	(CMD-2 Collab.)	REFID=52268
Translated from ZETFP 88 93.					
FLOREZ-BAEZ	08	PR D78 077301	F.V. Florez-Baez, G. Lopez Castro		REFID=52584
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51616
AMBROSINO	07A	PL B648 267	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51646
DUBYSNKIY	07	PR D75 113001	S. Dubynskiy <i>et al.</i>		REFID=51719
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AKHMETSHIN	06	PL B642 203	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=51465
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AMBROSINO	05	PL B608 199	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=50453
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50824
AKHMETSHIN	04N	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
AULCHENKO	03B	JETP 97 24	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49613
Translated from ZETF 124 28.					
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48549
ACHASOV	02D	JETPL 75 449	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48814
Translated from ZETFP 75 539.					
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48823
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
ALOISIO	02E	PL B541 45	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48825
FISCHBACH	02	PL B526 355	E. Fischbach, A.W. Overhauser, B. Woodahl		REFID=48575
GOKALP	02	JP G28 2783	A. Gokalp <i>et al.</i>		REFID=49167
ACHASOV	01B	PL B504 275	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48111
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)	REFID=48312
ACHASOV	01G	PRL 86 1698	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48315
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
AKHMETSHIN	01	PL B501 191	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48110
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
AKHMETSHIN	01C	PL B503 237	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48323
BENAYOUN	01	EPJ C22 503	M. Benayoun, H.B. O'Connell		REFID=48570
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00B	JETP 90 17	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47425
Translated from ZETF 117 22.					
ACHASOV	00C	PL B474 188	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47431
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
Translated from ZETFP 72 411.					
ACHASOV	00E	NP B569 158	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47927
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	00B	PL B473 337	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47422
AKHMETSHIN	00E	PL B491 81	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47936
AKHMETSHIN	00F	PL B494 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47937
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
Translated from ZETF 117 1067.					
BRAMON	00	PL B486 406	A. Bramon <i>et al.</i>		REFID=47969
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ACHASOV	99	PL B449 122	M.N. Achasov <i>et al.</i>		REFID=46896
ACHASOV	99C	PL B456 304	M.N. Achasov <i>et al.</i>		REFID=46939
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47393
AKHMETSHIN	99D	PL B466 385	R.R. Akhmetshin <i>et al.</i>		REFID=47397
Also		PL B508 217 (errata.)	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48328
AKHMETSHIN	99F	PL B460 242	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47473
AULCHENKO	99	JETPL 69 97	V.M. Aulchenko <i>et al.</i>		REFID=46920
Translated from ZETFP 69 87.					

ACHASOV	98B	PL B438 441	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=46317
ACHASOV	98F	JETPL 68 573	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=46321
ACHASOV	98I	PL B440 442	M.N. Achasov et al.		REFID=46600
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin et al.	(CMD-2 Collab.)	REFID=46325
AULCHENKO	98	PL B436 199	V.M. Aulchenko et al.	(Novosibirsk SND Collab.)	REFID=46336
BARBERIS	98	PL B432 436	D. Barberis et al.	(Omega Expt.)	REFID=46344
AKHMETSHIN	97B	PL B415 445	R.R. Akhmetshin et al.	(NOVO, BOST, PITT+)	REFID=45801
AKHMETSHIN	97C	PL B415 452	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)	REFID=45802
BENAYOUN	96	ZPHY C72 221	M. Benayoun et al.	(IPNP, NOVO)	REFID=45753
AKHMETSHIN	95	PL B364 199	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)	REFID=44617
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky et al.	(NOVO)	REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn et al.	(MPIM)	REFID=45862
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky et al.	(NOVO)	REFID=41003
BARKOV	88	SJNP 47 248	L.M. Barkov et al.	(NOVO)	REFID=41024
		Translated from YAF 47 393.			
DOLINSKY	88	SJNP 48 277	S.I. Dolinsky et al.	(NOVO)	REFID=41022
		Translated from YAF 48 442.			
DRUZHININ	87	ZPHY C37 1	V.P. Druzhinin et al.	(NOVO)	REFID=40448
ARMSTRONG	86	PL 166B 245	T.A. Armstrong et al.	(ATHU, BARI, BIRM+)	REFID=20563
ATKINSON	86	ZPHY C30 521	M. Atkinson et al.	(BONN, CERN, GLAS+)	REFID=20564
BEBEK	86	PRL 56 1893	C. Bebek et al.	(CLEO Collab.)	REFID=11540
DAVENPORT	86	PR D33 2519	T.F. Davenport	(TUFTS, ARIZ, FNAL, FSU, NDAM+)	REFID=20567
DIJKSTRA	86	ZPHY C31 375	H. Dijkstra et al.	(ANIK, BRIS, CERN+)	REFID=20568
FRAME	86	NP B276 667	D. Frame et al.	(GLAS)	REFID=20569
GOLUBEV	86	SJNP 44 409	V.B. Golubev et al.	(NOVO)	REFID=40449
		Translated from YAF 44 633.			
ALBRECHT	85D	PL 153B 343	H. Albrecht et al.	(ARGUS Collab.)	REFID=20562
GOLUBEV	85	SJNP 41 756	V.B. Golubev et al.	(NOVO)	REFID=40450
		Translated from YAF 41 1183.			
DRUZHININ	84B	PL 144B 136	V.P. Druzhinin et al.	(NOVO)	REFID=20561
ARMSTRONG	83B	NP B224 193	T.A. Armstrong et al.	(BARI, BIRM, CERN+)	REFID=20558
BARATE	83	PL 121B 449	R. Barate et al.	(SACL, LOIC, SHMP, IND)	REFID=12177
KURDADZE	83C	JETPL 38 366	L.M. Kurdadze et al.	(NOVO)	REFID=20560
		Translated from ZETFP 38 306.			
ARENTON	82	PR D25 2241	M.W. Arenton et al.	(ANL, ILL)	REFID=20556
PELLINEN	82	PS 25 599	A. Pellinen, M. Roos	(HELS)	REFID=20557
DAUM	81	PL 100B 439	C. Daum et al.	(AMST, BRIS, CERN, CRAC+)	REFID=20552
IVANOV	81	PL 107B 297	P.M. Ivanov et al.	(NOVO)	REFID=20553
		Also Private Comm.	S.I. Eidelman	(NOVO)	REFID=20554
VASSERMAN	81	PL 99B 62	I.B. Vasserman et al.	(NOVO)	REFID=20555
		Also SJNP 35 240	L.M. Kurdadze et al.		REFID=47475
		Translated from YAF 35 352.			
CORDIER	80	NP B172 13	A. Cordier et al.	(LALO)	REFID=20240
CORDIER	79	PL 81B 389	A. Cordier et al.	(LALO)	REFID=20549
BUKIN	78B	SJNP 27 521	A.D. Bukin et al.	(NOVO)	REFID=20545
		Translated from YAF 27 985.			
BUKIN	78C	SJNP 27 516	A.D. Bukin et al.	(NOVO)	REFID=20544
		Translated from YAF 27 976.			
COOPER	78B	NP B146 1	A.M. Cooper et al.	(TATA, CERN, CDEF+)	REFID=20235
LOSTY	78	NP B133 38	M.J. Losty et al.	(CERN, AMST, NIJM+)	REFID=20547
AKERLOF	77	PRL 39 861	C.W. Akerlof et al.	(FNAL, MICH, PURD)	REFID=20534
ANDREWS	77	PRL 38 198	D.E. Andrews et al.	(ROCH)	REFID=20120
BALDI	77	PL 68B 381	R. Baldi et al.	(GEVA)	REFID=20536
CERRADA	77B	NP B126 241	M. Cerrada et al.	(AMST, CERN, NIJM+)	REFID=20537
COHEN	77	PRL 38 269	D. Cohen et al.	(ANL)	REFID=20538
LAVEN	77	NP B127 43	H. Laven et al.	(AACH3, BERL, CERN, LOIC+)	REFID=20541
LYONS	77	NP B125 207	L. Lyons, A.M. Cooper, A.G. Clark	(OXF)	REFID=20232
COSME	76	PL 63B 352	G. Cosme et al.	(ORSAY)	REFID=20529
KALBFLEISCH	76	PR D13 22	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20531
PARROUR	76	PL 63B 357	G. Parrou et al.	(ORSAY)	REFID=20532
PARROUR	76B	PL 63B 362	G. Parrou et al.	(ORSAY)	REFID=20533
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AYRES	74	PRL 32 1463	D.S. Ayres et al.	(ANL)	REFID=20522
BESCH	74	NP B70 257	H.J. Besch et al.	(BONN)	REFID=20523
COSME	74	PL 48B 155	G. Cosme et al.	(ORSAY)	REFID=20525
COSME	74B	PL 48B 159	G. Cosme et al.	(ORSAY)	REFID=20526
DEGROOT	74	NP B74 77	A.J. de Groot et al.	(AMST, NIJM)	REFID=20527
AUGUSTIN	73	PRL 30 462	J.E. Augustin et al.	(ORSAY)	REFID=47515
BALLAM	73	PR D7 3150	J. Ballam et al.	(SLAC, LBL)	REFID=20520
BINNIE	73B	PR D8 2789	D.M. Binnie et al.	(LOIC, SHMP)	REFID=20216
AGUILAR...	72B	PR D6 29	M. Aguilar-Benitez et al.	(BNL)	REFID=20205
ALVENSLEB...	72	PRL 28 66	H. Alvensleben et al.	(MIT, DESY)	REFID=20514
BORENSTEIN	72	PR D5 1559	S.R. Borenstein et al.	(BNL, MICH)	REFID=20215
COLLEY	72	NP B50 1	D.C. Colley et al.	(BIRM, GLAS)	REFID=20519
BALAKIN	71	PL 34B 328	V.E. Balakin et al.	(NOVO)	REFID=20507
CHATELUS	71	Thesis LAL 1247	Y. Chatelus	(STRB)	REFID=20508
		Also PL 32B 416	J.C. Bizot et al.	(ORSAY)	REFID=20501
HAYES	71	PR D4 899	S. Hayes et al.	(CORN)	REFID=20511
STOTTLE...	71	Thesis ORO 2504 170	A.R. Stottlemyer	(UMD)	REFID=20512
BIZOT	70	PL 32B 416	J.C. Bizot et al.	(ORSAY)	REFID=20501
		Also Liverpool Sym. 69	J.P. Perez-y-Jorba		REFID=20502
EARLES	70	PRL 25 1312	D.R. Earles et al.	(NEAS)	REFID=20504
LINDSEY	66	PR 147 913	J.S. Lindsey, G. Smith	(LRL)	REFID=20481
LONDON	66	PR 143 1034	G.W. London et al.	(BNL, SYRA) IGJPC	REFID=11774
BADIER	65B	PL 17 337	J. Badier et al.	(EPOL, SACL, AMST)	REFID=20253
LINDSEY	65	PRL 15 221	J.S. Lindsey, G.A. Smith	(LRL)	REFID=20478
		LINDSEY 65 data included in LINDSEY 66.			
SCHLEIN	63	PRL 10 368	P.E. Schlein et al.	(UCLA) IGJP	REFID=20474

$h_1(1170)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

NODE=M030

 $h_1(1170)$ MASS

NODE=M030M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1170±20 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1168± 4	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1166± 5±3	¹ ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1190±60	² DANKOWY...	81	SPEC 0	8 $\pi p \rightarrow 3\pi n$
¹ Average and spread of values using 2 variants of the model of BOWLER 75.				
² Uses the model of BOWLER 75.				

NODE=M030M

→ UNCHECKED ←

OCCUR=2

NODE=M030M;LINKAGE=B
NODE=M030M;LINKAGE=C **$h_1(1170)$ WIDTH**

NODE=M030W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
360±40 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
345± 6	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
375± 6±34	³ ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
320±50	⁴ DANKOWY...	81	SPEC 0	8 $\pi p \rightarrow 3\pi n$
³ Average and spread of values using 2 variants of the model of BOWLER 75.				
⁴ Uses the model of BOWLER 75.				

NODE=M030W

→ UNCHECKED ←

OCCUR=2

NODE=M030W;LINKAGE=B
NODE=M030W;LINKAGE=C **$h_1(1170)$ DECAY MODES**

NODE=M030215;NODE=M030

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\pi$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

 $h_1(1170)$ BRANCHING RATIOS

NODE=M030220

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
seen	ATKINSON	84	OMEG	20-70 $\gamma p \rightarrow \pi^+ \pi^- \pi^0 p$
seen	DANKOWY...	81	SPEC	8 $\pi p \rightarrow 3\pi n$

NODE=M030R1
NODE=M030R1 **$h_1(1170)$ REFERENCES**

NODE=M030

ANDO	92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
DANKOWY...	81	PRL 46 580	J.A. Dankowych <i>et al.</i>	(TNTO, BNL, CARL+)
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)

REFID=43171
REFID=20574
REFID=20572
REFID=20571

$b_1(1235)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

NODE=M011


 $b_1(1235)$ MASS

NODE=M011M

NODE=M011M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1229.5 ± 3.2 OUR AVERAGE		Error includes scale factor of 1.6.			See the ideogram below.
1225 ± 5		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
1235 ± 15		ALDE 92C	GAM2		38,100 $\pi^- p \rightarrow \omega \pi^0 n$
1236 ± 16		FUKUI 91	SPEC		8.95 $\pi^- p \rightarrow \omega \pi^0 n$
1222 ± 6		ATKINSON 84E	OMEG ±		25-55 $\gamma p \rightarrow \omega \pi X$
1237 ± 7		ATKINSON 84E	OMEG 0		25-55 $\gamma p \rightarrow \omega \pi X$
1239 ± 5		EVANGELIS...	81 OMEG -		12 $\pi^- p \rightarrow \omega \pi p$
1251 ± 8	450	GESSAROLI 77	HBC -		11 $\pi^- p \rightarrow \pi^- \omega p$
1245 ± 11	890	FLATTE 76C	HBC -		4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
1222 ± 4	1400	CHALOUPKA 74	HBC -		3.9 $\pi^- p$
1220 ± 7	600	KARSHON 74B	HBC +		4.9 $\pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1190 ± 10		AUGUSTIN 89	DM2 ±		$e^+ e^- \rightarrow 5\pi$
1213 ± 5		ATKINSON 84C	OMEG 0		20-70 γp
1271 ± 11		COLLICK 84	SPEC +		200 $\pi^+ Z \rightarrow Z \pi \omega$

OCCUR=2

 **$b_1(1235)$ WIDTH**

NODE=M011W

NODE=M011W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
142 ± 9 OUR AVERAGE		Error includes scale factor of 1.2.			
113 ± 12		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
160 ± 30		ALDE 92C	GAM2		38,100 $\pi^- p \rightarrow \omega \pi^0 n$
151 ± 31		FUKUI 91	SPEC		8.95 $\pi^- p \rightarrow \omega \pi^0 n$
170 ± 15		EVANGELIS...	81 OMEG -		12 $\pi^- p \rightarrow \omega \pi p$
170 ± 50	225	BALTAY 78B	HBC +		15 $\pi^+ p \rightarrow p 4\pi$
155 ± 32	450	GESSAROLI 77	HBC -		11 $\pi^- p \rightarrow \pi^- \omega p$
182 ± 45	890	FLATTE 76C	HBC -		4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
135 ± 20	1400	CHALOUPKA 74	HBC -		3.9 $\pi^- p$
156 ± 22	600	KARSHON 74B	HBC +		4.9 $\pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
210 ± 19		AUGUSTIN 89	DM2 ±		$e^+ e^- \rightarrow 5\pi$
231 ± 14		ATKINSON 84C	OMEG 0		20-70 γp
232 ± 29		COLLICK 84	SPEC +		200 $\pi^+ Z \rightarrow Z \pi \omega$

$b_1(1235)$ DECAY MODES

NODE=M011215;NODE=M011

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\omega\pi$ [D/S amplitude ratio = 0.277 ± 0.027]	dominant	
Γ_2 $\pi^\pm\gamma$	$(1.6 \pm 0.4) \times 10^{-3}$	
Γ_3 $\eta\rho$	seen	
Γ_4 $\pi^+\pi^+\pi^-\pi^0$	< 50 %	84%
Γ_5 $K^*(892)^\pm K^\mp$	seen	
Γ_6 $(K\bar{K})^\pm\pi^0$	< 8 %	90%
Γ_7 $K_S^0 K_L^0 \pi^\pm$	< 6 %	90%
Γ_8 $K_S^0 K_S^0 \pi^\pm$	< 2 %	90%
Γ_9 $\phi\pi$	< 1.5 %	84%

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=9
DESIG=8;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=74
DESIG=71;OUR EST;→ UNCHECKED ←
DESIG=73;OUR EST;→ UNCHECKED ←
DESIG=72;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←

$b_1(1235)$ PARTIAL WIDTHS

NODE=M011220

$\Gamma(\pi^\pm\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2
	230±60	COLLICK	84	SPEC	+	200 $\pi^+ Z \rightarrow Z\pi\omega$

NODE=M011W3
NODE=M011W3


$b_1(1235)$ D-wave/S-wave AMPLITUDE RATIO IN DECAY OF $b_1(1235) \rightarrow \omega\pi$

NODE=M011DS

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.277±0.027 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.			
0.269±0.009±0.010		NOZAR	02	MPS	- 18 $\pi^- p \rightarrow \omega\pi^- p$
0.23 ±0.03		AMSLER	94C	CBAR	0.0 $\bar{p}p \rightarrow \omega\eta\pi^0$
0.45 ±0.04		AMSLER	93B	CBAR	0.0 $\bar{p}p \rightarrow \omega\pi^0\pi^0$
0.235±0.047		ATKINSON	84C	OMEG	20-70 γp
0.4 $\begin{smallmatrix} +0.1 \\ -0.1 \end{smallmatrix}$		GESSAROLI	77	HBC	- 11 $\pi^- p \rightarrow \pi^- \omega p$
0.21 ±0.08		CHUNG	75B	HBC	+ 7.1 $\pi^+ p$
0.3 ±0.1		CHALOUPKA	74	HBC	- 3.9-7.5 $\pi^- p$
0.35 ±0.25	600	KARSHON	74B	HBC	+ 4.9 $\pi^+ p$

NODE=M011DS

WEIGHTED AVERAGE
0.277±0.027 (Error scaled by 2.4)



$b_1(1235)$ D-wave/S-wave amplitude ratio in decay of $b_1(1235) \rightarrow \omega\pi$

$b_1(1235)$ D-wave/S-wave AMPLITUDE PHASE DIFFERENCE IN DECAY OF $b_1(1235) \rightarrow \omega\pi$

NODE=M011PH

VALUE (°)	DOCUMENT ID	TECN	CHG	COMMENT
10.5±2.4±3.9	NOZAR	02	MPS	- 18 $\pi^- p \rightarrow \omega\pi^- p$

NODE=M011PH

$b_1(1235)$ BRANCHING RATIOS

NODE=M011230

 $\Gamma(\eta\rho)/\Gamma(\omega\pi)$ Γ_3/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
<0.10	ATKINSON	84D	OMEG 20-70 γp

NODE=M011R9
NODE=M011R9 $\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(\omega\pi)$ Γ_4/Γ_1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.5	ABOLINS	63	HBC	+ 3.5 $\pi^+ p$

NODE=M011R1
NODE=M011R1 $\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	10E	BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

NODE=M011R10
NODE=M011R10¹ From a fit including ten additional resonances and energy-independent Breit-Wigner width.

NODE=M011R10;LINKAGE=AB

 $\Gamma((K\bar{K})^\pm \pi^0)/\Gamma(\omega\pi)$ Γ_6/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.08	90	BALTAY	67	HBC	\pm 0.0 $\bar{p} p$

NODE=M011R6
NODE=M011R6 $\Gamma(K_S^0 K_L^0 \pi^\pm)/\Gamma(\omega\pi)$ Γ_7/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.06	90	BALTAY	67	HBC	\pm 0.0 $\bar{p} p$

NODE=M011R8
NODE=M011R8 $\Gamma(K_S^0 K_S^0 \pi^\pm)/\Gamma(\omega\pi)$ Γ_8/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.02	90	BALTAY	67	HBC	\pm 0.0 $\bar{p} p$

NODE=M011R7
NODE=M011R7 $\Gamma(\phi\pi)/\Gamma(\omega\pi)$ Γ_9/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.004	95	VIKTOROV	96	SPEC	0 32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$

NODE=M011R4
NODE=M011R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	95	BIZZARRI	69	HBC	\pm 0.0 $\bar{p} p$
<0.015		DAHL	67	HBC	1.6-4.2 $\pi^- p$

 $b_1(1235)$ REFERENCES

NODE=M011

ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
NOZAR	02	PL B541 35	M. Nozar <i>et al.</i>		REFID=48850
VIKTOROV	96	PAN 59 1184	V.A. Viktorov <i>et al.</i>	(SERP)	REFID=45203
		Translated from YAF 59 1239.			
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
AMSLER	93B	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43602
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41859
FUKUI	91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41581
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
ATKINSON	84C	NP B243 1	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+) JP	REFID=20625
ATKINSON	84D	NP B242 269	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20623
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20624
COLLICK	84	PRL 53 2374	B. Collick <i>et al.</i>	(MINN, ROCH, FNAL)	REFID=20626
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+) JP	REFID=20230
FLATTE	76C	PL 64B 225	S.M. Flatte <i>et al.</i>	(CERN, AMST, NIJM+) JP	REFID=20615
CHUNG	75B	PR D11 2426	S.U. Chung <i>et al.</i>	(BNL, LBL, UCSC) JP	REFID=20613
CHALOUPKA	74	PL 51B 407	V. Chaloupka <i>et al.</i>	(CERN) JP	REFID=20611
KARSHON	74B	PR D10 3608	U. Karshon <i>et al.</i>	(REHO) JP	REFID=20612
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20171
BALTAY	67	PRL 18 93	C. Baltay <i>et al.</i>	(COLU)	REFID=20159
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL)	REFID=20321
ABOLINS	63	PRL 11 381	M.A. Abolins <i>et al.</i>	(UCSD)	REFID=20006

$a_1(1260)$

$$I^G(J^{PC}) = 1^-(1^{++})$$

See also our review under the $a_1(1260)$ in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M010

NODE=M010

NODE=M010M

NODE=M010M
→ UNCHECKED ←

 $a_1(1260)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1230±40 OUR ESTIMATE				
1255± 6⁺⁷₋₁₇	420k	ALEKSEEV	10	COMP 190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1243±12±20		¹ AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow \rho^0 \rho^\pm \pi^\mp \gamma$
1230-1270	6360	² LINK	07A	FOCS $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1203± 3		³ GOMEZ-DUM..04	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$
1330±24	90k	SALVINI	04	OBLX $\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
1331±10± 3	37k	⁴ ASNER	00	CLE2 10.6 $e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
1255± 7± 6	5904	⁵ ABREU	98G	DLPH $e^+ e^-$
1207± 5± 8	5904	⁶ ABREU	98G	DLPH $e^+ e^-$
1196± 4± 5	5904	^{7,8} ABREU	98G	DLPH $e^+ e^-$
1240±10		BARBERIS	98B	450 $pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$
1262± 9± 7		^{5,9} ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94$, $\tau \rightarrow 3\pi\nu$
1210± 7± 2		^{6,9} ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94$, $\tau \rightarrow 3\pi\nu$
1211± 7 ⁺⁵⁰ ₋₀		⁶ ALBRECHT	93C	ARG $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1121± 8		¹⁰ ANDO	92	SPEC $8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1242±37		¹¹ IVANOV	91	RVUE $\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1260±14		¹² IVANOV	91	RVUE $\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1250± 9		¹³ IVANOV	91	RVUE $\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1208±15		ARMSTRONG	90	OMEG $300.0 pp \rightarrow pp \pi^+ \pi^- \pi^0$
1220±15		¹⁴ ISGUR	89	RVUE $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1260±25		¹⁵ BOWLER	88	RVUE
1166±18±11		BAND	87	MAC $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1164±41±23		BAND	87	MAC $\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$
1250±40		¹⁴ TORNQVIST	87	RVUE
1046±11		ALBRECHT	86B	ARG $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1056±20±15		RUCKSTUHL	86	DLCO $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1194±14±10		SCHMIDKE	86	MRK2 $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1255±23		BELLINI	85	SPEC $40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1240±80		¹⁶ DANKOWY...	81	SPEC $8.45 \pi^- p \rightarrow n 3\pi$
1280±30		¹⁶ DAUM	81B	CNTR $63,94 \pi^- p \rightarrow p 3\pi$
1041±13		¹⁷ GAVILLET	77	HBC $4.2 K^- p \rightarrow \Sigma 3\pi$

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M010M;LINKAGE=AU

NODE=M010M;LINKAGE=LI

NODE=M010M;LINKAGE=GO

NODE=M010M;LINKAGE=B6

NODE=M010M;LINKAGE=KS

NODE=M010M;LINKAGE=IM

NODE=M010M;LINKAGE=A1

NODE=M010M;LINKAGE=F1

NODE=M010M;LINKAGE=X

NODE=M010M;LINKAGE=P

NODE=M010M;LINKAGE=I

NODE=M010M;LINKAGE=L

NODE=M010M;LINKAGE=M

NODE=M010M;LINKAGE=K

NODE=M010M;LINKAGE=G

NODE=M010M;LINKAGE=D

NODE=M010M;LINKAGE=F

¹ The $\rho^\pm \pi^\mp$ state can be also due to the $\pi(1300)$.

² Using the Breit-Wigner parameterization; strong correlation between mass and width.

³ Using the data of BARATE 98R.

⁴ From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.

⁵ Uses the model of KUHN 90.

⁶ Uses the model of ISGUR 89.

⁷ Includes the effect of a possible a_1' state.

⁸ Uses the model of FEINDT 90.

⁹ Supersedes AKERS 95P.

¹⁰ Average and spread of values using 2 variants of the model of BOWLER 75.

¹¹ Reanalysis of RUCKSTUHL 86.

¹² Reanalysis of SCHMIDKE 86.

¹³ Reanalysis of ALBRECHT 86B.

¹⁴ From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

¹⁵ From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

¹⁶ Uses the model of BOWLER 75.

¹⁷ Produced in K^- backward scattering.

 $a_1(1260)$ WIDTH

NODE=M010W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
250 to 600 OUR ESTIMATE				
367± 9⁺²⁸₋₂₅	420k	ALEKSEEV	10	COMP 190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$

NODE=M010W
→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

410± 31± 30		18	AUBERT	07AU	BABR	10.6	$e^+e^- \rightarrow \rho^0 \rho^\pm \pi^\mp \gamma$	
520-680	6360	19	LINK	07A	FOCS		$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$	
480± 20		20	GOMEZ-DUM..	04	RVUE		$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$	
580± 41	90k		SALVINI	04	OBLX		$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$	
460± 85	205	21	DRUTSKOY	02	BELL		$B \rightarrow D^{(*)} K^- K^{*0}$	
814± 36± 13	37k	22	ASNER	00	CLE2	10.6	$e^+e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$	
450± 50	22k	23	AKHMETSHIN	99E	CMD2	1.05-1.38	$e^+e^- \rightarrow$ $\pi^+ \pi^- \pi^0 \pi^0$	
570± 10		24	BONDAR	99	RVUE		$e^+e^- \rightarrow 4\pi, \tau \rightarrow 3\pi \nu_\tau$	
587± 27± 21	5904	25	ABREU	98G	DLPH		e^+e^-	OCCUR=2
478± 3± 15	5904	26	ABREU	98G	DLPH		e^+e^-	OCCUR=3
425± 14± 8	5904	27,28	ABREU	98G	DLPH		e^+e^-	
400± 35			BARBERIS	98B		450	$pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$	
621± 32± 58		25,29	ACKERSTAFF	97R	OPAL	$E_{cm}^{ee} = 88-94$,	$\tau \rightarrow 3\pi \nu$	
457± 15± 17		26,29	ACKERSTAFF	97R	OPAL	$E_{cm}^{ee} = 88-94$,	$\tau \rightarrow 3\pi \nu$	OCCUR=2
446± 21 ⁺¹⁴⁰ ₋₀		26	ALBRECHT	93C	ARG		$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
239± 11			ANDO	92	SPEC	8	$\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
266± 13± 4		30	ANDO	92	SPEC	8	$\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	OCCUR=3
465 ⁺²²⁸ ₋₁₄₃		31	IVANOV	91	RVUE		$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	
298 ⁺⁴⁰ ₋₃₄		32	IVANOV	91	RVUE		$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=2
488± 32		33	IVANOV	91	RVUE		$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=3
430± 50			ARMSTRONG	90	OMEG	300.0	$pp \rightarrow pp \pi^+ \pi^- \pi^0$	
420± 40		34	ISGUR	89	RVUE		$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
396± 43		35	BOWLER	88	RVUE			
405± 75± 25			BAND	87	MAC		$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
419± 108± 57			BAND	87	MAC		$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$	OCCUR=2
521± 27			ALBRECHT	86B	ARG		$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
476 ⁺¹³² ₋₁₂₀ ± 54			RUCKSTUHL	86	DLCO		$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
462± 56± 30			SCHMIDKE	86	MRK2		$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
292± 40			BELLINI	85	SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
380± 100		36	DANKOWY...	81	SPEC	8.45	$\pi^- p \rightarrow n 3\pi$	
300± 50		36	DAUM	81B	CNTR	63,94	$\pi^- p \rightarrow p 3\pi$	
230± 50		37	GAVILLET	77	HBC	4.2	$K^- p \rightarrow \Sigma 3\pi$	

18 The $\rho^\pm \pi^\mp$ state can be also due to the $\pi(1300)$.

19 Using the Breit-Wigner parameterization; strong correlation between mass and width.

20 Using the data of BARATE 98R.

21 From a fit of the $K^- K^{*0}$ distribution assuming $m_{a_1} = 1230$ MeV and purely resonant production of the $K^- K^{*0}$ system.

22 From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.

23 Using the $a_1(1260)$ mass of 1230 MeV.

24 From AKHMETSHIN 99E and ASNER 00 data using the $a_1(1260)$ mass of 1230 MeV.

25 Uses the model of KUHN 90.

26 Uses the model of ISGUR 89.

27 Includes the effect of a possible a_1' state.

28 Uses the model of FEINDT 90.

29 Supersedes AKERS 95P.

30 Average and spread of values using 2 variants of the model of BOWLER 75.

31 Reanalysis of RUCKSTUHL 86.

32 Reanalysis of SCHMIDKE 86.

33 Reanalysis of ALBRECHT 86B.

34 From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

35 From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

36 Uses the model of BOWLER 75.

37 Produced in K^- backward scattering.

NODE=M010W;LINKAGE=AU

NODE=M010W;LINKAGE=LI

NODE=M010W;LINKAGE=GO

NODE=M010W;LINKAGE=DR

NODE=M010W;LINKAGE=B6

NODE=M010W;LINKAGE=WE

NODE=M010W;LINKAGE=WB

NODE=M010W;LINKAGE=KS

NODE=M010W;LINKAGE=IM

NODE=M010W;LINKAGE=A1

NODE=M010W;LINKAGE=F1

NODE=M010W;LINKAGE=X

NODE=M010W;LINKAGE=P

NODE=M010W;LINKAGE=I

NODE=M010W;LINKAGE=L

NODE=M010W;LINKAGE=M

NODE=M010W;LINKAGE=K

NODE=M010W;LINKAGE=G

NODE=M010W;LINKAGE=D

NODE=M010W;LINKAGE=F

$a_1(1260)$ DECAY MODES

NODE=M010215;NODE=M010

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^+ \pi^- \pi^0$	
Γ_2 $\pi^0 \pi^0 \pi^0$	
Γ_3 $(\rho\pi)_{S\text{-wave}}$	seen
Γ_4 $(\rho\pi)_{D\text{-wave}}$	seen
Γ_5 $(\rho(1450)\pi)_{S\text{-wave}}$	seen
Γ_6 $(\rho(1450)\pi)_{D\text{-wave}}$	seen
Γ_7 $\sigma \pi$	seen
Γ_8 $f_0(980) \pi$	not seen
Γ_9 $f_0(1370) \pi$	seen
Γ_{10} $f_2(1270) \pi$	seen
Γ_{11} $K \bar{K}^*(892) + \text{c.c.}$	seen
Γ_{12} $\pi \gamma$	seen

DESIG=22
 DESIG=23
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=8;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←
 DESIG=10;OUR EST;→ UNCHECKED ←
 DESIG=16;OUR EST;→ UNCHECKED ←
 DESIG=11;OUR EST;→ UNCHECKED ←
 DESIG=12;OUR EST;→ UNCHECKED ←
 DESIG=13;OUR EST;→ UNCHECKED ←
 DESIG=14;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←

$a_1(1260)$ PARTIAL WIDTHS

NODE=M010220

$\Gamma(\pi\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_{12}
	640 ± 246	ZIELINSKI	84C SPEC	200 $\pi^+ Z \rightarrow Z 3\pi$	

NODE=M010W4
 NODE=M010W4

D-wave/S-wave AMPLITUDE RATIO IN DECAY OF $a_1(1260) \rightarrow \rho\pi$


NODE=M010DS

VALUE	DOCUMENT ID	TECN	COMMENT
-0.062 ± 0.020 OUR AVERAGE	Error includes scale factor of 2.3. See the ideogram below.		
$-0.043 \pm 0.009 \pm 0.005$	LINK	07A FOCS	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
$-0.14 \pm 0.04 \pm 0.07$	³⁸ CHUNG	02 B852	$18.3 \pi^- \rho \rightarrow \pi^+ \pi^- \pi^- \rho$
$-0.10 \pm 0.02 \pm 0.02$	^{39,40} ACKERSTAFF	97R OPAL	$E_{cm}^{e^+e^-} = 88-94, \tau \rightarrow 3\pi\nu$
-0.11 ± 0.02	³⁹ ALBRECHT	93C ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$

NODE=M010DS

³⁸ Deck-type background not subtracted.
³⁹ Uses the model of ISGUR 89.
⁴⁰ Supersedes AKERS 95P.

NODE=M010DS;LINKAGE=C
 NODE=M010DS;LINKAGE=IM
 NODE=M010DS;LINKAGE=X



D-wave/S-wave AMPLITUDE RATIO IN DECAY OF $a_1(1260) \rightarrow \rho\pi$

$a_1(1260)$ BRANCHING RATIOS

NODE=M010225

$\Gamma((\rho\pi)_{S\text{-wave}})/\Gamma_{\text{total}}$ Γ_3/Γ NODE=M010R5
NODE=M010R5

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
60.19	37k	41 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma((\rho\pi)_{D\text{-wave}})/\Gamma_{\text{total}}$ Γ_4/Γ NODE=M010R6
NODE=M010R6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.30 \pm 0.60 \pm 0.22$	37k	41 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma((\rho(1450)\pi)_{S\text{-wave}})/\Gamma_{\text{total}}$ Γ_5/Γ NODE=M010R7
NODE=M010R7

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.56 \pm 0.84 \pm 0.32$	37k	41,42 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma((\rho(1450)\pi)_{D\text{-wave}})/\Gamma_{\text{total}}$ Γ_6/Γ NODE=M010R8
NODE=M010R8

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2.04 \pm 1.20 \pm 0.28$	37k	41,42 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma(\sigma\pi)/\Gamma_{\text{total}}$ Γ_7/Γ NODE=M010R9
NODE=M010R9

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen		CHUNG	02	B852 18.3 $\pi^-p \rightarrow$ $\pi^+\pi^-\pi^-p$
$18.76 \pm 4.29 \pm 1.48$	37k	41,43 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma(f_0(980)\pi)/\Gamma_{\text{total}}$ Γ_8/Γ NODE=M010R10
NODE=M010R10

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	37k	ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma(f_0(1370)\pi)/\Gamma_{\text{total}}$ Γ_9/Γ NODE=M010R11
NODE=M010R11

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$7.40 \pm 2.71 \pm 1.26$	37k	41,44 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma(f_2(1270)\pi)/\Gamma_{\text{total}}$ Γ_{10}/Γ NODE=M010R12
NODE=M010R12

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.19 \pm 0.49 \pm 0.17$	37k	41,45 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

 $\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ NODE=M010R13
NODE=M010R13

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.2 ± 0.5	2255	46 COAN	04	CLEO $\tau^- \rightarrow K^-\pi^-K^+\nu_\tau$
8 to 15	205	47 DRUTSKOY	02	BELL $B \rightarrow D^{(*)}K^-K^{*0}$
$3.3 \pm 0.5 \pm 0.1$	37k	48 ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
2.6 ± 0.3		49 BARATE	99R	ALEP $\tau \rightarrow K\bar{K}\pi\nu_\tau$

$\Gamma(\sigma\pi)/\Gamma((\rho\pi)S\text{-wave})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_3
0.06 ± 0.05	90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$	
~ 0.3	28k	AKHMETSHIN	99E	CMD2 $1.05\text{--}1.38 e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$	
0.003 ± 0.003		⁵⁰ LONGACRE	82	RVUE	

NODE=M010R4
 NODE=M010R4

 $\Gamma(\pi^0 \pi^0 \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE	CL%	DOCUMENT ID	COMMENT	Γ_2/Γ_1
< 0.008	90	⁵¹ BARBERIS	01 450 $pp \rightarrow p_f 3\pi^0 p_s$	
<p>41 From a fit to the Dalitz plot.</p> <p>42 Assuming for $\rho(1450)$ mass and width of 1370 and 386 MeV respectively.</p> <p>43 Assuming for σ mass and width of 860 and 880 MeV respectively.</p> <p>44 Assuming for $f_0(1370)$ mass and width of 1186 and 350 MeV respectively.</p> <p>45 Assuming for $f_2(1270)$ mass and width of 1275 and 185 MeV respectively.</p> <p>46 Using structure functions from KUHN 92 and DECKER 93A and $B(\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau) = (0.155 \pm 0.006 \pm 0.009)\%$ from BRIERE 03.</p> <p>47 From a comparison to ALAM 94 assuming purely resonant production of the $K^- K^{*0}$ system.</p> <p>48 From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.</p> <p>49 Assuming $a_1(1260)$ dominance and taking $B(\tau \rightarrow a_1(1260)\nu_\tau)$ from BUSKULIC 96.</p> <p>50 Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from GAVILLET 77, DAUM 80, and DANKOWYCH 81.</p> <p>51 Inconsistent with observations of $\sigma\pi$, $f_0(1370)\pi$, and $f_2(1270)\pi$ decay modes.</p>				

NODE=M010R15
 NODE=M010R15

NODE=M010R;LINKAGE=B1
 NODE=M010R;LINKAGE=B2
 NODE=M010R;LINKAGE=B3
 NODE=M010R;LINKAGE=B4
 NODE=M010R;LINKAGE=B5
 NODE=M010R13;LINKAGE=CO

NODE=M010R;LINKAGE=DR

NODE=M010R;LINKAGE=B6
 NODE=M010R13;LINKAGE=BA
 NODE=M010R4;LINKAGE=E

NODE=M010R;LINKAGE=RB

 $a_1(1260)$ REFERENCES

ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
LINK	07A	PR D75 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51713
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
GOMEZ-DUMM...	04	PR D69 073002	D. Gomez Dumm, A. Pich, J. Portoles		REFID=49771
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)	REFID=53226
BRIERE	03	PRL 90 181802	R. A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=49360
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=48780
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>		REFID=48324
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
AKHMETSHIN	99E	PL B466 392	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47411
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47366
BONDAR	99	PL B466 403	A.E. Bondar <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47358
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=45909
BARATE	98R	EPJ C4 409	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=46147
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46345
ACKERSTAFF	97R	ZPHY C75 593	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45616
BUSKULIC	96	ZPHY C70 579	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44588
AKERS	95P	ZPHY C67 45	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44366
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i>	(CLEO Collab.)	REFID=43738
ALBRECHT	93C	ZPHY C58 61	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=43310
DECKER	93A	ZPHY C58 445	R. Decker <i>et al.</i>		REFID=51577
ANDO	92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)	REFID=43171
KUHN	92	ZPHY C56 661	J.H. Kuhn, E. Mirkes		REFID=51576
IVANOV	91	ZPHY C49 563	Y.P. Ivanov, A.A. Osipov, M.K. Volkov	(JINR)	REFID=41750
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch	(WA76 Coll.)	REFID=41375
FEINDT	90	ZPHY C48 681	M. Feindt	(HAMB)	REFID=45912
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ISGUR	89	PR D39 1357	N. Isgur, C. Morningstar, C. Reader	(TNTO)	REFID=40730
BOWLER	88	PL B209 99	M.G. Bowler	(OXF)	REFID=40578
BAND	87	PL B198 297	H.R. Band <i>et al.</i>	(MAC Collab.)	REFID=40263
TORNQVIST	87	ZPHY C36 695	N.A. Tornqvist	(HELS)	REFID=40030
ALBRECHT	86B	ZPHY C33 7	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=20884
RUCKSTUHL	86	PRL 56 2132	W. Ruckstuhl <i>et al.</i>	(DELCO Collab.)	REFID=10349
SCHMIDKE	86	PRL 57 527	W.B. Schmidke <i>et al.</i>	(Mark II Collab.)	REFID=10350
BELLINI	85	JNP 41 781	D. Bellini <i>et al.</i>		REFID=47490
Translated from YAF 41 1223.					
ZIELINSKI	84C	PRL 52 1195	M. Zielinski <i>et al.</i>	(ROCH, MINN, FNAL)	REFID=20882
LONGACRE	82	PR D26 82	R.S. Longacre	(BNL)	REFID=20878
DANKOWY...	81	PRL 46 580	J.A. Dankowych <i>et al.</i>	(TNTO, BNL, CARL+)	REFID=20572
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=20872
DAUM	80	PL 89B 281	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=20868
GAVILLET	77	PL 69B 119	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+)	REFID=20852
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)	REFID=20571

NODE=M010

$f_2(1270)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M005

 $f_2(1270)$ MASS

NODE=M005M

NODE=M005M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1275.5 ± 0.8 OUR AVERAGE				
1275.8 ± 1.0 ± 0.4		¹ BOGOLYUB...	13 SPEC	$7\pi^+(K^+,p)A \rightarrow n\gamma + X$
1262 $\pm \frac{1}{2} \pm 8$		ABLIKIM	06v BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1275 ± 15		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
1283 ± 5		ALDE	98 GAM4	$100\pi^-p \rightarrow \pi^0\pi^0n$
1278 ± 5		² BERTIN	97c OBLX	$0.0\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1272 ± 8	200k	PROKOSHKIN	94 GAM2	$38\pi^-p \rightarrow \pi^0\pi^0n$
1269.7 ± 5.2	5730	AUGUSTIN	89 DM2	$e^+e^- \rightarrow 5\pi$
1283 ± 8	400	³ ALDE	87 GAM4	$100\pi^-p \rightarrow 4\pi^0n$
1274 ± 5		³ AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1283 ± 6		⁴ LONGACRE	86 MPS	$22\pi^-p \rightarrow n2K_S^0$
1276 ± 7		COURAU	84 DLCO	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
1273.3 ± 2.3		⁵ CHABAUD	83 ASPK	$17\pi^-p$ polarized
1280 ± 4		⁶ CASON	82 STRC	$8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$
1281 ± 7	11600	GIDAL	81 MRK2	J/ψ decay
1282 ± 5		⁷ CORDEN	79 OMEG	$12-15\pi^-p \rightarrow n2\pi$
1269 ± 4	10k	APEL	75 NICE	$40\pi^-p \rightarrow n2\pi^0$
1272 ± 4	4600	ENGLER	74 DBC	$6\pi^+n \rightarrow \pi^+\pi^-p$
1277 ± 4	5300	FLATTE	71 HBC	$7.0\pi^+p$
1273 ± 8		³ STUNTEBECK	70 HBC	$8\pi^-p, 5.4\pi^+d$
1265 ± 8		BOESEBECK	68 HBC	$8\pi^+p$

••• We do not use the following data for averages, fits, limits, etc. •••

1259 ± 4 ± 4	1.7k	^{8,9} DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1267 ± 4 ± 3	1.5k	^{8,9} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1270 ± 8		¹⁰ ANISOVICH	09 RVUE	$0.0\bar{p}p, \pi N$
1277 ± 6	870	¹¹ SCHEGELSKY	06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
1251 ± 10		TIKHOMIROV	03 SPEC	$40.0\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$
1260 ± 10		¹² ALDE	97 GAM2	$450pp \rightarrow pp\pi^0\pi^0$
1278 ± 6		¹² GRYGOREV	96 SPEC	$40\pi^-N \rightarrow K_S^0 K_S^0 X$
1262 ± 11		AGUILAR...	91 EHS	$400pp$
1275 ± 10		AKER	91 CBAR	$0.0\bar{p}p \rightarrow 3\pi^0$
1220 ± 10		BREAKSTONE	90 SFM	$pp \rightarrow pp\pi^+\pi^-$
1288 ± 12		ABACHI	86B HRS	$e^+e^- \rightarrow \pi^+\pi^-X$
1284 ± 30	3k	BINON	83 GAM2	$38\pi^-p \rightarrow n2\eta$
1280 ± 20	3k	APEL	82 CNTR	$25\pi^-p \rightarrow n2\pi^0$
1284 ± 10	16000	DEUTSCH...	76 HBC	$16\pi^+p$
1258 ± 10	600	TAKAHASHI	72 HBC	$8\pi^-p \rightarrow n2\pi$
1275 ± 13		ARMENISE	70 HBC	$9\pi^+n \rightarrow p\pi^+\pi^-$
1261 ± 5	1960	³ ARMENISE	68 DBC	$5.1\pi^+n \rightarrow p\pi^+MM^-$
1270 ± 10	360	³ ARMENISE	68 DBC	$5.1\pi^+n \rightarrow p\pi^0MM$
1268 ± 6		¹³ JOHNSON	68 HBC	$3.7-4.2\pi^-p$

OCCUR=2

OCCUR=2

OCCUR=2

¹ Averaged over six nuclear targets, no statistically significant dependence from target nucleus observed.

² T-matrix pole.

³ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

⁵ From an energy-independent partial-wave analysis.

⁶ From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$.

⁷ From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^+\pi^-$ scattering data.

⁸ Using CLEO-c data but not authored by the CLEO Collaboration.

⁹ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 185$ MeV.

¹⁰ 4-poles, 5-channel K matrix fit.

¹¹ From analysis of L3 data at 91 and 183–209 GeV.

¹² Systematic uncertainties not estimated.

¹³ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

NODE=M005M;LINKAGE=B

NODE=M005M;LINKAGE=A

NODE=M005M;LINKAGE=T

NODE=M005M;LINKAGE=L

NODE=M005M;LINKAGE=O

NODE=M005M;LINKAGE=P

NODE=M005M;LINKAGE=S

NODE=M005M;LINKAGE=C

NODE=M005M;LINKAGE=D

NODE=M005M;LINKAGE=AN

NODE=M005M;LINKAGE=SC

NODE=M005M;LINKAGE=QQ

NODE=M005M;LINKAGE=J

$f_2(1270)$ WIDTH

NODE=M005W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W

186.7^{+2.2}_{-2.5} OUR FIT Error includes scale factor of 1.4.

185.9^{+2.8}_{-2.1} OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

190.3 ± 1.9 ± 1.8		¹ BOGOLYUB...	13	SPEC	$7\pi^+(K^+,p)A \rightarrow n\gamma + X$	
175 ⁺⁶ ₋₄ ± 10		ABLIKIM	06v	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
190 ± 20		ABLIKIM	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$	
171 ± 10		ALDE	98	GAM4	$100\pi^-p \rightarrow \pi^0\pi^0n$	
204 ± 20		² BERTIN	97c	OBLX	$0.0\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
192 ± 5	200k	PROKOSHKIN	94	GAM2	$38\pi^-p \rightarrow \pi^0\pi^0n$	
180 ± 24		AGUILAR-...	91	EHS	400 pp	
169 ± 9	5730	³ AUGUSTIN	89	DM2	$e^+e^- \rightarrow 5\pi$	
150 ± 30	400	³ ALDE	87	GAM4	$100\pi^-p \rightarrow 4\pi^0n$	
186 ⁺⁹ ₋₂		⁴ LONGACRE	86	MPS	$22\pi^-p \rightarrow n2K_S^0$	
179.2 ^{+6.9} _{-6.6}		⁵ CHABAUD	83	ASPK	17 π^-p polarized	
160 ± 11		DENNEY	83	LASS	10 π^+N	
196 ± 10	3k	APEL	82	CNTR	$25\pi^-p \rightarrow n2\pi^0$	
152 ± 9		⁶ CASON	82	STRC	$8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$	
186 ± 27	11600	GIDAL	81	MRK2	J/ψ decay	
216 ± 13		⁷ CORDEN	79	OMEG	$12-15\pi^-p \rightarrow n2\pi$	
190 ± 10	10k	APEL	75	NICE	$40\pi^-p \rightarrow n2\pi^0$	
192 ± 16	4600	ENGLER	74	DBC	$6\pi^+n \rightarrow \pi^+\pi^-p$	
183 ± 15	5300	FLATTE	71	HBC	$7\pi^+p \rightarrow \Delta^{++}f_2$	
196 ± 30		³ STUNTEBECK	70	HBC	$8\pi^-p, 5.4\pi^+d$	
216 ± 20	1960	³ ARMENISE	68	DBC	$5.1\pi^+n \rightarrow p\pi^+MM^-$	OCCUR=2
128 ± 27		³ BOESEBECK	68	HBC	$8\pi^+p$	
176 ± 21		^{3,8} JOHNSON	68	HBC	$3.7-4.2\pi^-p$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
194 ± 36		⁹ ANISOVICH	09	RVUE	$0.0\bar{p}p, \pi N$	
195 ± 15	870	¹⁰ SCHEGELSKY	06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
121 ± 26		TIKHOMIROV	03	SPEC	$40.0\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$	
187 ± 20		¹¹ ALDE	97	GAM2	$450pp \rightarrow pp\pi^0\pi^0$	
184 ± 10		¹¹ GRYGOREV	96	SPEC	$40\pi^-N \rightarrow K_S^0 K_S^0 X$	
200 ± 10		AKER	91	CBAR	$0.0\bar{p}p \rightarrow 3\pi^0$	
240 ± 40	3k	BINON	83	GAM2	$38\pi^-p \rightarrow n2\eta$	
187 ± 30	650	³ ANTIPOV	77	CIBS	$25\pi^-p \rightarrow p3\pi$	
225 ± 38	16000	DEUTSCH...	76	HBC	$16\pi^+p$	
166 ± 28	600	³ TAKAHASHI	72	HBC	$8\pi^-p \rightarrow n2\pi$	
173 ± 53		³ ARMENISE	70	HBC	$9\pi^+n \rightarrow p\pi^+\pi^-$	OCCUR=2

¹ Averaged over six nuclear targets, no statistically significant dependence from target nucleus observed.

² T-matrix pole.

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

⁵ From an energy-independent partial-wave analysis.

⁶ From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$.

⁷ From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^+\pi^-$ scattering data.

⁸ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

⁹ 4-poles, 5-channel K matrix fit.

¹⁰ From analysis of L3 data at 91 and 183–209 GeV.

¹¹ Systematic uncertainties not estimated.

NODE=M005W;LINKAGE=C

NODE=M005W;LINKAGE=QA

NODE=M005W;LINKAGE=T

NODE=M005W;LINKAGE=L

NODE=M005W;LINKAGE=R

NODE=M005W;LINKAGE=Q


NODE=M005W;LINKAGE=U

NODE=M005W;LINKAGE=J

NODE=M005W;LINKAGE=AN

NODE=M005W;LINKAGE=SC

NODE=M005W;LINKAGE=QQ



$f_2(1270)$ DECAY MODES

NODE=M005215;NODE=M005

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 $\pi\pi$	(84.2 $^{+2.9}_{-0.9}$) %	S=1.1	DESIG=1
Γ_2 $\pi^+\pi^-2\pi^0$	(7.7 $^{+1.1}_{-3.2}$) %	S=1.2	DESIG=3
Γ_3 $K\bar{K}$	(4.6 $^{+0.5}_{-0.4}$) %	S=2.7	DESIG=4
Γ_4 $2\pi^+2\pi^-$	(2.8 ± 0.4) %	S=1.2	DESIG=2
Γ_5 $\eta\eta$	(4.0 ± 0.8) $\times 10^{-3}$	S=2.1	DESIG=7
Γ_6 $4\pi^0$	(3.0 ± 1.0) $\times 10^{-3}$		DESIG=9
Γ_7 $\gamma\gamma$	(1.42 ± 0.24) $\times 10^{-5}$	S=1.4	DESIG=8
Γ_8 $\eta\pi\pi$	< 8 $\times 10^{-3}$	CL=95%	DESIG=6
Γ_9 $K^0K^-\pi^+$ + c.c.	< 3.4 $\times 10^{-3}$	CL=95%	DESIG=5
Γ_{10} e^+e^-	< 6 $\times 10^{-10}$	CL=90%	DESIG=10

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 45 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 83.0$ for 38 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-90						
x_3	10	-39					
x_4	10	-38	1				
x_5	1	-6	0	0			
x_6	0	-7	0	0	0		
x_7	3	1	-15	0	0	0	
Γ	-71	65	-10	-7	-1	0	-6
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode	Rate (MeV)	Scale factor
Γ_1 $\pi\pi$	157.2 $^{+4.0}_{-1.1}$	
Γ_2 $\pi^+\pi^-2\pi^0$	14.4 $^{+2.1}_{-6.0}$	1.2
Γ_3 $K\bar{K}$	8.5 ± 0.8	2.8
Γ_4 $2\pi^+2\pi^-$	5.2 ± 0.7	1.2
Γ_5 $\eta\eta$	0.75 ± 0.14	2.1
Γ_6 $4\pi^0$	0.56 ± 0.19	
Γ_7 $\gamma\gamma$	0.0026 ± 0.0005	1.4

$f_2(1270)$ PARTIAL WIDTHS

NODE=M005220

$\Gamma(\pi\pi)$

 Γ_1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W1
NODE=M005W1

157.2 $^{+4.0}_{-1.1}$ OUR FIT

157.0 $^{+6.0}_{-1.0}$ ¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

152 ± 8 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(K\bar{K})$

 Γ_3

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W4
NODE=M005W4

8.5 ± 0.8 OUR FIT Error includes scale factor of 2.8.

9.0 $^{+0.7}_{-0.3}$ ¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.5 ± 2.0 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(\eta\eta)$

 Γ_5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W7
NODE=M005W7

0.75 ± 0.14 OUR FIT Error includes scale factor of 2.1.

1.0 ± 0.1 ¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 0.4 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(\gamma\gamma)$

 Γ_7

The value of this width depends on the theoretical model used. Unitary approaches with scalars typically (with exception of PENNINGTON 08) give values clustering around 2.6 keV; without an *S*-wave contribution, values are systematically higher (typically around 3 keV).

NODE=M005W8
NODE=M005W8

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W8

2.6 ± 0.5 OUR FIT Error includes scale factor of 1.4.

2.93 ± 0.40 ³ DAI 14A RVUE Compilation

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.14±0.20		4,5	PENNINGTON 08	RVUE	Compilation		OCCUR=2
3.82±0.30		5,6	PENNINGTON 08	RVUE	Compilation		OCCUR=3
2.55±0.15	870	2	SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$		
2.84±0.35			BOGLIONE 99	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$		
2.93±0.23±0.32		7	YABUKI 95	VNS			
2.58±0.13 ^{+0.36} _{-0.27}		8	BEHREND 92	CELL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
3.10±0.35±0.35		9	BLINOV 92	MD1	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
2.27±0.47±0.11			ADACHI 90D	TOPZ	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
3.15±0.04±0.39			BOYER 90	MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
3.19±0.16 ^{+0.29} _{-0.28}			MARSISKE 90	CBAL	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$		
2.35±0.65		10	MORGAN 90	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$		
3.19±0.09 ^{+0.22} _{-0.38}	2177		OEST 90	JADE	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$		
3.2 ±0.1 ±0.4		11	AIHARA 86B	TPC	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
2.5 ±0.1 ±0.5			BEHREND 84B	CELL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
2.85±0.25±0.5		12	BERGER 84	PLUT	$e^+e^- \rightarrow e^+e^-2\pi$		
2.70±0.05±0.20			COURAU 84	DLCO	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
2.52±0.13±0.38		13	SMITH 84C	MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
2.7 ±0.2 ±0.6			EDWARDS 82F	CBAL	$e^+e^- \rightarrow e^+e^-2\pi^0$		
2.9 ^{+0.6} _{-0.4} ±0.6		14	EDWARDS 82F	CBAL	$e^+e^- \rightarrow e^+e^-2\pi^0$		OCCUR=2
3.2 ±0.2 ±0.6			BRANDELIK 81B	TASS	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
3.6 ±0.3 ±0.5			ROUSSARIE 81	MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$		
2.3 ±0.8		15	BERGER 80B	PLUT	e^+e^-		

$\Gamma(e^+e^-)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{10}
<0.11	90	ACHASOV 00K	SND	$e^+e^- \rightarrow \pi^0\pi^0$	NODE=M005W9 NODE=M005W9

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7	90	VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^0\pi^0$	
------	----	-------------	----	---------------------------------	--

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

³ Based on a K-matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis. Supersedes PENNINGTON 08.

⁴ Solution A (preferred solution based on χ^2 -analysis).

⁵ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

⁶ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

⁷ With a narrow scalar state around 1220 MeV.

⁸ Using a unitarized model with a 300 - 500 keV wide scalar at 1100 MeV.

⁹ Using the unitarized model of LYTH 85.

¹⁰ Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with $\gamma\gamma$ width of $f_0(1370) : \Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$ KeV.

¹¹ Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes 2.66 ± 0.21 in the calculation of LANDRO 86.

¹² Using the MENNESSIER 83 model.

¹³ Superseded by BOYER 90.

¹⁴ If helicity = 2 assumption is not made.

¹⁵ Using mass, width and $B(f_2(1270) \rightarrow 2\pi)$ from PDG 78.

NODE=M005PW;LINKAGE=L
NODE=M005W1;LINKAGE=SC
NODE=M005W8;LINKAGE=A

NODE=M005W8;LINKAGE=P1
NODE=M005W8;LINKAGE=P3

NODE=M005W8;LINKAGE=P2

NODE=M005W8;LINKAGE=YA
NODE=M005W;LINKAGE=B
NODE=M005W;LINKAGE=A
NODE=M005PW;LINKAGE=C

NODE=M005PW;LINKAGE=B

NODE=M005PW;LINKAGE=X
NODE=M005PW;LINKAGE=V
NODE=M005PW;LINKAGE=H
NODE=M005PW;LINKAGE=A

$f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_3\Gamma_7/\Gamma$
0.121±0.020 OUR FIT			Error includes scale factor of 1.3.	NODE=M005G1 NODE=M005G1
0.091±0.007±0.027	¹ ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.104±0.007±0.072	² ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$	OCCUR=2
-------------------	---------------------------	-----	-----------------------------------	---------

¹ Using an incoherent background.

² Using a coherent background.

OCCUR=2

NODE=M005G1;LINKAGE=A
NODE=M005G1;LINKAGE=K

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_7/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$11.5^{+1.8+4.5}_{-2.0-3.7}$	¹ UEHARA	10A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

NODE=M005G02
 NODE=M005G02

¹ Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_0(Y)$.

NODE=M005G02;LINKAGE=UE

Helicity-0/Helicity-2 RATIO IN $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi\pi$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$3.7 \pm 0.3^{+15.9}_{-2.9}$	UEHARA	08A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$

NODE=M005HR0
 NODE=M005HR0

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5 ± 1.8	¹ DAI	14A	RVUE	Compilation
13	^{2,3} PENNINGTON	08	RVUE	Compilation
26	^{3,4} PENNINGTON	08	RVUE	Compilation

OCCUR=2
 OCCUR=3

¹ Based on a K -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis.

NODE=M005HR0;LINKAGE=A

² Solution A (preferred solution based on χ^2 -analysis).

NODE=M005HR0;LINKAGE=P1
 NODE=M005HR0;LINKAGE=P3

³ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

⁴ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

NODE=M005HR0;LINKAGE=P2

 $f_2(1270)$ BRANCHING RATIOS

NODE=M005225

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R10
 NODE=M005R10

$0.842^{+0.029}_{-0.009}$ OUR FIT Error includes scale factor of 1.1.

0.837 ± 0.020 OUR AVERAGE

0.849 ± 0.025		CHABAUD	83	ASPK	17 $\pi^- p$ polarized
0.85 ± 0.05	250	BEAUPRE	71	HBC	8 $\pi^+ p \rightarrow \Delta^{++} f_2$
0.8 ± 0.04	600	OH	70	HBC	1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$

 $\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(\pi\pi)$ Γ_2/Γ_1

Should be twice $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$ if decay is $\rho\rho$. (See ASCOLI 68D.)

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R2
 NODE=M005R2
 NODE=M005R2

$0.091^{+0.014}_{-0.040}$ OUR FIT Error includes scale factor of 1.2.

0.15 ± 0.06 600 EISENBERG 74 HBC 4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 EMMS 75D DBC 4 $\pi^+ n \rightarrow p f_2$

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_3/Γ_1

We average only experiments which either take into account $f_2(1270)$ - $a_2(1320)$ interference explicitly or demonstrate that $a_2(1320)$ production is negligible.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R3
 NODE=M005R3
 NODE=M005R3

$0.054^{+0.005}_{-0.006}$ OUR FIT Error includes scale factor of 2.7.

$0.041^{+0.004}_{-0.005}$ OUR AVERAGE

0.045 ± 0.01		¹ BARGIOTTI	03	OBLX	$\bar{p}p$
$0.037^{+0.008}_{-0.021}$		ETKIN	82B	MPS	23 $\pi^- p \rightarrow n2K_S^0$
0.045 ± 0.009		CHABAUD	81	ASPK	17 $\pi^- p$ polarized
0.039 ± 0.008		LOVERRE	80	HBC	4 $\pi^- p \rightarrow K\bar{K}N$
0.052 ± 0.025		ABLIKIM	04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
0.036 ± 0.005		² COSTA...	80	OMEG	1-2.2 $\pi^- p \rightarrow K^+ K^- n$
0.030 ± 0.005		³ MARTIN	79	RVUE	
0.027 ± 0.009		⁴ POLYCHRO...	79	STRC	7 $\pi^- p \rightarrow n2K_S^0$
0.025 ± 0.015		EMMS	75D	DBC	4 $\pi^+ n \rightarrow p f_2$
0.031 ± 0.012	20	ADERHOLZ	69	HBC	8 $\pi^+ p \rightarrow K^+ K^- \pi^+ p$

$\Gamma(2\pi^+2\pi^-)/\Gamma(\pi\pi)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.033±0.005 OUR FIT	Error	includes scale factor of 1.2.		
0.033±0.004 OUR AVERAGE	Error	includes scale factor of 1.1.		
0.024±0.006	160	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$
0.051±0.025	70	EISENBERG	74 HBC	$4.9 \pi^+ p \rightarrow \Delta^{++} f_2$
0.043 ^{+0.007} _{-0.011}	285	LOUIE	74 HBC	$3.9 \pi^- p \rightarrow n f_2$
0.037±0.007	154	ANDERSON	73 DBC	$6 \pi^+ n \rightarrow p f_2$
0.047±0.013		OH	70 HBC	$1.26 \pi^- p \rightarrow \pi^+ \pi^- n$

NODE=M005R1
 NODE=M005R1

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.0±0.8 OUR FIT	Error	includes scale factor of 2.1.	
2.9±0.5 OUR AVERAGE			
2.7±0.7	BINON	05 GAMS	$33 \pi^- p \rightarrow \eta\eta n$
2.8±0.7	ALDE	86D GAM4	$100 \pi^- p \rightarrow 2\eta n$
5.2±1.7	BINON	83 GAM2	$38 \pi^- p \rightarrow 2\eta n$

NODE=M005R7
 NODE=M005R7

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$ Γ_5/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.003±0.001		BARBERIS	00E	$450 pp \rightarrow p_f \eta \eta p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.05	95	EDWARDS	82F CBAL	$e^+ e^- \rightarrow e^+ e^- 2\eta$
<0.016	95	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$
<0.09	95	EISENBERG	74 HBC	$4.9 \pi^+ p \rightarrow \Delta^{++} f_2$

NODE=M005R6
 NODE=M005R6

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0030±0.0010 OUR FIT				
0.003 ±0.001	400 ± 50	ALDE	87 GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$

NODE=M005R11
 NODE=M005R11

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.57±0.01 ^{+1.39} _{-0.14}	UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$

NODE=M005R13
 NODE=M005R13

 $\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$ Γ_8/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	95	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$

NODE=M005R5
 NODE=M005R5

 $\Gamma(K^0 K^- \pi^+ + \text{c.c.})/\Gamma(\pi\pi)$ Γ_9/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.004	95	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$

NODE=M005R4
 NODE=M005R4

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-10})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	ACHASOV	00K SND	$e^+ e^- \rightarrow \pi^0 \pi^0$

NODE=M005R12
 NODE=M005R12

¹ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

² Re-evaluated by CHABAUD 83.

³ Includes PAWLICKI 77 data.

⁴ Takes into account the $f_2(1270)$ - $f_2'(1525)$ interference.

NODE=M005R;LINKAGE=BG
 NODE=M005R3;LINKAGE=D
 NODE=M005R3;LINKAGE=F
 NODE=M005R3;LINKAGE=M

 $f_2(1270)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
BOGOLYUB...	13	PAN 76 1324	M.Yu. Bogolyubsky <i>et al.</i>	(HYPERON-M Collab.)
		Translated from YAF 76 1389.		
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
PENNINGTON	08	EPJ C56 1	M.R. Pennington <i>et al.</i>	
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		

NODE=M005

REFID=56805
 REFID=55923
 REFID=55585

REFID=55592
 REFID=53641
 REFID=52719
 REFID=52761
 REFID=52166
 REFID=52303
 REFID=52309
 REFID=51652
 REFID=51507
 REFID=51185
 REFID=50450
 REFID=50780

ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860			
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446			
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
GRYGOREV	96	PAN 59 2105	V.K. Grigoriev, O.N. Baloshin, B.P. Barkov	(ITEP)	REFID=45566
		Translated from YAF 59 2187			
YABUKI	95	JPSJ 64 435	F. Yabuki <i>et al.</i>	(VENUS Collab.)	REFID=46384
PROKOSHKIN	94	SPD 39 420	Y.D. Prokoshkin, A.A. Kondashov	(SERP)	REFID=44094
		Translated from DANS 336 613			
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
BLINOV	92	ZPHY C53 33	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41858
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)	REFID=41587
ADACHI	90D	PL B234 185	I. Adachi <i>et al.</i>	(TOPAZ Collab.)	REFID=41345
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436			
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)	REFID=20394
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=20764
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LANDRO	86	PL B172 445	M. Landro, K.J. Mork, H.A. Olsen	(UTRO)	REFID=20767
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
LYTH	85	JP G11 459	D.H. Lyth		REFID=42169
BEHREND	84B	ZPHY C23 223	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20757
BERGER	84	ZPHY C26 199	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20760
COURAU	84	PL 147B 227	A. Courau <i>et al.</i>	(CIT, SLAC)	REFID=20758
SMITH	84C	PR D30 851	J.R. Smith <i>et al.</i>	(SLAC, LBL, HARV)	REFID=20759
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
Also		SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20751
		Translated from YAF 38 934			
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
MENNESSIER	83	ZPHY C16 241	G. Mennessier	(MONP)	REFID=20393
APEL	82	NP B201 197	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20745
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BRANDELIK	81B	ZPHY C10 117	R. Brandelik <i>et al.</i>	(TASSO Collab.)	REFID=20741
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ROUSSARIE	81	PL 105B 304	A. Roussarie <i>et al.</i>	(SLAC, LBL)	REFID=20388
BERGER	80B	PL 94B 254	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20736
COSTA...	80	NP B175 402	G. Costa de Beaugregard <i>et al.</i>	(BARI, BONN+)	REFID=20737
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+)	REFID=20382
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
PDG	78	PL 75B 1	C. Bricman <i>et al.</i>		REFID=40124
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL)	REFID=20367
DEUTSCH...	76	NP B103 426	M. Deuschmann <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20119
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20720
EMMS	75D	NP B96 155	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL)	REFID=20721
EISENBERG	74	PL 52B 239	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=20715
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
LOUIE	74	PL 48B 385	J. Louie <i>et al.</i>	(SACL, CERN)	REFID=20719
ANDERSON	73	PRL 31 562	J.C. Anderson <i>et al.</i>	(CMU, CASE)	REFID=20710
TAKAHASHI	72	PR D6 1266	K. Takahashi <i>et al.</i>	(TOHOK, PENN, NDAM+)	REFID=20103
BEAUPRE	71	NP B28 77	J.V. Beaupre <i>et al.</i>	(AACH, BERL, CERN)	REFID=20698
FLATTE	71	PL 34B 551	S.M. Flatte <i>et al.</i>	(LBL)	REFID=20700
ARMENISE	70	LCN 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
OH	70	PR D1 2494	B.Y. Oh <i>et al.</i>	(WISC, TNTO) JP	REFID=20335
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
ASCOLI	68D	PRL 21 1712	G. Ascoli <i>et al.</i>	(ILL)	REFID=20681
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
JOHNSON	68	PR 176 1651	P.B. Johnson <i>et al.</i>	(NDAM, PURD, SLAC)	REFID=20065
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
DERADO	65	PRL 14 872	I. Derado <i>et al.</i>	(NDAM)	REFID=20668
LEE	64	PRL 12 342	Y.Y. Lee <i>et al.</i>	(MICH)	REFID=20663
BONDAR	63	PL 5 153	L. Bondar <i>et al.</i>	(AACH, BIRM, BONN, DESY+)	REFID=20657

$f_1(1285)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

NODE=M008

 $f_1(1285)$ MASS

NODE=M008M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1282.0 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below. [1281.9 ± 0.5 MeV OUR 2015 AVERAGE Scale factor = 1.8]		
1287.4 ± 3.0	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
1281.16 ± 0.39 ± 0.45		¹ LEES	12X BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$
1285.1 ± 1.0 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.6 \\ 0.3 \end{smallmatrix}$		² ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
1281 ± 2 ± 1		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
1276.1 ± 8.1 ± 8.0	203	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
1274 ± 6	237	ABDALLAH	03H DLPH	91.2 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
1280 ± 4		ACCIARRI	01G L3	
1288 ± 4 ± 5	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1284 ± 6	1400	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta\pi^0\pi^0 n$
1281 ± 1		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
1281 ± 1		BARBERIS	97C OMEG	450 $pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
1280 ± 2		³ ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
1282.2 ± 1.5		LEE	94 MPS2	18 $\pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
1279 ± 5		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
1278 ± 2	140	ARMSTRONG	89 OMEG	300 $pp \rightarrow K \bar{K} \pi pp$
1278 ± 2		ARMSTRONG	89G OMEG	85 $\pi^+ p \rightarrow 4\pi pp, pp \rightarrow 4\pi pp$
1280.1 ± 2.1	60	RATH	89 MPS	21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1285 ± 1	4750	⁴ BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1280 ± 1	504	BITYUKOV	88 SPEC	32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1280 ± 4		ANDO	86 SPEC	8 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
1277 ± 2	420	REEVES	86 SPEC	6.6 $p\bar{p} \rightarrow K K \pi X$
1285 ± 2		CHUNG	85 SPEC	8 $\pi^- p \rightarrow N K \bar{K} \pi$
1279 ± 2	604	ARMSTRONG	84 OMEG	85 $\pi^+ p \rightarrow K \bar{K} \pi \pi p, pp \rightarrow K \bar{K} \pi pp$
1286 ± 1		CHAUVAT	84 SPEC	ISR 31.5 pp
1278 ± 4		EVANGELIS...	81 OMEG	12 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
1283 ± 3	103	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
1282 ± 2	320	NACASCH	78 HBC	0.7,0.76 $\bar{p}p \rightarrow K \bar{K} 3\pi$
1279 ± 5	210	GRASSLER	77 HBC	16 $\pi^\mp p$
1286 ± 3	180	DUBOC	72 HBC	1.2 $\bar{p}p \rightarrow 2K4\pi$
1283 ± 5		DAHL	67 HBC	1.6-4.2 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1284.2 ± 2.2		⁵ AAIJ	14Y LHCB	$\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+\pi^-)$
1281.9 ± 0.5		⁵ SOSA	99 SPEC	$pp \rightarrow p_{slow} (K_S^0 K^+ \pi^-) p_{fast}$
1282.8 ± 0.6		⁵ SOSA	99 SPEC	$pp \rightarrow p_{slow} (K_S^0 K^- \pi^+) p_{fast}$
1270 ± 10		AMELIN	95 VES	37 $\pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
1280 ± 2		ABATZIS	94 OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
1282 ± 4		ARMSTRONG	93C E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1270 ± 6 ± 10		ARMSTRONG	92C OMEG	300 $pp \rightarrow pp\pi^+\pi^-\gamma$
1281 ± 1		ARMSTRONG	89E OMEG	300 $pp \rightarrow pp2(\pi^+\pi^-)$
1279 ± 6 ± 10	16	BECKER	87 MRK3	$e^+e^- \rightarrow \phi K \bar{K} \pi$
1286 ± 9		GIDAL	87 MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
1287 ± 5	353	BITYUKOV	84B SPEC	32 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
~ 1279		⁶ TORNQVIST	82B RVUE	
1275 ± 6	31	BROMBERG	80 SPEC	100 $\pi^- p \rightarrow K \bar{K} \pi X$
1288 ± 9	200	GURTU	79 HBC	4.2 $K^- p \rightarrow n\eta 2\pi$

NODE=M008M

NODE=M008M
NEW


OCCUR=2

~ 1275.0	46	⁷ STANTON	79	CNTR	$8.5 \pi^- p \rightarrow n2\gamma2\pi$
1271 ± 10	34	CORDEN	78	OMEG	$12-15 \pi^- p \rightarrow K^+ K^- \pi n$
1295 ± 12	85	CORDEN	78	OMEG	$12-15 \pi^- p \rightarrow n5\pi$
1292 ± 10	150	DEFOIX	72	HBC	$0.7 \bar{p} p \rightarrow 7\pi$
1280 ± 3	500	⁸ THUN	72	MMS	$13.4 \pi^- p$
1303 ± 8		BARDADIN-...	71	HBC	$8 \pi^+ p \rightarrow p6\pi$
1283 ± 6		BOESEBECK	71	HBC	$16.0 \pi p \rightarrow p5\pi$
1270 ± 10		CAMPBELL	69	DBC	$2.7 \pi^+ d$
1285 ± 7		LORSTAD	69	HBC	$0.7 \bar{p} p, 4,5\text{-body}$
1290 ± 7		D'ANDLAU	68	HBC	$1.2 \bar{p} p, 5-6 \text{ body}$

OCCUR=2

- 1 Using the $2\pi^+ 2\pi^-$ and $\pi^+ \pi^- \eta$ modes of $f_1(1285)$ decay.
- 2 The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.
- 3 Supersedes ABATZIS 94, ARMSTRONG 89E.
- 4 From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.
- 5 No systematic error given.
- 6 From a unitarized quark-model calculation.
- 7 From phase shift analysis of $\eta \pi^+ \pi^-$ system.
- 8 Seen in the missing mass spectrum.

NODE=M008M;LINKAGE=LE
 NODE=M008M;LINKAGE=BL
 NODE=M008M;LINKAGE=B
 NODE=M008M;LINKAGE=A
 NODE=M008M;LINKAGE=N1
 NODE=M008M;LINKAGE=T
 NODE=M008M;LINKAGE=P
 NODE=M008M;LINKAGE=S



$f_1(1285)$ WIDTH

NODE=M008W

Only experiments giving width error less than 20 MeV are kept for averaging.

NODE=M008W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
24.1 ± 1.0 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
[24.2 ± 1.1 MeV OUR 2015 AVERAGE Scale factor = 1.3]				
18.3 ± 6.3	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
22.0 ± 3.1 ⁺ _{-1.5}		¹ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
35 ± 6 ± 4		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
40.0 ± 8.6 ± 9.3	203	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
29 ± 12	237	ABDALLAH	03H DLPH	91.2 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
45 ± 9 ± 7	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
55 ± 18	1400	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta\pi^0\pi^0 n$
24 ± 3		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
20 ± 2		BARBERIS	97C OMEG	450 $pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
36 ± 5		² ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
29.0 ± 4.1		LEE	94 MPS2	18 $\pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
25 ± 4	140	ARMSTRONG	89 OMEG	300 $pp \rightarrow K \bar{K} \pi p p$
22 ± 2	4750	³ BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
25 ± 4	504	BITYUKOV	88 SPEC	32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
19 ± 5		ANDO	86 SPEC	8 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
32 ± 8	420	REEVES	86 SPEC	6.6 $p\bar{p} \rightarrow K K \pi X$
22 ± 2		CHUNG	85 SPEC	8 $\pi^- p \rightarrow N K \bar{K} \pi$
32 ± 3	604	ARMSTRONG	84 OMEG	85 $\pi^+ p \rightarrow K \bar{K} \pi \pi p,$ $pp \rightarrow K \bar{K} \pi p p$
24 ± 3		CHAUVAT	84 SPEC	ISR 31.5 pp
29 ± 10	103	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
28.3 ± 6.7	320	NACASCH	78 HBC	0.7,0.76 $\bar{p}p \rightarrow K \bar{K} 3\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
32.4 ± 5.8		⁴ AAIJ	14Y LHCb	$\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+\pi^-)$
18.2 ± 1.2		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$ p_{fast}
19.4 ± 1.5		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+)$ p_{fast}
40 ± 5		ABATZIS	94 OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
31 ± 5		ARMSTRONG	89E OMEG	300 $pp \rightarrow pp2(\pi^+\pi^-)$
41 ± 12		ARMSTRONG	89G OMEG	85 $\pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$
17.9 ± 10.9	60	RATH	89 MPS	21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
14 ⁺²⁰ ₋₁₄ ± 10	16	BECKER	87 MRK3	$e^+e^- \rightarrow \phi K \bar{K} \pi$
26 ± 12		EVANGELIS...	81 OMEG	12 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
25 ± 15	200	GURTU	79 HBC	4.2 $K^- p \rightarrow n\eta 2\pi$
~ 10		⁵ STANTON	79 CNTR	8.5 $\pi^- p \rightarrow n 2\gamma 2\pi$
24 ± 18	210	GRASSLER	77 HBC	16 $\pi^\mp p$
28 ± 5	150	⁶ DEFOIX	72 HBC	0.7 $\bar{p}p \rightarrow 7\pi$
46 ± 9	180	⁶ DUBOC	72 HBC	1.2 $\bar{p}p \rightarrow 2K 4\pi$
37 ± 5	500	⁷ THUN	72 MMS	13.4 $\pi^- p$
10 ± 10		BOESEBECK	71 HBC	16.0 $\pi p \rightarrow p 5\pi$
30 ± 15		CAMPBELL	69 DBC	2.7 $\pi^+ d$
60 ± 15		⁶ LORSTAD	69 HBC	0.7 $\bar{p}p, 4,5\text{-body}$
35 ± 10		⁶ DAHL	67 HBC	1.6-4.2 $\pi^- p$

¹ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

² Supersedes ABATZIS 94, ARMSTRONG 89E.

³ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.

⁴ No systematic error given.

⁵ From phase shift analysis of $\eta\pi^+\pi^-$ system.

⁶ Resolution is not unfolded.

⁷ Seen in the missing mass spectrum.

NODE=M008W

NEW

OCCUR=2

NODE=M008W;LINKAGE=BL

NODE=M008W;LINKAGE=B

NODE=M008W;LINKAGE=A


NODE=M008W;LINKAGE=N1

NODE=M008W;LINKAGE=P

NODE=M008W;LINKAGE=R

NODE=M008W;LINKAGE=S

WEIGHTED AVERAGE
24.1±1.0 (Error scaled by 1.3)



$f_1(1285)$ DECAY MODES

NODE=M008215;NODE=M008

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 4π	$(33.1^{+2.1}_{-1.8})\%$	S=1.3	DESIG=21
Γ_2 $\pi^0\pi^0\pi^+\pi^-$	$(22.0^{+1.4}_{-1.2})\%$	S=1.3	DESIG=22
Γ_3 $2\pi^+2\pi^-$	$(11.0^{+0.7}_{-0.6})\%$	S=1.3	DESIG=20
Γ_4 $\rho^0\pi^+\pi^-$	$(11.0^{+0.7}_{-0.6})\%$	S=1.3	DESIG=191
Γ_5 $\rho^0\rho^0$	seen		DESIG=23
Γ_6 $4\pi^0$	$< 7 \times 10^{-4}$	CL=90%	DESIG=7
Γ_7 $\eta\pi^+\pi^-$	$(35 \pm 15)\%$		DESIG=198
Γ_8 $\eta\pi\pi$	$(52.4^{+1.9}_{-2.2})\%$	S=1.2	DESIG=3
Γ_9 $a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$]	$(36 \pm 7)\%$		DESIG=4
Γ_{10} $\eta\pi\pi$ [excluding $a_0(980)\pi$]	$(16 \pm 7)\%$		DESIG=5
Γ_{11} $K\bar{K}\pi$	$(9.0 \pm 0.4)\%$	S=1.1	DESIG=1
Γ_{12} $K\bar{K}^*(892)$	not seen		DESIG=6
Γ_{13} $\pi^+\pi^-\pi^0$	$(3.0 \pm 0.9) \times 10^{-3}$		DESIG=197
Γ_{14} $\rho^\pm\pi^\mp$	$< 3.1 \times 10^{-3}$	CL=95%	DESIG=199
Γ_{15} $\gamma\rho^0$	$(5.5 \pm 1.3)\%$	S=2.8	DESIG=13
Γ_{16} $\phi\gamma$	$(7.4 \pm 2.6) \times 10^{-4}$		DESIG=10
Γ_{17} $\gamma\gamma^*$			DESIG=9
Γ_{18} $\gamma\gamma$			DESIG=8

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 16 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 24.7$ for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_9	-17			
x_{10}	-8	-95		
x_{11}	46	-9	-4	
x_{15}	-36	-4	-2	-34
	x_1	x_9	x_{10}	x_{11}

$f_1(1285) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

NODE=M008217

$$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$$

$$\Gamma_8 \Gamma_{18} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{18} / \Gamma$$
NODE=M008G2
NODE=M008G2

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.62	95	GIDAL	87	MRK2 $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$

$$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*) / \Gamma_{\text{total}}$$

$$\Gamma_8 \Gamma_{17} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{17} / \Gamma$$
NODE=M008G3
NODE=M008G3

VALUE (keV)	EVTs	DOCUMENT ID	TECN	COMMENT
1.4 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.4.
1.18 ± 0.25 ± 0.20	26	^{1,2} AIHARA	88B	TPC $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
2.30 ± 0.61 ± 0.42		^{1,3} GIDAL	87	MRK2 $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 0.3 ± 0.3	420	⁴ ACHARD	02B	L3 $183-209 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
-----------------	-----	---------------------	-----	---

¹ Assuming a ρ -pole form factor.

² Published value multiplied by $\eta\pi\pi$ branching ratio 0.49.

³ Published value divided by 2 and multiplied by the $\eta\pi\pi$ branching ratio 0.49.

⁴ Published value multiplied by the $\eta\pi\pi$ branching ratio 0.52.

NODE=M008G3;LINKAGE=A
NODE=M008G3;LINKAGE=F
NODE=M008G3;LINKAGE=B
NODE=M008G3;LINKAGE=AC

$f_1(1285)$ BRANCHING RATIOS

NODE=M008220

$$\Gamma(K\bar{K}\pi) / \Gamma(4\pi)$$

$$\Gamma_{11} / \Gamma_1$$
NODE=M008R1
NODE=M008R1

VALUE	DOCUMENT ID	TECN	COMMENT
0.271 ± 0.016 OUR FIT			Error includes scale factor of 1.3.
0.271 ± 0.016 OUR AVERAGE			Error includes scale factor of 1.2.
0.265 ± 0.014	¹ BARBERIS	97C	OMEG 450 $pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
0.28 ± 0.05	² ARMSTRONG	89E	OMEG 300 $pp \rightarrow pp f_1(1285)$
0.37 ± 0.03 ± 0.05	³ ARMSTRONG	89G	OMEG 85 $\pi p \rightarrow 4\pi X$

¹ Using $2(\pi^+ \pi^-)$ data from BARBERIS 97B.

² Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.

³ 4π consistent with being entirely $\rho\pi\pi$.

NODE=M008R1;LINKAGE=B
NODE=M008R1;LINKAGE=M
NODE=M008R1;LINKAGE=A
$$\Gamma(\pi^0 \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$$

$$\Gamma_2 / \Gamma = \frac{2}{3} \Gamma_1 / \Gamma$$
NODE=M008R18
NODE=M008R18

0.220 $^{+0.014}_{-0.012}$ OUR FIT	Error includes scale factor of 1.3.
--	-------------------------------------

$$\Gamma(2\pi^+ 2\pi^-) / \Gamma_{\text{total}}$$

$$\Gamma_3 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$$
NODE=M008R17
NODE=M008R17

0.110 $^{+0.007}_{-0.006}$ OUR FIT	Error includes scale factor of 1.3.
--	-------------------------------------

$$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$$

$$\Gamma_4 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$$
NODE=M008R19
NODE=M008R19

0.110 $^{+0.007}_{-0.006}$ OUR FIT	Error includes scale factor of 1.3.
--	-------------------------------------

$$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma(2\pi^+ 2\pi^-)$$

 Γ_4 / Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M008R6
NODE=M008R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.0 ± 0.4	GRASSLER	77	HBC 16 GeV $\pi^\pm p$
-----------	----------	----	------------------------

$$\Gamma(\rho^0 \rho^0) / \Gamma_{\text{total}}$$

 Γ_5 / Γ

VALUE	DOCUMENT ID	COMMENT
-------	-------------	---------

NODE=M008R21
NODE=M008R21

seen BARBERIS 00C 450 $pp \rightarrow p_f 4\pi p_s$

$$\Gamma(4\pi^0) / \Gamma_{\text{total}}$$

 Γ_6 / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M008R8
NODE=M008R8

<7 90 ALDE 87 GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$

$$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma(\eta \pi^+ \pi^-)$$

 Γ_{13} / Γ_7

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

NODE=M008R02
NODE=M008R02

0.86 ± 0.16 ± 0.20 2.3k ¹DOROFEEV 11 VES $\pi^- N \rightarrow \pi^- f_1(1285) N$

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+ \pi^-$ mass spectrum.

NODE=M008R02;LINKAGE=DO

$$\Gamma(\eta \pi \pi) / \Gamma_{\text{total}}$$

 $\Gamma_8 / \Gamma = (\Gamma_9 + \Gamma_{10}) / \Gamma$

VALUE	DOCUMENT ID
-------	-------------

NODE=M008R22
NODE=M008R22

0.524^{+0.019}_{-0.022} OUR FIT Error includes scale factor of 1.2.

$$\Gamma(4\pi) / \Gamma(\eta \pi \pi)$$

 $\Gamma_1 / \Gamma_8 = \Gamma_1 / (\Gamma_9 + \Gamma_{10})$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M008R4
NODE=M008R4

0.63 ± 0.06 OUR FIT Error includes scale factor of 1.2.

0.41 ± 0.14 OUR AVERAGE

0.37 ± 0.11 ± 0.11	BOLTON	92	MRK3 $J/\psi \rightarrow \gamma f_1(1285)$
--------------------	--------	----	--

0.64 ± 0.40	GURTU	79	HBC 4.2 $K^- p$
-------------	-------	----	-----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.93 ± 0.30	¹ GRASSLER	77	HBC 16 $\pi^\mp p$
-------------	-----------------------	----	--------------------

¹ Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.

NODE=M008R4;LINKAGE=M

$$\Gamma(2\pi^+ 2\pi^-) / \Gamma(\eta \pi \pi)$$

 Γ_3 / Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M008R04
NODE=M008R04

0.28 ± 0.02 ± 0.02 ¹LEES 12X BABR $\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$

¹ Assuming $B(f_1(1285) \rightarrow \pi\pi\eta) = 3/2 B(f_1(1285) \rightarrow \pi^+ \pi^- \eta)$.

NODE=M008R04;LINKAGE=LE

$$\Gamma(a_0(980)\pi \text{ [ignoring } a_0(980) \rightarrow K\bar{K}]) / \Gamma(\eta \pi \pi)$$

 $\Gamma_9 / \Gamma_8 = \Gamma_9 / (\Gamma_9 + \Gamma_{10})$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

NODE=M008R3
NODE=M008R3

0.69 ± 0.13 OUR FIT

0.69^{+0.13}_{-0.12} OUR AVERAGE

0.72 ± 0.15	GURTU	79	HBC 4.2 $K^- p$
-------------	-------	----	-----------------

0.6 ^{+0.3} _{-0.2}	CORDEN	78	OMEG 12-15 $\pi^- p$
-------------------------------------	--------	----	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.69	95	318	ACHARD	02B	L3	183-209 $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
-------	----	-----	--------	-----	----	--

0.28 ± 0.07	1400	ALDE	97B	GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
-------------	------	------	-----	------	--

1.0 ± 0.3	GRASSLER	77	HBC 16 $\pi^\mp p$
-----------	----------	----	--------------------

$$\Gamma(K\bar{K}\pi) / \Gamma(\eta \pi \pi)$$

 $\Gamma_{11} / \Gamma_8 = \Gamma_{11} / (\Gamma_9 + \Gamma_{10})$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M008R2
NODE=M008R2

0.171 ± 0.013 OUR FIT Error includes scale factor of 1.1.

0.170 ± 0.012 OUR AVERAGE

0.166 ± 0.01 ± 0.008	BARBERIS	98C	OMEG 450 $pp \rightarrow p_f f_1(1285) p_s$
----------------------	----------	-----	---

0.42 ± 0.15	GURTU	79	HBC 4.2 $K^- p$
-------------	-------	----	-----------------

0.5 ± 0.2	¹ CORDEN	78	OMEG 12-15 $\pi^- p$
-----------	---------------------	----	----------------------

0.20 ± 0.08	² DEFOIX	72	HBC 0.7 $\bar{p} p \rightarrow 7\pi$
-------------	---------------------	----	--------------------------------------

0.16 ± 0.08	CAMPBELL	69	DBC 2.7 $\pi^+ d$
-------------	----------	----	-------------------

¹ CORDEN 78 assumes low-mass $\eta\pi\pi$ region is dominantly 1^{++} . See BARBERIS 98C and MANAK 00A for discussion.

NODE=M008R2;LINKAGE=CD

² $K\bar{K}$ system characterized by the $l = 1$ threshold enhancement. (See under $a_0(980)$).

NODE=M008R2;LINKAGE=K

$\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	NACASCH	78	HBC 0.7,0.76 $\bar{p}p \rightarrow K\bar{K}3\pi$
----------	---------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	¹ ACHARD	07	L3 183-209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
------	---------------------	----	---

¹ A clear signal of 19.8 ± 4.4 events observed at high Q^2 .

NODE=M008R5
NODE=M008R5

NODE=M008R5;LINKAGE=CH

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.30±0.055±0.074	2.3k	¹ DOROFEEV	11	VES $\pi^-N \rightarrow \pi^-f_1(1285)N$
-------------------------	------	-----------------------	----	--

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum. The systematic error includes the uncertainty on the partial width $f_1 \rightarrow \eta\pi\pi$ obtained from PDG 10 data.

NODE=M008R01
NODE=M008R01

NODE=M008R01;LINKAGE=DO

 $\Gamma(\rho^\pm\pi^\mp)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

<0.31	95	DOROFEEV	11	VES $\pi^-N \rightarrow \pi^-f_1(1285)N$
-----------------	----	----------	----	--

NODE=M008R03
NODE=M008R03

 $\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

5.5±1.3 OUR FIT Error includes scale factor of 2.8.

2.8±0.7±0.6		AMELIN	95	VES $37 \pi^-N \rightarrow \pi^-\pi^+\pi^-\gamma N$
--------------------	--	--------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5	95	BITYUKOV	91B	SPEC $32 \pi^-p \rightarrow \pi^+\pi^-\gamma n$
----	----	----------	-----	---

NODE=M008R15
NODE=M008R15

 $\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.50±0.13 OUR FIT Error includes scale factor of 2.5.

0.45±0.18	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
------------------	----------------------	----	--

¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.

NODE=M008R13
NODE=M008R13

NODE=M008R13;LINKAGE=E

 $\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$ $\Gamma_8/\Gamma_{15} = (\Gamma_9+\Gamma_{10})/\Gamma_{15}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

9.5±2.0 OUR FIT Error includes scale factor of 2.5.

7.9±0.9 OUR AVERAGE

10.0±1.0±2.0	BARBERIS	98C	OMEG 450 $pp \rightarrow p_f f_1(1285) p_s$
--------------	----------	-----	---

7.5±1.0	¹ ARMSTRONG	92C	OMEG 300 $pp \rightarrow pp\pi^+\pi^-\gamma, pp\eta\pi^+\pi^-$
---------	------------------------	-----	--

¹ Published value multiplied by 1.5.

NODE=M008R16
NODE=M008R16

NODE=M008R16;LINKAGE=B

 $\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$ Γ_{15}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.035	90	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
--------	----	----------------------	----	--

¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) = 0.72 \times 10^{-3}$.

NODE=M008R12
NODE=M008R12

NODE=M008R12;LINKAGE=F

 $\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$ Γ_{16}/Γ_{11}

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.82±0.21±0.20		19	BITYUKOV	88	SPEC $32.5 \pi^-p \rightarrow K^+K^-\pi^0 n$
-----------------------	--	----	----------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.50	95	BARBERIS	98C	OMEG 450 $pp \rightarrow p_f f_1(1285) p_s$
-------	----	----------	-----	---

<0.93	95	AMELIN	95	VES $37 \pi^-N \rightarrow \pi^-\pi^+\pi^-\gamma N$
-------	----	--------	----	---

NODE=M008R9
NODE=M008R9

f₁(1285) REFERENCES

NODE=M008

ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56781
AAIJ	14Y	PRL 112 091802	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55837
LEES	12X	PR D86 092010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54714
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53931
DOROFEEV	11	EPJ A47 68	V. Dorofeev <i>et al.</i>	(SERP, MIPT)	REFID=16755
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)	REFID=53229
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=51698
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49548
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48574
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47989
SOSA	99	PRL 83 913	M. Sosa <i>et al.</i>		REFID=46937
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45396
		Translated from YAF 60 458.			
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
AMELIN	95	ZPHY C66 71	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=44376
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44437
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44090
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)	REFID=44092
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
ARMSTRONG	92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=42097
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BITYUKOV	91B	SJNP 54 318	S.I. Bitjukov <i>et al.</i>	(SERP)	REFID=41864
		Translated from YAF 54 529.			
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
ARMSTRONG	89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+)	REFID=40729
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
ARMSTRONG	89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)	REFID=40930
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)	REFID=40924
AIHARA	88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40572
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
BITYUKOV	88	PL B203 327	S.I. Bitjukov <i>et al.</i>	(SERP)	REFID=40569
MIR	88	Photon-Photon 88, 126	R. Mir	(Mark III Collab.)	REFID=41574
Conference					
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA) IJP	REFID=20891
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP	REFID=20936
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP	REFID=20934
ARMSTRONG	84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP	REFID=20929
BITYUKOV	84B	PL 144B 133	S.I. Bitjukov <i>et al.</i>	(SERP)	REFID=20468
CHAUVAT	84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)	REFID=20932
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)	REFID=20573
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
BROMBERG	80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)	REFID=20922
DIONISI	80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+)	REFID=20924
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)	REFID=20456
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP	REFID=20887
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20452
NACASCH	78	NP B135 203	R. Nacasch <i>et al.</i>	(PARIS, MADR, CERN)	REFID=20919
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20447
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
DUBOC	72	NP B46 429	J. Duboc <i>et al.</i>	(PARIS, LVP)	REFID=20339
THUN	72	PRL 28 1733	R. Thun <i>et al.</i>	(STON, NEAS)	REFID=20911
BARDADIN...	71	PR D4 2711	M. Bardadin-Otwinowska <i>et al.</i>	(WARS)	REFID=20196
BOESEBECK	71	PL 34B 659	K. Boesebeck	(AACH, BERL, BONN, CERN, CRAC+)	REFID=20905
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)	REFID=20419
LORSTAD	69	NP B14 63	B. Lorstad <i>et al.</i>	(CDEF, CERN) JP	REFID=20901
D'ANDLAU	68	NP B5 693	C. d'Andlau <i>et al.</i>	(CDEF, CERN, IRAD+) IJP	REFID=20897
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP	REFID=20321

$\eta(1295)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See also the mini-review under $\eta(1405)$

NODE=M037

NODE=M037

NODE=M037M


NODE=M037M

 $\eta(1295)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1294±4 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
1302±9±8	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1282±5	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1299±4	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1295±4		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1264±8		¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~ 1275		STANTON	79 CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$



¹ PWA analysis of AUGUSTIN 92 assigns 0^{-+} quantum numbers to this state rather than 1^{++} as before.

NODE=M037M;LINKAGE=AG

 $\eta(1295)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
55± 5 OUR AVERAGE				
57±23±21	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
66±13	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
53± 6		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<40	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
44±20		² AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~ 70		STANTON	79 CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$

² PWA analysis of AUGUSTIN 92 assigns 0^{-+} quantum numbers to this state rather than 1^{++} as before.

NODE=M037W

NODE=M037W

NODE=M037W;LINKAGE=AG

$\eta(1295)$ DECAY MODES

NODE=M037215;NODE=M037

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi^+\pi^-$	seen
Γ_2 $a_0(980)\pi$	seen
Γ_3 $\gamma\gamma$	
Γ_4 $\eta\pi^0\pi^0$	seen
Γ_5 $\eta(\pi\pi)S\text{-wave}$	seen
Γ_6 $\sigma\eta$	
Γ_7 $K\bar{K}\pi$	

DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=3
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6
DESIG=7

 $\eta(1295)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M037220

 $\Gamma(\eta\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_3/\Gamma$ NODE=M037G2
NODE=M037G2

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.066	95	ACCIARRI	01G L3	183-202 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.6	90	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
<0.3		ANTREASYAN	87 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi\pi$

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_3/\Gamma$ NODE=M037G3
NODE=M037G3

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.014	90	3,4 AHOHE	05 CLE2	10.6 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$

³ Using $\eta(1295)$ mass and width 1294 MeV and 55 MeV, respectively.⁴ Assuming three-body phase-space decay to $K_S^0K^\pm\pi^\mp$.NODE=M037G3;LINKAGE=AH
NODE=M037G3;LINKAGE=B3 $\eta(1295)$ BRANCHING RATIOS

NODE=M037225

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M037R1
NODE=M037R1

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
not seen	BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
seen	BIRMAN	88 MPS	8 $\pi^-p \rightarrow K^+\bar{K}^0\pi^-n$
large	ANDO	86 SPEC	8 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
large	STANTON	79 CNTR	8.4 $\pi^-p \rightarrow n\eta2\pi$

 $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi^0\pi^0)$ Γ_2/Γ_4 NODE=M037R2
NODE=M037R2

VALUE	DOCUMENT ID	TECN	COMMENT
0.65±0.10	⁵ ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

⁵ Assuming that $a_0(980)$ decays only to $\eta\pi$.

NODE=M037R2;LINKAGE=A

 $\Gamma(\eta(\pi\pi)S\text{-wave})/\Gamma(\eta\pi^0\pi^0)$ Γ_5/Γ_4 NODE=M037R4
NODE=M037R4

VALUE	DOCUMENT ID	TECN	COMMENT
0.35±0.10	ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

 $\Gamma(a_0(980)\pi)/\Gamma(\sigma\eta)$ Γ_2/Γ_6 NODE=M037R5
NODE=M037R5

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.48±0.22	9082	MANAK	00A MPS	18 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

 $\eta(1295)$ REFERENCES

NODE=M037

AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)	REFID=50764
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47989
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45396
Translated from YAF 60 458.					
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40564
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=40008
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) IJP	REFID=20891
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP	REFID=20887

$\pi(1300)$

$$I^G(J^{PC}) = 1^-(0^{-+})$$

NODE=M058

 $\pi(1300)$ MASS

NODE=M058M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M058M

1300±100 OUR ESTIMATE

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

1345± 8±10	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
1200± 40	90k	SALVINI 04	OBLX	$\bar{p}p \rightarrow 2\pi^+2\pi^-$
1343± 15±24		CHUNG 02	B852	$18.3 \pi^-p \rightarrow \pi^+\pi^-\pi^-p$
1375± 40		ABELE 01	CBAR	$0.0 \bar{p}d \rightarrow \pi^-4\pi^0p$
1275± 15		BERTIN 97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+2\pi^-$
~ 1114		ABELE 96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
1190± 30		ZIELINSKI 84	SPEC	$200 \pi^+Z \rightarrow Z3\pi$
1240± 30		BELLINI 82	SPEC	$40 \pi^-A \rightarrow A3\pi$
1273± 50		² AARON 81	RVUE	
1342± 20		BONESINI 81	OMEG	$12 \pi^-p \rightarrow p3\pi$
~ 1400		DAUM 81B	SPEC	$63,94 \pi^-p$

¹ From analysis of L3 data at 183–209 GeV.

NODE=M058M;LINKAGE=SC

² Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

NODE=M058M;LINKAGE=E

 $\pi(1300)$ WIDTH

NODE=M058W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M058W

200 to 600 OUR ESTIMATE

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

260± 20±30	18k	³ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
470±120	90k	SALVINI 04	OBLX	$\bar{p}p \rightarrow 2\pi^+2\pi^-$
449± 39±47		CHUNG 02	B852	$18.3 \pi^-p \rightarrow \pi^+\pi^-\pi^-p$
268± 50		ABELE 01	CBAR	$0.0 \bar{p}d \rightarrow \pi^-4\pi^0p$
218±100		BERTIN 97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+2\pi^-$
~ 340		ABELE 96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
440± 80		ZIELINSKI 84	SPEC	$200 \pi^+Z \rightarrow Z3\pi$
360±120		BELLINI 82	SPEC	$40 \pi^-A \rightarrow A3\pi$
580±100		⁴ AARON 81	RVUE	
220± 70		BONESINI 81	OMEG	$12 \pi^-p \rightarrow p3\pi$
~ 600		DAUM 81B	SPEC	$63,94 \pi^-p$

³ From analysis of L3 data at 183–209 GeV.

NODE=M058W;LINKAGE=SC

⁴ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

NODE=M058W;LINKAGE=E

 $\pi(1300)$ DECAY MODES

NODE=M058215;NODE=M058

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\pi(\pi\pi)S$ -wave	seen
Γ_3 $\gamma\gamma$	

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

 $\pi(1300)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M058218

$\Gamma(\rho\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_3/\Gamma$
---	---------------------------

NODE=M058G1

NODE=M058G1

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

<0.085	90	ACCIARRI 97T	L3	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
--------	----	--------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	95	⁵ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
<0.54	90	ALBRECHT 97B	ARG	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$

⁵ From analysis of L3 data at 183–209 GeV.

NODE=M058G1;LINKAGE=SC

$\pi(1300)$ BRANCHING RATIOS

NODE=M058220

 $\Gamma(\pi\pi)_{S\text{-wave}}/\Gamma(\rho\pi)$ Γ_2/Γ_1

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.4		90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
seen			CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+ 2\pi^- p$
<0.15	90		ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
2.12			⁶ AARON	81	RVUE

NODE=M058R1
NODE=M058R1

⁶ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

NODE=M058R1;LINKAGE=E

 $\pi(1300)$ REFERENCES

NODE=M058

SCHEGELSKY 06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	REFID=51186
SALVINI 04	EPJ C35 21	P. Salvini <i>et al.</i>	REFID=53226
CHUNG 02	PR D65 072001	S.U. Chung <i>et al.</i>	REFID=48837
ABELE 01	EPJ C19 667	A. Abele <i>et al.</i>	REFID=48334
ACCIARRI 97T	PL B413 147	M. Acciarri <i>et al.</i>	REFID=45761
ALBRECHT 97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	REFID=45418
BERTIN 97D	PL B414 220	A. Bertin <i>et al.</i>	REFID=45763
ABELE 96	PL B380 453	A. Abele <i>et al.</i>	REFID=45011
ZIELINSKI 84	PR D30 1855	M. Zielinski <i>et al.</i>	REFID=20881
BELLINI 82	PRL 48 1697	G. Bellini <i>et al.</i>	REFID=21134
AARON 81	PR D24 1207	R.A. Aaron, R.S. Longacre	REFID=20870
BONESINI 81	PL 103B 75	M. Bonesini <i>et al.</i>	REFID=21130
DANKOWYCH... 81	PRL 46 580	J.A. Dankowych <i>et al.</i>	REFID=20572
DAUM 81B	NP B182 269	C. Daum <i>et al.</i>	REFID=20872
DAUM 80	PL 89B 281	C. Daum <i>et al.</i>	REFID=20868
BOWLER 75	NP B97 227	M.G. Bowler <i>et al.</i>	REFID=20571

NODE=M012

 $a_2(1320)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

 $a_2(1320)$ MASS

NODE=M012205

VALUE (MeV) DOCUMENT ID

NODE=M012M0

1318.3^{+0.5}_{-0.6} OUR AVERAGE Includes data from the 4 datablocks that follow this one.
Error includes scale factor of 1.2.

3 π MODENODE=M012M1
NODE=M012M1

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

1319.0^{+1.0}_{-1.3} OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

1321 ± 1	⁺⁰ ₋₇	420k	ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1326 ± 2	± 2		CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1317 ± 3			BARBERIS	98B		450 $pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$
1323 ± 4	± 3		ACCIARRI	97T	L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
1320 ± 7			ALBRECHT	97B	ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
1311.3 ± 1.6 ± 3.0	72.4k		AMELIN	96	VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1310 ± 5			ARMSTRONG	90	OMEG 0	300.0 $pp \rightarrow pp \pi^+ \pi^- \pi^0$
1323.8 ± 2.3	4022		AUGUSTIN	89	DM2 ±	$J/\psi \rightarrow \rho^\pm a_2^\mp$
1320.6 ± 3.1	3562		AUGUSTIN	89	DM2 0	$J/\psi \rightarrow \rho^0 a_2^0$
1317 ± 2	25k		¹ DAUM	80C	SPEC -	63,94 $\pi^- p \rightarrow 3\pi p$
1320 ± 10	1097		¹ BALTAY	78B	HBC +0	15 $\pi^+ p \rightarrow p 4\pi$
1306 ± 8			FERRERSORIA	78	OMEG -	9 $\pi^- p \rightarrow p 3\pi$
1318 ± 7	1.6k		¹ EMMS	75	DBC 0	4 $\pi^+ n \rightarrow p(3\pi)^0$
1315 ± 5			¹ ANTIPOV	73C	CNTR -	25,40 $\pi^- p \rightarrow p \eta \pi^-$
1306 ± 9	1580		CHALOUKKA	73	HBC -	3.9 $\pi^- p$

OCCUR=2


• • • We do not use the following data for averages, fits, limits, etc. • • •

1300 ± 2 ± 4	18k	² SCHEGELSKY 06	RVUE	0	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	
1305 ± 14		CONDO 93	SHF		$\gamma p \rightarrow n\pi^+ \pi^+ \pi^-$	
1310 ± 2		¹ EVANGELIS...	81	OMEG -	$12 \pi^- p \rightarrow 3\pi p$	
1343 ± 11	490	BALTAY	78B	HBC 0	$15 \pi^+ p \rightarrow \Delta 3\pi$	OCCUR=2
1309 ± 5	5k	BINNIE	71	MMS -	$\pi^- p$ near a_2 thresh- old	OCCUR=2
1299 ± 6	28k	BOWEN	71	MMS -	$5 \pi^- p$	
1300 ± 6	24k	BOWEN	71	MMS +	$5 \pi^+ p$	OCCUR=2
1309 ± 4	17k	BOWEN	71	MMS -	$7 \pi^- p$	OCCUR=3
1306 ± 4	941	ALSTON-...	70	HBC +	$7.0 \pi^+ p \rightarrow 3\pi p$	

¹From a fit to $J^P = 2^+ \rho\pi$ partial wave.

²From analysis of L3 data at 183–209 GeV.

NODE=M012M1;LINKAGE=P
NODE=M012M1;LINKAGE=SC



K \bar{K} MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M2
NODE=M012M2

1318.1 ± 0.7 OUR AVERAGE

1319 ± 5	4700	^{1,2} CLELAND	82B	SPEC +	$50 \pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
1324 ± 6	5200	^{1,2} CLELAND	82B	SPEC -	$50 \pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
1320 ± 2	4000	CHABAUD	80	SPEC -	$17 \pi^- A \rightarrow K_S^0 K^- A$	
1312 ± 4	11000	CHABAUD	78	SPEC -	$9.8 \pi^- p \rightarrow K^- K_S^0 p$	
1316 ± 2	4730	CHABAUD	78	SPEC -	$18.8 \pi^- p \rightarrow K^- K_S^0 p$	OCCUR=2
1318 ± 1		^{1,3} MARTIN	78D	SPEC -	$10 \pi^- p \rightarrow K_S^0 K^- p$	
1320 ± 2	2724	MARGULIE	76	SPEC -	$23 \pi^- p \rightarrow K^- K_S^0 p$	
1313 ± 4	730	FOLEY	72	CNTR -	$20.3 \pi^- p \rightarrow K^- K_S^0 p$	
1319 ± 3	1500	³ GRAYER	71	ASPK -	$17.2 \pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1304 ± 10	870	⁴ SCHEGELSKY 06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
1330 ± 11	1000	^{1,2} CLELAND	82B	SPEC +	$30 \pi^+ p \rightarrow K_S^0 K^+ p$	
1324 ± 5	350	HYAMS	78	ASPK +	$12.7 \pi^+ p \rightarrow K^+ K_S^0 p$	

¹From a fit to $J^P = 2^+$ partial wave.

²Number of events evaluated by us.

³Systematic error in mass scale subtracted.

⁴From analysis of L3 data at 91 and 183–209 GeV.

NODE=M012M2;LINKAGE=P
NODE=M012M2;LINKAGE=W
NODE=M012M2;LINKAGE=S
NODE=M012M2;LINKAGE=SC

$\eta\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M3
NODE=M012M3**1317.7± 1.4 OUR AVERAGE**

1308 ± 9		BARBERIS	00H		450 $p p \rightarrow p_f \eta \pi^0 p_s$
1316 ± 9		BARBERIS	00H		450 $p p \rightarrow \Delta_f^{++} \eta \pi^- p_s$
1317 ± 1 ± 2		THOMPSON	97	MPS	18 $\pi^- p \rightarrow \eta \pi^- p$
1315 ± 5 ± 2		¹ AMSLER	94D	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$
1325.1 ± 5.1		AOYAGI	93	BKEI	$\pi^- p \rightarrow \eta \pi^- p$
1317.7 ± 1.4 ± 2.0		BELADIDZE	93	VES	37 $\pi^- N \rightarrow \eta \pi^- N$
1323 ± 8	1000	² KEY	73	OSPK -	6 $\pi^- p \rightarrow p \pi^- \eta$

OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1315 ± 12		³ ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(l)} \pi^- p$
1309 ± 4		ANISOVICH	09	RVUE	$\bar{p} p, \pi N$
1324 ± 5		ARMSTRONG	93C	E760 0	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1336.2 ± 1.7	2561	DELFOSE	81	SPEC +	$\pi^\pm p \rightarrow p \pi^\pm \eta$
1330.7 ± 2.4	1653	DELFOSE	81	SPEC -	$\pi^\pm p \rightarrow p \pi^\pm \eta$
1324 ± 8	6200	^{2,4} CONFORTO	73	OSPK -	6 $\pi^- p \rightarrow p \text{MM}^-$

OCCUR=2

¹ The systematic error of 2 MeV corresponds to the spread of solutions.² Error includes 5 MeV systematic mass-scale error.³ ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the $\eta\pi$ and $\rho\pi$ channels into account.⁴ Missing mass with enriched MMS = $\eta\pi^-$, $\eta = 2\gamma$.

NODE=M012M3;LINKAGE=DD

NODE=M012M3;LINKAGE=E

NODE=M012M3;LINKAGE=A

NODE=M012M3;LINKAGE=M

 $\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M4
NODE=M012M4**1322 ± 7 OUR AVERAGE**

1318 ± 8 $\begin{smallmatrix} +3 \\ -5 \end{smallmatrix}$		IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
1327.0 ± 10.7		BELADIDZE	93	VES	37 $\pi^- N \rightarrow \eta' \pi^- N$

 $a_2(1320)$ WIDTH

NODE=M012210

3 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M012W1
NODE=M012W1**105.0 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.6 \\ 1.9 \end{smallmatrix}$ OUR AVERAGE**

110 ± 2 $\begin{smallmatrix} +2 \\ -15 \end{smallmatrix}$	420k	ALEKSEEV	10	COMP	190 $\pi^- P b \rightarrow \pi^- \pi^- \pi^+ P b'$
108 ± 3 ± 15		CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
120 ± 10		BARBERIS	98B		450 $p p \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$
105 ± 10 ± 11		ACCIARRI	97T	L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
120 ± 10		ALBRECHT	97B	ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
103.0 ± 6.0 ± 3.3	72.4k	AMELIN	96	VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
120 ± 10		ARMSTRONG	90	OMEG 0	300.0 $p p \rightarrow p p \pi^+ \pi^- \pi^0$
107.0 ± 9.7	4022	AUGUSTIN	89	DM2 ±	$J/\psi \rightarrow \rho^\pm a_2^\mp$
118.5 ± 12.5	3562	AUGUSTIN	89	DM2 0	$J/\psi \rightarrow \rho^0 a_2^0$
97 ± 5		¹ EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow 3\pi p$
96 ± 9	25k	¹ DAUM	80C	SPEC -	63,94 $\pi^- p \rightarrow 3\pi p$
110 ± 15	1097	¹ BALTAY	78B	HBC +0	15 $\pi^+ p \rightarrow p 4\pi$
112 ± 18	1.6k	¹ EMMS	75	DBC 0	4 $\pi^+ n \rightarrow p(3\pi)^0$
122 ± 14	1.2k	^{1,2} WAGNER	75	HBC 0	7 $\pi^+ p \rightarrow \Delta^{++} (3\pi)^0$
115 ± 15		¹ ANTIPOV	73C	CNTR -	25,40 $\pi^- p \rightarrow p \eta \pi^-$
99 ± 15	1580	CHALOUPKA	73	HBC -	3.9 $\pi^- p$
105 ± 5	28k	BOWEN	71	MMS -	5 $\pi^- p$
99 ± 5	24k	BOWEN	71	MMS +	5 $\pi^+ p$
103 ± 5	17k	BOWEN	71	MMS -	7 $\pi^- p$

OCCUR=2

OCCUR=2

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

117 ± 6 ±20	18k	³ SCHEGELSKY 06	RVUE	0	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$	
120 ±40		CONDO 93	SHF		$\gamma p \rightarrow n\pi^+\pi^+\pi^-$	
115 ±14	490	BALTAY 78B	HBC	0	$15 \pi^+ p \rightarrow \Delta 3\pi$	OCCUR=2
72 ±16	5k	BINNIE 71	MMS	-	$\pi^- p$ near a_2 thresh- old	OCCUR=2
79 ±12	941	ALSTON-... 70	HBC	+	$7.0 \pi^+ p \rightarrow 3\pi p$	

¹ From a fit to $J^P = 2^+$ $\rho\pi$ partial wave.

² Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

³ From analysis of L3 data at 183–209 GeV.

NODE=M012W1;LINKAGE=P
NODE=M012W1;LINKAGE=S
NODE=M012W1;LINKAGE=SC

$K\bar{K}$ AND $\eta\pi$ MODES

VALUE (MeV) DOCUMENT ID

107 ±5 OUR ESTIMATE

110.4±1.7 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

NODE=M012W0
NODE=M012W0
→ UNCHECKED ←

$K\bar{K}$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M012W2
NODE=M012W2

109.8± 2.4 OUR AVERAGE

112 ±20	4700	^{1,2} CLELAND	82B	SPEC	+	$50 \pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
120 ±25	5200	^{1,2} CLELAND	82B	SPEC	-	$50 \pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
106 ± 4	4000	CHABAUD	80	SPEC	-	$17 \pi^- A \rightarrow K_S^0 K^- A$	
126 ±11	11000	CHABAUD	78	SPEC	-	$9.8 \pi^- p \rightarrow K^- K_S^0 p$	
101 ± 8	4730	CHABAUD	78	SPEC	-	$18.8 \pi^- p \rightarrow K^- K_S^0 p$	OCCUR=2
113 ± 4		^{1,3} MARTIN	78D	SPEC	-	$10 \pi^- p \rightarrow K_S^0 K^- p$	
105 ± 8	2724	³ MARGULIE	76	SPEC	-	$23 \pi^- p \rightarrow K^- K_S^0 p$	
113 ±19	730	FOLEY	72	CNTR	-	$20.3 \pi^- p \rightarrow K^- K_S^0 p$	
123 ±13	1500	³ GRAYER	71	ASPK	-	$17.2 \pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

120 ±15	870	⁴ SCHEGELSKY 06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$		
121 ±51	1000	^{1,2} CLELAND	82B	SPEC	+	$30 \pi^+ p \rightarrow K_S^0 K^+ p$	
110 ±18	350	HYAMS 78	ASPK	+	$12.7 \pi^+ p \rightarrow K^+ K_S^0 p$		

¹ From a fit to $J^P = 2^+$ partial wave.

² Number of events evaluated by us.

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁴ From analysis of L3 data at 91 and 183–209 GeV.

NODE=M012W2;LINKAGE=P
NODE=M012W2;LINKAGE=W
NODE=M012W2;LINKAGE=S
NODE=M012W2;LINKAGE=SC

$\eta\pi$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M012W3
NODE=M012W3

111.1± 2.4 OUR AVERAGE

115 ±20		BARBERIS	00H			$450 p p \rightarrow p_f \eta \pi^0 p_S$	
112 ±14		BARBERIS	00H			$450 p p \rightarrow$ $\Delta_f^{++} \eta \pi^- p_S$	OCCUR=2
112 ± 3 ±2		¹ AMSLER	94D	CBAR		$0.0 \bar{p} p \rightarrow \pi^0 \pi^0 \eta$	
103 ± 6 ±3		BELADIDZE	93	VES		$37 \pi^- N \rightarrow \eta \pi^- N$	
112.2± 5.7	2561	DELFOSSÉ	81	SPEC	+	$\pi^\pm p \rightarrow p \pi^\pm \eta$	
116.6± 7.7	1653	DELFOSSÉ	81	SPEC	-	$\pi^\pm p \rightarrow p \pi^\pm \eta$	OCCUR=2
108 ± 9	1000	KEY	73	OSPK	-	$6 \pi^- p \rightarrow p \pi^- \eta$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

119 ±14		² ADOLPH	15	COMP		$191 \pi^- p \rightarrow$ $\eta^{(\prime)} \pi^- p$	
110 ± 4		ANISOVICH	09	RVUE		$\bar{p} p, \pi N$	
127 ± 2 ±2		³ THOMPSON	97	MPS		$18 \pi^- p \rightarrow \eta \pi^- p$	
118 ±10		ARMSTRONG	93C	E760	0	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
104 ± 9	6200	⁴ CONFORTO	73	OSPK	-	$6 \pi^- p \rightarrow p M M^-$	

¹ The systematic error of 2 MeV corresponds to the spread of solutions.

² ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the $\eta\pi$ and $\rho\pi$ channels into account.

³ Resolution is not unfolded.

⁴ Missing mass with enriched MMS = $\eta\pi^-$, $\eta = 2\gamma$.

NODE=M012W3;LINKAGE=DD
NODE=M012W3;LINKAGE=B
NODE=M012W3;LINKAGE=A
NODE=M012W3;LINKAGE=M

$\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
119±25 OUR AVERAGE			
140±35±20	IVANOV 01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
106±32	BELADIDZE 93	VES	37 $\pi^- N \rightarrow \eta' \pi^- N$

NODE=M012W4
 NODE=M012W4

 $a_2(1320)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 3π	(70.1 ± 2.7) %	S=1.2
Γ_2 $\rho(770)\pi$		
Γ_3 $f_2(1270)\pi$		
Γ_4 $\rho(1450)\pi$		
Γ_5 $\eta\pi$	(14.5 ± 1.2) %	
Γ_6 $\omega\pi\pi$	(10.6 ± 3.2) %	S=1.3
Γ_7 $K\bar{K}$	(4.9 ± 0.8) %	
Γ_8 $\eta'(958)\pi$	(5.5 ± 0.9) × 10 ⁻³	
Γ_9 $\pi^\pm\gamma$	(2.91 ± 0.27) × 10 ⁻³	
Γ_{10} $\gamma\gamma$	(9.4 ± 0.7) × 10 ⁻⁶	
Γ_{11} e^+e^-	< 5 × 10 ⁻⁹	CL=90%

NODE=M012215;NODE=M012

DESIG=1
 DESIG=11
 DESIG=12
 DESIG=13
 DESIG=3
 DESIG=4
 DESIG=2
 DESIG=8
 DESIG=7
 DESIG=9
 DESIG=10

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 18 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 9.3$ for 15 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_5	10		
x_6	-89	-46	
x_7	-1	-2	-24
	x_1	x_5	x_6

 $a_2(1320)$ PARTIAL WIDTHS **$\Gamma(\eta\pi)$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
18.5±3.0	870	¹ SCHEGELSKY 06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$ keV and SU(3) relations.

 Γ_5

NODE=M012220

NODE=M012W6
 NODE=M012W6

NODE=M012W6;LINKAGE=SC

 $\Gamma(K\bar{K})$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
7.0 ^{+2.0} _{-1.5}	870	¹ SCHEGELSKY 06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$ keV and SU(3) relations.

 Γ_7

NODE=M012W5
 NODE=M012W5

NODE=M012W5;LINKAGE=SC

 $\Gamma(\pi^\pm\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
311± 25 OUR AVERAGE					

358± 6±42		¹ ADOLPH 14	COMP	-	190 $\pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$
284± 25±25	7.1k	MOLCHANOV 01	SELX		600 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
295± 60		CIHANGIR 82	SPEC	+	200 $\pi^+ A$

• • • We do not use the following data for averages, fits, limits, etc. • • •

461±110		² MAY 77	SPEC	±	9.7 γA
---------	--	---------------------	------	---	----------------

¹ Primakoff reaction using $a_2(1320) \rightarrow 3\pi$ branching ratio of 70.1%.

² Assuming one-pion exchange.

 Γ_9

NODE=M012W7
 NODE=M012W7

NODE=M012W7;LINKAGE=AD
 NODE=M012W;LINKAGE=M2

$\Gamma(\gamma\gamma)$ Γ_{10}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.00±0.06 OUR AVERAGE					
0.98±0.05±0.09		ACCIARRI	97T	L3	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
0.96±0.03±0.13		ALBRECHT	97B	ARG	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.26±0.26±0.18	36	BARU	90	MD1	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.00±0.07±0.15	415	BEHREND	90C	CELL 0	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.03±0.13±0.21		BUTLER	90	MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.01±0.14±0.22	85	OEST	90	JADE	$e^+e^- \rightarrow e^+e^-\pi^0\eta$
0.90±0.27±0.15	56	¹ ALTHOFF	86	TASS 0	$e^+e^- \rightarrow e^+e^-3\pi$
1.14±0.20±0.26		² ANTREASYAN	86	CBAL 0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$
1.06±0.18±0.19		BERGER	84C	PLUT 0	$e^+e^- \rightarrow e^+e^-3\pi$

••• We do not use the following data for averages, fits, limits, etc. •••

0.81±0.19 ^{+0.42} _{-0.11}	35	¹ BEHREND	83B	CELL 0	$e^+e^- \rightarrow e^+e^-3\pi$
0.77±0.18±0.27	22	² EDWARDS	82F	CBAL 0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$

¹ From $\rho\pi$ decay mode.

² From $\eta\pi^0$ decay mode.

NODE=M012W9
NODE=M012W9

NODE=M012W;LINKAGE=F
NODE=M012W;LINKAGE=G

 $\Gamma(e^+e^-)$ Γ_{11}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.56	90	ACHASOV	00K	SND $e^+e^- \rightarrow \pi^0\pi^0$
<25	90	VOROBYEV	88	ND $e^+e^- \rightarrow \pi^0\eta$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M012W10
NODE=M012W10

 $a_2(1320) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M012223

 $\Gamma(3\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{10}/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.65±0.02±0.02	18k	¹ SCHEGELSKY	06	RVUE $\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

¹ From analysis of L3 data at 183–209 GeV.

NODE=M012G2
NODE=M012G2

NODE=M012G2;LINKAGE=SC

 $\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_{10}/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.145 ^{+0.097} _{-0.034}	¹ UEHARA	09A	BELL $e^+e^- \rightarrow e^+e^-\eta\pi^0$

¹ From the D_2 -wave. The fraction of the D_0 -wave is 3.4^{+2.3}_{-1.1}%.

NODE=M012G01
NODE=M012G01

NODE=M012G01;LINKAGE=UE

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_{10}/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.126±0.007±0.028	¹ ALBRECHT	90G	ARG $e^+e^- \rightarrow e^+e^-K^+K^-$
0.081±0.006±0.027	² ALBRECHT	90G	ARG $e^+e^- \rightarrow e^+e^-K^+K^-$

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Using an incoherent background.

² Using a coherent background.

NODE=M012G1
NODE=M012G1

OCCUR=2

NODE=M012G1;LINKAGE=A
NODE=M012G1;LINKAGE=B

 $a_2(1320) \text{ BRANCHING RATIOS}$

NODE=M012225

 $[\Gamma(f_2(1270)\pi) + \Gamma(\rho(1450)\pi)]/\Gamma(\rho(770)\pi)$ $(\Gamma_3+\Gamma_4)/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.12	90	ABRAMOVI...	70B	HBC	— 3.93 π^-p

NODE=M012R9
NODE=M012R9

$\Gamma(\eta\pi)/\Gamma(3\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.207±0.018 OUR FIT					
0.213±0.020 OUR AVERAGE					
0.18 ±0.05		FORINO 76	HBC		11 $\pi^- p$
0.22 ±0.05	52	ANTIPOV 73	CNTR	-	40 $\pi^- p$
0.211±0.044	149	CHALOUPKA 73	HBC	-	3.9 $\pi^- p$
0.246±0.042	167	ALSTON-... 71	HBC	+	7.0 $\pi^+ p$
0.25 ±0.09	15	BOECKMANN 70	HBC	+	5.0 $\pi^+ p$
0.23 ±0.08	22	ASCOLI 68	HBC	-	5 $\pi^- p$
0.12 ±0.08		CHUNG 68	HBC	-	3.2 $\pi^- p$
0.22 ±0.09		CONTE 67	HBC	-	11.0 $\pi^- p$

Γ_5/Γ_1

NODE=M012R3
NODE=M012R3

$\Gamma(\omega\pi\pi)/\Gamma(3\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.15±0.05 OUR FIT					Error includes scale factor of 1.3.
0.15±0.05 OUR AVERAGE					Error includes scale factor of 1.3. See the ideogram below.
0.28±0.09	60	DIAZ 74	DBC	0	6 $\pi^+ n$
0.18±0.08		¹ KARSHON 74	HBC		Avg. of above two
0.10±0.05	279	² CHALOUPKA 73	HBC	-	3.9 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.29±0.08	140	¹ KARSHON 74	HBC	0	4.9 $\pi^+ p$
0.10±0.04	60	¹ KARSHON 74	HBC	+	4.9 $\pi^+ p$
0.19±0.08		DEFOIX 73	HBC	0	0.7 $\bar{p}p$

Γ_6/Γ_1

NODE=M012R12
NODE=M012R12

¹ KARSHON 74 suggest an additional $I = 0$ state strongly coupled to $\omega\pi\pi$ which could explain discrepancies in branching ratios and masses. We use a central value and a systematic spread.


² Decays to $b_1(1040)\pi, b_1 \rightarrow \omega\pi$. Error increased to account for possible systematic errors of complicated analysis.

OCCUR=3

OCCUR=2

NODE=M012R12;LINKAGE=K

NODE=M012R12;LINKAGE=01



$\Gamma(K\bar{K})/\Gamma(3\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.070±0.012 OUR FIT					
0.078±0.017		CHABAUD 78	RVUE		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.011±0.003		¹ BERTIN 98B	OBLX		0.0 $\bar{p}p \rightarrow K^\pm K_s \pi^\mp$
0.056±0.014	50	² CHALOUPKA 73	HBC	-	3.9 $\pi^- p$
0.097±0.018	113	² ALSTON-... 71	HBC	+	7.0 $\pi^+ p$
0.06 ±0.03		² ABRAMOVI... 70B	HBC	-	3.93 $\pi^- p$
0.054±0.022		² CHUNG 68	HBC	-	3.2 $\pi^- p$

Γ_7/Γ_1

NODE=M012R1
NODE=M012R1

¹ Using 4π data from BERTIN 97D.

² Included in CHABAUD 78 review.

NODE=M012R1;LINKAGE=BE
NODE=M012R1;LINKAGE=C

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$ Γ_7/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.08 ± 0.02	¹ BERTIN	98B	OBLX 0.0 $\bar{p}p \rightarrow K^\pm K_S \pi^\mp$
-------------	---------------------	-----	---

¹ Using $\eta\pi\pi$ data from AMSLER 94D.

NODE=M012R14
NODE=M012R14

NODE=M012R14;LINKAGE=BE

 $\Gamma(\eta\pi)/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$ $\Gamma_5/(\Gamma_1+\Gamma_5+\Gamma_7)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

0.162 ± 0.012 OUR FIT

0.140 ± 0.028 OUR AVERAGE

0.13 ± 0.04		ESPIGAT	72	HBC	± 0.0 $\bar{p}p$
0.15 ± 0.04	34	BARNHAM	71	HBC	+ 3.7 $\pi^+ p$

NODE=M012R2
NODE=M012R2

 $\Gamma(K\bar{K})/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$ $\Gamma_7/(\Gamma_1+\Gamma_5+\Gamma_7)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

0.054 ± 0.009 OUR FIT

0.048 ± 0.012 OUR AVERAGE

0.05 ± 0.02		TOET	73	HBC	+ 5 $\pi^+ p$
0.09 ± 0.04		TOET	73	HBC	0 5 $\pi^+ p$
0.03 ± 0.02	8	¹ DAMERI	72	HBC	- 11 $\pi^- p$
0.06 ± 0.03	17	BARNHAM	71	HBC	+ 3.7 $\pi^+ p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.020 ± 0.004		² ESPIGAT	72	HBC	± 0.0 $\bar{p}p$
---------------	--	----------------------	----	-----	------------------

¹ Montanet agrees. Vlada.

² Not averaged because of discrepancy between masses from $K\bar{K}$ and $\rho\pi$ modes.

NODE=M012R8
NODE=M012R8

OCCUR=2

NODE=M012R8;LINKAGE=01
NODE=M012R8;LINKAGE=A

 $\Gamma(\eta'(958)\pi)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.006	95	ALDE	92B	GAM2	38,100 $\pi^- p \rightarrow \eta' \pi^0 n$
<0.02	97	BARNHAM	71	HBC	+ 3.7 $\pi^+ p$
0.004 ± 0.004		¹ BOESEBECK	68	HBC	+ 8 $\pi^+ p$

¹ No longer valid since $\Gamma(K\bar{K})/\Gamma(3\pi)$ value has changed (MORRISON 71).

NODE=M012R4
NODE=M012R4

NODE=M012R4;LINKAGE=B

 $\Gamma(\eta'(958)\pi)/\Gamma(3\pi)$ Γ_8/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.011	90	EISENSTEIN	73	HBC	- 5 $\pi^- p$
<0.04		ALSTON-...	71	HBC	+ 7.0 $\pi^+ p$
0.04 $\begin{smallmatrix} +0.03 \\ -0.04 \end{smallmatrix}$		BOECKMANN	70	HBC	0 5.0 $\pi^+ p$

NODE=M012R5
NODE=M012R5

 $\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$ Γ_8/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.038 ± 0.005 OUR AVERAGE

0.05 ± 0.02	ADOLPH	15	COMP 191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
0.032 ± 0.009	ABELE	97C	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta'$
0.047 ± 0.010 ± 0.004	¹ BELADIDZE	93	VES 37 $\pi^- N \rightarrow a_2^- N$
0.034 ± 0.008 ± 0.005	BELADIDZE	92	VES 36 $\pi^- C \rightarrow a_2^- C$

¹ Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = 0.441$, $B(\eta \rightarrow \gamma\gamma) = 0.389$ and $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 0.236$.

NODE=M012R13
NODE=M012R13

NODE=M012R13;LINKAGE=A

 $\Gamma(\pi^\pm \gamma)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.005 $\begin{smallmatrix} +0.005 \\ -0.003 \end{smallmatrix}$	¹ EISENBERG	72	HBC 4.3,5.25,7.5 γp
--	------------------------	----	-----------------------------

¹ Pion-exchange model used in this estimation.

NODE=M012R11
NODE=M012R11

NODE=M012R11;LINKAGE=R

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90	ACHASOV	00K	SND $e^+ e^- \rightarrow \pi^0 \pi^0$
----	----	---------	-----	---------------------------------------

NODE=M012R15
NODE=M012R15

a₂(1320) REFERENCES

NODE=M012

ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=55911
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48317
MOLCHANOV	01	PL B521 171	V.V. Molchanov <i>et al.</i>	(FNAL SELEX Collab.)	REFID=48559
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46345
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
ABELE	97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45531
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45761
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=45418
THOMPSON	97	PRL 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)	REFID=45584
AMELIN	96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44649
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
AOYAGI	93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)	REFID=43599
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BELADIDZE	93	PL B313 276	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43598
CONDO	93	PR D48 3045	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=43600
ALDE	92B	ZPHY C54 549	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41852
BELADIDZE	92	ZPHY C54 235	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42171
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch	(WA76 Coll.)	REFID=41375
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41356
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
Translated from YAF 48 436.					
ALTHOFF	86	ZPHY C31 537	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21287
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
BERGER	84C	PL 149B 427	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=21286
BEHREND	83B	PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
CIHANGIR	82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)	REFID=21280
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
DELFOSSÉ	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
CHABAUD	80	NP B175 189	V. Chabaud <i>et al.</i>	(CERN, MPIM, AMST)	REFID=21274
DAUM	80C	PL 89B 276	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21275
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
CHABAUD	78	NP B145 349	V. Chabaud <i>et al.</i>	(CERN, MPIM)	REFID=21267
FERRERSORIA	78	PL 74B 287	A. Ferrer Soria <i>et al.</i>	(ORSAY, CERN, CDEF+)	REFID=21270
HYAMS	78	NP B146 303	B.D. Hyams <i>et al.</i>	(CERN, MPIM, ATEN)	REFID=21271
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA) JP	REFID=21272
MAY	77	PR D16 1983	E.N. May <i>et al.</i>	(ROCH, CORN)	REFID=20450
FORINO	76	NC 35A 465	A. Forino <i>et al.</i>	(BGNA, FIRZ, GENO, MILA+)	REFID=21259
MARGULIE	76	PR D14 667	M. Margulies <i>et al.</i>	(BNL, CUNY)	REFID=21261
EMMS	75	PL 58B 117	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL) JP	REFID=21254
WAGNER	75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP	REFID=20843
DIAZ	74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)	REFID=21248
KARSHON	74	PRL 32 852	U. Karshon <i>et al.</i>	(REHO)	REFID=21249
ANTIPOV	73	NP B63 175	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=21238
ANTIPOV	73C	NP B63 153	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=20817
CHALOUPKA	73	PL 44B 211	V. Chaloupka <i>et al.</i>	(CERN)	REFID=21242
CONFORTO	73	PL 45B 154	G. Conforto <i>et al.</i>	(EFI, FNAL, TNTO+)	REFID=21243
DEFOIX	73	PL 43B 141	C. Defoix <i>et al.</i>	(CDEF)	REFID=21244
EISENSTEIN	73	PR D7 278	L. Eisenstein <i>et al.</i>	(ILL)	REFID=21245
KEY	73	PRL 30 503	A.W. Key <i>et al.</i>	(TNTO, EFI, FNAL, WISC)	REFID=21246
TOET	73	NP B63 248	D.Z. Toet <i>et al.</i>	(NIJM, BONN, DURH, TORI)	REFID=20714
DAMERI	72	NC 9A 1	M. Dameri <i>et al.</i>	(GENO, MILA, SACL)	REFID=20338
EISENBERG	72	PR D5 15	Y. Eisenberg <i>et al.</i>	(REHO, SLAC, TELA)	REFID=20098
ESPIGAT	72	NP B36 93	P. Espigat <i>et al.</i>	(CERN, CDEF)	REFID=21232
FOLEY	72	PR D6 747	K.J. Foley <i>et al.</i>	(BNL, CUNY)	REFID=21233
ALSTON-...	71	PL 34B 156	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21214
BARNHAM	71	PRL 26 1494	K.W.J. Barnham <i>et al.</i>	(LBL)	REFID=21215
BINNIE	71	PL 36B 257	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=21217
BOWEN	71	PRL 26 1663	D.R. Bowen <i>et al.</i>	(NEAS, STON)	REFID=21219
GRAYR	71	PL 34B 333	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=21223
ABRAMOVI...	70B	NP B23 466	M. Abramovich <i>et al.</i>	(CERN) JP	REFID=21195
ALSTON-...	70	PL 33B 607	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21196
BOECKMANN	70	NP B16 221	K. Boeckmann <i>et al.</i>	(BONN, DURH, NIJM+)	REFID=21202
ASCOLI	68	PRL 20 1321	G. Ascoli <i>et al.</i>	(ILL) JP	REFID=21171
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
CHUNG	68	PR 165 1491	S.U. Chung <i>et al.</i>	(LRL)	REFID=20059
CONTE	67	NC 51A 175	F. Conte <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=21166

$f_0(1370)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See also the mini-reviews on scalar mesons under $f_0(500)$ (see the index for the page number) and on non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M147

NODE=M147

 $f_0(1370)$ T-MATRIX POLE POSITION

NODE=M147PP

Note that $\Gamma \approx 2 \text{Im}(\sqrt{s_{\text{pole}}})$.

NODE=M147PP

NODE=M147PP

→ UNCHECKED ←

VALUE (MeV) DOCUMENT ID TECN COMMENT

(1200–1500)– i (150–250) OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(1290 \pm 50) - i(170^{+20}_{-40})$	1 ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$	
$(1373 \pm 15) - i(137 \pm 10)$	2 BARGIOTTI	03	OBLX	$\bar{p}p$	
$(1302 \pm 17) - i(166 \pm 18)$	3 BARBERIS	00C		$450 \bar{p}p \rightarrow p_f 4\pi p_s$	
$(1312 \pm 25 \pm 10) - i(109 \pm 22 \pm 15)$	BARBERIS	99D	OMEG	$450 \bar{p}p \rightarrow K^+ K^-, \pi^+ \pi^-$	
$(1406 \pm 19) - i(80 \pm 6)$	4 KAMINSKI	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$	OCCUR=2
$(1300 \pm 20) - i(120 \pm 20)$	ANISOVICH	98B	RVUE	Compilation	
$(1290 \pm 15) - i(145 \pm 15)$	BARBERIS	97B	OMEG	$450 \bar{p}p \rightarrow p\bar{p}2(\pi^+ \pi^-)$	
$(1548 \pm 40) - i(560 \pm 40)$	BERTIN	97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
$(1380 \pm 40) - i(180 \pm 25)$	ABELE	96B	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$	
$(1300 \pm 15) - i(115 \pm 8)$	BUGG	96	RVUE		
$(1330 \pm 50) - i(150 \pm 40)$	5 AMSLER	95B	CBAR	$\bar{p}p \rightarrow 3\pi^0$	
$(1360 \pm 35) - i(150-300)$	5 AMSLER	95C	CBAR	$\bar{p}p \rightarrow \pi^0 \eta\eta$	
$(1390 \pm 30) - i(190 \pm 40)$	6 AMSLER	95D	CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$	OCCUR=2
1346 – i 249	7,8 JANSSEN	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	OCCUR=2
1214 – i 168	8,9 TORNQVIST	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$	
1364 – i 139	AMSLER	94D	CBAR	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta$	
$(1365^{+20}_{-55}) - i(134 \pm 35)$	ANISOVICH	94	CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$	
$(1340 \pm 40) - i(127^{+30}_{-20})$	10 BUGG	94	RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0 \pi^0$	OCCUR=2
$(1430 \pm 5) - i(73 \pm 13)$	11 KAMINSKI	94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
1420 – i 220	12 AU	87	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	

1 Another pole is found at $(1510 \pm 130) - i(800^{+100}_{-150})$ MeV.

2 Coupled channel analysis of $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0,$ and $K^\pm K_S^0 \pi^\mp$.

3 Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

4 T-matrix pole on sheet – – –.

5 Supersedes ANISOVICH 94.

6 Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta,$ and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

7 Analysis of data from FALVARD 88.

8 The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

9 Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

10 Reanalysis of ANISOVICH 94 data.

11 T-matrix pole on sheet III.

12 Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

NODE=M147PP;LINKAGE=AO

NODE=M147PP;LINKAGE=BG

NODE=M147PP;LINKAGE=PC

NODE=M147PP;LINKAGE=TK

NODE=M147PP;LINKAGE=K

NODE=M147PP;LINKAGE=A

NODE=M147PP;LINKAGE=C

NODE=M147PP;LINKAGE=DD

NODE=M147PP;LINKAGE=BB

NODE=M147PP;LINKAGE=C1

NODE=M147PP;LINKAGE=KM

NODE=M147PP;LINKAGE=H

 $f_0(1370)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETER

NODE=M147205

VALUE (MeV) DOCUMENT ID

1200 to 1500 OUR ESTIMATE

NODE=M147M

→ UNCHECKED ←

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1400±40		¹ AUBERT 09L	BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
1470 ⁺⁶⁺⁷² ₋₇₋₂₅₅		² UEHARA 08A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1259±55	2.6k	BONVICINI 07	CLEO	$D^+ \rightarrow \pi^-\pi^+\pi^+$
1309±1±15		³ BUGG 07A	RVUE	0.0 $p\bar{p} \rightarrow 3\pi^0$
1449±13	4.3k	⁴ GARMASH 06	BELL	$B^+ \rightarrow K^+\pi^+\pi^-$
1350±50		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
1265±30 ⁺²⁰ ₋₃₅		ABLIKIM 05Q	BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
1434±18±9	848	AITALA 01A	E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$
1308±10		BARBERIS 99B	OMEG	450 $pp \rightarrow p_s p_f \pi^+\pi^-$
1315±50		BELLAZZINI 99	GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
1315±30		ALDE 98	GAM4	100 $\pi^-p \rightarrow \pi^0\pi^0n$
1280±55		BERTIN 98	OBLX	0.05-0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
1186		^{5,6} TORNQVIST 95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472±12		ARMSTRONG 91	OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275±20		BREAKSTONE90	SFM	62 $pp \rightarrow pp\pi^+\pi^-$
1420±20		AKESSON 86	SPEC	63 $pp \rightarrow pp\pi^+\pi^-$
1256		FROGGATT 77	RVUE	$\pi^+\pi^-\text{ channel}$

¹ Breit-Wigner mass.² Breit-Wigner mass. May also be the $f_0(1500)$.³ Reanalysis of ABELE 96C data.⁴ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0\pi^+\pi^-$ decays. Supersedes GARMASH 05.⁵ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.⁶ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decaysNODE=M147M1
NODE=M147M1NODE=M147M1;LINKAGE=BW
NODE=M147M1;LINKAGE=UE
NODE=M147M1;LINKAGE=BU
NODE=M147M1;LINKAGE=GR
NODE=M147M1;LINKAGE=BB

NODE=M147M1;LINKAGE=FF

 $K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1360±31±28	430	^{1,2} DOBBS 15		$J/\psi \rightarrow \gamma K^+K^-$
1350±48±15	168	^{1,2} DOBBS 15		$\psi(2S) \rightarrow \gamma K^+K^-$
1440±6		VLADIMIRSK...06	SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1391±10		TIKHOMIROV 03	SPEC	40.0 $\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440±50		BOLONKIN 88	SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1463±9		ETKIN 82B	MPS	23 $\pi^-p \rightarrow n2K_S^0$
1425±15		WICKLUND 80	SPEC	6 $\pi N \rightarrow K^+K^-N$
~1300		POLYCHRO... 79	STRC	7 $\pi^-p \rightarrow n2K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² From a fit to a Breit-Wigner line shape with fixed $\Gamma = 346$ MeV.NODE=M147M2
NODE=M147M2

OCCUR=2

NODE=M147M2;LINKAGE=A
NODE=M147M2;LINKAGE=B **4π MODE $2(\pi\pi)_S + \rho\rho$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1395±40		ABELE 01	CBAR	0.0 $\bar{p}d \rightarrow \pi^-4\pi^0p$
1374±38		AMSLER 94	CBAR	0.0 $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
1345±12		ADAMO 93	OBLX	$\bar{n}p \rightarrow 3\pi^+2\pi^-$
1386±30		GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$
~1410	5751	¹ BETTINI 66	DBC	0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$

¹ $\rho\rho$ dominant.NODE=M147M3
NODE=M147M3

NODE=M147M3;LINKAGE=BE

 $\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1262 ⁺⁵¹⁺⁸² ₋₇₈₋₁₀₃	¹ UEHARA 10A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
1430	AMSLER 92	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
1220±40	ALDE 86D	GAM4	100 $\pi^-p \rightarrow n2\eta$

¹ Breit-Wigner mass. May also be the $f_0(1500)$.NODE=M147M4
NODE=M147M4

NODE=M147M4;LINKAGE=UE

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN
-------------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1306±20	¹ ANISOVICH	03	RVUE
---------	------------------------	----	------

¹K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

NODE=M147M5
NODE=M147M5

NODE=M147M;LINKAGE=KM

 $f_0(1370)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID
-------------	-------------

200 to 500 OUR ESTIMATE

NODE=M147210

NODE=M147W
→ UNCHECKED ←

 $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

300± 80		¹ AUBERT	09L	BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
90 ⁺ ₋ 2 ⁺ ₁₋ 50 ₂₂		² UEHARA	08A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
298± 21	2.6k	BONVICINI	07	CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
126± 25	4286	³ GARMASH	06	BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
265± 40		ABLIKIM	05	BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
350±100 ⁺ ₋ 105 ₆₀		ABLIKIM	05Q	BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
173± 32± 6	848	AITALA	01A	E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
222± 20		BARBERIS	99B	OMEG	450 $pp \rightarrow p_s p_f \pi^+ \pi^-$
255± 60		BELLAZZINI	99	GAM4	450 $pp \rightarrow pp \pi^0 \pi^0$
190± 50		ALDE	98	GAM4	100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
323± 13		BERTIN	98	OBLX	0.05–0.405 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
350		^{4,5} TORNQVIST	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
195± 33		ARMSTRONG	91	OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$
285± 60		BREAKSTONE	90	SFM	62 $pp \rightarrow pp\pi^+ \pi^-$
460± 50		AKESSON	86	SPEC	63 $pp \rightarrow pp\pi^+ \pi^-$
~ 400		⁶ FROGGATT	77	RVUE	$\pi^+ \pi^-$ channel

¹ The systematic errors are not reported.

² Breit-Wigner width. May also be the $f_0(1500)$.

³ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.

⁴ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CA-SON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

⁵ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays

⁶ Width defined as distance between 45 and 135° phase shift.

NODE=M147W1;LINKAGE=NS
NODE=M147W1;LINKAGE=UE
NODE=M147W1;LINKAGE=GR
NODE=M147W1;LINKAGE=BB

NODE=M147W1;LINKAGE=FF
NODE=M147W1;LINKAGE=E

 $K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

121± 15		VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
55± 26		TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
250± 80		BOLONKIN 88	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
118 ⁺ ₋ 138 ₁₆		ETKIN 82B	MPS	23 $\pi^- p \rightarrow n 2K_S^0$
160± 30		WICKLUND 80	SPEC	6 $\pi N \rightarrow K^+ K^- N$
~ 150		POLYCHRO... 79	STRC	7 $\pi^- p \rightarrow n 2K_S^0$

NODE=M147W2
NODE=M147W2

 4π MODE $2(\pi\pi)_S + \rho\rho$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

275±55		ABELE	01	CBAR	0.0 $\bar{p} d \rightarrow \pi^- 4\pi^0 p$
375±61		AMSLER	94	CBAR	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- 3\pi^0$
398±26		ADAMO	93	OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
310±50		GASPERO	93	DBC	0.0 $\bar{p} n \rightarrow 2\pi^+ 3\pi^-$
~ 90	5751	¹ BETTINI	66	DBC	0.0 $\bar{p} n \rightarrow 2\pi^+ 3\pi^-$

¹ $\rho\rho$ dominant.

NODE=M147W3
NODE=M147W3

NODE=M147W3;LINKAGE=BE

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$484^{+246+246}_{-170-263}$	¹ UEHARA	10A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
250	AMSLER	92	CBAR 0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
320 ± 40	ALDE	86D	GAM4 100 $\pi^-p \rightarrow n2\eta$

¹Breit-Wigner width. May also be the $f_0(1500)$.

NODE=M147W4
NODE=M147W4

NODE=M147W4;LINKAGE=UE

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN
-------------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

147^{+30}_{-50}	¹ ANISOVICH	03	RVUE
-------------------	------------------------	----	------

¹K-matrix pole from combined analysis of $\pi^-p \rightarrow \pi^0\pi^0n$, $\pi^-p \rightarrow K\bar{K}n$, $\pi^+\pi^- \rightarrow \pi^+\pi^-$, $\bar{p}p \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$, $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, $K_S^0K_S^0\pi^0$, $K^+K_S^0\pi^-$ at rest, $\bar{p}n \rightarrow \pi^-\pi^-\pi^+$, $K_S^0K^-\pi^0$, $K_S^0K_S^0\pi^-$ at rest.

NODE=M147W5
NODE=M147W5

NODE=M147W;LINKAGE=KM

 $f_0(1370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 4π	seen
Γ_3 $4\pi^0$	seen
Γ_4 $2\pi^+2\pi^-$	seen
Γ_5 $\pi^+\pi^-2\pi^0$	seen
Γ_6 $\rho\rho$	dominant
Γ_7 $2(\pi\pi)_{S\text{-wave}}$	seen
Γ_8 $\pi(1300)\pi$	seen
Γ_9 $a_1(1260)\pi$	seen
Γ_{10} $\eta\eta$	seen
Γ_{11} $K\bar{K}$	seen
Γ_{12} $K\bar{K}n\pi$	not seen
Γ_{13} 6π	not seen
Γ_{14} $\omega\omega$	not seen
Γ_{15} $\gamma\gamma$	seen
Γ_{16} e^+e^-	not seen

NODE=M147215;NODE=M147

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=10;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←
DESIG=14;OUR EST;→ UNCHECKED ←
DESIG=15;OUR EST;→ UNCHECKED ←
DESIG=16;OUR EVAL;→ UNCHECKED ←
DESIG=17;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=11;OUR EST;→ UNCHECKED ←
DESIG=18;OUR EVAL;→ UNCHECKED ←
DESIG=19;OUR EVAL;→ UNCHECKED ←
DESIG=20;OUR EVAL;→ UNCHECKED ←
DESIG=12;OUR EST;→ UNCHECKED ←
DESIG=13;OUR EST;→ UNCHECKED ←

 $f_0(1370)$ PARTIAL WIDTHS **$\Gamma(\gamma\gamma)$**

See $\gamma\gamma$ widths under $f_0(500)$ and MORGAN 90.

 Γ_{15}

NODE=M147217

NODE=M147W11
NODE=M147W11
NODE=M147W11

 $\Gamma(e^+e^-)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	VOROBYEV	88	ND $e^+e^- \rightarrow \pi^0\pi^0$

 Γ_{16}

NODE=M147W12
NODE=M147W12

 $f_0(1370)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M147225

 $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$121^{+133+169}_{-53-106}$	¹ UEHARA	10A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
----------------------------	---------------------	-----	---

¹Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

NODE=M147G01
NODE=M147G01

NODE=M147G01;LINKAGE=UE

$f_0(1370)$ BRANCHING RATIOS

NODE=M147220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ **Γ_1/Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R3
NODE=M147R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.09	BUGG	96	RVUE
< 0.15	¹ AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
< 0.06	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$

¹ Using AMSLER 95B ($3\pi^0$).

NODE=M147R3;LINKAGE=B

 $\Gamma(4\pi)/\Gamma_{\text{total}}$ **$\Gamma_2/\Gamma = (\Gamma_3 + \Gamma_4 + \Gamma_5)/\Gamma$**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R4
NODE=M147R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 0.72	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
----------	---------	----	---

 $\Gamma(4\pi^0)/\Gamma(4\pi)$ **Γ_3/Γ_2**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R12
NODE=M147R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ABELE	96	CBAR $0.0 \bar{p}p \rightarrow 5\pi^0$
0.068 ± 0.005	¹ GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$

¹ Model-dependent evaluation.

NODE=M147R12;LINKAGE=GA

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma(4\pi)$ **$\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3 + \Gamma_4 + \Gamma_5)$**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R5
NODE=M147R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.420 ± 0.014	¹ GASPERO	93	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
-------------------	----------------------	----	--

¹ Model-dependent evaluation.

NODE=M147R5;LINKAGE=A

 $\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(4\pi)$ **$\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3 + \Gamma_4 + \Gamma_5)$**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R6
NODE=M147R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.512 ± 0.019	¹ GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
-------------------	----------------------	----	---

¹ Model-dependent evaluation.

NODE=M147R6;LINKAGE=A

 $\Gamma(\rho\rho)/\Gamma(4\pi)$ **Γ_6/Γ_2**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R17
NODE=M147R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.07	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi p$
-----------------	-------	-----	--

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(\pi\pi)$ **Γ_7/Γ_1**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R15
NODE=M147R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.6 ± 2.6	¹ ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
---------------	--------------------	----	--

¹ From the combined data of ABELE 96 and ABELE 96C.

NODE=M147R;LINKAGE=KZ

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$ **Γ_7/Γ_2**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R16
NODE=M147R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.51 ± 0.09	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi p$
-----------------	-------	-----	--

 $\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{S\text{-wave}})$ **Γ_6/Γ_7**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R10
NODE=M147R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

large	BARBERIS	00C	$450 pp \rightarrow p_f 4\pi p_s$
1.6 ± 0.2	AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
~ 0.65	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$

OCCUR=2

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ **Γ_8/Γ_2**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R18
NODE=M147R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ± 0.06	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi p$
-----------------	-------	-----	--

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.06±0.02	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$
-----------	-------	-----	--

 Γ_9/Γ_2

NODE=M147R19
NODE=M147R19

 $\Gamma(\eta\eta)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(28 \pm 11) \times 10^{-3}$	¹ ANISOVICH	02D	SPEC Combined fit
$(4.7 \pm 2.0) \times 10^{-3}$	BARBERIS	00E	450 $p\bar{p} \rightarrow p_f \eta \eta p_S$

¹From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta \eta n$, $\eta \eta' n$), and BNL ($\pi p \rightarrow K \bar{K} n$) data.

NODE=M147R14
NODE=M147R14

NODE=M147R14;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.35±0.13	BUGG	96	RVUE
-----------	------	----	------

 Γ_{11}/Γ

NODE=M147R11
NODE=M147R11

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.08±0.08	ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$, $\phi K^+ K^-$
0.91±0.20	¹ BARGIOTTI	03	OBLX $\bar{p}p$
0.12±0.06	² ANISOVICH	02D	SPEC Combined fit
0.46±0.15±0.11	BARBERIS	99D	OMEG 450 $p\bar{p} \rightarrow K^+ K^-$, $\pi^+ \pi^-$

¹Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K_S^0 \pi^+ \pi^-$.

²From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta \eta n$, $\eta \eta' n$), and BNL ($\pi p \rightarrow K \bar{K} n$) data.

 Γ_{11}/Γ_1

NODE=M147R13
NODE=M147R13

NODE=M147R;LINKAGE=BG
NODE=M147R;LINKAGE=CH

 $\Gamma(K\bar{K}n\pi)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.03	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	--

 Γ_{12}/Γ

NODE=M147R20
NODE=M147R20

 $\Gamma(6\pi)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.22	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	--

 Γ_{13}/Γ

NODE=M147R21
NODE=M147R21

 $\Gamma(\omega\omega)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	--

 Γ_{14}/Γ

NODE=M147R22
NODE=M147R22

 $f_0(1370)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		REFID=52719
AUBERT	09L	PR D79 072006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52723
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=51721
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>		REFID=53252
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51594
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51162
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=50641
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
		Translated from YAF 65 1583.			

NODE=M147

ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48334
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
ANISOVICH	98B	Translated from SPU 41 419	YAF 62 446	V.V. Anisovich <i>et al.</i>	REFID=46331
		Translated from UFN 168 481.			
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45011
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45038
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45076
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)	REFID=45094
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
TORNQVIST	95	ZPHY C68 647	N.A. Tornqvist	(HELS)	REFID=44529
AMSLER	94	PL B322 431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC	REFID=43660
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) JPC	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC	REFID=43657
GASPERO	93	NP A562 407	M. Gaspero	(ROMAI) JPC	REFID=43658
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)	REFID=40064
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)	REFID=21123
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20383
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)	REFID=11004
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)	REFID=20349
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)	REFID=44530
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)	REFID=21361

$h_1(1380)$

$$I^G(J^{PC}) = ?^-(1^{+-})$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K\bar{K}\pi$ system. Needs confirmation.

NODE=M109

NODE=M109

 $h_1(1380)$ MASS

NODE=M109M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1407 ± 12 OUR AVERAGE 2015 AVERAGE]	Error includes scale factor of 1.5. [1386 ± 19 MeV OUR		
1412 ± 4 ± 8	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1,2} \rightarrow \gamma K^* \bar{K}$
1440 ± 60	ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$
1380 ± 20	ASTON	88C LASS	11 $K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

NODE=M109M

NEW

 $h_1(1380)$ WIDTH

NODE=M109W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
89 ± 23 OUR AVERAGE [91 ± 30 MeV OUR 2015 AVERAGE Scale factor = 1.1]			
84 ± 12 ± 40	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1,2} \rightarrow \gamma K^* \bar{K}$
170 ± 80	ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$
80 ± 30	ASTON	88C LASS	11 $K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

NODE=M109W

NEW

 $h_1(1380)$ DECAY MODES

NODE=M109215;NODE=M109

Mode

 $\Gamma_1 \quad K\bar{K}^*(892) + \text{c.c.}$

DESIG=1

 $h_1(1380)$ REFERENCES

ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56778
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45765
ASTON	88C	PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40282

NODE=M109

REFID=56778
REFID=45765
REFID=40282 **$\pi_1(1400)$**

$$I^G(J^{PC}) = 1^-(1^{-+})$$

NODE=M111

NODE=M111

See also the mini-review under non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006). **$\pi_1(1400)$ MASS**


NODE=M111M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1354 ± 25 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.			
1257 ± 20 ± 25	23.5k	ADAMS	07B B852		18 $\pi^- p \rightarrow \eta \pi^0 n$
1384 ± 20 ± 35	90k	SALVINI	04 OBLX		$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
1360 ± 25		ABELE	99 CBAR		0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
1400 ± 20 ± 20		ABELE	98B CBAR		0.0 $\bar{p}n \rightarrow \pi^- \pi^0 \eta$
1370 ± 16 $\begin{smallmatrix} +50 \\ -30 \end{smallmatrix}$		¹ THOMPSON	97 MPS		18 $\pi^- p \rightarrow \eta \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1323.1 ± 4.6		² AOYAGI	93 BKEI		$\pi^- p \rightarrow \eta \pi^- p$
1406 ± 20		³ ALDE	88B GAM4 0		100 $\pi^- p \rightarrow \eta \pi^0 n$

NODE=M111M

- ¹ Natural parity exchange, questioned by DZIERBA 03.
- ² Unnatural parity exchange.
- ³ Seen in the P_0 -wave intensity of the $\eta\pi^0$ system, unnatural parity exchange.

NODE=M111M;LINKAGE=B
 NODE=M111M;LINKAGE=C
 NODE=M111M;LINKAGE=A



$\pi_1(1400)$ WIDTH

NODE=M111W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
330 ±35	OUR AVERAGE				
354 ±64 ± 58	23.5k	ADAMS	07B	B852	18 $\pi^- p \rightarrow \eta\pi^0 n$
378 ±50 ± 50	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
220 ±90		ABELE	99	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\eta$
310 ±50 + 50 - 30		ABELE	98B	CBAR	0.0 $\bar{p}n \rightarrow \pi^- \pi^0 \eta$
385 ±40 + 65 - 105		⁴ THOMPSON	97	MPS	18 $\pi^- p \rightarrow \eta\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

143.2±12.5		⁵ AOYAGI	93	BKEI	$\pi^- p \rightarrow \eta\pi^- p$
180 ±20		⁶ ALDE	88B	GAM4 0	100 $\pi^- p \rightarrow \eta\pi^0 n$

NODE=M111W

- ⁴ Resolution is not unfolded, natural parity exchange, questioned by DZIERBA 03.
- ⁵ Unnatural parity exchange.
- ⁶ Seen in the P_0 -wave intensity of the $\eta\pi^0$ system, unnatural parity exchange.

NODE=M111W;LINKAGE=QQ
 NODE=M111W;LINKAGE=C
 NODE=M111W;LINKAGE=A

$\pi_1(1400)$ DECAY MODES

NODE=M111215;NODE=M1111

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi^0$	seen
Γ_2 $\eta\pi^-$	seen
Γ_3 $\eta'\pi$	

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=3

$\pi_1(1400)$ BRANCHING RATIOS

NODE=M111220

$\Gamma(\eta\pi^0)/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_i/Γ
not seen	PROKOSHKIN 95B	GAM4		100 $\pi^- p \rightarrow \eta\pi^0 n$	
not seen	⁷ BUGG	94	RVUE	$\bar{p}p \rightarrow \eta 2\pi^0$	
not seen	⁸ APEL	81	NICE 0	40 $\pi^- p \rightarrow \eta\pi^0 n$	

NODE=M111R1
 NODE=M111R1

- ⁷ Using Crystal Barrel data.
- ⁸ A general fit allowing S , D , and P waves (including $m=0$) is not done because of limited statistics.

NODE=M111R1;LINKAGE=C
 NODE=M111R1;LINKAGE=B

$\Gamma(\eta\pi^-)/\Gamma_{total}$

Γ_2/Γ

VALUE DOCUMENT ID TECN COMMENT

NODE=M111R4
NODE=M111R4

••• We do not use the following data for averages, fits, limits, etc. •••

possibly seen BELADIDZE 93 VES $37\pi^- N \rightarrow \eta\pi^- N$

$\Gamma(\eta'\pi)/\Gamma(\eta\pi^0)$

Γ_3/Γ_1

VALUE CL% DOCUMENT ID TECN COMMENT

NODE=M111R3
NODE=M111R3

••• We do not use the following data for averages, fits, limits, etc. •••

<0.80 95 BOUTEMEUR 90 GAM4 $100\pi^- p \rightarrow 4\gamma n$

$\pi_1(1400)$ REFERENCES

NODE=M111

ADAMS	07B	PL B657 27	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=52048
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)	REFID=53226
DZIERBA	03	PR D67 094015	A.R. Dzierba <i>et al.</i>		REFID=49412
ABELE	99	PL B446 349	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46602
ABELE	98B	PL B423 175	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45864
THOMPSON	97	PRL 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)	REFID=45584
PROKOSHKIN	95B	PAN 58 606	Y.D. Prokoshkin, S.A. Sadovsky	(SERP)	REFID=44619
		Translated from YAF 58 662.			
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AOYAGI	93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)	REFID=43599
BELADIDZE	93	PL B313 276	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43598
BOUTEMEUR	90	Hadron 89 Conf. p 119	M. BoutemEUR, M. Poulet	(SERP, BELG, LANL+)	REFID=41751
ALDE	88B	PL B205 397	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP) IGJPC	REFID=40558
APEL	81	NP B193 269	W.D. Apel <i>et al.</i>	(SERP, CERN)	REFID=22913

$\eta(1405)$

$I^G(J^{PC}) = 0^+(0^-+)$

NODE=M027

See also the $\eta(1475)$.

A REVIEW GOES HERE – Check our WWW List of Reviews


NODE=M027
NODE=M027

$\eta(1405)$ MASS

NODE=M027205

VALUE (MeV) DOCUMENT ID
1408.8±1.8 OUR AVERAGE Includes data from the 2 datablocks that follow this one.
Error includes scale factor of 2.1. See the ideogram below.

NODE=M027MX



$\eta\pi\pi$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT


NODE=M027M1
NODE=M027M1

The data in this block is included in the average printed for a previous datablock.

1406.2 ± 2.3 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

1409.0 ± 1.7	743	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$	
1407.0 ± 3.5	198	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$	OCCUR=2
1399.8 ± 2.2 ^{+2.8} _{-0.1}		1 ABLIKIM	11J	BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$	
1392 ± 14	900 ± 375	AMSLER	04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$	
1394 ± 8	6.6 ± 2.0k	AMSLER	04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$	OCCUR=2
1404 ± 6	9082	MANAK	00A	MPS	$18 \pi^-p \rightarrow \eta\pi^+\pi^-n$	
1424 ± 6	2200	ALDE	97B	GAM4	$100 \pi^-p \rightarrow \eta\pi^0\pi^0n$	
1409 ± 3		AMSLER	95F	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$	
1400 ± 6		2 BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	
1388 ± 4		FUKUI	91C	SPEC	$8.95 \pi^-p \rightarrow \eta\pi^+\pi^-n$	
1398 ± 6	261	3 AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	
1420 ± 5		ANDO	86	SPEC	$8 \pi^-p \rightarrow \eta\pi^+\pi^-n$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1385 ± 7		BAI	99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	

WEIGHTED AVERAGE
1406.2 ± 2.3 (Error scaled by 2.2)

 **$K\bar{K}\pi$ MODE ($a_0(980)\pi$ or direct $K\bar{K}\pi$)**

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M027M4
NODE=M027M4

The data in this block is included in the average printed for a previous datablock.

1413.9 ± 1.7 OUR AVERAGE Error includes scale factor of 1.1.

1413 ± 14	3651	4 NICHITIU	02	OBLX		
1416 ± 4 ± 2	20k	ADAMS	01B	B852	$18 \text{ GeV } \pi^-p \rightarrow K^+K^-\pi^0n$	
1405 ± 5		5 CICALO	99	OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$	
1407 ± 5		5 BERTIN	97	OBLX	$0 \bar{p}p \rightarrow K^\pm(K^0) \pi^\mp \pi^+ \pi^-$	
1416 ± 2		5 BERTIN	95	OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$	
1416 ± 8 ⁺⁷ ₋₅	700	6 BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	OCCUR=2
1413 ± 5		6 RATH	89	MPS	$21.4 \pi^-p \rightarrow n K_S^0 K_S^0 \pi^0$	OCCUR=3
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1459 ± 5		7 AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$	

 $\pi\pi\gamma$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M027M2
NODE=M027M2

1390 ± 12 235 ± 91 AMSLER 04B CBAR $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1424 ± 10 ± 11	547	BAI	04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	
1401 ± 18		8.9 AUGUSTIN	90	DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$	OCCUR=4
1432 ± 8		9 COFFMAN	90	MRK3	$J/\psi \rightarrow \pi^+\pi^-2\gamma$	

4 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1420 \pm 20		BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1489 \pm 12	3270	¹⁰ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi \gamma$

NODE=M027M3
 NODE=M027M3

K \bar{K} π MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1452.7 \pm 3.3	191	^{11,12} ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$
1437.6 \pm 3.2	249 \pm 35	^{11,12} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$
1445.9 \pm 5.7	62 \pm 18	^{11,12} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
1442 \pm 10	410	¹¹ BAI	98C BES	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1445 \pm 8	693	¹¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1433 \pm 8	296	¹¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1413 \pm 8	500	¹¹ DUCH	89 ASTE	$\bar{p} p \rightarrow \pi^+ \pi^- K^\pm \pi^\mp K^0$
1453 \pm 7	170	¹¹ RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1419 \pm 1	8800	¹¹ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ K^0 \pi^- n$
1424 \pm 3	620	¹¹ REEVES	86 SPEC	$6.6 p \bar{p} \rightarrow K \bar{K} \pi X$
1421 \pm 2		¹¹ CHUNG	85 SPEC	$8 \pi^- p \rightarrow K \bar{K} \pi n$
1440 $\begin{smallmatrix} +20 \\ -15 \end{smallmatrix}$	174	¹¹ EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1440 $\begin{smallmatrix} +10 \\ -15 \end{smallmatrix}$		¹¹ SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1425 \pm 7	800	^{11,13} BAILLON	67 HBC	$0 \bar{p} p \rightarrow K \bar{K} \pi \pi \pi$

NODE=M027M6
 NODE=M027M6

OCCUR=2

OCCUR=2

OCCUR=2

- ¹ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.
- ² From fit to the $a_0(980) \pi 0^-+$ partial wave.
- ³ Best fit with a single Breit Wigner.
- ⁴ Decaying dominantly directly to $K^+ K^- \pi^0$.
- ⁵ Decaying into $(K \bar{K})_S \pi$, $(K \pi)_S \bar{K}$, and $a_0(980) \pi$.
- ⁶ From fit to the $a_0(980) \pi 0^-+$ partial wave. Cannot rule out a $a_0(980) \pi 1^++$ partial wave.
- ⁷ Excluded from averaging because averaging would be meaningless.
- ⁸ Best fit with a single Breit Wigner.
- ⁹ This peak in the $\gamma \rho$ channel may not be related to the $\eta(1405)$.
- ¹⁰ Estimated by us from various fits.
- ¹¹ These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.
- ¹² Systematic uncertainty not evaluated.
- ¹³ From best fit of 0^-+ partial wave, 50% $K^*(892) K$, 50% $a_0(980) \pi$.

NODE=M027M1;LINKAGE=BL

NODE=M027M1;LINKAGE=J1

NODE=M027M1;LINKAGE=A1

NODE=M027M;LINKAGE=NC

NODE=M027M4;LINKAGE=FX

NODE=M027M4;LINKAGE=C2

NODE=M027M4;LINKAGE=AA

NODE=M027M2;LINKAGE=E

NODE=M027M2;LINKAGE=X

NODE=M027M3;LINKAGE=E

NODE=M027M;LINKAGE=NP

NODE=M027M;LINKAGE=NS

NODE=M027M6;LINKAGE=H


 $\eta(1405)$ WIDTH

NODE=M027210

VALUE (MeV) _____ DOCUMENT ID _____

51.0 \pm 2.9 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.8. See the ideogram below.

NODE=M027WX



ηππ MODE


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M027W1
NODE=M027W1

The data in this block is included in the average printed for a previous datablock.

54 ± 4 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

48.3 ± 5.2	743	ABLIKIM	12E BES3	J/ψ → γ(π ⁺ π ⁻ π ⁰)	
55.0 ± 11.0	198	ABLIKIM	12E BES3	J/ψ → γ(π ⁰ π ⁰ π ⁰)	OCCUR=2
52.8 ± 7.6 ^{+0.1} _{-7.6}	14	ABLIKIM	11J BES3	J/ψ → ω(ηπ ⁺ π ⁻)	
55 ± 11	900 ± 375	AMSLER	04B CBAR	0 p̄p → π ⁺ π ⁻ π ⁺ π ⁻ η	
55 ± 12	6.6 ± 2.0k	AMSLER	04B CBAR	0 p̄p → π ⁺ π ⁻ π ⁰ π ⁰ γ	OCCUR=2
80 ± 21	9082	MANAK	00A MPS	18 π ⁻ p → ηπ ⁺ π ⁻ n	
85 ± 18	2200	ALDE	97B GAM4	100 π ⁻ p → ηπ ⁰ π ⁰ n	
86 ± 10		AMSLER	95F CBAR	0 p̄p → π ⁺ π ⁻ π ⁰ π ⁰ η	
47 ± 13	15	BOLTON	92B MRK3	J/ψ → γηπ ⁺ π ⁻	
59 ± 4		FUKUI	91C SPEC	8.95 π ⁻ p → ηπ ⁺ π ⁻ n	
53 ± 11	16	AUGUSTIN	90 DM2	J/ψ → γηπ ⁺ π ⁻	
31 ± 7		ANDO	86 SPEC	8 π ⁻ p → ηπ ⁺ π ⁻ n	



$K\bar{K}\pi$ MODE ($a_0(980)\pi$ or direct $K\bar{K}\pi$)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M027W4
 NODE=M027W4


The data in this block is included in the average printed for a previous datablock.

48 ± 4 OUR AVERAGE

Error includes scale factor of 2.1. See the ideogram below.

51 ± 6	3651	17 NICHITIU	02	OBLX		
42 ± 10 ± 9	20k	ADAMS	01B	B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$	
50 ± 4		CICALO	99	OBLX	0 $\bar{p} p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$	
48 ± 5		18 BERTIN	97	OBLX	0.0 $\bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$	
50 ± 4		18 BERTIN	95	OBLX	0 $\bar{p} p \rightarrow K\bar{K}\pi\pi\pi$	
75 ± 9		AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$	
91 ⁺⁶⁷ ₋₃₁ ± 15 ₋₃₈		19 BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	OCCUR=2
19 ± 7		19 RATH	89	MPS	21.4 $\pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$	OCCUR=3

WEIGHTED AVERAGE
 48±4 (Error scaled by 2.1)



$\eta(1405)$ width $K\bar{K}\pi$ mode ($a_0(980)\pi$ dominant)

 $\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M027W2
 NODE=M027W2

64 ± 18 235 ± 91 AMSLER 04B CBAR 0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

101.0 ± 8.8 ± 8.8	547	BAI	04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	OCCUR=2
174 ± 44		AUGUSTIN	90	DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$	
90 ± 26		20 COFFMAN	90	MRK3	$J/\psi \rightarrow \pi^+\pi^- 2\gamma$	

4 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M027W3
 NODE=M027W3

• • • We do not use the following data for averages, fits, limits, etc. • • •

160 ± 30		BUGG	95	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
144 ± 13	3270	21 BISELLO	89B	DM2	$J/\psi \rightarrow 4\pi\gamma$

 $K\bar{K}\pi$ MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M027W6
 NODE=M027W6

• • • We do not use the following data for averages, fits, limits, etc. • • •

45.9 ± 8.2	191	22,23 ABLIKIM	13M	BES3	$\psi(2S) \rightarrow \omega K K \pi$	
48.9 ± 9.0	249 ± 35	22,23 ABLIKIM	08E	BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$	
34.2 ± 18.5	62 ± 18	22,23 ABLIKIM	08E	BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$	OCCUR=2
93 ± 14	296	22 AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$	OCCUR=2
105 ± 10	693	22 AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	OCCUR=3
62 ± 16	500	22 DUCH	89	ASTE	$\bar{p} p \rightarrow K\bar{K}\pi\pi\pi$	

100 ±11	170	22 RATH	89	MPS	21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
66 ± 2	8800	22 BIRMAN	88	MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
60 ±10	620	22 REEVES	86	SPEC	6.6 $p\bar{p} \rightarrow K K \pi X$
60 ±10		22 CHUNG	85	SPEC	8 $\pi^- p \rightarrow K \bar{K} \pi n$
55 $\begin{smallmatrix} +20 \\ -30 \end{smallmatrix}$	174	22 EDWARDS	82E	CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
50 $\begin{smallmatrix} +30 \\ -20 \end{smallmatrix}$		22 SCHARRE	80	MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
80 ±10	800	22,24 BAILLON	67	HBC	0.0 $\bar{p}p \rightarrow K \bar{K} \pi \pi \pi$

OCCUR=2

14 The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.15 From fit to the $a_0(980) \pi 0^-+$ partial wave.16 From $\eta \pi^+ \pi^-$ mass distribution - mainly $a_0(980) \pi$ - no spin-parity determination available.17 Decaying dominantly directly to $K^+ K^- \pi^0$.18 Decaying into $(K \bar{K})_S \pi$, $(K \pi)_S \bar{K}$, and $a_0(980) \pi$.19 From fit to the $a_0(980) \pi 0^-+$ partial wave, but $a_0(980) \pi 1^++$ cannot be excluded.20 This peak in the $\gamma \rho$ channel may not be related to the $\eta(1405)$.

21 Estimated by us from various fits.

22 These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

23 Systematic uncertainty not evaluated.

24 From best fit to 0^-+ partial wave, 50% $K^*(892) K$, 50% $a_0(980) \pi$.

NODE=M027W1;LINKAGE=BL

NODE=M027W1;LINKAGE=A1

NODE=M027W1;LINKAGE=D1

NODE=M027W;LINKAGE=NC

NODE=M027W4;LINKAGE=FX

NODE=M027W4;LINKAGE=C

NODE=M027W2;LINKAGE=X

NODE=M027W3;LINKAGE=F2

NODE=M027W;LINKAGE=NP

NODE=M027W;LINKAGE=NS

NODE=M027W6;LINKAGE=H1

NODE=M027215;NODE=M027

 $\eta(1405)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K \bar{K} \pi$	seen	
Γ_2 $\eta \pi \pi$	seen	
Γ_3 $a_0(980) \pi$	seen	
Γ_4 $\eta(\pi\pi)_S$ -wave	seen	
Γ_5 $f_0(980) \eta$	seen	
Γ_6 4π	seen	
Γ_7 $\rho \rho$	<58 %	99.85%
Γ_8 $\gamma \gamma$		
Γ_9 $\rho^0 \gamma$	seen	
Γ_{10} $\phi \gamma$		
Γ_{11} $K^*(892) K$	seen	

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=9;OUR EST;→ UNCHECKED ←

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=12

DESIG=7

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=13

DESIG=11;OUR EST;→ UNCHECKED ←

 $\eta(1405) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

NODE=M027220

$\Gamma(K \bar{K} \pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$					$\Gamma_1 \Gamma_8 / \Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027G3

NODE=M027G3

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.035	90	25,26 AHOHE	05	CLE2	10.6 $e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$
--------	----	-------------	----	------	--

$\Gamma(\eta \pi \pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$					$\Gamma_2 \Gamma_8 / \Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027G5

NODE=M027G5

<0.095	95	ACCIARRI	01G	L3	183–202 $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
--------	----	----------	-----	----	--

$\Gamma(\rho^0 \gamma) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$					$\Gamma_9 \Gamma_8 / \Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027G8

NODE=M027G8

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	95	ALTHOFF	84E	TASS	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \gamma$
------	----	---------	-----	------	--

25 Using $\eta(1405)$ mass and width 1410 MeV and 51 MeV, respectively.26 Assuming three-body phase-space decay to $K_S^0 K^\pm \pi^\mp$.

NODE=M027G3;LINKAGE=AH

NODE=M027G3;LINKAGE=B3

 $\eta(1405)$ BRANCHING RATIOS

NODE=M027225

$\Gamma(\eta \pi \pi) / \Gamma(K \bar{K} \pi)$					Γ_2 / Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027R3

NODE=M027R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.09 ± 0.48		27 AMSLER	04B	CBAR	0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \eta$
-------------	--	-----------	-----	------	---

<0.5	90	EDWARDS	83B	CBAL	$J/\psi \rightarrow \eta \pi \pi \gamma$
------	----	---------	-----	------	--

<1.1	90	SCHARRE	80	MRK2	$J/\psi \rightarrow \eta \pi \pi \gamma$
------	----	---------	----	------	--

<1.5	95	FOSTER	68B	HBC	0.0 $\bar{p}p$
------	----	--------	-----	-----	----------------

$\Gamma(\rho^0\gamma)/\Gamma(\eta\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_2
0.111±0.064	AMSLER	04B	CBAR 0 $\bar{p}p$	

NODE=M027R12
NODE=M027R12

 $\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}\pi)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
••• We do not use the following data for averages, fits, limits, etc. •••					
~ 0.15		28 BERTIN	95	OBLX 0 $\bar{p}p \rightarrow K\bar{K}\pi\pi$	
~ 0.8	500	28 DUCH	89	ASTE $\bar{p}p \rightarrow \pi^+\pi^-K^\pm\pi^\mp K^0$	
~ 0.75		28 REEVES	86	SPEC 6.6 $p\bar{p} \rightarrow KK\pi X$	

NODE=M027R4
NODE=M027R4

 $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
••• We do not use the following data for averages, fits, limits, etc. •••					
0.29±0.10		ABELE	98E	CBAR 0 $p\bar{p} \rightarrow \eta\pi^0\pi^0\pi^0$	
0.19±0.04	2200	29 ALDE	97B	GAM4 100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$	
0.56±0.04±0.03		29 AMSLER	95F	CBAR 0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$	

NODE=M027R2
NODE=M027R2

 $\Gamma(a_0(980)\pi)/\Gamma(\eta(\pi\pi)s\text{-wave})$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_4
••• We do not use the following data for averages, fits, limits, etc. •••					
0.91±0.12		ANISOVICH	01	SPEC 0.0 $\bar{p}p \rightarrow \eta\pi^+\pi^-\pi^+\pi^-$	
0.15±0.04	9082	30 MANAK	00A	MPS 18 $\pi^-p \rightarrow \eta\pi^+\pi^-n$	
0.70±0.12±0.20		31 BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	

NODE=M027R9
NODE=M027R9

 $\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_1
0.0152±0.0038	32 COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	

NODE=M027R7
NODE=M027R7

 $\Gamma(\eta(\pi\pi)s\text{-wave})/\Gamma(\eta\pi\pi)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_2
••• We do not use the following data for averages, fits, limits, etc. •••					
0.81±0.04	2200	ALDE	97B	GAM4 100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$	

NODE=M027R8
NODE=M027R8

 $\Gamma(f_0(980)\eta)/\Gamma(\eta\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
••• We do not use the following data for averages, fits, limits, etc. •••				
0.32±0.07	33 ANISOVICH	00	SPEC 0.9–1.2 $\bar{p}p \rightarrow \eta 3\pi^0$	

NODE=M027R10
NODE=M027R10

 $\Gamma(\rho\rho)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
<0.58	99.85	27,34 AMSLER	04B	CBAR 0 $\bar{p}p$	

NODE=M027R13
NODE=M027R13

 $\Gamma(K^*(892)K)/\Gamma(a_0(980)\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ_3
••• We do not use the following data for averages, fits, limits, etc. •••				
0.084±0.024	30 ADAMS	01B	B852 18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$	

NODE=M027R11
NODE=M027R11

 $\Gamma(\phi\gamma)/\Gamma(\rho^0\gamma)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ_9
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.77	95	35 BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$	

NODE=M027R14
NODE=M027R14

27 Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$.

28 Assuming that the $a_0(980)$ decays only into $K\bar{K}\pi$.

29 Assuming that the $a_0(980)$ decays only into $\eta\pi$.

30 Statistical error only.

31 Assuming that the $a_0(980)$ decays only into $\eta\pi$.

32 Using $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi) = 4.2 \times 10^{-3}$ and $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma\gamma\rho^0) = 6.4 \times 10^{-5}$ and assuming that the $\gamma\rho^0$ signal does not come from the $f_1(1420)$.

33 Using preliminary Crystal Barrel data.

34 Assuming that the $\eta(1405)$ decays are saturated by the $\pi\pi\eta$, $K\bar{K}\pi$ and $\rho\rho$ modes.

35 Calculated by us from $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \phi\gamma\gamma) < 0.82 \times 10^{-4}$ and $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \rho^0\gamma\gamma) = (1.07 \pm 0.17 \pm 0.11) \times 10^{-4}$.

NODE=M027R3;LINKAGE=AM
NODE=M027R4;LINKAGE=C
NODE=M027R2;LINKAGE=A
NODE=M027R;LINKAGE=K3
NODE=M027R9;LINKAGE=BK
NODE=M027R7;LINKAGE=D

NODE=M027R10;LINKAGE=D
NODE=M027R13;LINKAGE=AM
NODE=M027R14;LINKAGE=BA

$\eta(1405)$ REFERENCES

Author	Year	Pub	Collab	REFID
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i> (BES III Collab.)	REFID=55386
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i> (BES III Collab.)	REFID=54270
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i> (BES III Collab.)	REFID=53931
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i> (BES Collab.)	REFID=52143
AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i> (CLEO Collab.)	REFID=50764
AMSLER	04B	EPJ C33 23	C. Amsler <i>et al.</i> (Crystal Barrel Collab.)	REFID=51079
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i> (BES Collab.)	REFID=50167
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i> (OBELIX Collab.)	REFID=48848
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i> (L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i> (BNL E852 Collab.)	REFID=49649
ANISOVICH	01	NP A690 567	A.V. Anisovich <i>et al.</i>	REFID=48308
ANISOVICH	00	PL B472 168	A.V. Anisovich <i>et al.</i>	REFID=47429
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i> (BNL E852 Collab.)	REFID=47989
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i> (BES Collab.)	REFID=46606
CICALO	99	PL B462 453	C. Cicalo <i>et al.</i> (OBELIX Collab.)	REFID=47394
ABELE	98E	NP B514 45	A. Abele <i>et al.</i> (Crystal Barrel Collab.)	REFID=46314
BAI	98C	PL B440 217	J.Z. Bai <i>et al.</i> (BES Collab.)	REFID=46337
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i> (GAMS Collab.)	REFID=45396
Translated from YAF 60 458.				
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i> (OBELIX Collab.)	REFID=45417
AMSLER	95F	PL B358 389	C. Amsler <i>et al.</i> (Crystal Barrel Collab.)	REFID=44613
BERTIN	95	PL B361 187	A. Bertin <i>et al.</i> (OBELIX Collab.)	REFID=44614
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i> (LOQM, PNPI, WASH)	REFID=44438
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme (DM2 Collab.)	REFID=41584
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i> (Mark III Collab.)	REFID=42176
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i> (SUGI, NAGO, KEK, KYOT+)	REFID=41748
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i> (DM2 Collab.)	REFID=41352
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i> (Mark III Collab.)	REFID=41578
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i> (Mark III Collab.)	REFID=41350
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i> (DM2 Collab.)	REFID=40575
DUCH	89	ZPHY C45 223	K.D. Duch <i>et al.</i> (ASTERIX Collab.) JP	REFID=41016
RATH	89	PR D40 693	M.G. Rath <i>et al.</i> (NDAM, BRAN, BNL, CUNY+)	REFID=40924
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i> (BNL, FSU, IND, MASD) JP	REFID=40568
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i> (KEK, KYOT, NIRS, SAGA+) IJP	REFID=20891
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i> (FLOR, BNL, IND+) JP	REFID=20936
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i> (BNL, FLOR, IND+) JP	REFID=20934
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i> (TASSO Collab.)	REFID=20305
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i> (CIT, HARV, PRIN+)	REFID=21318
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i> (CIT, HARV, PRIN+)	REFID=21314
Also		PRL 50 219	C. Edwards <i>et al.</i> (CIT, HARV, PRIN+)	REFID=21315
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i> (SLAC, LBL)	REFID=21329
FOSTER	68B	NP B8 174	M. Foster <i>et al.</i> (CERN, CDEF)	REFID=21179
BAILLON	67	NC 50A 393	P.H. Baillon <i>et al.</i> (CERN, CDEF, IRAD)	REFID=20407

NODE=M006

 $f_1(1420)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the minireview under $\eta(1405)$.

NODE=M006

 $f_1(1420)$ MASS

NODE=M006M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1426.4 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.1.		
1434 ± 5 ± 5	133	¹ ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
1426 ± 6	711	ABDALLAH	03H DLPH	91.2 $e^+e^- \rightarrow K_S^0K^\pm\pi^\mp + X$
1420 ± 14	3651	NICHITIU	02 OBLX	
1428 ± 4 ± 2	20k	ADAMS	01B B852	18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$
1426 ± 1		BARBERIS	97C OMEG	450 $pp \rightarrow p\rho K_S^0K^\pm\pi^\mp$
1425 ± 8		BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
1435 ± 9		PROKOSHKIN	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$
1430 ± 4		² ARMSTRONG	92E OMEG	85,300 $\pi^+p, pp \rightarrow \pi^+p, p\rho(K\bar{K}\pi)$
1462 ± 20		³ AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
1443 + 7 + 3 - 6 - 2	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0K^\pm\pi^\mp$
1425 ± 10	17	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
1442 ± 5 + 10 - 17	111	BECKER	87 MRK3	$e^+e^- \rightarrow \omega K\bar{K}\pi$
1423 ± 4		GIDAL	87B MRK2	$e^+e^- \rightarrow e^+e^-K\bar{K}\pi$
1417 ± 13	13	AIHARA	86C TPC	$e^+e^- \rightarrow e^+e^-K\bar{K}\pi$
1422 ± 3		CHAUVAT	84 SPEC	ISR 31.5 pp
1440 ± 10		⁴ BROMBERG	80 SPEC	100 $\pi^-p \rightarrow K\bar{K}\pi X$
1426 ± 6	221	DIONISI	80 HBC	4 $\pi^-p \rightarrow K\bar{K}\pi n$
1420 ± 20		DAHL	67 HBC	1.6-4.2 π^-p

NODE=M006M2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1430.8 ± 0.9	⁵ SOSA	99	SPEC	$pp \rightarrow P_{\text{slow}} (K_S^0 K^+ \pi^-) P_{\text{fast}}$
1433.4 ± 0.8	⁵ SOSA	99	SPEC	$pp \rightarrow P_{\text{slow}} (K_S^0 K^- \pi^+) P_{\text{fast}}$
1429 ± 3	389	ARMSTRONG	89	OMEG 300 $pp \rightarrow K \bar{K} \pi pp$
1425 ± 2	1520	ARMSTRONG	84	OMEG 85 $\pi^+ p, pp \rightarrow (\pi^+, p)(K \bar{K} \pi)p$
~ 1420		BITYUKOV	84	SPEC 32 $K^- p \rightarrow K^+ K^- \pi^0 \gamma$

OCCUR=2

¹ From a fit with a width fixed at 55 MeV.

² This result supersedes ARMSTRONG 84, ARMSTRONG 89.

³ From fit to the $K^*(892)K 1^{++}$ partial wave.

⁴ Mass error increased to account for $a_0(980)$ mass cut uncertainties.

⁵ No systematic error given.

NODE=M006M2;LINKAGE=CH
 NODE=M006M2;LINKAGE=C
 NODE=M006M2;LINKAGE=B
 NODE=M006M2;LINKAGE=A
 NODE=M006M2;LINKAGE=N1

$f_1(1420)$ WIDTH

NODE=M006W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
54.9 ± 2.6 OUR AVERAGE				
51 ± 14	711	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
61 ± 8	3651	NICHITIU	02 OBLX	
$38 \pm 9 \pm 6$	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
58 ± 4		BARBERIS	97C OMEG	450 $pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
45 ± 10		BERTIN	97 OBLX	0.0 $\bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
90 ± 25		PROKOSHKIN	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
58 ± 10		⁶ ARMSTRONG	92E OMEG	85,300 $\pi^+ p, pp \rightarrow \pi^+ p, pp(K \bar{K} \pi)$
129 ± 41		⁷ AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$68 \begin{smallmatrix} +29 \\ -18 \end{smallmatrix} \begin{smallmatrix} +8 \\ -9 \end{smallmatrix}$	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
42 ± 22	17	BEHREND	89 CELL	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
$40 \begin{smallmatrix} +17 \\ -13 \end{smallmatrix} \pm 5$	111	BECKER	87 MRK3	$e^+ e^- \rightarrow \omega K \bar{K} \pi$
$35 \begin{smallmatrix} +47 \\ -20 \end{smallmatrix}$	13	AIHARA	86C TPC	$e^+ e^- \rightarrow e^+ e^- K \bar{K} \pi$
47 ± 10		CHAUVAT	84 SPEC	ISR 31.5 pp
62 ± 14		BROMBERG	80 SPEC	100 $\pi^- p \rightarrow K \bar{K} \pi X$
40 ± 15	221	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
60 ± 20		DAHL	67 HBC	1.6-4.2 $\pi^- p$

NODE=M006W

• • • We do not use the following data for averages, fits, limits, etc. • • •

68.7 ± 2.9	⁸ SOSA	99	SPEC	$pp \rightarrow P_{\text{slow}} (K_S^0 K^+ \pi^-) P_{\text{fast}}$
58.8 ± 3.3	⁸ SOSA	99	SPEC	$pp \rightarrow P_{\text{slow}} (K_S^0 K^- \pi^+) P_{\text{fast}}$
58 ± 8	389	ARMSTRONG	89	OMEG 300 $pp \rightarrow K \bar{K} \pi pp$
62 ± 5	1520	ARMSTRONG	84	OMEG 85 $\pi^+ p, pp \rightarrow (\pi^+, p)(K \bar{K} \pi)p$
~ 50		BITYUKOV	84	SPEC 32 $K^- p \rightarrow K^+ K^- \pi^0 \gamma$

OCCUR=2

⁶ This result supersedes ARMSTRONG 84, ARMSTRONG 89.

⁷ From fit to the $K^*(892)K 1^{++}$ partial wave.

⁸ No systematic error given.

NODE=M006W;LINKAGE=C
 NODE=M006W;LINKAGE=B
 NODE=M006W;LINKAGE=N1

$f_1(1420)$ DECAY MODES

NODE=M006215;NODE=M006

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K} \pi$	dominant
Γ_2 $K \bar{K}^*(892) + \text{c.c.}$	dominant
Γ_3 $\eta \pi \pi$	possibly seen
Γ_4 $a_0(980) \pi$	

DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=4

Γ_5 $\pi\pi\rho$
 Γ_6 4π
 Γ_7 $\rho^0\gamma$
 Γ_8 $\phi\gamma$

seen

DESIG=3
DESIG=6
DESIG=8
DESIG=9;OUR EST;→ UNCHECKED ←

 $f_1(1420) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M006220

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}}$

NODE=M006G2
NODE=M006G2

VALUE (keV)	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
1.9±0.4 OUR AVERAGE					
3.2±0.6±0.7		133	9,10 ACHARD	07 L3	183–209 $e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
3.0±0.9±0.7			11,12 BEHREND	89 CELL	$e^+e^- \rightarrow e^+e^- K_S^0 K\pi$
2.3 ^{+1.0} _{-0.9} ±0.8			HILL	89 JADE	$e^+e^- \rightarrow e^+e^- K^\pm K_S^0 \pi^\mp$
1.3±0.5±0.3			AIHARA	88B TPC	$e^+e^- \rightarrow e^+e^- K^\pm K_S^0 \pi^\mp$
1.6±0.7±0.3			11,13 GIDAL	87B MRK2	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
••• We do not use the following data for averages, fits, limits, etc. •••					
<8.0	95		JENNI	83 MRK2	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$

⁹From a fit with a width fixed at 55 MeV.¹⁰The form factor parameter from the fit is 926 ± 78 MeV.¹¹Assume a ρ -pole form factor.¹²A ϕ -pole form factor gives considerably smaller widths.¹³Published value divided by 2.

NODE=M006G2;LINKAGE=CH
NODE=M006G2;LINKAGE=CR
NODE=M006G2;LINKAGE=A
NODE=M006G2;LINKAGE=D
NODE=M006G2;LINKAGE=B

 $f_1(1420)$ BRANCHING RATIOS

NODE=M006225

 $\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(K\bar{K}\pi)$ **Γ_2/Γ_1**

VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
0.76±0.06	BROMBERG 80	SPEC	100 $\pi^- p \rightarrow K\bar{K}\pi X$
0.86±0.12	DIONISI 80	HBC	4 $\pi^- p \rightarrow K\bar{K}\pi n$

NODE=M006R1
NODE=M006R1

 $\Gamma(\pi\pi\rho)/\Gamma(K\bar{K}\pi)$ **Γ_5/Γ_1**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.3	95	CORDEN 78	OMEG	12–15 $\pi^- p$
<2.0		DAHL 67	HBC	1.6–4.2 $\pi^- p$

NODE=M006R2
NODE=M006R2

 $\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$ **Γ_3/Γ_1**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	95	ARMSTRONG 91B	OMEG	300 $p\bar{p} \rightarrow p\rho\eta\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••				
1.35±0.75		KOPKE 89	MRK3	$J/\psi \rightarrow \omega\eta\pi\pi(K\bar{K}\pi)$
<0.6	90	GIDAL 87	MRK2	$e^+e^- \rightarrow e^+e^- \eta\pi^+\pi^-$
<0.5	95	CORDEN 78	OMEG	12–15 $\pi^- p$
1.5 ±0.8		DEFOIX 72	HBC	0.7 $\bar{p}p$

NODE=M006R3
NODE=M006R3

 $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$ **Γ_4/Γ_3**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.1	90	PROKOSHKIN 97B	GAM4	100 $\pi^- p \rightarrow \eta\pi^0\pi^0 n$
••• We do not use the following data for averages, fits, limits, etc. •••				
not seen in either mode		ANDO 86	SPEC	8 $\pi^- p$
not seen in either mode		CORDEN 78	OMEG	12–15 $\pi^- p$
0.4±0.2		DEFOIX 72	HBC	0.7 $\bar{p}p \rightarrow 7\pi$

NODE=M006R4
NODE=M006R4

 $\Gamma(4\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$ **Γ_6/Γ_2**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.90	95	DIONISI 80	HBC	4 $\pi^- p$

NODE=M006R5
NODE=M006R5

$$\frac{\Gamma(K\bar{K}\pi)}{[\Gamma(K\bar{K}^*(892)+c.c.)+\Gamma(a_0(980)\pi)]} \quad \Gamma_1/(\Gamma_2+\Gamma_4)$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.65±0.27	¹⁴ DIONISI	80	HBC 4 π ⁻ p
-----------	-----------------------	----	------------------------

¹⁴ Calculated using $\Gamma(K\bar{K})/\Gamma(\eta\pi) = 0.24 \pm 0.07$ for $a_0(980)$ fractions.

NODE=M006R6
NODE=M006R6

NODE=M006R6;LINKAGE=C

$$\frac{\Gamma(a_0(980)\pi)}{\Gamma(K\bar{K}^*(892)+c.c.)} \quad \Gamma_4/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.04±0.01±0.01 BARBERIS 98C OMEG 450 pp →
p_f f₁(1420) p_s

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	68	ARMSTRONG 84	OMEG 85 π ⁺ p
-------	----	--------------	--------------------------

NODE=M006R7
NODE=M006R7

$$\frac{\Gamma(4\pi)}{\Gamma(K\bar{K}\pi)} \quad \Gamma_6/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.62 ARMSTRONG 89G OMEG 85 π p → 4π X

NODE=M006R8
NODE=M006R8

$$\frac{\Gamma(\rho^0\gamma)}{\Gamma_{\text{total}}} \quad \Gamma_7/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.08 95 ¹⁵ ARMSTRONG 92C SPEC 300 pp → ppπ⁺π⁻γ

¹⁵ Using the data on the $\bar{K}K\pi$ mode from ARMSTRONG 89.

NODE=M006R9
NODE=M006R9

NODE=M006R9;LINKAGE=A

$$\frac{\Gamma(\rho^0\gamma)}{\Gamma(K\bar{K}\pi)} \quad \Gamma_7/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.02 95 BARBERIS 98C OMEG 450 pp →
p_f f₁(1420) p_s

NODE=M006R10
NODE=M006R10

$$\frac{\Gamma(\phi\gamma)}{\Gamma(K\bar{K}\pi)} \quad \Gamma_8/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.003±0.001±0.001 BARBERIS 98C OMEG 450 pp →
p_f f₁(1420) p_s

NODE=M006R11
NODE=M006R11

f₁(1420) REFERENCES

ACHARD 07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=51698
ABDALLAH 03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49548
NICHITIU 02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
ADAMS 01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
SOSA 99	PRL 83 913	M. Sosa <i>et al.</i>		REFID=46937
BARBERIS 98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
BARBERIS 97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
BERTIN 97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
PROKOSHKIN 97B	SPD 42 298	Yu.D. Prokoshkin, S.A. Sadovsky		REFID=45549
	Translated from DANS 354 751.			
ARMSTRONG 92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=42097
ARMSTRONG 92E	ZPHY C56 29	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JPC	REFID=43173
AUGUSTIN 92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
ARMSTRONG 91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BAI 90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
ARMSTRONG 89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+) JPC	REFID=40729
ARMSTRONG 89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)	REFID=40930
BEHREND 89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40732
HILL 89	ZPHY C42 355	P. Hill <i>et al.</i>	(JADE Collab.) JP	REFID=40741
KOPKE 89	PRPL 174 67	L. Kopke <i>et al.</i>	(CERN)	REFID=41863
AIHARA 88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2γ Collab.)	REFID=40572
BECKER 87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.) JP	REFID=40015
GIDAL 87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
GIDAL 87B	PRL 59 2016	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40224
AIHARA 86C	PRL 57 2500	H. Aihara <i>et al.</i>	(TPC-2γ Collab.) JP	REFID=21326
ANDO 86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)	REFID=20891
ARMSTRONG 84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP	REFID=20929
BITYUKOV 84	SJNP 39 735	S. Bitjukov <i>et al.</i>	(SERP)	REFID=45856
	Translated from YAF 39 1165.			
CHAUVAT 84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)	REFID=20932
JENNI 83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)	REFID=20304
BROMBERG 80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)	REFID=20922
DIONISI 80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+) IJP	REFID=20924
CORDEN 78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20452
DEFOIX 72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
DAHL 67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP	REFID=20321
Also	PRL 14 1074	D.H. Miller <i>et al.</i>	(LRL, UCB)	REFID=21291

NODE=M006

$\omega(1420)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M125

 $\omega(1420)$ MASS

NODE=M125M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M125M

(1400–1450) OUR ESTIMATE

→ UNCHECKED ←

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1470 ± 50	13.1k	¹ AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1382 ± 23 ± 70		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1350 ± 20 ± 20		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1400 ± 50 ± 130	1.2M	² ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1450 ± 10		³ HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1373 ± 70	177	⁴ AKHMETSHIN	00D CMD2	1.2–1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
1370 ± 25	5095	ANISOVICH	00H SPEC	0.0 $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
1400 ⁺¹⁰⁰ ₋₂₀₀		⁵ ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
~ 1400		⁶ ACHASOV	98H RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
~ 1460		⁷ ACHASOV	98H RVUE	$e^+e^- \rightarrow K^+K^-$
1440 ± 70		⁸ CLEGG	94 RVUE	
1419 ± 31	315	⁹ ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi$

OCCUR=2

OCCUR=3

¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.³ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.⁴ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.⁵ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.⁶ Using the data from ANTONELLI 92.⁷ Using the data from IVANOV 81 and BISELLO 88B.⁸ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.⁹ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

NODE=M125M;LINKAGE=E

NODE=M125M;LINKAGE=VH

NODE=M125M;LINKAGE=AB

NODE=M125M;LINKAGE=KL

NODE=M125M;LINKAGE=L1

NODE=M125M;LINKAGE=L2

NODE=M125M;LINKAGE=L3

NODE=M125M;LINKAGE=AD

NODE=M125M;LINKAGE=B

 $\omega(1420)$ WIDTH

NODE=M125W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M125W

(180–250) OUR ESTIMATE

→ UNCHECKED ←

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

880 ± 170	13.1k	¹⁰ AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
130 ± 50 ± 100		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
450 ± 70 ± 70		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
870 ⁺⁵⁰⁰ ₋₃₀₀ ± 450	1.2M	¹¹ ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
199 ± 15		¹² HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
188 ± 45	177	¹³ AKHMETSHIN	00D CMD2	1.2–1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
360 ⁺¹⁰⁰ ₋₆₀	5095	ANISOVICH	00H SPEC	0.0 $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
240 ± 70		¹⁴ CLEGG	94 RVUE	
174 ± 59	315	¹⁵ ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi$

¹⁰ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.¹¹ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.¹² Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.¹³ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.¹⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.¹⁵ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

NODE=M125W;LINKAGE=E

NODE=M125W;LINKAGE=VH

NODE=M125W;LINKAGE=AB

NODE=M125W;LINKAGE=KL

NODE=M125W;LINKAGE=AD

NODE=M125W;LINKAGE=B

$\omega(1420)$ DECAY MODES

NODE=M125215;NODE=M125

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	dominant
Γ_2 $\omega\pi\pi$	seen
Γ_3 $b_1(1235)\pi$	seen
Γ_4 e^+e^-	seen
Γ_5 $\pi^0\gamma$	

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=6

 $\omega(1420)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M125230

 $\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_4/\Gamma$ NODE=M125G3
NODE=M125G3

VALUE (units 10^{-6})	EVTs	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.73 ± 0.08	13.1k	¹⁶ AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.82 ± 0.05 ± 0.06		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
0.65 ± 0.13 ± 0.21	1.2M	^{17,18} ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.625 ± 0.160		^{19,20} CLEGG	94 RVUE	
0.466 ± 0.178		^{21,22} ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi$

¹⁶ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.¹⁷ Calculated by us from the cross section at the peak.¹⁸ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.¹⁹ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.²⁰ From the partial and leptonic width given by the authors.²¹ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.²² From the product of the leptonic width and partial branching ratio given by the authors.NODE=M125G3;LINKAGE=A
NODE=M125G;LINKAGE=AW
NODE=M125G;LINKAGE=VH

NODE=M125G;LINKAGE=AD

NODE=M125G;LINKAGE=SE
NODE=M125G;LINKAGE=A

NODE=M125G;LINKAGE=ES

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_4/\Gamma$ NODE=M125G4
NODE=M125G4

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
19.7 ± 5.7	AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1.9 ± 1.9	²³ AKHMETSHIN	00D CMD2	1.2–2.4 $e^+e^- \rightarrow \omega\pi^+\pi^-$

²³ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

NODE=M125G;LINKAGE=KL

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma \times \Gamma_4/\Gamma$ NODE=M125G5
NODE=M125G5

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2.03 ^{+0.70} _{-0.75}	²⁴ AKHMETSHIN	05 CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$

²⁴ Using 1420 MeV and 220 MeV for the $\omega(1420)$ mass and width.

NODE=M125G5;LINKAGE=AK

 $\omega(1420)$ BRANCHING RATIOS

NODE=M125225

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M125R2
NODE=M125R2

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.301 ± 0.029	²⁵ HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
possibly seen	AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$

 $\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$ Γ_2/Γ_3 NODE=M125R1
NODE=M125R1

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.60 ± 0.16	5095	ANISOVICH	00H SPEC	0.0 $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ NODE=M125R3
NODE=M125R3

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.699 ± 0.029	²⁵ HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-7})	EVTs	DOCUMENT ID	TECN	COMMENT
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
~ 6.6	1.2M	^{26,27} ACHASOV	03D	RVUE $0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
23 \pm 1		²⁵ HENNER	02	RVUE $1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
²⁵ Assuming that the $\omega(1420)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.				
²⁶ Calculated by us from the cross section at the peak.				
²⁷ Assuming that the $\omega(1420)$ decays into $\rho\pi$ only.				

NODE=M125R4
NODE=M125R4NODE=M125R;LINKAGE=AC
NODE=M125R;LINKAGE=AW
NODE=M125R;LINKAGE=GS $\omega(1420)$ REFERENCES

AULCHENKO	15A	JETP 121 27 Translated from ZETF 148 34.	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>	
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164 Translated from ZETFP 46 132.	L.M. Barkov <i>et al.</i>	(NOVO)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)

NODE=M125

REFID=56843

REFID=52049

REFID=50330

REFID=50184

REFID=49577

REFID=48815

REFID=49177

REFID=48311

REFID=47935

REFID=47948

REFID=47391

REFID=46323

REFID=44081

REFID=43168

REFID=41369

REFID=40581

REFID=40280

NODE=M066

 $f_2(1430)$

$$J^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry lists nearby peaks observed in the D wave of the $K\bar{K}$ and $\pi^+\pi^-$ systems. Needs confirmation.

NODE=M066

 $f_2(1430)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1430 OUR ESTIMATE			
●●● We do not use the following data for averages, fits, limits, etc. ●●●			
1453 ± 4	¹ VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1421 ± 5	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
1480 ± 50	AKESSON	86	SPEC $pp \rightarrow pp \pi^+ \pi^-$
1436^{+26}_{-16}	DAUM	84	CNTR $17-18 \pi^- p \rightarrow K^+ K^- n$
1412 ± 3	DAUM	84	CNTR $63 \pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
1439^{+5}_{-6}	² BEUSCH	67	OSPK $5,7,12 \pi^- p \rightarrow K_S^0 K_S^0 n$
¹ $J^{PC} = 0^{++}$ or 2^{++} .			
² Not seen by WETZEL 76.			

NODE=M066M1

NODE=M066M1

→ UNCHECKED ←

OCCUR=2

NODE=M066M;LINKAGE=AC
NODE=M066M;LINKAGE=C $f_2(1430)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
●●● We do not use the following data for averages, fits, limits, etc. ●●●			
13 ± 5	³ VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
30 ± 9	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
150 ± 50	AKESSON	86	SPEC $pp \rightarrow pp \pi^+ \pi^-$
81^{+56}_{-29}	DAUM	84	CNTR $17-18 \pi^- p \rightarrow K^+ K^- n$
14 ± 6	DAUM	84	CNTR $63 \pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
43^{+17}_{-18}	⁴ BEUSCH	67	OSPK $5,7,12 \pi^- p \rightarrow K_S^0 K_S^0 n$
³ $J^{PC} = 0^{++}$ or 2^{++} .			
⁴ Not seen by WETZEL 76.			

NODE=M066W1

NODE=M066W1

OCCUR=2

NODE=M066W;LINKAGE=AC
NODE=M066W;LINKAGE=C

$f_2(1430)$ DECAY MODES

NODE=M066215;NODE=M066

Mode	
Γ_1	$K\bar{K}$
Γ_2	$\pi\pi$

DESIG=1

DESIG=2

 $f_2(1430)$ REFERENCES

NODE=M066

VLADIMIRSK... 01	PAN 64 1895 Translated from YAF 64 1979.	V.V. Vladimirov <i>et al.</i>	
AUGUSTIN 87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
AKESSON 86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
DAUM 84	ZPHY C23 339	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
WETZEL 76	NP B115 208	W. Wetzels <i>et al.</i>	(ETH, CERN, LOIC)
BEUSCH 67	PL 25B 357	W. Beusch <i>et al.</i>	(ETH, CERN)

REFID=48571

REFID=40268

REFID=21123

REFID=21372

REFID=20362

REFID=20320

NODE=M149

 $a_0(1450)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

See minireview on scalar mesons under $f_0(500)$.

NODE=M149

 $a_0(1450)$ MASS

NODE=M149M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1474 ±19	OUR AVERAGE			
1480 ±30		ABELE 98	CBAR	0.0 $\bar{p}p \rightarrow K^0 K^\pm \pi^\mp$
1470 ±25		¹ AMSLER 95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0,$ $\pi^0 \eta \eta, \pi^0 \pi^0 \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1515 ±30		² ANISOVICH 09	RVUE	0.0 $\bar{p}p, \pi N$
1316.8 ⁺ _{-1.0} 0.7 ⁺ _{-4.6}		³ UEHARA 09A	BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
1432 ±13 ±25		⁴ BUGG 08A	RVUE	$\bar{p}p$
1477 ±10	80k	⁵ UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
1441 ⁺ ₋₁₅ +40	35280	² BAKER 03	SPEC	$\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
1303 ±16		⁶ BARGIOTTI 03	OBLX	$\bar{p}p$
1296 ±10		⁷ AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
1565 ±30		⁷ ANISOVICH 98B	RVUE	Compilation
1290 ±10		⁸ BERTIN 98B	OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp$
1450 ±40		AMSLER 94D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
1410 ±25		ETKIN 82C	MPS	23 $\pi^- p \rightarrow n 2K_S^0$
~ 1300		MARTIN 78	SPEC	10 $K^\pm p \rightarrow K_S^0 \pi p$
1255 ± 5		⁹ CASON 76		

NODE=M149M

¹ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.² From the pole position.³ May be a different state.⁴ Using data from AMSLER 94D, ABELE 98, and BAKER 03. Supersedes BUGG 94.⁵ Statistical error only.⁶ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.⁷ T-matrix pole.⁸ Not confirmed by BUGG 08A.⁹ Isospin 0 not excluded.

NODE=M149M;LINKAGE=AB

NODE=M149M;LINKAGE=PP

NODE=M149M;LINKAGE=UE

NODE=M149M;LINKAGE=BU

NODE=M149M;LINKAGE=ST

NODE=M149M;LINKAGE=BG

NODE=M149M;LINKAGE=AN

NODE=M149M;LINKAGE=BE

NODE=M149M;LINKAGE=CC

 $a_0(1450)$ WIDTH

NODE=M149W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
265 ±13	OUR AVERAGE			
265 ±15		ABELE 98	CBAR	0.0 $\bar{p}p \rightarrow K^0 K^\pm \pi^\mp$
265 ±30		¹⁰ AMSLER 95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0,$ $\pi^0 \eta \eta, \pi^0 \pi^0 \eta$

NODE=M149W

- • • We do not use the following data for averages, fits, limits, etc. • • •

230 ±36		11 ANISOVICH	09	RVUE	0.0 $\bar{p}p, \pi N$
65.0 ⁺ ₋ 2.1+99.1 5.4-32.6		12 UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
196 ±10 ±10		13 BUGG	08A	RVUE	$\bar{p}p$
267 ±11	80k	14 UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
110 ±14	35280	11 BAKER	03	SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$
92 ±16		15 BARGIOTTI	03	OBLX	$\bar{p}p$
81 ±21		16 AMSLER	02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta$
292 ±40		16 ANISOVICH	98B	RVUE	Compilation
80 ± 5		17 BERTIN	98B	OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp$
270 ±40		AMSLER	94D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\eta$
230 ±30		ETKIN	82C	MPS	23 $\pi^-p \rightarrow n2K_S^0$
~ 250		MARTIN	78	SPEC	10 $K^\pm p \rightarrow K_S^0 \pi p$
79 ±10		18 CASON	76		

¹⁰ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

¹¹ From the pole position.

¹² May be a different state.

¹³ Using data from AMSLER 94D, ABELE 98, and BAKER 03. Supersedes BUGG 94.

¹⁴ Statistical error only.

¹⁵ Coupled channel analysis of $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, and $K^\pm K_S^0 \pi^\mp$.

¹⁶ T-matrix pole.

¹⁷ Not confirmed by BUGG 08A.

¹⁸ Isospin 0 not excluded.

NODE=M149W;LINKAGE=AB
 NODE=M149W;LINKAGE=PP
 NODE=M149W;LINKAGE=UE
 NODE=M149W;LINKAGE=BU
 NODE=M149W;LINKAGE=ST
 NODE=M149W;LINKAGE=BG
 NODE=M149W;LINKAGE=AN
 NODE=M149W;LINKAGE=BE
 NODE=M149W;LINKAGE=CC

$a_0(1450)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\eta$	seen
Γ_2 $\pi\eta'(958)$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $\omega\pi\pi$	seen
Γ_5 $a_0(980)\pi\pi$	seen
Γ_6 $\gamma\gamma$	seen

NODE=M149215;NODE=M149

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=5
 DESIG=6

$a_0(1450)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_6/\Gamma$
VALUE (eV)				

NODE=M149225

NODE=M149G01
 NODE=M149G01

- • • We do not use the following data for averages, fits, limits, etc. • • •

432±6 ⁺ ₋ 1073 256	19 UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
---	-----------	-----	------	--------------------------------------

¹⁹ May be a different state.

NODE=M149G01;LINKAGE=UE

$a_0(1450)$ BRANCHING RATIOS

$\Gamma(\pi\eta'(958))/\Gamma(\pi\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
VALUE				

0.35±0.16 ²⁰ ABELE 98 CBAR 0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.43±0.19 ABELE 97C CBAR 0.0 $\bar{p}p \rightarrow \pi^0\pi^0\eta'$

²⁰ Using $\pi^0\eta$ from AMSLER 94D.

NODE=M149R1;LINKAGE=A

$\Gamma(K\bar{K})/\Gamma(\pi\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
VALUE				

0.88±0.23 ²¹ ABELE 98 CBAR 0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$

²¹ Using $\pi^0\eta$ from AMSLER 94D.

NODE=M149R2
 NODE=M149R2

NODE=M149R2;LINKAGE=A

$\Gamma(\omega\pi\pi)/\Gamma(\pi\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
VALUE				

- • • We do not use the following data for averages, fits, limits, etc. • • •

10.7±2.3 35280 ²² BAKER 03 SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

²² Using results on $\bar{p}p \rightarrow a_0(1450)^0\pi^0$, $a_0(1450) \rightarrow \eta\pi^0$ from ABELE 96C and assuming the $\omega\rho$ mechanism for the $\omega\pi\pi$ state.

NODE=M149R3
 NODE=M149R3

NODE=M149R;LINKAGE=PP

$\Gamma(a_0(980)\pi\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
seen	BUGG	08A	RVUE $\bar{p}p$	

NODE=M149R01
 NODE=M149R01

 $\Gamma(a_0(980)\pi\pi)/\Gamma(\pi\eta)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_1
-------	-------------	------	-----	---------	---------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

≤ 4.3 ANISOVICH 01 RVUE 0 $\bar{p}p \rightarrow \eta 2\pi^+ 2\pi^-$

NODE=M149R02
 NODE=M149R02

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
-------	-------------	------	---------	-------------------

seen 23 UEHARA 09A BELL $\gamma\gamma \rightarrow \pi^0\eta$

23 May be a different state.

NODE=M149R03
 NODE=M149R03

NODE=M149R03;LINKAGE=UE

 $a_0(1450)$ REFERENCES

ANISOVICH 09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
UEHARA 09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)
BUGG 08A	PR D78 074023	D.V. Bugg	(LOQM)
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
BAKER 03	PL B563 140	C.A. Baker <i>et al.</i>	
BARGIOTTI 03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH 01	NP A690 567	A.V. Anisovich <i>et al.</i>	
ABELE 98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH 98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
	Translated from UFN 168 481.		
BERTIN 98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE 97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE 96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) IGJPC
BUGG 94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
ETKIN 82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
MARTIN 78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)
CASON 76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL)

NODE=M149

REFID=52719
 REFID=53002
 REFID=52578
 REFID=51063
 REFID=49414
 REFID=49217
 REFID=48580
 REFID=48308
 REFID=45863
 REFID=46331

REFID=46351
 REFID=45531
 REFID=45076
 REFID=44377
 REFID=44440
 REFID=44441
 REFID=44093
 REFID=44078
 REFID=20391
 REFID=22446
 REFID=21064

NODE=M105

 $\rho(1450)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

See our mini-review under the $\rho(1700)$.

NODE=M105

 $\rho(1450)$ MASS

VALUE (MeV) DOCUMENT ID
1465 ± 25 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

NODE=M105205

NODE=M105M0
 → UNCHECKED ←

 $\eta\rho^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1497 ± 14	¹ AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1421 ± 15	² AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1470 ± 20	ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1446 ± 10	FUKUI 88	SPEC	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$

¹ Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$.

² Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

NODE=M105M1
 NODE=M105M1

NODE=M105M;LINKAGE=SW

NODE=M105M1;LINKAGE=KL

 $\omega\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1544 ± 22 ⁺¹¹ ₋₄₆	821	¹ MATVIENKO 15	BELL	$\bar{B}^0 \rightarrow D^{*0}\omega\pi^-$
1491 ± 19	7815	² ACHASOV 13	SND	$1.05-2.00 e^+e^- \rightarrow \pi^0\pi^0\gamma$
1582 ± 17 ± 25	2382	³ AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1349 ± 25 ⁺¹⁰ ₋₅	341	⁴ ALEXANDER 01B	CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
1523 ± 10		⁵ EDWARDS 00A	CLE2	$\tau^- \rightarrow \omega\pi^- \nu_\tau$
1463 ± 25		⁶ CLEGG 94	RVUE	
1250		⁷ ASTON 80C	OMEG	$20-70 \gamma p \rightarrow \omega\pi^0 p$
1290 ± 40		⁷ BARBER 80C	SPEC	$3-5 \gamma p \rightarrow \omega\pi^0 p$

NODE=M105M3
 NODE=M105M3

- ¹ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.
- ² From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.
- ³ Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the $\omega\pi^0$ and $\pi^+\pi^-$ mass dependence of the total width. $\rho(1700)$ mass and width fixed at 1700 MeV and 240 MeV, respectively.
- ⁴ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming the $\omega\pi^-$ mass dependence for the total width.
- ⁵ Mass-independent width parameterization. $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
- ⁶ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.
- ⁷ Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.

NODE=M105M3;LINKAGE=C
 NODE=M105M3;LINKAGE=AC
 NODE=M105M3;LINKAGE=HK
 NODE=M105M3;LINKAGE=3Z
 NODE=M105M;LINKAGE=E1
 NODE=M105M3;LINKAGE=B
 NODE=M105M3;LINKAGE=A
 NODE=M105M6
 NODE=M105M6

4 π MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1435 \pm 40	ABELE 01B	CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0 \pi^+$
1350 \pm 50	ACHASOV 97	RVUE	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1449 \pm 4	¹ ARMSTRONG 89E	OMEG	300 $pp \rightarrow pp2(\pi^+\pi^-)$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- ¹ Not clear whether this observation has $l=1$ or 0.

NODE=M105M6;LINKAGE=A

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1350 \pm 20 $^{+20}_{-30}$	63.5k	¹ ABRAMOWICZ12	ZEUS	$e p \rightarrow e \pi^+ \pi^- p$
1493 \pm 15		² LEES	12G BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
1446 \pm 7 \pm 28	5.4M	^{3,4} FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
1328 \pm 15		⁵ SCHAEEL	05C ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
1406 \pm 15	87k	^{3,6} ANDERSON	00A CLE2	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
\sim 1368		⁷ ABELE	99C CBAR	0.0 $\bar{p}d \rightarrow \pi^+ \pi^- \pi^- p$
1348 \pm 33		BERTIN	98 OBLX	0.05–0.405 $\bar{n}p \rightarrow 2\pi^+ \pi^-$
1411 \pm 14		⁸ ABELE	97 CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
1370 $^{+90}_{-70}$		ACHASOV	97 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
1359 \pm 40		⁶ BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
1282 \pm 37		BERTIN	97D OBLX	0.05 $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
1424 \pm 25		BISELLO	89 DM2	$e^+ e^- \rightarrow \pi^+ \pi^-$
1265.5 \pm 75.3		DUBNICKA	89 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
1292 \pm 17		⁹ KURDADZE	83 OLYA	0.64–1.4 $e^+ e^- \rightarrow \pi^+ \pi^-$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- ¹ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
- ² Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
- ³ From the GOUNARIS 68 parametrization of the pion form factor.
- ⁴ $|F_\pi(0)|^2$ fixed to 1.
- ⁵ From the combined fit of the τ^- data from ANDERSON 00A and SCHAEEL 05C and e^+e^- data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. $\rho(1700)$ mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.
- ⁶ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV, respectively.
- ⁷ $\rho(1700)$ mass and width fixed at 1780 MeV and 275 MeV respectively.
- ⁸ T-matrix pole.
- ⁹ Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

NODE=M105M5
 NODE=M105M5

NODE=M105M5;LINKAGE=AB
 NODE=M105M5;LINKAGE=LE

NODE=M105M;LINKAGE=1K
 NODE=M105M5;LINKAGE=FU
 NODE=M105M5;LINKAGE=SC

NODE=M105M5;LINKAGE=A
 NODE=M105M5;LINKAGE=C5
 NODE=M105M5;LINKAGE=QQ
 NODE=M105M5;LINKAGE=KD

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1422.8 \pm 6.5	27k	¹ ABELE	99D	CBAR \pm	0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$

- ¹ K-matrix pole. Isospin not determined, could be $\omega(1420)$.

NODE=M105M7
 NODE=M105M7

NODE=M105M7;LINKAGE=AN

$K\bar{K}^*(892) + \text{c.c.}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1505 \pm 19 \pm 7	AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K\bar{K}^*(892)\gamma$

- • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M105M8
 NODE=M105M8

$\rho(1450)$ WIDTH

VALUE (MeV)

DOCUMENT ID

400±60 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

NODE=M105210

NODE=M105W0

→ UNCHECKED ←

 $\eta\rho^0$ MODE

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

226±44		¹ AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
211±31		² AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
230±30		ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
60±15		FUKUI 88	SPEC	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$

¹ Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$.² Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

NODE=M105W1

NODE=M105W1

NODE=M105W;LINKAGE=SW

NODE=M105W1;LINKAGE=KL

 $\omega\pi$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$303^{+31}_{-52} +^{69}_{-7}$	821	¹ MATVIENKO 15	BELL	$\bar{B}^0 \rightarrow D^{*0}\omega\pi^-$
429±42±10	2382	² AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
547±86 ⁺⁴⁶ ₋₄₅	341	³ ALEXANDER 01B	CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
400±35		⁴ EDWARDS 00A	CLE2	$\tau^- \rightarrow \omega\pi^- \nu_\tau$
311±62		⁵ CLEGG 94	RVUE	
300		⁶ ASTON 80C	OMEG	20-70 $\gamma p \rightarrow \omega\pi^0 p$
320±100		⁶ BARBER 80C	SPEC	3-5 $\gamma p \rightarrow \omega\pi^0 p$

¹ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.² Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the $\omega\pi^0$ and $\pi^+\pi^-$ mass dependence of the total width. $\rho(1700)$ mass and width fixed at 1700 MeV and 240 MeV, respectively.³ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming the $\omega\pi^-$ mass dependence for the total width.⁴ Mass-independent width parameterization. $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.⁵ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.⁶ Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.

NODE=M105W3

NODE=M105W3

NODE=M105W3;LINKAGE=C

NODE=M105W3;LINKAGE=HK

NODE=M105W3;LINKAGE=3Z

NODE=M105W;LINKAGE=E1

NODE=M105W3;LINKAGE=B

NODE=M105W3;LINKAGE=A

4 π MODE

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

325±100		ABELE 01B	CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0 \pi^+$
---------	--	-----------	------	--

NODE=M105W66

NODE=M105W66

 $\pi\pi$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

460±30 ⁺⁴⁰ ₋₄₅	63.5k	¹ ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+\pi^- p$
427±31		² LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
434±16±60	5.4M	^{3,4} FUJIKAWA 08	BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
468±41		⁵ SCHAEEL 05C	ALEP	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
455±41	87k	^{3,6} ANDERSON 00A	CLE2	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
~ 374		⁷ ABELE 99C	CBAR	0.0 $\bar{p}d \rightarrow \pi^+\pi^-\pi^- p$
275±10		BERTIN 98	OBLX	0.05-0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
343±20		⁸ ABELE 97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
310±40		⁶ BERTIN 97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
236±36		BERTIN 97D	OBLX	0.05 $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
269±31		BISELLO 89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$
391±70		DUBNICKA 89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
218±46		⁹ KURDADZE 83	OLYA	0.64-1.4 $e^+e^- \rightarrow \pi^+\pi^-$

NODE=M105W5

NODE=M105W5

- ¹ Using the KUHN 90 parametrization of the pion form factor, neglecting ρ - ω interference.
² Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
³ From the GOUNARIS 68 parametrization of the pion form factor.
⁴ $|F_\pi(0)|^2$ fixed to 1.
⁵ From the combined fit of the τ^- data from ANDERSON 00A and SCHAELE 05C and e^+e^- data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. $\rho(1700)$ mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.
⁶ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV, respectively.
⁷ $\rho(1700)$ mass and width fixed at 1780 MeV and 275 MeV respectively.
⁸ T-matrix pole.
⁹ Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

NODE=M105W5;LINKAGE=AB
 NODE=M105W5;LINKAGE=LE

NODE=M105W5;LINKAGE=1K
 NODE=M105W5;LINKAGE=FU
 NODE=M105W5;LINKAGE=SC

NODE=M105W5;LINKAGE=A
 NODE=M105W5;LINKAGE=C5
 NODE=M105W5;LINKAGE=QQ
 NODE=M105W5;LINKAGE=KD

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

146.5 ± 10.5	27k	¹ ABELE	99D	CBAR ±	0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$
--------------	-----	--------------------	-----	--------	--

¹ K-matrix pole. Isospin not determined, could be $\omega(1420)$.

NODE=M105W7
 NODE=M105W7

NODE=M105W7;LINKAGE=AN

$K\bar{K}^*(892) + c.c.$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

418 ± 25 ± 4	AUBERT	08S	BABR 10.6 $e^+e^- \rightarrow K\bar{K}^*(892)\gamma$
--------------	--------	-----	--

NODE=M105W8
 NODE=M105W8

$\rho(1450)$ DECAY MODES

NODE=M105215;NODE=M105

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 4π	seen
Γ_3 $\omega\pi$	
Γ_4 $a_1(1260)\pi$	
Γ_5 $h_1(1170)\pi$	
Γ_6 $\pi(1300)\pi$	
Γ_7 $\rho\rho$	
Γ_8 $\rho(\pi\pi)$ S-wave	
Γ_9 e^+e^-	seen
Γ_{10} $\eta\rho$	seen
Γ_{11} $a_2(1320)\pi$	not seen
Γ_{12} $K\bar{K}$	not seen
Γ_{13} $K\bar{K}^*(892) + c.c.$	possibly seen
Γ_{14} $\eta\gamma$	seen
Γ_{15} $f_0(500)\gamma$	not seen
Γ_{16} $f_0(980)\gamma$	not seen
Γ_{17} $f_0(1370)\gamma$	not seen
Γ_{18} $f_2(1270)\gamma$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=6
 DESIG=10
 DESIG=11
 DESIG=12
 DESIG=13
 DESIG=14
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=3
 DESIG=8;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EVAL;→ UNCHECKED ←
 DESIG=15;OUR EST;→ UNCHECKED ←
 DESIG=9
 DESIG=16;OUR EST;→ UNCHECKED ←
 DESIG=17;OUR EST;→ UNCHECKED ←
 DESIG=18;OUR EST;→ UNCHECKED ←
 DESIG=19;OUR EST;→ UNCHECKED ←

$\rho(1450) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M105220

$\Gamma(\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_9/\Gamma$
--	-------------	------	---------	---------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

0.12	¹ DIEKMAN	88	RVUE $e^+e^- \rightarrow \pi^+\pi^-$	
0.027 ^{+0.015} _{-0.010}	² KURDADZE	83	OLYA $0.64-1.4 e^+e^- \rightarrow \pi^+\pi^-$	

NODE=M105G3
 NODE=M105G3

$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_{10}\Gamma_9/\Gamma$
--	-------------	------	---------	------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

74 ± 20	³ AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$	
91 ± 19	ANTONELLI	88	DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$	

NODE=M105G4
 NODE=M105G4

$$\Gamma(\eta\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_9/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16.4	⁴ AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
2.2±0.5±0.3	⁵ AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$

NODE=M105G6
NODE=M105G6

$$\Gamma(K\bar{K}^*(892)+c.c.) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{13}\Gamma_9/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

127±15±6	AUBERT	08s	BABR 10.6 $e^+e^- \rightarrow K\bar{K}^*(892)\gamma$
----------	--------	-----	--

¹ Using total width = 235 MeV.

² Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

³ Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

⁴ From 2γ decay mode of η using 1465 MeV and 310 MeV for the $\rho(1450)$ mass and width. Recalculated by us.

⁵ Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$. Recalculated by us using width of 226 MeV.

NODE=M105G8
NODE=M105G8

NODE=M105G3;LINKAGE=B
NODE=M105G3;LINKAGE=KD
NODE=M105G4;LINKAGE=KL

NODE=M105G6;LINKAGE=AK

NODE=M105G;LINKAGE=SW

$$\rho(1450) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+e^-)/\Gamma(\text{total})$$

NODE=M105230

$$\Gamma(\omega\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.3±0.4	7815	¹ ACHASOV	13	SND 1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
---------	------	----------------------	----	---

NODE=M105R05
NODE=M105R05

$$\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.3 ^{+1.1} _{-0.9} ±0.2	4.9k	² AULCHENKO	15	SND 1.22-2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
--	------	------------------------	----	---

NODE=M105R00
NODE=M105R00

$$\Gamma(f_0(500)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<4.0	90	ACHASOV	11	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
------	----	---------	----	---

NODE=M105R01
NODE=M105R01

$$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2.6	90	ACHASOV	11	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
------	----	---------	----	---

NODE=M105R02
NODE=M105R02

$$\Gamma(f_0(1370)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<3.5	90	ACHASOV	11	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
------	----	---------	----	---

NODE=M105R03
NODE=M105R03

$$\Gamma(f_2(1270)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.8	90	³ ACHASOV	11	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
------	----	----------------------	----	---

NODE=M105R04
NODE=M105R04

¹ From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

² From a fit to the $e^+e^- \rightarrow \eta\pi^+\pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta\rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

³ Using Breit-Wigner parametrization of the $\rho(1450)$ with mass and width of 1465 MeV and 400 MeV, respectively.

NODE=M105R05;LINKAGE=AC

NODE=M105R00;LINKAGE=A

NODE=M105R01;LINKAGE=AC

$\rho(1450)$ BRANCHING RATIOS

NODE=M105225

$$\Gamma(\pi\pi)/\Gamma(4\pi) \quad \Gamma_1/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.37±0.10	^{1,2} ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	----------------------	-----	--------------------------------------

NODE=M105R15
NODE=M105R15

$\Gamma(\omega\pi)/\Gamma_{\text{total}}$					Γ_3/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
seen	821	³ MATVIENKO	15	BELL	$\bar{B}^0 \rightarrow D^{*0}\omega\pi^-$	
seen	1.6k	ACHASOV	12	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
~ 0.21		CLEGG	94	RVUE		
$\Gamma(\pi\pi)/\Gamma(\omega\pi)$					Γ_1/Γ_3	
VALUE		DOCUMENT ID	TECN			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
~ 0.32		CLEGG	94	RVUE		
$\Gamma(\omega\pi)/\Gamma(4\pi)$					Γ_3/Γ_2	
VALUE		DOCUMENT ID	TECN			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 0.14		CLEGG	88	RVUE		
$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$					Γ_4/Γ_2	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.27 ± 0.08		¹ ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$	
$\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$					Γ_5/Γ_2	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.08 ± 0.04		¹ ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$	
$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$					Γ_6/Γ_2	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.37 ± 0.13		¹ ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$	
$\Gamma(\rho\rho)/\Gamma(4\pi)$					Γ_7/Γ_2	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.11 ± 0.05		¹ ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$	
$\Gamma(\rho(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$					Γ_8/Γ_2	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.17 ± 0.09		¹ ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$	
$\Gamma(\eta\rho)/\Gamma_{\text{total}}$					Γ_{10}/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
seen	35	⁴ ACHASOV	14	SND	1.15–2.00 $e^+e^- \rightarrow \eta\gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 0.04		DONNACHIE	87B	RVUE		
$\Gamma(\eta\rho)/\Gamma(\omega\pi)$					Γ_{10}/Γ_3	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.081 ± 0.020		^{5,6} AULCHENKO	15	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$	
~ 0.24		⁷ DONNACHIE	91	RVUE		
> 2		FUKUI	91	SPEC	8.95 $\pi^-p \rightarrow \omega\pi^0n$	
$\Gamma(\pi\pi)/\Gamma(\eta\rho)$					Γ_1/Γ_{10}	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
1.3 ± 0.4		⁵ AULCHENKO	15	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$	
$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$					Γ_{11}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
not seen		AMELIN	00	VES	37 $\pi^-p \rightarrow \eta\pi^+\pi^-n$	

$\Gamma(K\bar{K})/\Gamma(\omega\pi)$

VALUE	DOCUMENT ID	TECN
-------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •
 <0.08 ⁷ DONNACHIE 91 RVUE

 Γ_{12}/Γ_3

NODE=M105R8
 NODE=M105R8

 $\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •
 possibly seen COAN 04 CLEO $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$

 Γ_{13}/Γ

NODE=M105R16
 NODE=M105R16

 $\Gamma(\eta\gamma)/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 35 ⁴ ACHASOV 14 SND 1.15–2.00 $e^+ e^- \rightarrow \eta\gamma$

 Γ_{14}/Γ

NODE=M105R06
 NODE=M105R06

¹ $\omega\pi$ not included.

² Using ABELE 97.

³ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.

⁴ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

⁵ From a fit to the $e^+ e^- \rightarrow \eta\pi^+ \pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta\rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

⁶ Reports the inverse of the quoted value as 12.3 ± 3.1 .

⁷ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

NODE=M105R;LINKAGE=BL
 NODE=M105R;LINKAGE=LK
 NODE=M105R5;LINKAGE=A

NODE=M105R2;LINKAGE=A

NODE=M105R4;LINKAGE=A

NODE=M105R4;LINKAGE=B
 NODE=M105R;LINKAGE=A

 $\rho(1450)$ REFERENCES

NODE=M105

AULCHENKO	15	PR D91 052013	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56793
MATVIENKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55912
PDG	14	CPC 38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
		Translated from ZETFP 94 796.			
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ACHASOV	11	JETP 113 75	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=16721
		Translated from ZETF 140 87.			
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
ALOISIO	05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=51112
SCHAEEL	05C	PRPL 421 191	S. Schaeel <i>et al.</i>	(ALEPH Collab.)	REFID=50845
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
ALEXANDER	01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48391
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)	REFID=47468
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE	99C	PL B450 275	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46916
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BARATE	97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45622
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
BERTIN	97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45763
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
BISELLO	91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)	REFID=41752
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=41632
FUKUI	91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41581
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
CLEGG	88	ZPHY C40 313	A.B. Clegg, A. Donnachie	(MCHS, LANC)	REFID=40922
DIEKMANN	88	PRPL 159 99	B. Diekmann	(BONN)	REFID=40272
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40418
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=20246
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
		Translated from ZETFP 37 613.			
ASTON	80C	PL 92B 211	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=20652
BARBER	80C	ZPHY C4 169	D.P. Barber <i>et al.</i>	(DARE, LANC, SHEF)	REFID=20653
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054

η(1475)

$$I^G(J^{PC}) = 0^+(0^-+)$$

See also the η(1405).

NODE=M175

NODE=M175

NODE=M175205

NODE=M175M5
NODE=M175M5

η(1475) MASS

K $\bar{K}\pi$ MODE (K*(892) K dominant)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1476 ± 4 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
1469 ± 14 ± 13	74	ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
1460 ± 19	3651	NICHITIU	02 OBLX	
1485 ± 8 ± 5	20k	ADAMS	01B B852	18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$
1500 ± 10		CICALO	99 OBLX	$0\bar{p}p \rightarrow K^\pm K_S^0\pi^\mp\pi^+\pi^-$
1464 ± 10		BERTIN	97 OBLX	$0\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
1460 ± 10		BERTIN	95 OBLX	$0\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
1490 ⁺¹⁴⁺³ ₋₈₋₁₆	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0K^\pm\pi^\mp$
1475 ± 4		RATH	89 MPS	21.4 $\pi^-p \rightarrow nK_S^0K_S^0\pi^0$
1565 ± 8 ⁺⁰ ₋₆₃		¹ ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0K_S^0\eta$
1421 ± 14		AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$


OCCUR=2
OCCUR=2

OCCUR=2

NODE=M175M5;LINKAGE=A

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Could also be the η(1405).



η(1475) WIDTH

K $\bar{K}\pi$ MODE (K*(892) K dominant)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
85 ± 9 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
67 ± 18 ± 7	74	ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
120 ± 19	3651	NICHITIU	02 OBLX	
98 ± 18 ± 3	20k	ADAMS	01B B852	18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$
100 ± 20		CICALO	99 OBLX	$0\bar{p}p \rightarrow K^\pm K_S^0\pi^\mp\pi^+\pi^-$
105 ± 15		BERTIN	97 OBLX	$0.0\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
105 ± 15		BERTIN	95 OBLX	$0\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
63 ± 18		AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
54 ⁺³⁷⁺¹³ ₋₂₁₋₂₄		BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0K^\pm\pi^\mp$
51 ± 13		RATH	89 MPS	21.4 $\pi^-p \rightarrow nK_S^0K_S^0\pi^0$
54 ⁺¹⁴⁺²¹ ₋₁₃₋₂₈		¹ ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0K_S^0\eta$

NODE=M175210


NODE=M175W5
NODE=M175W5

OCCUR=2
OCCUR=2
OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Could also be the $\eta(1405)$.

NODE=M175W5;LINKAGE=A



$\eta(1475)$ DECAY MODES

NODE=M175215;NODE=M175

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}\pi$	dominant
Γ_2 $K\bar{K}^*(892) + c.c.$	seen
Γ_3 $a_0(980)\pi$	seen
Γ_4 $\gamma\gamma$	seen
Γ_5 $K_S^0 K_S^0 \eta$	possibly seen

DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=8;OUR EVAL;→ UNCHECKED ←

$\eta(1475)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M175220

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_1\Gamma_4/\Gamma$
VALUE (keV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
0.23 ± 0.05 ± 0.05	74	¹ ACHARD	07 L3	$183-209 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$

NODE=M175G2
 NODE=M175G2

••• We do not use the following data for averages, fits, limits, etc. •••

< 0.089	90	^{2,3} AHOHE	05 CLE2	$10.6 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
---------	----	----------------------	---------	--

¹ Supersedes ACCIARRI 01G. Compatible with K^*K decay. Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6895$.

² Using $\eta(1475)$ mass of 1481 MeV and width of 48 MeV. The upper limit increases to 0.140 keV if the world average value, 87 MeV, of the width is used.

³ Assuming three-body phase-space decay to $K_S^0 K^\pm \pi^\mp$.

NODE=M175G2;LINKAGE=CH

NODE=M175G2;LINKAGE=AH

NODE=M175G2;LINKAGE=B3

$\eta(1475)$ BRANCHING RATIOS

NODE=M175225

$\Gamma(K\bar{K}^*(892) + c.c.)/\Gamma(K\bar{K}\pi)$		Γ_2/Γ_1	
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M175R1
 NODE=M175R1

••• We do not use the following data for averages, fits, limits, etc. •••

0.50 ± 0.10	¹ BAILLON	67 HBC	$0.0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
-------------	----------------------	--------	--

¹ Data could also refer to $\eta(1405)$.

NODE=M175R;LINKAGE=BL

$\Gamma(K\bar{K}^*(892) + c.c.)/[\Gamma(K\bar{K}^*(892) + c.c.) + \Gamma(a_0(980)\pi)]$		$\Gamma_2/(\Gamma_2 + \Gamma_3)$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT

NODE=M175R6
 NODE=M175R6

••• We do not use the following data for averages, fits, limits, etc. •••

< 0.25	90	EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
--------	----	---------	----------	---

$\eta(1475)$ REFERENCES

ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56785
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=51698
AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)	REFID=50764
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
CICALO	99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)	REFID=47394
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
BERTIN	95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=44614
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)	REFID=40924
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
BAILLON	67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)	REFID=20407

NODE=M175

REFID=56785
REFID=51698
REFID=50764
REFID=48848
REFID=48319
REFID=49649
REFID=47394
REFID=45417
REFID=44614
REFID=41584
REFID=41578
REFID=40924
REFID=21314
REFID=20407

NODE=M152

 $f_0(1500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See also the mini-reviews on scalar mesons under $f_0(500)$ (see the index for the page number) and on non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M152

 $f_0(1500)$ MASS

NODE=M152M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1504 ± 6 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		

NODE=M152M

NODE=M152M

1468 ⁺¹⁴⁺²³ ₋₁₅₋₇₄	5.5k	¹ ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
1466 ± 6 ± 20		ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1515 ± 12		² BARBERIS	00A	450 $pp \rightarrow p_f\eta\eta p_S$
1511 ± 9		^{2,3} BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_S$
1510 ± 8		² BARBERIS	00E	450 $pp \rightarrow p_f\eta\eta p_S$
1522 ± 25		BERTIN	98 OBLX	0.05–0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
1449 ± 20		² BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1515 ± 20		ABELE	96B CBAR	0.0 $\bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
1500 ± 15		⁴ AMSLER	95B CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$
1505 ± 15		⁵ AMSLER	95C CBAR	0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1447 ± 16 ± 13	163	^{6,7} DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1442 ± 9 ± 4	261	^{6,7} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1486 ± 10		² ANISOVICH	09 RVUE	0.0 $\bar{p}p, \pi N$
1470 ± 60	568	⁸ KLEMP	08 E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$
1470 ⁺⁶⁺⁷² ₋₇₋₂₅₅		⁹ UEHARA	08A BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1495 ± 4		AMSLER	06 CBAR	0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
1539 ± 20	9.9k	AUBERT	06O BABR	$B^+ \rightarrow K^+K^+K^-$
1473 ± 5	80k	^{10,11} UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
1478 ± 6		VLADIMIRSK...	06 SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1493 ± 7		¹⁰ BINON	05 GAMS	33 $\pi^-p \rightarrow \eta\eta n$
1524 ± 14	1400	¹² GARMASH	05 BELL	$B^+ \rightarrow K^+K^+K^-$
1489 ⁺⁸ ₋₄		¹³ ANISOVICH	03 RVUE	
1490 ± 30		¹⁰ ABELE	01 CBAR	0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
1497 ± 10		¹⁰ BARBERIS	99 OMEG	450 $pp \rightarrow p_S p_f K^+ K^-$
1502 ± 10		¹⁰ BARBERIS	99B OMEG	450 $pp \rightarrow p_S p_f \pi^+ \pi^-$
1502 ± 12 ± 10		¹⁴ BARBERIS	99D OMEG	450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$
1530 ± 45		¹⁰ BELLAZZINI	99 GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
1505 ± 18		¹⁰ FRENCH	99	300 $pp \rightarrow p_f(K^+K^-)p_S$
1447 ± 27		¹⁵ KAMINSKI	99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
1580 ± 80		¹⁰ ALDE	98 GAM4	100 $\pi^-p \rightarrow \pi^0\pi^0 n$
1499 ± 8		² ANISOVICH	98B RVUE	Compilation
~ 1520		REYES	98 SPEC	800 $pp \rightarrow p_S p_f K_S^0 K_S^0$
1510 ± 20		² BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
~ 1475		FRABETTI	97D E687	$D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 1505		ABELE	96 CBAR	0.0 $\bar{p}p \rightarrow 5\pi^0$
1500 ± 8		² ABELE	96C RVUE	Compilation

OCCUR=2

1460±20	120	10	AMELIN	96B	VES	37	$\pi^- A \rightarrow \eta\eta\pi^- A$
1500± 8			BUGG	96	RVUE		
1500±10		16	AMSLER	95D	CBAR	0.0	$\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta,$ $\pi^0\pi^0\eta$
1445± 5		17	ANTINORI	95	OMEG	300,450	$pp \rightarrow pp2(\pi^+\pi^-)$
1497±30		10	ANTINORI	95	OMEG	300,450	$pp \rightarrow pp\pi^+\pi^-$
~ 1505			BUGG	95	MRK3		$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1446± 5		10	ABATZIS	94	OMEG	450	$pp \rightarrow pp2(\pi^+\pi^-)$
1545±25		10	AMSLER	94E	CBAR	0.0	$\bar{p}p \rightarrow \pi^0\eta\eta'$
1520±25		2,18	ANISOVICH	94	CBAR	0.0	$\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
1505±20		2,19	BUGG	94	RVUE		$\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0\pi^0$
1560±25		10	AMSLER	92	CBAR	0.0	$\bar{p}p \rightarrow \pi^0\eta\eta$
1550±45± 30		10	BELADIDZE	92C	VES	36	$\pi^- Be \rightarrow \pi^- \eta' \eta Be$
1449± 4		10	ARMSTRONG	89E	OMEG	300	$pp \rightarrow pp2(\pi^+\pi^-)$
1610±20		10	ALDE	88	GAM4	300	$\pi^- N \rightarrow \pi^- N2\eta$
~ 1525			ASTON	88D	LASS	11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1570±20	600	10	ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$
1575±45		20	ALDE	86D	GAM4	100	$\pi^- p \rightarrow 2\eta n$
1568±33		10	BINON	84C	GAM2	38	$\pi^- p \rightarrow \eta\eta' n$
1592±25		10	BINON	83	GAM2	38	$\pi^- p \rightarrow 2\eta n$
1525± 5		10	GRAY	83	DBC	0.0	$\bar{p}N \rightarrow 3\pi$

OCCUR=2


- 1 From partial wave analysis including all possible combinations of $0^{++}, 2^{++},$ and 4^{++} resonances.
- 2 T-matrix pole.
- 3 Average between $\pi^+\pi^-2\pi^0$ and $2(\pi^+\pi^-)$.
- 4 T-matrix pole, supersedes ANISOVICH 94.
- 5 T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.
- 6 Using CLEO-c data but not authored by the CLEO Collaboration.
- 7 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 109$ MeV.
- 8 Reanalysis of AITALA 01A data. This state could also be $f_0(1370)$.
- 9 Breit-Wigner mass. May also be the $f_0(1370)$.
- 10 Breit-Wigner mass.
- 11 Statistical error only.
- 12 Breit-Wigner, solution 1, PWA ambiguous.
- 13 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0\pi^0 n, \pi^- p \rightarrow K\bar{K}n,$
 $\pi^+\pi^- \rightarrow \pi^+\pi^-, \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta, \pi^+\pi^-\pi^0, K^+K^-\pi^0, K_S^0 K_S^0 \pi^0,$
 $K^+K_S^0\pi^-$ at rest, $\bar{p}n \rightarrow \pi^-\pi^-\pi^+, K_S^0 K^-\pi^0, K_S^0 K_S^0 \pi^-$ at rest.
- 14 Supersedes BARBERIS 99 and BARBERIS 99B.
- 15 T-matrix pole on sheet $--+$.
- 16 T-matrix pole. Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.
- 17 Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.
- 18 From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$.
- 19 Reanalysis of ANISOVICH 94 data.
- 20 From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152M;LINKAGE=C

NODE=M152M;LINKAGE=PP
 NODE=M152M;LINKAGE=PC
 NODE=M152M;LINKAGE=D
 NODE=M152M;LINKAGE=D1
 NODE=M152M;LINKAGE=D1
 NODE=M152M;LINKAGE=F
 NODE=M152M;LINKAGE=G
 NODE=M152M;LINKAGE=KL
 NODE=M152M;LINKAGE=UE
 NODE=M152M;LINKAGE=E
 NODE=M152M;LINKAGE=ST
 NODE=M152M;LINKAGE=GA
 NODE=M152M;LINKAGE=KM

NODE=M152M;LINKAGE=BD
 NODE=M152M;LINKAGE=TK
 NODE=M152M;LINKAGE=AB

NODE=M152M;LINKAGE=B
 NODE=M152M;LINKAGE=A
 NODE=M152M;LINKAGE=C1
 NODE=M152M;LINKAGE=AZ



$f_0(1500)$ WIDTH

NODE=M152W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
109± 7 OUR AVERAGE				
136 ⁺ ₋ 41 ⁺ ₋ 28 ⁺ ₋ 26 ⁻ ₋ 100	5.5k	21 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
108 ⁺ ₋ 14 ⁺ ₋ ± 25		ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
110± 24		22 BARBERIS	00A	450 $pp \rightarrow p_f\eta\eta p_s$
102± 18		22,23 BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
110± 16		22 BARBERIS	00E	450 $pp \rightarrow p_f\eta\eta p_s$
108± 33		BERTIN	98 OBLX	0.05-0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
114± 30		22 BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
105± 15		ABELE	96B CBAR	0.0 $\bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
120± 25		24 AMSLER	95B CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$
120± 30		25 AMSLER	95C CBAR	0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
114± 10		22 ANISOVICH	09 RVUE	0.0 $\bar{p}p, \pi N$
90 ⁺ ₋ 2 ⁺ ₋ 50 ⁺ ₋ 1 ⁻ ₋ 22		26 UEHARA	08A BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
121± 8		AMSLER	06 CBAR	0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
257± 33	9.9k	AUBERT	06O BABR	$B^+ \rightarrow K^+K^+K^-$
108± 9	80k	27,28 UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
119± 10		VLADIMIRSK...	06 SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
90± 15		27 BINON	05 GAMS	33 $\pi^-p \rightarrow \eta\eta n$
136± 23	1400	29 GARMASH	05 BELL	$B^+ \rightarrow K^+K^+K^-$
102± 10		30 ANISOVICH	03 RVUE	
140± 40		27 ABELE	01 CBAR	0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
104± 25		27 BARBERIS	99 OMEG	450 $pp \rightarrow p_s p_f K^+ K^-$
131± 15		27 BARBERIS	99B OMEG	450 $pp \rightarrow p_s p_f \pi^+ \pi^-$
98± 18 ± 16		31 BARBERIS	99D OMEG	450 $pp \rightarrow K^+K^-, \pi^+\pi^-$
160± 50		27 BELLAZZINI	99 GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
100± 33		27 FRENCH	99	300 $pp \rightarrow p_f(K^+K^-)p_s$
108± 46		32 KAMINSKI	99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
280± 100		27 ALDE	98 GAM4	100 $\pi^-p \rightarrow \pi^0\pi^0 n$
130± 20		22 ANISOVICH	98B RVUE	Compilation
120± 35		22 BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
~ 100		FRABETTI	97D E687	$D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 169		ABELE	96 CBAR	0.0 $\bar{p}p \rightarrow 5\pi^0$
100± 30	120	27 AMELIN	96B VES	37 $\pi^- A \rightarrow \eta\eta\pi^- A$
132± 15		BUGG	96 RVUE	
154± 30		33 AMSLER	95D CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$
65± 10		34 ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
199± 30		27 ANTINORI	95 OMEG	300,450 $pp \rightarrow pp\pi^+\pi^-$
56± 12		27 ABATZIS	94 OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
100± 40		27 AMSLER	94E CBAR	0.0 $\bar{p}p \rightarrow \pi^0\eta\eta'$
148 ⁺ ₋ 20 ⁺ ₋ 25		22,35 ANISOVICH	94 CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
150± 20		22,36 BUGG	94 RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0\pi^0$
245± 50		27 AMSLER	92 CBAR	0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
153± 67 ± 50		27 BELADIDZE	92C VES	36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$
78± 18		27 ARMSTRONG	89E OMEG	300 $pp \rightarrow pp2(\pi^+\pi^-)$
170± 40		27 ALDE	88 GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$
150± 20	600	27 ALDE	87 GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
265± 65		37 ALDE	86D GAM4	100 $\pi^- p \rightarrow 2\eta n$
260± 60		27 BINON	84C GAM2	38 $\pi^- p \rightarrow \eta\eta' n$
210± 40		27 BINON	83 GAM2	38 $\pi^- p \rightarrow 2\eta n$
101± 13		27 GRAY	83 DBC	0.0 $\bar{p}N \rightarrow 3\pi$

NODE=M152W

OCCUR=2

OCCUR=2

NODE=M152W;LINKAGE=C

NODE=M152W;LINKAGE=PP
 NODE=M152W;LINKAGE=PC
 NODE=M152W;LINKAGE=D
 NODE=M152W;LINKAGE=D1
 NODE=M152W;LINKAGE=UE
 NODE=M152W;LINKAGE=E

²¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

²² T-matrix pole.

²³ Average between $\pi^+\pi^-2\pi^0$ and $2(\pi^+\pi^-)$.

²⁴ T-matrix pole, supersedes ANISOVICH 94.

²⁵ T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

²⁶ Breit-Wigner width. May also be the $f_0(1370)$.

²⁷ Breit-Wigner width.

- 28 Statistical error only.
 29 Breit-Wigner, solution 1, PWA ambiguous.
 30 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$,
 $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$,
 $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.
 31 Supersedes BARBERIS 99 and BARBERIS 99B.
 32 T-matrix pole on sheet $- - +$.
 33 T-matrix pole. Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AM-
 SLER 94D.
 34 Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.
 35 From a simultaneous analysis of the annihilations $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta \eta$.
 36 Reanalysis of ANISOVICH 94 data.
 37 From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152W;LINKAGE=ST
 NODE=M152W;LINKAGE=GA
 NODE=M152W;LINKAGE=KM

NODE=M152W;LINKAGE=BD
 NODE=M152W;LINKAGE=TK
 NODE=M152W;LINKAGE=AB

NODE=M152W;LINKAGE=B
 NODE=M152W;LINKAGE=A
 NODE=M152W;LINKAGE=C1
 NODE=M152W;LINKAGE=AZ

$f_0(1500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 $\pi \pi$	(34.9±2.3) %	1.2
Γ_2 $\pi^+ \pi^-$	seen	
Γ_3 $2\pi^0$	seen	
Γ_4 4π	(49.5±3.3) %	1.2
Γ_5 $4\pi^0$	seen	
Γ_6 $2\pi^+ 2\pi^-$	seen	
Γ_7 $2(\pi\pi)_{S\text{-wave}}$	seen	
Γ_8 $\rho\rho$	seen	
Γ_9 $\pi(1300)\pi$	seen	
Γ_{10} $a_1(1260)\pi$	seen	
Γ_{11} $\eta\eta$	(5.1±0.9) %	1.4
Γ_{12} $\eta\eta'(958)$	(1.9±0.8) %	1.7
Γ_{13} $K\bar{K}$	(8.6±1.0) %	1.1
Γ_{14} $\gamma\gamma$	not seen	

NODE=M152215;NODE=M152

DESIG=8
 DESIG=9
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=7
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=11;OUR EST;→ UNCHECKED ←
 DESIG=12;OUR EST;→ UNCHECKED ←
 DESIG=13;OUR EST;→ UNCHECKED ←
 DESIG=14;OUR EST;→ UNCHECKED ←
 DESIG=1
 DESIG=2
 DESIG=4
 DESIG=10;OUR EST;→ UNCHECKED ←

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 10 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 11.4$ for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-83			
x_{11}	11	-52		
x_{12}	-5	-31	29	
x_{13}	39	-67	33	6
	x_1	x_4	x_{11}	x_{12}

$f_0(1500) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M152217

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1 \Gamma_{14}/\Gamma$
--	-----	-------------	------	---------	-------------------------------

NODE=M152G1
 NODE=M152G1

- • • We do not use the following data for averages, fits, limits, etc. • • •

$33_{-6}^{+12} + 1809_{-21}$	38	UEHARA	08A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
not seen		ACCIARRI	01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0$, $E_{\text{cm}}^{\text{ee}} = 91$, 183–209 GeV
<460	95	BARATE	00E	ALEP	$\gamma\gamma \rightarrow \pi^+ \pi^-$

- 38 May also be the $f_0(1370)$. Multiplied by us by 3 to obtain the $\pi\pi$ value.

NODE=M152G1;LINKAGE=UE

$f_0(1500)$ BRANCHING RATIOS

NODE=M152220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ NODE=M152R8
NODE=M152R8

VALUE	DOCUMENT ID	TECN	COMMENT
0.454±0.104	BUGG	96	RVUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M152R10
NODE=M152R10

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BERTIN	98	OBLX 0.05–0.405 $\bar{p}p \rightarrow \pi^+\pi^+\pi^-$
possibly seen	FRABETTI	97D	E687 $D_s^\pm \rightarrow \pi^\mp\pi^\pm\pi^\pm$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(4\pi)/\Gamma(\pi\pi)$ Γ_4/Γ_1 NODE=M152R6
NODE=M152R6

VALUE	DOCUMENT ID	TECN	COMMENT
1.42±0.18 OUR FIT			Error includes scale factor of 1.2.
1.42±0.18 OUR AVERAGE			Error includes scale factor of 1.2.
1.37±0.16	BARBERIS	00D	450 $pp \rightarrow p_f 4\pi p_s$
2.1 ±0.6	³⁹ AMSLER	98	RVUE
2.1 ±0.2	⁴⁰ ANISOVICH	02D	SPEC Combined fit
3.4 ±0.8	³⁹ ABELE	96	CBAR 0.0 $\bar{p}p \rightarrow 5\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(\pi\pi)$ Γ_7/Γ_1 NODE=M152R14
NODE=M152R14

VALUE	DOCUMENT ID	TECN	COMMENT
0.42±0.26	⁴¹ ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^-\pi^0 p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$ Γ_7/Γ_4 NODE=M152R15
NODE=M152R15

VALUE	DOCUMENT ID	TECN	COMMENT
0.26±0.07	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\rho\rho)/\Gamma(4\pi)$ Γ_8/Γ_4 NODE=M152R16
NODE=M152R16

VALUE	DOCUMENT ID	TECN	COMMENT
0.13±0.08	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{S\text{-wave}})$ Γ_8/Γ_7 NODE=M152R11
NODE=M152R11

VALUE	DOCUMENT ID	COMMENT
3.3±0.5	BARBERIS	00C 450 $pp \rightarrow p_f \pi^+ \pi^- 2\pi^0 p_s$
2.6±0.4	BARBERIS	00C 450 $pp \rightarrow p_f 2(\pi^+ \pi^-) p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

OCCUR=2

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_9/Γ_4 NODE=M152R17
NODE=M152R17

VALUE	DOCUMENT ID	TECN	COMMENT
0.50±0.25	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_{10}/Γ_4 NODE=M152R18
NODE=M152R18

VALUE	DOCUMENT ID	TECN	COMMENT
0.12±0.05	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_{11}/Γ NODE=M152R1
NODE=M152R1

VALUE	DOCUMENT ID	TECN	COMMENT
large	ALDE	88	GAM4 300 $\pi^- N \rightarrow \eta\eta\pi^- N$
large	BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$

• • • We do not use the following data for averages, fits, limits, etc. • • •


$\Gamma(\eta\eta)/\Gamma(\pi\pi)$

Γ_{11}/Γ_1

NODE=M152R13
NODE=M152R13

VALUE	DOCUMENT ID	TECN	COMMENT
0.145±0.027 OUR FIT			Error includes scale factor of 1.5.
0.14 ±0.04 OUR AVERAGE			Error includes scale factor of 1.7. See the ideogram below.
0.080±0.033	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$
0.18 ±0.03	BARBERIS	00E	450 $pp \rightarrow p_f \eta \eta p_s$
0.230±0.097	42 AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta \eta \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.11 ±0.03	40 ANISOVICH	02D	SPEC Combined fit
0.078±0.013	43 ABELE	96C	RVUE Compilation
0.157±0.060	44 AMSLER	95D	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$

WEIGHTED AVERAGE
0.14±0.04 (Error scaled by 1.7)



$\Gamma(4\pi^0)/\Gamma(\eta\eta)$

Γ_5/Γ_{11}

NODE=M152R5
NODE=M152R5

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.8±0.3	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$

$\Gamma(\eta\eta'(958))/\Gamma(\pi\pi)$

Γ_{12}/Γ_1

NODE=M152R12
NODE=M152R12

VALUE	DOCUMENT ID	TECN	COMMENT
0.055±0.024 OUR FIT			Error includes scale factor of 1.8.
0.095±0.026	BARBERIS	00A	450 $pp \rightarrow p_f \eta \eta p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.005±0.003	40 ANISOVICH	02D	SPEC Combined fit

$\Gamma(\eta\eta'(958))/\Gamma(\eta\eta)$

Γ_{12}/Γ_{11}

NODE=M152R2
NODE=M152R2

VALUE	DOCUMENT ID	TECN	COMMENT
0.38±0.16 OUR FIT			Error includes scale factor of 1.9.
0.29±0.10	45 AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta \eta \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.05±0.03	40 ANISOVICH	02D	SPEC Combined fit
0.84±0.23	ABELE	96C	RVUE Compilation
2.7 ±0.8	BINON	84C	GAM2 38 $\pi^- p \rightarrow \eta \eta' n$

$\Gamma(K\bar{K})/\Gamma_{total}$

Γ_{13}/Γ

NODE=M152R9
NODE=M152R9

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.044±0.021	BUGG	96	RVUE

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_{13}/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.246±0.026 OUR FIT			
0.241±0.028 OUR AVERAGE			
0.25 ±0.03	⁴⁶ BARGIOTTI 03	OBLX	$\bar{p}p$
0.19 ±0.07	⁴⁷ ABELE 98	CBAR	$0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.16 ±0.05	⁴⁰ ANISOVICH 02D	SPEC	Combined fit
0.33 ±0.03 ±0.07	BARBERIS 99D	OMEG	$450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$
0.20 ±0.08	⁴⁸ ABELE 96B	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$

NODE=M152R7
 NODE=M152R7

 $\Gamma(K\bar{K})/\Gamma(\eta\eta)$ Γ_{13}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
1.69±0.33 OUR FIT				Error includes scale factor of 1.4.
1.85±0.41		BARBERIS 00E		$450 pp \rightarrow p_f \eta \eta p_s$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.5 ±0.6		⁴⁰ ANISOVICH 02D	SPEC	Combined fit
<0.4	90	⁴⁹ PROKOSHKIN 91	GAM4	$300 \pi^- p \rightarrow \pi^- p \eta \eta$
<0.6		⁵⁰ BINON 83	GAM2	$38 \pi^- p \rightarrow 2 \eta n$
³⁹ Excluding $\rho\rho$ contribution to 4π .				
⁴⁰ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.				
⁴¹ From the combined data of ABELE 96 and ABELE 96C.				
⁴² Using AMSLER 95B ($3\pi^0$).				
⁴³ 2π width determined to be 60 ± 12 MeV.				
⁴⁴ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.				
⁴⁵ Using AMSLER 94E ($\eta \eta' \pi^0$).				
⁴⁶ Coupled channel analysis of $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.				
⁴⁷ Using $\pi^0 \pi^0$ from AMSLER 95B.				
⁴⁸ Using AMSLER 95B ($3\pi^0$), AMSLER 94C ($2\pi^0 \eta$) and SU(3).				
⁴⁹ Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production.				
⁵⁰ Using ETKIN 82B and COHEN 80.				

NODE=M152R4
 NODE=M152R4

NODE=M152R6;LINKAGE=C
 NODE=M152R;LINKAGE=CH

NODE=M152R;LINKAGE=KZ
 NODE=M152R3;LINKAGE=A
 NODE=M152R3;LINKAGE=CM
 NODE=M152R3;LINKAGE=AB
 NODE=M152R2;LINKAGE=A
 NODE=M152R;LINKAGE=BG
 NODE=M152R7;LINKAGE=A
 NODE=M152R7;LINKAGE=D
 NODE=M152R4;LINKAGE=BZ
 NODE=M152R4;LINKAGE=A

 $f_0(1500)$ REFERENCES

DOBBS 15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
ABLIKIM 13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)
ANISOVICH 09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
KLEMPPT 08	EPJ C55 39	E. Klempt, M. Matveev, A.V. Sarantsev	(BONN+)
UEHARA 08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM 06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
AMSLER 06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
AUBERT 06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG 06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSK... 06	PAN 69 493	V.V. Vladimirov <i>et al.</i>	(ITEP, Moscow)
	Translated from YAF 69 515.		
BINON 05	PAN 68 960	F. Binon <i>et al.</i>	
	Translated from YAF 68 998.		
GARMASH 05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ANISOVICH 03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BARGIOTTI 03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH 02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
	Translated from YAF 65 1583.		
ABELE 01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE 01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACCIARRI 01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
AITALA 01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BARATE 00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARBERIS 00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 00D	PL B474 423	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS 99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS 99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
BELLAZZINI 99	PL B467 296	R. Bellazzini <i>et al.</i>	
FRENCH 99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)
KAMINSKI 99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)
ABELE 98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ALDE 98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also	PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
	Translated from YAF 62 446.		
AMSLER 98	RMP 70 1293	C. Amsler	
ANISOVICH 98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
	Translated from UFN 168 481.		

NODE=M152

REFID=56805
 REFID=55387
 REFID=52719
 REFID=52286
 REFID=52309
 REFID=51507
 REFID=51136
 REFID=51141
 REFID=51004
 REFID=51063
 REFID=51191
 REFID=50780
 REFID=50641
 REFID=49401
 REFID=49217
 REFID=48580
 REFID=48831
 REFID=48334
 REFID=48356
 REFID=48321
 REFID=48004
 REFID=47428
 REFID=47957
 REFID=47959
 REFID=47960
 REFID=47961
 REFID=46921
 REFID=46922
 REFID=47395
 REFID=47400
 REFID=47491
 REFID=46927
 REFID=45863
 REFID=46605
 REFID=46914
 REFID=46601
 REFID=46331

BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
REYES	98	PRL 81 4079	M.A. Reyes <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMELIN	96B	PAN 59 976	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
		Translated from YAF 59 1021.		
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94E	PL B340 259	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)
		Translated from YAF 55 2748.		
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2, GAM4 Collab.)
		Translated from DANS 316 900.		
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
ALDE	88	PL B201 160	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
		Also SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
		Translated from YAF 38 934.		
GRAY	83	PR D27 307	L. Gray <i>et al.</i>	(SYRA)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL)

REFID=45782
REFID=46378
REFID=45758
REFID=45701
REFID=45554
REFID=45011
REFID=45038
REFID=45076
REFID=44725
REFID=45094
REFID=44377
REFID=44440
REFID=44441
REFID=44437
REFID=44438
REFID=44090
REFID=44091
REFID=44093
REFID=44080
REFID=43659
REFID=44078
REFID=43169
REFID=43175
REFID=41719
REFID=41011
REFID=40283
REFID=40330
REFID=40221
REFID=20765
REFID=21418
REFID=20750
REFID=20751
REFID=21370
REFID=20390
REFID=20381

$f_1(1510)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

OMITTED FROM SUMMARY TABLE
See the minireview under $\eta(1405)$.

NODE=M084

NODE=M084

$f_1(1510)$ MASS


NODE=M084M

NODE=M084M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1518 ± 5 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
1530 ± 10		ASTON	88C LASS	11 $K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
1512 ± 4	600	¹ BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ K^0 \pi^- n$
1526 ± 6	271	GAVILLET	82 HBC	4.2 $K^- p \rightarrow \Lambda K K \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1525		² BAUER	93B	$\gamma \gamma^* \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

¹ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ state.
² Not seen by AIHARA 88C in the $K_S^0 K^\pm \pi^\mp$ final state.


NODE=M084M;LINKAGE=A
 NODE=M084M;LINKAGE=C



$f_1(1510)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
73 ± 25 OUR AVERAGE	Error includes scale factor of 2.5. See the ideogram below.			
100 ± 40		ASTON	88C LASS	$11 K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
35 ± 15	600	³ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
107 ± 15	271	GAVILLET	82 HBC	$4.2 K^- p \rightarrow \Lambda K K \pi$

³From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ state.



NODE=M084W

NODE=M084W

NODE=M084W;LINKAGE=A

 $f_1(1510)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K}^*(892) + \text{c.c.}$	seen
Γ_2 $\pi^+ \pi^- \eta'$	seen

NODE=M084215;NODE=M084

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2 **$f_1(1510)$ BRANCHING RATIOS**

$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	230	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$	

NODE=M084225

NODE=M084R01
NODE=M084R01 **$f_1(1510)$ REFERENCES**

ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53684
BAUER	93B	PR D48 3976	D.A. Bauer <i>et al.</i>	(SLAC)	REFID=43678
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40564
ASTON	88C	PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP	REFID=40282
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
GAVILLET	82	ZPHY C16 119	P. Gavillet <i>et al.</i>	(CERN, CDEF, PADO+)	REFID=20877

NODE=M084

NODE=M013

 $f'_2(1525)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

 $f'_2(1525)$ MASS

NODE=M013205

VALUE (MeV)

DOCUMENT ID

1525±5 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

NODE=M013MX

→ UNCHECKED ←

PRODUCED BY PION BEAM

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1521±13		TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1547 ⁺¹⁰ ₋₂		¹ LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1496 ⁺⁹ ₋₈		² CHABAUD 81	ASPK	6 $\pi^- p \rightarrow K^+ K^- n$
1497 ⁺⁸ ₋₉		CHABAUD 81	ASPK	18.4 $\pi^- p \rightarrow K^+ K^- n$
1492±29		GORLICH 80	ASPK	17 $\pi^- p$ polarized $\rightarrow K^+ K^- n$
1502±25		³ CORDEN 79	OMEG	12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$
1480	14	CRENNELL 66	HBC	6.0 $\pi^- p \rightarrow K_S^0 K_S^0 n$

NODE=M013M1
NODE=M013M1

OCCUR=2

PRODUCED BY K^\pm BEAM

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

1523.3± 1.1 OUR AVERAGE Includes data from the datablock that follows this one.
Error includes scale factor of 1.1.NODE=M013M2
NODE=M013M2

1526.8± 4.3		ASTON 88D	LASS	11 $K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1504 ±12		BOLONKIN 86	SPEC	40 $K^- p \rightarrow K_S^0 K_S^0 Y$
1529 ± 3		ARMSTRONG 83B	OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$
1521 ± 6	650	AGUILAR-... 81B	HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
1521 ± 3	572	ALHARRAN 81	HBC	8.25 $K^- p \rightarrow \Lambda K \bar{K}$
1522 ± 6	123	BARREIRO 77	HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
1528 ± 7	166	EVANGELIS... 77	OMEG	10 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1527 ± 3	120	BRANDENB... 76C	ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1519 ± 7	100	AGUILAR-... 72B	HBC	3.9,4.6 $K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1514 ± 8	61	BINON 07	GAMS	32.5 $K^- p \rightarrow \eta \eta (\Lambda / \Sigma^0)$
1513 ±10		⁴ BARKOV 99	SPEC	40 $K^- p \rightarrow K_S^0 K_S^0 y$

PRODUCED IN e^+e^- ANNIHILATION AND PARTICLE DECAYS

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M013M3
NODE=M013M3**1521.9^{+1.8}_{-1.5} OUR AVERAGE** Error includes scale factor of 1.1.

1522.2± 2.8 ^{+5.3} _{-2.0}		AAIJ 13AN	LHCB	$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$
1513 ± 5 ⁺⁴ ₋₁₀	5.5k	⁵ ABLIKIM 13N	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1525.3 ^{+1.2+3.7} _{-1.4-2.1}		UEHARA 13	BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
1521 ± 5		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi K^+ K^-$
1518 ± 1 ± 3		ABE 04	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
1519 ± 2 ⁺¹⁵ ₋₅		BAI 03G	BES	$J/\psi \rightarrow \gamma K \bar{K}$
1523 ± 6	331	⁶ ACCIARRI 01H	L3	91, 183-209 $e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$
1535 ± 5 ± 4		ABREU 96C	DLPH	$Z^0 \rightarrow K^+ K^- + X$
1516 ± 5 ⁺⁹ ₋₁₅		BAI 96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
1531.6±10.0		AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
1515 ± 5		⁷ FALVARD 88	DM2	$J/\psi \rightarrow \phi K^+ K^-$
1525 ±10 ±10		BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1532 ± 3 ± 6	644	^{8,9} DOBBS 15		$J/\psi \rightarrow \gamma K^+ K^-$
1557 ± 9 ± 3	113	^{8,9} DOBBS 15		$\psi(2S) \rightarrow \gamma K^+ K^-$
1523 ± 5	870	¹⁰ SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$
1496 ± 2		¹¹ FALVARD 88	DM2	$J/\psi \rightarrow \phi K^+ K^-$

OCCUR=2

OCCUR=2

PRODUCED IN $\bar{p}p$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1530 ± 12	12 ANISOVICH	09	RVUE 0.0 $\bar{p}p, \pi N$
1513 ± 4	AMSLER	06	CBAR 0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
1508 ± 9	13 AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$

NODE=M013M9
 NODE=M013M9

CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1515 ± 15	BARBERIS	99	OMEG 450 $pp \rightarrow p_s p_f K^+ K^-$

NODE=M013M4
 NODE=M013M4

PRODUCED IN $e p$ COLLISIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1512 ± 3 ^{+1.4} _{-0.5}		14 CHEKANOV	08	ZEUS $e p \rightarrow K_S^0 K_S^0 X$

NODE=M013M10
 NODE=M013M10

• • • We do not use the following data for averages, fits, limits, etc. • • •

1537 ⁺⁹ ₋₈	84	15 CHEKANOV	04	ZEUS $e p \rightarrow K_S^0 K_S^0 X$
----------------------------------	----	-------------	----	--------------------------------------

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² CHABAUD 81 is a reanalysis of PAWLICKI 77 data.

³ From an amplitude analysis where the $f_2'(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious.

⁴ Systematic errors not estimated.

⁵ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

⁶ Supersedes ACCIARRI 95J.

⁷ From an analysis ignoring interference with $f_0(1710)$.

⁸ Using CLEO-c data but not authored by the CLEO Collaboration.

⁹ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 73$ MeV.

¹⁰ From analysis of L3 data at 91 and 183–209 GeV.

¹¹ From an analysis including interference with $f_0(1710)$.

¹² 4-poles, 5-channel K matrix fit.

¹³ T-matrix pole.

¹⁴ In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

¹⁵ Systematic errors not estimated.

NODE=M013M;LINKAGE=L
 NODE=M013M;LINKAGE=D
 NODE=M013M;LINKAGE=N

NODE=M013M2;LINKAGE=SK
 NODE=M013M3;LINKAGE=A

NODE=M013M;LINKAGE=HA
 NODE=M013M;LINKAGE=F1

NODE=M013M3;LINKAGE=B
 NODE=M013M3;LINKAGE=C

NODE=M013M3;LINKAGE=SC
 NODE=M013M;LINKAGE=F2

NODE=M013M9;LINKAGE=AN
 NODE=M013M;LINKAGE=TT

NODE=M013M10;LINKAGE=HE

NODE=M013M10;LINKAGE=CH

 $f_2'(1525)$ WIDTH

NODE=M013210

VALUE (MeV)	DOCUMENT ID	COMMENT
-------------	-------------	---------

NODE=M013WX

73⁺⁶₋₅ OUR FIT

76 ± 10	PDG	90	For fitting
---------	-----	----	-------------

PRODUCED BY PION BEAM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
102 ± 42	TIKHOMIROV	03	SPEC 40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
108 ⁺⁵ ₋₂	16 LONGACRE	86	MPS 22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
69 ⁺²² ₋₁₆	17 CHABAUD	81	ASPK 6 $\pi^- p \rightarrow K^+ K^- n$
137 ⁺²³ ₋₂₁	CHABAUD	81	ASPK 18.4 $\pi^- p \rightarrow K^+ K^- n$
150 ⁺⁸³ ₋₅₀	GORLICH	80	ASPK 17 $\pi^- p$ polarized $\rightarrow K^+ K^- n$
165 ± 42	18 CORDEN	79	OMEG 12–15 $\pi^- p \rightarrow \pi^+ \pi^- n$
92 ⁺³⁹ ₋₂₂	19 POLYCHRO...	79	STRC 7 $\pi^- p \rightarrow n K_S^0 K_S^0$

NODE=M013W1
 NODE=M013W1

• • • We do not use the following data for averages, fits, limits, etc. • • •

OCCUR=2

PRODUCED BY K^\pm BEAM

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M013W2
NODE=M013W2

81.4 $^{+2.2}_{-1.9}$ OUR AVERAGE Includes data from the datablock that follows this one.

90 ± 12		ASTON	88D	LASS	11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
73 ± 18		BOLONKIN	86	SPEC	40	$K^- p \rightarrow K_S^0 K_S^0 \Upsilon$
83 ± 15		ARMSTRONG	83B	OMEG	18.5	$K^- p \rightarrow K^- K^+ \Lambda$
85 ± 16	650	AGUILAR-...	81B	HBC	4.2	$K^- p \rightarrow \Lambda K^+ K^-$
80 $^{+14}_{-11}$	572	ALHARRAN	81	HBC	8.25	$K^- p \rightarrow \Lambda K \bar{K}$
72 ± 25	166	EVANGELIS...	77	OMEG	10	$K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
69 ± 22	100	AGUILAR-...	72B	HBC	3.9,4.6	$K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
92 $^{+25}_{-16}$	61	BINON	07	GAMS	32.5	$K^- p \rightarrow \eta \eta (\Lambda / \Sigma^0)$
75 ± 20	20	BARKOV	99	SPEC	40	$K^- p \rightarrow K_S^0 K_S^0 \Upsilon$
62 $^{+19}_{-14}$	123	BARREIRO	77	HBC	4.15	$K^- p \rightarrow \Lambda K_S^0 K_S^0$
61 ± 8	120	BRANDENB...	76C	ASPK	13	$K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$

PRODUCED IN $e^+ e^-$ ANNIHILATION AND PARTICLE DECAYS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M013W3
NODE=M013W3

The data in this block is included in the average printed for a previous datablock.

81.4 $^{+2.4}_{-2.0}$ OUR AVERAGE

84 ± 6	$^{+10}_{-5}$	AAIJ	13AN	LHCB	$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$	
75 $^{+12}_{-10}$	$^{+16}_{-8}$	5.5k	21	ABLIKIM	13N BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$	
82.9 $^{+2.1}_{-2.2}$	$^{+3.3}_{-2.0}$	UEHARA	13	BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$	
77 ± 15		ABLIKIM	05	BES2	$J/\psi \rightarrow \phi K^+ K^-$	
82 ± 2	± 3	ABE	04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
75 ± 4	$^{+15}_{-5}$	BAI	03G	BES	$J/\psi \rightarrow \gamma K \bar{K}$	
100 ± 15	331	22 ACCIARRI	01H	L3	$91, 183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
60 ± 20	± 19	ABREU	96C	DLPH	$Z^0 \rightarrow K^+ K^- + X$	
60 ± 23	$^{+13}_{-20}$	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$	
103 ± 30		AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$	
62 ± 10		23 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$	
85 ± 35		BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
104 ± 10	870	24 SCHEGELSKY	06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	
100 ± 3		25 FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$	

OCCUR=2

PRODUCED IN $\bar{p} p$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M013W9
NODE=M013W9

79 ± 8	26	AMSLER	02	CBAR	$0.9 \bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
128 ± 20	27	ANISOVICH	09	RVUE	$0.0 \bar{p} p, \pi N$
76 ± 6		AMSLER	06	CBAR	$0.9 \bar{p} p \rightarrow K^+ K^- \pi^0$

CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M013W4
NODE=M013W4

70 ± 25	BARBERIS	99	OMEG	450	$p p \rightarrow p_s p_f K^+ K^-$
-------------------------------	----------	----	------	-----	-----------------------------------

PRODUCED IN $e p$ COLLISIONS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M013W10
NODE=M013W10

83 ± 9	$^{+5}_{-4}$	28	CHEKANOV	08	ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
------------------------------	--------------	----	----------	----	------	---------------------------------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

50 $^{+34}_{-22}$	84	29	CHEKANOV	04	ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
-------------------	----	----	----------	----	------	---------------------------------

- 16 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- 17 CHABAUD 81 is a reanalysis of PAWLICKI 77 data.
- 18 From an amplitude analysis where the $f_2'(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious.
- 19 From a fit to the D with $f_2(1270)$ - $f_2'(1525)$ interference. Mass fixed at 1516 MeV.
- 20 Systematic errors not estimated.
- 21 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.
- 22 Supersedes ACCIARRI 95J.
- 23 From an analysis ignoring interference with $f_0(1710)$.
- 24 From analysis of L3 data at 91 and 183–209 GeV.
- 25 From an analysis including interference with $f_0(1710)$.
- 26 T-matrix pole.
- 27 4-poles, 5-channel K matrix fit.
- 28 In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.
- 29 Systematic errors not estimated.

NODE=M013W;LINKAGE=L
 NODE=M013W;LINKAGE=D
 NODE=M013W;LINKAGE=N

 NODE=M013W;LINKAGE=M
 NODE=M013W2;LINKAGE=SK
 NODE=M013W3;LINKAGE=A

 NODE=M013W;LINKAGE=HA
 NODE=M013W;LINKAGE=F1
 NODE=M013W3;LINKAGE=SC
 NODE=M013W;LINKAGE=F2
 NODE=M013W;LINKAGE=TT
 NODE=M013W9;LINKAGE=AN
 NODE=M013W10;LINKAGE=HE

 NODE=M013W10;LINKAGE=CH

$f_2'(1525)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	(88.7 \pm 2.2) %
Γ_2 $\eta\eta$	(10.4 \pm 2.2) %
Γ_3 $\pi\pi$	(8.2 \pm 1.5) $\times 10^{-3}$
Γ_4 $K\bar{K}^*(892) + c.c.$	
Γ_5 $\pi K\bar{K}$	
Γ_6 $\pi\pi\eta$	
Γ_7 $\pi^+\pi^+\pi^-\pi^-$	
Γ_8 $\gamma\gamma$	(1.10 \pm 0.14) $\times 10^{-6}$

NODE=M013215;NODE=M013

DESIG=2
 DESIG=4
 DESIG=1
 DESIG=3
 DESIG=6
 DESIG=5
 DESIG=7
 DESIG=8

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 2 partial widths, a combination of partial widths obtained from integrated cross sections, and 3 branching ratios uses 17 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 14.3$ for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100			
x_3	-6	-1		
x_8	-6	6	1	
Γ	-23	23	-1	-56
	x_1	x_2	x_3	x_8

Mode	Rate (MeV)
Γ_1 $K\bar{K}$	65 $\begin{matrix} +5 \\ -4 \end{matrix}$
Γ_2 $\eta\eta$	7.6 \pm 1.8
Γ_3 $\pi\pi$	0.60 \pm 0.12
Γ_8 $\gamma\gamma$	(8.1 \pm 0.9) $\times 10^{-5}$

DESIG=2
 DESIG=4
 DESIG=1
 DESIG=8

$f_2'(1525)$ PARTIAL WIDTHS

NODE=M013220

$\Gamma(K\bar{K})$	VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	Γ_1
--------------------	-------------	-------------	------	---------	------------

NODE=M013W6
 NODE=M013W6

65⁺⁵₋₄ OUR FIT

63⁺⁶₋₅

³⁰ LONGACRE 86 MPS 22 $\pi^- p \rightarrow K_S^0 K_S^0 n$

$\Gamma(\eta\eta)$ Γ_2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

7.6±1.8 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.0±0.8	870	³¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
24 $\begin{smallmatrix} +3 \\ -1 \end{smallmatrix}$		³⁰ LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$

NODE=M013W7
 NODE=M013W7

 $\Gamma(\pi\pi)$ Γ_3

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.60±0.12 OUR FIT

1.4 $\begin{smallmatrix} +1.0 \\ -0.5 \end{smallmatrix}$		³⁰ LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
--	--	---------------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2 $\begin{smallmatrix} +1.0 \\ -0.2 \end{smallmatrix}$	870	³¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
--	-----	------------------------------	------	--

NODE=M013W5
 NODE=M013W5

 $\Gamma(\gamma\gamma)$ Γ_8

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.081±0.009 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 ±0.03	870	³¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
------------	-----	------------------------------	------	--

³⁰ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.³¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f_2'(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.

NODE=M013W8
 NODE=M013W8

NODE=M013PW;LINKAGE=L
 NODE=M013W8;LINKAGE=SC

 $f_2'(1525) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M013223

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_8/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.072 ±0.007 OUR FIT**0.072 ±0.007 OUR AVERAGE**

0.048 $\begin{smallmatrix} +0.067 \\ -0.008 \end{smallmatrix}$	$\begin{smallmatrix} +0.108 \\ -0.012 \end{smallmatrix}$	UEHARA 13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
0.0564±0.0048±0.0116		ABE 04	BELL	10.6 $e^+e^- \rightarrow e^+e^- K^+K^-$
0.076 ±0.006 ±0.011	331	³² ACCIARRI 01H	L3	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.067 ±0.008 ±0.015		³³ ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
0.11 $\begin{smallmatrix} +0.03 \\ -0.02 \end{smallmatrix}$	±0.02	BEHREND 89c	CELL	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.10 $\begin{smallmatrix} +0.04 \\ -0.03 \end{smallmatrix}$	$\begin{smallmatrix} +0.03 \\ -0.02 \end{smallmatrix}$	BERGER 88	PLUT	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.12 ±0.07 ±0.04		³³ AIHARA 86B	TPC	$e^+e^- \rightarrow e^+e^- K^+K^-$
0.11 ±0.02 ±0.04		³³ ALTHOFF 83	TASS	$e^+e^- \rightarrow e^+e^- K\bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0314±0.0050±0.0077		³⁴ ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
----------------------	--	----------------------------	-----	------------------------------------

NODE=M013G1
 NODE=M013G1

OCCUR=2

³² Supersedes ACCIARRI 95J. From analysis of L3 data at 91 and 183–209 GeV,³³ Using an incoherent background.³⁴ Using a coherent background.

NODE=M013G;LINKAGE=HA
 NODE=M013G1;LINKAGE=A
 NODE=M013G1;LINKAGE=B

 $f_2'(1525) \text{ BRANCHING RATIOS}$

NODE=M013225

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	UEHARA 10A	BELL	10.6 $e^+e^- \rightarrow e^+e^- \eta\eta$
0.10±0.03	³⁵ PROKOSHKIN 91	GAM4	300 $\pi^- p \rightarrow \pi^- p \eta\eta$

³⁵ Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.

NODE=M013R8
 NODE=M013R8

NODE=M013R8;LINKAGE=B

$\Gamma(\eta\eta)/\Gamma(K\bar{K})$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

0.118±0.028 OUR FIT**0.115±0.028 OUR AVERAGE**

0.119±0.015±0.036	61	36	BINON	07	GAMS 32.5 $K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$
0.11 ±0.04		37	PROKOSHKIN	91	GAM4 300 $\pi^- p \rightarrow \pi^- p \eta\eta$
< 0.14	90		BARBERIS	00E	450 $p p \rightarrow p_f \eta \eta p_s$
< 0.50			BARNES	67	HBC 4.6,5.0 $K^- p$

³⁶ Using the compilation of the cross sections for $f'_2(1525)$ production in $K^- p$ collisions from ASTON 88D.

³⁷ Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.

NODE=M013R3
NODE=M013R3

NODE=M013R3;LINKAGE=BI

NODE=M013R3;LINKAGE=B

 $\Gamma(\pi\pi)/\Gamma_{total}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.0082±0.0016 OUR FIT**0.0075±0.0016 OUR AVERAGE**

0.007 ±0.002			COSTA...	80	OMEG 10 $\pi^- p \rightarrow K^+ K^- n$
0.027 ^{+0.071} _{-0.013}		38	GORLICH	80	ASPK 17,18 $\pi^- p$
0.0075±0.0025		38,39	MARTIN	79	RVUE
<0.06	95		AGUILAR-...	81B	HBC 4.2 $K^- p \rightarrow \Lambda K^+ K^-$
0.19 ±0.03			CORDEN	79	OMEG 12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$
<0.045	95		BARREIRO	77	HBC 4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
0.012 ±0.004		38	PAWLICKI	77	SPEC 6 $\pi N \rightarrow K^+ K^- N$
<0.063	90		BRANDENB...	76C	ASPK 13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
<0.0086		38	BEUSCH	75B	OSPK 8.9 $\pi^- p \rightarrow K^0 \bar{K}^0 n$

³⁸ Assuming that the $f'_2(1525)$ is produced by an one-pion exchange production mechanism.

³⁹ MARTIN 79 uses the PAWLICKI 77 data with different input value of the $f'_2(1525) \rightarrow K\bar{K}$ branching ratio.

NODE=M013R1
NODE=M013R1

NODE=M013R1;LINKAGE=C

NODE=M013R1;LINKAGE=D

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0092±0.0018 OUR FIT**0.075 ±0.035**

	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$
--	----------	----	---

 $[\Gamma(K\bar{K}^*(892) + c.c.) + \Gamma(\pi K\bar{K})]/\Gamma(K\bar{K})$ $(\Gamma_4 + \Gamma_5)/\Gamma_1$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.35	95	AGUILAR-...	72B	HBC 3.9,4.6 $K^- p$
<0.4	67	AMMAR	67	HBC

NODE=M013R7
NODE=M013R7

NODE=M013R5
NODE=M013R5

 $\Gamma(\pi\pi\eta)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.41	95	AGUILAR-...	72B	HBC 3.9,4.6 $K^- p$
<0.3	67	AMMAR	67	HBC

NODE=M013R4
NODE=M013R4

 $\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.32	95	AGUILAR-...	72B	HBC 3.9,4.6 $K^- p$
-------	----	-------------	-----	---------------------

NODE=M013R6
NODE=M013R6

$f'_2(1525)$ REFERENCES

NODE=M013

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
AAJ	13AN	PR D87 072004	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55137
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55387
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		REFID=52719
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52275
BINON	07	PAN 70 1713	F. Binon <i>et al.</i>	(GAMS Collab.)	REFID=52057
		Translated from YAF 70 1758.			
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)	REFID=51136
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=49672
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>		REFID=48580
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARKOV	99	JETPL 70 248	B.P. Barkov <i>et al.</i>		REFID=47379
		Translated from ZETFP 70 242.			
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
ACCIARRI	95J	PL B363 118	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=44615
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2, GAM4 Collab.)	REFID=41719
		Translated from DANS 316 900.			
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
PDG	90	PL B239 1	J.J. Hernandez <i>et al.</i>	(IFIC, BOST, CIT+)	REFID=40744
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40330
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BERGER	88	ZPHY C37 329	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=40566
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=20764
BOLONKIN	86	SJNP 43 776	B.V. Bolonkin <i>et al.</i>	(ITEP) JP	REFID=44646
		Translated from YAF 43 1211.			
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	83	PL 121B 216	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21408
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=20558
AGUILAR-...	81B	ZPHY C8 313	M. Aguilar-Benitez <i>et al.</i>	(CERN, CDEF+)	REFID=21104
ALHARRAN	81	NP B191 26	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=21403
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
COSTA...	80	NP B175 402	G. Costa de Beauregard <i>et al.</i>	(BARI, BONN+)	REFID=20737
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)	REFID=20738
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
BARREIRO	77	NP B121 237	F. Barreiro <i>et al.</i>	(CERN, AMST, NIJM+)	REFID=21392
EVANGELIS...	77	NP B127 384	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20540
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJP	REFID=20367
BRANDENB...	76C	NP B104 413	G.W. Brandenburg <i>et al.</i>	(SLAC)	REFID=20225
BEUSCH	75B	PL 60B 101	W. Beusch <i>et al.</i>	(CERN, ETH)	REFID=21390
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
AMMAR	67	PRL 19 1071	R. Ammar <i>et al.</i>	(NWES, ANL) JP	REFID=21382
BARNES	67	PRL 19 964	V.E. Barnes <i>et al.</i>	(BNL, SYRA) IJPC	REFID=21383
CRENNELL	66	PRL 16 1025	D.J. Crennell <i>et al.</i>	(BNL) I	REFID=20317

$f_2(1565)$

$I^G(J^{PC}) = 0^+(2^{++})$

OMITTED FROM SUMMARY TABLE

Seen mostly in antinucleon-nucleon annihilation. Needs confirmation in other channels.

NODE=M123

NODE=M123

NODE=M123M

NODE=M123M

$f_2(1565)$ MASS


VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1562±13 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.		
1590±10	1 AMELIN	06 VES	36 $\pi^- p \rightarrow \omega \omega n$
1552±13	2 AMSLER	02 CBAR	0.9 $\bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
1550±10±20	AMELIN	00 VES	37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1575±18	BERTIN	98 OBLX	0.05-0.405 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1507±15	2 BERTIN	97C OBLX	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
1565±20	MAY	90 ASTE	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1560±15	3 ANISOVICH	09 RVUE	0.0 $\bar{p} p, \pi N$
1598±11± 9	BAKER	99B SPEC	0 $\bar{p} p \rightarrow \omega \omega \pi^0$
1534±20	4 ABELE	96C RVUE	Compilation
~ 1552	5 AMSLER	95D CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
1598±72	BALOSHIN	95 SPEC	40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
1566 ⁺⁸⁰ ₋₅₀	6 ANISOVICH	94 CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0, \eta \eta \pi^0$
1502± 9	ADAMO	93 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1488±10	7 ARMSTRONG	93C E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1508±10	7 ARMSTRONG	93D E760	$\bar{p} p \rightarrow 3\pi^0 \rightarrow 6\gamma$
1525±10	7 ARMSTRONG	93D E760	$\bar{p} p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 1504	8 WEIDENAUER	93 ASTE	0.0 $\bar{p} N \rightarrow 3\pi^- 2\pi^+$
1540±15	7 ADAMO	92 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1515±10	9 AKER	91 CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$
1477± 5	BRIDGES	86C DBC	0.0 $\bar{p} N \rightarrow 3\pi^- 2\pi^+$

OCCUR=2

NODE=M123M;LINKAGE=AM
 NODE=M123M;LINKAGE=G
 NODE=M123M;LINKAGE=AN
 NODE=M123M;LINKAGE=AA
 NODE=M123M;LINKAGE=AB
 NODE=M123M;LINKAGE=C

NODE=M123M;LINKAGE=E
 NODE=M123M;LINKAGE=F
 NODE=M123M;LINKAGE=BA

- 1 Supersedes the $\omega \omega$ state of BELADIDZE 92B earlier assigned to the $f_2(1640)$.
- 2 T-matrix pole.
- 3 On sheet II in a two-pole solution.
- 4 T-matrix pole, large coupling to $\rho \rho$ and $\omega \omega$, could be $f_2(1640)$.
- 5 Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.
- 6 From a simultaneous analysis of the annihilations $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta \eta$ including AKER 91 data.
- 7 J^P not determined, could be partly $f_0(1500)$.
- 8 J^P not determined.
- 9 Superseded by AMSLER 95B.



$f_2(1565)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
134± 8 OUR AVERAGE			
140± 11	¹⁰ AMELIN	06 VES	$36 \pi^- p \rightarrow \omega \omega n$
113± 23	¹¹ AMSLER	02 CBAR	$0.9 \bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
130± 20±40	AMELIN	00 VES	$37 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
119± 24	BERTIN	98 OBLX	$0.05-0.405 \bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
130± 20	¹¹ BERTIN	97C OBLX	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
170± 40	MAY	90 ASTE	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
280± 40	¹² ANISOVICH	09 RVUE	$0.0 \bar{p} p, \pi N$
180± 60	¹³ ABELE	96C RVUE	Compilation
~ 142	¹⁴ AMSLER	95D CBAR	$0.0 \bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
263±101	BALOSHIN	95 SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$
166 ⁺ ₋₂₀	¹⁵ ANISOVICH	94 CBAR	$0.0 \bar{p} p \rightarrow 3\pi^0, \eta \eta \pi^0$
130± 10	¹⁶ ADAMO	93 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
148± 27	¹⁷ ARMSTRONG	93C E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
103± 15	¹⁷ ARMSTRONG	93D E760	$\bar{p} p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111± 10	¹⁷ ARMSTRONG	93D E760	$\bar{p} p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 206	¹⁸ WEIDENAUER	93 ASTE	$0.0 \bar{p} N \rightarrow 3\pi^- 2\pi^+$
132± 37	¹⁷ ADAMO	92 OBLX	$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
120± 10	¹⁹ AKER	91 CBAR	$0.0 \bar{p} p \rightarrow 3\pi^0$
116± 9	BRIDGES	86C DBC	$0.0 \bar{p} N \rightarrow 3\pi^- 2\pi^+$
¹⁰ Supersedes the $\omega \omega$ state of BELADIDZE 92B earlier assigned to the $f_2(1640)$.			
¹¹ T-matrix pole.			
¹² On sheet II in a two-pole solution.			
¹³ T-matrix pole, large coupling to $\rho \rho$ and $\omega \omega$, could be $f_2(1640)$.			
¹⁴ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.			
¹⁵ From a simultaneous analysis of the annihilations $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta \eta$ including AKER 91 data.			
¹⁶ Supersedes ADAMO 92.			
¹⁷ J^P not determined, could be partly $f_0(1500)$.			
¹⁸ J^P not determined.			
¹⁹ Superseded by AMSLER 95B.			

NODE=M123W

NODE=M123W

OCCUR=2

NODE=M123W;LINKAGE=AM
 NODE=M123W;LINKAGE=G
 NODE=M123W;LINKAGE=AN
 NODE=M123W;LINKAGE=CC
 NODE=M123W;LINKAGE=AB
 NODE=M123W;LINKAGE=D

NODE=M123W;LINKAGE=C
 NODE=M123W;LINKAGE=E
 NODE=M123W;LINKAGE=F
 NODE=M123W;LINKAGE=BA

 $f_2(1565)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi \pi$	seen
Γ_2 $\pi^+ \pi^-$	seen
Γ_3 $\pi^0 \pi^0$	seen
Γ_4 $\rho^0 \rho^0$	seen
Γ_5 $2\pi^+ 2\pi^-$	seen
Γ_6 $\eta \eta$	seen
Γ_7 $a_2(1320) \pi$	
Γ_8 $\omega \omega$	seen
Γ_9 $K \bar{K}$	
Γ_{10} $\gamma \gamma$	

NODE=M123215;NODE=M123

DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=8
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=9
 DESIG=10

 $f_2(1565)$ PARTIAL WIDTHS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.2±0.3	870	²⁰ SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	Γ_6
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2.0±1.0	870	²⁰ SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	Γ_9

NODE=M123225

NODE=M123W3
 NODE=M123W3

NODE=M123W1
 NODE=M123W1

$\Gamma(\gamma\gamma)$ Γ_{10}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.70±0.14	870	²⁰ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
-----------	-----	------------------------------	------	--

²⁰ From analysis of L3 data at 91 and 183–209 GeV, using $f_2(1565)$ mass of 1570 MeV, width of 160 MeV, $\Gamma(\pi\pi) = 25$ MeV, and SU(3) relations.

NODE=M123W2
NODE=M123W2

NODE=M123W1;LINKAGE=SC

 $f_2(1565)$ BRANCHING RATIOS

NODE=M123220

 $\Gamma(\pi\pi)/\Gamma_{total}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	BAKER	99B	SPEC $0 \bar{p}p \rightarrow \omega\omega\pi^0$
------	-------	-----	---

NODE=M123R5
NODE=M123R5

 $\Gamma(\pi^+\pi^-)/\Gamma_{total}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	BERTIN	98	OBLX $0.05-0.405 \bar{p}p \rightarrow$
------	--------	----	--

not seen	²¹ ANISOVICH	94B	RVUE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
----------	-------------------------	-----	---

seen	MAY	89	ASTE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
------	-----	----	---

²¹ ANISOVICH 94B is from a reanalysis of MAY 90.

NODE=M123R1
NODE=M123R1

NODE=M123R1;LINKAGE=A

 $\Gamma(\pi^0\pi^0)/\Gamma_{total}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	AMSLER	95B	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0$
------	--------	-----	--

NODE=M123R3
NODE=M123R3

 $\Gamma(\pi^+\pi^-)/\Gamma(\rho^0\rho^0)$ Γ_2/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.042±0.013	BRIDGES	86B	DBC $\bar{p}N \rightarrow 3\pi^- 2\pi^+$
-------------	---------	-----	--

NODE=M123R2
NODE=M123R2

 $\Gamma(\eta\eta)/\Gamma(\pi^0\pi^0)$ Γ_6/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.024±0.005±0.012	²² ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
-------------------	-------------------------	-----	---

²² J^P not determined, could be partly $f_0(1500)$.

NODE=M123R4
NODE=M123R4

NODE=M123R4;LINKAGE=E

 $\Gamma(\omega\omega)/\Gamma_{total}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	BAKER	99B	SPEC $0 \bar{p}p \rightarrow \omega\omega\pi^0$
------	-------	-----	---

NODE=M123R6
NODE=M123R6

 $f_2(1565)$ REFERENCES

NODE=M123

ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
BAKER	99B	PL B467 147	C.A. Baker <i>et al.</i>	
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94B	PR D50 1972	V.V. Anisovich <i>et al.</i>	(LOQM)
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)
MAY	90	ZPHY C46 203	B. May <i>et al.</i>	(ASTERIX Collab.)
MAY	89	PL B225 450	B. May <i>et al.</i>	(ASTERIX Collab.) IJP
BRIDGES	86B	PRL 56 215	D.L. Bridges <i>et al.</i>	(SYRA, CASE)
BRIDGES	86C	PRL 57 1534	D.L. Bridges <i>et al.</i>	(SYRA)

REFID=52719
REFID=51574

REFID=51185
REFID=48580
REFID=47432
REFID=47398
REFID=45782
REFID=45701
REFID=45076
REFID=44377
REFID=44440
REFID=44441
REFID=44621

REFID=44093
REFID=43659
REFID=44071
REFID=43657
REFID=43587
REFID=43596
REFID=43585
REFID=42177
REFID=42172
REFID=41587
REFID=41365
REFID=40921
REFID=21376
REFID=21377

$\rho(1570)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

May be an OZI-violating decay mode of $\rho(1700)$. See our mini-review under the $\rho(1700)$.

NODE=M188

NODE=M188

 $\rho(1570)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1570±36±62	54	¹ AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
1480±40		² BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$
¹ From the fit with two resonances.				
² Systematic errors not estimated.				

NODE=M188M

NODE=M188M

NODE=M188M;LINKAGE=AU
NODE=M188M;LINKAGE=BI **$\rho(1570)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
144±75±43	54	³ AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
130±60		⁴ BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$
³ From the fit with two resonances.				
⁴ Systematic errors not estimated.				

NODE=M188W

NODE=M188W

NODE=M188W;LINKAGE=AU
NODE=M188W;LINKAGE=BI **$\rho(1570)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	
Γ_2 $\phi\pi$	not seen
Γ_3 $\omega\pi$	

DESIG=1

DESIG=2

DESIG=3

NODE=M188215;NODE=M188

 $\rho(1570)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
3.5±0.9±0.3		54	⁵ AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$	
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
<70	90		⁶ AULCHENKO	87B ND	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$	
⁵ From the fit with two resonances.						
⁶ Using mass and width of BITYUKOV 87.						

NODE=M188225

NODE=M188G01
NODE=M188G01NODE=M188G01;LINKAGE=AU
NODE=M188G01;LINKAGE=AL **$\rho(1570)$ BRANCHING RATIOS**

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
not seen	ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$	
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
<0.01	⁷ DONNACHIE	91 RVUE		
⁷ Using data from BISELLO 91B, DOLINSKY 86, and ALBRECHT 87L.				

NODE=M188220

NODE=M188R01
NODE=M188R01

NODE=M188R01;LINKAGE=DO

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_3
>0.5	95	BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$	

NODE=M188R02
NODE=M188R02 **$\rho(1570)$ REFERENCES**

AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45765
BISELLO	91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)	REFID=41752
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=41632
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40418
AULCHENKO	87B	JETPL 45 145	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=41373
		Translated from ZETFP 45 118.			
BITYUKOV	87	PL B188 383	S.I. Bitjukov <i>et al.</i>	(SERP)	REFID=40011
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=20246

NODE=M188

$h_1(1595)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

OMITTED FROM SUMMARY TABLE

Seen in a partial-wave analysis of the $\omega\eta$ system produced in the reaction $\pi^- p \rightarrow \omega\eta n$ at 18 GeV/c.

NODE=M166

NODE=M166

 $h_1(1595)$ MASS

NODE=M166M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$1594 \pm 15^{+10}_{-60}$	EUGENIO	01	SPEC 18 $\pi^- p \rightarrow \omega\eta n$

NODE=M166M

 $h_1(1595)$ WIDTH

NODE=M166W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$384 \pm 60^{+70}_{-100}$	EUGENIO	01	SPEC 18 $\pi^- p \rightarrow \omega\eta n$

NODE=M166W

 $h_1(1595)$ DECAY MODES

NODE=M166215;NODE=M166

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega\eta$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

 $h_1(1595)$ REFERENCESEUGENIO 01 PL B497 190 P. Eugenio *et al.*

NODE=M166

REFID=48010

NODE=M164

 $\pi_1(1600)$

$$I^G(J^{PC}) = 1^-(1^{-+})$$

 $\pi_1(1600)$ MASS

NODE=M164M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1662^{+8}_{-9} OUR AVERAGE				

NODE=M164M

$1660 \pm 10^{+0}_{-64}$	420k	ALEKSEEV	10	COMP 190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$1664 \pm 8 \pm 10$	145k	¹ LU	05	B852 18 $\pi^- p \rightarrow \omega\pi^- \pi^0 p$
$1709 \pm 24 \pm 41$	69k	² KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$
$1597 \pm 10^{+45}_{-10}$		² IVANOV	01	B852 18 $\pi^- p \rightarrow \eta' \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1593 \pm 8^{+29}_{-47}$		^{2,3} ADAMS	98B	B852 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
--------------------------	--	----------------------	-----	---

¹ May be a different state: natural and unnatural parity exchanges.² Natural parity exchange.³ Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with 2.6 M events of $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data.

NODE=M164M;LINKAGE=LU

NODE=M164M;LINKAGE=A

NODE=M164M;LINKAGE=DZ

 $\pi_1(1600)$ WIDTH

NODE=M164W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
241 ± 40 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.

NODE=M164W


$269 \pm 21^{+42}_{-64}$	420k	ALEKSEEV	10	COMP 190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$185 \pm 25 \pm 28$	145k	⁴ LU	05	B852 18 $\pi^- p \rightarrow \omega\pi^- \pi^0 p$
$403 \pm 80 \pm 115$	69k	⁵ KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$
$340 \pm 40 \pm 50$		⁵ IVANOV	01	B852 18 $\pi^- p \rightarrow \eta' \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$168 \pm 20^{+150}_{-12}$		^{5,6} ADAMS	98B	B852 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
---------------------------	--	----------------------	-----	---

- ⁴ May be a different state: natural and unnatural parity exchanges.
⁵ Natural parity exchange.
⁶ Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with 2.6 M events of $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data.

NODE=M164W;LINKAGE=LU
 NODE=M164W;LINKAGE=A
 NODE=M164W;LINKAGE=DZ



$\pi_1(1600)$ DECAY MODES

NODE=M164215;NODE=M164

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi \pi \pi$	not seen
Γ_2 $\rho^0 \pi^-$	not seen
Γ_3 $f_2(1270) \pi^-$	not seen
Γ_4 $b_1(1235) \pi$	seen
Γ_5 $\eta'(958) \pi^-$	seen
Γ_6 $f_1(1285) \pi$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2
 DESIG=4
 DESIG=5
 DESIG=3
 DESIG=6;OUR EST;→ UNCHECKED ←

$\pi_1(1600)$ BRANCHING RATIOS

NODE=M164220

$\Gamma(\rho^0 \pi^-)/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	NOZAR	09	CLAS $\gamma p \rightarrow 2\pi^+ \pi^- n$	
not seen	⁷ DZIERBA	06	B852 $18 \pi^- p$	

NODE=M164R1
 NODE=M164R1

⁷ From the PWA analysis of 2.6 M $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data. Supersedes ADAMS 98B.

NODE=M164R1;LINKAGE=DZ

$\Gamma(f_2(1270) \pi^-)/\Gamma_{\text{total}}$				Γ_3/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	⁸ DZIERBA	06	B852 $18 \pi^- p$	

NODE=M164R3
 NODE=M164R3

⁸ From the PWA analysis of 2.6 M $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data. Supersedes CHUNG 02.

NODE=M164R3;LINKAGE=DZ

$\Gamma(b_1(1235) \pi)/\Gamma_{\text{total}}$				Γ_4/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	35280	⁹ BAKER	03	SPEC $\bar{p} p \rightarrow \omega \pi^+ \pi^- \pi^0$

NODE=M164R4
 NODE=M164R4

• • • We do not use the following data for averages, fits, limits, etc. • • •
 seen 145k LU 05 B852 $18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$

⁹ $B((b_1 \pi)_{D\text{-wave}})/B((b_1 \pi)_{S\text{-wave}}) = 0.3 \pm 0.1$.

NODE=M164R4;LINKAGE=RB

$\Gamma(\eta'(958) \pi^-)/\Gamma_{\text{total}}$				Γ_5/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	IVANOV	01	B852 $18 \pi^- p \rightarrow \eta' \pi^- p$	

NODE=M164R2
 NODE=M164R2

$\Gamma(f_1(1285)\pi)/\Gamma(\eta'(958)\pi^-)$

Γ_6/Γ_5

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
3.80±0.78	69k	¹⁰ KUHN	04 B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

¹⁰ Using $\eta'(958)\pi$ data from IVANOV 01.

NODE=M164R5
NODE=M164R5

NODE=M164R;LINKAGE=KU

$\pi_1(1600)$ REFERENCES

ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
NOZAR	09	PRL 102 102002	M. Nozar <i>et al.</i>	(JLab CLAS Collab.)
DZIERBA	06	PR D73 072001	A.R. Dzierba <i>et al.</i>	(BNL E852 Collab.)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)
ADAMS	98B	PRL 81 5760	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)

NODE=M164

REFID=53356
REFID=52758
REFID=51077
REFID=50459
REFID=49773
REFID=49414
REFID=48837
REFID=48317
REFID=46610

NODE=M161

$a_1(1640)$

$I^G(J^{PC}) = 1^-(1^{++})$

OMITTED FROM SUMMARY TABLE

Seen in the amplitude analysis of the $3\pi^0$ system produced in $\bar{p}p \rightarrow 4\pi^0$. Possibly seen in the study of the hadronic structure in decay $\tau \rightarrow 3\pi\nu_\tau$ (ABREU 98G and ASNER 00). Needs confirmation.


NODE=M161

$a_1(1640)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1647±22 OUR AVERAGE		Error includes scale factor of 1.4.		See the ideogram below.
1630±20	35280	¹ BAKER	03 SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$
1714± 9±36		CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
1640±12±30		BAKER	99 SPEC	1.94 $\bar{p}p \rightarrow 4\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1670±90		BELLINI	85 SPEC	40 $\pi^- A \rightarrow \pi^-\pi^+\pi^- A$

NODE=M161M

NODE=M161M



¹ Using the $a_1(1260)$ mass and width results of BOWLER 88.

NODE=M161M;LINKAGE=KB

$a_1(1640)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
254± 27 OUR AVERAGE		Error includes scale factor of 1.1.		
225± 30	35280	² BAKER	03 SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$
308± 37±62		CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
300± 22±40		BAKER	99 SPEC	1.94 $\bar{p}p \rightarrow 4\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
300±100		BELLINI	85 SPEC	40 $\pi^- A \rightarrow \pi^-\pi^+\pi^- A$

NODE=M161W

NODE=M161W

²Using the $a_1(1260)$ mass and width results of BOWLER 88.

NODE=M161W;LINKAGE=KB

$a_1(1640)$ DECAY MODES

NODE=M161215;NODE=M161

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi\pi$	seen
Γ_2 $f_2(1270)\pi$	seen
Γ_3 $\sigma\pi$	seen
Γ_4 $\rho\pi S\text{-wave}$	seen
Γ_5 $\rho\pi D\text{-wave}$	seen
Γ_6 $\omega\pi\pi$	seen
Γ_7 $f_1(1285)\pi$	seen
Γ_8 $a_1(1260)\eta$	not seen

DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=8

$a_1(1640)$ BRANCHING RATIOS

NODE=M161220

$\Gamma(f_2(1270)\pi)/\Gamma(\sigma\pi)$ Γ_2/Γ_3
 VALUE DOCUMENT ID TECN COMMENT

NODE=M161R1
 NODE=M161R1

••• We do not use the following data for averages, fits, limits, etc. •••

0.24±0.07 BAKER 99 SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$

$\Gamma(\rho\pi D\text{-wave})/\Gamma_{\text{total}}$ Γ_5/Γ
 VALUE DOCUMENT ID TECN COMMENT

NODE=M161R2
 NODE=M161R2

••• We do not use the following data for averages, fits, limits, etc. •••

seen CHUNG 02 B852 18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$

seen AMELIN 95B VES 36 $\pi^-A \rightarrow \pi^+\pi^-\pi^-A$

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_6/Γ
 VALUE EVTS DOCUMENT ID TECN COMMENT

NODE=M161R3
 NODE=M161R3

••• We do not use the following data for averages, fits, limits, etc. •••

seen 35280 ³BAKER 03 SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

$\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$ Γ_7/Γ
 VALUE DOCUMENT ID TECN COMMENT

NODE=M161R4
 NODE=M161R4

••• We do not use the following data for averages, fits, limits, etc. •••

not seen KUHN 04 B852 18 $\pi^-p \rightarrow \eta\pi^+\pi^-\pi^-p$

seen LEE 94 MPS2 18 $\pi^-p \rightarrow K^+\bar{K}^0\pi^-\pi^-p$

$\Gamma(a_1(1260)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ
 VALUE DOCUMENT ID TECN COMMENT

NODE=M161R5
 NODE=M161R5

not seen KUHN 04 B852 18 $\pi^-p \rightarrow \eta\pi^+\pi^-\pi^-p$

³Assuming the $\omega\rho$ mechanism for the $\omega\pi\pi$ state.

NODE=M161R;LINKAGE=KB

$a_1(1640)$ REFERENCES

NODE=M161

KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)
BOWLER	88	PL B209 99	M.G. Bowler	(OXF)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	

Translated from YAF 41 1223.

REFID=49773
 REFID=49414
 REFID=48837
 REFID=47339
 REFID=46888
 REFID=45909
 REFID=44433
 REFID=44092
 REFID=40578
 REFID=47490

$f_2(1640)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M117

OMITTED FROM SUMMARY TABLE

 $f_2(1640)$ MASS

NODE=M117M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1639± 6 OUR AVERAGE	Error includes scale factor of 1.2.		
1620±16	BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1647± 7	ADAMO	92	OBLX $\bar{n}p \rightarrow 3\pi^+2\pi^-$
1635± 7	ALDE	90	GAM2 $38\pi^-p \rightarrow \omega\omega n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1640± 5	AMSLER	06	CBAR $0.9\bar{p}p \rightarrow K^+K^-\pi^0$
1659± 6	VLADIMIRSK...06	SPEC	40 $\pi^-p \rightarrow K_S^0K_S^0n$
1643± 7	¹ ALDE	89B	GAM2 $38\pi^-p \rightarrow \omega\omega n$

NODE=M117M

¹Superseded by ALDE 90.

NODE=M117M;LINKAGE=BB

 $f_2(1640)$ WIDTH

NODE=M117W

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
99⁺⁶⁰₋₄₀ OUR AVERAGE		Error includes scale factor of 2.9.		
140 ⁺⁶⁰ ₋₂₀		BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
58±20		ADAMO	92	OBLX $\bar{n}p \rightarrow 3\pi^+2\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
44± 9		AMSLER	06	CBAR $0.9\bar{p}p \rightarrow K^+K^-\pi^0$
152±18		VLADIMIRSK...06	SPEC	40 $\pi^-p \rightarrow K_S^0K_S^0n$
< 70	90	ALDE	90	GAM2 $38\pi^-p \rightarrow \omega\omega n$

NODE=M117W

 $f_2(1640)$ DECAY MODES

NODE=M117215;NODE=M117

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega\omega$	seen
Γ_2 4π	seen
Γ_3 $K\bar{K}$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3 **$f_2(1640)$ BRANCHING RATIOS**

NODE=M117220

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
seen	AMSLER	06	CBAR $0.9\bar{p}p \rightarrow K^+K^-\pi^0$	

NODE=M117R2
NODE=M117R2 **$f_2(1640)$ REFERENCES**

NODE=M117

AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)	REFID=51136
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH) JP	REFID=44438
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)	REFID=42177
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=40935
ALDE	89B	PL B216 451	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+) IGJPC	REFID=40735

$\eta_2(1645)$

$$I^G(J^{PC}) = 0^+(2^-+)$$

NODE=M154

 $\eta_2(1645)$ MASS

NODE=M154M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1617± 5 OUR AVERAGE				
1613± 8	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
1617± 8	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
1620±20	BARBERIS	97B	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1645±14±15	ADOMEIT	96	CBAR 0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1645± 6±20	ANISOVICH	00E	SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M154M

 $\eta_2(1645)$ WIDTH

NODE=M154W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
181±11 OUR AVERAGE				
185±17	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
177±18	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
180±25	BARBERIS	97B	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
180 ⁺⁴⁰ ₋₂₁ ±25	ADOMEIT	96	CBAR 0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
200±25	ANISOVICH	00E	SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M154W

 $\eta_2(1645)$ DECAY MODES

NODE=M154215;NODE=M154

Mode	Fraction (Γ_i/Γ)
Γ_1 $a_2(1320)\pi$	seen
Γ_2 $K\bar{K}\pi$	seen
Γ_3 $K^*\bar{K}$	seen
Γ_4 $\eta\pi^+\pi^-$	seen
Γ_5 $a_0(980)\pi$	seen
Γ_6 $f_2(1270)\eta$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←

 $\eta_2(1645)$ BRANCHING RATIOS

NODE=M154220

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.07±0.03	¹ BARBERIS	97C	OMEG 450 $pp \rightarrow ppK\bar{K}\pi$	

NODE=M154R1
NODE=M154R1

¹ Using $2(\pi^+\pi^-)$ data from BARBERIS 97B.

NODE=M154R1;LINKAGE=A

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_5
13.1±2.3 OUR AVERAGE				
13.5±4.6	² ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$	
13.0±2.7	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$	

NODE=M154R3
NODE=M154R3

² Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M154R3;LINKAGE=AN

VALUE	DOCUMENT ID	COMMENT	Γ_6/Γ
not seen	BARBERIS	00B 450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$	

NODE=M154R4
NODE=M154R4

 $\eta_2(1645)$ REFERENCES

NODE=M154

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47958
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202

$\omega(1650)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M126

 $\omega(1650)$ MASS

NODE=M126M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1670± 30 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1680± 10	13.1k	¹ AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1667± 13± 6		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1645± 8	13	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
1660± 10± 2		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1770± 50± 60	1.2M	² ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1619± 5		³ HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1700± 20		EUGENIO	01 SPEC	18 $\pi^-p \rightarrow \omega\eta n$
1705± 26	612	⁴ AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1820 ⁺¹⁹⁰ ₋₁₅₀		⁵ ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1840 ⁺¹⁰⁰ ₋₇₀		⁶ ACHASOV	98H RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1780 ⁺¹⁷⁰ ₋₃₀₀		⁷ ACHASOV	98H RVUE	$e^+e^- \rightarrow K^+K^-$
~ 2100		⁸ ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606± 9		⁹ CLEGG	94 RVUE	
1662± 13	750	¹⁰ ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1670± 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
1657± 13		CORDIER	81 DM1	$e^+e^- \rightarrow \omega 2\pi$
1679± 34	21	ESPOSITO	80 FRAM	$e^+e^- \rightarrow 3\pi$
1652± 17		COSME	79 OSPK	$e^+e^- \rightarrow 3\pi$

NODE=M126M

NODE=M126M
→ UNCHECKED ←

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=4

¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.³ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.⁴ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.⁵ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.⁶ Using the data from ANTONELLI 92.⁷ Using the data from IVANOV 81 and BISELLO 88B.⁸ Using the data from BISELLO 91C.⁹ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.¹⁰ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.NODE=M126M;LINKAGE=A
NODE=M126M;LINKAGE=VH

NODE=M126M;LINKAGE=AB

NODE=M126M;LINKAGE=KI

NODE=M126M;LINKAGE=L1
NODE=M126M;LINKAGE=L2
NODE=M126M;LINKAGE=L3
NODE=M126M;LINKAGE=L4
NODE=M126M;LINKAGE=AD

NODE=M126M;LINKAGE=AE

 $\omega(1650)$ WIDTH

NODE=M126W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
315± 35 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
310± 30	13.1k	¹¹ AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
222± 25± 20		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
114± 14	13	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
230± 30± 20		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
490 ⁺²⁰⁰ ₋₁₅₀ ± 130	1.2M	¹² ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
250± 14		¹³ HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
250± 50		EUGENIO	01 SPEC	18 $\pi^-p \rightarrow \omega\eta n$
370± 25	612	¹⁴ AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
113± 20		¹⁵ CLEGG	94 RVUE	
280± 24	750	¹⁶ ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
160± 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
136± 46		CORDIER	81 DM1	$e^+e^- \rightarrow \omega 2\pi$
99± 49	21	ESPOSITO	80 FRAM	$e^+e^- \rightarrow 3\pi$
42± 17		COSME	79 OSPK	$e^+e^- \rightarrow 3\pi$

NODE=M126W

NODE=M126W
→ UNCHECKED ←

OCCUR=2

OCCUR=5

OCCUR=4

- 11 From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.
 12 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
 13 Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.
 14 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.
 15 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
 16 From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

NODE=M126W;LINKAGE=A
 NODE=M126W;LINKAGE=VH

NODE=M126W;LINKAGE=AB

NODE=M126W;LINKAGE=KI

NODE=M126W;LINKAGE=AD

NODE=M126W;LINKAGE=AE

NODE=M126215;NODE=M126

$\omega(1650)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\omega\pi\pi$	seen
Γ_3 $\omega\eta$	seen
Γ_4 e^+e^-	seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

$\omega(1650)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M126230

$\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_4/\Gamma$

NODE=M126G3
 NODE=M126G3

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.56 ± 0.23	13.1k	17 AULCHENKO	15A SND	$1.05\text{--}1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$1.3 \pm 0.1 \pm 0.1$		AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
$1.2^{+0.4}_{-0.1} \pm 0.8$	1.2M	18,19 ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.921 ± 0.230		20,21 CLEGG	94 RVUE	
0.479 ± 0.050	750	22,23 ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_4/\Gamma$

NODE=M126G4
 NODE=M126G4

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
7.0 ± 0.5		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
$4.1 \pm 0.9 \pm 1.3$	1.2M	18,19 ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 ± 0.95		24 AKHMETSHIN 00D	CMD2	$1.2\text{--}1.38 e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 ± 0.80		20,21 CLEGG	94 RVUE	
6.07 ± 0.61	750	22,23 ANTONELLI	92 DM2	$1.34\text{--}2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_4/\Gamma$

NODE=M126G5
 NODE=M126G5

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.57 ± 0.06		13	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
<6		90	25 AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$

- 17 From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.
 18 Calculated by us from the cross section at the peak.
 19 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
 20 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
 21 From the partial and leptonic width given by the authors.
 22 From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.
 23 From the product of the leptonic width and partial branching ratio given by the authors.
 24 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.
 25 $\omega(1650)$ mass and width fixed at 1700 MeV and 250 MeV, respectively.

NODE=M126G3;LINKAGE=A
 NODE=M126G;LINKAGE=AW
 NODE=M126G;LINKAGE=VH

NODE=M126G;LINKAGE=AD

NODE=M126G;LINKAGE=SE

NODE=M126G;LINKAGE=AE

NODE=M126G;LINKAGE=ES

NODE=M126G;LINKAGE=KL

NODE=M126G5;LINKAGE=KH

$\omega(1650)$ BRANCHING RATIOS

NODE=M126225

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
~ 0.35	1.2M	26 ACHASOV	03D	RVUE	$0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.620±0.014		27 HENNER	02	RVUE	$1.2-2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$

NODE=M126R2
NODE=M126R2

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
~ 0.65	1.2M	26 ACHASOV	03D	RVUE	$0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.380±0.014		27 HENNER	02	RVUE	$1.2-2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$

NODE=M126R3
NODE=M126R3

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
~ 18	1.2M	27,28 ACHASOV	03D	RVUE	$0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
32±1		27 HENNER	02	RVUE	$1.2-2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$

NODE=M126R4
NODE=M126R4

²⁶ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega\pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

NODE=M126R;LINKAGE=VH

²⁷ Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

NODE=M126R;LINKAGE=AC

²⁸ Calculated by us from the cross section at the peak.

NODE=M126R;LINKAGE=AW

 $\omega(1650)$ REFERENCES

NODE=M126

AULCHENKO	15A	JETP 121 27 Translated from ZETF 148 34.	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56843
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>		REFID=49177
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>		REFID=48010
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		REFID=46323
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41867
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40581
BARKOV	87	JETPL 46 164 Translated from ZETFP 46 132.	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21502
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)	REFID=21586
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
ESPOSITO	80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=21584
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)	REFID=21475

NODE=M045

 $\omega_3(1670)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

 $\omega_3(1670)$ MASS

NODE=M045M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1667 ± 4 OUR AVERAGE				
1665.3 ± 5.2 ± 4.5	23400	AMELIN	96 VES	$36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1685 ± 20	60	BAUBILLIER	79 HBC	$8.2 K^- p$ backward
1673 ± 12	430	^{1,2} BALTAY	78E HBC	$15 \pi^+ p \rightarrow \Delta 3\pi$
1650 ± 12		CORDEN	78B OMEG	$8-12 \pi^- p \rightarrow N 3\pi$
1669 ± 11	600	² WAGNER	75 HBC	$7 \pi^+ p \rightarrow \Delta^{++} 3\pi$
1678 ± 14	500	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 3\pi^0$
1660 ± 13	200	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p \omega \pi^0 \pi^0$
1679 ± 17	200	MATTHEWS	71D DBC	$7.0 \pi^+ n \rightarrow p 3\pi^0$
1670 ± 20		KENYON	69 DBC	$8 \pi^+ n \rightarrow p 3\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 1700	110	¹ CERRADA	77B HBC	$4.2 K^- p \rightarrow \Lambda 3\pi$
1695 ± 20		BARNES	69B HBC	$4.6 K^- p \rightarrow \omega 2\pi X$
1636 ± 20		ARMENISE	68B DBC	$5.1 \pi^+ n \rightarrow p 3\pi^0$

NODE=M045M

OCCUR=2

¹ Phase rotation seen for $J^P = 3^- \rho\pi$ wave.² From a fit to $I(J^P) = 0(3^-) \rho\pi$ partial wave.NODE=M045M;LINKAGE=E
NODE=M045M;LINKAGE=P $\omega_3(1670)$ WIDTH

NODE=M045W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
168 ± 10 OUR AVERAGE				
149 ± 19 ± 7	23400	AMELIN	96 VES	$36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
160 ± 80	60	³ BAUBILLIER	79 HBC	$8.2 K^- p$ backward
173 ± 16	430	^{4,5} BALTAY	78E HBC	$15 \pi^+ p \rightarrow \Delta 3\pi$
253 ± 39		CORDEN	78B OMEG	$8-12 \pi^- p \rightarrow N 3\pi$
173 ± 28	600	^{3,5} WAGNER	75 HBC	$7 \pi^+ p \rightarrow \Delta^{++} 3\pi$
167 ± 40	500	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 3\pi^0$
122 ± 39	200	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p \omega \pi^0 \pi^0$
155 ± 40	200	³ MATTHEWS	71D DBC	$7.0 \pi^+ n \rightarrow p 3\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
90 ± 20		BARNES	69B HBC	$4.6 K^- p \rightarrow \omega 2\pi$
100 ± 40		KENYON	69 DBC	$8 \pi^+ n \rightarrow p 3\pi^0$
112 ± 60		ARMENISE	68B DBC	$5.1 \pi^+ n \rightarrow p 3\pi^0$

NODE=M045W

OCCUR=2

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.⁴ Phase rotation seen for $J^P = 3^- \rho\pi$ wave.⁵ From a fit to $I(J^P) = 0(3^-) \rho\pi$ partial wave.NODE=M045W;LINKAGE=S
NODE=M045W;LINKAGE=E
NODE=M045W;LINKAGE=P $\omega_3(1670)$ DECAY MODES

NODE=M045215;NODE=M045

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \rho\pi$	seen
$\Gamma_2 \quad \omega\pi\pi$	seen
$\Gamma_3 \quad b_1(1235)\pi$	possibly seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

 $\omega_3(1670)$ BRANCHING RATIOS

NODE=M045220

$\Gamma(\omega\pi\pi)/\Gamma(\rho\pi)$					Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.71 ± 0.27	100	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 5\pi^0$	
$\Gamma(b_1(1235)\pi)/\Gamma(\rho\pi)$					Γ_3/Γ_1
VALUE		DOCUMENT ID	TECN	COMMENT	
possibly seen		DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 5\pi^0$	

NODE=M045R3
NODE=M045R3NODE=M045R4
NODE=M045R4

$\Gamma(b_1(1235)\pi)/\Gamma(\omega\pi\pi)$ Γ_3/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.75	68	BAUBILLIER	79	HBC	8.2 $K^- p$ backward
-------	----	------------	----	-----	----------------------

NODE=M045R5
NODE=M045R5

 $\omega_3(1670)$ REFERENCES

AMELIN	96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BAUBILLIER	79	PL 89B 131	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
BALTAY	78E	PRL 40 87	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
CORDEN	78B	NP B138 235	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+) JP
WAGNER	75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP
DIAZ	74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)
MATTHEWS	71D	PR D3 2561	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC)
BARNES	69B	PRL 23 142	V.E. Barnes <i>et al.</i>	(BNL)
KENYON	69	PRL 23 146	I.R. Kenyon <i>et al.</i>	(BNL, UCND, ORNL)
ARMENISE	68B	PL 26B 336	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)

NODE=M045

REFID=44649
REFID=21522
REFID=21520
REFID=21269
REFID=20537
REFID=20843
REFID=21248
REFID=21515
REFID=21512
REFID=20800
REFID=20783

NODE=M034

 $\pi_2(1670)$

$$I^G(J^{PC}) = 1^-(2^-+)$$

 $\pi_2(1670)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

1672.2 ± 3.0 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

1658 ± 3	$^{+24}_{-8}$	420k	ALEKSEEV	10	COMP	190 $\pi^- p b \rightarrow \pi^- \pi^- \pi^+ p b'$
1749 ± 10	±100	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
1676 ± 3	± 8		1 CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1685 ± 10	± 30		2 BARBERIS	01		450 $p p \rightarrow p_f 3\pi^0 p_s$
1687 ± 9	± 15		AMELIN	99	YES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
1669 ± 4			BARBERIS	98B		450 $p p \rightarrow p_f \rho \pi p_s$
1670 ± 4			BARBERIS	98B		450 $p p \rightarrow p_f f_2(1270) \pi p_s$
1730 ± 20			3 AMELIN	95B	YES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
1690 ± 14			4 BERDNIKOV	94	YES	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
1710 ± 20	700		ANTIPOV	87	SIGM -	50 $\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
1676 ± 6			4 EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow 3\pi p$
1657 ± 14			4,5 DAUM	80D	SPEC -	63-94 $\pi p \rightarrow 3\pi X$
1662 ± 10	2000		4 BALTAY	77	HBC +	15 $\pi^+ p \rightarrow p 3\pi$

NODE=M034M

NODE=M034M

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1742 ± 31	± 49		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
1624 ± 21			1 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1622 ± 35			6 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1693 ± 28			7 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1710 ± 20			8 DAUM	81B	SPEC -	63,94 $\pi^- p$
1660 ± 10			4 ASCOLI	73	HBC -	5-25 $\pi^- p \rightarrow p \pi_2$

OCCUR=2

OCCUR=3

¹ From $f_2(1270)\pi$ decay.

² From a fit to the invariant mass distribution.

³ From a fit to $J^{PC} = 2^-+ f_2(1270)\pi, f_0(1370)\pi$ waves.

⁴ From a fit to $J^P = 2^- S$ -wave $f_2(1270)\pi$ partial wave.

⁵ Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.


⁶ From $\rho\pi$ decay.

⁷ From $\sigma\pi$ decay.

⁸ From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.

NODE=M034M;LINKAGE=F2
NODE=M034M;LINKAGE=BR
NODE=M034M;LINKAGE=AX
NODE=M034M;LINKAGE=P
NODE=M034M;LINKAGE=D

NODE=M034M;LINKAGE=R2
NODE=M034M;LINKAGE=S2
NODE=M034M;LINKAGE=L



$\pi_2(1670)$ WIDTH

NODE=M034W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
260± 9	OUR AVERAGE	Error includes scale factor of 1.2.			
271± 9 ⁺ ₂₄	420k	ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
408± 60±250	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
254± 3± 31		9 CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
265± 30± 40		10 BARBERIS	01		450 $pp \rightarrow p_f 3\pi^0 p_s$
168± 43± 53		AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
268± 15		BARBERIS	98B		450 $pp \rightarrow p_f \rho \pi p_s$
256± 15		BARBERIS	98B		450 $pp \rightarrow p_f f_2(1270) \pi p_s$
310± 20		11 AMELIN	95B	VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
190± 50		12 BERDNIKOV	94	VES	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
170± 80	700	ANTIPOV	87	SIGM	- 50 $\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
260± 20		12 EVANGELIS...	81	OMEG	- 12 $\pi^- p \rightarrow 3\pi p$
219± 20		12,13 DAUM	80D	SPEC	- 63-94 $\pi p \rightarrow 3\pi X$
285± 60	2000	12 BALTAY	77	HBC	+ 15 $\pi^+ p \rightarrow p 3\pi$
236± 49± 36		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
304± 22		9 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
404±108		14 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
330± 90		15 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
312± 50		16 DAUM	81B	SPEC	- 63,94 $\pi^- p$
270± 60		12 ASCOLI	73	HBC	- 5-25 $\pi^- p \rightarrow p \pi_2$

NODE=M034W

OCCUR=2

OCCUR=2

OCCUR=3

9 From $f_2(1270)\pi$ decay.
 10 From a fit to the invariant mass distribution.
 11 From a fit to $J^{PC} = 2^- + f_2(1270)\pi, f_0(1370)\pi$ waves.
 12 From a fit to $J^P = 2^- f_2(1270)\pi$ partial wave.
 13 Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.
 14 From $\rho\pi$ decay.
 15 From $\sigma\pi$ decay.
 16 From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.

NODE=M034W;LINKAGE=F2
 NODE=M034W;LINKAGE=BR
 NODE=M034W;LINKAGE=AX
 NODE=M034W;LINKAGE=P
 NODE=M034W;LINKAGE=D

NODE=M034W;LINKAGE=R2
 NODE=M034W;LINKAGE=S2
 NODE=M034W;LINKAGE=L

$\pi_2(1670)$ DECAY MODES

NODE=M034215;NODE=M034

Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1 3π	(95.8±1.4) %		DESIG=20
Γ_2 $\pi^+\pi^-\pi^0$			DESIG=22
Γ_3 $\pi^0\pi^0\pi^0$			DESIG=23
Γ_4 $f_2(1270)\pi$	(56.3±3.2) %		DESIG=8
Γ_5 $\rho\pi$	(31 ±4) %		DESIG=2
Γ_6 $\sigma\pi$	(10.9±3.4) %		DESIG=13
Γ_7 $\pi(\pi\pi)_{S\text{-wave}}$	(8.7±3.4) %		DESIG=11
Γ_8 $K\bar{K}^*(892)+\text{c.c.}$	(4.2±1.4) %		DESIG=5
Γ_9 $\omega\rho$	(2.7±1.1) %		DESIG=14
Γ_{10} $\pi^\pm\gamma$	(7.0±1.1) $\times 10^{-4}$		DESIG=27
Γ_{11} $\gamma\gamma$	< 2.8 $\times 10^{-7}$	90%	DESIG=12
Γ_{12} $\eta\pi$			DESIG=3
Γ_{13} $\pi^\pm 2\pi^+ 2\pi^-$			DESIG=4
Γ_{14} $\rho(1450)\pi$	< 3.6 $\times 10^{-3}$	97.7%	DESIG=15
Γ_{15} $b_1(1235)\pi$	< 1.9 $\times 10^{-3}$	97.7%	DESIG=16
Γ_{16} $\eta 3\pi$			DESIG=24
Γ_{17} $f_1(1285)\pi$	possibly seen		DESIG=25
Γ_{18} $a_2(1320)\pi$	not seen		DESIG=26

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.9$ for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_5	-53		
x_7	-29	-59	
x_8	-8	-21	-9
	x_4	x_5	x_7

 $\pi_2(1670)$ PARTIAL WIDTHS

NODE=M034217

 $\Gamma(\pi^\pm\gamma)$ Γ_{10}

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
181±11±27	17 ADOLPH	14	COMP	- 190 $\pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$

NODE=M034W2
NODE=M034W2

¹⁷ Primakoff reaction. Assumes incoherent $f_2(1270)\pi$ contribution to 3π final state and uses $B(\pi_2(1670) \rightarrow f_2\pi) = 56\%$.

NODE=M034W2;LINKAGE=AAD

 $\Gamma(\gamma\gamma)$ Γ_{11}

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.072	90	18 ACCIARRI	97T	L3	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$

NODE=M034W1
NODE=M034W1

••• We do not use the following data for averages, fits, limits, etc. •••

<0.19	90	18 ALBRECHT	97B	ARG	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.41 ±0.23±0.28		ANTREASYAN 90	CBAL	0	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0\pi^0$
0.8 ±0.3 ±0.12		19 BEHREND	90C	CELL	0
1.3 ±0.3 ±0.2		20 BEHREND	90C	CELL	0

OCCUR=2

¹⁸ Decaying into $f_2(1270)\pi$ and $\rho\pi$.

¹⁹ Constructive interference between $f_2(1270)\pi, \rho\pi$ and background.

²⁰ Incoherent Ansatz.

NODE=M034W1;LINKAGE=QQ
NODE=M034W1;LINKAGE=C
NODE=M034W1;LINKAGE=G

$\pi_2(1670) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M034230

$$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_{11}/\Gamma$$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	95	²¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

NODE=M034G01
NODE=M034G01²¹ From analysis of L3 data at 183–209 GeV.

NODE=M034G01;LINKAGE=SC

 $\pi_2(1670) \text{ BRANCHING RATIOS}$

NODE=M034220

$$\Gamma(3\pi)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma = (\Gamma_4+\Gamma_5+\Gamma_7)/\Gamma$$

VALUE	DOCUMENT ID
0.958±0.014 OUR FIT	

NODE=M034R20
NODE=M034R20

$$\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_3/\Gamma_2$$

VALUE	DOCUMENT ID	COMMENT
0.29±0.03±0.05	²² BARBERIS 01	450 $p p \rightarrow p_f 3\pi^0 p_s$

NODE=M034R21
NODE=M034R21

$$\Gamma(\rho\pi)/0.565\Gamma(f_2(1270)\pi) \quad \Gamma_5/0.565\Gamma_4$$

(With $f_2(1270) \rightarrow \pi^+\pi^-$.)

VALUE	DOCUMENT ID	TECN	COMMENT
0.97±0.09 OUR AVERAGE			Error includes scale factor of 1.9.

NODE=M034R16
NODE=M034R16
NODE=M034R16

0.76±0.07±0.10	CHUNG 02	B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
1.01±0.05	BARBERIS 98B		450 $p p \rightarrow p_f \pi^+\pi^-\pi^0 p_s$

$$\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi) \quad \Gamma_6/\Gamma_4$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.19±0.06 OUR AVERAGE			

NODE=M034R15
NODE=M034R15

0.17±0.02±0.07	CHUNG 02	B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
0.24±0.10	^{23,24} BAKER 99	SPEC	1.94 $\bar{p} p \rightarrow 4\pi^0$

$$\frac{1}{2}\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-) \quad \frac{1}{2}\Gamma_5/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.29±0.04 OUR FIT				

NODE=M034R2
NODE=M034R2

0.29±0.05	²⁵ DAUM 81B	SPEC		63,94 $\pi^- p$
------------------	------------------------	------	--	-----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.3	BARTSCH 68	HBC	+	8 $\pi^+ p \rightarrow 3\pi p$
------	------------	-----	---	--------------------------------

$$0.565\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-) \quad 0.565\Gamma_4/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$$

(With $f_2(1270) \rightarrow \pi^+\pi^-$.)

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.604±0.035 OUR FIT				

NODE=M034R3
NODE=M034R3
NODE=M034R3

0.60 ±0.05 OUR AVERAGE				Error includes scale factor of 1.3.
-------------------------------	--	--	--	-------------------------------------

0.61 ±0.04	²⁵ DAUM 81B	SPEC		63,94 $\pi^- p$
------------	------------------------	------	--	-----------------

0.76 $\begin{smallmatrix} +0.24 \\ -0.34 \end{smallmatrix}$	ARMENISE 69	DBC	+	5.1 $\pi^+ d \rightarrow d3\pi$
---	-------------	-----	---	---------------------------------

0.35 ±0.20	BALTAY 68	HBC	+	7–8.5 $\pi^+ p$
------------	-----------	-----	---	-----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.59	BARTSCH 68	HBC	+	8 $\pi^+ p \rightarrow 3\pi p$
------	------------	-----	---	--------------------------------

$$0.624\Gamma(\pi(\pi\pi)_{S\text{-wave}})/\Gamma(\pi^\pm\pi^+\pi^-) \quad 0.624\Gamma_7/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$$

(With $(\pi\pi)_{S\text{-wave}} \rightarrow \pi^+\pi^-$.)

VALUE	DOCUMENT ID	TECN	COMMENT
0.10±0.04 OUR FIT			

NODE=M034R11
NODE=M034R11
NODE=M034R11

0.10±0.05	²⁵ DAUM 81B	SPEC		63,94 $\pi^- p$
------------------	------------------------	------	--	-----------------

$$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(f_2(1270)\pi) \quad \Gamma_8/\Gamma_4$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.075±0.025 OUR FIT				

NODE=M034R13
NODE=M034R13

0.075±0.025	²⁶ ARMSTRONG 82B	OMEG	–	16 $\pi^- p \rightarrow K^+ K^- \pi^- p$
--------------------	-----------------------------	------	---	--

$$\Gamma(\omega\rho)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.027±0.004±0.010	²⁷ AMELIN 99	VES	37 $\pi^- A \rightarrow \omega\pi^-\pi^0 A^*$

NODE=M034R17
NODE=M034R17

$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$ (All η decays.) $\Gamma_{12}/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.09	BALTAY	68	HBC +	7-8.5 π^+p
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.10	CRENNELL	70	HBC -	6 $\pi^-p \rightarrow f_2\pi^-N$

NODE=M034R5
 NODE=M034R5
 NODE=M034R5

 $\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$ $\Gamma_{13}/(0.565\Gamma_4+\frac{1}{2}\Gamma_5+0.624\Gamma_7)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.10	CRENNELL	70	HBC -	6 $\pi^-p \rightarrow f_2\pi^-N$
<0.1	BALTAY	68	HBC +	7,8.5 π^+p

NODE=M034R6
 NODE=M034R6;CHECK LIMITS

 $\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0036	97.7	AMELIN	99	37 $\pi^-A \rightarrow \omega\pi^-\pi^0A^*$

NODE=M034R18
 NODE=M034R18

 $\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0019	97.7	AMELIN	99	37 $\pi^-A \rightarrow \omega\pi^-\pi^0A^*$

NODE=M034R19
 NODE=M034R19

 $\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
possibly seen	69k	KUHN	04	B852 18 $\pi^-p \rightarrow \eta\pi^+\pi^-\pi^-p$

NODE=M034R23
 NODE=M034R23

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
not seen	69k	KUHN	04	B852 18 $\pi^-p \rightarrow \eta\pi^+\pi^-\pi^-p$

NODE=M034R24
 NODE=M034R24

D-wave/S-wave RATIO FOR $\pi_2(1670) \rightarrow f_2(1270)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.18±0.06	²³ BAKER	99	SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$
0.22±0.10	²⁵ DAUM	81B	SPEC 63,94 π^-p

NODE=M034R14
 NODE=M034R14

F-wave/P-wave RATIO FOR $\pi_2(1670) \rightarrow \rho\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.72±0.07±0.14	CHUNG	02	B852 18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$

NODE=M034R22
 NODE=M034R22

²² Using BARBERIS 98B.

²³ Using preliminary CBAR data.

²⁴ With the $\sigma\pi$ in $L=2$ and the $f_2(1270)\pi$ in $L=0$.

²⁵ From a two-resonance fit to four 2^-0^+ waves.

²⁶ From a partial-wave analysis of $K^+K^-\pi^-$ system.

²⁷ Normalized to the $B(\pi_2(1670) \rightarrow f_2\pi)$.

NODE=M034R;LINKAGE=RB
 NODE=M034R;LINKAGE=BK
 NODE=M034R15;LINKAGE=A
 NODE=M034R;LINKAGE=L
 NODE=M034R13;LINKAGE=M
 NODE=M034R;LINKAGE=DM

 $\pi_2(1670)$ REFERENCES

ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>	
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)
ANTREASYAN	90	ZPHY C48 561	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
ANTIPOV	87	EPL 4 403	Y.M. Antipov <i>et al.</i>	(SERP, JINR, INRM+)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
ARMSTRONG	82B	NP B202 1	T.A. Armstrong, B. Baccari	(AACH3, BARI, BONN+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
Also		NP B186 594	C. Evangelista	
DAUM	80D	PL 89B 285	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
ASCOLI	73	PR D7 669	G. Ascoli	(ILL, TINTO, GENO, HAMB, MILA+) JP
CRENNELL	70	PRL 24 781	D.J. Crennell <i>et al.</i>	(BNL)
ARMENISE	69	LCN 2 501	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I
BARTSCH	68	NP B7 345	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN) JP

NODE=M034

REFID=55911
 REFID=53356
 REFID=51186
 REFID=50459
 REFID=49773
 REFID=48837
 REFID=48324
 REFID=46910
 REFID=46888
 REFID=46345
 REFID=45761
 REFID=45418
 REFID=44433
 REFID=44073
 REFID=41372
 REFID=41356
 REFID=40004
 REFID=47490
 REFID=20874
 REFID=20872
 REFID=20462
 REFID=21576
 REFID=21573
 REFID=20847
 REFID=21553
 REFID=20805
 REFID=20689
 REFID=21531
 REFID=21532

$\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M067

 $\phi(1680)$ MASS

NODE=M067205

 e^+e^- PRODUCTION

NODE=M067M1

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M067M1

1680±20 OUR ESTIMATE

→ UNCHECKED ←

●●● We do not use the following data for averages, fits, limits, etc. ●●●

1689±7±10	4.8k	¹ SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^- \pi^+ \pi^- \gamma$
1709±20±43		² AUBERT	08s BABR	10.6 $e^+e^- \rightarrow$ hadrons
1623±20	948	³ AKHMETSHIN	03 CMD2	1.05-1.38 $e^+e^- \rightarrow K_L^0 K_S^0$
~1500		⁴ ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0, \omega \pi^+ \pi^-,$ $K^+ K^-$
~1900		⁵ ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1700±20		⁶ CLEGG	94 RVUE	$e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$
1657±27	367	⁷ BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655±17		⁷ BISELLO	88B DM2	$e^+e^- \rightarrow K^+ K^-$
1680±10		⁸ BUON	82 DM1	$e^+e^- \rightarrow$ hadrons
1677±12		⁹ MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K \pi$

OCCUR=4

¹ From a fit with two incoherent Breit-Wigners.

NODE=M067M1;LINKAGE=SH

² From the simultaneous fit to the $K\bar{K}^*(892)+$ c.c. and $\phi\eta$ data from AUBERT 08s using the results of AUBERT 07AK.

NODE=M067M1;LINKAGE=AU

³ From the combined fit of AKHMETSHIN 03 and MANE 81 also including $\rho, \omega,$ and ϕ . Neither isospin nor flavor structure known.

NODE=M067M;LINKAGE=HK

⁴ Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.

NODE=M067M1;LINKAGE=L1

⁵ Using the data from BISELLO 91C.

NODE=M067M1;LINKAGE=L4

⁶ Using BISELLO 88B and MANE 82 data.

NODE=M067M;LINKAGE=A

⁷ From global fit including ρ, ω, ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation.

NODE=M067M;LINKAGE=E

⁸ From global fit of ρ, ω, ϕ and their radial excitations to channels $\omega \pi^+ \pi^-, K^+ K^-, K_S^0 K_L^0, K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

NODE=M067M;LINKAGE=C

⁹ Fit to one channel only, neglecting interference with $\omega, \rho(1700)$.

NODE=M067M;LINKAGE=D

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M067M2

NODE=M067M2

●●● We do not use the following data for averages, fits, limits, etc. ●●●

1753±3	¹⁰ LINK	02K FOCS	20-160 $\gamma p \rightarrow K^+ K^- p$
1726±22	¹⁰ BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$
1760±20	¹⁰ ATKINSON	85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$
1690±10	¹⁰ ASTON	81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$

¹⁰ We list here a state decaying into $K^+ K^-$ possibly different from $\phi(1680)$.

NODE=M067M2;LINKAGE=LK

 $p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M067M3

NODE=M067M3

●●● We do not use the following data for averages, fits, limits, etc. ●●●

1700±8	¹¹ AMSLER	06 CBAR	0.9 $p\bar{p} \rightarrow K^+ K^- \pi^0$
--------	----------------------	---------	--

¹¹ Could also be $\rho(1700)$.

NODE=M067M3;LINKAGE=AM

 $\phi(1680)$ WIDTH

NODE=M067210

 e^+e^- PRODUCTION

NODE=M067W1

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M067W1

150±50 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

→ UNCHECKED ←

●●● We do not use the following data for averages, fits, limits, etc. ●●●

211±14±19	4.8k	¹² SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
322±77±160		¹³ AUBERT	08s BABR	10.6 $e^+e^- \rightarrow$ hadrons
139±60	948	¹⁴ AKHMETSHIN	03 CMD2	1.05-1.38 $e^+e^- \rightarrow K_L^0 K_S^0$
300±60		¹⁵ CLEGG	94 RVUE	$e^+e^- \rightarrow K^+ K^-, K_S^0 K \pi$
146±55	367	⁷ BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
207±45		¹⁶ BISELLO	88B DM2	$e^+e^- \rightarrow K^+ K^-$
185±22		¹⁷ BUON	82 DM1	$e^+e^- \rightarrow$ hadrons
102±36		¹⁸ MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K \pi$

- ¹² From a fit with two incoherent Breit-Wigners.
¹³ From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.
¹⁴ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known.
¹⁵ Using BISELLO 88B and MANE 82 data.
¹⁶ From global fit including ρ , ω , ϕ and $\rho(1700)$
¹⁷ From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0K_L^0$, $K_S^0K^\pm\pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.
¹⁸ Fit to one channel only, neglecting interference with ω , $\rho(1700)$.

NODE=M067W1;LINKAGE=SH
 NODE=M067W1;LINKAGE=AU

NODE=M067W;LINKAGE=HK

NODE=M067W;LINKAGE=A
 NODE=M067W;LINKAGE=E
 NODE=M067W;LINKAGE=C

NODE=M067W;LINKAGE=D

NODE=M067W2
 NODE=M067W2

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

122±63	¹⁹ LINK	02K	FOCS 20–160 $\gamma p \rightarrow K^+K^-\rho$
121±47	¹⁹ BUSENITZ	89	TPS $\gamma p \rightarrow K^+K^-X$
80±40	¹⁹ ATKINSON	85C	OMEG 20–70 $\gamma p \rightarrow K\bar{K}X$
100±40	¹⁹ ASTON	81F	OMEG 25–70 $\gamma p \rightarrow K^+K^-X$

¹⁹ We list here a state decaying into K^+K^- possibly different from $\phi(1680)$.

NODE=M067W2;LINKAGE=LK

$p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

143±24	²⁰ AMSLER	06	CBAR 0.9 $p\bar{p} \rightarrow K^+K^-\pi^0$
--------	----------------------	----	---

²⁰ Could also be $\rho(1700)$.

NODE=M067W3
 NODE=M067W3

NODE=M067W3;LINKAGE=AM

$\phi(1680)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}^*(892) + \text{c.c.}$	dominant
Γ_2 $K_S^0K\pi$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $K_L^0K_S^0$	
Γ_5 e^+e^-	seen
Γ_6 $\omega\pi\pi$	not seen
Γ_7 $\phi\pi\pi$	
Γ_8 $K^+K^-\pi^+\pi^-$	seen
Γ_9 $\eta\phi$	seen
Γ_{10} $\eta\gamma$	seen
Γ_{11} $K^+K^-\pi^0$	

DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=9
 DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=11
 DESIG=12;OUR EVAL;→ UNCHECKED ←
 DESIG=10
 DESIG=13
 DESIG=2

NODE=M067215;NODE=M067

$\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

This combination of a branching ratio into channel (i) and branching ratio into e^+e^- is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into (i) or e^+e^- .

NODE=M067223

NODE=M067223

$\Gamma(K_L^0K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.131±0.059	948	²¹ AKHMETSHIN 03	CMD2	1.05–1.38 $e^+e^- \rightarrow K_L^0K_S^0$
-------------	-----	-----------------------------	------	---

²¹ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known. Recalculated by us.

NODE=M067G5
 NODE=M067G5

NODE=M067G;LINKAGE=GK

$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.15±0.16±0.01	²² AUBERT	08S	BABR	10.6 $e^+e^- \rightarrow K\bar{K}^*(892)\gamma + \text{c.c.}$
3.29±1.57	²³ BISELLO	91C	DM2	1.35–2.40 $e^+e^- \rightarrow K_S^0K^\pm\pi^\mp$

²² From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

²³ Recalculated by us with the published value of $B(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)$.

NODE=M067G6
 NODE=M067G6

NODE=M067G6;LINKAGE=AU

NODE=M067G;LINKAGE=GL

$$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.86 ± 0.14 ± 0.21	4.8k	24 SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
--------------------	------	---------	---------	--

²⁴ Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.

NODE=M067G01
NODE=M067G01

NODE=M067G01;LINKAGE=SH

$$\Gamma(\eta\phi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.43 ± 0.10 ± 0.09	25 AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
--------------------	-----------	----------	--

²⁵ From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

NODE=M067G7
NODE=M067G7

NODE=M067G7;LINKAGE=AU

$\phi(1680)$ BRANCHING RATIOS

$$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K\pi) \quad \Gamma_1/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

dominant	MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$
----------	------	--------	---

NODE=M067225

NODE=M067R3
NODE=M067R3

$$\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892) + \text{c.c.}) \quad \Gamma_3/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 ± 0.01	BUON	82 DM1	e^+e^-
-------------	------	--------	----------

NODE=M067R2
NODE=M067R2

$$\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892) + \text{c.c.}) \quad \Gamma_6/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<0.10	BUON	82 DM1	e^+e^-
-------	------	--------	----------

NODE=M067R1
NODE=M067R1

$$\Gamma(\eta\phi)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	35	26 ACHASOV	14 SND	1.15–2.00 $e^+e^- \rightarrow \eta\gamma$
------	----	------------	--------	---

²⁶ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

NODE=M067R01
NODE=M067R01

NODE=M067R01;LINKAGE=A

$$\Gamma(\eta\phi)/\Gamma(K\bar{K}^*(892) + \text{c.c.}) \quad \Gamma_9/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

≈ 0.37	27 AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \text{hadrons}$
--------	-----------	----------	--

²⁷ From the fit including data from AUBERT 07AK.

NODE=M067R5
NODE=M067R5

NODE=M067R5;LINKAGE=AU

$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	35	28 ACHASOV	14 SND	1.15–2.00 $e^+e^- \rightarrow \eta\gamma$
------	----	------------	--------	---

²⁸ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

NODE=M067R02
NODE=M067R02

NODE=M067R02;LINKAGE=A

$\phi(1680)$ REFERENCES

ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AMSLER	06	PL B639 165	C. Amisler <i>et al.</i>	(CBAR Collab.)
AKHMETSHIN	03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
Also		PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin	
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BUSENITZ	89	PR D40 1	J.K. Busenitz <i>et al.</i>	(ILL, FNAL)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
ATKINSON	85C	ZPHY C27 233	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)
MANE	82	PL 112B 178	F. Mane <i>et al.</i>	(LALO)
ASTON	81F	PL 104B 231	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ORSAY)

NODE=M067

REFID=55912
REFID=54066
REFID=53000
REFID=52242
REFID=51908
REFID=51136
REFID=49172
REFID=48827

REFID=48845
REFID=46323
REFID=44081
REFID=43168
REFID=41867
REFID=41369
REFID=40927
REFID=40581
REFID=40280

REFID=21596
REFID=21494
REFID=21590
REFID=21585
REFID=20553
REFID=21588

$\rho_3(1690)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

NODE=M015

 $\rho_3(1690)$ MASS

NODE=M015205

VALUE (MeV)

DOCUMENT ID

1688.8±2.1 OUR AVERAGE Includes data from the 5 datablocks that follow this one.

NODE=M015M

2 π MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015M1
NODE=M015M1**1686± 4 OUR AVERAGE**

1677±14		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow 2\pi p$
1679±11	476	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow$ $\pi^+ \pi^- n$
1678±12	175	¹ ANTIPOV	77	CIBS	0	25 $\pi^- p \rightarrow p3\pi$
1690± 7	600	¹ ENGLER	74	DBC	0	6 $\pi^+ n \rightarrow$ $\pi^+ \pi^- p$
1693± 8		² GRAYER	74	ASPK	0	17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$
1678±12		MATTHEWS	71C	DBC	0	7 $\pi^+ N$
1734±10		³ CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow$ $n2\pi$
1692±12		^{2,4} ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$
1737±23		ARMENISE	70	DBC	0	9 $\pi^+ N$
1650±35	122	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\pi$
1687±21		STUNTEBECK	70	HDBC	0	8 $\pi^- p, 5.4 \pi^+ d$
1683±13		ARMENISE	68	DBC	0	5.1 $\pi^+ d$
1670±30		GOLDBERG	65	HBC	0	6 $\pi^+ d, 8 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.² Uses same data as HYAMS 75.³ From a phase shift solution containing a $f_2'(1525)$ width two times larger than the $K\bar{K}$

result.

⁴ From phase-shift analysis. Error takes account of spread of different phase-shift solutions.NODE=M015M1;LINKAGE=E
NODE=M015M1;LINKAGE=G
NODE=M015M1;LINKAGE=M
NODE=M015M1;LINKAGE=I **$K\bar{K}$ AND $K\bar{K}\pi$ MODES**

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015M2
NODE=M015M2**1696± 4 OUR AVERAGE**

1699± 5		ALPER	80	CNTR	0	62 $\pi^- p \rightarrow$ $K^+ K^- n$
1698±12	6k	^{5,6} MARTIN	78D	SPEC		10 $\pi p \rightarrow$ $K_S^0 K^- p$
1692± 6		BLUM	75	ASPK	0	18.4 $\pi^- p \rightarrow$ $nK^+ K^-$
1690±16		ADERHOLZ	69	HBC	+	8 $\pi^+ p \rightarrow K\bar{K}\pi$
1694± 8		⁷ COSTA...	80	OMEG		10 $\pi^- p \rightarrow$ $K^+ K^- n$

⁵ From a fit to $J^P = 3^-$ partial wave.⁶ Systematic error on mass scale subtracted.⁷ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.NODE=M015M2;LINKAGE=P
NODE=M015M2;LINKAGE=S
NODE=M015M2;LINKAGE=L**(4 π) $^\pm$ MODE**

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015M3
NODE=M015M3**1686± 5 OUR AVERAGE** Error includes scale factor of 1.1.

1694± 6		⁸ EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow p4\pi$
1665±15	177	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
1670±10		THOMPSON	74	HBC	+	13 $\pi^+ p$
1687±20		CASON	73	HBC	-	8,18.5 $\pi^- p$
1685±14		⁹ CASON	73	HBC	-	8,18.5 $\pi^- p$
1680±40	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N4\pi$
1689±20	102	⁹ BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\rho$
1705±21		CASO	70	HBC	-	11.2 $\pi^- p \rightarrow$ $n\rho2\pi$

OCCUR=2

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

1718±10		¹⁰ EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow p4\pi$
1673± 9		¹¹ EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow p4\pi$
1733± 9	66	⁹ KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow p4\pi$
1630±15		HOLMES	72	HBC	+	10-12 $K^+ p$
1720±15		BALTAY	68	HBC	+	7, 8.5 $\pi^+ p$

OCCUR=2

OCCUR=3

⁸ From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.

⁹ From $\rho^\pm \rho^0$ mode.

¹⁰ From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.

¹¹ From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

NODE=M015M3;LINKAGE=A

NODE=M015M3;LINKAGE=F

NODE=M015M3;LINKAGE=B

NODE=M015M3;LINKAGE=C

$\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

1681± 7 OUR AVERAGE

1670±25		¹² ALDE	95	GAM2		38 $\pi^- p \rightarrow \omega\pi^0 n$
1690±15		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow \omega\pi p$
1666±14		GESSAROLI	77	HBC		11 $\pi^- p \rightarrow \omega\pi p$
1686± 9		THOMPSON	74	HBC	+	13 $\pi^+ p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1654±24		BARNHAM	70	HBC	+	10 $K^+ p \rightarrow \omega\pi X$
---------	--	---------	----	-----	---	------------------------------------

¹² Supersedes ALDE 92C.

NODE=M015M5;LINKAGE=A

$\eta\pi^+\pi^-$ MODE

(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015M6

NODE=M015M6

NODE=M015M6

1682±12 OUR AVERAGE

1685±10±20		AMELIN	00	VES		37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
1680±15		FUKUI	88	SPEC	0	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1700±47		¹³ ANDERSON	69	MMS	-	16 $\pi^- p$ backward
1632±15		^{13,14} FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow \rho MM$
1700±15		^{13,14} FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow \rho MM$
1748±15		^{13,14} FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow \rho MM$

OCCUR=2

OCCUR=3

¹³ Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+2\pi^-$, with 0, 1, 2 $\pi^+\pi^-$ pairs in ρ band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1976)

NODE=M015M6;LINKAGE=R

¹⁴ Not seen by BOWEN 72.

NODE=M015M6;LINKAGE=N

$\rho_3(1690)$ WIDTH

NODE=M015210


2 π , $K\bar{K}$, AND $K\bar{K}\pi$ MODES

VALUE (MeV)	DOCUMENT ID
-------------	-------------

161±10 OUR AVERAGE Includes data from the 5 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.

NODE=M015W

NODE=M015W



$\rho_3(1690)$ width, 2π , $K\bar{K}$, and $K\bar{K}\pi$ modes (MeV)

2 π MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W1
NODE=M015W1

186±14 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

220±29		DENNEY	83	LASS		10 $\pi^+ N$
246±37		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow 2\pi p$
116±30	476	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow$ $\pi^+ \pi^- n$
162±50	175	¹⁵ ANTIPOV	77	CIBS	0	25 $\pi^- p \rightarrow p3\pi$
167±40	600	ENGLER	74	DBC	0	6 $\pi^+ n \rightarrow$ $\pi^+ \pi^- p$
200±18		¹⁶ GRAYER	74	ASPK	0	17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$
156±36		MATTHEWS	71C	DBC	0	7 $\pi^+ N$
171±65		ARMENISE	70	DBC	0	9 $\pi^+ d$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
322±35		¹⁷ CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow$ $n2\pi$
240±30		^{16,18} ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$
180±30	122	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\pi$
267 ⁺⁷² -46		STUNTEBECK	70	HDBC	0	8 $\pi^- p, 5.4 \pi^+ d$
188±49		ARMENISE	68	DBC	0	5.1 $\pi^+ d$
180±40		GOLDBERG	65	HBC	0	6 $\pi^+ d, 8 \pi^- p$


¹⁵ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

¹⁶ Uses same data as HYAMS 75 and BECKER 79.

¹⁷ From a phase shift solution containing a $f'_2(1525)$ width two times larger than the $K\bar{K}$ result.

¹⁸ From phase-shift analysis. Error takes account of spread of different phase-shift solutions.

NODE=M015W1;LINKAGE=T
NODE=M015W1;LINKAGE=G
NODE=M015W1;LINKAGE=M
NODE=M015W1;LINKAGE=I



$K\bar{K}$ AND $K\bar{K}\pi$ MODES

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W2
NODE=M015W2

204±18 OUR AVERAGE

199±40	6000	¹⁹ MARTIN	78D	SPEC	10	$\pi^+ p \rightarrow K_S^0 K^- p$
205±20		BLUM	75	ASPK 0	18.4	$\pi^- p \rightarrow n K^+ K^-$
219±4		ALPER	80	CNTR 0	62	$\pi^- p \rightarrow K^+ K^- n$
186±11		²⁰ COSTA...	80	OMEG	10	$\pi^- p \rightarrow K^+ K^- n$
112±60		ADERHOLZ	69	HBC +	8	$\pi^+ p \rightarrow K\bar{K}\pi$

¹⁹ From a fit to $J^P = 3^-$ partial wave.

²⁰ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

NODE=M015W2;LINKAGE=P
NODE=M015W2;LINKAGE=L

$(4\pi)^\pm$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W3
NODE=M015W3

129±10 OUR AVERAGE

123±13		²¹ EVANGELIS...	81	OMEG -	12	$\pi^- p \rightarrow p4\pi$
105±30	177	BALTAY	78B	HBC +	15	$\pi^+ p \rightarrow p4\pi$
169 ⁺⁷⁰ -48		CASON	73	HBC -	8,18.5	$\pi^- p$
135±30	144	BARTSCH	70B	HBC +	8	$\pi^+ p \rightarrow N4\pi$
160±30	102	BARTSCH	70B	HBC +	8	$\pi^+ p \rightarrow N2\rho$

OCCUR=3

••• We do not use the following data for averages, fits, limits, etc. •••

230±28		²² EVANGELIS...	81	OMEG -	12	$\pi^- p \rightarrow p4\pi$
184±33		²³ EVANGELIS...	81	OMEG -	12	$\pi^- p \rightarrow p4\pi$
150	66	²⁴ KLIGER	74	HBC -	4.5	$\pi^- p \rightarrow p4\pi$
106±25		THOMPSON	74	HBC +	13	$\pi^+ p$
125 ⁺⁸³ -35		²⁴ CASON	73	HBC -	8,18.5	$\pi^- p$
130±30		HOLMES	72	HBC +	10-12	$K^+ p$
180±30	90	²⁴ BARTSCH	70B	HBC +	8	$\pi^+ p \rightarrow N a_2 \pi$
100±35		BALTAY	68	HBC +	7, 8.5	$\pi^+ p$

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

²¹ From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.

²² From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.

²³ From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

²⁴ From $\rho^\pm \rho^0$ mode.

NODE=M015W3;LINKAGE=A
NODE=M015W3;LINKAGE=B
NODE=M015W3;LINKAGE=C
NODE=M015W3;LINKAGE=F

$\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

190±40 OUR AVERAGE

230±65	²⁵ ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$
190±65	EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow \omega\pi p$
160±56	GESSAROLI	77	HBC	11 $\pi^- p \rightarrow \omega\pi p$
89±25	THOMPSON	74	HBC +	13 $\pi^+ p$
130 ⁺⁷³ ₋₄₃	BARNHAM	70	HBC +	10 $K^+ p \rightarrow \omega\pi X$

²⁵Supersedes ALDE 92C.

NODE=M015W5;LINKAGE=A

 $\eta\pi^+\pi^-$ MODE(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

126±40 OUR AVERAGE Error includes scale factor of 1.8.

220±30±50	AMELIN	00	VES	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
106±27	FUKUI	88	SPEC 0	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
195	²⁶ ANDERSON	69	MMS -	16 $\pi^- p$ backward
< 21	^{26,27} FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow \rho MM$
< 30	^{26,27} FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow \rho MM$
< 38	^{26,27} FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow \rho MM$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

195	²⁶ ANDERSON	69	MMS -	16 $\pi^- p$ backward
< 21	^{26,27} FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow \rho MM$
< 30	^{26,27} FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow \rho MM$
< 38	^{26,27} FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow \rho MM$

²⁶Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+2\pi^-$, with 0, 1, 2 $\pi^+\pi^-$ pairs in ρ^0 band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1979)²⁷Not seen by BOWEN 72.

NODE=M015W6

NODE=M015W6

NODE=M015W6

OCCUR=2

OCCUR=3

NODE=M015W6;LINKAGE=R

NODE=M015W6;LINKAGE=N

 $\rho_3(1690)$ DECAY MODES

NODE=M015215;NODE=M015

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 4π	(71.1 ± 1.9) %	
Γ_2 $\pi^\pm\pi^+\pi^-\pi^0$	(67 ± 22) %	
Γ_3 $\omega\pi$	(16 ± 6) %	
Γ_4 $\pi\pi$	(23.6 ± 1.3) %	
Γ_5 $K\bar{K}\pi$	(3.8 ± 1.2) %	
Γ_6 $K\bar{K}$	(1.58 ± 0.26) %	1.2
Γ_7 $\eta\pi^+\pi^-$	seen	
Γ_8 $\rho(770)\eta$	seen	
Γ_9 $\pi\pi\rho$	seen	
Excluding 2ρ and $a_2(1320)\pi$.		
Γ_{10} $a_2(1320)\pi$	seen	
Γ_{11} $\rho\rho$	seen	
Γ_{12} $\phi\pi$		
Γ_{13} $\eta\pi$		
Γ_{14} $\pi^\pm 2\pi^+ 2\pi^- \pi^0$		

DESIG=2

DESIG=11

DESIG=7

DESIG=1

DESIG=3

DESIG=4

DESIG=13

DESIG=14;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=10

DESIG=12

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 10 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 14.7$ for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-77			
x_5	-74	17		
x_6	-15	2	0	
	x_1	x_4	x_5	

$\rho_3(1690)$ BRANCHING RATIOS

NODE=M015220

$\Gamma(\pi\pi) / \Gamma_{\text{total}}$ Γ_4 / Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.236 ± 0.013 OUR FIT**0.243 ± 0.013 OUR AVERAGE**

0.259 ^{+0.018} _{-0.019}	BECKER	79	ASPK	0	17 $\pi^- p$ polarized
0.23 ± 0.02	CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow$ $n^2\pi$
0.22 ± 0.04	²⁸ MATTHEWS	71C	HDBC	0	7 $\pi^+ n \rightarrow \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.245 ± 0.006	²⁹ ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow$ $\pi^+ \pi^- n$

²⁸ One-pion-exchange model used in this estimation.²⁹ From phase-shift analysis of HYAMS 75 data.NODE=M015R1
NODE=M015R1

$\Gamma(\pi\pi) / \Gamma(\pi^\pm \pi^+ \pi^- \pi^0)$ Γ_4 / Γ_2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.35 ± 0.11

CASON	73	HBC	-	8,18.5 $\pi^- p$
-------	----	-----	---	------------------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<0.2	HOLMES	72	HBC	+	10-12 $K^+ p$
<0.12	BALLAM	71B	HBC	-	16 $\pi^- p$

NODE=M015R2
NODE=M015R2

$\Gamma(\pi\pi) / \Gamma(4\pi)$ Γ_4 / Γ_1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.332 ± 0.026 OUR FIT Error includes scale factor of 1.1.**0.30 ± 0.10** BALTAY 78B HBC 0 15 $\pi^+ p \rightarrow p 4\pi$ NODE=M015R3
NODE=M015R3


$\Gamma(K\bar{K}) / \Gamma(\pi\pi)$ Γ_6 / Γ_4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.067 ± 0.011 OUR FIT Error includes scale factor of 1.2.**0.118^{+0.040}_{-0.032} OUR AVERAGE** Error includes scale factor of 1.7. See the ideogram below.

0.191 ^{+0.040} _{-0.037}	GORLICH	80	ASPK	0	17,18 $\pi^- p$ polarized
0.08 ± 0.03	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.08 ^{+0.08} _{-0.03}	CRENNELL	68B	HBC		6.0 $\pi^- p$

NODE=M015R4
NODE=M015R4



$\Gamma(K\bar{K}\pi)/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_4
0.16±0.05 OUR FIT					
0.16±0.05	³⁰ BARTSCH	70B	HBC	+	8 $\pi^+ p$

NODE=M015R5
NODE=M015R5

³⁰ Increased by us to correspond to $B(\rho_3(1690) \rightarrow \pi\pi)=0.24$.

NODE=M015R5;LINKAGE=A

$[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$(\Gamma_9+\Gamma_{10}+\Gamma_{11})/\Gamma_2$
0.94±0.09 OUR AVERAGE					
0.96±0.21	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.88±0.15	BALLAM	71B	HBC	-	16 $\pi^- p$
1 ±0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$
consistent with 1	CASO	68	HBC	-	11 $\pi^- p$

NODE=M015R6
NODE=M015R6

$\Gamma(\rho\rho)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{11}/Γ_2
0.12±0.11		BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.56	66	KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow p4\pi$
0.13±0.09		³¹ THOMPSON	74	HBC	+	13 $\pi^+ p$
0.7 ±0.15		BARTSCH	70B	HBC	+	8 $\pi^+ p$

NODE=M015R7
NODE=M015R7

³¹ $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

NODE=M015R7;LINKAGE=T

$\Gamma(\rho\rho)/[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_{11}/(\Gamma_9+\Gamma_{10}+\Gamma_{11})$
0.48±0.16	CASO	68	HBC	-	11 $\pi^- p$

NODE=M015R8
NODE=M015R8

$\Gamma(a_2(1320)\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{10}/Γ_2
0.66±0.08	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.36±0.14	³² THOMPSON	74	HBC	+	13 $\pi^+ p$
not seen	CASON	73	HBC	-	8,18.5 $\pi^- p$
0.6 ±0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.6	BALTAY	68	HBC	+	7,8.5 $\pi^+ p$

NODE=M015R9
NODE=M015R9

³² $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

NODE=M015R9;LINKAGE=T

$\Gamma(\omega\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$ Γ_3/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
0.23±0.05 OUR AVERAGE		Error includes scale factor of 1.2.			
0.33±0.07		THOMPSON	74	HBC	+ 13 $\pi^+ p$
0.12±0.07		BALLAM	71B	HBC	- 16 $\pi^- p$
0.25±0.10		BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$
0.25±0.10		JOHNSTON	68	HBC	- 7.0 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.11	95	BALTAY	78B	HBC	+ 15 $\pi^+ p \rightarrow p4\pi$
<0.09		KLIGER	74	HBC	- 4.5 $\pi^- p \rightarrow p4\pi$

NODE=M015R10
NODE=M015R10 $\Gamma(\phi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$ Γ_{12}/Γ_2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.11	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R11
NODE=M015R11 $\Gamma(\pi^\pm 2\pi^+ 2\pi^- \pi^0)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$ Γ_{14}/Γ_2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.15	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R12
NODE=M015R12 $\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$ Γ_{13}/Γ_2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.02	THOMPSON	74	HBC	+ 13 $\pi^+ p$

NODE=M015R13
NODE=M015R13 $\Gamma(K\bar{K})/\Gamma_{total}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.0158±0.0026 OUR FIT	Error includes scale factor of 1.2.			
0.0130±0.0024 OUR AVERAGE				
0.013 ±0.003	COSTA...	80	OMEG	0 10 $\pi^- p \rightarrow K^+ K^- n$
0.013 ±0.004	³³ MARTIN	78B	SPEC	- 10 $\pi p \rightarrow K_S^0 K^- p$

NODE=M015R14
NODE=M015R14³³From $(\Gamma_4\Gamma_6)^{1/2} = 0.056 \pm 0.034$ assuming $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.

NODE=M015R14;LINKAGE=B

 $\Gamma(\omega\pi)/[\Gamma(\omega\pi) + \Gamma(\rho\rho)]$ $\Gamma_3/(\Gamma_3+\Gamma_{11})$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.22±0.08	CASON	73	HBC	- 8,18.5 $\pi^- p$

NODE=M015R16
NODE=M015R16 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{total}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	FUKUI	88	SPEC 8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M015R17
NODE=M015R17 $\Gamma(a_2(1320)\pi)/\Gamma(\rho(770)\eta)$ Γ_{10}/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
5.5±2.0	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M015R18
NODE=M015R18 $\rho_3(1690)$ REFERENCES

NODE=M015

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP	REFID=44371
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41859
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21665
COSTA...	80	NP B175 402	G. Costa de Beaugard <i>et al.</i>	(BARI, BONN+)	REFID=20737
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)	REFID=20738
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
MARTIN	78B	NP B140 158	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=21273
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=21272
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)	REFID=20230
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP	REFID=21651
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20642
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110

GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
KLIGER	74	SJNP 19 428 Translated from YAF 19 839.	G.K. Kliger <i>et al.</i>	(ITEP)	REFID=21648
OREN	74	NP B71 189	Y. Oren <i>et al.</i>	(ANL, OXF)	REFID=20221
THOMPSON	74	NP B69 220	G. Thompson <i>et al.</i>	(PURD)	REFID=21650
CASON	73	PR D7 1971	N.M. Cason <i>et al.</i>	(NDAM)	REFID=20606
BOWEN	72	PRL 29 890	D.R. Bowen <i>et al.</i>	(NEAS, STON)	REFID=21711
HOLMES	72	PR D6 3336	R. Holmes <i>et al.</i>	(ROCH)	REFID=21639
BALLAM	71B	PR D3 2606	J. Ballam <i>et al.</i>	(SLAC)	REFID=21630
MATTHEWS	71C	NP B33 1	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC) JP	REFID=21633
ARMENISE	70	LNC 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
BARNHAM	70	PRL 24 1083	K.W.J. Barnham <i>et al.</i>	(BIRM)	REFID=21624
BARTSCH	70B	NP B22 109	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN)	REFID=21625
CASO	70	LNC 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20590
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ANDERSON	69	PRL 22 1390	E.W. Anderson <i>et al.</i>	(BNL, CMU)	REFID=20795
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+) I	REFID=20054
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I	REFID=21531
CASO	68	NC 54A 983	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20586
CRENNELL	68B	PL 28B 136	D.J. Crennell <i>et al.</i>	(BNL)	REFID=21616
JOHNSTON	68	PRL 20 1414	T.F. Johnston <i>et al.</i>	(TNTO, WISC) IJP	REFID=21617
FOCACCI	66	PRL 17 890	M.N. Focacci <i>et al.</i>	(CERN)	REFID=20402
GOLDBERG	65	PL 17 354	M. Goldberg <i>et al.</i>	(CERN, EPOL, ORSAY+)	REFID=21601

NODE=M065

$\rho(1700)$

$$I^G(J^{PC}) = 1^+(1^-)$$

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=M065

$\rho(1700)$ MASS

NODE=M065205

$\eta\rho^0$ AND $\pi^+\pi^-$ MODES

VALUE (MeV)

DOCUMENT ID

1720±20 OUR ESTIMATE

NODE=M065M0

NODE=M065M0

→ UNCHECKED ←

$\eta\rho^0$ MODE

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M065M6

NODE=M065M6

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1740±20

ANTONELLI 88 DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$

1701±15

¹ FUKUI 88 SPEC 8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

¹ Assuming $\rho^+ f_0(1370)$ decay mode interferes with $a_1(1260)^+\pi$ background. From a two Breit-Wigner fit.

NODE=M065M;LINKAGE=B

$\pi\pi$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M065M1

NODE=M065M1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1780 ±20 $^{+15}_{-20}$

63.5k

² ABRAMOWICZ12 ZEUS $e p \rightarrow e\pi^+\pi^-p$

1861 ±17

³ LEES 12G BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$

1728 ±17 ±89

5.4M

^{4,5} FUJIKAWA 08 BELL $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ 1780 $^{+37}_{-29}$ ⁶ ABELE 97 CBAR $\bar{p}n \rightarrow \pi^-\pi^0\pi^0$

1719 ±15

⁶ BERTIN 97C OBLX 0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$

1730 ±30

CLEGG 94 RVUE $e^+e^- \rightarrow \pi^+\pi^-$

1768 ±21

BISELLO 89 DM2 $e^+e^- \rightarrow \pi^+\pi^-$

1745.7±91.9

DUBNICKA 89 RVUE $e^+e^- \rightarrow \pi^+\pi^-$

1546 ±26

GESHKEN... 89 RVUE

1650

⁷ ERKAL 85 RVUE 20–70 $\gamma p \rightarrow \gamma\pi$

1550 ±70

ABE 84B HYBR 20 $\gamma p \rightarrow \pi^+\pi^-p$

1590 ±20

⁸ ASTON 80 OMEG 20–70 $\gamma p \rightarrow p2\pi$

1600 ±10

⁹ ATIYA 79B SPEC 50 $\gamma C \rightarrow C2\pi$ 1598 $^{+24}_{-22}$ BECKER 79 ASPK 17 π^-p polarized

1659 ±25

⁷ LANG 79 RVUE

1575

⁷ MARTIN 78C RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$

1610 ±30

⁷ FROGGATT 77 RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$

1590 ±20

¹⁰ HYAMS 73 ASPK 17 $\pi^-p \rightarrow \pi^+\pi^-n$

- ² Using the KUHN 90 parametrization of the pion form factor, neglecting ρ - ω interference.
³ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
⁴ $|F_\pi(0)|^2$ fixed to 1.
⁵ From the GOUNARIS 68 parametrization of the pion form factor.
⁶ T-matrix pole.
⁷ From phase shift analysis of HYAMS 73 data.
⁸ Simple relativistic Breit-Wigner fit with constant width.
⁹ An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.
¹⁰ Included in BECKER 79 analysis.

NODE=M065M1;LINKAGE=AB
 NODE=M065M1;LINKAGE=LE

NODE=M065M1;LINKAGE=FU
 NODE=M065M1;LINKAGE=GO
 NODE=M065M;LINKAGE=QQ
 NODE=M065M;LINKAGE=P
 NODE=M065M;LINKAGE=M
 NODE=M065M;LINKAGE=R

NODE=M065M;LINKAGE=H

NODE=M065M8
 NODE=M065M8

$\pi\omega$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1708±41	7815	¹¹ ACHASOV	13	SND 1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1550 to 1620		¹² ACHASOV	00i	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1580 to 1710		¹³ ACHASOV	00i	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1710±90		ACHASOV	97	RVUE $e^+e^- \rightarrow \omega\pi^0$

OCCUR=2

NODE=M065M8;LINKAGE=AC

NODE=M065M;LINKAGE=I1

NODE=M065M;LINKAGE=I2

- ¹¹ From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.
¹² Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00i on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.
¹³ Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00i on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$.

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1740.8±22.2	27k	¹⁴ ABELE	99D	CBAR ±	0.0 $\bar{p}p \rightarrow K^+K^-\pi^0$
1582 ±36	1600	CLELAND	82B	SPEC ±	50 $\pi p \rightarrow K_S^0 K^\pm p$

NODE=M065M2
 NODE=M065M2

- ¹⁴ K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

NODE=M065M2;LINKAGE=AN

$2(\pi^+\pi^-)$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1851 ⁺²⁷ ₋₂₄		ACHASOV	97	RVUE $e^+e^- \rightarrow 2(\pi^+\pi^-)$
1570±20		¹⁵ CORDIER	82	DM1 $e^+e^- \rightarrow 2(\pi^+\pi^-)$
1520±30		¹⁶ ASTON	81E	OMEG 20-70 $\gamma p \rightarrow p4\pi$
1654±25		¹⁷ DIBIANCA	81	DBC $\pi^+d \rightarrow pp2(\pi^+\pi^-)$
1666±39		¹⁵ BACCI	80	FRAG $e^+e^- \rightarrow 2(\pi^+\pi^-)$
1780	34	KILLIAN	80	SPEC 11 $e^-p \rightarrow 2(\pi^+\pi^-)$
1500		¹⁸ ATIYA	79B	SPEC 50 $\gamma C \rightarrow C4\pi^\pm$
1570±60	65	¹⁹ ALEXANDER	75	HBC 7.5 $\gamma p \rightarrow p4\pi$
1550±60		¹⁶ CONVERSI	74	OSPK $e^+e^- \rightarrow 2(\pi^+\pi^-)$
1550±50	160	SCHACHT	74	STRC 5.5-9 $\gamma p \rightarrow p4\pi$
1450±100	340	SCHACHT	74	STRC 9-18 $\gamma p \rightarrow p4\pi$
1430±50	400	BINGHAM	72B	HBC 9.3 $\gamma p \rightarrow p4\pi$

NODE=M065M4
 NODE=M065M4

OCCUR=2

NODE=M065M;LINKAGE=A
 NODE=M065M4;LINKAGE=M
 NODE=M065M;LINKAGE=O
 NODE=M065M;LINKAGE=C
 NODE=M065M;LINKAGE=D

$\pi^+\pi^-\pi^0\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1660±30	ATKINSON	85B	OMEG 20-70 γp

NODE=M065M5
 NODE=M065M5

$3(\pi^+\pi^-)$ AND $2(\pi^+\pi^-\pi^0)$ MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1730±34	²⁰ FRABETTI	04	E687 $\gamma p \rightarrow 3\pi^+3\pi^-p$
1783±15	CLEGG	90	RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

NODE=M065M7
 NODE=M065M7

- ²⁰ From a fit with two resonances with the JACOB 72 continuum.

NODE=M065M;LINKAGE=PI

$\rho(1700)$ WIDTH $\eta\rho^0$ AND $\pi^+\pi^-$ MODES

VALUE (MeV) _____ DOCUMENT ID _____
250±100 OUR ESTIMATE

NODE=M065210

NODE=M065W0
 NODE=M065W0

→ UNCHECKED ←

 $\eta\rho^0$ MODE

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ COMMENT _____

NODE=M065W6
 NODE=M065W6

The data in this block is included in the average printed for a previous datablock.

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

150±30 ANTONELLI 88 DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$
 282±44 21 FUKUI 88 SPEC 8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
 21 Assuming $\rho^+ f_0(1370)$ decay mode interferes with $a_1(1260)^+\pi$ background. From a two Breit-Wigner fit.

NODE=M065W;LINKAGE=B

 $\pi\pi$ MODE

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ COMMENT _____

NODE=M065W1
 NODE=M065W1

The data in this block is included in the average printed for a previous datablock.

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

310 ± 30 $^{+25}_{-35}$ 63.5k 22 ABRAMOWICZ12 ZEUS $ep \rightarrow e\pi^+\pi^-p$
 316 ± 26 23 LEES 12G BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$
 164 ± 21 $^{+89}_{-26}$ 5.4M 24,25 FUJIKAWA 08 BELL $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
 275 ± 45 26 ABELE 97 CBAR $\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
 310 ± 40 26 BERTIN 97C OBLX 0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
 400 ±100 CLEGG 94 RVUE $e^+e^- \rightarrow \pi^+\pi^-$
 224 ± 22 BISELLO 89 DM2 $e^+e^- \rightarrow \pi^+\pi^-$
 242.5±163.0 DUBNICKA 89 RVUE $e^+e^- \rightarrow \pi^+\pi^-$
 620 ± 60 GESHKEN... 89 RVUE
 <315 27 ERKAL 85 RVUE 20-70 $\gamma p \rightarrow \gamma\pi$
 280 $^{+30}_{-80}$ ABE 84B HYBR 20 $\gamma p \rightarrow \pi^+\pi^-p$
 230 ± 80 28 ASTON 80 OMEG 20-70 $\gamma p \rightarrow p2\pi$
 283 ± 14 29 ATIYA 79B SPEC 50 $\gamma C \rightarrow C2\pi$
 175 $^{+98}_{-53}$ BECKER 79 ASPK 17 π^-p polarized
 232 ± 34 27 LANG 79 RVUE
 340 27 MARTIN 78C RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$
 300 ±100 27 FROGGATT 77 RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$
 180 ± 50 30 HYAMS 73 ASPK 17 $\pi^-p \rightarrow \pi^+\pi^-n$

NODE=M065W1;LINKAGE=AB

NODE=M065W1;LINKAGE=LE

22 Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.23 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.24 $|F_\pi(0)|^2$ fixed to 1.

25 From the GOUNARIS 68 parametrization of the pion form factor.

26 T-matrix pole.

27 From phase shift analysis of HYAMS 73 data.

28 Simple relativistic Breit-Wigner fit with constant width.

29 An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

30 Included in BECKER 79 analysis.

NODE=M065W1;LINKAGE=FU

NODE=M065W1;LINKAGE=GO

NODE=M065W;LINKAGE=QQ

NODE=M065W;LINKAGE=P

NODE=M065W;LINKAGE=M

NODE=M065W;LINKAGE=R

NODE=M065W;LINKAGE=H

 $K\bar{K}$ MODE

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____

NODE=M065W2
 NODE=M065W2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

187.2± 26.7 27k 31 ABELE 99D CBAR ± 0.0 $\bar{p}p \rightarrow K^+K^-\pi^0$
 265 ±120 1600 CLELAND 82B SPEC ± 50 $\pi p \rightarrow K_S^0 K^\pm p$

31 K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

NODE=M065W2;LINKAGE=AN

2($\pi^+\pi^-$) MODE

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
510±40		32 CORDIER	82 DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
400±50		33 ASTON	81E OMEG	20-70 $\gamma p \rightarrow p4\pi$
400±146		34 DIBIANCA	81 DBC	$\pi^+d \rightarrow p\rho 2(\pi^+\pi^-)$
700±160		32 BACCI	80 FRAG	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
100	34	KILLIAN	80 SPEC	11 $e^-p \rightarrow 2(\pi^+\pi^-)$
600		35 ATIYA	79B SPEC	50 $\gamma C \rightarrow C4\pi^\pm$
340±160	65	36 ALEXANDER	75 HBC	7.5 $\gamma p \rightarrow p4\pi$
360±100		33 CONVERSI	74 OSPK	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
400±120	160	37 SCHACHT	74 STRC	5.5-9 $\gamma p \rightarrow p4\pi$
850±200	340	37 SCHACHT	74 STRC	9-18 $\gamma p \rightarrow p4\pi$
650±100	400	BINGHAM	72B HBC	9.3 $\gamma p \rightarrow p4\pi$

³² Simple relativistic Breit-Wigner fit with model-dependent width.

³³ Simple relativistic Breit-Wigner fit with constant width.

³⁴ One peak fit result.

³⁵ Parameters roughly estimated, not from a fit.

³⁶ Skew mass distribution compensated by Ross-Stodolsky factor.

³⁷ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

NODE=M065W4
NODE=M065W4

OCCUR=2

NODE=M065W;LINKAGE=A
NODE=M065W4;LINKAGE=M
NODE=M065W;LINKAGE=O
NODE=M065W;LINKAGE=C
NODE=M065W;LINKAGE=D
NODE=M065W;LINKAGE=E

 $\pi^+\pi^-\pi^0\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300±50	ATKINSON	85B OMEG	20-70 γp

 $\omega\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
350 to 580	38 ACHASOV	00I SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
490 to 1040	39 ACHASOV	00I SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

³⁸ Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00I on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.

³⁹ Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00I on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$.

NODE=M065W9
NODE=M065W9

OCCUR=2

NODE=M065W;LINKAGE=I1

NODE=M065W;LINKAGE=I2

3($\pi^+\pi^-$) AND 2($\pi^+\pi^-\pi^0$) MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
315±100	⁴⁰ FRABETTI	04 E687	$\gamma p \rightarrow 3\pi^+ 3\pi^- p$
285±20	CLEGG	90 RVUE	$e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

⁴⁰ From a fit with two resonances with the JACOB 72 continuum.

NODE=M065W7
NODE=M065W7

NODE=M065W;LINKAGE=PI

 $\rho(1700)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 4π	
Γ_2 $2(\pi^+\pi^-)$	large
Γ_3 $\rho\pi\pi$	dominant
Γ_4 $\rho^0\pi^+\pi^-$	large
Γ_5 $\rho^0\pi^0\pi^0$	
Γ_6 $\rho^\pm\pi^\mp\pi^0$	large
Γ_7 $a_1(1260)\pi$	seen
Γ_8 $h_1(1170)\pi$	seen
Γ_9 $\pi(1300)\pi$	seen
Γ_{10} $\rho\rho$	seen
Γ_{11} $\pi^+\pi^-$	seen
Γ_{12} $\pi\pi$	seen
Γ_{13} $K\bar{K}^*(892) + c.c.$	seen
Γ_{14} $\eta\rho$	seen
Γ_{15} $a_2(1320)\pi$	not seen
Γ_{16} $K\bar{K}$	seen
Γ_{17} e^+e^-	seen
Γ_{18} $\pi^0\omega$	seen

NODE=M065215;NODE=M065

DESIG=20

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=12;OUR EST;→ UNCHECKED ←

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=7

DESIG=9;OUR EST;→ UNCHECKED ←

DESIG=15;OUR EST;→ UNCHECKED ←

DESIG=16;OUR EST;→ UNCHECKED ←

DESIG=17;OUR EST;→ UNCHECKED ←

DESIG=18;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=13;OUR EST;→ UNCHECKED ←

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=11;OUR EST;→ UNCHECKED ←

DESIG=14;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=6;OUR EST;→ UNCHECKED ←

$\rho(1700) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M065225

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the cross-section into channel i in e^+e^- annihilation.

NODE=M065225

 $\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_{17}/\Gamma$ NODE=M065G2
NODE=M065G2

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.2	DEL COURT	81B	DM1 $e^+e^- \rightarrow 2(\pi^+\pi^-)$
2.83 ± 0.42	BACCI	80	FRAG $e^+e^- \rightarrow 2(\pi^+\pi^-)$

 $\Gamma(\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}\Gamma_{17}/\Gamma$ NODE=M065G4
NODE=M065G4

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13	⁴¹ DIEKMAN	88	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
0.029 ^{+0.016} _{-0.012}	KURDADZE	83	OLYA 0.64–1.4 $e^+e^- \rightarrow \pi^+\pi^-$

⁴¹ Using total width = 220 MeV.

NODE=M065G4;LINKAGE=B

 $\Gamma(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{13}\Gamma_{17}/\Gamma$ NODE=M065G10
NODE=M065G10

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.305 ± 0.071	⁴² BIZOT	80	DM1 e^+e^-
---------------	---------------------	----	--------------

⁴² Model dependent.

NODE=M065G;LINKAGE=M

 $\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_{17}/\Gamma$ NODE=M065G11
NODE=M065G11

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7 ± 3	ANTONELLI	88	DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$
-------	-----------	----	---

 $\Gamma(K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}\Gamma_{17}/\Gamma$ NODE=M065G5
NODE=M065G5

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.035 ± 0.029	⁴³ BIZOT	80	DM1 e^+e^-
---------------	---------------------	----	--------------

⁴³ Model dependent.

NODE=M065G5;LINKAGE=M

 $\Gamma(\rho\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_3\Gamma_{17}/\Gamma$ NODE=M065G12
NODE=M065G12

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.510 ± 0.090	⁴⁴ BIZOT	80	DM1 e^+e^-
---------------	---------------------	----	--------------

⁴⁴ Model dependent.

NODE=M065G12;LINKAGE=M

 $\rho(1700) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M065240

 $\Gamma(\pi^0\omega)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{17}/\Gamma$ NODE=M065R01
NODE=M065R01

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.7 ± 0.4	7815	⁴⁵ ACHASOV	13	SND 1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
-----------	------	-----------------------	----	---

⁴⁵ From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M065R01;LINKAGE=AC

 $\rho(1700) \text{ BRANCHING RATIOS}$

NODE=M065230

 $\Gamma(\rho\pi\pi)/\Gamma(4\pi) \quad \Gamma_3/\Gamma_1$ NODE=M065R19
NODE=M065R19

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.28 ± 0.06	⁴⁶ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-------------	---------------------	-----	--------------------------------------

⁴⁶ $\omega\pi$ not included.

NODE=M065R;LINKAGE=BL

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$ Γ_4/Γ_2

VALUE	EPTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1.0		DEL COURT	81B	DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
0.7 ± 0.1	500	SCHACHT	74	STRC	5.5–18 $\gamma p \rightarrow p4\pi$
0.80		47 BINGHAM	72B	HBC	9.3 $\gamma p \rightarrow p4\pi$

⁴⁷ The $\pi\pi$ system is in S -wave.

NODE=M065R1
NODE=M065R1

NODE=M065R1;LINKAGE=S

 $\Gamma(\rho^0\pi^0\pi^0)/\Gamma(\rho^\pm\pi^\mp\pi^0)$ Γ_5/Γ_6

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.10	ATKINSON	85B	OMEG	20–70 γp
< 0.15	ATKINSON	82	OMEG 0	20–70 $\gamma p \rightarrow p4\pi$

NODE=M065R6
NODE=M065R6

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_7/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.16 ± 0.05	48 ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$
-------------	----------	-----	------	---------------------------------

⁴⁸ $\omega\pi$ not included.

NODE=M065R15
NODE=M065R15

NODE=M065R15;LINKAGE=BL

 $\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$ Γ_8/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ± 0.06	49 ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$
-------------	----------	-----	------	---------------------------------

⁴⁹ $\omega\pi$ not included.

NODE=M065R16
NODE=M065R16

NODE=M065R16;LINKAGE=BL

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_9/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.30 ± 0.10	50 ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$
-------------	----------	-----	------	---------------------------------

⁵⁰ $\omega\pi$ not included.

NODE=M065R17
NODE=M065R17

NODE=M065R17;LINKAGE=BL

 $\Gamma(\rho\rho)/\Gamma(4\pi)$ Γ_{10}/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.09 ± 0.03	51 ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 5\pi$
-------------	----------	-----	------	---------------------------------

⁵¹ $\omega\pi$ not included.

NODE=M065R18
NODE=M065R18

NODE=M065R18;LINKAGE=BL

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.287 ^{+0.043} _{-0.042}	BECKER	79	ASPK	17 $\pi^- p$ polarized
0.15 to 0.30	52 MARTIN	78C	RVUE	17 $\pi^- p \rightarrow \pi^+\pi^- n$
< 0.20	53 COSTA...	77B	RVUE	$e^+e^- \rightarrow 2\pi, 4\pi$
0.30 ± 0.05	52 FROGGATT	77	RVUE	17 $\pi^- p \rightarrow \pi^+\pi^- n$
< 0.15	54 EISENBERG	73	HBC	5 $\pi^+ p \rightarrow \Delta^{++} 2\pi$
0.25 ± 0.05	55 HYAMS	73	ASPK	17 $\pi^- p \rightarrow \pi^+\pi^- n$

⁵² From phase shift analysis of HYAMS 73 data.

⁵³ Estimate using unitarity, time reversal invariance, Breit-Wigner.

⁵⁴ Estimated using one-pion-exchange model.

⁵⁵ Included in BECKER 79 analysis.

NODE=M065R5
NODE=M065R5

NODE=M065R5;LINKAGE=P
NODE=M065R5;LINKAGE=C
NODE=M065R5;LINKAGE=E
NODE=M065R5;LINKAGE=H

 $\Gamma(\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$ Γ_{11}/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 ± 0.05	ASTON	80	OMEG	20–70 $\gamma p \rightarrow p2\pi$
< 0.14	56 DAVIER	73	STRC	6–18 $\gamma p \rightarrow p4\pi$
< 0.2	57 BINGHAM	72B	HBC	9.3 $\gamma p \rightarrow p2\pi$

⁵⁶ Upper limit is estimate.

⁵⁷ 2σ upper limit.

NODE=M065R3
NODE=M065R3

NODE=M065R3;LINKAGE=E
NODE=M065R3;LINKAGE=S

$\Gamma(\pi\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.16±0.04	58,59 ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
			⁵⁸ Using ABELE 97.
			⁵⁹ $\omega\pi$ not included.

 Γ_{12}/Γ_1 NODE=M065R20
NODE=M065R20NODE=M065R;LINKAGE=LK
NODE=M065R20;LINKAGE=BL $\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	COAN	04	CLEO $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$
---------------	------	----	--

 Γ_{13}/Γ NODE=M065R21
NODE=M065R21 $\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(2(\pi^+\pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.15±0.03	⁶⁰ DELCOURT	81B	DM1 $e^+e^- \rightarrow \bar{K}K\pi$
-----------	------------------------	-----	--------------------------------------

⁶⁰ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

 Γ_{13}/Γ_2 NODE=M065R9
NODE=M065R9

NODE=M065R9;LINKAGE=D

 $\Gamma(\eta\rho)/\Gamma_{total}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen		AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
<0.04		DONNACHIE 87B	RVUE	
<0.02	58	ATKINSON 86B	OMEG	20-70 γp

 Γ_{14}/Γ NODE=M065R12
NODE=M065R12 $\Gamma(\eta\rho)/\Gamma(2(\pi^+\pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123±0.027	DELCOURT	82	DM1 $e^+e^- \rightarrow \pi^+\pi^- MM$
~0.1	ASTON	80	OMEG 20-70 γp

 Γ_{14}/Γ_2 NODE=M065R8
NODE=M065R8 $\Gamma(\pi^+\pi^- \text{ neutrals})/\Gamma(2(\pi^+\pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6±0.4	⁶¹ BALLAM	74	HBC 9.3 γp
---------	----------------------	----	--------------------

⁶¹ Upper limit. Background not subtracted.

 $(\Gamma_5+\Gamma_6+0.714\Gamma_{14})/\Gamma_2$ NODE=M065R7
NODE=M065R7

NODE=M065R7;LINKAGE=U

 $\Gamma(a_2(1320)\pi)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
----------	--------	----	---

 Γ_{15}/Γ NODE=M065R14
NODE=M065R14 $\Gamma(K\bar{K})/\Gamma(2(\pi^+\pi^-))$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.015±0.010		⁶² DELCOURT	81B	DM1	$e^+e^- \rightarrow \bar{K}K$
<0.04	95	BINGHAM	72B	HBC	0 9.3 γp

⁶² Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

 Γ_{16}/Γ_2 NODE=M065R4
NODE=M065R4

NODE=M065R4;LINKAGE=D

 $\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+c.c.)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.052±0.026	BUON	82	DM1 $e^+e^- \rightarrow \text{hadrons}$
-------------	------	----	---

 Γ_{16}/Γ_{13} NODE=M065R10
NODE=M065R10 $\Gamma(\pi^0\omega)/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		MATVIENKO	15	BELL $\bar{B}^0 \rightarrow D^{*0}\omega\pi^-$
seen	1.6k	ACHASOV	12	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
not seen	2382	AKHMETSHIN	03B	CMD2 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
seen		ACHASOV	97	RVUE $e^+e^- \rightarrow \omega\pi^0$

 Γ_{18}/Γ NODE=M065R13
NODE=M065R13

$\rho(1700)$ REFERENCES

NODE=M065

MATVIENKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
		Translated from ZETFP 94 796.			
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
FRABETTI	04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=49614
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
ACHASOV	001	PL B486 29	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47931
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=41355
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
GESHKEN...	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)	REFID=41017
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
DIEKMAN	88	PRPL 159 99	B. Diekmann	(BONN)	REFID=40272
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
ATKINSON	86B	ZPHY C30 531	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21508
ATKINSON	85B	ZPHY C26 499	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21506
ERKAL	85	ZPHY C29 485	C. Erkal, M.G. Olsson	(WISC)	REFID=20136
ABE	84B	PRL 53 751	K. Abe <i>et al.</i>	(SLAC HFP Collab.)	REFID=21503
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
		Translated from ZETFP 37 613.			
ATKINSON	82	PL 108B 55	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21493
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)	REFID=21494
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
CORDIER	82	PL 109B 129	A. Cordier <i>et al.</i>	(LALO)	REFID=21495
DELCOURT	82	PL 113B 93	B. Delcourt <i>et al.</i>	(LALO)	REFID=21496
ASTON	81E	NP B189 15	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=21487
DELCOURT	81B	Bonn Conf. 205	B. Delcourt	(ORSAY)	REFID=21490
Also		PL 109B 129	A. Cordier <i>et al.</i>	(LALO)	REFID=21495
DIBIANCA	81	PR D23 595	F.A. di Bianca <i>et al.</i>	(CASE, CMU)	REFID=21492
ASTON	80	PL 92B 215	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=21478
BACCI	80	PL 95B 139	C. Bacci <i>et al.</i>	(ROMA, FRAS)	REFID=21481
BIZOT	80	Madison Conf. 546	J.C. Bizot <i>et al.</i>	(LALO, MONP)	REFID=21482
KILLIAN	80	PR D21 3005	T.J. Killian <i>et al.</i>	(CORN)	REFID=21484
ATIYA	79B	PRL 43 1691	M.S. Atiya <i>et al.</i>	(COLU, ILL, FNAL)	REFID=21470
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parada	(GRAZ)	REFID=20126
MARTIN	78C	ANP 114 1	A.D. Martin, M.R. Pennington	(CERN)	REFID=21661
COSTA...	77B	PL 71B 345	B. Costa de Beauregard, B. Pire, T.N. Truong	(EPOL)	REFID=21465
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
ALEXANDER	75	PL 57B 487	G. Alexander <i>et al.</i>	(TELA)	REFID=21450
BALLAM	74	NP B76 375	J. Ballam <i>et al.</i>	(SLAC, LBL, MPIM)	REFID=20610
CONVERSI	74	PL 52B 493	M. Conversi <i>et al.</i>	(ROMA, FRAS)	REFID=20637
SCHACHT	74	NP B81 205	P. Schacht <i>et al.</i>	(MPIM)	REFID=21449
DAVIER	73	NP B58 31	M. Davier <i>et al.</i>	(SLAC)	REFID=21434
EISENBERG	73	PL 43B 149	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=21435
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
BINGHAM	72B	PL 41B 635	H.H. Bingham <i>et al.</i>	(LBL, UCB, SLAC) IGJP	REFID=21426
JACOB	72	PR D5 1847	M. Jacob, R. Slansky		REFID=49668
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054

$a_2(1700)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

NODE=M162

OMITTED FROM SUMMARY TABLE

 $a_2(1700)$ MASS

NODE=M162M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1732±16 OUR AVERAGE					
Error includes scale factor of 1.9.					
1737± 5± 7		ABE	04	BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
1698±44		¹ AMSLER	02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0\eta\eta$
1660±40		ABELE	99B	CBAR	1.94 $\bar{p}p \rightarrow \pi^0\eta\eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1675±25		ANISOVICH	09	RVUE	0.0 $\bar{p}p, \pi N$
1722± 9±15	18k	² SCHEGELSKY	06	RVUE 0	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
1702± 7	80k	³ UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
1721±13±44	145k	LU	05	B852	18 $\pi^-p \rightarrow \omega\pi^-\pi^0p$
1767±14	221	⁴ ACCIARRI	01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$
~ 1775		⁵ GRYGOREV	99	SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1752±21± 4		ACCIARRI	97T	L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

NODE=M162M

¹ T-matrix pole.² From analysis of L3 data at 183–209 GeV.³ Statistical error only.⁴ Spin 2 dominant, isospin not determined, could also be $I=1$.⁵ Possibly two $J^P = 2^+$ resonances with isospins 0 and 1.

NODE=M162M;LINKAGE=TT

NODE=M162M;LINKAGE=SC

NODE=M162M;LINKAGE=ST

NODE=M162M;LINKAGE=HA

NODE=M162M;LINKAGE=GR

 $a_2(1700)$ WIDTH

NODE=M162W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
194± 40 OUR AVERAGE					
Error includes scale factor of 1.6. See the ideogram below.					
151± 22±24		ABE	04	BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
265± 55		⁶ AMSLER	02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0\eta\eta$
280± 70		ABELE	99B	CBAR	1.94 $\bar{p}p \rightarrow \pi^0\eta\eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
270 ⁺⁵⁰ ₋₂₀		ANISOVICH	09	RVUE	0.0 $\bar{p}p, \pi N$
336± 20±20	18k	⁷ SCHEGELSKY	06	RVUE 0	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
417± 19	80k	⁸ UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
279± 49±66	145k	LU	05	B852	18 $\pi^-p \rightarrow \omega\pi^-\pi^0p$
187± 60	221	⁹ ACCIARRI	01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$
150±110±34		ACCIARRI	97T	L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

NODE=M162W


⁶ T-matrix pole.⁷ From analysis of L3 data at 183–209 GeV.⁸ Statistical error only.⁹ Spin 2 dominant, isospin not determined, could also be $I=1$.

NODE=M162W;LINKAGE=TT

NODE=M162W;LINKAGE=SC

NODE=M162W;LINKAGE=ST

NODE=M162W;LINKAGE=HA



a₂(1700) DECAY MODES

NODE=M162215;NODE=M162

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi$	seen
Γ_2 $\gamma\gamma$	
Γ_3 $\rho\pi$	
Γ_4 $f_2(1270)\pi$	
Γ_5 $K\bar{K}$	seen
Γ_6 $\omega\pi^-\pi^0$	seen
Γ_7 $\omega\rho$	seen

DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=1
DESIG=2
DESIG=3
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EVAL;→ UNCHECKED ←
DESIG=7;OUR EVAL;→ UNCHECKED ←

a₂(1700) PARTIAL WIDTHS

NODE=M162220

$\Gamma(\eta\pi)$						Γ_1
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT		
9.5±2.0	870	¹⁰ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$		
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
$\Gamma(\gamma\gamma)$						Γ_2
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT		
0.30±0.05	870	¹⁰ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$		
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
$\Gamma(K\bar{K})$						Γ_5
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT		
5.0±3.0	870	¹⁰ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$		
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
¹⁰ From analysis of L3 data at 91 and 183–209 GeV, using a ₂ (1700) mass of 1730 MeV and width of 340 MeV, and SU(3) relations.						

NODE=M162W3
NODE=M162W3

NODE=M162W2
NODE=M162W2

NODE=M162W1
NODE=M162W1

NODE=M162W1;LINKAGE=SC

a₂(1700) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M162225

$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$						$(\Gamma_3 + \Gamma_4)\Gamma_2/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT		
0.29±0.04±0.02		ACCIARRI	97T L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$		
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
0.37 ^{+0.12} _{-0.08} ±0.10	18k	¹¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$		

NODE=M162G1
NODE=M162G1

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_2/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$20.6 \pm 4.2 \pm 4.6$	¹² ABE	04 BELL	$10.6 e^+e^- \rightarrow e^+e^-K^+K^-$
$49 \pm 11 \pm 13$	¹³ ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183-209 \text{ GeV}$

¹¹ From analysis of L3 data at 183–209 GeV.¹² Assuming spin 2.¹³ Spin 2 dominant, isospin not determined, could also be $I=1$.NODE=M162G2
NODE=M162G2NODE=M162G1;LINKAGE=SC
NODE=M162G2;LINKAGE=AB
NODE=M162G;LINKAGE=HA $a_2(1700)$ BRANCHING RATIOS $\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$ Γ_3/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$3.4 \pm 0.4 \pm 0.1$	18k	¹⁴ SCHEGELSKY	06 RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

¹⁴ From analysis of L3 data at 183–209 GeV.

NODE=M162235

NODE=M162R01
NODE=M162R01

NODE=M162R01;LINKAGE=SC

 $a_2(1700)$ REFERENCES

ANISOVICH 09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev
SCHEGELSKY 06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>
SCHEGELSKY 06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>
UMAN 06	PR D73 052009	I. Uman <i>et al.</i> (FNAL E835)
LU 05	PRL 94 032002	M. Lu <i>et al.</i> (BNL E852 Collab.)
ABE 04	EPJ C32 323	K. Abe <i>et al.</i> (BELLE Collab.)
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>
ACCIARRI 01H	PL B501 173	M. Acciarri <i>et al.</i> (L3 Collab.)
ABELE 99B	EPJ C8 67	A. Abele <i>et al.</i> (Crystal Barrel Collab.)
GRYGOREV 99	PAN 62 470	V.K. Grygorev <i>et al.</i>
	Translated from YAF 62 513.	
ACCIARRI 97T	PL B413 147	M. Acciarri <i>et al.</i> (L3 Collab.)

NODE=M162

REFID=52719
REFID=51186
REFID=51185
REFID=51063
REFID=50459
REFID=49650
REFID=48580
REFID=48321
REFID=46904
REFID=46909

REFID=45761

NODE=M068

 $f_0(1710)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See our mini-review in the 2004 edition of this *Review*, Physics Letters **B592** 1 (2004). See also the mini-review on scalar mesons under $f_0(500)$ (see the index for the page number).

NODE=M068

 $f_0(1710)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1723^{+6}_{-5}		OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.
1759 ± 6	$^{+14}_{-25}$ 5.5k	¹ ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
1750^{+6}_{-7}	$^{+29}_{-18}$	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
1701 ± 5	$^{+9}_{-2}$ 4k	² CHEKANOV	08 ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
1765^{+4}_{-3}	± 13	ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1760 ± 15	$^{+15}_{-10}$	³ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
1738 ± 30		ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+K^-$
1740 ± 4	$^{+10}_{-25}$	⁴ BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
1740^{+30}_{-25}		⁴ BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
1698 ± 18		⁵ BARBERIS	00E	$450 pp \rightarrow p_f \eta \eta p_S$
1710 ± 12	± 11	⁶ BARBERIS	99D OMEG	$450 pp \rightarrow K^+K^-, \pi^+\pi^-$
1710 ± 25		⁷ FRENCH	99	$300 pp \rightarrow p_f(K^+K^-)p_S$
1707 ± 10		⁸ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+K^-, K_S^0 K_S^0$
1698 ± 15		⁸ AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1720 ± 10	± 10	⁹ BALTRUSAIT..	87 MRK3	$J/\psi \rightarrow \gamma K^+K^-$
1742 ± 15		⁸ WILLIAMS	84 MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$
1670 ± 50		BLOOM	83 CBAL	$J/\psi \rightarrow \gamma 2\eta$

NODE=M068M


NODE=M068M

• • • We do not use the following data for averages, fits, limits, etc. • • •

1744 ± 7 ± 5	381	10,11	DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$	
1705 ± 11 ± 5	237	10,11	DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$	OCCUR=2
1706 ± 4 ± 5	1.0k	10,11	DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$	OCCUR=3
1690 ± 8 ± 3	349	10,11	DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$	OCCUR=4
1750 ± 13			AMSLER	06	CBAR $1.64 \bar{p} p \rightarrow K^+ K^- \pi^0$	
1747 ± 5	80k	12,13	UMAN	06	E835 $5.2 \bar{p} p \rightarrow \eta \eta \pi^0$	
1776 ± 15			VLADIMIRSK...	06	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
1790 ⁺⁴⁰ ₋₃₀			3 ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$	
1670 ± 20			12 BINON	05	GAMS $33 \pi^- p \rightarrow \eta \eta n$	
1726 ± 7	74		13 CHEKANOV	04	ZEUS $e p \rightarrow K_S^0 K_S^0 X$	
1732 ± 15			14 ANISOVICH	03	RVUE	
1682 ± 16			TIKHOMIROV	03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
1670 ± 26	3.6k	4,15	NICHITIU	02	OBLX	
1770 ± 12			16,17 ANISOVICH	99B	SPEC $0.6-1.2 p \bar{p} \rightarrow \eta \eta \pi^0$	
1730 ± 15			4 BARBERIS	99	OMEG $450 p p \rightarrow p_S p_f K^+ K^-$	
1750 ± 20			4 BARBERIS	99B	OMEG $450 p p \rightarrow p_S p_f \pi^+ \pi^-$	
1750 ± 30			18 ANISOVICH	98B	RVUE Compilation	
1720 ± 39			BAI	98H	BES $J/\psi \rightarrow \gamma \pi^0 \pi^0$	
1775 ± 1.5	57		19 BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$	
1690 ± 11			20 ABREU	96C	DLPH $Z^0 \rightarrow K^+ K^- + X$	
1696 ± 5 ⁺⁹ ₋₃₄			9 BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$	
1781 ± 8 ⁺¹⁰ ₋₃₁			4 BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$	OCCUR=2
1768 ± 14			BALOSHIN	95	SPEC $40 \pi^- C \rightarrow K_S^0 K_S^0 X$	
1750 ± 15			21 BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$	
1620 ± 16			9 BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$	OCCUR=2
1748 ± 10			8 ARMSTRONG	93C	E760 $\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
~ 1750			BREAKSTONE	93	SFM $p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$	
1744 ± 15			22 ALDE	92D	GAM2 $38 \pi^- p \rightarrow \eta \eta n$	
1713 ± 10			23 ARMSTRONG	89D	OMEG $300 p p \rightarrow p p K^+ K^-$	
1706 ± 10			23 ARMSTRONG	89D	OMEG $300 p p \rightarrow p p K_S^0 K_S^0$	OCCUR=2
1700 ± 15			9 BOLONKIN	88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
1720 ± 60			4 BOLONKIN	88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$	OCCUR=2
1638 ± 10			24 FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	
1690 ± 4			25 FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	OCCUR=2
1755 ± 8			26 ALDE	86C	GAM2 $38 \pi^- p \rightarrow n 2 \eta$	
1730 ⁺² ₋₁₀			27 LONGACRE	86	RVUE $22 \pi^- p \rightarrow n 2 K_S^0$	
1650 ± 50			BURKE	82	MRK2 $J/\psi \rightarrow \gamma 2 \rho$	
1640 ± 50			28,29 EDWARDS	82D	CBAL $J/\psi \rightarrow \gamma 2 \eta$	
1730 ± 10 ± 20			30 ETKIN	82C	MPS $23 \pi^- p \rightarrow n 2 K_S^0$	
1					From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.	NODE=M068M;LINKAGE=D
2					In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.	NODE=M068M;LINKAGE=HE
3					This state may be different from $f_0(1710)$, see CLOSE 05.	
4					$J^P = 0^+$.	NODE=M068M;LINKAGE=AB
5					T-matrix pole.	NODE=M068M;LINKAGE=A8
6					Supersedes BARBERIS 99 and BARBERIS 99B.	NODE=M068M;LINKAGE=TP
7					$J^P = 0^+$, supersedes by ARMSTRONG 89D.	NODE=M068M;LINKAGE=BD
8					No J^{PC} determination.	NODE=M068M;LINKAGE=C3
9					$J^P = 2^+$.	NODE=M068M;LINKAGE=A1
10					Using CLEO-c data but not authored by the CLEO Collaboration.	NODE=M068M;LINKAGE=A3
11					From a fit to a Breit-Wigner line shape with fixed $\Gamma = 135$ MeV.	NODE=M068M;LINKAGE=F
12					Breit-Wigner mass.	NODE=M068M;LINKAGE=G
13					Systematic errors not estimated.	NODE=M068M;LINKAGE=BW
14					K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.	NODE=M068M;LINKAGE=CH
15					Decaying to $f_0(1370) \pi \pi$.	NODE=M068M;LINKAGE=KM
16					$J^P = 0^+$.	NODE=M068M;LINKAGE=NC
17					Not seen by AMSLER 02.	NODE=M068M;LINKAGE=AV
18					T-matrix pole, assuming $J^P = 0^+$	NODE=M068M;LINKAGE=NS
19					No J^{PC} determination.	NODE=M068M;LINKAGE=AN
						NODE=M068M;LINKAGE=4A

- 20 No J^{PC} determination, width not determined.
- 21 From a fit to the 0^+ partial wave.
- 22 ALDE 92D combines all the GAMS-2000 data.
- 23 $J^P = 2^+$, superseded by FRENCH 99.
- 24 From an analysis ignoring interference with $f'_2(1525)$.
- 25 From an analysis including interference with $f'_2(1525)$.
- 26 Superseded by ALDE 92D.
- 27 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
- 28 $J^P = 2^+$ preferred.
- 29 From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.
- 30 Superseded by LONGACRE 86.

NODE=M068M;LINKAGE=A4
 NODE=M068M;LINKAGE=Q0
 NODE=M068M;LINKAGE=AA
 NODE=M068M;LINKAGE=C
 NODE=M068M;LINKAGE=A
 NODE=M068M;LINKAGE=B
 NODE=M068M;LINKAGE=BB
 NODE=M068M;LINKAGE=A9
 NODE=M068M;LINKAGE=B2
 NODE=M068M;LINKAGE=E
 NODE=M068M;LINKAGE=B1



$f_0(1710)$ WIDTH

NODE=M068W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
139 ± 8	OUR AVERAGE	Error includes scale factor of 1.1.		
172 ± 10	+32 -16	5.5k	1 ABLIKIM 13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
139 +11 -12	+96 -50		UEHARA 13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
100 ± 24	+7 -22	4k	2 CHEKANOV 08 ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
145 ± 8	±69		ABLIKIM 06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
125 ± 25	+10 -15		3 ABLIKIM 05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^- K^+ K^-$
125 ± 20			ABLIKIM 04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
166 +5 -8	+15 -10		4 BAI 03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
120 +50 -40			4 BAI 00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
120 ± 26			5 BARBERIS 00E	450 $pp \rightarrow p_f \eta \eta p_s$
126 ± 16	±18		6 BARBERIS 99D OMEG	450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$
105 ± 34			7 FRENCH 99	300 $pp \rightarrow p_f(K^+ K^-) p_s$
166.4 ± 33.2			8 AUGUSTIN 88 DM2	$J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0$
136 ± 28			8 AUGUSTIN 87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
130 ± 20			9 BALTRUSAIT..87 MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
57 ± 38			10 WILLIAMS 84 MPSF	200 $\pi^- N \rightarrow 2K_S^0 X$
160 ± 80			BLOOM 83 CBAL	$J/\psi \rightarrow \gamma 2\eta$

NODE=M068W

• • • We do not use the following data for averages, fits, limits, etc. • • •

148	$\begin{matrix} + 40 \\ - 30 \end{matrix}$		AMSLER	06	CBAR	1.64	$\bar{p}p \rightarrow K^+ K^- \pi^0$	
188	± 13	80k	3,11 UMAN	06	E835	5.2	$\bar{p}p \rightarrow \eta\eta\pi^0$	
250	± 30		VLADIMIRSK...	06	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$	
270	$\begin{matrix} + 60 \\ - 30 \end{matrix}$		12 ABLIKIM	05	BES2		$J/\psi \rightarrow \phi\pi^+\pi^-$	
260	± 50		3 BINON	05	GAMS	33	$\pi^- p \rightarrow \eta\eta n$	
38	$\begin{matrix} + 20 \\ - 14 \end{matrix}$	74	11 CHEKANOV	04	ZEUS		$e p \rightarrow K_S^0 K_S^0 X$	
144	± 30		13,14 ANISOVICH	03	RVUE			
320	$\begin{matrix} + 50 \\ - 20 \end{matrix}$		14,15 ANISOVICH	03	RVUE			OCCUR=2
102	± 26		TIKHOMIROV	03	SPEC	40.0	$\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
267	± 44	3651	4,16 NICHITIU	02	OBLX			
220	± 40		17,18 ANISOVICH	99B	SPEC	0.6-1.2	$p\bar{p} \rightarrow \eta\eta\pi^0$	
100	± 25		4 BARBERIS	99	OMEG	450	$p p \rightarrow p_S p_f K^+ K^-$	
160	± 30		4 BARBERIS	99B	OMEG	450	$p p \rightarrow p_S p_f \pi^+ \pi^-$	
250	± 140		19 ANISOVICH	98B	RVUE		Compilation	
30	± 7	57	20 BARKOV	98			$\pi^- p \rightarrow K_S^0 K_S^0 n$	
103	± 18	$\begin{matrix} + 30 \\ - 11 \end{matrix}$	9 BAI	96C	BES		$J/\psi \rightarrow \gamma K^+ K^-$	
85	± 24	$\begin{matrix} + 22 \\ - 19 \end{matrix}$	4 BAI	96C	BES		$J/\psi \rightarrow \gamma K^+ K^-$	OCCUR=2
56	± 19		BALOSHIN	95	SPEC	40	$\pi^- C \rightarrow K_S^0 K_S^0 X$	
160	± 40		21 BUGG	95	MRK3		$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$	
160	$\begin{matrix} + 60 \\ - 20 \end{matrix}$		9 BUGG	95	MRK3		$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$	OCCUR=2
264	± 25		8 ARMSTRONG	93C	E760		$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$	
200	to 300		BREAKSTONE	93	SFM		$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$	
< 80	90% CL		22 ALDE	92D	GAM2	38	$\pi^- p \rightarrow \eta\eta N^*$	
181	± 30		23 ARMSTRONG	89D	OMEG	300	$p p \rightarrow p p K^+ K^-$	
104	± 30		23 ARMSTRONG	89D	OMEG	300	$p p \rightarrow p p K_S^0 K_S^0$	OCCUR=2
30	± 20		9 BOLONKIN	88	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$	
350	± 150		4 BOLONKIN	88	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$	OCCUR=2
148	± 17		24 FALVARD	88	DM2		$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	
184	± 6		25 FALVARD	88	DM2		$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	OCCUR=2
122	$\begin{matrix} + 74 \\ - 15 \end{matrix}$		26 LONGACRE	86	RVUE	22	$\pi^- p \rightarrow n 2 K_S^0$	
200	± 100		BURKE	82	MRK2		$J/\psi \rightarrow \gamma 2 p$	
220	$\begin{matrix} + 100 \\ - 70 \end{matrix}$		27,28 EDWARDS	82D	CBAL		$J/\psi \rightarrow \gamma 2 \eta$	
200	$\begin{matrix} + 156 \\ - 9 \end{matrix}$		29 ETKIN	82B	MPS	23	$\pi^- p \rightarrow n 2 K_S^0$	

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

³ Breit-Wigner width.

⁴ $J^P = 0^+$.

⁵ T-matrix pole.

⁶ Supersedes BARBERIS 99 and BARBERIS 99B.

⁷ $J^P = 0^+$, superseded by ARMSTRONG 89D.

⁸ No J^{PC} determination.

⁹ $J^P = 2^+$.

¹⁰ No J^{PC} determination.

¹¹ Systematic errors not estimated.

¹² This state may be different from $f_0(1710)$, see CLOSE 05.

¹³ (Solution I)

¹⁴ K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

¹⁵ (Solution I)

¹⁶ Decaying to $f_0(1370)\pi\pi$.

¹⁷ $J^P = 0^+$.

¹⁸ Not seen by AMSLER 02.

¹⁹ T-matrix pole, assuming $J^P = 0^+$

²⁰ No J^{PC} determination.

²¹ From a fit to the 0^+ partial wave.

NODE=M068W;LINKAGE=F

NODE=M068W;LINKAGE=HE

NODE=M068W;LINKAGE=BW

NODE=M068W;LINKAGE=A8

NODE=M068W;LINKAGE=TP

NODE=M068W;LINKAGE=BD

NODE=M068W;LINKAGE=C3

NODE=M068W;LINKAGE=A1

NODE=M068W;LINKAGE=A3

NODE=M068M;LINKAGE=WI

NODE=M068W;LINKAGE=CH

NODE=M068W;LINKAGE=AB

NODE=M068W;LINKAGE=K1

NODE=M068W;LINKAGE=KM

NODE=M068W;LINKAGE=K2

NODE=M068W;LINKAGE=NC

NODE=M068W;LINKAGE=AV

NODE=M068W;LINKAGE=NS

NODE=M068W;LINKAGE=AN

NODE=M068W;LINKAGE=4A

NODE=M068W;LINKAGE=Q0

- 22 ALDE 92D combines all the GAMS-2000 data.
 23 $J^P = 2^+, (0^+ \text{ excluded})$.
 24 From an analysis ignoring interference with $f'_2(1525)$.
 25 From an analysis including interference with $f'_2(1525)$.
 26 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
 27 $J^P = 2^+$ preferred.
 28 From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.
 29 From an amplitude analysis of the $K_S^0 K_S^0$ system, superseded by LONGACRE 86.

NODE=M068W;LINKAGE=AA
 NODE=M068W;LINKAGE=B
 NODE=M068W;LINKAGE=C
 NODE=M068W;LINKAGE=D
 NODE=M068W;LINKAGE=A9
 NODE=M068W;LINKAGE=B2
 NODE=M068W;LINKAGE=E
 NODE=M068W;LINKAGE=A

$f_0(1710)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $\eta\eta$	seen
Γ_3 $\pi\pi$	seen
Γ_4 $\gamma\gamma$	
Γ_5 $\omega\omega$	seen

NODE=M068215;NODE=M068

DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6
 DESIG=4

$f_0(1710)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_4/\Gamma$
$12_{-2}^{+3} - 22_{-8}^{+7}$		UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M068220

NODE=M068G2
 NODE=M068G2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<480	95	ALBRECHT	90G	ARG	$\gamma\gamma \rightarrow K^+ K^-$
<110	95	¹ BEHREND	89C	CELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
<280	95	¹ ALTHOFF	85B	TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

¹ Assuming helicity 2.

NODE=M068G2;LINKAGE=F

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_3\Gamma_4/\Gamma$
<0.82	95	¹ BARATE	00E	ALEP $\gamma\gamma \rightarrow \pi^+ \pi^-$	

NODE=M068G3
 NODE=M068G3

¹ Assuming spin 0.

NODE=M068G;LINKAGE=Z

$f_0(1710)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ	
seen	1004	¹ DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$		
seen	349	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$		
0.36 ± 0.12		ALBALADEJO	08	RVUE		
$0.38_{-0.19}^{+0.09}$		² LONGACRE	86	MPS $22 \pi^- p \rightarrow n 2 K_S^0$		

NODE=M068225

NODE=M068R2
 NODE=M068R2

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

NODE=M068R2;LINKAGE=A
 NODE=M068R;LINKAGE=L

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	Γ_2/Γ
0.22 ± 0.12	ALBALADEJO 08	RVUE	
$0.18_{-0.13}^{+0.03}$	¹ LONGACRE 86	RVUE	

NODE=M068R1
 NODE=M068R1

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

NODE=M068R1;LINKAGE=L

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen	381	¹ DOBBS 15		$J/\psi \rightarrow \gamma\pi^+\pi^-$
seen	237	¹ DOBBS 15		$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
not seen		AMSLER 02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
$0.039^{+0.002}_{-0.024}$		² LONGACRE 86	RVUE	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

NODE=M068R5
NODE=M068R5

OCCUR=2

NODE=M068R5;LINKAGE=A
NODE=M068R5;LINKAGE=L

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$ Γ_3/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$0.41^{+0.11}_{-0.17}$		ABLIKIM 06V	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
0.32±0.14		ALBALADEJO 08	RVUE	
< 0.11	95	¹ ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+K^-$
$5.8^{+9.1}_{-5.5}$		² ANISOVICH 02D	SPEC	Combined fit
$0.2 \pm 0.024 \pm 0.036$		BARBERIS 99D	OMEG 450	$p\bar{p} \rightarrow K^+K^-, \pi^+\pi^-$
0.39 ± 0.14		ARMSTRONG 91	OMEG 300	$p\bar{p} \rightarrow p\rho\pi\pi, p\rho K\bar{K}$

¹ Using data from ABLIKIM 04A.

² From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

NODE=M068R6
NODE=M068R6

NODE=M068R;LINKAGE=AB
NODE=M068R;LINKAGE=CH

 $\Gamma(\eta\eta)/\Gamma(K\bar{K})$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.48 ± 0.15		BARBERIS 00E		$450 p\bar{p} \rightarrow p_f\eta\eta p_s$
$0.46^{+0.70}_{-0.38}$		¹ ANISOVICH 02D	SPEC	Combined fit
< 0.02	90	² PROKOSHKIN 91	GA24	$300 \pi^- p \rightarrow \pi^- p\eta\eta$

¹ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

² Combining results of GAM4 with those of ARMSTRONG 89D.

NODE=M068R7
NODE=M068R7

NODE=M068R7;LINKAGE=CH

NODE=M068R;LINKAGE=A

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen	180	ABLIKIM 06H	BES	$J/\psi \rightarrow \gamma\omega\omega$

NODE=M068R3
NODE=M068R3

 $f_0(1710)$ REFERENCES

NODE=M068

DOBBS 15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM 13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55387
UEHARA 13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ALBALADEJO 08	PRL 101 252002	M. Albaladejo, J.A. Oller		REFID=52656
CHEKANOV 08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52275
ABLIKIM 06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM 06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
AMSLER 06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)	REFID=51136
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK... 06	PAN 69 493	V.V. Vladimirov <i>et al.</i>	(ITEP, Moscow)	REFID=51191
	Translated from YAF 69 515.			
ABLIKIM 05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM 05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
BINON 05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
	Translated from YAF 68 998.			
CLOSE 05	PR D71 094022	F.E. Close, Q. Zhao		REFID=50788
ABLIKIM 04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49740
ABLIKIM 04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
CHEKANOV 04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=49672
PDG 04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
ANISOVICH 03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BAI 03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
TIKHOMIROV 03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
	Translated from YAF 66 860.			
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>		REFID=48580
ANISOVICH 02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
	Translated from YAF 65 1583.			

NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47428
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>		REFID=46886
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)	REFID=47491
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BARKOV	98	JETPL 68 764	B.P. Barkov <i>et al.</i>		REFID=46616
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=44621
		Translated from YAF 58 50.			
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone <i>et al.</i>	(IOWA, CERN, DORT+)	REFID=43312
ALDE	92D	PL B284 457	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=41591
Also		SJNP 54 451	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=44696
		Translated from YAF 54 745.			
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2, GAM4 Collab.)	REFID=41719
		Translated from DANS 316 900.			
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41010
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=21694
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>	(VAND, NDAM, TUFTS+)	REFID=21693
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20391

NODE=M114

$\eta(1760)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Seen by DM2 in the $\rho\rho$ system (BISELLO 89B). Structure in this region has been reported before in the same system (BALTRUSAITIS 86B) and in the $\omega\omega$ system (BALTRUSAITIS 85C, BISELLO 87).

NODE=M114

$\eta(1760)$ MASS

NODE=M114M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1751 ± 15 OUR AVERAGE				
1768 ⁺²⁴ ₋₂₅ ± 10	465	¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
1744 ± 10 ± 15	1045	² ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1703 ⁺¹² ₋₁₁ ± 2		³ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
1760 ± 11	320	⁴ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

NODE=M114M

OCCUR=2

- ¹ From a single-resonance fit.
- ² From a partial wave analysis including $\eta(1760)$, $f_0(1710)$, $f_2(1640)$, and $f_2(1910)$.
- ³ From a two-resonance fit.
- ⁴ Estimated by us from various fits. Systematic uncertainties not estimated.

NODE=M114M;LINKAGE=ZA
 NODE=M114M;LINKAGE=MA
 NODE=M114M;LINKAGE=ZH
 NODE=M114M;LINKAGE=A

$\eta(1760)$ WIDTH

NODE=M114W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
240 ± 30 OUR AVERAGE				
224 ⁺⁶² ₋₅₆ ± 25	465	⁵ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
244 ⁺²⁴ ₋₂₁ ± 25	1045	⁶ ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
42 ⁺³⁶ ₋₂₂ ± 15		⁷ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
60 ± 16	320	⁸ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

NODE=M114W

OCCUR=2

- ⁵ From a single-resonance fit.
- ⁶ From a partial wave analysis including $\eta(1760)$, $f_0(1710)$, $f_2(1640)$, and $f_2(1910)$.
- ⁷ From a two-resonance fit.
- ⁸ Estimated by us from various fits. Systematic uncertainties not estimated.

NODE=M114W;LINKAGE=ZA
 NODE=M114W;LINKAGE=MA
 NODE=M114W;LINKAGE=ZH
 NODE=M114W;LINKAGE=B

$\eta(1760)$ DECAY MODES

NODE=M114215;NODE=M114

Mode	Fraction (Γ_i/Γ)
Γ_1 4π	
Γ_2 $2\pi^+2\pi^-$	seen
Γ_3 $\pi^+\pi^-2\pi^0$	seen
Γ_4 $\rho^0\rho^0$	seen
Γ_5 $\rho^+\rho^-$	seen
Γ_6 $2(\pi^+\pi^-\pi^0)$	
Γ_7 $\omega\omega$	seen
Γ_8 $\eta'\pi^+\pi^-$	seen
Γ_9 $\gamma\gamma$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8;OUR EVAL;→ UNCHECKED ←

DESIG=9;OUR EVAL;→ UNCHECKED ←

 $\eta(1760)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M114225

$\Gamma(\eta'\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_8\Gamma_9/\Gamma$
VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT	
$28.2^+_{-7.5} \pm 3.7$	465	⁹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	

NODE=M114G01

NODE=M114G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.0^+_{-1.2} \pm 0.8$	52	¹⁰ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	OCCUR=2
$18^{+13}_{-10} \pm 5$	315	¹¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	OCCUR=3

⁹ From a single-resonance fit.¹⁰ From a two-resonance fit. For constructive interference with the X(1835).¹¹ From a two-resonance fit. For destructive interference with the X(1835).

NODE=M114G01;LINKAGE=ZH

NODE=M114G01;LINKAGE=ZA

NODE=M114G01;LINKAGE=ZN

 $\eta(1760)$ BRANCHING RATIOS

NODE=M114210

$\Gamma(2\pi^+2\pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M114R01

NODE=M114R01

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$	Γ_3/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^- 2\pi^0$

NODE=M114R02

NODE=M114R02

$\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$	Γ_4/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma \rho^0 \rho^0$
seen	BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \gamma \rho^0 \rho^0$

NODE=M114R03

NODE=M114R03

$\Gamma(\rho^+\rho^-)/\Gamma_{\text{total}}$	Γ_5/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma \rho^+ \rho^-$
seen	BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \gamma \rho^+ \rho^-$

NODE=M114R04

NODE=M114R04

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$	Γ_7/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	87 DM2	$J/\psi \rightarrow \omega\omega$
seen	BALTRUSAIT...85C	MRK3	$J/\psi \rightarrow \gamma\omega\omega$

NODE=M114R06

NODE=M114R06

 $\eta(1760)$ REFERENCES

NODE=M114

ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)	REFID=54763
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
BALTRUSAIT... 86	PR	D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BALTRUSAIT... 86B	PR	D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT... 85C	PRL	55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095

$\pi(1800)$

$$J^{PC} = 1^-(0^-+)$$

See also minireview under non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M075

NODE=M075

$\pi(1800)$ MASS

NODE=M075M


NODE=M075M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1812±12 OUR AVERAGE		Error includes scale factor of 2.3. See the ideogram below.			
1785±9 ⁺¹² ₋₆	420k	ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1876±18±16	4k	¹ EUGENIO	08	B852	- 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
1774±18±20		² CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1863±9±10		³ CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1840±10±10	1200	AMELIN	96B	VES	- 37 $\pi^- A \rightarrow \eta \eta \pi^- A$
1775±7±10		⁴ AMELIN	95B	VES	- 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
1790±14		⁵ BERDNIKOV	94	VES	- 37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
1873±33±20		BELADIDZE	92C	VES	- 36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$
1814±10±23	426±57	BITYUKOV	91	VES	- 36 $\pi^- C \rightarrow \pi^- \eta \eta C$
1770±30	1100	BELLINI	82	SPEC	- 40 $\pi^- A \rightarrow 3\pi A$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1737±5±15		AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$

OCCUR=2

NODE=M075M;LINKAGE=SP
 NODE=M075M;LINKAGE=C1
 NODE=M075M;LINKAGE=C2
 NODE=M075M;LINKAGE=AX
 NODE=M075M;LINKAGE=A

- ¹ From a single-pole fit.
- ² In the $f_0(980)\pi$ wave.
- ³ In the $f_0(500)\pi$ wave.
- ⁴ From a fit to $J^{PC} = 0^-+ f_0(980)\pi, f_0(1370)\pi$ waves.
- ⁵ From a fit to $J^{PC} = 0^-+ K_0^*(1430)K^-$ and $f_0(980)\pi^-$ waves.



$\pi(1800)$ WIDTH

NODE=M075W

NODE=M075W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
208±22⁺²¹₋₃₇	420k	ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
221±26±38	4k	⁶ EUGENIO	08	B852	- 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
223±48±50		⁷ CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

191±21±20		⁸ CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	OCCUR=2
210±30±30	1200	AMELIN	96B	VES	- 37 $\pi^- A \rightarrow \eta \eta \pi^- A$	
190±15±15		⁹ AMELIN	95B	VES	- 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
210±70		¹⁰ BERDNIKOV	94	VES	- 37 $\pi^- A \rightarrow K^+ K^- \pi^- A$	
225±35±20		BELADIDZE	92C	VES	- 36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$	
205±18±32	426 ± 57	BITYUKOV	91	VES	- 36 $\pi^- C \rightarrow \pi^- \eta \eta C$	
310±50	1100	BELLINI	82	SPEC	- 40 $\pi^- A \rightarrow 3\pi A$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
259±19± 6		AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$	

⁶ From a single-pole fit.⁷ In the $f_0(980)\pi$ wave.⁸ In the $f_0(500)\pi$ wave.⁹ From a fit to $J^{PC} = 0^{-+} f_0(980)\pi, f_0(1370)\pi$ waves.¹⁰ From a fit to $J^{PC} = 0^{-+} K_0^*(1430)K^-$ and $f_0(980)\pi^-$ waves.

NODE=M075W;LINKAGE=SP
 NODE=M075W;LINKAGE=C1
 NODE=M075W;LINKAGE=C2
 NODE=M075W;LINKAGE=AX
 NODE=M075W;LINKAGE=A

 $\pi(1800)$ DECAY MODES

NODE=M075215;NODE=M075

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^+ \pi^- \pi^-$	seen
Γ_2 $f_0(500)\pi^-$	seen
Γ_3 $f_0(980)\pi^-$	seen
Γ_4 $f_0(1370)\pi^-$	seen
Γ_5 $f_0(1500)\pi^-$	not seen
Γ_6 $\rho\pi^-$	not seen
Γ_7 $\eta\eta\pi^-$	seen
Γ_8 $a_0(980)\eta$	seen
Γ_9 $a_2(1320)\eta$	not seen
Γ_{10} $f_2(1270)\pi$	not seen
Γ_{11} $f_0(1370)\pi^-$	not seen
Γ_{12} $f_0(1500)\pi^-$	seen
Γ_{13} $\eta\eta'(958)\pi^-$	seen
Γ_{14} $K_0^*(1430)K^-$	seen
Γ_{15} $K^*(892)K^-$	not seen

DESIG=10;OUR EST;→ UNCHECKED ←
 DESIG=11;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=1
 DESIG=12
 DESIG=2
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=13
 DESIG=14
 DESIG=15
 DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=8;OUR EST;→ UNCHECKED ←
 DESIG=4
 DESIG=9

 $\pi(1800)$ BRANCHING RATIOS

NODE=M075220

$\Gamma(f_0(980)\pi^-)/\Gamma(f_0(500)\pi^-)$	Γ_3/Γ_2			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.44±0.08±0.38	¹¹ CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

NODE=M075R11
 NODE=M075R11

$\Gamma(f_0(980)\pi^-)/\Gamma(f_0(1370)\pi^-)$	Γ_3/Γ_4			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.7±1.3	¹² AMELIN	95B	VES	- 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$

NODE=M075R5
 NODE=M075R5

$\Gamma(f_0(1370)\pi^-)/\Gamma_{\text{total}}$	Γ_4/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	BELLINI	82	SPEC	- 40 $\pi^- A \rightarrow 3\pi A$

NODE=M075R1
 NODE=M075R1

$\Gamma(f_0(1500)\pi^-)/\Gamma_{\text{total}}$	Γ_5/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
not seen	CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

NODE=M075R12
 NODE=M075R12

$\Gamma(\rho\pi^-)/\Gamma_{\text{total}}$	Γ_6/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
not seen	BELLINI	82	SPEC	- 40 $\pi^- A \rightarrow 3\pi A$

NODE=M075R2
 NODE=M075R2

$\Gamma(\rho\pi^-)/\Gamma(f_0(980)\pi^-)$ Γ_6/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

NODE=M075R6
 NODE=M075R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.25		CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
<0.14	90	AMELIN	95B	VES -	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$

 $\Gamma(\eta\eta\pi^-)/\Gamma(\pi^+\pi^-\pi^-)$ Γ_7/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M075R8
 NODE=M075R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.5±0.1	1200	¹² AMELIN	96B	VES -	37 $\pi^- A \rightarrow \eta\eta\pi^- A$
---------	------	----------------------	-----	-------	--

 $\Gamma(a_2(1320)\eta)/\Gamma_{total}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M075R13
 NODE=M075R13

not seen EUGENIO 08 B852 18 $\pi^- p \rightarrow \eta\eta\pi^- p$

 $\Gamma(f_2(1270)\pi)/\Gamma_{total}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M075R14
 NODE=M075R14

not seen EUGENIO 08 B852 18 $\pi^- p \rightarrow \eta\eta\pi^- p$

 $\Gamma(f_0(1370)\pi^-)/\Gamma_{total}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M075R15
 NODE=M075R15

not seen EUGENIO 08 B852 18 $\pi^- p \rightarrow \eta\eta\pi^- p$

 $\Gamma(f_0(1500)\pi^-)/\Gamma(a_0(980)\eta)$ Γ_{12}/Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M075R7
 NODE=M075R7

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.48 ±0.17	4k	^{12,13} EUGENIO	08	B852 -	18 $\pi^- p \rightarrow \eta\eta\pi^- p$
0.030 ^{+0.014} _{-0.011}		¹² ANISOVICH	01B	SPEC 0	0.6-1.94 $p\bar{p} \rightarrow \eta\eta\pi^0\pi^0$
0.08 ±0.03	1200	^{12,14} AMELIN	96B	VES -	37 $\pi^- A \rightarrow \eta\eta\pi^- A$

 $\Gamma(\eta\eta'(958)\pi^-)/\Gamma(\eta\eta\pi^-)$ Γ_{13}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M075R10
 NODE=M075R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.29±0.07		¹² BELADIDZE	92C	VES -	36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$
0.3 ±0.1	426 ± 57	¹² BITYUKOV	91	VES -	36 $\pi^- C \rightarrow \pi^- \eta \eta C$

 $\Gamma(K_0^*(1430)K^-)/\Gamma_{total}$ Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

NODE=M075R4
 NODE=M075R4

seen BERDNIKOV 94 VES - 37 $\pi^- A \rightarrow K^+ K^- \pi^- A$

 $\Gamma(K^*(892)K^-)/\Gamma_{total}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

NODE=M075R9
 NODE=M075R9

not seen BERDNIKOV 94 VES - 37 $\pi^- A \rightarrow K^+ K^- \pi^- A$

¹¹ Assuming that $f_0(980)$ decays only to $\pi\pi$.

¹² Systematic errors not estimated.

¹³ From a single-pole fit.

¹⁴ Assuming that $f_0(1500)$ decays only to $\eta\eta$ and $a_0(980)$ decays only to $\eta\pi$.

NODE=M075R;LINKAGE=CK
 NODE=M075R5;LINKAGE=NS
 NODE=M075R7;LINKAGE=SP
 NODE=M075R7;LINKAGE=A

 $\pi(1800)$ REFERENCES

NODE=M075

ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)	REFID=52160
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>		REFID=48318
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=46910
		Translated from YAF 62 487.			
AMELIN	96B	PAN 59 976	D.V. Amelin <i>et al.</i>	(SERP, TBIL) IGJPC	REFID=44725
		Translated from YAF 59 1021.			
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44433
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)	REFID=44073
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)	REFID=43175
		Translated from YAF 55 2748.			
BITYUKOV	91	PL B268 137	S.I. Bitjukov <i>et al.</i>	(SERP, TBIL)	REFID=41749
BELLINI	82	PRL 48 1697	G. Bellini <i>et al.</i>	(MILA, BGNA, JINR)	REFID=21134

NODE=M038

$f_2(1810)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE
Needs confirmation.

NODE=M038

NODE=M038M

NODE=M038M

$f_2(1810)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1815 ± 12 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
1822 ⁺²⁹ ₋₂₄ ± 66 ₋₅₇	5.5k	1 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
1737 ± 9 ⁺¹⁹⁸ ₋₆₅		2 UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
1800 ± 30	40	ALDE	88D GAM4	300 $\pi^-p \rightarrow \pi^-p4\pi^0$
1806 ± 10	1600	ALDE	87 GAM4	100 $\pi^-p \rightarrow 4\pi^0n$
1870 ± 40		3 ALDE	86D GAM4	100 $\pi^-p \rightarrow \eta\eta n$
1857 ⁺³⁵ ₋₂₄		4 COSTA...	80 OMEG	10 $\pi^-p \rightarrow K^+K^-n$
1858 ⁺¹⁸ ₋₇₁		5 LONGACRE	86 RVUE	Compilation
1799 ± 15		6 CASON	82 STRC	8 $\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$


• • • We do not use the following data for averages, fits, limits, etc. • • •

- ¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.
- ² Breit-Wigner mass.
- ³ Seen in only one solution.
- ⁴ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.
- ⁵ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.
- ⁶ From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

NODE=M038M;LINKAGE=B

NODE=M038M;LINKAGE=UE
NODE=M038M;LINKAGE=F
NODE=M038M;LINKAGE=A

NODE=M038M;LINKAGE=L



$f_2(1810)$ WIDTH

NODE=M038W

NODE=M038W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
197 ± 22 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
229 ⁺⁵² ₋₄₂ ± 88 ₋₁₅₅	5.5k	7 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
228 ⁺²¹ ₋₂₀ ± 234 ₋₁₅₃		8 UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
160 ± 30	40	ALDE	88D GAM4	300 $\pi^-p \rightarrow \pi^-p4\pi^0$

190 ± 20	1600	ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$
250 ± 30		⁹ ALDE	86D	GAM4	100	$\pi^- p \rightarrow \eta\eta n$
185 ⁺¹⁰² ₋₁₃₉		¹⁰ COSTA...	80	OMEG	10	$\pi^- p \rightarrow K^+ K^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •


388 ⁺¹⁵ ₋₂₁		¹¹ LONGACRE	86	RVUE	Compilation	
280 ⁺⁴² ₋₃₅		¹² CASON	82	STRC	8	$\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$

- ⁷ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.
- ⁸ Breit-Wigner width.
- ⁹ Seen in only one solution.
- ¹⁰ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.
- ¹¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.
- ¹² From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

NODE=M038W;LINKAGE=B

NODE=M038W;LINKAGE=UE
 NODE=M038W;LINKAGE=F
 NODE=M038W;LINKAGE=A
 NODE=M038W;LINKAGE=L

NODE=M038W;LINKAGE=P1



$f_2(1810)$ DECAY MODES

NODE=M038215;NODE=M038

Mode	Fraction (Γ_i/Γ)	
Γ_1 $\pi\pi$		DESIG=2
Γ_2 $\eta\eta$	seen	DESIG=3
Γ_3 $4\pi^0$	seen	DESIG=4;OUR EST;→ UNCHECKED ←
Γ_4 $K^+ K^-$		DESIG=1
Γ_5 $\gamma\gamma$	seen	DESIG=5;OUR EST;→ UNCHECKED ←

$f_2(1810)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M038225

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_5/\Gamma$	
VALUE (eV)	DOCUMENT ID	TECN COMMENT
5.2^{+0.9+37.3}_{-0.8-4.5}	¹³ UEHARA	10A BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$

NODE=M038G01
 NODE=M038G01

¹³ Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

NODE=M038G01;LINKAGE=UE

$f_2(1810)$ BRANCHING RATIOS

NODE=M038220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
not seen	PROKOSHKIN	97	GAM2 $38 \pi^- p \rightarrow \pi^0 \pi^0 n$
$0.21^{+0.02}_{-0.03}$	14 LONGACRE	86	RVUE Compilation
0.44 ± 0.03	15 CASON	82	STRC $8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$

¹⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

¹⁵ Included in LONGACRE 86 global analysis.

NODE=M038R2
NODE=M038R2

NODE=M038R2;LINKAGE=L

NODE=M038R;LINKAGE=C

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen

seen	ABLIKIM	13N	BES3 PWA of $J/\psi \rightarrow \gamma \eta \eta$
------	---------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.008^{+0.028}_{-0.003}$	16 LONGACRE	86	RVUE Compilation
---------------------------	-------------	----	------------------

¹⁶ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R3
NODE=M038R3

NODE=M038R3;LINKAGE=L

 $\Gamma(\pi\pi)/\Gamma(4\pi^0)$ Γ_1/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.75	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$
-------	------	----	---

NODE=M038R4
NODE=M038R4 **$\Gamma(4\pi^0)/\Gamma(\eta\eta)$ Γ_3/Γ_2**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.8 ± 0.3	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$
---------------	------	----	---

NODE=M038R5
NODE=M038R5 **$\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_4/Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.003^{+0.019}_{-0.002}$	17 LONGACRE	86	RVUE Compilation
seen	COSTA...	80	OMEG $10 \pi^- p \rightarrow K^+ K^- n$

¹⁷ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R1
NODE=M038R1

NODE=M038R1;LINKAGE=L

 $f_2(1810)$ REFERENCES

NODE=M038

ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55387
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>		REFID=48580
PROKOSHKIN	97	SPD 42 117	Y.D. Prokoshkin <i>et al.</i>	(SERP)	REFID=45386
ALDE	88D	Translated from DANS 353 323. SJNP 47 810	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=44652
ALDE	87	Translated from YAF 47 1273. PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
COSTA...	80	NP B175 402	G. Costa de Beauregard <i>et al.</i>	(BARI, BONN+)	REFID=20737

X(1835)

$$I^G(J^{PC}) = ?(0^{-+})$$

NODE=M085

OMITTED FROM SUMMARY TABLE

Could be a superposition of two states, one with small width appearing as threshold enhancement in $p\bar{p}$, the other one with a larger width, decaying into $\pi^+\pi^-\eta'$ and $K_S^0 K_S^0 \eta$. For the former ABLIKIM 12D determine $J^{PC} = 0^{-+}$.

NODE=M085

X(1835) MASS

NODE=M085M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M085M

1835.8^{+4.0}_{-3.2} OUR AVERAGE

NEW

[1835.7^{+5.0}_{-3.2} MeV OUR 2015 AVERAGE]

1844 ± 9 ⁺¹⁶ ₋₂₅		ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
1836.5 ± 3.0 ^{+5.6} _{-2.1}	4265	¹ ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
1833.7 ± 6.1 ± 2.7	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1832 ⁺¹⁹ ₋₅ ± 26		² ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
1877.3 ± 6.3 ^{+3.4} _{-7.4}		³ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
1837 ⁺¹⁰ ₋₁₂ ⁺⁹ ₋₇	231	^{4,5} ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
1831 ± 7		^{5,6} ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p\bar{p}$
1859 ⁺³ ₋₁₀ ⁺⁵ ₋₂₅		⁵ BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

OCCUR=2

¹ From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0 \pi^+ \pi^- \eta'$.

NODE=M085M;LINKAGE=AI

² From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.

NODE=M085M;LINKAGE=AK

³ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$.

NODE=M085M;LINKAGE=BL

⁴ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

NODE=M085M;LINKAGE=AE

⁵ Evidence for a threshold enhancement in the $p\bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in $B^+ \rightarrow p\bar{p}K^+$, WANG 05A in $B^0 \rightarrow p\bar{p}K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p\bar{p}D^0$, DEL-AMO-SANCHEZ 12 in $B \rightarrow D(D^*)p\bar{p}(\pi)$, and WEI 08 in $B^+ \rightarrow p\bar{p}\pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p\bar{p}\gamma$.

NODE=M085M;LINKAGE=HF

⁶ From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A. Systematic errors not estimated.

NODE=M085M;LINKAGE=AB

X(1835) WIDTH

NODE=M085W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	-----	------	-------------	------	---------

NODE=M085W

112 ± 40 OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.
[99 ± 50 MeV OUR 2015 AVERAGE Scale factor = 2.8]


NEW

192 ⁺²⁰ ₋₁₇ ⁺⁶² ₋₄₃		ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
190 ± 9 ⁺³⁸ ₋₃₆	4265	¹ ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
67.7 ± 20.3 ± 7.7	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 76	90	² ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
57 ± 12 ⁺¹⁹ ₋₄		³ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
0 ⁺⁴⁴ ₋₀	231	^{4,5} ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
< 153	90	^{5,6} ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p\bar{p}$
< 30		⁵ BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

OCCUR=2

- ¹ From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.
- ² From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.
- ³ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$.
- ⁴ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.
- ⁵ Evidence for a threshold enhancement in the $p\bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in $B^+ \rightarrow p\bar{p}K^+$, WANG 05A in $B^0 \rightarrow p\bar{p}K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p\bar{p}D^0$, DEL-AMO-SANCHEZ 12 in $B \rightarrow D(D^*)p\bar{p}(\pi)$, and WEI 08 in $B^+ \rightarrow p\bar{p}\pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p\bar{p}\gamma$.
- ⁶ From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A. Systematic errors not estimated.

NODE=M085W;LINKAGE=A1
 NODE=M085W;LINKAGE=AK
 NODE=M085W;LINKAGE=BL
 NODE=M085W;LINKAGE=AE
 NODE=M085W;LINKAGE=HF
 NODE=M085W;LINKAGE=AB



X(1835) DECAY MODES

NODE=M085215;NODE=M085

Mode	Fraction (Γ_i/Γ)
Γ_1 $p\bar{p}$	seen
Γ_2 $\eta'\pi^+\pi^-$	seen
Γ_3 $\gamma\gamma$	
Γ_4 $K_S^0 K_S^0 \eta$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=4
 DESIG=5;OUR EVAL;→ UNCHECKED ←

X(1835) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M085225

$\Gamma(\eta'\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$\Gamma_2\Gamma_3/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT

NODE=M085G01
 NODE=M085G01

••• We do not use the following data for averages, fits, limits, etc. •••

<35.6	90	¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
<83	90	² ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

OCCUR=2

- ¹ From a two-resonance fit and constructive interference of the $\eta(1760)$ and $X(1835)$, a significance of 2.8σ .
- ² From a two-resonance fit and destructive interference of the $\eta(1760)$ and $X(1835)$, a significance of 2.8σ .

NODE=M085G01;LINKAGE=ZH

NODE=M085G01;LINKAGE=ZA

X(1835) BRANCHING RATIOS

NODE=M085220

$\Gamma(p\bar{p})/\Gamma(\eta'\pi^+\pi^-)$		Γ_1/Γ_2	
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M085R01
 NODE=M085R01

••• We do not use the following data for averages, fits, limits, etc. •••

0.333	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
-------	---------	----------	--

$\Gamma(\eta'\pi^+\pi^-)/\Gamma(K_S^0 K_S^0 \eta)$ Γ_2/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M085R00
 NODE=M085R00

••• We do not use the following data for averages, fits, limits, etc. •••

6.7±1.8	¹ ABLIKIM	15T	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
---------	----------------------	-----	------	--

¹ Using results from ABLIKIM 05R.

NODE=M085R00;LINKAGE=A

X(1835) REFERENCES

NODE=M085

ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56785
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.) JPC	REFID=54269
DEL-AMO-SA...	12	PR D85 092017	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=54286
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)	REFID=54763
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53684
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53931
ABLIKIM	10G	CPC 34 421	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55685
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
WEI	08	PL B659 80	J.-T. Wei <i>et al.</i>	(BELLE Collab.)	REFID=52086
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50993
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50985
AUBERT,B	05L	PR D72 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50827
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer		REFID=51038
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)	REFID=50651
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)	REFID=49473
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48690
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48980

NODE=M215

X(1840)

$$I^G(J^{PC}) = ?^?(???)$$

OMITTED FROM SUMMARY TABLE

X(1840) MASS

NODE=M215M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1842.2±4.2^{+7.1}_{-2.6}	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

NODE=M215M

X(1840) WIDTH

NODE=M215W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83±14±11	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

NODE=M215W

X(1840) DECAY MODES

NODE=M215215;NODE=M215

Mode	Fraction (Γ_i/Γ)
Γ_1 $3(\pi^+\pi^-)$	seen

DESIG=1

X(1840) BRANCHING RATIOS

NODE=M215225

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_1/Γ
seen	0.6k

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

NODE=M215R01
 NODE=M215R01

X(1840) REFERENCES

NODE=M215

ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BES III Collab.)
---------	-----	---------------	--------------------------	-------------------

REFID=55582

$a_1(1420)$

$$I^G(J^{PC}) = 1^-(1^{++})$$

NODE=M230

OMITTED FROM SUMMARY TABLE

 $a_1(1420)$ MASS

NODE=M230M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1414^{+15}_{-13}	¹ ADOLPH	15C	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

NODE=M230M

¹ Using the isobar model and partial-wave analysis with 88 waves.

NODE=M230M;LINKAGE=A

 $a_1(1420)$ WIDTH

NODE=M230W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
153^{+8}_{-23}	¹ ADOLPH	15C	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

NODE=M230W

¹ Using the isobar model and partial-wave analysis with 88 waves.

NODE=M230W;LINKAGE=A

 $a_1(1420)$ DECAY MODES

NODE=M230215;NODE=M230

Mode	Fraction (Γ_i/Γ)
Γ_1 $f_0(980)\pi$	seen

DESIG=1

 $a_1(1420)$ BRANCHING RATIOS

NODE=M230220

$\Gamma(f_0(980)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ADOLPH	15C	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

NODE=M230R01
NODE=M230R01¹ Using the isobar model and partial-wave analysis with 88 waves.

NODE=M230R01;LINKAGE=A

 $a_1(1420)$ REFERENCES

NODE=M230

ADOLPH 15C PRL 115 082001 C. Adolph *et al.* (COMPASS Collab.)

REFID=56790

NODE=M054

 $\phi_3(1850)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

 $\phi_3(1850)$ MASS

NODE=M054M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1854 ± 7 OUR AVERAGE				
1855 ± 10		ASTON	88E	LASS 11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
1870 ⁺³⁰ ₋₂₀	430	ARMSTRONG	82	OMEG 18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
1850 ± 10	123	ALHARRAN	81B	HBC 8.25 $K^- p \rightarrow K \bar{K} \Lambda$

NODE=M054M

 $\phi_3(1850)$ WIDTH

NODE=M054W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
87⁺²⁸₋₂₃ OUR AVERAGE	Error includes scale factor of 1.2.			
64 ± 31		ASTON	88E	LASS 11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
160 ⁺⁹⁰ ₋₅₀	430	ARMSTRONG	82	OMEG 18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
80 ⁺⁴⁰ ₋₃₀	123	ALHARRAN	81B	HBC 8.25 $K^- p \rightarrow K \bar{K} \Lambda$

NODE=M054W

$\phi_3(1850)$ DECAY MODES

NODE=M054215;NODE=M054

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $K\bar{K}^*(892)+c.c.$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

 $\phi_3(1850)$ BRANCHING RATIOS

NODE=M054220

$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(K\bar{K})$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
$0.55^{+0.85}_{-0.45}$	ASTON	88E LASS	11 $K^-p \rightarrow K^-K^+\Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
0.8 ± 0.4	ALHARRAN	81B HBC	8.25 $K^-p \rightarrow K\bar{K}\pi\Lambda$

NODE=M054R1
NODE=M054R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\phi_3(1850)$ REFERENCES

NODE=M054

ASTON	88E	PL B208 324	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)IGJPC
ARMSTRONG	82	PL 110B 77	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)JP
ALHARRAN	81B	PL 101B 357	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)

REFID=40577
REFID=21405
REFID=21702 $\eta_2(1870)$

$$I^G(J^{PC}) = 0^+(2^-+)$$

NODE=M101

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M101

 $\eta_2(1870)$ MASS

NODE=M101M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1842 ± 8 OUR AVERAGE				
1835 ± 12		BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
1844 ± 13		BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
1840 ± 25		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1875 ± 20 ± 35		ADOMEIT	96 CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
1881 ± 32 ± 40	26	KARCH	92 CBAL	$e^+e^- \rightarrow e^+e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1860 ± 5 ± 15		ANISOVICH	00E SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
1840 ± 15		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M101M

 $\eta_2(1870)$ WIDTH

NODE=M101W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
225 ± 14 OUR AVERAGE				
235 ± 22		BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
228 ± 23		BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
200 ± 40		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
200 ± 25 ± 45		ADOMEIT	96 CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
221 ± 92 ± 44	26	KARCH	92 CBAL	$e^+e^- \rightarrow e^+e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
250 ± 25 $^{+50}_{-35}$		ANISOVICH	00E SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
170 ± 40		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M101W

 $\eta_2(1870)$ DECAY MODES

NODE=M101225;NODE=M101

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi\pi$	
Γ_2 $a_2(1320)\pi$	
Γ_3 $f_2(1270)\eta$	
Γ_4 $a_0(980)\pi$	
Γ_5 $\gamma\gamma$	seen

DESIG=1

DESIG=4

DESIG=8

DESIG=2

DESIG=9

$\eta_2(1870)$ BRANCHING RATIOS $\Gamma(a_2(1320)\pi)/\Gamma(f_2(1270)\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_3
1.7 ± 0.4 OUR AVERAGE				
1.60 ± 0.40	¹ ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$	
20.4 ± 6.6	BARBERIS	00B	450 $p\bar{p} \rightarrow p_f \eta \pi^+ \pi^- p_s$	
4.1 ± 2.3	ADOMEIT	96	CBAR 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$	

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M101230

NODE=M101R2
NODE=M101R2

NODE=M101R2;LINKAGE=AN

 $\Gamma(a_2(1320)\pi)/\Gamma(a_0(980)\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_4
32.6 ± 12.6	BARBERIS	00B	450 $p\bar{p} \rightarrow p_f \eta \pi^+ \pi^- p_s$	

NODE=M101R4
NODE=M101R4 $\Gamma(a_0(980)\pi)/\Gamma(f_2(1270)\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_3
0.48 ± 0.45	² ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$	

² Reanalysis of ADOMEIT 96 and ANISOVICH 00E.NODE=M101R01
NODE=M101R01

NODE=M101R01;LINKAGE=AN

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
seen	KARCH	92	CBAL $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$	

NODE=M101R02
NODE=M101R02 $\eta_2(1870)$ REFERENCES

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)

NODE=M101

REFID=53631
REFID=47945
REFID=47958
REFID=47959
REFID=46606
REFID=45758
REFID=45202
REFID=42170

NODE=M185

 $\pi_2(1880)$

$$I^G(J^{PC}) = 1^-(2^-+)$$

 $\pi(1880)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1895 ± 16 OUR AVERAGE					
1929 ± 24 ± 18	4k	EUGENIO	08	B852	– 18 $\pi^- p \rightarrow \eta\eta\pi^- p$
1876 ± 11 ± 67	145k	LU	05	B852	– 18 $\pi^- p \rightarrow \omega\pi^- \pi^0 p$
2003 ± 88 ± 148	69k	KUHN	04	B852	– 18 $\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$
1880 ± 20		ANISOVICH	01B	SPEC 0	0.6–1.94 $\bar{p}p \rightarrow \eta\eta\pi^0\pi^0$

NODE=M185M

NODE=M185M

 $\pi(1880)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
235 ± 34 OUR AVERAGE					
323 ± 87 ± 43	4k	EUGENIO	08	B852	– 18 $\pi^- p \rightarrow \eta\eta\pi^- p$
146 ± 17 ± 62	145k	LU	05	B852	– 18 $\pi^- p \rightarrow \omega\pi^- \pi^0 p$
306 ± 132 ± 121	69k	KUHN	04	B852	– 18 $\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$
255 ± 45		ANISOVICH	01B	SPEC 0	0.6–1.94 $\bar{p}p \rightarrow \eta\eta\pi^0\pi^0$

NODE=M185W

NODE=M185W

 $\pi_2(1880)$ DECAY MODES

Mode	DESIG
Γ_1 $\eta\eta\pi^-$	DESIG=1
Γ_2 $a_0(980)\eta$	DESIG=2
Γ_3 $a_2(1320)\eta$	DESIG=3
Γ_4 $f_0(1500)\pi$	DESIG=4
Γ_5 $f_1(1285)\pi$	DESIG=5
Γ_6 $\omega\pi^- \pi^0$	DESIG=6

NODE=M185215;NODE=M185

 $\Gamma(a_2(1320)\eta)/\Gamma(f_1(1285)\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_5
22.7 ± 7.3	69k	KUHN	04	B852	– 18 $\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$	

NODE=M185R01
NODE=M185R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(f_0(1500)\pi)/\Gamma(a_0(980)\eta)$ Γ_4/Γ_2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.28^{+0.20}_{-0.15}$	¹ ANISOVICH	01B	SPEC	0	0.6–1.94 $\bar{p}p \rightarrow \eta\eta\pi^0\pi^0$
------------------------	------------------------	-----	------	---	--

¹ Systematic errors not estimated.

NODE=M185R02
NODE=M185R02

NODE=M185R02;LINKAGE=NS

 $\pi_2(1880)$ REFERENCES

EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>	

NODE=M185

REFID=52160
REFID=50459
REFID=49773
REFID=48318

NODE=M170

 $\rho(1900)$

$$I^G(J^{PC}) = 1^+(1^--)$$

OMITTED FROM SUMMARY TABLE

See our mini-review under the $\rho(1700)$.

NODE=M170

 $\rho(1900)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1909 \pm 17 \pm 25$	54	¹ AUBERT	08S	BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
1880 ± 30		AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
1860 ± 20		AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
1910 ± 10		^{2,3} FRABETTI	04	E687	$\gamma p \rightarrow 3\pi^+3\pi^-\rho$
1870 ± 10		ANTONELLI	96	SPEC	$e^+e^- \rightarrow$ hadrons

NODE=M170M

NODE=M170M

OCCUR=2

¹ From the fit with two resonances.

² From a fit with two resonances with the JACOB 72 continuum.

³ Supersedes FRABETTI 01.

NODE=M170M;LINKAGE=AU
NODE=M170M;LINKAGE=PI
NODE=M170M;LINKAGE=RS

 $\rho(1900)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$48 \pm 17 \pm 2$	54	⁴ AUBERT	08S	BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
130 ± 30		AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
160 ± 20		AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
37 ± 13		^{5,6} FRABETTI	04	E687	$\gamma p \rightarrow 3\pi^+3\pi^-\rho$
10 ± 5		ANTONELLI	96	SPEC	$e^+e^- \rightarrow$ hadrons

NODE=M170W

NODE=M170W

OCCUR=2

⁴ From the fit with two resonances.

⁵ From a fit with two resonances with the JACOB 72 continuum.

⁶ Supersedes FRABETTI 01.

NODE=M170W;LINKAGE=AU
NODE=M170W;LINKAGE=PI
NODE=M170W;LINKAGE=RS

 $\rho(1900)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$
 $\Gamma(\phi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$
 $\Gamma_4/\Gamma \times \Gamma_6/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.2 \pm 1.2 \pm 0.8$	54	⁷ AUBERT	08S	BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
-----------------------	----	---------------------	-----	------	---

⁷ From the fit with two resonances.

NODE=M170215

NODE=M170B01
NODE=M170B01

NODE=M170B01;LINKAGE=AU

 $\rho(1900)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 6π	seen
Γ_2 $3\pi^+3\pi^-$	seen
Γ_3 $2\pi^+2\pi^-\pi^0$	
Γ_4 $\phi\pi$	
Γ_5 hadrons	seen
Γ_6 e^+e^-	seen
Γ_7 $\bar{N}N$	not seen

NODE=M170225;NODE=M170

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=6

DESIG=7

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

$\rho(1900)$ BRANCHING RATIOS

NODE=M170230

$\Gamma(6\pi)/\Gamma_{total}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	8k	AKHMETSHIN 13	CMD3	$e^+e^- \rightarrow 3\pi^+3\pi^-$	
not seen		AGNELLO 02	OBLX	$\bar{n}p \rightarrow 3\pi^+2\pi^-\pi^0$	
seen		FRABETTI 01	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$	
seen		ANTONELLI 96	SPEC	$e^+e^- \rightarrow \text{hadrons}$	

NODE=M170R1
NODE=M170R1

$\rho(1900)$ REFERENCES

NODE=M170

AKHMETSHIN 13	PL B723 82	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
AUBERT 08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
FRABETTI 04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AGNELLO 02	PL B527 39	M. Agnello <i>et al.</i>	(OBELIX Collab.)
FRABETTI 01	PL B514 240	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ANTONELLI 96	PL B365 427	A. Antonelli <i>et al.</i>	(FENICE Collab.)
JACOB 72	PR D5 1847	M. Jacob, R. Slansky	

REFID=55370
REFID=52242
REFID=51047
REFID=49614
REFID=48576
REFID=48350
REFID=44633
REFID=49668

$f_2(1910)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M142

OMITTED FROM SUMMARY TABLE

We list here three different peaks with close masses and widths seen in the mass distributions of $\omega\omega$, $\eta\eta'$, and K^+K^- final states. ALDE 91B argues that they are of different nature.

NODE=M142

$f_2(1910)$ MASS

NODE=M142205

NODE=M142MX


$f_2(1910)$ $\omega\omega$ MODE

NODE=M142M2
NODE=M142M2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1903 ± 9 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
1890 ± 10	¹ AMELIN 06	VES	$36 \pi^- p \rightarrow \omega\omega n$
1934 ± 20	ANISOVICH 00J	SPEC	
1897 ± 11	BARBERIS 00F		$450 pp \rightarrow p_f\omega\omega p_s$
1924 ± 14	ALDE 90	GAM2	$38 \pi^- p \rightarrow \omega\omega n$

¹Supersedes BELADIDZE 92B.

NODE=M142M2;LINKAGE=AM



$f_2(1910) \eta\eta'$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1934±16	² BARBERIS	00A	450 $p\bar{p} \rightarrow p_f \eta\eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1911±10	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$
² Also compatible with $J^{PC}=1^-+$.			

NODE=M142M3
NODE=M142M3

NODE=M142M3;LINKAGE=KS

$f_2(1910) K^+ K^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1941±18	AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+ K^- \pi^0$

NODE=M142M4
NODE=M142M4

$f_2(1910)$ WIDTH

NODE=M142210


NODE=M142WX

$f_2(1910) \omega\omega$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
196±31 OUR AVERAGE	Error includes scale factor of 2.3. See the ideogram below.		
165±19	³ AMELIN	06	VES 36 $\pi^- p \rightarrow \omega\omega n$
271±25	ANISOVICH	00J	SPEC
202±32	BARBERIS	00F	450 $p\bar{p} \rightarrow p_f \omega\omega p_s$
91±50	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$
³ Supersedes BELADIDZE 92B.			

NODE=M142W2
NODE=M142W2

NODE=M142W2;LINKAGE=AM



$f_2(1910) \eta\eta'$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
141±41	⁴ BARBERIS	00A	450 $p\bar{p} \rightarrow p_f \eta\eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
90±35	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$
⁴ Also compatible with $J^{PC}=1^-+$.			

NODE=M142W3
NODE=M142W3

NODE=M142W3;LINKAGE=KS

$f_2(1910) K^+ K^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
120±40	AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+ K^- \pi^0$

NODE=M142W4
NODE=M142W4

$f_2(1910)$ DECAY MODES

NODE=M142215;NODE=M142

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^0\pi^0$	
Γ_2 K^+K^-	seen
Γ_3 $K_S^0K_S^0$	
Γ_4 $\eta\eta$	seen
Γ_5 $\omega\omega$	seen
Γ_6 $\eta\eta'$	seen
Γ_7 $\eta'\eta'$	
Γ_8 $\rho\rho$	seen
Γ_9 $a_2(1320)\pi$	seen
Γ_{10} $f_2(1270)\eta$	seen

DESIG=6

DESIG=11

DESIG=8

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=12;OUR EST;→ UNCHECKED ←

DESIG=13;OUR EST;→ UNCHECKED ←

 $f_2(1910)$ BRANCHING RATIOS

NODE=M142225

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	AMSLER	06	CBAR	1.64 $\bar{p}p \rightarrow K^+K^-\pi^0$	

NODE=M142R11

NODE=M142R11

$\Gamma(\pi^0\pi^0)/\Gamma(\eta\eta')$					Γ_1/Γ_6
VALUE	DOCUMENT ID	TECN	COMMENT		

NODE=M142R4

NODE=M142R4

••• We do not use the following data for averages, fits, limits, etc. •••

<0.1 ALDE 89 GAM2 $38\pi^-p \rightarrow \eta\eta'n$

$\Gamma(K_S^0K_S^0)/\Gamma(\eta\eta')$					Γ_3/Γ_6
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M142R7

NODE=M142R7

••• We do not use the following data for averages, fits, limits, etc. •••

<0.066 90 BALOSHIN 86 SPEC $40\pi^+p \rightarrow K_S^0K_S^0n$

$\Gamma(\eta\eta)/\Gamma(\eta\eta')$					Γ_4/Γ_6
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M142R6

NODE=M142R6

••• We do not use the following data for averages, fits, limits, etc. •••

<0.05 90 ALDE 91B GAM2 $38\pi^-p \rightarrow \eta\eta'n$

$\Gamma(\omega\omega)/\Gamma(\eta\eta')$					Γ_5/Γ_6
VALUE		DOCUMENT ID	COMMENT		

NODE=M142R10

NODE=M142R10

••• We do not use the following data for averages, fits, limits, etc. •••

2.6±0.6 BARBERIS 00F $450pp \rightarrow p_f\omega\omega p_S$

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		

NODE=M142R8

NODE=M142R8

••• We do not use the following data for averages, fits, limits, etc. •••

probably not seen BARBERIS 00A $450pp \rightarrow p_f\eta'\eta' p_S$ possibly seen BELADIDZE 92D VES $37\pi^-p \rightarrow \eta'\eta'n$

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$					Γ_8/Γ_5
VALUE	DOCUMENT ID	COMMENT			

NODE=M142R9

NODE=M142R9

••• We do not use the following data for averages, fits, limits, etc. •••

2.6±0.4 BARBERIS 00F $450pp \rightarrow p_f\omega\omega p_S$

$\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$					Γ_{10}/Γ_9
VALUE	DOCUMENT ID	TECN	COMMENT		

NODE=M142R12

NODE=M142R12

0.09±0.05 ⁵ ANISOVICH 11 SPEC 0.9–1.94 $p\bar{p}$ ⁵ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M142R12;LINKAGE=AN

f₂(1910) REFERENCES

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 69 715.		
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)
BELADIDZE	92D	ZPHY C57 13	G.M. Beladidze <i>et al.</i>	(VES Collab.)
ALDE	91B	SJNP 54 455	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
		Translated from YAF 54 751.		
Also		PL B276 375	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ALDE	89	PL B216 447	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
Also		SJNP 48 1035	D.M. Alde <i>et al.</i>	(BELG, SERP, LANL, LAPP)
		Translated from YAF 48 1724.		
BALOSHIN	86	SJNP 43 959	O.N. Baloshin <i>et al.</i>	(ITEP)
		Translated from YAF 43 1487.		

NODE=M142

REFID=53631
REFID=51574REFID=51136
REFID=47945
REFID=47950
REFID=47957
REFID=47962
REFID=45202
REFID=42172
REFID=43309
REFID=41844REFID=41911
REFID=40935
REFID=40727
REFID=44697

REFID=40734

a₀(1950)

$$I^G(J^{PC}) = 1^-(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. Seen in $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$ by LEES 16A with significance 2.5σ in $K_S^0 K^\pm \pi^\mp$ and 4.2σ in $K^+ K^- \pi^0$. Spin-2 explanation ($a_2(1950)$) is not compatible with data.

NODE=M227

NODE=M227

a₀(1950) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1931 ± 14 ± 22	12k	^{1,2} LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1949 ± 32 ± 76	8k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
1927 ± 15 ± 23	4k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$
¹ From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating width.				
² Weighted average of the $K_S^0 K^\pm$ and $K^+ K^-$ decay modes.				

NODE=M227M

NODE=M227M

OCCUR=3

OCCUR=2

NODE=M227M;LINKAGE=A

NODE=M227M;LINKAGE=B

a₀(1950) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
271 ± 22 ± 29	12k	^{1,2} LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
265 ± 36 ± 110	8k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
274 ± 28 ± 30	4k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$
¹ From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating mass.				
² Weighted average of the $K_S^0 K^\pm$ and $K^+ K^-$ decay modes.				

NODE=M227W

NODE=M227W

OCCUR=3

OCCUR=2

NODE=M227W;LINKAGE=A

NODE=M227W;LINKAGE=B

a₀(1950) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen

NODE=M227215;NODE=M227

DESIG=1

a₀(1950) BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
¹ From a model-independent partial wave analysis.				

NODE=M227225

NODE=M227R01
NODE=M227R01

NODE=M227R01;LINKAGE=A

a₀(1950) REFERENCES

LEES	16A	PR D93 012005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
------	-----	---------------	-------------------------	-----------------

NODE=M227

REFID=57125

$f_2(1950)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M135

$f_2(1950)$ MASS

NODE=M135M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1944 ± 12 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
1930 ± 25	¹ BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
2010 ± 25	ANISOVICH	00J	SPEC
1940 ± 50	BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
1980 ± 22	² BARBERIS	00C	450 $pp \rightarrow pp4\pi$
1940 ± 22	³ BARBERIS	00C	450 $pp \rightarrow pp2\pi2\pi^0$
1980 ± 50	ANISOVICH	99B	SPEC 1.35-1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
1960 ± 30	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1918 ± 12	ANTINORI	95	OMEG 300,450 $pp \rightarrow pp2(\pi^+ \pi^-)$

NODE=M135M

NODE=M135M

• • • We do not use the following data for averages, fits, limits, etc. • • •

2038 ⁺¹³⁺¹² ₋₁₁₋₇₃	⁴ UEHARA	09	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1980 ± 2 ± 14	ABE	04	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
1867 ± 46	⁵ AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
~ 1990	⁶ OAKDEN	94	RVUE 0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
1950 ± 15	⁷ ASTON	91	LASS 11 $K^- p \rightarrow \Lambda K \bar{K} \pi \pi$

OCCUR=2

¹ First solution, PWA is ambiguous.

² Decaying into $\pi^+ \pi^- 2\pi^0$.

³ Decaying into $2(\pi^+ \pi^-)$.

⁴ Taking into account $f_4(2050)$.

⁵ T-matrix pole.

⁶ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

⁷ Cannot determine spin to be 2.

NODE=M135M;LINKAGE=BI

NODE=M135M;LINKAGE=A4


NODE=M135M;LINKAGE=B4

NODE=M135M;LINKAGE=UE

NODE=M135M;LINKAGE=TT

NODE=M135M;LINKAGE=BB

NODE=M135M;LINKAGE=A



$f_2(1950)$ WIDTH

NODE=M135W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
472 ± 18 OUR AVERAGE			
450 ± 50	⁸ BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
495 ± 35	ANISOVICH	00J	SPEC
380 ⁺¹²⁰ ₋₉₀	BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
520 ± 50	⁹ BARBERIS	00C	450 $pp \rightarrow pp4\pi$

NODE=M135W

485 ± 55	¹⁰ BARBERIS	00C	450 $pp \rightarrow pp4\pi$
500 ± 100	ANISOVICH	99B SPEC	1.35–1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
460 ± 40	BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
390 ± 60	ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
441 ⁺ ₂₅ 27 ⁺ ₁₉₂	¹¹ UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
297 ± 12 ± 6	ABE	04 BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
385 ± 58	¹² AMSLER	02 CBAR	0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
~ 100	¹³ OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
250 ± 50	¹⁴ ASTON	91 LASS	11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$
⁸ First solution, PWA is ambiguous.			
⁹ Decaying into $\pi^+\pi^-2\pi^0$.			
¹⁰ Decaying into $2(\pi^+\pi^-)$.			
¹¹ Taking into account $f_4(2050)$.			
¹² T-matrix pole.			
¹³ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.			
¹⁴ Cannot determine spin to be 2.			

OCCUR=2

NODE=M135W;LINKAGE=BI
 NODE=M135W;LINKAGE=A4
 NODE=M135W;LINKAGE=B4
 NODE=M135W;LINKAGE=UE
 NODE=M135W;LINKAGE=TT
 NODE=M135W;LINKAGE=BB

NODE=M135W;LINKAGE=A

NODE=M135215;NODE=M135

 $f_2(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\bar{K}^*(892)$	seen
Γ_2 $\pi\pi$	
Γ_3 $\pi^+\pi^-$	seen
Γ_4 $\pi^0\pi^0$	seen
Γ_5 4π	seen
Γ_6 $\pi^+\pi^-\pi^+\pi^-$	
Γ_7 $a_2(1320)\pi$	
Γ_8 $f_2(1270)\pi\pi$	
Γ_9 $\eta\eta$	seen
Γ_{10} $K\bar{K}$	seen
Γ_{11} $\gamma\gamma$	seen
Γ_{12} $p\bar{p}$	seen

DESIG=1
 DESIG=11
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=10;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=3
 DESIG=4
 DESIG=5
 DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=8;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←
 DESIG=12

 $f_2(1950)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{10}\Gamma_{11}/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT

NODE=M135225

NODE=M135G1
 NODE=M135G1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

122 ± 4 ± 26	¹⁵ ABE	04 BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
--------------	-------------------	---------	--

¹⁵ Assuming spin 2.

NODE=M135G1;LINKAGE=AB

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_{11}/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M135G2
 NODE=M135G2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

162 ⁺ ₄₂ 69 ⁺ ₂₀₄	¹⁶ UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
---	----------------------	---------	--

¹⁶ Taking into account $f_4(2050)$.

NODE=M135G2;LINKAGE=UE

 $f_2(1950)$ BRANCHING RATIOS

NODE=M135220

$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT

NODE=M135R1
 NODE=M135R1

seen ASTON 91 LASS 0 11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$	Γ_7/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M135R3
 NODE=M135R3

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

not seen	BARBERIS	00B	450 $pp \rightarrow p_f\eta\pi^+\pi^-p_s$
not seen	BARBERIS	00C	450 $pp \rightarrow p_f4\pi p_s$
possibly seen	BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$

$\Gamma(\eta\eta)/\Gamma(4\pi)$

VALUE	CL%	DOCUMENT ID	COMMENT
-------	-----	-------------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<5.0 \times 10^{-3}$	90	BARBERIS	00E 450 $p\bar{p} \rightarrow p_f \eta \eta p_S$
-----------------------	----	----------	--

 Γ_9/Γ_5

NODE=M135R5
NODE=M135R5

 $\Gamma(\eta\eta)/\Gamma(\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.14 ± 0.05	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
--------------------	--------	----	--

 Γ_9/Γ_3

NODE=M135R6
NODE=M135R6

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	111	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$
-------------	-----	-----------	----	---

 Γ_{12}/Γ

NODE=M135R07
NODE=M135R07

 $f_2(1950)$ REFERENCES

ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+) JP
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>	(LASS Collab.)

NODE=M135

REFID=53525
REFID=52761
REFID=50780

REFID=49650
REFID=48580
REFID=47950
REFID=47426
REFID=47958
REFID=47959
REFID=47961
REFID=46886
REFID=45758
REFID=45212
REFID=44437
REFID=45210
REFID=41746

 $\rho_3(1990)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

OMITTED FROM SUMMARY TABLE

NODE=M167

 $\rho_3(1990)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1982 ± 14	¹ ANISOVICH	02	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2007	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

¹From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167M

NODE=M167M

NODE=M167M;LINKAGE=AY

 $\rho_3(1990)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

188 ± 24	² ANISOVICH	02	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 287	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

²From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167W

NODE=M167W

NODE=M167W;LINKAGE=AY

 $\rho_3(1990)$ REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)

NODE=M167

REFID=48828
REFID=48327
REFID=48349
REFID=47950
REFID=44103

$f_2(2010)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M106

 $f_2(2010)$ MASS

NODE=M106M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2011⁺₋₇₆	¹ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2005 \pm 12	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1980 \pm 20	² BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2050 ⁺ ₋₅₀	ETKIN	85	MPS 22 $\pi^- p \rightarrow 2\phi n$
2120 ⁺ ₋₁₂₀	LINDENBAUM	84	RVUE
2160 \pm 50	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

NODE=M106M

¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi 2^{++} S_2$, D_2 , and D_0 is 98_{-3}^{+1} , 0_{-0}^{+1} , and 2_{-1}^{+2} , respectively.

NODE=M106M;LINKAGE=C

² Statistically very weak, only 1.4 s.d.

NODE=M106M;LINKAGE=E

 $f_2(2010)$ WIDTH

NODE=M106W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
202⁺₋₆₂	³ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
209 \pm 32	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
145 \pm 50	⁴ BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
200 ⁺ ₋₅₀	ETKIN	85	MPS 22 $\pi^- p \rightarrow 2\phi n$
300 ⁺ ₋₅₀	LINDENBAUM	84	RVUE
310 \pm 70	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

NODE=M106W

³ Includes data of ETKIN 85.

NODE=M106W;LINKAGE=C

⁴ Statistically very weak, only 1.4 s.d.

NODE=M106W;LINKAGE=E

 $f_2(2010)$ DECAY MODES

NODE=M106215;NODE=M106

Mode	Fraction (Γ_i/Γ)
Γ_1 $\phi\phi$	seen
Γ_2 $K\bar{K}$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2 **$f_2(2010)$ BRANCHING RATIOS**

NODE=M106230

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$	

NODE=M106R01
NODE=M106R01 **$f_2(2010)$ REFERENCES**

NODE=M106

VLADIMIRSK... 06	PAN 69 493 Translated from YAF 69 515.	V.V. Vladimisky <i>et al.</i>	(ITEP, Moscow)
BOLONKIN 88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ETKIN 88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)
ETKIN 85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)
LINDENBAUM 84	CNPP 13 285	S.J. Lindenbaum	(CUNY)
ETKIN 82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)
Also	Brighton Conf. 351	S.J. Lindenbaum	(BNL, CUNY)

REFID=51191

REFID=40580
REFID=40285
REFID=21871
REFID=21869
REFID=21866
REFID=21867

$f_0(2020)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M156

NODE=M156

NODE=M156M

NODE=M156M

NODE=M156M;LINKAGE=PC

NODE=M156M;LINKAGE=PP

NODE=M156M;LINKAGE=ST

NODE=M156W

NODE=M156W

NODE=M156W;LINKAGE=PC

NODE=M156W;LINKAGE=PP

NODE=M156W;LINKAGE=ST

NODE=M156215;NODE=M156

NODE=M156220

NODE=M156R1
NODE=M156R1NODE=M156R01
NODE=M156R01

NODE=M156

REFID=51063
REFID=47950
REFID=47959
REFID=47962
REFID=46605
REFID=46914
REFID=45758 **$f_0(2020)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1992±16		^{1,2} BARBERIS	00C	450 $p p \rightarrow p_f 4\pi p_s$
2037±8	80k	³ UMAN	06 E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
2040±38		ANISOVICH	00J SPEC	
2010±60		ALDE	98 GAM4	100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
2020±35		BARBERIS	97B OMEG	450 $p p \rightarrow p p 2(\pi^+ \pi^-)$

¹ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.² T-matrix pole.³ Statistical error only. **$f_0(2020)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
442±60		^{4,5} BARBERIS	00C	450 $p p \rightarrow p_f 4\pi p_s$
296±17	80k	⁶ UMAN	06 E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
405±40		ANISOVICH	00J SPEC	
240±100		ALDE	98 GAM4	100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
410±50		BARBERIS	97B OMEG	450 $p p \rightarrow p p 2(\pi^+ \pi^-)$

⁴ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.⁵ T-matrix pole.⁶ Statistical error only. **$f_0(2020)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi\pi$	seen
Γ_2 $\pi^0\pi^0$	seen
Γ_3 $\rho\rho$	seen
Γ_4 $\omega\omega$	seen
Γ_5 $\eta\eta$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5

 $f_0(2020)$ BRANCHING RATIOS

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$	Γ_3/Γ_4	
VALUE	DOCUMENT ID	COMMENT
~ 3	BARBERIS	00F 450 $p p \rightarrow p_f \omega \omega p_s$

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	Γ_5/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	UMAN	06 E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$

 $f_0(2020)$ REFERENCES

UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446		
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)

$a_4(2040)$

$$I^G(J^{PC}) = 1^-(4^{++})$$

NODE=M017

 $a_4(2040)$ MASS

NODE=M017M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1995^{+10}_{-8}					OUR AVERAGE Error includes scale factor of 1.1.
1900^{+80}_{-20}		ADOLPH	15	COMP	$191 \pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
$1885 \pm 13^{+50}_{-2}$	420k	ALEKSEEV	10	COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$1985 \pm 10 \pm 13$	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
$1996 \pm 25 \pm 43$		CHUNG	02	B852	$18.3 \pi^- p \rightarrow 3\pi p$
2005^{+25}_{-45}		¹ ANISOVICH	01F	SPEC	$2.0 \bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
$2000 \pm 40^{+60}_{-20}$		IVANOV	01	B852	$18 \pi^- p \rightarrow \eta' \pi^- p$
$1944 \pm 8 \pm 50$		² AMELIN	99	VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
2010 ± 20		³ DONSKOV	96	GAM2 0	$38 \pi^- p \rightarrow \eta \pi^0 n$
2040 ± 30		⁴ CLELAND	82B	SPEC \pm	$50 \pi p \rightarrow K_S^0 K^\pm p$
2030 ± 50		⁵ CORDEN	78C	OMEG 0	$15 \pi^- p \rightarrow 3\pi n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2004 ± 6	80k	⁶ UMAN	06	E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
1903 ± 10		⁷ BALDI	78	SPEC -	$10 \pi^- p \rightarrow p K_S^0 K^-$

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.² May be a different state.³ From a simultaneous fit to the G_+ and G_0 wave intensities.⁴ From an amplitude analysis.⁵ $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.⁶ Statistical error only.⁷ From a fit to the Y_8^0 moment. Limited by phase space.

NODE=M017M;LINKAGE=AN
 NODE=M017M;LINKAGE=DM
 NODE=M017M;LINKAGE=A
 NODE=M017M;LINKAGE=C
 NODE=M017M;LINKAGE=M
 NODE=M017M;LINKAGE=ST
 NODE=M017M;LINKAGE=Y

 $a_4(2040)$ WIDTH


NODE=M017W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
257^{+25}_{-23}					OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.
300^{+80}_{-100}		ADOLPH	15	COMP	$191 \pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
$294 \pm 25^{+46}_{-19}$	420k	ALEKSEEV	10	COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$231 \pm 30 \pm 46$	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
$298 \pm 81 \pm 85$		CHUNG	02	B852	$18.3 \pi^- p \rightarrow 3\pi p$
180 ± 30		¹ ANISOVICH	01F	SPEC	$2.0 \bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
$350 \pm 100^{+70}_{-50}$		IVANOV	01	B852	$18 \pi^- p \rightarrow \eta' \pi^- p$
$324 \pm 26 \pm 75$		² AMELIN	99	VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
370 ± 80		³ DONSKOV	96	GAM2 0	$38 \pi^- p \rightarrow \eta \pi^0 n$
380 ± 150		⁴ CLELAND	82B	SPEC \pm	$50 \pi p \rightarrow K_S^0 K^\pm p$
510 ± 200		⁵ CORDEN	78C	OMEG 0	$15 \pi^- p \rightarrow 3\pi n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
401 ± 16	80k	⁶ UMAN	06	E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
166 ± 43		⁷ BALDI	78	SPEC -	$10 \pi^- p \rightarrow p K_S^0 K^-$

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.² May be a different state.³ From a simultaneous fit to the G_+ and G_0 wave intensities.⁴ From an amplitude analysis.⁵ $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.⁶ Statistical error only.⁷ From a fit to the Y_8^0 moment. Limited by phase space.

NODE=M017W

NODE=M017W;LINKAGE=AN
 NODE=M017W;LINKAGE=DM
 NODE=M017W;LINKAGE=A
 NODE=M017W;LINKAGE=C
 NODE=M017W;LINKAGE=M
 NODE=M017W;LINKAGE=ST
 NODE=M017W;LINKAGE=Y



a₄(2040) DECAY MODES

NODE=M017215;NODE=M017

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $\pi^+\pi^-\pi^0$	seen
Γ_3 $\rho\pi$	seen
Γ_4 $f_2(1270)\pi$	seen
Γ_5 $\omega\pi^-\pi^0$	seen
Γ_6 $\omega\rho$	seen
Γ_7 $\eta\pi$	seen
Γ_8 $\eta'(958)\pi$	seen

DESIG=1
DESIG=2
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←
DESIG=7;OUR EST;→ UNCHECKED ←
DESIG=8
DESIG=3
DESIG=4;OUR EST;→ UNCHECKED ←

a₄(2040) BRANCHING RATIOS

NODE=M017220

$\Gamma(K\bar{K})/\Gamma_{total}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	BALDI	78	SPEC	± 10 $\pi^- p \rightarrow K_S^0 K^- p$

NODE=M017R1
NODE=M017R1

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	CORDEN	78C	OMEG	0 15 $\pi^- p \rightarrow 3\pi n$

NODE=M017R2
NODE=M017R2

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$ Γ_3/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
1.1±0.2±0.2	CHUNG	02 B852	18.3 $\pi^- p \rightarrow 3\pi p$

NODE=M017R4
NODE=M017R4

$\Gamma(\eta\pi)/\Gamma_{total}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	DONSKOV	96 GAM2	0	38 $\pi^- p \rightarrow \eta\pi^0 n$

NODE=M017R3
NODE=M017R3

$\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$ Γ_8/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
0.23±0.07	ADOLPH	15 COMP	191 $\pi^- p \rightarrow \eta^{(\prime)}\pi^- p$

NODE=M017R01
NODE=M017R01

$\Gamma(\omega\rho)/\Gamma_{total}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	145k	LU	05 B852	18 $\pi^- p \rightarrow \omega\pi^-\pi^0 p$

NODE=M017R5
NODE=M017R5

$a_4(2040)$ REFERENCES

ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48317
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=46910
		Translated from YAF 62 487.			
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>		REFID=46902
DONSKOV	96	PAN 59 982	S.V. Donskov <i>et al.</i>	(GAMS Collab.) IGJPC	REFID=45207
		Translated from YAF 59 1027.			
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
BALDI	78	PL 74B 413	R. Baldi <i>et al.</i>	(GEVA) JP	REFID=21783
CORDEN	78C	NP B136 77	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20859

NODE=M016

 $f_4(2050)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

 $f_4(2050)$ MASS

NODE=M016M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2018±11 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
1960±15		AMELIN	06 VES	$36 \pi^- p \rightarrow \omega \omega n$
2005±10		¹ BINON	05 GAMS	$33 \pi^- p \rightarrow \eta \eta n$
1998±15		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
2060±20		ALDE	90 GAM2	$38 \pi^- p \rightarrow \omega \omega n$
2038±30		AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
2086±15		BALTRUSAIT..	87 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
2000±60		ALDE	86D GAM4	$100 \pi^- p \rightarrow n2\eta$
2020±20	40k	² BINON	84B GAM2	$38 \pi^- p \rightarrow n2\pi^0$
2015±28		³ CASON	82 STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
2031 ⁺²⁵ ₋₃₆		ETKIN	82B MPS	$23 \pi^- p \rightarrow n2K_S^0$
2020±30	700	APEL	75 NICE	$40 \pi^- p \rightarrow n2\pi^0$
2050±25		BLUM	75 ASPK	$18.4 \pi^- p \rightarrow nK^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1966±25		⁴ ANISOVICH	09 RVUE	$0.0 \bar{p} p, \pi N$
1885 ⁺¹⁴⁺²¹⁸ ₋₁₃₋₂₅		⁵ UEHARA	09 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
2018± 6		ANISOVICH	00J SPEC	$2.0 \bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$ $\eta \eta, \eta \eta', \pi \pi$
~ 2000		⁶ MARTIN	98 RVUE	$N\bar{N} \rightarrow \pi \pi$
~ 2010		⁷ MARTIN	97 RVUE	$\bar{N}N \rightarrow \pi \pi$
~ 2040		⁸ OAKDEN	94 RVUE	$0.36-1.55 \bar{p} p \rightarrow \pi \pi$
~ 1990		⁹ OAKDEN	94 RVUE	$0.36-1.55 \bar{p} p \rightarrow \pi \pi$
1978± 5		¹⁰ ALPER	80 CNTR	$62 \pi^- p \rightarrow K^+ K^- n$
2040±10		¹⁰ ROZANSKA	80 SPRK	$18 \pi^- p \rightarrow p \bar{p} n$
1935±13		¹⁰ CORDEN	79 OMEG	$12-15 \pi^- p \rightarrow n2\pi$
1988± 7		EVANGELIS...	79B OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
1922±14		¹¹ ANTIPOV	77 CIBS	$25 \pi^- p \rightarrow p3\pi$


OCCUR=2

¹ From the first PWA solution.² From a partial-wave analysis of the data.³ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.⁴ K matrix pole.⁵ Taking into account the $f_2(1950)$. Helicity-2 production favored.⁶ Energy-dependent analysis.⁷ Single energy analysis.⁸ From solution A of amplitude analysis of data on $\bar{p} p \rightarrow \pi \pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.⁹ From solution B of amplitude analysis of data on $\bar{p} p \rightarrow \pi \pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.¹⁰ $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.¹¹ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

NODE=M016M;LINKAGE=BI
 NODE=M016M;LINKAGE=NN
 NODE=M016M;LINKAGE=NN
 NODE=M016M;LINKAGE=KM
 NODE=M016M;LINKAGE=UE
 NODE=M016M;LINKAGE=RB
 NODE=M016M;LINKAGE=BR
 NODE=M016M;LINKAGE=B

NODE=M016M;LINKAGE=BB

NODE=M016M;LINKAGE=M
 NODE=M016M;LINKAGE=T



f₄(2050) WIDTH

NODE=M016W

NODE=M016W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
237± 18 OUR AVERAGE		Error includes scale factor of 1.9. See the ideogram below.		
290± 20		AMELIN	06	VES 36 $\pi^- p \rightarrow \omega \omega n$
340± 80		12 BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
395± 40		ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
170± 60		ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega \omega n$
304± 60		AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
210± 63		BALTRUSAIT..	87	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
400±100		ALDE	86D	GAM4 100 $\pi^- p \rightarrow n 2 \eta$
240± 40	40k	13 BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2 \pi^0$
190± 14		DENNEY	83	LASS 10 $\pi^+ n/\pi^+ p$
186 ⁺¹⁰³ ₋₅₈		14 CASON	82	STRC 8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
305 ⁺³⁶ ₋₁₁₉		ETKIN	82B	MPS 23 $\pi^- p \rightarrow n 2 K_S^0$
180± 60	700	APEL	75	NICE 40 $\pi^- p \rightarrow n 2 \pi^0$
225 ⁺¹²⁰ ₋₇₀		BLUM	75	ASPK 18.4 $\pi^- p \rightarrow n K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

260± 40		15 ANISOVICH	09	RVUE 0.0 $\bar{p} p, \pi N$
453± 20 ⁺³¹ ₋₁₂₉		16 UEHARA	09	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
182± 7		ANISOVICH	00J	SPEC 2.0 $\bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$ $\eta \eta, \eta \eta', \pi \pi$
~ 170		17 MARTIN	98	RVUE $N \bar{N} \rightarrow \pi \pi$
~ 200		18 MARTIN	97	RVUE $\bar{N} N \rightarrow \pi \pi$
~ 60		19 OAKDEN	94	RVUE 0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
~ 80		20 OAKDEN	94	RVUE 0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
243± 16		21 ALPER	80	CNTR 62 $\pi^- p \rightarrow K^+ K^- n$
140± 15		21 ROZANSKA	80	SPRK 18 $\pi^- p \rightarrow p \bar{p} n$
263± 57		21 CORDEN	79	OMEG 12-15 $\pi^- p \rightarrow n 2 \pi$
100± 28		EVANGELIS...	79B	OMEG 10 $\pi^- p \rightarrow K^+ K^- n$
107± 56		22 ANTIPOV	77	CIBS 25 $\pi^- p \rightarrow p 3 \pi$

OCCUR=2

12 From the first PWA solution.

13 From a partial-wave analysis of the data.

14 From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2 \pi^0$.

15 K matrix pole.

16 Taking into account the $f_2(1950)$. Helicity-2 production favored.

17 Energy-dependent analysis.

18 Single energy analysis.


NODE=M016W;LINKAGE=BI
 NODE=M016W;LINKAGE=N
 NODE=M016W;LINKAGE=NN
 NODE=M016W;LINKAGE=KM
 NODE=M016W;LINKAGE=UE
 NODE=M016W;LINKAGE=RB
 NODE=M016W;LINKAGE=BR

- 19 From solution A of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
- 20 From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
- 21 $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.
- 22 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

NODE=M016W;LINKAGE=BW

NODE=M016W;LINKAGE=BB

NODE=M016W;LINKAGE=M
NODE=M016W;LINKAGE=T



$f_4(2050)$ DECAY MODES

NODE=M016215;NODE=M016

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega\omega$	seen
Γ_2 $\pi\pi$	$(17.0 \pm 1.5) \%$
Γ_3 $K\bar{K}$	$(6.8^{+3.4}_{-1.8}) \times 10^{-3}$
Γ_4 $\eta\eta$	$(2.1 \pm 0.8) \times 10^{-3}$
Γ_5 $4\pi^0$	$< 1.2 \%$
Γ_6 $\gamma\gamma$	
Γ_7 $a_2(1320)\pi$	seen

DESIG=6
DESIG=1
DESIG=2
DESIG=3
DESIG=5
DESIG=4
DESIG=7

$f_4(2050)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M016220

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_6/\Gamma$
VALUE (keV) CL% DOCUMENT ID TECN COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••	
<0.29 95 ALTHOFF 85B TASS $\gamma\gamma \rightarrow K\bar{K}\pi$	

NODE=M016G2
NODE=M016G2

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_6/\Gamma$
VALUE (eV) CL% EVTS DOCUMENT ID TECN COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••	
$23.1^{+3.6+70.5}_{-3.3-15.6}$ 23 UEHARA 09 BELL $10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	
<1100 95 13 ± 4 OEST 90 JADE $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	

NODE=M016G3
NODE=M016G3

²³ Taking into account the $f_2(1950)$. Helicity-2 production favored.

NODE=M016G3;LINKAGE=UE

$f_4(2050)$ BRANCHING RATIOS

NODE=M016225

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE DOCUMENT ID TECN COMMENT	
seen AMELIN 06 VES $36 \pi^-p \rightarrow \omega\omega n$	
••• We do not use the following data for averages, fits, limits, etc. •••	
not seen BARBERIS 00F $450 pp \rightarrow pf\omega\omega p_S$	

NODE=M016R7
NODE=M016R7

$\Gamma(\omega\omega)/\Gamma(\pi\pi)$				Γ_1/Γ_2	
VALUE	DOCUMENT ID	TECN	COMMENT		
1.5±0.3	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$		NODE=M016R5 NODE=M016R5
$\Gamma(\pi\pi)/\Gamma_{total}$				Γ_2/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
0.170±0.015 OUR AVERAGE					NODE=M016R1 NODE=M016R1
0.18 ±0.03	24 BINON	83C	GAM2 38 $\pi^- p \rightarrow n4\gamma$		
0.16 ±0.03	24 CASON	82	STRC 8 $\pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$		
0.17 ±0.02	24 CORDEN	79	OMEG 12-15 $\pi^- p \rightarrow n2\pi$		
²⁴ Assuming one pion exchange.					NODE=M016R1;LINKAGE=A
$\Gamma(K\bar{K})/\Gamma(\pi\pi)$				Γ_3/Γ_2	
VALUE	DOCUMENT ID	TECN	COMMENT		
0.04^{+0.02}_{-0.01}	ETKIN	82B	MPS 23 $\pi^- p \rightarrow n2K_S^0$		NODE=M016R2 NODE=M016R2
$\Gamma(\eta\eta)/\Gamma_{total}$				Γ_4/Γ	
VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT		
2.1±0.8	ALDE	86D	GAM4 100 $\pi^- p \rightarrow n4\gamma$		NODE=M016R3 NODE=M016R3
$\Gamma(4\pi^0)/\Gamma_{total}$				Γ_5/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
<0.012	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$		NODE=M016R4 NODE=M016R4
$\Gamma(a_2(1320)\pi)/\Gamma_{total}$				Γ_7/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$		NODE=M016R6 NODE=M016R6

$f_4(2050)$ REFERENCES

				NODE=M016
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	REFID=52719
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.) REFID=52761
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.) REFID=51574
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	REFID=50780
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.) REFID=47432
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	REFID=47950
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.) REFID=47962
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.) REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.) REFID=46914
MARTIN	98	PR C57 3492	B.R. Martin <i>et al.</i>	REFID=46373
MARTIN	97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH) REFID=45685
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD) REFID=45212
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH) REFID=45210
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+) REFID=40935
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.) REFID=41358
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP) REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+) REFID=40268
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.) REFID=40010
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+) REFID=20765
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.) REFID=21349
BINON	84B	LNC 39 41	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP) REFID=21780
BINON	83C	SJNP 38 723	F.G. Binon <i>et al.</i>	(SERP, BRUX+) REFID=40288
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH) REFID=20754
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL) REFID=20746
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND) REFID=20390
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) REFID=21665
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN) REFID=21774
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) REFID=20374
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+) REFID=21967
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA) REFID=20728
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+) REFID=20720
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) REFID=21651

$\pi_2(2100)$

$$I^G(J^{PC}) = 1^-(2^-+)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M020

NODE=M020

NODE=M020M

NODE=M020M

NODE=M020M;LINKAGE=AX
NODE=M020M;LINKAGE=L

NODE=M020W

NODE=M020W

NODE=M020W;LINKAGE=AX
NODE=M020W;LINKAGE=L

NODE=M020215;NODE=M020

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←

NODE=M020220

NODE=M020R1
NODE=M020R1NODE=M020R2
NODE=M020R2NODE=M020R3
NODE=M020R3NODE=M020R4
NODE=M020R4

NODE=M020R;LINKAGE=L

NODE=M020

REFID=44433
REFID=20872 $\pi_2(2100)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2090± 29 OUR AVERAGE			
2090± 30	¹ AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
2100± 150	² DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$
¹ From a fit to $J^{PC} = 2^-+ f_2(1270)\pi, (\pi\pi)_S\pi$ waves.			
² From a two-resonance fit to four 2^-0^+ waves.			

 $\pi_2(2100)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
625± 50 OUR AVERAGE	Error includes scale factor of 1.2.		
520± 100	³ AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
651± 50	⁴ DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$
³ From a fit to $J^{PC} = 2^-+ f_2(1270)\pi, (\pi\pi)_S\pi$ waves.			
⁴ From a two-resonance fit to four 2^-0^+ waves.			

 $\pi_2(2100)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	seen
Γ_2 $\rho\pi$	seen
Γ_3 $f_2(1270)\pi$	seen
Γ_4 $(\pi\pi)_S\pi$	seen

 $\pi_2(2100)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(3\pi)$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.19±0.05	⁵ DAUM	81B CNTR	63,94 $\pi^- p$
$\Gamma(f_2(1270)\pi)/\Gamma(3\pi)$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.36±0.09	⁵ DAUM	81B CNTR	63,94 $\pi^- p$
$\Gamma((\pi\pi)_S\pi)/\Gamma(3\pi)$	Γ_4/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.45±0.07	⁵ DAUM	81B CNTR	63,94 $\pi^- p$
D-wave/S-wave RATIO FOR $\pi_2(2100) \rightarrow f_2(1270)\pi$			
VALUE	DOCUMENT ID	TECN	COMMENT
0.39±0.23	⁵ DAUM	81B CNTR	63,94 $\pi^- p$
⁵ From a two-resonance fit to four 2^-0^+ waves.			

 $\pi_2(2100)$ REFERENCES

AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

$f_0(2100)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M168

NODE=M168

 $f_0(2100)$ MASS

NODE=M168M

NODE=M168M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2101 ± 7 OUR AVERAGE				
2081 ± 13 ⁺²⁴ ₋₃₆	5.5k	¹ ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
2102 ± 13		² ANISOVICH	00J SPEC	$2.0 \bar{p}p \rightarrow \eta\pi^0\pi^0, \pi^0\pi^0,$ $\eta\eta, \eta\eta', \pi^+\pi^-$
2090 ± 30		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
2105 ± 10		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p}p \rightarrow \eta\eta, \eta\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2090 ± 10 ± 6	529	^{3,4} DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
2099 ± 17 ± 8	283	^{3,4} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
2105 ± 8	80k	⁵ UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
~ 2104		BUGG	95	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
~ 2122		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$

OCCUR=2

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² Includes the data of ANISOVICH 00B indicating to exotic decay pattern.

³ Using CLEO-c data but not authored by the CLEO Collaboration.

⁴ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 209$ MeV.

⁵ Statistical error only.

NODE=M168M;LINKAGE=A

NODE=M168M;LINKAGE=AN

NODE=M168M;LINKAGE=B

NODE=M168M;LINKAGE=C

NODE=M168M;LINKAGE=ST

 $f_0(2100)$ WIDTH

NODE=M168W

NODE=M168W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
224⁺²³₋₂₁ OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
273 ⁺²⁷ ₋₂₄ ⁺⁷⁰ ₋₂₃	5.5k	⁶ ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
211 ± 29		⁷ ANISOVICH	00J SPEC	$2.0 \bar{p}p \rightarrow \eta\pi^0\pi^0, \pi^0\pi^0,$ $\eta\eta, \eta\eta', \pi^+\pi^-$
330 ± 100		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
200 ± 25		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p}p \rightarrow \eta\eta, \eta\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
236 ± 14	80k	⁸ UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
~ 203		BUGG	95	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
~ 273		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$

⁶ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.


⁷ Includes the data of ANISOVICH 00B indicating to exotic decay pattern.

⁸ Statistical error only.

NODE=M168W;LINKAGE=A

NODE=M168W;LINKAGE=AN

NODE=M168W;LINKAGE=ST



f₀(2100) REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>	
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)

NODE=M168

REFID=56805
REFID=55387
REFID=51063
REFID=47942
REFID=47950
REFID=47426
REFID=47472
REFID=44438
REFID=44103

f₂(2150)

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called T₀.

NODE=M042

NODE=M042

f₂(2150) MASS

f₂(2150) MASS, COMBINED MODES (MeV)

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

2157 ± 12 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

••• We do not use the following data for averages, fits, limits, etc. •••

2170 ± 6 80k ¹UMAN 06 E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$

¹ Statistical error only.

NODE=M042205

NODE=M042M
NODE=M042M

NODE=M042M;LINKAGE=ST

ηη MODE

VALUE (MeV) DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

2157 ± 12 OUR AVERAGE

2151 ± 16 BARBERIS 00E 450 $pp \rightarrow p_f \eta \eta p_s$
2175 ± 20 PROKOSHKIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$,
450 $pp \rightarrow pp 2\eta$
2130 ± 35 SINGOVSKI 94 GAM4 450 $pp \rightarrow pp 2\eta$

••• We do not use the following data for averages, fits, limits, etc. •••

2140 ± 30 ²ABELE 99B CBAR

2104 ± 20 ³ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$

² Spin not determined.

³ No J^{PC} determination.

NODE=M042M3
NODE=M042M3

NODE=M042M3;LINKAGE=K3
NODE=M042M3;LINKAGE=A

$\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2135±20±45	⁴ ADOMEIT	96	CBAR	0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
	⁴ ANISOVICH 00E				recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

NODE=M042M4
NODE=M042M4

NODE=M042M4;LINKAGE=AD

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2090	⁵ OAKDEN	94	RVUE	0.36-1.55	$\bar{p}p \rightarrow \pi\pi$
~ 2120	⁶ OAKDEN	94	RVUE	0.36-1.55	$\bar{p}p \rightarrow \pi\pi$
~ 2170	⁷ MARTIN	80B	RVUE		
~ 2150	⁷ MARTIN	80C	RVUE		
~ 2150	⁸ DULUDE	78B	OSPK	1-2	$\bar{p}p \rightarrow \pi^0\pi^0$

NODE=M042M1
NODE=M042M1

OCCUR=2

⁵OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.

NODE=M042M1;LINKAGE=B

⁶From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.

NODE=M042M1;LINKAGE=BB

⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

NODE=M042M1;LINKAGE=P

⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

NODE=M042M1;LINKAGE=L

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2139 ⁺⁸ ₋₉	⁹ EVANGELIS...	97	SPEC		0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
~ 2190	⁹ CUTTS	78B	CNTR		0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155±15	^{9,10} COUPLAND	77	CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2193±2	^{9,11} ALSPECTOR	73	CNTR		$\bar{p}p$ S channel

⁹Isospins 0 and 1 not separated.

NODE=M042M2;LINKAGE=I

¹⁰From a fit to the total elastic cross section.

NODE=M042M2;LINKAGE=E

¹¹Referred to as T or T region by ALSPECTOR 73.

NODE=M042M2;LINKAGE=M

 $K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2200±13	VLADIMIRSK...06	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$
2150±20	ABLIKIM	04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
2130±35	BARBERIS	99	OMEG	450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$

NODE=M042M5
NODE=M042M5

 $f_2(2150)$ WIDTH

NODE=M042210

 $f_2(2150)$ WIDTH, COMBINED MODES (MeV)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

152±30 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.4. See the ideogram below.


NODE=M042W
NODE=M042W

• • • We do not use the following data for averages, fits, limits, etc. • • •

182±11	80k	¹² UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
--------	-----	--------------------	----	------	--

¹²Statistical error only.

NODE=M042W;LINKAGE=ST



$\eta\eta$ MODE

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ COMMENT _____
The data in this block is included in the average printed for a previous datablock.

NODE=M042W3
NODE=M042W3

152±30 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.


280±70 BARBERIS 00E 450 $pp \rightarrow p_f \eta \eta p_s$
150±35 PROKOSHKIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$,
450 $pp \rightarrow pp 2\eta$
130±30 SINGOVSKI 94 GAM4 450 $pp \rightarrow pp 2\eta$

••• We do not use the following data for averages, fits, limits, etc. •••

310±50 ¹³ ABELE 99B CBAR
203±10 ¹⁴ ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$

¹³ Spin not determined.
¹⁴ No J^{PC} determination.

NODE=M042W3;LINKAGE=K3
NODE=M042W3;LINKAGE=A



$\eta\pi\pi$ MODE

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____
The data in this block is included in the average printed for a previous datablock.

NODE=M042W4
NODE=M042W4

••• We do not use the following data for averages, fits, limits, etc. •••

250±25±45 ¹⁵ ADOMEIT 96 CBAR 0 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

¹⁵ ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

NODE=M042W4;LINKAGE=AD

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

250 OUR ESTIMATE

••• We do not use the following data for averages, fits, limits, etc. •••

~ 70	¹⁶ OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	¹⁷ MARTIN	80B	RVUE
~ 250	¹⁷ MARTIN	80C	RVUE
~ 250	¹⁸ DULUDE	78B	OSPK 1–2 $\bar{p}p \rightarrow \pi^0\pi^0$

NODE=M042W1
 NODE=M042W1
 → UNCHECKED ←

¹⁶ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

NODE=M042W1;LINKAGE=CC

¹⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

NODE=M042W1;LINKAGE=P

¹⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

NODE=M042W1;LINKAGE=L

S-CHANNEL $\bar{p}p, \bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

56^{+31}_{-16}	¹⁹ EVANGELIS...	97	SPEC	0.6–2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
135 ± 75	^{20,21} COUPLAND	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98 ± 8	²¹ ALSPECTOR	73	CNTR	$\bar{p}p$ S channel

NODE=M042W2
 NODE=M042W2

¹⁹ Isospin 0 and 2 not separated.

NODE=M042W2;LINKAGE=F

²⁰ From a fit to the total elastic cross section.

NODE=M042W2;LINKAGE=E

²¹ Isospins 0 and 1 not separated.

NODE=M042W2;LINKAGE=I

 $K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

91 ± 62	VLADIMIRSK..06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
150 ± 30	ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
270 ± 50	BARBERIS	99	OMEG 450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$

NODE=M042W5
 NODE=M042W5

 $f_2(2150)$ DECAY MODES

NODE=M042215;NODE=M042

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	
Γ_2 $\eta\eta$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $f_2(1270)\eta$	seen
Γ_5 $a_2(1320)\pi$	seen
Γ_6 $\rho\bar{\rho}$	seen

DESIG=1
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6

 $f_2(2150)$ BRANCHING RATIOS

NODE=M042220

 $\Gamma(K\bar{K})/\Gamma(\eta\eta)$ **Γ_3/Γ_2**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

1.28 ± 0.23 BARBERIS 00E 450 $p\bar{p} \rightarrow p_f \eta \eta p_S$

NODE=M042R1
 NODE=M042R1

••• We do not use the following data for averages, fits, limits, etc. •••

<0.1	95	²² PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p\rho 2\eta$
------	----	------------------------------	------	---

NODE=M042R1;LINKAGE=A

²² Using data from ARMSTRONG 89D.

 $\Gamma(\pi\pi)/\Gamma(\eta\eta)$ **Γ_1/Γ_2**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.33	95	²³ PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p\rho 2\eta$
-------	----	------------------------------	------	---

NODE=M042R2
 NODE=M042R2

²³ Derived from a $\pi^0\pi^0/\eta\eta$ limit.

NODE=M042R2;LINKAGE=A

 $\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$ **Γ_4/Γ_5**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.79 ± 0.11 ²⁴ ADOMEIT 96 CBAR 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M042R3
 NODE=M042R3

²⁴ Using $B(a_2(1320) \rightarrow \eta\pi) = 0.145$

NODE=M042R3;LINKAGE=A

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	73	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M042R04
 NODE=M042R04

 $f_2(2150)$ REFERENCES

ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
EVANGELIS...	97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
MARTIN	97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
PROKOSHKIN	95D	SPD 40 495	Y.D. Prokoshkin	(SERP) IGJPC
		Translated from DANS 344 469.		
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
SINGOVSKI	94	NC 107A 1911	A.V. Singovsky	(SERP)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE	78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)

NODE=M042

REFID=53525
 REFID=51063
 REFID=51191

REFID=50174
 REFID=47945
 REFID=47961
 REFID=46904
 REFID=46921
 REFID=45687
 REFID=45685
 REFID=45202
 REFID=45212
 REFID=44647

REFID=44103
 REFID=45210
 REFID=44648
 REFID=43587
 REFID=41010
 REFID=21838
 REFID=21837
 REFID=21733
 REFID=21850
 REFID=21830
 REFID=21813

 $\rho(2150)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $T_1(2190)$. See our mini-review under the $\rho(1700)$.

NODE=M032

NODE=M032

 $\rho(2150)$ MASS **e^+e^- PRODUCED**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
●●● We do not use the following data for averages, fits, limits, etc. ●●●			
2254 ± 22	¹ LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
$2150 \pm 40 \pm 50$	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
1990 ± 80	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$
2153 ± 37	BIAGINI	91 RVUE	$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-$
2110 ± 50	² CLEGG	90 RVUE	$e^+e^- \rightarrow 3(\pi^+\pi^-), 2(\pi^+\pi^-\pi^0)$

NODE=M032205

NODE=M032M3
 NODE=M032M3

OCCUR=2

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
●●● We do not use the following data for averages, fits, limits, etc. ●●●			
~ 2191	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2070	³ OAKDEN	94 RVUE	$0.36-1.55 \bar{p}p \rightarrow \pi\pi$
~ 2170	⁴ MARTIN	80B RVUE	
~ 2100	⁴ MARTIN	80C RVUE	

NODE=M032M1
 NODE=M032M1

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
●●● We do not use the following data for averages, fits, limits, etc. ●●●			
2110 ± 35	⁵ ANISOVICH	02 SPEC	$0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2190	⁶ CUTTS	78B CNTR	$0.97-3 \bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{6,7} COUPLAND	77 CNTR	$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{6,8} ALSPECTOR	73 CNTR	$\bar{p}p$ S channel
2190 ± 10	⁹ ABRAMS	70 CNTR	S channel $\bar{p}N$

NODE=M032M2
 NODE=M032M2

$\pi^- p \rightarrow \omega \pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2155±21 OUR AVERAGE			
2140±30	ALDE	95	GAM2 38 $\pi^- p \rightarrow \omega \pi^0 n$
2170±30	ALDE	92C	GAM4 100 $\pi^- p \rightarrow \omega \pi^0 n$

¹ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

² Includes ATKINSON 85.

³ See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

⁴ $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

⁵ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

⁶ Isospins 0 and 1 not separated.

⁷ From a fit to the total elastic cross section.

⁸ Referred to as T or T region by ALSPECTOR 73.

⁹ Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M032M4
NODE=M032M4

NODE=M032M3;LINKAGE=LE

NODE=M032M3;LINKAGE=A
NODE=M032M1;LINKAGE=CC

NODE=M032M;LINKAGE=P
NODE=M032M;LINKAGE=AY

NODE=M032M;LINKAGE=I
NODE=M032M;LINKAGE=E
NODE=M032M;LINKAGE=M
NODE=M032M;LINKAGE=B

 $\rho(2150)$ WIDTH

NODE=M032210

 $e^+ e^-$ PRODUCED

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
109± 76	¹⁰ LEES	12G	BABR $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
350± 40±50	AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
310±140	AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow \eta' \pi^+ \pi^- \gamma$
389± 79	BIAGINI	91	RVUE $e^+ e^- \rightarrow \pi^+ \pi^-, K^+ K^-$
410±100	¹¹ CLEGG	90	RVUE $e^+ e^- \rightarrow 3(\pi^+ \pi^-), 2(\pi^+ \pi^- \pi^0)$

NODE=M032W3
NODE=M032W3

OCCUR=2

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 296	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 40	¹² OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	¹³ MARTIN	80B	RVUE
~ 200	¹³ MARTIN	80C	RVUE

NODE=M032W1
NODE=M032W1

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
230±50	¹⁴ ANISOVICH	02	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega \pi^0, \omega \eta \pi^0, \pi^+ \pi^-$
135±75	^{15,16} COUPLAND	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98± 8	¹⁶ ALSPECTOR	73	CNTR $\bar{p}p$ S channel
~ 85	¹⁷ ABRAMS	70	CNTR S channel $\bar{p}N$

NODE=M032W2
NODE=M032W2

 $\pi^- p \rightarrow \omega \pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
320±70	ALDE	95	GAM2 38 $\pi^- p \rightarrow \omega \pi^0 n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 300	ALDE	92C	GAM4 100 $\pi^- p \rightarrow \omega \pi^0 n$

NODE=M032W4
NODE=M032W4

¹⁰ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

¹¹ Includes ATKINSON 85.

¹² See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

¹³ $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

¹⁴ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

¹⁵ From a fit to the total elastic cross section.

¹⁶ Isospins 0 and 1 not separated.

¹⁷ Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M032W3;LINKAGE=LE

NODE=M032W3;LINKAGE=A
NODE=M032W1;LINKAGE=CC

NODE=M032W;LINKAGE=P
NODE=M032W;LINKAGE=AY

NODE=M032W;LINKAGE=E
NODE=M032W;LINKAGE=I
NODE=M032W;LINKAGE=B

$\rho(2150)$ DECAY MODES

NODE=M032215;NODE=M032

Mode	Fraction (Γ_i/Γ)
Γ_1 $e^+ e^-$	
Γ_2 $\pi^+ \pi^-$	seen
Γ_3 $K^+ K^-$	seen
Γ_4 $3(\pi^+ \pi^-)$	seen
Γ_5 $2(\pi^+ \pi^- \pi^0)$	seen
Γ_6 $\eta' \pi^+ \pi^-$	seen
Γ_7 $f_1(1285) \pi^+ \pi^-$	seen
Γ_8 $\omega \pi^0$	seen
Γ_9 $\omega \pi^0 \eta$	seen
Γ_{10} $p \bar{p}$	

DESIG=1
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=5;OUR EVAL;→ UNCHECKED ←
DESIG=6;OUR EVAL;→ UNCHECKED ←
DESIG=7;OUR EVAL;→ UNCHECKED ←
DESIG=8;OUR EVAL;→ UNCHECKED ←
DESIG=9;OUR EVAL;→ UNCHECKED ←
DESIG=10

 $\rho(2150) \Gamma(i)\Gamma(e^+ e^-)/\Gamma^2(\text{total})$

NODE=M032230

$\Gamma(f_1(1285)\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma \times \Gamma_1/\Gamma$
VALUE (units 10^{-7})	DOCUMENT ID TECN COMMENT
$3.1 \pm 0.6 \pm 0.5$	¹⁸ AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$

NODE=M032G01
NODE=M032G01

¹⁸ Calculated by us from the reported value of cross section at the peak.

NODE=M032G01;LINKAGE=AU

$\Gamma(\eta' \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma \times \Gamma_1/\Gamma$
VALUE (units 10^{-8})	DOCUMENT ID TECN COMMENT

NODE=M032G02
NODE=M032G02

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.9 ± 1.9	¹⁹ AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow \eta' \pi^+ \pi^- \gamma$
---------------	--

¹⁹ Calculated by us from the reported value of cross section at the peak.

NODE=M032G02;LINKAGE=AU

 $\rho(2150)$ REFERENCES

NODE=M032

LEES 12G PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT 07AU PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
ANISOVICH 02 PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01D PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01E PL B513 281	A.V. Anisovich <i>et al.</i>	
ANISOVICH 00J PL B491 47	A.V. Anisovich <i>et al.</i>	
KLOET 96 PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ALDE 95 ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
HASAN 94 PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN 94 NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ALDE 92C ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
BIAGINI 91 NC 104A 363	M.E. Biagini <i>et al.</i>	(FRAS, PRAG)
CLEGG 90 ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ATKINSON 85 ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
MARTIN 80B NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS 78B PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
COUPLAND 77 PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE 75 PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR 73 PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS 70 PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER 68 PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)
GOUNARIS 68 PRL 21 244	G.J. Gounaris, J.J. Sakurai	

REFID=54299
REFID=52049
REFID=48828
REFID=48327
REFID=48349
REFID=47950
REFID=45212
REFID=44371
REFID=44103
REFID=45210
REFID=41859
REFID=41894
REFID=41355
REFID=22000
REFID=21838
REFID=21837
REFID=21733
REFID=21830
REFID=21824
REFID=21813
REFID=21807
REFID=21805
REFID=48054

$\phi(2170)$

$$J^G(J^{PC}) = 0^-(1^{--})$$

Observed by AUBERT, BE 06D in the initial-state radiation process
 $e^+e^- \rightarrow \phi f_0(980)\gamma$.

NODE=M103

NODE=M103

NODE=M103M

NODE=M103M

NEW

OCCUR=2


NODE=M103M;LINKAGE=AB

NODE=M103M;LINKAGE=SH

NODE=M103M;LINKAGE=AU

$\phi(2170)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2189±11 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below. [2175 ± 15 MeV OUR 2015 AVERAGE Scale factor = 1.6]
2200±6±5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
2186±10±6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
2125±22±10	483	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
2175±10±15	201	¹ AUBERT, BE	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi\pi\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2079±13 ⁺⁷⁹ ₋₂₈	4.8k	² SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2192±14	116	³ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2169±20	149	³ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$

¹ From the $\phi f_0(980)$ component.² From a fit with two incoherent Breit-Wigners.³ From the $K^+K^-f_0(980)$ component.

$\phi(2170)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
79±14 OUR AVERAGE				[61 ± 18 MeV OUR 2015 AVERAGE]
104±15±15	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
65±23±17	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
61±50±13	483	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
58±16±20	201	⁴ AUBERT, BE	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi\pi\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
192±23 ⁺²⁵ ₋₆₁	4.8k	⁵ SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
71±21	116	⁶ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
102±27	149	⁶ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$

⁴ From the $\phi f_0(980)$ component.⁵ From a fit with two incoherent Breit-Wigners.⁶ From the $K^+K^-f_0(980)$ component.

NODE=M103W

NODE=M103W

NEW

OCCUR=2

NODE=M103W;LINKAGE=AB

NODE=M103W;LINKAGE=SH

NODE=M103W;LINKAGE=AU

$\phi(2170)$ DECAY MODES

NODE=M103215;NODE=M103

Mode	Fraction (Γ_i/Γ)
Γ_1 $e^+ e^-$	seen
Γ_2 $\phi\eta$	
Γ_3 $\phi\pi\pi$	
Γ_4 $\phi f_0(980)$	seen
Γ_5 $K^+ K^- \pi^+ \pi^-$	
Γ_6 $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
Γ_7 $K^+ K^- \pi^0 \pi^0$	
Γ_8 $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
Γ_9 $K^{*0} K^\pm \pi^\mp$	not seen
Γ_{10} $K^*(892)^0 \bar{K}^*(892)^0$	not seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=5
DESIG=9
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3
DESIG=6
DESIG=4
DESIG=7
DESIG=8
DESIG=10

 $\phi(2170) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

NODE=M103230

$\Gamma(\phi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_1/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT

NODE=M103G2
NODE=M103G2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.7±0.7±1.3 483 AUBERT 08S BABR 10.6 $e^+ e^- \rightarrow \phi\eta\gamma$

$\Gamma(\phi f_0(980)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_4\Gamma_1/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT

NODE=M103G1
NODE=M103G1

2.5±0.8±0.4 201 ⁷AUBERT,BE 06D BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi\pi\gamma$

⁷ From the $\phi f_0(980)$ component.

NODE=M103G1;LINKAGE=AB

 $\phi(2170) \Gamma(i)\Gamma(e^+ e^-)/\Gamma^2(\text{total})$

NODE=M103220

$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma \times \Gamma_1/\Gamma$			
VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT

NODE=M103G01
NODE=M103G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.65±0.15±0.18 4.8k ⁸ SHEN 09 BELL 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

⁸ Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

NODE=M103G01;LINKAGE=SH

 $\phi(2170)$ BRANCHING RATIOS

NODE=M103225

$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_6/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M103R01
NODE=M103R01

seen AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$	Γ_8/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M103R02
NODE=M103R02

seen AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

$\Gamma(K^{*0} K^\pm \pi^\mp)/\Gamma_{\text{total}}$	Γ_9/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M103R03
NODE=M103R03

not seen AUBERT 07AK BABR 10.6 GeV $e^+ e^-$

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$	Γ_{10}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M103R04
NODE=M103R04

not seen ABLIKIM 10C BES2 $J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

 $\phi(2170)$ REFERENCES

NODE=M103

ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56773
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511

$f_0(2200)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

NODE=M112

OMITTED FROM SUMMARY TABLE

Seen in $K_S^0 K_S^0$ (AUGUSTIN 88), $K^+ K^-$ (ABLIKIM 05Q) and $\eta\eta$ (BINON 05) system. Not seen in $\Upsilon(1S)$ radiative decays (BARU 89).

NODE=M112

 $f_0(2200)$ MASS

NODE=M112M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M112M

2189±13 OUR AVERAGE2170±20⁺¹⁰₋₁₅ABLIKIM 05Q BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^- K^+ K^-$

2210±50

¹ BINON 05 GAMS 33 $\pi^- p \rightarrow \eta\eta n$

2197±17

² AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2206±12± 8

381 3,4 DOBBS 15 $J/\psi \rightarrow \gamma K^+ K^-$

~ 2122

203 3,4 DOBBS 15 $\psi(2S) \rightarrow \gamma K^+ K^-$

~ 2321

HASAN 94 RVUE $\bar{p}p \rightarrow \pi\pi$ HASAN 94 RVUE $\bar{p}p \rightarrow \pi\pi$

OCCUR=2

OCCUR=2

¹ First solution, PWA is ambiguous.² Cannot determine spin to be 0.³ Using CLEO-c data but not authored by the CLEO Collaboration.⁴ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 238$ MeV.

NODE=M112M;LINKAGE=BI

NODE=M112M;LINKAGE=A

NODE=M112M;LINKAGE=B

NODE=M112M;LINKAGE=C

 $f_0(2200)$ WIDTH

NODE=M112W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M112W

238±50 OUR AVERAGE

Error includes scale factor of 1.2.

220±60⁺⁴⁰₋₄₅ABLIKIM 05Q BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^- K^+ K^-$

380±90

⁵ BINON 05 GAMS 33 $\pi^- p \rightarrow \eta\eta n$

201±51

⁶ AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 273

HASAN 94 RVUE $\bar{p}p \rightarrow \pi\pi$

~ 223

HASAN 94 RVUE $\bar{p}p \rightarrow \pi\pi$

OCCUR=2

⁵ First solution, PWA is ambiguous.⁶ Cannot determine spin to be 0.

NODE=M112W;LINKAGE=BI

NODE=M112W;LINKAGE=A

 $f_0(2200)$ REFERENCES

NODE=M112

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)	REFID=40917
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574

$f_J(2220)$

$$J^G(J^{PC}) = 0^+(2^{++} \text{ or } 4^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. See our mini-review in the 2004 edition of this Review, PDG 04.

NODE=M082

NODE=M082

 $f_J(2220)$ MASS

NODE=M082M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2231.1 ± 3.5 OUR AVERAGE				
2235 ± 4 ± 6	74	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
2230 $^{+6}_{-7}$ ±16	46	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
2232 $^{+8}_{-7}$ ±15	23	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
2235 ± 4 ± 5	32	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
2209 $^{+17}_{-15}$ ±10		ASTON	88F LASS	$11 K^-p \rightarrow K^+K^- \Lambda$
2230 ± 20		BOLONKIN	88 SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n$
2220 ± 10	41	¹ ALDE	86B GA24	$38-100 \pi p \rightarrow n\eta\eta'$
2230 ± 6 ± 14	93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
2232 ± 7 ± 7	23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0 K_S^0$
2223.9 ± 2.5		² VLADIMIRSK...08	SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n + m\pi^0$
2246 ± 36		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$

NODE=M082M

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ALDE 86B uses data from both the GAMS-2000 and GAMS-4000 detectors.² $J^{PC} = 2^{++}$. Systematic uncertainties not evaluatedNODE=M082M;LINKAGE=A
NODE=M082M;LINKAGE=VL $f_J(2220)$ WIDTH

NODE=M082W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
23 $^{+8}_{-7}$ OUR AVERAGE					
19 $^{+13}_{-11}$ ±12		74	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
20 $^{+20}_{-15}$ ±17		46	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
20 $^{+25}_{-16}$ ±14		23	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
15 $^{+12}_{-9}$ ± 9		32	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
60 $^{+107}_{-57}$			ASTON	88F LASS	$11 K^-p \rightarrow K^+K^- \Lambda$
80 ± 30			BOLONKIN	88 SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n$
26 $^{+20}_{-16}$ ±17		93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
18 $^{+23}_{-15}$ ±10		23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0 K_S^0$
8.6 ± 2.5			¹ VLADIMIRSK...08	SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n + m\pi^0$
<80	90		ALDE	87C GAM2	$38 \pi^-p \rightarrow \eta'\eta n$

NODE=M082W

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ $J^{PC} = 2^{++}$. Systematic uncertainties not evaluated

NODE=M082W;LINKAGE=VL

 $f_J(2220)$ DECAY MODES

NODE=M082215;NODE=M082

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	not seen
Γ_2 $\pi^+\pi^-$	not seen
Γ_3 $K\bar{K}$	not seen
Γ_4 $p\bar{p}$	not seen
Γ_5 $\gamma\gamma$	not seen
Γ_6 $\eta\eta'(958)$	seen
Γ_7 $\phi\phi$	not seen
Γ_8 $\eta\eta$	not seen

DESIG=5

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=1

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

$f_J(2220) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M082220

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_5/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	95	1 ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183-209 \text{ GeV}$

NODE=M082G1
NODE=M082G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.6	95	1 GODANG	97 CLE2	$\gamma\gamma \rightarrow K_S^0 K_S^0$
< 86	95	1 ALBRECHT	90G ARG	$\gamma\gamma \rightarrow K^+ K^-$
<1000	95	2 ALTHOFF	85B TASS	$\gamma\gamma, K\bar{K}\pi$

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_5/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	95	ALAM	98C CLE2	$\gamma\gamma \rightarrow \pi^+ \pi^-$

NODE=M082G3
NODE=M082G3¹ Assuming $J^P = 2^+$.² True for $J^P = 0^+$ and $J^P = 2^+$.NODE=M082G1;LINKAGE=D
NODE=M082G1;LINKAGE=C $f_J(2220) \Gamma(i)\Gamma(\rho\bar{\rho})/\Gamma^2(\text{total})$

NODE=M082223

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<18	95	1 AMSLER	01 CBAR	$1.4-1.5 \rho\bar{\rho} \rightarrow \pi^0 \pi^0$

NODE=M082GG1
NODE=M082GG1

• • • We do not use the following data for averages, fits, limits, etc. • • •

<(11-42)	99	2 HASAN	96 SPEC	$1.35-1.55 \rho\bar{\rho} \rightarrow \pi^+ \pi^-$
----------	----	---------	---------	--

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_7/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6	95	3 EVANGELIS...	98 SPEC	$1.1-2.0 \rho\bar{\rho} \rightarrow \phi\phi$

NODE=M082GG2
NODE=M082GG2 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<4	95	1 AMSLER	01 CBAR	$1.4-1.5 \rho\bar{\rho} \rightarrow \eta\eta$

NODE=M082GG3
NODE=M082GG3¹ For $J^P = 2^+$ in the mass range 2222-2240 MeV and the total width between 10 and 20 MeV.² For $J^P = 2^+$ and $J^P = 4^+$ in the mass range 2220-2245 MeV and the total width of 15 MeV.³ For $J^P = 2^+$, the mass of 2235 MeV and the total width of 15 MeV.

NODE=M082GG;LINKAGE=A

NODE=M082GG;LINKAGE=B

NODE=M082GG;LINKAGE=C

 $f_J(2220) \text{ BRANCHING RATIOS}$

NODE=M082225

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	COMMENT
not seen	1 DOBBS 15	$J/\psi \rightarrow \gamma\pi\pi$
not seen	1 DOBBS 15	$\psi(2S) \rightarrow \gamma\pi\pi$

NODE=M082R00
NODE=M082R00¹ Using CLEO-c data but not authored by the CLEO Collaboration.

OCCUR=2

NODE=M082R00;LINKAGE=A

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	COMMENT
not seen	1 DOBBS 15	$J/\psi \rightarrow \gamma K\bar{K}$
not seen	1 DOBBS 15	$\psi(2S) \rightarrow \gamma K\bar{K}$

NODE=M082R01
NODE=M082R01¹ Using CLEO-c data but not authored by the CLEO Collaboration.

OCCUR=2

NODE=M082R01;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$ Γ_1/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
1.0±0.5	BAI 96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma 2\pi, K\bar{K}$

NODE=M082R2
NODE=M082R2 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
not seen		1 AUBERT	07AV BABR	$B \rightarrow \rho\bar{\rho}K^*$
not seen		WANG	05A BELL	$B^+ \rightarrow \bar{p}\rho K^+$
<3.0	95	2 EVANGELIS...	97 SPEC	$1.96-2.40 \rho\bar{\rho} \rightarrow K_S^0 K_S^0$
<1.1	99.7	3 BARNES	93 SPEC	$1.3-1.57 \rho\bar{\rho} \rightarrow K_S^0 K_S^0$
<2.6	99.7	3 BARDIN	87 CNTR	$1.3-1.5 \rho\bar{\rho} \rightarrow K^+ K^-$
<3.6	99.7	3 SCULLI	87 CNTR	$1.29-1.55 \rho\bar{\rho} \rightarrow K^+ K^-$

NODE=M082R1
NODE=M082R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

- ¹ Assuming $\Gamma < 30$ MeV.
² Assuming $\Gamma \sim 20$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.
³ Assuming $\Gamma = 30\text{-}35$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.

NODE=M082R1;LINKAGE=AU

NODE=M082R1;LINKAGE=C

NODE=M082R1;LINKAGE=B

 $\Gamma(p\bar{p})/\Gamma(K\bar{K})$ Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.09	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}, K\bar{K}$

NODE=M082R3

NODE=M082R3

 $f_J(2220)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
VLADIMIRSK...	08	PAN 71 2129	V.V. Vladimirov <i>et al.</i>	(ITEP)
		Translated from YAF 71 2166.		
AUBERT	07AV	PR D76 092004	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
AMSLER	01	PL B520 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ALAM	98C	PRL 81 3328	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
EVANGELIS...	98	PR D57 5370	C. Evangelista <i>et al.</i>	(JETSET Collab.)
EVANGELIS...	97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
GODANG	97	PRL 79 3829	R. Godang <i>et al.</i>	(CLEO Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ASTON	88F	PL B215 199	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ALDE	87C	SJNP 45 255	D. Alde <i>et al.</i>	
		Translated from YAF 45 405.		
BARDIN	87	PL B195 292	G. Bardin <i>et al.</i>	(SACL, FERR, CERN, PADO+)
SCULLI	87	PRL 58 1715	J. Sculli <i>et al.</i>	(NYU, BNL)
ALDE	86B	PL B177 120	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)

NODE=M082

REFID=56805

REFID=52681

REFID=51990

REFID=50651

REFID=49653

REFID=48321

REFID=48558

REFID=46326

REFID=46342

REFID=46365

REFID=45687

REFID=45760

REFID=44736

REFID=45197

REFID=43601

REFID=41374

REFID=40585

REFID=40580

REFID=47474

REFID=40235

REFID=40023

REFID=21864

REFID=21865

REFID=21349

OTHER RELATED PAPERS

DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
---------------	-----	----------------	----------------------------------	-----------------

REFID=53533

 $\eta(2225)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Seen in $J/\psi \rightarrow \gamma\phi\phi$. Possibly seen in $B \rightarrow \phi\phi K$ by LEES 11A.

NODE=M115

NODE=M115

 $\eta(2225)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2226±16 OUR AVERAGE				
2240 ⁺³⁰⁺³⁰ ₋₂₀₋₂₀	196 ± 19	ABLIKIM	08I	BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
2230±25±15		BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
2214±20±13		BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

NODE=M115M

NODE=M115M

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 2220		BISELLO	86B	DM2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
--------	--	---------	-----	---

OCCUR=2

 $\eta(2225)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
185⁺⁷⁰₋₄₀ OUR AVERAGE				
190± 30 ⁺⁶⁰ ₋₄₀	196 ± 19	ABLIKIM	08I	BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
150 ⁺³⁰⁰ ₋₆₀ ± 60		BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
~ 80		BISELLO	86B	DM2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M115W

NODE=M115W

 $\eta(2225)$ REFERENCES

LEES	11A	PR D84 012001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)

NODE=M115

REFID=16595

REFID=52255

REFID=41354

REFID=22101

$\rho_3(2250)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

NODE=M044

OMITTED FROM SUMMARY TABLE

Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also $\rho(2150)$, $f_2(2150)$, $f_4(2300)$, $\rho_5(2350)$.

NODE=M044

 $\rho_3(2250)$ MASS

NODE=M044205

 $\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 2232	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2090	¹ OAKDEN	94	RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2250	² MARTIN	80B	RVUE	
~ 2300	² MARTIN	80C	RVUE	
~ 2140	³ CARTER	78B	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2150	⁴ CARTER	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

NODE=M044M1
NODE=M044M1

¹ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

² $I(J^P) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

³ $l = 0, 1, J^P = 3^-$ from Barrelet-zero analysis.

⁴ $I(J^P) = 1(3^-)$ from amplitude analysis.

NODE=M044M1;LINKAGE=CC

NODE=M044M1;LINKAGE=P
NODE=M044M1;LINKAGE=K
NODE=M044M1;LINKAGE=J**S-CHANNEL $\bar{N}N$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2260 ± 20	⁵ ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2190	⁶ CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{6,7} COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{6,8} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
2190 ± 10	⁹ ABRAMS	70	CNTR	S channel $\bar{p}N$

NODE=M044M2
NODE=M044M2

⁵ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

⁶ Isospins 0 and 1 not separated.

⁷ From a fit to the total elastic cross section.

⁸ Referred to as T or \bar{T} region by ALSPECTOR 73.

⁹ Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M044M;LINKAGE=AY

NODE=M044M2;LINKAGE=I
NODE=M044M2;LINKAGE=E
NODE=M044M2;LINKAGE=M
NODE=M044M2;LINKAGE=B **$\pi^- p \rightarrow \eta\pi\pi$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2290 \pm 20 \pm 30$	AMELIN	00	VES	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M044M3
NODE=M044M3 **$\rho_3(2250)$ WIDTH**

NODE=M044210

 $\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 220	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 60	¹⁰ OAKDEN	94	RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	¹¹ MARTIN	80B	RVUE	
~ 200	¹¹ MARTIN	80C	RVUE	
~ 150	¹² CARTER	78B	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 200	¹³ CARTER	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

NODE=M044W1
NODE=M044W1

¹⁰ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

¹¹ $I(J^P) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

¹² $l = 0, 1, J^P = 3^-$ from Barrelet-zero analysis.

¹³ $I(J^P) = 1(3^-)$ from amplitude analysis.

NODE=M044W1;LINKAGE=CC

NODE=M044W1;LINKAGE=P
NODE=M044W1;LINKAGE=K
NODE=M044W1;LINKAGE=J

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
160±25	¹⁴ ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
135±75	^{15,16} COUPLAND	77	CNTR	0 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
98±8	¹⁶ ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
~ 85	¹⁷ ABRAMS	70	CNTR	S channel $\bar{p}N$

- ¹⁴ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.
¹⁵ From a fit to the total elastic cross section.
¹⁶ Isospins 0 and 1 not separated.
¹⁷ Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M044W2
 NODE=M044W2

NODE=M044W;LINKAGE=AY

NODE=M044W2;LINKAGE=E
 NODE=M044W2;LINKAGE=I
 NODE=M044W2;LINKAGE=B

 $\pi^- p \rightarrow \eta\pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
230±50±80	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M044W3
 NODE=M044W3

 $\rho_3(2250)$ REFERENCES

ANISOVICH 02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
AMELIN 00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
KLOET 96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN 94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER 78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
CARTER 77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE 75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS 70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER 68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)

NODE=M044

REFID=48828
 REFID=48327
 REFID=48349
 REFID=47432
 REFID=47950
 REFID=45212
 REFID=44103
 REFID=45210
 REFID=21838
 REFID=21837
 REFID=21964
 REFID=21733
 REFID=21963
 REFID=21830
 REFID=21824
 REFID=21813
 REFID=21807
 REFID=21805

 $f_2(2300)$

$$I^G(J^PC) = 0^+(2^{++})$$

NODE=M107

 $f_2(2300)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2297±28	¹ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi\phi n$
2243 ⁺⁷ ₋₆ ⁺³ ₋₂₉	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
2270±12	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2327±9±6	ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^- K^+K^-$
2231±10	BOOTH	86	OMEG 85 $\pi^- Be \rightarrow 2\phi Be$
2220 ⁺⁹⁰ ₋₂₀	LINDENBAUM	84	RVUE
2320±40	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

NODE=M107M

NODE=M107M

- ¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi 2^{++} S_2$, D_2 , and D_0 is 6^{+15}_{-5} , 25^{+18}_{-14} , and 69^{+16}_{-27} , respectively.

NODE=M107M;LINKAGE=C

 $f_2(2300)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
149±41	² ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi\phi n$
145±12 ⁺²⁷ ₋₃₄	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
90±29	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
275±36±20	ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^- K^+K^-$
133±50	BOOTH	86	OMEG 85 $\pi^- Be \rightarrow 2\phi Be$
200±50	LINDENBAUM	84	RVUE
220±70	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

NODE=M107W

NODE=M107W

- ² Includes data of ETKIN 85.

NODE=M107W;LINKAGE=C

$f_2(2300)$ DECAY MODES

NODE=M107215;NODE=M107

Mode	Fraction (Γ_i/Γ)
Γ_1 $\phi\phi$	seen
Γ_2 $K\bar{K}$	seen
Γ_3 $\gamma\gamma$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←

 $f_2(2300) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M107225

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_3/\Gamma$
VALUE (eV)	DOCUMENT ID TECN COMMENT

NODE=M107G1
NODE=M107G1

••• We do not use the following data for averages, fits, limits, etc. •••

$3.2^{+0.5+1.3}_{-0.4-2.2}$	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
$44 \pm 6 \pm 12$	³ ABE	04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$

³ Assuming spin 2.

NODE=M107G1;LINKAGE=AB

 $f_2(2300)$ REFERENCES

NODE=M107

UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)
BOOTH	86	NP B273 677	P.S.L. Booth <i>et al.</i>	(LIVP, GLAS, CERN)
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)
ETKIN	82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)

REFID=55592
REFID=51191
REFID=49650
REFID=40285
REFID=21870
REFID=21871
REFID=21869
REFID=21866

NODE=M041

 $f_4(2300)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_0(2350)$. Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $\rho_5(2350)$.

NODE=M041

 $f_4(2300)$ MASS

NODE=M041205

NODE=M041M

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
~ 2314	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 2300	¹ MARTIN	80B	RVUE
~ 2300	¹ MARTIN	80C	RVUE
~ 2340	² CARTER	78B	CNTR $0.7-2.4 \bar{p}p \rightarrow K^- K^+$
~ 2330	DULUDE	78B	OSPK $1-2 \bar{p}p \rightarrow \pi^0 \pi^0$
~ 2310	³ CARTER	77	CNTR $0.7-2.4 \bar{p}p \rightarrow \pi\pi$

NODE=M041M1
NODE=M041M1

¹ $I(J^P) = 0(4^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

² $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.

³ $I(J^P) = 0(4^+)$ from amplitude analysis.

NODE=M041M1;LINKAGE=P
NODE=M041M1;LINKAGE=K
NODE=M041M1;LINKAGE=J

S-CHANNEL $\bar{p}p$ or $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2283 ± 17	⁴ ANISOVICH	00J	SPEC
~ 2380	⁵ CUTTS	78B	CNTR $0.97-3 \bar{p}p \rightarrow \bar{N}N$
2345 ± 15	^{5,6} COUPLAND	77	CNTR $0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
2359 ± 2	^{5,7} ALSPECTOR	73	CNTR $\bar{p}p$ S channel
2375 ± 10	ABRAMS	70	CNTR S channel $\bar{N}N$

NODE=M041M2
NODE=M041M2

⁴ From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on $\bar{p}p \rightarrow \eta\pi^0\pi^0$, $\pi^0\pi^0$, $\eta\eta$, $\eta\eta'$, $\pi^+\pi^-$.

⁵ Isospins 0 and 1 not separated.

⁶ From a fit to the total elastic cross section.

⁷ Referred to as U or U region by ALSPECTOR 73.

NODE=M041M2;LINKAGE=AN

NODE=M041M2;LINKAGE=I
NODE=M041M2;LINKAGE=E
NODE=M041M2;LINKAGE=M

$\pi^- p \rightarrow \eta \pi \pi n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2330 \pm 20 \pm 40$	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

NODE=M041M3
 NODE=M041M3

 $\rho\rho$ CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	COMMENT
2320 ± 60 OUR ESTIMATE		
2332 ± 15	BARBERIS	00F 450 $\rho\rho \rightarrow p_f \omega \omega p_s$

NODE=M041M4
 NODE=M041M4
 → UNCHECKED ←

 $f_4(2300)$ WIDTH

NODE=M041210

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
~ 278	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 200	⁸ MARTIN	80C	RVUE
~ 150	⁹ CARTER	78B	CNTR 0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 210	¹⁰ CARTER	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \pi\pi$
⁸ $I(J^P) = 0(4^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.			
⁹ $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.			
¹⁰ $I(J^P) = 0(4^+)$ from amplitude analysis.			

NODE=M041W1
 NODE=M041W1

NODE=M041W1;LINKAGE=P
 NODE=M041W1;LINKAGE=K
 NODE=M041W1;LINKAGE=J

S-CHANNEL $\bar{p}p$ or $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
310 ± 25	¹¹ ANISOVICH	00J	SPEC
135^{+150}_{-65}	^{12,13} COUPLAND	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
165^{+18}_{-8}	¹³ ALSPECTOR	73	CNTR $\bar{p}p$ S channel
~ 190	ABRAMS	70	CNTR S channel $\bar{N}N$
¹¹ From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on $\bar{p}p \rightarrow \eta \pi^0 \pi^0$, $\pi^0 \pi^0$, $\eta\eta$, $\eta\eta'$, $\pi^+ \pi^-$.			
¹² From a fit to the total elastic cross section.			
¹³ Isospins 0 and 1 not separated.			

NODE=M041W2
 NODE=M041W2

NODE=M041W2;LINKAGE=AN

NODE=M041W2;LINKAGE=E
 NODE=M041W2;LINKAGE=I

 $\pi^- p \rightarrow \eta \pi \pi n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$235 \pm 50 \pm 40$	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

NODE=M041W3
 NODE=M041W3

 $\rho\rho$ CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	COMMENT
250 ± 80 OUR ESTIMATE		
260 ± 57	BARBERIS	00F 450 $\rho\rho \rightarrow p_f \omega \omega p_s$

NODE=M041W4
 NODE=M041W4
 → UNCHECKED ←

 $f_4(2300)$ DECAY MODES

NODE=M041215;NODE=M041

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\rho$	seen
Γ_2 $\omega\omega$	seen
Γ_3 $\eta\pi\pi$	seen
Γ_4 $\pi\pi$	seen
Γ_5 $K\bar{K}$	seen
Γ_6 $N\bar{N}$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6;OUR EST;→ UNCHECKED ←

 $f_4(2300)$ BRANCHING RATIOS

NODE=M041220

 $\Gamma(\rho\rho)/\Gamma(\omega\omega)$ **Γ_1/Γ_2**

VALUE	DOCUMENT ID	COMMENT
2.8 ± 0.5	BARBERIS	00F 450 $\rho\rho \rightarrow p_f \omega \omega p_s$

NODE=M041R1
 NODE=M041R1

$f_4(2300)$ REFERENCES

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>	
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER	78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE	78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)

NODE=M041

REFID=47432
 REFID=47950
 REFID=47962
 REFID=46903
 REFID=46926
 REFID=44103
 REFID=21838
 REFID=21837
 REFID=21964
 REFID=21733
 REFID=21850
 REFID=21963
 REFID=21830
 REFID=21813
 REFID=21807

 $f_0(2330)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

NODE=M169

 $f_0(2330)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2314 ± 25	¹ BUGG	04A	RVUE
2337 ± 14	ANISOVICH	00J	SPEC 2.0 $\bar{p}p \rightarrow \pi\pi, \eta\eta$
~ 2321	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
¹ Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.			

NODE=M169M

NODE=M169M

NODE=M169M;LINKAGE=BU

 $f_0(2330)$ WIDTH

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
144 ± 20	² BUGG	04A	RVUE
217 ± 33	ANISOVICH	00J	SPEC 2.0 $\bar{p}p \rightarrow \pi\pi, \eta\eta$
~ 223	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
² Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.			

NODE=M169W

NODE=M169W

NODE=M169W;LINKAGE=BU

 $f_0(2330)$ REFERENCES

BUGG	04A	EPJ C36 161	D.V. Bugg	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
BARNES	00	PR C62 055203	P.D. Barnes <i>et al.</i>	
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)

NODE=M169

REFID=50158
 REFID=47950
 REFID=47965
 REFID=44103

$f_2(2340)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M108

 $f_2(2340)$ MASS

NODE=M108M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M108M

2345⁺⁵⁰₋₄₀ OUR AVERAGE

2362 ⁺³¹⁺¹⁴⁰ ₋₃₀₋₆₃	5.5k	1 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
2339 \pm 55		2 ETKIN	88 MPS	$22 \pi^- p \rightarrow \phi\phi n$
2350 \pm 7	80k	3 UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
2392 \pm 10		BOOTH	86 OMEG	$85 \pi^- Be \rightarrow 2\phi Be$
2360 \pm 20		LINDENBAUM	84 RVUE	

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M108M;LINKAGE=A

² Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi$ 2^{++} S_2 , D_2 , and D_0 is 37 ± 19 , 4_{-4}^{+12} , and 59_{-19}^{+21} , respectively.

NODE=M108M;LINKAGE=C

³ Statistical error only.

NODE=M108M;LINKAGE=ST

 $f_2(2340)$ WIDTH

NODE=M108W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M108W

322⁺⁷⁰₋₆₀ OUR AVERAGE

334 ⁺⁶²⁺¹⁶⁵ ₋₅₄₋₁₀₀	5.5k	4 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
319 ⁺⁸¹ ₋₆₉		5 ETKIN	88 MPS	$22 \pi^- p \rightarrow \phi\phi n$
218 \pm 16	80k	6 UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
198 \pm 50		BOOTH	86 OMEG	$85 \pi^- Be \rightarrow 2\phi Be$
150 ⁺¹⁵⁰ ₋₅₀		LINDENBAUM	84 RVUE	

⁴ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M108W;LINKAGE=A

⁵ Includes data of ETKIN 85.

NODE=M108W;LINKAGE=C

⁶ Statistical error only.

NODE=M108W;LINKAGE=ST

 $f_2(2340)$ DECAY MODES

NODE=M108215;NODE=M108

Mode	Fraction (Γ_i/Γ)
Γ_1 $\phi\phi$	seen
Γ_2 $\eta\eta$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2 $f_2(2340)$ BRANCHING RATIOS

NODE=M108220

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$

NODE=M108R01
NODE=M108R01 $f_2(2340)$ REFERENCES

NODE=M108

ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55387
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=40285
BOOTH	86	NP B273 677	P.S.L. Booth <i>et al.</i>	(LIVP, GLAS, CERN)	REFID=21870
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21871
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)	REFID=21869

$\rho_5(2350)$

$$I^G(J^{PC}) = 1^+(5^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_1(2400)$. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $f_4(2300)$.

NODE=M033

NODE=M033

 $\rho_5(2350)$ MASS

NODE=M033205

NODE=M033M

 $\pi^- p \rightarrow \omega \pi^0 n$ NODE=M033M3
NODE=M033M3

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2330±35	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$

NODE=M033M1

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 2303	HASAN	94	RVUE	$\bar{p} p \rightarrow \pi \pi$
~ 2300	¹ MARTIN	80B	RVUE	
~ 2250	¹ MARTIN	80C	RVUE	
~ 2500	² CARTER	78B	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow K^- K^+$
~ 2480	³ CARTER	77	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow \pi \pi$

S-CHANNEL $\bar{N} N$ NODE=M033M2
NODE=M033M2

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2300±45	⁴ ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega \pi^0$, $\omega \eta \pi^0, \pi^+ \pi^-$
2295±30	ANISOVICH	00J	SPEC	
~ 2380	⁵ CUTTS	78B	CNTR	0.97-3 $\bar{p} p \rightarrow \bar{N} N$
2345±15	^{5,6} COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow \bar{p} p$
2359± 2	^{5,7} ALSPECTOR	73	CNTR	$\bar{p} p$ S channel
2350±10	⁸ ABRAMS	70	CNTR	S channel $\bar{N} N$
2360±25	⁹ OH	70B	HDBC -0	$\bar{p}(p n), K^* K 2\pi$

 $\pi^- p \rightarrow K^+ K^- n$ NODE=M033M4
NODE=M033M4

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2307±6	ALPER	80	CNTR 0	62 $\pi^- p \rightarrow K^+ K^- n$
	¹			$I(J^P) = 1(5^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.
	²			$I = 0(1); J^P = 5^-$ from Barrelet-zero analysis.
	³			$I(J^P) = 1(5^-)$ from amplitude analysis.
	⁴			From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.
	⁵			Isospins 0 and 1 not separated.
	⁶			From a fit to the total elastic cross section.
	⁷			Referred to as U or U region by ALSPECTOR 73.
	⁸			For $I = 1 \bar{N} N$.
	⁹			No evidence for this bump seen in the $\bar{p} p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.

NODE=M033M1;LINKAGE=P
NODE=M033M1;LINKAGE=K
NODE=M033M1;LINKAGE=J
NODE=M033M2;LINKAGE=AYNODE=M033M2;LINKAGE=I
NODE=M033M2;LINKAGE=E
NODE=M033M2;LINKAGE=M
NODE=M033M2;LINKAGE=A
NODE=M033M2;LINKAGE=N **$\rho_5(2350)$ WIDTH**

NODE=M033210

 $\pi^- p \rightarrow \omega \pi^0 n$ NODE=M033W3
NODE=M033W3

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
400±100	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$

 $\bar{p} p \rightarrow \pi \pi$ or $\bar{K} K$ NODE=M033W1
NODE=M033W1

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 169	HASAN	94	RVUE	$\bar{p} p \rightarrow \pi \pi$
~ 250	¹⁰ MARTIN	80B	RVUE	
~ 300	¹⁰ MARTIN	80C	RVUE	
~ 150	¹¹ CARTER	78B	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow K^- K^+$
~ 210	¹² CARTER	77	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow \pi \pi$

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
260 ± 75	13 ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
235 ⁺⁶⁵ ₋₄₀	ANISOVICH	00J	SPEC	
135 ⁺¹⁵⁰ ₋₆₅	14,15 COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
165 ⁺¹⁸ ₋₈	15 ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
< 60	16 OH	70B	HDBC -0	$\bar{p}(pn), K^*K2\pi$
~ 140	ABRAMS	67C	CNTR	S channel $\bar{p}N$

NODE=M033W2
NODE=M033W2

$\pi^-p \rightarrow K^+K^-n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
245 ± 20	ALPER	80	CNTR 0	62 $\pi^-p \rightarrow K^+K^-n$
¹⁰ $I(J^P) = 1(5^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$. ¹¹ $I = 0(1); J^P = 5^-$ from Barrelet-zero analysis. ¹² $I(J^P) = 1(5^-)$ from amplitude analysis. ¹³ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02. ¹⁴ From a fit to the total elastic cross section. ¹⁵ Isospins 0 and 1 not separated. ¹⁶ No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.				

NODE=M033W4
NODE=M033W4

OCCUR=2

NODE=M033W1;LINKAGE=P
NODE=M033W1;LINKAGE=K
NODE=M033W1;LINKAGE=J
NODE=M033W2;LINKAGE=AY

NODE=M033W2;LINKAGE=E
NODE=M033W2;LINKAGE=I
NODE=M033W2;LINKAGE=N

$\rho_5(2350)$ REFERENCES

ANISOVICH 02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
ALDE 95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
ALPER 80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER 78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
CARTER 77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
OH 73	NP B51 57	B.Y. Oh <i>et al.</i>	(MSU)
CHAPMAN 71B	PR D4 1275	J.W. Chapman <i>et al.</i>	(MICH)
ABRAMS 70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
OH 70B	PRL 24 1257	B.Y. Oh <i>et al.</i>	(MSU)
ABRAMS 67C	PRL 18 1209	R.J. Abrams <i>et al.</i>	(BNL)

NODE=M033

REFID=48828
REFID=48327
REFID=48349
REFID=47950
REFID=44371
REFID=44103
REFID=21665
REFID=21838
REFID=21837
REFID=21964
REFID=21733
REFID=21963
REFID=21830
REFID=21813
REFID=21931
REFID=21926
REFID=21807
REFID=21925
REFID=21804

$a_6(2450)$

$$I^G(J^{PC}) = 1^-(6^{++})$$

OMITTED FROM SUMMARY TABLE
Needs confirmation.

NODE=M024

NODE=M024

$a_6(2450)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2450 ± 130	¹ CLELAND	82B	SPEC ±	50 $\pi p \rightarrow K_S^0 K^\pm p$
¹ From an amplitude analysis.				

NODE=M024M

NODE=M024M

NODE=M024M;LINKAGE=C

$a_6(2450)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
400 ± 250	² CLELAND	82B	SPEC ±	50 $\pi p \rightarrow K_S^0 K^\pm p$
² From an amplitude analysis.				

NODE=M024W

NODE=M024W

NODE=M024W;LINKAGE=C

$a_6(2450)$ DECAY MODES

NODE=M024215;NODE=M024

Mode	
Γ_1	$K\bar{K}$

DESIG=1

$a_6(2450)$ REFERENCES

CLELAND 82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
-------------	-------------	----------------------------	---------------------

NODE=M024

REFID=21281

$f_6(2510)$

$$I^G(J^{PC}) = 0^+(6^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M089

NODE=M089

NODE=M089M


NODE=M089M

NODE=M089M;LINKAGE=AN

 $f_6(2510)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2469±29 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
2485±40	¹ ANISOVICH	00J	SPEC 1.92-2.41 $p\bar{p}$
2420±30	ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
2510±30	BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2\pi^0$

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B.

 **$f_6(2510)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
283±40 OUR AVERAGE	Error includes scale factor of 1.1.		
410±90	² ANISOVICH	00J	SPEC 1.92-2.41 $p\bar{p}$
270±60	ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
240±60	BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2\pi^0$

² From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B.

NODE=M089W

NODE=M089W

NODE=M089W;LINKAGE=AN

 $f_6(2510)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi\pi$	(6.0±1.0) %

NODE=M089215;NODE=M089

DESIG=1

 $f_6(2510)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.06±0.01	³ BINON	83C	GAM2 38 $\pi^- p \rightarrow n 4\gamma$	

NODE=M089220

NODE=M089R1
NODE=M089R1

³ Assuming one pion exchange and using data of BOLOTOV 74.

NODE=M089R1;LINKAGE=A

f₀(2510) REFERENCES

ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99J	PL B471 271	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>	
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446.		
BINON	84B	LNC 39 41	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP) JP
BINON	83C	SJNP 38 723	F.G. Binon <i>et al.</i>	(SERP, BRUX+)
		Translated from YAF 38 1199.		
BOLOTOV	74	PL 52B 489	V.N. Bolotov <i>et al.</i>	(SERP)

NODE=M089

REFID=47942
REFID=47950
REFID=46903
REFID=46926
REFID=47416
REFID=47472
REFID=46605
REFID=46914REFID=21780
REFID=40288

REFID=44705

OTHER LIGHT MESONS

NODE=MXXX015

Further States

NODE=M300

OMITTED FROM SUMMARY TABLE

This section contains states observed by a single group or states poorly established that thus need confirmation.

NODE=M300

QUANTUM NUMBERS, MASSES, WIDTHS, AND BRANCHING RATIOS

X(360) $I^G(J^{PC}) = \text{?}^?(\text{?}^?+)$					
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
360 ± 7 ± 9	64 ± 18	2.3k	¹ ABRAAMYAN 09	CNTR	2.75 d C → γ γ X
¹ Not seen in p C → γ γ X at 5.5 GeV/c.					

NODE=M300K08
NODE=M300K08

NODE=M300K08;LINKAGE=AB

X(1070) $I^G(J^{PC}) = \text{?}^?(0^{++})$					
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>		<u>DOCUMENT ID</u>		<u>COMMENT</u>
1072 ± 1	3.5 ± 0.5		² VLADIMIRSK...08		40 π ⁻ p → K _S ⁰ K _S ⁰ n + m π ⁰
² Supersedes GRIGOR'EV 05.					

NODE=M300J07
NODE=M300J07

NODE=M300J07;LINKAGE=VL

X(1110) $I^G(J^{PC}) = 0^+(\text{even}^{++})$					
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1107 ± 4	111 ± 8 ± 15		DAFTARI	87	DBC 0. $\bar{p} n \rightarrow \rho^- \pi^+ \pi^-$

NODE=M300J30
NODE=M300J30

f₀(1200-1600) $I^G(J^{PC}) = 0^+(0^{++})$					
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1323 ± 8	237 ± 20		VLADIMIRSK...06	SPEC	40 π ⁻ p → K _S ⁰ K _S ⁰ n
1480 ⁺¹⁰⁰ ₋₁₅₀	1030 ⁺⁸⁰ ₋₁₇₀		³ ANISOVICH	03	SPEC
1530 ⁺⁹⁰ ₋₂₅₀	560 ± 40		⁴ ANISOVICH	03	SPEC

NODE=M300J98
NODE=M300J98

OCCUR=2

³ K-matrix pole from combined analysis of π⁻ p → π⁰ π⁰ n, π⁻ p → K \bar{K} n, π⁺ π⁻ → π⁺ π⁻, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta, \pi^+ \pi^- \pi^0, K^+ K^- \pi^0, K_S^0 K_S^0 \pi^0, K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+, K_S^0 K^- \pi^0, K_S^0 K_S^0 \pi^-$ at rest.

NODE=M300;LINKAGE=KM

⁴ K-matrix pole from combined analysis of π⁻ p → π⁰ π⁰ n, π⁻ p → K \bar{K} n, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$ at rest.

NODE=M300;LINKAGE=MK

X(1420) $I^G(J^{PC}) = 2^+(0^{++})$					
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1420 ± 20	160 ± 10		FILIPPI	00	OBLX 0 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$

NODE=M300J61
NODE=M300J61

X(1545) $I^G(J^{PC}) = \text{?}^?(\text{?}^?+)$					
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>		<u>DOCUMENT ID</u>		<u>COMMENT</u>
1545 ± 3	6.0 ± 2.5		⁵ VLADIMIRSK...08		40 π ⁻ p → K _S ⁰ K _S ⁰ n + m π ⁰

NODE=M300K07
NODE=M300K07

⁵Supersedes VLADIMIRSKII 00.

NODE=M300K07;LINKAGE=VL

X(1575) $I^G(J^{PC}) = ?^?(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1576^{+49+98}_{-55-91}	$818^{+22+64}_{-23-133}$	⁶ ABLIKIM	06S BES	$J/\psi \rightarrow K^+ K^- \pi^0$	

NODE=M300J08
NODE=M300J08

⁶A broad peak observed at $K^+ K^-$ invariant mass. Mass and width above are its pole position. The observed branching ratio is $B(J/\psi \rightarrow X \pi^0) B(X \rightarrow K^+ K^-) = (8.5 \pm 0.6^{+2.7}_{-3.6}) \times 10^{-4}$.

NODE=M300J08;LINKAGE=AB

X(1600) $I^G(J^{PC}) = 2^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1600 ± 100	400 ± 200	⁷ ALBRECHT	91F ARG	$10.2 e^+ e^- \rightarrow e^+ e^- 2(\pi^+ \pi^-)$	

NODE=M300J99
NODE=M300J99⁷Our estimate.

NODE=M300J99;LINKAGE=A

X(1650) $I^G(J^{PC}) = 0^-(?^?-)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1652 ± 7	<50	100	PROKOSHKIN 96	GAM2	$32,38 \pi p \rightarrow \omega \eta n$

NODE=M300J62
NODE=M300J62

X(1730) $I^G(J^{PC}) = ?^?(?^?+)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$1731.0 \pm 1.2 \pm 2.0$	$3.2 \pm 0.8 \pm 1.3$	58	VLADIMIRSK...07	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 X$

NODE=M300K06
NODE=M300K06

X(1750) $I^G(J^{PC}) = ?^?(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1753.5 \pm 1.5 \pm 2.3$	$122.2 \pm 6.2 \pm 8.0$	LINK	02K FOCS	$20-160 \gamma p \rightarrow K^+ K^- p$	

NODE=M300J94
NODE=M300J94
$$B(X(1750) \rightarrow \bar{K}^*(892)^0 K^0 \rightarrow K^\pm \pi^\mp K_S^0) / B(X(1750) \rightarrow K^+ K^-)$$

VALUE	CL%	DOCUMENT ID	TECN
<0.065	90	LINK	02K FOCS

NODE=M300B5
NODE=M300B5
$$B(X(1750) \rightarrow \bar{K}^*(892)^\pm K^\mp \rightarrow K^\pm \pi^\mp K_S^0) / B(X(1750) \rightarrow K^+ K^-)$$

VALUE	CL%	DOCUMENT ID	TECN
<0.183	90	LINK	02K FOCS

NODE=M300B6
NODE=M300B6

f₂(1750) $I^G(J^{PC}) = 0^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1755 ± 10	67 ± 12	870	⁸ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300JAM
NODE=M300JAM **$\Gamma(K\bar{K})$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
17 ± 5	870	⁹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300JA1
NODE=M300JA1 **$\Gamma(\gamma\gamma)$**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.04	870	⁹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300JA2
NODE=M300JA2 **$\Gamma(\pi\pi)$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.3 ± 1.0	870	⁹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300JA3
NODE=M300JA3 **$\Gamma(\eta\eta)$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.0 ± 0.5	870	⁹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300JA4
NODE=M300JA4

⁸From analysis of L3 data at 91 and 183–209 GeV.

⁹From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M300JAM;LINKAGE=SC
NODE=M300JA;LINKAGE=SC

X(1775) $I^G(J^{PC}) = 1^-(?^-+)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1763±20	192 ± 60	CONDO	91	SHF $\gamma p \rightarrow (p\pi^+)(\pi^+\pi^-\pi^-)$	
1787±18	118 ± 60	CONDO	91	SHF $\gamma p \rightarrow n\pi^+\pi^+\pi^-$	

NODE=M300J60
NODE=M300J60

OCCUR=2

f₀(1800) $I^G(J^{PC}) = 0^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1795± 7 ⁺²³ ₋₂₀	95 ± 10 ⁺⁷⁸ ₋₈₂	ABLIKIM	13J	BES3 $J/\psi \rightarrow \gamma\omega\phi$	
1812 ⁺¹⁹ ₋₂₆ ±18	105 ± 20 ± 28	¹⁰ ABLIKIM	06J	BES2 $J/\psi \rightarrow \gamma\omega\phi$	

NODE=M300K29
NODE=M300K29

¹⁰Not seen by LIU 09 in $B^\pm \rightarrow K^\pm\omega\phi$.

NODE=M300K29;LINKAGE=AB

X(1850 - 3100) $I^G(J^{PC}) = ?^?(1^{--})$					
$\Gamma(e^+e^-)B(X \rightarrow \text{hadrons})$ (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<120	90	¹¹ ANASHIN	11	KEDR $e^+e^- \rightarrow \text{hadrons}$	

NODE=M300K28
NODE=M300K28

¹¹This limit is center-of-mass energy dependent. We quote the most stringent one.

NODE=M300K28;LINKAGE=AN

X(1855) $I^G(J^{PC}) = ?^?(???)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1856.6±5	20 ± 5	BRIDGES	86D	SPEC $0. \bar{p}d \rightarrow \pi\pi N$	

NODE=M300J31
NODE=M300J31

X(1870) $I^G(J^{PC}) = ?^?(2^{??})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1870±40	250 ± 30	ALDE	86D	GAM4 $100 \pi^- p \rightarrow 2\eta X$	

NODE=M300J45
NODE=M300J45

a₃(1875) $I^G(J^{PC}) = 1^-(3^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1874±43±96	385 ± 121 ± 114	CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^-p$	

NODE=M300J95
NODE=M300J95

B(a₃(1875) → f₂(1270)π)/B(a₃(1875) → ρπ)					
VALUE	DOCUMENT ID	TECN	COMMENT		
0.8±0.2	¹² CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^-p$		

NODE=M300B7
NODE=M300B7

¹²Using the observable fractions of 50.0% ρπ, 56.5% f₂π, and 11.8% ρ₃π.

NODE=M300B;LINKAGE=C1

B(a₃(1875) → ρ₃(1690)π)/B(a₃(1875) → ρπ)					
VALUE	DOCUMENT ID	TECN	COMMENT		
0.9±0.3	¹³ CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^-p$		

NODE=M300B8
NODE=M300B8

¹³Using the observable fractions of 50.0% ρπ, 56.5% f₂π, and 11.8% ρ₃π.

NODE=M300B8;LINKAGE=C1

a₁(1930) $I^G(J^{PC}) = 1^-(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1930 ⁺³⁰ ₋₇₀	155 ± 45	ANISOVICH	01F	SPEC $2.0 \bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$	

NODE=M300J92
NODE=M300J92

X(1935) $I^G(J^{PC}) = 1^+(1^{-?})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1935±20	215 ± 30	EVANGELIS...	79	OMEG $10,16 \pi^- p \rightarrow \bar{p}pn$	

NODE=M300J33
NODE=M300J33

ρ₂(1940) $I^G(J^{PC}) = 1^+(2^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1940±40	155 ± 40	¹⁴ ANISOVICH	02	SPEC $0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$	

NODE=M300J85
NODE=M300J85

¹⁴ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J85;LINKAGE=AY

$\omega_3(1945)$ $I^G(J^{PC}) = 0^-(3^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1945±20	115 ± 22	15 ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J65
NODE=M300J65

¹⁵ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J65;LINKAGE=AZ

$a_2(1950)$ $I^G(J^{PC}) = 1^-(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1950 ⁺³⁰ ₋₇₀	180 ⁺³⁰ ₋₇₀	16 ANISOVICH	01F	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K24
NODE=M300K24

¹⁶ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K24;LINKAGE=AN

$\omega(1960)$ $I^G(J^{PC}) = 0^-(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1960±25	195 ± 60	17 ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J79
NODE=M300J79

¹⁷ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J79;LINKAGE=AZ

$b_1(1960)$ $I^G(J^{PC}) = 1^+(1^{+-})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1960±35	230 ± 50	18 ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J67
NODE=M300J67

¹⁸ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J67;LINKAGE=AY

$h_1(1965)$ $I^G(J^{PC}) = 0^-(1^{+-})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1965±45	345 ± 75	19 ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J64
NODE=M300J64

¹⁹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J64;LINKAGE=AZ

$f_1(1970)$ $I^G(J^{PC}) = 0^+(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1971±15	240 ± 45	ANISOVICH	00J	SPEC	

NODE=M300J1
NODE=M300J1

$X(1970)$ $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1970±10	40 ± 20	CHLIAPNIK...	80	HBC	32 $K^+p \rightarrow 2K_S^0 2\pi X$

NODE=M300J46
NODE=M300J46

$X(1975)$ $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1973±15	80	30	CASO	70	HBC 11.2 $\pi^-p \rightarrow \rho 2\pi$

NODE=M300J47
NODE=M300J47

$\omega_2(1975)$ $I^G(J^{PC}) = 0^-(2^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1975±20	175 ± 25	20 ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J81
NODE=M300J81

²⁰ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J81;LINKAGE=AZ

$a_2(1990)$ $I^G(J^{PC}) = 1^-(2^{++})$					
MASS (MeV)	WIDTH (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2050±10±40	190 ± 22 ± 100	18k	21 SCHEGELSKY	06	RVUE $\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
2003±10±19	249 ± 23 ± 32		LU	05	B852 18 $\pi^-p \rightarrow \omega\pi^-\pi^0 p$

NODE=M300J2
NODE=M300J2

²¹ From analysis of L3 data at 183–209 GeV.

NODE=M300J2;LINKAGE=SC

$\Gamma(\gamma\gamma) \Gamma(\pi^+ \pi^- \pi^0) / \Gamma(\text{total})$

VALUE (keV)	EVTs	DOCUMENT ID	TECN	COMMENT
$0.11 \pm 0.04 \pm 0.05$	18k	²² SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

²² From analysis of L3 data at 183–209 GeV.

NODE=M300J2G
 NODE=M300J2G

NODE=M300J2G;LINKAGE=SC

$\rho(2000)$ $I^G(J^{PC}) = 1^+(1^{--})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2000 ± 30	260 ± 45	²³ BUGG	04C	RVUE Compilation
~ 1988	~ 244	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

²³ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J77
 NODE=M300J77

NODE=M300;LINKAGE=AY

$f_2(2000)$ $I^G(J^{PC}) = 0^+(2^{++})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2001 ± 10	312 ± 32	ANISOVICH	00J	SPEC
~ 1996	~ 134	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

NODE=M300J25
 NODE=M300J25

$X(2000)$ $I^G(J^{PC}) = 1^-(?^{?+})$		DOCUMENT ID	TECN	CHG	COMMENT
MASS (MeV)	WIDTH (MeV)				
1964 ± 35	225 ± 50	²⁴ ARMSTRONG	93D	E760	$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
~ 2100	~ 500	²⁴ ANTIPOV	77	CIBS	- $25 \pi^- p \rightarrow p\pi^- \rho_3$
2214 ± 15	355 ± 21	²⁵ BALTAY	77	HBC	0 $15 \pi^- p \rightarrow \Delta^{++} 3\pi$
2080 ± 40	340 ± 80	KALELKAR	75	HBC	+ $15 \pi^+ p \rightarrow p\pi^+ \rho_3$

NODE=M300K01
 NODE=M300K01

²⁴ Cannot determine spin to be 3.

²⁵ BALTAY 77 favors $J^P = ,3^+$.

NODE=M300K01;LINKAGE=AA
 NODE=M300K01;LINKAGE=B

$X(2000)$ $I^G(J^{PC}) = ?^?(4^{++})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
$1998 \pm 3 \pm 5$	<15	VLADIMIRSK...03	SPEC	$\pi^- p \rightarrow K_S^0 K_S^0 M M$

NODE=M300J97
 NODE=M300J97

$\pi_2(2005)$ $I^G(J^{PC}) = 1^-(2^{-+})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)	EVTs		
$1974 \pm 14 \pm 83$	$341 \pm 61 \pm 139$	145k	LU	05 B852 $18 \pi^- p \rightarrow \omega\pi^- \pi^0 p$
2005 ± 15	200 ± 40		ANISOVICH	01F SPEC $2.0 \bar{p}p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$

NODE=M300J71
 NODE=M300J71

$\eta(2010)$ $I^G(J^{PC}) = 0^+(0^{-+})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2010^{+35}_{-60}	270 ± 60	ANISOVICH	00J	SPEC

NODE=M300J5
 NODE=M300J5

$\pi_1(2015)$ $I^G(J^{PC}) = 1^-(1^{-+})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)	EVTs		
$2014 \pm 20 \pm 16$	$230 \pm 32 \pm 73$	145k	LU	05 B852 $18 \pi^- p \rightarrow \omega\pi^- \pi^0 p$
$2001 \pm 30 \pm 92$	$333 \pm 52 \pm 49$	69k	KUHN	04 B852 $18 \pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$

NODE=M300J05
 NODE=M300J05

$a_0(2020)$ $I^G(J^{PC}) = 1^-(0^{++})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2025 ± 30	330 ± 75	ANISOVICH	99C	SPEC

NODE=M300J6
 NODE=M300J6

$X(2020)$ $I^G(J^{PC}) = ?^?(???)$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2015 ± 3	10 ± 4	FERRER	99	RVUE $\pi p \rightarrow p\rho\bar{p}\pi(\pi)$

NODE=M300J34
 NODE=M300J34

$h_3(2025)$ $I^G(J^{PC}) = 0^-(3^{+-})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2025 ± 20	145 ± 30	²⁶ ANISOVICH	02B	SPEC $0.6-1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J78
 NODE=M300J78

²⁶ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J78;LINKAGE=AZ

$b_3(2030)$		$I^G(J^{PC}) = 1^+(3^-)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2032±12	117 ± 11	²⁷ ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$	

NODE=M300J69
NODE=M300J69

²⁷ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J69;LINKAGE=AY

$a_2(2030)$		$I^G(J^{PC}) = 1^-(2^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2030±20	205 ± 30	²⁸ ANISOVICH 01F	SPEC	1.96–2.41 $\bar{p}p$	

NODE=M300K23
NODE=M300K23

²⁸ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K23;LINKAGE=AN

$a_3(2030)$		$I^G(J^{PC}) = 1^-(3^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2031±12	150 ± 18	²⁹ ANISOVICH 01F	SPEC	1.96–2.41 $\bar{p}p$	

NODE=M300K20
NODE=M300K20

²⁹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K20;LINKAGE=AN

$\eta_2(2030)$		$I^G(J^{PC}) = 0^+(2^{-+})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2030±5±15	205 ± 10 ± 15	ANISOVICH 00E	SPEC		

NODE=M300J8
NODE=M300J8

$B(a_2\pi)_{L=0}/B(a_2\pi)_{L=2}$					
VALUE		DOCUMENT ID	TECN	COMMENT	
0.05±0.03		³⁰ ANISOVICH 11	SPEC	0.9–1.94 $p\bar{p}$	

NODE=M300B1
NODE=M300B1

³⁰ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B1;LINKAGE=AN

$B(a_0\pi)/B(a_2\pi)_{L=2}$					
VALUE		DOCUMENT ID	TECN	COMMENT	
0.10±0.08		³¹ ANISOVICH 11	SPEC	0.9–1.94 $p\bar{p}$	

NODE=M300B2
NODE=M300B2

³¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B2;LINKAGE=AN

$B(f_2\eta)/B(a_2\pi)_{L=2}$					
VALUE		DOCUMENT ID	TECN	COMMENT	
0.13±0.06		³² ANISOVICH 11	SPEC	0.9–1.94 $p\bar{p}$	

NODE=M300B3
NODE=M300B3

³² Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B3;LINKAGE=AN

$f_3(2050)$		$I^G(J^{PC}) = 0^+(3^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2048±8	213 ± 34	ANISOVICH 00J	SPEC	2.0 $p\bar{p} \rightarrow \eta\pi^0\pi^0$	

NODE=M300J7
NODE=M300J7

$f_0(2060)$		$I^G(J^{PC}) = 0^+(0^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
~ 2050	~ 120	³³ OAKDEN 94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$	
~ 2060	~ 50	³³ OAKDEN 94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$	

NODE=M300J59
NODE=M300J59

³³ See SEMENOV 99 and KLOET 96.

OCCUR=2

NODE=M300J;LINKAGE=A

$\pi(2070)$		$I^G(J^{PC}) = 1^-(0^{-+})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2070±35	310 $^{+100}_{-50}$	ANISOVICH 01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$	

NODE=M300J91
NODE=M300J91

$X(2075)$		$I^G(J^{PC}) = ??(???)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2075±12±5	90 ± 35 ± 9	³⁴ ABLIKIM 04J	BES2	$J/\psi \rightarrow K^- p\bar{\Lambda}$	

NODE=M300J01
NODE=M300J01

³⁴ From a fit in the region $M_{p\bar{\Lambda}} - M_p - M_{\Lambda} < 150$ MeV. S-wave in the $p\bar{\Lambda}$ system preferred.

A similar near-threshold enhancement in the $p\bar{\Lambda}$ system is observed in $B^+ \rightarrow p\bar{\Lambda}\bar{D}^0$ by CHEN 11F.

NODE=M300J01;LINKAGE=AB

X(2080)		$I^G(J^{PC}) = ??(???)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2080 ± 10	110 ± 20	KREYMER	80	STRC	13 $\pi^- d \rightarrow p\bar{p}n(n_s)$

NODE=M300J35
NODE=M300J35

X(2080)		$I^G(J^{PC}) = ??(3^{-?})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2080 ± 10	190 ± 15	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p\bar{p}n$

NODE=M300J37
NODE=M300J37

a₁(2095)		$I^G(J^{PC}) = 1^-(1^{++})$			
MASS (MeV)	WIDTH (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2096 ± 17 ± 121	451 ± 41 ± 81	69k	KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^-p$

NODE=M300J04
NODE=M300J04

B(a₁(2095) → f₁(1285)π) / B(a₁(2095) → a₁(1260))					
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
3.18 ± 0.64	69k	KUHN	04	B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^-p$

NODE=M300B03
NODE=M300B03

η(2100)		$I^G(J^{PC}) = 0^+(0^{-+})$			
MASS (MeV)	WIDTH (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2103 ± 50	187 ± 75	586	³⁵ BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

NODE=M300J48
NODE=M300J48

³⁵ ASTON 81B sees no peak, has 850 events in Ajinenko+Barth bins. ARESTOV 80 sees no peak.

NODE=M300J;LINKAGE=A1

X(2100)		$I^G(J^{PC}) = ??(0^{??})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2100 ± 40	250 ± 40	ALDE	86D	GAM4	100 $\pi^- p \rightarrow 2\eta X$

NODE=M300J49
NODE=M300J49

X(2110)		$I^G(J^{PC}) = 1^+(3^{-?})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2110 ± 10	330 ± 20	EVANGELIS...	79	OMEG	10,16 $\pi^- p \rightarrow \bar{p}pn$

NODE=M300J36
NODE=M300J36

f₂(2140)		$I^G(J^{PC}) = 0^+(2^{++})$			
MASS (MeV)	WIDTH (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2141 ± 12	49 ± 28	389	GREEN	86	MPSF 400 $pA \rightarrow 4KX$

NODE=M300J50
NODE=M300J50

X(2150)		$I^G(J^{PC}) = ??(2^{+?})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2150 ± 10	260 ± 10	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p\bar{p}n$

NODE=M300J38
NODE=M300J38

a₂(2175)		$I^G(J^{PC}) = 1^-(2^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2175 ± 40	310 ⁺⁹⁰ ₋₄₅	ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J88
NODE=M300J88

η(2190)		$I^G(J^{PC}) = 0^+(0^{-+})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2190 ± 50	850 ± 100	BUGG	99	BES	

NODE=M300J13
NODE=M300J13

ω₂(2195)		$I^G(J^{PC}) = 0^-(2^{--})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2195 ± 30	225 ± 40	³⁶ ANISOVICH	02B	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J82
NODE=M300J82

³⁶ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J82;LINKAGE=AZ

$\omega(2205)$		$I^G(J^{PC}) = 0^-(1^--)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2205 ± 30	350 ± 90	³⁷ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J80
NODE=M300J80

³⁷ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J80;LINKAGE=AZ

X(2210)		$I^G(J^{PC}) = ??(???)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2210 ⁺⁷⁹ ₋₂₁	203 ⁺⁴³⁷ ₋₈₇	EVANGELIS...	79B	OMEG	10 $\pi^- p \rightarrow K^+ K^- n$

NODE=M300J39
NODE=M300J39

X(2210)		$I^G(J^{PC}) = ??(???)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2207 ± 22	130	CASO	70	HBC	11.2 $\pi^- p$

NODE=M300J51
NODE=M300J51

$h_1(2215)$		$I^G(J^{PC}) = 0^-(1^+-)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2215 ± 40	325 ± 55	³⁸ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J27
NODE=M300J27

³⁸ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J27;LINKAGE=AZ

$\rho_2(2225)$		$I^G(J^{PC}) = 1^+(2^--)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2225 ± 35	335 ⁺¹⁰⁰ ₋₅₀	³⁹ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J70
NODE=M300J70

³⁹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J70;LINKAGE=AY

$\rho_4(2230)$		$I^G(J^{PC}) = 1^+(4^--)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2230 ± 25	210 ± 30	⁴⁰ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J74
NODE=M300J74

⁴⁰ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J74;LINKAGE=AY

$b_1(2240)$		$I^G(J^{PC}) = 1^+(1^+-)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2240 ± 35	320 ± 85	⁴¹ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J87
NODE=M300J87

⁴¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J87;LINKAGE=AY

$f_2(2240)$		$I^G(J^{PC}) = 0^+(2^++)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2240 ± 15	241 ± 30	⁴² ANISOVICH	00J	SPEC	1.92–2.41 $p\bar{p}$

NODE=M300K26
NODE=M300K26

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2226 ~ 226 HASAN 94 RVUE $p\bar{p} \rightarrow \pi\pi$

⁴² From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

NODE=M300K26;LINKAGE=AN

$b_3(2245)$		$I^G(J^{PC}) = 1^+(3^+-)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2245 ± 50	320 ± 70	⁴³ BUGG	04C	RVUE	

NODE=M300K10
NODE=M300K10

⁴³ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300K10;LINKAGE=AY

$\eta_2(2250)$ $I^G(J^{PC}) = 0^+(2^-+)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2248 ± 20	280 ± 20	ANISOVICH 00I	SPEC	
2267 ± 14	290 ± 50	ANISOVICH 00J	SPEC	

NODE=M300J17
NODE=M300J17

$\pi_4(2250)$ $I^G(J^{PC}) = 1^-(4^-+)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 15	215 ± 25	ANISOVICH 01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J73
NODE=M300J73

$\omega_4(2250)$ $I^G(J^{PC}) = 0^-(4^{--})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 30	150 ± 50	⁴⁴ ANISOVICH 02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J84
NODE=M300J84

⁴⁴ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J84;LINKAGE=AZ

$\omega_5(2250)$ $I^G(J^{PC}) = 0^-(5^{--})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 70	320 ± 95	⁴⁵ BUGG 04	RVUE	

NODE=M300K11
NODE=M300K11

⁴⁵ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300;LINKAGE=AZ

$\omega_3(2255)$ $I^G(J^{PC}) = 0^-(3^{--})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255 ± 15	175 ± 30	⁴⁶ ANISOVICH 02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J66
NODE=M300J66

⁴⁶ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J66;LINKAGE=AZ

$a_4(2255)$ $I^G(J^{PC}) = 1^-(4^{++})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2237 ± 5	291 ± 12	UMAN	06	E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
2255 ± 40	330 ⁺¹¹⁰ ₋₅₀	⁴⁷ ANISOVICH 01F	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K21
NODE=M300K21

⁴⁷ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K21;LINKAGE=AN

$a_2(2255)$ $I^G(J^{PC}) = 1^-(2^{++})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255 ± 20	230 ± 15	⁴⁸ ANISOVICH 01G	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K22
NODE=M300K22

⁴⁸ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.

NODE=M300K22;LINKAGE=AN

$X(2260)$ $I^G(J^{PC}) = 0^+(4^{+?})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2260 ± 20	400 ± 100	EVANGELIS... 79	OMEG	10,16 $\pi^- p \rightarrow \bar{p}pn$

NODE=M300J40
NODE=M300J40

$\rho(2270)$ $I^G(J^{PC}) = 1^+(1^{--})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2265 ± 40	325 ± 80	⁴⁹ ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
2280 ± 50	440 ± 110	ATKINSON 85	OMEG	20–70 $\gamma p \rightarrow p\omega\pi^+\pi^-\pi^0$

NODE=M300J86
NODE=M300J86

⁴⁹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J86;LINKAGE=AY

a₁(2270) $I^G(J^{PC}) = 1^-(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2270 ⁺⁵⁵ ₋₄₀	305 ⁺⁷⁰ ₋₄₀	ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J72
NODE=M300J72

h₃(2275) $I^G(J^{PC}) = 0^-(3^{+-})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2275±25	190 ± 45	⁵⁰ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J28
NODE=M300J28

⁵⁰ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J28;LINKAGE=AZ

a₃(2275) $I^G(J^{PC}) = 1^-(3^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2275±35	350 ⁺¹⁰⁰ ₋₅₀	⁵¹ ANISOVICH	01G	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K19
NODE=M300K19

⁵¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.

NODE=M300K19;LINKAGE=AN

π₂(2285) $I^G(J^{PC}) = 1^-(2^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2285±20±25	250 ± 20 ± 25	⁵² ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$

NODE=M300K25
NODE=M300K25

⁵² Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300K25;LINKAGE=AN

ω₃(2285) $I^G(J^{PC}) = 0^-(3^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2278±28	224 ± 50	⁵³ BUGG	04A	RVUE	
2285±60	230 ± 40	⁵⁴ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J83
NODE=M300J83

⁵³ Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.

NODE=M300J83;LINKAGE=BU

⁵⁴ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J83;LINKAGE=AZ

ω(2290) $I^G(J^{PC}) = 0^-(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2290±20	275 ± 35	⁵⁵ BUGG	04A	RVUE	

NODE=M300J02
NODE=M300J02

⁵⁵ Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.

NODE=M300J02;LINKAGE=BU

f₂(2295) $I^G(J^{PC}) = 0^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2293±13	216 ± 37	⁵⁶ ANISOVICH	00J	SPEC	1.92–2.41 $p\bar{p}$

NODE=M300K27
NODE=M300K27

⁵⁶ From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

NODE=M300K27;LINKAGE=AN

f₃(2300) $I^G(J^{PC}) = 0^+(3^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2334±25	200 ± 20	⁵⁷ BUGG	04A	RVUE	

NODE=M300J19
NODE=M300J19

⁵⁷ Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.

NODE=M300J19;LINKAGE=BU

f₁(2310) $I^G(J^{PC}) = 0^+(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2310±60	255 ± 70	ANISOVICH	00J	SPEC	

NODE=M300J23
NODE=M300J23

η(2320) $I^G(J^{PC}) = 0^+(0^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2320±15	230 ± 35	⁵⁸ ANISOVICH	00M	SPEC	

NODE=M300J18
NODE=M300J18

⁵⁸ From the combined analysis of $\bar{p}p \rightarrow \eta\eta\eta$ from ANISOVICH 00M and $\bar{p}p \rightarrow \eta\pi^0\pi^0$ from ANISOVICH 00J.

NODE=M300;LINKAGE=B

$\eta_4(2330)$		$I^G(J^{PC}) = 0^+(4^-)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2328±38	240 ± 90	ANISOVICH	00J	SPEC	2.0 $p\bar{p} \rightarrow \eta\pi^0\pi^0$	

NODE=M300J22
NODE=M300J22

$\omega(2330)$		$I^G(J^{PC}) = 0^-(1^-)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2330±30	435 ± 75	ATKINSON	88	OMEG	25-50 $\gamma p \rightarrow \rho^\pm \rho^0 \pi^\mp$	

NODE=M300J53
NODE=M300J53

$X(2340)$		$I^G(J^{PC}) = ??(??)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2340±20	180 ± 60	126	⁵⁹ BALTAY	75	HBC	15 $\pi^+ p \rightarrow p 5\pi$

NODE=M300J54
NODE=M300J54

⁵⁹ Dominant decay into $\rho^0 \rho^0 \pi^+$. BALTAY 78 finds confirmation in $2\pi^+ \pi^- 2\pi^0$ events which contain $\rho^+ \rho^0 \pi^0$ and $2\rho^+ \pi^-$.

NODE=M300J;LINKAGE=B1

$\pi(2360)$		$I^G(J^{PC}) = 1^-(0^-)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2360±25	300^{+100}_{-50}	ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$	

NODE=M300J90
NODE=M300J90

$X(2360)$		$I^G(J^{PC}) = ??(4^?)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2360±10	430 ± 30	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p\bar{p}n$	

NODE=M300J42
NODE=M300J42

$X(2440)$		$I^G(J^{PC}) = ??(5^-)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2440±10	310 ± 20	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p\bar{p}n$	

NODE=M300J43
NODE=M300J43

$X(2540)$		$I^G(J^{PC}) = 0^+(0^+)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2539±14 ⁺³⁸ ₋₁₄	$274^{+77+126}_{-61-163}$	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M300K30
NODE=M300K30

$\Gamma(\gamma\gamma) \times B(K\bar{K})$						
VALUE (eV)	DOCUMENT ID	TECN	COMMENT			
40^{+9+17}_{-7-40}	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$		

NODE=M300K3G
NODE=M300K3G

$X(2632)$		$I^G(J^{PC}) = ??(??)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2635.2±3.3		⁶⁰ EVDOKIMOV	04	SELX	$X(2632) \rightarrow D_s^+ \eta$	
2631.6±2.1	< 17	⁶¹ EVDOKIMOV	04	SELX	$X(2632) \rightarrow D_s^0 K^+$	

NODE=M300J03
NODE=M300J03

OCCUR=2

⁶⁰ From a mass difference to D_s^+ of 666.9 ± 3.3 MeV.

NODE=M300J03;LINKAGE=EV
NODE=M300J03;LINKAGE=ED

⁶¹ From a mass difference to D_s^0 of 767.0 ± 2.0 MeV.

$B(X(2632) \rightarrow D_s^0 K^+)/B(X(2632) \rightarrow D_s^+ \eta)$						
VALUE	DOCUMENT ID	TECN	COMMENT			
0.14±0.06	⁶² EVDOKIMOV	04	SELX			

NODE=M300B01
NODE=M300B01

⁶² Possible interpretation of this decay pattern is discussed by YASUI 07.

NODE=M300B01;LINKAGE=YA

$X(2680)$		$I^G(J^{PC}) = ??(??)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2676±27	150	CASO	70	HBC	11.2 $\pi^- p \rightarrow \rho^- \pi^+ \pi^- p$	

NODE=M300J55
NODE=M300J55

X(2710) $I^G(J^{PC}) = ??(6^{+?})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2710±20	170 ± 40	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p \bar{p} n$

NODE=M300J44
NODE=M300J44

X(2750) $I^G(J^{PC}) = ??(7^{-?})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2747±32	195 ± 75	DENNEY	83	LASS	10 $\pi^+ p \rightarrow K^+ K^- \pi^+ p$

NODE=M300J56
NODE=M300J56

f₆(3100) $I^G(J^{PC}) = 0^+(6^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
3100±100	700 ± 130	BINON	05	GAMS	33 $\pi^- p \rightarrow \eta \eta n$

NODE=M300J06
NODE=M300J06

X(3250) $I^G(J^{PC}) = ??(???)$ 3-Body Decays					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
3250±8±20	45 ± 18	ALEEV	93	BIS2	$X(3250) \rightarrow \Lambda \bar{p} K^+$
3265±7±20	40 ± 18	ALEEV	93	BIS2	$X(3250) \rightarrow \bar{\Lambda} p K^-$

NODE=M300J57
NODE=M300J57

OCCUR=2

X(3250) $I^G(J^{PC}) = ??(???)$ 4-Body Decays					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
3245±8±20	25 ± 11	ALEEV	93	BIS2	$X(3250) \rightarrow \Lambda \bar{p} K^+ \pi^\pm$
3250±9±20	50 ± 20	ALEEV	93	BIS2	$X(3250) \rightarrow \bar{\Lambda} p K^- \pi^\mp$
3270±8±20	25 ± 11	ALEEV	93	BIS2	$X(3250) \rightarrow K_S^0 p \bar{p} K^\pm$

NODE=M300J58
NODE=M300J58

OCCUR=2

OCCUR=3

X(3350) $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3350 ⁺¹⁰ ₋₂₀ ±20	70 ⁺⁴⁰ ₋₃₀ ± 40	50 ± 10	⁶³ GABYSHEV	06A	BELL $B^- \rightarrow \Lambda_c^+ \bar{p} \pi^-$

NODE=M300J09
NODE=M300J09

⁶³A similar enhancement in the $\Lambda_c^+ \bar{p}$ final state is also reported by BABAR collaboration in AUBERT 10H.

NODE=M300J09;LINKAGE=AU

REFERENCES for Further States

ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54955
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ANISOVICH	12	PR D85 014001	A.V. Anisovich <i>et al.</i>		REFID=53961
ANASHIN	11	PL B703 543	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53932
ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
CHEN	11F	PR D84 071501	P. Chen <i>et al.</i>	(BELLE Collab.)	REFID=53814
AUBERT	10H	PR D82 031102	B. AUBERT <i>et al.</i>	(BABAR Collab.)	REFID=53363
ABRAAMYAN	09	PR C80 034001	Kh.U. Abraamyan <i>et al.</i>		REFID=53100
LIU	09	PR D79 071102	C. Liu <i>et al.</i>	(BELLE Collab.)	REFID=52752
VLADIMIRSK...	08	PAN 71 2129	V.V. Vladimirovsky <i>et al.</i>	(ITEP)	REFID=52681
VLADIMIRSK...	07	Translated from YAF 71 2166. PAN 70 1706	V. Vladimirovsky <i>et al.</i>		REFID=52058
YASUI	07	PR D76 034009	S. Yasui, M. Oka		REFID=51907
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM	06S	PRL 97 142002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51451
GABYSHEV	06A	PRL 97 242001	N. Gabyshev <i>et al.</i>	(BELLE Collab.)	REFID=51565
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
BINON	05	Translated from YAF 69 515. PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
GRIGOR'EV	05	Translated from YAF 68 998. PAN 68 1271	V.K. Grigor'ev <i>et al.</i>	(ITEP)	REFID=50844
LU	05	Translated from YAF 68 1324. PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ABLIKIM	04J	PRL 93 112002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50196
BUGG	04	PL B595 556 (err.)	D.V. Bugg		REFID=49763
BUGG	04A	EPJ C36 161	D.V. Bugg		REFID=50158
BUGG	04C	PRPL 397 257	D.V. Bugg		REFID=50203
EVDOKIMOV	04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>	(SELEX Collab.)	REFID=50337
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)	REFID=49773
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
VLADIMIRSK...	03	PAN 66 700	V.V. Vladimirovsky <i>et al.</i>		REFID=49419
		Translated from YAF 66 729.			

NODE=M300

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>		REFID=48828
ANISOVICH	02B	PL B542 19	A.V. Anisovich <i>et al.</i>		REFID=48829
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=48845
ANISOVICH	01C	PL B507 23	A.V. Anisovich <i>et al.</i>		REFID=48325
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>		REFID=48327
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>		REFID=48349
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352
ANISOVICH	01G	PL B517 273	A.V. Anisovich <i>et al.</i>		REFID=48353
ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>		REFID=47942
ANISOVICH	00D	PL B476 15	A.V. Anisovich <i>et al.</i>		REFID=47944
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
ANISOVICH	00I	PL B491 40	A.V. Anisovich <i>et al.</i>		REFID=47949
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>		REFID=47950
ANISOVICH	00M	PL B496 145	A.V. Anisovich <i>et al.</i>		REFID=48009
BARNES	00	PR C62 055203	P.D. Barnes <i>et al.</i>		REFID=47965
FILIPPI	00	PL B495 284	A. Filippi <i>et al.</i>	(OBELIX Experiment)	REFID=48006
VLADIMIRSKII	00	JETPL 72 486	V.V. Vladimirkii <i>et al.</i>		REFID=47997
		Translated from ZETFP 72 698.			
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>		REFID=46902
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>		REFID=46926
ANISOVICH	99J	PL B471 271	A.V. Anisovich <i>et al.</i>		REFID=47416
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>		REFID=47472
BUGG	99	PL B458 511	D.V. Bugg <i>et al.</i>		REFID=46938
FERRER	99	EPJ C10 249	A. Ferrer <i>et al.</i>		REFID=47404
SEME NOV	99	SPU 42 847	S.V. Semenov		REFID=47363
		Translated from UFN 42 937.			
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORDE)	REFID=45212
PROKOSHKIN	96	SPD 41 247	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=45182
		Translated from DANS 348 481.			
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ALEEV	93	PAN 56 1358	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)	REFID=43668
		Translated from YAF 56 100.			
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43596
ALBRECHT	91F	ZPHY C50 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41658
CONDO	91	PR D43 2787	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=41588
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
ATKINSON	88	ZPHY C38 535	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=40556
DAFTARI	87	PRL 58 859	I.K. Daftari <i>et al.</i>	(SYRA)	REFID=40412
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
BRIDGES	86D	PL B180 313	D.L. Bridges <i>et al.</i>	(SYRA, BNL, CASE+)	REFID=21984
GREEN	86	PRL 56 1639	D.R. Green <i>et al.</i>	(FNAL, ARIZ, FSU+)	REFID=21872
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=22000
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
ASTON	81B	NP B189 205	D. Aston <i>et al.</i>	(BONN, CERN, EPOL, GLAS+)	REFID=11553
ARESTOV	80	IHEP 80-165	Y.I. Arestov <i>et al.</i>	(SERP)	REFID=22312
CHLIAPNIK...	80	ZPHY C3 285	P.V. Chliapnikov <i>et al.</i>	(SERP, BRUX, MONS)	REFID=21996
KREYMER	80	PR D22 36	A.E. Kreymer <i>et al.</i>	(IND, PURD, SLAC+)	REFID=21970
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN)	REFID=21774
EVANGELIS...	79	NP B153 253	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21966
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21967
BALTAY	78	PR D17 52	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21569
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU)	REFID=20847
BALTAY	75	PRL 35 891	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21994
KALEL KAR	75	Thesis Nevis 207	M.S. Kalelkar	(COLU)	REFID=21564
CASO	70	LNC 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20590

STRANGE MESONS

($S = \pm 1, C = B = 0$)

$K^+ = u\bar{s}, K^0 = d\bar{s}, \bar{K}^0 = \bar{d}s, K^- = \bar{u}s$, similarly for K^{*} 's

$K_0^*(800)$

or κ

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. See the mini-review on scalar mesons under $f_0(500)$ (see the index for the page number).

$K_0^*(800)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
682 ±29	OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.		
826 ±49	+49 -34	1338	11B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
849 ±77	+18 -14	1421	2,3 ABLIKIM	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
841 ±30	+81 -73	25k	4,5 ABLIKIM	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
658 ±13			6 DESCOTES-G..06	RVUE $\pi K \rightarrow \pi K$
797 ±19 ±43	15k	7,8 AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
663 ± 8 ±34			9 BUGG	10 RVUE S-matrix pole
706.0 ± 1.8 ±22.8	141k	10 BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
856 ±17 ±13	54k	11 LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
750	+30 -55		12 BUGG	06 RVUE
855 ±15	0.6k	13 CAWLFIELD	06A CLEO	$D^0 \rightarrow K^+ K^- \pi^0$
694 ±53		3,14 ZHOU	06 RVUE	$K p \rightarrow K^- \pi^+ n$
753 ±52		15 PELAEZ	04A RVUE	$K \pi \rightarrow K \pi$
594 ±79		14 ZHENG	04 RVUE	$K^- p \rightarrow K^- \pi^+ n$
722 ±60		16 BUGG	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
905	+65 -30	17 ISHIDA	97B RVUE	$11 K^- p \rightarrow K^- \pi^+ n$

¹ The Breit-Wigner parameters from a fit with seven intermediate resonances. The S-matrix pole position is $(764 \pm 63^{+71}_{-54}) - i(306 \pm 149^{+143}_{-85})$ MeV.

² From a fit including ten additional resonances and energy-independent Breit-Wigner width.

³ S-matrix pole.

⁴ S-matrix pole. GUO 06 in a chiral unitary approach report a mass of 757 ± 33 MeV and a width of 558 ± 82 MeV.

⁵ A fit in the $K_0^*(800) + K^*(892) + K^*(1410)$ model with mass and width of the $K_0^*(800)$ from ABLIKIM 06C well describes the left slope of the $K_S^0 \pi^-$ invariant mass spectrum in $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$ decay studied by EPIFANOV 07.

⁶ S-matrix pole. Using Roy-Steiner equations (ROY 71) as well as unitarity, analyticity and crossing symmetry constraints.

⁷ Not seen by KOPP 01 using 7070 events of $D^0 \rightarrow K^- \pi^+ \pi^0$. LINK 02E and LINK 05I show clear evidence for a constant non-resonant scalar amplitude rather than $K_0^*(800)$ in their high statistics analysis of $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$.

⁸ AUBERT 07T does not find evidence for the charged $K_0^*(800)$ using 11k events of $D^0 \rightarrow K^- K^+ \pi^0$.

⁹ S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K \pi$ and $K \eta'$, and the Adler zero near thresholds.

¹⁰ T-matrix pole.

¹¹ A Breit-Wigner mass and width.

¹² S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C using for the κ an s-dependent width with an Adler zero near threshold.

¹³ Breit-Wigner parameters. A significant S-wave can be also modeled as a non-resonant contribution.

NODE=MXXX020

NODE=MXXX020

NODE=M174

NODE=M174

NODE=M174M

NODE=M174M

NODE=M174M;LINKAGE=LI

NODE=M174M;LINKAGE=BL

NODE=M174M;LINKAGE=SM

NODE=M174M;LINKAGE=AB

NODE=M174M;LINKAGE=EP

NODE=M174M;LINKAGE=DE

NODE=M174M;LINKAGE=A

NODE=M174M;LINKAGE=AU

NODE=M174M;LINKAGE=BG

NODE=M174M;LINKAGE=TM


NODE=M174M;LINKAGE=BW

NODE=M174M;LINKAGE=BU

NODE=M174M;LINKAGE=CA

- 14 Using ASTON 88.
- 15 T-matrix pole. Reanalysis of data from LINGLIN 73, ESTABROOKS 78, and ASTON 88 in the unitarized ChPT model.
- 16 T-matrix pole. Reanalysis of ASTON 88 data.
- 17 Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes.

NODE=M174M;LINKAGE=ZH
 NODE=M174M;LINKAGE=PE
 NODE=M174M;LINKAGE=A1
 NODE=M174M;LINKAGE=IS



$K_0^*(800)$ WIDTH

NODE=M174W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
547 ± 24	OUR AVERAGE	Error includes scale factor of 1.1.		
449 ± 156	$^{+144}_{-81}$	1338	18 ABLIKIM	11B BES2 $J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
512 ± 80	$^{+92}_{-44}$	1421	19,20 ABLIKIM	10E BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
618 ± 90	$^{+96}_{-144}$	25k	19,21 ABLIKIM	06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
557 ± 24		22	DESCOTES-G..06	RVUE $\pi K \rightarrow \pi K$
410 ± 43	± 87	15k	23,24 AITALA	02 E791 $D^+ \rightarrow K^- \pi^+ \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
658 ± 10	± 44	25	BUGG	10 RVUE S-matrix pole
638.8 ± 4.4	± 40.4	141k	26 BONVICINI	08A CLEO $D^+ \rightarrow K^- \pi^+ \pi^+$
464 ± 28	± 22	54k	27 LINK	07B FOCS $D^+ \rightarrow K^- \pi^+ \pi^+$
684 ± 120		28	BUGG	06 RVUE
251 ± 48	0.6k	29	CAWLFIELD	06A CLEO $D^0 \rightarrow K^+ K^- \pi^0$
606 ± 59		19,30	ZHOU	06 RVUE $K p \rightarrow K^- \pi^+ n$
470 ± 66		31	PELAEZ	04A RVUE $K \pi \rightarrow K \pi$
724 ± 332		30	ZHENG	04 RVUE $K^- p \rightarrow K^- \pi^+ n$
772 ± 100		32	BUGG	03 RVUE $11 K^- p \rightarrow K^- \pi^+ n$
545	$^{+235}_{-110}$	33	ISHIDA	97B RVUE $11 K^- p \rightarrow K^- \pi^+ n$

NODE=M174W

- 18 The Breit-Wigner parameters from a fit with seven intermediate resonances. The S-matrix pole position is $(764 \pm 63^{+71}_{-54}) - i(306 \pm 149^{+143}_{-85})$ MeV.
- 19 S-matrix pole.
- 20 From a fit including ten additional resonances and energy-independent Breit-Wigner width.
- 21 A fit in the $K_0^*(800) + K^*(892) + K^*(1410)$ model with mass and width of the $K_0^*(800)$ from ABLIKIM 06C well describes the left slope of the $K_S^0 \pi^-$ invariant mass spectrum in $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$ decay studied by EPIFANOV 07.
- 22 S-matrix pole. Using Roy-Steiner equations (ROY 71) as well as unitarity, analyticity and crossing symmetry constraints.
- 23 Not seen by KOPP 01 using 7070 events of $D^0 \rightarrow K^- \pi^+ \pi^0$. LINK 02E and LINK 05I show clear evidence for a constant non-resonant scalar amplitude rather than $K_0^*(800)$ in their high statistics analysis of $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$.

NODE=M174W;LINKAGE=LI

NODE=M174W;LINKAGE=AB
 NODE=M174W;LINKAGE=BL

NODE=M174W;LINKAGE=EP

NODE=M174W;LINKAGE=DE

NODE=M174W;LINKAGE=A

- 24 AUBERT 07T does not find evidence for the charged $K_0^*(800)$ using 11k events of $D^0 \rightarrow K^- K^+ \pi^0$.
 25 S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.
 26 T-matrix pole.
 27 A Breit-Wigner mass and width.
 28 S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C using for the κ an s-dependent width with an Adler zero near threshold.
 29 Statistical error only. A fit to the Dalitz plot including the $K_0^*(800)^\pm$, $K^*(892)^\pm$, and ϕ resonances modeled as Breit-Wigners. A significant S-wave can be also modeled as a non-resonant contribution.
 30 Using ASTON 88.
 31 T-matrix pole. Reanalysis of data from LINGLIN 73, ESTABROOKS 78, and ASTON 88 in the unitarized ChPT model.
 32 T-matrix pole. Reanalysis of ASTON 88 data.
 33 Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes.

NODE=M174W;LINKAGE=AU

NODE=M174W;LINKAGE=BG

NODE=M174W;LINKAGE=TM

NODE=M174W;LINKAGE=BW

NODE=M174W;LINKAGE=BU

NODE=M174W;LINKAGE=CA

NODE=M174W;LINKAGE=ZH

NODE=M174W;LINKAGE=PE

NODE=M174W;LINKAGE=A1

NODE=M174W;LINKAGE=IS

 $K_0^*(800)$ REFERENCES

NODE=M174

ABLIKIM	11B	PL B698 183	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53683
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
BUGG	10	PR D81 014002	D.V. Bugg	(LOQM)	REFID=53213
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=53056
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=52426
AUBERT	07T	PR D76 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51726
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)	REFID=51929
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51875
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51051
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51458
BUGG	06	PL B632 471	D.V. Bugg	(LOQM)	REFID=50996
CAWLFIELD	06A	PR D74 031108	C. Cawfield <i>et al.</i>	(CLEO Collab.)	REFID=51153
DESCOTES-G...	06	EPJ C48 553	S. Descotes-Genon, B. Moussallam		REFID=51518
GUO	06	NP A773 78	F.K. Guo <i>et al.</i>		REFID=51164
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng		REFID=51198
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=50679
PELAEZ	04A	MPL A19 2879	J.R. Pelaez		REFID=50347
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>		REFID=50165
BUGG	03	PL B572 1	D.V. Bugg		REFID=49586
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48807
LINK	02E	PL B535 43	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=48728
KOPP	01	PR D63 092001	S. Kopp <i>et al.</i>	(CLEO Collab.)	REFID=48134
ISHIDA	97B	PTP 98 621	S. Ishida <i>et al.</i>		REFID=48655
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
ROY	71	PL 36B 353	S.M. Roy		REFID=51107

$K^*(892)$

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M018

 $K^*(892)$ MASS

NODE=M018205

CHARGED ONLY, HADROPRODUCED

NODE=M018M1

NODE=M018M1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
891.66±0.26 OUR AVERAGE						
892.6 ±0.5	5840	BAUBILLIER 84B	HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
888 ±3		NAPIER 84	SPEC	+	200 $\pi^- p \rightarrow 2K_S^0 X$	
891 ±1		NAPIER 84	SPEC	-	200 $\pi^- p \rightarrow 2K_S^0 X$	
891.7 ±2.1	3700	BARTH 83	HBC	+	70 $K^+ p \rightarrow K^0 \pi^+ X$	
891 ±1	4100	TOAFF 81	HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
892.8 ±1.6		AJINENKO 80	HBC	+	32 $K^+ p \rightarrow K^0 \pi^+ X$	
890.7 ±0.9	1800	AGUILAR-...	78B	HBC	±	0.76 $\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$
886.6 ±2.4	1225	BALAND 78	HBC	±	12 $\bar{p} p \rightarrow (K\pi)^\pm X$	
891.7 ±0.6	6706	COOPER 78	HBC	±	0.76 $\bar{p} p \rightarrow (K\pi)^\pm X$	
891.9 ±0.7	9000	¹ PALER 75	HBC	-	14.3 $K^- p \rightarrow (K\pi)^- X$	
892.2 ±1.5	4404	AGUILAR-...	71B	HBC	-	3.9,4.6 $K^- p \rightarrow (K\pi)^- p$
891 ±2	1000	CRENNELL 69D	DBC	-	3.9 $K^- N \rightarrow K^0 \pi^- X$	
890 ±3.0	720	BARLOW 67	HBC	±	1.2 $\bar{p} p \rightarrow (K^0 \pi)^\pm K^\mp$	
889 ±3.0	600	BARLOW 67	HBC	±	1.2 $\bar{p} p \rightarrow (K^0 \pi)^\pm K\pi$	
891 ±2.3	620	² DEBAERE 67B	HBC	+	3.5 $K^+ p \rightarrow K^0 \pi^+ p$	
891.0 ±1.2	1700	³ WOJCICKI 64	HBC	-	1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
893.5 ±1.1	27k	⁴ ABELE 99D	CBAR	±	0.0 $\bar{p} p \rightarrow K^+ K^- \pi^0$	
890.4 ±0.2 ±0.5	80±0.8k	⁵ BIRD 89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
890.0 ±2.3	800	^{2,3} CLELAND 82	SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$	
896.0 ±1.1	3200	^{2,3} CLELAND 82	SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$	
893 ±1	3600	^{2,3} CLELAND 82	SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$	
896.0 ±1.9	380	DELFOSSÉ 81	SPEC	+	50 $K^\pm p \rightarrow K^\pm \pi^0 p$	
886.0 ±2.3	187	DELFOSSÉ 81	SPEC	-	50 $K^\pm p \rightarrow K^\pm \pi^0 p$	
894.2 ±2.0	765	² CLARK 73	HBC	-	3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
894.3 ±1.5	1150	^{2,3} CLARK 73	HBC	-	3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
892.0 ±2.6	341	² SCHWEING...68	HBC	-	5.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$	

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=2

CHARGED ONLY, PRODUCED IN τ LEPTON DECAYS

NODE=M018MCT

NODE=M018MCT

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
895.47±0.20±0.74				
	53k	⁶ EPIFANOV 07	BELL	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
892.0 ±0.5		⁷ BOITO 10	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
892.0 ±0.9		^{8,9} BOITO 09	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
895.3 ±0.2		^{8,10} JAMIN 08	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
896.4 ±0.9	11970	¹¹ BONVICINI 02	CLEO	$\tau^- \rightarrow K^- \pi^0 \nu_\tau$
895 ±2		¹² BARATE 99R	ALEP	$\tau^- \rightarrow K^- \pi^0 \nu_\tau$

NEUTRAL ONLY


NODE=M018M2

NODE=M018M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
895.81±0.19 OUR AVERAGE				
Error includes scale factor of 1.4. See the ideogram below.				
895.4 ±0.2 ±0.2	243k	¹³ DEL-AMO-SA..11i	BABR	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
895.7 ±0.2 ±0.3	141k	¹⁴ BONVICINI 08A	CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
895.41±0.32 ^{+0.35} _{-0.43}	18k	¹⁵ LINK 05i	FOCS	$D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
896 ±2		BARBERIS 98E	OMEG	450 $pp \rightarrow p_f p_s K^* \bar{K}^*$
895.9 ±0.5 ±0.2		ASTON 88	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
894.52±0.63	25k	¹ ATKINSON 86	OMEG	20-70 γp
894.63±0.76	20k	¹ ATKINSON 86	OMEG	20-70 γp
897 ±1	28k	EVANGELIS... 80	OMEG	10 $\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$
898.4 ±1.4	1180	AGUILAR-... 78B	HBC	0.76 $\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$
894.9 ±1.6		WICKLUND 78	ASPK	3,4,6 $K^\pm N \rightarrow (K\pi)^0 N$
897.6 ±0.9		BOWLER 77	DBC	5.4 $K^+ d \rightarrow K^+ \pi^- pp$
895.5 ±1.0	3600	MCCUBBIN 75	HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$
897.1 ±0.7	22k	¹ PALER 75	HBC	14.3 $K^- p \rightarrow (K\pi)^0 X$
896.0 ±0.6	10k	FOX 74	RVUE	2 $K^- p \rightarrow K^- \pi^+ n$

OCCUR=2

896.0 ±0.6		FOX	74	RVUE	2	$K^+ n \rightarrow K^+ \pi^- p$	OCCUR=2
896 ±2		¹⁶ MATISON	74	HBC	12	$K^+ p \rightarrow K^+ \pi^- \Delta$	
896 ±1	3186	LEWIS	73	HBC	2.1-2.7	$K^+ p \rightarrow K \pi \pi p$	
894.0 ±1.3		¹⁶ LINGLIN	73	HBC	2-13	$K^+ p \rightarrow K^+ \pi^- \pi^+ p$	
898.4 ±1.3	1700	² BUCHNER	72	DBC	4.6	$K^+ n \rightarrow K^+ \pi^- p$	
897.9 ±1.1	2934	² AGUILAR-...	71B	HBC	3.9,4.6	$K^- p \rightarrow K^- \pi^+ n$	
898.0 ±0.7	5362	² AGUILAR-...	71B	HBC	3.9,4.6	$K^- p \rightarrow K^- \pi^+ \pi^- p$	OCCUR=2
895 ±1	4300	³ HABER	70	DBC	3	$K^- N \rightarrow K^- \pi^+ X$	
893.7 ±2.0	10k	DAVIS	69	HBC	12	$K^+ p \rightarrow K^+ \pi^- \pi^+ p$	
894.7 ±1.4	1040	² DAUBER	67B	HBC	2.0	$K^- p \rightarrow K^- \pi^+ \pi^- p$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●							
895.53±0.17		LEES	13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$		
894.9 ±0.5 ±0.7	14.4k	¹⁷ MITCHELL	09A	CLEO	$D_s^+ \rightarrow K^+ K^- \pi^+$		
896.2 ±0.3	20k	⁸ AUBERT	07AK	BABR	$10.6 e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$		
900.7 ±1.1	5900	BARTH	83	HBC	70	$K^+ p \rightarrow K^+ \pi^- X$	



- 1 Inclusive reaction. Complicated background and phase-space effects.
- 2 Mass errors enlarged by us to Γ/\sqrt{N} . See note.
- 3 Number of events in peak reevaluated by us.
- 4 K-matrix pole.
- 5 From a partial wave amplitude analysis.
- 6 From a fit in the $K_0^*(800) + K^*(892) + K^*(1410)$ model.
- 7 From the pole position of the $K \pi$ vector form factor using EPIFANOV 07 and constraints from K_{J3} decays in ANTONELLI 10.
- 8 Systematic uncertainties not estimated.
- 9 From the pole position of the $K \pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.
- 10 Reanalysis of EPIFANOV 07 using resonance chiral theory.
- 11 Calculated by us from the shift by 4.7 ± 0.9 MeV (statistical uncertainty only) reported in BONVICINI 02 with respect to the world average value from PDG 00.
- 12 With mass and width of the $K^*(1410)$ fixed at 1412 MeV and 227 MeV, respectively.
- 13 Taking into account the $K^*(892)^0$, S -wave and P -wave ($K^*(1410)^0$).
- 14 From the isobar model with a complex pole for the κ .
- 15 Fit to $K \pi$ mass spectrum includes a non-resonant scalar component.

NODE=M018M;LINKAGE=I
 NODE=M018M;LINKAGE=D
 NODE=M018M;LINKAGE=W
 NODE=M018M1;LINKAGE=AN
 NODE=M018M1;LINKAGE=F
 NODE=M018MCT;LINKAGE=EF
 NODE=M018MCT;LINKAGE=BT
 NODE=M018M2;LINKAGE=NS
 NODE=M018MCT;LINKAGE=BI
 NODE=M018MCT;LINKAGE=JA
 NODE=M018MCT;LINKAGE=BO
 NODE=M018MCT;LINKAGE=BA
 NODE=M018M2;LINKAGE=DE
 NODE=M018M2;LINKAGE=BO
 NODE=M018W2;LINKAGE=LI

¹⁶ From pole extrapolation.¹⁷ This value comes from a fit with χ^2 of 178/117.NODE=M018M;LINKAGE=C
NODE=M018M2;LINKAGE=MI

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=M018209

 $m_{K^*(892)^0} - m_{K^*(892)^\pm}$

NODE=M018D

NODE=M018D

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
6.7±1.2 OUR AVERAGE					
7.7±1.7	2980	AGUILAR-...	78B	HBC	±0 0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
5.7±1.7	7338	AGUILAR-...	71B	HBC	-0 3.9,4.6 $K^- p$
6.3±4.1	283	¹⁸ BARASH	67B	HBC	0.0 $\bar{p}p$

¹⁸ Number of events in peak reevaluated by us.

NODE=M018D;LINKAGE=W

 $K^*(892)$ RANGE PARAMETER

NODE=M018R

NODE=M018R

All from partial wave amplitude analyses.

NODE=M018R

VALUE (GeV ⁻¹)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.1 ±0.5 ±0.5	243k	¹⁹ DEL-AMO-SA.11i	BABR	0	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
3.96±0.54 ^{+1.31} _{-0.90}	18k	²⁰ LINK	05i	FOCS	0 $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
3.4 ±0.7		ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
12.1 ±3.2 ±3.0		BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

¹⁹ Taking into account the $K^*(892)^0$, S -wave and P -wave ($K^*(1410)^0$).²⁰ Fit to $K \pi$ mass spectrum includes a non-resonant scalar component.NODE=M018R;LINKAGE=DE
NODE=M018R;LINKAGE=LI $K^*(892)$ WIDTH

NODE=M018215

CHARGED ONLY, HADROPRODUCED

NODE=M018W1
NODE=M018W1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
50.8±0.9 OUR FIT					
50.8±0.9 OUR AVERAGE					
49 ±2	5840	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
56 ±4		NAPIER	84	SPEC	- 200 $\pi^- p \rightarrow 2K_S^0 X$
51 ±2	4100	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
50.5±5.6		AJINENKO	80	HBC	+ 32 $K^+ p \rightarrow K^0 \pi^+ X$
45.8±3.6	1800	AGUILAR-...	78B	HBC	± 0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
52.0±2.5	6706	²¹ COOPER	78	HBC	± 0.76 $\bar{p}p \rightarrow (K\pi)^\pm X$
52.1±2.2	9000	²² PALER	75	HBC	- 14.3 $K^- p \rightarrow (K\pi)^- X$
46.3±6.7	765	²¹ CLARK	73	HBC	- 3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$
48.2±5.7	1150	^{21,23} CLARK	73	HBC	- 3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$
54.3±3.3	4404	²¹ AGUILAR-...	71B	HBC	- 3.9,4.6 $K^- p \rightarrow (K\pi)^- p$
46 ±5	1700	^{21,23} WOJCICKI	64	HBC	- 1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
54.8±1.7	27k	²⁴ ABELE	99D	CBAR	± 0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$
45.2±1 ±2	79.7±0.8k	²⁵ BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
42.8±7.1	3700	BARTH	83	HBC	+ 70 $K^+ p \rightarrow K^0 \pi^+ X$
64.0±9.2	800	^{21,23} CLELAND	82	SPEC	+ 30 $K^+ p \rightarrow K_S^0 \pi^+ p$
62.0±4.4	3200	^{21,23} CLELAND	82	SPEC	+ 50 $K^+ p \rightarrow K_S^0 \pi^+ p$
55 ±4	3600	^{21,23} CLELAND	82	SPEC	- 50 $K^+ p \rightarrow K_S^0 \pi^- p$
62.6±3.8	380	DELFOSSSE	81	SPEC	+ 50 $K^\pm p \rightarrow K^\pm \pi^0 p$
50.5±3.9	187	DELFOSSSE	81	SPEC	- 50 $K^\pm p \rightarrow K^\pm \pi^0 p$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

CHARGED ONLY, PRODUCED IN τ LEPTON DECAYSNODE=M018W5
NODE=M018W5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
46.2±0.6±1.2				
	53k	²⁶ EPIFANOV	07	BELL $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
46.5±1.1		²⁷ BOITO	10	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
46.2±0.4		^{28,29} BOITO	09	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
47.5±0.4		^{28,30} JAMIN	08	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
55 ±8		³¹ BARATE	99R	ALEP $\tau^- \rightarrow K^- \pi^0 \nu_\tau$

NEUTRAL ONLYNODE=M018W2
NODE=M018W2


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
47.4 ±0.6	OUR FIT	Error includes scale factor of 2.2.		
47.4 ±0.6	OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.		
46.5 ±0.3 ±0.2	243k	32 DEL-AMO-SA..11I	BABR	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
45.3 ±0.5 ±0.6	141k	33 BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
47.79 ±0.86 ^{+1.32} _{-1.06}	18k	34 LINK	05I FOCS	$D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
54 ±3		BARBERIS	98E OMEG	450 $pp \rightarrow p_f p_s K^* K^*$
50.8 ±0.8 ±0.9		ASTON	88 LASS	11 $K^- p \rightarrow K^- \pi^+ n$
46.5 ±4.3	5900	BARTH	83 HBC	70 $K^+ p \rightarrow K^+ \pi^- X$
54 ±2	28k	EVANGELIS...	80 OMEG	10 $\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$
45.9 ±4.8	1180	AGUILAR-...	78B HBC	0.76 $\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$
51.2 ±1.7		WICKLUND	78 ASPK	3,4,6 $K^\pm N \rightarrow (K\pi)^0 N$
48.9 ±2.5		BOWLER	77 DBC	5.4 $K^+ d \rightarrow K^+ \pi^- pp$
48 ⁺³ ₋₂	3600	MCCUBBIN	75 HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$
50.6 ±2.5	22k	22 PALER	75 HBC	14.3 $K^- p \rightarrow (K\pi)^0 X$
47 ±2	10k	FOX	74 RVUE	2 $K^- p \rightarrow K^- \pi^+ n$
51 ±2		FOX	74 RVUE	2 $K^+ n \rightarrow K^+ \pi^- p$
46.0 ±3.3	3186	21 LEWIS	73 HBC	2.1-2.7 $K^+ p \rightarrow K\pi\pi p$
51.4 ±5.0	1700	21 BUCHNER	72 DBC	4.6 $K^+ n \rightarrow K^+ \pi^- p$
55.8 ^{+4.2} _{-3.4}	2934	21 AGUILAR-...	71B HBC	3.9,4.6 $K^- p \rightarrow K^- \pi^+ n$
48.5 ±2.7	5362	AGUILAR-...	71B HBC	3.9,4.6 $K^- p \rightarrow$ $K^- \pi^+ \pi^- p$
54.0 ±3.3	4300	21,23 HABER	70 DBC	3 $K^- N \rightarrow K^- \pi^+ X$
53.2 ±2.1	10k	21 DAVIS	69 HBC	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$
44 ±5.5	1040	21 DAUBER	67B HBC	2.0 $K^- p \rightarrow K^- \pi^+ \pi^- p$

OCCUR=2

OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

44.90 ±0.30		LEES	13F BABR	$D^+ \rightarrow K^+ K^- \pi^+$
45.7 ±1.1 ±0.5	14.4k	35 MITCHELL	09A CLEO	$D_s^+ \rightarrow K^+ K^- \pi^+$
50.6 ±0.9	20k	28 AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow$ $K^{*0} K^\pm \pi^\mp \gamma$



NEUTRAL ONLY (MeV)

- 21 Width errors enlarged by us to $4 \times \Gamma / \sqrt{N}$; see note.
- 22 Inclusive reaction. Complicated background and phase-space effects.
- 23 Number of events in peak reevaluated by us.
- 24 K-matrix pole.
- 25 From a partial wave amplitude analysis.
- 26 From a fit in the $K_0^*(800) + K^*(892) + K^*(1410)$ model.
- 27 From the pole position of the $K\pi$ vector form factor using EPIFANOV 07 and constraints from K_{J3} decays in ANTONELLI 10.
- 28 Systematic uncertainties not estimated.
- 29 From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.
- 30 Reanalysis of EPIFANOV 07 using resonance chiral theory.
- 31 With mass and width of the $K^*(1410)$ fixed at 1412 MeV and 227 MeV, respectively.
- 32 Taking into account the $K^*(892)^0$, S -wave and P -wave ($K^*(1410)^0$).
- 33 From the isobar model with a complex pole for the κ .
- 34 Fit to $K\pi$ mass spectrum includes a non-resonant scalar component.
- 35 This value comes from a fit with χ^2 of 178/117.

NODE=M018W;LINKAGE=D
 NODE=M018W;LINKAGE=I
 NODE=M018W;LINKAGE=W
 NODE=M018W1;LINKAGE=AN
 NODE=M018W1;LINKAGE=F
 NODE=M018W5;LINKAGE=EF
 NODE=M018W5;LINKAGE=BT

 NODE=M018W2;LINKAGE=NS
 NODE=M018W5;LINKAGE=BI

 NODE=M018W2;LINKAGE=JA
 NODE=M018W5;LINKAGE=BA
 NODE=M018W2;LINKAGE=DE
 NODE=M018W2;LINKAGE=BO
 NODE=M018M2;LINKAGE=LI
 NODE=M018W2;LINKAGE=MI

$K^*(892)$ DECAY MODES

NODE=M018220;NODE=M018

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K\pi$	~ 100	%
Γ_2 $(K\pi)^\pm$	$(99.901 \pm 0.009) \%$	
Γ_3 $(K\pi)^0$	$(99.754 \pm 0.021) \%$	
Γ_4 $K^0\gamma$	$(2.46 \pm 0.21) \times 10^{-3}$	
Γ_5 $K^\pm\gamma$	$(9.9 \pm 0.9) \times 10^{-4}$	
Γ_6 $K\pi\pi$	< 7	$\times 10^{-4}$ 95%

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=11
 DESIG=12
 DESIG=4
 DESIG=3
 DESIG=2

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 7.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc}
 x_5 & -100 & \\
 \Gamma & 19 & -19 \\
 & x_2 & x_5
 \end{array}$$

	Mode	Rate (MeV)	
Γ_2	$(K\pi)^\pm$	50.7 \pm 0.9	DESIG=11
Γ_5	$K^\pm \gamma$	0.050 \pm 0.005	DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 22 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 66.8$ for 20 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc}
 x_4 & -100 & \\
 \Gamma & 15 & -15 \\
 & x_3 & x_4
 \end{array}$$

	Mode	Rate (MeV)	Scale factor	
Γ_3	$(K\pi)^0$	47.3 \pm 0.6	2.1	DESIG=12
Γ_4	$K^0 \gamma$	0.116 \pm 0.010		DESIG=4

K*(892) PARTIAL WIDTHS

$\Gamma(K^0 \gamma)$							Γ_4
VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
116 \pm 10 OUR FIT							
116.5 \pm 9.9	584	CARLSMITH	86	SPEC	0	$K_L^0 A \rightarrow K_S^0 \pi^0 A$	NODE=M018225

$\Gamma(K^\pm \gamma)$							Γ_5
VALUE (keV)		DOCUMENT ID	TECN	CHG	COMMENT		
50 \pm 5 OUR FIT							
50 \pm 5 OUR AVERAGE							
48 \pm 11		BERG	83	SPEC	-	156 $K^- A \rightarrow \bar{K} \pi A$	NODE=M018W3
51 \pm 5		CHANDLEE	83	SPEC	+	200 $K^+ A \rightarrow K \pi A$	NODE=M018W3

K*(892) BRANCHING RATIOS

$\Gamma(K^0 \gamma) / \Gamma_{\text{total}}$							Γ_4 / Γ
VALUE (units 10^{-3})		DOCUMENT ID	TECN	CHG	COMMENT		
2.46 \pm 0.21 OUR FIT							
1.5 \pm 0.7		CARITHERS	75B	CNTR	0	8-16 $\bar{K}^0 A$	NODE=M018230

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K^\pm\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
--------------------------	-----	-------------	------	-----	---------

0.99±0.09 OUR FIT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<1.6	95	BEMPORAD	73	CNTR +	10-16 K^+A
------	----	----------	----	--------	--------------

 Γ_5/Γ NODE=M018R2
NODE=M018R2 $\Gamma(K\pi\pi)/\Gamma((K\pi)^\pm)$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

< 7×10^{-4}	95	JONGEJANS	78	HBC	$4 K^- p \rightarrow p \bar{K}^0 2\pi$
----------------------	----	-----------	----	-----	--

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 20×10^{-4}		WOJCICKI	64	HBC -	$1.7 K^- p \rightarrow \bar{K}^0 \pi^- p$
-----------------------	--	----------	----	-------	---

 Γ_6/Γ_2 NODE=M018R1
NODE=M018R1**K*(892) REFERENCES**

NODE=M018

LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55127
DEL-AMO-SA...	11I	PR D83 072001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16493
ANTONELLI	10	EPJ C69 399	M. Antonelli <i>et al.</i>	(FlaviaNet Working Group)	REFID=53448
BOITO	10	JHEP 1009 031	D.R. Boito, R. Escribano, M. Jamin	(BARC)	REFID=53632
BOITO	09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin		REFID=52728
MITCHELL	09A	PR D79 072008	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52756
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=52426
JAMIN	08	PL B664 78	M. Jamin, A. Pich, J. Portoles		REFID=52285
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)	REFID=51929
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=50679
BONVICINI	02	PRL 88 111803	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=48701
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47366
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46348
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20564
CARLSMITH	86	PRL 56 18	D. Carlsmith <i>et al.</i>	(EFI, SACL)	REFID=22461
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
NAPIER	84	PL 149B 514	A. Napier <i>et al.</i>	(TUFTS, ARIZ, FNAL, FLOR+)	REFID=22460
BARTH	83	NP B223 296	M. Barth <i>et al.</i>	(BRUX, CERN, GENO, MONS+)	REFID=22456
BERG	83	Thesis UMI 83-21652	D.M. Berg	(ROCH)	REFID=22457
CHANDLEE	83	PRL 51 168	C. Handlee <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=22458
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
DELFOSSSE	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
AJINENKO	80	ZPHY C5 177	I.V. Ajinenko <i>et al.</i>	(SERP, BRUX, MONS+)	REFID=22449
EVANGELIS...	80	NP B165 383	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=22450
AGUILAR...	78B	NP B141 101	M. Aguilar-Benitez <i>et al.</i>	(MADR, TATA+)	REFID=22438
BALAND	78	NP B140 220	J.F. Baland <i>et al.</i>	(MONS, BELG, CERN+)	REFID=20369
COOPER	78	NP B136 365	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=22441
JONGEJANS	78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)	REFID=22445
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
BOWLER	77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)	REFID=22437
CARITHERS	75B	PRL 35 349	W.C.J. Carithers <i>et al.</i>	(ROCH, MCGI)	REFID=22433
MCCUBBIN	75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)	REFID=22434
PALER	75	NP B96 1	K. Paler <i>et al.</i>	(RHEL, SACL, EPOL)	REFID=22435
FOX	74	NP B80 403	G.C. Fox, M.L. Griss	(CIT)	REFID=22430
MATISON	74	PR D9 1872	M.J. Matison <i>et al.</i>	(LBL)	REFID=22431
BEMPORAD	73	NP B51 1	C. Bemporad <i>et al.</i>	(CERN, ETH, LOIC)	REFID=22416
CLARK	73	NP B54 432	A.G. Clark, L. Lyons, D. Radojicic	(OXF)	REFID=22426
LEWIS	73	NP B60 283	P.H. Lewis <i>et al.</i>	(LOWC, LOIC, CDEF)	REFID=22427
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
BUCHNER	72	NP B45 333	K. Buchner <i>et al.</i>	(MPIM, CERN, BRUX)	REFID=22418
AGUILAR...	71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)	REFID=22408
HABER	70	NP B17 289	B. Haber <i>et al.</i>	(REHO, SACL, BGNA, EPOL)	REFID=22406
CRENNELL	69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)	REFID=22399
DAVIS	69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)	REFID=22400
SCHWEING...	68	PR 166 1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)	REFID=22398
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)	REFID=20160
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
DAUBER	67B	PR 153 1403	P.M. Dauber <i>et al.</i>	(UCLA)	REFID=22389
DEBAERE	67B	NC 51A 401	W. de Baere <i>et al.</i>	(BRUX, CERN)	REFID=22390
WOJCICKI	64	PR 135 B484	S.G. Wojcicki	(LRL)	REFID=22379

$K_1(1270)$

$$I(J^P) = \frac{1}{2}(1^+)$$

NODE=M028

 $K_1(1270)$ MASS

NODE=M028205

VALUE (MeV) DOCUMENT ID**1272±7 OUR AVERAGE** Includes data from the 2 datablocks that follow this one.

NODE=M028MX

PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGENODE=M028M2
NODE=M028M2VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

1275±10 700 GAVILLET 78 HBC + 4.2 $K^- p \rightarrow \Xi^- (K \pi \pi)^+$ **PRODUCED BY K BEAMS**NODE=M028M3
NODE=M028M3VALUE (MeV) DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

1270±10 ¹ DAUM 81C CNTR - 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1276 ² TORNQVIST 82B RVUE~ 1300 VERGEEST 79 HBC - 4.2 $K^- p \rightarrow (\bar{K} \pi \pi)^- p$ 1289±25 ³ CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$ ~ 1300 BRANDENB... 76 ASPK ± 13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$ ~ 1270 OTTER 76 HBC - 10,14,16 $K^- p \rightarrow (\bar{K} \pi \pi)^- p$ 1260 DAVIS 72 HBC + 12 $K^+ p$ 1234±12 FIRESTONE 72B DBC + 12 $K^+ d$ ¹ Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.² From a unitarized quark-model calculation.³ From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

NODE=M028M3;LINKAGE=DA

NODE=M028M3;LINKAGE=T
NODE=M028M3;LINKAGE=E**PRODUCED BY BEAMS OTHER THAN K MESONS**NODE=M028M1
NODE=M028M1VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT**1248.1± 3.3±1.4** GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1279 ±10 25k ⁴ ABLIKIM 06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$ 1294 ±10 310 RODEBACK 81 HBC 4 $\pi^- p \rightarrow \Lambda K 2\pi$ 1300 40 CRENNELL 72 HBC 0 4.5 $\pi^- p \rightarrow \Lambda K 2\pi$ 1242 $\begin{smallmatrix} +9 \\ -10 \end{smallmatrix}$ ⁵ ASTIER 69 HBC 0 $\bar{p} p$ 1300 45 CRENNELL 67 HBC 0 6 $\pi^- p \rightarrow \Lambda K 2\pi$ ⁴ Systematic errors not estimated.⁵ This was called the C meson.NODE=M028M1;LINKAGE=AB
NODE=M028M1;LINKAGE=A**PRODUCED IN τ LEPTON DECAYS**NODE=M028MT
NODE=M028MTVALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT**1254±33±34** 7k ASNER 00B CLEO ± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ **$K_1(1270)$ WIDTH**

NODE=M028210

VALUE (MeV) DOCUMENT ID**90±20 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values.NODE=M028WX
→ UNCHECKED ←**87± 7 OUR AVERAGE** Includes data from the 2 datablocks that follow this one.**PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE**NODE=M028W2
NODE=M028W2VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

75±15 700 GAVILLET 78 HBC + 4.2 $K^- p \rightarrow \Xi^- K \pi \pi$

PRODUCED BY K BEAMS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

90 ± 8	⁶ DAUM	81C	CNTR	-	$63 K^- p \rightarrow K^- 2\pi p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
~ 150	VERGEEST	79	HBC	-	$4.2 K^- p \rightarrow (\bar{K} \pi \pi)^- p$
150 ± 71	⁷ CARNEGIE	77	ASPK	±	$13 K^\pm p \rightarrow (K \pi \pi)^\pm p$
~ 200	BRANDENB...	76	ASPK	±	$13 K^\pm p \rightarrow (K \pi \pi)^\pm p$
120	DAVIS	72	HBC	+	$12 K^+ p$
188 ± 21	FIRESTONE	72B	DBC	+	$12 K^+ d$

⁶Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

⁷From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

NODE=M028W3
NODE=M028W3

NODE=M028W3;LINKAGE=DA

NODE=M028W3;LINKAGE=E

PRODUCED BY BEAMS OTHER THAN K MESONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

$119.5 \pm 5.2 \pm 6.7$		GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
131 ± 21	25k	⁸ ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
66 ± 15	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
60	40	CRENNELL	72	HBC	0 $4.5 \pi^- p \rightarrow \Lambda K 2\pi$
$127 \begin{smallmatrix} +7 \\ -25 \end{smallmatrix}$		ASTIER	69	HBC	0 $\bar{p} p$
60	45	CRENNELL	67	HBC	0 $6 \pi^- p \rightarrow \Lambda K 2\pi$

⁸Systematic errors not estimated.

NODE=M028W1
NODE=M028W1

NODE=M028W1;LINKAGE=AB

PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

$260 \begin{smallmatrix} +90 \\ -70 \end{smallmatrix} \pm 80$	7k	ASNER	00B	CLEO	± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
---	----	-------	-----	------	---

NODE=M028WT
NODE=M028WT

 $K_1(1270)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \rho$	(42 ± 6) %
Γ_2 $K_0^*(1430) \pi$	(28 ± 4) %
Γ_3 $K^*(892) \pi$	(16 ± 5) %
Γ_4 $K \omega$	(11.0 ± 2.0) %
Γ_5 $K f_0(1370)$	(3.0 ± 2.0) %
Γ_6 γK^0	seen

NODE=M028215;NODE=M028

DESIG=2

DESIG=7

DESIG=1

DESIG=5

DESIG=8

DESIG=9;OUR EST;→ UNCHECKED ←

 $K_1(1270)$ PARTIAL WIDTHS **$\Gamma(K \rho)$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
57 ± 5	MAZZUCATO	79	HBC	+	$4.2 K^- p \rightarrow \Xi^- (K \pi \pi)^+$
75 ± 6	CARNEGIE	77B	ASPK	±	$13 K^\pm p \rightarrow (K \pi \pi)^\pm p$

Γ_1

NODE=M028W5
NODE=M028W5

 $\Gamma(K_0^*(1430) \pi)$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
26 ± 6	CARNEGIE	77B	ASPK	±	$13 K^\pm p \rightarrow (K \pi \pi)^\pm p$

Γ_2

NODE=M028W7
NODE=M028W7

 $\Gamma(K^*(892) \pi)$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
14 ± 11	MAZZUCATO	79	HBC	+	$4.2 K^- p \rightarrow \Xi^- (K \pi \pi)^+$
2 ± 2	CARNEGIE	77B	ASPK	±	$13 K^\pm p \rightarrow (K \pi \pi)^\pm p$

Γ_3

NODE=M028W4
NODE=M028W4

$\Gamma(K\omega)$ Γ_4

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4±4	MAZZUCATO 79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
24±3	CARNEGIE 77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W6
NODE=M028W6

 $\Gamma(K f_0(1370))$ Γ_5

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

22±5	CARNEGIE 77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
------	--------------	------	---	--

NODE=M028W8
NODE=M028W8

 $\Gamma(\gamma K^0)$ Γ_6

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

73.2±6.1±28.3	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$
----------------------	-----------------	------	-----------------------------

NODE=M028W9
NODE=M028W9

 $K_1(1270)$ BRANCHING RATIOS

NODE=M028225

 $\Gamma(K\rho)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.42 ±0.06	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.584±0.043	¹⁰ GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
dominant	RODEBACK	81	HBC $4 \pi^- p \rightarrow \Lambda K 2\pi$

NODE=M028R2
NODE=M028R2

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.28 ±0.04	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0201±0.0064	¹⁰ GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
---------------	---------------------	----	---

NODE=M028R4
NODE=M028R4

 $\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.16 ±0.05	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.171±0.023	¹⁰ GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
-------------	---------------------	----	---

NODE=M028R1
NODE=M028R1

 $\Gamma(K\omega)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.11 ±0.02	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-------------------	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.225±0.052	¹⁰ GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
-------------	---------------------	----	---

NODE=M028R3
NODE=M028R3

 $\Gamma(K\omega)/\Gamma(K\rho)$ Γ_4/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.30	95	RODEBACK	81	HBC $4 \pi^- p \rightarrow \Lambda K 2\pi$
-------	----	----------	----	--

NODE=M028R6
NODE=M028R6

 $\Gamma(K f_0(1370))/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.03±0.02	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
------------------	-------------------	-----	--

NODE=M028R5
NODE=M028R5

 D -wave/ S -wave RATIO FOR $K_1(1270) \rightarrow K^*(892)\pi$

NODE=M028R9
NODE=M028R9

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.0±0.7	⁹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
----------------	-------------------	-----	--

⁹ Average from low and high t data.

¹⁰ Assuming that decays are saturated by the $K\rho$, $K_0^*(1430)\pi$, $K^*(892)\pi$, $K\omega$ decay modes and neglecting interference between them. The values $B(\omega \rightarrow \pi^+\pi^-) = (1.53^{+0.11}_{-0.13})\%$ and $B(K_0^*(1430) \rightarrow K\pi) = (93 \pm 10)\%$ are used. Systematic uncertainties not estimated.

NODE=M028R;LINKAGE=F
NODE=M028R1;LINKAGE=GU

$K_1(1270)$ REFERENCES

GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
GENG	07	PR D75 014017	L.S. Geng <i>et al.</i>	
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELSE)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
RODEBACK	81	ZPHY C9 9	S. Rodeback <i>et al.</i>	(CERN, CDEF, MADR+)
MAZZUCATO	79	NP B156 532	M. Mazzucato <i>et al.</i>	(CERN, ZEEM, NIJM+)
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
GAVILLET	78	PL 76B 517	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+) JP
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)
CARNEGIE	77B	PL 68B 287	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
OTTER	76	NP B106 77	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP
CRENNELL	72	PR D6 1220	D.J. Crennell <i>et al.</i>	(BNL)
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)
ASTIER	69	NP B10 65	A. Astier <i>et al.</i>	(CDEF, CERN, IPNP, LIVP) IJP
CRENNELL	67	PRL 19 44	D.J. Crennell <i>et al.</i>	(BNL) I

NODE=M028

REFID=53668
 REFID=51623
 REFID=51037
 REFID=48822
 REFID=47766
 REFID=20573
 REFID=22548
 REFID=22550
 REFID=20867
 REFID=22542
 REFID=22538
 REFID=22535
 REFID=22536
 REFID=22532
 REFID=22533
 REFID=22419
 REFID=22505
 REFID=22506
 REFID=22482
 REFID=22473

 $K_1(1400)$

$$I(J^P) = \frac{1}{2}(1^+)$$

NODE=M064

 $K_1(1400)$ MASS

NODE=M064M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1403 ± 7 OUR AVERAGE					
1463 ± 64 ± 68	7k	ASNER	00B	CLEO	± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
1373 ± 14 ± 18		¹ ASTON	87	LASS	0 $11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1392 ± 18		BAUBILLIER	82B	HBC	0 $8.25 K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
1410 ± 25		DAUM	81C	CNTR	- $63 K^- p \rightarrow K^- 2\pi p$
1415 ± 15		ETKIN	80	MPS	0 $6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1404 ± 10		² CARNEGIE	77	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1418 ± 8	25k	³ ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 1350		⁴ TORNQVIST	82B	RVUE	
~ 1400		VERGEEST	79	HBC	- $4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 1400		BRANDENB...	76	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
1420		DAVIS	72	HBC	+ $12 K^+ p$
1368 ± 18		FIRESTONE	72B	DBC	+ $12 K^+ d$

NODE=M064M

¹ From partial-wave analysis of $K^0 \pi^+ \pi^-$ system.² From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.³ Systematic errors not estimated.⁴ From a unitarized quark-model calculation.

NODE=M064M;LINKAGE=P
 NODE=M064M;LINKAGE=E
 NODE=M064M;LINKAGE=AB
 NODE=M064M;LINKAGE=T

 $K_1(1400)$ WIDTH


NODE=M064W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
174 ± 13 OUR AVERAGE					
Error includes scale factor of 1.6. See the ideogram below.					
300 ⁺³⁷⁰ ₋₁₁₀ ± 140	7k	ASNER	00B	CLEO	± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
188 ± 54 ± 60		⁵ ASTON	87	LASS	0 $11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
276 ± 65		BAUBILLIER	82B	HBC	0 $8.25 K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
195 ± 25		DAUM	81C	CNTR	- $63 K^- p \rightarrow K^- 2\pi p$
180 ± 10		ETKIN	80	MPS	0 $6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
142 ± 16		⁶ CARNEGIE	77	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
152 ± 16	25k	⁷ ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 200		VERGEEST	79	HBC	- $4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 160		BRANDENB...	76	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
80		DAVIS	72	HBC	+ $12 K^+ p$
241 ± 30		FIRESTONE	72B	DBC	+ $12 K^+ d$

NODE=M064W

- ⁵ From partial-wave analysis of $K^0 \pi^+ \pi^-$ system.
- ⁶ From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.
- ⁷ Systematic errors not estimated.

NODE=M064W;LINKAGE=P
 NODE=M064W;LINKAGE=E
 NODE=M064W;LINKAGE=AB



$K_1(1400)$ DECAY MODES

NODE=M064215;NODE=M064

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	(94 ± 6) %
Γ_2 $K\rho$	(3.0 ± 3.0) %
Γ_3 $Kf_0(1370)$	(2.0 ± 2.0) %
Γ_4 $K\omega$	(1.0 ± 1.0) %
Γ_5 $K_0^*(1430)\pi$	not seen
Γ_6 γK^0	seen

DESIG=1
 DESIG=2
 DESIG=8
 DESIG=5
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←

$K_1(1400)$ PARTIAL WIDTHS

NODE=M064220

Γ	VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	Γ_i
$\Gamma(K^*(892)\pi)$	117 ± 10	CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	Γ_1
$\Gamma(K\rho)$	2 ± 1	CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	Γ_2
$\Gamma(K\omega)$	23 ± 12	CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	Γ_4
$\Gamma(\gamma K^0)$	280.8 ± 23.2 ± 40.4	ALAVI-HARATI02B	KTEV		$K + A \rightarrow K^* + A$	Γ_6

NODE=M064W1
 NODE=M064W1
 NODE=M064W2
 NODE=M064W2
 NODE=M064W5
 NODE=M064W5
 NODE=M064W6
 NODE=M064W6

$K_1(1400)$ BRANCHING RATIOS

NODE=M064225

Ratio	VALUE	DOCUMENT ID	TECN	COMMENT	Ratio
$\Gamma(K^*(892)\pi)/\Gamma_{total}$	0.94 ± 0.06	⁸ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	Γ_1/Γ
$\Gamma(K\rho)/\Gamma_{total}$	0.03 ± 0.03	⁸ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$	Γ_2/Γ

NODE=M064R1
 NODE=M064R1
 NODE=M064R2
 NODE=M064R2

$\Gamma(K f_0(1370))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.02±0.02	⁸ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

NODE=M064R5
 NODE=M064R5

 $\Gamma(K\omega)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.01±0.01	⁸ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

NODE=M064R3
 NODE=M064R3

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	⁸ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

NODE=M064R4
 NODE=M064R4

D-wave/S-wave RATIO FOR $K_1(1400) \rightarrow K^*(892)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.04±0.01	⁸ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

NODE=M064R9
 NODE=M064R9

⁸ Average from low and high t data.

NODE=M064R;LINKAGE=F

 $K_1(1400)$ REFERENCES

ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)

NODE=M064

REFID=51037
 REFID=48822
 REFID=47766
 REFID=40234
 REFID=22551
 REFID=20573
 REFID=22548
 REFID=22545
 REFID=22542
 REFID=22535
 REFID=22532
 REFID=22505
 REFID=22506

$K^*(1410)$

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M094

 $K^*(1410)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1414±15 OUR AVERAGE	Error includes scale factor of 1.3.			

NODE=M094M

NODE=M094M

1380±21±19	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
1420±7±10	ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

••• We do not use the following data for averages, fits, limits, etc. •••

1276 ⁺⁷² ₋₇₇	^{1,2} BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
1367±54	BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1474±25	BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
1500±30	ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

¹ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

² Systematic uncertainties not estimated.

NODE=M094M;LINKAGE=BI

NODE=M094M;LINKAGE=NS

 $K^*(1410)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
232±21 OUR AVERAGE	Error includes scale factor of 1.1.			

NODE=M094W

NODE=M094W

176±52±22	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
240±18±12	ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

••• We do not use the following data for averages, fits, limits, etc. •••

198 ⁺⁶¹ ₋₈₇	^{3,4} BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
114±101	BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
275±65	BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
500±100	ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

³ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

⁴ Systematic uncertainties not estimated.

NODE=M094W;LINKAGE=BI

NODE=M094W;LINKAGE=NS

$K^*(1410)$ DECAY MODES

NODE=M094215;NODE=M094

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K^*(892)\pi$	> 40 %	95%
Γ_2 $K\pi$	(6.6 ± 1.3) %	
Γ_3 $K\rho$	< 7 %	95%
Γ_4 γK^0	seen	

DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=1
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←

 $K^*(1410)$ PARTIAL WIDTHS

NODE=M094217

$\Gamma(\gamma K^0)$				Γ_4	
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
<52.9	90	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$	

NODE=M094W1
NODE=M094W1

 $K^*(1410)$ BRANCHING RATIOS

NODE=M094220

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$				Γ_3/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.17	95	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

NODE=M094R1
NODE=M094R1

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$				Γ_2/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.16	95	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

NODE=M094R2
NODE=M094R2

$\Gamma(K\pi)/\Gamma_{\text{total}}$				Γ_2/Γ	
VALUE		DOCUMENT ID	TECN	CHG	COMMENT
0.066 ± 0.010 ± 0.008		ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M094R3
NODE=M094R3

 $K^*(1410)$ REFERENCES

NODE=M094

BOITO	09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin			REFID=52728
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>		(BELLE Collab.)	REFID=51929
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>		(FNAL KTeV Collab.)	REFID=48822
BIRD	89	SLAC-332	P.F. Bird		(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>		(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	87	NP B292 693	D. Aston <i>et al.</i>		(SLAC, NAGO, CINC, INUS)	REFID=40234
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>		(SLAC, CARL, OTTA) JP	REFID=22689
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>		(BIRM, CERN, GLAS+)	REFID=22551
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>		(BNL, CUNY) JP	REFID=22545

NODE=M019

$K_0^*(1430)$

$$I(J^P) = \frac{1}{2}(0^+)$$

See our minireview in the 1994 edition and in this edition under the $f_0(500)$.

NODE=M019

 $K_0^*(1430)$ MASS

NODE=M019M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1425 ± 50	OUR ESTIMATE			

NODE=M019M

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

1438 ± 8 ± 4	5.4k	1 LEES	14E	BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
1427 ± 4 ± 13		2 BUGG	10	RVUE	S-matrix pole
1466.6 ± 0.7 ± 3.4	141k	3 BONVICINI	08A	CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1412		4 LINK	07	FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
1461.0 ± 4.0 ± 2.1	54k	5 LINK	07B	FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
1406 ± 29		6 BUGG	06	RVUE	
1435 ± 6		7 ZHOU	06	RVUE	$K\rho \rightarrow K^- \pi^+ n$
1455 ± 20 ± 15		ABLIKIM	05Q	BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
1456 ± 8		8 ZHENG	04	RVUE	$K^- p \rightarrow K^- \pi^+ n$
~ 1419		9 BUGG	03	RVUE	11 $K^- p \rightarrow K^- \pi^+ n$

~ 1440		10 LI	03 RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
1459 ± 9	15k	11 AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1440		12 JAMIN	00 RVUE	$K p \rightarrow K p$
1436 ± 8		13 BARBERIS	98E OMEG	450 $p p \rightarrow$ $p_f p_s K^+ K^- \pi^+ \pi^-$
1415 ± 25		9 ANISOVICH	97C RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
~ 1450		14 TORNQVIST	96 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, K \pi$
1412 ± 6		15 ASTON	88 LASS	11 $K^- p \rightarrow K^- \pi^+ n$
~ 1430		BAUBILLIER	84B HBC	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 1425		16 ESTABROOKS	78 ASPK	13 $K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$
~ 1450.0		MARTIN	78 SPEC	10 $K^\pm p \rightarrow K^0 \pi p$

¹ Using both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed width $\Gamma(K_0^*(1430)) = 210$ MeV.

² S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K \pi$ and $K \eta'$, and the Adler zero near thresholds.

³ From the isobar model with a complex pole for the κ .

⁴ From a non-parametric analysis.

⁵ A Breit-Wigner mass and width.

⁶ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s-dependent width and an Adler zero near threshold.

⁷ S-matrix pole. Using ASTON 88 and assuming $K_0^*(800)$, $K_0^*(1950)$.

⁸ Using ASTON 88 and assuming $K_0^*(800)$.

⁹ T-matrix pole. Reanalysis of ASTON 88 data.

¹⁰ Breit-Wigner fit. Using ASTON 88.

¹¹ Assuming a low-mass scalar $K \pi$ resonance, $\kappa(800)$.

¹² T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.

¹³ J^P not determined, could be $K_2^*(1430)$.

¹⁴ T-matrix pole.

¹⁵ Uses a model for the background, without this background they get a mass 1340 MeV, where the phase shift passes 90° .

¹⁶ Mass defined by pole position. From elastic $K \pi$ partial-wave analysis.

NODE=M019M;LINKAGE=LE

NODE=M019M;LINKAGE=BG

NODE=M019M;LINKAGE=BO

NODE=M019M;LINKAGE=LI

NODE=M019M;LINKAGE=BW

NODE=M019M;LINKAGE=BU

NODE=M019M;LINKAGE=ZU

NODE=M019M;LINKAGE=ZH

NODE=M019M;LINKAGE=A1

NODE=M019M;LINKAGE=E

NODE=M019M;LINKAGE=A0

NODE=M019M;LINKAGE=JM

NODE=M019M;LINKAGE=JP

NODE=M019M;LINKAGE=TT

NODE=M019M;LINKAGE=D

NODE=M019M;LINKAGE=A

$K_0^*(1430)$ WIDTH

NODE=M019W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
270 ± 80	OUR ESTIMATE			

NODE=M019W

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

210 ± 20 ± 12	5.4k	1 LEES	14E BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta / \pi^0$
270 ± 10 ± 40		2 BUGG	10 RVUE	S-matrix pole
174.2 ± 1.9 ± 3.2	141k	3 BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 500		4 LINK	07 FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
177.0 ± 8.0 ± 3.4	54k	5 LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
350 ± 40		6 BUGG	06 RVUE	
288 ± 22		7 ZHOU	06 RVUE	$K p \rightarrow K^- \pi^+ n$
270 ± 45 ⁺³⁰ ₋₃₅		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
217 ± 31		8 ZHENG	04 RVUE	$K^- p \rightarrow K^- \pi^+ n$
~ 316		9 BUGG	03 RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
~ 350		10 LI	03 RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
175 ± 17	15k	11 AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 300		12 JAMIN	00 RVUE	$K p \rightarrow K p$
196 ± 45		13 BARBERIS	98E OMEG	450 $p p \rightarrow$ $p_f p_s K^+ K^- \pi^+ \pi^-$
330 ± 50		9 ANISOVICH	97C RVUE	11 $K^- p \rightarrow K^- \pi^+ n$
~ 320		14 TORNQVIST	96 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, K \pi$
294 ± 23		ASTON	88 LASS	11 $K^- p \rightarrow K^- \pi^+ n$
~ 200		BAUBILLIER	84B HBC	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
200 to 300		15 ESTABROOKS	78 ASPK	13 $K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$

¹ Using both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed mass $M(K_0^*(1430)) = 1435$ MeV.

NODE=M019W;LINKAGE=LE

² S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K \pi$ and $K \eta'$, and the Adler zero near thresholds.

NODE=M019W;LINKAGE=BG

³ From the isobar model with a complex pole for the κ .

NODE=M019W;LINKAGE=BO

⁴ From a non-parametric analysis.

NODE=M019W;LINKAGE=LI

- ⁵ A Breit-Wigner mass and width.
⁶ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s -dependent width and an Adler zero near threshold.
⁷ S-matrix pole. Using ASTON 88 and assuming $K_0^*(800)$, $K_0^*(1950)$.
⁸ Using ASTON 88 and assuming $K_0^*(800)$.
⁹ T-matrix pole. Reanalysis of ASTON 88 data.
¹⁰ Breit-Wigner fit. Using ASTON 88.
¹¹ Assuming a low-mass scalar $K\pi$ resonance, $\kappa(800)$.
¹² T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.
¹³ J^P not determined, could be $K_2^*(1430)$.
¹⁴ T-matrix pole.
¹⁵ From elastic $K\pi$ partial-wave analysis.

NODE=M019W;LINKAGE=BW
 NODE=M019W;LINKAGE=BU
 NODE=M019W;LINKAGE=ZU
 NODE=M019W;LINKAGE=ZH
 NODE=M019W;LINKAGE=A1
 NODE=M019W;LINKAGE=E
 NODE=M019W;LINKAGE=A0
 NODE=M019W;LINKAGE=JM
 NODE=M019W;LINKAGE=JP
 NODE=M019W;LINKAGE=TT
 NODE=M019W;LINKAGE=C

$K_0^*(1430)$ DECAY MODES

NODE=M019215;NODE=M019

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(93 \pm 10) %
Γ_2 $K\eta$	(8.6 ⁺ ₋ 2.7 ⁺ _{3.4}) %
Γ_3 $K\eta'(958)$	seen

DESIG=1
 DESIG=2
 DESIG=3

$K_0^*(1430)$ BRANCHING RATIOS

NODE=M019220

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.93\pm0.04\pm0.09	ASTON	88	LASS	0 11 $K^-p \rightarrow K^- \pi^+ n$

NODE=M019R1
 NODE=M019R1

$\Gamma(K\eta)/\Gamma(K\pi)$	Γ_2/Γ_1			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.2\pm2.5⁺₋2.5	5.4k	¹ LEES	14E	BABR $\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$

NODE=M019R01
 NODE=M019R01

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a Dalitz analysis in the presence of several interfering scalar-meson resonances.

NODE=M019R01;LINKAGE=LE

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$	Γ_3/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M019R00
 NODE=M019R00

$K_0^*(1430)$ REFERENCES

NODE=M019

ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55901
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55937
BUGG	10	PR D81 014002	D.V. Bugg	(LOQM)	REFID=53213
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=53056
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=52426
LINK	07	PL B648 156	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51702
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51875
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51051
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51458
BUGG	06	PL B632 471	D.V. Bugg	(LOQM)	REFID=50996
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng		REFID=51198
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>		REFID=50165
BUGG	03	PL B572 1	D.V. Bugg		REFID=49586
LI	03	PR D67 034025	L. Li, B. Zou, G. Li		REFID=49192
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48807
JAMIN	00	NP B587 331	M. Jamin <i>et al.</i>		REFID=47983
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46348
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev		REFID=45815
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=22446

$K_2^*(1430)$

$$I(J^P) = \frac{1}{2}(2^+)$$

We consider that phase-shift analyses provide more reliable determinations of the mass and width.

NODE=M022

NODE=M022

 $K_2^*(1430)$ MASS

NODE=M022205

CHARGED ONLY, WITH FINAL STATE $K\pi$ NODE=M022M1
NODE=M022M1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1425.6 ± 1.5 OUR AVERAGE		Error includes scale factor of 1.1.			
1420 ± 4	1587	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1436 ± 5.5	400	^{1,2} CLELAND	82	SPEC	+ 30 $K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1500	^{1,2} CLELAND	82	SPEC	+ 50 $K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1200	^{1,2} CLELAND	82	SPEC	- 50 $K^+ p \rightarrow K_S^0 \pi^- p$
1423 ± 5	935	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1428.0 ± 4.6		³ MARTIN	78	SPEC	+ 10 $K^\pm p \rightarrow K_S^0 \pi p$
1423.8 ± 4.6		³ MARTIN	78	SPEC	- 10 $K^\pm p \rightarrow K_S^0 \pi p$
1420.0 ± 3.1	1400	AGUILAR-...	71B	HBC	- 3.9,4.6 $K^- p$
1425 ± 8.0	225	^{1,2} BARNHAM	71C	HBC	+ $K^+ p \rightarrow K^0 \pi^+ p$
1416 ± 10	220	CRENNELL	69D	DBC	- 3.9 $K^- N \rightarrow \bar{K}^0 \pi^- N$
1414 ± 13.0	60	¹ LIND	69	HBC	+ 9 $K^+ p \rightarrow K^0 \pi^+ p$
1427 ± 12	63	¹ SCHWEING...	68	HBC	- 5.5 $K^- p \rightarrow \bar{K} \pi N$
1423 ± 11.0	39	¹ BASSANO	67	HBC	- 4.6-5.0 $K^- p \rightarrow \bar{K}^0 \pi^- p$

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1423.4 ± 2 ± 3	24809 ± 820	⁴ BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
----------------	-------------	-------------------	----	------	--

NEUTRAL ONLYNODE=M022M4
NODE=M022M4

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1432.4 ± 1.3 OUR AVERAGE				
1431.2 ± 1.8 ± 0.7		⁵ ASTON	88	LASS 11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 4 ± 6		⁵ ASTON	87	LASS 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1433 ± 6 ± 10		⁵ ASTON	84B	LASS 11 $K^- p \rightarrow \bar{K}^0 2\pi n$
1471 ± 12		⁵ BAUBILLIER	82B	HBC 8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
1428 ± 3		⁵ ASTON	81C	LASS 11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 2		⁵ ESTABROOKS	78	ASPK 13 $K^\pm p \rightarrow pK\pi$
1440 ± 10		⁵ BOWLER	77	DBC 5.5 $K^+ d \rightarrow K\pi pp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1428.5 ± 3.9	1786 ± 127	⁶ AUBERT	07AK	BABR 10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
1420 ± 7	300	HENDRICK	76	DBC 8.25 $K^+ N \rightarrow K^+ \pi N$
1421.6 ± 4.2	800	MCCUBBIN	75	HBC 3.6 $K^- p \rightarrow K^- \pi^+ n$
1420.1 ± 4.3		⁷ LINGLIN	73	HBC 2-13 $K^+ p \rightarrow K^+ \pi^- X$
1419.1 ± 3.7	1800	AGUILAR-...	71B	HBC 3.9,4.6 $K^- p$
1416 ± 6	600	CORDS	71	DBC 9 $K^+ n \rightarrow K^+ \pi^- p$
1421.1 ± 2.6	2200	DAVIS	69	HBC 12 $K^+ p \rightarrow K^+ \pi^- X$

¹ Errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

² Number of events in peak re-evaluated by us.

³ Systematic error added by us.

⁴ From a partial wave amplitude analysis.

⁵ From phase shift or partial-wave analysis.

⁶ Systematic errors not estimated.

⁷ From pole extrapolation, using world $K^+ p$ data summary tape.

NODE=M022M;LINKAGE=D

NODE=M022M;LINKAGE=W

NODE=M022M;LINKAGE=B

NODE=M022M;LINKAGE=F

NODE=M022M;LINKAGE=P

NODE=M022M4;LINKAGE=NS

NODE=M022M;LINKAGE=C

K₂^{*}(1430) WIDTH

NODE=M022210

CHARGED ONLY, WITH FINAL STATE Kπ

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
98.5 ± 2.7 OUR FIT	Error includes scale factor of 1.1.				
98.5 ± 2.9 OUR AVERAGE	Error includes scale factor of 1.1.				
109 ± 22	400	8,9 CLELAND	82	SPEC +	30 K ⁺ p → K _S ⁰ π ⁺ p
124 ± 12.8	1500	8,9 CLELAND	82	SPEC +	50 K ⁺ p → K _S ⁰ π ⁺ p
113 ± 12.8	1200	8,9 CLELAND	82	SPEC -	50 K ⁺ p → K _S ⁰ π ⁻ p
85 ± 16	935	TOAFF	81	HBC -	6.5 K ⁻ p → $\bar{K}^0\pi^-p$
96.5 ± 3.8		MARTIN	78	SPEC +	10 K [±] p → K _S ⁰ πp
97.7 ± 4.0		MARTIN	78	SPEC -	10 K [±] p → K _S ⁰ πp
94.7 ^{+15.1} _{-12.5}	1400	AGUILAR-...	71B	HBC -	3.9,4.6 K ⁻ p
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
98 ± 4 ± 4	25k	¹⁰ BIRD	89	LASS -	11 K ⁻ p → $\bar{K}^0\pi^-p$

NODE=M022W1
NODE=M022W1

OCCUR=2
OCCUR=3


OCCUR=2

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
109 ± 5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.			
116.5 ± 3.6 ± 1.7		¹¹ ASTON	88	LASS 11 K ⁻ p → K ⁻ π ⁺ n
129 ± 15 ± 15		¹¹ ASTON	87	LASS 11 K ⁻ p → $\bar{K}^0\pi^+\pi^-n$
131 ± 24 ± 20		¹¹ ASTON	84B	LASS 11 K ⁻ p → $\bar{K}^02\pi n$
143 ± 34		¹¹ BAUBILLIER	82B	HBC 8.25 K ⁻ p → NK _S ⁰ ππ
98 ± 8		¹¹ ASTON	81C	LASS 11 K ⁻ p → K ⁻ π ⁺ n
140 ± 30		¹¹ ETKIN	80	SPEC 6 K ⁻ p → $\bar{K}^0\pi^+\pi^-n$
98 ± 5		¹¹ ESTABROOKS	78	ASPK 13 K [±] p → pKπ
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
113.7 ± 9.2	1786 ± 127	¹² AUBERT	07AK	BABR 10.6 e ⁺ e ⁻ → K ^{*0} K [±] π [∓] γ
125 ± 29	300	⁸ HENDRICK	76	DBC 8.25 K ⁺ N → K ⁺ πN
116 ± 18	800	MCCUBBIN	75	HBC 3.6 K ⁻ p → K ⁻ π ⁺ n
61 ± 14		¹³ LINGLIN	73	HBC 2-13 K ⁺ p → K ⁺ π ⁻ X
116.6 ^{+10.3} _{-15.5}	1800	AGUILAR-...	71B	HBC 3.9,4.6 K ⁻ p
144 ± 24.0	600	⁸ CORDS	71	DBC 9 K ⁺ n → K ⁺ π ⁻ p
101 ± 10	2200	DAVIS	69	HBC 12 K ⁺ p → K ⁺ π ⁻ π ⁺ p

NODE=M022W4
NODE=M022W4

WEIGHTED AVERAGE
109±5 (Error scaled by 1.9)



⁸ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the K^{*}(892) mass.
⁹ Number of events in peak re-evaluated by us.
¹⁰ From a partial wave amplitude analysis.
¹¹ From phase shift or partial-wave analysis.
¹² Systematic errors not estimated.
¹³ From pole extrapolation, using world K⁺p data summary tape.

NODE=M022W;LINKAGE=D
 NODE=M022W;LINKAGE=W
 NODE=M022W;LINKAGE=F
 NODE=M022W;LINKAGE=P
 NODE=M022W4;LINKAGE=NS
 NODE=M022W;LINKAGE=C

$K_2^*(1430)$ DECAY MODES

NODE=M022215;NODE=M022

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $K\pi$	$(49.9 \pm 1.2) \%$	
Γ_2 $K^*(892)\pi$	$(24.7 \pm 1.5) \%$	
Γ_3 $K^*(892)\pi\pi$	$(13.4 \pm 2.2) \%$	
Γ_4 $K\rho$	$(8.7 \pm 0.8) \%$	S=1.2
Γ_5 $K\omega$	$(2.9 \pm 0.8) \%$	
Γ_6 $K^+\gamma$	$(2.4 \pm 0.5) \times 10^{-3}$	S=1.1
Γ_7 $K\eta$	$(1.5^{+3.4}_{-1.0}) \times 10^{-3}$	S=1.3
Γ_8 $K\omega\pi$	$< 7.2 \times 10^{-4}$	CL=95%
Γ_9 $K^0\gamma$	$< 9 \times 10^{-4}$	CL=90%

DESIG=1
DESIG=2
DESIG=6
DESIG=3
DESIG=4
DESIG=8
DESIG=5
DESIG=7
DESIG=10;OUR EVAL;→ UNCHECKED ←

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 10 branching ratios uses 31 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 20.2$ for 24 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-9						
x_3	-40	-73					
x_4	-8	36	-52				
x_5	-11	-3	-26	-7			
x_6	-1	-1	-1	-1	0		
x_7	-4	-7	-5	-5	-2	0	
Γ	0	0	0	0	0	-13	0
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode	Rate (MeV)	Scale factor
Γ_1 $K\pi$	49.1 ± 1.8	
Γ_2 $K^*(892)\pi$	24.3 ± 1.6	
Γ_3 $K^*(892)\pi\pi$	13.2 ± 2.2	
Γ_4 $K\rho$	8.5 ± 0.8	1.2
Γ_5 $K\omega$	2.9 ± 0.8	
Γ_6 $K^+\gamma$	0.24 ± 0.05	1.1
Γ_7 $K\eta$	$0.15^{+0.33}_{-0.10}$	1.3

NODE=M022220

 $K_2^*(1430)$ PARTIAL WIDTHS **$\Gamma(K^+\gamma)$** **Γ_6**

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
241 ± 50 OUR FIT				Error includes scale factor of 1.1.
240 ± 45	CIHANGIR	82	SPEC +	$200 K^+ Z \rightarrow Z K^+ \pi^0,$ $Z K_S^0 \pi^+$

NODE=M022W8
NODE=M022W8 **$\Gamma(K^0\gamma)$** **Γ_9**

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 5.4	90	ALAVI-HARATI02B	KTEV		$K + A \rightarrow K^* + A$
•••					We do not use the following data for averages, fits, limits, etc. •••
<84	90	CARLSMITH	87	SPEC 0	$60-200 K_L^0 A \rightarrow$ $K_S^0 \pi^0 A$

NODE=M022W9
NODE=M022W9

K^{*}(1430) BRANCHING RATIOS

NODE=M022225

Γ(Kπ)/Γ_{total}

Γ₁/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.499±0.012 OUR FIT				
0.488±0.014 OUR AVERAGE				
0.485±0.006±0.020	¹⁴ ASTON	88	LASS	0 11 K ⁻ p → K ⁻ π ⁺ n
0.49 ±0.02	¹⁴ ESTABROOKS	78	ASPK	± 13 K [±] p → pKπ

NODE=M022R1
NODE=M022R1

Γ(K*(892)π)/Γ(Kπ)

Γ₂/Γ₁

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.496±0.034 OUR FIT				
0.47 ±0.04 OUR AVERAGE				
0.44 ±0.09	ASTON	84B	LASS	0 11 K ⁻ p → \bar{K}^0 2πn
0.62 ±0.19	LAUSCHER	75	HBC	0 10,16 K ⁻ p → K ⁻ π ⁺ n
0.54 ±0.16	DEHM	74	DBC	0 4.6 K ⁺ N
0.47 ±0.08	AGUILAR-...	71B	HBC	3.9,4.6 K ⁻ p
0.47 ±0.10	BASSANO	67	HBC	-0 4.6,5.0 K ⁻ p
0.45 ±0.13	BADIER	65C	HBC	- 3 K ⁻ p

NODE=M022R4
NODE=M022R4

Γ(Kω)/Γ(Kπ)

Γ₅/Γ₁

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.059±0.017 OUR FIT				
0.070±0.035 OUR AVERAGE				
0.05 ±0.04	AGUILAR-...	71B	HBC	3.9,4.6 K ⁻ p
0.13 ±0.07	BASSOMPIE...	69	HBC	0 5 K ⁺ p

NODE=M022R5
NODE=M022R5

OCCUR=2

Γ(Kρ)/Γ(Kπ)

Γ₄/Γ₁

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.174±0.017 OUR FIT	Error includes scale factor of 1.2.			
0.150^{+0.029}_{-0.017} OUR AVERAGE				
0.18 ±0.05	ASTON	84B	LASS	0 11 K ⁻ p → \bar{K}^0 2πn
0.02 ^{+0.10} _{-0.02}	DEHM	74	DBC	0 4.6 K ⁺ N
0.16 ±0.05	AGUILAR-...	71B	HBC	3.9,4.6 K ⁻ p
0.14 ±0.10	BASSANO	67	HBC	-0 4.6,5.0 K ⁻ p
0.14 ±0.07	BADIER	65C	HBC	- 3 K ⁻ p

NODE=M022R6
NODE=M022R6


Γ(Kρ)/Γ(K*(892)π)

Γ₄/Γ₂

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.350±0.031 OUR FIT	Error includes scale factor of 1.4.			
0.354±0.033 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
0.293±0.032±0.020	ASTON	87	LASS	0 11 K ⁻ p → \bar{K}^0 π ⁺ π ⁻ n
0.38 ±0.09	BAUBILLIER	82B	HBC	0 8.25 K ⁻ p → NK _S ⁰ ππ
0.39 ±0.03	DAUM	81C	CNTR	63 K ⁻ p → K ⁻ 2πp

NODE=M022R7
NODE=M022R7

WEIGHTED AVERAGE
0.354±0.033 (Error scaled by 1.4)



$\Gamma(K\omega)/\Gamma(K^*(892)\pi)$						Γ_5/Γ_2	
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			
0.118±0.034 OUR FIT							NODE=M022R8
0.10 ±0.04	FIELD	67	HBC	—	3.8 $K^- p$		NODE=M022R8
$\Gamma(K\eta)/\Gamma(K^*(892)\pi)$						Γ_7/Γ_2	
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			
0.006^{+0.014}_{-0.004} OUR FIT Error includes scale factor of 1.2.							NODE=M022R9
0.07 ±0.04	FIELD	67	HBC	—	3.8 $K^- p$		NODE=M022R9
$\Gamma(K\eta)/\Gamma(K\pi)$						Γ_7/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT		
0.0030^{+0.0070}_{-0.0020} OUR FIT Error includes scale factor of 1.3.							NODE=M022R10
0 ±0.0056		15 ASTON	88B	LASS	—	11 $K^- p \rightarrow K^- \eta p$	NODE=M022R10
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●							
<0.04	95	AGUILAR-...	71B	HBC		3.9,4.6 $K^- p$	
<0.065		16 BASSOMPIE...	69	HBC		5.0 $K^+ p$	
<0.02		BISHOP	69	HBC		3.5 $K^+ p$	
$\Gamma(K^*(892)\pi\pi)/\Gamma_{total}$						Γ_3/Γ	
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			
0.134±0.022 OUR FIT							NODE=M022R11
0.12 ±0.04	17	GOLDBERG	76	HBC	—	3 $K^- p \rightarrow p \bar{K}^0 \pi \pi \pi$	NODE=M022R11
$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$						Γ_3/Γ_1	
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			
0.27±0.05 OUR FIT							NODE=M022R12
0.21±0.08	16,17	JONGEJANS	78	HBC	—	4 $K^- p \rightarrow p \bar{K}^0 \pi \pi \pi$	NODE=M022R12
$\Gamma(K\omega\pi)/\Gamma_{total}$						Γ_8/Γ	
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		
<0.72	95	0	JONGEJANS	78	HBC	4 $K^- p \rightarrow p \bar{K}^0 4\pi$	NODE=M022R13
14 From phase shift analysis.							
15 ASTON 88B quote < 0.0092 at CL=95%. We convert this to a central value and 1 sigma error in order to be able to use it in our constrained fit.							
16 Restated by us.							
17 Assuming $\pi\pi$ system has isospin 1, which is supported by the data.							

$K_2^*(1430)$ REFERENCES

AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)	REFID=48822
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40281
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
CARLSMITH	87	PR D36 3502	D. Carlsmith <i>et al.</i>	(EFI, SACL)	REFID=40557
ASTON	84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)	REFID=22763
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
CIHANGIR	82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)	REFID=21280
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22821
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=22548
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
JONGEJANS	78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)	REFID=22445
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=22446
BOWLER	77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)	REFID=22437
GOLDBERG	76	LCN 17 253	J. Goldberg	(HAIF)	REFID=22742
HENDRICK	76	NP B112 189	K. Hendrickx <i>et al.</i>	(MONS, SACL, PARIS+)	REFID=22743
LAUSCHER	75	NP B86 189	P. Lauscher <i>et al.</i>	(ABCLV Collab.) JP	REFID=22582
MCCUBBIN	75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)	REFID=22434
DEHM	74	NP B75 47	G. Dehm <i>et al.</i>	(MPIM, BRUX, MONS, CERN)	REFID=22736
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
AGUILAR-...	71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)	REFID=22408
BARNHAM	71C	NP B28 171	K.W.J. Barnham <i>et al.</i>	(BIRM, GLAS)	REFID=22409
CORDS	71	PR D4 1974	D. Cords <i>et al.</i>	(PURD, UCD, IUPU)	REFID=22411
BASSOMPIE...	69	NP B13 189	G. Bassompierre <i>et al.</i>	(CERN, BRUX) JP	REFID=22710
BISHOP	69	NP B9 403	J.M. Bishop <i>et al.</i>	(WISC)	REFID=22485
CRENNELL	69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)	REFID=22399
DAVIS	69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)	REFID=22400
LIND	69	NP B14 1	V.G. Lind <i>et al.</i>	(LRL) JP	REFID=22404
SCHWEING... Also	68	PR 166 1317 Thesis	F. Schweingruber <i>et al.</i>	(ANL, NWES)	REFID=22398
BASSANO	67	PRL 19 968	F.L. Schweingruber	(NWES, NWES)	REFID=22709
FIELD	67	PL 24B 638	D. Bassano <i>et al.</i>	(BNL, SYRA)	REFID=22695
BADIER	65C	PL 19 612	J.H. Field <i>et al.</i>	(UCSD)	REFID=22701
			J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=22690

NODE=M022

K(1460)

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE
Observed in $K\pi\pi$ partial-wave analysis.

K(1460) MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
~ 1460	DAUM	81C	CNTR	- 63 $K^- p \rightarrow K^- 2\pi p$
~ 1400	¹ BRANDENB...	76B	ASPK	± 13 $K^\pm p \rightarrow K^\pm 2\pi p$
¹ Coupled mainly to $K f_0(1370)$. Decay into $K^*(892)\pi$ seen.				

K(1460) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
~ 260	DAUM	81C	CNTR	- 63 $K^- p \rightarrow K^- 2\pi p$
~ 250	² BRANDENB...	76B	ASPK	± 13 $K^\pm p \rightarrow K^\pm 2\pi p$
² Coupled mainly to $K f_0(1370)$. Decay into $K^*(892)\pi$ seen.				

K(1460) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K\rho$	seen
Γ_3 $K_0^*(1430)\pi$	seen

K(1460) PARTIAL WIDTHS **$\Gamma(K^*(892)\pi)$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
~ 109	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

 $\Gamma(K\rho)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
~ 34	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

 $\Gamma(K_0^*(1430)\pi)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
~ 117	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

K(1460) REFERENCES

DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BRANDENB...	76B	PRL 36 1239	G.W. Brandenburg <i>et al.</i>	(SLAC)JP

NODE=M021

NODE=M021

NODE=M021M

NODE=M021M

NODE=M021M;LINKAGE=A

NODE=M021W

NODE=M021W

NODE=M021W;LINKAGE=A

NODE=M021215;NODE=M021

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

NODE=M021220

NODE=M021W1
NODE=M021W1NODE=M021W2
NODE=M021W2NODE=M021W3
NODE=M021W3

NODE=M021

REFID=22548
REFID=22767

$K_2(1580)$

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+ \pi^-$ system. Needs confirmation.

NODE=M039

NODE=M039

 $K_2(1580)$ MASS

NODE=M039M

VALUE (MeV)	DOCUMENT ID	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 1580	OTTER	79	- 10,14,16 $K^- p$

NODE=M039M

 $K_2(1580)$ WIDTH

NODE=M039W

VALUE (MeV)	DOCUMENT ID	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 110	OTTER	79	- 10,14,16 $K^- p$

NODE=M039W

 $K_2(1580)$ DECAY MODES

NODE=M039215;NODE=M039

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K_2^*(1430)\pi$	possibly seen

DESIG=1

DESIG=2

 $K_2(1580)$ BRANCHING RATIOS

NODE=M039220

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
seen	OTTER	79	HBC	- 10,14,16 $K^- p$	
••• We do not use the following data for averages, fits, limits, etc. •••					
possibly seen	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$	

NODE=M039R1
NODE=M039R1

$\Gamma(K_2^*(1430)\pi)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
possibly seen	OTTER	79	HBC	- 10,14,16 $K^- p$	

NODE=M039R2
NODE=M039R2 **$K_2(1580)$ REFERENCES**

NODE=M039

GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
OTTER	79	NP B147 I	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP

REFID=53668
REFID=22772

K(1630)

$$I(J^P) = \frac{1}{2}(?^?)$$

OMITTED FROM SUMMARY TABLE

Seen as a narrow peak, compatible with the experimental resolution, in the invariant mass of the $K_S^0 \pi^+ \pi^-$ system produced in $\pi^- p$ interactions at high momentum transfers.

NODE=M160

NODE=M160

K(1630) MASS

NODE=M160M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1629±7	~ 75	KARNAUKHOV98	BC	16.0 $\pi^- p \rightarrow (K_S^0 \pi^+ \pi^-)$ $X^+ \pi^- X^0$

NODE=M160M

K(1630) WIDTH

NODE=M160W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
16⁺¹⁹₋₁₆	~ 75	¹ KARNAUKHOV98	BC	16.0 $\pi^- p \rightarrow (K_S^0 \pi^+ \pi^-)$ $X^+ \pi^- X^0$

NODE=M160W

¹ Compatible with an experimental resolution of 14 ± 1 MeV.

NODE=M160W;LINKAGE=A

K(1630) DECAY MODES

NODE=M160215;NODE=M160

Mode

 $\Gamma_1 \quad K_S^0 \pi^+ \pi^-$

DESIG=1

K(1630) REFERENCES

KARNAUKHOV 98 PAN 61 203 V.M. Karnaukhov, C. Coca, V.I. Moroz
Translated from YAF 61 252.

NODE=M160

REFID=46371

NODE=M099

K₁(1650)

$$I(J^P) = \frac{1}{2}(1^+)$$

OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems ($K^+ \phi$, $K \pi \pi$) reported in partial-wave analysis in the 1600–1900 mass region.

NODE=M099

K₁(1650) MASS

NODE=M099M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1650±50	FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 1840	ARMSTRONG	83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 1800	DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

NODE=M099M

K₁(1650) WIDTH

NODE=M099W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
150±50	FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 250	DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

NODE=M099W

K₁(1650) DECAY MODES

NODE=M099215;NODE=M099

Mode

 $\Gamma_1 \quad K \pi \pi$
 $\Gamma_2 \quad K \phi$

DESIG=1

DESIG=2

K₁(1650) REFERENCES

NODE=M099

FRAME 86 NP B276 667 D. Frame *et al.* (GLAS)
 ARMSTRONG 83 NP B221 1 T.A. Armstrong *et al.* (BARI, BIRM, CERN+)
 DAUM 81C NP B187 1 C. Daum *et al.* (AMST, CERN, CRAC, MPIM+)

REFID=20569
REFID=22801
REFID=22548

$K^*(1680)$

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M095

 $K^*(1680)$ MASS

NODE=M095M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1717±27 OUR AVERAGE	Error includes scale factor of 1.4.			
1677±10±32	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
1735±10±20	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1678±64	BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1800±70	ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
~ 1650	ESTABROOKS	78	ASPK	0 13 $K^\pm p \rightarrow K^\pm \pi^\pm n$

NODE=M095M

 $K^*(1680)$ WIDTH

NODE=M095W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
322±110 OUR AVERAGE	Error includes scale factor of 4.2.			
205± 16±34	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
423± 18±30	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
454±270	BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
170± 30	ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
250 to 300	ESTABROOKS	78	ASPK	0 13 $K^\pm p \rightarrow K^\pm \pi^\pm n$

NODE=M095W

 $K^*(1680)$ DECAY MODES

NODE=M095215;NODE=M095

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(38.7±2.5) %
Γ_2 $K\rho$	(31.4 ^{+5.0} _{-2.1}) %
Γ_3 $K^*(892)\pi$	(29.9 ^{+2.2} _{-5.0}) %

DESIG=1

DESIG=3

DESIG=2

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 2.9$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_2 & -36 & \\ x_3 & -39 & -72 \\ \hline & x_1 & x_2 \end{array}$$

 $K^*(1680)$ BRANCHING RATIOS

NODE=M095220

$\Gamma(K\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.387±0.026 OUR FIT				
0.388±0.014±0.022	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M095R4

NODE=M095R4

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$ Γ_1/Γ_3

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.30^{+0.23}_{-0.14} OUR FIT				
2.8 ±1.1	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

NODE=M095R2

NODE=M095R2

$\Gamma(K\rho)/\Gamma(K\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.81^{+0.14}_{-0.09} OUR FIT

1.2 ± 0.4	ASTON	84	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
------------------	-------	----	------	---	---

 $\Gamma(K\rho)/\Gamma(K^*(892)\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

1.05^{+0.27}_{-0.11} OUR FIT

0.97 ± 0.09^{+0.30}_{-0.10}	ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
--	-------	----	------	---	--

NODE=M095R3
NODE=M095R3NODE=M095R1
NODE=M095R1**K*(1680) REFERENCES**

BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP

NODE=M095

REFID=41002
REFID=40262
REFID=40234
REFID=22689
REFID=22545
REFID=22443

NODE=M023

K₂(1770)

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this *Review*, PDG 04.

NODE=M023

K₂(1770) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1773 ± 8		¹ ASTON	93	LASS	11 $K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1743 ± 15		TIKHOMIROV	03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1810 ± 20		FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
~ 1730		ARMSTRONG	83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 1780		² DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$
1710 ± 15	60	CHUNG	74	HBC -	7.3 $K^- p \rightarrow K^- \omega p$
1767 ± 6		BLIEDEN	72	MMS -	11-16 $K^- p$
1730 ± 20	306	³ FIRESTONE	72B	DBC +	12 $K^+ d$
1765 ± 40		⁴ COLLEY	71	HBC +	10 $K^+ p \rightarrow K 2\pi N$
1740		DENEGRI	71	DBC -	12.6 $K^- d \rightarrow \bar{K} 2\pi d$
1745 ± 20		AGUILAR-...	70C	HBC -	4.6 $K^- p$
1780 ± 15		BARTSCH	70C	HBC -	10.1 $K^- p$
1760 ± 15		LUDLAM	70	HBC -	12.6 $K^- p$

NODE=M023M

NODE=M023M

¹ From a partial wave analysis of the $K^- \omega$ system.² From a partial wave analysis of the $K^- 2\pi$ system.³ Produced in conjunction with excited deuteron.⁴ Systematic errors added correspond to spread of different fits.NODE=M023M;LINKAGE=A
NODE=M023M;LINKAGE=B
NODE=M023M;LINKAGE=P
NODE=M023M;LINKAGE=X**K₂(1770) WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
186 ± 14		⁵ ASTON	93	LASS	11 $K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
147 ± 70		TIKHOMIROV	03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
140 ± 40		FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
~ 220		ARMSTRONG	83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 210		⁶ DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$
110 ± 50	60	CHUNG	74	HBC -	7.3 $K^- p \rightarrow K^- \omega p$
100 ± 26		BLIEDEN	72	MMS -	11-16 $K^- p$
210 ± 30	306	⁷ FIRESTONE	72B	DBC +	12 $K^+ d$
90 ± 70		⁸ COLLEY	71	HBC +	10 $K^+ p \rightarrow K 2\pi N$
130		DENEGRI	71	DBC -	12.6 $K^- d \rightarrow \bar{K} 2\pi d$
100 ± 50		AGUILAR-...	70C	HBC -	4.6 $K^- p$
138 ± 40		BARTSCH	70C	HBC -	10.1 $K^- p$
50 ⁺⁴⁰ ₋₂₀		LUDLAM	70	HBC -	12.6 $K^- p$

NODE=M023W

NODE=M023W

- ⁵ From a partial wave analysis of the $K^- \omega$ system.
⁶ From a partial wave analysis of the $K^- 2\pi$ system.
⁷ Produced in conjunction with excited deuteron.
⁸ Systematic errors added correspond to spread of different fits.

NODE=M023W;LINKAGE=B
 NODE=M023W;LINKAGE=C
 NODE=M023W;LINKAGE=P
 NODE=M023W;LINKAGE=X

$K_2(1770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi \pi$	
Γ_2 $K_2^*(1430)\pi$	dominant
Γ_3 $K^*(892)\pi$	seen
Γ_4 $K f_2(1270)$	seen
Γ_5 $K f_0(980)$	
Γ_6 $K \phi$	seen
Γ_7 $K \omega$	seen

NODE=M023215;NODE=M023

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←
 DESIG=11
 DESIG=10
 DESIG=8

$K_2(1770)$ BRANCHING RATIOS

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$ Γ_2/Γ_1
 $(K_2^*(1430) \rightarrow K\pi)$

NODE=M023220

NODE=M023R1

NODE=M023R1
 NODE=M023R1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

~ 0.03	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$
~ 1.0	⁹ FIRESTONE	72B	DBC +	12 $K^+ d$
<1.0	COLLEY	71	HBC	10 $K^+ p$
0.2 ± 0.2	AGUILAR-...	70C	HBC -	4.6 $K^- p$
<1.0	BARTSCH	70C	HBC -	10.1 $K^- p$
1.0	BARBARO-...	69	HBC +	12.0 $K^+ p$

⁹ Produced in conjunction with excited deuteron.

NODE=M023R1;LINKAGE=P

$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$ Γ_3/Γ_1

NODE=M023R3
 NODE=M023R3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

~ 0.23	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$
--------	------	-----	------	-----------------------------------

$\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$ Γ_4/Γ_1
 $(f_2(1270) \rightarrow \pi\pi)$

NODE=M023R4

NODE=M023R4
 NODE=M023R4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

~ 0.74	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$
--------	------	-----	------	-----------------------------------

$\Gamma(K f_0(980))/\Gamma_{total}$ Γ_5/Γ

NODE=M023R6
 NODE=M023R6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

possibly seen TIKHOMIROV 03 SPEC 40.0 $\frac{\pi^- C}{K_S^0 K_S^0 K_L^0 X}$

$\Gamma(K\phi)/\Gamma_{total}$ Γ_6/Γ

NODE=M023R5
 NODE=M023R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

seen ARMSTRONG 83 OMEG - 18.5 $K^- p \rightarrow K^- \phi N$

$\Gamma(K\omega)/\Gamma_{total}$ Γ_7/Γ

NODE=M023R2
 NODE=M023R2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

seen OTTER 81 HBC ± 8.25,10,16 $K^\pm p$

seen CHUNG 74 HBC - 7.3 $K^- p \rightarrow K^- \omega p$

K₂(1770) REFERENCES

PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
TIKHOMIROV	03	PAN 66 828 Translated from YAF 66 860.	G.D. Tikhomirov <i>et al.</i>	
ASTON	93	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
OTTER	81	NP B181 1	G. Otter	(AACH3, BERL, LOIC, VIEN, BIRM+)
CHUNG	74	PL 51B 413	S.U. Chung <i>et al.</i>	(BNL)
BLIEDEN	72	PL 39B 668	H.R. Blieden <i>et al.</i>	(STON, NEAS)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)
COLLEY	71	NP B26 71	D.C. Colley <i>et al.</i>	(BIRM, GLAS)
DENEGRI	71	NP B28 13	D. Denegri <i>et al.</i>	(JHU) JP
AGUILAR...	70C	PRL 25 54	M. Aguilar-Benitez <i>et al.</i>	(BNL)
BARTSCH	70C	PL 33B 186	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN+)
LUDLAM	70	PR D2 1234	T. Ludlam, J. Sandweiss, A.J. Slaughter	(YALE)
BARBARO...	69	PRL 22 1207	A. Barbaro-Galtieri <i>et al.</i>	(LRL)

NODE=M023

REFID=49653
REFID=49423REFID=43597
REFID=20569
REFID=22801
REFID=22548
REFID=22549
REFID=22735
REFID=22788
REFID=22506
REFID=22785
REFID=22497
REFID=22782
REFID=22783
REFID=22784
REFID=22483**K₃^{*}(1780)**

$$I(J^P) = \frac{1}{2}(3^-)$$

NODE=M060

K₃^{*}(1780) MASS

NODE=M060M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
1776 ± 7 OUR AVERAGE Error includes scale factor of 1.1.						
1781 ± 8 ± 4		¹ ASTON	88	LASS	0	11 K ⁻ p → K ⁻ π ⁺ n
1740 ± 14 ± 15		¹ ASTON	87	LASS	0	11 K ⁻ p → $\bar{K}^0 \pi^+ \pi^- n$
1779 ± 11		² BALDI	76	SPEC	+	10 K ⁺ p → K ⁰ π ⁺ p
1776 ± 26		³ BRANDENB...	76D	ASPK	0	13 K [±] p → K [±] π [∓] N
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
1720 ± 10 ± 15	6111	⁴ BIRD	89	LASS	-	11 K ⁻ p → $\bar{K}^0 \pi^- p$
1749 ± 10		ASTON	88B	LASS	-	11 K ⁻ p → K ⁻ η p
1780 ± 9	300	BAUBILLIER	84B	HBC	-	8.25 K ⁻ p → $\bar{K}^0 \pi^- p$
1790 ± 15		BAUBILLIER	82B	HBC	0	8.25 K ⁻ p → K _S ⁰ 2π N
1784 ± 9	2060	CLELAND	82	SPEC	±	50 K ⁺ p → K _S ⁰ π [±] p
1786 ± 15		⁵ ASTON	81D	LASS	0	11 K ⁻ p → K ⁻ π ⁺ n
1762 ± 9	190	TOAFF	81	HBC	-	6.5 K ⁻ p → $\bar{K}^0 \pi^- p$
1850 ± 50		ETKIN	80	MPS	0	6 K ⁻ p → $\bar{K}^0 \pi^+ \pi^-$
1812 ± 28		BEUSCH	78	OMEG		10 K ⁻ p → $\bar{K}^0 \pi^+ \pi^- n$
1786 ± 8		CHUNG	78	MPS	0	6 K ⁻ p → K ⁻ π ⁺ n

NODE=M060M

¹ From energy-independent partial-wave analysis.² From a fit to Y₆² moment. J^P = 3⁻ found.³ Confirmed by phase shift analysis of ESTABROOKS 78, yields J^P = 3⁻.⁴ From a partial wave amplitude analysis.⁵ From a fit to the Y₆⁰ moment.

NODE=M060M;LINKAGE=K

NODE=M060M;LINKAGE=M

NODE=M060M;LINKAGE=A

NODE=M060M;LINKAGE=F

NODE=M060M;LINKAGE=J

K₃^{*}(1780) WIDTH

NODE=M060W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
159 ± 21 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.						
203 ± 30 ± 8		⁶ ASTON	88	LASS	0	11 K ⁻ p → K ⁻ π ⁺ n
171 ± 42 ± 20		⁶ ASTON	87	LASS	0	11 K ⁻ p → $\bar{K}^0 \pi^+ \pi^- n$
135 ± 22		⁷ BALDI	76	SPEC	+	10 K ⁺ p → K ⁰ π ⁺ p

NODE=M060W

• • • We do not use the following data for averages, fits, limits, etc. • • •

187±31±20	6111	⁸ BIRD	89	LASS	-	11	$K^- p \rightarrow \bar{K}^0 \pi^- p$
193 ⁺⁵¹ ₋₃₇		ASTON	88B	LASS	-	11	$K^- p \rightarrow K^- \eta p$
99±30	300	BAUBILLIER	84B	HBC	-	8.25	$K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 130		BAUBILLIER	82B	HBC	0	8.25	$K^- p \rightarrow K_S^0 2\pi N$
191±24	2060	CLELAND	82	SPEC	±	50	$K^+ p \rightarrow K_S^0 \pi^\pm p$
225±60		⁹ ASTON	81D	LASS	0	11	$K^- p \rightarrow K^- \pi^+ n$
~ 80	190	TOAFF	81	HBC	-	6.5	$K^- p \rightarrow \bar{K}^0 \pi^- p$
240±50		ETKIN	80	MPS	0	6	$K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$
181±44		¹⁰ BEUSCH	78	OMEG		10	$K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
96±31		CHUNG	78	MPS	0	6	$K^- p \rightarrow K^- \pi^+ n$
270±70		¹¹ BRANDENB...	76D	ASPK	0	13	$K^\pm p \rightarrow K^\pm \pi^\mp N$

⁶ From energy-independent partial-wave analysis.

⁷ From a fit to Y_6^2 moment. $J^P = 3^-$ found.


⁸ From a partial wave amplitude analysis.

⁹ From a fit to Y_6^0 moment.

¹⁰ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

¹¹ ESTABROOKS 78 find that BRANDENBURG 76D data are consistent with 175 MeV width. Not averaged.

NODE=M060W;LINKAGE=K
 NODE=M060W;LINKAGE=M
 NODE=M060W;LINKAGE=F
 NODE=M060W;LINKAGE=J
 NODE=M060W;LINKAGE=D
 NODE=M060W;LINKAGE=E



$K_3^*(1780)$ DECAY MODES

NODE=M060215;NODE=M060

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K \rho$	(31 ± 9) %	DESIG=3
Γ_2 $K^*(892) \pi$	(20 ± 5) %	DESIG=2
Γ_3 $K \pi$	(18.8 ± 1.0) %	DESIG=1
Γ_4 $K \eta$	(30 ± 13) %	DESIG=6
Γ_5 $K_2^*(1430) \pi$	< 16 %	95% DESIG=4

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 4 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 0.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	85		
x_3	18	21	
x_4	-98	-94	-27
	x_1	x_2	x_3

K₃^{*}(1780) BRANCHING RATIOS

NODE=M060220

 $\Gamma(K\rho)/\Gamma(K^*(892)\pi)$
 Γ_1/Γ_2 NODE=M060R5
NODE=M060R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.52±0.23 OUR FIT				
1.52±0.21±0.10	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

 $\Gamma(K^*(892)\pi)/\Gamma(K\pi)$
 Γ_2/Γ_3 NODE=M060R7
NODE=M060R7

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.09±0.26 OUR FIT				
1.09±0.26	ASTON	84B	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

 $\Gamma(K\pi)/\Gamma_{\text{total}}$
 Γ_3/Γ NODE=M060R4
NODE=M060R4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.188±0.010 OUR FIT				
0.188±0.010 OUR AVERAGE				
0.187±0.008±0.008	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
0.19 ±0.02	ESTABROOKS	78	ASPK	0 13 $K^\pm p \rightarrow K\pi N$

 $\Gamma(K\eta)/\Gamma(K\pi)$
 Γ_4/Γ_3 NODE=M060R8
NODE=M060R8

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.6 ±0.7 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.41±0.050	¹² BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
0.50±0.18	ASTON	88B	LASS	- 11 $K^- p \rightarrow K^- \eta p$

¹²This result supersedes ASTON 88B.

NODE=M060R8;LINKAGE=H

 $\Gamma(K_2^*(1430)\pi)/\Gamma(K^*(892)\pi)$
 Γ_5/Γ_2 NODE=M060R6
NODE=M060R6

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.78	95	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

K₃^{*}(1780) REFERENCES

NODE=M060

BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP	REFID=40281
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
ASTON	84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)	REFID=22763
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81D	PL 99B 502	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22820
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
BEUSCH	78	PL 74B 282	W. Beusch <i>et al.</i>	(CERN, AACH3, ETH) JP	REFID=22537
CHUNG	78	PRL 40 355	S.U. Chung <i>et al.</i>	(BNL, BRAN, CUNY+) JP	REFID=22814
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP	REFID=22443
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
BALDI	76	PL 63B 344	R. Baldi <i>et al.</i>	(GEVA) JP	REFID=22807
BRANDENB...	76D	PL 60B 478	G.W. Brandenburg <i>et al.</i>	(SLAC) JP	REFID=22808

$K_2(1820)$

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this *Review* (PDG 04) under $K_2(1770)$.

NODE=M146

NODE=M146

 $K_2(1820)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1816±13	¹ ASTON	93 LASS	11 $K^- p \rightarrow K^- \omega p$
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 1840	² DAUM	81C CNTR	63 $K^- p \rightarrow K^- 2\pi p$
¹ From a partial wave analysis of the $K^- \omega$ system.			
² From a partial wave analysis of the $K^- 2\pi$ system.			

NODE=M146M

NODE=M146M

NODE=M146M;LINKAGE=A
 NODE=M146M;LINKAGE=C

 $K_2(1820)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
276±35	³ ASTON	93 LASS	11 $K^- p \rightarrow K^- \omega p$
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 230	⁴ DAUM	81C CNTR	63 $K^- p \rightarrow K^- 2\pi p$
³ From a partial wave analysis of the $K^- \omega$ system.			
⁴ From a partial wave analysis of the $K^- 2\pi$ system.			

NODE=M146W

NODE=M146W

NODE=M146W;LINKAGE=B
 NODE=M146W;LINKAGE=C

 $K_2(1820)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi\pi$	
Γ_2 $K_2^*(1430)\pi$	seen
Γ_3 $K^*(892)\pi$	seen
Γ_4 $K f_2(1270)$	seen
Γ_5 $K\omega$	seen

NODE=M146215;NODE=M146

DESIG=5

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

 $K_2(1820)$ BRANCHING RATIOS

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$				Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 0.77	DAUM	81C CNTR	63 $K^- p \rightarrow \bar{K} 2\pi p$	
$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$				Γ_3/Γ_1
VALUE	DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 0.05	DAUM	81C CNTR	63 $K^- p \rightarrow \bar{K} 2\pi p$	
$\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$				Γ_4/Γ_1
VALUE	DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 0.18	DAUM	81C CNTR	63 $K^- p \rightarrow \bar{K} 2\pi p$	

NODE=M146220

NODE=M146R1
 NODE=M146R1

NODE=M146R2
 NODE=M146R2

NODE=M146R3
 NODE=M146R3

 $K_2(1820)$ REFERENCES

PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ASTON	93	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

NODE=M146

REFID=49653
 REFID=43597
 REFID=22548

K(1830)

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of $K^- \phi$ system. Needs confirmation.

NODE=M088

NODE=M088

K(1830) MASS

NODE=M088M

NODE=M088M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 1830	ARMSTRONG 83	OMEG	-	18.5 $K^- p \rightarrow 3K p$

K(1830) WIDTH

NODE=M088W

NODE=M088W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 250	ARMSTRONG 83	OMEG	-	18.5 $K^- p \rightarrow 3K p$

K(1830) DECAY MODES

NODE=M088215;NODE=M088

Mode

 Γ_1 $K \phi$

DESIG=1

K(1830) REFERENCES

NODE=M088

REFID=22801

NODE=M134

ARMSTRONG 83 NP B221 1 T.A. Armstrong *et al.* (BARI, BIRM, CERN+) JP**K₀^{*}(1950)**

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+$ system. Needs confirmation.

NODE=M134

K₀^{*}(1950) MASS

NODE=M134M

NODE=M134M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1945 ± 10 ± 20	¹ ASTON 88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
••• We do not use the following data for averages, fits, limits, etc. •••				
1917 ± 12	² ZHOU 06	RVUE		$K p \rightarrow K^- \pi^+ n$
1820 ± 40	³ ANISOVICH 97C	RVUE		11 $K^- p \rightarrow K^- \pi^+ n$

¹We take the central value of the two solutions and the larger error given.²S-matrix pole. Using ASTON 88 and assuming $K_0^*(800)$, $K_0^*(1430)$.³T-matrix pole. Reanalysis of ASTON 88 data.

NODE=M134M;LINKAGE=A

NODE=M134M;LINKAGE=ZU

NODE=M134M;LINKAGE=A1

K₀^{*}(1950) WIDTH

NODE=M134W

NODE=M134W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
201 ± 34 ± 79	⁴ ASTON 88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
••• We do not use the following data for averages, fits, limits, etc. •••				
145 ± 38	⁵ ZHOU 06	RVUE		$K p \rightarrow K^- \pi^+ n$
250 ± 100	⁶ ANISOVICH 97C	RVUE		11 $K^- p \rightarrow K^- \pi^+ n$

⁴We take the central value of the two solutions and the larger error given.⁵S-matrix pole. Using ASTON 88 and assuming $K_0^*(800)$, $K_0^*(1430)$.⁶T-matrix pole. Reanalysis of ASTON 88 data.

NODE=M134W;LINKAGE=A

NODE=M134W;LINKAGE=ZU

NODE=M134W;LINKAGE=A1

K₀^{*}(1950) DECAY MODES

NODE=M134215;NODE=M134

Mode

Fraction (Γ_i/Γ) Γ_1 $K \pi$

(52 ± 14) %

DESIG=1

$K_0^*(1950)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
$0.52 \pm 0.08 \pm 0.12$	⁷ ASTON	88	LASS	0	11	$K^- p \rightarrow K^- \pi^+ n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
~ 0.60	⁸ ZHOU	06	RVUE			$K p \rightarrow K^- \pi^+ n$
⁷ We take the central value of the two solutions and the larger error given.						
⁸ S-matrix pole. Using ASTON 88 and assuming $K_0^*(800)$, $K_0^*(1430)$.						

NODE=M134220

NODE=M134R1
NODE=M134R1NODE=M134R1;LINKAGE=A
NODE=M134R1;LINKAGE=ZU **$K_0^*(1950)$ REFERENCES**

ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev
ASTON	88	NP B296 493	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)

NODE=M134

REFID=51198
REFID=45815
REFID=40262

NODE=M104

$K_2^*(1980)$

$$I(J^P) = \frac{1}{2}(2^+)$$

OMITTED FROM SUMMARY TABLE
Needs confirmation.

NODE=M104

 $K_2^*(1980)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
$1973 \pm 8 \pm 25$		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
2020 \pm 20		TIKHOMIROV	03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1978 \pm 40	241 \pm 47	BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

NODE=M104M

NODE=M104M

 $K_2^*(1980)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
$373 \pm 33 \pm 60$		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
180 \pm 70		TIKHOMIROV	03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
398 \pm 47	241 \pm 47	BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

NODE=M104W

NODE=M104W

 $K_2^*(1980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	possibly seen
Γ_2 $K\rho$	possibly seen
Γ_3 $K f_2(1270)$	possibly seen

NODE=M104215;NODE=M104

DESIG=2

DESIG=3

DESIG=4

 $K_2^*(1980)$ BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
possibly seen	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$		
$\Gamma(K\rho)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
possibly seen	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$		
$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$						Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
$1.49 \pm 0.24 \pm 0.09$	ASTON	87	LASS	0	11	$K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

NODE=M104220

NODE=M104R01
NODE=M104R01NODE=M104R02
NODE=M104R02NODE=M104R1
NODE=M104R1

$\Gamma(K f_2(1270))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$

NODE=M104R3
 NODE=M104R3

 $K_2^*(1980)$ REFERENCES

GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

NODE=M104
 REFID=53668
 REFID=49423
 REFID=41002
 REFID=40234

NODE=M035

 $K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

 $K_4^*(2045)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2045 ± 9 OUR AVERAGE		Error includes scale factor of 1.1.			
2062 ± 14 ± 13		¹ ASTON	86	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
2039 ± 10	400	^{2,3} CLELAND	82	SPEC ±	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
2070 ⁺¹⁰⁰ ₋₄₀		⁴ ASTON	81C	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2079 ± 7	431	TORRES	86	MPSF	400 $pA \rightarrow 4KX$
2088 ± 20	650	BAUBILLIER	82	HBC -	8.25 $K^- p \rightarrow K_S^0 \pi^- p$
2115 ± 46	488	CARMONY	77	HBC 0	9 $K^+ d \rightarrow K^+ \pi^+ s X$

NODE=M035M

NODE=M035M

¹ From a fit to all moments.² From a fit to 8 moments.³ Number of events evaluated by us.⁴ From energy-independent partial-wave analysis.

NODE=M035M;LINKAGE=E
 NODE=M035M;LINKAGE=B
 NODE=M035M;LINKAGE=W
 NODE=M035M;LINKAGE=D

 $K_4^*(2045)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
198 ± 30 OUR AVERAGE					
221 ± 48 ± 27		⁵ ASTON	86	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
189 ± 35	400	^{6,7} CLELAND	82	SPEC ±	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
61 ± 58	431	TORRES	86	MPSF	400 $pA \rightarrow 4KX$
170 ⁺¹⁰⁰ ₋₅₀	650	BAUBILLIER	82	HBC -	8.25 $K^- p \rightarrow K_S^0 \pi^- p$
240 ⁺⁵⁰⁰ ₋₁₀₀		⁸ ASTON	81C	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
300 ± 200		CARMONY	77	HBC 0	9 $K^+ d \rightarrow K^+ \pi^+ s X$

NODE=M035W

NODE=M035W

⁵ From a fit to all moments.⁶ From a fit to 8 moments.⁷ Number of events evaluated by us.⁸ From energy-independent partial-wave analysis.

NODE=M035W;LINKAGE=E
 NODE=M035W;LINKAGE=B
 NODE=M035W;LINKAGE=W
 NODE=M035W;LINKAGE=D

 $K_4^*(2045)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi$	(9.9 ± 1.2) %
Γ_2 $K^*(892) \pi \pi$	(9 ± 5) %
Γ_3 $K^*(892) \pi \pi \pi$	(7 ± 5) %
Γ_4 $\rho K \pi$	(5.7 ± 3.2) %
Γ_5 $\omega K \pi$	(5.0 ± 3.0) %
Γ_6 $\phi K \pi$	(2.8 ± 1.4) %
Γ_7 $\phi K^*(892)$	(1.4 ± 0.7) %

NODE=M035215;NODE=M035

DESIG=1

DESIG=2

DESIG=5

DESIG=3

DESIG=4

DESIG=6

DESIG=7

$K_4^*(2045)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{total}$						Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
0.099±0.012	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	NODE=M035220 NODE=M035R1 NODE=M035R1
$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$						Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
0.89±0.53	BAUBILLIER	82	HBC	-	8.25 $K^- p \rightarrow p K_S^0 3\pi$	NODE=M035R2 NODE=M035R2
$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$						Γ_3/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
0.75±0.49	BAUBILLIER	82	HBC	-	8.25 $K^- p \rightarrow p K_S^0 3\pi$	NODE=M035R5 NODE=M035R5
$\Gamma(\rho K\pi)/\Gamma(K\pi)$						Γ_4/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
0.58±0.32	BAUBILLIER	82	HBC	-	8.25 $K^- p \rightarrow p K_S^0 3\pi$	NODE=M035R3 NODE=M035R3
$\Gamma(\omega K\pi)/\Gamma(K\pi)$						Γ_5/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
0.50±0.30	BAUBILLIER	82	HBC	-	8.25 $K^- p \rightarrow p K_S^0 3\pi$	NODE=M035R4 NODE=M035R4
$\Gamma(\phi K\pi)/\Gamma_{total}$						Γ_6/Γ
VALUE	DOCUMENT ID	TECN	COMMENT			
0.028±0.014	⁹ TORRES	86	MPSF	400 $pA \rightarrow 4KX$		NODE=M035R6 NODE=M035R6
$\Gamma(\phi K^*(892))/\Gamma_{total}$						Γ_7/Γ
VALUE	DOCUMENT ID	TECN	COMMENT			
0.014±0.007	⁹ TORRES	86	MPSF	400 $pA \rightarrow 4KX$		NODE=M035R7 NODE=M035R7

⁹ Error determination is model dependent.

NODE=M035R;LINKAGE=A

$K_4^*(2045)$ REFERENCES

ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=22462
TORRES	86	PR D34 707	S. Torres <i>et al.</i>	(VPI, ARIZ, FNAL, FSU+)	REFID=22845
BAUBILLIER	82	PL 118B 447	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22842
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22821
CARMONY	77	PR D16 1251	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)	REFID=22811

NODE=M040

$K_2(2250)$

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems reported in the 2150–2260 MeV region, as well as enhancements seen in the antihyperon-nucleon system, either in the mass spectra or in the $J^P = 2^-$ wave.

NODE=M040

$K_2(2250)$ MASS

NODE=M040M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
2247±17 OUR AVERAGE						
2200±40		¹ ARMSTRONG	83c	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
2235±50		¹ BAUBILLIER	81	HBC	-	8 $K^- p \rightarrow \Lambda \bar{p} X$
2260±20		¹ CLELAND	81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
••• We do not use the following data for averages, fits, limits, etc. •••						
2280±20		TIKHOMIROV	03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
2147±4	37	CHLIAPNIK...	79	HBC	+	32 $K^+ p \rightarrow \bar{\Lambda} p X$
2240±20	20	LISSAUER	70	HBC		9 $K^+ p$

NODE=M040M

$1 J^P = 2^-$ from moments analysis.

NODE=M040M;LINKAGE=Q

 $K_2(2250)$ WIDTH

NODE=M040W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
180±30 OUR AVERAGE		Error includes scale factor of 1.4.			
150±30		² ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
210±30		² CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
180±60		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
~ 200		² BAUBILLIER 81	HBC	-	8 $K^- p \rightarrow \Lambda \bar{p} X$
~ 40	37	CHLIAPNIK... 79	HBC	+	32 $K^+ p \rightarrow \bar{\Lambda} p X$
80±20	20	LISSAUER 70	HBC		9 $K^+ p$
$2 J^P = 2^-$ from moments analysis.					

NODE=M040W

NODE=M040W;LINKAGE=Q

 $K_2(2250)$ DECAY MODES

NODE=M040215;NODE=M040

Mode	
Γ_1	$K \pi \pi$
Γ_2	$K f_2(1270)$
Γ_3	$K^*(892) f_0(980)$
Γ_4	$p \bar{\Lambda}$

DESIG=1

DESIG=3

DESIG=4

DESIG=2

 $K_2(2250)$ REFERENCES

NODE=M040

TIKHOMIROV 03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
	Translated from YAF 66 860.		
ARMSTRONG 83C	NP B227 365	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
BAUBILLIER 81	NP B183 1	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+) JP
CLELAND 81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+) JP
CHLIAPNIK... 79	NP B158 253	P.V. Chliapnikov <i>et al.</i>	(CERN, BELG, MONS)
LISSAUER 70	NP B18 491	D. Lissauer <i>et al.</i>	(LBL)

REFID=49423

REFID=22852

REFID=22850

REFID=22851

REFID=22849

REFID=22847

NODE=M090

 $K_3(2320)$

$$I(J^P) = \frac{1}{2}(3^+)$$

OMITTED FROM SUMMARY TABLE

Seen in the $J^P = 3^+$ wave of the antihyperon-nucleon system.

Needs confirmation.

NODE=M090

 $K_3(2320)$ MASS

NODE=M090M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2324±24 OUR AVERAGE				
2330±40	¹ ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
2320±30	¹ CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
$1 J^P = 3^+$ from moments analysis.				

NODE=M090M

NODE=M090M;LINKAGE=P

 $K_3(2320)$ WIDTH

NODE=M090W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
150±30	² ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 250	² CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
$2 J^P = 3^+$ from moments analysis.				

NODE=M090W

NODE=M090W;LINKAGE=P

 $K_3(2320)$ DECAY MODES

NODE=M090215;NODE=M090

Mode	
Γ_1	$p \bar{\Lambda}$

DESIG=1

 $K_3(2320)$ REFERENCES

NODE=M090

ARMSTRONG 83C	NP B227 365	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
CLELAND 81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+)

REFID=22852

REFID=22851

$K_5^*(2380)$

$$I(J^P) = \frac{1}{2}(5^-)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M098

NODE=M098

 $K_5^*(2380)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2382±14±19	¹ ASTON	86	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

¹ From a fit to all the moments.

NODE=M098M

NODE=M098M

NODE=M098M;LINKAGE=E

 $K_5^*(2380)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
178±37±32	² ASTON	86	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

² From a fit to all the moments.

NODE=M098W

NODE=M098W

NODE=M098W;LINKAGE=E

 $K_5^*(2380)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(6.1±1.2) %

NODE=M098215;NODE=M098

DESIG=1

 $K_5^*(2380)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
0.061±0.012	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$	

NODE=M098220

NODE=M098R1
NODE=M098R1 **$K_5^*(2380)$ REFERENCES**

ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

NODE=M098

REFID=40262
REFID=22462

NODE=M091

 $K_4(2500)$

$$I(J^P) = \frac{1}{2}(4^-)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M091

 $K_4(2500)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2490±20	¹ CLELAND	81	SPEC	± 50 $K^+ p \rightarrow \Lambda \bar{p}$

¹ $J^P = 4^-$ from moments analysis.

NODE=M091M

NODE=M091M

NODE=M091M;LINKAGE=R

 $K_4(2500)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. ••• ~ 250	² CLELAND	81	SPEC	± 50 $K^+ p \rightarrow \Lambda \bar{p}$

² $J^P = 4^-$ from moments analysis.

NODE=M091W

NODE=M091W

NODE=M091W;LINKAGE=R

 $K_4(2500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $p\bar{\Lambda}$	

NODE=M091215;NODE=M091

DESIG=1

 $K_4(2500)$ REFERENCES

CLELAND	81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+)
---------	----	-----------	----------------------------	---------------------

NODE=M091

REFID=22851

$K(3100)$

$$I^G(J^{PC}) = ?^?(?^{??})$$

NODE=M129

OMITTED FROM SUMMARY TABLE

Narrow peak observed in several ($\Lambda\bar{p}$ + pions) and ($\bar{\Lambda}p$ + pions) states in Σ^- Be reactions by BOURQUIN 86 and in np and nA reactions by ALEEV 93. Not seen by BOEHNLEIN 91. If due to strong decays, this state has exotic quantum numbers ($B=0, Q=+1, S=-1$ for $\Lambda\bar{p}\pi^+\pi^+$ and $I \geq 3/2$ for $\Lambda\bar{p}\pi^-$). Needs confirmation.

NODE=M129

 $K(3100)$ MASS

NODE=M129205

VALUE (MeV)DOCUMENT ID **≈ 3100 OUR ESTIMATE**

NODE=M129M

→ UNCHECKED ←

3-BODY DECAYSVALUE (MeV)DOCUMENT IDTECNCOMMENT

NODE=M129M1

NODE=M129M1

3054 ± 11 OUR AVERAGE

3060 ± 7 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^+$

3056 ± 7 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^-$

OCCUR=2

3055 ± 8 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^-$

OCCUR=3

3045 ± 8 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^+$

OCCUR=4

4-BODY DECAYSVALUE (MeV)DOCUMENT IDTECNCOMMENT

NODE=M129M2

NODE=M129M2

3059 ± 11 OUR AVERAGE

3067 ± 6 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$

3060 ± 8 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

OCCUR=2

3055 ± 7 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$

OCCUR=3

3052 ± 8 ± 20

¹ ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$

OCCUR=4

• • • We do not use the following data for averages, fits, limits, etc. • • •

3105 ± 30

BOURQUIN 86 SPEC $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$

3115 ± 30

BOURQUIN 86 SPEC $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

OCCUR=2

5-BODY DECAYSVALUE (MeV)DOCUMENT IDTECNCOMMENT

NODE=M129M3

NODE=M129M3

• • • We do not use the following data for averages, fits, limits, etc. • • •

3095 ± 30

BOURQUIN 86 SPEC $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$ ¹ Supersedes ALEEV 90.

NODE=M129M;LINKAGE=A

 $K(3100)$ WIDTH

NODE=M129210

3-BODY DECAYSVALUE (MeV)DOCUMENT IDTECNCOMMENT

NODE=M129W1

NODE=M129W1

• • • We do not use the following data for averages, fits, limits, etc. • • •

42 ± 16

² ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^+$

36 ± 15

² ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^-$

OCCUR=2

50 ± 18

² ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^-$

OCCUR=3

30 ± 15

² ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^+$

OCCUR=4

4-BODY DECAYSVALUE (MeV)CL%DOCUMENT IDTECNCOMMENT

NODE=M129W2

NODE=M129W2

• • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 8

² ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$

28 ± 12

² ALEEV 93 BIS2 $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

OCCUR=2

32 ± 15

² ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$

OCCUR=3

30 ± 15

² ALEEV 93 BIS2 $K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$

OCCUR=4

<30

90

BOURQUIN 86 SPEC $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$

<80

90

BOURQUIN 86 SPEC $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

OCCUR=2

5-BODY DECAYSVALUE (MeV)CL%DOCUMENT IDTECNCOMMENT

NODE=M129W3

NODE=M129W3

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30

90

BOURQUIN 86 SPEC $K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$ ² Supersedes ALEEV 90.

NODE=M129W;LINKAGE=A

K(3100) DECAY MODES

NODE=M129215;NODE=M129

Mode		
Γ_1	$K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+$	DESIG=1
Γ_2	$K(3100)^{-} \rightarrow \Lambda \bar{p} \pi^-$	DESIG=2
Γ_3	$K(3100)^{-} \rightarrow \Lambda \bar{p} \pi^+ \pi^-$	DESIG=3
Γ_4	$K(3100)^+ \rightarrow \Lambda \bar{p} \pi^+ \pi^+$	DESIG=4
Γ_5	$K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+ \pi^+ \pi^-$	DESIG=5
Γ_6	$K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$	DESIG=6

$\Gamma(\Sigma(1385)^+ \bar{p})/\Gamma(\Lambda \bar{p} \pi^+)$					Γ_6/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.04	90	ALEEV	93	BIS2	$K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$

NODE=M129R1
NODE=M129R1**K(3100) REFERENCES**

ALEEV	93	PAN 56 1358	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)
		Translated from YAF 56 100.		
BOEHNLEIN	91	NPBPS B21 174	A. Boehnlein <i>et al.</i>	(FLOR, BNL, IND+)
ALEEV	90	ZPHY C47 533	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)
BOURQUIN	86	PL B172 113	M.H. Bourquin <i>et al.</i>	(GEVA, RAL, HEIDP+)

NODE=M129

REFID=43668
REFID=41743
REFID=42173
REFID=22928

CHARMED MESONS

(C = ±1)

$$D^+ = c\bar{d}, D^0 = c\bar{u}, \bar{D}^0 = \bar{c}u, D^- = \bar{c}d, \text{ similarly for } D^{*'}\text{'s}$$

NODE=MXXX035

NODE=MXXX035

NODE=M061

 $D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

J consistent with 1, value 0 ruled out (NGUYEN 77).

NODE=M061

 $D^*(2007)^0$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M061M

NODE=M061M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2006.85 ± 0.05 OUR FIT	Error includes scale factor of 1.1.	[2006.97 ± 0.08 MeV OUR 2015 FIT]	

NODE=M061M

NEW

••• We do not use the following data for averages, fits, limits, etc. •••

2006 ± 1.5 ¹ GOLDHABER 77 MRK1 e^+e^-

¹ From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+,$ and D^0 .

NODE=M061M;LINKAGE=G

 $m_{D^*(2007)^0} - m_{D^0}$

NODE=M061DM

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M061DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
142.016 ± 0.030 OUR FIT	Error includes scale factor of 1.5.	[142.12 ± 0.07 MeV OUR 2015 FIT]		

NODE=M061DM

NEW

142.016 ± 0.030 OUR AVERAGE Error includes scale factor of 1.5. [142.12 ± 0.07 MeV OUR 2015 AVERAGE]

NEW

142.007 ± 0.015 ± 0.014 10K ² TOMARADZE 15 CLEO $e^+e^- \rightarrow$ hadrons

142.2 ± 0.3 ± 0.2 145 ALBRECHT 95F ARG $e^+e^- \rightarrow$ hadrons

142.12 ± 0.05 ± 0.05 1176 BORTOLETTO92B CLE2 $e^+e^- \rightarrow$ hadrons

••• We do not use the following data for averages, fits, limits, etc. •••

142.2 ± 2.0 SADROZINSKI 80 CBAL $D^{*0} \rightarrow D^0 \pi^0$

142.7 ± 1.7 ³ GOLDHABER 77 MRK1 e^+e^-

² Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration . This value comes from the average of the results for two decay modes, $D^0 \rightarrow K^- \pi^+$ and $D^0 \rightarrow K^- \pi^+ \pi^- \pi^+$.

³ From simultaneous fit to $D^*(2010)^+$, $D^*(2007)^0$, D^+ , and D^0 .

NODE=M061DM;LINKAGE=A

NODE=M061DM;LINKAGE=G

NODE=M061W

NODE=M061W

NODE=M061W;LINKAGE=A

NODE=M061220;NODE=M061

NODE=M061

DESIG=1

DESIG=2

$D^*(2007)^0$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	⁴ ABACHI	88B HRS	$D^{*0} \rightarrow D^+ \pi^-$

⁴ Assuming $m_{D^{*0}} = 2007.2 \pm 2.1$ MeV/ c^2 .

$D^*(2007)^0$ DECAY MODES

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 \pi^0$	(61.9±2.9) %
Γ_2 $D^0 \gamma$	(38.1±2.9) %

CONSTRAINED FIT INFORMATION

An overall fit to a branching ratio uses 3 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 0.5$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_2 \begin{vmatrix} & -100 \\ & x_1 \end{vmatrix}$$

$D^*(2007)^0$ BRANCHING RATIOS

NODE=M061225

$\Gamma(D^0 \pi^0) / \Gamma(D^0 \gamma)$					Γ_1 / Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
1.85±0.07 OUR AVERAGE					
1.90±0.07±0.05	4.9k	ABLIKIM	15B BES3	10.6 $e^+ e^- \rightarrow$ hadrons	
1.74±0.02±0.13		AUBERT,BE	05G BABR	10.6 $e^+ e^- \rightarrow$ hadrons	

NODE=M061R3
NODE=M061R3

$\Gamma(D^0 \pi^0) / \Gamma_{\text{total}}$					Γ_1 / Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.619±0.029 OUR FIT					

NODE=M061R2
NODE=M061R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.655±0.008±0.005	3.2k	⁵ ABLIKIM	15B BES3	$e^+ e^- \rightarrow$ hadrons	
0.635±0.003±0.017	69k	⁵ AUBERT,BE	05G BABR	10.6 $e^+ e^- \rightarrow$ hadrons	
0.596±0.035±0.028	858	⁶ ALBRECHT	95F ARG	$e^+ e^- \rightarrow$ hadrons	
0.636±0.023±0.033	1097	⁶ BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons	

$\Gamma(D^0 \gamma) / \Gamma_{\text{total}}$					Γ_2 / Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.381±0.029 OUR FIT					
0.381±0.029 OUR AVERAGE					

NODE=M061R1
NODE=M061R1

0.404±0.035±0.028	456	⁶ ALBRECHT	95F ARG	$e^+ e^- \rightarrow$ hadrons	
0.364±0.023±0.033	621	⁶ BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons	
0.37 ±0.08 ±0.08		ADLER	88D MRK3	$e^+ e^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.345±0.008±0.005	1.8k	⁵ ABLIKIM	15B BES3	$e^+ e^- \rightarrow$ hadrons	
0.365±0.003±0.017	68k	⁵ AUBERT,BE	05G BABR	10.6 $e^+ e^- \rightarrow$ hadrons	
0.47 ±0.23		LOW	87 HRS	29 GeV $e^+ e^-$	
0.53 ±0.13		BARTEL	85G JADE	$e^+ e^-$, hadrons	
0.47 ±0.12		COLES	82 MRK2	$e^+ e^-$	
0.45 ±0.15		GOLDHABER	77 MRK1	$e^+ e^-$	

⁵ Derived from the ratio $\Gamma(D^0\pi^0) / \Gamma(D^0\gamma)$ assuming that the branching fractions of $D^{*0} \rightarrow D^0\pi^0$ and $D^{*0} \rightarrow D^0\gamma$ decays sum to 100%

NODE=M061R;LINKAGE=AU

⁶ The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

NODE=M061R;LINKAGE=A

$D^*(2007)^0$ REFERENCES

ABLIKIM 15B PR D91 031101	M. Ablikim <i>et al.</i>	(BES III Collab.)
TOMARADZE 15 PR D91 011102	A. Tomaradze <i>et al.</i>	(NWES)
AUBERT,BE 05G PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ALBRECHT 95F ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO 92B PRL 69 2046	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
BUTLER 92 PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)
ABACHI 88B PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)
ADLER 88D PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)
LOW 87 PL B183 232	E.H. Low <i>et al.</i>	(HRS Collab.)
BARTEL 85G PL 161B 197	W. Bartel <i>et al.</i>	(JADE Collab.)
COLES 82 PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)
SADROZINSKI 80 Madison Conf. 681	H.F.W. Sadrozinski <i>et al.</i>	(PRIN, CIT+)
GOLDHABER 77 PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)
NGUYEN 77 PRL 39 262	H.K. Nguyen <i>et al.</i>	(LBL, SLAC) J

NODE=M061

REFID=56375
REFID=57142
REFID=50942
REFID=44374
REFID=43116
REFID=43170
REFID=40584
REFID=40579
REFID=40017
REFID=22880
REFID=22866
REFID=22877
REFID=11434
REFID=11543

$D^*(2010)^\pm$

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

NODE=M062

$D^*(2010)^\pm$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M062M

NODE=M062M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

2010.26±0.05 OUR FIT

[2010.27 ± 0.05 MeV OUR 2015 FIT]

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2008 ±3	¹ GOLDHABER 77	MRK1 ±	e^+e^-
2008.6 ±1.0	² PERUZZI 77	LGW ±	e^+e^-

¹ From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+,$ and D^0 ; not independent of FELDMAN 77B mass difference below.

² PERUZZI 77 mass not independent of FELDMAN 77B mass difference below and PERUZZI 77 D^0 mass value.

NODE=M062M

NEW

NODE=M062M;LINKAGE=G

NODE=M062M;LINKAGE=P

$m_{D^*(2010)^+} - m_{D^+}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M062MD

NODE=M062MD

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

140.68±0.08 OUR FIT

[140.66 ± 0.08 MeV OUR 2015 FIT]

140.64±0.08±0.06	620	BORTOLETTO92B	CLE2	$e^+e^- \rightarrow$ hadrons
------------------	-----	---------------	------	------------------------------

NODE=M062MD

NEW

$m_{D^*(2010)^+} - m_{D^0}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M062DM

NODE=M062DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

145.4257±0.0017 OUR FIT

145.4258±0.0020 OUR AVERAGE

Error includes scale factor of 1.2.

145.4259±0.0004±0.0017	312.8k	LEES	13X	BABR $D^{*\pm} \rightarrow D^0\pi^\pm \rightarrow (K\pi, K3\pi)\pi^\pm$
145.412 ±0.002 ±0.012		ANASTASSOV 02	CLE2	$D^{*\pm} \rightarrow D^0\pi^\pm \rightarrow (K\pi)\pi^\pm$
145.54 ±0.08	611	³ ADINOLFI 99	BEAT	$D^{*\pm} \rightarrow D^0\pi^\pm$
145.45 ±0.02		³ BREITWEG 99	ZEUS	$D^{*\pm} \rightarrow D^0\pi^\pm \rightarrow (K\pi)\pi^\pm$
145.42 ±0.05		³ BREITWEG 99	ZEUS	$D^{*\pm} \rightarrow D^0\pi^\pm \rightarrow (K^-3\pi)\pi^\pm$
145.5 ±0.15	103	⁴ ADLOFF 97B	H1	$D^{*\pm} \rightarrow D^0\pi^\pm$
145.44 ±0.08	152	⁴ BREITWEG 97	ZEUS	$D^{*\pm} \rightarrow D^0\pi^\pm, D^0 \rightarrow K^-3\pi$

NODE=M062DM

OCCUR=3

OCCUR=2

145.42 ±0.11	199	⁴ BREITWEG	97	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm,$ $D^0 \rightarrow K^- \pi^\pm$	OCCUR=2
145.4 ±0.2	48	⁴ DERRICK	95	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.39 ±0.06 ±0.03		BARLAG	92B	ACCM	π^- 230 GeV	
145.5 ±0.2	115	⁴ ALEXANDER	91B	OPAL	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.30 ±0.06		⁴ DECAMP	91J	ALEP	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.40 ±0.05 ±0.10		ABACHI	88B	HRS	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.46 ±0.07 ±0.03		ALBRECHT	85F	ARG	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.5 ±0.3	28	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$	OCCUR=2
145.5 ±0.3	60	FITCH	81	SPEC	$\pi^- A$	
145.3 ±0.5	30	FELDMAN	77B	MRK1	$D^{*+} \rightarrow D^0 \pi^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

145.4256±0.0006±0.0017	138.5k	LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- \pi^+) \pi^\pm$	OCCUR=2
145.4266±0.0005±0.0019	174.3k	LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- 2\pi^+ \pi^-) \pi^\pm$	OCCUR=2
145.44 ±0.09	122	⁴ BREITWEG	97B	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm,$ $D^0 \rightarrow K^- \pi^+$	
145.8 ±1.5	16	AHLEN	83	HRS	$D^{*+} \rightarrow D^0 \pi^+$	
145.1 ±1.8	12	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$	OCCUR=3
145.1 ±0.5	14	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$	OCCUR=3
145.5 ±0.5	14	YELTON	82	MRK2	$29 e^+ e^- \rightarrow$ $K^- \pi^+$	
~ 145.5		AVERY	80	SPEC	γA	
145.2 ±0.6	2	BLIETSCHAU	79	BEBC	νp	

³Statistical errors only.

⁴Systematic error not evaluated.

NODE=M062DM;LINKAGE=AV
NODE=M062DM;LINKAGE=A

$m_{D^*(2010)^+} = m_{D^*(2007)^0}$

NODE=M062EM

VALUE (MeV) DOCUMENT ID TECN COMMENT

NODE=M062EM

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6±1.8		⁵ PERUZZI	77	LGW	$e^+ e^-$	
---------	--	----------------------	----	-----	-----------	--

⁵Not independent of FELDMAN 77B mass difference above, PERUZZI 77 D^0 mass, and GOLDHABER 77 $D^*(2007)^0$ mass.

NODE=M062EM;LINKAGE=P

$D^*(2010)^\pm$ WIDTH

NODE=M062W

VALUE (keV) CL% EVTS DOCUMENT ID TECN COMMENT

NODE=M062W

83.4±1.8 OUR AVERAGE

83.3±1.2± 1.4	312.8k	⁶ LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K \pi, K 3\pi) \pi^\pm$	OCCUR=3
96 ±4 ±22		⁶ ANASTASSOV	02	CLE2	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K \pi) \pi^\pm$	
83.4±1.7± 1.5	138.5k	⁶ LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- \pi^+) \pi^\pm$	OCCUR=2
83.2±1.5± 2.6	174.3k	⁶ LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- 2\pi^+ \pi^-) \pi^\pm$	OCCUR=2
<131	90 110	BARLAG	92B	ACCM	π^- 230 GeV	

⁶Ignoring the electromagnetic contribution from $D^{*\pm} \rightarrow D^\pm \gamma$.

NODE=M062W;LINKAGE=LE

$D^*(2010)^\pm$ DECAY MODES

NODE=M062225;NODE=M062

$D^*(2010)^-$ modes are charge conjugates of the modes below.

NODE=M062

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 \pi^+$	(67.7±0.5) %
Γ_2 $D^+ \pi^0$	(30.7±0.5) %
Γ_3 $D^+ \gamma$	(1.6±0.4) %

DESIG=1
DESIG=3
DESIG=2

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 6 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 0.3$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-62	
x_3	-43	-44
	x_1	x_2

$D^*(2010)^+$ BRANCHING RATIOS

NODE=M062230

$\Gamma(D^0 \pi^+) / \Gamma_{\text{total}}$ Γ_1 / Γ

NODE=M062R1
NODE=M062R1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.677 ± 0.005 OUR FIT**0.677 ± 0.006 OUR AVERAGE**

0.6759 ± 0.0029 ± 0.0064	7,8,9	BARTELT	98 CLE2 $e^+ e^-$
0.688 ± 0.024 ± 0.013		ALBRECHT	95F ARG $e^+ e^- \rightarrow$ hadrons
0.681 ± 0.010 ± 0.013	7	BUTLER	92 CLE2 $e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.57 ± 0.04 ± 0.04		ADLER	88D MRK3 $e^+ e^-$
0.44 ± 0.10		COLES	82 MRK2 $e^+ e^-$
0.6 ± 0.15	9	GOLDHABER	77 MRK1 $e^+ e^-$

$\Gamma(D^+ \pi^0) / \Gamma_{\text{total}}$ Γ_2 / Γ

NODE=M062R3
NODE=M062R3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.307 ± 0.005 OUR FIT**0.3073 ± 0.0013 ± 0.0062**

0.312 ± 0.011 ± 0.008	1404	ALBRECHT	95F ARG $e^+ e^- \rightarrow$ hadrons
0.308 ± 0.004 ± 0.008	410	7 BUTLER	92 CLE2 $e^+ e^- \rightarrow$ hadrons
0.26 ± 0.02 ± 0.02		ADLER	88D MRK3 $e^+ e^-$
0.34 ± 0.07		COLES	82 MRK2 $e^+ e^-$

$\Gamma(D^+ \gamma) / \Gamma_{\text{total}}$ Γ_3 / Γ

NODE=M062R2
NODE=M062R2

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

0.016 ± 0.004 OUR FIT**0.016 ± 0.005 OUR AVERAGE**

0.0168 ± 0.0042 ± 0.0029		7,8	BARTELT	98 CLE2 $e^+ e^-$	
0.011 ± 0.014 ± 0.016	12	7	BUTLER	92 CLE2 $e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.052	90		ALBRECHT	95F ARG $e^+ e^- \rightarrow$ hadrons	
0.17 ± 0.05 ± 0.05			ADLER	88D MRK3 $e^+ e^-$	
0.22 ± 0.12		10	COLES	82 MRK2 $e^+ e^-$	

⁷ The branching ratios are not independent, they have been constrained by the authors to sum to 100%.

NODE=M062R;LINKAGE=A

⁸ Systematic error includes theoretical error on the prediction of the ratio of hadronic modes.

NODE=M062R;LINKAGE=B

⁹ Assuming that isospin is conserved in the decay.

NODE=M062R;LINKAGE=G

¹⁰ Not independent of $\Gamma(D^0 \pi^+) / \Gamma_{\text{total}}$ and $\Gamma(D^+ \pi^0) / \Gamma_{\text{total}}$ measurement.

NODE=M062R;LINKAGE=C

$D^*(2010)^\pm$ REFERENCES

LEES	13X	PRL 111 111801	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55564
Also		PR D88 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55547
Also		PR D88 079902 (errat.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55695
ANASTASSOV	02	PR D65 032003	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=48550
ADINOLFI	99	NP B547 3	M. Adinolfi <i>et al.</i>	(Beatrice Collab.)	REFID=46925
BREITWEG	99	EPJ C6 67	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=46604
BARTELT	98	PRL 80 3919	J. Bartelt <i>et al.</i>	(CLEO Collab.)	REFID=46349
ADLOFF	97B	ZPHY C72 593	C. Adloff <i>et al.</i>	(H1 Collab.)	REFID=45421
BREITWEG	97	PL B401 192	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45520
BREITWEG	97B	PL B407 402	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45699
ALBRECHT	95F	ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44374
DERRICK	95	PL B349 225	M. Derrick <i>et al.</i>	(ZEUS Collab.)	REFID=44373
BARLAG	92B	PL B278 480	S. Barlag <i>et al.</i>	(ACCMOR Collab.)	REFID=42174
BORTOLETTO	92B	PRL 69 2046	D. Bortoletto <i>et al.</i>	(CLEO Collab.)	REFID=43116
BUTLER	92	PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43170
ALEXANDER	91B	PL B262 341	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=41553
DECAMP	91J	PL B266 218	D. Decamp <i>et al.</i>	(ALEPH Collab.)	REFID=41614
ABACHI	88B	PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)	REFID=40584
ADLER	88D	PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40579
ALBRECHT	85F	PL 150B 235	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=11527
AHLEN	83	PRL 51 1147	S.P. Ahlen <i>et al.</i>	(ANL, IND, LBL+)	REFID=22868
BAILEY	83	PL 132B 230	R. Bailey <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)	REFID=22870
COLES	82	PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)	REFID=22866
YELTON	82	PRL 49 430	J.M. Yelton <i>et al.</i>	(SLAC, LBL, UCB+)	REFID=22867
FITCH	81	PRL 46 761	V.L. Fitch <i>et al.</i>	(PRIN, SACL, TORI+)	REFID=22863
AVERY	80	PRL 44 1309	P. Avery <i>et al.</i>	(ILL, FNAL, COLU)	REFID=11498
BLIETSCHAU	79	PL 86B 108	J. Blietschau <i>et al.</i>	(AACH3, BONN, CERN+)	REFID=22861
FELDMAN	77B	PRL 38 1313	G.J. Feldman <i>et al.</i>	(Mark I Collab.)	REFID=22858
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)	REFID=11434
PERUZZI	77	PRL 39 1301	I. Peruzzi <i>et al.</i>	(LGW Collab.)	REFID=11435

NODE=M062

REFID=55564
REFID=55547
REFID=55695
REFID=48550
REFID=46925
REFID=46604
REFID=46349
REFID=45421
REFID=45520
REFID=45699
REFID=44374
REFID=44373
REFID=42174
REFID=43116
REFID=43170
REFID=41553
REFID=41614
REFID=40584
REFID=40579
REFID=11527
REFID=22868
REFID=22870
REFID=22866
REFID=22867
REFID=22863
REFID=11498
REFID=22861
REFID=22858
REFID=11434
REFID=11435

$$D_0^*(2400)^0$$

$$I(J^P) = \frac{1}{2}(0^+)$$

$J^P = 0^+$ assignment favored (ABE 04D).

NODE=M178


NODE=M178

$D_0^*(2400)^0$ MASS

NODE=M178M

NODE=M178M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2318 ± 29 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
2297 ± 8 ± 20	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
2308 ± 17 ± 32		ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$
2407 ± 21 ± 35	9.8k	LINK	04A FOCS	γA



$D_0^*(2400)^0$ WIDTH

NODE=M178W

NODE=M178W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
267 ± 40 OUR AVERAGE				
273 ± 12 ± 48	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
276 ± 21 ± 63		ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$
240 ± 55 ± 59	9.8k	LINK	04A FOCS	γA

$D_0^*(2400)^0$ DECAY MODES

NODE=M178215;NODE=M178

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^+\pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D_0^*(2400)^0$ REFERENCES

AUBERT	09AB PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04D PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
LINK	04A PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)

NODE=M178

REFID=52941
REFID=50011
REFID=49775

NODE=M179

 $D_0^*(2400)^\pm$

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

 J, P need confirmation.

NODE=M179

 $D_0^*(2400)^\pm$ MASS

NODE=M179M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M179M

2351 ± 7 OUR AVERAGE

NEW

[2403 ± 40 MeV OUR 2015 AVERAGE]

2360 ± 15 ± 30		¹ AAIJ	15X LHCB	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2349 ± 6 ± 4		² AAIJ	15Y LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2403 ± 14 ± 35	18.8k	LINK	04A FOCS	γA

• • • We do not use the following data for averages, fits, limits, etc. • • •

2354 ± 7 ± 11		³ AAIJ	15Y LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
---------------	--	-------------------	----------	---

OCCUR=2

¹ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S -wave and the $D\pi$ S - and P -waves.

NODE=M179M;LINKAGE=A

² Modeling the $\pi^+\pi^-$ S -wave with the Isobar formalism.

NODE=M179M;LINKAGE=B

³ Modeling the $\pi^+\pi^-$ S -wave with the K-matrix formalism.

NODE=M179M;LINKAGE=C

 $D_0^*(2400)^\pm$ WIDTH

NODE=M179W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M179W

230 ± 17 OUR AVERAGE Error includes scale factor of 1.1. [283 ± 40 MeV OUR 2015 AVERAGE]

NEW

255 ± 26 ± 51		¹ AAIJ	15X LHCB	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
217 ± 13 ± 13		² AAIJ	15Y LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
283 ± 24 ± 34	18.8k	LINK	04A FOCS	γA

• • • We do not use the following data for averages, fits, limits, etc. • • •

230 ± 15 ± 21		³ AAIJ	15Y LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
---------------	--	-------------------	----------	---

OCCUR=2

¹ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S -wave and the $D\pi$ S - and P -waves.

NODE=M179W;LINKAGE=A

² Modeling the $\pi^+\pi^-$ S -wave with the Isobar formalism.

NODE=M179W;LINKAGE=B

³ Modeling the $\pi^+\pi^-$ S -wave with the K-matrix formalism.

NODE=M179W;LINKAGE=C

 $D_0^*(2400)^\pm$ DECAY MODES

NODE=M179215;NODE=M179

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0\pi^+$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D_0^*(2400)^\pm$ REFERENCES

NODE=M179

AAIJ	15X PR D92 012012	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15Y PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.)
LINK	04A PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)

REFID=56588
REFID=56609
REFID=49775

NODE=M097

D₁(2420)⁰

$I(J^P) = \frac{1}{2}(1^+)$
I needs confirmation.

D₁(2420)⁰ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0,$
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M097M

NODE=M097M

NODE=M097M

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
2420.8±0.5 OUR FIT Error includes scale factor of 1.3. [2421.4 ± 0.6 MeV OUR 2015
FIT Scale factor = 1.2]
2420.5±0.6 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.
[2421.1 ± 0.7 MeV OUR 2015 AVERAGE Scale factor = 1.2]

NEW

NEW

2419.6±0.1±0.7	210k	AAIJ	13CC	LHCB	$pp \rightarrow D^{*+} \pi^- X$
2423.1±1.5 ^{+0.4} _{-1.0}	2.7k	¹ ABRAMOWICZ13	ZEUS		$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2420.1±0.1±0.8	103k	DEL-AMO-SA..10P	BABR		$e^+ e^- \rightarrow D^{*+} \pi^- X$
2426 ±3 ±1	151	ABE	05A	BELL	$B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
2421.4±1.5±0.9		² ABE	04D	BELL	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2421 ⁺¹ ₋₂ ±2	286	AVERY	94C	CLE2	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2422 ±2 ±2	51	FRABETTI	94B	E687	$\gamma Be \rightarrow D^{*+} \pi^- X$
2428 ±3 ±2	279	AVERY	90	CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2414 ±2 ±5	171	ALBRECHT	89H	ARG	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2428 ±8 ±5	171	ANJOS	89C	TPS	$\gamma N \rightarrow D^{*+} \pi^- X$

••• We do not use the following data for averages, fits, limits, etc. •••

2420.5±2.1±0.9	3110 ± 340	³ CHEKANOV	09	ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
2421.7±0.7±0.6	7.5k	ABULENCIA	06A	CDF	1900 $p\bar{p} \rightarrow D^{*+} \pi^- X$
2425 ±3	235	⁴ ABREU	98M	DLPH	$e^+ e^-$

NODE=M097M;LINKAGE=AR

NODE=M097M;LINKAGE=AB

NODE=M097M;LINKAGE=CH


¹ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

² Fit includes the contribution from $D_1^*(2430)^0$.

³ Calculated using the mass difference $m(D_1^0) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of 0.9 MeV.

⁴ No systematic error given.

NODE=M097M;LINKAGE=K



$$m_{D_1^0} - m_{D^{*+}}$$

NODE=M097DM

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0,$
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M097DM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
410.6±0.5 OUR FIT				Error includes scale factor of 1.3. [411.1 ± 0.6 OUR 2015 FIT Scale factor = 1.2]
411.5±0.8 OUR AVERAGE				
410.2±2.1±0.9	3110 ± 340	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
411.7±0.7±0.4	7.5k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^{*+} \pi^- X$

NODE=M097DM
NEW

$D_1(2420)^0$ WIDTH

NODE=M097W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7± 2.5 OUR AVERAGE				Error includes scale factor of 3.5. See the ideogram below. [27.4 ± 2.5 MeV OUR 2015 AVERAGE Scale factor = 2.3]
35.2± 0.4± 0.9	210k	AAIJ	13CC LHCb	$pp \rightarrow D^{*+} \pi^- X$
38.8± 5.0 ^{+1.9} _{-5.4}	2.7k	¹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
31.4± 0.5± 1.3	103k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$
20.0± 1.7± 1.3	7.5k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^{*+} \pi^- X$
24 ± 7 ± 8	151	ABE	05A BELL	$B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
23.7± 2.7± 4.0		² ABE	04D BELL	$B^- \rightarrow D^{*+} \pi^- \pi^-$
20 ⁺⁶ ₋₅ ± 3	286	AVERY	94C CLE2	$e^+ e^- \rightarrow D^{*+} \pi^- X$
15 ± 8 ± 4	51	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} \pi^- X$
23 ⁺⁸ ₋₆ ⁺¹⁰ ₋₃	279	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
13 ± 6 ⁺¹⁰ ₋₅	171	ALBRECHT	89H ARG	$e^+ e^- \rightarrow D^{*+} \pi^- X$

NODE=M097W
NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •


53.2± 7.2 ^{+3.3} _{-4.9}	3110 ± 340	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
58 ± 14 ± 10	171	ANJOS	89C TPS	$\gamma N \rightarrow D^{*+} \pi^- X$

¹ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

² Fit includes the contribution from $D_1^*(2430)^0$.

NODE=M097W;LINKAGE=AR

NODE=M097W;LINKAGE=AB



$D_1(2420)^0$ DECAY MODES

$\bar{D}_1(2420)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2010)^+ \pi^-$	seen
Γ_2 $D^0 \pi^+ \pi^-$	seen
Γ_3 $D^0 \rho^0$	
Γ_4 $D^0 f_0(500)$	
Γ_5 $D_0^*(2400)^+ \pi^-$	
Γ_6 $D^+ \pi^-$	not seen
Γ_7 $D^{*0} \pi^+ \pi^-$	not seen

NODE=M097215;NODE=M097

NODE=M097

DESIG=1

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

DESIG=5

DESIG=6

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

 $D_1(2420)^0$ BRANCHING RATIOS

$\Gamma(D^*(2010)^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	ACKERSTAFF	97W	OPAL $e^+ e^- \rightarrow D^{*+} \pi^- X$
seen	AVERY	90	CLEO $e^+ e^- \rightarrow D^{*+} \pi^- X$
seen	ALBRECHT	89H	ARG $e^+ e^- \rightarrow D^* \pi^- X$
seen	ANJOS	89C	TPS $\gamma N \rightarrow D^{*+} \pi^- X$

NODE=M097220

NODE=M097R1

NODE=M097R1

$\Gamma(D^+ \pi^-)/\Gamma(D^*(2010)^+ \pi^-)$	Γ_6/Γ_1			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	AVERY	90	CLEO $e^+ e^- \rightarrow D^+ \pi^- X$

NODE=M097R2

NODE=M097R2

 $D_1(2420)^0$ POLARIZATION AMPLITUDE A_{D_1}

A polarization amplitude A_{D_1} is a parameter that depends on the initial polarization of the D_1 and is sensitive to a possible S -wave contribution to its decay. For D_1 decays the helicity angle, θ_h , distribution varies like $1 + A_{D_1} \cos^2 \theta_h$, where θ_h is the angle in the D^* rest frame between the two pions emitted by the $D_1 \rightarrow D^* \pi$ and the $D^* \rightarrow D \pi$.

Unpolarized D_1 decaying purely via D -wave is predicted to give $A_{D_1} = 3$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
5.73 ± 0.25 OUR AVERAGE				
7.8 $^{+6.7}_{-2.7}$ $^{+4.6}_{-1.8}$	2.7k	1 ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
5.72 ± 0.25	103k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$
5.9 $^{+3.0}_{-1.7}$ $^{+2.4}_{-1.0}$		CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.30 ± 0.48	210k	2 AAJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
$3.8 \pm 0.6 \pm 0.8$		3 AUBERT	09Y BABR	$B^+ \rightarrow D_1^0 \ell^+ \nu_\ell$
$2.74^{+1.40}_{-0.93}$		4 AVERY	94C CLE2	$e^+ e^- \rightarrow D^{*+} \pi^- X$

NODE=M097PAH

NODE=M097PAH

NODE=M097PAH

¹ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 . A pure D -wave not excluded although some S -wave mixing possible.

² Systematic uncertainty not estimated. Resonance parameters fixed.

³ Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+ B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths and helicity angle distributions for charged and neutral D_1 mesons.

⁴ Systematic uncertainties not estimated.

NODE=M097PAH;LINKAGE=AR

NODE=M097PAH;LINKAGE=A

NODE=M097PAH;LINKAGE=AU

NODE=M097PAH;LINKAGE=AV

 $D_1(2420)^0$ REFERENCES

AAJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABRAMOWICZ	13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
DEL-AMO-SA...	10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABULENCIA	06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE	05A	PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)

NODE=M097

REFID=55581

REFID=54743

REFID=53534

REFID=52929

REFID=52733

REFID=51054

REFID=50755

REFID=50011

REFID=46315

REFID=45788

REFID=44096

REFID=43687

REFID=41013

REFID=41001

REFID=40737

$D_1(2420)^\pm$

$I(J^P) = \frac{1}{2}(?^?)$
I needs confirmation.

OMITTED FROM SUMMARY TABLE
 Seen in $D^*(2007)^0 \pi^+$. $J^P = 0^+$ ruled out.

NODE=M120

NODE=M120

NODE=M120M


NODE=M120M

$D_1(2420)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2423.2±2.4 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
2421.9±4.7 ^{+3.4} _{-1.2}	759	¹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)0} \pi^+ X$
2421 ±2 ±1	124	ABE	05A BELL	$\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
2425 ±2 ±2	146	BERGFELD	94B CLE2	$e^+ e^- \rightarrow D^{*0} \pi^+ X$
2443 ±7 ±5	190	ANJOS	89C TPS	$\gamma N \rightarrow D^0 \pi^+ X^0$

¹ From the fit of the $M(D^0 \pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1, respectively.

NODE=M120M;LINKAGE=AB



$m_{D_1^*(2420)^\pm} - m_{D_1^*(2420)^0}$

NODE=M120DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4^{+2}_{-3} \pm 3$	BERGFELD	94B CLE2	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M120DM

$D_1(2420)^\pm$ WIDTH

NODE=M120W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
25± 6 OUR AVERAGE				
21± 5±8	124	ABE	05A BELL	$\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
26 ⁺⁸ ₋₇ ±4	146	BERGFELD	94B CLE2	$e^+ e^- \rightarrow D^{*0} \pi^+ X$
41±19±8	190	ANJOS	89C TPS	$\gamma N \rightarrow D^0 \pi^+ X^0$

NODE=M120W

$D_1(2420)^\pm$ DECAY MODES $D_1^*(2420)^-$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2007)^0 \pi^+$	seen
Γ_2 $D^+ \pi^+ \pi^-$	seen
Γ_3 $D^+ \rho^0$	
Γ_4 $D^+ f_0(500)$	
Γ_5 $D_0^*(2400)^0 \pi^+$	
Γ_6 $D^0 \pi^+$	not seen
Γ_7 $D^{*+} \pi^+ \pi^-$	not seen

NODE=M120215;NODE=M120

NODE=M120

DESIG=1

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

DESIG=5

DESIG=6

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

 $D_1(2420)^\pm$ BRANCHING RATIOS

$\Gamma(D^*(2007)^0 \pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	ANJOS	89C	TPS $\gamma N \rightarrow D^0 \pi^+ \chi^0$

NODE=M120220

NODE=M120R1

NODE=M120R1

$\Gamma(D^0 \pi^+)/\Gamma(D^*(2007)^0 \pi^+)$	Γ_6/Γ_1			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.18	90	BERGFELD	94B	CLE2 $e^+ e^- \rightarrow \text{hadrons}$

NODE=M120R2

NODE=M120R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $D_1(2420)^\pm$ POLARIZATION AMPLITUDE A_{D_1}

A polarization amplitude A_{D_1} is a parameter that depends on the initial polarization of the D_1 and is sensitive to a possible S -wave contribution to its decay. For D_1 decays the helicity angle, θ_h , distribution varies like $1 + A_{D_1} \cos^2 \theta_h$, where θ_h is the angle in the D^* rest frame between the two pions emitted by the $D_1 \rightarrow D^* \pi$ and the $D^* \rightarrow D \pi$.

Unpolarized D_1 decaying purely via D -wave is predicted to give $A_{D_1} = 3$.

VALUE	DOCUMENT ID	TECN	COMMENT
$3.8 \pm 0.6 \pm 0.8$	² AUBERT	09Y	BABR $B^0 \rightarrow D_1^- \ell^+ \nu_\ell$

NODE=M120PAH

NODE=M120PAH

• • • We do not use the following data for averages, fits, limits, etc. • • •

² Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+ B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths and helicity angle distributions for charged and neutral D_1 mesons.

NODE=M120PAH

NODE=M120PAH;LINKAGE=AU

 $D_1(2420)^\pm$ REFERENCES

ABRAMOWICZ 13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
AUBERT	09Y PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	05A PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)
BERGFELD	94B PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
ANJOS	89C PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)

NODE=M120

REFID=54743

REFID=52929

REFID=50755

REFID=44099

REFID=40737

$D_1(2430)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

OMITTED FROM SUMMARY TABLE
 $J = 1^+$ assignment favored (ABE 04D).

 $D_1(2430)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2427 ± 26 ± 25	ABE	04D	BELL $B^- \rightarrow D^{*+} \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2477 ± 28	¹ AUBERT	06L	BABR $\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$
¹ Systematic errors not estimated.			

 $D_1(2430)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
384⁺¹⁰⁷₋₇₅ ± 74	ABE	04D	BELL $B^- \rightarrow D^{*+} \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
266 ± 97	² AUBERT	06L	BABR $\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$
² Systematic errors not estimated.			

 $D_1(2430)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2010)^+ \pi^-$	seen

 $D_1(2430)^0$ REFERENCES

AUBERT	06L	PR D74 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)

 $D_2^*(2460)^0$

$$I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored (ALBRECHT 89B, ALBRECHT 89H), natural parity confirmed by the helicity analysis (DEL-AMO-SANCHEZ 10P). AAIJ 13CC confirms $J^P = 2^+$ and natural parity.

 $D_2^*(2460)^0$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2460.57 ± 0.15 OUR FIT	Error includes scale factor of 1.1. [2462.6 ± 0.6 MeV OUR 2015 FIT Scale factor = 1.2]			
2460.47 ± 0.21 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below. [2461.8 ± 0.7 MeV OUR 2015 AVERAGE Scale factor = 1.1]			
2460.4 ± 0.4 ± 1.2	82k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
2460.4 ± 0.1 ± 0.1	675k	AAIJ	13CC LHCB	$pp \rightarrow D^+ \pi^- X$
2462.5 ± 2.4 ^{+1.3} _{-1.1}	2.3k	¹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2462.2 ± 0.1 ± 0.8	243k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$
2460.4 ± 1.2 ± 2.2	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
2461.6 ± 2.1 ± 3.3		² ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$
2464.5 ± 1.1 ± 1.9	5.8k	² LINK	04A FOCS	γA
2465 ± 3 ± 3	486	AVERY	94C CLE2	$e^+ e^- \rightarrow D^+ \pi^- X$
2453 ± 3 ± 2	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+ \pi^- X$
2461 ± 3 ± 1	440	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2455 ± 3 ± 5	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$
2459 ± 3 ± 2	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$

NODE=M180

NODE=M180

NODE=M180M

NODE=M180M

NODE=M180M;LINKAGE=AU

NODE=M180W

NODE=M180W

NODE=M180W;LINKAGE=AU

NODE=M180215;NODE=M180

DESIG=1;OUR EVAL;→ UNCHECKED ←

NODE=M180

REFID=51140
REFID=50011

NODE=M119

NODE=M119

NODE=M119M

NODE=M119M

NODE=M119M

NEW

NEW

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

2469.1 ± 3.7	$^{+1.2}_{-1.3}$	1.5k	³ CHEKANOV	09	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2463.3 ± 0.6	± 0.8	20k	ABULENCIA	06A	CDF	1900 $p\bar{p} \rightarrow D^+ \pi^- X$
2461 ± 6		126	⁴ ABREU	98M	DLPH	$e^+ e^-$
2466 ± 7		1	ASRATYAN	95	BEBC	53,40 $\nu(\bar{\nu}) \rightarrow pX, dX$

¹ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

² Fit includes the contribution from $D_0^*(2400)^0$.

³ Calculated using the mass difference $m(D_2^{*0}) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of $^{+1.2}_{-1.3}$ MeV.


⁴ No systematic error given.

NODE=M119M;LINKAGE=AR

NODE=M119M;LINKAGE=LI

NODE=M119M;LINKAGE=CH

NODE=M119M;LINKAGE=K



$D_2^*(2460)^0$ mass (MeV)

$m_{D_2^{*0}} - m_{D^{*+}}$

NODE=M119DM

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0,$ and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M119DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
590.98 ± 0.18 OUR FIT	Error includes scale factor of 1.1.			[593.0 ± 0.6 MeV OUR 2015 FIT Scale factor = 1.2]
593.9 ± 0.6 ± 0.5	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+ \pi^- X$

NODE=M119DM

NEW

$m_{D_2^{*0}} - m_{D^{*++}}$

NODE=M119DM2

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0,$ and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M119DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
450.31 ± 0.16 OUR FIT	Error includes scale factor of 1.1.			[452.3 ± 0.6 MeV OUR 2015 FIT Scale factor = 1.2]
458.8 ± 3.7	$^{+1.2}_{-1.3}$ 1560 ± 230	CHEKANOV	09	ZEUS $e^\pm p \rightarrow D^{(*)+} \pi^- X$

NODE=M119DM2

NEW

$D_2^*(2460)^0$ WIDTH

NODE=M119W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
47.7 ± 1.3 OUR AVERAGE				Error includes scale factor of 2.0. See the ideogram below. [49.0 ± 1.3 MeV OUR 2015 AVERAGE Scale factor = 1.5]
43.2 ± 1.2 ± 3.0	82k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
45.6 ± 0.4 ± 1.1	675k	AAIJ	13CC LHCB	$pp \rightarrow D^+ \pi^- X$
46.6 ± 8.1 ⁺ ₋ 5.9 3.8	2.3k	⁵ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
50.5 ± 0.6 ± 0.7	243k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$
41.8 ± 2.5 ± 2.9	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
49.2 ± 2.3 ± 1.3	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+ \pi^- X$
45.6 ± 4.4 ± 6.7		⁶ ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$
38.7 ± 5.3 ± 2.9	5.8k	⁶ LINK	04A FOCS	γA
28 ⁺ ₋ 8 7 ± 6	486	AVERY	94C CLE2	$e^+ e^- \rightarrow D^+ \pi^- X$
25 ± 10 ± 5	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+ \pi^- X$
20 ⁺ ₋ 9 12 ± 9 -10	440	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
15 ⁺ ₋ 13 10 ± 5 -10	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$
20 ± 10 ± 5	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$


NODE=M119W
NEW
OCCUR=2

⁵ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

⁶ Fit includes the contribution from $D_0^{*0}(2400)^0$.

NODE=M119W;LINKAGE=AR

NODE=M119W;LINKAGE=LI



$D_2^{*0}(2460)^0$ DECAY MODES

NODE=M119215;NODE=M119

$\bar{D}_2^{*0}(2460)^0$ modes are charge conjugates of modes below.

NODE=M119

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^+ \pi^-$	seen
Γ_2 $D^{*0}(2010)^+ \pi^-$	seen
Γ_3 $D^0 \pi^+ \pi^-$	not seen
Γ_4 $D^{*0} \pi^+ \pi^-$	not seen

CLUMP=A;DESIG=1

DESIG=2

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

$D_2^{*0}(2460)^0$ BRANCHING RATIOS

NODE=M119220

$\Gamma(D^+ \pi^-)/\Gamma_{total}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$	
seen		337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$	
seen			ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$	

NODE=M119R1

NODE=M119R1

$\Gamma(D^*(2010)^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ACKERSTAFF 97W	OPAL	$e^+e^- \rightarrow D^{*+}\pi^- X$
seen	AVERY 90	CLEO	$e^+e^- \rightarrow D^{*+}\pi^- X$
seen	ALBRECHT 89H	ARG	$e^+e^- \rightarrow D^*\pi^- X$

NODE=M119R2
NODE=M119R2 $\Gamma(D^+\pi^-)/\Gamma(D^*(2010)^+\pi^-)$ Γ_1/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.54±0.15 OUR AVERAGE				
1.4 ±0.3 ±0.3	2.3k	⁷ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+}\pi^- X$
1.47±0.03±0.16	379k	DEL-AMO-SA...10P	BABR	$e^+e^- \rightarrow D^{(*)+}\pi^- X$
2.8 ±0.8 $^{+0.5}_{-0.6}$	1560 ± 230	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{(*)+}\pi^- X$
2.2 ±0.7 ±0.6		AVERY 94C	CLE2	$e^+e^- \rightarrow D^{*+}\pi^- X$
2.3 ±0.8		AVERY 90	CLEO	e^+e^-
3.0 ±1.1 ±1.5		ALBRECHT 89H	ARG	$e^+e^- \rightarrow D^*\pi^- X$

NODE=M119R3
NODE=M119R3

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1.9 ±0.5		ABE 04D	BELL	$B^- \rightarrow D^{(*)+}\pi^- \pi^-$
----------	--	---------	------	---------------------------------------

⁷ From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 .

NODE=M119R3;LINKAGE=AR

 $\Gamma(D^+\pi^-)/[\Gamma(D^+\pi^-) + \Gamma(D^*(2010)^+\pi^-)]$ $\Gamma_1/(\Gamma_1+\Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.62±0.03±0.02	8414	⁸ AUBERT 09Y	BABR	$B^+ \rightarrow D_2^{*0}\ell^+\nu_\ell$
----------------	------	-------------------------	------	--

⁸ Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths for charged and neutral D_2^* mesons.

NODE=M119R01
NODE=M119R01

NODE=M119R01;LINKAGE=AU

 $D_2^*(2460)^0$ POLARIZATION AMPLITUDE A_{D_2}

NODE=M119PAM

A polarization amplitude A_{D_2} is a parameter that depends on the initial polarization of the D_2 . For D_2 decays the helicity angle, θ_H , distribution varies like $1 + A_{D_2} \cos^2(\theta_H)$, where θ_H is the angle in the D^* rest frame between the two pions emitted by the $D_2 \rightarrow D^*\pi$ and $D^* \rightarrow D\pi$.

NODE=M119PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

-1.16±0.35	2.3k	⁹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+}\pi^- X$
consistent with -1	243k	DEL-AMO-SA...10P	BABR	$e^+e^- \rightarrow D^+\pi^- X$
-0.74 $^{+0.49}_{-0.38}$		¹⁰ AVERY 94C	CLE2	$e^+e^- \rightarrow D^{*+}\pi^- X$

⁹ From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions.¹⁰ Systematic uncertainties not estimated.

NODE=M119PAM

NODE=M119PAM;LINKAGE=AB

NODE=M119PAM;LINKAGE=AV

 $D_2^*(2460)^0$ REFERENCES

NODE=M119

AAIJ 13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
ABRAMOWICZ 13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54743
DEL-AMO-SA... 10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534
AUBERT 09AB	PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52941
AUBERT 09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52929
CHEKANOV 09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
ABULENCIA 06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)	REFID=51054
ABE 04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50011
LINK 04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)	REFID=49775
ABREU 98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46315
ACKERSTAFF 97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ASRATYAN 95	ZPHY C68 43	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)	REFID=44439
AVERY 94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)	REFID=44096
FRABETTI 94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
AVERY 90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT 89B	PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=40736
ALBRECHT 89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=41001
ANJOS 89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)	REFID=40737

$D_2^*(2460)^\pm$

$I(J^P) = \frac{1}{2}(2^+)$

$J^P = 2^+$ assignment strongly favored(ALBRECHT 89B).

NODE=M150

NODE=M150

NODE=M150M

NODE=M150M

NEW

$D_2^*(2460)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2465.4±1.3 OUR AVERAGE		Error includes scale factor of 3.1. See the ideogram below. [2464.3 ± 1.6 MeV OUR 2015 AVERAGE Scale factor = 1.7]		
2465.6±1.8±1.3		1 AAIJ	15X LHCB	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2468.6±0.6±0.3		2 AAIJ	15Y LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2463.1±0.2±0.6	342k	AAIJ	13CC LHCB	$p p \rightarrow D^0 \pi^+ X$
2460.6±4.4 ^{+3.6} _{-0.8}	1371	3 ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)0} \pi^+ X$
2465.4±0.2±1.1	111k	4 DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^0 \pi^+ X$
2465.7±1.8 ^{+1.4} _{-4.8}	2909	KUZMIN	07 BELL	$e^+ e^- \rightarrow$ hadrons
2463 ±3 ±3	310	BERGFELD	94B CLE2	$e^+ e^- \rightarrow D^0 \pi^+ X$
2453 ±3 ±2	185	FRABETTI	94B E687	$\gamma Be \rightarrow D^0 \pi^+ X$
2469 ±4 ±6		ALBRECHT	89F ARG	$e^+ e^- \rightarrow D^0 \pi^+ X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2468.1±0.6±0.5		5 AAIJ	15Y LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2467.6±1.5±0.8	3.5k	6 LINK	04A FOCS	γA

OCCUR=2

NODE=M150M;LINKAGE=A

NODE=M150M;LINKAGE=B

NODE=M150M;LINKAGE=AB

¹ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S-wave and the $D\pi$ S- and P-waves.

² Modeling the $\pi^+\pi^-$ S-wave with the Isobar formalism.

³ From the fit of the $M(D^0\pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1, respectively.

⁴ At a fixed width of 50.5 MeV.


⁵ Modeling the $\pi^+\pi^-$ S-wave with the K-matrix formalism.

⁶ Fit includes the contribution from $D_0^*(2400)^\pm$. Not independent of the corresponding mass difference measurement, $(m_{D_2^*(2460)^\pm}) - (m_{D_2^*(2460)^0})$.

NODE=M150M;LINKAGE=DE

NODE=M150M;LINKAGE=C

NODE=M150M;LINKAGE=LI



$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0}$

NODE=M150DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2.4±1.7 OUR AVERAGE			
3.1±1.9±0.9	LINK	04A	FOCS γ A
- 2 ±4 ±4	BERGFELD	94B	CLE2 $e^+e^- \rightarrow$ hadrons
0 ±4	FRABETTI	94B	E687 γ Be \rightarrow $D\pi X$
14 ±5 ±8	ALBRECHT	89F	ARG $e^+e^- \rightarrow D^0\pi^+X$

NODE=M150DM

 $D_2^*(2460)^\pm$ WIDTH

NODE=M150W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
46.7± 1.2 OUR AVERAGE				
[37 ± 6 MeV OUR 2015 AVERAGE Scale factor = 1.4]				
46.0± 3.4±3.2		¹ AAIJ	15X	LHCB $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
47.3± 1.5±0.7		² AAIJ	15Y	LHCB $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
48.6± 1.3±1.9	342k	AAIJ	13CC	LHCB $p p \rightarrow D^0 \pi^+ X$
49.7± 3.8±6.4	2909	KUZMIN	07	BELL $e^+e^- \rightarrow$ hadrons
34.1± 6.5±4.2	3.5k	³ LINK	04A	FOCS γ A
27 ⁺¹¹ ₋₈ ±5	310	BERGFELD	94B	CLE2 $e^+e^- \rightarrow D^0 \pi^+ X$
23 ± 9 ±5	185	FRABETTI	94B	E687 γ Be $\rightarrow D^0 \pi^+ X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
46.0± 1.4±1.8		⁴ AAIJ	15Y	LHCB $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
¹ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S -wave and the $D\pi$ S - and P -waves.				
² Modeling the $\pi^+\pi^-$ S -wave with the Isobar formalism.				
³ Fit includes the contribution from $D_0^*(2400)^\pm$.				
⁴ Modeling the $\pi^+\pi^-$ S -wave with the K-matrix formalism.				

NODE=M150W
NEW

OCCUR=2

NODE=M150W;LINKAGE=A

NODE=M150W;LINKAGE=B

NODE=M150W;LINKAGE=LI

NODE=M150W;LINKAGE=C

 $D_2^*(2460)^\pm$ DECAY MODES

NODE=M150215;NODE=M150

 $D_2^*(2460)^-$ modes are charge conjugates of modes below.

NODE=M150

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 \pi^+$	seen
Γ_2 $D^{*0} \pi^+$	seen
Γ_3 $D^+ \pi^+ \pi^-$	not seen
Γ_4 $D^{*+} \pi^+ \pi^-$	not seen

DESIG=1

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

 $D_2^*(2460)^\pm$ BRANCHING RATIOS

NODE=M150220

$\Gamma(D^0 \pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		ALBRECHT	89F	ARG $e^+e^- \rightarrow D^0 \pi^+ X$	

NODE=M150R1
NODE=M150R1

$\Gamma(D^0 \pi^+)/\Gamma(D^{*0} \pi^+)$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
1.2±0.4 OUR AVERAGE						

NODE=M150R2
NODE=M150R2

1.1±0.4 ^{+0.3} _{-0.2}	1371	¹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)0} \pi^+ X$
1.9±1.1±0.3		BERGFELD	94B	CLE2 $e^+e^- \rightarrow$ hadrons

¹ From the fit of the $M(D^0 \pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1, respectively.

NODE=M150R2;LINKAGE=AB

$\Gamma(D^0 \pi^+)/[\Gamma(D^0 \pi^+) + \Gamma(D^{*0} \pi^+)]$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/(\Gamma_1+\Gamma_2)$
--	-------	------	-------------	------	---------	--------------------------------

NODE=M150R01
NODE=M150R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.62±0.03±0.02	3361	¹ AUBERT	09Y	BABR $\bar{B}^0 \rightarrow D_2^{*+} \ell^- \nu_\ell$
----------------	------	---------------------	-----	---

¹ Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+ B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths for charged and neutral D_2^* mesons.

NODE=M150R01;LINKAGE=AU

$D_2^*(2460)^\pm$ REFERENCES

AAIJ	15X	PR D92 012012	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56588
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56609
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
ABRAMOWICZ	13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54743
DEL-AMO-SA...	10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52929
KUZMIN	07	PR D76 012006	A. Kuzmin <i>et al.</i>	(BELLE Collab.)	REFID=51854
LINK	04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)	REFID=49775
BERGFELD	94B	PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)	REFID=44099
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
ALBRECHT	89B	PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40736
ALBRECHT	89F	PL B231 208	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40931

NODE=M150

 $D(2550)^0$

$$I(J^P) = \frac{1}{2}(?^?)$$

OMITTED FROM SUMMARY TABLE

Unnatural parity according to the helicity analysis of DEL-AMO-SANCHEZ 10P and AAIJ 13CC. DEL-AMO-SANCHEZ 10P suggests $J^P = 0^-$.

NODE=M198

NODE=M198

 $D(2550)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2564 ± 20	OUR AVERAGE	Error includes scale factor of 3.9.		[2539 ± 8 MeV OUR 2015 AVERAGE]
2579.5 ± 3.4 ± 5.5	60k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
2539.4 ± 4.5 ± 6.8	34k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$

NODE=M198M

NODE=M198M
NEW $D(2550)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
135 ± 17	OUR AVERAGE	[130 ± 18 MeV OUR 2015 AVERAGE]		
177.5 ± 17.8 ± 46.0	60k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
130 ± 12 ± 13	34k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$

NODE=M198W

NODE=M198W
NEW $D(2550)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^{*+} \pi^-$	seen

NODE=M198215;NODE=M198

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D(2550)^0$ POLARIZATION AMPLITUDE A_{D_J}

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$ decays.

NODE=M198PAM

NODE=M198PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 1.3	60k	¹ AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
¹ Systematic uncertainty not estimated.				

NODE=M198PAM

NODE=M198PAM;LINKAGE=A

 $D(2550)^0$ REFERENCES

AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
DEL-AMO-SA...	10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534

NODE=M198

NODE=M199

$D_J^*(2600)$
was $D(2600)$,

$$I(J^P) = \frac{1}{2}(??)$$

OMITTED FROM SUMMARY TABLE

J^P consistent with natural parity (DEL-AMO-SANCHEZ 10P, AAIJ 13CC).

NODE=M199

$D_J^*(2600)$ MASS

NODE=M199M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2622 ± 12	OUR AVERAGE	Error includes scale factor of 4.7. See the ideogram below. [2612 ± 6 MeV OUR 2015 AVERAGE Scale factor = 1.9]			
2649.2 ± 3.5 ± 3.5	51k	AAIJ	13CC	LHCB	$pp \rightarrow D^{*+} \pi^- X$
2608.7 ± 2.4 ± 2.5	26k	DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^+ \pi^- X$
2621.3 ± 3.7 ± 4.2	13k	¹ DEL-AMO-SA..10P	BABR	+	$e^+ e^- \rightarrow D^0 \pi^+ X$


NODE=M199M

NEW

OCCUR=2

NODE=M199M;LINKAGE=DE

¹ At a fixed width of 93 MeV.



$D_J^*(2600)$ WIDTH

NODE=M199W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
104 ± 20	OUR AVERAGE	Error includes scale factor of 1.6. [93 ± 14 MeV OUR 2015 AVERAGE]		
140.2 ± 17.1 ± 18.6	51k	AAIJ	13CC	LHCB $pp \rightarrow D^{*+} \pi^- X$
93 ± 6 ± 13	26k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$

NODE=M199W

NEW

$D_J^*(2600)$ DECAY MODES

NODE=M199215;NODE=M199

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\pi$	seen
Γ_2 $D^+ \pi^-$	seen
Γ_3 $D^0 \pi^\pm$	seen
Γ_4 $D^* \pi$	seen
Γ_5 $D^{*+} \pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4;OUR EVAL;→ UNCHECKED ←

DESIG=5;OUR EVAL;→ UNCHECKED ←

$D^*(2600)$ BRANCHING RATIOS

$\Gamma(D^+ \pi^-) / \Gamma(D^{*+} \pi^-)$					Γ_2 / Γ_5
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
0.32 ± 0.02 ± 0.09	76k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{(*)+} \pi^- X$	

NODE=M199220

NODE=M199R01
NODE=M199R01

$D^*(2600)$ REFERENCES

AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA...10P	PR D82	111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)

NODE=M199

REFID=55581
REFID=53534

NODE=M158

$D^*(2640)^\pm$

$$I(J^P) = \frac{1}{2}(?^?)$$

OMITTED FROM SUMMARY TABLE

Seen in Z decays by ABREU 98M. Not seen by ABBIENDI 01N and CHEKANOV 09. Needs confirmation.

NODE=M158

$D^*(2640)^\pm$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2637 ± 2 ± 6	66 ± 14	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

NODE=M158M

NODE=M158M

$D^*(2640)^\pm$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<15	95	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

NODE=M158W

NODE=M158W

$D^*(2640)^+$ DECAY MODES

$D^*(2640)^-$ modes are charge conjugates of modes below.

NODE=M158215;NODE=M158

NODE=M158

Mode	Fraction (Γ_i / Γ)
Γ_1 $D^*(2010)^+ \pi^+ \pi^-$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

$D^*(2640)^\pm$ REFERENCES

CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABBIENDI	01N	EPJ C20 445	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)

NODE=M158

REFID=52733
REFID=48296
REFID=46315

$D(2740)^0$

$$I(J^P) = \frac{1}{2}(?^?)$$

OMITTED FROM SUMMARY TABLE

 J^P consistent with unnatural parity (AAIJ 13CC).

NODE=M228

NODE=M228

 $D(2740)^0$ MASS

NODE=M228M

NODE=M228M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2737.0±3.5±11.2	7.7k	AAIJ	13CC LHCB	$\rho\rho \rightarrow D^{*+}\pi^- X$

 $D(2740)^0$ WIDTH

NODE=M228W

NODE=M228W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
73.2±13.4±25.0	7.7k	AAIJ	13CC LHCB	$\rho\rho \rightarrow D^{*+}\pi^- X$

 $D(2740)^0$ DECAY MODES

NODE=M228215;NODE=M228

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^{*+}\pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D(2740)^0$ POLARIZATION AMPLITUDE A_{D_J}

NODE=M228PAM

NODE=M228PAM

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^*\pi$ and $D^* \rightarrow D\pi$ decays.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±2.2	7.7k	¹ AAIJ	13CC LHCB	$\rho\rho \rightarrow D^{*+}\pi^- X$

¹Systematic uncertainty not estimated.

NODE=M228PAM

NODE=M228PAM;LINKAGE=A

 $D(2740)^0$ REFERENCESAAIJ 13CC JHEP 1309 145 R. Aaij *et al.* (LHCb Collab.)

NODE=M228

REFID=55581

NODE=M203

 $D(2750)$

$$I(J^P) = \frac{1}{2}(3^-)$$

OMITTED FROM SUMMARY TABLE

 J^P determined by AAIJ 15Y from the Dalitz plot analysis of $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$ decays. J^P consistent with natural parity (AAIJ 13CC).

NODE=M203

 $D(2750)$ MASS

NODE=M203M

NODE=M203M

NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2763 ± 4 OUR AVERAGE					Error includes scale factor of 2.3. See the ideogram below. [2761 ± 5 MeV OUR 2015 AVERAGE Scale factor = 2.5]
2798 ± 7 ± 7		¹ AAIJ	15Y LHCB		$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2761.1± 5.1± 6.5	14k	AAIJ	13CC LHCB	0	$\rho\rho \rightarrow D^{*+}\pi^- X$
2760.1± 1.1± 3.7	56k	AAIJ	13CC LHCB	0	$\rho\rho \rightarrow D^+\pi^- X$
2771.7± 1.7± 3.8	20k	AAIJ	13CC LHCB	+	$\rho\rho \rightarrow D^0 \pi^+ X$
2752.4± 1.7± 2.7	23.5k	² DEL-AMO-SA..10P	BABR	0	$e^+e^- \rightarrow D^{*+}\pi^- X$
2763.3± 2.3± 2.3	11.3k	² DEL-AMO-SA..10P	BABR	0	$e^+e^- \rightarrow D^+\pi^- X$
2769.7± 3.8± 1.5	5.7k	^{2,3} DEL-AMO-SA..10P	BABR	+	$e^+e^- \rightarrow D^0 \pi^+ X$
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
2802 ±11 ±10		⁴ AAIJ	15Y LHCB		$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$

NODE=M203M

NEW

OCCUR=2

OCCUR=3


OCCUR=2

OCCUR=3

OCCUR=2

- 1 Modeling the $\pi^+\pi^-$ S-wave with the Isobar formalism.
- 2 The states observed in the $D^*\pi$ and $D\pi$ final states are not necessarily the same.
- 3 At a fixed width of 60.9 MeV.
- 4 Modeling the $\pi^+\pi^-$ S-wave with the K-matrix formalism.

NODE=M203M;LINKAGE=A
 NODE=M203M;LINKAGE=DE
 NODE=M203M;LINKAGE=DA
 NODE=M203M;LINKAGE=B



D(2750) WIDTH

NODE=M203W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
65 ± 5 OUR AVERAGE					
[63 ± 6 MeV OUR 2015 AVERAGE]					
105 ±18 ±24		⁵ AAIJ	15Y	LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
74.4± 3.4±37.0	14k	AAIJ	13CC	LHCB 0	$pp \rightarrow D^{*+} \pi^- X$
74.4± 3.4±19.1	56k	AAIJ	13CC	LHCB 0	$pp \rightarrow D^+ \pi^- X$
66.7± 6.6±10.5	20k	AAIJ	13CC	LHCB +	$pp \rightarrow D^0 \pi^+ X$
71 ± 6 ±11	23.5k	⁶ DEL-AMO-SA..10P	BABR		$e^+ e^- \rightarrow D^{*+} \pi^- X$
60.9± 5.1± 3.6	11.3k	⁶ DEL-AMO-SA..10P	BABR		$e^+ e^- \rightarrow D^+ \pi^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
154 ±27 ±16		⁷ AAIJ	15Y	LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$

NODE=M203W
 NEW

OCCUR=2
 OCCUR=4

OCCUR=2

OCCUR=2

NODE=M203W;LINKAGE=A
 NODE=M203W;LINKAGE=DE
 NODE=M203W;LINKAGE=B

D(2750) DECAY MODES

NODE=M203215;NODE=M203

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\pi$	seen
Γ_2 $D^+ \pi^-$	seen
Γ_3 $D^0 \pi^\pm$	seen
Γ_4 $D^* \pi$	seen
Γ_5 $D^{*+} \pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=3;OUR EVAL;→ UNCHECKED ←
 DESIG=4;OUR EVAL;→ UNCHECKED ←
 DESIG=5;OUR EVAL;→ UNCHECKED ←

D(2750) BRANCHING RATIOS

NODE=M203220

$\Gamma(D^+ \pi^-)/\Gamma(D^{*+} \pi^-)$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_5
	0.42±0.05±0.11	34.8k	⁸ DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{(*)+} \pi^- X$	

NODE=M203R01
 NODE=M203R01

⁸The states observed in the $D^*\pi$ and $D\pi$ final states are not necessarily the same.

$D(2750)$ POLARIZATION AMPLITUDE A_D

A polarization amplitude A_D is a parameter that depends on the initial polarization of the $D(2750)$. For $D(2750)$ decays the helicity angle, θ_H , distribution varies like $1 + A_D \cos(\theta_H)$, where θ_H is the angle in the D^* rest frame between the two pions emitted by the $D(2750) \rightarrow D^*\pi$ and $D^* \rightarrow D\pi$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

-0.33 ± 0.28	23.5k	⁹ DEL-AMO-SA...10P	BABR	$e^+e^- \rightarrow D^{*+}\pi^- X$
------------------	-------	-------------------------------	------	------------------------------------

⁹Systematic uncertainties not estimated. The states observed in the $D^*\pi$ and $D\pi$ final states are not necessarily the same.

$D(2750)$ REFERENCES

AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA...10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	

$D(3000)^0$

$$I(J^P) = \frac{1}{2}(??)$$

OMITTED FROM SUMMARY TABLE

Both natural- and unnatural-parity components observed depending on the decay mode (AAIJ 13CC).

$D(3000)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

2971.8 ± 8.7	9.5k	^{1,2} AAIJ	13CC LHCB	$pp \rightarrow D^{*+}\pi^- X$
3008.1 ± 4.0	17.6k	^{1,3} AAIJ	13CC LHCB	$pp \rightarrow D^+\pi^- X$

¹Systematic uncertainty not estimated.

²Unnatural parity preferred.

³Natural parity state. A state $D(3000)^+$ is possibly seen in $D^0\pi^+$ final state.

$D(3000)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

188.1 ± 44.8	9.5k	^{4,5} AAIJ	13CC LHCB	$pp \rightarrow D^{*+}\pi^- X$
110.5 ± 11.5	17.6k	^{4,6} AAIJ	13CC LHCB	$pp \rightarrow D^+\pi^- X$

⁴Systematic uncertainty not estimated.

⁵Unnatural parity preferred.

⁶Natural parity state. A state $D(3000)^+$ is possibly seen in $D^0\pi^+$ final state.

$D(3000)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^{*+}\pi^-$	seen

$D(3000)^0$ POLARIZATION AMPLITUDE A_{D_J}

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^*\pi$ and $D^* \rightarrow D\pi$ decays.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

1.5 ± 0.9	9.5k	⁷ AAIJ	13CC LHCB	$pp \rightarrow D^{*+}\pi^- X$
---------------	------	-------------------	-----------	--------------------------------

NODE=M203R01;LINKAGE=DE

NODE=M203PAM

NODE=M203PAM

NODE=M203PAM

NODE=M203PAM;LINKAGE=DE

NODE=M203

REFID=56609

REFID=55581

REFID=53534

NODE=M229

NODE=M229

NODE=M229M

NODE=M229M

OCCUR=2

NODE=M229M;LINKAGE=A

NODE=M229M;LINKAGE=B

NODE=M229M;LINKAGE=C

NODE=M229W

NODE=M229W

OCCUR=2

NODE=M229W;LINKAGE=A

NODE=M229W;LINKAGE=C

NODE=M229W;LINKAGE=B

NODE=M229215;NODE=M229

DESIG=1;OUR EVAL;→ UNCHECKED ←

NODE=M229PAM

NODE=M229PAM

NODE=M229PAM

⁷Systematic uncertainty not estimated.

NODE=M229PAM;LINKAGE=A

 $D(3000)^0$ REFERENCES

AAIJ

13CC JHEP 1309 145

R. Aaij *et al.*

(LHCb Collab.)

NODE=M229

REFID=55581

**CHARMED, STRANGE MESONS
($C = S = \pm 1$)**

$$D_s^+ = c\bar{s}, D_s^- = \bar{c}s, \text{ similarly for } D_s^{*\pm}$$

 $D_s^{*\pm}$

$$I(J^P) = 0(?^?)$$

 J^P is natural, width and decay modes consistent with 1^- .

NODE=MXXX040

NODE=MXXX040

NODE=S074

NODE=S074

 $D_s^{*\pm}$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=S074M

NODE=S074M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2112.1 ± 0.4 OUR FIT			
2106.6 ± 2.1 ± 2.7	¹ BLAYLOCK	87	MRK3 $e^+ e^- \rightarrow D_s^\pm \gamma X$
¹ Assuming D_s^\pm mass = 1968.7 ± 0.9 MeV.			

NODE=S074M

NODE=S074M;LINKAGE=E

 $m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=S074DM

NODE=S074DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
143.8 ± 0.4 OUR FIT				
143.9 ± 0.4 OUR AVERAGE				
143.76 ± 0.39 ± 0.40		GRONBERG	95	CLE2 $e^+ e^-$
144.22 ± 0.47 ± 0.37		BROWN	94	CLE2 $e^+ e^-$
142.5 ± 0.8 ± 1.5		² ALBRECHT	88	ARG $e^+ e^- \rightarrow D_s^\pm \gamma X$
139.5 ± 8.3 ± 9.7	60	AIHARA	84D	TPC $e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
143.0 ± 18.0	8	ASRATYAN	85	HLBC FNAL 15-ft, ν - ² H
110 ± 46		BRANDELIK	79	DASP $e^+ e^- \rightarrow D_s^\pm \gamma X$
² Result includes data of ALBRECHT 84B.				

NODE=S074DM

NODE=S074DM;LINKAGE=A

 $D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.9	90	GRONBERG	95	CLE2 $e^+ e^-$
< 4.5	90	ALBRECHT	88	ARG $E_{cm}^{ee} = 10.2$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 4.9	90	BROWN	94	CLE2 $e^+ e^-$
< 22	90	BLAYLOCK	87	MRK3 $e^+ e^- \rightarrow D_s^\pm \gamma X$

NODE=S074W

NODE=S074W

D_s^{*+} DECAY MODES

NODE=S074215;NODE=S074

 D_s^{*-} modes are charge conjugates of the modes below.

NODE=S074

Mode	Fraction (Γ_i/Γ)
Γ_1 $D_s^+ \gamma$	(93.5±0.7) %
Γ_2 $D_s^+ \pi^0$	(5.8±0.7) %
Γ_3 $D_s^+ e^+ e^-$	(6.7±1.6) × 10 ⁻³

DESIG=1

DESIG=2

DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 3 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 0.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-97	
x_3	-19	-4
	x_1	x_2

 D_s^{*+} BRANCHING RATIOS

NODE=S074220

$\Gamma(D_s^+ \gamma) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.935±0.007 OUR FIT				

NODE=S074R1
NODE=S074R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ASRATYAN	91	HLBC	$\bar{\nu}_\mu \text{Ne}$
seen	ALBRECHT	88	ARG	$e^+ e^- \rightarrow D_s^\pm \gamma X$
seen	AIHARA	84D		
seen	ALBRECHT	84B		
seen	BRANDELIK	79		

$\Gamma(D_s^+ \pi^0) / \Gamma(D_s^+ \gamma)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.062±0.008 OUR FIT				

NODE=S074R2
NODE=S074R2

0.062±0.008 OUR AVERAGE				
0.062±0.005±0.006	AUBERT,BE	05G	BABR	10.6 $e^+ e^- \rightarrow$ hadrons
0.062 ^{+0.020} _{-0.018} ±0.022	GRONBERG	95	CLE2	$e^+ e^-$

$\Gamma(D_s^+ e^+ e^-) / \Gamma(D_s^+ \gamma)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
7.2±1.7 OUR FIT				

NODE=S074R01
NODE=S074R01

7.2^{+1.5}_{-1.3}±1.0	38	CRONIN-HEN..12	CLEO	4.17 $e^+ e^- \rightarrow$ hadrons
--	----	----------------	------	------------------------------------

 D_s^{*+} REFERENCES

NODE=S074

CRONIN-HEN...12	PR D86 072005	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=54627
AUBERT,BE 05G	PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50942
GRONBERG 95	PRL 75 3232	J. Gronberg <i>et al.</i>	(CLEO Collab.)	REFID=44568
BROWN 94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)	REFID=43868
ASRATYAN 91	PL B257 525	A.E. Asratyan <i>et al.</i>	(ITEP, BELG, SACL+)	REFID=41582
ALBRECHT 88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40269
BLAYLOCK 87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)	REFID=40005
ASRATYAN 85	PL 156B 441	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)	REFID=22887
AIHARA 84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=11561
ALBRECHT 84B	PL 146B 111	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22886
BRANDELIK 79	PL 80B 412	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=11442

$D_{s0}^*(2317)^\pm$
 $I(J^P) = 0(0^+)$
 J, P need confirmation.

AUBERT 06P and CHOI 15A do not observe neutral and doubly charged partners of the $D_{s0}^*(2317)^\pm$.

NODE=M172

NODE=M172

 $D_{s0}^*(2317)^\pm$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M172M

NODE=M172M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2317.7±0.6 OUR FIT	Error includes scale factor of 1.1.			
2318.0±1.0 OUR AVERAGE	Error includes scale factor of 1.4.			
2319.6±0.2±1.4	3180	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^0 X$
2317.3±0.4±0.8	1022	¹ AUBERT	04E BABR	10.6 $e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2317.2±1.3	88	² AUBERT,B	04S BABR	$B \rightarrow D_{s0}^{(*)}(2317)^+ \bar{D}^{(*)}$
2317.2±0.5±0.9	761	³ MIKAMI	04 BELL	10.6 $e^+ e^-$
2316.8±0.4±3.0	1267 ± 53	^{3,4} AUBERT	03G BABR	10.6 $e^+ e^-$
2317.6±1.3	273 ± 33	^{3,5} AUBERT	03G BABR	10.6 $e^+ e^-$
2319.8±2.1±2.0	24	³ KROKOVNY	03B BELL	10.6 $e^+ e^-$
¹ Supersedes AUBERT 03G.				
² Systematic errors not evaluated.				
³ Not independent of the corresponding $m_{D_{s0}^*(2317)} - m_{D_s}$.				
⁴ From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.				
⁵ From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.				

NODE=M172M

OCCUR=2

NODE=M172M;LINKAGE=AU

NODE=M172M;LINKAGE=AB

NODE=M172M;LINKAGE=B1

NODE=M172M;LINKAGE=A1

NODE=M172M;LINKAGE=A2

 $m_{D_{s0}^*(2317)^\pm} = m_{D_s^\pm}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M172DM

NODE=M172DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
349.4±0.6 OUR FIT	Error includes scale factor of 1.1.			
349.2±0.7 OUR AVERAGE				
348.7±0.5±0.7	761	MIKAMI	04 BELL	10.6 $e^+ e^-$
350.0±1.2±1.0	135	BESSION	03 CLE2	10.6 $e^+ e^-$
351.3±2.1±1.9	24	⁶ KROKOVNY	03B BELL	10.6 $e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
349.6±0.4±3.0	1267	^{7,8} AUBERT	03G BABR	10.6 $e^+ e^-$
350.2±1.3	273	^{9,10} AUBERT	03G BABR	10.6 $e^+ e^-$
⁶ Recalculated by us using $m_{D_s^+} = 1968.5 \pm 0.6$ MeV.				
⁷ From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.				
⁸ Recalculated by us using $m_{D_s^+} = 1967.20 \pm 0.03$ MeV.				
⁹ From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.				
¹⁰ Recalculated by us using $m_{D_s^+} = 1967.4 \pm 0.2$ MeV. Systematic errors not estimated.				

NODE=M172DM

OCCUR=2

NODE=M172DM;LINKAGE=K3

NODE=M172DM;LINKAGE=A1

NODE=M172DM;LINKAGE=C1

NODE=M172DM;LINKAGE=A2

NODE=M172DM;LINKAGE=C2

 $D_{s0}^*(2317)^\pm$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.8	95	3180	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^0 X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 4.6	90	761	MIKAMI	04 BELL	10.6 $e^+ e^-$
< 10			AUBERT	03G BABR	10.6 $e^+ e^-$
< 7	90	135	BESSION	03 CLE2	10.6 $e^+ e^-$

NODE=M172W

NODE=M172W

$D_{s0}^*(2317)^\pm$ DECAY MODES

NODE=M172215;NODE=M172

 $D_{s0}^*(2317)^-$ modes are charge conjugates of modes below.

NODE=M172

Mode	Fraction (Γ_i/Γ)
Γ_1 $D_s^+ \pi^0$	seen
Γ_2 $D_s^+ \gamma$	
Γ_3 $D_s^*(2112)^+ \gamma$	
Γ_4 $D_s^+ \gamma \gamma$	
Γ_5 $D_s^*(2112)^+ \pi^0$	
Γ_6 $D_s^+ \pi^+ \pi^-$	
Γ_7 $D_s^+ \pi^0 \pi^0$	not seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7;OUR EVAL;→ UNCHECKED ←

 $D_{s0}^*(2317)^\pm$ BRANCHING RATIOS

NODE=M172220

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	1540 ± 62	AUBERT	03G BABR	10.6 $e^+ e^-$	

NODE=M172R1
NODE=M172R1

$\Gamma(D_s^+ \gamma)/\Gamma(D_s^+ \pi^0)$					Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.05	90	MIKAMI	04 BELL	10.6 $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.14	95	AUBERT	06P BABR	10.6 $e^+ e^-$	
<0.052	90	BESSION	03 CLE2	10.6 $e^+ e^-$	

NODE=M172R5
NODE=M172R5

$\Gamma(D_s^*(2112)^+ \gamma)/\Gamma(D_s^+ \pi^0)$					Γ_3/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.059	90	BESSION	03 CLE2	10.6 $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.16	95	AUBERT	06P BABR	10.6 $e^+ e^-$	
<0.18	90	MIKAMI	04 BELL	10.6 $e^+ e^-$	

NODE=M172R6
NODE=M172R6

$\Gamma(D_s^+ \gamma \gamma)/\Gamma(D_s^+ \pi^0)$					Γ_4/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.18	95	AUBERT	06P BABR	10.6 $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen		AUBERT	03G BABR	10.6 $e^+ e^-$	

NODE=M172R7
NODE=M172R7

$\Gamma(D_s^*(2112)^+ \pi^0)/\Gamma(D_s^+ \pi^0)$					Γ_5/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.11	90	BESSION	03 CLE2	10.6 $e^+ e^-$	

NODE=M172R8
NODE=M172R8

$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma(D_s^+ \pi^0)$					Γ_6/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.004	90	MIKAMI	04 BELL	10.6 $e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.005	95	AUBERT	06P BABR	10.6 $e^+ e^-$	
<0.019	90	BESSION	03 CLE2	10.6 $e^+ e^-$	

NODE=M172R9
NODE=M172R9

$\Gamma(D_s^+ \pi^0 \pi^0)/\Gamma(D_s^+ \pi^0)$					Γ_7/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.25	95	AUBERT	06P BABR	10.6 $e^+ e^-$	

NODE=M172R10
NODE=M172R10 $D_{s0}^*(2317)^\pm$ REFERENCES

NODE=M172

CHOI	15A	PR D91 092011	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=56577
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49747
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50195
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)	REFID=49629
AUBERT	03G	PRL 90 242001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49417
BESSION	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=49583
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=49615

NODE=M173

 $D_{s1}(2460)^\pm$

$$I(J^P) = 0(1^+)$$

 $D_{s1}(2460)^\pm$ MASS

NODE=M173M

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M173M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M173M

2459.5±0.6 OUR FIT Error includes scale factor of 1.1.**2459.6±0.9 OUR AVERAGE** Error includes scale factor of 1.3.2460.1±0.2±0.8 ¹ AUBERT 06P BABR 10.6 e⁺e⁻2458.0±1.0±1.0 195 AUBERT 04E BABR 10.6 e⁺e⁻

• • • We do not use the following data for averages, fits, limits, etc. • • •

2459.5±1.2±3.7 920 AUBERT 06P BABR 10.6 e⁺e⁻ → D_s⁺γX

OCCUR=2

2458.6±1.0±2.5 560 AUBERT 06P BABR 10.6 e⁺e⁻ → D_s⁺π⁰γX

OCCUR=3

2460.2±0.2±0.8 123 AUBERT 06P BABR 10.6 e⁺e⁻ → D_s⁺π⁺π⁻X

OCCUR=4

2458.9±1.5 112 ² AUBERT,B 04S BABR B → D_{s1}(2460)⁺ $\bar{D}^{(*)}$ 2461.1±1.6 139 ³ AUBERT,B 04S BABR B → D_{s1}(2460)⁺ $\bar{D}^{(*)}$

OCCUR=2

2456.5±1.3±1.3 126 ^{4,5} MIKAMI 04 BELL 10.6 e⁺e⁻2459.5±1.3±2.0 152 ^{6,7} MIKAMI 04 BELL 10.6 e⁺e⁻

OCCUR=2

2459.9±0.9±1.6 60 ^{6,7} MIKAMI 04 BELL 10.6 e⁺e⁻

OCCUR=3

2459.2±1.6±2.0 57 KROKOVNY 03B BELL 10.6 e⁺e⁻¹ The average of the values obtained from the D_s⁺γ, D_s⁺π⁰γ, D_s⁺π⁺π⁻ final state.

NODE=M173M;LINKAGE=UB

² Systematic errors not evaluated. From the decay to D_s⁺π⁰.

NODE=M173M;LINKAGE=AU

³ Systematic errors not evaluated. From the decay to D_s⁺γ.

NODE=M173M;LINKAGE=AB

⁴ Not independent of the corresponding m_{D_{s1}(2460)[±]} - m_{D_s[±]*}.

NODE=M173M;LINKAGE=B1

⁵ Using m_{D_s⁺*} = 2112.4 ± 0.7 MeV.

NODE=M173M;LINKAGE=B2

⁶ Not independent of the corresponding m_{D_{s1}(2460)[±]} - m_{D_s[±]*}.

NODE=M173M;LINKAGE=B3

⁷ Using m_{D_s⁺} = 1968.5 ± 0.6 MeV.

NODE=M173M;LINKAGE=B4

 $m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$

NODE=M173MD

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.


NODE=M173MD

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M173MD

347.3±0.7 OUR FIT Error includes scale factor of 1.2.**347.1±2.2 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.344.1±1.3±1.1 126 MIKAMI 04 BELL 10.6 e⁺e⁻351.2±1.7±1.0 41 BESSON 03 CLE2 10.6 e⁺e⁻346.8±1.6±1.9 57 ⁸ KROKOVNY 03B BELL 10.6 e⁺e⁻⁸ Recalculated by us using m_{D_s⁺*} = 2112.4 ± 0.7 MeV.

NODE=M173MD;LINKAGE=K3



$m_{D_{s1}(2460)^{\pm}} - m_{D_s^{\pm}}$

The fit includes $D^{\pm}, D^0, D_s^{\pm}, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^{\pm}$ mass and mass difference measurements.

NODE=M173DM

NODE=M173DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
491.2±0.6 OUR FIT				Error includes scale factor of 1.1.
491.3±1.4 OUR AVERAGE				
491.0±1.3±1.9	152	⁹ MIKAMI	04 BELL	10.6 e^+e^-
491.4±0.9±1.5	60	¹⁰ MIKAMI	04 BELL	10.6 e^+e^-

⁹ From the decay to $D_s^{\pm}\gamma$.
¹⁰ From the decay to $D_s^{\pm}\pi^+\pi^-$.

NODE=M173DM

OCCUR=2

NODE=M173DM;LINKAGE=M1
 NODE=M173DM;LINKAGE=M2

$D_{s1}(2460)^{\pm}$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.5	95	123	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow D_s^+\pi^+\pi^-X$
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
< 6.3	95	560	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow D_s^+\pi^0\gamma X$
<10		195	AUBERT	04E BABR	10.6 e^+e^-
< 5.5	90	126	MIKAMI	04 BELL	10.6 e^+e^-
< 7	90	41	BESSON	03 CLE2	10.6 e^+e^-

NODE=M173W

NODE=M173W

OCCUR=2

$D_{s1}(2460)^+$ DECAY MODES

$D_{s1}(2460)^-$ modes are charge conjugates of the modes below.

NODE=M173215;NODE=M173

NODE=M173

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $D_s^{*+}\pi^0$	(48 ± 11) %	
Γ_2 $D_s^+\gamma$	(18 ± 4) %	
Γ_3 $D_s^+\pi^+\pi^-$	(4.3± 1.3) %	S=1.1
Γ_4 $D_s^{*+}\gamma$	< 8 %	CL=90%
Γ_5 $D_{s0}^*(2317)^+\gamma$	(3.7 ⁺ _{-2.4}) %	
Γ_6 $D_s^+\pi^0$		
Γ_7 $D_s^+\pi^0\pi^0$		
Γ_8 $D_s^+\gamma\gamma$		

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=7

DESIG=8

DESIG=9

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 8 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 3.4$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	80		
x_3	68	62	
x_5	-3	25	26
	x_1	x_2	x_3

$D_{s1}(2460)^\pm$ BRANCHING RATIOS

NODE=M173220

$\Gamma(D_s^{*+} \pi^0) / \Gamma_{\text{total}}$ Γ_1 / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.48 ± 0.11 OUR FIT**0.56 ± 0.13 ± 0.09**¹¹ AUBERT 06N BABR $B \rightarrow D_{s1}(2460)^- \bar{D}^{(*)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 41 BESSON 03 CLE2 10.6 $e^+ e^-$ ¹¹ Evaluated in AUBERT 06N including measurements from AUBERT,B 04s.NODE=M173R1
NODE=M173R1

NODE=M173R1;LINKAGE=AU

$\Gamma(D_s^+ \gamma) / \Gamma_{\text{total}}$ Γ_2 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.18 ± 0.04 OUR FIT**0.16 ± 0.04 ± 0.03**¹² AUBERT 06N BABR $B \rightarrow D_{s1}(2460)^- \bar{D}^{(*)}$ ¹² Evaluated in AUBERT 06N including measurements from AUBERT,B 04s.NODE=M173R6
NODE=M173R6

NODE=M173R6;LINKAGE=AU

$\Gamma(D_s^+ \gamma) / \Gamma(D_s^{*+} \pi^0)$ Γ_2 / Γ_1

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

0.38 ± 0.05 OUR FIT**0.44 ± 0.09 OUR AVERAGE**0.55 ± 0.13 ± 0.08 152 MIKAMI 04 BELL 10.6 $e^+ e^-$ 0.38 ± 0.11 ± 0.04 38 KROKOVNY 03B BELL 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.274 ± 0.045 ± 0.020 251 ¹³ AUBERT,B 04s BABR $B \rightarrow D_{s1}(2460)^+ \bar{D}^{(*)}$ < 0.49 90 BESSON 03 CLE2 10.6 $e^+ e^-$ ¹³ Used by AUBERT 06N in their measurement of $B(D_s^{*-} \pi^0)$ and $B(D_s^- \gamma)$.NODE=M173R2
NODE=M173R2

NODE=M173R2;LINKAGE=AU

$\Gamma(D_s^+ \pi^+ \pi^-) / \Gamma(D_s^{*+} \pi^0)$ Γ_3 / Γ_1

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

0.090 ± 0.020 OUR FIT Error includes scale factor of 1.2.**0.14 ± 0.04 ± 0.02** 60 MIKAMI 04 BELL 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.08 90 BESSON 03 CLE2 10.6 $e^+ e^-$ NODE=M173R3
NODE=M173R3

$\Gamma(D_s^{*+} \gamma) / \Gamma(D_s^{*+} \pi^0)$ Γ_4 / Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< **0.16** 90 BESSON 03 CLE2 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.31 90 MIKAMI 04 BELL 10.6 $e^+ e^-$ NODE=M173R4
NODE=M173R4

$\Gamma(D_{s0}^*(2317)^+ \gamma) / \Gamma(D_s^{*+} \pi^0)$ Γ_5 / Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< **0.22** 95 AUBERT 04E BABR 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.58 90 BESSON 03 CLE2 10.6 $e^+ e^-$ NODE=M173R5
NODE=M173R5

$\Gamma(D_s^{*+} \pi^0) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_1 / (\Gamma_1 + \Gamma_5)$	
VALUE	DOCUMENT ID	TECN	COMMENT			NODE=M173R7 NODE=M173R7
0.93±0.09 OUR FIT						
0.97±0.09±0.05	AUBERT	06P	BABR	10.6 e ⁺ e ⁻		
$\Gamma(D_s^+ \gamma) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_2 / (\Gamma_1 + \Gamma_5)$	
VALUE	DOCUMENT ID	TECN	COMMENT			NODE=M173R8 NODE=M173R8
0.35 ±0.04 OUR FIT						
0.337±0.036±0.038	AUBERT	06P	BABR	10.6 e ⁺ e ⁻		
$\Gamma(D_s^+ \pi^+ \pi^-) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_3 / (\Gamma_1 + \Gamma_5)$	
VALUE	DOCUMENT ID	TECN	COMMENT			NODE=M173R9 NODE=M173R9
0.083±0.017 OUR FIT						
				Error includes scale factor of 1.2.		
0.077±0.013±0.008	AUBERT	06P	BABR	10.6 e ⁺ e ⁻		
$\Gamma(D_s^{*+} \gamma) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_4 / (\Gamma_1 + \Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M173R10 NODE=M173R10
<0.24	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	
$\Gamma(D_{s0}^*(2317)^+ \gamma) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_5 / (\Gamma_1 + \Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M173R11 NODE=M173R11
<0.25	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	
$\Gamma(D_s^+ \pi^0) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_6 / (\Gamma_1 + \Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M173R12 NODE=M173R12
<0.042	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	
$\Gamma(D_s^+ \pi^0 \pi^0) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_7 / (\Gamma_1 + \Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M173R13 NODE=M173R13
<0.68	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	
$\Gamma(D_s^+ \gamma \gamma) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$					$\Gamma_8 / (\Gamma_1 + \Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M173R14 NODE=M173R14
<0.33	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	

$D_{s1}(2460)^\pm$ REFERENCES

					NODE=M173
AUBERT	06N	PR D74 031103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51142
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49747
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50195
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)	REFID=49629
BESSION	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=49583
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=49615

$D_{s1}(2536)^\pm$
 $I(J^P) = 0(1^+)$
 J, P need confirmation.

Seen in $D^*(2010)^+ K^0$, $D^*(2007)^0 K^+$, and $D_s^+ \pi^+ \pi^-$. Not seen in $D^+ K^0$ or $D^0 K^+$. $J^P = 1^+$ assignment strongly favored.

NODE=M121

NODE=M121

 $D_{s1}(2536)^\pm$ MASS

NODE=M121M

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M121M

2535.10±0.06 OUR FIT

NEW

[2535.11 ± 0.06 MeV OUR 2015 FIT]

2535.18±0.24 OUR AVERAGE

2535.7 ± 0.6 ± 0.5	46 ± 9	1 ABAZOV	09G D0	$B_s^0 \rightarrow D_{s1}^- \mu^+ \nu_\mu X$
2534.78 ± 0.31 ± 0.40	182	AUBERT	08B BABR	$B \rightarrow \bar{D}^{(*)} D^* K$
2534.6 ± 0.3 ± 0.7	193	AUBERT	06P BABR	$10.6 e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$
2535.3 ± 0.7	92	2 HEISTER	02B ALEP	$e^+ e^- \rightarrow D^{*+} K^0 X, D^{*0} K^+ X$
2534.2 ± 1.2	9	ASRATYAN	94 BEBC	$\nu N \rightarrow D^* K^0 X, D^{*0} K^\pm X$
2535 ± 0.6 ± 1	75	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} K^0 X, D^{*0} K^+ X$
2535.3 ± 0.2 ± 0.5	134	ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*0} K^+ X$
2534.8 ± 0.6 ± 0.6	44	ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535.2 ± 0.5 ± 1.5	28	ALBRECHT	92R ARG	$10.4 e^+ e^- \rightarrow D^{*0} K^+ X$
2536.6 ± 0.7 ± 0.4		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535.9 ± 0.6 ± 0.2		ALBRECHT	89E ARG	$D_{s1}^- \rightarrow D^*(2010) K^0$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

2534.1 ± 0.6	116	3 AUSHEV	11 BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
2535.08 ± 0.01 ± 0.15	8038	4 LEES	11B BABR	$10.6 e^+ e^- \rightarrow D^{*+} K_S^0 X$
2535.57 ^{+0.44} _{-0.41} ± 0.10	236 ± 30	5 CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X, D^{*0} K^+ X$
2535 ± 28		6 ASRATYAN	88 HLBC	$\nu N \rightarrow D_s \gamma \gamma X$

¹ Using the $D^*(2010)^\pm$ mass of 2010.0 ± 0.4 MeV from PDG 06.

² Calculated using $m(D^*(2010)^\pm) = 2010.0 \pm 0.5$ MeV, $m(D^*(2007)^0) = 2006.7 \pm 0.5$ MeV, and the mass difference below.

³ Systematic uncertainties not evaluated.

⁴ Calculated using the mass difference $m(D_{s1}^+) - m(D^{*+})_{PDG}$ below and $m(D^{*+})_{PDG} = 2010.25 \pm 0.14$ MeV. Assuming S-wave decay of the $D_{s1}(2536)$ to $D^{*+} K_S^0$, using a Breit-Wigner line shape corresponding to L=0.

⁵ Calculated using the mass difference $m(D_{s1}^+) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV.

⁶ Not seen in $D^* K$.

NODE=M121M;LINKAGE=AB

NODE=M121M;LINKAGE=HI

NODE=M121M;LINKAGE=AU

NODE=M121M;LINKAGE=LE

NODE=M121M;LINKAGE=CH

NODE=M121M;LINKAGE=B

 $m_{D_{s1}(2536)^\pm} - m_{D_s^*(2111)}$

NODE=M121DM

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M121DM

423.0± 0.4 OUR FIT**424 ± 28**ASRATYAN 88 HLBC $D_s^{*\pm} \gamma$ **$m_{D_{s1}(2536)^\pm} - m_{D^*(2010)^\pm}$**

NODE=M121DN

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121DN

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
524.84±0.04 OUR FIT				
524.84±0.04 OUR AVERAGE				
524.83±0.01±0.04	8038	⁷ LEES	11B BABR	10.6 e ⁺ e ⁻ → D ^{*+} K _S ⁰ X
525.30 ^{+0.44} _{-0.41} ±0.10	236 ± 30	CHEKANOV 09	ZEUS	e [±] p → D ^{*+} K _S ⁰ X,
525.3 ±0.6 ±0.1	41	HEISTER	02B ALEP	e ⁺ e ⁻ → D ^{*0} K ⁺ X D ^{*0} K ⁺ X

⁷ Assuming S-wave decay of the D_{s1}(2536) to D^{*+}K_S⁰, using a Breit-Wigner line shape corresponding to L=0.

NODE=M121DN

NODE=M121DN;LINKAGE=LE

$m_{D_{s1}(2536)^\pm} - m_{D^*(2007)^0}$

The fit includes D[±], D⁰, D_S[±], D^{*±}, D^{*0}, D_S^{*±}, D₁(2420)⁰, D₂^{*}(2460)⁰, and D_{s1}(2536)[±] mass and mass difference measurements.

NODE=M121DP

NODE=M121DP

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
528.25±0.05 OUR FIT Error includes scale factor of 1.1. [528.14 ± 0.08 MeV OUR 2015 FIT]				
528.1 ±1.5 OUR AVERAGE				
528.7 ±1.9 ±0.5	51	HEISTER	02B ALEP	e ⁺ e ⁻ → D ^{*0} K ⁺ X
527.3 ±2.2	29	ACKERSTAFF	97W OPAL	e ⁺ e ⁻ → D ^{*0} K ⁺ X

NODE=M121DP

NEW

D_{s1}(2536)[±] WIDTH

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
0.92±0.03±0.04	8038	⁸ LEES	11B BABR	10.6 e ⁺ e ⁻ → D ^{*+} K _S ⁰ X
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.75±0.23	116	⁹ AUSHEV	11 BELL	B → D _{s1} (2536) ⁺ D ^(*)
< 2.5	95 193	AUBERT	06P BABR	10.6 e ⁺ e ⁻ → D _S ⁺ π ⁺ π ⁻ X
< 3.2	90 75	FRABETTI	94B E687	γBe → D ^{*+} K ⁰ X, D ^{*0} K ⁺ X
< 2.3	90	ALEXANDER	93 CLEO	e ⁺ e ⁻ → D ^{*0} K ⁺ X
< 3.9	90	ALBRECHT	92R ARG	10.4 e ⁺ e ⁻ → D ^{*0} K ⁺ X
< 5.44	90	AVERY	90 CLEO	e ⁺ e ⁻ → D ^{*+} K ⁰ X
< 4.6	90	ALBRECHT	89E ARG	D _{s1} [*] → D [*] (2010)K ⁰

NODE=M121W

NODE=M121W

⁸ Assuming S-wave decay of the D_{s1}(2536) to D^{*+}K_S⁰, using a Breit-Wigner line shape corresponding to L=0.

NODE=M121W;LINKAGE=LE

⁹ Systematic uncertainties not evaluated.

NODE=M121W;LINKAGE=AU

D_{s1}(2536)⁺ DECAY MODES

NODE=M121215;NODE=M121

D_{s1}(2536)⁻ modes are charge conjugates of the modes below.

NODE=M121

Mode	Fraction (Γ _i /Γ)	Confidence level
Γ ₁ D [*] (2010) ⁺ K ⁰	0.85 ±0.12	
Γ ₂ (D [*] (2010) ⁺ K ⁰) _{S-wave}	0.61 ±0.09	
Γ ₃ (D [*] (2010) ⁺ K ⁰) _{D-wave}		
Γ ₄ D ⁺ π ⁻ K ⁺	0.028±0.005	
Γ ₅ D [*] (2007) ⁰ K ⁺	DEFINED AS 1	
Γ ₆ D ⁺ K ⁰	<0.34	90%
Γ ₇ D ⁰ K ⁺	<0.12	90%
Γ ₈ D _S ^{*+} γ	possibly seen	
Γ ₉ D _S ⁺ π ⁺ π ⁻	seen	

DESIG=1

DESIG=7

DESIG=9

DESIG=8

DESIG=4

DESIG=2

DESIG=5

DESIG=3

DESIG=6

$D_{s1}(2536)^+$ BRANCHING RATIOS

NODE=M121220

$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^*(2010)^+ K^0)$

 Γ_5/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.18±0.16 OUR AVERAGE				
0.88±0.24±0.08	116	AUSHEV 11	BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
2.3 ±0.6 ±0.3	236 ± 30	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X,$ $D^{*0} K^+ X$
1.32±0.47±0.23	92	¹⁰ HEISTER 02B	ALEP	$e^+ e^- \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
1.9 $\begin{smallmatrix} +1.1 \\ -0.9 \end{smallmatrix}$ ±0.4	35	¹⁰ ACKERSTAFF 97W	OPAL	$e^+ e^- \rightarrow D^{*0} K^+ X,$ $D^{*+} K^0 X$
1.1 ±0.3		ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X, D^{*+} K^0 X$
1.4 ±0.3 ±0.2		¹¹ ALBRECHT 92R	ARG	10.4 $e^+ e^- \rightarrow D^{*0} K^+ X, D^{*+} K^0 X$

NODE=M121R6
NODE=M121R6¹⁰ Ratio of the production rates measured in Z^0 decays.¹¹ Evaluated by us from published inclusive cross-sections.NODE=M121R6;LINKAGE=6A
NODE=M121R6;LINKAGE=A

$\Gamma((D^*(2010)^+ K^0)_{S-wave})/\Gamma(D^*(2010)^+ K^0)$

 Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.72±0.05±0.01	5485	BALAGURA 08	BELL	10.6 $e^+ e^- \rightarrow D^{*+} K^0 X$

NODE=M121R8
NODE=M121R8

$\Gamma(D^+ \pi^- K^+)/\Gamma(D^*(2010)^+ K^0)$

 Γ_4/Γ_1

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.27±0.18±0.37	1264	BALAGURA 08	BELL	10.6 $e^+ e^- \rightarrow D^+ \pi^- K^+ X$

NODE=M121R9
NODE=M121R9

$\Gamma(D^+ K^0)/\Gamma(D^*(2010)^+ K^0)$

 Γ_6/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.40	90	ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
<0.43	90	ALBRECHT 89E	ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$

NODE=M121R1
NODE=M121R1

$\Gamma(D^0 K^+)/\Gamma(D^*(2007)^0 K^+)$

 Γ_7/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.12	90	ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$

NODE=M121R4
NODE=M121R4

$\Gamma(D_s^{*+} \gamma)/\Gamma_{total}$

 Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	ASRATYAN 88	HLBC	$\nu N \rightarrow D_s \gamma \gamma X$

NODE=M121R3
NODE=M121R3

$\Gamma(D_s^{*+} \gamma)/\Gamma(D^*(2007)^0 K^+)$

 Γ_8/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42	90	ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$

NODE=M121R5
NODE=M121R5

$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma_{total}$

 Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 06P	BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$

NODE=M121R7
NODE=M121R7 $D_{s1}(2536)^\pm$ REFERENCES

NODE=M121

AUSHEV 11	PR D83 051102	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=16505
LEES 11B	PR D83 072003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16773
ABAZOV 09G	PRL 102 051801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52652
CHEKANOV 09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
AUBERT 08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BALAGURA 08	PR D77 032001	V. Balagura <i>et al.</i>	(BELLE Collab.)	REFID=52133
AUBERT 06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
PDG 06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
HEISTER 02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
ACKERSTAFF 97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ASRATYAN 94	ZPHY C61 563	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)	REFID=43667
FRABETTI 94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
ALEXANDER 93	PL B303 377	J. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=43316
ALBRECHT 92R	PL B297 425	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=43179
AVERY 90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT 89E	PL B230 162	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40914
ASRATYAN 88	ZPHY C40 483	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)	REFID=40916

NODE=M148

$D_{s2}^*(2573)$

$$I(J^P) = 0(2^+)$$

J^P is natural, width and decay modes consistent with 2^+ .
AAIJ 14BJ confirms $J^P = 2^+$.

NODE=M148

$D_{s2}^*(2573)$ MASS


NODE=M148M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2569.1 ± 0.8 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below. [2571.9 ± 0.8 MeV OUR 2015 AVERAGE]		
2568.39 ± 0.29 ± 0.26		AAIJ	14AW LHC B	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
2569.4 ± 1.6 ± 0.5	82	AAIJ	11A LHC B	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$
2572.2 ± 0.3 ± 1.0		AUBERT, BE	06E BABR	$e^+ e^- \rightarrow DKX$
2574.5 ± 3.3 ± 1.6		ALBRECHT	96 ARG	$e^+ e^- \rightarrow D^0 K^+ X$
2573.2 $^{+1.7}_{-1.6}$ ± 0.9	217	KUBOTA	94 CLE2	$e^+ e^- \sim 10.5$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2570.0 ± 4.3	25	¹ EVDOKIMOV	04 SELX	600 $\Sigma^- A \rightarrow D^0 K^+ X$
2568.6 ± 3.2	64	² HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$

NODE=M148M
NEW

¹ Not independent of the mass difference below.
² Calculated using $m_{D^0} = 1864.5 \pm 0.5$ MeV and the mass difference below.

NODE=M148M; LINKAGE=EV
NODE=M148M; LINKAGE=HI



$m_{D_{s2}^*(2573)} - m_{D^0}$

NODE=M148DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
704 ± 3 ± 1	64	HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
705.4 ± 4.3	25	¹ EVDOKIMOV	04 SELX	600 $\Sigma^- A \rightarrow D^0 K^+ X$
¹ Systematic errors not estimated.				

NODE=M148DM

NODE=M148DM; LINKAGE=EV

$D_{s2}^*(2573)$ WIDTH

NODE=M148W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.9 ± 0.8 OUR AVERAGE				

NODE=M148W
NEW

[17 ± 4 MeV OUR 2015 AVERAGE Scale factor = 1.3]

16.9±0.5±0.6		AAIJ	14AW	LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
12.1±4.5±1.6	82	AAIJ	11A	LHCB	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$
27.1±0.6±5.6		AUBERT,BE	06E	BABR	$e^+ e^- \rightarrow DKX$
10.4±8.3±3.0		ALBRECHT	96	ARG	$e^+ e^- \rightarrow D^0 K^+ X$
16 $\begin{smallmatrix} +5 \\ -4 \end{smallmatrix}$ ±3	217	KUBOTA	94	CLE2	$e^+ e^- \sim 10.5 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 $\begin{smallmatrix} +9 \\ -6 \end{smallmatrix}$	25	¹ EVDOKIMOV	04	SELX	600 $\Sigma^- A \rightarrow D^0 K^+ X$
---	----	------------------------	----	------	--

¹Systematic errors not estimated.

NODE=M148W;LINKAGE=EV

 $D_{s2}^*(2573)^+$ DECAY MODES $D_{s2}^*(2573)^-$ modes are charge conjugates of the modes below.

NODE=M148215;NODE=M148

NODE=M148

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 K^+$	seen
Γ_2 $D^*(2007)^0 K^+$	not seen

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

 $D_{s2}^*(2573)^+$ BRANCHING RATIOS

NODE=M148220

$\Gamma(D^0 K^+)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	217	KUBOTA	94	CLE2	\pm $e^+ e^- \sim 10.5 \text{ GeV}$	

NODE=M148R2
NODE=M148R2

$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^0 K^+)$						Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.33	90	KUBOTA	94	CLE2	$+$ $e^+ e^- \sim 10.5 \text{ GeV}$	

NODE=M148R1
NODE=M148R1 **$D_{s2}^*(2573)$ REFERENCES**

NODE=M148

AAIJ	14AW	PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56105
AAIJ	14BJ	PRL 113 242002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=56258
AAIJ	11A	PL B698 14	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=16665
AUBERT,BE	06E	PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51512
EVDOKIMOV	04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>	(SELEX Collab.)	REFID=50337
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
ALBRECHT	96	ZPHY C69 405	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44631
KUBOTA	94	PRL 72 1972	Y. Kubota <i>et al.</i>	(CLEO Collab.)	REFID=43781

NODE=M182

 $D_{s1}^*(2700)^\pm$ $I(J^P) = 0(1^-)$ **$D_{s1}^*(2700)^+$ MASS**

NODE=M182M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M182M

2708.3 $\begin{smallmatrix} +4.0 \\ -3.4 \end{smallmatrix}$ OUR AVERAGE

NEW

[2709 ± 4 MeV OUR 2015 AVERAGE]

2699 $\begin{smallmatrix} +14 \\ -7 \end{smallmatrix}$		¹ LEES	15C	BABR	$B \rightarrow DD^0 K^+$
2709.2± 1.9± 4.5	52k	² AAIJ	12AU	LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
2710 ± 2 $\begin{smallmatrix} +12 \\ -7 \end{smallmatrix}$	10.4k	³ AUBERT	09AR	BABR	$e^+ e^- \rightarrow D^{(*)} KX$
2708 ± 9 $\begin{smallmatrix} +11 \\ -10 \end{smallmatrix}$	182	BRODZICKA	08	BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2694 ± 8 $\begin{smallmatrix} +13 \\ -3 \end{smallmatrix}$		LEES	15C	BABR	$B^0 \rightarrow D^- D^0 K^+$
2707 ± 8 ± 8		LEES	15C	BABR	$B^+ \rightarrow \bar{D}^0 D^0 K^+$
2688 ± 4 ± 3		⁴ AUBERT,BE	06E	BABR	10.6 $e^+ e^- \rightarrow DKX$

OCCUR=2

OCCUR=3

- ¹ From a combined analysis of $B^0 \rightarrow D^- D^0 K^+$ and $B^+ \rightarrow \bar{D}^0 D^0 K^+$.
² From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{s,J}^*(2860)^+$.
³ From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.
⁴ Superseded by AUBERT 09AR.

NODE=M182M;LINKAGE=B
 NODE=M182M;LINKAGE=AA
 NODE=M182M;LINKAGE=AB
 NODE=M182M;LINKAGE=AU

$D_{s1}^*(2700)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

122 \pm 11 \pm 8 OUR AVERAGE

[117 \pm 13 MeV OUR 2015 AVERAGE]

127 \pm 24 \pm 10		⁵ LEES	15C BABR	$B \rightarrow D D^0 K^+$
115.8 \pm 7.3 \pm 12.1	52k	⁶ AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
149 \pm 7 \pm 39 \pm 52	10.4k	⁷ AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$
108 \pm 23 \pm 36 \pm 31	182	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$
145 \pm 24 \pm 22 \pm 14		LEES	15C BABR	$B^0 \rightarrow D^- D^0 K^+$
113 \pm 21 \pm 20 \pm 16		LEES	15C BABR	$B^+ \rightarrow \bar{D}^0 D^0 K^+$
112 \pm 7 \pm 36		⁸ AUBERT,BE	06E BABR	10.6 $e^+ e^- \rightarrow DK X$

NODE=M182W
 NODE=M182W
 NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

OCCUR=2
 OCCUR=3

- ⁵ From a combined analysis of $B^0 \rightarrow D^- D^0 K^+$ and $B^+ \rightarrow \bar{D}^0 D^0 K^+$.
⁶ From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{s,J}^*(2860)^+$.
⁷ From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.
⁸ Superseded by AUBERT 09AR.

NODE=M182W;LINKAGE=A
 NODE=M182W;LINKAGE=AA
 NODE=M182W;LINKAGE=AB
 NODE=M182W;LINKAGE=AU

$D_{s1}^*(2700)^\pm$ DECAY MODES

Mode	DESIG
Γ_1 DK	DESIG=2
Γ_2 $D^0 K^+$	DESIG=1
Γ_3 $D^+ K_S^0$	DESIG=3
Γ_4 $D^* K$	DESIG=4
Γ_5 $D^{*0} K^+$	DESIG=5
Γ_6 $D^{*+} K_S^0$	DESIG=6

NODE=M182215;NODE=M182

$D_{s1}^*(2700)^\pm$ BRANCHING RATIOS

$\Gamma(D^* K)/\Gamma(DK)$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
----------------------------	------	-------------	------	---------	---------------------

0.91 \pm 0.13 \pm 0.12 10.4k ⁹ AUBERT 09AR BABR $e^+ e^- \rightarrow D^{(*)} K X$

⁹ From the average of the corresponding ratios with $D^{(*)0} K^+$ and $D^{(*)+} K_S^0$.

NODE=M182225
 NODE=M182R01
 NODE=M182R01

$\Gamma(D^{*0} K^+)/\Gamma(D^0 K^+)$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
--------------------------------------	------	-------------	------	---------	---------------------

0.88 \pm 0.14 \pm 0.14 7716 ¹⁰ AUBERT 09AR BABR $e^+ e^- \rightarrow D^{(*)} K X$

¹⁰ From the $D^{*0} K^+$ and $D^0 K^+$, where $D^{*0} \rightarrow D^0 \pi^0$.

NODE=M182R01;LINKAGE=AU
 NODE=M182R02
 NODE=M182R02

$\Gamma(D^{*+} K_S^0)/\Gamma(D^+ K_S^0)$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_3
--	------	-------------	------	---------	---------------------

1.14 \pm 0.39 \pm 0.23 2700 ¹¹ AUBERT 09AR BABR $e^+ e^- \rightarrow D^{(*)} K X$

¹¹ From the $D^{*+} K_S^0$ and $D^+ K_S^0$, where $D^{*+} \rightarrow D^+ \pi^0$.

NODE=M182R03
 NODE=M182R03

NODE=M182R03;LINKAGE=AU

$D_{s1}^*(2700)^\pm$ REFERENCES

LEES	15C	PR D91 052002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAJ	12AU	JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT	09AR	PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)
AUBERT,BE	06E	PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M182

REFID=56412
 REFID=54735
 REFID=53135
 REFID=52144
 REFID=51512

$D_{s1}^*(2860)^\pm$

$I(J^P) = 0(1^-)$

NODE=M196

OMITTED FROM SUMMARY TABLE

J^P consistent with 1^- from angular analysis of AAIJ 14AW. Observed by AUBERT, BE 06E and AUBERT 09AR in inclusive production of DK and D^*K in e^+e^- annihilation.

NODE=M196

 $D_{s1}^*(2860)^+$ MASS

NODE=M196M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2859 ±27 OUR AVERAGE		[2863.2 ^{+4.0} _{-2.6} MeV OUR 2015 AVERAGE]		
2859 ±12 ±24		¹ AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
2866.1 ± 1.0 ± 6.3	36k	^{2,3} AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
2862 ± 2 ± $\frac{5}{2}$	3122	^{3,4} AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)} K X$
2856.6 ± 1.5 ± 5.0		⁵ AUBERT, BE	06E BABR	$e^+e^- \rightarrow DKX$

NODE=M196M

NEW

¹ Separated from the spin-3 component $D_{s3}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M196M;LINKAGE=A

² From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{s,J}^*(2860)^+$.

NODE=M196M;LINKAGE=AA

³ Possible contribution from the $D_{s3}^*(2860)$ state.

NODE=M196M;LINKAGE=B

⁴ From simultaneous fits to the two DK mass spectra and to the total D^*K mass spectrum.

NODE=M196M;LINKAGE=AB

⁵ Superseded by AUBERT 09AR.

NODE=M196M;LINKAGE=AU

 $D_{s1}^*(2860)^+$ WIDTH

NODE=M196W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
159 ±80 OUR AVERAGE		[58 ± 11 MeV OUR 2015 AVERAGE Scale factor = 2.2]		
159 ±23 ±77		¹ AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
69.9 ± 3.2 ± 6.6	36k	^{2,3} AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
48 ± 3 ± 6	3122	^{3,4} AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)} K X$
47 ± 7 ± 10		⁵ AUBERT, BE	06E BABR	$e^+e^- \rightarrow DKX$

NODE=M196W

NEW

¹ Separated from the spin-3 component $D_{s3}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M196W;LINKAGE=A

² From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{s,J}^*(2860)^+$.

NODE=M196W;LINKAGE=AA

³ Possible contribution from the $D_{s3}^*(2860)$ state.

NODE=M196W;LINKAGE=B

⁴ From simultaneous fits to the two DK mass spectra and to the total D^*K mass spectrum.

NODE=M196W;LINKAGE=AB

⁵ Superseded by AUBERT 09AR.

NODE=M196W;LINKAGE=AU

 $D_{s1}^*(2860)^\pm$ DECAY MODES

NODE=M196215;NODE=M196

Mode	
Γ_1	DK
Γ_2	$D^0 K^+$
Γ_3	$D^+ K_S^0$
Γ_4	$D^* K$
Γ_5	$D^{*0} K^+$
Γ_6	$D^{*+} K_S^0$

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

$D_{s1}^*(2860)^\pm$ BRANCHING RATIOS **$\Gamma(D^*K)/\Gamma(DK)$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.10 \pm 0.15 \pm 0.19$	3122	¹ AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$

¹ From the average of the corresponding ratios with $D^{(*)0} K^+$ and $D^{(*)+} K_S^0$. **Γ_4/Γ_1**

NODE=M196225

NODE=M196R01
NODE=M196R01

NODE=M196R01;LINKAGE=AU

 $\Gamma(D^{*0} K^+)/\Gamma(D^0 K^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.04 \pm 0.17 \pm 0.20$	2241	¹ AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$

¹ From the $D^{*0} K^+$ and $D^0 K^+$, where $D^{*0} \rightarrow D^0 \pi^0$. **Γ_5/Γ_2** NODE=M196R02
NODE=M196R02

NODE=M196R02;LINKAGE=AU

 $\Gamma(D^{*+} K_S^0)/\Gamma(D^+ K_S^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.38 \pm 0.35 \pm 0.49$	881	¹ AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$

¹ From the $D^{*+} K_S^0$ and $D^+ K_S^0$, where $D^{*+} \rightarrow D^+ \pi^0$. **Γ_6/Γ_3** NODE=M196R03
NODE=M196R03

NODE=M196R03;LINKAGE=AU

 $D_{s1}^*(2860)^\pm$ REFERENCES

AAIJ	14AW PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	12AU JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06E PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M196

REFID=56105
REFID=54735
REFID=53135
REFID=51512

NODE=M226

 $D_{s3}^*(2860)^\pm$ $I(J^P) = 0(3^-)$

OMITTED FROM SUMMARY TABLE

 J^P consistent with 3^- from angular analysis of AAIJ 14AW.

NODE=M226

 $D_{s3}^*(2860)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2860.5 \pm 2.6 \pm 6.5$	¹ AAIJ	14AW LHCB	$B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$

¹ Separated from the spin-1 component $D_{s1}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M226M

NODE=M226M

NODE=M226M;LINKAGE=A

 $D_{s3}^*(2860)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$53 \pm 7 \pm 7$	¹ AAIJ	14AW LHCB	$B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$

¹ Separated from the spin-1 component $D_{s1}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M226W

NODE=M226W

NODE=M226W;LINKAGE=A

 $D_{s3}^*(2860)^\pm$ REFERENCES

AAIJ	14AW PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
------	---------------------	-----------------------	-------------------

NODE=M226

REFID=56105

$D_{sJ}(3040)^\pm$

$$I(J^P) = 0(?^?)$$

OMITTED FROM SUMMARY TABLE

Observed by AUBERT 09AR in inclusive production of $D^* K$ in $e^+ e^-$ annihilation.

NODE=M197

NODE=M197

 $D_{sJ}(3040)^+$ MASS

NODE=M197M

NODE=M197M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$3044 \pm 8_{-5}^{+30}$	AUBERT	09AR BABR	$e^+ e^- \rightarrow D^* K X$

 $D_{sJ}(3040)^+$ WIDTH

NODE=M197W

NODE=M197W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$239 \pm 35_{-42}^{+46}$	AUBERT	09AR BABR	$e^+ e^- \rightarrow D^* K X$

 $D_{sJ}(3040)^\pm$ DECAY MODES

NODE=M197215;NODE=M197

Mode

Γ_1	$D^* K$
Γ_2	$D^{*0} K^+$
Γ_3	$D^{*+} K_S^0$

DESIG=1

DESIG=2

DESIG=3

 $D_{sJ}(3040)^\pm$ REFERENCES

NODE=M197

AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
--------	--------------------	-------------------------	-----------------

REFID=53135

OTHER RELATED PAPERS

SUN	09 PR D80 074037	Z.-F. Sun, X. Lin
-----	------------------	-------------------

REFID=53128

BOTTOM MESONS

($B = \pm 1$)

NODE=MXXX045

$$B^+ = u\bar{b}, B^0 = d\bar{b}, \bar{B}^0 = \bar{d}b, B^- = \bar{u}b, \text{ similarly for } B^{*'}\text{'s}$$

NODE=MXXX045

NODE=M218

 $B_1(5721)^+$

$$I(J^P) = \frac{1}{2}(1^+) \text{ Status: } **$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M218

 $B_1(5721)^+$ MASS

NODE=M218M

OUR FIT uses $m_{B^{*0}}$ and $m_{B_1^+} - m_{B^{*0}}$ to determine $m_{B_1(5721)^+}$.

NODE=M218M

NODE=M218M

VALUE (MeV)	DOCUMENT ID
$5725.9_{-2.7}^{+2.5}$ OUR FIT	
[5726.8 $_{-4.0}^{+3.2}$ MeV OUR 2015 FIT]	

NEW

 $m_{B_1^+} - m_{B^{*0}}$

NODE=M218DM

NODE=M218DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$401.2_{-2.7}^{+2.4}$ OUR FIT				

NEW

[402.0^{+3.2}_{-4.0} MeV OUR 2015 FIT]**401.2^{+2.4}_{-2.7} OUR AVERAGE**[402^{+3.2}_{-4.2} MeV OUR 2015 AVERAGE]400.5 ± 1.8 ± 3.1 8K ¹ AAIJ 15AB LHCB $p\bar{p}$ at 7, 8 TeV402 ± 3 ⁺¹₋₃ ² AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV¹ AAIJ 15AB reports $[m_{B_1^+} - m_{B^0}] - (m_{B^{*0}} - m_{B^0}) - m_{\pi^+} = 260.9 \pm 1.8 \pm 3.1$ MeV which we adjust by the π^+ mass and assume $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV. The masses inside the square brackets were measured for each candidate event.² AALTONEN 14I reports $m_{B_1(5721)^+} - m_{B^{*0}} - m_{\pi^+} = 262 \pm 3 ⁺¹₋₃$ MeV which we adjusted by the π^+ mass.

NEW

NODE=M218DM;LINKAGE=A

NODE=M218DM;LINKAGE=AA

 $B_1(5721)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
31 ± 6 OUR AVERAGE				Error includes scale factor of 1.1. [49 ⁺¹² ₋₁₆ MeV OUR 2015 AVERAGE]

29.1 ± 3.6 ± 4.3 8K AAIJ 15AB LHCB $p\bar{p}$ at 7, 8 TeV49 ⁺¹²₋₁₀ ⁺²₋₁₃ AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

NODE=M218W

NODE=M218W

NEW

 $B_1(5721)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 B^{*0} \pi^+$	seen

NODE=M218215;NODE=M218

DESIG=1

 $B_1(5721)^+$ BRANCHING RATIOS

$\Gamma(B^{*0} \pi^+)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	8K	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV	
seen		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV	

NODE=M218220

NODE=M218R01
NODE=M218R01 **$B_1(5721)^+$ REFERENCES**AAIJ 15AB JHEP 1504 024 R. Aaij *et al.* (LHCb Collab.)
AALTONEN 14I PR D90 012013 T. Aaltonen *et al.* (CDF Collab.)

NODE=M218

REFID=56628
REFID=56029

NODE=M183

 $B_1(5721)^0$ $I(J^P) = \frac{1}{2}(1^+)$ Status: ***
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M183

 $B_1(5721)^0$ MASSOUR FIT uses mass differences measurements listed below to determine the mass $m_{B_1(5721)^0}$.

NODE=M183M

NODE=M183M

VALUE (MeV)	DOCUMENT ID
5726.0 ± 1.3 OUR FIT	Error includes scale factor of 1.2. [5724.9 ± 2.4 MeV OUR 2015 FIT Scale factor = 1.8]

NODE=M183M

NEW

 $m_{B_1^0} = m_{B^+}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
446.7 ± 1.3 OUR FIT	Error includes scale factor of 1.2. [445.6 ± 2.4 MeV OUR 2015 FIT Scale factor = 1.8]		

NODE=M183DM

NODE=M183DM

NEW

441.5 ± 2.4 ± 1.3 ¹ ABAZOV 07T D0 $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

446.2 ^{+1.9+1.0}_{-2.1-1.2} ¹ AALTONEN 09D CDF Repl. by AALTONEN 14I¹ Observed in $B_1^0 \rightarrow B^{*+} \pi^-$.

NODE=M183DM;LINKAGE=AA

$m_{B_1^0} - m_{B^{*+}}$

NODE=M183DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
401.4±1.2 OUR FIT				Error includes scale factor of 1.2. [400.1 ± 2.4 MeV OUR 2015 FIT Scale factor = 1.9]

NODE=M183DM2

NEW

402.8±1.1 OUR AVERAGE

NEW

[402.3 ± 1.5 MeV OUR 2015 AVERAGE]

403.4±0.7±1.5 35K ² AAIJ 15AB LHCB $p\bar{p}$ at 7, 8 TeV402.3±0.9^{+1.1}_{-1.2} ³ AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV² AAIJ 15AB reports $[m_{B_1^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 263.9 \pm 0.7 \pm 1.5$

NODE=M183DM2;LINKAGE=B

MeV which we adjust by the π^- mass and $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV. The masses inside the square brackets were measured for each candidate event.³ AALTONEN 14I reports $m_{B_1(5721)^0} - m_{B^{*+}} - m_{\pi^-} = 262.7 \pm 0.9^{+1.1}_{-1.2}$ MeV which we adjusted by the π^- mass.

NODE=M183DM2;LINKAGE=AA

 $B_1(5721)^0$ WIDTH

NODE=M183W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.5±3.4 OUR AVERAGE				Error includes scale factor of 1.1. [23 ± 5 MeV OUR 2015 AVERAGE]

NODE=M183W

NEW

30.1±1.5±3.5 35k AAIJ 15AB LHCB $p\bar{p}$ at 7, 8 TeV23 ±3 ±4 AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV $B_1(5721)^0$ DECAY MODES

NODE=M183215;NODE=M183

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^{*+} \pi^-$	dominant

DESIG=1

 $B_1(5721)^0$ BRANCHING RATIOS

NODE=M183220

$\Gamma(B^{*+} \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	
dominant	
dominant	

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	35K	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
dominant		AALTONEN	09D CDF	$p\bar{p}$ at 1.96 TeV
dominant		⁴ ABAZOV	07T D0	$p\bar{p}$ at 1.96 TeV

NODE=M183R01
NODE=M183R01⁴ Observed in $B_1^0 \rightarrow B^{*+} \pi^-$ with $B^{*+} \rightarrow B^+ \gamma$ and $B^+ \rightarrow J/\psi \pi^+$.

NODE=M183R01;LINKAGE=AB

 $B_1(5721)^0$ REFERENCES

NODE=M183

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56628
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=56029
AALTONEN	09D PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52700
ABAZOV	07T PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52014

$B_J^*(5732)$
or B^{**}

$I(J^P) = ?(??)$
 I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Signal can be interpreted as stemming from several narrow and broad resonances. Needs confirmation.

NODE=M151

NODE=M151

$B_J^*(5732)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5698 ± 8 OUR AVERAGE	Error includes scale factor of 1.2.			
5710 ± 20		¹ AFFOLDER	01F CDF	$p\bar{p}$ at 1.8 TeV
5695 ⁺¹⁷ ₋₁₉		² BARATE	98L ALEP	$e^+e^- \rightarrow Z$
5704 ± 4 ± 10	1944	³ BUSKULIC	96D ALEP	$E_{cm}^{ee} = 88-94$ GeV
5732 ± 5 ± 20	2157	ABREU	95B DLPH	$E_{cm}^{ee} = 88-94$ GeV
5681 ± 11	1738	AKERS	95E OPAL	$E_{cm}^{ee} = 88-94$ GeV

NODE=M151M

NODE=M151M

• • • We do not use the following data for averages, fits, limits, etc. • • •

5713 ± 2		⁴ ACCIARRI	99N L3	$e^+e^- \rightarrow Z$
¹ AFFOLDER 01F uses the reconstructed B meson through semileptonic decay channels. The fraction of light B mesons that are produced at $L=1$ B^{**} states is measured to be $0.28 \pm 0.06 \pm 0.03$.				
² BARATE 98L uses fully reconstructed B mesons to search for B^{**} production in the $B\pi^\pm$ system. In the framework of heavy quark symmetry (HQS), they also measured the mass of B_2^* to be $5739_{-11}^{+8+6}_{-4}$ MeV/ c^2 and the relative production rate of $B(b \rightarrow B_2^* \rightarrow B^{(*)}\pi)/B(b \rightarrow B_{u,d}) = (31 \pm 9_{-5}^{+6})\%$.				
³ Using $m_{B\pi} - m_B = 424 \pm 4 \pm 10$ MeV.				
⁴ ACCIARRI 99N uses inclusive reconstructed B mesons to search for B^{**} production in the $B^{(*)}\pi^\pm$ system. In the framework of HQET, they measured the mass of B_1^* and B_2^* to be $5670 \pm 10 \pm 13$ MeV and $5768 \pm 5 \pm 6$ with the $B(b \rightarrow B^{**}) = (32 \pm 3 \pm 6) \times 10^{-2}$. They also reported the evidence for the existence of an excited B -meson state or mixture of states in the region 5.9–6.0 GeV.				

NODE=M151M;LINKAGE=MF

NODE=M151M;LINKAGE=B

NODE=M151M;LINKAGE=A

NODE=M151M;LINKAGE=N

$B_J^*(5732)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
128 ± 18 OUR AVERAGE				
145 ± 28	2157	ABREU	95B DLPH	$E_{cm}^{ee} = 88-94$ GeV
116 ± 24	1738	AKERS	95E OPAL	$E_{cm}^{ee} = 88-94$ GeV

NODE=M151W

NODE=M151W

$B_J^*(5732)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^*\pi + B\pi$	dominant
Γ_2 $B^*\pi(X)$	[a] (85 ± 29) %

NODE=M151215;NODE=M151

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2

[a] X refers to decay modes with or without additional accompanying decay particles.

LINKAGE=151

$B_J^*(5732)$ BRANCHING RATIOS

X refers to decay modes with or without additional accompanying decay particles.

NODE=M151220

NODE=M151220

$\Gamma(B^*\pi(X))/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.85^{+0.26}_{-0.27} ± 0.12	ABBIENDI	02E OPAL	$e^+e^- \rightarrow Z$

NODE=M151R1
NODE=M151R1

$B_2^*(5732)$ REFERENCES

ABBIENDI	02E	EPJ C23 437	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	01F	PR D64 072002	T. Affolder <i>et al.</i>	(CDF Collab.)
ACCIARRI	99N	PL B465 323	M. Acciarri <i>et al.</i>	(L3 Collab.)
BARATE	98L	PL B425 215	R. Barate <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96D	ZPHY C69 393	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	95B	PL B345 598	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	95E	ZPHY C66 19	R. Akers <i>et al.</i>	(OPAL Collab.)

NODE=M151

REFID=48742
 REFID=48369
 REFID=47247
 REFID=46082
 REFID=44677
 REFID=44131
 REFID=44182

 $B_2^*(5747)^+$

$I(J^P) = \frac{1}{2}(2^+)$ Status: **
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M219

NODE=M219

 $B_2^*(5747)^+$ MASSOUR FIT uses m_{B^0} and $m_{B_2^{*+}} - m_{B^0}$ to determine $m_{B_2^*(5747)^+}$.

NODE=M219M

NODE=M219M

VALUE (MeV)	DOCUMENT ID
5737.2 ± 0.7 OUR FIT	
[5736.9 ^{+1.3} _{-1.6} MeV OUR 2015 FIT]	

NODE=M219M

NODE=M219M

NEW

 $m_{B_2^{*+}} - m_{B^0}$

NODE=M219DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
457.5 ± 0.7 OUR FIT				
[457.3 ^{+1.3} _{-1.6} MeV OUR 2015 FIT]				
457.5 ± 0.7 OUR AVERAGE				
[456.3 ^{+1.3} _{-1.6} MeV OUR 2015 AVERAGE]				
457.62 ± 0.72 ± 0.40	4K	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV
457.3 ± 1.3 ^{+0.3} _{-0.9}		² AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M219DM

NEW

NEW

¹ AAIJ 15AB reports $[m_{B_2^{*+}} - m_{B^0}] - m_{\pi^+} = 318.1 \pm 0.7 \pm 0.4$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14I reports $m_{B_2^*(5747)^+} - m_{B^0} - m_{\pi^+} = 317.7 \pm 1.2^{+0.3}_{-0.9}$ MeV which we adjusted by the π^+ mass.

NODE=M219DM;LINKAGE=B

NODE=M219DM;LINKAGE=A

 $B_2^*(5747)^+$ WIDTH

NODE=M219W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20 ± 5 OUR AVERAGE				Error includes scale factor of 2.2. [11 ± 5 MeV OUR 2015 AVERAGE]
23.6 ± 2.0 ± 2.1	4K	AAIJ	15AB LHCB	pp at 7, 8 TeV
11 ⁺⁴ ₋₃ ⁺³ ₋₄		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M219W

NEW

 $B_2^*(5747)^+$ DECAY MODES

NODE=M219215;NODE=M219

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^0 \pi^+$	seen
Γ_2 $B^{*0} \pi^+$	seen

DESIG=1

DESIG=2

 $B_2^*(5747)^+$ BRANCHING RATIOS

NODE=M219220

$\Gamma(B^0 \pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	4K	AAIJ	15AB LHCB	pp at 7, 8 TeV
seen		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
$\Gamma(B^{*0} \pi^+)/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	4k	AAIJ	15AB LHCB	pp at 7, 8 TeV

NODE=M219R01

NODE=M219R01

NODE=M219R02

NODE=M219R02

$\Gamma(B^{*0}\pi^+)/\Gamma(B^0\pi^+)$ Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.5±0.8	4k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV

NODE=M219R03
 NODE=M219R03

 $B_2^*(5747)^+$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)

NODE=M219

REFID=56628
 REFID=56029

NODE=M184

 $B_2^*(5747)^0$

$I(J^P) = \frac{1}{2}(2^+)$ Status: ***
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M184

 $B_2^*(5747)^0$ MASS

OUR FIT uses m_{B^+} , $m_{B_1^0} - m_{B^+}$, and $m_{B_2^{*0}} - m_{B_1^0}$ to determine $m_{B_2^*(5747)^0}$. The -0.659 correlation between statistical uncertainties of $m_{B_1^0} - m_{B^+}$ and $m_{B_2^{*0}} - m_{B_1^0}$ measurements reported by ABAZOV 07T is taken into account.

NODE=M184M

NODE=M184M

VALUE (MeV)	DOCUMENT ID
5739.5±0.7 OUR FIT	Error includes scale factor of 1.4. [5739 ± 5 MeV OUR 2015 FIT Scale factor = 4.0]

NODE=M184M

NEW

$$m_{B_2^{*0}} = m_{B_1^0}$$

NODE=M184DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
13.5±1.4 OUR FIT	Error includes scale factor of 1.3. [14 ± 6 MeV OUR 2015 FIT Scale factor = 3.4]		

NODE=M184DM

NEW

26.2±3.1±0.9	¹ ABAZOV	07T D0	$p\bar{p}$ at 1.96 TeV
14.9 ^{+2.2+1.2} _{-2.5-1.4}	¹ AALTONEN	09D CDF	Repl. by AALTONEN 14I

¹ Observed in $B_2^{*0} \rightarrow B^{*+}\pi^-$ and $B_2^{*0} \rightarrow B^+\pi^-$.

$$m_{B_2^{*0}} - m_{B^+}$$

NODE=M184DM;LINKAGE=AB

NODE=M184DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
460.2 ±0.6 OUR FIT		Error includes scale factor of 1.4. [460 ± 5 MeV OUR 2015 FIT Scale factor = 4.0]		

NODE=M184DM2

NEW

459.9 ±0.8 OUR AVERAGE	Error includes scale factor of 1.8. [457.5 ± 1.5 MeV OUR 2015 AVERAGE]
-------------------------------	--

NEW

460.18±0.37±0.33	17K	² AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
457.5 ±1.2 ^{+0.8} _{-0.9}		³ AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

² AAIJ 15AB reports $[m_{B_2^{*0}} - m_{B^+}] - m_{\pi^-} = 320.6 \pm 0.4 \pm 0.3$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event.

³ AALTONEN 14I reports $m_{B_2^*(5747)^0} - m_{B^+} - m_{\pi^-} = 317.9 \pm 1.2^{+0.8}_{-0.9}$ MeV which we adjusted by the π^- mass.

NODE=M184DM2;LINKAGE=A

NODE=M184DM2;LINKAGE=AA

 $B_2^*(5747)^0$ WIDTH

NODE=M184W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
24.2±1.7 OUR AVERAGE		[22 ± 5 MeV OUR 2015 AVERAGE]		

NODE=M184W

NEW

24.5±1.0±1.5	17K	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
22 ⁺³⁺⁴ ₋₂₋₅		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

22.7 ^{+3.8+3.2} _{-3.2-10.2}		AALTONEN	09D CDF	Repl. by AALTONEN 14I
---	--	----------	---------	-----------------------

$B_2^*(5747)^0$ DECAY MODES

NODE=M184215;NODE=M184

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^+ \pi^-$	dominant
Γ_2 $B^{*+} \pi^-$	dominant

DESIG=1
DESIG=2 $B_2^*(5747)^0$ BRANCHING RATIOS

NODE=M184220

$\Gamma(B^+ \pi^-)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	17K	AAIJ	15AB LHCB	pp at 7, 8 TeV	
dominant		AALTONEN	09D CDF	$\rho\bar{p}$ at 1.96 TeV	
dominant		ABAZOV	07T D0	$\rho\bar{p}$ at 1.96 TeV	

NODE=M184R01
NODE=M184R01

$\Gamma(B^{*+} \pi^-)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	17K	AAIJ	15AB LHCB	pp at 7, 8 TeV	
dominant		AALTONEN	09D CDF	$\rho\bar{p}$ at 1.96 TeV	
dominant		ABAZOV	07T D0	$\rho\bar{p}$ at 1.96 TeV	

NODE=M184R02
NODE=M184R02

$\Gamma(B^{*+} \pi^-)/\Gamma(B^+ \pi^-)$					Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.82 ± 0.28 OUR AVERAGE					
[1.1 ± 0.5 OUR 2015 AVERAGE]					
0.71 ± 0.14 ± 0.30	17K	AAIJ	15AB LHCB	pp at 7, 8 TeV	
1.10 ± 0.42 ± 0.31		⁴ ABAZOV	07T D0	$\rho\bar{p}$ at 1.96 TeV	

NODE=M184R03
NODE=M184R03
NEW

⁴ Converted from measured ratio of $R = B(B_2^{*0} \rightarrow B^{*+} \pi^-) / B(B_2^{*0} \rightarrow B^{(*)+} \pi^-)$
= 0.475 ± 0.095 ± 0.069.

NODE=M184R03;LINKAGE=AB

 $B_2^*(5747)^0$ REFERENCES

NODE=M184

AAIJ	15AB	JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56628
AALTONEN	14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=56029
AALTONEN	09D	PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52700
ABAZOV	07T	PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52014

NODE=M224

 $B_J(5840)^+$
 $I(J^P) = \frac{1}{2}(?^?)$ Status: **
I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M224

 $B_J(5840)^+$ MASS

NODE=M224M

OUR FIT uses m_{B^0} and $m_{B_J(5840)^+} - m_{B^0}$ to determine $m_{B_J(5840)^+}$.

NODE=M224M

VALUE (MeV)	DOCUMENT ID
5851 ± 19 OUR FIT	

NODE=M224M

 $m_{B_J(5840)^+} - m_{B^0}$

NODE=M224DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
571 ± 19 OUR FIT				
571 ± 13 ± 14	7k	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV

NODE=M224DM

• • • We do not use the following data for averages, fits, limits, etc. • • •

595 ± 26 ± 14	7k	² AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------	----	-------------------	-----------	------------------

OCCUR=2

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 431 \pm 13 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224DM;LINKAGE=A

² AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 455 \pm 26 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224DM;LINKAGE=B

$m_{B_J(5840)^+} - m_{B^{*0}}$

NODE=M224DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M224DM2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$565 \pm 15 \pm 14$	7k	³ AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------------	----	-------------------	-----------	------------------

³ AAIJ 15AB reports $[m_{B^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 425 \pm 15 \pm 14$

NODE=M224DM2;LINKAGE=A

MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

 $B_J(5840)^+$ WIDTH

NODE=M224W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M224W

$224 \pm 24 \pm 80$	7k	⁴ AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------------------------------	----	-------------------	-----------	------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$215 \pm 27 \pm 80$	7k	⁵ AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------------	----	-------------------	-----------	------------------

OCCUR=2

$229 \pm 27 \pm 80$	7k	⁶ AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------------	----	-------------------	-----------	------------------

OCCUR=3

⁴ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224W;LINKAGE=A

⁵ Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224W;LINKAGE=B

⁶ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224W;LINKAGE=C

 $B_J(5840)^+$ DECAY MODES

NODE=M224215;NODE=M224

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^{*0} \pi^+$	seen
Γ_2 $B^0 \pi^+$	possibly seen

DESIG=1

DESIG=2

 $B_J(5840)^+$ BRANCHING RATIOS

NODE=M224220

$\Gamma(B^{*0} \pi^+)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
--	-------	------	-------------	------	---------	-------------------

NODE=M224R01
NODE=M224R01

seen	7k	AAIJ	15AB LHCB	pp at 7, 8 TeV	
-------------	----	------	-----------	------------------	--

$\Gamma(B^0 \pi^+)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
---	-------	------	-------------	------	---------	-------------------

NODE=M224R02
NODE=M224R02

possibly seen	7k	⁷ AAIJ	15AB LHCB	pp at 7, 8 TeV	
----------------------	----	-------------------	-----------	------------------	--

⁷ A $B \pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^* \pi$ is allowed.

NODE=M224R02;LINKAGE=A

 $B_J(5840)^+$ REFERENCES

NODE=M224

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	--------------------	-----------------------	----------------

REFID=56628

$B_J(5840)^0$
 $I(J^P) = \frac{1}{2}(?)^?$ Status: **
 I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M225

NODE=M225

NODE=M225M

NODE=M225M

NODE=M225M

 $B_J(5840)^0$ MASSOUR FIT uses m_{B^+} and $m_{B_J(5840)^0} - m_{B^+}$ to determine $m_{B_J(5840)^0}$.

VALUE (MeV)

DOCUMENT ID

584 ± 9 OUR FIT **$m_{B_J(5840)^0} - m_{B^+}$**

NODE=M225DM

NODE=M225DM

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

584 ± 9 OUR FIT**584 ± 5 ± 7**

12k

1 AAIJ

15AB LHCB pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

610 ± 22 ± 7

12k

2 AAIJ

15AB LHCB pp at 7, 8 TeV

OCCUR=2

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M225DM;LINKAGE=A

² AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 471 \pm 22 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M225DM;LINKAGE=B

 $m_{B_J(5840)^0} - m_{B^{*+}}$

NODE=M225DM2

NODE=M225DM2

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

584 ± 5 ± 7

12k

3 AAIJ

15AB LHCB pp at 7, 8 TeV

³ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M225DM2;LINKAGE=A

 $B_J(5840)^0$ WIDTH

NODE=M225W

NODE=M225W

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

127 ± 17 ± 34

12k

4 AAIJ

15AB LHCB pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

107 ± 20 ± 34

12k

5 AAIJ

15AB LHCB pp at 7, 8 TeV

OCCUR=2

119 ± 17 ± 34

12k

6 AAIJ

15AB LHCB pp at 7, 8 TeV

OCCUR=3

⁴ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M225W;LINKAGE=A

⁵ Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M225W;LINKAGE=B

⁶ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M225W;LINKAGE=C

 $B_J(5840)^0$ DECAY MODES

NODE=M225215;NODE=M225

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^{*+} \pi^-$	seen
Γ_2 $B^+ \pi^-$	possibly seen

DESIG=1

DESIG=2

$B_J(5840)^0$ BRANCHING RATIOS $\Gamma(B^{*+}\pi^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12k	AAIJ	15AB LHCB	pp at 7, 8 TeV

 $\Gamma(B^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	⁷ AAIJ	15AB LHCB	pp at 7, 8 TeV

⁷ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.

 $B_J(5840)^0$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	--------------------	-----------------------	----------------

 $B_J(5970)^+$
 $I(J^P) = \frac{1}{2}(?^?)$ Status: **
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

 $B_J(5970)^+$ MASS

OUR FIT uses m_{B^0} and $m_{B_J(5970)^+} - m_{B^0}$ to determine $m_{B_J(5970)^+}$.

VALUE (MeV)	DOCUMENT ID
5964 ± 5 OUR FIT	
[5961 ± 13 MeV OUR 2015 FIT]	

 $m_{B_J(5970)^+} - m_{B^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
685 ± 5 OUR FIT				
[681 ± 13 MeV OUR 2015 FIT]				
685 ± 5 OUR AVERAGE				
[681 ± 13 MeV OUR 2015 AVERAGE]				
685.3 ± 4.1 ± 2.5	2K	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV
681 ± 5 ± 12		² AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
686.8 ± 4.5 ± 2.5	2K	³ AAIJ	15AB LHCB	pp at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 545.8 \pm 4.1 \pm 2.5$ MeV which we adjust by

the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AALTONEN 14I reports $m_{B_J(5970)^+} - m_{B^0} - m_{\pi^+} = 541 \pm 5 \pm 12$ MeV which we adjusted by the π^+ mass.

³ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 547 \pm 5 \pm 3$ MeV which we adjust by

the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$$m_{B_J(5970)^+} = m_{B^0}$$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
686.0 ± 4.0 ± 2.5	2k	⁴ AAIJ	15AB LHCB	pp at 7, 8 TeV
⁴ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 547 \pm 4 \pm 3$ MeV which				
we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.				

 $B_J(5970)^+$ WIDTH

NODE=M225220

NODE=M225R01
NODE=M225R01NODE=M225R02
NODE=M225R02

NODE=M225R02;LINKAGE=A

NODE=M225

REFID=56628

NODE=M220

NODE=M220

NODE=M220M

NODE=M220M

NODE=M220M

NEW

NODE=M220DM

NODE=M220DM

NEW

NEW

OCCUR=2

NODE=M220DM;LINKAGE=B

NODE=M220DM;LINKAGE=A

NODE=M220DM;LINKAGE=C

NODE=M220DM2

NODE=M220DM2

NODE=M220DM2;LINKAGE=A

NODE=M220W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62±20 OUR AVERAGE				
[60 ⁺⁵⁰ ₋₄₀ MeV OUR 2015 AVERAGE]				
63±15±17	2K	⁵ AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
60 ⁺³⁰ ₋₂₀ ±40		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
61±14±17	2K	⁶ AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
61±15±17	2K	⁷ AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
⁵ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.				
⁶ Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.				
⁷ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.				

NODE=M220W
NEW

$B_J(5970)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^0\pi^+$	possibly seen
Γ_2 $B^{*0}\pi^+$	seen

OCCUR=2
OCCUR=3
NODE=M220W;LINKAGE=A
NODE=M220W;LINKAGE=B
NODE=M220W;LINKAGE=C

NODE=M220215;NODE=M220

DESIG=1
DESIG=2

$B_J(5970)^+$ BRANCHING RATIOS

$\Gamma(B^0\pi^+)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
possibly seen [seen OUR 2015 BEST LIMIT]					
possibly seen	2K	⁸ AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV	
possibly seen		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV	
⁸ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.					
$\Gamma(B^{*0}\pi^+)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	2k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV	
seen		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV	

NODE=M220220

NODE=M220R01
NODE=M220R01

NODE=M220R01;LINKAGE=A

NODE=M220R02
NODE=M220R02

$B_J(5970)^+$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)

NODE=M220

REFID=56628
REFID=56029

$B_J(5970)^0$
 $I(J^P) = \frac{1}{2}(?)^?$ Status: **
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M221

NODE=M221

 $B_J(5970)^0$ MASSOUR FIT uses m_{B^+} and $m_{B_J(5970)^0} - m_{B^+}$ to determine $m_{B_J(5970)^0}$.

NODE=M221M

NODE=M221M

NODE=M221M

NEW

VALUE (MeV)	DOCUMENT ID
5971 ± 5 OUR FIT	
[5977 ± 13 MeV OUR 2015 FIT]	

 $m_{B_J(5970)^0} - m_{B^+}$

NODE=M221DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
691 ± 5 OUR FIT				
[698 ± 13 MeV OUR 2015 FIT]				
691 ± 5 OUR AVERAGE				
[698 ± 13 MeV OUR 2015 AVERAGE]				

NODE=M221DM

NEW

NEW

689.9 ± 2.9 ± 5.1	10K	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV
698 ± 5 ± 12		² AALTONEN	14l CDF	$p\bar{p}$ at 1.96 TeV

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

714.3 ± 6.4 ± 5.1	10K	³ AAIJ	15AB LHCB	pp at 7, 8 TeV
-------------------	-----	-------------------	-----------	------------------

OCCUR=2

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 550.4 \pm 2.9 \pm 5.1$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M221DM;LINKAGE=B

² AALTONEN 14l reports $m_{B_J(5970)^0} - m_{B^+} - m_{\pi^-} = 558 \pm 5 \pm 12$ MeV which we adjusted by the π^- mass.

NODE=M221DM;LINKAGE=A

³ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 575 \pm 6 \pm 5$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M221DM;LINKAGE=C

 $m_{B_J(5970)^0} - m_{B^{*+}}$

NODE=M221DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
691.6 ± 3.7 ± 5.1	10k	⁴ AAIJ	15AB LHCB	pp at 7, 8 TeV
⁴ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 552 \pm 4 \pm 5$ MeV				
which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.				

NODE=M221DM2

NODE=M221DM2;LINKAGE=A

 $B_J(5970)^0$ WIDTH

NODE=M221W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
81 ± 12 OUR AVERAGE				
[70 ± 40 MeV OUR 2015 AVERAGE]				
82 ± 8 ± 9	10K	⁵ AAIJ	15AB LHCB	pp at 7, 8 TeV
70 ⁺³⁰ ₋₂₀ ± 30		AALTONEN	14l CDF	$p\bar{p}$ at 1.96 TeV

NODE=M221W

NEW

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

56 ± 7 ± 9	10K	⁶ AAIJ	15AB LHCB	pp at 7, 8 TeV
82 ± 10 ± 9	10K	⁷ AAIJ	15AB LHCB	pp at 7, 8 TeV

OCCUR=2

OCCUR=3

⁵ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M221W;LINKAGE=A

⁶ Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M221W;LINKAGE=B

⁷ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M221W;LINKAGE=C

$B_J(5970)^0$ DECAY MODES

NODE=M221215;NODE=M221

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^+\pi^-$	possibly seen
Γ_2 $B^{*+}\pi^-$	seen

DESIG=1

DESIG=2

 $B_J(5970)^0$ BRANCHING RATIOS

NODE=M221220

 $\Gamma(B^+\pi^-)/\Gamma_{\text{total}}$ **Γ_1/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

possibly seen [seen OUR 2015 BEST LIMIT]possibly seen 10K ⁸ AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV**possibly seen** AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV⁸ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.

NODE=M221R01

NODE=M221R01

OCCUR=2

NODE=M221R01;LINKAGE=A

 $\Gamma(B^{*+}\pi^-)/\Gamma_{\text{total}}$ **Γ_2/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 10K AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV**seen** AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

NODE=M221R02

NODE=M221R02

OCCUR=2

 $B_J(5970)^0$ REFERENCES

NODE=M221

AAIJ 15AB JHEP 1504 024
AALTONEN 14I PR D90 012013R. Aaij *et al.*
T. Aaltonen *et al.*(LHCb Collab.)
(CDF Collab.)

REFID=56628

REFID=56029

BOTTOM, STRANGE MESONS

$(B = \pm 1, S = \mp 1)$

NODE=MXXX046

$$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b, \text{ similarly for } B_s^{*+}$$

NODE=MXXX046

 $B_{s1}(5830)^0$ $I(J^P) = 0(1^+)$ Status: ***
 I, J, P need confirmation.

NODE=M187

Quantum numbers shown are quark-model predictions.

NODE=M187

 $B_{s1}(5830)^0$ MASS

NODE=M187M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

 5828.63 ± 0.27 OUR FIT[5828.78 \pm 0.35 MeV OUR 2015 FIT Scale factor = 1.2] **$5828.40 \pm 0.04 \pm 0.41$** ¹ AAIJ 130 LHCb $p\bar{p}$ at 7 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

5829.4 \pm 0.7 ² AALTONEN 08K CDF Repl. by AALTONEN 14I¹ Uses $B_{s1}(5830)^0 \rightarrow B^{*+}K^-$ decay.² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$,
 $J/\psi \rightarrow \mu^+\mu^-$ or $B^+ \rightarrow \bar{D}^0\pi^+$, $\bar{D}^0 \rightarrow K^+\pi^-$.

NODE=M187M

NEW

NODE=M187M;LINKAGE=AI

NODE=M187M;LINKAGE=AA

$$m_{B_{s1}^0} - m_{B^{*+}}$$

NODE=M187DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

 503.98 ± 0.18 OUR FIT[503.95 \pm 0.23 MeV OUR 2015 FIT Scale factor = 1.3] **$504.03 \pm 0.12 \pm 0.15$** ³ AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

504.41 \pm 0.21 \pm 0.14 ⁴ AALTONEN 08K CDF Repl. by AALTONEN 14I

NODE=M187DM

NEW

³ AALTONEN 14I reports $m_{B_{s1}(5830)^0} - m_{B^{*+}} - m_{K^-} = 10.35 \pm 0.12 \pm 0.15$ MeV

which we adjusted by the K^- mass.

⁴ Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.

NODE=M187DM;LINKAGE=AL

NODE=M187DM;LINKAGE=AA

$B_{s1}(5830)^0$ WIDTH

NODE=M187W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
0.5±0.3±0.3	AALTONEN	14I	CDF $\rho\bar{p}$ at 1.96 TeV

NODE=M187W

$B_{s1}(5830)^0$ DECAY MODES

NODE=M187215;NODE=M187

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^{*+} K^-$	dominant

DESIG=1

$B_{s1}(5830)^0$ BRANCHING RATIOS

NODE=M187220

$\Gamma(B^{*+} K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
dominant	AALTONEN	08K	CDF $\rho\bar{p}$ at 1.96 TeV	

NODE=M187R01
NODE=M187R01

$B_{s1}(5830)^0$ REFERENCES

NODE=M187

AALTONEN	14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=56029
AAIJ	130	PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54968
AALTONEN	08K	PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52235

NODE=M186

$B_{s2}^*(5840)^0$

$I(J^P) = 0(2^+)$ Status: ***
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M186

$B_{s2}^*(5840)^0$ MASS

NODE=M186M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5839.84±0.18 OUR FIT			Error includes scale factor of 1.1. [5839.83 ± 0.19 MeV OUR 2015 FIT Scale factor = 1.2]

NODE=M186M
NEW

5839.98±0.20 OUR AVERAGE

5839.99±0.05±0.20	AAIJ	130	LHCB $p\bar{p}$ at 7 TeV
5839.6 ±1.1 ±0.7	¹ ABAZOV	08E	D0 $\rho\bar{p}$ at 1.96 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

5839.7 ±0.7	² AALTONEN	08K	CDF Repl. by AALTONEN 14I
-------------	-----------------------	-----	---------------------------

¹ Observed in $B_{s2}^{*0} \rightarrow B^+ K^-$. Measured production rate of B_{s2}^{*0} relative to B^+ to be $(1.15 \pm 0.23 \pm 0.13)\%$.

NODE=M186M;LINKAGE=AB

² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.

NODE=M186M;LINKAGE=AA

$m_{B_{s2}^{*0}} - m_{B_{s1}^0}$

NODE=M186DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10.5±0.6	³ AALTONEN	08K	CDF Repl. by AALTONEN 14I

NODE=M186DM

³ Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.

NODE=M186DM;LINKAGE=AA

$m_{B_{s2}^{*0}} - m_{B^+}$

NODE=M186DM2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
560.53±0.18 OUR FIT			Error includes scale factor of 1.1. [560.54 ± 0.19 MeV OUR 2015 FIT Scale factor = 1.2]

NODE=M186DM2
NEW

560.41±0.13±0.14	⁴ AALTONEN	14I	CDF $\rho\bar{p}$ at 1.96 TeV
-------------------------	-----------------------	-----	-------------------------------

⁴AALTONEN 14I reports $m_{B_{s2}^*(5840)^0} - m_{B^+} - m_{K^-} = 66.73 \pm 0.13 \pm 0.14$ MeV which we adjusted by the K^- mass.

NODE=M186DM2;LINKAGE=AL

 $B_{s2}^*(5840)^0$ WIDTH

NODE=M186W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1.47±0.33 OUR AVERAGE			
1.4 ±0.4 ±0.2	AALTONEN 14I	CDF	$p\bar{p}$ at 1.96 TeV
1.56±0.13±0.47	⁵ AAIJ 130	LHCB	pp at 7 TeV

⁵Uses $B_{s2}^*(5840)^0 \rightarrow B^{*+} K^-$ decays.

NODE=M186W

NODE=M186W;LINKAGE=AI

 $B_{s2}^*(5840)^0$ DECAY MODES

NODE=M186215;NODE=M186

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^+ K^-$	dominant
Γ_2 $B^{*+} K^-$	

DESIG=1

DESIG=2

 $B_{s2}^*(5840)^0$ BRANCHING RATIOS

NODE=M186220

$\Gamma(B^+ K^-)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
dominant		AALTONEN 08K	CDF	$p\bar{p}$ at 1.96 TeV	
dominant		⁶ ABAZOV 08E	D0	$p\bar{p}$ at 1.96 TeV	

⁶Measured production rate of B_{s2}^{*0} relative to B^+ to be $(1.15 \pm 0.23 \pm 0.13)\%$.

NODE=M186R01
NODE=M186R01

NODE=M186R01;LINKAGE=AB

$\Gamma(B^{*+} K^-)/\Gamma(B^+ K^-)$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.093±0.013±0.012		AAIJ 130	LHCB	pp at 7 TeV	

NODE=M186R02
NODE=M186R02 **$B_{s2}^*(5840)^0$ REFERENCES**

NODE=M186

AALTONEN 14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ 130	PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN 08K	PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV 08E	PRL 100 082002	V.M. Abazov <i>et al.</i>	(D0 Collab.)

REFID=56029
REFID=54968
REFID=52235
REFID=52232

NODE=M153

 $B_{sJ}^*(5850)$

$I(J^P) = ?(??)$
I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Signal can be interpreted as coming from $\bar{b}s$ states. Needs confirmation.

NODE=M153

 $B_{sJ}^*(5850)$ MASS

NODE=M153M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5853±15	141	AKERS 95E	OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

NODE=M153M

 $B_{sJ}^*(5850)$ WIDTH

NODE=M153W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
47±22	141	AKERS 95E	OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

NODE=M153W

 $B_{sJ}^*(5850)$ REFERENCES

NODE=M153

AKERS 95E	ZPHY C66 19	R. Akers <i>et al.</i>	(OPAL Collab.)
-----------	-------------	------------------------	----------------

REFID=44182

BOTTOM, CHARMED MESONS ($B = C = \pm 1$)

$$B_c^+ = c\bar{b}, B_c^- = \bar{c}b, \text{ similarly for } B_c^{*'}\text{'s}$$

$B_c(2S)^\pm$

$$I(J^P) = 0(0^-)$$

OMITTED FROM SUMMARY TABLE

Quantum numbers neither measured nor confirmed.

$B_c(2S)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$6842 \pm 4 \pm 5$	57	¹ AAD	14AQ ATLS	pp at 7, 8 TeV

¹ Observed in the decay mode $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$ ($B_c^+ \rightarrow J/\psi \pi^+$) with 5.2 standard deviations significance.

$B_c(2S)^\pm$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B_c^+ \pi^+ \pi^-$	seen

$B_c(2S)^\pm$ BRANCHING RATIOS

$\Gamma(B_c^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	57	¹ AAD	14AQ ATLS	pp at 7, 8 TeV	

¹ Observed with 5.2 standard deviations significance.

$B_c(2S)^\pm$ REFERENCES

AAD 14AQ PRL 113 212004 G. Aad *et al.* (ATLAS Collab.)

A REVIEW GOES HERE – Check our WWW List of Reviews

$c\bar{c}$ MESONS

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=MXXX049

NODE=MXXX049

NODE=M217

NODE=M217

NODE=M217M

NODE=M217M

NODE=M217M;LINKAGE=AA

NODE=M217215;NODE=M217

DESIG=1

NODE=M217225

NODE=M217R01
NODE=M217R01

NODE=M217R01;LINKAGE=AA

NODE=M217

REFID=56117

NODE=M209

NODE=MXXX025

NODE=M826

$\eta_c(1S)$

$$J^{PC} = 0^+(0^-+)$$

NODE=M026

$\eta_c(1S)$ MASS

NODE=M026M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2983.4 ± 0.5	OUR AVERAGE	Error includes scale factor of 1.2. [2983.6 ± 0.6 MeV OUR 2015 AVERAGE Scale factor = 1.2]		
2982.2 ± 1.5 ± 0.1	2.0k	1 AAIJ	15BI LHCB	$pp \rightarrow \eta_c(1S)X$
2983.5 ± 1.4 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 1.6 \\ 3.6 \end{smallmatrix}$		2 ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
2979.8 ± 0.8 ± 3.5	4.5k	3,4 LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
2984.1 ± 1.1 ± 2.1	900	3,4,5 LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
2984.3 ± 0.6 ± 0.6		6,7 ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$
2984.49 ± 1.16 ± 0.52	832	3 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma$ hadrons
2982.7 ± 1.8 ± 2.2	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
2984.5 ± 0.8 ± 3.1	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
2985.4 ± 1.5 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 0.5 \\ 2.0 \end{smallmatrix}$	920	7 VINOKUROVA	11 BELL	$B^\pm \rightarrow K_S^\pm(K_S^0K^\pm\pi^\mp)$
2982.2 ± 0.4 ± 1.6	14k	8 LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
2985.8 ± 1.5 ± 3.1	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K(*) \rightarrow K\bar{K}\pi K(*)$
2986.1 ± 1.0 ± 2.5	7.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow$ hadrons
2970 ± 5 ± 6	501	9 ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
2971 ± 3 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}$	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
2974 ± 7 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 2 \\ 1 \end{smallmatrix}$	20	WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
2981.8 ± 1.3 ± 1.5	592	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0K^\pm\pi^\mp$
2984.1 ± 2.1 ± 1.0	190	10 AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2982.5 ± 0.4 ± 1.4	12k	11 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
2982.2 ± 0.6		12 MITCHELL	09 CLEO	$e^+e^- \rightarrow \gamma X$
2982 ± 5	270	13 AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X c\bar{c}$
2982.5 ± 1.1 ± 0.9	2.5k	14 AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
2977.5 ± 1.0 ± 1.2		12,15 BAI	03 BES	$J/\psi \rightarrow \gamma\eta_c$
2979.6 ± 2.3 ± 1.6	180	16 FANG	03 BELL	$B \rightarrow \eta_c K$
2976.3 ± 2.3 ± 1.2		12,17 BAI	00F BES	$J/\psi, \psi(2S) \rightarrow \gamma\eta_c$
2976.6 ± 2.9 ± 1.3	140	12,18 BAI	00F BES	$J/\psi \rightarrow \gamma\eta_c$
2980.4 ± 2.3 ± 0.6		19 BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$
2975.8 ± 3.9 ± 1.2		18 BAI	99B BES	Sup. by BAI 00F
2999 ± 8	25	ABREU	98O DLPH	$e^+e^- \rightarrow e^+e^- +$ hadrons
2988.3 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 3.3 \\ 3.1 \end{smallmatrix}$		ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
2974.4 ± 1.9		12,20 BISELLO	91 DM2	$J/\psi \rightarrow \eta_c\gamma$
2969 ± 4 ± 4	80	12 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^-K^+K^-$
2956 ± 12 ± 12		12 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^-K_S^0K_L^0$
2982.6 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ $\begin{smallmatrix} 2.7 \\ 2.3 \end{smallmatrix}$	12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
2980.2 ± 1.6		12,20 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$
2984 ± 2.3 ± 4.0		12 GAISER	86 CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
2976 ± 8		12,21 BALTRUSAIT..84	MRK3	$J/\psi \rightarrow 2\phi\gamma$
2982 ± 8	18	22 HIMEL	80B MRK2	e^+e^-
2980 ± 9		22 PARTRIDGE	80B CBAL	e^+e^-

NODE=M026M

NEW

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

- ¹ AAIJ 15BI reports $m_{J/\psi} - m_{\eta_c(1S)} = 114.7 \pm 1.5 \pm 0.1$ MeV from a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. We have used current value of $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to arrive at the quoted $m_{\eta_c(1S)}$ result.
- ² Taking into account an asymmetric photon lineshape.
- ³ With floating width.
- ⁴ Ignoring possible interference with the non-resonant 0^- amplitude.
- ⁵ Using both, $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.
- ⁶ From a simultaneous fit to six decay modes of the η_c .
- ⁷ Accounts for interference with non-resonant continuum.
- ⁸ Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.
- ⁹ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.
- ¹⁰ Using mass of $\psi(2S) = 3686.00$ MeV.
- ¹¹ Not independent from the measurements reported by LEES 10.
- ¹² MITCHELL 09 observes a significant asymmetry in the lineshapes of $\psi(2S) \rightarrow \gamma\eta_c$ and $J/\psi \rightarrow \gamma\eta_c$ transitions. If ignored, this asymmetry could lead to significant bias whenever the mass and width are measured in $\psi(2S)$ or J/ψ radiative decays.
- ¹³ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.
- ¹⁴ Superseded by LEES 10.
- ¹⁵ From a simultaneous fit of five decay modes of the η_c .
- ¹⁶ Superseded by VINOKUROVA 11.
- ¹⁷ Weighted average of the $\psi(2S)$ and $J/\psi(1S)$ samples. Using an η_c width of 13.2 MeV.
- ¹⁸ Average of several decay modes. Using an η_c width of 13.2 MeV.
- ¹⁹ Superseded by ASNER 04.
- ²⁰ Average of several decay modes.
- ²¹ $\eta_c \rightarrow \phi\phi$.
- ²² Mass adjusted by us to correspond to $J/\psi(1S)$ mass = 3097 MeV.

NODE=M026M;LINKAGE=D

NODE=M026M;LINKAGE=E
 NODE=M026M;LINKAGE=AL
 NODE=M026M;LINKAGE=LS
 NODE=M026M;LINKAGE=EL
 NODE=M026M;LINKAGE=BL
 NODE=M026M;LINKAGE=VA
 NODE=M026M;LINKAGE=LE
 NODE=M026M;LINKAGE=EB
 NODE=M026M;LINKAGE=BG
 NODE=M026M;LINKAGE=DE
 NODE=M026M;LINKAGE=MI

NODE=M026M;LINKAGE=AU
 NODE=M026M;LINKAGE=UB
 NODE=M026M;LINKAGE=AK
 NODE=M026M;LINKAGE=FA
 NODE=M026M;LINKAGE=KZ
 NODE=M026M;LINKAGE=C1
 NODE=M026M;LINKAGE=NN
 NODE=M026M;LINKAGE=A
 NODE=M026M;LINKAGE=B
 NODE=M026M;LINKAGE=M

 $\eta_c(1S)$ WIDTH

NODE=M026W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.8 ± 0.8 OUR FIT				
31.9 ± 1.0 OUR AVERAGE				Error includes scale factor of 1.2. [32.0 ± 1.0 MeV OUR 2015 AVERAGE Scale factor = 1.2]
27.2 ± 3.1 ^{+5.4} _{-2.6}		¹ ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
25.2 ± 2.6 ± 2.4	4.5k	^{2,3} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
34.8 ± 3.1 ± 4.0	900	^{2,3,4} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
32.0 ± 1.2 ± 1.0		^{5,6} ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$
36.4 ± 3.2 ± 1.7	832	² ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma$ hadrons
37.8 ⁺ _{-5.3} ± 5.8 ± 3.1	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
36.2 ± 2.8 ± 3.0	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
35.1 ± 3.1 ^{+1.0} _{-1.6}	920	⁶ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm\pi^\mp)$
31.7 ± 1.2 ± 0.8	14k	⁷ LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0 K^\pm\pi^\mp$
36.3 ⁺ _{-3.6} ± 3.7 ± 4.4	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K(*) \rightarrow K\bar{K}\pi K(*)$
28.1 ± 3.2 ± 2.2	7.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow$ hadrons
48 ⁺ ₋₇ ± 8 ± 5	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
40 ± 19 ± 5	20	WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
24.8 ± 3.4 ± 3.5	592	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm\pi^\mp$
20.4 ⁺ _{-6.7} ± 7.7 ± 2.0	190	AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
23.9 ⁺ _{-7.1} ± 12.6		ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
32.1 ± 1.1 ± 1.3	12k	⁸ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm\pi^\mp$
34.3 ± 2.3 ± 0.9	2.5k	⁹ AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
17.0 ± 3.7 ± 7.4		¹⁰ BAI	03 BES	$J/\psi \rightarrow \gamma\eta_c$
29 ± 8 ± 6	180	¹¹ FANG	03 BELL	$B \rightarrow \eta_c K$
11.0 ± 8.1 ± 4.1		¹² BAI	00F BES	$J/\psi \rightarrow \gamma\eta_c$ and $\psi(2S) \rightarrow \gamma\eta_c$
27.0 ± 5.8 ± 1.4		¹³ BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$
7.0 ⁺ _{-7.0} ± 7.5 ± 7.0	12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
10.1 ⁺ _{-8.2} ± 33.0 ± 8.2	23	¹⁴ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma p\bar{p}$
11.5 ± 4.5		GAISER	86 CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
< 40 90% CL	18	HIMEL	80B MRK2	e^+e^-
< 20 90% CL		PARTRIDGE	80B CBAL	e^+e^-

NODE=M026W

NEW

OCCUR=2

OCCUR=2

OCCUR=2

- 1 Taking into account an asymmetric photon lineshape.
- 2 With floating mass.
- 3 Ignoring possible interference with the non-resonant 0^- amplitude.
- 4 Using both, $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.
- 5 From a simultaneous fit to six decay modes of the η_c .
- 6 Accounts for interference with non-resonant continuum.
- 7 Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.
- 8 Not independent from the measurements reported by LEES 10.
- 9 Superseded by LEES 10.
- 10 From a simultaneous fit of five decay modes of the η_c .
- 11 Superseded by VINOKUROVA 11.
- 12 From a fit to the 4-prong invariant mass in $\psi(2S) \rightarrow \gamma\eta_c$ and $J/\psi(1S) \rightarrow \gamma\eta_c$ decays.
- 13 Superseded by ASNER 04.
- 14 Positive and negative errors correspond to 90% confidence level.

NODE=M026W;LINKAGE=A
 NODE=M026W;LINKAGE=AL
 NODE=M026W;LINKAGE=LS
 NODE=M026W;LINKAGE=EL
 NODE=M026W;LINKAGE=BL
 NODE=M026W;LINKAGE=VA
 NODE=M026W;LINKAGE=LE
 NODE=M026W;LINKAGE=DE
 NODE=M026W;LINKAGE=UB
 NODE=M026W;LINKAGE=AK
 NODE=M026W;LINKAGE=FA
 NODE=M026W;LINKAGE=KZ
 NODE=M026W;LINKAGE=NN
 NODE=M026W;LINKAGE=L

$\eta_c(1S)$ DECAY MODES

NODE=M026215;NODE=M026

Mode	Fraction (Γ_i/Γ)	Confidence level
Decays involving hadronic resonances		
Γ_1 $\eta'(958)\pi\pi$	(4.1 \pm 1.7) %	
Γ_2 $\rho\rho$	(1.8 \pm 0.5) %	
Γ_3 $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(2.0 \pm 0.7) %	
Γ_4 $K^*(892) K^*(892)$	(7.0 \pm 1.3) $\times 10^{-3}$	
Γ_5 $K^*(892)^0 K^*(892)^0 \pi^+ \pi^-$	(1.1 \pm 0.5) %	
Γ_6 $\phi K^+ K^-$	(2.9 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\phi\phi$	(1.75 \pm 0.20) $\times 10^{-3}$	
Γ_8 $\phi 2(\pi^+ \pi^-)$	< 4 $\times 10^{-3}$	90%
Γ_9 $a_0(980)\pi$	< 2 %	90%
Γ_{10} $a_2(1320)\pi$	< 2 %	90%
Γ_{11} $K^*(892) \bar{K} + \text{c.c.}$	< 1.28 %	90%
Γ_{12} $f_2(1270)\eta$	< 1.1 %	90%
Γ_{13} $\omega\omega$	< 3.1 $\times 10^{-3}$	90%
Γ_{14} $\omega\phi$	< 1.7 $\times 10^{-3}$	90%
Γ_{15} $f_2(1270) f_2(1270)$	(9.8 \pm 2.5) $\times 10^{-3}$	
Γ_{16} $f_2(1270) f_2'(1525)$	(9.7 \pm 3.2) $\times 10^{-3}$	
Γ_{17} $f_0(980)\eta$	seen	
Γ_{18} $f_0(1500)\eta$	seen	
Γ_{19} $f_0(2200)\eta$	seen	
Γ_{20} $a_0(980)\pi$	seen	
Γ_{21} $a_0(1320)\pi$	seen	
Γ_{22} $a_0(1450)\pi$	seen	
Γ_{23} $a_0(1950)\pi$	seen	
Γ_{24} $a_2(1950)\pi$	not seen	
Γ_{25} $K_0^*(1430) \bar{K}$	seen	
Γ_{26} $K_2^*(1430) \bar{K}$	seen	
Γ_{27} $K_0^*(1950) \bar{K}$	seen	
Decays into stable hadrons		
Γ_{28} $K \bar{K} \pi$	(7.3 \pm 0.5) %	
Γ_{29} $K K \eta$	(1.35 \pm 0.16) %	
Γ_{30} $\eta \pi^+ \pi^-$	(1.7 \pm 0.5) %	
Γ_{31} $\eta 2(\pi^+ \pi^-)$	(4.4 \pm 1.3) %	
Γ_{32} $K^+ K^- \pi^+ \pi^-$	(6.9 \pm 1.1) $\times 10^{-3}$	
Γ_{33} $K^+ K^- \pi^+ \pi^- \pi^0$	(3.5 \pm 0.6) %	
Γ_{34} $K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}$	(5.6 \pm 1.5) %	
Γ_{35} $K^+ K^- 2(\pi^+ \pi^-)$	(7.5 \pm 2.4) $\times 10^{-3}$	
Γ_{36} $2(K^+ K^-)$	(1.46 \pm 0.30) $\times 10^{-3}$	
Γ_{37} $\pi^+ \pi^- \pi^0 \pi^0$	(4.7 \pm 1.0) %	
Γ_{38} $2(\pi^+ \pi^-)$	(9.7 \pm 1.2) $\times 10^{-3}$	
Γ_{39} $2(\pi^+ \pi^- \pi^0)$	(17.4 \pm 3.3) %	
Γ_{40} $3(\pi^+ \pi^-)$	(1.8 \pm 0.4) %	
Γ_{41} $p \bar{p}$	(1.50 \pm 0.16) $\times 10^{-3}$	
Γ_{42} $p \bar{p} \pi^0$	(3.6 \pm 1.3) $\times 10^{-3}$	
Γ_{43} $\Lambda \bar{\Lambda}$	(1.09 \pm 0.24) $\times 10^{-3}$	
Γ_{44} $\Sigma^+ \bar{\Sigma}^-$	(2.1 \pm 0.6) $\times 10^{-3}$	
Γ_{45} $\Xi^- \bar{\Xi}^+$	(8.9 \pm 2.7) $\times 10^{-4}$	
Γ_{46} $\pi^+ \pi^- p \bar{p}$	(5.3 \pm 1.8) $\times 10^{-3}$	

NODE=M026;CLUMP=A
 DESIG=24
 DESIG=19
 DESIG=26
 DESIG=18
 DESIG=57
 DESIG=28
 DESIG=17
 DESIG=58
 DESIG=21
 DESIG=22
 DESIG=40
 DESIG=23
 DESIG=20
 DESIG=47
 DESIG=46
 DESIG=59
 DESIG=70
 DESIG=71
 DESIG=72
 DESIG=73
 DESIG=74
 DESIG=75
 DESIG=79
 DESIG=80
 DESIG=76
 DESIG=77
 DESIG=78

NODE=M026;CLUMP=B
 DESIG=14
 DESIG=25
 DESIG=16
 DESIG=61
 DESIG=15
 DESIG=60
 DESIG=62
 DESIG=55
 DESIG=27
 DESIG=63
 DESIG=11
 DESIG=64
 DESIG=56
 DESIG=12
 DESIG=65
 DESIG=45
 DESIG=66
 DESIG=67
 DESIG=13

Radiative decays

$\Gamma_{47} \quad \gamma\gamma \quad (1.59 \pm 0.13) \times 10^{-4}$

NODE=M026;CLUMP=C
DESIG=31

**Charge conjugation (C), Parity (P),
Lepton family number (LF) violating modes**

NODE=M026;CLUMP=D

Γ_{48}	$\pi^+ \pi^-$	$P, CP < 1.1$	$\times 10^{-4}$	90%
Γ_{49}	$\pi^0 \pi^0$	$P, CP < 4$	$\times 10^{-5}$	90%
Γ_{50}	$K^+ K^-$	$P, CP < 6$	$\times 10^{-4}$	90%
Γ_{51}	$K_S^0 K_S^0$	$P, CP < 3.1$	$\times 10^{-4}$	90%

DESIG=51
DESIG=52
DESIG=53
DESIG=54

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 8 combinations of partial widths obtained from integrated cross section, and 19 branching ratios uses 85 measurements and one constraint to determine 13 parameters. The overall fit has a $\chi^2 = 118.3$ for 73 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	18									
x_{15}	3	6								
x_{28}	22	41	7							
x_{29}	12	22	4	54						
x_{32}	11	21	4	25	13					
x_{36}	9	16	3	25	14	10				
x_{38}	14	25	5	30	16	16	12			
x_{41}	14	26	5	36	19	16	13	20		
x_{43}	3	6	1	9	5	4	3	5	25	
x_{47}	-29	-54	-10	-66	-35	-34	-27	-41	-46	-11
Γ	-2	-3	-1	-4	-2	-2	-1	-2	7	2
	x_4	x_7	x_{15}	x_{28}	x_{29}	x_{32}	x_{36}	x_{38}	x_{41}	x_{43}

Γ	-28
	x_{47}

Mode	Rate (MeV)	
$\Gamma_4 \quad K^*(892) \bar{K}^*(892)$	0.22 ± 0.04	DESIG=18
$\Gamma_7 \quad \phi\phi$	0.056 ± 0.007	DESIG=17
$\Gamma_{15} \quad f_2(1270) \bar{f}_2(1270)$	0.31 ± 0.08	DESIG=46
$\Gamma_{28} \quad K \bar{K} \pi$	2.31 ± 0.16	DESIG=14
$\Gamma_{29} \quad K \bar{K} \eta$	0.43 ± 0.05	DESIG=25
$\Gamma_{32} \quad K^+ K^- \pi^+ \pi^-$	0.219 ± 0.034	DESIG=15
$\Gamma_{36} \quad 2(K^+ K^-)$	0.046 ± 0.010	DESIG=27
$\Gamma_{38} \quad 2(\pi^+ \pi^-)$	0.31 ± 0.04	DESIG=11
$\Gamma_{41} \quad p \bar{p}$	0.048 ± 0.005	DESIG=12
$\Gamma_{43} \quad \Lambda \bar{\Lambda}$	0.034 ± 0.008	DESIG=45
$\Gamma_{47} \quad \gamma\gamma$	0.0051 ± 0.0004	DESIG=31

$\eta_c(1S)$ PARTIAL WIDTHS

NODE=M026217

$\Gamma(\gamma\gamma)$

Γ_{47}

VALUE (keV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ COMMENT _____

NODE=M026W1
NODE=M026W1

5.1 ± 0.4 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

5.8 ± 1.1	486	¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
5.2 ± 1.2	273 ± 43	^{2,3} AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
5.5 ± 1.2 ± 1.8	157 ± 33	⁴ KUO	05 BELL	$\gamma\gamma \rightarrow p\bar{p}$
7.4 ± 0.4 ± 2.3		⁵ ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
13.9 ± 2.0 ± 3.0	41	⁶ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \eta_c$
3.8 ⁺ _{-1.0} 1.1 ⁺ _{-1.0}	190	⁷ AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
7.6 ± 0.8 ± 2.3		^{5,8} BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$
6.9 ± 1.7 ± 2.1	76	⁹ ACCIARRI	99T L3	$e^+e^- \rightarrow e^+e^-\eta_c$
27 ± 16 ± 10	5	⁵ SHIRAI	98 AMY	58 e^+e^-
6.7 ⁺ _{-1.7} 2.4 ± 2.3		⁴ ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
11.3 ± 4.2		¹⁰ ALBRECHT	94H ARG	$e^+e^- \rightarrow e^+e^-\eta_c$
8.0 ± 2.3 ± 2.4	17	¹¹ ADRIANI	93N L3	$e^+e^- \rightarrow e^+e^-\eta_c$
5.9 ⁺ _{-1.8} 2.1 ± 1.9		⁷ CHEN	90B CLEO	$e^+e^- \rightarrow e^+e^-\eta_c$
6.4 ⁺ _{-3.4} 5.0		¹² AIHARA	88D TPC	$e^+e^- \rightarrow e^+e^-X$
4.3 ⁺ _{-3.7} 3.4 ± 2.4		⁴ BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
28 ± 15		^{5,13} BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$

¹ Assuming there is no interference with the non-resonant background.

² Calculated by us using $\Gamma(\eta_c \rightarrow K\bar{K}\pi) \times \Gamma(\eta_c \rightarrow \gamma\gamma) / \Gamma = 0.44 \pm 0.05$ keV from PDG 06 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

³ Systematic errors not evaluated.

⁴ Normalized to $B(\eta_c \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$.

⁵ Normalized to $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$.

⁶ Average of $K_S^0 K^\pm \pi^\mp$, $\pi^+ \pi^- K^+ K^-$, and $2(K^+ K^-)$ decay modes.

⁷ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

⁸ Superseded by ASNER 04.

⁹ Normalized to the sum of 9 branching ratios.

¹⁰ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow \phi\phi)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

¹¹ Superseded by ACCIARRI 99T.

¹² Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow 2K^+ 2K^-)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

¹³ Re-evaluated by AIHARA 88D.

NODE=M026W1;LINKAGE=ZH
NODE=M026W1;LINKAGE=AU

NODE=M026W1;LINKAGE=NS
NODE=M026W1;LINKAGE=N3
NODE=M026W1;LINKAGE=N2
NODE=M026W;LINKAGE=FF
NODE=M026W1;LINKAGE=N6

NODE=M026W1;LINKAGE=NN
NODE=M026W1;LINKAGE=N1
NODE=M026W1;LINKAGE=N5

NODE=M026W1;LINKAGE=WD
NODE=M026W1;LINKAGE=N4

NODE=M026W1;LINKAGE=A

$\eta_c(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M026220

$\Gamma(\eta'(958)\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_{47}/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
75.8 ⁺ _{-6.2} 6.3 ± 8.4	486	¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	NODE=M026G10 NODE=M026G10

¹ Assuming there is no interference with the non-resonant background.

NODE=M026G10;LINKAGE=ZH

$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_{47}/\Gamma$	
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<39	90	< 1556	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+ \pi^-)$	NODE=M026G09 NODE=M026G09

$\Gamma(K^*(892)\bar{K}^*(892)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_{47}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
35 ± 6 OUR FIT					NODE=M026G08 NODE=M026G08 NEW
32.4 ± 4.2 ± 5.8	882 ± 115	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+ \pi^- K^+ K^-$	

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_{47}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
8.9 ± 0.8 OUR FIT					NODE=M026G07 NODE=M026G07
7.75 ± 0.66 ± 0.62	386 ± 31	¹ LIU	12B BELL	$\gamma\gamma \rightarrow 2(K^+ K^-)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.8 ± 1.2 ± 1.3 132 ± 23 UEHARA 08 BELL $\gamma\gamma \rightarrow 2(K^+ K^-)$

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

NODE=M026G07;LINKAGE=L1

$\Gamma(\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_{47}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.67±2.86±0.96	85 ± 29	¹ LIU	12B BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M026G03
NODE=M026G03

NODE=M026G03;LINKAGE=LI

 $\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_{47}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.49	90	¹ LIU	12B BELL	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.49 90 ¹ LIU 12B BELL $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹ Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M026G04
NODE=M026G04

NODE=M026G04;LINKAGE=LI

 $\Gamma(f_2(1270)f_2(1270)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_{47}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
50±13 OUR FIT				
69±17±12	3182±766	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

50±13 OUR FIT

69±17±12 3182±766 UEHARA 08 BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

NODE=M026G19
NODE=M026G19

 $\Gamma(f_2(1270)f_2'(1525)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_{47}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
49±9±13	1128±206	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

49±9±13 1128±206 UEHARA 08 BELL $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

NODE=M026G20
NODE=M026G20

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{28}\Gamma_{47}/\Gamma$

VALUE (keV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
0.368±0.021 OUR FIT				
0.407±0.027 OUR AVERAGE				Error includes scale factor of 1.2.
0.374±0.009±0.031	14k	¹ LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
0.407±0.022±0.028		^{2,3} ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0K^\pm\pi^\mp$
0.60 ±0.12 ±0.09	41	^{3,4} ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
1.47 ±0.87 ±0.27		³ SHIRAI	98 AMY	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$
0.84 ±0.21		³ ALBRECHT	94H ARG	$\gamma\gamma \rightarrow K^\pm K_S^0\pi^\mp$
0.60 ^{+0.23} _{-0.20}		³ CHEN	90B CLEO	$\gamma\gamma \rightarrow \eta_c K^\pm K_S^0\pi^\mp$
1.06 ±0.41 ±0.27	11	³ BRAUNSCH...	89 TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$
1.5 ^{+0.60} _{-0.45} ±0.3	7	³ BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$

0.368±0.021 OUR FIT

0.407±0.027 OUR AVERAGE Error includes scale factor of 1.2.

0.374±0.009±0.031 14k ¹ LEES 10 BABR $10.6 e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$

0.407±0.022±0.028 ^{2,3} ASNER 04 CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0K^\pm\pi^\mp$

0.60 ±0.12 ±0.09 41 ^{3,4} ABDALLAH 03J DLPH $\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$

1.47 ±0.87 ±0.27 ³ SHIRAI 98 AMY $\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$

0.84 ±0.21 ³ ALBRECHT 94H ARG $\gamma\gamma \rightarrow K^\pm K_S^0\pi^\mp$

0.60 ^{+0.23}_{-0.20} ³ CHEN 90B CLEO $\gamma\gamma \rightarrow \eta_c K^\pm K_S^0\pi^\mp$

1.06 ±0.41 ±0.27 11 ³ BRAUNSCH... 89 TASS $\gamma\gamma \rightarrow K\bar{K}\pi$

1.5 ^{+0.60}_{-0.45} ±0.3 7 ³ BERGER 86 PLUT $\gamma\gamma \rightarrow K\bar{K}\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.386±0.008±0.021 12k ⁵ DEL-AMO-SA...11M BABR $\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$

0.418±0.044±0.022 ^{3,6} BRANDENB... 00B CLE2 $\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$

<0.63 95 ³ BEHREND 89 CELL $\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$

<4.4 95 ALTHOFF 85B TASS $\gamma\gamma \rightarrow K\bar{K}\pi$

¹ From the corrected and unfolded mass spectrum.

² Calculated by us from the value reported in ASNER 04 that assumes $B(\eta_c \rightarrow K\bar{K}\pi) = 5.5 \pm 1.7\%$

³ We have multiplied $K^\pm K_S^0\pi^\mp$ measurement by 3 to obtain $K\bar{K}\pi$.

⁴ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow K_S^0K^\pm\pi^\mp) = (1.5 \pm 0.4)\%$.

⁵ Not independent from the measurements reported by LEES 10.

⁶ Superseded by ASNER 04.

NODE=M026G14;LINKAGE=LE
NODE=M026G14;LINKAGE=AA

NODE=M026G14;LINKAGE=C
NODE=M026G;LINKAGE=BB

NODE=M026G14;LINKAGE=DE
NODE=M026G14;LINKAGE=NN

 $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{32}\Gamma_{47}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35 ± 5 OUR FIT				
27 ± 6 OUR AVERAGE				
25.7 ± 3.2 ± 4.9	2019 ± 248	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$
280 ± 100 ± 60	42	¹ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$
170 ± 80 ± 20	13.9 ± 6.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

35 ± 5 OUR FIT

27 ± 6 OUR AVERAGE

25.7 ± 3.2 ± 4.9 2019 ± 248 UEHARA 08 BELL $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

280 ± 100 ± 60 42 ¹ ABDALLAH 03J DLPH $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

170 ± 80 ± 20 13.9 ± 6.6 ALBRECHT 94H ARG $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

¹ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow \pi^+\pi^-K^+K^-) = (2.0 \pm 0.7)\%$.

NODE=M026G15
NODE=M026G15

NODE=M026G;LINKAGE=CC

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{33}\Gamma_{47}/\Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.190 ± 0.006 ± 0.028	11k	¹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
-----------------------	-----	------------------------------	------	--

¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M026G02
NODE=M026G02

NODE=M026G02;LINKAGE=DE

$$\Gamma(2(K^+K^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_{47}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

7.4 ± 1.5 OUR FIT

5.8 ± 1.9 OUR AVERAGE

5.6 ± 1.1 ± 1.6	216 ± 42	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$
350 ± 90 ± 60	46	¹ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow 2(K^+K^-)$
231 ± 90 ± 23	9.1 ± 3.3	² ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(K^+K^-)$

¹ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow 2(K^+K^-)) = (2.1 \pm 1.2)\%$.

² Includes all topological modes except $\eta_c \rightarrow \phi\phi$.

NODE=M026G27
NODE=M026G27

NODE=M026G;LINKAGE=DD

NODE=M026G;LINKAGE=EE

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{38}\Gamma_{47}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

49 ± 6 OUR FIT

42 ± 6 OUR AVERAGE

40.7 ± 3.7 ± 5.3	5381 ± 492	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$
180 ± 70 ± 20	21.4 ± 8.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

NODE=M026G11
NODE=M026G11

$$\Gamma(p\bar{p}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{41}\Gamma_{47}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

7.6 ± 0.7 OUR FIT

7.20 ± 1.53^{+0.67}_{-0.75}	157 ± 33	¹ KUO	05 BELL	$\gamma\gamma \rightarrow p\bar{p}$
--	----------	------------------	---------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.6 ^{+1.3} _{-1.1} ± 0.4	190	¹ AMBROGIANI	03 E835	$\bar{p}p \rightarrow \gamma\gamma$
---	-----	-------------------------	---------	-------------------------------------

8.1 ^{+2.9} _{-2.0}		¹ ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
-------------------------------------	--	------------------------	----------	-------------------------------------

¹ Not independent from the $\Gamma_{\gamma\gamma}$ reported by the same experiment.

NODE=M026G01
NODE=M026G01

NODE=M026G01;LINKAGE=GG

$$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_{47}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.6	90	¹ UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
----------------	----	---------------------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.29	90	² UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
-------	----	---------------------	---------	--

¹ Taking into account interference with the non-resonant continuum.

² Neglecting interference with the non-resonant continuum.

NODE=M026G05
NODE=M026G05

OCCUR=2

NODE=M026G05;LINKAGE=U1

NODE=M026G05;LINKAGE=U2

$\eta_c(1S)$ BRANCHING RATIOS

NODE=M026225

HADRONIC DECAYS

NODE=M026305

$$\Gamma(\eta'(958)\pi\pi)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.041 ± 0.017	14	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$
----------------------	----	-----------------------------	------	-----------------------------------

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R14
NODE=M026R14

NODE=M026R14;LINKAGE=E

$$\Gamma(\rho\rho)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

18 ± 5 OUR AVERAGE

12.6 ± 3.8 ± 5.1	72	¹ ABLIKIM	05L BES2	$J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
26.0 ± 2.4 ± 8.8	113	¹ BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^0\rho^0$
23.6 ± 10.6 ± 8.2	32	¹ BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^+\rho^-$

NODE=M026R9
NODE=M026R9

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$
-----	----	-----------------------------	------	-----------------------------------

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

NODE=M026R9;LINKAGE=E

$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.02±0.007	63	^{1,2} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

NODE=M026R16
 NODE=M026R16

¹ BALTRUSAITIS 86 has an error according to Partridge.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R;LINKAGE=03
 NODE=M026R16;LINKAGE=E

 $\Gamma(K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

70±13 OUR FIT**91±26 OUR AVERAGE**

108±25±44	60	¹ ABLIKIM	05L BES2	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
82±28±27	14	¹ BISELLO	91 DM2	$e^+ e^- \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
90±50	9	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

NODE=M026R8
 NODE=M026R8

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

NODE=M026R8;LINKAGE=E

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

113±47±25

	45	¹ ABLIKIM	06A BES2	$J/\psi \rightarrow K^{*0} \bar{K}^{*0} \pi^+ \pi^- \gamma$
--	----	----------------------	----------	---

¹ ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.91 \pm 0.64 \pm 0.48) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R25
 NODE=M026R25

NODE=M026R25;LINKAGE=AB

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.9^{+0.9}_{-0.8}±1.1	14.1 ^{+4.4} _{-3.7}	¹ HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
--	--------------------------------------	--------------------	---------	--------------------------------------

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R21
 NODE=M026R21

NODE=M026R;LINKAGE=BB

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

17.5± 2.0 OUR FIT**30 ± 5 OUR AVERAGE**

25.3± 5.1± 9.1	72	¹ ABLIKIM	05L BES2	$J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$
26 ± 9	357 ± 64	¹ BAI	04 BES	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
31 ± 7 ± 10	19	¹ BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
30 ⁺¹⁸ ₋₁₂ ± 10	5	¹ BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
74 ± 18 ± 24	80	¹ BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
67 ± 21 ± 24		¹ BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

NODE=M026R7
 NODE=M026R7

OCCUR=2

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

18 ⁺⁸ ₋₆ ± 7	7.0 ^{+3.0} _{-2.3}	² HUANG	03 BELL	$B^+ \rightarrow (\phi\phi) K^+$
------------------------------------	-------------------------------------	--------------------	---------	----------------------------------

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

² Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R;LINKAGE=E

NODE=M026R7;LINKAGE=BB

 $\Gamma(\phi\phi)/\Gamma(K \bar{K} \pi)$ Γ_7/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0240±0.0026 OUR FIT**0.044 ^{+0.012}_{-0.010} OUR AVERAGE**

0.055 ± 0.014 ± 0.005		AUBERT,B	04B BABR	$B^\pm \rightarrow K^\pm \eta_c$
0.032 ^{+0.014} _{-0.010} ± 0.009	7	¹ HUANG	03 BELL	$B^\pm \rightarrow K^\pm \phi\phi$

NODE=M026R39
 NODE=M026R39

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R39;LINKAGE=BB

$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<40	90	¹ ABLIKIM 06A	BES2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
¹ ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow \phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.603 \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.				

NODE=M026R26
NODE=M026R26

NODE=M026R26;LINKAGE=AB

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	90	^{1,2} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.				
² We are assuming $B(a_0(980) \rightarrow \eta \pi) > 0.5$.				

NODE=M026R11
NODE=M026R11NODE=M026R11;LINKAGE=E
NODE=M026R11;LINKAGE=F $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.				

NODE=M026R12
NODE=M026R12

NODE=M026R12;LINKAGE=E

 $\Gamma(K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0128	90	BISELLO 91	DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
<0.0132	90	¹ BISELLO 91	DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.				

NODE=M026R17
NODE=M026R17

OCCUR=2

NODE=M026R17;LINKAGE=E

 $\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.011	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.				

NODE=M026R13
NODE=M026R13

NODE=M026R13;LINKAGE=E

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0031	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0063	90	¹ ABLIKIM 05L	BES2	$J/\psi \rightarrow \pi^+ \pi^- \pi^0 \pi^+ \pi^- \pi^0 \gamma$
<0.0063		¹ BISELLO 91	DM2	$J/\psi \rightarrow \gamma \omega \omega$
¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.				

NODE=M026R10
NODE=M026R10

NODE=M026R10;LINKAGE=E

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0017	90	¹ ABLIKIM 05L	BES2	$J/\psi \rightarrow \pi^+ \pi^- \pi^0 K^+ K^- \gamma$
¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.				

NODE=M026R22
NODE=M026R22

NODE=M026R22;LINKAGE=E

 $\Gamma(f_2(1270) f_2(1270))/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.25 OUR FIT				
0.77^{+0.25}_{-0.30} ± 0.17	91.2 ± 19.8	¹ ABLIKIM 04M	BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$
¹ ABLIKIM 04M reports $[\Gamma(\eta_c(1S) \rightarrow f_2(1270) f_2(1270))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.3 \pm 0.3+0.3-0.4) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M026R19
NODE=M026R19

NODE=M026R19;LINKAGE=AB

 $\Gamma(f_0(980)\eta)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$

NODE=M026R41
NODE=M026R41 $\Gamma(f_0(1500)\eta)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$

NODE=M026R42
NODE=M026R42 $\Gamma(f_0(2200)\eta)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$

NODE=M026R43
NODE=M026R43

$\Gamma(a_0(980)\pi)/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

NODE=M026R44
NODE=M026R44

 $\Gamma(a_0(1320)\pi)/\Gamma_{\text{total}} \quad \Gamma_{21}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

NODE=M026R45
NODE=M026R45

 $\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}} \quad \Gamma_{22}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

NODE=M026R46
NODE=M026R46

 $\Gamma(a_0(1950)\pi)/\Gamma_{\text{total}} \quad \Gamma_{23}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12k	¹ LEES 16A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

NODE=M026R00
NODE=M026R00

¹ From a model-independant partial wave analysis.

NODE=M026R00;LINKAGE=A

 $\Gamma(a_2(1950)\pi)/\Gamma_{\text{total}} \quad \Gamma_{24}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
not seen	12k	¹ LEES 16A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

NODE=M026R50
NODE=M026R50

¹ From a model-independent partial wave analysis assuming the existence of a hypothetical tensor isovector $a_2(1950)$.

NODE=M026R50;LINKAGE=A

 $\Gamma(K_0^*(1430)\bar{K})/\Gamma_{\text{total}} \quad \Gamma_{25}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12k	¹ LEES 16A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
seen		LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta/\pi^0$

NODE=M026R47
NODE=M026R47

¹ From a model-independant partial wave analysis.

NODE=M026R47;LINKAGE=A

 $\Gamma(K_2^*(1430)\bar{K})/\Gamma_{\text{total}} \quad \Gamma_{26}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

NODE=M026R48
NODE=M026R48

 $\Gamma(K_0^*(1950)\bar{K})/\Gamma_{\text{total}} \quad \Gamma_{27}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12K	¹ LEES 16A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
seen		LEES 14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta/\pi^0$

NODE=M026R49
NODE=M026R49

¹ From a Dalitz plot analysis using an isobar model.

NODE=M026R49;LINKAGE=A

 $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}} \quad \Gamma_{28}/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.3 ± 0.5 OUR FIT
6.5 ± 0.6 OUR AVERAGE

6.3 ± 1.3 ± 0.6	55	^{1,2} ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^0$	
7.9 ± 1.4 ± 0.7	107	^{3,4} ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\pm$	OCCUR=2
8.5 ± 1.8		⁵ AUBERT	06E	BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$	
5.1 ± 2.1	0.6k	⁶ BAI	04	BES	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$	
6.90 ± 1.42 ± 1.32	33	⁶ BISELLO	91	DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$	
5.43 ± 0.94 ± 0.94	68	⁶ BISELLO	91	DM2	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$	OCCUR=2
4.8 ± 1.7	95	^{6,7} BALTRUSAIT..86	MRK3		$J/\psi \rightarrow \eta_c \gamma$	
16.1 ^{+9.2} / _{-7.3}		^{8,9} HIMEL	80B	MRK2	$\psi(2S) \rightarrow \eta_c \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 10.7 90% CL ^{6,10} PARTRIDGE 80B CBAL $J/\psi \rightarrow \eta_c \gamma$

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (4.54 \pm 0.76 \pm 0.48) \times 10^{-6}$ which we multiply by 6 to account for isospin symmetry.

NODE=M026R4;LINKAGE=BK

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (27.24 \pm 4.56 \pm 2.88) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R4;LINKAGE=CK

³ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (11.35 \pm 1.25 \pm 1.50) \times 10^{-6}$ which we multiply by 3 to account for isospin symmetry.

NODE=M026R4;LINKAGE=BL

⁴ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (34.05 \pm 3.75 \pm 4.50) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R4;LINKAGE=CL

⁵ Determined from the ratio of $B(B^\pm \rightarrow K^\pm \eta_c) B(\eta_c \rightarrow K\bar{K}\pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$ reported in AUBERT, B 04B and $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$ reported in AUBERT 06E.

NODE=M026R4;LINKAGE=AB

⁶ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

NODE=M026R4;LINKAGE=E

⁷ Average from $K^+ K^- \pi^0$ and $K^\pm K_S^0 \pi^\mp$ decay channels.

NODE=M026R4;LINKAGE=D

⁸ $K^\pm K_S^0 \pi^\mp$ corrected to $K\bar{K}\pi$ by factor 3. KS, MR.

NODE=M026R4;LINKAGE=01

⁹ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

NODE=M026R4;LINKAGE=A

¹⁰ $K^+ K^- \pi^0$ corrected to $K\bar{K}\pi$ by factor 6. KS, MR

NODE=M026R4;LINKAGE=02

$\Gamma(\phi K^+ K^-)/\Gamma(K\bar{K}\pi)$

 Γ_6/Γ_{28}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.052^{+0.016}_{-0.014} ± 0.014	7	¹ HUANG	03	BELL $B^\pm \rightarrow K^\pm \phi \phi$

NODE=M026R02
NODE=M026R02

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R02;LINKAGE=BB

$\Gamma(K\bar{K}\eta)/\Gamma_{\text{total}}$

 Γ_{29}/Γ

VALUE (units 10^{-2})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
1.35 ± 0.16 OUR FIT					
1.0 ± 0.5 ± 0.2		7	^{1,2} ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta K^+ K^-$

NODE=M026R15
NODE=M026R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

OCCUR=2

<3.1 90 ³ BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \eta) = (2.11 \pm 1.01 \pm 0.32) \times 10^{-6}$ which we multiply by 2 to account for isospin symmetry.

NODE=M026R15;LINKAGE=AK

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] \times [B(h_c(1P) \rightarrow \eta_c(1S)\gamma)] = (4.22 \pm 2.02 \pm 0.64) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$, $B(h_c(1P) \rightarrow \eta_c(1S)\gamma) = (51 \pm 6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R15;LINKAGE=AM

³ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R15;LINKAGE=E

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$

 Γ_{29}/Γ_{28}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.186 ± 0.018 OUR FIT				
0.190 ± 0.008 ± 0.017	5.4k	¹ LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \eta/\pi^0$

NODE=M026R40
NODE=M026R40

¹ LEES 14E reports $B(\eta_c(1S) \rightarrow K^+ K^- \eta)/B(\eta_c(1S) \rightarrow K^+ K^- \pi^0) = 0.571 \pm 0.025 \pm 0.051$, which we divide by 3 to account for isospin symmetry. It uses both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

NODE=M026R40;LINKAGE=LE

$\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

 Γ_{30}/Γ

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 ± 0.1	33	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta \pi^+ \pi^-$

NODE=M026R6
NODE=M026R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.4 ± 2.0 75 ² BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$

3.7 ± 1.3 ± 2.0 18 ² PARTRIDGE 80B CBAL $J/\psi \rightarrow \eta \pi^+ \pi^- \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (7.22 \pm 1.47 \pm 1.11) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R6;LINKAGE=AB

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

NODE=M026R6;LINKAGE=E

$\Gamma(\eta 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

4.4±1.2±0.4 39 1 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma \eta 2(\pi^+ \pi^-)$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (19.17 \pm 3.77 \pm 3.72) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{31}/Γ NODE=M026R05
NODE=M026R05

NODE=M026R05;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

6.9± 1.1 OUR FIT

11.2± 1.9 OUR AVERAGE

9.7± 2.2±0.9 38 1 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^+ \pi^-$ 12 ± 4 0.4k 2 BAI 04 BES $J/\psi \rightarrow \gamma K^+ K^- \pi^+ \pi^-$ 21 ± 7 110 2 BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$ 14 +22
- 9 3 HIMEL 80B MRK2 $\psi(2S) \rightarrow \eta_c \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (4.16 \pm 0.76 \pm 0.59) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

 Γ_{32}/Γ NODE=M026R5
NODE=M026R5

NODE=M026R5;LINKAGE=AB

NODE=M026R5;LINKAGE=E

NODE=M026R5;LINKAGE=A

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$

VALUE EVTS

DOCUMENT ID TECN COMMENT

0.477±0.017±0.070 11k 1 DEL-AMO-SA..11M BABR $\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 Γ_{33}/Γ_{28} NODE=M026R01
NODE=M026R01

NODE=M026R01;LINKAGE=DE

 $\Gamma(K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

5.6±1.4±0.5 43 1,2 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\mp 2\pi^\pm$

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^- \pi^- 2\pi^+) = (12.01 \pm 2.22 \pm 2.04) \times 10^{-6}$ which we multiply by 2 to take c.c. into account.

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (24.02 \pm 4.44 \pm 4.08) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{34}/Γ NODE=M026R06
NODE=M026R06

NODE=M026R06;LINKAGE=AA

NODE=M026R06;LINKAGE=AB

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

7.5±2.4 OUR AVERAGE

8 ± 4 ± 1 10 1 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- 2(\pi^+ \pi^-)$ 7.2±2.4±1.6 100 2 ABLIKIM 06A BES2 $J/\psi \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (3.60 \pm 1.71 \pm 0.64) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.21 \pm 0.32 \pm 0.24) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{35}/Γ NODE=M026R23
NODE=M026R23

NODE=M026R23;LINKAGE=AL

NODE=M026R23;LINKAGE=AB

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{36}/Γ VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

1.46 ± 0.30 OUR FIT**2.2 ± 0.9 ± 0.2**

7

¹ ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma 2(K^+ K^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.4 $\begin{matrix} +0.5 \\ -0.4 \end{matrix}$ ± 0.6 14.5 $\begin{matrix} +4.6 \\ -3.0 \end{matrix}$ ² HUANG 03 BELL $B^+ \rightarrow 2(K^+ K^-) K^+$ 21 ± 10 ± 6 ³ ALBRECHT 94H ARG $\gamma \gamma \rightarrow K^+ K^- K^+ K^-$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (0.94 \pm 0.37 \pm 0.14) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 \begin{matrix} +0.10 \\ -0.12 \end{matrix}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

³ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow \phi \phi)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

NODE=M026R20
NODE=M026R20

NODE=M026R20;LINKAGE=AB

NODE=M026R20;LINKAGE=BB

NODE=M026R20;LINKAGE=AL

 $\Gamma(2(K^+ K^-))/\Gamma(K \bar{K} \pi)$ Γ_{36}/Γ_{28}

VALUE EVTS

DOCUMENT ID TECN COMMENT

0.020 ± 0.004 OUR FIT**0.024 ± 0.007 OUR AVERAGE**

0.023 ± 0.007 ± 0.006

AUBERT,B 04B BABR $B^\pm \rightarrow K^\pm \eta_c$ 0.026 $\begin{matrix} +0.009 \\ -0.007 \end{matrix}$ ± 0.007

15

¹ HUANG 03 BELL $B^\pm \rightarrow K^\pm (2K^+ 2K^-)$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 \begin{matrix} +0.10 \\ -0.12 \end{matrix}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R38
NODE=M026R38

NODE=M026R38;LINKAGE=BB

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

4.7 ± 0.9 ± 0.4

118

¹ ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma \pi^+ \pi^- 2\pi^0$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (20.31 \pm 2.20 \pm 3.33) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R07
NODE=M026R07

NODE=M026R07;LINKAGE=AB

 $\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{38}/Γ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

0.97 ± 0.12 OUR FIT**1.35 ± 0.21 OUR AVERAGE**

1.74 ± 0.32 ± 0.15

100

¹ ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^-)$

1.0 ± 0.5

542 ± 75

² BAI 04 BES $J/\psi \rightarrow \gamma 2(\pi^+ \pi^-)$

1.05 ± 0.17 ± 0.34

137

² BISELLO 91 DM2 $J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

1.3 ± 0.6

25

² BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$ 2.0 $\begin{matrix} +1.5 \\ -1.0 \end{matrix}$ ³ HIMEL 80B MRK2 $\psi(2S) \rightarrow \eta_c \gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (7.51 \pm 0.85 \pm 1.11) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

NODE=M026R1
NODE=M026R1

NODE=M026R1;LINKAGE=AB

NODE=M026R1;LINKAGE=E

NODE=M026R1;LINKAGE=A

 $\Gamma(2(\pi^+ \pi^- \pi^0))/\Gamma_{\text{total}}$ Γ_{39}/Γ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

17.4 ± 2.9 ± 1.5

175

¹ ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^- 2\pi^0)$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^- \pi^0))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (75.13 \pm 7.42 \pm 9.99) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R08
NODE=M026R08

NODE=M026R08;LINKAGE=AB

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{40}/Γ VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

18 ± 4 OUR AVERAGE

20 ± 5 ± 2	51	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 3(\pi^+\pi^-)$
15.3 ± 3.4 ± 3.3	479	2 ABLIKIM	06A BES2	$J/\psi \rightarrow 3(\pi^+\pi^-)\gamma$

NODE=M026R24
NODE=M026R24

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (8.82 \pm 1.57 \pm 1.59) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R24;LINKAGE=AL

² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (2.59 \pm 0.32 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R24;LINKAGE=AB

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{41}/Γ VALUE (units 10^{-4}) EVTS

DOCUMENT ID TECN COMMENT

15.0 ± 1.6 OUR FIT**13.2 ± 2.7 OUR AVERAGE**

15 ± 5 ± 1	15	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}$
15 ± 6	213 ± 33	2 BAI	04 BES	$J/\psi \rightarrow \gamma p\bar{p}$
10 ± 3 ± 4	18	2 BISELLO	91 DM2	$J/\psi \rightarrow \gamma p\bar{p}$
11 ± 6	23	2 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
29 +29 -15		3 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$

NODE=M026R2
NODE=M026R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.8 + 2.0 + 1.7 - 2.4 - 1.8	195	4 WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
---------------------------------	-----	------	---------	-------------------------------

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (0.65 \pm 0.19 \pm 0.10) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=AB

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

NODE=M026R2;LINKAGE=E

³ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

NODE=M026R2;LINKAGE=A

⁴ WU 06 reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (1.42 \pm 0.11^{+0.16}_{-0.20}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=WU

 $\Gamma(p\bar{p})/\Gamma(K\bar{K}\pi)$ Γ_{41}/Γ_{28}

VALUE EVTS

DOCUMENT ID TECN COMMENT

0.0207 ± 0.0021 OUR FIT**0.021 ± 0.002** +0.004
-0.006

195	1 WU	06 BELL	$B^\pm \rightarrow K^\pm p\bar{p}$
-----	------	---------	------------------------------------

NODE=M026R03
NODE=M026R03

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R03;LINKAGE=BB

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$ $\Gamma_{41}/\Gamma \times \Gamma_7/\Gamma$ VALUE (units 10^{-5})

DOCUMENT ID TECN COMMENT

0.26 ± 0.05 OUR FIT**4.0** +3.5
-3.2

	BAGLIN	89 SPEC	$\bar{p}p \rightarrow K^+ K^- K^+ K^-$
--	--------	---------	--

NODE=M026R33
NODE=M026R33 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

0.36 ± 0.13 ± 0.03

14	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}\pi^0$
----	-----------	----------	---

NODE=M026R09
NODE=M026R09

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (1.53 \pm 0.49 \pm 0.23) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R09;LINKAGE=AB

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

10.9±2.4 OUR FIT**11.7±2.3±2.6**¹ ABLIKIM 12B BES3

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.9^{+2.7}_{-2.6} \pm 1.2$	20	² WU	06	BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
-----------------------------	----	-----------------	----	------	---

<20	90	³ BISELLO	91	DM2	$e^+e^- \rightarrow \gamma\Lambda\bar{\Lambda}$
-----	----	----------------------	----	-----	---

¹ ABLIKIM 12B reports $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (0.198 \pm 0.021 \pm 0.032) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² WU 06 reports $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (0.95^{+0.25+0.08}_{-0.22-0.11}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R18
NODE=M026R18

NODE=M026R18;LINKAGE=AB

NODE=M026R18;LINKAGE=WU

NODE=M026R18;LINKAGE=E

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma(p\bar{p})$ Γ_{43}/Γ_{41}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.72±0.16 OUR FIT**0.67^{+0.19}_{-0.16}±0.12**¹ WU 06 BELL $B^+ \rightarrow p\bar{p}K^+, \Lambda\bar{\Lambda}K^+$

¹ Not independent from other $\eta_c \rightarrow \Lambda\bar{\Lambda}, p\bar{p}$ branching ratios reported by WU 06.

NODE=M026R27
NODE=M026R27

NODE=M026R27;LINKAGE=WU

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.1±0.3±0.5112 ¹ ABLIKIM 13C BES3 $J/\psi \rightarrow \gamma p\bar{p}\pi^0\pi^0$

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.60 \pm 0.48 \pm 0.31) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R28
NODE=M026R28

NODE=M026R28;LINKAGE=AB

 $\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.89±0.18±0.1978 ¹ ABLIKIM 13C BES3 $J/\psi \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.51 \pm 0.27 \pm 0.14) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R29
NODE=M026R29

NODE=M026R29;LINKAGE=AB

 $\Gamma(\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

5.3±1.7±0.519 ¹ ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0\gamma p\bar{p}\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	HIMEL	80B	MRK2	$\psi(2S) \rightarrow \eta_c\gamma$
-----	----	-------	-----	------	-------------------------------------

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (2.30 \pm 0.65 \pm 0.36) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R3
NODE=M026R3

NODE=M026R3;LINKAGE=AB

————— RADIATIVE DECAYS —————

NODE=M026310

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.59±0.13 OUR FIT[(1.59 ± 0.12) × 10⁻⁴ OUR 2015 FIT]**1.9^{+0.7}_{-0.6} OUR AVERAGE**

2.7 ± 0.8 ± 0.6

¹ ABLIKIM 13I BES31.4^{+0.7}_{-0.5} ± 0.3 1.2^{+2.8}_{-1.1}² ADAMS 08 CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ NODE=M026R31
NODE=M026R31

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3	$\begin{smallmatrix} +1.0 \\ -0.8 \end{smallmatrix} \pm 0.3$	13	3	WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
2.80	$\begin{smallmatrix} +0.67 \\ -0.58 \end{smallmatrix} \pm 1.0$		4	ARMSTRONG	95F	E760	$\bar{p} p \rightarrow \gamma \gamma$
< 9	90		5	BISELLO	91	DM2	$J/\psi \rightarrow \gamma \gamma \gamma$
6	$\begin{smallmatrix} +4 \\ -3 \end{smallmatrix} \pm 4$		4	BAGLIN	87B	SPEC	$\bar{p} p \rightarrow \gamma \gamma$
< 18	90		6	BLOOM	83	CBAL	$J/\psi \rightarrow \eta_c \gamma$

¹ ABLIKIM 13I reports $[\Gamma(\eta_c(1S) \rightarrow \gamma \gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (4.5 \pm 1.2 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R31;LINKAGE=AL

² ADAMS 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma \gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (2.4 \begin{smallmatrix} +1.1 \\ -0.8 \end{smallmatrix} \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R31;LINKAGE=AD

³ WICHT 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma \gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (2.2 \begin{smallmatrix} +0.9+0.4 \\ -0.7-0.2 \end{smallmatrix}) \times 10^{-7}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R31;LINKAGE=WI

⁴ Not independent from the values of the total and two-photon width quoted by the same experiment.

NODE=M026R31;LINKAGE=AB

⁵ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R31;LINKAGE=E

⁶ Using $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R31;LINKAGE=C

$\Gamma(\gamma \gamma)/\Gamma(K \bar{K} \pi)$

Γ_{47}/Γ_{28}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.19 ± 0.29 OUR FIT				

NODE=M026R04
NODE=M026R04

3.2	$\begin{smallmatrix} +1.3 & +0.8 \\ -1.0 & -0.6 \end{smallmatrix}$	13	1	WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
-----	--	----	---	-------	----	------	---

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 \begin{smallmatrix} +0.10 \\ -0.12 \end{smallmatrix}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R04;LINKAGE=BB

$\Gamma(\bar{p} p)/\Gamma_{\text{total}} \times \Gamma(\gamma \gamma)/\Gamma_{\text{total}}$

$\Gamma_{41}/\Gamma \times \Gamma_{47}/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
0.240 ± 0.024 OUR FIT				
[(0.239 ± 0.024) × 10 ⁻⁶ OUR 2015 FIT]				
0.26 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.4.

NODE=M026R32
NODE=M026R32
NEW

0.224	$\begin{smallmatrix} +0.038 \\ -0.037 \end{smallmatrix} \pm 0.020$	190		AMBROGIANI	03	E835	$\bar{p} p \rightarrow \eta_c \rightarrow \gamma \gamma$
-------	--	-----	--	------------	----	------	--

0.336	$\begin{smallmatrix} +0.080 \\ -0.070 \end{smallmatrix}$			ARMSTRONG	95F	E760	$\bar{p} p \rightarrow \gamma \gamma$
-------	--	--	--	-----------	-----	------	---------------------------------------

0.68	$\begin{smallmatrix} +0.42 \\ -0.31 \end{smallmatrix}$	12		BAGLIN	87B	SPEC	$\bar{p} p \rightarrow \gamma \gamma$
------	--	----	--	--------	-----	------	---------------------------------------

————— Charge conjugation (C), Parity (P), —————
————— Lepton family number (LF) violating modes —————

NODE=M026320

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{48}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	1	ABLIKIM	11G BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^-$

NODE=M026R34
NODE=M026R34

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 70	90	2	ABLIKIM	06B BES2 $J/\psi \rightarrow \pi^+ \pi^- \gamma$
------	----	---	---------	--

¹ ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 1.82 \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

NODE=M026R34;LINKAGE=AL

² ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 1.1 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

NODE=M026R34;LINKAGE=AB

$\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$

Γ_{49}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4	90	1	ABLIKIM	11G BES3 $J/\psi \rightarrow \gamma \pi^0 \pi^0$

NODE=M026R35
NODE=M026R35

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 40	90	2	ABLIKIM	06B BES2 $J/\psi \rightarrow \pi^0 \pi^0 \gamma$
------	----	---	---------	--

¹ ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 6.0 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

NODE=M026R35;LINKAGE=AL

² ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.71 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

NODE=M026R35;LINKAGE=AB

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT
<60	90	¹ ABLIKIM	06B	BES2 $J/\psi \rightarrow K^+K^-\gamma$

¹ ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K^+K^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.96 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.

NODE=M026R36
NODE=M026R36

NODE=M026R36;LINKAGE=AB

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT
<31	90	¹ ABLIKIM	06B	BES2 $J/\psi \rightarrow K_S^0 K_S^0 \gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

<32	90	² UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
< 5.6	90	³ UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M026R37
NODE=M026R37

OCCUR=2

¹ ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.53 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.

NODE=M026R37;LINKAGE=AB

² Taking into account interference with the non-resonant continuum.

NODE=M026R37;LINKAGE=U1

³ Neglecting interference with the non-resonant continuum.

NODE=M026R37;LINKAGE=U2

$\eta_c(1S)$ REFERENCES

NODE=M026

LEES	16A	PR D93 012005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57125
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57147
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56130
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55937
ABLIKIM	13C	PR D87 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54878
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54954
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54267
ABLIKIM	12F	PRL 108 222002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54271
ABLIKIM	12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54741
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)	REFID=54763
ABLIKIM	11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53711
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=53927
LEES	10	PR D81 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53236
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=52261
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52267
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
ABLIKIM	06A	PL B633 19	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50987
ABLIKIM	06B	EPJ C45 337	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50988
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM	05L	PR D72 072005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50837
KUO	05	PL B621 41	C.C. Kuo <i>et al.</i>	(BELLE Collab.)	REFID=50801
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=49745
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49746
AUBERT,B	04B	PR D70 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50043
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49620
ABDALLAH	03J	EPJ C31 481	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49625
AMBROGIANI	03	PL B566 45	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=49465
BAI	03	PL B555 174	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49185
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)	REFID=49206
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
BAI	00F	PR D62 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48546
BRANDENB...	00B	PRL 85 3095	G. Brandenburg <i>et al.</i>	(CLEO Collab.)	REFID=48553
ACCIARRI	99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47476
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46553
SHIRAI	98	PL B424 405	M. Shirai <i>et al.</i>	(AMY Collab.)	REFID=46381
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=44623
ALBRECHT	94H	PL B338 390	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44098
ADRIANI	93N	PL B318 575	O. Adriani <i>et al.</i>	(L3 Collab.)	REFID=43670
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41668
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
CHEN	90B	PL B243 169	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=41360
BAGLIN	89	PL B231 557	C. Baglin, S. Baird, G. Bassompierre	(R704 Collab.)	REFID=41015
BEHREND	89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40732
BRAUNSCH...	89	ZPHY C41 533	W. Braunschweig <i>et al.</i>	(TASSO Collab.)	REFID=40728
AIHARA	88D	PRL 60 2355	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=40588
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BERGER	86	PL 167B 120	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22010
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BALTRUSAIT...	84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22006
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
HIMEL	80B	PRL 45 1146	T.M. Himel <i>et al.</i>	(SLAC, LBL, UCB)	REFID=22003
PARTRIDGE	80B	PRL 45 1150	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22004

J/ψ(1S)

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M070

J/ψ(1S) MASS

NODE=M070M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3096.900±0.006 OUR AVERAGE				
[3096.916 ± 0.011 MeV OUR 2015 AVERAGE]				
3096.66 ± 0.19 ± 0.02	6.1k	¹ AAIJ	15Bl LHCb	$pp \rightarrow J/\psi X$
3096.900±0.002±0.006		² ANASHIN	15 KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ± 0.09	502	³ ARTAMONOV	00 OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ± 0.03 ± 0.01		⁴ ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ± 0.1 ± 0.3	193	BAGLIN	87 SPEC	$\bar{p}p \rightarrow e^+e^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3096.917±0.010±0.007		AULCHENKO	03 KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ± 0.3		GRIBUSHIN	96 FMPS	$515 \pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ± 2.0	38k	LEMOIGNE	82 GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
3096.93 ± 0.09	502	⁵ ZHOLENTZ	80 REDE	e^+e^-
3097.0 ± 1		⁶ BRANDELIK	79C DASP	e^+e^-

NODE=M070M
NEW¹ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays.² Supersedes AULCHENKO 03.³ Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).⁴ Mass central value and systematic error recalculated by us according to Eq.(16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁵ Superseded by ARTAMONOV 00.⁶ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.NODE=M070M;LINKAGE=B
NODE=M070M;LINKAGE=A
NODE=M070M;LINKAGE=AR

NODE=M070M;LINKAGE=NW

NODE=M070M;LINKAGE=RZ
NODE=M070M;LINKAGE=F**J/ψ(1S) WIDTH**

NODE=M070W

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
92.9± 2.8 OUR AVERAGE Error includes scale factor of 1.1.				
96.1± 3.2	13k	¹ ADAMS	06A CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4± 8.9		BAI	95B BES	e^+e^-
91 ± 11 ± 6		² ARMSTRONG	93B E760	$\bar{p}p \rightarrow e^+e^-$
85.5 ⁺ ₋ 6.1 5.8		³ HSUEH	92 RVUE	See Υ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
94.1± 2.7		⁴ ANASHIN	10 KEDR	$3.097 e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7± 3.5	7.8k	¹ AUBERT	04 BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

NODE=M070W

¹ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(e^+e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].³ Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.⁴ Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

NODE=M070W;LINKAGE=AA

NODE=M070W;LINKAGE=AN
NODE=M070W;LINKAGE=A

NODE=M070W;LINKAGE=AS

J/ψ(1S) DECAY MODES

NODE=M070215;NODE=M070

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 ± 0.5) %	DESIG=3
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.50 ± 0.30) %	DESIG=4
Γ_3 ggg	(64.1 ± 1.0) %	DESIG=249
Γ_4 γgg	(8.8 ± 1.1) %	DESIG=250
Γ_5 e^+e^-	(5.971±0.032) %	DESIG=1
Γ_6 $e^+e^-\gamma$	[a] (8.8 ± 1.4) × 10 ⁻³	DESIG=5
Γ_7 $\mu^+\mu^-$	(5.961±0.033) %	DESIG=2

Decays involving hadronic resonances

				NODE=M070;CLUMP=A
Γ_8	$\rho\pi$	(1.69 \pm 0.15) %	S=2.4	DESIG=20
Γ_9	$\rho^0\pi^0$	(5.6 \pm 0.7) $\times 10^{-3}$		DESIG=21
Γ_{10}	$a_2(1320)\rho$	(1.09 \pm 0.22) %		DESIG=43
Γ_{11}	$\omega\pi^+\pi^+\pi^-\pi^-$	(8.5 \pm 3.4) $\times 10^{-3}$		DESIG=26
Γ_{12}	$\omega\pi^+\pi^-\pi^0$	(4.0 \pm 0.7) $\times 10^{-3}$		DESIG=211
Γ_{13}	$\omega\pi^+\pi^-$	(8.6 \pm 0.7) $\times 10^{-3}$	S=1.1	DESIG=24
Γ_{14}	$\omega f_2(1270)$	(4.3 \pm 0.6) $\times 10^{-3}$		DESIG=28
Γ_{15}	$K^*(892)^0\bar{K}^*(892)^0$	(2.3 \pm 0.7) $\times 10^{-4}$		DESIG=46
Γ_{16}	$K^*(892)^\pm K^*(892)^\mp$	(1.00 $^{+0.22}_{-0.40}$) $\times 10^{-3}$		DESIG=256
Γ_{17}	$K^*(892)^\pm K^*(800)^\mp$	(1.1 $^{+1.0}_{-0.6}$) $\times 10^{-3}$		DESIG=257
Γ_{18}	$\eta K^*(892)^0\bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$		DESIG=252
Γ_{19}	$K^*(892)^0\bar{K}_2^*(1430)^0 + c.c.$	(6.0 \pm 0.6) $\times 10^{-3}$		DESIG=48
Γ_{20}	$K^*(892)^0\bar{K}_2^*(1770)^0 + c.c. \rightarrow$ $K^*(892)^0 K^-\pi^+ + c.c.$	(6.9 \pm 0.9) $\times 10^{-4}$		DESIG=235
Γ_{21}	$\omega K^*(892)\bar{K} + c.c.$	(6.1 \pm 0.9) $\times 10^{-3}$		DESIG=102
Γ_{22}	$K^+ K^*(892)^- + c.c.$	(5.12 \pm 0.30) $\times 10^{-3}$		DESIG=121
Γ_{23}	$K^+ K^*(892)^- + c.c. \rightarrow$ $K^+ K^-\pi^0$	(1.97 \pm 0.20) $\times 10^{-3}$		DESIG=231
Γ_{24}	$K^+ K^*(892)^- + c.c. \rightarrow$ $K^0 K^\pm\pi^\mp + c.c.$	(3.0 \pm 0.4) $\times 10^{-3}$		DESIG=232
Γ_{25}	$K^0\bar{K}^*(892)^0 + c.c.$	(4.39 \pm 0.31) $\times 10^{-3}$		DESIG=122
Γ_{26}	$K^0\bar{K}^*(892)^0 + c.c. \rightarrow$ $K^0 K^\pm\pi^\mp + c.c.$	(3.2 \pm 0.4) $\times 10^{-3}$		DESIG=233
Γ_{27}	$K_1(1400)^\pm K^\mp$	(3.8 \pm 1.4) $\times 10^{-3}$		DESIG=132
Γ_{28}	$\bar{K}^*(892)^0 K^+\pi^- + c.c.$	seen		DESIG=214
Γ_{29}	$\omega\pi^0\pi^0$	(3.4 \pm 0.8) $\times 10^{-3}$		DESIG=140
Γ_{30}	$b_1(1235)^\pm\pi^\mp$	[b] (3.0 \pm 0.5) $\times 10^{-3}$		DESIG=49
Γ_{31}	$\omega K^\pm K_S^0\pi^\mp$	[b] (3.4 \pm 0.5) $\times 10^{-3}$		DESIG=101
Γ_{32}	$b_1(1235)^0\pi^0$	(2.3 \pm 0.6) $\times 10^{-3}$		DESIG=160
Γ_{33}	$\eta K^\pm K_S^0\pi^\mp$	[b] (2.2 \pm 0.4) $\times 10^{-3}$		DESIG=230
Γ_{34}	$\phi K^*(892)\bar{K} + c.c.$	(2.18 \pm 0.23) $\times 10^{-3}$		DESIG=104
Γ_{35}	$\omega K\bar{K}$	(1.70 \pm 0.32) $\times 10^{-3}$		DESIG=27
Γ_{36}	$\omega f_0(1710) \rightarrow \omega K\bar{K}$	(4.8 \pm 1.1) $\times 10^{-4}$		DESIG=130
Γ_{37}	$\phi 2(\pi^+\pi^-)$	(1.66 \pm 0.23) $\times 10^{-3}$		DESIG=35
Γ_{38}	$\Delta(1232)^{++}\bar{p}\pi^-$	(1.6 \pm 0.5) $\times 10^{-3}$		DESIG=70
Γ_{39}	$\omega\eta$	(1.74 \pm 0.20) $\times 10^{-3}$	S=1.6	DESIG=30
Γ_{40}	$\phi K\bar{K}$	(1.83 \pm 0.24) $\times 10^{-3}$	S=1.5	DESIG=36
Γ_{41}	$\phi f_0(1710) \rightarrow \phi K\bar{K}$	(3.6 \pm 0.6) $\times 10^{-4}$		DESIG=129
Γ_{42}	$\phi f_2(1270)$	(7.2 \pm 1.3) $\times 10^{-4}$		DESIG=39
Γ_{43}	$\Delta(1232)^{++}\bar{\Delta}(1232)^{--}$	(1.10 \pm 0.29) $\times 10^{-3}$		DESIG=66
Γ_{44}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+ (or c.c.)$	[b] (1.10 \pm 0.12) $\times 10^{-3}$		DESIG=67
Γ_{45}	$\phi f_2'(1525)$	(8 \pm 4) $\times 10^{-4}$	S=2.7	DESIG=40
Γ_{46}	$\phi\pi^+\pi^-$	(9.4 \pm 0.9) $\times 10^{-4}$	S=1.2	DESIG=34
Γ_{47}	$\phi\pi^0\pi^0$	(5.6 \pm 1.6) $\times 10^{-4}$		DESIG=76
Γ_{48}	$\phi K^\pm K_S^0\pi^\mp$	[b] (7.2 \pm 0.8) $\times 10^{-4}$		DESIG=103
Γ_{49}	$\omega f_1(1420)$	(6.8 \pm 2.4) $\times 10^{-4}$		DESIG=105
Γ_{50}	$\phi\eta$	(7.5 \pm 0.8) $\times 10^{-4}$	S=1.5	DESIG=37
Γ_{51}	$\Xi^0\Xi^0$	(1.20 \pm 0.24) $\times 10^{-3}$		DESIG=248
Γ_{52}	$\Xi(1530)^-\Xi^+$	(5.9 \pm 1.5) $\times 10^{-4}$		DESIG=107
Γ_{53}	$\rho K^-\bar{\Sigma}(1385)^0$	(5.1 \pm 3.2) $\times 10^{-4}$		DESIG=74
Γ_{54}	$\omega\pi^0$	(4.5 \pm 0.5) $\times 10^{-4}$	S=1.4	DESIG=32
Γ_{55}	$\phi\eta'(958)$	(4.0 \pm 0.7) $\times 10^{-4}$	S=2.1	DESIG=38
Γ_{56}	$\phi f_0(980)$	(3.2 \pm 0.9) $\times 10^{-4}$	S=1.9	DESIG=41
Γ_{57}	$\phi f_0(980) \rightarrow \phi\pi^+\pi^-$	(1.8 \pm 0.4) $\times 10^{-4}$		DESIG=236
Γ_{58}	$\phi f_0(980) \rightarrow \phi\pi^0\pi^0$	(1.7 \pm 0.7) $\times 10^{-4}$		DESIG=237
Γ_{59}	$\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-$	(4.5 \pm 1.0) $\times 10^{-6}$		DESIG=278
Γ_{60}	$\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0$	(1.7 \pm 0.6) $\times 10^{-6}$		DESIG=279

Г61	$\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	(3.2 ±1.0) × 10 ⁻⁴		DESIG=229
Г62	$\phi a_0(980)^0 \rightarrow \phi\eta\pi^0$	(5 ±4) × 10 ⁻⁶		DESIG=258
Г63	$\Xi(1530)^0 \Xi^0$	(3.2 ±1.4) × 10 ⁻⁴		DESIG=108
Г64	$\Sigma(1385)^- \bar{\Sigma}^+$ (or c.c.)	[b] (3.1 ±0.5) × 10 ⁻⁴		DESIG=68
Г65	$\phi f_1(1285)$	(2.6 ±0.5) × 10 ⁻⁴		DESIG=106
Г66	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow$ $\phi\pi^0\pi^+\pi^-$	(9.4 ±2.8) × 10 ⁻⁷		DESIG=280
Г67	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow$ $\phi\pi^0\pi^0\pi^0$	(2.1 ±2.2) × 10 ⁻⁷		DESIG=281
Г68	$\eta\pi^+\pi^-$	(4.0 ±1.7) × 10 ⁻⁴		DESIG=239
Г69	$\eta\rho$	(1.93 ±0.23) × 10 ⁻⁴		DESIG=22
Г70	$\omega\eta'(958)$	(1.82 ±0.21) × 10 ⁻⁴		DESIG=31
Г71	$\omega f_0(980)$	(1.4 ±0.5) × 10 ⁻⁴		DESIG=150
Г72	$\rho\eta'(958)$	(1.05 ±0.18) × 10 ⁻⁴		DESIG=23
Г73	$a_2(1320)^\pm\pi^\mp$	[b] < 4.3 × 10 ⁻³	CL=90%	DESIG=42
Г74	$K K_2^*(1430) + c.c.$	< 4.0 × 10 ⁻³	CL=90%	DESIG=45
Г75	$K_1(1270)^\pm K^\mp$	< 3.0 × 10 ⁻³	CL=90%	DESIG=131
Г76	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	< 2.9 × 10 ⁻³	CL=90%	DESIG=47
Г77	$\phi\pi^0$	3 × 10 ⁻⁶ or 1 × 10 ⁻⁷		DESIG=33;OUR EVAL;→ UNCHECKED ←
Г78	$\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-$	(2.0 ±1.0) × 10 ⁻⁵		DESIG=128
Г79	$\omega f_2'(1525)$	< 2.2 × 10 ⁻⁴	CL=90%	DESIG=29
Г80	$\omega X(1835) \rightarrow \omega p\bar{p}$	< 3.9 × 10 ⁻⁶	CL=95%	DESIG=263
Г81	$\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-$	< 2.8 × 10 ⁻⁴	CL=90%	DESIG=288
Г82	$\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-$	< 6.13 × 10 ⁻⁵	CL=90%	DESIG=289
Г83	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow$ $\eta\phi\pi^+\pi^-$	(1.2 ±0.4) × 10 ⁻⁴		DESIG=287
Г84	$\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0 \bar{K}^*(892)^0$	< 2.52 × 10 ⁻⁴	CL=90%	DESIG=253
Г85	$\Sigma(1385)^0 \bar{\Lambda} + c.c.$	< 8.2 × 10 ⁻⁶	CL=90%	DESIG=111
Г86	$\Delta(1232)^+ \bar{p}$	< 1 × 10 ⁻⁴	CL=90%	DESIG=112
Г87	$\Lambda(1520) \bar{\Lambda} + c.c. \rightarrow \gamma\Lambda\bar{\Lambda}$	< 4.1 × 10 ⁻⁶	CL=90%	DESIG=260
Г88	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + c.c.$	< 1.1 × 10 ⁻⁵	CL=90%	DESIG=205
Г89	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 2.1 × 10 ⁻⁵	CL=90%	DESIG=206
Г90	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 1.6 × 10 ⁻⁵	CL=90%	DESIG=207
Г91	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 5.6 × 10 ⁻⁵	CL=90%	DESIG=208
Г92	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	< 1.1 × 10 ⁻⁵	CL=90%	DESIG=209
Г93	$\Sigma^0 \bar{\Lambda}$	< 9 × 10 ⁻⁵	CL=90%	DESIG=110

Decays into stable hadrons

Г94	$2(\pi^+\pi^-)\pi^0$	(4.1 ±0.5) %	S=2.4	NODE=M070;CLUMP=B DESIG=9
Г95	$3(\pi^+\pi^-)\pi^0$	(2.9 ±0.6) %		DESIG=11
Г96	$\pi^+\pi^-\pi^0$	(2.11 ±0.07) %	S=1.5	DESIG=7
Г97	$\pi^+\pi^-\pi^0 K^+ K^-$	(1.79 ±0.29) %	S=2.2	DESIG=18
Г98	$4(\pi^+\pi^-)\pi^0$	(9.0 ±3.0) × 10 ⁻³		DESIG=12
Г99	$\pi^+\pi^- K^+ K^-$	(6.6 ±0.5) × 10 ⁻³		DESIG=16
Г100	$\pi^+\pi^- K^+ K^- \eta$	(1.84 ±0.28) × 10 ⁻³		DESIG=238
Г101	$\pi^0\pi^0 K^+ K^-$	(2.45 ±0.31) × 10 ⁻³		DESIG=234
Г102	$K\bar{K}\pi$	(6.1 ±1.0) × 10 ⁻³		DESIG=15
Г103	$2(\pi^+\pi^-)$	(3.57 ±0.30) × 10 ⁻³		DESIG=8
Г104	$3(\pi^+\pi^-)$	(4.3 ±0.4) × 10 ⁻³		DESIG=10
Г105	$2(\pi^+\pi^-\pi^0)$	(1.62 ±0.21) %		DESIG=210
Г106	$2(\pi^+\pi^-)\eta$	(2.29 ±0.24) × 10 ⁻³		DESIG=201
Г107	$3(\pi^+\pi^-)\eta$	(7.2 ±1.5) × 10 ⁻⁴		DESIG=202
Г108	$\rho\bar{p}$	(2.120 ±0.029) × 10 ⁻³		DESIG=50
Г109	$\rho\bar{p}\pi^0$	(1.19 ±0.08) × 10 ⁻³	S=1.1	DESIG=52
Г110	$\rho\bar{p}\pi^+\pi^-$	(6.0 ±0.5) × 10 ⁻³	S=1.3	DESIG=54
Г111	$\rho\bar{p}\pi^+\pi^-\pi^0$	[c] (2.3 ±0.9) × 10 ⁻³	S=1.9	DESIG=55
Г112	$\rho\bar{p}\eta$	(2.00 ±0.12) × 10 ⁻³		DESIG=56
Г113	$\rho\bar{p}\rho$	< 3.1 × 10 ⁻⁴	CL=90%	DESIG=57
Г114	$\rho\bar{p}\omega$	(9.8 ±1.0) × 10 ⁻⁴	S=1.3	DESIG=58

Γ ₁₁₅	$p\bar{p}\eta'(958)$	$(2.1 \pm 0.4) \times 10^{-4}$		DESIG=59
Γ ₁₁₆	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$	$(6.8 \pm 1.8) \times 10^{-5}$		DESIG=276
Γ ₁₁₇	$p\bar{p}\phi$	$(4.5 \pm 1.5) \times 10^{-5}$		DESIG=127
Γ ₁₁₈	$n\bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$		DESIG=64
Γ ₁₁₉	$n\bar{n}\pi^+\pi^-$	$(4 \pm 4) \times 10^{-3}$		DESIG=65
Γ ₁₂₀	$\Sigma^+\bar{\Sigma}^-$	$(1.50 \pm 0.24) \times 10^{-3}$		DESIG=247
Γ ₁₂₁	$\Sigma^0\bar{\Sigma}^0$	$(1.29 \pm 0.09) \times 10^{-3}$		DESIG=63
Γ ₁₂₂	$2(\pi^+\pi^-)K^+K^-$	$(4.7 \pm 0.7) \times 10^{-3}$	S=1.3	DESIG=17
Γ ₁₂₃	$p\bar{n}\pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$		DESIG=53
Γ ₁₂₄	$nN(1440)$	seen		DESIG=215;OUR EST;→ UNCHECKED ←
Γ ₁₂₅	$nN(1520)$	seen		DESIG=216;OUR EST;→ UNCHECKED ←
Γ ₁₂₆	$nN(1535)$	seen		DESIG=217;OUR EST;→ UNCHECKED ←
Γ ₁₂₇	$\Xi^-\bar{\Xi}^+$	$(8.6 \pm 1.1) \times 10^{-4}$	S=1.2	DESIG=62
Γ ₁₂₈	$\Lambda\bar{\Lambda}$	$(1.61 \pm 0.15) \times 10^{-3}$	S=1.9	DESIG=60
Γ ₁₂₉	$\Lambda\bar{\Sigma}^-\pi^+$ (or c.c.)	[b] $(8.3 \pm 0.7) \times 10^{-4}$	S=1.2	DESIG=71
Γ ₁₃₀	$\rho K^-\bar{\Lambda}$	$(8.9 \pm 1.6) \times 10^{-4}$		DESIG=72
Γ ₁₃₁	$2(K^+K^-)$	$(7.6 \pm 0.9) \times 10^{-4}$		DESIG=19
Γ ₁₃₂	$\rho K^-\bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$		DESIG=73
Γ ₁₃₃	K^+K^-	$(2.86 \pm 0.21) \times 10^{-4}$		DESIG=13
Γ ₁₃₄	$K_S^0 K_L^0$	$(2.1 \pm 0.4) \times 10^{-4}$	S=3.2	DESIG=75
Γ ₁₃₅	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$		DESIG=261
Γ ₁₃₆	$\Lambda\bar{\Lambda}\eta$	$(1.62 \pm 0.17) \times 10^{-4}$		DESIG=228
Γ ₁₃₇	$\Lambda\bar{\Lambda}\pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$		DESIG=109
Γ ₁₃₈	$\bar{\Lambda}nK_S^0$ + c.c.	$(6.5 \pm 1.1) \times 10^{-4}$		DESIG=225
Γ ₁₃₉	$\pi^+\pi^-$	$(1.47 \pm 0.14) \times 10^{-4}$		DESIG=6
Γ ₁₄₀	$\Lambda\bar{\Sigma}^+$ + c.c.	$(2.83 \pm 0.23) \times 10^{-5}$		DESIG=61
Γ ₁₄₁	$K_S^0 K_S^0$	$< 1 \times 10^{-6}$	CL=95%	DESIG=14

Radiative decays

				NODE=M070;CLUMP=C
Γ ₁₄₂	3γ	$(1.16 \pm 0.22) \times 10^{-5}$		DESIG=81
Γ ₁₄₃	4γ	$< 9 \times 10^{-6}$	CL=90%	DESIG=244
Γ ₁₄₄	5γ	$< 1.5 \times 10^{-5}$	CL=90%	DESIG=245
Γ ₁₄₅	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$		DESIG=283
Γ ₁₄₆	$\gamma\eta_c(1S)$	$(1.7 \pm 0.4)\%$	S=1.5	DESIG=85
Γ ₁₄₇	$\gamma\eta_c(1S) \rightarrow 3\gamma$	$(3.8 \pm 1.3 \mp 1.0) \times 10^{-6}$	S=1.1	DESIG=246
Γ ₁₄₈	$\gamma\pi^+\pi^-2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$		DESIG=99
Γ ₁₄₉	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$		DESIG=96
Γ ₁₅₀	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$		DESIG=142
Γ ₁₅₁	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	[d] $(2.8 \pm 0.6) \times 10^{-3}$	S=1.6	DESIG=89
Γ ₁₅₂	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8	DESIG=171
Γ ₁₅₃	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$		DESIG=170
Γ ₁₅₄	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	$< 8.2 \times 10^{-5}$	CL=95%	DESIG=212
Γ ₁₅₅	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$		DESIG=94
Γ ₁₅₆	$\gamma\rho\omega$	$< 5.4 \times 10^{-4}$	CL=90%	DESIG=226
Γ ₁₅₇	$\gamma\rho\phi$	$< 8.8 \times 10^{-5}$	CL=90%	DESIG=227
Γ ₁₅₈	$\gamma\eta'(958)$	$(5.15 \pm 0.16) \times 10^{-3}$	S=1.2	DESIG=84
Γ ₁₅₉	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9	DESIG=95
Γ ₁₆₀	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$		DESIG=203
Γ ₁₆₁	$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$		DESIG=204
Γ ₁₆₂	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$		DESIG=143
Γ ₁₆₃	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$		DESIG=100
Γ ₁₆₄	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$		DESIG=97
Γ ₁₆₅	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3	DESIG=124
Γ ₁₆₆	$\gamma f_2(1270)$	$(1.64 \pm 0.12) \times 10^{-3}$	S=1.3	DESIG=86
Γ ₁₆₇	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$		DESIG=284
Γ ₁₆₈	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(1.00 \pm 0.11 \mp 0.09) \times 10^{-3}$	S=1.5	DESIG=91
Γ ₁₆₉	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$		DESIG=135
Γ ₁₇₀	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$		DESIG=221

Γ ₁₇₁	$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$(2.4 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix}) \times 10^{-4}$		DESIG=266
Γ ₁₇₂	$\gamma \eta$	$(1.104 \pm 0.034) \times 10^{-3}$		DESIG=83
Γ ₁₇₃	$\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi$	$(7.9 \pm 1.3) \times 10^{-4}$		DESIG=175
Γ ₁₇₄	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$		DESIG=88
Γ ₁₇₅	$\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$		DESIG=141
Γ ₁₇₆	$\gamma f'_2(1525)$	$(5.7 \begin{smallmatrix} +0.8 \\ -0.5 \end{smallmatrix}) \times 10^{-4}$	S=1.5	DESIG=87
Γ ₁₇₇	$\gamma f'_2(1525) \rightarrow \gamma \eta \eta$	$(3.4 \pm 1.4) \times 10^{-5}$		DESIG=268
Γ ₁₇₈	$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	$(2.8 \pm 1.8) \times 10^{-4}$		DESIG=222
Γ ₁₇₉	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$(2.0 \pm 1.4) \times 10^{-4}$		DESIG=223
Γ ₁₈₀	$\gamma f_0(1800) \rightarrow \gamma \omega \phi$	$(2.5 \pm 0.6) \times 10^{-4}$		DESIG=262
Γ ₁₈₁	$\gamma f_2(1810) \rightarrow \gamma \eta \eta$	$(5.4 \begin{smallmatrix} +3.5 \\ -2.4 \end{smallmatrix}) \times 10^{-5}$		DESIG=269
Γ ₁₈₂	$\gamma f_2(1950) \rightarrow$ $\gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$		DESIG=144
Γ ₁₈₃	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$		DESIG=145
Γ ₁₈₄	$\gamma \phi \phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1	DESIG=98
Γ ₁₈₅	$\gamma \rho \bar{\rho}$	$(3.8 \pm 1.0) \times 10^{-4}$		DESIG=90
Γ ₁₈₆	$\gamma \eta(2225)$	$(3.3 \pm 0.5) \times 10^{-4}$		DESIG=126
Γ ₁₈₇	$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$		DESIG=125
Γ ₁₈₈	$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	$(1.98 \pm 0.33) \times 10^{-3}$		DESIG=224
Γ ₁₈₉	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.6 \pm 0.4) \times 10^{-4}$		DESIG=213
Γ ₁₉₀	$\gamma X(1835) \rightarrow \gamma \rho \bar{\rho}$	$(7.7 \begin{smallmatrix} +1.5 \\ -0.9 \end{smallmatrix}) \times 10^{-5}$		DESIG=254
Γ ₁₉₁	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \begin{smallmatrix} +2.0 \\ -1.3 \end{smallmatrix}) \times 10^{-5}$		DESIG=282
Γ ₁₉₂	$\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \begin{smallmatrix} +0.7 \\ -0.8 \end{smallmatrix}) \times 10^{-5}$		DESIG=264
Γ ₁₉₃	$\gamma (K \bar{K} \pi) [J^{PC} = 0^- +]$	$(7 \pm 4) \times 10^{-4}$	S=2.1	DESIG=176
Γ ₁₉₄	$\gamma \pi^0$	$(3.49 \begin{smallmatrix} +0.33 \\ -0.30 \end{smallmatrix}) \times 10^{-5}$		DESIG=82
Γ ₁₉₅	$\gamma \rho \bar{\rho} \pi^+ \pi^-$	$< 7.9 \times 10^{-4}$	CL=90%	DESIG=93
Γ ₁₉₆	$\gamma \Lambda \bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=200
Γ ₁₉₇	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \begin{smallmatrix} +0.60 \\ -0.30 \end{smallmatrix}) \times 10^{-4}$		DESIG=267
Γ ₁₉₈	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$		DESIG=286
Γ ₁₉₉	$\gamma f_0(2200)$			DESIG=123
Γ ₂₀₀	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$		DESIG=285
Γ ₂₀₁	$\gamma f_J(2220)$			DESIG=92
Γ ₂₀₂	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 3.9 \times 10^{-5}$	CL=90%	DESIG=136
Γ ₂₀₃	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 4.1 \times 10^{-5}$	CL=90%	DESIG=137
Γ ₂₀₄	$\gamma f_J(2220) \rightarrow \gamma \rho \bar{\rho}$	$(1.5 \pm 0.8) \times 10^{-5}$		DESIG=138
Γ ₂₀₅	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \begin{smallmatrix} +2.4 \\ -2.2 \end{smallmatrix}) \times 10^{-5}$		DESIG=270
Γ ₂₀₆	$\gamma f_0(1500) \rightarrow \gamma \pi \pi$	$(1.09 \pm 0.24) \times 10^{-4}$		DESIG=172
Γ ₂₀₇	$\gamma f_0(1500) \rightarrow \gamma \eta \eta$	$(1.7 \begin{smallmatrix} +0.6 \\ -1.4 \end{smallmatrix}) \times 10^{-5}$		DESIG=265
Γ ₂₀₈	$\gamma A \rightarrow \gamma \text{invisible}$	$[e] < 6.3 \times 10^{-6}$	CL=90%	DESIG=251
Γ ₂₀₉	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	$[f] < 2.1 \times 10^{-5}$	CL=90%	DESIG=259
Dalitz decays				
Γ ₂₁₀	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$		NODE=M070;CLUMP=G DESIG=271
Γ ₂₁₁	$\eta e^+ e^-$	$(1.16 \pm 0.09) \times 10^{-5}$		DESIG=272
Γ ₂₁₂	$\eta'(958) e^+ e^-$	$(5.81 \pm 0.35) \times 10^{-5}$		DESIG=273
Weak decays				
Γ ₂₁₃	$D^- e^+ \nu_e + \text{c.c.}$	$< 1.2 \times 10^{-5}$	CL=90%	DESIG=218
Γ ₂₁₄	$\bar{D}^0 e^+ e^- + \text{c.c.}$	$< 1.1 \times 10^{-5}$	CL=90%	DESIG=219
Γ ₂₁₅	$D_s^- e^+ \nu_e + \text{c.c.}$	$< 1.3 \times 10^{-6}$	CL=90%	DESIG=220
Γ ₂₁₆	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	$< 1.8 \times 10^{-6}$	CL=90%	DESIG=290
Γ ₂₁₇	$D^- \pi^+ + \text{c.c.}$	$< 7.5 \times 10^{-5}$	CL=90%	DESIG=241
Γ ₂₁₈	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	$< 1.7 \times 10^{-4}$	CL=90%	DESIG=242
Γ ₂₁₉	$\bar{D}^0 \bar{K}^{*0} + \text{c.c.}$	$< 2.5 \times 10^{-6}$	CL=90%	DESIG=275
Γ ₂₂₀	$D_s^- \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=243
Γ ₂₂₁	$D_s^- \rho^+ + \text{c.c.}$	$< 1.3 \times 10^{-5}$	CL=90%	DESIG=274

**Charge conjugation (C), Parity (P),
Lepton Family number (LF) violating modes**

Γ_{222}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$	CL=90%
Γ_{223}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{224}	$e^\pm\mu^\mp$	LF	< 1.6	$\times 10^{-7}$	CL=90%
Γ_{225}	$e^\pm\tau^\mp$	LF	< 8.3	$\times 10^{-6}$	CL=90%
Γ_{226}	$\mu^\pm\tau^\mp$	LF	< 2.0	$\times 10^{-6}$	CL=90%

NODE=M070;CLUMP=D

DESIG=80

DESIG=277

DESIG=177

DESIG=178

DESIG=179

Other decays

Γ_{227}	invisible		< 7	$\times 10^{-4}$	CL=90%
----------------	-----------	--	-----	------------------	--------

NODE=M070;CLUMP=F

DESIG=240

[a] For $E_\gamma > 100$ MeV.

LINKAGE=EGM

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

LINKAGE=SG

[c] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.

LINKAGE=MF

[d] See the "Note on the $\eta(1405)$ " in the $\eta(1405)$ Particle Listings.

LINKAGE=MG

[e] For a narrow state A with mass less than 960 MeV.

LINKAGE=NSA

[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

LINKAGE=NAO

$J/\psi(1S)$ PARTIAL WIDTHS

NODE=M070220

$\Gamma(\text{hadrons})$

 Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
74.1 ± 8.1	BAI	95B	BES e^+e^-
59 ± 24	BALDINI-...	75	FRAG e^+e^-
59 ± 14	BOYARSKI	75	MRK1 e^+e^-
50 ± 25	ESPOSITO	75B	FRAM e^+e^-

NODE=M070W3

NODE=M070W3

$\Gamma(e^+e^-)$

 Γ_5

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.55±0.14±0.02 OUR EVALUATION				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.71 ± 0.16	13k	¹ ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57 ± 0.19	7.8k	¹ AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14 ± 0.39		BAI	95B	BES e^+e^-
$5.36^{+0.29}_{-0.28}$		² HSUEH	92	RVUE See Υ mini-review
4.72 ± 0.35		ALEXANDER	89	RVUE See Υ mini-review
4.4 ± 0.6		² BRANDELIK	79C	DASP e^+e^-
4.6 ± 0.8		³ BALDINI-...	75	FRAG e^+e^-
4.8 ± 0.6		BOYARSKI	75	MRK1 e^+e^-
4.6 ± 1.0		ESPOSITO	75B	FRAM e^+e^-

NODE=M070W1

NODE=M070W1

→ UNCHECKED ←

¹ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

NODE=M070W1;LINKAGE=AA

² From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

NODE=M070W1;LINKAGE=F

³ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

NODE=M070W1;LINKAGE=B

$\Gamma(\mu^+\mu^-)$

 Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
5.13 ± 0.52	BAI	95B	BES e^+e^-
4.8 ± 0.6	BOYARSKI	75	MRK1 e^+e^-
5 ± 1	ESPOSITO	75B	FRAM e^+e^-

NODE=M070W2

NODE=M070W2

$\Gamma(\gamma\gamma)$

 Γ_{222}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C	DASP e^+e^-

NODE=M070W70

NODE=M070W70

$J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M070225

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel_i in the e^+e^- annihilation.

NODE=M070225

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M070G3
NODE=M070G3

••• We do not use the following data for averages, fits, limits, etc. •••

4 ± 0.8	¹ BALDINI-...	75	FRAG e^+e^-
---------	--------------------------	----	---------------

3.9 ± 0.8	¹ ESPOSITO	75B	FRAM e^+e^-
-----------	-----------------------	-----	---------------

¹Data redundant with branching ratios or partial widths above.

NODE=M070G3;LINKAGE=S

 $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M070G1
NODE=M070G1

332.3 ± 6.4 ± 4.8	ANASHIN	10	KEDR 3.097 $e^+e^- \rightarrow e^+e^-$
--------------------------	---------	----	--

••• We do not use the following data for averages, fits, limits, etc. •••

350 ± 20	BRANDELIK	79C	DASP e^+e^-
----------	-----------	-----	---------------

320 ± 70	¹ BALDINI-...	75	FRAG e^+e^-
----------	--------------------------	----	---------------

340 ± 90	¹ ESPOSITO	75B	FRAM e^+e^-
----------	-----------------------	-----	---------------

360 ± 100	¹ FORD	75	SPEC e^+e^-
-----------	-------------------	----	---------------

¹Data redundant with branching ratios or partial widths above.

NODE=M070G1;LINKAGE=S

 $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M070G2
NODE=M070G2

334 ± 5 OUR AVERAGE				
----------------------------	--	--	--	--

331.8 ± 5.2 ± 6.3		ANASHIN	10	KEDR 3.097 $e^+e^- \rightarrow \mu^+\mu^-$
-------------------	--	---------	----	--

338.4 ± 5.8 ± 7.1	13k	ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
-------------------	-----	-------	-----	--

330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
-------------------	------	--------	----	--

••• We do not use the following data for averages, fits, limits, etc. •••

510 ± 90		DASP	75	DASP e^+e^-
----------	--	------	----	---------------

380 ± 50		¹ ESPOSITO	75B	FRAM e^+e^-
----------	--	-----------------------	-----	---------------

¹Data redundant with branching ratios or partial widths above.

NODE=M070G2;LINKAGE=S

 $\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{12}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------------------	------	-------------	------	---------

NODE=M070G8
NODE=M070G8

2.2 ± 0.3 ± 0.2	170	AUBERT	06D	BABR 10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$
------------------------	-----	--------	-----	--

 $\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{13}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M070G24
NODE=M070G24

53.6 ± 5.0 ± 0.4	788	¹ AUBERT	07AU	BABR 10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
-------------------------	-----	---------------------	------	---

¹AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$] = $47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G24;LINKAGE=AU

 $\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M070G02
NODE=M070G02

33 ± 4 ± 1	317 ± 23	^{1,2} AUBERT	07AK	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
-------------------	----------	-----------------------	------	---

¹Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+\pi^-) = 2/3$.

²AUBERT 07AK reports [$\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(K_2^*(1430) \rightarrow K\pi)$] = $16.4 \pm 1.1 \pm 1.4$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G02;LINKAGE=AE
NODE=M070G02;LINKAGE=UB $\Gamma(K^*(892)^0\bar{K}_2(1770)^0 + \text{c.c.}) \Rightarrow K^*(892)^0 K^-\pi^+ + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M070G03
NODE=M070G03

3.8 ± 0.4 ± 0.3	110 ± 14	¹ AUBERT	07AK	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
------------------------	----------	---------------------	------	---

¹Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+\pi^-) = 2/3$.

NODE=M070G03;LINKAGE=AE

 $\Gamma(K^+K^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M070G18
NODE=M070G18

29.0 ± 1.7 ± 1.3	AUBERT	08S	BABR 10.6 $e^+e^- \rightarrow K^+K^*(892)^-\gamma$
-------------------------	--------	-----	--

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \rightarrow K^+ K^- \pi^0 \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{23} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.96 ± 0.85 ± 0.70	155	AUBERT	08S	BABR 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁰ γ

NODE=M070G20
NODE=M070G20

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{24} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.76 ± 1.70 ± 1.00	89	AUBERT	08S	BABR 10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ

NODE=M070G21
NODE=M070G21

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{25} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
26.6 ± 2.5 ± 1.5	AUBERT	08S	BABR 10.6 e ⁺ e ⁻ → K ⁰ $\bar{K}^*(892)^0$ γ

NODE=M070G19
NODE=M070G19

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{26} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
17.70 ± 1.70 ± 1.00	94	AUBERT	08S	BABR 10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ

NODE=M070G22
NODE=M070G22

$$\Gamma(\omega K \bar{K}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{35} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.70 ± 1.98 ± 0.03	24	¹ AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → ω K ⁺ K ⁻ γ

NODE=M070G29
NODE=M070G29

¹ AUBERT 07AU reports [Γ(J/ψ(1S) → ω K \bar{K}) × Γ(J/ψ(1S) → e⁺e⁻) / Γ_{total}] × [B(ω(782) → π⁺π⁻π⁰)] = 3.3 ± 1.3 ± 1.2 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.2 ± 0.7) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G29;LINKAGE=AU

$$\Gamma(\phi 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{37} \Gamma_5 / \Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.96 ± 0.19 ± 0.01	35	¹ AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → φ 2(π ⁺ π ⁻) γ

NODE=M070G10
NODE=M070G10

¹ AUBERT 06D reports [Γ(J/ψ(1S) → φ 2(π⁺π⁻)) × Γ(J/ψ(1S) → e⁺e⁻) / Γ_{total}] × [B(φ(1020) → K⁺K⁻)] = (0.47 ± 0.09 ± 0.03) × 10⁻² keV which we divide by our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G10;LINKAGE=AU

$$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{46} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.8 ± 0.4 OUR AVERAGE				
4.52 ± 0.48 ± 0.04	254 ± 23	¹ SHEN	09	BELL 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ
5.33 ± 0.71 ± 0.05	103	² AUBERT, BE	06D	BABR 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ

NODE=M070G14
NODE=M070G14

¹ SHEN 09 reports 4.50 ± 0.41 ± 0.26 eV from a measurement of [Γ(J/ψ(1S) → φ π⁺π⁻) × Γ(J/ψ(1S) → e⁺e⁻) / Γ_{total}] × [B(φ(1020) → K⁺K⁻)] assuming B(φ(1020) → K⁺K⁻) = (49.2 ± 0.6) × 10⁻², which we rescale to our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G14;LINKAGE=SH

² AUBERT, BE 06D reports [Γ(J/ψ(1S) → φ π⁺π⁻) × Γ(J/ψ(1S) → e⁺e⁻) / Γ_{total}] × [B(φ(1020) → K⁺K⁻)] = 2.61 ± 0.30 ± 0.18 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G14;LINKAGE=AU

$$\Gamma(\phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{47} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.15 ± 0.88 ± 0.03	23	¹ AUBERT, BE	06D	BABR 10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁰ π ⁰ γ

NODE=M070G15
NODE=M070G15

¹ AUBERT, BE 06D reports [Γ(J/ψ(1S) → φ π⁰π⁰) × Γ(J/ψ(1S) → e⁺e⁻) / Γ_{total}] × [B(φ(1020) → K⁺K⁻)] = 1.54 ± 0.40 ± 0.16 eV which we divide by our best value B(φ(1020) → K⁺K⁻) = (48.9 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G15;LINKAGE=AU

$$\Gamma(\phi \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{50} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.1 ± 2.7 ± 0.4	6	¹ AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → φ η γ

NODE=M070G28
NODE=M070G28

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi \eta) \cdot B(\phi \rightarrow K^+ K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.

NODE=M070G28;LINKAGE=AU

$$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{57} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.21 ± 0.23 OUR AVERAGE Error includes scale factor of 1.2.

1.48 ± 0.27 ± 0.09	60 ± 11	¹ SHEN 09 BELL	10.6	$e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
1.02 ± 0.24 ± 0.01	20 ± 5	² AUBERT 07AK BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Multiplied by 2/3 to take into account the $\phi \pi^+ \pi^-$ mode only. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G05
NODE=M070G05

NODE=M070G05;LINKAGE=SH

NODE=M070G05;LINKAGE=UB

$$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{58} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.96 ± 0.40 ± 0.01 7.0 ± 2.8 ¹ AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

¹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G06
NODE=M070G06

NODE=M070G06;LINKAGE=UB

$$\Gamma(\eta \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{68} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.23 ± 0.97 ± 0.03 9 ¹ AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+ \pi^- \pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best value $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (22.92 \pm 0.28) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G25
NODE=M070G25

NODE=M070G25;LINKAGE=AU

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{15} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.28 ± 0.40 ± 0.11 25 ± 8 ¹ AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

NODE=M070G01
NODE=M070G01

NODE=M070G01;LINKAGE=AE

$$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{42} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.05 ± 0.73^{+0.04}_{-0.14} 44 ± 7 ^{1,2} AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Using $B(\phi \rightarrow (K + K)^-) = (49.3 \pm 0.6)\%$.

² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = 3.41 \pm 0.55 \pm 0.28$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G07
NODE=M070G07

NODE=M070G07;LINKAGE=AE
NODE=M070G07;LINKAGE=UB

$$\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{94} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

303 ± 5 ± 18 4990 AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$

NODE=M070G23
NODE=M070G23

$$\Gamma(\pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{96} \Gamma_5 / \Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

0.122 ± 0.005 ± 0.008 AUBERT,B 04N BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$

NODE=M070G5
NODE=M070G5

$$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{97} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

107.0 ± 4.3 ± 6.4 768 AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$

NODE=M070G27
NODE=M070G27

$$\Gamma(\pi^+ \pi^- K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{99} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

36.3 ± 1.3 ± 2.1 1586 ± 58 AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

33.6 ± 2.7 ± 2.7 233 ¹ AUBERT 05D BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹ Superseded by AUBERT 07AK.

NODE=M070G12
NODE=M070G12

NODE=M070G12;LINKAGE=AU

$\Gamma(\pi^+\pi^-K^+K^-\eta) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{100}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9±3.9±0.1	73	¹ AUBERT 07AU	BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ
¹ AUBERT 07AU reports [Γ(J/ψ(1S) → π ⁺ π ⁻ K ⁺ K ⁻ η) × Γ(J/ψ(1S) → e ⁺ e ⁻)/Γ _{total}] × [B(η → 2γ)] = 10.2 ± 1.3 ± 0.8 eV which we divide by our best value B(η → 2γ) = (39.41 ± 0.20) × 10 ⁻² . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G30
NODE=M070G30

NODE=M070G30;LINKAGE=AU

$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{101}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.6±1.1±1.3	203 ± 16	AUBERT 07AK	BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ

NODE=M070G04
NODE=M070G04

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{103}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4±0.9±0.4		LEES 12E	BABR	10.6 e ⁺ e ⁻ → 2π ⁺ 2π ⁻ γ
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
19.5±1.4±1.3	270	¹ AUBERT 05D	BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)γ
¹ Superseded by LEES 12E.				

NODE=M070G11
NODE=M070G11

NODE=M070G11;LINKAGE=AU

$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{104}\Gamma_5/\Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.16±0.14	496	AUBERT 06D	BABR	10.6 e ⁺ e ⁻ → 3(π ⁺ π ⁻)γ

NODE=M070G6
NODE=M070G6

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{105}\Gamma_5/\Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9±0.5±1.0	761	AUBERT 06D	BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻ π ⁰)γ

NODE=M070G7
NODE=M070G7

$\Gamma(2(\pi^+\pi^-\eta)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{106}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	¹ AUBERT 07AU	BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ
¹ AUBERT 07AU reports [Γ(J/ψ(1S) → 2(π ⁺ π ⁻)η) × Γ(J/ψ(1S) → e ⁺ e ⁻)/Γ _{total}] × [B(η → 2γ)] = 5.16 ± 0.85 ± 0.39 eV which we divide by our best value B(η → 2γ) = (39.41 ± 0.20) × 10 ⁻² . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G26
NODE=M070G26

NODE=M070G26;LINKAGE=AU


$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{108}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.9±0.5 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
11.3±0.4±0.3	821	LEES 13O	BABR	e ⁺ e ⁻ → p \bar{p} γ
12.9±0.4±0.4	918	LEES 13Y	BABR	e ⁺ e ⁻ → p \bar{p} γ
12.0±0.6±0.5	438	AUBERT 06B		e ⁺ e ⁻ → p \bar{p} γ
9.7±1.7		¹ ARMSTRONG 93B	E760	p \bar{p} → e ⁺ e ⁻

NODE=M070G4
NODE=M070G4

¹ Using Γ_{total} = 85.5^{+6.1}_{-5.8} MeV.

NODE=M070G;LINKAGE=A



$$\Gamma(\Sigma^0 \bar{\Sigma}^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{121} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
6.4 ± 1.2 ± 0.6	AUBERT	07BD BABR	10.6 e ⁺ e ⁻ → Σ ⁰ Σ ⁰ γ

NODE=M070G17
NODE=M070G17

$$\Gamma(2(\pi^+ \pi^-) K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{122} \Gamma_5 / \Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.75 ± 0.23 ± 0.17	205	AUBERT	06D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ 2(π ⁺ π ⁻) γ

NODE=M070G9
NODE=M070G9

$$\Gamma(\Lambda \bar{\Lambda}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{128} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
10.7 ± 0.9 ± 0.7	AUBERT	07BD BABR	10.6 e ⁺ e ⁻ → Λ Λ̄ γ

NODE=M070G16
NODE=M070G16

$$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{131} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.11 ± 0.39 ± 0.30	156 ± 15	AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻) γ

NODE=M070G13
NODE=M070G13

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.0 ± 0.7 ± 0.6	38	¹ AUBERT	05D BABR	10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻) γ
-----------------	----	---------------------	----------	--

¹ Superseded by AUBERT 07AK.

NODE=M070G13;LINKAGE=AU

$$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{133} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78 ± 0.11 ± 0.05	462	¹ LEES	15J BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ
--------------------	-----	-------------------	----------	---

1.94 ± 0.11 ± 0.05	462	² LEES	15J BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ
--------------------	-----	-------------------	----------	---

1.42 ± 0.23 ± 0.08	51	³ LEES	13Q BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ
--------------------	----	-------------------	----------	---

¹ sin φ > 0.

² sin φ < 0.

³ Interference with non-resonant K⁺ K⁻ production not taken into account.

NODE=M070G08
NODE=M070G08

OCCUR=2

NODE=M070G08;LINKAGE=A
NODE=M070G08;LINKAGE=B
NODE=M070G08;LINKAGE=BA

J/ψ(1S) BRANCHING RATIOS

NODE=M070230

For the first four branching ratios, see also the partial widths, and (partial widths) × Γ(e⁺ e⁻)/Γ_{total} above.

NODE=M070300

$$\Gamma(\text{hadrons}) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.877 ± 0.005 OUR AVERAGE

0.878 ± 0.005	BAI	95B BES	e ⁺ e ⁻
---------------	-----	---------	-------------------------------

0.86 ± 0.02	BOYARSKI	75 MRK1	e ⁺ e ⁻
-------------	----------	---------	-------------------------------

NODE=M070R3
NODE=M070R3

$$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons}) / \Gamma_{\text{total}} \quad \Gamma_2 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.135 ± 0.003 ^{1,2} SETH 04 RVUE e⁺ e⁻

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ± 0.02	¹ BOYARSKI	75 MRK1	e ⁺ e ⁻
-------------	-----------------------	---------	-------------------------------

¹ Included in Γ(hadrons)/Γ_{total}.

² Using B(J/ψ → ℓ⁺ ℓ⁻) = (5.90 ± 0.09)% from RPP-2002 and R = 2.28 ± 0.04 determined by a fit to data from BAI 00 and BAI 02C.

NODE=M070R4
NODE=M070R4

NODE=M070R4;LINKAGE=C
NODE=M070R4;LINKAGE=SE

$$\Gamma(g g g) / \Gamma_{\text{total}} \quad \Gamma_3 / \Gamma$$

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

64.1 ± 1.0 6 M ¹ BESSON 08 CLEO ψ(2S) → π⁺ π⁻ + hadrons

¹ Calculated using the value Γ(γγg)/Γ(ggg) = 0.137 ± 0.001 ± 0.016 ± 0.004 from BESSON 08 and the PDG 08 values of B(ℓ⁺ ℓ⁻), B(virtual γ → hadrons), and B(γη_c). The statistical error is negligible and the systematic error is partially correlated with that of Γ(γγg)/Γ_{total} measurement of BESSON 08.

NODE=M070S65
NODE=M070S65

NODE=M070S65;LINKAGE=BE

$$\Gamma(\gamma g g) / \Gamma_{\text{total}} \quad \Gamma_4 / \Gamma$$

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

8.79 ± 1.05 200 k ¹ BESSON 08 CLEO ψ(2S) → π⁺ π⁻ γ + hadrons

¹ Calculated using the value Γ(γγg)/Γ(ggg) = 0.137 ± 0.001 ± 0.016 ± 0.004 from BESSON 08 and the value of Γ(ggg)/Γ_{total}. The statistical error is negligible and the systematic error is partially correlated with that of Γ(ggg)/Γ_{total} measurement of BESSON 08.

NODE=M070S66
NODE=M070S66

NODE=M070S66;LINKAGE=BE

$\Gamma(\gamma g g)/\Gamma(g g g)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_3
13.7±0.1±0.7	6 M	BESSION	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

NODE=M070S67
NODE=M070S67

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
5.971±0.032 OUR AVERAGE					
5.983±0.007±0.037	720k	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945±0.067±0.042	15k	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ±0.05 ±0.10		BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ±0.33		BAI	95B	BES	$e^+ e^-$
5.92 ±0.15 ±0.20		COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1	$e^+ e^-$

NODE=M070R1
NODE=M070R1

 $\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
8.8±1.3±0.4	¹ ARMSTRONG	96	E760	$\bar{p} p \rightarrow e^+ e^- \gamma$

NODE=M070S33
NODE=M070S33

¹ For $E_\gamma > 100$ MeV.

NODE=M070S33;LINKAGE=A

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
5.961±0.033 OUR AVERAGE					
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960±0.065±0.050	17k	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ±0.06 ±0.10		BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ±0.33		BAI	95B	BES	$e^+ e^-$
5.90 ±0.15 ±0.19		COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1	$e^+ e^-$

NODE=M070R2
NODE=M070R2

 $\Gamma(e^+ e^-)/\Gamma(\mu^+ \mu^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_7
1.0016±0.0031 OUR AVERAGE				
1.0022±0.0044±0.0048	¹ AULCHENKO	14	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017±0.0017±0.0033	² ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 ±0.021 ±0.013	³ ANASHIN	10	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 ±0.012 ±0.006	LI	05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.011 ±0.013 ±0.016	BAI	98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 ±0.07	BAI	95B	BES	$e^+ e^-$
1.00 ±0.05	BOYARSKI	75	MRK1	$e^+ e^-$
0.91 ±0.15	ESPOSITO	75B	FRAM	$e^+ e^-$
0.93 ±0.10	FORD	75	SPEC	$e^+ e^-$

NODE=M070R5
NODE=M070R5

¹ From 235.3k $J/\psi \rightarrow e^+ e^-$ and 156.6k $J/\psi \rightarrow \mu^+ \mu^-$ observed events.

² Not independent of the corresponding measurements of $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$.

³ Not independent of the corresponding measurements of $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$.

NODE=M070R5;LINKAGE=A
NODE=M070R5;LINKAGE=AB

NODE=M070R5;LINKAGE=AN

———— HADRONIC DECAYS ————

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
1.69 ±0.15 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.	
2.18 ±0.19		^{1,2} AUBERT,B	04N	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
2.184±0.005±0.201	220k	^{2,3} BAI	04H	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
2.091±0.021±0.116		^{2,4} BAI	04H	BES	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
1.21 ±0.20		BAI	96D	BES	$e^+ e^- \rightarrow \rho\pi$
1.42 ±0.01 ±0.19		COFFMAN	88	MRK3	$e^+ e^-$
1.3 ±0.3	150	FRANKLIN	83	MRK2	$e^+ e^-$
1.6 ±0.4	183	ALEXANDER	78	PLUT	$e^+ e^-$
1.33 ±0.21		BRANDELIK	78B	DASP	$e^+ e^-$
1.0 ±0.2	543	BARTEL	76	CNTR	$e^+ e^-$
1.3 ±0.3	153	JEAN-MARIE	76	MRK1	$e^+ e^-$


NODE=M070305

NODE=M070R20
NODE=M070R20

OCCUR=2

- ¹ From the ratio of $\Gamma(e^+e^-)B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-)B(\mu^+\mu^-)$ (AUBERT 04).
- ² Not independent of their $B(\pi^+\pi^-\pi^0)$.
- ³ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.
- ⁴ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

NODE=M070R20;LINKAGE=AU
 NODE=M070R20;LINKAGE=BU
 NODE=M070R20;LINKAGE=BA
 NODE=M070R20;LINKAGE=BI



$\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_8
$0.328 \pm 0.005 \pm 0.027$	COFFMAN	88	MRK3	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.35 ± 0.08	ALEXANDER	78	PLUT	e^+e^-
0.32 ± 0.08	BRANDELIK	78B	DASP	e^+e^-
0.39 ± 0.11	BARTEL	76	CNTR	e^+e^-
0.37 ± 0.09	JEAN-MARIE	76	MRK1	e^+e^-

NODE=M070R21
 NODE=M070R21

$\Gamma(a_2(1320)\rho)/\Gamma_{total}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
10.9 ± 2.2	OUR AVERAGE				
11.7 ± 0.7 ± 2.5	7584	AUGUSTIN	89	DM2	$J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$
8.4 ± 4.5	36	VANNUCCI	77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

NODE=M070R43
 NODE=M070R43

$\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{total}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
85 ± 34	140	VANNUCCI	77	MRK1	$e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$

NODE=M070R26
 NODE=M070R26

$\Gamma(\omega\pi^+\pi^-\pi^0)/\Gamma_{total}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
$0.40 \pm 0.06 \pm 0.04$	170	¹ AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$
¹ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.					

NODE=M070R76
 NODE=M070R76

NODE=M070R76;LINKAGE=EE

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{total}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
8.6 ± 0.7	OUR AVERAGE	Error includes scale factor of 1.1.			
9.7 ± 0.6 ± 0.6	788	¹ AUBERT	07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
7.0 ± 1.6	18058	AUGUSTIN	89	DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8 ± 1.6	215	BURMESTER	77D	PLUT	e^+e^-
6.8 ± 1.9	348	VANNUCCI	77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

NODE=M070R24
 NODE=M070R24

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 47.8 \pm 3.1 \pm 3.2$ eV.

NODE=M070R24;LINKAGE=AU

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{14}/Γ VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

4.3±0.6 OUR AVERAGE4.3±0.2±0.6 5860 AUGUSTIN 89 DM2 e^+e^- 4.0±1.6 70 BURMESTER 77D PLUT e^+e^-

••• We do not use the following data for averages, fits, limits, etc. •••

1.9±0.8 81 VANNUCCI 77 MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$ NODE=M070R28
NODE=M070R28 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ VALUE (units 10^{-4}) CL% EVTS DOCUMENT ID TECN COMMENT**2.3±0.7±0.1** 25 ± 8 ¹ AUBERT 07AK BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^- K^+ K^- \gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

<5 90 VANNUCCI 77 MRK1 $e^+e^- \rightarrow \pi^+\pi^- K^+ K^-$ ¹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (1.28 \pm 0.40 \pm 0.11) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.NODE=M070R46
NODE=M070R46

NODE=M070R46;LINKAGE=BE

 $\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$ Γ_{16}/Γ VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT**1.00±0.19^{+0.11}_{-0.32}** 323 ABLIKIM 10E BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$ NODE=M070S73
NODE=M070S73 $\Gamma(K^*(892)^\pm K^*(800)^\mp)/\Gamma_{\text{total}}$ Γ_{17}/Γ VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT**1.09±0.18^{+0.94}_{-0.54}** 655 ABLIKIM 10E BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$ NODE=M070S74
NODE=M070S74 $\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT**1.15±0.13±0.22** 209 ABLIKIM 10C BES2 $J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$ NODE=M070S69
NODE=M070S69 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{19}/Γ VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT**6.0±0.6 OUR AVERAGE**5.9±0.6±0.2 317 ± 23 ^{1,2} AUBERT 07AK BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^- K^+ K^- \gamma$ 6.7±2.6 40 VANNUCCI 77 MRK1 $e^+e^- \rightarrow \pi^+\pi^- K^+ K^-$ ¹ Using $B(K_2^*(1430)^0 \rightarrow K\pi) = (49.9 \pm 1.2)\%$.² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (32.9 \pm 2.3 \pm 2.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.NODE=M070R48
NODE=M070R48NODE=M070R48;LINKAGE=AU
NODE=M070R48;LINKAGE=BE $\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{21}/Γ VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT**61 ± 9 OUR AVERAGE**62.0 ± 6.8 ± 10.6 899 ± 98 ABLIKIM 08E BES2 $J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$ 65.3 ± 10.2 ± 13.5 176 ± 28 ABLIKIM 08E BES2 $J/\psi \rightarrow \omega K^+ K^- \pi^0$ 53 ± 14 ± 14 530 ± 140 BECKER 87 MRK3 $e^+e^- \rightarrow \text{hadrons}$ NODE=M070S2
NODE=M070S2

OCCUR=2

 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{22}/Γ VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT**5.12±0.30 OUR AVERAGE**5.2 ± 0.4 ± 0.1 ¹ AUBERT 08S BABR 10.6 $e^+e^- \rightarrow K^+ K^*(892)^- \gamma$ 4.57±0.17±0.70 2285 JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$ 5.26±0.13±0.53 COFFMAN 88 MRK3 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$

••• We do not use the following data for averages, fits, limits, etc. •••

2.6 ± 0.6 24 FRANKLIN 83 MRK2 $J/\psi \rightarrow K^+ K^- \pi^0$ 3.2 ± 0.6 48 VANNUCCI 77 MRK1 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$ 4.1 ± 1.2 39 BRAUNSCH... 76 DASP $J/\psi \rightarrow K^\pm X$ NODE=M070S15
NODE=M070S15

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (29.0 \pm 1.7 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S15;LINKAGE=AU

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.97±0.20±0.05	155	¹ AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$

NODE=M070R09
NODE=M070R09

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (10.96 \pm 0.85 \pm 0.70) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R09;LINKAGE=AU

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0±0.4±0.1	89	¹ AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

NODE=M070S58
NODE=M070S58

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (16.76 \pm 1.70 \pm 1.00) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S58;LINKAGE=AU

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.39±0.31 OUR AVERAGE				

NODE=M070S16
NODE=M070S16

4.8 ±0.5 ±0.1		¹ AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$
3.96 ±0.15 ±0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
4.33 ±0.12 ±0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ±0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
----------	----	----------	---------	--

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (26.6 \pm 2.5 \pm 1.5) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S16;LINKAGE=AU

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma(K^+ K^*(892)^- + \text{c.c.})$ Γ_{25}/Γ_{22}

VALUE	DOCUMENT ID	TECN	COMMENT
0.82±0.05±0.09	COFFMAN 88 MRK3		$J/\psi \rightarrow K \bar{K}^*(892) + \text{c.c.}$

NODE=M070S17
NODE=M070S17

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2±0.4±0.1	94	¹ AUBERT 08S	BABR	10.6 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

NODE=M070S59
NODE=M070S59

¹AUBERT 08S reports $[\Gamma(J/\psi(1S) \rightarrow K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (17.70 \pm 1.70 \pm 1.00) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S59;LINKAGE=AU

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.8±0.8±1.2	¹ BAI 99C	BES	$e^+ e^-$

NODE=M070S35
NODE=M070S35

¹Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

NODE=M070S35;LINKAGE=M3

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM 06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

NODE=M070S52
NODE=M070S52

¹A $K_0^*(800)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

NODE=M070S52;LINKAGE=AB

$\Gamma(\omega \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.4±0.3±0.7	509	AUGUSTIN 89	DM2	$J/\psi \rightarrow \pi^+ \pi^- 3\pi^0$

NODE=M070S26
NODE=M070S26

$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**30 ± 5 OUR AVERAGE**31 ± 6 4600
29 ± 7 87

DOCUMENT ID TECN COMMENT

AUGUSTIN 89 DM2 $J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
BURMESTER 77D PLUT $e^+ e^-$ Γ_{30}/Γ NODE=M070R49
NODE=M070R49 $\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**34 ± 5 OUR AVERAGE**37.7 ± 0.8 ± 5.8 1972 ± 41
29.5 ± 1.4 ± 7.0 879 ± 41

DOCUMENT ID TECN COMMENT

ABLIKIM 08E BES2 $e^+ e^- \rightarrow J/\psi$
BECKER 87 MRK3 $e^+ e^- \rightarrow \text{hadrons}$ Γ_{31}/Γ NODE=M070S1
NODE=M070S1 $\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**23 ± 3 ± 5**

229

DOCUMENT ID TECN COMMENT

AUGUSTIN 89 DM2 $e^+ e^-$ Γ_{32}/Γ NODE=M070S28
NODE=M070S28 $\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**21.8 ± 2.2 ± 3.4**

232 ± 23

DOCUMENT ID TECN COMMENT

ABLIKIM 08E BES2 $e^+ e^- \rightarrow J/\psi$ Γ_{33}/Γ NODE=M070S57
NODE=M070S57 $\Gamma(\phi K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**21.8 ± 2.3 OUR AVERAGE**20.8 ± 2.7 ± 3.9 195 ± 25
29.6 ± 3.7 ± 4.7 238 ± 30
20.7 ± 2.4 ± 3.0
20 ± 3 ± 3 155 ± 20

DOCUMENT ID TECN COMMENT

ABLIKIM 08E BES2 $J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
ABLIKIM 08E BES2 $J/\psi \rightarrow \phi K^+ K^- \pi^0$
FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$
BECKER 87 MRK3 $e^+ e^- \rightarrow \text{hadrons}$ Γ_{34}/Γ NODE=M070S4
NODE=M070S4

OCCUR=2

 $\Gamma(\omega K \bar{K})/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**17.0 ± 3.2 OUR AVERAGE**13.6 ± 5.0 ± 1.0 24
19.8 ± 2.1 ± 3.9
16 ± 10 22

DOCUMENT ID TECN COMMENT

¹ AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow \omega K^+ K^- \gamma$
² FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$
FELDMAN 77 MRK1 $e^+ e^-$ Γ_{35}/Γ NODE=M070R27
NODE=M070R27¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \omega K^+ K^-) \cdot B(\eta \rightarrow 3\pi) = 3.3 \pm 1.3 \pm 0.2 \text{ eV}$.² Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.NODE=M070R27;LINKAGE=AU
NODE=M070R27;LINKAGE=B $\Gamma(\omega f_0(1710) \rightarrow \omega K \bar{K})/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**4.8 ± 1.1 ± 0.3**1,2 FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

DOCUMENT ID TECN COMMENT

1,2 FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$ Γ_{36}/Γ NODE=M070S25
NODE=M070S25¹ Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios.NODE=M070S25;LINKAGE=F
NODE=M070S25;LINKAGE=G $\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**16.6 ± 2.3 OUR AVERAGE**17.3 ± 3.3 ± 1.2 35
16.0 ± 1.0 ± 3.0

DOCUMENT ID TECN COMMENT

¹ AUBERT 06D BABR 10.6 $e^+ e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$ Γ_{37}/Γ NODE=M070R35
NODE=M070R35¹ Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04 \text{ keV}$.

NODE=M070R35;LINKAGE=EE

 $\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS**1.58 ± 0.23 ± 0.40**

332

DOCUMENT ID TECN COMMENT

EATON 84 MRK2 $e^+ e^-$ Γ_{38}/Γ NODE=M070R70
NODE=M070R70 $\Gamma(\omega \eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS**1.74 ± 0.20 OUR AVERAGE**2.352 ± 0.273 5k
1.44 ± 0.40 ± 0.14 13
1.43 ± 0.10 ± 0.21 378
1.71 ± 0.08 ± 0.20

DOCUMENT ID TECN COMMENT


Error includes scale factor of 1.6. See the ideogram below.
¹ ABLIKIM 06F BES2 $J/\psi \rightarrow \omega \eta$
² AUBERT 06D BABR 10.6 $e^+ e^- \rightarrow \omega \eta \gamma$
JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$
COFFMAN 88 MRK3 $e^+ e^- \rightarrow 3\pi \eta$ Γ_{39}/Γ NODE=M070R30
NODE=M070R30

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

² Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

NODE=M070R30;LINKAGE=BL

NODE=M070R30;LINKAGE=EE



$\Gamma(\phi K \bar{K})/\Gamma_{\text{total}}$

Γ_{40}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
18.3± 2.4 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
21.4± 0.4±2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
48 $^{+20}_{-16}$ ±6	9.0 $^{+3.7}_{-3.0}$	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6± 0.8±2.1		³ FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
18 ± 8	14	FELDMAN	77 MRK1	e^+e^-


NODE=M070R36
NODE=M070R36

¹ We have multiplied K^+K^- measurement by 2 to obtain $K\bar{K}$.

² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

³ Addition of ϕK^+K^- and $\phi K^0\bar{K}^0$ branching ratios.

NODE=M070R36;LINKAGE=AA
NODE=M070R36;LINKAGE=CC
NODE=M070R36;LINKAGE=A



$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K})/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.6±0.2±0.6	1,2 FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S24
 NODE=M070S24

¹ Including interference with $f_2'(1525)$.

² Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.

NODE=M070S24;LINKAGE=D
 NODE=M070S24;LINKAGE=E

 $\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.72±0.13±0.02	44 ± 7	1,2	AUBERT 07AK	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M070R39
 NODE=M070R39

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.45	90	FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$
< 0.37	90	VANNUCCI	77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

¹ Using $B(f_2(1270) \rightarrow \pi \pi) = (84.8_{-1.2}^{+2.4})\%$

² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (4.02 \pm 0.65 \pm 0.33) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R39;LINKAGE=AU
 NODE=M070R39;LINKAGE=BE

 $\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10±0.09±0.28	233	EATON	84	MRK2 $e^+ e^-$

NODE=M070R66
 NODE=M070R66

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ (\text{or c.c.}))/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10±0.12 OUR AVERAGE				
1.23±0.07±0.30	0.8k	ABLIKIM 12P	BES2	$J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
1.50±0.08±0.38	1k	ABLIKIM 12P	BES2	$J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
1.00±0.04±0.21	0.6k	HENRARD 87	DM2	$e^+ e^- \rightarrow \Sigma^{*-}$
1.19±0.04±0.25	0.7k	HENRARD 87	DM2	$e^+ e^- \rightarrow \Sigma^{*+}$
0.86±0.18±0.22	56	EATON 84	MRK2	$e^+ e^- \rightarrow \Sigma^{*-}$
1.03±0.24±0.25	68	EATON 84	MRK2	$e^+ e^- \rightarrow \Sigma^{*+}$

NODE=M070R67
 NODE=M070R67

OCCUR=2

OCCUR=2

OCCUR=2

 $\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4 OUR AVERAGE				Error includes scale factor of 2.7.
12.3±0.6±2.0		1,2 FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
4.8±1.8	46	1 GIDAL 81	MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

NODE=M070R40
 NODE=M070R40

¹ Re-evaluated using $B(f_2'(1525) \rightarrow K \bar{K}) = 0.713$.

² Including interference with $f_0(1710)$.

NODE=M070R40;LINKAGE=B
 NODE=M070R40;LINKAGE=C

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.94±0.09 OUR AVERAGE				Error includes scale factor of 1.2.
0.96±0.13	103	1 AUBERT, BE 06D	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
1.09±0.02±0.13		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
0.78±0.03±0.12		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ± 0.9	23	FELDMAN 77	MRK1	$e^+ e^-$

NODE=M070R34
 NODE=M070R34

¹ Derived by us. AUBERT, BE 06D measures $\Gamma(J/\psi \rightarrow e^+ e^-) \times B(J/\psi \rightarrow \phi \pi^+ \pi^-) \times B(\phi \rightarrow K^+ K^-) = (2.61 \pm 0.30 \pm 0.18)$ eV

NODE=M070R34;LINKAGE=AU

 $\Gamma(\phi \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.56±0.16	23	1 AUBERT, BE 06D	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

NODE=M070S44
 NODE=M070S44

¹ Derived by us. AUBERT, BE 06D measures $\Gamma(J/\psi \rightarrow e^+ e^-) \times B(J/\psi \rightarrow \phi \pi^0 \pi^0) \times B(\phi \rightarrow K^+ K^-) = (1.54 \pm 0.40 \pm 0.16)$ eV

NODE=M070S44;LINKAGE=AU

 $\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2±0.8 OUR AVERAGE				
7.4±0.6±1.4	227 ± 19	ABLIKIM 08E	BES2	$e^+ e^- \rightarrow J/\psi$
7.4±0.9±1.1		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
7 ± 0.6±1.0	163 ± 15	BECKER 87	MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S3
 NODE=M070S3

$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.8^{+1.9}_{-1.6} \pm 1.7$	111^{+31}_{-26}	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S5
 NODE=M070S5


 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
$1.4 \pm 0.6 \pm 0.1$	6	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \phi\eta\gamma$
$0.898 \pm 0.024 \pm 0.089$		ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$
$0.64 \pm 0.04 \pm 0.11$	346	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$0.661 \pm 0.045 \pm 0.078$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \eta$

NODE=M070R37
 NODE=M070R37

$$^1 \text{AUBERT 07AU quotes } \Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+ K^-) \cdot B(\eta \rightarrow \gamma\gamma) = 0.84 \pm 0.37 \pm 0.05 \text{ eV.}$$

NODE=M070R37;LINKAGE=AU

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.20 \pm 0.12 \pm 0.21$	206	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S64
 NODE=M070S64

 $\Gamma(\Xi(1530)^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.59 \pm 0.09 \pm 0.12$	75 ± 11	HENRARD	87 DM2	$e^+ e^-$

NODE=M070S9
 NODE=M070S9

 $\Gamma(pK^- \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.51 \pm 0.26 \pm 0.18$	89	EATON	84 MRK2	$e^+ e^-$

NODE=M070R74
 NODE=M070R74


 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.05 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
$0.538 \pm 0.012 \pm 0.065$	2090	¹ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\pi^0$
$0.360 \pm 0.028 \pm 0.054$	222	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$0.482 \pm 0.019 \pm 0.064$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^0 \pi^+ \pi^- \pi^0$

NODE=M070R32
 NODE=M070R32

$$^1 \text{Using } B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%.$$

NODE=M070R32;LINKAGE=BL




$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$

Γ_{55}/Γ

NODE=M070R38
NODE=M070R38

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.40 ±0.07 OUR AVERAGE					Error includes scale factor of 2.1. See the ideogram below.
0.546±0.031±0.056			ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadr}$
0.41 ±0.03 ±0.08		167	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.308±0.034±0.036			COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\eta'$
••• We do not use the following data for averages, fits, limits, etc. •••					
< 1.3		90	VANNUCCI	77 MRK1	e^+e^-



$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$

Γ_{56}/Γ

NODE=M070R41
NODE=M070R41

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2±0.9 OUR AVERAGE				Error includes scale factor of 1.9.
4.6±0.4±0.8		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6±0.6	50	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+K^-K^+K^-$
¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.				

NODE=M070R41;LINKAGE=A

$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{57} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.182 ± 0.042 ± 0.005	19.5 ± 4.5	1,2 AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M070S02
NODE=M070S02

¹ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (1.01 \pm 0.22 \pm 0.08) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S02;LINKAGE=AU
NODE=M070S02;LINKAGE=BE

 $\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{58} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.171 ± 0.073 ± 0.004	7.0 ± 2.8	1,2 AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

NODE=M070S03
NODE=M070S03

¹ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

² AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (0.95 \pm 0.39 \pm 0.10) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S03;LINKAGE=AU
NODE=M070S03;LINKAGE=BE

 $\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{59} / Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.50 ± 0.80 ± 0.61	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S97
NODE=M070S97

 $\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{60} / Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67 ± 0.50 ± 0.24	70	ABLIKIM	15P BESE	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S98
NODE=M070S98

 $\Gamma(\eta \phi f_0(980) \rightarrow \eta \phi \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{61} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.23 ± 0.75 ± 0.73	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$

NODE=M070R08
NODE=M070R08

 $\Gamma(\phi a_0(980)^0 \rightarrow \phi \eta \pi^0) / \Gamma_{\text{total}}$ Γ_{62} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
5.0 ± 2.7 ± 2.5	1 ABLIKIM	11D BES3	$J/\psi \rightarrow \phi \eta \pi^0$

NODE=M070S75
NODE=M070S75

¹ Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops.

NODE=M070S75;LINKAGE=AB

 $\Gamma(\Xi(1530)^0 \Xi^0) / \Gamma_{\text{total}}$ Γ_{63} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.12 ± 0.07	24 ± 9	HENRARD	87 DM2	$e^+ e^-$

NODE=M070S10
NODE=M070S10

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.})) / \Gamma_{\text{total}}$ Γ_{64} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.05 OUR AVERAGE				

NODE=M070R68
NODE=M070R68

0.30 ± 0.03 ± 0.07 74 ± 8 HENRARD 87 DM2 $e^+ e^- \rightarrow \Sigma^{*-}$

0.34 ± 0.04 ± 0.07 77 ± 9 HENRARD 87 DM2 $e^+ e^- \rightarrow \Sigma^{*+}$

0.29 ± 0.11 ± 0.10 26 EATON 84 MRK2 $e^+ e^- \rightarrow \Sigma^{*-}$

0.31 ± 0.11 ± 0.11 28 EATON 84 MRK2 $e^+ e^- \rightarrow \Sigma^{*+}$

OCCUR=2

OCCUR=2

 $\Gamma(\phi f_1(1285)) / \Gamma_{\text{total}}$ Γ_{65} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.5 OUR AVERAGE				

NODE=M070S6
NODE=M070S6

[(2.6 ± 0.5) × 10⁻⁴ OUR 2015 AVERAGE Scale factor = 1.1]

3.4 ± 1.8 ± 1.5 1.1k ¹ ABLIKIM 15H BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

3.2 ± 0.6 ± 0.4 JOUSSET 90 DM2 $J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$

2.1 ± 0.5 ± 0.4 25 ² JOUSSET 90 DM2 $J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NEW

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.2 ± 0.1 16 BECKER 87 MRK3 $J/\psi \rightarrow \phi K \bar{K} \pi$

¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285)) / \Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta \pi^+ \pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta \pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+ \pi^- \eta$ invariant mass distribution at 1297 MeV.

NODE=M070S6;LINKAGE=A

NODE=M070S6;LINKAGE=Q

$\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{66} / Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
9.36 ± 2.31 ± 1.54	78	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S99
NODE=M070S99

 $\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{67} / Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
2.08 ± 1.63 ± 1.47	9	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S00
NODE=M070S00

 $\Gamma(\eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{68} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.40 ± 0.17 ± 0.03	9	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \eta \pi^+ \pi^- \gamma$

NODE=M070S05
NODE=M070S05

¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \eta \pi^+ \pi^-) \cdot B(\eta \rightarrow 3\pi) = 0.51 \pm 0.22 \pm 0.03$ eV.

NODE=M070S05;LINKAGE=AU

 $\Gamma(\eta \rho) / \Gamma_{\text{total}}$ Γ_{69} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.193 ± 0.023 OUR AVERAGE				
0.194 ± 0.017 ± 0.029	299	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.193 ± 0.013 ± 0.029		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^+ \pi^- \eta$

NODE=M070R22
NODE=M070R22

 $\Gamma(\omega \eta'(958)) / \Gamma_{\text{total}}$ Γ_{70} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.182 ± 0.021 OUR AVERAGE				
0.226 ± 0.043	218	¹ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \eta'$
0.18 ^{+0.10} _{-0.08} ± 0.03	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.166 ± 0.017 ± 0.019		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi \eta'$

NODE=M070R31
NODE=M070R31

¹ Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+ \pi^- \gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

NODE=M070R31;LINKAGE=BL

 $\Gamma(\omega f_0(980)) / \Gamma_{\text{total}}$ Γ_{71} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.41 ± 0.27 ± 0.47	¹ AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M070S27
NODE=M070S27

¹ Assuming $B(f_0(980) \rightarrow \pi \pi) = 0.78$.

NODE=M070S27;LINKAGE=K

 $\Gamma(\rho \eta'(958)) / \Gamma_{\text{total}}$ Γ_{72} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.105 ± 0.018 OUR AVERAGE				
0.083 ± 0.030 ± 0.012	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.114 ± 0.014 ± 0.016		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

NODE=M070R23
NODE=M070R23

 $\Gamma(a_2(1320)^\pm \pi^\mp) / \Gamma_{\text{total}}$ Γ_{73} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<43	90	BRAUNSCH...	76 DASP	$e^+ e^-$

NODE=M070R42
NODE=M070R42

 $\Gamma(K \bar{K}_2^*(1430) + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{74} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<40	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$

NODE=M070R45
NODE=M070R45

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	BRAUNSCH...	76 DASP	$e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$
-----	----	-------------	---------	--

 $\Gamma(K_1(1270)^\pm K^\mp) / \Gamma_{\text{total}}$ Γ_{75} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	¹ BAI	99C BES	$e^+ e^-$

NODE=M070S34
NODE=M070S34

¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

NODE=M070S34;LINKAGE=M2

 $\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0) / \Gamma_{\text{total}}$ Γ_{76} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<29	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

NODE=M070R47
NODE=M070R47

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$

The two different fit values of ABLIKIM 15k below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	¹ ABLIKIM	15k BES3	$e^+e^- \rightarrow J/\psi \rightarrow K^+K^-\gamma\gamma$
$0.124 \pm 0.033 \pm 0.030$	35 ± 9		² ABLIKIM	15k BES3	$e^+e^- \rightarrow J/\psi \rightarrow K^+K^-\gamma\gamma$
<6.4	90		³ ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$
<6.8	90		COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+K^-\pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+K^-\pi^0$ contributions.

³ Superseded by ABLIKIM 15k.

 Γ_{π}/Γ

NODE=M070R33

NODE=M070R33

NODE=M070R33

OCCUR=2

NODE=M070R33;LINKAGE=A

NODE=M070R33;LINKAGE=C

NODE=M070R33;LINKAGE=B

 $\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.01 \pm 0.58 \pm 0.82$		172	¹ ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17 90 ² FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ With 3.6σ significance.

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

 Γ_{78}/Γ

NODE=M070S23

NODE=M070S23

NODE=M070S23;LINKAGE=B

NODE=M070S23;LINKAGE=A

 $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	¹ VANNUCCI 77 MRK1		$e^+e^- \rightarrow \pi^+\pi^-\pi^0 K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.8 90 ¹ FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

 Γ_{79}/Γ

NODE=M070R29

NODE=M070R29

NODE=M070R29;LINKAGE=C

 $\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.9	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma\pi^0 p\bar{p}$

 Γ_{80}/Γ

NODE=M070S81

NODE=M070S81

 $\Gamma(\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.8 $\times 10^{-4}$	90	ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 Γ_{81}/Γ

NODE=M070B10

NODE=M070B10

 $\Gamma(\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.13 $\times 10^{-5}$	90	ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 Γ_{82}/Γ

NODE=M070B11

NODE=M070B11

 $\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.20 \pm 0.14 \pm 0.37$	471	ABLIKIM	15H BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 Γ_{83}/Γ

NODE=M070B12

NODE=M070B12

 $\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.52	90	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

 Γ_{84}/Γ

NODE=M070S70

NODE=M070S70

 $\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.82	90	ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20 90 HENRARD 87 DM2 e^+e^-

 Γ_{85}/Γ

NODE=M070S13

NODE=M070S13

 $\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	90	HENRARD	87 DM2	e^+e^-

 Γ_{86}/Γ

NODE=M070S14

NODE=M070S14

$\Gamma(\Lambda(1520)\bar{\Lambda} + c.c. \rightarrow \gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.1	90	ABLIKIM	12B	BES3 $J/\psi \rightarrow \Lambda\bar{\Lambda}\gamma$

NODE=M070S77
NODE=M070S77

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 \rho K^- \bar{n} + c.c.)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI	04G	BES2 e^+e^-

NODE=M070S47
NODE=M070S47

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	BAI	04G	BES2 e^+e^-

NODE=M070S48
NODE=M070S48

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	BAI	04G	BES2 e^+e^-

NODE=M070S49
NODE=M070S49

$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5.6	90	BAI	04G	BES2 e^+e^-

NODE=M070S50
NODE=M070S50

$\Gamma(\bar{\Theta}(1540)K_S^0 \rho \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	BAI	04G	BES2 e^+e^-

NODE=M070S51
NODE=M070S51

$\Gamma(\Sigma^0 \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	HENRARD	87	DM2 e^+e^-

NODE=M070S12
NODE=M070S12

STABLE HADRONS

NODE=M070307


$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.1 ± 0.5 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.		
5.46 ± 0.34 ± 0.14	4990	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+ \pi^-)\pi^0 \gamma$
3.25 ± 0.49	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
3.17 ± 0.42	147	FRANKLIN	83 MRK2	$e^+e^- \rightarrow$ hadrons
3.64 ± 0.52	1500	BURMESTER	77D PLUT	e^+e^-
4 ± 1	675	JEAN-MARIE	76 MRK1	e^+e^-

NODE=M070R9
NODE=M070R9

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = 0.303 \pm 0.005 \pm 0.018$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R9;LINKAGE=AU



$\Gamma(\omega\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-)\pi^0)$

Γ_{13}/Γ_{94}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M070R25
NODE=M070R25

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3 ¹ JEAN-MARIE 76 MRK1 e^+e^-

¹ Final state $(\pi^+\pi^-\pi^0)$ under the assumption that $\pi\pi$ is isospin 0.

NODE=M070R25;LINKAGE=J

$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{total}$

Γ_{95}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M070R11
NODE=M070R11

0.029±0.006 OUR AVERAGE

0.028±0.009	11	FRANKLIN	83	MRK2 $e^+e^- \rightarrow$ hadrons
0.029±0.007	181	JEAN-MARIE	76	MRK1 e^+e^-

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$

Γ_{96}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070R7
NODE=M070R7

21.1 ±0.7 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

21.37±0.04 ^{+0.64} _{-0.62}	1.8M	1,2 ABLIKIM	12H	BES3 $e^+e^- \rightarrow J/\psi$
23.0 ±2.0 ±0.4	256	³ AUBERT	07AU	BABR 10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
21.8 ±1.9		4,5 AUBERT,B	04N	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
21.84±0.05±2.01	220k	1,5 BAI	04H	BES e^+e^-
20.91±0.21±1.16		5,6 BAI	04H	BES e^+e^-
15 ±2	168	FRANKLIN	83	MRK2 e^+e^-

OCCUR=2

¹ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.

² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.

³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total} = 0.807 \pm 0.013$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ From the ratio of $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-) B(\mu^+\mu^-)$ (AUBERT 04).


⁵ Mostly $\rho\pi$, see also $\rho\pi$ subsection.

⁶ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

NODE=M070R;LINKAGE=BA
NODE=M070R7;LINKAGE=AB

NODE=M070R7;LINKAGE=AU

NODE=M070R;LINKAGE=AU
NODE=M070R;LINKAGE=BU
NODE=M070R;LINKAGE=BI



$\Gamma(\pi^+\pi^-\pi^0 K^+K^-)/\Gamma_{total}$

Γ_{97}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070R18
NODE=M070R18

1.79±0.29 OUR AVERAGE Error includes scale factor of 2.2.

1.93±0.14±0.05	768	¹ AUBERT	07AU	BABR 10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$
1.2 ±0.3	309	VANNUCCI	77	MRK1 e^+e^-

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0 K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = 0.1070 \pm 0.0043 \pm 0.0064$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R18;LINKAGE=AU

$\Gamma(4(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{98}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
90±30	13	JEAN-MARIE 76	MRK1	$e^+ e^-$	NODE=M070R12 NODE=M070R12

$\Gamma(\pi^+\pi^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{99}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
6.6±0.5 OUR AVERAGE					NODE=M070R16 NODE=M070R16

6.5±0.4±0.2 1.6k ¹AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+\pi^- K^+ K^- \gamma$
7.2±2.3 205 VANNUCCI 77 MRK1 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1±0.7±0.2 233 ²AUBERT 05D BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (36.3 \pm 1.3 \pm 2.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R16;LINKAGE=BE

²Superseded by AUBERT 07AK. AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (33.6 \pm 2.7 \pm 2.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R16;LINKAGE=AU

$\Gamma(\pi^+\pi^- K^+ K^- \eta)/\Gamma_{\text{total}}$					Γ_{100}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.84±0.28±0.05	73	¹ AUBERT 07AU BABR		10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$	NODE=M070S04 NODE=M070S04

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^- K^+ K^- \eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (10.2 \pm 1.3 \pm 0.8) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S04;LINKAGE=AU

$\Gamma(\pi^0 \pi^0 K^+ K^-)/\Gamma_{\text{total}}$					Γ_{101}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.45±0.31±0.06	203 ± 16	¹ AUBERT 07AK BABR		10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$	NODE=M070S01 NODE=M070S01

¹AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \pi^0 \pi^0 K^+ K^-)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (13.6 \pm 1.1 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S01;LINKAGE=BE

$\Gamma(K \bar{K} \pi)/\Gamma_{\text{total}}$					Γ_{102}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
61 ± 10 OUR AVERAGE					NODE=M070R15 NODE=M070R15
55.2±12.0	25	FRANKLIN 83	MRK2	$e^+ e^- \rightarrow K^+ K^- \pi^0$	
78.0±21.0	126	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$	

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_{103}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.57±0.30 OUR AVERAGE					NODE=M070R8 NODE=M070R8

3.53±0.12±0.29 1107 ¹ABLİKIM 05H BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \pi^+ \pi^-, J/\psi \rightarrow 2(\pi^+ \pi^-)$

4.0 ± 1.0 76 JEAN-MARIE 76 MRK1 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.51±0.34±0.09 270 ²AUBERT 05D BABR 10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \gamma$

¹Computed using $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

²AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+ e^-)] = (19.5 \pm 1.4 \pm 1.3) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+ e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.

NODE=M070R8;LINKAGE=AB

NODE=M070R8;LINKAGE=AU

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
43 ± 4 OUR AVERAGE				
43.0 ± 2.9 ± 2.8	496	¹ AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$
40 ± 20	32	JEAN-MARIE	76 MRK1	e^+e^-

NODE=M070R10
NODE=M070R10¹ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

NODE=M070R10;LINKAGE=EE

 $\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.62 ± 0.09 ± 0.19	761	¹ AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

NODE=M070R69
NODE=M070R69¹ Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

NODE=M070R69;LINKAGE=EE

 $\Gamma(2(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.29 ± 0.24 OUR AVERAGE				
2.35 ± 0.39 ± 0.20	85	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\eta)\gamma$
2.26 ± 0.08 ± 0.27	4839	ABLIKIM	05C BES2	$e^+e^- \rightarrow 2(\pi^+\pi^-\eta)$

NODE=M070S42
NODE=M070S42¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow 2(\pi^+\pi^-\eta)) \cdot B(\eta \rightarrow \gamma\gamma) = 5.16 \pm 0.85 \pm 0.39$ eV.

NODE=M070S42;LINKAGE=AU

 $\Gamma(3(\pi^+\pi^-\eta))/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.24 ± 0.96 ± 1.11	616	ABLIKIM	05C BES2	$e^+e^- \rightarrow 3(\pi^+\pi^-\eta)$

NODE=M070S43
NODE=M070S43 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.120 ± 0.029 OUR AVERAGE				
2.112 ± 0.004 ± 0.031	314k	ABLIKIM	12C BES3	e^+e^-
2.15 ± 0.16 ± 0.06	317	¹ WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
2.26 ± 0.01 ± 0.14	63316	BAI	04E BES2	$e^+e^- \rightarrow J/\psi$
1.97 ± 0.22	99	BALDINI	98 FENI	e^+e^-
1.91 ± 0.04 ± 0.30		PALLIN	87 DM2	e^+e^-
2.16 ± 0.07 ± 0.15	1420	EATON	84 MRK2	e^+e^-
2.5 ± 0.4	133	BRANDELIK	79C DASP	e^+e^-
2.0 ± 0.5		BESCH	78 BONA	e^+e^-
2.2 ± 0.2	331	² PERUZZI	78 MRK1	e^+e^-

NODE=M070R50
NODE=M070R50

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3 48 ANTONELLI 93 SPEC e^+e^- ¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R50;LINKAGE=WU

² Assuming angular distribution $(1+\cos^2\theta)$.

NODE=M070R50;LINKAGE=A

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{109}/Γ


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.08 OUR AVERAGE				Error includes scale factor of 1.1.
1.33 ± 0.02 ± 0.11	11k	ABLIKIM	09B BES2	e^+e^-
1.13 ± 0.09 ± 0.09	685	EATON	84 MRK2	e^+e^-
1.4 ± 0.4		BRANDELIK	79C DASP	e^+e^-
1.00 ± 0.15	109	PERUZZI	78 MRK1	e^+e^-

NODE=M070R52
NODE=M070R52 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
6.46 ± 0.17 ± 0.43	1435	EATON	84 MRK2	e^+e^-
3.8 ± 1.6	48	BESCH	81 BONA	e^+e^-
5.5 ± 0.6	533	PERUZZI	78 MRK1	e^+e^-

NODE=M070R54
NODE=M070R54

WEIGHTED AVERAGE
6.0±0.5 (Error scaled by 1.3)



$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{total}$

Γ_{111}/Γ

Including $p\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

NODE=M070R55
NODE=M070R55
NODE=M070R55

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ±0.9 OUR AVERAGE				Error includes scale factor of 1.9.
3.36±0.65±0.28	364	EATON	84 MRK2	e^+e^-
1.6 ±0.6	39	PERUZZI	78 MRK1	e^+e^-

$\Gamma(p\bar{p}\eta)/\Gamma_{total}$

Γ_{112}/Γ

NODE=M070R56
NODE=M070R56

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00±0.12 OUR AVERAGE				
1.91±0.02±0.17	13k	¹ ABLIKIM	09 BES2	e^+e^-
2.03±0.13±0.15	826	EATON	84 MRK2	e^+e^-
2.5 ±1.2		BRANDELIK	79C DASP	e^+e^-
2.3 ±0.4	197	PERUZZI	78 MRK1	e^+e^-

¹ From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.

NODE=M070R56;LINKAGE=AB

$\Gamma(p\bar{p}\rho)/\Gamma_{total}$

Γ_{113}/Γ

NODE=M070R57
NODE=M070R57


VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.31	90	EATON	84 MRK2	$e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{total}$

Γ_{114}/Γ

NODE=M070R58
NODE=M070R58

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98±0.10 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.90±0.02±0.09	2670	ABLIKIM	13P BES3	e^+e^-
0.98±0.03±0.14	2449	ABLIKIM	08 BES2	e^+e^-
1.10±0.17±0.18	486	EATON	84 MRK2	e^+e^-
1.6 ±0.3	77	PERUZZI	78 MRK1	e^+e^-

 **$\Gamma(\rho\bar{\rho}\eta(958))/\Gamma_{\text{total}}$** **$\Gamma_{115}/\Gamma$**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.21 ± 0.04 OUR AVERAGE				
0.200 ± 0.023 ± 0.028	265 ± 31	¹ ABLIKIM	09 BES2	e^+e^-
0.68 ± 0.23 ± 0.17	19	EATON	84 MRK2	e^+e^-
1.8 ± 0.6	19	PERUZZI	78 MRK1	e^+e^-

NODE=M070R59
NODE=M070R59¹ From the combination of $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\pi^+\pi^-\eta$ and $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\gamma\rho^0$ channels.

NODE=M070R59;LINKAGE=AB

 $\Gamma(\rho\bar{\rho}a_0(980) \rightarrow \rho\bar{\rho}\pi^0\eta)/\Gamma_{\text{total}}$ **Γ_{116}/Γ**

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
6.8 ± 1.2 ± 1.3	ABLIKIM	14N BES3	$e^+e^- \rightarrow J/\psi$

NODE=M070S94
NODE=M070S94 **$\Gamma(\rho\bar{\rho}\phi)/\Gamma_{\text{total}}$** **$\Gamma_{117}/\Gamma$**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.13 ± 0.07	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S22
NODE=M070S22 **$\Gamma(n\bar{n})/\Gamma_{\text{total}}$** **$\Gamma_{118}/\Gamma$**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.09 ± 0.16 OUR AVERAGE				
2.07 ± 0.01 ± 0.17	36k	ABLIKIM	12C BES3	e^+e^-
2.31 ± 0.49	79	BALDINI	98 FENI	e^+e^-
1.8 ± 0.9		BESCH	78 BONA	e^+e^-

NODE=M070R64
NODE=M070R64

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.90 ± 0.55	40	ANTONELLI	93 SPEC	e^+e^-
-------------	----	-----------	---------	----------

 $\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{119}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 3.6	5	BESCH	81 BONA	e^+e^-

NODE=M070R65
NODE=M070R65 **$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$** **$\Gamma_{120}/\Gamma$**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.50 ± 0.10 ± 0.22	399	ABLIKIM	080 BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070S09
NODE=M070S09 **$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$** **$\Gamma_{121}/\Gamma$**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.29 ± 0.09 OUR AVERAGE				
1.15 ± 0.24 ± 0.03		¹ AUBERT	07BD BABR	$10.6 e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$
1.33 ± 0.04 ± 0.11	1779	ABLIKIM	06 BES2	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 ± 0.04 ± 0.23	884 ± 30	PALLIN	87 DM2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 ± 0.16 ± 0.25	90	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

NODE=M070R63
NODE=M070R63

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.4 ±2.6 3 BESCH 81 BONA $e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

¹AUBERT 07BD reports $[\Gamma(J/\psi(1S) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (6.4 \pm 1.2 \pm 0.6) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R63;LINKAGE=AU

$\Gamma(2(\pi^+\pi^-)K^+K^-)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

47 ± 7 OUR AVERAGE Error includes scale factor of 1.3.

49.8 ± 4.2 ± 3.4	205	¹ AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega K^+K^- 2(\pi^+\pi^-)\gamma$
------------------	-----	---------------------	----------	---

31 ± 13	30	VANNUCCI	77 MRK1	e^+e^-
---------	----	----------	---------	----------

¹Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

NODE=M070R17
NODE=M070R17

NODE=M070R17;LINKAGE=EE

$\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.12 ± 0.09 OUR AVERAGE

2.36 ± 0.02 ± 0.21	59k	ABLIKIM	06K BES2	$J/\psi \rightarrow \rho\pi^-\bar{n}$
--------------------	-----	---------	----------	---------------------------------------

2.47 ± 0.02 ± 0.24	55k	ABLIKIM	06K BES2	$J/\psi \rightarrow \bar{\rho}\pi^+n$
--------------------	-----	---------	----------	---------------------------------------

OCCUR=2

2.02 ± 0.07 ± 0.16	1288	EATON	84 MRK2	$e^+e^- \rightarrow \rho\pi^-$
--------------------	------	-------	---------	--------------------------------

OCCUR=2

1.93 ± 0.07 ± 0.16	1191	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\rho}\pi^+$
--------------------	------	-------	---------	--------------------------------------

1.7 ± 0.7	32	BESCH	81 BONA	$e^+e^- \rightarrow \rho\pi^-$
-----------	----	-------	---------	--------------------------------

1.6 ± 1.2	5	BESCH	81 BONA	$e^+e^- \rightarrow \bar{\rho}\pi^+$
-----------	---	-------	---------	--------------------------------------

OCCUR=2

2.16 ± 0.29	194	PERUZZI	78 MRK1	$e^+e^- \rightarrow \rho\pi^-$
-------------	-----	---------	---------	--------------------------------

2.04 ± 0.27	204	PERUZZI	78 MRK1	$e^+e^- \rightarrow \bar{\rho}\pi^+$
-------------	-----	---------	---------	--------------------------------------

OCCUR=2

$\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.86 ± 0.11 OUR AVERAGE Error includes scale factor of 1.2.

0.90 ± 0.03 ± 0.18	961 ± 35	ABLIKIM	12P BES2	$J/\psi \rightarrow \Xi^-\bar{\Xi}^+$
--------------------	----------	---------	----------	---------------------------------------

0.70 ± 0.06 ± 0.12	132 ± 11	HENRARD	87 DM2	$e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$
--------------------	----------	---------	--------	---------------------------------------

1.14 ± 0.08 ± 0.20	194	EATON	84 MRK2	$e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$
--------------------	-----	-------	---------	---------------------------------------

1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$
-----------	----	---------	---------	---------------------------------------

NODE=M070R62
NODE=M070R62

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.61 ± 0.15 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.

1.93 ± 0.21 ± 0.05		¹ AUBERT	07BD BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
--------------------	--	---------------------	-----------	--

2.03 ± 0.03 ± 0.15	8887	ABLIKIM	06 BES2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$
--------------------	------	---------	---------	---

1.9 $^{+0.5}_{-0.4}$ ± 0.1	46	² WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
----------------------------	----	-----------------	---------	---

1.08 ± 0.06 ± 0.24	631	BAI	98G BES	e^+e^-
--------------------	-----	-----	---------	----------

1.38 ± 0.05 ± 0.20	1847	PALLIN	87 DM2	e^+e^-
--------------------	------	--------	--------	----------

1.58 ± 0.08 ± 0.19	365	EATON	84 MRK2	e^+e^-
--------------------	-----	-------	---------	----------

2.6 ± 1.6	5	BESCH	81 BONA	e^+e^-
-----------	---	-------	---------	----------

1.1 ± 0.2	196	PERUZZI	78 MRK1	e^+e^-
-----------	-----	---------	---------	----------


NODE=M070R60
NODE=M070R60

¹AUBERT 07BD reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (10.7 \pm 0.9 \pm 0.7) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R60;LINKAGE=AU

²WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00 $^{+0.34}_{-0.29}$ ± 0.34) × 10⁻⁶ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.$

NODE=M070R60;LINKAGE=WU

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma(p\bar{p})$ $\Gamma_{128}/\Gamma_{108}$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.90^{+0.15}_{-0.14} \pm 0.10$	¹ WU	06	BELL $B^+ \rightarrow p\bar{p}K^+, \Lambda\bar{\Lambda}K^+$

NODE=M070R79
NODE=M070R79

¹ Not independent of other $J/\psi \rightarrow \Lambda\bar{\Lambda}, p\bar{p}$ branching ratios reported by WU 06.

NODE=M070R79;LINKAGE=WU

 $\Gamma(\Lambda\bar{\Sigma}^- \pi^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.2.
0.770 ± 0.051 ± 0.083	335	¹ ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^+ \pi^-$
0.747 ± 0.056 ± 0.076	254	¹ ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
0.90 ± 0.06 ± 0.16	225 ± 15	HENRARD	87 DM2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^+ \pi^-$
1.11 ± 0.06 ± 0.20	342 ± 18	HENRARD	87 DM2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$
1.53 ± 0.17 ± 0.38	135	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^+ \pi^-$
1.38 ± 0.21 ± 0.35	118	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^- \pi^+$

NODE=M070R71
NODE=M070R71

OCCUR=2

OCCUR=2

OCCUR=2

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

NODE=M070R71;LINKAGE=AB

 $\Gamma(pK^-\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.89 ± 0.07 ± 0.14	307	EATON	84	MRK2 e^+e^-

NODE=M070R72
NODE=M070R72

 $\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.76 ± 0.09 OUR AVERAGE				
0.74 ± 0.09 ± 0.02	156 ± 15	¹ AUBERT	07AK	BABR 10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$
1.4 $^{+0.5}_{-0.4} \pm 0.2$	11.0 $^{+4.3}_{-3.5}$	² HUANG	03	BELL $B^+ \rightarrow 2(K^+K^-)K^+$
0.7 ± 0.3		VANNUCCI	77	MRK1 e^+e^-

NODE=M070R19
NODE=M070R19

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.72 ± 0.17 ± 0.02	38	³ AUBERT	05D	BABR 10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$
--------------------	----	---------------------	-----	--

¹ AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow 2(K^+K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (4.11 \pm 0.39 \pm 0.30) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R19;LINKAGE=BE

² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

NODE=M070R19;LINKAGE=CC

³ Superseded by AUBERT 07AK. AUBERT 05D reports $[\Gamma(J/\psi(1S) \rightarrow 2(K^+K^-))/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] = (4.0 \pm 0.7 \pm 0.6) \times 10^{-3}$ keV which we divide by our best value $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.55 \pm 0.14 \pm 0.02$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R19;LINKAGE=AU

$\Gamma(pK-\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.06±0.05	90	EATON	84	MRK2 e^+e^-

NODE=M070R73
 NODE=M070R73

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.86±0.21 OUR AVERAGE		[(2.70 ± 0.17) × 10 ⁻⁴ OUR 2015 AVERAGE]		

NODE=M070R13
 NODE=M070R13

2.86±0.09±0.19 1k ¹METREVELI 12 $\psi(2S) \rightarrow \pi^+\pi^-K^+K^-$

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.22±0.20±0.12	462	^{2,3} LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
3.50±0.20±0.12	462	^{3,4} LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
2.39±0.24±0.22	107	⁵ BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ±0.9	6	⁵ BRANDELIK	79C	DASP e^+e^-

OCCUR=2

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(J/\psi \rightarrow e^+e^-) = (5.55 \pm 0.14)$ keV.

⁴ $\sin\phi < 0$.

⁵ Interference with non-resonant K^+K^- production not taken into account.

NODE=M070R13;LINKAGE=ME
 NODE=M070R13;LINKAGE=A
 NODE=M070R13;LINKAGE=B
 NODE=M070R13;LINKAGE=C
 NODE=M070R13;LINKAGE=BA

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{134}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1 ±0.4 OUR AVERAGE		Error includes scale factor of 3.2.		

NODE=M070R75
 NODE=M070R75

2.62±0.15±0.14 0.3k ¹METREVELI 12 $\psi(2S) \rightarrow \pi^+\pi^-K_S^0 K_L^0$

1.82±0.04±0.13 2.1k ²BAI 04A BES2 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.18±0.12±0.18		JOUSSET	90	DM2 $J/\psi \rightarrow$ hadrons
1.01±0.16±0.09	74	BALTRUSAIT..85D	MRK3	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$.

NODE=M070R75;LINKAGE=ME
 NODE=M070R;LINKAGE=HZ

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{135}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30±0.13±0.99	2.4k	ABLIKIM	12P	BES2 J/ψ

NODE=M070S78
 NODE=M070S78

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{136}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2±1.7 OUR AVERAGE				

NODE=M070R07
 NODE=M070R07

15.7±0.80±1.54 454 ¹ABLIKIM 13F BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

26.2±6.0 ±4.4 44 ²ABLIKIM 07H BES2 $e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^-\rho) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^-\rho) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

NODE=M070R07;LINKAGE=AL
 NODE=M070R07;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{137}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

NODE=M070S11
 NODE=M070S11

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4 90 ²ABLIKIM 07H BES2 $e^+e^- \rightarrow \psi(2S)$

23 ±7 ±8 11 BAI 98G BES e^+e^-

22 ±5 ±5 19 HENRARD 87 DM2 e^+e^-

¹ Using $B(\Lambda \rightarrow \pi^-\rho) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^-\rho) = 63.9\%$.

NODE=M070S11;LINKAGE=AL
 NODE=M070S11;LINKAGE=AB

 $\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{138}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46±0.20±1.07	1058	¹ ABLIKIM	08C	BES2 $e^+e^- \rightarrow J/\psi$

NODE=M070S56
 NODE=M070S56

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

NODE=M070S56;LINKAGE=AB

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{139}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE				
1.47±0.13±0.13	140	¹ METREVELI 12		$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	e^+e^-
1.0 ±0.5	5	BRANDELIK 78B	DASP	e^+e^-
1.6 ±1.6	1	VANNUCCI 77	MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070R6
NODE=M070R6

NODE=M070R6;LINKAGE=ME

 $\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83±0.23 OUR AVERAGE					
2.74±0.24±0.22		234 ± 21	¹ ABLIKIM 12B	BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92±0.22±0.24		308 ± 24	² ABLIKIM 12B	BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<15 90 PERUZZI 78 MRK1 $e^+e^- \rightarrow \Lambda X$

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B quotes $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

NODE=M070R61
NODE=M070R61

OCCUR=2

NODE=M070R61;LINKAGE=AB
NODE=M070R61;LINKAGE=AC

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{141}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	¹ BAI 04D	BES	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.052	90	¹ BALTRUSAIT..85C	MRK3	e^+e^-

¹ Forbidden by CP.

NODE=M070R14
NODE=M070R14

NODE=M070R14;LINKAGE=C

————— RADIATIVE DECAYS —————

 $\Gamma(3\gamma)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.6±2.2 OUR AVERAGE					
11.3±1.8±2.0		113 ± 18	ABLIKIM 13I	BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ±3 ±2		24.2 ^{+7.2} _{-6.0}	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<55	90	PARTRIDGE 80	CBAL	e^+e^-	

NODE=M070R81
NODE=M070R81

NODE=M070310

 $\Gamma(4\gamma)/\Gamma_{\text{total}}$ Γ_{143}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

NODE=M070S06
NODE=M070S06

 $\Gamma(5\gamma)/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<15	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

NODE=M070S07
NODE=M070S07

 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{145}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.15±0.05	¹ ABLIKIM 15AE	BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ The uncertainty is systematic as statistical is negligible.

NODE=M070B00
NODE=M070B00

NODE=M070B00;LINKAGE=A

 $\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7 ±0.4 OUR AVERAGE				Error includes scale factor of 1.5.
2.01±0.32±0.02		¹ MITCHELL 09	CLEO	$e^+e^- \rightarrow \gamma X$
1.27±0.36		GAISER 86	CBAL	$J/\psi \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen		ANASHIN 14	KEDR	$J/\psi \rightarrow \gamma\eta_c$
0.79±0.20	273 ± 43	² AUBERT 06E	BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
seen	16	BALTRUSAIT..84	MRK3	$J/\psi \rightarrow 2\phi\gamma$

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by the authors using an average of $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$ from BALTRUSAITIS 86, BISELLO 91, BAI 04 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

NODE=M070R85
NODE=M070R85

NODE=M070R85;LINKAGE=MI

NODE=M070R85;LINKAGE=AU

$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070S08
NODE=M070S08

$3.8^{+1.3}_{-1.0}$ OUR AVERAGE Error includes scale factor of 1.1.

$4.5 \pm 1.2 \pm 0.6$	33 ± 9	ABLIKIM	13I	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$1.2^{+2.7}_{-1.1} \pm 0.3$	$1.2^{+2.8}_{-1.1}$	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $\Gamma(\gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070R99
NODE=M070R99

$8.3 \pm 0.2 \pm 3.1$

¹ BALTRUSAIT..86B MRK3 $J/\psi \rightarrow 4\pi\gamma$

¹ 4π mass less than 2.0 GeV.

NODE=M070R99;LINKAGE=M

 $\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070R96
NODE=M070R96

6.1 ± 1.0 OUR AVERAGE

$5.85 \pm 0.3 \pm 1.05$	¹ EDWARDS	83B	CBAL	$J/\psi \rightarrow \eta\pi^+\pi^-$
$7.8 \pm 1.2 \pm 2.4$	¹ EDWARDS	83B	CBAL	$J/\psi \rightarrow \eta 2\pi^0$

OCCUR=2

¹ Broad enhancement at 1700 MeV.

NODE=M070R96;LINKAGE=M

 $\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070S37
NODE=M070S37

$6.2 \pm 2.2 \pm 0.9$

BAI 99 BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070R89
NODE=M070R89

2.8 ± 0.6 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

$1.66 \pm 0.1 \pm 0.58$	^{1,2} BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$3.8 \pm 0.3 \pm 0.6$	³ AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$4.0 \pm 0.7 \pm 1.0$	³ EDWARDS	82E	CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	^{3,4} SCHARRE	80	MRK2	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.78 \pm 0.21 \pm 0.33$	^{3,5,6} AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.83 \pm 0.13 \pm 0.18$	^{3,7,8} AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.66^{+0.17+0.24}_{-0.16-0.15}$	^{3,6,9} BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
$1.03^{+0.21+0.26}_{-0.18-0.19}$	^{3,8,10} BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

OCCUR=2

OCCUR=2

¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K \bar{K} \pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

NODE=M070R89;LINKAGE=BD

² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

NODE=M070R89;LINKAGE=BE

³ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

NODE=M070R89;LINKAGE=BE

⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.

NODE=M070R89;LINKAGE=C

⁵ From fit to the $a_0(980)\pi 0^-+$ partial wave.

NODE=M070R89;LINKAGE=H

⁶ $a_0(980)\pi$ mode.

NODE=M070R89;LINKAGE=K9

⁷ From fit to the $K^*(892)K 0^-+$ partial wave.

NODE=M070R89;LINKAGE=J

⁸ $K^* K$ mode.


NODE=M070R89;LINKAGE=K8

⁹ From $a_0(980)\pi$ final state.

NODE=M070R89;LINKAGE=D

¹⁰ From $K^*(890)K$ final state.

NODE=M070R89;LINKAGE=E



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0) / \Gamma_{\text{total}}$

Γ_{152} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
1.07 ± 0.17 ± 0.11	¹ BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
0.64 ± 0.12 ± 0.07	¹ COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

NODE=M070S30
NODE=M070S30

¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

NODE=M070S30;LINKAGE=C

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-) / \Gamma_{\text{total}}$

Γ_{153} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ± 0.5 OUR AVERAGE				
2.6 ± 0.7 ± 0.4		BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
3.38 ± 0.33 ± 0.64		¹ BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M070S29
NODE=M070S29

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0 ± 0.6 ± 1.1	261	² AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-----------------	-----	-----------------------	--------	---

¹ Via $a_0(980)\pi$.

² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

NODE=M070S29;LINKAGE=RR
NODE=M070S29;LINKAGE=R

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi) / \Gamma_{\text{total}}$

Γ_{154} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.82	95	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma K^+ K^-$

NODE=M070R77
NODE=M070R77

$\Gamma(\gamma\rho\rho) / \Gamma_{\text{total}}$

Γ_{155} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
4.5 ± 0.8 OUR AVERAGE				
4.7 ± 0.3 ± 0.9		¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
3.75 ± 1.05 ± 1.20		² BURKE	82 MRK2	$J/\psi \rightarrow 4\pi\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.09	90	³ BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
-------	----	----------------------	-----	---------------------------------

¹ 4π mass less than 2.0 GeV.

² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .

³ 4π mass in the range 2.0–25 GeV.

NODE=M070R94;LINKAGE=N
NODE=M070R94;LINKAGE=M
NODE=M070R94;LINKAGE=A

$\Gamma(\gamma\rho\omega) / \Gamma_{\text{total}}$

Γ_{156} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070R05
NODE=M070R05

$\Gamma(\gamma\rho\phi) / \Gamma_{\text{total}}$

Γ_{157} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<8.8	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070R06
NODE=M070R06

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.15±0.16 OUR AVERAGE		Error includes scale factor of 1.2.		
4.82±0.23±0.08		¹ ABLIKIM	11 BES3	$J/\psi \rightarrow \eta'\gamma$
5.24±0.12±0.11		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta'\gamma$
5.55±0.44	35k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta'\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.50±0.14±0.53		BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$
4.30±0.31±0.71		BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$
4.04±0.16±0.85	622	AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
4.39±0.09±0.66	2420	AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
4.1 ±0.3 ±0.6		BLOOM	83 CBAL	$e^+e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 ±1.1	6	BRANDELIK	79C DASP	$e^+e^- \rightarrow 3\gamma$
2.4 ±0.7	57	BARTEL	76 CNTR	$e^+e^- \rightarrow 2\gamma\rho$

NODE=M070R84
 NODE=M070R84

OCCUR=2

OCCUR=2

NODE=M070R84;LINKAGE=AB

¹ ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.9 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ±0.5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
4.32±0.14±0.73	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
2.08±0.13±0.35	² BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
3.05±0.08±0.45	² BALTRUSAIT..	86B MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85±0.45±1.20	³ BURKE	82 MRK2	e^+e^-


NODE=M070R95
 NODE=M070R95

OCCUR=2

NODE=M070R95;LINKAGE=A
 NODE=M070R95;LINKAGE=B
 NODE=M070R95;LINKAGE=M

- ¹ 4π mass less than 3.0 GeV.
² 4π mass less than 2.0 GeV.
³ 4π mass less than 2.5 GeV.

WEIGHTED AVERAGE
 2.8±0.5 (Error scaled by 1.9)

 $\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$ Γ_{160}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.5±0.7±1.6	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M070S45
 NODE=M070S45

 $\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant}))/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.2±0.8±1.7	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M070S46
 NODE=M070S46

¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

NODE=M070S46;LINKAGE=AB

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.1±0.6	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

NODE=M070B05
 NODE=M070B05

 $\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.7±0.5±0.5	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

NODE=M070S7
 NODE=M070S7

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi \pi / \text{total} = 0.167$.

NODE=M070S7;LINKAGE=V

 $\Gamma(\gamma \omega \omega)/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61±0.33 OUR AVERAGE				
6.0 ±4.8 ±1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma \omega \pi^+ \pi^-$
1.41±0.2 ±0.42	120 ± 17	BISELLO	87 SPEC	$e^+ e^-$, hadrons γ
1.76±0.09±0.45		BALTRUSAIT..85C	MRK3	$e^+ e^- \rightarrow \text{hadrons } \gamma$

NODE=M070R97
 NODE=M070R97

 $\Gamma(\gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.7 ±0.4 OUR AVERAGE	Error includes scale factor of 1.3.		
2.1 ±0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1.36±0.38	^{1,2} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi \gamma$

NODE=M070S19
 NODE=M070S19

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0 \rho^0$.

NODE=M070S19;LINKAGE=A
 NODE=M070S19;LINKAGE=B

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.64±0.12 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
[(1.43 ± 0.11) × 10 ⁻³ OUR 2015 AVERAGE]				
2.07±0.16 ^{+0.02} _{-0.07}	2.4k	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
1.63±0.26 ^{+0.02} _{-0.06}		³ ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.42±0.21 ^{+0.01} _{-0.05}		⁴ ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
1.33±0.05±0.20		⁵ AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.36±0.09±0.23		⁵ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.48±0.25±0.30	178	EDWARDS	82B CBAL	$e^+ e^- \rightarrow 2\pi^0 \gamma$
2.0 ±0.7	35	ALEXANDER	78 PLUT	$e^+ e^-$
1.2 ±0.6	30	⁶ BRANDELIK	78B DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

NODE=M070R86
 NODE=M070R86

NEW

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(15) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 06V reports $[\Gamma(J/\psi(15) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(15) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi \pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Estimated using $B(f_2(1270) \rightarrow \pi \pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁶ Restated by us to take account of spread of E1, M2, E3 transitions.

NODE=M070R86;LINKAGE=A
 NODE=M070R86;LINKAGE=DO


NODE=M070R86;LINKAGE=A1

NODE=M070R86;LINKAGE=AL

NODE=M070R86;LINKAGE=X

NODE=M070R86;LINKAGE=T

OCCUR=2



$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{total}$ **Γ_{167}/Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	COMMENT
4.19±0.73±1.34	478	¹ DOBBS 15	$J/\psi \rightarrow \gamma K \bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070R00
NODE=M070R00

NODE=M070R00;LINKAGE=A

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{total}$ **Γ_{168}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
10.0 \pm 1.1 $\overline{-0.9}$					OUR AVERAGE Error includes scale factor of 1.5. See the ideogram

below. $[(8.5^{+1.2}_{-0.9}) \times 10^{-4}]$ OUR 2015 AVERAGE Scale factor = 1.2]

11.76±	0.54±0.94	1.2k	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
9.62±029	$\begin{matrix} +3.51 \\ -1.86 \end{matrix}$		² BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
5.0 ±	$\begin{matrix} 0.8 \\ +1.8 \\ -0.4 \end{matrix}$		^{3,4} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
9.2 ±	1.4 ±1.4		⁴ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
10.4 ±	1.2 ±1.6		⁴ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
9.6 ±	1.2 ±1.8		⁴ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6 ±	0.2 $\begin{matrix} +0.6 \\ -0.2 \end{matrix}$		^{4,5} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	90		⁶ BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
1.6 ±	0.4 ±0.3		⁷ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.8 ±	1.6		⁸ EDWARDS	82D CBAL	$e^+ e^- \rightarrow \eta \eta \gamma$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.

³ Assuming $J^P = 2^+$ for $f_0(1710)$.

⁴ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K \bar{K}$ result.

⁵ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁶ Includes unknown branching fraction to $\rho^0 \rho^0$.

⁷ Includes unknown branching fraction to $\pi^+ \pi^-$.

⁸ Includes unknown branching fraction to $\eta \eta$.

NODE=M070R91
NODE=M070R91

NEW


OCCUR=2

OCCUR=2

OCCUR=2

NODE=M070R91;LINKAGE=D
NODE=M070R91;LINKAGE=K9
NODE=M070R91;LINKAGE=A1
NODE=M070R91;LINKAGE=B

NODE=M070R91;LINKAGE=A2
NODE=M070R91;LINKAGE=C
NODE=M070R91;LINKAGE=Z
NODE=M070R91;LINKAGE=A



$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{169} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5 OUR AVERAGE				
[(4.0 ± 1.0) × 10 ⁻⁴ OUR 2015 AVERAGE]				
3.72 ± 0.30 ± 0.43	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
3.96 ± 0.06 ± 1.12		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.99 ± 0.15 ± 2.64		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.5 ± 1.6 ± 0.8		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$

NODE=M070B01
NODE=M070B01
NEW

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi \pi$.

NODE=M070B01;LINKAGE=A
NODE=M070B01;LINKAGE=AB

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{170} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.06 ± 0.08	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$

NODE=M070R01
NODE=M070R01

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{171} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35^{+0.13+1.24}_{-0.11-0.74}	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

NODE=M070S84
NODE=M070S84

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S84;LINKAGE=A

$\Gamma(\gamma \eta) / \Gamma_{\text{total}}$ Γ_{172} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.104 ± 0.034 OUR AVERAGE				
1.101 ± 0.029 ± 0.022		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta \gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.88 ± 0.08 ± 0.11		BLOOM	83 CBAL	$e^+ e^-$
0.82 ± 0.10		BRANDELIK	79C DASP	$e^+ e^-$
1.3 ± 0.4	21	BARTEL	77 CNTR	$e^+ e^-$

NODE=M070R83
NODE=M070R83

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{173} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.79 ± 0.13 OUR AVERAGE			
0.68 ± 0.04 ± 0.24	BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.76 ± 0.15 ± 0.21	^{1,2} AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
0.87 ± 0.14 ^{+0.14} _{-0.11}	¹ BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

NODE=M070S31
NODE=M070S31

OCCUR=2

¹ Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.

² From fit to the $K^*(892)K 1^{++}$ partial wave.

NODE=M070S31;LINKAGE=A
NODE=M070S31;LINKAGE=D

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$ Γ_{174}/Γ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.61 ± 0.08 OUR AVERAGE

0.69 ± 0.16 ± 0.20

1 BAI 04J BES2 $J/\psi \rightarrow \gamma \gamma \rho^0$

0.61 ± 0.04 ± 0.21

2 BAI 00D BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

0.45 ± 0.09 ± 0.17

3 BAI 99 BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

0.625 ± 0.063 ± 0.103

4 BOLTON 92 MRK3 $J/\psi \rightarrow \gamma f_1(1285)$

0.70 ± 0.08 ± 0.16

5 BOLTON 92B MRK3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$ ¹ Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.² Assuming $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$.³ Assuming $\Gamma(f_1(1285) \rightarrow \eta \pi \pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$.⁴ Obtained summing the sequential decay channels $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$; $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$; $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$; $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.⁵ Using $B(f_1(1285) \rightarrow a_0(980) \pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta \pi$.NODE=M070R88
NODE=M070R88NODE=M070R88;LINKAGE=BI
NODE=M070R88;LINKAGE=BD
NODE=M070R88;LINKAGE=BA
NODE=M070R88;LINKAGE=B

NODE=M070R88;LINKAGE=A

 $\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{175}/Γ VALUE (units 10^{-4})

DOCUMENT ID TECN COMMENT

4.5 ± 1.0 ± 0.7BAI 99 BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$ NODE=M070S36
NODE=M070S36 $\Gamma(\gamma f_2'(1525))/\Gamma_{\text{total}}$ Γ_{176}/Γ VALUE (units 10^{-4})

CL% EVTS


DOCUMENT ID TECN COMMENT

5.7 $\pm_{-0.5}^{+0.8}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogrambelow. $[(4.5 \pm_{-0.4}^{+0.7}) \times 10^{-4}]$ OUR 2015 AVERAGE8.0 ± 0.9 ± 0.2 750 ^{1,2} DOBBS 15 $J/\psi \rightarrow \gamma K \bar{K}$ 3.85 ± 0.17 $\pm_{-0.73}^{+1.91}$ ³ BAI 03G BES $J/\psi \rightarrow \gamma K \bar{K}$ 3.6 ± 0.4 $\pm_{-0.4}^{+1.4}$ ³ BAI 96C BES $J/\psi \rightarrow \gamma K^+ K^-$ 5.6 ± 1.4 ± 0.9 ³ AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K^+ K^-$ 4.5 ± 0.4 ± 0.9 ³ AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$ 6.8 ± 1.6 ± 1.4 ³ BALTRUSAIT..87 MRK3 $J/\psi \rightarrow \gamma K^+ K^-$ NODE=M070R87
NODE=M070R87

NEW

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<3.4 90 4 ⁴ BRANDELIK 79C DASP $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$ <2.3 90 3 ALEXANDER 78 PLUT $e^+ e^- \rightarrow K^+ K^- \gamma$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2'(1525))/\Gamma_{\text{total}}] \times [B(f_2'(1525) \rightarrow K \bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f_2'(1525) \rightarrow K \bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.³ Using $B(f_2'(1525) \rightarrow K \bar{K}) = 0.888$.⁴ Assuming isotropic production and decay of the $f_2'(1525)$ and isospin.NODE=M070R87;LINKAGE=B
NODE=M070R87;LINKAGE=DONODE=M070R87;LINKAGE=A1
NODE=M070R87;LINKAGE=I



$\Gamma(\gamma f'_2(1525) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{177}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma\eta\eta$

NODE=M070S86
NODE=M070S86

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S86;LINKAGE=A

$\Gamma(\gamma f_2(1640) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{178}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma\omega\omega$

NODE=M070R02
NODE=M070R02

$\Gamma(\gamma f_2(1910) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{179}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma\omega\omega$

NODE=M070R03
NODE=M070R03

$\Gamma(\gamma f_0(1800) \rightarrow \gamma\omega\phi)/\Gamma_{\text{total}}$ Γ_{180}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.6 OUR AVERAGE				
$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J	BES3 $J/\psi \rightarrow \gamma\omega\phi$
$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J	BES2 $J/\psi \rightarrow \gamma\omega\phi$

NODE=M070S79
NODE=M070S79

$\Gamma(\gamma f_2(1810) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{181}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N $J/\psi \rightarrow \gamma\eta\eta$

NODE=M070S87
NODE=M070S87

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S87;LINKAGE=A

$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{182}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.7 \pm 0.1 \pm 0.2$	BAI	00B	BES $J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

NODE=M070B06
NODE=M070B06

$\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{183}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.0 \pm 0.3 \pm 1.3$	320	¹ BAI	00B	BES $J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

NODE=M070B07
NODE=M070B07

¹ Summed over all charges.


NODE=M070R;LINKAGE=B7

$\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$ Γ_{184}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.2 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
$7.5 \pm 0.6 \pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
$3.4 \pm 0.8 \pm 0.6$	33 ± 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.1 \pm 0.7 \pm 0.4$		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ $\phi\phi$ mass less than 2.9 GeV, η_C excluded.NODE=M070R98
NODE=M070R98

NODE=M070R98;LINKAGE=C

 $\Gamma(\gamma\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{185}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.38 \pm 0.07 \pm 0.07$		49	EATON	84	MRK2 $e^+ e^-$
< 0.11	90		PERUZZI	78	MRK1 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M070R90
NODE=M070R90 $\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$ Γ_{186}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.05 OUR AVERAGE				
$0.44 \pm 0.04 \pm 0.08$	196 ± 19	¹ ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$0.33 \pm 0.08 \pm 0.05$		¹ BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$0.27 \pm 0.06 \pm 0.06$		¹ BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$0.24^{+0.15}_{-0.10}$		^{2,3} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Includes unknown branching fraction to $\phi\phi$.² Estimated by us from various fits.³ Includes unknown branching fraction to $\rho^0\rho^0$.NODE=M070S21
NODE=M070S21

OCCUR=2

NODE=M070S21;LINKAGE=U
NODE=M070S21;LINKAGE=A
NODE=M070S21;LINKAGE=B $\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{187}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.09	^{1,2} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.² Includes unknown branching fraction to $\rho^0\rho^0$.NODE=M070S20
NODE=M070S20NODE=M070S20;LINKAGE=A
NODE=M070S20;LINKAGE=B $\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{188}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

NODE=M070R04
NODE=M070R04

$\Gamma(\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{189}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 \pm 0.4 OUR AVERAGE				
$2.87 \pm 0.09^{+0.49}_{-0.52}$	4265	¹ ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
$2.2 \pm 0.4 \pm 0.4$	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M070R78
 NODE=M070R78

¹ From a fit of the $\pi^+ \pi^- \eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0 \pi^+ \pi^- \eta'$.

NODE=M070R78;LINKAGE=AI

 $\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{190}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.77$^{+0.15}_{-0.09}$ OUR AVERAGE				
$0.90^{+0.04+0.27}_{-0.11-0.55}$		¹ ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
$1.14^{+0.43+0.42}_{-0.30-0.26}$	231	² ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
$0.70 \pm 0.04^{+0.19}_{-0.08}$		BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

NODE=M070S71
 NODE=M070S71

¹ From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A.

NODE=M070S71;LINKAGE=AK

² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

NODE=M070S71;LINKAGE=AL

 $\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$ Γ_{191}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
3.31$^{+0.33+1.96}_{-0.30-1.29}$ OUR AVERAGE			
	ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

NODE=M070S96
 NODE=M070S96

 $\Gamma(\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.44\pm 0.36$^{+0.60}_{-0.74}$				
	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$

NODE=M070S82
 NODE=M070S82

 $\Gamma(\gamma(K\bar{K}\pi)[J^{PC}=0^{-+}])/ \Gamma_{\text{total}}$ Γ_{193}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 \pm 0.4 OUR AVERAGE Error includes scale factor of 2.1.			
$0.58 \pm 0.03 \pm 0.20$	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$2.1 \pm 0.1 \pm 0.7$	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

NODE=M070S38
 NODE=M070S38

¹ For a broad structure around 1800 MeV.

OCCUR=2

² For a broad structure around 2040 MeV.

NODE=M070S38;LINKAGE=BD
 NODE=M070S38;LINKAGE=BE

 $\Gamma(\gamma \pi^0)/\Gamma_{\text{total}}$ Γ_{194}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.49$^{+0.33}_{-0.30}$ OUR AVERAGE				
$3.63 \pm 0.36 \pm 0.13$		PEDLAR	09 CLE3	$J/\psi \rightarrow \pi^0 \gamma$
$3.13^{+0.65}_{-0.47}$	586	ABLIKIM	06E BES2	$J/\psi \rightarrow \pi^0 \gamma$

NODE=M070R82
 NODE=M070R82

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.6 \pm 1.1 \pm 0.7$	BLOOM	83 CBAL	$e^+ e^-$
7.3 ± 4.7	10	BRANDELIK	79C DASP $e^+ e^-$

 $\Gamma(\gamma p\bar{p}\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{195}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.79				
	90	EATON	84 MRK2	$e^+ e^-$

NODE=M070R93
 NODE=M070R93

 $\Gamma(\gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{196}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.13				
	90	HENRARD	87 DM2	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.16	90	BAI	98G BES	$e^+ e^-$

NODE=M070S8
 NODE=M070S8

$\Gamma(\gamma f_0(2100) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{197} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

NODE=M070S85
NODE=M070S85¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S85;LINKAGE=A

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{198} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$

NODE=M070B08
NODE=M070B08¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070B08;LINKAGE=A

 $\Gamma(\gamma f_0(2200)) / \Gamma_{\text{total}}$ Γ_{199} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070S18
NODE=M070S18

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ¹ AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$ ¹ Includes unknown branching fraction to $K_S^0 K_S^0$.

NODE=M070S18;LINKAGE=A

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{200} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.86 \pm 0.49 \pm 1.20$	490	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$

NODE=M070B09
NODE=M070B09¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070B09;LINKAGE=A

 $\Gamma(\gamma f_J(2220)) / \Gamma_{\text{total}}$ Γ_{201} / Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M070R92
NODE=M070R92

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300 ¹ BAI 96B BES $e^+ e^- \rightarrow \gamma \bar{p} p, K \bar{K}$ >250 99.9 ² HASAN 96 SPEC $\bar{p} p \rightarrow \pi^+ \pi^-$ < 2.3 95 ³ AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K^+ K^-$ < 1.6 95 ³ AUGUSTIN 88 DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

OCCUR=2

 $12.4^{+6.4}_{-5.2} \pm 2.8$ 23 ³ BALTRUSAIT..86D MRK3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$ $8.4^{+3.4}_{-2.8} \pm 1.6$ 93 ³ BALTRUSAIT..86D MRK3 $J/\psi \rightarrow \gamma K^+ K^-$

OCCUR=2

¹ Using BARNES 93.² Using BAI 96B.³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.NODE=M070R92;LINKAGE=A
NODE=M070R92;LINKAGE=M
NODE=M070R92;LINKAGE=W $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{202} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M070B02
NODE=M070B02< 3.9 90 ^{1,2} DOBBS 15 $J/\psi \rightarrow \gamma \pi \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $14 \pm 8 \pm 4$ BAI 98H BES $J/\psi \rightarrow \gamma \pi^0 \pi^0$ $8.4 \pm 2.6 \pm 3.0$ BAI 96B BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.NODE=M070B02;LINKAGE=A
NODE=M070B02;LINKAGE=D0 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{203} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M070B03
NODE=M070B03< 4.1 (CL = 90%) [$< 3.6 \times 10^{-5}$ OUR 2015 BEST LIMIT]< 4.1 90 ^{1,2} DOBBS 15 $J/\psi \rightarrow \gamma K \bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6 ³ DEL-AMO-SA..100 BABR $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$ < 2.9 ³ DEL-AMO-SA..100 BABR $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

OCCUR=2

 $6.6 \pm 2.9 \pm 2.4$ BAI 96B BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$ $10.8 \pm 4.0 \pm 3.2$ BAI 96B BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.NODE=M070B03;LINKAGE=A
NODE=M070B03;LINKAGE=D0³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

NODE=M070B03;LINKAGE=D0

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{204}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.6 \pm 0.5$	BAI	96B	BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma p \bar{p}$

NODE=M070B04
 NODE=M070B04

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{205}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.60^{+0.62+2.37}_{-0.65-2.07}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

NODE=M070S88
 NODE=M070S88

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S88;LINKAGE=A

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{206}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.24 OUR AVERAGE [(1.01 ± 0.32) × 10 ⁻⁴ OUR 2015 AVERAGE]				

NODE=M070S32
 NODE=M070S32

NEW

1.21 ± 0.29 ± 0.24	174	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
1.00 ± 0.03 ± 0.45		² ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.02 ± 0.09 ± 0.45		² ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.7 ± 0.8		^{3,4} BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
-----------	--	---------------------	----	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi \pi$.

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.

⁴ Assuming that $f_0(1500)$ decays only to two S-wave dipions.

NODE=M070S32;LINKAGE=C
 NODE=M070S32;LINKAGE=AB
 NODE=M070S32;LINKAGE=A
 NODE=M070S32;LINKAGE=B

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{207}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.65^{+0.26+0.51}_{-0.31-1.40}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

NODE=M070S83
 NODE=M070S83

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S83;LINKAGE=A

 $\Gamma(\gamma A \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$
(narrow state A with $m_A < 960$ MeV) Γ_{208}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.3	90	¹ INSLER	10	CLEO $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

NODE=M070S68
 NODE=M070S68

¹ The limit varies with mass m_A of a narrow state A and is 4.3×10^{-6} for $m_A = 0$ MeV, reaches its largest value of 6.3×10^{-6} at $m_A = 500$ MeV, and is 3.6×10^{-6} at $m_A = 960$ MeV.

NODE=M070S68;LINKAGE=IN

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{209}/Γ (narrow state A^0 with $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$)

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	¹ ABLIKIM	12	BES3 $J/\psi \rightarrow \gamma \mu^+ \mu^-$

NODE=M070S76
 NODE=M070S76

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

NODE=M070S76;LINKAGE=AB

———— DALITZ DECAYS ————

NODE=M070330

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{210}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM	14I	BES3 $J/\psi \rightarrow \pi^0 e^+ e^-$

NODE=M070S89
 NODE=M070S89

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{211}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM	14I	BES3 $J/\psi \rightarrow \eta e^+ e^-$

NODE=M070S90
 NODE=M070S90

¹ Using both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

NODE=M070S90;LINKAGE=A

 $\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$ Γ_{212}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.81 \pm 0.16 \pm 0.31$	1.4k	¹ ABLIKIM	14I	BES3 $J/\psi \rightarrow \eta' e^+ e^-$

NODE=M070S91
 NODE=M070S91

¹ Using both $\eta' \rightarrow \gamma \pi^+ \pi^-$ and $\eta' \rightarrow \pi^+ \pi^- \eta$ decays.

NODE=M070S91;LINKAGE=A

WEAK DECAYS

NODE=M070320

 $\Gamma(D^- e^+ \nu_e + c.c.)/\Gamma_{\text{total}}$ Γ_{213}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S53
NODE=M070S53 $\Gamma(\bar{D}^0 e^+ e^- + c.c.)/\Gamma_{\text{total}}$ Γ_{214}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S54
NODE=M070S54 $\Gamma(D_s^- e^+ \nu_e + c.c.)/\Gamma_{\text{total}}$ Γ_{215}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.3 (CL = 90%) [$<3.6 \times 10^{-5}$ (CL = 90%) OUR 2015 BEST LIMIT]				

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.3	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$

••• We do not use the following data for averages, fits, limits, etc. •••

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<36	90	¹ ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$

¹ Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5\%$.NODE=M070S55
NODE=M070S55

NODE=M070S55;LINKAGE=AB

 $\Gamma(D_s^{*-} e^+ \nu_e + c.c.)/\Gamma_{\text{total}}$ Γ_{216}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.8 $\times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070B13
NODE=M070B13 $\Gamma(D^- \pi^+ + c.c.)/\Gamma_{\text{total}}$ Γ_{217}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.5 $\times 10^{-5}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S61
NODE=M070S61 $\Gamma(\bar{D}^0 \bar{K}^0 + c.c.)/\Gamma_{\text{total}}$ Γ_{218}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 $\times 10^{-4}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S62
NODE=M070S62 $\Gamma(\bar{D}^0 \bar{K}^{*0} + c.c.)/\Gamma_{\text{total}}$ Γ_{219}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.5 $\times 10^{-6}$	90	ABLIKIM	14K BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S93
NODE=M070S93 $\Gamma(D_s^- \pi^+ + c.c.)/\Gamma_{\text{total}}$ Γ_{220}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 $\times 10^{-4}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S63
NODE=M070S63 $\Gamma(D_s^- \rho^+ + c.c.)/\Gamma_{\text{total}}$ Γ_{221}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 $\times 10^{-5}$	90	ABLIKIM	14K BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S92
NODE=M070S92

CHARGE CONJUGATION (C), PARITY (P),

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

NODE=M070315

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{222}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.7	90	ABLIKIM	14Q BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

NODE=M070R80
NODE=M070R80

••• We do not use the following data for averages, fits, limits, etc. •••

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 50	90	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1600	90	¹ WICHT	08 BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 220	90	ABLIKIM	07J BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5000	90	BARTEL	77 CNTR	$e^+ e^-$

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.026 \times 10^{-3}$.

NODE=M070R80;LINKAGE=WI

 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{223}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 $\times 10^{-6}$	90	ABLIKIM	14Q BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

NODE=M070S95
NODE=M070S95

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{224}/Γ	
< 1.6	90	ABLIKIM	13L	BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S39 NODE=M070S39
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<11	90	BAI	03D	BES	$e^+ e^- \rightarrow J/\psi$	

 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{225}/Γ	
<8.3	90	ABLIKIM	04	BES	$e^+ e^- \rightarrow J/\psi$	NODE=M070S40 NODE=M070S40

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{226}/Γ	
<2.0	90	ABLIKIM	04	BES	$e^+ e^- \rightarrow J/\psi$	NODE=M070S41 NODE=M070S41

OTHER DECAYS

 $\Gamma(\text{invisible})/\Gamma(e^+ e^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{227}/Γ_5	
<6.6 $\times 10^{-2}$	90	LEES	13I	BABR	$B \rightarrow K^{(*)} J/\psi$	NODE=M070S80 NODE=M070S80

 $\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{227}/Γ_7	
<1.2 $\times 10^{-2}$	90	ABLIKIM	08G	BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	NODE=M070S60 NODE=M070S60

 $J/\psi(1S)$ REFERENCES

AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57147
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56984
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56773
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56776
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56781
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56785
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55900
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55902
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55905
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56238
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56388
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56130
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=55655
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54920
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54954
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54955
ABLIKIM	13L	PR D87 112007	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55300
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55387
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55392
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55402
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55582
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54265
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54267
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54268
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54269
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54273
ABLIKIM	12P	CPC 36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54863
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53646
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53684
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16715
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53220
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53533
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)	REFID=53359
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52718
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53099
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52047
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52128
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52143
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52253
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52255
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52256
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52571


ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=52261
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=52685
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52072
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50986
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51058
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51128
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51130
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51036
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
AUBERT_BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50496
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50507
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50985
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)	REFID=50802
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer		REFID=51038
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49739
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49611
AUBERT_B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49620
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49607
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49751
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49754
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49403
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)	REFID=49473
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47427
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47954
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46606
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46341
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)	REFID=46608
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45146
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44736
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45198
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)	REFID=44739
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)	REFID=45197
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44434
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)	REFID=43314
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)	REFID=43601
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42176
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41866
HSUEH	92	PR D45 R2181	S. Hsueh, S. Palestini	(FNAL, TORI)	REFID=41899
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41668
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41359
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)	REFID=41349
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=40346
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)	REFID=40002
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
HENRRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40261
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)	REFID=40243
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)	REFID=21865
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=22101
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22097
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033

BALTRUSAIT... 84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22006
EATON 84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)	REFID=22092
BLOOM 83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS 83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21318
FRANKLIN 83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
BURKE 82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS 82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22080
EDWARDS 82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
Also	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS 82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
LEMOIGNE 82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
BESCH 81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)	REFID=22077
GIDAL 81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
PARTRIDGE 80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22073
SCHARRE 80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ 80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also	SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
Translated from YAF 34 1471.				
BRANDELIK 79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
ALEXANDER 78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22065
BESCH 78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)	REFID=22066
BRANDELIK 78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22067
PERUZZI 78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)	REFID=22068
BARTEL 77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22058
BURMESTER 77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22060
FELDMAN 77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
VANNUCCI 77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)	REFID=22063
BARTEL 76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
BRAUNSCH... 76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22054
JEAN-MARIE 76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG	REFID=22056
BALDINI-... 75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)	REFID=22026
BOYARSKI 75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC	REFID=22030
DASP 75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22036
ESPOSITO 75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=22038
FORD 75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)	REFID=22039

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=M870

NODE=M056



$$I^G(J^{PC}) = 0^+(0^{++})$$

χc0(1P) MASS

NODE=M056M

NODE=M056M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3414.75 ± 0.31 OUR AVERAGE				
3414.2 ± 0.5 ± 2.3	5.4k	UEHARA 08	BELL	γγ → χc0 → hadrons
3406 ± 7 ± 6	230	¹ ABE 07	BELL	e ⁺ e ⁻ → J/ψ(c \bar{c})
3414.21 ± 0.39 ± 0.27		ABLIKIM 05G	BES2	ψ(2S) → γχc0
3414.7 + 0.7 - 0.6 ± 0.2		² ANDREOTTI 03	E835	p̄p → χc0 → π ⁰ π ⁰
3415.5 ± 0.4 ± 0.4	392	³ BAGNASCO 02	E835	p̄p → χc0 → J/ψγ
3417.4 + 1.8 - 1.9 ± 0.2		² AMBROGIANI 99B	E835	p̄p → e ⁺ e ⁻ γ
3414.1 ± 0.6 ± 0.8		BAI 99B	BES	ψ(2S) → γX
3417.8 ± 0.4 ± 4		² GAISER 86	CBAL	ψ(2S) → γX
3416 ± 3 ± 4		⁴ TANENBAUM 78	MRK1	e ⁺ e ⁻
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3414.6 ± 1.1	266	UEHARA 13	BELL	γγ → K _S ⁰ K _S ⁰
3416.5 ± 3.0		EISENSTEIN 01	CLE2	e ⁺ e ⁻ → e ⁺ e ⁻ χc0
3422 ± 10		⁴ BARTEL 78B	CNTR	e ⁺ e ⁻ → J/ψ2γ
3415 ± 9		⁴ BIDDICK 77	CNTR	e ⁺ e ⁻ → γX

NODE=M056M;LINKAGE=EB
 NODE=M056M;LINKAGE=C
 NODE=M056M;LINKAGE=NW
 NODE=M056M;LINKAGE=D

¹ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.
² Using mass of ψ(2S) = 3686.0 MeV.
³ Recalculated by ANDREOTTI 05A, using the value of ψ(2S) mass from AULCHENKO 03.
⁴ Mass value shifted by us by amount appropriate for ψ(2S) mass = 3686 MeV and J/ψ(1S) mass = 3097 MeV.

χc0(1P) WIDTH

NODE=M056W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±0.6 OUR FIT				
10.5±0.8 OUR AVERAGE Error includes scale factor of 1.1.				
10.6±1.9±2.6	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
12.6 ^{+1.5+0.9} _{-1.6-1.1}		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
8.6 ^{+1.7} _{-1.3} ±0.1		ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
9.7±1.0	392	¹ BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
16.6 ^{+5.2} _{-3.7} ±0.1		AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
14.3±2.0±3.0		BAI	981 BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
13.5±3.3±4.2		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X, \gamma\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
13.2±2.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹Recalculated by ANDREOTTI 05A.

NODE=M056W

NODE=M056W;LINKAGE=AN

NODE=M056215;NODE=M056

$\chi_{c0}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic decays		
Γ_1	$2(\pi^+\pi^-)$	(2.24±0.18) %
Γ_2	$\rho^0\pi^+\pi^-$	(8.7 ±2.8) × 10 ⁻³
Γ_3	$\rho^0\rho^0$	
Γ_4	$f_0(980)f_0(980)$	(6.5 ±2.1) × 10 ⁻⁴
Γ_5	$\pi^+\pi^-\pi^0\pi^0$	(3.3 ±0.4) %
Γ_6	$\rho^+\pi^-\pi^0 + \text{c.c.}$	(2.8 ±0.4) %
Γ_7	$4\pi^0$	(3.2 ±0.4) × 10 ⁻³
Γ_8	$\pi^+\pi^-K^+K^-$	(1.75±0.14) %
Γ_9	$K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-$	(9.6 ^{+3.5} _{-2.8}) × 10 ⁻⁴
Γ_{10}	$K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	(7.8 ^{+1.9} _{-2.4}) × 10 ⁻⁴
Γ_{11}	$K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	(6.1 ±1.9) × 10 ⁻³
Γ_{12}	$K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	< 2.6 × 10 ⁻³ CL=90%
Γ_{13}	$f_0(980)f_0(980)$	(1.6 ^{+1.0} _{-0.9}) × 10 ⁻⁴
Γ_{14}	$f_0(980)f_0(2200)$	(7.8 ^{+2.0} _{-2.5}) × 10 ⁻⁴
Γ_{15}	$f_0(1370)f_0(1370)$	< 2.7 × 10 ⁻⁴ CL=90%
Γ_{16}	$f_0(1370)f_0(1500)$	< 1.7 × 10 ⁻⁴ CL=90%
Γ_{17}	$f_0(1370)f_0(1710)$	(6.6 ^{+3.5} _{-2.3}) × 10 ⁻⁴
Γ_{18}	$f_0(1500)f_0(1370)$	< 1.3 × 10 ⁻⁴ CL=90%
Γ_{19}	$f_0(1500)f_0(1500)$	< 5 × 10 ⁻⁵ CL=90%
Γ_{20}	$f_0(1500)f_0(1710)$	< 7 × 10 ⁻⁵ CL=90%
Γ_{21}	$K^+K^-\pi^+\pi^-\pi^0$	(8.6 ±0.9) × 10 ⁻³
Γ_{22}	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	(4.2 ±0.4) × 10 ⁻³
Γ_{23}	$K^+K^-\pi^0\pi^0$	(5.4 ±0.9) × 10 ⁻³
Γ_{24}	$K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(2.44±0.33) %
Γ_{25}	$\rho^+K^-K^0 + \text{c.c.}$	(1.18±0.21) %
Γ_{26}	$K^*(892)^-K^+\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(4.5 ±1.1) × 10 ⁻³
Γ_{27}	$K_S^0 K_S^0 \pi^+ \pi^-$	(5.6 ±1.0) × 10 ⁻³
Γ_{28}	$K^+K^-\eta\pi^0$	(3.0 ±0.7) × 10 ⁻³
Γ_{29}	$3(\pi^+\pi^-)$	(1.20±0.18) %
Γ_{30}	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	(7.2 ±1.6) × 10 ⁻³
Γ_{31}	$K^*(892)^0\bar{K}^*(892)^0$	(1.7 ±0.6) × 10 ⁻³
Γ_{32}	$\pi\pi$	(8.33±0.35) × 10 ⁻³
Γ_{33}	$\pi^0\eta$	< 1.8 × 10 ⁻⁴
Γ_{34}	$\pi^0\eta'$	< 1.1 × 10 ⁻³

NODE=M056;CLUMP=A
DESIG=3

DESIG=9

DESIG=54

DESIG=20

DESIG=61

DESIG=62

DESIG=70

DESIG=5

DESIG=31

DESIG=32

DESIG=33

DESIG=34

DESIG=23

DESIG=24

DESIG=25

DESIG=26

DESIG=27

DESIG=28

DESIG=29

DESIG=30

DESIG=75

DESIG=87

DESIG=63

DESIG=65

DESIG=66

DESIG=67

DESIG=41

DESIG=68

DESIG=4

DESIG=10

DESIG=21

DESIG=18

DESIG=35

DESIG=36

Γ ₃₅	$\pi^0 \eta_c$	$< 1.6 \times 10^{-3}$	CL=90%	DESIG=86
Γ ₃₆	$\eta \eta$	$(2.95 \pm 0.19) \times 10^{-3}$		DESIG=13
Γ ₃₇	$\eta \eta'$	$< 2.3 \times 10^{-4}$	CL=90%	DESIG=37
Γ ₃₈	$\eta' \eta'$	$(1.96 \pm 0.21) \times 10^{-3}$		DESIG=46
Γ ₃₉	$\omega \omega$	$(9.5 \pm 1.1) \times 10^{-4}$		DESIG=22
Γ ₄₀	$\omega \phi$	$(1.16 \pm 0.21) \times 10^{-4}$		DESIG=76
Γ ₄₁	$\omega K^+ K^-$	$(1.94 \pm 0.21) \times 10^{-3}$		DESIG=88
Γ ₄₂	$K^+ K^-$	$(5.91 \pm 0.32) \times 10^{-3}$		DESIG=2
Γ ₄₃	$K_S^0 K_S^0$	$(3.10 \pm 0.18) \times 10^{-3}$		DESIG=15
Γ ₄₄	$\pi^+ \pi^- \eta$	$< 1.9 \times 10^{-4}$	CL=90%	DESIG=50
Γ ₄₅	$\pi^+ \pi^- \eta'$	$< 3.5 \times 10^{-4}$	CL=90%	DESIG=53
Γ ₄₆	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$< 9 \times 10^{-5}$	CL=90%	DESIG=17
Γ ₄₇	$K^+ K^- \pi^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=47
Γ ₄₈	$K^+ K^- \eta$	$< 2.2 \times 10^{-4}$	CL=90%	DESIG=51
Γ ₄₉	$K^+ K^- K_S^0 K_S^0$	$(1.4 \pm 0.5) \times 10^{-3}$		DESIG=42
Γ ₅₀	$K^+ K^- K^+ K^-$	$(2.75 \pm 0.28) \times 10^{-3}$		DESIG=14
Γ ₅₁	$K^+ K^- \phi$	$(9.5 \pm 2.4) \times 10^{-4}$		DESIG=44
Γ ₅₂	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(3.7 \pm 0.6) \times 10^{-3}$		DESIG=91
Γ ₅₃	$K^+ K^- \pi^0 \phi$	$(1.90 \pm 0.35) \times 10^{-3}$		DESIG=92
Γ ₅₄	$\phi \pi^+ \pi^- \pi^0$	$(1.18 \pm 0.15) \times 10^{-3}$		DESIG=89
Γ ₅₅	$\phi \phi$	$(7.7 \pm 0.7) \times 10^{-4}$		DESIG=16
Γ ₅₆	$\rho \bar{\rho}$	$(2.25 \pm 0.09) \times 10^{-4}$		DESIG=11
Γ ₅₇	$\rho \bar{\rho} \pi^0$	$(6.8 \pm 0.7) \times 10^{-4}$	S=1.3	DESIG=48
Γ ₅₈	$\rho \bar{\rho} \eta$	$(3.5 \pm 0.4) \times 10^{-4}$		DESIG=52
Γ ₅₉	$\rho \bar{\rho} \omega$	$(5.1 \pm 0.6) \times 10^{-4}$		DESIG=69
Γ ₆₀	$\rho \bar{\rho} \phi$	$(5.9 \pm 1.4) \times 10^{-5}$		DESIG=74
Γ ₆₁	$\rho \bar{\rho} \pi^+ \pi^-$	$(2.1 \pm 0.7) \times 10^{-3}$	S=1.4	DESIG=8
Γ ₆₂	$\rho \bar{\rho} \pi^0 \pi^0$	$(1.02 \pm 0.27) \times 10^{-3}$		DESIG=64
Γ ₆₃	$\rho \bar{\rho} K^+ K^- (\text{non-resonant})$	$(1.19 \pm 0.26) \times 10^{-4}$		DESIG=71
Γ ₆₄	$\rho \bar{\rho} K_S^0 K_S^0$	$< 8.8 \times 10^{-4}$	CL=90%	DESIG=40
Γ ₆₅	$\rho \bar{n} \pi^-$	$(1.24 \pm 0.11) \times 10^{-3}$		DESIG=43
Γ ₆₆	$\bar{\rho} n \pi^+$	$(1.34 \pm 0.12) \times 10^{-3}$		DESIG=82
Γ ₆₇	$\rho \bar{n} \pi^- \pi^0$	$(2.29 \pm 0.21) \times 10^{-3}$		DESIG=83
Γ ₆₈	$\bar{\rho} n \pi^+ \pi^0$	$(2.16 \pm 0.18) \times 10^{-3}$		DESIG=84
Γ ₆₉	$\Lambda \bar{\Lambda}$	$(3.21 \pm 0.25) \times 10^{-4}$		DESIG=19
Γ ₇₀	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(1.15 \pm 0.13) \times 10^{-3}$		DESIG=38
Γ ₇₁	$\Lambda \bar{\Lambda} \pi^+ \pi^- (\text{non-resonant})$	$< 5 \times 10^{-4}$	CL=90%	DESIG=77
Γ ₇₂	$\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}$	$< 5 \times 10^{-4}$	CL=90%	DESIG=78
Γ ₇₃	$\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}$	$< 5 \times 10^{-4}$	CL=90%	DESIG=79
Γ ₇₄	$K^+ \bar{p} \Lambda + \text{c.c.}$	$(1.22 \pm 0.12) \times 10^{-3}$	S=1.3	DESIG=49
Γ ₇₅	$K^+ \bar{p} \Lambda(1520) + \text{c.c.}$	$(2.9 \pm 0.7) \times 10^{-4}$		DESIG=72
Γ ₇₆	$\Lambda(1520) \bar{\Lambda}(1520)$	$(3.1 \pm 1.2) \times 10^{-4}$		DESIG=73
Γ ₇₇	$\Sigma^0 \bar{\Sigma}^0$	$(4.4 \pm 0.4) \times 10^{-4}$		DESIG=58
Γ ₇₈	$\Sigma^+ \bar{\Sigma}^-$	$(3.9 \pm 0.7) \times 10^{-4}$	S=1.7	DESIG=59
Γ ₇₉	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$(1.6 \pm 0.6) \times 10^{-4}$		DESIG=80
Γ ₈₀	$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$(2.3 \pm 0.6) \times 10^{-4}$		DESIG=81
Γ ₈₁	$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(1.90 \pm 0.34) \times 10^{-4}$		DESIG=85
Γ ₈₂	$\Xi^0 \bar{\Xi}^0$	$(3.1 \pm 0.8) \times 10^{-4}$		DESIG=60
Γ ₈₃	$\Xi^- \bar{\Xi}^+$	$(4.7 \pm 0.7) \times 10^{-4}$		DESIG=39
Γ ₈₄	$\eta_c \pi^+ \pi^-$	$< 7 \times 10^{-4}$	CL=90%	DESIG=90
Radiative decays				
Γ ₈₅	$\gamma J/\psi(1S)$	$(1.27 \pm 0.06) \%$		NODE=M056;CLUMP=B DESIG=6
Γ ₈₆	$\gamma \rho^0$	$< 9 \times 10^{-6}$	CL=90%	DESIG=55
Γ ₈₇	$\gamma \omega$	$< 8 \times 10^{-6}$	CL=90%	DESIG=56
Γ ₈₈	$\gamma \phi$	$< 6 \times 10^{-6}$	CL=90%	DESIG=57
Γ ₈₉	$\gamma \gamma$	$(2.23 \pm 0.13) \times 10^{-4}$		DESIG=7

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 240 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 342.4$ for 191 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_2	25									
x_8	14	4								
x_{30}	7	2	29							
x_{32}	15	4	16	5						
x_{36}	8	2	9	3	21					
x_{42}	13	3	14	5	28	17				
x_{43}	12	3	13	4	27	16	22			
x_{50}	9	2	8	3	14	8	12	11		
x_{55}	10	3	10	4	14	9	12	12	7	
x_{56}	7	2	9	3	13	6	15	15	8	8
x_{69}	8	2	9	3	20	12	17	16	8	9
x_{85}	4	1	5	1	14	9	10	10	5	5
x_{89}	-16	-4	-9	-6	5	4	2	4	-2	-4
Γ	-22	-6	-17	-8	-16	-9	-15	-14	-10	-13
	x_1	x_2	x_8	x_{30}	x_{32}	x_{36}	x_{42}	x_{43}	x_{50}	x_{55}
x_{69}	11									
x_{85}	-21	8								
x_{89}	2	4	9							
Γ	-7	-9	-8	-48						
	x_{56}	x_{69}	x_{85}	x_{89}						

 $\chi_{c0}(1P)$ PARTIAL WIDTHS

$$\text{--- } \chi_{c0}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total}) \text{ ---}$$

NODE=M056217

NODE=M056223

$$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}} \quad \Gamma_{56} \Gamma_{85} / \Gamma$$

NODE=M056G1
NODE=M056G1

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

30.0 ± 2.3 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

26.6 ± 2.6 ± 1.4	392	1,2 BAGNASCO	02	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$
------------------	-----	--------------	----	------	--

48.7 ^{+11.3} _{-8.9} ± 2.4		1,2 AMBROGIANI	99B	E835	$\bar{p}p \rightarrow \gamma J/\psi$
---	--	----------------	-----	------	--------------------------------------

¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

² Values in $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$ and $(\Gamma(p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$ are not independent. The latter is used in the fit since it is less correlated to the total width.

NODE=M056G;LINKAGE=7A
NODE=M056G;LINKAGE=KS

$$\text{--- } \chi_{c0}(1P) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total}) \text{ ---}$$

NODE=M056224

$$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}} \quad \Gamma_1 \Gamma_{89} / \Gamma$$

NODE=M056G2
NODE=M056G2
NEW

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

53 ± 4 OUR FIT

[52 ± 4 eV OUR 2015 FIT]

49 ± 10 OUR AVERAGE Error includes scale factor of 1.8.

44.7 ± 3.6 ± 4.9	3.6k	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+ \pi^-)$
------------------	------	--------	----	------	---

75 ± 13 ± 8		EISENSTEIN	01	CLE2	$e^+ e^- \rightarrow e^+ e^- \chi_{c0}$
-------------	--	------------	----	------	---

$$\Gamma(\rho^0 \rho^0) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}} \quad \Gamma_3 \Gamma_{89} / \Gamma$$

NODE=M056G07
NODE=M056G07

VALUE (eV)	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	<252	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+ \pi^-)$
-----	----	------	--------	----	------	---

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
41 ± 4 OUR FIT				
38.8±3.7±4.7	1.7k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M056G08
NODE=M056G08

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{21}\Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
26±4±4	1094	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M056G01
NODE=M056G01

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{30}\Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
17 ± 4 OUR FIT				
16.7±6.1±3.0	495 ± 182	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M056G09
NODE=M056G09

$$\Gamma(K^*(892)^0\bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{31}\Gamma_{89}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

NODE=M056G10
NODE=M056G10

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<6	90	<148	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{32}\Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M056G3
NODE=M056G3

19.5± 1.4 OUR FIT

23 ± 5 OUR AVERAGE

29.7 ^{+17.4} _{-12.0} ± 4.8	103 ⁺⁶⁰ ₋₄₂	¹ UEHARA	09	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
22.7 ± 3.2 ± 3.5	129 ± 18	² NAKAZAWA	05	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

¹We multiplied the measurement by 3 to convert from $\pi^0\pi^0$ to $\pi\pi$. Interference with the continuum included.

²We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

NODE=M056G3;LINKAGE=UE

NODE=M056G;LINKAGE=NA

$$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M056G06
NODE=M056G06

9.4±2.3±1.2 22 ¹ UEHARA 10A BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

¹Interference with the continuum not included.

NODE=M056G06;LINKAGE=UE

$$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{39}\Gamma_{89}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M056G02
NODE=M056G02

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.9 90 ¹ LIU 12B BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

¹Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M056G02;LINKAGE=LI

$$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{40}\Gamma_{89}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M056G03
NODE=M056G03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.34 90 ¹ LIU 12B BELL $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M056G03;LINKAGE=LI

$$\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M056G4
NODE=M056G4

13.9±1.1 OUR FIT

14.3±1.6±2.3 153 ± 17 NAKAZAWA 05 BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$

$$\Gamma(K_S^0K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{43}\Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M056G5
NODE=M056G5

7.3 ± 0.6 OUR FIT

8.7 ± 1.7 ± 0.9 266 ¹ UEHARA 13 BELL $\gamma\gamma \rightarrow K_S^0K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.00 ± 0.65 ± 0.71 134 ± 12 CHEN 07B BELL $e^+e^- \rightarrow e^+e^-\chi_{c0}$

¹Supersedes CHEN 07B.

NODE=M056G5;LINKAGE=UE

$$\Gamma(K^+K^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{50}\Gamma_{89}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M056G11
NODE=M056G11

6.4±0.7 OUR FIT

7.9±1.3±1.1 215 ± 36 UEHARA 08 BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+K^-)$

$$\frac{\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\Gamma_{55} \Gamma_{89}/\Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.82±0.19 OUR FIT

1.72±0.33±0.14 56 ± 11 ¹ LIU 12B BELL $\gamma\gamma \rightarrow 2(K^+ K^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ± 0.9 ± 0.4 23.6 ± 9.6 UEHARA 08 BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+ K^-)$

¹Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

NODE=M056G12
NODE=M056G12

NODE=M056G12;LINKAGE=LI

$\chi_{c0}(1P)$ BRANCHING RATIOS

NODE=M056220

HADRONIC DECAYS

NODE=M056305

$$\frac{\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}}{\Gamma_1/\Gamma}$$

VALUE	DOCUMENT ID
-------	-------------

0.0224±0.0018 OUR FIT

NODE=M056R2
NODE=M056R2

$$\frac{\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-))}{\Gamma_2/\Gamma_1}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.39±0.12 OUR FIT

0.39±0.12

TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R54
NODE=M056R54

$$\frac{\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}}{\Gamma_2/\Gamma}$$

VALUE	DOCUMENT ID
-------	-------------

0.0087±0.0028 OUR FIT

NODE=M056R9
NODE=M056R9

$$\frac{\Gamma(f_0(980) f_0(980))/\Gamma_{\text{total}}}{\Gamma_4/\Gamma}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.5±2.1±0.2 36 ± 9 ¹ ABLIKIM 04G BES $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ABLIKIM 04G reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980) f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(6.5 \pm 1.6 \pm 1.3) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R24
NODE=M056R24

NODE=M056R24;LINKAGE=AB

$$\frac{\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}}{\Gamma_5/\Gamma}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

3.3±0.4±0.1 1751.4 ¹ HE 08B CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹HE 08B reports $3.54 \pm 0.10 \pm 0.43 \pm 0.18$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R62
NODE=M056R62

NODE=M056R62;LINKAGE=HE

$$\frac{\Gamma(\rho^+ \pi^- \pi^0 + \text{c.c.})/\Gamma_{\text{total}}}{\Gamma_6/\Gamma}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

2.8±0.4±0.1 1358.5 ^{1,2} HE 08B CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹HE 08B reports $3.04 \pm 0.18 \pm 0.42 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+ \pi^- \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Calculated by us. We have added the values from HE 08B for $\rho^+ \pi^- \pi^0$ and $\rho^- \pi^+ \pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M056R63
NODE=M056R63

NODE=M056R63;LINKAGE=HE

NODE=M056R63;LINKAGE=OC

$$\frac{\Gamma(4\pi^0)/\Gamma_{\text{total}}}{\Gamma_7/\Gamma}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.2±0.4±0.1 3296 ¹ ABLIKIM 11A BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

¹ABLIKIM 11A reports $(3.34 \pm 0.06 \pm 0.44) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R71
NODE=M056R71

NODE=M056R71;LINKAGE=AB

$$\frac{\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}}{\Gamma_8/\Gamma}$$

VALUE (units 10^{-3})	DOCUMENT ID
--------------------------	-------------

17.5±1.4 OUR FIT

NODE=M056R3
NODE=M056R3

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma(\pi^+\pi^-K^+K^-)$$

 Γ_{30}/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M056R55
NODE=M056R55

0.41±0.09 OUR FIT**0.41±0.10**TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c0}$

$$\Gamma(K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$$

 Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R36
NODE=M056R36

9.6^{+3.5}_{-2.8}±0.3	83	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
--	----	----------------------	----------	---

¹ ABLIKIM 05Q reports $(10.44 \pm 2.37^{+3.05}_{-1.90}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R36;LINKAGE=AB

$$\Gamma(K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$$

 Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R37
NODE=M056R37

7.8^{+1.9}_{-2.4}±0.2	62	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
--	----	----------------------	----------	---

¹ ABLIKIM 05Q reports $(8.49 \pm 1.66^{+1.32}_{-1.99}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R37;LINKAGE=AB

$$\Gamma(K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$$

 Γ_{11}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R38
NODE=M056R38

6.1^{+1.9}_{-1.8}±0.2	68	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
--	----	----------------------	----------	---

¹ ABLIKIM 05Q reports $(6.66 \pm 1.31^{+1.60}_{-1.51}) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The measurement assumes $B(K_1(1270) \rightarrow K\rho(770)) = 42 \pm 6\%$.

NODE=M056R38;LINKAGE=AB

$$\Gamma(K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$$

 Γ_{12}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M056R39
NODE=M056R39

<2.6	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
----------------	----	----------------------	----------	---

¹ ABLIKIM 05Q reports $< 2.85 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$. The measurement assumes $B(K_1(1400) \rightarrow K^*(892)\pi) = 94 \pm 6\%$.

NODE=M056R39;LINKAGE=AB

$$\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}}$$

 Γ_{13}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R28
NODE=M056R28

15.9^{+10.2}_{-8.8}±0.4	28	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
--	----	----------------------	----------	---

¹ ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.59 \pm 0.50^{+0.89}_{-0.72}) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. One of the $f_0(980)$ mesons is identified via decay to $\pi^+\pi^-$ while the other via K^+K^- decay.

NODE=M056R28;LINKAGE=AB

$$\Gamma(f_0(980)f_0(2200))/\Gamma_{\text{total}}$$

 Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R29
NODE=M056R29

7.8^{+2.0}_{-2.5}±0.2	77	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
--	----	----------------------	----------	---

¹ ABLIKIM 05Q reports $(8.42 \pm 1.42^{+1.65}_{-2.29}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(2200))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(980) \rightarrow \pi^+\pi^-$ and $f_0(2200) \rightarrow K^+K^-$ decays.

NODE=M056R29;LINKAGE=AB

$\Gamma(f_0(1370)f_0(1370))/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R30
NODE=M056R30

¹ ABLIKIM 05Q reports $< 2.9 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$. One of the $f_0(1370)$ mesons is identified via decay to $\pi^+\pi^-$ while the other via K^+K^- decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R30;LINKAGE=AB

$\Gamma(f_0(1370)f_0(1500))/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R31
NODE=M056R31

¹ ABLIKIM 05Q reports $< 1.8 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+\pi^-$ and $f_0(1500) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R31;LINKAGE=AB

$\Gamma(f_0(1370)f_0(1710))/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.6^{+3.5}_{-2.3} \pm 0.2$	61	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R32
NODE=M056R32

¹ ABLIKIM 05Q reports $(7.12 \pm 1.85^{+3.28}_{-1.68}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+\pi^-$ and $f_0(1710) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R32;LINKAGE=AB

$\Gamma(f_0(1500)f_0(1370))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R33
NODE=M056R33

¹ ABLIKIM 05Q reports $< 1.4 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+\pi^-$ and $f_0(1370) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R33;LINKAGE=AB

$\Gamma(f_0(1500)f_0(1500))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R34
NODE=M056R34

¹ ABLIKIM 05Q reports $< 0.55 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$. One of the $f_0(1500)$ is identified via decay to $\pi^+\pi^-$ while the other via K^+K^- decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R34;LINKAGE=AB

$\Gamma(f_0(1500) f_0(1710))/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
¹ ABLIKIM 05Q reports $< 0.73 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500) f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+ \pi^-$ and $f_0(1710) \rightarrow K^+ K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

NODE=M056R35
NODE=M056R35

NODE=M056R35;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.61±0.13±0.94	9.0k	¹ ABLIKIM	13B	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_c^0$
¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_c^0 \gamma) = (9.68 \pm 0.31)\%$.				

NODE=M056R85
NODE=M056R85

NODE=M056R85;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.22±0.10±0.43	2.7k	¹ ABLIKIM	13B	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_c^0$
¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_c^0 \gamma) = (9.68 \pm 0.31)\%$.				

NODE=M056R86
NODE=M056R86

NODE=M056R86;LINKAGE=A

 $\Gamma(K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.54±0.09±0.01	213.5	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹ HE 08B reports $0.59 \pm 0.05 \pm 0.08 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R64
NODE=M056R64

NODE=M056R64;LINKAGE=HE

 $\Gamma(K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.44±0.32±0.07	401.7	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹ HE 08B reports $2.64 \pm 0.15 \pm 0.31 \pm 0.14$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R66
NODE=M056R66

NODE=M056R66;LINKAGE=HE

 $\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.18±0.20±0.03	179.7	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹ HE 08B reports $1.28 \pm 0.16 \pm 0.15 \pm 0.07$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R67
NODE=M056R67

NODE=M056R67;LINKAGE=HE

 $\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.45±0.11±0.01	64.1	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹ HE 08B reports $0.49 \pm 0.10 \pm 0.07 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R68
NODE=M056R68

NODE=M056R68;LINKAGE=HE

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.6±1.0±0.2	152 ± 14	¹ ABLIKIM	05o	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$
¹ ABLIKIM 05o reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(0.558 \pm 0.051 \pm 0.089) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R47
NODE=M056R47

NODE=M056R47;LINKAGE=AB

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{28}/Γ
0.30±0.07±0.01	56.4	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	NODE=M056R69 NODE=M056R69

¹ HE 08B reports $0.32 \pm 0.05 \pm 0.05 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R69;LINKAGE=HE

 $\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_{29}/Γ
12.0±1.8 OUR EVALUATION	Treating systematic error as correlated.			NODE=M056R4 NODE=M056R4
12.0±1.7 OUR AVERAGE				→ UNCHECKED ←

11.7±1.0±1.9

¹ BAI 99B BES $\psi(2S) \rightarrow \gamma \chi_{c0}$

12.5±2.9±0.5

¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

NODE=M056R;LINKAGE=X1

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	Γ_{30}/Γ
0.0072±0.0016 OUR FIT		NODE=M056R10 NODE=M056R10

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{31}/Γ
1.68^{+0.59}_{-0.53}±0.05	64	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	NODE=M056R26 NODE=M056R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.53±0.39±0.04 30±6 ^{2,3} ABLIKIM 04H BES Repl. by ABLIKIM 05Q

¹ ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.168 \pm 0.035^{+0.047}_{-0.040}) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R26;LINKAGE=A1

² Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

NODE=M056R;LINKAGE=AL
NODE=M056R26;LINKAGE=AB

³ ABLIKIM 04H reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.53 \pm 0.29 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	Γ_{32}/Γ
8.33±0.35 OUR FIT		NODE=M056R22 NODE=M056R22

 $\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{35}/Γ
<1.6 × 10⁻³	90	¹ ABLIKIM	15N BES3	$\psi(2S) e^+ e^- \rightarrow \gamma \pi^0 \eta_c$	NODE=M056R00 NODE=M056R00

¹ Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$.

NODE=M056R00;LINKAGE=A

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	Γ_{36}/Γ
2.95±0.19 OUR FIT		NODE=M056R13 NODE=M056R13

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{36}/Γ_{32}
0.354±0.025 OUR FIT				NODE=M056R20 NODE=M056R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.09 ^{+0.03}_{-0.02} ¹ ANDREOTTI 05C E835 $\bar{p} p \rightarrow 2$ mesons0.24 ± 0.10 ± 0.08 ¹ BAI 03C BES $\psi(2S) \rightarrow 5\gamma$

¹ We have multiplied $\pi^0 \pi^0$ measurement by 3 to obtain $\pi\pi$.

NODE=M056R;LINKAGE=D1

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<0.23	90	35 ± 13	¹ ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta'\eta$
-----------------	----	---------	--------------------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.5	90		² ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
------	----	--	--------------------	---------	--

¹ ASNER 09 reports $< 0.25 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R03
NODE=M056R03

NODE=M056R03;LINKAGE=AS

NODE=M056R03;LINKAGE=AD

² Superseded by ASNER 09. ADAMS 07 reports $< 0.5 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.96 ± 0.20 ± 0.05	0.4k	¹ ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta'\eta'$
---------------------------	------	--------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.57 ± 0.40 ± 0.04	23	² ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
--------------------	----	--------------------	---------	--

¹ ASNER 09 reports $(2.12 \pm 0.13 \pm 0.21) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R04
NODE=M056R04

NODE=M056R04;LINKAGE=AS

NODE=M056R04;LINKAGE=AD

² Superseded by ASNER 09. ADAMS 07 reports $(1.7 \pm 0.4 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 0.0922 \pm 0.0011 \pm 0.0046$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.95 ± 0.11 OUR AVERAGE

0.91 ± 0.11 ± 0.02	991	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
--------------------	-----	----------------------	----------	---------------------------------------

2.1 ± 0.6 ± 0.1	38.1 ± 9.6	² ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma 6\pi$
-----------------	------------	----------------------	----------	--

¹ ABLIKIM 11K reports $(0.95 \pm 0.03 \pm 0.11) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R27
NODE=M056R27

NODE=M056R27;LINKAGE=AL

² ABLIKIM 05N reports $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (0.212 \pm 0.053 \pm 0.037) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R27;LINKAGE=AB

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.16 ± 0.21 ± 0.03	76	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
---------------------------	----	----------------------	----------	---------------------------------------

¹ ABLIKIM 11K reports $(1.2 \pm 0.1 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R76
NODE=M056R76

NODE=M056R76;LINKAGE=AL

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.94 ± 0.06 ± 0.20	1.4k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_c^0$
---------------------------	------	----------------------	----------	---

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_c^0 \gamma) = (9.68 \pm 0.31)\%$.

NODE=M056R87
NODE=M056R87

NODE=M056R87;LINKAGE=A

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	DOCUMENT ID
--------------------------	-------------

5.91 ± 0.32 OUR FIT	
----------------------------	--

NODE=M056R6
NODE=M056R6

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID

 Γ_{43}/Γ NODE=M056R15
NODE=M056R15**3.10±0.18 OUR FIT** $\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$

VALUE

DOCUMENT ID

TECN

COMMENT

 Γ_{43}/Γ_{32} NODE=M056R53
NODE=M056R53**0.372±0.023 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ±0.05 ±0.05 1,2 CHEN 07B BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c0}$ ¹ Using $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from the $\pi^+ \pi^-$ measurement of NAKAZAWA 05 rescaled by 3/2 to convert to $\pi\pi$.

NODE=M056R53;LINKAGE=CH

² Not independent from other measurements.

NODE=M056R53;LINKAGE=NI

 $\Gamma(K_S^0 K_S^0)/\Gamma(K^+ K^-)$

VALUE

DOCUMENT ID

TECN

COMMENT

 Γ_{43}/Γ_{42} NODE=M056R52
NODE=M056R52**0.52±0.04 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.49±0.07±0.08 1,2 CHEN 07B BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c0}$ ¹ Using $\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from NAKAZAWA 05.

NODE=M056R52;LINKAGE=CH

² Not independent from other measurements.

NODE=M056R52;LINKAGE=NI

 $\Gamma(\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{44}/Γ NODE=M056R08
NODE=M056R08**<0.19** 90 1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.0 90 2 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$ ¹ ATHAR 07 reports $< 0.21 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R08;LINKAGE=AT

² ABLIKIM 06R reports $< 1.1 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R08;LINKAGE=AB

 $\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{45}/Γ NODE=M056R51
NODE=M056R51**<0.35** 90 1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ ¹ ATHAR 07 reports $< 0.38 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R51;LINKAGE=AT

 $\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{46}/Γ NODE=M056R17
NODE=M056R17**<0.09** 90 1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6 90 2,3 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$ <0.7 90 3,4 BAI 99B BES $\psi(2S) \rightarrow \gamma \chi_{c0}$ ¹ ATHAR 07 reports $< 0.10 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R17;LINKAGE=AT

² ABLIKIM 06R reports $< 0.70 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R17;LINKAGE=AB

³ We have multiplied the $K_S^0 K^+ \pi^-$ measurement by a factor of 2 to convert to $K^0 K^+ \pi^-$.

NODE=M056R17;LINKAGE=BA

⁴ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

NODE=M056R17;LINKAGE=X1

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	¹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M056R05
NODE=M056R05

¹ ATHAR 07 reports $< 0.06 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R05;LINKAGE=AT

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	¹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M056R09
NODE=M056R09

¹ ATHAR 07 reports $< 0.24 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R09;LINKAGE=AT

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.38 \pm 0.46 \pm 0.04$	16.8 ± 4.8	¹ ABLIKIM	050	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R48
NODE=M056R48

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.138 \pm 0.039 \pm 0.025) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R48;LINKAGE=AB

 $\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	DOCUMENT ID
2.75 ± 0.28 OUR FIT	

NODE=M056R14
NODE=M056R14 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.95 \pm 0.24 \pm 0.03$	38	¹ ABLIKIM	06T	BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M056R01
NODE=M056R01

¹ ABLIKIM 06T reports $(1.03 \pm 0.22 \pm 0.15) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R01;LINKAGE=AB

 $\Gamma(K^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$3.68 \pm 0.30 \pm 0.50$	ABLIKIM	15M	BES3 $\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R90
NODE=M056R90 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.90 \pm 0.14 \pm 0.32$	ABLIKIM	15M	BES3 $\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R91
NODE=M056R91 $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.18 \pm 0.07 \pm 0.13$	538	¹ ABLIKIM	13B	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_c^0$

NODE=M056R88
NODE=M056R88

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_c^0 \gamma) = (9.68 \pm 0.31)\%$.

NODE=M056R88;LINKAGE=A

 $\Gamma(\phi \phi)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-3})	DOCUMENT ID
0.77 ± 0.07 OUR FIT	

NODE=M056R16
NODE=M056R16 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-4})	DOCUMENT ID
2.25 ± 0.09 OUR FIT	

NODE=M056R11
NODE=M056R11

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{57}/Γ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.68±0.07 OUR AVERAGE Error includes scale factor of 1.3.

0.72±0.06±0.02

¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$

0.54±0.11±0.01

² ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(7.76 \pm 0.37 \pm 0.51 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.59 \pm 0.10 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R06
NODE=M056R06

NODE=M056R06;LINKAGE=ON

NODE=M056R06;LINKAGE=AT

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{58}/Γ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.35±0.04 OUR AVERAGE

0.34±0.04±0.01

¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$

0.36±0.11±0.01

² ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(3.73 \pm 0.38 \pm 0.28 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.39 \pm 0.11 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R50
NODE=M056R50

NODE=M056R50;LINKAGE=ON

NODE=M056R50;LINKAGE=AT

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{59}/Γ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.51±0.05±0.01¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$

¹ ONYISI 10 reports $(5.57 \pm 0.48 \pm 0.42 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R70
NODE=M056R70

NODE=M056R70;LINKAGE=ON

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{60}/Γ VALUE (units 10^{-5})

EVTS

DOCUMENT ID TECN COMMENT

5.9±1.4±0.2

42 ± 8

¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$

¹ ABLIKIM 11F reports $(6.12 \pm 1.18 \pm 0.86) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R75
NODE=M056R75

NODE=M056R75;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

2.1 ±0.7 OUR EVALUATION Error includes scale factor of 1.4. Treating systematic error as correlated.**2.1 ±1.0 OUR AVERAGE** Error includes scale factor of 2.0.

1.57±0.21±0.53

¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c0}$

4.20±1.15±0.18

¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M056R7
NODE=M056R7

→ UNCHECKED ←

NODE=M056R7;LINKAGE=X1

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.102±0.027±0.003	39.5	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M056R65
 NODE=M056R65

¹ HE 08B reports $0.11 \pm 0.02 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R65;LINKAGE=HE

 $\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19±0.26±0.03	48 ± 8	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M056R72
 NODE=M056R72

¹ ABLIKIM 11F reports $(1.24 \pm 0.20 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R72;LINKAGE=AB

 $\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8.8	90	¹ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$

NODE=M056R46
 NODE=M056R46

¹ Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

NODE=M056R;LINKAGE=AB

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±1.1 OUR AVERAGE				
12.6±1.1±0.3	5150	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
11.0±3.0±0.3		² ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$

NODE=M056R49
 NODE=M056R49

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.26 \pm 0.02 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R49;LINKAGE=AL

² ABLIKIM 06I reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.10 \pm 0.24 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R49;LINKAGE=AB

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
13.4±1.1±0.4	5808	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+$

NODE=M056R82
 NODE=M056R82

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.34 \pm 0.03 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R82;LINKAGE=AL

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
22.9±2.0±0.6	2480	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

NODE=M056R83
 NODE=M056R83

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.29 \pm 0.08 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R83;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.6±1.7±0.6	2757	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+\pi^0$

NODE=M056R84
 NODE=M056R84

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.16 \pm 0.07 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R84;LINKAGE=AL

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	DOCUMENT ID
3.21±0.25 OUR FIT	

NODE=M056R23
 NODE=M056R23

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
115±12±3		426	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M056R44
 NODE=M056R44

••• We do not use the following data for averages, fits, limits, etc. •••

<400	90		² ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$
------	----	--	----------------------	----------	--

¹ ABLIKIM 12I reports $(119.0 \pm 6.4 \pm 11.4) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R44;LINKAGE=AL

² Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

NODE=M056R44;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M056R77
 NODE=M056R77

¹ ABLIKIM 12I reports $< 54 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R77;LINKAGE=AL

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

NODE=M056R78
 NODE=M056R78

¹ ABLIKIM 12I reports $< 55 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R78;LINKAGE=AL

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$

NODE=M056R79
 NODE=M056R79

¹ ABLIKIM 12I reports $< 50 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R79;LINKAGE=AL

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.22±0.12 OUR AVERAGE		Error includes scale factor of 1.3.		
1.28±0.09±0.03	9k	^{1,2} ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$
0.99±0.19±0.03		³ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M056R07
 NODE=M056R07

¹ ABLIKIM 13D reports $(1.32 \pm 0.03 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R07;LINKAGE=AB

² Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.

NODE=M056R07;LINKAGE=LB

³ ATHAR 07 reports $(1.07 \pm 0.17 \pm 0.12) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R07;LINKAGE=AT

 $\Gamma(K^+\bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9±0.7±0.1	62 ± 12	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M056R73
 NODE=M056R73

¹ ABLIKIM 11F reports $(3.00 \pm 0.58 \pm 0.50) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R73;LINKAGE=AB

$\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±1.2±0.1	28 ± 10	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NODE=M056R74
NODE=M056R74

¹ ABLIKIM 11F reports $(3.18 \pm 1.11 \pm 0.53) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R74;LINKAGE=AB

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±0.4 OUR AVERAGE				
4.6±0.5±0.1	243	¹ ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0\bar{\Sigma}^0$
4.1±0.6±0.1	78 ± 10	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0\bar{\Sigma}^0$

NODE=M056R59
NODE=M056R59

¹ ABLIKIM 13H reports $(4.78 \pm 0.34 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R59;LINKAGE=AB

² NAIK 08 reports $(4.41 \pm 0.56 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R59;LINKAGE=NA

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9±0.7 OUR AVERAGE				Error includes scale factor of 1.7.
4.4±0.5±0.1	148	¹ ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+\bar{\Sigma}^-$
3.0±0.6±0.1	39 ± 7	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+\bar{\Sigma}^-$

NODE=M056R60
NODE=M056R60

¹ ABLIKIM 13H reports $(4.54 \pm 0.42 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R60;LINKAGE=AB

² NAIK 08 reports $(3.25 \pm 0.57 \pm 0.43) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R60;LINKAGE=NA

 $\Gamma(\Sigma(1385)^+\bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
15.9±5.7±0.4	27	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

NODE=M056R80
NODE=M056R80

¹ ABLIKIM 12I reports $(16.4 \pm 5.7 \pm 1.6) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+\bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R80;LINKAGE=AL

 $\Gamma(\Sigma(1385)^-\bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
23±6±1	33	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

NODE=M056R81
NODE=M056R81

¹ ABLIKIM 12I reports $(23.5 \pm 6.2 \pm 2.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^-\bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R81;LINKAGE=AL

$\Gamma(K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.34±0.05	57	¹ ABLIKIM 15i	BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \Xi^+ + \text{c.c.}$

NODE=M056R92
 NODE=M056R92

¹ ABLIKIM 15i reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.90 \pm 0.30 \pm 0.16) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R92;LINKAGE=A

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±0.8±0.1	23.3 ± 4.9	¹ NAIK 08	CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$

NODE=M056R61
 NODE=M056R61

¹ NAIK 08 reports $(3.34 \pm 0.70 \pm 0.48) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R61;LINKAGE=NA

 $\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.7±0.7±0.1		95 ± 11	¹ NAIK 08	CLEO	$\psi(2S) \rightarrow \gamma \Xi^- \Xi^+$

NODE=M056R45
 NODE=M056R45

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10.3	90	² ABLIKIM 06D	BES2	$\psi(2S) \rightarrow \chi_{c0} \gamma$
-------	----	--------------------------	------	---

¹ NAIK 08 reports $(5.14 \pm 0.60 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R45;LINKAGE=NA

² Using $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.2 \pm 0.5)\%$

NODE=M056R45;LINKAGE=AB

 $\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 7 × 10⁻⁴	90	^{1,2} ABLIKIM 13B	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_c^0$

NODE=M056R89
 NODE=M056R89

• • • We do not use the following data for averages, fits, limits, etc. • • •

<41 × 10 ⁻⁴	90	^{1,3} ABLIKIM 13B	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_c^0$
------------------------	----	----------------------------	------	--

OCCUR=2

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_c^0 \gamma) = (9.68 \pm 0.31)\%$.

NODE=M056R89;LINKAGE=A

² From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

NODE=M056R89;LINKAGE=B

³ From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

NODE=M056R89;LINKAGE=C

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$ $\Gamma_{56}/\Gamma \times \Gamma_{32}/\Gamma$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
18.8±1.2 OUR FIT	¹ ANDREOTTI 03	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0 \pi^0$

NODE=M056R21
 NODE=M056R21

15.3±2.4±0.8

¹ We have multiplied $B(p\bar{p}) \cdot B(\pi^0 \pi^0)$ measurement by 3 to obtain $B(p\bar{p}) \cdot B(\pi\pi)$.

NODE=M056R;LINKAGE=AD

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0 \eta)/\Gamma_{\text{total}}$ $\Gamma_{56}/\Gamma \times \Gamma_{33}/\Gamma$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
<0.4	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$

NODE=M056R41
 NODE=M056R41

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0 \eta')/\Gamma_{\text{total}}$ $\Gamma_{56}/\Gamma \times \Gamma_{34}/\Gamma$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
<2.5	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$

NODE=M056R42
 NODE=M056R42

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$ $\Gamma_{56}/\Gamma \times \Gamma_{36}/\Gamma$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
6.6±0.5 OUR FIT	¹ ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \eta\eta$

NODE=M056R40
 NODE=M056R40

4.0±1.2^{+0.5}_{-0.3}

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta')/\Gamma_{\text{total}}$ $\Gamma_{56}/\Gamma \times \Gamma_{37}/\Gamma$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
2.1^{+2.3}_{-1.5}	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$

NODE=M056R43
 NODE=M056R43

• • • We do not use the following data for averages, fits, limits, etc. • • •

———— RADIATIVE DECAYS ————

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})

DOCUMENT ID TECN COMMENT

127 ± 6 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

200 ± 20 ± 20 ¹ ADAM 05A CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma\chi_{c0})$ from ATHAR 04.

NODE=M056310

NODE=M056R8
NODE=M056R8

NODE=M056R8;LINKAGE=AD

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6}) CL% EVTS DOCUMENT ID TECN COMMENT

< 9 90 1.2 ± 4.5 ¹ BENNETT 08A CLEO $\psi(2S) \rightarrow \gamma\gamma\rho^0$

••• We do not use the following data for averages, fits, limits, etc. •••

< 10 90 6 ± 12 ² ABLIKIM 11E BES3 $\psi(2S) \rightarrow \gamma\gamma\rho^0$

¹ BENNETT 08A reports $< 9.6 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

² ABLIKIM 11E reports $< 10.5 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R56
NODE=M056R56

NODE=M056R56;LINKAGE=BE

NODE=M056R56;LINKAGE=AB

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-6}) CL% EVTS DOCUMENT ID TECN COMMENT

< 8 90 0.0 ± 2.8 ¹ BENNETT 08A CLEO $\psi(2S) \rightarrow \gamma\gamma\omega$

••• We do not use the following data for averages, fits, limits, etc. •••

< 12 90 5 ± 11 ² ABLIKIM 11E BES3 $\psi(2S) \rightarrow \gamma\gamma\omega$

¹ BENNETT 08A reports $< 8.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

² ABLIKIM 11E reports $< 12.9 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R57
NODE=M056R57

NODE=M056R57;LINKAGE=BE

NODE=M056R57;LINKAGE=AB

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-6}) CL% EVTS DOCUMENT ID TECN COMMENT

< 6 90 0.1 ± 1.6 ¹ BENNETT 08A CLEO $\psi(2S) \rightarrow \gamma\gamma\phi$

••• We do not use the following data for averages, fits, limits, etc. •••

< 16 90 15 ± 7 ² ABLIKIM 11E BES3 $\psi(2S) \rightarrow \gamma\gamma\phi$

¹ BENNETT 08A reports $< 6.4 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

² ABLIKIM 11E reports $< 16.2 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.99 \times 10^{-2}$.

NODE=M056R58
NODE=M056R58

NODE=M056R58;LINKAGE=BE

NODE=M056R58;LINKAGE=AB

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-4}) CL% DOCUMENT ID TECN COMMENT

2.23 ± 0.13 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

< 7 90 ¹ WICHT 08 BELL $B^\pm \rightarrow K^\pm\gamma\gamma$

¹ WICHT 08 reports $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^\pm \rightarrow \chi_{c0}(1P)K^\pm)] < 0.11 \times 10^{-6}$ which we divide by our best value $B(B^\pm \rightarrow \chi_{c0}(1P)K^\pm) = 1.50 \times 10^{-4}$.

NODE=M056R1
NODE=M056R1

NODE=M056R1;LINKAGE=WI

$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$

VALUE (units 10^{-2}) DOCUMENT ID TECN COMMENT

1.76 ± 0.13 OUR FIT

2.0 ± 0.4 OUR AVERAGE

2.2 ± 0.4 ^{+0.1}_{-0.2} ¹ ANDREOTTI 04 E835 $p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$

1.45 ± 0.74 ² AMBROGIANI 00B E835 $p\bar{p} \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$

¹ The values of $B(p\bar{p})B(\gamma\gamma)$ and $B(\gamma\gamma)B(\gamma J/\psi)$ measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

² Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

Γ_{89}/Γ_{85}

NODE=M056R18
NODE=M056R18

NODE=M056R;LINKAGE=AN

NODE=M056R;LINKAGE=7A

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_{56}/\Gamma \times \Gamma_{85}/\Gamma$$

VALUE (units 10^{-7}) EVTS DOCUMENT ID TECN COMMENT

28.5±1.6 OUR FIT

28.2±2.1 OUR AVERAGE

28.0±1.9±1.3 392 1,2,3 BAGNASCO 02 E835 $p\bar{p} \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$

29.3^{+5.7}_{-4.7}±1.5 89 1,2 AMBROGIANI 99B $p\bar{p} \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$

¹ Values in $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$ and $(\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$ are not independent. The latter is used in the fit since it is less correlated to the total width.

² Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

³ Recalculated by ANDREOTTI 05A.

NODE=M056R19
NODE=M056R19

NODE=M056R;LINKAGE=KS

NODE=M056R19;LINKAGE=7A
NODE=M056R19;LINKAGE=AN

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{56}/\Gamma \times \Gamma_{89}/\Gamma$$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

5.0 ±0.4 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.52±1.18^{+0.48}_{-0.72} ¹ ANDREOTTI 04 E835 $p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$

¹ The values of $B(p\bar{p})B(\gamma\gamma)$ and $B(\gamma\gamma)B(\gamma J/\psi)$ measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

NODE=M056R25
NODE=M056R25

NODE=M056R25;LINKAGE=AN

$\chi_{c0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M056230

$$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{56}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT

22.5±0.9 OUR FIT

23.7±1.0 OUR AVERAGE

23.7±0.8±0.9 1222 ABLIKIM 13v BES3 $\psi(2S) \rightarrow \gamma p\bar{p}$

23.7±1.4±1.4 383 ± 22 ¹ NAIK 08 CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$

23.6^{+3.7}_{-3.4}±3.4 89.5⁺¹⁴₋₁₃ BAI 04F BES $\psi(2S) \rightarrow \gamma\chi_{c0}(1P) \rightarrow \gamma p\bar{p}$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow p\bar{p}) = (25.7 \pm 1.5 \pm 1.5 \pm 1.3) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056B6
NODE=M056B6

NODE=M056B6;LINKAGE=NA

$$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{56}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

6.53±0.27 OUR FIT

4.6 ±1.9 ¹ BAI 98i BES $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma p\bar{p}$

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow p\bar{p})$ reported in BAI 98i is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B1
NODE=M056B1

NODE=M056B;LINKAGE=B1

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{69}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT

32.0±2.3 OUR FIT

31.7±2.3 OUR AVERAGE

32.0±1.9±2.2 369 ¹ ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

31.2±3.3±2.0 131 ± 12 ² NAIK 08 CLEO $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ Calculated by us. ABLIKIM 13H reports $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.3 \pm 2.0 \pm 2.6) \times 10^{-5}$ from a measurement of $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c0})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.62 \pm 0.31)\%$.

² Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.8 \pm 3.6 \pm 2.2 \pm 1.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056B20
NODE=M056B20

NODE=M056B20;LINKAGE=AB

NODE=M056B20;LINKAGE=NA

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{69}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

9.3±0.7 OUR FIT

13.0^{+3.6}_{-3.5}±2.5 15.2^{+4.2}_{-4.0} ¹ BAI 03E BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

NODE=M056B21
NODE=M056B21

¹ BAI 03E reports [$B(\chi_{c0} \rightarrow \Lambda \bar{\Lambda}) B(\psi(2S) \rightarrow \gamma \chi_{c0}) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p \bar{p})] = (2.45^{+0.68}_{-0.65} \pm 0.46)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p \bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

NODE=M056B21;LINKAGE=BA

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \times \Gamma_{85} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.127±0.006 OUR FIT[(0.126 ± 0.006) × 10⁻² OUR 2015 FIT]**0.131±0.035 OUR AVERAGE** Error includes scale factor of 3.9.

0.151±0.003±0.010	4.3k	ABLIKIM	120	BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.069±0.018		¹ OREGLIA	82	CBAL	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.4 ± 0.3		² BRANDELIK	79B	DASP	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.16 ± 0.11		² BARTEL	78B	CNTR	$\psi(2S) \rightarrow \gamma \chi_{c0}$
3.3 ± 1.7		³ BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.125±0.007±0.013	560	⁴ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.18 ± 0.01 ± 0.02	172	⁵ ADAM	05A	CLEO	Repl. by MENDEZ 08

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.³ Assumes isotropic gamma distribution.⁴ Not independent from other measurements of MENDEZ 08.⁵ Not independent from other values reported by ADAM 05A.NODE=M056B2
NODE=M056B2

NEW

NODE=M056B;LINKAGE=3Q
NODE=M056B;LINKAGE=2Q
NODE=M056B;LINKAGE=EA
NODE=M056B2;LINKAGE=ME
NODE=M056B2;LINKAGE=AD

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything}) \times \Gamma_{85} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

$$\Gamma_{85} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{\psi(2S)} = \Gamma_{85} / \Gamma \times \Gamma_{132}^{\psi(2S)} / (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + 0.339 \Gamma_{133}^{\psi(2S)} + 0.192 \Gamma_{134}^{\psi(2S)})$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.207±0.011 OUR FIT[(0.208 ± 0.011) × 10⁻² OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.201±0.011±0.021	560	¹ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.31 ± 0.02 ± 0.03	172	ADAM	05A	CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

NODE=M056B7

NODE=M056B7

NODE=M056B7

NEW

NODE=M056B7;LINKAGE=ME

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma_{85} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.367±0.019 OUR FIT**0.358±0.020±0.037** 560 MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma \chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.55 ± 0.04 ± 0.06	172	¹ ADAM	05A	CLEO	Repl. by MENDEZ 08
--------------------	-----	-------------------	-----	------	--------------------

¹ Not independent from other values reported by ADAM 05A.NODE=M056B8
NODE=M056B8

NODE=M056B;LINKAGE=AD

$$\Gamma(\chi_{c0}(1P) \Rightarrow \gamma \gamma) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \Rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \times \Gamma_{89} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.23±0.14 OUR FIT**2.18±0.18 OUR AVERAGE**

2.17±0.17±0.12	0.8k	ABLIKIM	12A	BES3	$\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$
2.17±0.32±0.10	0.2k	ECKLUND	08A	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$
3.7 ± 1.8 ± 1.0		LEE	85	CBAL	$\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056B3
NODE=M056B3

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{32}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.32±0.29 OUR FIT**8.80±0.34 OUR AVERAGE**

9.11±0.08±0.65	17k	1 ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
8.81±0.11±0.43	8.9k	2 ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
8.13±0.19±0.89	2.8k	3 ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

NODE=M056B22
NODE=M056B22

OCCUR=2

NODE=M056B22;LINKAGE=AB

NODE=M056B22;LINKAGE=AS

NODE=M056B22;LINKAGE=AN

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (3.23 \pm 0.03 \pm 0.23 \pm 0.14) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^+\pi^-) = (6.37 \pm 0.08 \pm 0.31 \pm 0.32) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

³ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (2.94 \pm 0.07 \pm 0.32 \pm 0.15) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{32}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

24.1±0.8 OUR FIT**20.7±1.7 OUR AVERAGE**

23.9±2.7±4.1	97 ± 11	1 BAI	03C BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^0\pi^0$
20.2±1.1±1.5	720 ± 32	2 BAI	98i BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^+\pi^-$

¹ We have multiplied $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. The value for $B(\chi_{c0} \rightarrow \pi^+\pi^-)$ reported in BAI 98i is derived using $B(\psi' \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi' \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

NODE=M056B5
NODE=M056B5

NODE=M056B;LINKAGE=D1

NODE=M056B;LINKAGE=D2

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{36}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.95±0.18 OUR FIT**3.12±0.19 OUR AVERAGE**

3.23±0.09±0.23	2132	1 ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
2.93±0.12±0.29	0.9k	2 ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.86±0.46±0.37	48	3 ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
----------------	----	---------	---------	--

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c0} \rightarrow \eta\eta) = (3.44 \pm 0.10 \pm 0.24 \pm 0.13) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$.

² Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \eta\eta) = (3.18 \pm 0.13 \pm 0.31 \pm 0.16) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

³ Superseded by ASNER 09. Calculated by us. The value of $B(\chi_{c0}(1P) \rightarrow \eta\eta)$ reported by ADAMS 07 was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46)\%$ (ATHAR 04).

NODE=M056B11
NODE=M056B11

NODE=M056B11;LINKAGE=AB

NODE=M056B11;LINKAGE=AS

NODE=M056B11;LINKAGE=AD

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{36}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.85 ±0.05 OUR FIT**0.578±0.241±0.158**

BAI	03C BES	$\psi(2S) \rightarrow \gamma\eta\eta$
-----	---------	---------------------------------------

NODE=M056B10
NODE=M056B10

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{42}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.91±0.28 OUR FIT[(5.90 ± 0.28) × 10⁻⁴ OUR 2015 FIT]

5.97±0.07±0.32	8.1k	1 ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma K^+K^-$
----------------	------	---------	---------	--------------------------------------

¹ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow K^+K^-) = (6.47 \pm 0.08 \pm 0.35 \pm 0.32) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056B23
NODE=M056B23

NEW

NODE=M056B23;LINKAGE=AS

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{42} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

1.71±0.08 OUR FIT

1.63±0.10±0.15	774 ± 38	¹ BAI	98I BES	$\psi(2S) \rightarrow \gamma K^+ K^-$
----------------	----------	------------------	---------	---------------------------------------

NODE=M056B9
NODE=M056B9

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow K^+ K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B9;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_{43} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

3.09±0.16 OUR FIT**3.18±0.17 OUR AVERAGE**

3.22±0.07±0.17	2.1k	¹ ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
----------------	------	--------------------	---------	---

3.02±0.19±0.33	322	ABLIKIM	05O BES2	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
----------------	-----	---------	----------	---

¹ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow K_S^0 K_S^0) = (3.49 \pm 0.08 \pm 0.18 \pm 0.17) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056B12
NODE=M056B12

NODE=M056B12;LINKAGE=AS

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{43} / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

9.0±0.5 OUR FIT**5.6±0.8±1.3**

¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
------------------	---------	---

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B13
NODE=M056B13

NODE=M056B13;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_1 / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

6.5±0.5 OUR FIT**6.9±2.4 OUR AVERAGE** Error includes scale factor of 3.8.

4.4±0.1±0.9	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
-------------	------------------	---------	---

9.3±0.9	² TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$
---------	------------------------	---------	---

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

² The value $B(\psi(1S) \rightarrow \gamma \chi_{c0}) \times B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

NODE=M056B4
NODE=M056B4

NODE=M056B;LINKAGE=B2

NODE=M056B;LINKAGE=J1

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_8 / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

1.75±0.14 OUR FIT**1.64±0.05±0.2**

ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
---------	----------	---

NODE=M056B18
NODE=M056B18

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_8 / \Gamma \times \Gamma_{132}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

5.1 ±0.4 OUR FIT**5.8 ±1.6 OUR AVERAGE** Error includes scale factor of 2.3.

4.22±0.20±0.97	BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
----------------	-----	---------	---

7.4 ±1.0	¹ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$
----------	------------------------	---------	---

¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

NODE=M056B19
NODE=M056B19

NODE=M056B19;LINKAGE=TA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{50}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{\psi(2S)}}{\Gamma_{132}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
2.74±0.28 OUR FIT				
3.20±0.11±0.41	278	1 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M056B14
NODE=M056B14

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

NODE=M056B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{50}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
8.0±0.8 OUR FIT			
6.1±0.8±0.9	1 BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M056B15
NODE=M056B15

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi \phi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{55}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{\psi(2S)}}{\Gamma_{132}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
0.77±0.07 OUR FIT				
0.78±0.08 OUR AVERAGE				
0.77±0.03±0.08	612	1 ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
0.86±0.19±0.12	26	2 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M056B16
NODE=M056B16

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi \phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31)\%$.

NODE=M056B16;LINKAGE=AL

² Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi \phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

NODE=M056B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi \phi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{55}/\Gamma \times \Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma_{132}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
2.24±0.21 OUR FIT			
2.6 ±1.0 ±1.1	1 BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M056B17
NODE=M056B17

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi \phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B17;LINKAGE=BA

$\chi_{c0}(1P)$ REFERENCES

ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56778
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56779
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55583
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54266
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54742
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53647
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16719
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53940
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53347
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126

NODE=M056

ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
ANDREOTTI	05C	PR D72 112002	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50991
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ABLIKIM	04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50187
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ANDREOTTI	04	PL B584 16	M. Andreotti <i>et al.</i>	(E835 Collab.)	REFID=49744
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
ANDREOTTI	03	PRL 91 091801	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=49578
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
BAGNASCO	02	PL B533 237	S. Bagnasco <i>et al.</i>	(FNAL E835 Collab.)	REFID=48833
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
AMBROGIANI	99B	PRL 83 2902	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47389
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059

$\chi_{c1}(1P)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the Review on " $\psi(2S)$ and χ_c branching ratios" before the $\chi_{c0}(1P)$ Listings.

$\chi_{c1}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3510.66 ± 0.07	OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
3510.30 ± 0.14 ± 0.16		ABLIKIM 05G	BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3510.719 ± 0.051 ± 0.019		ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
3509.4 ± 0.9		BAI 99B	BES	$\psi(2S) \rightarrow \gamma X$
3510.60 ± 0.087 ± 0.019	513	¹ ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
3511.3 ± 0.4 ± 0.4	30	BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+ e^- X$
3512.3 ± 0.3 ± 4.0		² GAISER 86	CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 ± 1.7	91	³ LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- \text{A}$
3510.4 ± 0.6		OREGLIA 82	CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
3510.1 ± 1.1	254	⁴ HIMEL 80	MRK2	$e^+ e^- \rightarrow J/\psi 2\gamma$
3509 ± 11	21	BRANDELIK 79B	DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
3507 ± 3		⁴ BARTEL 78B	CNTR	$e^+ e^- \rightarrow J/\psi 2\gamma$
3505.0 ± 4 ± 4		^{4,5} TANENBAUM 78	MRK1	$e^+ e^-$
3513 ± 7	367	⁴ BIDDICK 77	CNTR	$\psi(2S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3500 ± 10 40 TANENBAUM 75 MRK1 Hadrons γ

¹ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

² Using mass of $\psi(2S) = 3686.0$ MeV.

³ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁴ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁵ From a simultaneous fit to radiative and hadronic decay channels.

NODE=M055

NODE=M055

NODE=M055M

NODE=M055M

OCCUR=2


NODE=M055M;LINKAGE=NW

NODE=M055M;LINKAGE=C

NODE=M055M;LINKAGE=P

NODE=M055M;LINKAGE=D

NODE=M055M;LINKAGE=M



chi_c1(1P) WIDTH

NODE=M055W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.84 ±0.04					OUR FIT
0.88 ±0.05					OUR AVERAGE
1.39 +0.40 +0.26 -0.38 -0.77			ABLIKIM 05G BES2		$\psi(2S) \rightarrow \gamma \chi_{c1}$
0.876±0.045±0.026			ANDREOTTI 05A E835		$p\bar{p} \rightarrow e^+ e^- \gamma$
0.87 ±0.11 ±0.08		513	¹ ARMSTRONG 92 E760		$\bar{p}p \rightarrow e^+ e^- \gamma$
<1.3	95		BAGLIN 86B SPEC		$\bar{p}p \rightarrow e^+ e^- X$
<3.8	90		GAISER 86 CBAL		$\psi(2S) \rightarrow \gamma X$

NODE=M055W

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Recalculated by ANDREOTTI 05A.

NODE=M055W;LINKAGE=AN

chi_c1(1P) DECAY MODES

NODE=M055215;NODE=M055

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic decays		
Γ_1 $3(\pi^+ \pi^-)$	$(5.8 \pm 1.4) \times 10^{-3}$	S=1.2
Γ_2 $2(\pi^+ \pi^-)$	$(7.6 \pm 2.6) \times 10^{-3}$	
Γ_3 $\pi^+ \pi^- \pi^0 \pi^0$	$(1.22 \pm 0.16) \%$	
Γ_4 $\rho^+ \pi^- \pi^0 + c.c.$	$(1.48 \pm 0.25) \%$	
Γ_5 $\rho^0 \pi^+ \pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$	
Γ_6 $4\pi^0$	$(5.5 \pm 0.8) \times 10^{-4}$	
Γ_7 $\pi^+ \pi^- K^+ K^-$	$(4.5 \pm 1.0) \times 10^{-3}$	
Γ_8 $K^+ K^- \pi^0 \pi^0$	$(1.14 \pm 0.28) \times 10^{-3}$	
Γ_9 $K^+ K^- \pi^+ \pi^- \pi^0$	$(1.15 \pm 0.13) \%$	
Γ_{10} $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(7.5 \pm 0.8) \times 10^{-3}$	
Γ_{11} $K^+ \pi^- \bar{K}^0 \pi^0 + c.c.$	$(8.7 \pm 1.4) \times 10^{-3}$	
Γ_{12} $\rho^- K^+ K^0 + c.c.$	$(5.1 \pm 1.2) \times 10^{-3}$	
Γ_{13} $K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + c.c.$	$(2.4 \pm 0.7) \times 10^{-3}$	
Γ_{14} $K^+ K^- \eta \pi^0$	$(1.14 \pm 0.35) \times 10^{-3}$	
Γ_{15} $\pi^+ \pi^- K_S^0 K_S^0$	$(7.0 \pm 3.0) \times 10^{-4}$	
Γ_{16} $K^+ K^- \eta$	$(3.2 \pm 1.0) \times 10^{-4}$	
Γ_{17} $\bar{K}^0 K^+ \pi^- + c.c.$	$(7.1 \pm 0.6) \times 10^{-3}$	
Γ_{18} $K^*(892)^0 \bar{K}^0 + c.c.$	$(1.0 \pm 0.4) \times 10^{-3}$	
Γ_{19} $K^*(892)^+ K^- + c.c.$	$(1.5 \pm 0.7) \times 10^{-3}$	

NODE=M055;CLUMP=A

DESIG=6

DESIG=5

DESIG=51

DESIG=52

DESIG=9

DESIG=60

DESIG=7

DESIG=53

DESIG=79

DESIG=84

DESIG=55

DESIG=56

DESIG=57

DESIG=58

DESIG=28

DESIG=42

DESIG=17

DESIG=32

DESIG=33

Г20	$K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$	$< 8 \times 10^{-4}$	CL=90%	DESIG=34
Г21	$K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$	$< 2.2 \times 10^{-3}$	CL=90%	DESIG=35
Г22	$K^+ K^- \pi^0$	$(1.85 \pm 0.25) \times 10^{-3}$		DESIG=38
Г23	$\eta \pi^+ \pi^-$	$(4.9 \pm 0.5) \times 10^{-3}$		DESIG=31
Г24	$a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-$	$(1.8 \pm 0.6) \times 10^{-3}$		DESIG=36
Г25	$f_2(1270) \eta$	$(2.7 \pm 0.8) \times 10^{-3}$		DESIG=37
Г26	$\pi^+ \pi^- \eta'$	$(2.3 \pm 0.5) \times 10^{-3}$		DESIG=44
Г27	$K^+ K^- \eta'(958)$	$(8.8 \pm 0.9) \times 10^{-4}$		DESIG=85
Г28	$K_0^*(1430)^+ K^- + \text{c.c.}$	$(6.4 \pm_{-2.8}^{+2.2}) \times 10^{-4}$		DESIG=86
Г29	$f_0(980) \eta'(958)$	$(1.6 \pm_{-0.7}^{+1.4}) \times 10^{-4}$		DESIG=87
Г30	$f_0(1710) \eta'(958)$	$(7 \pm_{-5}^{+7}) \times 10^{-5}$		DESIG=88
Г31	$f_2'(1525) \eta'(958)$	$(9 \pm 6) \times 10^{-5}$		DESIG=89
Г32	$\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-$	$< 6 \times 10^{-6}$	CL=90%	DESIG=61
Г33	$K^+ K^*(892)^0 \pi^- + \text{c.c.}$	$(3.2 \pm 2.1) \times 10^{-3}$		DESIG=10
Г34	$K^*(892)^0 \bar{K}^*(892)^0$	$(1.5 \pm 0.4) \times 10^{-3}$		DESIG=21
Г35	$K^+ K^- K_S^0 K_S^0$	$< 4 \times 10^{-4}$	CL=90%	DESIG=29
Г36	$K^+ K^- K^+ K^-$	$(5.5 \pm 1.1) \times 10^{-4}$		DESIG=14
Г37	$K^+ K^- \phi$	$(4.2 \pm 1.6) \times 10^{-4}$		DESIG=30
Г38	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(3.3 \pm 0.5) \times 10^{-3}$		DESIG=90
Г39	$K^+ K^- \pi^0 \phi$	$(1.62 \pm 0.30) \times 10^{-3}$		DESIG=91
Г40	$\phi \pi^+ \pi^- \pi^0$	$(7.5 \pm 1.0) \times 10^{-4}$		DESIG=82
Г41	$\omega \omega$	$(5.8 \pm 0.7) \times 10^{-4}$		DESIG=66
Г42	$\omega K^+ K^-$	$(7.8 \pm 0.9) \times 10^{-4}$		DESIG=81
Г43	$\omega \phi$	$(2.1 \pm 0.6) \times 10^{-5}$		DESIG=67
Г44	$\phi \phi$	$(4.2 \pm 0.5) \times 10^{-4}$		DESIG=68
Г45	$\rho \bar{\rho}$	$(7.72 \pm 0.35) \times 10^{-5}$		DESIG=11
Г46	$\rho \bar{\rho} \pi^0$	$(1.59 \pm 0.19) \times 10^{-4}$		DESIG=39
Г47	$\rho \bar{\rho} \eta$	$(1.48 \pm 0.25) \times 10^{-4}$		DESIG=43
Г48	$\rho \bar{\rho} \omega$	$(2.16 \pm 0.31) \times 10^{-4}$		DESIG=59
Г49	$\rho \bar{\rho} \phi$	$< 1.8 \times 10^{-5}$	CL=90%	DESIG=65
Г50	$\rho \bar{\rho} \pi^+ \pi^-$	$(5.0 \pm 1.9) \times 10^{-4}$		DESIG=8
Г51	$\rho \bar{\rho} \pi^0 \pi^0$			DESIG=54
Г52	$\rho \bar{\rho} K^+ K^-$ (non-resonant)	$(1.30 \pm 0.23) \times 10^{-4}$		DESIG=62
Г53	$\rho \bar{\rho} K_S^0 K_S^0$	$< 4.5 \times 10^{-4}$	CL=90%	DESIG=25
Г54	$\rho \bar{n} \pi^-$	$(3.9 \pm 0.5) \times 10^{-4}$		DESIG=74
Г55	$\bar{\rho} n \pi^+$	$(4.0 \pm 0.5) \times 10^{-4}$		DESIG=75
Г56	$\rho \bar{n} \pi^- \pi^0$	$(1.05 \pm 0.12) \times 10^{-3}$		DESIG=76
Г57	$\bar{\rho} n \pi^+ \pi^0$	$(1.03 \pm 0.12) \times 10^{-3}$		DESIG=77
Г58	$\Lambda \bar{\Lambda}$	$(1.16 \pm 0.12) \times 10^{-4}$		DESIG=19
Г59	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$		DESIG=24
Г60	$\Lambda \bar{\Lambda} \pi^+ \pi^-$ (non-resonant)	$(2.5 \pm 0.6) \times 10^{-4}$		DESIG=69
Г61	$\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=70
Г62	$\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=71
Г63	$K^+ \bar{p} \Lambda$	$(4.2 \pm 0.4) \times 10^{-4}$	S=1.1	DESIG=40
Г64	$K^+ \bar{p} \Lambda(1520) + \text{c.c.}$	$(1.7 \pm 0.5) \times 10^{-4}$		DESIG=63
Г65	$\Lambda(1520) \bar{\Lambda}(1520)$	$< 1.0 \times 10^{-4}$	CL=90%	DESIG=64
Г66	$\Sigma^0 \bar{\Sigma}^0$	$< 4 \times 10^{-5}$	CL=90%	DESIG=48
Г67	$\Sigma^+ \bar{\Sigma}^-$	$< 6 \times 10^{-5}$	CL=90%	DESIG=49
Г68	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$< 1.0 \times 10^{-4}$	CL=90%	DESIG=72
Г69	$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$< 5 \times 10^{-5}$	CL=90%	DESIG=73
Г70	$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(1.38 \pm 0.25) \times 10^{-4}$		DESIG=92
Г71	$\Xi^0 \bar{\Xi}^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=50
Г72	$\Xi^- \bar{\Xi}^+$	$(8.2 \pm 2.2) \times 10^{-5}$		DESIG=26
Г73	$\pi^+ \pi^- + K^+ K^-$	$< 2.1 \times 10^{-3}$		DESIG=23
Г74	$K_S^0 K_S^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=27
Г75	$\eta_c \pi^+ \pi^-$	$< 3.2 \times 10^{-3}$	CL=90%	DESIG=83

Radiative decays

Γ_{76}	$\gamma J/\psi(1S)$	$(33.9 \pm 1.2) \%$
Γ_{77}	$\gamma \rho^0$	$(2.20 \pm 0.18) \times 10^{-4}$
Γ_{78}	$\gamma \omega$	$(6.9 \pm 0.8) \times 10^{-5}$
Γ_{79}	$\gamma \phi$	$(2.5 \pm 0.5) \times 10^{-5}$
Γ_{80}	$\gamma \gamma$	

NODE=M055;CLUMP=B
DESIG=1
DESIG=45
DESIG=46
DESIG=47
DESIG=4

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 240 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 342.4$ for 191 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_{36}	6				
x_{45}	8	3			
x_{58}	13	5	7		
x_{76}	31	13	6	26	
Γ	-19	-8	-62	-16	-51
	x_{17}	x_{36}	x_{45}	x_{58}	x_{76}

 $\chi_{c1}(1P)$ PARTIAL WIDTHS

$$\chi_{c1}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$$

NODE=M055220

NODE=M055223

$$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}} \quad \Gamma_{45} \Gamma_{76} / \Gamma$$

NODE=M055G1
NODE=M055G1

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
21.9 ± 0.8 OUR FIT			
21.4 ± 0.9 OUR AVERAGE			
21.5 ± 0.5 ± 0.8	¹ ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
21.4 ± 1.5 ± 2.2	^{1,2} ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
19.9 ^{+4.4} _{-4.0}	¹ BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+ e^- X$

¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

² Recalculated by ANDREOTTI 05A.

NODE=M055G;LINKAGE=7A
NODE=M055G;LINKAGE=AN

 $\chi_{c1}(1P)$ BRANCHING RATIOS

NODE=M055225

HADRONIC DECAYS

NODE=M055305

$$\Gamma(3(\pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$

NODE=M055R6
NODE=M055R6

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.8 ± 1.4 OUR EVALUATION	Error includes scale factor of 1.2. Treating systematic error as correlated.		
5.8 ± 1.1 OUR AVERAGE			

→ UNCHECKED ←

5.4 ± 0.7 ± 0.9	¹ BAI 99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
16.0 ± 5.9 ± 0.8	¹ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R;LINKAGE=X2

$$\Gamma(2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \quad \Gamma_2 / \Gamma$$

NODE=M055R4
NODE=M055R4

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
7.6 ± 2.6 OUR EVALUATION	Treating systematic error as correlated.		
8 ± 4 OUR AVERAGE	Error includes scale factor of 1.5.		

→ UNCHECKED ←

4.6 ± 2.1 ± 2.6	¹ BAI 99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
12.5 ± 4.2 ± 0.6	¹ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R4;LINKAGE=X2

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.22±0.15±0.04	604.7	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R35
NODE=M055R35

¹ HE 08B reports $1.28 \pm 0.06 \pm 0.15 \pm 0.08$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R35;LINKAGE=HE

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.48±0.24±0.05	712.3	^{1,2} HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R36
NODE=M055R36

¹ HE 08B reports $1.56 \pm 0.13 \pm 0.22 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R36;LINKAGE=HE

² Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M055R36;LINKAGE=OC

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
39±35	¹ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R8
NODE=M055R8

¹ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R;LINKAGE=T

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.55±0.08±0.02	608	¹ ABLIKIM 11A	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R44
NODE=M055R44

¹ ABLIKIM 11A reports $(0.57 \pm 0.03 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R44;LINKAGE=AB

 $\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.5±1.0 OUR EVALUATION	Treating systematic error as correlated.		
4.5±0.9 OUR AVERAGE			

NODE=M055R5
NODE=M055R5
→ UNCHECKED ←

4.2±0.4±0.9	¹ BAI 99B	BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$
7.3±3.0±0.4	¹ TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R5;LINKAGE=X2

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.114±0.028±0.004	45.1	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R37
NODE=M055R37

¹ HE 08B reports $0.12 \pm 0.02 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R37;LINKAGE=HE

 $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
11.46±0.12±1.29	12k	¹ ABLIKIM 13B	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R00
NODE=M055R00

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R00;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^\pm \pi^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.52±0.11±0.79	5.1k	¹ ABLIKIM 13B	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R60
NODE=M055R60

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R60;LINKAGE=A

$\Gamma(K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.87±0.14±0.03	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R39
NODE=M055R39

¹ HE 08B reports $0.92 \pm 0.09 \pm 0.11 \pm 0.06$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R39;LINKAGE=HE

 $\Gamma(\rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.51±0.12±0.02	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R40
NODE=M055R40

¹ HE 08B reports $0.54 \pm 0.11 \pm 0.07 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R40;LINKAGE=HE

 $\Gamma(K^*(892)^0\bar{K}^0\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.24±0.06±0.01	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R41
NODE=M055R41

¹ HE 08B reports $0.25 \pm 0.06 \pm 0.03 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0\bar{K}^0\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R41;LINKAGE=HE

 $\Gamma(K^+K^-\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.114±0.035±0.004	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R42
NODE=M055R42

¹ HE 08B reports $0.12 \pm 0.03 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\eta\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R42;LINKAGE=HE

 $\Gamma(\pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.0±3.0±0.2	19.8 ± 7.7	¹ ABLIKIM	05o BES2	$\psi(2S) \rightarrow \chi_{c1}\gamma$

NODE=M055R05
NODE=M055R05

¹ ABLIKIM 05o reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ $= (0.67 \pm 0.26 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R05;LINKAGE=AB

 $\Gamma(K^+K^-\eta)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.32±0.10±0.01	¹ ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M055R25
NODE=M055R25

¹ ATHAR 07 reports $(0.34 \pm 0.10 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R25;LINKAGE=AT

 $\Gamma(\bar{K}^0K^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-3})	DOCUMENT ID
7.1±0.6 OUR FIT	

NODE=M055R17
NODE=M055R17

 $\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.00±0.37±0.03	22	¹ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R09
NODE=M055R09

¹ ABLIKIM 06R reports $(1.1 \pm 0.4 \pm 0.1) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R09;LINKAGE=AB

$\Gamma(K^*(892)^+ K^- + c.c.)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.46±0.66±0.05	27	1 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R10
NODE=M055R10

¹ ABLIKIM 06R reports $(1.6 \pm 0.7 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R10;LINKAGE=AB

 $\Gamma(K_J^*(1430)^0 \bar{K}^0 + c.c. \rightarrow K_S^0 K^+ \pi^- + c.c.)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	1 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R12
NODE=M055R12

¹ ABLIKIM 06R reports $< 0.9 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^0 \bar{K}^0 + c.c. \rightarrow K_S^0 K^+ \pi^- + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R12;LINKAGE=AB

 $\Gamma(K_J^*(1430)^+ K^- + c.c. \rightarrow K_S^0 K^+ \pi^- + c.c.)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	1 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R13
NODE=M055R13

¹ ABLIKIM 06R reports $< 2.4 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^+ K^- + c.c. \rightarrow K_S^0 K^+ \pi^- + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R13;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.85±0.24±0.06	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M055R20
NODE=M055R20

¹ ATHAR 07 reports $(1.95 \pm 0.16 \pm 0.23) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R20;LINKAGE=AT

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.9±0.5 OUR AVERAGE				
4.7±0.5±0.2		1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
5.4±0.9±0.2	222	2 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R08
NODE=M055R08

¹ ATHAR 07 reports $(5.0 \pm 0.3 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R08;LINKAGE=AT

² ABLIKIM 06R reports $(5.9 \pm 0.7 \pm 0.8) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R08;LINKAGE=AB

 $\Gamma(a_0(980)^+ \pi^- + c.c. \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.8±0.6±0.1	58	1 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R15
NODE=M055R15

¹ ABLIKIM 06R reports $(2.0 \pm 0.5 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+ \pi^- + c.c. \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R15;LINKAGE=AB

$\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.7 ± 0.8 ± 0.1	53	¹ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R16
 NODE=M055R16

¹ ABLIKIM 06R reports $(3.0 \pm 0.7 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R16;LINKAGE=AB

 $\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.5 ± 0.1	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M055R28
 NODE=M055R28

¹ ATHAR 07 reports $(2.4 \pm 0.4 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R28;LINKAGE=AT

 $\Gamma(K^+K^-\eta'(958))/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.75 ± 0.87	310	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^-\eta'(958)$

NODE=M055R64
 NODE=M055R64

¹ Derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.2 \pm 0.4)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.

NODE=M055R64;LINKAGE=A

 $\Gamma(K_0^*(1430)^+K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
6.41 ± 0.57^{+2.09}_{-2.71}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^-\eta'(958)$

NODE=M055R65
 NODE=M055R65

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^-\eta'(958))$ branching fraction.

NODE=M055R65;LINKAGE=A

 $\Gamma(f_0(980)\eta'(958))/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.65 ± 0.47^{+1.32}_{-0.56}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^-\eta'(958)$

NODE=M055R66
 NODE=M055R66

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^-\eta'(958))$ branching fraction.

NODE=M055R66;LINKAGE=A

 $\Gamma(f_0(1710)\eta'(958))/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.71 ± 0.22^{+0.68}_{-0.48}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^-\eta'(958)$

NODE=M055R67
 NODE=M055R67

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^-\eta'(958))$ branching fraction.

NODE=M055R67;LINKAGE=A

 $\Gamma(f_2'(1525)\eta'(958))/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.23^{+0.55}_{-0.51}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^-\eta'(958)$

NODE=M055R68
 NODE=M055R68

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^-\eta'(958))$ branching fraction.

NODE=M055R68;LINKAGE=A

 $\Gamma(\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6 × 10⁻⁶	90	¹ ABLIKIM	11D BES3	$\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

NODE=M055R18
 NODE=M055R18

¹ ABLIKIM 11D reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] < 6.0 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R18;LINKAGE=BR

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
32 ± 21	¹ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R9
 NODE=M055R9

¹ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R9;LINKAGE=T

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.47±0.36±0.05	28.4 ± 5.5	^{1,2} ABLIKIM	04H BES	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
-----------------------	------------	------------------------	---------	---

NODE=M055R26
NODE=M055R26

¹ ABLIKIM 04H reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.40 \pm 0.27 \pm 0.22) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R26;LINKAGE=AB

² Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

NODE=M055R26;LINKAGE=AL

 $\Gamma(K^+ K^- K_S^0 \bar{K}_S^0)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<4	90	3.2 ± 2.4	¹ ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$
--------------	----	-----------	----------------------	----------	---

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K_S^0 \bar{K}_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 4.2 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R06
NODE=M055R06

NODE=M055R06;LINKAGE=AB

 $\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-3})	DOCUMENT ID
--------------------------	-------------

0.55±0.11 OUR FITNODE=M055R14
NODE=M055R14 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.42±0.15±0.01	17	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
-----------------------	----	----------------------	----------	---

¹ ABLIKIM 06T reports $(0.46 \pm 0.16 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R07
NODE=M055R07

NODE=M055R07;LINKAGE=AB

 $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

3.27±0.28±0.46	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
-----------------------	---------	----------	---

NODE=M055R69
NODE=M055R69 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.62±0.12±0.28	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
-----------------------	---------	----------	---

NODE=M055R70
NODE=M055R70 $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.75±0.06±0.08	373	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$
-----------------------	-----	----------------------	----------	---

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R62
NODE=M055R62

NODE=M055R62;LINKAGE=A

 $\Gamma(\omega)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.8±0.7±0.2	597	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
--------------------	-----	----------------------	----------	---------------------------------------

¹ ABLIKIM 11K reports $(6.0 \pm 0.3 \pm 0.7) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R49
NODE=M055R49

NODE=M055R49;LINKAGE=AL

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.78±0.04±0.08	628	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$
-----------------------	-----	----------------------	----------	---

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R61
NODE=M055R61

NODE=M055R61;LINKAGE=A

 $\Gamma(\omega \phi)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.21±0.06±0.01	15	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
-----------------------	----	----------------------	----------	---------------------------------------

¹ ABLIKIM 11K reports $(0.22 \pm 0.06 \pm 0.02) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R50
NODE=M055R50

NODE=M055R50;LINKAGE=AL

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
4.2±0.5±0.1	366	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons

NODE=M055R51
 NODE=M055R51

¹ ABLIKIM 11K reports $(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R51;LINKAGE=AL

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	DOCUMENT ID
0.772±0.035 OUR FIT	

NODE=M055R11
 NODE=M055R11

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.159±0.019 OUR AVERAGE			

NODE=M055R21
 NODE=M055R21

0.166±0.020±0.005

¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$

0.114±0.048±0.004

² ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(1.75 \pm 0.16 \pm 0.13 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R21;LINKAGE=ON

² ATHAR 07 reports $(1.2 \pm 0.5 \pm 0.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R21;LINKAGE=AT

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
0.148±0.025±0.005		¹ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$

NODE=M055R27
 NODE=M055R27

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.15 90 ² ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(1.56 \pm 0.22 \pm 0.14 \pm 0.10) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R27;LINKAGE=ON

² ATHAR 07 reports $< 0.16 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R27;LINKAGE=AT

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.216±0.031±0.007	¹ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$

NODE=M055R43
 NODE=M055R43

¹ ONYISI 10 reports $(2.28 \pm 0.28 \pm 0.16 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R43;LINKAGE=ON

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$

NODE=M055R48
 NODE=M055R48

¹ ABLIKIM 11F reports $< 1.82 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R48;LINKAGE=AB

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{50}/Γ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.50±0.19 OUR EVALUATION Treating systematic error as correlated.NODE=M055R7
NODE=M055R7

→ UNCHECKED ←

0.50±0.19 OUR AVERAGE

0.46±0.12±0.15

¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c1}$

1.08±0.77±0.05

¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$ ¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R7;LINKAGE=X2

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (%)

CL%

DOCUMENT ID TECN COMMENT

<0.05 90 ¹ HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ NODE=M055R38
NODE=M055R38¹ HE 08B reports < 0.05 % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R38;LINKAGE=HE

 $\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{52}/Γ VALUE (units 10^{-4})

EVTS

DOCUMENT ID TECN COMMENT

1.30±0.23±0.04 82 ± 9 ¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$ NODE=M055R45
NODE=M055R45¹ ABLIKIM 11F reports $(1.35 \pm 0.15 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R45;LINKAGE=AB

 $\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{53}/Γ VALUE (units 10^{-4})

CL%

DOCUMENT ID TECN COMMENT

<4.5 90 ¹ ABLIKIM 06D BES2 $\psi(2S) \rightarrow \gamma\chi_{c1}$ NODE=M055R02
NODE=M055R02¹ Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.1 \pm 0.6)\%$.

NODE=M055R;LINKAGE=AB

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{54}/Γ VALUE (units 10^{-4})

EVTS

DOCUMENT ID TECN COMMENT

3.9±0.5±0.1 1412 ¹ ABLIKIM 12J BES3 $\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$ NODE=M055R56
NODE=M055R56¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.37 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R56;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{55}/Γ VALUE (units 10^{-4})

EVTS

DOCUMENT ID TECN COMMENT

4.0±0.5±0.1 1625 ¹ ABLIKIM 12J BES3 $\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$ NODE=M055R57
NODE=M055R57¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.38 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R57;LINKAGE=AL

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{56}/Γ VALUE (units 10^{-4})

EVTS

DOCUMENT ID TECN COMMENT

10.5±1.2±0.3 1082 ¹ ABLIKIM 12J BES3 $\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$ NODE=M055R58
NODE=M055R58¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (1.00 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R58;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{57}/Γ VALUE (units 10^{-4})

EVTS

DOCUMENT ID TECN COMMENT

10.3±1.2±0.3 1261 ¹ ABLIKIM 12J BES3 $\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$ NODE=M055R59
NODE=M055R59¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.98 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R59;LINKAGE=AL

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

DOCUMENT ID

1.16±0.12 OUR FIT Γ_{58}/Γ NODE=M055R23
NODE=M055R23 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-5}) CL% EVTS

DOCUMENT ID

TECN

COMMENT

30±5±1

105

¹ ABLIKIM 12I BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<150 90

² ABLIKIM 06D BES2 $\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ ABLIKIM 12I reports $(31.1 \pm 3.4 \pm 3.9) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.1 \pm 0.6)\%$.

 Γ_{59}/Γ NODE=M055R01
NODE=M055R01 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

EVTS

DOCUMENT ID

TECN

COMMENT

25±6±1

13

¹ ABLIKIM 12I BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

¹ ABLIKIM 12I reports $(26.2 \pm 5.5 \pm 3.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{60}/Γ NODE=M055R19
NODE=M055R19 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

<13

90

¹ ABLIKIM 12I BES3 $\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

 Γ_{61}/Γ NODE=M055R52
NODE=M055R52 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

<13

90

¹ ABLIKIM 12I BES3 $\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$

¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

 Γ_{62}/Γ NODE=M055R53
NODE=M055R53 $\Gamma(K^+\bar{p}\Lambda)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

EVTS

DOCUMENT ID

TECN

COMMENT

4.2±0.4 OUR AVERAGE Error includes scale factor of 1.1.

4.3±0.4±0.1

3k

^{1,2} ABLIKIM 13D BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$

3.1±0.9±0.1

³ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ABLIKIM 13D reports $(4.5 \pm 0.2 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.

³ ATHAR 07 reports $(3.3 \pm 0.9 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{63}/Γ NODE=M055R22
NODE=M055R22

NODE=M055R22;LINKAGE=AB

NODE=M055R22;LINKAGE=LB
NODE=M055R22;LINKAGE=AT

$\Gamma(K^+ \bar{p} \Lambda(1520) + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{64} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 ± 0.1	48 ± 10	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$
¹ ABLIKIM 11F reports $(1.81 \pm 0.38 \pm 0.28) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M055R46
 NODE=M055R46

NODE=M055R46;LINKAGE=AB

 $\Gamma(\Lambda(1520) \bar{\Lambda}(1520)) / \Gamma_{\text{total}}$ Γ_{65} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$
¹ ABLIKIM 11F reports $< 1.00 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.				

NODE=M055R47
 NODE=M055R47

NODE=M055R47;LINKAGE=AB

 $\Gamma(\Sigma^0 \bar{\Sigma}^0) / \Gamma_{\text{total}}$ Γ_{66} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.4	90	3.8 ± 2.5	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.6	90		² ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
¹ NAIK 08 reports $< 0.44 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.					
² ABLIKIM 13H reports $< 0.62 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.					

NODE=M055R32
 NODE=M055R32

NODE=M055R32;LINKAGE=NA

NODE=M055R32;LINKAGE=AB

 $\Gamma(\Sigma^+ \bar{\Sigma}^-) / \Gamma_{\text{total}}$ Γ_{67} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.6	90	4.3 ± 2.3	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.8	90		² ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
¹ NAIK 08 reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.					
² ABLIKIM 13H reports $< 0.87 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.					

NODE=M055R33
 NODE=M055R33

NODE=M055R33;LINKAGE=NA

NODE=M055R33;LINKAGE=AB

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-) / \Gamma_{\text{total}}$ Γ_{68} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<10	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
¹ ABLIKIM 12I reports $< 10 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.				

NODE=M055R54
 NODE=M055R54

NODE=M055R54;LINKAGE=AL

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+) / \Gamma_{\text{total}}$ Γ_{69} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
¹ ABLIKIM 12I reports $< 5.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.				

NODE=M055R55
 NODE=M055R55

NODE=M055R55;LINKAGE=AL

$\Gamma(K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.38±0.24±0.05	49	¹ ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \Xi^+ + \text{c.c.}$

NODE=M055R71
NODE=M055R71

¹ ABLIKIM 15I reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.32 \pm 0.20 \pm 0.12) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R71;LINKAGE=A

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.6	90	1.7 ± 2.4	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$

NODE=M055R34
NODE=M055R34

¹ NAIK 08 reports $< 0.60 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R34;LINKAGE=NA

 $\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.82±0.22±0.03	16.4 ± 4.3		¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^+ \Xi^-$

NODE=M055R03
NODE=M055R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.4	90		² ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
-------	----	--	----------------------	----------	---

¹ NAIK 08 reports $(0.86 \pm 0.22 \pm 0.08) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R03;LINKAGE=NA

² Using $B(\psi(2S) \rightarrow \chi_{c1} \gamma) (9.1 \pm 0.6)\%$.

NODE=M055R03;LINKAGE=AB

 $[\Gamma(\pi^+ \pi^-) + \Gamma(K^+ K^-)]/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<21		¹ FELDMAN	77 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R2
NODE=M055R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<38	90	¹ BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c1}$
-----	----	------------------------	----------	---

¹ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R2;LINKAGE=T

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.6	90	¹ ABLIKIM	05O BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

NODE=M055R04
NODE=M055R04

¹ ABLIKIM 05O reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $< 0.6 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R04;LINKAGE=AB

 $\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻³	90	^{1,2} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R63
NODE=M055R63

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.4 × 10 ⁻³	90	^{1,3} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$
-------------------------	----	------------------------	----------	---

OCCUR=2

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R63;LINKAGE=A

² Using the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

NODE=M055R63;LINKAGE=B

³ Using the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

NODE=M055R63;LINKAGE=C

————— RADIATIVE DECAYS —————

NODE=M055310

 $\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.339±0.012 OUR FIT			

NODE=M055R1
NODE=M055R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.379±0.008±0.021	¹ ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$
-------------------	-------------------	----------	---

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c1} \rightarrow \gamma \gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma \chi_{c1})$ from ATHAR 04.

NODE=M055R1;LINKAGE=AD

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

220 ± 18 OUR AVERAGE

220 ± 23 ± 7	432 ± 25	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
221 ± 24 ± 7	186 ± 15	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$

¹ ABLIKIM 11E reports $(228 \pm 13 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BENNETT 08A reports $(243 \pm 19 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R29
NODE=M055R29

NODE=M055R29;LINKAGE=AB

NODE=M055R29;LINKAGE=BE

 $\Gamma(\gamma\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

69 ± 8 OUR AVERAGE

67 ± 9 ± 2	136 ± 14	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
76 ± 17 ± 2	39 ± 7	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$

¹ ABLIKIM 11E reports $(69.7 \pm 7.2 \pm 6.6) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BENNETT 08A reports $(83 \pm 15 \pm 12) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R30
NODE=M055R30

NODE=M055R30;LINKAGE=AB

NODE=M055R30;LINKAGE=BE

 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

25 ± 5 ± 1		43 ± 9	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
-------------------	--	--------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<24	90	5.2 ± 3.1	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
-----	----	-----------	----------------------	----------	---

¹ ABLIKIM 11E reports $(25.8 \pm 5.2 \pm 2.3) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BENNETT 08A reports $< 26 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.55 \times 10^{-2}$.

NODE=M055R31
NODE=M055R31

NODE=M055R31;LINKAGE=AB

NODE=M055R31;LINKAGE=BE

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.5	90	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$
<150	90	¹ YAMADA	77 DASP	$e^+e^- \rightarrow 3\gamma$

¹ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R3
NODE=M055R3

NODE=M055R;LINKAGE=T1

 $\chi_{c1}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow$

$J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{45}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

2.14 ± 0.11 OUR FIT

1.1 ± 1.0	¹ BAI	98I BES	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma p\bar{p}$
------------------	------------------	---------	--

¹ Calculated by us. The value for $B(\chi_{c1} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M055230

NODE=M055B1
NODE=M055B1

NODE=M055B;LINKAGE=J2

$$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{58}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
11.1±1.1 OUR FIT				
10.9±1.1 OUR AVERAGE				

11.2±1.0±0.9	136	¹ ABLIKIM	13H	BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
10.5±1.6±0.6	46 ± 7	² NAIK	08	CLEO $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ Calculated by us. ABLIKIM 13H reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (12.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ from a measurement of $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c1})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.2 \pm 0.4)\%$.

² Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (11.6 \pm 1.8 \pm 0.7 \pm 0.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

NODE=M055B10
NODE=M055B10

NODE=M055B10;LINKAGE=AB

NODE=M055B10;LINKAGE=NA

$$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{58}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.22±0.31 OUR FIT				

7.1 $^{+2.8}_{-2.4}$ ±1.3	9.0 $^{+3.5}_{-3.1}$	¹ BAI	03E	BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
---------------------------	----------------------	------------------	-----	---

¹ BAI 03E reports $[B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c1}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^-p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.52}_{-0.46} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^-p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

NODE=M055B11
NODE=M055B11

NODE=M055B11;LINKAGE=BA

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{76}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.24 ±0.07 OUR FIT				
2.93 ±0.15 OUR AVERAGE				

3.377±0.009±0.183	142k	ABLIKIM	120	BES3 $\psi(2S) \rightarrow \gamma\chi_{c1}$
2.81 ±0.05 ±0.23	13k	BAI	04I	BES2 $\psi(2S) \rightarrow J/\psi\gamma\gamma$
2.56 ±0.12 ±0.20		GAISER	86	CBAL $\psi(2S) \rightarrow \gamma X$
2.78 ±0.30		¹ OREGLIA	82	CBAL $\psi(2S) \rightarrow \gamma\chi_{c1}$
2.2 ±0.5		² BRANDELIK	79B	DASP $\psi(2S) \rightarrow \gamma\chi_{c1}$
2.9 ±0.5		² BARTEL	78B	CNTR $\psi(2S) \rightarrow \gamma\chi_{c1}$
5.0 ±1.5		³ BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$
2.8 ±0.9		¹ WHITAKER	76	MRK1 e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.56 ±0.03 ±0.12	24.9k	⁴ MENDEZ	08	CLEO $\psi(2S) \rightarrow \gamma\chi_{c1}$
3.44 ±0.06 ±0.13	3.7k	⁵ ADAM	05A	CLEO Repl. by MENDEZ 08

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.


³ Assumes isotropic gamma distribution.

⁴ Not independent from other measurements of MENDEZ 08.

⁵ Not independent from other values reported by ADAM 05A.

NODE=M055B2
NODE=M055B2

NODE=M055B;LINKAGE=3Q
NODE=M055B;LINKAGE=2Q
NODE=M055B;LINKAGE=EA
NODE=M055B2;LINKAGE=ME
NODE=M055B;LINKAGE=AD



$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}} \text{ (units } 10^{-2})$$

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{76} / \Gamma \times \Gamma_{133}^{\psi(2S)} / \Gamma_9^{\psi(2S)} = \Gamma_{76} / \Gamma \times \Gamma_{133}^{\psi(2S)} / (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + \Gamma_{134}^{\psi(2S)})$$

$$0.339 \Gamma_{133}^{\psi(2S)} + 0.192 \Gamma_{134}^{\psi(2S)}$$

NODE=M055B7
NODE=M055B7

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

5.31±0.11 OUR FIT
[(5.32 ± 0.11) × 10^{-2} OUR 2015 FIT]

NODE=M055B7
NEW

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 5.70 ± 0.04 ± 0.15 24.9k ¹ MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma \chi_{c1}$
- 5.77 ± 0.10 ± 0.12 3.7k ADAM 05A CLEO Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

NODE=M055B7;LINKAGE=ME

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{76} / \Gamma \times \Gamma_{133}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

NODE=M055B3
NODE=M055B3

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

9.40±0.21 OUR FIT
10.15±0.28 OUR AVERAGE

- 10.17 ± 0.07 ± 0.27 24.9k MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma \chi_{c1}$
- 12.6 ± 0.3 ± 3.8 3k ¹ ABLIKIM 04B BES $\psi(2S) \rightarrow J/\psi X$
- 8.5 ± 2.1 ² HIMEL 80 MRK2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 10.24 ± 0.17 ± 0.23 3.7k ³ ADAM 05A CLEO Repl. by MENDEZ 08

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$ quoted in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

³ Not independent from other values reported by ADAM 05A.

NODE=M055B;LINKAGE=AB
NODE=M055B;LINKAGE=J3

NODE=M055B3;LINKAGE=AD

$$\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{17} / \Gamma \times \Gamma_{133}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

NODE=M055B16
NODE=M055B16

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

6.8±0.5 OUR FIT
7.2±0.6 OUR AVERAGE

- 7.3 ± 0.5 ± 0.5 ¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
- 7.0 ± 0.5 ± 0.9 ² ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K^0 K^+ \pi^- + \text{c.c.})$ reported by ATHAR 07 was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54)\%$.

² Calculated by us. ABLIKIM 06R reports $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-) = (4.0 \pm 0.3 \pm 0.5) \times 10^{-3}$. We use $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.7 \pm 0.4) \times 10^{-2}$.

NODE=M055B16;LINKAGE=AT

NODE=M055B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{17}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}$$

NODE=M055B17
NODE=M055B17

VALUE (units 10⁻⁴) DOCUMENT ID TECN COMMENT

19.7±1.6 OUR FIT

13.2±2.4±3.2 ¹BAI 99B BES $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M055B17;LINKAGE=BA

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{36}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}$$

NODE=M055B14
NODE=M055B14

VALUE (units 10⁻⁴) EVTS DOCUMENT ID TECN COMMENT

0.52±0.11 OUR FIT

0.61±0.11±0.08 54 ¹ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$.

NODE=M055B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{36}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}$$

NODE=M055B15
NODE=M055B15

VALUE (units 10⁻⁴) DOCUMENT ID TECN COMMENT

1.52±0.31 OUR FIT

1.13±0.40±0.29 ¹BAI 99B BES $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M055B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{45}/\Gamma \times \Gamma_{133}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}$$

NODE=M055B6
NODE=M055B6

VALUE (units 10⁻⁶) EVTS DOCUMENT ID TECN COMMENT


7.4±0.4 OUR FIT

7.8±0.6 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

7.9±0.4±0.3	453	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
8.2±0.7±0.4	141 ± 13	¹ NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
4.8 ^{+1.4} _{-1.3} ±0.6	18.2 ^{+5.5} _{-4.9}	BAI	04F	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}(1P) \rightarrow \gamma p\bar{p}$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow p\bar{p}) = (9.0 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

NODE=M055B6;LINKAGE=NA




$$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}} \text{ (units } 10^{-6}\text{)}$$

MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

NODE=M055240

 $a_2 = M2/\sqrt{E1^2 + M2^2}$ Magnetic quadrupole fractional transition amplitude
NODE=M055A1
NODE=M055A1

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-5.4 \pm 1.2				
-1.5				
OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.				
-6.26 \pm 0.63 \pm 0.24	39k	ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
0.2 \pm 3.2 \pm 0.4	2090	AMBROGIANI	02	E835 $p\bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi\gamma$
-0.2 \pm 0.8	921	OREGLIA	82	CBAL $\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$
-2.0				


MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$ RADIATIVE DECAY

NODE=M055250

 $b_2 = M2/\sqrt{E1^2 + M2^2}$ Magnetic quadrupole fractional transition amplitude
NODE=M055QB2
NODE=M055QB2

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 \pm 0.8				
OUR AVERAGE				
2.76 \pm 0.73 \pm 0.23	39k	ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
7.7 \pm 5.0	921	OREGLIA	82	CBAL $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-4.5				

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS
 $\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$ and $\chi_{c1} \rightarrow \gamma J/\psi(1S)$

NODE=M055260

 a_2/b_2 Magnetic quadrupole transition amplitude ratio
NODE=M055QAR
NODE=M055QAR

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-2.27 \pm 0.57				
-0.99				
-2.27 \pm 0.57 \pm 0.99	39k	¹ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹Statistical and systematic errors combined. Not independent of $a_2(\chi_{c1})$ and $b_2(\chi_{c1})$ values from ARTUSO 09.

NODE=M055QAR;LINKAGE=AR

 $\chi_{c1}(1P)$ REFERENCES

NODE=M055

ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56778
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55901
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55583
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54742
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53647
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16715
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16719
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53940
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206

BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
Also		Private Comm.	M.J. Oreglia	(EFI)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
TANENBAUM	75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)

REFID=52575
 REFID=52583
 REFID=52588
 REFID=52684
 REFID=52301
 REFID=51618
 REFID=51049
 REFID=51447
 REFID=51453
 REFID=50756
 REFID=50846
 REFID=50763
 REFID=50769
 REFID=49741
 REFID=50188
 REFID=50331
 REFID=49752
 REFID=49755
 REFID=49579
 REFID=49416
 REFID=48552
 REFID=47385
 REFID=46338
 REFID=46343
 REFID=41865
 REFID=41907
 REFID=22145
 REFID=22012
 REFID=22084
 REFID=22120
 REFID=22143
 REFID=22119
 REFID=22113
 REFID=22115
 REFID=22111
 REFID=22112
 REFID=22113
 REFID=22059
 REFID=22062
 REFID=22064
 REFID=22151
 REFID=22106

NODE=M144

$h_c(1P)$

$$I^G(J^{PC}) = ?^?(1+ -)$$

Quantum numbers are quark model prediction, $C = -$ established by $\eta_c \gamma$ decay.

NODE=M144

$h_c(1P)$ MASS

NODE=M144M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3525.38 ± 0.11 OUR AVERAGE				
3525.31 ± 0.11 ± 0.14	832	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
3525.40 ± 0.13 ± 0.18	3679	ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
3525.20 ± 0.18 ± 0.12	1282	² DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3525.8 ± 0.2 ± 0.2	13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3525.6 ± 0.5	92 ⁺²³ ₋₂₂	ADAMS	09 CLEO	$\psi(2S) \rightarrow 2(\pi^+ \pi^- \pi^0)$
3524.4 ± 0.6 ± 0.4	168 ± 40	³ ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3527 ± 8	42	ANTONIAZZI	94 E705	$300 \pi^\pm, pLi \rightarrow J/\psi \pi^0 X$
3526.28 ± 0.18 ± 0.19	59	⁴ ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$
3525.4 ± 0.8 ± 0.4	5	BAGLIN	86 SPEC	$\bar{p}p \rightarrow J/\psi X$

NODE=M144M

¹ With floating width.

² Combination of exclusive and inclusive analyses for the reaction $\psi(2S) \rightarrow \pi^0 h_c \rightarrow \pi^0 \eta_c \gamma$. This result is the average of DOBBS 08A and ROSNER 05.

³ Superseded by DOBBS 08A.

⁴ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.

NODE=M144M;LINKAGE=AB

NODE=M144M;LINKAGE=DO

NODE=M144M;LINKAGE=RO

NODE=M144M;LINKAGE=NW

$h_c(1P)$ WIDTH

NODE=M144W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.70 ± 0.28 ± 0.22					
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 1.44	90	3679	⁶ ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
< 1		13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
< 1.1	90	59	ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$

NODE=M144W

⁵ With floating mass.

⁶ The central value is $\Gamma = 0.73 \pm 0.45 \pm 0.28$ MeV.

NODE=M144W;LINKAGE=AL

NODE=M144W;LINKAGE=AB

$h_c(1P)$ DECAY MODES

NODE=M144215;NODE=M144

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $J/\psi(1S)\pi^0$		
Γ_2 $J/\psi(1S)\pi\pi$	not seen	
Γ_3 $\rho\bar{\rho}$	$< 1.5 \times 10^{-4}$	90%
Γ_4 $\eta_c(1S)\gamma$	$(51 \pm 6) \%$	
Γ_5 $\pi^+\pi^-\pi^0$	$< 2.2 \times 10^{-3}$	
Γ_6 $2\pi^+2\pi^-\pi^0$	$(2.2^{+0.8}_{-0.7}) \%$	
Γ_7 $3\pi^+3\pi^-\pi^0$	$< 2.9 \%$	

DESIG=1

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

 $h_c(1P)$ PARTIAL WIDTHS

NODE=M144220

 $h_c(1P) \Gamma(i)\Gamma(\bar{\rho}\rho)/\Gamma(\text{total})$

NODE=M144223

$\Gamma(\eta_c(1S)\gamma) \times \Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_3/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
12.0 ± 4.5	13	⁷ ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c\gamma$	
⁷ Assuming $\Gamma = 1$ MeV.					

NODE=M144G1
NODE=M144G1

NODE=M144G1;LINKAGE=AN

 $h_c(1P)$ BRANCHING RATIOS

NODE=M144225

$\Gamma(J/\psi(1S)\pi\pi)/\Gamma(J/\psi(1S)\pi^0)$					Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.18	90	ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi\pi^0$	

NODE=M144R1
NODE=M144R1

$\Gamma(\eta_c(1S)\gamma)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
51 ± 6 OUR AVERAGE					
$54.3 \pm 6.7 \pm 5.2$	3679	ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0\gamma\eta_c$	
$48 \pm 6 \pm 7$		⁸ DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0\eta_c\gamma$	
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
$48 \pm 6 \pm 7$	1282	⁹ DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0\eta_c\gamma$	
$46 \pm 12 \pm 7$	168	¹⁰ ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0\eta_c\gamma$	

NODE=M144R2
NODE=M144R2

OCCUR=2

⁸ Average of DOBBS 08A and ROSNER 05. DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (4.16 \pm 0.30 \pm 0.37) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹ DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (4.19 \pm 0.32 \pm 0.45) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰ ROSNER 05 reports $[\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (4.0 \pm 0.8 \pm 0.7) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=DB

NODE=M144R2;LINKAGE=DO

NODE=M144R2;LINKAGE=RO

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT	
<2.2		¹¹ ADAMS	09 CLEO	$\psi(2S) \rightarrow \pi^0\gamma\eta_c$	
¹¹ ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] < 0.19 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.					

NODE=M144R01
NODE=M144R01

NODE=M144R01;LINKAGE=AD

$\Gamma(2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.2^{+0.8}_{-0.6} \pm 0.3$	92	¹² ADAMS	09 CLEO	$\psi(2S) \rightarrow \pi^0\gamma\eta_c$	
¹² ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (1.88^{+0.48+0.47}_{-0.45-0.30}) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M144R02
NODE=M144R02

NODE=M144R02;LINKAGE=AD

$\Gamma(3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
<2.9	13 ADAMS	09	CLEO $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
¹³ ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))]$ $< 2.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.			

NODE=M144R03
NODE=M144R03

NODE=M144R03;LINKAGE=AD

 $\Gamma(h_c(1P) \rightarrow \eta_c(1S)\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}$
 $\Gamma_4/\Gamma \times \Gamma_{15}^{\psi(2S)}/\Gamma_{\psi(2S)}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3 ± 0.4 OUR AVERAGE				
4.58 ± 0.40 ± 0.50	3679	14 ABLIKIM	10B	BES3 $\psi(2S) \rightarrow \pi^0 \gamma X$
4.16 ± 0.30 ± 0.37	1430	15 DOBBS	08A	CLEO $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

NODE=M144R04
NODE=M144R04NODE=M144R04;LINKAGE=AB
NODE=M144R04;LINKAGE=DO
 $\Gamma(h_c(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}$
 $\Gamma_3/\Gamma \times \Gamma_{15}^{\psi(2S)}/\Gamma_{\psi(2S)}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10⁻⁷	90	ABLIKIM	13V	BES3 $\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M144R05
NODE=M144R05 $h_c(1P)$ REFERENCES

ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55583
ABLIKIM	12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54741
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53348
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=53103
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52579
ANDREOTTI	05B	PR D72 032001	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50768
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=50812
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
ARMSTRONG	92D	PRL 69 2337	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43174
BAGLIN	86	PL B171 135	C. Baglin <i>et al.</i>	(LAPP, CERN, TORI, STRB+)	REFID=43180

NODE=M144

 $\chi_{c2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See the Review on " $\psi(2S)$ and χ_c branching ratios" before the $\chi_{c0}(1P)$ Listings.

NODE=M057

NODE=M057

 $\chi_{c2}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3556.20 ± 0.09 OUR AVERAGE				
3555.3 ± 0.6 ± 2.2	2.5k	UEHARA	08	BELL $\gamma\gamma \rightarrow \text{hadrons}$
3555.70 ± 0.59 ± 0.39		ABLIKIM	05G	BES2 $\psi(2S) \rightarrow \gamma \chi_{c2}$
3556.173 ± 0.123 ± 0.020		ANDREOTTI	05A	E835 $p\bar{p} \rightarrow e^+ e^- \gamma$
3559.9 ± 2.9		EISENSTEIN	01	CLE2 $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
3556.4 ± 0.7		BAI	99B	BES $\psi(2S) \rightarrow \gamma X$
3556.22 ± 0.131 ± 0.020	585	¹ ARMSTRONG	92	E760 $\bar{p}p \rightarrow e^+ e^- \gamma$
3556.9 ± 0.4 ± 0.5	50	BAGLIN	86B	SPEC $\bar{p}p \rightarrow e^+ e^- X$
3557.8 ± 0.2 ± 4		² GAISER	86	CBAL $\psi(2S) \rightarrow \gamma X$
3553.4 ± 2.2	66	³ LEMOIGNE	82	GOLI $185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
3555.9 ± 0.7		⁴ OREGLIA	82	CBAL $e^+ e^- \rightarrow J/\psi 2\gamma$
3557 ± 1.5	69	⁵ HIMEL	80	MRK2 $e^+ e^- \rightarrow J/\psi 2\gamma$
3551 ± 11	15	BRANDELIK	79B	DASP $e^+ e^- \rightarrow J/\psi 2\gamma$
3553 ± 4		⁵ BARTEL	78B	CNTR $e^+ e^- \rightarrow J/\psi 2\gamma$
3553 ± 4 ± 4		^{5,6} TANENBAUM	78	MRK1 $e^+ e^-$
3563 ± 7	360	⁵ BIDDICK	77	CNTR $e^+ e^- \rightarrow \gamma X$

NODE=M057M

NODE=M057M

• • • We do not use the following data for averages, fits, limits, etc. • • •

3555.4 ± 1.3	53	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
3543 ± 10	4	WHITAKER	76	MRK1 $e^+ e^- \rightarrow J/\psi 2\gamma$

- ¹ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.
² Using mass of $\psi(2S) = 3686.0$ MeV.
³ $J/\psi(1S)$ mass constrained to 3097 MeV.
⁴ Assuming $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.
⁵ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.
⁶ From a simultaneous fit to radiative and hadronic decay channels.

NODE=M057M;LINKAGE=NW
 NODE=M057M;LINKAGE=C
 NODE=M057M;LINKAGE=P
 NODE=M057M;LINKAGE=E
 NODE=M057M;LINKAGE=D
 NODE=M057M;LINKAGE=M

$\chi_{c2}(1P)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.93 ±0.11				OUR FIT
1.95 ±0.13				OUR AVERAGE
1.915 ±0.188 ±0.013		ANDREOTTI 05A	E835	$\rho\bar{\rho} \rightarrow e^+e^-\gamma$
1.96 ±0.17 ±0.07	585	¹ ARMSTRONG 92	E760	$\bar{\rho}\rho \rightarrow e^+e^-\gamma$
2.6 ^{+1.4} _{-1.0}	50	BAGLIN 86B	SPEC	$\bar{\rho}\rho \rightarrow e^+e^-X$
2.8 ^{+2.1} _{-2.0}		² GAISER 86	CBAL	$\psi(2S) \rightarrow \gamma X$

NODE=M057W
 NODE=M057W

- ¹ Recalculated by ANDREOTTI 05A.
² Errors correspond to 90% confidence level; authors give only width range.

NODE=M057W;LINKAGE=AN
 NODE=M057W;LINKAGE=E

$\chi_{c2}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Hadronic decays		
Γ_1 $2(\pi^+\pi^-)$	(1.07 ±0.10) %	
Γ_2 $\rho\rho$		
Γ_3 $\pi^+\pi^-\pi^0\pi^0$	(1.91 ±0.25) %	
Γ_4 $\rho^+\pi^-\pi^0 + c.c.$	(2.3 ±0.4) %	
Γ_5 $4\pi^0$	(1.16 ±0.16) × 10 ⁻³	
Γ_6 $K^+K^-\pi^0\pi^0$	(2.2 ±0.4) × 10 ⁻³	
Γ_7 $K^+\pi^-\bar{K}^0\pi^0 + c.c.$	(1.44 ±0.21) %	
Γ_8 $\rho^-K^+\bar{K}^0 + c.c.$	(4.3 ±1.3) × 10 ⁻³	
Γ_9 $K^*(892)^0K^-\pi^+ \rightarrow$ $K^-\pi^+K^0\pi^0 + c.c.$	(3.1 ±0.8) × 10 ⁻³	
Γ_{10} $K^*(892)^0\bar{K}^0\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + c.c.$	(4.0 ±0.9) × 10 ⁻³	
Γ_{11} $K^*(892)^-K^+\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + c.c.$	(3.9 ±0.9) × 10 ⁻³	
Γ_{12} $K^*(892)^+K^0\pi^-\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + c.c.$	(3.1 ±0.8) × 10 ⁻³	
Γ_{13} $K^+K^-\eta\pi^0$	(1.3 ±0.5) × 10 ⁻³	
Γ_{14} $K^+K^-\pi^+\pi^-$	(8.9 ±1.0) × 10 ⁻³	
Γ_{15} $K^+K^-\pi^+\pi^-\pi^0$	(1.17 ±0.13) %	
Γ_{16} $K_S^0K^\pm\pi^\mp\pi^+\pi^-$	(7.3 ±0.8) × 10 ⁻³	
Γ_{17} $K^+\bar{K}^*(892)^0\pi^- + c.c.$	(2.2 ±1.1) × 10 ⁻³	
Γ_{18} $K^*(892)^0K^*(892)^0$	(2.4 ±0.5) × 10 ⁻³	
Γ_{19} $3(\pi^+\pi^-)$	(8.6 ±1.8) × 10 ⁻³	
Γ_{20} $\phi\phi$	(1.12 ±0.10) × 10 ⁻³	
Γ_{21} $\omega\omega$	(8.8 ±1.1) × 10 ⁻⁴	
Γ_{22} ωK^+K^-	(7.3 ±0.9) × 10 ⁻⁴	
Γ_{23} $\omega\phi$		
Γ_{24} $\pi\pi$	(2.33 ±0.12) × 10 ⁻³	
Γ_{25} $\rho^0\pi^+\pi^-$	(3.8 ±1.6) × 10 ⁻³	
Γ_{26} $\pi^+\pi^-\eta$	(5.0 ±1.3) × 10 ⁻⁴	
Γ_{27} $\pi^+\pi^-\eta'$	(5.2 ±1.9) × 10 ⁻⁴	
Γ_{28} $\eta\eta$	(5.7 ±0.5) × 10 ⁻⁴	
Γ_{29} K^+K^-	(1.05 ±0.07) × 10 ⁻³	
Γ_{30} $K_S^0K_S^0$	(5.5 ±0.4) × 10 ⁻⁴	
Γ_{31} $\bar{K}^0K^+\pi^- + c.c.$	(1.34 ±0.19) × 10 ⁻³	
Γ_{32} $K^+K^-\pi^0$	(3.2 ±0.8) × 10 ⁻⁴	
Γ_{33} $K^+K^-\eta$	< 3.4 × 10 ⁻⁴	90%
Γ_{34} $K^+K^-\eta'(958)$	(1.94 ±0.34) × 10 ⁻⁴	

NODE=M057215;NODE=M057

NODE=M057;CLUMP=A
 DESIG=3

DESIG=43

DESIG=50

DESIG=51

DESIG=62

DESIG=52

DESIG=54

DESIG=55

DESIG=60

DESIG=56

DESIG=57

DESIG=58

DESIG=59

DESIG=5

DESIG=67

DESIG=78

DESIG=10

DESIG=21

DESIG=4

DESIG=16

DESIG=25

DESIG=79

DESIG=68

DESIG=22

DESIG=9

DESIG=39

DESIG=42

DESIG=14

DESIG=2

DESIG=15

DESIG=17

DESIG=36

DESIG=40

DESIG=82

Γ ₃₅	$\eta\eta'$	$< 6 \times 10^{-5}$	90%	DESIG=34
Γ ₃₆	$\eta'\eta'$	$< 1.0 \times 10^{-4}$	90%	DESIG=35
Γ ₃₇	$\pi^+\pi^-K_S^0K_S^0$	$(2.3 \pm 0.6) \times 10^{-3}$		DESIG=29
Γ ₃₈	$K^+K^-K_S^0K_S^0$	$< 4 \times 10^{-4}$	90%	DESIG=30
Γ ₃₉	$K^+K^-K^+K^-$	$(1.73 \pm 0.21) \times 10^{-3}$		DESIG=24
Γ ₄₀	$K^+K^-\phi$	$(1.48 \pm 0.31) \times 10^{-3}$		DESIG=32
Γ ₄₁	$\bar{K}^0K^+\pi^-\phi + \text{c.c.}$	$(4.8 \pm 0.7) \times 10^{-3}$		DESIG=83
Γ ₄₂	$K^+K^-\pi^0\phi$	$(2.7 \pm 0.5) \times 10^{-3}$		DESIG=84
Γ ₄₃	$\phi\pi^+\pi^-\pi^0$	$(9.3 \pm 1.2) \times 10^{-4}$		DESIG=80
Γ ₄₄	$p\bar{p}$	$(7.5 \pm 0.4) \times 10^{-5}$		DESIG=11
Γ ₄₅	$p\bar{p}\pi^0$	$(4.9 \pm 0.4) \times 10^{-4}$		DESIG=37
Γ ₄₆	$p\bar{p}\eta$	$(1.82 \pm 0.26) \times 10^{-4}$		DESIG=41
Γ ₄₇	$p\bar{p}\omega$	$(3.8 \pm 0.5) \times 10^{-4}$		DESIG=61
Γ ₄₈	$p\bar{p}\phi$	$(2.9 \pm 0.9) \times 10^{-5}$		DESIG=66
Γ ₄₉	$p\bar{p}\pi^+\pi^-$	$(1.32 \pm 0.34) \times 10^{-3}$		DESIG=8
Γ ₅₀	$p\bar{p}\pi^0\pi^0$	$(8.2 \pm 2.5) \times 10^{-4}$		DESIG=53
Γ ₅₁	$p\bar{p}K^+K^-$ (non-resonant)	$(2.00 \pm 0.34) \times 10^{-4}$		DESIG=63
Γ ₅₂	$p\bar{p}K_S^0K_S^0$	$< 7.9 \times 10^{-4}$	90%	DESIG=28
Γ ₅₃	$p\bar{n}\pi^-$	$(8.9 \pm 1.0) \times 10^{-4}$		DESIG=31
Γ ₅₄	$\bar{p}n\pi^+$	$(9.3 \pm 0.9) \times 10^{-4}$		DESIG=75
Γ ₅₅	$p\bar{n}\pi^-\pi^0$	$(2.27 \pm 0.19) \times 10^{-3}$		DESIG=76
Γ ₅₆	$\bar{p}n\pi^+\pi^0$	$(2.21 \pm 0.20) \times 10^{-3}$		DESIG=77
Γ ₅₇	$\Lambda\bar{\Lambda}$	$(1.92 \pm 0.16) \times 10^{-4}$		DESIG=19
Γ ₅₈	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(1.31 \pm 0.17) \times 10^{-3}$		DESIG=27
Γ ₅₉	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$(6.9 \pm 1.6) \times 10^{-4}$		DESIG=70
Γ ₆₀	$\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$	$< 4 \times 10^{-4}$	90%	DESIG=71
Γ ₆₁	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	$< 6 \times 10^{-4}$	90%	DESIG=72
Γ ₆₂	$K^+\bar{p}\Lambda + \text{c.c.}$	$(8.1 \pm 0.6) \times 10^{-4}$		DESIG=38
Γ ₆₃	$K^+\bar{p}\Lambda(1520) + \text{c.c.}$	$(2.9 \pm 0.7) \times 10^{-4}$		DESIG=64
Γ ₆₄	$\Lambda(1520)\bar{\Lambda}(1520)$	$(4.8 \pm 1.5) \times 10^{-4}$		DESIG=65
Γ ₆₅	$\Sigma^0\bar{\Sigma}^0$	$< 6 \times 10^{-5}$	90%	DESIG=47
Γ ₆₆	$\Sigma^+\bar{\Sigma}^-$	$< 7 \times 10^{-5}$	90%	DESIG=48
Γ ₆₇	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$< 1.6 \times 10^{-4}$	90%	DESIG=73
Γ ₆₈	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$< 8 \times 10^{-5}$	90%	DESIG=74
Γ ₆₉	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(1.84 \pm 0.34) \times 10^{-4}$		DESIG=85
Γ ₇₀	$\Xi^0\bar{\Xi}^0$	$< 1.1 \times 10^{-4}$	90%	DESIG=49
Γ ₇₁	$\Xi^-\bar{\Xi}^+$	$(1.48 \pm 0.33) \times 10^{-4}$		DESIG=26
Γ ₇₂	$J/\psi(1S)\pi^+\pi^-\pi^0$	$< 1.5 \%$	90%	DESIG=12
Γ ₇₃	$\pi^0\eta_c$	$< 3.2 \times 10^{-3}$	90%	DESIG=81
Γ ₇₄	$\eta_c(1S)\pi^+\pi^-$	$< 5.4 \times 10^{-3}$	90%	DESIG=69
Radiative decays				
Γ ₇₅	$\gamma J/\psi(1S)$	$(19.2 \pm 0.7) \%$		NODE=M057;CLUMP=B DESIG=6
Γ ₇₆	$\gamma\rho^0$	$< 2.0 \times 10^{-5}$	90%	DESIG=44
Γ ₇₇	$\gamma\omega$	$< 6 \times 10^{-6}$	90%	DESIG=45
Γ ₇₈	$\gamma\phi$	$< 8 \times 10^{-6}$	90%	DESIG=46
Γ ₇₉	$\gamma\gamma$	$(2.74 \pm 0.14) \times 10^{-4}$		DESIG=7

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 240 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 342.4$ for 191 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_{14}	13									
x_{17}	3	21								
x_{18}	8	7	1							
x_{20}	14	12	3	7						
x_{24}	19	16	3	10	24					
x_{25}	19	3	1	2	3	4				
x_{28}	11	9	2	6	14	27	2			
x_{29}	14	12	3	7	17	33	3	19		
x_{30}	13	11	2	6	15	28	3	17	20	
x_{31}	7	6	1	4	8	16	1	9	11	10
x_{39}	9	8	2	5	10	18	2	10	13	11
x_{44}	16	13	3	8	16	24	4	14	17	15
x_{57}	11	9	2	6	14	28	2	16	20	17
x_{75}	24	21	4	12	29	55	5	32	40	34
x_{79}	-8	-6	-1	-3	1	19	-2	13	13	10
Γ	-28	-23	-5	-14	-28	-43	-6	-25	-32	-28
	x_1	x_{14}	x_{17}	x_{18}	x_{20}	x_{24}	x_{25}	x_{28}	x_{29}	x_{30}
x_{39}	6									
x_{44}	8	10								
x_{57}	9	11	14							
x_{75}	19	22	19	33						
x_{79}	6	4	26	13	30					
Γ	-15	-19	-54	-25	-61	-52				
	x_{31}	x_{39}	x_{44}	x_{57}	x_{75}	x_{79}				

$\chi_{c2}(1P)$ PARTIAL WIDTHS

$$\chi_{c2}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$$

NODE=M057220

NODE=M057223

$$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{44} \Gamma_{75} / \Gamma$$

NODE=M057G1
NODE=M057G1

VALUE (eV)

DOCUMENT ID

TECN

COMMENT

27.9 ± 1.3 OUR FIT**27.5 ± 1.5 OUR AVERAGE**

27.0 ± 1.5 ± 1.1

¹ ANDREOTTI 05A E835 $p\bar{p} \rightarrow e^+ e^- \gamma$

27.7 ± 1.5 ± 2.0

^{1,2} ARMSTRONG 92 E760 $p\bar{p} \rightarrow e^+ e^- \gamma$

36 ± 8

¹ BAGLIN 86B SPEC $p\bar{p} \rightarrow e^+ e^- X$ ¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.² Recalculated by ANDREOTTI 05A.

NODE=M057G;LINKAGE=7A

NODE=M057G;LINKAGE=AN

$$\Gamma(\gamma\gamma) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{79} \Gamma_{75} / \Gamma$$

NODE=M057G2
NODE=M057G2

VALUE (eV)

EVTS

DOCUMENT ID

TECN

COMMENT

102 ± 5 OUR FIT**117 ± 10 OUR AVERAGE**

111 ± 12 ± 9 147 ± 15

¹ DOBBS 06 CLE3 10.4 $e^+ e^- \rightarrow$ $e^+ e^- \chi_{c2}$

114 ± 11 ± 9 136 ± 13.3

^{1,2} ABE 02T BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

139 ± 55 ± 21

^{1,3} ACCIARRI 99E L3 $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

242 ± 65 ± 51	1,4 ACKER.,K...	98	OPAL	$e^+e^- \rightarrow e^+e^- \chi_{c2}$
150 ± 42 ± 36	1,5 DOMINICK	94	CLE2	$e^+e^- \rightarrow e^+e^- \chi_{c2}$
470 ± 240 ± 120	1,6 BAUER	93	TPC	$e^+e^- \rightarrow e^+e^- \chi_{c2}$

¹ Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1187 \pm 0.0008$.

² All systematic errors added in quadrature.

³ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACCIARRI 99E is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.0162 \pm 0.0014$.

⁴ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACKERSTAFF,K 98 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1203 \pm 0.0038$.

⁵ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in DOMINICK 94 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

⁶ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in BAUER 93 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

————— $\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ —————

NODE=M057G;LINKAGE=LL
NODE=M057G;LINKAGE=GT
NODE=M057G;LINKAGE=J4

NODE=M057G;LINKAGE=J5

NODE=M057G;LINKAGE=J6

NODE=M057G;LINKAGE=J7

NODE=M057224

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{79}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

5.7 ± 0.5 OUR FIT

5.2 ± 0.7 OUR AVERAGE

5.01 ± 0.44 ± 0.55	1597 ± 138	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
6.4 ± 1.8 ± 0.8		EISENSTEIN	01	CLE2 $e^+e^- \rightarrow e^+e^- \chi_{c2}$

NODE=M057G3
NODE=M057G3

$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_{79}/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7.8	90	<598	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
------	----	------	--------	----	---

NODE=M057G08
NODE=M057G08

$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_{79}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.7 ± 0.5 OUR FIT

4.42 ± 0.42 ± 0.53	780 ± 74	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$
---------------------------	----------	--------	----	--

NODE=M057G09
NODE=M057G09

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_{79}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

6.5 ± 0.9 ± 1.5 1250 DEL-AMO-SA..11M BABR $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M057G02
NODE=M057G02

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{18}\Gamma_{79}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.26 ± 0.24 OUR FIT

0.8 ± 0.17 ± 0.27 151 ± 30 UEHARA 08 BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M057G10
NODE=M057G10

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_{79}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.59 ± 0.05 OUR FIT

0.62 ± 0.07 ± 0.05 89 ± 11 ¹ LIU 12B BELL $\gamma\gamma \rightarrow 2(K^+K^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.58 ± 0.18 ± 0.16 26.5 ± 8.1 UEHARA 08 BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+K^-)$

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.

NODE=M057G12;LINKAGE=LI

$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{21}\Gamma_{79}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.64 90 ¹ LIU 12B BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M057G03
NODE=M057G03

NODE=M057G03;LINKAGE=LI

$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{23}\Gamma_{79}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04 90 ¹ LIU 12B BELL $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹ Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M057G04
NODE=M057G04

NODE=M057G04;LINKAGE=LI

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_{79}/\Gamma$
 VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M057G4
 NODE=M057G4

1.23±0.08 OUR FIT

1.18±0.25 OUR AVERAGE

1.44±0.54±0.47 34 ± 13 ¹UEHARA 09 BELL 10.6 e⁺e⁻ → e⁺e⁻π⁰π⁰
 1.14±0.21±0.17 54 ± 10 ²NAKAZAWA 05 BELL 10.6 e⁺e⁻ → e⁺e⁻π⁺π⁻

¹We multiplied the measurement by 3 to convert from π⁰π⁰ to ππ. Interference with the continuum included.

NODE=M057G4;LINKAGE=UE

²We have multiplied π⁺π⁻ measurement by 3/2 to obtain ππ.

NODE=M057G;LINKAGE=NA

$\Gamma(\rho^0\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_{79}/\Gamma$
 VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M057G07
 NODE=M057G07

2.0±0.9 OUR FIT

3.2±1.9±0.5 986 ± 578 UEHARA 08 BELL γγ → χ_{c2} → 2(π⁺π⁻)

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{28}\Gamma_{79}/\Gamma$
 VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M057G13
 NODE=M057G13

0.53±0.22±0.09

8 ¹UEHARA 10A BELL 10.6 e⁺e⁻ → e⁺e⁻ηη

¹Interference with the continuum not included.

NODE=M057G13;LINKAGE=UE

$\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{29}\Gamma_{79}/\Gamma$
 VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M057G5
 NODE=M057G5

0.56±0.04 OUR FIT

0.44±0.11±0.07 33 ± 8 NAKAZAWA 05 BELL 10.6 e⁺e⁻ → e⁺e⁻K⁺K⁻

$\Gamma(K_S^0K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{30}\Gamma_{79}/\Gamma$
 VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M057G6
 NODE=M057G6

0.291±0.025 OUR FIT

0.27 ^{+0.07}_{-0.06} ±0.03 53 ¹UEHARA 13 BELL γγ → K_S⁰K_S⁰

••• We do not use the following data for averages, fits, limits, etc. •••

0.31 ±0.05 ±0.03 38 ± 7 CHEN 07B BELL e⁺e⁻ → e⁺e⁻χ_{c2}

¹Supersedes CHEN 07B.

NODE=M057G6;LINKAGE=UE

$\Gamma(\bar{K}^0K^+\pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{31}\Gamma_{79}/\Gamma$
 VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M057G01
 NODE=M057G01

0.71±0.11 OUR FIT

1.20±0.33±0.13 126 ¹DEL-AMO-SA..11M BABR γγ → K_S⁰K[±]π[∓]

¹We have multiplied $\bar{K}K\pi$ by 2/3 to obtain $\bar{K}^0K^+\pi^- + \text{c.c.}$

NODE=M057G01;LINKAGE=DE

$\Gamma(K^+K^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{39}\Gamma_{79}/\Gamma$
 VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M057G11
 NODE=M057G11

0.91±0.12 OUR FIT

1.10±0.21±0.15 126 ± 24 UEHARA 08 BELL γγ → χ_{c2} → 2(K⁺K⁻)

$\Gamma(\eta_c(1S)\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{74}\Gamma_{79}/\Gamma$
 VALUE (eV) CL% DOCUMENT ID TECN COMMENT

NODE=M057G05
 NODE=M057G05

<15.7

90 LEES 12AE BABR e⁺e⁻ → e⁺e⁻π⁺π⁻η_c

χ_{c2}(1P) BRANCHING RATIOS

NODE=M057225

HADRONIC DECAYS

NODE=M057305

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_1/Γ
 VALUE DOCUMENT ID

NODE=M057R2
 NODE=M057R2

0.0107±0.0010 OUR FIT

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$ Γ_{25}/Γ_1
 VALUE DOCUMENT ID TECN COMMENT

NODE=M057R38
 NODE=M057R38

0.36±0.15 OUR FIT

0.31±0.17 TANENBAUM 78 MRK1 ψ(2S) → γχ_{c2}

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
1.91±0.25 OUR AVERAGE		[(1.92 ± 0.25)% OUR 2015 AVERAGE]		
1.91±0.24±0.07	903.5	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R46
NODE=M057R46

NEW

NODE=M057R46;LINKAGE=HE

¹ HE 08B reports $1.87 \pm 0.07 \pm 0.22 \pm 0.13$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
2.28±0.35±0.08	1031.9	^{1,2} HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R47
NODE=M057R47

NODE=M057R47;LINKAGE=HE

¹ HE 08B reports $2.23 \pm 0.11 \pm 0.32 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M057R47;LINKAGE=OC

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT
1.16±0.15±0.04	1164	¹ ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R58
NODE=M057R58

NODE=M057R58;LINKAGE=AB

¹ ABLIKIM 11A reports $(1.21 \pm 0.05 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
0.22±0.04±0.01	76.9	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R48
NODE=M057R48

NODE=M057R48;LINKAGE=HE

¹ HE 08B reports $0.21 \pm 0.03 \pm 0.03 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
1.44±0.20±0.05	211.6	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R50
NODE=M057R50

NODE=M057R50;LINKAGE=HE

¹ HE 08B reports $1.41 \pm 0.11 \pm 0.16 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
0.43±0.13±0.01	62.9	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R51
NODE=M057R51

NODE=M057R51;LINKAGE=HE

¹ HE 08B reports $0.42 \pm 0.11 \pm 0.06 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^0K^-\pi^+ \rightarrow K^-\pi^+K^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
0.31±0.08±0.01	38.7	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R57
NODE=M057R57

NODE=M057R57;LINKAGE=HE

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0K^-\pi^+ \rightarrow K^-\pi^+K^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.40±0.09±0.01	63.0	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R52
NODE=M057R52

¹ HE 08B reports $0.39 \pm 0.07 \pm 0.05 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R52;LINKAGE=HE

$$\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.39±0.08±0.01	51.1	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R53
NODE=M057R53

¹ HE 08B reports $0.38 \pm 0.07 \pm 0.04 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R53;LINKAGE=HE

$$\Gamma(K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.31±0.08±0.01	39.3	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R54
NODE=M057R54

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R54;LINKAGE=HE

$$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.133±0.046±0.005	22.9	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R55
NODE=M057R55

¹ HE 08B reports $0.13 \pm 0.04 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R55;LINKAGE=HE

$$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID
8.9±1.0 OUR FIT	

NODE=M057R3
NODE=M057R3

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
11.69±0.13±1.31	11k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R00
NODE=M057R00

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $\text{B}(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R00;LINKAGE=A

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.30±0.11±0.75	4.5k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R73
NODE=M057R73

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $\text{B}(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R73;LINKAGE=A

$$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(K^+ K^- \pi^+ \pi^-) \quad \Gamma_{17}/\Gamma_{14}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.25±0.13 OUR FIT			
0.25±0.13	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R39
NODE=M057R39

$$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID
22±11 OUR FIT	

NODE=M057R9
NODE=M057R9

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID
2.4±0.5 OUR FIT	

NODE=M057R26
NODE=M057R26

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

8.6±1.8 OUR EVALUATION Treating systematic error as correlated.**8.6±1.8 OUR AVERAGE**

8.6±0.9±1.6

¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$

8.7±5.9±0.4

¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

 Γ_{19}/Γ

NODE=M057R4

NODE=M057R4

→ UNCHECKED ←

NODE=M057R;LINKAGE=X3

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID

1.12±0.10 OUR FIT Γ_{20}/Γ

NODE=M057R20

NODE=M057R20

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

EVTS

DOCUMENT ID

TECN

COMMENT

0.88±0.11 OUR AVERAGE

0.85±0.10±0.03

762

¹ ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

1.8 ±0.6 ±0.1 27.7 ± 7.4

² ABLIKIM 05N BES2 $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma 6\pi$

¹ ABLIKIM 11K reports $(8.9 \pm 0.3 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 05N reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{21}/Γ

NODE=M057R28

NODE=M057R28

NODE=M057R28;LINKAGE=AL

NODE=M057R28;LINKAGE=AB

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

EVTS

DOCUMENT ID

TECN

COMMENT

0.73±0.04±0.08

512

¹ ABLIKIM 13B BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

 Γ_{22}/Γ

NODE=M057R74

NODE=M057R74

NODE=M057R74;LINKAGE=A

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

<1.9

90

¹ ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

¹ ABLIKIM 11K reports $< 2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

 Γ_{23}/Γ

NODE=M057R63

NODE=M057R63

NODE=M057R63;LINKAGE=AL

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID

2.33±0.12 OUR FIT Γ_{24}/Γ

NODE=M057R27

NODE=M057R27

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

DOCUMENT ID

38±16 OUR FIT Γ_{25}/Γ

NODE=M057R8

NODE=M057R8

 $\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

CL%

DOCUMENT ID

TECN

COMMENT

0.50±0.13±0.02¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5

90

² ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma\chi_{c2}$

¹ ATHAR 07 reports $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06R reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

 Γ_{26}/Γ

NODE=M057R08

NODE=M057R08

NODE=M057R08;LINKAGE=AT

NODE=M057R08;LINKAGE=AB

$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.52±0.19±0.02¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{27}/Γ NODE=M057R35
NODE=M057R35

NODE=M057R35;LINKAGE=AT

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

DOCUMENT ID

5.7±0.5 OUR FIT Γ_{28}/Γ NODE=M057R16
NODE=M057R16 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID

1.05±0.07 OUR FIT Γ_{29}/Γ NODE=M057R11
NODE=M057R11 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID

0.55±0.04 OUR FIT Γ_{30}/Γ NODE=M057R19
NODE=M057R19 $\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$

VALUE

DOCUMENT ID TECN COMMENT

0.235±0.019 OUR FIT Γ_{30}/Γ_{24} NODE=M057R36
NODE=M057R36

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.27 ±0.07 ±0.04 ^{1,2} CHEN 07B BELL $e^+e^- \rightarrow e^+e^-\chi_{c2}$

¹ Using $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from the $\pi^+\pi^-$ measurement of NAKAZAWA 05 rescaled by 3/2 to convert to $\pi\pi$.

² Not independent from other measurements.

NODE=M057R36;LINKAGE=CH

NODE=M057R36;LINKAGE=NI

 $\Gamma(K_S^0 K_S^0)/\Gamma(K^+K^-)$

VALUE

DOCUMENT ID TECN COMMENT

0.52±0.05 OUR FIT Γ_{30}/Γ_{29} NODE=M057R37
NODE=M057R37

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.70±0.21±0.12 ^{1,2} CHEN 07B BELL $e^+e^- \rightarrow e^+e^-\chi_{c2}$

¹ Using $\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from NAKAZAWA 05.

² Not independent from other measurements.

NODE=M057R37;LINKAGE=CH

NODE=M057R37;LINKAGE=NI

 $\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.32±0.08±0.01¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(0.31 \pm 0.07 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{32}/Γ NODE=M057R05
NODE=M057R05

NODE=M057R05;LINKAGE=AT

 $\Gamma(K^+K^-\eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})

CL%

DOCUMENT ID TECN COMMENT

<0.34

90

¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $< 0.33 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

 Γ_{33}/Γ NODE=M057R09
NODE=M057R09

NODE=M057R09;LINKAGE=AT

 $\Gamma(K^+K^-\eta'(958))/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

EVTS

DOCUMENT ID TECN COMMENT

1.94±0.34

107

¹ ABLIKIM 14J BES3 $\psi(2S) \rightarrow \gamma K^+K^-\eta'(958)$

¹ Derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.72 \pm 0.34)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.

 Γ_{34}/Γ NODE=M057R78
NODE=M057R78

NODE=M057R78;LINKAGE=A

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<0.6	90	3.3 ± 8.0	¹ ASNER	09	CLEO $\psi(2S) \rightarrow \gamma\eta\eta'$
----------------	----	---------------	--------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.4	90		² ADAMS	07	CLEO $\psi(2S) \rightarrow \gamma\chi_{c2}$
------	----	--	--------------------	----	---

¹ ASNER 09 reports $< 0.6 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

² Superseded by ASNER 09. ADAMS 07 reports $< 2.3 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R03
NODE=M057R03

NODE=M057R03;LINKAGE=AS

NODE=M057R03;LINKAGE=AD

 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<1.0	90	12 ± 7	¹ ASNER	09	CLEO $\psi(2S) \rightarrow \gamma\eta'\eta'$
----------------	----	------------	--------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2	90		² ADAMS	07	CLEO $\psi(2S) \rightarrow \gamma\chi_{c2}$
------	----	--	--------------------	----	---

¹ ASNER 09 reports $< 1.0 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

² Superseded by ASNER 09. ADAMS 07 reports $< 3.1 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R04
NODE=M057R04

NODE=M057R04;LINKAGE=AS

NODE=M057R04;LINKAGE=AD

 $\Gamma(\pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$2.3 \pm 0.6 \pm 0.1$	57 ± 11	¹ ABLIKIM	050	BES2 $\psi(2S) \rightarrow \gamma\chi_{c2}$
---	-------------	----------------------	-----	---

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R31
NODE=M057R31

NODE=M057R31;LINKAGE=AB

 $\Gamma(K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<4	90	2.3 ± 2.2	¹ ABLIKIM	050	BES2 $e^+e^- \rightarrow \chi_{c2}\gamma$
--------------	----	---------------	----------------------	-----	---

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ $< 3.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R32
NODE=M057R32

NODE=M057R32;LINKAGE=AB

 $\Gamma(K^+K^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-3})	DOCUMENT ID
--------------------------	-------------

1.73 ± 0.21 OUR FIT

NODE=M057R18
NODE=M057R18

 $\Gamma(K^+K^-\phi)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.48 \pm 0.31 \pm 0.05$	52	¹ ABLIKIM	06T	BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
--	----	----------------------	-----	--

¹ ABLIKIM 06T reports $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R01
NODE=M057R01

NODE=M057R01;LINKAGE=AB

 $\Gamma(\bar{K}^0K^+\pi^-\phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$4.83 \pm 0.32 \pm 0.66$	ABLIKIM	15M	BES3 $\psi(2S) \rightarrow \gamma\chi_{c2}$
--	---------	-----	---

NODE=M057R79
NODE=M057R79

 $\Gamma(K^+K^-\pi^0\phi)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.74 \pm 0.16 \pm 0.44$	ABLIKIM	15M	BES3 $\psi(2S) \rightarrow \gamma\chi_{c2}$
--	---------	-----	---

NODE=M057R80
NODE=M057R80

$\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{43}/Γ VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

0.93±0.06±0.10

408

¹ ABLIKIM

13B

BES3

 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ NODE=M057R75
NODE=M057R75¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R75;LINKAGE=A

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{44}/Γ VALUE (units 10^{-4})

DOCUMENT ID

0.75±0.04 OUR FITNODE=M057R12
NODE=M057R12 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ VALUE (units 10^{-3})

DOCUMENT ID

TECN

COMMENT

0.49±0.04 OUR AVERAGE

0.49±0.04±0.02

¹ ONYISI

10

CLE3

 $\psi(2S) \rightarrow \gamma p\bar{p}X$

0.45±0.09±0.02

² ATHAR

07

CLEO

 $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ NODE=M057R06
NODE=M057R06¹ ONYISI 10 reports $(4.83 \pm 0.25 \pm 0.35 \pm 0.31) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R06;LINKAGE=ON

² ATHAR 07 reports $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R06;LINKAGE=AT

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{46}/Γ VALUE (units 10^{-3})

DOCUMENT ID

TECN

COMMENT

0.182±0.026 OUR AVERAGE

0.180±0.027±0.006

¹ ONYISI

10

CLE3

 $\psi(2S) \rightarrow \gamma p\bar{p}X$

0.19 ±0.07 ±0.01

² ATHAR

07

CLEO

 $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ NODE=M057R34
NODE=M057R34¹ ONYISI 10 reports $(1.76 \pm 0.23 \pm 0.14 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R34;LINKAGE=ON

² ATHAR 07 reports $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R34;LINKAGE=AT

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{47}/Γ VALUE (units 10^{-3})

DOCUMENT ID

TECN

COMMENT

0.38±0.04±0.01¹ ONYISI

10

CLE3

 $\psi(2S) \rightarrow \gamma p\bar{p}X$ NODE=M057R56
NODE=M057R56¹ ONYISI 10 reports $(3.68 \pm 0.35 \pm 0.26 \pm 0.24) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R56;LINKAGE=ON

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{48}/Γ VALUE (units 10^{-5})

EVTS

DOCUMENT ID

TECN

COMMENT

2.9±0.9±0.1

24 ± 7

¹ ABLIKIM

11F

BES3

 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$ NODE=M057R62
NODE=M057R62¹ ABLIKIM 11F reports $(3.04 \pm 0.85 \pm 0.43) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R62;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{49}/Γ VALUE (units 10^{-3})

DOCUMENT ID

TECN

COMMENT

1.32±0.34 OUR EVALUATION

Treating systematic error as correlated.

1.3 ±0.4 OUR AVERAGE Error includes scale factor of 1.3.

1.17±0.19±0.30

¹ BAI

99B

BES

 $\psi(2S) \rightarrow \gamma\chi_{c2}$

2.64±1.03±0.14

¹ TANENBAUM

78

MRK1

 $\psi(2S) \rightarrow \gamma\chi_{c2}$ NODE=M057R6
NODE=M057R6

→ UNCHECKED ←

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

NODE=M057R6;LINKAGE=X3

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.082±0.024±0.003	29.2	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R49
NODE=M057R49

¹ HE 08B reports $0.08 \pm 0.02 \pm 0.01 \pm 0.01\%$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R49;LINKAGE=HE

$\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00±0.33±0.07	131 ± 12	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M057R59
NODE=M057R59

¹ ABLIKIM 11F reports $(2.08 \pm 0.19 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R59;LINKAGE=AB

$\Gamma(p\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7.9	90	¹ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$

NODE=M057R30
NODE=M057R30

¹ Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

NODE=M057R;LINKAGE=AB

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.9±1.0 OUR AVERAGE				
8.8±1.0±0.3	3309	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
10.6±3.6±0.4		² ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$

NODE=M057R33
NODE=M057R33

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(0.80 \pm 0.02 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R33;LINKAGE=AL

² ABLIKIM 06I reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R33;LINKAGE=AB

$\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±0.8±0.3	3732	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$

NODE=M057R70
NODE=M057R70

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(0.85 \pm 0.02 \pm 0.07) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R70;LINKAGE=AL

$\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
22.7±1.8±0.8	2128	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

NODE=M057R71
NODE=M057R71

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(2.07 \pm 0.06 \pm 0.15) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R71;LINKAGE=AL

$\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
22.1±1.9±0.8	2352	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

NODE=M057R72
NODE=M057R72

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ = $(2.01 \pm 0.06 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R72;LINKAGE=AL

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**1.92±0.16 OUR FIT**

DOCUMENT ID

 Γ_{57}/Γ NODE=M057R25
NODE=M057R25 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-5})**131±16±5**

CL%

371

EVTS

DOCUMENT ID

¹ ABLIKIM

TECN

12I

BES3

COMMENT

 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$ Γ_{58}/Γ NODE=M057R29
NODE=M057R29

• • • We do not use the following data for averages, fits, limits, etc. • • •

<350

90

² ABLIKIM

06D

BES2

 $\psi(2S) \rightarrow \chi_{c2}\gamma$

¹ ABLIKIM 12I reports $(137.0 \pm 7.6 \pm 15.7) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R29;LINKAGE=AL

² Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

NODE=M057R29;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ VALUE (units 10^{-5})**69±16±2**

EVTS

36

DOCUMENT ID

¹ ABLIKIM

TECN

12I

BES3

COMMENT

 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$ Γ_{59}/Γ NODE=M057R65
NODE=M057R65

¹ ABLIKIM 12I reports $(71.8 \pm 14.5 \pm 8.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R65;LINKAGE=AL

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

<40

CL%

90

DOCUMENT ID

¹ ABLIKIM

TECN

12I

BES3

COMMENT

 $\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$ Γ_{60}/Γ NODE=M057R66
NODE=M057R66

¹ ABLIKIM 12I reports $< 42 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R66;LINKAGE=AL

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

<60

CL%

90

DOCUMENT ID

¹ ABLIKIM

TECN

12I

BES3

COMMENT

 $\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$ Γ_{61}/Γ NODE=M057R67
NODE=M057R67

¹ ABLIKIM 12I reports $< 61 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R67;LINKAGE=AL

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**8.1±0.6 OUR AVERAGE**

EVTS

5k

DOCUMENT ID

^{1,2} ABLIKIM

TECN

13D

BES3

COMMENT

 $\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$ Γ_{62}/Γ NODE=M057R07
NODE=M057R07

8.0±0.6±0.3

8.7±1.7±0.3

³ ATHAR

07

CLEO

 $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ABLIKIM 13D reports $(8.4 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R07;LINKAGE=AB

² Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.

NODE=M057R07;LINKAGE=LB

³ ATHAR 07 reports $(8.5 \pm 1.4 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R07;LINKAGE=AT

$\Gamma(K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.9 ± 0.7 ± 0.1	79 ± 13	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$
------------------------	---------	----------------------	----------	---

¹ ABLIKIM 11F reports $(3.06 \pm 0.50 \pm 0.54) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R60
NODE=M057R60

NODE=M057R60;LINKAGE=AB

 $\Gamma(\Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.8 ± 1.5 ± 0.2	29 ± 7	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$
------------------------	--------	----------------------	----------	---

¹ ABLIKIM 11F reports $(5.05 \pm 1.29 \pm 0.93) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R61
NODE=M057R61

NODE=M057R61;LINKAGE=AB

 $\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<0.6	90		¹ ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
----------------	----	--	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	90	7.5 ± 3.4	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
------	----	-----------	-------------------	---------	---

¹ ABLIKIM 13H reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R43
NODE=M057R43

NODE=M057R43;LINKAGE=AB

² NAIK 08 reports $< 0.75 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R43;LINKAGE=NA

 $\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<0.7	90	4.0 ± 3.5	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
----------------	----	-----------	-------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	90		² ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
------	----	--	----------------------	----------	---

¹ NAIK 08 reports $< 0.67 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R44
NODE=M057R44

NODE=M057R44;LINKAGE=NA

² ABLIKIM 13H reports $< 0.88 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R44;LINKAGE=AB

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<16	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
---------------	----	----------------------	----------	---

¹ ABLIKIM 12I reports $< 17 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R68
NODE=M057R68

NODE=M057R68;LINKAGE=AL

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<8	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Sigma}(1385)^+$
--------------	----	----------------------	----------	---

¹ ABLIKIM 12I reports $< 8.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R69
NODE=M057R69

NODE=M057R69;LINKAGE=AL

$\Gamma(K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.84 ± 0.33 ± 0.06	51	¹ ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \Xi^+ + \text{c.c.}$

NODE=M057R81
NODE=M057R81

¹ ABLIKIM 15I reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (1.68 \pm 0.26 \pm 0.15) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R81;LINKAGE=A

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.1	90	2.9 ± 1.7	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$

NODE=M057R45
NODE=M057R45

¹ NAIK 08 reports $< 1.06 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R45;LINKAGE=NA

 $\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.48 ± 0.33 ± 0.05	29 ± 5		¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^+ \Xi^-$

NODE=M057R17
NODE=M057R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.7	90		² ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2} \gamma$
-------	----	--	----------------------	----------	---

¹ NAIK 08 reports $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R17;LINKAGE=NA

² Using $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$.

NODE=M057R17;LINKAGE=AB

 $\Gamma(J/\psi(1S) \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.015	90	BARATE	81 SPEC	190 GeV $\pi^- \text{Be} \rightarrow 2\pi 2\mu$

NODE=M057R13
NODE=M057R13

 $\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻³	90	¹ ABLIKIM	15N BES3	$\psi(2S) e^+ e^- \rightarrow \gamma \pi^0 \eta_c$

NODE=M057R77
NODE=M057R77

¹ Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$.

NODE=M057R77;LINKAGE=A

 $\Gamma(\eta_c(1S) \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.54 × 10⁻²	90	^{1,2} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R76
NODE=M057R76

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2 × 10 ⁻²	90	^{1,3} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$
-------------------------	----	------------------------	----------	---

OCCUR=2

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R76;LINKAGE=A

² From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

NODE=M057R76;LINKAGE=B

³ From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

NODE=M057R76;LINKAGE=C

 $\Gamma(\eta_c(1S) \pi^+ \pi^-)/\Gamma(K^0 K^+ \pi^- + \text{c.c.})$ Γ_{74}/Γ_{31}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<16.4	90	¹ LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

NODE=M057R64
NODE=M057R64

¹ We divided the reported limit by 2 to take into account the $K_L^0 K^+ \pi^-$ mode.

NODE=M057R64;LINKAGE=LE

————— RADIATIVE DECAYS —————

NODE=M057310

 $\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.192 ± 0.007 OUR FIT			

NODE=M057R7
NODE=M057R7

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.199 ± 0.005 ± 0.012	¹ ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$
-----------------------	-------------------	----------	---

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma \gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma \chi_{c2})$ from ATHAR 04.

NODE=M057R7;LINKAGE=AD

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<20	90	13 ± 11	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
-----	----	---------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<40	90	17.2 ± 6.8	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
-----	----	------------	----------------------	----------	---

¹ ABLIKIM 11E reports $< 20.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R40
NODE=M057R40

NODE=M057R40;LINKAGE=AB

NODE=M057R40;LINKAGE=BE

 $\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<6	90	1 ± 6	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
----	----	-------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90	0.0 ± 1.8	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$
----	----	-----------	----------------------	----------	---

¹ ABLIKIM 11E reports $< 6.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R41
NODE=M057R41

NODE=M057R41;LINKAGE=AB

NODE=M057R41;LINKAGE=BE

 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 8	90	5 ± 5	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
-----	----	-------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	1.3 ± 2.5	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
-----	----	-----------	----------------------	----------	---

¹ ABLIKIM 11E reports $< 8.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.11 \times 10^{-2}$.

NODE=M057R42
NODE=M057R42

NODE=M057R42;LINKAGE=AB

NODE=M057R42;LINKAGE=BE

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-4})	DOCUMENT ID
--------------------------	-------------

2.74 ± 0.14 OUR FIT

NODE=M057R1
NODE=M057R1

 $\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$ Γ_{79}/Γ_{75}

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.43 ± 0.08 OUR FIT

0.99 ± 0.18

¹ AMBROGIANI 00B E835 $\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$

¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

NODE=M057R23
NODE=M057R23

NODE=M057R;LINKAGE=7A

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(\bar{p}p)/\Gamma_{\text{total}}$ $\Gamma_{79}/\Gamma \times \Gamma_{44}/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

2.06 ± 0.16 OUR FIT

1.7 ± 0.4 OUR AVERAGE

1.60 ± 0.42

ARMSTRONG 93 E760 $\bar{p}p \rightarrow \gamma\gamma X$

9.9 ± 4.5

BAGLIN 87B SPEC $\bar{p}p \rightarrow \gamma\gamma X$

NODE=M057R24
NODE=M057R24

 $\chi_{c2}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M057230

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{14}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

2.34 ± 0.26 OUR FIT

2.5 ± 0.9 OUR AVERAGE Error includes scale factor of 2.3.

1.90 ± 0.14 ± 0.44

BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$

3.8 ± 0.67

¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+\pi^-J/\psi) \times B(J/\psi \rightarrow \ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

NODE=M057B18
NODE=M057B18

NODE=M057B18;LINKAGE=TA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{18} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma^{\psi(2S)}}{\Gamma_{\text{total}}}$$

VALUE (units 10⁻⁴) DOCUMENT ID TECN COMMENT

2.2 ± 0.4 OUR FIT

3.11 ± 0.36 ± 0.48 ABLIKIM 04H BES2 $\psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057B19
NODE=M057B19

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{44} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10⁻⁵) DOCUMENT ID TECN COMMENT

1.98 ± 0.10 OUR FIT

[(1.99 ± 0.10) × 10⁻⁵ OUR 2015 FIT]

1.4 ± 1.1 ¹BAI 98I BES $\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma \bar{p} p$

NODE=M057B1
NODE=M057B1
NEW

¹ Calculated by us. The value for B($\chi_{c2} \rightarrow p \bar{p}$) reported in BAI 98I is derived using B($\psi(2S) \rightarrow \gamma \chi_{c2}$) = (7.8 ± 0.8)% and B($\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-$) = (32.4 ± 2.6)% [BAI 98D].

NODE=M057B;LINKAGE=J8

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p \bar{p}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{44} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma^{\psi(2S)}}{\Gamma_{\text{total}}}$$

VALUE (units 10⁻⁶) EVTS DOCUMENT ID TECN COMMENT

6.85 ± 0.33 OUR FIT

NODE=M057B6
NODE=M057B6

7.1 ± 0.5 OUR AVERAGE Error includes scale factor of 1.2.

7.3 ± 0.4 ± 0.3 405 ABLIKIM 13V BES3 $\psi(2S) \rightarrow \gamma p \bar{p}$

7.2 ± 0.7 ± 0.4 121 ± 12 ¹NAIK 08 CLEO $\psi(2S) \rightarrow \gamma p \bar{p}$

4.4 ^{+1.6}_{-1.4} ± 0.6 14.3 ^{+5.2}_{-4.7} BAI 04F BES $\psi(2S) \rightarrow \gamma \chi_{c2}(1P) \rightarrow \gamma \bar{p} p$

¹ Calculated by us. NAIK 08 reports B($\chi_{c2} \rightarrow p \bar{p}$) = (7.7 ± 0.8 ± 0.4 ± 0.5) × 10⁻⁵ using B($\psi(2S) \rightarrow \gamma \chi_{c2}$) = (9.33 ± 0.14 ± 0.61)%.

NODE=M057B6;LINKAGE=NA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{57} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma^{\psi(2S)}}{\Gamma_{\text{total}}}$$

VALUE (units 10⁻⁶) EVTS DOCUMENT ID TECN COMMENT

NODE=M057B10
NODE=M057B10

17.5 ± 1.3 OUR FIT

17.4 ± 1.4 OUR AVERAGE

18.2 ± 1.4 ± 0.9 207 ¹ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

15.9 ± 2.1 ± 1.0 71 ± 9 ²NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

¹ Calculated by us. ABLIKIM 13H reports B($\chi_{c2} \rightarrow \Lambda \bar{\Lambda}$) = (20.8 ± 1.6 ± 2.3) × 10⁻⁵ from a measurement of B($\chi_{c2} \rightarrow \Lambda \bar{\Lambda}$) × B($\psi(2S) \rightarrow \gamma \chi_{c2}$) assuming B($\psi(2S) \rightarrow \gamma \chi_{c2}$) = (8.74 ± 0.35)%.

NODE=M057B10;LINKAGE=AB

² Calculated by us. NAIK 08 reports B($\chi_{c2} \rightarrow \Lambda \bar{\Lambda}$) = (17.0 ± 2.2 ± 1.1 ± 1.1) × 10⁻⁵ using B($\psi(2S) \rightarrow \gamma \chi_{c2}$) = (9.33 ± 0.14 ± 0.61)%.

NODE=M057B10;LINKAGE=NA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{57} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10⁻⁵) EVTS DOCUMENT ID TECN COMMENT

5.1 ± 0.4 OUR FIT

NODE=M057B11
NODE=M057B11

7.1 ^{+3.1}_{-2.9} ± 1.3 8.3 ^{+3.7}_{-3.4} ¹BAI 03E BES $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

¹ BAI 03E reports [B($\chi_{c2} \rightarrow \Lambda \bar{\Lambda}$) B($\psi(2S) \rightarrow \gamma \chi_{c2}$) / B($\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$)] × [B²($\Lambda \rightarrow \pi^- p$) / B($J/\psi \rightarrow p \bar{p}$)] = (1.33 ^{+0.59}_{-0.55} ± 0.25)%. We calculate from this measurement the presented value using B($\Lambda \rightarrow \pi^- p$) = (63.9 ± 0.5)% and B($J/\psi \rightarrow p \bar{p}$) = (2.17 ± 0.07) × 10⁻³.

NODE=M057B11;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi \pi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{24} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma^{\psi(2S)}}{\Gamma_{\text{total}}}$$

VALUE (units 10⁻⁴) EVTS DOCUMENT ID TECN COMMENT

NODE=M057B02
NODE=M057B02

2.12 ± 0.08 OUR FIT

2.17 ± 0.09 OUR AVERAGE

2.19 ± 0.05 ± 0.15 4.5k ¹ABLIKIM 10A BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

2.23 ± 0.06 ± 0.10 2.5k ²ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$

1.90 ± 0.08 ± 0.20 0.8k ³ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$

OCCUR=2

- ¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \pi^0 \pi^0) = (0.88 \pm 0.02 \pm 0.06 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi\pi$.
- ² Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^+ \pi^-) = (1.59 \pm 0.04 \pm 0.07 \pm 0.10) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.
- ³ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^0 \pi^0) = (0.68 \pm 0.03 \pm 0.07 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi\pi$.

NODE=M057B02;LINKAGE=AB

NODE=M057B02;LINKAGE=AS

NODE=M057B02;LINKAGE=AN

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{24}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.615±0.023 OUR FIT[(0.616 ± 0.023) × 10⁻³ OUR 2015 FIT]**0.54 ± 0.06 OUR AVERAGE**0.66 ± 0.18 ± 0.37 21 ± 6 ¹ BAI 03C BES $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$ 0.54 ± 0.05 ± 0.04 185 ± 16 ² BAI 98I BES $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$ ¹ We have multiplied $\pi^0 \pi^0$ measurement by 3 to obtain $\pi\pi$.² Calculated by us. The value for $B(\chi_{c2} \rightarrow \pi^+ \pi^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.NODE=M057B9
NODE=M057B9

NEW

NODE=M057B;LINKAGE=BM

NODE=M057B;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{28}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.52±0.04 OUR FIT**0.52±0.04 OUR AVERAGE**0.54±0.03±0.04 386 ¹ ABLIKIM 10A BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$ 0.47±0.05±0.05 156 ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \eta\eta$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 0.44 90 ² ADAMS 07 CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$ < 3 90 BAI 03C BES $\psi(2S) \rightarrow \gamma \eta\eta \rightarrow 5\gamma$ 0.62±0.31±0.19 LEE 85 CBAL $\psi(2S) \rightarrow \text{photons}$ ¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \eta\eta) = (0.65 \pm 0.04 \pm 0.05 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$.² Superseded by ASNER 09.NODE=M057B04
NODE=M057B04

NODE=M057B04;LINKAGE=AB

NODE=M057B04;LINKAGE=AD

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{29}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.6±0.6 OUR FIT**10.5±0.3±0.6**1.6k ¹ ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K^+ K^-$ ¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K^+ K^-) = (1.13 \pm 0.03 \pm 0.06 \pm 0.07) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.NODE=M057B03
NODE=M057B03

NODE=M057B03;LINKAGE=AS

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{29}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.277±0.017 OUR FIT[(0.278 ± 0.017) × 10⁻³ OUR 2015 FIT]**0.190±0.034±0.019** 115 ± 13 ¹ BAI 98I BES $\psi(2S) \rightarrow \gamma K^+ K^-$ ¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow K^+ K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].NODE=M057B8
NODE=M057B8

NEW

NODE=M057B;LINKAGE=BI

$$\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{30} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

5.0 ± 0.4 OUR FIT

5.0 ± 0.4 OUR AVERAGE

4.9 ± 0.3 ± 0.3	373 ± 20	¹ ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
5.72 ± 0.76 ± 0.63	65	ABLIKIM	050	BES2	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K_S^0 K_S^0) = (0.53 \pm 0.03 \pm 0.03 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

NODE=M057B12
NODE=M057B12

NODE=M057B12;LINKAGE=AS

$$\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{30} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

14.5 ± 1.1 OUR FIT

14.7 ± 4.1 ± 3.3

		¹ BAI	99B	BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
--	--	------------------	-----	-----	---

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M057B13
NODE=M057B13

NODE=M057B13;LINKAGE=BA

$$\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{31} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

1.22 ± 0.17 OUR FIT

1.15 ± 0.18 OUR AVERAGE

1.21 ± 0.19 ± 0.09	37	¹ ATHAR	07	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
0.97 ± 0.32 ± 0.13	28	² ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Calculated by us. ATHAR 07 reports $B(\chi_{c2} \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (1.3 \pm 0.2 \pm 0.1 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

² Calculated by us. ABLIKIM 06R reports $B(\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp) = (0.6 \pm 0.2 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.1 \pm 0.6)\%$. We have multiplied by 2 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$ from $K_S^0 K^\pm \pi^\mp$.

NODE=M057B05
NODE=M057B05

NODE=M057B05;LINKAGE=AT

NODE=M057B05;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_1 / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

2.83 ± 0.27 OUR FIT

3.1 ± 1.0 OUR AVERAGE Error includes scale factor of 2.5.

2.3 ± 0.1 ± 0.5	¹ BAI	99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
4.3 ± 0.6	² TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S)\ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

NODE=M057B5
NODE=M057B5

NODE=M057B;LINKAGE=K1

NODE=M057B;LINKAGE=K2

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{39} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

1.57 ± 0.19 OUR FIT

1.76 ± 0.16 ± 0.24 160 ¹ ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

NODE=M057B14
NODE=M057B14

NODE=M057B14;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{39} / \Gamma \times \Gamma_{134}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

4.6 ± 0.5 OUR FIT

3.6 ± 0.6 ± 0.6 ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M057B15
NODE=M057B15

NODE=M057B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{20}}{\Gamma} \times \frac{\Gamma_{134}^{\psi(2S)}}{\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.02±0.08 OUR FIT				
0.98±0.13 OUR AVERAGE				Error includes scale factor of 1.3.
0.94±0.03±0.10	849	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
1.38±0.24±0.23	41	² ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35)\%$.
² Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

NODE=M057B16
 NODE=M057B16

NODE=M057B16;LINKAGE=AL

NODE=M057B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)} \times \frac{\Gamma_{20}}{\Gamma} \times \frac{\Gamma_{134}^{\psi(2S)}}{\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.95±0.24 OUR FIT			
4.8 ±1.3 ±1.3	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98b].

NODE=M057B17
 NODE=M057B17

NODE=M057B17;LINKAGE=BA


$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{75}}{\Gamma} \times \frac{\Gamma_{134}^{\psi(2S)}}{\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.75 ±0.04 OUR FIT				
1.52 ±0.15 OUR AVERAGE				Error includes scale factor of 2.6. See the ideogram below.
1.874±0.007±0.102	76k	ABLIKIM	120 BES3	$\psi(2S) \rightarrow \gamma\chi_{c2}$
1.62 ±0.04 ±0.12	5.8k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
0.99 ±0.10 ±0.08		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
1.47 ±0.17		¹ OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\chi_{c2}$
1.8 ±0.5		² BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma\chi_{c2}$
1.2 ±0.2		² BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma\chi_{c2}$
2.2 ±1.2		³ BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$
1.2 ±0.7		¹ WHITAKER	76 MRK1	e^+e^-
••• We do not use the following data for averages, fits, limits, etc. •••				
1.95 ±0.02 ±0.07	12.4k	⁴ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2}$
1.85 ±0.04 ±0.07	1.9k	⁵ ADAM	05A CLEO	Repl. by MENDEZ 08

NODE=M057B2
 NODE=M057B2

NODE=M057B;LINKAGE=3Q
 NODE=M057B;LINKAGE=2Q
 NODE=M057B;LINKAGE=EA
 NODE=M057B2;LINKAGE=ME
 NODE=M057B;LINKAGE=AD

- ¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- ² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.
- ³ Assumes isotropic gamma distribution.
- ⁴ Not independent from other measurements of MENDEZ 08.
- ⁵ Not independent from other values reported by ADAM 05A.



$$10^{-2} \times \frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))}{\Gamma_{\text{total}}}$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\frac{\Gamma_{75}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_9^{\psi(2S)}}{\Gamma_{75}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + 0.339\Gamma_{133}^{\psi(2S)} + 0.192\Gamma_{134}^{\psi(2S)}}$$

$$\frac{\Gamma_{75}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_9^{\psi(2S)}}{\Gamma_{75}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

NODE=M057B7

NODE=M057B7

NODE=M057B7

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.87±0.07 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.12±0.03±0.09	12.4k	¹ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
3.11±0.07±0.07	1.9k	ADAM	05A	CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

NODE=M057B7;LINKAGE=ME

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\frac{\Gamma_{75}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}{\Gamma_{75}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

NODE=M057B3

NODE=M057B3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.08±0.12 OUR FIT**5.53±0.17 OUR AVERAGE**

5.56±0.05±0.16	12.4k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
6.0 ±2.8	1.3k	¹ ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
3.9 ±1.2		² HIMEL	80	MRK2	$\psi(2S) \rightarrow \gamma \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.52±0.13±0.13	1.9k	³ ADAM	05A	CLEO	Repl. by MENDEZ 08
----------------	------	-------------------	-----	------	--------------------

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

NODE=M057B;LINKAGE=AB

NODE=M057B;LINKAGE=H8

NODE=M057B3;LINKAGE=AD

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma \gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\frac{\Gamma_{79}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma^{\psi(2S)}}{\Gamma_{79}/\Gamma \times \Gamma_{134}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

NODE=M057B4

NODE=M057B4

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.50±0.13 OUR FIT**2.78±0.18 OUR AVERAGE**

2.81±0.17±0.15	1.1k	¹ ABLIKIM	12A	BES3	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow 3\gamma$
2.68±0.28±0.15	0.3k	ECKLUND	08A	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow 3\gamma$
7.0 ±2.1 ±2.0		LEE	85	CBAL	$\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ ABLIKIM 12A measures the ratio of two-photon partial widths for the helicity $\lambda = 0$ and helicity $\lambda = 2$ components to be $f_{0/2} = \Gamma_{\gamma\gamma}^{\lambda=0}/\Gamma_{\gamma\gamma}^{\lambda=2} = 0.00 \pm 0.02 \pm 0.02$.

NODE=M057B4;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma \gamma)/\Gamma(\chi_{c0}(1P) \rightarrow \gamma \gamma)$$

$$\frac{\Gamma_{79}/\Gamma_{89}^{\chi_{c0}(1P)}}{\Gamma_{79}/\Gamma_{89}^{\chi_{c0}(1P)}}$$

NODE=M057B06

NODE=M057B06

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.273±0.035 OUR AVERAGE

0.271±0.029±0.030	1.9k	¹ ABLIKIM	12A	BES3	$\psi(2S) \rightarrow \gamma \chi_{cJ} \rightarrow 3\gamma$
0.278±0.050±0.036	0.5k	¹ ECKLUND	08A	CLEO	$\psi(2S) \rightarrow \gamma \chi_{cJ} \rightarrow 3\gamma$

¹ Not independent from the values of $\Gamma(\chi_{c0}, \chi_{c2})$ and $B(\psi(2S) \rightarrow \chi_{c0}, \chi_{c2})$.

NODE=M057B06;LINKAGE=AB

MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

NODE=M057240

 $a_2 = M_2/\sqrt{E_1^2 + M_2^2 + E_3^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

-10.0± 1.5 OUR AVERAGE

- 9.3± 1.6±0.3	19.8k	¹ ARTUSO	09	CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
- 9.3 ⁺ ₋ 3.9 ⁺ _{4.1} ±0.6	5.9k	² AMBROGIANI	02	E835	$\rho \bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi \gamma$
-14 ± 6	1.9k	² ARMSTRONG	93E	E760	$\rho \bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi \gamma$
-33.3 ⁺ ₋ 11.6 ⁺ _{29.2}	441	² OREGLIA	82	CBAL	$\psi(2S) \rightarrow \chi_{c1} \gamma \rightarrow J/\psi \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

- 7.9± 1.9±0.3	19.8k	³ ARTUSO	09	CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
----------------	-------	---------------------	----	------	--

OCCUR=2

¹ From a fit with floating M_2 amplitudes a_2 and b_2 , and fixed E_3 amplitudes $a_3=b_3=0$.

² Assuming $a_3=0$.

³ From a fit with floating M_2 and E_3 amplitudes a_2 , b_2 , and a_3 , and b_3 .

NODE=M057A1;LINKAGE=AR

NODE=M057A1;LINKAGE=A

NODE=M057A1;LINKAGE=AT

$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.6±1.3 OUR AVERAGE				
1.7±1.4±0.3	19.8k	¹ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.0 ^{+5.5} _{-4.4} ±0.9	5908	AMBROGIANI	02 E835	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
0 ⁺⁶ ₋₅	1904	ARMSTRONG	93E E760	$\rho\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$

¹ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

NODE=M057A2
NODE=M057A2

NODE=M057A2;LINKAGE=AR

MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ RADIATIVE DECAY

NODE=M057250

$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±1.8 OUR AVERAGE				
Error includes scale factor of 1.7. See the ideogram below.				
4.6±1.0±1.3	13.8k	¹ ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
0.2±1.5±0.4	19.8k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
- 5.1 ^{+5.4} _{-3.6}	721	¹ ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
13.2 ^{+9.8} _{-7.5}	441	³ OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M057QB2
NODE=M057QB2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •


1.0±1.3±0.3	19.8k	³ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-------------	-------	---------------------	---------	---

¹ From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

² From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

³ From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$.

NODE=M057QB2;LINKAGE=AB
NODE=M057QB2;LINKAGE=AT
NODE=M057QB2;LINKAGE=AR



$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude (units 10^{-2})

$b_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-0.3±1.0 OUR AVERAGE				
1.5±0.8±1.8	13.8k	¹ ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
-0.8±1.2±0.2	19.8k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-2.7 ^{+4.3} _{-2.9}	721	¹ ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$

¹ From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

NODE=M057QB3
NODE=M057QB3

NODE=M057QB3;LINKAGE=AB

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$$\psi(2S) \rightarrow \gamma \chi_{c2}(1P) \text{ and } \chi_{c2} \rightarrow \gamma J/\psi(1S)$$

 b_2/a_2 Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-11^{+14}_{-15}	19.8k	¹ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Statistical and systematic errors combined. From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$. Not independent of values for $a_2(\chi_{c2}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

NODE=M057260

NODE=M057QAR
NODE=M057QAR

NODE=M057QAR;LINKAGE=AR

 $\chi_{c2}(1P)$ REFERENCES

NODE=M057

ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56778
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56779
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55901
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55583
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54266
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54742
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53647
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=16719
ABLIKIM	11I	PR D84 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53930
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53940
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53347
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
DOBBS	06	PR D73 071101	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51062
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ABLIKIM	04I	PR D70 092004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50189
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE	02T	PL B540 33	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48813
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48552
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
ACCIARRI	99E	PL B453 73	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46943
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ACKER...K...	98	PL B439 197	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46324
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
DOMINICK	94	PR D50 4265	J. Dominick <i>et al.</i>	(CLEO Collab.)	REFID=44077
ARMSTRONG	93	PRL 70 2988	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43306
ARMSTRONG	93E	PR D48 3037	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=48616
BAUER	93	PL B302 345	D.A. Bauer <i>et al.</i>	(TPC Collab.)	REFID=43315
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41865
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41907
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)	REFID=22145
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
Also		Private Comm.	M.J. Oreglia	(EFI)	REFID=22143
BARATE	81	PR D24 2994	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, CERN+)	REFID=22164
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BRANDELIC	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151

$$I^G(J^{PC}) = 0^+(0^-+)$$

Quantum numbers are quark model predictions.

NODE=M059

NODE=M059

NODE=M059M

NODE=M059M

OCCUR=2

OCCUR=2

$\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3639.2±1.2 OUR AVERAGE				
3637.0±5.7±3.4	178	^{1,2} LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^0$
3635.1±5.8±2.1	47	^{1,3} LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \eta$
3646.9±1.6±3.6	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow$ $\gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6±2.9±1.6	127 ± 18	⁴ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$
3638.5±1.5±0.8	624	¹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5±3.2±2.5	1201	¹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^\mp K^- \pi^+ \pi^- \pi^0$
3636.1 ^{+3.9+0.7} -4.2-2.0	128	⁵ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	⁶ ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
3645.0±5.5 ^{+4.9} -7.8	121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
3642.9±3.1±1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3639 ± 7	98 ± 52	⁷ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
3630.8±3.4±1.0	112 ± 24	⁸ AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
3654 ± 6 ± 8	39 ± 11	⁹ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		¹⁰ EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ Ignoring possible interference with continuum.

² With a width fixed to 11.3 MeV.

³ With a width fixed to 11.3 MeV. Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁴ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

⁵ Accounts for interference with non-resonant continuum.

⁶ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

⁷ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

⁸ Superseded by DEL-AMO-SANCHEZ 11M.

⁹ Superseded by VINOKUROVA 11.

¹⁰ Assuming mass of $\psi(2S) = 3686$ MeV.

NODE=M059M;LINKAGE=DE

NODE=M059M;LINKAGE=LE

NODE=M059M;LINKAGE=LS

NODE=M059M;LINKAGE=AB

NODE=M059M;LINKAGE=VA

NODE=M059M;LINKAGE=EB

NODE=M059M;LINKAGE=AU

NODE=M059M;LINKAGE=AR

NODE=M059M;LINKAGE=CH

NODE=M059M;LINKAGE=A

$\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.3⁺_{-2.9} OUR AVERAGE					
9.9 ± 4.8 ± 2.9		57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow$ $\gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 ± 6.4 ± 4.8		127 ± 18	¹¹ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi,$ $K K \pi^0$
13.4 ± 4.6 ± 3.2		624	¹² DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
6.6 ⁺ ₋ 8.4+2.6 5.1-0.9		128	¹³ VINOKUROVA	11 BELL	$B^\pm \rightarrow$ $K^\pm (K_S^0 K^\pm \pi^\mp)$
6.3 ± 12.4 ± 4.0		61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow$ $K_S^0 K^\pm \pi^\mp$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 23 90 98 ± 52 ¹⁴AUBERT 06E BABR $B^\pm \rightarrow K^\pm X_{c\bar{c}}$

22 ± 14 121 ± 27 AUBERT 05C BABR $e^+ e^- \rightarrow J/\psi c\bar{c}$

17.0 ± 8.3 ± 2.5 112 ± 24 ¹⁵AUBERT 04D BABR $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow$

<55 90 39 ± 11 ¹⁶CHOI 02 BELL $B \rightarrow K K_S K^- \pi^+$

<8.0 95 ¹⁷EDWARDS 82C CBAL $e^+ e^- \rightarrow \gamma X$

¹¹ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

¹² Ignoring possible interference with continuum.

¹³ Accounts for interference with non-resonant continuum.

¹⁴ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁵ Superseded by DEL-AMO-SANCHEZ 11M.

¹⁶ For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11.

¹⁷ For a mass value of 3594 ± 5 MeV

NODE=M059W

NODE=M059W

NODE=M059W;LINKAGE=AB

NODE=M059W;LINKAGE=DE

NODE=M059W;LINKAGE=VA

NODE=M059W;LINKAGE=AU

NODE=M059W;LINKAGE=AR

NODE=M059W;LINKAGE=W2

NODE=M059W;LINKAGE=W

$\eta_c(2S)$ DECAY MODES

NODE=M059215;NODE=M059

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	not seen	
Γ_2 $K\bar{K}\pi$	(1.9±1.2) %	
Γ_3 $K\bar{K}\eta$	(5 ± 4) × 10 ⁻³	
Γ_4 $2\pi^+2\pi^-$	not seen	
Γ_5 $\rho^0\rho^0$	not seen	
Γ_6 $3\pi^+3\pi^-$	not seen	
Γ_7 $K^+K^-\pi^+\pi^-$	not seen	
Γ_8 $K^{*0}\bar{K}^{*0}$	not seen	
Γ_9 $K^+K^-\pi^+\pi^-\pi^0$	(1.4±1.0) %	
Γ_{10} $K^+K^-2\pi^+2\pi^-$	not seen	
Γ_{11} $K_S^0K^-2\pi^+\pi^- + c.c.$	seen	
Γ_{12} $2K^+2K^-$	not seen	
Γ_{13} $\phi\phi$	not seen	
Γ_{14} $p\bar{p}$	< 2.0 × 10 ⁻³	90%
Γ_{15} $\gamma\gamma$	(1.9±1.3) × 10 ⁻⁴	
Γ_{16} $\pi^+\pi^-\eta$	not seen	
Γ_{17} $\pi^+\pi^-\eta'$	not seen	
Γ_{18} $\pi^+\pi^-\eta_c(1S)$	< 25 %	90%

DESIG=1

DESIG=4

DESIG=20

DESIG=5

DESIG=16

DESIG=8;OUR EVAL;→ UNCHECKED ←

DESIG=6

DESIG=17

DESIG=9

DESIG=10;OUR EVAL;→ UNCHECKED ←

DESIG=11

DESIG=7

DESIG=18

DESIG=3

DESIG=2

DESIG=12;OUR EVAL;→ UNCHECKED ←

DESIG=13;OUR EVAL;→ UNCHECKED ←

DESIG=15

 $\eta_c(2S)$ PARTIAL WIDTHS

NODE=M059216

 $\Gamma(\gamma\gamma)$ Γ_{15}

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M059W1

NODE=M059W1

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3±0.6 18 ASNER 04 CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$

¹⁸They measure $\Gamma(\eta_c(2S)\gamma\gamma)$ $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02)$ $\Gamma(\eta_c(1S)\gamma\gamma)$ $B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S^0 K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.

NODE=M059W1;LINKAGE=AS

 $\eta_c(2S)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M059218

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<6.5 90 UEHARA 08 BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

 $\Gamma_4\Gamma_{15}/\Gamma$

NODE=M059G01

NODE=M059G01

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

41±4±6 624 ¹⁹ DEL-AMO-SA..11M BABR $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

 $\Gamma_2\Gamma_{15}/\Gamma$

NODE=M059G04

NODE=M059G04

¹⁹Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059G04;LINKAGE=DE

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<5.0 90 UEHARA 08 BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

 $\Gamma_7\Gamma_{15}/\Gamma$

NODE=M059G02

NODE=M059G02

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

30±6±5 1201 ²⁰ DEL-AMO-SA..11M BABR $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

 $\Gamma_9\Gamma_{15}/\Gamma$

NODE=M059G05

NODE=M059G05

²⁰Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059G05;LINKAGE=DE

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<2.9 90 UEHARA 08 BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+K^-)$

 $\Gamma_{12}\Gamma_{15}/\Gamma$

NODE=M059G03

NODE=M059G03

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<133 90 LEES 12AE BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

 $\Gamma_{18}\Gamma_{15}/\Gamma$

NODE=M059G06

NODE=M059G06

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

NODE=M059217

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{14}/\Gamma \times \Gamma_{15}/\Gamma$

VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90 ^{21,22,23}	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	90 ^{21,22,24}	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
< 12.0	90 ^{22,24}	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

NODE=M059G1
NODE=M059G1OCCUR=2
OCCUR=3²¹ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.²² For a total width $\Gamma=5$ MeV.²³ For the resonance mass region 3589–3599 MeV/ c^2 .²⁴ For the resonance mass region 3575–3660 MeV/ c^2 .NODE=M059G1;LINKAGE=A
NODE=M059G1;LINKAGE=B
NODE=M059G1;LINKAGE=C1
NODE=M059G1;LINKAGE=C2 $\eta_c(2S)$ BRANCHING RATIOS

NODE=M059220

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABREU 98O	DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	²⁵ EDWARDS 82C	CBAL	$e^+e^- \rightarrow \gamma X$
²⁵ For a mass value of 3594 ± 5 MeV			

NODE=M059R1
NODE=M059R1

NODE=M059R;LINKAGE=W

 $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.4 \pm 1.1$	59 ± 12	²⁶ AUBERT 08AB	BABR	$B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	127 ± 18	ABLIKIM 13K	BES3	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$
seen	39 ± 11	²⁷ CHOI 02	BELL	$B \rightarrow K K_S K^- \pi^+$

NODE=M059R3
NODE=M059R3

NODE=M059R3;LINKAGE=AU

²⁶ Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$.²⁷ For a mass value of 3654 ± 6 MeV

NODE=M059R;LINKAGE=W2

 $\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$ Γ_3/Γ_2

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$27.3 \pm 7.0 \pm 5.7$	225	²⁸ LEES 14E	BABR	$\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$
²⁸ LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.17$, which we divide by 3 to account for isospin symmetry.				

NODE=M059R26
NODE=M059R26

NODE=M059R26;LINKAGE=LE

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

NODE=M059R01
NODE=M059R01 $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M059R15
NODE=M059R15 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA 08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

NODE=M059R02
NODE=M059R02 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ Γ_9/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.73 \pm 0.17 \pm 0.17$	1201	²⁹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M059R21
NODE=M059R21²⁹ We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059R21;LINKAGE=DE

 $\Gamma(K^*0 \bar{K}^*0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

NODE=M059R16
NODE=M059R16

$$\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$

NODE=M059R22
NODE=M059R22

$$\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

NODE=M059R03
NODE=M059R03

$$\Gamma(\phi\phi)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M059R17
NODE=M059R17

$$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5 × 10 ⁻⁴	90	30 WICHT	08 BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
not seen		AMBROGIANI	01 E835	$\bar{p}p \rightarrow \gamma\gamma$
<0.01	90	LEE	85 CBAL	$\psi' \rightarrow \text{photons}$

NODE=M059R2
NODE=M059R2

³⁰WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c(2S) K^+) = 3.4 \times 10^{-4}$.

NODE=M059R2;LINKAGE=W1

$$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma(K\bar{K}\pi) \quad \Gamma_{18}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.33	90	31 LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

NODE=M059R23
NODE=M059R23

³¹We divided the reported limit by 3 to take into account isospin relations.

NODE=M059R23;LINKAGE=L1

$\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M059230

$$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<14.6 × 10 ⁻⁶	90	32 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M059R05
NODE=M059R05

³²Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R05;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<12.7 × 10 ⁻⁷	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M059R18
NODE=M059R18

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<13.2 × 10 ⁻⁶	90	33 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$

NODE=M059R06
NODE=M059R06

³³Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R06;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9.6 × 10 ⁻⁶	90	34 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

NODE=M059R07
NODE=M059R07

³⁴Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R07;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<19.6 × 10 ⁻⁷	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

NODE=M059R19
NODE=M059R19

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_9 / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	35 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M059R08
NODE=M059R08

³⁵ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R08;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{10} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	36 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

NODE=M059R09
NODE=M059R09

³⁶ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R09;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{11} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60		ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

NODE=M059R10
NODE=M059R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

<15.2	90	37 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
---------	----	-------------------	------	--

³⁷ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R10;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{13} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M059R20
NODE=M059R20

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{16} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-6}$	90	38 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta$

NODE=M059R11
NODE=M059R11

³⁸ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R11;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta') / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{17} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<14.2 \times 10^{-6}$	90	39 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M059R12
NODE=M059R12

³⁹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R12;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K \bar{K} \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_3 / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<11.8 \times 10^{-6}$	90	40 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$

NODE=M059R25
NODE=M059R25

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁴⁰ CRONIN-HENNESSY 10 reports a limit of $< 5.9 \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 account for isospin symmetry. It assumes $\Gamma(\eta_c(2S)) = 14$ MeV. It also gives the analytic dependence of limits on width.

NODE=M059R25;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{18} / \Gamma \times \Gamma_{136}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	41 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$

NODE=M059R14
NODE=M059R14

⁴¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R14;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{14}/\Gamma \times \Gamma_{136}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M059R24
 NODE=M059R24

$\eta_c(2S)$ REFERENCES

LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55937
ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54956
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55583
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54272
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53929
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=53927
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=53233
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52267
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50773
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=49745
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49746
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=48760
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48340
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46553
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=44623
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22173

NODE=M059

$\psi(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the Review on " $\psi(2S)$ and χ_c branching ratios" before the $\chi_{c0}(1P)$ Listings.

NODE=M071

NODE=M071

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

NODE=M071M

NODE=M071M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3686.097 ± 0.025 OUR FIT				Error includes scale factor of 2.6. [3686.109 ^{+0.012} _{-0.014} MeV
OUR 2015 FIT]				
3686.097 ± 0.010 OUR AVERAGE				
[3686.108 ^{+0.011} _{-0.014} MeV				OUR 2015 AVERAGE]
3686.099 ± 0.004 ± 0.009		¹ ANASHIN	15	KEDR $e^+e^- \rightarrow$ hadrons
3686.12 ± 0.06 ± 0.10	4k	AAIJ	12H	LHCB $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3685.95 ± 0.10	413	² ARTAMONOV	00	OLYA $e^+e^- \rightarrow$ hadrons
3685.98 ± 0.09 ± 0.04		³ ARMSTRONG	93B	E760 $p\bar{p} \rightarrow e^+e^-$
3686.114 ± 0.007 ^{+0.011} _{-0.016}		⁴ ANASHIN	12	KEDR $e^+e^- \rightarrow$ hadrons
3686.111 ± 0.025 ± 0.009		AULCHENKO	03	KEDR $e^+e^- \rightarrow$ hadrons
3686.00 ± 0.10	413	⁵ ZHOLENTZ	80	OLYA e^+e^-

NODE=M071M

NEW

NEW

¹ Supersedes AULCHENKO 03 and ANASHIN 12.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

⁴ From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$ MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

⁵ Superseded by ARTAMONOV 00.

NODE=M071M;LINKAGE=A
 NODE=M071M;LINKAGE=AR

NODE=M071M;LINKAGE=NW

NODE=M071M;LINKAGE=AN

NODE=M071M;LINKAGE=RZ

$$m_{\psi(2S)} - m_{J/\psi(1S)}$$

NODE=M071DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
589.188 ± 0.028 OUR AVERAGE			
589.194 ± 0.027 ± 0.011	¹ AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
589.7 ± 1.2	LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- \text{A}$
589.07 ± 0.13	¹ ZHOLENTZ 80	OLYA	e^+e^-
588.7 ± 0.8	LUTH 75	MRK1	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
588 ± 1	² BAI 98E	BES	e^+e^-
¹ Redundant with data in mass above.			
² Systematic errors not evaluated.			

NODE=M071DM

NODE=M071DM;LINKAGE=R
NODE=M071DM;LINKAGE=BD **$\psi(2S)$ WIDTH**

VALUE (keV)	EVTs	DOCUMENT ID	TECN	COMMENT
296 ± 8 OUR FIT				
[298 ± 8 keV OUR 2015 FIT]				
286 ± 16 OUR AVERAGE				
358 ± 88 ± 4		ABLIKIM 08B	BES2	$e^+e^- \rightarrow \text{hadrons}$
290 ± 25 ± 4	2.7k	ANDREOTTI 07	E835	$p\bar{p} \rightarrow e^+e^-, J/\psi X$
331 ± 58 ± 2		ABLIKIM 06L	BES2	$e^+e^- \rightarrow \text{hadrons}$
264 ± 27		¹ BAI 02B	BES2	e^+e^-
287 ± 37 ± 16		² ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
¹ From a simultaneous fit to the hadronic and $\mu^+\mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.				
² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].				

NODE=M071W

NODE=M071W
NEW

NODE=M071W;LINKAGE=BC

NODE=M071W;LINKAGE=AN

 $\psi(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(97.85 ± 0.13) %	
Γ_2 virtual $\gamma \rightarrow \text{hadrons}$	(1.73 ± 0.14) %	S=1.5
Γ_3 ggg	(10.6 ± 1.6) %	
Γ_4 γgg	(1.03 ± 0.29) %	
Γ_5 light hadrons	(15.4 ± 1.5) %	
Γ_6 e^+e^-	(7.89 ± 0.17) × 10 ⁻³	
Γ_7 $\mu^+\mu^-$	(7.9 ± 0.9) × 10 ⁻³	
Γ_8 $\tau^+\tau^-$	(3.1 ± 0.4) × 10 ⁻³	

NODE=M071220;NODE=M071

DESIG=3

DESIG=4

DESIG=255

DESIG=256

DESIG=226

DESIG=1

DESIG=2

DESIG=68

Decays into $J/\psi(1S)$ and anything

Γ_9 $J/\psi(1S)$ anything	(61.0 ± 0.6) %
Γ_{10} $J/\psi(1S)$ neutrals	(25.14 ± 0.33) %
Γ_{11} $J/\psi(1S) \pi^+ \pi^-$	(34.49 ± 0.30) %
Γ_{12} $J/\psi(1S) \pi^0 \pi^0$	(18.16 ± 0.31) %
Γ_{13} $J/\psi(1S) \eta$	(3.36 ± 0.05) %
Γ_{14} $J/\psi(1S) \pi^0$	(1.268 ± 0.032) × 10 ⁻³

NODE=M071;CLUMP=A

DESIG=11

DESIG=12

DESIG=13

DESIG=14

DESIG=15

DESIG=18

Hadronic decays

Γ_{15} $\pi^0 h_c(1P)$	(8.6 ± 1.3) × 10 ⁻⁴	
Γ_{16} $3(\pi^+ \pi^-) \pi^0$	(3.5 ± 1.6) × 10 ⁻³	
Γ_{17} $2(\pi^+ \pi^-) \pi^0$	(2.9 ± 1.0) × 10 ⁻³	S=4.7
Γ_{18} $\rho a_2(1320)$	(2.6 ± 0.9) × 10 ⁻⁴	
Γ_{19} $p\bar{p}$	(2.88 ± 0.09) × 10 ⁻⁴	
Γ_{20} $\Delta^{++} \bar{\Delta}^{--}$	(1.28 ± 0.35) × 10 ⁻⁴	
Γ_{21} $\Lambda \bar{\Lambda} \pi^0$	< 2.9 × 10 ⁻⁶	CL=90%
Γ_{22} $\Lambda \bar{\Lambda} \eta$	(2.5 ± 0.4) × 10 ⁻⁵	
Γ_{23} $\Lambda \bar{p} K^+$	(1.00 ± 0.14) × 10 ⁻⁴	
Γ_{24} $\Lambda \bar{p} K^+ \pi^+ \pi^-$	(1.8 ± 0.4) × 10 ⁻⁴	
Γ_{25} $\Lambda \bar{\Lambda} \pi^+ \pi^-$	(2.8 ± 0.6) × 10 ⁻⁴	
Γ_{26} $\Lambda \bar{\Lambda}$	(3.57 ± 0.18) × 10 ⁻⁴	
Γ_{27} $\Lambda \bar{\Sigma}^+ \pi^- + \text{c.c.}$	(1.40 ± 0.13) × 10 ⁻⁴	
Γ_{28} $\Lambda \bar{\Sigma}^- \pi^+ + \text{c.c.}$	(1.54 ± 0.14) × 10 ⁻⁴	
Γ_{29} $\Sigma^0 \bar{p} K^+ + \text{c.c.}$	(1.67 ± 0.18) × 10 ⁻⁵	

NODE=M071;CLUMP=B

DESIG=254

DESIG=37

DESIG=25

DESIG=65

DESIG=27

DESIG=70

DESIG=238

DESIG=239

DESIG=214

DESIG=215

DESIG=213

DESIG=28

DESIG=280

DESIG=281

DESIG=274

Г30	$\Sigma^+ \bar{\Sigma}^-$	$(2.51 \pm 0.21) \times 10^{-4}$		DESIG=223
Г31	$\Sigma^0 \bar{\Sigma}^0$	$(2.32 \pm 0.16) \times 10^{-4}$		DESIG=71
Г32	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$(1.1 \pm 0.4) \times 10^{-4}$		DESIG=72
Г33	$\Xi^- \bar{\Xi}^+$	$(2.64 \pm 0.18) \times 10^{-4}$		DESIG=29
Г34	$\Xi^0 \bar{\Xi}^0$	$(2.07 \pm 0.23) \times 10^{-4}$		DESIG=224
Г35	$\Xi(1530)^0 \bar{\Xi}(1530)^0$	$(5.2 \pm_{-1.2}^{+3.2}) \times 10^{-5}$		DESIG=73
Г36	$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(3.9 \pm 0.4) \times 10^{-5}$		DESIG=293
Г37	$\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(5.2 \pm 1.6) \times 10^{-6}$		DESIG=294
Г38	$\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(1.20 \pm 0.32) \times 10^{-5}$		DESIG=295
Г39	$K^- \Sigma^0 \bar{\Xi}^+ + \text{c.c.}$	$(3.7 \pm 0.4) \times 10^{-5}$		DESIG=296
Г40	$\Omega^- \bar{\Omega}^+$	$(4.7 \pm 1.0) \times 10^{-5}$		DESIG=74
Г41	$\pi^0 p \bar{p}$	$(1.53 \pm 0.07) \times 10^{-4}$		DESIG=35
Г42	$N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(6.4 \pm_{-1.3}^{+1.8}) \times 10^{-5}$		DESIG=267
Г43	$N(1440) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(7.3 \pm_{-1.5}^{+1.7}) \times 10^{-5}$	S=2.5	DESIG=261
Г44	$N(1520) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(6.4 \pm_{-1.8}^{+2.3}) \times 10^{-6}$		DESIG=268
Г45	$N(1535) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(2.5 \pm 1.0) \times 10^{-5}$		DESIG=269
Г46	$N(1650) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(3.8 \pm_{-1.7}^{+1.4}) \times 10^{-5}$		DESIG=270
Г47	$N(1720) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(1.79 \pm_{-0.70}^{+0.26}) \times 10^{-5}$		DESIG=271
Г48	$N(2300) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(2.6 \pm_{-0.7}^{+1.2}) \times 10^{-5}$		DESIG=272
Г49	$N(2570) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p}$	$(2.13 \pm_{-0.31}^{+0.40}) \times 10^{-5}$		DESIG=273
Г50	$\pi^0 f_0(2100) \rightarrow \pi^0 p \bar{p}$	$(1.1 \pm 0.4) \times 10^{-5}$		DESIG=262
Г51	$\eta p \bar{p}$	$(6.0 \pm 0.4) \times 10^{-5}$		DESIG=200
Г52	$\eta f_0(2100) \rightarrow \eta p \bar{p}$	$(1.2 \pm 0.4) \times 10^{-5}$		DESIG=263
Г53	$N(1535) \bar{p} \rightarrow \eta p \bar{p}$	$(4.4 \pm 0.7) \times 10^{-5}$		DESIG=264
Г54	$\omega p \bar{p}$	$(6.9 \pm 2.1) \times 10^{-5}$		DESIG=77
Г55	$\phi p \bar{p}$	$< 2.4 \times 10^{-5}$	CL=90%	DESIG=80
Г56	$\pi^+ \pi^- p \bar{p}$	$(6.0 \pm 0.4) \times 10^{-4}$		DESIG=31
Г57	$p \bar{n} \pi^- \text{ or c.c.}$	$(2.48 \pm 0.17) \times 10^{-4}$		DESIG=227
Г58	$p \bar{n} \pi^- \pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$		DESIG=228
Г59	$2(\pi^+ \pi^- \pi^0)$	$(4.8 \pm 1.5) \times 10^{-3}$		DESIG=221
Г60	$\eta \pi^+ \pi^-$	$< 1.6 \times 10^{-4}$	CL=90%	DESIG=202
Г61	$\eta \pi^+ \pi^- \pi^0$	$(9.5 \pm 1.7) \times 10^{-4}$		DESIG=203
Г62	$2(\pi^+ \pi^-) \eta$	$(1.2 \pm 0.6) \times 10^{-3}$		DESIG=251
Г63	$\eta' \pi^+ \pi^- \pi^0$	$(4.5 \pm 2.1) \times 10^{-4}$		DESIG=204
Г64	$\omega \pi^+ \pi^-$	$(7.3 \pm 1.2) \times 10^{-4}$	S=2.1	DESIG=75
Г65	$b^\pm \pi^\mp$	$(4.0 \pm 0.6) \times 10^{-4}$	S=1.1	DESIG=40
Г66	$b_1^0 \pi^0$	$(2.4 \pm 0.6) \times 10^{-4}$		DESIG=193
Г67	$\omega f_2(1270)$	$(2.2 \pm 0.4) \times 10^{-4}$		DESIG=64
Г68	$\pi^+ \pi^- K^+ K^-$	$(7.5 \pm 0.9) \times 10^{-4}$	S=1.9	DESIG=26
Г69	$\rho^0 K^+ K^-$	$(2.2 \pm 0.4) \times 10^{-4}$		DESIG=205
Г70	$K^*(892)^0 \bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$		DESIG=66
Г71	$K^+ K^- \pi^+ \pi^- \eta$	$(1.3 \pm 0.7) \times 10^{-3}$		DESIG=252
Г72	$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$		DESIG=240
Г73	$K^+ K^- 2(\pi^+ \pi^-)$	$(1.9 \pm 0.9) \times 10^{-3}$		DESIG=222
Г74	$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$		DESIG=41
Г75	$K_S^0 K_S^0 \pi^+ \pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$		DESIG=225
Г76	$\rho^0 p \bar{p}$	$(5.0 \pm 2.2) \times 10^{-5}$		DESIG=210
Г77	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$		DESIG=34
Г78	$2(\pi^+ \pi^-)$	$(2.4 \pm 0.6) \times 10^{-4}$	S=2.2	DESIG=24
Г79	$\rho^0 \pi^+ \pi^-$	$(2.2 \pm 0.6) \times 10^{-4}$	S=1.4	DESIG=33
Г80	$K^+ K^- \pi^+ \pi^- \pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$		DESIG=206

Г81	$\omega f_0(1710) \rightarrow \omega K^+ K^-$	$(5.9 \pm 2.2) \times 10^{-5}$		DESIG=216
Г82	$K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$		DESIG=217
Г83	$K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$		DESIG=218
Г84	$K^*(892)^+ K^- \rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$		DESIG=219
Г85	$K^*(892)^0 K^- \rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$		DESIG=220
Г86	$\eta K^+ K^-$, no $\eta\phi$	$(3.1 \pm 0.4) \times 10^{-5}$		DESIG=207
Г87	$\omega K^+ K^-$	$(1.62 \pm 0.11) \times 10^{-4}$	S=1.1	DESIG=76
Г88	$\omega K^*(892)^+ K^- + \text{c.c.}$	$(2.07 \pm 0.26) \times 10^{-4}$		DESIG=276
Г89	$\omega K_2^*(1430)^+ K^- + \text{c.c.}$	$(6.1 \pm 1.2) \times 10^{-5}$		DESIG=277
Г90	$\omega \bar{K}^*(892)^0 K^0$	$(1.68 \pm 0.30) \times 10^{-4}$		DESIG=278
Г91	$\omega \bar{K}_2^*(1430)^0 K^0$	$(5.8 \pm 2.2) \times 10^{-5}$		DESIG=279
Г92	$\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-5}$		DESIG=282
Г93	$\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$	$(1.09 \pm 0.26) \times 10^{-5}$		DESIG=283
Г94	$\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.}$	$(3.0 \pm 1.0) \times 10^{-6}$		DESIG=284
Г95	$\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$	$(1.2 \pm 0.7) \times 10^{-6}$		DESIG=285
Г96	$3(\pi^+ \pi^-)$	$(3.5 \pm 2.0) \times 10^{-4}$	S=2.8	DESIG=32
Г97	$p\bar{p}\pi^+ \pi^- \pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$		DESIG=211
Г98	$K^+ K^-$	$(7.5 \pm 0.5) \times 10^{-5}$		DESIG=23
Г99	$K_S^0 K_L^0$	$(5.34 \pm 0.33) \times 10^{-5}$		DESIG=85
Г100	$\pi^+ \pi^- \pi^0$	$(2.01 \pm 0.17) \times 10^{-4}$	S=1.7	DESIG=36
Г101	$\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$	$(1.9 \pm 1.2) \times 10^{-4}$		DESIG=201
Г102	$\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$	$(3.2 \pm 1.2) \times 10^{-5}$	S=1.8	DESIG=22
Г103	$\pi^+ \pi^-$	$(7.8 \pm 2.6) \times 10^{-6}$		DESIG=21
Г104	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%	DESIG=42
Г105	$K_2^*(1430)^\pm K^\mp$	$(7.1 \pm 1.3) \times 10^{-5}$		DESIG=265
Г106	$K^+ K^- \pi^0$	$(4.07 \pm 0.31) \times 10^{-5}$		DESIG=38
Г107	$K^+ K^*(892)^- + \text{c.c.}$	$(2.9 \pm 0.4) \times 10^{-5}$	S=1.2	DESIG=39
Г108	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$		DESIG=194
Г109	$\phi \pi^+ \pi^-$	$(1.17 \pm 0.29) \times 10^{-4}$	S=1.7	DESIG=78
Г110	$\phi f_0(980) \rightarrow \pi^+ \pi^-$	$(6.8 \pm 2.5) \times 10^{-5}$	S=1.2	DESIG=81
Г111	$2(K^+ K^-)$	$(6.0 \pm 1.4) \times 10^{-5}$		DESIG=208
Г112	$\phi K^+ K^-$	$(7.0 \pm 1.6) \times 10^{-5}$		DESIG=79
Г113	$2(K^+ K^-) \pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$		DESIG=209
Г114	$\phi \eta$	$(3.10 \pm 0.31) \times 10^{-5}$		DESIG=89
Г115	$\phi \eta'$	$(3.1 \pm 1.6) \times 10^{-5}$		DESIG=90
Г116	$\omega \eta'$	$(3.2 \pm 2.5) \times 10^{-5}$		DESIG=91
Г117	$\omega \pi^0$	$(2.1 \pm 0.6) \times 10^{-5}$		DESIG=92
Г118	$\rho \eta'$	$(1.9 \pm 1.7) \times 10^{-5}$		DESIG=93
Г119	$\rho \eta$	$(2.2 \pm 0.6) \times 10^{-5}$	S=1.1	DESIG=94
Г120	$\omega \eta$	$< 1.1 \times 10^{-5}$	CL=90%	DESIG=95
Г121	$\phi \pi^0$	$< 4 \times 10^{-7}$	CL=90%	DESIG=96
Г122	$\eta_c \pi^+ \pi^- \pi^0$	$< 1.0 \times 10^{-3}$	CL=90%	DESIG=229
Г123	$p\bar{p}K^+ K^-$	$(2.7 \pm 0.7) \times 10^{-5}$		DESIG=212
Г124	$\bar{\Lambda} n K_S^0 + \text{c.c.}$	$(8.1 \pm 1.8) \times 10^{-5}$		DESIG=237
Г125	$\phi f_2'(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$		DESIG=67
Г126	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	$< 8.8 \times 10^{-6}$	CL=90%	DESIG=195
Г127	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$< 1.0 \times 10^{-5}$	CL=90%	DESIG=196
Г128	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$< 7.0 \times 10^{-6}$	CL=90%	DESIG=197
Г129	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$< 2.6 \times 10^{-5}$	CL=90%	DESIG=198
Г130	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$< 6.0 \times 10^{-6}$	CL=90%	DESIG=199
Г131	$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$		DESIG=86

Radiative decays

Γ ₁₃₂	$\gamma\chi_{c0}(1P)$	(9.99 ± 0.27) %		NODE=M071;CLUMP=C
Γ ₁₃₃	$\gamma\chi_{c1}(1P)$	(9.55 ± 0.31) %		DESIG=56
Γ ₁₃₄	$\gamma\chi_{c2}(1P)$	(9.11 ± 0.31) %		DESIG=58
Γ ₁₃₅	$\gamma\eta_c(1S)$	(3.4 ± 0.5) × 10 ⁻³	S=1.3	DESIG=59
Γ ₁₃₆	$\gamma\eta_c(2S)$	(7 ± 5) × 10 ⁻⁴		DESIG=61
Γ ₁₃₇	$\gamma\pi^0$	(1.6 ± 0.4) × 10 ⁻⁶		DESIG=63
Γ ₁₃₈	$\gamma\eta'(958)$	(1.23 ± 0.06) × 10 ⁻⁴		DESIG=52
Γ ₁₃₉	$\gamma f_2(1270)$	(2.73 ^{+0.29} _{-0.25}) × 10 ⁻⁴	S=1.8	DESIG=54
Γ ₁₄₀	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	(3.1 ± 1.7) × 10 ⁻⁵		DESIG=82
Γ ₁₄₁	$\gamma f_0(1500)$	(9.2 ± 1.9) × 10 ⁻⁵		DESIG=286
Γ ₁₄₂	$\gamma f_2'(1525)$	(3.3 ± 0.8) × 10 ⁻⁵		DESIG=287
Γ ₁₄₃	$\gamma f_0(1710)$			DESIG=288
Γ ₁₄₄	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	(3.5 ± 0.6) × 10 ⁻⁵		DESIG=236
Γ ₁₄₅	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	(6.6 ± 0.7) × 10 ⁻⁵		DESIG=83
Γ ₁₄₆	$\gamma f_0(2100) \rightarrow \gamma\pi\pi$	(4.8 ± 1.0) × 10 ⁻⁶		DESIG=84
Γ ₁₄₇	$\gamma f_0(2200) \rightarrow \gamma K\bar{K}$	(3.2 ± 1.0) × 10 ⁻⁶		DESIG=289
Γ ₁₄₈	$\gamma f_J(2220) \rightarrow \gamma\pi\pi$	< 5.8 × 10 ⁻⁶	CL=90%	DESIG=290
Γ ₁₄₉	$\gamma f_J(2220) \rightarrow \gamma K\bar{K}$	< 9.5 × 10 ⁻⁶	CL=90%	DESIG=291
Γ ₁₅₀	$\gamma\gamma$	< 1.5 × 10 ⁻⁴	CL=90%	DESIG=292
Γ ₁₅₁	$\gamma\eta$	(1.4 ± 0.5) × 10 ⁻⁶		DESIG=51
Γ ₁₅₂	$\gamma\eta\pi^+\pi^-$	(8.7 ± 2.1) × 10 ⁻⁴		DESIG=53
Γ ₁₅₃	$\gamma\eta(1405)$			DESIG=230
Γ ₁₅₄	$\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi$	< 9 × 10 ⁻⁵	CL=90%	DESIG=231
Γ ₁₅₅	$\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-$	(3.6 ± 2.5) × 10 ⁻⁵		DESIG=62
Γ ₁₅₆	$\gamma\eta(1475)$			DESIG=232
Γ ₁₅₇	$\gamma\eta(1475) \rightarrow K\bar{K}\pi$	< 1.4 × 10 ⁻⁴	CL=90%	DESIG=233
Γ ₁₅₈	$\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-$	< 8.8 × 10 ⁻⁵	CL=90%	DESIG=234
Γ ₁₅₉	$\gamma 2(\pi^+\pi^-)$	(4.0 ± 0.6) × 10 ⁻⁴		DESIG=235
Γ ₁₆₀	$\gamma K^{*0}K^+\pi^- + c.c.$	(3.7 ± 0.9) × 10 ⁻⁴		DESIG=241
Γ ₁₆₁	$\gamma K^{*0}\bar{K}^{*0}$	(2.4 ± 0.7) × 10 ⁻⁴		DESIG=242
Γ ₁₆₂	$\gamma K_S^0K^+\pi^- + c.c.$	(2.6 ± 0.5) × 10 ⁻⁴		DESIG=243
Γ ₁₆₃	$\gamma K^+K^-\pi^+\pi^-$	(1.9 ± 0.5) × 10 ⁻⁴		DESIG=244
Γ ₁₆₄	$\gamma\rho\bar{\rho}$	(3.9 ± 0.5) × 10 ⁻⁵	S=2.0	DESIG=245
Γ ₁₆₅	$\gamma f_2(1950) \rightarrow \gamma\rho\bar{\rho}$	(1.20 ± 0.22) × 10 ⁻⁵		DESIG=246
Γ ₁₆₆	$\gamma f_2(2150) \rightarrow \gamma\rho\bar{\rho}$	(7.2 ± 1.8) × 10 ⁻⁶		DESIG=257
Γ ₁₆₇	$\gamma X(1835) \rightarrow \gamma\rho\bar{\rho}$	(4.6 ^{+1.8} _{-4.0}) × 10 ⁻⁶		DESIG=258
Γ ₁₆₈	$\gamma X \rightarrow \gamma\rho\bar{\rho}$	[a] < 2 × 10 ⁻⁶	CL=90%	DESIG=259
Γ ₁₆₉	$\gamma\pi^+\pi^-\rho\bar{\rho}$	(2.8 ± 1.4) × 10 ⁻⁵		DESIG=260
Γ ₁₇₀	$\gamma 2(\pi^+\pi^-)K^+K^-$	< 2.2 × 10 ⁻⁴	CL=90%	DESIG=247
Γ ₁₇₁	$\gamma 3(\pi^+\pi^-)$	< 1.7 × 10 ⁻⁴	CL=90%	DESIG=248
Γ ₁₇₂	$\gamma K^+K^-K^+K^-$	< 4 × 10 ⁻⁵	CL=90%	DESIG=249
Γ ₁₇₃	$\gamma\gamma J/\psi$	(3.1 ^{+1.0} _{-1.2}) × 10 ⁻⁴		DESIG=250

Other decays

Γ ₁₇₄	invisible	< 1.6 %	CL=90%	NODE=M071;CLUMP=D
------------------	-----------	---------	--------	-------------------

[a] For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

LINKAGE=NMR

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 240 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 342.4$ for 191 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_7	3									
x_8	1	0								
x_{11}	30	8	2							
x_{12}	29	5	1	49						
x_{13}	13	3	1	36	16					
x_{19}	0	0	0	5	3	2				
x_{132}	1	0	0	3	1	1	0			
x_{133}	2	0	0	4	1	1	0	0		
x_{134}	1	0	0	4	1	1	0	0	0	
Γ	-81	-3	-1	-39	-35	-17	-9	-1	-2	-2
	x_6	x_7	x_8	x_{11}	x_{12}	x_{13}	x_{19}	x_{132}	x_{133}	x_{134}

$\psi(2S)$ PARTIAL WIDTHS

NODE=M071225

$\Gamma(\text{hadrons})$

 Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

258 ± 26	BAI	02B	BES2 e^+e^-
224 ± 56	LUTH	75	MRK1 e^+e^-

NODE=M071W3
NODE=M071W3

$\Gamma(e^+e^-)$

 Γ_6

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

2.34 \pm 0.04 OUR FIT

[2.35 \pm 0.04 keV OUR 2015 FIT]

2.30 \pm 0.06 OUR AVERAGE

[2.33 \pm 0.07 keV OUR 2015 AVERAGE]

$2.24 \pm 0.10 \pm 0.02$	¹ ABLIKIM	15V	BES3	4.0-4.4 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
$2.338 \pm 0.037 \pm 0.096$	ABLIKIM	08B	BES2	$e^+e^- \rightarrow \text{hadrons}$
$2.330 \pm 0.036 \pm 0.110$	ABLIKIM	06L	BES2	$e^+e^- \rightarrow \text{hadrons}$
2.44 ± 0.21	² BAI	02B	BES2	e^+e^-
2.14 ± 0.21	ALEXANDER	89	RVUE	See Υ mini-review

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3	BRANDELIK	79C	DASP	e^+e^-
2.1 ± 0.3	³ LUTH	75	MRK1	e^+e^-

¹ ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau / 0.38847$.

³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

NODE=M071W1
NODE=M071W1

NEW

NEW

NODE=M071W1;LINKAGE=A

NODE=M071W;LINKAGE=BB

NODE=M071W1;LINKAGE=F

$\Gamma(\gamma\gamma)$

 Γ_{150}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<43	90	BRANDELIK	79C	DASP e^+e^-
---------------	----	-----------	-----	---------------

NODE=M071W51
NODE=M071W51

$\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(i) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

NODE=M071230

NODE=M071230

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_6/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

2.233±0.015±0.042	¹ ANASHIN	12	KEDR $e^+e^- \rightarrow \text{hadrons}$
--------------------------	----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.2 ±0.4	ABRAMS	75	MRK1 e^+e^-
----------	--------	----	---------------

¹ ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

NODE=M071G3
NODE=M071G3

NODE=M071G3;LINKAGE=AN

 $\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.0±2.6	79	¹ ANASHIN	07	KEDR $e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$
---------	----	----------------------	----	---

¹ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

NODE=M071G9
NODE=M071G9

NODE=M071G9;LINKAGE=AN

 $\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}\Gamma_6/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.807±0.013 OUR FIT

[0.811 ± 0.013 keV OUR 2015 FIT]

0.837±0.025 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

0.837±0.028±0.005	¹ LEES	12E	BABR	10.6 $e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$
-------------------	-------------------	-----	------	--

0.852±0.010±0.026	19.5k	ADAM	06	CLEO $e^+e^- \rightarrow \gamma\psi(2S)$
-------------------	-------	------	----	--

0.68 ±0.09	² BAI	98E	BES	e^+e^-
------------	------------------	-----	-----	----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ±0.08 ±0.03	256	³ AUBERT	07AU	BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
------------------	-----	---------------------	------	------	--

0.755±0.048±0.004	544	⁴ AUBERT	05D	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$
-------------------	-----	---------------------	-----	------	--

¹ LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

³ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.11 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.


NODE=M071G1
NODE=M071G1
NEW

NODE=M071G1;LINKAGE=LE

NODE=M071G1;LINKAGE=A

NODE=M071G1;LINKAGE=UB

NODE=M071G1;LINKAGE=AU



$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{12}\Gamma_6/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.425±0.009 OUR FIT				
[0.427 ± 0.009 keV OUR 2015 FIT]				
0.411±0.008±0.018	3.6k±96	ADAM	06	CLEO 3.773 e ⁺ e ⁻ → $\gamma\psi(2S)$

NODE=M071G6
NODE=M071G6
NEW

$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{13}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.6± 1.6 OUR FIT				
[79.1 ± 1.7 eV OUR 2015 FIT]				
87 ± 9 OUR AVERAGE				
83 ±25 ±5	14	¹ AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → $J/\psi\pi^+\pi^-\pi^0\gamma$
88 ± 6 ±7	291 ± 24	ADAM	06	CLEO 3.773 e ⁺ e ⁻ → $\gamma\psi(2S)$
¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow J/\psi\eta) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07$ eV.				

NODE=M071G7
NODE=M071G7
NEW

NODE=M071G7;LINKAGE=UB

$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{14}\Gamma_6/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8	90	<37	ADAM	06	CLEO 3.773 e ⁺ e ⁻ → $\gamma\psi(2S)$

NODE=M071G8
NODE=M071G8

$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{19}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.674±0.023 OUR FIT				
[0.676 ± 0.023 eV OUR 2015 FIT]				
0.64 ±0.04 OUR AVERAGE				
0.67 ±0.12 ±0.02	43	LEES	130	BABR e ⁺ e ⁻ → $p\bar{p}\gamma$
0.74 ±0.07 ±0.04	142	LEES	13Y	BABR e ⁺ e ⁻ → $p\bar{p}\gamma$
0.579±0.038±0.036	2.7k	ANDREOTTI	07	E835 $p\bar{p} \rightarrow e^+e^-, J/\psi X$
0.70 ±0.17 ±0.03	22	AUBERT	06B	e ⁺ e ⁻ → $p\bar{p}\gamma$

NODE=M071G2
NODE=M071G2
NEW

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{26}\Gamma_6/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.5±0.4±0.1	AUBERT	07BD	BABR 10.6 e ⁺ e ⁻ → $\Lambda\bar{\Lambda}\gamma$

NODE=M071G11
NODE=M071G11

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{59}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.2±3.3±1.3	43	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → $2(\pi^+\pi^-\pi^0)\gamma$

NODE=M071G4
NODE=M071G4

$\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{73}\Gamma_6/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±2.1±0.3	26	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → $K^+K^-2(\pi^+\pi^-)\gamma$

NODE=M071G5
NODE=M071G5

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{68}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.56±0.42±0.16	85	AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

NODE=M071G12
NODE=M071G12

$$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{110}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.347±0.169±0.003	6 ± 3	¹ AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

NODE=M071G13
NODE=M071G13

¹ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.17 \pm 0.08 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G13;LINKAGE=AU

$$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{109}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.57±0.23±0.01	10	¹ AUBERT, BE	06D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ γ

NODE=M071G10
NODE=M071G10

¹ AUBERT, BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.28 \pm 0.11 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (48.9 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G10;LINKAGE=AU

$$\Gamma(2(\pi^+\pi^-)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
29.7±2.2±1.8	410	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)π ⁰ γ

NODE=M071G01
NODE=M071G01

$$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{64}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.01±0.84±0.02	37	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ

NODE=M071G02
NODE=M071G02

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 2.69 \pm 0.73 \pm 0.16$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G02;LINKAGE=UB

$$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{62}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.87±1.41±0.01	16	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ

NODE=M071G03
NODE=M071G03

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G03;LINKAGE=UB

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{80}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±1.3±0.3	32	AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ

NODE=M071G04
NODE=M071G04

$$\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{71}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.04±1.79±0.02	7	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ

NODE=M071G05
NODE=M071G05

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G05;LINKAGE=UB

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{98}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M071G06
NODE=M071G06

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.147±0.035±0.005	66	¹ LEES	15J BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ
0.197±0.035±0.005	66	² LEES	15J BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ
0.35 ± 0.14 ± 0.03	11	³ LEES	13Q BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ

OCCUR=2

¹ sinφ > 0.

² sinφ < 0.

³ Interference with non-resonant K⁺K⁻ production not taken into account.

NODE=M071G06;LINKAGE=A
NODE=M071G06;LINKAGE=B
NODE=M071G06;LINKAGE=BA

$\psi(2S)$ BRANCHING RATIOS

NODE=M071235

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.9785 ± 0.0013 OUR AVERAGE			
0.9779 ± 0.0015	¹ BAI	02B	BES2 $e^+ e^-$
0.981 ± 0.003	¹ LUTH	75	MRK1 $e^+ e^-$

NODE=M071R3
NODE=M071R3¹ Includes cascade decay into $J/\psi(1S)$.

NODE=M071R;LINKAGE=P

 $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0173 ± 0.0014 OUR AVERAGE			Error includes scale factor of 1.5.
0.0166 ± 0.0010	^{1,2} SETH	04	RVUE $e^+ e^-$
0.0199 ± 0.0019	¹ BAI	02B	BES2 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.029 ± 0.004	¹ LUTH	75	MRK1 $e^+ e^-$

NODE=M071R5
NODE=M071R5¹ Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.NODE=M071R;LINKAGE=Z
NODE=M071R5;LINKAGE=SE² Using $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. $\Gamma(g g g)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
10.58 ± 1.62	2.9 M	¹ LIBBY	09	CLEO $\psi(2S) \rightarrow \text{hadrons}$

NODE=M071S43
NODE=M071S43¹ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow X J/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma \eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+ \ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

NODE=M071S43;LINKAGE=LI

 $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.025 ± 0.288	200 k	¹ LIBBY	09	CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$

NODE=M071S44
NODE=M071S44¹ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

NODE=M071S44;LINKAGE=LI

 $\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.7 ± 2.6 ± 1.6	2.9 M	LIBBY	09	CLEO $\psi(2S) \rightarrow (\gamma +) \text{hadrons}$

NODE=M071S45
NODE=M071S45 $\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.154 ± 0.015	¹ MENDEZ	08	CLEO $e^+ e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.169 ± 0.026	² ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S)$

NODE=M071S27
NODE=M071S27¹ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.

NODE=M071S27;LINKAGE=ME

² Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ} \gamma)$, $B(\eta_c \gamma)$ from ATHAR 04 and $B(\ell^+ \ell^-)$ from PDG 04. Superseded by MENDEZ 08.

NODE=M071S27;LINKAGE=AD

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
78.9 ± 1.7 OUR FIT			

NODE=M071R1
NODE=M071R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

88 ± 13	¹ FELDMAN	77	RVUE $e^+ e^-$
---------	----------------------	----	----------------

¹ From an overall fit assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

NODE=M071R;LINKAGE=L

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	DOCUMENT ID
79 ± 9 OUR FIT	

NODE=M071R2
NODE=M071R2

$\Gamma(\mu^+ \mu^-) / \Gamma(e^+ e^-)$

Γ_7 / Γ_6

NODE=M071R4
NODE=M071R4

VALUE DOCUMENT ID TECN COMMENT

1.00 ± 0.11 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89 ± 0.16 BOYARSKI 75C MRK1 $e^+ e^-$

$\Gamma(\tau^+ \tau^-) / \Gamma_{total}$

Γ_8 / Γ

NODE=M071R75
NODE=M071R75

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

31 ± 4 OUR FIT

30.8 ± 2.1 ± 3.8 ¹ ABLIKIM 06W BES $e^+ e^- \rightarrow \psi(2S)$

¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

NODE=M071R75;LINKAGE=AB

———— DECAYS INTO $J/\psi(1S)$ AND ANYTHING ————

NODE=M071305

$\Gamma(J/\psi(1S)\text{anything}) / \Gamma_{total}$

Γ_9 / Γ

NODE=M071R10
NODE=M071R10

VALUE EVTS DOCUMENT ID TECN COMMENT

0.610 ± 0.006 OUR FIT

0.55 ± 0.07 OUR AVERAGE

0.51 ± 0.12 BRANDELIK 79C DASP $e^+ e^- \rightarrow \mu^+ \mu^- X$

0.57 ± 0.08 ABRAMS 75B MRK1 $e^+ e^- \rightarrow \mu^+ \mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6254 ± 0.0016 ± 0.0155 1.1M ¹ MENDEZ 08 CLEO $\psi(2S) \rightarrow \ell^+ \ell^- X$

0.5950 ± 0.0015 ± 0.0190 151k ADAM 05A CLEO Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

NODE=M071R10;LINKAGE=ME

$\Gamma(e^+ e^-) / \Gamma(J/\psi(1S)\text{anything})$

$\Gamma_6 / \Gamma_9 = \Gamma_6 / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.339\Gamma_{133} + 0.192\Gamma_{134})$

NODE=M071R72

NODE=M071R72

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

1.294 ± 0.026 OUR FIT

[(1.295 ± 0.026) × 10^{-2} OUR 2015 FIT]

1.28 ± 0.04 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

1.22 ± 0.02 ± 0.05 5097 ± 73 ¹ ANDREOTTI 05 E835 $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+ e^-$

1.28 ± 0.03 ± 0.02 ¹ AMBROGIANI 00A E835 $p\bar{p} \rightarrow \psi(2S)$


1.44 ± 0.08 ± 0.02 ¹ ARMSTRONG 97 E760 $p\bar{p} \rightarrow \psi(2S)$

¹ Using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

NEW

NODE=M071R;LINKAGE=7A

WEIGHTED AVERAGE
1.28 ± 0.04 (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

			χ^2
ANDREOTTI	05	E835	1.3
AMBROGIANI	00A	E835	0.0
ARMSTRONG	97	E760	3.7
			5.0

(Confidence Level = 0.082)

$\Gamma(e^+ e^-) / \Gamma(J/\psi(1S)\text{anything})$ (units 10^{-2})

$\Gamma(\mu^+ \mu^-) / \Gamma(J/\psi(1S)\text{anything})$

$\Gamma_7 / \Gamma_9 = \Gamma_7 / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.339\Gamma_{133} + 0.192\Gamma_{134})$

NODE=M071R74

NODE=M071R74

VALUE DOCUMENT ID TECN COMMENT

0.0130 ± 0.0014 OUR FIT

0.014 ± 0.003

HILGER 75 SPEC $e^+ e^-$

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$

VALUE

DOCUMENT ID

 Γ_{10}/Γ

NODE=M071R18
 NODE=M071R18
 NEW

0.2514±0.0033 OUR FIT

[0.2511 ± 0.0033 OUR 2015 FIT]

 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{11}/Γ

NODE=M071R12
 NODE=M071R12
 NEW

0.3449±0.0030 OUR FIT

[0.3446 ± 0.0030 OUR 2015 FIT]

0.348 ±0.005 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

0.3498±0.0002±0.0045 20M

ABLIKIM

13R

BES3

 $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$

0.3504±0.0007±0.0077 565k

MENDEZ

08

CLEO

 $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$

0.323 ±0.014

BAI

02B

BES2

 e^+e^-

0.32 ±0.04

ABRAMS

75B


MRK1

 $e^+e^- \rightarrow J/\psi\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3354±0.0014±0.0110 60k ¹ADAM 05A CLEO Repl. by MENDEZ 08¹Not independent from other values reported by ADAM 05A.

NODE=M071R;LINKAGE=AD

 $\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE

DOCUMENT ID

TECN

COMMENT

 Γ_6/Γ_{11}

NODE=M071R73
 NODE=M071R73

0.0229±0.0005 OUR FIT**0.0252±0.0028±0.0011**¹AUBERT

02B

BABR

 e^+e^- ¹Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

NODE=M071R73;LINKAGE=7A

 $\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE

DOCUMENT ID

TECN

COMMENT

 Γ_7/Γ_{11}

NODE=M071R63
 NODE=M071R63

0.0229±0.0025 OUR FIT**0.0224±0.0029 OUR AVERAGE**

0.0216±0.0026±0.0014

¹AUBERT

02B

BABR

 e^+e^-

0.0327±0.0077±0.0072

¹GRIBUSHIN

96

FMPS

515 $\pi^- \text{Be} \rightarrow 2\mu X$ ¹Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

NODE=M071R;LINKAGE=Q2

 $\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ VALUE (units 10^{-3})

DOCUMENT ID

TECN

COMMENT

 Γ_8/Γ_{11}

NODE=M071R76
 NODE=M071R76

8.9 ±1.1 OUR FIT**8.73±1.39±1.57**

BAI

02

BES

 e^+e^-

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$ Γ_{11}/Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.5653±0.0026 OUR FIT

[0.5654 ± 0.0026 OUR 2015 FIT]

0.554 ± 0.008 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

0.5604±0.0009±0.0062	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$
0.525 ± 0.009 ± 0.022	4k	ANDREOTTI	05	E835 $\psi(2S) \rightarrow J/\psi X$
0.536 ± 0.007 ± 0.016	20k	^{1,2} ABLIKIM	04B	BES $\psi(2S) \rightarrow J/\psi X$
0.496 ± 0.037		ARMSTRONG	97	E760 $\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •


0.5637±0.0027±0.0046 60k ADAM 05A CLEO Repl. by MENDEZ 08

¹ From a fit to the J/ψ recoil mass spectra.² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$.

NODE=M071R70
 NODE=M071R70
 NEW

NODE=M071R;LINKAGE=AB
 NODE=M071R;LINKAGE=AL

WEIGHTED AVERAGE
 0.554±0.008 (Error scaled by 1.3)

 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$ Γ_{11}/Γ_9 $\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$ $\Gamma_{10}/\Gamma_{11} = (0.9761\Gamma_{12} + 0.719\Gamma_{13} + 0.339\Gamma_{133} + 0.192\Gamma_{134})/\Gamma_{11}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.729±0.008 OUR FIT**0.73 ± 0.09**TANENBAUM 76 MRK1 e^+e^-

NODE=M071R11
 NODE=M071R11

 $\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.1816±0.0031 OUR FIT

[0.1814 ± 0.0031 OUR 2015 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1769±0.0008±0.0053	61k	¹ MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$
0.1652±0.0014±0.0058	13.4k	² ADAM	05A	CLEO Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.² Not independent from other values reported by ADAM 05A.

NODE=M071R17
 NODE=M071R17
 NEW

NODE=M071R17;LINKAGE=ME
 NODE=M071R17;LINKAGE=AD

 $\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$ Γ_{12}/Γ_9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.2977±0.0031 OUR FIT

[0.2976 ± 0.0031 OUR 2015 FIT]

0.320 ± 0.012 OUR AVERAGE

0.300 ± 0.008 ± 0.022	1655 ± 44	ANDREOTTI	05	E835 $\psi(2S) \rightarrow J/\psi X$
0.328 ± 0.013 ± 0.008		AMBROGIANI	00A	E835 $p\bar{p} \rightarrow \psi(2S)$
0.323 ± 0.033		ARMSTRONG	97	E760 $\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2829±0.0012±0.0056	61k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-2\pi^0$
0.2776±0.0025±0.0043	13.4k	ADAM	05A	CLEO Repl. by MENDEZ 08

NODE=M071R69
 NODE=M071R69
 NEW

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{12}/Γ_{11}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.527 ± 0.008 OUR FIT				
[0.526 ± 0.008 OUR 2015 FIT]				
0.513 ± 0.022 OUR AVERAGE				Error includes scale factor of 2.2.
0.5047 ± 0.0022 ± 0.0102	61k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.570 ± 0.009 ± 0.026	14k	¹ ABLIKIM	04B	BES $\psi(2S) \rightarrow J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.4924 ± 0.0047 ± 0.0086	73k	^{2,3} ADAM	05A	CLEO Repl. by MENDEZ 08
0.571 ± 0.018 ± 0.044		⁴ ANDREOTTI	05	E835 $\psi(2S) \rightarrow J/\psi X$
0.53 ± 0.06		TANENBAUM	76	MRK1 $e^+ e^-$
0.64 ± 0.15		⁵ HILGER	75	SPEC $e^+ e^-$

NODE=M071R14
 NODE=M071R14
 NEW

- ¹ From a fit to the J/ψ recoil mass spectra.
- ² Not independent from other values reported by ADAM 05A.
- ³ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.
- ⁴ Not independent from other values reported by ANDREOTTI 05.
- ⁵ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

NODE=M071R14;LINKAGE=AB
 NODE=M071R14;LINKAGE=AD
 NODE=M071R14;LINKAGE=AM
 NODE=M071R;LINKAGE=AN
 NODE=M071R;LINKAGE=I

$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$


Γ_{13}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
33.6 ± 0.5 OUR FIT				
32.9 ± 1.7 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
33.75 ± 0.17 ± 0.86	68.2k	ABLIKIM	12M	BES3 $e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
29.8 ± 0.9 ± 2.3	5.7k	BAI	04I	BES2 $\psi(2S) \rightarrow J/\psi\gamma\gamma$
25.5 ± 2.9	386	¹ OREGLIA	80	CBAL $e^+ e^- \rightarrow J/\psi 2\gamma$
45 ± 12	17	² BRANDELIK	79B	DASP $e^+ e^- \rightarrow J/\psi 2\gamma$
42 ± 6	164	² BARTEL	78B	CNTR $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
34.3 ± 0.4 ± 0.9	18.4k	³ MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- \eta$
32.5 ± 0.6 ± 1.1	2.8k	⁴ ADAM	05A	CLEO Repl. by MENDEZ 08
43 ± 8	44	TANENBAUM	76	MRK1 $e^+ e^-$

NODE=M071R15
 NODE=M071R15

- ¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.
- ² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.
- ³ Not independent from other measurements of MENDEZ 08.
- ⁴ Not independent from other values reported by ADAM 05A.

NODE=M071R;LINKAGE=3Q
 NODE=M071R;LINKAGE=2Q
 NODE=M071R15;LINKAGE=ME
 NODE=M071R15;LINKAGE=AD



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{13}/Γ_9

VALUE EVTS DOCUMENT ID TECN COMMENT

NODE=M071R68
NODE=M071R68

0.0551±0.0008 OUR FIT

0.058 ±0.007 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

0.050 ±0.006 ±0.003	298 ± 20	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
0.072 ±0.009		AMBROGIANI 00A	E835	$\rho\bar{\rho} \rightarrow \psi(2S)$
0.061 ±0.015		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$


OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

0.0549±0.0006±0.0009	18.4k	¹ MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.0546±0.0010±0.0007	2.8k	ADAM 05A	CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

NODE=M071R68;LINKAGE=ME



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{13}/Γ_{11}

VALUE EVTS DOCUMENT ID TECN COMMENT

NODE=M071R71
NODE=M071R71
NEW

0.0974±0.0014 OUR FIT

[0.0975 ± 0.0014 OUR 2015 FIT]

0.0979±0.0018 OUR AVERAGE

0.0979±0.0010±0.0015	18.4k	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.098 ±0.005 ±0.010	2k	¹ ABLIKIM 04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ±0.021		² HIMEL 80	MRK2	$e^+ e^- \rightarrow \psi(2S) X$

••• We do not use the following data for averages, fits, limits, etc. •••

0.0968±0.0019±0.0013	2.8k	³ ADAM 05A	CLEO	Repl. by MENDEZ 08
0.095 ±0.007 ±0.007		⁴ ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

⁴ Not independent from other values reported by ANDREOTTI 05.

NODE=M071R71;LINKAGE=AB
NODE=M071R;LINKAGE=8H

NODE=M071R71;LINKAGE=AD
NODE=M071R71;LINKAGE=AN

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{total}$

Γ_{14}/Γ

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

NODE=M071R16
NODE=M071R16

12.68±0.32 OUR AVERAGE

12.6 ±0.2 ±0.3	4.1k	ABLIKIM 12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
13.3 ±0.8 ±0.3	530	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$
14.3 ±1.4 ±1.2	280	BAI 04i	BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
14 ±6	7	HIMEL 80	MRK2	$e^+ e^-$
9 ±2 ±1	23	¹ OREGLIA 80	CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

13 ±1 ±1	88	ADAM 05A	CLEO	Repl. by MENDEZ 08
----------	----	----------	------	--------------------

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

NODE=M071R16;LINKAGE=3Q

$$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$$

$$\Gamma_{14}/\Gamma_9 = \Gamma_{14}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13}+0.339\Gamma_{133}+0.192\Gamma_{134})$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.213 \pm 0.012 \pm 0.003$	527	¹ MENDEZ	08	CLEO $e^+e^- \rightarrow J/\psi\gamma\gamma$
$0.22 \pm 0.02 \pm 0.01$		² ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

NODE=M071S25
NODE=M071S25

NODE=M071S25;LINKAGE=ME
NODE=M071S25;LINKAGE=AD

$$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{14}/\Gamma_{11}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.380 \pm 0.022 \pm 0.005$	527	¹ MENDEZ	08	CLEO $e^+e^- \rightarrow J/\psi\gamma\gamma$
$0.39 \pm 0.04 \pm 0.01$		² ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

NODE=M071S26
NODE=M071S26

NODE=M071S26;LINKAGE=ME
NODE=M071S26;LINKAGE=AD

HADRONIC DECAYS

$$\Gamma(\pi^0 h_c(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{15}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.6 ± 1.3 OUR AVERAGE

$9.0 \pm 1.5 \pm 1.3$	3k	¹ GE	11	CLEO $\psi(2S) \rightarrow \pi^0$ anything
$8.4 \pm 1.3 \pm 1.0$	11k	ABLIKIM	10B	BES3 $\psi(2S) \rightarrow \pi^0 h_c$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	92^{+23}_{-22}	ADAMS	09	CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$
seen	1282	DOBBS	08A	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
seen	168 ± 40	ROSNER	05	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86 \text{ MeV} \equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P))/\Gamma_0) \times 10^{-4}$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

NODE=M071S42
NODE=M071S42

NODE=M071310

NODE=M071S42;LINKAGE=GE

$$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$$

$$\Gamma_{16}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

35 ± 16	6	FRANKLIN	83	MRK2 $e^+e^- \rightarrow \text{hadrons}$
----------------	---	----------	----	--

NODE=M071R37
NODE=M071R37

$$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$$

$$\Gamma_{17}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

29 ± 10 OUR AVERAGE Error includes scale factor of 4.7. See the ideogram below.
 $[(29 \pm 10) \times 10^{-4} \text{ OUR 2015 AVERAGE Scale factor} = 4.7]$


$24.9 \pm 0.7 \pm 3.6$	2173	ABLIKIM	07D	BES2 $e^+e^- \rightarrow \psi(2S)$
$127 \pm 12 \pm 2$	410	¹ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
$26.1 \pm 0.7 \pm 3.0$	1703	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$
30 ± 8	42	FRANKLIN	83	MRK2 e^+e^-

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (297 \pm 22 \pm 18) \times 10^{-4} \text{ keV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.34 \pm 0.04 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R22
NODE=M071R22

NEW

NODE=M071R22;LINKAGE=UB



$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$

Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.55±0.73±0.47		112 ± 31	BAI	04c BES2	$\psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$
<2.3	90		BAI	98J BES	e^+e^-

NODE=M071R65
NODE=M071R65

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$

Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.88±0.09 OUR FIT				
[(2.87 ± 0.09) × 10 ⁻⁴ OUR 2015 FIT]				
3.00±0.13 OUR AVERAGE		Error includes scale factor of 1.1.		
3.08±0.05±0.18	4.5k	¹ DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.36±0.09±0.25	1.6k	ABLIKIM	07C BES	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.87±0.12±0.15	557	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
1.4 ±0.8	4	BRANDELIK	79C DASP	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.3 ±0.7		FELDMAN	77 MRK1	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

NODE=M071R25
NODE=M071R25
NEW

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071R25;LINKAGE=A

$\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{19}/Γ_{11}

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.35±0.28 OUR FIT			
[(8.34 ± 0.28) × 10 ⁻⁴ OUR 2015 FIT]			
6.98±0.49±0.97	BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

NODE=M071S40
NODE=M071S40
NEW

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$

Γ_{20}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
12.8±1.0±3.4	157	¹ BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

NODE=M071R50
NODE=M071R50

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

NODE=M071R50;LINKAGE=PP

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.29	90	¹ ABLIKIM	13F BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
<12	90	² ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R6
NODE=M071R6

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

NODE=M071R6;LINKAGE=AL

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

NODE=M071R6;LINKAGE=AB

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.48±0.34±0.19		60	¹ ABLIKIM	13F BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

NODE=M071R7
 NODE=M071R7

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.9 90 ² ABLIKIM 07H BES2 $e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

NODE=M071R7;LINKAGE=AL
 NODE=M071R7;LINKAGE=AB

 $\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.1±0.1	74.0	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^-$

NODE=M071S18
 NODE=M071S18

 $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.8±0.3±0.3	45.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$

NODE=M071S19
 NODE=M071S19

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8±0.4±0.5	73.4	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$

NODE=M071S17
 NODE=M071S17

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.57±0.18 OUR AVERAGE					

NODE=M071R28
 NODE=M071R28

3.75±0.09±0.23	1.9k	¹ DOBBS	14		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
3.39±0.20±0.32	337	ABLIKIM	07C BES		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
6.4 ±1.8 ±0.1		² AUBERT	07BD BABR	10.6	$e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
3.28±0.23±0.25	208	PEDLAR	05 CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.81±0.20±0.27 80 ³ BAI 01 BES $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
 < 4 90 FELDMAN 77 MRK1 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

NODE=M071R28;LINKAGE=A
 NODE=M071R28;LINKAGE=AU

NODE=M071R28;LINKAGE=PP

 $\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.40±0.03±0.13	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

NODE=M071S65
 NODE=M071S65

 $\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.54±0.04±0.13	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$

NODE=M071S66
 NODE=M071S66

 $\Gamma(\Sigma^0\bar{p}K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67±0.13±0.12	276	¹ ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$

NODE=M071S63
 NODE=M071S63

¹ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$.

NODE=M071S63;LINKAGE=AB

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.51±0.21 OUR AVERAGE					
2.51±0.15±0.16	281	¹ DOBBS	14		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.57±0.44±0.68	35	PEDLAR	05 CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071R47
 NODE=M071R47

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071R47;LINKAGE=A

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.32±0.16 OUR AVERAGE				
2.25±0.11±0.16	439	¹ DOBBS 14		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.35±0.36±0.32	59	ABLIKIM 07C	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.63±0.35±0.21	58	PEDLAR 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 ±0.4 ±0.4	8	² BAI 01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
---------------	---	---------------------	-----	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

Γ₃₁/Γ
NODE=M071R51
NODE=M071R51

NODE=M071R51;LINKAGE=A
NODE=M071R51;LINKAGE=PP

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
11±3±3	14	¹ BAI 01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

Γ₃₂/Γ
NODE=M071R52
NODE=M071R52

NODE=M071R52;LINKAGE=PP

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.64±0.18 OUR AVERAGE					
2.66±0.12±0.20	548	¹ DOBBS 14		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
3.03±0.40±0.32	67	ABLIKIM 07C	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
2.38±0.30±0.21	63	PEDLAR 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.94±0.27±0.15	12	² BAI 01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<2	90	FELDMAN 77	MRK1	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

Γ₃₃/Γ
NODE=M071R29
NODE=M071R29

NODE=M071R29;LINKAGE=A
NODE=M071R29;LINKAGE=PP

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.07±0.23 OUR AVERAGE				
2.02±0.19±0.15	112	¹ DOBBS 14		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.75±0.64±0.61	19	PEDLAR 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

Γ₃₄/Γ
NODE=M071R48
NODE=M071R48

NODE=M071R48;LINKAGE=A

 $\Gamma(\Xi(1530)^0 \bar{\Xi}(1530)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.2±0.3^{+3.2}_{-1.2}	527	¹ ABLIKIM 13S	BES3	$\psi(2S) \rightarrow \eta p \bar{p}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32	90	PEDLAR 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 8.1	90	² BAI 01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ With $N(1535)$ decaying to $p\eta$.

² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

Γ₃₅/Γ
NODE=M071R53
NODE=M071R53

NODE=M071R53;LINKAGE=A
NODE=M071R53;LINKAGE=PP

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.86±0.27±0.32	236	ABLIKIM 15I	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$

Γ₃₆/Γ
NODE=M071S82
NODE=M071S82

 $\Gamma(\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
5.21±1.48±0.57	74	ABLIKIM 15I	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$

Γ₃₇/Γ
NODE=M071S83
NODE=M071S83

 $\Gamma(\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
12.03±2.94±1.22	136	ABLIKIM 15I	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$

Γ₃₈/Γ
NODE=M071S84
NODE=M071S84

$\Gamma(K^- \Sigma^0 \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{39}/Γ
$3.67 \pm 0.33 \pm 0.28$	142	ABLIKIM	15I	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Sigma^0 \Xi^+ + \text{c.c.}$	

NODE=M071S85
NODE=M071S85

 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{40}/Γ
$0.47 \pm 0.09 \pm 0.05$		27	¹ DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

NODE=M071R54
NODE=M071R54

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	90	ABLIKIM	12Q	BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<1.6	90	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<0.73	90	² BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

NODE=M071R54;LINKAGE=A
NODE=M071R54;LINKAGE=PP

 $\Gamma(\pi^0 p \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{41}/Γ
1.53 ± 0.07 OUR AVERAGE					
$1.65 \pm 0.03 \pm 0.15$	4.5k	ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$	
$1.54 \pm 0.06 \pm 0.06$	948	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \pi^0 p \bar{p}$	
$1.32 \pm 0.10 \pm 0.15$	256	¹ ABLIKIM	05E	BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \gamma \gamma$	
1.4 ± 0.5	9	FRANKLIN	83	MRK2 $e^+ e^-$	

NODE=M071R35
NODE=M071R35

¹ Computed using $B(\pi^0 \rightarrow \gamma \gamma) = (98.80 \pm 0.03)\%$.

NODE=M071R35;LINKAGE=AB

 $\Gamma(N(940) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{42}/Γ
$6.42 \pm 0.20^{+1.78}_{-1.28}$	1.9k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$	

NODE=M071S56
NODE=M071S56

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

NODE=M071S56;LINKAGE=AB

 $\Gamma(N(1440) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{43}/Γ
$7.3^{+1.7}_{-1.5}$ OUR AVERAGE				Error includes scale factor of 2.5.	

NODE=M071S50
NODE=M071S50

$3.58 \pm 0.25^{+1.59}_{-0.84}$	1.1k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$
$8.1 \pm 0.7 \pm 0.3$	474	² ALEXANDER	10	CLEO $\psi(2S) \rightarrow \pi^0 p \bar{p}$

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

² From a fit of the $p \bar{p}$ and $p \pi^0$ mass distributions to a combination of $N(1440) \bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

NODE=M071S50;LINKAGE=AB
NODE=M071S50;LINKAGE=AL

 $\Gamma(N(1520) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{44}/Γ
$0.64 \pm 0.05^{+0.22}_{-0.17}$	0.2k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$	

NODE=M071S57
NODE=M071S57

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

NODE=M071S57;LINKAGE=AB

 $\Gamma(N(1535) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{45}/Γ
$2.47 \pm 0.28^{+0.99}_{-0.97}$	0.7k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$	

NODE=M071S58
NODE=M071S58

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

NODE=M071S58;LINKAGE=AB

 $\Gamma(N(1650) \bar{p} + \text{c.c.} \rightarrow \pi^0 p \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{46}/Γ
$3.76 \pm 0.28^{+1.37}_{-1.66}$	1.1k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p \bar{p} \pi^0$	

NODE=M071S59
NODE=M071S59

¹ From a fit of $\pi^0 p \bar{p}$ data to eight distinct intermediate $N \bar{p}$ resonant states.

NODE=M071S59;LINKAGE=AB

$\Gamma(N(1720)\bar{p} + c.c. \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.79 \pm 0.10^{+0.24}_{-0.71}$	0.5k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow \rho\bar{p}\pi^0$

NODE=M071S60
NODE=M071S60¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

NODE=M071S60;LINKAGE=AB

 $\Gamma(N(2300)\bar{p} + c.c. \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.62 \pm 0.28^{+1.12}_{-0.64}$	0.9k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow \rho\bar{p}\pi^0$

NODE=M071S61
NODE=M071S61¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

NODE=M071S61;LINKAGE=AB

 $\Gamma(N(2570)\bar{p} + c.c. \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.13 \pm 0.08^{+0.40}_{-0.30}$	0.8k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow \rho\bar{p}\pi^0$

NODE=M071S62
NODE=M071S62¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

NODE=M071S62;LINKAGE=AB

 $\Gamma(\pi^0 f_0(2100) \rightarrow \pi^0 \rho\bar{p})/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.4 \pm 0.1$	76	¹ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 \rho\bar{p}$

NODE=M071S51
NODE=M071S51¹ From a fit of the $\rho\bar{p}$ and $\rho\pi^0$ mass distributions to a combination of $N_1^*(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

NODE=M071S51;LINKAGE=AL

 $\Gamma(\eta\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.4 OUR AVERAGE				
$6.4 \pm 0.2 \pm 0.6$	679	¹ ABLIKIM	13S BES3	$\psi(2S) \rightarrow \eta\rho\bar{p}$
$5.6 \pm 0.6 \pm 0.3$	154	¹ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta\rho\bar{p}$
$5.8 \pm 1.1 \pm 0.7$	44.8 ± 8.5	² ABLIKIM	05E BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ $\rho\bar{p}\gamma\gamma$
$8 \pm 3 \pm 3$	9.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ $\rho\bar{p}\pi^+ \pi^- \pi^0$

NODE=M071R56
NODE=M071R56¹ With $N(1535)$ decaying to $\rho\eta$.² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$.NODE=M071R56;LINKAGE=A
NODE=M071R56;LINKAGE=AB $\Gamma(\eta f_0(2100) \rightarrow \eta\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.2 \pm 0.4 \pm 0.1$	31	¹ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta\rho\bar{p}$

NODE=M071S52
NODE=M071S52¹ From a fit of the $\rho\bar{p}$ and $\rho\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

NODE=M071S52;LINKAGE=AL

 $\Gamma(N(1535)\bar{p} \rightarrow \eta\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 0.6 \pm 0.3$	123	¹ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta\rho\bar{p}$

NODE=M071S53
NODE=M071S53¹ From a fit of the $\rho\bar{p}$ and $\rho\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

NODE=M071S53;LINKAGE=AL

 $\Gamma(\omega\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.69 ± 0.21 OUR AVERAGE				
$0.6 \pm 0.2 \pm 0.2$	21.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ $\rho\bar{p}\pi^+ \pi^- \pi^0$
$0.8 \pm 0.3 \pm 0.1$	14.9 ± 0.1	¹ BAI	03B BES	$\psi(2S) \rightarrow \rho\bar{p}\pi^+ \pi^- \pi^0$

NODE=M071R79
NODE=M071R79¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

NODE=M071R;LINKAGE=B3

 $\Gamma(\phi\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ $\rho\bar{p}K^+ K^-$

NODE=M071R82
NODE=M071R82

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.26	90	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \rho\bar{p}$
-------	----	------------------	---------	--

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

NODE=M071R82;LINKAGE=B3

$\Gamma(\pi^+ \pi^- \rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0±0.4 OUR AVERAGE				
5.9±0.2±0.4	904.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-$
8 ± 2		¹ TANENBAUM	78	MRK1 $e^+ e^-$

NODE=M071R31
 NODE=M071R31

¹ Assuming entirely strong decay.

NODE=M071R;LINKAGE=K

 $\Gamma(\rho\bar{\rho}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.48±0.17 OUR AVERAGE				
2.45±0.11±0.21	851	ABLIKIM	06I	BES2 $e^+ e^- \rightarrow \rho\pi^- X$
2.52±0.12±0.22	849	ABLIKIM	06I	BES2 $e^+ e^- \rightarrow \bar{\rho}\pi^+ X$

NODE=M071R01
 NODE=M071R01

OCCUR=2

 $\Gamma(\rho\bar{\rho}\pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.18±0.50±0.50				
135 ± 21		ABLIKIM	06I	BES2 $e^+ e^- \rightarrow \rho\pi^- \pi^0 X$

NODE=M071R02
 NODE=M071R02

 $\Gamma(\eta\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6				
90		BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M071S06
 NODE=M071S06

 $\Gamma(\eta\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.5±0.7±1.5				
		¹ BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.3±0.8±1.4	201.7	² BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow \gamma\gamma)$
8.1±1.4±1.6	50.0	² BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow 3\pi)$

NODE=M071S07
 NODE=M071S07

OCCUR=2

OCCUR=3

¹ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

² Not independent from other values reported by BRIERE 05.

NODE=M071S07;LINKAGE=BR
 NODE=M071S07;LINKAGE=BI

 $\Gamma(2(\pi^+ \pi^-)\eta)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.6±0.1				
16		¹ AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-)\eta\gamma$

NODE=M071S38
 NODE=M071S38

¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$.

NODE=M071S38;LINKAGE=UB

 $\Gamma(\eta'\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.5±1.6±1.3				
12.8		BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$

NODE=M071S08
 NODE=M071S08

 $\Gamma(\omega\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±1.2 OUR AVERAGE				
Error includes scale factor of 2.1. See the ideogram below.				
8.4±0.5±1.2	386	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$
12.2±2.2±0.7	37	¹ AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow \omega\pi^+\pi^-\gamma$
8.2±0.5±0.7	391	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0$
4.8±0.6±0.7	100 ± 22	² BAI	03B	BES $\psi(2S) \rightarrow 2(\pi^+ \pi^-)\pi^0$


NODE=M071R77
 NODE=M071R77

¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16 \text{ eV}$.

NODE=M071R77;LINKAGE=UB

² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

NODE=M071R77;LINKAGE=B3



$\Gamma(b_1^\pm \pi^\mp) / \Gamma_{total}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{65}/Γ
4.0 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.1.	
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$	
4.18 ^{+0.43} _{-0.42} ± 0.92	170	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$	
3.2 ± 0.6 ± 0.5	61 ± 11	^{1,2} BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
5.2 ± 0.8 ± 1.0		¹ BAI	99C BES	Repl. by BAI 03B	

¹ Assuming $B(b_1 \rightarrow \omega \pi) = 1$.

² Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R40
NODE=M071R40

NODE=M071R;LINKAGE=M1
NODE=M071R40;LINKAGE=B3

$\Gamma(b_1^0 \pi^0) / \Gamma_{total}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{66}/Γ
2.35^{+0.47}_{-0.42} ± 0.40	45	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$	

NODE=M071R21
NODE=M071R21

$\Gamma(\omega f_2(1270)) / \Gamma_{total}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{67}/Γ
2.2 ± 0.4 OUR AVERAGE						
2.3 ± 0.5 ± 0.4		57	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$	
2.05 ± 0.41 ± 0.38		62 ± 12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
<1.5	90		¹ BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
<1.7	90		BAI	98J BES	Repl. by BAI 03B	

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R64
NODE=M071R64

NODE=M071R64;LINKAGE=B3

$\Gamma(\pi^+ \pi^- K^+ K^-) / \Gamma_{total}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{68}/Γ
7.5 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.9. [(7.5 ± 0.9) × 10 ⁻⁴ OUR 2015 AVERAGE Scale factor = 1.9]	
10.9 ± 1.9 ± 0.2	85	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	
7.1 ± 0.3 ± 0.4	817.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$	
16 ± 4		² TANENBAUM	78 MRK1	$e^+ e^-$	

¹ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assuming entirely strong decay.

NODE=M071R24
NODE=M071R24

NEW

NODE=M071R24;LINKAGE=BE

NODE=M071R24;LINKAGE=K

$\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{69}/Γ
2.2±0.2±0.4	223.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$	NODE=M071S09 NODE=M071S09

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{70}/Γ
1.86±0.32±0.43	93 ± 16	BAI	04C	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$	NODE=M071R66 NODE=M071R66	
• • •				We do not use the following data for averages, fits, limits, etc. • • •		
<1.2	90	BAI	98J	BES $e^+ e^-$		

 $\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{71}/Γ
1.3±0.7±0.1	7	¹ AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$	NODE=M071S39 NODE=M071S39
		¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^- \eta)) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1$ eV.			NODE=M071S39;LINKAGE=UB

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{72}/Γ
10.0±2.5±1.8	65	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$	NODE=M071R09 NODE=M071R09

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{74}/Γ
10.0±1.8±2.1	¹ BAI	99C	BES $e^+ e^-$	NODE=M071R41 NODE=M071R41
	¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$			NODE=M071R;LINKAGE=M2

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{75}/Γ
2.20±0.25±0.37	83 ± 9	ABLIKIM	050	BES2 $e^+ e^- \rightarrow \psi(2S)$	NODE=M071R49 NODE=M071R49

 $\Gamma(\rho^0 p \bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{76}/Γ
0.5±0.1±0.2	61.1	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \pi^+ \pi^-$	NODE=M071S14 NODE=M071S14

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{77}/Γ
6.7±2.5	TANENBAUM 78	MRK1	$e^+ e^-$	NODE=M071R34 NODE=M071R34

 $\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{78}/Γ
2.4±0.6 OUR AVERAGE	Error includes scale factor of 2.2.				NODE=M071R27 NODE=M071R27
2.2±0.2±0.2	308	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$	
4.5±1.0		TANENBAUM 78	MRK1	$e^+ e^-$	

 $\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{79}/Γ
2.2±0.6 OUR AVERAGE	Error includes scale factor of 1.4.				NODE=M071R33 NODE=M071R33
2.0±0.2±0.4	285.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$	
4.2±1.5		TANENBAUM 78	MRK1	$e^+ e^-$	

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{80}/Γ
12.6±0.9 OUR AVERAGE					NODE=M071S10 NODE=M071S10
18.8±5.7±0.3	32	¹ AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$	
11.7±1.0±1.5	597	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
12.7±0.5±1.0	711.6	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

¹AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071S10;LINKAGE=UB

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-) / \Gamma_{\text{total}}$ Γ_{81} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.9±2.0±0.9	19	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S20
NODE=M071S20

 $\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{82} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6±1.3±1.8	238	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S21
NODE=M071S21

 $\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{83} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.6±2.2±1.7	133	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S22
NODE=M071S22

 $\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{84} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±2.2±1.4	78	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S23
NODE=M071S23

 $\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{85} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±1.3±1.2	125	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S24
NODE=M071S24

 $\Gamma(\eta K^+ K^-, \text{ no } \eta\phi) / \Gamma_{\text{total}}$ Γ_{86} / Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.08±0.29±0.25	0.3k	1	ABLIKIM	12L	BES3 $\psi(2S) \rightarrow K^+ K^- \gamma\gamma$
••• We do not use the following data for averages, fits, limits, etc. •••					
<13	90		BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S11
NODE=M071S11

¹ Excluding $\eta\phi$.

NODE=M071S11;LINKAGE=AB

 $\Gamma(\omega K^+ K^-) / \Gamma_{\text{total}}$ Γ_{87} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.62±0.11 OUR AVERAGE				Error includes scale factor of 1.1.
1.56±0.04±0.11	2.8k	ABLIKIM	14G	BES3 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
2.38±0.37±0.29	78	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.9 ±0.3 ±0.3	76.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.5 ±0.3 ±0.2	23	¹ BAI	03B	BES $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071R78
NODE=M071R78

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R78;LINKAGE=B3

 $\Gamma(\omega K^*(892)^+ K^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{88} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
20.7±2.6 OUR AVERAGE				
18.9±2.9±2.2	396	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
22.6±3.0±2.4	535	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

NODE=M071S67
NODE=M071S67

OCCUR=2

 $\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{89} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1 ±1.2 OUR AVERAGE				
6.39±1.50±0.78	128	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
5.86±1.61±0.83	143	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

NODE=M071S68
NODE=M071S68

OCCUR=2

 $\Gamma(\omega \bar{K}^*(892)^0 K^0) / \Gamma_{\text{total}}$ Γ_{90} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.8±2.5±1.6	356	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

NODE=M071S69
NODE=M071S69

 $\Gamma(\omega \bar{K}_2^*(1430)^0 K^0) / \Gamma_{\text{total}}$ Γ_{91} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.82±2.08±0.72	116	ABLIKIM	13M	BES3 $\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

NODE=M071S70
NODE=M071S70

$\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.60 ± 0.27 ± 0.24	109	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

NODE=M071S71
NODE=M071S71

NODE=M071S71;LINKAGE=AB

 $\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.20 ± 0.16	82	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.

NODE=M071S72
NODE=M071S72

NODE=M071S72;LINKAGE=AB

 $\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
0.302 ± 0.098 ± 0.027	22	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ Statistical significance 4.5 σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.

NODE=M071S73
NODE=M071S73

NODE=M071S73;LINKAGE=AB

 $\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
0.125 ± 0.070 ± 0.013	10	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ Statistical significance 3.2 σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.

NODE=M071S74
NODE=M071S74

NODE=M071S74;LINKAGE=AB

 $\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ± 2.0 OUR AVERAGE	Error	includes scale factor of 2.8.		

5.45 ± 0.42 ± 0.87 671 ABLIKIM 05H BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$

1.5 ± 1.0 ¹ TANENBAUM 78 MRK1 $e^+ e^-$

¹ Assuming entirely strong decay.

NODE=M071R32
NODE=M071R32

NODE=M071R32;LINKAGE=K

 $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3 ± 0.4 ± 0.6	434.9	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

NODE=M071S15
NODE=M071S15

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.5 ± 0.5 OUR AVERAGE			[(7.1 ± 0.5) × 10 ⁻⁵ OUR 2015 AVERAGE Scale factor = 1.5]		

7.48 ± 0.23 ± 0.39 1.3k ¹ METREVELI 12 $\psi(2S) \rightarrow K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.2 ± 1.5 ± 0.2 66 ^{2,3} LEES 15J BABR $e^+ e^- \rightarrow K^+ K^- \gamma$

8.3 ± 1.5 ± 0.2 66 ^{3,4} LEES 15J BABR $e^+ e^- \rightarrow K^+ K^- \gamma$

6.3 ± 0.6 ± 0.3 ⁵ DOBBS 06A CLEO $e^+ e^-$

10 ± 7 ⁵ BRANDELIK 79C DASP $e^+ e^-$

< 5 90 FELDMAN 77 MRK1 $e^+ e^-$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin\phi > 0$.

³ Using $\Gamma(\psi(2S) \rightarrow e^+ e^-) = (2.37 \pm 0.04) \text{ keV}$.

⁴ $\sin\phi < 0$.

⁵ Interference with non-resonant $K^+ K^-$ production not taken into account.

NODE=M071R23
NODE=M071R23

NEW

OCCUR=2

NODE=M071R23;LINKAGE=ME
NODE=M071R23;LINKAGE=A
NODE=M071R23;LINKAGE=B
NODE=M071R23;LINKAGE=C
NODE=M071R23;LINKAGE=BA

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.34 ± 0.33 OUR AVERAGE				

5.28 ± 0.25 ± 0.34 478 ± 23 ¹ METREVELI 12 $\psi(2S) \rightarrow K_S^0 K_L^0$

5.8 ± 0.8 ± 0.4 DOBBS 06A CLEO $e^+ e^-$

5.24 ± 0.47 ± 0.48 156 ± 14 ² BAI 04B BES2 $\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.

NODE=M071R87
NODE=M071R87

NODE=M071R87;LINKAGE=ME
NODE=M071R;LINKAGE=KZ

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ **Γ_{100}/Γ**


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.01±0.17 OUR AVERAGE				Error includes scale factor of 1.7. See the ideogram below.
2.14±0.03 ^{+0.12} _{-0.11}	7k	¹ ABLIKIM	12H BES3	$e^+e^- \rightarrow \psi(2S)$
1.81±0.18±0.19	260 ± 19	² ABLIKIM	05J BES2	$e^+e^- \rightarrow \psi(2S)$
1.88 ^{+0.16} _{-0.15} ±0.28	194	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
0.85±0.46	4	FRANKLIN	83 MRK2	$e^+e^- \rightarrow$ hadrons

NODE=M071R36
NODE=M071R36

¹ From $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.
² From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

NODE=M071R36;LINKAGE=AB

NODE=M071R;LINKAGE=AK



$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{total}$ **Γ_{101}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.94±0.25^{+1.15}_{-0.34}	¹ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$

NODE=M071R57
NODE=M071R57

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

NODE=M071R57;LINKAGE=AK

$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{total}$ **Γ_{102}/Γ**

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.32±0.12 OUR AVERAGE					Error includes scale factor of 1.8.
0.51±0.07±0.11			¹ ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
0.24 ^{+0.08} _{-0.07} ±0.02		22	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R26
NODE=M071R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1	FRANKLIN	83 MRK2	e^+e^-
<10	90		BARTEL	76 CNTR	e^+e^-
<10	90	²	ABRAMS	75 MRK1	e^+e^-

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.
² Final state $\rho^0\pi^0$.

NODE=M071R26;LINKAGE=AK
NODE=M071R;LINKAGE=N

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ **Γ_{103}/Γ**

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.78±0.26 OUR AVERAGE					
0.76±0.25±0.06		30	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-$
8 ± 5			BRANDELIK	79C DASP	e^+e^-

NODE=M071R20
NODE=M071R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90		DOBBS	06A CLEO	$e^+e^- \rightarrow \psi(2S)$
<5	90		FELDMAN	77 MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+\pi^-$ for continuum subtraction.

NODE=M071R20;LINKAGE=ME

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	¹ BAI	99C BES	$e^+ e^-$

NODE=M071R45
NODE=M071R45¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

NODE=M071R;LINKAGE=M3

 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.12 \pm 0.62^{+1.13}_{-0.61}$	251 ± 22	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S54
NODE=M071S54 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.07 \pm 0.16 \pm 0.26$	0.9k	ABLIKIM	12L BES3	BES3	$e^+ e^- \rightarrow \psi(2S)$
<8.9	90	1	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M071R38
NODE=M071R38 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.4 OUR AVERAGE					Error includes scale factor of 1.2.
$3.18 \pm 0.30^{+0.26}_{-0.31}$		0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
$2.9^{+1.3}_{-1.7} \pm 0.4$		9.6 ± 4.2	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
$1.3^{+1.0}_{-0.7} \pm 0.3$		7	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
<5.4	90		FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M071R39
NODE=M071R39 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9 ± 2.0 OUR AVERAGE				
$13.3^{+2.4}_{-2.8} \pm 1.7$	65.6 ± 9.0	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
$9.2^{+2.7}_{-2.2} \pm 0.9$	25	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R30
NODE=M071R30 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$ $\Gamma_{107}/\Gamma_{108}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.06 OUR AVERAGE			
$0.22^{+0.10}_{-0.14}$	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
$0.14^{+0.08}_{-0.06}$	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R46
NODE=M071R46 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.17 ± 0.29 OUR AVERAGE				Error includes scale factor of 1.7.
$2.44 \pm 0.96 \pm 0.04$	10 ± 4	^{1,2} AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
$0.9 \pm 0.2 \pm 0.1$	47.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
$1.5 \pm 0.2 \pm 0.2$	51.5 ± 8.3	³ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M071R80
NODE=M071R80

¹ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R80;LINKAGE=BE

NODE=M071R80;LINKAGE=UB
NODE=M071R80;LINKAGE=B3 $\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.68 ± 0.25 OUR AVERAGE				Error includes scale factor of 1.2. $[(0.68 \pm 0.24) \times 10^{-4}]$
$1.45 \pm 0.70 \pm 0.02$	6 ± 3	^{1,2} AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
$0.6 \pm 0.2 \pm 0.1$	18.4 ± 6.4	³ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M071R83
NODE=M071R83

NEW

¹ AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.34 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R83;LINKAGE=BE

$\Gamma(2(K^+ K^-)) / \Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{111} / Γ	
0.6 ± 0.1 ± 0.1	59.2	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$	NODE=M071S12 NODE=M071S12

$\Gamma(\phi K^+ K^-) / \Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{112} / Γ
0.70 ± 0.16 OUR AVERAGE					NODE=M071R81 NODE=M071R81
0.8 ± 0.2 ± 0.1	36.8	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
0.6 ± 0.2 ± 0.1	16.1 ± 5.0	¹ BAI	03B	BES	$\psi(2S) \rightarrow 2(K^+ K^-)$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R81;LINKAGE=B3

$\Gamma(2(K^+ K^-) \pi^0) / \Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{113} / Γ	
1.1 ± 0.2 ± 0.2	44.7	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-) \pi^0$	NODE=M071S13 NODE=M071S13

$\Gamma(\phi \eta) / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{114} / Γ
3.10 ± 0.31 OUR AVERAGE					NODE=M071R89 NODE=M071R89
3.14 ± 0.23 ± 0.23	0.2k	ABLIKIM	12L	BES3	$e^+ e^- \rightarrow \psi(2S)$
2.0 $^{+1.5}_{-1.1}$ ± 0.4	6	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.3 ± 1.1 ± 0.5	17	ABLIKIM	04K	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\phi \eta') / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{115} / Γ	
3.1 ± 1.4 ± 0.7	8	¹ ABLIKIM	04K	BES	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071R90 NODE=M071R90

¹ Calculated combining $\eta' \rightarrow \gamma \rho$ and $\eta \pi^+ \pi^-$ channels.

NODE=M071R;LINKAGE=AI

$\Gamma(\omega \eta') / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{116} / Γ	
3.2 $^{+2.4}_{-2.0}$ ± 0.7	4	¹ ABLIKIM	04K	BES	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071R91 NODE=M071R91

¹ Calculated combining $\eta' \rightarrow \gamma \rho$ and $\eta \pi^+ \pi^-$ channels.

NODE=M071R91;LINKAGE=AI

$\Gamma(\omega \pi^0) / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{117} / Γ
2.1 ± 0.6 OUR AVERAGE					NODE=M071R92 NODE=M071R92
2.5 $^{+1.2}_{-1.0}$ ± 0.2	14	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$
1.87 $^{+0.68}_{-0.62}$ ± 0.28	14	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\rho \eta') / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{118} / Γ	
1.87 $^{+1.64}_{-1.11}$ ± 0.33	2	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071R93 NODE=M071R93

$\Gamma(\rho \eta) / \Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{119} / Γ
2.2 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.1.	NODE=M071R94 NODE=M071R94
3.0 $^{+1.1}_{-0.9}$ ± 0.2	18	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$
1.78 $^{+0.67}_{-0.62}$ ± 0.17	13	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					Γ_{120}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.1	90	ADAM	05	CLEO	$e^+e^- \rightarrow \psi(2S)$	NODE=M071R95 NODE=M071R95
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<3.1	90	ABLIKIM	04K	BES	$e^+e^- \rightarrow \psi(2S)$	
$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$					Γ_{121}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.04	90	ABLIKIM	12L	BES3	$e^+e^- \rightarrow \psi(2S)$	NODE=M071R96 NODE=M071R96
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.7	90	ADAM	05	CLEO	$e^+e^- \rightarrow \psi(2S)$	
<0.4	90	ABLIKIM	04K	BES	$e^+e^- \rightarrow \psi(2S)$	
$\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{122}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.0	90	PEDLAR	07	CLEO	$e^+e^- \rightarrow \psi(2S)$	NODE=M071R03 NODE=M071R03
$\Gamma(\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$					Γ_{123}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
$2.7 \pm 0.6 \pm 0.4$	30.1	BRIERE	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}K^+K^-$	NODE=M071S16 NODE=M071S16
$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{124}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.81 \pm 0.11 \pm 0.14$	50	¹ ABLIKIM	08C	BES2	$e^+e^- \rightarrow J/\psi$	NODE=M071R08 NODE=M071R08
¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.						
$\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$					Γ_{125}/Γ	
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.44 \pm 0.12 \pm 0.11$	20 ± 6	BAI	04C		$\psi(2S) \rightarrow 2(K^+K^-)$	NODE=M071R67 NODE=M071R67
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.45	90	BAI	98J	BES	$e^+e^- \rightarrow 2(K^+K^-)$	
$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0\rho K^-\bar{n} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{126}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.88	90	BAI	04G	BES2	e^+e^-	NODE=M071S01 NODE=M071S01
$\Gamma(\Theta(1540)K^-\bar{n} \rightarrow K_S^0\rho K^-\bar{n})/\Gamma_{\text{total}}$					Γ_{127}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.0	90	BAI	04G	BES2	e^+e^-	NODE=M071S02 NODE=M071S02
$\Gamma(\Theta(1540)K_S^0\bar{p} \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$					Γ_{128}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.70	90	BAI	04G	BES2	e^+e^-	NODE=M071S03 NODE=M071S03
$\Gamma(\bar{\Theta}(1540)K^+n \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$					Γ_{129}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<2.6	90	BAI	04G	BES2	e^+e^-	NODE=M071S04 NODE=M071S04
$\Gamma(\bar{\Theta}(1540)K_S^0\rho \rightarrow K_S^0\rho K^-\bar{n})/\Gamma_{\text{total}}$					Γ_{130}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.60	90	BAI	04G	BES2	e^+e^-	NODE=M071S05 NODE=M071S05
$\Gamma(K_S^0K_S^0)/\Gamma_{\text{total}}$					Γ_{131}/Γ	
VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT		
<0.046		¹ BAI	04D	BES	e^+e^-	NODE=M071R88 NODE=M071R88
¹ Forbidden by CP.						
						NODE=M071R;LINKAGE=BA

————— **RADIATIVE DECAYS** —————

 $\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$
 Γ_{132}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.99±0.27 OUR FIT
9.2 ±0.4 OUR AVERAGE

9.22±0.11±0.46	72600	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
9.9 ±0.5 ±0.8		¹ GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.2 ±2.3		¹ BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$
7.5 ±2.6		¹ WHITAKER	76	MRK1 e^+e^-

¹ Angular distribution $(1+\cos^2\theta)$ assumed.

NODE=M071315

NODE=M071R55
NODE=M071R55

NODE=M071R;LINKAGE=A

 $\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$
 Γ_{133}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.55±0.31 OUR FIT
8.9 ±0.5 OUR AVERAGE

9.07±0.11±0.54	76700	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
9.0 ±0.5 ±0.7		¹ GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.1 ±1.9		² BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$

¹ Angular distribution $(1-0.189 \cos^2\theta)$ assumed.

² Valid for isotropic distribution of the photon.
NODE=M071R58
NODE=M071R58NODE=M071R;LINKAGE=G
NODE=M071R;LINKAGE=B
 $\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$
 Γ_{134}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.11±0.31 OUR FIT
8.8 ±0.5 OUR AVERAGE Error includes scale factor of 1.1.

9.33±0.14±0.61	79300	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
8.0 ±0.5 ±0.7		¹ GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.0 ±2.0		² BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$

¹ Angular distribution $(1-0.052 \cos^2\theta)$ assumed.

² Valid for isotropic distribution of the photon.
NODE=M071R59
NODE=M071R59NODE=M071R;LINKAGE=F
NODE=M071R59;LINKAGE=B
 $[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]/\Gamma_{\text{total}} (\Gamma_{132} + \Gamma_{133} + \Gamma_{134})/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

27.6±0.3±2.0	¹ ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
--------------	--------------------	----	------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.
NODE=M071R19
NODE=M071R19

NODE=M071R;LINKAGE=AH

 $\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$
 $\Gamma_{132}/\Gamma_{133}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

1.02±0.01±0.07	¹ ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
----------------	--------------------	----	------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.
NODE=M071R97
NODE=M071R97

NODE=M071R97;LINKAGE=AH

 $\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$
 $\Gamma_{134}/\Gamma_{133}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

1.03±0.02±0.03	¹ ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
----------------	--------------------	----	------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.
NODE=M071R98
NODE=M071R98

NODE=M071R98;LINKAGE=AH

 $\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$
 $\Gamma_{132}/\Gamma_{134}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.99±0.02±0.08	¹ ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
----------------	--------------------	----	------------------------------------

¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.
NODE=M071R99
NODE=M071R99

NODE=M071R99;LINKAGE=AH

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{135}/Γ


VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.05	OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
0.432 ± 0.016 ± 0.060		MITCHELL	09	CLEO $e^+e^- \rightarrow \gamma X$
0.32 ± 0.04 ± 0.06	2560	¹ ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
0.28 ± 0.06		² GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$

¹ ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.

² GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.

NODE=M071R60
NODE=M071R60

NODE=M071R60;LINKAGE=AT
NODE=M071R60;LINKAGE=GA

 $\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{136}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
7 ± 2 ± 4		¹ ABLIKIM	12G	BES3 $\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8	90	² CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K \bar{K} \pi$
< 20	90	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
20-130	95	EDWARDS	82C	CBAL $e^+e^- \rightarrow \gamma X$

¹ ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M071R62
NODE=M071R62

NODE=M071R62;LINKAGE=AB

NODE=M071R62;LINKAGE=CR

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{137}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.58 ± 0.40 ± 0.13		37	ABLIKIM	10F	BES3 $\psi(2S) \rightarrow \gamma\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 5	90		PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
< 5400	95		¹ LIBERMAN	75	SPEC e^+e^-
< 1×10^4	90		WIIK	75	DASP e^+e^-

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+\mu^-) = 0.0077$.

NODE=M071R42
NODE=M071R42

NODE=M071R;LINKAGE=U

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.23±0.06 OUR AVERAGE					
1.26±0.03±0.08		2226	¹ ABLIKIM	10F BES3	$\psi(2S) \rightarrow 3\gamma\pi^+\pi^-$, $2\gamma\pi^+\pi^-$
1.19±0.08±0.03			PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
1.24±0.27±0.15		23	ABLIKIM	06R BES2	$e^+e^- \rightarrow \psi(2S)$
1.54±0.31±0.20		~ 43	BAI	98F BES	$\psi(2S) \rightarrow \pi^+\pi^-2\gamma$, $\pi^+\pi^-3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 60	90		² BRAUNSCH...	77 DASP	e^+e^-
< 11	90		³ BARTEL	76 CNTR	e^+e^-

¹ Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

NODE=M071R44
NODE=M071R44

NODE=M071R44;LINKAGE=AB
NODE=M071R;LINKAGE=R
NODE=M071R;LINKAGE=C

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.73^{+0.29}_{-0.25} OUR AVERAGE Error includes scale factor of 1.8. $[(2.1 \pm 0.4) \times 10^{-4}$ OUR 2015 AVERAGE]				

2.84±0.15 ^{+0.03} _{-0.10}	1.9k	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
2.12±0.19±0.32		^{3,4} BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.08±0.19±0.33	200.6 ± 18.8	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
2.90±1.08±1.07	29.9 ± 11.1	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

NODE=M071R84
NODE=M071R84

NEW

OCCUR=2
OCCUR=3

NODE=M071R84;LINKAGE=A
NODE=M071R84;LINKAGE=B

NODE=M071R;LINKAGE=3B
NODE=M071R;LINKAGE=B9

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±1.0±1.4				
	175	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S75
NODE=M071S75

NODE=M071S75;LINKAGE=A

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
9.2±1.8±0.6				
	274	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi\pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi\pi) = (34.9 \pm 2.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S76
NODE=M071S76

NODE=M071S76;LINKAGE=A

NODE=M071S76;LINKAGE=B

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.3±0.8±0.1				
	136	^{1,2} DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S77
NODE=M071S77

NODE=M071S77;LINKAGE=A

NODE=M071S77;LINKAGE=B

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ± 0.6 OUR AVERAGE				
[(0.30 ± 0.13) × 10 ⁻⁴ OUR 2015 AVERAGE]				

3.6 ± 0.4 ± 0.5	290	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
3.01±0.41±1.24	35.6 ± 4.8	² BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

NODE=M071R85
NODE=M071R85

NEW

NODE=M071R85;LINKAGE=A
NODE=M071R85;LINKAGE=3B

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{145}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

6.6 ± 0.7 OUR AVERAGE[(0.60 ± 0.16) × 10⁻⁴ OUR 2015 AVERAGE]6.7 ± 0.6 ± 0.6 375 ¹ DOBBS 15 $\psi(2S) \rightarrow \gamma K \bar{K}$ 6.04 ± 0.90 ± 1.32 39.6 ± 5.9 ^{2,3} BAI 03C BES $\psi(2S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.6 90 6.8 ± 3.1 ^{2,3} BAI 03C BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Includes unknown branching fractions to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied the $K^+ K^-$ result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K \bar{K}$ result.³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R86

NODE=M071R86

NEW

OCCUR=2

NODE=M071R86;LINKAGE=A

NODE=M071R;LINKAGE=CK

NODE=M071R86;LINKAGE=3B

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	COMMENT
--------------------------	------	-------------	---------

4.8 ± 0.5 ± 0.9 373 ¹ DOBBS 15 $\psi(2S) \rightarrow \gamma \pi \pi$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S78

NODE=M071S78

NODE=M071S78;LINKAGE=A

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	COMMENT
--------------------------	------	-------------	---------

3.2 ± 0.6 ± 0.8 207 ¹ DOBBS 15 $\psi(2S) \rightarrow \gamma K \bar{K}$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S79

NODE=M071S79

NODE=M071S79;LINKAGE=A

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
-------	-----	-------------	---------

< **5.8 × 10⁻⁶** 90 ^{1,2} DOBBS 15 $\psi(2S) \rightarrow \gamma \pi \pi$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively.

NODE=M071S80

NODE=M071S80

NODE=M071S80;LINKAGE=A

NODE=M071S80;LINKAGE=DO

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
-------	-----	-------------	---------

< **9.5 × 10⁻⁶** 90 ^{1,2} DOBBS 15 $\psi(2S) \rightarrow \gamma K \bar{K}$ ¹ Using CLEO-c data but not authored by the CLEO Collaboration.² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively.

NODE=M071S81

NODE=M071S81

NODE=M071S81;LINKAGE=A

NODE=M071S81;LINKAGE=DO

 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.38 ± 0.48 ± 0.09 13 ¹ ABLIKIM 10F BES3 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2 90 PEDLAR 09 CLE3 $\psi(2S) \rightarrow \gamma X$ < 90 90 BAI 98F BES $\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$ < 200 90 YAMADA 77 DASP $e^+ e^- \rightarrow 3\gamma$ ¹ Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

NODE=M071R43

NODE=M071R43

NODE=M071R43;LINKAGE=AB

 $\Gamma(\gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.71 ± 1.25 ± 1.64 418 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M071R04

NODE=M071R04

 $\Gamma(\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< **0.9** 90 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3 90 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ < 1.2 90 ¹ SCHARRE 80 MRK1 $e^+ e^-$ ¹ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

NODE=M071R61

NODE=M071R61

OCCUR=2

NODE=M071R;LINKAGE=E

 $\Gamma(\gamma \eta(1405) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.36 ± 0.25 ± 0.05 10 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M071R05

NODE=M071R05

$\Gamma(\gamma\eta(1475) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.5	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

NODE=M071R06
 NODE=M071R06
 OCCUR=2

 $\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.88	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M071R07
 NODE=M071R07

 $\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
39.6 ± 2.8 ± 5.0	583	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S28
 NODE=M071S28

 $\Gamma(\gamma K^{*0} K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{160}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
37.0 ± 6.1 ± 7.2	237	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S29
 NODE=M071S29

 $\Gamma(\gamma K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
24.0 ± 4.5 ± 5.0	41	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S30
 NODE=M071S30

 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
25.6 ± 3.6 ± 3.6	115	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S31
 NODE=M071S31

 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{163}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
19.1 ± 2.7 ± 4.3	132	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S32
 NODE=M071S32

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9 ± 0.5 OUR AVERAGE				Error includes scale factor of 2.0.
4.18 ± 0.26 ± 0.18	348	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
2.9 ± 0.4 ± 0.4	142	ABLIKIM 07D	BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S33
 NODE=M071S33

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

NODE=M071S33;LINKAGE=AL

 $\Gamma(\gamma f_2(1950) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2 ± 0.2 ± 0.1	111	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M071S46
 NODE=M071S46

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

NODE=M071S46;LINKAGE=AL

 $\Gamma(\gamma f_2(2150) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.18 ± 0.03	73	¹ ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M071S47
 NODE=M071S47

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

NODE=M071S47;LINKAGE=AL

 $\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$ Γ_{167}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
4.57 ± 0.36 +1.77 -4.26		ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$

NODE=M071S48
 NODE=M071S48

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
<5.4	90	ABLIKIM 07D	BES	$\psi(2S) \rightarrow \gamma p\bar{p}$

$\Gamma(\gamma X \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{168}/Γ For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.

NODE=M071S49

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

NODE=M071S49

NODE=M071S49

 $\Gamma(\gamma \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{169}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 1.2 \pm 0.7$	17	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S34

NODE=M071S34

 $\Gamma(\gamma 2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$ Γ_{170}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<22	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S35

NODE=M071S35

 $\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{171}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<17	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S36

NODE=M071S36

 $\Gamma(\gamma K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{172}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S37

NODE=M071S37

 $\Gamma(\gamma \gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{173}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.1 \pm 0.6^{+0.8}_{-1.0}$	1.1k	ABLIKIM 120	BES3	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S55

NODE=M071S55

OTHER DECAYS

NODE=M071320

 $\Gamma(\text{invisible})/\Gamma(e^+ e^-)$ Γ_{174}/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.0	90	LEES 13I	BABR	$B \rightarrow K^{(*)} \psi(2S)$

NODE=M071S64

NODE=M071S64

 $\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M071240

For measurements involving $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

NODE=M071240

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

NODE=M071250

 $\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$ $a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

NODE=M071QAR

NODE=M071QAR

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
67^{+19}_{-13}	59k	¹ ARTUSO 09	CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.

NODE=M071QAR;LINKAGE=AR

 $b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

NODE=M071QBR

NODE=M071QBR

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
37^{+53}_{-47}	59k	¹ ARTUSO 09	CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

NODE=M071QBR;LINKAGE=AR

$\psi(2S)$ REFERENCES

NODE=M071

ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56774
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56787
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14G	PR D89 112006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55898
DOBBS	14	PL B739 90	S. Dobbs <i>et al.</i>	(NWES, WAYN)	REFID=56333
ABLIKIM	13A	PRL 110 022001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54834
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54879
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54920
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55386
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55402
ABLIKIM	13S	PR D88 032010	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55403
ABLIKIM	13W	PR D88 112007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55634
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54056
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54269
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54272
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54273
ABLIKIM	12L	PR D86 072011	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54739
ABLIKIM	12M	PR D86 092008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54740
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54742
ABLIKIM	12Q	CPC 36 1040	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54864
ANASHIN	12	PL B711 280	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=54038
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)	REFID=53960
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53348
ABLIKIM	10F	PRL 105 261801	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=53630
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=53233
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=53103
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=53124
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52579
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51636
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51725
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=51655
		Translated from ZETFP 85 429.			
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)	REFID=52717; ERROR=3
PEDLAR	07	PR D75 011102	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=51630
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51048
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51560
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT, BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
DOBBS	06A	PR D74 011105	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51158
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50757
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50758
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50760
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50451
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50497
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50785
PEDLAR	05	PR D72 051108	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=50808
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=50812
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50327
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50328
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49608
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49749
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49186

BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
AUBERT	02B	PR D65 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48548
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48578
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49171
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)	REFID=48632
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48003
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47939
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46340
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46554
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45416
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 Collab., E706 Collab.)	REFID=44739
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
		Translated from YAF 41 733.			
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22173
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22207
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
		Translated from YAF 34 1471.			
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22197
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22198
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)	REFID=22064
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG	REFID=22194
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)	REFID=22176
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)	REFID=22177
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)	REFID=22179
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)	REFID=22186
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)	REFID=22046
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC	REFID=22188
WIIK	75	Stanford Symp. 69	B.H. Wiik	(DESY)	REFID=22050

$\psi(3770)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M053

$\psi(3770)$ MASS (MeV)

NODE=M053M

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

NODE=M053M

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
3773.13 ± 0.35 OUR FIT Error includes scale factor of 1.1. [3773.15 ± 0.33 MeV OUR 2015 FIT]

NODE=M053M
NEW

3778.1 ± 1.2 OUR AVERAGE

3779.2	+1.8 -1.7	+0.6 -0.8	1	ANASHIN	12A	KEDR	$e^+e^- \rightarrow D\bar{D}$
3775.5	±2.4	±0.5	57	AUBERT	08B	BABR	$B \rightarrow D\bar{D}K$
3776	±5	±4	68	BRODZICKA	08	BELL	$B^+ \rightarrow D^0\bar{D}^0K^+$
3778.8	±1.9	±0.9		AUBERT	07BE	BABR	$e^+e^- \rightarrow D\bar{D}\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3772.0	±1.9		2,3	ABLIKIM	08D	BES2	$e^+e^- \rightarrow$ hadrons
3778.4	±3.0	±1.3	34	CHISTOV	04	BELL	Sup. by BRODZICKA 08

NODE=M053M;LINKAGE=AN
 NODE=M053M;LINKAGE=AB

¹ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

² Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

³ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

NODE=M053M;LINKAGE=NI

$m_{\psi(3770)} - m_{\psi(2S)}$

NODE=M053DM

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

NODE=M053DM

VALUE (MeV) DOCUMENT ID TECN COMMENT
87.04 ± 0.35 OUR FIT Error includes scale factor of 1.1. [87.04 ± 0.33 MeV OUR 2015 FIT]

NODE=M053DM
NEW


86.6 ± 0.7 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.

86.9	±0.4		4	ABLIKIM	07E	BES2	$e^+e^- \rightarrow$ hadrons
86.7	±0.7			ABLIKIM	06L	BES2	$e^+e^- \rightarrow$ hadrons
80	±2			SCHINDLER	80	MRK2	e^+e^-
86	±2		5	BACINO	78	DLCO	e^+e^-
88	±3			RAPIDIS	77	LGW	e^+e^-

⁴ BES-II $\psi(2S)$ mass subtracted (see ABLIKIM 06L).

⁵ SPEAR $\psi(2S)$ mass subtracted (see SCHINDLER 80).

NODE=M053DM;LINKAGE=AK
 NODE=M053DM;LINKAGE=S



$\psi(3770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.2\pm 1.0 OUR FIT				
27.5\pm 0.9 OUR AVERAGE				
24.9 $^{+4.6+0.5}_{-4.0-1.1}$		⁶ ANASHIN	12A KEDR	$e^+ e^- \rightarrow D\bar{D}$
30.4 \pm 8.5		^{7,8} ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
27 \pm 10 \pm 5	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$
28.5 \pm 1.2 \pm 0.2		⁸ ABLIKIM	07E BES2	$e^+ e^- \rightarrow$ hadrons
23.5 \pm 3.7 \pm 0.9		AUBERT	07BE BABR	$e^+ e^- \rightarrow D\bar{D}\gamma$
26.9 \pm 2.4 \pm 0.3		⁸ ABLIKIM	06L BES2	$e^+ e^- \rightarrow$ hadrons
24 \pm 5		⁸ SCHINDLER	80 MRK2	$e^+ e^-$
24 \pm 5		⁸ BACINO	78 DLCO	$e^+ e^-$
28 \pm 5		⁸ RAPIDIS	77 LGW	$e^+ e^-$

⁶ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

⁷ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

⁸ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

NODE=M053W

NODE=M053W

NODE=M053W;LINKAGE=AN

NODE=M053W;LINKAGE=AB

NODE=M053W;LINKAGE=NI

 $\psi(3770)$ DECAY MODES

In addition to the dominant decay mode to $D\bar{D}$, $\psi(3770)$ was found to decay into the final states containing the J/ψ (BAI 05, ADAM 06). ADAMS 06 and HUANG 06A searched for various decay modes with light hadrons and found a statistically significant signal for the decay to $\phi\eta$ only (ADAMS 06).

NODE=M053220;NODE=M053

NODE=M053

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $D\bar{D}$	(93 $^{+8}_{-9}$) %	S=2.0
Γ_2 $D^0 \bar{D}^0$	(52 $^{+4}_{-5}$) %	S=2.0
Γ_3 $D^+ D^-$	(41 ± 4) %	S=2.0
Γ_4 $J/\psi \pi^+ \pi^-$	$(1.93 \pm 0.28) \times 10^{-3}$	
Γ_5 $J/\psi \pi^0 \pi^0$	$(8.0 \pm 3.0) \times 10^{-4}$	
Γ_6 $J/\psi \eta$	$(9 \pm 4) \times 10^{-4}$	
Γ_7 $J/\psi \pi^0$	$< 2.8 \times 10^{-4}$	CL=90%
Γ_8 $e^+ e^-$	$(9.6 \pm 0.7) \times 10^{-6}$	S=1.3

DESIG=2

DESIG=5

DESIG=6

DESIG=4

DESIG=46

DESIG=47

DESIG=48

DESIG=1

Decays to light hadrons

Γ_9 $b_1(1235)\pi$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{10} $\phi\eta'$	$< 7 \times 10^{-4}$	CL=90%
Γ_{11} $\omega\eta'$	$< 4 \times 10^{-4}$	CL=90%
Γ_{12} $\rho^0\eta'$	$< 6 \times 10^{-4}$	CL=90%
Γ_{13} $\phi\eta$	$(3.1 \pm 0.7) \times 10^{-4}$	
Γ_{14} $\omega\eta$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{15} $\rho^0\eta$	$< 5 \times 10^{-4}$	CL=90%
Γ_{16} $\phi\pi^0$	$< 3 \times 10^{-5}$	CL=90%
Γ_{17} $\omega\pi^0$	$< 6 \times 10^{-4}$	CL=90%
Γ_{18} $\pi^+ \pi^- \pi^0$	$< 5 \times 10^{-6}$	CL=90%
Γ_{19} $\rho\pi$	$< 5 \times 10^{-6}$	CL=90%
Γ_{20} $K^+ K^-$	$< 2 \times 10^{-5}$	CL=90%
Γ_{21} $K^*(892)^+ K^- + \text{c.c.}$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{22} $K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$< 1.2 \times 10^{-3}$	CL=90%
Γ_{23} $K_S^0 K_L^0$	$< 1.2 \times 10^{-5}$	CL=90%
Γ_{24} $2(\pi^+ \pi^-)$	$< 1.12 \times 10^{-3}$	CL=90%
Γ_{25} $2(\pi^+ \pi^-)\pi^0$	$< 1.06 \times 10^{-3}$	CL=90%
Γ_{26} $2(\pi^+ \pi^- \pi^0)$	< 5.85 %	CL=90%
Γ_{27} $\omega \pi^+ \pi^-$	$< 6.0 \times 10^{-4}$	CL=90%
Γ_{28} $3(\pi^+ \pi^-)$	$< 9.1 \times 10^{-3}$	CL=90%
Γ_{29} $3(\pi^+ \pi^-)\pi^0$	< 1.37 %	CL=90%
Γ_{30} $3(\pi^+ \pi^-)2\pi^0$	< 11.74 %	CL=90%

NODE=M053;CLUMP=H

DESIG=20

DESIG=17

DESIG=16

DESIG=15

DESIG=8

DESIG=14

DESIG=13

DESIG=12

DESIG=11

DESIG=9

DESIG=10

DESIG=234

DESIG=19

DESIG=18

DESIG=3

DESIG=21

DESIG=22

DESIG=208

DESIG=24

DESIG=52

DESIG=55

DESIG=210

Г31	$\eta\pi^+\pi^-$	< 1.24	$\times 10^{-3}$	CL=90%	DESIG=23
Г32	$\pi^+\pi^-2\pi^0$	< 8.9	$\times 10^{-3}$	CL=90%	DESIG=206
Г33	$\rho^0\pi^+\pi^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=64
Г34	$\eta3\pi$	< 1.34	$\times 10^{-3}$	CL=90%	DESIG=25
Г35	$\eta2(\pi^+\pi^-)$	< 2.43	%	CL=90%	DESIG=53
Г36	$\eta\rho^0\pi^+\pi^-$	< 1.45	%	CL=90%	DESIG=221
Г37	$\eta'3\pi$	< 2.44	$\times 10^{-3}$	CL=90%	DESIG=26
Г38	$K^+K^-\pi^+\pi^-$	< 9.0	$\times 10^{-4}$	CL=90%	DESIG=27
Г39	$\phi\pi^+\pi^-$	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=28
Г40	$K^+K^-2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%	DESIG=207
Г41	$4(\pi^+\pi^-)$	< 1.67	%	CL=90%	DESIG=62
Г42	$4(\pi^+\pi^-)\pi^0$	< 3.06	%	CL=90%	DESIG=63
Г43	$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%	DESIG=29
Г44	$K^+K^-\pi^+\pi^-\pi^0$	< 2.36	$\times 10^{-3}$	CL=90%	DESIG=30
Г45	$K^+K^-\rho^0\pi^0$	< 8	$\times 10^{-4}$	CL=90%	DESIG=67
Г46	$K^+K^-\rho^+\pi^-$	< 1.46	%	CL=90%	DESIG=68
Г47	ωK^+K^-	< 3.4	$\times 10^{-4}$	CL=90%	DESIG=32
Г48	$\phi\pi^+\pi^-\pi^0$	< 3.8	$\times 10^{-3}$	CL=90%	DESIG=69
Г49	$K^{*0}K^-\pi^+\pi^0 + \text{c.c.}$	< 1.62	%	CL=90%	DESIG=70
Г50	$K^{*+}K^-\pi^+\pi^- + \text{c.c.}$	< 3.23	%	CL=90%	DESIG=71
Г51	$K^+K^-\pi^+\pi^-2\pi^0$	< 2.67	%	CL=90%	DESIG=209
Г52	$K^+K^-2(\pi^+\pi^-)$	< 1.03	%	CL=90%	DESIG=57
Г53	$K^+K^-2(\pi^+\pi^-)\pi^0$	< 3.60	%	CL=90%	DESIG=58
Г54	ηK^+K^-	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=31
Г55	$\eta K^+K^-\pi^+\pi^-$	< 1.24	%	CL=90%	DESIG=222
Г56	$\rho^0 K^+K^-$	< 5.0	$\times 10^{-3}$	CL=90%	DESIG=65
Г57	$2(K^+K^-)$	< 6.0	$\times 10^{-4}$	CL=90%	DESIG=33
Г58	ϕK^+K^-	< 7.5	$\times 10^{-4}$	CL=90%	DESIG=34
Г59	$2(K^+K^-)\pi^0$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=35
Г60	$2(K^+K^-)\pi^+\pi^-$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=59
Г61	$K_S^0 K^-\pi^+$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=200
Г62	$K_S^0 K^-\pi^+\pi^0$	< 1.33	%	CL=90%	DESIG=201
Г63	$K_S^0 K^-\rho^+$	< 6.6	$\times 10^{-3}$	CL=90%	DESIG=214
Г64	$K_S^0 K^-\pi^+\pi^-$	< 8.7	$\times 10^{-3}$	CL=90%	DESIG=202
Г65	$K_S^0 K^-\pi^+\rho^0$	< 1.6	%	CL=90%	DESIG=215
Г66	$K_S^0 K^-\pi^+\eta$	< 1.3	%	CL=90%	DESIG=216
Г67	$K_S^0 K^-\pi^+\pi^-\pi^0$	< 4.18	%	CL=90%	DESIG=203
Г68	$K_S^0 K^-\pi^+\pi^-\eta$	< 4.8	%	CL=90%	DESIG=217
Г69	$K_S^0 K^-\pi^+2(\pi^+\pi^-)$	< 1.22	%	CL=90%	DESIG=204
Г70	$K_S^0 K^-\pi^+2\pi^0$	< 2.65	%	CL=90%	DESIG=205
Г71	$K_S^0 K^-\pi^+K^-\pi^+$	< 4.9	$\times 10^{-3}$	CL=90%	DESIG=218
Г72	$K_S^0 K^-\pi^+K^-\pi^+\pi^0$	< 3.0	%	CL=90%	DESIG=219
Г73	$K_S^0 K^-\pi^+K^-\pi^+\eta$	< 2.2	%	CL=90%	DESIG=220
Г74	$K^{*0} K^-\pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%	DESIG=60
Г75	$\rho\bar{\rho}$				DESIG=233
Г76	$\rho\bar{\rho}\pi^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=54
Г77	$\rho\bar{\rho}\pi^+\pi^-$	< 5.8	$\times 10^{-4}$	CL=90%	DESIG=36
Г78	$\Lambda\bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%	DESIG=42
Г79	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	< 1.85	$\times 10^{-3}$	CL=90%	DESIG=37
Г80	$\omega\rho\bar{\rho}$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=39
Г81	$\Lambda\bar{\Lambda}\pi^0$	< 7	$\times 10^{-5}$	CL=90%	DESIG=72
Г82	$\rho\bar{\rho}2(\pi^+\pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%	DESIG=61
Г83	$\eta\rho\bar{\rho}$	< 5.4	$\times 10^{-4}$	CL=90%	DESIG=38
Г84	$\eta\rho\bar{\rho}\pi^+\pi^-$	< 3.3	$\times 10^{-3}$	CL=90%	DESIG=223
Г85	$\rho^0\rho\bar{\rho}$	< 1.7	$\times 10^{-3}$	CL=90%	DESIG=66
Г86	$\rho\bar{\rho}K^+K^-$	< 3.2	$\times 10^{-4}$	CL=90%	DESIG=40
Г87	$\eta\rho\bar{\rho}K^+K^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=224

Γ_{88}	$\pi^0 p \bar{p} K^+ K^-$	< 1.2	$\times 10^{-3}$	CL=90%	DESIG=225
Γ_{89}	$\phi p \bar{p}$	< 1.3	$\times 10^{-4}$	CL=90%	DESIG=41
Γ_{90}	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	< 2.5	$\times 10^{-4}$	CL=90%	DESIG=43
Γ_{91}	$\Lambda \bar{p} K^+$	< 2.8	$\times 10^{-4}$	CL=90%	DESIG=44
Γ_{92}	$\Lambda \bar{p} K^+ \pi^+ \pi^-$	< 6.3	$\times 10^{-4}$	CL=90%	DESIG=45
Γ_{93}	$\Lambda \bar{\Lambda} \eta$	< 1.9	$\times 10^{-4}$	CL=90%	DESIG=226
Γ_{94}	$\Sigma^+ \bar{\Sigma}^-$	< 1.0	$\times 10^{-4}$	CL=90%	DESIG=227
Γ_{95}	$\Sigma^0 \bar{\Sigma}^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=228
Γ_{96}	$\Xi^+ \bar{\Xi}^-$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=229
Γ_{97}	$\Xi^0 \bar{\Xi}^0$	< 1.4	$\times 10^{-4}$	CL=90%	DESIG=230

Radiative decays

Γ_{98}	$\gamma \chi_{c2}$	< 6.4	$\times 10^{-4}$	CL=90%	NODE=M053;CLUMP=R DESIG=51
Γ_{99}	$\gamma \chi_{c1}$	(2.48 ± 0.23)	$\times 10^{-3}$		DESIG=50
Γ_{100}	$\gamma \chi_{c0}$	(7.0 ± 0.6)	$\times 10^{-3}$		DESIG=49
Γ_{101}	$\gamma \eta_c$	< 7	$\times 10^{-4}$	CL=90%	DESIG=231
Γ_{102}	$\gamma \eta_c(2S)$	< 9	$\times 10^{-4}$	CL=90%	DESIG=232
Γ_{103}	$\gamma \eta'$	< 1.8	$\times 10^{-4}$	CL=90%	DESIG=213
Γ_{104}	$\gamma \eta$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=212
Γ_{105}	$\gamma \pi^0$	< 2	$\times 10^{-4}$	CL=90%	DESIG=211

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 3 branching ratios uses 23 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 20.1$ for 19 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	99		
x_8	0	0	
Γ	0	0	-44
	x_2	x_3	x_8

	Mode	Rate (MeV)	Scale factor
Γ_2	$D^0 \bar{D}^0$	14.0 ± 1.4	1.8
Γ_3	$D^+ D^-$	11.2 ± 1.1	1.7
Γ_8	$e^+ e^-$	$(2.62 \pm 0.18) \times 10^{-4}$	1.4

DESIG=5
DESIG=6
DESIG=1

$\psi(3770)$ PARTIAL WIDTHS

NODE=M053225

$\Gamma(e^+ e^-)$

Γ_8

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.262 ± 0.018 OUR FIT	Error includes scale factor of 1.4.			
0.256 ± 0.016 OUR AVERAGE	Error includes scale factor of 1.2.			
$0.154^{+0.079+0.021}_{-0.058-0.027}$	9,10	ANASHIN	12A KEDR	$e^+ e^- \rightarrow D \bar{D}$
0.22 ± 0.05	11,12	ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
$0.277 \pm 0.011 \pm 0.013$	12	ABLIKIM	07E BES2	$e^+ e^- \rightarrow$ hadrons
$0.203 \pm 0.003^{+0.041}_{-0.027}$	1.4M 12,13	BESSION	06 CLEO	$e^+ e^- \rightarrow$ hadrons
0.276 ± 0.050	12	SCHINDLER	80 MRK2	$e^+ e^-$
0.18 ± 0.06	12	BACINO	78 DLCO	$e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.414^{+0.072+0.093}_{-0.080-0.028}$	10,14	ANASHIN	12A KEDR	$e^+ e^- \rightarrow D \bar{D}$
0.37 ± 0.09	15	RAPIDIS	77 LGW	$e^+ e^-$

NODE=M053W1
NODE=M053W1

OCCUR=2

- ⁹ Solution I of the two solutions.
¹⁰ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.
¹¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.
¹² Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.
¹³ BESSON 06 (as corrected in BESSON 10) measure $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30}$ nb at $\sqrt{s} = 3773 \pm 1$ MeV, and obtain Γ_{ee} from the Born-level cross section calculated using $\psi(3770)$ mass and width from our 2004 edition, PDG 04.
¹⁴ Solution II of the two solutions.
¹⁵ See also $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ below.

NODE=M053W1;LINKAGE=A1
 NODE=M053W1;LINKAGE=AN
 NODE=M053W1;LINKAGE=AB

NODE=M053W1;LINKAGE=NI

NODE=M053W1;LINKAGE=BE

NODE=M053W1;LINKAGE=A2

NODE=M053W1;LINKAGE=R

$\psi(3770)$ BRANCHING RATIOS

NODE=M053230

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma = (\Gamma_2+\Gamma_3)/\Gamma$
 VALUE EVTS DOCUMENT ID TECN COMMENT

NODE=M053R1
 NODE=M053R1

0.93 $^{+0.08}_{-0.09}$ OUR FIT Error includes scale factor of 2.0.

0.93 $^{+0.08}_{-0.09}$ OUR AVERAGE Error includes scale factor of 2.1.

0.849 ± 0.056 ± 0.018 ¹⁶ ABLIKIM 08B BES2 $e^+e^- \rightarrow \text{non-}D\bar{D}$

1.033 ± 0.014 $^{+0.048}_{-0.066}$ 1.427M ¹⁷ BESSON 06 CLEO $e^+e^- \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.866 ± 0.050 ± 0.036 ^{18,19} ABLIKIM 07K BES2 $e^+e^- \rightarrow \text{non-}D\bar{D}$

0.836 ± 0.073 ± 0.042 ¹⁹ ABLIKIM 06L BES2 $e^+e^- \rightarrow D\bar{D}$

0.855 ± 0.017 ± 0.058 ^{19,20} ABLIKIM 06N BES2 $e^+e^- \rightarrow D\bar{D}$

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_2/Γ
 VALUE DOCUMENT ID TECN COMMENT

NODE=M053R46
 NODE=M053R46

0.52 $^{+0.04}_{-0.05}$ OUR FIT Error includes scale factor of 2.0. [0.52 ± 0.05 OUR 2015 FIT
 Scale factor = 2.0]

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.467 ± 0.047 ± 0.023 ABLIKIM 06L BES2 $e^+e^- \rightarrow D^0\bar{D}^0$

0.499 ± 0.013 ± 0.038 ²⁰ ABLIKIM 06N BES2 $e^+e^- \rightarrow D^0\bar{D}^0$

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$ Γ_3/Γ
 VALUE DOCUMENT ID TECN COMMENT

NODE=M053R47
 NODE=M053R47

0.41 ± 0.04 OUR FIT Error includes scale factor of 2.0. [0.41 ± 0.04 OUR 2015 FIT
 Scale factor = 2.0]

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.369 ± 0.037 ± 0.028 ABLIKIM 06L BES2 $e^+e^- \rightarrow D^+D^-$

0.357 ± 0.011 ± 0.034 ²⁰ ABLIKIM 06N BES2 $e^+e^- \rightarrow D^+D^-$

$\Gamma(D^0\bar{D}^0)/\Gamma(D^+D^-)$ Γ_2/Γ_3
 VALUE EVTS DOCUMENT ID TECN COMMENT

NODE=M053R5
 NODE=M053R5

1.253 ± 0.016 OUR FIT
 [1.260 ± 0.021 OUR 2015 FIT]

NEW

1.253 ± 0.016 OUR AVERAGE
 [1.260 ± 0.021 OUR 2015 AVERAGE]

NEW

1.252 ± 0.009 ± 0.013 5.3M BONVICINI 14 CLEO $e^+e^- \rightarrow D\bar{D}$

1.39 ± 0.31 ± 0.12 PAKHLOVA 08 BELL 10.6 $e^+e^- \rightarrow D\bar{D}\gamma$

1.78 ± 0.33 ± 0.24 AUBERT 07BE BABR $e^+e^- \rightarrow D\bar{D}\gamma$

1.27 ± 0.12 ± 0.08 ABLIKIM 06L BES2 $e^+e^- \rightarrow D\bar{D}$

2.43 ± 1.50 ± 0.43 34 ²¹ CHISTOV 04 BELL $B^+ \rightarrow \psi(3770)K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.258 ± 0.016 ± 0.014 ²² DOBBS 07 CLEO $e^+e^- \rightarrow D\bar{D}$

$\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_4/Γ
 VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

NODE=M053R4
 NODE=M053R4

1.93 ± 0.28 OUR AVERAGE

1.89 ± 0.20 ± 0.20 231 ± 33 ADAM 06 CLEO $e^+e^- \rightarrow \psi(3770)$

3.4 ± 1.4 ± 0.9 17.8 ± 4.8 BAI 05 BES2 $e^+e^- \rightarrow \psi(3770)$

$\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_5/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT			
0.080±0.025±0.016	39 ± 14	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R7 NODE=M053R7	
$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$					Γ_6/Γ		
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT			
87±33±22	22 ± 10	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R8 NODE=M053R8	
$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$					Γ_7/Γ		
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		
<28	90	<10	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R9 NODE=M053R9
$\Gamma(e^+e^-)/\Gamma_{\text{total}}$					Γ_8/Γ		
VALUE (units 10^{-5})	DOCUMENT ID TECN COMMENT						
0.96±0.07 OUR FIT	Error includes scale factor of 1.3.						
1.3 ±0.2	RAPIDIS	77	LGW	e^+e^-	NODE=M053R2 NODE=M053R2		
¹⁶ Neglecting interference. ¹⁷ Obtained by comparing a measurement of the total cross section (corrected in BESSON 10) with that of $D\bar{D}$ reported by CLEO in DOBBS 07. ¹⁸ Using $\sigma^{obs} = 7.07 \pm 0.58$ nb and neglecting interference. ¹⁹ Not independent of ABLIKIM 08B. ²⁰ From a measurement of $\sigma(e^+e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L. ²¹ See ADLER 88C for older measurements of this quantity. ²² Superseded by BONVICINI 14.							

————— DECAYS TO LIGHT HADRONS —————

$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$					Γ_9/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.4	90	²³ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R82 NODE=M053R82
$\Gamma(\phi\eta')/\Gamma_{\text{total}}$					Γ_{10}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<7	90	²³ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R83 NODE=M053R83
$\Gamma(\omega\eta')/\Gamma_{\text{total}}$					Γ_{11}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<4	90	²³ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R84 NODE=M053R84
$\Gamma(\rho^0\eta')/\Gamma_{\text{total}}$					Γ_{12}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<6	90	²³ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R85 NODE=M053R85
$\Gamma(\phi\eta)/\Gamma_{\text{total}}$					Γ_{13}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
3.1±0.6±0.3		²³ ADAMS	06	CLEO	$3.773 e^+e^- \rightarrow \phi\eta$	NODE=M053R6 NODE=M053R6
<ul style="list-style-type: none"> • • • We do not use the following data for averages, fits, limits, etc. • • • 						
<19	90	²⁴ ABLIKIM	07B	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					Γ_{14}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.4	90	²³ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R86 NODE=M053R86
$\Gamma(\rho^0\eta)/\Gamma_{\text{total}}$					Γ_{15}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<5	90	²³ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R87 NODE=M053R87
$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$					Γ_{16}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
< 3	90	²³ ADAMS	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	NODE=M053R11 NODE=M053R11
<ul style="list-style-type: none"> • • • We do not use the following data for averages, fits, limits, etc. • • • 						
<50	90	²⁴ ABLIKIM	07B	BES2	$e^+e^- \rightarrow \psi(3770)$	

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					Γ_{17}/Γ	NODE=M053R88 NODE=M053R88
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<6	90	23 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{18}/Γ	NODE=M053R89 NODE=M053R89
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<5	90	23,25 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					Γ_{19}/Γ	NODE=M053R90 NODE=M053R90
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<5	90	23,25 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(K^+K^-)/\Gamma_{\text{total}}$					Γ_{20}/Γ	NODE=M053R00 NODE=M053R00
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 2×10^{-5}	90	26 DRUZHININ	15 RVUE	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(K^*(892)^+K^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{21}/Γ	NODE=M053R91 NODE=M053R91
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.4	90	23 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{22}/Γ	NODE=M053R92 NODE=M053R92
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.2	90	23 ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$					Γ_{23}/Γ	NODE=M053R3 NODE=M053R3
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 1.2	90	27 CRONIN-HEN..06	CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<21	90	28 ABLIKIM	04F BES	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_{24}/Γ	NODE=M053R21 NODE=M053R21
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<11.2	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<48	90	24 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{25}/Γ	NODE=M053R22 NODE=M053R22
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<10.6	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<62	90	24 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{26}/Γ	NODE=M053R72 NODE=M053R72
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<58.5	90	305	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{27}/Γ	NODE=M053R24 NODE=M053R24
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 6.0	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<55	90	24 ABLIKIM	07I BES2	$3.77 e^+e^-$		
$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_{28}/Γ	NODE=M053R07 NODE=M053R07
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<91 (CL = 90%)	[< 91×10^{-4} OUR 2015 BEST LIMIT]					
<91	90	24 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{29}/Γ	NODE=M053R10 NODE=M053R10
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<137 (CL = 90%)	[< 137×10^{-4} OUR 2015 BEST LIMIT]					
<137	90	24 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		

$\Gamma(3(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$						Γ_{30}/Γ	NODE=M053R74 NODE=M053R74
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<117.4	90	59	ABLIKIM	08N	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{31}/Γ	NODE=M053R23 NODE=M053R23
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.24	90		29 HUANG	06A	CLEO	$e^+e^- \rightarrow \psi(3770)$	
••• We do not use the following data for averages, fits, limits, etc. •••							
<2.3	90		24 ABLIKIM	10D	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$						Γ_{32}/Γ	NODE=M053R70 NODE=M053R70
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<8.9	90	218	ABLIKIM	08N	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{33}/Γ	NODE=M053R53 NODE=M053R53
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<6.9	90		24 ABLIKIM	07F	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\eta3\pi)/\Gamma_{\text{total}}$						Γ_{34}/Γ	NODE=M053R25 NODE=M053R25
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<13.4	90		29 HUANG	06A	CLEO	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\eta2(\pi^+\pi^-))/\Gamma_{\text{total}}$						Γ_{35}/Γ	NODE=M053R08 NODE=M053R08
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<243 (CL = 90%)			[<243 × 10 ⁻⁴ OUR 2015 BEST LIMIT]				
<243	90		24 ABLIKIM	07B	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\eta\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{36}/Γ	NODE=M053R77 NODE=M053R77
<u>VALUE (units 10⁻²)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.45	90		24 ABLIKIM	10D	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\eta'3\pi)/\Gamma_{\text{total}}$						Γ_{37}/Γ	NODE=M053R26 NODE=M053R26
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<24.4	90		29 HUANG	06A	CLEO	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{38}/Γ	NODE=M053R27 NODE=M053R27
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 9.0	90		29 HUANG	06A	CLEO	$e^+e^- \rightarrow \psi(3770)$	
••• We do not use the following data for averages, fits, limits, etc. •••							
<48	90		24 ABLIKIM	07B	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{39}/Γ	NODE=M053R28 NODE=M053R28
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 4.1	90		29 HUANG	06A	CLEO	$e^+e^- \rightarrow \psi(3770)$	
••• We do not use the following data for averages, fits, limits, etc. •••							
<16	90		24 ABLIKIM	07B	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(K^+K^-2\pi^0)/\Gamma_{\text{total}}$						Γ_{40}/Γ	NODE=M053R71 NODE=M053R71
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<4.2	90	14	ABLIKIM	08N	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(4(\pi^+\pi^-))/\Gamma_{\text{total}}$						Γ_{41}/Γ	NODE=M053R50 NODE=M053R50
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<16.7	90		24 ABLIKIM	07F	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$						Γ_{42}/Γ	NODE=M053R52 NODE=M053R52
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<30.6	90		24 ABLIKIM	07F	BES2	$e^+e^- \rightarrow \psi(3770)$	
$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$						Γ_{43}/Γ	NODE=M053R29 NODE=M053R29
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<4.5	90		29 HUANG	06A	CLEO	$e^+e^- \rightarrow \psi(3770)$	

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{44}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
< 23.6	90	29 HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R30 NODE=M053R30
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<111	90	24 ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$		
$\Gamma(K^+ K^- \rho^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{45}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<8	90	24 ABLIKIM	07I	BES2 $3.77 e^+ e^-$		NODE=M053R58 NODE=M053R58
$\Gamma(K^+ K^- \rho^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{46}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<146	90	24 ABLIKIM	07I	BES2 $3.77 e^+ e^-$		NODE=M053R59 NODE=M053R59
$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$					Γ_{47}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
< 3.4	90	29 HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R32 NODE=M053R32
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<66	90	24 ABLIKIM	07I	BES2 $3.77 e^+ e^-$		
$\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{48}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<38	90	24 ABLIKIM	07I	BES2 $3.77 e^+ e^-$		NODE=M053R60 NODE=M053R60
$\Gamma(K^{*0} K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{49}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<162	90	24 ABLIKIM	07I	BES2 $3.77 e^+ e^-$		NODE=M053R61 NODE=M053R61
$\Gamma(K^{*+} K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{50}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<323	90	24 ABLIKIM	07I	BES2 $3.77 e^+ e^-$		NODE=M053R62 NODE=M053R62
$\Gamma(K^+ K^- \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$					Γ_{51}/Γ	
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<26.7	90	24	ABLIKIM	08N	BES2 $e^+ e^- \rightarrow \psi(3770)$	NODE=M053R73 NODE=M053R73
$\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$					Γ_{52}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT		
<10.3	90	24 ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R57 NODE=M053R57
$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$					Γ_{53}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT		
<36.0	90	24 ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R51 NODE=M053R51
$\Gamma(\eta K^+ K^-)/\Gamma_{\text{total}}$					Γ_{54}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
< 4.1	90	29 HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R31 NODE=M053R31
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<31	90	24 ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$		
$\Gamma(\eta K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{55}/Γ	
VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.24	90	24 ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R78 NODE=M053R78
$\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$					Γ_{56}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT		
<5.0	90	24 ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R54 NODE=M053R54
$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$					Γ_{57}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
< 6.0	90	29 HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R33 NODE=M053R33
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<17	90	24 ABLIKIM	07B	BES2 $e^+ e^- \rightarrow \psi(3770)$		

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 7.5	90	29 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<24	90	24 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R34
NODE=M053R34 $\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	29 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<46	90	24 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R35
NODE=M053R35 $\Gamma(2(K^+ K^-)\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.2	90	24 ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R48
NODE=M053R48 $\Gamma(K_S^0 K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.2	90	18	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R64
NODE=M053R64 $\Gamma(K_S^0 K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<13.3	90	40	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R65
NODE=M053R65 $\Gamma(K_S^0 K^- \rho^+)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R15
NODE=M053R15 $\Gamma(K_S^0 K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8.7	90	39	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R66
NODE=M053R66 $\Gamma(K_S^0 K^- \pi^+ \rho^0)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R16
NODE=M053R16 $\Gamma(K_S^0 K^- \pi^+ \eta)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R17
NODE=M053R17 $\Gamma(K_S^0 K^- 2\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<41.8	90	23	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R67
NODE=M053R67 $\Gamma(K_S^0 K^- 2\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<4.8	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R18
NODE=M053R18 $\Gamma(K_S^0 K^- \pi^+ 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<12.2	90	4	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R68
NODE=M053R68 $\Gamma(K_S^0 K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<26.5	90	17	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R69
NODE=M053R69 $\Gamma(K_S^0 K^- K^+ K^- \pi^+)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4.9	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R19
NODE=M053R19

$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$						Γ_{72}/Γ	
VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT			
<3.0	90	ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R75 NODE=M053R75
$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \eta)/\Gamma_{\text{total}}$						Γ_{73}/Γ	
VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT			
<2.2	90	ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R76 NODE=M053R76
$\Gamma(K^{*0} K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{74}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<9.7	90	24 ABLIKIM	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R55 NODE=M053R55
$\Gamma(\rho\bar{p})/\Gamma_{\text{total}}$						Γ_{75}/Γ	
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT			
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$7.1^{+8.6}_{-2.9}$	684	30 ABLIKIM	14L	BES3	$e^+ e^- \rightarrow \psi(3770)$		
310 ±30	684	31 ABLIKIM	14L	BES3	$e^+ e^- \rightarrow \psi(3770)$		OCCUR=2
$\Gamma(\rho\bar{p}\pi^0)/\Gamma_{\text{total}}$						Γ_{76}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT			
< 0.4	90	32 ABLIKIM	14O	BES3	Sol. I, $e^+ e^- \rightarrow \psi(3770)$		NODE=M053R09 NODE=M053R09
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$59^{+3}_{-2} \pm 5$		32 ABLIKIM	14O	BES3	Sol. II, $e^+ e^- \rightarrow \psi(3770)$		OCCUR=2
<12	90	24 ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$		
$\Gamma(\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{77}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT			
< 5.8	90	29 HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R36 NODE=M053R36
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<16	90	24 ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$		
$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$						Γ_{78}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT			
<1.2	90	29 HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R42 NODE=M053R42
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<4	90	24 ABLIKIM	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$		
$\Gamma(\rho\bar{p}\pi^+\pi^- \pi^0)/\Gamma_{\text{total}}$						Γ_{79}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT			
<18.5	90	29 HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R37 NODE=M053R37
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<73	90	24 ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$		
$\Gamma(\omega\rho\bar{p})/\Gamma_{\text{total}}$						Γ_{80}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT			
< 2.9	90	29 HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R39 NODE=M053R39
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<30	90	33 ABLIKIM	07I	BES2	$3.77 e^+ e^-$		
$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$						Γ_{81}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT			
< 0.7	90	34 ABLIKIM	13Q	BES3	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R63 NODE=M053R63
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<12	90	24 ABLIKIM	07I	BES2	$3.77 e^+ e^-$		
$\Gamma(\rho\bar{p}2(\pi^+\pi^-))/\Gamma_{\text{total}}$						Γ_{82}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<2.6	90	24 ABLIKIM	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$		NODE=M053R49 NODE=M053R49

$\Gamma(\eta\rho\bar{p})/\Gamma_{\text{total}}$					Γ_{83}/Γ	NODE=M053R38 NODE=M053R38
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 5.4	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<11	90	24 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\eta\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{84}/Γ	NODE=M053R79 NODE=M053R79
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<3.3	90	24 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\rho^0\rho\bar{p})/\Gamma_{\text{total}}$					Γ_{85}/Γ	NODE=M053R56 NODE=M053R56
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.7	90	24 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$					Γ_{86}/Γ	NODE=M053R40 NODE=M053R40
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 3.2	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<11	90	24 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\eta\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$					Γ_{87}/Γ	NODE=M053R80 NODE=M053R80
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<6.9	90	24 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\pi^0\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$					Γ_{88}/Γ	NODE=M053R81 NODE=M053R81
<u>VALUE (units 10⁻³)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.2	90	24 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\phi\rho\bar{p})/\Gamma_{\text{total}}$					Γ_{89}/Γ	NODE=M053R41 NODE=M053R41
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.3	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<9	90	24 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{90}/Γ	NODE=M053R43 NODE=M053R43
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
< 2.5	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 4.7	90	34 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$		
<39	90	24 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$					Γ_{91}/Γ	NODE=M053R44 NODE=M053R44
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<2.8	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{92}/Γ	NODE=M053R45 NODE=M053R45
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<6.3	90	29 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$					Γ_{93}/Γ	NODE=M053R93 NODE=M053R93
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.9	90	34 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$					Γ_{94}/Γ	NODE=M053R94 NODE=M053R94
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.0	90	34 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$		
$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$					Γ_{95}/Γ	NODE=M053R95 NODE=M053R95
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.4	90	34 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$		

$\Gamma(\Xi^+\Xi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{96}/Γ
<1.5	90	34 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(3770)$	NODE=M053R96 NODE=M053R96

 $\Gamma(\Xi^0\Xi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{97}/Γ
<1.4	90	34 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(3770)$	NODE=M053R97 NODE=M053R97

²³ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

²⁴ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

²⁵ Data suggest possible destructive interference with continuum.

²⁶ DRUZHININ 15 uses measurements derived from BABAR and CLEO data.

²⁷ Using $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = (6.38 \pm 0.08^{+0.41}_{-0.30})$ nb from BESSON 06 and $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6895 \pm 0.0014$.

²⁸ Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$.

²⁹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

³⁰ Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

³¹ Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

³² Solution I or II of two equivalent solutions in a fit with a resonance interfering with continuum. Calculated by the authors using $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30}$ nb from BESSON 10.

³³ Using $\sigma^{obs} = 7.15 \pm 0.27 \pm 0.27$ nb and neglecting interference.

³⁴ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R6;LINKAGE=AD

NODE=M053R10;LINKAGE=AK

NODE=M053R89;LINKAGE=AD

NODE=M053R00;LINKAGE=A

NODE=M053R3;LINKAGE=CR

NODE=M053R3;LINKAGE=AB

NODE=M053R;LINKAGE=HU

NODE=M053R98;LINKAGE=A

NODE=M053R98;LINKAGE=B

NODE=M053R09;LINKAGE=A

NODE=M053R24;LINKAGE=AB

NODE=M053R43;LINKAGE=A

NODE=M053240

RADIATIVE DECAYS

 $\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{98}/Γ
<0.64 (CL = 90%)				[<0.9 $\times 10^{-3}$ (CL = 90%) OUR 2015 BEST LIMIT]	NODE=M053R03 NODE=M053R03

<0.64	90	35 ABLIKIM	15J	BES3 $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	
-------	----	------------	-----	--	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0	90	36 BRIERE	06	CLEO $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
------	----	-----------	----	--	--

<0.9	90	37 COAN	06A	CLEO $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	
------	----	---------	-----	--	--

 $\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
2.48 \pm 0.23 OUR AVERAGE					NODE=M053R02 NODE=M053R02 NEW

[(2.7 \pm 0.5) $\times 10^{-3}$ OUR 2015 AVERAGE]

1.9 \pm 0.4 \pm 0.6	202	38 ABLIKIM	16B	BES3 $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
-------------------------	-----	------------	-----	--	--

2.48 \pm 0.15 \pm 0.23	0.6k	ABLIKIM	15J	BES3 $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	
----------------------------	------	---------	-----	--	--

2.4 \pm 0.8 \pm 0.2		39 ABLIKIM	14H	BES3 $e^+e^- \rightarrow \psi(3770) \rightarrow K_S^0 K^\pm \pi^\mp$	
-------------------------	--	------------	-----	--	--

2.9 \pm 0.5 \pm 0.4		40 BRIERE	06	CLEO $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}, \gamma\gamma J/\psi$	OCCUR=2
-------------------------	--	-----------	----	---	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.9 \pm 1.4 \pm 0.6	54	41 BRIERE	06	CLEO $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
-------------------------	----	-----------	----	--	--

2.8 \pm 0.5 \pm 0.4	53	37 COAN	06A	CLEO $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	
-------------------------	----	---------	-----	--	--

 $\Gamma(\gamma\chi_{c1})/\Gamma(J/\psi\pi^+\pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ_4
1.49 \pm 0.31 \pm 0.26	53 \pm 10	42 COAN	06A	CLEO $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	NODE=M053R04 NODE=M053R04

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ
--------------------------	-----	------	-------------	------	---------	-----------------------

7.0±0.6 OUR AVERAGE[(7.3 ± 0.9) × 10⁻³ OUR 2015 AVERAGE]

6.9±0.3±0.7	2.2K	43	ABLIKIM	16B	BES3	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
7.3±0.7±0.6	274		BRIERE	06	CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
< 44	90	37	COAN	06A	CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

 $\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c2})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ_{98}
-------	-----	-------------	------	---------	----------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

>8	90	44	BRIERE	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	
----	----	----	--------	----	------	---------------------------------	--

 $\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c1})$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ_{99}
-------	-------------	------	---------	----------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

2.5±0.6	44	BRIERE	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	
---------	----	--------	----	------	---------------------------------	--

 $\Gamma(\gamma\eta_c)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{101}/Γ
-------	-----	-------------	------	---------	-----------------------

<7 × 10 ⁻⁴	90	45	ABLIKIM	14H	BES3	
-----------------------	----	----	---------	-----	------	--

 $\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{102}/Γ
-------	-----	-------------	------	---------	-----------------------

<9 × 10 ⁻⁴	90	46	ABLIKIM	14H	BES3	
-----------------------	----	----	---------	-----	------	--

 $\Gamma(\gamma\eta')/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{103}/Γ
---------------------------------	-----	-------------	------	---------	-----------------------

<1.8	90	47	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
------	----	----	--------	----	------	---------------------------------

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{104}/Γ
---------------------------------	-----	-------------	------	---------	-----------------------

<1.5	90	47	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
------	----	----	--------	----	------	---------------------------------

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{105}/Γ
---------------------------------	-----	-------------	------	---------	-----------------------

<2	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$	
----	----	--------	----	------	---------------------------------	--

³⁵ This limit is equivalent to $(0.25 \pm 0.21 \pm 0.18) \times 10^{-3}$ branching fraction value.

³⁶ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = 9.22 \pm 0.11 \pm 0.46\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

³⁷ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+e^- \rightarrow \psi(3770))$.

³⁸ ABLIKIM 16B reports $(1.94 \pm 0.42 \pm 0.64) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$.

³⁹ ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)] = (8.51 \pm 2.39 \pm 1.42) \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp) = 0.00356 \pm 0.00030$. Our first error is their experiment's error and our second error is the systematic error from using our best value. We have calculated the best value of $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/2 of $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.1 \pm 0.6) \times 10^{-3}$.

⁴⁰ Averages the two measurements from COAN 06A and BRIERE 06.

⁴¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = 9.07 \pm 0.11 \pm 0.54\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

⁴² Using $B(\psi(3770) \rightarrow J/\psi \pi^+ \pi^-) = (1.89 \pm 0.20 \pm 0.20) \times 10^{-3}$ from ADAM 06.

⁴³ ABLIKIM 16B reports $(6.88 \pm 0.28 \pm 0.67) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c0})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$.

⁴⁴ Not independent of other results in BRIERE 06.

⁴⁵ ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 16 \times 10^{-6}$ which we divide by our best value $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp) = 2.43 \times 10^{-2}$.

NODE=M053R01
 NODE=M053R01
 NEW

NODE=M053R06
 NODE=M053R06

NODE=M053R05
 NODE=M053R05

NODE=M053R99
 NODE=M053R99

NODE=M053R20
 NODE=M053R20

NODE=M053R14
 NODE=M053R14

NODE=M053R13
 NODE=M053R13

NODE=M053R12
 NODE=M053R12

NODE=M053R03;LINKAGE=A
 NODE=M053R03;LINKAGE=BR

NODE=M053R0;LINKAGE=CO

NODE=M053R02;LINKAGE=A

NODE=M053R02;LINKAGE=AB

NODE=M053R02;LINKAGE=BI
 NODE=M053R02;LINKAGE=BR

NODE=M053R04;LINKAGE=CO
 NODE=M053R01;LINKAGE=B

NODE=M053R05;LINKAGE=BR
 NODE=M053R99;LINKAGE=AB

We have calculated the best value of $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(1S) \rightarrow K \bar{K} \pi) = 7.3 \times 10^{-2}$.

46 ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 5.6 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp) = 6 \times 10^{-3}$.

We have calculated the best value of $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(2S) \rightarrow K \bar{K} \pi) = 1.9 \times 10^{-2}$.

47 Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.

NODE=M053R20;LINKAGE=AB

NODE=M053R13;LINKAGE=PE

$\psi(3770)$ REFERENCES

NODE=M053

ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=57126
ABLIKIM	15J	PR D91 092009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56775
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)	REFID=56962
ABLIKIM	14H	PR D89 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55899
ABLIKIM	14L	PL B735 101	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55903
ABLIKIM	14O	PR D90 032007	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55906
BONVICINI	14	PR D89 072002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=55798
ABLIKIM	13Q	PR D87 112011	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55393
ANASHIN	12A	PL B711 292	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=54055
ABLIKIM	10D	EPJ C66 11	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53350
BESSON	10	PRL 104 159901 (err.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
ABLIKIM	09C	EPJ C64 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=53134
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
ABLIKIM	08M	PL B670 179	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52569
ABLIKIM	08N	PL B670 184	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52570
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)	REFID=52144
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
ABLIKIM	07B	PL B650 111	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51704
ABLIKIM	07E	PL B652 238	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51882
ABLIKIM	07F	PL B656 30	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51940
ABLIKIM	07I	EPJ C52 805	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52045
ABLIKIM	07K	PR D76 122002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52073
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52074
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52075
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06N	PL B641 145	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51131
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
ADAMS	06	PR D73 012002	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50990
BESSON	06	PRL 96 092002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51041
Also		PRL 104 159901 (err.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
BRIERE	06	PR D74 031106	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=51149
COAN	06A	PRL 96 182002	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51155
CRONIN-HEN...	06	PR D74 012005	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=51156
HUANG	06A	PRL 96 032003	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=50999
BAI	05	PL B605 63	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50332
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=50924
Also		PRL 96 199903 (err.)	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=51211
ABLIKIM	04F	PR D70 077101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50185
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40361
SCHINDLER	80	PR D21 2716	R.H. Schindler <i>et al.</i>	(Mark II Collab.)	REFID=22222
BACINO	78	PRL 40 671	W.J. Bacino <i>et al.</i>	(SLAC, UCLA, UCI)	REFID=11437
RAPIDIS	77	PRL 39 526	P.A. Rapidis <i>et al.</i>	(LGW Collab.)	REFID=22220

$\psi(3823)$ was $X(3823)$,

$$J^{PC} = 2^{--}$$

 J, P need confirmation.

Seen by BHARDWAJ 13 in $B \rightarrow \chi_{c1} \gamma K$ and ABLIKIM 15S in $e^+ e^- \rightarrow \pi^+ \pi^- \gamma \chi_{c1}$ decays as a narrow peak in the invariant mass distribution of the $\chi_{c1} \gamma$ system. Properties consistent with the $\psi_2(1^3D_2) c \bar{c}$ state.

NODE=M212

NODE=M212

$\psi(3823)$ MASS

NODE=M212M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M212M

3822.2 ± 1.2 OUR AVERAGE

NEW

[3823.1 ± 1.9 MeV OUR 2015 AVERAGE]

3821.7 ± 1.3 ± 0.7	19 ± 5	¹ ABLIKIM	15S	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
3823.1 ± 1.8 ± 0.7	33 ± 10	² BHARDWAJ	13	BELL	$B \rightarrow \chi_{c1} \gamma K$

¹ From a simultaneous unbinned maximum likelihood fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1} \gamma$ and $\psi(3823) \rightarrow \chi_{c1} \gamma$ together, with floating mass scale offset for $\psi(2S)$, floating $\psi(3823)$ mass, and zero $\psi(3823)$ width, resulting in a significance of 5.9σ when including systematic uncertainties.

NODE=M212M;LINKAGE=B

² From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$ and $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$ with significance 4.0σ including systematics. Corrected for the measured $\psi(2S)$ mass using $B \rightarrow \psi(2S) K \rightarrow (\gamma \chi_{c1}) K$ decays.

NODE=M212M;LINKAGE=A

$\psi(3823)$ WIDTH

NODE=M212W

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

NODE=M212W

<16 (CL = 90%) [<24 MeV (CL = 90%) OUR 2015 BEST LIMIT]

<16	90	¹ ABLIKIM	15S	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
---------------	----	----------------------	-----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<24	90	² BHARDWAJ	13	BELL	$B \rightarrow \chi_{c1} \gamma K$
-------	----	-----------------------	----	------	------------------------------------

¹ From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to a Breit-Wigner function with the mass fixed from the likelihood fit above, Gaussian resolution smearing, and floating width.

NODE=M212W;LINKAGE=B

² From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$ and $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$ with significance 4.0σ including systematics.

NODE=M212W;LINKAGE=A

$\psi(3823)$ DECAY MODES

NODE=M212215;NODE=M212

Mode	Fraction (Γ_i/Γ)
Γ_1 $\chi_{c1} \gamma$	seen
Γ_2 $\chi_{c2} \gamma$	not seen

DESIG=1

DESIG=2

$\psi(3823)$ BRANCHING RATIOS

NODE=M212225

$\Gamma(\chi_{c1} \gamma)/\Gamma_{\text{total}}$	Γ_1/Γ
--	-------------------

NODE=M212R01

NODE=M212R01

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	33 ± 10	¹ BHARDWAJ	13	BELL	$B^+ \rightarrow \chi_{c1} \gamma K^+$
-------------	---------	-----------------------	----	------	--

¹ Reported $B(B^\pm \rightarrow \psi(3823) K^\pm) \times B(\psi(3823) \rightarrow \gamma \chi_{c1}) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$ with statistical significance 3.8σ .

NODE=M212R01;LINKAGE=A

$\Gamma(\chi_{c2} \gamma)/\Gamma_{\text{total}}$	Γ_2/Γ
--	-------------------

NODE=M212R02

NODE=M212R02

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	¹ ABLIKIM	15S	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c2} \gamma$
----------	----------------------	-----	------	--

not seen	² BHARDWAJ	13	BELL	$B^+ \rightarrow \chi_{c2} \gamma K^+$
-----------------	-----------------------	----	------	--

¹ From a simultaneous unbinned maximum likelihood fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c2} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c2} \gamma$ and $\psi(3823) \rightarrow \chi_{c2} \gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi(3823)$ mass floating (fixed to that above), and zero $\psi(3823)$ width.

NODE=M212R02;LINKAGE=B

² Reported $B(B^\pm \rightarrow \psi(3823) K^\pm) \times B(\psi(3823) \rightarrow \gamma \chi_{c2}) < 3.6 \times 10^{-6}$ at 90% CL.

NODE=M212R02;LINKAGE=A

$\Gamma(\chi_{c2}\gamma)/\Gamma(\chi_{c1}\gamma)$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.41	90	BHARDWAJ 13	BELL	$B^+ \rightarrow \chi_{c1/c2}\gamma K^+$

NODE=M212R03
 NODE=M212R03

••• We do not use the following data for averages, fits, limits, etc. •••

<0.42	90	¹ ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
-------	----	----------------------	----------	--

¹ From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1(2)}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1(2)}\gamma$ and $\psi(3823) \rightarrow \chi_{c1(2)}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi(3823)$ mass floating (fixed to that above), and zero $\psi(3823)$ width.

NODE=M212R03;LINKAGE=A

$\psi(3823)$ REFERENCES

ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BES III Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)

NODE=M212

REFID=56784
 REFID=55412

NODE=M176

X(3872)

$$J^{PC} = 0^+(1^{++})$$

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^- J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^- J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

NODE=M176

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow X(3872)K^+$ decays, where $X(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to $< 4\%$ at 95% CL.

See our note on "Developments in Heavy Quarkonium Spectroscopy".

X(3872) MASS FROM $J/\psi X$ MODE

NODE=M176M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3871.69 ± 0.17 OUR AVERAGE				
3871.9 ± 0.7 ± 0.2	20 ± 5	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
3871.95 ± 0.48 ± 0.12	0.6k	AAIJ	12H LHCB	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3871.85 ± 0.27 ± 0.19	~ 170	¹ CHOI	11 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3873 $\begin{smallmatrix} +1.8 \\ -1.6 \end{smallmatrix}$ ± 1.3	27 ± 8	² DEL-AMO-SA.10B	BABR	$B \rightarrow \omega J/\psi K$
3871.61 ± 0.16 ± 0.19	6k	^{2,3} AALTONEN	09AU CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3871.4 ± 0.6 ± 0.1	93.4	AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
3868.7 ± 1.5 ± 0.4	9.4	AUBERT	08Y BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.8 ± 3.1 ± 3.0	522	^{2,4} ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
••• We do not use the following data for averages, fits, limits, etc. •••				
3868.6 ± 1.2 ± 0.2	8	⁵ AUBERT	06 BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.3 ± 0.6 ± 0.1	61	⁵ AUBERT	06 BABR	$B^- \rightarrow K^- J/\psi\pi^+\pi^-$
3873.4 ± 1.4	25	⁶ AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
3871.3 ± 0.7 ± 0.4	730	^{2,7} ACOSTA	04 CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3872.0 ± 0.6 ± 0.5	36	⁸ CHOI	03 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3836 ± 13	58	^{2,9} ANTONIAZZI	94 E705	$300 \pi^\pm Li \rightarrow J/\psi\pi^+\pi^- X$

NODE=M176M

OCCUR=2

OCCUR=2

¹ The mass difference for the $X(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.

² Width consistent with detector resolution.

³ A possible equal mixture of two states with a mass difference greater than 3.6 MeV/ c^2 is excluded at 95% CL.

⁴ Calculated from the corresponding $m_{X(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.

⁵ Calculated from the corresponding $m_{X(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.

NODE=M176M;LINKAGE=CO

NODE=M176M;LINKAGE=AC

NODE=M176M;LINKAGE=AA

NODE=M176M;LINKAGE=AB

NODE=M176M;LINKAGE=AE

⁶ Calculated from the corresponding $m_{X(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96\text{MeV}$.

Superseded by AUBERT 06.

⁷ Superseded by AALTONEN 09AU.

⁸ Superseded by CHOI 11.

⁹ A lower mass value can be due to an incorrect momentum scale for soft pions.

NODE=M176M;LINKAGE=AU

NODE=M176M;LINKAGE=AT

NODE=M176M;LINKAGE=CH

NODE=M176M;LINKAGE=AN

X(3872) MASS FROM $\bar{D}^{*0} D^0$ MODE

NODE=M176MD0

NODE=M176MD0

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3872.9^{+0.6+0.4}_{-0.4-0.5}$	50	^{1,2} AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
--------------------------------	----	-----------------------	----	---

$3875.1^{+0.7}_{-0.5} \pm 0.5$	33 ± 6	² AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$
--------------------------------	------------	---------------------	-----	---

$3875.2 \pm 0.7^{+0.9}_{-1.8}$	24 ± 6	^{2,3} GOKHROO	06	BELL $B \rightarrow D^0 \bar{D}^0 \pi^0 K$
--------------------------------	------------	------------------------	----	--

¹ Calculated from the measured $m_{X(3872)} - m_{D^{*0}} - m_{\bar{D}^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$ MeV.

² Experiments report $D^{*0} \bar{D}^0$ invariant mass above $D^{*0} \bar{D}^0$ threshold because D^{*0} decay products are kinematically constrained to the D^{*0} mass, even though the D^{*0} may decay off-shell.

³ Superseded by AUSHEV 10.

NODE=M176MD0;LINKAGE=AS

NODE=M176MD0;LINKAGE=AU

NODE=M176MD0;LINKAGE=GO

$m_{X(3872)} - m_{J/\psi}$

NODE=M176207

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

$774.9 \pm 3.1 \pm 3.0$	522	ABAZOV	04F D0	$\rho \bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
---	-----	--------	--------	---

NODE=M176DM

$m_{X(3872)} - m_{\psi(2S)}$

NODE=M176DM2

NODE=M176DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

187.4 ± 1.4	25	¹ AUBERT	05R	BABR $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
-----------------	----	---------------------	-----	---

¹ Superseded by AUBERT 06.

NODE=M176DM2;LINKAGE=AU

X(3872) WIDTH

NODE=M176W

NODE=M176W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	-----	------	-------------	------	---------

<1.2	90	CHOI	11	BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
----------------	----	------	----	------	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.4	90	ABLIKIM	14	BES3	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
------	----	---------	----	------	---

<3.3	90	AUBERT	08Y	BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
------	----	--------	-----	------	--

<4.1	90	69	AUBERT	06	BABR $B \rightarrow K \pi^+ \pi^- J/\psi$
------	----	----	--------	----	---

<2.3	90	36	¹ CHOI	03	BELL $B \rightarrow K \pi^+ \pi^- J/\psi$
------	----	----	-------------------	----	---

¹ Superseded by CHOI 11.

NODE=M176W;LINKAGE=CH

X(3872) WIDTH FROM $\bar{D}^{*0} D^0$ MODE

NODE=M176WD0

NODE=M176WD0

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.9^{+2.8+0.2}_{-1.4-1.1}$	50	¹ AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
-----------------------------	----	---------------------	----	---

$3.0^{+1.9}_{-1.4} \pm 0.9$	33 ± 6	AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$
-----------------------------	------------	--------	-----	---

¹ With a measured value of $B(B \rightarrow X(3872) K) \times B(X(3872) \rightarrow D^{*0} \bar{D}^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.

NODE=M176WD0;LINKAGE=AU

X(3872) DECAY MODES

NODE=M176215;NODE=M176

Mode	Fraction (Γ_i/Γ)	
Γ_1 e^+e^-		DESIG=1
Γ_2 $\pi^+\pi^- J/\psi(1S)$	> 2.6 %	DESIG=2
Γ_3 $\rho^0 J/\psi(1S)$		DESIG=10
Γ_4 $\omega J/\psi(1S)$	> 1.9 %	DESIG=13
Γ_5 $D^0\bar{D}^0\pi^0$	>32 %	DESIG=8
Γ_6 $\bar{D}^{*0}D^0$	>24 %	DESIG=12
Γ_7 $\gamma\gamma$		DESIG=5
Γ_8 $D^0\bar{D}^0$		DESIG=6
Γ_9 D^+D^-		DESIG=7
Γ_{10} γX_{c1}		DESIG=3
Γ_{11} γX_{c2}		DESIG=15
Γ_{12} $\gamma J/\psi$	> 6×10^{-3}	DESIG=9
Γ_{13} $\gamma\psi(2S)$	> 3.0 %	DESIG=11
Γ_{14} $\pi^+\pi^-\eta_c(1S)$	not seen	DESIG=14;OUR EVAL;→ UNCHECKED ←
Γ_{15} $p\bar{p}$	not seen	DESIG=16;OUR EVAL;→ UNCHECKED ←
C-violating decays		
Γ_{16} $\eta J/\psi$		NODE=M176;CLUMP=A DESIG=4

X(3872) PARTIAL WIDTHS

NODE=M176220

 $\Gamma(e^+e^-)$ **Γ_1**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M176W1
NODE=M176W1

••• We do not use the following data for averages, fits, limits, etc. •••

< 4.3	90	¹ ABLIKIM	15V BES3	4.0-4.4 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
<280	90	² YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ ABLIKIM 15V reports this limit from the measurement of $\Gamma(X(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) \times \Gamma(X(3872) \rightarrow e^+e^-)/\Gamma < 0.13$ eV using $\Gamma(X(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma = 3\%$.

NODE=M176W1;LINKAGE=B

² Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. Assuming that $\Gamma(\pi^+\pi^- J/\psi)$ of X(3872) is the same as that of $\psi(2S)$ (85.4 keV).

NODE=M176W1;LINKAGE=A

X(3872) $\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M176230

 $\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_2\Gamma_1/\Gamma$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M176G1
NODE=M176G1

< **0.13 (CL = 90%)** [<6.2 eV (CL = 90%) OUR 2015 BEST LIMIT]

< 0.13	90	ABLIKIM	15V BES3	4.0-4.4 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
---------------	----	---------	----------	--

••• We do not use the following data for averages, fits, limits, etc. •••

< 6.2	90	^{1,2} AUBERT	05D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
-------	----	-----------------------	----------	--

< 8.3	90	² DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
-------	----	--------------------	---------	--

<10	90	³ YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
-----	----	-------------------	---------	--

¹ Using $B(X(3872) \rightarrow J/\psi\pi^+\pi^-) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot \Gamma(X(3872) \rightarrow e^+e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.

NODE=M176G1;LINKAGE=AU

² Assuming X(3872) has $J^{PC} = 1^{--}$.

NODE=M176G1;LINKAGE=DO

³ Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10)\%$.

NODE=M176G1;LINKAGE=A

X(3872) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M176232

 $\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_2\Gamma_7/\Gamma$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M176H1
NODE=M176H1

••• We do not use the following data for averages, fits, limits, etc. •••

<12.9	90	¹ DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi\gamma$
-------	----	--------------------	---------	--

¹ Assuming X(3872) has positive C parity and spin 0.

NODE=M176H1;LINKAGE=DO

$$\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M176G01
NODE=M176G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7 90 1 LEES 12AD BABR $e^+e^- \rightarrow e^+e^-\omega J/\psi$

¹ Assuming $X(3872)$ has spin 2.

NODE=M176G01;LINKAGE=LE

$$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M176G02
NODE=M176G02

<11.1 90 LEES 12AE BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

X(3872) BRANCHING RATIOS

NODE=M176235

$$\Gamma(\pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M176R6
NODE=M176R6

>0.026 93 ± 17 1 AUBERT 08Y BABR $B \rightarrow X(3872)K$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 151 2 BALA 15 BELL $B \rightarrow X(3872)K\pi$

>0.04 30 3 AUBERT 05R BABR $B^+ \rightarrow K^+\pi^+\pi^-J/\psi$

>0.04 36 ± 7 4 CHOI 03 BABR $B^+ \rightarrow K^+\pi^+\pi^-J/\psi$

NODE=M176R6;LINKAGE=AB

¹ AUBERT 08Y reports $[\Gamma(X(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

² BALA 15 reports $B(X(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^0 \rightarrow X(3872)K^+\pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(X(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^+ \rightarrow X(3872)K^0\pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.

NODE=M176R6;LINKAGE=A

³ Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(X(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R6;LINKAGE=AE

⁴ CHOI 03 reports $[\Gamma(X(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] / [B(B^+ \rightarrow \psi(2S)K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S)K^+) = (6.26 \pm 0.24) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$.

NODE=M176R6;LINKAGE=CH

$$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M176R14
NODE=M176R14

>0.019 21 ± 7 1 DEL-AMO-SA..10B BABR $B^+ \rightarrow \omega J/\psi K^+$

¹ DEL-AMO-SANCHEZ 10B reports $[\Gamma(X(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow X(3872)K^0) \times B(X(3872) \rightarrow J/\psi\omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.

NODE=M176R14;LINKAGE=DE

$$\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+\pi^-J/\psi(1S)) \quad \Gamma_4/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M176R15
NODE=M176R15

0.8 ± 0.3 1 DEL-AMO-SA..10B BABR $B \rightarrow \omega J/\psi K$

¹ Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow X(3872)K) \times B(X(3872) \rightarrow J/\psi\pi^+\pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.

NODE=M176R15;LINKAGE=DE

$$\Gamma(D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M176R12
NODE=M176R12

>0.32 17 ± 5 1 GOKHROO 06 BELL $B^+ \rightarrow D^0\bar{D}^0\pi^0 K^+$

¹ GOKHROO 06 reports $[\Gamma(X(3872) \rightarrow D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R12;LINKAGE=GO

$$\Gamma(\bar{D}^{*0}D^0)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M176R13
NODE=M176R13

>0.24 41 \pm $\frac{9}{8}$ 1 AUSHEV 10 BELL $B^+ \rightarrow D^{*0}\bar{D}^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.5 27 ± 6 2 AUBERT 08B BABR $B^+ \rightarrow \bar{D}^{*0}D^0 K^+$

¹ AUSHEV 10 reports $[\Gamma(X(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R13;LINKAGE=AS

² AUBERT 08B reports $[\Gamma(X(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R13;LINKAGE=AU

$\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_5/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ GOKHROO 06	BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$

NODE=M176R5
NODE=M176R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	AUSHEV 10	BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$
------	-----------	------	---------------------------------------

¹ May not necessarily be the same state as that observed in the $J/\psi \pi^+ \pi^-$ mode. Superseded CHISTOV 04.

NODE=M176R5;LINKAGE=GO

$\Gamma(D^0 \bar{D}^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_8/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CHISTOV 04	BELL	$B \rightarrow K D^0 \bar{D}^0$

NODE=M176R3
NODE=M176R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	CHISTOV 04	BELL	$B \rightarrow K D^0 \bar{D}^0$
----------	------------	------	---------------------------------

$\Gamma(D^+ D^-)/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_9/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CHISTOV 04	BELL	$B \rightarrow K D^+ D^-$

NODE=M176R4
NODE=M176R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	CHISTOV 04	BELL	$B \rightarrow K D^+ D^-$
----------	------------	------	---------------------------

$\Gamma(\gamma \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{10}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
not seen		¹ BHARDWAJ 13	BELL	$B^+ \rightarrow \chi_{c1} \gamma K^+$
<0.89	90	CHOI 03	BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$

NODE=M176R1
NODE=M176R1

¹ Reported $B(B^\pm \rightarrow X(3872)K^\pm) \times B(X(3872) \rightarrow \gamma \chi_{c1}) < 1.9 \times 10^{-6}$ at 90% CL.

NODE=M176R1;LINKAGE=A

$\Gamma(\gamma \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{11}/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ BHARDWAJ 13	BELL	$B^\pm \rightarrow \chi_{c2} \gamma K^\pm$

NODE=M176R01
NODE=M176R01

¹ Reported $B(B^\pm \rightarrow X(3872)K^\pm) \times B(X(3872) \rightarrow \gamma \chi_{c2}) < 6.7 \times 10^{-6}$ at 90% CL.

NODE=M176R01;LINKAGE=A

$\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
>6 $\times 10^{-3}$		¹ BHARDWAJ 11	BELL	$B^\pm \rightarrow \gamma J/\psi K^\pm$

NODE=M176R7
NODE=M176R7

• • • We do not use the following data for averages, fits, limits, etc. • • •

>9 $\times 10^{-3}$	20	² AUBERT 09B	BABR	$B^+ \rightarrow \gamma J/\psi K^+$
---------------------	----	-------------------------	------	-------------------------------------

>0.010	19	³ AUBERT, BE 06M	BABR	$B^+ \rightarrow \gamma J/\psi K^+$
--------	----	-----------------------------	------	-------------------------------------

¹ BHARDWAJ 11 reports $[\Gamma(X(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R7;LINKAGE=BA

² AUBERT 09B reports $[\Gamma(X(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R7;LINKAGE=AB

³ Superseded by AUBERT 09B. AUBERT, BE 06M reports $[\Gamma(X(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R7;LINKAGE=AU

$\Gamma(\gamma \psi(2S))/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	36 \pm 9	¹ AAIJ 14AH	LHCB	$B^+ \rightarrow \gamma \psi(2S) K^+$
>0.030	25 \pm 7	² AUBERT 09B	BABR	$B^+ \rightarrow \gamma \psi(2S) K^+$

NODE=M176R10
NODE=M176R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		³ BHARDWAJ 11	BELL	$B^+ \rightarrow \gamma \psi(2S) K^+$
----------	--	--------------------------	------	---------------------------------------

¹ From 36.4 ± 9 events of $X(3872) \rightarrow J/\psi \gamma$ decays with a statistical significance of 4.4σ .

NODE=M176R10;LINKAGE=A

² AUBERT 09B reports $[\Gamma(X(3872) \rightarrow \gamma \psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow X(3872)K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+) < 3.2 \times 10^{-4}$.

NODE=M176R10;LINKAGE=AU

³ BHARDWAJ 11 reports $B(B^+ \rightarrow K^+ X(3872)) \times B(X \rightarrow \gamma \psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.

NODE=M176R10;LINKAGE=BH

$\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$ Γ_{13}/Γ_{12}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

2.6 ± 0.6 OUR AVERAGE2.46 ± 0.64 ± 0.29 36 ± 9 ¹ AAIJ 14AH LHCB $B^+ \rightarrow \gamma\psi(2S)K^+$ 3.4 ± 1.4 AUBERT 09B BABR $B^+ \rightarrow \gamma c\bar{c}K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1 90 BHARDWAJ 11 BELL $B^+ \rightarrow \gamma\psi(2S)K^+$ ¹ From 36.4 ± 9 events of $X(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ.NODE=M176R11
NODE=M176R11

NODE=M176R11;LINKAGE=A

 $\Gamma(\rho\bar{\rho})/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_{15}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10⁻³ 95 ¹ AAIJ 13S LHCB $B^+ \rightarrow \rho\bar{\rho}K^+$ ¹ AAIJ 13S reports $[\Gamma(X(3872) \rightarrow \rho\bar{\rho})/\Gamma(X(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] \times [B(B^+ \rightarrow X(3872)K^+, X \rightarrow J/\psi\pi^+\pi^-)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+, X \rightarrow J/\psi\pi^+\pi^-) = 8.6 \times 10^{-6}$.NODE=M176R02
NODE=M176R02

NODE=M176R02;LINKAGE=A

C-violating decays $\Gamma(\eta J/\psi)/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_{16}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.4 (CL = 90%) [<0.6 (CL = 90%) OUR 2015 BEST LIMIT]<0.4 90 ^{1,2} IWASHITA 14 BELL $B \rightarrow K\eta J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6 90 AUBERT 04Y BABR $B \rightarrow K\eta J/\psi$ ¹ IWASHITA 14 reports $[\Gamma(X(3872) \rightarrow \eta J/\psi)/\Gamma(X(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] \times [B(B^+ \rightarrow X(3872)K^+, X \rightarrow J/\psi\pi^+\pi^-)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow X(3872)K^+, X \rightarrow J/\psi\pi^+\pi^-) = 8.6 \times 10^{-6}$.² IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow X(3872)K^\pm) \times B(X(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals.NODE=M176R2
NODE=M176R2

NODE=M176405

NODE=M176R2;LINKAGE=A

NODE=M176R2;LINKAGE=C

X(3872) REFERENCES

AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56771
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56787
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)	REFID=56408
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55897
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55647
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)	REFID=55925
AAIJ	13Q	PRL 110 222001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=54985
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55008
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=55412
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54056
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54751
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=53779
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=53934
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=53225
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53362
AALTONEN	09AU	PRL 103 152001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53098
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52722
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52265
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017
AUBERT_BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51454
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)	REFID=51432
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50498
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50627
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=50458
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=50200
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=49742
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49997
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>		REFID=49677
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=49628
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074

NODE=M176

X(3900)

$$I^G(J^{PC}) = 1^+(1^{+-})$$

Charged X(3900) seen as a peak in the invariant mass distribution of the $J/\psi\pi^\pm$ system by BES III (ABLIKIM 13T) in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at c.m. energy of 4.26 GeV and by radiative return from e^+e^- collisions at \sqrt{s} from 9.46 to 10.86 GeV at Belle (LIU 13B). Angular analysis of ABLIKIM 14A and ABLIKIM 15AC favor the $J^P = 1^+$ assignment. Neutral X(3900) seen in the $J/\psi\pi^0$ invariant mass distribution in $e^+e^- \rightarrow \pi^0\pi^0J/\psi$ at c.m. energies of 4.23, 4.26, and 4.36 GeV by BES III (ABLIKIM 15U) and at 4.17 GeV by XIAO 13A. Peaks in $(D\bar{D}^*)^{0,\pm}$ reported by BES III (ABLIKIM 14A, ABLIKIM 15AB) are assumed to be related.

NODE=M210

NODE=M210

X(3900) MASS

NODE=M210M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3886.6 ± 2.4 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.			
[3888.7 ± 3.4 MeV OUR 2015 AVERAGE		Scale factor = 1.3]			
3885.7 ^{+4.3} _{-5.7} ± 8.4		¹ ABLIKIM 15AB	BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
3881.7 ± 1.6 ± 1.6	1248	¹ ABLIKIM 15AC	BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3894.8 ± 2.3 ± 3.2	356	¹ ABLIKIM 15U	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0J/\psi$
3883.9 ± 1.5 ± 4.2	1212	¹ ABLIKIM 14A	BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3899.0 ± 3.6 ± 4.9	307	¹ ABLIKIM 13T	BES3	±	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
3894.5 ± 6.6 ± 4.5	159	¹ LIU 13B	BELL	±	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
3886 ± 4 ± 2	81	^{1,2} XIAO 13A		±	4.17 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$
3904 ± 9 ± 5	25	^{1,2} XIAO 13A		0	4.17 $e^+e^- \rightarrow \pi^0\pi^0J/\psi$

NODE=M210M

NEW


OCCUR=2

¹ Neglecting interference between the X(3900) and non-resonant continuum.

² For $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

NODE=M210M;LINKAGE=A

NODE=M210M;LINKAGE=B



X(3900) WIDTH

NODE=M210W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
28.1 ± 2.6 OUR AVERAGE		[35 ± 7 MeV OUR 2015 AVERAGE]			
35 ⁺¹¹ ₋₁₂ ± 15		¹ ABLIKIM 15AB	BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
26.6 ± 2.0 ± 2.1	1248	¹ ABLIKIM 15AC	BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$

NODE=M210W

NEW

29.6 ± 8.2 ± 8.2	356	¹ ABLIKIM	15U	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
24.8 ± 3.3 ± 11.0	1212	¹ ABLIKIM	14A	BES3	±	$e^+e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$
46 ± 10 ± 20	307	¹ ABLIKIM	13T	BES3	±	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
63 ± 24 ± 26	159	¹ LIU	13B	BELL	±	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
37 ± 4 ± 8	81	^{1,2} XIAO	13A		±	$4.17 e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ Neglecting interference between the $X(3900)$ and non-resonant continuum.

² For $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

NODE=M210W;LINKAGE=A
NODE=M210W;LINKAGE=B

X(3900) DECAY MODES

NODE=M210215;NODE=M210

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\pi$	seen
Γ_2 $h_c\pi^\pm$	not seen
Γ_3 $(D\bar{D}^*)^\pm$	seen
Γ_4 $D^0 D^{*-} + \text{c.c.}$	seen
Γ_5 $D^- D^{*0} + \text{c.c.}$	seen
Γ_6 $\omega\pi^\pm$	not seen
Γ_7 $J/\psi\eta$	not seen
Γ_8 $D^+ D^{*-} + \text{c.c.}$	seen
Γ_9 $D^0 \bar{D}^{*0} + \text{c.c.}$	seen

DESIG=1
DESIG=2
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=8
DESIG=9
DESIG=4
DESIG=5
DESIG=6
DESIG=7

X(3900) BRANCHING RATIOS

NODE=M210225

$\Gamma(J/\psi\pi)/\Gamma_{\text{total}}$							Γ_1/Γ	
VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
seen		356	ABLIKIM	15U	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$	
seen		307	ABLIKIM	13T	BES3	±	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
seen		25	¹ XIAO	13A	0	$4.17 e^+e^- \rightarrow \pi^0\pi^0 J/\psi$		

NODE=M210R01
NODE=M210R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen 90 ² ADOLPH 15D COMP ± $\gamma N \rightarrow J/\psi\pi^\pm N$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² ADOLPH 15D measure $B(X(3900)^\pm \rightarrow J/\psi\pi^\pm)\sigma(\gamma N \rightarrow X(3900)^\pm N)/\sigma(\gamma N \rightarrow J/\psi N) < 3.7 \times 10^{-3}$ at 90% CL.

NODE=M210R01;LINKAGE=XI
NODE=M210R01;LINKAGE=A

$\Gamma(h_c\pi^\pm)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM	13X	BES3	±	$e^+e^- \rightarrow h_c\pi^+\pi^-$	

NODE=M210R02
NODE=M210R02

$\Gamma((D\bar{D}^*)^\pm)/\Gamma(J/\psi\pi)$						Γ_3/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
6.2 ± 1.1 ± 2.7	¹ ABLIKIM	14A	BES3	±	$e^+e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$	

NODE=M210R03
NODE=M210R03

¹ Assuming the same origin of the $(D\bar{D}^*)^\pm$ and $\pi^\pm J/\psi$ decay modes.

NODE=M210R03;LINKAGE=A

$\Gamma(D^0 D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_4/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
seen	ABLIKIM	15AC	BES3	±	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$	
seen	ABLIKIM	14A	BES3	±	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$	

NODE=M210R09
NODE=M210R09

$\Gamma(D^- D^{*0} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_5/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
seen	ABLIKIM	15AC	BES3	±	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$	
seen	ABLIKIM	14A	BES3	±	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$	

NODE=M210R10
NODE=M210R10

$\Gamma(\omega\pi^\pm)/\Gamma_{\text{total}}$						Γ_6/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM	15R	BES3	±	$e^+e^- \rightarrow \omega\pi^+\pi^-$	

NODE=M210R00
NODE=M210R00

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$						Γ_7/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM	15Q	BES3	0	$4.0-4.6 e^+e^- \rightarrow J/\psi\eta\pi^0$	

NODE=M210R04
NODE=M210R04

$\Gamma(J/\psi\eta)/\Gamma(J/\psi\pi)$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.15	90	ABLIKIM	15Q	BES3	0 4.226 $e^+e^- \rightarrow J/\psi\eta\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.65	90	ABLIKIM	15Q	BES3	0 4.257 $e^+e^- \rightarrow J/\psi\eta\pi^0$

 Γ_7/Γ_1 NODE=M210R05
NODE=M210R05

OCCUR=2

 $\Gamma(D^+D^{*-} + c.c.)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AB	BES3	0 $e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

 Γ_8/Γ NODE=M210R06
NODE=M210R06 $\Gamma(D^0\bar{D}^{*0} + c.c.)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AB	BES3	0 $e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

 Γ_9/Γ NODE=M210R07
NODE=M210R07 $\Gamma(D^+D^{*-} + c.c.)/\Gamma(D^0\bar{D}^{*0} + c.c.)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.96±0.18±0.12	ABLIKIM	15AB	BES3	0 $e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

 Γ_8/Γ_9 NODE=M210R08
NODE=M210R08

X(3900) REFERENCES

ABLIKIM	15AB	PRL 115 222002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56954
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BES III Collab.) JP	REFID=56967
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56782
ABLIKIM	15R	PR D92 032009	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56783
ABLIKIM	15U	PRL 115 112003	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56786
ADOLPH	15D	PL B742 330	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56791
ABLIKIM	14A	PRL 112 022001	M. Ablikim <i>et al.</i>	(BES III Collab.) JP	REFID=55648
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55409
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55635
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=55410
XIAO	13A	PL B727 366	T. Xiao <i>et al.</i>	(NWES)	REFID=55593

NODE=M210

REFID=56954
REFID=56967
REFID=56782
REFID=56783
REFID=56786
REFID=56791
REFID=55648
REFID=55409
REFID=55635
REFID=55410
REFID=55593

NODE=M159

X(3915)
was $\chi_{c0}(3915)$

$$J^{PC} = 0^+(0^{++})$$

The experimental analysis prefers $J^{PC} = 0^{++}$. However, a re-analysis presented in ZHOU 15C shows that if helicity-2 dominance assumption is abandoned and a sizable helicity-0 component is allowed, a $J^{PC} = 2^{++}$ assignment is possible.

NODE=M159

X(3915) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3918.4±1.9 OUR AVERAGE				
3919.4±2.2±1.6	59±10	LEES	12AD	BABR $e^+e^- \rightarrow e^+e^- \omega J/\psi$
3919.1 ^{+3.8} _{-3.4} ±2.0		DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3915 ± 3 ± 2	49±15	UEHARA	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- \omega J/\psi$
3943 ±11 ±13	58±11	¹ CHOI	05	BELL $B \rightarrow \omega J/\psi K$
3914.6 ^{+3.8} _{-3.4} ±2.0		¹ AUBERT	08W	BABR Superseded by DEL-AMO-SANCHEZ 10B

NODE=M159M

NODE=M159M

¹ $\omega J/\psi$ threshold enhancement fitted as an S-wave Breit-Wigner resonance.

NODE=M159M;LINKAGE=CH

X(3915) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20±5 OUR AVERAGE	Error	includes scale factor of 1.1.		
13±6±3	59±10	LEES	12AD	BABR $e^+e^- \rightarrow e^+e^- \omega J/\psi$
31 ⁺¹⁰ ₋₈ ±5		DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
17±10±3	49±15	UEHARA	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- \omega J/\psi$
87±22±26	58±11	² CHOI	05	BELL $B \rightarrow \omega J/\psi K$
34 ⁺¹² ₋₈ ±5		² AUBERT	08W	BABR Superseded by DEL-AMO-SANCHEZ 10B

NODE=M159W

NODE=M159W

² $\omega J/\psi$ threshold enhancement fitted as an S-wave Breit-Wigner resonance.

NODE=M159W;LINKAGE=CH

X(3915) DECAY MODES

NODE=M159215;NODE=M159

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega J/\psi$	seen
Γ_2 $\bar{D}^{*0} D^0$	
Γ_3 $\pi^+ \pi^- \eta_c(1S)$	not seen
Γ_4 $\eta_c \eta$	not seen
Γ_5 $\eta_c \pi^0$	not seen
Γ_6 $K \bar{K}$	not seen
Γ_7 $\gamma \gamma$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=3
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=6
DESIG=7
DESIG=5;OUR EVAL;→ UNCHECKED ←
DESIG=2

X(3915) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M159220

$\Gamma(\omega J/\psi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_1\Gamma_7/\Gamma$**

NODE=M159G01
NODE=M159G01

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

54± 9 OUR AVERAGE

52±10±3	59 ± 10	³ LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
61±17±8	49 ± 15	³ UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

18± 5±2	49 ± 15	⁴ UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
---------	---------	---------------------	---------	--

NODE=M159G01;LINKAGE=UH
NODE=M159G01;LINKAGE=UR

³For $J^P = 0^+$.

⁴For $J^P = 2^+$, helicity-2.

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_3\Gamma_7/\Gamma$**

NODE=M159G02
NODE=M159G02

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<16	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$
-----	----	------	-----------	--

$\Gamma(K \bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_6\Gamma_7/\Gamma$**

NODE=M159G03
NODE=M159G03

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.96	90	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
-------	----	--------	---------	--

X(3915) BRANCHING RATIOS

NODE=M159225

$\Gamma(\omega J/\psi)/\Gamma_{\text{total}}$ **Γ_1/Γ**

NODE=M159R03
NODE=M159R03

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	⁵ DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
------	------------------------------	------	---------------------------------

seen	⁶ CHOI	05 BELL	$B \rightarrow \omega J/\psi K$
------	-------------------	---------	---------------------------------

⁵DEL-AMO-SANCHEZ 10B reports $B(B^\pm \rightarrow X(3915)K^\pm) \times B(X(3915) \rightarrow J/\psi\omega) = (3.0_{-0.6}^{+0.7+0.5}) \times 10^{-5}$ and $B(B^0 \rightarrow X(3915)K^0) \times B(X(3915) \rightarrow J/\psi\omega) = (2.1 \pm 0.9 \pm 0.3) \times 10^{-5}$.

NODE=M159R03;LINKAGE=DE

⁶CHOI 05 reports $B(B \rightarrow X(3915)K) \times B(X(3915) \rightarrow J/\psi\omega) = (7.1 \pm 1.3 \pm 3.1) \times 10^{-5}$.

NODE=M159R03;LINKAGE=CH

$\Gamma(\omega J/\psi)/\Gamma(\bar{D}^{*0} D^0)$ **Γ_1/Γ_2**

NODE=M159R02
NODE=M159R02

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

>0.71	90	⁷ AUSHEV	10 BELL	$B \rightarrow \bar{D}^{*0} D^0 K$
-------	----	---------------------	---------	------------------------------------

⁷By combining the upper limit $B(B \rightarrow X(3915)K) \times B(X(3915) \rightarrow \bar{D}^{*0} D^0) < 0.67 \times 10^{-4}$ from AUSHEV 10 with the average of CHOI 05 and AUBERT 08W measurements $B(B \rightarrow X(3915)K) \times B(X(3915) \rightarrow \omega J/\psi) = (0.51 \pm 0.11) \times 10^{-4}$.

NODE=M159R02;LINKAGE=AU

$\Gamma(\eta_c \eta)/\Gamma_{\text{total}}$ **Γ_4/Γ**

NODE=M159R00
NODE=M159R00

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	⁸ VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \eta$
----------	----------------------------	------	-----------------------------------

⁸VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ X(3915)^0) \times B(X \rightarrow \eta_c \eta) < 3.3 \times 10^{-5}$ at 90% CL.

NODE=M159R00;LINKAGE=VI

$\Gamma(\eta_c \pi^0)/\Gamma_{\text{total}}$ **Γ_5/Γ**

NODE=M159R04
NODE=M159R04

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	⁹ VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \pi^0$
----------	----------------------------	------	------------------------------------

⁹VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ X(3915)^0) \times B(X \rightarrow \eta_c \pi^0) < 1.8 \times 10^{-5}$ at 90% CL.

NODE=M159R04;LINKAGE=VI

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	59 ± 10	LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
seen		UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$

 Γ_7/Γ

NODE=M159R01
 NODE=M159R01

X(3915) REFERENCES

VINOKUROVA 15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
ZHOU 15C	PRL 115 022001	Z.-Y. Zhou, Z. Xiao, H.-Q. Zhou	(BEIJT, NANJ)
UEHARA 13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
LEES 12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES 12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUSHEV 10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
UEHARA 10	PRL 104 092001	S. Uehara <i>et al.</i>	(BELLE Collab.)
AUBERT 08W	PRL 101 082001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHOI 05	PRL 94 182002	S.-K. Choi <i>et al.</i>	(BELLE Collab.)

NODE=M159

REFID=56706
 REFID=56842
 REFID=55592
 REFID=54751
 REFID=54752
 REFID=53225
 REFID=53362
 REFID=53232
 REFID=52263
 REFID=50737

 $\chi_{c2}(2P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M050

 $\chi_{c2}(2P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3927.2 ± 2.6 OUR AVERAGE				
3926.7 ± 2.7 ± 1.1	76 ± 17	AUBERT	10G BABR	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$
3929 ± 5 ± 2	64	UEHARA	06 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$

NODE=M050M

NODE=M050M

 $\chi_{c2}(2P)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
24 ± 6 OUR AVERAGE				
21.3 ± 6.8 ± 3.6	76 ± 17	AUBERT	10G BABR	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$
29 ± 10 ± 2	64	UEHARA	06 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$

NODE=M050W

NODE=M050W

 $\chi_{c2}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\gamma\gamma$	seen
Γ_2 $K\bar{K}\pi$	
Γ_3 $K^+K^-\pi^+\pi^-\pi^0$	
Γ_4 $D\bar{D}$	seen
Γ_5 D^+D^-	seen
Γ_6 $D^0\bar{D}^0$	seen
Γ_7 $\pi^+\pi^-\eta_c(1S)$	not seen
Γ_8 $K\bar{K}$	not seen

NODE=M050215;NODE=M050

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=5
 DESIG=6
 DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=3;OUR EVAL;→ UNCHECKED ←
 DESIG=4;OUR EVAL;→ UNCHECKED ←
 DESIG=7;OUR EVAL;→ UNCHECKED ←
 DESIG=8;OUR EVAL;→ UNCHECKED ←

 $\chi_{c2}(2P)$ PARTIAL WIDTHS

$$\chi_{c2}(2P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$$

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	DEL-AMO-SA...11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

 $\Gamma_2\Gamma_1/\Gamma$

NODE=M050G01
 NODE=M050G01

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<3.4	90	DEL-AMO-SA...11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

 $\Gamma_3\Gamma_1/\Gamma$

NODE=M050G02
 NODE=M050G02

 $\Gamma(D\bar{D}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.21 ± 0.04 OUR AVERAGE				
0.24 ± 0.05 ± 0.04	76 ± 17	AUBERT	10G BABR	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$
0.18 ± 0.05 ± 0.03	64	¹ UEHARA	06 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$

 $\Gamma_4\Gamma_1/\Gamma$

NODE=M050G1
 NODE=M050G1

¹ Assuming $B(D^+D^-) = 0.89 B(D^0\bar{D}^0)$.

NODE=M050G1;LINKAGE=UE

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<18	90	LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

NODE=M050G03
NODE=M050G03 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_8\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.256	90	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M050G04
NODE=M050G04 $\chi_{c2}(2P)$ BRANCHING RATIOS

NODE=M050225

 $\Gamma(D^+D^-)/\Gamma(D^0\bar{D}^0)$ Γ_5/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.74 \pm 0.43 \pm 0.16$	64	UEHARA	06 BELL	$10.6 e^+e^- \rightarrow e^+e^-D\bar{D}$

NODE=M050R01
NODE=M050R01 $\chi_{c2}(2P)$ REFERENCES

NODE=M050

UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
AUBERT	10G	PR D81 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53357
UEHARA	06	PRL 96 082003	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=51039

NODE=M029

X(3940)

$$J^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

Reported by ABE 07, observed in $e^+e^- \rightarrow J/\psi X$.

NODE=M029

X(3940) MASS

NODE=M029M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3942^{+7}_{-6} \pm 6$	52	PAKHLOV	08 BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M029M

••• We do not use the following data for averages, fits, limits, etc. •••

$3943 \pm 6 \pm 6$	25	¹ ABE	07 BELL	$e^+e^- \rightarrow J/\psi X$
3936 ± 14	266	² ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$

OCCUR=2

¹ From a fit to $D^{*+}D^-$ and $D^{*0}\bar{D}^0$ events.² From the inclusive fit. Not independent of the exclusive measurement by ABE 07.NODE=M029M;LINKAGE=EB
NODE=M029M;LINKAGE=EM**X(3940) WIDTH**

NODE=M029W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$37^{+26}_{-15} \pm 8$		52	PAKHLOV	08 BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M029W

••• We do not use the following data for averages, fits, limits, etc. •••

<52	90	25	ABE	07 BELL	$e^+e^- \rightarrow J/\psi X$
-----	----	----	-----	---------	-------------------------------

X(3940) DECAY MODES

NODE=M029215;NODE=M029

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\bar{D}^* + \text{c.c.}$	seen
Γ_2 $D\bar{D}$	not seen
Γ_3 $J/\psi\omega$	not seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

X(3940) BRANCHING RATIOS

NODE=M029225

 $\Gamma(D\bar{D}^* + \text{c.c.})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
>0.45	90	25	^{3,4} ABE	07 BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M029R01
NODE=M029R01

••• We do not use the following data for averages, fits, limits, etc. •••

³ For X(3940) decaying to final states with more than two tracks.⁴ PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c² may consist of several states.NODE=M029R01;LINKAGE=AB
NODE=M029R01;LINKAGE=AE

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.41	90	5,6 ABE	07	BELL $e^+e^- \rightarrow J/\psi X$
-------	----	---------	----	------------------------------------

⁵ For $X(3940)$ decaying to final states with more than two tracks.

⁶ PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c² may consist of several states.

NODE=M029R02
NODE=M029R02

NODE=M029R02;LINKAGE=AB
NODE=M029R02;LINKAGE=AE

 $\Gamma(J/\psi\omega)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.26	90	7,8 ABE	07	BELL $e^+e^- \rightarrow J/\psi X$
-------	----	---------	----	------------------------------------

⁷ For $X(3940)$ decaying to final states with more than two tracks.

⁸ PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c² may consist of several states.

NODE=M029R03
NODE=M029R03

NODE=M029R03;LINKAGE=AB
NODE=M029R03;LINKAGE=AE

X(3940) REFERENCES

PAKHLOV	08	PRL 100 202001	P. Pakhlov <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M029

REFID=52302
REFID=51627

NODE=M213

X(4020)

$$I(J^P) = 1(?^?)$$

Charged X(4020) seen by ABLIKIM 13X from $e^+e^- \rightarrow \pi^+\pi^-h_c(1P)$ at c.m. energy from 3.90 to 4.42 GeV as a peak in the invariant mass distribution of the $\pi^\pm h_c(1P)$ system, and by ABLIKIM 14B from $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ events in $(D^*\bar{D}^*)^\pm$ mass. A neutral X(4020) seen by ABLIKIM 14P at three c.m. energies in the same range in $e^+e^- \rightarrow \pi^0\pi^0h_c(1P)$ as a peak in the larger of the two masses recoiling against a π^0 . ABLIKIM 15AA observes a 5.9σ signal in $(D^*\bar{D}^*)^0$ in $e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$ events using collisions at two c.m. energies. Production rates and mass values support grouping neutral and charged X(4020) together as manifestations of a single $I = 1$ particle.

NODE=M213

X(4020) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

4024.1 ± 1.9 OUR AVERAGE

[4023.9 ± 2.4 MeV OUR 2015 AVERAGE]

4025.5 ^{+2.0} _{-4.7} ± 3.1	116	¹ ABLIKIM	15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
--	-----	----------------------	------	------	---	--

4026.3 ± 2.6 ± 3.7	401	¹ ABLIKIM	14B	BES3	±	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
--------------------	-----	----------------------	-----	------	---	--

4023.9 ± 2.2 ± 3.8	61	^{1,2} ABLIKIM	14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0h_c$
--------------------	----	------------------------	-----	------	---	------------------------------------

4022.9 ± 0.8 ± 2.7	253	¹ ABLIKIM	13X	BES3	±	$e^+e^- \rightarrow \pi^+\pi^-h_c$
--------------------	-----	----------------------	-----	------	---	------------------------------------

¹ Neglecting interference between the X(4020) and non-resonant continuum.

² Assuming $J^P = 1^+$ and width of 7.9 ± 2.6 MeV.

NODE=M213M

NODE=M213M
NEW

NODE=M213M;LINKAGE=AB
NODE=M213M;LINKAGE=B

X(4020) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

13 ± 5 OUR AVERAGE

Error includes scale factor of 1.7. See the ideogram below.
[10 ± 6 MeV OUR 2015 AVERAGE Scale factor = 1.7]

23.0 ± 6.0 ± 1.0	116	¹ ABLIKIM	15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
------------------	-----	----------------------	------	------	---	--

24.8 ± 5.6 ± 7.7	401	¹ ABLIKIM	14B	BES3	±	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
------------------	-----	----------------------	-----	------	---	--


7.9 ± 2.7 ± 2.6	253	¹ ABLIKIM	13X	BES3	±	$e^+e^- \rightarrow \pi^+\pi^-h_c$
-----------------	-----	----------------------	-----	------	---	------------------------------------

NODE=M213W

NODE=M213W
NEW

¹ Neglecting interference between the $X(4020)$ and non-resonant continuum.

NODE=M213W;LINKAGE=AB



X(4020) DECAY MODES

NODE=M213215;NODE=M213

Mode	Fraction (Γ_i/Γ)
Γ_1 $h_c(1P)\pi$	seen
Γ_2 $D^*\bar{D}^*$	seen
Γ_3 $D\bar{D}^* + c.c.$	not seen
Γ_4 $\eta_c\pi^+\pi^-$	not seen

DESIG=1

DESIG=2

DESIG=4

DESIG=3

X(4020) BRANCHING RATIOS

NODE=M213225

$\Gamma(h_c(1P)\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
seen	61	ABLIKIM	14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 h_c$
seen	253	ABLIKIM	13X	BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^- h_c$

NODE=M213R01
NODE=M213R01

$\Gamma(D^*\bar{D}^*)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
seen	116	¹ ABLIKIM	15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
seen	401	¹ ABLIKIM	14B	BES3	\pm	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$

NODE=M213R02
NODE=M213R02

¹ Neglecting interference between the $X(4020)$ and non-resonant continuum.

NODE=M213R02;LINKAGE=A

$\Gamma(D\bar{D}^* + c.c.)/\Gamma_{\text{total}}$						Γ_3/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	15AC	BES3	\pm	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$

NODE=M213R03
NODE=M213R03

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_4/Γ
VALUE		DOCUMENT ID	TECN		COMMENT	
not seen		¹ VINOKUROVA	15	BELL	$B^+ \rightarrow K^+\eta_c\pi^+\pi^-$	

NODE=M213R00
NODE=M213R00

¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+X(4020)^0) \times B(X \rightarrow \eta_c\pi^+\pi^-) < 1.6 \times 10^{-5}$ at 90% CL.

NODE=M213R00;LINKAGE=V1

X(4020) REFERENCES

NODE=M213

ABLIKIM	15AA	PRL 115 182002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56951
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56967
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
ABLIKIM	14B	PRL 112 132001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55654
ABLIKIM	14P	PRL 113 212002	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56118
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=55635

$\psi(4040)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M072

 $\psi(4040)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4039 ± 1 OUR ESTIMATE			
4039.6 ± 4.3	¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4034 ± 6	² MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4037 ± 2	³ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4040 ± 1	⁴ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4040 ± 10	BRANDELIK	78C DASP	e^+e^-
¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
³ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁴ From a fit to BES (BAI 02C) data.			

NODE=M072M

NODE=M072M
→ UNCHECKED ←

OCCUR=2

NODE=M072M;LINKAGE=AB

NODE=M072M;LINKAGE=MO

NODE=M072M;LINKAGE=ST
NODE=M072M;LINKAGE=SE $\psi(4040)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
80 ± 10 OUR ESTIMATE			
84.5 ± 12.3	⁵ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
87 ± 11	⁶ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
85 ± 10	⁷ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
89 ± 6	⁸ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
52 ± 10	BRANDELIK	78C DASP	e^+e^-
⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
⁶ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁷ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁸ From a fit to BES (BAI 02C) data.			

NODE=M072W

NODE=M072W
→ UNCHECKED ←

OCCUR=2

NODE=M072W;LINKAGE=AB

NODE=M072W;LINKAGE=MO

NODE=M072W;LINKAGE=ST
NODE=M072W;LINKAGE=SE $\psi(4040)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

NODE=M072215;NODE=M072

NODE=M072

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 e^+e^-	$(1.07 \pm 0.16) \times 10^{-5}$	DESIG=5
Γ_2 $D\bar{D}$	seen	DESIG=17;OUR EST;→ UNCHECKED ←
Γ_3 $D^0\bar{D}^0$	seen	DESIG=1
Γ_4 D^+D^-	seen	DESIG=18
Γ_5 $D^*\bar{D} + \text{c.c.}$	seen	DESIG=19;OUR EST;→ UNCHECKED ←
Γ_6 $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	DESIG=2
Γ_7 $D^*(2010)^+D^- + \text{c.c.}$	seen	DESIG=20
Γ_8 $D^*\bar{D}^*$	seen	DESIG=21;OUR EST;→ UNCHECKED ←
Γ_9 $D^*(2007)^0\bar{D}^*(2007)^0$	seen	DESIG=3
Γ_{10} $D^*(2010)^+D^*(2010)^-$	seen	DESIG=22
Γ_{11} $D\bar{D}\pi$ (excl. $D^*\bar{D}$)		DESIG=23
Γ_{12} $D^0D^-\pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0\bar{D}^0 + \text{c.c.}$, $D^*(2010)^+D^- + \text{c.c.}$)	not seen	DESIG=24
Γ_{13} $D\bar{D}^*\pi$ (excl. $D^*\bar{D}^*$)	not seen	DESIG=25

Γ ₁₄	$D^0 \bar{D}^{*-} \pi^+ + c.c. (excl. D^*(2010)^+ D^*(2010)^-)$	seen			DESIG=26
Γ ₁₅	$D_s^+ D_s^-$	seen			DESIG=27
Γ ₁₆	$J/\psi(1S)$ hadrons				DESIG=4
Γ ₁₇	$J/\psi \pi^+ \pi^-$	< 4	$\times 10^{-3}$	90%	DESIG=7
Γ ₁₈	$J/\psi \pi^0 \pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=8
Γ ₁₉	$J/\psi \eta$	(5.2 ± 0.7)	$\times 10^{-3}$		DESIG=9
Γ ₂₀	$J/\psi \pi^0$	< 2.8	$\times 10^{-4}$	90%	DESIG=10
Γ ₂₁	$J/\psi \pi^+ \pi^- \pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=11
Γ ₂₂	$\chi_{c1} \gamma$	< 3.4	$\times 10^{-3}$	90%	DESIG=12
Γ ₂₃	$\chi_{c2} \gamma$	< 5	$\times 10^{-3}$	90%	DESIG=13
Γ ₂₄	$\chi_{c1} \pi^+ \pi^- \pi^0$	< 1.1	%	90%	DESIG=14
Γ ₂₅	$\chi_{c2} \pi^+ \pi^- \pi^0$	< 3.2	%	90%	DESIG=15
Γ ₂₆	$h_c(1P) \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%	DESIG=28
Γ ₂₇	$\phi \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%	DESIG=16
Γ ₂₈	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	< 2.9	$\times 10^{-4}$	90%	DESIG=29
Γ ₂₉	$\Lambda \bar{\Lambda} \pi^0$	< 9	$\times 10^{-5}$	90%	DESIG=30
Γ ₃₀	$\Lambda \bar{\Lambda} \eta$	< 3.0	$\times 10^{-4}$	90%	DESIG=31
Γ ₃₁	$\Sigma^+ \bar{\Sigma}^-$	< 1.3	$\times 10^{-4}$	90%	DESIG=32
Γ ₃₂	$\Sigma^0 \bar{\Sigma}^0$	< 7	$\times 10^{-5}$	90%	DESIG=33
Γ ₃₃	$\Xi^+ \bar{\Xi}^-$	< 1.6	$\times 10^{-4}$	90%	DESIG=34
Γ ₃₄	$\Xi^0 \bar{\Xi}^0$	< 1.8	$\times 10^{-4}$	90%	DESIG=35
Γ ₃₅	$\mu^+ \mu^-$				DESIG=6

ψ(4040) PARTIAL WIDTHS

NODE=M072220

Γ(e⁺e⁻)

Γ₁

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.07 OUR ESTIMATE			
0.83 ± 0.20	⁹ ABLIKIM	08D BES2	e ⁺ e ⁻ → hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.6 to 1.4	¹⁰ MO	10 RVUE	e ⁺ e ⁻ → hadrons
0.88 ± 0.11	¹¹ SETH	05A RVUE	e ⁺ e ⁻ → hadrons
0.91 ± 0.13	¹² SETH	05A RVUE	e ⁺ e ⁻ → hadrons
0.75 ± 0.15	BRANDELIK	78C DASP	e ⁺ e ⁻

NODE=M072W5
NODE=M072W5
→ UNCHECKED ←

OCCUR=2

⁹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the ψ(3770), ψ(4040), ψ(4160), and ψ(4415) resonances. Phase angle fixed in the fit to δ = (130 ± 46)°.

NODE=M072W5;LINKAGE=AB

¹⁰ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the ψ(4040), ψ(4160) and ψ(4415) resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e⁺e⁻ partial widths. We quote only the range of values.

NODE=M072W5;LINKAGE=MO

¹¹ From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M072W5;LINKAGE=ST

¹² From a fit to BES (BAI 02C) data.

NODE=M072W5;LINKAGE=SE

ψ(4040) Γ(i) × Γ(e⁺e⁻)/Γ(total)

NODE=M072235

Γ(χ_{c1}γ) × Γ(e⁺e⁻)/Γ_{total}

Γ₂₂Γ₁/Γ

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	¹³ HAN	15 BELL	10.58 e ⁺ e ⁻ → χ _{c1} γ

NODE=M072G01
NODE=M072G01

¹³ Using B(η → γγ) = (39.41 ± 0.21)%.

NODE=M072G01;LINKAGE=A

Γ(χ_{c2}γ) × Γ(e⁺e⁻)/Γ_{total}

Γ₂₃Γ₁/Γ

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<4.6	90	¹⁴ HAN	15 BELL	10.58 e ⁺ e ⁻ → χ _{c2} γ

NODE=M072G02
NODE=M072G02

¹⁴ Using B(η → γγ) = (39.41 ± 0.21)%.

NODE=M072G02;LINKAGE=A

$\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$

NODE=M072230

 $\Gamma(J/\psi\eta) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{19} / \Gamma \times \Gamma_1 / \Gamma$ NODE=M072R25
NODE=M072R25VALUE (units 10^{-8})

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $5.1 \pm 1.4 \pm 1.5$

15 WANG

13B

BELL

 $e^+ e^- \rightarrow J/\psi \eta \gamma$ $12.8 \pm 2.1 \pm 1.9$

16 WANG

13B

BELL

 $e^+ e^- \rightarrow J/\psi \eta \gamma$

OCCUR=2

¹⁵ Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

NODE=M072R25;LINKAGE=A

¹⁶ Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

NODE=M072R25;LINKAGE=B

 $\psi(4040)$ BRANCHING RATIOS

NODE=M072225

 $\Gamma(e^+ e^-) / \Gamma_{\text{total}}$ Γ_1 / Γ NODE=M072R4
NODE=M072R4VALUE (units 10^{-5})

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

 ~ 1.0

FELDMAN

77

MRK1

 $e^+ e^-$ $\Gamma(D^0 \bar{D}^0) / \Gamma_{\text{total}}$ Γ_3 / Γ NODE=M072R14
NODE=M072R14

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^0 \bar{D}^0$

seen

PAKHLOVA

08

BELL

 $e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$ $\Gamma(D^+ D^-) / \Gamma_{\text{total}}$ Γ_4 / Γ NODE=M072R15
NODE=M072R15

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^+ D^- \gamma$

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^+ D^-$

seen

PAKHLOVA

08

BELL

 $e^+ e^- \rightarrow D^+ D^- \gamma$ $\Gamma(D \bar{D}) / \Gamma(D^* \bar{D} + \text{c.c.})$ Γ_2 / Γ_5 NODE=M072R12
NODE=M072R12

VALUE

DOCUMENT ID

TECN

COMMENT

 $0.24 \pm 0.05 \pm 0.12$

AUBERT

09M

BABR

 $e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}$ $\Gamma(D^0 \bar{D}^0) / \Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.})$ Γ_3 / Γ_6 NODE=M072R1
NODE=M072R1

VALUE

DOCUMENT ID

TECN

COMMENT

 0.05 ± 0.03

17 GOLDHABER

77

MRK1

 $e^+ e^-$ ¹⁷ Phase-space factor (p^3) explicitly removed.

NODE=M072R;LINKAGE=P

 $\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_6 / Γ NODE=M072R16
NODE=M072R16

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^{*0} \bar{D}^0 \gamma$

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^{*0} \bar{D}^0$ $\Gamma(D^*(2010)^+ D^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_7 / Γ NODE=M072R17
NODE=M072R17

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^{*+} D^- \gamma$

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^{*+} D^-$

seen

PAKHLOVA

07

BELL

 $e^+ e^- \rightarrow D^{*+} D^- \gamma$ $\Gamma(D^*(2010)^+ D^- + \text{c.c.}) / \Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.})$ Γ_7 / Γ_6 NODE=M072R11
NODE=M072R11

VALUE

DOCUMENT ID

TECN

COMMENT

 $0.95 \pm 0.09 \pm 0.10$

AUBERT

09M

BABR

 $e^+ e^- \rightarrow \gamma D^* \bar{D}$ $\Gamma(D^* \bar{D}^*) / \Gamma(D^* \bar{D} + \text{c.c.})$ Γ_8 / Γ_5 NODE=M072R13
NODE=M072R13

VALUE

DOCUMENT ID

TECN

COMMENT

 $0.18 \pm 0.14 \pm 0.03$

AUBERT

09M

BABR

 $e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$ $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0) / \Gamma_{\text{total}}$ Γ_9 / Γ NODE=M072R18
NODE=M072R18

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$

$$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0) / \Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.}) \quad \Gamma_9 / \Gamma_6$$

VALUE	DOCUMENT ID	TECN	COMMENT
32.0 ± 12.0	¹⁸ GOLDHABER 77	MRK1	$e^+ e^-$

NODE=M072R2
NODE=M072R2

¹⁸ Phase-space factor (p^3) explicitly removed.

NODE=M072R2;LINKAGE=P

$$\Gamma(D^*(2010)^+ D^*(2010)^-) / \Gamma_{\text{total}} \quad \Gamma_{10} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

NODE=M072R19
NODE=M072R19

$$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.})) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

NODE=M072R20
NODE=M072R20

$$\Gamma(D \bar{D}^* \pi \text{ (excl. } D^* \bar{D}^*)) / \Gamma_{\text{total}} \quad \Gamma_{13} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

NODE=M072R21
NODE=M072R21

$$\Gamma(D^0 \bar{D}^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-) / \Gamma_{\text{total}} \quad \Gamma_{14} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

NODE=M072R22
NODE=M072R22

$$\Gamma(D_s^+ D_s^-) / \Gamma_{\text{total}} \quad \Gamma_{15} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

NODE=M072R23
NODE=M072R23

$$\Gamma(J/\psi \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{17} / \Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	COAN 06	CLEO	3.97–4.06 $e^+ e^- \rightarrow$ hadrons

NODE=M072R01
NODE=M072R01

$$\Gamma(J/\psi \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{18} / \Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN 06	CLEO	3.97–4.06 $e^+ e^- \rightarrow$ hadrons

NODE=M072R02
NODE=M072R02

$$\Gamma(J/\psi \eta) / \Gamma_{\text{total}} \quad \Gamma_{19} / \Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
5.2 ± 0.5 ± 0.5		¹⁹ ABLIKIM 12K	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$

NODE=M072R03
NODE=M072R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	COAN 06	CLEO	3.97–4.06 $e^+ e^- \rightarrow$ hadrons
----	----	---------	------	---

¹⁹ ABLIKIM 12K measure $\sigma(e^+ e^- \rightarrow J/\psi \eta) = 32.1 \pm 2.8 \pm 1.3$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.

NODE=M072R03;LINKAGE=AB

$$\Gamma(J/\psi \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{20} / \Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.28	90	²⁰ ABLIKIM 12K	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$

NODE=M072R04
NODE=M072R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2	90	COAN 06	CLEO	3.97–4.06 $e^+ e^- \rightarrow$ hadrons
----	----	---------	------	---

²⁰ ABLIKIM 12K measure $\sigma(e^+ e^- \rightarrow J/\psi \pi^0) < 1.6$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.

NODE=M072R04;LINKAGE=AB

$$\Gamma(J/\psi \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{21} / \Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN 06	CLEO	3.97–4.06 $e^+ e^- \rightarrow$ hadrons

NODE=M072R05
NODE=M072R05

$$\Gamma(\chi_{c1} \gamma) / \Gamma_{\text{total}} \quad \Gamma_{22} / \Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	COAN 06	CLEO	3.97–4.06 $e^+ e^- \rightarrow$ hadrons

NODE=M072R06
NODE=M072R06

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17	90	COAN	06	CLEO $3.97\text{--}4.06 e^+ e^- \rightarrow \text{hadrons}$
-----	----	------	----	---

NODE=M072R07
NODE=M072R07

 $\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<11	90	COAN	06	CLEO $3.97\text{--}4.06 e^+ e^- \rightarrow \text{hadrons}$
-----	----	------	----	---

NODE=M072R08
NODE=M072R08

 $\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<32	90	COAN	06	CLEO $3.97\text{--}4.06 e^+ e^- \rightarrow \text{hadrons}$
-----	----	------	----	---

NODE=M072R09
NODE=M072R09

 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<3	90	²¹ PEDLAR	11	CLEO $e^+ e^- \rightarrow h_c(1P)\pi^+\pi^-$
----	----	----------------------	----	--

NODE=M072R24
NODE=M072R24

²¹ From several values of \sqrt{s} near the peak of the $\psi(4040)$, PEDLAR 11 measures $\sigma(e^+ e^- \rightarrow h_c(1P)\pi^+\pi^-) = 1.0 \pm 8.0 \pm 5.4 \pm 0.2$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M072R24;LINKAGE=PE

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<3	90	COAN	06	CLEO $3.97\text{--}4.06 e^+ e^- \rightarrow \text{hadrons}$
----	----	------	----	---

NODE=M072R10
NODE=M072R10

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2.9	90	²² ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$
------	----	-----------------------	-----	---------------------------------------

NODE=M072R26
NODE=M072R26

²² Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R26;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.9	90	²³ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$
------	----	-----------------------	-----	---------------------------------------

NODE=M072R27
NODE=M072R27

²³ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R27;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<3.0	90	²⁴ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$
------	----	-----------------------	-----	---------------------------------------

NODE=M072R28
NODE=M072R28

²⁴ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R28;LINKAGE=A

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.3	90	²⁵ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$
------	----	-----------------------	-----	---------------------------------------

NODE=M072R29
NODE=M072R29

²⁵ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R29;LINKAGE=A

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.7	90	²⁶ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$
------	----	-----------------------	-----	---------------------------------------

NODE=M072R30
NODE=M072R30

²⁶ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R30;LINKAGE=A

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.6	90	²⁷ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$
------	----	-----------------------	-----	---------------------------------------

NODE=M072R31
NODE=M072R31

²⁷ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R31;LINKAGE=A

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.8	90	²⁸ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$
------	----	-----------------------	-----	---------------------------------------

NODE=M072R32
NODE=M072R32

²⁸ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R32;LINKAGE=A

$\psi(4040)$ REFERENCES

HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
ABLIKIM	13Q	PR D87 112011	Ablikim M. <i>et al.</i>	(BES III Collab.)	REFID=55393
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
ABLIKIM	12K	PR D86 071101	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=54738
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=53114
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51075
SETH	05A	PR D72 017501	K.K. Seth		REFID=50813
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=51064
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22232
Also		ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)	REFID=11434

NODE=M072

 $X(4050)^\pm$

$$I(J^P) = ?(??)$$

OMITTED FROM SUMMARY TABLE

Observed by MIZUK 08 in the $\pi^+ \chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure.

NODE=M191

NODE=M191

 $X(4050)^\pm$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4051 \pm 14^{+20}_{-41}$	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

NODE=M191M

NODE=M191M

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M191M;LINKAGE=MI

 $X(4050)^\pm$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
82^{+21+47}_{-17-22}	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

NODE=M191W

NODE=M191W

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M191W;LINKAGE=MI

 $X(4050)^\pm$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+ \chi_{c1}(1P)$	seen

NODE=M191215;NODE=M191

DESIG=1

 $X(4050)^\pm$ BRANCHING RATIOS

$\Gamma(\pi^+ \chi_{c1}(1P))/\Gamma_{\text{total}}$	Γ_1/Γ		
seen	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$
not seen	² LEES	12B	BABR $B \rightarrow K\pi \chi_{c1}(1P)$

NODE=M191225

NODE=M191R01
NODE=M191R01

¹ With a product branching fraction measurement of $B(\bar{B}^0 \rightarrow K^- X(4050)^+) \times B(X(4050)^+ \rightarrow \pi^+ \chi_{c1}(1P)) = (3.0^{+1.5+3.7}_{-0.8-1.6}) \times 10^{-5}$.

NODE=M191R01;LINKAGE=MI

² With a product branching fraction limit of $B(\bar{B}^0 \rightarrow X(4050)^+ K^-) \times B(X(4050)^+ \rightarrow \chi_{c1} \pi^+) < 1.8 \times 10^{-5}$ at 90% CL.

NODE=M191R01;LINKAGE=LE

 $X(4050)^\pm$ REFERENCES

LEES	12B	PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54042
MIZUK	08	PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=52535

NODE=M191

X(4055)[±]

$$I(J^P) = ?(?^?)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. Seen by WANG 15A in the $\psi(2S)\pi^+$ invariant mass distribution in $X(4360) \rightarrow \psi(2S)\pi^+\pi^-$ decay.

NODE=M223

NODE=M223

X(4055)[±] MASS

NODE=M223M

NODE=M223M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4054±3±1	¹ WANG	15A BELL	10.58 e ⁺ e ⁻ → $\gamma\pi^+\pi^-\psi(2S)$

¹ Statistical significance of 3.5 σ .

NODE=M223M;LINKAGE=A

X(4055)[±] WIDTH

NODE=M223W

NODE=M223W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
45±11±6	¹ WANG	15A BELL	10.58 e ⁺ e ⁻ → $\gamma\pi^+\pi^-\psi(2S)$

¹ Statistical significance of 3.5 σ .

NODE=M223W;LINKAGE=A

X(4055)[±] DECAY MODES

NODE=M223215;NODE=M223

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+\psi(2S)$	seen

DESIG=1

X(4055)[±] BRANCHING RATIOS

NODE=M223225

$\Gamma(\pi^+\psi(2S))/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ WANG	15A BELL	10.58 e ⁺ e ⁻ → $\gamma\pi^+\pi^-\psi(2S)$

¹ Statistical significance of 3.5 σ .NODE=M223R01
NODE=M223R01

NODE=M223R01;LINKAGE=A

X(4055)[±] REFERENCES

WANG 15A PR D91 112007 X.L. Wang et al. (BELLE Collab.)

NODE=M223

REFID=56839

NODE=M193

X(4140)

$$I^G(J^{PC}) = 0^+(?^?+)$$

Seen by AALTONEN 09AH, ABAZOV 14A, CHATRCHYAN 14M in $B^+ \rightarrow XK^+$, $X \rightarrow J/\psi\phi$, and by ABAZOV 15M separately in both prompt (4.7 σ) and non-prompt (5.6 σ) production in $p\bar{p} \rightarrow J/\psi\phi + \text{anything}$. Not seen by SHEN 10 in $\gamma\gamma \rightarrow J/\psi\phi$, AAIJ 12AA in $B^+ \rightarrow J/\psi\phi K^+$, and ABLIKIM 15 in $e^+e^- \rightarrow \gamma J/\psi\phi$ at $\sqrt{s} = 4.23, 4.26, 4.36$ GeV.

NODE=M193

X(4140) MASS

NODE=M193M

NODE=M193M


NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4146.9±3.1 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below. [4143.0 ± 3.1 MeV OUR 2012 AVERAGE]

4152.5±1.7 ^{+6.2} _{-5.4}	1.4k	¹ ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi\phi + \text{anything}$
4159.0±4.3±6.6	52	² ABAZOV	14A D0	$B^+ \rightarrow J/\psi\phi K^+$
4148.0±2.4±6.3	0.3k	³ CHATRCHYAN	14M CMS	$B^+ \rightarrow J/\psi\phi K^+$
4143.0±2.9±1.2	14	⁴ AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi\phi K^+$

- 1 Statistical significance of more than 6σ .
- 2 Statistical significance of 3.1σ .
- 3 From a fit assuming an S -wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5σ .
- 4 Statistical significance of 3.8σ .

NODE=M193M;LINKAGE=C
 NODE=M193M;LINKAGE=A
 NODE=M193M;LINKAGE=B
 NODE=M193M;LINKAGE=AA



X(4140) WIDTH

NODE=M193W
 NODE=M193W
 NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
15 \pm $\frac{6}{5}$				OUR AVERAGE
[12 \pm $\frac{9}{6}$ MeV OUR 2012 AVERAGE]				
16.3± 5.6±11.4	1.4k	1 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi\phi + \text{anything}$
20 ±13 \pm $\frac{3}{8}$	52	2 ABAZOV	14A D0	$B^+ \rightarrow J/\psi\phi K^+$
28 \pm $\frac{15}{11}$ ±19	0.3k	3 CHATRCHYAN	14M CMS	$B^+ \rightarrow J/\psi\phi K^+$
11.7 \pm $\frac{8.3}{5.0}$ ± 3.7	14	4 AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi\phi K^+$

- 1 Statistical significance of more than 6σ .
- 2 Statistical significance of 3.1σ .
- 3 From a fit assuming an S -wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5σ .
- 4 Statistical significance of 3.8σ .

NODE=M193W;LINKAGE=C
 NODE=M193W;LINKAGE=A
 NODE=M193W;LINKAGE=B
 NODE=M193W;LINKAGE=AA

X(4140) DECAY MODES

NODE=M193215;NODE=M193

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\phi$	seen
Γ_2 $\gamma\gamma$	not seen

DESIG=1
 DESIG=2

X(4140) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M193220

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
<41	90	1 SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
< 6	90	2 SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	
1 For $J^P = 0^+$.					
2 For $J^P = 2^+$.					

NODE=M193G01
 NODE=M193G01
 OCCUR=2
 NODE=M193G01;LINKAGE=S0
 NODE=M193G01;LINKAGE=S2

X(4140) BRANCHING RATIOS

NODE=M193225

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
seen [not seen OUR 2015 BEST LIMIT]						
seen	1.9k	1 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi\phi + \text{anything}$		
seen	52	2 ABAZOV	14A D0	$B^+ \rightarrow J/\psi\phi K^+$		
seen	0.3k	3 CHATRCHYAN	14M CMS	$B^+ \rightarrow J/\psi\phi K^+$		
seen	14	4 AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi\phi K^+$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		5 ABLIKIM	15 BES3	$e^+e^- \rightarrow \gamma\phi J/\psi$		
not seen		6 AAIJ	12AA LHCB	$pp \rightarrow B^+ X$ at 7 TeV		
1 Statistical significance of more than 6σ .						
2 ABAZOV 14A reports $B(B^+ \rightarrow X(4140)K^+ \rightarrow J/\psi\phi K^+)/B(B^+ \rightarrow J/\psi\phi K^+) = (19 \pm 7 \pm 4)\%$ with 3.1σ significance.						
3 From a fit assuming an S-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5σ .						
4 Statistical significance of 3.8σ .						
5 Reported $\sigma(e^+e^- \rightarrow \gamma X(4140)) \cdot B(X(4140) \rightarrow J/\psi\phi) < 0.35, 0.28, \text{ and } 0.33 \text{ pb at } 4.23, 4.26, \text{ and } 4.36 \text{ GeV, respectively, at } 90\% \text{ CL.}$						
6 Reported $B(B^+ \rightarrow X(4140)K^+) \cdot B(X(4140) \rightarrow J/\psi\phi)/B(B^+ \rightarrow J/\psi\phi K^+) < 0.07$ at $90\% \text{ CL.}$						

NODE=M193R01
NODE=M193R01NODE=M193R01;LINKAGE=D
NODE=M193R01;LINKAGE=A

NODE=M193R01;LINKAGE=C

NODE=M193R01;LINKAGE=AA
NODE=M193R01;LINKAGE=B

NODE=M193R01;LINKAGE=AI

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		SHEN	10 BELL	$10.6 e^+e^- \rightarrow e^+e^- J/\psi\phi$	

NODE=M193R02
NODE=M193R02**X(4140) REFERENCES**

ABAZOV	15M	PRL 115 232001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	15	PR D91 032002	M. Ablikim <i>et al.</i>	(BES III Collab.)
ABAZOV	14A	PR D89 012004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	14M	PL B734 261	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AAIJ	12AA	PR D85 091103	R. Aaij <i>et al.</i>	(LHCb Collab.)
SHEN	10	PRL 104 112004	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AALTONEN	09AH	PRL 102 242002	T. Aaltonen <i>et al.</i>	(CDF Collab.)

NODE=M193

REFID=56957
REFID=56368
REFID=55650
REFID=55753
REFID=54263
REFID=53235
REFID=52968

NODE=M025

 $\psi(4160)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $\psi(4160)$ MASS

NODE=M025M

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
4191 \pm 5 OUR AVERAGE				
4191 $\begin{smallmatrix} +9 \\ -8 \end{smallmatrix}$		AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$
4191.7 \pm 6.5		1 ABLIKIM	08D BES2	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4193 \pm 7		2 MO	10 RVUE	$e^+e^- \rightarrow \text{hadrons}$
4151 \pm 4		3 SETH	05A RVUE	$e^+e^- \rightarrow \text{hadrons}$
4155 \pm 5		4 SETH	05A RVUE	$e^+e^- \rightarrow \text{hadrons}$
4159 \pm 20		BRANDELIK	78C DASP	e^+e^-

NODE=M025M

OCCUR=2

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

NODE=M025M;LINKAGE=AB

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

NODE=M025M;LINKAGE=MO

³ From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M025M;LINKAGE=ST

⁴ From a fit to BES (BAI 02C) data.

NODE=M025M;LINKAGE=SE

 $\psi(4160)$ WIDTH

NODE=M025W

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
70 \pm 10 OUR AVERAGE				
65 $\begin{smallmatrix} +22 \\ -16 \end{smallmatrix}$		AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$
71.8 \pm 12.3		5 ABLIKIM	08D BES2	$e^+e^- \rightarrow \text{hadrons}$

NODE=M025W

• • • We do not use the following data for averages, fits, limits, etc. • • •

79 ±14	⁶ MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
107 ±10	⁷ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
107 ±16	⁸ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
78 ±20	BRANDELIK	78C DASP	$e^+ e^-$

OCCUR=2

⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

NODE=M025W;LINKAGE=AB

⁶ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

NODE=M025W;LINKAGE=MO

⁷ From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M025W;LINKAGE=ST

⁸ From a fit to BES (BAI 02C) data.

NODE=M025W;LINKAGE=SE

$\psi(4160)$ DECAY MODES

NODE=M025215;NODE=M025

NODE=M025

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$	
Γ_2 $\mu^+ \mu^-$	seen	
Γ_3 $D\bar{D}$	seen	
Γ_4 $D^0\bar{D}^0$	seen	
Γ_5 D^+D^-	seen	
Γ_6 $D^*\bar{D} + \text{c.c.}$	seen	
Γ_7 $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
Γ_8 $D^*(2010)^+D^- + \text{c.c.}$	seen	
Γ_9 $D^*\bar{D}^*$	seen	
Γ_{10} $D^*(2007)^0\bar{D}^*(2007)^0$	seen	
Γ_{11} $D^*(2010)^+D^*(2010)^-$	seen	
Γ_{12} $D^0D^-\pi^+ + \text{c.c. (excl. } D^*(2007)^0\bar{D}^0 + \text{c.c., } D^*(2010)^+D^- + \text{c.c.)}$	not seen	
Γ_{13} $D\bar{D}^*\pi + \text{c.c. (excl. } D^*\bar{D}^*)$	seen	
Γ_{14} $D^0D^{*-}\pi^+ + \text{c.c. (excl. } D^*(2010)^+D^*(2010)^-)$	not seen	
Γ_{15} $D_s^+D_s^-$	not seen	
Γ_{16} $D_s^{*+}D_s^- + \text{c.c.}$	seen	
Γ_{17} $J/\psi\pi^+\pi^-$	< 3 $\times 10^{-3}$	90%
Γ_{18} $J/\psi\pi^0\pi^0$	< 3 $\times 10^{-3}$	90%
Γ_{19} $J/\psi K^+K^-$	< 2 $\times 10^{-3}$	90%
Γ_{20} $J/\psi\eta$	< 8 $\times 10^{-3}$	90%
Γ_{21} $J/\psi\pi^0$	< 1 $\times 10^{-3}$	90%
Γ_{22} $J/\psi\eta'$	< 5 $\times 10^{-3}$	90%
Γ_{23} $J/\psi\pi^+\pi^-\pi^0$	< 1 $\times 10^{-3}$	90%
Γ_{24} $\psi(2S)\pi^+\pi^-$	< 4 $\times 10^{-3}$	90%
Γ_{25} $\chi_{c1}\gamma$	< 5 $\times 10^{-3}$	90%
Γ_{26} $\chi_{c2}\gamma$	< 1.3 %	90%
Γ_{27} $\chi_{c1}\pi^+\pi^-\pi^0$	< 2 $\times 10^{-3}$	90%
Γ_{28} $\chi_{c2}\pi^+\pi^-\pi^0$	< 8 $\times 10^{-3}$	90%
Γ_{29} $h_c(1P)\pi^+\pi^-$	< 5 $\times 10^{-3}$	90%
Γ_{30} $h_c(1P)\pi^0\pi^0$	< 2 $\times 10^{-3}$	90%
Γ_{31} $h_c(1P)\eta$	< 2 $\times 10^{-3}$	90%
Γ_{32} $h_c(1P)\pi^0$	< 4 $\times 10^{-4}$	90%
Γ_{33} $\phi\pi^+\pi^-$	< 2 $\times 10^{-3}$	90%

DESIG=1

DESIG=33

DESIG=15;OUR EVAL;→ UNCHECKED ←

DESIG=16

DESIG=17

DESIG=18;OUR EVAL;→ UNCHECKED ←

DESIG=19

DESIG=20

DESIG=21;OUR EVAL;→ UNCHECKED ←

DESIG=22

DESIG=23

DESIG=24

DESIG=25

DESIG=26

DESIG=27

DESIG=28

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

DESIG=9

DESIG=10

DESIG=11

DESIG=12

DESIG=13

DESIG=29

DESIG=30

DESIG=31

DESIG=32

DESIG=14

Γ_{34}	$\gamma X(3872) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 6.8	$\times 10^{-5}$	90%	DESIG=34
Γ_{35}	$\gamma X(3915) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.36	$\times 10^{-4}$	90%	DESIG=35
Γ_{36}	$\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.18	$\times 10^{-4}$	90%	DESIG=36
Γ_{37}	$\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.47	$\times 10^{-4}$	90%	DESIG=37
Γ_{38}	$\gamma X(3872) \rightarrow \gamma \gamma J/\psi$	< 1.05	$\times 10^{-4}$	90%	DESIG=38
Γ_{39}	$\gamma X(3915) \rightarrow \gamma \gamma J/\psi$	< 1.26	$\times 10^{-4}$	90%	DESIG=39
Γ_{40}	$\gamma X(3930) \rightarrow \gamma \gamma J/\psi$	< 8.8	$\times 10^{-5}$	90%	DESIG=40
Γ_{41}	$\gamma X(3940) \rightarrow \gamma \gamma J/\psi$	< 1.79	$\times 10^{-4}$	90%	DESIG=41

 $\psi(4160)$ PARTIAL WIDTHS

NODE=M025220

 $\Gamma(e^+ e^-)$ **Γ_1**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.48±0.22	⁹ ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.4 to 1.1	¹⁰ MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
0.83±0.08	¹¹ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.84±0.13	¹² SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.77±0.23	BRANDELIK	78C DASP	$e^+ e^-$

NODE=M025W1
NODE=M025W1

OCCUR=2

⁹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.

NODE=M025W1;LINKAGE=AB

¹⁰ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.

NODE=M025W1;LINKAGE=MO

¹¹ From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M025W1;LINKAGE=ST

¹² From a fit to BES (BAI 02C) data.

NODE=M025W1;LINKAGE=SE

 $\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

NODE=M025235

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{25} \Gamma_1 / \Gamma$
<2.2	90	¹³ HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c1} \gamma$	NODE=M025G01 NODE=M025G01

¹³ Using $B(\eta \rightarrow \gamma \gamma) = (39.41 \pm 0.21)\%$.

NODE=M025G01;LINKAGE=A

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{26} \Gamma_1 / \Gamma$
<6.1	90	¹⁴ HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c2} \gamma$	NODE=M025G02 NODE=M025G02

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<6.1 90 ¹⁴ HAN 15 BELL $10.58 e^+ e^- \rightarrow \chi_{c2} \gamma$

¹⁴ Using $B(\eta \rightarrow \gamma \gamma) = (39.41 \pm 0.21)\%$.

NODE=M025G02;LINKAGE=A

 $\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$

NODE=M025230

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT	$\Gamma_{20} / \Gamma \times \Gamma_1 / \Gamma$
2.8±0.9±0.9	¹⁵ WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$	NODE=M025R32 NODE=M025R32
12.8±1.7±2.0	¹⁶ WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$	OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2.8±0.9±0.9 ¹⁵ WANG 13B BELL $e^+ e^- \rightarrow J/\psi \eta \gamma$

12.8±1.7±2.0 ¹⁶ WANG 13B BELL $e^+ e^- \rightarrow J/\psi \eta \gamma$

OCCUR=2

¹⁵ Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

NODE=M025R32;LINKAGE=A

¹⁶ Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

NODE=M025R32;LINKAGE=B

 $\psi(4160)$ BRANCHING RATIOS

NODE=M025225

 $\Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}}$ **Γ_2 / Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹⁷ AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$

NODE=M025R31
NODE=M025R31

¹⁷ AAIJ 13BC report $B(B^+ \rightarrow K^+ \psi(4160)) B(\psi(4160) \rightarrow \mu^+ \mu^-) = (3.5_{-0.8}^{+0.9}) \times 10^{-9}$.

NODE=M025R31;LINKAGE=A

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$				Γ_3/Γ_9	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R14 NODE=M025R14
0.02±0.03±0.02	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	
$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$				Γ_4/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R16 NODE=M025R16
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^0\bar{D}^0$	
seen	PAKHLOVA 08	BELL		$e^+e^- \rightarrow D^0\bar{D}^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$	
$\Gamma(D^+D^-)/\Gamma_{\text{total}}$				Γ_5/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R17 NODE=M025R17
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^+D^-$	
seen	PAKHLOVA 08	BELL		$e^+e^- \rightarrow D^+D^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^+D^-\gamma$	
$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_7/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R18 NODE=M025R18
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$	
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*0}\bar{D}^0$	
$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_8/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R19 NODE=M025R19
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$	
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*+}D^-$	
seen	PAKHLOVA 07	BELL		$e^+e^- \rightarrow D^{*+}D^-\gamma$	
$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$				Γ_6/Γ_9	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R15 NODE=M025R15
0.34±0.14±0.05	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	
$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$				Γ_{10}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R20 NODE=M025R20
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$	
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$	
$\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$				Γ_{11}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R21 NODE=M025R21
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$	
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*+}D^{*-}$	
seen	PAKHLOVA 07	BELL		$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$	
$\Gamma(D^0D^-\pi^+ + \text{c.c. (excl. } D^*(2007)^0\bar{D}^0 + \text{c.c., } D^*(2010)^+D^- + \text{c.c.}))/\Gamma_{\text{total}}$				Γ_{12}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R22 NODE=M025R22
not seen	PAKHLOVA 08A	BELL		$e^+e^- \rightarrow D^0D^-\pi^+\gamma$	
$\Gamma(D\bar{D}^*\pi + \text{c.c. (excl. } D^*\bar{D}^*))/\Gamma_{\text{total}}$				Γ_{13}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R23 NODE=M025R23
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D\bar{D}^*\pi$	
$\Gamma(D^0D^{*-}\pi^+ + \text{c.c. (excl. } D^*(2010)^+D^*(2010)^-))/\Gamma_{\text{total}}$				Γ_{14}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R24 NODE=M025R24
not seen	PAKHLOVA 09	BELL		$e^+e^- \rightarrow D^0D^{*-}\pi^+\gamma$	
$\Gamma(D_s^+D_s^-)/\Gamma_{\text{total}}$				Γ_{15}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M025R25 NODE=M025R25
not seen	PAKHLOVA 11	BELL		$e^+e^- \rightarrow D_s^+D_s^-\gamma$	
not seen	DEL-AMO-SA..10N	BABR		$e^+e^- \rightarrow D_s^+D_s^-\gamma$	
not seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D_s^+D_s^-$	

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	19 PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

NODE=M025R27
NODE=M025R27

¹⁹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M025R27;LINKAGE=PE

 $\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	20 PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$

NODE=M025R28
NODE=M025R28

²⁰ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M025R28;LINKAGE=PE

 $\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	21 PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\eta$

NODE=M025R29
NODE=M025R29

²¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M025R29;LINKAGE=PE

 $\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	22 PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\pi^0$

NODE=M025R30
NODE=M025R30

²² At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M025R30;LINKAGE=PE

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06 CLEO	$4.12-4.2 e^+e^- \rightarrow \text{hadrons}$

NODE=M025R13
NODE=M025R13

 $\Gamma(\gamma X(3872) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<0.68 $\times 10^{-4}$	90	23 XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

NODE=M025R34
NODE=M025R34

²³ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R34;LINKAGE=A

 $\Gamma(\gamma X(3915) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.36 $\times 10^{-4}$	90	24 XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

NODE=M025R35
NODE=M025R35

²⁴ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R35;LINKAGE=A

 $\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.18 $\times 10^{-4}$	90	25 XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

NODE=M025R36
NODE=M025R36

²⁵ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R36;LINKAGE=A

 $\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.47 $\times 10^{-4}$	90	26 XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

NODE=M025R37
NODE=M025R37

²⁶ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R37;LINKAGE=A

 $\Gamma(\gamma X(3872) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.05 $\times 10^{-4}$	90	27 XIAO	13 $\psi(4160) \rightarrow \gamma\gamma J/\psi$

NODE=M025R38
NODE=M025R38

²⁷ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R38;LINKAGE=A

 $\Gamma(\gamma X(3915) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.26 $\times 10^{-4}$	90	28 XIAO	13 $\psi(4160) \rightarrow \gamma\gamma J/\psi$

NODE=M025R39
NODE=M025R39

²⁸ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R39;LINKAGE=A

$\Gamma(\gamma X(3930) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{40}/Γ
$<0.88 \times 10^{-4}$	90	29 XIAO	13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$	

²⁹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R40
 NODE=M025R40

NODE=M025R40;LINKAGE=A

 $\Gamma(\gamma X(3940) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{41}/Γ
$<1.79 \times 10^{-4}$	90	30 XIAO	13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$	

³⁰ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R41
 NODE=M025R41

NODE=M025R41;LINKAGE=A

 $\psi(4160)$ REFERENCES

ABLIKIM 15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BES III Collab.)
HAN 15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
AAIJ 13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
WANG 13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
XIAO 13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)
PAKHLOVA 11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR 11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA... 10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO 10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT 09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN... 09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA 09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM 08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA 08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN 06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH 05A	PR D72 017501	K.K. Seth	
BAI 02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI 00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD 86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK 78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)

NODE=M025

REFID=56777
 REFID=56816
 REFID=55229
 REFID=55377
 REFID=55381
 REFID=53638
 REFID=16787
 REFID=53532
 REFID=53532
 REFID=53540
 REFID=52724
 REFID=53114
 REFID=53143
 REFID=52142
 REFID=52132
 REFID=52134
 REFID=51628
 REFID=51075
 REFID=50813
 REFID=50506
 REFID=50503
 REFID=51064
 REFID=22232

NODE=M190

X(4160)

$$I^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

Seen by PAKHLOV 08 in $e^+e^- \rightarrow J/\psi X$, $X \rightarrow D^*\bar{D}^*$

NODE=M190

 $X(4160)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4156^{+25}_{-20} \pm 15$	24	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M190M

NODE=M190M

 $X(4160)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$139^{+111}_{-61} \pm 21$	24	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M190W

NODE=M190W

 $X(4160)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\bar{D}$	not seen
Γ_2 $D^*\bar{D} + \text{c.c.}$	not seen
Γ_3 $D^*\bar{D}^*$	seen

NODE=M190215;NODE=M190

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

 $X(4160)$ BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$					Γ_1/Γ_3
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.09	90	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$	

NODE=M190225

NODE=M190R01
 NODE=M190R01

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$					Γ_2/Γ_3
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.22	90	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$	

NODE=M190R02
 NODE=M190R02

 $X(4160)$ REFERENCES

PAKHLOV 08	PRL 100 202001	P. Pakhlov <i>et al.</i>	(BELLE Collab.)
------------	----------------	--------------------------	-----------------

NODE=M190

REFID=52302

X(4200)[±]

$$I(J^P) = ?(1^+)$$

OMITTED FROM SUMMARY TABLE

Reported by CHILIKIN 14 in $J/\psi\pi^+$ at a significance of 6.2σ . Assignments of 0^- , 1^- , 2^- , and 2^+ excluded at 6.1σ , 7.4σ , 4.4σ , and 7.0σ level, respectively. Needs confirmation.

NODE=M231

NODE=M231

X(4200)[±] MASS

NODE=M231M

NODE=M231M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4196^{+31+17}_{-29-13}	CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

X(4200)[±] WIDTH

NODE=M231W

NODE=M231W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$370 \pm 70^{+70}_{-132}$	CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

X(4200)[±] DECAY MODES

NODE=M231215;NODE=M231

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\pi^+$	seen

DESIG=1

X(4200)[±] BRANCHING RATIOS

NODE=M231220

$\Gamma(J/\psi\pi^+)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow J/\psi K^- \pi^+$	

NODE=M231R01
NODE=M231R01**X(4200)[±] REFERENCES**

NODE=M231

CHILIKIN 14 PR D90 112009 K. Chilikin *et al.* (BELLE Collab.)

REFID=56344

NODE=M222

X(4230)

$$I^G(J^{PC}) = ?^?(1^{--})$$

OMITTED FROM SUMMARY TABLE

Enhancement reported by ABLIKIM 15C in $e^+e^- \rightarrow \omega\chi_{c0}$ at $\sqrt{s} = 4.23\text{--}4.26$ GeV at 9σ significance. Lineshape found to be inconsistent with origination from X(4260). NEEDS CONFIRMATION.

NODE=M222

X(4230) MASS

NODE=M222M

NODE=M222M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4230 \pm 8 \pm 6$	180	¹ ABLIKIM 15C	BES3	$e^+e^- \rightarrow \omega\chi_{c0}$

¹From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+\pi^-$, $\chi_{c0} \rightarrow K^+K^-$, and $\omega \rightarrow \pi^+\pi^-\pi^0$.

NODE=M222M;LINKAGE=A

X(4230) WIDTH

NODE=M222W

NODE=M222W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$38 \pm 12 \pm 2$	180	¹ ABLIKIM 15C	BES3	$e^+e^- \rightarrow \omega\chi_{c0}$

¹From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+\pi^-$, $\chi_{c0} \rightarrow K^+K^-$, and $\omega \rightarrow \pi^+\pi^-\pi^0$.

NODE=M222W;LINKAGE=A

X(4230) DECAY MODES

NODE=M222215;NODE=M222

Mode	Fraction (Γ_i/Γ)
Γ_1 $e^+ e^-$	
Γ_2 $\omega \chi_{c0}$	seen

DESIG=1

DESIG=2

X(4230) $\Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

NODE=M222220

$$\Gamma(\omega \chi_{c0}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_2 \Gamma_1 / \Gamma$$

NODE=M222G01

NODE=M222G01

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
2.7±0.5±0.4	180	¹ ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega \chi_{c0}$

¹ From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$, $\chi_{c0} \rightarrow K^+ K^-$, and $\omega \rightarrow \pi^+ \pi^- \pi^0$.

NODE=M222G01;LINKAGE=A

X(4230) BRANCHING RATIOS

NODE=M222225

$$\Gamma(\omega \chi_{c0}) / \Gamma_{\text{total}} \quad \Gamma_2 / \Gamma$$

NODE=M222R01

NODE=M222R01

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen	180	¹ ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega \chi_{c0}$

¹ From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$, $\chi_{c0} \rightarrow K^+ K^-$, and $\omega \rightarrow \pi^+ \pi^- \pi^0$.

NODE=M222R01;LINKAGE=A

X(4230) REFERENCES

NODE=M222

ABLIKIM 15C PRL 114 092003 M. Ablikim *et al.* (BES III Collab.)

REFID=56401

NODE=M216

X(4240)[±]

$$I^G(J^P) = ?^?(0^-)$$

OMITTED FROM SUMMARY TABLE

Spin and parity assignment $J^P = 0^-$ is favored over 1^- , 2^- , and 2^+ by 8σ and over 1^+ by 1σ , according to the four-dimensional amplitude analysis of AAIJ 14AG.

NODE=M216

X(4240)[±] MASS

NODE=M216M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4239±18⁺⁴⁵₋₁₀	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

NODE=M216M

¹ From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $X(4430)^\pm$ fit, with significance 6σ including systematic variations.

NODE=M216M;LINKAGE=AA

X(4240)[±] WIDTH

NODE=M216W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
220±47⁺¹⁰⁸₋₇₄	² AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

NODE=M216W

² From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $X(4430)^\pm$ fit, with significance 6σ including systematic variations.

NODE=M216W;LINKAGE=AA

X(4240)[±] DECAY MODES

NODE=M216215;NODE=M216

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^- \psi(2S)$	seen

DESIG=1

X(4240)[±] BRANCHING RATIOS $\Gamma(\pi^- \psi(2S))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	³ AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

³From a 4-dimensional analysis when a second, lower mass resonance is allowed in the X(4430)[±] fit. No partial branching fraction quoted.

NODE=M216225

NODE=M216R01
NODE=M216R01

NODE=M216R01;LINKAGE=AA

X(4240)[±] REFERENCES

AAIJ	14AG PRL 112 222002	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	---------------------	-----------------------	----------------

NODE=M216

REFID=55896

NODE=M192

X(4250)[±] $I(J^P) = ?(??)$ **OMITTED FROM SUMMARY TABLE**

Observed by MIZUK 08 in the $\pi^+ \chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure.

NODE=M192

X(4250)[±] MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4248^{+44+180}_{-29-35}$	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

NODE=M192M

NODE=M192M

¹From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192M;LINKAGE=MI

X(4250)[±] WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$177^{+54+316}_{-39-61}$	² MIZUK	08	BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

NODE=M192W

NODE=M192W

²From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192W;LINKAGE=MI

X(4250)[±] DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+ \chi_{c1}(1P)$	seen

NODE=M192215;NODE=M192

DESIG=1

X(4250)[±] BRANCHING RATIOS $\Gamma(\pi^+ \chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	³ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

NODE=M192225

NODE=M192R01
NODE=M192R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	⁴ LEES	12B	BABR $B \rightarrow K \pi \chi_{c1}(1P)$
----------	-------------------	-----	--

³With a product branching fraction measurement of $B(\bar{B}^0 \rightarrow K^- X(4250)^+) \times B(X(4250)^+ \rightarrow \pi^+ \chi_{c1}(1P)) = (4.0^{+2.3+19.7}_{-0.9-0.5}) \times 10^{-5}$.

NODE=M192R01;LINKAGE=MI

⁴With a product branching fraction limit of $B(\bar{B}^0 \rightarrow X(4250)^+ K^-) \times B(X(4250)^+ \rightarrow \chi_{c1} \pi^+) < 4.0 \times 10^{-5}$ at 90% CL.

NODE=M192R01;LINKAGE=LE

X(4250)[±] REFERENCES

LEES	12B PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08 PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M192

REFID=54042
REFID=52535

X(4260)

$$I^G(J^{PC}) = ?(1^{--})$$

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by AUBERT,B 05I, HE 06B, and YUAN 07, and in e^+e^- collisions at $\sqrt{s} \approx 4.26$ GeV by COAN 06. Possibly seen by AUBERT 06 in $B^- \rightarrow K^- \pi^+ \pi^- J/\psi$. See also the mini-review under the X(3872). (See the index for the page number.)

NODE=M074

NODE=M074

X(4260) MASS

NODE=M074M

NODE=M074M

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

4251 ± 9 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

4258.6 ± 8.3 ± 12.1	1	LIU	13B	BELL	$e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
4245 ± 5 ± 4	2	LEES	12AC	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
4284 $^{+17}_{-16}$ ± 413.6		HE	06B	CLEO	9.4–10.6 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4247 ± 12 $^{+17}_{-32}$	1,3	YUAN	07	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
4259 ± 8 $^{+2}_{-6}$ 125	4	AUBERT,B	05I	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$

¹ From a two-resonance fit.

² From a single-resonance fit. Supersedes AUBERT,B 05I.

³ Superseded by LIU 13B.


⁴ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074M;LINKAGE=YU

NODE=M074M;LINKAGE=LE

NODE=M074M;LINKAGE=YN

NODE=M074M;LINKAGE=AU

**X(4260) WIDTH**

NODE=M074W

NODE=M074W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

120 ± 12 OUR AVERAGE Error includes scale factor of 1.1.

134.1 ± 16.4 ± 5.5	1	LIU	13B	BELL	$e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
114 $^{+16}_{-15}$ ± 7	2	LEES	12AC	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
73 $^{+39}_{-25}$ ± 5 13.6		HE	06B	CLEO	9.4–10.6 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

108 ± 19 ± 10	1,3	YUAN	07	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
88 ± 23 $^{+6}_{-4}$ 125	4	AUBERT,B	05I	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$

¹ From a two-resonance fit.

² From a single-resonance fit. Supersedes AUBERT,B 05l.

³ Superseded by LIU 13B.

⁴ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074W;LINKAGE=YU

NODE=M074W;LINKAGE=LE

NODE=M074W;LINKAGE=YN

NODE=M074W;LINKAGE=AU

X(4260) DECAY MODES

NODE=M074215;NODE=M074

Mode	Fraction (Γ_i/Γ)	
Γ_1 $e^+ e^-$		DESIG=1
Γ_2 $J/\psi \pi^+ \pi^-$	seen	DESIG=2;OUR EVAL;→ UNCHECKED ←
Γ_3 $J/\psi f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$	seen	DESIG=41;OUR EVAL;→ UNCHECKED ←
Γ_4 $X(3900)^\pm \pi^\mp, X^\pm \rightarrow J/\psi \pi^\pm$	seen	DESIG=43;OUR EVAL;→ UNCHECKED ←
Γ_5 $J/\psi \pi^0 \pi^0$	seen	DESIG=4;OUR EVAL;→ UNCHECKED ←
Γ_6 $J/\psi K^+ K^-$	seen	DESIG=5;OUR EVAL;→ UNCHECKED ←
Γ_7 $J/\psi K_S^0 K_S^0$	not seen	DESIG=44
Γ_8 $X(3872) \gamma$	seen	DESIG=42
Γ_9 $J/\psi \eta$	not seen	DESIG=6;OUR EVAL;→ UNCHECKED ←
Γ_{10} $J/\psi \pi^0$	not seen	DESIG=7;OUR EVAL;→ UNCHECKED ←
Γ_{11} $J/\psi \eta'$	not seen	DESIG=8;OUR EVAL;→ UNCHECKED ←
Γ_{12} $J/\psi \pi^+ \pi^- \pi^0$	not seen	DESIG=9;OUR EVAL;→ UNCHECKED ←
Γ_{13} $J/\psi \eta \pi^0$	not seen	DESIG=45
Γ_{14} $J/\psi \eta \eta$	not seen	DESIG=10;OUR EVAL;→ UNCHECKED ←
Γ_{15} $\psi(2S) \pi^+ \pi^-$	not seen	DESIG=11;OUR EVAL;→ UNCHECKED ←
Γ_{16} $\psi(2S) \eta$	not seen	DESIG=12;OUR EVAL;→ UNCHECKED ←
Γ_{17} $\chi_{c0} \omega$	not seen	DESIG=13;OUR EVAL;→ UNCHECKED ←
Γ_{18} $\chi_{c1} \gamma$	not seen	DESIG=14;OUR EVAL;→ UNCHECKED ←
Γ_{19} $\chi_{c2} \gamma$	not seen	DESIG=15;OUR EVAL;→ UNCHECKED ←
Γ_{20} $\chi_{c1} \pi^+ \pi^- \pi^0$	not seen	DESIG=16;OUR EVAL;→ UNCHECKED ←
Γ_{21} $\chi_{c2} \pi^+ \pi^- \pi^0$	not seen	DESIG=17;OUR EVAL;→ UNCHECKED ←
Γ_{22} $h_c(1P) \pi^+ \pi^-$	not seen	DESIG=40;OUR EVAL;→ UNCHECKED ←
Γ_{23} $\phi \pi^+ \pi^-$	not seen	DESIG=18;OUR EVAL;→ UNCHECKED ←
Γ_{24} $\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	not seen	DESIG=22;OUR EVAL;→ UNCHECKED ←
Γ_{25} $D \bar{D}$	not seen	DESIG=19;OUR EVAL;→ UNCHECKED ←
Γ_{26} $D^0 \bar{D}^0$	not seen	DESIG=31
Γ_{27} $D^+ D^-$	not seen	DESIG=32
Γ_{28} $D^* \bar{D} + c.c.$	not seen	DESIG=23;OUR EVAL;→ UNCHECKED ←
Γ_{29} $D^*(2007)^0 \bar{D}^0 + c.c.$	not seen	DESIG=33
Γ_{30} $D^*(2010)^+ D^- + c.c.$	not seen	DESIG=34
Γ_{31} $D^* \bar{D}^*$	not seen	DESIG=24;OUR EVAL;→ UNCHECKED ←
Γ_{32} $D^*(2007)^0 \bar{D}^*(2007)^0$	not seen	DESIG=35
Γ_{33} $D^*(2010)^+ D^*(2010)^-$	not seen	DESIG=36
Γ_{34} $D \bar{D} \pi + c.c.$		DESIG=37
Γ_{35} $D^0 D^- \pi^+ + c.c. (excl. D^*(2007)^0 \bar{D}^{*0} + c.c., D^*(2010)^+ D^- + c.c.)$	not seen	DESIG=38
Γ_{36} $D \bar{D}^* \pi + c.c. (excl. D^* \bar{D}^*)$	not seen	DESIG=25
Γ_{37} $D^0 D^{*-} \pi^+ + c.c. (excl. D^*(2010)^+ D^*(2010)^-)$	not seen	DESIG=39
Γ_{38} $D^0 D^*(2010)^- \pi^+ + c.c.$	not seen	DESIG=30;OUR EVAL;→ UNCHECKED ←
Γ_{39} $D^* \bar{D}^* \pi$	not seen	DESIG=26
Γ_{40} $D_s^+ D_s^-$	not seen	DESIG=27
Γ_{41} $D_s^{*+} D_s^- + c.c.$	not seen	DESIG=28
Γ_{42} $D_s^{*+} D_s^{*-}$	not seen	DESIG=29
Γ_{43} $p \bar{p}$	not seen	DESIG=3;OUR EVAL;→ UNCHECKED ←
Γ_{44} $K_S^0 K^\pm \pi^\mp$	not seen	DESIG=20;OUR EVAL;→ UNCHECKED ←
Γ_{45} $K^+ K^- \pi^0$	not seen	DESIG=21;OUR EVAL;→ UNCHECKED ←

$X(4260) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

NODE=M074230

 $\Gamma(J/\psi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_2 \Gamma_1 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

9.2±1.0 OUR AVERAGE

9.2±0.8±0.7		¹ LEES	12AC BABR	10.58 e ⁺ e ⁻ → γπ ⁺ π ⁻ J/ψ
8.9 ^{+3.9} _{-3.1} ±1.8	8.1	HE	06B CLEO	9.4-10.6 e ⁺ e ⁻ → γπ ⁺ π ⁻ J/ψ

••• We do not use the following data for averages, fits, limits, etc. •••

6.4±0.8±0.6		² LIU	13B BELL	e ⁺ e ⁻ → γπ ⁺ π ⁻ J/ψ
20.5±1.4±2.0		³ LIU	13B BELL	e ⁺ e ⁻ → γπ ⁺ π ⁻ J/ψ
6.0±1.2 ^{+4.7} _{-0.5}		^{2,4} YUAN	07 BELL	10.58 e ⁺ e ⁻ → γπ ⁺ π ⁻ J/ψ
20.6±2.3 ^{+9.1} _{-1.7}		^{3,4} YUAN	07 BELL	10.58 e ⁺ e ⁻ → γπ ⁺ π ⁻ J/ψ
5.5±1.0 ^{+0.8} _{-0.7}	125	⁵ AUBERT,B	05I BABR	10.58 e ⁺ e ⁻ → γπ ⁺ π ⁻ J/ψ

¹ From a single-resonance fit. Supersedes AUBERT,B 05I.

² Solution I of two equivalent solutions in a fit using two interfering resonances.

³ Solution II of two equivalent solutions in a fit using two interfering resonances.

⁴ Superseded by LIU 13B.

⁵ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074G1
NODE=M074G1

OCCUR=2

OCCUR=2

NODE=M074G1;LINKAGE=LE
NODE=M074G1;LINKAGE=YOU
NODE=M074G1;LINKAGE=YA
NODE=M074G1;LINKAGE=YN
NODE=M074G1;LINKAGE=AU $\Gamma(J/\psi K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_6 \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.7	90	¹ SHEN	14 BELL	9.4-10.9 e ⁺ e ⁻ → γK ⁺ K ⁻ J/ψ
------	----	-------------------	---------	---

••• We do not use the following data for averages, fits, limits, etc. •••

<1.2	90	² YUAN	08 BELL	e ⁺ e ⁻ → γK ⁺ K ⁻ J/ψ
------	----	-------------------	---------	--

¹ From a fit of the broad K⁺K⁻J/ψ enhancement including a coherent X(4260) amplitude with mass and width from LIU 13B. Supersedes YUAN 08.

² From a fit of the broad K⁺K⁻J/ψ enhancement including a coherent X(4260) amplitude with mass and width from YUAN 07.

NODE=M074G3
NODE=M074G3

NODE=M074G3;LINKAGE=A

NODE=M074G3;LINKAGE=YOU

 $\Gamma(J/\psi K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_7 \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.85	90	¹ SHEN	14 BELL	9.4-10.9 e ⁺ e ⁻ → γK _S ⁰ K _S ⁰ J/ψ
-------	----	-------------------	---------	---

¹ From a fit of the K_S⁰K_S⁰J/ψ mass range from 4.4 to 5.5 GeV including a coherent X(4260) amplitude with mass and width from LIU 13B.

NODE=M074G02
NODE=M074G02

NODE=M074G02;LINKAGE=A

 $\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_9 \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<14.2	90	WANG	13B BELL	e ⁺ e ⁻ → J/ψηγ
-------	----	------	----------	---------------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M074G01
NODE=M074G01 $\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{15} \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<4.3	90	¹ LIU	08H RVUE	10.58 e ⁺ e ⁻ → ψ(2S)π ⁺ π ⁻ γ
7.4 ^{+2.1} _{-1.7}		² LIU	08H RVUE	10.58 e ⁺ e ⁻ → ψ(2S)π ⁺ π ⁻ γ

••• We do not use the following data for averages, fits, limits, etc. •••

¹ For constructive interference with the X(4360) in a combined fit of AUBERT 07S and WANG 07D data with three resonances.

² For destructive interference with the X(4360) in a combined fit of AUBERT 07S and WANG 07D data with three resonances.

NODE=M074G7
NODE=M074G7

OCCUR=2

NODE=M074G7;LINKAGE=LI

NODE=M074G7;LINKAGE=LU

 $\Gamma(\chi_{c1} \gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{18} \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.4	90	¹ HAN	15 BELL	10.58 e ⁺ e ⁻ → χ _{c1} γ
------	----	------------------	---------	---

¹ Using B(η → γγ) = (39.41 ± 0.21)%.

NODE=M074G03
NODE=M074G03

NODE=M074G03;LINKAGE=A

 $\Gamma(\chi_{c2} \gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{19} \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<4.0	90	¹ HAN	15 BELL	10.58 e ⁺ e ⁻ → χ _{c2} γ
------	----	------------------	---------	---

¹ Using B(η → γγ) = (39.41 ± 0.21)%.

NODE=M074G04
NODE=M074G04

NODE=M074G04;LINKAGE=A

$$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{23}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	AUBERT,BE	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

NODE=M074G2
NODE=M074G2

$$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.29	90	¹ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

NODE=M074G6
NODE=M074G6

¹ AUBERT 07AK reports $[\Gamma(X(4260) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(X(4260) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] < 0.14$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = 48.9 \times 10^{-2}$.

NODE=M074G6;LINKAGE=AU

$$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{44}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.5	90	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$
------	----	--------	----------	--

NODE=M074G4
NODE=M074G4

$$\Gamma(K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{45}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.6	90	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\gamma$
------	----	--------	----------	---

NODE=M074G5
NODE=M074G5

X(4260) BRANCHING RATIOS

NODE=M074240

$$\Gamma(J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-) \quad \Gamma_3/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.17±0.13	¹ LEES	12AC BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
-----------	-------------------	-----------	---

¹ Systematic uncertainties not estimated.

NODE=M074R02
NODE=M074R02

NODE=M074R02;LINKAGE=LE

$$\Gamma(X(3900)^\pm \pi^\mp, X^\pm \rightarrow J/\psi\pi^\pm)/\Gamma(J/\psi\pi^+\pi^-) \quad \Gamma_4/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.215±0.033±0.075 ¹ ABLIKIM 13T BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$

••• We do not use the following data for averages, fits, limits, etc. •••

0.29 ±0.08	² LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
------------	------------------	----------	---

¹ Assuming that the cross section of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ is fully due to the X(4260).

² Systematic error not evaluated.

NODE=M074R01
NODE=M074R01

NODE=M074R01;LINKAGE=AB
NODE=M074R01;LINKAGE=A

$$\Gamma(J/\psi K_S^0 K_S^0)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen SHEN 14 BELL 9.4–10.9 $e^+e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$ |

NODE=M074R27
NODE=M074R27

$$\Gamma(X(3872)\gamma)/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 20 ± 5 ABLIKIM 14 BES3 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$

NODE=M074R26
NODE=M074R26

$$\Gamma(J/\psi\eta\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen ABLIKIM 15Q BES3 4.0–4.6 $e^+e^- \rightarrow J/\psi\eta\pi^0$ |

NODE=M074R28
NODE=M074R28

$$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-) \quad \Gamma_{22}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.0 90 ¹ PEDLAR 11 CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

¹ At $\sqrt{s} = 4260$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 32 \pm 17 \pm 6 \pm 6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M074R25
NODE=M074R25

NODE=M074R25;LINKAGE=PE

$$\Gamma(D\bar{D})/\Gamma(J/\psi\pi^+\pi^-) \quad \Gamma_{25}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.0 90 ¹ AUBERT 07BE BABR $e^+e^- \rightarrow D\bar{D}\gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

<4.0	90	CRONIN-HEN..09	CLEO	e^+e^-
------	----	----------------	------	----------

¹ Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of X(4260).

NODE=M074R2
NODE=M074R2

NODE=M074R2;LINKAGE=AU

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{26}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$	
••• We do not use the following data for averages, fits, limits, etc. •••				
not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^0\bar{D}^0\gamma$	
not seen	PAKHLOVA	08	BELL $e^+e^- \rightarrow D^0\bar{D}^0\gamma$	

NODE=M074R12
NODE=M074R12

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{27}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^+D^-$	
••• We do not use the following data for averages, fits, limits, etc. •••				
not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^+D^-\gamma$	
not seen	PAKHLOVA	08	BELL $e^+e^- \rightarrow D^+D^-\gamma$	

NODE=M074R13
NODE=M074R13

 $\Gamma(D^*\bar{D}^0+c.c.)/\Gamma(J/\psi\pi^+\pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{28}/Γ_2
<34	90	AUBERT	09M	BABR $e^+e^- \rightarrow \gamma D^*\bar{D}^0$	
••• We do not use the following data for averages, fits, limits, etc. •••					
<45	90	CRONIN-HEN..09	CLEO	e^+e^-	

NODE=M074R03
NODE=M074R03

 $\Gamma(D^*(2007)^0\bar{D}^0+c.c.)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{29}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^0$	
••• We do not use the following data for averages, fits, limits, etc. •••				
not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$	

NODE=M074R14
NODE=M074R14

 $\Gamma(D^*(2010)^+D^-+c.c.)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{30}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^-$	
not seen	PAKHLOVA	07	BELL $e^+e^- \rightarrow D^{*+}D^-\gamma$	
••• We do not use the following data for averages, fits, limits, etc. •••				
not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*+}D^-\gamma$	

NODE=M074R15
NODE=M074R15

 $\Gamma(D^*\bar{D}^*)/\Gamma(J/\psi\pi^+\pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{31}/Γ_2
<11	90	CRONIN-HEN..09	CLEO	e^+e^-	
••• We do not use the following data for averages, fits, limits, etc. •••					
<40	90	AUBERT	09M	BABR $e^+e^- \rightarrow \gamma D^*\bar{D}^*$	

NODE=M074R04
NODE=M074R04

 $\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{32}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$	
••• We do not use the following data for averages, fits, limits, etc. •••				
not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$	

NODE=M074R17
NODE=M074R17

 $\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{33}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^{*-}$	
not seen	PAKHLOVA	07	BELL $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$	
••• We do not use the following data for averages, fits, limits, etc. •••				
not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$	

NODE=M074R18
NODE=M074R18

 $\Gamma(D^0D^-\pi^++c.c. (\text{excl. } D^*(2007)^0\bar{D}^{*0}+c.c., D^*(2010)^+D^-+c.c.))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{35}/Γ
not seen	PAKHLOVA	08A	BELL $10.6 e^+e^- \rightarrow D^0D^-\pi^+\gamma$	

NODE=M074R16
NODE=M074R16

 $\Gamma(D\bar{D}^*\pi+c.c. (\text{excl. } D^*\bar{D}^*))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{36}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^*\bar{D}\pi$	

NODE=M074R22
NODE=M074R22

 $\Gamma(D\bar{D}^*\pi+c.c. (\text{excl. } D^*\bar{D}^*))/\Gamma(J/\psi\pi^+\pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{36}/Γ_2
<15	90	CRONIN-HEN..09	CLEO	e^+e^-	

NODE=M074R05
NODE=M074R05

$$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}} \quad \Gamma_{37}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

NODE=M074R23
NODE=M074R23

$$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{38}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M074R10
NODE=M074R10

$$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{38}/\Gamma \times \Gamma_1/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42 × 10 ⁻⁶	90	¹ PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M074R11
NODE=M074R11

¹ Using 4263⁺⁸₋₉ MeV for the mass of X(4260).

NODE=M074R11;LINKAGE=PA

$$\Gamma(D^* \bar{D}^* \pi)/\Gamma_{\text{total}} \quad \Gamma_{39}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D}^* \pi$

NODE=M074R24
NODE=M074R24

$$\Gamma(D^* \bar{D}^* \pi)/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{39}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

NODE=M074R06
NODE=M074R06

$$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}} \quad \Gamma_{40}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$

NODE=M074R19
NODE=M074R19

$$\Gamma(D_s^+ D_s^-)/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{40}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	95	DEL-AMO-SA..10N	BABR	10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.3	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

NODE=M074R07
NODE=M074R07

$$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{41}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$

NODE=M074R20
NODE=M074R20

$$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{41}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.8	90	CRONIN-HEN..09	CLEO	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<44	95	DEL-AMO-SA..10N	BABR	10.6 $e^+ e^-$

NODE=M074R08
NODE=M074R08

$$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}} \quad \Gamma_{42}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$

NODE=M074R21
NODE=M074R21

$$\Gamma(D_s^{*+} D_s^{*-})/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{42}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 9.5	90	CRONIN-HEN..09	CLEO	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<30	95	DEL-AMO-SA..10N	BABR	10.6 $e^+ e^-$

NODE=M074R09
NODE=M074R09

$\Gamma(p\bar{p})/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{43}/Γ_2

VALUE	CL%	DOCUMENT ID	COMMENT
<0.13	90	¹ AUBERT	06B $e^+e^- \rightarrow p\bar{p}\gamma$

¹ Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $X(4260)$.NODE=M074R1
NODE=M074R1

NODE=M074R1;LINKAGE=AU

X(4260) REFERENCES

ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BES III Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BES III Collab.)
SHEN	14	PR D89 072015	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BES III Collab.)
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	12AC	PR D86 051102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
YUAN	08	PR D77 011105	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07S	PR D80 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)
YUAN	07	PRL 99 182004	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
HE	06B	PR D74 091104	Q. He <i>et al.</i>	(CLEO Collab.)
AUBERT,B	05I	PRL 95 142001	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M074

REFID=56782
REFID=56816
REFID=55647
REFID=55944
REFID=55409
REFID=55410
REFID=55377
REFID=54750
REFID=53638
REFID=16787
REFID=53532
REFID=52724
REFID=53114
REFID=53143
REFID=52242
REFID=52296
REFID=52132
REFID=52134
REFID=52135
REFID=51908
REFID=52074
REFID=51724
REFID=51628
REFID=51959
REFID=51960
REFID=51017
REFID=51026
REFID=51511
REFID=51075
REFID=51523
REFID=50776**X(4350)**

$$I^G(J^{PC}) = 0^+(?^{?+})$$

OMITTED FROM SUMMARY TABLE

Seen by SHEN 10 in the $\gamma\gamma \rightarrow J/\psi\phi$. Needs confirmation.

NODE=M194

NODE=M194

X(4350) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4350.6^{+4.6}_{-5.1} \pm 0.7$	$8.8^{+4.2}_{-3.2}$	¹ SHEN	10	BELL $10.6 e^+e^- \rightarrow e^+e^- J/\psi\phi$

¹ Statistical significance of 3.2σ .

NODE=M194M

NODE=M194M

NODE=M194M;LINKAGE=SH

X(4350) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$13^{+18}_{-9} \pm 4$	$8.8^{+4.2}_{-3.2}$	¹ SHEN	10	BELL $10.6 e^+e^- \rightarrow e^+e^- J/\psi\phi$

¹ Statistical significance of 3.2σ .

NODE=M194W

NODE=M194W

NODE=M194W;LINKAGE=SH

X(4350) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\phi$	seen
Γ_2 $\gamma\gamma$	seen

NODE=M194215;NODE=M194

DESIG=1

DESIG=2

X(4350) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\gamma\gamma) \times \Gamma(J/\psi\phi)/\Gamma_{\text{total}}$	VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
$6.7^{+3.2}_{-2.4} \pm 1.1$	$8.8^{+4.2}_{-3.2}$	¹ SHEN	10	BELL	$10.6 e^+e^- \rightarrow e^+e^- J/\psi\phi$	

••• We do not use the following data for averages, fits, limits, etc. •••

$1.5^{+0.7}_{-0.6} \pm 0.3$	$8.8^{+4.2}_{-3.2}$	² SHEN	10	BELL	$10.6 e^+e^- \rightarrow e^+e^- J/\psi\phi$	
-----------------------------	---------------------	-------------------	----	------	---	--

¹ For $J^P = 0^+$. Statistical significance of 3.2σ .² For $J^P = 2^+$. Statistical significance of 3.2σ .

NODE=M194220

NODE=M194G01
NODE=M194G01

OCCUR=2

NODE=M194G01;LINKAGE=S0
NODE=M194G01;LINKAGE=S2

X(4350) BRANCHING RATIOS

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	¹ SHEN	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	

¹ Statistical significance of 3.2 σ .

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	¹ SHEN	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	

¹ Statistical significance of 3.2 σ .

X(4350) REFERENCES

SHEN 10 PRL 104 112004 C.P. Shen *et al.* (BELLE Collab.)

X(4360)

$$I^G(J^{PC}) = ?^?(1^{--})$$

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by AUBERT 07S, WANG 07D, and LEES 14F. See also the review under the X(3872) particle listings. (See the index for the page number.)

X(4360) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4346 ± 6 OUR AVERAGE				
[4354 ± 10 MeV OUR 2015 AVERAGE]				
4347 ± 6 ± 3	279	¹ WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4340 ± 16 ± 9	37	² LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4355 ⁺ ₋₁₀ ± 9	74	³ LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4324 ± 24		⁴ AUBERT	07S BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4361 ± 9 ± 9	47	² WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
¹ From a two-resonance fit. Supersedes WANG 07D.				
² From a two-resonance fit.				
³ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.				
⁴ From a single-resonance fit. Systematic errors not estimated.				

X(4360) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
102 ± 10 OUR AVERAGE				
[78 ± 16 MeV OUR 2015 AVERAGE]				
103 ± 9 ± 5	279	¹ WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
94 ± 32 ± 13	37	² LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
103 ⁺ ₋₁₅ ± 11	74	³ LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
172 ± 33		⁴ AUBERT	07S BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
74 ± 15 ± 10	47	² WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
¹ From a two-resonance fit. Supersedes WANG 07D.				
² From a two-resonance fit.				
³ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.				
⁴ From a single-resonance fit. Systematic errors not estimated.				

NODE=M194225

NODE=M194R01
NODE=M194R01

NODE=M194R01;LINKAGE=SH

NODE=M194R02
NODE=M194R02

NODE=M194R02;LINKAGE=SH

NODE=M194

REFID=53235

NODE=M181

NODE=M181

NODE=M181M

NODE=M181M

NEW

NODE=M181M;LINKAGE=A
NODE=M181M;LINKAGE=WA
NODE=M181M;LINKAGE=LI
NODE=M181M;LINKAGE=AU

NODE=M181W

NODE=M181W

NEW

NODE=M181W;LINKAGE=A
NODE=M181W;LINKAGE=WA
NODE=M181W;LINKAGE=LI
NODE=M181W;LINKAGE=AU

X(4360) DECAY MODES

NODE=M181215;NODE=M181

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	
Γ_2 $\psi(2S)\pi^+\pi^-$	seen
Γ_3 $\psi(3823)\pi^+\pi^-$	possibly seen
Γ_4 $J/\psi\eta$	
Γ_5 $D^0D^{*-}\pi^+$	
Γ_6 $\chi_{c1}\gamma$	
Γ_7 $\chi_{c2}\gamma$	

DESIG=1
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=5
DESIG=4
DESIG=3
DESIG=6
DESIG=7

X(4360) $\Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M181230

$\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_2\Gamma_1/\Gamma$**

NODE=M181G1
NODE=M181G1

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

9.2±0.6±0.6	279	¹ WANG	15A	BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
10.9±0.6±0.7	279	² WANG	15A	BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
6.0±1.0±0.5	37	³ LEES	14F	BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
7.2±1.0±0.6	37	⁴ LEES	14F	BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
11.1 ^{+1.3} _{-1.2}	74	⁵ LIU	08H	RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
12.3±1.2	74	⁶ LIU	08H	RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
10.4±1.7±1.5	47	³ WANG	07D	BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
11.8±1.8±1.4	47	⁴ WANG	07D	BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

¹ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

² Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

³ Solution I of two equivalent solutions in a fit using two interfering resonances.

⁴ Solution II of two equivalent solutions in a fit using two interfering resonances.

⁵ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

⁶ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M181G1;LINKAGE=A

NODE=M181G1;LINKAGE=B

NODE=M181G1;LINKAGE=WA

NODE=M181G1;LINKAGE=WN

NODE=M181G1;LINKAGE=LI

NODE=M181G1;LINKAGE=LU

$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_4\Gamma_1/\Gamma$**

NODE=M181G01
NODE=M181G01

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<6.8	90	WANG	13B	BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$
------	----	------	-----	------	---------------------------------------

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_6\Gamma_1/\Gamma$**

NODE=M181G02
NODE=M181G02

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.57	90	¹ HAN	15	BELL	10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$
-------	----	------------------	----	------	--

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M181G02;LINKAGE=A

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_7\Gamma_1/\Gamma$**

NODE=M181G03
NODE=M181G03

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.9	90	¹ HAN	15	BELL	10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$
------	----	------------------	----	------	--

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M181G03;LINKAGE=A

X(4360) BRANCHING RATIOS

NODE=M181225

$\Gamma(D^0D^{*-}\pi^+)/\Gamma(\psi(2S)\pi^+\pi^-)$ **Γ_5/Γ_2**

NODE=M181R01
NODE=M181R01

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<8	90	PAKHLOVA	09	BELL	$e^+e^- \rightarrow X(4360) \rightarrow D^0D^{*-}\pi^+$
----	----	----------	----	------	---

$\Gamma(\psi(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_3/Γ**

NODE=M181R03
NODE=M181R03

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

possibly seen	19	¹ ABLIKIM	15S	BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
---------------	----	----------------------	-----	------	--

¹ From a fit of $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$, $\psi(3823) \rightarrow \chi_{c1}\gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the X(4360) line shape.

NODE=M181R03;LINKAGE=A

$$\frac{\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}}{\text{VALUE}} \times \frac{\Gamma_5/\Gamma \times \Gamma_1/\Gamma}{\text{COMMENT}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.72 \times 10^{-6}$	90	¹ PAKHLOVA	09	BELL $e^+ e^- \rightarrow X(4360) \rightarrow D^0 D^{*-} \pi^+$

¹ Using $4355^{+9}_{-10} \pm 9$ MeV for the mass of X(4360).

NODE=M181R02
NODE=M181R02

NODE=M181R02;LINKAGE=PA

X(4360) REFERENCES

ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56784
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=56839
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55938
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		REFID=52296
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51724
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=51959

NODE=M181

NODE=M073

$\psi(4415)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\psi(4415)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4421 ± 4 OUR ESTIMATE			
4415.1 ± 7.9	¹ ABLIKIM	08D	BES2 $e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4412 ± 15	² MO	10	RVUE $e^+ e^- \rightarrow$ hadrons
4411 ± 7	³ PAKHLOVA	08A	BELL $10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
4425 ± 6	⁴ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
4429 ± 9	⁵ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
4417 ± 10	BRANDELIK	78C	DASP $e^+ e^-$
4414 ± 7	SIEGRIST	76	MRK1 $e^+ e^-$

NODE=M073M

NODE=M073M
→ UNCHECKED ←

OCCUR=2

¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

³ Systematic uncertainties not estimated.

⁴ From a fit to Crystal Ball (OSTERHELD 86) data.

⁵ From a fit to BES (BAI 02C) data.

NODE=M073M;LINKAGE=AB

NODE=M073M;LINKAGE=MO

NODE=M073M;LINKAGE=NS

NODE=M073M;LINKAGE=ST

NODE=M073M;LINKAGE=SE

$\psi(4415)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
62 ± 20 OUR ESTIMATE			
71.5 ± 19.0	⁶ ABLIKIM	08D	BES2 $e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118 ± 32	⁷ MO	10	RVUE $e^+ e^- \rightarrow$ hadrons
77 ± 20	⁸ PAKHLOVA	08A	BELL $10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
119 ± 16	⁹ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
118 ± 35	¹⁰ SETH	05A	RVUE $e^+ e^- \rightarrow$ hadrons
66 ± 15	BRANDELIK	78C	DASP $e^+ e^-$
33 ± 10	SIEGRIST	76	MRK1 $e^+ e^-$

NODE=M073W

NODE=M073W
→ UNCHECKED ←

OCCUR=2

⁶ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

⁷ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

⁸ Systematic uncertainties not estimated.

⁹ From a fit to Crystal Ball (OSTERHELD 86) data.

¹⁰ From a fit to BES (BAI 02C) data.

NODE=M073W;LINKAGE=AB

NODE=M073W;LINKAGE=MO

NODE=M073W;LINKAGE=NS

NODE=M073W;LINKAGE=ST

NODE=M073W;LINKAGE=SE

$\psi(4415)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, "seen" ("not seen") means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle's central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

NODE=M073215;NODE=M073

NODE=M073

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $D\bar{D}$	seen	
Γ_2 $D^0\bar{D}^0$	seen	
Γ_3 D^+D^-	seen	
Γ_4 $D^*\bar{D} + \text{c.c.}$	seen	
Γ_5 $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
Γ_6 $D^*(2010)^+D^- + \text{c.c.}$	seen	
Γ_7 $D^*\bar{D}^*$	seen	
Γ_8 $D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.}$	seen	
Γ_9 $D^*(2010)^+D^*(2010)^- + \text{c.c.}$	seen	
Γ_{10} $D^0D^-\pi^+$ (excl. $D^*(2007)^0\bar{D}^0$ +c.c., $D^*(2010)^+D^-$ +c.c.)	< 2.3 %	90%
Γ_{11} $D\bar{D}_2^*(2460) \rightarrow D^0D^-\pi^+$ +c.c.	(10 \pm 4) %	
Γ_{12} $D^0D^{*-}\pi^+$ +c.c.	< 11 %	90%
Γ_{13} $D_s^+D_s^-$	not seen	
Γ_{14} $\omega\chi_{c2}$	possibly seen	
Γ_{15} $D_s^{*+}D_s^-$ +c.c.	seen	
Γ_{16} $D_s^{*+}D_s^{*-}$	not seen	
Γ_{17} $\psi(3823)\pi^+\pi^-$	possibly seen	
Γ_{18} $J/\psi\eta$	< 6 $\times 10^{-3}$	90%
Γ_{19} $\chi_{c1}\gamma$	< 8 $\times 10^{-4}$	90%
Γ_{20} $\chi_{c2}\gamma$	< 4 $\times 10^{-3}$	90%
Γ_{21} e^+e^-	(9.4 \pm 3.2) $\times 10^{-6}$	

DESIG=7;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=8
DESIG=9
DESIG=10;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=11
DESIG=12
DESIG=13;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=14
DESIG=15
DESIG=4
DESIG=5
DESIG=6
DESIG=16
DESIG=20
DESIG=17
DESIG=18
DESIG=21
DESIG=19
DESIG=22
DESIG=23
DESIG=1

 $\psi(4415)$ PARTIAL WIDTHS

NODE=M073220

 $\Gamma(e^+e^-)$ **Γ_{21}**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.58\pm0.07 OUR ESTIMATE			
0.35\pm0.12	11 ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.4 to 0.8	12 MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
0.72 \pm 0.11	13 SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.64 \pm 0.23	14 SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.49 \pm 0.13	BRANDELIK	78C DASP	e^+e^-
0.44 \pm 0.14	SIEGRIST	76 MRK1	e^+e^-

NODE=M073W1
NODE=M073W1
 \rightarrow UNCHECKED \leftarrow

OCCUR=2

¹¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

NODE=M073W1;LINKAGE=AB

¹² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e^+e^- partial widths. We quote only the range of values.

NODE=M073W1;LINKAGE=MO

¹³ From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M073W1;LINKAGE=ST

¹⁴ From a fit to BES (BAI 02C) data.

NODE=M073W1;LINKAGE=SE

 $\psi(4415)$ $\Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M073230

 $\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{18}\Gamma_{21}/\Gamma$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<3.6	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

NODE=M073G01
NODE=M073G01

$$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{19}\Gamma_{21}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.47	90	¹⁵ HAN	15	BELL 10.58 e ⁺ e ⁻ → χ _{c1} γ
¹⁵ Using B(η → γγ) = (39.41 ± 0.21)%.				

NODE=M073G02
NODE=M073G02

NODE=M073G02;LINKAGE=A

$$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{20}\Gamma_{21}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	¹⁶ HAN	15	BELL 10.58 e ⁺ e ⁻ → χ _{c2} γ
¹⁶ Using B(η → γγ) = (39.41 ± 0.21)%.				

NODE=M073G03
NODE=M073G03

NODE=M073G03;LINKAGE=A

ψ(4415) BRANCHING RATIOS

NODE=M073225

$$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 08	BELL	e ⁺ e ⁻ → D ⁰ \bar{D}^0 γ
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M	BABR	e ⁺ e ⁻ → D ⁰ \bar{D}^0 γ

NODE=M073R04
NODE=M073R04

$$\Gamma(D^+D^-)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 08	BELL	e ⁺ e ⁻ → D ⁺ D ⁻ γ
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M	BABR	e ⁺ e ⁻ → D ⁺ D ⁻ γ

NODE=M073R05
NODE=M073R05

$$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*) \quad \Gamma_1/\Gamma_7$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.14±0.12±0.03	AUBERT 09M	BABR	e ⁺ e ⁻ → γD ^(*) $\bar{D}^{(*)}$

NODE=M073R02
NODE=M073R02

$$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	e ⁺ e ⁻ → D ^{*0} \bar{D}^0 γ

NODE=M073R06
NODE=M073R06

$$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	e ⁺ e ⁻ → D ^{*+} D ⁻ γ
seen	PAKHLOVA 07	BELL	e ⁺ e ⁻ → D ^{*+} D ⁻ γ

NODE=M073R07
NODE=M073R07

$$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*) \quad \Gamma_4/\Gamma_7$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.25±0.03	AUBERT 09M	BABR	e ⁺ e ⁻ → γD ^(*) $\bar{D}^{(*)}$

NODE=M073R03
NODE=M073R03

$$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	e ⁺ e ⁻ → D ^{*0} \bar{D}^{*0} γ

NODE=M073R08
NODE=M073R08

$$\Gamma(D^*(2010)^+D^*(2010)^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	e ⁺ e ⁻ → D ^{*+} D ^{*-} γ
seen	PAKHLOVA 07	BELL	e ⁺ e ⁻ → D ^{*+} D ^{*-} γ

NODE=M073R09
NODE=M073R09

$$\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0D^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
10.5±2.4±3.8	¹⁷ PAKHLOVA 08A	BELL	10.6 e ⁺ e ⁻ → D ⁰ D ⁻ π ⁺ γ

NODE=M073R3
NODE=M073R3

¹⁷ Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of ψ(4415).

NODE=M073R3;LINKAGE=PA

$$\Gamma(D^0D^-\pi^+ (\text{excl. } D^*(2007)^0\bar{D}^0 + \text{c.c.}, D^*(2010)^+D^- + \text{c.c.})/$$

$$\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0D^-\pi^+ + \text{c.c.}) \quad \Gamma_{10}/\Gamma_{11}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	¹⁸ PAKHLOVA 08A	BELL	10.6 e ⁺ e ⁻ → D ⁰ D ⁻ π ⁺ γ

NODE=M073R4
NODE=M073R4

¹⁸ Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of ψ(4415).

NODE=M073R4;LINKAGE=PA

$\Gamma(D^0 D^{*-} \pi^+ + c.c.)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_{12}/\Gamma \times \Gamma_{21}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<0.99 \times 10^{-6}$	90	19 PAKHLOVA	09 BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$	
¹⁹ Using 4421 \pm 4 MeV for the mass of $\psi(4415)$.					

NODE=M073R01
NODE=M073R01

NODE=M073R01;LINKAGE=PA

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$				Γ_{13}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	
not seen		DEL-AMO-SA...10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	

NODE=M073R10
NODE=M073R10

$\Gamma(\omega \chi_{c2})/\Gamma_{\text{total}}$				Γ_{14}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT	
possibly seen		ABLIKIM	16A BES3	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$	

NODE=M073R00
NODE=M073R00

$\Gamma(D_s^{*+} D_s^- + c.c.)/\Gamma_{\text{total}}$				Γ_{15}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$	
seen		DEL-AMO-SA...10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$	

NODE=M073R11
NODE=M073R11

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$				Γ_{16}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$	
not seen		DEL-AMO-SA...10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$	

NODE=M073R12
NODE=M073R12

$\Gamma(\psi(3823) \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{17}/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
possibly seen	19	²⁰ ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$	

NODE=M073R13
NODE=M073R13

²⁰ From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3823)$, $\psi(3823) \rightarrow \chi_{c1} \gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4415)$ line shape.

NODE=M073R13;LINKAGE=A

$\psi(4415)$ REFERENCES

ABLIKIM	16A	PR D93 011102	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=57122
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BES III Collab.)	REFID=56784
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
DEL-AMO-SA...10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532	REFID=53540
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=52724
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53143
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52142
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52132
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=50813
SETH	05A	PR D72 017501	K.K. Seth		REFID=50506
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=51064
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=22232
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22243
SIEGRIST	76	PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)	

NODE=M073

X(4430)[±]

$$I(J^P) = ?(1^+)$$

First seen by CHOI 08 in $B \rightarrow K\pi^+\psi(2S)$ decays, confirmed by AAIJ 14AG, and confirmed in a model-independent way by AAIJ 15BH. Also seen by CHILIKIN 14 in $B \rightarrow K^+\pi J/\psi$ decays. J^P was determined by CHILIKIN 13 and AAIJ 14AG.

NODE=M195

NODE=M195

X(4430)[±] MASS

NODE=M195M

NODE=M195M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

4478⁺¹⁵₋₁₈ OUR AVERAGE

4475 ± 7 ⁺¹⁵ ₋₂₅	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
--	-------------------	-----------	------------------------------------

4485 ± 22 ⁺²⁸ ₋₁₁	¹ CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
---	-----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4443 ⁺¹⁵⁺¹⁹ ₋₁₂₋₁₃	² MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
--	--------------------	---------	--------------------------------

4433 ± 4 ± 2	³ CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
--------------	-------------------	---------	--------------------------------

¹ From a four-dimensional amplitude analysis.

² From a Dalitz plot analysis. Superseded by CHILIKIN 13.

³ Superseded by MIZUK 09 and CHILIKIN 13.

NODE=M195M;LINKAGE=A
NODE=M195M;LINKAGE=MI
NODE=M195M;LINKAGE=CH

X(4430)[±] WIDTH

NODE=M195W

NODE=M195W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

181 ± 31 OUR AVERAGE

172 ± 13 ⁺³⁷ ₋₃₄	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
--	-------------------	-----------	------------------------------------

200 ⁺⁴¹⁺²⁶ ₋₄₆₋₃₅	¹ CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
---	-----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

107 ⁺⁸⁶⁺⁷⁴ ₋₄₃₋₅₆	² MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
---	--------------------	---------	--------------------------------

45 ⁺¹⁸⁺³⁰ ₋₁₃₋₁₃	³ CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
--	-------------------	---------	--------------------------------

¹ From a four-dimensional amplitude analysis.

² From a Dalitz plot analysis. Superseded by CHILIKIN 13.

³ Superseded by MIZUK 09 and CHILIKIN 13.

NODE=M195W;LINKAGE=A
NODE=M195W;LINKAGE=MI
NODE=M195W;LINKAGE=CH

X(4430)[±] DECAY MODES

NODE=M195215;NODE=M195

Mode	Fraction (Γ_i/Γ)
------	--------------------------------

Γ_1 $\pi^+\psi(2S)$	seen
----------------------------	------

DESIG=1

Γ_2 $\pi^+ J/\psi$	seen
---------------------------	------

DESIG=2

X(4430)[±] BRANCHING RATIOS

NODE=M195225

$\Gamma(\pi^+\psi(2S))/\Gamma_{\text{total}}$	Γ_1/Γ
---	-------------------

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
------	-------------------	-----------	------------------------------------

seen	² CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
------	-----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	³ AUBERT	09AA BABR	$B \rightarrow K\pi^+\psi(2S)$
----------	---------------------	-----------	--------------------------------

seen	⁴ MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
------	--------------------	---------	--------------------------------

¹ From a four-dimensional amplitude analysis. No product of branching fractions quoted.

² From a four-dimensional amplitude analysis. Measured a product of branching fractions $B(B^0 \rightarrow X(4430)^- K^+) \times B(X(4430)^- \rightarrow \psi(2S)\pi^-) = (6.0^{+1.7+2.5}_{-2.0-1.4}) \times 10^{-5}$.

³ AUBERT 09AA quotes $B(B^+ \rightarrow \bar{K}^0 X(4430)^+) \times B(X(4430)^+ \rightarrow \pi^+\psi(2S)) < 4.7 \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^- X(4430)^+) \times B(X(4430)^+ \rightarrow \pi^+\psi(2S)) < 3.1 \times 10^{-5}$ at 95% CL.

⁴ Measured a product of branching fractions $B(\bar{B}^0 \rightarrow K^- X(4430)^+) \times B(X(4430)^+ \rightarrow \pi^+\psi(2S)) = (3.2^{+1.8+5.3}_{-0.9-1.6}) \times 10^{-5}$. Superseded by CHILIKIN 13.

NODE=M195R01
NODE=M195R01

NODE=M195R01;LINKAGE=AA
NODE=M195R01;LINKAGE=A

NODE=M195R01;LINKAGE=AU

NODE=M195R01;LINKAGE=MI

$\Gamma(\pi^+ J/\psi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen [not seen OUR 2015 BEST LIMIT]			
seen	¹ CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow K^- \pi^+ J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	² AUBERT 09AA	BABR	$B \rightarrow K \pi^+ J/\psi$
¹ CHILIKIN 14 reports $B(\bar{B}^0 \rightarrow X(4430)^+ K^-) \times B(X(4430)^+ \rightarrow J/\psi \pi^+) = (5.4^{+4.0+1.1}_{-1.0-0.9}) \times 10^{-6}$.			
² AUBERT 09AA quotes $B(B^+ \rightarrow \bar{K}^0 X(4430)^+) \times B(X(4430)^+ \rightarrow \pi^+ J/\psi) < 1.5 \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^- X(4430)^+) \times B(X(4430)^+ \rightarrow \pi^+ J/\psi) < 0.4 \times 10^{-5}$ at 95% CL.			

NODE=M195R02
NODE=M195R02

NODE=M195R02;LINKAGE=A

NODE=M195R02;LINKAGE=AU

X(4430)[±] REFERENCES

AAIJ 15BH PR D92 112009	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ 14AG PRL 112 222002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
CHILIKIN 14 PR D90 112009	K. Chilikin <i>et al.</i>	(BELLE Collab.)
CHILIKIN 13 PR D88 074026	K. Chilikin <i>et al.</i>	(BELLE Collab.) JP
AUBERT 09AA PR D79 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
MIZUK 09 PR D80 031104	R. Mizuk <i>et al.</i>	(BELLE Collab.)
CHOI 08 PRL 100 142001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)

NODE=M195

REFID=57110
REFID=55896
REFID=56344
REFID=55551
REFID=52940
REFID=52960
REFID=52178

NODE=M189

X(4660)

$$I^G(J^{PC}) = ?^?(1^{--})$$

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review under the X(3872) particle listings. (See the index for the page number.)

NODE=M189

X(4660) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4643 ± 9 OUR AVERAGE		Error includes scale factor of 1.2.		[4665 ± 10 MeV OUR 2015 AVERAGE]
4652 ± 10 ± 11	279	¹ WANG 15A	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4669 ± 21 ± 3	37	² LEES 14F	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4634 $^{+8+5}_{-7-8}$	142	³ PAKHLOVA 08B	BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4661 $^{+9}_{-8} \pm 6$	44	⁴ LIU 08H	RVUE	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4664 ± 11 ± 5	44	WANG 07D	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
¹ From a two-resonance fit. Supersedes WANG 07D.				
² From a two-resonance fit.				
³ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.				
⁴ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.				

NODE=M189M

NODE=M189M
NEWNODE=M189M;LINKAGE=A
NODE=M189M;LINKAGE=LE
NODE=M189M;LINKAGE=PA
NODE=M189M;LINKAGE=LI**X(4660) WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
72 ± 11 OUR AVERAGE				[53 ± 16 MeV OUR 2015 AVERAGE Scale factor = 1.1]
68 ± 11 ± 5	279	¹ WANG 15A	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
104 ± 48 ± 10	37	² LEES 14F	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
92 $^{+40+10}_{-24-21}$	142	³ PAKHLOVA 08B	BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
42 $^{+17}_{-12} \pm 6$	44	⁴ LIU 08H	RVUE	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
48 ± 15 ± 3	44	WANG 07D	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
¹ From a two-resonance fit. Supersedes WANG 07D.				
² From a two-resonance fit.				
³ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.				
⁴ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.				

NODE=M189W

NODE=M189W
NEWNODE=M189W;LINKAGE=A
NODE=M189W;LINKAGE=LE
NODE=M189W;LINKAGE=B
NODE=M189W;LINKAGE=LI

X(4660) DECAY MODES

NODE=M189215;NODE=M189

Mode	Fraction (Γ_i/Γ)
Γ_1 $e^+ e^-$	
Γ_2 $\psi(2S)\pi^+\pi^-$	seen
Γ_3 $J/\psi\eta$	
Γ_4 $D^0 D^{*-}\pi^+$	
Γ_5 $\chi_{c1}\gamma$	
Γ_6 $\chi_{c2}\gamma$	
Γ_7 $\Lambda_c^+ \Lambda_c^-$	

DESIG=1
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=4
DESIG=3
DESIG=6
DESIG=7
DESIG=5

X(4660) $\Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

NODE=M189230

 $\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M189G1
NODE=M189G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.0 \pm 0.3 \pm 0.2$	279	1 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$8.1 \pm 1.1 \pm 1.0$	279	2 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$2.7 \pm 1.3 \pm 0.5$	37	3 LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$7.5 \pm 1.7 \pm 0.7$	37	4 LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$2.2^{+0.7}_{-0.6}$	44	5 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
5.9 ± 1.6	44	6 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$3.0 \pm 0.9 \pm 0.3$	44	3 WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
$7.6 \pm 1.8 \pm 0.8$	44	4 WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

¹ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

NODE=M189G1;LINKAGE=A

² Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

NODE=M189G1;LINKAGE=B

³ Solution I of two equivalent solutions in a fit using two interfering resonances.

NODE=M189G1;LINKAGE=WA

⁴ Solution II of two equivalent solutions in a fit using two interfering resonances.

NODE=M189G1;LINKAGE=WN

⁵ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189G1;LINKAGE=LI

⁶ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189G1;LINKAGE=LU

 $\Gamma(J/\psi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M189G01
NODE=M189G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.94	90	WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$
-------	----	------	----------	--

 $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M189G02
NODE=M189G02

<0.45

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M189G02;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_6\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M189G03
NODE=M189G03

<2.1

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M189G03;LINKAGE=A

X(4660) BRANCHING RATIOS

NODE=M189225

 $\Gamma(D^0 D^{*-}\pi^+)/\Gamma(\psi(2S)\pi^+\pi^-)$ Γ_4/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M189R01
NODE=M189R01

<10

 $\Gamma(D^0 D^{*-}\pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_1/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M189R02
NODE=M189R02

<0.37 $\times 10^{-6}$

¹ Using $4664 \pm 11 \pm 5$ MeV for the mass of X(4660).

NODE=M189R02;LINKAGE=PA

$$\Gamma(\Lambda_c^+ \Lambda_c^-) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_7 / \Gamma \times \Gamma_1 / \Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.68^{+0.16+0.29}_{-0.15-0.30}$	142	¹ PAKHLOVA	08B BELL	$e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$

¹ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

NODE=M189R03
NODE=M189R03

NODE=M189R03;LINKAGE=A

X(4660) REFERENCES

HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
PAKHLOVA	08B	PRL 101 172001	C. Pakhlova <i>et al.</i>	(BELLE Collab.)
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)

NODE=M189

REFID=56816
REFID=56839
REFID=55938
REFID=55377
REFID=53143
REFID=52296
REFID=52596
REFID=51724
REFID=51959

$b\bar{b}$ MESONS

NODE=MXXX030

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=M849

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=M849

$\eta_b(1S)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

NODE=M171

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$.

NODE=M171

$\eta_b(1S)$ MASS

NODE=M171M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9399.0 ± 2.3 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
		[9398.0 \pm 3.2 MeV OUR 2015 AVERAGE Scale factor = 1.8]		
$9400.7 \pm 1.7 \pm 1.6$	33.1k	TAMPONI	15 BELL	$e^+ e^- \rightarrow \eta_b(1S) \gamma \eta$
$9402.4 \pm 1.5 \pm 1.8$	34k	¹ MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- +$
$9391.8 \pm 6.6 \pm 2.0$	2.3k	² BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
$9394.2^{+4.8}_{-4.9} \pm 2.0$	13k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
$9388.9^{+3.1}_{-2.3} \pm 2.7$	19k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$

NODE=M171M
NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9393.2 \pm 3.4 \pm 2.3$	10	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
$9300 \pm 20 \pm 20$		HEISTER	02D ALEP	181–209 $e^+ e^-$

¹ With floating width. Not independent of the corresponding mass difference measurement.


² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass difference measurements.

³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M;LINKAGE=MI

NODE=M171M;LINKAGE=AU

NODE=M171M;LINKAGE=DO



$m_{\Upsilon(1S)} - m_{\eta_b}$

NODE=M171M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62.3±3.2 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.
57.9±1.5±1.8	34k	⁴ MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-$ + hadrons
68.5±6.6±2.0	2.3±0.5k	⁵ BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
66.1 ^{+4.8} _{-4.9} ±2.0	13±5k	⁵ AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
71.4 ^{+2.3} _{-3.1} ±2.7	19±3k	⁵ AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
67.1±3.4±2.3	10 ⁺⁵ ₋₄	^{5,6} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons

NODE=M171M2

⁴With floating width. Not independent of the corresponding mass measurement.


⁵Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass measurements.

⁶Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M2;LINKAGE=MI

NODE=M171M2;LINKAGE=AU

NODE=M171M2;LINKAGE=DO



γ ENERGY IN $\Upsilon(3S)$ DECAY

NODE=M171DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M171DM

920.6^{+2.8}_{-3.2} OUR AVERAGE

918.6 \pm 6.0 \pm 1.9	2.3 \pm 0.5k	⁷ BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
921.2 ^{+2.1} _{-2.8} \pm 2.4	19 \pm 3k	⁷ AUBERT	08V	BABR $\Upsilon(3S) \rightarrow \gamma X$

⁷ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171DM;LINKAGE=BO

 γ ENERGY IN $\Upsilon(2S)$ DECAY

NODE=M171U2S

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M171U2S

609.3^{+4.6}_{-4.5} \pm1.9	13 \pm 5k	⁸ AUBERT	09AQ	BABR $\Upsilon(2S) \rightarrow \gamma X$
--	-------------	---------------------	------	--

⁸ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171U2S;LINKAGE=AU

 $\eta_b(1S)$ WIDTH

NODE=M171W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M171W

10 ⁺⁵₋₄ OUR AVERAGE

NEW

[11⁺⁶₋₄ MeV OUR 2015 AVERAGE]

8 ⁺⁶ ₋₅ \pm 5	33.1k	TAMPONI	15	BELL $e^+e^- \rightarrow \eta_b(1S)\gamma\eta$
10.8 ^{+4.0+4.5} _{-3.7-2.0}	34k	⁹ MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^- +$ hadrons

⁹With floating mass.

NODE=M171W;LINKAGE=MI

 $\eta_b(1S)$ DECAY MODES

NODE=M171225;NODE=M171

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $3h^+3h^-$	not seen	
Γ_3 $2h^+2h^-$	not seen	
Γ_4 $4h^+4h^-$		
Γ_5 $\gamma\gamma$	not seen	
Γ_6 $\mu^+\mu^-$	$<9 \times 10^{-3}$	90%
Γ_7 $\tau^+\tau^-$	$<8\%$	90%

DESIG=7

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=4

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=5

DESIG=6

 $\eta_b(1S)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M171230

$\Gamma(3h^+3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_5/\Gamma$
--	---------------------------

NODE=M171G1

NODE=M171G1

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<470	95	ABDALLAH	06	DLPH 161–209 e^+e^-
<132	95	HEISTER	02D	ALEP 181–209 e^+e^-

$\Gamma(2h^+2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_5/\Gamma$
--	---------------------------

NODE=M171G2

NODE=M171G2

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<190	95	ABDALLAH	06	DLPH 161–209 e^+e^-
<48	95	HEISTER	02D	ALEP 181–209 e^+e^-

$\Gamma(4h^+4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_4\Gamma_5/\Gamma$
--	---------------------------

NODE=M171G3

NODE=M171G3

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<660	95	ABDALLAH	06	DLPH 161–209 e^+e^-
------	----	----------	----	-----------------------

$\eta_b(1S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	34k	MIZUK	12	BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$

NODE=M171235

NODE=M171R03
NODE=M171R03

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<9 \times 10^{-3}$	90	¹⁰ AUBERT	09Z	BABR	$e^+e^- \rightarrow \Upsilon(2S, 3S) \rightarrow \gamma\eta_b$
¹⁰ Obtained using $B(\Upsilon(2S) \rightarrow \gamma\eta_b) = (4.2^{+1.1}_{-1.0} \pm 0.9) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma\eta_b) = (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$. This limit is equivalent to $B(\eta_b \rightarrow \mu^+\mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$ measurement.					

NODE=M171R01
NODE=M171R01

NODE=M171R01;LINKAGE=AU

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8 \times 10^{-2}$	90	AUBERT	09P	BABR	$e^+e^- \rightarrow \gamma\tau^+\tau^-$

NODE=M171R02
NODE=M171R02 $\eta_b(1S)$ REFERENCES

TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER	02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)

NODE=M171

REFID=56996
REFID=54288
REFID=54718
REFID=53231
REFID=53106
REFID=53062
REFID=52930
REFID=52262
REFID=51042
REFID=48577

NODE=M049

 $\Upsilon(1S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $\Upsilon(1S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9460.30 ± 0.26 OUR AVERAGE	Error includes scale factor of 3.3.		
9460.51 ± 0.09 ± 0.05	¹ ARTAMONOV 00	MD1	$e^+e^- \rightarrow \text{hadrons}$
9459.97 ± 0.11 ± 0.07	MACKAY 84	REDE	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9460.60 ± 0.09 ± 0.05	^{2,3} BARU	92B	REDE $e^+e^- \rightarrow \text{hadrons}$
9460.59 ± 0.12	BARU	86	REDE $e^+e^- \rightarrow \text{hadrons}$
9460.6 ± 0.4	^{3,4} ARTAMONOV 84	REDE	$e^+e^- \rightarrow \text{hadrons}$

NODE=M049M

NODE=M049M

¹ Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).² Superseding BARU 86.³ Superseded by ARTAMONOV 00.⁴ Value includes data of ARTAMONOV 82.NODE=M049M;LINKAGE=AR
NODE=M049M;LINKAGE=A
NODE=M049M;LINKAGE=RZ
NODE=M049M;LINKAGE=G $\Upsilon(1S)$ WIDTH

VALUE (keV)	DOCUMENT ID
54.02 ± 1.25 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

NODE=M049W

NODE=M049W
→ UNCHECKED ← $\Upsilon(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\tau^+\tau^-$	(2.60 ± 0.10) %	DESIG=3
Γ_2 e^+e^-	(2.38 ± 0.11) %	DESIG=2
Γ_3 $\mu^+\mu^-$	(2.48 ± 0.05) %	DESIG=1

NODE=M049215;NODE=M049

Hadronic decays

				NODE=M049;CLUMP=A
Γ ₄	ggg	(81.7 ± 0.7)%		DESIG=117
Γ ₅	$\gamma g g$	(2.2 ± 0.6)%		DESIG=118
Γ ₆	$\eta'(958)$ anything	(2.94 ± 0.24)%		DESIG=73
Γ ₇	$J/\psi(1S)$ anything	(6.5 ± 0.7) × 10 ⁻⁴		DESIG=12
Γ ₈	$J/\psi(1S)\eta_c$	< 2.2 × 10 ⁻⁶	90%	DESIG=146
Γ ₉	$J/\psi(1S)\chi_{c0}$	< 3.4 × 10 ⁻⁶	90%	DESIG=147
Γ ₁₀	$J/\psi(1S)\chi_{c1}$	(3.9 ± 1.2) × 10 ⁻⁶		DESIG=148
Γ ₁₁	$J/\psi(1S)\chi_{c2}$	< 1.4 × 10 ⁻⁶	90%	DESIG=149
Γ ₁₂	$J/\psi(1S)\eta_c(2S)$	< 2.2 × 10 ⁻⁶	90%	DESIG=150
Γ ₁₃	$J/\psi(1S)X(3940)$	< 5.4 × 10 ⁻⁶	90%	DESIG=151
Γ ₁₄	$J/\psi(1S)X(4160)$	< 5.4 × 10 ⁻⁶	90%	DESIG=152
Γ ₁₅	χ_{c0} anything	< 5 × 10 ⁻³	90%	DESIG=5
Γ ₁₆	χ_{c1} anything	(2.3 ± 0.7) × 10 ⁻⁴		DESIG=6
Γ ₁₇	χ_{c2} anything	(3.4 ± 1.0) × 10 ⁻⁴		DESIG=7
Γ ₁₈	$\psi(2S)$ anything	(2.7 ± 0.9) × 10 ⁻⁴		DESIG=8
Γ ₁₉	$\psi(2S)\eta_c$	< 3.6 × 10 ⁻⁶	90%	DESIG=153
Γ ₂₀	$\psi(2S)\chi_{c0}$	< 6.5 × 10 ⁻⁶	90%	DESIG=154
Γ ₂₁	$\psi(2S)\chi_{c1}$	< 4.5 × 10 ⁻⁶	90%	DESIG=155
Γ ₂₂	$\psi(2S)\chi_{c2}$	< 2.1 × 10 ⁻⁶	90%	DESIG=156
Γ ₂₃	$\psi(2S)\eta_c(2S)$	< 3.2 × 10 ⁻⁶	90%	DESIG=157
Γ ₂₄	$\psi(2S)X(3940)$	< 2.9 × 10 ⁻⁶	90%	DESIG=158
Γ ₂₅	$\psi(2S)X(4160)$	< 2.9 × 10 ⁻⁶	90%	DESIG=159
Γ ₂₆	$\rho\pi$	< 3.68 × 10 ⁻⁶	90%	DESIG=11
Γ ₂₇	$\omega\pi^0$	< 3.90 × 10 ⁻⁶	90%	DESIG=131
Γ ₂₈	$\pi^+\pi^-$	< 5 × 10 ⁻⁴	90%	DESIG=23
Γ ₂₉	K^+K^-	< 5 × 10 ⁻⁴	90%	DESIG=24
Γ ₃₀	$\rho\bar{\rho}$	< 5 × 10 ⁻⁴	90%	DESIG=25
Γ ₃₁	$\pi^+\pi^-\pi^0$	(2.1 ± 0.8) × 10 ⁻⁶		DESIG=72
Γ ₃₂	ϕK^+K^-	(2.4 ± 0.5) × 10 ⁻⁶		DESIG=136
Γ ₃₃	$\omega\pi^+\pi^-$	(4.5 ± 1.0) × 10 ⁻⁶		DESIG=137
Γ ₃₄	$K^*(892)^0 K^- \pi^+ + c.c.$	(4.4 ± 0.8) × 10 ⁻⁶		DESIG=138
Γ ₃₅	$\phi f'_2(1525)$	< 1.63 × 10 ⁻⁶	90%	DESIG=139
Γ ₃₆	$\omega f_2(1270)$	< 1.79 × 10 ⁻⁶	90%	DESIG=140
Γ ₃₇	$\rho(770) a_2(1320)$	< 2.24 × 10 ⁻⁶	90%	DESIG=141
Γ ₃₈	$K^*(892)^0 K_2^*(1430)^0 + c.c.$	(3.0 ± 0.8) × 10 ⁻⁶		DESIG=142
Γ ₃₉	$K_1(1270)^\pm K^\mp$	< 2.41 × 10 ⁻⁶	90%	DESIG=143
Γ ₄₀	$K_1(1400)^\pm K^\mp$	(1.0 ± 0.4) × 10 ⁻⁶		DESIG=144
Γ ₄₁	$b_1(1235)^\pm \pi^\mp$	< 1.25 × 10 ⁻⁶	90%	DESIG=145
Γ ₄₂	$\pi^+\pi^-\pi^0\pi^0$	(1.28 ± 0.30) × 10 ⁻⁵		DESIG=132
Γ ₄₃	$K_S^0 K^+ \pi^- + c.c.$	(1.6 ± 0.4) × 10 ⁻⁶		DESIG=133
Γ ₄₄	$K^*(892)^0 \bar{K}^0 + c.c.$	(2.9 ± 0.9) × 10 ⁻⁶		DESIG=134
Γ ₄₅	$K^*(892)^- K^+ + c.c.$	< 1.11 × 10 ⁻⁶	90%	DESIG=135
Γ ₄₆	$D^*(2010)^\pm$ anything	(2.52 ± 0.20)%		DESIG=30
Γ ₄₇	2H anything	(2.85 ± 0.25) × 10 ⁻⁵		DESIG=107
Γ ₄₈	Sum of 100 exclusive modes	(1.200 ± 0.017)%		DESIG=128

Radiative decays

				NODE=M049;CLUMP=B
Γ ₄₉	$\gamma\pi^+\pi^-$	(6.3 ± 1.8) × 10 ⁻⁵		DESIG=70
Γ ₅₀	$\gamma\pi^0\pi^0$	(1.7 ± 0.7) × 10 ⁻⁵		DESIG=71
Γ ₅₁	$\gamma\pi^0\eta$	< 2.4 × 10 ⁻⁶	90%	DESIG=111
Γ ₅₂	γK^+K^-	[a] (1.14 ± 0.13) × 10 ⁻⁵		DESIG=102
Γ ₅₃	$\gamma\rho\bar{\rho}$	[b] < 6 × 10 ⁻⁶	90%	DESIG=103
Γ ₅₄	$\gamma 2h^+ 2h^-$	(7.0 ± 1.5) × 10 ⁻⁴		DESIG=20
Γ ₅₅	$\gamma 3h^+ 3h^-$	(5.4 ± 2.0) × 10 ⁻⁴		DESIG=21
Γ ₅₆	$\gamma 4h^+ 4h^-$	(7.4 ± 3.5) × 10 ⁻⁴		DESIG=22
Γ ₅₇	$\gamma\pi^+\pi^- K^+ K^-$	(2.9 ± 0.9) × 10 ⁻⁴		DESIG=14
Γ ₅₈	$\gamma 2\pi^+ 2\pi^-$	(2.5 ± 0.9) × 10 ⁻⁴		DESIG=13
Γ ₅₉	$\gamma 3\pi^+ 3\pi^-$	(2.5 ± 1.2) × 10 ⁻⁴		DESIG=17
Γ ₆₀	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	(2.4 ± 1.2) × 10 ⁻⁴		DESIG=18
Γ ₆₁	$\gamma\pi^+\pi^-\rho\bar{\rho}$	(1.5 ± 0.6) × 10 ⁻⁴		DESIG=15
Γ ₆₂	$\gamma 2\pi^+ 2\pi^- \rho\bar{\rho}$	(4 ± 6) × 10 ⁻⁵		DESIG=19

Γ ₆₃	$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$			DESIG=16
Γ ₆₄	$\gamma \eta'(958)$	$< 1.9 \times 10^{-6}$	90%		DESIG=55
Γ ₆₅	$\gamma \eta$	$< 1.0 \times 10^{-6}$	90%		DESIG=54
Γ ₆₆	$\gamma f_0(980)$	$< 3 \times 10^{-5}$	90%		DESIG=105
Γ ₆₇	$\gamma f'_2(1525)$	$(3.8 \pm 0.9) \times 10^{-5}$			DESIG=52
Γ ₆₈	$\gamma f_2(1270)$	$(1.01 \pm 0.09) \times 10^{-4}$			DESIG=51
Γ ₆₉	$\gamma \eta(1405)$	$< 8.2 \times 10^{-5}$	90%		DESIG=65
Γ ₇₀	$\gamma f_0(1500)$	$< 1.5 \times 10^{-5}$	90%		DESIG=108
Γ ₇₁	$\gamma f_0(1710)$	$< 2.6 \times 10^{-4}$	90%		DESIG=53
Γ ₇₂	$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$	$< 7 \times 10^{-6}$	90%		DESIG=112
Γ ₇₃	$\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0$	$< 1.4 \times 10^{-6}$	90%		DESIG=109
Γ ₇₄	$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$< 1.8 \times 10^{-6}$	90%		DESIG=110
Γ ₇₅	$\gamma f_4(2050)$	$< 5.3 \times 10^{-5}$	90%		DESIG=104
Γ ₇₆	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	$< 2 \times 10^{-4}$	90%		DESIG=69
Γ ₇₇	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	$< 8 \times 10^{-7}$	90%		DESIG=60
Γ ₇₈	$\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-$	$< 6 \times 10^{-7}$	90%		DESIG=61
Γ ₇₉	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$< 1.1 \times 10^{-6}$	90%		DESIG=62
Γ ₈₀	$\gamma \eta(2225) \rightarrow \gamma \phi \phi$	$< 3 \times 10^{-3}$	90%		DESIG=68
Γ ₈₁	$\gamma \eta_c(1S)$	$< 5.7 \times 10^{-5}$	90%		DESIG=119
Γ ₈₂	$\gamma \chi_{c0}$	$< 6.5 \times 10^{-4}$	90%		DESIG=120
Γ ₈₃	$\gamma \chi_{c1}$	$< 2.3 \times 10^{-5}$	90%		DESIG=121
Γ ₈₄	$\gamma \chi_{c2}$	$< 7.6 \times 10^{-6}$	90%		DESIG=122
Γ ₈₅	$\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi$	$< 1.6 \times 10^{-6}$	90%		DESIG=123
Γ ₈₆	$\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	$< 2.8 \times 10^{-6}$	90%		DESIG=124
Γ ₈₇	$\gamma X(3915) \rightarrow \omega J/\psi$	$< 3.0 \times 10^{-6}$	90%		DESIG=125
Γ ₈₈	$\gamma X(4140) \rightarrow \phi J/\psi$	$< 2.2 \times 10^{-6}$	90%		DESIG=126
Γ ₈₉	γX	[c] $< 4.5 \times 10^{-6}$	90%		DESIG=66
Γ ₉₀	$\gamma X \bar{X} (m_X < 3.1 \text{ GeV})$	[d] $< 1 \times 10^{-3}$	90%		DESIG=67
Γ ₉₁	$\gamma X \bar{X} (m_X < 4.5 \text{ GeV})$	[e] $< 2.4 \times 10^{-4}$	90%		DESIG=127
Γ ₉₂	$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	[f] $< 1.78 \times 10^{-4}$	95%		DESIG=113
Γ ₉₃	$\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$	[g] $< 9 \times 10^{-6}$	90%		DESIG=114
Γ ₉₄	$\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$	[a] $< 1.30 \times 10^{-4}$	90%		DESIG=115
Γ ₉₅	$\gamma a_1^0 \rightarrow \gamma g g$	[h] $< 1 \%$	90%		DESIG=129
Γ ₉₆	$\gamma a_1^0 \rightarrow \gamma s \bar{s}$	[h] $< 1 \times 10^{-3}$	90%		DESIG=130
Lepton Family number (LF) violating modes					
Γ ₉₇	$\mu^\pm \tau^\mp$	LF $< 6.0 \times 10^{-6}$	95%		NODE=M049;CLUMP=C DESIG=116
Other decays					
Γ ₉₈	invisible	$< 3.0 \times 10^{-4}$	90%		NODE=M049;CLUMP=D DESIG=106

[a] $2m_\tau < M(\tau^+ \tau^-) < 9.2 \text{ GeV}$

[b] $2 \text{ GeV} < m_{K^+ K^-} < 3 \text{ GeV}$

[c] $X = \text{scalar with } m < 8.0 \text{ GeV}$

[d] $X \bar{X} = \text{vectors with } m < 3.1 \text{ GeV}$

[e] $X \text{ and } \bar{X} = \text{zero spin with } m < 4.5 \text{ GeV}$

[f] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

[g] $201 \text{ MeV} < M(\mu^+ \mu^-) < 3565 \text{ MeV}$

[h] $0.5 \text{ GeV} < m_X < 9.0 \text{ GeV}$, where m_X is the invariant mass of the hadronic final state.

$\Upsilon(1S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

NODE=M049218

$\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

$\Gamma_2 \Gamma_3 / \Gamma$

NODE=M049G1
NODE=M049G1

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
31.2±1.6±1.7	KOBEL	92	CBAL $e^+ e^- \rightarrow \mu^+ \mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_0\Gamma_2/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.240±0.016 OUR AVERAGE			
1.252±0.004±0.019	⁵ ROSNER 06	CLEO	9.5 e ⁺ e ⁻ → hadrons
1.187±0.023±0.031	⁵ BARU 92B	MD1	e ⁺ e ⁻ → hadrons
1.23 ±0.02 ±0.05	⁵ JAKUBOWSKI 88	CBAL	e ⁺ e ⁻ → hadrons
1.37 ±0.06 ±0.09	⁶ GILES 84B	CLEO	e ⁺ e ⁻ → hadrons
1.23 ±0.08 ±0.04	⁶ ALBRECHT 82	DASP	e ⁺ e ⁻ → hadrons
1.13 ±0.07 ±0.11	⁶ NICZYPORUK 82	LENA	e ⁺ e ⁻ → hadrons
1.09 ±0.25	⁶ BOCK 80	CNTR	e ⁺ e ⁻ → hadrons
1.35 ±0.14	⁷ BERGER 79	PLUT	e ⁺ e ⁻ → hadrons

⁵ Radiative corrections evaluated following KURAEV 85.⁶ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.⁷ Radiative corrections reevaluated by ALEXANDER 89 using B($\mu\mu$) = 0.026.NODE=M049G2
NODE=M049G2NODE=M049G2;LINKAGE=B
NODE=M049G2;LINKAGE=R
NODE=M049G2;LINKAGE=P $\Upsilon(1S)$ PARTIAL WIDTHS

NODE=M049220

 $\Gamma(e^+e^-)$ Γ_2

VALUE (keV)	DOCUMENT ID
1.340±0.018 OUR EVALUATION	

NODE=M049W2
NODE=M049W2
→ UNCHECKED ← $\Upsilon(1S)$ BRANCHING RATIOS

NODE=M049225

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
2.60±0.10 OUR AVERAGE				
2.53±0.13±0.05	60k	⁸ BESSON 07	CLEO	e ⁺ e ⁻ → $\Upsilon(1S)$ → $\tau^+\tau^-$
2.61±0.12 ^{+0.09} _{-0.13}	25k	CINABRO 94B	CLE2	e ⁺ e ⁻ → $\tau^+\tau^-$
2.7 ±0.4 ±0.2		⁹ ALBRECHT 85C	ARG	$\Upsilon(2S)$ → $\pi^+\pi^-\tau^+\tau^-$
3.4 ±0.4 ±0.4		GILES 83	CLEO	e ⁺ e ⁻ → $\tau^+\tau^-$

NODE=M049R3
NODE=M049R3⁸ BESSON 07 reports $[\Gamma(\Upsilon(1S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = 1.02 \pm 0.02 \pm 0.05$ which we multiply by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.⁹ Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.

NODE=M049R3;LINKAGE=BE

NODE=M049R3;LINKAGE=A

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
2.38±0.11 OUR AVERAGE				
2.29±0.08±0.11		ALEXANDER 98	CLE2	$\Upsilon(2S)$ → $\pi^+\pi^-e^+e^-$
2.42±0.14±0.14	307	ALBRECHT 87	ARG	$\Upsilon(2S)$ → $\pi^+\pi^-e^+e^-$
2.8 ±0.3 ±0.2	826	BESSON 84	CLEO	$\Upsilon(2S)$ → $\pi^+\pi^-e^+e^-$
5.1 ±3.0		BERGER 80C	PLUT	e ⁺ e ⁻ → e ⁺ e ⁻

NODE=M049R2
NODE=M049R2 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0248±0.0005 OUR AVERAGE				
0.0249±0.0002±0.0007	345k	ADAMS 05	CLEO	e ⁺ e ⁻ → $\mu^+\mu^-$
0.0249±0.0008±0.0013		ALEXANDER 98	CLE2	$\Upsilon(2S)$ → $\pi^+\pi^-\mu^+\mu^-$
0.0212±0.0020±0.0010		¹⁰ BARU 92	MD1	e ⁺ e ⁻ → $\mu^+\mu^-$
0.0231±0.0012±0.0010		¹⁰ KOBEL 92	CBAL	e ⁺ e ⁻ → $\mu^+\mu^-$
0.0252±0.0007±0.0007		CHEN 89B	CLEO	e ⁺ e ⁻ → $\mu^+\mu^-$
0.0261±0.0009±0.0011		KAARSBERG 89	CSB2	e ⁺ e ⁻ → $\mu^+\mu^-$
0.0230±0.0025±0.0013	86	ALBRECHT 87	ARG	$\Upsilon(2S)$ → $\pi^+\pi^-\mu^+\mu^-$
0.029 ±0.003 ±0.002	864	BESSON 84	CLEO	$\Upsilon(2S)$ → $\pi^+\pi^-\mu^+\mu^-$
0.027 ±0.003 ±0.003		ANDREWS 83	CLEO	e ⁺ e ⁻ → $\mu^+\mu^-$
0.032 ±0.013 ±0.003		ALBRECHT 82	DASP	e ⁺ e ⁻ → $\mu^+\mu^-$
0.038 ±0.015 ±0.002		NICZYPORUK 82	LENA	e ⁺ e ⁻ → $\mu^+\mu^-$
0.014 ^{+0.034} _{-0.014}		BOCK 80	CNTR	e ⁺ e ⁻ → $\mu^+\mu^-$
0.022 ±0.020		BERGER 79	PLUT	e ⁺ e ⁻ → $\mu^+\mu^-$

NODE=M049R1
NODE=M049R1¹⁰ Taking into account interference between the resonance and continuum.

NODE=M049R1;LINKAGE=G

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_1/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.008±0.023 OUR AVERAGE				
1.005±0.013±0.022	0.7M	¹¹ DEL-AMO-SA..10C	BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\Upsilon(1S)$
1.02 ±0.02 ±0.05	60k	BESSION 07	CLEO	$e^+e^- \rightarrow \Upsilon(1S)$

NODE=M049R43
NODE=M049R43¹¹ Allows any number of extra photons with total energy < 500 MeV.

NODE=M049R43;LINKAGE=DE

 $\Gamma(ggg)/\Gamma_{total}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
81.7±0.7	20M	¹² BESSION 06A	CLEO	$\Upsilon(1S) \rightarrow$ hadrons

NODE=M049R35
NODE=M049R35¹² Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ from BESSION 06A and PDG 08 values of $B(\mu^+\mu^-) = (2.48 \pm 0.05)\%$ and $R_{hadrons} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{total}$ measurement of BESSION 06A.

NODE=M049R35;LINKAGE=BE

 $\Gamma(\gamma gg)/\Gamma_{total}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20±0.60	400k	¹³ BESSION 06A	CLEO	$\Upsilon(1S) \rightarrow \gamma +$ hadrons

NODE=M049R36
NODE=M049R36¹³ Calculated using BESSION 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ and $\Gamma(ggg)/\Gamma_{total}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{total}$ measurement of BESSION 06A.

NODE=M049R36;LINKAGE=BE

 $\Gamma(\gamma gg)/\Gamma(ggg)$ Γ_5/Γ_4

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.70±0.01±0.27	20M	BESSION 06A	CLEO	$\Upsilon(1S) \rightarrow (\gamma +)$ hadrons

NODE=M049R37
NODE=M049R37 $\Gamma(\eta'(958) \text{ anything})/\Gamma_{total}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0294±0.0024 OUR AVERAGE			
0.030 ±0.002 ±0.002	AQUINES 06A	CLE3	$\Upsilon(1S) \rightarrow \eta'$ anything
0.028 ±0.004 ±0.002	ARTUSO 03	CLE2	$\Upsilon(1S) \rightarrow \eta'$ anything

NODE=M049R73
NODE=M049R73 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{total}$ Γ_7/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.65±0.07 OUR AVERAGE					
0.64±0.04±0.06		730 ± 40	BRIERE 04	CLEO	$e^+e^- \rightarrow J/\psi X$
1.1 ±0.4 ±0.2			¹⁴ FULTON 89	CLEO	$e^+e^- \rightarrow \mu^+\mu^- X$

NODE=M049R12
NODE=M049R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.68	90	ALBRECHT 92J	ARG	$e^+e^- \rightarrow e^+e^- X, \mu^+\mu^- X$
<1.7	90	MASCHMANN 90	CBAL	$e^+e^- \rightarrow$ hadrons
<20	90	NICZYPORUK 83	LENA	

¹⁴ Using $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$.

NODE=M049R12;LINKAGE=K

 $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{total}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10⁻⁶	90	YANG 14	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M049R85
NODE=M049R85 $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{total}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.4 × 10⁻⁶	90	YANG 14	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M049R86
NODE=M049R86 $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{total}$ Γ_{10}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.90±1.21±0.23	20	YANG 14	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M049R87
NODE=M049R87 $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{total}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10⁻⁶	90	YANG 14	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M049R88
NODE=M049R88 $\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{total}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10⁻⁶	90	YANG 14	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M049R89
NODE=M049R89

$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$					Γ_{13}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<5.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M049R90 NODE=M049R90
$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$					Γ_{14}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<5.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M049R91 NODE=M049R91
$\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{15}/Γ_7	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<7.4	90	BRIERE	04	CLEO	$e^+ e^- \rightarrow J/\psi X$	NODE=M049R25 NODE=M049R25
$\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{16}/Γ_7	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.35 \pm 0.08 \pm 0.06$	52 ± 12	BRIERE	04	CLEO	$e^+ e^- \rightarrow J/\psi X$	NODE=M049R26 NODE=M049R26
$\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{17}/Γ_7	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.52 \pm 0.12 \pm 0.09$	47 ± 11	BRIERE	04	CLEO	$e^+ e^- \rightarrow J/\psi X$	NODE=M049R27 NODE=M049R27
$\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{18}/Γ_7	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
$0.41 \pm 0.11 \pm 0.08$	42 ± 11	BRIERE	04	CLEO	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- X$	NODE=M049R28 NODE=M049R28
$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$					Γ_{19}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.6 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M049R92 NODE=M049R92
$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$					Γ_{20}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<6.5 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M049R93 NODE=M049R93
$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$					Γ_{21}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<4.5 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M049R94 NODE=M049R94
$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$					Γ_{22}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<2.1 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M049R95 NODE=M049R95
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$					Γ_{23}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.2 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M049R96 NODE=M049R96
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$					Γ_{24}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<2.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M049R97 NODE=M049R97
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$					Γ_{25}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<2.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M049R98 NODE=M049R98
$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					Γ_{26}/Γ	
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<3.68	90	SHEN	13	BELL	$\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0$	NODE=M049R11 NODE=M049R11
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<1 \times 10^3$	90	BLINOV	90	MD1	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$	
$<2 \times 10^2$	90	FULTON	90B		$\Upsilon(1S) \rightarrow \rho^0 \pi^0$	
$<2.1 \times 10^3$	90	NICZYPORUK	83	LENA	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$	
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					Γ_{27}/Γ	
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<3.90	90	SHEN	13	BELL	$\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	NODE=M049R05 NODE=M049R05

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{28}/Γ	NODE=M049R57 NODE=M049R57
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<5	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow \pi^+\pi^-$			
$\Gamma(K^+K^-)/\Gamma_{\text{total}}$						Γ_{29}/Γ	NODE=M049R58 NODE=M049R58
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<5	90	BARU	92 MD1	$\Upsilon(1S) \rightarrow K^+K^-$			
$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$						Γ_{30}/Γ	NODE=M049R59 NODE=M049R59
<u>VALUE (units 10⁻⁴)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<5	90	¹⁵ BARU	96 MD1	$\Upsilon(1S) \rightarrow \rho\bar{\rho}$			
¹⁵ Supersedes BARU 92 in this node.							NODE=M049R59;LINKAGE=A
$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						Γ_{31}/Γ	NODE=M049R72 NODE=M049R72
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
2.14±0.72±0.34		26 ± 9	SHEN	13 BELL	$\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0$		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<18.4	90		ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$		
$\Gamma(\phi K^+K^-)/\Gamma_{\text{total}}$						Γ_{32}/Γ	NODE=M049R75 NODE=M049R75
<u>VALUE (units 10⁻⁶)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
2.36±0.37±0.29	56	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(K^+K^-)$			
$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{33}/Γ	NODE=M049R76 NODE=M049R76
<u>VALUE (units 10⁻⁶)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
4.46±0.67±0.72	64	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$			
$\Gamma(K^*(892)^0 K^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{34}/Γ	NODE=M049R77 NODE=M049R77
<u>VALUE (units 10⁻⁶)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
4.42±0.50±0.58	173	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$			
$\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$						Γ_{35}/Γ	NODE=M049R78 NODE=M049R78
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<1.63	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(K^+K^-)$			
$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$						Γ_{36}/Γ	NODE=M049R79 NODE=M049R79
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<1.79	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$			
$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$						Γ_{37}/Γ	NODE=M049R80 NODE=M049R80
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<2.24	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$			
$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{38}/Γ	NODE=M049R81 NODE=M049R81
<u>VALUE (units 10⁻⁶)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
3.02±0.68±0.34	42	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$			
$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$						Γ_{39}/Γ	NODE=M049R82 NODE=M049R82
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<2.41	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$			
$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$						Γ_{40}/Γ	NODE=M049R83 NODE=M049R83
<u>VALUE (units 10⁻⁶)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
1.02±0.35±0.22	24	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$			
$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$						Γ_{41}/Γ	NODE=M049R84 NODE=M049R84
<u>VALUE (units 10⁻⁶)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<1.25	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$			
$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$						Γ_{42}/Γ	NODE=M049R06 NODE=M049R06
<u>VALUE (units 10⁻⁶)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
12.8±2.0±2.3	143 ± 22	SHEN	13 BELL	$\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0$			

$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.59±0.33±0.18		37 ± 8	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R07
 NODE=M049R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	¹⁶ DOBBS	12A		$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$
------	----	---------------------	-----	--	--

¹⁶ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M049R07;LINKAGE=DO

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
2.92±0.85±0.37	16 ± 5	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R08
 NODE=M049R08

 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.11	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R09
 NODE=M049R09

 $\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
25.2±1.3±1.5		≈ 2k	¹⁷ AUBERT	10C	BABR $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

NODE=M049R32
 NODE=M049R32

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	¹⁸ ALBRECHT	92J	ARG	$e^+ e^- \rightarrow D^0 \pi^\pm X$
-----	----	------------------------	-----	-----	-------------------------------------

¹⁷ For $x_p > 0.1$.

¹⁸ For $x_p > 0.2$.

NODE=M049R32;LINKAGE=AU
 NODE=M049R32;LINKAGE=B

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.85±0.25 OUR AVERAGE				
[(2.86 ± 0.28) × 10 ⁻⁵ OUR 2015 AVERAGE]				
2.81±0.49 ^{+0.20} _{-0.24}		LEES	14G	BABR $e^+ e^- \rightarrow \overline{2H} X$
2.86±0.19±0.21	455	ASNER	07	CLEO $e^+ e^- \rightarrow \overline{2H} X$

NODE=M049R33
 NODE=M049R33

NEW

OCCUR=2

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.200±0.017	^{19,20} DOBBS	12A	$\Upsilon(1S) \rightarrow \text{hadrons}$

NODE=M049R02
 NODE=M049R02

¹⁹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

NODE=M049R02;LINKAGE=DO

²⁰ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M049R02;LINKAGE=NC

 $\Gamma(ggg, \gamma gg \rightarrow \bar{d} \text{ anything})/\Gamma(ggg, \gamma gg \rightarrow \text{anything})$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.36±0.23±0.25	455	ASNER	07	CLEO $e^+ e^- \rightarrow \bar{d} X$

NODE=M049R34
 NODE=M049R34

 $\Gamma(\gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
6.3±1.2±1.3	²¹ ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R70
 NODE=M049R70

²¹ For $m_{\pi\pi} > 1 \text{ GeV}$.

NODE=M049R70;LINKAGE=A

 $\Gamma(\gamma \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.7±0.6±0.3	²² ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R71
 NODE=M049R71

²² For $m_{\pi\pi} > 1 \text{ GeV}$.

NODE=M049R71;LINKAGE=A

 $\Gamma(\gamma \pi^0 \eta)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.4	90	²³ BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S)$

NODE=M049R47
 NODE=M049R47

²³ BESSON 07A obtained this limit for $0.7 < m_{\pi^0 \eta} < 3 \text{ GeV}$.

NODE=M049R47;LINKAGE=BE

 $\Gamma(\gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
1.14±0.08±0.10	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R24

NODE=M049R24
 NODE=M049R24

$\Gamma(\gamma\rho\bar{\rho})/\Gamma_{\text{total}}$ ($2 < m_{\rho\bar{\rho}} < 3 \text{ GeV}$)					Γ_{53}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<0.6	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma\rho\bar{\rho}$		NODE=M049R29 NODE=M049R29 NODE=M049R29
$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$					Γ_{54}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
7.0±1.1±1.0	80 ± 12	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R20 NODE=M049R20
$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$					Γ_{55}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
5.4±1.5±1.3	39 ± 11	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R21 NODE=M049R21
$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$					Γ_{56}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
7.4±2.5±2.5	36 ± 12	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R22 NODE=M049R22
$\Gamma(\gamma\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$					Γ_{57}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
2.9±0.7±0.6	29 ± 8	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R14 NODE=M049R14
$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$					Γ_{58}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
2.5±0.7±0.5	26 ± 7	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R13 NODE=M049R13
$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$					Γ_{59}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
2.5±0.9±0.8	17 ± 5	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R17 NODE=M049R17
$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{60}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
2.4±0.9±0.8	18 ± 7	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R18 NODE=M049R18
$\Gamma(\gamma\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$					Γ_{61}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.5±0.5±0.3	22 ± 6	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R15 NODE=M049R15
$\Gamma(\gamma 2\pi^+ 2\pi^- \rho\bar{\rho})/\Gamma_{\text{total}}$					Γ_{62}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.4±0.4±0.4	7 ± 6	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R19 NODE=M049R19
$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$					Γ_{63}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.2±0.2	2 ± 2	FULTON	90B CLEO	$e^+e^- \rightarrow \text{hadrons}$		NODE=M049R16 NODE=M049R16
$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$					Γ_{64}/Γ	
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
< 1.9	90	ATHAR	07A CLEO	$\Upsilon(1S) \rightarrow \gamma\eta' \rightarrow \gamma\pi^+\pi^-\eta, \gamma\rho$		NODE=M049R55 NODE=M049R55
••• We do not use the following data for averages, fits, limits, etc. •••						
<16	90	RICHICHI	01B CLE2	$\Upsilon(1S) \rightarrow \gamma\eta' \rightarrow \gamma\eta\pi^+\pi^-$		
$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$					Γ_{65}/Γ	
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
< 1.0	90	ATHAR	07A CLEO	$\Upsilon(1S) \rightarrow \gamma\eta \rightarrow \gamma\gamma\gamma, \gamma\pi^+\pi^-\pi^0, \gamma 3\pi^0$		NODE=M049R54 NODE=M049R54
••• We do not use the following data for averages, fits, limits, etc. •••						
<21	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma\eta$		
$\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$					Γ_{66}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<3	90	²⁴ ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$		NODE=M049R31 NODE=M049R31
²⁴ Assuming $B(f_0(980) \rightarrow \pi\pi) = 1$.						
						NODE=M049R31;LINKAGE=AT

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{67}/Γ
3.8±0.9 OUR AVERAGE						
4.0±1.4±0.1		17 ± 5	²⁵ BESSON	11 CLEO	$\Upsilon(1S) \rightarrow K_S^0 K_S^0$	
3.7 ^{+0.9} _{-0.7} ±0.8			ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14	90		²⁶ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
<19.4	90		²⁶ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
²⁵ BESSON 11 reports $(4.0 \pm 1.3 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(\Upsilon(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})]$ assuming $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 3.1) \times 10^{-2}$, which we rescale to our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The result also assumes $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ and $B(f'_2(1525) \rightarrow K\bar{K}) = 4 B(f'_2(1525) \rightarrow K_S^0 K_S^0)$.						

²⁶ Assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.

NODE=M049R52
NODE=M049R52

NODE=M049R52;LINKAGE=BE

NODE=M049R52;LINKAGE=D

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{68}/Γ
10.1±0.9 OUR AVERAGE					
10.5±1.6 ^{+1.9} _{-1.8}		²⁷ BESSON	07A CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$	
10.2±0.8±0.7		ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
8.1±2.3 ^{+2.9} _{-2.7}		²⁸ ANASTASSOV	99 CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90		²⁸ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
<13	90		²⁸ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
<81	90		SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$	
²⁷ Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = B(f_2(1270) \rightarrow \pi\pi)/3$ and $B(f_2(1270) \rightarrow \pi\pi) = (0.845 \pm 0.025)_{-0.012}^+$ %.						

²⁸ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.

NODE=M049R51
NODE=M049R51

NODE=M049R51;LINKAGE=BE

NODE=M049R51;LINKAGE=C

 $\Gamma(\gamma \eta(1405))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{69}/Γ
<8.2	90	²⁹ FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$	

²⁹ Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$.

NODE=M049R23
NODE=M049R23

NODE=M049R23;LINKAGE=J

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{70}/Γ
<1.5	90	³⁰ BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.1	90	³¹ BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$	OCCUR=2	
³⁰ Using $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi\pi)/3$ and $B(f_0(1500) \rightarrow \pi\pi) = (0.349 \pm 0.023)\%$.						
³¹ Calculated by us using $B(f_0(1500) \rightarrow \eta\eta) = (5.1 \pm 0.9)\%$.						

NODE=M049R44
NODE=M049R44

NODE=M049R44;LINKAGE=BE

NODE=M049R44;LINKAGE=BS

 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{71}/Γ	
< 2.6	90	³² ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 6.3	90	³² FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$		
<19	90	³² FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$	OCCUR=2	
< 8	90	³³ ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	OCCUR=2	
<24	90	³⁴ SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$		

³² Assuming $B(f_0(1710) \rightarrow K\bar{K}) = 0.38$.

³³ Assuming $B(f_0(1710) \rightarrow \pi\pi) = 0.04$.

³⁴ Assuming $B(f_0(1710) \rightarrow \eta\eta) = 0.18$.

NODE=M049R53
NODE=M049R53

NODE=M049R53;LINKAGE=E

NODE=M049R53;LINKAGE=F

NODE=M049R53;LINKAGE=A

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{72}/Γ
<0.7	90	ATHAR	06 CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$	

NODE=M049R50
NODE=M049R50

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{73} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

NODE=M049R45
 NODE=M049R45

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{74} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$

NODE=M049R46
 NODE=M049R46

 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{75} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5.3	90	³⁵ ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M049R30
 NODE=M049R30

³⁵ Assuming $B(f_4(2050) \rightarrow \pi \pi) = 0.17$.

NODE=M049R30;LINKAGE=AT

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{76} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0002	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R63
 NODE=M049R63

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{77} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 8	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R56
 NODE=M049R56

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 160	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 150	90	FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 290	90	ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<2000	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{78} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 6	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M049R41
 NODE=M049R41

• • • We do not use the following data for averages, fits, limits, etc. • • •

<120	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
------	----	-------	---------	---

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \rho \bar{\rho}) / \Gamma_{\text{total}}$ Γ_{79} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \rho \bar{\rho}$

NODE=M049R42
 NODE=M049R42

• • • We do not use the following data for averages, fits, limits, etc. • • •

<160	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma \rho \bar{\rho}$
------	----	-------	---------	---

 $\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi \phi) / \Gamma_{\text{total}}$ Γ_{80} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M049R62
 NODE=M049R62

 $\Gamma(\gamma \eta_c(1S)) / \Gamma_{\text{total}}$ Γ_{81} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5.7	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R38
 NODE=M049R38

 $\Gamma(\gamma \chi_{c0}) / \Gamma_{\text{total}}$ Γ_{82} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6.5	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R39
 NODE=M049R39

 $\Gamma(\gamma \chi_{c1}) / \Gamma_{\text{total}}$ Γ_{83} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R40
 NODE=M049R40

 $\Gamma(\gamma \chi_{c2}) / \Gamma_{\text{total}}$ Γ_{84} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<7.6	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R48
 NODE=M049R48

 $\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi) / \Gamma_{\text{total}}$ Γ_{85} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R49
 NODE=M049R49

$\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R68
NODE=M049R68 $\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R69
NODE=M049R69 $\Gamma(\gamma X(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$

NODE=M049R74
NODE=M049R74 $\Gamma(\gamma X)/\Gamma_{\text{total}}$ Γ_{89}/Γ (X = scalar with $m < 8.0$ GeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.5	90	³⁶ DEL-AMO-SA..11J	BABR	$e^+ e^- \rightarrow \gamma + X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30	90	³⁷ BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X$
-----	----	----------------------	---------	----------------------------------

³⁶ For a noninteracting scalar X with mass $m < 8.0$ GeV.³⁷ For a noninteracting pseudoscalar X with mass < 7.2 GeV.NODE=M049R60
NODE=M049R60
NODE=M049R60NODE=M049R60;LINKAGE=DA
NODE=M049R60;LINKAGE=A $\Gamma(\gamma X \bar{X}(m_X < 3.1 \text{ GeV}))/\Gamma_{\text{total}}$ Γ_{90}/Γ (X \bar{X} = vectors with $m < 3.1$ GeV)

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	³⁸ BALEST	95 CLEO	$e^+ e^- \rightarrow \gamma + X \bar{X}$

³⁸ For a noninteracting vector X with mass < 3.1 GeV.NODE=M049R61
NODE=M049R61
NODE=M049R61

NODE=M049R61;LINKAGE=A

 $\Gamma(\gamma X \bar{X}(m_X < 4.5 \text{ GeV}))/\Gamma_{\text{total}}$ Γ_{91}/Γ X and \bar{X} = zero spin with $m < 4.5$ GeV

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	³⁹ DEL-AMO-SA..11J	BABR	$e^+ e^- \rightarrow \gamma + X \bar{X}$

³⁹ For a noninteracting scalar X with mass $m < 4.5$ GeV.NODE=M049R01
NODE=M049R01
NODE=M049R01

NODE=M049R01;LINKAGE=DA

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{92}/Γ (1.5 GeV $< m_X < 5.0$ GeV)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.78	95	ROSNER	07A CLEO	$e^+ e^- \rightarrow \gamma X$

NODE=M049R64
NODE=M049R64
NODE=M049R64 $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{93}/Γ (201 $< M(\mu^+ \mu^-) < 3565$ MeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	⁴⁰ LOVE	08 CLEO	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.7	90	⁴¹ LEES	13C BABR	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$
------	----	--------------------	----------	---

⁴⁰ For a narrow scalar or pseudoscalar a_1^0 with 201 $< M(\mu^+ \mu^-) < 3565$ MeV, excluding J/ψ . Measured 90% CL limits as a function of $M(\mu^+ \mu^-)$ range from $1-9 \times 10^{-6}$.⁴¹ For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9200 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from $0.28-9.7 \times 10^{-6}$.NODE=M049R65
NODE=M049R65
NODE=M049R65

NODE=M049R65;LINKAGE=LO

NODE=M049R65;LINKAGE=LE

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{94}/Γ (2 $m_\tau < M(\tau^+ \tau^-) < 9.2$ GeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<130	90	⁴² LEES	13R BABR	$\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 50	90	⁴³ LOVE	08 CLEO	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$
------	----	--------------------	---------	---

⁴² For a narrow scalar a_1^0 with $2m_\tau < M(a_1^0) < 9.2$ GeV, which result in a 90% CL upper limits of 0.9×10^{-5} at $M(a_1^0) = 2m_\tau$, $\approx 1.5 \times 10^{-5}$ at $M(a_1^0) = 7.5$ GeV, and 13×10^{-5} at $M(a_1^0) = 9.2$ GeV.⁴³ For a narrow scalar or pseudoscalar a_1^0 with $2m_\tau < M(a_1^0) < 7.5$ GeV, which result in a 90% CL limits ranging from 1×10^{-5} at $M(a_1^0) = 2m_\tau$ to 5×10^{-5} at $M(a_1^0) = 7.5$ GeV.NODE=M049R66
NODE=M049R66
NODE=M049R66

NODE=M049R66;LINKAGE=A

NODE=M049R66;LINKAGE=LO

$\Gamma(\gamma a_1^0 \rightarrow \gamma g g) / \Gamma_{\text{total}}$ Γ_{95} / Γ
(0.5 GeV < m < 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1 \times 10^{-2}$	90	44 LEES	13L BABR	$\Upsilon(1S) \rightarrow \gamma X$

⁴⁴ For a narrow, CP-odd pseudoscalar a_1^0 searched for in 26 hadronic decay modes with invariant mass 0.5 GeV < m_X < 9.0 GeV. Measured 90% CL limit as a function of m_X range from 10^{-6} to 10^{-2} .

NODE=M049R03

NODE=M049R03
NODE=M049R03

NODE=M049R03;LINKAGE=A

$\Gamma(\gamma a_1^0 \rightarrow \gamma s \bar{s}) / \Gamma_{\text{total}}$ Γ_{96} / Γ
(0.5 GeV < m < 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1 \times 10^{-3}$	90	45 LEES	13L BABR	$\Upsilon(1S) \rightarrow \gamma X$

⁴⁵ For a narrow, CP-odd pseudoscalar a_1^0 searched for in 14 hadronic decay modes with invariant mass 1.5 GeV < m_X < 9.0 GeV. Measured 90% CL limit as a function of m_X range from 10^{-5} to 10^{-3} .

NODE=M049R04

NODE=M049R04
NODE=M049R04

NODE=M049R04;LINKAGE=A

———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

NODE=M049230

$\Gamma(\mu^\pm \tau^\mp) / \Gamma_{\text{total}}$ Γ_{97} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	95	LOVE	08A CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M049R67
NODE=M049R67

———— OTHER DECAYS ————

NODE=M049235

$\Gamma(\text{invisible}) / \Gamma_{\text{total}}$ Γ_{98} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.0	90	AUBERT	09AX BABR	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
< 39	90	RUBIN	07 CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
< 25	90	TAJIMA	07 BELL	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M049R10
NODE=M049R10

———— $\Upsilon(1S)$ REFERENCES ————

NODE=M049

LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)	REFID=56345
LEES	13C	PR D87 031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54949
LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55167
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55451
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55395
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>		REFID=54746
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54314
BESSION	11	PR D83 037101	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=16737
DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16495
AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53211
DEL-AMO-SA...	10C	PRL 104 191801	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53354
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53545
AUBERT	09AX	PRL 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53201
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52565
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51945
BESSION	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
BESSION	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51638
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)	REFID=51629
TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)	REFID=51645
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51510
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50993
BESSION	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50183
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=49395
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)	REFID=48846
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)	REFID=48345
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=46609
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)	REFID=44146
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)	REFID=44102
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=42167
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)	REFID=41860
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)	REFID=42168

KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41361
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=41012
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)	REFID=40917
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=40919
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=40918
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding, World Scientific, Singapore					
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.)	REFID=40742
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)	REFID=40582
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22284
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22282
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22279
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)	REFID=22281
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)	REFID=22273
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)	REFID=22274
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=12488
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)	REFID=22270
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22271
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22272
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22263
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)	REFID=22264
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22259

$\chi_{b0}(1P)$

$I^G(J^{PC}) = 0^+(0^{++})$
J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore *C* = +. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore *P* = +.

$\chi_{b0}(1P)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	
9859.44 ± 0.42 ± 0.31 OUR EVALUATION	From average γ energy below, using $\Upsilon(2S)$	
mass = 10023.26 ± 0.31 MeV		→ UNCHECKED ←

γ ENERGY IN $\Upsilon(2S)$ DECAY

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
162.5 ± 0.4 OUR AVERAGE			
162.56 ± 0.19 ± 0.42	ARTUSO	05	CLEO $\Upsilon(2S) \rightarrow \gamma X$
162.0 ± 0.8 ± 1.2	EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
162.1 ± 0.5 ± 1.4	ALBRECHT	85E	ARG $\Upsilon(2S) \rightarrow \text{conv. } \gamma X$
163.8 ± 1.6 ± 2.7	NERNST	85	CBAL $\Upsilon(2S) \rightarrow \gamma X$
158.0 ± 7 ± 1	HAAS	84	CLEO $\Upsilon(2S) \rightarrow \text{conv. } \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
149.4 ± 0.7 ± 5.0	KLOPFEN...	83	CUSB $\Upsilon(2S) \rightarrow \gamma X$

$\chi_{b0}(1P)$ DECAY MODES

<u>Mode</u>	<u>Fraction (Γ_i/Γ)</u>	<u>Confidence level</u>
Γ_1 $\gamma \Upsilon(1S)$	(1.76 ± 0.35) %	DESIG=1
Γ_2 $D^0 X$	< 10.4 %	90% DESIG=2
Γ_3 $\pi^+ \pi^- K^+ K^- \pi^0$	< 1.6 × 10 ⁻⁴	90% DESIG=3
Γ_4 $2\pi^+ \pi^- K^- K_S^0$	< 5 × 10 ⁻⁵	90% DESIG=4
Γ_5 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 5 × 10 ⁻⁴	90% DESIG=5
Γ_6 $2\pi^+ 2\pi^- 2\pi^0$	< 2.1 × 10 ⁻⁴	90% DESIG=6
Γ_7 $2\pi^+ 2\pi^- K^+ K^-$	(1.1 ± 0.6) × 10 ⁻⁴	DESIG=7
Γ_8 $2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.7 × 10 ⁻⁴	90% DESIG=8
Γ_9 $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 5 × 10 ⁻⁴	90% DESIG=9

Γ_{10}	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 1.6	$\times 10^{-4}$	90%	DESIG=10
Γ_{11}	$3\pi^+ 3\pi^-$	< 8	$\times 10^{-5}$	90%	DESIG=11
Γ_{12}	$3\pi^+ 3\pi^- 2\pi^0$	< 6	$\times 10^{-4}$	90%	DESIG=12
Γ_{13}	$3\pi^+ 3\pi^- K^+ K^-$	(2.4 ± 1.2)	$\times 10^{-4}$		DESIG=13
Γ_{14}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 1.0	$\times 10^{-3}$	90%	DESIG=14
Γ_{15}	$4\pi^+ 4\pi^-$	< 8	$\times 10^{-5}$	90%	DESIG=15
Γ_{16}	$4\pi^+ 4\pi^- 2\pi^0$	< 2.1	$\times 10^{-3}$	90%	DESIG=16
Γ_{17}	$J/\psi J/\psi$	< 7	$\times 10^{-5}$	90%	DESIG=17
Γ_{18}	$J/\psi \psi(2S)$	< 1.2	$\times 10^{-4}$	90%	DESIG=18
Γ_{19}	$\psi(2S) \psi(2S)$	< 3.1	$\times 10^{-5}$	90%	DESIG=19

$\chi_{b0}(1P)$ BRANCHING RATIOS

NODE=M076220

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_1/Γ

NODE=M076R1
NODE=M076R1

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.76 \pm 0.30 \pm 0.18$		87	^{1,2} KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.6	90		³ LEES	11J	BABR $\Upsilon(2S) \rightarrow X \gamma$
< 6	90		WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
< 11	90		PAUSS	83	CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

² KORNICER 11 reports $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (6.59 \pm 0.96 \pm 0.60) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$.

NODE=M076R1;LINKAGE=KA
NODE=M076R1;LINKAGE=KR

NODE=M076R1;LINKAGE=LE

$\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_2/Γ

NODE=M076R01
NODE=M076R01

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 10.4 \times 10^{-2}$	90	^{4,5} BRIERE	08	CLEO $\Upsilon(2S) \rightarrow \gamma D^0 X$

⁴ For $p_{D^0} > 2.5$ GeV/c.

⁵ The authors also present their result as $(5.6 \pm 3.6 \pm 0.5) \times 10^{-2}$.

NODE=M076R01;LINKAGE=BR
NODE=M076R01;LINKAGE=R1

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

NODE=M076R02
NODE=M076R02

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.6	90	⁶ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

⁶ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M076R02;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ Γ_4/Γ

NODE=M076R03
NODE=M076R03

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.5	90	⁷ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

⁷ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 2 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M076R03;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

NODE=M076R04
NODE=M076R04

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 5	90	⁸ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

⁸ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 18 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M076R04;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

NODE=M076R05
NODE=M076R05

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.1	90	⁹ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

⁹ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 8 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M076R05;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.6±0.1	7	10 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$
<p>¹⁰ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (4 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M076R06
NODE=M076R06

NODE=M076R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	11 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
<p>¹¹ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 10 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R07
NODE=M076R07

NODE=M076R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	12 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
<p>¹² ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 20 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R08
NODE=M076R08

NODE=M076R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	13 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
<p>¹³ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R09
NODE=M076R09

NODE=M076R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	14 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
<p>¹⁴ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R10
NODE=M076R10

NODE=M076R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	15 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
<p>¹⁵ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 22 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R11
NODE=M076R11

NODE=M076R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±1.2±0.2	9	16 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
<p>¹⁶ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (9 \pm 4 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M076R12
NODE=M076R12

NODE=M076R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<10	90	17 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
<p>¹⁷ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 37 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R13
NODE=M076R13

NODE=M076R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	18 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<p>¹⁸ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R14
NODE=M076R14

NODE=M076R14;LINKAGE=AS

$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<21	90	19 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$ 19 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ < 77×10^{-6} which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M076R15
NODE=M076R15

NODE=M076R15;LINKAGE=AS

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	20 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$ 20 SHEN 12 reports < 7.1×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}]$ $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.

NODE=M076R16
NODE=M076R16

NODE=M076R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	21 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$ 21 SHEN 12 reports < 12×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}]$ $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.

NODE=M076R17
NODE=M076R17

NODE=M076R17;LINKAGE=SH

 $\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	22 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$ 22 SHEN 12 reports < 3.1×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}]$ $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.

NODE=M076R18
NODE=M076R18

NODE=M076R18;LINKAGE=SH

 $\chi_{b0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M076230

 $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$
 $\Gamma_1/\Gamma \times \Gamma_{47}^{\Upsilon(2S)}/\Gamma \Upsilon(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 $\times 10^{-3}$	90	23 LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$ 23 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}} < 4.6\%$ using $B(\Upsilon(4S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4)\%$.

NODE=M076B02
NODE=M076B02

NODE=M076B02;LINKAGE=LE

 $B(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	COMMENT
1.63 $\pm 0.24 \pm 0.15$	87	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M076B01
NODE=M076B01 $\chi_{b0}(1P)$ REFERENCES

NODE=M076

SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54313
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)	REFID=22290
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)	REFID=22286

$\chi_{b1}(1P)$

$I^G(J^{PC}) = 0^+(1^{++})$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 1$ from SKWARNICKI 87.

NODE=M077

NODE=M077

$\chi_{b1}(1P)$ MASS

NODE=M077M

VALUE (MeV) DOCUMENT ID
9892.78 ± 0.26 ± 0.31 OUR EVALUATION From average γ energy below, using $\Upsilon(2S)$
 mass = 10023.26 ± 0.31 MeV


NODE=M077M
 → UNCHECKED ←

γ ENERGY IN $\Upsilon(2S)$ DECAY

NODE=M077DM

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
129.63 ± 0.33 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
129.58 ± 0.09 ± 0.29	ARTUSO	05	CLEO $\Upsilon(2S) \rightarrow \gamma X$
128.8 ± 0.4 ± 0.6	EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
131.7 ± 0.9 ± 1.3	WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
131.7 ± 0.3 ± 1.1	ALBRECHT	85E	ARG $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
130.6 ± 0.8 ± 2.4	NERNST	85	CBAL $\Upsilon(2S) \rightarrow \gamma X$
129 ± 0.8 ± 1	HAAS	84	CLEO $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
128.1 ± 0.4 ± 3.0	KLOPFEN...	83	CUSB $\Upsilon(2S) \rightarrow \gamma X$
130.6 ± 3.0	PAUSS	83	CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M077DM



$\chi_{b1}(1P)$ DECAY MODES

NODE=M077215; NODE=M077

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\gamma \Upsilon(1S)$	(33.9 ± 2.2) %	DESIG=1
Γ_2 $D^0 X$	(12.6 ± 2.2) %	DESIG=2
Γ_3 $\pi^+ \pi^- K^+ K^- \pi^0$	(2.0 ± 0.6) × 10 ⁻⁴	DESIG=3
Γ_4 $2\pi^+ \pi^- K^- K_S^0$	(1.3 ± 0.5) × 10 ⁻⁴	DESIG=4
Γ_5 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 6 × 10 ⁻⁴	90% DESIG=5
Γ_6 $2\pi^+ 2\pi^- 2\pi^0$	(8.0 ± 2.5) × 10 ⁻⁴	DESIG=6
Γ_7 $2\pi^+ 2\pi^- K^+ K^-$	(1.5 ± 0.5) × 10 ⁻⁴	DESIG=7
Γ_8 $2\pi^+ 2\pi^- K^+ K^- \pi^0$	(3.5 ± 1.2) × 10 ⁻⁴	DESIG=8
Γ_9 $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(8.6 ± 3.2) × 10 ⁻⁴	DESIG=9

Γ_{10}	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$(9.3 \pm 3.3) \times 10^{-4}$		DESIG=10
Γ_{11}	$3\pi^+ 3\pi^-$	$(1.9 \pm 0.6) \times 10^{-4}$		DESIG=11
Γ_{12}	$3\pi^+ 3\pi^- 2\pi^0$	$(1.7 \pm 0.5) \times 10^{-3}$		DESIG=12
Γ_{13}	$3\pi^+ 3\pi^- K^+ K^-$	$(2.6 \pm 0.8) \times 10^{-4}$		DESIG=13
Γ_{14}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(7.5 \pm 2.6) \times 10^{-4}$		DESIG=14
Γ_{15}	$4\pi^+ 4\pi^-$	$(2.6 \pm 0.9) \times 10^{-4}$		DESIG=15
Γ_{16}	$4\pi^+ 4\pi^- 2\pi^0$	$(1.4 \pm 0.6) \times 10^{-3}$		DESIG=16
Γ_{17}	$J/\psi J/\psi$	$< 2.7 \times 10^{-5}$	90%	DESIG=17
Γ_{18}	$J/\psi \psi(2S)$	$< 1.7 \times 10^{-5}$	90%	DESIG=18
Γ_{19}	$\psi(2S)\psi(2S)$	$< 6 \times 10^{-5}$	90%	DESIG=19

$\chi_{b1}(1P)$ BRANCHING RATIOS

NODE=M077220

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_1/Γ

NODE=M077R1
NODE=M077R1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.339 ± 0.022 OUR AVERAGE				
0.331 ± 0.018 ± 0.017	3222	^{1,2} KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
0.350 ± 0.023 ± 0.018	13k	³ LEES	11J	BABR $\Upsilon(2S) \rightarrow X \gamma$
0.32 ± 0.06 ± 0.07		WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
0.47 ± 0.18		KLOPFEN...	83	CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

² KORNICER 11 reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(22.8 \pm 0.4 \pm 1.2) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$ = $(6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ LEES 11J reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (24.1 \pm 0.6 \pm 1.5) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=KA
NODE=M077R1;LINKAGE=KR

NODE=M077R1;LINKAGE=LE

$\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_2/Γ

NODE=M077R01
NODE=M077R01

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
12.6 ± 1.9 ± 1.1	2310	⁴ BRIERE	08	CLEO $\Upsilon(2S) \rightarrow \gamma D^0 X$

⁴ For $p_{D^0} > 2.5$ GeV/c.

NODE=M077R01;LINKAGE=BR

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

NODE=M077R02
NODE=M077R02

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.0 ± 0.6 ± 0.1	18	⁵ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

⁵ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (14 \pm 3 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R02;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ Γ_4/Γ

NODE=M077R03
NODE=M077R03

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.3 ± 0.5 ± 0.1	11	⁶ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

⁶ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (9 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R03;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

NODE=M077R04
NODE=M077R04

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 6	90	⁷ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

⁷ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] < 42 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

NODE=M077R04;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.0±2.4±0.4	46	⁸ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

NODE=M077R05
NODE=M077R05

⁸ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (55 \pm 9 \pm 14) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R05;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.5±0.5±0.1	18	⁹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

NODE=M077R06
NODE=M077R06

⁹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (10 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5±1.2±0.2	22	¹⁰ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

NODE=M077R07
NODE=M077R07

¹⁰ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (24 \pm 6 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6±3.2±0.4	26	¹¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

NODE=M077R08
NODE=M077R08

¹¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (59 \pm 14 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±3.3±0.5	21	¹² ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$

NODE=M077R09
NODE=M077R09

¹² ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (64 \pm 16 \pm 16) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.9±0.6±0.1	25	¹³ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$

NODE=M077R10
NODE=M077R10

¹³ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (13 \pm 3 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17±5±1	56	¹⁴ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$

NODE=M077R11
NODE=M077R11

¹⁴ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (119 \pm 18 \pm 32) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6±0.8±0.1	21	¹⁵ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$

NODE=M077R12
NODE=M077R12

¹⁵ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (18 \pm 4 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$
 $= (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R12;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.5±2.6±0.4	28	16 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
<p>¹⁶ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(52 \pm 11 \pm 14) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M077R13
NODE=M077R13

NODE=M077R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6±0.9±0.1	24	17 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<p>¹⁷ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(18 \pm 4 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M077R14
NODE=M077R14

NODE=M077R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
14±5±1	26	18 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
<p>¹⁸ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(96 \pm 24 \pm 29) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M077R15
NODE=M077R15

NODE=M077R15;LINKAGE=AS

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	19 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
<p>¹⁹ SHEN 12 reports $< 2.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$.</p>				

NODE=M077R16
NODE=M077R16

NODE=M077R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	20 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
<p>²⁰ SHEN 12 reports $< 1.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$.</p>				

NODE=M077R17
NODE=M077R17

NODE=M077R17;LINKAGE=SH

 $\Gamma(\psi(2S) \psi(2S))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	21 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
<p>²¹ SHEN 12 reports $< 6.2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow \psi(2S) \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$.</p>				

NODE=M077R18
NODE=M077R18

NODE=M077R18;LINKAGE=SH

 $\chi_{b1}(1P)$ Cross-Particle Branching Ratios

NODE=M077230

 $\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_{45}^{\Upsilon(2S)}/\Gamma \Upsilon(2S)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
24.1±0.6±1.5	13k	LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$

NODE=M077B03
NODE=M077B03 $B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$ NODE=M077B01
NODE=M077B01

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.65±0.11±0.27	3222	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

 $B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$ NODE=M077B02
NODE=M077B02

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.33±0.30±0.23	50	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

 $B(\chi_{b2}(1P) \rightarrow p X + \bar{p} X)/B(\chi_{b1}(1P) \rightarrow p X + \bar{p} X)$ NODE=M077R20
NODE=M077R20

VALUE	DOCUMENT ID	TECN	COMMENT
1.068±0.010±0.040	BRIERE	07 CLEO	$\Upsilon(2S) \rightarrow \gamma \chi_{bJ}(1P)$

$B(\chi_{b0}(1P) \rightarrow pX + \bar{p}X)/B(\chi_{b1}(1P) \rightarrow pX + \bar{p}X)$

VALUE	DOCUMENT ID	TECN	COMMENT
1.11±0.15±0.20	BRIERE	07 CLEO	$\Upsilon(2S) \rightarrow \gamma\chi_{bJ}(1P)$

NODE=M077R21
 NODE=M077R21

 $\chi_{b1}(1P)$ REFERENCES

SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.) J
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)

NODE=M077

REFID=54313
 REFID=16769
 REFID=53936
 REFID=52574
 REFID=52577
 REFID=51887
 REFID=50454
 REFID=46612
 REFID=40019
 REFID=22290
 REFID=22288
 REFID=22289
 REFID=22287
 REFID=22285
 REFID=22286

 $h_b(1P)$

$$I^G(J^{PC}) = ?^?(1^{+-})$$

Quantum numbers are quark model predictions, $C = -$ established by $\eta_b\gamma$ decay.

NODE=M204

NODE=M204

 $h_b(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9899.3±0.8 OUR AVERAGE				
[9899.3 ± 1.0 MeV OUR 2015 AVERAGE]				
9899.3±0.4±1.0	112k	TAMPONI	15 BELL	$e^+e^- \rightarrow h_b(1P)\eta$
9899.1±0.4±1.0	70k	MIZUK	12 BELL	$e^+e^- \rightarrow \pi^+\pi^-$ hadrons
9902 ±4 ±2	10.8k	LEES	11k BABR	$\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
9898.2 ^{+1.1+1.0} _{-1.0-1.1}	50.0k	¹ ADACHI	12 BELL	10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM
¹ Superseded by MIZUK 12.				

NODE=M204M

NODE=M204M
 NEW

NODE=M204M;LINKAGE=AD

 $h_b(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \eta_b(1S)\gamma$	(52 ⁺⁶ ₋₅) %

NODE=M204215;NODE=M204

DESIG=1

 $h_b(1P)$ BRANCHING RATIOS

$\Gamma(\eta_b(1S)\gamma)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
52⁺⁶₋₅ OUR AVERAGE				
[(49 ⁺⁸ ₋₇) × 10 ⁻² OUR 2015 AVERAGE]				
56 ±8 ±4	33.1k	¹ TAMPONI	15 BELL	$e^+e^- \rightarrow \eta_b(1S)\gamma\eta$
49.2±5.7 ^{+5.6} _{-3.3}	24k	MIZUK	12 BELL	$e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadrons
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
seen	10.8k	LEES	11k BABR	$\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$
¹ Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$.				

NODE=M204225

NODE=M204R01
 NODE=M204R01

NEW

NODE=M204R01;LINKAGE=A

 $h_b(1P)$ REFERENCES

TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)

NODE=M204

REFID=56996
 REFID=53962
 REFID=54718
 REFID=53937

$\chi_{b2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 2$ from SKWARNICKI 87.

NODE=M078

NODE=M078

 $\chi_{b2}(1P)$ MASS

NODE=M078M

VALUE (MeV) DOCUMENT ID
9912.21 ± 0.26 ± 0.31 OUR EVALUATION From average γ energy below, using $\Upsilon(2S)$
 mass = 10023.26 ± 0.31 MeV

NODE=M078M
 → UNCHECKED ←

 $m_{\chi_{b2}(1P)} - m_{\chi_{b1}(1P)}$

NODE=M078DM2

VALUE (MeV) DOCUMENT ID TECN COMMENT
19.81 ± 0.65 ± 0.20 ¹ AAIJ 14BG LHCB $pp \rightarrow \gamma \mu^+ \mu^- X$
¹ From the $\chi_{bj}(1P) \rightarrow \Upsilon(1S)\gamma$ transition.

NODE=M078DM2

NODE=M078DM2;LINKAGE=A

 γ ENERGY IN $\Upsilon(2S)$ DECAY

NODE=M078DM

VALUE (MeV) DOCUMENT ID TECN COMMENT
110.44 ± 0.29 OUR AVERAGE Error includes scale factor of 1.1.
 110.58 ± 0.08 ± 0.30 ARTUSO 05 CLEO $\Upsilon(2S) \rightarrow \gamma X$
 110.8 ± 0.3 ± 0.6 EDWARDS 99 CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
 107.0 ± 1.1 ± 1.3 WALK 86 CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
 110.6 ± 0.3 ± 0.9 ALBRECHT 85E ARG $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
 110.4 ± 0.8 ± 2.2 NERNST 85 CBAL $\Upsilon(2S) \rightarrow \gamma X$
 109.5 ± 0.7 ± 1.0 HAAS 84 CLEO $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
 108.2 ± 0.3 ± 2.0 KLOPFEN... 83 CUSB $\Upsilon(2S) \rightarrow \gamma X$
 108.8 ± 4.0 PAUSS 83 CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M078DM

 $\chi_{b2}(1P)$ DECAY MODES

NODE=M078215;NODE=M078

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\gamma \Upsilon(1S)$	(19.1 ± 1.2) %	
Γ_2 $D^0 X$	< 7.9 %	90%
Γ_3 $\pi^+ \pi^- K^+ K^- \pi^0$	(8 ± 5) × 10 ⁻⁵	
Γ_4 $2\pi^+ \pi^- K^- K_S^0$	< 1.0 × 10 ⁻⁴	90%
Γ_5 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	(5.3 ± 2.4) × 10 ⁻⁴	
Γ_6 $2\pi^+ 2\pi^- 2\pi^0$	(3.5 ± 1.4) × 10 ⁻⁴	
Γ_7 $2\pi^+ 2\pi^- K^+ K^-$	(1.1 ± 0.4) × 10 ⁻⁴	
Γ_8 $2\pi^+ 2\pi^- K^+ K^- \pi^0$	(2.1 ± 0.9) × 10 ⁻⁴	
Γ_9 $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(3.9 ± 1.8) × 10 ⁻⁴	
Γ_{10} $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 5 × 10 ⁻⁴	90%
Γ_{11} $3\pi^+ 3\pi^-$	(7.0 ± 3.1) × 10 ⁻⁵	
Γ_{12} $3\pi^+ 3\pi^- 2\pi^0$	(1.0 ± 0.4) × 10 ⁻³	
Γ_{13} $3\pi^+ 3\pi^- K^+ K^-$	< 8 × 10 ⁻⁵	90%
Γ_{14} $3\pi^+ 3\pi^- K^+ K^- \pi^0$	(3.6 ± 1.5) × 10 ⁻⁴	
Γ_{15} $4\pi^+ 4\pi^-$	(8 ± 4) × 10 ⁻⁵	
Γ_{16} $4\pi^+ 4\pi^- 2\pi^0$	(1.8 ± 0.7) × 10 ⁻³	
Γ_{17} $J/\psi J/\psi$	< 4 × 10 ⁻⁵	90%
Γ_{18} $J/\psi \psi(2S)$	< 5 × 10 ⁻⁵	90%
Γ_{19} $\psi(2S) \psi(2S)$	< 1.6 × 10 ⁻⁵	90%

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

DESIG=9

DESIG=10

DESIG=11

DESIG=12

DESIG=13

DESIG=14

DESIG=15

DESIG=16

DESIG=17

DESIG=18

DESIG=19

$\chi_{b2}(1P)$ BRANCHING RATIOS

NODE=M078220

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.191 ± 0.012 OUR AVERAGE				
0.186 ± 0.011 ± 0.009	1770	^{2,3} KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
0.194 ^{+0.014} _{-0.017} ± 0.009	8k	⁴ LEES	11J	BABR $\Upsilon(2S) \rightarrow X \gamma$
0.27 ± 0.06 ± 0.06		WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
0.20 ± 0.05		KLOPFEN...	83	CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M078R1
NODE=M078R1² Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.³ KORNICER 11 reports $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.33 \pm 0.04 \pm 0.07) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.NODE=M078R1;LINKAGE=KA
NODE=M078R1;LINKAGE=KR⁴ LEES 11J reports $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (13.9 \pm 0.5^{+0.9}_{-1.1}) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=LE

 $\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 7.9 × 10⁻²				
	90	^{5,6} BRIERE	08	CLEO $\Upsilon(2S) \rightarrow \gamma D^0 X$

⁵ For $p_{D^0} > 2.5$ GeV/c.⁶ The authors also present their result as $(5.4 \pm 1.9 \pm 0.5) \times 10^{-2}$.NODE=M078R01
NODE=M078R01NODE=M078R01;LINKAGE=BR
NODE=M078R01;LINKAGE=RI $\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
0.84 ± 0.50 ± 0.04				
	8	⁷ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

⁷ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.NODE=M078R02
NODE=M078R02

NODE=M078R02;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.0				
	90	⁸ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

⁸ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] < 7 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.NODE=M078R03
NODE=M078R03

NODE=M078R03;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
5.3 ± 2.4 ± 0.3				
	11	⁹ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

⁹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (38 \pm 14 \pm 10) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.NODE=M078R04
NODE=M078R04

NODE=M078R04;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ± 1.4 ± 0.2				
	19	¹⁰ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

¹⁰ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (25 \pm 8 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.NODE=M078R05
NODE=M078R05

NODE=M078R05;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
1.1 ± 0.4 ± 0.1				
	14	¹¹ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

¹¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (8 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.NODE=M078R06
NODE=M078R06

NODE=M078R06;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.9±0.1	13	12 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
<p>12 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ = $(15 \pm 5 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M078R07
NODE=M078R07

NODE=M078R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9±1.8±0.2	11	13 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
<p>13 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ = $(28 \pm 11 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M078R08
NODE=M078R08

NODE=M078R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	14 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
<p>14 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ < 36×10^{-6} which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.</p>				

NODE=M078R09
NODE=M078R09

NODE=M078R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.31±0.03	9	15 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
<p>15 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ = $(5 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M078R10
NODE=M078R10

NODE=M078R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.2±3.6±0.5	34	16 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
<p>16 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ = $(73 \pm 16 \pm 20) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M078R11
NODE=M078R11

NODE=M078R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	17 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
<p>17 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ < 6×10^{-6} which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.</p>				

NODE=M078R12
NODE=M078R12

NODE=M078R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.6±1.5±0.2	14	18 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
<p>18 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ = $(26 \pm 8 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M078R13
NODE=M078R13

NODE=M078R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.84±0.40±0.04	7	19 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<p>19 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ = $(6 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M078R14
NODE=M078R14

NODE=M078R14;LINKAGE=AS

$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$18 \pm 7 \pm 1$	29	20 ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
20 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ $= (132 \pm 31 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))$ $= (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M078R15
NODE=M078R15

NODE=M078R15;LINKAGE=AS

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	21 SHEN	12	BELL $\Upsilon(2S) \rightarrow \gamma \psi X$
21 SHEN 12 reports $< 4.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}]$ $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

NODE=M078R16
NODE=M078R16

NODE=M078R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	22 SHEN	12	BELL $\Upsilon(2S) \rightarrow \gamma \psi X$
22 SHEN 12 reports $< 4.9 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}]$ $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

NODE=M078R17
NODE=M078R17

NODE=M078R17;LINKAGE=SH

 $\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	23 SHEN	12	BELL $\Upsilon(2S) \rightarrow \gamma \psi X$
23 SHEN 12 reports $< 1.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}]$ $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

NODE=M078R18
NODE=M078R18

NODE=M078R18;LINKAGE=SH

 $\chi_{b2}(1P)$ Cross-Particle Branching Ratios

NODE=M078230

 $\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}$
 $\Gamma_1/\Gamma \times \Gamma_{46}^{\Upsilon(2S)}/\Gamma^{\Upsilon(2S)}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$13.9 \pm 0.5^{+0.9}_{-1.1}$	8k	LEES	11J	BABR $\Upsilon(2S) \rightarrow X \gamma$

NODE=M078B03
NODE=M078B03 $B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$ NODE=M078B01
NODE=M078B01

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.29 \pm 0.09 \pm 0.16$	1770	KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

 $B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$ NODE=M078B02
NODE=M078B02

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.56 \pm 0.40 \pm 0.41$	126	KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

 $\chi_{b2}(1P)$ REFERENCES

NODE=M078

AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56199
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54313
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.)	REFID=40019
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)	REFID=22290
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)	REFID=22286

$\eta_b(2S)$

$$J^{PC} = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M200

NODE=M200

NODE=M200M

NODE=M200M

 $\eta_b(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$9999.0 \pm 3.5^{+2.8}_{-1.9}$	26k	¹ MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^- +$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$9974.6 \pm 2.3 \pm 2.1$	11 ± 4	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
¹ Assuming $\Gamma_{\eta_b(2S)} = 4.9$ MeV. Not independent of the corresponding mass difference measurement.				
² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.				
³ Assuming $\Gamma_{\eta_b(2S)} = 5$ MeV. Not independent of the corresponding mass difference measurement.				

NODE=M200M;LINKAGE=MI

NODE=M200M;LINKAGE=DO

NODE=M200M;LINKAGE=NI

 $m\Upsilon(2S) - m_{\eta_b(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$24.3 \pm 3.5^{+2.8}_{-1.9}$	26k	⁴ MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^- +$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$48.7 \pm 2.3 \pm 2.1$	11 ± 4	^{5,6} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
⁴ Assuming $\Gamma_{\eta_b(2S)} = 4.9$ MeV. Not independent of the corresponding mass measurement.				
⁵ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.				
⁶ Assuming $\Gamma_{\eta_b(2S)} = 5$ MeV. Not independent of the corresponding mass measurement.				

NODE=M200DM

NODE=M200DM

NODE=M200DM;LINKAGE=MI

NODE=M200DM;LINKAGE=DO

NODE=M200DM;LINKAGE=NI

 $\eta_b(2S)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^-$ hadrons

NODE=M200W

NODE=M200W

 $\eta_b(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	seen

NODE=M200215;NODE=M200

DESIG=1

 $\eta_b(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		26k	MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^-$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
seen			⁷ DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons	
⁷ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.						

NODE=M200225

NODE=M200R01

NODE=M200R01

NODE=M200R01;LINKAGE=DO

 $\eta_b(2S)$ REFERENCES

DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M200

REFID=54288

REFID=54718

$\Upsilon(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M052

 $\Upsilon(2S)$ MASS

NODE=M052M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10023.26 ± 0.31 OUR AVERAGE			
10023.5 ± 0.5	¹ ARTAMONOV 00	MD1	$e^+e^- \rightarrow$ hadrons
10023.1 ± 0.4	BARBER 84	REDE	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10023.6 ± 0.5	^{2,3} BARU	86B REDE	$e^+e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 86B using new electron mass (COHEN 87).			
² Reanalysis of ARTAMONOV 84.			
³ Superseded by ARTAMONOV 00.			

NODE=M052M

NODE=M052M;LINKAGE=AR
 NODE=M052M;LINKAGE=C
 NODE=M052M;LINKAGE=RZ

 $m\Upsilon(3S) - m\Upsilon(2S)$

NODE=M052DM3

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50 ± 0.02 ± 0.13	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M052DM3

 $\Upsilon(2S)$ WIDTH

NODE=M052W

VALUE (keV)	DOCUMENT ID	COMMENT
31.98 ± 2.63 OUR EVALUATION		See the Note on "Width Determinations of the Υ States"

NODE=M052W
 → UNCHECKED ←

 $\Upsilon(2S)$ DECAY MODES

NODE=M052215;NODE=M052

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 $\Upsilon(1S)\pi^+\pi^-$	(17.85 ± 0.26) %		DESIG=4
Γ_2 $\Upsilon(1S)\pi^0\pi^0$	(8.6 ± 0.4) %		DESIG=5
Γ_3 $\tau^+\tau^-$	(2.00 ± 0.21) %		DESIG=3
Γ_4 $\mu^+\mu^-$	(1.93 ± 0.17) %	S=2.2	DESIG=1
Γ_5 e^+e^-	(1.91 ± 0.16) %		DESIG=2
Γ_6 $\Upsilon(1S)\pi^0$	< 4	$\times 10^{-5}$ CL=90%	DESIG=10
Γ_7 $\Upsilon(1S)\eta$	(2.9 ± 0.4)	$\times 10^{-4}$ S=2.0	DESIG=6
Γ_8 $J/\psi(1S)$ anything	< 6	$\times 10^{-3}$ CL=90%	DESIG=20
Γ_9 $J/\psi(1S)\eta_c$	< 5.4	$\times 10^{-6}$ CL=90%	DESIG=143
Γ_{10} $J/\psi(1S)\chi_{c0}$	< 3.4	$\times 10^{-6}$ CL=90%	DESIG=144
Γ_{11} $J/\psi(1S)\chi_{c1}$	< 1.2	$\times 10^{-6}$ CL=90%	DESIG=145
Γ_{12} $J/\psi(1S)\chi_{c2}$	< 2.0	$\times 10^{-6}$ CL=90%	DESIG=146
Γ_{13} $J/\psi(1S)\eta_c(2S)$	< 2.5	$\times 10^{-6}$ CL=90%	DESIG=147
Γ_{14} $J/\psi(1S)X(3940)$	< 2.0	$\times 10^{-6}$ CL=90%	DESIG=148
Γ_{15} $J/\psi(1S)X(4160)$	< 2.0	$\times 10^{-6}$ CL=90%	DESIG=149
Γ_{16} $\psi(2S)\eta_c$	< 5.1	$\times 10^{-6}$ CL=90%	DESIG=150
Γ_{17} $\psi(2S)\chi_{c0}$	< 4.7	$\times 10^{-6}$ CL=90%	DESIG=151
Γ_{18} $\psi(2S)\chi_{c1}$	< 2.5	$\times 10^{-6}$ CL=90%	DESIG=152
Γ_{19} $\psi(2S)\chi_{c2}$	< 1.9	$\times 10^{-6}$ CL=90%	DESIG=153
Γ_{20} $\psi(2S)\eta_c(2S)$	< 3.3	$\times 10^{-6}$ CL=90%	DESIG=154
Γ_{21} $\psi(2S)X(3940)$	< 3.9	$\times 10^{-6}$ CL=90%	DESIG=155
Γ_{22} $\psi(2S)X(4160)$	< 3.9	$\times 10^{-6}$ CL=90%	DESIG=156
Γ_{23} ${}^2\overline{H}$ anything	(2.78 ⁺ _{-0.26})	$\times 10^{-5}$ S=1.2	DESIG=16
Γ_{24} hadrons	(94 ± 11) %		DESIG=101
Γ_{25} ggg	(58.8 ± 1.2) %		DESIG=105
Γ_{26} γgg	(1.87 ± 0.28) %		DESIG=106
Γ_{27} $\phi K^+ K^-$	(1.6 ± 0.4)	$\times 10^{-6}$	DESIG=133

Γ ₂₈	$\omega\pi^+\pi^-$	< 2.58	$\times 10^{-6}$	CL=90%	DESIG=134
Γ ₂₉	$K^*(892)^0 K^-\pi^+ + \text{c.c.}$	(2.3 ± 0.7)	$\times 10^{-6}$		DESIG=135
Γ ₃₀	$\phi f'_2(1525)$	< 1.33	$\times 10^{-6}$	CL=90%	DESIG=136
Γ ₃₁	$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$	CL=90%	DESIG=137
Γ ₃₂	$\rho(770) a_2(1320)$	< 8.8	$\times 10^{-7}$	CL=90%	DESIG=138
Γ ₃₃	$K^*(892)^0 K_2^*(1430)^0 + \text{c.c.}$	(1.5 ± 0.6)	$\times 10^{-6}$		DESIG=139
Γ ₃₄	$K_1(1270)^\pm K^\mp$	< 3.22	$\times 10^{-6}$	CL=90%	DESIG=140
Γ ₃₅	$K_1(1400)^\pm K^\mp$	< 8.3	$\times 10^{-7}$	CL=90%	DESIG=141
Γ ₃₆	$b_1(1235)^\pm \pi^\mp$	< 4.0	$\times 10^{-7}$	CL=90%	DESIG=142
Γ ₃₇	$\rho\pi$	< 1.16	$\times 10^{-6}$	CL=90%	DESIG=126
Γ ₃₈	$\pi^+\pi^-\pi^0$	< 8.0	$\times 10^{-7}$	CL=90%	DESIG=127
Γ ₃₉	$\omega\pi^0$	< 1.63	$\times 10^{-6}$	CL=90%	DESIG=128
Γ ₄₀	$\pi^+\pi^-\pi^0\pi^0$	(1.30 ± 0.28)	$\times 10^{-5}$		DESIG=129
Γ ₄₁	$K_S^0 K^+\pi^- + \text{c.c.}$	(1.14 ± 0.33)	$\times 10^{-6}$		DESIG=130
Γ ₄₂	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	< 4.22	$\times 10^{-6}$	CL=90%	DESIG=131
Γ ₄₃	$K^*(892)^- K^+ + \text{c.c.}$	< 1.45	$\times 10^{-6}$	CL=90%	DESIG=132
Γ ₄₄	Sum of 100 exclusive modes	(2.90 ± 0.30)	$\times 10^{-3}$		DESIG=121
Radiative decays					
Γ ₄₅	$\gamma\chi_{b1}(1P)$	(6.9 ± 0.4)	%		DESIG=8
Γ ₄₆	$\gamma\chi_{b2}(1P)$	(7.15 ± 0.35)	%		DESIG=7
Γ ₄₇	$\gamma\chi_{b0}(1P)$	(3.8 ± 0.4)	%		DESIG=9
Γ ₄₈	$\gamma f_0(1710)$	< 5.9	$\times 10^{-4}$	CL=90%	DESIG=13
Γ ₄₉	$\gamma f'_2(1525)$	< 5.3	$\times 10^{-4}$	CL=90%	DESIG=12
Γ ₅₀	$\gamma f_2(1270)$	< 2.41	$\times 10^{-4}$	CL=90%	DESIG=11
Γ ₅₁	$\gamma f_J(2220)$				DESIG=14
Γ ₅₂	$\gamma\eta_c(1S)$	< 2.7	$\times 10^{-5}$	CL=90%	DESIG=111
Γ ₅₃	$\gamma\chi_{c0}$	< 1.0	$\times 10^{-4}$	CL=90%	DESIG=112
Γ ₅₄	$\gamma\chi_{c1}$	< 3.6	$\times 10^{-6}$	CL=90%	DESIG=113
Γ ₅₅	$\gamma\chi_{c2}$	< 1.5	$\times 10^{-5}$	CL=90%	DESIG=114
Γ ₅₆	$\gamma X(3872) \rightarrow \pi^+\pi^- J/\psi$	< 8	$\times 10^{-7}$	CL=90%	DESIG=115
Γ ₅₇	$\gamma X(3872) \rightarrow \pi^+\pi^-\pi^0 J/\psi$	< 2.4	$\times 10^{-6}$	CL=90%	DESIG=116
Γ ₅₈	$\gamma X(3915) \rightarrow \omega J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%	DESIG=117
Γ ₅₉	$\gamma X(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%	DESIG=118
Γ ₆₀	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%	DESIG=119
Γ ₆₁	$\gamma\eta_b(1S)$	(3.9 ± 1.5)	$\times 10^{-4}$		DESIG=102
Γ ₆₂	$\gamma\eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	< 3.7	$\times 10^{-6}$	CL=90%	DESIG=124
Γ ₆₃	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	< 4.9	$\times 10^{-6}$	CL=90%	DESIG=125
Γ ₆₄	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95	$\times 10^{-4}$	CL=95%	DESIG=103
Γ ₆₅	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$	CL=90%	DESIG=108
Γ ₆₆	$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	< 8.3	$\times 10^{-6}$	CL=90%	DESIG=123
Lepton Family number (LF) violating modes					
Γ ₆₇	$e^\pm\tau^\mp$	LF	< 3.2	$\times 10^{-6}$	CL=90% DESIG=107
Γ ₆₈	$\mu^\pm\tau^\mp$	LF	< 3.3	$\times 10^{-6}$	CL=90% DESIG=104

[a] 1.5 GeV < m_X < 5.0 GeV

LINKAGE=C52

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 11.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_7 \begin{array}{|c} \hline 2 \\ \hline x_1 \end{array}$$

$\Upsilon(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M052218

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
6.5±1.5±1.0	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$	

NODE=M052G1
NODE=M052G1

$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
105.4±1.0±4.2	11.8K	¹ AUBERT	08BP	BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$

NODE=M052G03
NODE=M052G03

¹Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

NODE=M052G03;LINKAGE=AU

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{24}\Gamma_5/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
0.577±0.009 OUR AVERAGE					

NODE=M052G2
NODE=M052G2

0.581±0.004±0.009	¹ ROSNER	06	CLEO	10.0	$e^+e^- \rightarrow \text{hadrons}$
0.552±0.031±0.017	¹ BARU	96	MD1		$e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.04 ±0.02	¹ JAKUBOWSKI	88	CBAL		$e^+e^- \rightarrow \text{hadrons}$
0.58 ±0.03 ±0.04	² GILES	84B	CLEO		$e^+e^- \rightarrow \text{hadrons}$
0.60 ±0.12 ±0.07	² ALBRECHT	82	DASP		$e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.07 ^{+0.09} -0.05	² NICZYPORUK	81C	LENA		$e^+e^- \rightarrow \text{hadrons}$
0.41 ±0.18	² BOCK	80	CNTR		$e^+e^- \rightarrow \text{hadrons}$

¹Radiative corrections evaluated following KURAEV 85.

²Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

NODE=M052G2;LINKAGE=P
NODE=M052G2;LINKAGE=R $\Upsilon(2S)$ PARTIAL WIDTHS

NODE=M052220

$\Gamma(e^+e^-)$					Γ_5
VALUE (keV)	DOCUMENT ID				
0.612±0.011 OUR EVALUATION					

NODE=M052W2
NODE=M052W2
→ UNCHECKED ← $\Upsilon(2S)$ BRANCHING RATIOS

NODE=M052225

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_1/Γ
--	--	--	--	--	-------------------

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
17.85±0.26 OUR FIT					
17.92±0.26 OUR AVERAGE					
16.8 ±1.1 ±1.3	906k	¹ LEES	11C	BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
17.80±0.05±0.37	170k	² LEES	11L	BABR	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02±0.02±0.61	851k	³ BHARI	09	CLEO	$e^+e^- \rightarrow \pi^+\pi^-MM$
17.22±0.17±0.75	11.8K	⁴ AUBERT	08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
19.2 ±0.2 ±1.0	52.6k	⁵ ALEXANDER	98	CLE2	$\pi^+\pi^-\ell^+\ell^-, \pi^+\pi^-MM$
18.1 ±0.5 ±1.0	11.6k	ALBRECHT	87	ARG	$e^+e^- \rightarrow \pi^+\pi^-MM$
16.9 ±4.0		GELPHMAN	85	CBAL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
19.1 ±1.2 ±0.6		BESSON	84	CLEO	$\pi^+\pi^-MM$
18.9 ±2.6		FONSECA	84	CUSB	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$
21 ±7	7	NICZYPORUK	81B	LENA	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$

NODE=M052R4
NODE=M052R4
NODE=M052R4

¹LEES 11c reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³A weighted average of the inclusive and exclusive results.

⁴Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

⁵Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

NODE=M052R4;LINKAGE=ES

NODE=M052R4;LINKAGE=LE
NODE=M052R4;LINKAGE=BH
NODE=M052R4;LINKAGE=AU

NODE=M052R4;LINKAGE=T

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{total}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 ± 0.4 OUR AVERAGE				
8.43 ± 0.16 ± 0.42	38k	¹ BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.2 ± 0.6 ± 0.8	275	² ALEXANDER	98	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.5 ± 1.9 ± 1.9	25	ALBRECHT	87	ARG $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 ± 1.5		GELPHMAN	85	CBAL $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 ± 2.3		FONSECA	84	CUSB $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

Γ_2/Γ

NODE=M052R5
NODE=M052R5

¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

NODE=M052R5;LINKAGE=BH
NODE=M052R5;LINKAGE=T

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

Γ_2/Γ_1

NODE=M052R21
NODE=M052R21

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.462 ± 0.037	¹ BHARI	09	CLEO $e^+e^- \rightarrow \Upsilon(2S)$
---------------	--------------------	----	--

¹ Not independent of other values reported by BHARI 09.

NODE=M052R21;LINKAGE=BH

$\Gamma(\tau^+\tau^-)/\Gamma_{total}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

Γ_3/Γ

NODE=M052R3
NODE=M052R3

2.00 ± 0.21 OUR AVERAGE

2.00 ± 0.12 ± 0.18	22k	¹ BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$
1.7 ± 1.5 ± 0.6		HAAS	84B	CLEO $e^+e^- \rightarrow \tau^+\tau^-$

¹ BESSON 07 reports $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{total}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M052R3;LINKAGE=BE

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

Γ_4/Γ

NODE=M052R1
NODE=M052R1

0.0193 ± 0.0017 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

0.0203 ± 0.0003 ± 0.0008	120k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0122 ± 0.0028 ± 0.0019		¹ KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$
0.0138 ± 0.0025 ± 0.0015		KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
0.009 ± 0.006 ± 0.006		² ALBRECHT	85	ARG $e^+e^- \rightarrow \mu^+\mu^-$
0.018 ± 0.008 ± 0.005		HAAS	84B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$


• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.038	90	NICZYPORUK	81C	LENA $e^+e^- \rightarrow \mu^+\mu^-$
---------	----	------------	-----	--------------------------------------

¹ Taking into account interference between the resonance and continuum.

² Re-evaluated using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$.

NODE=M052R1;LINKAGE=A
NODE=M052R1;LINKAGE=R



$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_3/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.04±0.04±0.05	22k	BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(2S)$

NODE=M052R17
 NODE=M052R17

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R10
 NODE=M052R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	90	¹ TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$
< 18	90	² HE	08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<110	90	ALEXANDER	98	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<800	90	LURZ	87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

¹ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$.

NODE=M052R10;LINKAGE=TA

² Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

NODE=M052R10;LINKAGE=HE

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_6/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	TAMPONI	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

NODE=M052R09
 NODE=M052R09

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M052R6
 NODE=M052R6

2.9 ±0.4 OUR FIT Error includes scale factor of 2.0.

2.9 ±0.4 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.

2.39±0.31±0.14	112	¹ LEES	11L	BABR	$\Upsilon(2S) \rightarrow \ell^+\ell^-\eta$
2.1 $\begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix}$ ±0.3	14	² HE	08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$

• • • We use the following data for averages but not for fits. • • •

3.55±0.32±0.05	241	³ TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\eta$
----------------	-----	----------------------	----	------	---------------------------------------

NOTFITTED

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90	^{1,4} AUBERT	08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
< 28	90	ALEXANDER	98	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\eta$
< 50	90	ALBRECHT	87	ARG	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-MM$
< 70	90	LURZ	87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-MM$
< 20	90	FONSECA	84	CUSB	$e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, \pi^+\pi^-\pi^0)$

¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

NODE=M052R6;LINKAGE=AU

² Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.


NODE=M052R6;LINKAGE=HE

³ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (17.85 \pm 0.26) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M052R6;LINKAGE=TA

⁴ Using $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

NODE=M052R6;LINKAGE=UB



$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

Γ_7/Γ_1

NODE=M052R22
NODE=M052R22

VALUE (units 10^{-3}) CL% EVTS DOCUMENT ID TECN COMMENT

1.64±0.25 OUR FIT Error includes scale factor of 2.0.

1.99±0.14±0.11 241 TAMPONI 13 BELL $e^+e^- \rightarrow \Upsilon(1S)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35±0.17±0.08 ¹ LEES 11L BABR $\Upsilon(2S) \rightarrow (\pi^+\pi^-(\gamma\gamma))\mu^+\mu^-$

< 5.2 90 ² AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

¹ Not independent of other values reported by LEES 11L.

² Not independent of other values reported by AUBERT 08BP.

NODE=M052R22;LINKAGE=LE
NODE=M052R22;LINKAGE=AU

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$

Γ_6/Γ_7

NODE=M052R23
NODE=M052R23

VALUE CL% DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13 90 TAMPONI 13 BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

Γ_8/Γ

NODE=M052R16
NODE=M052R16

VALUE CL% DOCUMENT ID TECN COMMENT

<0.006 90 MASCHMANN 90 CBAL $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$

Γ_9/Γ

NODE=M052R53
NODE=M052R53

VALUE CL% DOCUMENT ID TECN COMMENT

<5.4 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$

Γ_{10}/Γ

NODE=M052R54
NODE=M052R54

VALUE CL% DOCUMENT ID TECN COMMENT

<3.4 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$

Γ_{11}/Γ

NODE=M052R55
NODE=M052R55

VALUE CL% DOCUMENT ID TECN COMMENT

<1.2 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$

Γ_{12}/Γ

NODE=M052R56
NODE=M052R56

VALUE CL% DOCUMENT ID TECN COMMENT

<2.0 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$

Γ_{13}/Γ

NODE=M052R57
NODE=M052R57

VALUE CL% DOCUMENT ID TECN COMMENT

<2.5 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$

Γ_{14}/Γ

NODE=M052R58
NODE=M052R58

VALUE CL% DOCUMENT ID TECN COMMENT

<2.0 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$					Γ_{15}/Γ	NODE=M052R59 NODE=M052R59
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.0 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$					Γ_{16}/Γ	NODE=M052R60 NODE=M052R60
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<5.1 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$					Γ_{17}/Γ	NODE=M052R61 NODE=M052R61
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<4.7 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$					Γ_{18}/Γ	NODE=M052R62 NODE=M052R62
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.5 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$					Γ_{19}/Γ	NODE=M052R63 NODE=M052R63
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<1.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$					Γ_{20}/Γ	NODE=M052R64 NODE=M052R64
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<3.3 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$					Γ_{21}/Γ	NODE=M052R65 NODE=M052R65
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S)X$	
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$					Γ_{22}/Γ	NODE=M052R66 NODE=M052R66
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S)X$	
$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$					Γ_{23}/Γ	NODE=M052R18 NODE=M052R18
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.78^{+0.30}_{-0.26}$ OUR AVERAGE				Error includes scale factor of 1.2. $[(3.4 \pm 0.6) \times 10^{-5}$ OUR 2015 AVERAGE]		NEW
$2.64 \pm 0.11^{+0.26}_{-0.21}$		LEES	14G	BABR	$e^+ e^- \rightarrow \overline{2H} X$	
$3.37 \pm 0.50 \pm 0.25$	58	ASNER	07	CLEO	$e^+ e^- \rightarrow \overline{2H} X$	
$\Gamma(ggg)/\Gamma_{\text{total}}$					Γ_{25}/Γ	NODE=M052R01 NODE=M052R01
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
58.8 ± 1.2	6M	¹ BESSON	06A	CLEO	$\Upsilon(2S) \rightarrow \text{hadrons}$	
¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+ \pi^- \Upsilon(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0 \pi^0 \Upsilon(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.						NODE=M052R01;LINKAGE=BE
$\Gamma(\gamma gg)/\Gamma(ggg)$					Γ_{26}/Γ_{25}	NODE=M052R03 NODE=M052R03
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$3.18 \pm 0.04 \pm 0.47$	6M	BESSON	06A	CLEO	$\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$	
$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$					Γ_{27}/Γ	NODE=M052R43 NODE=M052R43
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$1.58 \pm 0.33 \pm 0.18$	58	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$	
$\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{28}/Γ	NODE=M052R44 NODE=M052R44
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<2.58	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{29}/Γ	NODE=M052R45 NODE=M052R45
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.32 \pm 0.40 \pm 0.54$	135	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	

$\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$	Γ_{30}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R46 NODE=M052R46
<1.33 90	SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(K^+ K^-)$	
$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$	Γ_{31}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R47 NODE=M052R47
<0.57 90	SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$	Γ_{32}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R48 NODE=M052R48
<0.88 90	SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{33}/Γ	
<u>VALUE (units 10^{-6})</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R49 NODE=M052R49
1.53±0.52±0.19 32	SHEN 12A BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$	Γ_{34}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R50 NODE=M052R50
<3.22 90	SHEN 12A BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$	Γ_{35}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R51 NODE=M052R51
<0.83 90	SHEN 12A BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$	Γ_{36}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R52 NODE=M052R52
<0.40 90	SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	Γ_{37}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R27 NODE=M052R27
<1.16 90	SHEN 13 BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$	
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$	Γ_{38}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R28 NODE=M052R28
<0.80 90	SHEN 13 BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$	
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$	Γ_{39}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R29 NODE=M052R29
<1.63 90	SHEN 13 BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	
$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$	Γ_{40}/Γ	
<u>VALUE (units 10^{-6})</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R30 NODE=M052R30
13.0±1.9±2.1 261 ± 37	SHEN 13 BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	
$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{41}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R40 NODE=M052R40
1.14±0.30±0.13 40 ± 10	SHEN 13 BELL $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •		
<3.2 90	¹ DOBBS 12A $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$	
¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.		NODE=M052R40;LINKAGE=DO
$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{42}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R41 NODE=M052R41
<4.22 90	SHEN 13 BELL $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$	
$\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{43}/Γ	
<u>VALUE (units 10^{-6})</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	NODE=M052R42 NODE=M052R42
<1.45 90	SHEN 13 BELL $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$	

$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$ Γ_{44}/Γ VALUE (units 10^{-2})

DOCUMENT ID

COMMENT

0.29 ± 0.031,2 DOBBS 12A $\Upsilon(2S) \rightarrow \text{hadrons}$ NODE=M052R08
NODE=M052R08¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

NODE=M052R08;LINKAGE=DO

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M052R08;LINKAGE=NC

 $\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

0.069 ± 0.004 OUR AVERAGE

0.0693 ± 0.0012 ± 0.0041 407k

ARTUSO 05 CLEO $e^+e^- \rightarrow \gamma X$

0.069 ± 0.005 ± 0.009

EDWARDS 99 CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$

0.091 ± 0.018 ± 0.022

ALBRECHT 85E ARG $e^+e^- \rightarrow \gamma\text{conv. } X$

0.065 ± 0.007 ± 0.012

NERNST 85 CBAL $e^+e^- \rightarrow \gamma X$

0.080 ± 0.017 ± 0.016

HAAS 84 CLEO $e^+e^- \rightarrow \gamma\text{conv. } X$

0.059 ± 0.014

KLOPFEN... 83 CUSB $e^+e^- \rightarrow \gamma X$ NODE=M052R8
NODE=M052R8 $\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

0.0715 ± 0.0035 OUR AVERAGE

0.0724 ± 0.0011 ± 0.0040 410k

ARTUSO 05 CLEO $e^+e^- \rightarrow \gamma X$

0.074 ± 0.005 ± 0.008

EDWARDS 99 CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$

0.098 ± 0.021 ± 0.024

ALBRECHT 85E ARG $e^+e^- \rightarrow \gamma\text{conv. } X$

0.058 ± 0.007 ± 0.010

NERNST 85 CBAL $e^+e^- \rightarrow \gamma X$

0.102 ± 0.018 ± 0.021

HAAS 84 CLEO $e^+e^- \rightarrow \gamma\text{conv. } X$

0.061 ± 0.014

KLOPFEN... 83 CUSB $e^+e^- \rightarrow \gamma X$ NODE=M052R7
NODE=M052R7 $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

0.038 ± 0.004 OUR AVERAGE

0.0375 ± 0.0012 ± 0.0047 198k

ARTUSO 05 CLEO $e^+e^- \rightarrow \gamma X$

0.034 ± 0.005 ± 0.006

EDWARDS 99 CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$

0.064 ± 0.014 ± 0.016

ALBRECHT 85E ARG $e^+e^- \rightarrow \gamma\text{conv. } X$

0.036 ± 0.008 ± 0.009

NERNST 85 CBAL $e^+e^- \rightarrow \gamma X$

0.044 ± 0.023 ± 0.009

HAAS 84 CLEO $e^+e^- \rightarrow \gamma\text{conv. } X$

0.035 ± 0.014

KLOPFEN... 83 CUSB $e^+e^- \rightarrow \gamma X$ NODE=M052R9
NODE=M052R9 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$ Γ_{48}/Γ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

<59

90

¹ ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.9

90

² ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$ ¹ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.² Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+\pi^-$.NODE=M052R13
NODE=M052R13

OCCUR=2

NODE=M052R13;LINKAGE=M
NODE=M052R13;LINKAGE=N $\Gamma(\gamma f_2'(1525))/\Gamma_{\text{total}}$ Γ_{49}/Γ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

<53

90

¹ ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$ ¹ Re-evaluated assuming $B(f_2'(1525) \rightarrow K\bar{K}) = 0.71$.NODE=M052R12
NODE=M052R12

NODE=M052R12;LINKAGE=L

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{50}/Γ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

<24.1

90

¹ ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-$ ¹ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.NODE=M052R11
NODE=M052R11

NODE=M052R11;LINKAGE=K

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{51}/Γ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

<6.8

90

¹ ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$ ¹ Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.NODE=M052R14
NODE=M052R14

NODE=M052R14;LINKAGE=S

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$					Γ_{52}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R31 NODE=M052R31
$<2.7 \times 10^{-5}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$					Γ_{53}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R32 NODE=M052R32
$<1.0 \times 10^{-4}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$					Γ_{54}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R33 NODE=M052R33
$<3.6 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$					Γ_{55}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R34 NODE=M052R34
$<1.5 \times 10^{-5}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$					Γ_{56}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R35 NODE=M052R35
$<0.8 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$					Γ_{57}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R36 NODE=M052R36
$<2.4 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$					Γ_{58}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R37 NODE=M052R37
$<2.8 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$					Γ_{59}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R38 NODE=M052R38
$<1.2 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$					Γ_{60}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R39 NODE=M052R39
$<1.3 \times 10^{-6}$	90	WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$		
$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$					Γ_{61}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M052R15 NODE=M052R15
$3.9 \pm 1.1_{-0.9}^{+1.1}$		$13 \pm 5k$	¹ AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<21	90		LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$	
< 8.4	90		¹ BONVICINI	10 CLEO	$\Upsilon(2S) \rightarrow \gamma X$	
< 5.1	90		² ARTUSO	05 CLEO	$e^+e^- \rightarrow \gamma X$	
¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.						
² Superseded by BONVICINI 10.						
$\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$					Γ_{62}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R25 NODE=M052R25
$<3.7 \times 10^{-6}$	90	SANDILYA	13 BELL	$\Upsilon(2S) \rightarrow \gamma$ hadrons		
$\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$					Γ_{63}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M052R26 NODE=M052R26
< 4.9	90		SANDILYA	13 BELL	$\Upsilon(2S) \rightarrow \gamma$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$46.2_{-14.2}^{+29.7} \pm 10.6$		10	¹ DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons	
¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.						
$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ (1.5 GeV $< m_X < 5.0$ GeV)					Γ_{64}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M052R19 NODE=M052R19 NODE=M052R19
<1.95	95	ROSNER	07A CLEO	$e^+e^- \rightarrow \gamma X$		

$$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$$

(0.3 GeV < m_{A^0} < 7 GeV)

 Γ_{65}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-5}$	90	¹ LEES	11H BABR	$\Upsilon(2S) \rightarrow \gamma \text{hadrons}$

NODE=M052R06

NODE=M052R06
NODE=M052R06

¹ For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

NODE=M052R06;LINKAGE=LE

$$\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$$
 Γ_{66}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 8.3	90	¹ AUBERT	09Z BABR	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

NODE=M052R24
NODE=M052R24

¹ For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from 0.26–8.3 $\times 10^{-6}$.

NODE=M052R24;LINKAGE=AU

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

$$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$$
 Γ_{67}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	LEES	10B BABR	$e^+ e^- \rightarrow e^\pm \tau^\mp$

NODE=M052230

NODE=M052R04
NODE=M052R04
$$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$$
 Γ_{68}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.3	90	LEES	10B BABR	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M052R20
NODE=M052R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 14.4	95	LOVE	08A CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$
----------	----	------	----------	--

$\Upsilon(2S)$ Cross-Particle Branching Ratios

NODE=M052240

$$B(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.78 \pm 0.02 \pm 0.11$	906k	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

NODE=M052R05
NODE=M052R05

$\Upsilon(2S)$ REFERENCES

NODE=M052

LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)	REFID=56345
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)	REFID=55590
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55395
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=54919
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>		REFID=54288
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>		REFID=54746
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54314
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53877
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53938
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=53939
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53355
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)	REFID=52662
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52587
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731

KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding, World Scientific, Singapore					
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC	REFID=40742
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)	REFID=40021
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22338
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22334
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
GELPHMAN	85	PR D32 2893	D. Gelpman <i>et al.</i>	(Crystal Ball Collab.)	REFID=22336
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	(DESY, ARGUS Collab.+)	REFID=22327
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22279
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)	REFID=22329
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22332
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)	REFID=22270
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22319
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22318
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)	REFID=22264

NODE=M177

 $\Upsilon(1D)$

$$J^G(J^{PC}) = 0^-(2^{--})$$

First observed by BONVICINI 04 in the decay to $\gamma\gamma \Upsilon(1S)$ and confirmed by DEL-AMO-SANCHEZ 10R in the decay to $\pi^+\pi^- \Upsilon(1S)$.

Data consistent with $J^P = 2^-$. The states with $J = 1$ and 3 also possibly seen, but need confirmation.

NODE=M177

 $\Upsilon(1D)$ MASS

NODE=M177M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
10163.7 ± 1.4 OUR AVERAGE		Error includes scale factor of 1.7.		
10164.5 ± 0.8 ± 0.5		DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$
10161.1 ± 0.6 ± 1.6	38	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

NODE=M177M

 $\Upsilon(1D)$ DECAY MODES

NODE=M177215;NODE=M177

Mode	Fraction (Γ_i/Γ)
Γ_1 $\gamma\gamma \Upsilon(1S)$	seen
Γ_2 $\gamma\chi_{bJ}(1P)$	seen
Γ_3 $\eta \Upsilon(1S)$	not seen
Γ_4 $\pi^+\pi^- \Upsilon(1S)$	$(6.6 \pm 1.6) \times 10^{-3}$

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4

 $\Upsilon(1D)$ BRANCHING RATIOS

NODE=M177225

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
<0.25	90	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	

NODE=M177R01
NODE=M177R01

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
$0.66^{+0.15}_{-0.14} \pm 0.06$	¹ DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$	

NODE=M177R03
NODE=M177R03

¹ Using theoretical predictions for $B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(1D))$.

NODE=M177R03;LINKAGE=DE

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
<1.2	90	² BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	

NODE=M177R02
NODE=M177R02

² Assuming $J = 2$.

NODE=M177R02;LINKAGE=BO

 $\Upsilon(1D)$ REFERENCES

NODE=M177

DEL-AMO-SA... 10R	PR D82 111102	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53634
BONVICINI 04	PR D70 032001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=49759

$\chi_{b0}(2P)$

$I^G(J^{PC}) = 0^+(0^{++})$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

NODE=M079

NODE=M079

$\chi_{b0}(2P)$ MASS

VALUE (MeV)	DOCUMENT ID
10232.5 ± 0.4 ± 0.5 OUR EVALUATION	From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

NODE=M079M

NODE=M079M
 → UNCHECKED ←

γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
121.9 ± 0.4 OUR EVALUATION		Treating systematic errors as correlated		
122.2 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
121.55 ± 0.16 ± 0.46		ARTUSO 05	CLEO	$\Upsilon(3S) \rightarrow \gamma X$
123.0 ± 0.8	4959	¹ HEINTZ 92	CSB2	$e^+ e^- \rightarrow \gamma X$
124.6 ± 1.4	17	² HEINTZ 92	CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
122.3 ± 0.3 ± 0.6	9903	MORRISON 91	CLE2	$e^+ e^- \rightarrow \gamma X$

NODE=M079DM


NODE=M079DM
 → UNCHECKED ←

OCCUR=2

¹ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.
² A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M079DM;LINKAGE=A

NODE=M079DM;LINKAGE=B



$\chi_{b0}(2P)$ DECAY MODES

NODE=M079215;NODE=M079

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \quad \gamma \Upsilon(2S)$	(4.6 ± 2.1) %	DESIG=2
$\Gamma_2 \quad \gamma \Upsilon(1S)$	(9 ± 6) × 10 ⁻³	DESIG=1
$\Gamma_3 \quad D^0 X$	< 8.2 %	90% DESIG=3
$\Gamma_4 \quad \pi^+ \pi^- K^+ K^- \pi^0$	< 3.4 × 10 ⁻⁵	90% DESIG=4
$\Gamma_5 \quad 2\pi^+ \pi^- K^- K_S^0$	< 5 × 10 ⁻⁵	90% DESIG=5
$\Gamma_6 \quad 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 2.2 × 10 ⁻⁴	90% DESIG=6
$\Gamma_7 \quad 2\pi^+ 2\pi^- 2\pi^0$	< 2.4 × 10 ⁻⁴	90% DESIG=7
$\Gamma_8 \quad 2\pi^+ 2\pi^- K^+ K^-$	< 1.5 × 10 ⁻⁴	90% DESIG=8

Γ_9	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.2	$\times 10^{-4}$	90%	DESIG=9
Γ_{10}	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 1.1	$\times 10^{-3}$	90%	DESIG=10
Γ_{11}	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=11
Γ_{12}	$3\pi^+ 3\pi^-$	< 7	$\times 10^{-5}$	90%	DESIG=12
Γ_{13}	$3\pi^+ 3\pi^- 2\pi^0$	< 1.2	$\times 10^{-3}$	90%	DESIG=13
Γ_{14}	$3\pi^+ 3\pi^- K^+ K^-$	< 1.5	$\times 10^{-4}$	90%	DESIG=14
Γ_{15}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=15
Γ_{16}	$4\pi^+ 4\pi^-$	< 1.7	$\times 10^{-4}$	90%	DESIG=16
Γ_{17}	$4\pi^+ 4\pi^- 2\pi^0$	< 6	$\times 10^{-4}$	90%	DESIG=17

$\chi_{b0}(2P)$ BRANCHING RATIOS

NODE=M079220

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$

 Γ_1/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.046 ± 0.020 ± 0.007 ³ HEINTZ 92 CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.028 90 ⁴ LEES 11J BABR $\Upsilon(3S) \rightarrow X \gamma$

< 0.089 90 ⁵ CRAWFORD 92B CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

³ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (6.0 \pm 0.4 \pm 0.6)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

⁴ LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (-0.3 \pm 0.2_{-0.4}^{+0.5})\%$.

⁵ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) < 1.19 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P) \gamma) = 0.049$.

NODE=M079R2
NODE=M079R2

NODE=M079R2;LINKAGE=C

NODE=M079R2;LINKAGE=LE

NODE=M079R2;LINKAGE=B

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$

 Γ_2/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.009 ± 0.006 ± 0.001 ⁶ HEINTZ 92 CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.012 90 ⁷ LEES 11J BABR $\Upsilon(3S) \rightarrow X \gamma$

< 0.025 90 ⁸ CRAWFORD 92B CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

⁶ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (6.0 \pm 0.4 \pm 0.6)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

⁷ LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (3.9 \pm 2.2_{-0.6}^{+1.2}) \times 10^{-4}$.

⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) < 0.63 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P) \gamma) = 0.049$.

NODE=M079R1
NODE=M079R1

NODE=M079R1;LINKAGE=C

NODE=M079R1;LINKAGE=LE

NODE=M079R1;LINKAGE=B

$\Gamma(D^0 X)/\Gamma_{\text{total}}$

 Γ_3/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 8.2 × 10⁻² 90 ^{9,10} BRIERE 08 CLEO $\Upsilon(3S) \rightarrow \gamma D^0 X$

⁹ For $p_{D^0} > 2.5$ GeV/c.

¹⁰ The authors also present their result as $(4.1 \pm 3.0 \pm 0.4) \times 10^{-2}$.

NODE=M079R01
NODE=M079R01NODE=M079R01;LINKAGE=BR
NODE=M079R01;LINKAGE=RI

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

 Γ_4/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

< 0.34 90 ¹¹ ASNER 08A CLEO $\Upsilon(3S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

¹¹ ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 2 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R02
NODE=M079R02

NODE=M079R02;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$

 Γ_5/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

< 0.5 90 ¹² ASNER 08A CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

¹² ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R03
NODE=M079R03

NODE=M079R03;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	13 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$ 13 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 13×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R04
NODE=M079R04

NODE=M079R04;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.4	90	14 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$ 14 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 14×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R05
NODE=M079R05

NODE=M079R05;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	15 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$ 15 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 9×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R06
NODE=M079R06

NODE=M079R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	16 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$ 16 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 13×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R07
NODE=M079R07

NODE=M079R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	17 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$ 17 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 63×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R08
NODE=M079R08

NODE=M079R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	18 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$ 18 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 39×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R09
NODE=M079R09

NODE=M079R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	19 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$ 19 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 4×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R10
NODE=M079R10

NODE=M079R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	20 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$ 20 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 72×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R11
NODE=M079R11

NODE=M079R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	21 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$ 21 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 9×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R12
NODE=M079R12

NODE=M079R12;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{total}$ Γ_{15}/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	22 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
22 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 43×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R13
NODE=M079R13

NODE=M079R13;LINKAGE=AS

$\Gamma(4\pi^+ 4\pi^-)/\Gamma_{total}$ Γ_{16}/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	23 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
23 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{total}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 10×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R14
NODE=M079R14

NODE=M079R14;LINKAGE=AS

$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{total}$ Γ_{17}/Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	24 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
24 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{total}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 38×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R15
NODE=M079R15

NODE=M079R15;LINKAGE=AS

$\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{total}$
 $\Gamma_2/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	25 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
25 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{total} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) < 1.2\%$ using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$.				

NODE=M079B01
NODE=M079B01

NODE=M079B01;LINKAGE=LE

$\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{total}$
 $\Gamma_1/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$

VALUE (units 10 ⁻³)	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	26 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
26 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{total} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{total} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) < 2.8\%$ using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$.				

NODE=M079B02
NODE=M079B02

NODE=M079B02;LINKAGE=LE

$\chi_{b0}(2P)$ REFERENCES

LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)

NODE=M079

REFID=53936
REFID=52574
REFID=52577
REFID=50454
REFID=43177
REFID=43604
REFID=41580
REFID=41634
REFID=41586

$\chi_{b1}(2P)$

$I^G(J^{PC}) = 0^+(1^{++})$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

NODE=M080

NODE=M080

$\chi_{b1}(2P)$ MASS

NODE=M080M

VALUE (MeV) DOCUMENT ID
10255.46 ± 0.22 ± 0.50 OUR EVALUATION From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

NODE=M080M
 → UNCHECKED ←

$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

NODE=M080M2

VALUE (MeV) DOCUMENT ID TECN COMMENT
23.5 ± 0.7 ± 0.7 ¹ HEINTZ 92 CSB2 $e^+e^- \rightarrow \gamma X, \ell^+\ell^-\gamma\gamma$

NODE=M080M2

¹From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

NODE=M080M2;LINKAGE=A

γ ENERGY IN $\Upsilon(3S)$ DECAY

NODE=M080DM

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
99.26 ± 0.22 OUR EVALUATION Treating systematic errors as correlated

NODE=M080DM
 → UNCHECKED ←

99.53 ± 0.23 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

99.15 ± 0.07 ± 0.25		ARTUSO	05	CLEO	$\Upsilon(3S) \rightarrow \gamma X$
99 ± 1	169	CRAWFORD	92B	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
100.1 ± 0.4	11147	² HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma X$
100.2 ± 0.5	223	³ HEINTZ	92	CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
99.5 ± 0.1 ± 0.5	25759	MORRISON	91	CLE2	$e^+e^- \rightarrow \gamma X$


OCCUR=2

²A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

NODE=M080DM;LINKAGE=A

³A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M080DM;LINKAGE=B



$\chi_{b1}(2P)$ DECAY MODES

NODE=M080215;NODE=M080

Mode	Fraction (Γ_i/Γ)	Scale factor	
Γ_1 $\omega \Upsilon(1S)$	$(1.63^{+0.40}_{-0.34})\%$		DESIG=3
Γ_2 $\gamma \Upsilon(2S)$	$(19.9 \pm 1.9)\%$		DESIG=2
Γ_3 $\gamma \Upsilon(1S)$	$(9.2 \pm 0.8)\%$	1.1	DESIG=1
Γ_4 $\pi\pi\chi_{b1}(1P)$	$(9.1 \pm 1.3) \times 10^{-3}$		DESIG=4
Γ_5 $D^0 X$	$(8.8 \pm 1.7)\%$		DESIG=5
Γ_6 $\pi^+\pi^-K^+K^-\pi^0$	$(3.1 \pm 1.0) \times 10^{-4}$		DESIG=6
Γ_7 $2\pi^+\pi^-K^-K_S^0$	$(1.1 \pm 0.5) \times 10^{-4}$		DESIG=7
Γ_8 $2\pi^+\pi^-K^-K_S^0 2\pi^0$	$(7.7 \pm 3.2) \times 10^{-4}$		DESIG=8
Γ_9 $2\pi^+2\pi^-2\pi^0$	$(5.9 \pm 2.0) \times 10^{-4}$		DESIG=9
Γ_{10} $2\pi^+2\pi^-K^+K^-$	$(10 \pm 4) \times 10^{-5}$		DESIG=10
Γ_{11} $2\pi^+2\pi^-K^+K^-\pi^0$	$(5.5 \pm 1.8) \times 10^{-4}$		DESIG=11
Γ_{12} $2\pi^+2\pi^-K^+K^-2\pi^0$	$(10 \pm 4) \times 10^{-4}$		DESIG=12
Γ_{13} $3\pi^+2\pi^-K^-K_S^0\pi^0$	$(6.7 \pm 2.6) \times 10^{-4}$		DESIG=13
Γ_{14} $3\pi^+3\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$		DESIG=14
Γ_{15} $3\pi^+3\pi^-2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$		DESIG=15
Γ_{16} $3\pi^+3\pi^-K^+K^-$	$(2.0 \pm 0.8) \times 10^{-4}$		DESIG=16
Γ_{17} $3\pi^+3\pi^-K^+K^-\pi^0$	$(6.1 \pm 2.2) \times 10^{-4}$		DESIG=17
Γ_{18} $4\pi^+4\pi^-$	$(1.7 \pm 0.6) \times 10^{-4}$		DESIG=18
Γ_{19} $4\pi^+4\pi^-2\pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$		DESIG=19

 $\chi_{b1}(2P)$ BRANCHING RATIOS

NODE=M080220

 $\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_1/Γ NODE=M080R3
NODE=M080R3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.63^{+0.35}_{-0.31} + 0.16_{-0.15}$	$32.6^{+6.9}_{-6.1}$	4 CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma\omega \Upsilon(1S)$

⁴Using $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (11.3 \pm 0.6)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = 2$
 $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 2 (2.48 \pm 0.06)\%$.

NODE=M080R3;LINKAGE=CR

 $\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M080R2
NODE=M080R2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.199 ± 0.019 OUR AVERAGE				
$0.190 \pm 0.018 \pm 0.017$	4.3k	⁵ LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
$0.356 \pm 0.042 \pm 0.092$		⁶ CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
$0.199 \pm 0.020 \pm 0.022$		⁷ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

⁵LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (2.4 \pm 0.1 \pm 0.2) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R2;LINKAGE=LE

⁶Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma\gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (10.23 \pm 1.20 \pm 1.26) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

NODE=M080R2;LINKAGE=B

⁷Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M080R2;LINKAGE=C

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_3/Γ NODE=M080R1
NODE=M080R1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.092 ± 0.008 OUR AVERAGE				Error includes scale factor of 1.1.
$0.098 \pm 0.005 \pm 0.009$	15k	⁸ LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
$0.120 \pm 0.021 \pm 0.021$		⁹ CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
$0.080 \pm 0.009 \pm 0.007$		¹⁰ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

⁸LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (12.4 \pm 0.3 \pm 0.6) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R1;LINKAGE=LE

⁹Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma\gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (6.47 \pm 1.12 \pm 0.82) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

NODE=M080R1;LINKAGE=B

¹⁰Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M080R1;LINKAGE=C

$\Gamma(\pi\pi\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±1.3 OUR AVERAGE				
9.2±1.1±0.8	31k	11 LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
8.6±2.3±2.1		12 CAWLFIELD	06 CLE3	$\Upsilon(3S) \rightarrow 2(\gamma\pi\ell)$

NODE=M080R4
NODE=M080R4

¹¹LEES 11C measures $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) \times B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) = (1.16 \pm 0.07 \pm 0.12) \times 10^{-3}$. We derive the value assuming $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) = B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)\gamma) = (12.6 \pm 1.2) \times 10^{-2}$.

NODE=M080R4;LINKAGE=LE

¹²CAWLFIELD 06 quote $\Gamma(\chi_{b1}(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming l-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

NODE=M080R4;LINKAGE=CA

 $\Gamma(D^0X)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.8±1.5±0.8	2243	13 BRIERE	08 CLEO	$\Upsilon(3S) \rightarrow \gamma D^0X$

NODE=M080R01
NODE=M080R01

¹³For $p_{D^0} > 2.5$ GeV/c.

NODE=M080R01;LINKAGE=BR

 $\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±1.0±0.3	30	14 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$

NODE=M080R02
NODE=M080R02

¹⁴ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (39 \pm 8 \pm 9) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R02;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.5±0.1	10	15 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$

NODE=M080R03
NODE=M080R03

¹⁵ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (14 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R03;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.7±3.1±0.7	15	16 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

NODE=M080R04
NODE=M080R04

¹⁶ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (97 \pm 30 \pm 26) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R04;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.9±2.0±0.5	36	17 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$

NODE=M080R05
NODE=M080R05

¹⁷ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (74 \pm 16 \pm 19) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R05;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.4±0.1	12	18 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$

NODE=M080R06
NODE=M080R06

¹⁸ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R06;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.5±1.7±0.5	38	19 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$

NODE=M080R07
NODE=M080R07

¹⁹ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (69 \pm 13 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R07;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.6±3.5±0.9	27	20 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
<p>20 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(121 \pm 29 \pm 33) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R08
NODE=M080R08

NODE=M080R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±2.5±0.6	17	21 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
<p>21 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(85 \pm 23 \pm 22) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R09
NODE=M080R09

NODE=M080R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.4±0.1	18	22 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
<p>22 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(15 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R10
NODE=M080R10

NODE=M080R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12±4±1	44	23 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
<p>23 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(150 \pm 30 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R11
NODE=M080R11

NODE=M080R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.0±0.7±0.2	16	24 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
<p>24 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(25 \pm 7 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R12
NODE=M080R12

NODE=M080R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±2.1±0.6	25	25 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
<p>25 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(77 \pm 17 \pm 21) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R13
NODE=M080R13

NODE=M080R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7±0.6±0.2	16	26 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<p>26 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(22 \pm 6 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R14
NODE=M080R14

NODE=M080R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19±7±2	41	27 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
<p>27 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(241 \pm 47 \pm 72) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R15
NODE=M080R15

NODE=M080R15;LINKAGE=AS

$\chi_{b1}(2P)$ Cross-Particle Branching Ratios

NODE=M080230

$$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma \times \Gamma_{21}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$

NODE=M080B01
NODE=M080B01

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±0.3±0.6	15k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

$$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma \times \Gamma_{21}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$

NODE=M080B02
NODE=M080B02

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±0.1±0.2	4.3k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

$$B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X)$$

NODE=M080R16
NODE=M080R16

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.16±0.07±0.12	31k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

$$B(\chi_{b2}(2P) \rightarrow \rho X + \bar{\rho} X)/B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$$

NODE=M080R20
NODE=M080R20

VALUE	DOCUMENT ID	TECN	COMMENT
1.109±0.007±0.040	BRIERE 07	CLEO	$\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

$$B(\chi_{b0}(2P) \rightarrow \rho X + \bar{\rho} X)/B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$$

NODE=M080R21
NODE=M080R21

VALUE	DOCUMENT ID	TECN	COMMENT
1.082±0.025±0.060	BRIERE 07	CLEO	$\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

 $\chi_{b1}(2P)$ REFERENCES

NODE=M080

LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=51887
CAWLFIELD	06	PR D73 012003	C. Cawlfild <i>et al.</i>	(CLEO Collab.)	REFID=50997
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
CRONIN-HEN...04	PRL	92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=49766
CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)	REFID=43177
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586

NODE=M205

 $h_b(2P)$

$$I^G(J^{PC}) = ?^?(1^{+-})$$

OMITTED FROM SUMMARY TABLE

Quantum numbers are quark model predictions.

NODE=M205

 $h_b(2P)$ MASS

NODE=M205M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10259.8±0.5±1.1	90k	MIZUK	12 BELL	$e^+e^- \rightarrow \pi^+\pi^-$ hadrons

NODE=M205M

• • • We do not use the following data for averages, fits, limits, etc. • • •

10259.8±0.6 ^{+1.4} _{-1.0}	83.9k	¹ ADACHI	12 BELL	10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM
---	-------	---------------------	---------	--

¹Superseded by MIZUK 12.

NODE=M205M;LINKAGE=AD

 $h_b(2P)$ DECAY MODES

NODE=M205215;NODE=M205

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	not seen
Γ_2 $\eta_b(1S)\gamma$	(22±5) %
Γ_3 $\eta_b(2S)\gamma$	(48±13) %

DESIG=1

DESIG=2

DESIG=3

 $h_b(2P)$ BRANCHING RATIOS

NODE=M205225

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
not seen	83.9k	ADACHI	12 BELL	10.86	$e^+e^- \rightarrow \pi^+\pi^-$ MM	

NODE=M205R01
NODE=M205R01

$\Gamma(\eta_b(1S)\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

 $22.3 \pm 3.8^{+3.1}_{-3.3}$ 10kMIZUK 12 BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadronsNODE=M205R02
NODE=M205R02 $\Gamma(\eta_b(2S)\gamma)/\Gamma_{\text{total}}$ Γ_3/Γ VALUE (units 10^{-2}) EVTS

DOCUMENT ID TECN COMMENT

 $47.5 \pm 10.5^{+6.8}_{-7.7}$ 26kMIZUK 12 BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadronsNODE=M205R03
NODE=M205R03 **$h_b(2P)$ REFERENCES**

NODE=M205

ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

REFID=53962
REFID=54718 **$\chi_{b2}(2P)$**

$$J^G(J^{PC}) = 0^+(2^{++})$$

J needs confirmation.

NODE=M081

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

NODE=M081

 $\chi_{b2}(2P)$ MASS

NODE=M081M

VALUE (MeV) DOCUMENT ID

 $10268.65 \pm 0.22 \pm 0.50$ OUR EVALUATION From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeVNODE=M081M
→ UNCHECKED ← **$m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$**

NODE=M081M2

VALUE (MeV)

DOCUMENT ID TECN COMMENT

 13.4 ± 0.6 OUR AVERAGE $12.3 \pm 2.6 \pm 0.6$ ¹ AAIJ 14BG LHCB $pp \rightarrow \gamma\mu^+\mu^-X$ $13.5 \pm 0.4 \pm 0.5$ ² HEINTZ 92 CSB2 $e^+e^- \rightarrow \gamma X, \ell^+\ell^-\gamma\gamma$ ¹ From the $\chi_{bj}(2P) \rightarrow \Upsilon(1S)\gamma$ transition.² From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

NODE=M081M2

NODE=M081M2;LINKAGE=B
NODE=M081M2;LINKAGE=A **γ ENERGY IN $\Upsilon(3S)$ DECAY**

NODE=M081DM

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

 86.19 ± 0.22 OUR EVALUATION Treating systematic errors as correlated **86.40 ± 0.18 OUR AVERAGE** $86.04 \pm 0.06 \pm 0.27$ ARTUSO 05 CLEO $\Upsilon(3S) \rightarrow \gamma X$ 86 ± 1

101

CRAWFORD 92B CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ 86.7 ± 0.4

10319

³ HEINTZ 92 CSB2 $e^+e^- \rightarrow \gamma X$ 86.9 ± 0.4

157

⁴ HEINTZ 92 CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ $86.4 \pm 0.1 \pm 0.4$

30741

MORRISON 91 CLE2 $e^+e^- \rightarrow \gamma X$ NODE=M081DM
→ UNCHECKED ←

OCCUR=2

³ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

NODE=M081DM;LINKAGE=A

⁴ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M081DM;LINKAGE=B

 $\chi_{b2}(2P)$ DECAY MODES

NODE=M081215;NODE=M081

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\omega \Upsilon(1S)$	$(1.10^{+0.34}_{-0.30})\%$	
Γ_2 $\gamma \Upsilon(2S)$	$(10.6 \pm 2.6)\%$	S=2.0
Γ_3 $\gamma \Upsilon(1S)$	$(7.0 \pm 0.7)\%$	
Γ_4 $\pi\pi\chi_{b2}(1P)$	$(5.1 \pm 0.9) \times 10^{-3}$	
Γ_5 $D^0 X$	$< 2.4\%$	CL=90%
Γ_6 $\pi^+\pi^-K^+K^-\pi^0$	$< 1.1 \times 10^{-4}$	CL=90%
Γ_7 $2\pi^+\pi^-K^-K_S^0$	$< 9 \times 10^{-5}$	CL=90%
Γ_8 $2\pi^+\pi^-K^-K_S^0 2\pi^0$	$< 7 \times 10^{-4}$	CL=90%

DESIG=3

DESIG=2

DESIG=1

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

Γ_9	$2\pi^+ 2\pi^- 2\pi^0$	$(3.9 \pm 1.6) \times 10^{-4}$	DESIG=9
Γ_{10}	$2\pi^+ 2\pi^- K^+ K^-$	$(9 \pm 4) \times 10^{-5}$	DESIG=10
Γ_{11}	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(2.4 \pm 1.1) \times 10^{-4}$	DESIG=11
Γ_{12}	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(4.7 \pm 2.3) \times 10^{-4}$	DESIG=12
Γ_{13}	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 4 \times 10^{-4}$	DESIG=13
Γ_{14}	$3\pi^+ 3\pi^-$	$(9 \pm 4) \times 10^{-5}$	DESIG=14
Γ_{15}	$3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$	DESIG=15
Γ_{16}	$3\pi^+ 3\pi^- K^+ K^-$	$(1.4 \pm 0.7) \times 10^{-4}$	DESIG=16
Γ_{17}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(4.2 \pm 1.7) \times 10^{-4}$	DESIG=17
Γ_{18}	$4\pi^+ 4\pi^-$	$(9 \pm 5) \times 10^{-5}$	DESIG=18
Γ_{19}	$4\pi^+ 4\pi^- 2\pi^0$	$(1.3 \pm 0.5) \times 10^{-3}$	DESIG=19

CL=90%

$\chi_{b2}(2P)$ BRANCHING RATIOS

NODE=M081220

$\Gamma(\omega \Upsilon(1S))/\Gamma_{total}$ Γ_1/Γ

NODE=M081R3
NODE=M081R3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.10^{+0.32+0.11}_{-0.28-0.10}$	$20.1^{+5.8}_{-5.1}$	⁵ CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

⁵Using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$.

NODE=M081R3;LINKAGE=CR

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{total}$ Γ_2/Γ

NODE=M081R2
NODE=M081R2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.106 ± 0.026 OUR AVERAGE		Error includes scale factor of 2.0. See the ideogram below.		
$0.084 \pm 0.011 \pm 0.010$	2.5k	⁶ LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
$0.135 \pm 0.025 \pm 0.035$		⁷ CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
$0.173 \pm 0.021 \pm 0.019$		⁸ HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

⁶LEES 11J reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{total}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (1.1 \pm 0.1 \pm 0.1) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (4.98 \pm 0.94 \pm 0.62) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$.


⁸Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M081R2;LINKAGE=LE

NODE=M081R2;LINKAGE=B

NODE=M081R2;LINKAGE=C

WEIGHTED AVERAGE
 0.106 ± 0.026 (Error scaled by 2.0)



$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.070±0.007 OUR AVERAGE				
0.070±0.004±0.008	11k	⁹ LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
0.072±0.014±0.013		¹⁰ CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
0.070±0.010±0.006		¹¹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

NODE=M081R1
NODE=M081R1

⁹ LEES 11J reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (9.2 \pm 0.3 \pm 0.4) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R1;LINKAGE=LE

¹⁰ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma\gamma \Upsilon(2S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (5.03 \pm 0.94 \pm 0.63) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$.

NODE=M081R1;LINKAGE=B

¹¹ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M081R1;LINKAGE=C

 $\Gamma(\pi\pi\chi_{b2}(1P))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.1±0.9 OUR AVERAGE				
4.9±0.7±0.6	17k	¹² LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
6.0±1.6±1.4		¹³ CAWLFIELD	06 CLE3	$\Upsilon(3S) \rightarrow 2(\gamma\pi\ell)$

NODE=M081R4
NODE=M081R4

¹² $(0.64 \pm 0.05 \pm 0.08) \times 10^{-3}$. We derive the value assuming $B(\Upsilon(3S) \rightarrow \chi_{b2}(2P)X) = B(\Upsilon(3S) \rightarrow \chi_{b2}(2P)\gamma) = (13.1 \pm 1.6) \times 10^{-2}$.

NODE=M081R4;LINKAGE=LE

¹³ CAWLFIELD 06 quote $\Gamma(\chi_{b2}(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming l-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

NODE=M081R4;LINKAGE=CA

 $\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.4 × 10⁻²	90	^{14,15} BRIERE	08 CLEO	$\Upsilon(3S) \rightarrow \gamma D^0 X$

NODE=M081R01
NODE=M081R01

¹⁴ For $p_{D^0} > 2.5$ GeV/c.

NODE=M081R01;LINKAGE=BR
NODE=M081R01;LINKAGE=RI

¹⁵ The authors also present their result as $(0.2 \pm 1.4 \pm 0.1) \times 10^{-2}$.

 $\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	¹⁶ ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$

NODE=M081R02
NODE=M081R02

¹⁶ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

NODE=M081R02;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	¹⁷ ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$

NODE=M081R03
NODE=M081R03

¹⁷ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 12 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

NODE=M081R03;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	¹⁸ ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

NODE=M081R04
NODE=M081R04

¹⁸ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 87 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

NODE=M081R04;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9±1.6±0.5	23	¹⁹ ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

NODE=M081R05
NODE=M081R05

¹⁹ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (51 \pm 16 \pm 13) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R05;LINKAGE=AS

$\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	11	20 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$
<p>20 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ = $(12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R06
NODE=M081R06

NODE=M081R06;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±1.0±0.3	16	21 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$
<p>21 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ = $(32 \pm 11 \pm 8) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R07
NODE=M081R07

NODE=M081R07;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.7±2.2±0.6	14	22 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$
<p>22 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ = $(62 \pm 23 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R08
NODE=M081R08

NODE=M081R08;LINKAGE=AS

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	23 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$
<p>23 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ < 58×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = 13.1 \times 10^{-2}$.</p>				

NODE=M081R09
NODE=M081R09

NODE=M081R09;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	14	24 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+3\pi^-$
<p>24 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ = $(12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R10
NODE=M081R10

NODE=M081R10;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12±4±1	45	25 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+3\pi^-2\pi^0$
<p>25 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ = $(159 \pm 33 \pm 43) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R11
NODE=M081R11

NODE=M081R11;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.4±0.7±0.2	12	26 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+3\pi^-K^+K^-$
<p>26 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+3\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ = $(19 \pm 7 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R12
NODE=M081R12

NODE=M081R12;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2±1.7±0.5	16	27 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+3\pi^-K^+K^-\pi^0$
<p>27 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+3\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P))]$ = $(55 \pm 16 \pm 15) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R13
NODE=M081R13

NODE=M081R13;LINKAGE=AS

$\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	9	28 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<p>²⁸ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (12 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$ $(13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R14
NODE=M081R14

NODE=M081R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
13±5±2	27	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
<p>²⁹ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (165 \pm 46 \pm 50) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$ $(13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M081R15
NODE=M081R15

NODE=M081R15;LINKAGE=AS

 $\chi_{b2}(2P)$ Cross-Particle Branching Ratios

NODE=M081230

$$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
9.2±0.3±0.4	11k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M081B01
NODE=M081B01

$$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.1±0.1	2.5k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M081B02
NODE=M081B02 $B(\chi_{b2}(2P) \rightarrow \chi_{b2}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b2}(2P)X)$ NODE=M081R16
NODE=M081R16

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.64±0.05±0.08	17k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

 $\chi_{b2}(2P)$ REFERENCES

NODE=M081

AAJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56199
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
CAWLFIELD	06	PR D73 012003	C. Cawlfeld <i>et al.</i>	(CLEO Collab.)	REFID=50997
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
CRONIN-HEN...	04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=49766
CRAWFORD	92B	PL B294 139	G. Crawford, R. Fulton	(CLEO Collab.)	REFID=43177
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586

$\Upsilon(3S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M048

 $\Upsilon(3S)$ MASS

NODE=M048M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10355.2±0.5	¹ ARTAMONOV 00	MD1	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10355.3±0.5	^{2,3} BARU	86B REDE	$e^+e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 86B using new electron mass (COHEN 87).			
² Reanalysis of ARTAMONOV 84.			
³ Superseded by ARTAMONOV 00.			

NODE=M048M

NODE=M048M;LINKAGE=AR
NODE=M048M;LINKAGE=C
NODE=M048M;LINKAGE=RZ **$m\Upsilon(3S) - m\Upsilon(2S)$**

NODE=M048DM2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50±0.02±0.13	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M048DM2

 $\Upsilon(3S)$ WIDTH

NODE=M048W

VALUE (keV)	DOCUMENT ID	COMMENT
20.32±1.85 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"	

NODE=M048W
→ UNCHECKED ← **$\Upsilon(3S)$ DECAY MODES**

NODE=M048215;NODE=M048

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\Upsilon(2S)$ anything	(10.6 ± 0.8) %	
Γ_2 $\Upsilon(2S)\pi^+\pi^-$	(2.82±0.18) %	S=1.6
Γ_3 $\Upsilon(2S)\pi^0\pi^0$	(1.85±0.14) %	
Γ_4 $\Upsilon(2S)\gamma\gamma$	(5.0 ± 0.7) %	
Γ_5 $\Upsilon(2S)\pi^0$	< 5.1 × 10 ⁻⁴	CL=90%
Γ_6 $\Upsilon(1S)\pi^+\pi^-$	(4.37±0.08) %	
Γ_7 $\Upsilon(1S)\pi^0\pi^0$	(2.20±0.13) %	
Γ_8 $\Upsilon(1S)\eta$	< 1 × 10 ⁻⁴	CL=90%
Γ_9 $\Upsilon(1S)\pi^0$	< 7 × 10 ⁻⁵	CL=90%
Γ_{10} $h_b(1P)\pi^0$	< 1.2 × 10 ⁻³	CL=90%
Γ_{11} $h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$	(4.3 ± 1.4) × 10 ⁻⁴	
Γ_{12} $h_b(1P)\pi^+\pi^-$	< 1.2 × 10 ⁻⁴	CL=90%
Γ_{13} $\tau^+\tau^-$	(2.29±0.30) %	
Γ_{14} $\mu^+\mu^-$	(2.18±0.21) %	S=2.1
Γ_{15} e^+e^-	seen	
Γ_{16} hadrons		
Γ_{17} ggg	(35.7 ± 2.6) %	
Γ_{18} γgg	(9.7 ± 1.8) × 10 ⁻³	
Γ_{19} 2H anything	(2.33±0.33) × 10 ⁻⁵	

DESIG=8
DESIG=4
DESIG=10
DESIG=12
DESIG=107
DESIG=3
DESIG=11
DESIG=9
DESIG=106
DESIG=112
DESIG=113
DESIG=114
DESIG=16
DESIG=1
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=101
DESIG=109
DESIG=110
DESIG=117**Radiative decays**

NODE=M048;CLUMP=B

Γ_{20} $\gamma\chi_{b2}(2P)$	(13.1 ± 1.6) %	S=3.4
Γ_{21} $\gamma\chi_{b1}(2P)$	(12.6 ± 1.2) %	S=2.4
Γ_{22} $\gamma\chi_{b0}(2P)$	(5.9 ± 0.6) %	S=1.4
Γ_{23} $\gamma\chi_{b2}(1P)$	(9.9 ± 1.3) × 10 ⁻³	S=2.0
Γ_{24} $\gamma A^0 \rightarrow \gamma$ hadrons	< 8 × 10 ⁻⁵	CL=90%
Γ_{25} $\gamma\chi_{b1}(1P)$	(9 ± 5) × 10 ⁻⁴	S=1.9
Γ_{26} $\gamma\chi_{b0}(1P)$	(2.7 ± 0.4) × 10 ⁻³	
Γ_{27} $\gamma\eta_b(2S)$	< 6.2 × 10 ⁻⁴	CL=90%
Γ_{28} $\gamma\eta_b(1S)$	(5.1 ± 0.7) × 10 ⁻⁴	
Γ_{29} $\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 × 10 ⁻⁴	CL=95%
Γ_{30} $\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	< 5.5 × 10 ⁻⁶	CL=90%
Γ_{31} $\gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$	[b] < 1.6 × 10 ⁻⁴	CL=90%

DESIG=108

Lepton Family number (LF) violating modes

Γ_{32}	$e^\pm \tau^\mp$	LF	< 4.2	$\times 10^{-6}$	CL=90%
Γ_{33}	$\mu^\pm \tau^\mp$	LF	< 3.1	$\times 10^{-6}$	CL=90%

NODE=M048;CLUMP=C

DESIG=111

DESIG=105

[a] $1.5 \text{ GeV} < m_\chi < 5.0 \text{ GeV}$

LINKAGE=C48

[b] For $m_{\tau^+ \tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV.

LINKAGE=MRG

 $\Upsilon(3S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

NODE=M048218

 $\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ **$\Gamma_{16} \Gamma_{15} / \Gamma$**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M048G2

NODE=M048G2

0.414 ± 0.007 OUR AVERAGE0.413 ± 0.004 ± 0.006 ROSNER 06 CLEO 10.4 $e^+ e^- \rightarrow$ hadrons0.45 ± 0.03 ± 0.03 ⁴ GILES 84B CLEO $e^+ e^- \rightarrow$ hadrons⁴ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

NODE=M048G2;LINKAGE=R

 $\Gamma(\Upsilon(1S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ **$\Gamma_6 \Gamma_{15} / \Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M048G01

NODE=M048G01

18.46 ± 0.27 ± 0.77 6.4K ⁵ AUBERT 08BP BABR $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$ ⁵ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

NODE=M048G01;LINKAGE=AU

 $\Upsilon(3S)$ PARTIAL WIDTHS

NODE=M048220

 $\Gamma(e^+ e^-)$ **Γ_{15}**

VALUE (keV)	DOCUMENT ID
-------------	-------------

NODE=M048W2

NODE=M048W2

0.443 ± 0.008 OUR EVALUATION

→ UNCHECKED ←

 $\Upsilon(3S)$ BRANCHING RATIOS

NODE=M048225

 $\Gamma(\Upsilon(2S) \text{ anything}) / \Gamma_{\text{total}}$ **Γ_1 / Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M048R8

NODE=M048R8

0.106 ± 0.008 OUR AVERAGE0.1023 ± 0.0105 4625 ^{6,7,8} BUTLER 94B CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- X$ 0.111 ± 0.012 4891 ^{7,8,9} BROCK 91 CLEO $e^+ e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$ ⁶ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma \gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-)$.

NODE=M048R;LINKAGE=A

⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

NODE=M048R;LINKAGE=B

⁸ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-) = (18.5 \pm 0.8)\%$.

NODE=M048R;LINKAGE=D

⁹ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma \gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

NODE=M048R;LINKAGE=C

 $\Gamma(\Upsilon(2S) \pi^+ \pi^-) / \Gamma_{\text{total}}$ **Γ_2 / Γ**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M048R4

NODE=M048R4

2.82 ± 0.18 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.3.00 ± 0.02 ± 0.14 543k LEES 11C BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$ 2.40 ± 0.10 ± 0.26 800 ¹⁰ AUBERT 08BP BABR $e^+ e^- \rightarrow \gamma \pi^+ \pi^- e^+ e^-$ 3.12 ± 0.49 980 ^{11,12} BUTLER 94B CLE2 $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ 2.13 ± 0.38 974 ¹³ BROCK 91 CLEO $e^+ e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

4.82 ± 0.65 ± 0.53 138 ¹³ WU 93 CUSB $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ 3.1 ± 2.0 5 MAGERAS 82 CUSB $\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ ¹⁰ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008 \text{ keV}$.

NODE=M048R4;LINKAGE=AU

¹¹ From the exclusive mode.


NODE=M048R;LINKAGE=M

¹² Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma \gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-)$.

NODE=M048R4;LINKAGE=A

¹³ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \gamma \gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^0 \pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

NODE=M048R4;LINKAGE=C



$\Gamma(\Upsilon(2S)\pi^0\pi^0)/\Gamma_{total}$ Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.85±0.14 OUR AVERAGE				
1.82±0.09±0.12	4391	¹⁴ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.16±0.39		^{15,16} BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.7 ±0.5 ±0.2	10	¹⁷ HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

¹⁴ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.
¹⁵ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.
¹⁶ From the exclusive mode.
¹⁷ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M048R10
NODE=M048R10

NODE=M048R10;LINKAGE=BH
NODE=M048R;LINKAGE=K
NODE=M048R10;LINKAGE=M
NODE=M048R;LINKAGE=G

$\Gamma(\Upsilon(2S)\gamma\gamma)/\Gamma_{total}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0502±0.0069	¹⁸ BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

¹⁸ From the exclusive mode.

NODE=M048R12
NODE=M048R12

NODE=M048R12;LINKAGE=M

$\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{total}$ Γ_5/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.51	90	¹⁹ HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

¹⁹ Authors assume $B(\Upsilon(2S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

NODE=M048R25
NODE=M048R25

NODE=M048R25;LINKAGE=HE

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{total}$ Γ_6/Γ

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.37±0.08 OUR AVERAGE				
4.32±0.07±0.13	90k	²⁰ LEES	11L BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.01±0.13	190k	²¹ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^-$ MM
4.17±0.06±0.19	6.4K	²² AUBERT	08BP BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
4.52±0.35	11830	²³ BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$
4.46±0.34±0.50	451	²³ WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.30	11221	²³ BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

NODE=M048R3
NODE=M048R3
NODE=M048R3

••• We do not use the following data for averages, fits, limits, etc. •••

4.9 ±1.0	22	GREEN	82 CLEO	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.9 ±1.3	26	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

²⁰ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

²¹ A weighted average of the inclusive and exclusive results.

²² Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

²³ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

NODE=M048R3;LINKAGE=LE

NODE=M048R3;LINKAGE=BH
NODE=M048R3;LINKAGE=AU

NODE=M048R3;LINKAGE=B

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M048R28
 NODE=M048R28

••• We do not use the following data for averages, fits, limits, etc. •••

0.577±0.026±0.060 800 ²⁴ AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$

²⁴ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$,
 $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$.
 Not independent of other values reported by AUBERT 08BP.

NODE=M048R28;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{total}$ Γ_7/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M048R11
 NODE=M048R11

2.20±0.13 OUR AVERAGE

2.24±0.09±0.11 6584 25 BHARI 09 CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

1.99±0.34 56 26 BUTLER 94B CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

2.2 ±0.4 ±0.3 33 27 HEINTZ 92 CSB2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

²⁵ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

²⁶ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality.

²⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M048R11;LINKAGE=BH
 NODE=M048R11;LINKAGE=B
 NODE=M048R;LINKAGE=I

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M048R26
 NODE=M048R26

••• We do not use the following data for averages, fits, limits, etc. •••

0.501±0.043 ²⁸ BHARI 09 CLEO $e^+e^- \rightarrow \Upsilon(3S)$

²⁸ Not independent of other values reported by BHARI 09.

NODE=M048R26;LINKAGE=BH

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{total}$ Γ_8/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R9
 NODE=M048R9

<0.1 90 ²⁹ LEES 11L BABR $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.8 90 ^{29,30} AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$

<0.18 90 ³¹ HE 08A CLEO $e^+e^- \rightarrow \ell^+\ell^-\eta$

<2.2 90 BROCK 91 CLEO $e^+e^- \rightarrow \ell^+\ell^-\eta$

²⁹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³⁰ Using $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

³¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

NODE=M048R9;LINKAGE=LE
 NODE=M048R9;LINKAGE=AU
 NODE=M048R9;LINKAGE=HE

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_8/Γ_6

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R27
 NODE=M048R27

<0.23 90 ³² LEES 11L BABR $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<1.9 90 ³³ AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

³² Not independent of other values reported by LEES 11L.

³³ Not independent of other values reported by AUBERT 08BP.

NODE=M048R27;LINKAGE=LE
 NODE=M048R27;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{total}$ Γ_9/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R24
 NODE=M048R24

<0.07 90 ³⁴ HE 08A CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

³⁴ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

NODE=M048R24;LINKAGE=HE

 $\Gamma(h_b(1P)\pi^0)/\Gamma_{total}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M048R03
 NODE=M048R03

<1.2 × 10⁻³ 90 ³⁵ GE 11 CLEO $\Upsilon(3S) \rightarrow \pi^0$ anything

³⁵ Assuming $M(h_b(1P)) = 9900$ MeV and $\Gamma(h_b(1P)) = 0$ MeV, and allowing $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$ to vary from 0–100%.

NODE=M048R03;LINKAGE=GE

 $\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{total}$ Γ_{11}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M048R33
 NODE=M048R33

4.3±1.1±0.9 LEES 11K BABR $\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$

$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{total}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	³⁶ LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<18		³⁶ BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^- X$
<15		³⁶ BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^- X$
³⁶ For $M(h_b(1P)) = 9900$ MeV.				

NODE=M048R34
NODE=M048R34

NODE=M048R34;LINKAGE=MH

$\Gamma(\tau^+\tau^-)/\Gamma_{total}$ Γ_{13}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.29±0.21±0.22	15k	³⁷ BESSON	07 CLEO	$e^+e^- \rightarrow \Upsilon(3S) \rightarrow \tau^+\tau^-$
³⁷ BESSON 07 reports $[\Gamma(\Upsilon(3S) \rightarrow \tau^+\tau^-)/\Gamma_{total}] / [B(\Upsilon(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$ which we multiply by our best value $B(\Upsilon(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M048R18
NODE=M048R18

NODE=M048R18;LINKAGE=BE

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_{13}/Γ_{14}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.05±0.08±0.05	15k	BESSON	07 CLEO	$e^+e^- \rightarrow \Upsilon(3S)$


NODE=M048R19
NODE=M048R19

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0218±0.0021 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
0.0239±0.0007±0.0010	81k	ADAMS	05 CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0202±0.0019±0.0033		CHEN	89B CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0173±0.0015±0.0011		KAARSBERG	89 CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.033 ±0.013 ±0.007	1096	ANDREWS	83 CLEO	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M048R1
NODE=M048R1

WEIGHTED AVERAGE
0.0218±0.0021 (Error scaled by 2.1)



$\Gamma(g g g)/\Gamma_{total}$ Γ_{17}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
35.7±2.6	3M	³⁸ BESSON	06A CLEO	$\Upsilon(3S) \rightarrow \text{hadrons}$

NODE=M048R30
NODE=M048R30

NODE=M048R30;LINKAGE=BE

³⁸ Calculated using BESSON 06A value of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and the PDG 08 values of $B(\Upsilon(2S) + \text{anything}) = (10.6 \pm 0.8)\%$, $B(\pi^+\pi^-\Upsilon(1S)) = (4.40 \pm 0.10)\%$, $B(\pi^0\pi^0\Upsilon(1S)) = (2.20 \pm 0.13)\%$, $B(\gamma\chi_{b2}(2P)) = (13.1 \pm 1.6)\%$, $B(\gamma\chi_{b1}(2P)) = (12.6 \pm 1.2)\%$, $B(\gamma\chi_{b0}(2P)) = (5.9 \pm 0.6)\%$, $B(\gamma\chi_{b0}(1P)) = (0.30 \pm 0.11)\%$, $B(\mu^+\mu^-) = (2.18 \pm 0.21)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(\gamma g g)/\Gamma_{total}$ BESSON 06A value.

$\Gamma(\gamma g g)/\Gamma_{total}$

Γ_{18}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97±0.18	60k	³⁹ BESSON	06A CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

NODE=M048R31
NODE=M048R31

³⁹ Calculated using BESSON 06A values of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and $\Gamma(g g g)/\Gamma_{total}$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(g g g)/\Gamma_{total}$ BESSON 06A value.

NODE=M048R31;LINKAGE=BE

$\Gamma(\gamma g g)/\Gamma(g g g)$

Γ_{18}/Γ_{17}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.72±0.06±0.49	3M	BESSON	06A CLEO	$\Upsilon(3S) \rightarrow (\gamma +) \text{hadrons}$

NODE=M048R32
NODE=M048R32

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{total}$

Γ_{19}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
2.33±0.15^{+0.31}_{-0.28}	LEES	14G BABR	$e^+ e^- \rightarrow \overline{2H} X$

NODE=M048R00
NODE=M048R00

$\Gamma(\gamma \chi_{b2}(2P))/\Gamma_{total}$


Γ_{20}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.131 ±0.016 OUR AVERAGE				Error includes scale factor of 3.4. See the ideogram below.
0.1579±0.0017±0.0073	568k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
0.111 ±0.005 ±0.004	10319	⁴⁰ HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
0.135 ±0.003 ±0.017	30741	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$

NODE=M048R5
NODE=M048R5

⁴⁰ Supersedes NARAIN 91.

NODE=M048R;LINKAGE=H



$\Gamma(\gamma \chi_{b1}(2P))/\Gamma_{total}$


Γ_{21}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.126 ±0.012 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.
0.1454±0.0018±0.0073	537k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
0.115 ±0.005 ±0.005	11147	⁴¹ HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
0.105 ^{+0.003} _{-0.002} ±0.013	25759	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$

NODE=M048R6
NODE=M048R6

⁴¹ Supersedes NARAIN 91.

NODE=M048R6;LINKAGE=H




$\Gamma(\gamma\chi_{b0}(2P))/\Gamma_{total}$ **Γ_{22}/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.059 ±0.006 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.0677±0.0020±0.0065	225k	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$
0.060 ±0.004 ±0.006	4959	⁴² HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma X$
0.049 ^{+0.003} _{-0.004} ±0.006	9903	MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma X$

NODE=M048R7
NODE=M048R7

⁴²Supersedes NARAIN 91.

NODE=M048R7;LINKAGE=H



$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{total}$ **Γ_{23}/Γ**

VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.9±1.3 OUR AVERAGE					Error includes scale factor of 2.0.
7.5±1.2±0.5	126	^{43,44} KORNICER	11	CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
10.5±0.3 ^{+0.7} _{-0.6}	9.7k	LEES	11J	BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M048R21
NODE=M048R21

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	⁴⁵ ASNER	08A	CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
seen		⁴⁶ HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

⁴³ Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

⁴⁴ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))]$
 $= (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$ which we divide by our best value $B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) = (19.1 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁵ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$
 $< 27.1 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

⁴⁶ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma \Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\Upsilon(1S) \rightarrow \ell^+ \ell^-$.

NODE=M048R21;LINKAGE=KA
 NODE=M048R21;LINKAGE=KR

NODE=M048R21;LINKAGE=AS

NODE=M048R21;LINKAGE=HE

$\Gamma(\gamma \chi_{b1}(1P))/\Gamma_{\text{total}}$

Γ_{25}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.9±0.5 OUR AVERAGE Error includes scale factor of 1.9.

1.6±0.5±0.1		50	^{47,48} KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
-------------	--	----	---------------------------	----	--

0.5±0.3 ^{+0.2} _{-0.1}			LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$
---	--	--	------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7	90		⁴⁹ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
seen			⁵⁰ HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

⁴⁷ Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

⁴⁸ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))] = (5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$ which we divide by our best value $B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) = (33.9 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁹ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] < 2.5 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

⁵⁰ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma \Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\Upsilon(1S) \rightarrow \ell^+ \ell^-$.

NODE=M048R22
 NODE=M048R22

NODE=M048R22;LINKAGE=KA
 NODE=M048R22;LINKAGE=KR

NODE=M048R22;LINKAGE=AS

NODE=M048R22;LINKAGE=HE

$\Gamma(\gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.27±0.04 OUR AVERAGE

0.27±0.04±0.02		2.3k	LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$
----------------	--	------	------	-----	---

0.30±0.04±0.10		8.7k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
----------------	--	------	--------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	90		⁵¹ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
------	----	--	---------------------	-----	---

⁵¹ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 21.9 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M048R15
 NODE=M048R15

NODE=M048R15;LINKAGE=AS

$\Gamma(\gamma \eta_b(2S))/\Gamma_{\text{total}}$

Γ_{27}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 6.2	90		ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
-------	----	--	--------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90		LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$
-----	----	--	------	-----	---

NODE=M048R16
 NODE=M048R16

$\Gamma(\gamma \eta_b(1S))/\Gamma_{\text{total}}$

Γ_{28}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

5.1±0.7 OUR AVERAGE

7.1±1.8±1.3		2.3 ± 0.5k	⁵² BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
-------------	--	------------	-------------------------	----	--

4.8±0.5±0.6		19 ± 3k	⁵² AUBERT	09AQ	BABR $\Upsilon(3S) \rightarrow \gamma X$
-------------	--	---------	----------------------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8.5	90		LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$
------	----	--	------	-----	---

4.8±0.5±1.2		19 ± 3k	^{52,53} AUBERT	08V	BABR $\Upsilon(3S) \rightarrow \gamma X$
-------------	--	---------	-------------------------	-----	--

<4.3	90		⁵⁴ ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
------	----	--	----------------------	----	-------------------------------------

⁵² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

⁵³ Systematic error re-evaluated by AUBERT 09AQ.

⁵⁴ Superseded by BONVICINI 10.

NODE=M048R17
 NODE=M048R17

NODE=M048R17;LINKAGE=BO
 NODE=M048R17;LINKAGE=AU
 NODE=M048R17;LINKAGE=SU

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ (1.5 GeV < m_X < 5.0 GeV) Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	95	ROSNER	07A	CLEO $e^+e^- \rightarrow \gamma X$

NODE=M048R20

NODE=M048R20

NODE=M048R20

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<5.5	90	55 AUBERT	09Z	BABR $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

NODE=M048R04

NODE=M048R04

⁵⁵ For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from 0.27–5.5 $\times 10^{-6}$.

NODE=M048R04;LINKAGE=AU

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 $\times 10^{-4}$	90	56 AUBERT	09P	BABR $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$

NODE=M048R29

NODE=M048R29

⁵⁶ For a narrow scalar or pseudoscalar a_1^0 with $M(\tau^+ \tau^-)$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. Measured 90% CL limits as a function of $M(\tau^+ \tau^-)$ range from 1.5–16 $\times 10^{-5}$.

NODE=M048R29;LINKAGE=AU

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$ (0.3 GeV < m_{A^0} < 7 GeV) Γ_{24}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8 $\times 10^{-5}$	90	57 LEES	11H	BABR $\Upsilon(3S) \rightarrow \gamma \text{hadrons}$

NODE=M048R02

NODE=M048R02

NODE=M048R02

⁵⁷ For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1 $\times 10^{-6}$ to 8 $\times 10^{-5}$.

NODE=M048R02;LINKAGE=LE

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.2	90	LEES	10B	BABR $e^+e^- \rightarrow e^\pm \tau^\mp$

NODE=M048230

NODE=M048R01

NODE=M048R01

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.1	90	LEES	10B	BABR $e^+e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M048R23

NODE=M048R23

••• We do not use the following data for averages, fits, limits, etc. •••

<20.3	95	LOVE	08A	CLEO $e^+e^- \rightarrow \mu^\pm \tau^\mp$
-------	----	------	-----	--

 $\Upsilon(3S)$ REFERENCES

NODE=M048

LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)	REFID=53960
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53877
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53937
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53938
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53355
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53062
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)	REFID=52662
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52262
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52587
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
BESSION	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
BESSION	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	(CLEO Collab.)	REFID=47424

BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)

REFID=43799
 REFID=43313
 REFID=43604
 REFID=41579
 REFID=41580
 REFID=41634
 REFID=41586
 REFID=40919
 REFID=40733
 REFID=40034

REFID=11616
 REFID=22338
 REFID=40033

REFID=22278
 REFID=22280
 REFID=22273
 REFID=22321
 REFID=22359

NODE=M206

$\chi_{b1}(3P)$

$$J^G(J^{PC}) = 0^+(1^{++})$$

Observed in the radiative decay to $\Upsilon(1S, 2S, 3S)$, therefore $C = +$.
 J needs confirmation.

NODE=M206

$\chi_{b1}(3P)$ MASS

NODE=M206M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10512.1 ± 2.1 ± 0.9	351	¹ AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
10515.7 ⁺ ₋ 2.2 ⁺ ₋ 1.5 ₋ 3.9 ₋ 2.1	169	² AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10511.3 ± 1.7 ± 2.5	182	³ AAIJ	14BI LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10530 ± 5 ± 9		⁴ AAD	12A ATLS	$pp \rightarrow \gamma \mu^+ \mu^- X$
10551 ± 14 ± 17		⁴ ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma \mu^+ \mu^- X$

NODE=M206M

OCCUR=2

¹ The mass of the $\chi_{b1}(3P)$ state obtained by combining the results of AAIJ 14BG with that of AAIJ 14BI. The first uncertainty is experimental and the second attributable to the unknown mass splitting, assumed to be $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

NODE=M206M;LINKAGE=B

² From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

NODE=M206M;LINKAGE=A

³ From $\chi_{b1}(3P) \rightarrow \Upsilon(3S)\gamma$ transition assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

NODE=M206M;LINKAGE=C

⁴ The mass barycenter of the merged lineshapes from the $J = 1$ and 2 states.

NODE=M206M;LINKAGE=AA

$\chi_{b1}(3P)$ DECAY MODES

NODE=M206215;NODE=M206

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \Upsilon(1S)\gamma$	seen
$\Gamma_2 \quad \Upsilon(2S)\gamma$	seen
$\Gamma_3 \quad \Upsilon(3S)\gamma$	seen

DESIG=1

DESIG=2

DESIG=3

$\chi_{b1}(3P)$ BRANCHING RATIOS

NODE=M206225

$\Gamma(\Upsilon(1S)\gamma)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	169	⁵ AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
seen		AAD	12A ATLS	$pp \rightarrow \gamma \mu^+ \mu^- X$	
seen		ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma \mu^+ \mu^- X$	

NODE=M206R01

NODE=M206R01

⁵ From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

NODE=M206R01;LINKAGE=A

$\Gamma(\Upsilon(2S)\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	169	⁶ AAIJ	14BG LHCB	$pp \rightarrow \gamma\mu^+\mu^- X$
seen		AAD	12A ATLS	$pp \rightarrow \gamma\mu^+\mu^- X$

NODE=M206R02
 NODE=M206R02

⁶From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

NODE=M206R02;LINKAGE=A

 $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	182	AAIJ	14BI LHCB	$pp \rightarrow \gamma\mu^+\mu^- X$

NODE=M206R03
 NODE=M206R03

 $\chi_{b1}(3P)$ REFERENCES

AAIJ	14BG JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BI EPJ C74 3092	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	12A PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABAZOV	12Q PR D86 031103	V.M. Abazov <i>et al.</i>	(D0 Collab.)

NODE=M206

REFID=56199
 REFID=56235
 REFID=54037
 REFID=54264

NODE=M047

$\Upsilon(4S)$
 or $\Upsilon(10580)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $\Upsilon(4S)$ MASS

NODE=M047M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10579.4\pm1.2 OUR AVERAGE			
10579.3 \pm 0.4 \pm 1.2	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
10580.0 \pm 3.5	¹ BEBEK	87 CLEO	$e^+e^- \rightarrow$ hadrons
10577.4 \pm 1.0	² LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Reanalysis of BESSON 85.
² No systematic error given.

NODE=M047M

NODE=M047M;LINKAGE=C
 NODE=M047M;LINKAGE=B

 $\Upsilon(4S)$ WIDTH

NODE=M047W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
20.5\pm2.5 OUR AVERAGE			
20.7 \pm 1.6 \pm 2.5	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
20 \pm 2 \pm 4	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
25 \pm 2.5	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M047W

 $\Upsilon(4S)$ DECAY MODES

NODE=M047215;NODE=M047

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $B\bar{B}$	> 96 %	95%
Γ_2 B^+B^-	(51.4 \pm 0.6) %	
Γ_3 D_s^+ anything + c.c.	(17.8 \pm 2.6) %	
Γ_4 $B^0\bar{B}^0$	(48.6 \pm 0.6) %	
Γ_5 $J/\psi K_S^0 + (J/\psi, \eta_c) K_S^0$	< 4 $\times 10^{-7}$	90%
Γ_6 non- $B\bar{B}$	< 4 %	95%
Γ_7 e^+e^-	(1.57 \pm 0.08) $\times 10^{-5}$	
Γ_8 $\rho^+\rho^-$	< 5.7 $\times 10^{-6}$	90%
Γ_9 $K^*(892)^0\bar{K}^0$	< 2.0 $\times 10^{-6}$	90%
Γ_{10} $J/\psi(1S)$ anything	< 1.9 $\times 10^{-4}$	95%
Γ_{11} D^{*+} anything + c.c.	< 7.4 %	90%

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=10

DESIG=12

DESIG=11

DESIG=15

DESIG=6

DESIG=1

DESIG=16

DESIG=22

DESIG=2

DESIG=3

Γ_{12}	ϕ anything	$(7.1 \pm 0.6) \%$			DESIG=4
Γ_{13}	$\phi\eta$	< 1.8	$\times 10^{-6}$	90%	DESIG=13
Γ_{14}	$\phi\eta'$	< 4.3	$\times 10^{-6}$	90%	DESIG=18
Γ_{15}	$\rho\eta$	< 1.3	$\times 10^{-6}$	90%	DESIG=19
Γ_{16}	$\rho\eta'$	< 2.5	$\times 10^{-6}$	90%	DESIG=20
Γ_{17}	$\Upsilon(1S)$ anything	< 4	$\times 10^{-3}$	90%	DESIG=5
Γ_{18}	$\Upsilon(1S)\pi^+\pi^-$	$(8.1 \pm 0.6) \times 10^{-5}$			DESIG=7
Γ_{19}	$\Upsilon(1S)\eta$	$(1.96 \pm 0.28) \times 10^{-4}$			DESIG=17
Γ_{20}	$\Upsilon(2S)\pi^+\pi^-$	$(8.6 \pm 1.3) \times 10^{-5}$			DESIG=9
Γ_{21}	$h_b(1P)\pi^+\pi^-$	not seen			DESIG=21
Γ_{22}	$h_b(1P)\eta$	$(2.18 \pm 0.21) \times 10^{-3}$			DESIG=23
Γ_{23}	2H anything	< 1.3	$\times 10^{-5}$	90%	DESIG=14

$\Upsilon(4S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

Γ_7


VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.272 ± 0.029 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
$0.321 \pm 0.017 \pm 0.029$	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
$0.28 \pm 0.05 \pm 0.01$	³ ALBRECHT	95E ARG	$e^+e^- \rightarrow$ hadrons
$0.192 \pm 0.007 \pm 0.038$	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
0.283 ± 0.037	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

³Using LEYAOUANC 77 parametrization of $\Gamma(s)$.

NODE=M047220

NODE=M047W1
NODE=M047W1

NODE=M047W1;LINKAGE=A



$\Upsilon(4S)$ BRANCHING RATIOS

$B\bar{B}$ DECAYS

The ratio of branching fraction to charged and neutral B mesons is often derived assuming isospin invariance in the decays, and relies on the knowledge of the B^+/B^0 lifetime ratio. "OUR EVALUATION" is obtained based on averages of rescaled data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>. The averaging/rescaling procedure takes into account the common dependence of the measurement on the value of the lifetime ratio.

NODE=M047230

NODE=M047BBD

NODE=M047BBD

$\Gamma(B^+B^-)/\Gamma_{total}$

Γ_2/Γ

VALUE	DOCUMENT ID
0.514 ± 0.006 OUR EVALUATION	Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$

NODE=M047R11
NODE=M047R11
→ UNCHECKED ←

$\Gamma(D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.178±0.021±0.016	⁴ ARTUSO	05B CLE3	$e^+e^- \rightarrow D_s^+ X$

NODE=M047R13
NODE=M047R13

⁴ ARTUSO 05B reports $[\Gamma(\Upsilon(4S) \rightarrow D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (8.0 \pm 0.2 \pm 0.9) \times 10^{-3}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R13;LINKAGE=AR

 $\Gamma(B^0\bar{B}^0)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.486±0.006 OUR EVALUATION	Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$		

NODE=M047R12
NODE=M047R12
→ UNCHECKED ←

••• We do not use the following data for averages, fits, limits, etc. •••

0.487±0.010±0.008	⁵ AUBERT,B	05H BABR	$\Upsilon(4S) \rightarrow \bar{B}B \rightarrow D^*\ell\nu\ell$
-------------------	-----------------------	----------	--

⁵ Direct measurement. This value is averaged with the value extracted from the $\Gamma(B^+B^-) / \Gamma(B^0\bar{B}^0)$ measurements.

NODE=M047R12;LINKAGE=AU

 $\Gamma(B^+B^-)/\Gamma(B^0\bar{B}^0)$ Γ_2/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
1.058±0.024 OUR EVALUATION			
1.006±0.036±0.031	⁶ AUBERT	04F BABR	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K$
1.01 ±0.03 ±0.09	⁶ HASTINGS	03 BELL	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow \text{dileptons}$
1.058±0.084±0.136	⁷ ATHAR	02 CLEO	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow D^*\ell\nu$
1.10 ±0.06 ±0.05	⁸ AUBERT	02 BABR	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow (c\bar{c})K^*$
1.04 ±0.07 ±0.04	⁹ ALEXANDER	01 CLEO	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K^*$

NODE=M047R10
NODE=M047R10
→ UNCHECKED ←⁶ HASTINGS 03 and AUBERT 04F assume $\tau(B^+) / \tau(B^0) = 1.083 \pm 0.017$.⁷ ATHAR 02 assumes $\tau(B^+) / \tau(B^0) = 1.074 \pm 0.028$. Supersedes BARISH 95.⁸ AUBERT 02 assumes $\tau(B^+) / \tau(B^0) = 1.062 \pm 0.029$.⁹ ALEXANDER 01 assumes $\tau(B^+) / \tau(B^0) = 1.066 \pm 0.024$.NODE=M047R10;LINKAGE=F
NODE=M047R10;LINKAGE=D
NODE=M047R10;LINKAGE=E
NODE=M047R10;LINKAGE=C $[\Gamma(J/\psi K_S^0) + \Gamma((J/\psi, \eta_c) K_S^0)]/\Gamma_{\text{total}}$ Γ_5/Γ Forbidden by CP invariance.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	¹⁰ TAJIMA	07A BELL	$\Upsilon(4S) \rightarrow B^0\bar{B}^0$

¹⁰ $\Upsilon(4S)$ with $CP = +1$ decays to the final state with $CP = -1$.NODE=M047R16
NODE=M047R16
NODE=M047R16

NODE=M047R16;LINKAGE=TA

non- $B\bar{B}$ DECAYS

NODE=M047NBB

 $\Gamma(\text{non-}B\bar{B})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	95	BARISH	96B CLEO	e^+e^-

NODE=M047R6
NODE=M047R6 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.57±0.08 OUR AVERAGE			

1.55±0.04±0.07 AUBERT 05Q BABR $e^+e^- \rightarrow \text{hadrons}$ 2.77±0.50±0.49 ¹¹ ALBRECHT 95E ARG $e^+e^- \rightarrow \text{hadrons}$ ¹¹ Using LEYAOUANANC 77 parametrization of $\Gamma(s)$.NODE=M047R5
NODE=M047R5

NODE=M047R5;LINKAGE=A

 $\Gamma(\rho^+\rho^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.7 × 10⁻⁶	90	AUBERT	08BO BABR	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0$

NODE=M047R17
NODE=M047R17 $\Gamma(K^*(892)^0\bar{K}^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.0 × 10⁻⁶	90	SHEN	13A BELL	$e^+e^- \rightarrow K^*(892)^0\bar{K}^0$

NODE=M047R02
NODE=M047R02 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	95	¹² ABE	02D BELL	$e^+e^- \rightarrow J/\psi X \rightarrow \ell^+\ell^- X$

••• We do not use the following data for averages, fits, limits, etc. •••

<4.7 90 ¹² AUBERT 01C BABR $e^+e^- \rightarrow J/\psi X \rightarrow \ell^+\ell^- X$ ¹² Uses $B(J/\psi \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.NODE=M047R1
NODE=M047R1

NODE=M047R;LINKAGE=AC

$\Gamma(D^{*+} \text{ anything} + \text{ c.c.})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
<0.074	90	¹³ ALEXANDER 90C	CLEO	$e^+ e^-$	NODE=M047R2 NODE=M047R2

¹³ For $x > 0.473$.

NODE=M047R2;LINKAGE=A

 $\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
7.1 \pm 0.1 \pm 0.6		HUANG 07	CLEO	$\Upsilon(4S) \rightarrow \phi X$	NODE=M047R3 NODE=M047R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.23 90 ¹⁴ ALEXANDER 90C CLEO $e^+ e^-$ ¹⁴ For $x > 0.52$.

NODE=M047R3;LINKAGE=A

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
<1.8	90	¹⁵ BELOUS 09	BELL	$e^+ e^- \rightarrow \phi\eta$	NODE=M047R14 NODE=M047R14

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.5 90 AUBERT,BE 06F BABR $e^+ e^- \rightarrow \phi\eta$ ¹⁵ Using all intermedite branching fraction values from PDG 08.

NODE=M047R14;LINKAGE=BE

 $\Gamma(\phi\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
<4.3	90	¹⁶ BELOUS 09	BELL	$e^+ e^- \rightarrow \phi\eta'$	NODE=M047R21 NODE=M047R21

¹⁶ Using all intermedite branching fraction values from PDG 08.

NODE=M047R21;LINKAGE=BE

 $\Gamma(\rho\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
<1.3	90	¹⁷ BELOUS 09	BELL	$e^+ e^- \rightarrow \rho\eta$	NODE=M047R22 NODE=M047R22

¹⁷ Using all intermedite branching fraction values from PDG 08.

NODE=M047R22;LINKAGE=BE

 $\Gamma(\rho\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
<2.5	90	¹⁸ BELOUS 09	BELL	$e^+ e^- \rightarrow \rho\eta'$	NODE=M047R23 NODE=M047R23

¹⁸ Using all intermedite branching fraction values from PDG 08.

NODE=M047R23;LINKAGE=BE

 $\Gamma(\Upsilon(1S) \text{ anything})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
<0.004	90	ALEXANDER 90C	CLEO	$e^+ e^-$	NODE=M047R4 NODE=M047R4

 $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
8.1 \pm 0.6 OUR AVERAGE						NODE=M047R7 NODE=M047R7

8.5 \pm 1.3 \pm 0.2 113 \pm 16 ¹⁹ SOKOLOV 09 BELL $e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$ 8.00 \pm 0.64 \pm 0.27 430 ²⁰ AUBERT 08BP BABR $\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.8 \pm 4.0 \pm 0.3 ^{21,22} SOKOLOV 07 BELL $e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$ 9.0 \pm 1.5 \pm 0.2 167 \pm 19 ²³ AUBERT 06R BABR $e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$ <12 90 GLENN 99 CLE2 $e^+ e^-$ ¹⁹ SOKOLOV 09 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (0.211 \pm 0.030 \pm 0.014) \times 10^{-5}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R7;LINKAGE=SK

²⁰ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

NODE=M047R7;LINKAGE=UB

²¹ SOKOLOV 07 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (4.42 \pm 0.81 \pm 0.56) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R7;LINKAGE=SO

²² According to the authors, systematic errors were underestimated.

NODE=M047R7;LINKAGE=US

²³ Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (2.23 \pm 0.25 \pm 0.27) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R7;LINKAGE=AU

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.96±0.26±0.09		56	24 AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\ell^+\ell^-$

NODE=M047R18
 NODE=M047R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.7 90 25 TAMPONI 15 BELL $e^+e^- \rightarrow \Upsilon(1S)\eta$ |

24 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

NODE=M047R18;LINKAGE=UB

25 Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$. |

NODE=M047R18;LINKAGE=A

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_{19}/Γ_{18}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
2.41±0.40±0.12	56	26 AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-(\pi^0)\ell^+\ell^-$

NODE=M047R19
 NODE=M047R19

• • • We do not use the following data for averages, fits, limits, etc. • • •

26 Not independent of other values reported by AUBERT 08BP.

NODE=M047R19;LINKAGE=UB

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.11±0.07		220	27 AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

NODE=M047R9
 NODE=M047R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88±0.17±0.08 97 ± 15 28 AUBERT 06R BABR $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$

<3.9 90 GLENN 99 CLE2 e^+e^-

NODE=M047R9;LINKAGE=UB

27 Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$.

NODE=M047R9;LINKAGE=AU

28 Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = (1.69 \pm 0.26 \pm 0.20) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_{20}/Γ_{18}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.16±0.16±0.14	220	29 AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

NODE=M047R20
 NODE=M047R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

29 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$,

NODE=M047R20;LINKAGE=UB

$B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
not seen	35 ± 21k	30 ADACHI	12 BELL	$10.58 e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

NODE=M047R01
 NODE=M047R01

30 From the upper limit on the ratio of $\sigma(e^+e^- \rightarrow h_b(1P)\pi^+\pi^-)$ at the $\Upsilon(4S)$ to that at the $\Upsilon(5S)$ of 0.27.

NODE=M047R01;LINKAGE=AD

 $\Gamma(h_b(1P)\eta)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.18±0.11±0.18	112k	31 TAMPONI	15 BELL	$e^+e^- \rightarrow h_b(1P)\eta$

NODE=M047R00
 NODE=M047R00

31 Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$. |

NODE=M047R00;LINKAGE=A

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	ASNER	07 CLEO	$e^+e^- \rightarrow \overline{d}X$

NODE=M047R15
 NODE=M047R15

 $\Upsilon(4S)$ REFERENCES

TAMPONI 15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=56996
SHEN 13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55591
ADACHI 12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=53962
BELOUS 09	PL B681 400	K. Belous <i>et al.</i>	(BELLE Collab.)	REFID=53107
SOKOLOV 09	PR D79 051103	A. Sokolov <i>et al.</i>	(BELLE Collab.)	REFID=52760
AUBERT 08BO	PR D78 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52659
AUBERT 08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
PDG 08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER 07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
HUANG 07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=51624
SOKOLOV 07	PR D75 071103	A. Sokolov <i>et al.</i>	(BELLE Collab.)	REFID=51715
TAJIMA 07A	PRL 99 211601	O. Tajima <i>et al.</i>	(BELLE Collab.)	REFID=52066
AUBERT 06R	PRL 96 232001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51143
AUBERT,BE 06F	PR D74 111103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51563

NODE=M047

ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50992
AUBERT	05Q	PR D72 032005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50774
AUBERT,B	05H	PRL 95 042001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50777
AUBERT	04F	PR D69 071101	B.Aubert <i>et al.</i>		REFID=49748
HASTINGS	03	PR D67 052004	N.C. Hastings <i>et al.</i>	(BELLE Collab.)	REFID=49209
ABE	02D	PRL 88 052001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48557
ATHAR	02	PR D66 052003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=48832
AUBERT	02	PR D65 032001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48514
ALEXANDER	01	PRL 86 2737	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48316
AUBERT	01C	PRL 87 162002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48346
GLENN	99	PR D59 052003	S. Glenn <i>et al.</i>		REFID=46890
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44693
ALBRECHT	95E	ZPHY C65 619	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44372
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44139
ALEXANDER	90C	PRL 64 2226	J. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=41346
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)	REFID=40270
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369
LEYAOUANC	77	PL B71 397	A. Le Yaouanc <i>et al.</i>	(ORSAY)	REFID=44695

NODE=M207

 $X(10610)^\pm$

$$I^G(J^P) = 1^+(1^+)$$

OMITTED FROM SUMMARY TABLE

Observed by BONDAR 12 in $\Upsilon(5S)$ decays to $\Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses. Isospin = 1 is favored due to observation by KROKOVNY 13 of a corresponding neutral state produced in $\Upsilon(10860) \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$ decays at a consistent mass.

NODE=M207

 $X(10610)^\pm$ MASS

NODE=M207M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10607.2±2.0	¹ BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10608.5±3.4 ^{+3.7} _{-1.4}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10608.1±1.2 ^{+1.5} _{-0.2}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10607.4±1.5 ^{+0.8} _{-0.2}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10611 ±4 ±3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10609 ±2 ±3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10608 ±2 ±3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10605 ±2 ⁺³ ₋₁	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10599 ⁺⁶ ₋₃ ⁺⁵ ₋₄	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M207M

NODE=M207M

¹ Average of the BONDAR 12 measurements in separate channels.² Correlated with the corresponding result from BONDAR 12.³ Superseded by the average measurement of BONDAR 12.

NODE=M207M;LINKAGE=BO

NODE=M207M;LINKAGE=A

NODE=M207M;LINKAGE=BN

 $X(10610)^\pm$ WIDTH

NODE=M207W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
18.4± 2.4	⁴ BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
18.5± 5.3 ^{+6.1} _{-2.3}	⁵ GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
20.8± 2.5 ^{+0.3} _{-2.1}	⁵ GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
18.7±13.4 ^{+2.5} _{-1.3}	⁵ GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
22.3± 7.7 ^{+3.0} _{-4.0}	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
24.2± 3.1 ^{+2.0} _{-3.0}	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
17.6± 3.0±3.0	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
11.4 ^{+ 4.5} _{- 3.9} ^{+2.1} _{-1.2}	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
13 ⁺¹⁰ _{- 8} ⁺⁹ _{- 7}	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M207W

⁴ Average of the BONDAR 12 measurements in separate channels.⁵ Correlated with the corresponding result from BONDAR 12.⁶ Superseded by the average measurement of BONDAR 12.

NODE=M207W;LINKAGE=BO

NODE=M207W;LINKAGE=A

NODE=M207W;LINKAGE=BN

X(10610)⁺ DECAY MODES

NODE=M207215;NODE=M207

X(10610)⁻ decay modes are charge conjugates of the modes below.

NODE=M207

Mode	Fraction (Γ_i/Γ)
Γ_1 $\gamma(1S)\pi^+$	seen
Γ_2 $\gamma(2S)\pi^+$	seen
Γ_3 $\gamma(3S)\pi^+$	seen
Γ_4 $h_b(1P)\pi^+$	seen
Γ_5 $h_b(2P)\pi^+$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

X(10610)[±] BRANCHING RATIOS

NODE=M207225

 $\Gamma(\gamma(1S)\pi^+)/\Gamma_{\text{total}}$ Γ_1/Γ NODE=M207R01
NODE=M207R01

VALUE	DOCUMENT ID	TECN	COMMENT
seen	GARMASH 15	BELL	$e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$
seen	BONDAR 12	BELL	$e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$

 $\Gamma(\gamma(2S)\pi^+)/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M207R02
NODE=M207R02

VALUE	DOCUMENT ID	TECN	COMMENT
seen	GARMASH 15	BELL	$e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$
seen	BONDAR 12	BELL	$e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$

 $\Gamma(\gamma(3S)\pi^+)/\Gamma_{\text{total}}$ Γ_3/Γ NODE=M207R03
NODE=M207R03

VALUE	DOCUMENT ID	TECN	COMMENT
seen	GARMASH 15	BELL	$e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$
seen	BONDAR 12	BELL	$e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$

 $\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$ Γ_4/Γ NODE=M207R04
NODE=M207R04

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

 $\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$ Γ_5/Γ NODE=M207R05
NODE=M207R05

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

X(10610)[±] REFERENCES

NODE=M207

GARMASH 15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)
KROKOVNY 13	PR D88 052016	P. Krokovny <i>et al.</i>	(BELLE Collab.)
BONDAR 12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)

REFID=56811

REFID=55588

REFID=53963

X(10610)⁰

$$I^G(J^P) = 1^+(1^+)$$

NODE=M214

OMITTED FROM SUMMARY TABLE

Observed by KROKOVNY 13 in $\Upsilon(10860) \rightarrow \Upsilon(nS)\pi^0\pi^0$ ($n=2,3$).
 Isospin 1 is favored from the proximity in mass to $X(10610)^\pm$ and
 their similarity of observed decay modes and cross sections. J^P
 $= 1^+$ is favored from angular analysis of $X(10610)^\pm$ decays by
 BONDAR 12.

NODE=M214

X(10610)⁰ MASS

NODE=M214M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10609±4±4	¹ KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$

NODE=M214M

¹From a simultaneous fit to the KROKOVNY 13 Dalitz analysis of $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$ decays with fixed width $\Gamma(X(10610)^0) = 18.4$ MeV.

NODE=M214M;LINKAGE=A

X(10610)⁰ DECAY MODES

NODE=M214215;NODE=M214

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \Upsilon(1S)\pi^0$	not seen
$\Gamma_2 \quad \Upsilon(2S)\pi^0$	seen
$\Gamma_3 \quad \Upsilon(3S)\pi^0$	seen

DESIG=1

DESIG=2

DESIG=3

X(10610)⁰ BRANCHING RATIOS

NODE=M214225

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
not seen		KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0\pi^0$	

NODE=M214R01
NODE=M214R01

$\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen		² KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^0\pi^0$	
² Combined significance in $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$, including systematics, of 6.5σ .					

NODE=M214R02
NODE=M214R02

NODE=M214R02;LINKAGE=A

$\Gamma(\Upsilon(3S)\pi^0)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
seen		³ KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^0\pi^0$	
³ Combined significance in $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$, including systematics, of 6.5σ .					

NODE=M214R03
NODE=M214R03

NODE=M214R03;LINKAGE=A

X(10610)⁰ REFERENCES

NODE=M214

KROKOVNY 13	PR D88 052016	P. Krokovny <i>et al.</i>	(BELLE Collab.)
BONDAR 12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)

REFID=55588
REFID=53963

X(10650)[±]

$$I^G(J^P) = ?^+(1^+)$$

NODE=M208

OMITTED FROM SUMMARY TABLE

Observed by BONDAR 12 in $\Upsilon(5S)$ decays to $\Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses.

NODE=M208

X(10650)[±] MASS

NODE=M208M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10652.2±1.5	¹ BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10656.7±5.0 ^{+1.1} _{-3.1}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10650.7±1.5 ^{+0.5} _{-0.2}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10651.2±1.0 ^{+0.4} _{-0.3}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10657 ±6 ±3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10651 ±2 ±3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10652 ±1 ±2	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10654 ±3 ⁺¹ ₋₂	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10651 ⁺² ₋₃ ⁺³ ₋₂	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M208M

¹ Average of the BONDAR 12 measurements in separate channels.² Correlated with the corresponding result from BONDAR 12.³ Superseded by the average measurement of BONDAR 12.
 NODE=M208M;LINKAGE=BO
 NODE=M208M;LINKAGE=A
 NODE=M208M;LINKAGE=BN
X(10650)[±] WIDTH

NODE=M208W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
11.5± 2.2	⁴ BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
12.1 ^{+11.3} _{-4.8} ^{+2.7} _{-0.6}	⁵ GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
14.2± 3.7 ^{+0.9} _{-0.4}	⁵ GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
9.3± 2.2 ^{+0.3} _{-0.5}	⁵ GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
16.3± 9.8 ^{+6.0} _{-2.0}	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
13.3± 3.3 ^{+4.0} _{-3.0}	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
8.4± 2.0± 2.0	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
20.9 ^{+5.4} _{-4.7} ^{+2.1} _{-5.7}	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
19 ± 7 ⁺¹¹ ₋₇	⁶ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M208W

⁴ Average of the BONDAR 12 measurements in separate channels.⁵ Correlated with the corresponding result from BONDAR 12.⁶ Superseded by the average measurement of BONDAR 12.
 NODE=M208W;LINKAGE=BO
 NODE=M208W;LINKAGE=A
 NODE=M208W;LINKAGE=BN
X(10650)⁺ DECAY MODES

NODE=M208215;NODE=M208

X(10650)⁻ decay modes are charge conjugates of the modes below.

NODE=M208

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Upsilon(1S)\pi^+$	seen
Γ_2 $\Upsilon(2S)\pi^+$	seen
Γ_3 $\Upsilon(3S)\pi^+$	seen
Γ_4 $h_b(1P)\pi^+$	seen
Γ_5 $h_b(2P)\pi^+$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

X(10650)[±] BRANCHING RATIOS

				Γ_1/Γ		
$\Gamma(\Upsilon(1S)\pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT		
seen		GARMASH	15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$	NODE=M208R01
seen		BONDAR	12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$	NODE=M208R01
				Γ_2/Γ		
$\Gamma(\Upsilon(2S)\pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT		
seen		GARMASH	15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$	NODE=M208R02
seen		BONDAR	12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$	NODE=M208R02
				Γ_3/Γ		
$\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT		
seen		GARMASH	15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$	NODE=M208R03
seen		BONDAR	12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$	NODE=M208R03
				Γ_4/Γ		
$\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT		
seen		BONDAR	12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$	NODE=M208R04
						NODE=M208R04
				Γ_5/Γ		
$\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT		
seen		BONDAR	12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$	NODE=M208R05
						NODE=M208R05

X(10650)[±] REFERENCES

GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=56811
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)	REFID=53963

NODE=M208

 $\Upsilon(10860)$

$$J^{PC} = 0^-(1^{--})$$

NODE=M092M

 $\Upsilon(10860)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT		
10891 ± 4 OUR AVERAGE	[10.865 ± 0.008 GeV OUR 2010 AVERAGE Scale factor = 1.1]			NODE=M092M	
10891.1 ± 3.2^{+1.2}_{-2.0}	¹ SANTEL	16	BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$	NODE=M092M
10881.8 ^{+1.0} _{-1.1} ± 1.2	^{2,3} SANTEL	16	BELL	$e^+e^- \rightarrow \text{hadrons}$	NEW
10879 ± 3	^{4,5} CHEN	10	BELL	$e^+e^- \rightarrow \text{hadrons}$	OCCUR=2
10888.4 ^{+2.7} _{-2.6} ± 1.2	⁶ CHEN	10	BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$	OCCUR=2
10876 ± 2	⁴ AUBERT	09E	BABR	$e^+e^- \rightarrow \text{hadrons}$	OCCUR=2
10869 ± 2	⁷ AUBERT	09E	BABR	$e^+e^- \rightarrow \text{hadrons}$	
10868 ± 6 ± 5	⁸ BESSON	85	CLEO	$e^+e^- \rightarrow \text{hadrons}$	
10845 ± 20	⁹ LOVELOCK	85	CUSB	$e^+e^- \rightarrow \text{hadrons}$	

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M092M;LINKAGE=C

² From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).

NODE=M092M;LINKAGE=A

³ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

NODE=M092M;LINKAGE=B

⁴ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M092M;LINKAGE=AU

⁵ The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.

NODE=M092M;LINKAGE=CH

⁶In a model where a flat nonresonant $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.

NODE=M092M;LINKAGE=CE

⁷In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.

NODE=M092M;LINKAGE=UB

⁸Assuming four Gaussians with radiative tails and a single step in R .

NODE=M092M;LINKAGE=BE

⁹In a coupled-channel model with three resonances and a smooth step in R .

NODE=M092M;LINKAGE=LO

$\Upsilon(10860)$ WIDTH

NODE=M092W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
54 ± 7 OUR AVERAGE	[110 ± 13 MeV OUR 2010 AVERAGE]		
53.7^{+7.1+1.3}_{-5.6-5.4}	¹⁰ SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••			
48.5 ^{+1.9+2.0} _{-1.8-2.8}	11,12 SANTEL	16	BELL $e^+e^- \rightarrow$ hadrons
46 ⁺⁹ ₋₇	13,14 CHEN	10	BELL $e^+e^- \rightarrow$ hadrons
30.7 ^{+8.3} _{-7.0} ± 3.1	¹⁵ CHEN	10	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
43 ± 4	¹³ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
74 ± 4	¹⁶ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
112 ± 17 ± 23	¹⁷ BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
110 ± 15	¹⁸ LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons
¹⁰ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.			
¹¹ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).			
¹² Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.			
¹³ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.			
¹⁴ The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.			
¹⁵ In a model where a flat nonresonant $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.			
¹⁶ In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.			
¹⁷ Assuming four Gaussians with radiative tails and a single step in R .			
¹⁸ In a coupled-channel model with three resonances and a smooth step in R .			

NODE=M092W

NEW

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M092W;LINKAGE=C

NODE=M092W;LINKAGE=A

NODE=M092W;LINKAGE=B

NODE=M092W;LINKAGE=AU

NODE=M092W;LINKAGE=CH

NODE=M092W;LINKAGE=CE

NODE=M092W;LINKAGE=UB

NODE=M092W;LINKAGE=BE

NODE=M092W;LINKAGE=LO

$\Upsilon(10860)$ DECAY MODES

NODE=M092215;NODE=M092

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $B\bar{B}X$	(76.2 ^{+2.7} _{-4.0}) %	DESIG=9
Γ_2 $B\bar{B}$	(5.5 ± 1.0) %	DESIG=2
Γ_3 $B\bar{B}^* + \text{c.c.}$	(13.7 ± 1.6) %	DESIG=3
Γ_4 $B^*\bar{B}^*$	(38.1 ± 3.4) %	DESIG=4
Γ_5 $B\bar{B}^{(*)}\pi$	< 19.7 %	90% DESIG=10
Γ_6 $B\bar{B}\pi$	(0.0 ± 1.2) %	DESIG=23
Γ_7 $B^*\bar{B}\pi + B\bar{B}^*\pi$	(7.3 ± 2.3) %	DESIG=24
Γ_8 $B^*\bar{B}^*\pi$	(1.0 ± 1.4) %	DESIG=25
Γ_9 $B\bar{B}\pi\pi$	< 8.9 %	90% DESIG=11
Γ_{10} $B_s^{(*)}\bar{B}_s^{(*)}$	(20.1 ± 3.1) %	DESIG=16
Γ_{11} $B_s\bar{B}_s$	(5 ± 5) × 10 ⁻³	DESIG=5
Γ_{12} $B_s\bar{B}_s^* + \text{c.c.}$	(1.35 ± 0.32) %	DESIG=7
Γ_{13} $B_s^*\bar{B}_s^*$	(17.6 ± 2.7) %	DESIG=8

Γ_{14}	no open-bottom	(3.8 $^{+5.0}_{-0.5}$) %		DESIG=28
Γ_{15}	$e^+ e^-$	(5.7 ± 1.5) $\times 10^{-6}$		DESIG=1
Γ_{16}	$K^*(892)^0 \bar{K}^0$	< 1.0 $\times 10^{-5}$	90%	DESIG=29
Γ_{17}	$\Upsilon(1S) \pi^+ \pi^-$	(5.3 ± 0.6) $\times 10^{-3}$		DESIG=17
Γ_{18}	$\Upsilon(2S) \pi^+ \pi^-$	(7.8 ± 1.3) $\times 10^{-3}$		DESIG=18
Γ_{19}	$\Upsilon(3S) \pi^+ \pi^-$	(4.8 $^{+1.9}_{-1.7}$) $\times 10^{-3}$		DESIG=19
Γ_{20}	$\Upsilon(1S) K^+ K^-$	(6.1 ± 1.8) $\times 10^{-4}$		DESIG=20
Γ_{21}	$h_b(1P) \pi^+ \pi^-$	(3.5 $^{+1.0}_{-1.3}$) $\times 10^{-3}$		DESIG=26
Γ_{22}	$h_b(2P) \pi^+ \pi^-$	(6.0 $^{+2.1}_{-1.8}$) $\times 10^{-3}$		DESIG=27
Γ_{23}	$\chi_{b0}(1P) \pi^+ \pi^- \pi^0$	< 6.3 $\times 10^{-3}$	90%	DESIG=30
Γ_{24}	$\chi_{b0}(1P) \omega$	< 3.9 $\times 10^{-3}$	90%	DESIG=31
Γ_{25}	$\chi_{b0}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	< 4.8 $\times 10^{-3}$	90%	DESIG=32
Γ_{26}	$\chi_{b1}(1P) \pi^+ \pi^- \pi^0$	(1.85 ± 0.33) $\times 10^{-3}$		DESIG=33
Γ_{27}	$\chi_{b1}(1P) \omega$	(1.57 ± 0.30) $\times 10^{-3}$		DESIG=34
Γ_{28}	$\chi_{b1}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	(5.2 ± 1.9) $\times 10^{-4}$		DESIG=35
Γ_{29}	$\chi_{b2}(1P) \pi^+ \pi^- \pi^0$	(1.17 ± 0.30) $\times 10^{-3}$		DESIG=36
Γ_{30}	$\chi_{b2}(1P) \omega$	(6.0 ± 2.7) $\times 10^{-4}$		DESIG=37
Γ_{31}	$\chi_{b2}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	(6 ± 4) $\times 10^{-4}$		DESIG=38
Γ_{32}	$\gamma \chi_b \rightarrow \gamma \Upsilon(1S) \omega$	< 3.8 $\times 10^{-5}$	90%	DESIG=39

Inclusive Decays.

NODE=M092;CLUMP=I

These decay modes are submodes of one or more of the decay modes above.

NODE=M092

Γ_{33}	ϕ anything	(13.8 $^{+2.4}_{-1.7}$) %		DESIG=12
Γ_{34}	D^0 anything + c.c.	(108 ± 8) %		DESIG=13
Γ_{35}	D_s anything + c.c.	(46 ± 6) %		DESIG=6
Γ_{36}	J/ψ anything	(2.06 ± 0.21) %		DESIG=14
Γ_{37}	B^0 anything + c.c.	(77 ± 8) %		DESIG=21
Γ_{38}	B^+ anything + c.c.	(72 ± 6) %		DESIG=22

 $\Upsilon(10860)$ PARTIAL WIDTHS

NODE=M092220

 $\Gamma(e^+ e^-)$ **Γ_{15}**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.3.		
0.22 $\pm 0.05 \pm 0.07$	BESSON	85	CLEO $e^+ e^- \rightarrow$ hadrons
0.365 ± 0.070	LOVELOCK	85	CUSB $e^+ e^- \rightarrow$ hadrons

NODE=M092W1
NODE=M092W1 **$\Upsilon(10860)$ BRANCHING RATIOS**

NODE=M092230

"OUR EVALUATION" is obtained based on averages of rescaled data listed below. The averages and rescaling were performed by the Heavy Flavor Averaging Group (HFAG) and are described at <http://www.slac.stanford.edu/xorg/hfag/>.

NODE=M092230

 $\Gamma(B\bar{B}X)/\Gamma_{\text{total}}$ **Γ_1/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.762 $^{+0.027}_{-0.043}$ OUR EVALUATION				
0.71 ± 0.06 OUR AVERAGE				
0.737 $\pm 0.032 \pm 0.051$	1063	¹⁹ DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^+ X, B^0 X$
0.589 $\pm 0.100 \pm 0.092$		²⁰ HUANG	07	CLEO $\Upsilon(5S) \rightarrow$ hadrons

NODE=M092R13
NODE=M092R13

→ UNCHECKED ←

 $\Gamma(B\bar{B})/\Gamma_{\text{total}}$ **Γ_2/Γ**

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
5.5 $^{+1.0}_{-0.9} \pm 0.4$		²¹ DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^+ X, B^0 X$

NODE=M092R16
NODE=M092R16

••• We do not use the following data for averages, fits, limits, etc. •••

<13.8	90	²⁰ HUANG	07	CLEO $\Upsilon(5S) \rightarrow$ hadrons
-------	----	---------------------	----	---

$\Gamma(B\bar{B})/\Gamma(B\bar{B}X)$					Γ_2/Γ_1	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R05 NODE=M092R05
<0.22	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma_{\text{total}}$					Γ_3/Γ	
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R15 NODE=M092R15
0.137±0.016 OUR AVERAGE						
0.137±0.013±0.011		21 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$	
0.143±0.053±0.027		20 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma(B\bar{B}X)$					Γ_3/Γ_1	
<u>VALUE</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R06 NODE=M092R06
0.24±0.09±0.03	10	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B^* \bar{B}^*)/\Gamma_{\text{total}}$					Γ_4/Γ	
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R14 NODE=M092R14
0.381±0.034 OUR AVERAGE						
0.375 ^{+0.021} _{-0.019} ±0.030		21 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$	
0.436±0.083±0.072		20 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B^* \bar{B}^*)/\Gamma(B\bar{B}X)$					Γ_4/Γ_1	
<u>VALUE</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R07 NODE=M092R07
0.74±0.15±0.08	31	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B\bar{B}^{(*)}\pi)/\Gamma_{\text{total}}$					Γ_5/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R17 NODE=M092R17
<0.197	90	20 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B\bar{B}^{(*)}\pi)/\Gamma(B\bar{B}X)$					Γ_5/Γ_1	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R08 NODE=M092R08
<0.32	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B\bar{B}\pi)/\Gamma_{\text{total}}$					Γ_6/Γ	
<u>VALUE (units 10⁻²)</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R28 NODE=M092R28
0.0±1.2±0.3	0	21 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$	
$[\Gamma(B^* \bar{B}\pi) + \Gamma(B\bar{B}^* \pi)]/\Gamma_{\text{total}}$					Γ_7/Γ	
<u>VALUE (units 10⁻²)</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R29 NODE=M092R29
7.3^{+2.3}_{-2.1}±0.8	38	21 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$	
$\Gamma(B^* \bar{B}^* \pi)/\Gamma_{\text{total}}$					Γ_8/Γ	
<u>VALUE (units 10⁻²)</u>	<u>EVTs</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R30 NODE=M092R30
1.0^{+1.4}_{-1.3}±0.4	5	21 DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$	
$\Gamma(B\bar{B}\pi\pi)/\Gamma_{\text{total}}$					Γ_9/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R18 NODE=M092R18
<0.089	90	20 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B\bar{B}\pi\pi)/\Gamma(B\bar{B}X)$					Γ_9/Γ_1	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R09 NODE=M092R09
<0.14	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow$ hadrons	
$\Gamma(B_s^{(*)} \bar{B}_s^{(*)})/\Gamma_{\text{total}}$					$\Gamma_{10}/\Gamma = (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})/\Gamma$	
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R01 NODE=M092R01
0.201^{+0.030}_{-0.031} OUR EVALUATION						→ UNCHECKED ←
0.189^{+0.027}_{-0.021} OUR AVERAGE						
0.172±0.030		22 ESEN	13	BELL	$\Upsilon(5S) \rightarrow D^0 X, D_s X$	
0.21 ^{+0.06} _{-0.03}		23 HUANG	07	CLEO	$\Upsilon(5S) \rightarrow D_s X$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.180±0.013±0.032		24 DRUTSKOY	07	BELL	$\Upsilon(5S) \rightarrow D^0 X, D_s X$	
0.160±0.026±0.058		25 ARTUSO	05B	CLEO	$e^+ e^- \rightarrow D_X X$	

$$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma(B\bar{B}X)$$

VALUE

DOCUMENT ID

 Γ_{10}/Γ_1 NODE=M092R34
NODE=M092R34

$$0.264^{+0.052}_{-0.045} \text{ OUR EVALUATION}$$

→ UNCHECKED ←

$$\Gamma(B_s^*\bar{B}_s^*)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$$

VALUE (units 10⁻²)

EVTS

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{13}/\Gamma_{10} = \Gamma_{13}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$ NODE=M092R19
NODE=M092R19

87.8±1.5 OUR AVERAGE

87.0±1.7

26,27

ESEN

13

BELL

 $B_s^0 \rightarrow D_s^- \pi^+$

90.5±3.2±0.1

227

27,28

LI

12

BELL

 $B_s^0 \rightarrow J/\psi \eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

90.1^{+3.8}_{-4.0}±0.2

29

LOUVOT

09

BELL

 $10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$ 93⁺⁷₋₉ ±1

29

DRUTSKOY

07A

BELL

Superseded by LOUVOT 09

$$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$$

VALUE (units 10⁻²)

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{11}/\Gamma_{10} = \Gamma_{11}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$ NODE=M092R24
NODE=M092R242.6^{+2.6}_{-2.5}

LOUVOT

09

BELL

 $10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

$$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^*\bar{B}_s^*)$$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{11}/Γ_{13} NODE=M092R03
NODE=M092R03

<0.16

90

BONVICINI

06

CLE3

 $e^+ e^-$

$$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$$

VALUE (units 10⁻²)

EVTS

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{12}/\Gamma_{10} = \Gamma_{12}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$ NODE=M092R25
NODE=M092R25

6.7±1.2 OUR AVERAGE

7.3±1.4

26,27

ESEN

13

BELL

 $B_s^0 \rightarrow D_s^- \pi^+$

4.9±2.5±0.0

227

27,28

LI

12

BELL

 $B_s^0 \rightarrow J/\psi \eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.3^{+3.3}_{-3.0}±0.1

LOUVOT

09

BELL

 $10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

$$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^*\bar{B}_s^*)$$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{12}/Γ_{13} NODE=M092R04
NODE=M092R04

<0.16

90

BONVICINI

06

CLE3

 $e^+ e^-$

$$\Gamma(\text{no open-bottom})/\Gamma_{\text{total}}$$

VALUE

DOCUMENT ID

 Γ_{14}/Γ NODE=M092R33
NODE=M092R33

$$0.038^{+0.051}_{-0.005} \text{ OUR EVALUATION}$$

→ UNCHECKED ←

$$\Gamma(K^*(892)^0\bar{K}^0)/\Gamma_{\text{total}}$$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{16}/Γ NODE=M092R35
NODE=M092R35<1.0 × 10⁻⁵

90

SHEN

13A

BELL

 $e^+ e^- \rightarrow K^*(892)^0\bar{K}^0$

$$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$$

VALUE (units 10⁻³)

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{17}/Γ NODE=M092R20
NODE=M092R20

5.3±0.3±0.5

325

30

CHEN

08

BELL

 $10.87 e^+ e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$

$$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$$

VALUE (units 10⁻³)

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{18}/Γ NODE=M092R21
NODE=M092R21

7.8±0.6±1.1

186

30

CHEN

08

BELL

 $10.87 e^+ e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

$$\Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$$

VALUE (units 10⁻³)

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{19}/Γ NODE=M092R22
NODE=M092R224.8^{+1.8}_{-1.5}±0.7

10

30

CHEN

08

BELL

 $10.87 e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

$$\Gamma(\Upsilon(1S)K^+K^-)/\Gamma_{\text{total}}$$

VALUE (units 10⁻⁴)

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{20}/Γ NODE=M092R23
NODE=M092R236.1^{+1.6}_{-1.4}±1.0

20

30

CHEN

08

BELL

 $10.87 e^+ e^- \rightarrow \Upsilon(1S)K^+K^-$

$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$					Γ_{21}/Γ_{18}	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M092R31 NODE=M092R31
$0.45 \pm 0.08^{+0.07}_{-0.12}$	ADACHI	12	BELL	10.86 e ⁺ e ⁻ → hadrons		
$\Gamma(h_b(2P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$					Γ_{22}/Γ_{18}	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M092R32 NODE=M092R32
$0.77 \pm 0.08^{+0.22}_{-0.17}$	ADACHI	12	BELL	10.86 e ⁺ e ⁻ → hadrons		
$\Gamma(\chi_{b0}(1P)\pi^+\pi^-\pi^0)/\Gamma_{total}$					Γ_{23}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R36 NODE=M092R36
$<6.3 \times 10^{-3}$	90	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b0}(1P)\omega)/\Gamma_{total}$					Γ_{24}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R37 NODE=M092R37
$<3.9 \times 10^{-3}$	90	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{non-\omega})/\Gamma_{total}$					Γ_{25}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R38 NODE=M092R38
$<4.8 \times 10^{-3}$	90	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{total}$					Γ_{26}/Γ	
<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R39 NODE=M092R39
$1.85 \pm 0.23 \pm 0.23$	80	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b1}(1P)\omega)/\Gamma_{total}$					Γ_{27}/Γ	
<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R40 NODE=M092R40
$1.57 \pm 0.22 \pm 0.21$	60	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{non-\omega})/\Gamma_{total}$					Γ_{28}/Γ	
<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R41 NODE=M092R41
$0.52 \pm 0.15 \pm 0.11$	24	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{total}$					Γ_{29}/Γ	
<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R42 NODE=M092R42
$1.17 \pm 0.27 \pm 0.14$	29	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b2}(1P)\omega)/\Gamma_{total}$					Γ_{30}/Γ	
<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R43 NODE=M092R43
$0.60 \pm 0.23 \pm 0.15$	13	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b2}(1P)\omega)/\Gamma(\chi_{b1}(1P)\omega)$					Γ_{30}/Γ_{27}	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M092R44 NODE=M092R44
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$0.38 \pm 0.16 \pm 0.09$	32 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{non-\omega})/\Gamma_{total}$					Γ_{31}/Γ	
<u>VALUE (units 10⁻³)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R45 NODE=M092R45
$0.61 \pm 0.22 \pm 0.28$	16	31 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{non-\omega})/\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{non-\omega})$					Γ_{31}/Γ_{28}	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M092R46 NODE=M092R46
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$1.20 \pm 0.55 \pm 0.65$	32 HE	14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\gamma\chi_b \rightarrow \gamma\Upsilon(1S)\omega)/\Gamma_{total}$					Γ_{32}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M092R47 NODE=M092R47
$<3.8 \times 10^{-5}$	90	33 HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
$\Gamma(\phi \text{ anything})/\Gamma_{total}$					Γ_{33}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M092R12 NODE=M092R12
$0.138 \pm 0.007^{+0.023}_{-0.015}$	HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \phi X$		

$\Gamma(D^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{34}/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT			
1.076 ± 0.040 ± 0.068	DRUTSKOY	07	BELL	$\Upsilon(5S) \rightarrow D^0 X$		NODE=M092R10 NODE=M092R10
$\Gamma(D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{35}/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT			
0.46 ± 0.06 OUR AVERAGE						NODE=M092R02 NODE=M092R02
0.472 ± 0.024 ± 0.072	24 DRUTSKOY	07	BELL	$\Upsilon(5S) \rightarrow D_s X$		
0.44 ± 0.09 ± 0.04	34 ARTUSO	05B	CLE3	$e^+ e^- \rightarrow D_s X$		
$\Gamma(J/\psi \text{ anything})/\Gamma_{\text{total}}$					Γ_{36}/Γ	
VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT			
2.060 ± 0.160 ± 0.134	DRUTSKOY	07	BELL	$\Upsilon(5S) \rightarrow J/\psi X$		NODE=M092R11 NODE=M092R11
$\Gamma(B^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{37}/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
0.770^{+0.058}_{-0.056} ± 0.061	352	DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^0 X$	NODE=M092R26 NODE=M092R26
$\Gamma(B^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{38}/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
0.721^{+0.039}_{-0.038} ± 0.050	711	DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+ X$	NODE=M092R27 NODE=M092R27

¹⁹ Not independent of DRUTSKOY 10 values for $\Upsilon(5S) \rightarrow B^{\pm,0}$ anything.

²⁰ Using measurements or limits from AQUINES 06.

²¹ Assuming isospin conservation.

²² Supersedes DRUTSKOY 07.

²³ Supersedes ARTUSO 05B. Combining inclusive ϕ , D_s , and B measurements. Using

$$B(D_s^+ \rightarrow \phi \pi^+) = 4.4 \pm 0.6\% \text{ from PDG 06.}$$

²⁴ Using $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6)\%$ from PDG 06.

²⁵ Uses a model-dependent estimate $B(B_s \rightarrow D_s X) = (92 \pm 11)\%$.

²⁶ Supersedes LOUVOT 09.

²⁷ With $N(B_s^{(*)} \bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$.

²⁸ The ratios $N(B_s^* \bar{B}_s^*) / N(B_s^{(*)} \bar{B}_s^{(*)})$ and $N(B_s^* \bar{B}_s^0) / N(B_s^{(*)} \bar{B}_s^{(*)})$ are measured with a correlation coefficient of -0.72 .

²⁹ From a measurement of $\sigma(e^+ e^- \rightarrow B_s^* \bar{B}_s^*) / \sigma(e^+ e^- \rightarrow B_s^{(*)} \bar{B}_s^{(*)})$ at $\sqrt{s} = 10.86$ GeV.

³⁰ Assuming that the observed events are solely due to the $\Upsilon(5S)$ resonance.

³¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

³² Accounting for correlated systematics.

³³ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14. For a state X_b with mass between 10.55 GeV/ c^2 and 10.65 GeV/ c^2 , the obtained 90% upper limit as a function of m_{X_b} varies from 2.6×10^{-5} to 3.8×10^{-5} .

³⁴ ARTUSO 05B reports $[\Gamma(\Upsilon(10860) \rightarrow D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = 0.0198 \pm 0.0019 \pm 0.0038$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M092R13;LINKAGE=DR
NODE=M092R;LINKAGE=HU
NODE=M092R15;LINKAGE=DR
NODE=M092R01;LINKAGE=ES
NODE=M092R01;LINKAGE=HU

NODE=M092R02;LINKAGE=DR
NODE=M092R01;LINKAGE=AR
NODE=M092R19;LINKAGE=ES
NODE=M092R19;LINKAGE=IL
NODE=M092R19;LINKAGE=LI

NODE=M092R19;LINKAGE=DR
NODE=M092R20;LINKAGE=CH
NODE=M092R36;LINKAGE=A
NODE=M092R44;LINKAGE=A
NODE=M092R47;LINKAGE=A

NODE=M092R02;LINKAGE=AR

$\Upsilon(10860)$ REFERENCES

SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57121
HE	14	PRL 113 142001	X.H. He <i>et al.</i>	(BELLE Collab.)	REFID=55927
ESEN	13	PR D87 031101	S. Esen <i>et al.</i>	(BELLE Collab.)	REFID=54894
SHEN	13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55591
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=53962
LI	12	PRL 108 181808	J. Li <i>et al.</i>	(BELLE Collab.)	REFID=54116
CHEN	10	PR D82 091106	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=53531
DRUTSKOY	10	PR D81 112003	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=53358
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52661
LOUVOT	09	PRL 102 021801	R. Louvot <i>et al.</i>	(BELLE Collab.)	REFID=52646
CHEN	08	PRL 100 112001	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=52153
DRUTSKOY	07	PRL 98 052001	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51621
DRUTSKOY	07A	PR D76 012002	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51852
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=51624
AQUINES	06	PRL 96 152001	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51106
BONVICINI	06	PRL 96 022002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=50995
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50992
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369

NODE=M092

$\Upsilon(11020)$

$$J^{PC} = 0^{-}(1^{-}-)$$

NODE=M093

 $\Upsilon(11020)$ MASS

NODE=M093M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M093M

10987.5^{+11.0}_{-3.4} OUR AVERAGE [11019 ± 8 MeV OUR 2015 AVERAGE]

NEW

10987.5^{+6.4+9.1}_{-2.5-2.3}	¹ SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
--	---------------------	----	--

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

11003.0 ± 1.1 ^{+0.9} _{-1.0}	^{2,3} SANTEL	16	BELL $e^+e^- \rightarrow$ hadrons
---	-----------------------	----	-----------------------------------

10996 ± 2	⁴ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
-----------	---------------------	-----	-----------------------------------

11019 ± 5 ± 7	BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
---------------	--------	----	-----------------------------------

11020 ± 30	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons
------------	----------	----	-----------------------------------

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6$ –11.05 GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M093M;LINKAGE=C

² From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82$ –11.05 GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).

NODE=M093M;LINKAGE=A

³ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

NODE=M093M;LINKAGE=B

⁴ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M093M;LINKAGE=AU

 $\Upsilon(11020)$ WIDTH

NODE=M093W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M093W

61⁺⁹₋₂₈ OUR AVERAGE [79 ± 16 MeV OUR 2015 AVERAGE]

NEW

61⁺⁹⁺²₋₁₉₋₂₀	⁵ SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
---	---------------------	----	--

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

39.3 ^{+1.7+1.3} _{-1.6-2.4}	^{6,7} SANTEL	16	BELL $e^+e^- \rightarrow$ hadrons
--	-----------------------	----	-----------------------------------

37 ± 3	⁸ AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
--------	---------------------	-----	-----------------------------------

61 ± 13 ± 22	BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
--------------	--------	----	-----------------------------------

90 ± 20	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons
---------	----------	----	-----------------------------------

⁵ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n=1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6$ –11.05 GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M093W;LINKAGE=C

⁶ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82$ –11.05 GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).

NODE=M093W;LINKAGE=A

⁷ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

NODE=M093W;LINKAGE=B

⁸ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M093W;LINKAGE=AU

 $\Upsilon(11020)$ DECAY MODES

NODE=M093215;NODE=M093

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	$(2.1^{+1.1}_{-0.6}) \times 10^{-6}$

DESIG=1

$\Upsilon(11020)$ PARTIAL WIDTHS $\Gamma(e^+e^-)$ Γ_1

VALUE (keV)

DOCUMENT ID TECN COMMENT

0.130 ± 0.030 OUR AVERAGE

0.095 ± 0.03 ± 0.035

BESSON 85 CLEO $e^+e^- \rightarrow$ hadrons

0.156 ± 0.040

LOVELOCK 85 CUSB $e^+e^- \rightarrow$ hadrons

NODE=M093220

NODE=M093W1
NODE=M093W1 $\Upsilon(11020)$ REFERENCES

SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)

NODE=M093

REFID=57121
REFID=52661
REFID=22368
REFID=22369