

Supersymmetric Particle Searches

The exclusion of particle masses within a mass range (m_1, m_2) will be denoted with the notation “none m_1 – m_2 ” in the VALUE column of the following Listings. The latest unpublished results are described in the “Supersymmetry: Experiment” review.

NODE=S046

NODE=S046

See the related review(s):

[Supersymmetry, Part I \(Theory\)](#)

[Supersymmetry, Part II \(Experiment\)](#)

CONTENTS:

NODE=S046CNT

NODE=S046CNT

- $\tilde{\chi}_1^0$ (Lightest Neutralino) mass limit
 - Accelerator limits for stable $\tilde{\chi}_1^0$
 - Bounds on $\tilde{\chi}_1^0$ from dark matter searches
 - $\tilde{\chi}_1^0$ - p elastic cross section
 - Spin-dependent interactions
 - Spin-independent interactions
 - Other bounds on $\tilde{\chi}_1^0$ from astrophysics and cosmology
 - Unstable $\tilde{\chi}_1^0$ (Lightest Neutralino) mass limit
- $\tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0$ (Neutralinos) mass limits
- $\tilde{\chi}_1^\pm, \tilde{\chi}_2^\pm$ (Charginos) mass limits
- Long-lived $\tilde{\chi}^\pm$ (Chargino) mass limit
- $\tilde{\nu}$ (Sneutrino) mass limit
- Charged sleptons
 - R-parity conserving \tilde{e} (Selectron) mass limit
 - R-parity violating \tilde{e} (Selectron) mass limit
 - R-parity conserving $\tilde{\mu}$ (Smuon) mass limit
 - R-parity violating $\tilde{\mu}$ (Smuon) mass limit
 - R-parity conserving $\tilde{\tau}$ (Stau) mass limit
 - R-parity violating $\tilde{\tau}$ (Stau) mass limit
 - Long-lived $\tilde{\ell}$ (Slepton) mass limit
- \tilde{q} (Squark) mass limit
 - R-parity conserving \tilde{q} (Squark) mass limit
 - R-parity violating \tilde{q} (Squark) mass limit
- Long-lived \tilde{q} (Squark) mass limit
- \tilde{b} (Sbottom) mass limit
 - R-parity conserving \tilde{b} (Sbottom) mass limit
 - R-parity violating \tilde{b} (Sbottom) mass limit
- \tilde{t} (Stop) mass limit
 - R-parity conserving \tilde{t} (Stop) mass limit
 - R-parity violating \tilde{t} (Stop) mass limit
- Heavy \tilde{g} (Gluino) mass limit
 - R-parity conserving heavy \tilde{g} (Gluino) mass limit
 - R-parity violating heavy \tilde{g} (Gluino) mass limit
- Long-lived \tilde{g} (Gluino) mass limit
- Light \tilde{G} (Gravitino) mass limits from collider experiments
- Supersymmetry miscellaneous results

NODE=S046CNT

NODE=S046100

The results shown below, unless stated otherwise, are based on the Minimal Supersymmetric Standard Model (MSSM), as described in the Note on Supersymmetry. Unless otherwise indicated, this includes the assumption of common gaugino and scalar masses at the scale of Grand Unification (GUT), and use of the resulting relations in the spectrum and decay branching ratios. Unless otherwise indicated, it is also assumed that R -parity (R) is conserved and that:

- 1) The $\tilde{\chi}_1^0$ is the lightest supersymmetric particle (LSP),

- 2) $m_{\tilde{f}_L} = m_{\tilde{f}_R}$, where $\tilde{f}_{L,R}$ refer to the scalar partners of left- and right-handed fermions.

Limits involving different assumptions are identified in the Comments or in the Footnotes, in particular also the many simplified models, see definitions below. We summarize here the notations used in this Chapter to characterize some of the most common deviations from the MSSM (for further details, see the Note on Supersymmetry).

Theories with R -parity violation (RPV) are characterized by a superpotential of the form: $\lambda_{ijk} L_i L_j e_k^c + \lambda'_{ijk} L_i Q_j d_k^c + \lambda''_{ijk} u_i^c d_j^c d_k^c$, where i, j, k are generation indices. The presence of any of these couplings is often identified in the following by the symbols $LL\bar{E}$, $LQ\bar{D}$, and \overline{UDD} . Mass limits in the presence of RPV will often refer to “direct” and “indirect” decays. Direct refers to RPV decays of the particle in consideration. Indirect refers to cases where RPV appears in the decays of the LSP. The LSP need not be the $\tilde{\chi}_1^0$.

In several models, most notably in theories with so-called Gauge Mediated Supersymmetry Breaking (GMSB), the gravitino (\tilde{G}) is the LSP. It is usually much lighter than any other massive particle in the spectrum, and $m_{\tilde{G}}$ is then neglected in all decay processes involving gravitinos. In these scenarios, particles other than the neutralino are sometimes considered as the next-to-lightest supersymmetric particle (NLSP), and are assumed to decay to their even- R partner plus \tilde{G} . If the lifetime is short enough for the decay to take place within the detector, \tilde{G} is assumed to be undetected and to give rise to missing energy (\cancel{E}) or missing transverse energy (\cancel{E}_T) signatures.

When needed, specific assumptions on the eigenstate content of $\tilde{\chi}^0$ and $\tilde{\chi}^\pm$ states are indicated, using the notation $\tilde{\gamma}$ (photino), \tilde{H} (higgsino), \tilde{W} (wino), and \tilde{Z} (zino) to signal that the limit of pure states was used. The term gaugino is also used, to generically indicate wino-like charginos and zino-like neutralinos.

In the listings we have made use of the following abbreviations for simplified models employed by the experimental collaborations in supersymmetry searches published in the past year.

WARNING: Experimental lower mass limits determined within simplified models are to be treated with extreme care as they might not be directly applicable to realistic models. This is outlined in detail in the publications and we recommend

consulting them before using bounds. For example, branching ratios, typically fixed to specific values in simplified models, can vary substantially in more elaborate models.

Simplified Models Table

- Tglu1A:** gluino pair production with $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$.
- Tglu1B:** gluino pair production with $\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$.
- Tglu1C:** gluino pair production with a 2/3 probability of having a $\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$ decay and a 1/3 probability of having a $\tilde{g} \rightarrow qq\tilde{\chi}_2^0$, $\tilde{\chi}_2^0 \rightarrow Z^\pm\tilde{\chi}_1^0$ decay.
- Tglu1D:** gluino pair production with one gluino decaying to $qq'\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm + \tilde{G}$, and the other gluino decaying to $q\bar{q}\tilde{\chi}_1^0$ with $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$.
- Tglu1E:** gluino pair production with $\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_2^0$ and $\tilde{\chi}_2^0 \rightarrow Z^\pm\tilde{\chi}_1^0$ where $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_2^0} = (m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})/2$.
- Tglu1F:** gluino pair production with $\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm$ or $\tilde{g} \rightarrow qq\tilde{\chi}_2^0$ with equal branching ratios, where $\tilde{\chi}_1^\pm$ decays through an intermediate scalar tau lepton or sneutrino to $\tau\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate scalar tau lepton or sneutrino to $\tau^+\tau^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$; the mass hierarchy is such that $m_{\tilde{\chi}_1^\pm} \sim m_{\tilde{\chi}_2^0} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$ and $m_{\tilde{\tau},\tilde{\nu}} = (m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})/2$.
- Tglu1G:** gluino pair production with $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_2^0$, and $\tilde{\chi}_2^0$ decaying through an intermediate slepton or sneutrino to $l^+l^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$ where $m_{\tilde{\chi}_2^0} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$ and $m_{\tilde{l},\tilde{\nu}} = (m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})/2$.
- Tglu1H:** gluino pair production with $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_2^0$, and $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 Z^{0(*)}$.
- Tglu1I:** gluino pair production with $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_2^0$, and $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 H$.
- Tglu1J:** gluino pair production with $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_2^0$, and $\text{BR}(\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 Z^{0(*)}) = \text{BR}(\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 H) = 0.5$.
- Tglu1LL** gluino pair production where $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ happens with 1/3 probability and $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^\pm$ happens with 2/3 probability. The $\tilde{\chi}_1^\pm$ is assumed to be few hundreds of MeV heavier than the $\tilde{\chi}_1^0$, and decays to $\tilde{\chi}_1^0$ via a pion.
- Tglu2A:** gluino pair production with $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$.
- Tglu3A:** gluino pair production with $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$.
- Tglu3B:** gluino pair production with $\tilde{g} \rightarrow t\bar{t}$ where \tilde{t} decays exclusively to $t\tilde{\chi}_1^0$.
- Tglu3C:** gluino pair production with $\tilde{g} \rightarrow t\bar{t}$ where \tilde{t} decays exclusively to $c\tilde{\chi}_1^0$.
- Tglu3D:** gluino pair production with $\tilde{g} \rightarrow t\bar{b}\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$.
- Tglu3E:** gluino pair production where the gluino decays 25% of the time through $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$, 25% of the time through $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$ and 50% of the time through $\tilde{g} \rightarrow t\bar{b}\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$.
- Tglu3F:** gluino pair production with wino-like couplings to electroweakinos, that is: $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_{1,2}^0$ with BR 17%, $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_{1,2}^0$ with BR 17%, $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^\pm$ with BR 66%.
- Tglu3G:** gluino pair production with higgsino-like couplings to electroweakinos, that is: $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_{1,2}^0$ with BR 50%, $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^\pm$ with BR 50%.
- Tglu4A:** gluino pair production with one gluino decaying to $qq'\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm + \tilde{G}$, and the other gluino decaying to $q\bar{q}\tilde{\chi}_1^0$ with $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$.
- Tglu4B:** gluino pair production with gluinos decaying to $q\bar{q}\tilde{\chi}_1^0$ and $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$.
- Tglu4C:** gluino pair production with gluinos decaying to $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ and $\tilde{\chi}_1^0 \rightarrow Z + \tilde{G}$.

- Tglu4D:** gluino pair production with $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ where the $\tilde{\chi}_1^0$ decays with equal probability to $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$ or to $\tilde{\chi}_1^0 \rightarrow H + \tilde{G}$.
- Tglu4E:** gluino pair production with $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$ where the $\tilde{\chi}_1^0$ decays with equal probability to $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$ or to $\tilde{\chi}_1^0 \rightarrow Z + \tilde{G}$.
- Tglu4F:** gluino pair production with $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ where the $\tilde{\chi}_1^0$ decays with equal probability to $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$ or to $\tilde{\chi}_1^0 \rightarrow Z + \tilde{G}$.
- Tglu4G:** gluino pair production with $\tilde{g} \rightarrow qq\tilde{\chi}_1^0$ where the $\tilde{\chi}_1^0$ decays with equal probability to $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$ or to $\tilde{\chi}_1^0 \rightarrow Z + \tilde{G}$.
- Tglu1RPV:** gluino pair production with $\tilde{g} \rightarrow uds$ via RPV coupling λ''_{112} .
- Tglu2RPV:** gluino pair production with $\tilde{g} \rightarrow (tbd, tbs)$ via RPV coupling λ''_{313} or λ''_{323} .

-
- Tsqk1:** squark pair production with $\tilde{q} \rightarrow q\tilde{\chi}_1^0$.
- Tsqk1LL** squark pair production where $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ and $\tilde{q} \rightarrow q\tilde{\chi}_1^\pm$ each happen with 50% probability. The $\tilde{\chi}_1^\pm$ is assumed to be few hundreds of MeV heavier than the $\tilde{\chi}_1^0$, and decays to $\tilde{\chi}_1^0$ via a pion.
- Tsqk2:** squark pair production with $\tilde{q} \rightarrow q\tilde{\chi}_2^0$ and $\tilde{\chi}_2^0 \rightarrow Z + \tilde{\chi}_1^0$.
- Tsqk2A:** squark pair production with $\tilde{q} \rightarrow q\tilde{\chi}_2^0$, where one of the $\tilde{\chi}_2^0 \rightarrow Z^{(*)}\tilde{\chi}_1^0 \rightarrow f\bar{f}\tilde{\chi}_1^0$ and the other $\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\ell^+ \rightarrow \ell^+\ell^-\tilde{\chi}_1^0$.
- Tsqk3:** squark pair production with $\tilde{q} \rightarrow q'\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$ (like Tglu1B but for squarks)
- Tsqk4:** squark pair production with squarks decaying to $q\tilde{\chi}_1^0$ and $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$.
- Tsqk4A:** squark pair production with one squark decaying to $q\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm + \tilde{G}$, and the other squark decaying to $q\tilde{\chi}_1^0$ with $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$.
- Tsqk4B:** squark pair production with squarks decaying to $q\tilde{\chi}_1^0$ and $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$.

-
- Tstop1:** stop pair production with $\tilde{t} \rightarrow t\tilde{\chi}_1^0$.
- Tstop1LL** stop pair production where $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ and $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ each happen with 50% probability. The $\tilde{\chi}_1^\pm$ is assumed to be few hundreds of MeV heavier than the $\tilde{\chi}_1^0$, and decays to $\tilde{\chi}_1^0$ via a pion.
- Tstop2:** stop pair production with $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$.
- Tstop3:** stop pair production with the subsequent four-body decay $\tilde{t} \rightarrow bff'\tilde{\chi}_1^0$ where f represents a lepton or a quark.
- Tstop4:** stop pair production with $\tilde{t} \rightarrow c\tilde{\chi}_1^0$.
- Tstop5:** stop pair production with $\tilde{t} \rightarrow b\bar{\nu}\tilde{\tau}$ with $\tilde{\tau} \rightarrow \tau\tilde{G}$.
- Tstop6:** stop pair production with $\tilde{t} \rightarrow t + \tilde{\chi}_2^0$, where $\tilde{\chi}_2^0 \rightarrow Z + \tilde{\chi}_1^0$ or $H + \tilde{\chi}_1^0$ each with BR 50%.
- Tstop7:** stop pair production with $\tilde{t}_2 \rightarrow \tilde{t}_1 + H/Z$, where $\tilde{t}_1 \rightarrow t + \tilde{\chi}_1^0$.
- Tstop8:** stop pair production with equal probability of the stop decaying via $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ or via $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$.
- Tstop9:** stop pair production with equal probability of the stop decaying via $\tilde{t} \rightarrow c\tilde{\chi}_1^0$ or via the four-body decay $\tilde{t} \rightarrow bff'\tilde{\chi}_1^0$ where f represents a lepton or a quark.
- Tstop10:** stop pair production with $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^\pm \rightarrow W^{\pm*}\tilde{\chi}_1^0 \rightarrow (f\bar{f}') + \tilde{\chi}_1^0$ with a virtual W -boson.
- Tstop11:** stop pair production with $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm$ decaying through an intermediate slepton to $l\nu\tilde{\chi}_1^0$
- Tstop12:** stop pair production with $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ and $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$
- Tstop13:** stop pair production with $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ where the $\tilde{\chi}_1^0$ can decay with equal probability to $\tilde{\chi}_1^0 \rightarrow \gamma + \tilde{G}$ or to $\tilde{\chi}_1^0 \rightarrow Z + \tilde{G}$.
- Tstop14:** stop pair production with wino-like couplings to electroweakinos, that is: $\tilde{t} \rightarrow t\tilde{\chi}_{1,2}^0$ with BR 33%, $\tilde{g} \rightarrow b\tilde{\chi}_1^\pm$ with BR 67%.

- Tstop15:** stop pair production with higgsino-like couplings to electroweakinos, that is: $\tilde{t} \rightarrow t\tilde{\chi}_{1,2}^0$ with BR 50%, $\tilde{g} \rightarrow b\tilde{\chi}_1^\pm$ with BR 50%.
- Tstop16:** stop pair production with $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$, followed either by $\tilde{\chi}_1^\pm \rightarrow \nu_\tau\tilde{\tau}_1$ and $\tilde{\tau}_1 \rightarrow \tau\tilde{\chi}_1^0$, or by $\tilde{\chi}_1^\pm \rightarrow \tau\tilde{\nu}_\tau$ and $\tilde{\nu}_\tau \rightarrow \nu\tilde{\chi}_1^0$, each with BR 50%.
- Tstop1RPV:** stop pair production with $\tilde{t} \rightarrow \bar{b}\bar{s}$ via RPV coupling λ''_{323} .
- Tstop2RPV:** stop pair production with $\tilde{t} \rightarrow b\ell$, via RPV coupling λ''_{i33} .
- Tstop3RPV:** stop pair production with $\tilde{t} \rightarrow q\mu$, via RPV coupling λ''_{23k} .
- Tstop4RPV:** stop pair production with $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow bbs$ via RPV coupling λ''_{323} .
- Tstop5RPV:** stop pair production with $\tilde{t} \rightarrow t\tilde{\chi}_{1,2}^0$, $\tilde{\chi}_{1,2}^0 \rightarrow tbs$ via RPV coupling λ''_{323} .
-
- Tsbot1:** sbottom pair production with $\tilde{b} \rightarrow b\tilde{\chi}_1^0$.
- Tsbot2:** sbottom pair production with $\tilde{b} \rightarrow t\chi_1^-, \chi_1^- \rightarrow W^-\tilde{\chi}_1^0$.
- Tsbot3:** sbottom pair production with $\tilde{b} \rightarrow b\tilde{\chi}_2^0$, where one of the $\tilde{\chi}_2^0 \rightarrow Z^{(*)}\tilde{\chi}_1^0 \rightarrow f\bar{f}\tilde{\chi}_1^0$ and the other $\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\ell^+ \rightarrow \ell^+\ell^-\tilde{\chi}_1^0$.
- Tsbot4:** sbottom pair production with $\tilde{b} \rightarrow b\tilde{\chi}_2^0$, with $\tilde{\chi}_2^0 \rightarrow H\tilde{\chi}_1^0$.
-
- Tchi1chi1A:** electroweak pair and associated production of nearly mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$, where $\tilde{\chi}_1^\pm$ decays to $\tilde{\chi}_1^0$ plus soft radiation, and where one of the $\tilde{\chi}_1^0$ decays to $\gamma + \tilde{G}$ while the other one decays to $Z/H + \tilde{G}$ (with equal probability).
- Tchi1chi1B:** electroweak pair production of charginos $\tilde{\chi}_1^\pm$, where $\tilde{\chi}_1^\pm$ decays through an intermediate slepton or sneutrino to $l\nu\tilde{\chi}_1^0$ and where the slepton or sneutrino mass is 5%, 25%, 50%, 75% and 95% of the $\tilde{\chi}_1^\pm$ mass.
- Tchi1chi1C:** electroweak pair production of charginos $\tilde{\chi}_1^\pm$, where $\tilde{\chi}_1^\pm$ decays through an intermediate slepton or sneutrino to $l\nu\tilde{\chi}_1^0$ and where $m_{\tilde{\ell},\tilde{\nu}} = (m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})/2$.
- Tchi1chi1D:** electroweak associated pair production of charginos $\tilde{\chi}_1^\pm$, where $\tilde{\chi}_1^\pm$ decays through an intermediate scalar tau lepton or sneutrino to $\tau\nu\tilde{\chi}_1^0$ and where $m_{\tilde{\tau},\tilde{m}_\nu} = (m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})/2$.
- Tchi1chi1F:** electroweak pair and associated production of nearly mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$ (*i.e.* $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^\pm\tilde{\chi}_1^0$ production) where the $\tilde{\chi}_1^\pm$ decays exclusively to $\tilde{\chi}_1^0$ plus soft radiation and the $\tilde{\chi}_1^0$ decays to $\gamma/Z + \tilde{G}$.
- Tchi1chi1G:** electroweak pair production of charginos $\tilde{\chi}_1^\pm$, which are nearly mass-degenerate with neutralinos $\tilde{\chi}_1^0$. The $\tilde{\chi}_1^\pm$ decays either to $W^\pm + \tilde{G}$, or to $\tilde{\chi}_1^0$ plus soft radiation. The $\tilde{\chi}_1^0$ decays exclusively to $\gamma + \tilde{G}$.
- Tchi1chi1H:** electroweak pair production of charginos $\tilde{\chi}_1^\pm$, with $\tilde{\chi}_1^\pm \rightarrow W^\pm + \tilde{\chi}_1^0$ and $W^\pm \rightarrow \ell^\pm + \nu$.
- Tchi1chi1I:** electroweak pair production of charginos $\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$ and $W^\pm \rightarrow qq'$.
-
- Tchi1n1A:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$, where $\tilde{\chi}_1^\pm$ decays exclusively to $W^\pm + \tilde{G}$ and $\tilde{\chi}_1^0$ decays exclusively to $\gamma + \tilde{G}$.
- Tchi1n2A:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate slepton or sneutrino to $l\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate slepton or sneutrino to $l^+l^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$.

- Tchi1n2B:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate slepton or sneutrino to $l\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate slepton or sneutrino to $l^+l^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$ and where the slepton or sneutrino mass is 5%, 25%, 50%, 75% and 95% of the $\tilde{\chi}_1^\pm$ mass.
- Tchi1n2C:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate slepton or sneutrino to $l\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate slepton or sneutrino to $l^+l^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$ and where $m_{\tilde{\ell},\tilde{\nu}} = (m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})/2$.
- Tchi1n2D:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate scalar tau lepton or sneutrino to $\tau\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate scalar tau lepton or sneutrino to $\tau^+\tau^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$ and where $m_{\tilde{\tau},\tilde{\nu}} = (m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})/2$.
- Tchi1n2E:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm \rightarrow W^\pm + \tilde{\chi}_1^0$ and $\tilde{\chi}_2^0 \rightarrow H + \tilde{\chi}_1^0$.
- Tchi1n2F:** electroweak associated production of mass-degenerate wino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate $W^{\pm*}$ to $l\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate Z^* to $l^+l^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$.
- Tchi1n2Fa:** electroweak associated production of mass-degenerate wino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate $W^{\pm*}$ to $q\bar{q}\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate Z^* to $l^+l^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$.
- Tchi1n2Fb:** electroweak associated production of mass-degenerate wino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate $W^{(*)}$ to $q\bar{q}\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate $Z^{(*)}$ to $q\bar{q}\tilde{\chi}_1^0$.
- Tchi1n2Fc:** electroweak associated production of mass-degenerate wino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate $W^{(*)}$ to $q\bar{q}\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate $H^{(*)}$ to $q\bar{q}\tilde{\chi}_1^0$.
- Tchi1n2G:** electroweak associated production of Higgsino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, and electroweak associated production of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$, where $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})/2$ and where $\tilde{\chi}_1^\pm$ decays through an intermediate $W^{\pm*}$ to $q\bar{q}\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate Z^* to $l^+l^-\tilde{\chi}_1^0$.
- Tchi1n2Ga:** electroweak associated production of Higgsino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, and electroweak associated production of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$, where $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})/2$ and where $\tilde{\chi}_1^\pm$ decays through an intermediate $W^{\pm*}$ to $l\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate Z^* to $l^+l^-\tilde{\chi}_1^0$.
- Tchi1n2H:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays through an intermediate slepton or sneutrino to $l\nu\tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays through an intermediate scalar tau lepton or sneutrino to $\tau^+\tau^-\tilde{\chi}_1^0$ or $\nu\bar{\nu}\tilde{\chi}_1^0$.
- Tchi1n2I:** electroweak associated production of mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ decays to $W^\pm + \tilde{\chi}_1^0$ and where $\tilde{\chi}_2^0$ decays 50% of the time to $Z + \tilde{\chi}_1^0$ and 50% of the time to $H + \tilde{\chi}_1^0$.
- Tchi1n2_GGM:** in the framework of General Gauge Mediation (GGM): electroweak pair and associated production of nearly mass-degenerate charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0, \tilde{\chi}_2^0$ (*i.e.* $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm\tilde{\chi}_1^0$).

$\tilde{\chi}_1^\pm \tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ production) where the $\tilde{\chi}_1^\pm$ decays exclusively to $W^\pm + \tilde{G}$, the $\tilde{\chi}_2^0$ decays to $Z/H + \tilde{G}$ and the $\tilde{\chi}_1^0$ decays to $\gamma/Z + \tilde{G}$. The branching ratios depend on the composition of the gauge eigenstates of the neutralinos in the GGM scenario.

TwinoLSPBL: Electroweak pair production of wino-like $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^0$ (i.e. $\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^0 \tilde{\chi}_1^0$). The $\tilde{\chi}_1^\pm$ can decay via bi-linear RPV into $Z\ell$, $H\ell$ or $W\nu$; the $\tilde{\chi}_1^0$ can decay into $Z\nu$, $H\nu$ or $W\ell$.

Tn1n1A: electroweak pair and associated production of nearly mass-degenerate Higgsino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decay to $\tilde{\chi}_1^0$ plus soft radiation and where both of the $\tilde{\chi}_1^0$ decay to $H + \tilde{G}$.

Tn1n1B: electroweak pair and associated production of nearly mass-degenerate Higgsino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decay to $\tilde{\chi}_1^0$ plus soft radiation and where the $\tilde{\chi}_1^0$ decays 50% of the time to $H + \tilde{G}$ and 50 % of the time to $Z + \tilde{G}$.

Tn1n1C: electroweak pair and associated production of nearly mass-degenerate Higgsino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$, where $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decay to $\tilde{\chi}_1^0$ plus soft radiation and where both of the $\tilde{\chi}_1^0$ decay to $Z + \tilde{G}$.

Tn1n1D: electroweak pair and associated production of nearly mass-degenerate Higgsino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$, $\tilde{\chi}_2^0$.

Tn1n1E: electroweak pair and associated production of nearly mass-degenerate wino-like charginos $\tilde{\chi}_1^\pm$ and neutralinos $\tilde{\chi}_1^0$.

Tn1n2A: electroweak associated production of nearly mass-degenerate neutralinos $\tilde{\chi}_1^0$ and $\tilde{\chi}_2^0$, where the $\tilde{\chi}_2^0$ always decays to $\gamma + \tilde{G}$ and $\tilde{\chi}_1^0$ 50% of the time to $H + \tilde{G}$ and 50 % of the time to $Z + \tilde{G}$.

Tn2n3A: electroweak associated production of mass-degenerate neutralinos $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$, where $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ decay through intermediate sleptons to $l^+l^- \tilde{\chi}_1^0$ and where the slepton mass is 5%, 25%, 50%, 75% and 95% of the $\tilde{\chi}_2^0$ mass.

Tn2n3B: electroweak associated production of mass-degenerate neutralinos $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$, where $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ decay through intermediate sleptons to $l^+l^- \tilde{\chi}_1^0$ and where $m_{\tilde{l}} = (m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})/2$.

TWinoBinoA: electroweak pair production of mass-degenerate wino-like doublet ($\tilde{\chi}_2^0, \tilde{\chi}_1^\pm$) (including all pair-production mechanisms) decaying into a bino singlet ($\tilde{\chi}_1^0$). Decays happen via Standard Model bosons, assumed to decay via hadrons.

TWinoHinoA: electroweak pair production of mass-degenerate wino-like doublet ($\tilde{\chi}_3^0, \tilde{\chi}_2^\pm$) (including all possible pair-production mechanisms) decaying into a quasi-mass-degenerate Higgsino triplet ($\tilde{\chi}_1^0, \tilde{\chi}_2^0, \tilde{\chi}_1^\pm$). Decays happen via Standard Model bosons, assumed to decay via hadrons.

THinoBinoA: electroweak pair production of quasi-mass-degenerate higgsino-like triplet ($\tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_1^\pm$) (including all possible pair-production mechanisms) decaying into a bino singlet ($\tilde{\chi}_1^0$). Decays happen via Standard Model bosons, assumed to decay via hadrons.

THinoWinoA: electroweak pair production of quasi-mass-degenerate higgsino-like triplet ($\tilde{\chi}_2^0, \tilde{\chi}_2^0, \tilde{\chi}_2^\pm$) (including all possible pair-production mechanisms) decaying into a mass-degenerate wino doublet ($\tilde{\chi}_1^0, \tilde{\chi}_1^\pm$). Decays happen via Standard Model bosons, assumed to decay via hadrons.

$\tilde{\chi}_1^0$ is often assumed to be the lightest supersymmetric particle (LSP). See also the $\tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0$ section below.

NODE=S046PHO

We have divided the $\tilde{\chi}_1^0$ listings below into five sections:

- 1) Accelerator limits for stable $\tilde{\chi}_1^0$,
- 2) Bounds on $\tilde{\chi}_1^0$ from dark matter searches,
- 3) $\tilde{\chi}_1^0 - p$ elastic cross section (spin-dependent, spin-independent interactions),
- 4) Other bounds on $\tilde{\chi}_1^0$ from astrophysics and cosmology, and
- 5) Unstable $\tilde{\chi}_1^0$ (Lightest Neutralino) mass limit.

———— Accelerator limits for stable $\tilde{\chi}_1^0$ ————

NODE=S046PHA

Unless otherwise stated, results in this section assume spectra, production rates, decay modes, and branching ratios as evaluated in the MSSM, with gaugino and sfermion mass unification at the GUT scale. These papers generally study production of $\tilde{\chi}_i^0 \tilde{\chi}_j^0$ ($i \geq 1, j \geq 2$), $\tilde{\chi}_1^+ \tilde{\chi}_1^-$, and (in the case of hadronic collisions) $\tilde{\chi}_1^+ \tilde{\chi}_2^0$ pairs. The mass limits on $\tilde{\chi}_1^0$ are either direct, or follow indirectly from the constraints set by the non-observation of $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ states on the gaugino and higgsino MSSM parameters M_2 and μ . In some cases, information is used from the nonobservation of slepton decays.

NODE=S046PHA

Obsolete limits obtained from e^+e^- collisions up to $\sqrt{s}=184$ GeV have been removed from this compilation and can be found in the 2000 Edition (The European Physical Journal **C15** 1 (2000)) of this Review.

$$\Delta m = m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}.$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 0.5–4.29	95	1 LEES	23C BABR	$B +$ charged track, RPV $B \rightarrow \tilde{\chi}_1^0 p, \lambda''_{113}$ of order $10^{-7}-10^{-6}$
>150	95	2 AAD	22E ATLS	$t\tilde{\mu}_L$ production, RPV, $\tilde{\mu}_L \rightarrow$ $\mu\tilde{\chi}_1^0, \lambda'_{231} = 1, 200 \text{ GeV} <$ $m_{\tilde{\mu}_L} < 600 \text{ GeV}.$
none 125–175	95	3 TUMASYAN	22S CMS	2 same-sign e or μ , 3 or 4 leptons, Tn1n1A, $m_{\tilde{G}} = 1 \text{ GeV}$
none 125–415	95	3 TUMASYAN	22S CMS	2 same-sign e or μ , 3 or 4 leptons, Tn1n1B, $m_{\tilde{G}} = 1 \text{ GeV}$
none 100–625	95	3 TUMASYAN	22S CMS	2 same-sign e or μ , 3 or 4 leptons, Tn1n1C, $m_{\tilde{G}} = 1 \text{ GeV}$
none 175–1025	95	4 TUMASYAN	22V CMS	3, 4 b -tag jets or 2 large-radius jets, \cancel{E}_T ; Tn1n1A; $m_{\tilde{G}}=1 \text{ GeV}$
none 450–930	95	5 AAD	21AX ATLS	jets + large-R jets + \cancel{E}_T , Tn1n1C
none 200–320	95	6 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tn1n1D, RPV, λ''_{323} elec- troweakino decay, degenerate Higgsino triplet
none 200–370	95	6 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tn1n1E, RPV, λ''_{323} elec- troweakino decay, degenerate Wino doublet
> 40	95	7 DREINER	09 THEO	
		8 ABBIENDI	04H OPAL	all $\tan\beta, \Delta m > 5 \text{ GeV},$ $m_0 > 500 \text{ GeV}, A_0 = 0$
> 42.4	95	9 HEISTER	04 ALEP	all $\tan\beta, \text{ all } \Delta m, \text{ all } m_0$
> 39.2	95	10 ABDALLAH	03M DLPH	all $\tan\beta, m_{\tilde{\nu}} > 500 \text{ GeV}$
> 46	95	11 ABDALLAH	03M DLPH	all $\tan\beta, \text{ all } \Delta m, \text{ all } m_0$
> 32.5	95	12 ACCIARRI	00D L3	$\tan\beta > 0.7, \Delta m > 3 \text{ GeV}, \text{ all } m_0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		13 AAD	14K ATLS	
> 24		14 CALIBBI	13	thermal relic abundance, MSSM particle content

NODE=S046PHA;CHECK LIMITS

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

- ¹ LEES 23C search in 398 fb^{-1} of e^+e^- annihilations at 10.58 GeV for SUSY in events with a tagged B meson and one and only one charged track that must be consistent with the hypothesis of being a proton. The results are interpreted in an RPV SUSY model, where a neutralino is produced in the decay of a B meson into a neutralino and a proton with the RPV coupling λ''_{113} . A branching fraction upper limit is determined for the λ''_{113} coupling, divided by the relevant squark mass squared as a function of the neutralino mass, see their figure 6. They also search for a new dark sector antibaryon that could be produced in decays of B mesons. NODE=S046PHA;LINKAGE=L
- ² AAD 22E searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry by measuring the yield asymmetry between events containing $e^- \mu^+$ and those containing $e^+ \mu^-$. This was found in agreement with the standard model prediction of 1. Limits are set on the RPV production of $t \tilde{\mu}_L$ events with $\tilde{\mu}_L \rightarrow \mu \tilde{\chi}_1^0$ for various values of λ'_{231} , see their figures 6 and 7. NODE=S046PHA;LINKAGE=I
- ³ TUMASYAN 22S searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino pair production in events with three or four leptons, with up to two hadronically decaying τ leptons, or two same-sign light leptons (e or μ). No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tchi1n2B (in flavory-democratic and tau-enriched or -dominated scenarios), Tchi1n2E, Tchi1n2F, see their Figures 16–20, and on the mass of the higgsino-triplet $\tilde{\chi}_2^0$, $\tilde{\chi}_1^\pm$, and $\tilde{\chi}_1^0$ in the models Tn1n1A, Tn1n1B, and Tn1n1C, see their Figure 21. NODE=S046PHA;LINKAGE=H
- ⁴ TUMASYAN 22V searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino pair production with decay to two Higgs bosons H , with $H \rightarrow b\bar{b}$, resulting either in 4 resolved b -jets or two large-radius jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tn1n1A, see their Figures 11 and 12, or in a model where higgsino-like nearly mass degenerate $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ are pair produced and each decay to H and a bino-like $\tilde{\chi}_1^0$, see their Figure 13. Limits are also set on the gluino mass in the model Tglu1l, see their Figure 14. NODE=S046PHA;LINKAGE=J
- ⁵ AAD 21AX searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of electroweakinos decaying to the LSP via the emission of Standard Model bosons (Higgs, W , Z) decaying into hadrons. The final state in all cases characterised by the presence of \cancel{E}_T , jets, and large- R jets tagged according to the boson of interest. Different assumptions (Higgsino, Wino, Bino) are made for the pair produced electroweakinos and for the LSP multiplet. No significant excess above the Standard Model predictions is observed. Limits are set on the electroweakino masses as a function of the model parameters (in particular $m_{\tilde{\chi}_1^0}$). See Fig. 16. NODE=S046PHA;LINKAGE=G
- ⁶ AAD 21BF searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of gluinos, stops, electroweakinos decaying RPV either directly or indirectly via the LSP. The final state in all cases is one or two leptons, many jets (up to fifteen) and b -jets. Different models with different branching fractions of the gluino or stop follow from the assumptions on the nature of the electroweakinos. No significant excess above the Standard Model predictions is observed. Limits are set on the *gluino*, \tilde{t}_1 , electroweakino masses as a function of the $\tilde{\chi}_1^0$ mass in several scenarios of gluino, stop and electroweakino pair production. NODE=S046PHA;LINKAGE=F
- ⁷ DREINER 09 show that in the general MSSM with non-universal gaugino masses there exists no model-independent laboratory bound on the mass of the lightest neutralino. An essentially massless $\tilde{\chi}_1^0$ is allowed by the experimental and observational data, imposing some constraints on other MSSM parameters, including M_2 , μ and the slepton and squark masses. NODE=S046PHA;LINKAGE=DR
- ⁸ ABBIENDI 04H search for charginos and neutralinos in events with acoplanar leptons+jets and multi-jet final states in the 192–209 GeV data, combined with the results on leptonic final states from ABBIENDI 04. The results hold for a scan over the parameter space covering the region $0 < M_2 < 5000 \text{ GeV}$, $-1000 < \mu < 1000 \text{ GeV}$ and $\tan\beta$ from 1 to 40. This limit supersedes ABBIENDI 00H. NODE=S046PHA;LINKAGE=AN
- ⁹ HEISTER 04 data collected up to 209 GeV. Updates earlier analysis of selectrons from HEISTER 02E, includes a new analysis of charginos and neutralinos decaying into stau and uses results on charginos with initial state radiation from HEISTER 02J. The limit is based on the direct search for charginos and neutralinos, the constraints from the slepton search and the Higgs mass limits from HEISTER 02 using a top mass of 175 GeV, interpreted in a framework with universal gaugino and sfermion masses. Assuming the mixing in the stau sector to be negligible, the limit improves to 43.1 GeV. Under the assumption of MSUGRA with unification of the Higgs and sfermion masses, the limit improves to 50 GeV, and reaches 53 GeV for $A_0 = 0$. These limits include and update the results of BARATE 01. NODE=S046PHA;LINKAGE=HE
- ¹⁰ ABDALLAH 03M uses data from $\sqrt{s} = 192\text{--}208 \text{ GeV}$. A limit on the mass of $\tilde{\chi}_1^0$ is derived from direct searches for neutralinos combined with the chargino search. Neutralinos are searched in the production of $\tilde{\chi}_1^0 \tilde{\chi}_2^0$, $\tilde{\chi}_1^0 \tilde{\chi}_3^0$, as well as $\tilde{\chi}_2^0 \tilde{\chi}_3^0$ and $\tilde{\chi}_2^0 \tilde{\chi}_4^0$ giving rise to cascade decays, and $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ and $\tilde{\chi}_1^0 \tilde{\chi}_3^0$, followed by the decay $\tilde{\chi}_2^0 \rightarrow \tilde{\tau} \tau$. The results hold for the parameter space defined by values of $M_2 < 1 \text{ TeV}$, $|\mu| \leq 2 \text{ TeV}$ with the $\tilde{\chi}_1^0$ as LSP. The limit is obtained for $\tan\beta = 1$ and large m_0 , where $\tilde{\chi}_2^0 \tilde{\chi}_4^0$ and chargino NODE=S046PHA;LINKAGE=AD

pair production are important. If the constraint from Higgs searches is also imposed, the limit improves to 49.0 GeV in the m_h^{\max} scenario with $m_t=174.3$ GeV. These limits update the results of ABREU 00J.

- 11 ABDALLAH 03M uses data from $\sqrt{s} = 192\text{--}208$ GeV. An indirect limit on the mass of $\tilde{\chi}_1^0$ is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays and $\tilde{\tau}\tau$ final states), for charginos (for all Δm_{\pm}) and for sleptons, stop and sbottom. The results hold for the full parameter space defined by values of $M_2 < 1$ TeV, $|\mu| \leq 2$ TeV with the $\tilde{\chi}_1^0$ as LSP. Constraints from the Higgs search in the m_h^{\max} scenario assuming $m_t=174.3$ GeV are included. The limit is obtained for $\tan\beta \geq 5$ when stau mixing leads to mass degeneracy between $\tilde{\tau}_1$ and $\tilde{\chi}_1^0$ and the limit is based on $\tilde{\chi}_2^0$ production followed by its decay to $\tilde{\tau}_1\tau$. In the pathological scenario where m_0 and $|\mu|$ are large, so that the $\tilde{\chi}_2^0$ production cross section is negligible, and where there is mixing in the stau sector but not in stop nor sbottom, the limit is based on charginos with soft decay products and an ISR photon. The limit then degrades to 39 GeV. See Figs. 40–42 for the dependence of the limit on $\tan\beta$ and $m_{\tilde{\nu}}$. These limits update the results of ABREU 00W.
- 12 ACCIARRI 00D data collected at $\sqrt{s}=189$ GeV. The results hold over the full parameter space defined by $0.7 \leq \tan\beta \leq 60$, $0 \leq M_2 \leq 2$ TeV, $m_0 \leq 500$ GeV, $|\mu| \leq 2$ TeV. The minimum mass limit is reached for $\tan\beta=1$ and large m_0 . The results of slepton searches from ACCIARRI 99W are used to help set constraints in the region of small m_0 . The limit improves to 48 GeV for $m_0 \gtrsim 200$ GeV and $\tan\beta \gtrsim 10$. See their Figs. 6–8 for the $\tan\beta$ and m_0 dependence of the limits. Updates ACCIARRI 98F.
- 13 AAD 14K sets limits on the χ -nucleon spin-dependent and spin-independent cross sections out to $m_\chi = 10$ TeV.
- 14 CALIBBI 13 use the fact that if the relic abundance of $\tilde{\chi}_1^0$ does not overclose the universe, scalar lepton and Higgsino masses must be relatively small. Using 8 TeV ATLAS constraints on the scalar tau mass and on invisible Higgs decays, they estimate a lower bound for the $\tilde{\chi}_1^0$ mass.

NODE=S046PHA;LINKAGE=AL

NODE=S046PHA;LINKAGE=DL

NODE=S046PHA;LINKAGE=DA

NODE=S046PHA;LINKAGE=E

Bounds on $\tilde{\chi}_1^0$ from dark matter searches

These papers generally exclude regions in the $M_2 - \mu$ parameter plane assuming that $\tilde{\chi}_1^0$ is the dominant form of dark matter in the galactic halo. These limits are based on the lack of detection in laboratory experiments, telescopes, or by the absence of a signal in underground neutrino detectors. The latter signal is expected if $\tilde{\chi}_1^0$ accumulates in the Sun or the Earth and annihilates into high-energy ν 's.

NODE=S046PHB

NODE=S046PHB

NODE=S046PHB

VALUE DOCUMENT ID TECN

••• We do not use the following data for averages, fits, limits, etc. •••

1	ABE	23B	MGIC
2	FOSTER	23	FLAT
3	GUO	23A	ICCB
4	ABBASI	22B	ICCB
5	ABDALLA	22	HESS
6	ABDALLAH	21	HESS
7	ABAZAJIAN	20	FLAT
8	ABDALLAH	20	HESS
9	ABE	20G	SKAM
10	ALBERT	20	HAWC
11	ALBERT	20A	ANTR
12	ALBERT	20C	ANIC
13	ALVAREZ	20	FLAT
14	HOOF	20	FLAT
15	DI-MAURO	19	FLAT
16	JOHNSON	19	FLAT
17	LI	19D	FLAT
18	AHNEN	18	MGIC
19	ALBERT	18B	HAWC
20	ALBERT	18C	HAWC
21	AARTSEN	17	ICCB
22	AARTSEN	17A	ICCB
23	AARTSEN	17C	ICCB
24	ARCHAMBAU..	17	VRTS
25	ADRIAN-MAR..	16	ANTR
26	AHNEN	16	MGFL
27	AVRORIN	16	BAIK
28	CIRELLI	16	THEO

	28	LEITE	16	THEO	
	29	ACKERMANN	15	FLAT	
	30	ACKERMANN	15A	FLAT	
	31	ACKERMANN	15B	FLAT	
	32	BUCKLEY	15	THEO	
	33	CHOI	15	SKAM	
	34	ALEKSIC	14	MGIC	
	35	AVRORIN	14	BAIK	
	36	AARTSEN	13C	ICCB	
	37	BERGSTROM	13	COSM	
	38	BOLIEV	13	BAKS	
	37	JIN	13	ASTR	
	37	KOPP	13	COSM	
	39	ACKERMANN	10	FLAT	
	40	ACHTERBERG	06	AMND	
	41	ACKERMANN	06	AMND	
	42	DEBOER	06	RVUE	
	43	DESAI	04	SKAM	
	43	AMBROSIO	99	MCRO	
	44	LOSECCO	95	RVUE	
	45	MORI	93	KAMI	
	46	BOTTINO	92	COSM	
	47	BOTTINO	91	RVUE	
	48	GELMINI	91	COSM	
	49	KAMIONKOW	91	RVUE	
	50	MORI	91B	KAMI	
none 4–15 GeV	51	OLIVE	88	COSM	
1		ABE 23B sets limits on the dark matter annihilation cross section from line-like features in TeV gamma-rays in the direction of the Galactic center using the MAGIC stereoscopic telescope.			NODE=S046PHB;LINKAGE=YA
2		FOSTER 23 sets limits on the dark matter annihilation cross section from monochromatic gamma-rays in the inner Milky Way using 14 years of data from Fermi-LAT.			NODE=S046PHB;LINKAGE=CB
3		GUO 23A sets limits on the dark matter annihilation cross section from 10 years of IceCube muon-track data from 18 dwarf spheroidal galaxies.			NODE=S046PHB;LINKAGE=EB
4		ABBASI 22B presents 7 years of data from a search of neutrinos from dark matter annihilations in the sun using the DeepCore sub-array of IceCube. Annihilation cross section limits applies to dark matter masses between 5–100 GeV.			NODE=S046PHB;LINKAGE=VA
5		ABDALLA 22 uses gamma-ray observations in the Galactic center to constrain the dark matter annihilation cross section for annihilations into WW and $\tau\tau$ for dark matter masses between 200 GeV to 70 TeV. This updates ABDALLAH 18.			NODE=S046PHB;LINKAGE=XA
6		ABDALLAH 21 places constraints on the dark matter annihilation cross section for annihilations into gamma-rays from the dwarf irregular galaxy WLM for masses between 0.15 to 10 TeV.			NODE=S046PHB;LINKAGE=UA
7		ABAZAJIAN 20 sets constraints on the dark matter annihilation from gamma-ray searches from Fermi LAT observations of the Galactic center.			NODE=S046PHB;LINKAGE=NA
8		ABDALLAH 20 places constraints on the dark matter annihilation cross section for annihilations into gamma-rays from Milky Way dwarf galaxy satellites for masses between 0.2 to 40 TeV.			NODE=S046PHB;LINKAGE=RA
9		ABE 20G is based on SuperKamiokande data taken from 1996 to 2016 searching for neutrinos produced from dark matter annihilations in the galactic center or halo. They place constraints on the dark matter-nucleon scattering cross section for dark matter masses between 1 GeV and 10 TeV.			NODE=S046PHB;LINKAGE=SA
10		ALBERT 20 sets limits on the annihilation cross section of dark matter with mass between 1 and 100 TeV from gamma-ray observations of the local dwarf spheroidal galaxies.			NODE=S046PHB;LINKAGE=MA
11		ALBERT 20A set limits on the dark matter annihilation cross section from neutrinos observations in the Galactic center using 11 years of ANTARES data.			NODE=S046PHB;LINKAGE=OA
12		ALBERT 20C set limits on the dark matter annihilation cross section from neutrinos observations in the Galactic center combining Antares and IceCube data.			NODE=S046PHB;LINKAGE=TA
13		ALVAREZ 20 set limits on the dark matter annihilation from gamma-ray searches from Fermi LAT observations in the directions of dwarf spheroidal galaxies.			NODE=S046PHB;LINKAGE=PA
14		HOOF 20 set limits on the dark matter annihilation from gamma-ray searches from Fermi LAT observations in the directions of dwarf spheroidal galaxies.			NODE=S046PHB;LINKAGE=QA
15		DI-MAURO 19 sets limits on the dark matter annihilation from gamma-ray searches in M31 and M33 galaxies using Fermi LAT data.			NODE=S046PHB;LINKAGE=JA
16		JOHNSON 19 sets limits on p-wave dark matter annihilations in the galactic center using Fermi data.			NODE=S046PHB;LINKAGE=KA
17		LI 19D sets limits on dark matter annihilation cross sections searching for line-like signals in the all-sky Fermi data.			NODE=S046PHB;LINKAGE=LA
18		AHNEN 18 uses observations of the dwarf satellite galaxy Ursa Major II to obtain upper limits on annihilation cross sections for dark matter in various channels for masses between 0.1–100 TeV.			NODE=S046PHB;LINKAGE=GA
19		ALBERT 18B sets limits on the annihilation cross section of dark matter with mass between 1 and 100 TeV from gamma-ray observations of the Andromeda galaxy.			NODE=S046PHB;LINKAGE=HA

- 20 ALBERT 18C sets limits on the spin-dependent coupling of dark matter to protons from dark matter annihilation in the Sun. NODE=S046PHB;LINKAGE=IA
- 21 AARTSEN 17 is based on data collected during 327 days of detector livetime with IceCube. They looked for interactions of ν 's resulting from neutralino annihilations in the Earth over a background of atmospheric neutrinos and set 90% CL limits on the spin independent neutralino-proton cross section for neutralino masses in the range 10–10000 GeV. NODE=S046PHB;LINKAGE=X
- 22 AARTSEN 17A is based on data collected during 532 days of livetime with the IceCube 86-string detector including the DeepCore sub-array. They looked for interactions of ν 's from neutralino annihilations in the Sun over a background of atmospheric neutrinos and set 90% CL limits on the spin dependent neutralino-proton cross section for neutralino masses in the range 10–10000 GeV. This updates AARTSEN 16C. NODE=S046PHB;LINKAGE=Y
- 23 AARTSEN 17C is based on 1005 days of running with the IceCube detector. They set a limit on the annihilation cross section for dark matter with masses between 10–1000 GeV annihilating in the Galactic center assuming an NFW profile. The limit is of $1.2 \times 10^{23} \text{ cm}^3 \text{ s}^{-1}$ in the $\tau^+ \tau^-$ channel. Supercedes AARTSEN 15E. NODE=S046PHB;LINKAGE=EA
- 24 ARCHAMBAULT 17 performs a joint statistical analysis of four dwarf galaxies with VERITAS looking for gamma-ray emission from neutralino annihilation. They set limits on the neutralino annihilation cross section. NODE=S046PHB;LINKAGE=DA
- 25 ADRIAN-MARTINEZ 16 is based on data from the ANTARES neutrino telescope. They looked for interactions of ν 's from neutralino annihilations in the Sun over a background of atmospheric neutrinos and set 90% CL limits on the muon neutrino flux. They also obtain limits on the spin dependent and spin independent neutralino-proton cross section for neutralino masses in the range 50 to 5,000 GeV. This updates ADRIAN-MARTINEZ 13. NODE=S046PHB;LINKAGE=R
- 26 AHNEN 16 combines 158 hours of Segue 1 observations with MAGIC with 6 year observations of 15 dwarf satellite galaxies by Fermi-LAT to set limits on annihilation cross sections for dark matter masses between 10 GeV and 100 TeV. NODE=S046PHB;LINKAGE=T
- 27 AVRORIN 16 is based on 2.76 years with Lake Baikal neutrino telescope. They derive 90% upper limits on the annihilation cross section from dark matter annihilations in the Galactic center. NODE=S046PHB;LINKAGE=U
- 28 CIRELLI 16 and LEITE 16 derive bounds on the annihilation cross section from radio observations. NODE=S046PHB;LINKAGE=W
- 29 ACKERMANN 15 is based on 5.8 years of data with Fermi-LAT and search for monochromatic gamma-rays in the energy range of 0.2–500 GeV from dark matter annihilations. This updates ACKERMANN 13A. NODE=S046PHB;LINKAGE=D
- 30 ACKERMANN 15A is based on 50 months of data with Fermi-LAT and search for dark matter annihilation signals in the isotropic gamma-ray background as well as galactic subhalos in the energy range of a few GeV to a few tens of TeV. NODE=S046PHB;LINKAGE=E
- 31 ACKERMANN 15B is based on 6 years of data with Fermi-LAT observations of Milky Way dwarf spheroidal galaxies. Set limits on the annihilation cross section from $m_\chi = 2 \text{ GeV}$ to 10 TeV. This updates ACKERMANN 14. NODE=S046PHB;LINKAGE=L
- 32 BUCKLEY 15 is based on 5 years of Fermi-LAT data searching for dark matter annihilation signals from Large Magellanic Cloud. NODE=S046PHB;LINKAGE=G
- 33 CHOI 15 is based on 3903 days of SuperKamiokande data searching for neutrinos produced from dark matter annihilations in the sun. They place constraints on the dark matter-nucleon scattering cross section for dark matter masses between 4–200 GeV. NODE=S046PHB;LINKAGE=I
- 34 ALEKSIC 14 is based on almost 160 hours of observations of Segue 1 satellite dwarf galaxy using the MAGIC telescopes between 2011 and 2013. Sets limits on the annihilation cross section out to $m_\chi = 10 \text{ TeV}$. NODE=S046PHB;LINKAGE=AL
- 35 AVRORIN 14 is based on almost 2.76 years with Lake Baikal neutrino telescope. They derive 90% upper limits on the fluxes of muons and muon neutrinos from dark matter annihilations in the Sun. NODE=S046PHB;LINKAGE=AV
- 36 AARTSEN 13C is based on data collected during 339.8 effective days with the IceCube 59-string detector. They looked for interactions of ν_μ 's from neutralino annihilations in nearby galaxies and galaxy clusters. They obtain limits on the neutralino annihilation cross section for neutralino masses in the range 30–100,000 GeV. NODE=S046PHB;LINKAGE=RT
- 37 BERGSTROM 13, JIN 13, and KOPP 13 derive limits on the mass and annihilation cross section using AMS-02 data. JIN 13 also sets a limit on the lifetime of the dark matter particle. NODE=S046PHB;LINKAGE=KO
- 38 BOLIEV 13 is based on data collected during 24.12 years of live time with the Bakson Underground Scintillator Telescope. They looked for interactions of ν_μ 's from neutralino annihilations in the Sun over a background of atmospheric neutrinos and set 90% CL limits on the muon flux. They also obtain limits on the spin dependent and spin independent neutralino-proton cross section for neutralino masses in the range 10–1000 GeV. NODE=S046PHB;LINKAGE=BO
- 39 ACKERMANN 10 place upper limits on the annihilation cross section with $b\bar{b}$ or $\mu^+ \mu^-$ final states. NODE=S046PHB;LINKAGE=AC
- 40 ACHTERBERG 06 is based on data collected during 421.9 effective days with the AMANDA detector. They looked for interactions of ν_μ 's from the centre of the Earth over a background of atmospheric neutrinos and set 90% CL limits on the muon flux. Their limit is compared with the muon flux expected from neutralino annihilations into $W^+ W^-$ and $b\bar{b}$ at the centre of the Earth for MSSM parameters compatible with the relic dark matter density, see their Fig. 7. NODE=S046PHB;LINKAGE=AH
- 41 ACKERMANN 06 is based on data collected during 143.7 days with the AMANDA-II detector. They looked for interactions of ν_μ 's from the Sun over a background of atmospheric neutrinos and set 90% CL limits on the muon flux. Their limit is compared

- with the muon flux expected from neutralino annihilations into $W^+ W^-$ in the Sun for SUSY model parameters compatible with the relic dark matter density, see their Fig. 3.
- 42 DEBOER 06 interpret an excess of diffuse Galactic gamma rays observed with the EGRET satellite as originating from π^0 decays from the annihilation of neutralinos into quark jets. They analyze the corresponding parameter space in a supergravity inspired MSSM model with radiative electroweak symmetry breaking, see their Fig. 3 for the preferred region in the $(m_0, m_{1/2})$ plane of a scenario with large $\tan\beta$. NODE=S046PHB;LINKAGE=DB
- 43 AMBROSIO 99 and DESAI 04 set new neutrino flux limits which can be used to limit the parameter space in supersymmetric models based on neutralino annihilation in the Sun and the Earth. NODE=S046PHB;LINKAGE=BI
- 44 LOSECCO 95 reanalyzed the IMB data and places lower limit on $m_{\tilde{\chi}_1^0}$ of 18 GeV if the LSP is a photino and 10 GeV if the LSP is a higgsino based on LSP annihilation in the sun producing high-energy neutrinos and the limits on neutrino fluxes from the IMB detector. NODE=S046PHB;LINKAGE=LC
- 45 MORI 93 excludes some region in $M_2-\mu$ parameter space depending on $\tan\beta$ and lightest scalar Higgs mass for neutralino dark matter $m_{\tilde{\chi}_1^0} > m_W$, using limits on upgoing muons produced by energetic neutrinos from neutralino annihilation in the Sun and the Earth. NODE=S046PHB;LINKAGE=P
- 46 BOTTINO 92 excludes some region $M_2-\mu$ parameter space assuming that the lightest neutralino is the dark matter, using upgoing muons at Kamiokande, direct searches by Ge detectors, and by LEP experiments. The analysis includes top radiative corrections on Higgs parameters and employs two different hypotheses for nucleon-Higgs coupling. Effects of rescaling in the local neutralino density according to the neutralino relic abundance are taken into account. NODE=S046PHB;LINKAGE=S
- 47 BOTTINO 91 excluded a region in $M_2-\mu$ plane using upgoing muon data from Kamioka experiment, assuming that the dark matter surrounding us is composed of neutralinos and that the Higgs boson is not too heavy. NODE=S046PHB;LINKAGE=ZA
- 48 GELMINI 91 exclude a region in $M_2-\mu$ plane using dark matter searches. NODE=S046PHB;LINKAGE=S4
- 49 KAMIONKOWSKI 91 excludes a region in the $M_2-\mu$ plane using IMB limit on upgoing muons originated by energetic neutrinos from neutralino annihilation in the sun, assuming that the dark matter is composed of neutralinos and that $m_{H_1^0} \lesssim 50$ GeV. See Fig. 8 in the paper. NODE=S046PHB;LINKAGE=B
- 50 MORI 91B exclude a part of the region in the $M_2-\mu$ plane with $m_{\tilde{\chi}_1^0} \lesssim 80$ GeV using a limit on upgoing muons originated by energetic neutrinos from neutralino annihilation in the earth, assuming that the dark matter surrounding us is composed of neutralinos and that $m_{H_1^0} \lesssim 80$ GeV. NODE=S046PHB;LINKAGE=BB
- 51 OLIVE 88 result assumes that photinos make up the dark matter in the galactic halo. Limit is based on annihilations in the sun and is due to an absence of high energy neutrinos detected in underground experiments. The limit is model dependent. NODE=S046PHB;LINKAGE=F

———— $\tilde{\chi}_1^0$ -p elastic cross section ————

Experimental results on the $\tilde{\chi}_1^0$ -p elastic cross section are evaluated at $m_{\tilde{\chi}_1^0}=100$ GeV. The experimental results on the cross section are often mass dependent. Therefore, the mass and cross section results are also given where the limit is strongest, when appropriate. Results are quoted separately for spin-dependent interactions (based on an effective 4-Fermi Lagrangian of the form $\bar{\chi}\gamma^\mu\gamma^5\chi\bar{q}\gamma_\mu\gamma^5q$) and spin-independent interactions ($\bar{\chi}\chi\bar{q}q$). For calculational details see GRIEST 88B, ELLIS 88D, BARBIERI 89C, DREES 93B, ARNOWITT 96, BERGSTROM 96, and BAER 97 in addition to the theory papers listed in the Tables. For a description of the theoretical assumptions and experimental techniques underlying most of the listed papers, see the review on "Dark matter" in this "Review of Particle Physics," and references therein. Most of the following papers use galactic halo and nuclear interaction assumptions from (LEWIN 96).

NODE=S046303

NODE=S046303

Spin-dependent interactions

VALUE (pb)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 1.9 × 10 ⁻⁴	90	1 AALBERS	23 LZ	Xe
< 3.3 × 10 ⁻⁴	90	2 APRILE	23A XENT	Xe
< 2 × 10 ⁻⁴	90	3 HUANG	22 PNDX	Xe
< 4 × 10 ⁻⁵	90	4 AMOLE	19 PICO	C ₃ F ₈
< 5 × 10 ⁻⁴	90	5 APRILE	19A XE1T	Xe
< 8 × 10 ⁻⁴	90	6 AKERIB	17A LUX	Xe
< 0.28	90	7 BATTAT	17 DRFT	CS ₂ ; CF ₄
< 0.027	90	8 BEHNKE	17 PICA	C ₄ F ₁₀
< 5 × 10 ⁻⁴	90	9 AMOLE	16 PICO	CF ₃ I
< 6.8 × 10 ⁻³	90	10 APRILE	16B X100	Xe
< 6.3 × 10 ⁻³	90	11 FELIZARDO	14 SMPL	C ₂ ClF ₅

NODE=S046DM1

NODE=S046DM1

< 0.01	90	12 AKIMOV	12 ZEP3	Xe
< 7 × 10 ⁻³		13 BEHNKE	12 COUP	CF ₃ I
< 8.5 × 10 ⁻³		14 FELIZARDO	12 SMPL	C ₂ ClF ₅
< 0.016	90	15 KIM	12 KIMS	CsI
5 × 10 ⁻¹⁰ to 10 ⁻⁵	95	16 BUCHMUEL...	11B THEO	
< 1	90	17 ANGLE	08A XE10	Xe
< 0.055		18 BEDNYAKOV	08 HDMS	Ge
< 0.33	90	19 BEHNKE	08 COUP	CF ₃ I
< 5		20 AKERIB	06 CDMS	Ge
< 2		21 SHIMIZU	06A CNTR	CaF ₂
< 0.4		22 ALNER	05 NAIA	NaI Spin Dep.
< 2		23 BARNABE-HE.	05 PICA	C
2 × 10 ⁻¹¹ to 1 × 10 ⁻⁴		24 ELLIS	04 THEO	μ > 0
< 0.8		25 AHMED	03 NAIA	NaI Spin Dep.
< 40		26 TAKEDA	03 BOLO	NaF Spin Dep.
< 10		27 ANGLOHER	02 CRES	Saphire
8 × 10 ⁻⁷ to 2 × 10 ⁻⁵		28 ELLIS	01C THEO	tanβ ≤ 10
< 3.8		29 BERNABEI	00D DAMA	Xe
< 0.8		SPOONER	00 UKDM	NaI
< 4.8		30 BELLI	99C DAMA	F
<100		31 OOTANI	99 BOLO	LiF
< 0.6		BERNABEI	98C DAMA	Xe
< 5		30 BERNABEI	97 DAMA	F

- 1 The strongest upper limit is 4.2×10^{-5} pb at 32 GeV. The limit for scattering on neutrons is 4×10^{-6} pb at 100 GeV and is 1.5×10^{-6} pb at 30 GeV. NODE=S046DM1;LINKAGE=S
- 2 The strongest upper limit is 1.4×10^{-4} pb at 28 GeV. The limit for scattering on neutrons is 1.1×10^{-5} pb at 100 GeV and is 4.3×10^{-6} pb at 28 GeV. NODE=S046DM1;LINKAGE=T
- 3 The strongest limit is $< 1.7 \times 10^{-4}$ pb at $m_\chi = 40$ GeV. This updates FU 17 and XIA 19A. NODE=S046DM1;LINKAGE=R
- 4 The strongest limit is $< 3.2 \times 10^{-5}$ pb at $m_\chi = 25$ GeV. This updates AMOLE 17. NODE=S046DM1;LINKAGE=O
- 5 The strongest limit is $< 2 \times 10^{-4}$ pb at $m_\chi = 30$ GeV. For scatterings on neutrons, the strongest limit is $< 6.3 \times 10^{-6}$ at $m_\chi = 30$ GeV. NODE=S046DM1;LINKAGE=P
- 6 The strongest limit is 5×10^{-4} pb at $m_\chi = 35$ GeV. The limit for scattering on neutrons is 3×10^{-5} pb at 100 GeV and is 1.6×10^{-5} pb at 35 GeV. This updates AKERIB 16A. NODE=S046DM1;LINKAGE=H
- 7 Directional recoil detector. This updates DAW 12. NODE=S046DM1;LINKAGE=L
- 8 This result updates ARCHAMBAULT 12. The strongest limit is 0.013 pb at $m_\chi = 20$ GeV. NODE=S046DM1;LINKAGE=M
- 9 The strongest limit is 5×10^{-4} pb at $m_\chi = 80$ GeV. NODE=S046DM1;LINKAGE=D
- 10 The strongest limit is 5.2×10^{-3} pb at 50 GeV. The limit for scattering on neutrons is 2.8×10^{-4} pb at 100 GeV and the strongest limit is 2.0×10^{-4} pb at 50 GeV. This updates APRILE 13. NODE=S046DM1;LINKAGE=G
- 11 The strongest limit is 0.0043 pb and occurs at $m_\chi = 35$ GeV. FELIZARDO 14 also presents limits for the scattering on neutrons. At $m_\chi = 100$ GeV, the upper limit is 0.13 pb and the strongest limit is 0.066 pb at $m_\chi = 35$ GeV. NODE=S046DM1;LINKAGE=FL
- 12 This result updates LEBEDENKO 09A. The strongest limit is 8×10^{-3} pb at $m_\chi = 50$ GeV. Limit applies to the neutralino neutron elastic cross section. NODE=S046DM1;LINKAGE=KI
- 13 The strongest limit is 6×10^{-3} at $m_\chi = 60$ GeV. NODE=S046DM1;LINKAGE=BN
- 14 The strongest limit is 5.7×10^{-3} at $m_\chi = 35$ GeV. NODE=S046DM1;LINKAGE=FE
- 15 This result updates LEE 07A. The strongest limit is at $m_\chi = 80$ GeV. NODE=S046DM1;LINKAGE=KM
- 16 Predictions for the spin-dependent elastic cross section based on a frequentist approach to electroweak observables in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046DM1;LINKAGE=MU
- 17 The strongest limit is 0.6 pb and occurs at $m_\chi = 30$ GeV. The limit for scattering on neutrons is 0.01 pb at $m_\chi = 100$ GeV, and the strongest limit is 0.0045 pb at $m_\chi = 30$ GeV. NODE=S046DM1;LINKAGE=AG
- 18 Limit applies to neutron elastic cross section. NODE=S046DM1;LINKAGE=BE
- 19 The strongest upper limit is 0.25 pb and occurs at $m_\chi \simeq 40$ GeV. NODE=S046DM1;LINKAGE=BU
- 20 The strongest upper limit is 4 pb and occurs at $m_\chi \simeq 60$ GeV. The limit on the neutron spin-dependent elastic cross section is 0.07 pb. This latter limit is improved in AHMED 09, where a limit of 0.02 pb is obtained at $m_\chi = 100$ GeV. The strongest limit in AHMED 09 is 0.018 pb and occurs at $m_\chi = 60$ GeV. NODE=S046DM1;LINKAGE=AK
- 21 The strongest upper limit is 1.2 pb and occurs at $m_\chi \simeq 40$ GeV. The limit on the neutron spin-dependent cross section is 35 pb. NODE=S046DM1;LINKAGE=SH
- 22 The strongest upper limit is 0.35 pb and occurs at $m_\chi \simeq 60$ GeV. NODE=S046DM1;LINKAGE=AN
- 23 The strongest upper limit is 1.2 pb and occurs $m_\chi \simeq 30$ GeV. NODE=S046DM1;LINKAGE=BA

- 24 ELLIS 04 calculates the $\chi\rho$ elastic scattering cross section in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry, but without universal scalar masses. In the case of universal squark and slepton masses, but non-universal Higgs masses, the limit becomes 2×10^{-4} , see ELLIS 03E.
- 25 The strongest upper limit is 0.75 pb and occurs at $m_\chi \approx 70$ GeV.
- 26 The strongest upper limit is 30 pb and occurs at $m_\chi \approx 20$ GeV.
- 27 The strongest upper limit is 8 pb and occurs at $m_\chi \approx 30$ GeV.
- 28 ELLIS 01C calculates the χ - p elastic scattering cross section in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. In models with nonuniversal Higgs masses, the upper limit to the cross section is 6×10^{-4} .
- 29 The strongest upper limit is 3 pb and occurs at $m_\chi \simeq 60$ GeV. The limits are for inelastic scattering $X^0 + {}^{129}\text{Xe} \rightarrow X^0 + {}^{129}\text{Xe}^*$ (39.58 keV).
- 30 The strongest upper limit is 4.4 pb and occurs at $m_\chi \simeq 60$ GeV.
- 31 The strongest upper limit is about 35 pb and occurs at $m_\chi \simeq 15$ GeV.

NODE=S046DM1;LINKAGE=EI

NODE=S046DM1;LINKAGE=AH

NODE=S046DM1;LINKAGE=TA

NODE=S046DM1;LINKAGE=FF

NODE=S046DM1;LINKAGE=EL

NODE=S046DM1;LINKAGE=A

NODE=S046DM1;LINKAGE=BL

NODE=S046DM1;LINKAGE=I

NODE=S046DM2

NODE=S046DM2

Spin-independent interactions

VALUE (pb)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$< 3 \times 10^{-11}$	90	1 AALBERS	23 LZ	Xe
$< 6.1 \times 10^{-11}$	90	2 APRILE	23A XENT	Xe
$< 6.5 \times 10^{-11}$	90	3 MENG	21B PNDX	Xe
$< 5 \times 10^{-10}$	90	4 WANG	20G PNDX	Xe
$< 2.5 \times 10^{-8}$	90	5 ABE	19 XMAS	Xe
$< 3.9 \times 10^{-9}$	90	6 AJAJ	19 DEAP	Ar
$< 2 \times 10^{-8}$	90	7 AMOLE	19 PICO	C_3F_8
$< 2.25 \times 10^{-6}$	90	8 ADHIKARI	18 C100	Nal
$< 1.14 \times 10^{-8}$	90	9 AGNES	18A DS50	Ar
$< 1.6 \times 10^{-8}$	90	10 AGNESE	18A CDMS	Ge
$< 9 \times 10^{-11}$	90	11 APRILE	18 XE1T	Xe
$< 1.8 \times 10^{-10}$	90	12 AKERIB	17 LUX	Xe
$< 1.5 \times 10^{-9}$	90	13 APRILE	16B X100	Xe
$< 1.5 \times 10^{-9}$	90	14 AKERIB	14 LUX	Xe
10^{-11} - 10^{-7}	95	15 BUCHMUEL...	14A THEO	
$< 4.6 \times 10^{-6}$	90	16 FELIZARDO	14 SMPL	C_2ClF_5
10^{-11} - 10^{-8}	95	17 ROSZKOWSKI	14 THEO	
$< 2.2 \times 10^{-6}$	90	18 AGNESE	13 CDMS	Si
$< 5 \times 10^{-8}$	90	19 AKIMOV	12 ZEP3	Xe
1.6×10^{-6} ; 3.7×10^{-5}		20 ANGLOHER	12 CRES	CaWO_4
3×10^{-12} to 3×10^{-9}	95	21 BECHTLE	12 THEO	
$< 1.6 \times 10^{-7}$		22 BEHNKE	12 COUP	CF_3I
$< 2.3 \times 10^{-7}$	90	23 KIM	12 KIMS	Csl
$< 3.3 \times 10^{-8}$	90	24 AHMED	11A	Ge
$< 4.4 \times 10^{-8}$	90	25 ARMENGAUD	11 EDE2	Ge
$< 1 \times 10^{-7}$	90	26 ANGLE	08 XE10	Xe
$< 1 \times 10^{-6}$	90	BENETTI	08 WARP	Ar
$< 7.5 \times 10^{-7}$	90	27 ALNER	07A ZEP2	Xe
$< 2 \times 10^{-7}$		28 AKERIB	06A CDMS	Ge
$< 90 \times 10^{-7}$		ALNER	05 NAIA	Nal Spin Indep.
$< 12 \times 10^{-7}$		29 ALNER	05A ZEPL	
$< 14 \times 10^{-7}$		SANGLARD	05 EDEL	Ge
$< 4 \times 10^{-7}$		30 AKERIB	04 CDMS	Ge
2×10^{-11} to 1.5×10^{-7}	95	31 BALTZ	04 THEO	
2×10^{-11} to 8×10^{-6}	32,33	ELLIS	04 THEO	$\mu > 0$
$< 5 \times 10^{-8}$		34 PIERCE	04A THEO	
$< 2 \times 10^{-5}$		35 AHMED	03 NAIA	Nal Spin Indep.
$< 3 \times 10^{-6}$		36 AKERIB	03 CDMS	Ge
2×10^{-13} to 2×10^{-7}		37 BAER	03A THEO	
$< 1.4 \times 10^{-5}$		38 KLAPDOR-K...	03 HDMS	Ge
$< 6 \times 10^{-6}$		39 ABRAMS	02 CDMS	Ge
1×10^{-12} to 7×10^{-6}		32 KIM	02B THEO	
$< 3 \times 10^{-5}$		40 MORALES	02B CSME	Ge
$< 1 \times 10^{-5}$		41 MORALES	02C IGEX	Ge
$< 1 \times 10^{-6}$		BALTZ	01 THEO	
$< 3 \times 10^{-5}$		42 BAUDIS	01 HDMS	Ge
$< 7 \times 10^{-6}$		43 BOTTINO	01 THEO	
$< 1 \times 10^{-8}$		44 CORSETTI	01 THEO	$\tan\beta \leq 25$

5×10^{-10} to 1.5×10^{-8}	45 ELLIS	01c THEO	$\tan\beta \leq 10$
$< 4 \times 10^{-6}$	44 GOMEZ	01 THEO	
2×10^{-10} to 1×10^{-7}	44 LAHANAS	01 THEO	
$< 3 \times 10^{-6}$	ABUSAIDI	00 CDMS	Ge, Si
$< 6 \times 10^{-7}$	46 ACCOMANDO	00 THEO	
	47 BERNABEI	00 DAMA	Nal
2.5×10^{-9} to 3.5×10^{-8}	48 FENG	00 THEO	$\tan\beta=10$
$< 1.5 \times 10^{-5}$	MORALES	00 IGEX	Ge
$< 4 \times 10^{-5}$	SPOONER	00 UKDM	Nal
$< 7 \times 10^{-6}$	BAUDIS	99 HDMO	^{76}Ge
$< 7 \times 10^{-6}$	BERNABEI	98c DAMA	Xe

- 1 The strongest upper limit is 9.2×10^{-12} pb at 36 GeV.
- 2 The strongest upper limit is 2.6×10^{-11} pb at 28 GeV.
- 3 Commissioning Run for PandaX-4T. The strongest limit is 3.8×10^{-11} pb at $m_\chi = 40$ GeV.
- 4 WANG 20G strongest limit is 2.2×10^{-10} pb at 30 GeV using 132 ton-day full exposure of PandaX-II. This updates CUI 17A, though the results here provide weaker constraints.
- 5 The strongest upper limit is 2.2×10^{-8} pb at 60 GeV.
- 6 This updates AMAUDRUZ 18.
- 7 This updates AMOLE 16.
- 8 The strongest limit is 2.05×10^{-6} at $m = 60$ GeV.
- 9 The strongest limit is 1.09×10^{-8} pb at $m_\chi = 126$ GeV. This updates AGNES 15.
- 10 The strongest limit is 1.0×10^{-8} pb at $m_\chi = 46$ GeV. This updates AGNESE 15B.
- 11 Based on 278.8 days of data collection. The strongest limit is 4.1×10^{-11} pb at $m_\chi = 30$ GeV. This updates APRILE 17G.
- 12 AKERIB 17. The strongest limit is 1.1×10^{-10} pb at 50 GeV. This updates AKERIB 16.
- 13 The strongest limit is 1.1×10^{-9} pb at 50 GeV. This updates APRILE 12.
- 14 The strongest upper limit is 7.6×10^{-10} at $m_\chi = 33$ GeV.
- 15 Predictions for the spin-independent elastic cross section based on a frequentist approach to electroweak observables in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using the 20 fb^{-1} 8 TeV and the 5 fb^{-1} 7 TeV LHC data and the LUX data.
- 16 The strongest limit is 3.6×10^{-6} pb and occurs at $m_\chi = 35$ GeV. Felizardo 2014 updates Felizardo 2012.
- 17 Predictions for the spin-independent elastic cross section based on a Bayesian approach to electroweak observables in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using the 20 fb^{-1} LHC data and LUX.
- 18 AGNESE 13 presents 90% CL limits on the elastic cross section for masses in the range 7–100 GeV using the Si based detector. The strongest upper limit is 1.8×10^{-6} pb at $m_\chi = 50$ GeV. This limit is improved to 7×10^{-7} pb in AGNESE 13A.
- 19 This result updates LEBEDENKO 09. The strongest limit is 3.9×10^{-8} pb at $m_\chi = 52$ GeV.
- 20 ANGLOHER 12 presents results of 730 kg days from the CRESST-II dark matter detector. They find two maxima in the likelihood function corresponding to best fit WIMP masses of 25.3 and 11.6 GeV with elastic cross sections of 1.6×10^{-6} and 3.7×10^{-5} pb respectively, see their Table 4. The statistical significance is more than 4σ . ANGLOHER 12 updates ANGLOHER 09
- 21 Predictions for the spin-independent elastic cross section based on a frequentist approach to electroweak observables in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using the 5 fb^{-1} LHC data and XENON100.
- 22 The strongest limit is 1.4×10^{-7} at $m_\chi = 60$ GeV.
- 23 This result updates LEE 07A. The strongest limit is 2.1×10^{-7} at $m_\chi = 70$ GeV.
- 24 AHMED 11A gives combined results from CDMS and EDELWEISS. The strongest limit is at $m_\chi = 90$ GeV.
- 25 ARMENGAUD 11 updates result of ARMENGAUD 10. Strongest limit at $m_\chi = 85$ GeV.
- 26 The strongest upper limit is 5.1×10^{-8} pb and occurs at $m_\chi \simeq 30$ GeV. The values quoted here are based on the analysis performed in ANGLE 08 with the update from SORENSEN 09.
- 27 The strongest upper limit is 6.6×10^{-7} pb and occurs at $m_\chi \simeq 65$ GeV.
- 28 AKERIB 06A updates the results of AKERIB 05. The strongest upper limit is 1.6×10^{-7} pb and occurs at $m_\chi \approx 60$ GeV.
- 29 The strongest upper limit is also close to 1.0×10^{-6} pb and occurs at $m_\chi \simeq 70$ GeV. BENOIT 06 claim that the discrimination power of ZEPLIN-I measurement (ALNER 05A) is not reliable enough to obtain a limit better than 1×10^{-3} pb. However, SMITH 06 do not agree with the criticisms of BENOIT 06.
- 30 AKERIB 04 is incompatible with BERNABEI 00 most likely value, under the assumption of standard WIMP-halo interactions. The strongest upper limit is 4×10^{-7} pb and occurs at $m_\chi \simeq 60$ GeV.

NODE=S046DM2;LINKAGE=IA
 NODE=S046DM2;LINKAGE=JA
 NODE=S046DM2;LINKAGE=HA
 NODE=S046DM2;LINKAGE=GA
 NODE=S046DM2;LINKAGE=W
 NODE=S046DM2;LINKAGE=Z
 NODE=S046DM2;LINKAGE=EA
 NODE=S046DM2;LINKAGE=Y
 NODE=S046DM2;LINKAGE=X
 NODE=S046DM2;LINKAGE=V
 NODE=S046DM2;LINKAGE=U
 NODE=S046DM2;LINKAGE=Q
 NODE=S046DM2;LINKAGE=R
 NODE=S046DM2;LINKAGE=AB
 NODE=S046DM2;LINKAGE=UH
 NODE=S046DM2;LINKAGE=FL
 NODE=S046DM2;LINKAGE=RO
 NODE=S046DM2;LINKAGE=AS
 NODE=S046DM2;LINKAGE=KI
 NODE=S046DM2;LINKAGE=NL
 NODE=S046DM2;LINKAGE=BL
 NODE=S046DM2;LINKAGE=BN
 NODE=S046DM2;LINKAGE=KM
 NODE=S046DM2;LINKAGE=ME
 NODE=S046DM2;LINKAGE=RM
 NODE=S046DM2;LINKAGE=NG
 NODE=S046DM2;LINKAGE=LN
 NODE=S046DM2;LINKAGE=AI
 NODE=S046DM2;LINKAGE=AL
 NODE=S046DM2;LINKAGE=AE

- 31 Predictions for the spin-independent elastic cross section in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046DM2;LINKAGE=TZ
- 32 KIM 02 and ELLIS 04 calculate the χp elastic scattering cross section in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry, but without universal scalar masses. NODE=S046DM2;LINKAGE=EK
- 33 In the case of universal squark and slepton masses, but non-universal Higgs masses, the limit becomes 2×10^{-6} (2×10^{-11} when constraint from the BNL $g-2$ experiment are included), see ELLIS 03E. ELLIS 05 display the sensitivity of the elastic scattering cross section to the π -Nucleon Σ term. NODE=S046DM2;LINKAGE=EL
- 34 PIERCE 04A calculates the χp elastic scattering cross section in the framework of models with very heavy scalar masses. See Fig. 2 of the paper. NODE=S046DM2;LINKAGE=PI
- 35 The strongest upper limit is 1.8×10^{-5} pb and occurs at $m_{\chi} \approx 80$ GeV. NODE=S046DM2;LINKAGE=AH
NODE=S046DM2;LINKAGE=AK
- 36 Under the assumption of standard WIMP-halo interactions, Akerib 03 is incompatible with BERNABEI 00 most likely value at the 99.98% CL. See Fig. 4. NODE=S046DM2;LINKAGE=BA
- 37 BAER 03A calculates the χp elastic scattering cross section in several models including the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046DM2;LINKAGE=BA
- 38 The strongest upper limit is 7×10^{-6} pb and occurs at $m_{\chi} \approx 30$ GeV. NODE=S046DM2;LINKAGE=A
NODE=S046DM2;LINKAGE=B
- 39 ABRAMS 02 is incompatible with the DAMA most likely value at the 99.9% CL. The strongest upper limit is 3×10^{-6} pb and occurs at $m_{\chi} \approx 30$ GeV. NODE=S046DM2;LINKAGE=E
NODE=S046DM2;LINKAGE=C
- 40 The strongest upper limit is 2×10^{-5} pb and occurs at $m_{\chi} \approx 40$ GeV. NODE=S046DM2;LINKAGE=H
NODE=S046DM2;LINKAGE=BT
- 41 The strongest upper limit is 7×10^{-6} pb and occurs at $m_{\chi} \approx 46$ GeV. NODE=S046DM2;LINKAGE=LH
- 42 The strongest upper limit is 1.8×10^{-5} pb and occurs at $m_{\chi} \approx 32$ GeV. NODE=S046DM2;LINKAGE=EM
- 43 BOTTINO 01 calculates the χ - p elastic scattering cross section in the framework of the following supersymmetric models: $N=1$ supergravity with the radiative breaking of the electroweak gauge symmetry, $N=1$ supergravity with nonuniversal scalar masses and an effective MSSM model at the electroweak scale. NODE=S046DM2;LINKAGE=J
- 44 Calculates the χ - p elastic scattering cross section in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046DM2;LINKAGE=NR
- 45 ELLIS 01C calculates the χ - p elastic scattering cross section in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. ELLIS 02B find a range 2×10^{-8} – 1.5×10^{-7} at $\tan\beta=50$. In models with nonuniversal Higgs masses, the upper limit to the cross section is 4×10^{-7} . NODE=S046DM2;LINKAGE=FG
- 46 ACCOMANDO 00 calculate the χ - p elastic scattering cross section in the framework of minimal $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. The limit is relaxed by at least an order of magnitude when models with nonuniversal scalar masses are considered. A subset of the authors in ARNOWITT 02 updated the limit to $< 9 \times 10^{-8}$ ($\tan\beta < 55$). NODE=S046DM2;LINKAGE=PHC
- 47 BERNABEI 00 search for annual modulation of the WIMP signal. The data favor the hypothesis of annual modulation at 4σ and are consistent, for a particular model framework quoted there, with $m_{\chi_0} = 44^{+12}_{-9}$ GeV and a spin-independent X^0 -proton cross section of $(5.4 \pm 1.0) \times 10^{-6}$ pb. See also BERNABEI 01 and BERNABEI 00c. NODE=S046DM2;LINKAGE=PHC
- 48 FENG 00 calculate the χ - p elastic scattering cross section in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry with a particular emphasis on focus point models. At $\tan\beta=50$, the range is 8×10^{-8} – 4×10^{-7} . NODE=S046DM2;LINKAGE=PHC

Other bounds on $\tilde{\chi}_1^0$ from astrophysics and cosmology

Most of these papers generally exclude regions in the $M_2 - \mu$ parameter plane by requiring that the $\tilde{\chi}_1^0$ contribution to the overall cosmological density is less than some maximal value to avoid overclosure of the Universe. Those not based on the cosmological density are indicated. Many of these papers also include LEP and/or other bounds.

VALUE	DOCUMENT ID	TECN	COMMENT
>46 GeV	1 ELLIS	00	RVUE
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	2 ATHRON	17B	COSM
	3 BECHTLE	16	COSM
	4 BAGNASCHI	15	COSM
	5 BUCHMUEL...	14	COSM
	6 BUCHMUEL...	14A	COSM
	7 ROSZKOWSKI	14	COSM
	8 CABRERA	13	COSM
	9 ELLIS	13B	COSM
	8 STREGE	13	COSM
	5 AKULA	12	COSM
	5 ARBEY	12A	COSM
	5 BAER	12	COSM
	10 BALAZS	12	COSM
	11 BECHTLE	12	COSM

NODE=S046PHC

NODE=S046PHC

NODE=S046PHC

	12	BESKIDT	12	COSM	
> 18 GeV	13	BOTTINO	12	COSM	
	5	BUCHMUEL...	12	COSM	
	5	CAO	12A	COSM	
	5	ELLIS	12B	COSM	
	14	FENG	12B	COSM	
	5	KADASTIK	12	COSM	
	10	STREGE	12	COSM	
	15	BUCHMUEL...	11	COSM	
	16	ROSZKOWSKI	11	COSM	
	17	ELLIS	10	COSM	
	18	BUCHMUEL...	09	COSM	
	19	DREINER	09	THEO	
	20	BUCHMUEL...	08	COSM	
	16	ELLIS	08	COSM	
	21	CALIBBI	07	COSM	
	22	ELLIS	07	COSM	
	23	ALLANACH	06	COSM	
	24	DE-AUSTRI	06	COSM	
	16	BAER	05	COSM	
	25	BALTZ	04	COSM	
> 6 GeV	13,26	BELANGER	04	THEO	
	27	ELLIS	04B	COSM	
	28	PIERCE	04A	COSM	
	29	BAER	03	COSM	
> 6 GeV	13	BOTTINO	03	COSM	
	29	CHATTOPAD...	03	COSM	
	30	ELLIS	03	COSM	
	16	ELLIS	03B	COSM	
	29	ELLIS	03C	COSM	
	29	LAHANAS	03	COSM	
	31	LAHANAS	02	COSM	
	32	BARGER	01C	COSM	
	33	ELLIS	01B	COSM	
	30	BOEHM	00B	COSM	
	34	FENG	00	COSM	
< 600 GeV	35	ELLIS	98B	COSM	
	36	EDSJO	97	COSM	Co-annihilation
	37	BAER	96	COSM	
	16	BEREZINSKY	95	COSM	
	38	FALK	95	COSM	CP-violating phases
	39	DREES	93	COSM	Minimal supergravity
	40	FALK	93	COSM	Sfermion mixing
	39	KELLEY	93	COSM	Minimal supergravity
	41	MIZUTA	93	COSM	Co-annihilation
	42	LOPEZ	92	COSM	Minimal supergravity, $m_0=A=0$
	43	MCDONALD	92	COSM	
	44	GRIEST	91	COSM	
	45	NOJIRI	91	COSM	Minimal supergravity
	46	OLIVE	91	COSM	
	47	ROSZKOWSKI	91	COSM	
	48	GRIEST	90	COSM	
	46	OLIVE	89	COSM	
none 100 eV – 15 GeV		SREDNICKI	88	COSM	$\tilde{\gamma}; m_{\tilde{f}}=100$ GeV
none 100 eV–5 GeV		ELLIS	84	COSM	$\tilde{\gamma};$ for $m_{\tilde{f}}=100$ GeV
		GOLDBERG	83	COSM	$\tilde{\gamma}$
	49	KRAUSS	83	COSM	$\tilde{\gamma}$
		VYSOTSKII	83	COSM	$\tilde{\gamma}$

OCCUR=2

OCCUR=2

OCCUR=2

¹ ELLIS 00 updates ELLIS 98. Uses LEP e^+e^- data at $\sqrt{s}=202$ and 204 GeV to improve bound on neutralino mass to 51 GeV when scalar mass universality is assumed and 46 GeV when Higgs mass universality is relaxed. Limits on $\tan\beta$ improve to > 2.7 ($\mu > 0$), > 2.2 ($\mu < 0$) when scalar mass universality is assumed and > 1.9 (both signs of μ) when Higgs mass universality is relaxed.

NODE=S046PHC;LINKAGE=EP

² ATHRON 17B places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using all Run I and the 13 fb^{-1} 13 TeV Run II LHC searches and other experimental data.

NODE=S046PHC;LINKAGE=J

³ BECHTLE 16 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using all Run I LHC searches.

NODE=S046PHC;LINKAGE=H

- 4 BAGNASCHI 15 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using all Run I LHC searches. NODE=S046PHC;LINKAGE=G
- 5 Implications of the LHC result on the Higgs mass and on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046PHC;LINKAGE=BR
- 6 BUCHMUELLER 14A places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using indirect experimental searches using the 20 fb^{-1} 8 TeV and the 5 fb^{-1} 7 TeV LHC and the LUX data. NODE=S046PHC;LINKAGE=UH
- 7 ROSZKOWSKI 14 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using Bayesian statistics and indirect experimental searches using the 20 fb^{-1} LHC and the LUX data. NODE=S046PHC;LINKAGE=RO
- 8 CABRERA 13 and STREGE 13 place constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry with and without non-universal Higgs masses using the 5.8 fb^{-1} , $\sqrt{s} = 7 \text{ TeV}$ ATLAS supersymmetry searches and XENON100 results. NODE=S046PHC;LINKAGE=SR
- 9 ELLIS 13B places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry with and without Higgs mass universality. Models with universality below the GUT scale are also considered. NODE=S046PHC;LINKAGE=LI
- 10 BALAZS 12 and STREGE 12 place constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using the 1 fb^{-1} LHC supersymmetry searches, the 5 fb^{-1} Higgs mass constraints, both with $\sqrt{s} = 7 \text{ TeV}$, and XENON100 results. NODE=S046PHC;LINKAGE=ST
- 11 BECHTLE 12 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using indirect experimental searches, using the 5 fb^{-1} LHC and XENON100 data. NODE=S046PHC;LINKAGE=BL
- 12 BESKIDT 12 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using indirect experimental searches, the 5 fb^{-1} LHC and the XENON100 data. NODE=S046PHC;LINKAGE=BS
- 13 BELANGER 04 and BOTTINO 12 (see also BOTTINO 03, BOTTINO 03A and BOTTINO 04) do not assume gaugino or scalar mass unification. NODE=S046PHC;LINKAGE=BO
- 14 FENG 12B places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry and large sfermion masses using the 1 fb^{-1} LHC supersymmetry searches, the 5 fb^{-1} LHC Higgs mass constraints both with $\sqrt{s} = 7 \text{ TeV}$, and XENON100 results. NODE=S046PHC;LINKAGE=FE
- 15 BUCHMUELLER 11 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using indirect experimental searches and including supersymmetry breaking relations between A and B parameters. NODE=S046PHC;LINKAGE=UC
- 16 Places constraints on the SUSY parameter space in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry but non-Universal Higgs masses. NODE=S046PHC;LINKAGE=EL
- 17 ELLIS 10 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry with universality above the GUT scale. NODE=S046PHC;LINKAGE=IS
- 18 BUCHMUELLER 09 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using indirect experimental searches. NODE=S046PHC;LINKAGE=BC
- 19 DREINER 09 show that in the general MSSM with non-universal gaugino masses there exists no model-independent laboratory bound on the mass of the lightest neutralino. An essentially massless χ_1^0 is allowed by the experimental and observational data, imposing some constraints on other MSSM parameters, including M_2 , μ and the slepton and squark masses. NODE=S046PHC;LINKAGE=DI
- 20 BUCHMUELLER 08 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry using indirect experimental searches. NODE=S046PHC;LINKAGE=BU
- 21 CALIBBI 07 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry with universality above the GUT scale including the effects of right-handed neutrinos. NODE=S046PHC;LINKAGE=CA
- 22 ELLIS 07 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry with universality below the GUT scale. NODE=S046PHC;LINKAGE=LL
- 23 ALLANACH 06 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046PHC;LINKAGE=AL
- 24 DE-AUSTRI 06 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046PHC;LINKAGE=DA
- 25 BALTZ 04 places constraints on the SUSY parameter space in the framework of $N = 1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046PHC;LINKAGE=TZ
- 26 Limit assumes a pseudo scalar mass $< 200 \text{ GeV}$. For larger pseudo scalar masses, $m_{\chi} > 18(29) \text{ GeV}$ for $\tan\beta = 50(10)$. Bounds from WMAP, $(g - 2)_{\mu}$, $b \rightarrow s\gamma$, LEP. NODE=S046PHC;LINKAGE=BV

- 27 ELLIS 04B places constraints on the SUSY parameter space in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry including supersymmetry breaking relations between A and B parameters. See also ELLIS 03D. NODE=S046PHC;LINKAGE=EI
- 28 PIERCE 04A places constraints on the SUSY parameter space in the framework of models with very heavy scalar masses. NODE=S046PHC;LINKAGE=PI
- 29 BAER 03, CHATTOPADHYAY 03, ELLIS 03C and LAHANAS 03 place constraints on the SUSY parameter space in the framework of $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry based on WMAP results for the cold dark matter density. NODE=S046PHC;LINKAGE=BA
- 30 BOEHM 00B and ELLIS 03 place constraints on the SUSY parameter space in the framework of minimal $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. Includes the effect of $\chi-\tilde{t}$ co-annihilations. NODE=S046PHC;LINKAGE=EB
- 31 LAHANAS 02 places constraints on the SUSY parameter space in the framework of minimal $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. Focuses on the role of pseudo-scalar Higgs exchange. NODE=S046PHC;LINKAGE=LH
- 32 BARGER 01C use the cosmic relic density inferred from recent CMB measurements to constrain the parameter space in the framework of minimal $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046PHC;LINKAGE=PA
- 33 ELLIS 01B places constraints on the SUSY parameter space in the framework of minimal $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. Focuses on models with large $\tan\beta$. NODE=S046PHC;LINKAGE=PD
- 34 FENG 00 explores cosmologically allowed regions of MSSM parameter space with multi-TeV masses. NODE=S046PHC;LINKAGE=FG
- 35 ELLIS 98B assumes a universal scalar mass and radiative supersymmetry breaking with universal gaugino masses. The upper limit to the LSP mass is increased due to the inclusion of $\chi-\tilde{\tau}_R$ coannihilations. NODE=S046PHC;LINKAGE=C8
- 36 EDSJO 97 included all coannihilation processes between neutralinos and charginos for any neutralino mass and composition. NODE=S046PHC;LINKAGE=EJ
- 37 Notes the location of the neutralino Z resonance and h resonance annihilation corridors in minimal supergravity models with radiative electroweak breaking. NODE=S046PHC;LINKAGE=BE
- 38 Mass of the bino (=LSP) is limited to $m_{\tilde{B}} \lesssim 350$ GeV for $m_t = 174$ GeV. NODE=S046PHC;LINKAGE=A
- 39 DREES 93, KELLEY 93 compute the cosmic relic density of the LSP in the framework of minimal $N=1$ supergravity models with radiative breaking of the electroweak gauge symmetry. NODE=S046PHC;LINKAGE=DR
- 40 FALK 93 relax the upper limit to the LSP mass by considering sfermion mixing in the MSSM. NODE=S046PHC;LINKAGE=FK
- 41 MIZUTA 93 include coannihilations to compute the relic density of Higgsino dark matter. NODE=S046PHC;LINKAGE=MZ
- 42 LOPEZ 92 calculate the relic LSP density in a minimal SUSY GUT model. NODE=S046PHC;LINKAGE=LZ
- 43 MCDONALD 92 calculate the relic LSP density in the MSSM including exact tree-level annihilation cross sections for all two-body final states. NODE=S046PHC;LINKAGE=MC
- 44 GRIEST 91 improve relic density calculations to account for coannihilations, pole effects, and threshold effects. NODE=S046PHC;LINKAGE=RG
- 45 NOJIRI 91 uses minimal supergravity mass relations between squarks and sleptons to narrow cosmologically allowed parameter space. NODE=S046PHC;LINKAGE=NJ
- 46 Mass of the bino (=LSP) is limited to $m_{\tilde{B}} \lesssim 350$ GeV for $m_t \leq 200$ GeV. Mass of the higgsino (=LSP) is limited to $m_{\tilde{H}} \lesssim 1$ TeV for $m_t \leq 200$ GeV. NODE=S046PHC;LINKAGE=S2
- 47 ROSZKOWSKI 91 calculates LSP relic density in mixed gaugino/higgsino region. NODE=S046PHC;LINKAGE=R9
- 48 Mass of the bino (=LSP) is limited to $m_{\tilde{B}} \lesssim 550$ GeV. Mass of the higgsino (=LSP) is limited to $m_{\tilde{H}} \lesssim 3.2$ TeV. NODE=S046PHC;LINKAGE=S3
- 49 KRAUSS 83 finds $m_{\tilde{\gamma}}$ not 30 eV to 2.5 GeV. KRAUSS 83 takes into account the gravitino decay. Find that limits depend strongly on reheated temperature. For example a new allowed region $m_{\tilde{\gamma}} = 4\text{--}20$ MeV exists if $m_{\text{gravitino}} < 40$ TeV. See figure 2. NODE=S046PHC;LINKAGE=D

Unstable $\tilde{\chi}_1^0$ (Lightest Neutralino) mass limit

Unless otherwise stated, results in this section assume spectra and production rates as evaluated in the MSSM. Unless otherwise stated, the goldstino or gravitino mass $m_{\tilde{G}}$ is assumed to be negligible relative to all other masses. In the following, \tilde{G} is assumed to be undetected and to give rise to a missing energy (\cancel{E}) signature.

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 900	95	1 AAD	23AE ATLS	2 SFOS ℓ , jets, \cancel{E}_T , Tn1n1C, $m_{\tilde{\chi}_1^0} = 1$ GeV
> 365	95	2 AAD	23AMATLS	long-lived $\tilde{\chi}_1^0$, displaced diphoton vertex, Tn1n1A, $\tau = 2$ ns

NODE=S046UPH

NODE=S046UPH

NODE=S046UPH;CHECK LIMITS

> 605	95	2 AAD	23AMATLS	long-lived $\tilde{\chi}_1^0$, displaced diphoton vertex, Tn1n1B, $\tau = 2$ ns	OCCUR=2
> 705	95	2 AAD	23AMATLS	long-lived $\tilde{\chi}_1^0$, displaced diphoton vertex, Tn1n1C, $\tau = 2$ ns	OCCUR=3
> 440	95	3 AAD	23CP ATLS	2 same-sign or 3 ℓ , Tn1n1D, bRPV higgsino decays to νW , ℓW	
>1180	95	4 TUMASYAN	23AO CMS	long-lived $\tilde{\chi}_1^0$, ≥ 2 trackless delayed jets + \cancel{E}_T , Tn1n1B, $c\tau = 0.5$ m	
> 990	95	4 TUMASYAN	23AO CMS	long-lived $\tilde{\chi}_1^0$, ≥ 2 trackless delayed jets + \cancel{E}_T , Tn1n1B, $c\tau = 3$ m	OCCUR=2
> 540	95	5 AAD	21Y ATLS	$\geq 4\ell$, Tchi1n12-GGM, $\tilde{\chi}_1^0 \rightarrow Z \tilde{G}$	
none 7-50	95	6 AAIJ	21V LHCb	$e^\pm \mu^\mp$, RPV $\tilde{\chi}_1^0 \rightarrow e^\pm \mu^\mp \nu$, 2 ps $< \tau < 50$ ps	
>1100	95	7 SIRUNYAN	21AF CMS	long-lived $\tilde{\chi}_1^0$, RPV $\tilde{\chi}_1^0 \rightarrow tbs$, λ_{323}'' coupling, 0.6 mm $< c\tau < 70$ mm	
> 800	95	8 SIRUNYAN	21M CMS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tn1n1C	
> 650	95	8 SIRUNYAN	21M CMS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tn1n1B	OCCUR=2
> 380	95	9 AAD	20AN ATLS	$2\gamma + \cancel{E}_T$, Tn1n1A, GMSB	
> 525	95	10 SIRUNYAN	19CA CMS	$\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$, GMSB, SPS8, $c\tau=1$ m	
> 290	95	11 SIRUNYAN	19CI CMS	$\geq 1 H (\rightarrow \gamma\gamma) +$ jets + \cancel{E}_T , Tn1n1A, GMSB	
> 230	95	11 SIRUNYAN	19CI CMS	$\geq 1 H (\rightarrow \gamma\gamma) +$ jets + \cancel{E}_T , Tn1n1B, GMSB	OCCUR=2
> 930	95	12 SIRUNYAN	19K CMS	$\gamma +$ lepton + \cancel{E}_T , Tchi1n1A	
none 130-230, 290-880	95	13 AABOUD	18CK ATLS	$2H (\rightarrow bb) + \cancel{E}_T$, Tn1n1A, GMSB	
> 295	95	14 AABOUD	18Z ATLS	$\geq 4\ell$, GMSB, Tn1n1C	
> 180	95	15 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tn1n1A	
> 260	95	15 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tn1n1B	OCCUR=2
> 450	95	15 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tn1n1C	OCCUR=3
> 750	95	16 SIRUNYAN	18AP CMS	Combination of searches, GMSB, Tn1n1A	
> 650	95	16 SIRUNYAN	18AP CMS	Combination of searches, GMSB, Tn1n1B	OCCUR=2
> 690	95	16 SIRUNYAN	18AP CMS	Combination of searches, GMSB, Tn1n1C	OCCUR=3
> 500	95	17 SIRUNYAN	18AR CMS	$\ell^\pm \ell^\mp +$ jets + \cancel{E}_T , GMSB, Tn1n1B	
> 650	95	17 SIRUNYAN	18AR CMS	$\ell^\pm \ell^\mp +$ jets + \cancel{E}_T , GMSB, Tn1n1C	OCCUR=2
none 230-770	95	18 SIRUNYAN	18O CMS	$2H (\rightarrow bb) + \cancel{E}_T$, Tn1n1A, GMSB	
> 205	95	19 SIRUNYAN	18X CMS	$\geq 1 H (\rightarrow \gamma\gamma) +$ jets + \cancel{E}_T , Tn1n1A, GMSB	
> 130	95	19 SIRUNYAN	18X CMS	$\geq 1 H (\rightarrow \gamma\gamma) +$ jets + \cancel{E}_T , Tn1n1B, GMSB	OCCUR=2
> 380	95	20 KHACHATRY...14L	CMS	$\tilde{\chi}_1^0 \rightarrow Z \tilde{G}$ simplified models, GMSB, RPV	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
		21 AAD	20D	$\tilde{q} \rightarrow q \tilde{\chi}_1^0$, $\tilde{\chi}_1^0 \rightarrow \ell\ell\nu$, RPV, λ_{121} or $\lambda_{122} \neq 0$	
none 300-1000	95	22 AABOUD	19G ATLS	$\tilde{\chi}_1^0 \rightarrow Z \tilde{G}$ from gluinos as in Tglu1A, GMSB, depending on $c\tau$	
		23 AAIJ	17Z	displaced vertex with associated μ	
		24 KHACHATRY...16BX	CMS	$\geq 3\ell^\pm$, RPV, λ or λ' couplings, wino- or higgsino-like neutralinos	
		25 AAD	14BH ATLS	$2\gamma + \cancel{E}_T$, GMSB, SPS8	
		26 AAD	13AP ATLS	$2\gamma + \cancel{E}_T$, GMSB, SPS8	
none 220-380	95	27 AAD	13Q ATLS	$\gamma + b + \cancel{E}_T$, higgsino-like neutralino, GMSB	
		28 AAD	13R ATLS	$\tilde{\chi}_1^0 \rightarrow \mu jj$, RPV, $\lambda'_{211} \neq 0$	
		29 AALTONEN	13I CDF	$\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$, \cancel{E}_T , GMSB	
> 220	95	30 CHATRCHYAN	13AH CMS	$\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$, GMSB, SPS8, $c\tau < 500$ mm	
		31 AAD	12CP ATLS	$2\gamma + \cancel{E}_T$, GMSB	
		32 AAD	12CT ATLS	$\geq 4\ell^\pm$, RPV	
		33 AAD	12R ATLS	$\tilde{\chi}_1^0 \rightarrow \mu jj$, RPV, $\lambda'_{211} \neq 0$	
		34 ABAZOV	12AD D0	$\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow \gamma Z \tilde{G}$, GMSB	
		35 CHATRCHYAN	12BK CMS	$2\gamma + \cancel{E}_T$, GMSB	

> 149	95	36 CHATRCHYAN11B	10	CMS	$\tilde{W}^0 \rightarrow \gamma \tilde{G}, \tilde{W}^\pm \rightarrow \ell^\pm \tilde{G}, \text{GMSB}$	
		37 AALTONEN	10	CDF	$p\bar{p} \rightarrow \tilde{\chi}\tilde{\chi}, \tilde{\chi}=\tilde{\chi}_2^0, \tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}, \text{GMSB}$	
> 175	95	38 ABAZOV	10P	D0	$\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}, \text{GMSB}$	
> 125	95	39 ABAZOV	08F	D0	$p\bar{p} \rightarrow \tilde{\chi}\tilde{\chi}, \tilde{\chi}=\tilde{\chi}_2^0, \tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}, \text{GMSB}$	
> 96.8	95	40 ABULENCIA	07H	CDF	RPV, $LL\bar{E}$	
		41 ABBIENDI	06B	OPAL	$e^+e^- \rightarrow \tilde{B}\tilde{B}, (\tilde{B} \rightarrow \tilde{G}\gamma)$	
		42 ABDALLAH	05B	DLPH	$e^+e^- \rightarrow \tilde{G}\tilde{\chi}_1^0, (\tilde{\chi}_1^0 \rightarrow \tilde{G}\gamma)$	
> 96	95	43 ABDALLAH	05B	DLPH	$e^+e^- \rightarrow \tilde{B}\tilde{B}, (\tilde{B} \rightarrow \tilde{G}\gamma)$	OCCUR=2
		1 AAD 23AE			searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 2 ℓ with same flavour and opposite sign, plus jets and \cancel{E}_T , defining signal region with the dilepton invariant mass both on- and off-shell with respect to the Z boson. No significant excess above the Standard Model predictions is observed. Limits are set on models of strong and electroweak production. In this case, limits are placed on production of mass-degenerate, higgsino triplet NLSP with $\tilde{\chi}_1^0 \rightarrow Z\tilde{G}$ in a GGM-like scenario, see figure 15.	NODE=S046UPH;LINKAGE=RA
		2 AAD 23AM			searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing electron/photon pairs with invariant mass compatible with h/Z and originating from a common displaced vertex. No significant excess above the Standard Model predictions is observed. Limits are set on a model where members of a nearly degenerate higgsino triplet are pair-produced, yielding long-lived $\tilde{\chi}_1^0$ followed by $\tilde{\chi}_1^0 \rightarrow h/Z\tilde{G}$. Limits are set on $m_{\tilde{\chi}_1^0}$ as a function of its lifetime and of the $B(\tilde{\chi}_1^0 \rightarrow h\tilde{G})$ assuming $B(\tilde{\chi}_1^0 \rightarrow h\tilde{G}) + B(\tilde{\chi}_1^0 \rightarrow Z\tilde{G}) = 1$, see Figure 10.	NODE=S046UPH;LINKAGE=UA
		3 AAD 23CP			searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 2 ℓ with same charge or 3 ℓ plus at least one jet and \cancel{E}_T , defining signal region based on 'stransverse mass' of the dilepton system, \cancel{E}_T significance and effective mass. No significant excess above the Standard Model predictions is observed. Limits are set on the mass of a mass-degenerate higgsino triplet decaying into a lepton (neutral or charged) and a W via a bilinear RPV coupling, see figure 14.	NODE=S046UPH;LINKAGE=QA
		4 TUMASYAN 23AO			searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of neutralino-chargino production in events with nearly trackless and out-of-time jets that are used to identify decays of long-lived particles. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the long-lived $\tilde{\chi}_1^0$ in the model Tn1n1B, see their figures 8–10.	NODE=S046UPH;LINKAGE=PA
		5 AAD 21Y			searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with four or more leptons (electrons, muons and tau-leptons). No significant excess above the Standard Model expectations is observed. Limits are set on Tchi1n12-GGM, and RPV models similar to Tchi1n2l, Tglu1A (with $q = u, d, s, c, b$, with equal branching fractions), and $\tilde{\ell}_L/\tilde{\nu} \rightarrow \ell/\nu\tilde{\chi}_1^0$ (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations), all with $\tilde{\chi}_1^0 \rightarrow \ell^\pm\ell^\mp\nu$ via λ_{12k} or λ_{i33} (where $i, k \in 1, 2$), see their Figure 11.	NODE=S046UPH;LINKAGE=HA
		6 AAIJ 21V			searched in 5.38 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived particles (LLP) decaying to $e^\pm\mu^\mp\nu$. The LLP can be a $\tilde{\chi}_1^0$ in RPV SUSY, or a right-handed neutrino, and can be produced in pairs, in the decay of the Higgs boson, or from charged current processes. No significant excess above the Standard Model expectations is observed. Limits are set on the cross section times branching ratio for all three production mechanisms, see their Figures 6–8.	NODE=S046UPH;LINKAGE=IA
		7 SIRUNYAN 21AF			searched in 140 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with with two displaced vertices from long-lived particles decaying into multijet or dijet final states. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu2RPV with λ''_{323} coupling, on the $\tilde{\chi}_1^0$ mass in an RPV model with $\tilde{\chi}_1^0$ pair production and the RPV decay $\tilde{\chi}_1^0 \rightarrow tbs$ with λ''_{323} coupling and on the \tilde{t} mass in an RPV model with top squark pair production and the RPV decay $\tilde{t} \rightarrow \bar{d}_i\bar{d}_j$ with λ''_{3ij} coupling, see their Figure 7.	NODE=S046UPH;LINKAGE=NA
		8 SIRUNYAN 21M			searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0}=m_{\tilde{\chi}_1^\pm}=m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tsbob3, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14.	NODE=S046UPH;LINKAGE=MA
		9 AAD 20AN			searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons and missing transverse momentum. Events are further categorised in terms of lepton or jet multiplicity. No significant excess over the expected background is observed.	NODE=S046UPH;LINKAGE=FA

- Limits at 95% C.L. are set on the Higgsino mass in the T1n1n1A simplified model, see their Figure 11.
- 10 SIRUNYAN 19CA searched in 77.4 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing delayed photons in both single and diphoton plus \cancel{E}_T final states. No excess is observed above the background expected from Standard Model processes. The results are used to set 95% C.L. exclusion limits in the context of GMSB, using the SPS8 benchmark model. For neutralino proper decay lengths of 0.1, 1, 10, and 100 m, masses up to about 320, 525, 360, and 215 GeV are excluded, respectively. See their Fig. 5. The searches involve the simplified models Tglu1D, Tglu4A,B,C, Tsqk4,4A,4B. NODE=S046UPH;LINKAGE=Z
 - 11 SIRUNYAN 19CI searched in 77.5 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more high-momentum Higgs bosons, decaying to pairs of photons, jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t4 simplified model, see Figure 3, and on the wino mass in the Tchi1n2E simplified model, see their Figure 4. Limits are also set on the higgsino mass in the Tn1n1A and Tn1n1B simplified models, see their Figure 5. NODE=S046UPH;LINKAGE=Y
 - 12 SIRUNYAN 19K searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a photon, an electron or muon, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. In the framework of GMSB, limits are set on the chargino and neutralino mass in the Tchi1n1A simplified model, see their Figure 6. Limits are also set on the gluino mass in the Tglu4A simplified model, and on the squark mass in the Tsqk4A simplified model, see their Figure 7. NODE=S046UPH;LINKAGE=X
 - 13 AABOUD 18CK searched for events with at least 3 b -jets and large missing transverse energy in two datasets of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ of 36.1 fb^{-1} and 24.3 fb^{-1} depending on the trigger requirements. The analyses aimed to reconstruct two Higgs bosons decaying to pairs of b -quarks. No significant excess above the Standard Model expectations is observed. Limits are set on the Higgsino mass in the Tn1n1A simplified model, see their Figure 15(a). Constraints are also presented as a function of the BR of Higgsino decaying into an higgs boson and a gravitino, see their Figure 15(b). NODE=S046UPH;LINKAGE=V
 - 14 AABOUD 18Z searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more charged leptons (electrons, muons and up to two hadronically decaying taus). No significant deviation from the expected SM background is observed. Limits are set on the Higgsino mass in simplified models of general gauge mediated supersymmetry Tn1n1A/Tn1n1B/Tn1n1C, see their Figure 9. Limits are also set on the wino, slepton, sneutrino and gluino mass in a simplified model of NLSP pair production with R-parity violating decays of the LSP via λ_{12k} or λ_{j33} to charged leptons, see their Figures 7, 8. NODE=S046UPH;LINKAGE=Q
 - 15 SIRUNYAN 18AO searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and neutralinos in events with either two or more leptons (electrons or muons) of the same electric charge, or with three or more leptons, which can include up to two hadronically decaying tau leptons. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino/neutralino mass in the Tchi1n2A, Tchi1n2H, Tchi1n2D, Tchi1n2E and Tchi1n2F simplified models, see their Figures 14, 15, 16, 17 and 18. Limits are also set on the higgsino mass in the Tn1n1A, Tn1n1B and Tn1n1C simplified models, see their Figure 19. NODE=S046UPH;LINKAGE=S
 - 16 SIRUNYAN 18AP searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and neutralinos by combining a number of previous and new searches. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino/neutralino mass in the Tchi1n2E, Tchi1n2F and Tchi1n2I simplified models, see their Figures 7, 8, 9 and 10. Limits are also set on the higgsino mass in the Tn1n1A, Tn1n1B and Tn1n1C simplified models, see their Figure 11, 12, 13 and 14. NODE=S046UPH;LINKAGE=T
 - 17 SIRUNYAN 18AR searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two opposite-charge, same-flavour leptons (electrons or muons), jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4C simplified model, see their Figure 7. Limits are also set on the chargino/neutralino mass in the Tchi1n2F simplified models, see their Figure 8, and on the higgsino mass in the Tn1n1B and Tn1n1C simplified models, see their Figure 9. Finally, limits are set on the sbottom mass in the Tsb0t3 simplified model, see their Figure 10. NODE=S046UPH;LINKAGE=U
 - 18 SIRUNYAN 18O searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two Higgs bosons, decaying to pairs of b -quarks, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the Higgsino mass in the T1n1n1A simplified model, see their Figure 9. NODE=S046UPH;LINKAGE=R
 - 19 SIRUNYAN 18X searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more high-momentum Higgs bosons, decaying to pairs of photons, jets and \cancel{E}_T . The razor variables (M_R and R^2) are used to categorise the events. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t4 simplified model and on the wino mass in the Tchi1n2E simplified model, see their Figure 5. Limits are also set on the higgsino mass in the Tn1n1A and Tn1n1B simplified models, see their Figure 6. NODE=S046UPH;LINKAGE=P
 - 20 KHACHATRYAN 14L searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of direct pair production of neutralinos with Higgs or Z -bosons in the decay chain, leading to HH , HZ and ZZ final states with missing transverse energy. The decays of 16–20. a Higgs boson to a b -quark pair, to a photon pair, and to final states with leptons are considered in conjunction with hadronic and leptonic decay modes of the Z and W bosons. No significant excesses over the expected SM backgrounds are observed. The results are interpreted in the context of GMSB simplified models where the decays $\tilde{\chi}_1^0 \rightarrow H\tilde{G}$ or $\tilde{\chi}_1^0 \rightarrow Z\tilde{G}$ take place either 100% or 50% of the time, see Figs. 16–20. NODE=S046UPH;LINKAGE=D

- 21 AAD 20D searched in 32.8 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing an oppositely charge lepton pair ($ee, \mu\mu$ or $e\mu$) coming from long-lived neutralinos decaying through the R-parity-violating decay $\tilde{\chi}_1^0 \rightarrow \ell\ell\nu$ with $\lambda_{121} \neq 0$ or $\lambda_{122} \neq 0$. No excess over the expected background is observed. Limits are derived for decay lengths of the neutralino between 1 mm and 10 m in a scenario where a squark-antisquark pair is produced, with the squark decaying to a quark and a $\tilde{\chi}_1^0$, with either $\tilde{\chi}_1^0 \rightarrow ee\nu/e\mu\nu$ ($\lambda_{121} \neq 0$) or $\tilde{\chi}_1^0 \rightarrow e\mu\nu/\mu\mu\nu$ ($\lambda_{122} \neq 0$), see their Figures 4 and 5. NODE=S046UPH;LINKAGE=BA
- 22 AABOUD 19G searched in 32.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of neutralinos decaying into a Z-boson and a gravitino, in events characterized by the presence of dimuon vertices with displacements from the pp interaction point in the range of 1400 cm. Neutralinos are assumed to be produced in the decay chain of gluinos as in Tglu1A models. No significant excess is observed in the number of vertices relative to the predicted background. In GGM with a gluino mass of 1100 GeV, neutralino masses in the range 300–1000 GeV are excluded for certain values of $c\tau$, see their Figure 7. NODE=S046UPH;LINKAGE=W
- 23 AAIJ 17Z searched in 1 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and in 2 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing a displaced vertex with one associated high transverse momentum μ . No excess is observed above the background expected from Standard Model processes. The results are used to set 95% C.L. upper limits on the cross section times branching fractions of pair-produced neutralinos decaying non-promptly into a muon and two quarks. Long-lived particles in a mass range 23–198 GeV are considered, see their Fig. 5 and Fig. 6. NODE=S046UPH;LINKAGE=O
- 24 KHACHATRYAN 16BX searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing 3 or more leptons coming from the electroweak production of wino- or higgsino-like neutralinos, assuming non-zero R-parity-violating leptonic couplings λ_{122} , λ_{123} , and λ_{233} or semileptonic couplings λ'_{131} , λ'_{233} , λ'_{331} , and λ'_{333} . No excess over the expected background is observed and limits are derived on the neutralino mass, see Figs. 24 and 25. NODE=S046UPH;LINKAGE=I
- 25 AAD 14BH searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing non-pointing photons in a diphoton plus missing transverse energy final state. No excess is observed above the background expected from Standard Model processes. The results are used to set 95% C.L. exclusion limits in the context of gauge-mediated supersymmetric breaking models, with the lightest neutralino being the next-to-lightest supersymmetric particle and decaying with a lifetime in the range from 0.25 ns to about 100 ns into a photon and a gravitino. For limits on the NLSP lifetime versus Λ plane, for the SPS8 model, see their Fig. 7. NODE=S046UPH;LINKAGE=H
- 26 AAD 13AP searched in 4.8 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing non-pointing photons in a diphoton plus missing transverse energy final state. No excess is observed above the background expected from Standard Model processes. The results are used to set 95% C.L. exclusion limits in the context of gauge-mediated supersymmetric breaking models, with the lightest neutralino being the next-to-lightest supersymmetric particle and decaying with a lifetime in excess of 0.25 ns into a photon and a gravitino. For limits in the NLSP lifetime versus Λ plane, for the SPS8 model, see their Fig. 8. NODE=S046UPH;LINKAGE=LS
- 27 AAD 13Q searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing a high- p_T isolated photon, at least one jet identified as originating from a bottom quark, and high missing transverse momentum. Such signatures may originate from supersymmetric models with gauge-mediated supersymmetry breaking in events in which one of a pair of higgsino-like neutralinos decays into a photon and a gravitino while the other decays into a Higgs boson and a gravitino. No significant excess above the expected background was found and limits were set on the neutralino mass in a generalized GMSB model (GGM) with a higgsino-like neutralino NLSP, see their Fig. 4. Intermediate neutralino masses between 220 and 380 GeV are excluded at 95% C.L., regardless of the squark and gluino masses, purely on the basis of the expected weak production. NODE=S046UPH;LINKAGE=SA
- 28 AAD 13R looked in 4.4 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing new, heavy particles that decay at a significant distance from their production point into a final state containing a high-momentum muon and charged hadrons. No excess over the expected background is observed and limits are placed on the production cross-section of neutralinos via squarks for various $m_{\tilde{q}}, m_{\tilde{\chi}_1^0}$ in an R-parity violating scenario with $\lambda'_{211} \neq 0$, as a function of the neutralino lifetime, see their Fig. 6. NODE=S046UPH;LINKAGE=GA
- 29 AALTONEN 13I searched in 6.3 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for events containing \cancel{E}_T and a delayed photon that arrives late in the detector relative to the time expected from prompt production. No evidence of delayed photon production is observed. NODE=S046UPH;LINKAGE=LO
- 30 CHATRCHYAN 13AH searched in 4.9 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing \cancel{E}_T and a delayed photon that arrives late in the detector relative to the time expected from prompt production. No significant excess above the expected background was found and limits were set on the pair production of $\tilde{\chi}_1^0$ depending on the neutralino proper decay length, see Fig. 8. Supersedes CHATRCHYAN 12BK. NODE=S046UPH;LINKAGE=CA
- 31 AAD 12CP searched in 4.8 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with two photons and large \cancel{E}_T due to $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$ decays in a GMSB framework. No significant excess above the expected background was found and limits were set on the neutralino mass in a generalized GMSB model (GGM) with a bino-like neutralino NLSP, see Figs. 6 and 7. The other sparticle masses were decoupled, $\tan\beta = 2$ and $c\tau_{NLSP} < 0.1 \text{ mm}$. Also, in the framework of the SPS8 model, limits are presented in Fig. 8. NODE=S046UPH;LINKAGE=GT

- 32 AAD 12CT searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing four or more leptons (electrons or muons) and either moderate values of missing transverse momentum or large effective mass. No significant excess is found in the data. Limits are presented in a simplified model of R -parity violating supersymmetry in which charginos are pair-produced and then decay into a W -boson and a $\tilde{\chi}_1^0$, which in turn decays through an RPV coupling into two charged leptons ($e^\pm e^\mp$ or $\mu^\pm \mu^\mp$) and a neutrino. In this model, limits are set on the neutralino mass as a function of the chargino mass, see Fig. 3a. Limits are also set in an R -parity violating mSUGRA model, see Fig. 3b. NODE=S046UPH;LINKAGE=DG
- 33 AAD 12R looked in 33 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing new, heavy particles that decay at a significant distance from their production point into a final state containing a high-momentum muon and charged hadrons. No excess over the expected background is observed and limits are placed on the production cross-section of neutralinos via squarks for various $(m_{\tilde{q}}, m_{\tilde{\chi}_1^0})$ in an R -parity violating scenario with $\lambda'_{211} \neq 0$, as a function of the neutralino lifetime, see their Fig. 8. Superseded by AAD 13R. NODE=S046UPH;LINKAGE=GD
- 34 ABAZOV 12AD looked in 6.2 fb^{-1} of pp collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for events with a photon, a Z -boson, and large \cancel{E}_T in the final state. This topology corresponds to a GMSB model where pairs of neutralino NLSPs are either pair produced promptly or from decays of other supersymmetric particles and then decay to either $Z\tilde{G}$ or $\gamma\tilde{G}$. No significant excess over the SM expectation is observed and a limit at 95% C.L. on the cross section is derived as a function of the effective SUSY breaking scale Λ , see Fig. 3. Assuming $N_{mes} = 2$, $M_{mes} = 3 \Lambda$, $\tan\beta = 3$, $\mu = 0.75 M_1$, and $C_{grav} = 1$, the model is excluded at 95% C.L. for values of $\Lambda < 87 \text{ TeV}$. NODE=S046UPH;LINKAGE=AV
- 35 CHATRCHYAN 12BK searched in 2.23 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with two photons and large \cancel{E}_T due to $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$ decays in a GMSB framework. No significant excess above the expected background was found and limits were set on the pair production of $\tilde{\chi}_1^0$ depending on the neutralino lifetime, see Fig. 6. NODE=S046UPH;LINKAGE=CH
- 36 CHATRCHYAN 11B looked in 35 pb^{-1} of pp collisions at $\sqrt{s}=7 \text{ TeV}$ for events with an isolated lepton (e or μ), a photon and \cancel{E}_T which may arise in a generalized gauge mediated model from the decay of Wino-like NLSPs. No evidence for an excess over the expected background is observed. Limits are derived in the plane of squark/gluino mass versus Wino mass (see Fig. 4). Mass degeneracy of the produced squarks and gluinos is assumed. NODE=S046UPH;LINKAGE=C1
- 37 AALTONEN 10 searched in 2.6 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for diphoton events with large \cancel{E}_T . They may originate from the production of $\tilde{\chi}^\pm$ in pairs or associated to a $\tilde{\chi}_2^0$, decaying into $\tilde{\chi}_1^0$ which itself decays in GMSB to $\gamma\tilde{G}$. There is no excess of events beyond expectation. An upper limit on the cross section is calculated in the GMSB model as a function of the $\tilde{\chi}_1^0$ mass and lifetime, see their Fig. 2. A limit is derived on the $\tilde{\chi}_1^0$ mass of 149 GeV for $\tau_{\tilde{\chi}_1^0} \ll 1 \text{ ns}$, which improves the results of previous searches. NODE=S046UPH;LINKAGE=LT
- 38 ABAZOV 10P looked in 6.3 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for events with at least two isolated γ s and large \cancel{E}_T . These could be the signature of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ production, decaying to $\tilde{\chi}_1^0$ and finally $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$ in a GMSB framework. No significant excess over the SM expectation is observed, and a limit at 95% C.L. on the cross section is derived for $N_{mes} = 1$, $\tan\beta = 15$ and $\mu > 0$, see their Fig. 2. This allows them to set a limit on the effective SUSY breaking scale $\Lambda > 124 \text{ TeV}$, from which the excluded $\tilde{\chi}_1^0$ mass range is obtained. NODE=S046UPH;LINKAGE=AZ
- 39 ABAZOV 08F looked in 1.1 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for diphoton events with large \cancel{E}_T . They may originate from the production of $\tilde{\chi}^\pm$ in pairs or associated to a $\tilde{\chi}_2^0$, decaying to a $\tilde{\chi}_1^0$ which itself decays promptly in GMSB to $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$. No significant excess was found compared to the background expectation. A limit is derived on the masses of SUSY particles in the GMSB framework for $M = 2\Lambda$, $N = 1$, $\tan\beta = 15$ and $\mu > 0$, see Figure 2. It also excludes $\Lambda < 91.5 \text{ TeV}$. Supersedes the results of ABAZOV 05A. Superseded by ABAZOV 10P. NODE=S046UPH;LINKAGE=ZO
- 40 ABULENCIA 07H searched in 346 pb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for events with at least three leptons (e or μ) from the decay of $\tilde{\chi}_1^0$ via $LL\bar{E}$ couplings. The results are consistent with the hypothesis of no signal. Upper limits on the cross-section are extracted and a limit is derived in the framework of mSUGRA on the masses of $\tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm$, see e.g. their Fig. 3 and Tab. II. NODE=S046UPH;LINKAGE=BL
- 41 ABBIENDI 06B use 600 pb^{-1} of data from $\sqrt{s} = 189\text{--}209 \text{ GeV}$. They look for events with diphotons + \cancel{E} final states originating from prompt decays of pair-produced neutralinos in a GMSB scenario with $\tilde{\chi}_1^0$ NLSP. Limits on the cross-section are computed as a function of $m(\tilde{\chi}_1^0)$, see their Fig. 14. The limit on the $\tilde{\chi}_1^0$ mass is for a pure Bino state assuming a prompt decay, with lifetimes up to 10^{-9}s . Supersedes the results of ABBIENDI 04N. NODE=S046UPH;LINKAGE=BI
- 42 ABDALLAH 05B use data from $\sqrt{s} = 180\text{--}209 \text{ GeV}$. They look for events with single photons + \cancel{E} final states. Limits are computed in the plane $(m(\tilde{G}), m(\tilde{\chi}_1^0))$, shown in their Fig. 9b for a pure Bino state in the GMSB framework and in Fig. 9c for a no-scale supergravity model. Supersedes the results of ABREU 00Z. NODE=S046UPH;LINKAGE=AA

⁴³ ABDALLAH 05B use data from $\sqrt{s} = 130\text{--}209$ GeV. They look for events with diphotons + \cancel{E} final states and single photons not pointing to the vertex, expected in GMSB when the $\tilde{\chi}_1^0$ is the NLSP. Limits are computed in the plane ($m(\tilde{G}), m(\tilde{\chi}_1^0)$), see their Fig. 10. The lower limit is derived on the $\tilde{\chi}_1^0$ mass for a pure Bino state assuming a prompt decay and $m_{\tilde{e}_R} = m_{\tilde{e}_L} = 2 m_{\tilde{\chi}_1^0}$. It improves to 100 GeV for $m_{\tilde{e}_R} = m_{\tilde{e}_L} = 1.1 m_{\tilde{\chi}_1^0}$. and the limit in the plane ($m(\tilde{\chi}_1^0), m(\tilde{e}_R)$) is shown in Fig. 10b. For long-lived neutralinos, cross-section limits are displayed in their Fig 11. Supersedes the results of ABREU 00Z.

NODE=S046UPH;LINKAGE=AL

$\tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0$ (Neutralinos) mass limits

Neutralinos are unknown mixtures of photinos, z-inos, and neutral higgsinos (the supersymmetric partners of photons and of Z and Higgs bosons). The limits here apply only to $\tilde{\chi}_2^0, \tilde{\chi}_3^0$, and $\tilde{\chi}_4^0$. $\tilde{\chi}_1^0$ is the lightest supersymmetric particle (LSP); see $\tilde{\chi}_1^0$ Mass Limits. It is not possible to quote rigorous mass limits because they are extremely model dependent; i.e. they depend on branching ratios of various $\tilde{\chi}^0$ decay modes, on the masses of decay products ($\tilde{e}, \tilde{\gamma}, \tilde{q}, \tilde{g}$), and on the \tilde{e} mass exchanged in $e^+e^- \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0$. Limits arise either from direct searches, or from the MSSM constraints set on the gaugino and higgsino mass parameters M_2 and μ through searches for lighter charginos and neutralinos. Often limits are given as contour plots in the $m_{\tilde{\chi}^0} - m_{\tilde{e}}$ plane vs other parameters. When specific assumptions are made, e.g. the neutralino is a pure photino ($\tilde{\gamma}$), pure z-ino (\tilde{Z}), or pure neutral higgsino (\tilde{H}^0), the neutralinos will be labelled as such.

NODE=S046ZNO

NODE=S046ZNO

Limits obtained from e^+e^- collisions at energies up to 136 GeV, as well as other limits from different techniques, are now superseded and have not been included in this compilation. They can be found in the 1998 Edition (The European Physical Journal **C3** 1 (1998)) of this Review. Some later papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT		
> 820	95	¹ AAD	23AE ATLS	2 SFOS ℓ , jets, \cancel{E}_T , Tchi1n2Fa, $m_{\tilde{\chi}_1^0} = 1$ GeV		NODE=S046ZNO
none 260–420	95	² AAD	23CI ATLS	$1\ell + \text{jets} + \cancel{E}_T$, Tchi1n2J, $m_{\tilde{\chi}_1^0} = 0$ GeV		
> 230	95	³ AAD	23CI ATLS	$1\ell + \text{jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 133$ GeV		OCCUR=2
> 450	95	³ AAD	23CI ATLS	$1\ell + \text{jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 260$ GeV		OCCUR=3
> 525	95	⁴ AAD	23CP ATLS	2 same-sign ℓ , Tchi1n2E, wino-bino, $m_{\tilde{\chi}_1^0} = 1$ GeV		
none 200–250	95	⁴ AAD	23CP ATLS	2 same-sign ℓ , Tchi1n2F, wino-bino, $m_{\tilde{\chi}_1^0} = 1$ GeV		OCCUR=2
none 200–585	95	⁵ AAD	23CR ATLS	RPV, 2 same-sign, 3, 4 ℓ , 1, 2 b -jets, higgsino production with $\tilde{\chi} \rightarrow b + \ell/\nu + t/b$ via λ'_{i33} coupling		
none 200–670	95	⁵ AAD	23CR ATLS	RPV, 2 same-sign, 3, 4 ℓ , 1, 2 b -jets, wino production with $\tilde{\chi} \rightarrow b + \ell/\nu + t/b$ via λ'_{i33} coupling		OCCUR=2
>1050	95	⁶ HAYRAPETY...23E	CMS	$\gamma + \text{jets} + \cancel{E}_T$, Tchi1chi1A		
> 450	95	⁶ HAYRAPETY...23E	CMS	$\gamma + \text{jets} + \cancel{E}_T$, Tn1n2A		OCCUR=2
none 290–670	95	⁷ TUMASYAN	23B CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , Tchi1chi1l, $m_{\tilde{\chi}_1^0} = 1$ GeV		
none 230–760	95	⁷ TUMASYAN	23B CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , Tchi1n2Fb, $m_{\tilde{\chi}_1^0} = 1$ GeV		OCCUR=2
none 240–970	95	⁷ TUMASYAN	23B CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , Tchi1n2Fc, $m_{\tilde{\chi}_1^0} = 1$ GeV		OCCUR=3
none 300–650	95	⁷ TUMASYAN	23B CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , THinoBinoA, $m_{\tilde{\chi}_1^0} = 1$ GeV		OCCUR=4
> 275	95	⁸ TUMASYAN	22Q CMS	2 or 3 ℓ (soft), \cancel{E}_T ; Tchi1n2F, wino-bino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10$ GeV		

> 205	95	⁸	TUMASYAN	22Q	CMS	2 or 3 ℓ (soft), \cancel{E}_T ; higgsino model with $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ prod., $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 7.5$ GeV	OCCUR=2
> 150	95	⁸	TUMASYAN	22Q	CMS	2 or 3 ℓ (soft), \cancel{E}_T ; higgsino model with $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ prod., $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 3$ GeV	OCCUR=3
>1450	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 850$ GeV	OCCUR=2
>1360	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
>1290	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 0.05m_{\tilde{\chi}_1^\pm} + 0.95m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
>1440	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 0.95m_{\tilde{\chi}_1^\pm} + 0.05m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>1140	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (lepton in $\tilde{\chi}_1^\pm$ decay is τ), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
>1110	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (lepton in $\tilde{\chi}_1^\pm$ decay is τ), $m_{\tilde{\ell}} = 0.05m_{\tilde{\chi}_1^\pm} + 0.95m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=6
>1140	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (lepton in $\tilde{\chi}_1^\pm$ decay is τ), $m_{\tilde{\ell}} = 0.95m_{\tilde{\chi}_1^\pm} + 0.05m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=7
> 980	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (leptons in $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decays are τ), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=8
> 905	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (leptons in $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decays are τ), $m_{\tilde{\ell}} = 0.05m_{\tilde{\chi}_1^\pm} + 0.95m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=9
> 875	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (leptons in $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decays are τ), $m_{\tilde{\ell}} = 0.95m_{\tilde{\chi}_1^\pm} + 0.05m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=10
> 650	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=11
> 260	95	⁹	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=12
none 265–305	95	¹⁰	TUMASYAN	22V	CMS	3, 4 b -tagged or 2 large-radius jets, \cancel{E}_T ; higgsino $\tilde{\chi}_2^0 \tilde{\chi}_3^0$ prod. with $\tilde{\chi}_{2,3}^0 \rightarrow H\tilde{\chi}_1^0$; $m_{\tilde{\chi}_1^0} = 1$ GeV	
> 640	95	¹¹	AAD	21BG	ATLS	3 ℓ + \cancel{E}_T , Tchi1n2F, wino cross section, $m_{\tilde{\chi}_1^0} = 0$ GeV	

> 300	95	11 AAD	21BG ATLS	$3\ell + \cancel{E}_T$, Tchi1n2F, wino cross section, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = m_Z$	OCCUR=2
> 240	95	11 AAD	21BG ATLS	$3\ell + \cancel{E}_T$, Tchi1n2F, wino cross section, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10$ GeV	OCCUR=3
> 195	95	11 AAD	21BG ATLS	$3\ell + \cancel{E}_T$, Tchi1n2Ga, higgsino cross section, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10$ GeV	OCCUR=4
> 190	95	11 AAD	21BG ATLS	$3\ell + \cancel{E}_T$, Tchi1n2E, wino cross section, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
>1600	95	12 AAD	21Y ATLS	$\geq 4\ell$, RPV Tchi1n2l with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, $\lambda_{12k} \neq 0$, $m_{\tilde{\chi}_1^0} = 1200$ GeV	
>1100	95	12 AAD	21Y ATLS	$\geq 4\ell$, RPV Tchi1n2l with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, $\lambda_{i33} \neq 0$, $m_{\tilde{\chi}_1^0} = 1000$ GeV	OCCUR=2
> 750	95	13 SIRUNYAN	21M CMS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tchi1n2Fa, $m_{\tilde{\chi}_1^0} < 100$ GeV	
none 400–820	95	14 TUMASYAN	21C CMS	$1 \ell^\pm + 2b\text{-jets} + \cancel{E}_T$, Tchi1n2E, $\tilde{\chi}_1^0 = 200$ GeV	
none 160–820	95	14 TUMASYAN	21C CMS	$1 \ell^\pm + 2b\text{-jets} + \cancel{E}_T$, Tchi1n2E, $\tilde{\chi}_1^0 = 0$ GeV	OCCUR=2
> 380	95	15 AAD	20AN ATLS	$2\gamma + \cancel{E}_T$, Tn1n1A, GMSB	OCCUR=2
> 193	95	16 AAD	20I ATLS	2ℓ (soft), jets, \cancel{E}_T ; Tchi1n2Ga, higgsino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 9.3$ GeV	
> 240	95	17 AAD	20I ATLS	2ℓ (soft), jets, \cancel{E}_T ; Tchi1n2Fa, wino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 7$ GeV	OCCUR=2
> 345	95	18 AAD	20K ATLS	$3\ell + \cancel{E}_T$, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 740	95	19 AAD	20R ATLS	$1\ell + 2b\text{-jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 290	95	20 SIRUNYAN	20AU CMS	soft τ + jet + \cancel{E}_T , Tchi1n2D, wino, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 50$ GeV	
> 680	95	21 AABOUD	19AU ATL	0, 1, 2 or more ℓ , H ($\rightarrow \gamma\gamma, bb, WW^*, ZZ^*, \tau\tau$) (various searches), Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 112	95	22 SIRUNYAN	19BU CMS	$pp \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_2^0 + 2$ jets, $\tilde{\chi}_2^0 \rightarrow \ell^+ \ell^- \tilde{\chi}_1^0$, heavy sleptons, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 1$ GeV, $m_{\tilde{\chi}_2^0} = \tilde{m}_{\tilde{\chi}_1^+}$	
> 215	95	22 SIRUNYAN	19BU CMS	$pp \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_2^0 + 2$ jets, $\tilde{\chi}_2^0 \rightarrow \ell^+ \ell^- \tilde{\chi}_1^0$, heavy sleptons, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 30$ GeV, $m_{\tilde{\chi}_2^0} = \tilde{m}_{\tilde{\chi}_1^+}$	OCCUR=2
> 760	95	23 AABOUD	18AY ATLS	$2\tau + \cancel{E}_T$, Tchi1n2D and $\tilde{\tau}_L$ -only, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>1125	95	24 AABOUD	18BT ATLS	$2,3\ell + \cancel{E}_T$, Tchi1n2C, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 580	95	25 AABOUD	18BT ATLS	$2,3\ell + \cancel{E}_T$, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
none 130–230, 290–880	95	26 AABOUD	18CK ATLS	$2H$ ($\rightarrow bb$) + \cancel{E}_T , Tn1n1A, GMSB	
none 220–600	95	27 AABOUD	18CO ATLS	$2,3\ell + \cancel{E}_T$, recursive jigsaw, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 145	95	28 AABOUD	18R ATLS	2ℓ (soft) + \cancel{E}_T , Tchi1n2G, higgsino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 5$ GeV	
> 175	95	29 AABOUD	18R ATLS	2ℓ (soft) + \cancel{E}_T , Tchi1n2F, wino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10$ GeV	OCCUR=2
>1060	95	30 AABOUD	18U ATLS	$2\gamma + \cancel{E}_T$, GGM, Tchi1chi1A, any NLSP mass	
> 167	95	31 SIRUNYAN	18AJ CMS	2ℓ (soft) + \cancel{E}_T , Tchi1n2G, higgsino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 15$ GeV	

> 710	95	32	SIRUNYAN	18DP CMS	$2\tau + \cancel{E}_T$, Tchi1n2D, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 220–490	95	33	SIRUNYAN	17AW CMS	$1\ell + 2 b\text{-jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 600	95	34	AAD	16AA ATLS	$3,4\ell + \cancel{E}_T$, Tn2n3A, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 670	95	34	AAD	16AA ATLS	$3,4\ell + \cancel{E}_T$, Tn2n3B, $m_{\tilde{\chi}_1^0} < 200$ GeV	OCCUR=2
> 250	95	35	AAD	15BA ATLS	$m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 380	95	36	AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow \tau^\pm \nu \tilde{\chi}_1^0 \tau^\pm \tau^\mp \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 700	95	36	AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow \ell^\pm \nu \tilde{\chi}_1^0 \ell^\pm \ell^\mp \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 345	95	36	AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow W \tilde{\chi}_1^0 Z \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
> 148	95	36	AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow W \tilde{\chi}_1^0 H \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
> 620	95	37	AAD	14X ATLS	$\geq 4\ell^\pm$, $\tilde{\chi}_{2,3}^0 \rightarrow \ell^\pm \ell^\mp \tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 0$ GeV	
		38	AAD	13 ATLS	$3\ell^\pm + \cancel{E}_T$, pMSSM, SMS	
> 62.4	95	40	ABREU	00W DLPH	$\tilde{\chi}_2^0$, $1 \leq \tan\beta \leq 40$, all Δm , all m_0	
> 99.9	95	40	ABREU	00W DLPH	$\tilde{\chi}_3^0$, $1 \leq \tan\beta \leq 40$, all Δm , all m_0	OCCUR=2
> 116.0	95	40	ABREU	00W DLPH	$\tilde{\chi}_4^0$, $1 \leq \tan\beta \leq 40$, all Δm , all m_0	OCCUR=3
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
> 310	95	41	AAD	20AN ATLS	$2\gamma + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 180–355	95	42	AAD	14G ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow W \tilde{\chi}_1^0 Z \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	
		43	KHACHATRY...14I	CMS	$\tilde{\chi}_2^0 \rightarrow (Z, H) \tilde{\chi}_1^0 \tilde{\ell}\ell$, simplified model	
		44	AAD	12AS ATLS	$3\ell^\pm + \cancel{E}_T$, pMSSM	
		45	AAD	12T ATLS	$\ell^\pm \ell^\pm + \cancel{E}_T$, $pp \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$	

¹ AAD 23AE searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with 2 ℓ with same flavour and opposite sign, plus jets and \cancel{E}_T , defining signal region with the dilepton invariant mass both on- and off-shell with respect to the Z boson. No significant excess above the Standard Model predictions is observed. Limits are set on models of strong and electroweak production. For electroweak production, limits are placed on production of mass-degenerate, wino-like $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ with $\tilde{\chi}_2^0 \rightarrow Z \tilde{\chi}_1^0$ and $\tilde{\chi}_1^0 \rightarrow W \tilde{\chi}_1^0$, see figure 15.

NODE=S046ZNO;LINKAGE=IB

² AAD 23CI searched in 139 fb^{-1} of pp collisions for events containing 1 ℓ (e or μ), jets, and \cancel{E}_T . Final states consistent with the production of a diboson system plus \cancel{E}_T were identified also by making use of large-R jet tagging techniques. No excess on top of the Standard Model background was observed. Limits were set on the production of $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm$ (assuming wino cross sections) decaying to $W Z \tilde{\chi}_1^0 \tilde{\chi}_1^0$ or $W W \tilde{\chi}_1^0 \tilde{\chi}_1^0$. See their figure 9.

NODE=S046ZNO;LINKAGE=OB

³ AAD 23CI searched in 139 fb^{-1} of pp collisions for events containing 1 ℓ (e or μ), jets, and \cancel{E}_T . Final states consistent with the production of a boson + Higgs system plus \cancel{E}_T were identified via a BDT. No excess on top of the Standard Model background was observed. Limits were set on the production of degenerate $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ (assuming wino cross sections) decaying into $W h \tilde{\chi}_1^0 \tilde{\chi}_1^0$. See their figure 10.

NODE=S046ZNO;LINKAGE=PB

⁴ AAD 23CP searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with 2 ℓ with same charge plus at least one jet and \cancel{E}_T , defining signal region based on 'stransverse mass' of the dilepton system, \cancel{E}_T significance and effective mass. No significant excess above the Standard Model predictions is observed. Limits are set on the mass of mass-degenerate $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ for the wino-like production of $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ followed by the decay into either $W Z \tilde{\chi}_1^0 \tilde{\chi}_1^0$ or $W h \tilde{\chi}_1^0 \tilde{\chi}_1^0$, see figure 13.

NODE=S046ZNO;LINKAGE=NB

- 5 AAD 23CR searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for RPV SUSY in final states with multiple leptons and b -tagged jets. No significant excess above the Standard Model expectations is observed. Limits are set on the production of electroweakinos (wino or higgsino) that decay via RPV coupling λ'_{i33} to a charged lepton or a neutrino, a b quark, and an additional t or b quark, see their figure 16. A second model addresses direct $\tilde{\mu}_{L,R}$ production and decay to a muon and a bino-like neutralino, which decays in the same way as in the first model, see their figure 17.
- 6 HAYRAPETYAN 23E searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of gluino, top squark and electroweakino pair production in events with at least one photon, multiple jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set in models for strong production, Tglu4D, Tglu4E, Tglu4F and Tstop13, see their figure 9. They also interpret the results in the models for electroweak production, shown in their figure 10. Tchi1n1A assumes wino-like $\tilde{\chi}_1^\pm \tilde{\chi}_1^0$ production, while Tchi1chi1A assumes higgsino-like cross sections and includes $\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm$, $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ and $\tilde{\chi}_{1,2}^0 \tilde{\chi}_1^\pm$ production. For $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ alone no mass point can be excluded in the model Tchi1chi1A, but in another model for $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ production, Tn1n2A.
- 7 TUMASYAN 23B searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino pair production with decays including hadronically decaying bosons, WW , WZ , WH , or ZH , identified with a DNN classifying large-area (AK8) jets. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the nearly mass degenerate wino-like $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tchi1chi1I, Tchi1n2Fb, and Tchi1n2Fc, see their figure 4. They also consider a model that contains both $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm$ production, see their figure 5 (upper). Results are also interpreted in the model THinoBinoA with nearly mass-degenerate higgsino-like $\tilde{\chi}_3^0$, $\tilde{\chi}_2^0$, $\tilde{\chi}_1^\pm$, and a lighter bino-like $\tilde{\chi}_1^0$, see their figure 5 (lower).
- 8 TUMASYAN 22Q searched in up to 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino and top squark pair production with a small mass difference between the produced supersymmetric particles and the lightest neutralino in events with two or three low-momentum leptons and missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the model Tchi1n2F, see their Figure 8. Limits are also set in a higgsino simplified model with both $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ production, where $\tilde{\chi}_2^0 \rightarrow Z \tilde{\chi}_1^0$ and $m_{\tilde{\chi}_1^\pm} = 1/2(m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})$. A model inspired by the pMSSM is used for further interpretations in the case of a higgsino LSP, see their Figure 9. Limits are also set on the mass of the top squark in the models Tstop2 and Tstop3, see their Figure 10.
- 9 TUMASYAN 22S searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino pair production in events with three or four leptons, with up to two hadronically decaying τ leptons, or two same-sign light leptons (e or μ). No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tchi1n2B (in flavory-democratic and tau-enriched or -dominated scenarios), Tchi1n2E, Tchi1n2F, see their Figures 16–20, and on the mass of the higgsino-triplet $\tilde{\chi}_2^0$, $\tilde{\chi}_1^\pm$, and $\tilde{\chi}_1^0$ in the models Tn1n1A, Tn1n1B, and Tn1n1C, see their Figure 21.
- 10 TUMASYAN 22V searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino pair production with decay to two Higgs bosons H , with $H \rightarrow b\bar{b}$, resulting either in 4 resolved b -jets or two large-radius jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tn1n1A, see their Figures 11 and 12, or in a model where higgsino-like nearly mass degenerate $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ are pair produced and each decay to H and a bino-like $\tilde{\chi}_1^0$, see their Figure 13. Limits are also set on the gluino mass in the model Tglu1I, see their Figure 14.
- 11 AAD 21BG searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ in final states with three leptons, with and without assuming the presence of a $Z \rightarrow \ell\ell$ decay. No significant excess above the Standard Model predictions is observed. Limits are set on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2E, Tchi1n2F and Tchi1n2Ga. See their Fig. 16.
- 12 AAD 21Y searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with four or more leptons (electrons, muons and tau-leptons). No significant excess above the Standard Model expectations is observed. Limits are set on Tchi1n12-GGM, and RPV models similar to Tchi1n2I, Tglu1A (with $q = u, d, s, c, b$, with equal branching fractions), and $\tilde{\ell}_L/\tilde{\nu} \rightarrow \ell/\nu \tilde{\chi}_1^0$ (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations), all with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$ via λ_{12k} or λ_{i33} (where $i, k \in 1, 2$), see their Figure 11.
- 13 SIRUNYAN 21M searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$

NODE=S046ZNO;LINKAGE=LB

NODE=S046ZNO;LINKAGE=MB

NODE=S046ZNO;LINKAGE=GB

NODE=S046ZNO;LINKAGE=CB

NODE=S046ZNO;LINKAGE=DB

NODE=S046ZNO;LINKAGE=EB

NODE=S046ZNO;LINKAGE=BB

NODE=S046ZNO;LINKAGE=VA

NODE=S046ZNO;LINKAGE=XA

- mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tsb0t3, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14.
- 14 TUMASYAN 21C searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with one lepton, a Higgs boson decaying to a pair of bottom quarks, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Lower limits are set on the masses of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the simplified model Tchi1n2E, see their Figure 6. NODE=S046ZNO;LINKAGE=YA
- 15 AAD 20AN searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons and missing transverse momentum. Events are further categorised in terms of lepton or jet multiplicity. No significant excess over the expected background is observed. Limits at 95% C.L. are set on the Higgsino mass in the T1n1n1A simplified model, see their Figure 11. NODE=S046ZNO;LINKAGE=UA
- 16 AAD 20I reported on ATLAS searches for electroweak production in models with compressed mass spectra as Tchi1n2Ga. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Events with \cancel{E}_T , two same-flavour, opposite-charge, low-transverse-momentum leptons, and jets from initial-state radiation or characteristic of vector-boson fusion production are selected. Constraints at 95% C.L. are placed in Higgsino models on the mass of the $\tilde{\chi}_2^0$ (the $\tilde{\chi}_1^\pm$ mass is halfway between the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$ masses) at 193 GeV for a mass splitting between $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$ of 9.3 GeV and extend down to a mass splitting of 2.4 GeV at the LEP chargino mass limit. See their Fig. 14(a). NODE=S046ZNO;LINKAGE=RA
- 17 AAD 20I reported on ATLAS searches for electroweak production in models with compressed mass spectra as Tchi1n2Fa. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Events with \cancel{E}_T , two same-flavour, opposite-charge, low-transverse-momentum leptons, and jets from initial-state radiation or characteristic of vector-boson fusion production are selected. Constraints at 95% C.L. are placed in Wino-Bino models on the mass of the $\tilde{\chi}_2^0$ (degenerate with $\tilde{\chi}_1^\pm$) at 240 GeV for a mass splitting between $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^0$ of 7 GeV and extend down to a mass splitting of 1.5 GeV at the LEP chargino mass limit of 92.4 GeV. See their Fig. 14(b,c). NODE=S046ZNO;LINKAGE=SA
- 18 AAD 20K reported on a search for electroweak production in models with mass splittings near the electroweak scale as Tchi1n2F and exploiting three-lepton final state events with an emulated recursive jigsaw reconstruction method. The analysis uses a dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} . Exclusion limits at 95% C.L. are derived on next-to-lightest neutralinos and charginos with masses up to 345 GeV for a massless lightest neutralino, see their Fig. 7. NODE=S046ZNO;LINKAGE=OA
- 19 AAD 20R searched for electroweak production in the model Tchi1n2E, selecting events with a pair of b -tagged jets consistent with those from a Higgs boson decay, either an electron or a muon from the W boson decay and \cancel{E}_T . The analysis uses a dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} . Exclusion limits at 95% C.L. are derived on next-to-lightest neutralinos and charginos with masses up to 740 GeV for a massless lightest neutralino, assuming pure wino cross-sections. See their Fig. 6. NODE=S046ZNO;LINKAGE=PA
- 20 SIRUNYAN 20AU searched in 77.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing one soft, hadronically decaying tau lepton, one energetic jet from initial-state radiation, and large \cancel{E}_T . No excess over the expected background is observed. Limits are derived on the wino mass in the Tchi1n2D simplified model, see their Figure 2. NODE=S046ZNO;LINKAGE=QA
- 21 AABOUD 19AU searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and next-to-lightest neutralinos decaying into lightest neutralinos and a W and a Higgs boson, respectively. Fully hadronic, semileptonic, diphoton, and multilepton (electrons, muons) final states with missing transverse momentum are considered in this search. Observations are consistent with the Standard Model expectations, and 95% confidence-level limits of up to 680 GeV on the chargino/next-to-lightest neutralino masses are set (Tchi1n2E model). See their Figure 14 for an overlay of exclusion contours from all searches. NODE=S046ZNO;LINKAGE=NA
- 22 SIRUNYAN 19BU searched for pair production of gauginos via vector boson fusion assuming the gaugino spectrum is compressed, in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$. The final states explored included zero leptons plus two jets, one lepton plus two jets, and one hadronic tau plus two jets. A similar bound is obtained in the light slepton limit. NODE=S046ZNO;LINKAGE=MA
- 23 AABOUD 18AY searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and neutralinos as in Tchi1n2D models, in events characterised by the presence of at least two hadronically decaying tau leptons and large missing transverse energy. No significant deviation from the expected SM background is observed. Assuming decays via intermediate $\tilde{\tau}_L$ and $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, the observed limits rule out $\tilde{\chi}_2^0$ masses up to 760 GeV for a massless $\tilde{\chi}_1^0$. See their Fig.7 (right). Interpretations are also provided in Fig 8 (bottom) for different assumptions on the ratio between $m_{\tilde{\tau}}$ and $m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0}$. NODE=S046ZNO;LINKAGE=Z
- 24 AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons NODE=S046ZNO;LINKAGE=FA

- in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the next-to-lightest neutralino mass up to 1100 GeV for massless $\tilde{\chi}_1^0$ in the Tchi1n2C simplified model exploiting the 3ℓ signature, see their Figure 8(c).
- 25 AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the next-to-lightest neutralino mass up to 580 GeV for massless $\tilde{\chi}_1^0$ in the Tchi1n2F simplified model exploiting the $2\ell+2$ jets and 3ℓ signatures, see their Figure 8(d). NODE=S046ZNO;LINKAGE=HA
- 26 AABOUD 18CK searched for events with at least 3 b -jets and large missing transverse energy in two datasets of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ of 36.1 fb^{-1} and 24.3 fb^{-1} depending on the trigger requirements. The analyses aimed to reconstruct two Higgs bosons decaying to pairs of b -quarks. No significant excess above the Standard Model expectations is observed. Limits are set on the Higgsino mass in the T1n1n1A simplified model, see their Figure 15(a). Constraints are also presented as a function of the BR of Higgsino decaying into a higgs boson and a gravitino, see their Figure 15(b). NODE=S046ZNO;LINKAGE=KA
- 27 AABOUD 18CO searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of mass-degenerate charginos and next-to-lightest neutralinos in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. The search channels are based on recursive jigsaw reconstruction. Limits are set on the next-to-lightest neutralinos mass up to 600 GeV for massless neutralinos in the Tchi1n2F simplified model exploiting the statistical combination of $2\ell+2$ jets and 3ℓ channels. Next-to-lightest neutralinos masses below 220 GeV are not excluded due to an excess of events above the SM prediction in the dedicated regions. See their Figure 13(d). NODE=S046ZNO;LINKAGE=IA
- 28 AABOUD 18R searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for electroweak production in scenarios with compressed mass spectra in final states with two low-momentum leptons and missing transverse momentum. The data are found to be consistent with the SM prediction. Results are interpreted in Tchi1n2G higgsino models, and $\tilde{\chi}_2^0$ masses are excluded up to 145 GeV for $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$. The exclusion limits extend down to mass splittings of 2.5 GeV, see their Fig. 10 (top). Results are also interpreted in terms of exclusion bounds on the production cross-sections for the NUHM2 scenario as a function of the universal gaugino mass $m_{1/2}$ and $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$, see their Fig. 12. NODE=S046ZNO;LINKAGE=AA
- 29 AABOUD 18R searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for electroweak production in scenarios with compressed mass spectra in final states with two low-momentum leptons and missing transverse momentum. The data are found to be consistent with the SM prediction. Results are interpreted in Tchi1n2F wino models, and $\tilde{\chi}_2^0$ masses are excluded up to 175 GeV for $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10 \text{ GeV}$. The exclusion limits extend down to mass splittings of 2 GeV, see their Fig. 10 (bottom). Results are also interpreted in terms of exclusion bounds on the production cross-sections for the NUHM2 scenario as a function of the universal gaugino mass $m_{1/2}$ and $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$, see their Fig. 12. NODE=S046ZNO;LINKAGE=DA
- 30 AABOUD 18U searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with at least one isolated photon, possibly jets and significant transverse momentum targeting generalised models of gauge-mediated SUSY breaking. No significant excess of events is observed above the SM prediction. Results of the diphoton channel are interpreted in terms of lower limits on the masses of gauginos Tchi1chi1A models, which reach as high as 1.3 TeV. Gaugino masses below 1060 GeV are excluded for any NLSP mass, see their Fig. 10. NODE=S046ZNO;LINKAGE=EA
- 31 SIRUNYAN 18AJ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two low-momentum, oppositely charged leptons (electrons or muons) and \cancel{E}_T . No excess over the expected background is observed. Limits are derived on the wino mass in the Tchi1n2F simplified model, see their Figure 5. Limits are also set on the stop mass in the Tstop10 simplified model, see their Figure 6. Finally, limits are set on the Higgsino mass in the Tchi1n2G simplified model, see Figure 8 and in the pMSSM, see Figure 7. NODE=S046ZNO;LINKAGE=Y
- 32 SIRUNYAN 18DP searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and neutralinos or of chargino pairs in events with a tau lepton pair and significant missing transverse momentum. Both hadronic and leptonic decay modes are considered for the tau lepton. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino mass in the Tchi1chi1D and Tchi1n2 simplified models, see their Figures 14 and 15. Also, excluded stau pair production cross sections are shown in Figures 11, 12, and 13. NODE=S046ZNO;LINKAGE=LA
- 33 SIRUNYAN 17AW searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a charged lepton (electron or muon), two jets identified as originating from a b -quark, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the chargino and the next-to-lightest neutralino in the Tchi1n2E simplified model, see their Figure 6. NODE=S046ZNO;LINKAGE=W
- 34 AAD 16AA summarized and extended ATLAS searches for electroweak supersymmetry in final states containing several charged leptons, \cancel{E}_T , with or without hadronic jets, in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$. The paper reports the results of new interpretations NODE=S046ZNO;LINKAGE=V

- and statistical combinations of previously published analyses, as well as new analyses. Exclusion limits at 95% C.L. are set on mass-degenerate $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ masses in the Tn2n3A and Tn2n3B simplified models. See their Fig. 15.
- 35 AAD 15BA searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of charginos and neutralinos decaying to a final state containing a W boson and a 125 GeV Higgs boson, plus missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of direct chargino and next-to-lightest neutralino production, with the decays $\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ and $\tilde{\chi}_2^0 \rightarrow H \tilde{\chi}_1^0$ having 100% branching fraction, see Fig. 8. A combination of the multiple final states for the Higgs decay yields the best limits (Fig. 8d). NODE=S046ZNO;LINKAGE=Q
- 36 AAD 14H searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of charginos and neutralinos decaying to a final state with three leptons and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of direct chargino and next-to-lightest neutralino production, with decays to the lightest neutralino via either all three generations of leptons, staus only, gauge bosons, or Higgs bosons, see Fig. 7. An interpretation in the pMSSM is also given, see Fig. 8. NODE=S046ZNO;LINKAGE=O
- 37 AAD 14X searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least four leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the neutralino mass in an R-parity conserving simplified model where the decay $\tilde{\chi}_{2,3}^0 \rightarrow \ell^\pm \ell^\mp \tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 10. NODE=S046ZNO;LINKAGE=L
- 38 AAD 13 searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for charginos and neutralinos decaying to a final state with three leptons (e and μ) and missing transverse energy. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in the phenomenological MSSM, see Fig. 2 and 3, and in simplified models, see Fig. 4. For the simplified models with intermediate slepton decays, degenerate $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ masses up to 500 GeV are excluded at 95% C.L. for very large mass differences with the $\tilde{\chi}_1^0$. Supersedes AAD 12AS. NODE=S046ZNO;LINKAGE=GA
- 39 CHATRCHYAN 12BJ searched in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for direct electroweak production of charginos and neutralinos in events with at least two leptons, jets and missing transverse momentum. No significant excesses over the expected SM backgrounds are observed and 95% C.L. limits on the production cross section of $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ pair production were set in a number of simplified models, see Figs. 7 to 12. Most limits are for exactly 3 jets. NODE=S046ZNO;LINKAGE=CH
- 40 ABREU 00W combines data collected at $\sqrt{s}=189 \text{ GeV}$ with results from lower energies. The mass limit is obtained by constraining the MSSM parameter space with gaugino and sfermion mass universality at the GUT scale, using the results of negative direct searches for neutralinos (including cascade decays and $\tilde{\tau}\tau$ final states) from ABREU 01, for charginos from ABREU 00J and ABREU 00T (for all Δm_{\pm}), and for charged sleptons from ABREU 01B. The results hold for the full parameter space defined by all values of M_2 and $|\mu| \leq 2 \text{ TeV}$ with the $\tilde{\chi}_1^0$ as LSP. NODE=S046ZNO;LINKAGE=WA
- 41 AAD 20AN searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons and missing transverse momentum. Events are further categorised in terms of lepton or jet multiplicity. No significant excess over the expected background is observed. Limits at 95% C.L. are derived in Tchi1n2E simplified models. Next-to-lightest neutralinos and charginos with masses up to 310 GeV for a massless lightest neutralino are excluded. See their Fig. 10. NODE=S046ZNO;LINKAGE=TA
- 42 AAD 14G searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of chargino-neutralino pairs, decaying to a final state with two leptons (e and μ) and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of chargino and next-to-lightest neutralino production, with decays to the lightest neutralino via gauge bosons, see Fig. 7. An interpretation in the pMSSM is also given, see Fig. 10. NODE=S046ZNO;LINKAGE=P
- 43 KHACHATRYAN 14I searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of charginos and neutralinos decaying to a final state with three leptons (e or μ) and missing transverse momentum, or with a Z -boson, dijets and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models, see Figs. 12–16. NODE=S046ZNO;LINKAGE=N
- 44 AAD 12AS searched in 2.06 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for charginos and neutralinos decaying to a final state with three leptons (e and μ) and missing transverse energy. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in the phenomenological MSSM, see Fig. 2 (top), and in simplified models, see Fig. 2 (bottom). NODE=S046ZNO;LINKAGE=GL
- 45 AAD 12T looked in 1 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for the production of supersymmetric particles decaying into final states with missing transverse momentum and exactly two isolated leptons (e or μ). Same-sign dilepton events were separately studied. Additionally, in opposite-sign events, a search was made for an excess of same-flavor over different-flavor lepton pairs. No excess over the expected background is observed and limits are placed on the effective production cross section of opposite-sign dilepton events with $\cancel{E}_T > 250 \text{ GeV}$ and on same-sign dilepton events with $\cancel{E}_T > 100 \text{ GeV}$. The latter limit is interpreted in a simplified electroweak gaugino production model. NODE=S046ZNO;LINKAGE=AG

$\tilde{\chi}_1^\pm, \tilde{\chi}_2^\pm$ (Charginos) mass limits

Charginos are unknown mixtures of w -inos and charged higgsinos (the supersymmetric partners of W and Higgs bosons). A lower mass limit for the lightest chargino ($\tilde{\chi}_1^\pm$) of approximately 45 GeV, independent of the field composition and of the decay mode, has been obtained by the LEP experiments from the analysis of the Z width and decays. These results, as well as other now superseded limits from e^+e^- collisions at energies below 136 GeV, and from hadronic collisions, can be found in the 1998 Edition (The European Physical Journal **C3** 1 (1998)) of this Review.

Unless otherwise stated, results in this section assume spectra, production rates, decay modes and branching ratios as evaluated in the MSSM, with gaugino and sfermion mass unification at the GUT scale. These papers generally study production of $\tilde{\chi}_1^0\tilde{\chi}_2^0$, $\tilde{\chi}_1^+\tilde{\chi}_1^-$ and (in the case of hadronic collisions) $\tilde{\chi}_1^+\tilde{\chi}_2^0$ pairs, including the effects of cascade decays. The mass limits on $\tilde{\chi}_1^\pm$ are either direct, or follow indirectly from the constraints set by the non-observation of $\tilde{\chi}_2^0$ states on the gaugino and higgsino MSSM parameters M_2 and μ . For generic values of the MSSM parameters, limits from high-energy e^+e^- collisions coincide with the highest value of the mass allowed by phase-space, namely $m_{\tilde{\chi}_1^\pm} \lesssim \sqrt{s}/2$. The still unpublished combination of the results of the four LEP collaborations from the 2000 run of LEP2 at \sqrt{s} up to $\simeq 209$ GeV yields a lower mass limit of 103.5 GeV valid for general MSSM models. The limits become however weaker in certain regions of the MSSM parameter space where the detection efficiencies or production cross sections are suppressed. For example, this may happen when: (i) the mass differences $\Delta m_+ = m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}$ or $\Delta m_\nu = m_{\tilde{\chi}_1^\pm} - m_{\tilde{\nu}}$ are very small, and the detection efficiency is reduced; (ii) the electron sneutrino mass is small, and the $\tilde{\chi}_1^\pm$ production rate is suppressed due to a destructive interference between s and t channel exchange diagrams. The regions of MSSM parameter space where the following limits are valid are indicated in the comment lines or in the footnotes.

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	NODE=S046WNO;CHECK LIMITS
> 820	95	1 AAD	23AE ATLS	2 SFOS ℓ , jets, \cancel{E}_T , Tchi1n2Fa, $m_{\tilde{\chi}_1^0} = 1$ GeV	
none 260–420	95	2 AAD	23CI ATLS	1 ℓ + jets + \cancel{E}_T , Tchi1n2J, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 260–520	95	2 AAD	23CI ATLS	1 ℓ + jets + \cancel{E}_T , Tchi1chi1J, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 230	95	3 AAD	23CI ATLS	1 ℓ + jets + \cancel{E}_T , Tchi1n2E, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 133$ GeV	OCCUR=3
> 450	95	3 AAD	23CI ATLS	1 ℓ + jets + \cancel{E}_T , Tchi1n2E, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 260$ GeV	OCCUR=4
none 200–250	95	4 AAD	23CP ATLS	2 same-sign ℓ , Tchi1n2F, wino-bino, $m_{\tilde{\chi}_1^0} = 1$ GeV	
> 525	95	4 AAD	23CP ATLS	2 same-sign ℓ , Tchi1n2E, wino-bino, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=2
none 200–585	95	5 AAD	23CR ATLS	RPV, 2 same-sign, 3, 4 ℓ , 1, 2 b -jets, higgsino production with $\tilde{\chi} \rightarrow b + \ell/\nu + t/b$ via λ'_{j33} coupling	
none 200–670	95	5 AAD	23CR ATLS	RPV, 2 same-sign, 3, 4 ℓ , 1, 2 b -jets, wino production with $\tilde{\chi} \rightarrow b + \ell/\nu + t/b$ via λ'_{j33} coupling	OCCUR=2
> 150	95	6 AAD	23M ATLS	2 ℓ , Tchi1chi1H, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} > 110$ GeV	
> 104	95	6 AAD	23M ATLS	2 ℓ , Tchi1chi1H, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} > 90$ GeV	OCCUR=2
>1230	95	7 HAYRAPETY...23E	CMS	γ + jets + \cancel{E}_T , Tchi1n1A	
>1050	95	7 HAYRAPETY...23E	CMS	γ + jets + \cancel{E}_T , Tchi1chi1A	OCCUR=2
none 290–670	95	8 TUMASYAN	23B CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , Tchi1chi1l, $m_{\tilde{\chi}_1^0} = 1$ GeV	
none 230–760	95	8 TUMASYAN	23B CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , Tchi1n2Fb, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=2

none 240–970	95	⁸ TUMASYAN	23B	CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , Tchi1n2Fc, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=3
none 300–650	95	⁸ TUMASYAN	23B	CMS	2 AK8 jets + 2–6 AK4 jets + \cancel{E}_T , THinoBinoA, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=4
> 275	95	⁹ TUMASYAN	22Q	CMS	2 or 3 ℓ (soft), \cancel{E}_T ; Tchi1n2F, wino-bino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10$ GeV	OCCUR=2
> 205	95	⁹ TUMASYAN	22Q	CMS	2 or 3 ℓ (soft), \cancel{E}_T ; higgsino model with $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ prod., $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 7.5$ GeV	OCCUR=3
> 150	95	⁹ TUMASYAN	22Q	CMS	2 or 3 ℓ (soft), \cancel{E}_T ; higgsino model with $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ prod., $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 3$ GeV	OCCUR=3
>1450	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 850$ GeV	OCCUR=2
>1360	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
>1290	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 0.05m_{\tilde{\chi}_1^\pm} + 0.95m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>1440	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (flavor-democratic), $m_{\tilde{\ell}} = 0.95m_{\tilde{\chi}_1^\pm} + 0.05m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
>1140	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (lepton in $\tilde{\chi}_1^\pm$ decay is τ), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=6
>1110	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (lepton in $\tilde{\chi}_1^\pm$ decay is τ), $m_{\tilde{\ell}} = 0.05m_{\tilde{\chi}_1^\pm} + 0.95m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=7
>1140	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (lepton in $\tilde{\chi}_1^\pm$ decay is τ), $m_{\tilde{\ell}} = 0.95m_{\tilde{\chi}_1^\pm} + 0.05m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=8
> 980	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (leptons in $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decays are τ), $m_{\tilde{\ell}} = 1/2(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=9
> 905	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (leptons in $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decays are τ), $m_{\tilde{\ell}} = 0.05m_{\tilde{\chi}_1^\pm} + 0.95m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=10
> 875	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2B (leptons in $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ decays are τ), $m_{\tilde{\ell}} = 0.95m_{\tilde{\chi}_1^\pm} + 0.05m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=11
> 650	95	¹⁰ TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	

> 260	95	10	TUMASYAN	22S	CMS	2 same-sign e or μ , 3 or 4 leptons, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=12
>1080	95	11	AAD	21AX	ATLS	jets + large-R jets + \cancel{E}_T , TWinoBinoA, nearly independent of $B(\tilde{\chi}_2^0 \rightarrow Z\tilde{\chi}_1^0)$, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>1060	95	11	AAD	21AX	ATLS	jets + large-R jets + \cancel{E}_T , TWinoHinoA, $\tan\beta = 10$, $\mu > 0$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 900	95	11	AAD	21AX	ATLS	jets + large-R jets + \cancel{E}_T , THinoBinoA, nearly independent of $B(\tilde{\chi}_2^0 \rightarrow Z\tilde{\chi}_1^0)$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
> 900	95	11	AAD	21AX	ATLS	jets + large-R jets + \cancel{E}_T , THinoWinoA, $\tan\beta = 10$, $\mu > 0$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>1060	95	11	AAD	21AX	ATLS	jets + large-R jets + \cancel{E}_T , Tchi1n2E, full hadronic final state, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
> 960	95	11	AAD	21AX	ATLS	jets + large-R jets + \cancel{E}_T , Tchi1n2Fb, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=6
none 620–740	95	11	AAD	21AX	ATLS	jets + large-R jets + \cancel{E}_T , Tchi1chi1l, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=7
> 640	95	12	AAD	21BG	ATLS	$3\ell + \cancel{E}_T$, Tchi1n2F, wino cross section, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 300	95	12	AAD	21BG	ATLS	$3\ell + \cancel{E}_T$, Tchi1n2F, wino cross section, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = m_Z$	OCCUR=2
> 240	95	12	AAD	21BG	ATLS	$3\ell + \cancel{E}_T$, Tchi1n2F, wino cross section, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10$ GeV	OCCUR=3
> 190	95	12	AAD	21BG	ATLS	$3\ell + \cancel{E}_T$, Tchi1n2E, wino cross section, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>1100	95	13	AAD	21E	ATLS	3ℓ , $Z\ell$ resonances, Twinol-SPBL, RPV, $B(\tilde{\chi}_1^\pm \rightarrow Ze)$ $= B(\tilde{\chi}_1^0 \rightarrow Z\nu) = 1$	OCCUR=2
>1050	95	13	AAD	21E	ATLS	3ℓ , $Z\ell$ resonances, Twinol-SPBL, RPV, $B(\tilde{\chi}_1^\pm \rightarrow Z\mu)$ $= B(\tilde{\chi}_1^0 \rightarrow Z\nu) = 1$	OCCUR=3
> 625	95	13	AAD	21E	ATLS	3ℓ , $Z\ell$ resonances, Twinol-SPBL, RPV, $B(\tilde{\chi}_1^\pm \rightarrow Z\tau)$ $= B(\tilde{\chi}_1^0 \rightarrow Z\nu) = 1$	OCCUR=4
> 975	95	13	AAD	21E	ATLS	3ℓ , $Z\ell$ resonances, Twinol-SPBL, RPV, $B(\tilde{\chi}_1^\pm \rightarrow Z\ell)$ $= B(\tilde{\chi}_1^0 \rightarrow Z\nu) = 1$ and $\ell = e, \mu, \tau$	
>1600	95	14	AAD	21Y	ATLS	$\geq 4\ell$, RPV Tchi1n2l with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, $\lambda_{12k} \neq 0$, $m_{\tilde{\chi}_1^0} = 1200$ GeV	OCCUR=2
>1100	95	14	AAD	21Y	ATLS	$\geq 4\ell$, RPV Tchi1n2l with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, $\lambda_{i33} \neq 0$, $m_{\tilde{\chi}_1^0} = 1000$ GeV	
> 750	95	15	SIRUNYAN	21M	CMS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tchi1n2Fa, $m_{\tilde{\chi}_1^0} < 100$ GeV	
none 400–820	95	16	TUMASYAN	21C	CMS	$1\ell^\pm + 2b$ -jets + \cancel{E}_T , Tchi1n2E, $\tilde{\chi}_1^0 = 200$ GeV	
none 160–820	95	16	TUMASYAN	21C	CMS	$1\ell^\pm + 2b$ -jets + \cancel{E}_T , Tchi1n2E, $\tilde{\chi}_1^0 = 0$ GeV	OCCUR=2
> 380	95	17	AAD	20AN	ATLS	$2\gamma + \cancel{E}_T$, Tn1n1A, GMSB	OCCUR=2
> 240	95	18	AAD	20I	ATLS	2ℓ (soft), jets, \cancel{E}_T ; Tchi1n2Fa, wino, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 7$ GeV	
> 345	95	19	AAD	20K	ATLS	$3\ell + \cancel{E}_T$, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 420	95	20	AAD	20O	ATLS	$2\ell + \cancel{E}_T$, Tchi1chi1H, $m_{\tilde{\chi}_1^0} = 0$ GeV	

>1000	95	21 AAD	20O ATLS	$2\ell + \cancel{E}_T$, Tchi1chi1C, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 740	95	22 AAD	20R ATLS	$1\ell + 2b\text{-jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 290	95	23 SIRUNYAN	20AU CMS	soft $\tau + \text{jet} + \cancel{E}_T$, Tchi1n2D, wino, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 50$ GeV	
>1050	95	24 SIRUNYAN	20B CMS	$\geq 1\gamma + \cancel{E}_T$, Tchi1chi1F, $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$	
> 825	95	24 SIRUNYAN	20B CMS	$\geq 1\gamma + \cancel{E}_T$, Tchi1chi1G, $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 + \text{soft}$	OCCUR=2
> 840	95	24 SIRUNYAN	20B CMS	$\geq 1\gamma + \cancel{E}_T$, Tchi1n12-GGM, 120 GeV $< m_{\tilde{\chi}_1^0} < 720$ GeV	OCCUR=3
> 680	95	25 AABOUD	19AU ATL	0, 1, 2 or more ℓ , $H (\rightarrow \gamma\gamma, bb, WW^*, ZZ^*, \tau\tau)$ (various searches), Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 112	95	26 SIRUNYAN	19BU CMS	$pp \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_2^0 + 2 \text{ jets}$, $\tilde{\chi}_1^+ \rightarrow \ell^+ \nu \tilde{\chi}_1^0$, heavy sleptons, $m_{\tilde{\chi}_1^+} - m_{\tilde{\chi}_1^0} = 1$ GeV, $m_{\tilde{\chi}_1^+} = m_{\tilde{\chi}_2^0}$	
> 215	95	26 SIRUNYAN	19BU CMS	$pp \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_2^0 + 2 \text{ jets}$, $\tilde{\chi}_1^+ \rightarrow \ell^+ \nu \tilde{\chi}_1^0$, heavy sleptons, $m_{\tilde{\chi}_1^+} - m_{\tilde{\chi}_1^0} = 30$ GeV, $m_{\tilde{\chi}_1^+} = m_{\tilde{\chi}_2^0}$	OCCUR=2
> 235	95	27 SIRUNYAN	19CI CMS	$\geq 1 H (\rightarrow \gamma\gamma) + \text{jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 1$ GeV	
> 930	95	28 SIRUNYAN	19K CMS	$\gamma + \text{lepton} + \cancel{E}_T$, Tchi1n1A	
> 630	95	29 AABOUD	18AY ATLS	$2\tau + \cancel{E}_T$, Tchi1chi1D and $\tilde{\tau}_L$ -only, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 760	95	30 AABOUD	18AY ATLS	$2\tau + \cancel{E}_T$, Tchi1n2D and $\tilde{\tau}_L$ -only, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 740	95	31 AABOUD	18BT ATLS	$2\ell + \cancel{E}_T$, Tchi1chi1C, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>1125	95	32 AABOUD	18BT ATLS	$2,3\ell + \cancel{E}_T$, Tchi1n2C, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 580	95	33 AABOUD	18BT ATLS	$2,3\ell + \cancel{E}_T$, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
none 130–230, 290–880 none 220–600	95	34 AABOUD	18CK ATLS	$2H (\rightarrow bb) + \cancel{E}_T$, Tn1n1A, GMSB	
	95	35 AABOUD	18CO ATLS	$2,3\ell + \cancel{E}_T$, recursive jigsaw, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 175	95	36 AABOUD	18R ATLS	$2\ell (\text{soft}) + \cancel{E}_T$, Tchi1n2F, wino, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 10$ GeV	
> 145	95	37 AABOUD	18R ATLS	$2\ell (\text{soft}) + \cancel{E}_T$, Tchi1n2G, higgsino, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5$ GeV	OCCUR=2
>1060	95	38 AABOUD	18U ATLS	$2\gamma + \cancel{E}_T$, GGM, Tchi1chi1A, any NLSP mass	
>1400	95	39 AABOUD	18Z ATLS	$\geq 4\ell$, RPV, $\lambda_{12k} \neq 0$, $m_{\tilde{\chi}_1^0} > 500$ GeV	
>1320	95	39 AABOUD	18Z ATLS	$\geq 4\ell$, RPV, $\lambda_{12k} \neq 0$, $m_{\tilde{\chi}_1^0} > 50$ GeV	OCCUR=2
> 980	95	39 AABOUD	18Z ATLS	$\geq 4\ell$, RPV, $\lambda_{j33} \neq 0$, 400 GeV $< m_{\tilde{\chi}_1^0} < 700$ GeV	OCCUR=3
> 980	95	40 SIRUNYAN	18AA CMS	$\geq 1\gamma + \cancel{E}_T$, GGM, wino-like $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ pair production, nearly degenerate wino and bino masses	
> 780	95	40 SIRUNYAN	18AA CMS	$\geq 1\gamma + \cancel{E}_T$, Tchi1n1A	OCCUR=2
> 950	95	40 SIRUNYAN	18AA CMS	$\geq 1\gamma + \cancel{E}_T$, Tchi1chi1A	OCCUR=3
> 230	95	41 SIRUNYAN	18AJ CMS	$2\ell (\text{soft}) + \cancel{E}_T$, Tchi1n2F, wino, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 20$ GeV	

>1150	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2A, $m_{\tilde{\ell}}^-$ $= m_{\tilde{\nu}} = m_{\tilde{\chi}_1^0} + 0.5 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>1120	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2A, $m_{\tilde{\ell}}^-$ $= m_{\tilde{\nu}} = m_{\tilde{\chi}_1^0} + 0.05 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
>1050	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2A, $m_{\tilde{\ell}}^-$ $= m_{\tilde{\nu}} = m_{\tilde{\chi}_1^0} + 0.95 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
>1080	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2H, $m_{\tilde{\ell}}^-$ $= m_{\tilde{\chi}_1^0} + 0.5 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>1030	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2H, $m_{\tilde{\ell}}^-$ $= m_{\tilde{\chi}_1^0} + 0.05 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
>1050	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2H, $m_{\tilde{\ell}}^-$ $= m_{\tilde{\chi}_1^0} + 0.95 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=6
> 625	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2D, $m_{\tilde{\tau}}^-$ $= m_{\tilde{\chi}_1^0} + 0.5 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=7
> 180	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=8
> 450	95	42 SIRUNYAN	18AO CMS	$\ell^\pm \ell^\pm$ or $\geq 3\ell$, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=9
> 480	95	43 SIRUNYAN	18AP CMS	Combination of searches, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 650	95	43 SIRUNYAN	18AP CMS	Combination of searches, Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 535	95	43 SIRUNYAN	18AP CMS	Combination of searches, Tchi1n2I, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
none 160–610	95	44 SIRUNYAN	18AR CMS	$\ell^\pm \ell^\mp$ + jets + \cancel{E}_T , Tchi1n2F, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 170–200	95	45 SIRUNYAN	18DN CMS	$\ell^\pm \ell^\mp$, Tchi1chi1E, $m_{\tilde{\chi}_1^0} = 1$ GeV	
> 810	95	45 SIRUNYAN	18DN CMS	$\ell^\pm \ell^\mp$, Tchi1chi1C, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 630	95	46 SIRUNYAN	18DP CMS	$2\tau + \cancel{E}_T$, Tchi1chi1D, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 710	95	46 SIRUNYAN	18DP CMS	$2\tau + \cancel{E}_T$, Tchi1n2D, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 170	95	47 SIRUNYAN	18X CMS	$\geq 1 H (\rightarrow \gamma\gamma) + \text{jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_1^0} < 25$ GeV	
> 420	95	48 KHACHATRY...17L	CMS	$2\tau + \cancel{E}_T$, Tchi1chi1C and $\tilde{\tau}$ -only, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 220–490	95	49 SIRUNYAN	17AW CMS	$1\ell + 2b\text{-jets} + \cancel{E}_T$, Tchi1n2E, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 500	95	50 AAD	16AA ATLS	$2\ell^\pm + \cancel{E}_T$, Tchi1chi1B, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 220	95	50 AAD	16AA ATLS	$2\ell^\pm + \cancel{E}_T$, Tchi1chi1C, low Δm for $\tilde{\chi}_1^\pm, \tilde{\chi}_1^0$	OCCUR=2
> 700	95	51 AAD	16AA ATLS	$3,4\ell + \cancel{E}_T$, Tchi1n2B, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
> 700	95	51 AAD	16AA ATLS	$3,4\ell + \cancel{E}_T$, Tchi1n2C, $m_{\tilde{\ell}}^- = m_{\tilde{\chi}_1^0} + 0.5$ (or 0.95) $(m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})$	OCCUR=5

> 400	95	51 AAD	16AA ATLS	2 hadronic $\tau + \cancel{E}_T$ & $3\ell + \cancel{E}_T$ combination, $T_{\text{chi1n2D}}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=6
> 540	95	52 KHACHATRY...16R	CMS	$\geq 1\gamma + 1 e$ or $\mu + \cancel{E}_T$, T_{chi1n1A}	
> 250	95	53 AAD	15BA ATLS	$m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0} = 0$ GeV	
> 590	95	54 AAD	15CA ATLS	$\geq 2\gamma + \cancel{E}_T$, GGM, bino-like NLSP, any NLSP mass	
none 124–361	95	54 AAD	15CA ATLS	$\geq 1\gamma + e, \mu + \cancel{E}_T$, GGM, wino-like NLSP	OCCUR=2
> 700	95	55 AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow \ell^\pm \nu \tilde{\chi}_1^0 \ell^\mp \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0} = 0$ GeV	
> 345	95	55 AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow W \tilde{\chi}_1^0 Z \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 148	95	55 AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow W \tilde{\chi}_1^0 H \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
> 380	95	55 AAD	14H ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow \tau^\pm \nu \tilde{\chi}_1^0 \tau^\mp \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
> 750	95	56 AAD	14X ATLS	RPV, $\geq 4\ell^\pm, \tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$	
> 210	95	57 KHACHATRY...14L	CMS	$\tilde{\chi}_2^0 \rightarrow H \tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ simplified models, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm}, m_{\tilde{\chi}_1^0} = 0$ GeV	
> 540	95	58 AAD	13 ATLS	$3\ell^\pm + \cancel{E}_T$, pMSSM, SMS	
		59 AAD	13B ATLS	$2\ell^\pm + \cancel{E}_T$, pMSSM, SMS	
		60 AAD	12CT ATLS	$\geq 4\ell^\pm$, RPV, $m_{\tilde{\chi}_1^0} > 300$ GeV	
> 94	95	61 CHATRCHYAN	12BJ CMS	$\geq 2\ell$, jets + \cancel{E}_T , $pp \rightarrow \tilde{\chi}_1^\pm \tilde{\chi}_2^0$	
		62 ABDALLAH	03M DLPH	$\tilde{\chi}_1^\pm, \tan\beta \leq 40, \Delta m_+ > 3$ GeV, all m_0	OCCUR=4
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
> 310	95	63 AAD	20AN ATLS	$2\gamma + \cancel{E}_T$, $T_{\text{chi1n2E}}, m_{\tilde{\chi}_1^0} = 0$ GeV	
> 570	95	64 KHACHATRY...16AA	CMS	$\geq 1\gamma + \text{jets} + \cancel{E}_T$, $T_{\text{chi1chi1A}}$	
> 680	95	64 KHACHATRY...16AA	CMS	$\geq 1\gamma + \text{jets} + \cancel{E}_T$, T_{chi1n1A}	OCCUR=2
> 710	95	64 KHACHATRY...16AA	CMS	$\geq 1\gamma + \text{jets} + \cancel{E}_T$, GGM, $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ pair production, wino-like NLSP	OCCUR=3
>1000	95	65 KHACHATRY...16R	CMS	$\geq 1\gamma + 1 e$ or $\mu + \cancel{E}_T$, $T_{\text{glu1F}}, m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0} > 200$ GeV	OCCUR=2
> 307	95	66 KHACHATRY...16Y	CMS	1,2 soft $\ell^\pm + \text{jets} + \cancel{E}_T$, $T_{\text{chi1n2A}}, m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 20$ GeV	OCCUR=2
> 410	95	67 AAD	14AV ATLS	$\geq 2\tau + \cancel{E}_T$, direct $\tilde{\chi}_1^\pm \tilde{\chi}_2^0, \tilde{\chi}_1^\pm \tilde{\chi}_1^\mp$ production, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm}, m_{\tilde{\chi}_1^0} = 0$ GeV	
> 345	95	68 AAD	14AV ATLS	$\geq 2\tau + \cancel{E}_T$, direct $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp$ production, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
none 100–105, 120–135, 145–160	95	69 AAD	14G ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp \rightarrow W^+ \tilde{\chi}_1^0 W^- \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 140–465	95	69 AAD	14G ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp \rightarrow \ell^+ \nu \tilde{\chi}_1^0 \ell^- \bar{\nu} \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
none 180–355	95	69 AAD	14G ATLS	$\tilde{\chi}_1^\pm \tilde{\chi}_2^0 \rightarrow W \tilde{\chi}_1^0 Z \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3

> 168	95	70 AALTONEN	14	CDF	$3\ell^\pm + \cancel{E}_T, \tilde{\chi}_1^\pm \rightarrow \ell\nu\tilde{\chi}_1^0$, mSUGRA with $m_0=60$ GeV
		71 KHACHATRY...	14I	CMS	$\tilde{\chi}_1^\pm \rightarrow W\tilde{\chi}_1^0, \ell\tilde{\nu}, \ell\nu$, simplified model
		72 AALTONEN	13Q	CDF	$\tilde{\chi}_1^\pm \rightarrow \tau X$, simplified gravity- and gauge-mediated models
		73 AAD	12AS	ATLS	$3\ell^\pm + \cancel{E}_T$, pMSSM
		74 AAD	12T	ATLS	$\ell^\pm\ell^\mp + \cancel{E}_T, \ell^\pm\ell^\pm + \cancel{E}_T$, $pp \rightarrow \tilde{\chi}_1^\pm\tilde{\chi}_2^0$
		75 CHATRCHYAN	11B	CMS	$\tilde{W}^0 \rightarrow \gamma\tilde{G}, \tilde{W}^\pm \rightarrow \ell^\pm\tilde{G}$, GMSB
> 163	95	76 CHATRCHYAN	11V	CMS	$\tan\beta=3, m_0=60$ GeV, $A_0=0$, $\mu > 0$

¹ AAD 23AE searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with 2 ℓ with same flavour and opposite sign, plus jets and \cancel{E}_T , defining signal region with the dilepton invariant mass both on- and off-shell with respect to the Z boson. No significant excess above the Standard Model predictions is observed. Limits are set on models of strong and electroweak production. For electroweak production, limits are placed on production of mass-degenerate, wino-like $\tilde{\chi}_2^0\tilde{\chi}_1^0$ with $\tilde{\chi}_2^0 \rightarrow Z\tilde{\chi}_1^0$ and $\tilde{\chi}_1^0 \rightarrow W\tilde{\chi}_1^0$, see figure 15.

NODE=S046WNO;LINKAGE=ID

² AAD 23CI searched in 139 fb^{-1} of pp collisions for events containing 1 ℓ (e or μ), jets, and \cancel{E}_T . Final states consistent with the production of a diboson system plus \cancel{E}_T were identified also by making use of large-R jet tagging techniques. No excess on top of the Standard Model background was observed. Limits were set on the production of $\tilde{\chi}_1^\pm\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm$ (assuming wino cross sections) decaying to $WZ\tilde{\chi}_1^0\tilde{\chi}_1^0$ or $WW\tilde{\chi}_1^0\tilde{\chi}_1^0$. See their figure 9.

NODE=S046WNO;LINKAGE=MD

³ AAD 23CI searched in 139 fb^{-1} of pp collisions for events containing 1 ℓ (e or μ), jets, and \cancel{E}_T . Final states consistent with the production of a boson + Higgs system plus \cancel{E}_T were identified via a BDT. No excess on top of the Standard Model background was observed. Limits were set on the production of degenerate $\tilde{\chi}_1^\pm\tilde{\chi}_2^0$ (assuming wino cross sections) decaying into $Wh\tilde{\chi}_1^0\tilde{\chi}_1^0$. See their figure 10.

NODE=S046WNO;LINKAGE=ND

⁴ AAD 23CP searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with 2 ℓ with same charge plus at least one jet and \cancel{E}_T , defining signal region based on 'stransverse mass' of the dilepton system, \cancel{E}_T significance and effective mass. No significant excess above the Standard Model predictions is observed. Limits are set on the mass of mass-degenerate $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ for the wino-like production of $\tilde{\chi}_1^\pm\tilde{\chi}_2^0$ followed by the decay into either $WZ\tilde{\chi}_1^0\tilde{\chi}_1^0$ or $Wh\tilde{\chi}_1^0\tilde{\chi}_1^0$, see figure 13.

NODE=S046WNO;LINKAGE=LD

⁵ AAD 23CR searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for RPV SUSY in final states with multiple leptons and b -tagged jets. No significant excess above the Standard Model expectations is observed. Limits are set on the production of electroweakinos (wino or higgsino) that decay via RPV coupling λ'_{i33} to a charged lepton or a neutrino, a b quark, and an additional t or b quark, see their figure 16. A second model addresses direct $\tilde{\mu}_{L,R}$ production and decay to a muon and a bino-like neutralino, which decays in the same way as in the first model, see their figure 17.

NODE=S046WNO;LINKAGE=JD

⁶ AAD 23M searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for $\tilde{\chi}_1^\pm$ pair production, followed by $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0 \rightarrow \ell^\pm\nu\tilde{\chi}_1^0$ in events with two leptons. The focus is on models where $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}$ is close to the W mass. No significant excess above the Standard Model predictions is observed. Limits are set on the $\tilde{\chi}_1^\pm$ mass as a function of $m_{\tilde{\chi}_1^0}$, see Figure 9.

NODE=S046WNO;LINKAGE=GD

⁷ HAYRAPETYAN 23E searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of gluino, top squark and electroweakino pair production in events with at least one photon, multiple jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set in models for strong production, Tglu4D, Tglu4E, Tglu4F and Tstop13, see their figure 9. They also interpret the results in the models for electroweak production, shown in their figure 10. Tchi1n1A assumes wino-like $\tilde{\chi}_1^\pm\tilde{\chi}_1^0$ production, while Tchi1chi1A assumes higgsino-like cross sections and includes $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^0\tilde{\chi}_2^0$ and $\tilde{\chi}_{1,2}^0\tilde{\chi}_1^\pm$ production. For $\tilde{\chi}_1^0\tilde{\chi}_2^0$ alone no mass point can be excluded in the model Tchi1chi1A, but in another model for $\tilde{\chi}_1^0\tilde{\chi}_2^0$ production, Tn1n2A.

NODE=S046WNO;LINKAGE=KD

⁸ TUMASYAN 23B searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of electroweakino pair production with decays including hadronically decaying bosons, WW , WZ , WH , or ZH , identified with a DNN classifying large-area (AK8) jets. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the nearly mass degenerate wino-like $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tchi1chi1I, Tchi1n2Fb, and Tchi1n2Fc, see their figure 4. They also consider a model that contains both $\tilde{\chi}_2^0\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^\pm\tilde{\chi}_1^\pm$ production, see their figure 5 (upper). Results are also interpreted in the model THinoBinoA with nearly mass-degenerate higgsino-like $\tilde{\chi}_3^0, \tilde{\chi}_2^0, \tilde{\chi}_1^\pm$, and a lighter bino-like $\tilde{\chi}_1^0$, see their figure 5 (lower).

NODE=S046WNO;LINKAGE=HD

- 9 TUMASYAN 22Q searched in up to 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino and top squark pair production with a small mass difference between the produced supersymmetric particles and the lightest neutralino in events with two or three low-momentum leptons and missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the model Tchi1n2F, see their Figure 8. Limits are also set in a higgsino simplified model with both $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ production, where $\tilde{\chi}_2^0 \rightarrow Z \tilde{\chi}_1^0$ and $m_{\tilde{\chi}_1^\pm} = 1/2(m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})$. A model inspired by the pMSSM is used for further interpretations in the case of a higgsino LSP, see their Figure 9. Limits are also set on the mass of the top squark in the models Tstop2 and Tstop3, see their Figure 10.
- 10 TUMASYAN 22S searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino pair production in events with three or four leptons, with up to two hadronically decaying τ leptons, or two same-sign light leptons (e or μ). No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tchi1n2B (in flavory-democratic and tau-enriched or -dominated scenarios), Tchi1n2E, Tchi1n2F, see their Figures 16–20, and on the mass of the higgsino-triplet $\tilde{\chi}_2^0$, $\tilde{\chi}_1^\pm$, and $\tilde{\chi}_1^0$ in the models Tn1n1A, Tn1n1B, and Tn1n1C, see their Figure 21.
- 11 AAD 21AX searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of electroweakinos decaying to the LSP via the emission of Standard Model bosons (Higgs, W , Z) decaying into hadrons. The final state in all cases characterised by the presence of \cancel{E}_T , jets, and large-R jets tagged according to the boson of interest. Different assumptions (Higgsino, Wino, Bino) are made for the pair produced electroweakinos and for the LSP multiplet. No significant excess above the Standard Model predictions is observed. Limits are set on the electroweakino masses as a function of the model parameters (in particular $m_{\tilde{\chi}_1^0}$). See Figs. 12, 14, 15.
- 12 AAD 21BG searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ in final states with three leptons, with and without assuming the presence of a $Z \rightarrow \ell\ell$ decay. No significant excess above the Standard Model predictions is observed. Limits are set on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2E, Tchi1n2F and Tchi1n2Ga. See their Fig. 16.
- 13 AAD 21E searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for production of wino-like $\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^\pm \tilde{\chi}_1^0$, followed by the RPV decay of $\tilde{\chi}_1^\pm$ into $Z\ell$, $H\ell$ or $W\nu$ and of $\tilde{\chi}_1^0$ into $Z\nu$, $H\nu$ or $W\ell$, in events with three leptons, looking for $Z\ell$ resonances. No significant excess above the Standard Model predictions is observed. Limits are set on the common $m_{\tilde{\chi}_1^\pm}/m_{\tilde{\chi}_1^0}$ mass in the TwinoLSPRPV simplified model, as a function of the common $\tilde{\chi}_1^\pm/\tilde{\chi}_1^0$ branching fraction to a Z boson. See Figure 9.
- 14 AAD 21Y searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with four or more leptons (electrons, muons and tau-leptons). No significant excess above the Standard Model expectations is observed. Limits are set on Tchi1n12-GGM, and RPV models similar to Tchi1n2I, Tglu1A (with $q = u, d, s, c, b$, with equal branching fractions), and $\tilde{\ell}_L/\tilde{\nu} \rightarrow \ell/\nu \tilde{\chi}_1^0$ (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations), all with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$ via λ_{12k} or λ_{i33} (where $i, k \in 1, 2$), see their Figure 11.
- 15 SIRUNYAN 21M searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tsbob3, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14.
- 16 TUMASYAN 21C searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with with one lepton, a Higgs boson decaying to a pair of bottom quarks, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Lower limits are set on the masses of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the simplified model Tchi1n2E, see their Figure 6.
- 17 AAD 20AN searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons and missing transverse momentum. Events are further categorised in terms of lepton or jet multiplicity. No significant excess over the expected background is observed. Limits at 95% C.L. are set on the Higgsino mass in the T1n1n1A simplified model, see their Figure 11.
- 18 AAD 20I reported on ATLAS searches for electroweak production in models with compressed mass spectra as Tchi1n2Fa. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Events with \cancel{E}_T , two same-flavour, opposite-charge, low-transverse-momentum leptons, and jets from initial-state radiation or characteristic of vector-boson fusion production are selected. Constraints at

NODE=S046WNO;LINKAGE=ED

NODE=S046WNO;LINKAGE=FD

NODE=S046WNO;LINKAGE=DD

NODE=S046WNO;LINKAGE=ZC

NODE=S046WNO;LINKAGE=XC

NODE=S046WNO;LINKAGE=VC

NODE=S046WNO;LINKAGE=YC

NODE=S046WNO;LINKAGE=WC

NODE=S046WNO;LINKAGE=UC

NODE=S046WNO;LINKAGE=MC

- 95% C.L. are placed on the mass of the $\tilde{\chi}_1^\pm$ (degenerate with $\tilde{\chi}_2^0$) at 240 GeV for a mass splitting between $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^0$ of 7 GeV and extend down to a mass splitting of 1.5 GeV at the LEP chargino mass limit of 92.4 GeV. See their Fig. 14(b,c).
- 19 AAD 20K reported on a search for electroweak production in models with mass splittings near the electroweak scale as Tchi1n2F and exploiting three-lepton final state events with an emulated recursive jigsaw reconstruction method. The analysis uses a dataset of pp collisions at $\sqrt{s} = 13$ TeV corresponding to an integrated luminosity of 139 fb^{-1} . Exclusion limits at 95% C.L. are derived on next-to-lightest neutralinos and charginos with masses up to 345 GeV for a massless lightest neutralino, see their Fig. 7. NODE=S046WNO;LINKAGE=JC
- 20 AAD 200 reported on a search for electroweak production in models with charginos and sleptons decaying into final states with exactly two oppositely charged leptons and missing transverse momentum. A dataset of pp collisions at $\sqrt{s} = 13$ TeV corresponding to an integrated luminosity of 139 fb^{-1} was used. Exclusion limits at 95% C.L. are derived on $m_{\tilde{\chi}_1^\pm}$ decaying according to the Tchi1chi1H simplified model. Chargino masses up to 420 GeV are excluded for a massless lightest neutralino, see their Fig. 7(a). NODE=S046WNO;LINKAGE=PC
- 21 AAD 200 reported on a search for electroweak production in models with charginos and sleptons decaying into final states with exactly two oppositely charged leptons and missing transverse momentum. A dataset of pp collisions at $\sqrt{s} = 13$ TeV corresponding to an integrated luminosity of 139 fb^{-1} was used. Exclusion limits at 95% C.L. are derived on $m_{\tilde{\chi}_1^\pm}$ decaying according to the Tchi1chi1C simplified model. Chargino masses up to 1000 GeV are excluded for a massless lightest neutralino, see their Fig. 7(b). NODE=S046WNO;LINKAGE=QC
- 22 AAD 20R searched for electroweak production in the model Tchi1n2E, selecting events with a pair of b -tagged jets consistent with those from a Higgs boson decay, either an electron or a muon from the W boson decay and \cancel{E}_T . The analysis uses a dataset of pp collisions at $\sqrt{s} = 13$ TeV corresponding to an integrated luminosity of 139 fb^{-1} . Exclusion limits at 95% C.L. are derived on next-to-lightest neutralinos and charginos with masses up to 740 GeV for a massless lightest neutralino, assuming pure wino cross-sections. See their Fig. 6. NODE=S046WNO;LINKAGE=KC
- 23 SIRUNYAN 20AU searched in 77.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing one soft, hadronically decaying tau lepton, one energetic jet from initial-state radiation, and large \cancel{E}_T . No excess over the expected background is observed. Limits are derived on the wino mass in the Tchi1n2D simplified model, see their Figure 2. NODE=S046WNO;LINKAGE=LC
- 24 SIRUNYAN 20B searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one photon and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on chargino masses in a general gauge-mediated SUSY breaking (GGM) scenario Tchi1n12-GGM, see Figure 4. Limits are also set on the NLSP mass in the Tchi1chi1F and Tchi1chi1G simplified models, see their Figure 5. Finally, limits are set on the gluino mass in the Tglu4A simplified model, see Figure 6. NODE=S046WNO;LINKAGE=IC
- 25 AABOUD 19AU searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for direct electroweak production of charginos and next-to-lightest neutralinos decaying into lightest neutralinos and a W , and a Higgs boson, respectively. Fully hadronic, semileptonic, diphoton, and multilepton (electrons, muons) final states with missing transverse momentum are considered in this search. Observations are consistent with the Standard Model expectations, and 95% confidence-level limits of up to 680 GeV on the chargino/next-to-lightest neutralino masses are set (Tchi1n2E model). See their Figure 14 for an overlay of exclusion contours from all searches. NODE=S046WNO;LINKAGE=EC
- 26 SIRUNYAN 19BU searched for pair production of gauginos via vector boson fusion assuming the gaugino spectrum is compressed, in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV. The final states explored included zero leptons plus two jets, one lepton plus two jets, and one hadronic tau plus two jets. A similar bound is obtained in the light slepton limit. NODE=S046WNO;LINKAGE=BC
- 27 SIRUNYAN 19CI searched in 77.5 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with one or more high-momentum Higgs bosons, decaying to pairs of photons, jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t4 simplified model, see Figure 3, and on the wino mass in the Tchi1n2E simplified model, see their Figure 4. Limits are also set on the higgsino mass in the Tn1n1A and Tn1n1B simplified models, see their Figure 5. NODE=S046WNO;LINKAGE=GC
- 28 SIRUNYAN 19K searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with a photon, an electron or muon, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. In the framework of GMSB, limits are set on the chargino and neutralino mass in the Tchi1n1A simplified model, see their Figure 6. Limits are also set on the gluino mass in the Tglu4A simplified model, and on the squark mass in the Tsqk4A simplified model, see their Figure 7. NODE=S046WNO;LINKAGE=CC
- 29 AABOUD 18AY searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for direct electroweak production of charginos as in Tchi1chi1D models in events characterised by the presence of at least two hadronically decaying tau leptons and large missing transverse energy. No significant deviation from the expected SM background is observed. In the Tchi1chi1D model, assuming decays via intermediate $\tilde{\tau}_L$, the observed limits rule out $\tilde{\chi}_1^\pm$ masses up to 630 GeV for a massless $\tilde{\chi}_1^0$. See their Fig.7 (left). Interpretations are also provided in Fig 8 (top) for different assumptions on the ratio between $m_{\tilde{\tau}}$ and $m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0}$. NODE=S046WNO;LINKAGE=LB
- 30 AABOUD 18AY searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for direct electroweak production of charginos and neutralinos as in Tchi1n2D models, in events NODE=S046WNO;LINKAGE=MB

characterised by the presence of at least two hadronically decaying tau leptons and large missing transverse energy. No significant deviation from the expected SM background is observed. Assuming decays via intermediate $\tilde{\tau}_L$ and $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$, the observed

limits rule out $\tilde{\chi}_1^\pm$ masses up to 760 GeV for a massless $\tilde{\chi}_1^0$. See their Fig.7 (right). Interpretations are also provided in Fig 8 (bottom) for different assumptions on the ratio between $m_{\tilde{\tau}}$ and $m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0}$.

- 31 AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons in events with two or three leptons (electrons or muons), with or without jets and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino mass up to 750 GeV for massless neutralinos in the Tchi1chi1C simplified model exploiting $2\ell + 0$ jets signatures, see their Figure 8(a). NODE=S046WNO;LINKAGE=VB
- 32 AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino mass up to 1100 GeV for massless neutralinos in the Tchi1n2C simplified model exploiting 3ℓ signature, see their Figure 8(c). NODE=S046WNO;LINKAGE=WB
- 33 AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino mass up to 580 GeV for massless neutralinos in the Tchi1n2F simplified model exploiting $2\ell+2$ jets and 3ℓ signatures, see their Figure 8(d). NODE=S046WNO;LINKAGE=XB
- 34 AABOUD 18CK searched for events with at least 3 b -jets and large missing transverse energy in two datasets of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ of 36.1 fb^{-1} and 24.3 fb^{-1} depending on the trigger requirements. The analyses aimed to reconstruct two Higgs bosons decaying to pairs of b -quarks. No significant excess above the Standard Model expectations is observed. Limits are set on the Higgsino mass in the T1n1n1A simplified model, see their Figure 15(a). Constraints are also presented as a function of the BR of Higgsino decaying into an higgs boson and a gravitino, see their Figure 15(b). NODE=S046WNO;LINKAGE=ZB
- 35 AABOUD 18CO searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of mass-degenerate charginos and next-to-lightest neutralinos in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. The search channels are based on recursive jigsaw reconstruction. Limits are set on the chargino mass up to 600 GeV for massless neutralinos in the Tchi1n2F simplified model exploiting the statistical combination of $2\ell+2$ jets and 3ℓ channels. Chargino masses below 220 GeV are not excluded due to an excess of events above the SM prediction in the dedicated regions. See their Figure 13(d). NODE=S046WNO;LINKAGE=YB
- 36 AABOUD 18R searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for electroweak production in scenarios with compressed mass spectra in final states with two low-momentum leptons and missing transverse momentum. The data are found to be consistent with the SM prediction. Results are interpreted in Tchi1n2G wino models and $\tilde{\chi}_1^\pm$ masses are excluded up to 175 GeV for $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 10 \text{ GeV}$. The exclusion limits extend down to mass splittings of 2 GeV, see their Fig. 10 (bottom). NODE=S046WNO;LINKAGE=NB
- 37 AABOUD 18R searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for electroweak production in scenarios with compressed mass spectra in final states with two low-momentum leptons and missing transverse momentum. The data are found to be consistent with the SM prediction. Results are interpreted in Tchi1n2G higgsino models and $\tilde{\chi}_1^\pm$ masses are excluded up to 145 GeV for $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$. The exclusion limits extend down to mass splittings of 2.5 GeV, see their Fig. 10 (top). NODE=S046WNO;LINKAGE=OB
- 38 AABOUD 18U searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with at least one isolated photon, possibly jets and significant transverse momentum targeting generalised models of gauge-mediated SUSY breaking. No significant excess of events is observed above the SM prediction. Results of the diphoton channel are interpreted in terms of lower limits on the masses of gauginos Tchi1chi1A models, which reach as high as 1.3 TeV. Gaugino masses below 1060 GeV are excluded for any NLSP mass, see their Fig. 10. NODE=S046WNO;LINKAGE=PB
- 39 AABOUD 18Z searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more charged leptons (electrons, muons and up to two hadronically decaying taus). No significant deviation from the expected SM background is observed. Limits are set on the Higgsino mass in simplified models of general gauge mediated supersymmetry Tn1n1A/Tn1n1B/Tn1n1C, see their Figure 9. Limits are also set on the wino, slepton, sneutrino and gluino mass in a simplified model of NLSP pair production with R-parity violating decays of the LSP via λ_{12k} or λ_{j33} to charged leptons, see their Figures 7, 8. NODE=S046WNO;LINKAGE=QB
- 40 SIRUNYAN 18AA searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one photon and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on wino masses in a general gauge-mediated SUSY breaking (GGM) scenario with bino-like $\tilde{\chi}_1^0$ and wino-like $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$, see Figure 7. Limits are also set on the NLSP mass in the Tchi1n1A and Tchi1chi1A simplified models, see their Figure 8. Finally, limits are set on the gluino mass in the Tglu4A and NODE=S046WNO;LINKAGE=RA

- Tglu4B simplified models, see their Figure 9, and on the squark mass in the Tskq4A and Tskq4B simplified models, see their Figure 10.
- 41 SIRUNYAN 18AJ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two low-momentum, oppositely charged leptons (electrons or muons) and \cancel{E}_T . No excess over the expected background is observed. Limits are derived on the wino mass in the Tchi1n2F simplified model, see their Figure 5. Limits are also set on the stop mass in the Tstop10 simplified model, see their Figure 6. Finally, limits are set on the Higgsino mass in the Tchi1n2G simplified model, see Figure 8 and in the pMSSM, see Figure 7. NODE=S046WNO;LINKAGE=DB
- 42 SIRUNYAN 18AO searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and neutralinos in events with either two or more leptons (electrons or muons) of the same electric charge, or with three or more leptons, which can include up to two hadronically decaying tau leptons. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino/neutralino mass in the Tchi1n2A, Tchi1n2H, Tchi1n2D, Tchi1n2E and Tchi1n2F simplified models, see their Figures 14, 15, 16, 17 and 18. Limits are also set on the higgsino mass in the Tn1n1A, Tn1n1B and Tn1n1C simplified models, see their Figure 19. NODE=S046WNO;LINKAGE=GB
- 43 SIRUNYAN 18AP searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and neutralinos by combining a number of previous and new searches. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino/neutralino mass in the Tchi1n2E, Tchi1n2F and Tchi1n2I simplified models, see their Figures 7, 8, 9 and 10. Limits are also set on the higgsino mass in the Tn1n1A, Tn1n1B and Tn1n1C simplified models, see their Figure 11, 12, 13 and 14. NODE=S046WNO;LINKAGE=JB
- 44 SIRUNYAN 18AR searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two opposite-charge, same-flavour leptons (electrons or muons), jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4C simplified model, see their Figure 7. Limits are also set on the chargino/neutralino mass in the Tchi1n2F simplified models, see their Figure 8, and on the higgsino mass in the Tn1n1B and Tn1n1C simplified models, see their Figure 9. Finally, limits are set on the sbottom mass in the Tsb0t3 simplified model, see their Figure 10. NODE=S046WNO;LINKAGE=KB
- 45 SIRUNYAN 18DN searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and for pair production of top squarks in events with two leptons (electrons or muons) of the opposite electric charge. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino mass in the Tchi1chi1C and Tchi1chi1E simplified models, see their Figure 8. Limits are also set on the stop mass in the Tstop1 and Tstop2 simplified models, see their Figure 9. NODE=S046WNO;LINKAGE=RB
- 46 SIRUNYAN 18DP searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and neutralinos or of chargino pairs in events with a tau lepton pair and significant missing transverse momentum. Both hadronic and leptonic decay modes are considered for the tau lepton. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino mass in the Tchi1chi1D and Tchi1n2 simplified models, see their Figures 14 and 15. Also, excluded stau pair production cross sections are shown in Figures 11, 12, and 13. NODE=S046WNO;LINKAGE=TB
- 47 SIRUNYAN 18X searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more high-momentum Higgs bosons, decaying to pairs of photons, jets and \cancel{E}_T . The razor variables (M_R and R^2) are used to categorise the events. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t4 simplified model and on the wino mass in the Tchi1n2E simplified model, see their Figure 5. Limits are also set on the higgsino mass in the Tn1n1A and Tn1n1B simplified models, see their Figure 6. NODE=S046WNO;LINKAGE=XA
- 48 KHACHATRYAN 17L searched in about 19 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with two τ (at least one decaying hadronically) and \cancel{E}_T . In the Tchi1chi1C model, assuming decays via intermediate $\tilde{\tau}$ or $\tilde{\nu}_\tau$ with equivalent mass, the observed limits rule out $\tilde{\chi}_1^\pm$ masses up to 420 GeV for a massless $\tilde{\chi}_1^0$. See their Fig.5. NODE=S046WNO;LINKAGE=PA
- 49 SIRUNYAN 17AW searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a charged lepton (electron or muon), two jets identified as originating from a b -quark, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the chargino and the next-to-lightest neutralino in the Tchi1n2E simplified model, see their Figure 6. NODE=S046WNO;LINKAGE=QA
- 50 AAD 16AA summarized and extended ATLAS searches for electroweak supersymmetry in final states containing several charged leptons, \cancel{E}_T , with or without hadronic jets, in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$. The paper reports the results of new interpretations and statistical combinations of previously published analyses, as well as new analyses. Exclusion limits at 95% C.L. are set on the $\tilde{\chi}_1^\pm$ mass in the Tchi1chi1B and Tchi1chi1C simplified models. See their Fig. 13. NODE=S046WNO;LINKAGE=VA
- 51 AAD 16AA summarized and extended ATLAS searches for electroweak supersymmetry in final states containing several charged leptons, \cancel{E}_T , with or without hadronic jets, in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$. The paper reports the results of new interpretations and statistical combinations of previously published analyses, as well as new analyses. Exclusion limits at 95% C.L. are set on mass-degenerate $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ masses in the Tchi1n2B, Tchi1n2C, and Tchi1n2D simplified models. See their Figs. 16, 17, and 18. Interpretations in phenomenological-MSSM, two-parameter Non Universal Higgs Masses (NUHM2), and gauge-mediated symmetry breaking (GMSB) models are also given in their Figs. 20, 21 and 22. NODE=S046WNO;LINKAGE=BB

- 52 KHACHATRYAN 16R searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with one or more photons, one electron or muon, and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on wino masses in a general gauge-mediated SUSY breaking model (GGM), for a wino-like neutralino NLSP scenario, see Fig. 5. Limits are also set in the Tglu1D and Tchi1n1A simplified models, see Fig. 6. The Tchi1n1A limit is reduced to 340 GeV for a branching ratio reduced by the weak mixing angle. NODE=S046WNO;LINKAGE=SA
- 53 AAD 15BA searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of charginos and neutralinos decaying to a final state containing a W boson and a 125 GeV Higgs boson, plus missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of direct chargino and next-to-lightest neutralino production, with the decays $\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ and $\tilde{\chi}_2^0 \rightarrow H \tilde{\chi}_1^0$ having 100% branching fraction, see Fig. 8. A combination of the multiple final states for the Higgs decay yields the best limits (Fig. 8d). NODE=S046WNO;LINKAGE=KA
- 54 AAD 15CA searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with one or more photons and \cancel{E}_T , with or without leptons (e, μ). No significant excess above the Standard Model expectations is observed. Limits are set on wino masses in the general gauge-mediated SUSY breaking model (GGM), for wino-like NLSP, see Fig. 9, 12. NODE=S046WNO;LINKAGE=OA
- 55 AAD 14H searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of charginos and neutralinos decaying to a final state with three leptons and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of direct chargino and next-to-lightest neutralino production, with decays to the lightest neutralino via either all three generations of leptons, taus only, gauge bosons, or Higgs bosons, see Fig. 7. An interpretation in the pMSSM is also given, see Fig. 8. NODE=S046WNO;LINKAGE=HA
- 56 AAD 14X searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least four leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the wino-like chargino mass in an R-parity violating simplified model where the decay $\tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_1^0$, with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, takes place with a branching ratio of 100%, see Fig. 8. NODE=S046WNO;LINKAGE=X
- 57 KHACHATRYAN 14L searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of chargino-neutralino $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ pair production with Higgs or W -bosons in the decay chain, leading to HW final states with missing transverse energy. The decays of a Higgs boson to a photon pair are considered in conjunction with hadronic and leptonic decay modes of the W bosons. No significant excesses over the expected SM backgrounds are observed. The results are interpreted in the context of simplified models where the decays $\tilde{\chi}_2^0 \rightarrow H \tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ take place 100% of the time, see Figs. 22–23. NODE=S046WNO;LINKAGE=Y
- 58 AAD 13 searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for charginos and neutralinos decaying to a final state with three leptons (e and μ) and missing transverse energy. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in the phenomenological MSSM, see Fig. 2 and 3, and in simplified models, see Fig. 4. For the simplified models with intermediate slepton decays, degenerate $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$ masses up to 500 GeV are excluded at 95% C.L. for very large mass differences with the $\tilde{\chi}_1^0$. Supersedes AAD 12AS. NODE=S046WNO;LINKAGE=GA
- 59 AAD 13B searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for gauginos decaying to a final state with two leptons (e and μ) and missing transverse energy. No excess beyond the Standard Model expectation is observed. Limits are derived in a simplified model of wino-like chargino pair production, where the chargino always decays to the lightest neutralino via an intermediate on-shell charged slepton, see Fig. 2(b). Chargino masses between 110 and 340 GeV are excluded at 95% C.L. for $m_{\tilde{\chi}_1^0} = 10 \text{ GeV}$. Exclusion limits are also derived in the phenomenological MSSM, see Fig. 3. NODE=S046WNO;LINKAGE=CD
- 60 AAD 12CT searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing four or more leptons (electrons or muons) and either moderate values of missing transverse momentum or large effective mass. No significant excess is found in the data. Limits are presented in a simplified model of R-parity violating supersymmetry in which charginos are pair-produced and then decay into a W -boson and a $\tilde{\chi}_1^0$, which in turn decays through an RPV coupling into two charged leptons ($e^\pm e^\mp$ or $e^\pm \mu^\mp$) and a neutrino. In this model, chargino masses up to 540 GeV are excluded at 95% C.L. for $m_{\tilde{\chi}_1^0}$ above 300 GeV, see Fig. 3a. The limit deteriorates for lighter $\tilde{\chi}_1^0$. Limits are also set in an R-parity violating mSUGRA model, see Fig. 3b. NODE=S046WNO;LINKAGE=DA
- 61 CHATRCHYAN 12BJ searched in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for direct electroweak production of charginos and neutralinos in events with at least two leptons, jets and missing transverse momentum. No significant excesses over the expected SM backgrounds are observed and 95% C.L. limits on the production cross section of $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ pair production were set in a number of simplified models, see Figs. 7 to 12. NODE=S046WNO;LINKAGE=CH
- 62 ABDALLAH 03M uses data from $\sqrt{s} = 192\text{--}208 \text{ GeV}$ to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass of charginos is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays), for charginos and for sleptons. These limits are valid for values of $M_2 < 1 \text{ TeV}$, $|\mu| \leq 2 \text{ TeV}$ with the $\tilde{\chi}_1^0$ as LSP. Constraints from the Higgs search in the m_h^{max} scenario assuming $m_t =$ NODE=S046WNO;LINKAGE=AH

- 174.3 GeV are included. The quoted limit applies if there is no mixing in the third family or when $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} > 6$ GeV. If mixing is included the limit degrades to 90 GeV. See Fig. 43 for the mass limits as a function of $\tan\beta$. These limits update the results of ABREU 00W.
- 63 AAD 20AN searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with two photons and missing transverse momentum. Events are further categorised in terms of lepton or jet multiplicity. No significant excess over the expected background is observed. Limits at 95% C.L. are derived in Tchi1n2E simplified models. Next-to-lightest neutralinos and charginos with masses up to 310 GeV for a massless lightest neutralino are excluded. See their Fig. 10. NODE=S046WNO;LINKAGE=TC
- 64 KHACHATRYAN 16AA searched in 7.4 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with one or more photons, hadronic jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on wino masses in the general gauge-mediated SUSY breaking model (GGM), for a wino-like neutralino NLSP scenario and with the wino mass fixed at 10 GeV above the bino mass, see Fig. 4. Limits are also set in the Tchi1chi1A and Tchi1n1A simplified models, see Fig. 3. NODE=S046WNO;LINKAGE=NA
- 65 KHACHATRYAN 16R searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with one or more photons, one electron or muon, and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are also set in the Tglu1F simplified model, see Fig. 6. NODE=S046WNO;LINKAGE=TA
- 66 KHACHATRYAN 16Y searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with one or two soft isolated leptons, hadronic jets, and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the $\tilde{\chi}_1^\pm$ mass (which is degenerate with the $\tilde{\chi}_2^0$) in the Tchi1n2A simplified model, see Fig. 4. NODE=S046WNO;LINKAGE=UA
- 67 AAD 14AV searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for the direct production of charginos, neutralinos and staus in events containing at least two hadronically decaying τ -leptons, large missing transverse momentum and low jet activity. The quoted limit was derived for direct $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp$ production with $\tilde{\chi}_2^0 \rightarrow \tilde{\tau}\tau \rightarrow \tau\tau\tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm \rightarrow \tilde{\tau}\nu(\tilde{\nu}_\tau\tau) \rightarrow \tau\nu\tilde{\chi}_1^0$, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm}$, $m_{\tilde{\tau}} = 0.5(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV. No excess over the expected SM background is observed. Exclusion limits are set in simplified models of $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp$ and $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ pair production, see their Figure 7. Upper limits on the cross section and signal strength for direct di-stau production are derived, see Figures 8 and 9. Also, limits are derived in a pMSSM model where the only light slepton is the $\tilde{\tau}_R$, see Figure 10. NODE=S046WNO;LINKAGE=A1
- 68 AAD 14AV searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for the direct production of charginos, neutralinos and staus in events containing at least two hadronically decaying τ -leptons, large missing transverse momentum and low jet activity. The quoted limit was derived for direct $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp$ production with $\tilde{\chi}_1^\pm \rightarrow \tilde{\tau}\nu(\tilde{\nu}_\tau\tau) \rightarrow \tau\nu\tilde{\chi}_1^0$, $m_{\tilde{\tau}} = 0.5(m_{\tilde{\chi}_1^\pm} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV. No excess over the expected SM background is observed. Exclusion limits are set in simplified models of $\tilde{\chi}_1^\pm \tilde{\chi}_1^\mp$ and $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$ pair production, see their Figure 7. Upper limits on the cross section and signal strength for direct di-stau production are derived, see Figures 8 and 9. Also, limits are derived in a pMSSM model where the only light slepton is the $\tilde{\tau}_R$, see Figure 10. NODE=S046WNO;LINKAGE=A2
- 69 AAD 14G searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for electroweak production of chargino pairs, or chargino-neutralino pairs, decaying to a final state with two leptons (e and μ) and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of chargino pair production, with chargino decays to the lightest neutralino via either sleptons or gauge bosons, see Fig 5.; or in simplified models of chargino and next-to-lightest neutralino production, with decays to the lightest neutralino via gauge bosons, see Fig. 7. An interpretation in the pMSSM is also given, see Fig. 10. NODE=S046WNO;LINKAGE=IA
- 70 AALTONEN 14 searched in 5.8 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV for evidence of chargino and next-to-lightest neutralino associated production in final states consisting of three leptons (electrons, muons or taus) and large missing transverse momentum. The results are consistent with the Standard Model predictions within 1.85σ . Limits on the chargino mass are derived in an mSUGRA model with $m_0 = 60$ GeV, $\tan\beta = 3$, $A_0 = 0$ and $\mu > 0$, see their Fig. 2. NODE=S046WNO;LINKAGE=W
- 71 KHACHATRYAN 14I searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for electroweak production of chargino pairs decaying to a final state with opposite-sign lepton pairs (e or μ) and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models, see Fig. 18. NODE=S046WNO;LINKAGE=FA
- 72 AALTONEN 13Q searched in 6.0 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV for evidence of chargino-neutralino associated production in like-sign dilepton final states. One lepton is identified as the hadronic decay of a tau lepton, while the other is an electron or muon. Good agreement with the Standard Model predictions is observed and limits are set on the chargino-neutralino cross section for simplified gravity- and gauge-mediated models, see their Figs. 2 and 3. NODE=S046WNO;LINKAGE=TO
- 73 AAD 12AS searched in 2.06 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for charginos and neutralinos decaying to a final state with three leptons (e and μ) and missing transverse energy. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in the phenomenological MSSM, see Fig. 2 (top), and in simplified models, see Fig. 2 (bottom). NODE=S046WNO;LINKAGE=GL

- ⁷⁴ AAD 12T looked in 1 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for the production of supersymmetric particles decaying into final states with missing transverse momentum and exactly two isolated leptons (e or μ). Opposite-sign and same-sign dilepton events were separately studied. Additionally, in opposite-sign events, a search was made for an excess of same-flavor over different-flavor lepton pairs. No excess over the expected background is observed and limits are placed on the effective production cross section of opposite-sign dilepton events with $\cancel{E}_T > 250 \text{ GeV}$ and on same-sign dilepton events with $\cancel{E}_T > 100 \text{ GeV}$. The latter limit is interpreted in a simplified electroweak gaugino production model as a lower chargino mass limit.
- ⁷⁵ CHATRCHYAN 11B looked in 35 pb^{-1} of pp collisions at $\sqrt{s}=7 \text{ TeV}$ for events with an isolated lepton (e or μ), a photon and \cancel{E}_T which may arise in a generalized gauge mediated model from the decay of Wino-like NLSPs. No evidence for an excess over the expected background is observed. Limits are derived in the plane of squark/gluino mass versus Wino mass (see Fig. 4). Mass degeneracy of the produced squarks and gluinos is assumed.
- ⁷⁶ CHATRCHYAN 11V looked in 35 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with ≥ 3 isolated leptons (e , μ or τ), with or without jets and \cancel{E}_T . No evidence for an excess over the expected background is observed. Limits are derived in the CMSSM $(m_0, m_{1/2})$ plane for $\tan\beta = 3$ (see Fig. 5).

NODE=S046WNO;LINKAGE=AG

NODE=S046WNO;LINKAGE=C1

NODE=S046WNO;LINKAGE=C2

Long-lived $\tilde{\chi}^\pm$ (Chargino) mass limit

Limits on charginos which leave the detector before decaying.					NODE=S046SWN
VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	NODE=S046SWN NODE=S046SWN
>1050	95	1 AAD	23G ATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, wino LSP, $\tau=20 \text{ ns}$	OCCUR=3
>1050	95	1 AAD	23G ATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, wino LSP, stable	OCCUR=4
> 660	95	2 AAD	22U ATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, wino LSP, AMSB, $\tan\beta = 5, \mu > 0, \tau = 0.2 \text{ ns}$	
> 860	95	2 AAD	22U ATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, wino LSP, AMSB, $\tan\beta = 5, \mu > 0, \tau = 1.5 \text{ ns}$	OCCUR=2
> 220	95	2 AAD	22U ATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, higgsino LSP, $\tau=0.04 \text{ ns}$	OCCUR=3
> 710	95	2 AAD	22U ATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, higgsino LSP, $\tau=1 \text{ ns}$	OCCUR=4
> 884	95	3 SIRUNYAN	20N CMS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, wino LSP, AMSB, $\tan\beta = 5, \mu > 0, \tau = 3 \text{ ns}$	
> 474	95	3 SIRUNYAN	20N CMS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, wino LSP, AMSB, $\tan\beta = 5, \mu > 0, \tau = 0.2 \text{ ns}$	OCCUR=2
> 750	95	3 SIRUNYAN	20N CMS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, higgsino LSP, AMSB, $\tan\beta=5, \mu > 0, \tau=3 \text{ ns}$	OCCUR=3
> 175	95	3 SIRUNYAN	20N CMS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, higgsino LSP, AMSB, $\tan\beta=5, \mu > 0, \tau=0.05 \text{ ns}$	OCCUR=4
>1090	95	4 AABOUD	19AT ATLS	long-lived $\tilde{\chi}_1^\pm$ mAMSB	
> 460	95	5 AABOUD	18AS ATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, lifetime 0.2 ns, $m_{\tilde{\chi}^\pm} - m_{\tilde{\chi}_1^0} = 160 \text{ MeV}$	
> 715	95	6 SIRUNYAN	18BR CMS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, AMSB, $\tan\beta = 5$ and $\mu > 0, \tau = 3 \text{ ns}$	
> 695	95	6 SIRUNYAN	18BR CMS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, AMSB, $\tan\beta = 5$ and $\mu > 0, \tau = 7 \text{ ns}$	OCCUR=2
> 505	95	6 SIRUNYAN	18BR CMS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, AMSB, $\tan\beta = 5, \mu > 0, 0.5 \text{ ns} > \tau > 60 \text{ ns}$	OCCUR=3
> 620	95	7 AAD	15AE ATLS	stable $\tilde{\chi}^\pm$	
> 534	95	8 AAD	15BMATLS	stable $\tilde{\chi}^\pm$	
> 239	95	8 AAD	15BMATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, lifetime 1 ns, $m_{\tilde{\chi}^\pm} - m_{\tilde{\chi}_1^0} = 0.14 \text{ GeV}$	OCCUR=2
> 482	95	8 AAD	15BMATLS	$\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, lifetime 15 ns, $m_{\tilde{\chi}^\pm} - m_{\tilde{\chi}_1^0} = 0.14 \text{ GeV}$	OCCUR=3
> 103	95	9 AAD	13H ATLS	long-lived $\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, mAMSB, $\Delta m_{\tilde{\chi}_1^0} = 160 \text{ MeV}$	
> 92	95	10 AAD	12BJ ATLS	long-lived $\tilde{\chi}^\pm \rightarrow \pi^\pm \tilde{\chi}_1^0$, mAMSB	
> 171	95	11 ABAZOV	09M D0	\tilde{H}	
> 102	95	12 ABBIENDI	03L OPAL	$m_{\tilde{\nu}} > 500 \text{ GeV}$	
none 2-93.0	95	13 ABREU	00T DLPH	\tilde{H}^\pm or $m_{\tilde{\nu}} > m_{\tilde{\chi}^\pm}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 260	95	14	KHACHATRY...15AB CMS	$\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm, \tau_{\tilde{\chi}_1^\pm} = 0.2\text{ns}$, AMSB	
> 800	95	15	KHACHATRY...15AO CMS	long-lived $\tilde{\chi}_1^\pm$, mAMSB, $\tau > 100\text{ns}$	
> 100	95	15	KHACHATRY...15AO CMS	long-lived $\tilde{\chi}_1^\pm$, mAMSB, $\tau > 3\text{ ns}$	OCCUR=2
		16	KHACHATRY...15W CMS	long-lived $\tilde{\chi}^0, \tilde{q} \rightarrow q\tilde{\chi}^0, \tilde{\chi}^0 \rightarrow \ell^+ \ell^- \nu$, RPV	
> 270	95	17	AAD 13BD ATLS	disappearing-track signature, AMSB	
> 278	95	18	ABAZOV 13B D0	long-lived $\tilde{\chi}^\pm$, gaugino-like	
> 244	95	18	ABAZOV 13B D0	long-lived $\tilde{\chi}^\pm$, higgsino-like	OCCUR=2
1	AAD 23G searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$ for chargino/neutralino pair production (wino-like LSP) in events with high-pt tracks with large ionisation in the pixel detector. No significant excess above the Standard Model predictions is observed. Limits are set on the chargino mass as a function of its lifetime, see Figure 19.				NODE=S046SWN;LINKAGE=M
2	AAD 22U searched for the signature of disappearing track from a long-lived chargino in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$. Long-lived charginos decay into quasi-degenerate neutralino emitting a low-momentum particle whose identification is not attempted. The signal is identified by requiring short tracklets in the four pixel layers with no continuation in the SCT (strip) detector. The main background from fake tracklets is estimated directly with the data. No significant excess above the background prediction is found. The results are interpreted in an AMSB scenario (wino LSP), on $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\pm$ and $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}_1^0$, assuming $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm) = 100\%$, see their figure 7. Results are also interpreted in a higgsino-LSP model, with $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\mp$, and $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}_{1,2}^0$, assuming $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm) = 95.5\%$, $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 e^\pm) = 3\%$, $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \mu^\pm) = 1.5\%$, see their figure 8. Finally, results are interpreted in a simplified model of gluino pair production, with $pp \rightarrow \tilde{g}\tilde{g}$ and $B(\tilde{g} \rightarrow qq\tilde{\chi}_1^0) = B(\tilde{g} \rightarrow qq\tilde{\chi}^+) = B(\tilde{g} \rightarrow qq\tilde{\chi}^-) = 1/3$, see their figure 9.				NODE=S046SWN;LINKAGE=L
3	SIRUNYAN 20N searched in 101 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$ for direct electroweak production of long-lived charginos in events containing isolated tracks with missing hits in the outer layer of the silicon tracker and little or no associated calorimetric energy deposits (disappearing tracks). No significant excess above the Standard Model expectations is observed. In an AMSB context and assuming a wino LSP, limits are set on the cross section of direct chargino production through $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\mp$ and $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}_1^0$, assuming $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm) = 100\%$, as a function of the chargino mass and mean proper lifetime, see Figure 2. In the case of a Higgsino LSP, limits are set on the cross section of direct chargino production through $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\mp$ and $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}_{1,2}^0$, assuming $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm) = 95.5\%$, $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 e^\pm) = 3\%$, $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \mu^\pm) = 1.5\%$, as a function of the chargino mass and mean proper lifetime, see Figure 3.				NODE=S046SWN;LINKAGE=K
4	AABOUD 19AT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$ for metastable R -hadrons. Multiple search strategies for a wide range of lifetimes, corresponding to path lengths of a few meters, are defined. No significant deviations from the expected Standard Model background are observed. Results are interpreted in terms of direct electroweak production of long-lived charginos in the context of mAMSB scenarios. Chargino masses are excluded at 95% C.L. below 1090 GeV. See their Figure 10 (right).				NODE=S046SWN;LINKAGE=J
5	AABOUD 18AS searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$ for direct electroweak production of long-lived charginos in the context of AMSB or phenomenological MSSM scenarios with wino-like LSP. Events with a disappearing track due to a low-momentum pion accompanied by at least one jet with high transverse momentum from initial-state radiation are considered. No significant excess above the Standard Model expectations is observed. Exclusion limits are set at 95% confidence level on the mass of charginos for different chargino lifetimes. For a pure wino with a lifetime of about 0.2 ns, corresponding to a mass-splitting between the charged and neutral wino of around 160 MeV, chargino masses up to 460 GeV are excluded, see their Fig. 8.				NODE=S046SWN;LINKAGE=G
6	SIRUNYAN 18BR searched in 38.4 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$ for direct electroweak production of long-lived charginos in events containing isolated tracks with missing hits in the outer layer of the silicon tracker and little or no associated calorimetric energy deposits (disappearing tracks). No significant excess above the Standard Model expectations is observed. In an AMSB context, limits are set on the cross section of direct chargino production through $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\mp$ and $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}_1^0$, assuming $BR(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm) = 100\%$, as a function of the chargino mass and mean proper lifetime, see Figures 3, 4 and 5.				NODE=S046SWN;LINKAGE=I
7	AAD 15AE searched in 19.1 fb^{-1} of pp collisions at $\sqrt{s} = 8\text{ TeV}$ for heavy long-lived charged particles, measured through their specific ionization energy loss in the ATLAS pixel detector or their time-of-flight in the ALTAS muon system. In the absence of an excess of events above the expected backgrounds, limits are set on stable charginos, see Fig. 10.				NODE=S046SWN;LINKAGE=A
8	AAD 15BM searched in 18.4 fb^{-1} of pp collisions at $\sqrt{s} = 8\text{ TeV}$ for stable and metastable non-relativistic charged particles through their anomalous specific ionization energy loss in the ATLAS pixel detector. In absence of an excess of events above the				NODE=S046SWN;LINKAGE=B

expected backgrounds, limits are set on stable charginos (see Table 5) and on metastable charginos decaying to $\tilde{\chi}_1^0 \pi^\pm$, see Fig. 11.

- 9 AAD 13H searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for direct electroweak production of long-lived charginos in the context of AMSB scenarios. The search is based on the signature of a high-momentum isolated track with few associated hits in the outer part of the tracking system, arising from a chargino decay into a neutralino and a low-momentum pion. The p_T spectrum of the tracks was found to be consistent with the SM expectations. Constraints on the lifetime and the production cross section were obtained, see Fig. 6. In the minimal AMSB framework with $\tan\beta = 5$, and $\mu > 0$, a chargino having a mass below 103 (85) GeV for a chargino-neutralino mass splitting $\Delta m_{\tilde{\chi}_1^0}$ of 160 (170) MeV is excluded at the 95% C.L. See Fig. 7 for more precise bounds. NODE=S046SWN;LINKAGE=GA
- 10 AAD 12BJ looked in 1.02 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for signatures of decaying charginos resulting in isolated tracks with few associated hits in the outer region of the tracking system. The p_T spectrum of the tracks was found to be consistent with the SM expectations. Constraints on the lifetime and the production cross section were obtained. In the minimal AMSB framework with $m_{3/2} < 32 \text{ TeV}$, $m_0 < 1.5 \text{ TeV}$, $\tan\beta = 5$, and $\mu > 0$, a chargino having a mass below 92 GeV and a lifetime between 0.5 ns and 2 ns is excluded at the 95% C.L. See their Fig. 8 for more precise bounds. NODE=S046SWN;LINKAGE=DA
- 11 ABAZOV 09M searched in 1.1 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for events with direct production of a pair of charged massive stable particles identified by their TOF. The number of the observed events is consistent with the predicted background. The data are used to constrain the production cross section as a function of the $\tilde{\chi}_1^\pm$ mass, see their Fig. 2. The quoted limit improves to 206 GeV for gaugino-like charginos. NODE=S046SWN;LINKAGE=AZ
- 12 ABBIENDI 03L used e^+e^- data at $\sqrt{s} = 130\text{--}209 \text{ GeV}$ to select events with two high momentum tracks with anomalous dE/dx . The excluded cross section is compared to the theoretical expectation as a function of the heavy particle mass in their Fig. 3. The bounds are valid for colorless fermions with lifetime longer than 10^{-6} s . Supersedes the results from ACKERSTAFF 98P. NODE=S046SWN;LINKAGE=AB
- 13 ABREU 00T searches for the production of heavy stable charged particles, identified by their ionization or Cherenkov radiation, using data from $\sqrt{s} = 130$ to 189 GeV. These limits include and update the results of ABREU 98P. NODE=S046SWN;LINKAGE=WT
- 14 KHACHATRYAN 15AB searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing tracks with little or no associated calorimeter energy deposits and with missing hits in the outer layers of the tracking system (disappearing-track signature). Such disappearing tracks can result from the decay of charginos that are nearly mass degenerate with the lightest neutralino. The number of observed events is in agreement with the background expectation. Limits are set on the cross section of electroweak chargino production in terms of the chargino mass and mean proper lifetime, see Fig. 4. In the minimal AMSB model, a chargino mass below 260 GeV is excluded at 95% C.L., see their Fig. 5. NODE=S046SWN;LINKAGE=E
- 15 KHACHATRYAN 15O searched in 18.8 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of long-lived charginos in the context of AMSB and pMSSM scenarios. The results are based on a previously published search for heavy stable charged particles at 7 and 8 TeV. In the minimal AMSB framework with $\tan\beta = 5$ and $\mu \geq 0$, constraints on the chargino mass and lifetime were placed, see Fig. 5. Charginos with a mass below 800 (100) GeV are excluded at the 95% C.L. for lifetimes above 100 ns (3 ns). Constraints are also placed on the pMSSM parameter space, see Fig. 3. NODE=S046SWN;LINKAGE=D
- 16 KHACHATRYAN 15W searched in up to 20.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of long-lived neutralinos produced through \tilde{q} -pair production, with $\tilde{q} \rightarrow q\tilde{\chi}^0$ and $\tilde{\chi}^0 \rightarrow \ell^+\ell^-\nu$ (RPV: $\lambda_{121}, \lambda_{122} \neq 0$). 95% C.L. exclusion limits on cross section times branching ratio are set as a function of mean proper decay length of the neutralino, see Figs. 6 and 9. NODE=S046SWN;LINKAGE=F
- 17 AAD 13BD searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing tracks with no associated hits in the outer region of the tracking system resulting from the decay of charginos that are nearly mass degenerate with the lightest neutralino, as is often the case in AMSB scenarios. No significant excess above the background expectation is observed for candidate tracks with large transverse momentum. Constraints on chargino properties are obtained and in the minimal AMSB model, a chargino mass below 270 GeV is excluded at 95% C.L., see their Fig. 7. NODE=S046SWN;LINKAGE=AA
- 18 ABAZOV 13B looked in 6.3 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for charged massive long-lived particles in events with muon-like particles that have both speed and ionization energy loss inconsistent with muons produced in beam collisions. In the absence of an excess, limits are set at 95% C.L. on gaugino- and higgsino-like charginos, see their Table 20 and Fig. 23. NODE=S046SWN;LINKAGE=AV

$\tilde{\nu}$ (Sneutrino) mass limit

The limits may depend on the number, $N(\tilde{\nu})$, of sneutrinos assumed to be degenerate in mass. Only $\tilde{\nu}_L$ (not $\tilde{\nu}_R$) is assumed to exist. It is possible that $\tilde{\nu}$ could be the lightest supersymmetric particle (LSP).

We report here, but do not include in the Listings, the limits obtained from the fit of the final results obtained by the LEP Collaborations on the invisible width of the Z boson ($\Delta\Gamma_{\text{inv.}} < 2.0 \text{ MeV}$, LEP-SLC 06): $m_{\tilde{\nu}} > 43.7 \text{ GeV}$ ($N(\tilde{\nu})=1$) and $m_{\tilde{\nu}} > 44.7 \text{ GeV}$ ($N(\tilde{\nu})=3$).

NODE=S046SNU
NODE=S046SNU

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	NODE=S046SNU;CHECK LIMITS
>3900	95	1 AAD	23CB ATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $\lambda_{312} = \lambda_{321} = 0.07$, $\lambda'_{311} = 0.11$	
>2800	95	1 AAD	23CB ATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\tau$, $\lambda_{313} = 0.07$, $\lambda'_{311} = 0.11$	OCCUR=2
>2700	95	1 AAD	23CB ATLS	RPV, $\tilde{\nu}_\tau \rightarrow \mu\tau$, $\lambda_{323} = 0.07$, $\lambda'_{311} = 0.11$	OCCUR=3
>4200	95	2 TUMASYAN	23H CMS	$1e + 1\mu$, RPV $\nu_\tau \rightarrow e\mu$, $\lambda = \lambda' = 0.1$	
>3700	95	2 TUMASYAN	23H CMS	$1e + 1\tau$, RPV $\nu_\tau \rightarrow e\tau$, $\lambda = \lambda' = 0.1$	OCCUR=2
>3600	95	2 TUMASYAN	23H CMS	$1\mu + 1\tau$, RPV $\nu_\tau \rightarrow \mu\tau$, $\lambda = \lambda' = 0.1$	OCCUR=3
>2200	95	2 TUMASYAN	23H CMS	$1e + 1\mu$, RPV $\nu_\tau \rightarrow e\mu$, $\lambda = \lambda' = 0.01$	OCCUR=4
>1600	95	2 TUMASYAN	23H CMS	$1e + 1\tau$, RPV $\nu_\tau \rightarrow e\tau$, $\lambda = \lambda' = 0.01$	OCCUR=5
>1600	95	2 TUMASYAN	23H CMS	$1\mu + 1\tau$, RPV $\nu_\tau \rightarrow \mu\tau$, $\lambda = \lambda' = 0.01$	OCCUR=6
>3400	95	3 AABOUD	18CMATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $\lambda_{312} = \lambda_{321} = 0.07$, $\lambda'_{311} = 0.11$	
>2900	95	4 AABOUD	18CMATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\tau$, $\lambda_{313} = \lambda_{331} = 0.07$, $\lambda'_{311} = 0.11$	OCCUR=2
>2600	95	5 AABOUD	18CMATLS	RPV, $\tilde{\nu}_\tau \rightarrow \mu\tau$, $\lambda_{323} = \lambda_{332} = 0.07$, $\lambda'_{311} = 0.11$	OCCUR=3
>1060	95	6 AABOUD	18Z ATLS	RPV, $\geq 4\ell$, $\lambda_{12k} \neq 0$, $m_{\tilde{\chi}_1^0} = 600$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)	
> 780	95	6 AABOUD	18Z ATLS	RPV, $\geq 4\ell$, $\lambda_{i33} \neq 0$, $m_{\tilde{\chi}_1^0} = 300$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)	OCCUR=2
>1700	95	7 SIRUNYAN	18AT CMS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $\lambda_{132} = \lambda_{231} = \lambda'_{311} = 0.01$	
>3800	95	7 SIRUNYAN	18AT CMS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $\lambda_{132} = \lambda_{231} = \lambda'_{311} = 0.1$	OCCUR=2
>2300	95	8 AABOUD	16P ATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $\lambda'_{311} = 0.11$	
>2200	95	8 AABOUD	16P ATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\tau$, $\lambda'_{311} = 0.11$	OCCUR=2
>1900	95	8 AABOUD	16P ATLS	RPV, $\tilde{\nu}_\tau \rightarrow \mu\tau$, $\lambda'_{311} = 0.11$	OCCUR=3
> 400	95	9 AAD	14X ATLS	RPV, $\geq 4\ell^\pm$, $\tilde{\nu} \rightarrow \nu\tilde{\chi}_1^0$, $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$	
> 94	95	10 AAD 11 ABDALLAH	11Z ATLS 03M DLPH	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$ $1 \leq \tan\beta \leq 40$, $m_{\tilde{e}_R} - m_{\tilde{\chi}_1^0} > 10$ GeV	
> 84	95	12 HEISTER	02N ALEP	$\tilde{\nu}_e$, any Δm	
> 41	95	13 DECAMP	92 ALEP	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=3$, model independent	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
		14 SIRUNYAN	19AO	RPV, $\mu^\pm \mu^\pm + \geq 2\text{jets}$, $\lambda'_{211} \neq 0$, $\tilde{\nu}_\mu \rightarrow \mu\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow \mu q \bar{q} q \bar{q}$	
>1280	95	15 KHACHATRY...16BE	CMS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $\lambda_{132} = \lambda_{231} = \lambda'_{311} = 0.01$	
>2300	95	15 KHACHATRY...16BE	CMS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $\lambda_{132} = \lambda_{231} = 0.07$, $\lambda'_{311} = 0.11$	OCCUR=2
>2000	95	16 AAD	15O ATLS	RPV ($e\mu$), $\tilde{\nu}_\tau$, $\lambda'_{311} = 0.11$, $\lambda_{i3k} = 0.07$	
>1700	95	16 AAD	15O ATLS	RPV ($\tau\mu$, $e\tau$), $\tilde{\nu}_\tau$, $\lambda'_{311} = 0.11$, $\lambda_{i3k} = 0.07$	OCCUR=2
		17 AAD	13AI ATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$, $e\tau$, $\mu\tau$	
		18 AAD	11H ATLS	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$	

		19	AALTONEN	10Z	CDF	RPV, $\tilde{\nu}_\tau \rightarrow e\mu, e\tau, \mu\tau$		
		20	ABAZOV	10M	D0	RPV, $\tilde{\nu}_\tau \rightarrow e\mu$		
>	95	95	21	ABDALLAH	04H	DLPH	AMSB, $\mu > 0$	
>	37.1	95	22	ADRIANI	93M	L3	$\Gamma(Z \rightarrow \text{invisible}); N(\tilde{\nu})=1$	OCCUR=2
>	36	95		ABREU	91F	DLPH	$\Gamma(Z \rightarrow \text{invisible}); N(\tilde{\nu})=1$	
>	31.2	95	23	ALEXANDER	91F	OPAL	$\Gamma(Z \rightarrow \text{invisible}); N(\tilde{\nu})=1$	
			1	AAD 23CB			searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for heavy particles decaying into an $e\mu, e\tau, \mu\tau$ final state. No significant deviation from the expected SM background is observed. Limits are set on the mass of a stau neutrino with R-parity-violating couplings, with decays $\tilde{\nu}_\tau \rightarrow e\mu, \tilde{\nu}_\tau \rightarrow e\tau, \tilde{\nu}_\tau \rightarrow \mu\tau$, see figures 4b, 5b, 6b.	NODE=S046SNU;LINKAGE=Z
			2	TUMASYAN 23H			searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of resonant $\tilde{\nu}_\tau$ production in events with two charged leptons, $e\mu, e\tau, \mu\tau$. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\nu}_\tau$ in an RPV model for resonant sneutrino production, where all RPV couplings vanish, except for those that are connected to the production and decay of the $\tilde{\nu}_\tau$, considering a SUSY mass hierarchy with $\tilde{\nu}_\tau$ as the LSP. The $\tilde{\nu}_\tau$ is produced resonantly through λ'_{311} coupling, and decays via λ_{i3k} coupling to two leptons, see their figure 3 for couplings of 0.1 and 0.01. Exclusion limits are also shown in the plane of $\tilde{\nu}_\tau$ mass and λ' coupling, for four values of λ couplings, see their figure 6. In addition, limits are set on heavy Z' gauge bosons with lepton flavor violating decays, see their figure 4, and on nonresonant quantum black hole production in models with extra spatial dimensions, see their figure 5. Model-independent upper limits on the product of the cross section, the branching fraction, acceptance, and efficiency are given as well, see their figure 7.	NODE=S046SNU;LINKAGE=Y
			3	AABOUD 18CM			searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for heavy particles decaying into an $e\mu, e\tau, \mu\tau$ final state. No significant deviation from the expected SM background is observed. Limits are set on the mass of a stau neutrino with R-parity-violating couplings. For $\tilde{\nu}_\tau \rightarrow e\mu$, masses below 3.4 TeV are excluded at 95% CL, see their Figure 4(b). Upper limits on the RPV couplings $ \lambda_{312} $ versus $ \lambda'_{311} $ are also performed, see their Figure 8(a-b).	NODE=S046SNU;LINKAGE=T
			4	AABOUD 18CM			searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for heavy particles decaying into an $e\mu, e\tau, \mu\tau$ final state. No significant deviation from the expected SM background is observed. Limits are set on the mass of a stau neutrino with R-parity-violating couplings. For $\tilde{\nu}_\tau \rightarrow e\tau$, masses below 2.9 TeV are excluded at 95% CL, see their Figure 5(b). Upper limits on the RPV couplings $ \lambda_{313} $ versus $ \lambda'_{311} $ are also performed, see their Figure 8(c).	NODE=S046SNU;LINKAGE=U
			5	AABOUD 18CM			searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for heavy particles decaying into an $e\mu, e\tau, \mu\tau$ final state. No significant deviation from the expected SM background is observed. Limits are set on the mass of a stau neutrino with R-parity-violating couplings. For $\tilde{\nu}_\tau \rightarrow \mu\tau$, masses below 2.6 TeV are excluded at 95% CL, see their Figure 6(b). Upper limits on the RPV couplings $ \lambda_{323} $ versus $ \lambda'_{311} $ are also performed, see their Figure 8(d).	NODE=S046SNU;LINKAGE=W
			6	AABOUD 18Z			searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more charged leptons (electrons, muons and up to two hadronically decaying taus). No significant deviation from the expected SM background is observed. Limits are set on the Higgsino mass in simplified models of general gauge mediated supersymmetry Tn1n1A/Tn1n1B/Tn1n1C, see their Figure 9. Limits are also set on the wino, slepton, sneutrino and gluino mass in a simplified model of NLSP pair production with R-parity violating decays of the LSP via λ_{12k} or λ_{i33} to charged leptons, see their Figures 7, 8.	NODE=S046SNU;LINKAGE=S
			7	SIRUNYAN 18AT			searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for heavy resonances decaying into $e\mu$ final states. No significant excess above the Standard Model expectation is observed and 95% C.L. exclusions are placed on the cross section times branching ratio for the R-parity-violating production and decay of a supersymmetric tau sneutrino, see their Fig. 3.	NODE=S046SNU;LINKAGE=R
			8	AABOUD 16P			searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with different flavour dilepton pairs ($e\mu, e\tau, \mu\tau$) from the production of $\tilde{\nu}_\tau$ via an RPV λ'_{311} coupling and followed by a decay via $\lambda_{312} = \lambda_{321} = 0.07$ for $e + \mu$, via $\lambda_{313} = \lambda_{331} = 0.07$ for $e + \tau$ and via $\lambda_{323} = \lambda_{332} = 0.07$ for $\mu + \tau$. No evidence for a dilepton resonance over the SM expectation is observed, and limits are derived on $m_{\tilde{\nu}}$ at 95% CL, see their Figs. 2(b), 3(b), 4(b), and Table 3.	NODE=S046SNU;LINKAGE=V
			9	AAD 14X			searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least four leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the sneutrino mass in an R-parity violating simplified model where the decay $\tilde{\nu} \rightarrow \nu \tilde{\chi}_1^0$, with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, takes place with a branching ratio of 100%, see Fig. 9.	NODE=S046SNU;LINKAGE=O
			10	AAD 11Z			looked in 1.07 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with one electron and one muon of opposite charge from the production of $\tilde{\nu}_\tau$ via an RPV λ'_{311} coupling and followed by a decay via λ_{312} into $e + \mu$. No evidence for an (e, μ) resonance over the SM expectation is observed, and a limit is derived in the plane of λ'_{311} versus $m_{\tilde{\nu}}$ for three values of λ_{312} , see their Fig. 2. Masses $m_{\tilde{\nu}} < 1.32$ (1.45) TeV are excluded for $\lambda'_{311} = 0.10$ and $\lambda_{312} = 0.05$ ($\lambda'_{311} = 0.11$ and $\lambda_{312} = 0.07$).	NODE=S046SNU;LINKAGE=D1

- 11 ABDALLAH 03M uses data from $\sqrt{s} = 192\text{--}208$ GeV to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays) and for sleptons. These limits are valid for values of $M_2 < 1$ TeV, $|\mu| \leq 1$ TeV with the $\tilde{\chi}_1^0$ as LSP. The quoted limit is obtained when there is no mixing in the third family. See Fig. 43 for the mass limits as a function of $\tan\beta$. These limits update the results of ABREU 00W. NODE=S046SNU;LINKAGE=AA
- 12 HEISTER 02N derives a bound on $m_{\tilde{\nu}_e}$ by exploiting the mass relation between the $\tilde{\nu}_e$ and \tilde{e} , based on the assumption of universal GUT scale gaugino and scalar masses $m_{1/2}$ and m_0 and the search described in the \tilde{e} section. In the MSUGRA framework with radiative electroweak symmetry breaking, the limit improves to $m_{\tilde{\nu}_e} > 130$ GeV, assuming a trilinear coupling $A_0=0$ at the GUT scale. See Figs. 5 and 7 for the dependence of the limits on $\tan\beta$. NODE=S046SNU;LINKAGE=HN
- 13 DECAMP 92 limit is from $\Gamma(\text{invisible})/\Gamma(\ell\ell) = 5.91 \pm 0.15$ ($N_\nu = 2.97 \pm 0.07$). NODE=S046SNU;LINKAGE=YY
- 14 SIRUNYAN 19AO searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing two same-sign muons and at last two jets, originating from resonant production of second-generation sleptons ($\tilde{\mu}_L, \tilde{\nu}_\mu$) via the R-parity violating coupling λ'_{211} to quarks. No significant excess above the Standard Model expectations is observed. Upper limits on cross sections are derived in the context of two simplified models, see their Figure 4. The cross section limits are translated into limits on λ'_{211} for a modified CMSSM, see their Figure 5. NODE=S046SNU;LINKAGE=X
- 15 KHACHATRYAN 16BE searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for evidence of narrow resonances decaying into $e\mu$ final states. No significant excess above the Standard Model expectation is observed and 95% C.L. exclusions are placed on the cross section times branching ratio for the production of an R-parity-violating supersymmetric tau sneutrino, see their Fig. 3. NODE=S046SNU;LINKAGE=Q
- 16 AAD 15O searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for evidence of heavy particles decaying into $e\mu, e\tau$ or $\mu\tau$ final states. No significant excess above the Standard Model expectation is observed, and 95% C.L. exclusions are placed on the cross section times branching ratio for the production of an R-parity-violating supersymmetric tau sneutrino, applicable to any sneutrino flavour, see their Fig. 2. NODE=S046SNU;LINKAGE=P
- 17 AAD 13AI searched in 4.6 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for evidence of heavy particles decaying into $e\mu, e\tau$ or $\mu\tau$ final states. No significant excess above the Standard Model expectation is observed, and 95% C.L. exclusions are placed on the cross section times branching ratio for the production of an R-parity-violating supersymmetric tau sneutrino, see their Fig. 2. For couplings $\lambda'_{311} = 0.10$ and $\lambda_{i3k} = 0.05$, the lower limits on the $\tilde{\nu}_\tau$ mass are 1610, 1110, 1100 GeV in the $e\mu, e\tau$, and $\mu\tau$ channels, respectively. NODE=S046SNU;LINKAGE=YA
- 18 AAD 11H looked in 35 pb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with one electron and one muon of opposite charge from the production of $\tilde{\nu}_\tau$ via an RPV λ'_{311} coupling and followed by a decay via λ_{312} into $e + \mu$. No evidence for an excess over the SM expectation is observed, and a limit is derived in the plane of λ'_{311} versus $m_{\tilde{\nu}_\tau}$ for several values of λ_{312} , see their Fig. 2. Superseded by AAD 11Z. NODE=S046SNU;LINKAGE=A1
- 19 AALTONEN 10Z searched in 1 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV for events from the production $d\bar{d} \rightarrow \tilde{\nu}_\tau$ with the subsequent decays $\tilde{\nu}_\tau \rightarrow e\mu, \mu\tau, e\tau$ in the MSSM framework with RPV. Two isolated leptons of different flavor and opposite charges are required, with τ s identified by their hadronic decay. No statistically significant excesses are observed over the SM background. Upper limits on λ'_{311} times the branching ratio are listed in their Table III for various $\tilde{\nu}_\tau$ masses. Limits on the cross section times branching ratio for $\lambda'_{311} = 0.10$ and $\lambda_{i3k} = 0.05$, displayed in Fig. 2, are used to set limits on the $\tilde{\nu}_\tau$ mass of 558 GeV for the $e\mu$, 441 GeV for the $\mu\tau$ and 442 GeV for the $e\tau$ channels. NODE=S046SNU;LINKAGE=LT
- 20 ABAZOV 10M looked in 5.3 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV for events with exactly one pair of high p_T isolated $e\mu$ and a veto against hard jets. No evidence for an excess over the SM expectation is observed, and a limit at 95% C.L. on the cross section times branching ratio is derived, see their Fig. 3. These limits are translated into limits on couplings as a function of $m_{\tilde{\nu}_\tau}$ as shown on their Fig. 4. As an example, for $m_{\tilde{\nu}_\tau} = 100$ GeV and $\lambda_{312} \leq 0.07$, couplings $\lambda'_{311} > 7.7 \times 10^{-4}$ are excluded. NODE=S046SNU;LINKAGE=BZ
- 21 ABDALLAH 04H use data from LEP 1 and $\sqrt{s} = 192\text{--}208$ GeV. They re-use results or re-analyze the data from ABDALLAH 03M to put limits on the parameter space of anomaly-mediated supersymmetry breaking (AMSB), which is scanned in the region $1 < m_{3/2} < 50$ TeV, $0 < m_0 < 1000$ GeV, $1.5 < \tan\beta < 35$, both signs of μ . The constraints are obtained from the searches for mass degenerate chargino and neutralino, for SM-like and invisible Higgs, for leptonically decaying charginos and from the limit on non-SM Z width of 3.2 MeV. The limit is for $m_{\tilde{t}} = 174.3$ GeV (see Table 2 for other $m_{\tilde{t}}$ values). The limit improves to 114 GeV for $\mu < 0$. NODE=S046SNU;LINKAGE=BD
- 22 ADRIANI 93M limit from $\Delta\Gamma(Z)(\text{invisible}) < 16.2$ MeV. NODE=S046SNU;LINKAGE=DM
- 23 ALEXANDER 91F limit is for one species of $\tilde{\nu}$ and is derived from $\Gamma(\text{invisible, new})/\Gamma(\ell\ell) < 0.38$. NODE=S046SNU;LINKAGE=I

Charged sleptons

This section contains limits on charged scalar leptons ($\tilde{\ell}$, with $\ell=e,\mu,\tau$). Studies of width and decays of the Z boson (use is made here of $\Delta\Gamma_{\text{inv}} < 2.0 \text{ MeV}$, LEP 00) conclusively rule out $m_{\tilde{\ell}_R} < 40 \text{ GeV}$ (41

GeV for $\tilde{\ell}_L$), independently of decay modes, for each individual slepton. The limits improve to 43 GeV (43.5 GeV for $\tilde{\ell}_L$) assuming all 3 flavors to be degenerate. Limits on higher mass sleptons depend on model assumptions and on the mass splitting $\Delta m = m_{\tilde{\ell}} - m_{\tilde{\chi}_1^0}$. The mass and composition

of $\tilde{\chi}_1^0$ may affect the selectron production rate in e^+e^- collisions through t-channel exchange diagrams. Production rates are also affected by the potentially large mixing angle of the lightest mass eigenstate $\tilde{\ell}_1 = \tilde{\ell}_R \sin\theta_\ell + \tilde{\ell}_L \cos\theta_\ell$. It is generally assumed that only $\tilde{\tau}$ may have significant mixing. The coupling to the Z vanishes for $\theta_\ell=0.82$. In the high-energy limit of e^+e^- collisions the interference between γ and Z exchange leads to a minimal cross section for $\theta_\ell=0.91$, a value which is sometimes used in the following entries relative to data taken at LEP2. When limits on $m_{\tilde{\ell}_R}$ are quoted, it is understood that limits on $m_{\tilde{\ell}_L}$ are usually at least as strong.

Possibly open decays involving gauginos other than $\tilde{\chi}_1^0$ will affect the detection efficiencies. Unless otherwise stated, the limits presented here result from the study of $\tilde{\ell}^+\tilde{\ell}^-$ production, with production rates and decay properties derived from the MSSM. Limits made obsolete by the recent analyses of e^+e^- collisions at high energies can be found in previous Editions of this Review.

For decays with final state gravitinos (\tilde{G}), $m_{\tilde{G}}$ is assumed to be negligible relative to all other masses.

NODE=S046210

NODE=S046210

R-parity conserving \tilde{e} (Selectron) mass limit

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

NODE=S046SE

NODE=S046SE

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>270	95	¹ AAD	23M ATLS	$2\ell, \tilde{\ell}$ pair production, $m_{\tilde{e}_L} = m_{\tilde{e}_R}$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
> 90	95	¹ AAD	23M ATLS	$2\ell, \tilde{\ell}$ pair production, $m_{\tilde{e}_L} = m_{\tilde{e}_R}$, $m_{\tilde{e}} - m_{\tilde{\chi}_1^0} = 26 \text{ GeV}$
>700	95	² SIRUNYAN	21M CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
>700	95	³ AAD	200 ATLS	$2\ell + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
>250	95	⁴ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{e}_R, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
>310	95	⁴ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{e}_L, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
>350	95	⁴ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, m_{\tilde{e}_R} = m_{\tilde{e}_L}, m_{\tilde{\chi}_1^0}$ $= 0 \text{ GeV}$
>290	95	⁴ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{\ell}_R$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
>400	95	⁴ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{\ell}_L$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0}$ $= 0 \text{ GeV}$
>450	95	⁴ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
>500	95	⁵ AABOUD	18BT ATLS	$2\ell + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e},$ $\tilde{\mu}, \tilde{\tau}$, with $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
>190	95	⁶ AABOUD	18R ATLS	2ℓ (soft) + $\cancel{E}_T, m_{\tilde{e}} = m_{\tilde{\mu}}, m_{\tilde{e}} -$ $m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$
		⁷ CHATRCHYAN14R	CMS	$\geq 3\ell^\pm, \tilde{\ell} \rightarrow \ell^\pm \tau^\mp \tau^\mp \tilde{G}$ sim- plified model, GMSB, stau (N)NLSP scenario

NODE=S046SE;CHECK LIMITS

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

		8	AAD	13B	ATLS	$2\ell^\pm + \cancel{E}_T$, SMS, pMSSM	
> 97.5		9	ABBIENDI	04	OPAL	$\tilde{e}_R, \Delta m > 11 \text{ GeV}, \mu > 100 \text{ GeV}, \tan\beta=1.5$	
> 94.4		10	ACHARD	04	L3	$\tilde{e}_R, \Delta m > 10 \text{ GeV}, \mu > 200 \text{ GeV}, \tan\beta \geq 2$	
> 71.3		10	ACHARD	04	L3	\tilde{e}_R , all Δm	OCCUR=2
none 30–94	95	11	ABDALLAH	03M	DLPH	$\Delta m > 15 \text{ GeV}, \tilde{e}_R^+ \tilde{e}_R^-$	
> 94	95	12	ABDALLAH	03M	DLPH	$\tilde{e}_R, 1 \leq \tan\beta \leq 40, \Delta m > 10 \text{ GeV}$	OCCUR=2
> 95	95	13	HEISTER	02E	ALEP	$\Delta m > 15 \text{ GeV}, \tilde{e}_R^+ \tilde{e}_R^-$	
> 73	95	14	HEISTER	02N	ALEP	\tilde{e}_R , any Δm	
>107	95	14	HEISTER	02N	ALEP	\tilde{e}_L , any Δm	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●							
>101	95	15	AAD	20I	ATLS	2ℓ (soft), jets, $\cancel{E}_T, \tilde{e}_R$ only, $m_{\tilde{e}_R} - m_{\tilde{\chi}_1^0} = 7.5 \text{ GeV}$	
>169	95	16	AAD	20I	ATLS	2ℓ (soft), jets, $\cancel{E}_T, \tilde{e}_L$ only, $m_{\tilde{e}_L} - m_{\tilde{\chi}_1^0} = 7.1 \text{ GeV}$	OCCUR=2
none 90–325	95	17	AAD	14G	ATLS	$\tilde{\ell}\tilde{\ell} \rightarrow \ell^+ \tilde{\chi}_1^0 \ell^- \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\ell}_L} = m_{\tilde{\ell}_R}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
		18	KHACHATRY...	14I	CMS	$\tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$, simplified model	
<p>¹ AAD 23M searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for $\tilde{\ell}^\pm$ pair production, followed by $\tilde{\ell}^\pm \rightarrow \ell^\pm \tilde{\chi}_1^0$ in events with two leptons. The focus is on models where $m_{\tilde{\ell}^\pm} - m_{\tilde{\chi}_1^0}$ is close to the W mass. No significant excess above the Standard Model predictions is observed. Limits were set on the $\tilde{\ell}$ mass (assuming $\tilde{e} - \tilde{\mu}$ and $L - R$ degeneracy), as a function of $m_{\tilde{\chi}_1^0}$, see Figure 6. Limits were also derived for single \tilde{e} or $\tilde{\mu}$, and for L and R independently, see Figure 7.</p>							NODE=S046SE;LINKAGE=RA
<p>² SIRUNYAN 21M searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tsb0t3, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14.</p>							NODE=S046SE;LINKAGE=QA
<p>³ AAD 200 reported on a search for electroweak production in models with charginos and sleptons decaying into final states with exactly two oppositely charged leptons and missing transverse momentum. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Light-flavour sleptons \tilde{e} and $\tilde{\mu}$ are constrained at 95% C.L. to have masses above 700 GeV for massless lightest neutralino, see their Fig. 7(c). Exclusion limits are also set for selectrons and smuons separately, considering either right- or left-handed components, by including only the di-electron and di-muon same-flavour signal regions defined in the search, see their Fig. 8.</p>							NODE=S046SE;LINKAGE=NA
<p>⁴ SIRUNYAN 19AW searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak pair production of selectrons or smuons in events with two leptons (electrons or muons) of the opposite electric charge and same flavour, no jets and large \cancel{E}_T. No significant excess above the Standard Model expectations is observed. Limits are set on the selectron mass assuming left-handed, right-handed or both left- and right-handed (mass degenerate) production, see their Figure 6. Similarly, limits are set on the smuon mass, see their Figure 7. Limits are also set on slepton masses under the assumption that the selectron and smuon are mass degenerate, see their Figure 5.</p>							NODE=S046SE;LINKAGE=JA
<p>⁵ AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the slepton mass up to 500 GeV for massless $\tilde{\chi}_1^0$, assuming degeneracy of $\tilde{e}, \tilde{\mu}$, and $\tilde{\tau}$ and exploiting the 2ℓ signature, see their Figure 8(b).</p>							NODE=S046SE;LINKAGE=GA
<p>⁶ AABOUD 18R searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for electroweak production in scenarios with compressed mass spectra in final states with two low-momentum leptons and missing transverse momentum. The data are found to be consistent with the SM prediction. Results are interpreted in slepton pair production models with a fourfold degeneracy assumed in selectron and smuon masses. The \tilde{e} masses are excluded up to 190 GeV for $m_{\tilde{e}} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$. The exclusion limits extend down to mass splittings of 1 GeV, see their Fig. 11.</p>							NODE=S046SE;LINKAGE=FA
<p>⁷ CHATRCHYAN 14R searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least three leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the slepton</p>							NODE=S046SE;LINKAGE=K

- mass in a stau (N)NLSP simplified model (GMSB) where the decay $\tilde{\ell} \rightarrow \ell^\pm \tau^\pm \tau^\mp \tilde{G}$ takes place with a branching ratio of 100%, see Fig. 8.
- 8 AAD 13B searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for sleptons decaying to a final state with two leptons (e and μ) and missing transverse energy. No excess beyond the Standard Model expectation is observed. Limits are derived in a simplified model of direct left-handed slepton pair production, where left-handed slepton masses between 85 and 195 GeV are excluded at 95% C.L. for $m_{\tilde{\chi}_1^0} = 20 \text{ GeV}$. See also Fig. 2(a). Exclusion limits are also derived in the phenomenological MSSM, see Fig. 3.
- 9 ABBIENDI 04 search for $\tilde{e}_R \tilde{e}_R$ production in acoplanar di-electron final states in the 183–208 GeV data. See Fig. 13 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$ and for the limit at $\tan\beta=35$. This limit supersedes ABBIENDI 00G.
- 10 ACHARD 04 search for $\tilde{e}_R \tilde{e}_L$ and $\tilde{e}_R \tilde{e}_R$ production in single- and acoplanar di-electron final states in the 192–209 GeV data. Absolute limits on $m_{\tilde{e}_R}$ are derived from a scan over the MSSM parameter space with universal GUT scale gaugino and scalar masses $m_{1/2}$ and m_0 , $1 \leq \tan\beta \leq 60$ and $-2 \leq \mu \leq 2 \text{ TeV}$. See Fig. 4 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$. This limit supersedes ACCIARRI 99W.
- 11 ABDALLAH 03M looked for acoplanar dielectron + \cancel{E} final states at $\sqrt{s} = 189\text{--}208 \text{ GeV}$. The limit assumes $\mu = -200 \text{ GeV}$ and $\tan\beta=1.5$ in the calculation of the production cross section and $B(\tilde{e} \rightarrow e \tilde{\chi}_1^0)$. See Fig. 15 for limits in the $(m_{\tilde{e}_R}, m_{\tilde{\chi}_1^0})$ plane. These limits include and update the results of ABREU 01.
- 12 ABDALLAH 03M uses data from $\sqrt{s} = 192\text{--}208 \text{ GeV}$ to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays) and for sleptons. These limits are valid for values of $M_2 < 1 \text{ TeV}$, $|\mu| \leq 1 \text{ TeV}$ with the $\tilde{\chi}_1^0$ as LSP. The quoted limit is obtained when there is no mixing in the third family. See Fig. 43 for the mass limits as a function of $\tan\beta$. These limits update the results of ABREU 00W.
- 13 HEISTER 02E looked for acoplanar dielectron + \cancel{E}_T final states from $e^+ e^-$ interactions between 183 and 209 GeV. The mass limit assumes $\mu < -200 \text{ GeV}$ and $\tan\beta=2$ for the production cross section and $B(\tilde{e} \rightarrow e \tilde{\chi}_1^0)=1$. See their Fig. 4 for the dependence of the limit on Δm . These limits include and update the results of BARATE 01.
- 14 HEISTER 02N search for $\tilde{e}_R \tilde{e}_L$ and $\tilde{e}_R \tilde{e}_R$ production in single- and acoplanar di-electron final states in the 183–208 GeV data. Absolute limits on $m_{\tilde{e}_R}$ are derived from a scan over the MSSM parameter space with universal GUT scale gaugino and scalar masses $m_{1/2}$ and m_0 , $1 \leq \tan\beta \leq 50$ and $-10 \leq \mu \leq 10 \text{ TeV}$. The region of small $|\mu|$, where cascade decays are important, is covered by a search for $\tilde{\chi}_1^0 \tilde{\chi}_3^0$ in final states with leptons and possibly photons. Limits on $m_{\tilde{e}_L}$ are derived by exploiting the mass relation between the \tilde{e}_L and \tilde{e}_R , based on universal m_0 and $m_{1/2}$. When the constraint from the mass limit of the lightest Higgs from HEISTER 02 is included, the bounds improve to $m_{\tilde{e}_R} > 77(75) \text{ GeV}$ and $m_{\tilde{e}_L} > 115(115) \text{ GeV}$ for a top mass of 175(180) GeV. In the MSUGRA framework with radiative electroweak symmetry breaking, the limits improve further to $m_{\tilde{e}_R} > 95 \text{ GeV}$ and $m_{\tilde{e}_L} > 152 \text{ GeV}$, assuming a trilinear coupling $A_0=0$ at the GUT scale. See Figs. 4, 5, 7 for the dependence of the limits on $\tan\beta$.
- 15 AAD 20i reported on ATLAS searches for slepton pair production in models with compressed mass spectra. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Events with \cancel{E}_T , two same-flavour, opposite-charge, low-transverse-momentum leptons, and jets from initial-state radiation or characteristic of vector-boson fusion production are selected. Light-flavour sleptons \tilde{e} and $\tilde{\mu}$ are constrained at 95% C.L. to have masses above 251 GeV for a mass splitting slepton– $\tilde{\chi}_1^0$ of 10 GeV, with constraints extending down to mass splittings of 550 MeV at the LEP slepton limits (73 GeV), see their Fig. 16(a). If only selectrons are considered, and $\tilde{e} = \tilde{e}_R$, masses below 101 GeV are excluded for mass splitting $\tilde{e}_R, \tilde{\chi}_1^0$ of 7.5 GeV. See their Fig. 16(b).
- 16 AAD 20i reported on ATLAS searches for slepton pair production in models with compressed mass spectra. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Events with \cancel{E}_T , two same-flavour, opposite-charge, low-transverse-momentum leptons, and jets from initial-state radiation or characteristic of vector-boson fusion production are selected. Light-flavour sleptons \tilde{e} and $\tilde{\mu}$ are constrained at 95% C.L. to have masses above 251 GeV for a mass splitting slepton– $\tilde{\chi}_1^0$ of 10 GeV, with constraints extending down to mass splittings of 550 MeV at the LEP slepton limits (73 GeV). See their Fig. 16(a). If only selectron are considered, and $\tilde{e} = \tilde{e}_L$, masses below 169 GeV are excluded for mass splitting $\tilde{e}_L, \tilde{\chi}_1^0$ of 7.1 GeV. See their Fig. 16(b).
- 17 AAD 14G searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of slepton pairs, decaying to a final state with two leptons (e and μ) and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of slepton pair production, see Fig. 8. An interpretation in the pMSSM is also given, see Fig. 10.
- 18 KHACHATRYAN 14i searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for electroweak production of slepton pairs decaying to a final state with opposite-sign lepton

pairs (e or μ) and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models, see Fig. 18.

R-parity violating \tilde{e} (Selectron) mass limit

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1200	95	¹ AAD	21Y ATLS	$\geq 4\ell, \lambda_{12k} \neq 0, m_{\tilde{\chi}_1^0} = 900$ GeV (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations)
> 870	95	¹ AAD	21Y ATLS	$\geq 4\ell, \lambda_{i33} \neq 0, m_{\tilde{\chi}_1^0} = 450$ GeV (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations)
>1065	95	² AABOUD	18Z ATLS	$\geq 4\ell, \lambda_{12k} \neq 0, m_{\tilde{\chi}_1^0} = 600$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)
> 780	95	² AABOUD	18Z ATLS	$\geq 4\ell, \lambda_{i33} \neq 0, m_{\tilde{\chi}_1^0} = 300$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)
> 410	95	³ AAD	14X ATLS	$\geq 4\ell^\pm, \tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 89	95	⁴ ABBIENDI	04F OPAL	\tilde{e}_L
> 92	95	⁵ ABDALLAH	04M DLPH	\tilde{e}_R , indirect, $\Delta m > 5$ GeV

¹AAD 21Y searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for supersymmetry in events with four or more leptons (electrons, muons and tau-leptons). No significant excess above the Standard Model expectations is observed. Limits are set on Tchi1n12-GGM, and RPV models similar to Tchi1n2l, Tglu1A (with $q = u, d, s, c, b$, with equal branching fractions), and $\tilde{\ell}_L/\tilde{\nu} \rightarrow \ell/\nu \tilde{\chi}_1^0$ (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations), all with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$ via λ_{12k} or λ_{i33} (where $i, k \in 1, 2$), see their Figure 11.

²AABOUD 18Z searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing four or more charged leptons (electrons, muons and up to two hadronically decaying taus). No significant deviation from the expected SM background is observed. Limits are set on the Higgsino mass in simplified models of general gauge mediated supersymmetry Tn1n1A/Tn1n1B/Tn1n1C, see their Figure 9. Limits are also set on the wino, slepton, sneutrino and gluino mass in a simplified model of NLSP pair production with R-parity violating decays of the LSP via λ_{12k} or λ_{i33} to charged leptons, see their Figures 7, 8.

³AAD 14X searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with at least four leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the slepton mass in an R-parity violating simplified model where the decay $\tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$, with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, takes place with a branching ratio of 100%, see Fig. 9.

⁴ABBIENDI 04F use data from $\sqrt{s} = 189\text{--}209$ GeV. They derive limits on sparticle masses under the assumption of RPV with $LL\bar{E}$ or $LQ\bar{D}$ couplings. The results are valid for $\tan\beta = 1.5$, $\mu = -200$ GeV, with, in addition, $\Delta m > 5$ GeV for indirect decays via $LQ\bar{D}$. The limit quoted applies to direct decays via $LL\bar{E}$ or $LQ\bar{D}$ couplings. For indirect decays, the limits on the \tilde{e}_R mass are respectively 99 and 92 GeV for $LL\bar{E}$ and $LQ\bar{D}$ couplings and $m_{\tilde{\chi}_1^0} = 10$ GeV and degrade slightly for larger $\tilde{\chi}_1^0$ mass. Supersedes the results of ABBIENDI 00.

⁵ABDALLAH 04M use data from $\sqrt{s} = 192\text{--}208$ GeV to derive limits on sparticle masses under the assumption of RPV with $LL\bar{E}$ or $UD\bar{D}$ couplings. The results are valid for $\mu = -200$ GeV, $\tan\beta = 1.5$, $\Delta m > 5$ GeV and assuming a BR of 1 for the given decay. The limit quoted is for indirect $UD\bar{D}$ decays using the neutralino constraint of 39.5 GeV for $LL\bar{E}$ and of 38.0 GeV for $UD\bar{D}$ couplings, also derived in ABDALLAH 04M. For indirect decays via $LL\bar{E}$ the limit improves to 95 GeV if the constraint from the neutralino is used and to 94 GeV if it is not used. For indirect decays via $UD\bar{D}$ couplings it remains unchanged when the neutralino constraint is not used. Supersedes the result of ABREU 00U.

R-parity conserving $\tilde{\mu}$ (Smuon) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 220–460	95	¹ AAD	23CR ATLS	2 same-sign, 3, 4 ℓ , 1, 2 b -jets, $\tilde{\mu}_{L,R}$ pair production with $\tilde{\mu}_{L,R} \rightarrow \mu \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow b + \ell/\nu + t/b$ via λ'_{i33} coupling

NODE=S046SEV

NODE=S046SEV

NODE=S046SEV;CHECK LIMITS

OCCUR=2

OCCUR=2

NODE=S046SEV;LINKAGE=B

NODE=S046SEV;LINKAGE=A

NODE=S046SEV;LINKAGE=BA

NODE=S046SEV;LINKAGE=IA

NODE=S046SEV;LINKAGE=DH

NODE=S046SMU

NODE=S046SMU;CHECK LIMITS

>240	95	² AAD	23M ATLS	$2\ell, \tilde{\ell}$ pair production, $m_{\tilde{\mu}_L} = m_{\tilde{\mu}_R}, m_{\tilde{\chi}_1^0} = 0$ GeV	
> 90	95	² AAD	23M ATLS	$2\ell, \tilde{\ell}$ pair production, $m_{\tilde{\mu}_L} = m_{\tilde{\mu}_R}, m_{\tilde{\mu}} - m_{\tilde{\chi}_1^0} = 32$ GeV	OCCUR=2
>700	95	³ SIRUNYAN	21M CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0$ GeV	
>150	95	⁴ AAD	20I ATLS	2ℓ (soft), jets, $\cancel{E}_T, \tilde{\mu}_R$ only, $m_{\tilde{\mu}_R} - m_{\tilde{\chi}_1^0} = 8.2$ GeV	
>216	95	⁵ AAD	20I ATLS	2ℓ (soft), jets, $\cancel{E}_T, \tilde{\mu}_L$ only, $m_{\tilde{\mu}_L} - m_{\tilde{\chi}_1^0} = 10$ GeV	OCCUR=2
>700	95	⁶ AAD	200 ATLS	$2\ell + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0$ GeV	
>210	95	⁷ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{\mu}_R, m_{\tilde{\chi}_1^0} = 0$ GeV	
>280	95	⁷ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{\mu}_L, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
>290	95	⁷ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{\ell}_R$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
>400	95	⁷ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{\ell}_L$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>450	95	⁷ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
>310	95	⁷ SIRUNYAN	19AW CMS	$\ell^\pm \ell^\mp + \cancel{E}_T, m_{\tilde{\mu}_R} = m_{\tilde{\mu}_L}, m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=6
>190	95	⁸ AABOUD	18R ATLS	2ℓ (soft) + $\cancel{E}_T, m_{\tilde{e}} = m_{\tilde{\mu}}, m_{\tilde{\mu}} - m_{\tilde{\chi}_1^0} = 5$ GeV	
		⁹ CHATRCHYAN	14R CMS	$\geq 3\ell^\pm, \tilde{\ell} \rightarrow \ell^\pm \tau^\mp \tau^\mp \tilde{G}$ simplified model, GMSB, stau (N)NLSP scenario	
		¹⁰ AAD	13B ATLS	$2\ell^\pm + \cancel{E}_T, \text{SMS, pMSSM}$	
> 91.0		¹¹ ABBIENDI	04 OPAL	$\Delta m > 3$ GeV, $\tilde{\mu}_R^+ \tilde{\mu}_R^-$, $ \mu > 100$ GeV, $\tan\beta = 1.5$	
> 86.7		¹² ACHARD	04 L3	$\Delta m > 10$ GeV, $\tilde{\mu}_R^+ \tilde{\mu}_R^-$, $ \mu > 200$ GeV, $\tan\beta \geq 2$	
none 30–88	95	¹³ ABDALLAH	03M DLPH	$\Delta m > 5$ GeV, $\tilde{\mu}_R^+ \tilde{\mu}_R^-$	
> 94	95	¹⁴ ABDALLAH	03M DLPH	$\tilde{\mu}_R, 1 \leq \tan\beta \leq 40, \Delta m > 10$ GeV	OCCUR=2
> 88	95	¹⁵ HEISTER	02E ALEP	$\Delta m > 15$ GeV, $\tilde{\mu}_R^+ \tilde{\mu}_R^-$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>500	95	¹⁶ AABOUD	18BT ATLS	$2\ell + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$ and $\tilde{\ell} = \tilde{e}, \tilde{\mu}, \tilde{\tau}$, with $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 90–325	95	¹⁷ AAD	14G ATLS	$\tilde{\ell}\tilde{\ell} \rightarrow \ell^+ \tilde{\chi}_1^0 \ell^- \tilde{\chi}_1^0$, simplified model, $m_{\tilde{\ell}_L} = m_{\tilde{\ell}_R}, m_{\tilde{\chi}_1^0} = 0$ GeV	
> 80	95	¹⁸ KHACHATRY... ¹⁹ ABREU	14I CMS 00V DLPH	$\tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$, simplified model $\tilde{\mu}_R \tilde{\mu}_R (\tilde{\mu}_R \rightarrow \mu \tilde{G}), m_{\tilde{G}} > 8$ eV	OCCUR=2

¹ AAD 23CR searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for RPV SUSY in final states with multiple leptons and b-tagged jets. No significant excess above the Standard Model expectations is observed. Limits are set on the production of electroweakinos (wino or higgsino) that decay via RPV coupling λ'_{i33} to a charged lepton or a neutrino, a b quark, and an additional t or b quark, see their figure 16. A second model addresses direct $\tilde{\mu}_{L,R}$ production and decay to a muon and a bino-like neutralino, which decays in the same way as in the first model, see their figure 17.

² AAD 23M searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for $\tilde{\ell}^\pm$ pair production, followed by $\tilde{\ell}^\pm \rightarrow \ell^\pm \tilde{\chi}_1^0$ in events with two leptons. The focus is on models where $m_{\tilde{\ell}^\pm} - m_{\tilde{\chi}_1^0}$ is close to the W mass. No significant excess above the Standard Model predictions is observed. Limits were set on the $\tilde{\ell}$ mass (assuming $\tilde{e} - \tilde{\mu}$ and $L - R$ degeneracy), as a function of $m_{\tilde{\chi}_1^0}$, see Figure 6. Limits were also derived for single \tilde{e} or $\tilde{\mu}$, and for L and R independently, see Figure 7.

NODE=S046SMU;LINKAGE=S

NODE=S046SMU;LINKAGE=Q

- 3 SIRUNYAN 21M searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tsbob3, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14.
- 4 AAD 20i reported on ATLAS searches for slepton pair production in models with compressed mass spectra. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Events with \cancel{E}_T , two same-flavour, opposite-charge, low-transverse-momentum leptons, and jets from initial-state radiation or characteristic of vector-boson fusion production are selected. Light-flavour sleptons \tilde{e} and $\tilde{\mu}$ are constrained at 95% C.L. to have masses above 251 GeV for a mass splitting slepton- $\tilde{\chi}_1^0$ of 10 GeV, with constraints extending down to mass splittings of 550 MeV at the LEP slepton limits (73 GeV). See their Fig. 16(a). If only smuon are considered, and $\tilde{\mu} = \tilde{\mu}_R$, masses below 150 GeV are excluded for mass splitting $\tilde{\mu}_R, \tilde{\chi}_1^0$ of 8.2 GeV. See their Fig. 16(b).
- 5 AAD 20i reported on ATLAS searches for slepton pair production in models with compressed mass spectra. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Events with \cancel{E}_T , two same-flavour, opposite-charge, low-transverse-momentum leptons, and jets from initial-state radiation or characteristic of vector-boson fusion production are selected. Light-flavour sleptons \tilde{e} and $\tilde{\mu}$ are constrained at 95% C.L. to have masses above 251 GeV for a mass splitting slepton- $\tilde{\chi}_1^0$ of 10 GeV, with constraints extending down to mass splittings of 550 MeV at the LEP slepton limits (73 GeV). See their Fig. 16(a). If only smuon are considered, and $\tilde{\mu} = \tilde{\mu}_L$, masses below 216 GeV are excluded for mass splitting $\tilde{\mu}_L, \tilde{\chi}_1^0$ of 10 GeV. See their Fig. 16(b).
- 6 AAD 200 reported on a search for electroweak production in models with charginos and sleptons decaying into final states with exactly two oppositely charged leptons and missing transverse momentum. A dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} was used. Light-flavour sleptons \tilde{e} and $\tilde{\mu}$ are constrained at 95% C.L. to have masses above 700 GeV for massless lightest neutralino, see their Fig. 7(c). Exclusion limits are also set for selectrons and smuons separately, considering either right- or left-handed components, by including only the di-electron and di-muon same-flavour signal regions defined in the search, see their Fig. 8.
- 7 SIRUNYAN 19AW searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak pair production of selectrons or smuons in events with two leptons (electrons or muons) of the opposite electric charge and same flavour, no jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the selectron mass assuming left-handed, right-handed or both left- and right-handed (mass degenerate) production, see their Figure 6. Similarly, limits are set on the smuon mass, see their Figure 7. Limits are also set on slepton masses under the assumption that the selectron and smuon are mass degenerate, see their Figure 5.
- 8 AABOUD 18R searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for electroweak production in scenarios with compressed mass spectra in final states with two low-momentum leptons and missing transverse momentum. The data are found to be consistent with the SM prediction. Results are interpreted in slepton pair production models with a fourfold degeneracy assumed in selectron and smuon masses. The $\tilde{\mu}$ masses are excluded up to 190 GeV for $m_{\tilde{\mu}} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$. The exclusion limits extend down to mass splittings of 1 GeV, see their Fig. 11.
- 9 CHATRCHYAN 14R searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least three leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the slepton mass in a stau (N)NLSP simplified model (GMSB) where the decay $\tilde{\ell} \rightarrow \ell^\pm \tau^\pm \tilde{\tau} \tilde{G}$ takes place with a branching ratio of 100%, see Fig. 8.
- 10 AAD 13B searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for sleptons decaying to a final state with two leptons (e and μ) and missing transverse energy. No excess beyond the Standard Model expectation is observed. Limits are derived in a simplified model of direct left-handed slepton pair production, where left-handed slepton masses between 85 and 195 GeV are excluded at 95% C.L. for $m_{\tilde{\chi}_1^0} = 20 \text{ GeV}$. See also Fig. 2(a). Exclusion limits are also derived in the phenomenological MSSM, see Fig. 3.
- 11 ABBIENDI 04 search for $\tilde{\mu}_R \tilde{\mu}_R$ production in acoplanar di-muon final states in the 183–208 GeV data. See Fig. 14 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$ and for the limit at $\tan\beta=35$. Under the assumption of 100% branching ratio for $\tilde{\mu}_R \rightarrow \mu \tilde{\chi}_1^0$, the limit improves to 94.0 GeV for $\Delta m > 4 \text{ GeV}$. See Fig. 11 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$ at several values of the branching ratio. This limit supersedes ABBIENDI 00G.
- 12 ACHARD 04 search for $\tilde{\mu}_R \tilde{\mu}_R$ production in acoplanar di-muon final states in the 192–209 GeV data. Limits on $m_{\tilde{\mu}_R}$ are derived from a scan over the MSSM parameter space with universal GUT scale gauging and scalar masses $m_{1/2}$ and m_0 , $1 \leq$

NODE=S046SMU;LINKAGE=R

NODE=S046SMU;LINKAGE=N

NODE=S046SMU;LINKAGE=O

NODE=S046SMU;LINKAGE=P

NODE=S046SMU;LINKAGE=M

NODE=S046SMU;LINKAGE=K

NODE=S046SMU;LINKAGE=G

NODE=S046SMU;LINKAGE=CD

NODE=S046SMU;LINKAGE=AB

NODE=S046SMU;LINKAGE=AC

$\tan\beta \leq 60$ and $-2 \leq \mu \leq 2$ TeV. See Fig. 4 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$.

This limit supersedes ACCIARRI 99W.

- 13 ABDALLAH 03M looked for acoplanar dimuon $+ \cancel{E}$ final states at $\sqrt{s} = 189\text{--}208$ GeV. The limit assumes $B(\tilde{\mu} \rightarrow \mu \tilde{\chi}_1^0) = 100\%$. See Fig. 16 for limits on the $(m_{\tilde{\mu}_R}, m_{\tilde{\chi}_1^0})$ plane. These limits include and update the results of ABREU 01. NODE=S046SMU;LINKAGE=AA
- 14 ABDALLAH 03M uses data from $\sqrt{s} = 192\text{--}208$ GeV to obtain limits in the framework of the MSSM with gaugino and sfermion mass universality at the GUT scale. An indirect limit on the mass is derived by constraining the MSSM parameter space by the results from direct searches for neutralinos (including cascade decays) and for sleptons. These limits are valid for values of $M_2 < 1$ TeV, $|\mu| \leq 1$ TeV with the $\tilde{\chi}_1^0$ as LSP. The quoted limit is obtained when there is no mixing in the third family. See Fig. 43 for the mass limits as a function of $\tan\beta$. These limits update the results of ABREU 00W. NODE=S046SMU;LINKAGE=AL
- 15 HEISTER 02E looked for acoplanar dimuon $+ \cancel{E}_T$ final states from e^+e^- interactions between 183 and 209 GeV. The mass limit assumes $B(\tilde{\mu} \rightarrow \mu \tilde{\chi}_1^0) = 1$. See their Fig. 4 for the dependence of the limit on Δm . These limits include and update the results of BARATE 01. NODE=S046SMU;LINKAGE=HE
- 16 AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the slepton mass up to 500 GeV for massless $\tilde{\chi}_1^0$, assuming degeneracy of \tilde{e} , $\tilde{\mu}$, and $\tilde{\tau}$ and exploiting the 2ℓ signature, see their Figure 8(b). NODE=S046SMU;LINKAGE=L
- 17 AAD 14G searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for electroweak production of slepton pairs, decaying to a final state with two leptons (e and μ) and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models of slepton pair production, see Fig. 8. An interpretation in the pMSSM is also given, see Fig. 10. NODE=S046SMU;LINKAGE=J
- 18 KHACHATRYAN 14I searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for electroweak production of slepton pairs decaying to a final state with opposite-sign lepton pairs (e or μ) and missing transverse momentum. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in simplified models, see Fig. 18. NODE=S046SMU;LINKAGE=I
- 19 ABREU 00V use data from $\sqrt{s} = 130\text{--}189$ GeV to search for tracks with large impact parameter or visible decay vertices. Limits are obtained as function of $m_{\tilde{G}}$, after combining these results with the search for slepton pair production in the SUGRA framework from ABREU 01 to cover prompt decays and on stable particle searches from ABREU 00Q. For limits at different $m_{\tilde{G}}$, see their Fig. 12. NODE=S046SMU;LINKAGE=VU

R-parity violating $\tilde{\mu}$ (Smuon) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 120–645	95	1 AAD	22E ATLS	$t\tilde{\mu}_L$ production, RPV, $\tilde{\mu}_L \rightarrow \mu \tilde{\chi}_1^0$, $\lambda'_{231} = 1$, $m_{\tilde{\chi}_1^0} = 0$ GeV.
>1200	95	2 AAD	21Y ATLS	$\geq 4\ell$, $\lambda_{12k} \neq 0$, $m_{\tilde{\chi}_1^0} = 900$ GeV (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations)
> 870	95	2 AAD	21Y ATLS	$\geq 4\ell$, $\lambda_{i33} \neq 0$, $m_{\tilde{\chi}_1^0} = 450$ GeV (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations)
> 780	95	3 AABOUD	18Z ATLS	$\geq 4\ell$, $\lambda_{i33} \neq 0$, $m_{\tilde{\chi}_1^0} = 300$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)
>1060	95	3 AABOUD	18Z ATLS	$\geq 4\ell$, $\lambda_{12k} \neq 0$, $m_{\tilde{\chi}_1^0} = 600$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)
> 410	95	4 AAD	14X ATLS	RPV, $\geq 4\ell^\pm$, $\tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$, $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		5 SIRUNYAN	19AO	$\mu^\pm \mu^\pm + \geq 2\text{jets}$, $\lambda'_{211} \neq 0$, $\tilde{\mu}_L \rightarrow \mu \tilde{\chi}_1^0$, $\tilde{\chi}_1^0 \rightarrow \mu q \bar{q}$
> 87	95	6 ABDALLAH	04M DLPH	RPV, $\tilde{\mu}_R$, indirect, $\Delta m > 5$ GeV
> 81	95	7 HEISTER	03G ALEP	RPV, $\tilde{\mu}_L$

NODE=S046SMV
NODE=S046SMV;CHECK LIMITS

OCCUR=2

OCCUR=2

- ¹ AAD 22E searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry by measuring the yield asymmetry between events containing $e^- \mu^+$ and those containing $e^+ \mu^-$. This was found in agreement with the standard model prediction of 1. Limits are set on the RPV production of $t \tilde{\mu}_L$ events with $\tilde{\mu}_L \rightarrow \mu \tilde{\chi}_1^0$ for various values of λ'_{231} , see their figures 6 and 7.
- ² AAD 21Y searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with four or more leptons (electrons, muons and tau-leptons). No significant excess above the Standard Model expectations is observed. Limits are set on Tchi1n12-GGM, and RPV models similar to Tchi1n2l, Tglu1A (with $q = u, d, s, c, b$, with equal branching fractions), and $\tilde{\ell}_L/\tilde{\nu} \rightarrow \ell/\nu \tilde{\chi}_1^0$ (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations), all with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$ via λ_{12k} or λ_{i33} (where $i, k \in 1, 2$), see their Figure 11.
- ³ AABOUD 18Z searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more charged leptons (electrons, muons and up to two hadronically decaying taus). No significant deviation from the expected SM background is observed. Limits are set on the Higgsino mass in simplified models of general gauge mediated supersymmetry Tn1n1A/Tn1n1B/Tn1n1C, see their Figure 9. Limits are also set on the wino, slepton, sneutrino and gluino mass in a simplified model of NLSP pair production with R-parity violating decays of the LSP via λ_{12k} or λ_{i33} to charged leptons, see their Figures 7, 8.
- ⁴ AAD 14X searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least four leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the slepton mass in an R-parity violating simplified model where the decay $\tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$, with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, takes place with a branching ratio of 100%, see Fig. 9.
- ⁵ SIRUNYAN 19AO searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two same-sign muons and at last two jets, originating from resonant production of second-generation sleptons ($\tilde{\mu}_L, \tilde{\nu}_\mu$) via the R-parity violating coupling λ'_{211} to quarks. No significant excess above the Standard Model expectations is observed. Upper limits on cross sections are derived in the context of two simplified models, see their Figure 4. The cross section limits are translated into limits on λ'_{211} for a modified CMSSM, see their Figure 5.
- ⁶ ABDALLAH 04M use data from $\sqrt{s} = 192\text{--}208 \text{ GeV}$ to derive limits on sparticle masses under the assumption of RPV with $LL\bar{E}$ or $U\bar{D}\bar{D}$ couplings. The results are valid for $\mu = -200 \text{ GeV}$, $\tan\beta = 1.5$, $\Delta m > 5 \text{ GeV}$ and assuming a BR of 1 for the given decay. The limit quoted is for indirect $U\bar{D}\bar{D}$ decays using the neutralino constraint of 39.5 GeV for $LL\bar{E}$ and of 38.0 GeV for $U\bar{D}\bar{D}$ couplings, also derived in ABDALLAH 04M. For indirect decays via $LL\bar{E}$ the limit improves to 90 GeV if the constraint from the neutralino is used and remains at 87 GeV if it is not used. For indirect decays via $U\bar{D}\bar{D}$ couplings it degrades to 85 GeV when the neutralino constraint is not used. Supersedes the result of ABREU 00U.
- ⁷ HEISTER 03G searches for the production of smuons in the case of RPV prompt decays with $LL\bar{E}$, $LQ\bar{D}$ or $U\bar{D}\bar{D}$ couplings at $\sqrt{s} = 189\text{--}209 \text{ GeV}$. The search is performed for direct and indirect decays, assuming one coupling at a time to be non-zero. The limit holds for direct decays mediated by RPV $LQ\bar{D}$ couplings and improves to 90 GeV for indirect decays (for $\Delta m > 10 \text{ GeV}$). Limits are also given for $LL\bar{E}$ direct ($m_{\tilde{\mu}_R} > 87 \text{ GeV}$) and indirect decays ($m_{\tilde{\mu}_R} > 96 \text{ GeV}$ for $m(\tilde{\chi}_1^0) > 23 \text{ GeV}$ from BARATE 98S) and for $U\bar{D}\bar{D}$ indirect decays ($m_{\tilde{\mu}_R} > 85 \text{ GeV}$ for $\Delta m > 10 \text{ GeV}$). Supersedes the results from BARATE 01B.

NODE=S046SMV;LINKAGE=D

NODE=S046SMV;LINKAGE=C

NODE=S046SMV;LINKAGE=A

NODE=S046SMV;LINKAGE=H

NODE=S046SMV;LINKAGE=B

NODE=S046SMV;LINKAGE=DH

NODE=S046SMV;LINKAGE=HI

R-parity conserving $\tilde{\tau}$ (Stau) mass limit

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

NODE=S046STA

NODE=S046STA

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>400	95	¹ TUMASYAN	23AG CMS	2 hadronic $\tau + \cancel{E}_T, \tilde{\tau}_{R,L} \rightarrow \tau \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$
none 115–340	95	¹ TUMASYAN	23AG CMS	2 hadronic $\tau + \cancel{E}_T, \tilde{\tau}_L \rightarrow \tau \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$
none 120–390	95	² AAD	20H	2 hadronic $\tau + \cancel{E}_T, \tilde{\tau}_{R/L} \rightarrow \tau \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
none 90–150	95	³ SIRUNYAN	20P CMS	2 $\tau + \cancel{E}_T, \tau_h \tau_h$ and $\ell \tau_h, m_{\tilde{\tau}_R} = m_{\tilde{\tau}_L}, m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$
> 85.2		⁴ ABBIENDI	04 OPAL	$\Delta m > 6 \text{ GeV}, \theta_\tau = \pi/2, \mu > 100 \text{ GeV}, \tan\beta = 1.5$
> 78.3		⁵ ACHARD	04 L3	$\Delta m > 15 \text{ GeV}, \theta_\tau = \pi/2, \mu > 200 \text{ GeV}, \tan\beta \geq 2$
> 81.9	95	⁶ ABDALLAH	03M DLPH	$\Delta m > 15 \text{ GeV}$, all θ_τ
> 79	95	⁷ HEISTER	02E ALEP	$\Delta m > 15 \text{ GeV}, \theta_\tau = \pi/2$
> 76	95	⁷ HEISTER	02E ALEP	$\Delta m > 15 \text{ GeV}, \theta_\tau = 0.91$

NODE=S046STA;CHECK LIMITS

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

>500	95	⁸ AABOUD	18BT ATLS	$2\ell + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}, \tilde{\ell} = \tilde{e}, \tilde{\mu}, \tilde{\tau},$ $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
		⁹ KHACHATRYAN...17L	CMS	$2\tau + \cancel{E}_T, \tilde{\tau}_L \rightarrow \tau \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} =$ 0 GeV
none 109	95	¹⁰ AAD	16AA ATLS	$2 \text{ hadronic } \tau + \cancel{E}_T, \tilde{\tau}_{R/L} \rightarrow$ $\tau \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$
		¹¹ AAD	12AF ATLS	$2\tau + \text{jets} + \cancel{E}_T, \text{GMSB}$
		¹² AAD	12AG ATLS	$\geq 1\tau_h + \text{jets} + \cancel{E}_T, \text{GMSB}$
		¹³ AAD	12CMATLS	$\geq 1\tau + \text{jets} + \cancel{E}_T, \text{GMSB}$
> 87.4	95	¹⁴ ABBIENDI	06B OPAL	$\tilde{\tau}_R \rightarrow \tau \tilde{G}, \text{all } \tau(\tilde{\tau}_R)$
> 68	95	¹⁵ ABDALLAH	04H DLPH	AMSB, $\mu > 0$
none $m_{\tilde{\tau}} - 26.3$	95	⁶ ABDALLAH	03M DLPH	$\Delta m > m_{\tilde{\tau}}, \text{all } \theta_{\tilde{\tau}}$

OCCUR=2

NODE=S046STA;LINKAGE=K

¹ TUMASYAN 23AG searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for or direct pair production of tau sleptons in events with two hadronically decaying tau leptons. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the tau slepton in models with $\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0$ for mass-degenerate, pure left-handed and pure right-handed tau sleptons, see their figures 4–7. Limits are also set for the maximally mixed scenario with long-lived tau sleptons and $\tilde{\tau}$ lifetimes of 0.01 mm to 2.5 mm, see their figure 8.

NODE=S046STA;LINKAGE=J

² AAD 20H presented ATLAS searches for direct production for $\tilde{\tau}$ in final states with two hadronically decaying leptons and \cancel{E}_T . The analysis uses a dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 139 fb^{-1} . Exclusion limits at 95% C.L. are derived in scenarios of direct production of $\tilde{\tau}$ pairs with each $\tilde{\tau}$ decaying into a τ and the lightest neutralino $\tilde{\chi}_1^0$ in simplified models where the $\tilde{\tau}_R$ and $\tilde{\tau}_L$ mass eigenstates are degenerate. Stau masses from 120 GeV to 390 GeV are excluded for a massless lightest neutralino, see their Fig. 7(a). If $\tilde{\tau}_L$ -only pair production is considered, the exclusion region extends between 155 GeV to 310 GeV, see their Fig. 7(b).

NODE=S046STA;LINKAGE=I

³ SIRUNYAN 20P searched in 77.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct pair production of tau sleptons in events with a tau lepton pair and significant missing transverse momentum. Final states with two double hadronic decay of the tau leptons are considered, as well as where one of the tau leptons decays into an electron or a muon. No significant excess above the Standard Model expectations is observed. Limits are set on the stau mass in a simplified models where two tau sleptons are pair produced and decay to a tau lepton and the lightest neutralino, assuming either only left-handed stau production, see Figure 8, or assuming degenerate left- and right-handed stau production, see Figure 9.

NODE=S046STA;LINKAGE=AB

⁴ ABBIENDI 04 search for $\tilde{\tau}\tilde{\tau}$ production in acoplanar di-tau final states in the 183–208 GeV data. See Fig. 15 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$ and for the limit at $\tan\beta=35$. Under the assumption of 100% branching ratio for $\tilde{\tau}_R \rightarrow \tau \tilde{\chi}_1^0$, the limit improves to 89.8 GeV for $\Delta m > 8 \text{ GeV}$. See Fig. 12 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$ at several values of the branching ratio and for their dependence on $\theta_{\tilde{\tau}}$. This limit supersedes ABBIENDI 00G.

NODE=S046STA;LINKAGE=AC

⁵ ACHARD 04 search for $\tilde{\tau}\tilde{\tau}$ production in acoplanar di-tau final states in the 192–209 GeV data. Limits on $m_{\tilde{\tau}_R}$ are derived from a scan over the MSSM parameter space with universal GUT scale gaugino and scalar masses $m_{1/2}$ and m_0 , $1 \leq \tan\beta \leq 60$ and $-2 \leq \mu \leq 2 \text{ TeV}$. See Fig. 4 for the dependence of the limits on $m_{\tilde{\chi}_1^0}$.

NODE=S046STA;LINKAGE=AA

⁶ ABDALLAH 03M looked for acoplanar ditau + \cancel{E} final states at $\sqrt{s} = 130\text{--}208 \text{ GeV}$. A dedicated search was made for low mass $\tilde{\tau}$ s decoupling from the Z^0 . The limit assumes $B(\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0) = 100\%$. See Fig. 20 for limits on the $(m_{\tilde{\tau}}, m_{\tilde{\chi}_1^0})$ plane and as function of the $\tilde{\chi}_1^0$ mass and of the branching ratio. The limit in the low-mass region improves to 29.6 and 31.1 GeV for $\tilde{\tau}_R$ and $\tilde{\tau}_L$, respectively, at $\Delta m > m_{\tilde{\tau}}$. The limit in the high-mass region improves to 84.7 GeV for $\tilde{\tau}_R$ and $\Delta m > 15 \text{ GeV}$. These limits include and update the results of ABREU 01.

NODE=S046STA;LINKAGE=HE

⁷ HEISTER 02E looked for acoplanar ditau + \cancel{E}_T final states from e^+e^- interactions between 183 and 209 GeV. The mass limit assumes $B(\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0) = 1$. See their Fig. 4 for the dependence of the limit on Δm . These limits include and update the results of BARATE 01.

NODE=S046STA;LINKAGE=H

⁸ AABOUD 18BT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos, chargino and next-to-lightest neutralinos and sleptons in events with two or three leptons (electrons or muons), with or without jets, and large missing transverse energy. No significant excess above the Standard Model expectations is observed. Limits are set on the slepton mass up to 500 GeV for massless $\tilde{\chi}_1^0$, assuming degeneracy of $\tilde{e}, \tilde{\mu},$ and $\tilde{\tau}$ and exploiting the 2ℓ signature, see their Figure 8(b).

NODE=S046STA;LINKAGE=G

⁹ KHACHATRYAN 17L searched in about 19 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with two τ (at least one decaying hadronically) and \cancel{E}_T . Results were interpreted

to set constraints on the cross section for production of $\tilde{\tau}_L$ pairs for $m_{\tilde{\chi}_1^0}=1$ GeV. No mass constraints are set, see their Fig. 7.

- 10 AAD 16AA summarized and extended ATLAS searches for electroweak supersymmetry in final states containing several charged leptons, \cancel{E}_T , with or without hadronic jets, in 20 fb⁻¹ of pp collisions at $\sqrt{s} = 8$ TeV. The paper reports 95% C.L. exclusion limits on the cross-section for production of $\tilde{\tau}_R$ and $\tilde{\tau}_L$ pairs for various $m_{\tilde{\chi}_1^0}$, using the 2 hadronic $\tau + \cancel{E}_T$ analysis. The $m_{\tilde{\tau}_{R/L}} = 109$ GeV is excluded for $m_{\tilde{\chi}_1^0} = 0$ GeV, with the constraints being stronger for $\tilde{\tau}_R$. See their Fig. 12.
- 11 AAD 12AF searched in 2 fb⁻¹ of pp collisions at $\sqrt{s} = 7$ TeV for events with two tau leptons, jets and large \cancel{E}_T in a GMSB framework. No significant excess above the expected background was found and an upper limit on the visible cross section for new phenomena is set. A 95% C.L. lower limit of 32 TeV on the mGMSB breaking scale Λ is set for $M_{mess} = 250$ TeV, $N_S = 3$, $\mu > 0$ and $C_{grav} = 1$, independent of $\tan\beta$.
- 12 AAD 12AG searched in 2.05 fb⁻¹ of pp collisions at $\sqrt{s} = 7$ TeV for events with at least one hadronically decaying tau lepton, jets, and large \cancel{E}_T in a GMSB framework. No significant excess above the expected background was found and an upper limit on the visible cross section for new phenomena is set. A 95% C.L. lower limit of 30 TeV on the mGMSB breaking scale Λ is set for $M_{mess} = 250$ TeV, $N_S = 3$, $\mu > 0$ and $C_{grav} = 1$, independent of $\tan\beta$. For large values of $\tan\beta$, the limit on Λ increases to 43 TeV.
- 13 AAD 12CM searched in 4.7 fb⁻¹ of pp collisions at $\sqrt{s}=7$ TeV for events with at least one tau lepton, zero or one additional light lepton (e/μ) jets, and large \cancel{E}_T in a GMSB framework. No significant excess above the expected background was found and an upper limit on the visible cross section for new phenomena is set. A 95% C. L. lower limit of 54 TeV on the mGMSB breaking scale Λ is set for $M_{mess} = 250$ TeV, $N_S = 3$, $\mu > 0$ and $C_{grav} = 1$, for $\tan\beta > 20$. Here the $\tilde{\tau}_1$ is the NLSP.
- 14 ABBIENDI 06B use 600 pb⁻¹ of data from $\sqrt{s} = 189\text{--}209$ GeV. They look for events from pair-produced staus in a GMSB scenario with $\tilde{\tau}$ NLSP including prompt $\tilde{\tau}$ decays to ditaus + \cancel{E} final states, large impact parameters, kinked tracks and heavy stable charged particles. Limits on the cross-section are computed as a function of $m(\tilde{\tau})$ and the lifetime, see their Fig. 7. The limit is compared to the $\sigma \cdot BR^2$ from a scan over the GMSB parameter space.
- 15 ABDALLAH 04H use data from LEP 1 and $\sqrt{s} = 192\text{--}208$ GeV. They re-use results or re-analyze the data from ABDALLAH 03M to put limits on the parameter space of anomaly-mediated supersymmetry breaking (AMSB), which is scanned in the region $1 < m_{3/2} < 50$ TeV, $0 < m_0 < 1000$ GeV, $1.5 < \tan\beta < 35$, both signs of μ . The constraints are obtained from the searches for mass degenerate chargino and neutralino, for SM-like and invisible Higgs, for leptonically decaying charginos and from the limit on non-SM Z width of 3.2 MeV. The limit is for $m_t = 174.3$ GeV (see Table 2 for other m_t values). The limit improves to 75 GeV for $\mu < 0$.

NODE=S046STA;LINKAGE=FF

NODE=S046STA;LINKAGE=LS

NODE=S046STA;LINKAGE=GD

NODE=S046STA;LINKAGE=DA

NODE=S046STA;LINKAGE=BI

NODE=S046STA;LINKAGE=BD

R-parity violating $\tilde{\tau}$ (Stau) mass limit

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

NODE=S046STU

NODE=S046STU

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1200	95	¹ AAD	21Y ATLS	$\geq 4\ell, \lambda_{12k} \neq 0, m_{\tilde{\chi}_1^0} = 900$ GeV (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations)
> 870	95	¹ AAD	21Y ATLS	$\geq 4\ell, \lambda_{i33} \neq 0, m_{\tilde{\chi}_1^0} = 450$ GeV (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations)
>1060	95	² AABOUD	18Z ATLS	$\geq 4\ell, \lambda_{12k} \neq 0, m_{\tilde{\chi}_1^0} = 600$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)
> 780	95	² AABOUD	18Z ATLS	$\geq 4\ell, \lambda_{i33} \neq 0, m_{\tilde{\chi}_1^0} = 300$ GeV (mass-degenerate left-handed sleptons and sneutrinos of all 3 generations)
> 90	95	³ ABDALLAH	04M DLPH	$\tilde{\tau}_R$, indirect, $\Delta m > 5$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
> 74	95	⁴ ABBIENDI	04F OPAL	$\tilde{\tau}_L$

NODE=S046STU;CHECK LIMITS

OCCUR=2

OCCUR=2

NODE=S046STU;LINKAGE=B

- ¹ AAD 21Y searched in 139 fb⁻¹ of pp collisions at $\sqrt{s} = 13$ TeV for supersymmetry in events with four or more leptons (electrons, muons and tau-leptons). No significant excess above the Standard Model expectations is observed. Limits are set on Tchi1n12-GGM, and RPV models similar to Tchi1n2l, Tglu1A (with $q = u, d, s, c, b$, with equal branching fractions), and $\tilde{\ell}_L/\tilde{\nu} \rightarrow \ell/\nu\tilde{\chi}_1^0$ (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3

generations), all with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$ via λ_{12k} or λ_{i33} (where $i, k \in 1, 2$), see their Figure 11.

² AABOUD 18Z searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more charged leptons (electrons, muons and up to two hadronically decaying taus). No significant deviation from the expected SM background is observed. Limits are set on the Higgsino mass in simplified models of general gauge mediated supersymmetry Tn1n1A/Tn1n1B/Tn1n1C, see their Figure 9. Limits are also set on the wino, slepton, sneutrino and gluino mass in a simplified model of NLSP pair production with R-parity violating decays of the LSP via λ_{12k} or λ_{i33} to charged leptons, see their Figures 7, 8.

³ ABDALLAH 04M use data from $\sqrt{s} = 192\text{--}208 \text{ GeV}$ to derive limits on sparticle masses under the assumption of RPV with $LL\bar{E}$ couplings. The results are valid for $\mu = -200 \text{ GeV}$, $\tan\beta = 1.5$, $\Delta m > 5 \text{ GeV}$ and assuming a BR of 1 for the given decay. The limit quoted is for indirect decays using the neutralino constraint of 39.5 GeV, also derived in ABDALLAH 04M. For indirect decays via $LL\bar{E}$ the limit decreases to 86 GeV if the constraint from the neutralino is not used. Supersedes the result of ABREU 00U.

⁴ ABBIENDI 04F use data from $\sqrt{s} = 189\text{--}209 \text{ GeV}$. They derive limits on sparticle masses under the assumption of RPV with $LL\bar{E}$ or $LQ\bar{D}$ couplings. The results are valid for $\tan\beta = 1.5$, $\mu = -200 \text{ GeV}$, with, in addition, $\Delta m > 5 \text{ GeV}$ for indirect decays via $LQ\bar{D}$. The limit quoted applies to direct decays with $LL\bar{E}$ couplings and improves to 75 GeV for $LQ\bar{D}$ couplings. The limit on the $\tilde{\tau}_R$ mass for indirect decays is 92 GeV for $LL\bar{E}$ couplings at $m_{\tilde{\chi}_1^0} = 10 \text{ GeV}$ and no exclusion is obtained for $LQ\bar{D}$ couplings. Supersedes the results of ABBIENDI 00.

NODE=S046STU;LINKAGE=A

NODE=S046STU;LINKAGE=DH

NODE=S046STU;LINKAGE=IA

Long-lived $\tilde{\ell}$ (Slepton) mass limit

Limits on scalar leptons which leave detector before decaying. Limits from Z decays are independent of lepton flavor. Limits from continuum e^+e^- annihilation are also independent of flavor for smuons and staus. Selectron limits from e^+e^- collisions in the continuum depend on MSSM parameters because of the additional neutralino exchange contribution.

NODE=S046SLP

NODE=S046SLP

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>520	95	¹ AAD	23BQ ATLS	2ℓ slightly displaced, long-lived $\tilde{\mu}, \tilde{\mu} \rightarrow \mu \tilde{G}$, $m_{\tilde{\mu}_R} = m_{\tilde{\mu}_L}$, $\tau_{\tilde{\mu}} = 10 \text{ ps}$
>190	95	¹ AAD	23BQ ATLS	2ℓ slightly displaced, long-lived $\tilde{\mu}, \tilde{\mu} \rightarrow \mu \tilde{G}$, $m_{\tilde{\mu}_R} = m_{\tilde{\mu}_L}$, $\tau_{\tilde{\mu}} = 1 \text{ ps}$
none 220–360	95	² AAD	23G ATLS	direct $\tilde{\tau}$ pair, $\tilde{\tau} \rightarrow \tau \tilde{G}$, $\tau = 10 \text{ ns}$
none 150–220	95	³ TUMASYAN	23AG CMS	2 hadronic $\tau + \cancel{E}_T$, $\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0$, maximally mixed scenario with $c\tau = 0.1 \text{ mm}$, $m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$
>610	95	⁴ TUMASYAN	22AF CMS	2ℓ displaced, long-lived $\tilde{e}, \tilde{e} \rightarrow e \tilde{G}$, $m_{\tilde{e}_R} = m_{\tilde{e}_L}$, $c\tau = 0.7 \text{ cm}$
>610	95	⁴ TUMASYAN	22AF CMS	2ℓ displaced, long-lived $\tilde{\mu}, \tilde{\mu} \rightarrow \mu \tilde{G}$, $m_{\tilde{\mu}_R} = m_{\tilde{\mu}_L}$, $c\tau = 3 \text{ cm}$
>405	95	⁴ TUMASYAN	22AF CMS	2ℓ displaced, long-lived $\tilde{\tau}, \tilde{\tau} \rightarrow \tau \tilde{G}$, $m_{\tilde{\tau}_R} = m_{\tilde{\tau}_L}$, $c\tau = 2 \text{ cm}$
>270	95	⁴ TUMASYAN	22AF CMS	2ℓ displaced, long-lived $\tilde{\ell}, \tilde{\ell} \rightarrow \ell \tilde{G}$, $m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$, $m_{\tilde{e}} = m_{\tilde{\mu}} = m_{\tilde{\tau}}$, $0.005 \text{ cm} < c\tau < 265 \text{ cm}$
>680	95	⁴ TUMASYAN	22AF CMS	2ℓ displaced, long-lived $\tilde{\ell}, \tilde{\ell} \rightarrow \ell \tilde{G}$, $m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$, $m_{\tilde{e}} = m_{\tilde{\mu}} = m_{\tilde{\tau}}$, $c\tau = 2 \text{ cm}$
>720	95	⁵ AAD	21AL ATLS	2ℓ displaced, long-lived $\tilde{e}, \tilde{e} \rightarrow e \tilde{G}$, $m_{\tilde{e}_R} = m_{\tilde{e}_L}$, $\tau_{\tilde{e}} = 0.1 \text{ ns}$
>680	95	⁵ AAD	21AL ATLS	2ℓ displaced, long-lived $\tilde{\mu}, \tilde{\mu} \rightarrow \mu \tilde{G}$, $m_{\tilde{\mu}_R} = m_{\tilde{\mu}_L}$, $\tau_{\tilde{\mu}} = 0.1 \text{ ns}$
>340	95	⁵ AAD	21AL ATLS	2ℓ displaced, long-lived $\tilde{\tau}, \tilde{\tau} \rightarrow \tau \tilde{G}$, mixing $\sin\theta_{\tilde{\tau}} = 0.95$, $\tau_{\tilde{\tau}} = 0.1 \text{ ns}$
>820	95	⁵ AAD	21AL ATLS	2ℓ displaced, long-lived $\tilde{\ell}, \tilde{\ell} \rightarrow \ell \tilde{G}$, $m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L}$, $m_{\tilde{e}} = m_{\tilde{\mu}} = m_{\tilde{\tau}}$, $\tau_{\tilde{\ell}} = 0.1 \text{ ns}$
>430	95	⁶ AABOUD	19AT ATLS	long-lived $\tilde{\tau}$, GMSB
>490	95	⁷ KHACHATRY...	16BWCMS	long-lived $\tilde{\tau}$ from inclusive production, mGMSB SPS line 7 scenario

NODE=S046SLP;CHECK LIMITS

OCCUR=2

OCCUR=2

OCCUR=4

OCCUR=6

OCCUR=7

OCCUR=8

OCCUR=2

OCCUR=3

OCCUR=4

>240	95	⁷ KHACHATRY...16BWCMS		long-lived $\tilde{\tau}$ from direct pair production, mGMSB SPS line 7 scenario	OCCUR=2
>440	95	⁸ AAD	15AE ATLS	mGMSB, $M_{m\tilde{e}s s} = 250$ TeV, $N_5 = 3$, $\mu > 0$, $C_{grav} = 5000$, $\tan\beta = 10$	
>385	95	⁸ AAD	15AE ATLS	mGMSB, $M_{m\tilde{e}s s} = 250$ TeV, $N_5 = 3$, $\mu > 0$, $C_{grav} = 5000$, $\tan\beta = 50$	OCCUR=2
>286	95	⁸ AAD	15AE ATLS	direct $\tilde{\tau}$ production	OCCUR=3
none 124–309	95	⁹ AAIJ	15BD LHCB	long-lived $\tilde{\tau}$, mGMSB, SPS7	
> 98	95	¹⁰ ABBIENDI	03L OPAL	$\tilde{\mu}_R, \tilde{\tau}_R$	
none 2–87.5	95	¹¹ ABREU	00Q DLPH	$\tilde{\mu}_R, \tilde{\tau}_R$	
> 81.2	95	¹² ACCIARRI	99H L3	$\tilde{\mu}_R, \tilde{\tau}_R$	
> 81	95	¹³ BARATE	98K ALEP	$\tilde{\mu}_R, \tilde{\tau}_R$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>300	95	¹⁴ AAD	13AA ATLS	long-lived $\tilde{\tau}$, GMSB, $\tan\beta = 5-20$	
		¹⁵ ABAZOV	13B D0	long-lived $\tilde{\tau}$, $100 < m_{\tilde{\tau}} < 300$ GeV	
>339	95	^{16,17} CHATRCHYAN	13AB CMS	long-lived $\tilde{\tau}$, direct $\tilde{\tau}_1$ pair prod., minimal GMSB, SPS line 7	
>500	95	^{16,18} CHATRCHYAN	13AB CMS	long-lived $\tilde{\tau}$, $\tilde{\tau}_1$ from direct pair prod. and from decay of heavier SUSY particles, minimal GMSB, SPS line 7	OCCUR=2
>314	95	¹⁹ CHATRCHYAN	12L CMS	long-lived $\tilde{\tau}$, $\tilde{\tau}_1$ from decay of heavier SUSY particles, minimal GMSB, SPS line 7	
>136	95	²⁰ AAD	11P ATLS	stable $\tilde{\tau}$, GMSB scenario, $\tan\beta=5$	
<p>¹ AAD 23BQ searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pair production of long-lived $\tilde{\mu}$ in events with muons with impact parameters in the millimeter range. No significant excess above the Standard Model predictions is observed. Limits are set on $m_{\tilde{\mu}}$ as a function of the $\tilde{\mu}$ lifetime, assuming the $\tilde{\mu} \rightarrow \mu \tilde{G}$ decay and mass-degenerate $\tilde{\mu}_L$ and $\tilde{\mu}_R$. See Figure 4.</p>					
<p>² AAD 23G searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for stau pair production in events with high-pt tracks with large ionisation in the pixel detector. No significant excess above the Standard Model predictions is observed. Limits are set on the stau mass as a function of its lifetime, see Figure 19.</p>					
<p>³ TUMASYAN 23AG searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for or direct pair production of tau sleptons in events with two hadronically decaying tau leptons. No significant excess above the Standard Model expectations is observed. Limits are set for the maximally mixed scenario with long-lived tau sleptons and $\tilde{\tau}$ lifetimes of 0.01 mm to 2.5 mm, see their figure 8. Limits are also set on the mass of the tau slepton in models with $\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0$ for mass-degenerate, pure left-handed and pure right-handed tau sleptons, see their figures 4–7.</p>					
<p>⁴ TUMASYAN 22AF searched for evidence of new long-lived particles decaying to leptons in pp collisions at $\sqrt{s} = 13$ TeV, corresponding to 118 (113) fb^{-1} in the $e\mu$ and $\mu\mu$ channels. The leptons are required to have transverse impact parameter values between 0.01 and 10 cm and are not required to form a common vertex. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the top squark in RPV models with top squark pair production and $\tilde{t} \rightarrow b\tilde{\ell}$ and $\tilde{t} \rightarrow d\tilde{\ell}$, see their Figure 4, which contains a wider range of lifetime limits. Limits are also set on a gauge-mediated SUSY breaking model, where the next-to-lightest SUSY particle is a slepton and the lightest SUSY particle a gravitino \tilde{G}, see their Figure 5, which also contains a wider range of lifetime limits. Limits are also set in a model that produces BSM Higgs bosons (H) with a mass of 125 GeV through gluon-gluon fusion, where the H decays to two long-lived scalars S, each of which decays to two oppositely charged and same-flavor leptons.</p>					
<p>⁵ AAD 21AL searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pair production of long-lived sleptons in events with highly displaced leptons. No significant excess above the Standard Model predictions is observed. Limits are set on $m_{\tilde{e}}$, $m_{\tilde{\mu}}$, $m_{\tilde{\tau}}$ as a function of the slepton lifetime, assuming the $\tilde{\ell} \rightarrow \ell \tilde{G}$ decay and mass-degenerate $\tilde{\ell}_L$ and $\tilde{\ell}_R$. See Figures 2.</p>					
<p>⁶ AABOUD 19AT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for metastable and stable R-hadrons. Multiple search strategies for a wide range of lifetimes, corresponding to path lengths of a few meters, are defined. No significant deviations from the expected Standard Model background are observed. Results are interpreted in terms of exclusion limits on long-lived stau in the context of GMSB models. Lower limits on the mass for direct production of staus are set at 430 GeV, see their Fig. 10 (left).</p>					
<p>⁷ KHACHATRYAN 16BW searched in 2.5 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with heavy stable charged particles, identified by their anomalously high energy deposits in the silicon tracker and/or long time-of-flight measurements by the muon system. No evidence for an excess over the expected background is observed. Limits are derived for pair production of tau sleptons as a function of mass, depending on their direct or</p>					

NODE=S046SLP;LINKAGE=L

NODE=S046SLP;LINKAGE=K

NODE=S046SLP;LINKAGE=J

NODE=S046SLP;LINKAGE=I

NODE=S046SLP;LINKAGE=H

NODE=S046SLP;LINKAGE=G

NODE=S046SLP;LINKAGE=E

- inclusive production in a minimal GMSB scenario along the Snowmass Points and Slopes (SPS) line 7, see Fig. 4 and Table 7.
- 8 AAD 15AE searched in 19.1 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for heavy long-lived charged particles, measured through their specific ionization energy loss in the ATLAS pixel detector or their time-of-flight in the ALTAS muon system. In the absence of an excess of events above the expected backgrounds, limits are set on stable $\tilde{\tau}$ sleptons in various scenarios, see Figs. 5-7. NODE=S046SLP;LINKAGE=D
- 9 AAJ 15BD searched in 3.0 fb^{-1} of pp collisions at $\sqrt{s} = 7$ and 8 TeV for evidence of Drell-Yan pair production of long-lived $\tilde{\tau}$ particles. No evidence for such particles is observed and 95% C.L. upper limits on the cross section of $\tilde{\tau}$ pair production are derived, see Fig. 7. In the mGMSB, assuming the SPS7 benchmark scenario $\tilde{\tau}$ masses between 124 and 309 GeV are excluded at 95% C.L. NODE=S046SLP;LINKAGE=F
- 10 ABBIENDI 03L used e^+e^- data at $\sqrt{s} = 130\text{--}209 \text{ GeV}$ to select events with two high momentum tracks with anomalous dE/dx . The excluded cross section is compared to the theoretical expectation as a function of the heavy particle mass in their Fig. 3. The limit improves to 98.5 GeV for $\tilde{\mu}_L$ and $\tilde{\tau}_L$. The bounds are valid for colorless spin 0 particles with lifetimes longer than 10^{-6} s . Supersedes the results from ACKERSTAFF 98P. NODE=S046SLP;LINKAGE=AB
- 11 ABREU 00Q searches for the production of pairs of heavy, charged stable particles in e^+e^- annihilation at $\sqrt{s} = 130\text{--}189 \text{ GeV}$. The upper bound improves to 88 GeV for $\tilde{\mu}_L$, $\tilde{\tau}_L$. These limits include and update the results of ABREU 98P. NODE=S046SLP;LINKAGE=QT
- 12 ACCIARRI 99H searched for production of pairs of back-to-back heavy charged particles at $\sqrt{s} = 130\text{--}183 \text{ GeV}$. The upper bound improves to 82.2 GeV for $\tilde{\mu}_L$, $\tilde{\tau}_L$. NODE=S046SLP;LINKAGE=HA
- 13 The BARATE 98K mass limit improves to 82 GeV for $\tilde{\mu}_L, \tilde{\tau}_L$. Data collected at $\sqrt{s} = 161\text{--}184 \text{ GeV}$. NODE=S046SLP;LINKAGE=TC
- 14 AAD 13AA searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing long-lived massive particles in a GMSB framework. No significant excess above the expected background was found. A 95% C.L. lower limit of 300 GeV is placed on long-lived $\tilde{\tau}$'s in the GMSB model with $M_{mess} = 250 \text{ TeV}$, $N_\zeta = 3$, $\mu > 0$, for $\tan\beta = 5\text{--}20$. The lower limit on the GMSB breaking scale Λ was found to be 99–110 TeV, for $\tan\beta$ values between 5 and 40, see Fig. 4 (top). Also, directly produced long-lived sleptons, or sleptons decaying to long-lived ones, are excluded at 95% C.L. up to a $\tilde{\tau}$ mass of 278 GeV for models with slepton splittings smaller than 50 GeV. NODE=S046SLP;LINKAGE=AA
- 15 ABAZOV 13B looked in 6.3 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for charged massive long-lived particles in events with muon-like particles that have both speed and ionization energy loss inconsistent with muons produced in beam collisions. In the absence of an excess, limits are set at 95% C.L. on the production cross section of stau leptons in the mass range 100–300 GeV, see their Table 20 and Fig. 23. NODE=S046SLP;LINKAGE=AV
- 16 CHATRCHYAN 13AB looked in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and in 18.8 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or additionally requiring that it be identified as muon in the muon chambers, from pair production of $\tilde{\tau}_1$'s. No evidence for an excess over the expected background is observed. Supersedes CHATRCHYAN 12L. NODE=S046SLP;LINKAGE=CA
- 17 CHATRCHYAN 13AB limits are derived for pair production of $\tilde{\tau}_1$ as a function of mass in minimal GMSB scenarios along the Snowmass Points and Slopes (SPS) line 7 (see Fig. 8 and Table 7). The limit given here is valid for direct pair $\tilde{\tau}_1$ production. NODE=S046SLP;LINKAGE=CT
- 18 CHATRCHYAN 13AB limits are derived for the production of $\tilde{\tau}_1$ as a function of mass in minimal GMSB scenarios along the Snowmass Points and Slopes (SPS) line 7 (see Fig. 8 and Table 7). The limit given here is valid for the production of $\tilde{\tau}_1$ from both direct pair production and from the decay of heavier supersymmetric particles. NODE=S046SLP;LINKAGE=CR
- 19 CHATRCHYAN 12L looked in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or additionally requiring that it be identified as muon in the muon chambers, from pair production of $\tilde{\tau}_1$'s. No evidence for an excess over the expected background is observed. Limits are derived for the production of $\tilde{\tau}_1$ as a function of mass in minimal GMSB scenarios along the Snowmass Points and Slopes (SPS) line 7 (see Fig. 3). The limit given here is valid for the production of $\tilde{\tau}_1$ in the decay of heavier supersymmetric particles. NODE=S046SLP;LINKAGE=CH
- 20 AAD 11P looked in 37 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with two heavy stable particles, reconstructed in the Inner tracker and the Muon System and identified by their time of flight in the Muon System. No evidence for an excess over the SM expectation is observed. Limits on the mass are derived, see Fig. 3, for $\tilde{\tau}$ in a GMSB scenario and for sleptons produced by electroweak processes only, in which case the limit degrades to 110 GeV. NODE=S046SLP;LINKAGE=A1

\tilde{q} (Squark) mass limit

NODE=S046225

For $m_{\tilde{q}} > 60\text{--}70 \text{ GeV}$, it is expected that squarks would undergo a cascade decay via a number of neutralinos and/or charginos rather than undergo a direct decay to photinos as assumed by some papers. Limits obtained when direct decay is assumed are usually higher than limits when cascade decays are included.

NODE=S046225

Limits from e^+e^- collisions depend on the mixing angle of the lightest mass eigenstate $\tilde{q}_1 = \tilde{q}_R \sin\theta_q + \tilde{q}_L \cos\theta_q$. It is usually assumed that only the sbottom and stop squarks have non-trivial mixing angles (see the stop and sbottom sections). Here, unless otherwise noted, squarks are always

taken to be either left/right degenerate, or purely of left or right type. Data from Z decays have set squark mass limits above 40 GeV, in the case of $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ decays if $\Delta m = m_{\tilde{q}} - m_{\tilde{\chi}_1^0} \gtrsim 5$ GeV. For smaller values of Δm , current constraints on the invisible width of the Z ($\Delta\Gamma_{\text{inv}} < 2.0$ MeV, LEP 00) exclude $m_{\tilde{u}_{L,R}} < 44$ GeV, $m_{\tilde{d}_R} < 33$ GeV, $m_{\tilde{d}_L} < 44$ GeV and, assuming all squarks degenerate, $m_{\tilde{q}} < 45$ GeV.

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

R-parity conserving \tilde{q} (Squark) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1550	95	1 AAD	23AE ATLS	2 SFOS ℓ , jets, \cancel{E}_T , Tsqk2, $m_{\tilde{\chi}_2^0} = (m_{\tilde{q}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 100$ GeV
none 1200–2500	95	2 TUMASYAN	23X CMS	2 AK8 jets + 1 AK4 jet, $\tilde{q} \rightarrow q\tilde{\chi}_2^0$ and $\tilde{\chi}_2^0 \rightarrow H_1\tilde{\chi}_S^0$, $40 < m_{H_1} < 120$ GeV
>1400	95	3 AAD	21AK ATLS	ℓ^\pm + jets + \cancel{E}_T , Tsqk3, 4 degenerate light \tilde{q}_ℓ , $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{q}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 200$ GeV
>1040	95	3 AAD	21AK ATLS	ℓ^\pm + jets + \cancel{E}_T , Tsqk3, 1 light \tilde{q}_ℓ , $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{q}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 200$ GeV
> 925	95	4 AAD	21F ATLS	≥ 1 jet + \cancel{E}_T , Tsqk1, $m_{\tilde{q}} - m_{\tilde{\chi}_1^0} = 5$ GeV
> 550	95	4 AAD	21F ATLS	≥ 1 jet + \cancel{E}_T , Tstop3, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 5$ GeV
> 550	95	4 AAD	21F ATLS	≥ 1 jet + \cancel{E}_T , Tstop4, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 5$ GeV
> 545	95	4 AAD	21F ATLS	≥ 1 jet + \cancel{E}_T , Tsb0t1, $m_{\tilde{b}} - m_{\tilde{\chi}_1^0} = 5$ GeV
>1850	95	5 AAD	21L ATLS	jets + \cancel{E}_T , Tsqk1, 8 degenerate \tilde{q} , $m_{\tilde{\chi}_1^0} = 0$ GeV
>1220	95	5 AAD	21L ATLS	jets + \cancel{E}_T , Tsqk1, 1 non-degenerate \tilde{q} , $m_{\tilde{\chi}_1^0} = 0$ GeV
>1310	95	5 AAD	21L ATLS	jets + \cancel{E}_T , Tsqk3, 4 degenerate \tilde{q}_l , $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{q}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$ GeV
>3000	95	5 AAD	21L ATLS	jets + \cancel{E}_T , combined $\tilde{g}\tilde{g}$, $\tilde{g}\tilde{q}$, $\tilde{q}\tilde{q}$ production, $\tilde{g} \rightarrow qq'\tilde{\chi}_1^0$, $\tilde{q} \rightarrow q\tilde{\chi}_1^0$, $m_{\tilde{q}} = m_{\tilde{g}}$, $m_{\tilde{\chi}_1^0} = 0$ GeV
>1800	95	6 SIRUNYAN	21M CMS	$\ell^\pm\ell^\mp$ + \cancel{E}_T , Tsqk2A, $m_{\tilde{\chi}_2^0} = 1500$ GeV, $m_{\tilde{\chi}_1^0} = 100$ GeV
>1590	95	7 SIRUNYAN	19AG CMS	2γ + \cancel{E}_T , Tsqk4B, 500 GeV $< m_{\tilde{\chi}_1^0} < 1500$ GeV
>1130	95	8 SIRUNYAN	19CH CMS	jets + \cancel{E}_T , Tsqk1, 1 light flavour, $m_{\tilde{\chi}_1^0} = 0$ GeV
>1630	95	8 SIRUNYAN	19CH CMS	jets + \cancel{E}_T , Tsqk1, 8 degenerate light flavours, $m_{\tilde{\chi}_1^0} = 0$ GeV
>1430	95	9 SIRUNYAN	19K CMS	γ + ℓ + \cancel{E}_T , Tsqk4A, $m_{\tilde{\chi}_1^0} = 1200$ GeV
>1200	95	10 AABOUD	18BJ ATLS	$\ell^\pm\ell^\mp$ + jets + \cancel{E}_T , Tsqk2, $m_{\tilde{\chi}_1^0} = 1$ GeV, any $m_{\tilde{\chi}_2^0}$
> 850	95	11 AABOUD	18BV ATLS	c-jets + \cancel{E}_T , Tsqk1 (charm only), $m_{\tilde{\chi}_1^0} = 0$ GeV

NODE=S046SQK
NODE=S046SQK;CHECK LIMITS

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

> 710	95	12 AABOUD	18I ATLS	$\geq 1 \text{ jets} + \cancel{E}_T, \text{ Tsqk1}, m_{\tilde{q}} \sim m_{\tilde{\chi}_1^0}$	
>1820	95	13 AABOUD	18U ATLS	$2 \gamma + \cancel{E}_T, \text{ GGM}, \text{ Tsqk4B}, \text{ any NLSP mass}$	
>1550	95	14 AABOUD	18V ATLS	$\text{jets} + \cancel{E}_T, \text{ Tsqk1}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1150	95	15 AABOUD	18V ATLS	$\text{jets} + \cancel{E}_T, \text{ Tsqk3}, m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{q}} + m_{\tilde{\chi}_1^0}), m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1650	95	16 SIRUNYAN	18AA CMS	$\geq 1 \gamma + \cancel{E}_T, \text{ Tsqk4A}$	
>1750	95	16 SIRUNYAN	18AA CMS	$\geq 1 \gamma + \cancel{E}_T, \text{ Tsqk4B}$	OCCUR=2
> 675	95	17 SIRUNYAN	18AY CMS	$\text{jets} + \cancel{E}_T, \text{ Tsqk1}, 1 \text{ light flavor state}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1320	95	17 SIRUNYAN	18AY CMS	$\text{jets} + \cancel{E}_T, \text{ Tsqk1}, 8 \text{ degenerate light flavor states}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1220	95	18 AABOUD	17AR ATLS	$1 \ell + \text{jets} + \cancel{E}_T, \text{ Tsqk3}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1000	95	19 AABOUD	17N ATLS	$2 \text{ same-flavour, opposite-sign } \ell + \text{jets} + \cancel{E}_T, \text{ Tsqk2}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1150	95	20 KHACHATRY...17P	CMS	$1 \text{ or more jets} + \cancel{E}_T, \text{ Tsqk1}, 4(\text{flavor}) \times 2(\text{isospin}) = 8 \text{ mass degenerate states}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 575	95	20 KHACHATRY...17P	CMS	$1 \text{ or more jets} + \cancel{E}_T, \text{ Tsqk1}, \text{ one light flavor state}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1370	95	21 KHACHATRY...17V	CMS	$2 \gamma + \cancel{E}_T, \text{ GGM}, \text{ Tsqk4}, \text{ any NLSP mass}$	
>1600	95	22 SIRUNYAN	17AY CMS	$\gamma + \text{jets} + \cancel{E}_T, \text{ Tsqk4B}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1370	95	22 SIRUNYAN	17AY CMS	$\gamma + \text{jets} + \cancel{E}_T, \text{ Tsqk4A}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1050	95	23 SIRUNYAN	17AZ CMS	$\geq 1 \text{ jets} + \cancel{E}_T, \text{ Tsqk1}, \text{ single light flavor state}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1550	95	23 SIRUNYAN	17AZ CMS	$\geq 1 \text{ jets} + \cancel{E}_T, \text{ Tsqk1}, 4(\text{flavor}) \times 2(\text{isospin}) = 8 \text{ degenerate mass states}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1390	95	24 SIRUNYAN	17P CMS	$\text{jets} + \cancel{E}_T, \text{ Tsqk1}, 4(\text{flavor}) \times 2(\text{isospin}) = 8 \text{ degenerate mass states}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 950	95	24 SIRUNYAN	17P CMS	$\text{jets} + \cancel{E}_T, \text{ Tsqk1}, \text{ one light flavor state}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
> 608	95	25 AABOUD	16D ATLS	$\geq 1 \text{ jet} + \cancel{E}_T, \text{ Tsqk1}, m_{\tilde{q}} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$	
>1030	95	26 AABOUD	16N ATLS	$\geq 2 \text{ jets} + \cancel{E}_T, \text{ Tsqk1}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 600	95	27 KHACHATRY...16BS	CMS	$\text{jets} + \cancel{E}_T, \text{ Tsqk1}, \text{ single light squark}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1260	95	27 KHACHATRY...16BS	CMS	$\text{jets} + \cancel{E}_T, \text{ Tsqk1}, 8 \text{ degenerate light squarks}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
> 850	95	28 AAD	15BV ATLS	$\text{jets} + \cancel{E}_T, \tilde{q} \rightarrow q \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
> 250	95	29 AAD	15CS ATLS	$\text{photon} + \cancel{E}_T, pp \rightarrow \tilde{q} \tilde{q}^* \gamma, \tilde{q} \rightarrow q \tilde{\chi}_1^0, m_{\tilde{q}} - m_{\tilde{\chi}_1^0} = m_c$	
> 490	95	30 AAD	15K ATLS	$\tilde{c} \rightarrow c \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 200 \text{ GeV}$	
> 875	95	31 KHACHATRY...15AF	CMS	$\tilde{q} \rightarrow q \tilde{\chi}_1^0, \text{ simplified model}, 8 \text{ degenerate light } \tilde{q}, m_{\tilde{\chi}_1^0} = 0$	
> 520	95	31 KHACHATRY...15AF	CMS	$\tilde{q} \rightarrow q \tilde{\chi}_1^0, \text{ simplified model}, \text{ single light squark}, m_{\tilde{\chi}_1^0} = 0$	OCCUR=2

>1450	95	31	KHACHATRY...15AF	CMS	CMSSM, $\tan\beta = 30$, $A_0 = -2\max(m_0, m_{1/2})$, $\mu > 0$	OCCUR=3
> 850	95	32	AAD	14AE ATLS	jets + \cancel{E}_T , $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ simplified model, mass degenerate first and second generation squarks, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 440	95	32	AAD	14AE ATLS	jets + \cancel{E}_T , $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ simplified model, single light-flavour squark, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
>1700	95	32	AAD	14AE ATLS	jets + \cancel{E}_T , mSUGRA/CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	OCCUR=3
> 800	95	33	CHATRCHYAN 14AH	CMS	jets + \cancel{E}_T , $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 50$ GeV	
> 780	95	34	CHATRCHYAN 14I	CMS	multijets + \cancel{E}_T , $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} < 200$ GeV	
>1360	95	35	AAD	13L ATLS	jets + \cancel{E}_T , CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	
>1200	95	36	AAD	13Q ATLS	$\gamma + b + \cancel{E}_T$, higgsino-like neutralino, $m_{\tilde{\chi}_1^0} > 220$ GeV, GMSB	
>1250	95	37	CHATRCHYAN 13	CMS	$\ell^\pm \ell^\mp +$ jets + \cancel{E}_T , CMSSM	
		38	CHATRCHYAN 13G	CMS	0,1,2, ≥ 3 b-jets + \cancel{E}_T , CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	
>1430	95	39	CHATRCHYAN 13H	CMS	$2\gamma + \geq 4$ jets + low \cancel{E}_T , stealth SUSY model	
> 750	95	40	CHATRCHYAN 13T	CMS	jets + \cancel{E}_T , $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 820	95	41	AAD	12AX ATLS	$\ell +$ jets + \cancel{E}_T , CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	
>1200	95	42	AAD	12CJ ATLS	$\ell^\pm +$ jets + \cancel{E}_T , CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	
> 870	95	43	AAD	12CP ATLS	$2\gamma + \cancel{E}_T$, GMSB, bino NLSP, $m_{\tilde{\chi}_1^0} > 50$ GeV	
> 950	95	44	AAD	12W ATLS	jets + \cancel{E}_T , CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	
		45	CHATRCHYAN 12	CMS	e, μ , jets, razor, CMSSM	
> 760	95	46	CHATRCHYAN 12AE	CMS	jets + \cancel{E}_T , $\tilde{q} \rightarrow q\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} < 200$ GeV	
>1110	95	47	CHATRCHYAN 12AT	CMS	jets + \cancel{E}_T , CMSSM	
>1180	95	47	CHATRCHYAN 12AT	CMS	jets + \cancel{E}_T , CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>1080	95	48	AABOUD	18V ATLS	jets + \cancel{E}_T , Tsqk5, $(m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}) / (m_{\tilde{q}} - m_{\tilde{\chi}_1^0}) < 0.95$, $m_{\tilde{\chi}_1^0} = 60$ GeV	OCCUR=3
> 300	95	49	KHACHATRY...16BT	CMS	19-parameter pMSSM model, global Bayesian analysis, flat prior	
>1650	95	50	AAD	15AI ATLS	$\ell^\pm +$ jets + \cancel{E}_T	
		28	AAD	15BV ATLS	jets + \cancel{E}_T , $m_{\tilde{g}} = m_{\tilde{q}}$, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=2
> 790	95	28	AAD	15BV ATLS	jets + \cancel{E}_T , $\tilde{q} \rightarrow qW\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=3
> 820	95	28	AAD	15BV ATLS	2 or 3 leptons + jets, \tilde{q} decays via sleptons, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=4
> 850	95	28	AAD	15BV ATLS	τ , \tilde{q} decays via staus, $m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=5
> 700	95	51	KHACHATRY...15AR	CMS	$\tilde{q} \rightarrow q\tilde{\chi}_1^0$, $\tilde{\chi}_1^0 \rightarrow \tilde{S}g$, $\tilde{S} \rightarrow S\tilde{G}$, $S \rightarrow gg$, $m_{\tilde{S}} = 100$ GeV, $m_S = 90$ GeV	
> 550	95	51	KHACHATRY...15AR	CMS	ℓ^\pm , $\tilde{q} \rightarrow q\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow \tilde{S}W^\pm$, $\tilde{S} \rightarrow S\tilde{G}$, $S \rightarrow gg$, $m_{\tilde{S}} = 100$ GeV, $m_S = 90$ GeV	OCCUR=2
>1500	95	52	KHACHATRY...15AZ	CMS	$\geq 2 \gamma$, ≥ 1 jet, (Razor), bino-like NLSP, $m_{\tilde{\chi}_1^0} = 375$ GeV	

>1000	95	⁵² KHACHATRY...15AZ	CMS	$\geq 1 \gamma, \geq 2 \text{ jet, wino-like NLSP, } m_{\tilde{\chi}_1^0} = 375 \text{ GeV}$	OCCUR=2
> 670	95	⁵³ AAD	14E ATLS	$\ell^\pm \ell^\mp (\ell^\mp) + \text{jets, } \tilde{q} \rightarrow q' \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow Z^{(*)} \tilde{\chi}_1^0 \text{ simplified model, } m_{\tilde{\chi}_1^0} < 300 \text{ GeV}$	
> 780	95	⁵³ AAD	14E ATLS	$\ell^\pm \ell^\mp (\ell^\mp) + \text{jets, } \tilde{q} \rightarrow q' \tilde{\chi}_1^\pm / \tilde{\chi}_2^0, \tilde{\chi}_1^\pm \rightarrow \ell^\pm \nu \tilde{\chi}_1^0, \tilde{\chi}_2^0 \rightarrow \ell^\pm \ell^\mp (\nu\nu) \tilde{\chi}_1^0 \text{ simplified model}$	OCCUR=2
> 700	95	⁵⁴ CHATRCHYAN 13AO	CMS	$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T, \text{ CMSSM, } m_0 < 700 \text{ GeV}$	
>1350	95	⁵⁵ CHATRCHYAN 13AV	CMS	$\text{jets (+ leptons)} + \cancel{E}_T, \text{ CMSSM, } m_{\tilde{g}} = m_{\tilde{q}}$	
> 800	95	⁵⁶ CHATRCHYAN 13W	CMS	$\geq 1 \text{ photons} + \text{jets} + \cancel{E}_T, \text{ GGM, wino-like NLSP, } m_{\tilde{\chi}_1^0} = 375 \text{ GeV}$	
>1000	95	⁵⁶ CHATRCHYAN 13W	CMS	$\geq 2 \text{ photons} + \text{jets} + \cancel{E}_T, \text{ GGM, bino-like NLSP, } m_{\tilde{\chi}_1^0} = 375 \text{ GeV}$	OCCUR=2
> 340	95	⁵⁷ DREINER	12A THEO	$m_{\tilde{q}} \sim m_{\tilde{\chi}_1^0}$	
> 650	95	⁵⁸ DREINER	12A THEO	$m_{\tilde{q}} = m_{\tilde{g}} \sim m_{\tilde{\chi}_1^0}$	OCCUR=2

¹ AAD 23AE searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 2 ℓ with same flavour and opposite sign, plus jets and \cancel{E}_T , defining signal region with the dilepton invariant mass both on- and off-shell with respect to the Z boson. No significant excess above the Standard Model predictions is observed. Limits are set on models of strong and electroweak production. In this case, limits are placed on the mass of pair-produced squarks, assuming a scenario like in Tsqk2, see figure 16.

NODE=S046SQK;LINKAGE=VB

² TUMASYAN 23X searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for squark pair production with cascade decays to CP -even singlet-like Higgs bosons (H_1), leading to final states with small missing transverse momentum. This search targets H_1 decays to $b\bar{b}$ -pairs that are reconstructed in large-area (AK8) jets. No significant excess above the Standard Model expectations is observed. Limits are set in the next-to-minimal supersymmetric extension of the SM, where a singlino of small mass leads to squark and gluino cascade decays that can predominantly end in a highly Lorentz-boosted singlet-like H_1 and a singlino-like neutralino $\tilde{\chi}_S^0$ of small transverse momentum. The eight first- and second-generation squarks are assumed mass-degenerate, and the gluino mass is set at 1% larger.

NODE=S046SQK;LINKAGE=WB

³ AAD 21AK searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of gluinos and squarks in events with a single isolated electron or muon, originating from the decay of a W boson, multiple jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1B simplified model and on the squark mass in the Tsqk3 simplified model, see their Figure 8.

NODE=S046SQK;LINKAGE=RB

⁴ AAD 21F searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of squarks in events with a high- p_T jet and \cancel{E}_T . No significant excess above the Standard Model predictions is observed. Limits are set on the \tilde{t} mass in the Tstop3 and Tstop4, on the \tilde{b} mass in the Tstop1, and on the \tilde{q} mass in the Tsqk1 simplified model (four-flavour, two chirality states degeneracy).

NODE=S046SQK;LINKAGE=TB

⁵ AAD 21L searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of gluinos and squarks in events with jets, large missing transverse momentum but no electrons or muons. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A and Tglu1B simplified models, on the squark mass in the Tsqk1 and Tsqk3 simplified models and in a simplified model for gluino-squark production, see their Figures 13-17.

NODE=S046SQK;LINKAGE=SB

⁶ SIRUNYAN 21M searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tstop3, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14.

NODE=S046SQK;LINKAGE=UB

⁷ SIRUNYAN 19AG searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4B simplified model and on the squark mass in the Tsqk4B simplified model, see their Figure 3.

NODE=S046SQK;LINKAGE=MB

- 8 SIRUNYAN 19CH searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A and Tglu3A simplified models, see their Figure 13. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tshot1, Tstop1 simplified models, see their Figure 14. NODE=S046SQK;LINKAGE=OB
- 9 SIRUNYAN 19K searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a photon, an electron or muon, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. In the framework of GMSB, limits are set on the chargino and neutralino mass in the Tchi1n1A simplified model, see their Figure 6. Limits are also set on the gluino mass in the Tglu4A simplified model, and on the squark mass in the Tsqk4A simplified model, see their Figure 7. NODE=S046SQK;LINKAGE=NB
- 10 AABOUD 18BJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with two opposite-sign charged leptons (electrons and muons), jets and missing transverse momentum, with various requirements to be sensitive to signals with different kinematic endpoint values in the dilepton invariant mass distribution. The data are found to be consistent with the SM expectation. Results are interpreted in the Tsqk2 model in case of $m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$: for any $m_{\tilde{\chi}_2^0}$, squark masses below 1200 GeV are excluded, see their Fig. 14(b). NODE=S046SQK;LINKAGE=GB
- 11 AABOUD 18BV searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one jet identified as c -jet, large missing transverse energy and no leptons. Good agreement is observed between the number of events in data and Standard Model predictions. The results are translated into exclusion limits in Tsqk1 models considering only \tilde{c}_1 . In scenarios with massless neutralinos, scharm masses below 850 GeV are excluded. If the differences of the \tilde{c}_1 and $\tilde{\chi}_1^0$ masses is below 100 GeV, scharm masses below 500 GeV are excluded. See their Fig.6 and Fig.7. NODE=S046SQK;LINKAGE=KB
- 12 AABOUD 18I searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one jet with a transverse momentum above 250 GeV and no leptons. Good agreement is observed between the number of events in data and Standard Model predictions. The results are translated into exclusion limits in Tsqk1 models. In the compressed scenario with similar squark and neutralino masses, squark masses below 710 GeV are excluded. See their Fig.10(b). NODE=S046SQK;LINKAGE=BB
- 13 AABOUD 18U searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with at least one isolated photon, possibly jets and significant transverse momentum targeting generalised models of gauge-mediated SUSY breaking. No significant excess of events is observed above the SM prediction. Results are interpreted in terms of lower limits on the masses of squark in Tsqk4B models. Masses below 1820 GeV are excluded for any NLSP mass, see their Fig. 9. NODE=S046SQK;LINKAGE=EB
- 14 AABOUD 18V searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with no charged leptons, jets and missing transverse momentum. The data are found to be consistent with the SM expectation. Results are interpreted in the Tsqk1 model: squark masses below 1550 GeV are excluded for massless LSP, see their Fig. 13(a). NODE=S046SQK;LINKAGE=HB
- 15 AABOUD 18V searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with no charged leptons, jets and missing transverse momentum. The data are found to be consistent with the SM expectation. Results are interpreted in the Tsqk3 model. Assuming that $m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{q}} + m_{\tilde{\chi}_1^0})$, squark masses below 1150 GeV are excluded for massless LSP, see their Fig. 14(a). Exclusions are also shown assuming $m_{\tilde{\chi}_1^0} = 60 \text{ GeV}$, see their Fig. 14(b). NODE=S046SQK;LINKAGE=IB
- 16 SIRUNYAN 18AA searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one photon and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on wino masses in a general gauge-mediated SUSY breaking (GGM) scenario with bino-like $\tilde{\chi}_1^0$ and wino-like $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$, see Figure 7. Limits are also set on the NLSP mass in the Tchi1n1A and Tchi1chi1A simplified models, see their Figure 8. Finally, limits are set on the gluino mass in the Tglu4A and Tglu4B simplified models, see their Figure 9, and on the squark mass in the Tskq4A and Tsqk4B simplified models, see their Figure 10. NODE=S046SQK;LINKAGE=AB
- 17 SIRUNYAN 18AY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing one or more jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A and Tglu3A simplified models, see their Figure 3. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tshot1, Tstop1 and Tstop4 simplified models, see their Figure 3. Finally, limits are set on long-lived gluino masses in a Tglu1A simplified model where the gluino is metastable or long-lived with proper decay lengths in the range $10^{-3} \text{ mm} < c\tau < 10^5 \text{ mm}$, see their Figure 4. NODE=S046SQK;LINKAGE=DB
- 18 AABOUD 17AR searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one isolated lepton, at least two jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.25 TeV are set on the 1st and 2nd generation squark masses in Tsqk3 simplified models, with $x = (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}) / (m_{\tilde{q}} - m_{\tilde{\chi}_1^0}) = 1/2$. Similar limits are obtained for variable x and fixed neutralino mass, $m_{\tilde{\chi}_1^0} = 60 \text{ GeV}$. See their Figure 13. NODE=S046SQK;LINKAGE=ZA
- 19 AABOUD 17N searched in 14.7 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 2 same-flavour, opposite-sign leptons (electrons or muons), jets and large missing transverse momentum. The results are interpreted as 95% C.L. limits in Tsqk2 models, NODE=S046SQK;LINKAGE=WA

- assuming $m_{\tilde{\chi}_1^0} = 0$ GeV and $m_{\tilde{\chi}_2^0} = 600$ GeV. See their Fig. 12 for exclusion limits as a function of $m_{\tilde{\chi}_2^0}$.
- 20 KHACHATRYAN 17P searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figures 7 and 8. Limits are also set on the squark mass in the Tsqk1 simplified model, see their Fig. 7, and on the sbottom mass in the Tsb0t1 simplified model, see Fig. 8. Finally, limits are set on the stop mass in the Tstop1, Tstop3, Tstop4, Tstop6 and Tstop7 simplified models, see Fig. 8. NODE=S046SQK;LINKAGE=VA
- 21 KHACHATRYAN 17V searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with two photons and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino and squark mass in the context of general gauge mediation models Tglu4B and Tsqk4, see their Fig. 4. NODE=S046SQK;LINKAGE=TA
- 22 SIRUNYAN 17AY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one photon, jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4A and Tglu4B simplified models, and on the squark mass in the Tskq4A and Tskq4B simplified models, see their Figure 6. NODE=S046SQK;LINKAGE=YA
- 23 SIRUNYAN 17AZ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A simplified models, see their Figures 6. Limits are also set on the squark mass in the Tsqk1 simplified model (for single light squark and for 8 degenerate light squarks), on the sbottom mass in the Tsb0t1 simplified model and on the stop mass in the Tstop1 simplified model, see their Fig. 7. Finally, limits are set on the stop mass in the Tstop2, Tstop4 and Tstop8 simplified models, see Fig. 8. NODE=S046SQK;LINKAGE=XA
- 24 SIRUNYAN 17P searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A, Tglu3A and Tglu3D simplified models, see their Fig. 12. Limits are also set on the squark mass in the Tsqk1 simplified model, on the stop mass in the Tstop1 simplified model, and on the sbottom mass in the Tsb0t1 simplified model, see Fig. 13. NODE=S046SQK;LINKAGE=UA
- 25 AABOUD 16D searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with an energetic jet and large missing transverse momentum. The results are interpreted as 95% C.L. limits on masses of first and second generation squarks decaying into a quark and the lightest neutralino in scenarios with $m_{\tilde{q}} - m_{\tilde{\chi}_1^0} < 25$ GeV. See their Fig. 6. NODE=S046SQK;LINKAGE=KA
- 26 AABOUD 16N searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing hadronic jets, large \cancel{E}_T , and no electrons or muons. No significant excess above the Standard Model expectations is observed. First- and second-generation squark masses below 1030 GeV are excluded at the 95% C.L. decaying to quarks and a massless lightest neutralino. See their Fig. 7a. NODE=S046SQK;LINKAGE=OA
- 27 KHACHATRYAN 16BS searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one energetic jet, no isolated leptons, and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the squark mass in the Tskq1 simplified model, both in the assumption of a single light squark and of 8 degenerate squarks, see Fig. 11 and Table 3. NODE=S046SQK;LINKAGE=PA
- 28 AAD 15BV summarized and extended ATLAS searches for gluinos and first- and second-generation squarks in final states containing jets and missing transverse momentum, with or without leptons or b -jets in the $\sqrt{s} = 8$ TeV data set collected in 2012. The paper reports the results of new interpretations and statistical combinations of previously published analyses, as well as new analyses. Exclusion limits at 95% C.L. are set on the squark mass in several R-parity conserving models. See their Figs. 9, 11, 18, 22, 24, 27, 28. NODE=S046SQK;LINKAGE=FA
- 29 AAD 15CS searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for evidence of pair production of squarks, decaying into a quark and a neutralino, where a photon was radiated either from an initial-state quark, from an intermediate squark, or from a final-state quark. No evidence was found for an excess above the expected level of Standard Model background and a 95% C.L. exclusion limit was set on the squark mass as a function of the squark-neutralino mass difference, see Fig. 19. NODE=S046SQK;LINKAGE=NA
- 30 AAD 15K searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing at least two jets, where the two leading jets are each identified as originating from c -quarks, and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the mass of superpartners of charm quarks (\tilde{c}). Assuming that the decay $\tilde{c} \rightarrow c\tilde{\chi}_1^0$ takes place 100% of the time, a scalar charm mass below 490 GeV is excluded for $m_{\tilde{\chi}_1^0} < 200$ GeV. For more details, see their Fig. 2. NODE=S046SQK;LINKAGE=HA
- 31 KHACHATRYAN 15AF searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with at least two energetic jets and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the squark mass in simplified models where the decay $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, both for the case of a single light squark or 8 degenerate squarks, see Fig. 12. NODE=S046SQK;LINKAGE=Y

See also Table 5. Exclusions in the CMSSM, assuming $\tan\beta = 30$, $A_0 = -2 \max(m_0, m_{1/2})$ and $\mu > 0$, are also presented, see Fig. 15.

- 32 AAD 14AE searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for strongly produced supersymmetric particles in events containing jets and large missing transverse momentum, and no electrons or muons. No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing squarks that decay via $\tilde{q} \rightarrow q\tilde{\chi}_1^0$, where either a single light state or two degenerate generations of squarks are assumed, see Fig. 10. NODE=S046SQK;LINKAGE=V
- 33 CHATRCHYAN 14AH searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the razor variables (M_R and R^2) to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on squark masses in simplified models where the decay $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 28. Exclusions in the CMSSM, assuming $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, are also presented, see Fig. 26. NODE=S046SQK;LINKAGE=X
- 34 CHATRCHYAN 14I searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing multijets and large \cancel{E}_T . No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing squarks that decay via $\tilde{q} \rightarrow q\tilde{\chi}_1^0$, where either a single light state or two degenerate generations of squarks are assumed, see Fig. 7a. NODE=S046SQK;LINKAGE=I
- 35 AAD 13L searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for the production of squarks and gluinos in events containing jets, missing transverse momentum and no high- p_T electrons or muons. No excess over the expected SM background is observed. In mSUGRA/CMSSM models with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, squarks and gluinos of equal mass are excluded for masses below 1360 GeV at 95% C.L. In a simplified model containing only squarks of the first two generations, a gluino octet and a massless neutralino, squark masses below 1320 GeV are excluded at 95% C.L. for gluino masses below 2 TeV. See Figures 10–15 for more precise bounds. NODE=S046SQK;LINKAGE=GL
- 36 AAD 13Q searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing a high- p_T isolated photon, at least one jet identified as originating from a bottom quark, and high missing transverse momentum. Such signatures may originate from supersymmetric models with gauge-mediated supersymmetry breaking in events in which one of a pair of higgsino-like neutralinos decays into a photon and a gravitino while the other decays into a Higgs boson and a gravitino. No significant excess above the expected background was found and limits were set on the squark mass as a function of the neutralino mass in a generalized GMSB model (GGM) with a higgsino-like neutralino NLSP, see their Fig. 4. For neutralino masses greater than 220 GeV, squark masses below 1020 GeV are excluded at 95% C.L. NODE=S046SQK;LINKAGE=SA
- 37 CHATRCHYAN 13 looked in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with two opposite-sign leptons (e, μ, τ), jets and missing transverse energy. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in the mSUGRA/CMSSM model with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, see Fig. 6. NODE=S046SQK;LINKAGE=CY
- 38 CHATRCHYAN 13G searched in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for the production of squarks and gluinos in events containing 0,1,2, ≥ 3 b -jets, missing transverse momentum and no electrons or muons. No excess over the expected SM background is observed. In mSUGRA/CMSSM models with $\tan\beta = 10$, $A_0 = 0$, and $\mu > 0$, squarks and gluinos of equal mass are excluded for masses below 1250 GeV at 95% C.L. Exclusions are also derived in various simplified models, see Fig. 7. NODE=S046SQK;LINKAGE=CR
- 39 CHATRCHYAN 13H searched in 4.96 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with two photons, ≥ 4 jets and low \cancel{E}_T due to $\tilde{q} \rightarrow \gamma\tilde{\chi}_1^0$ decays in a stealth SUSY framework, where the $\tilde{\chi}_1^0$ decays through a singlino (\tilde{S}) intermediate state to $\gamma\tilde{S}\tilde{G}$, with the singlet state \tilde{S} decaying to two jets. No significant excess above the expected background was found and limits were set in a particular R-parity conserving stealth SUSY model. The model assumes $m_{\tilde{\chi}_1^0} = 0.5 m_{\tilde{q}}$, $m_{\tilde{S}} = 100 \text{ GeV}$ and $m_{\tilde{G}} = 90 \text{ GeV}$. Under these assumptions, squark masses less than 1430 GeV were excluded at the 95% C.L. NODE=S046SQK;LINKAGE=CN
- 40 CHATRCHYAN 13T searched in 11.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the α_T variable to discriminate between processes with genuine and misreconstructed \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on squark masses in simplified models where the decay $\tilde{q} \rightarrow q\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, assuming an eightfold degeneracy of the masses of the first two generation squarks, see Fig. 8 and Table 9. Also limits in the case of a single light squark are given. NODE=S046SQK;LINKAGE=CS
- 41 AAD 12AX searched in 1.04 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for supersymmetry in events containing jets, missing transverse momentum and one isolated electron or muon. No excess over the expected SM background is observed and model-independent limits are set on the cross section of new physics contributions to the signal regions. In mSUGRA/CMSSM models with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, squarks and gluinos of equal mass are excluded for masses below 820 GeV at 95% C.L. Limits are also set on simplified models for squark production and decay via an intermediate chargino and on supersymmetric models with bilinear R-parity violation. Supersedes AAD 11G. NODE=S046SQK;LINKAGE=AX
- 42 AAD 12CJ searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing one or more isolated leptons (electrons or muons), jets and \cancel{E}_T . The observations are in good agreement with the SM expectations and exclusion limits have been set in number NODE=S046SQK;LINKAGE=GA

- of SUSY models. In the mSUGRA/CMSSM model with $\tan\beta = 10$, $A_0 = 0$, and $\mu > 0$, 95% C.L. exclusion limits have been derived for $m_{\tilde{q}} < 1200$ GeV, assuming equal squark and gluino masses. In minimal GMSB, values of the effective SUSY breaking scale $\Lambda < 50$ TeV are excluded at 95% C.L. for $\tan\beta < 45$. Also exclusion limits in a number of simplified models have been presented, see Figs. 10 and 12.
- 43 AAD 12CP searched in 4.8 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with two photons and large \cancel{E}_T due to $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$ decays in a GMSB framework. No significant excess above the expected background was found and limits were set on the squark mass as a function of the neutralino mass in a generalized GMSB model (GGM) with a bino-like neutralino NLSP. The other sparticle masses were decoupled, $\tan\beta = 2$ and $c\tau_{NLSP} < 0.1$ mm. Also, in the framework of the SPS8 model, a 95% C.L. lower limit was set on the breaking scale Λ of 196 TeV. NODE=S046SQK;LINKAGE=GD
- 44 AAD 12W searched in 1.04 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for the production of squarks and gluinos in events containing jets, missing transverse momentum and no electrons or muons. No excess over the expected SM background is observed. In mSUGRA/CMSSM models with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, squarks and gluinos of equal mass are excluded for masses below 950 GeV at 95% C.L. In a simplified model containing only squarks of the first two generations, a gluino octet and a massless neutralino, squark masses below 875 GeV are excluded at 95% C.L. NODE=S046SQK;LINKAGE=TG
- 45 CHATRCHYAN 12 looked in 35 pb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with e and/or μ and/or jets, a large total transverse energy, and \cancel{E}_T . The event selection is based on the dimensionless razor variable R , related to the \cancel{E}_T and M_R , an indicator of the heavy particle mass scale. No evidence for an excess over the expected background is observed. Limits are derived in the CMSSM ($m_0, m_{1/2}$) plane for $\tan\beta = 3, 10$ and 50 (see Fig. 7 and 8). Limits are also obtained for Simplified Model Spectra. NODE=S046SQK;LINKAGE=C9
- 46 CHATRCHYAN 12AE searched in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with at least three jets and large missing transverse momentum. No significant excesses over the expected SM backgrounds are observed and 95% C.L. limits on the production cross section of squarks in a scenario where $\tilde{q} \rightarrow q \tilde{\chi}_1^0$ with a 100% branching ratio, see Fig. 3. For $m_{\tilde{\chi}_1^0} < 200$ GeV, values of $m_{\tilde{q}}$ below 760 GeV are excluded at 95% C.L. Also limits in the CMSSM are presented, see Fig. 2. NODE=S046SQK;LINKAGE=HT
- 47 CHATRCHYAN 12AT searched in 4.73 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for the production of squarks and gluinos in events containing jets, missing transverse momentum and no electrons or muons. No excess over the expected SM background is observed. In mSUGRA/CMSSM models with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, squarks with masses below 1110 GeV are excluded at 95% C.L. Squarks and gluinos of equal mass are excluded for masses below 1180 GeV at 95% C.L. Exclusions are also derived in various simplified models, see Fig. 6. NODE=S046SQK;LINKAGE=CA
- 48 AABOUD 18V searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV in events with no charged leptons, jets and missing transverse momentum. The data are found to be consistent with the SM expectation. Results are interpreted in the Tsqk5 model. Squark masses below 1100 GeV are excluded if $(m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0})/(m_{\tilde{q}} - m_{\tilde{\chi}_1^0}) < 0.95$ and $m_{\tilde{\chi}_1^0} = 60$ GeV, see their Fig. 16(a). NODE=S046SQK;LINKAGE=JB
- 49 KHACHATRYAN 16BT performed a global Bayesian analysis of a wide range of CMS results obtained with data samples corresponding to 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV and in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV. The set of searches considered, both individually and in combination, includes those with all-hadronic final states, same-sign and opposite-sign dileptons, and multi-lepton final states. An interpretation was given in a scan of the 19-parameter pMSSM. No scan points with a gluino mass less than 500 GeV survived and 98% of models with a squark mass less than 300 GeV were excluded. NODE=S046SQK;LINKAGE=QA
- 50 AAD 15AI searched in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing at least one isolated lepton (electron or muon), jets, and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the squark masses in the CMSSM/mSUGRA, see Fig. 15, in the NUHMG, see Fig. 16, and in various simplified models, see Figs. 19–21. NODE=S046SQK;LINKAGE=EA
- 51 KHACHATRYAN 15AR searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing jets, either a charged lepton or a photon, and low missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the squark mass in a stealth SUSY model where the decays $\tilde{q} \rightarrow q \tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow \tilde{S} W^\pm$, $\tilde{S} \rightarrow S \tilde{G}$ and $S \rightarrow gg$, with $m_{\tilde{S}} = 100$ GeV and $m_S = 90$ GeV, take place with a branching ratio of 100%. See Fig. 6 for γ or Fig. 7 for ℓ^\pm analyses. NODE=S046SQK;LINKAGE=Z
- 52 KHACHATRYAN 15AZ searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with either at least one photon, hadronic jets and \cancel{E}_T (single photon channel) or with at least two photons and at least one jet and using the razor variables. No significant excess above the Standard Model expectations is observed. Limits are set on gluino masses in the general gauge-mediated SUSY breaking model (GGM), for both a bino-like and wino-like neutralino NLSP scenario, see Fig. 8 and 9. NODE=S046SQK;LINKAGE=LA
- 53 AAD 14E searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for strongly produced supersymmetric particles in events containing jets and two same-sign leptons or three leptons. The search also utilises jets originating from b -quarks, missing transverse momentum and other variables. No excess over the expected SM background is observed. NODE=S046SQK;LINKAGE=H

Exclusion limits are derived in simplified models containing gluinos and squarks, see Figures 5 and 6. In the $\tilde{q} \rightarrow q' \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow Z^{(*)} \tilde{\chi}_1^0$ simplified model, the following assumptions have been made: $m_{\tilde{\chi}_1^\pm} = 0.5 m_{\tilde{\chi}_1^0} + m_{\tilde{g}}$, $m_{\tilde{\chi}_2^0} = 0.5 (m_{\tilde{\chi}_1^0} + m_{\tilde{g}})$. In the $\tilde{q} \rightarrow q' \tilde{\chi}_1^\pm$ or $\tilde{q} \rightarrow q' \tilde{\chi}_2^0, \tilde{\chi}_1^\pm \rightarrow \ell^\pm \nu \tilde{\chi}_1^0$ or $\tilde{\chi}_2^0 \rightarrow \ell^\pm \ell^\mp (\nu \nu) \tilde{\chi}_1^0$ simplified model, the following assumptions have been made: $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0} = 0.5 (m_{\tilde{\chi}_1^0} + m_{\tilde{g}})$, $m_{\tilde{\chi}_1^0} < 460$ GeV. Limits are also derived in the mSUGRA/CMSSM, bRPV and GMSB models, see their Fig. 8.

- 54 CHATRCHYAN 13A0 searched in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with two opposite-sign isolated leptons accompanied by hadronic jets and \cancel{E}_T . No significant excesses over the expected SM backgrounds are observed and 95% C.L. exclusion limits are derived in the mSUGRA/CMSSM model with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, see Fig. 8. NODE=S046SQK;LINKAGE=NC
- 55 CHATRCHYAN 13AV searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for new heavy particle pairs decaying into jets (possibly b -tagged), leptons and \cancel{E}_T using the Razor variables. No significant excesses over the expected SM backgrounds are observed and 95% C.L. exclusion limits are derived in the mSUGRA/CMSSM model with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, see Fig. 3. The results are also interpreted in various simplified models, see Fig. 4. NODE=S046SQK;LINKAGE=MC
- 56 CHATRCHYAN 13W searched in 4.93 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with one or more photons, hadronic jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on squark masses in the general gauge-mediated SUSY breaking model (GGM), for both a wino-like and bino-like neutralino NLSP scenario, see Fig. 5. NODE=S046SQK;LINKAGE=MS
- 57 DREINER 12A reassesses constraints from CMS (at 7 TeV, $\sim 4.4 \text{ fb}^{-1}$) under the assumption that the first and second generation squarks and the lightest SUSY particle are quasi-degenerate in mass (compressed spectrum). NODE=S046SQK;LINKAGE=DR
- 58 DREINER 12A reassesses constraints from CMS (at 7 TeV, $\sim 4.4 \text{ fb}^{-1}$) under the assumption that the first and second generation squarks, the gluino, and the lightest SUSY particle are quasi-degenerate in mass (compressed spectrum). NODE=S046SQK;LINKAGE=DE

R-parity violating \tilde{q} (Squark) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 100–720	95	¹ SIRUNYAN	18EA CMS	2 large jets with four-parton substructure, $\tilde{q} \rightarrow 4q$
>1600	95	² KHACHATRYAN...16BX	CMS	$\tilde{q} \rightarrow q \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell \ell \nu$, $\lambda_{121} \neq 0$ or $\lambda_{122} \neq 0$, $m_{\tilde{g}} = 2400$ GeV
>1000	95	³ AAD	15CB ATLS	jets, $\tilde{q} \rightarrow q \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell q q$, $m_{\tilde{\chi}_1^0} = 108$ GeV and $2.5 < c\tau_{\tilde{\chi}_1^0} < 200$ mm
		⁴ AAD	12AX ATLS	ℓ + jets + \cancel{E}_T , CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$
		⁵ CHATRCHYAN 12AL	CMS	$\geq 3\ell^\pm$

NODE=S046SQV
NODE=S046SQV

- ¹ SIRUNYAN 18EA searched in 38.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for the pair production of resonances, each decaying to at least four quarks. Reconstructed particles are clustered into two large jets of similar mass, each consistent with four-parton substructure. No statistically significant excess over the Standard Model expectation is observed. Limits are set on the squark and gluino mass in RPV supersymmetry models where squarks (gluinos) decay, through intermediate higgsinos, to four (five) quarks, see their Figure 4. NODE=S046SQV;LINKAGE=A
- ² KHACHATRYAN 16BX searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing 4 leptons coming from R-parity-violating decays of $\tilde{\chi}_1^0 \rightarrow \ell \ell \nu$ with $\lambda_{121} \neq 0$ or $\lambda_{122} \neq 0$. No excess over the expected background is observed. Limits are derived on the gluino, squark and stop masses, see Fig. 23. NODE=S046SQV;LINKAGE=RA
- ³ AAD 15CB searched for events containing at least one long-lived particle that decays at a significant distance from its production point (displaced vertex, DV) into two leptons or into five or more charged particles in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV. The dilepton signature is characterised by DV formed from at least two lepton candidates. Four different final states were considered for the multitrack signature, in which the DV must be accompanied by a high-transverse momentum muon or electron candidate that originates from the DV, jets or missing transverse momentum. No events were observed in any of the signal regions. Results were interpreted in SUSY scenarios involving R-parity violation, split supersymmetry, and gauge mediation. See their Fig. 14–20. NODE=S046SQV;LINKAGE=JA
- ⁴ AAD 12AX searched in 1.04 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for supersymmetry in events containing jets, missing transverse momentum and one isolated electron or muon. No excess over the expected SM background is observed and model-independent limits are set on the cross section of new physics contributions to the signal regions. In mSUGRA/CMSSM models with $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, squarks and gluinos of equal mass are excluded for masses below 820 GeV at 95% C.L. Limits are also set on simplified models for squark production and decay via an intermediate chargino and on supersymmetric models with bilinear R-parity violation. Supersedes AAD 11G. NODE=S046SQV;LINKAGE=AX

⁵ CHATRCHYAN 12AL looked in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for anomalous production of events with three or more isolated leptons. Limits on squark and gluino masses are set in RPV SUSY models with leptonic $LL\bar{E}$ couplings, $\lambda_{123} > 0.05$, and hadronic $U\bar{D}\bar{D}$ couplings, $\lambda''_{112} > 0.05$, see their Fig. 5. In the $U\bar{D}\bar{D}$ case the leptons arise from supersymmetric cascade decays. A very specific supersymmetric spectrum is assumed. All decays are prompt.

NODE=S046SQV;LINKAGE=HR

Long-lived \tilde{q} (Squark) mass limit

The following are bounds on long-lived scalar quarks, assumed to hadronise into hadrons with lifetime long enough to escape the detector prior to a possible decay. Limits may depend on the mixing angle of mass eigenstates: $\tilde{q}_1 = \tilde{q}_L \cos\theta_q + \tilde{q}_R \sin\theta_q$.

The coupling to the Z^0 boson vanishes for up-type squarks when $\theta_u=0.98$, and for down type squarks when $\theta_d=1.17$.

NODE=S046SSQ

NODE=S046SSQ

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1250	95	1 AABOUD	19AT ATLS	\tilde{b} R-hadrons
>1340	95	2 AABOUD	19AT ATLS	\tilde{t} R-hadrons
>1600	95	3 SIRUNYAN	19BH CMS	long-lived \tilde{t} , RPV, $\tilde{t} \rightarrow \bar{d}\bar{d}$, 10 mm < $c\tau$ < 110 mm
>1350	95	3 SIRUNYAN	19BH CMS	long-lived \tilde{t} , RPV, $\tilde{t} \rightarrow b\bar{l}$, 7 mm < $c\tau$ < 110 mm
> 805	95	4 AABOUD	16B ATLS	\tilde{b} R-hadrons
> 890	95	5 AABOUD	16B ATLS	\tilde{t} R-hadrons
>1040	95	6 KHACHATRY...16BWCMS		\tilde{t} R-hadrons, cloud interaction model
>1000	95	6 KHACHATRY...16BWCMS		\tilde{t} R-hadrons, charge-suppressed interaction model
> 845	95	7 AAD	15AE ATLS	\tilde{b} R-hadron, stable, Regge model
> 900	95	7 AAD	15AE ATLS	\tilde{t} R-hadron, stable, Regge model
>1500	95	7 AAD	15AE ATLS	\tilde{g} decaying to 300 GeV stable sleptons, LeptoSUSY model
> 751	95	8 AAD	15BMATLS	\tilde{b} R-hadron, stable, Regge model
> 766	95	8 AAD	15BMATLS	\tilde{t} R-hadron, stable, Regge model
> 525	95	9 KHACHATRY...15AK CMS		\tilde{t} R-hadrons, 10 $\mu\text{s} < \tau < 1000 \text{ s}$
> 470	95	9 KHACHATRY...15AK CMS		\tilde{t} R-hadrons, 1 $\mu\text{s} < \tau < 1000 \text{ s}$

NODE=S046SSQ;CHECK LIMITS

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 683	95	10 AAD	13AA ATLS	\tilde{t} , R-hadrons, generic interaction model
> 612	95	11 AAD	13AA ATLS	\tilde{b} , R-hadrons, generic interaction model
> 344	95	12 AAD	13BC ATLS	R-hadrons, $\tilde{t} \rightarrow b\tilde{\chi}_1^0$, Regge model, lifetime between 10^{-5} and 10^3 s , $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$
> 379	95	13 AAD	13BC ATLS	R-hadrons, $\tilde{t} \rightarrow t\tilde{\chi}_1^0$, Regge model, lifetime between 10^{-5} and 10^3 s , $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$
> 935	95	14 CHATRCHYAN 13AB CMS		long-lived \tilde{t} forming R-hadrons, cloud interaction model

NODE=S046SSQ;LINKAGE=H

¹ AABOUD 19AT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for metastable and stable R-hadrons. Multiple search strategies for a wide range of lifetimes, corresponding to path lengths of a few meters, are defined. No significant deviations from the expected Standard Model background are observed. Stopped R-hadrons are excluded at 95% C.L. for masses below 1250 GeV. Less stringent constraints are achieved with the muon-spectrometer agnostic analysis. See their Figure 9 (bottom-left).

NODE=S046SSQ;LINKAGE=I

² AABOUD 19AT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for metastable and stable R-hadrons. Multiple search strategies for a wide range of lifetimes, corresponding to path lengths of a few meters, are defined. No significant deviations from the expected Standard Model background are observed. Stopped R-hadrons are excluded at 95% C.L. for masses below 1340 GeV. Similar constraints are achieved with the muon-spectrometer agnostic analysis. See their Figure 9 (bottom-right).

NODE=S046SSQ;LINKAGE=G

³ SIRUNYAN 19BH searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived particles decaying into jets, with each long-lived particle having a decay vertex well displaced from the production vertex. The selected events are found to be consistent with standard model predictions. Limits are set on the gluino mass in a GMSB model where the gluino is decaying via $\tilde{g} \rightarrow g\tilde{G}$, see their Figure 4 and in an RPV model of supersymmetry where the gluino is decaying via $\tilde{g} \rightarrow \tilde{t}\bar{b}\bar{s}$, see their Figures 5. Limits are also set on the stop mass in two RPV models, see their Figure 6 (for $\tilde{t} \rightarrow b\bar{l}$ decays) and Figure 7 (for $\tilde{t} \rightarrow \bar{d}\bar{d}$ decays).

NODE=S046SSQ;LINKAGE=B

⁴ AABOUD 16B searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived R-hadrons using observables related to large ionization losses and slow propagation velocities, which are signatures of heavy charged particles traveling significantly slower than

- the speed of light. Exclusion limits at 95% C.L. are set on the long-lived sbottom masses exceeding 805 GeV. See their Fig. 5.
- 5 AABOUD 16B searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived R -hadrons using observables related to large ionization losses and slow propagation velocities, which are signatures of heavy charged particles traveling significantly slower than the speed of light. Exclusion limits at 95% C.L. are set on the long-lived stop masses exceeding 890 GeV. See their Fig. 5. NODE=S046SSQ;LINKAGE=D
- 6 KHACHATRYAN 16BW searched in 2.5 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with heavy stable charged particles, identified by their anomalously high energy deposits in the silicon tracker and/or long time-of-flight measurements by the muon system. No evidence for an excess over the expected background is observed. Limits are derived for pair production of top squarks as a function of mass, depending on the interaction model, see Fig. 4 and Table 7. NODE=S046SSQ;LINKAGE=F
- 7 AAD 15AE searched in 19.1 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for heavy long-lived charged particles, measured through their specific ionization energy loss in the ATLAS pixel detector or their time-of-flight in the ATLAS muon system. In the absence of an excess of events above the expected backgrounds, limits are set on R -hadrons in various scenarios, see Fig. 11. Limits are also set in LeptoSUSY models where the gluino decays to stable 300 GeV leptons, see Fig. 9. NODE=S046SSQ;LINKAGE=C
- 8 AAD 15BM searched in 18.4 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for stable and metastable non-relativistic charged particles through their anomalous specific ionization energy loss in the ATLAS pixel detector. In absence of an excess of events above the expected backgrounds, limits are set on stable bottom and top squark R -hadrons, see Table 5. NODE=S046SSQ;LINKAGE=E
- 9 KHACHATRYAN 15AK looked in a data set corresponding to fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$, and a search interval corresponding to 281 h of trigger lifetime, for long-lived particles that have stopped in the CMS detector. No evidence for an excess over the expected background in a cloud interaction model is observed. Assuming the decay $\tilde{t} \rightarrow t \tilde{\chi}_1^0$ and lifetimes between $1 \mu\text{s}$ and 1000 s, limits are derived on \tilde{t} production as a function of $m_{\tilde{\chi}_1^0}$, see Figs. 4 and 7. The exclusions require that $m_{\tilde{\chi}_1^0}$ is kinematically consistent with the minimum values of the jet energy thresholds used. NODE=S046SSQ;LINKAGE=A
- 10 AAD 13AA searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing colored long-lived particles that hadronize forming R -hadrons. No significant excess above the expected background was found. Long-lived R -hadrons containing a \tilde{t} are excluded for masses up to 683 GeV at 95% C.L in a general interaction model. Also, limits independent of the fraction of R -hadrons that arrive charged in the muon system were derived, see Fig. 6. NODE=S046SSQ;LINKAGE=D1
- 11 AAD 13AA searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing colored long-lived particles that hadronize forming R -hadrons. No significant excess above the expected background was found. Long-lived R -hadrons containing a \tilde{b} are excluded for masses up to 612 GeV at 95% C.L in a general interaction model. Also, limits independent of the fraction of R -hadrons that arrive charged in the muon system were derived, see Fig. 6. NODE=S046SSQ;LINKAGE=D2
- 12 AAD 13BC searched in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and in 22.9 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for bottom squark R -hadrons that have come to rest within the ATLAS calorimeter and decay at some later time to hadronic jets and a neutralino. In absence of an excess of events above the expected backgrounds, limits are set on sbottom masses for the decay $\tilde{b} \rightarrow b \tilde{\chi}_1^0$, for different lifetimes, and for a neutralino mass of 100 GeV, see their Table 6 and Fig 10. NODE=S046SSQ;LINKAGE=DD
- 13 AAD 13BC searched in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and in 22.9 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for bottom squark R -hadrons that have come to rest within the ATLAS calorimeter and decay at some later time to hadronic jets and a neutralino. In absence of an excess of events above the expected backgrounds, limits are set on stop masses for the decay $\tilde{t} \rightarrow t \tilde{\chi}_1^0$, for different lifetimes, and for a neutralino mass of 100 GeV, see their Table 6 and Fig 10. NODE=S046SSQ;LINKAGE=DA
- 14 CHATRCHYAN 13AB looked in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and in 18.8 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or additionally requiring that it be identified as muon in the muon chambers, from pair production of \tilde{t}_1 's. No evidence for an excess over the expected background is observed. Limits are derived for pair production of stops as a function of mass in the cloud interaction model (see Fig. 8 and Table 6). In the charge-suppressed model, the limit decreases to 818 GeV. NODE=S046SSQ;LINKAGE=CH

\tilde{b} (Sbottom) mass limit

NODE=S046228

Limits in e^+e^- depend on the mixing angle of the mass eigenstate $\tilde{b}_1 = \tilde{b}_L \cos\theta_b + \tilde{b}_R \sin\theta_b$. Coupling to the Z vanishes for $\theta_b \sim 1.17$. As a consequence, no absolute constraint in the mass region $\lesssim 40 \text{ GeV}$ is available in the literature at this time from e^+e^- collisions. In the Listings below, we use $\Delta m = m_{\tilde{b}_1} - m_{\tilde{\chi}_1^0}$.

NODE=S046228

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

R-parity conserving \tilde{b} (Sbottom) mass limitNODE=S046SBT
NODE=S046SBT;CHECK LIMITS

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	
> 850	95	1 AAD	21AMATLS	τ^\pm 's + b -jets + \cancel{E}_T , Tsb04, $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 130$ GeV, $m_{\tilde{\chi}_2^0} < 180$ GeV	
>1270	95	2 AAD	21S ATLS	b -jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 660	95	2 AAD	21S ATLS	b -jets + \cancel{E}_T , Tsb01, $m_{\tilde{b}_1} - m_{\tilde{\chi}_1^0}$ $= 10$ GeV	OCCUR=2
>1600	95	3 SIRUNYAN	21M CMS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tsb03, $m_{\tilde{\chi}_2^0} = 1500$ GeV, $m_{\tilde{\chi}_1^0} = 100$ GeV	
> 750	95	4 AAD	20V ATLS	same-sign $\ell^\pm \ell^\pm$ + jets, Tsb02, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 100$ GeV, $m_{\tilde{\chi}_1^0} \sim 50$ GeV	
> 850	95	5 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tsb02, $m_{\tilde{\chi}_1^\pm} < 800$ GeV, $m_{\tilde{\chi}_1^0}$ $= 50$ GeV	
>1500	95	6 AAD	19H ATLS	≥ 3 b -jets + \cancel{E}_T , Tsb04, ≥ 1 $h(\rightarrow b\bar{b})$, $m_{\tilde{\chi}_1^0} = 60$ GeV	
>1300	95	7 AAD	19H ATLS	≥ 3 b -jets + \cancel{E}_T , Tsb04, $\geq 1h(\rightarrow$ $b\bar{b})$, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 130$ GeV	OCCUR=2
>1220	95	8 SIRUNYAN	19CH CMS	jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 530	95	9 SIRUNYAN	19CI CMS	≥ 1 $H(\rightarrow \gamma\gamma)$ + jets + \cancel{E}_T , Ts- bot4, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 130$ GeV, $m_{\tilde{\chi}_1^0} = 1$ GeV	
> 430	95	10 AABOUD	18I ATLS	≥ 1 jets + \cancel{E}_T , Tsb01, $m_{\tilde{b}} -$ $m_{\tilde{\chi}_1^0} \sim m_b$	
> 840	95	11 SIRUNYAN	18AL CMS	$\geq 3\ell^\pm$ + jets + \cancel{E}_T , Tsb02, $m_{\tilde{\chi}_1^0}$ $= 50$ GeV	
> 975	95	12 SIRUNYAN	18AR CMS	$\ell^\pm \ell^\mp$ + jets + \cancel{E}_T , Tsb03, $m_{\tilde{\ell}} =$ $(m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 100$ GeV	
>1060	95	13 SIRUNYAN	18AY CMS	jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>1230	95	14 SIRUNYAN	18B CMS	jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 420	95	15 SIRUNYAN	18X CMS	≥ 1 $H(\rightarrow \gamma\gamma)$ + jets + \cancel{E}_T , Ts- bot4, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 130$ GeV, $m_{\tilde{\chi}_1^0} < 225$ GeV	
> 700	95	16 AABOUD	17AJ ATLS	same-sign $\ell^\pm \ell^\pm$ / 3ℓ + jets + \cancel{E}_T , Tsb02, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 950	95	17 AABOUD	17AX ATLS	2 b -jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 880	95	18 AABOUD	17AX ATLS	2 b -jets + \cancel{E}_T , mixture Tsb01 and Tsb02 BR=50%, $m_{\tilde{\chi}_1^0} =$ 0 GeV, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=2
> 315	95	19 KHACHATRY...17A	CMS	2 VBF jets + \cancel{E}_T , Tsb01, $m_{\tilde{b}} -$ $m_{\tilde{\chi}_1^0} = 5$ GeV	
> 450	95	20 KHACHATRY...17AW	CMS	$\geq 3\ell^\pm$, 2 jets, Tsb02, $m_{\tilde{\chi}_1^0} = 50$ GeV, $m_{\tilde{\chi}_1^\pm} = 200$ GeV	
> 800	95	21 KHACHATRY...17P	CMS	1 or more jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0}$ $= 0$ GeV	
>1175	95	22 SIRUNYAN	17AZ CMS	≥ 1 jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 890	95	23 SIRUNYAN	17K CMS	jets + \cancel{E}_T , Tsb01, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 810	95	24 SIRUNYAN	17S CMS	same-sign $\ell^\pm \ell^\pm$ + jets + \cancel{E}_T , Ts- bot2, $m_{\tilde{\chi}_1^0} = 50$ GeV, $m_{\tilde{\chi}_1^\pm} =$ 100 GeV	

> 323	95	25	AABOUD	16D	ATLS	$\geq 1 \text{ jet} + \cancel{E}_T, \text{Tsbot1}, m_{\tilde{b}_1} - m_{\tilde{\chi}_1^0}$ $= 5 \text{ GeV}$	
> 840	95	26	AABOUD	16Q	ATLS	$2 \text{ } b\text{-jets} + \cancel{E}_T, \text{Tsbot1}, m_{\tilde{\chi}_1^0} = 100$ GeV	
> 540	95	27	AAD	16BB	ATLS	$2 \text{ same-sign}/3\ell + \text{jets} + \cancel{E}_T, \text{Tsbot2}, m_{\tilde{\chi}_1^0} < 55 \text{ GeV}$	
> 680	95	28	KHACHATRY...16BJ	CMS		same-sign $\ell^\pm \ell^\pm, \text{Tsbot2}, m_{\tilde{\chi}_1^\pm} <$ $550 \text{ GeV}, m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$	
> 500	95	28	KHACHATRY...16BJ	CMS		same-sign $\ell^\pm \ell^\pm, \text{Tsbot2}, m_{\tilde{b}} -$ $m_{\tilde{\chi}_1^\pm} < 100 \text{ GeV}, m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$	OCCUR=2
> 880	95	29	KHACHATRY...16BS	CMS		jets + $\cancel{E}_T, \text{Tsbot1}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 550	95	30	KHACHATRY...16BY	CMS		opposite-sign $\ell^\pm \ell^\pm, \text{Tsbot3}, m_{\tilde{\chi}_1^0}$ $= 100 \text{ GeV}$	
> 600	95	31	AAD	15CJ	ATLS	$\tilde{b} \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 250 \text{ GeV}$	
> 440	95	31	AAD	15CJ	ATLS	$\tilde{b} \rightarrow t\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^{(*)}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0}$ $= 60 \text{ GeV}, m_{\tilde{b}} - m_{\tilde{\chi}_1^\pm} < m_t$	OCCUR=2
none 300–650	95	31	AAD	15CJ	ATLS	$\tilde{b} \rightarrow \tilde{b}b\tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow h\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} =$ $60 \text{ GeV}, m_{\tilde{\chi}_2^0} > 250 \text{ GeV}$	OCCUR=3
> 640	95	32	KHACHATRY...15AF	CMS		$\tilde{b} \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0$	
> 650	95	33	KHACHATRY...15AH	CMS		$\tilde{b} \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0$	
> 250	95	33	KHACHATRY...15AH	CMS		$\tilde{b} \rightarrow b\tilde{\chi}_1^0, m_{\tilde{b}} - m_{\tilde{\chi}_1^0} < 10 \text{ GeV}$	OCCUR=2
> 570	95	34	KHACHATRY...15I	CMS		$\tilde{b} \rightarrow t\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0}$ $= 50 \text{ GeV}, 150 < m_{\tilde{\chi}_1^\pm} < 300 \text{ GeV}$	
> 255	95	35	AAD	14T	ATLS	$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, m_{\tilde{b}_1} - m_{\tilde{\chi}_1^0} \approx m_b$	
> 400	95	36	CHATRCHYAN14AH	CMS		jets + $\cancel{E}_T, \tilde{b} \rightarrow b\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$	
		37	CHATRCHYAN14R	CMS		$\geq 3\ell^\pm, \tilde{b} \rightarrow t\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow$ $W^\pm\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0}$ $= 50 \text{ GeV}$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●							
		38	KHACHATRY...15AD	CMS		$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T, \tilde{b} \rightarrow$ $b\ell^\pm \ell^\mp \tilde{\chi}_1^0$	
none 340–600	95	39	AAD	14AX	ATLS	$\geq 3 \text{ } b\text{-jets} + \cancel{E}_T, \tilde{b} \rightarrow b\tilde{\chi}_2^0$ sim- plified model with $\tilde{\chi}_2^0 \rightarrow h\tilde{\chi}_1^0,$ $m_{\tilde{\chi}_1^0} = 60 \text{ GeV}, m_{\tilde{\chi}_2^0} = 300 \text{ GeV}$	
> 440	95	40	AAD	14E	ATLS	$\ell^\pm \ell^\pm (\ell^\mp) + \text{jets}, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^{(*)}\tilde{\chi}_1^0$ sim- plified model, $m_{\tilde{\chi}_1^\pm} = 2 m_{\tilde{\chi}_1^0}$	
> 500	95	41	CHATRCHYAN14H	CMS		same-sign $\ell^\pm \ell^\pm, \tilde{b} \rightarrow t\tilde{\chi}_1^\pm,$ $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^\pm} = 2 \text{ GeV}, m_{\tilde{\chi}_1^0} =$ 100 GeV	
> 620	95	42	AAD	13AU	ATLS	$2 \text{ } b\text{-jets} + \cancel{E}_T, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} <$ 120 GeV	
> 550	95	43	CHATRCHYAN13AT	CMS		jets + $\cancel{E}_T, \tilde{b} \rightarrow b\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$	
> 600	95	44	CHATRCHYAN13T	CMS		jets + $\cancel{E}_T, \tilde{b} \rightarrow b\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 450	95	45	CHATRCHYAN13V	CMS		same-sign $\ell^\pm \ell^\pm + \geq 2 \text{ } b\text{-jets},$ $\tilde{b} \rightarrow t\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$ sim- plified model, $m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$	
> 390		46	AAD	12AN	ATLS	$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, \text{simplified model},$ $m_{\tilde{\chi}_1^0} < 60 \text{ GeV}$	

		47	CHATRCHYAN12AI	CMS	$\ell^\pm \ell^\pm + b\text{-jets} + \cancel{E}_T$
> 410	95	48	CHATRCHYAN12BO	CMS	$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$, simplified model, $m_{\tilde{\chi}_1^0} = 50$ GeV
> 294	95	49	AAD	11K ATLS	stable b
		50	AAD	11O ATLS	$\tilde{g} \rightarrow \tilde{b}_1 b, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 60$ GeV
> 230	95	51	CHATRCHYAN11D	CMS	$\tilde{b}, \tilde{t} \rightarrow b$
		52	AALTONEN	10R CDF	$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 70$ GeV
> 247	95	53	ABAZOV	10L D0	$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0$ GeV

- ¹ AAD 21AM searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pair production of bottom squarks in events with hadronically decaying τ^\pm -leptons, b -tagged jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the bottom squark mass in the Tsb04 simplified model, assuming $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 130$ GeV, see their Figure 8. NODE=S046SBT;LINKAGE=ZA
- ² AAD 21S searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pair production of sbottoms, LQ or dark matter in events with b -jets and \cancel{E}_T , also using dedicated secondary-vertex-finding techniques. No significant excess above the Standard Model predictions is observed. Limits are set on $m_{\tilde{b}_1}$ in the Tsb01 simplified model, on the LQ masses depending on the BR in $b\nu$, on scalar and pseudoscalar dark matter mediator masses. See Figures 8, 9, 10. NODE=S046SBT;LINKAGE=BB
- ³ SIRUNYAN 21M searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tsb03, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14. NODE=S046SBT;LINKAGE=CB
- ⁴ AAD 20V searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with two same-sign charged leptons (electrons or muons) and jets. No significant excess above the Standard Model expectations is observed. Exclusion limits at 95% C.L. are set on the bottom squark masses in the Tsb02 simplified model for $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 100$ GeV, see their Fig. 8(a). NODE=S046SBT;LINKAGE=XA
- ⁵ SIRUNYAN 20T searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least two jets, and two isolated same-sign or three or more charged leptons (electrons or muons). No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figure 7, and in the Tglu1C and Tglu1B simplified models, see their Figures 8 and 9. Limits are also set on the sbottom mass in the Tsb02 simplified model, see their Figure 10, and on the stop mass in the Tstop7 simplified model, see their Figure 11. Finally, limits are set on the gluino mass in RPV simplified models where the gluino decays either via $\tilde{g} \rightarrow qq\bar{q} + e/\mu/\tau$ or via $\tilde{g} \rightarrow tbs$, see Figure 12. NODE=S046SBT;LINKAGE=WA
- ⁶ AAD 19H searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with no charged leptons, three or more b -jets, and large \cancel{E}_T . Higgs boson candidates are reconstructed as b -jet pairs. No significant excess above the Standard Model expectations is observed. Limits up to 1500 GeV are set on the sbottom mass in the Tsb04 simplified model, see Figure 8(a), for fixed $m_{\tilde{\chi}_1^0} = 60$ GeV and for $m_{\tilde{\chi}_2^0}$ up to 1200 GeV. NODE=S046SBT;LINKAGE=TA
- ⁷ AAD 19H searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with no charged leptons, three or more b -jets, and large \cancel{E}_T . Higgs boson candidates are reconstructed as b -jet pairs. No significant excess above the Standard Model expectations is observed. Limits up to 1300 GeV are set on the sbottom mass in the Tsb04 simplified model, see Figure 8(b), for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 130$ GeV and $m_{\tilde{\chi}_2^0}$ from 200 to 750 GeV. NODE=S046SBT;LINKAGE=UA
- ⁸ SIRUNYAN 19CH searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A and Tglu3A simplified models, see their Figure 13. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tsb01, Tstop1 simplified models, see their Figure 14. NODE=S046SBT;LINKAGE=RA
- ⁹ SIRUNYAN 19Ci searched in 77.5 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with one or more high-momentum Higgs bosons, decaying to pairs of photons, jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb04 simplified model, see Figure 3, and on the wino mass in the Tchi1n2E simplified model, see their Figure 4. Limits are also set on the higgsino mass in the Tn1n1A and Tn1n1B simplified models, see their Figure 5. NODE=S046SBT;LINKAGE=QA
- ¹⁰ AABOUD 18I searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one jet with a transverse momentum above 250 GeV and no leptons. Good agreement is observed between the number of events in data and Standard Model predictions. NODE=S046SBT;LINKAGE=MA

The results are translated into exclusion limits in Tsb0t1 models. In the compressed scenario with sbottom and neutralino masses differing by m_b , sbottom masses below 430 GeV are excluded. For $m_{\tilde{\chi}_1^0} = 0$ they exclude sbottom masses up to 610 GeV. See their Fig.10(a).

- 11 SIRUNYAN 18AL searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least three charged leptons, in any combination of electrons and muons, jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu1C simplified models, see their Figure 5. Limits are also set on the sbottom mass in the Tsb0t2 simplified model, see their Figure 6, and on the stop mass in the Tstop7 simplified model, see their Figure 7. NODE=S046SBT;LINKAGE=NA
- 12 SIRUNYAN 18AR searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two opposite-charge, same-flavour leptons (electrons or muons), jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4C simplified model, see their Figure 7. Limits are also set on the chargino/neutralino mass in the Tchi1n2F simplified model, see their Figure 8, and on the neutralino mass in the Tn1n1B and Tn1n1C simplified models, see their Figure 9. Finally, limits are set on the sbottom mass in the Tsb0t3 simplified model, see their Figure 10. NODE=S046SBT;LINKAGE=OA
- 13 SIRUNYAN 18AY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing one or more jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A and Tglu3A simplified models, see their Figure 3. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tsb0t1, Tstop1 and Tstop4 simplified models, see their Figure 3. Finally, limits are set on long-lived gluino masses in a Tglu1A simplified model where the gluino is metastable or long-lived with proper decay lengths in the range $10^{-3} \text{ mm} < c\tau < 10^5 \text{ mm}$, see their Figure 4. NODE=S046SBT;LINKAGE=PA
- 14 SIRUNYAN 18B searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for the pair production of third-generation squarks in events with jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t1 simplified model, see their Figure 5, and on the stop mass in the Tstop4 simplified model, see their Figure 6. NODE=S046SBT;LINKAGE=BA
- 15 SIRUNYAN 18X searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more high-momentum Higgs bosons, decaying to pairs of photons, jets and \cancel{E}_T . The razor variables (M_R and R^2) are used to categorise the events. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t4 simplified model and on the wino mass in the Tchi1n2E simplified model, see their Figure 5. Limits are also set on the higgsino mass in the Tn1n1A and Tn1n1B simplified models, see their Figure 6. NODE=S046SBT;LINKAGE=KA
- 16 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 700 GeV are set on the bottom squark mass in Tsb0t2 simplified models assuming $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$. See their Figure 4(d). NODE=S046SBT;LINKAGE=IA
- 17 AABOUD 17AX searched in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two jets identified as originating from b -quarks and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of bottom squarks. In the Tsb0t1 simplified model, a \tilde{b}_1 mass below 950 GeV is excluded for $m_{\tilde{\chi}_1^0} = 0$ (<420) GeV. See their Fig. 7(a). NODE=S046SBT;LINKAGE=GA
- 18 AABOUD 17AX searched in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two jets identified as originating from b -quarks and large missing transverse momentum, with or without leptons. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of bottom squarks. Assuming 50% BR for Tsb0t1 and Tsb0t2 simplified models, a \tilde{b}_1 mass below 880 (860) GeV is excluded for $m_{\tilde{\chi}_1^0} = 0$ (<250) GeV. See their Fig. 7(b). NODE=S046SBT;LINKAGE=HA
- 19 KHACHATRYAN 17A searched in 18.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with two forward jets, produced through vector boson fusion, and missing transverse momentum. No significant excess above the Standard Model expectations is observed. A limit is set on sbottom masses in the Tsb0t1 simplified model, see Fig. 3. NODE=S046SBT;LINKAGE=S
- 20 KHACHATRYAN 17AW searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least three charged leptons, in any combination of electrons and muons, and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu1C simplified models, and on the sbottom mass in the Tsb0t2 simplified model, see their Figure 4. NODE=S046SBT;LINKAGE=Z
- 21 KHACHATRYAN 17P searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figures 7 and 8. Limits are also set on the squark mass in the Tsqk1 simplified model, see their Fig. 7, and on the sbottom mass in the Tsb0t1 simplified model, see Fig. 8. Finally, limits are set on the stop mass in the Tstop1, Tstop3, Tstop4, Tstop6 and Tstop7 simplified models, see Fig. 8. NODE=S046SBT;LINKAGE=W
- 22 SIRUNYAN 17AZ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model NODE=S046SBT;LINKAGE=X

- expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A simplified models, see their Figures 6. Limits are also set on the squark mass in the Tsqk1 simplified model (for single light squark and for 8 degenerate light squarks), on the sbottom mass in the Tsb0t1 simplified model and on the stop mass in the Tstop1 simplified model, see their Fig. 7. Finally, limits are set on the stop mass in the Tstop2, Tstop4 and Tstop8 simplified models, see Fig. 8.
- 23 SIRUNYAN 17K searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct production of stop or sbottom pairs in events with multiple jets and significant \cancel{E}_T . A second search also requires an isolated lepton and is combined with the all-hadronic search. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1, Tstop8 and Tstop4 simplified models, see their Figures 7, 8 and 9 (for the Tstop4 limits, only the results of the all-hadronic search are used). Limits are also set on the sbottom mass in the Tsb0t1 simplified model, see Fig. 10 (also here, only the results of the all-hadronic search are used). NODE=S046SBT;LINKAGE=V
- 24 SIRUNYAN 17S searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two isolated same-sign leptons, jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the gluino mass in the Tglu3A, Tglu3B, Tglu3C, Tglu3D and Tglu1B simplified models, see their Figures 5 and 6, and on the sbottom mass in the Tsb0t2 simplified model, see their Figure 6. NODE=S046SBT;LINKAGE=Y
- 25 AABOUD 16D searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with an energetic jet and large missing transverse momentum. The results are interpreted as 95% C.L. limits on mass of sbottom decaying into a b -quark and the lightest neutralino in scenarios with $m_{\tilde{b}_1} - m_{\tilde{\chi}_1^0}$ between 5 and 20 GeV. See their Fig. 6. NODE=S046SBT;LINKAGE=J
- 26 AABOUD 16Q searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two jets identified as originating from b -quarks and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of third-generation squarks. Assuming that the decay $\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$ (Tsb0t1) takes place 100% of the time, a \tilde{b}_1 mass below 840 (800) GeV is excluded for $m_{\tilde{\chi}_1^0} < 100$ (360) GeV. Differences in mass above 100 GeV between the \tilde{b}_1 and the $\tilde{\chi}_1^0$ are excluded up to a \tilde{b}_1 mass of 500 GeV. For more details, see their Fig. 4. NODE=S046SBT;LINKAGE=U
- 27 AAD 16BB searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with exactly two same-sign leptons or at least three leptons, multiple hadronic jets, b -jets, and \cancel{E}_T . No significant excess over the Standard Model expectation is found. Exclusion limits at 95% C.L. are set on the sbottom mass for the Tsb0t2 model, assuming $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 100 \text{ GeV}$. See their Fig. 4c. NODE=S046SBT;LINKAGE=N
- 28 KHACHATRYAN 16BJ searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t2 simplified model, see Fig. 6. NODE=S046SBT;LINKAGE=O
- 29 KHACHATRYAN 16BS searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one energetic jet, no isolated leptons, and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t1 simplified model, see Fig. 11 and Table 3. NODE=S046SBT;LINKAGE=P
- 30 KHACHATRYAN 16BY searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two opposite-sign, same-flavour leptons, jets, and missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4C simplified model, see Fig. 4, and on sbottom masses in the Tsb0t3 simplified model, see Fig. 5. NODE=S046SBT;LINKAGE=R
- 31 AAD 15CJ searched in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of third generation squarks by combining a large number of searches covering various final states. Limits on the sbottom mass are shown, either assuming the $\tilde{b} \rightarrow b\tilde{\chi}_1^0$ decay, see Fig. 11, or assuming the $\tilde{b} \rightarrow t\tilde{\chi}_1^\pm$ decay, with $\tilde{\chi}_1^\pm \rightarrow W^{(*)}\tilde{\chi}_1^0$, see Fig. 12a, or assuming the $\tilde{b} \rightarrow b\tilde{\chi}_2^0$ decay, with $\tilde{\chi}_2^0 \rightarrow h\tilde{\chi}_1^0$, see Fig. 12b. Interpretations in the pMSSM are also discussed, see Figures 13–15. NODE=S046SBT;LINKAGE=M
- 32 KHACHATRYAN 15AF searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in simplified models where the decay $\tilde{b} \rightarrow b\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 12. See also Table 5. Exclusions in the CMSSM, assuming $\tan\beta = 30$, $A_0 = -2 \max(m_0, m_{1/2})$ and $\mu > 0$, are also presented, see Fig. 15. NODE=S046SBT;LINKAGE=G
- 33 KHACHATRYAN 15AH searched in 19.4 or 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing either a fully reconstructed top quark, or events containing dijets requiring one or both jets to originate from b -quarks, or events containing a mono-jet. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in simplified models where the decay $\tilde{b} \rightarrow b\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 12. Limits are also set in a simplified model where the decay $\tilde{b} \rightarrow c\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 12. NODE=S046SBT;LINKAGE=H
- 34 KHACHATRYAN 15I searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events in which b -jets and four W -bosons are produced. Five individual search channels are NODE=S046SBT;LINKAGE=L

- combined (fully hadronic, single lepton, same-sign dilepton, opposite-sign dilepton, multilepton). No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in a simplified model where the decay $\tilde{b} \rightarrow t\tilde{\chi}_1^\pm$, with $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$, takes place with a branching ratio of 100%, see Fig. 7.
- 35 AAD 14T searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for monojet-like events. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of third-generation squarks in simplified models which assume that the decay $\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$ takes place 100% of the time, see Fig. 12. NODE=S046SBT;LINKAGE=B
- 36 CHATRCHYAN 14AH searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the razor variables (M_R and R^2) to discriminate between signal and background processes. A second analysis requires at least one of the jets to be originating from a b -quark. No significant excess above the Standard Model expectations is observed. Limits are set on sbottom masses in simplified models where the decay $\tilde{b} \rightarrow b\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Figs. 28 and 29. Exclusions in the CMSSM, assuming $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, are also presented, see Fig. 26. NODE=S046SBT;LINKAGE=E
- 37 CHATRCHYAN 14R searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least three leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in a simplified model where the decay $\tilde{b} \rightarrow t\tilde{\chi}_1^\pm$, with $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$, takes place with a branching ratio of 100%, see Fig. 11. NODE=S046SBT;LINKAGE=A
- 38 KHACHATRYAN 15AD searched in 19.4 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with two opposite-sign same flavor isolated leptons featuring either a kinematic edge, or a peak at the Z -boson mass, in the invariant mass spectrum. No evidence for a statistically significant excess over the expected SM backgrounds is observed and 95% C.L. exclusion limits are derived in a simplified model of sbottom pair production where the sbottom decays into a b -quark, two opposite-sign dileptons and a neutralino LSP, through an intermediate state containing either an off-shell Z -boson or a slepton, see Fig. 8. NODE=S046SBT;LINKAGE=F
- 39 AAD 14AX searched in 20.1 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for the strong production of supersymmetric particles in events containing either zero or at least one high p_T lepton, large missing transverse momentum, high jet multiplicity and at least three jets identified as originating from b -quarks. No excess over the expected SM background is observed. Limits are derived in mSUGRA/CMSSM models with $\tan\beta = 30$, $A_0 = -2 m_0$ and $\mu > 0$, see their Fig. 14. Also, exclusion limits are set in simplified models containing scalar bottom quarks, where the decay $\tilde{b} \rightarrow b\tilde{\chi}_2^0$ and $\tilde{\chi}_2^0 \rightarrow h\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see their Figures 11. NODE=S046SBT;LINKAGE=DA
- 40 AAD 14E searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for strongly produced supersymmetric particles in events containing jets and two same-sign leptons or three leptons. The search also utilises jets originating from b -quarks, missing transverse momentum and other variables. No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing bottom, see Fig. 7. Limits are also derived in the mSUGRA/CMSSM, bRPV and GMSB models, see their Fig. 8. NODE=S046SBT;LINKAGE=D
- 41 CHATRCHYAN 14H searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in a simplified models where the decay $\tilde{b} \rightarrow t\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, with varying mass of the $\tilde{\chi}_1^\pm$, for $m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$, see Fig. 6. NODE=S046SBT;LINKAGE=C
- 42 AAD 13AU searched in 20.1 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing two jets identified as originating from b -quarks and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of third-generation squarks. Assuming that the decay $\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$ takes place 100% of the time, a \tilde{b}_1 mass below 620 GeV is excluded for $m_{\tilde{\chi}_1^0} < 120 \text{ GeV}$. For more details, see their Fig. 5. NODE=S046SBT;LINKAGE=DT
- 43 CHATRCHYAN 13AT provides interpretations of various searches for supersymmetry by the CMS experiment based on $4.73\text{--}4.98 \text{ fb}^{-1}$ of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ in the framework of simplified models. Limits are set on the sbottom mass in a simplified models where sbottom quarks are pair-produced and the decay $\tilde{b} \rightarrow b\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 4. NODE=S046SBT;LINKAGE=YA
- 44 CHATRCHYAN 13T searched in 11.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the α_T variable to discriminate between processes with genuine and misreconstructed \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on sbottom masses in simplified models where the decay $\tilde{b} \rightarrow b\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 8 and Table 9. NODE=S046SBT;LINKAGE=CS
- 45 CHATRCHYAN 13V searched in 10.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with two isolated same-sign dileptons and at least two b -jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the bottom mass in a simplified models where the decay $\tilde{b} \rightarrow t\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$ takes place

with a branching ratio of 100%, with varying mass of the $\tilde{\chi}_1^\pm$, for $m_{\tilde{\chi}_1^0} = 50$ GeV, see

Fig. 4.

- 46 AAD 12AN searched in 2.05 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for scalar bottom quarks in events with large missing transverse momentum and two b -jets in the final state. The data are found to be consistent with the Standard Model expectations. Limits are set in an R-parity conserving minimal supersymmetric scenario, assuming $B(\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0) = 100\%$, see their Fig. 2. NODE=S046SBT;LINKAGE=DG
- 47 CHATRCHYAN 12AI looked in 4.98 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with two same-sign leptons (e, μ), but not necessarily same flavor, at least 2 b -jets and missing transverse energy. No excess beyond the Standard Model expectation is observed. Exclusion limits are derived in a simplified model for sbottom pair production, where the sbottom decays through $\tilde{b}_1 \rightarrow t\tilde{\chi}_1 W$, see Fig. 8. NODE=S046SBT;LINKAGE=TR
- 48 CHATRCHYAN 12BO searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for scalar bottom quarks in events with large missing transverse momentum and two b -jets in the final state. The data are found to be consistent with the Standard Model expectations. Limits are set in an R-parity conserving minimal supersymmetric scenario, assuming $B(\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0) = 100\%$, see their Fig. 2. NODE=S046SBT;LINKAGE=CH
- 49 AAD 11K looked in 34 pb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or time of flight in the tile calorimeter, from pair production of \tilde{b} . No evidence for an excess over the SM expectation is observed and limits on the mass are derived for pair production of sbottom, see Fig. 4. NODE=S046SBT;LINKAGE=A2
- 50 AAD 11O looked in 35 pb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with jets, of which at least one is a b -jet, and \cancel{E}_T . No excess above the Standard Model was found. Limits are derived in the $(m_{\tilde{g}}, m_{\tilde{b}_1})$ plane (see Fig. 2) under the assumption of 100% branching ratios and \tilde{b}_1 being the lightest squark. The quoted limit is valid for $m_{\tilde{b}_1} < 500$ GeV. A similar approach for \tilde{t}_1 as the lightest squark with $\tilde{g} \rightarrow \tilde{t}_1 t$ and $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ with 100% branching ratios leads to a gluino mass limit of 520 GeV for $130 < m_{\tilde{t}_1} < 300$ GeV. Limits are also derived in the CMSSM $(m_0, m_{1/2})$ plane for $\tan\beta = 40$, see Fig. 4, and in scenarios based on the gauge group $SO(10)$. NODE=S046SBT;LINKAGE=A6
- 51 CHATRCHYAN 11D looked in 35 pb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with ≥ 2 jets, at least one of which is b -tagged, and \cancel{E}_T , where the b -jets are decay products of \tilde{t} or \tilde{b} . No evidence for an excess over the expected background is observed. Limits are derived in the CMSSM $(m_0, m_{1/2})$ plane for $\tan\beta = 50$ (see Fig. 2). NODE=S046SBT;LINKAGE=C1
- 52 AALTONEN 10R searched in 2.65 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV for events with \cancel{E}_T and exactly two jets, at least one of which is b -tagged. The results are in agreement with the SM prediction, and a limit on the cross section of 0.1 pb is obtained for the range of masses $80 < m_{\tilde{b}_1} < 280$ GeV assuming that the sbottom decays exclusively to $b\tilde{\chi}_1^0$. The excluded mass region in the framework of conserved R_p is shown in a plane of $(m_{\tilde{b}_1}, m_{\tilde{\chi}_1^0})$, see their Fig.2. NODE=S046SBT;LINKAGE=LA
- 53 ABAZOV 10L looked in 5.2 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV for events with at least 2 b -jets and \cancel{E}_T from the production of $\tilde{b}_1\tilde{b}_1$. No evidence for an excess over the SM expectation is observed, and a limit on the cross section is derived under the assumption of 100% branching ratio. The excluded mass region in the framework of conserved R_p is shown in a plane of $(m_{\tilde{b}_1}, m_{\tilde{\chi}_1^0})$, see their Fig. 3b. The exclusion also extends to $m_{\tilde{\chi}_1^0} = 110$ GeV for $160 < m_{\tilde{b}_1} < 200$ GeV. NODE=S046SBT;LINKAGE=AB

R-parity violating \tilde{b} (Sbottom) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>307	95	¹ KHACHATRY...16BX CMS		RPV, $\tilde{b} \rightarrow td$ or ts , λ''_{332} or λ''_{331} coupling

• • • We do not use the following data for averages, fits, limits, etc. • • •

² AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell\bar{\ell}) + \text{jets}, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_1^0$ sim- plified model, $m_{\tilde{\chi}_1^\pm} = 2 m_{\tilde{\chi}_1^0}$
------------------	----------	---

¹ KHACHATRYAN 16BX searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing 2 leptons coming from R-parity-violating decays of supersymmetric particles. No excess over the expected background is observed. Limits are derived on the sbottom mass, assuming the RPV $\tilde{b} \rightarrow td$ or $\tilde{b} \rightarrow ts$ decay, see Fig. 15. NODE=S046SBV;LINKAGE=T

² AAD 14E searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for strongly produced supersymmetric particles in events containing jets and two same-sign leptons or three leptons. The search also utilises jets originating from b -quarks, missing transverse momentum and other variables. No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing bottom, see Fig. 7. Limits are also derived in the mSUGRA/CMSSM, bRPV and GMSB models, see their Fig. 8. NODE=S046SBV;LINKAGE=D

NODE=S046SBV
NODE=S046SBV

\tilde{t} (Stop) mass limit

NODE=S046227

Limits depend on the decay mode. In e^+e^- collisions they also depend on the mixing angle of the mass eigenstate $\tilde{t}_1 = \tilde{t}_L \cos\theta_t + \tilde{t}_R \sin\theta_t$. The coupling to the Z vanishes when $\theta_t = 0.98$. In the Listings below, we use $\Delta m \equiv m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$ or $\Delta m \equiv m_{\tilde{t}_1} - m_{\tilde{\nu}}$, depending on relevant decay mode. See also bounds in “ \tilde{q} (Squark) MASS LIMIT.”

NODE=S046227

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

R-parity conserving \tilde{t} (Stop) mass limit

NODE=S046STP

NODE=S046STP;CHECK LIMITS

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	
>1430	95	1 HAYRAPETY...23E	CMS	$\gamma + \text{jets} + \cancel{E}_T$, Tstop13, $m_{\tilde{\chi}_1^0}$	
				= 1170 GeV	
>1150	95	2 TUMASYAN 23AB	CMS	$\geq 1 \tau^\pm + \cancel{E}_T$, Tstop16, $m_{\tilde{\chi}_1^0}$	
				= 1 GeV	
> 480	95	3 TUMASYAN 23K	CMS	1 high- p_T jet, 1 low- p_T e or μ , Tstop3, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 10$	
				GeV	
> 700	95	3 TUMASYAN 23K	CMS	1 high- p_T jet, 1 low- p_T e or μ , Tstop3, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 80$	OCCUR=2
				GeV	
> 480	95	4 TUMASYAN 22Q	CMS	2 or 3 ℓ (soft), \cancel{E}_T ; Tstop2, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 30$ GeV	
> 540	95	4 TUMASYAN 22Q	CMS	2 or 3 ℓ (soft), \cancel{E}_T ; Tstop3, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 30$ GeV	OCCUR=2
>1400	95	5 AAD 21AW	ATLS	$\tau^\pm + \text{jets} + b\text{-jets} + \cancel{E}_T$, Tstop5, $m_{\tilde{\tau}_1} = 1200$ GeV	OCCUR=2
>1200	95	6 AAD 21O	ATLS	$\ell^\pm + \text{jet} + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0}$	
				= 0 GeV	
> 710	95	6 AAD 21O	ATLS	$\ell^\pm + \text{jet} + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0}$	OCCUR=2
				= 580 GeV	
> 640	95	6 AAD 21O	ATLS	$\ell^\pm + \text{jet} + \cancel{E}_T$, Tstop3, $m_{\tilde{\chi}_1^0}$	OCCUR=3
				= 580 GeV	
>1000	95	7 AAD 21P	ATLS	$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
				= 580 GeV	
> 600	95	7 AAD 21P	ATLS	$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T$, Tstop2, $m_{\tilde{\chi}_1^0} = 500$ GeV	OCCUR=2
				= 580 GeV	
> 550	95	7 AAD 21P	ATLS	$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T$, Tstop3, $m_{\tilde{\chi}_1^0} = 500$ GeV	OCCUR=3
				= 580 GeV	
>1310	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} < 300$	
				GeV	
>1170	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 100$	OCCUR=2
				GeV	
>1150	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop1 (50%) or Tstop2 (50%), $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5$ GeV, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=3
> 640	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop3, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=4
> 620	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop3, 10 GeV < $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} < 60$ GeV	OCCUR=5
> 740	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop2, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 80$ GeV	OCCUR=6
> 720	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop2, 40 GeV < $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} < 80$ GeV	OCCUR=7
> 595	95	8 SIRUNYAN 21AD	CMS	jets + \cancel{E}_T , Tstop2, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 10$ GeV	OCCUR=8

> 630	95	8 SIRUNYAN	21AD CMS	jets + \cancel{E}_T , Tstop4, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0}$	OCCUR=9
none 200–920	95	9 SIRUNYAN	21B CMS	$\ell^\pm \ell^\mp + b$ -jets + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 250–810		9 SIRUNYAN	21B CMS	$\ell^\pm \ell^\mp + b$ -jets + \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
>1300	95	9 SIRUNYAN	21B CMS	$\ell^\pm \ell^\mp + b$ -jets + \cancel{E}_T , Tstop11, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\ell}} = (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0})/2 + m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
none 400–1180	95	9 SIRUNYAN	21B CMS	$\ell^\pm \ell^\mp + b$ -jets + \cancel{E}_T , Tstop11, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\ell}} = 0.05 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}) + m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=4
>1400	95	9 SIRUNYAN	21B CMS	$\ell^\pm \ell^\mp + b$ -jets + \cancel{E}_T , Tstop11, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\ell}} = 0.95 (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}) + m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=5
>1325	95	10 TUMASYAN	21I CMS	≥ 2 jets + \cancel{E}_T + 0,1,2 ℓ , Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
>1150	95	10 TUMASYAN	21I CMS	≥ 2 jets + \cancel{E}_T + 0,1,2 ℓ , Tstop1, $m_{\tilde{\chi}_1^0} = 700$ GeV	OCCUR=4
>1260	95	10 TUMASYAN	21I CMS	≥ 2 jets + \cancel{E}_T + 0,1,2 ℓ , Tstop2, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
>1000	95	10 TUMASYAN	21I CMS	≥ 2 jets + \cancel{E}_T + 0,1,2 ℓ , Tstop2, $m_{\tilde{\chi}_1^0} < 575$ GeV	OCCUR=6
>1175	95	10 TUMASYAN	21I CMS	≥ 2 jets + \cancel{E}_T + 0,1,2 ℓ , Tstop1 (50%) or Tstop2 (50%), $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=7
>1000	95	10 TUMASYAN	21I CMS	≥ 2 jets + \cancel{E}_T + 0,1,2 ℓ , Tstop1 (50%) or Tstop2 (50%), $m_{\tilde{\chi}_1^0} = 570$ GeV	OCCUR=8
none 145–295	95	10 TUMASYAN	21I CMS	$\ell^\pm \ell^\mp +$ jets + \cancel{E}_T , Tstop1, $ m_{\tilde{t}} - m_{\tilde{\chi}_1^0} - 175 \text{ GeV} < 30 \text{ GeV}$	OCCUR=9
none, 170–230	95	11 AABOUD	20 ATLS	$e^\pm \mu^\mp +$ $\geq 1b$ -jet, Tstop1, $m_{\tilde{\chi}_1^0} = 0.5$ GeV	
none, 170–220	95	11 AABOUD	20 ATLS	$e^\pm \mu^\mp +$ $\geq 1b$ -jet, Tstop1, $m_{\tilde{\chi}_1^0} < 62$ GeV	OCCUR=2
>1220	95	12 AAD	20AS ATLS	$\ell^\pm \ell^\mp$ or 2 b -jets and \cancel{E}_T , Tstop6, $m_{\tilde{\chi}_2^0} = 900$ GeV	
> 860	95	13 AAD	20AS ATLS	$\ell^\pm \ell^\mp$ or 2 b -jets and \cancel{E}_T , \tilde{t}_2 with $\tilde{t}_2 \rightarrow \tilde{t}_1 Z$, $\tilde{t}_1 \rightarrow b f f' \tilde{\chi}_1^0$, $\Delta m(\tilde{t}_1, \tilde{\chi}_1^0) = 40$ GeV	OCCUR=2
none 400–1250	95	14 AAD	20S ATLS	jets + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 300–660	95	15 AAD	20S ATLS	jets + \cancel{E}_T , Tstop3, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 765	95	16 AAD	20V ATLS	same-sign $\ell^\pm \ell^\pm +$ jets, $\tilde{t}_1 \rightarrow t \tilde{\chi}_2^0$, $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^\pm W$, $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 W$, $m_{\tilde{\chi}_1^\pm} \sim m_{\tilde{\chi}_1^0}$	
>1200	95	17 SIRUNYAN	20AH CMS	$\ell^\pm +$ jet + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>1175	95	17 SIRUNYAN	20AH CMS	$\ell^\pm +$ jet + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} < 425$ GeV	OCCUR=2

none 230–1140	95	17 SIRUNYAN	20AH CMS	$\ell^\pm + \text{jet} + \cancel{E}_T, T_{\text{stop}2}, m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2, m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
>1100	95	17 SIRUNYAN	20AH CMS	$\ell^\pm + \text{jet} + \cancel{E}_T, T_{\text{stop}2}, m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2, 50 < m_{\tilde{\chi}_1^0} < 425 \text{ GeV}$	OCCUR=4
>1070	95	17 SIRUNYAN	20AH CMS	$\ell^\pm + \text{jet} + \cancel{E}_T, T_{\text{stop}8}, m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=5
>1050	95	17 SIRUNYAN	20AH CMS	$\ell^\pm + \text{jet} + \cancel{E}_T, T_{\text{stop}8}, m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}, m_{\tilde{\chi}_1^0} < 350 \text{ GeV}$	OCCUR=6
> 730	95	18 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm + \text{jets}, T_{\text{stop}7}, m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}, m_{\tilde{t}_1} = 200 \text{ GeV}, B(\tilde{t}_2 \rightarrow \tilde{t}_1 H) = 100\%$	OCCUR=2
> 890	95	18 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm + \text{jets}, T_{\text{stop}7}, m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}, m_{\tilde{t}_1} = 200 \text{ GeV}, B(\tilde{t}_2 \rightarrow \tilde{t}_1 Z) = 100\%$	OCCUR=3
> 760	95	18 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm + \text{jets}, T_{\text{stop}7}, m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}, m_{\tilde{t}_1} = 200 \text{ GeV}, B(\tilde{t}_2 \rightarrow \tilde{t}_1 Z) = B(\tilde{t}_2 \rightarrow \tilde{t}_1 H) = 50\%$	OCCUR=2
>1100	95	19 SIRUNYAN	20U CMS	$\tau^\pm \tau^\mp + b\text{-jets} + \cancel{E}_T, T_{\text{stop}11}, m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{t}} + m_{\tilde{\chi}_1^0}), m_{\tilde{\tau}} = 0.5 m_{\tilde{\chi}_1^\pm}, m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
>1110	95	20 SIRUNYAN	19AU CMS	$\gamma + \text{jets} + b\text{-jets} + \cancel{E}_T, T_{\text{stop}13}, m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$	OCCUR=2
>1230	95	20 SIRUNYAN	19AU CMS	$\gamma + \text{jets} + b\text{-jets} + \cancel{E}_T, T_{\text{stop}13}, m_{\tilde{\chi}_1^0} = 800 \text{ GeV}$	OCCUR=2
>1190	95	21 SIRUNYAN	19CH CMS	$\text{jets} + \cancel{E}_T, T_{\text{stop}1}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1140	95	22 SIRUNYAN	19S CMS	$1 \text{ or } 2 \ell + \text{jets} + \cancel{E}_T, T_{\text{stop}1}, m_{\tilde{\chi}_1^0} < 200 \text{ GeV}$	OCCUR=2
> 208	95	23 SIRUNYAN	19U CMS	$e^\pm \mu^\mp + \geq 1b\text{-jet}, T_{\text{stop}1}, m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}$	OCCUR=2
> 235	95	23 SIRUNYAN	19U CMS	$e^\pm \mu^\mp + \geq 1b\text{-jet}, T_{\text{stop}1}, m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 182.5 \text{ GeV}$	OCCUR=2
> 242	95	23 SIRUNYAN	19U CMS	$e^\pm \mu^\mp + \geq 1b\text{-jet}, T_{\text{stop}1}, m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 167.5 \text{ GeV}$	OCCUR=3
> 940	95	24 AABOUD	18AQ ATLS	$1\ell + \text{jets} + \cancel{E}_T, T_{\text{stop}1}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
> 270	95	25 AABOUD	18AQ ATLS	$1\ell + \text{jets} + \cancel{E}_T, T_{\text{stop}3}, m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 20 \text{ GeV}$	OCCUR=2
> 840	95	26 AABOUD	18AQ ATLS	$1\ell + \text{jets} + \cancel{E}_T, T_{\text{stop}2}, m_{\tilde{t}} - m_{\tilde{\chi}_1^\pm} = 10 \text{ GeV}$	OCCUR=3
> 500	95	27 AABOUD	18BV ATLS	$c\text{-jets} + \cancel{E}_T, T_{\text{stop}4}, m_{\tilde{t}} - m_{\tilde{\chi}_1^0} < 100 \text{ GeV}$	OCCUR=2
> 850	95	28 AABOUD	18BV ATLS	$c\text{-jets} + \cancel{E}_T, T_{\text{stop}4}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
> 390	95	29 AABOUD	18l ATLS	$\geq 1 \text{ jets} + \cancel{E}_T, T_{\text{stop}3}, m_{\tilde{t}} \sim m_{\tilde{\chi}_1^0}$	OCCUR=2

> 430	95	30	AABOUD	18I	ATLS	≥ 1 jets + \cancel{E}_T , Tstop4, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 5$ GeV	OCCUR=2
>1160	95	31	AABOUD	18Y	ATLS	2ℓ (≥ 1 hadronic τ) + b -jets + \cancel{E}_T , Tstop5, $m_{\tilde{\tau}} \sim 800$ GeV	
> 450	95	32	SIRUNYAN	18AJ	CMS	2ℓ (soft) + \cancel{E}_T , Tstop10, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 40$ GeV	
> 720	95	33	SIRUNYAN	18AL	CMS	$\geq 3\ell^\pm$ + jets + \cancel{E}_T , Tstop7, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175$ GeV, $m_{\tilde{t}_1} = 200$ GeV, BR($\tilde{t}_2 \rightarrow \tilde{t}_1 H$) = 100%	
> 780	95	33	SIRUNYAN	18AL	CMS	$\geq 3\ell^\pm$ + jets + \cancel{E}_T , Tstop7, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175$ GeV, $m_{\tilde{t}_1} = 200$ GeV, BR($\tilde{t}_2 \rightarrow \tilde{t}_1 Z$) = 100%	OCCUR=2
> 710	95	33	SIRUNYAN	18AL	CMS	$\geq 3\ell^\pm$ + jets + \cancel{E}_T , Tstop7, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175$ GeV, $m_{\tilde{t}_1} = 200$ GeV, BR($\tilde{t}_2 \rightarrow \tilde{t}_1 Z$) = BR($\tilde{t}_2 \rightarrow \tilde{t}_1 H$) = 50%	OCCUR=3
> 730	95	34	SIRUNYAN	18AN	CMS	1 or 2 γ + ℓ + jets, GGM, Tstop12, $m_{\tilde{\chi}_1^0} = 150$ GeV	
> 650	95	34	SIRUNYAN	18AN	CMS	1 or 2 γ + ℓ + jets, GGM, Tstop12, $m_{\tilde{\chi}_1^0} = 500$ GeV	OCCUR=2
>1000	95	35	SIRUNYAN	18AY	CMS	jets + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 500	95	35	SIRUNYAN	18AY	CMS	jets + \cancel{E}_T , Tstop4, $m_{\tilde{\chi}_1^0} = 420$ GeV	OCCUR=2
> 510	95	36	SIRUNYAN	18B	CMS	jets + \cancel{E}_T , Tstop4, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 10$ GeV	
> 800	95	37	SIRUNYAN	18C	CMS	$\ell^\pm \ell^\mp$ + b -jets + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0$	
> 750	95	37	SIRUNYAN	18C	CMS	$\ell^\pm \ell^\mp$ + b -jets + \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=2
>1050	95	37	SIRUNYAN	18C	CMS	Combination of all-hadronic, 1 ℓ^\pm and $\ell^\pm \ell^\mp$ searches, Tstop1, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
>1000	95	37	SIRUNYAN	18C	CMS	Combination of all-hadronic, 1 ℓ^\pm and $\ell^\pm \ell^\mp$ searches, Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=4
>1200	95	37	SIRUNYAN	18C	CMS	$\ell^\pm \ell^\mp$ + b -jets + \cancel{E}_T , Tstop11, $m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\ell}} = 0.5 m_{\tilde{\chi}_1^\pm}$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=5
>1300	95	37	SIRUNYAN	18C	CMS	$\ell^\pm \ell^\mp$ + b -jets + \cancel{E}_T , Tstop11, $m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\ell}} = 0.95 m_{\tilde{\chi}_1^\pm}$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=6
none 460–1060	95	37	SIRUNYAN	18C	CMS	$\ell^\pm \ell^\mp$ + b -jets + \cancel{E}_T , Tstop11, $m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\ell}} = 0.05 m_{\tilde{\chi}_1^\pm}$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=7
>1020	95	38	SIRUNYAN	18D	CMS	top quark (hadronically decaying) + jets + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 420	95	39	SIRUNYAN	18DI	CMS	ℓ^\pm + jet + \cancel{E}_T , Tstop3, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 10$ GeV	
> 560	95	39	SIRUNYAN	18DI	CMS	ℓ^\pm + jet + \cancel{E}_T , Tstop3, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 80$ GeV	OCCUR=2

> 540	95	39 SIRUNYAN	18DI CMS	ℓ^\pm , Tstop10, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 40$ GeV	OCCUR=3
> 590	95	39 SIRUNYAN	18DI CMS	Combination of all-hadronic and 1 ℓ^\pm searches, Tstop3, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 30$ GeV	OCCUR=4
> 670	95	39 SIRUNYAN	18DI CMS	Combination of all-hadronic and 1 ℓ^\pm searches, Tstop10, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 60$ GeV	OCCUR=5
> 450	95	40 SIRUNYAN	18DN CMS	$\ell^\pm \ell^\mp$, Tstop1, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = m_W$	
none 225–325	95	40 SIRUNYAN	18DN CMS	$\ell^\pm \ell^\mp$, Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 2 m_W$	OCCUR=2
none 210–690	95	40 SIRUNYAN	18DN CMS	$\ell^\pm \ell^\mp$, Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
none 250–600	95	40 SIRUNYAN	18DN CMS	$\ell^\pm \ell^\mp$, Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
> 700	95	41 AABOUD	17AJ ATLS	same-sign $\ell^\pm \ell^\pm / 3 \ell + \text{jets} + \cancel{E}_T$, Tstop11, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 100$ GeV	
> 880	95	42 AABOUD	17AX ATLS	$b\text{-jets} + \cancel{E}_T$, mixture Tstop1 and Tstop2 with BR=50%, $m_{\tilde{\chi}_1^0} = 0$ GeV, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 1$ GeV	
none 250–1000	95	43 AABOUD	17AY ATLS	$\text{jets} + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
none 450–850	95	44 AABOUD	17AY ATLS	$\text{jets} + \cancel{E}_T$, mixture of Tstop1 and Tstop2 with BR=50%, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=2
> 720	95	45 AABOUD	17BE ATLS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 400	95	46 AABOUD	17BE ATLS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tstop3, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 40$ GeV	OCCUR=2
> 430	95	47 AABOUD	17BE ATLS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tstop1 (offshell t), $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} \sim m_W$	OCCUR=3
> 700	95	48 AABOUD	17BE ATLS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tstop2, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^\pm} = 10$ GeV, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
> 750	95	49 KHACHATRY...17	CMS	$\text{jets} + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0} = 100$ GeV	
none 250–740	95	50 KHACHATRY...17AD	CMS	$\text{jets} + b\text{-jets} + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 610	95	51 KHACHATRY...17AD	CMS	$\text{jets} + b\text{-jets} + \cancel{E}_T$, mixture Tstop1 and Tstop2 with BR=50%, $m_{\tilde{\chi}_1^0} = 60$ GeV	OCCUR=2
> 590	95	52 KHACHATRY...17P	CMS	1 or more $\text{jets} + \cancel{E}_T$, Tstop8, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5$ GeV, $m_{\tilde{\chi}_1^0} = 100$ GeV	
none 280–640	95	52 KHACHATRY...17P	CMS	1 or more $\text{jets} + \cancel{E}_T$, Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 350	95	52 KHACHATRY...17P	CMS	1 or more $\text{jets} + \cancel{E}_T$, Tstop4, $10 \text{ GeV} < m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 80$ GeV	OCCUR=3
> 280	95	52 KHACHATRY...17P	CMS	1 or more $\text{jets} + \cancel{E}_T$, Tstop3, $10 \text{ GeV} < m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 80$ GeV	OCCUR=4
> 320	95	52 KHACHATRY...17P	CMS	1 or more $\text{jets} + \cancel{E}_T$, Tstop9, $10 \text{ GeV} < m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 80$ GeV	OCCUR=5
> 240	95	53 KHACHATRY...17S	CMS	$\text{jets} + \cancel{E}_T$, Tstop4, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 10$ GeV	

> 225	95	54	KHACHATRY...17S	CMS	jets+ \cancel{E}_T , Tstop3, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 10 \text{ GeV}$	OCCUR=2	
> 325	95	55	KHACHATRY...17S	CMS	jets+ \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = 0.25 m_{\tilde{t}} + 0.75 m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 225 \text{ GeV}$	OCCUR=3	
> 400	95	56	KHACHATRY...17S	CMS	jets+ \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = 0.75 m_{\tilde{t}} + 0.25 m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=4	
> 500	95	57	KHACHATRY...17S	CMS	jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=5	
>1120	95	58	SIRUNYAN	17AS	CMS	1 ℓ +jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1000	95	58	SIRUNYAN	17AS	CMS	1 ℓ +jets+ \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
> 980	95	58	SIRUNYAN	17AS	CMS	1 ℓ +jets+ \cancel{E}_T , Tstop8, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=5
>1040	95	59	SIRUNYAN	17AT	CMS	jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 750	95	59	SIRUNYAN	17AT	CMS	jets+ \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
> 940	95	59	SIRUNYAN	17AT	CMS	jets+ \cancel{E}_T , Tstop8, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=3
> 540	95	59	SIRUNYAN	17AT	CMS	jets+ \cancel{E}_T , Tstop3, $10 \text{ GeV} < m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 80 \text{ GeV}$	OCCUR=4
> 480	95	59	SIRUNYAN	17AT	CMS	jets+ \cancel{E}_T , Tstop4, $10 \text{ GeV} < m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 80 \text{ GeV}$	OCCUR=5
> 530	95	59	SIRUNYAN	17AT	CMS	jets+ \cancel{E}_T , Tstop10, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $10 \text{ GeV} < m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 80 \text{ GeV}$	OCCUR=6
>1070	95	60	SIRUNYAN	17AZ	CMS	≥ 1 jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 900	95	60	SIRUNYAN	17AZ	CMS	≥ 1 jets+ \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1020	95	60	SIRUNYAN	17AZ	CMS	≥ 1 jets+ \cancel{E}_T , Tstop8, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=3
> 540	95	60	SIRUNYAN	17AZ	CMS	≥ 1 jets+ \cancel{E}_T , Tstop4, $10 \text{ GeV} < m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 80 \text{ GeV}$	OCCUR=4
none 280–830	95	61	SIRUNYAN	17K	CMS	0, 1 ℓ^\pm +jets+ \cancel{E}_T (combination), Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 700	95	61	SIRUNYAN	17K	CMS	0, 1 ℓ^\pm +jets+ \cancel{E}_T (combination), Tstop8, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=2
> 160	95	61	SIRUNYAN	17K	CMS	jets+ \cancel{E}_T , Tstop4, $10 < m_{\tilde{t}} - m_{\tilde{\chi}_1^0} < 80 \text{ GeV}$	OCCUR=3
none 230–960	95	62	SIRUNYAN	17P	CMS	jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 990	95	62	SIRUNYAN	17P	CMS	jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
> 323	95	63	AABOUD	16D	ATLS	≥ 1 jet + \cancel{E}_T , Tstop4, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$	
none, 745–780	95	64	AABOUD	16J	ATLS	1 ℓ^\pm + ≥ 4 jets + \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
> 490–650	95	65	AAD	16AY	ATLS	2 ℓ (including hadronic τ) + \cancel{E}_T , Tstop5, $87 \text{ GeV} < m_{\tilde{\tau}} < m_{\tilde{t}_1}$	

> 700	95	66	KHACHATRY...16AV CMS	1 or 2 ℓ^\pm +jets+b-jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} < 250$ GeV	
> 700	95	66	KHACHATRY...16AV CMS	1 or 2 ℓ^\pm +jets+b-jets \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^0} = 0$ GeV, $m_{\tilde{\chi}_1^\pm}$ $= 0.75 m_{\tilde{t}_1} + 0.25 m_{\tilde{\chi}_1^0}$	OCCUR=2
> 775	95	67	KHACHATRY...16BK CMS	jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} < 200$ GeV	
> 620	95	67	KHACHATRY...16BK CMS	jets+ \cancel{E}_T , Tstop2, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
> 800	95	68	KHACHATRY...16BS CMS	jets+ \cancel{E}_T , Tstop1, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 316	95	69	KHACHATRY...16Y CMS	1 or 2 soft ℓ^\pm + jets + \cancel{E}_T , Tstop3, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 25$ GeV	
> 250	95	70	AAD	15CJ ATLS B($\tilde{t} \rightarrow c\tilde{\chi}_1^0$)+B($\tilde{t} \rightarrow bf f'\tilde{\chi}_1^0$) $= 1$, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 10$ GeV	
> 270	95	70	AAD	15CJ ATLS $\tilde{t} \rightarrow c\tilde{\chi}_1^0$, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 80$ GeV	OCCUR=2
none, 200–700	95	70	AAD	15CJ ATLS $\tilde{t} \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
> 500	95	70	AAD	15CJ ATLS B($\tilde{t} \rightarrow t\tilde{\chi}_1^0$) + B($\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$) $= 1$, $\tilde{\chi}_1^\pm \rightarrow W^{(*)}\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^\pm}$ $= 2m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} < 160$ GeV	OCCUR=4
> 600	95	70	AAD	15CJ ATLS $\tilde{t}_2 \rightarrow Z\tilde{t}_1$, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 180$ GeV, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=5
> 600	95	70	AAD	15CJ ATLS $\tilde{t}_2 \rightarrow h\tilde{t}_1$, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 180$ GeV, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=6
none, 172.5–191	95	71	AAD	15J ATLS $\tilde{t} \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 1$ GeV	
> 450	95	72	KHACHATRY...15AF CMS	$\tilde{t} \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 0$, $m_{\tilde{t}} > m_t$ $+ m_{\tilde{\chi}_1^0}$	
> 560	95	73	KHACHATRY...15AH CMS	$\tilde{t} \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 0$, $m_{\tilde{t}} > m_t$ $+ m_{\tilde{\chi}_1^0}$	
> 250	95	74	KHACHATRY...15AH CMS	$\tilde{t} \rightarrow c\tilde{\chi}_1^0$, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} < 10$ GeV	OCCUR=2
> 730	95	75	KHACHATRY...15X CMS	$\tilde{t} \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 100$ GeV, $m_{\tilde{t}} > m_t + m_{\tilde{\chi}_1^0}$	
none 400–645	95	75	KHACHATRY...15X CMS	$\tilde{t} \rightarrow t\tilde{\chi}_1^0$ or $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$, $m_{\tilde{\chi}_1^0}$ $= 100$ GeV, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} =$ 5 GeV	OCCUR=2
none 270–645	95	76	AAD	14AJ ATLS ≥ 4 jets + \cancel{E}_T , $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} < 30$ GeV	
none 250–550	95	76	AAD	14AJ ATLS ≥ 4 jets + \cancel{E}_T , B($\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$) $= 50\%$, $m_{\tilde{\chi}_1^\pm} = 2 m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} < 60$ GeV	OCCUR=2
none 210–640	95	77	AAD	14BD ATLS ℓ^\pm + jets + \cancel{E}_T , $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 500	95	77	AAD	14BD ATLS ℓ^\pm + jets + \cancel{E}_T , $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$, $m_{\tilde{\chi}_1^\pm} = 2 m_{\tilde{\chi}_1^0}$, 100 GeV < $m_{\tilde{\chi}_1^0} < 150$ GeV	OCCUR=2
none 150–445	95	78	AAD	14F ATLS $\ell^\pm \ell^\mp$ final state, $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^\pm} = 10$ GeV, $m_{\tilde{\chi}_1^0}$ $= 1$ GeV	
none 215–530	95	78	AAD	14F ATLS $\ell^\pm \ell^\mp$ final state, $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=2
> 270	95	79	AAD	14T ATLS $\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 200$ GeV	
> 240	95	79	AAD	14T ATLS $\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} < 85$ GeV	OCCUR=2

> 255	95	79 AAD	14T ATLS	$\tilde{t}_1 \rightarrow b f f' \tilde{\chi}_1^0, m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} \approx m_b$	OCCUR=3
> 400	95	80 CHATRCHYAN 14AH	CMS	jets + $\cancel{E}_T, \tilde{t} \rightarrow t \tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 50$ GeV	
		81 CHATRCHYAN 14R	CMS	$\geq 3\ell^\pm, \tilde{t} \rightarrow (b \tilde{\chi}_1^\pm / t \tilde{\chi}_1^0), \tilde{\chi}_1^\pm \rightarrow (q q' / \ell \nu) \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow (H/Z) \tilde{G}, \text{GMSB, natural higgsino NLSP scenario}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
> 850	95	82 AABOUD	17AF ATLS	$2\ell + \text{jets} + b\text{-jets} + \cancel{E}_T, T_{\text{stop}6}, m_{\tilde{\chi}_1^0} = 0$	
> 800	95	83 AABOUD	17AF ATLS	$2\ell + \text{jets} + b\text{-jets} + \cancel{E}_T, T_{\text{stop}7}$ with 100% decays via Z, $m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=2
> 880	95	84 AABOUD	17AF ATLS	$2\ell + \text{jets} + b\text{-jets} + \cancel{E}_T, T_{\text{stop}7}$ with 100% decays via higgs, $m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=3
> 230		85 AABOUD ROLBIECKI	17AY ATLS 15 THEO	jets + \cancel{E}_T , pMSSM-inspired $W W$ xsection, $\tilde{t}_1 \rightarrow b W \tilde{\chi}_1^0, m_{\tilde{t}_1} \simeq m_b + m_W + m_{\tilde{\chi}_1^0}$	OCCUR=3
> 600	95	86 AAD	14B ATLS	$Z + b \cancel{E}_T, \tilde{t}_2 \rightarrow Z \tilde{t}_1, \tilde{t}_1 \rightarrow t \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 200$ GeV	
> 540	95	86 AAD	14B ATLS	$Z + b \cancel{E}_T, \tilde{t}_1 \rightarrow t \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow Z \tilde{G}, \text{natural GMSB, } 100$ GeV $< m_{\tilde{\chi}_1^0} < m_{\tilde{t}_1} - 10$ GeV	OCCUR=2
> 360	95	87 CHATRCHYAN 14U	CMS	$\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm r, \tilde{\chi}_1^\pm \rightarrow f f' \tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow H \tilde{G}$ simplified model, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5$ GeV, GMSB	
> 215	95	CZAKON	14	$\tilde{t} \rightarrow t \tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 10$ GeV	
		88 KHACHATRY...14C	CMS	$\tilde{t}_2 \rightarrow H \tilde{t}_1$ or $\tilde{t}_2 \rightarrow Z \tilde{t}_1$ simplified model	

¹ HAYRAPETYAN 23E searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of gluino, top squark and electroweakino pair production in events with at least one photon, multiple jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set in models for strong production, Tglu4D, Tglu4E, Tglu4F and Tstop13, see their figure 9. They also interpret the results in the models for electroweak production, shown in their figure 10. Tchi1n1A assumes wino-like $\tilde{\chi}_1^\pm \tilde{\chi}_1^0$ production, while Tchi1chi1A assumes higgsino-like cross sections and includes $\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \tilde{\chi}_2^0$ and $\tilde{\chi}_{1,2}^0 \tilde{\chi}_1^\pm$ production. For $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ alone no mass point can be excluded in the model Tchi1chi1A, but in another model for $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ production, Tn1n2A.

NODE=S046STP;LINKAGE=XD

² TUMASYAN 23AB searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of top squark pair production in a final state with at least one hadronically decaying tau lepton and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of \tilde{t} for the model Tstop16, see their Figure 9. The exclusion limits are not very sensitive to the choice of the $\tilde{\tau}$ mass parameter, chosen between $0.25 < (m_{\tilde{\tau}^\pm} - m_{\tilde{\chi}_1^0}) / (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}) < 0.75$ because of the complementary nature of the signal diagrams.

NODE=S046STP;LINKAGE=VD

³ TUMASYAN 23K searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of top squark pair production in events with a high-momentum jet, an electron or muon with low transverse momentum, and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the top squark mass in the simplified model Tstop3 for $10 \text{ GeV} < m_{\tilde{t}} - m_{\tilde{\chi}_1^0} < 80$ GeV, see their Figure 10.

NODE=S046STP;LINKAGE=WD

⁴ TUMASYAN 22Q searched in up to 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of electroweakino and top squark pair production with a small mass difference between the produced supersymmetric particles and the lightest neutralino in events with two or three low-momentum leptons and missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the model Tchi1n2F, see their Figure 8. Limits are also set in a higgsino simplified model with both $\tilde{\chi}_2^0 \tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0 \tilde{\chi}_1^0$ production, where $\tilde{\chi}_2^0 \rightarrow Z \tilde{\chi}_1^0$ and $m_{\tilde{\chi}_1^\pm} = 1/2(m_{\tilde{\chi}_2^0} + m_{\tilde{\chi}_1^0})$. A model inspired by the pMSSM is used for further interpretations

NODE=S046STP;LINKAGE=UD

- in the case of a higgsino LSP, see their Figure 9. Limits are also set on the mass of the top squark in the models Tstop2 and Tstop3, see their Figure 10.
- 5 AAD 21AW searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of stops in events with one or two hadronically decaying τ leptons, jets, b -jets and \cancel{E}_T . No significant excess above the Standard Model predictions is observed. Limits are set on the \tilde{t}_1 mass as a function of the $\tilde{\tau}_1$ in the Tstop5 scenario. See their Fig. 8. NODE=S046STP;LINKAGE=OD
- 6 AAD 21O searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of top squarks in events with one electron or muon, jets, and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the top squark mass in the Tstop1 and Tstop3 simplified models and dark matter models, see their Figures 13, 14 and 15. NODE=S046STP;LINKAGE=MD
- 7 AAD 21P searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of top squarks in events with two opposite-sign leptons, jets, and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the top squark mass in the Tstop1, Tstop2, and Tstop3 simplified models, see their Figures 14. NODE=S046STP;LINKAGE=LD
- 8 SIRUNYAN 21AD searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with multiple jets, no leptons, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the top squark mass in the simplified models Tstop1, Tstop2 with $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, and a 50:50 mixture of these with $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, see their Figure 8. Limits are also set on the top squark mass for $10 \text{ GeV} < m_{\tilde{t}} - m_{\tilde{\chi}_1^\pm} < 80 \text{ GeV}$ in the simplified models Tstop2, Tstop 3, and Tstop4, see their Figure 9. For indirect top squark production, limits are set on the gluino mass in the simplified models Tglu3A, Tglu3C with $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 20 \text{ GeV}$, and Tglu3D with $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, see their Figure 10. NODE=S046STP;LINKAGE=ND
- 9 SIRUNYAN 21B searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for the pair production of top squarks in events with two oppositely charged leptons (electrons or muons), jets identified as originating from a b -quark and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1, Tstop2 and Tstop11 simplified models, see their Figures 6 and 7. NODE=S046STP;LINKAGE=ID
- 10 TUMASYAN 21I searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of top squarks in events with at least two jets and large \cancel{E}_T , categorized into events with 0, 1, or 2 leptons. No significant excess above the Standard Model expectations is observed. Limits are set on the top squark mass in the simplified model Tstop1 in the top corridor $|m_{\tilde{t}} - m_{\tilde{\chi}_1^0} - 175 \text{ GeV}| < 30 \text{ GeV}$ using dilepton events, see their Figure 7. Limits are also set for a combination of earlier searches with 0, 1, and 2 leptons in the models Tstop1, Tstop2 and a 50:50 mixture of these models, see their Figure 9. The results are interpreted in an alternative signal model of dark matter production via a spin-0 mediator in association with a top quark pair as well. NODE=S046STP;LINKAGE=TD
- 11 AABOUD 20 searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing one electron-muon pair with opposite charge. The search targets a region of parameter space where the kinematics of top squark pair production and top quark pair production is very similar and makes use of the double-differential angular distributions of the leptons. No excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1 model, see Figures 16 and 17. NODE=S046STP;LINKAGE=DD
- 12 AAD 20AS searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of top squarks in events containing either a pair of jets consistent with SM Higgs boson decay into b -quarks or a same-flavour opposite-sign dilepton pair with an invariant mass consistent with a Z boson. No significant excess over the expected background is observed. Limits at 95% C.L. are set in Tstop6 simplified model. Assuming $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$, \tilde{t}_1 masses up to 1220 GeV are excluded for $m_{\tilde{\chi}_2^0}$ around 900 GeV. Limits reduce down to \tilde{t}_1 masses up to 900 GeV for $m_{\tilde{\chi}_2^0} = 130 \text{ GeV}$. See their Fig. 10. Limits are presented also in case of $B(\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 h) = 0$ and 1, see their Fig. 11. NODE=S046STP;LINKAGE=JD
- 13 AAD 20AS searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of top squarks in events containing either a pair of jets consistent with SM Higgs boson decay into b -quarks or a same-flavour opposite-sign dilepton pair with an invariant mass consistent with a Z boson. No significant excess over the expected background is observed. Limits at 95% C.L. are set in simplified model featuring \tilde{t}_2 pair production, $\tilde{t}_2 \rightarrow \tilde{t}_1 Z$ and $\tilde{t}_1 \rightarrow b f f' \tilde{\chi}_1^0$. Assuming $m_{\tilde{\chi}_1^0} = 300 \text{ GeV}$, and a mass difference between \tilde{t}_1 and $\tilde{\chi}_1^0$ of 40 GeV, \tilde{t}_2 masses up to 860 GeV are excluded. See their Fig. 12. NODE=S046STP;LINKAGE=KD
- 14 AAD 20S searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Exclusion limits at 95% C.L. are set on top squark masses in the Tstop1 model up to 1250 GeV for lightest neutralino masses below 200 GeV. Additional constraints are set in the case where $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} \sim m_{\tilde{t}}$ for which top squark masses in the range 300–630 GeV are excluded. See their Fig. 13. NODE=S046STP;LINKAGE=ED
- 15 AAD 20S searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Exclusion limits at 95% C.L. are set on top squark masses in the Tstop3 NODE=S046STP;LINKAGE=FD

model in the range 300–660 GeV. In case $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} \sim 5$ GeV or above, $m_{\tilde{t}}$ below 500 GeV are excluded. See their Fig. 13(b).

- 16 AAD 20V searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with two same-sign charged leptons (electrons or muons) and jets. No significant excess above the Standard Model expectations is observed. Exclusion limits at 95% C.L. are set on the top squark mass up to 765 GeV assuming $\tilde{t}_1 \rightarrow t\tilde{\chi}_2^0$ with $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^\pm W$ and $\tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 W$. Masses of the charginos and lightest neutralinos are set as $m_{\tilde{\chi}_1^0} = m_{\tilde{t}_1} - 275$ GeV, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 100$ GeV and $m_{\tilde{\chi}_1^\pm} \sim m_{\tilde{\chi}_1^0}$. See their Fig. 8(b). NODE=S046STP;LINKAGE=HD
- 17 SIRUNYAN 20AH searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pair production of top squarks in events with a single isolated electron or muon, multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1, Tstop2 and Tstop8 simplified models, see Figures 6, 7 and 8, respectively. NODE=S046STP;LINKAGE=BD
- 18 SIRUNYAN 20T searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least two jets, and two isolated same-sign or three or more charged leptons (electrons or muons). No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figure 7, and in the Tglu1C and Tglu1B simplified models, see their Figures 8 and 9. Limits are also set on the sbottom mass in the Tsbott2 simplified model, see their Figure 10, and on the stop mass in the Tstop7 simplified model, see their Figure 11. Finally, limits are set on the gluino mass in RPV simplified models where the gluino decays either via $\tilde{g} \rightarrow qq\bar{q}\bar{q} + e/\mu/\tau$ or via $\tilde{g} \rightarrow tbs$, see Figure 12. NODE=S046STP;LINKAGE=YC
- 19 SIRUNYAN 20U searched in 77.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for the pair production of top squarks in events with two hadronically decaying taus, jets identified as originating from a b -quark and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop11 simplified model assuming the final state leptons are taus. Different values of the scalar tau mass are considered; the impact on the lower bound is negligible. NODE=S046STP;LINKAGE=XC
- 20 SIRUNYAN 19AU searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at last one photon, jets, some of which are identified as originating from b -quarks, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. In the framework of GMSB, limits are set on the gluino mass in the Tglu4C, Tglu4D and Tglu4E simplified models, and on the top squark mass in the Tstop13 simplified model, see their Figure 5. NODE=S046STP;LINKAGE=RC
- 21 SIRUNYAN 19CH searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A and Tglu3A simplified models, see their Figure 13. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tsbott1, Tstop1 simplified models, see their Figure 14. NODE=S046STP;LINKAGE=VC
- 22 SIRUNYAN 19S searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with zero or one charged leptons, jets and \cancel{E}_T . The razor variables (M_R and R^2) are used to categorize the events. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu3C simplified models, see Figures 22 and 23, and on the stop mass in the Tstop1 simplified model, see their Figure 24. NODE=S046STP;LINKAGE=SC
- 23 SIRUNYAN 19U searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing one electron-muon pair with opposite charge. The search targets a region of parameter space where the kinematics of top squark pair production and top quark pair production is very similar, due to the mass difference between the top squark and the neutralino being close to the top quark mass. No excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1 model, with $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$ close to m_t , see Figure 5. NODE=S046STP;LINKAGE=UC
- 24 AABOUD 18AQ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for top squark pair production in final states with one isolated electron or muon, several energetic jets, and missing transverse momentum. No significant excess over the Standard Model prediction is observed. In case of Tstop1 models, top squark masses up to 940 GeV are excluded assuming $m_{\tilde{\chi}_1^0} = 0$ GeV, see their Fig. 20. If the top quark is not on-shell (3-body) decay, exclusions up to 500 GeV are obtained for $m_{\tilde{\chi}_1^0} = 300$ GeV. Exclusions as a function of $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$ are given in their Fig. 21. NODE=S046STP;LINKAGE=JC
- 25 AABOUD 18AQ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for top squark pair production in final states with one isolated electron or muon, several energetic jets, and missing transverse momentum. No significant excess over the Standard Model prediction is observed. In case of Tstop3 models (4-body), top squark masses up to 370 GeV are excluded for $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$ as low as 20 GeV. Top squark masses below 195 GeV are excluded for all $m_{\tilde{\chi}_1^0}$, see their Fig. 20 and Fig. 21. NODE=S046STP;LINKAGE=KC
- 26 AABOUD 18AQ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for top squark pair production in final states with one isolated electron or muon, several energetic jets, and missing transverse momentum. No significant excess over the Standard Model prediction is observed. In case of Tstop2 models, top squark masses up to 840 GeV are excluded NODE=S046STP;LINKAGE=LC

- for $m_{\tilde{\tau}} - m_{\tilde{\chi}_1^\pm} = 10$ GeV. See their Fig. 23. Exclusion limits for this decay mode are presented also in the context of Higgsino-LSP phenomenological MSSM models, where $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5$ GeV, see their Fig 26.
- 27 AABOUD 18BV searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one jet identified as c -jet, large missing transverse energy and no leptons. Good agreement is observed between the number of events in data and Standard Model predictions. The results are translated into exclusion limits in Tstop4 models. In scenarios with differences of the stop and neutralino masses below 100 GeV, stop masses below 500 GeV are excluded. See their Fig.6 and Fig.7. NODE=S046STP;LINKAGE=PC
- 28 AABOUD 18BV searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one jet identified as c -jet, large missing transverse energy and no leptons. Good agreement is observed between the number of events in data and Standard Model predictions. The results are translated into exclusion limits in Tstop1 models. In scenarios with massless neutralinos, top squark masses below 850 GeV are excluded. See their Fig.6. NODE=S046STP;LINKAGE=QC
- 29 AABOUD 18I searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one jet with a transverse momentum above 250 GeV and no leptons. Good agreement is observed between the number of events in data and Standard Model predictions. The results are translated into exclusion limits in Tstop3 models. Stop masses below 390 GeV are excluded for $m_{\tilde{\tau}} - m_{\tilde{\chi}_1^0} = m_b$. See their Fig.9(b). NODE=S046STP;LINKAGE=DC
- 30 AABOUD 18I searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least one jet with a transverse momentum above 250 GeV and no leptons. Good agreement is observed between the number of events in data and Standard Model predictions. The results are translated into exclusion limits in Tstop4 models. In scenarios with differences of the stop and neutralino masses around 5 GeV, stop masses below 430 GeV are excluded. See their Fig.9(a). NODE=S046STP;LINKAGE=EC
- 31 AABOUD 18Y searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for direct pair production of top squarks in final states with two tau leptons, b -jets, and missing transverse momentum. At least one hadronic τ is required. No significant deviation from the SM predictions is observed in the data. The analysis results are interpreted in Tstop5 models with a nearly massless gravitino. Top squark masses up to 1.16 TeV and tau slepton masses up to 1 TeV are excluded, see their Fig 7. NODE=S046STP;LINKAGE=HC
- 32 SIRUNYAN 18AJ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing two low-momentum, oppositely charged leptons (electrons or muons) and \cancel{E}_T . No excess over the expected background is observed. Limits are derived on the wino mass in the Tchi1n2F simplified model, see their Figure 5. Limits are also set on the stop mass in the Tstop10 simplified model, see their Figure 6. Finally, limits are set on the Higgsino mass in the Tchi1n2G simplified model, see Figure 8 and in the pMSSM, see Figure 7. NODE=S046STP;LINKAGE=CC
- 33 SIRUNYAN 18AL searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least three charged leptons, in any combination of electrons and muons, jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu1C simplified models, see their Figure 5. Limits are also set on the sbottom mass in the Tsb0t2 simplified model, see their Figure 6, and on the stop mass in the Tstop7 simplified model, see their Figure 7. NODE=S046STP;LINKAGE=GC
- 34 SIRUNYAN 18AN searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing one or two photons and a pair of top quarks from the decay of a pair of top squark in a natural gauge-mediated scenario. The final state consists of a lepton (electron or muon), jets and one or two photons. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop12 simplified model, see their Figure 6. NODE=S046STP;LINKAGE=FC
- 35 SIRUNYAN 18AY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing one or more jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A and Tglu3A simplified models, see their Figure 3. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tsb0t1, Tstop1 and Tstop4 simplified models, see their Figure 3. Finally, limits are set on long-lived gluino masses in a Tglu1A simplified model where the gluino is metastable or long-lived with proper decay lengths in the range $10^{-3} \text{ mm} < c\tau < 10^5 \text{ mm}$, see their Figure 4. NODE=S046STP;LINKAGE=IC
- 36 SIRUNYAN 18B searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for the pair production of third-generation squarks in events with jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the sbottom mass in the Tsb0t1 simplified model, see their Figure 5, and on the stop mass in the Tstop4 simplified model, see their Figure 6. NODE=S046STP;LINKAGE=MB
- 37 SIRUNYAN 18C searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for the pair production of top squarks in events with two oppositely charged leptons (electrons or muons), jets identified as originating from a b -quark and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1, Tstop2 and Tstop11 simplified models, see their Figures 11 and 12. The Tstop1 and Tstop2 results are combined with complementary searches in the all-hadronic and single lepton channels, see their Figures 13 and 14. NODE=S046STP;LINKAGE=NB
- 38 SIRUNYAN 18D searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events containing identified hadronically decaying top quarks, no leptons, and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop NODE=S046STP;LINKAGE=OB

- mass in the Tstop1 simplified model, see their Figure 8, and on the gluino mass in the Tglu3A, Tglu3B, Tglu3C and Tglu3E simplified models, see their Figure 9.
- 39 SIRUNYAN 18DI searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of top squarks in events with a low transverse momentum lepton (electron or muon), a high-momentum jet and significant missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop3 and Tstop10 simplified models, see their Figures 7 and 8. A combination of this search with the all-hadronic search is presented in Figure 9. NODE=S046STP;LINKAGE=OC
- 40 SIRUNYAN 18DN searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct electroweak production of charginos and for pair production of top squarks in events with two leptons (electrons or muons) of the opposite electric charge. No significant excess above the Standard Model expectations is observed. Limits are set on the chargino mass in the Tchi1chi1C and Tchi1chi1E simplified models, see their Figure 8. Limits are also set on the stop mass in the Tstop1 and Tstop2 simplified models, see their Figure 9. NODE=S046STP;LINKAGE=NC
- 41 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 700 GeV are set on the top squark mass in Tstop11 simplified models, assuming $m_{\tilde{\chi}_1^0} = m_{\tilde{t}} - 275 \text{ GeV}$ and $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 100 \text{ GeV}$. See their Figure 4(e). NODE=S046STP;LINKAGE=BC
- 42 AABOUD 17AX searched in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two jets identified as originating from b -quarks and large missing transverse momentum, with or without leptons. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of top squarks. Assuming 50% BR for Tstop1 and Tstop2 simplified models, a \tilde{t}_1 mass below 880 (860) GeV is excluded for $m_{\tilde{\chi}_1^0} = 0$ (<250) GeV. See their Fig. 7(b). NODE=S046STP;LINKAGE=QB
- 43 AABOUD 17AY searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least four jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits in the range 250–1000 GeV are set on the top squark mass in Tstop1 simplified models. For the first time, additional constraints are set for the region $m_{\tilde{t}_1} \sim m_{\tilde{t}} + m_{\tilde{\chi}_1^0}$, with exclusion of the \tilde{t}_1 mass range 235–590 GeV. See their Figure 8. NODE=S046STP;LINKAGE=XB
- 44 AABOUD 17AY searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least four jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits in the range 450–850 GeV are set on the top squark mass in a mixture of Tstop1 and Tstop2 simplified models with BR=50% and assuming $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$ and $m_{\tilde{\chi}_1^0} < 240 \text{ GeV}$. Constraints are given for various values of the BR. See their Figure 9. NODE=S046STP;LINKAGE=YB
- 45 AABOUD 17BE searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two opposite-charge leptons (electrons and muons) and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 720 GeV are set on the top squark mass in Tstop1 simplified models, assuming massless neutralinos. See their Figure 9 (2-body area). NODE=S046STP;LINKAGE=RB
- 46 AABOUD 17BE searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two opposite-charge leptons (electrons and muons) and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 400 GeV are set on the top squark mass in Tstop3 simplified models, assuming $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 40 \text{ GeV}$. See their Figure 9 (4-body area). NODE=S046STP;LINKAGE=TB
- 47 AABOUD 17BE searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two opposite-charge leptons (electrons and muons) and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 430 GeV are set on the top squark mass in Tstop1 simplified models where top quarks are offshell, assuming $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$ close to the W mass. See their Figure 9 (3-body area). NODE=S046STP;LINKAGE=UB
- 48 AABOUD 17BE searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two opposite-charge leptons (electrons and muons) and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 700 GeV are set on the top squark mass in Tstop2 simplified models, assuming $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^\pm} = 10 \text{ GeV}$ and massless neutralinos. See their Figure 10. NODE=S046STP;LINKAGE=WB
- 49 KHACHATRYAN 17 searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more jets, no more than one lepton, and missing transverse momentum, using the razor variables (M_R and R^2) to discriminate between signal and background processes. No evidence for an excess over the expected background is observed. Limits are derived on the stop mass in the Tstop1 simplified model, see Fig. 17. NODE=S046STP;LINKAGE=QA
- 50 KHACHATRYAN 17AD searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing at least four jets (including b -jets), missing transverse momentum and tagged top quarks. No evidence for an excess over the expected background is observed. Top squark masses in the range 250–740 GeV and neutralino masses up to 240 GeV are excluded at 95% C.L. See Fig. 12. NODE=S046STP;LINKAGE=ZA
- 51 KHACHATRYAN 17AD searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing at least four jets (including b -jets), missing transverse momentum and tagged top quarks. No evidence for an excess over the expected background is observed. Limits are derived on the \tilde{t} mass in simplified models that are a mixture of Tstop1 and Tstop2 NODE=S046STP;LINKAGE=BB

- with branching fractions 50% for each of the two decay modes: top squark masses of up to 610 GeV and neutralino masses up to 190 GeV are excluded at 95% C.L. The $\tilde{\chi}_1^\pm$ and the $\tilde{\chi}_1^0$ are assumed to be nearly degenerate in mass, with a 5 GeV difference between their masses. See Fig. 12.
- 52 KHACHATRYAN 17P searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figures 7 and 8. Limits are also set on the squark mass in the Tsqk1 simplified model, see their Fig. 7, and on the sbottom mass in the Tsb0t1 simplified model, see Fig. 8. Finally, limits are set on the stop mass in the Tstop1, Tstop3, Tstop4, Tstop6 and Tstop7 simplified models, see Fig. 8. NODE=S046STP;LINKAGE=UA
- 53 KHACHATRYAN 17S searched in 18.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing multiple jets and missing transverse momentum, using the α_T variable to discriminate between signal and background processes. No evidence for an excess over the expected background is observed. Limits are derived on the stop mass in the Tstop4 model: for $\Delta m = m_{\tilde{t}} - m_{\tilde{\chi}_1^0}$ equal to 10 and 80 GeV, masses of stop below 240 and 260 GeV are excluded, respectively. See their Fig.3. NODE=S046STP;LINKAGE=CB
- 54 KHACHATRYAN 17S searched in 18.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing multiple jets and missing transverse momentum, using the α_T variable to discriminate between signal and background processes. No evidence for an excess over the expected background is observed. Limits are derived on the stop mass in the Tstop3 model: for $\Delta m = m_{\tilde{t}} - m_{\tilde{\chi}_1^0}$ equal to 10 and 80 GeV, masses of stop below 225 and 130 GeV are excluded, respectively. See their Fig.3. NODE=S046STP;LINKAGE=EB
- 55 KHACHATRYAN 17S searched in 18.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing multiple jets and missing transverse momentum, using the α_T variable to discriminate between signal and background processes. No evidence for an excess over the expected background is observed. Limits are derived on the stop mass in the Tstop2 model: assuming $m_{\tilde{\chi}_1^\pm} = 0.25 m_{\tilde{t}} + 0.75 m_{\tilde{\chi}_1^0}$, masses of stop up to 325 GeV and masses of the neutralino up to 225 GeV are excluded. See their Fig.3. NODE=S046STP;LINKAGE=FB
- 56 KHACHATRYAN 17S searched in 18.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing multiple jets and missing transverse momentum, using the α_T variable to discriminate between signal and background processes. No evidence for an excess over the expected background is observed. Limits are derived on the stop mass in the Tstop2 model: assuming $m_{\tilde{\chi}_1^\pm} = 0.75 m_{\tilde{t}} + 0.25 m_{\tilde{\chi}_1^0}$, masses of stop up to 400 GeV are excluded for low neutralino masses. See their Fig.3. NODE=S046STP;LINKAGE=GB
- 57 KHACHATRYAN 17S searched in 18.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing multiple jets and missing transverse momentum, using the α_T variable to discriminate between signal and background processes. No evidence for an excess over the expected background is observed. Limits are derived on the stop mass in the Tstop1 model: assuming masses of stop up to 500 GeV and masses of the neutralino up to 105 GeV are excluded. See their Fig.3. NODE=S046STP;LINKAGE=HB
- 58 SIRUNYAN 17AS searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a single lepton (electron or muon), jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1, Tstop2 and Tstop8 simplified models, see their Figures 5, 6 and 7. NODE=S046STP;LINKAGE=JB
- 59 SIRUNYAN 17AT searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct production of top squarks in events with jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1, Tstop2, Tstop3, Tstop4, Tstop8 and Tstop10 simplified models, see their Figures 9 to 14. NODE=S046STP;LINKAGE=LB
- 60 SIRUNYAN 17AZ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A simplified models, see their Figures 6. Limits are also set on the squark mass in the Tsqk1 simplified model (for single light squark and for 8 degenerate light squarks), on the sbottom mass in the Tsb0t1 simplified model and on the stop mass in the Tstop1 simplified model, see their Fig. 7. Finally, limits are set on the stop mass in the Tstop2, Tstop4 and Tstop8 simplified models, see Fig. 8. NODE=S046STP;LINKAGE=OA
- 61 SIRUNYAN 17K searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for direct production of stop or sbottom pairs in events with multiple jets and significant \cancel{E}_T . A second search also requires an isolated lepton and is combined with the all-hadronic search. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1, Tstop8 and Tstop4 simplified models, see their Figures 7, 8 and 9 (for the Tstop4 limits, only the results of the all-hadronic search are used). Limits are also set on the sbottom mass in the Tsb0t1 simplified model, see Fig. 10 (also here, only the results of the all-hadronic search are used). NODE=S046STP;LINKAGE=CA
- 62 SIRUNYAN 17P searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A, Tglu3A and Tglu3D simplified models, see their Fig. 12. Limits are also set on the squark mass in the Tsqk1 simplified model, on the stop mass in the Tstop1 simplified model, and on the sbottom mass in the Tsb0t1 simplified model, see Fig. 13. NODE=S046STP;LINKAGE=SA

- 63 AABOUD 16D searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with an energetic jet and large missing transverse momentum. The results are interpreted as 95% C.L. limits on mass of stop decaying into a charm-quark and the lightest neutralino in scenarios with $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0}$ between 5 and 20 GeV. See their Fig. 5. NODE=S046STP;LINKAGE=FA
- 64 AABOUD 16J searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with one isolated electron or muon, jets, and missing transverse momentum. For the direct stop pair production model where the stop decays via top and lightest neutralino, the results exclude at 95% C.L. stop masses between 745 GeV and 780 GeV for a massless $\tilde{\chi}_1^0$. See their Fig. 8. NODE=S046STP;LINKAGE=JA
- 65 AAD 16AY searched in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with either two hadronically decaying tau leptons, one hadronically decaying tau and one light lepton, or two light leptons. No significant excess over the Standard Model expectation is found. Exclusion limits at 95% C.L. on the mass of top squarks decaying via $\tilde{\tau}$ to a nearly massless gravitino are placed depending on $m_{\tilde{\tau}}$ which is ranging from the 87 GeV LEP limit to $m_{\tilde{t}_1}$. See their Figs. 9 and 10. NODE=S046STP;LINKAGE=LA
- 66 KHACHATRYAN 16AV searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with one or two isolated leptons, hadronic jets, b -jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1 and Tstop2 simplified models, see Fig. 11. NODE=S046STP;LINKAGE=PA
- 67 KHACHATRYAN 16BK searched in 18.9 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with hadronic jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1 and Tstop2 simplified models, see Fig. 16. NODE=S046STP;LINKAGE=NA
- 68 KHACHATRYAN 16BS searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one energetic jet, no isolated leptons, and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1 simplified model, see Fig. 11 and Table 3. NODE=S046STP;LINKAGE=Y
- 69 KHACHATRYAN 16Y searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with one or two soft isolated leptons, hadronic jets, and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop3 simplified model, see Fig. 3. NODE=S046STP;LINKAGE=Z
- 70 AAD 15CJ searched in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of third generation squarks by combining a large number of searches covering various final states. Stop decays with and without charginos in the decay chain are considered and summaries of all ATLAS Run 1 searches for direct stop production can be found in Fig. 4 (no intermediate charginos) and Fig. 7 (intermediate charginos). Limits are set on stop masses in compressed mass regions, with $B(\tilde{t} \rightarrow c\tilde{\chi}_1^0) + B(\tilde{t} \rightarrow bf'\tilde{\chi}_1^0) = 1$, see Fig. 5. Limits are also set on stop masses assuming that both the decay $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ and $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ are possible, with both their branching ratios summing up to 1, assuming $\tilde{\chi}_1^\pm \rightarrow W^{(*)}\tilde{\chi}_1^0$ and $m_{\tilde{\chi}_1^\pm} = 2 m_{\tilde{\chi}_1^0}$, see Fig. 6. Limits on the mass of the next-to-lightest stop \tilde{t}_2 , decaying either to $Z\tilde{t}_1$, $h\tilde{t}_1$ or $t\tilde{\chi}_1^0$, are also presented, see Figs. 9 and 10. Interpretations in the pMSSM are also discussed, see Figs 13–15. NODE=S046STP;LINKAGE=X
- 71 AAD 15J interpreted the measurement of spin correlations in $t\bar{t}$ production using 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ in exclusion limits on the pair production of light \tilde{t}_1 squarks with masses similar to the top quark mass. The \tilde{t}_1 is assumed to decay through $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$ with predominantly right-handed top and a 100% branching ratio. The data are found to be consistent with the Standard Model expectations and masses between the top quark mass and 191 GeV are excluded, see their Fig. 2. NODE=S046STP;LINKAGE=V
- 72 KHACHATRYAN 15AF searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in simplified models where the decay $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 12. See also Table 5. Exclusions in the CMSSM, assuming $\tan\beta = 30$, $A_0 = -2 \max(m_0, m_{1/2})$ and $\mu > 0$, are also presented, see Fig. 15. NODE=S046STP;LINKAGE=N
- 73 KHACHATRYAN 15AH searched in 19.4 or 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing either a fully reconstructed top quark, or events containing dijets requiring one or both jets to originate from b -quarks, or events containing a mono-jet. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in simplified models where the decay $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 9. Limits are also set in simplified models where the decays $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ and $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$, with $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, each take place with a branching ratio of 50%, see Fig. 10, or with other fractions, see Fig. 11. Finally, limits are set in a simplified model where the decay $\tilde{t} \rightarrow c\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Figs. 9, 10 and 11. NODE=S046STP;LINKAGE=O
- 74 KHACHATRYAN 15AH searched in 19.4 or 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing either a fully reconstructed top quark, or events containing dijets requiring one or both jets to originate from b -quarks, or events containing a mono-jet. No significant excess above the Standard Model expectations is observed. Limits are set NODE=S046STP;LINKAGE=P

- on the stop mass in simplified models where the decay $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 9. Limits are also set in simplified models where the decays $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ and $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$, with $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5$ GeV, each take place with a branching ratio of 50%, see Fig. 10, or with other fractions, see Fig. 11. Finally, limits are set in a simplified model where the decay $\tilde{t} \rightarrow c\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Figs. 9, 10, and 11.
- 75 KHACHATRYAN 15X searched in 19.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with at least two energetic jets, at least one of which is required to originate from a b quark, possibly a lepton, and significant \cancel{E}_T , using the razor variables (M_R and R^2) to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in simplified models where the decay $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ and the decay $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$, with $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5$ GeV, take place with branching ratios varying between 0 and 100%, see Figs. 15, 16 and 17. NODE=S046STP;LINKAGE=S
- 76 AAD 14AJ searched in 20.1 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing four or more jets and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of third-generation squarks in simplified models which either assume that the decay $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$ takes place 100% of the time, see Fig. 8, or that this decay takes place 50% of the time, while the decay $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ takes place the other 50% of the time, see Fig. 9. NODE=S046STP;LINKAGE=K
- 77 AAD 14BD searched in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing one isolated lepton, jets and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of third-generation squarks in simplified models which either assume that the decay $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$ takes place 100% of the time, see Fig. 15, or the decay $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ takes place 100% of the time, see Fig. 16–22. For the mixed decay scenario, see Fig. 23. NODE=S046STP;LINKAGE=L
- 78 AAD 14F searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing two leptons (e or μ), and possibly jets and missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of third-generation squarks in simplified models which either assume that the decay $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ takes place 100% of the time, see Figs. 14–17 and 20, or that the decay $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$ takes place 100% of the time, see Figs. 18 and 19. NODE=S046STP;LINKAGE=J
- 79 AAD 14T searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for monojet-like and c -tagged events. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the masses of third-generation squarks in simplified models which assume that the decay $\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$ takes place 100% of the time, see Fig. 9 and 10. The results of the monojet-like analysis are also interpreted in terms of stop pair production in the four-body decay $\tilde{t}_1 \rightarrow bff'\tilde{\chi}_1^0$, see Fig. 11. NODE=S046STP;LINKAGE=G
- 80 CHATRCHYAN 14AH searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV for events with at least two energetic jets and significant \cancel{E}_T , using the razor variables (M_R and R^2) to discriminate between signal and background processes. A second analysis requires at least one of the jets to be originating from a b -quark. No significant excess above the Standard Model expectations is observed. Limits are set on sbottom masses in simplified models where the decay $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Figs. 28 and 29. Exclusions in the CMSSM, assuming $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, are also presented, see Fig. 26. NODE=S046STP;LINKAGE=M
- 81 CHATRCHYAN 14R searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with at least three leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in a natural higgsino NLSP simplified model (GMSB) where the decay $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$, with $\tilde{\chi}_1^\pm \rightarrow (qq'/\ell\nu)H$, $Z\tilde{G}$, takes place with a branching ratio of 100% (the particles between brackets have a soft p_T spectrum), see Figs. 4–6. NODE=S046STP;LINKAGE=F
- 82 AABOUD 17AF searched in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of top squarks in events containing 2 leptons, jets, b -jets and \cancel{E}_T . In Tstop6 model, assuming $m_{\tilde{\chi}_1^0} = 0$ GeV, \tilde{t}_1 masses up to 850 GeV are excluded for $m_{\tilde{\chi}_2^0} > 200$ GeV. NODE=S046STP;LINKAGE=WA
- 83 AABOUD 17AF searched in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of \tilde{t}_2 in events containing 2 leptons, jets, b -jets and \cancel{E}_T . In Tstop7 model, assuming $m_{\tilde{\chi}_1^0} = 50$ GeV and 100% decays via Z boson, \tilde{t}_2 masses up to 800 GeV are excluded. Exclusion limits are also shown as a function of the \tilde{t}_2 branching ratios in their Figure 7. NODE=S046STP;LINKAGE=XA
- 84 AABOUD 17AF searched in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for evidence of \tilde{t}_2 in events containing 2 leptons, jets, b -jets and \cancel{E}_T . In Tstop7 model, assuming $m_{\tilde{\chi}_1^0} = 50$ GeV and 100% decays via higgs boson, \tilde{t}_2 masses up to 880 GeV are excluded. Exclusion limits are also shown as a function of the \tilde{t}_2 branching ratios in their Figure 7. NODE=S046STP;LINKAGE=YA

- 85 AABOUD 17AY searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least four jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the top squark mass assuming three pMSSM-inspired models. The first one, referred to as Higgsino LSP model, assumes $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$ and $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = 10 \text{ GeV}$, with a mixture of decay modes as in Tstop1, Tstop2 and Tstop6. See their Figure 10. The second and third models are referred to as Wino NLSP and well-tempered pMSSM models, respectively. See their Figure 11 and Figure 12, and text for details on assumptions. NODE=S046STP;LINKAGE=ZB
- 86 AAD 14B searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing a Z boson, with or without additional leptons, plus jets originating from b -quarks and significant missing transverse momentum. No excess over the expected SM background is observed. Limits are derived in simplified models featuring \tilde{t}_2 production, with $\tilde{t}_2 \rightarrow Z\tilde{t}_1$, $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$ with a 100% branching ratio, see Fig. 4, and in the framework of natural GMSB, see Fig. 6. NODE=S046STP;LINKAGE=I
- 87 CHATRCHYAN 14U searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of direct pair production of top squarks, with Higgs bosons in the decay chain. The search is performed using a selection of events containing two Higgs bosons, each decaying to a photon pair, missing transverse energy and possibly b -quark jets. No significant excesses over the expected SM backgrounds are observed. The results are interpreted in the context of a “natural SUSY” simplified model where the decays $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$, with $\tilde{\chi}_1^\pm \rightarrow f\tilde{\chi}_1^0$, and $\tilde{\chi}_1^0 \rightarrow H\tilde{G}$, all happen with 100% branching ratio, see Fig. 4. NODE=S046STP;LINKAGE=H
- 88 KHACHATRYAN 14C searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for evidence of direct pair production of top squarks, with Higgs or Z-bosons in the decay chain. The search is performed using a selection of events containing leptons and b -quark jets. No significant excesses over the expected SM backgrounds are observed. The results are interpreted in the context of a simplified model with pair production of a heavier top-squark mass eigenstate \tilde{t}_2 decaying to a lighter top-squark eigenstate \tilde{t}_1 via either $\tilde{t}_2 \rightarrow H\tilde{t}_1$ or $\tilde{t}_2 \rightarrow Z\tilde{t}_1$, followed in both cases by $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$. The interpretation is performed in the region where the mass difference between the \tilde{t}_1 and $\tilde{\chi}_1^0$ is approximately equal to the top-quark mass, which is not probed by searches for direct \tilde{t}_1 pair production, see Figs. 5 and 6. The analysis excludes top squarks with masses $m_{\tilde{t}_2} < 575 \text{ GeV}$ and $m_{\tilde{t}_1} < 400 \text{ GeV}$ at 95% C.L. NODE=S046STP;LINKAGE=D

R-parity violating \tilde{t} (Stop) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	
none 500–520, 580–770	95	1 TUMASYAN	23L CMS	4 jets with dijet masses $> 350 \text{ GeV}$, Tstop1aRPV	NODE=S046STV NODE=S046STV;CHECK LIMITS
>1500	95	2 TUMASYAN	22AF CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow b\bar{\ell}$, $c\tau = 2 \text{ cm}$	OCCUR=3
>1500	95	2 TUMASYAN	22AF CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow d\bar{\ell}$, $c\tau = 2 \text{ cm}$	OCCUR=5
> 460	95	2 TUMASYAN	22AF CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow b\bar{\ell}$, $0.01\text{cm} < c\tau < 1000 \text{ cm}$	OCCUR=6
> 460	95	2 TUMASYAN	22AF CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow d\bar{\ell}$, $0.01\text{cm} < c\tau < 1000 \text{ cm}$	OCCUR=6
>1100	95	3 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tstop14, λ_{323}'' electroweakino decay, $500 \text{ GeV} < m_{\tilde{\chi}_1^0} < 800 \text{ GeV}$	OCCUR=2
>1150	95	3 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tstop15, λ_{323}'' electroweakino decay, $600 \text{ GeV} < m_{\tilde{\chi}_1^0} < 900 \text{ GeV}$	OCCUR=2
>1300	95	3 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tstop1, λ_{323}'' , electroweakino decay, $500 \text{ GeV} < m_{\tilde{\chi}_1^0} < 1000 \text{ GeV}$	OCCUR=3
>1600	95	4 SIRUNYAN	21AF CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow \bar{d}\bar{d}$, λ_{3i3}'' coupling, $0.4 \text{ mm} < c\tau < 80 \text{ mm}$	OCCUR=3
>1600	95	5 SIRUNYAN	21U CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow b\bar{\ell}$, $5 < c\tau < 240 \text{ mm}$	OCCUR=3
>1600	95	5 SIRUNYAN	21U CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow d\bar{\ell}$, λ_{3i3}'' coupling, $3 < c\tau < 360 \text{ mm}$	OCCUR=5
>1600	95	5 SIRUNYAN	21U CMS	long-lived \tilde{t} , $\tilde{t} \rightarrow \bar{d}\bar{d}$, η_{311}'' coupling, $2 < c\tau < 1320 \text{ mm}$	OCCUR=5
> 670	95	6 SIRUNYAN	21V CMS	$\ell^\pm + \geq 7$ jets, Tstop1 with $\tilde{\chi}_1^0 \rightarrow qqq$, λ_{abc}'' coupling, $a,b,c \in 1,2$	

> 870	95	6 SIRUNYAN	21V CMS	$\ell^\pm + \geq 7$ jets, stealth SYY model	OCCUR=2
>1700	95	7 AAD	20M ATLS	$\tilde{t} \rightarrow q\mu$, long-lived, $T_{\text{stop3RPV}}, \tau = 0.1$ ns	
>1150	95	8 SIRUNYAN	19BI ATLS	$\tilde{t} \rightarrow b\mu$, long-lived, $T_{\text{stop2RPV}}, c\tau = 0.1$ cm	
>1100	95	9 SIRUNYAN	19BJ CMS	$\tilde{t} \rightarrow be$, T_{stop2RPV} , prompt	
none 100–410	95	10 AABOUD	18BB ATLS	4 jets, T_{stop1RPV} with $\tilde{t} \rightarrow ds, \lambda''_{312}$ coupling	
none 100–470, 480–610	95	11 AABOUD	18BB ATLS	4 jets, $T_{\text{stop1RPV}}, \lambda''_{323}$ coupling	OCCUR=2
≥ 600 –1500	95	12 AABOUD	18P ATLS	$2\ell + b$ -jets, T_{stop2RPV} , depending on λ'_{i33} coupling ($i = 1, 2, 3$)	
>1130	95	13 SIRUNYAN	18AD CMS	$\tilde{t} \rightarrow b\ell$, long-lived, $c\tau = 70$ –100 mm	
> 550	95	13 SIRUNYAN	18AD CMS	$\tilde{t} \rightarrow b\ell$, long-lived, $c\tau = 1$ –1000 mm	OCCUR=2
>1400	95	14 SIRUNYAN	18DV CMS	long-lived $\tilde{t}, \tilde{t} \rightarrow \bar{d}\bar{d}$, 0.6 mm $< c\tau < 80$ mm	
none 80–520	95	15 SIRUNYAN	18DY CMS	2, 4 jets, $T_{\text{stop3RPV}}, \lambda''_{312}$ coupling	
none 80–270, 285–340, 400–525	95	15 SIRUNYAN	18DY CMS	2, 4 jets, $T_{\text{stop1RPV}}, \lambda''_{323}$ coupling	OCCUR=2
>1200	95	16 AABOUD	17AI ATLS	$\geq 1\ell + \geq 8$ jets, T_{stop1} with $\tilde{\chi}_1^0 \rightarrow tbs, \lambda''_{323}$ coupling, $m_{\tilde{\chi}_1^0} = 500$ GeV	
none, 100–315	95	17 AAD	16AMATLS	2 large-radius jets, T_{stop1RPV}	
none, 200–350	95	18 KHACHATRY...15L	CMS	$\tilde{t} \rightarrow qq, \lambda''_{312} \neq 0$	
none, 200–385	95	18 KHACHATRY...15L	CMS	$\tilde{t} \rightarrow qb, \lambda''_{323} \neq 0$	OCCUR=2
> 740	95	19 KHACHATRY...14T	CMS	$\tau + b$ -jets, $LQ\bar{D}, \lambda'_{333} \neq 0$, $\tilde{t} \rightarrow \tau b$ simplified model	
> 580	95	19 KHACHATRY...14T	CMS	$\tau + b$ -jets, $LQ\bar{D}, \lambda'_{3jk} \neq 0$ ($j \neq 3$), $\tilde{t} \rightarrow \tilde{\chi}^\pm b, \tilde{\chi}^\pm \rightarrow qq\tau^\pm$ simplified model	OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 770	95	20 AAD	21B ATLS	≥ 8 jets, ≥ 5 b -jets, T_{stop4RPV}	
> 890	95	21 KHACHATRY...16AC	CMS	$e^+e^- + \geq 5$ jets; $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$; $\tilde{\chi}_1^\pm \rightarrow \ell^\pm jj, \lambda'_{ijk}$	
>1000	95	21 KHACHATRY...16AC	CMS	$\mu^+\mu^- + \geq 5$ jets; $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$; $\tilde{\chi}_1^\pm \rightarrow \ell^\pm jj, \lambda'_{ijk}$	OCCUR=2
> 950	95	22 KHACHATRY...16BX	CMS	$\tilde{t} \rightarrow t\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell\ell\nu, \lambda_{121}$ or $\lambda_{122} \neq 0$	
> 790	95	23 KHACHATRY...15E	CMS	$\tilde{t}_1 \rightarrow b\ell, c\tau = 2$ cm	

¹ TUMASYAN 23L searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pairs of dijet resonances with the same mass in final states with at least four jets, for the case where the four-jet production proceeds via an intermediate resonant state and for nonresonant production. No significant excess above the Standard Model expectations is observed. Limits are set in the nonresonant search on the top squark mass in the simplified model $T_{\text{stop1aRPV}}$ with λ_{312} coupling, assuming $B(ds) = 1$, see their figure 12. Limits are also set on resonant pair production of dijet resonances via high mass intermediate states and compared to a signal model of diquarks that decay into pairs of vector-like quarks, see their figures 10 and 11.

NODE=S046STV;LINKAGE=T

² TUMASYAN 22AF searched for evidence of new long-lived particles decaying to leptons in pp collisions at $\sqrt{s} = 13$ TeV, corresponding to 118 (113) fb^{-1} in the ee channel ($e\mu$ and $\mu\mu$) channels. The leptons are required to have transverse impact parameter values between 0.01 and 10 cm and are not required to form a common vertex. No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the top squark in RPV models with top squark pair production and $\tilde{t} \rightarrow b\bar{\ell}$ and $\tilde{t} \rightarrow d\bar{\ell}$, see their Figure 4, which contains a wider range of lifetime limits. Limits are also set on a gauge-mediated SUSY breaking model, where the next-to-lightest SUSY particle is a slepton and the lightest SUSY particle a gravitino \tilde{G} , see their Figure 5, which also contains a wider range of lifetime limits. Limits are also set in a model that produces BSM Higgs bosons (H) with a mass of 125 GeV through gluon-gluon fusion, where the H decays to two long-lived scalars S , each of which decays to two oppositely charged and same-flavor leptons.

NODE=S046STV;LINKAGE=S

³ AAD 21BF searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pair production of gluinos, stops, electroweakinos decaying RPV either directly or indirectly via the LSP. The final state in all cases is one or two leptons, many jets (up to fifteen) and b -jets.

NODE=S046STV;LINKAGE=PD

Different models with different branching fractions of the gluino or stop follow from the assumptions on the nature of the electroweakinos. No significant excess above the Standard Model predictions is observed. Limits are set on the *gluino*, \tilde{t}_1 , electroweakino masses as a function of the $\tilde{\chi}_1^0$ mass in several scenarios of gluino, stop and electroweakino pair production.

- 4 SIRUNYAN 21AF searched in 140 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two displaced vertices from long-lived particles decaying into multijet or dijet final states. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu2RPV with λ''_{323} coupling, on the $\tilde{\chi}_1^0$ mass in an RPV model with $\tilde{\chi}_1^0$ pair production and the RPV decay $\tilde{\chi}_1^0 \rightarrow tbs$ with λ''_{323} coupling and on the \tilde{t} mass in an RPV model with top squark pair production and the RPV decay $\tilde{t} \rightarrow \bar{d}_i \bar{d}_j$ with λ''_{3ij} coupling, see their Figure 7. NODE=S046STV;LINKAGE=P
- 5 SIRUNYAN 21U searched in 132 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with displaced tracks and displaced vertices associated with a dijet system. No significant excess above the Standard Model expectations is observed. Limits are set on long-lived gluinos in an RPC GMSB SUSY model of gluino pair production, with $\tilde{g} \rightarrow g \tilde{G}$, see their Figure 9, in Tglu1A in a mini-split model, see their Figure 10, and in an RPV model of gluino pair production, with $\tilde{g} \rightarrow tbs$ with coupling λ''_{323} , see their Figure 11. Limits are also set on long-lived top squarks in Tstop2RPV, see their Figure 12, in an RPV model with $\tilde{t} \rightarrow d\bar{\ell}$ and $\lambda'_{\chi 31}$ coupling, see their Figure 13, and in a dynamical RPV model with $\tilde{t} \rightarrow \bar{d}\bar{d}$ via a nonholomorphic RPV coupling η''_{311} , see their Figure 14. The best mass limit is achieved in all cases at $c\tau = 30 \text{ mm}$. NODE=S046STV;LINKAGE=M
- 6 SIRUNYAN 21V searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with one charged lepton (e^\pm or μ^\pm) and ≥ 7 jets. No significant excess above the Standard Model expectations is observed. Limits are set on an RPV SUSY model like Tstop1 with the additional decay $\tilde{\chi}_1^0 \rightarrow qqq$ with coupling λ''_{abc} , with $a, b, c \in 1, 2$, and on a stealth SUSY model called SYY, with one scalar particle S with even R-parity and its superpartner \tilde{S} , both singlets under all SM interactions, and with a portal mediated by loop interactions involving a new vectorlike messenger field (Y), where pair produced top squarks decay as $\tilde{t} \rightarrow tg\tilde{S}$, and $\tilde{S} \rightarrow \tilde{G}S$, and $S \rightarrow gg$, see their Figure 6 and 7. NODE=S046STV;LINKAGE=O
- 7 AAD 20M searched for long-lived particles decaying into hadrons and at least one muon in events containing a displaced muon track and a displaced vertex. The analysis uses a dataset of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ corresponding to an integrated luminosity of 136 fb^{-1} . Using the Tstop3RPV simplified model, top squarks with masses up to 1.7 TeV are excluded for a lifetime of 0.1 ns, and masses below 1.3 TeV are excluded for lifetimes between 0.01 ns and 30 ns, see their Fig. 7. The dependence on the RPV coupling λ_{23k} multiplied by $\cos\theta_t$, with θ_t the mixing angle between the left- and right-handed \tilde{t} squarks, is also shown, see their Fig. 7. NODE=S046STV;LINKAGE=K
- 8 SIRUNYAN 19BI searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with two muons and two jets, or with one muon, two jets, and missing transverse momentum. Limits are set in a model of pair-produced, prompt or long-lived top squarks with R-parity violating decays to a b -quark and a lepton (Tstop2RPV), branching fraction of $\tilde{t} \rightarrow b\mu$ equal to 1/3 and $c\tau$ between 0.1 cm and 10 cm in the case of long-lived top squarks. See their Fig. 10. NODE=S046STV;LINKAGE=I
- 9 SIRUNYAN 19BJ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with two electrons and two jets, or with one electron, two jets, and missing transverse momentum. Limits are set in a model of pair-produced, prompt top squarks with R-parity violating decays to a b -quark and a lepton (Tstop2RPV), assuming branching fraction of $\tilde{t} \rightarrow be$ equal to 1/3 and $c\tau = 0 \text{ cm}$. See their Fig.10. NODE=S046STV;LINKAGE=J
- 10 AABOUD 18BB searched in 36.7 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for massive colored resonances which are pair-produced and decay into two jets. No significant deviation from the background prediction is observed. Results are interpreted in a SUSY simplified model as Tstop1RPV with $\tilde{t} \rightarrow ds$. Top squarks with masses in the range 100–410 GeV are excluded, see their Figure 9(a). The λ''_{312} coupling is assumed to be sufficiently large for the decays to be prompt, but small enough to neglect the single-top-squark resonant production through RPV couplings. NODE=S046STV;LINKAGE=C
- 11 AABOUD 18BB searched in 36.7 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for massive coloured resonances which are pair-produced and decay into two jets. No significant deviation from the background prediction is observed. Results are interpreted in Tstop1RPV. Top squarks with masses in the range 100–470 GeV or 480–610 GeV are excluded, see their Figure 9(b). The λ''_{323} coupling is assumed to be sufficiently large for the decays to be prompt, but small enough to neglect the single-top-squark resonant production through RPV couplings. NODE=S046STV;LINKAGE=D
- 12 AABOUD 18P searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair-produced top squarks that decay through RPV λ'_{i33} ($i = 1, 2, 3$) couplings to a final state with two leptons and two jets, at least one of which is identified as a b -jet. No significant excess is observed over the SM background. In the Tstop2RPV model, lower limits on the top squark masses between 600 and 1500 GeV are set depending on the branching fraction to be , $b\mu$, and $b\tau$ final states. See their Figs 6 and 7. NODE=S046STV;LINKAGE=E
- 13 SIRUNYAN 18AD searched in 2.6 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived particles by exploiting the multiplicity of displaced jets to search for the presence of NODE=S046STV;LINKAGE=B

signal decays occurring at distances between 1 and 1000 mm. Limits are set in a model of pair-produced, long-lived top squarks with R-parity violating decays to a b -quark and a lepton, see their Figure 3.

- 14 SIRUNYAN 18DV searched in 38.5 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived particles in events with multiple jets and two displaced vertices composed of many tracks. No events with two well-separated high-track-multiplicity vertices were observed. Limits are set on the stop and the gluino mass in RPV models of supersymmetry where the stop (gluino) is decaying solely into dijet (multijet) final states, see their Figures 6 and 7. NODE=S046STV;LINKAGE=F
- 15 SIRUNYAN 18DY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for the pair production of resonances, each decaying to two quarks. The search is conducted separately in a boosted (two-jet) and resolved (four-jet) jet topology. The mass spectra are found to be consistent with the Standard Model expectations. Limits are set on the stop mass in the Tstop3RPV and Tstop1RPV simplified models, see their Figure 11. NODE=S046STV;LINKAGE=H
- 16 AABOUD 17AI searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more isolated lepton, at least eight jets, either zero or many b -jets, for evidence of R-parity violating decays of the top squark. No significant excess above the Standard Model expectations is observed. Limits up to 1.25 (1.10) TeV are set on the top squark mass in R-parity-violating supersymmetry models where \tilde{t}_1 decays for a bino LSP as: $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ and for a higgsino LSP as $\tilde{t} \rightarrow t\tilde{\chi}_{1,2}^0/b\tilde{\chi}_1^\pm$. These is followed by the decays through the non-zero λ_{323}'' coupling $\tilde{\chi}_{1,2}^0 \rightarrow tbs$, $\tilde{\chi}_1^\pm \rightarrow bbs$. See their Figure 10 and text for details on model assumptions. NODE=S046STV;LINKAGE=A
- 17 AAD 16AM searched in 17.4 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing two large-radius hadronic jets. No deviation from the background prediction is observed. Top squarks with masses between 100 and 315 GeV are excluded at 95% C.L. in the hypothesis that they both decay via R-parity violating coupling λ_{323}'' to b - and s -quarks. See their Fig. 10. NODE=S046STV;LINKAGE=KA
- 18 KHACHATRYAN 15L searched in 19.4 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for pair production of heavy resonances decaying to pairs of jets in four jet events. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in R-parity-violating supersymmetry models where $\tilde{t} \rightarrow qq$ ($\lambda_{312}'' \neq 0$), see Fig. 6 (top) and $\tilde{t} \rightarrow qb$ ($\lambda_{323}'' \neq 0$), see Fig. 6 (bottom). NODE=S046STV;LINKAGE=U
- 19 KHACHATRYAN 14T searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with τ -leptons and b -quark jets, possibly with extra light-flavour jets. No excess above the Standard Model expectations is observed. Limits are set on stop masses in RPV SUSY models with $LQ\bar{D}$ couplings, in two simplified models. In the first model, the decay $\tilde{t} \rightarrow \tau b$ is considered, with $\lambda_{333}' \neq 0$, see Fig. 3. In the second model, the decay $\tilde{t} \rightarrow \tilde{\chi}^\pm b$, with the subsequent decay $\tilde{\chi}^\pm \rightarrow qq\tau^\pm$ is considered, with $\lambda_{3jk}' \neq 0$ and the mass splitting between the top squark and the charging chosen to be 100 GeV, see Fig. 4. NODE=S046STV;LINKAGE=EK
- 20 AAD 21B searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least eight jets and at least 5 b -jets, for evidence of R-parity violating decays of the top squark. No significant excess above the Standard Model expectations is observed. Limits up to 950 GeV are set on the top squark mass in Tstop4RPV simplified model. See their Figure 7 for more detailed mass bounds. NODE=S046STV;LINKAGE=L
- 21 KHACHATRYAN 16AC searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with low missing transverse momentum, two oppositely charged electrons or muons, and at least five jets, at least one of which is a b -jet, for evidence of R-parity violating, charging-mediated decays of the top squark. No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in R-parity-violating supersymmetry models where $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ with $\tilde{\chi}_1^\pm \rightarrow \ell^\pm jj$, $\lambda_{ijk}' \neq 0$ ($ij, k \leq 2$), and with $m_{\tilde{t}} - m_{\tilde{\chi}_1^\pm} = 100 \text{ GeV}$, see Fig. 3. NODE=S046STV;LINKAGE=R
- 22 KHACHATRYAN 16BX searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing 4 leptons coming from R-parity-violating decays of $\tilde{\chi}_1^0 \rightarrow \ell\ell\nu$ with $\lambda_{121} \neq 0$ or $\lambda_{122} \neq 0$. No excess over the expected background is observed. Limits are derived on the gluino, squark and stop masses, see Fig. 23. NODE=S046STV;LINKAGE=RA
- 23 KHACHATRYAN 15E searched for long-lived particles decaying to leptons in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$. Events were selected with an electron and muon with opposite charges and each with transverse impact parameter values between 0.02 and 2 cm. Limits are set on SUSY benchmark models with pair production of top squarks decaying into an $e\mu$ final state via RPV interactions. See their Fig. 2 NODE=S046STV;LINKAGE=W

Heavy \tilde{g} (Gluino) mass limit

For $m_{\tilde{g}} > 60\text{--}70 \text{ GeV}$, it is expected that gluinos would undergo a cascade decay via a number of neutralinos and/or charginos rather than undergo a direct decay to photinos as assumed by some papers. Limits obtained when direct decay is assumed are usually higher than limits when cascade decays are included.

NODE=S046230

NODE=S046230

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

R-parity conserving heavy \tilde{g} (Gluino) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT		NODE=S046GNO NODE=S046GNO;CHECK LIMITS
>2200	95	1 AAD	23AB ATLS	$\geq 1 \gamma + \text{jets} + \cancel{E}_T$, GGM-like, Tglu4D, $\tilde{\chi}_1^0$ NLSP, $m_{\tilde{\chi}_1^0} > 300 \text{ GeV}$		
>2200	95	1 AAD	23AB ATLS	$\geq 1 \gamma + \text{jets} + \cancel{E}_T$, GGM-like, Tglu4G, $\tilde{\chi}_1^0$ NLSP, $m_{\tilde{\chi}_1^0} > 350 \text{ GeV}$		OCCUR=2
>2250	95	2 AAD	23AE ATLS	2 SFOS ℓ , jets, \cancel{E}_T , Tglu1G, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$		
>1950	95	3 AAD	23AE ATLS	2 SFOS ℓ , jets, \cancel{E}_T , Tglu1H, $m_{\tilde{\chi}_2^0} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$		OCCUR=2
>2440	95	4 AAD	23AL ATLS	At least 3 b -tagged jets, 0 or 1 lepton, Tglu3B, $m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$		
>2350	95	4 AAD	23AL ATLS	At least 3 b -tagged jets, 0 or 1 lepton, Tglu2A, $m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$		OCCUR=2
>2050	95	5 AAD	23AL ATLS	At least 3 b -tagged jets, 0 or 1 lepton, Tglu3E, $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 2 \text{ GeV}$, $m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$		OCCUR=3
>2320	95	6 HAYRAPETY...23E	CMS	$\gamma + \text{jets} + \cancel{E}_T$, Tglu4E, $m_{\tilde{\chi}_1^0} = 1700 \text{ GeV}$		
>2375	95	6 HAYRAPETY...23E	CMS	$\gamma + \text{jets} + \cancel{E}_T$, Tglu4D, $m_{\tilde{\chi}_1^0} = 1700 \text{ GeV}$		OCCUR=2
>2260	95	6 HAYRAPETY...23E	CMS	$\gamma + \text{jets} + \cancel{E}_T$, Tglu4F, $m_{\tilde{\chi}_1^0} = 1700 \text{ GeV}$		OCCUR=3
>2120	95	7 TUMASYAN	23AY CMS	$\ell^\pm + \geq 6 \text{ jets} + \geq 1 b\text{-jet}$, Tglu3A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$		
>2050	95	7 TUMASYAN	23AY CMS	$\ell^\pm + \geq 5 \text{ jets}$, 0 b -jets, Tglu1B, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$, $m_{\tilde{\chi}_1^\pm} = 0.5(m_{\tilde{g}} + m_{\tilde{\chi}_1^0})$		OCCUR=2
>2200	95	8 AAD	22U ATLS	$\tilde{g} \rightarrow q q \tilde{\chi}_1^0$, $q q \tilde{\chi}_1^\pm$, $m_{\tilde{\chi}_1^\pm} = 1000 \text{ GeV}$, $\tau(\tilde{\chi}_1^\pm) = 1 \text{ ns}$		
>2330	95	9 TUMASYAN	22V CMS	3 or 4 b -tagged jets or 2 large-radius jets, \cancel{E}_T ; Tglu1I; $m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$		
>2200	95	10 AAD	21AK ATLS	$\ell^\pm + \text{jets} + \cancel{E}_T$, Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 400 \text{ GeV}$		OCCUR=2
none 1300–2050	95	10 AAD	21AK ATLS	$\ell^\pm + \text{jets} + \cancel{E}_T$, Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 1000 \text{ GeV}$		OCCUR=2
>2300	95	11 AAD	21L ATLS	jets + \cancel{E}_T , Tglu1A, $m_{\tilde{\chi}_1^0} < 200 \text{ GeV}$		
>3000	95	11 AAD	21L ATLS	jets + \cancel{E}_T , combined $\tilde{g}\tilde{g}$, $\tilde{g}\tilde{q}$, $\tilde{q}\tilde{q}$ production, $\tilde{g} \rightarrow q q' \tilde{\chi}_1^0$, $\tilde{q} \rightarrow q \tilde{\chi}_1^0$, $m_{\tilde{q}} = m_{\tilde{g}}$, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$		OCCUR=2
>2200	95	11 AAD	21L ATLS	jets + \cancel{E}_T , Tglu1B, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$		OCCUR=3
>1400	95	12 AAD	21X ATLS	jets in empty bunch crossings, Tglu1A, long-lived R-hadron, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$, $10^{-5} \text{ s} < \tau_{\text{R-hadron}} < 10^3 \text{ s}$		

> 870	95	12 AAD	21X ATLS	jets in empty bunch crossings, Tglu1A, long-lived R-hadron, $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} = 100 \text{ GeV}, 10^{-5}$	OCCUR=2
				$s < \tau_{R\text{-hadron}} < 10^3 \text{ s}$	
>2260	95	13 SIRUNYAN	21AD CMS	jets + \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} < 1050 \text{ GeV}$	
>2150	95	13 SIRUNYAN	21AD CMS	jets + \cancel{E}_T , Tglu3C, $m_{\tilde{\chi}_1^0} = 600 \text{ GeV}, m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 20 \text{ GeV}$	OCCUR=2
>2250	95	13 SIRUNYAN	21AD CMS	jets + \cancel{E}_T , Tglu3D, $m_{\tilde{\chi}_1^0} = 700 \text{ GeV}, m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$	OCCUR=3
>1870	95	14 SIRUNYAN	21M CMS	$\ell^\pm \ell^\mp + \cancel{E}_T$, Tglu4C, $m_{\tilde{\chi}_1^0} = 1100 \text{ GeV}$	
>1980	95	15 AAD	20AL ATLS	8 or more jets + \cancel{E}_T , Tglu1E, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
>1820	95	15 AAD	20AL ATLS	8 or more jets + \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=2
>1600	95	16 AAD	20V ATLS	same-sign $\ell^\pm \ell^\pm$ + jets, Tglu1E, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
>1975	95	17 SIRUNYAN	20B CMS	$\geq 1\gamma + \cancel{E}_T$, Tglu4A, $\text{BR}(\tilde{g} \rightarrow qq\tilde{\chi}_1^\pm) = 0.5, m_{\tilde{\chi}_1^0} \simeq m_{\tilde{g}}$	OCCUR=9
>1920	95	18 SIRUNYAN	20BJ CMS	jets + \cancel{E}_T , Tglu1H, $m_{\tilde{g}} - m_{\tilde{\chi}_2^0} = 50 \text{ GeV}, m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$	
>2150	95	19 SIRUNYAN	20E CMS	1ℓ + jets, Tglu3A, $m_{\tilde{\chi}_1^0} < 700 \text{ GeV}$	
>2050	95	19 SIRUNYAN	20E CMS	1ℓ + jets, Tglu3A, $m_{\tilde{\chi}_1^0} < 1100 \text{ GeV}$	OCCUR=2
>1650	95	19 SIRUNYAN	20E CMS	1ℓ + jets, Tglu3C, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}, m_{\tilde{\chi}_1^0} < 1150 \text{ GeV}$	OCCUR=3
>1700	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu3A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1610	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu3B, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1300	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu3C, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 20 \text{ GeV}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=3
>1500	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu3D, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 5 \text{ GeV}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=4
>1350	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu1C, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=5
>1250	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu1C, $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 20 \text{ GeV}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=6
>1425	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=7
>1425	95	20 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm$ + jets, Tglu1B, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 20 \text{ GeV}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=8
>2000	95	21 AABOUD	19I ATLAS	≥ 2 jets + 1 or 2 τ + \cancel{E}_T , Tglu1F, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
>1860	95	22 SIRUNYAN	19AG CMS	$2\gamma + \cancel{E}_T$, Tglu4B, 500 GeV $< m_{\tilde{\chi}_1^0} < 1500 \text{ GeV}$	

>1920	95	23	SIRUNYAN	19AU CMS	γ + jets + b -jets + \cancel{E}_T , Tglu4D, $m_{\tilde{\chi}_1^0} = 127$ GeV	
>1950	95	23	SIRUNYAN	19AU CMS	γ + jets + b -jets + \cancel{E}_T , Tglu4E, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=2
>1800	95	23	SIRUNYAN	19AU CMS	γ + jets + b -jets + \cancel{E}_T , Tglu4F, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=3
>2090	95	23	SIRUNYAN	19AU CMS	γ + jets + b -jets + \cancel{E}_T , Tglu4D, $m_{\tilde{\chi}_1^0} = 1200$ GeV	OCCUR=4
>2120	95	23	SIRUNYAN	19AU CMS	γ + jets + b -jets + \cancel{E}_T , Tglu4E, $m_{\tilde{\chi}_1^0} = 1200$ GeV	OCCUR=5
>1970	95	23	SIRUNYAN	19AU CMS	γ + jets + b -jets + \cancel{E}_T , Tglu4F, $m_{\tilde{\chi}_1^0} = 1200$ GeV	OCCUR=6
>1700	95	24	SIRUNYAN	19CE CMS	2 jets, Stealth SUSY, Tglu1A and $\tilde{\chi}_1^0 \rightarrow \tilde{S} \gamma$ ($\tilde{S} \rightarrow S \tilde{G}$), $m_{\tilde{\chi}_1^0}$ $= 200$ GeV	
>2000	95	25	SIRUNYAN	19CH CMS	jets+ \cancel{E}_T , Tglu1A, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>2030	95	25	SIRUNYAN	19CH CMS	jets+ \cancel{E}_T , Tglu1C, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0} =$ $0.5(m_{\tilde{g}} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2
>2270	95	25	SIRUNYAN	19CH CMS	jets+ \cancel{E}_T , Tglu2A, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=3
>2180	95	25	SIRUNYAN	19CH CMS	jets+ \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>1750	95	26	SIRUNYAN	19K CMS	$\gamma + \ell + \cancel{E}_T$, Tglu4A, $m_{\tilde{\chi}_1^0} = 1500$ GeV	
>2000	95	27	SIRUNYAN	19S CMS	1 or 2 ℓ + jets + \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} < 700$ GeV	
>1900	95	27	SIRUNYAN	19S CMS	1 or 2 ℓ + jets + \cancel{E}_T , Tglu3C, 150 GeV $< m_{\tilde{\chi}_1^0} < 950$ GeV	OCCUR=2
>1970	95	28	AABOUD	18AR ATLS	jets+ $\geq 3b$ -jets+ \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} < 300$ GeV	
>1920	95	29	AABOUD	18AR ATLS	jets+ $\geq 3b$ -jets+ \cancel{E}_T , Tglu2A, $m_{\tilde{\chi}_1^0} < 600$ GeV	OCCUR=2
>1650	95	30	AABOUD	18AS ATLS	≥ 4 jets and disappearing tracks from $\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm$, modified Tglu1A or Tglu1B, $\tilde{\chi}^\pm$ life- time 0.2 ns, $m_{\tilde{\chi}^\pm} = 460$ GeV	
>1850	95	31	AABOUD	18BJ ATLS	$\ell^\pm \ell^\mp$ + jets + \cancel{E}_T , Tglu1G, $m_{\tilde{\chi}_1^0} = 100$ GeV	
>1650	95	32	AABOUD	18BJ ATLS	$\ell^\pm \ell^\mp$ + jets + \cancel{E}_T , Tglu1H, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=2
>2150	95	33	AABOUD	18U ATLS	2 γ + \cancel{E}_T , GGM, Tglu4B, any NLSP mass	
>1600	95	34	AABOUD	18U ATLS	γ + jets + \cancel{E}_T , GGM higgsino- bino, mix of Tglu4B and Tglu4C, any NLSP mass	OCCUR=2
>2030	95	35	AABOUD	18V ATLS	jets+ \cancel{E}_T , Tglu1A, $m_{\tilde{\chi}_1^0} = 0$ GeV	
>1980	95	36	AABOUD	18V ATLS	jets+ \cancel{E}_T , Tglu1B, $m_{\tilde{\chi}_1^\pm} = 0.5(m_{\tilde{g}} + m_{\tilde{\chi}_1^0})$, $m_{\tilde{\chi}_1^0}$ $= 0$ GeV	OCCUR=2
>1750	95	37	AABOUD	18V ATLS	jets+ \cancel{E}_T , Tglu1C, $m_{\tilde{\chi}_1^0} = 1$ GeV, any $m_{\tilde{\chi}_2^0} > 100$ GeV	OCCUR=3
>2000	95	38	SIRUNYAN	18AA CMS	$\geq 1\gamma$ + \cancel{E}_T , Tglu4A	
>2100	95	38	SIRUNYAN	18AA CMS	$\geq 1\gamma$ + \cancel{E}_T , Tglu4B	OCCUR=2
>1800	95	39	SIRUNYAN	18AC CMS	1 ℓ +jets, Tglu3A, $m_{\tilde{\chi}_1^0} < 650$ GeV	
>1700	95	39	SIRUNYAN	18AC CMS	1 ℓ +jets, Tglu3A, $m_{\tilde{\chi}_1^0} < 1040$ GeV	OCCUR=2
>1900	95	39	SIRUNYAN	18AC CMS	1 ℓ + jets, Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}}$ $+ m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 300$ GeV	OCCUR=3

>1250	95	39	SIRUNYAN	18AC CMS	$1\ell + \text{jets, Tglu1B, } m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2, m_{\tilde{\chi}_1^0} < 950 \text{ GeV}$	OCCUR=4
>1610	95	40	SIRUNYAN	18AL CMS	$\geq 3\ell^\pm + \text{jets} + \cancel{E}_T, \text{Tglu3A, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1160	95	40	SIRUNYAN	18AL CMS	$\geq 3\ell^\pm + \text{jets} + \cancel{E}_T, \text{Tglu1C, } m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1500	95	41	SIRUNYAN	18AR CMS	$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T, \text{GMSB, Tglu4C, } m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
>1770	95	41	SIRUNYAN	18AR CMS	$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T, \text{GMSB, Tglu4C, } m_{\tilde{\chi}_1^0} = 1400 \text{ GeV}$	OCCUR=2
>1625	95	42	SIRUNYAN	18AY CMS	$\text{jets} + \cancel{E}_T, \text{Tglu1A, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1825	95	42	SIRUNYAN	18AY CMS	$\text{jets} + \cancel{E}_T, \text{Tglu2A, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1625	95	42	SIRUNYAN	18AY CMS	$\text{jets} + \cancel{E}_T, \text{Tglu3A, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=3
>2040	95	43	SIRUNYAN	18D CMS	$\text{top quark (hadronically decaying)} + \text{jets} + \cancel{E}_T, \text{Tglu3A, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1930	95	43	SIRUNYAN	18D CMS	$\text{top quark (hadronically decaying)} + \text{jets} + \cancel{E}_T, \text{Tglu3B, } m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV, } m_{\tilde{\chi}_1^0} = 200 \text{ GeV}$	OCCUR=2
>1690	95	43	SIRUNYAN	18D CMS	$\text{top quark (hadronically decaying)} + \text{jets} + \cancel{E}_T, \text{Tglu3C, } m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 20 \text{ GeV, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=3
>1990	95	43	SIRUNYAN	18D CMS	$\text{top quark (hadronically decaying)} + \text{jets} + \cancel{E}_T, \text{Tglu3E, } m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 5 \text{ GeV, } m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=4
>2010	95	44	SIRUNYAN	18M CMS	$\geq 1 H (\rightarrow bb) + \cancel{E}_T, \text{Tglu1I}$	
>1825	95	44	SIRUNYAN	18M CMS	$\geq 1 H (\rightarrow bb) + \cancel{E}_T, \text{Tglu1J}$	OCCUR=2
>1750	95	45	AABOUD	17AJ ATLS	$\text{same-sign } \ell^\pm \ell^\pm / 3 \ell + \text{jets} + \cancel{E}_T, \text{Tglu3A, } m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
>1570	95	46	AABOUD	17AJ ATLS	$\text{same-sign } \ell^\pm \ell^\pm / 3 \ell + \text{jets} + \cancel{E}_T, \text{Tglu1E, } m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=2
>1860	95	47	AABOUD	17AJ ATLS	$\text{same-sign } \ell^\pm \ell^\pm / 3 \ell + \text{jets} + \cancel{E}_T, \text{Tglu1G, } m_{\tilde{\chi}_1^0} = 200 \text{ GeV}$	OCCUR=3
>2100	95	48	AABOUD	17AR ATLS	$1\ell + \text{jets} + \cancel{E}_T, \text{Tglu1B, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1740	95	49	AABOUD	17AR ATLS	$1\ell + \text{jets} + \cancel{E}_T, \text{Tglu1E, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1800	95	50	AABOUD	17AY ATLS	$\text{jets} + \cancel{E}_T, \text{Tglu3A, } m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$	
>1800	95	51	AABOUD	17AZ ATLS	$\geq 7 \text{ jets} + \cancel{E}_T, \text{large R-jets and/or } b\text{-jets, Tglu1E, } m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
>1540	95	52	AABOUD	17AZ ATLS	$\geq 7 \text{ jets} + \cancel{E}_T, \text{large R-jets and/or } b\text{-jets, Tglu3A, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1340	95	53	AABOUD	17N ATLS	$2 \text{ same-flavor, opposite-sign } \ell + \text{jets} + \cancel{E}_T, \text{Tglu1H, } m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1310	95	54	AABOUD	17N ATLS	$2 \text{ same-flavor, opposite-sign } \ell + \text{jets} + \cancel{E}_T, \text{Tglu1H, } m_{\tilde{\chi}_2^0} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2, m_{\tilde{\chi}_1^0} < 400 \text{ GeV}$	OCCUR=2
>1700	95	55	AABOUD	17N ATLS	$2 \text{ same-flavor, opposite-sign } \ell + \text{jets} + \cancel{E}_T, \text{Tglu1G, } m_{\tilde{\chi}_1^0} \sim 1 \text{ GeV}$	OCCUR=3
>1400	95	56	KHACHATRY...17	CMS	$\text{jets} + \cancel{E}_T, \text{Tglu1A, } m_{\tilde{\chi}_1^0} = 200 \text{ GeV}$	
>1650	95	56	KHACHATRY...17	CMS	$\text{jets} + \cancel{E}_T, \text{Tglu2A, } m_{\tilde{\chi}_1^0} = 200 \text{ GeV}$	OCCUR=2

>1600	95	56	KHACHATRY...17	CMS	jets+ \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0}=200\text{GeV}$	OCCUR=3	
>1550	95	57	KHACHATRY...17AD	CMS	jets+b-jets+ \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} =$		
>1450	95	58	KHACHATRY...17AD	CMS	0 GeV jets+b-jets+ \cancel{E}_T , Tglu3C, $200 <$ $m_{\tilde{\chi}_1^0} < 400 \text{ GeV}$	OCCUR=2	
>1570	95	59	KHACHATRY...17AS	CMS	1ℓ , Tglu3A, $m_{\tilde{\chi}_1^0} < 600 \text{ GeV}$		
>1500	95	59	KHACHATRY...17AS	CMS	1ℓ , Tglu3A, $m_{\tilde{\chi}_1^0} < 775 \text{ GeV}$	OCCUR=2	
>1400	95	59	KHACHATRY...17AS	CMS	1ℓ , Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} +$ $m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 725 \text{ GeV}$	OCCUR=3	
none 1050-1350	95	59	KHACHATRY...17AS	CMS	1ℓ , Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} +$ $m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} < 850 \text{ GeV}$	OCCUR=4	
>1175	95	60	KHACHATRY...17AW	CMS	$\geq 3\ell^\pm$, 2 jets, Tglu3A, $m_{\tilde{\chi}_1^0} = 0$		
> 825	95	60	KHACHATRY...17AW	CMS	GeV $\geq 3\ell^\pm$, 2 jets, Tglu1C, $m_{\tilde{\chi}_1^\pm}$ $= (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=2	
>1350	95	61	KHACHATRY...17P	CMS	1 or more jets+ \cancel{E}_T , Tglu1A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$		
>1545	95	61	KHACHATRY...17P	CMS	1 or more jets+ \cancel{E}_T , Tglu2A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2	
>1120	95	61	KHACHATRY...17P	CMS	1 or more jets+ \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=3	
>1300	95	61	KHACHATRY...17P	CMS	1 or more jets+ \cancel{E}_T , Tglu3D, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 5 \text{ GeV}$, $m_{\tilde{\chi}_1^0}$ $= 100 \text{ GeV}$	OCCUR=4	
> 780	95	61	KHACHATRY...17P	CMS	1 or more jets+ \cancel{E}_T , Tglu3B, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}$, $m_{\tilde{\chi}_1^0}$ $= 50 \text{ GeV}$	OCCUR=5	
> 790	95	61	KHACHATRY...17P	CMS	1 or more jets+ \cancel{E}_T , Tglu3C, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 20 \text{ GeV}$, $m_{\tilde{\chi}_1^0}$ $= 0 \text{ GeV}$	OCCUR=6	
>1650	95	62	KHACHATRY...17V	CMS	$2\gamma + \cancel{E}_T$, GGM, Tglu4B, any NLSP mass		
>1900	95	63	SIRUNYAN	17AF	CMS	$1\ell + \text{jets} + b\text{-jets} + \cancel{E}_T$, Tglu3A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1600	95	63	SIRUNYAN	17AF	CMS	$1\ell + \text{jets} + b\text{-jets} + \cancel{E}_T$, Tglu3B, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175 \text{ GeV}$, $m_{\tilde{\chi}_1^0}$ $= 50 \text{ GeV}$	OCCUR=2
>1800	95	64	SIRUNYAN	17AY	CMS	$\gamma + \text{jets} + \cancel{E}_T$, Tglu4B, $m_{\tilde{\chi}_1^0} = 0$	
>1600	95	64	SIRUNYAN	17AY	CMS	GeV $\gamma + \text{jets} + \cancel{E}_T$, Tglu4A, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=2
>1860	95	65	SIRUNYAN	17AZ	CMS	GeV $\geq 1 \text{ jets} + \cancel{E}_T$, Tglu1A, $m_{\tilde{\chi}_1^0} =$	
>2025	95	65	SIRUNYAN	17AZ	CMS	0 GeV $\geq 1 \text{ jets} + \cancel{E}_T$, Tglu2A, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=2
>1900	95	65	SIRUNYAN	17AZ	CMS	GeV $\geq 1 \text{ jets} + \cancel{E}_T$, Tglu3A, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
>1825	95	66	SIRUNYAN	17P	CMS	GeV jets+ \cancel{E}_T , Tglu1A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1950	95	66	SIRUNYAN	17P	CMS	GeV jets+ \cancel{E}_T , Tglu2A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1960	95	66	SIRUNYAN	17P	CMS	GeV jets+ \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=3
>1800	95	66	SIRUNYAN	17P	CMS	GeV jets+ \cancel{E}_T , Tglu1C, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0}$ $= (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=4
>1870	95	66	SIRUNYAN	17P	CMS	GeV jets+ \cancel{E}_T , Tglu3D, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$ $+ 5 \text{ GeV}$, $m_{\tilde{\chi}_1^0} = 1000 \text{ GeV}$	OCCUR=5
>1520	95	67	SIRUNYAN	17S	CMS	GeV same-sign $\ell^\pm \ell^\pm + \text{jets} + \cancel{E}_T$, Tglu3A, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	

>1200	95	67 SIRUNYAN	17S	CMS	same-sign $\ell^\pm \ell^\pm + \text{jets} + \cancel{E}_T$, Tglu3D, $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 5$ GeV, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=2
>1370	95	67 SIRUNYAN	17S	CMS	same-sign $\ell^\pm \ell^\pm + \text{jets} + \cancel{E}_T$, Tglu3B, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 175$ GeV, $m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=3
>1180	95	67 SIRUNYAN	17S	CMS	same-sign $\ell^\pm \ell^\pm + \text{jets} + \cancel{E}_T$, Tglu3C, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 20$ GeV, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=4
>1280	95	67 SIRUNYAN	17S	CMS	same-sign $\ell^\pm \ell^\pm + \text{jets} + \cancel{E}_T$, Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} +$ $m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=5
>1300	95	67 SIRUNYAN	17S	CMS	same-sign $\ell^\pm \ell^\pm + \text{jets} + \cancel{E}_T$, Tglu1B, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 20$ GeV, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=6
>1570	95	68 AABOUD	16AC	ATLS	≥ 2 jets + 1 or 2 $\tau + \cancel{E}_T$, Tglu1F, $m_{\tilde{\chi}_1^0} = 100$ GeV	
>1460	95	69 AABOUD	16J	ATLS	1 $\ell^\pm + \geq 4$ jets + \cancel{E}_T , Tglu3C, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 5$ GeV	
>1650	95	70 AABOUD	16M	ATLS	2 $\gamma + \cancel{E}_T$, Tglu1D, any NLSP mass	
>1510	95	71 AABOUD	16N	ATLS	≥ 4 jets + \cancel{E}_T , Tglu1A, $m_{\tilde{\chi}_1^0} =$ 0 GeV	
>1500	95	72 AABOUD	16N	ATLS	≥ 4 jets + \cancel{E}_T , Tglu1B, $m_{\tilde{\chi}_1^\pm} =$ $(m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 200$ GeV	OCCUR=2
>1780	95	73 AAD	16AD	ATLS	0 ℓ , ≥ 3 b-jets + \cancel{E}_T , Tglu2A, $m_{\tilde{\chi}_1^0} < 800$ GeV	
>1760	95	74 AAD	16AD	ATLS	1 ℓ , ≥ 3 b-jets + \cancel{E}_T , Tglu3A, $m_{\tilde{\chi}_1^0} < 700$ GeV	OCCUR=2
>1300	95	75 AAD	16BB	ATLS	2 same-sign/3 $\ell + \text{jets} + \cancel{E}_T$, Tglu1D, $m_{\tilde{\chi}_1^0} < 600$ GeV	
>1100	95	75 AAD	16BB	ATLS	2 same-sign/3 $\ell + \text{jets} + \cancel{E}_T$, Tglu1E, $m_{\tilde{\chi}_1^0} < 300$ GeV	OCCUR=2
>1200	95	75 AAD	16BB	ATLS	2 same-sign /3 $\ell + \text{jets} + \cancel{E}_T$, Tglu3A, $m_{\tilde{\chi}_1^0} < 600$ GeV	OCCUR=3
>1600		76 AAD	16BG	ATLS	1 ℓ , ≥ 4 jets, \cancel{E}_T , Tglu1B, $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$, $m_{\tilde{\chi}_1^0} = 100$ GeV	
>1400	95	77 AAD	16V	ATLS	≥ 7 to ≥ 10 jets + \cancel{E}_T , Tglu1E, $m_{\tilde{\chi}_1^0} < 200$ GeV	
>1400	95	77 AAD	16V	ATLS	≥ 7 to ≥ 10 jets + \cancel{E}_T , pMSSM $M_1 = 60$ GeV, M_2 $= 3$ TeV, $\tan\beta=10$, $\mu < 0$	OCCUR=2
>1100	95	78 KHACHATRY...16AM	CMS	boosted $W+b$, Tglu3C, $m_{\tilde{t}_1} -$ $m_{\tilde{\chi}_1^0} < 80$ GeV, $m_{\tilde{\chi}_1^0} < 400$ GeV		
> 700	95	78 KHACHATRY...16AM	CMS	boosted $W+b$, Tglu3B, $m_{\tilde{t}_1} -$ $m_{\tilde{\chi}_1^0} = 175$ GeV, $m_{\tilde{\chi}_1^0} = 0$ GeV	OCCUR=2	
>1050	95	79 KHACHATRY...16BJ	CMS	same-sign $\ell^\pm \ell^\pm$, Tglu3A, $m_{\tilde{\chi}_1^0} < 800$ GeV	OCCUR=2	
>1300	95	79 KHACHATRY...16BJ	CMS	same-sign $\ell^\pm \ell^\pm$, Tglu3A, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=3	
>1140	95	79 KHACHATRY...16BJ	CMS	same-sign $\ell^\pm \ell^\pm$, Tglu3B, $m_{\tilde{t}_1} -$ $m_{\tilde{\chi}_1^0} = 20$ GeV, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=5	
> 850	95	79 KHACHATRY...16BJ	CMS	same-sign $\ell^\pm \ell^\pm$, Tglu3B, $m_{\tilde{t}_1} -$ $m_{\tilde{\chi}_1^0} = 20$ GeV, $m_{\tilde{\chi}_1^0} < 700$ GeV	OCCUR=6	
> 950	95	79 KHACHATRY...16BJ	CMS	same-sign $\ell^\pm \ell^\pm$, Tglu3D, $m_{\tilde{\chi}_1^\pm}$ $= m_{\tilde{\chi}_1^0} + 5$ GeV	OCCUR=7	

>1100	95	79	KHACHATRY...16BJ CMS	same-sign $\ell^\pm \ell^\pm, \text{Tglu1B}, m_{\tilde{\chi}_1^\pm} = 0.5(m_{\tilde{g}} + m_{\tilde{\chi}_1^0}), m_{\tilde{\chi}_1^0} < 400 \text{ GeV}$	OCCUR=8
> 830	95	79	KHACHATRY...16BJ CMS	same-sign $\ell^\pm \ell^\pm, \text{Tglu1B}, m_{\tilde{\chi}_1^\pm} = 0.5(m_{\tilde{g}} + m_{\tilde{\chi}_1^0}), m_{\tilde{\chi}_1^0} < 700 \text{ GeV}$	OCCUR=9
>1300	95	79	KHACHATRY...16BJ CMS	same-sign $\ell^\pm \ell^\pm, \text{Tglu3B}, m_{\tilde{\tau}} - m_{\tilde{\chi}_1^0} = m_t, m_{\tilde{\chi}_1^0} = 0$	OCCUR=10
>1050	95	79	KHACHATRY...16BJ CMS	same-sign $\ell^\pm \ell^\pm, \text{Tglu3B}, m_{\tilde{\tau}} - m_{\tilde{\chi}_1^0} = m_t, m_{\tilde{\chi}_1^0} < 800 \text{ GeV}$	OCCUR=11
>1725	95	80	KHACHATRY...16BS CMS	jets + $\cancel{E}_T, \text{Tglu1A}, m_{\tilde{\chi}_1^0} = 0$	
>1750	95	80	KHACHATRY...16BS CMS	jets + $\cancel{E}_T, \text{Tglu2A}, m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
>1550	95	80	KHACHATRY...16BS CMS	jets + $\cancel{E}_T, \text{Tglu3A}, m_{\tilde{\chi}_1^0} = 0$	OCCUR=4
>1280	95	81	KHACHATRY...16BY CMS	opposite-sign $\ell^\pm \ell^\pm, \text{Tglu4C}, m_{\tilde{\chi}_1^0} = 1000 \text{ GeV}$	
>1030	95	81	KHACHATRY...16BY CMS	opposite-sign $\ell^\pm \ell^\pm, \text{Tglu4C}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	OCCUR=2
>1440	95	82	KHACHATRY...16V CMS	jets + $\cancel{E}_T, \text{Tglu1A}, m_{\tilde{\chi}_1^0} = 0$	
>1600	95	82	KHACHATRY...16V CMS	jets + $\cancel{E}_T, \text{Tglu2A}, m_{\tilde{\chi}_1^0} = 0$	OCCUR=2
>1550	95	82	KHACHATRY...16V CMS	jets + $\cancel{E}_T, \text{Tglu3A}, m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
>1450	95	82	KHACHATRY...16V CMS	jets + $\cancel{E}_T, \text{Tglu1C}, m_{\tilde{\chi}_1^0} = 0$	OCCUR=4
> 820	95	83	AAD 15BG ATLS	GGM, $\tilde{g} \rightarrow q\tilde{q}Z\tilde{G}, \tan\beta = 30, \mu > 600 \text{ GeV}$	
> 850	95	83	AAD 15BG ATLS	GGM, $\tilde{g} \rightarrow q\tilde{q}Z\tilde{G}, \tan\beta = 1.5, \mu > 450 \text{ GeV}$	OCCUR=2
>1150	95	84	AAD 15BV ATLS	general RPC \tilde{g} decays, $m_{\tilde{\chi}_1^0} < 100 \text{ GeV}$	
> 700	95	85	AAD 15BX ATLS	$\tilde{g} \rightarrow X\tilde{\chi}_1^0$, independent of $m_{\tilde{\chi}_1^0}$	
>1290	95	86	AAD 15CA ATLS	$\geq 2 \gamma + \cancel{E}_T$, GGM, bino-like NLSP, any NLSP mass	
>1260	95	86	AAD 15CA ATLS	$\geq 1 \gamma + b\text{-jets} + \cancel{E}_T$, GGM, higgsino-bino admix. NLSP and $\mu < 0, m(\text{NLSP}) > 450 \text{ GeV}$	OCCUR=2
>1140	95	86	AAD 15CA ATLS	$\geq 1 \gamma + \text{jets} + \cancel{E}_T$, GGM, higgsino-bino admixture NLSP, all $\mu > 0$	OCCUR=3
>1225	95	87	KHACHATRY...15AF CMS	$\tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0$	
>1300	95	87	KHACHATRY...15AF CMS	$\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0$	OCCUR=2
>1225	95	87	KHACHATRY...15AF CMS	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0$	OCCUR=3
>1550	95	87	KHACHATRY...15AF CMS	CMSSM, $\tan\beta=30, m_{\tilde{g}}=m_{\tilde{q}}, A_0=-2\max(m_0, m_{1/2}), \mu > 0$	OCCUR=4
>1150	95	87	KHACHATRY...15AF CMS	CMSSM, $\tan\beta=30, A_0=-2\max(m_0, m_{1/2}), \mu > 0$	OCCUR=5
>1280	95	88	KHACHATRY...15I CMS	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 0$	
>1310	95	89	KHACHATRY...15X CMS	$\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	
>1175	95	89	KHACHATRY...15X CMS	$\tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=2
>1330	95	90	AAD 14AE ATLS	jets + $\cancel{E}_T, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$	
>1700	95	90	AAD 14AE ATLS	jets + \cancel{E}_T , mSUGRA/CMSSM, $m_{\tilde{q}} = m_{\tilde{g}}$	OCCUR=2
>1090	95	91	AAD 14AG ATLS	$\tau + \text{jets} + \cancel{E}_T$, natural Gauge Mediation	
>1600	95	91	AAD 14AG ATLS	$\tau + \text{jets} + \cancel{E}_T$, mGMSB, $M_{mess} = 250 \text{ GeV}, N_5 = 3, \mu > 0, C_{grav} = 1$	OCCUR=2
> 640	95	92	AAD 14X ATLS	$\geq 4\ell^\pm, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \tilde{G}, \tan\beta = 30$, GGM	OCCUR=2
>1000	95	93	CHATRCHYAN 14AH CMS	jets + $\cancel{E}_T, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$	

>1350	95	93	CHATRCHYAN 14AH CMS	jets + \cancel{E}_T , CMSSM, $m_{\tilde{g}} = m_{\tilde{q}}$	OCCUR=2
>1000	95	94	CHATRCHYAN 14AH CMS	jets + \cancel{E}_T , $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=3
>1000	95	95	CHATRCHYAN 14AH CMS	jets + \cancel{E}_T , $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=4
>1160	95	96	CHATRCHYAN 14I CMS	jets + \cancel{E}_T , $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} < 100$ GeV	
>1130	95	96	CHATRCHYAN 14I CMS	multijets + \cancel{E}_T , $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} < 100$ GeV	OCCUR=2
>1210	95	96	CHATRCHYAN 14I CMS	multijets + \cancel{E}_T , $\tilde{g} \rightarrow q\bar{q}W/Z\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} < 100$ GeV	OCCUR=3
>1260	95	97	CHATRCHYAN 14N CMS	$1\ell^\pm + \text{jets} + \geq 2b\text{-jets}$, $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} = 0$ GeV, $m_{\tilde{t}} > m_{\tilde{g}}$	
		98	CHATRCHYAN 14R CMS	$\geq 3\ell^\pm$, $(\tilde{g}/\tilde{q}) \rightarrow q\ell^\pm\ell^\mp\tilde{G}$ simplified model, GMSB, slepton co-NLSP scenario	
		99	CHATRCHYAN 14R CMS	$\geq 3\ell^\pm$, $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ simplified model	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>1500	95	100	AABOUD 18BJ ATLS	$\ell^\pm\ell^\mp + \text{jets} + \cancel{E}_T$, Tglu1H, $m_{\tilde{\chi}_1^0} = 1$ GeV, any $m_{\tilde{\chi}_2^0}$	OCCUR=3
>1770	95	101	AABOUD 18V ATLS	jets+ \cancel{E}_T , Tglu1C-like, 1/2 BR per decay mode, any $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$, $m_{\tilde{\chi}_1^0} = 60$ GeV	OCCUR=4
>1600	95	102	AABOUD 17AZ ATLS	≥ 7 jets+ \cancel{E}_T , large R-jets and/or b-jets, pMSSM, $m_{\tilde{\chi}_1^\pm} = 200$ GeV	OCCUR=4
>1600	95	103	KHACHATRY...16AY CMS	$1\ell^\pm + \text{jets} + b\text{-jets} + \cancel{E}_T$, Tglu3A, $m_{\tilde{\chi}_1^0} = 0$ GeV	
> 500	95	104	KHACHATRY...16BT CMS	19-parameter pMSSM model, global Bayesian analysis, flat prior	
		105	AAD 15AB ATLS	$\tilde{g} \rightarrow \tilde{S}g$, $c\tau = 1$ m, $\tilde{S} \rightarrow S\tilde{G}$ and $S \rightarrow gg$, BR = 100%	
		106	AAD 15AI ATLS	$\ell^\pm + \text{jets} + \cancel{E}_T$	
>1600	95	84	AAD 15BV ATLS	pMSSM, $M_1 = 60$ GeV, $m_{\tilde{q}} < 1500$ GeV	OCCUR=2
>1280	95	84	AAD 15BV ATLS	mSUGRA, $m_0 > 2$ TeV	OCCUR=3
>1100	95	84	AAD 15BV ATLS	via $\tilde{\tau}$, natural GMSB, all $m_{\tilde{\tau}}$	OCCUR=4
>1330	95	84	AAD 15BV ATLS	jets + \cancel{E}_T , $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=5
>1500	95	84	AAD 15BV ATLS	jets + \cancel{E}_T , $\tilde{g} \rightarrow \tilde{q}q$, $\tilde{q} \rightarrow q\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=6
>1650	95	84	AAD 15BV ATLS	jets + \cancel{E}_T , $m_{\tilde{g}} = m_{\tilde{q}}$, $m_{\tilde{\chi}_1^0} = 1$ GeV	OCCUR=7
> 850	95	84	AAD 15BV ATLS	jets + \cancel{E}_T , $\tilde{g} \rightarrow g\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} < 550$ GeV	OCCUR=8
>1270	95	84	AAD 15BV ATLS	jets + \cancel{E}_T , $\tilde{g} \rightarrow q\bar{q}W\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=9
>1150	95	84	AAD 15BV ATLS	jets + $\ell^\pm\ell^\pm$, $\tilde{g} \rightarrow q\bar{q}WZ\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=10
>1320	95	84	AAD 15BV ATLS	jets + $\ell^\pm\ell^\pm$, \tilde{g} decays via sleptons, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=11
>1220	95	84	AAD 15BV ATLS	τ , \tilde{q} decays via staus, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=12
>1310	95	84	AAD 15BV ATLS	b-jets, $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} < 400$ GeV	OCCUR=13
>1220	95	84	AAD 15BV ATLS	b-jets, $\tilde{g} \rightarrow \tilde{t}_1 t$ and $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$, $m_{T_1} < 1000$ GeV	OCCUR=14

>1180	95	84 AAD	15BV ATLS	b -jets, $\tilde{g} \rightarrow \tilde{t}_1 t$ and $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm, m_{T_1} < 1000$ GeV, $m_{\tilde{\chi}_1^0} = 60$ GeV	OCCUR=15
>1260	95	84 AAD	15BV ATLS	b -jets, $\tilde{g} \rightarrow \tilde{t}_1 t$ and $\tilde{g} \rightarrow c\tilde{\chi}_1^0$	OCCUR=16
>1200	95	84 AAD	15BV ATLS	b -jets, $\tilde{g} \rightarrow \tilde{b}_1 b$ and $\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, m_{\tilde{b}_1} < 1000$ GeV	OCCUR=18
>1250	95	84 AAD	15BV ATLS	b -jets, $\tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} < 400$ GeV	OCCUR=19
none, 750-1250	95	84 AAD	15BV ATLS	b -jets, \tilde{g} decay via offshell \tilde{t}_1 and $\tilde{b}_1, m_{\tilde{\chi}_1^0} < 500$ GeV	OCCUR=20
>1100	95	107 AAD	15CB ATLS	jets, $\tilde{g} \rightarrow qq\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow Z\tilde{G}$, GGM, $m_{\tilde{\chi}_1^0} = 400$ GeV and $3 < c\tau_{\tilde{\chi}_1^0} < 500$ mm	OCCUR=3
>1400	95	107 AAD	15CB ATLS	jets or $\cancel{E}_T, \tilde{g} \rightarrow qq\tilde{\chi}_1^0$, Split SUSY, $m_{\tilde{\chi}_1^0} = 100$ GeV and $15 < c\tau < 300$ mm	OCCUR=4
>1500	95	107 AAD	15CB ATLS	$\cancel{E}_T, \tilde{g} \rightarrow qq\tilde{\chi}_1^0$, Split SUSY, $m_{\tilde{\chi}_1^0} = 100$ GeV and $20 < c\tau < 250$ mm	OCCUR=5
>1300	95	108 KHACHATRY...15AD CMS		$\ell^\pm \ell^\mp + \text{jets} + \cancel{E}_T$, GMSB, $\tilde{g} \rightarrow q\bar{q}Z\tilde{G} \geq 2 \gamma, \geq 1$ jet, (Razor), bino-like NLSP, $m_{\tilde{\chi}_1^0} = 375$ GeV	
> 800	95	109 KHACHATRY...15AZ CMS		$\geq 1 \gamma, \geq 2$ jet, wino-like NLSP, $m_{\tilde{\chi}_1^0} = 375$ GeV	OCCUR=2
>1280	95	110 AAD	14AX ATLS	≥ 3 b -jets + \cancel{E}_T , CMSSM	
>1250	95	110 AAD	14AX ATLS	≥ 3 b -jets + $\cancel{E}_T, \tilde{g} \rightarrow \tilde{b}_1 b\tilde{\chi}_1^0$ simplified model, $\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 60$ GeV, $m_{\tilde{b}_1} < 900$ GeV	OCCUR=2
>1190	95	110 AAD	14AX ATLS	≥ 3 b -jets + $\cancel{E}_T, \tilde{g} \rightarrow \tilde{t}_1 t\tilde{\chi}_1^0$ simplified model, $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0, m_{\tilde{\chi}_1^0} = 60$ GeV, $m_{\tilde{t}_1} < 1000$ GeV	OCCUR=3
>1180	95	110 AAD	14AX ATLS	≥ 3 b -jets + $\cancel{E}_T, \tilde{g} \rightarrow \tilde{t}_1 t\tilde{\chi}_1^0$ simplified model, $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm, m_{\tilde{\chi}_1^\pm} = 2m_{\tilde{\chi}_1^0}, m_{\tilde{\chi}_1^0} = 60$ GeV, $m_{\tilde{t}_1} < 1000$ GeV	OCCUR=4
>1250	95	110 AAD	14AX ATLS	≥ 3 b -jets + $\cancel{E}_T, \tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} < 400$ GeV	OCCUR=5
>1340	95	110 AAD	14AX ATLS	≥ 3 b -jets + $\cancel{E}_T, \tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} < 400$ GeV	OCCUR=6
>1300	95	110 AAD	14AX ATLS	≥ 3 b -jets + $\cancel{E}_T, \tilde{g} \rightarrow t\tilde{b}\tilde{\chi}_1^\pm$ simplified model, $\tilde{\chi}_1^\pm \rightarrow f f' \tilde{\chi}_1^0, m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 2$ GeV, $m_{\tilde{\chi}_1^0} < 300$ GeV	OCCUR=7
> 950	95	111 AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell^\mp) + \text{jets}, \tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$ simplified model	
>1000	95	111 AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell^\mp) + \text{jets}, \tilde{g} \rightarrow t\tilde{t}_1$ with $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$ simplified model, $m_{\tilde{t}_1} < 200$ GeV, $m_{\tilde{\chi}_1^\pm} = 118$ GeV, $m_{\tilde{\chi}_1^0} = 60$ GeV	OCCUR=2
> 640	95	111 AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell^\mp) + \text{jets}, \tilde{g} \rightarrow t\tilde{t}_1$ with $\tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$ simplified model, $m_{\tilde{t}_1} = m_{\tilde{\chi}_1^0} + 20$ GeV	OCCUR=3

> 860	95	111 AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell^\mp) + \text{jets}, \tilde{g} \rightarrow qq' \tilde{\chi}_1^\pm,$ $\tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^\pm} = 2 m_{\tilde{\chi}_1^0},$ $m_{\tilde{\chi}_1^0} < 400 \text{ GeV}$	OCCUR=5
>1040	95	111 AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell^\mp) + \text{jets}, \tilde{g} \rightarrow qq' \tilde{\chi}_1^\pm,$ $\tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow$ $Z^{(*)} \tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^0} < 520 \text{ GeV}$	OCCUR=6
>1200	95	111 AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell^\mp) + \text{jets}, \tilde{g} \rightarrow$ $qq' \tilde{\chi}_1^\pm / \tilde{\chi}_2^0, \tilde{\chi}_1^\pm \rightarrow \ell^\pm \nu \tilde{\chi}_1^0,$ $\tilde{\chi}_2^0 \rightarrow \ell^\pm \ell^\mp (\nu\nu) \tilde{\chi}_1^0$ simplified model	OCCUR=7
>1050	95	112 CHATRCHYAN 14H	CMS	same-sign $\ell^\pm \ell^\pm, \tilde{g} \rightarrow t \bar{t} \tilde{\chi}_1^0$ simplified model, massless $\tilde{\chi}_1^0$	
> 900	95	113 CHATRCHYAN 14H	CMS	same-sign $\ell^\pm \ell^\pm, \tilde{g} \rightarrow qq' \tilde{\chi}_1^\pm,$ $\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^\pm} = 0.5 m_{\tilde{g}},$ mass-less $\tilde{\chi}_1^0$	OCCUR=2
>1050	95	114 CHATRCHYAN 14H	CMS	same-sign $\ell^\pm \ell^\pm, \tilde{g} \rightarrow b \bar{t} \tilde{\chi}_1^\pm,$ $\tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ simplified model, $m_{\tilde{\chi}_1^\pm} = 300 \text{ GeV}, m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$	OCCUR=3

¹ AAD 23AB searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for an excess of events with one photon, jets and \cancel{E}_T . No significant excess above the Standard Model predictions is observed. Limits are set on the mass of pair produced gluinos decaying to $\tilde{g} \rightarrow qq \tilde{\chi}_1^0$ followed by $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$ or $\tilde{\chi}_1^0 \rightarrow X \tilde{G}$ with equal probability, see Figure 4. X can be Z (left figure) or h (right figure).

NODE=S046GNO;LINKAGE=LF

² AAD 23AE searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 2 ℓ with same flavour and opposite sign, plus jets and \cancel{E}_T , defining signal region with the dilepton invariant mass both on- and off-shell with respect to the Z boson. No significant excess above the Standard Model predictions is observed. Limits are set on models of strong and electroweak production. In this case, limits are placed on the mass of pair-produced gluinos, assuming a scenario like in Tglu1G, see figure 16.

NODE=S046GNO;LINKAGE=JF

³ AAD 23AE searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 2 ℓ with same flavour and opposite sign, plus jets and \cancel{E}_T , defining signal region with the dilepton invariant mass both on- and off-shell with respect to the Z boson. No significant excess above the Standard Model predictions is observed. Limits are set on models of strong and electroweak production. In this case, limits are placed on the gluino mass assuming gluino pair production, assuming a scenario like in Tglu1H, see figure 16.

NODE=S046GNO;LINKAGE=KF

⁴ AAD 23AL searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 0 or 1 lepton and at least three b-tagged jets. No significant excess above the Standard Model prediction is observed. Results are interpreted in terms of gluino pair production followed by the decay of gluinos into off-shell third generation squarks, yielding final states with top and bottom quarks, and missing transverse momentum from a $\tilde{\chi}_1^0$ LSP. Limits are set on the mass of the gluino as a function of the $\tilde{\chi}_1^0$ assuming $B(\tilde{g} \rightarrow \tilde{t} t) = 100\%$ or $B(\tilde{g} \rightarrow \tilde{b} b) = 100\%$, see figure 10.

NODE=S046GNO;LINKAGE=HF

⁵ AAD 23AL searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 0 or 1 lepton and at least three b-tagged jets. No significant excess above the Standard Model prediction is observed. Results are interpreted in terms of gluino pair production followed by the decay of gluinos into off-shell third generation squarks, yielding final states with top and bottom quarks, and missing transverse momentum from a $\tilde{\chi}_1^0$ LSP. Limits are set on the mass of the gluino as a function of $m_{\tilde{\chi}_1^0}$, assuming $B(\tilde{g} \rightarrow \tilde{t} t) + B(\tilde{g} \rightarrow \tilde{b} b) + B(\tilde{g} \rightarrow t b \tilde{\chi}_1^\pm) = 100\%$, and $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 2 \text{ GeV}$, see figures 11–13.

NODE=S046GNO;LINKAGE=IF

⁶ HAYRAPETYAN 23E searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of gluino, top squark and electroweakino pair production in events with at least one photon, multiple jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set in models for strong production, Tglu4D, Tglu4E, Tglu4F and Tstop13, see their figure 9. They also interpret the results in the models for electroweak production, shown in their figure 10. Tchi1n1A assumes wino-like $\tilde{\chi}_1^\pm \tilde{\chi}_1^0$ production, while Tchi1chi1A assumes higgsino-like cross sections and includes $\tilde{\chi}_1^\pm \tilde{\chi}_1^\pm, \tilde{\chi}_1^0 \tilde{\chi}_2^0$ and $\tilde{\chi}_{1,2}^0 \tilde{\chi}_1^\pm$ production. For $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ alone no mass point can be excluded in the model Tchi1chi1A, but in another model for $\tilde{\chi}_1^0 \tilde{\chi}_2^0$ production, Tn1n2A.

NODE=S046GNO;LINKAGE=NF

- ⁷ TUMASYAN 23AY searched in 138 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of gluino pair production in events with a single electron or muon and multiple hadronic jets. No significant excess above the Standard Model expectations is observed. Limits are set in the models Tglu3A and Tglu1B, see their figure 11. For Tglu1B, the chargino mass is set to $m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})$.
NODE=S046GNO;LINKAGE=OF
- ⁸ AAD 22U searched for the signature of disappearing track from a long-lived chargino in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$. Long-lived charginos decay into quasi-degenerate neutralino emitting a low-momentum particle whose identification is not attempted. The signal is identified by requiring short tracklets in the four pixel layers with no continuation in the SCT (strip) detector. The main background from fake tracklets is estimated directly with the data. No significant excess above the background prediction is found. The results are interpreted in an AMSB scenario (win LSP), on $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\pm$ and $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}_1^0$, assuming $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm) = 100\%$, see their figure 7. Results are also interpreted in a higgsino-LSP model, with $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}^\mp$, and $pp \rightarrow \tilde{\chi}^\pm \tilde{\chi}_{1,2}^0$, assuming $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \pi^\pm) = 95.5\%$, $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 e^\pm) = 3\%$, $B(\tilde{\chi}^\pm \rightarrow \tilde{\chi}_1^0 \mu^\pm) = 1.5\%$, see their figure 8. Finally, results are interpreted in a simplified model of gluino pair production, with $pp \rightarrow \tilde{g} \tilde{g}$ and $B(\tilde{g} \rightarrow qq \tilde{\chi}_1^0) = B(\tilde{g} \rightarrow qq \tilde{\chi}^\pm) = B(\tilde{g} \rightarrow qq \tilde{\chi}^\mp) = 1/3$, see their figure 9.
NODE=S046GNO;LINKAGE=FF
- ⁹ TUMASYAN 22V searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for evidence of electroweakino pair production with decay to two Higgs bosons H , with $H \rightarrow b \bar{b}$, resulting either in 4 resolved b-jets or two large-radius jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the mass of $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ in the models Tn1n1A, see their Figures 11 and 12, or in a model where higgsino-like nearly mass degenerate $\tilde{\chi}_2^0$ and $\tilde{\chi}_3^0$ are pair produced and each decay to H and a bino-like $\tilde{\chi}_1^0$, see their Figure 13. Limits are also set on the gluino mass in the model Tglu1I, see their Figure 14.
NODE=S046GNO;LINKAGE=GF
- ¹⁰ AAD 21AK searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of gluinos and squarks in events with a single isolated electron or muon, originating from the decay of a W boson, multiple jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1B simplified model and on the squark mass in the Tsqk3 simplified model, see their Figure 8.
NODE=S046GNO;LINKAGE=ZE
- ¹¹ AAD 21L searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for pair production of gluinos and squarks in events with jets, large missing transverse momentum but no electrons or muons. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A and Tglu1B simplified models, on the squark mass in the Tsqk1 and Tsqk3 simplified models and in a simplified model for gluino-squark production, see their Figures 13-17.
NODE=S046GNO;LINKAGE=AF
- ¹² AAD 21X searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for the decay of long-lived R-hadrons stopped by the calorimeter, producing high-momentum jets resulting in large out-of-time energy deposits in the calorimeters. These decays are detected using data collected during periods in the LHC bunch structure when collisions are absent. No significant excess above the predicted background is observed. Limits are set on the R-hadron mass in the Tglu1A simplified model as a function of the R-hadron lifetime, for different $m_{\tilde{\chi}_1^0}$. See Figures 9, 10.
NODE=S046GNO;LINKAGE=BF
- ¹³ SIRUNYAN 21AD searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with multiple jets, no leptons, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the top squark mass in the simplified models Tstop1, Tstop2 with $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{t}} + m_{\tilde{\chi}_1^0})/2$, and a 50:50 mixture of these with $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, see their Figure 8. Limits are also set on the top squark mass for $10 \text{ GeV} < m_{\tilde{t}} - m_{\tilde{\chi}_1^\pm} < 80 \text{ GeV}$ in the simplified models Tstop2, Tstop 3, and Tstop4, see their Figure 9. For indirect top squark production, limits are set on the gluino mass in the simplified models Tglu3A, Tglu3C with $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 20 \text{ GeV}$, and Tglu3D with $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$, see their Figure 10.
NODE=S046GNO;LINKAGE=DF
- ¹⁴ SIRUNYAN 21M searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with two opposite-sign same-flavor leptons (electrons, muons) and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu4C, see their Figure 10, on the $\tilde{\chi}_2^0$ and $\tilde{\chi}_1^\pm$ mass in Tchi1n2Fa, see their Figure 11, on the $\tilde{\chi}_1^0$ mass in Tn1n1C and Tn1n1B for $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0}$, see their Figure 12. Limits are also set on the light squark mass for the simplified model Tsqk2A, on the sbottom mass in Tsb0t3, see their Figure 13, and on the slepton mass in direct electroweak pair production of mass-degenerate left- and right-handed sleptons (selectrons and smuons), see their Figure 14.
NODE=S046GNO;LINKAGE=CF
- ¹⁵ AAD 20AL searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with 8 or more jets and moderate missing transverse momentum. The selection makes requirements according to the number of b -tagged jets and the scalar sum of masses of large-radius jets. No significant excess above the Standard Model expectations is observed. Limits up to about 2 TeV are set on the gluino mass in Tglu1E simplified model. Limits up
NODE=S046GNO;LINKAGE=XE

- to about 1.8 TeV are set on the gluino mass in Tglu3A simplified model. See their Fig. 10(a).
- 16 AAD 20V searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with same-sign charged leptons (electrons or muons) and jets. No significant excess over the Standard Model expectation is observed. In the Tglu1E model, considering off-shell intermediate W and Z bosons in the decay chains, gluino masses are excluded at 95% C.L. up to 1600 GeV for neutralino masses of 100 GeV or above (up to 1000 GeV). See their Fig. 7(a). NODE=S046GNO;LINKAGE=VE
- 17 SIRUNYAN 20B searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one photon and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on chargino masses in a general gauge-mediated SUSY breaking (GGM) scenario Tchi1n12-GGM, see Figure 4. Limits are also set on the NLSP mass in the Tchi1chi1F and Tchi1chi1G simplified models, see their Figure 5. Finally, limits are set on the gluino mass in the Tglu4A simplified model, see Figure 6. NODE=S046GNO;LINKAGE=SE
- 18 SIRUNYAN 20BJ searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two hadronically decaying, highly energetic Z bosons and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1H simplified model, see their Figure 9. NODE=S046GNO;LINKAGE=WE
- 19 SIRUNYAN 20E searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a single electron or muon and multiple jets, including at least one identified as originating from a b -quark, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A simplified model, see their Fig. 10, and the Tglu3C simplified model, see their Fig. 11. NODE=S046GNO;LINKAGE=TE
- 20 SIRUNYAN 20T searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least two jets, and two isolated same-sign or three or more charged leptons (electrons or muons). No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figure 7, and in the Tglu1C and Tglu1B simplified models, see their Figures 8 and 9. Limits are also set on the sbottom mass in the Tsbott2 simplified model, see their Figure 10, and on the stop mass in the Tstop7 simplified model, see their Figure 11. Finally, limits are set on the gluino mass in RPV simplified models where the gluino decays either via $\tilde{g} \rightarrow qq\bar{q}\bar{q} + e/\mu/\tau$ or via $\tilde{g} \rightarrow tbs$, see Figure 12. NODE=S046GNO;LINKAGE=UE
- 21 AABOUD 19I searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with hadronic jets, 1 or two hadronically decaying τ and \cancel{E}_T . In Tglu1F, gluino masses are excluded at 95% C.L. up to 2000 GeV for neutralino masses of 100 GeV or below. Neutralino masses up to 1000 GeV are excluded for all gluino masses below 1400 GeV. See their Fig. 9. Limits are also presented in the context of Gauge-Mediated Symmetry Breaking models: in this case, values of Λ below 110 TeV are excluded at the 95% CL for all values of $\tan\beta$ in the range $2 < \tan\beta < 60$, see their Fig 10. NODE=S046GNO;LINKAGE=NE
- 22 SIRUNYAN 19AG searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4B simplified model and on the squark mass in the Tsqk4B simplified model, see their Figure 3. NODE=S046GNO;LINKAGE=JE
- 23 SIRUNYAN 19AU searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at last one photon, jets, some of which are identified as originating from b -quarks, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. In the framework of GMSB, limits are set on the gluino mass in the Tglu4C, Tglu4D and Tglu4E simplified models, and on the top squark mass in the Tstop13 simplified model, see their Figure 5. NODE=S046GNO;LINKAGE=LE
- 24 SIRUNYAN 19CE searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for new particles decaying to a photon and two gluons in events with at least three large-radius jets of which two have substructure and are composed of a photon and two gluons. No statistically significant excess is observed above the SM background expectation. Upper limits at 95% confidence level on the cross section for gluino pair production are set, using a simplified Tglu1A-like stealth SUSY model. Gluino masses up to 1500-1700 GeV are excluded, depending on the neutralino mass, with the highest exclusion set for $m_{\tilde{\chi}_1^0} = 200 \text{ GeV}$. See their Fig 4. NODE=S046GNO;LINKAGE=RE
- 25 SIRUNYAN 19CH searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A and Tglu3A simplified models, see their Figure 13. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tsbott1, Tstop1 simplified models, see their Figure 14. NODE=S046GNO;LINKAGE=OE
- 26 SIRUNYAN 19K searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a photon, an electron or muon, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. In the framework of GMSB, limits are set on the chargino and neutralino mass in the Tchi1n1A simplified model, see their Figure 6. Limits are also set on the gluino mass in the Tglu4A simplified model, and on the squark mass in the Tsqk4A simplified model, see their Figure 7. NODE=S046GNO;LINKAGE=KE
- 27 SIRUNYAN 19S searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with zero or one charged leptons, jets and \cancel{E}_T . The razor variables (M_R and R^2) are used to categorize the events. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu3C simplified models, see Figures 22 and 23, and on the stop mass in the Tstop1 simplified model, see their Figure 24. NODE=S046GNO;LINKAGE=ME
- 28 AABOUD 18AR searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for gluino pair production in events containing large missing transverse momentum and several energetic NODE=S046GNO;LINKAGE=UD

- jets, at least three of which must be identified as originating from b -quarks. No excess is found above the predicted background. In Tglu3A models, gluino masses of less than 1.97 TeV are excluded for $m_{\tilde{\chi}_1^0}$ below 300 GeV, see their Fig. 10(a). Interpretations are also provided for scenarios where Tglu3A modes mix with Tglu2A and Tglu3D, see their Fig 11.
- 29 AABOUD 18AR searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for gluino pair production in events containing large missing transverse momentum and several energetic jets, at least three of which must be identified as originating from b -quarks. No excess is found above the predicted background. In Tglu2A models, gluino masses of less than 1.92 TeV are excluded for $m_{\tilde{\chi}_1^0}$ below 600 GeV, see their Fig. 10(b). Interpretations are also provided for scenarios where Tglu2A modes mix with Tglu3A and Tglu3D, see their Fig 11. NODE=S046GNO;LINKAGE=VD
- 30 AABOUD 18AS searched for in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for gluino pair production in the context of AMSB or phenomenological MSSM scenarios with wino-like LSP and long-lived charginos. Events with a disappearing track due to a low-momentum pion accompanied by at least four jets are considered. No significant excess above the Standard Model expectations is observed. Exclusion limits are set at 95% confidence level on the mass of gluinos for different chargino lifetimes. Gluino masses up to 1.65 TeV are excluded assuming a chargino mass of 460 GeV and lifetime of 0.2 ns, corresponding to a mass-splitting between the charged and neutral wino of around 160 MeV. See their Fig. 9. NODE=S046GNO;LINKAGE=ND
- 31 AABOUD 18BJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with two opposite-sign charged leptons (electrons and muons), jets and missing transverse momentum, with various requirements to be sensitive to signals with different kinematic endpoint values in the dilepton invariant mass distribution. The data are found to be consistent with the SM expectation. Results are interpreted in the Tglu1G model: gluino masses below 1850 GeV are excluded for $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$, see their Fig. 12(a). NODE=S046GNO;LINKAGE=YD
- 32 AABOUD 18BJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with two opposite-sign charged leptons (electrons and muons), jets and missing transverse momentum, with various requirements to be sensitive to signals with different kinematic endpoint values in the dilepton invariant mass distribution. The data are found to be consistent with the SM expectation. Results are interpreted in the Tglu1H model: gluino masses below 1650 GeV are excluded for $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$, see their Fig. 13(a). NODE=S046GNO;LINKAGE=ZD
- 33 AABOUD 18U searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with at least one isolated photon, possibly jets and significant transverse momentum targeting generalised models of gauge-mediated SUSY breaking. No significant excess of events is observed above the SM prediction. Results for the di-photon channel are interpreted in terms of lower limits on the masses of gluinos in Tglu4B models, which reach as high as 2.3 TeV. Gluinos with masses below 2.15 TeV are excluded for any NLSP mass, see their Fig. 8. NODE=S046GNO;LINKAGE=WD
- 34 AABOUD 18U searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with at least one isolated photon, possibly jets and significant transverse momentum targeting generalised models of gauge-mediated SUSY breaking. No significant excess of events is observed above the SM prediction. Results of the $\gamma + \text{jets} + \cancel{E}_T$ channel are interpreted in terms of lower limits on the masses of gluinos in GGM higgsino-bino models (mix of Tglu4B and Tglu4C), which reach as high as 2050 GeV. Gluino masses below 1600 GeV are excluded for any NLSP mass provided that $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} > 50 \text{ GeV}$. See their Fig. 11. NODE=S046GNO;LINKAGE=XD
- 35 AABOUD 18V searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with no charged leptons, jets and missing transverse momentum. The data are found to be consistent with the SM expectation. Results are interpreted in the Tglu1A model: gluino masses below 2030 GeV are excluded for massless LSP, see their Fig. 13(b). NODE=S046GNO;LINKAGE=BE
- 36 AABOUD 18V searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with no charged leptons, jets and missing transverse momentum. The data are found to be consistent with the SM expectation. Results are interpreted in the Tglu1B model. Assuming that $m_{\tilde{\chi}_1^\pm} = 0.5 (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})$, gluino masses below 1980 GeV are excluded for massless LSP, see their Fig. 14(c). Exclusions are also shown assuming $m_{\tilde{\chi}_1^0} = 60 \text{ GeV}$, see their Fig. 14(d). NODE=S046GNO;LINKAGE=CE
- 37 AABOUD 18V searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with no charged leptons, jets and missing transverse momentum. The data are found to be consistent with the SM expectation. Results are interpreted in the Tglu1E model: gluino masses below 1750 GeV are excluded for $m_{\tilde{\chi}_1^0} = 1 \text{ GeV}$ and any $m_{\tilde{\chi}_2^0}$ above 100 GeV, see their Fig. 15. Gluino mass exclusion up to 2 TeV is found for $m_{\tilde{\chi}_2^0} = 1 \text{ TeV}$. NODE=S046GNO;LINKAGE=EE
- 38 SIRUNYAN 18AA searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one photon and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on wino masses in a general gauge-mediated SUSY breaking (GGM) scenario with bino-like $\tilde{\chi}_1^0$ and wino-like $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_2^0$, see Figure 7. Limits are also set on the NLSP mass in the Tchi1n1A and Tchi1chi1A simplified models, see their Figure 8. Finally, limits are set on the gluino mass in the Tglu4A and Tglu4B simplified models, see their Figure 9, and on the squark mass in the Tskq4A and Tskq4B simplified models, see their Figure 10. NODE=S046GNO;LINKAGE=LD

- 39 SIRUNYAN 18AC searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a single electron or muon and multiple jets. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu1B simplified models, see their Figure 5. NODE=S046GNO;LINKAGE=MD
- 40 SIRUNYAN 18AL searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least three charged leptons, in any combination of electrons and muons, jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu1C simplified models, see their Figure 5. Limits are also set on the sbottom mass in the Tsb02 simplified model, see their Figure 6, and on the stop mass in the Tstop7 simplified model, see their Figure 7. NODE=S046GNO;LINKAGE=QD
- 41 SIRUNYAN 18AR searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing two opposite-charge, same-flavour leptons (electrons or muons), jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4C simplified model, see their Figure 7. Limits are also set on the chargino/neutralino mass in the Tchi1n2F simplified models, see their Figure 8, and on the higgsino mass in the Tn1n1B and Tn1n1C simplified models, see their Figure 9. Finally, limits are set on the sbottom mass in the Tsb03 simplified model, see their Figure 10. NODE=S046GNO;LINKAGE=RD
- 42 SIRUNYAN 18AY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing one or more jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A and Tglu3A simplified models, see their Figure 3. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tsb01, Tstop1 and Tstop4 simplified models, see their Figure 3. Finally, limits are set on long-lived gluino masses in a Tglu1A simplified model where the gluino is metastable or long-lived with proper decay lengths in the range $10^{-3} \text{ mm} < c\tau < 10^5 \text{ mm}$, see their Figure 4. NODE=S046GNO;LINKAGE=SD
- 43 SIRUNYAN 18D searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing identified hadronically decaying top quarks, no leptons, and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1 simplified model, see their Figure 8, and on the gluino mass in the Tglu3A, Tglu3B, Tglu3C and Tglu3E simplified models, see their Figure 9. NODE=S046GNO;LINKAGE=KC
- 44 SIRUNYAN 18M searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more high-momentum Higgs bosons, decaying to pairs of b -quarks, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1I and Tglu1J simplified models, see their Figure 3. NODE=S046GNO;LINKAGE=PD
- 45 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.75 TeV are set on the gluino mass in Tglu3A simplified models in case of off-shell top squarks and for $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$. See their Figure 4(a). NODE=S046GNO;LINKAGE=ZC
- 46 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.57 TeV are set on the gluino mass in Tglu1E simplified models (2-step models) for $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$. See their Figure 4(b). NODE=S046GNO;LINKAGE=BD
- 47 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.86 TeV are set on the gluino mass in Tglu1G simplified models for $m_{\tilde{\chi}_1^0} = 200 \text{ GeV}$. See their Figure 4(c). NODE=S046GNO;LINKAGE=CD
- 48 AABOUD 17AR searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one isolated lepton, at least two jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 2.1 TeV are set on the gluino mass in Tglu1B simplified models, with $x = (m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0}) / (m_{\tilde{g}} - m_{\tilde{\chi}_1^0}) = 1/2$. Similar limits are obtained for variable x and fixed neutralino mass, $m_{\tilde{\chi}_1^0} = 60 \text{ GeV}$. See their Figure 13. NODE=S046GNO;LINKAGE=QC
- 49 AABOUD 17AR searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one isolated lepton, at least two jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.74 TeV are set on the gluino mass in Tglu1E simplified model. Limits up to 1.7 TeV are also set on pMSSM models leading to similar signal event topologies. See their Figure 13. NODE=S046GNO;LINKAGE=TC
- 50 AABOUD 17AY searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least four jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.8 TeV are set on the gluino mass in Tglu3A simplified models assuming $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 5 \text{ GeV}$. See their Figure 13. NODE=S046GNO;LINKAGE=YC
- 51 AABOUD 17AZ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least seven jets and large missing transverse momentum. Selected events are further classified based on the presence of large R-jets or b -jets and no leptons. No significant excess above the Standard Model expectations is observed. Limits up to 1.8 TeV are set on the gluino mass in Tglu1E simplified models. See their Figure 6b. NODE=S046GNO;LINKAGE=UC
- 52 AABOUD 17AZ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least seven jets and large missing transverse momentum. Selected events are further NODE=S046GNO;LINKAGE=VC

classified based on the presence of large R-jets or b -jets and no leptons. No significant excess above the Standard Model expectations is observed. Limits up to 1.54 TeV are set on the gluino mass in Tglu3A simplified models. See their Figure 7a.

- 53 AABOUD 17N searched in 14.7 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with 2 same-flavor, opposite-sign leptons (electrons or muons), jets and large missing transverse momentum. In Tglu1J models, gluino masses are excluded at 95% C.L. up to 1300 GeV for $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$ and $m_{\tilde{\chi}_2^0} = 1100 \text{ GeV}$. See their Fig. 12 for exclusion limits as a function of $m_{\tilde{\chi}_2^0}$. Limits are also presented assuming $m_{\tilde{\chi}_2^0} = m_{\tilde{\chi}_1^0} + 100 \text{ GeV}$, see their Fig. 13. NODE=S046GNO;LINKAGE=WB
- 54 AABOUD 17N searched in 14.7 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with 2 same-flavor, opposite-sign leptons (electrons or muons), jets and large missing transverse momentum. In Tglu1H models, gluino masses are excluded at 95% C.L. up to 1310 GeV for $m_{\tilde{\chi}_1^0} < 400 \text{ GeV}$ and assuming $m_{\tilde{\chi}_2^0} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$. See their Fig. 15. NODE=S046GNO;LINKAGE=XB
- 55 AABOUD 17N searched in 14.7 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with 2 same-flavor, opposite-sign leptons (electrons or muons), jets and large missing transverse momentum. In Tglu1G models, gluino masses are excluded at 95% C.L. up to 1700 GeV for small $m_{\tilde{\chi}_1^0}$. The results probe kinematic endpoints as small as $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0} = (m_{\tilde{g}} - m_{\tilde{\chi}_1^0})/2 = 50 \text{ GeV}$. See their Fig. 14. NODE=S046GNO;LINKAGE=YB
- 56 KHACHATRYAN 17 searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more jets, no more than one lepton, and missing transverse momentum, using the razor variables (M_R and R^2) to discriminate between signal and background processes. No evidence for an excess over the expected background is observed. Limits are derived on the gluino mass in the Tglu1A, Tglu2A and Tglu3A simplified models, see Figs. 16 and 17. Also, assuming gluinos decay only via three-body processes involving third-generation quarks plus a neutralino/chargino, and assuming $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_1^0} + 5 \text{ GeV}$, a branching ratio-independent limit on the gluino mass is given, see Fig. 16. NODE=S046GNO;LINKAGE=PB
- 57 KHACHATRYAN 17AD searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing at least four jets (including b -jets), missing transverse momentum and tagged top quarks. No evidence for an excess over the expected background is observed. Gluino masses up to 1550 GeV and neutralino masses up to 900 GeV are excluded at 95% C.L. See Fig. 13. NODE=S046GNO;LINKAGE=AC
- 58 KHACHATRYAN 17AD searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing at least four jets (including b -jets), missing transverse momentum and tagged top quarks. No evidence for an excess over the expected background is observed. Gluino masses up to 1450 GeV and neutralino masses up to 820 GeV are excluded at 95% C.L. See Fig. 13. NODE=S046GNO;LINKAGE=BC
- 59 KHACHATRYAN 17AS searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a single electron or muon and multiple jets. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu1B simplified models, see their Fig. 7. NODE=S046GNO;LINKAGE=OB
- 60 KHACHATRYAN 17AW searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least three charged leptons, in any combination of electrons and muons, and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu1C simplified models, and on the sbottom mass in the Tsb0t2 simplified model, see their Figure 4. NODE=S046GNO;LINKAGE=IC
- 61 KHACHATRYAN 17P searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figures 7 and 8. Limits are also set on the squark mass in the Tsqk1 simplified model, see their Fig. 7, and on the sbottom mass in the Tsb0t1 simplified model, see Fig. 8. Finally, limits are set on the stop mass in the Tstop1, Tstop3, Tstop4, Tstop6 and Tstop7 simplified models, see Fig. 8. NODE=S046GNO;LINKAGE=VB
- 62 KHACHATRYAN 17V searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino and squark mass in the context of general gauge mediation models Tglu4B and Tsqk4, see their Fig. 4. NODE=S046GNO;LINKAGE=PA
- 63 SIRUNYAN 17AF searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with a single lepton (electron or muon), jets, including at least one jet originating from a b -quark, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A and Tglu3B simplified models, see their Figure 2. NODE=S046GNO;LINKAGE=EC
- 64 SIRUNYAN 17AY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one photon, jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4A and Tglu4B simplified models, and on the squark mass in the Tsqk4A and Tsqk4B simplified models, see their Figure 6. NODE=S046GNO;LINKAGE=FC
- 65 SIRUNYAN 17AZ searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A, Tglu3A simplified models, see their Figures 6. Limits are also set on the squark mass in the Tsqk1 simplified model (for single light squark and for 8 degenerate light squarks), on the sbottom mass in the Tsb0t1 simplified model and on the stop mass in the Tstop1 NODE=S046GNO;LINKAGE=DC

- simplified model, see their Fig. 7. Finally, limits are set on the stop mass in the Tstop2, Tstop4 and Tstop8 simplified models, see Fig. 8.
- 66 SIRUNYAN 17P searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with multiple jets and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A, Tglu3A and Tglu3D simplified models, see their Fig. 12. Limits are also set on the squark mass in the Tsqk1 simplified model, on the stop mass in the Tstop1 simplified model, and on the sbottom mass in the Tsb01 simplified model, see Fig. 13. NODE=S046GNO;LINKAGE=UB
- 67 SIRUNYAN 17S searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two isolated same-sign leptons, jets, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the mass of the gluino mass in the Tglu3A, Tglu3B, Tglu3C, Tglu3D and Tglu1B simplified models, see their Figures 5 and 6, and on the sbottom mass in the Tsb02 simplified model, see their Figure 6. NODE=S046GNO;LINKAGE=GC
- 68 AABOUD 16AC searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with hadronic jets, 1 or two hadronically decaying τ and \cancel{E}_T . In Tglu1F, gluino masses are excluded at 95% C.L. up to 1570 GeV for neutralino masses of 100 GeV or below. Neutralino masses up to 700 GeV are excluded for all gluino masses between 800 GeV and 1500 GeV, while the strongest neutralino-mass exclusion of 750 GeV is achieved for gluino masses around 1400 GeV. See their Fig. 8. Limits are also presented in the context of Gauge-Mediated Symmetry Breaking models: in this case, values of Λ below 92 TeV are excluded at the 95% CL, corresponding to gluino masses below 2000 GeV. See their Fig. 9. NODE=S046GNO;LINKAGE=SB
- 69 AABOUD 16J searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with one isolated electron or muon, hadronic jets, and \cancel{E}_T . Gluino-mediated pair production of stops with a nearly mass-degenerate stop and neutralino are targeted and gluino masses are excluded at 95% C.L. up to 1460 GeV. A 100% of stops decaying via charm + neutralino is assumed. The results are also valid in case of 4-body decays $\tilde{t}_1 \rightarrow f f' b \tilde{\chi}_1^0$. See their Fig. 8. NODE=S046GNO;LINKAGE=CB
- 70 AABOUD 16M searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two photons, hadronic jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Exclusion limits at 95% C.L. are set on gluino masses in the general gauge-mediated SUSY breaking model (GGM), for bino-like NLSP. See their Fig. 3. NODE=S046GNO;LINKAGE=DB
- 71 AABOUD 16N searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing hadronic jets, large \cancel{E}_T , and no electrons or muons. No significant excess above the Standard Model expectations is observed. Gluino masses below 1510 GeV are excluded at the 95% C.L. in a simplified model with only gluinos and the lightest neutralino. See their Fig. 7b. NODE=S046GNO;LINKAGE=EB
- 72 AABOUD 16N searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing hadronic jets, large \cancel{E}_T , and no electrons or muons. No significant excess above the Standard Model expectations is observed. Gluino masses below 1500 GeV are excluded at the 95% C.L. in a simplified model with gluinos decaying via an intermediate $\tilde{\chi}_1^\pm$ to two quarks, a W boson and a $\tilde{\chi}_1^0$, for $m_{\tilde{\chi}_1^0} = 200 \text{ GeV}$. See their Fig 8. NODE=S046GNO;LINKAGE=FB
- 73 AAD 16AD searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing several energetic jets, of which at least three must be identified as b -jets, large \cancel{E}_T and no electrons or muons. No significant excess above the Standard Model expectations is observed. For $\tilde{\chi}_1^0$ below 800 GeV, gluino masses below 1780 GeV are excluded at 95% C.L. for gluinos decaying via bottom squarks. See their Fig. 7a. NODE=S046GNO;LINKAGE=GB
- 74 AAD 16AD searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing several energetic jets, of which at least three must be identified as b -jets, large \cancel{E}_T and one electron or muon. Large-radius jets with a high mass are also used to identify highly boosted top quarks. No significant excess above the Standard Model expectations is observed. For $\tilde{\chi}_1^0$ below 700 GeV, gluino masses below 1760 GeV are excluded at 95% C.L. for gluinos decaying via top squarks. See their Fig. 7b. NODE=S046GNO;LINKAGE=HB
- 75 AAD 16BB searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with exactly two same-sign leptons or at least three leptons, multiple hadronic jets, b -jets, and \cancel{E}_T . No significant excess over the Standard Model expectation is found. Exclusion limits at 95% C.L. are set on the gluino mass in various simplified models (Tglu1D, Tglu1E, Tglu3A). See their Figs. 4.a, 4.b, and 4.d. NODE=S046GNO;LINKAGE=IB
- 76 AAD 16BG searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in final states with one isolated electron or muon, hadronic jets, and \cancel{E}_T . The data agree with the SM background expectation in the six signal selections defined in the search, and the largest deviation is a 2.1 standard deviation excess. Gluinos are excluded at 95% C.L. up to 1600 GeV assuming they decay via the lightest chargino to the lightest neutralino as in the model Tglu1B for $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$, assuming $m_{\tilde{\chi}_1^\pm} = (m_{\tilde{g}} + m_{\tilde{\chi}_1^0})/2$. See their Fig. 6. NODE=S046GNO;LINKAGE=TB
- 77 AAD 16V searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with \cancel{E}_T various hadronic jet multiplicities from ≥ 7 to ≥ 10 and with various b -jet multiplicity requirements. No significant excess over the Standard Model expectation is found. Exclusion limits at 95% C.L. are set on the gluino mass in one simplified model (Tglu1E) and a pMSSM-inspired model. See their Fig. 5. NODE=S046GNO;LINKAGE=JB
- 78 KHACHATRYAN 16AM searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with highly boosted W -bosons and b -jets, using the razor variables (M_R and R^2) to discriminate between signal and background processes. No significant excess above the NODE=S046GNO;LINKAGE=KB

- Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3C and Tglu3B simplified models, see Fig. 12.
- 79 KHACHATRYAN 16BJ searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the following simplified models: Tglu3A and Tglu3D, see Fig. 4, Tglu3B and Tglu3C, see Fig. 5, and Tglu1B, see Fig. 7. NODE=S046GNO;LINKAGE=NB
- 80 KHACHATRYAN 16BS searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least one energetic jet, no isolated leptons, and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A and Tglu3A simplified models, see Fig. 10 and Table 3. NODE=S046GNO;LINKAGE=MA
- 81 KHACHATRYAN 16BY searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two opposite-sign, same-flavour leptons, jets, and missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu4C simplified model, see Fig. 4, and on sbottom masses in the Tsb0t3 simplified model, see Fig. 5. NODE=S046GNO;LINKAGE=MB
- 82 KHACHATRYAN 16V searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least four energetic jets and significant \cancel{E}_T , no identified isolated electron or muon or charged track. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu1C, Tglu2A, and Tglu3A simplified models, see Fig. 8. NODE=S046GNO;LINKAGE=UA
- 83 AAD 15BG searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with jets, missing E_T , and two opposite-sign same flavor isolated leptons featuring either a kinematic edge, or a peak at the Z-boson mass, in the invariant mass spectrum. No evidence for a statistically significant excess over the expected SM backgrounds are observed and 95% C.L. exclusion limits are derived in a GGM simplified model of gluino pair production where the gluino decays into quarks, a Z-boson, and a massless gravitino LSP, see Fig. 12. Also, limits are set in simplified models with slepton/sneutrino intermediate states, see Fig. 13. NODE=S046GNO;LINKAGE=LA
- 84 AAD 15BV summarized and extended ATLAS searches for gluinos and first- and second-generation squarks in final states containing jets and missing transverse momentum, with or without leptons or b -jets in the $\sqrt{s} = 8 \text{ TeV}$ data set collected in 2012. The paper reports the results of new interpretations and statistical combinations of previously published analyses, as well as new analyses. Exclusion limits at 95% C.L. are set on the gluino mass in several R-parity conserving models, leading to a generalized constraint on gluino masses exceeding 1150 GeV for lightest supersymmetric particle masses below 100 GeV. See their Figs. 10, 19, 20, 21, 23, 25, 26, 29-37. NODE=S046GNO;LINKAGE=OA
- 85 AAD 15BX interpreted the results of a wide range of ATLAS direct searches for supersymmetry, during the first run of the LHC using the $\sqrt{s} = 7 \text{ TeV}$ and $\sqrt{s} = 8 \text{ TeV}$ data set collected in 2012, within the wider framework of the phenomenological MSSM (pMSSM). The integrated luminosity was up to 20.3 fb^{-1} . From an initial random sampling of 500 million pMSSM points, generated from the 19-parameter pMSSM, a total of 310,327 model points with $\tilde{\chi}_1^0$ LSP were selected each of which satisfies constraints from previous collider searches, precision measurements, cold dark matter energy density measurements and direct dark matter searches. The impact of the ATLAS Run 1 searches on this space was presented, considering the fraction of model points surviving, after projection into two-dimensional spaces of sparticle masses. Good complementarity is observed between different ATLAS analyses, with almost all showing regions of unique sensitivity. ATLAS searches have good sensitivity at LSP mass below 800 GeV. NODE=S046GNO;LINKAGE=QA
- 86 AAD 15CA searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with one or more photons, hadronic jets or b -jets and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on gluino masses in the general gauge-mediated SUSY breaking model (GGM), for bino-like or higgsino-bino admixtures NLSP, see Fig. 8, 10, 11. NODE=S046GNO;LINKAGE=VA
- 87 KHACHATRYAN 15AF searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the transverse mass variable M_{T2} to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in simplified models where the decay $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 13(a), or where the decay $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 13(b), or where the decay $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 13(c). See also Table 5. Exclusions in the CMSSM, assuming $\tan\beta = 30$, $A_0 = -2 \max(m_0, m_{1/2})$ and $\mu > 0$, are also presented, see Fig. 15. NODE=S046GNO;LINKAGE=EA
- 88 KHACHATRYAN 15I searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events in which b -jets and four W -bosons are produced. Five individual search channels are combined (fully hadronic, single lepton, same-sign dilepton, opposite-sign dilepton, multi-lepton). No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in a simplified model where the decay $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 5. Also a simplified model with gluinos decaying into on-shell top squarks is considered, see Fig. 6. NODE=S046GNO;LINKAGE=JA
- 89 KHACHATRYAN 15X searched in 19.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least two energetic jets, at least one of which is required to originate from a b quark, and significant \cancel{E}_T , using the razor variables (M_R) and R^2) to discriminate NODE=S046GNO;LINKAGE=KA

- between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in simplified models where the decay $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$ and the decay $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ take place with branching ratios varying between 0, 50 and 100%, see Figs. 13 and 14.
- 90 AAD 14AE searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for strongly produced supersymmetric particles in events containing jets and large missing transverse momentum, and no electrons or muons. No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing gluinos and squarks, see Figures 5, 6 and 7. Limits are also derived in the mSUGRA/CMSSM with parameters $\tan\beta = 30$, $A_0 = -2 m_0$ and $\mu > 0$, see their Fig. 8. NODE=S046GNO;LINKAGE=U
- 91 AAD 14AG searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing one hadronically decaying τ -lepton, zero or one additional light leptons (electrons or muons), jets and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set in several SUSY scenarios. For an interpretation in the minimal GMSB model, see their Fig. 8. For an interpretation in the mSUGRA/CMSSM with parameters $\tan\beta = 30$, $A_0 = -2 m_0$ and $\mu > 0$, see their Fig. 9. For an interpretation in the framework of natural Gauge Mediation, see Fig. 10. For an interpretation in the bRPV scenario, see their Fig. 11. NODE=S046GNO;LINKAGE=V
- 92 AAD 14X searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least four leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in a general gauge-mediation model (GGM) where the decay $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$, with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \tilde{G}$, takes place with a branching ratio of 100%, for two choices of $\tan\beta = 1.5$ and 30, see Fig. 11. Also some constraints on the higgsino mass parameter μ are discussed. NODE=S046GNO;LINKAGE=I
- 93 CHATRCHYAN 14AH searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the razor variables (M_R and R^2) to discriminate between signal and background processes. No significant excess above the Standard Model expectations is observed. Limits are set on sbottom masses in simplified models where the decay $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 28. Exclusions in the CMSSM, assuming $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, are also presented, see Fig. 26. NODE=S046GNO;LINKAGE=W
- 94 CHATRCHYAN 14AH searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the razor variables (M_R and R^2) to discriminate between signal and background processes. A second analysis requires at least one of the jets to be originating from a b -quark. No significant excess above the Standard Model expectations is observed. Limits are set on sbottom masses in simplified models where the decay $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Figs. 28 and 29. Exclusions in the CMSSM, assuming $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, are also presented, see Fig. 26. NODE=S046GNO;LINKAGE=X
- 95 CHATRCHYAN 14AH searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with at least two energetic jets and significant \cancel{E}_T , using the razor variables (M_R and R^2) to discriminate between signal and background processes. A second analysis requires at least one of the jets to be originating from a b -quark. No significant excess above the Standard Model expectations is observed. Limits are set on sbottom masses in simplified models where the decay $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Figs. 28 and 29. Exclusions in the CMSSM, assuming $\tan\beta = 10$, $A_0 = 0$ and $\mu > 0$, are also presented, see Fig. 26. NODE=S046GNO;LINKAGE=Y
- 96 CHATRCHYAN 14I searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing multijets and large \cancel{E}_T . No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing gluinos that decay via $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ with a 100% branching ratio, see Fig. 7b, or via $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ with a 100% branching ratio, see Fig. 7c, or via $\tilde{g} \rightarrow q\bar{q}W/Z\tilde{\chi}_1^0$, see Fig. 7d. NODE=S046GNO;LINKAGE=T
- 97 CHATRCHYAN 14N searched in 19.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing a single isolated electron or muon and multiple jets, at least two of which are identified as originating from a b -quark. No significant excesses over the expected SM backgrounds are observed. The results are interpreted in three simplified models of gluino pair production with subsequent decay into virtual or on-shell top squarks, where each of the top squarks decays in turn into a top quark and a $\tilde{\chi}_1^0$, see Fig. 4. The models differ in which masses are allowed to vary. NODE=S046GNO;LINKAGE=A
- 98 CHATRCHYAN 14R searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least three leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in a slepton co-NLSP simplified model (GMSB) where the decay $\tilde{g} \rightarrow q\ell^\pm \ell^\mp \tilde{G}$ takes place with a branching ratio of 100%, see Fig. 8. NODE=S046GNO;LINKAGE=B
- 99 CHATRCHYAN 14R searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least three leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in a simplified model where the decay $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ takes place with a branching ratio of 100%, see Fig. 11. NODE=S046GNO;LINKAGE=C
- 100 AABOUD 18BJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ in events with two opposite-sign charged leptons (electrons and muons), jets and missing transverse momentum, with various requirements to be sensitive to signals with different kinematic NODE=S046GNO;LINKAGE=AE

- endpoint values in the dilepton invariant mass distribution. The data are found to be consistent with the SM expectation. Results are interpreted in the Tglu1H model in case of $m_{\tilde{\chi}_1^0} = 1$ GeV: for any $m_{\tilde{\chi}_2^0}$, gluino masses below 1500 GeV are excluded, see their Fig. 14(a).
- 101 AABOUD 18V searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV in events with no charged leptons, jets and missing transverse momentum. The data are found to be consistent with the SM expectation. Results are interpreted in a Tglu1C-like model, assuming 50% BR for each gluino decay mode. Gluino masses below 1770 GeV are excluded for any $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$ and $m_{\tilde{\chi}_1^0} = 60$ GeV, see their Fig. 16(b). NODE=S046GNO;LINKAGE=FE
- 102 AABOUD 17AZ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with at least seven jets and large missing transverse momentum. Selected events are further classified based on the presence of large R-jets or b -jets and no leptons. No significant excess above the Standard Model expectations is observed. Limits are set for pMSSM models with $M_1 = 60$ GeV, $\tan(\beta) = 10$, $\mu < 0$ varying the soft-breaking parameters M_3 and μ . Gluino masses up to 1600 GeV are excluded for $m_{\tilde{\chi}_1^\pm} = 200$ GeV. See their Figure 6a and text for details on the model. NODE=S046GNO;LINKAGE=XC
- 103 KHACHATRYAN 16AY searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with one isolated high transverse momentum lepton (e or μ), hadronic jets of which at least one is identified as coming from a b -quark, and large \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A simplified model, see Fig. 10, and in the Tglu3B model, see Fig. 11. NODE=S046GNO;LINKAGE=FA
- 104 KHACHATRYAN 16BT performed a global Bayesian analysis of a wide range of CMS results obtained with data samples corresponding to 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV and in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV. The set of searches considered, both individually and in combination, includes those with all-hadronic final states, same-sign and opposite-sign dileptons, and multi-lepton final states. An interpretation was given in a scan of the 19-parameter pMSSM. No scan points with a gluino mass less than 500 GeV survived and 98% of models with a squark mass less than 300 GeV were excluded. NODE=S046GNO;LINKAGE=XA
- 105 AAD 15AB searched for the decay of neutral, weakly interacting, long-lived particles in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV. Signal events require at least two reconstructed vertices possibly originating from long-lived particles decaying to jets in the inner tracking detector and muon spectrometer. No significant excess of events over the expected background was found. Results were interpreted in Stealth SUSY benchmark models where a pair of gluinos decay to long-lived singlinos, \tilde{S} , which in turn each decay to a low-mass gravitino and a pair of jets. The 95% confidence-level limits are set on the cross section \times branching ratio for the decay $\tilde{g} \rightarrow \tilde{S}g$, as a function of the singlino proper lifetime ($c\tau$). See their Fig. 10(f). NODE=S046GNO;LINKAGE=ZA
- 106 AAD 15AI searched in 20 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events containing at least one isolated lepton (electron or muon), jets, and large missing transverse momentum. No excess of events above the expected level of Standard Model background was found. Exclusion limits at 95% C.L. are set on the gluino mass in the CMSSM/mSUGRA, see Fig. 15, in the NUHMG, see Fig. 16, and in various simplified models, see Figs. 18–22. NODE=S046GNO;LINKAGE=NA
- 107 AAD 15CB searched for events containing at least one long-lived particle that decays at a significant distance from its production point (displaced vertex, DV) into two leptons or into five or more charged particles in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV. The dilepton signature is characterised by DV formed from at least two lepton candidates. Four different final states were considered for the multitrak signature, in which the DV must be accompanied by a high-transverse momentum muon or electron candidate that originates from the DV, jets or missing transverse momentum. No events were observed in any of the signal regions. Results were interpreted in SUSY scenarios involving R -parity violation, split supersymmetry, and gauge mediation. See their Fig. 12–20. NODE=S046GNO;LINKAGE=TA
- 108 KHACHATRYAN 15AD searched in 19.4 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with two opposite-sign same flavor isolated leptons featuring either a kinematic edge, or a peak at the Z -boson mass, in the invariant mass spectrum. No evidence for a statistically significant excess over the expected SM backgrounds is observed and 95% C.L. exclusion limits are derived in a simplified model of gluino pair production where the gluino decays into quarks, a Z -boson, and a massless gravitino LSP, see Fig. 9. NODE=S046GNO;LINKAGE=Z
- 109 KHACHATRYAN 15AZ searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with either at least one photon, hadronic jets and \cancel{E}_T (single photon channel) or with at least two photons and at least one jet and using the razor variables. No significant excess above the Standard Model expectations is observed. Limits are set on gluino masses in the general gauge-mediated SUSY breaking model (GGM), for both a bino-like and wino-like neutralino NLSP scenario, see Fig. 8 and 9. NODE=S046GNO;LINKAGE=BB
- 110 AAD 14AX searched in 20.1 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for the strong production of supersymmetric particles in events containing either zero or at last one high high- p_T lepton, large missing transverse momentum, high jet multiplicity and at least three jets identified as originating from b -quarks. No excess over the expected SM background is observed. Limits are derived in mSUGRA/CMSSM models with $\tan\beta = 30$, $A_0 = -2m_0$ and $\mu > 0$, see their Fig. 14. Also, exclusion limits in simplified models containing gluinos and scalar top and bottom quarks are set, see their Figures 12, 13. NODE=S046GNO;LINKAGE=BA
- 111 AAD 14E searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for strongly produced supersymmetric particles in events containing jets and two same-sign leptons or three NODE=S046GNO;LINKAGE=B2

leptons. The search also utilises jets originating from b -quarks, missing transverse momentum and other variables. No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing gluinos and squarks, see Figures 5 and 6. In the $\tilde{g} \rightarrow qq' \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^{(*)\pm} \tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow Z^{(*)} \tilde{\chi}_1^0$ simplified model, the following assumptions have been made: $m_{\tilde{\chi}_1^\pm} = 0.5 m_{\tilde{\chi}_1^0} + m_{\tilde{g}}, m_{\tilde{\chi}_2^0} = 0.5 (m_{\tilde{\chi}_1^0} + m_{\tilde{\chi}_1^\pm}), m_{\tilde{\chi}_1^0} < 520$ GeV. In the $\tilde{g} \rightarrow qq' \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \ell^\pm \nu \tilde{\chi}_1^0$ or $\tilde{g} \rightarrow qq' \tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow \ell^\pm \ell^\mp (\nu\nu) \tilde{\chi}_1^0$ simplified model, the following assumptions have been made: $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0} = 0.5 (m_{\tilde{\chi}_1^0} + m_{\tilde{g}}), m_{\tilde{\chi}_1^0} < 660$ GeV. Limits are also derived in the mSUGRA/CMSSM, bRPV and GMSB models, see their Fig. 8.

- 112 CHATRCHYAN 14H searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in simplified models where the decay $\tilde{g} \rightarrow t \bar{t} \tilde{\chi}_1^0$ takes place with a branching ratio of 100%, or where the decay $\tilde{g} \rightarrow \tilde{t} t, \tilde{t} \rightarrow t \tilde{\chi}_1^0$ takes place with a branching ratio of 100%, with varying mass of the $\tilde{\chi}_1^0$, or where the decay $\tilde{g} \rightarrow \tilde{b} b, \tilde{b} \rightarrow t \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ takes place with a branching ratio of 100%, with varying mass of the $\tilde{\chi}_1^\pm$, see Fig. 5.
- 113 CHATRCHYAN 14H searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in simplified models where the decay $\tilde{g} \rightarrow qq' \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ takes place with a branching ratio of 100%, with varying mass of the $\tilde{\chi}_1^\pm$ and $\tilde{\chi}_1^0$, see Fig. 7.
- 114 CHATRCHYAN 14H searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in simplified models where the decay $\tilde{g} \rightarrow b \bar{t} \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$ takes place with a branching ratio of 100%, for two choices of $m_{\tilde{\chi}_1^\pm}$ and fixed $m_{\tilde{\chi}_1^0}$, see Fig. 6.

NODE=S046GNO;LINKAGE=J

NODE=S046GNO;LINKAGE=M

NODE=S046GNO;LINKAGE=N

R-parity violating heavy \tilde{g} (Gluino) mass limit

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>2200	95	1 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tglu3F, λ_{323}'' electroweakino decay, $500 \text{ GeV} < m_{\tilde{\chi}_1^0} < 1600$ GeV
>2250	95	1 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tglu3G, λ_{323}'' electroweakino decay, $600 \text{ GeV} < m_{\tilde{\chi}_1^0} < 1600$ GeV
>2200	95	1 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tglu3B, λ_{323}'' electroweakino decay, $600 \text{ GeV} < m_{\tilde{\chi}_1^0} < 1600$ GeV
>1800	95	1 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tglu3B, λ_{323}'' , \tilde{t} decay, $m_{\tilde{t}} < 1200$ GeV
>2200	95	1 AAD	21BF ATLS	$\ell^\pm + b$ -jets + many jets, Tglu1A, $\lambda_{11}^I, \tilde{\chi}_1^0$ decay with equal probability into e, μ, ν_e, ν_μ , $400 \text{ GeV} < m_{\tilde{\chi}_1^0} < 1700$ GeV
>2500	95	2 AAD	21Y ATLS	$\geq 4\ell$, Tglu1A with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu, \lambda_{12k} \neq 0, m_{\tilde{\chi}_1^0} = 2200$ GeV
>1900	95	2 AAD	21Y ATLS	$\geq 4\ell$, Tglu1A with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu, \lambda_{133} \neq 0, m_{\tilde{\chi}_1^0} = 1550$ GeV
>1600	95	3 AAD	20AL ATLS	8 or more jets+ \cancel{E}_T , Tglu2RPV
>1600	95	4 AAD	20V ATLS	same-sign $\ell^\pm \ell^\pm +$ jets, $\tilde{g} \rightarrow t b d$ simplified model
>2150	95	5 SIRUNYAN	20T CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm +$ jets, $\tilde{g} \rightarrow qq\bar{q}\bar{q} + e/\mu/\tau$ simplified model

NODE=S046GNV

NODE=S046GNV;CHECK LIMITS

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=2

>1725	95	5	SIRUNYAN	20T	CMS	same-sign $\ell^\pm \ell^\pm$ or $\geq 3\ell^\pm + \text{jets}$, $\tilde{g} \rightarrow tbs$ simplified model	OCCUR=2
>1500	95	6	SIRUNYAN	19F	CMS	$\tilde{g} \rightarrow jjj$	
>2260	95	7	AABOUD	18Z	ATLS	$\geq 4\ell, \lambda_{12k} \neq 0, m_{\tilde{\chi}_1^0} > 1000$ GeV	
>1650	95	7	AABOUD	18Z	ATLS	$\geq 4\ell, \lambda_{i33} \neq 0, m_{\tilde{\chi}_1^0} > 500$ GeV	OCCUR=2
>1610	95	8	SIRUNYAN	18AK	CMS	$\tilde{g} \rightarrow tbs, \lambda''_{332}$ coupling	
>1690	95	9	SIRUNYAN	18D	CMS	top quark (hadronically decay- ing) + jets + \cancel{E}_T , Tglu3C, $m_{\tilde{t}_1} - m_{\tilde{\chi}_1^0} = 20$ GeV, $m_{\tilde{\chi}_1^0} =$ 0 GeV	OCCUR=3
none 100–1410	95	10	SIRUNYAN	18EA	CMS	2 large jets with four-parton sub- structure, $\tilde{g} \rightarrow 5q$	
>2100	95	11	AABOUD	17AI	ATLS	$\geq 1\ell + \geq 8$ jets, Tglu3A and $\tilde{\chi}_1^0 \rightarrow uds, \lambda''_{112}$ coupling, $m_{\tilde{\chi}_1^0} = 1000$ GeV	
>1650	95	12	AABOUD	17AI	ATLS	$\geq 1\ell + \geq 8$ jets, $\tilde{g} \rightarrow t\tilde{t}, \tilde{t} \rightarrow$ bs, λ''_{323} coupling, $m_{\tilde{t}} = 1000$ GeV	OCCUR=2
>1800	95	13	AABOUD	17AI	ATLS	$\geq 1\ell + \geq 8$ jets, Tglu1A and $\tilde{\chi}_1^0 \rightarrow qq\ell, \lambda'$ coupling, $m_{\tilde{\chi}_1^0} = 1000$ GeV	OCCUR=3
>1800	95	14	AABOUD	17AJ	ATLS	same-sign $\ell^\pm \ell^\pm / 3\ell + \text{jets} +$ \cancel{E}_T , Tglu3A, λ''_{112} coupling, $m_{\tilde{\chi}_1^0} = 50$ GeV	OCCUR=4
>1750	95	15	AABOUD	17AJ	ATLS	same-sign $\ell^\pm \ell^\pm / 3\ell + \text{jets} +$ \cancel{E}_T , Tglu1A and $\tilde{\chi}_1^0 \rightarrow qq\ell,$ λ' coupling	OCCUR=6
>1450	95	16	AABOUD	17AJ	ATLS	same-sign $\ell^\pm \ell^\pm / 3\ell + \text{jets} +$ $\cancel{E}_T, \tilde{g} \rightarrow t\tilde{t}_1$ and $\tilde{t}_1 \rightarrow sd,$ λ''_{321} coupling	OCCUR=8
>1450	95	17	AABOUD	17AJ	ATLS	same-sign $\ell^\pm \ell^\pm / 3\ell + \text{jets} +$ $\cancel{E}_T, \tilde{g} \rightarrow t\tilde{t}_1$ and $\tilde{t}_1 \rightarrow bd,$ λ''_{313} coupling	OCCUR=9
> 400	95	18	AABOUD	17AJ	ATLS	same-sign $\ell^\pm \ell^\pm / 3\ell + \text{jets} +$ $\cancel{E}_T, \tilde{d}_R \rightarrow tb(ts), \lambda''_{313}$ (λ''_{321}) coupling	OCCUR=10
none 625–1375	95	19	AABOUD	17AZ	ATLS	≥ 7 jets + \cancel{E}_T , large R-jets and/or b-jets, $\tilde{g} \rightarrow t\tilde{t}_1$ and $\tilde{t}_1 \rightarrow bs, \lambda''_{323}$ coupling	OCCUR=3
none 600–650	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqqqq, \lambda''_{212}$ coupling, $m_{\tilde{q}} = 100$ GeV	
none 600–1030	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqqqq, \lambda''_{212}$ coupling, $m_{\tilde{q}} = 900$ GeV	OCCUR=2
none 600–650	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqqqb, \lambda''_{213}$ coupling, $m_{\tilde{q}} = 100$ GeV	OCCUR=3
none 600–1080	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqqqb, \lambda''_{213}$ coupling, $m_{\tilde{q}} = 900$ GeV	OCCUR=4
none 600–680	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqqqbb, \lambda''_{212}$ coupling, $m_{\tilde{q}} = 100$ GeV	OCCUR=5
none 600–1080	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqqqbb, \lambda''_{212}$ coupling, $m_{\tilde{q}} = 900$ GeV	OCCUR=6
none 600–650	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqbbbb, \lambda''_{213}$ coupling, $m_{\tilde{q}} = 100$ GeV	OCCUR=7
none 600–1100	95	20	KHACHATRY...17Y		CMS	$\tilde{g} \rightarrow qqbbbb, \lambda''_{213}$ coupling, $m_{\tilde{q}} = 900$ GeV	OCCUR=8
>1050	95	21	KHACHATRY...16BJ		CMS	same-sign $\ell^\pm \ell^\pm$, Tglu3A, $m_{\tilde{\chi}_1^0} < 800$ GeV	OCCUR=2

>1140	95	21	KHACHATRY...16BJ	CMS	same-sign $\ell^\pm \ell^\pm$, Tglu3B, $m_{\tilde{t}} - m_{\tilde{\chi}_1^0} = 20$ GeV, $m_{\tilde{\chi}_1^0} = 0$	OCCUR=5
>1030	95	22	KHACHATRY...16BX	CMS	$\tilde{g} \rightarrow tbs$, λ''_{332} coupling	
>1150	95	23	AAD	15BV ATLS	general RPC \tilde{g} decays, $m_{\tilde{\chi}_1^0} < 100$ GeV	
>1350	95	24	AAD	14X ATLS	$\geq 4\ell^\pm, \tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$	
> 650	95	25	CHATRCHYAN 14P	CMS	$\tilde{g} \rightarrow jjj$	
none 200–835	95	25	CHATRCHYAN 14P	CMS	$\tilde{g} \rightarrow bjj$	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
>1875	95	26	AABOUD	18CF ATLS	jets and large R-jets, Tglu2RPV and $\tilde{\chi}_1^0 \rightarrow qq\bar{q}$, λ'' coupling, $m_{\tilde{\chi}_1^0} = 1000$ GeV	
>1400	95	27	KHACHATRY...16BX	CMS	$\tilde{g} \rightarrow qq\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell\ell\nu, \lambda_{121}$ or $\lambda_{122} \neq 0, m_{\tilde{\chi}_1^0} > 400$ GeV	OCCUR=2
>1600	95	23	AAD	15BV ATLS	pMSSM, $M_1 = 60$ GeV, $m_{\tilde{q}} < 1500$ GeV	OCCUR=2
>1280	95	23	AAD	15BV ATLS	mSUGRA, $m_0 > 2$ TeV	OCCUR=3
>1100	95	23	AAD	15BV ATLS	via $\tilde{\tau}$, natural GMSB, all $m_{\tilde{\tau}}$	OCCUR=4
>1220	95	23	AAD	15BV ATLS	b -jets, $\tilde{g} \rightarrow \tilde{t}_1 t$ and $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$, $m_{T_1} < 1000$ GeV	OCCUR=14
>1180	95	23	AAD	15BV ATLS	b -jets, $\tilde{g} \rightarrow \tilde{t}_1 t$ and $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm, m_{T_1} < 1000$ GeV, $m_{\tilde{\chi}_1^0} = 60$ GeV	OCCUR=15
> 880	95	23	AAD	15BV ATLS	jets, $\tilde{g} \rightarrow \tilde{t}_1 t$ and $\tilde{t}_1 \rightarrow sb$, $400 < m_{\tilde{t}_1} < 1000$ GeV	OCCUR=17
> 600	95	28	AAD	15CB ATLS	$\ell, \tilde{g} \rightarrow (e/\mu)qq$, benchmark gluino, neutralino masses	
> 600	95	28	AAD	15CB ATLS	$\ell\ell/Z, \tilde{g} \rightarrow (ee/\mu\mu/e\mu)qq$, $m_{\tilde{\chi}_1^0} = 400$ GeV and $0.7 < c\tau_{\tilde{\chi}_1^0} < 3 \times 10^5$ mm	OCCUR=2
>1000	95	29	AAD	15X ATLS	≥ 10 jets, $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow qq\bar{q}$, $m_{\tilde{\chi}_1^0} = 500$ GeV	
> 917	95	29	AAD	15X ATLS	$\geq 6,7$ jets, $\tilde{g} \rightarrow qq\bar{q}$, (light-quark, λ'' couplings)	OCCUR=2
> 929	95	29	AAD	15X ATLS	$\geq 6,7$ jets, $\tilde{g} \rightarrow qq\bar{q}$, (b -quark, λ'' couplings)	OCCUR=3
>1180	95	30	AAD	14AX ATLS	≥ 3 b -jets + \cancel{E}_T , $\tilde{g} \rightarrow \tilde{t}_1 t\tilde{\chi}_1^0$ simplified model, $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$, $m_{\tilde{\chi}_1^\pm} = 2m_{\tilde{\chi}_1^0}, m_{\tilde{\chi}_1^0} = 60$ GeV, $m_{\tilde{t}_1} < 1000$ GeV	OCCUR=4
> 850	95	31	AAD	14E ATLS	$\ell^\pm \ell^\pm (\ell^\mp) +$ jets, $\tilde{g} \rightarrow t\tilde{t}_1$ with $\tilde{t}_1 \rightarrow bs$ simplified model	OCCUR=4
> 900	95	32	CHATRCHYAN 14H	CMS	same-sign $\ell^\pm \ell^\pm, \tilde{g} \rightarrow tbs$ simplified model	OCCUR=4

¹ AAD 21BF searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for pair production of gluinos, stops, electroweakinos decaying RPV either directly or indirectly via the LSP. The final state in all cases is one or two leptons, many jets (up to fifteen) and b -jets. Different models with different branching fractions of the gluino or stop follow from the assumptions on the nature of the electroweakinos. No significant excess above the Standard Model predictions is observed. Limits are set on the *gluino*, \tilde{t}_1 , electroweakino masses as a function of the $\tilde{\chi}_1^0$ mass in several scenarios of gluino, stop and electroweakino pair production.

² AAD 21Y searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for supersymmetry in events with four or more leptons (electrons, muons and tau-leptons). No significant excess above the Standard Model expectations is observed. Limits are set on Tchi1n12-GGM, and RPV models similar to Tchi1n2l, Tglu1A (with $q = u, d, s, c, b$, with equal branching fractions), and $\tilde{\ell}_L/\tilde{\nu} \rightarrow \ell/\nu\tilde{\chi}_1^0$ (mass-degenerate $\tilde{\ell}_L$ and $\tilde{\nu}$ of all 3 generations), all with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$ via λ_{12k} or λ_{i33} (where $i, k \in 1, 2$), see their Figure 11.

³ AAD 20AL searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV for events with 8 or more jets and moderate missing transverse momentum. The selection makes requirements according to the number of b -tagged jets and the scalar sum of masses of large-radius

NODE=S046GNV;LINKAGE=EF

NODE=S046GNV;LINKAGE=K

NODE=S046GNV;LINKAGE=H

- jets. No significant excess above the Standard Model expectations is observed. Exclusion limits at 95% C.L. are set on the gluino mass in RPV simplified models where the gluino decays via $\tilde{g} \rightarrow tbd$ or $\tilde{g} \rightarrow tbs$. They extend up to almost 1.6 TeV for a \tilde{t}_1 mass of 900 GeV. See their Fig. 10(c).
- 4 AAD 20V searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign charged leptons (electrons or muons) and jets. No significant excess above the Standard Model expectations is observed. Exclusion limits at 95% C.L. are set on the gluino mass in RPV simplified models where the gluino decays via $\tilde{g} \rightarrow tbd$, see Figure 7(b). NODE=S046GNV;LINKAGE=G
- 5 SIRUNYAN 20T searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least two jets, and two isolated same-sign or three or more charged leptons (electrons or muons). No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu3A, Tglu3B, Tglu3C and Tglu3D simplified models, see their Figure 7, and in the Tglu1C and Tglu1B simplified models, see their Figures 8 and 9. Limits are also set on the sbottom mass in the Tsbott2 simplified model, see their Figure 10, and on the stop mass in the Tstop7 simplified model, see their Figure 11. Finally, limits are set on the gluino mass in RPV simplified models where the gluino decays either via $\tilde{g} \rightarrow qq\bar{q}\bar{q} + e/\mu/\tau$ or via $\tilde{g} \rightarrow tbs$, see Figure 12. NODE=S046GNV;LINKAGE=F
- 6 SIRUNYAN 19F searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for three-jet resonances produced in the decay of a gluino in R-parity violating supersymmetric models. The mass range from 200 to 2000GeV is explored in four separate mass regions. The observations show agreement with standard model expectations. The results are interpreted within the framework of R-parity violating SUSY, where pair-produced gluinos decay to a six quark final state. Gluino masses below 1500GeV are excluded at 95% C.L. See their Fig.5. NODE=S046GNV;LINKAGE=E
- 7 AABOUD 18Z searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing four or more charged leptons (electrons, muons and up to two hadronically decaying taus). No significant deviation from the expected SM background is observed. Limits are set on the Higgsino mass in simplified models of general gauge mediated supersymmetry Tn1n1A/Tn1n1B/Tn1n1C, see their Figure 9. Limits are also set on the wino, slepton, sneutrino and gluino mass in a simplified model of NLSP pair production with R-parity violating decays of the LSP via λ_{12k} or λ_{j33} to charged leptons, see their Figures 7, 8. NODE=S046GNV;LINKAGE=GE
- 8 SIRUNYAN 18AK searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing a single lepton, large jet and b -quark jet multiplicities, coming from R-parity-violating decays of gluinos. No excess over the expected background is observed. Limits are derived on the gluino mass, assuming the RPV $\tilde{g} \rightarrow tbs$ decay, see their Figure 9. NODE=S046GNV;LINKAGE=OD
- 9 SIRUNYAN 18D searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing identified hadronically decaying top quarks, no leptons, and \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the stop mass in the Tstop1 simplified model, see their Figure 8, and on the gluino mass in the Tglu3A, Tglu3B, Tglu3C and Tglu3E simplified models, see their Figure 9. NODE=S046GNV;LINKAGE=KC
- 10 SIRUNYAN 18EA searched in 38.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for the pair production of resonances, each decaying to at least four quarks. Reconstructed particles are clustered into two large jets of similar mass, each consistent with four-parton substructure. No statistically significant excess over the Standard Model expectation is observed. Limits are set on the squark and gluino mass in RPV supersymmetry models where squarks (gluinos) decay, through intermediate higgsinos, to four (five) quarks, see their Figure 4. NODE=S046GNV;LINKAGE=IE
- 11 AABOUD 17Al searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more isolated lepton, at least eight jets, either zero or many b -jets, for evidence of R-parity violating decays of the gluino. No significant excess above the Standard Model expectations is observed. Limits up to 2.1 TeV are set on the gluino mass in R-parity-violating supersymmetry models as Tglu3A with LSP decay through the non-zero λ''_{112} coupling as $\tilde{\chi}_1^0 \rightarrow uds$. See their Figure 9. NODE=S046GNV;LINKAGE=LC
- 12 AABOUD 17Al searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more isolated lepton, at least eight jets, either zero or many b -jets, for evidence of R-parity violating decays of the gluino. No significant excess above the Standard Model expectations is observed. Limits up to 1.65 TeV are set on the gluino mass in R-parity-violating supersymmetry models with $\tilde{g} \rightarrow t\tilde{t}$, $\tilde{t} \rightarrow bs$ through the non-zero λ''_{323} coupling. See their Figure 9. NODE=S046GNV;LINKAGE=OC
- 13 AABOUD 17Al searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with one or more isolated lepton, at least eight jets, either zero or many b -jets, for evidence of R-parity violating decays of the gluino. No significant excess above the Standard Model expectations is observed. Limits up to 1.8 TeV are set on the gluino mass in R-parity-violating supersymmetry models as Tglu1A with the LSP decay through the non-zero λ' coupling as $\tilde{\chi}_1^0 \rightarrow qq\ell$. See their Figure 9. NODE=S046GNV;LINKAGE=PC
- 14 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.8 TeV are set on the gluino mass in R-parity-violating supersymmetry models as Tglu3A with LSP decaying through the non-zero λ''_{112} coupling as $\tilde{\chi}_1^0 \rightarrow uds$. See their Figure 5(d). NODE=S046GNV;LINKAGE=DD
- 15 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.75 TeV are NODE=S046GNV;LINKAGE=FD

- set on the gluino mass in R-parity-violating supersymmetry models as Tglu1A with LSP decaying through the non-zero λ' coupling as $\tilde{\chi}_1^0 \rightarrow qql$. See their Figure 5(c).
- 16 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.45 TeV are set on the gluino mass in R-parity-violating supersymmetry models where $\tilde{g} \rightarrow t\tilde{t}_1$ and $\tilde{t}_1 \rightarrow sd$ through the non-zero λ''_{321} coupling. See their Figure 5(b). NODE=S046GNV;LINKAGE=ID
- 17 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 1.45 TeV are set on the gluino mass in R-parity-violating supersymmetry models where $\tilde{g} \rightarrow t\tilde{t}_1$ and $\tilde{t}_1 \rightarrow bd$ through the non-zero λ''_{313} coupling. See their Figure 5(a). NODE=S046GNV;LINKAGE=JD
- 18 AABOUD 17AJ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two same-sign or three leptons, jets and large missing transverse momentum. No significant excess above the Standard Model expectations is observed. Limits up to 400 GeV are set on the down type squark (\tilde{d}_R mass in R-parity-violating supersymmetry models where $\tilde{d}_R \rightarrow tb$ through the non-zero λ''_{313} coupling or $\tilde{d}_R \rightarrow ts$ through the non-zero λ''_{321} . See their Figure 5(e) and 5(f). NODE=S046GNV;LINKAGE=KD
- 19 AABOUD 17AZ searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with at least seven jets and large missing transverse momentum. Selected events are further classified based on the presence of large R-jets or b -jets and no leptons. No significant excess above the Standard Model expectations is observed. Limits are set for R-parity violating decays of the gluino assuming $\tilde{g} \rightarrow t\tilde{t}_1$ and $\tilde{t}_1 \rightarrow bs$ through the non-zero λ''_{323} couplings. The range 625–1375 GeV is excluded for $m_{\tilde{t}_1} = 400 \text{ GeV}$. See their Figure 7b. NODE=S046GNV;LINKAGE=WC
- 20 KHACHATRYAN 17Y searched in 19.7 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing at least 8 or 10 jets, possibly b -tagged, coming from R-parity-violating decays of supersymmetric particles. No excess over the expected background is observed. Limits are derived on the gluino mass, assuming various RPV decay modes, see Fig. 7. NODE=S046GNV;LINKAGE=AB
- 21 KHACHATRYAN 16BJ searched in 2.3 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the following simplified models: Tglu3A and Tglu3D, see Fig. 4, Tglu3B and Tglu3C, see Fig. 5, and Tglu1B, see Fig. 7. NODE=S046GNV;LINKAGE=NB
- 22 KHACHATRYAN 16BX searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing 0 or 1 leptons and b -tagged jets, coming from R-parity-violating decays of supersymmetric particles. No excess over the expected background is observed. Limits are derived on the gluino mass, assuming the RPV $\tilde{g} \rightarrow tbs$ decay, see Fig. 7 and 10. NODE=S046GNV;LINKAGE=QB
- 23 AAD 15BV summarized and extended ATLAS searches for gluinos and first- and second-generation squarks in final states containing jets and missing transverse momentum, with or without leptons or b -jets in the $\sqrt{s} = 8 \text{ TeV}$ data set collected in 2012. The paper reports the results of new interpretations and statistical combinations of previously published analyses, as well as new analyses. Exclusion limits at 95% C.L. are set on the gluino mass in several R-parity conserving models, leading to a generalized constraint on gluino masses exceeding 1150 GeV for lightest supersymmetric particle masses below 100 GeV. See their Figs. 10, 19, 20, 21, 23, 25, 26, 29-37. NODE=S046GNV;LINKAGE=OA
- 24 AAD 14X searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with at least four leptons (electrons, muons, taus) in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in an R-parity violating simplified model where the decay $\tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$, with $\tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu$, takes place with a branching ratio of 100%, see Fig. 8. NODE=S046GNV;LINKAGE=D
- 25 CHATRCHYAN 14P searched in 19.4 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for three-jet resonances produced in the decay of a gluino in R-parity violating supersymmetric models. No excess over the expected SM background is observed. Assuming a 100% branching ratio for the gluino decay into three light-flavour jets, limits are set on the cross section of gluino pair production, see Fig. 7, and gluino masses below 650 GeV are excluded at 95% C.L. Assuming a 100% branching ratio for the gluino decaying to one b -quark jet and two light-flavour jets, gluino masses between 200 GeV and 835 GeV are excluded at 95% C.L. NODE=S046GNV;LINKAGE=HC
- 26 AABOUD 18CF searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with several jets, possibly b -jets, and large-radius jets for evidence of R-parity violating decays of the gluino. No significant excess above the Standard Model expectations is observed. Limits between 1000 and 1875 GeV are set on the gluino mass in R-parity-violating supersymmetry models as Tglu2RPV with the LSP decay through the non-zero λ'' coupling as $\tilde{\chi}_1^0 \rightarrow qq\bar{q}$. The most stringent limit is obtained for $m_{\tilde{\chi}_1^0} = 1000 \text{ GeV}$, the weakest for $m_{\tilde{\chi}_1^0} = 50 \text{ GeV}$. See their Figure 7(b). Figure 7(a) presents results for gluinos directly decaying into 3 quarks, Tglu1RPV. NODE=S046GNV;LINKAGE=HE
- 27 KHACHATRYAN 16BX searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing 4 leptons coming from R-parity-violating decays of $\tilde{\chi}_1^0 \rightarrow \ell\ell\nu$ with $\lambda_{121} \neq 0$ or $\lambda_{122} \neq 0$. No excess over the expected background is observed. Limits are derived on the gluino, squark and stop masses, see Fig. 23. NODE=S046GNV;LINKAGE=RB
- 28 AAD 15CB searched for events containing at least one long-lived particle that decays at a significant distance from its production point (displaced vertex, DV) into two leptons NODE=S046GNV;LINKAGE=TA

or into five or more charged particles in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$. The dilepton signature is characterised by DV formed from at least two lepton candidates. Four different final states were considered for the multitrak signature, in which the DV must be accompanied by a high-transverse momentum muon or electron candidate that originates from the DV, jets or missing transverse momentum. No events were observed in any of the signal regions. Results were interpreted in SUSY scenarios involving R -parity violation, split supersymmetry, and gauge mediation. See their Fig. 12–20.

- 29 AAD 15X searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events containing large number of jets, no requirements on missing transverse momentum and no isolated electrons or muons. The sensitivity of the search is enhanced by considering the number of b -tagged jets and the scalar sum of masses of large-radius jets in an event. No evidence was found for excesses above the expected level of Standard Model background. Exclusion limits at 95% C.L. are set on the gluino mass assuming the gluino decays to various quark flavors, and for various neutralino masses. See their Fig. 11–16. NODE=S046GNV;LINKAGE=WA
- 30 AAD 14AX searched in 20.1 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for the strong production of supersymmetric particles in events containing either zero or at last one high high- p_T lepton, large missing transverse momentum, high jet multiplicity and at least three jets identified as originating from b -quarks. No excess over the expected SM background is observed. Limits are derived in mSUGRA/CMSSM models with $\tan\beta = 30$, $A_0 = -2m_0$ and $\mu > 0$, see their Fig. 14. Also, exclusion limits in simplified models containing gluinos and scalar top and bottom quarks are set, see their Figures 12, 13. NODE=S046GNV;LINKAGE=BA
- 31 AAD 14E searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for strongly produced supersymmetric particles in events containing jets and two same-sign leptons or three leptons. The search also utilises jets originating from b -quarks, missing transverse momentum and other variables. No excess over the expected SM background is observed. Exclusion limits are derived in simplified models containing gluinos and squarks, see Figures 5 and 6. In the $\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^{(*)\pm}\tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow Z^{(*)}\tilde{\chi}_1^0$ simplified model, the following assumptions have been made: $m_{\tilde{\chi}_1^\pm} = 0.5 m_{\tilde{\chi}_1^0} + m_{\tilde{g}}$, $m_{\tilde{\chi}_2^0} = 0.5 (m_{\tilde{\chi}_1^0} + m_{\tilde{\chi}_1^\pm})$, $m_{\tilde{\chi}_1^0} < 520 \text{ GeV}$. In the $\tilde{g} \rightarrow qq'\tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow \ell^\pm \nu \tilde{\chi}_1^0$ or $\tilde{g} \rightarrow qq'\tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow \ell^\pm \ell^\mp (\nu\nu)\tilde{\chi}_1^0$ simplified model, the following assumptions have been made: $m_{\tilde{\chi}_1^\pm} = m_{\tilde{\chi}_2^0} = 0.5 (m_{\tilde{\chi}_1^0} + m_{\tilde{g}})$, $m_{\tilde{\chi}_1^0} < 660 \text{ GeV}$. Limits are also derived in the mSUGRA/CMSSM, bRPV and GMSB models, see their Fig. 8. NODE=S046GNV;LINKAGE=B2
- 32 CHATRCHYAN 14H searched in 19.5 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with two isolated same-sign dileptons and jets in the final state. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in simplified models where the R -parity violating decay $\tilde{g} \rightarrow tbs$ takes place with a branching ratio of 100%, see Fig. 8. NODE=S046GNV;LINKAGE=P

Long-lived \tilde{g} (Gluino) mass limit

Limits on light gluinos ($m_{\tilde{g}} < 5 \text{ GeV}$) were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT	
>2050	95	1 AAD	23G ATLS	R -hadrons, Tglu1A, stable, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	NODE=S046LGN
>2270	95	1 AAD	23G ATLS	R -hadrons, Tglu1A, $\tau = 20 \text{ ns}$, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	OCCUR=2
>2050	95	1 AAD	23G ATLS	R -hadrons, Tglu1A, stable, $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} = 30 \text{ GeV}$	OCCUR=3
>2050	95	1 AAD	23G ATLS	R -hadrons, Tglu1A, $\tau = 20 \text{ ns}$, $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} = 30 \text{ GeV}$	OCCUR=4
>2500	95	2 SIRUNYAN	21AF CMS	long-lived \tilde{g} , Tglu2RPV, λ_{323}'' coupling, $0.6 \text{ mm} < c\tau < 90 \text{ mm}$	
>2450	95	3 SIRUNYAN	21U CMS	long-lived \tilde{g} , $pp \rightarrow \tilde{g}\tilde{g}, \tilde{g} \rightarrow g\tilde{G}$, GMSB, $6 < c\tau < 550 \text{ mm}$	OCCUR=3
>2500	95	3 SIRUNYAN	21U CMS	long-lived \tilde{g} , $pp \rightarrow \tilde{g}\tilde{g}, \tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$, mini-split, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$, $7 < c\tau < 360 \text{ mm}$	OCCUR=3
>2500	95	3 SIRUNYAN	21U CMS	long-lived \tilde{g} , $pp \rightarrow \tilde{g}\tilde{g}, \tilde{g} \rightarrow tbs$, λ_{323}'' coupling, $3 < c\tau < 1000 \text{ mm}$	OCCUR=5
>1980	95	4 AABOUD	19AT ATLS	R -hadrons, Tglu1A, metastable	
>2060	95	5 AABOUD	19C ATLS	R -hadrons, Tglu1A, $\tau \geq 10 \text{ ns}$, $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$	

>1890	95	⁵ AABOUD	19C ATLS	R -hadrons, Tglu1A, stable	OCCUR=2
>2400	95	⁶ SIRUNYAN	19BH CMS	long-lived \tilde{g} , RPV, $\tilde{g} \rightarrow \bar{t}b\bar{s}$, 10 mm < $c\tau$ < 250 mm	
>2300	95	⁶ SIRUNYAN	19BH CMS	long-lived \tilde{g} , GMSB, $\tilde{g} \rightarrow$ $g\tilde{G}$, 20 mm < $c\tau$ < 110 mm	OCCUR=2
>2100	95	⁷ SIRUNYAN	19BT CMS	long-lived \tilde{g} , GMSB, $\tilde{g} \rightarrow$ $g\tilde{G}$, 0.3 m < $c\tau$ < 30 m	
>2500	95	⁷ SIRUNYAN	19BT CMS	long-lived \tilde{g} , GMSB, $\tilde{g} \rightarrow$ $g\tilde{G}$, $c\tau = 1$ m	OCCUR=2
>1900	95	⁷ SIRUNYAN	19BT CMS	long-lived \tilde{g} , GMSB, $\tilde{g} \rightarrow$ $g\tilde{G}$, $c\tau = 100$ m	OCCUR=3
>2370	95	⁸ AABOUD	18S ATLS	displaced vertex + \cancel{E}_T , long- lived Tglu1A, $m_{\tilde{\chi}_1^0} = 100$ GeV, and $\tau=0.17$ ns	
>1600	95	⁹ SIRUNYAN	18AY CMS	jets+ \cancel{E}_T , Tglu1A, $c\tau < 0.1$ mm, $m_{\tilde{\chi}_1^0} = 100$ GeV	
>1750	95	⁹ SIRUNYAN	18AY CMS	jets+ \cancel{E}_T , Tglu1A, $c\tau = 1$ mm, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=2
>1640	95	⁹ SIRUNYAN	18AY CMS	jets+ \cancel{E}_T , Tglu1A, $c\tau = 10$ mm, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=3
>1490	95	⁹ SIRUNYAN	18AY CMS	jets+ \cancel{E}_T , Tglu1A, $c\tau = 100$ mm, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=4
>1300	95	⁹ SIRUNYAN	18AY CMS	jets+ \cancel{E}_T , Tglu1A, $c\tau = 1$ m, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=5
> 960	95	⁹ SIRUNYAN	18AY CMS	jets+ \cancel{E}_T , Tglu1A, $c\tau = 10$ m, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=6
> 900	95	⁹ SIRUNYAN	18AY CMS	jets+ \cancel{E}_T , Tglu1A, $c\tau = 100$ m, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=7
>2200	95	¹⁰ SIRUNYAN	18DV CMS	long-lived \tilde{g} , RPV, $\tilde{g} \rightarrow \bar{t}b\bar{s}$, 0.6 mm < $c\tau$ < 80 mm	
>1000	95	¹¹ KHACHATRY...17AR	CMS	long-lived \tilde{g} , RPV, $\tilde{g} \rightarrow t\bar{b}\bar{s}$, $c\tau = 0.3$ mm	
>1300	95	¹¹ KHACHATRY...17AR	CMS	long-lived \tilde{g} , RPV, $\tilde{g} \rightarrow t\bar{b}\bar{s}$, $c\tau = 1.0$ mm	OCCUR=2
>1400	95	¹¹ KHACHATRY...17AR	CMS	long-lived \tilde{g} , RPV, $\tilde{g} \rightarrow t\bar{b}\bar{s}$, 2 mm < $c\tau$ < 30 mm	OCCUR=3
>1580	95	¹² AABOUD	16B ATLS	long-lived R -hadrons	
> 740–1590	95	¹³ AABOUD	16C ATLS	R -hadrons, Tglu1A, $\tau \geq 0.4$ ns, $m_{\tilde{\chi}_1^0} = 100$ GeV	
>1570	95	¹³ AABOUD	16C ATLS	R -hadrons, Tglu1A, stable	OCCUR=3
>1610	95	¹⁴ KHACHATRY...16BWCMS		long-lived \tilde{g} forming R - hadrons, $f = 0.1$, cloud interaction model	
>1580	95	¹⁴ KHACHATRY...16BWCMS		long-lived \tilde{g} forming R - hadrons, $f = 0.1$, charge- suppressed interaction model	OCCUR=2
>1520	95	¹⁴ KHACHATRY...16BWCMS		long-lived \tilde{g} forming R - hadrons, $f = 0.5$, cloud interaction model	OCCUR=3
>1540	95	¹⁴ KHACHATRY...16BWCMS		long-lived \tilde{g} forming R - hadrons, $f = 0.5$, charge- suppressed interaction model	OCCUR=4
>1270	95	¹⁵ AAD	15AE ATLS	\tilde{g} R -hadron, generic R -hadron model	
>1360	95	¹⁵ AAD	15AE ATLS	\tilde{g} decaying to 300 GeV stable sleptons, LeptoSUSY model	OCCUR=2
>1115	95	¹⁶ AAD	15BMATLS	\tilde{g} R -hadron, stable	
>1185	95	¹⁶ AAD	15BMATLS	$\tilde{g} \rightarrow (g/q\bar{q})\tilde{\chi}_1^0$, lifetime 10 ns, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=2
>1099	95	¹⁶ AAD	15BMATLS	$\tilde{g} \rightarrow (g/q\bar{q})\tilde{\chi}_1^0$, lifetime 10 ns, $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=3
>1182	95	¹⁶ AAD	15BMATLS	$\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$, lifetime 10 ns, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=4
>1157	95	¹⁶ AAD	15BMATLS	$\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$, lifetime 10 ns, $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} = 480$ GeV	OCCUR=5
> 869	95	¹⁶ AAD	15BMATLS	$\tilde{g} \rightarrow (g/q\bar{q})\tilde{\chi}_1^0$, lifetime 1 ns, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=6

> 821	95	16 AAD	15BMATLS	$\tilde{g} \rightarrow (g/q\bar{q})\tilde{\chi}_1^0$, lifetime 1 ns, $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=7
> 836	95	16 AAD	15BMATLS	$\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$, lifetime 1 ns, $m_{\tilde{\chi}_1^0} = 100$ GeV	OCCUR=8
> 836	95	16 AAD	15BMATLS	$\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$, lifetime 10 ns, $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} = 480$ GeV	OCCUR=9
>1000	95	17 KHACHATRY...15AK	CMS	\tilde{g} R-hadrons, $10 \mu\text{s} < \tau < 1000$ s	
> 880	95	17 KHACHATRY...15AK	CMS	\tilde{g} R-hadrons, $1 \mu\text{s} < \tau < 1000$ s	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
> 985	95	18 AAD	13AA ATLS	\tilde{g} , R-hadrons, generic interaction model	
> 832	95	19 AAD	13BC ATLS	R-hadrons, $\tilde{g} \rightarrow g/q\bar{q}\tilde{\chi}_1^0$, generic R-hadron model, lifetime between 10^{-5} and 10^3 s, $m_{\tilde{\chi}_1^0} = 100$ GeV	
>1322	95	20 CHATRCHYAN 13AB	CMS	long-lived \tilde{g} forming R-hadrons, $f = 0.1$, cloud interaction model	
none 200–341	95	21 AAD	12P ATLS	long-lived $\tilde{g} \rightarrow g\tilde{\chi}_1^0$, $m_{\tilde{\chi}_1^0} = 100$ GeV	
> 640	95	22 CHATRCHYAN 12AN	CMS	long-lived $\tilde{g} \rightarrow g\tilde{\chi}_1^0$	
>1098	95	23 CHATRCHYAN 12L	CMS	long-lived \tilde{g} forming R-hadrons, $f = 0.1$	
> 586	95	24 AAD	11K ATLS	stable \tilde{g}	
> 544	95	25 AAD	11P ATLS	stable \tilde{g} , GMSB scenario, $\tan\beta=5$	
> 370	95	26 KHACHATRY...11	CMS	long lived \tilde{g}	
> 398	95	27 KHACHATRY...11C	CMS	stable \tilde{g}	

¹ AAD 23G searched in 139 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for R-hadron pair production in events with high-pt tracks with large ionisation in the pixel detector. No significant excess above the Standard Model predictions is observed. Limits are set on the R-hadron mass for different masses of the LSP and for different R-hadron lifetimes, see Figure 18.

NODE=S046LGN;LINKAGE=PA

² SIRUNYAN 21AF searched in 140 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with with two displaced vertices from long-lived particles decaying into multijet or dijet final states. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the simplified model Tglu2RPV with λ''_{323} coupling, on the $\tilde{\chi}_1^0$ mass in an RPV model with $\tilde{\chi}_1^0$ pair production and the RPV decay $\tilde{\chi}_1^0 \rightarrow tbs$ with λ''_{323} coupling and on the \tilde{t} mass in an RPV model with top squark pair production and the RPV decay $\tilde{t} \rightarrow \bar{d}_i \bar{d}_j$ with λ''_{3ij} coupling, see their Figure 7.

NODE=S046LGN;LINKAGE=OA

³ SIRUNYAN 21U searched in 132 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for supersymmetry in events with displaced tracks and displaced vertices associated with a dijet system. No significant excess above the Standard Model expectations is observed. Limits are set on long-lived gluinos in an RPC GMSB SUSY model of gluino pair production, with $\tilde{g} \rightarrow g\tilde{G}$, see their Figure 9, in Tglu1A in a mini-split model, see their Figure 10, and in an RPV model of gluino pair production, with $\tilde{g} \rightarrow tbs$ with coupling λ''_{323} , see their Figure 11. Limits are also set on long-lived top squarks in Tstop2RPV, see their Figure 12, in an RPV model with $\tilde{t} \rightarrow d\bar{\ell}$ and λ'_{x31} coupling, see their Figure 13, and in a dynamical RPV model with $\tilde{t} \rightarrow \bar{d}\bar{d}$ via a nonholomorphic RPV coupling η''_{311} , see their Figure 14. The best mass limit is achieved in all cases at $c\tau = 30 \text{ mm}$.

NODE=S046LGN;LINKAGE=NA

⁴ AABOUD 19AT searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for metastable and stable R-hadrons. Multiple search strategies for a wide range of lifetimes, corresponding to path lengths of a few meters, are defined. No significant deviations from the expected Standard Model background are observed. Gluino R-hadrons with lifetimes of the order of 50 ns are excluded at 95% C.L. for masses below 1980 GeV using the muon-spectrometer agnostic analysis. Using the full-detector search, the observed lower limits on the mass are 2000 GeV. See their Figure 9 (top).

NODE=S046LGN;LINKAGE=IA

⁵ AABOUD 19C searched in 36.1 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for metastable and stable R-hadrons arising as excesses in the mass distribution of reconstructed tracks with high transverse momentum and large dE/dx . Gluino R-hadrons with lifetimes above 10 ns are excluded at 95% C.L. with lower mass limit range between 1000 GeV and 2060 GeV, see their Figure 5(a). Masses smaller than 1290 GeV are excluded for a lifetime of 1 ns, see their Figure 6. In the case of stable R-hadrons, the lower mass limit is 1890 GeV, see their Figure 5(b).

NODE=S046LGN;LINKAGE=EA

⁶ SIRUNYAN 19BH searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived particles decaying into jets, with each long-lived particle having a decay vertex well displaced from the production vertex. The selected events are found to be consistent

NODE=S046LGN;LINKAGE=FA

- with standard model predictions. Limits are set on the gluino mass in a GMSB model where the gluino is decaying via $\tilde{g} \rightarrow g\tilde{G}$, see their Figure 4 and in an RPV model of supersymmetry where the gluino is decaying via $\tilde{g} \rightarrow \tilde{t}\bar{b}s$, see their Figures 5. Limits are also set on the stop mass in two RPV models, see their Figure 6 (for $\tilde{t} \rightarrow b\ell$ decays) and Figure 7 (for $\tilde{t} \rightarrow \bar{d}\bar{d}$ decays).
- 7 SIRUNYAN 19BT searched in 137 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived particles decaying to displaced, nonprompt jets and missing transverse momentum. Candidate signal events are identified using the timing capabilities of the CMS electromagnetic calorimeter. The results of the search are found to be consistent with the background predictions. Limits are set on the gluino mass in a GMSB model where long-lived gluinos are pair produced and decaying via $\tilde{g} \rightarrow g\tilde{G}$, see their Figures 4 and 5. NODE=S046LGN;LINKAGE=LA
- 8 AABOUD 18S searched in 32.8 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived gluinos in final states with large missing transverse momentum and at least one high-mass displaced vertex with five or more tracks. The observed yield is consistent with the expected background. Exclusion limits are derived for Tglu1A models predicting the existence of long-lived gluinos reaching roughly $m(\tilde{g}) = 2000 \text{ GeV}$ to 2370 GeV for $m(\tilde{\chi}_1^0) = 100 \text{ GeV}$ and gluino lifetimes between 0.02 and 10 ns , see their Fig. 8. Limits are presented also as a function of the lifetime (for a fixed gluino-neutralino mass difference of 100 GeV) and of the gluino and neutralino masses (for a fixed lifetime of 1 ns). See their Fig. 9 and 10 respectively. NODE=S046LGN;LINKAGE=X
- 9 SIRUNYAN 18AY searched in 35.9 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events containing one or more jets and significant \cancel{E}_T . No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass in the Tglu1A, Tglu2A and Tglu3A simplified models, see their Figure 3. Limits are also set on squark, sbottom and stop masses in the Tsqk1, Tsb0t1, Tstop1 and Tstop4 simplified models, see their Figure 3. Finally, limits are set on long-lived gluino masses in a Tglu1A simplified model where the gluino is metastable or long-lived with proper decay lengths in the range $10^{-3} \text{ mm} < c\tau < 10^5 \text{ mm}$, see their Figure 4. NODE=S046LGN;LINKAGE=BA
- 10 SIRUNYAN 18DV searched in 38.5 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived particles in events with multiple jets and two displaced vertices composed of many tracks. No events with two well-separated high-track-multiplicity vertices were observed. Limits are set on the stop and the gluino mass in RPV models of supersymmetry where the stop (gluino) is decaying solely into dijet (multijet) final states, see their Figures 6 and 7. NODE=S046LGN;LINKAGE=CA
- 11 KHACHATRYAN 17AR searched in 17.6 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for R-parity-violating SUSY in which long-lived neutralinos or gluinos decay into multijet final states. No significant excess above the Standard Model expectations is observed. Limits are set on the gluino mass for a range of mean proper decay lengths ($c\tau$), see their Fig. 7. The upper limits on the production cross section times branching ratio squared (Fig. 7) are also applicable to long-lived neutralinos. NODE=S046LGN;LINKAGE=O
- 12 AABOUD 16B searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived R -hadrons using observables related to large ionization losses and slow propagation velocities, which are signatures of heavy charged particles traveling significantly slower than the speed of light. Exclusion limits at 95% C.L. are set on the long-lived gluino masses exceeding 1580 GeV . See their Fig. 5. NODE=S046LGN;LINKAGE=R
- 13 AABOUD 16C searched in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for long-lived and stable R -hadrons identified by anomalous specific ionization energy loss in the ATLAS Pixel detector. Gluino R -hadrons with lifetimes above 0.4 ns are excluded at 95% C.L. with lower mass limit range between 740 GeV and 1590 GeV . In the case of stable R -hadrons, the lower mass limit is 1570 GeV . See their Figs. 5 and 6. NODE=S046LGN;LINKAGE=K
- 14 KHACHATRYAN 16BW searched in 2.5 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$ for events with heavy stable charged particles, identified by their anomalously high energy deposits in the silicon tracker and/or long time-of-flight measurements by the muon system. No evidence for an excess over the expected background is observed. Limits are derived for pair production of gluinos as a function of mass, depending on the interaction model and on the fraction f , of produced gluinos hadronizing into a \tilde{g} -gluon state, see Fig. 4 and Table 7. NODE=S046LGN;LINKAGE=S
- 15 AAD 15AE searched in 19.1 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for heavy long-lived charged particles, measured through their specific ionization energy loss in the ATLAS pixel detector or their time-of-flight in the ALTAS muon system. In the absence of an excess of events above the expected backgrounds, limits are set R -hadrons in various scenarios, see Fig. 11. Limits are also set in LeptoSUSY models where the gluino decays to stable 300 GeV leptons, see Fig. 9. NODE=S046LGN;LINKAGE=L
- 16 AAD 15BM searched in 18.4 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for stable and metastable non-relativistic charged particles through their anomalous specific ionization energy loss in the ATLAS pixel detector. In absence of an excess of events above the expected backgrounds, limits are set within a generic R -hadron model, on stable gluino R -hadrons (see Table 5) and on metastable gluino R -hadrons decaying to $(g/q\bar{q})$ plus a light $\tilde{\chi}_1^0$ (see Fig. 7) and decaying to $t\bar{t}$ plus a light $\tilde{\chi}_1^0$ (see Fig. 9). NODE=S046LGN;LINKAGE=Q
- 17 KHACHATRYAN 15AK looked in a data set corresponding to 18.6 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$, and a search interval corresponding to 281 h of trigger lifetime, for long-lived particles that have stopped in the CMS detector. No evidence for an excess over the expected background in a cloud interaction model is observed. Assuming the decay $\tilde{g} \rightarrow g\tilde{\chi}_1^0$ and lifetimes between $1 \mu\text{s}$ and 1000 s , limits are derived on \tilde{g} production as a function of $m_{\tilde{\chi}_1^0}$, see Figs. 4 and 6. The exclusions require that $m_{\tilde{\chi}_1^0}$ is kinematically consistent with the minimum values of the jet energy thresholds used. NODE=S046LGN;LINKAGE=H

- 18 AAD 13AA searched in 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events containing colored long-lived particles that hadronize forming R -hadrons. No significant excess above the expected background was found. Long-lived R -hadrons containing a \tilde{g} are excluded for masses up to 985 GeV at 95% C.L. in a general interaction model. Also, limits independent of the fraction of R -hadrons that arrive charged in the muon system were derived, see Fig. 6. NODE=S046LGN;LINKAGE=DA
- 19 AAD 13BC searched in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and in 22.9 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for bottom squark R -hadrons that have come to rest within the ATLAS calorimeter and decay at some later time to hadronic jets and a neutralino. In absence of an excess of events above the expected backgrounds, limits are set on gluino masses for different decays, lifetimes, and neutralino masses, see their Table 6 and Fig. 10. NODE=S046LGN;LINKAGE=DD
- 20 CHATRCHYAN 13AB looked in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and in 18.8 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or additionally requiring that it be identified as muon in the muon chambers, from pair production of \tilde{g} 's. No evidence for an excess over the expected background is observed. Limits are derived for pair production of gluinos as a function of mass (see Fig. 8 and Table 5), depending on the fraction, f , of formation of $\tilde{g}-g$ (R -gluonball) states. The quoted limit is for $f = 0.1$, while for $f = 0.5$ it degrades to 1276 GeV. In the conservative scenario where every hadronic interaction causes it to become neutral, the limit decreases to 928 GeV for $f = 0.1$. NODE=S046LGN;LINKAGE=CH
- 21 AAD 12P looked in 31 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with pair production of long-lived gluinos. The hadronization of the gluinos leads to R -hadrons which may stop inside the detector and later decay via $\tilde{g} \rightarrow g \tilde{\chi}_1^0$ during gaps between the proton bunches. No significant excess over the expected background is observed. From a counting experiment, a limit at 95% C.L. on the cross section as a function of $m_{\tilde{g}}$ is derived for $m_{\tilde{\chi}_1^0} = 100 \text{ GeV}$, see Fig. 4. The limit is valid for lifetimes between 10^{-5} and 10^3 seconds and assumes the *Generic* matter interaction model for the production cross section. NODE=S046LGN;LINKAGE=GA
- 22 CHATRCHYAN 12AN looked in 4.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with pair production of long-lived gluinos. The hadronization of the gluinos leads to R -hadrons which may stop inside the detector and later decay via $\tilde{g} \rightarrow g \tilde{\chi}_1^0$ during gaps between the proton bunches. No significant excess over the expected background is observed. From a counting experiment, a limit at 95% C.L. on the cross section as a function of $m_{\tilde{g}}$ is derived, see Fig. 3. The mass limit is valid for lifetimes between 10^{-5} and 10^3 seconds, for what they call "the daughter gluon energy $E_g > 100 \text{ GeV}$ and assuming the *cloud* interaction model for R -hadrons. Supersedes KHACHATRYAN 11. NODE=S046LGN;LINKAGE=CT
- 23 CHATRCHYAN 12L looked in 5.0 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or additionally requiring that it be identified as muon in the muon chambers, from pair production of \tilde{g} 's. No evidence for an excess over the expected background is observed. Limits are derived for pair production of gluinos as a function of mass (see Fig. 3), depending on the fraction, f , of formation of $\tilde{g}-g$ (R -glueball) states. The quoted limit is for $f = 0.1$, while for $f = 0.5$ it degrades to 1046 GeV. In the conservative scenario where every hadronic interaction causes it to become neutral, the limit decreases to 928 GeV for $f=0.1$. Supersedes KHACHATRYAN 11C. NODE=S046LGN;LINKAGE=HR
- 24 AAD 11K looked in 34 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or time of flight in the tile calorimeter, from pair production of \tilde{g} . No evidence for an excess over the SM expectation is observed. Limits are derived for pair production of gluinos as a function of mass (see Fig. 4), for a fraction, $f = 10\%$, of formation of $\tilde{g}-g$ (R -gluonball). If instead of a phase space driven approach for the hadronic scattering of the R -hadrons, a triple-Regge model or a bag-model is used, the limit degrades to 566 and 562 GeV, respectively. NODE=S046LGN;LINKAGE=A2
- 25 AAD 11P looked in 37 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with heavy stable particles, reconstructed and identified by their time of flight in the Muon System. There is no requirement on their observation in the tracker to increase the sensitivity to cases where gluinos have a large fraction, f , of formation of neutral $\tilde{g}-g$ (R -gluonball). No evidence for an excess over the SM expectation is observed. Limits are derived as a function of mass (see Fig. 4), for $f=0.1$. For fractions $f = 0.5$ and 1.0 the limit degrades to 537 and 530 GeV, respectively. NODE=S046LGN;LINKAGE=A1
- 26 KHACHATRYAN 11 looked in 10 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with pair production of long-lived gluinos. The hadronization of the gluinos leads to R -hadrons which may stop inside the detector and later decay via $\tilde{g} \rightarrow g \tilde{\chi}_1^0$ during gaps between the proton bunches. No significant excess over the expected background is observed. From a counting experiment, a limit at 95% C.L. on the cross section times branching ratio is derived for $m_{\tilde{g}} - m_{\tilde{\chi}_1^0} > 100 \text{ GeV}$, see their Fig. 2. Assuming 100% branching ratio, lifetimes between 75 ns and $3 \times 10^5 \text{ s}$ are excluded for $m_{\tilde{g}} = 300 \text{ GeV}$. The \tilde{g} mass exclusion is obtained with the same assumptions for lifetimes between $10 \mu\text{s}$ and 1000 s , but shows some dependence on the model for R -hadron interactions with matter, illustrated in Fig. 3. From a time-profile analysis, the mass exclusion is 382 GeV for a lifetime of $10 \mu\text{s}$ under the same assumptions as above. NODE=S046LGN;LINKAGE=KH
- 27 KHACHATRYAN 11C looked in 3.1 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with heavy stable particles, identified by their anomalous dE/dx in the tracker or additionally

requiring that it be identified as muon in the muon chambers, from pair production of \tilde{g} . No evidence for an excess over the expected background is observed. Limits are derived for pair production of gluinos as a function of mass (see Fig. 3), depending on the fraction, f , of formation of $\tilde{g} - g$ (R-gluonball). The quoted limit is for $f=0.1$, while for $f=0.5$ it degrades to 357 GeV. In the conservative scenario where every hadronic interaction causes it to become neutral, the limit decreases to 311 GeV for $f=0.1$.

Light \tilde{G} (Gravitino) mass limits from collider experiments

The following are bounds on light ($\ll 1$ eV) gravitino indirectly inferred from its coupling to matter suppressed by the gravitino decay constant.

Unless otherwise stated, all limits assume that other supersymmetric particles besides the gravitino are too heavy to be produced. The gravitino is assumed to be undetected and to give rise to a missing energy (\cancel{E}) signature.

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$> 3.5 \times 10^{-4}$	95	¹ AAD	15BH ATLS	jet + \cancel{E}_T , $pp \rightarrow (\tilde{q}/\tilde{g})\tilde{G}$, $m_{\tilde{q}} = m_{\tilde{g}} = 500$ GeV
$> 3 \times 10^{-4}$	95	¹ AAD	15BH ATLS	jet + \cancel{E}_T , $pp \rightarrow (\tilde{q}/\tilde{g})\tilde{G}$, $m_{\tilde{q}} = m_{\tilde{g}} = 1000$ GeV
$> 2 \times 10^{-4}$	95	¹ AAD	15BH ATLS	jet + \cancel{E}_T , $pp \rightarrow (\tilde{q}/\tilde{g})\tilde{G}$, $m_{\tilde{q}} = m_{\tilde{g}} = 1500$ GeV
$> 1.09 \times 10^{-5}$	95	² ABDALLAH	05B DLPH	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
$> 1.35 \times 10^{-5}$	95	³ ACHARD	04E L3	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
$> 1.3 \times 10^{-5}$		⁴ HEISTER	03C ALEP	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
$> 11.7 \times 10^{-6}$	95	⁵ ACOSTA	02H CDF	$p\bar{p} \rightarrow \tilde{G}\tilde{G}\gamma$
$> 8.7 \times 10^{-6}$	95	⁶ ABBIENDI,G	00D OPAL	$e^+e^- \rightarrow \tilde{G}\tilde{G}\gamma$
¹ AAD 15BH searched in 20.3 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV for associated production of a light gravitino and a squark or gluino. The squark (gluino) is assumed to decay exclusively to a quark (gluon) and a gravitino. No evidence was found for an excess above the expected level of Standard Model background and 95% C.L. lower limits were set on the gravitino mass as a function of the squark/gluino mass, both in the case of degenerate and non-degenerate squark/gluino masses, see Figs. 14 and 15.				
² ABDALLAH 05B use data from $\sqrt{s} = 180\text{--}208$ GeV. They look for events with a single photon + \cancel{E} final states from which a cross section limit of $\sigma < 0.18 \text{ pb}$ at 208 GeV is obtained, allowing a limit on the mass to be set. Supersedes the results of ABREU 00Z.				
³ ACHARD 04E use data from $\sqrt{s} = 189\text{--}209$ GeV. They look for events with a single photon + \cancel{E} final states from which a limit on the Gravitino mass is set corresponding to $\sqrt{F} > 238$ GeV. Supersedes the results of ACCIARRI 99R.				
⁴ HEISTER 03C use the data from $\sqrt{s} = 189\text{--}209$ GeV to search for $\gamma\cancel{E}_T$ final states.				
⁵ ACOSTA 02H looked in 87 pb^{-1} of $p\bar{p}$ collisions at $\sqrt{s}=1.8$ TeV for events with a high- E_T photon and \cancel{E}_T . They compared the data with a GMSB model where the final state could arise from $q\bar{q} \rightarrow \tilde{G}\tilde{G}\gamma$. Since the cross section for this process scales as $1/ F ^4$, a limit at 95% CL is derived on $ F ^{1/2} > 221$ GeV. A model independent limit for the above topology is also given in the paper.				
⁶ ABBIENDI,G 00D searches for $\gamma\cancel{E}$ final states from $\sqrt{s}=189$ GeV.				

Supersymmetry miscellaneous results

Results that do not appear under other headings or that make nonminimal assumptions.

Some earlier papers are now obsolete and have been omitted. They were last listed in our PDG 14 edition: K. Olive, *et al.* (Particle Data Group), Chinese Physics **C38** 070001 (2014) (<http://pdg.lbl.gov>).

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		¹ AAD	20C ATLS	habemus MSSM, $m_{A-\tan\beta}$ plane
none 450–1400	95	² AAD	20L ATLS	heavy neutral Higgs bosons, hMSSM, $m_{A-\tan\beta}$ plane
> 65	95	³ AABOUD	16AF ATLS	selected ATLAS searches on EWK sector
none 0–2	95	⁴ AAD	16AG ATLS	dark photon, γ_d , in SUSY- and Higgs-portal models
		⁵ AAD	13P ATLS	dark γ , hidden valley

none 100–185	95	⁶ AALTONEN 12AB CDF hidden-valley Higgs ⁷ AAD 11AA ATLS scalar gluons ⁸ CHATRCHYAN 11E CMS $\mu\mu$ resonances ⁹ ABAZOV 10N D0 γ_D , hidden valley	
		¹ AAD 20C uses a statistical combination of six final states $b\bar{b}b\bar{b}$, $b\bar{b}WW$, $b\bar{b}\tau\tau$, $WWWW$, $b\bar{b}\gamma\gamma$, and $WW\gamma\gamma$ to search for non-resonant and resonant production of Higgs boson pairs. The search uses 36.1 fb^{-1} of pp collisions data at $\sqrt{s} = 13 \text{ TeV}$. Constraints in the habemus Minimal Supersymmetric Standard Model in the $(m_A, \tan\beta)$ parameter space are placed, see their Figure 7(b).	NODE=S046OTH;LINKAGE=G
		² AAD 20L used 27.8 fb^{-1} of pp collision data at $\sqrt{s} = 13 \text{ TeV}$ to search for heavy neutral Higgs bosons produced in association with at least one b -quark and decaying into a pair of b -quarks. The data are compatible with SM expectations, yielding no significant excess of events in the mass range 450–1400 GeV, see their Fig. 11. Exclusion limits at 95% C.L. were derived in hMSSM scenarios as a function of m_A and $\tan\beta$, see their Fig. 9 and 10.	NODE=S046OTH;LINKAGE=H
		³ AABOUD 16AF uses a selection of searches by ATLAS for the electroweak production of SUSY particles studying resulting constraints on dark matter candidates. They use 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$. A likelihood-driven scan of an effective model focusing on the gaugino-higgsino and Higgs sector of the pMSSM is performed. The ATLAS searches impact models where $m_{\chi_1^0} < 65 \text{ GeV}$, excluding 86% of them. See their Figs. 2, 4, and 6.	NODE=S046OTH;LINKAGE=D
		⁴ AAD 16AG searches for prompt lepton-jets using 20 fb^{-1} of pp collisions at $\sqrt{s} = 8 \text{ TeV}$ collected with the ATLAS detector. Lepton-jets are expected from decays of low-mass dark photons in SUSY-portal and Higgs-portal models. No significant excess of events is observed and 95% CL upper limits are computed on the production cross section times branching ratio for two prompt lepton-jets in models predicting 2 or 4 γ_D via SUSY-portal topologies, for γ_D mass values between 0 and 2 GeV. See their Figs 9 and 10. The results are also interpreted in terms of a 90% CL exclusion region in kinetic mixing and dark-photon mass parameter space. See their Fig. 13.	NODE=S046OTH;LINKAGE=F
		⁵ AAD 13P searched in 5 fb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for single lepton-jets with at least four muons; pairs of lepton-jets, each with two or more muons; and pairs of lepton-jets with two or more electrons. All of these could be signatures of Hidden Valley supersymmetric models. No statistically significant deviations from the Standard Model expectations are found. 95% C.L. limits are placed on the production cross section times branching ratio of dark photons for several parameter sets of a Hidden Valley model.	NODE=S046OTH;LINKAGE=DA
		⁶ AALTONEN 12AB looked in 5.1 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for anomalous production of multiple low-energy leptons in association with a W or Z boson. Such events may occur in hidden valley models in which a supersymmetric Higgs boson is produced in association with a W or Z boson, with $H \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0$ pair and with the $\tilde{\chi}_1^0$ further decaying into a dark photon (γ_D) and the unobservable lightest SUSY particle of the hidden sector. As the γ_D is expected to be light, it may decay into a lepton pair. No significant excess over the SM expectation is observed and a limit at 95% C.L. is set on the cross section for a benchmark model of supersymmetric hidden-valley Higgs production.	NODE=S046OTH;LINKAGE=AA
		⁷ AAD 11AA looked in 34 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with ≥ 4 jets originating from pair production of scalar gluons, each decaying to two gluons. No two-jet resonances are observed over the SM background. Limits are derived on the cross section times branching ratio (see Fig. 3). Assuming 100% branching ratio for the decay to two gluons, the quoted exclusion range is obtained, except for a 5 GeV mass window around 140 GeV.	NODE=S046OTH;LINKAGE=AD
		⁸ CHATRCHYAN 11E looked in 35 pb^{-1} of pp collisions at $\sqrt{s} = 7 \text{ TeV}$ for events with collimated μ pairs (leptonic jets) from the decay of hidden sector states. No evidence for new resonance production is found. Limits are derived and compared to various SUSY models (see Fig. 4) where the LSP, either the $\tilde{\chi}_1^0$ or a \tilde{q} , decays to dark sector particles.	NODE=S046OTH;LINKAGE=C2
		⁹ ABAZOV 10N looked in 5.8 fb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ for events from hidden valley models in which a $\tilde{\chi}_1^0$ decays into a dark photon, γ_D , and the unobservable lightest SUSY particle of the hidden sector. As the γ_D is expected to be light, it may decay into a tightly collimated lepton pair, called lepton jet. They searched for events with \cancel{E}_T and two isolated lepton jets observable by an opposite charged lepton pair $e\bar{e}$, $e\mu$ or $\mu\mu$. No significant excess over the SM expectation is observed, and a limit at 95% C.L. on the cross section times branching ratio is derived, see their Table I. They also examined the invariant mass of the lepton jets for a narrow resonance, see their Fig. 4, but found no evidence for a signal.	NODE=S046OTH;LINKAGE=AB

REFERENCES FOR Supersymmetric Particle Searches

AAD	23AB JHEP 2307 021	G. Aad <i>et al.</i>	(ATLAS Collab.)	NODE=S046
AAD	23AE EPJ C83 515	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62164
AAD	23AL EPJ C83 561	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62170
AAD	23AM PR D108 012012	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62182
AAD	23BQ PL B846 138172	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62295
AAD	23CB JHEP 2310 082	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62473
AAD	23CI JHEP 2312 167	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62540
AAD	23CP JHEP 2311 150	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62576
AAD	23CR JHEP 2312 081	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62611
AAD	23G JHEP 2306 158	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62616
AAD	23M JHEP 2306 031	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62120
				REFID=62134

AALBERS	23	PRL 131 041002	J. Aalbers <i>et al.</i>	(LZ Collab.)	REFID=62236
ABE	23B	PRL 130 061002	H. Abe <i>et al.</i>	(MAGIC Collab.)	REFID=62201
APRILE	23A	PRL 131 041003	E. Aprile <i>et al.</i>	(XENONnT Collab.)	REFID=62237
FOSTER	23	PR D107 103047	J.W. Foster <i>et al.</i>	(MIT, UCB, LBL+)	REFID=62332
GUO	23A	PR D108 043001	X.-K. Guo <i>et al.</i>		REFID=62359
HAYRAPETY...	23E	JHEP 2310 046	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62539
LEES	23C	PRL 131 201801	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=62465
TUMASYAN	23AB	JHEP 2307 110	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62194
TUMASYAN	23AG	PR D108 012011	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62294
TUMASYAN	23AO	JHEP 2307 210	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62372
TUMASYAN	23AY	JHEP 2309 149	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62531
TUMASYAN	23B	PL B842 137460	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62103
TUMASYAN	23H	JHEP 2305 227	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62114
TUMASYAN	23K	JHEP 2306 060	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62130
TUMASYAN	23L	JHEP 2307 161	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62139
TUMASYAN	23X	EPJ C83 571	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62179
AAD	22E	PL B830 137106	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61706
AAD	22U	EPJ C82 606	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61834
ABBASI	22B	PR D105 062004	R. Abbasi <i>et al.</i>	(IceCube Collab.)	REFID=61731
ABDALLA	22	PRL 129 111101	H. Abdalla <i>et al.</i>	(H.E.S.S. Collab.)	REFID=61913
HUANG	22	PL B834 137487	Z. Huang <i>et al.</i>	(PandaX-4T Collab.)	REFID=61935
TUMASYAN	22AF	EPJ C82 153	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61825
TUMASYAN	22Q	JHEP 2204 091	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61791
TUMASYAN	22S	JHEP 2204 147	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61793
TUMASYAN	22V	JHEP 2205 014	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61796
AAD	21AK	EPJ C81 600	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61341
AAD	21AL	PRL 127 051802	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61350
AAD	21AM	PR D104 032014	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61391
AAD	21AW	PR D104 112005	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61533
AAD	21AX	PR D104 112010	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61534
AAD	21B	EPJ C81 11	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61045
Also		EPJ C81 249 (errata.)	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61323
AAD	21BF	EPJ C81 1023	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61571
AAD	21BG	EPJ C81 1118	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61573
AAD	21E	PR D103 112003	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61245
AAD	21F	PR D103 112006	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61247
AAD	21L	JHEP 2102 143	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61282
AAD	21O	JHEP 2104 174	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61293
AAD	21P	JHEP 2104 165	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61295
AAD	21S	JHEP 2105 093	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61304
AAD	21X	JHEP 2107 173	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61318
AAD	21Y	JHEP 2107 167	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61319
AAJ	21V	EPJ C81 261	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61328
ABDALLAH	21	PR D103 102002	H. Abdallah <i>et al.</i>	(H.E.S.S. Collab.)	REFID=61240
MENG	21B	PRL 127 261802	Y. Meng <i>et al.</i>	(PandaX-4T Collab.)	REFID=61506
SIRUNYAN	21AD	PR D104 052001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=61513
SIRUNYAN	21AF	PR D104 052011	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=61518
SIRUNYAN	21B	EPJ C81 3	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60776
SIRUNYAN	21M	JHEP 2104 123	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=61299
SIRUNYAN	21U	PR D104 012015	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=61366
SIRUNYAN	21V	PR D104 032006	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=61376
TUMASYAN	21C	JHEP 2110 045	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61554
TUMASYAN	21I	EPJ C81 970	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61572
AABOUD	20	EPJ C80 754	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=60460
AAD	20AL	JHEP 2010 062	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60756
AAD	20AN	JHEP 2010 005	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60759
AAD	20AS	EPJ C80 1080	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60772
AAD	20C	PL B800 135103	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60147
AAD	20D	PL B801 135114	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60192
AAD	20H	PR D101 032009	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60257
AAD	20I	PR D101 052005	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60270
AAD	20K	PR D101 072001	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60288
AAD	20L	PR D102 032004	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60347
AAD	20M	PR D102 032006	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60348
AAD	20O	EPJ C80 123	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60414
AAD	20R	EPJ C80 691	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60448
AAD	20S	EPJ C80 737	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60456
AAD	20V	JHEP 2006 046	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60500
ABAZAJIAN	20	PR D102 043012	K.N. Abazajian <i>et al.</i>	(UCI, VPI, TOKY+)	REFID=60360
ABDALLAH	20	PR D102 062001	H. Abdallah <i>et al.</i>	(H.E.S.S. Collab.)	REFID=60721
ABE	20G	PR D102 072002	K. Abe <i>et al.</i>	(Super-Kamiokande Collab.)	REFID=60727
ALBERT	20	PR D101 103001	A. Albert <i>et al.</i>	(HAWC Collab.)	REFID=60362
ALBERT	20A	PL B805 135439	A. Albert <i>et al.</i>	(ANTARES Collab.)	REFID=60400
ALBERT	20C	PR D102 082002	A. Albert <i>et al.</i>	(ANTARES and IceCube Collab.)	REFID=60730
ALVAREZ	20	JCAP 2009 004	A. Alvarez <i>et al.</i>		REFID=60637
HOOF	20	JCAP 2002 012	S. Hoof, A. Geringer-Sameth, R. Trotta	(GOET+)	REFID=60365
SIRUNYAN	20AH	JHEP 2005 032	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60491
SIRUNYAN	20AU	PRL 124 041803	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60606
SIRUNYAN	20B	PL B801 135183	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60205
SIRUNYAN	20BJ	JHEP 2009 149	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60753
SIRUNYAN	20E	PR D101 052010	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60273
SIRUNYAN	20N	PL B806 135502	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60404
SIRUNYAN	20P	EPJ C80 189	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60418
SIRUNYAN	20T	EPJ C80 752	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60459
SIRUNYAN	20U	JHEP 2002 015	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60465
WANG	20G	CP C44 125001	Q. Wang <i>et al.</i>	(PandaX-II Collab.)	REFID=61260
AABOUD	19AT	PR D99 092007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59877
AABOUD	19AU	PR D100 012006	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59904
AABOUD	19C	PL B788 96	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59415
AABOUD	19G	PR D99 012001	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59518
AABOUD	19I	PR D99 012009	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59601
AAD	19H	JHEP 1912 060	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60097
ABE	19	PL B789 45	K. Abe <i>et al.</i>	(XMASS Collab.)	REFID=59423
AJAJ	19	PR D100 022004	R. Ajaj <i>et al.</i>	(DEAP-3600 Collab.)	REFID=59954
AMOLE	19	PR D100 022001	C. Amole <i>et al.</i>	(PICO Collab.)	REFID=59953
APRILE	19A	PRL 122 141301	E. Aprile <i>et al.</i>	(XENON1T Collab.)	REFID=59772
DI-MAURO	19	PR D99 123027	M. Di Mauro <i>et al.</i>		REFID=59949
JOHNSON	19	PR D99 103007	C. Johnson <i>et al.</i>		REFID=59946
LI	19D	PR D99 123519	S. Li <i>et al.</i>		REFID=59951

SIRUNYAN	19AG	JHEP 1906 143	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59694
SIRUNYAN	19AO	EPJ C79 305	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59719
SIRUNYAN	19AU	EPJ C79 444	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59727
SIRUNYAN	19AW	PL B790 140	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59742
SIRUNYAN	19BH	PR D99 032011	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59832
SIRUNYAN	19BI	PR D99 032014	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59834
SIRUNYAN	19BJ	PR D99 052002	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59839
SIRUNYAN	19BT	PL B797 134876	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59936
SIRUNYAN	19BU	JHEP 1908 150	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59938
SIRUNYAN	19CA	PR D100 112003	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60039
SIRUNYAN	19CE	PRL 123 241801	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60075
SIRUNYAN	19CH	JHEP 1910 244	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60091
SIRUNYAN	19CI	JHEP 1911 109	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60094
SIRUNYAN	19F	PR D99 012010	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59602
SIRUNYAN	19K	JHEP 1901 154	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59633
SIRUNYAN	19S	JHEP 1903 031	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59656
SIRUNYAN	19U	JHEP 1903 101	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59658
XIA	19A	PL B792 193	J. Xia <i>et al.</i>	(PandaX-II Collab.)	REFID=59926
AABOUD	18AQ	JHEP 1806 108	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59113
AABOUD	18AR	JHEP 1806 107	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59114
AABOUD	18AS	JHEP 1806 022	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59122
AABOUD	18AY	EPJ C78 154	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59153
AABOUD	18BB	EPJ C78 250	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59163
AABOUD	18BJ	EPJ C78 625	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59196
AABOUD	18BT	EPJ C78 995	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59331
AABOUD	18BV	JHEP 1809 050	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59338
AABOUD	18CF	PL B785 136	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59402
AABOUD	18CK	PR D98 092002	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59470
AABOUD	18CM	PR D98 092008	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59474
AABOUD	18CO	PR D98 092012	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59478
AABOUD	18I	JHEP 1801 126	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58792
AABOUD	18P	PR D97 032003	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58878
AABOUD	18R	PR D97 052010	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58902
AABOUD	18S	PR D97 052012	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58903
AABOUD	18U	PR D97 092006	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58939
AABOUD	18V	PR D97 112001	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58947
AABOUD	18Y	PR D98 032008	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58976
AABOUD	18Z	PR D98 032009	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58977
ABDALLAH	18	PRL 120 201101	H. Abdallah <i>et al.</i>	(H.E.S.S. Collab.)	REFID=58824
ADHIKARI	18	NAT 564 83	G. Adhikari <i>et al.</i>	(COSINE-100 Collab.)	REFID=59583
AGNES	18A	PR D98 102006	P. Agnes <i>et al.</i>	(DarkSide-50 Collab.)	REFID=59489
AGNESE	18A	PRL 120 061802	R. Agnese <i>et al.</i>	(SuperCDMS Collab.)	REFID=58728
AHNEN	18	JCAP 1803 009	M.L. Ahnen <i>et al.</i>	(MAGIC Collab.)	REFID=59058
ALBERT	18B	JCAP 1806 043	A. Albert <i>et al.</i>	(HAWC Collab.)	REFID=59062
ALBERT	18C	PR D98 123012	A. Albert <i>et al.</i>	(HAWC Collab.)	REFID=59510
AMAUDRUZ	18	PRL 121 071801	P.A. Amaudruz <i>et al.</i>	(DEAP-3600 Collab.)	REFID=58862
APRILE	18	PRL 121 111302	E. Aprile <i>et al.</i>	(XENON1T Collab.)	REFID=59288
SIRUNYAN	18AA	PL B780 118	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59005
SIRUNYAN	18AC	PL B780 384	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59008
SIRUNYAN	18AD	PL B780 432	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59009
SIRUNYAN	18AJ	PL B782 440	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59026
SIRUNYAN	18AK	PL B783 114	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59032
SIRUNYAN	18AL	JHEP 1802 067	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59076
SIRUNYAN	18AN	JHEP 1803 167	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59082
SIRUNYAN	18AO	JHEP 1803 166	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59083
SIRUNYAN	18AP	JHEP 1803 160	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59084
SIRUNYAN	18AR	JHEP 1803 076	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59089
SIRUNYAN	18AT	JHEP 1804 073	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59095
SIRUNYAN	18AY	JHEP 1805 025	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59106
SIRUNYAN	18B	PL B778 263	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58628
SIRUNYAN	18BR	JHEP 1808 016	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59147
SIRUNYAN	18C	PR D97 032009	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58718
SIRUNYAN	18D	PR D97 012007	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58719
SIRUNYAN	18DI	JHEP 1809 065	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59340
SIRUNYAN	18DN	JHEP 1811 079	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59362
SIRUNYAN	18DP	JHEP 1811 151	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59365
SIRUNYAN	18DV	PR D98 092011	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59477
SIRUNYAN	18DY	PR D98 112014	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59504
SIRUNYAN	18EA	PRL 121 141802	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59524
SIRUNYAN	18M	PRL 120 241801	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58841
SIRUNYAN	18O	PR D97 032007	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58880
SIRUNYAN	18X	PL B779 166	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59000
AABOUD	17AF	JHEP 1708 006	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58126
AABOUD	17AI	JHEP 1709 088	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58208
AABOUD	17AJ	JHEP 1709 084	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58209
Also		JHEP 1908 121 (errat.)	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59939
AABOUD	17AR	PR D96 112010	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58323
AABOUD	17AX	JHEP 1711 195	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58341
AABOUD	17AY	JHEP 1712 085	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58350
AABOUD	17AZ	JHEP 1712 034	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58351
AABOUD	17BE	EPJ C77 898	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58362
AABOUD	17N	EPJ C77 144	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57843
AAIJ	17Z	EPJ C77 224	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57852
AARTSEN	17	EPJ C77 82	M.G. Aartsen <i>et al.</i>	(IceCube Collab.)	REFID=57835
AARTSEN	17A	EPJ C77 146	M.G. Aartsen <i>et al.</i>	(IceCube Collab.)	REFID=57844
Also		EPJ C79 214 (errat.)	M.G. Aartsen <i>et al.</i>	(IceCube Collab.)	REFID=59712
AARTSEN	17C	EPJ C77 627	M.G. Aartsen <i>et al.</i>	(IceCube Collab.)	REFID=58190
AKERIB	17	PRL 118 021303	D.S. Akerib <i>et al.</i>	(LUX Collab.)	REFID=57652
AKERIB	17A	PRL 118 251302	D.S. Akerib <i>et al.</i>	(LUX Collab.)	REFID=57936
AMOLE	17	PRL 118 251301	C. Amole <i>et al.</i>	(PICO Collab.)	REFID=57935
APRILE	17G	PRL 119 181301	E. Aprile <i>et al.</i>	(XENON Collab.)	REFID=58269
ARCHAMBAU...17		PR D95 082001	S. Archambault <i>et al.</i>	(VERITAS Collab.)	REFID=57985
ATHRON	17B	EPJ C77 824	P. Athron <i>et al.</i>	(GAMBIT Collab.)	REFID=60656
BATTAT	17	ASP 91 65	J.B.R. Battat <i>et al.</i>	(DRIFT-IIld Collab.)	REFID=58037
BEHNKE	17	ASP 90 85	E. Behnke <i>et al.</i>	(PICASSO Collab.)	REFID=58036
CUI	17A	PRL 119 181302	X. Cui <i>et al.</i>	(PandaX-II Collab.)	REFID=58270
FU	17	PRL 118 071301	C. Fu <i>et al.</i>	(PandaX-II Collab.)	REFID=57912
Also		PRL 120 049902 (errat.)	C. Fu <i>et al.</i>	(PandaX-II Collab.)	REFID=59198
KHACHATRY...17		PR D95 012003	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57637

KHACHATRY... 17A	PRL 118 021802	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57654
KHACHATRY... 17AD	PR D96 012004	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=58012
KHACHATRY... 17AR	PR D95 012009	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=58104
KHACHATRY... 17AS	PR D95 012011	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=58106
KHACHATRY... 17AW	EPJ C77 635	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=58189
KHACHATRY... 17L	JHEP 1704 018	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57804
KHACHATRY... 17P	EPJ C77 294	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57856
KHACHATRY... 17S	PL B767 403	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57886
KHACHATRY... 17V	PL B769 391	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57898
KHACHATRY... 17Y	PL B770 257	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57904
SIRUNYAN 17AF	PRL 119 151802	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58264
SIRUNYAN 17AS	JHEP 1710 019	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58339
SIRUNYAN 17AT	JHEP 1710 005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58340
SIRUNYAN 17AW	JHEP 1711 029	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58346
SIRUNYAN 17AY	JHEP 1712 142	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58349
SIRUNYAN 17AZ	EPJ C77 710	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58356
SIRUNYAN 17K	EPJ C77 327	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=57860
SIRUNYAN 17P	PR D96 032003	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58028
SIRUNYAN 17S	EPJ C77 578	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58192
AABOUD 16AC	EPJ C76 683	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57687
AABOUD 16AF	JHEP 1609 175	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57730
AABOUD 16B	PL B760 647	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57245
AABOUD 16C	PR D93 112015	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57298
AABOUD 16D	PR D94 032005	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57310
AABOUD 16J	PR D94 052009	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57430
AABOUD 16M	EPJ C76 517	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57470
AABOUD 16N	EPJ C76 392	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57379
AABOUD 16P	EPJ C76 541	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57558
AABOUD 16Q	EPJ C76 547	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57559
AAD 16AA	PR D93 052002	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57264
AAD 16AD	PR D94 032003	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57308
AAD 16AG	JHEP 1602 062	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57317
AAD 16AM	JHEP 1606 067	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57336
AAD 16AY	EPJ C76 81	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57362
AAD 16BB	EPJ C76 259	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57371
AAD 16BG	EPJ C76 565	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57563
AAD 16V	PL B757 334	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57218
AARTSEN 16C	JCAP 1604 022	M.G. Aartsen <i>et al.</i>	(IceCube Collab.)	REFID=57476
ADRIAN-MAR...16	PL B759 69	S. Adrian-Martinez <i>et al.</i>	(ANTARES Collab.)	REFID=57232
AHNEN 16	JCAP 1602 039	M.L. Ahnen <i>et al.</i>	(MAGIC and Fermi-LAT Collab.)	REFID=57473
AKERIB 16	PRL 116 161301	D.S. Akerib <i>et al.</i>	(LUX Collab.)	REFID=57185
AKERIB 16A	PRL 116 161302	D.S. Akerib <i>et al.</i>	(LUX Collab.)	REFID=57186
AMOLE 16	PR D93 052014	C. Amole <i>et al.</i>	(PICO Collab.)	REFID=57270
APRILE 16B	PR D94 122001	E. Aprile <i>et al.</i>	(XENON100 Collab.)	REFID=57629
AVRORIN 16	ASP 81 12	A.D. Avrorin <i>et al.</i>	(BAIKAL Collab.)	REFID=57471
BECHTLE 16	EPJ C76 96	P. Bechtle <i>et al.</i>		REFID=60651
CIRELLI 16	JCAP 1607 041	M. Cirelli, M. Taoso	(LPNHE, MADE)	REFID=57579
KHACHATRY... 16AA	PL B759 479	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57237
KHACHATRY... 16AC	PL B760 178	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57242
KHACHATRY... 16AM	PR D93 092009	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57285
KHACHATRY... 16AV	JHEP 1607 027	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57341
KHACHATRY... 16AY	JHEP 1608 122	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57348
KHACHATRY... 16BE	EPJ C76 317	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57373
KHACHATRY... 16BJ	EPJ C76 439	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57382
KHACHATRY... 16BK	EPJ C76 460	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57384
KHACHATRY... 16BS	JHEP 1610 006	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57552
KHACHATRY... 16BT	JHEP 1610 129	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57554
KHACHATRY... 16BW	PR D94 112004	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57620
KHACHATRY... 16BX	PR D94 112009	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57622
KHACHATRY... 16BY	JHEP 1612 013	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57662
KHACHATRY... 16R	PL B757 6	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57214
KHACHATRY... 16V	PL B758 152	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57226
KHACHATRY... 16Y	PL B759 9	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57230
LEITE 16	JCAP 1611 021	N. Leite <i>et al.</i>		REFID=57743
AAD 15AB	PR D92 012010	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56600
AAD 15AE	JHEP 1501 068	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56616
AAD 15AI	JHEP 1504 116	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56620
AAD 15BA	EPJ C75 208	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56668
AAD 15BG	EPJ C75 318	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56674
Also	EPJ C75 463	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56971
AAD 15BH	EPJ C75 299	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56675
Also	EPJ C75 408 (err.)	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57177
AAD 15BM	EPJ C75 407	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56680
AAD 15BV	JHEP 1510 054	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56858
AAD 15BX	JHEP 1510 134	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56861
AAD 15CA	PR D92 072001	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56889
AAD 15CB	PR D92 072004	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56891
AAD 15CJ	EPJ C75 510	G. Aad	(ATLAS Collab.)	REFID=56938
AAD 15CS	PR D91 012008	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57143
Also	PR D92 059903 (err.)	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56964
AAD 15J	PRL 114 142001	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56466
AAD 15K	PRL 114 161801	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56469
AAD 15O	PRL 115 031801	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56481
AAD 15X	PR D91 112016	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56590
AAJ 15BD	EPJ C75 595	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57032
AARTSEN 15E	EPJ C75 492	M.G. Aartsen <i>et al.</i>	(IceCube Collab.)	REFID=56937
ACKERMANN 15	PR D91 122002	M. Ackermann <i>et al.</i>	(Fermi-LAT Collab.)	REFID=56595
ACKERMANN 15A	JCAP 1509 008	M. Ackermann <i>et al.</i>	(Fermi-LAT Collab.)	REFID=56845
ACKERMANN 15B	PRL 115 231301	M. Ackermann <i>et al.</i>	(Fermi-LAT Collab.)	REFID=57106
AGNES 15	PL B743 456	P. Agnes <i>et al.</i>	(DarkSide-50 Collab.)	REFID=56502
AGNESE 15B	PR D92 072003	R. Agnese <i>et al.</i>	(SuperCDMS Collab.)	REFID=56890
BAGNASCHI 15	EPJ C75 500	E.A. Bagnaschi <i>et al.</i>		REFID=60650
BUCKLEY 15	PR D91 102001	M.R. Buckley <i>et al.</i>		REFID=56580
CHOI 15	PRL 114 141301	K. Choi <i>et al.</i>	(Super-Kamiokande Collab.)	REFID=56455
KHACHATRY... 15AB	JHEP 1501 096	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56541
KHACHATRY... 15AD	JHEP 1504 124	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56543
KHACHATRY... 15AF	JHEP 1505 078	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56633
KHACHATRY... 15AH	JHEP 1506 116	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56635
KHACHATRY... 15AK	EPJ C75 151	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56655

KHACHATRY... 15AO	EPJ C75 325	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56659
KHACHATRY... 15AR	PL B743 503	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56751
KHACHATRY... 15AZ	PR D92 072006	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56893
KHACHATRY... 15E	PRL 114 061801	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56439
KHACHATRY... 15I	PL B745 5	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56507
KHACHATRY... 15L	PL B747 98	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56516
KHACHATRY... 15O	PL B748 255	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56528
KHACHATRY... 15W	PR D91 052012	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56536
KHACHATRY... 15X	PR D91 052018	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56537
ROLBIECKI 15	PL B750 247	K. Rolbiecki, J. Tattersall	(MADE, HEID)	REFID=56958
AAD 14AE	JHEP 1409 176	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56003
AAD 14AG	JHEP 1409 103	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56005
AAD 14AJ	JHEP 1409 015	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56008
AAD 14AV	JHEP 1410 096	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56166
AAD 14AX	JHEP 1410 024	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56170
AAD 14B	EPJ C74 2883	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55701
AAD 14BD	JHEP 1411 118	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56292
AAD 14BH	PR D90 112005	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56331
AAD 14E	JHEP 1406 035	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55717
AAD 14F	JHEP 1406 124	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55718
AAD 14G	JHEP 1405 071	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55719
AAD 14H	JHEP 1404 169	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55720
AAD 14K	PR D90 012004	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55782
AAD 14T	PR D90 052008	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55882
AAD 14X	PR D90 052001	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55886
AALTONEN 14	PR D90 012011	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=55802
ACKERMANN 14	PR D89 042001	M. Ackermann <i>et al.</i>	(Fermi-LAT Collab.)	REFID=55796
AKERIB 14	PRL 112 091303	D.S. Akerib <i>et al.</i>	(LUX Collab.)	REFID=55832
ALEKSIC 14	JCAP 1402 008	J. Aleksic <i>et al.</i>	(MAGIC Collab.)	REFID=55964
AVRORIN 14	ASP 62 12	A.D. Avrorin <i>et al.</i>	(BAIKAL Collab.)	REFID=56193
BUCHMUEL... 14	EPJ C74 2809	O. Buchmueller <i>et al.</i>		REFID=56414
BUCHMUEL... 14A	EPJ C74 2922	O. Buchmueller <i>et al.</i>		REFID=56415
CHATRCHYAN 14AH	PR D90 112001	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=56327
CHATRCHYAN 14H	JHEP 1401 163	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55714
CHATRCHYAN 14I	JHEP 1406 055	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55731
CHATRCHYAN 14N	PL B733 328	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55754
CHATRCHYAN 14P	PL B730 193	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55756
CHATRCHYAN 14R	PR D90 032006	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55825
CHATRCHYAN 14U	PRL 112 161802	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55840
CZAKON 14	PR 113 201803	M. Czakon <i>et al.</i>	(AACH, CAMB, UCB, LBL+)	REFID=56120
FELIZARDO 14	PR D89 072013	M. Felizardo <i>et al.</i>	(SIMPLE Collab.)	REFID=56028
KHACHATRY... 14C	PL B736 371	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=55757
KHACHATRY... 14I	EPJ C74 3036	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56099
KHACHATRY... 14L	PR D90 092007	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56155
KHACHATRY... 14T	PL B739 229	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56335
PDG 14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ROSZKOWSKI 14	JHEP 1408 067	L. Roszkowski, E.M. Sessolo, A.J. Williams	(WINR)	REFID=56413
AAD 13	PL B718 841	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54788
AAD 13AA	PL B720 277	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54967
AAD 13AI	PL B723 15	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55080
AAD 13AP	PR D88 012001	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55157
AAD 13AU	JHEP 1310 189	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55422
AAD 13B	PL B718 879	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54790
AAD 13BC	PR D88 112003	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55607
AAD 13BD	PR D88 112006	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55609
AAD 13H	JHEP 1301 131	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54908
AAD 13L	PR D87 012008	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54930
AAD 13P	PL B719 299	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54934
AAD 13Q	PL B719 261	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54935
AAD 13R	PL B719 280	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54936
AALTONEN 13I	PR D88 031103	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=55173
AALTONEN 13Q	PRL 110 201802	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=55213
AARTSEN 13C	PR D88 122001	M.G. Aartsen <i>et al.</i>	(IceCube Collab.)	REFID=55611
ABAZOV 13B	PR D87 052011	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=55130
ACKERMANN 13A	PR D88 082002	M. Ackermann <i>et al.</i>	(Fermi-LAT Collab.)	REFID=55466
ADRIAN-MAR...13	JCAP 1311 032	S. Adrian-Martinez <i>et al.</i>	(ANTARES Collab.)	REFID=55597
AGNESE 13	PR D88 031104	R. Agnese <i>et al.</i>	(CDMS Collab.)	REFID=55174
AGNESE 13A	PRL 111 251301	R. Agnese <i>et al.</i>	(CDMS Collab.)	REFID=55615
APRILE 13	PRL 111 021301	E. Aprile <i>et al.</i>	(XENON100 Collab.)	REFID=55218
BERGSTROM 13	PRL 111 171101	L. Bergstrom <i>et al.</i>		REFID=55469
BOLIEV 13	JCAP 1309 019	M. Boliev <i>et al.</i>		REFID=55514
CABRERA 13	JHEP 1307 182	M. Cabrera, J. Casas, R. de Austri		REFID=55046
CALIBBI 13	JHEP 1310 132	L. Calibbi <i>et al.</i>		REFID=60246
CHATRCHYAN 13	PL B718 815	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54769
CHATRCHYAN 13AB	JHEP 1307 122	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55042
Also	JHEP 2211 149 (errata.)	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=61972
CHATRCHYAN 13AH	PL B722 273	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55075
CHATRCHYAN 13AO	PR D87 072001	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55138
CHATRCHYAN 13AT	PR D88 052017	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55194
CHATRCHYAN 13AV	PRL 111 081802	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55222
CHATRCHYAN 13G	JHEP 1301 077	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54896
CHATRCHYAN 13H	PL B719 42	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54940
CHATRCHYAN 13T	EPJ C73 2568	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55016
CHATRCHYAN 13V	JHEP 1303 037	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55027
Also	JHEP 1307 041 (errata.)	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55039
CHATRCHYAN 13W	JHEP 1303 111	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55028
ELLIS 13B	EPJ C73 2403	J. Ellis <i>et al.</i>		REFID=55672
JIN 13	JCAP 1311 026	H.-B. Jin, Y.-L. Wu, Y.-F. Zhou		REFID=55596
KOPP 13	PR D88 076013	J. Kopp		REFID=55552
STREGE 13	JCAP 1304 013	C. Streve <i>et al.</i>		REFID=55663
AAD 12AF	PL B714 180	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54195
AAD 12AG	PL B714 197	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54196

AAD	12AN	PRL 108 181802	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54223
AAD	12AS	PRL 108 261804	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54228
AAD	12AX	PR D85 012006	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54353
Also		PR D87 099903 (err.)	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55154
AAD	12BJ	EPJ C72 1993	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54521
AAD	12CJ	PR D86 092002	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54698
AAD	12CM	EPJ C72 2215	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54720
AAD	12CP	PL B718 411	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54725
AAD	12CT	JHEP 1212 124	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54783
AAD	12P	EPJ C72 1965	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54132
AAD	12R	PL B707 478	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54150
AAD	12T	PL B709 137	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54154
AAD	12W	PL B710 67	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54157
AALTONEN	12AB	PR D85 092001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=54367
ABAZOV	12AD	PR D86 071701	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=54606
AKIMOV	12	PL B709 14	D.Yu. Akimov <i>et al.</i>	(ZEPLIN-III Collab.)	REFID=54153
AKULA	12	PR D85 075001	S. Akula <i>et al.</i>	(NEAS, MICH)	REFID=54983
ANGLOHER	12	EPJ C72 1971	G. Angloher <i>et al.</i>	(CRESST-II Collab.)	REFID=54137
APRILE	12	PRL 109 181301	E. Aprile <i>et al.</i>	(XENON100 Collab.)	REFID=54616
ARBHEY	12A	PL B708 162	A. Arbey <i>et al.</i>		REFID=54982
ARCHAMBAU...	12	PL B711 153	S. Archambault <i>et al.</i>	(PICASSO Collab.)	REFID=54166
BAER	12	JHEP 1205 091	H. Baer, V. Barger, A. Mustafayev	(OKLA, WISC+)	REFID=54977
BALAZS	12	EPJ C73 2563	C. Balazs <i>et al.</i>		REFID=55665
BECHTLE	12	JHEP 1206 098	P. Bechtle <i>et al.</i>		REFID=54577
BEHNKE	12	PR D86 052001	E. Behnke <i>et al.</i>	(COUPP Collab.)	REFID=54610
Also		PR D90 079902 (err.)	E. Behnke <i>et al.</i>	(COUPP Collab.)	REFID=56151
BESKIDT	12	EPJ C72 2166	C. Beskidt <i>et al.</i>	(KARLE, JINR, ITEP)	REFID=54678
BOTTINO	12	PR D85 095013	A. Bottino, N. Fornengo, S. Scopel	(TORI, SOGA)	REFID=54988
BUCHMUEL...	12	EPJ C72 2020	O. Buchmueller <i>et al.</i>		REFID=54978
CAO	12A	PL B710 665	J. Cao <i>et al.</i>		REFID=54542
CHATRCHYAN	12	PR D85 012004	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=53990
CHATRCHYAN	12AE	PRL 109 171803	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54617
CHATRCHYAN	12AI	JHEP 1208 110	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54560
CHATRCHYAN	12AL	JHEP 1206 169	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54569
CHATRCHYAN	12AN	JHEP 1208 026	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54570
CHATRCHYAN	12AT	JHEP 1210 018	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54623
CHATRCHYAN	12BJ	JHEP 1211 147	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54774
CHATRCHYAN	12BK	JHEP 1211 172	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54775
CHATRCHYAN	12BO	JHEP 1212 055	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54778
CHATRCHYAN	12L	PL B713 408	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54179
DAW	12	ASP 35 397	E. Daw <i>et al.</i>	(DRIFT-II-d Collab.)	REFID=54431
DREINER	12A	EPL 99 61001	H.K. Dreiner, M. Kramer, J. Tattersall	(BONN+)	REFID=54642
ELLIS	12B	EPJ C72 2005	J. Ellis, K. Olive		REFID=54981
FELIZARDO	12	PRL 108 201302	M. Felizardo <i>et al.</i>	(SIMPLE Collab.)	REFID=54236
FENG	12B	PR D85 075007	J. Feng, K. Matchev, D. Sanford		REFID=55664
KADASTIK	12	JHEP 1205 061	M. Kadastik <i>et al.</i>		REFID=54984
KIM	12	PRL 108 181301	S.C. Kim <i>et al.</i>	(KIMS Collab.)	REFID=54232
STREGE	12	JCAP 1203 030	C. Strege <i>et al.</i>	(LOIC, AMST, MADU, GRAN+)	REFID=54446
AAD	11AA	EPJ C71 1828	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=53942
AAD	11G	PRL 106 131802	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=16561
AAD	11H	PRL 106 251801	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=16563
AAD	11K	PL B701 1	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=16569
AAD	11O	PL B701 398	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=16577
AAD	11P	PL B703 428	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=53783
AAD	11Z	EPJ C71 1809	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=53941
AHMED	11A	PR D84 011102	Z. Ahmed <i>et al.</i>	(CDMS and EDELWEISS Collabs.)	REFID=16611
ARMENGAUD	11	PL B702 329	E. Armengaud <i>et al.</i>	(EDELWEISS-II Collab.)	REFID=16711
BUCHMUEL...	11	EPJ C71 1583	O. Buchmueller <i>et al.</i>		REFID=54109
BUCHMUEL...	11B	EPJ C71 1722	O. Buchmueller <i>et al.</i>		REFID=54111
CHATRCHYAN	11B	JHEP 1106 093	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=16349
CHATRCHYAN	11D	JHEP 1107 113	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=16353
CHATRCHYAN	11E	JHEP 1107 098	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=16355
CHATRCHYAN	11V	PL B704 411	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=53846
KHACHATRY...	11	PRL 106 011801	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=53577
KHACHATRY...	11C	JHEP 1103 024	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=16343
ROSZKOWSKI	11	PR D83 015014	L. Roszkowski <i>et al.</i>		REFID=54105
AALTONEN	10	PRL 104 011801	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53188
AALTONEN	10R	PRL 105 081802	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53367
AALTONEN	10Z	PRL 105 191801	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53439
ABAZOV	10L	PL B693 95	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=53425
ABAZOV	10M	PRL 105 191802	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=53440
ABAZOV	10N	PRL 105 211802	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=53441
ABAZOV	10P	PRL 105 221802	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=53443
ACKERMANN	10	JCAP 1005 025	M. Ackermann	(Fermi-LAT Collab.)	REFID=53379
ARMENGAUD	10	PL B687 294	E. Armengaud <i>et al.</i>	(EDELWEISS-II Collab.)	REFID=53281
ELLIS	10	EPJ C69 201	J. Ellis, A. Mustafayev, K. Olive		REFID=54107
ABAZOV	09M	PRL 102 161802	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52866
AHMED	09	PRL 102 011301	Z. Ahmed <i>et al.</i>	(CDMS Collab.)	REFID=52613
ANGLOHER	09	ASP 31 270	G. Angloher <i>et al.</i>	(CRESST Collab.)	REFID=52769
BUCHMUEL...	09	EPJ C64 391	O. Buchmueller <i>et al.</i>	(LOIC, FNAL, CERN+)	REFID=53250
DREINER	09	EPJ C62 547	H. Dreiner <i>et al.</i>		REFID=52909
LEBEDENKO	09	PR D80 052010	V.N. Lebedenko <i>et al.</i>	(ZEPLIN-III Collab.)	REFID=53037
LEBEDENKO	09A	PRL 103 151302	V.N. Lebedenko <i>et al.</i>	(ZEPLIN-III Collab.)	REFID=53093
SORENSEN	09	NIM A601 339	P. Sorensen <i>et al.</i>	(XENON10 Collab.)	REFID=53257
ABAZOV	08F	PL B659 856	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52239
ANGLE	08	PRL 100 021303	J. Angle <i>et al.</i>	(XENON10 Collab.)	REFID=52246
ANGLE	08A	PRL 101 091301	J. Angle <i>et al.</i>	(XENON10 Collab.)	REFID=52446
BEDNYAKOV	08	PAN 71 111	V.A. Bednyakov, H.P. Klapdor-Kleingrothaus, I.V. Krivosheina		REFID=52471;ERROR=1

Translated from YAF 71 112.

BEHNKE	08	SCI 319 933	E. Behnke	(COUPP Collab.)	REFID=53249
BENETTI	08	ASP 28 495	P. Benetti <i>et al.</i>	(WARP Collab.)	REFID=52310
BUCHMUEL...	08	JHEP 0809 117	O. Buchmueller <i>et al.</i>		REFID=52516
ELLIS	08	PR D78 075012	J. Ellis, K. Olive, P. Sandick	(CERN, MINN)	REFID=52709
ABULENCIA	07H	PRL 98 131804	A. Abulencia <i>et al.</i>	(CDF Collab.)	REFID=51748
ALNER	07A	ASP 28 287	G.J. Alner <i>et al.</i>	(ZEPLIN-II Collab.)	REFID=51961
CALIBBI	07	JHEP 0709 081	L. Calibbi <i>et al.</i>		REFID=54108
ELLIS	07	JHEP 0706 079	J. Ellis, K. Olive, P. Sandick	(CERN, MINN)	REFID=52708
LEE	07A	PRL 99 091301	H.S. Lee <i>et al.</i>	(KIMS Collab.)	REFID=51919
ABBIENDI	06B	EPJ C46 307	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=51388
ACHTERBERG	06	ASP 26 129	A. Achterberg <i>et al.</i>	(AMANDA Collab.)	REFID=51396
ACKERMANN	06	ASP 24 459	M. Ackermann <i>et al.</i>	(AMANDA Collab.)	REFID=51392
AKERIB	06	PR D73 011102	D.S. Akerib <i>et al.</i>	(CDMS Collab.)	REFID=51018
AKERIB	06A	PRL 96 011302	D.S. Akerib <i>et al.</i>	(CDMS Collab.)	REFID=51056
ALLANACH	06	PR D73 015013	B.C. Allanach <i>et al.</i>		REFID=54102
BENOIT	06	PL B637 156	A. Benoit <i>et al.</i>		REFID=51256
DE-AUSTRI	06	JHEP 0605 002	R.R. de Austri, R. Trotta, L. Roszkowski		REFID=54104
DEBOER	06	PL B636 13	W. de Boer <i>et al.</i>		REFID=51368
LEP-SLC	06	PRPL 427 257	ALEPH, DELPHI, L3, OPAL, SLD and working groups		REFID=51219; ERROR=2; ERROR=3
SHIMIZU	06A	PL B633 195	Y. Shimizu <i>et al.</i>		REFID=51200
SMITH	06	PL B642 567	N.J.T. Smith, A.S. Murphy, T.J. Sumner		REFID=51492
ABAZOV	05A	PRL 94 041801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=50489
ABDALLAH	05B	EPJ C38 395	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=50448
AKERIB	05	PR D72 052009	D.S. Akerib <i>et al.</i>	(CDMS Collab.)	REFID=50906
ALNER	05	PL B616 17	G.J. Alner <i>et al.</i>	(UK Dark Matter Collab.)	REFID=50608
ALNER	05A	ASP 23 444	G.J. Alner <i>et al.</i>	(UK Dark Matter Collab.)	REFID=50712
BAER	05	JHEP 0507 065	H. Baer <i>et al.</i>	(FSU, MSU, HAWA)	REFID=52707
BARNABE-HE...	05	PL B624 186	M. Barnabe-Heider <i>et al.</i>	(PICASSO Collab.)	REFID=50874
ELLIS	05	PR D71 095007	J. Ellis <i>et al.</i>		REFID=50727
SANGLARD	05	PR D71 122002	V. Sanglard <i>et al.</i>	(EDELWEISS Collab.)	REFID=50668
ABBIENDI	04	EPJ C32 453	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49690
ABBIENDI	04F	EPJ C33 149	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49914
ABBIENDI	04H	EPJ C35 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49966
ABBIENDI	04N	PL B602 167	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=50288
ABDALLAH	04H	EPJ C34 145	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49920
ABDALLAH	04M	EPJ C36 1	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49977
Also		EPJ C37 129 (errat.)	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=50312
ACHARD	04	PL B580 37	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49692
ACHARD	04E	PL B587 16	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49893
AKERIB	04	PRL 93 211301	D.S. Akerib <i>et al.</i>	(CDMS II Collab.)	REFID=50537
BALTZ	04	JHEP 0410 052	E. Baltz, P. Gondolo		REFID=54106
BELANGER	04	JHEP 0403 012	G. Belanger <i>et al.</i>		REFID=50020
BOTTINO	04	PR D69 037302	A. Bottino <i>et al.</i>		REFID=50035
DESAI	04	PR D70 083523	S. Desai <i>et al.</i>	(Super-Kamiokande Collab.)	REFID=50356
ELLIS	04	PR D69 015005	J. Ellis <i>et al.</i>		REFID=50032
ELLIS	04B	PR D70 055005	J. Ellis <i>et al.</i>		REFID=50535
HEISTER	04	PL B583 247	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=49821
PIERCE	04A	PR D70 075006	A. Pierce		REFID=50536
ABBIENDI	03L	PL B572 8	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49550
ABDALLAH	03M	EPJ C31 421	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49689
AHMED	03	ASP 19 691	B. Ahmed <i>et al.</i>	(UK Dark Matter Collab.)	REFID=49498
AKERIB	03	PR D68 082002	D.S. Akerib <i>et al.</i>	(CDMS Collab.)	REFID=49838
BAER	03	JCAP 0305 006	H. Baer, C. Balazs		REFID=49694
BAER	03A	JCAP 0309 007	H. Baer <i>et al.</i>		REFID=50533
BOTTINO	03	PR D68 043506	A. Bottino <i>et al.</i>		REFID=49564
BOTTINO	03A	PR D67 063519	A. Bottino, N. Fornengo, S. Scopel		REFID=49725
CHATTOPAD...	03	PR D68 035005	U. Chattopadhyay, A. Corsetti, P. Nath		REFID=49563
ELLIS	03	ASP 18 395	J. Ellis, K.A. Olive, Y. Santoso		REFID=49112
ELLIS	03B	NP B652 259	J. Ellis <i>et al.</i>		REFID=49334
ELLIS	03C	PL B565 176	J. Ellis <i>et al.</i>		REFID=49455
ELLIS	03D	PL B573 162	J. Ellis <i>et al.</i>		REFID=49602
ELLIS	03E	PR D67 123502	J. Ellis <i>et al.</i>		REFID=50534
HEISTER	03C	EPJ C28 1	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=49379
HEISTER	03G	EPJ C31 1	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=49605
KLAPDOR-K...	03	ASP 18 525	H.V. Klapdor-Kleingrothaus <i>et al.</i>		REFID=49113
LAHANAS	03	PL B568 55	A. Lahanas, D. Nanopoulos		REFID=49523
TAKEDA	03	PL B572 145	A. Takeda <i>et al.</i>		REFID=49551
ABRAMS	02	PR D66 122003	D. Abrams <i>et al.</i>	(CDMS Collab.)	REFID=49139
ACOSTA	02H	PRL 89 281801	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=49293
ANGLOHER	02	ASP 18 43	G. Angloher <i>et al.</i>	(CRESSST Collab.)	REFID=49046
ARNOWITT	02	hep-ph/0211417	R. Arnowitt, B. Dutta		REFID=49350
ELLIS	02B	PL B532 318	J. Ellis, A. Ferstl, K.A. Olive		REFID=48942
HEISTER	02	PL B526 191	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48539
HEISTER	02E	PL B526 206	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48584
HEISTER	02J	PL B533 223	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48725
HEISTER	02N	PL B544 73	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48915
KIM	02	PL B527 18	H.B. Kim <i>et al.</i>		REFID=48932
KIM	02B	JHEP 0212 034	Y.G. Kim <i>et al.</i>		REFID=49117
LAHANAS	02	EPJ C23 185	A. Lahanas, V.C. Spanos		REFID=48970
MORALES	02B	ASP 16 325	A. Morales <i>et al.</i>	(COSME Collab.)	REFID=49038
MORALES	02C	PL B532 8	A. Morales <i>et al.</i>	(IGEX Collab.)	REFID=49234
ABREU	01	EPJ C19 29	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=48012
ABREU	01B	EPJ C19 201	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=48013
BALTZ	01	PRL 86 5004	E. Baltz, P. Gondolo		REFID=48140
BARATE	01	PL B499 67	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=48014
BARATE	01B	EPJ C19 415	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=48015
BARGER	01C	PL B518 117	V. Barger, C. Kao		REFID=48444
BAUDIS	01	PR D63 022001	L. Baudis <i>et al.</i>	(Heidelberg-Moscow Collab.)	REFID=47899
BERNABEI	01	PL B509 197	R. Bernabei <i>et al.</i>	(DAMA Collab.)	REFID=48166
BOTTINO	01	PR D63 125003	A. Bottino <i>et al.</i>		REFID=49239
CORSETTI	01	PR D64 125010	A. Corsetti, P. Nath		REFID=49240
ELLIS	01B	PL B510 236	J. Ellis <i>et al.</i>		REFID=48268
ELLIS	01C	PR D63 065016	J. Ellis, A. Ferstl, K.A. Olive		REFID=49238

GOMEZ	01	PL B512 252	M.E. Gomez, J.D. Vergados	REFID=49241
LAHANAS	01	PL B518 94	A. Lahanas, D.V. Nanopoulos, V. Spanos	REFID=48442
ABBIENDI	00	EPJ C12 1	G. Abbiendi <i>et al.</i> (OPAL Collab.)	REFID=47369
ABBIENDI	00G	EPJ C14 51	G. Abbiendi <i>et al.</i> (OPAL Collab.)	REFID=47486
ABBIENDI	00H	EPJ C14 187	G. Abbiendi <i>et al.</i> (OPAL Collab.)	REFID=47495
Also		EPJ C16 707 (errata.)	G. Abbiendi <i>et al.</i> (OPAL Collab.)	REFID=47787
ABBIENDI,G	00D	EPJ C18 253	G. Abbiendi <i>et al.</i> (OPAL Collab.)	REFID=48020
ABREU	00J	PL B479 129	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=47507
ABREU	00Q	PL B478 65	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=47608
ABREU	00T	PL B485 95	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=47724
ABREU	00U	PL B487 36	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=47725
ABREU	00V	EPJ C16 211	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=47749
ABREU	00W	PL B489 38	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=47755
ABREU	00Z	EPJ C17 53	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=47790
ABUSAIDI	00	PRL R4 5699	R. Abusaidi <i>et al.</i> (CDMS Collab.)	REFID=47634
ACCIARRI	00D	PL B472 420	M. Acciarri <i>et al.</i> (L3 Collab.)	REFID=47494
ACCOMANDO	00	NP B585 124	E. Accomando <i>et al.</i>	REFID=49237
BERNABEI	00	PL B480 23	R. Bernabei <i>et al.</i> (DAMA Collab.)	REFID=47623
BERNABEI	00C	EPJ C18 283	R. Bernabei <i>et al.</i> (DAMA Collab.)	REFID=48032
BERNABEI	00D	NJP 2 15	R. Bernabei <i>et al.</i> (DAMA Collab.)	REFID=49104
BOEHM	00B	PR D62 035012	C. Boehm, A. Djouadi, M. Drees	REFID=49200
ELLIS	00	PR D62 075010	J. Ellis <i>et al.</i>	REFID=47768
FENG	00	PL B482 388	J.L. Feng, K.T. Matchev, F. Wilczek	REFID=48058
LEP	00	CERN-EP-2000-016	LEP Collabs. (ALEPH, DELPHI, L3, OPAL, SLD+)	REFID=47501
MORALES	00	PL B489 268	A. Morales <i>et al.</i> (IGEX Collab.)	REFID=47761
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i> (PDG Collab.)	REFID=47469
SPOONER	00	PL B473 330	N.J.C. Spooner <i>et al.</i> (UK Dark Matter Col.)	REFID=47577
ACCIARRI	99H	PL B456 283	M. Acciarri <i>et al.</i> (L3 Collab.)	REFID=47034
ACCIARRI	99R	PL B470 268	M. Acciarri <i>et al.</i> (L3 Collab.)	REFID=47322
ACCIARRI	99W	PL B471 280	M. Acciarri <i>et al.</i> (L3 Collab.)	REFID=47493
AMBROSIO	99	PR D60 082002	M. Ambrosio <i>et al.</i> (Macro Collab.)	REFID=47231
BAUDIS	99	PR D59 022001	L. Baudis <i>et al.</i> (Heidelberg-Moscow Collab.)	REFID=46702
BELLI	99C	NP B563 97	P. Belli <i>et al.</i> (DAMA Collab.)	REFID=47326
OOTANI	99	PL B461 371	W. Ootani <i>et al.</i>	REFID=47222
ABREU	98P	PL B444 491	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=46682
ACCIARRI	98F	EPJ C4 207	M. Acciarri <i>et al.</i> (L3 Collab.)	REFID=45950
ACKERSTAFF	98P	PL B433 195	K. Ackerstaff <i>et al.</i> (OPAL Collab.)	REFID=46077
BARATE	98K	PL B433 176	R. Barate <i>et al.</i> (ALEPH Collab.)	REFID=46076
BARATE	98S	EPJ C4 433	R. Barate <i>et al.</i> (ALEPH Collab.)	REFID=46148
BERNABEI	98C	PL B436 379	R. Bernabei <i>et al.</i> (DAMA Collab.)	REFID=46247
ELLIS	98	PR D58 095002	J. Ellis <i>et al.</i>	REFID=46480
ELLIS	98B	PL B444 367	J. Ellis, T. Falk, K. Olive	REFID=47152
PDG	98	EPJ C3 1	C. Caso <i>et al.</i> (PDG Collab.)	REFID=45838
BAER	97	PR D57 567	H. Baer, M. Brhlik	REFID=49233
BERNABEI	97	ASP 7 73	R. Bernabei <i>et al.</i> (DAMA Collab.)	REFID=45563
EDSJO	97	PR D56 1879	J. Edsjo, P. Gondolo	REFID=47557
ARNOWITT	96	PR D54 2374	R. Arnowitt, P. Nath	REFID=49231
BAER	96	PR D53 597	H. Baer, M. Brhlik	REFID=49726
BERGSTROM	96	ASP 5 263	L. Bergstrom, P. Gondolo	REFID=49232
LEWIN	96	ASP 6 87	J.D. Lewin, P.F. Smith	REFID=49381
BEREZINSKY	95	ASP 5 1	V. Berezinsky <i>et al.</i>	REFID=49199
FALK	95	PL B354 99	T. Falk, K.A. Olive, M. Srednicki (MINN, UCSB)	REFID=44356
LOSECCO	95	PL B342 392	J.M. LoSecco (NDAM)	REFID=44116
ADRIANI	93M	PRPL 236 1	O. Adriani <i>et al.</i> (L3 Collab.)	REFID=43644
DREES	93	PR D47 376	M. Drees, M.M. Nojiri (DESY, SLAC)	REFID=43806
DREES	93B	PR D48 3483	M. Drees, M.M. Nojiri	REFID=49230
FALK	93	PL B318 354	T. Falk <i>et al.</i> (UCB, UCSB, MINN)	REFID=43807
KELLEY	93	PR D47 2461	S. Kelley <i>et al.</i> (TAMU, ALAH)	REFID=43279
MIZUTA	93	PL B298 120	S. Mizuta, M. Yamaguchi (TOHO)	REFID=43808
MORI	93	PR D48 5505	M. Mori <i>et al.</i> (KEK, NIIG, TOKY, TOKA+)	REFID=43641
BOTTINO	92	MPL A7 733	A. Bottino <i>et al.</i> (TORI, ZARA)	REFID=42065
Also		PL B265 57	A. Bottino <i>et al.</i> (TORI, INFN)	REFID=41599
DECAMP	92	PRPL 216 253	D. Decamp <i>et al.</i> (ALEPH Collab.)	REFID=41877
LOPEZ	92	NP B370 445	J.L. Lopez, D.V. Nanopoulos, K.J. Yuan (TAMU)	REFID=41998
MCDONALD	92	PL B283 80	J. McDonald, K.A. Olive, M. Srednicki (LISB+)	REFID=43804
ABREU	91F	NP B367 511	P. Abreu <i>et al.</i> (DELPHI Collab.)	REFID=41840
ALEXANDER	91F	ZPHY C52 175	G. Alexander <i>et al.</i> (OPAL Collab.)	REFID=41737
BOTTINO	91	PL B265 57	A. Bottino <i>et al.</i> (TORI, INFN)	REFID=41599
GELMINI	91	NP B351 623	G.B. Gelmini, P. Gondolo, E. Roulet (UCLA, TRST)	REFID=41770
GRIEST	91	PR D43 3191	K. Griest, D. Seckel	REFID=48059
KAMIONKOW...	91	PR D44 3021	M. Kamionkowski (CHIC, FNAL)	REFID=41729
MORI	91B	PL B270 89	M. Mori <i>et al.</i> (Kamiokande Collab.)	REFID=41805
NOJIRI	91	PL B261 76	M.M. Nojiri (KEK)	REFID=41684
OLIVE	91	NP B355 208	K.A. Olive, M. Srednicki (MINN, UCSB)	REFID=41771
ROSZKOWSKI	91	PL B262 59	L. Roszkowski (CERN)	REFID=41688
GRIEST	90	PR D41 3565	K. Griest, M. Kamionkowski, M.S. Turner (UCB+)	REFID=41671
BARBIERI	89C	NP B313 725	R. Barbieri, M. Frigeni, G. Giudice	REFID=49229
OLIVE	89	PL B230 78	K.A. Olive, M. Srednicki (MINN, UCSB)	REFID=41812
ELLIS	88D	NP B307 883	J. Ellis, R. Flores	REFID=49228
GRIEST	88B	PR D38 2357	K. Griest	REFID=49227
OLIVE	88	PL B205 553	K.A. Olive, M. Srednicki (MINN, UCSB)	REFID=40755
SREDNICKI	88	NP B310 693	M. Srednicki, R. Watkins, K.A. Olive (MINN, UCSB)	REFID=40756
ELLIS	84	NP B238 453	J. Ellis <i>et al.</i> (CERN)	REFID=12520
GOLDBERG	83	PRL 50 1419	H. Goldberg (NEAS)	REFID=12508
KRAUSS	83	NP B227 556	L.M. Krauss (HARV)	REFID=12510
VYSOTSKII	83	SJNP 37 948	M.I. Vysotsky (ITEP)	REFID=12511

Translated from YAF 37 1597.