


JIŘÍ MLÍKOVSKÝ

CENOZOIC BIRDS OF THE WORLD

PART 1: EUROPE



PRAHA: NINOX PRESS

2002

Jiří Mlíkovský

Vršovická 11, CZ-101 00 Praha 10, Czech Republic
e-mail: mlikovsky@sendme.cz

Published by NINOX Press, Praha.
Date of publication: 11 January 2002.
Binding: PBtisk Příbram.
© by the author.

ISBN 80-901105-3-8

Contents

Introduction	6
Scope and arrangement of the catalogue	6
Stratigraphy	8
Names	8
Taxonomic background	14
Classification of birds	19
List of collections	21
Acknowledgments	22
History of the research	24
List of localities	26
Systematic list	58
Tinamiformes	58
Tinamidae	58
Struthioniformes	60
Palaeotididae	60
Struthionidae	60
Phaethontiformes	63
Gaviidae	63
Phaethontidae	64
Ardeiformes	66
Sulidae	66
Ardeidae	67
Phalacrocoracidae	70
Anhingidae	73
Bucerotiformes	75
Laurillardidae	75
Upupidae	76
Phoeniculidae	77
Procellariiformes	78
Procellariidae	78
Pelagornithidae	81
Ciconiiformes	85
Podicipedidae	85
Pelecanidae	86
Messelornithidae	87
Gruidae	88
Threskiornithidae	92
Gastornithidae	94
Eogruidae	96
Otididae	98
Ciconiidae	100
Cathartidae	101
Anseriformes	103
Phoenicopteridae	103
Recurvirostridae	107
Anseranatidae	107
Anatidae	109
Charadriiformes	126
Jacanidae	126
Rostratulidae	126

Scolopacidae	126
Glareolidae	132
Thinocoridae	133
Pterocletidae	133
Charadriidae.....	135
Laridae	136
Alcidae	138
Piciformes	141
Zygodactylidae	141
Coliidae	145
Psittacidae	146
Ramphastidae	147
Picidae	148
Galliformes	150
Megapodiidae	150
Cracidae	151
Phasianidae	151
Accipitriformes	171
Rallidae	171
Cariamidae	178
Cuculidae	185
Accipitridae	187
Columbiformes	197
Caprimulgidae	197
Aegialornithidae	200
Apodidae	203
Podargidae	205
Sophiornithidae	206
Strigidae	207
Columbidae	218
Falconidae	220
Coraciiformes	223
Coraciidae	223
Trogonidae	224
Alcedinidae	225
Todidae	225
Meropidae	226
Passeriformes	227
Eurylaimidae	227
Hirundinidae	227
Alaudidae	228
Bombycillidae	229
Laniidae	229
Sturnidae	230
Corvidae	230
Troglodytidae	236
Turdidae	237
Sylviidae	240
Prunellidae	241
Aegithalidae	242
Paridae	242
Certhiidae	243

Motacillidae	243
Passeridae	244
Fringillidae	244
Emberizidae	246
Family incertae sedis	247
Aves incertae sedis	250
Ichnotaxa	273
Taxa non avium	277
Nomina nuda	279
References	281
Index of authors	363
Index of countries	373
Index of localities	375
Index of collections	386
Index of stratigraphical units	388
Index of family-group names	389
Index of genus-group names	390
Index of species-group names	394

Introduction

Birds of the past entered the realm of human knowledge at the dawn of the 17th century, when Swiss naturalist Johann Jacob von Scheuchzer (1708a,b) described from a quarry near Öhningen in Switzerland a feather imprint of an unidentified bird¹. A century later, French naturalist Georges Cuvier (1822) founded the science of paleornithology. Today, fossil birds are known from almost 600 Tertiary and 1200 Pleistocene localities of Europe (Mlíkovský 1996a and the present paper, Tyrberg 1998, 2001), which represents roughly half of all avian localities known from the world.

First reviews of the fossil birds of Europe were presented already in the early 19th century by Hoff (1801), Cuvier (1822) and Krüger (1823). The rapidly growing knowledge was continuously mapped in a variety of subsequent reviews, the most important of which were those by Gervais (1844a), Milne-Edwards (1863, 1867-1868, 1869-1871), Lydekker (1891a), Lambrecht (1933), and Brodkorb (1963c, 1964, 1967, 1971b, 1978). Lists of the European fossil birds were presented also by Lambrecht (1921) and Bocheński (1997).

The aim of this paper is to present a comprehensive systematic review of the Cenozoic birds of Europe at the close of the 20th century. The review is based on my studies of the bones of fossil birds from 13 European countries, on the restudy of published figures and descriptions of type materials of most other taxa of fossil birds of Europe, and on the extensive scanning of literature.

Scope and arrangement of the catalogue

The catalogue lists all fossil and extinct bird species known from the European Cenozoic, and all modern species with fossil record older than the middle Pleistocene. It covers in full the period from the beginning of the Tertiary till the end of the early Pleistocene (i.e. ca. 65-0.6 Ma BP), while only records of fossil and extinct species are comprehensively listed for the last 0.6 Ma. The many other records from the latter time period can be found in Tyrberg (1998, 2001).

Europe is understood here in the geographical sense. It is bordered by the sea from the north, west and south. Its eastern border tracks the eastern foot of Ural mountains (i.e. approximately the 60° E) and Emba river, which enters in the Caspian Sea. Between the Caspian and Azov Seas, the border is formed by the Kuma-Manyč depression and the lower Don river, which opens into the Azov Sea. Islands northwest of the European continent are included here to Iceland and Jan Mayen, and most Mediterranean islands are included as well, excepting Cyprus and small islands laying near the continent of Africa. Macaronesian islands are not included. See also Mlíkovský (1996b). The area of Europe is ca. 10 527 000 km².

The catalogue is arranged systematically. Avian taxa of uncertain taxonomic position within families are listed at the end of each section, while taxa of uncertain position within the Aves, ichnotaxa, non-avian taxa formerly thought to represent birds, and nomina nuda are listed at the end of the catalogue. However, those nomina nuda, which could be unequivocally assigned to a valid taxon, are listed in the synonymy of the respective taxon, although they do not represent synonyms in the sense of the ICZN (1999).

The entries include the following data:

Fossil species:

- ☐ Reference to original description, incl. page number.
- ☐ Data on the type material, incl. its current place of deposition, and references to its figures. Note, that Brodkorb (1963c, 1964, 1967, 1971b, 1978) listed in his *Catalogue* figures of

¹ The bird was believed to have perished in the Biblical Flood. The locality, now in Germany, is known to be middle Miocene in age. The bird continues to be unidentified. For more data on this specimen see Footnote 2.

both typical and referred material, but limited this information to figures in that paper, in which the species was described.

- ❑ Nomenclatural status of each name, if different from the standard, incl. indications of senior and junior homonyms, and nomina nuda.
- ❑ Full synonymy, incl. new combinations, correct and incorrect emendations (but not subsequent incorrect spellings), and changes in rank.
- ❑ Full list of localities, incl. extralimital ones (where known). The localities are listed chronologically, and within stratigraphical units geographically from the northwest to the southeast. Relevant authors are cited for each locality, but authors who just listed the species or the locality in a catalogue or in another paper are not.
- ❑ Remarks, particularly those on the taxonomic status of the species.

Modern species:

- ❑ General reference to its original description. These papers are not cited in the References section.
- ❑ List of fossil synonyms, for which full data are given as above.
- ❑ Full list of pre-middle Pleistocene localities, and a list of younger localities, where the remains were attributed to a fossil taxon. The respective name of the fossil taxon is listed in parentheses. For a list of the middle and late Pleistocene localities of modern species see Tyrberg (1998, 2001).
- ❑ Remarks, particularly those on the taxonomic status of the species.

Fossil genus:

- ❑ Reference to original description, incl. page number.
- ❑ Type species, incl. data on its fixation as a type.
- ❑ Nomenclatural status of each name, if different from the standard, incl. indications of senior and junior homonyms, and nomina nuda.
- ❑ Full synonymy, incl. correct and incorrect emendations (but not subsequent incorrect spellings), and changes in rank.
- ❑ General data on the distribution if more than a single species is included in the genus, incl. data on extralimital localities.
- ❑ Remarks, particularly those on the taxonomic status of the genera.

Modern genus:

- ❑ General reference to original description. These papers are not cited in the References section.
- ❑ List of fossil synonyms, for which full data are given as above.
- ❑ Remarks, particularly those on the taxonomic status of the genera.

Fossil family:

- ❑ Reference to its original description, incl. page number.
- ❑ Type genus.
- ❑ Nomenclatural status of each name, if different from the standard. See also Mlíkovský (2000f).
- ❑ Full synonymy, incl. correct and incorrect emendations (but not subsequent incorrect spellings), and changes in rank.
- ❑ General data on the distribution if more than a single species is included in the family, incl. those on extralimital localities.
- ❑ Remarks, particularly those on the taxonomic status of the families, and the groupings of genera.

Modern family:

- ❑ General reference to original description. These papers are not cited in the References section.
- ❑ List of fossil synonyms, for which full data are given as above.

- Remarks, particularly those on the taxonomic status of the families, and the groupings of genera.

Modern order:

- General reference to original description. These papers are not cited in the References section. Note, that no fossil orders are recognized in this paper.
- List of fossil synonyms, for which full data are given as above.

Stratigraphy

The stratigraphical arrangement of the Paleogene follows Schmidt-Kittler (1987) and Legendre & Lévêque (1997). The period is divided into 30 Paleogene micromammal zones (MP), but no definitions are available for the first five MP-zones, which encompass most of the Paleocene. See Fig. 1 for additional data.

The stratigraphical divisioning of the Neogene follows Mein (1976), respecting subsequent amendments of the system (Fahlbusch 1976, 1991, Mein 1990, Steininger et al. 1987, Bruijn et al. 1992, Steininger et al. 1996, Steininger 1999). The period has been divided in 17 Neogene micromammal zones (MN). Guérin (1982) separated the youngest part of the MN 17 zone as MN 18. The latter proposal is supported by ornithological data (Mlíkovský 2001) and is used in the present paper. Some of the MN-zones (particularly MN 3, 4, 16, and 17) are now subdivided into two subzones each, labeled with letters (*a* being older than *b*). The MN-zones 11-13 are clearly separable in westernmost Europe, but are difficult to discern in southeastern Europe. See Fig. 1 for additional data.

Three numerical systems for the divisioning of the Pleistocene were proposed by Horáček (1981, Horáček & Ložek 1988; Q zones), Guérin (1982; MNQ zones), and Fejfar & Heinrich (1990a,b, Fejfar et al. 1997, 1998; MQ zones), respectively. A derived system developed by Mlíkovský (2001) is used in the present paper. See Fig. 1 for a comparison of these systems. Note, that Guérin's (1982) MNQ zones simply correspond with the main European glacials and interglacials (in the sense of Penck & Brückner 1901-1909), the identity of which is dubious (see Kukla 1975, 1977, 1978).

It should be noted, that general stratigraphical terms used in literature, such as Pannonian or Biharian are used in different sense by different authors, and cannot be interpreted without the knowledge of how the author understood it. General understanding of the stratigraphical divisioning of the European Cenozoic in the early 20th century is correlated with the modern understanding in Fig. 1 (reconstructed from Lambrecht 1933).

Absolute ages are ca. 65-24 Ma for the Paleogene, 24-5.5 Ma for the Miocene, 5.5-1.65 Ma for the Pliocene, 1.65-0.01 Ma for the Pleistocene, and 0.01-0 Ma for the Holocene (Legendre & Lévêque 1997, Steininger et al. 1996).

Names

Transliteration from non-Roman alphabets

Names originally written in non-Roman alphabets are converted here to the Roman (= Latin) alphabet using transliteration. Note, that these names were formerly often transcribed or even translated, which resulted in variant spellings (see Mlíkovský 1996b).

Names originally written in the Cyrillic alphabet are converted here to the Roman alphabet using the "Berlin" system of transliteration, adopted in 1978, which superceded the older "London" system of transliteration. See Mlíkovský (1996b: 523) for the transliteration table. The use of the Cyrillic alphabet is currently limited to several eastern Slavonic languages (Russian, Ukrainian, Belorussian) and southern Slavonic languages (Bulgarian, Macedonian, Serbo-Croatian in Serbia). In addition, it was used in the 1940-1980s in Moldovan, a Romance language, when Moldova was part of the then Soviet Union.

In converting the Greek alphabet to the Roman one, I applied the system adopted by the ICZN (1985). Greek alphabet is used only in the Greek language.

Lambrecht ¹		Modern ²		MQ ³	MQ ⁴	Q ⁵	MNQ ⁶	
<i>Holocene</i> ⁷		Holocene		2E				
<i>Pleistocene</i> ⁸		Pleistocene	late	2D	2	4	26	
				2C			25	
<i>Pliocene</i>	<i>late</i>	Pleistocene	middle	2B	1	3 _{2,3}	24	
				2A			3 ₁	23
			early	1b		2	21	
				1a		1	20	
<i>Miocene</i>	<i>early</i>	Pliocene	late	Villanyian	18	17	17 ₃	18
					17		17 _{1,2}	17
					16	16	16	16
<i>Miocene</i>	<i>late</i>	Pliocene	early	Ruscinian	15	15	15	15
					14	14		14
<i>Miocene</i>	<i>late</i>	Miocene	late	Turolian	13	13		13
					12	12		12
					11	11		11

<i>Miocene</i>		<i>late</i>	<i>Miocene</i>	<i>late</i>	Vallesian	10	10		10
						9	9		9
					Astaracian	8			
						7			
Orleanian	6								
	5								
<i>early</i>			<i>early</i>	Agenian	4				
					3				
					2				
<i>Oligocene</i>		<i>late</i>	<i>Oligocene</i>	<i>late</i>	Chattian	1			
						2			
						30			
						29			
						28			
						27			
					Stampian	26			
						25			
						24			
						23			
<i>middle</i>			<i>middle</i>		22				
					21				
<i>early</i>			<i>early</i>						

<i>Eocene</i>	—	<i>Eocene</i>	late	Priabonian	20			
					19			
					18			
				Bartonian	17			
					16			
					15			
	<i>middle</i>		Lutetian	14				
				13				
				12				
	<i>early</i>		Ypresian	11				
				10				
				9				
8								
<i>Paleocene</i>		<i>Paleocene</i>	late		7			
			early		6			
					1-5			

Fig. 1. Biochronological chart of the Cenozoic of Europe. 1 - state in the early the 20th century (reconstructed after Lambrecht 1933); 2 - current understanding (used in this paper); 3 - current understanding (used in this paper) after Mlíkovský (2001; MN 18 to Recent), Mein (1976; MN 1-17: Miocene and Pliocene), and Schmidt-Kittler (1987; MP 1-30: Paleocene, Eocene and Oligocene). 4 - after Fejfar & Heinrich (1990a,b), 5 - after Horáček (1981), 6 - after Guérin (1982). See text for additional references, where the systems used were amended (the amendments being taken into account here). 7 - often called 'Alluvium', 8 - often called 'Diluvium'.

Personal names and toponyms

Personal names and toponyms are subject to variation, which results from several factors. In all cases, the original spelling of the name (transliterated from non-Roman alphabets) is given precedence. For details see below.

Use of different systems of conversion of names from a non-Roman alphabet to the Roman one. For example, Курочкин (original spelling in the Russian version of the Cyrillic alphabet) → Kuročkin (correct transliteration using the Berlin system), Kurochkin (transliteration using the London system of transliteration, or English transcription), and Kuroczkin (Polish transcription).

Transcriptions between languages within the Roman alphabet. Transcriptions adapt the spelling so that the pronunciation remains approximately the same in the other language. This system was often applied within the Roman alphabet until first decades of the 20th century. For example, Novák (Czech) → Nowak (German), Rebiełice (Polish) → Rembielice (e.g. English). Original spellings are used in all cases, but variant spellings are listed in the respective Indexes.

Transcriptions between languages within the Cyrillic alphabet. Names continue to be transcribed from language to language within this alphabet. This makes it difficult to recognize which is the original spelling of a personal name or a toponym without additional data, such as nationality of the author or ethnical origin of a village. Such data are obviously unavailable in many cases, so I have chosen the standard name with respect to the official language accepted as the original spelling that used in the country of the author's main activity.

Translation of names, sometimes used till the first half of the 20th century, was applied mainly to first names. For example: Kálmán (Hungarian) → Koloman (German), Várhegy (Hungarian) → Castle Hill (English). Also Latinization, common till the early 18th century, belongs to this category: e.g. Linné (Swedish) → Linnaeus (Latin). Original names were used throughout this paper, but Linnaeus was given preference over Linné, because of current standards in zoological nomenclature.

Omission of diacritical marks, which is still very common. For example, Švec (Czech) → Svec, Ciobanița (Romanian) → Ciobanita.

Carelessness of authors. For example Mlíkovský (Czech) can be found in scientific literature with at least three typographical errors (in addition to omitted diacritical marks) → Milikowski (any combination of misprints possible).

Double or single surnames of Spanish/Catalan authors, e.g. Sánchez Marco ↔ Sánchez. Here I followed the prevailing use.

Changes of ladies' names, following marriage. To my best knowledge, this applies only to three researchers in European paleornithology: Brjuzgina → Umans'ka, Chauviré → Mourer-Chauviré, and Malez → Malez-Bačić. These authors are cited according to the name they used in the particular paper.

Use of explanatory names in composite toponyms: (variant spellings and alternative names being listed in the Index of localities). Many, particularly Quaternary localities have composite names, one or more of which are explanatory names, such as "hill" or "cave". The explanatory names are sometimes used in the original language, sometimes translated, and sometimes omitted. In the present paper, I omit the explanatory name if it is written separately, but I include it if it is written together with the proper name. For example, the Hungarian locality <Somssichhegy> (i.e. Somssich hill), is listed here under the name <Somssichhegy>, while the French locality <La Grotte du Lazaret> (i.e. the Lazaret cave), is listed under the name <Lazaret>. When the proper name is an adjective (this being the case particularly in the Slavonic languages), and the explanatory name forms thus part of the locality name, the locality is listed under the composite name, but the explanatory name is translated to English. For example: the Russian locality <Medvež'ja peščera> (i.e. Bear Cave) is listed under the name <Medvež'ja Cave>. Exceptions are allowed only if the name could cause any doubts.

Use of articles. Definite articles, such as <La>, <L'> or <Les>, often precede the proper

name of a locality or a person, particularly in the Romance languages. These articles are always omitted in toponyms, but included as alternative names in the respective Indexes.

Renaming of localities. Some localities were renamed, when changing state borders influenced their administrative position. For example Püspökfürdő → Betfia, Krottensee → Mokřina. These localities are listed under their new names, and the earlier names are listed in the Index of localities.

Note: Toponyms are cited in original languages up to the level of cities (Praha, not Prague), but names of the countries are translated to English (Germany, not Deutschland).

Order-group taxa

The names of the order-group taxa are not regulated by the International Code of Zoological Nomenclature (ICZN 1999). However, I follow here the ornithological standards, first adopted by Brodkorb (1963c, 1964, 1967, 1971b, 1978), and treat these names similarly as those of the family-group taxa, with the exception of endings, for which I follow Wetmore's (1930b) proposals.

Family-group taxa

The names of the family-group taxa are regulated by the International Code of Zoological Nomenclature from 1961 onwards (ICZN 1999, see also Bock 1994). The names adopted here are based on the lists by Brodkorb (1963c, 1964, 1967, 1971b, 1978), Bock (1994; but see Olson 1995b), and Mlíkovský (2000f). In synonymies, family-group names are cited in the original form. However, this may not be true for the few family-group names taken from Bock (1994), because he cited all names in the emended form, and with the ending *-idae*. I was not able to control the original spelling of these names before the deadline for the present paper.

Genus-group taxa

The rules of the International Code of Zoological Nomenclature (ICZN 1999) were applied. The only major change which resulted from the application of the Code regards the status of several genus names proposed by Lambrecht (1933). Type species were not fixed for some of them, which is mandatory after 1930 (ICZN 1999, Art. 13.3.), and are thus *nomina nuda*. Brodkorb (1952) fixed type species for these genera, becoming thus the author of all of them, although he attributed them to Lambrecht (1933).

Species-group names

The rules of the International Code of Zoological Nomenclature (ICZN 1999) were applied. The only relevant problem regards the identification of the name-bearing types of some of the species described by Milne-Edwards (1863, 1867-1868, 1869-1871) from the early Miocene of Saint-Gérard-le-Puy in France, because some were based on hundreds of specimens, and the type series were not unequivocally defined in the original papers. In the present paper, I listed in the type series all bones mentioned by Milne-Edwards, when he describing the species, and found in the early collections of MNHN. These lists probably differ from the proper types series, because: (1) Milne-Edwards sometimes indicated that he examined – and identified as such – also specimens from other, usually private collections, the present location of which is unknown; (2) some of the specimens from the original series deposited in MNHN are missing; and (3) considerable material was accumulated between the early 1860s and mid 1870s, and this material cannot be discerned at MNHN (this applies to the species described by Milne-Edwards in 1863). Hence, not all of the bones in MNHN formed the type series of these species, while also an unknown number of bones not deposited in MNHN (and probably lost in many cases) belonged to the respective type series.

In his revisions of Milne-Edwards's species, Cheneval (1983a,b,c, 1984a, 2000) presented lists of syntypes for each species, and selected relevant lectotypes, without observing the rules

of zoological nomenclature. His lists of the alleged lectotypes include bones which Milne-Edwards attributed to other species (even those which are holotypes of other Milne-Edwards's species), while they do *not* include bones which Milne-Edwards evidently assigned to the given species, but which Cheneval re-identified as belonging to some other species. Hence, his lists of the alleged syntypes are incorrect in these cases. Also Cheneval's selection of lectotypes of these species is often unhappy. He overlooked that most (if not all) of the drawings in Milne-Edwards (1867-1868, 1869-1871) are side-reversed, and selected the lectotypes to match the figures, i.e. unfigured specimens from the other side of body than intended. It is thus not certain, whether the lectotypes selected by Cheneval were part of the original type series or not. In spite of this, I consider his respective actions valid, until the opposite is proven.

Milne-Edwards (1863, 1867-1868, 1869-1871, 1874, 1892) described also a number of species on the basis of a few bones. They are usually clearly identifiable, but the way how the material was described caused much confusion as regards the decision whether a species was based on a holotype or on a type series. In many cases, Milne-Edwards based the new species on a single bone, adding other bones (often with descriptions) as assigned material, usually with a comment "I also assign to this species" or that alike. In such cases I consider the first bone the holotype of the species, following the intentions of Art. 72.4.6 of the ICZN (1999).

English names of modern species were accepted from Beaman (1994).

Taxonomic background

Recognition of taxa is always dependent on the underlying taxonomic philosophy, although practical paleontologists usually do not explain it. In paleornithology, most researchers evidently recognized taxa using purely subjective assessment of similarities and dissimilarities of the material they evaluated, looking for "gaps" between supposed taxa of any rank. Assignment of taxa to the rank was also purely subjective. Below, I understand *species* as units organized at the population level, *genera* and *families* as evolutionary-ecomorphological units (of lower and higher level, respectively), and *orders* as units of adaptive radiation. All the *sub-* and *super-* taxa are understood here as purely subsidiary units, which make the classification more lucid.

Order

Orders are understood here as units of the adaptive radiation of families. Because families are defined here in ecomorphological terms (plus monophyly), the orders include birds, which are often very different in appearance, but which have a common ancestor.

Family

For the history of the family notion see d'Hondt (1987) and Bock (1994). In the present paper I understand families as units similar to the genera, but occupying adaptive zones at a higher rank (Mlíkovský 1982d, 1983a, 1987b). Each adaptive zone of the family rank is composed of one or more adaptive zones of the genus level.

One or more phases of the adaptive radiation of the genera can occur within a family (see Müller 1955, 1961). Taxonomically, I interpret these phases as within-family units. For example, the Strigidae are represented in Europe from the late Paleocene to the end of the Eocene by ancestral forms, barn owls (Tytoninae) are prevailing in the Oligocene and the early Miocene, while modern types of owls (Striginae) appeared during the early Miocene (in MN 3), and prevail from the middle Miocene to the present (Mlíkovský 1998c,f).

Genus

The genus notion underwent considerable evolution (see e.g. Inger 1958, Smirnov 1960, Tintant 1984, Dubois 1988). In the present paper, the genus is understood as a monophyletic part of an adaptive zone (Mlíkovský 1987b). The notion of adaptive zones was introduced to the evolutionary biology by Simpson (1944, 1953), who believed that adaptive zones are

occupied by taxa. Mlíkovský (1982d, 1983a, 1987b) suggested that counterparts of adaptive zones are not taxa, but life forms, i.e. ecomorphological units. Each adaptive zone corresponds to a life form, which is defined in ecomorphological terms, and which does not reflect the evolutionary past of the organisms which belong to it. A taxon is then “a part of a phyletic line bordered by the limits of an adaptive zone, or, in other words, a taxon is a monophyletic part of a life form” (Mlíkovský 1983a: 311). Macroevolution is thus basically an ecological process (Wahlert 1965, 1973, Mlíkovský 1982d, 1983a,b, 1987b, Lemen & Freeman 1984), and this process is not reducible to speciation.

Life forms of birds are usually defined by body proportions and/or by bill shape (see e.g. Leisler 1980, 1985), and – if the material is sufficient – these data are available also for fossil forms. Accordingly, the recognition of ecomorphological units is possible in both modern and fossil birds. Bones bear sufficient characters to discern whether a group of organisms within an adaptive zone originated from a single invader, i.e. whether the group is monophyletic (see Mlíkovský 1982d, 1987b). Hence, taxa of the genus rank can be defined identically for both the modern and the fossil birds.

There are no theoretical limits for the stratigraphical and geographical distribution of the genera.

Species

The species notion goes back to the Ancient Greece (Uhlmann 1923, Balme 1962), and considerably changed since that time (e.g. Mayr 1963, 1982, Nowiński & Kuznicki 1965, Mlíkovský & Zemek 1983). In this paper I understand a species as an evolutionary unit organized at the population level, and existing in the evolutionary (historical) time. The evolution of species is realized in the particular (ecological) time through its particular forms, i.e. populations (see Mlíkovský & Zemek 1983). The relativity of time solves the aged problem of the relation between species and populations, and also between the horizontal (“biological”) and vertical (“phylogenetic”) understanding of species.

Species as sets: From the point of view of practical systematists, species (and other taxa) are usually understood as classical sets, i.e. sets to which elements either belong or not. (This does not compete with the opinion that species are functional units, i.e. individuals – see also Van Valen 1988) This approach is untenable from the evolutionary point of view, because it implies evolution by jumps, i.e. the discontinuous evolution, which is at variance with the available data. I understand species as fuzzy units in the sense of Zadeh (1965), which allow for the continuous evolution (Mlíkovský 1983a,b; see also Kitcher 1984, Stent 1986, Van Valen 1988). Hence, the famous “missing links” need not be unequivocally attributable to any species.

Phenons vs. species

Specimens are described using “characters” (auctorum), for which also the terms “phenes” (Johannsen 1909) and “biocharacters” (Osborn 1917) were applied. In a n-dimensional space (formed by both metrical and morphological variables) specimens form clusters with unimodal distribution, known as “phenons” (Camp & Gilly 1943). The relation between these phenons and species is ambiguous, because a single phenon can consist of more than one species, while a single species can include specimens from more than one phenon (Mlíkovský et al. 1985). Because of this, phenons cannot be simply interpreted as species. The situation is the more complicated, because different bones may show different results.

Example 1: Wing bones of *Lagopus lagopus* and *Lagopus mutus*, two grouse species frequently found in the late Pleistocene deposits of Europe, fall into a single phenon, while their hindlimb bones, particularly the tarsometatarsus, form two rather distinctive clusters (Kraft 1972). Hence these two modern (and good) species cannot be separated on the basis of their wing bones, but can be well separated on the basis of their tarsometatarsi.

Example 2: A similar picture, but at the intraspecific level, is shown by domestic chickens, where the wing bones of males and females of the European standard early Medieval race form a single phenon, while hindlimb bones of the two sexes fall into two distinct clusters (Benecke 1989, Prummel 1993, Mlíkovský 2001). In the latter case, two phenons belong to a single species.

Example 3: Tarsometatarsi of the two modern peacock species, *Pavo cristatus* and *Pavo muticus*, fall into three clusters, of which the smallest (= with shortest tarsometatarsi) is formed by female *cristatus*, the middle-sized includes male *cristatus* and female *muticus*, and the largest contains only male *muticus* (see Mourer-Chauviré 1989b for measurements). Here, three phenons are observed, which are formed from two species.

Phyletic lines vs. species

The separation of phyletic lines into species is basically the question of the tempo and mode of evolution, and of the longevity of species. The theories on the tempo and mode of evolution – which influenced avian paleontology – are as follows: (1) species do not evolve (prevailing pre-Darwinian opinion), (2) species do evolve at a constant rate (prevailing early Darwinian opinion), (3) species do evolve at a changing rate (Simpson 1953, 1961, and many others), (4) species (i.e. phyletic lines in this sense) alternate periods of stasis with periods of rapid change (Eldredge & Gould 1972, Gould & Eldredge 1977, and many others). In my opinion, the rate of evolution (i.e. the rate of morphological change in the restricted sense) ranges between zero (stasis) and high (rapid), and the periods of no change, slow change and rapid change can alternate (see also e.g. Hoffman 1982, Gingerich 1985, Fortey 1985, 1988, Haffer 1995).

Phyletic lines can be separated into species in one of the following ways: (1) arbitrarily (common approach; often subjectively using the criteria 2, 4, and 5, as listed below), (2) where a gap in the fossil record occurs (common approach; requires insufficient knowledge of the fossil record), (3) at each branching point (e.g. Hennig 1966, Cracraft 1983, Willmann 1985; requires the cladistic approach to evolution), (4) when the older and the next younger part of the phyletic line are sufficiently distinct morphologically (common approach; requires insufficient knowledge of the fossil record, and a subjective assumption or an expressive statement what does it mean "sufficiently distinct"), (5) according to stratigraphical units (common, cf. e.g. Gingerich 1979; requires forever defined stratigraphical units), (6) according to a given time period (generally identified with the criterion 5; requires constant rate of evolution), (7) when slow rate of evolution is interrupted with a period of rapid change (Simpson 1943, 1953, 1961), and (8) when the evolutionary stasis is interrupted with a period of rapid change (Eldredge & Gould 1972, Gould & Eldredge 1977, and many others; this is just the extreme case of the criterion 7). All of these criteria (with the possible exception of the criterion 3) have been applied to fossil birds, but only No. 7 and 8 delimitate natural species (No. 7 only if the difference between the slow and rapid change is marked), while species separated according to the criteria 1-6 are artificial (see also Simpson 1961, Zarenkov 1976, Wiley 1978, 1981).

I do not recognize artificial "species" created by arbitrary delimitations of phyletic lines, such as allochronous species (Mayr 1942), chronospecies (George 1956), or successional species (Imbrie 1957). Accordingly, I do not believe, that the existence of any species is a-priori time-limited.

For various on the speciation rate and/or species longevity in birds see also Brodkorb (1960a), Ballmann (1977) and Zink & Slowinski (1995).

Intraspecific variability

Both morphological and metrical variability are considered. The variability as such is usually small in strictly defined, homogenous samples, but sharply increases when the sample includes individuals from: (1) different age classes, (2) different sex classes, (3) geographically different localities, and (4) stratigraphically different localities.

Allometrical variability: Both morphological and metrical variability is strongly, though in a predictable way influenced by allometrical variability. This phenomenon is often neglected in paleornithological studies. Objects are allometrically similar, when their proportions change with their overall size in an given manner, which can be usually described using the following equation: $P = a.W^b$, where P = dimension of the part, W = dimension of the whole, and a and b are coefficients (Röhrs 1959, 1961, Gould 1966, Rayner 1985, La Barbera 1989 and many others). In avian bones, it is possible to predict, that larger bones (intraspecifically or intragenerically) will be: (1) more robust, (2) will have processes more pronounced, and (3) will have muscular attachments more deepened (e.g. Rensch 1940, Hoerschelmann 1966, Alexander 1983, Anderson et al. 1985, Campbell & Marcus 1992, Mlíkovský orig., see also McMahon 1973, 1975). Morphological differences between congeneric species are often described using exactly these characters (see e.g. the alleged morphological differences between the Miocene European *Palaelodus* species listed by Cheneval 1983c and Heizmann & Hesse 1995). When controlled for size, the alleged differences disappear, as is also the case with the European *Palaelodus* species (Mlíkovský, orig.). Jánossy (1972, 1974a,b, etc.) based many of his new species of birds on alleged allometrical differences. Unfortunately, he totally misinterpreted the meaning of allometry: what he called allometric *difference*, is in fact the allometric *similarity*.

Metrical variability: The degree of variability of metrical characters is generally expressed using the coefficient of variability (CV), calculated as $CV = 100.SD/M$, where SD = standard deviation, and M = arithmetical mean. Metrical characters are normally distributed as a rule (see e.g. Warheit 1992), but the value of CV sharply increases with lower homogeneity of samples (see above for possible factors), with the measurement error (Pankakoski et al. 1987), and when the accuracy of measurements is too close to the mean value (Soulé 1982, Rohlf et al. 1983, Pankakoski et al. 1987). The threshold is reached when the accuracy of measurement is less than 1-2% of the mean (Mlíkovský orig.), i.e. if the bone is measured with an accuracy to 0.1 mm (as is usual in paleornithological studies), then the mean size of the measured object should not be smaller than approximately 15 mm (which is often violated when widths or depths of bones are measured).

Considering that CV is measured properly, it usually reaches value 3-5 in modern populations of birds, if they are geographically restricted, and when only adult individuals of one sex are considered (Mlíkovský, in prep.). Samples of fossil bones of birds are usually far less homogenous, so that the values of CV will be higher, usually reaching values around 10 (Mlíkovský, in prep.). The values of CV can be even higher when body size of birds from a given phyletic line is consistently growing or decreasing over time.

Only bones of adult birds are normally included in biometrical analyses of fossil birds. The age of birds is identified according to the state of ossification of the bones. Nevertheless, Bacher (1967) and Northcote (1981) showed that long bones of subadult Mute Swans *Cygnus olor*, which were inseparable from fully ossified bones, were markedly shorter in mean than the corresponding elements of adult birds. This is another factor contributing to the non-homogeneity of samples. In addition, this observation could explain the occurrence of isolated bones which are smaller than expected for a given species.

Morphometrical characters are usually normally distributed, which allows for an estimation of the range of measurements in any given case. Generally, roughly 70 % of measurements will be found within $0.01 CV.M$ around the mean, and 95 % of them within $0.02 CV.M$ around the mean. Hence, bones which fall into the limits $\pm 0.1 CV.M$ are considered here as belonging to the same species, while bones within the limits $\pm 0.2 CV.M$ are considered as probably belonging to the same species. However, this criterion is not used strictly.

Discerning congeneric species: morphological vs. metrical characters

Intrageneric variation in size is much higher than the variation in shape (Lemen & Freeman 1984). Accordingly, morphological differences between congeneric species are usually

negligible. Bones of congeneric species use appear separable from each other if very few bones are at disposal (e.g. Bocheński & Tomek 1995), but the alleged differences disappear when large series of bones are compared (Mlíkovský orig., see also Warheit 1992). If an author described many morphological differences between two congeneric species, then it is probable, that he or she underestimated the intraspecific variability and/or that one of the species is misclassified as to the genus and/or that he or she did not consider the allometrical variability.

Generally, I believe that most congeneric species of birds cannot be properly discerned on the basis of postcranial bones. Hence, two or more congeneric and sympatric species, which are similar in body size, may remain undetected in fossil samples.

Fossil and modern species: a comparison

The fossil record offers much less characters useful for the recognition of species than modern specimens or even living birds. In particular, data on plumage color, song, breeding habits etc. are not available for fossil birds, and useful molecular data can be obtained only from late Quaternary specimens. Unlike in modern birds, it is thus highly probable that sympatric species, which fall into the same size class, will not be discerned in paleontological studies, especially when only a few bones or even bone fragments are available. Similarly, it is impossible to exactly discern allopatric species. These problems are the more important, that a large proportion of modern species (in the broad sense of the 1970s) belongs to superspecies (Vuilleumier 1976, see also Stepanjan 1983, Haffer 1989), and that a large proportion of speciation events occurs sympatrically (Chesser & Zink 1994). Hence, the fossil species resembles probably most closely the superspecies of Schilder (1952; see also Kiriakoff 1967) or the species group of Haffer (1986), which includes closely related species without respect whether their ranges overlap or not. Also, it approaches the very narrow genus concept of Wolters (1971). For further discussion of the species problem in paleontology see e.g. Simpson 1943, 1950, 1961, Sylvester-Bradley 1956, Imbrie 1957, Shaw 1969, Reif 1984, Mlíkovský et al. 1985, Remane 1985, and Haffer 1995.

Subspecific categories

A few paleornithologists described fossil subspecies. This is not substantiated, especially with respect to the unclear definition of fossil species (see above). Naming fossil subspecies thus only inflates the number of names. No subspecific taxa are recognized in this paper.

Extinction and extermination

While the origin of taxa is a frequently discussed issue, the end of their existence is not. In the present paper I recognize the notions of *extinction*, where the phyletic line (at the levels of families, genera, and species) ceased to exist, and *extermination*, where the phyletic line either changed into another taxon of the same rank during a speciation or a macroevolutionary event, or when it lumped with another phyletic line (see e.g. Rhymer & Simberloff 1996 for the latter possibility; valid only at the species level).

Psychological factors in taxonomic decisions

Psychological factors influence taxonomic decisions to a surprisingly large degree (Mlíkovský, in prep.). In particular:

- ❑ Larger bones tend to be more often separated at the species level, than small bones, i.e. the perception of differences is size-dependent.
- ❑ Aberrant bones, i.e. those laying near the border of a phenon, tend to be selected as holotypes if more similar bones are available, especially when no study of variability is done (see e.g. the position of holotypes of four alleged *Necrobyas* species in the figures in Mlíkovský 1998c).

- ❑ For some authors it is so tempting to describe new taxa, that they tend to underestimate the intraspecific variability, and base their new taxa on indeterminate scraps of bones.
- ❑ Strong support is generally required for newly created taxa (i.e. the type I error is given preference), but equally strong support is required for synonymizing taxa (i.e. the type II error is given preference). Both types of errors should be considered in both cases.
- ❑ Researchers studying fossil birds from younger strata tend to describe new taxa on insufficient material, when they receive for study bird bones from older strata. This is especially marked, when students of Quaternary (especially late Quaternary) birds start to describe Tertiary bird remains.
- ❑ Modern birds from Plio-Pleistocene deposits are often identified on the "in-this-area-it-must-be-that-species" basis (see Tyrberg 1998: 518 for this apt term). This principle is often used also for older birds, where bones are assigned to previously described taxa on the "in-this-time period-it-must-be-that-species" basis. Although this approach is understandable to some degree, it obscures the distributional patterns of taxa both in time and space.

Although taxonomic decisions should be as exact as possible, the scarcity of material often requires, that the sixth sense is used. The psychological factors listed above limit the use of the sixth sense, and should be considered in evaluating the validity of taxonomic conclusions.

Classification of birds

The evolution of avian classifications has a long history (see Gadow 1893 and Sibley & Ahlquist 1990 for summaries). Early classifications were based on the general appearance of the animals, closely resembling the pre-scientific ethnobiological classifications (see Mlíkovský 1983a, Berlin 1992). Subsequent research modified these classifications using less apparent (anatomical, molecular, etc.) characters, in attempt to make the classification more "natural". I understand here as natural such a classification, which reflects both the adaptive radiation and phylogenetic relationships of the taxa (see also Mlíkovský 1979, 1983a, 1987, and the above section of this paper).

The 20th century understanding of the avian classification evolved in the 19th century, when fossil birds were too insufficiently known to have any influence on the classification. Almost all the work was done in Europe, culminating with the papers by Fürbringer (1888, 1902), Gadow (1891, 1892, 1893), and Beddard (1898). In the early 20th century, the research in avian systematics moved from Europe to the USA, where Wetmore (1930b) transformed the results of the above mentioned comparative morphologists and systematists in a new, simple and lucid classification of birds. Subsequently, the Wetmorean classification of birds became the standard for the 20th century (Wetmore 1930b, 1934b, 1940b, 1951, 1960, see also Hachler 1944, Mayr & Amadon 1951, Stresemann 1959, Storer 1960, 1971, Bock 1992). The Wetmorean classification of birds was used also in the last two catalogues of fossil birds (Lambrecht 1933, Brodkorb 1963, 1964, 1967, 1971, 1978), though not in all details.

The advancements in avian systematics (with special respect to fossil birds) showed that the Wetmorean classification "amounts to little more than superstition and bears about as much relationship to a true phylogeny of the Class Aves as Greek mythology does to the theory of relativity" (Olson 1981b: 193). Starting in the 1960s, several alternative classifications of birds were published, based on different data sets and using different methodologies (Verheyen 1961, Wolters 1975-1982, Cracraft 1981, Mlíkovský 1982, 1985, Olson 1985, Sibley & Ahlquist 1988, 1990, Livezey & Zusi 2001). These classifications differ from each other as well as from the Wetmorean "standard" classification, but none was generally accepted. In view of this, Mayr & Bock (1994) suggested to use the Wetmorean classification until another classification of birds will be generally accepted. Nevertheless, the Wetmorean orders still resemble the ethnobiological classification, based on the overall similarity (see above), and some of the early Tertiary taxa of birds cannot be placed in this classification, because they

represent evolutionary links between families formerly placed far away from each other, and in different orders.

Hence, the birds are arranged here according to a modified version of my classification of birds (Mlíkovský 1982, 1985). Only European taxa are considered here, although all taxa of birds were taken into account during the preparation of this classification (Mlíkovský, in prep.). In constructing the classification, all types of data were used, ranging from morphological over molecular to behavioral and ecological ones. The structure of the telencephalon is considered especially important, because it separates the Cenozoic birds into four distinct groups (Stingelin 1958a,b, 1960, Mlíkovský 1977, and unpublished data), which do not follow the ecomorphological classification in the Wetmorean orders, but seem to excellently indicate groups, within which adaptive radiation took place (Mlíkovský, in prep.). This arrangement is supported by an enormous amount of further data, which will be presented elsewhere (Mlíkovský, in prep.). In spite of this, the taxonomic position of some taxa remains uncertain. Below, I present an abbreviated version of the new classification of birds, limited to the taxa recorded from the Cenozoic of Europe:

Class Aves

Superorder Struthioni

- Order Tinamiformes: Tinamidae
- Order Struthioniformes: Palaeotididae, Struthionidae

Superorder Gavii

- Order Phaethontiformes: Gaviidae, Phaethontidae
- Order Ardeiformes: Sulidae, Ardeidae, Phalacrocoracidae, Anhingidae
- Order Bucerotiformes: Laurillardidae, Bucerotidae, Upupidae, Phoeniculidae

Superorder Ciconii

- Order Procellariiformes: Procellariidae, Pelagornithidae
- Order Ciconiiformes: Podicipedidae, Pelecanidae, Messelornithidae, Gruidae, Threskiornithidae, Gastornithidae, Eogruidae, Otididae, Ciconiidae, Cathartidae,
- Order Anseriformes: Phoenicopteridae, Recurvirostridae, Anhimidae, Anatidae
- Order Charadriiformes: Jacanidae, Rostratulidae, Scolopacidae, Glareolidae, Thinocoridae, Pterocletidae, Charadriidae, Laridae, Alcidae
- Order Piciformes: Coliidae, Psittacidae, Ramphastidae, Picidae

Superorder Passeri

- Order Galliformes: Megapodiidae, Cracidae, Phasianidae
- Order Accipitriformes: Rallidae, Cariamidae (incl. Phorusrhacidae), Cuculidae (incl. Musophagidae), Accipitridae (incl. Sagittariidae)
- Order Columbiformes: Caprimulgidae (incl. Archaeotrogonidae, Nyctibiidae, Pterocinidae), Aegialornithidae, Apodidae, Podargidae, Sophiornithidae, Strigidae, Columbidae, Falconidae
- Order Coraciiformes: Sylphornithidae, Coraciidae, Trogonidae, Halcyornithidae, Alcedinidae, Todidae, Meropidae
- Order Passeriformes: several modern families.

The Struthioni is an ancient group of unknown systematic position. They probably represent remains of an early radiation of birds. All known forms are terrestrial and omnivorous. The adaptive radiation is very limited.

The Gavii is another ancient group of birds (see Olson 1992). The ancestral forms seem to have been aquatic, predatory birds, with limited attempts at colonizing dry ground (Ardeidae). A single offshoot adapted to the arboreal life habits (Bucerotiformes). The adaptive radiation is rather limited.

The Ciconii is a large group of birds, whose adaptive radiation probably took place mainly in the Paleogene. The ancestral forms seem to have been aquatic, predatory birds, but they

were successful in adapting to terrestrial life habits, probably in several independent attempts (various Ciconiiformes). They also evolved filter-feeding (primary adaptation of the Anseriformes), and an offshoot adapted to the arboreal life habits (Piciformes). Most of the Ciconii radiated before the Oligocene. The Anseriformes originated in the Eocene, but most of their adaptive radiation probably took place in the Oligo-Miocene. The adaptive radiation is broad. The delimitation and arrangement of families depend on the still insufficiently known data on the Paleocene and Eocene forms.

The Passeri is a large group of primarily terrestrial/arboreal, predatory birds. Their adaptive radiation took place mainly in the Paleogene, prior to the Oligocene. The Passeriformes possibly also originated in the Eocene (see Boles 1997), but most of their adaptive radiation took place later, probably in the Oligo-Miocene. The Passeri radiated into a great variety of adaptive zones, but only very few entered the aquatic ones. The taxonomic position of the Sylphornithidae and Coraciidae is uncertain.

List of collections

BaM	Naturhistorisches Museum, Basel, Switzerland
BMNH	Department of Paleontology, Natural History Museum, London, England [formerly British Museum (Natural History)]
BSP	Bayerische Staatssammlung für Paläontologie und Historische Geologie, München, Germany
CHI	Conservation Hall, Ipolytarnóc, Hungary
CMP	Carnegie Museum, Pittsburg, Pennsylvania, USA
DPFNSP	Institute of Geology and Paleontology [formerly Department of Paleontology], Faculty of Natural Sciences, Charles University, Praha, Czechia
FSL	Centre des Sciences de la Terre, Université Claude-Bernard, Lyon, France
FSUM	Faculté des Sciences, Université de Marseille, France
GIB	Geological Institute, Budapest, Hungary
GIH	Geologisches Institut, Hamburg, Germany
GIK	Geological Institute, Ukrainian Academy of Sciences, Kyjiv, Ukraine
GIKM	State Museum of National History, Chişinău, Moldova
GMH	Geiseltalmuseum, Halle an der Saale, Germany
GMK	Museum of Geology, Institute of Geological Sciences, Ukrainian Academy of Sciences, Kyjiv, Ukraine
GPIW	Geologisch-Paläontologisches Institut, Universität Würzburg, Würzburg, Germany
GUN	Novorosijsk State University, Novorosijsk, Ukraine
HLMD	Hessisches Landesmuseum für Naturkunde, Darmstadt, Germany
HUJ	Hebrew University of Jerusalem, Jerusalem, Israel
IGF	Museum of Geology and Paleontology, University of Firenze, Firenze, Italy
IGS	Institute of Geological Sciences [formerly Geological Survey Museum], London, England
IGGB	Institute of Geology and Geophysics, Bucureşti, Romania
IGGL	Institute of Geology and Geochemistry of Fossil Fuels, Ukrainian Academy of Sciences, L'vyv, Ukraine
IM	Ipswich Museum, Ipswich, England
IPS	Institute of Paleontology, Sabadell, Spain
IRScNB	Institut Royal des Sciences Naturelles, Bruxelles, Belgium
ISEZ	Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Kraków, Poland
IQW	Indtitut für Quartärpaläontologie, Weimar, Germany
IZAN	Institute of Zoology, Ukrainian Academy of Sciences, Kyjiv, Ukraine
IZC	Institute of Zoology, Moldovan Academy of Sciences, Chişinău, Moldova
LMB	Landesmuseum Braunschweig, Germany
LPLN	Laboratoire de Préhistoire du Lazaret, Nice, France

LPUB	Laboratory of Paleontology, University of București, București, Romania
LPUI	Laboratory of Paleontology, University of Iași, Iași, Romania
MB	Natural History Museum, Baku, Azerbaijan
MCI	Museum of Imola, Imola, Italy
MG	Landesmuseum Joanneum, Graz, Austria
MGPUB	Museum of Geology and Paleontology, University of Bologna, Bologna, Italy
MGPOT	Museum of Geology and Paleontology, University of Torino, Torino, Italy
MHNA	Musée d'Histoire Naturelle, Aix-en-Provence, France
MHNB	Muséum d'Histoire Naturelle, Bordeaux, France
MHNL	Muséum d'Histoire Naturelle, Lyon, France
ML	Musée Guimet d'Histoire Naturelle, Lyon, France
MMB	Anthropos Institute, Moravian Museum, Brno, Czechia
MNCN	National Museum of Natural Sciences, Madrid, Spain
MNHM	Muséum d'Histoire Naturelle, Montauban, France
MNHN	Laboratoire de Paléontologie, Muséum National d'Histoire Naturelle, Paris, France
MNHNS	National Museum of Natural History, Sofija, Bulgaria
MP	Museum of Petralona, Petralona, Greece
MPC	Musée de Paléontologie, Chilhac, France
MPUP	Museum of Paleontology, University of Padova, Padova, Italy
MS	Musée de Sartène, Corsica, France
MSNUP	Natural History Museum, University of Pisa, Pisa, Italy
NHMW	Geologisch-Paläontologische Abteilung, Naturhistorisches Museum, Wien, Austria
NKMB	Museum für Naturkunde, Humboldt-Universität, Berlin, Germany
NMB	National Museum, Budapest, Hungary
NMM	National Museum of Natural History, Valletta, Malta
NMP	Department of Paleontology, National Museum, Praha, Czechia
NMZ	Naturhistorisches Museum, Zürich, Switzerland
NRM	Department of Paleontology, Natural History Museum, Stockholm, Sweden
PIN	Paleontological Institute, Russian Academy of Sciences, Moskva, Russia
PINK	Paleontological Institute, Ukrainian Academy of Sciences, Kyjiv, Ukraine
RGM	State Museum of Geology and Paleontology, Leyden, Holland
RMO	Regional Museum, Oradea, Romania
RMT	Regional Museum, Teplice, Czechia
SMC	Sedgwick Museum [formerly Woodwardian Museum], Cambridge, England
SMF	Naturmuseum und Forschungsinstitut Senckenberg, Frankfurt am Main, Germany
SMNK	Staatliches Museum für Naturkunde, Karlsruhe, Germany
SMNS	Staatliches Museum für Naturkunde, Stuttgart, Germany
TGPI	Tiraspol State Pedagogical Institute, Tiraspol, Moldova
ULg	Université de Liège, Liège, Belgium
UNL	New University, Lisboa, Portugal
USNM	United States National Museum, Washington, D.C., USA
USTL	Université des Sciences et Techniques de Languedoc, Montpellier, France
UUT	University of Utrecht, Utrecht, Holland
UWPI	Institut für Paläontologie, Universität Wien, Wien, Austria
WDC	Wyoming Dinosaur Center, Thermopolis, Wyoming, USA
ZGIB	Zentrales Geologisches Institut, Berlin, Germany
ZIN	Zoological Institute, Russian Academy of Sciences, Sankt Peterburg, Russia

Private collections:

- coll. Aymard, France (present location unknown)
- coll. Barbey (present location unknown)
- coll. Bonifay (Lyon, France)
- coll. Croizet (part in BMNH, present location of the rest unknown)
- coll. Danninger, Allerding, Austria

coll. Delfortrie (present location unknown)
 coll. Dvořák, Bílina, Czechia
 coll. Fejfar, Praha, Czechia
 coll. Keferstein (present location unknown)
 coll. Kessler (Germany)
 coll. Kormos, Budapest, Hungary (probably in NMB)
 coll. Maschwitz, Germany
 coll. Michel, France (to be deposited in Musée d'Histoire Naturelle in Montbéliard, France)
 coll. Nouel, France (present location unknown)
 coll. Paupe, France (to be deposited in Musée d'Histoire Naturelle in Montbéliard, France)
 coll. Rummel, Weissenburg, Germany
 coll. Villalta, Spain (present location unknown)

Acknowledgments

Great number of persons kindly provided me with information necessary for the preparation of this paper and/or allowed me to work in collections under their care. I am much obliged to all of them. They include in ARGENTINA: Eduardo P. Tonni (Buenos Aires); in AUSTRALIA: Walter E. Boles (Sydney), Patricia Vickers-Rich (Clayton); in AUSTRIA: Gudrun Daxner-Höck (Wien), Florian A. Fladerer (Wien), Gernot Rabeder (Wien), Karl Rauscher (Wien), Ortwin Schultz (Wien); in AZERBAIJAN: Nikolaj Ivanovič Burčak-Abramovič (Baku); in BRAZIL: Herculano Maros Ferraz de Alvarenga (Taubaté); in BULGARIA: Zlatozar Boev (Sofija); in CHINA: Lian-Hai Hou (Beijing); in CROATIA: Vesna Malez-Bačić (Zagreb), in CZECHIA: Petr Benda (Praha), Josef Beneš (Praha), Ivana Boháčová (Praha), Dana Ernygrová (Teplice), Oldřich Fejfar (Praha), Ivan Horáček (Praha), Renata Mlíková (Praha), Rudolf Musil (Brno), Radko Samek (Nymburk), Luděk Seitl (Brno), Petr Stýblo (Praha), Petr Švec (Praha); in DENMARK: Ella Hoch (Kjøbenhavn); in FRANCE: Eric Buffetaut (Paris), Jacques Cheneval (Lyon), Colette Dechaseaux (Paris), Pierre Fiquet (Paris), Daniel Goujet (Paris), Christine Lefèvre (Paris), Cécile Mourer-Chauviré (Lyon), Eric Pelle (Paris); in GERMANY: Peter Ballmann (Bonn), Siegfried Eck (Dresden), Volker Fahlbusch (München), Karlheinz Fischer (Berlin), Ursula B. Göhlich (München), Hartmut Haubold (Halle an der Saale), Wolf-Dieter Heinrich (Berlin), Kurt Heissig (München), Angelika Hesse (Dessau), Erlend Martini (Stuttgart), Günther Mauersberger (Berlin), Lutz Maul (Weimar), Gerald Mayr (Frankfurt am Main), Dieter Stephan Peters (Frankfurt am Main), Rudolf Piechocki (Halle an der Saale), Harald Pieper (Kiel), Burkhard Stephan (Berlin); in HOLLAND: A. W. Janssen (Leiden), Peter D. M. Weesie (Leiden); in HUNGARY: Dénes Jánossy (Budapest); in ITALY: Laura Delle Cave (Firenze), Marco Pavia (Torino); in NEW ZEALAND: Ron J. Scarlett; in POLAND: Zbigniew Bocheński (Kraków), Zygmunt Bocheński (Kraków), Andrzej Elżanowski (Warszawa), Tereza Tomek (Kraków); in PORTUGAL: Cornelis J. Hazevoet (Lisboa); in ROMANIA: Erika Gál (Cluj), Eugen Kessler (Cluj); in RUSSIA: A. O. Aver'janov (Sankt Peterburg), Gennadij F. Baryšnikov (Sankt Peterburg), Aleksandr A. Karchu (Moskva), Evgenij Nikolaevič Kuročkin (Moskva); in SLOVENIA: Tea Kolar-Jurovšek (Ljubljana); in SPAIN: Josep Antoni Alcover (Madrid), Antonio Sánchez Marco (Madrid); in SWEDEN: S.-A. Bengtson (Stockholm), Per G. P. Ericson (Stockholm), Tommy Tyrberg (Kimstad); in SWITZERLAND: B. Engesser (Basel), Marc Weidmann (Zürich); in the UNITED KINGDOM: Colin James Oliver Harrison (London), Gillian & Vladimír Horálek (Swansea), E. Marjorie Northcote (Cambridge), Cyril A. Walker (London); in the USA: Allison Victor Andors (New York, NY), Jonathan J. Becker (Washington, DC), Walter J. Bock (New York, NY), Pierce Brodkorb (Tallahassee, FL), Kenneth E. Campbell (Los Angeles, CA), Charles T. Collins (Long Beach, CA), Joel Cracraft (Chicago, IL), Alan Feduccia (Chapel Hill, NC), Peter Houde (Washington, DC), Hildegard Howard (Los Angeles, CA), Helen F. James (Washington, DC), Bradley C. Livezey (Pittsburg, KS), Larry D. Martin (Lawrence, KS), Storrs L. Olson (Washington, DC), Pamela C. Rasmussen (East Lansing, MI), David W. Steadman (Tallahassee, FL), Robert W. Storer (Ann Arbor, MI).

History of the research

Avian paleontology evolved within the past 300 years and the research was done exclusively by European scientists in the first one-and-half century. In the 18th century, the research was limited to Switzerland, France and Germany, but it slowly spread over the Europe in the course of the 19th century, mainly in its second half. The history of research can be roughly separated into five periods, which are detailed below.

1700-1800: The understanding of fossils was weak, and they were usually explained as remnants of animals perished in the Biblical Deluge (see e.g. Zittel 1899, Rudwick 1972). The knowledge of avian morphology was negligible. Accordingly, most of the alleged records of fossil birds turned out to be based on non-avian material (see Cuvier 1822: 302-305, Krüger 1823: 709-715, and Bronn 1834 for reviews). Proper avian fossils were reported only from the Miocene of Öhningen, then in Switzerland (Scheuchzer 1708a,b², 1716, Razoumowsky 1789³), and from the Eocene of Montmartre in France (Lamanon 1782, 1783⁴, Camper 1786).




Fig. 2. An unidentified avian feather from the middle Miocene (MN 6) of Öhningen, Germany: the first record of a fossil bird from Europe (after Scheuchzer 1708a).

1800-1860: Accumulated evidence resulted in the abandonment of the single Deluge theory, in the formulation of the theory of multiple, more or less universal floods (e.g. Parkinson 1808-1811, Cuvier 1812), and finally in the formulation of the theory of evolution (Darwin 1859); see e.g. Zittel (1899), Glass (1959) and Laurent (1987). Avian paleontology started to develop, but remained to be limited to the western Europe (England, France, Germany, Switzerland, and marginally also Italy), although local paleontologists started to receive (in the 1820s) bones of fossil and subfossil birds also from "exotic" islands, particularly Sardinia (Wagner 1828, 1831, 1832, Marmora 1831), and later (in the 1830s and 1840s) New Zealand, Mauritius and Madagascar (see Lambrecht 1933). First comprehensive paper in avian paleontology is the description of the Eocene birds from Montmartre, France, by Georges Cuvier (1822). At the same time, English naturalist William Buckland (1822, 1823) published first account on the late Quaternary birds, describing their remains from the Kirkdale cave in England (see Harrison 1980c for their review). In 1825, German naturalist Carl Dietrich Eberhard König (1825) named first fossil bird: *Larus toliapicus*⁵ from the Eocene of the Isle of

² Scheuchzer's specimen (a feather imprint) was figured by Scheuchzer (1708a,b, pl. 2), Peyer (1957, fig. 5-6), and here (fig. 2). The original is deposited in NMZ (Peyer 1957).

³ Razoumowsky's specimen (a hindlimb in slab) was figured by Karg (1805, pl. 2, fig. 1) and is deposited in BMNH under the catalogue number A-152 (Lydekker 1891a: 172).

⁴ Lamanon's specimen is figured in Cuvier (1822, pl. 73, fig. 1).

⁵ *Bucklandium diluvii*, another alleged bird named by König (1825), turned out to be a fish (see below).

Sheppey, England, but naming of avian fossils started to be more common only in the 1840s with the work of Richard Owen (1840, 1841, 1846) in England, Paul Gervais (1844a,b, 1852) in France, and Christoph Gottlieb Andreas Giebel (1847) in Germany. Simultaneously, the knowledge of the comparative osteology of birds developed (Nitzsch 1811, Merrem 1816, Blainville 1821)⁶. Significant contributions to the avian paleontology were published particularly by Georges Cuvier (1769-1832) and Paul Gervais (1816-1879) in France, by Hermann von Meyer (1801-1869) and Christoph Gottlieb Giebel (1820-1881) in Germany, and by Richard Owen (1804-1892) in England.

1860-1935: With the establishment of the theory of evolution (Darwin 1859), the meaning of fossils changed, although the theory itself had to be improved from the paleontological point of view (e.g. Owen 1860, Lyell 1863, Rolle 1863; see also Zittel 1899, Davitašvili 1948, Wilson 1970, Rudwick 1972, Bowler 1976). New standards in avian paleontology were set in the 1860s and 1870s especially by the continuing work of Richard Owen in England, and by Alphonse Milne-Edwards (1835-1900) in France with his treatise *Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France* (see also d'Archiac et al. 1866). The science of paleornithology slowly spread over all the Europe, and became established also in New Zealand and in the Americas (see Lambrecht 1933 for the authors).

Toward the end of the 19th century, the knowledge of the comparative morphology of birds, incl. osteology, markedly increased (see e.g. Fürbringer 1888, Selenka 1891, Gadow 1891, Beddard 1902), which in turn supported the development of paleornithology. Although many authors contributed to the research in this time period, the most significant ones appear to be Richard Lydekker (1849-1915) in England, Claude Gaillard (1861-1946) in France, and Kálmán Lambrecht (1889-1936) in Hungary. This period culminated with the *Handbuch der Palaeornithologie* by Lambrecht (1933).

1935-1970: This is the dark age of the European paleornithology, the onset of which coincides with the economical crisis which struck the developed countries of Europe in the late 1930s, and with the subsequent World War II (1939-1945). In these decades, Europe lost its position of the world leader in avian paleontology, leaving it for the USA. Consistent interest in the Cenozoic birds of Europe started to show in this period only Nikolaj Iosifovič Burčák-Abramovič (1939 ff.) in the then Soviet Union, and Miklós Kretzoi (1941 ff.) and Dénes Jánossy (1954 ff.) in Hungary, although all of them were primarily interested in fossil mammals.

1970-2000: Renaissance of the European paleornithology. This era brought several important novelties, generally imported from the USA, incl. (1) efforts at revising taxa described by earlier researchers, (2) world-wide reviews of selected taxa, (3) use of much improved comparative collections of modern avian skeletons, which allow for a comparison of fossil bones with the corresponding elements of previously unavailable taxa, (4) application of biometrical studies, and (5) departure from the geological understanding of the avian paleontology toward the zoological one.

⁶ A much earlier study by Coiter (1575) fell into oblivion (Allen 1951, Stresemann 1951: 47).

List of localities

The following list contains all Tertiary and early Pleistocene localities of Europe, which yielded remains of birds, and of which I became aware. Younger localities are listed only if they are mentioned in the text. Details on the localities can be found in Mlíkovský (1996a) and Tyrberg (1998, 2001). The list is arranged chronologically, and within periods alphabetically. All Tertiary localities are limited to a single mammal zone. If more zones are indicated for a given locality (e.g. MP 7-10), then its exact age is unknown, being somewhere within the limits. On the other hand, Quaternary localities use to be multi-layered, which is indicated below. Note, that the biochronological position of localities may be subject of change, caused by continuing research.

For each locality, its general geographic position is given, followed – in parentheses – with the reference to the catalogues by Mlíkovský (1996a) and Tyrberg (1998, 2001). In addition, references to relevant paleornithological papers are listed in each case (simple mentioning of birds from the locality being not considered). In brackets, the type of material recovered is indicated. The abbreviations are as follows: B – bones, E – eggshells, F – feather imprints, and T – tracks (footprints). An asterisk indicates that one or more avian species were created on the basis of the material.

MP 1-5

Gelinden, Belgium (Cheneval 1996a, # 1): Misonne 1958. [B]

Limhamn, Sweden (Tyrberg & Ericson 1996, # 1): Dames 1890, Lambrecht 1933, Olson & Feduccia 1980a, Tyrberg & Ericson 1996. [B*]

Selk, Germany (new): Lierl 1993. [B]

MP 5

Menat, France (Mourer-Chauviré 1996a, # 1): Launay 1908, 1923, Pitton 1940. [B]

Sundby, island of Mors, Denmark (new): Hoch 1997. [B]

MP 6

Cernay-les-Reims, France (Mourer-Chauviré 1996a, # 2): Gervais 1873, 1877, Lemoine 1878a, 1880, 1881a,b,c, 1884, 1885, 1893, Meunier 1882, Repelin 1924, Andors 1992, Martin 1992, Mourer-Chauviré 1994, 1995a. This is a complex locality, which includes the sites of Chalons-sur-Vesle, Berru and Rilly. [B*]

Maret, Belgium (Cheneval 1996a, # 2): Misonne 1958. [B]

Mesvin, Belgium (Cheneval 1996a, # 3): Dollo 1883, 1909, Martin 1992. [B]

Walbeck, Germany (Mlíkovský & Hesse 1996, # 1): Weigelt 1939, 1942. [B]

MP 7

Basse Provence, France (Mourer-Chauviré 1996a, # 7): Dughi & Sirugue 1959, 1962, 1968, Fabre-Taxy & Touraine 1960, Touraine 1960, 1961, 1969, Rey 1969, Cailleux 1969, Kerourio & Aujard 1987, Michajlov 1991, Dauphin 1991, 1992, 1994. This is a complex locality, which includes the sites of Baraque, Cengle, Mauquiers, Pontevès, Saint-Antonin, Saint-Julien-le-Montagnié, Saint-Maurin, and Sillans-la-Cascade. [E*]

Croydon, England (Mlíkovský 1996q, # 1): Newton 1885a,b, 1886a,b, 1889, Harrison & Walker 1977a, Martin 1992. [B*]

Dormaal, Belgium (Cheneval 1996a, # 5): Misonne 1958. Smith & Smith (1995) did not list birds from this locality. [B]

Fur, Denmark (Mlíkovský 1996f, # 1): Hoch 1975, Houde & Olson 1992, Kristoffersen 1998, 1999. [B]

Meudon, France (Mourer-Chauviré 1996a, # 3): Prévost 1855a,b, Hébert 1855a,b,c,d, Lartet 1855, Valenciennes 1855, Duméril 1855, Owen 1856a,b, Milne-Edwards 1867a,b, 1872, Meunier 1882, Woodward 1886, Lavocat 1899, Repelin 1924, Russell et al. 1990, Martin 1992. [B*]



Fig. 3. Type localities of Paleocene (MP 1-6) birds of Europe. 1 - Limhamn, 2 - Cernay-les-Reims. Scale bar = 100 km.

Passy, France (Mourer-Chauviré 1996a, # 4): Milne-Edwards 1867a,b, 1872. [B]

Silverinha, Portugal (Sánchez Marco 1996a, # 1): Harrison 1983a. [B*]

MP 7-10

Aude, France (Mourer-Chauviré 1996a, # 8a-c): Plaziat 1964, Villatte 1966, Dughi et al. 1969, Cailleux 1969, Kerourio & Aujard 1987. This is a complex locality, which includes the sites of Gallié, Auzières, and Serre de Trigodinna. [E]

Erquelines, Belgium (Cheneval 1996a, # 4): Misonne 1958. [B]

Grange Farm, England (Mlíkovský 1996q, # 9): George & Vincent 1978, Harrison 1980a, 1983b. [B*]

Havighorst, Germany (Mlíkovský & Hesse 1996, # 3): Illies 1941, 1942. [F]

High Ongar, England (Mlíkovský 1996q, # 7): Harrison & Walker 1977a, Harrison 1984b, Houde 1988. [B*]

Katharinenhof, Germany (Mlíkovský & Hesse 1996, # 2): Hoch 1975. [B, brain casts]

Orp-le-Grand, Belgium (Cheneval 1996a, # 6): Misonne 1958. [B]

Primrose Hill, England (Mlíkovský 1996q, # 5): Owen 1846, Lydekker 1891a, Harrison & Walker 1977a, 1978a, Harrison 1979a. [B*]

Vinalmont, Belgium (Cheneval 1996a, # 7): Misonne 1958.

MP 8

Mariager Fjord, Denmark (Mlíkovský 1996f, # 2): Hoch 1975. [B, F]

Røsnaes, Denmark (Mlíkovský 1996e, # 3): Harrison 1984, Houde 1988. [B*]

Walton-on-the-Naze, England (Mlíkovský 1996q, # 11): Daniels 1979, 1987, 1988, 1989, 1990, 1991, 1993, 1994, 1997, 1999, Harrison 1982b,c, 1983b, 1984b,c, Houde 1988, Olson & Daniels 1994, Mayr 1998c, Mayr & Daniels 1998, Olson 1999a,b. [B*]



Fig. 4. Type localities of early Eocene (MP 7-10) birds of Europe. 1 - Croydon, Primrose Hill, 2 - Burnham-on-Crouch, Grange Farm, High Ongar, Walton-on-the-Naze, 3 - Abbey Wood, Herne Bay, Sheppey, 4 - Bognor Regis, 5 - Meudon, 6 - Monthelon, 7 - Silverinha, 8 - Basse Provence, 9 - Røsnaes. Scale bar = 100 km.

MP 8-9

Abbey Wood, England (Mlíkovský 1996q, # 2): Harrison & Walker 1977a. [B*]

Bognor Regis, England (Mlíkovský 1996q, # 4): Harrison & Walker 1975, 1977a, Olson & Feduccia 1979, Harrison 1982a, 1984a, Olson 1985, Baird & Vickers-Rich 1997. [B*]

Burnham-on-Crouch, England (Mlíkovský 1996q, # 6): Harrison & Walker 1977a, Olson & Feduccia 1979, Harrison 1982a, Martin & Mengel 1984, Baird & Vickers-Rich 1997, Olson 1999b: 127. [B*]

Herne Bay, England (Mlíkovský 1996q, # 10): Harrison & Walker 1977a, Harrison 1984b, 1985a. [B*]

Mutigny, France (Mourer-Chauviré 1996a, # 5): Martin 1992. [B]

Saint James Park, England (Mlíkovský 1996q, # 8): Harrison & Walker 1977a, Harrison 1984b. [B]

Sheppey, England (Mlíkovský 1996q, # 3): König 1825, Owen 1840, 1841, 1846, 1869, 1870, 1873, 1878, 1880, Bowerbank 1854, Seeley 1866, 1869, 1874, Martin 1874, Lydekker 1891a, 1896, Andrews 1899, Lambrecht 1933, Harrison & Walker 1971, 1972, 1975, 1976b,c, 1977a, Cracraft & Rich 1972, Harrison 1982a, 1984a,b,c, 1986, Houde 1988. [B*]

MP 10

Monthelon, France (Mourer-Chauviré 1996a, # 6): Schaub 1929a,b, Gaillard 1936, Andors 1992. [B*]

MP 11

Geiseltal XIII, Germany (Mlíkovský & Hesse 1996, # 5c): Fischer 1962, 1978. [B*]

Geiseltal XIV, Germany (Mlíkovský & Hesse 1996, # 5d): Fischer 1962, 1978. [B*]




Fig. 5. Type localities of middle Eocene (MP 11-13) birds of Europe. 1 - Lee-on-Solent, Yateley, 2 - Geiseltal (incl. Cecilie), 3 - Messel, 4 - Egerkingen, 5 - Monte Zuello, 6 - Cluj-Manastur. Scale bar = 100 km.

Geiseltal IL, Germany (Mlíkovský & Hesse 1996, # 5j): Mlíkovský, orig. [B]

Messel, Germany (Mlíkovský & Hesse 1996, # 4): Wittich 1899, Berg 1965, Hoch 1980, 1988, Peters 1983, 1985, 1987a,b, 1988a,b, 1989, 1991, 1992, 1994, 1995, 1997a,b,c, Houde 1988, Hesse 1988a,b, 1989, 1990, Eikamp 1990, Hesse & Habersetzer 1993, Peters & Storch 1993, 1994, 1995, Mayr 1997, 1998a,b,c,d, 1999b,c, 2000b,c,d,f, Mayr & Daniels 1998, Mayr & Peters 1998, 1999. [B*, F]

MP 11-13

Chamblon, Switzerland (Mlíkovský 1996o, # 2): Thenius 1959. [B]

Egerkingen, Switzerland (Mlíkovský 1996o, # 1): Schaub 1929b, 1940a,b, Mlíkovský 1992b. [B*]

Etterbeek, Belgium (Cheneval 1996a, # 8): Dollo 1909, Lambrecht 1933, Misonne 1958. [B]

Fontcouverte, France (Mourer-Chauviré 1996a, # 8d): Dughi & Sirugue 1962. [E]

Geiseltal, Germany (Mlíkovský & Hesse 1996, # 5): Lambrecht 1928, 1933, 1935, Heller 1932, Bachofen-Echt 1936a, Krumbiegel 1959, Fischer 1962, 1967, 1978, 1987, Matthes 1970, Haubold & Krumbiegel 1984, Houde 1986, 1988, Houde & Haubold 1987, Peters 1989, 1998, Mlíkovský 1992b, Kohring & Hirsch 1996, Mayr 2000e. This is a complex locality, which includes many fossiliferous sites. Three early sites are called Cecilie and Leonhard, respectively, while the newer ones Geiseltal. Vertebrate sites are numbered with Roman numbers (Krumbiegel 1959, 1962, 1966, 1968, 1977, Krumbiegel et al. 1983). [B*, E, F]

Lee-on-Solent, England (Mlíkovský 1996q, # 12): Harrison & Walker 1979a, Harrison 1982c. [B*]




Fig. 6. Type localities of late Eocene (MP 14-20) birds of Europe. 1 - Hordle, Milford, 2 - Montmartre, Romainville, 3 - Quercy (incl. Bouffie, Bretou, Escamps, Perrière, Rosières, Sainte-Néboule), 5 - Argentera, 6 - Verrua Savoia, 7 - Vestena Nova, 8 - Kyjiv. Scale bar = 100 km.

Meilleries-Saint-Sauveur, France (Mourer-Chauviré 1996a, # 11): Robertson 1834. This is a doubtful locality. The age may be late Cretaceous, and the coproliths (the only remains found) may have been produced by dinosaurs (see Mourer-Chauviré 1996a: 573). [Coproliths]

Monte Zuello, Italy (Delle Cave 1996, # 1): Zigno 1884, Portis 1884, Cracraft 1973. [B*]

Yateley, England (Mlíkovský 1996q, # 13): Harrison & Walker 1979a. [B*]

MP 12

Geiseltal L, Germany (new): Peters 1998 [B*]

MP 13

Bouxviller, France (Mourer-Chauviré 1996a, # 9): Kuntz 1983. [E]

Cecilie I, Germany (Mlíkovský & Hesse 1996, # 5a): Lambrecht 1928, 1933, 1935, Houde 1986, 1988, Houde & Haubold 1987, Mlíkovský 1992. [B*, E, F]

Cecilie II, Germany (Mlíkovský & Hesse 1996, # 5b): Weigelt 1933. [B]

Cluj-Manastur, Romania (Kessler 1996, # 1): Lambrecht 1929, Kessler 1986. [B*]

Geiseltal XX, Germany (Mlíkovský & Hesse 1996, # 5e): Fischer 1987, Mlíkovský 1992b. [B*]

Geiseltal XXII, Germany (Mlíkovský & Hesse 1996, # 5f): Mlíkovský, orig. [B]

Geiseltal XXXV, Germany (Mlíkovský & Hesse 1996, # 5g): Fischer 1962, 1978, 1987, Mlíkovský, orig. [B]

Geiseltal XXXVI (Mlíkovský & Hesse 1996, # 5): Fischer 1987, Mayr 2001e. [B]

Geiseltal XXXVII, Germany (Mlíkovský & Hesse 1996, # 5h): Mlíkovský, orig. [B]

Geiseltal XLI, Germany (Mlíkovský & Hesse 1996, # 5i): Fischer 1962, 1978, Mlíkovský, orig. [B]

Leonhard III, Germany (new): Lambrecht 1928, 1933. [B]

MP 14

Lissieu, France (Mourer-Chauviré 1996a, # 10): Gaillard 1936, 1937, 1939, Andors 1992. [B*]

MP 14-16

Barton, England (Mlíkovský 1996q, # 14): Harrison & Walker 1976a. [B*]

Highcliffe, England (Mlíkovský 1996q, # 15): Harrison & Walker 1976a. [B*]

Jantamyj, Russia (Mlíkovský 1996g, # 1): Berendt 1845, Giebel 1862, H. Meyer 1867, A.B. Meyer 1887, Andrée 1937, Bachofen-Echt 1929, 1936, 1944, 1949, Lambrecht 1933, Katinas 1983. This locality refers to a secondary deposit of the so-called Baltic amber, which originated in pine forests in the territory of the today's Finland (Katinas 1971, 1983). Washed-out pieces of fossiliferous amber can be found anywhere on the coasts of the eastern Baltic Sea. [F]

MP 14-20

Pin, France (Mourer-Chauviré 1996a, # 14): Meunier 1906. [T]

MP 16

Bretou, France (Mourer-Chauviré 1996a, # 12b): Mourer-Chauviré 1988a. [B*]

Lavergne, France (Mourer-Chauviré 1996a, # 12c): Mourer-Chauviré 1983, 1988b, 1989a, 1992b. [B]

MP 16-20

Garrigues, France (Mourer-Chauviré 1996a, # 12l): Mourer-Chauviré 1980, 1983. [B]

MP 16-28

Quercy, France (Mourer-Chauviré 1996a, # 12, 12a): Lydekker 1891a, Milne-Edwards 1892, 1893, Gaillard 1908, 1939, Lambrecht 1933, Lowe 1939, Brodkorb 1963b, 1964, 1967, 1971b, Cracraft & Rich 1972, Cracraft 1973, Collins 1976a, Jollie 1977c: 117, Olson 1977, 1985a, Mourer-Chauviré 1978, 1980a, 1981a, 1982a,b, 1983, 1985a,b, 1987, 1988a,b, 1989a, 1991, 1992a,b,c, 1993a, 1995a,b,c 1996a, Rich 1982, Mourer-Chauviré & Cheneval 1983, Harrison 1975c, 1984c, Hesse 1988, 1990, Karchu 1988, 1992, Fischer & Mauersberger 1989, Mlíkovský 1989a,b, 1998c, 1999d, Peters 1992, Sigé et al. 1998, Mourer-Chauviré et al. 1999, Mayr & Mourer-Chauviré 2000. This is a complex locality, consisting of numerous fissure fillings in a karstic area. Exact sites are unknown for the material from early excavations (henceforth called "Quercy"). Recently excavated sites are listed separately under the names Aubrelong, Baraval, Belgarric, Bouffie, Boussac, Bretou, Crabit, Desse, Escamps, Fonbonne, Fraysse, Garouillas, Garrigues, Gousnat, Itardiès, Lavergne, Lostanges, Mas de Got, Mounayne, Pécarel, Perrière, Phalip, Plante, Pradigues, Ravet-Lupovici, Roqueprune, Rosières, Sainte-Néboule, and Salême. [B*]

MP 17

Bouffie, France (Mourer-Chauviré 1996a, # 12e): Mourer-Chauviré 1983, 1988b, 1991, 1992b, 1995a, see also Mourer-Chauviré 1996a: 577. [B*]

Hordle, England (Mlíkovský 1996q, # 16): Lydekker 1891a, Storer 1956, Cracraft 1973, Olson 1977, 1981, Harrison 1971, 1976, 1986, Harrison & Walker 1976a, 1977a. [B*]

Milford, England (Mlíkovský 1996q, # 17): Lydekker 1891a, Harrison 1971, 1986, Harrison & Walker 1976a. [B*]

Perrière, France (Mourer-Chauviré 1996a, # 12f): Mourer-Chauviré 1980, 1983, 1987, 1988a,b, Peters 1992, Mlíkovský 1998c, Mayr & Mourer-Chauviré 2000. [B*]

Pradigues, France (Mourer-Chauviré 1996a, # 12d): Mourer-Chauviré 1988b. [B]

Salême, France (Mourer-Chauviré 1996a, # 12g): Mourer-Chauviré 1987, Peters 1992. [B]

MP 17-20

Argentera, Italy (Delle Cave 1996, # 4): Portis 1879, 1884. [T*]

Kyjiv, Ukraine (Mlíkovský 1996p, # 1): Rogovič 1875a,b, Aver'janov et al. 1990. [B*]

Verrua Savoia, Italy (Delle Cave 1996, # 3): Portis 1884, 1887, 1900. [T*]




Fig. 7. Type localities of early Oligocene (MP 21-22) and middle Oligocene (MP 23-24) birds of Europe. 1 - Hamstead, Yarmouth, 2 - Rupelmonde, 3 - Quercy (incl. Itardiès), 4 - Ronzon, 5 - Saignon, 6 - Céreste (incl. Lupéron?), 7 - Froidefontaine, 8 - Matt, 9 - Sieblos, 10 - Frauenweiler, 11 - Espenhain, 12 - Skalice, Varnsdorf, 13 - Cetaŕaie. Scale bar = 100 km.

Vestena Nova, Italy (Delle Cave 1996, # 2): Faujas-Saint-Fond 1801, Pictet 1853, Portis 1887, Omboni 1885. [F*]

MP 18

Gousnat, France (new): Mayr & Mourer-Chauviré 2000. [B]

Sainte-Néboule, France (Mourer-Chauviré 1996a, # 12h): Mourer-Chauviré 1978, 1992b. [B*]

MP 19

Escamps, France (Mourer-Chauviré 1996a, # 12i): Mourer-Chauviré 1978, 1980a, 1983, 1985b, 1988b, 1992b, 1999a, Karchu 1988, 1992). This is a complex locality, which includes also the sites Escamps III, Escamps IV, Escamps A, and Escamps C. [B*]

Lostanges, France (Mourer-Chauviré 1996a, # 12j): Mourer-Chauviré 1983. [B]

Möhren 6, Germany (Mlíkovský & Hesse 1996, # 6): Heissig 1970. [B]

Montmartre, France (Mourer-Chauviré 1996a, # 13): Lamanon 1782, 1783, Delamétherie 1800, 1802, Cuvier 1800a,b, 1807, 1809, 1822, 1825, 1836, Geoffroy-Saint-Hilaire 1842, Gervais 1844a,b, 1849, 1848-1852, 1859, Giebel 1847, Laurillard 1849, Reichenbach 1852, Bonaparte 1856c, Blanchard 1857, Desnoyers 1859a,b, Milne-Edwards 1863, 1867-1868, 1869-1871, 1872, Murie 1873, Woodward 1886, Flot 1892, Lydekker 1891a, Gaillard 1908, 1939, Oberholser 1917, Lambrecht 1933, Brodkorb 1963c, 1964, 1967, 1971b, 1978, Brunet 1970, Dechauseaux 1970a,b, Cracraft 1973, Olson 1974a, 1977, 1985a, Warter 1976, Harrison 1979b, Mlíkovský 1981, 1995a, Mayr 1998c, 2000a. [B*]

Pécarel, France (new): Mayr & Mourer-Chauviré 2000. [B]

Rosières, France (Mourer-Chauviré 1996a, # 12k): Mourer-Chauviré 1983, 1987, 1992b, 1996a. Mlíkovský 1998c, Mayr & Mourer-Chauviré 2000. This is a complex locality, which includes the sites Rosières 1 and Rosières 2. [B*]

MP 20

Frohnstetten, Germany (Mlíkovský & Hesse 1996, # 7): Meyer 1852: 305, Quenstedt 1853, Lambrecht 1933. [B]

Romainville, France (Mourer-Chauviré 1996a, # 15): Lebedinsky 1927, Brunet 1970, Harrison 1979a. [B*]

MP 21

Aubrelong 1, France (Mourer-Chauviré 1996a, # 12m): Mourer-Chauviré 1983, 1987, Mlíkovský 1998c. [B]

Möhren 19, Germany (Mlíkovský & Hesse 1996, # 9): Heissig 1978. [B]

Ravet-Lupovici, France (Mourer-Chauviré 1996a, # 12n): Mourer-Chauviré 1983, 1987, 1992b, Mlíkovský 1998c. [B]

Ronzon, France (Mourer-Chauviré 1996a, # 16): Gervais 1844a,b, 1849, 1848-1852, 1859, Aymard 1856, Milne-Edwards 1863, 1867-1868, Woodward 1886, Lydekker 1891a, Gaillard 1908, 1939, Cracraft & Rich 1972, Harrison 1975a, Olson 1978, 1985a. [B*, E, F]

Saignon, France (Mourer-Chauviré 1996a, # 17): Demathieu et al. 1984. [T*]

MP 21-22

Barros, Spain (Sánchez Marco 1996b, # 3): Sánchez Marco 1996b. [B]

MP 21-23

Hamstead, England (Mlíkovský 1996q, # 21): Harrison & Walker 1979b. [B*]

Yarmouth, England (Mlíkovský 1996q, # 20): Harrison & Walker 1979b. [B*]

MP 21-24

Agramunt, Spain (Sánchez Marco 1996b, # 2): Casanovas & Santafé 1982. [T]

Burnt Wood, England (Mlíkovský 1996q, # 18): Harrison & Walker 1976a, 1979b. [B]

Cadillac, France (Mourer-Chauviré 1996a, # 26): Gervais 1844a,b, 1849, 1848-1852. [B]

Frauenweiler, Germany (new): Mayr 2000a. [B*]

Gurnard Bay, England (Mlíkovský 1996q, # 19): Harrison & Walker 1979b. [B]

Hoeleden, Belgium (Cheneval 1996a, # 10): Misonne 1958. [B]

Hoogbutsel, Belgium (Cheneval 1996a, # 9): Misonne 1958. [B]

Liedena, Spain (new): Mangin 1962, de Raaf et al. 1965. [F]

Matt, Switzerland (Mlíkovský 1996o, # 3): Escher 1839, Meyer 1839c, 1844, 1865, Gervais 1844a, Heer 1865, 1879, Peyer 1949, 1950, 1957, Stüssi 1958, Baumann 1958, Olson 1976, Mlíkovský 1992b. [B*]

Oppertshofen 2, Germany (Mlíkovský & Hesse 1996, # 12): Dehm 1978. [B]

Sieblös, Germany (Mlíkovský & Hesse 1996, # 8): Hassenkamp 1858: 207, Martini 1967a,b, 1988, 1998, Martini & Tobien 1984. [B*, E]

Weissenburg 9, Germany (Mlíkovský & Hesse 1996, # 13): Dehm 1978. [B]

MP 21-25

Lupéron, France (new): Mayr 2000i. Mayr (2000i: 627) said that Lupéron is "probably" identical with Céreste without any substantiation. A locality named Lupéron was unknown to Mourer-Chauviré (1996a,b) and Cheneval (1996b). However, all localities from southern France, which yielded avian bones in plates, are early Oligocene (MP 21-25) in age (see Mourer-Chauviré 1996a), which makes it probable, that Lupéron is also early Oligocene in age, whether it is a hitherto unidentified locality or it is identical with one of the relevant localities listed in Mourer-Chauviré (1996a). [B*]

MP 21-28

Belgarite 4A, France (Mourer-Chauviré 1996a, # 12aa): Mourer-Chauviré 1980a, 1987. [B]

Boussac, France (Mourer-Chauviré 1996a, # 12z): Mourer-Chauviré 1980a, 1983. This is a complex locality, which includes the sites Boussac 1 and Boussac 2. [B]

Fonbonne 1, France (Mourer-Chauviré 1996a, # 12y): Mourer-Chauviré 1980a, 1985b, 1987, Mlíkovský 1998c. [B]

Phalip, France (Mourer-Chauviré 1996a, # 12bb): Mourer-Chauviré 1987, Mlíkovský 1998c. [B]

MP 21-30

- Alliers, France (Mourer-Chauviré 1996a, # 34): Milne-Edwards 1867-1868. [B]
 Aurillac, France (new): Gervais 1848-1852. [B]
 Peralta de la Sal, Spain (Sánchez Marco 1996b, # 1): Hernández-Pacheco 1929, Casanovas & Santafé 1982. [T]
 Sčigry, Ukraine (Mlíkovský 1996p, # 1): Nesov 1992. [B]
 Vorochta, Ukraine (Mlíkovský 1996p, # 2): Čerkašenko & Ljaščuk 1960, Ljaščuk 1962. [E]

MP 22

- Baraval, France (new): Sigé et al. 1998. [B]
 Bechlejovice, Czechia (Mlíkovský 1996e, # 1): Špinar 1986: 95, Mlíkovský 1992b. [F]
 Herrlingen 1, Germany (Mlíkovský & Hesse 1996, # 11): Palmowski & Wachendorf 1966. [B]
 Mas de Got, France (Mourer-Chauviré 1996a, # 12p): Mourer-Chauviré 1980a, 1983, 1987, 1992b, Mlíkovský 1998c. This is a complex locality, which includes the sites Mas de Got A and Mas de Got B. [B]
 Möhren 13, Germany (Mlíkovský & Hesse 1996, # 10): Heissig 1973. [B]
 Plante 2, France (Mourer-Chauviré 1996a, # 12o): Mourer-Chauviré 1980a, 1983. [B]

MP 23

- Céreste, France (Mourer-Chauviré 1996a, # 19): Lambrecht 1933, Stemvers-van-Bemmel 1984, Mourer-Chauviré 1992b, Mayr 1999a, 2000i. [B*, F]
 Crabit 1, France (Mourer-Chauviré 1996a, # 12s): Mourer-Chauviré 1978b, 1983, 1987, Karchu 1988, Mlíkovský 1998c. [B]
 Itardiès, France (Mourer-Chauviré 1996a, # 12q): Mourer-Chauviré 1980a, 1981, 1983, 1985b, 1987, 1992b, 1995a. [B*]
 Mounayne, France (Mourer-Chauviré 1996a, # 12r): Mourer-Chauviré 1980a. [B]
 Roqueprune 2, France (Mourer-Chauviré 1996a, # 12t): Mourer-Chauviré 1980a, 1983, Mourer-Chauviré & Cheneval 1983. [B]

MP 23-24

- Burgmagerbein 3, Germany (Mlíkovský & Hesse 1996, # 15): Dehm 1978. [B]
 Espenhain, Germany (Mlíkovský & Hesse 1996, # 16): Fischer 1982, 1983a,b, 1985, 1990. [B*]
 Froidefontaine, France (Mourer-Chauviré 1996a, # 25): Cheneval 1995. [B*]
 Kleinkems, Germany (Mlíkovský & Hesse 1996, # 14): Mieg et al. 1892: 379, Lambrecht 1933: 665. [F]
 Rupelmonde, Belgium (Cheneval 1996a, # 11): Beneden 1871, 1872, 1873a,b, 1883, Dollo 1909, Lambrecht 1931a, 1933, Miller & Sibley 1941, Brodkorb 1962a, Cheneval 1984a. [B*]
 Skalice, Czechia (Mlíkovský 1996e, # 3): Bayer 1882, 1883, Mlíkovský & Švec 1989. [B*]
 Varnsdorf, Czechia (Mlíkovský 1996e, # 2): Bayer 1882, 1883, Mlíkovský & Švec 1989. [B*]

MP 23-25

- Nassiet, France (Mourer-Chauviré 1996a, # 20): Viret 1938, Mourer-Chauviré 1996a. [B]

MP 24

- Cetățuie, Romania (Kessler 1996, # 2): Lambrecht 1933: 463, Cracraft 1973, Olson 1977. Dating corrected after Rusu (1989) and Kessler et al. 1998. [B*]
 Vachères, France (Mourer-Chauviré 1996a, # 18): Bessonat & Michaux 1973, Mayr 2000i: 635. [B]

MP 25

- Antoingt, France (Mourer-Chauviré 1996a, # 21): Lydekker 1891a. [B]
 Armissan, France (Mourer-Chauviré 1996a, # 22): Gervais 1862, Milne-Edwards 1863, 1869-1871, Woodward 1886, Lydekker 1891a, Eastman 1905, Schaub 1945, Brodkorb 1964: 313. [B*]
 Belgarric, France (Mourer-Chauviré 1996a, # 12u): Mourer-Chauviré 1983, 1987, 1992b. [B]




Fig. 8. Type localities of late Oligocene (MP 25-30) birds of Europe. 1 - Armissan, 2 - Chaptuzat, Gannat, Gergovie, 3 - Traun-Pucking, 4 - Câmpu lui Neag. Scale bar = 100 km.

Chaptuzat, France (Mourer-Chauviré 1996a, # 23): Cuvier 1825, Gervais 1844a, 1848-1852, 1849, 1859, Milne-Edwards 1863, 1867-1868, 1869-1871, 1872, Lydekker 1891a, Cheneval 1984a. [B*]

Garouillas, France (Mourer-Chauviré 1996a, # 12v): Mourer-Chauviré 1980a. [B]

Sauvetat, France (Mourer-Chauviré 1996a, # 24): Laizer 1828, Gervais 1844a, 1848-1852, Lydekker 1891a. [B, E]

MP 25-30

Bellingen, Germany (Mlíkovský & Hesse 1996: 627): Böhm 1896, 1898, Abel 1935, Mlíkovský 1992b. No avian remains. The footprints were reidentified as those of a mammal. [T]

Câmpu lui Neag, Romania (new): Rădan & Brustur 1993. [T*]

Perignat, France (Mourer-Chauviré 1996a, # 33): Lydekker 1891a. [B]

Tomerdingen, Germany (Mlíkovský & Hesse 1996, # 17): Seemann & Berckhemer 1930, Lambrecht 1933: 671. [B]

MP 27-30

Lausanne, Switzerland (Mlíkovský 1996o, # 9): Gaudin 1854. [E]

Thalberggraben, Germany (new): U.B. Göhlich in Darga et al. 1999. [B]

MP 28

Cournon, France (Mourer-Chauviré 1996a, # 27): Gervais 1844a, 1848-1852, Milne-Edwards 1867-1868, 1869-1871, 1872. [B, E]

Desse, France (Mourer-Chauviré 1996a, # 12w): Mourer-Chauviré 1980a, 1983, 1992b, 1993a, Mourer-Chauviré & Cheneval 1983. [B]

Fraysse, France (Mourer-Chauviré 1996a, # 12x): Crochet 1971, Mourer-Chauviré 1980a, 1983, 1992b, 1993a, Mourer-Chauviré & Cheneval 1983. [B]

Gaimersheim 1, Germany (Mlíkovský & Hesse 1996, # 18): Ballmann 1970. [B]




Fig. 9. Type localities of early Miocene (MN 1-4) birds of Europe. 1 - Léognan, Saucats, 2 - Orléanais, Touraine (age uncertain), 3 - Saint-Gérand-le-Puy, 4 - Weisenau, Wiesbaden, 5 - Harbatshofen, 6 - Grafenmühle, Wintershof-West, 7 - Dolnice, 8 - Břešťany, Merkur, Skyřice, 9 - Ipolytarnóc, 10 - Nižnij Strutyn', 11 - Piatra Neamț (incl. Cuejdi), Putna, 12 - Fallol, 13 - Armagnac, 14 - Ceva, 15 - San Angelo. Scale bar = 100 km.

MP 29

Herrlingen 9, Germany (new): Mayr 2001a. [B]

Pont-du-Château, France (Mourer-Chauviré 1996a, # 28): Gervais 1844a, 1848-1852, Milne-Edwards 1869-1871, Cheneval 1984a. [B, E]

MP 29 – MN 1

Chauderon, Switzerland (Mlíkovský 1996o, # 6): Weidmann & Reichel 1979. [T]

Crau Coulet, Switzerland (Mlíkovský 1996o, # 8): Weidmann & Reichel 1979. [T]

Goldau, Switzerland (Mlíkovský 1996o, # 5): Bräm 1954. [T]

Lammschlucht, Switzerland (Mlíkovský 1996o, # 4): Clerq & Holst 1971, Weidmann & Reichel 1979. [T]

Roches, Switzerland (Mlíkovský 1996o, # 7): Weidmann & Reichel 1979. [T]

MP 30

Aix-en-Provence, France (Mourer-Chauviré 1996a, # 35): Gervais 1844a, 1848-1852, 1859, Milne-Edwards 1863, 1869-1871, Bayan 1873, Woodward 1886). [F]

Coderet, France (Mourer-Chauviré 1996a, # 29): Mourer-Chauviré et al. 1989. [B]

Flörsheim, Germany (Mlíkovský & Hesse 1996, # 20): Lambrecht 1933: 671. [B]

Gannat, France (Mourer-Chauviré 1996a, # 30): Gervais 1844a, Milne-Edwards 1867-1868, Lydekker 1891a, Lambrecht 1933, Harrison 1975a,b, Cheneval 1984a, Olson 1985a, Mourer-Chauviré et al. 1989. [B*]

Gergovie, France (Mourer-Chauviré 1996a, # 31): Gervais 1844a, 1848-1852, 1849, Milne-Edwards 1863, 1867-1868, Cheneval 1984a. [B*]

Peulblanc, France (Mourer-Chauviré 1996a, # 32): Milne-Edwards 1867-1868. [B]
 Rott, Germany (Mlíkovský & Hesse 1996, # 19): Meyer 1859, 1867, Mlíkovský & Hesse 1996.
 [B, F]

Traun-Pucking, Austria (Mlíkovský 1996c, # 1): Mlíkovský 1987. [B*]

Trebovlje, Slovenia (new): Kolar-Jurovšek & Jurovšek 1996. [F]

MN 1

Oppenheim, Germany (Mlíkovský & Hesse 1996, # 24): Rothausen 1966, Olson & Feduccia
 1980a, Mlíkovský 1988, Cheneval & Escuillé 1992. [B]

Paulhiac, France (Cheneval 1996b, # 1): Mourer-Chauviré 1993a. [B]

Pyrimont-Challonges, France (Cheneval 1996b, # 2): Depéret & Douxami 1902, Cheneval
 1983c. [B]

Weisenau-AS, Germany (Mlíkovský & Hesse 1996, # 23): Meyer 1839b, 1843, Becker 1849,
 Milne-Edwards 1869-1871, Lambrecht 1933, Lydekker 1891a, Cheneval 1995. [B*, E]

Zell, Germany (Mlíkovský & Hesse 1996, # 22): Lambrecht 1933: 359. [E]

MN 1-2

Bachów, Poland (Bocheński 1996, # 2): Bocheński 1989, 1993, 1996. [B]

Ceva, Italy (Delle Cave 1996, # 5): Portis 1884. [B*]

Hłudno, Poland (Bocheński 1996, # 1): Bocheński & Szymczyk 1979, Bocheński 1989, 1993.
 [B]

MN 1-4

Kaltennordheim, Germany (Mlíkovský & Hesse 1996, # 21): Schlotheim 1820: 26. [B]

MN 1-8

Milićevo Brdo, Serbia (Mlíkovský 1996r, # 1): Laskarev 1936. [B]

MN 1-13

Ariuşd, Romania (Kessler 1996, # 3): Lambrecht 1929. [B]

Cevico de la Torre, Spain (Sánchez Marco 1996b, # 16): Olavarria 1898, E. Hernández-
 Pacheco 1914a, Lambrecht 1933. [E]

Heiligenstadt, Austria (Mlíkovský 1996c, # 7): Lambrecht 1933: 687. There are also Quatern-
 ary records from this locality (G. Rabeder, pers. communication 2001). [E]

Lisboa, Portugal (Sánchez Marco 1996a, # 4): Sánchez Marco 1996a. [B]

San Angelo, Italy (Delle Cave 1996, # 10): Procaccini Ricci 1840: 60, Portis 1887. [F*]

Setubal Peninsula, Portugal (Sánchez Marco 1996a, # 5): Sánchez Marco 1996a. [B]

Simeria, Romania (Kessler 1996, # 4): Lambrecht 1929. [F]

MN 2

Budenheim, Germany (Mlíkovský & Hesse 1996, # 25): Lambrecht 1933: 670. [B]

Büchelberg, Germany (Mlíkovský & Hesse 1996, # 32): Kuss 1960. [B]

Fischenbach, Switzerland (Mlíkovský 1996p, # 11): Gasser 1966, Weidmann & Reichel 1979.
 [F]

Ingelheim, Germany (Mlíkovský & Hesse 1996, # 34): Lambrecht 1933: 670. [B?]

Mombach, Germany (Mlíkovský & Hesse 1996, # 33): Hoeninghaus 1839. [B]

Monsheim, Germany (Mlíkovský & Hesse 1996, # 28): Lambrecht 1933: 341. [B]

Navarrete del Rio, Spain (Sánchez Marco 1996b, # 4): Adrover 1975. [B]

Ravolzhausen, Germany (Mlíkovský & Hesse 1996, # 35): Martini 1974, Eikamp 1980. [B]

Saint-Gérard-le-Puy, France (Cheneval 1996b, # 3): Milne-Edwards 1863, 1867-1868, 1869-
 1871, 1870, 1872, Lydekker 1891a,c, Shufeldt 1896, Lambrecht 1933, Gaillard 1939, Lowe
 1939, Storer 1956, Wetmore 1956, Brodkorb 1963b,c, 1964, 1967, 1970, 1971b, 1978,
 Ballmann 1969a, Cracraft 1973, Collins 1976b, Harrison & Walker 1976d, Olson 1977,
 1981, Harrison 1979a, 1984c, 1986, Mourer-Chauviré & Cheneval 1983, Cheneval 1979,
 1983,b,c, 1984a,b, 1987, 1989, Livezey & Martin 1988, Fischer & Mauersberger 1989,
 Cheneval & Escuillé 1992, Mourer-Chauviré 1992b, 1993a, 1995b, 2000, Mlíkovský
 1998c,e, 1999e: 35, 2000e, Mlíkovský & Göhlich 2000. [B*]

Senftenberg, Germany (Mlíkovský & Hesse 1996, # 36): Menzel 1906. [F]
 Treuchtlingen 2, Germany (Mlíkovský & Hesse 1996, # 30): Dehm 1978. [B]
 Ulm-Westtangente, Germany (Mlíkovský & Hesse 1996, # 29): Mlíkovský & Hesse 1996. [B]
 Weissenburg 6, Germany (Mlíkovský & Hesse 1996, # 31): Dehm 1978. [B]
 Wiesbaden 1, Germany (Mlíkovský & Hesse 1996, # 27): Meyer 1839b, Lambrecht 1933,
 Mlíkovský 1992b, 1995a. [B*]
 Wiesbaden-Hessler, Germany (Mlíkovský & Hesse 1996, # 26): Meyer 1839b, Lambrecht
 1931a, 1933, Rothausen 1966, Eikamp 1979. [B*, E]

MN 2-3

Armagnac, France (Cheneval 1996b, # 6): Lartet 1857, Milne-Edwards 1867-1868, 1874,
 Harrison & Walker 1996b. [B*]
 Grafenmühle 21, Germany (new): Mlíkovský 1998c. [B*]
 Léognan, France (Cheneval 1996b, # 5): Lartet 1857, Milne-Edwards 1867-1868, 1874,
 Brochon 1880, Lydekker 1891a, Lambrecht 1933, Lucazeau 1959, Brodkorb 1963c. [B]
 Saucats, France (Cheneval 1996b, # 4): Lartet 1857, Milne-Edwards 1874, Lambrecht 1933,
 Lucazeau 1959, Brodkorb 1963c. [B*]
 Solnhofen, Germany (Mlíkovský & Hesse 1996, # 37): Dehm 1935. [B]

MN 2-5

Touraine, France (Cheneval 1996b, # 7): Milne-Edwards 1869-1871. [B*]

MN 3

Ahnikov, Czechia (Mlíkovský 1996e, # 5): Mlíkovský 1992b. [B]
 Augine 7, Switzerland (Mlíkovský 1996o, # 14): Berger 1985. [T]
 Bré 8, Switzerland (Mlíkovský 1996o, # 17): Berger 1985. [T]
 Břešťany, Czechia (Mlíkovský 1996e, # 7): Laube 1901, 1909, Mlíkovský & Švec 1989,
 Mlíkovský 1992b, 1999c. [B*]
 Burg, Switzerland (Mlíkovský 1996o, # 19): Berger 1985. [T]
 Can Mas, Spain (Sánchez Marco 1996b, # 6): Villalta 1963. Age uncertain. [B]
 Deljatin, Ukraine (Mlíkovský 1996p, # 3): Vjalov & Flerov 1952, Vjalov 1965, 1966. [T]
 Fallol, Spain (Sánchez Marco 1996b, # 5): Crusafont & Villalta 1955, Villalta 1963. Age
 uncertain. [T*]
 Flon Morand, Switzerland (Mlíkovský 1996o, # 12): Weidmann & Reichel 1979. [T]
 Harbatshofen, Germany (Mlíkovský & Hesse 1996, # 38): Haushalter 1855, Brodkorb 1980.
 [B*]
 Heitenried 7, Switzerland (Mlíkovský 1996o, # 18): Berger 1985. [T]
 Ipolytarnóc, Hungary (Mlíkovský 1996j, # 1): Lambrecht 1912b, 1933: 675, Kordos 1983. [T*]
 Kosov, Ukraine (Mlíkovský 1996p, # 4): Vjalov & Flerov 1952, Tkačenko et al. 1967. [T]
 Liefrens 5, Switzerland (Mlíkovský 1996o, # 16): Berger 1985. [T]
 Mafflon 1, Switzerland (Mlíkovský 1996o, # 15): Berger 1985. [T]
 Mauvières, France (new): Ginsburg et al. 2000. [B]
 Merkur, Czechia (Mlíkovský 1996e, # 8): Mlíkovský 1998e,f. [B*]
 Nadvornaja, Ukraine (Mlíkovský 1996p, # 5): Dubrovo & Kapelist 1979. [T]
 Nižnij Strutyn', Ukraine (Mlíkovský 1996p, # 25): Vjalov 1960, 1965, 1966. [T*]
 Renggbach, Switzerland (Mlíkovský 1996o, # 21): Weidmann & Reichel 1979. [T]
 Skyřice, Czechia (Mlíkovský 1996e, # 4): Laube 1910, Mlíkovský, 2000d. [B*]
 Sloboda, Ukraine (Mlíkovský 1996p, # 6): Vjalov 1951. [T]
 Stubersheim, Germany (Mlíkovský & Hesse 1996, # 40): Mlíkovský & Hesse 1996. [T]
 Tuchořice, Czechia (Mlíkovský 1996e, # 6): Mlíkovský 1980, 1992b, 1999e: 35. [B, brain
 casts]
 Ussements, Switzerland (Mlíkovský 1996o, # 13): Weidmann & Reichel 1979. [T]
 Waldhalde, Switzerland (Mlíkovský 1996o, # 20): Speck 1945, 1952, Peyer 1950. [T]
 Wintershof-West, Germany (Mlíkovský & Hesse 1996, # 39): Dehm 1937, Ballmann 1966,
 1969b, Olson 1991, Mlíkovský 1998e, Mlíkovský & Göhlich 2000. [B*]

MN 3-4

- Andrașiu de Jos, Romania (Kessler 1996, # 7): Paucă 1942. [T]
 Călugăr, Romania (new): Panin 1964. [T]
 Cuejdi, Romania (new): Panin 1964. [T*]
 Limberg, Austria (Mlíkovský 1996c, # 2): Bonaparte 1856a: 781, Bachmayer 1980, Mlíkovský 1998c. [B]
 Mîngălar, Romania (new): Panin 1964. [T]
 Muntenia, Romania (new): Panin et al. 1966. [T]
 Petersbuch 28, Germany (new): Mlíkovský, orig. [B]
 Piatra Neamț, Romania (Kessler 1996, # 6): Panin 1964. This is a complex locality, where footprints were found in the valleys of several brooks, incl. Călugăr, Cuejdi, and Mîngălar. [T*]
 Putna, Romania (Kessler 1996, # 8): Panin 1961, 1964, Panin & Avram 1962. [T*]
 Schitu Frumusea, Romania (Kessler 1996, # 5): Grozescu 1914, Vjalov 1966. [T]

MN 4

- Allerding, Austria (new): Mlíkovský, orig. (unpublished material in coll. Danninger). [B]
 Alto de Ballester, Spain (Sánchez Marco 1996b, # 7): Sánchez Marco 1996b. [B]
 Artenay, France (new): Mlíkovský, orig. (unpublished material in MNHN). [B]
 Artesilla, Spain (Sánchez Marco 1996b, # 8): Sánchez Marco 1996b. [B]
 Casots, Spain (Sánchez Marco 1996b, # 11): Sánchez Marco 1996b. [B]
 Córcoles, Spain (Sánchez Marco 1996b, # 9): Sánchez Marco 1996b. [B]
 Dolnice, Czechia (Mlíkovský 1996e, # 9): Švec 1980, 1981, 1982, 1983, 1984, 1985, Švec & Mlíkovský 1986, Mlíkovský 1992b, 1999a,b, 2000c, Mourer-Chauviré 1999a. [B*, E]
 Erkerthshofen 1, Germany (Mlíkovský & Hesse 1996, # 44): Dehm 1978. [B]
 Estacion Imperial, Spain (Sánchez Marco 1996b, # 12): Sánchez Marco 1996b. [B]
 Kirchberg, Germany (Mlíkovský & Hesse 1996, # 43): Probst 1879, Lambrecht 1933. [B]
 Langenau, Germany (Mlíkovský & Hesse 1996, # 45): Mlíkovský & Hesse 1996. [B]
 Ořečov, Czechia (Mlíkovský 1996e, # 10): Mlíkovský 1992b. [B]
 Orleanais, France (Cheneval 1996b, # 8): Milne-Edwards 1867-1868, 1869-1871. [B*]
 Petersbuch 2, Germany (Mlíkovský & Hesse 1996, # 41): Heissig 1978. [B]
 Petersbuch 4, Germany (Mlíkovský & Hesse 1996, # 42): Mlíkovský & Hesse 1996. [B]
 Petersbuch 38, Germany (new): Mlíkovský & Göhlich 2000. [B]
 Retama, Spain (Sánchez Marco 1996b, # 10): Sánchez Marco 1996b. [B]

MN 4-5

- Oberdorf, Austria (Mlíkovský 1996c, # 3): Mlíkovský 1998a. [B]
 Raitenbuch 2, Germany (Mlíkovský & Hesse 1996, # 46): Heissig 1978. [B]
 Zelená, Czechia (Mlíkovský 1996d, # 11): Mlíkovský 1992b. [F]

MN 5

- Amôr 2, Portugal (Sánchez Marco 1996a, # 2): Zbyszewski & Ferreira 1967. [B]
 Františkovy Lázně, Czechia (Mlíkovský 1996e, # 13): Mlíkovský 1992b. [B]
 Mokřina, Czechia (Mlíkovský 1996e, # 14): Novák 1877, Mlíkovský 1992b. [B, F]
 Rothenstein 1-13, Germany (new): Mlíkovský, orig. (unpublished material in coll. Rummel). [B]
 Sandelzhausen, Germany (Mlíkovský & Hesse 1996, # 52): Fahlbusch & Gall 1970, Göhlich 2000. Dating corrected after Göhlich & Rössner (1999). [B]
 Sokolov, Czechia (Mlíkovský 1996e, # 12): Mlíkovský 1992b. [B]
 Vieux Collonges, France (Cheneval 1996b, # 9): Balmann 1972, Mlíkovský 1998f. [B*]

MN 5-6

- Emmenweid, Switzerland (Mlíkovský 1996p, # 10): Bachmann 1878. [E]
 Günzburg, Germany (Mlíkovský & Hesse 1996, # 49): Rühl 1896, Lambrecht 1933. [B]
 Heggbach, Germany (Mlíkovský & Hesse 1996, # 48): Probst 1879. [B]




Fig. 10. Type localities of middle Miocene (MN 5-8) birds of Europe. 1 - Sansan, 2 - Vieux Collonges, 3 - Grive-Saint-Alban, 4 - Antwerp, 5 - Öhningen, 6 - Steinheim, 7 - Nördlinger Ries (incl. Hahnenberg, Lierheim, Steinberg), 8 - Dechbetten, 9 - Sankt Margarethen, 10 - Radoboj, 11 - Ciobanița, Credința. Scale bar = 100 km.

Risitobel, Switzerland (Mlíkovský 1996o, # 23): Weidmann & Reichel 1979. [T]

Schönweg, Austria (Mlíkovský 1996c, # 5): Rauscher 1984. [E]

Sulzigotobel, Switzerland (Mlíkovský 1996o, # 22): Weidmann & Reichel 1979. [T]

Weingraben, Austria (Mlíkovský 1996c, # 4): Bachmayer 1964. [F]

MN 5-8

Kardam, Bulgaria (Boev 1996, # 2): Boev 1988, 1992, 1996. [B]

Schemmerberg, Germany (Mlíkovský & Hesse 1996, # 47): Probst 1879, Lambrecht 1933. [B]

MN 5-13

Bela Stena, Serbia (Mlíkovský 1996r, # 3): Stevanović 1959. [F]

Peskovita, Serbia (Mlíkovský 1996r, # 2): Stevanović 1959. [E]

MN 6

Devínská Nová Ves – fissure, Slovakia (Mlíkovský 1996n, # 1): Švec 1986, Mlíkovský 1996n. [B]

Goldberg, Germany (Mlíkovský & Hesse 1996, # 50a): Lambrecht 1933, Seemann 1941, Ballmann 1979. [B]

Hahnenberg, Germany (Mlíkovský & Hesse 1996, # 50b): Fraas 1870, Lydekker 1891a, Ammon 1918, Lambrecht 1933, Mlíkovský 1988. [B*, E]

Lierheim, Germany (Mlíkovský & Hesse 1996, # 50c): Lydekker 1891a, Ammon 1918, Lambrecht 1933. [B*, E, F]

Mátraszőlős 1, Hungary (new): Gál et al. 1999, 2000. [B]

Mátraszőlős 2, Hungary (new): Gál et al. 2000. [B]

Nördlinger Ries, Germany (Mlíkovský & Hesse 1996, # 50): Fraas 1879, Lydekker 1891a, Ammon 1918, Lambrecht 1933, Seemann 1941, Ballmann 1979, 1983, Heizmann & Fahlbusch 1983, Mlíkovský 1988, 1992b, Heizmann & Hesse 1995. This is a complex locality which includes the following sites: Goldberg, Hahnenberg, Lierheim, Steinberg, and Wallerstein. [B*, E, F]

Paracuellos 5, Spain (Sánchez Marco 1996b, # 13): Sánchez Marco 1996b. [B]

Petersbuch 39, Germany (new): Mlíkovský, orig. (unpublished material in coll. Rummel). [B]

Rothenstein, Germany (Mlíkovský & Hesse 1996, # 51): Mlíkovský & Hesse 1996. [B]

Sansan, France (Cheneval 1996b, # 10): Milne-Edwards 1867-1868, 1869-1871, 1872, Woodward 1886, Lydekker 1891a, Lambrecht 1933, Gaillard 1939, Brodkorb 1964, 1967, 1971b, Howard 1964, Cracraft 1973, Olson 1977, Rich & Haarhoff 1985, Cheneval 1987, 2000, Mlíkovský 1998e. [B*]

Steinberg, Germany (Mlíkovský & Hesse 1996, # 50d): Ballmann 1979, 1983, Heizmann & Fahlbusch 1983. [B*]

Wallerstein, Germany (Mlíkovský & Hesse 1996, # 50e): Ammon 1918. [B]

MN 6-7

Baltringen, Germany (Mlíkovský & Hesse 1996, # 54): Probst 1879. [B]

Minișu de Sus, Romania (new): Kessler & Codrea 1996. [B]

Storzingen, Germany (Mlíkovský & Hesse 1996, # 53): Probst 1879. [B]

MN 7

Attenfeld, Germany (Mlíkovský & Hesse 1996, # 58): Schlosser 1916, Lambrecht 1933. [B]

Böttingen, Germany (Mlíkovský & Hesse 1996, # 56): Zeuner 1931. [F]

Climbach 1, Germany (Mlíkovský & Hesse 1996, # 59): Dieffenbach 1853, Engelhardt & Schottler 1914. [B]

Lišov, Czechia (Mlíkovský 1996e, # 15): Mlíkovský 1996d. [B]

Öhningen, Germany (Mlíkovský & Hesse 1996, # 55): Scheuchzer 1708a,b, 1716, Karg 1805, Meyer 1838, 1839a, 1845, 1864, 1865, Heer 1865, Milne-Edwards 1867-1868, Lydekker 1891a, Lambrecht 1933, Mlíkovský 1992b. [B*, F]

Przeworno 2, Poland (Bocheński 1996, # 3): Bocheński 1987. [B]

Radoboj, Croatia (Mlíkovský 1996d, # 1): Meyer 1850, 1864, 1865, Mlíkovský 1997a. [B*]

Sankt Margarethen, Austria (Mlíkovský 1996c, # 6): Mlíkovský 1998b. [B*]

Steinheim, Germany (Mlíkovský & Hesse 1996, # 57): Fraas 1870, 1879, Lydekker 1891a, Lambrecht 1933, Bocheński 1987, Mlíkovský 1992b, Heizmann & Hesse 1995, Mlíkovský & Hesse 1996. [B*]

MN 7-8

Antwerp, Belgium (Cheneval 1996a, # 12): Beneden 1871, 1872, 1873a,b, 1883, Shufeldt 1909: 336, Lambrecht 1933, Brodkorb 1962a, Olson & Walker 1997. [B*]

Brașov, Romania (Kessler 1996, # 10): Andrae 1855, Lambrecht 1933. [F]

Cuesta del Rey, Spain (Sánchez Marco 1996b, # 19): Sánchez Marco 1996b. [B]

Dechbetten, Germany (Mlíkovský & Hesse 1996, # 60): Ammon 1918, Lambrecht 1933, Brodkorb 1980, Olson 1985a, Švec 1986, Bocheński 1987, Mlíkovský 1992b. [B*]

Escobosa de Calatañazor, Spain (Sánchez Marco 1996b, # 18): Sesé & López 1977, López et al. 1977, Sánchez Marco 1996b. [B]

Grive-Saint-Alban – early collections, France (Cheneval 1996b, # 11): Depéret 1887, Lydekker 1891a, 1893, Shufeldt 1896, Regàlia 1897, Ennouchi 1930, Lambrecht 1933, Gaillard 1939, Brodkorb 1964, 1967, 1971a, Ballmann 1969a, Cracraft 1973, Olson 1974a, Bocheński 1987, Cheneval 1987, Mlíkovský 1998c. This is a complex locality, consisting of several fissures. Relevant data are missing for the early collections, but recently excavated fissures (L-7, M) belong to MN 8 (see below). [B*]

Otero, Spain (Sánchez Marco 1996b, # 17): Hernández-Pacheco 1914, Hernández-Pacheco & Dantín 1915, Lambrecht 1933. [B]




Fig. 11. Type localities of late Miocene (MN 9-13) birds of Europe. 1 - Eppelsheim, 2 - Aumeister, 3 - Vösendorf, 4 - Kohfidisch, 5 - Sümeg, 6 - Polgárdi, 7 - Csákvár, 8 - Rudabánya, 9 - Tataruş-Brusturi, 10 - Chişinău, Golboçica, 11 - Bujoru, 12 - Čebotarevka, Hrebenyky, Nova Emetivka, Varnița, 13 - Novaja Slobodka, 14 - Novoelisavetovka, 15 - Malynivcy, 16 - Apostolovo, 17 - Libros, Mansuetos, 18 - Gabbro, 19 - Bamboli, 20 - Ancona, Senigallia, 21 - Chrabärsko, 22 - Chadžidimovo, 23 - Trojanovo, 24 - Pikermi, 25 - Samos. Scale bar = 100 km.

Paracuellos 3, Spain (Sánchez Marco 1996b, # 14): Sánchez Marco 1996b. [B]

Săcădat, Romania (Kessler 1996, # 9): Andrae 1855, Lambrecht 1933. [F]

Toril 3, Spain (Sánchez Marco 1996b, # 15): Sánchez Marco 1996b. [B]

MN 8

Ciobănița, Romania (Kessler 1996, # 12): Grigorescu & Kessler 1977, Kessler & Gál 1996. [B*]

Collet-Redon, France (Cheneval 1996b, # 12): Jullien et al. 1979. [B]

Credința, Romania (Kessler 1996, # 11): Grigorescu & Kessler 1977, 1988, Cheneval 1987, Kessler 1992. [B*]

Grive-Saint-Alban – fissure L-7, France (Cheneval 1996b, # 11): Ballmann 1969a. [B]

Grive-Saint-Alban – fissure M, France (Cheneval 1996, # 11): Ballmann 1969a, Olson 1974, Bocheński 1987. [B]

Petersbuch 6, Germany (new): Mlíkovský, orig. (unpublished material in coll. Rummel). [B]

Petersbuch 10, Germany (new): Mlíkovský, orig. (unpublished material in coll. Rummel). [B]

Petersbuch 18, Germany (new): Mlíkovský, orig. (unpublished material in coll. Rummel). [B]

MN 8-9

Hostalets de Pierola, Spain (Sánchez Marco 1996b, # 20): Villalta & Crusafont Pairó 1950, Villalta 1963. [B]

Maar am Kaltenbrunnen, Germany (Mlíkovský & Hesse 1996, # 62): Bräuhäuser 1922. [B]

Randecker Maar, Germany (Mlíkovský & Hesse 1996, # 61): Westphal 1963. [F, coprolithes]

MN 9

Aumeister, Germany (Mlíkovský & Hesse 1996, # 63): Stromer 1928, Lambrecht 1933, Mlíkovský 1992b. [B*]

Ballestar, Spain (Sánchez Marco 1996b, # 23): Roset 1974, Agustí 1988. [B]

Can Ponsic, Spain (Sánchez Marco 1996b, # 22): Sánchez Marco 1996b. [B]

Chişinău, Moldova (Mlíkovský 1996l, # 5): Macarovici & Oescu 1941, Kuročkin & Ganea 1972, Kessler 1984b, 1992, Kessler & Gál 1996. [B*]

Croix-Rousse, France (Cheneval 1996b, # 13): Depéret 1887. [B]

Eppelsheim, Germany (Mlíkovský & Hesse 1996, # 65): Lambrecht 1933, Fischer & Stephan 1971, Mourer-Chauviré et al. 1985, Mlíkovský 1992b. [B*]

Golbočica, Moldova (Mlíkovský 1996i, # 4): Kuročkin & Ganea 1972. Age uncertain. [B*]

Höwenegg, Germany (Mlíkovský & Hesse 1996, # 64): Jörg 1957. [B]

Kalfa, Moldova (Mlíkovský 1996l, # 3): Ganea 1965, Kuročkin & Ganea 1972. [B]

Rudabánya, Hungary (Mlíkovský 1996j, # 2): Jánossy 1976b, 1977, 1993. [B*]

Valles de Fuentidueña, Spain (Sánchez Marco 1996b, # 21): Villalta 1963, Sánchez Marco 1996b. [B]

Varnița, Moldova (Mlíkovský 1996l, # 2): Lungu 1966a,b, 1968, Roška 1967, Kuročkin & Lungu 1970. [B*, E]

MN 9-10

Apostolovo, Ukraine (Mlíkovský 1996p, # 13): Alekseev 1916, Roščin 1959, 1952, Michajlov 1988. [E*]

Aveiras de Baixo, Portugal (Sánchez Marco 1996a): Sánchez Marco 1996a. [B]

Jur'yvka, Ukraine (Mlíkovský 1996p, # 12): Pidopličko 1956. [E]

Kerč' Peninsula 1, Ukraine (Mlíkovský 1996p, # 8): Dubrovo & Kapelist 1979. [B]

Kolkotova Balka, Moldova (Mlíkovský 1996l, # 7): Laskarev 1908, 1912, Lambrecht 1933: 443, Burčak-Abramovič 1951, Burčak-Abramovič & Bendukidze 1971, Cracraft 1973, Kuročkin 1981, Michajlov 1988. [B, E]

Krivoj Roh, Ukraine (Mlíkovský 1996p, # 7): Dubrovo & Kapelist 1979. [B]

Kubanka, Ukraine (Mlíkovský 1996p, # 9): Vojinstvens'kyj 1967. [B]

Libros, Spain (Sánchez Marco 1996b, # 24): Navás 1922, Lambrecht 1933, Cracraft 1973, Olson 1995a. [B*]

Pokşeşti, Moldova (Mlíkovský 1996l, # 6): Michajlov 1988. [E]

Sokolov, Ukraine (Mlíkovský 1996p, # 10): Vojinstvens'kyj 1967. [B]

Suchomasty 3, Czechia (Mlíkovský 1996e, # 16): Mlíkovský 1996e. [B]

Tjaginka, Ukraine (Mlíkovský 1996p, # 11): Vojinstvens'kyj 1967. [B]

MN 9-11

Bujoru, Moldova (Mlíkovský 1996l, # 1): Kuročkin & Ganea 1972, Olson 1977. Age = MN 9 (Mein 1990) or MN 11 (Fejfar & Heinrich 1990). [B*]

MN 10

Batallones, Spain (Sánchez Marco 1996b, # 25): Sánchez Marco 1996b. [B]

Götzendorf, Austria (Mlíkovský 1996c, # 9): Mlíkovský 1992a. [B]

Kohfidisch, Austria (Mlíkovský 1996c, # 10): Mlíkovský 1998c, orig. (new taxa to be described) [B*]

Tataruş-Brusturi, Romania (Kessler 1996, # 13): Lambrecht 1916a, 1933. [B*]

Viladecavalls, Spain (Sánchez Marco 1996b, # 26): Villalta 1963. [B]

Vösendorf, Austria (Mlíkovský 1996c, # 8): Thenius 1954, Mlíkovský 1997b. [B*]

MN 11

Crevillente 2, Spain (Sánchez Marco 1996b, # 27): Sánchez Marco 1996b. [B]

Csákvár, Hungary (Mlíkovský 1996h, # 3): Lambrecht 1933, Kretzoi 1954b, 1957, Jánossy 1977, Mlíkovský 1992b, 1998f. [B*]

Hrebenyky, Ukraine (Mlíkovský 1996p, # 16): Voznesens'kyj 1937, Burčák-Abramovič 1939, 1949, 1951, 1953a,b, Umans'ka 1979a, Kuročkin 1981. [B*]

Puente Minero, Spain (Sánchez Marco 1996b, # 28): Sánchez Marco 1996b. [B]

Samos 1, Greece (Mlíkovský 1996h, # 1): Forsyth Major 1888, 1892, 1894, Lydekker 1891a, Martin 1903, Lambrecht 1933, Cracraft 1973, Harrison 1981, Kuročkin 1981, Solounias 1981, Karchu 1997. [B*]

MN 11-12

Chadžidimovo, Bulgaria (new): Boev & Kovačev 1998. [B*]

Novoelisavetovka, Ukraine (Mlíkovský 1996p, # 21): Alekseev 1916, Burčák-Abramovič 1939, Vojinstvens'kyj 1967, Kuročkin & Lungu 1970, Umans'ka 1981. [B*]

Sümeğ, Hungary (Mlíkovský 1996j, # 4): Jánossy 1976b, 1977, Mlíkovský 1992b. [B*]

MN 11-13

Ancona, Italy (Delle Cave 1996, # 11): Portis 1887. [F*]

Andreevka, Ukraine (Mlíkovský 1996p, # 14): see Dubrovo & Kapelist 1979. [B]

Belka, Ukraine (Mlíkovský 1996p, # 15): Vojinstvens'kyj 1967. [B]

Čebotarevka, Ukraine (Mlíkovský 1996p, # 24): Vojinstvens'kyj 1967, Umans'ka 1973, 1979b, 1981, Kuročkin 1981, Michajlov 1988. [B*, E]

Chrabářsko, Bulgaria (Boev 1996, # 3): Burčák-Abramovič & Nikolov 1984, Boev 1988, 1992, 1996, 2000d. [B*]

Çimişlia, Moldova (Mlíkovský 1996l, # 8): Michajlov 1988, Michajlov & Kuročkin 1988. [E]

Corund, Romania (Kessler 1996, # 14): Gheorghiu et al. 1965. [F]

Il'inka, Ukraine (Mlíkovský 1996p, # 18): Burčák-Abramovič 1953a, Vojinstvens'kyj 1967. [B, E]

Kujal'nik, Ukraine (Mlíkovský 1996p, # 19): Laskarev 1912, Pržemysskij 1912, 1914, Alekseev 1916, Korotkevič 1961, 1972, Kuročkin & Lungu 1970. [B]

Licata, Italy (Delle Cave 1996, # 7): Milne-Edwards 1869-1871, Portis 1879, 1887. [B]

Malynivcy, Ukraine (Mlíkovský 1996p, # 26): Brandt 1873, 1874, 1885, Mäklin 1873, Nathusius 1886, Bakulovskij 1892, Bensley 1921, Burčák-Abramovič 1953a, Schönwetter 1960, Roščin 1962. [E*]

Naumovka, Ukraine (Mlíkovský 1996p, # 27): Dubrovo & Kapelist 1979. [B]

Novaja Slobodka, Ukraine (Mlíkovský 1996p, # 28): Vidgal'm 1886, Tugarinov 1940a, Burčák-Abramovič 1958, Miller 1966, Ganea & Kuročkin 1967, Vojinstvens'kyj 1967, Burčák-Abramovič & Nikolov 1984. [B*]

Novoukrainka, Ukraine (Mlíkovský 1996p, # 22): Vojinstvens'kyj 1967. [B]

Odesa – surroundings, Ukraine (Mlíkovský 1996p, # 29): Nordmann 1847, Dubrovo & Kapelist 1979. This is not a locality in the strict sense. There are several late Miocene fossiliferous localities in the surroundings of Odesa, but they were not indicated for some old collections. [B]

Škodova Gora, Ukraine (Mlíkovský 1996p, # 30): Vojinstvens'kyj 1967. [B]

Snigirevka, Ukraine (Mlíkovský 1996p, # 23): Burčák-Abramovič 1953a. [E]

Sofija 1, Bulgaria (Boev 1996, # 4): Popov et al. 1921, Boev 1988 1992. [B]

Žovten', Ukraine (Mlíkovský 1996p, # 17): Ganea & Kuročkin 1967. [B]

MN 11-15

Trojanovo, Bulgaria (Boev 1996, # 6): Burčák-Abramovič & Nikolov 1984. [B*]

MN 12

Aljezar B, Spain (Sánchez Marco 1996b, # 32): Adrover 1986, Cheneval & Adrover 1993. [B]

Concud 1, Spain (Sánchez Marco 1996b, # 29): Sánchez Marco 1996b. [B]

Mansuetos, Spain (Sánchez Marco 1996b, # 30): Villalta 1963. [B*]

Tardosbánya, Hungary (Mlíkovský 1996j, # 5): Jánossy 1976b. [B]

Valdecebro 5, Spain (Sánchez Marco 1996b, # 31): Sánchez Marco 1996b. [B]

MN 12-13

- Kremnikovci, Bulgaria (Boev 1996, # 1): Boev 1996. [B]
 Pikermi, Greece (Mlíkovský 1996h, # 2): Roth & Wagner 1855, Gaudry 1862a,b,c, 1862-1867, Weithofer 1888, Lambrecht 1933, Bachmayer & Zapfe 1962, Cracraft 1973, Jánossy 1976b, Mourer-Chauviré et al. 1985, Mlíkovský 1996h. [B*]
 Veles, Macedonia (Mlíkovský 1996k): Schlosser 1921, Lambrecht 1933: 692; see also Cirić 1957. [B]

MN 13

- Amandoli, Italy (Delle Cave 1996, # 6a): Delle Cave 1996. [B, F]
 Arquillo, Spain (Sánchez Marco 1996b, # 33): Sánchez Marco 1996b. [B]
 Bamboli, Italy (Delle Cave 1996, # 12): Gastaldi 1866a,b, Portis 1884, Weithofer 1888, Ristori 1891, Howard 1964. [B*]
 Cherasco, Italy (Delle Cave 1996, # 12a): Cheneval 1993, see also Cavallo & Gaudant 1984, Cavallo et al. 1986. [B]
 Gabbro, Italy (Delle Cave 1996, # 8): Bosniaski 1879, Portis 1887, Pycraft 1909, Lambrecht 1933, Bradley & Landini 1984. [B*, F*]
 Mondaino, Italy (Delle Cave 1996, # 6c): Delle Cave 1996. [B, F]
 Nova Emetivka, Ukraine (Mlíkovský 1996p, # 20): Voznesens'kyj 1939, Burčák-Abramovič 1939, 1953a, Ganea & Kuročkin 1967, Vojinstev's'kyj 1967, Burčák-Abramovič & Vekua 1971, Umans'ka 1979a. [B*]
 Polgárdi, Hungary (Mlíkovský 1996j, # 6): Lambrecht 1912a,b, Jánossy 1976b, 1979, 1991, 1995, Mlíkovský 1998c. This is a complex locality, consisting of several fissures, incl. Polgárdi 2, 4 and 5. Relevant data are missing for some of the collections. [B*]
 Polgárdi 2, Hungary (Mlíkovský 1996j, # 6): V. Čapek in Lambrecht 1933: 780, Jánossy 1991. [B]
 Polgárdi 4, Hungary (Mlíkovský 1996j, # 6): Jánossy 1976b, 1991. [B*]
 Polgárdi 5, Hungary (Mlíkovský 1996j, # 6): Jánossy 1991, Mlíkovský 1998c. [B*]
 Roddi, Italy (Delle Cave 1996, # 12c): Cavallo et al. 1986. [B]
 Scaparoni, Italy (Delle Cave 1996, # 12b): Cavallo et al. 1986. [F]
 Senigallia, Italy (Delle Cave 1996, # 9): Procaccini Ricci 1840, Massalongo & Scarabelli-Gommi-Flamini 1859, Portis 1887, Olson 1977, Brodkorb 1978. [B*]
 Val Mala, Italy (Delle Cave 1996, # 6b): Delle Cave 1996. [F]
 Venta del Moro, Spain (Sánchez Marco 1996b, # 34): Aguirre et al. 1974, Morales & Aguirre 1976. [B]

MN 14

- Dorkovo, Bulgaria (Boev 1996, # 8): Boev 1988d, 1992, 1996. [B*]
 Kamenskoe, Ukraine (Mlíkovský 1996p, # 31): Vojinstvens'kyj 1967. [B]
 Montpellier, France (Mourer-Chauviré 1996b, # 1): Christol 1835, Serres 1838a,b, Serres et al. 1839, Gervais 1844a, 1849, 1848-1852, 1859, Milne-Edwards 1863, 1869-1871, Depéret 1897, Lambrecht 1933. [B]
 Osztramos 1, Hungary (Mlíkovský 1996j, # 7): Jánossy 1976b, 1979, Mlíkovský 1992b. [B*]
 Osztramos 9, Hungary (Mlíkovský 1996j, # 8): Jánossy 1979, Mlíkovský 1992b. [B*]
 Vojničevo, Ukraine (Mlíkovský 1996p, # 33): Burčák-Abramovič 1953a, Vojinstvens'kyj 1967, Bocheński & Kuročkin 1987a. [B, E]

MN 14-15

- Aegina, Greece (Mlíkovský 1996e, # 3): Lambrecht 1933: 689. [B]
 Biancone 1, Italy (Delle Cave 1996, # 13e): Ballmann 1973, 1976a. [B*]
 Biancone 2, Italy (Delle Cave 1996, # 13f): Ballmann 1973, 1976a. [B]
 Cantatore 3A, Italy (Delle Cave 1996, # 13g): Ballmann 1973, 1976a. [B]
 Chirò 2, 2N, 2S, Italy (Delle Cave 1996, # 13h,i,j): Ballmann 1973, 1976a. [B]
 Chirò 3, Italy (Delle Cave 1996, # 13k): Ballmann 1973, 1976a. [B]
 Chirò 4, Italy (Delle Cave 1996, # 13l): Ballmann 1973, 1976a. [B]




Fig. 12. Type localities of early Pliocene (MN 14-15) birds of Europe. 1 - Węże, 2 - Osztamos, 3 - Csarnóta, 4 - Malușteni-Berești, 5 - Odesa, 6 - Layna, 7 - Perpignan, 8 - Orciano Pisano (dating uncertain), 9 - Gargano (incl. Biancone, Chirò, San Giovannino), 10 - Lozenec, 11 - Dorkovo, 12 - Muselievo. Scale bar = 100 km.

Chirò 5, 5A, 5B, Italy (Delle Cave 1996, # 13m,n,o): Ballmann 1973, 1976a. [B]

Chirò 6, Italy (Delle Cave 1996, # 13p): Ballmann 1973, 1976a. [B]

Chirò 7A, 7C, Italy (Delle Cave 1996, # 13q,r): Ballmann 1973, 1976a. [B]

Chirò 9, Italy (Delle Cave 1996, #13s): Ballmann 1973, 1976a. [B]

Chirò 10A, 10C, Italy (Delle Cave 1996, # 13b,t): Ballmann 1973, 1976a. [B*]

Chirò 11A, 11B, Italy (Delle Cave 1996, # 13u,v): Ballmann 1973, 1976a. [B]

Chirò 12, Italy (Delle Cave 1996, # 13c): Ballmann 1973, 1976a. [B*]

Chirò 14A, 14B, Italy (Delle Cave 1996, # 13w,x): Ballmann 1973, 1976a. [B]

Chirò 15, Italy (Delle Cave 1996, # 13y): Ballmann 1973, 1976a. [B]

Chirò 20, 20A, 20B, 20C, 20D, 20E, Italy (Delle Cave 1996, # 13z,aa,bb,cc,dd,ee): Ballmann 1973, 1976a. [B]

Chirò 22, Italy (Delle Cave 1996, # 13 ff): Ballmann 1973, 1976a. [B]

Chirò 24, Italy (Delle Cave 1996, # 13d): Ballmann 1973, 1976a. [B*]

Chirò 27, Italy (Delle Cave 1996, #13gg): Ballmann 1973, 1976a. [B]

Empoli, Italy (new): Delle Cave et al. 1984. [B]

Falcone 2A, 2B, Italy (Delle Cave 1996, # 13hh,ii): Ballmann 1973, 1976a. [B]

Fina D, Italy (Delle Cave 1996, # 13jj): Ballmann 1973, 1976a. [B]

Fina E, Italy (Delle Cave 1996, # 13kk): Ballmann 1973, 1976a. [B]

Fina H, Italy (Delle Cave 1996, # 13ll): Ballmann 1973, 1976a. [B]

Gargano, Italy (Delle Cave 1996, # 13): Ballmann 1973, 1976a, Mlíkovský 1998c. This is a complex locality, composed of numerous fossiliferous fissures, which can be found under the following names: Biancone, Cantatore, Chirò, Falcone, Fina, Gervaisio, Monte Granata, Nazario, Pirro, Pizzioli, Posticchia, Rinascita, and San Giovannino. [B*]

Gervaisio 1, Italy (Delle Cave 1996, # 13mm): Ballmann 1973, 1976a. [B]
 Gervaisio 2, Italy (Delle Cave 1996, # 13nn): Ballmann 1973, 1976a. [B]
 Grotte, Italy (Delle Cave 1996, # 18): Delle Cave et al. 1984. [B]
 Kerč' Peninsula 2, Ukraine (Mlíkovský 1996q, # 32): Dubrovo & Kapelist 1979. [B]
 Lozenec, Bulgaria (new): Boev 1999a, 2000b. [B*]
 Monte Granata, Italy (Delle Cave 1996, # 13oo): Ballmann 1973, 1976a. [B]
 Nazario 2B, Italy (Delle Cave 1996, # 13pp): Ballmann 1973, 1976a. [B]
 Nazario 3, Italy (Delle Cave 1996, # 13qq): Ballmann 1973, 1976a. [B]
 Nazario 4, Italy (Delle Cave 1996, # 13rr): Ballmann 1973, 1976a. [B]
 Pirro 11A, 11C, Italy (Delle Cave 1996, # 13ss,t): Ballmann 1973, 1976a. [B]
 Pizzioli 4, Italy (Delle Cave 1996, # 13uu): Ballmann 1973, 1976a. [B]
 Posticchia 1B, Italy (Delle Cave 1996, # 13vv): Ballmann 1973, 1976a. [B]
 Rinascita 1, Italy (Delle Cave 1996, # 13ww): Ballmann 1973, 1976a. [B]
 San Giovannino, Italy (Delle Cave 1996, # 13a): Ballmann 1973, 1976a. [B*]
 Sofija 2, Bulgaria (Boev 1996, # 5): Boev 1996. [B]

MN 14-17

Gérce, Hungary (Mlíkovský 1996j, # 15): Fischer & Hably 1991. [F]
 Semenovka, Russia (Mlíkovský 1996m, # 4): Pidopličko & Goldin 1964. [E]

MN 15

Arquillo 3, Spain (Sánchez Marco 1996b, # 36): Adrover et al. 1978. [B]
 Ciupereni, Romania (Kessler 1996, # 16): Kessler 1979. [B]
 Csarnóta 1, Hungary (new): Kretzoi 1956, Jánossy 1979. [B]
 Csarnóta 2, Hungary (Mlíkovský 1996j, # 9): Jánossy 1976a, 1977, 1979, 1986, Mlíkovský 1992b, 1995b. [B*]
 Gundersheim, Germany (Mlíkovský & Hesse 1996, # 66): Heller 1936. [B]
 Ivanovce 1, Slovakia (Mlíkovský 1996n, # 2): P. Švec in Fejfar & Heinrich 1985, Mlíkovský 1996n, orig. [B]
 Layna, Spain (Sánchez Marco 1996b, # 35): Mourer-Chauviré & Sánchez Marco 1988 Sánchez Marco 1990a. [B*]
 Malușteni-Berești, Romania (Kessler 1996, # 15): Kessler 1979, 1984a, 1992. [B*]
 Megalo Emvolon, Greece (new): Boev & Koufos 2000. [B]
 Muselievo, Bulgaria (Boev 1996, # 7): Boev 1998e, 1999a, 2000a. [B*]
 Odesa – catacombs, Ukraine (Mlíkovský 1996p, # 34): Grycaj 1938, 1939, Zubareva 1939, 1948, Burčák-Abramovič 1939, 1953a,b, Tugarinov 1940a,b, Serebrov'skyj 1941b, Treťjakov 1941, Vojinstvens'kyj 1959, 1967, Kuročkin 1985, Michajlov 1988, Michajlov & Kuročkin 1988. [B*]
 Perpignan, France (Mourer-Chauviré 1996b, # 2): Depéret 1892, 1897, Gaillard 1908, 1939, Lambrecht 1933, Brodkorb 1964, 1978, Mourer-Chauviré 1989b, 1990. [B*]
 Sète, France (Mourer-Chauviré 1996b, # 3): Cuvier 1825, Serres 1830, 1838b, Gervais 1844a, 1848-1852, 1859, Milne-Edwards 1863, 1869-1871, Mourer-Chauviré & Sánchez Marco 1988. [B]
 Weże 1, Poland (Bocheński 1996, # 4): Jánossy 1974a, 1981, Bocheński 1991, Mlíkovský 1992b. [B*]

MN 15-16

Castelnuevo Berardenga Scalo, Italy (Delle Cave 1996, # 16): Delle Cave 1996. [B]
 Host'ovce 2, Slovakia (Mlíkovský 1996n, # 3): Ložek & Horáček 1992. [B]
 Orciano Pisano, Italy (Delle Cave 1996, # 14): Portis 1887, 1889, Regàlia 1902, 1907. This is a composite locality. [B*]

MN 16

Ardé, France (Mourer-Chauviré 1996b, # 4): Gervais 1844a, 1848-1852, 1859, Milne-Edwards 1863, 1869-1871, Lambrecht 1933, Gaillard 1939, Mourer-Chauviré 1989b, 1990. [B*]




Fig. 13. Type localities of late Pliocene (MN 16-18) birds of Europe. 1 - Foxhall, 2 - Beremend, Villány, 3 - Rębielice Królewskie, 4 - Ca Na Reia, 5 - Canet, 6 - Ardé, 7 - Chilhac, 8 - Montecarlo, 9 - Kisláng, 10 - Beremend, 11 - Slivnica, 12 - Väršec, 13 - Lučești, 14 - Etulia. See also Fig. 12 for Orciano Pisano, which is potentially late Pliocene in age. Scale bar = 100 km.

- Balaruc 2, France (Mourer-Chauviré 1996b, # 5): Mourer-Chauviré & Sánchez Marco 1988. [B]
- Beremend 5, Hungary (Mlíkovský 1996j, # 10): Kretzoi 1956, Jánossy 1976b, 1977. [B*]
- Beremend 15, Hungary (new): Jánossy 1987b, 1990b, 1991, 1992. [B*]
- Çişmikioi, Moldova (Mlíkovský 1996l, # 10): Bocheński & Kuročkin 1987b. [B]
- Escorihuela, Spain (Sánchez Marco 1996b, # 39): Sánchez Marco 1996b. [B]
- Espejos, Spain (Sánchez Marco 1996b, # 38): Adrover 1975. [B]
- Etulia, Moldova (Mlíkovský 1996l, # 9): Chozackij & Kuročkin 1966, Kuročkin & Chozackij 1972, Bocheński & Kuročkin 1987a,b, Michajlov & Kuročkin 1988, Michajlov 1988, Mourer-Chauviré 1990. [B*, E]
- Hajnáčka, Slovakia (Mlíkovský 1996n, # 4): P. Švec in Fejfar & Heinrich 1985, Mlíkovský 1996n. [B]
- Higueruelas, Spain (Sánchez Marco 1996b, # 42): Sánchez Marco 1996b. [B]
- Kotlovina, Ukraine (Mlíkovský 1996p, # 35): Kuročkin & Chozackij 1972, Bocheński & Kuročkin 1987b. [B]
- Lučești, Moldova (Mlíkovský 1996l, # 11): Bocheński & Kuročkin 1987a,b, Mourer-Chauviré 1989b, 1990. [B*]
- Moreda, Spain (Sánchez Marco 1996b, # 40): Sánchez Marco 1996b. [B]
- Orrios 3, Spain (Sánchez Marco 1996b, # 37): Adrover 1986. [B]
- Osztramos 7, Hungary (Mlíkovský 1996j, # 11): Jánossy 1976a,b, 1977, 1986, Mlíkovský 1992b, 1995b. [B]
- Puebla de Almoradiel, Spain (Sánchez Marco 1996b, # 41): Hernández-Pacheco 1914b. [E]

Rebiełice Królewskie 1, Poland (Bocheński 1996, # 5a): Jánossy 1974a, Mlíkovský 1992b, 1995b [B*]

Rebiełice Królewskie 2, Poland (Bocheński 1996, # 5b): Jánossy 1974a, Mlíkovský 1995b. [B]
Suvorovskoe, Moldova (new): Ganea 1972. [E]

Tatarești, Moldova (new): Ganea 1972 [E]

MN 16-17

Armaiolo, Italy (Delle Cave 1996, # 17): Delle Cave 1996. [B]

Beremend 4, Hungary (Tyrberg 1998, # HU-4): Kretzoi 1956, Jánossy 1977. Most of the bones from this site are late Pliocene in age, but early Pleistocene bones (MQ 1) are admixed (Kretzoi 1956). [B]

Dolinskoe, Ukraine (Mlíkovský 1996p, # 36): Dubrovo & Kapelist 1979. [B]

Foxhall, England (Mlíkovský 1996q, # 22): Lydekker 1886, 1891a,c,e, Newton 1891, Harrison & Walker 1978b. [B*]

Orford, England (new): Lydekker 1886, 1891a,b, Newton 1891, Harrison & Walker 1978b. [B]

Tourkobounia 1, Greece (new): Mourer-Chauviré 1980b, 1993. [B]

MN 16 – MQ 1

Beremend 1-3, Hungary (new): Lambrecht 1916b, 1933, Jánossy 1976b [B]

Bobila Ordis, Spain (Tyrberg 2001, # SP-104): Mayr & Gregor 1999. [B]

Montecarlo, Italy (Delle Cave 1996, # IT-19, Tyrberg 1998, # IT-121): Portis 1887, 1889, Lambrecht 1933, Brodkorb 1964. [B*]

Strette, Italy (Tyrberg 1998, # IT-124): Portis 1889. [B]

MN 17

Beremend 11, Hungary (Mlíkovský 1996j, # 14): Jánossy 1986. [B]

Koliňany 2, Slovakia (Mlíkovský 1996n, # 5): Mlíkovský 1992b, 1996n. [B]

Kryžanovka 1, Ukraine (Mlíkovský 1996p, # 37): Dubrovo & Kapelist 1979. [B]

Matveev Kurgan, Russia (Mlíkovský 1996m, # 2): Gromov 1948, Burčák-Abramovič 1953a. [B]

Obitočnoe, Ukraine (Mlíkovský 1996p, # 38): Vojinstvens'kyj 1967. [B]

Puebla de Valverde, Spain (Sánchez Marco 1996b, # 43): Adrover et al. 1974. [B]

Saint-Vallier, France (Mourer-Chauviré 1996b, # 8): Viret 1954, Mourer-Chauviré 1975a, 1989b, 1990. [B]

Šandalja 1 – layer 6, Croatia (Mlíkovský 1996d, # 2): Malez-Bačić 1979. [B]

Sant'Andrea, Italy (Delle Cave 1996, # 15): Delle Cave 1996. [B]

Schernfeld, Germany (Mlíkovský & Hesse 1996, # 68): Dehm 1962. [B]

Tegelen, Holland (Mlíkovský 1996i, # 1): Junge 1953. [B]

Väršec, Bulgaria (Boev 1996, # 9): Boev 1991, 1995a, 1996, 1997b, 1998a,b,f, 1999a,c,d,e,f, g,h, 2000e. [B*]

Včeláre 3, Slovakia (Mlíkovský 1996n; # 6): Horáček 1985. [B]

MN 18

Aldeby, England (Mlíkovský 1996q, # 23): Newton 1882a,b, 1887, Lydekker 1891d, Harrison 1979c. [B]

Almenara 1, Spain (Sánchez Marco 1996b, # 44): Mourer-Chauviré & Sánchez Marco 1988. [B]

Betfia 13, Romania (Tyrberg 1998, # RM-2): Kessler 1975. [B]

Binigaus, Menorca, Spain (Sánchez Marco 1996b, # 47): Mourer-Chauviré et al. 1980b, Alcover et al. 1981. [B]

Ca Na Reia, Eivissa, Spain (Sánchez Marco 1996b, # 48): Alcover 1989. [B*]

Canet, Mallorca, Spain (Sánchez Marco 1996b, # 45): Mourer-Chauviré et al. 1980b. [B*]

Chilhac 2b, France (Mourer-Chauviré 1996b, # 6a): Boeuf & Mourer-Chauviré 1992. [B]

Chilhac 3, France (Mourer-Chauviré 1996b, # 6b): Boeuf & Mourer-Chauviré 1992. [B*]

Chillesford, England (Mlíkovský 1996q, # 24; Tyrberg 1998, # UK-11): Harrison 1979d. [B]

Meda Gran, Spain (Sánchez Marco 1996b, # 49): Villalta 1965, Sánchez Marco 1996b. [B]




Fig. 14. Type localities of early Pleistocene (MQ 1) birds of Europe. 1 - Přezletice, 2 - Stránská skála, 3 - Včeláre, 4 - Beremend, Nagyharsányhegy, 5 - Betfia. Scale bar = 100 km.

- Kadzielnia 1, Poland (Bocheński 1996, # 7): Jánossy 1974a, Bocheński 1989. [B]
 Kielniki 3B, Poland (Bocheński 1996, # 6): Jánossy 1974a, Bocheński 1991. [B]
 Kisláng, Hungary (Mlíkovský 1996j, # 12): Kretzoi 1954a, 1955a, Jánossy 1986, Michajlov 1988, Michajlov & Kuročkin 1988. [B*, E]
 Liveneck, Russia (Mlíkovský 1996j, # 3, Tyrberg 1998, # RO-22): Jan'kova 1959. [B]
 Montoussé 5, France (Mourer-Chauviré 1996b, # 7): Clot et al. 1976a,b,. [B]
 Rippersroda 1, Germany (Mlíkovský & Hesse 1996, # 67): Giebel 1860, Brodkorb 1980, Mlíkovský 1992b. Age corrected following Heinrich (1990). [B*]
 Senèze, France (Mourer-Chauviré 1996b, # 9): Stehlin 1923, Lambrecht 1933, Gaillard 1939, Mourer-Chauviré 1989b, 1990. [B]
 Slivnica, Bulgaria (Boev 1996, # 10): Boev 1995a, 1998b,c. [B*]
 S'Onix, Mallorca, Spain (Sánchez Marco 1996b, # 46): Mourer-Chauviré et al. 1977, 1980b, Sondaar et al. 1995. [B]
 Tourkobounia 2, Greece (new): Mourer-Chauviré 1980b. [B]
 Villány 3, Hungary (Mlíkovský 1996j, # 13): Jánossy 1974b, 1976a,b, 1977, 1979, 1980, Mlíkovský 1981, 1982a, 1995b. [B*]

MQ 1a

- Balša, Bulgaria (new): Boev, in litt. [B]
 Beremend 16, Hungary (Tyrberg 1998, # HU-5 partim): Jánossy 1992, 1996. [B*]
 Beremend 17, Hungary (Tyrberg 1998, # HU-5): Jánossy 1992, 1996. [B]
 Betfia 1, Romania (Tyrberg 1998, # RM-2): Jánossy 1976a. [B]
 Betfia 2, Romania (Tyrberg 1998, # RM-2): Čapek 1917, Lambrecht 1933, Kretzoi 1941, 1962, Jánossy 1972, 1976b, 1977, 1979, 1980a. [B*]

Čerdženica – trench 6, Bulgaria (Tyrberg 2001, # BU-10): Boev 2000c. [B]
 Deutsch-Altenburg 2C₁, Austria (Tyrberg 1998, # AU-2): Jánossy 1981, Mlíkovský 1998d. [B]
 Deutsch-Altenburg 4B, Austria (Tyrberg 1998, # AU-2): Jánossy 1981, Mlíkovský 1998d. [B]
 Huéscar 1, Spain (new): Sánchez Marco 1989. [B]
 Kairy, Ukraine (Tyrberg 1998, UR-32): Vojinstvens'kyj 1967. [B]
 Kőröshegy, Hungary (Tyrberg 1998, # HU-26): Kretzoi & Krolopp 1977. [B]
 Kryžanovka, Ukraine (Tyrberg 1998, # UR-36): Dubrovo & Kapelist 1979. Note: Two horizons are distinguished: lower (Kryžanovka 1) and upper (Kryžanovka 2). [B]
 Mas Rambault, France (Tyrberg 1998, # FR-196): Mourer-Chauviré 1975a. [B]
 Osztramos 2, Hungary (Tyrberg 1998, # HU-33): Jánossy 1976a,b,c, 1977, 1978, 1980, 1986, Jánossy & Kordos 1976a. [B]
 Osztramos 5, Hungary (Tyrberg 1998, # HU-34): Jánossy & Kordos 1977, Jánossy 1976b, 1977. [B]
 Osztramos 8, Hungary (Tyrberg 1998, # HU-35): Jánossy 1976a,b,c, 1977, 1986, Jánossy & Kordos 1977. [B]
 Primorsk, Ukraine (new): Vojinstvens'kyj 1967. [B]
 Quibas, Spain (Tyrberg 2001, # SP-101): Montoya et al. 1999, 2001. [B]
 Tarchankut, Ukraine (Tyrberg 1998, # UR-35): Vojinstvens'kyj 1967. [B]
 Untermassfeld, Germany (Tyrberg 1998, # BRD-141): Jánossy 1988, 1997. [B]
 Valerots, France (Tyrberg 1998, # FR-189): Mourer-Chauviré 1975a. [B]
 Vallonnet, France (Tyrberg 1998, # FR-187): Mourer-Chauviré 1975a. [B]
 Żabia Cave, Poland (Tyrberg 1998, # PO-26): Bosák et al. 1982, Mlíkovský orig. [B]
 Ževachova Hill, Ukraine (Tyrberg 1998, # UR-33): Dubrovo & Kapelist 1979. [B]

MQ 1a-b

Betfia 7, Romania (Tyrberg 1998, # RM-2): Kessler 1975. [B]
 Betfia 10, Romania (new): Terzea & Jurcsák 1968. [B]
 Betfia – Aven, Romania (Tyrberg 1998, # RM-2): Kessler 1978. [B]
 Gènova, Mallorca, Spain (Tyrberg 1998, # SP-63): Mourer-Chauviré et al. 1977. [B]
 Kunino, Bulgaria (new): Boev 2001. [B]
 Monte Argentario, Italy (Tyrberg 1998, # IT-78): Cassoli 1980. [B]
 Zalesiaki 1A, Poland (Tyrberg 1998, # PO-28): Bocheński 1989. [B]

MQ 1b

Aldeby, England (Tyrberg 1998, # UK-1): Newton 1882b, Harrison 1979f. [B]
 Bacton, England (Tyrberg 1998, # UK-4): Harrison 1979f, 1985b. [B]
 Belle-Roche, Belgium (Tyrberg 1998, # BE-1): Cordy 1993. [B]
 Betfia 5, Romania (Tyrberg 1998, # RM-2): Kretzoi 1962, Jurcsák & Kessler 1973, Kessler 1975, Mourer-Chauviré 1975a, Jánossy 1976a, 1980. [B*]
 Betfia 7 (Tyrberg 1998, # RM-2): Kessler 1975. [B]
 Bourgade, France (Tyrberg 1998, # FR-51): Mourer-Chauviré 1976, 1980b. [B]
 Boxgrove, England (Tyrberg 2001, # UK-75): Harrison & Stewart 1999. [B]
 Červený Hill (new): Mlíkovský 1996t. [B]
 Chlum 6, Czechia (Tyrberg 1998, # CZ-54): Mlíkovský 1996t. [B]
 East Runton, England (Tyrberg 1998, # UK-23): Newton 1882b, Bell 1915, Harrison 1979f, 1985b. [B]
 Erpfingen, Germany (Tyrberg 1998, # BRD-30): Jánossy 1976a,b. [B]
 Holštejn 1, Czechia (Tyrberg 1998, # CZ-10): Musil 1966. [B]
 Koněprusy C-718, Czechia (Tyrberg 1998, # CZ-19 and CZ-55): Mlíkovský 1996t, see also Jánossy 1972, 1976c, 1977a, 1979. [B]
 Kövesvárad, Hungary (Tyrberg 1998, # HU-28): Jánossy 1963, 1977, 1986. [B]
 Kozi Grzbiet, Poland (Tyrberg 1998, # PO-10): Bocheński 1984. [B]




Fig. 15. Type localities of middle (MQ 2A-B) Pleistocene birds of Europe. 1 - Tornewton, West Runton, 2 - Ostend, 3 - Fage, 4 - Lunel-Viel, 5 - Saint-Estève-Janson, 6 - Lazaret, 7 - Castiglione, 8 - Hundsheim, 9 - Várhegy, 10 - Tarkó, 11 - Petralona. See Fig 16 for Monte Reale, which is potentially of this age. Scale bar = 100 km.

Nagyharsányhegy 1-4, Hungary (Tyrberg 1998, # HU-31 and HU-32): Lambrecht 1916b, Mourer-Chauviré 1975a, Jánossy 1976a,b, 1977, 1979, 1980a. Most bones from the early collections at Nagyharányhegy cannot be assigned to any one of the four fissures known at that time (see Kretzoi 1956, Jánossy 1986). [B*]

Overstrand, England (Tyrberg 1998, # UK-50): Harrison 1979f, 1985b. [B]

Petralona – layers 11, 13, 15, 24, 26, and "mausoleum", Greece (Tyrberg 1998, # GR-13): Kretzoi 1977, Kretzoi & Poulianos 1981, Poulianos 1996. The age is uncertain; these layers may well be early MQ 2 in age. [B]

Podumci, Croatia (Tyrberg 1998, # CR-9): Malez 1973, 1986. [B]

Přezletice, Czechia (Tyrberg 1998, # CZ-33): Jánossy 1982, 1983, Mlíkovský 1997c. [B*]

Razvođe, Croatia (Tyrberg 1998, # CR-10): Malez 1973, 1984b, 1986. [B]

Sackdilling, Germany (Tyrberg 1998, # BRD-154): Brunner 1933, Lambrecht 1933, Jánossy 1976a,b. [B]

Sima del Elefante, Spain (Tyrberg 2001, # SP-111): Rosas et al. 2001. [B]

Soave, Italy (Tyrberg 1998, # IT-22): Mourer-Chauviré 1980b, 1993b. [B]

Somssichhegy 2, Hungary (Tyrberg 1998, # HU-46): Toulou 1910, Jánossy 1983, 1986. [B]

Stránská skála – Absolon's 2nd cave, Czechia (Tyrberg 1998, # CZ-41): Skutil & Stehlík 1939, Musil 1965, Musil & Valoch 1968, Jánossy 1972. [B*]

Stránská skála – Čapek's site 4, Czechia (Tyrberg 1998, # CZ-41): Čapek 1925, Skutil & Stehlík 1939, Musil 1965, Musil & Valoch 1968, Jánossy 1972. [B]

Stránská skála – Čapek's site 5, Czechia (Tyrberg 1998, # CZ-41): Čapek 1925, Skutil & Stehlík 1939, Musil 1965, Musil & Valoch 1968, Jánossy 1972. [B]

- Stránská skála – Čapek's site 8, Czechia (Tyrberg 1998, # CZ-41): Čapek 1925, Skutil & Stehlík 1939, Musil 1965, Musil & Valoch 1968, Jánossy 1972. [B]
- Stránská skála – Čapek's site 9, Czechia (Tyrberg 1998, # CZ-41): Čapek 1925, Skutil & Stehlík 1939, Musil 1965, Musil & Valoch 1968, Jánossy 1972. [B]
- Stránská skála – Čapek's sites 1-3, Czechia (Tyrberg 1998, # CZ-41): Čapek 1925, Skutil & Stehlík 1939, Musil 1965, Musil & Valoch 1968, Jánossy 1972. These three sites cannot be separated from each other any more (Musil 1965, 1972). [B]
- Stránská skála – Musil's talus cone, layers 3-16, Czechia (Tyrberg 1998, # CZ-41): Mlíkovský 1995c, Musil 1995b. [B*, E]
- Stránská skála, Czechia (Tyrberg 1998, # CZ-41): (Čapek 1925, Skutil & Stehlík 1939, 1944, Musil 1965, 1995a,b, Musil & Valoch 1968, Jánossy 1972, Mlíkovský 1995b. This is a complex locality, consisting of several fissures and a talus cone, as follows: Čapek's sites 1-3, 4, 5, 7, 8, 9, Absolon's cave, Musil's talus cone. [B*, E]
- Tatinja Draga, Croatia (Tyrberg 1998, # CR-14): Malez 1973, 1984b. [B]
- Uppony 1 – layer 10, Hungary (Tyrberg 1998, # HU-58): Jánossy 1979, 1986. [B]
- Üromhegy, Hungary (new): Jánossy 1986. [B]
- Včeláre 1, Slovakia (Tyrberg 1998, # SK-5): Kretzoi 1975, Jánossy 1969, 1976a,b, 1977, 1979, Horáček 1985, Ložek & Horáček 1992. [B*]
- Včeláre 4E, Slovakia (new): Mlíkovský orig. [B]
- Venta Micena, Spain (new): Agustí et al. 1984, Sánchez Marco 1999a. [B]
- Villány 6, Hungary (new): Kretzoi 1956. [B]
- Villány 8, Hungary (new): Kretzoi 1956, Jánossy 1986. [B]
- Voigtstedt, Germany (Tyrberg 1998, # BRD-144): Jánossy 1965, 1976a, 1980a, Mlíkovský 1982c. [B]
- West Runton – Upper Freshwater Bed, England (Tyrberg 1998, # UK-68): Newton 1882a,b, 1887, 1891, Lydekker 1891a, Bell 1915, Harrison 1979d, Mlíkovský 1982c. [B*]

MQ 2A

- Ambroña, Spain (Tyrberg 1998, # SP-5): Jánossy 1983, Sánchez Marco 1990b, 1999c. [B]
- Arago, France (Tyrberg 1998, # FR-59): Mourer-Chauviré 1975a. [B]
- Čertkov 2, Ukraine (Tyrberg 1998, # UR-21): Vojinstvens'kyj 1967. [B]
- Cimay, France (Tyrberg 1998, # FR-67): Mourer-Chauviré 1975a. [B]
- Gombasek, Slovakia (Tyrberg 1998, # SK-4): Jánossy 1976a, 1980b. [B]
- Hundsheim, Austria (Tyrberg 1998, # AU-7): Freudenberg 1914, Jánossy 1974b, 1977, Mlíkovský 1998g. [B*]
- Monte Peglia, Italy (new): Cassoli 1980. [B]
- Montoussé 3, France (Tyrberg 1998, # FR-202): Clot et al. 1976a. [B]
- Montoussé 4, France (Tyrberg 1998, # FR-203): Clot et al. 1976a. [B]
- Mosbach, Germany (Tyrberg 1998, # BRD-94): Mourer-Chauviré 1977, Mlíkovský 1998g. [B]
- Ostend, England (Tyrberg 1998, # UK-49): Lydekker 1891a, Harrison 1978, 1979d, 1985b. [B*]
- Petalona – layers 2, 8 and 10, Greece (Tyrberg 1998, # GR-13): Kretzoi 1977, Kretzoi & Poulianos 1981, Poulianos 1996. [B*]
- Saint-Estève-Janson, France (Tyrberg 1998, # FR-157): Mourer-Chauviré 1975a. [B*]
- Spinagallo, Italy (Tyrberg 1998, # IT-123): Pavia 1999b. [B]
- Tarkő, Hungary (Tyrberg 1998, # HU-55): Jánossy 1972, 1974b, 1976a,c, 1977, 1979, 1980a, 1986. [B*]
- Terra Amata (Tyrberg 1998, # FR-242): Mourer-Chauviré 1975a. [B]
- Tourkobounia 5, Greece (Tyrberg 1998, # GR-19): Mourer-Chauviré 1980b. [B]
- Uppony 1 – layers 1, 3, 6 and 7, Hungary (Tyrberg 1998, # HU-58): Jánossy 1977. [B]
- Várhegy – Fortuna Street 25, Hungary (Tyrberg 1998, # HU-62): Jánossy 1969, 1976a,b, 1979, 1986. [B]




Fig. 16. Type localities of late Pleistocene (MQ 2C-D) and Holocene (MQ E) birds of Europe. 1 - Reach, 2 - Goyet, 3 - Nichet, 4 - Salzgitter-Lebenstedt, 5 - Westeregeln, 6 - Seveckenberg, 7 - Breitenberghöhle, 8 - Schusterlucke, 9 - Repolusthöhle, 10 - Trinka, 11 - Brynzeny, 12 - Eyzies, 13 - Massat, 14 - Figari, Pietro Tampoia Cave, 15 - Monte Reale (age uncertain), 16 - Ghar Dalam, Zebbug, 17 - Liko, 18 - Sozopol, 19 - Medvež'ja Cave (outside the picture at NE Ural Mountains). Not exactly localized site: Peschiera, Italy. Scale bar = 100 km.

Várhegy – Uri Street 72, Hungary (Tyrberg 1998, # HU-63): Jánossy 1969. [B*]

Vértesszőllős 2, Hungary (Tyrberg 1998, # HU-65): Jánossy 1976a,b, 1977, 1979, 1986, 1990a, Mlíkovský 1998g. [B]

MQ 2A-B

Rapaci, Sardinia, Italy (Tyrberg 1998, # IT-40): Mourer-Chauviré & Weesie 1986. [B]

MQ 2A-C

Capins, France (Tyrberg 1998, # FR-257): Mourer-Chauviré et al 1975. [B]

Monte Reale, Sardinia, Italy (Tyrberg 1998, # IT-122): Wagner 1828, 1832, 1833, Marmora 1831, Studiati 1857, Milne-Edwards 1869-1871, Mlíkovský 1995a, 2000b. [B*]

Son Bauçà, Mallorca, Spain (Tyrberg 1998, # SP-80): Ballmann & Adrover 1970, Mourer-Chauviré et al. 1980. [B]

MQ 2B

Aldène, France (Tyrberg 1998, FR-30): Mourer-Chauviré 1975a. [B]

Aridos 1, Spain (Tyrberg 1998, # SP-9): Mourer-Chauviré 1980c. [B]

Atapuerca, Spain (Tyrberg 1998, # SP-10): Sánchez Marco 1987a,b, 1995, 1999d. [B]

Azé 2, France (Tyrberg 1998, # FR-216): Mourer-Chauviré 1975a. [B]

Biehle, France (Tyrberg 1998, # FR-54): Mourer-Chauviré 1975a,b. [B]

Carrière, France (Tyrberg 1998, # FR-126): Mourer-Chauviré 1975a. [B]

Castiglione 3, Corsica, France (Tyrberg 2001, # FR-259): Mourer-Chauviré et al. 1997, Salotti et al. 1997. [B*]

- Combe Grenal, France (Tyrberg 1998, # FR-69): Mourer-Chauviré 1975a. [B]
 Fage, France (Tyrberg 1998, # FR-171): Mourer-Chauviré 1975a,b,c. [B*]
 Fournier Quarry, France (Tyrberg 1998, # FR-56): Mourer-Chauviré 1975a. [B]
 Hórvölgy, Hungary (Tyrberg 1998, # HU-19): Jánossy 1976a, 1977, 1979. [B]
 Hunas, Germany (Tyrberg 1998, # BRD-66): Jánossy 1983b. [B]
 Lazaret, France (Tyrberg 1998, # FR-147, 148): Mourer-Chauviré 1975a, Vilette 1994. Several sites within the cave yielded bird bones. [B*]
 Lunel-Viel, France (Tyrberg 1998, # FR-197): Serres et al. 1839, Milne-Edwards 1869-1871, Mourer-Chauviré 1975a. [B*]
 Orgnac 3, France (Tyrberg 1998, # FR-211): Mourer-Chauviré 1975a. [B]
 Saint-Sol-Belcastel, France (Tyrberg 1998, # FR-164): Philippe et al. 1980. [B]
 Suard, France (Tyrberg 1998, # FR-13): Mourer-Chauviré 1975a. [B]
 Tornewton, England (Tyrberg 1998, # UK-64): Harrison 1980b, Stewart 1996. [B*]
 Westbury-sub-Mendip, England (Tyrberg 1998, # 69): Andrews 1990. [B]

MQ 2C

- Archi, Italy (Tyrberg 1998, # IT-2): Ascenzi & Segre 1971, Cassoli & Segre 1985. [B]
 Avenc de Na Corna, Mallorca, Spain (Tyrberg 1998, # SP-11): Mourer-Chauviré et al. 1975, 1977, Northcote & Mourer-Chauviré 1988. [B]
 Balauzière, France (Tyrberg 1998, # FR-38): Mourer-Chauviré 1975a. [B]
 Benghisa, Malta (Tyrberg 1998, # MA-1): Lambrecht 1933. [B]
 Bordu Mare, Romania (Tyrberg 1998, RM-12): Jurcsák & Kessler 1988. [B]
 Brynzeny 1 – lower cultural layer, Moldova (Tyrberg 1998, # MO-1): Ganea & Ketraru 1965, Ganea 1969a,b, 1972. [B*]
 Butești – cultural layer, Moldova (Tyrberg 1998, # MO-2): Ganea 1972. [B]
 Calafrana, Malta (Tyrberg 1998, # MA-2): Lambrecht 1933. [B]
 Corbeddu, Sardinia, Italy (Tyrberg 1998, # IT-63): Mourer-Chauviré & Weesie 1986, Weesie 1999. [B]
 Coscia, France (Tyrberg 2001, # FR-260): Bonifay et al. 1998. [B]
 Crayford, England (Tyrberg 1998, # UK-16): Harrison & Walker 1977b. [B]
 Devetaška Cave, Bulgaria (Tyrberg 2001, # BU-11): Boev 1999a. [B]
 Devil's Tower, Gibraltar (Tyrberg 1998, # SP-53): Bate 1928. [B]
 Dorgali, Sardinia, Italy (Tyrberg 1998, # IT-15): Kotsakis 1980, Mourer-Chauviré & Weesie 1986. [B]
 Dürrloch, Germany (Tyrberg 1998, # BRD-26): Schlosser 1900. [B]
 Figari, Sardinia, Italy (Tyrberg 2001, # IT-127): Dehaut 1911, 1914, 1920, Comaschi Caria 1969, 1970, Pavia 1999a. [B*]
 Figueira Brava – layers 2-4, Portugal (Tyrberg 1998, # PG-3): Mourer-Chauviré & Antunes 1991, 2000. [B]
 Fontéchevade, France (Tyrberg 1998, # FR-84): Mourer-Chauviré 1975a. [B]
 Ghaqba, Malta (Tyrberg 1998, # MA-8): Northcote 1982a, 1984, 1992. [B]
 Ghar Dalam, Malta (Tyrberg 1998, # MA-4): Bate 1916, Fischer & Stephan 1974, Olson 1976b, Northcote 1992. [B*]
 Gnien, Malta (Tyrberg 1998, # MA-9): Harrison 1979, Harrison & Cowles 1977, Northcote 1983, 1984, 1985, 1992. [B]
 Gonvillard, France (Tyrberg 1998, # FR-42): Mourer-Chauviré 1975a. [B]
 Gorham's Cave, Gibraltar (Tyrberg 1998, # SP-64): Eastham 1968. [B]
 Goyet, Belgium (Tyrberg 1998, # BE-2): Friant 1950c, 1951, Mlíkovský 2000a. [B*]
 Grappin, France (Tyrberg 1998, # FR-96): Mourer-Chauviré 1975a. [B]
 Gumbes B, Greece (Tyrberg 1998, # GR-6): Weesie 1982, 1987a,b, 1988. [B]
 Gumbes C, Greece (Tyrberg 1998, # GR-7): Weesie 1982, 1987a,b, 1988. [B]
 Ilford, England (Tyrberg 1998, # UK-34): Bell 1915, Harrison & Cowles 1977, Harrison & Walker 1977c. [B]

- Kalafrana, Malta (new): Lambrecht 1933. [B]
- Kálmán Lambrecht Cave, Hungary (Tyrberg 1998, # HU-22): Jánossy 1964, 1977, 1986, 1989, Mlíkovský 1998g. [B]
- Kiik-Koba, Ukraine (Tyrberg 1998, # UR-9): Tugarinov 1937, Vojinstvens'kyj 1967, Baryšnikov & Potapova 1992. [B]
- Liko, Crete, Greece (Tyrberg 1998, # GR-10): Weesie 1982, 1987a,b, 1988. [B*]
- Macinaggio, Corsica, France (Tyrberg 1998, # FR-145): Mourer-Chauviré & Weesie 1986, Mourer-Chauviré in Tyrberg 1998: 136. [B]
- Mnajdra, Malta (Tyrberg 1998, # MA-6): Northcote 1982b, 1985, 1992. [B]
- Monte San Giovanni, Sardinia, Italy (Tyrberg 1998, # IT-79): Lydekker 1891b, Mourer-Chauviré & Weesie 1986. [B]
- Nichet – lower layer, France (Tyrberg 1998, # FR-118): Bastin 1933. [B*]
- Nižnekryvčansk, Ukraine (Tyrberg 1998, # UR-15): Marisova & Tatarinov 1962, Marisova 1968. [B]
- Petit Puymoyen, France (Tyrberg 1998, # FR-217): Mourer-Chauviré 1975a. [B]
- Prince, Italy (Tyrberg 1998, # IT-55): Boule 1910, Mourer-Chauviré 1975a. [B]
- Rapaci, Sardinia, Italy (Tyrberg 1998, # 39): Mourer-Chauviré & Weesie 1986. [B]
- Repolusthöhle, Austria (Tyrberg 1998, # AU-11): Mottl 1951, Jánossy 1989, Mlíkovský 1998g. [B*]
- Schusterlucke, Austria (Tyrberg 1998, # AU-16): Woldřich 1893, Mlíkovský & Lukáš 1991, Mlíkovský 1997d. [B*]
- Seveckenberg, Germany (Tyrberg 1998, # BRD 124): Giebel 1847, Nehring 1880, 1904. [B*]
- Simonelli, Crete, Greece (Tyrberg 1998, # GR-15): Suriano 1980, Weesie 1987a,b, 1988. [B]
- Soulabé (partim), France (Tyrberg 1998, # FR-240): Mourer-Chauviré 1975a. [B]
- Starýe Duruitorý – layers II-IV, Moldova (Tyrberg 1998, # MO-3): Ganea & Ketraru 1964, Ganea 1972. [B]
- Sungir, Moskva Province, Russia (Tyrberg 1998, # RO-35): Gromov 1966, Burčák-Abramovič 1972, 1975. [B]
- Ta'Kandja, Malta (Tyrberg 1998, # MA-3): Despott 1929, Lambrecht 1933. [B]
- Trinka 1 – middle Paleolithic layer, Moldova (Tyrberg 1998, # MO-4): Ganea 1969a, 1972. [B]
- Vindija, Croatia (Tyrberg 1998, # CR-19): Malez 1988. [B]
- Westeregeln, Germany (Tyrberg 1998, # BRD-150): Germar 1826, Kefenstein 1834, Giebel 1847, Nehring 1877, 1880, Mlíkovský 1998g. [B*]
- Zebbug, Malta (Tyrberg 1998, # MA-10): Parker 1865, 1869, Falconer 1868, Lydekker 1890, 1891a, Northcote 1982a, 1984, 1985, 1992. [B*]
- Zwergloch, Germany (Tyrberg 1998, # BRD-156): Ranke 1879. [B]
- MQ 2D**
- Arene Candide, Italy (Tyrberg 1998, # IT-3): Cassoli 1980. [B]
- Blomvåg, Norway (Tyrberg 1998, # NO-1): Lie 1986. [B]
- Breitenberghöhle, Germany (Tyrberg 1998, # BRD-14): Brunner 1958. This locality may contain also remains from older and younger strata (see Tyrberg 1998: 205). [B*]
- Castiglione 1, Corsica, France (Tyrberg 2001, # FR-258): Mourer-Chauviré et al. 1997. [B]
- Cave 16, Bulgaria (Tyrberg 1998, BU-2): Boev 1999a. [B]
- Colombière, France (Tyrberg 1998, FR-169): Mourer-Chauviré 1975a. [B]
- Cosquer, France (Tyrberg 1998, # FR-94): Clottes et al. 1992, Clottes & Courtin 1993, d'Errico 1994a,b,c, McDonald 1994. [B]
- Duruthy, France (Tyrberg 1998, # FR-76): Delpech 1968, 1983. [B]
- Eyzies, France (Tyrberg 1998, # FR-138): Milne-Edwards 1869-1871, 1875a,b. [B*]
- Gargas, Jersey, England (new): Andrews 1920, Breuil & Begouën 1936, d'Errico 1994a,c, McDonald 1994. The record is invalid. [B]
- Gouërris, France (Tyrberg 1998, # FR-107): Saint-Périer 1927. [B]

- Gourdan, France (Tyrberg 1998, # FR-108): Clot & Mourer-Chauviré 1986. [B]
- Grand Baille, France (Tyrberg 1998, # FR-173): Mourer-Chauviré 1975a. [B]
- Happurg, Germany (Tyrberg 1998, # BRD-62): Hörmann 1913. [B]
- Harpons, France (Tyrberg 1998, # FR-141): Saint-Périer 1920, Mourer-Chauviré 1975a, Clot & Mourer-Chauviré 1986. [B]
- Hermanstorp, Sweden (Tyrberg 1998, # SV-3): Munthe 1914, Löppenthin 1952. [E]
- Hohen Vielchen, Germany (new): Soergel 1961. [B]
- Kozarnika, Bulgaria (Tyrberg 2001, # BU-9): Boev 1999a. [B]
- Lafaye, France (Tyrberg 1998, # FR-8): Mourer-Chauviré 1975a. [B]
- Madelaine, France (Tyrberg 1998, # FR-174): Milne-Edwards 1875a,b. [B]
- Massat, France (Tyrberg 1998, # FR-116): Milne-Edwards 1875a,b, Mourer-Chauviré 1975a. [B*]
- Medvež'ja Cave, Russia (Tyrberg 1998, # RO-26): Potapova 1986, 1990. [B*]
- Mège, France (Tyrberg 1998, # FR-9): Capitan et al. 1906. [B]
- Morin, France (Tyrberg 1998, # FR-206): Delpech 1983. [B]
- Nerja, Spain (Tyrberg 1998, # SP-41): Eastham 1986. [B]
- Pageyral, France (Tyrberg 1998, # FR-17): Rivière 1887, 1892. [B]
- Paglicci, Italy (new): Mezzena & Palma di Cesnola 1992. [B]
- Pendo, Spain (Tyrberg 1998, # SP-57): Alcalde del Río et al. 1912, Eastham 1968, Gonzales Morales 1980, d'Errico 1994a,b,c, McDonald 1994. [Wall engravings.]
- Pietro Tampoia Cave, Sardinia, Italy (Tyrberg 1998, # IT-37): Shufeldt 1896, Regàlia 1897, Mayaud & Schaub 1950. [B*]
- Pouàs, Eivissa, Spain (Tyrberg 1998, # SP-61): J.A. Alcover in Tyrberg 1998. [B]
- Raymonden, France (new): Hardy 1891, Breuil et al. 1909, Alcalde del Río et al. 1912, Lambrecht 1933, d'Errico 1994a,c, McDonald 1994. [Sculpture.]
- Raziška Cave, Bulgaria (Tyrberg 2001, # BU-15): Boev 1999a. [B]
- Romanelli, Italy (Tyrberg 1998, # IT-40): Regàlia 1904, Blanck 1927, 1928, Cassoli & Segre 1985, Cassoli 1992, Cassoli & Tagliacozzo 1997, Gál 1999. [B]
- Salzgitter-Lebenstedt, Germany (Tyrberg 1998, # BRD-116): Kleinschmidt 1953a,b, Mlíkovský 1998g. [B]
- Soulabé (partim), France (Tyrberg 1998, # FR-240): Mourer-Chauviré 1975a. [B]
- Temnata Cave, Bulgaria (Tyrberg 1998, # BU-4): Boev 1999a. [B]
- Trois Frères, France (Tyrberg 1998, # FR-246): Mourer-Chauviré 1975a. [B]
- Urriaga, Spain (Tyrberg 1998, # SP-83): Eastham 1985, 1989. [B]
- Veyrier, France (Tyrberg 1998, # FR-159): Rüttimeyer 1873. [B]
- Vieulac, France (new): Rivière 1887, 1892. [B]
- MQ 2E**
- Burwell, England (new): Milne-Edwards 1869-1871: 108, Lydekker 1891c, Bell 1915, Friant 1950a,b, Northcote 1980, 1982d. [B]
- Ehrenstein, Germany (new): Soergel 1955. [B]
- Figueira Brava – "upper layers", Portugal (new): Mourer-Chauviré & Antunes 2000. [B]
- Peschiera, Italy (new): Portis 1884, Lambrecht 1933. [B*]
- Reach, England (new): Milne-Edwards 1869-1871: 108, Lydekker 1891c, Bell 1915, Friant 1950a,b, Northcote 1980, 1982d. [B*]
- Somerset, England (new): Harrison & Cowles 1977. [B]
- Sozopol, Bulgaria (new): Boev 1995c, Boev & Karajvanova 1998. [B*]
- Strathclyde, Isle of Jura, Scotland (new): Harrison & Cowles 1977. [B]

Order Tinamiformes Huxley

Tinamomorphae Huxley, 1872 [Modern order.]

Lithornithiformes Houde, 1988: 19 [Type genus: *Lithornis* Owen, 1840.]

Family Tinamidae Gray

Tinamidae Gray, 1840 [Modern family.]

Lithornithidae Houde, 1988: 19 [Type genus: *Lithornis* Owen, 1840.]

Lithornithinae Houde: Mlíkovský, this paper [New rank.]

Distribution: Early Eocene (MP 8-9) of England and Denmark. Extralimital record of the subfamily is known from the late Paleocene to the middle Eocene of Montana and Wyoming (Houde 1988, see also Houde 1981, 1986, Witmer 1989).

Remarks: I see no reason for placing the lithornithids in a separate family or even order, as suggested by Houde (1988). The lithornithids seem to represent an ancient radiation of the Tinamidae, and should be separated at the subfamily level. All European forms of tinamids belong to the Lithornithinae.

Genus *Lithornis* Owen

Lithornis Owen, 1840: 162 [Type by monotypy: *Lithornis vulturinus* Owen, 1840.]

Pediorallus Harrison & Walker, 1977a: 36 [Type by original designation: *Pediorallus barbarae* Harrison & Walker, 1977a.]

Promusophaga Harrison & Walker, 1977a: 40 [Type by original designation: *Promusophaga meini* Harrison & Walker, 1977a.]

Distribution: Early Eocene (MP 8-9) of England. Extralimital record is available from the late Paleocene of Montana (Houde 1988), and the early Eocene of Wyoming (Houde 1988).

Remarks: *Lithornis* Owen was described as a member of the Vulturidae, then comprising both New World and Old World vultures (Owen 1840, 1841, 1846). Cracraft & Rich (1972) doubted the vulturine relationships of the genus, Harrison & Walker (1977a) transferred it to the Threskiornithidae, but Houde (1988) showed, that it belongs to a separate family, related to the modern Tinamidae (here included in the Tinamidae as a subfamily).

Pediorallus Harrison & Walker was described in the Rallidae, and *Promusophaga* Harrison & Walker as a member of the Musophagidae (Harrison & Walker 1977a). Houde (1988) synonymized both these genera with *Lithornis* Owen.

Lithornis vulturinus Owen

Lithornis vulturinus Owen, 1840: 162 [Holotype from Sheppey: associated partial sternum, thoracic vertebra, ribs, distal end of left femur, and proximal end of left tibiotarsus; destroyed during World War II (Rich & Cracraft 1972: 281, Harrison & Walker 1977a: 29). Figured by Owen 1841, pl. 21, fig. 5-6, Owen 1846, fig. 232, Lambrecht 1933, fig. 134, Harrison & Walker 1977a, fig. 12, and Houde 1988, fig. 32. Neotype from Sheppey (designed by Houde 1988: 25): partial skeleton, incl. 5 cervical vertebrae, 1 thoracic vertebra, right scapula, right half of sternum, ribs, right humerus, radius and ulna, all lacking distal ends, distal ends of left radius and ulna, proximal end of left femur, and proximal end of right tibiotarsus; BMNH 38934 (formerly BMNH A-5204). Figured by Houde 1988, fig. 33.]

Pediorallus barbarae Harrison & Walker, 1977a: 36 [Holotype from High Ongar: proximal end of right tarsometatarsus; BMNH A-2681. Figured by Harrison & Walker 1977a, text-fig. 13a-f.]

Promusophaga magnifica Harrison & Walker, 1977a: 41 [Holotype from Sheppey: associated distal end of right coracoid, proximal end of left humerus, incomplete sternum, and ribs;

BMNH 33138. Figured by Lambrecht 1933, fig. 134a (sub *Lithornis vulturinus*), and Harrison & Walker 1977a, text-fig. 14a-b, pl. 9, fig. i-m.]

Pediorallus nasi Harrison 1984b:19 [Holotype from Walton-on-the-Naze: associated proximal end of left humerus, proximal end of right ulna, distal end of right femur, distal end of right tibiotarsus, and two vertebrae; BMNH A-5200. Figured by Harrison 1984b, fig. 3a-e, 4a-e, 5a-b.]

Lithornis nasi (Harrison): Houde 1988: 31 [New combination.]

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Harrison 1984b, Houde 1988), and Mariager Fjord, Denmark (Hoch 1975 sub Galliformes indet; was tentatively assigned to this species by Houde 1988); and early Eocene (MP 8-9) of Sheppey, England (Owen 1840, 1841, 1846, Hunter 1845, Lydekker 1891a, Cracraft & Rich 1972, Harrison & Walker 1977a [also sub *Neptuniavis minor* and *Parvigyps praecox*], Harrison 1984b, Houde 1988), and Saint James Park, England (Lydekker 1891a: 362 sub *Proherodius oweni*, Houde 1988).

Remarks: *Pediorallus barbarae* Harrison & Walker (described in the Rallidae), and *Promusophaga magnifica* Harrison & Walker (described in the Musophagidae) were synonymized with *Lithornis vulturinus* Owen by Houde (1988). *Lithornis nasi* (Harrison) was said to differ from *Lithornis vulturinus* Owen by size alone (Houde 1988), but Houde's tables of measurements convincingly show, that this is not the case. Hence, I synonymize here *Pediorallus* [= *Lithornis*] *nasi* Harrison with *Lithornis vulturinus* Owen.

***Lithornis hookeri* (Harrison)**

Pediorallus hookeri Harrison, 1984b: 21 [Holotype from Warden Point on the Isle of Sheppey: distal end of left tibiotarsus; BMNH A-5202. Figured by Houde 1988, fig. 25e.]

Lithornis hookeri (Harrison): Houde 1988: 32 [New combination.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison 1984b, Houde 1988), and Bognor Regis, England (Harrison & Walker 1977a sub *Pediorallus barbarae*, Houde 1988).

Order Struthioniformes Latham

Struthiones Latham, 1790 [Modern order.]

Family Palaeotididae Houde & Haubold

Palaeotididae Houde & Haubold, 1987: 32 [Type genus: *Palaeotis* Lambrecht, 1928.]

Distribution: Middle Eocene (MP 11-13) of Germany.

Genus *Palaeotis* Lambrecht

Palaeotis Lambrecht, 1928: 20 [Type by monotypy: *Palaeotis weigelti* Lambrecht, 1928.]

Palaeotis weigelti Lambrecht

Palaeotis Weigelti Lambrecht, 1928: 20 [Holotype from Cecilie I: distal end of left tarsometatarsus with associated phalanx 1 digiti II; GM 4415 (tarsometatarsus; present location unknown) and GM 4418 (phalanx). Figured by Lambrecht 1928, pl. 1, fig. 1-4, and Lambrecht 1933, fig. 157a.]

Palaeogrus geiseltalensis Lambrecht, 1935: 361 [Holotype from Cecilie I: associated right tibiotarsus, right tarsometatarsus, and two basal phalanges digitorum pedis; GM 5882. Figured by Lambrecht 1935, pl. 1, fig. 1. The phalanges were not described or figured by Lambrecht 1935.]

Ornitocnemus geiseltalensis (Lambrecht): Brodkorb 1967: 148 [New combination.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Houde & Haubold 1987, Peters 1988a), and middle Eocene (MP 13) of Geiseltal – Cecilie I, Cecilie II, and Leonhard III, Germany (Lambrecht 1928, 1933, Houde & Haubold 1987, Mlíkovský 1992b).

Remarks: *Palaeotis weigelti* Lambrecht was originally described as a bustard (Lambrecht 1928, 1933), while *Palaeogrus geiseltalensis* Lambrecht as a crane (Lambrecht 1935). *Palaeotis weigelti* Lambrecht was recognized as a paleognathous bird by Houde (1986, Houde & Haubold 1987), and *Palaeogrus geiseltalensis* Lambrecht was synonymized with it by Houde & Haubold (1987: 34).

Family Struthionidae Vigors

Struthionidae Vigors, 1825 [Modern family.]

Struthiolithidae Vjalov, 1971: 49 [Type genus: *Struthiolithus* Brandt, 1873. This family-group name is not available for nomenclatural purposes, because the genus name was created to denote fossil eggshell remains of *Struthio* Linnaeus; see Mlíkovský 2000d: 81.]

Genus *Struthio* Linnaeus

Struthio Linnaeus, 1758 [Modern genus.]

Struthiolithus Brandt, 1873: 158 [Type by monotypy: *Struthiolithus chersonensis* Brandt, 1873. Vjalov (1971: 44) argued that this name should be retained for the eggs of *Struthio* Linnaeus, which is unsubstantiated.]

Palaeostruthio Burčák-Abramovič, 1953a: 81 [Type by monotypy: *Palaeostruthio sternatus* Burčák-Abramovič, 1953a.]

Pachystruthio Kretzoi, 1954a: 233 [Type by monotypy: *Struthio pannonicus* Kretzoi, 1954a. Established as a subgenus of *Struthio* Linnaeus.]

Struthio chersonensis (Brandt)

Struthiolithus chersonensis Brandt, 1873: 158 [Holotype from Malynivcy: entire eggshell;

present location unknown. Not figured.]

Struthio novorossicus Alekseev, 1916: 388 [Syntypes from Novoelisavetovka: distal ends of three tarsometatarsi; GUN 1559-1561. Figured by Alekseev 1916, fig. 55-56, and Lambrecht 1933, fig. 43.]

Struthio chersonensis (Brandt): Lambrecht 1921: 2 [New combination.]

Struthio brachydactylus Burčák-Abramovič, 1939: 95 [Not seen. Probably based on the same material as *S. brachydactylus* Burčák-Abramovič, 1949, which would then be a younger objective synonym of this species. Note, that Burčák-Abramovič (1949: 141) listed *Struthio brachydactylus* Burčák-Abramovič, 1939 in the synonymy of *Struthio brachydactylus* Burčák-Abramovič, 1949.]

Struthio brachydactylus Burčák-Abramovič, 1949: 141 [Holotype from Hrebenyky: associated skull, 10 cervical vertebrae (1-2, 11-18), pelvis, left tibiotarsus with co-ossified left fibula, left tarsometatarsus, phalanges 1-4 digiti pedis III, and phalanges 1-4 digiti pedis IV; GMK 408/381(skull and cervical vertebrae), GMK 408/359 (tibiotarsus, fibula, tarsometatarsus, phalanges digiti pedis III), and GMK 408/352 (phalanges digiti pedis IV). All bones were transported away by German troops in 1943 (Burčák-Abramovič 1953a: 5) or in 1944 (Burčák-Abramovič 1949: 142); their present location is unknown. Figured by Burčák-Abramovič 1949, fig. 1 (skull), 2 (vertebrae 10-13), 3a (tibiotarsus and fibula), 3b (tarsometatarsus), 4 (phalanges pedis); Burčák-Abramovič 1953a, pl. 7, fig. 1 (skull), pl. 7, fig. 2-4 (vertebrae), pl. 8, fig. 1-2 (vertebrae), pl. 10, fig. 3-4 (tibiotarsus and fibula), pl. 10, fig. 5 (tibiotarsus), pl. 11, fig. 1-2 (tibiotarsus), pl. 12, fig. 1-4 (tarsometatarsus), pl. 13, fig. 1-6 (phalanges digiti III), pl. 14, fig. 1-4 (phalanges digiti III), pl. 14, fig. 5-7 (phalanges digiti IV), pl. 15, fig. 1-7 (phalanges digiti IV), pl. 15, fig. 8 (reconstructed foot); and Dement'ev 1964, fig. 670a (epistropheus), 670b,v (tibiotarsus), 670g,d (tarsometatarsus), and 670e,ž,z,i (pedal phalanges).]

Palaeostruthio sternatus Burčák-Abramovič, 1953a: 81 [Holotype from Hrebenyky: sternum; GIK 408/367. Figured by Burčák-Abramovič 1953a, pl. 18, fig. 1.]

Struthio (Pachystruthio) pannonicus Kretzoi, 1954a: 233 [Holotype from Kisláng: left pedal phalanx 1 digiti III; GIB, uncatalogued. Figured by Kretzoi 1954a, pl. 1, fig. 1., pl. 2, fig. 1-2.]

Struthiolithus alexejevi Roščin, 1962. [Holotype from Apostolovo: eggshell; present location unknown. Figured by Roščin 1962.]

Struthiolithus adzalycensis Roščin, 1962. [Holotype from Apostolovo: eggshell; present location unknown. Figured by Roščin 1959, p. 536.]

Struthio orlovi Kuročkin & Lungu, 1970: 119 [Holotype from Varnița: distal end of right tibiotarsus; TGPI 6-3. Figured by Kuročkin & Lungu 1970, text-fig. 1a,b,v, pl. 7, fig. 1.]

Distribution: Late Miocene (MN 9) of Varnița, Moldova (Lungu 1966a,b, Roška 1967, Kuročkin & Lungu 1970); late Miocene (MN 9-10) of Pokșești, Moldova (Michajlov 1988), Kolkotova Balka, Moldova (Michajlov 1988), Kubanka, Ukraine (Vojinstvens'kyj 1967), Tjaginka, Ukraine (Vojinstvens'kyj 1967), Jur'yvka, Ukraine (Pidopličko 1956), and Apostolovo, Ukraine (Alekseev 1916, Roščin 1959, 1962, Michajlov 1988);

late Miocene (MN 11) of Hrebenyky, Ukraine (Burčák-Abramovič 1939, 1949, 1951, 1953a,b), and Novoelisavetovka, Ukraine (Alekseev 1916, Burčák-Abramovič 1939, Vojinstvens'kyj 1967, Kuročkin & Lungu 1970); late Miocene (MN 11-13) of Čimișlia, Moldova (Michajlov 1988, Michajlov & Kuročkin 1988), Belka, Ukraine (Vojinstvens'kyj 1967), Žovten', Ukraine (Ganea & Kuročkin 1967), Il'inka, Ukraine (Burčák-Abramovič 1953, Vojinstvens'kyj 1967), Kujal'nik, Ukraine (Pržemysskij 1912, 1914, Alekseev 1916, Korotkevič 1961, Kuročkin & Lungu 1970), Nova Emetivka, Ukraine (Burčák-Abramovič 1939, 1953a, Ganea & Kuročkin 1967, Vojinstvens'kyj 1967, Burčák-Abramovič & Vekua 1971), Novoukrainka, Ukraine (Vojinstvens'kyj 1967), Snigirevka, Ukraine (Burčák-Abramovič 1953a), Čebotarevka, Ukraine (Vojinstvens'kyj 1967, Michajlov 1988), and Malynivcy, Ukraine (Brandt 1873, 1874, 1885, Mäklin 1873, Nathusius 1886, Bakulovskij

1892, Bensley 1921, Burčák-Abramovič 1953a, Schönwetter 1960: 8-10, Roščin 1962); late Miocene (MN 12) of Pikermi, Greece (Bachmayer & Zapfe 1962);

early Pliocene (MN 14) of Vojničevo, Ukraine (Burčák-Abramovič 1953a, Vojinstvens'kyj 1967); early Pliocene (MN 15) of Odesa – catacombs, Ukraine (Burčák-Abramovič 1953a,b, Vojinstvens'kyj 1967, Michajlov 1988, Michajlov & Kuročkin 1988);

late Pliocene (MN 16) of Etulia, Moldova (Michajlov 1988, Michajlov & Kuročkin 1988), Suvorovskoe, Molavia (Ganea 1972: 24), Tatarești, Moldova (Ganea 1972: 25), Kotlovina, Ukraine (Dubrovo & Kapelist 1979); late Pliocene (MN 16-17) of Dolinskoe, Ukraine (Dubrovo & Kapelist 1979); late Pliocene (MN 17) of Matveev Kurgan, Russia (Gromov 1948, Burčák-Abramovič 1953a), Liveneck, Russia (Jan'kova 1959); late Pliocene (MN 18) of Kisláng, Hungary (Kretzoi 1954a, 1955a, Jánossy 1986, Michajlov 1988, Michajlov & Kuročkin 1988);

undated Pliocene (MN 14-17) of Semenovka, Russia (Pidopličko & Goldin 1964); and doubtfully early Pleistocene (MQ 1a) of Kryžanovka 2, Ukraine (Dubrovo & Kapelist 1979).

Extralimital records of *chersonensis* from the Pleistocene of China (Eastman 1898a,b, Andersson 1923), based on eggshells, refer probably to another *Struthio* species (Lowe 1931, Lambrecht 1933).

Additional reference: Burčák-Abramovič & Kon'kova (1967).

Remarks: Until the opposite is proved, I believe that only one *Struthio* species inhabited Europe, that this species spread to it from the east in MN 9-10 and became extinct at the Plio-Pleistocene boundary. Brodkorb (1963: 197) included also *Struthio karatheodoris* Forsyth Major in the synonymy of *Struthio chersonensis* (Brandt), but I prefer to treat *karatheodoris* as a species separate from *chersonensis* for zoogeographical grounds. The areas of *Struthio chersonensis* (Brandt) and *Struthio karatheodoris* Forsyth Major were separated by Caucasus Mountains. Turkey and Greece were connected by land only in MN 13 (Steiniger & Rögl 1984), i.e. long after ostriches were recorded from both eastern Europe and southwestern Asia.

***Struthio karatheodoris* Forsyth Major**

Struthio karatheodoris Forsyth Major, 1888: 1178 [Holotype from Samos 1: femur. Said to be in coll. Barbey (Lambrecht 1933: 100); present location unknown. Figured by Martin 1903, fig. 30-31.]

Distribution: Late Miocene (MN 11) of Samos 1, Greece (Forsyth Major 1888, 1892, 1894, Lydekker 1891a, Martin 1903, Lowe 1931, Lambrecht 1933: 100-101, Solounias 1981).

Remarks: The island of Samos was connected with the Asian mainland during the late Miocene (Solounias 1981). Martin (1903) suggested, that *Struthio asiaticus* Milne-Edwards is a direct descendant of *Struthio karatheodoris* Forsyth Major. It is well possible, that *Struthio karatheodoris* Forsyth Major is identical with some or all of the Mio-Pliocene ostriches described from Siwalik Hills in NW India and adjacent parts of Pakistan (see Brodkorb 1963c: 197). If so, *Struthio asiaticus* Milne-Edwards, 1871 would be the oldest available name for the species. The case requires further study.

Order Phaethontiformes Sharpe

Phaethontes Sharpe, 1891 [Modern order.]

Gaviiformes Wetmore & Miller, 1926 [Modern order.]

Family Gaviidae Allen

Gaviidae Allen, 1897 [Modern family.]

Colymboididae Brodkorb, 1963c: 222 [Type genus: *Colymboides* Storer, 1956.]

Genus *Colymboides* Storer

Colymboides Milne-Edwards, 1867: 297 [Type by monotypy: *Colymboides minutus* Milne-Edwards, 1867.]

Hydrornis Milne-Edwards, 1867: 362 [Type by monotypy: *Hydrornis natator* Milne-Edwards, 1867b. This is a junior homonym of *Hydrornis* Blyth, 1843.]

Dyspetornis Oberholser, 1905: 60 [New name for *Hydrornis* Milne-Edwards, 1867.]

Distribution: Late Eocene (MP 17) of England, and early Miocene (MN 2-4) of France and Czechia.

Colymboides anglicus Lydekker

Colymboides anglicus Lydekker, 1891a: 43 [Holotype from Hordle: left coracoid; BMNH 303330. Figured by Lydekker 1891a, fig. 43, and Harrison & Walker 1976a, pl. 1, fig. m-p.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a: 192-193, Harrison 1976, Harrison & Walker 1976a).

Colymboides minutus Milne-Edwards

Colymboides minutus Milne-Edwards, 1867b: 297 [Holotype from Saint-Gérard-le-Puy: left humerus; MNHN Av-6581. Figured by Milne-Edwards 1867-1868, pl. 54, fig. 1-3. Cheneval (1984a: 40, 44) erroneously called this specimen lectotype.]

Hydrornis natator Milne-Edwards, 1867b: 362 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus. Figured by Milne-Edwards 1867-1868, pl. 57, fig. 18-22. Cheneval (1984a: 40) erroneously listed this specimen as a paralectotype of *Colymboides minutus* Milne-Edwards.]

Dyspetornis natator (Milne-Edwards): Oberholser, 1905: 60 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1867-1868, Storer 1956, Cheneval 1984a); early Miocene (MN 2-3) of Saucats, France (Lucazeau 1959); and early Miocene (MN 4b) of Dolnice, Czechia (Švec 1982, 1985).

Remarks: *Hydrornis natator* Milne-Edwards was synonymized with *Colymboides minutus* Milne-Edwards by Storer (1956: 423).

Genus *Gavia* Foster

Gavia Forster, 1788 [Modern genus.]

Gavia egeriana Švec

Gavia egeriana Švec, 1982: 251 [Holotype from Dolnice: distal end of left humerus; DP FNSP 4816. Figured by Švec 1982, pl. 1, fig. 1a, text-fig. 8a.]

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Švec 1982). Tentative extralimital records were reported from the middle Miocene of Virginia, Maryland, North Carolina and Delaware (Rasmussen 1998).

***Gavia schultzi* Mlíkovský**

Gavia schultzi Mlíkovský, 1998b: 332 [Holotype from Sankt Margarethen: right coracoid; NHMW 1986/92. Figured by Mlíkovský 1998b, pl. 1, fig. 1-3.]

Distribution: Middle Miocene (MN 7) of Sankt Margarethen, Austria (Mlíkovský 1998b).

***Gavia moldavica* Kessler**

Gavia moldavica Kessler, 1984b: 522 [Holotype from Chişinău: proximal end of right carpo-metacarpus; LPUB 273/1. Figured by Kessler 1984b, fig. 1-2.]

Distribution: Late Miocene (MN 9?) of Chişinău, Moldova (Kessler 1984b).

***Gavia paradoxa* Umans'ka**

Gavia paradoxa Umans'ka, 1981: 17 [Holotype from Čebotarevka: proximal end of left ulna; IZAN 45-4043. Figured by Umans'ka 1981, fig. 1a,b,v.]

Distribution: Late Miocene (MN 11-13) of Čebotarevka, Ukraine (Umans'ka 1981).

***Gavia* sp.**

Distribution: Middle Miocene (MN 7) of Steinheim, Germany (Heizmann & Hesse 1995: three different species); early Pliocene (MN 14-15) of Kerč' Peninsula 2, Ukraine (Kuročkin in Delle Cave et al. 1984: 89), and Empoli, Italy (Delle Cave et al. 1984 sub cf. *Gavia concinna*); and early Pleistocene (MQ 1a) of Kairy, Ukraine (Vojinstvens'kyj 1967). The record from the middle Eocene (MP 11-13) of Geiseltal, Germany (Mlíkovský in Delle Cave et al. 1984: 88) is invalid (Mlíkovský 1998b: 333).

Remarks: The record from Empoli (Delle Cave et al. 1984) is based on a skull, which was tentatively referred to *Gavia concinna* Wetmore, originally described from the late Miocene (late Hemphillian) of California (Wetmore 1940a), and later reported also from the late Miocene of Florida (Brodkorb 1953, Emslie 1995), and the late Pliocene of California (Howard 1949, Miller 1956). The Pliocene record from California is invalid according to Chandler (1990). The Italian skull fits the same element of the modern European *Gavia arctica* (Linnaeus, 1758) in size and morphology, and may belong to the latter species. There is no indication, that *Gavia concinna* Wetmore occurred in Europe.

Family Phaethontidae Brandt

Phaethontidae Brandt, 1840 [Modern family.]

Prophaethontidae Harrison & Walker, 1976c: 28 [Type genus: *Prophaethon* Andrews, 1899.]

Genus *Prophaethon* Andrews, 1899

Prophaethon Andrews, 1899: 776 [Type by monotypy: *Prophaethon shrubsolei* Andrews, 1899.]

Remarks: Some authors prefer to separate this genus at the family level (Harrison & Walker 1976c, 1977a, Olson 1985a).

***Prophaethon shrubsolei* Andrews**

Prophaethon shrubsolei Andrews, 1899: 776 [Holotype from Sheppey: associated bones, incl. imperfect skull, mandible, anterior fragment of sternum, distal end of left coracoid, left scapula, 11 vertebrae, synsacrum, right femur, proximal end of left femur, proximal end of right tibiotarsus; BMNH A-683. Figured by Andrews 1899, text-fig. 1-2, pl. 51, fig. 1-2, Harrison & Walker 1976c, 1977a, text-fig. 9-11, pl. 5, fig. A-E.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Andrews 1899, Harrison & Walker 1976c, 1977a).

Genus *Heliadornis* Olson

Heliadornis Olson, 1985b: 852 [Type by original designation: *Heliadornis ashbyi* Olson, 1985b.]

Distribution: Middle Miocene (MN 7-8) of Belgium and late Miocene (MN 10) of Austria. Extralimital record comes from the middle Miocene of Maryland (Olson 1985b).

***Heliadornis ashbyi* Olson**

Heliadornis ashbyi Olson, 1985b: 852 [Holotype from Parker Creek: associated proximal end of left humerus and proximal end of left scapula; USNM 237226. Figured by Olson 1985b, fig. 1a,c,e (humerus), and 1g,i (scapula).]

Distribution: Middle Miocene (MN 7-8) of Antwerp, Belgium (Olson & Walker 1997). An extralimital record is known from the middle Miocene (Langhian) of Parker Creek, Maryland (Olson 1985a, Olson & Walker 1997).

***Heliadornis paratethydicus* Mlíkovský**

Heliadornis paratethydicus Mlíkovský, 1997b: 152 [Holotype from Vösendorf: proximal end of right ulna; NHMW 1996/0184/0001. Figured by Mlíkovský 1997b, fig. 1.]

Distribution: Late Miocene (MN 10) of Vösendorf, Austria (Mlíkovský 1997b).

Order Ardeiformes Wagler

Ardeae Wagler, 1831 [Modern order.]

Family Sulidae Reichenbach

Sulidae Reichenbach, 1849 [Modern family.]

Pseudosulidae Harrison, 1975a: 53 [Type genus: *Pseudosula* Harrison, 1975a = *Enkurosula* Kašin, 1975a.]

Enkurosulidae Kašin, 1977: 139 [Type genus: *Enkurosula* Kašin, 1977.]

Remarks: Generic position of European Tertiary sulids is far from understood (Olson 1985a: 203-204). It is well possible, that at least some of the Oligo-Miocene sulids will be attributed to modern genera after a proper restudy.

If *Enkurosula* Kašin is separated at a suprageneric level, then the the family-group name Enkurosulidae Kašin has precedence over Pseudosulidae Harrison (Mlíkovský 2000d: 79).

Genus *Eostega* Lambrecht

Eostega Lambrecht, 1929: 1272 [Type by monotypy: *Eostega lebedinskyi* Lambrecht, 1929.]

Eostega lebedinskyi Lambrecht

Eostega Lebedinskyi Lambrecht, 1929: 1272 [Holotype from Cluj-Manastur: incomplete lower jaw; NHMW 1930/I/13. Figured by Lambrecht, 1929, fig. 12-13, Lambrecht 1933, fig. 103, and Jurcsák & Kessler 1973, fig. 8.]

Distribution: Middle Eocene (MP 13) of Cluj-Manastur, Romania (Lambrecht 1929, Kessler 1986, Mlíkovský in prep.).

Remarks: Lambrecht (1929) did not include the species in a family, indicating that it represents a "steganopode" (= a peleciform bird). Subsequently, the species was included in the family Elopterygidae (Lambrecht 1933, Brodkorb 1963c). I reexamined the fossil, concluding that it is a member of the Sulidae (Mlíkovský orig.).

Genus *Empheresula* Harrison

Parasula Harrison, 1975a: 53 [Type by original designation: *Sula arvernensis* Milne-Edwards, 1867b. This is junior homonym of *Parasula* Mathews, 1913.]

Empheresula Harrison, 1975b: 175 [Type by original designation: *Sula arvernensis* Milne-Edwards, 1867b.]

Empheresula arvernensis (Milne-Edwards)

Sula arvernensis Milne-Edwards, 1867b: 267 [Lectotype from Gannat (selected by Cheneval 1984a: 65): incomplete pelvis in slab; MNHN 1903-16. Figured by Milne-Edwards 1867-1868, pl. 43, fig. 12. Paralectotype from Gannat: incomplete sternum in slab; MNHN 1903-16. Figured by Milne-Edwards 1867-1868, pl. 42, fig. 13. Cheneval (1984a: 65) stated that the selection of the lectotype was done by Lydekker (1891a: 46), but the latter author just diagnosed *Sula arvernensis* on the basis of the morphology of its pelvis, not mentioning the syntypical sternum. This cannot be interpreted as a selection of a lectotype, so that this action has to be attributed to Cheneval (1984a) himself. Note that the catalogue number in NMNH is the same for both syntypes of *Sula arvernensis* Milne-Edwards and for *Tetrao pessieti* Gervais.]

Parasula arvernensis (Milne-Edwards): Harrison 1975a: 53 [New combination.]

Empheresula arvernensis (Milne-Edwards): Harrison 1975b: 175 [New combination.]

Distribution: Late Oligocene (MP 30) of Gannat, France (Milne-Edwards 1867-1868,

Lydekker 1891a, Harrison 1975a,b, Cheneval 1984a), and early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Cheneval 1984a, tentatively included).

Remarks: The paralectotypical sternum does not belong to the Sulidae according to Lambrecht (1933: 285).

Genus *Enkurosula* Kašín

Pseudosula Harrison, 1975a: 53 [Type by original designation: *Sula pygmaea* Milne-Edwards, 1874. This is a junior homonym of *Pseudosula* Boetticher, 1955]

Enkurosula Kašín, 1977: 139 [New name for *Pseudosula* Harrison, 1975a; hence its junior objective synonym.]

Enkurosula pygmaea (Milne-Edwards)

Sula pygmaea Milne-Edwards, 1874: 8 [Holotype from Léognan: left humerus; formerly in coll. Delfortrie, present location unknown. Figured by Milne-Edwards 1874, pl. 12, fig. 2.]

Microsula pygmaea (Milne-Edwards): Brodkorb 1963c: 259 [New combination.]

Pseudosula pygmaea (Milne-Edwards): Harrison 1975a: 53 [New combination.]

Enkurosula pygmaea (Milne-Edwards): Kašín 1977: 139 [New combination.]

Distribution: Early Miocene (MN 2-3) of Léognan, France (Milne-Edwards 1874, Harrison 1975a).

Genus *Sarmatosula* Grigorescu & Kessler

Sarmatosula Grigorescu & Kessler, 1977: 94 [Type by monotypy: *Sarmatosula dobrogensis* Grigorescu & Kessler, 1977.]

Sarmatosula dobrogensis Grigorescu & Kessler

Sarmatosula dobrogensis Grigorescu & Kessler, 1977: 94 [Holotype from Credița: proximal end of left humerus; LPUB 251. Figured by Grigorescu & Kessler 1977, pl. 1, fig. 1a-b, pl. 4, fig. B.]

Distribution: Middle Miocene (MN 8) of Credița, Romania (Grigorescu & Kessler 1977), and Ciobanița, Romania (Grigorescu & Kessler 1977).

Genus *Morus* Vieillot

Morus Vieillot, 1816 [Modern genus.]

Morus olsoni Grigorescu & Kessler

Morus olsoni Grigorescu & Kessler, 1988: 94 [Holotype from Credița: right carpometacarpus; LPUB 281. Figured by Grigorescu & Kessler 1998, fig. 2, 3a-b.]

Distribution: Middle Miocene (MN 8) of Credița, Romania (Grigorescu & Kessler 1988), and Ciobanița, Romania (Grigorescu & Kessler 1988).

Family Ardeidae Leach

Ardeidae Leach, 1820 [Modern family.]

Remarks: All Oligo-Miocene species included in this family require revision.

Genus *Proardeola* Harrison

Proardeola Harrison, 1979a: 14 [Type by original designation: *Proardeola walkeri* Harrison, 1979a.]

***Proardeola walkeri* Harrison**

Ardea formosa Milne-Edwards, 1871: 572 [Nomen nudum; no description or indication.]

Proardeola walkeri Harrison, 1979a: 14 [Holotype from Saint-Gérand-le-Puy: left tarsometatarsus; BMNH A-777. Figured by Harrison 1979a, fig. 2, and Cheneval 1984a, pl. 5, fig. 7a-b.]

Ardeola walkeri (Harrison): Mlíkovský & Švec 1989: 200 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérand-le-Puy, France (Harrison 1979a, Cheneval 1984a); and middle Miocene (MN 6) of Mátraszőlös 2, Hungary (Gál & Kessler in Gál et al. 2000).

Remarks: *Ardea formosa* Milne-Edwards was neither described nor figured by Milne-Edwards (1869-1871), hence it is a nomen nudum (see also Lydekker 1891a: vii, Lambrecht 1933: 312, Brodkorb 1978: 222; contra Cheneval 1984a: 77). Cheneval (1984a: 77-78) discovered in MNHN right coracoid (MNHN Av-2872, figured by Cheneval 1984a, pl. 5, fig. 6a-b) from Saint-Gérand-le-Puy, which was available to Milne-Edwards, and believed accordingly that *Ardea formosa* Milne-Edwards is an available name. This is not true (ICZN 1999, Art. 12.3), but Cheneval's discovery of the "type" allows to list *Ardea formosa* Milne-Edwards as a nomen nudum in the synonymy of *Proardeola walkeri* Harrison.

Proardeola Harrison is possibly synonymous with *Proardea* Lambrecht (C. Mourer-Chauviré in Olson 1985a: 167), and it is even possible that *Proardeola walkeri* Harrison is synonymous with *Ardea* (= *Proardea*) *amissa* Milne-Edwards (Mlíkovský orig.).

Genus *Proardea* Lambrecht

Proardea Lambrecht, 1933: 311 [Type by monotypy: *Ardea amissa* Milne-Edwards, 1892.]

***Proardea amissa* (Milne-Edwards)**

Ardea amissa Milne-Edwards, 1892: 73 [Holotype from Quercy: right tarsometatarsus; MNHN QU-15720. Not figured.]

Proardea amissa (Milne-Edwards): Lambrecht 1933: 311 [New combination.]

Egretta amissa (Milne-Edwards): Mlíkovský & Švec 1989: 200 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892).

Remarks: This species was said to belong in the subfamily Ardeinae (C. Mourer-Chauviré in Cheneval 1984a: 81), similarly as *Proardeola walkeri* Harrison. Both *Ardea amissa* Milne-Edwards and *Proardeola walkeri* Harrison were said to be rather similar to the modern *Ardeola* Boie, 1822, and their holotypical tarsometatarsi are similar in size. I suspect that further research will show that these two forms are conspecific.

Genus *Ardeagrandis* Kuročkin & Ganea

Ardeagrandis Kuročkin & Ganea, 1972: 51 [Type by original designation: *Ardeagrandis* [sic!] *arborea* Kuročkin & Ganea, 1972.]

***Ardeagrandis arborea* Kuročkin & Ganea**

Ardeagrandis arborea Kuročkin & Ganea, 1972: 52 [Holotype from Golbočica: proximal end of right tarsometatarsus; GIKM 12173/1. Figured by Kuročkin & Ganea 1972, text-fig. 3, pl. 1, fig. 4.]

Distribution: Late Miocene (MN 9) of Golbočica, Moldova (Kuročkin & Ganea 1972).

Genus *Ardea* Linnaeus

Ardea Linnaeus, 1758 [Modern genus.]

***Ardea aurelianensis* Milne-Edwards**

Ardea aurelianensis Milne-Edwards, 1871: 585 [Holotype from Orleanais: a humerus; present location unknown – Cheneval 1996: 604. Not figured.]

Distribution: Early Miocene (MN 4) of Orleanais, France (Milne-Edwards 1871).

Remarks: The holotypical humerus was said to be by 1/7 longer than the same element of the modern *Ixobrychus minutus* (Linnaeus, 1766), but still slightly shorter than that of the modern *Ardeola ralloides* (Scopoli, 1769) (Milne-Edwards 1871: 585). Hence *Ardea aurelianensis* falls in the same size class as *Proardea amissa* (Milne-Edwards) and *Proardeola walkeri* Harrison. It is possible that these three forms are conspecific.

***Ardea cinerea* Linnaeus – Grey Heron**

Ardea cinerea Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub. *A. cf. cinerea*).

***Ardea* sp.**

Remarks: Diaphysis of a tibiotarsus from the late Miocene (MN 9) of Kalfa, Moldova, was referred to *Ardea* sp. by Ganea (1965), but it represents an unidentified member of the Anatidae (Kuročkin & Ganea 1972).

Genus *Egretta* Forster

Egretta Forster, 1817 [Modern genus.]

***Egretta garzetta* Linnaeus – Little Egret**

Ardea Garzetta Linnaeus, 1766 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *E. cf. garzetta*).

Genus *Botaurus* Stephen

Botaurus Stephen, 1819 [Modern genus.]

***Botaurus stellaris* (Linnaeus) – Great Bittern**

Ardea stellaris Linnaeus, 1758 [Modern species.]

Nycticorax fenensis Friant, 1950a: 2126 [Lectotype from Reach (here selected): partial skeleton, incl. partial skull (SMC 753), mandible (SMC 790), coracoid, humerus, ulna, carpometacarpus, and femur; SMC Reach-1901. Figured by Friant 1950a, unnumbered fig. (skull and mandible), and Friant 1950b, fig. 2 (skull and mandible). Paralectotypes from Reach and Burwell: incomplete skeletons Reach-1902, incl. humerus and femur; Reach-1904, incl. coracoid, sternum, humerus, ulna, carpometacarpus, femur and tarsometatarsus; Burwell-1897, incl. skull, mandible (SMC 162), coracoid (SMC 149), sternum (SMC 136), ulna, carpometacarpus (SMC 146), femur, tibiotarsus and tarsometatarsus; Burwell-1900, incl. carpometacarpus, femur and tibiotarsus; Burwell-1903, incl. skull (SMC 1252); and Burwell-1907, incl. skull (SMC 1554), cervical vertebra (SMC 1568), coracoid, scapula (SMC 1569), furcula (SMC 1578), humerus (SMC 1555), ulna (SMC 1557), radius (SMC 1560), femur (SMC 1561), tibiotarsus (SMC 1564), tarsometatarsus (SMC 1565), and pedal phalanx (SMC 1572). Figured by Friant 1950b, pl. 1 (SMC 136, 146, 149, 162 and 1252), pl. 2 (SMC 1555, 1557, 1560, 1569 and 1578), and pl. 3 (SMC 1554, 1561, 1564, 1565, 1572, 1579 and 1586). List of specimens and catalogue numbers were derived from Friant 1950b. More bones and bone fragments than indicated are probably available per syntypical skeleton.]

Nycticorax fenensis Friant, 1950b: 333 [Same syntypes as for *Nycticorax fenensis* Friant, 1950a. This is both a junior objective synonym and a junior primary homonym of *Nycticorax fenensis* Friant, 1950a.]

Distribution: Holocene (Mesolithic) of Burwell and Reach, Cambridgeshire fens, England (Milne-Edwards 1869-1871: 108, Lydekker 1891c, Bell 1915, Friant 1950a,b sub *N. fenensis*, Northcote 1980, 1982d).

Remarks: In describing the species, Friant (1950a,b) was apparently misled by the length and shape of the bill of a juvenile Bittern found in the collections under the catalogue number Reach-1901. Northcote (1980, 1982d) overlooked Friant's identification of these bones and restudied them, finding that they belong to the modern *Botaurus stellaris* (Linnaeus). Based on her results I formally synonymize here *Nycticorax fenensis* Friant, 1950a and *Nycticorax fenensis* Friant, 1950b with the modern *Botaurus stellaris* (Linnaeus, 1758).

Genus incertae sedis

"*Anas*" *basaltica* Bayer

Anas basaltica Bayer, 1882: 20 [Holotype from Varnsdorf: cranial end of left coracoid, cranial end of left scapula, proximal end of left humerus and a rib fragment, all probably from a single individual, in slab and counter-slab; NMP ČM-119, cast in NHMW, uncatalogued. Figured by Bayer 1882, fig. 8/1, Bayer 1883, fig. 1, and Mlíkovský & Švec 1989, pl. 1, fig. a,c.]

Anas basaltica Bayer, 1883: 62 [Same holotype, repeated description.]

Distribution: Middle Oligocene (MP 23-24) of Varnsdorf, Czechia (Bayer 1882, 1883, Mlíkovský & Švec 1989).

Remarks: Described as a duck. This is an indeterminate heron of the subfamily Ardeinae (Mlíkovský & Švec 1989).

Family Phalacrocoracidae Reichenbach

Phalacrocoracidae Reichenbach, 1850 [Modern family.]

Genus *Nectornis* Cheneval

Nectornis Cheneval, 1984a: 56 [Type by original designation: *Graculus miocaenus* Milne-Edwards, 1867b.]

Distribution: Early Miocene (MN 2a-4b) of France, Germany and Czechia. Extralimital record is confined to *Nectornis anatolicus* (Mourer-Chauviré) from the middle Miocene (MN 6-8) of Bes-Konak in Turkey (Mourer-Chauviré 1978, Mlíkovský 1999c).

Nectornis miocaenus (Milne-Edwards)

Graculus miocaenus Milne-Edwards, 1867b: 255 [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1984a: 56): right tarsometatarsus; MNHN Av-9416. Figured by Milne-Edwards 1867-1868, pl. 39, fig. 5-9, Cheneval 1984a, pl. 3, fig. 7a,b. Paralectotypes from Saint-Gérard-le-Puy: sternum (MNHN Av-9323), right wing of furcula (MNHN Av-9442), right coracoid (MNHN Av-9326), scapula (MNHN Av-9375), left ulna (MNHN Av-9375), pelvis (MNHN Av-9452), left femur (MNHN Av-9406). Figured by Milne-Edwards 1867-1868, pl. 39, fig. 5, 13-16 (femur), 17-18 (pelvis), pl. 41, fig. 1-4 (sternum), 5-6 (furcula), 7-8 (coracoid), 9-10 (scapula), pl. 42, fig. 3-4 (ulna), and Cheneval 1984a, pl. 3, fig. 1 (furcula), 2a,b (coracoid), 4a,b (ulna), 5a,b (femur).

Phalacrocorax miocaenus (Milne-Edwards): Lydekker 1891a: 54 [New combination.]

Oligocorax miocaenus (Milne-Edwards): Lambrecht 1931b: 81 [New combination, but *Oligocorax* was nomen nudum at this point.]

Nectornis miocaenus (Milne-Edwards): Cheneval 1984a: 56 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards, 1867-1868, Cheneval 1984a), early Miocene (MN 2) of Ravelzhausen, Germany (Martini 1974 sub *Phalacrocorax littoralis*, Cheneval 1984a), and early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský orig.).

Genus *Phalacrocorax* Brisson

Phalacrocorax Brisson, 1760 [Modern genus.]

Botaurites Ammon, 1918: 31 [Type by monotypy: *Botaurites avitus* Ammon, 1918.]

Oligocorax Lambrecht, 1931b: 80 [Nomen nudum; type not fixed after 1930.]

Oligocorax Lambrecht, 1933: 290 [Nomen nudum; type not fixed after 1930.]

Paracorax Lambrecht, 1933: 292 [Type by monotypy: *Phalacrocorax destefanii* Regàlia, 1902.]

Pliocarbo Tugarinov, 1940a: 205 [Type by original designation: *Pliocarbo longipes* Tugarinov, 1940a.]

Oligocorax Brodkorb, 1952: 175 [Type by original designation: *Graculus littoralis* Milne-Edwards, 1863.]

Phalacrocorax littoralis (Milne-Edwards)

Graculus littoralis Milne-Edwards, 1863: 161 [Holotype from Saint-Gérard-le-Puy: proximal end of right tarsometatarsus, MNHN Av-9434. Figured by Milne-Edwards 1867-1868, pl. 42, fig. 5-8, Cheneval 1984a, pl. 2, fig. 7a-c. Cheneval (1984a: 53, 55) overlooked that Milne-Edwards (1863: 161) expressly based this species only on the tarsometatarsus, and erroneously considered all the five elements figured by Milne-Edwards (1867-1868) syntypes, selecting left humerus, MNHN Av-9421, as the lectotype of the species. His action is invalid. The "lectotype" was figured by Milne-Edwards 1867-1868, pl. 44, fig. 1-5, and Cheneval 1984a, pl. 2, fig. 3a-b.]

Phalacrocorax littoralis (Milne-Edwards): Lydekker 1891a: 56 [New combination.]

Oligocorax littoralis (Milne-Edwards): Lambrecht 1931b: 81 [New combination, but *Oligocorax* was nomen nudum at this point.]

Distribution: Early Miocene (MN 1) of Weisenau-AS, Germany (Lambrecht 1933: 290, locality uncertain); early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1867-1868, Cheneval 1984a); and early Miocene (MN 3) of Merkur, Czechia (Mlíkovský orig.). The record from the early Miocene (MN 2) of Ravelzhausen, Germany (Martini 1974), is invalid, because the material belongs to *Nectornis miocaenus* (Milne-Edwards) (Cheneval 1984a). The extralimital record from the late Miocene (MN 9) of Bled ed Dourah in Tunisia (Rich 1972) is dubious because of its age (Mlíkovský orig.).

Phalacrocorax intermedius (Milne-Edwards)

Graculus intermedius Milne-Edwards, 1867b: 266 [Holotype humerus from Orleanais: proximal end of left; formerly in coll. Nouel, present location unknown. Figured by Milne-Edwards 1867-1868, pl. 43, fig. 8-11. Brodkorb (1963: 251) stated that the holotypical humerus is from the right side of the body. However, bones are drawn side-reversed in Milne-Edwards 1867-1868, which makes it probable, that the specimen originated from the left side of body.]

Phalacrocorax intermedius (Milne-Edwards): Lydekker 1891a: 53 [New combination.]

Phalacrocorax praecarbo Ammon, 1918: 28 [Holotype from Dechbetten: humeral end of left coracoid, SMF, uncatalogued. Figured by Ammon 1918, fig. 3.]

Ardea brunhuberi Ammon, 1918: 30 [Holotype from Dechbetten: proximal end of left carpo-metacarpus, SMF, uncatalogued. Figured by Ammon 1918, fig. 4.]

Botaurites avitus Ammon, 1918: 31 [Holotype from Dechbetten: cervical vertebra, SMF, uncatalogued. Figured by Ammon 1918, fig. 5-6.]

Miocorax intermedius (Milne-Edwards): Lambrecht 1933: 292 [New combination.]

Phalacrocorax brunhuberi (Milne-Edwards): Brodkorb 1980: 91 [New combination.]

Distribution: Early Miocene (MN 3) of Břešřany, Czechia (Mlíkovský 1999c), early Miocene (MN 4) of Orleanais, France (Milne-Edwards 1867-1868, Mlíkovský 1992b); and middle Miocene (MN 7-8) of Dechbetten, Germany (Ammon 1918, Lambrecht 1933, Brodkorb 1980, Olson 1985a, Mlíkovský 1992b).

Remarks: *Phalacrocorax praecarbo* Ammon and *Botaurites avitus* Ammon were synonymized with *Phalacrocorax brunhuberi* (Ammon) by Brodkorb (1980) and Olson (1985a: 67). The latter species was subsequently synonymized with *Phalacrocorax intermedius* (Milne-Edwards) by Mlíkovský (1992b: 437).

***Phalacrocorax ibericus* Villalta**

Phalacrocorax ibericum Villalta, 1963: 267 [Holotype from Valles de Fuentidueña: distal end of right humerus; CV 3362. Figured by Villalta 1963, pl. 2, fig. 4, 4a.]

Phalacrocorax ibericus Villalta: Mlíkovský, this paper [Ending corrected.]

Distribution: Late Miocene (MN 9) of Valles de Fuentidueña, Spain (Villalta 1963).

Remarks: The figures show a highly abraded fragment, unsuitable for exact identification.

***Phalacrocorax lautus* Kuročkin & Ganea**

Phalacrocorax lautus Kuročkin & Ganea, 1972: 47 [Holotype from Golbočica: proximal end of right femur; GIKM 12173/3c. Figured by Kuročkin & Ganea 1972, text-fig. 1, pl. 1, fig. 1.]

Distribution: Late Miocene (MN ?9) of Golbočica, Moldova (Kuročkin & Ganea 1972). The partial carpometacarpus from the late Miocene (MN ?9) of Chișinău, Moldova, tentatively attributed to *Phalacrocorax lautus* Kuročkin & Ganea by Kessler (1984b), became later holotype of *Branta minor* Kessler & Gál (Kessler & Gál 1996).

Remarks: This species agrees in size and age with *Phalacrocorax ibericus* Villalta, with which it was not compared (but see above under *Phalacrocorax ibericus* Villalta).

***Phalacrocorax longipes* (Tugarinov)**

Pliocarbo longipes Tugarinov, 1940a: 205 [Lectotype from Novaja Slobodka (selected by Brodkorb 1963c: 254): right tarsometatarsus; PIN 228-2. Figured by Tugarinov 1940a, fig. 1a-b; and Dement'ev 1964, fig. 693b,v. Paralectotype from Novaja Slobodka: right femur; PIN 228-1. Figured by Tugarinov 1940a, fig. 2; and Dement'ev 1964, fig. 693a. Brodkorb (1963c: 254) stated that this species was based on a tarsometatarsus, not mentioning the syntypical femur. Nevertheless, this can be interpreted as a selection of the lectotype.]

Phalacrocorax longipes (Tugarinov): Mlíkovský, this paper [New combination.]

Distribution: Late Miocene (MN 11-13) of Novaja Slobodka, Ukraine (Vidgal'm 1886 sub *Haliaeetus fossilis* var. *odessana major*, Tugarinov 1940a, Burčak-Abramovič & Nikolov 1984), Škodova Hill, Ukraine (Vojinstens'kyj 1967), unspecified surroundings of Odesa, Ukraine (Dubrovo & Kapelist 1979); and early Pliocene (MN 15) of Odesa – catacombs, Ukraine (Vojinstves'kyj 1967).

Remarks: Burčak-Abramovič (in Burčak-Abramovič & Nikolov 1984: 24) concluded that *Pliocarbo longipes* is a typical cormorant, differing from *Phalacrocorax carbo* (Linnaeus) in being larger, but retained its generic position. Accordingly, I synonymize here *Pliocarbo* Tugarinov with the modern genus *Phalacrocorax* Brisson.

***Phalacrocorax serdicensis* Burčak-Abramovič & Nikolov**

Phalacrocorax serdicensis Burčak-Abramovič & Nikolov, 1984: 25 [Holotype from Chrabářsko: proximal end of right humerus; NMNHS 1-1/15. Figured by Burčak-

Abramovič & Nikolov 1984, fig. 1, 1a.]

Distribution: Late Miocene (MN 11-13) of Chrabärsko, Bulgaria (Burčák-Abramovič & Nikolov 1984).

Remarks: This species agrees in size, habitat and geographic distribution with the modern *Phalacrocorax aristotelis* (Linnaeus), with which it was not compared.

***Phalacrocorax aristotelis* (Linnaeus) – European Shag**

Pelecanus aristotelis Linnaeus, 1761 [Modern species.]

Phalacrocorax De Stefani Regàlia, 1902: 225 [Syntypes from Orciano Pisano: one cervical and two thoracic vertebrae, fragmentary furcula, right coracoid, proximal end of left humerus, distal end of left humerus, left ulna, proximal end of left femur, fragmentary shaft of tibiotarsus, right tarsometatarsus, pedal phalanx; present location unknown – Delle Cave 1996. Figured by Regàlia 1902, pl. 27, fig. 4 (cervical vertebra), 5-6 (furcula), 7 (coracoid), 8 (proximal end of humerus), 9 (distal end of humerus), 10 (ulna), 11 (femur), 12 (tibiotarsus), 13-14 (tarsometatarsus).]

Paracorax De Stefani (Regàlia): Lambrecht 1933: 292 [New combination.]

Phalacrocorax destefani Regàlia: Brodkorb 1963: 252 [Spelling emended.]

Phalacrocorax destefanii Regàlia: Mlíkovský this paper [Spelling emended.]

Distribution: Middle Pliocene (MN 15-16) of Orciano Pisano, Italy (Regàlia 1902).

Remarks: *Phalacrocorax destefanii* Regàlia was found in marine deposits and falls in the size class of the modern *Phalacrocorax aristotelis* (Linnaeus), which still inhabits the region. In absence of the evidence to the contrary, I synonymize here the former with the latter species.

***Phalacrocorax carbo* (Linnaeus) – Great Cormorant**

Pelecanus Carbo Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Overstrand, England (Harrison 1979d, 1985b), West Runton – unknown bed, England (Newton 1891, Harrison 1979d), and Přezletice, Czechia (Jánossy 1983 sub *P. cf. carbo*, Mlíkovský orig.).

Family Anhingidae Reichenbach

Anhingidae Reichenbach, 1849 [Modern family.]

Genus *Anhinga* Ridgway

Anhinga Ridgway, 1887 [Modern genus.]

***Anhinga pannonica* (Lambrecht)**

Plotus pannonicus Lambrecht, 1916a: 9 [Holotype from Tataruș-Brusturi: 6th cervical vertebra; ?GIB. Figured by Lambrecht 1916a, fig. 1, 3, 5. Rich (1972: 45) selected the 6th cervical vertebra from Tataruș-Brusturi as a lectotype of the species, erroneously believing that the species was based on the vertebra and on a carpometacarpus. Nevertheless, Lambrecht (1916a: 9) clearly based the species only on the vertebra, the carpometacarpus being a referred specimen (see also Brodkorb & Mourer-Chauviré 1982: 510). Rich's (1972) action is invalid.]

Anhinga pannonica (Lambrecht): Brodkorb 1963: 256 [New combination.]

Distribution: Late Miocene (MN 10) of Tataruș-Brusturi, Romania (Lambrecht 1916a), and Götzendorf, Austria (Mlíkovský 1992a). Extralimital record includes late Miocene (MN 9) of Bled ed Dourah 20, Tunisia (Rich 1972) and Miocene (MN 4-13) of Siwalik Mts., Pakistan (Harrison & Walker 1982).

***Anhinga* sp.**

Distribution: Middle Miocene (MN 6-8) of Mátraszőlös 1, Hungary (Gál & Kessler in Gál et al. 1999 sub aff. *Anhinga* sp.).

Order Bucerotiformes Fürbringer

Bucerotes Fürbringer, 1888 [Modern order.]

Family Laurillardiidae Harrison

Laurillardiidae Harrison, 1979b: 108 [Type genus: *Laurillardia* Milne-Edwards, 1871.]

Messelirrisoridae Mayr, 1998c: 13 [Type genus: *Messelirrisor* Mayr, 1998c.]

Distribution: Early Eocene (MP 8) of England, middle Eocene (MP 11 or 13) of Germany, and late Eocene (MP 19) of France. In addition to the record listed below, Mayr (1998c: 27) listed the family also from the early Eocene (MP 8) of Walton-on-the-Naze, England, and from the middle Eocene (MP 11 or 13) of Geiseltal, Germany.

Remarks: Messelirrisoridae Mayr is synonymized here with the Laurillardiidae Harrison, following the discussion in Mayr (1998c).

Genus *Messelirrisor* Mayr

Messelirrisor Mayr, 1998c: 24 [Type by original designation: *Messelirrisor parvus* Mayr, 1998c.]

Distribution: Middle Eocene (MP 11) of Germany.

Remarks: Differences between *Messelirrisor* Mayr and *Laurillardia* Milne-Edwards were not demonstrated by Mayr (1998c), and it is well possible that the former will fall in the synonymy of the latter genus.

Messelirrisor parvus Mayr

Messelirrisor parvus Mayr, 1998c: 25 [Holotype from Messel: skeleton in slab; SMF ME-2793. Figured by Mayr 1998c, pl. 2.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998c).

Messelirrisor halcyrostris Mayr

Messelirrisor halcyrostris Mayr, 1998c: 26 [Holotype from Messel: skeleton in slab; SMF ME-1883a,b. Figured by Mayr 1998c, pl. 1, pl. 3 left, and cover; and Mayr 2000g, fig. 1.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998c).

Messelirrisor grandis Mayr

Messelirrisor grandis Mayr, 2000f: 965 [Holotype from Messel: partial skeleton in slab, incl. sternum, pectoral girdle and both wings; SMNK PAL-3803. Figured by Mayr 2000f, fig. 5.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 2000f).

Genus *Laurillardia* Milne-Edwards

Laurillardia Milne-Edwards, 1871: 374 [Type by monotypy: *Laurillardia longirostris* Milne-Edwards, 1871.]

Laurillardia longirostris Milne-Edwards

Laurillardia longirostris Milne-Edwards, 1871: 374 [Holotype from Montmartre: partial skeleton in slab; MNHN 7964. Figured by Cuvier 1822, pl. 75, fig. 6 (sub 9^e espèce); Cuvier 1836, pl. 156, fig. 6 (sub 9^e espèce); Milne-Edwards 1869-1871, pl. 161, fig. 1; Zittel 1890, fig. 719; and Mayr 1998c, pl. 3 right.]

Laurillardia parisiensis Flot, 1892: 1 [Holotype from Montmartre: incomplete skeleton in a slab and counterslab; present location unknown. Figured by Flot 1892, pl. 18, fig. 1-2, and text-fig. 1.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Cuvier 1822, 1836 sub 9^e espèce, Milne-Edwards 1869-1871, Flot 1892 sub *Laurillardia parisiensis*, Brunet 1970, Harrison 1979b, Mayr 1998c).

Remarks: Milne-Edwards (1869-1871) placed this species in his "Passereaux", which included most arboreal birds, incl. the Wetmorean orders Cuculiformes, Coraciiformes, Piciformes and Passeriformes. Paris (1912: 284) and Brunet (1970: 47) included it in the Sturnidae (= Eulabeidae in Paris 1912) without explanation, Harrison (1979b: 107) believed that it is an ancient songbird (order Passeriformes), and Brodkorb (1978: 214) relegated this species in the category of Aves incertae sedis. Mayr (1998c: 13) identified the species as a relative of hoopoes.

Laurillardia parisiensis Flot was synonymized with *Laurillardia longirostris* Milne-Edwards by Brunet (1970), with whom I concur in this point.

***Laurillardia munieri* Flot**

Laurillardia Munieri Flot, 1892: 7 [Holotype from Montmartre: incomplete skeleton in a slab; present location unknown. Figured by Flot 1892, pl. 18, fig. 3, and text-fig. 3.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Flot 1892, Brunet 1970).

Remarks: Brunet (1970) synonymized this species with *Laurillardia longirostris* Milne-Edwards, which however is larger in dimensions (Mlíkovský, orig.).

Family Upupidae Leach

Upupidae Leach, 1819 [Modern family.]

Genus *Upupa* Linnaeus

Upupa Linnaeus, 1758 [Modern genus.]

***Upupa epops* Linnaeus – Eurasian Hoopoe**

Upupa epops Linnaeus, 1758 [Modern species]

Upupa phoeniculides Jánossy, 1974b: 231 [Holotype from Hundsheim: right coracoid; UWPI 1889/65. Not figured.]

Distribution: Middle Pleistocene (MQ 2A) of Hundsheim, Austria (Jánossy 1974b sub *Upupa phoeniculides*).

Remarks: Jánossy (1974b) separated this species from the modern *Upupa epops* Linnaeus on the basis of the shape of the ligamentum acrocoraco-procoracoideum, which is ossified in *Upupa* species, observing that the distal-medial edge of this ligament is smooth in *Upupa epops*, but edged in the holotype of *Upupa phoeniculides* and in the modern *Phoeniculus* Jarocki, 1821. I restudied the holotype and confirmed the condition for it and for *Phoeniculus* coracoids. However, I found that the edge of the ligament is smooth only in the modern *Upupa epops* from Europe, while it exhibits the *Phoeniculus* condition in all specimens of modern Hoopoes from Africa and southern Asia (ca. 20 specimens examined in USNM and two specimens in my comparative collection). Hence, I synonymize here *Upupa phoeniculides* Jánossy with the modern *Upupa epops* Linnaeus.

***Upupa* sp.**

Distribution: Early Miocene (MN 3) of Merkur, Czech Republic (Mlíkovský, orig.).

Family Phoeniculidae Bonaparte

Phoeniculidae Bonaparte, 1831 [Modern family.]

Genus *Phirriculus* Mlíkovský & Göhlich

Phirriculus Mlíkovský & Göhlich, 2000: 419 [Type by original designation: *Phirriculus pinicola* Mlíkovský & Göhlich, 2000.]

***Phirriculus pinicola* Mlíkovský & Göhlich**

Phirriculus pinicola Mlíkovský & Göhlich, 2000: 420 [Holotype from Wintershof-West: distal end of left humerus; BSP 1037-II-18188. Figured by Mlíkovský & Göhlich 2000, fig. 1a,b, 6a-c.]

Distribution: Early Miocene (MN 2a) of Saint-Gérand-le-Puy, France (Milne-Edwards 1871 sub *Limnatornis paludicola*, Ballmann 1969b sub *Limnatornis paludicola*, Mlíkovský & Göhlich 2000); early Miocene (MN 3a) of Wintershof-West, Germany (Ballmann 1969b sub Phoeniculidae indet., Mlíkovský & Göhlich 2000); and early Miocene (MN 4) of Petersbuch 38, Germany (Mlíkovský & Göhlich 2000).

Order Procellariiformes Fürbringer

Procellariiformes Fürbringer, 1888 [Modern order.]

Family Procellariidae Leach

Procellariidae Leach, 1820 [Modern family.]

Diomedeoididae Fischer, 1985: 113 [Type genus: *Diomedeoides* Fischer, 1985.]

Remarks: I regard all modern procellariiforms as belonging to a single family. In this broad sense, the Procellariidae encompass also the fossil Diomedeoididae. All fossil forms listed below are in need of revision.

Genus *Neptuniavis* Harrison & Walker

Neptuniavis Harrison & Walker, 1977a: 9 [Type by original designation: *Neptuniavis miranda* Harrison & Walker, 1977.]

Remarks: The taxonomic position of this genus is uncertain (see Cheneval 1995).

Neptuniavis miranda Harrison & Walker

Neptuniavis miranda Harrison & Walker, 1977a: 9 [Holotype from Sheppey: proximal end of right tarsometatarsus; BMNH A-3682. Figured by Harrison & Walker 1977a, fig. 2a-e.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Neptuniavis minor Harrison & Walker

Neptuniavis minor Harrison & Walker, 1977a: 11 [Holotype from Sheppey: proximal end of right tarsometatarsus; BMNH A-134. Figured by Harrison & Walker 1977a, pl. 1, fig. F-J.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Genus *Primodroma* Harrison & Walker

Primodroma Harrison & Walker, 1977a: 8 [Type by original designation: *Primodroma bournei* Harrison & Walker, 1977a.]

Primodroma bournei Harrison & Walker

Primodroma bournei Harrison & Walker, 1977a: 8 [Holotype from Sheppey: distal end of right humerus; BMNH A-3678. Figured by Harrison & Walker 1977a, pl. 1, fig. a-e.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Genus *Diomedeoides* Fischer

Diomedeoides Fischer, 1985: 114 [Type by original designation: *Diomedeoides minimus* Fischer, 1985.]

Diomedeoides minimus Fischer

Diomedeoides minimus Fischer, 1985: 114 [Holotype from Espenhain: right femur; NKMB Av-764. Figured by Fischer 1985, fig. 1-6, 12-14.]

Distribution: Middle Oligocene (MP 23-24) of Espenhain, Germany (Fischer 1985).

Genus *Frigidafons* Cheneval

Frigidafons Cheneval, 1995: 189 [Type by monotypy: *Frigidafons brodkorbi* Cheneval, 1995.]

Distribution: Extralimital record is limited to *Frigidafons babaheydariensis* Peters & Hamedani, 2000 from the Oligocene of Iran (Peters & Hamedani 2000).

***Frigidafons brodkorbi* Cheneval**

Frigidafons brodkorbi Cheneval, 1995: 189 [Holotype from Froidefontaine: almost complete skeleton in slab and counter-slab; coll. P. Paupe (slab) and coll. S. Michel (counter-slab). Figured by Cheneval & Pharisat 1995, text-fig. 1, pl. 1, fig. 1; and Cheneval 1995, fig. 1-2.]

Distribution: Middle Oligocene (MP 23-24) of Froidefontaine, France (Cheneval 1995); and early Miocene (MN 1) of Weisenau-AS, Germany (Cheneval 1995).

Genus *Puffinus* Brisson

Puffinus Brisson, 1760 [Modern genus.]

***Puffinus raemdonckii* (Beneden)**

Larus Raemdonckii Beneden, 1871: 258 [Holotype from Rupelmonde: left humerus lacking proximal end; IRScNB, uncatalogued. Figured by Beneden 1871, fig. 1. Brodkorb (1962a: 707) incorrectly believed that the species was based on two syntypical humeri, and selected the holotypical one as a lectotype of the species. His action is invalid.]

Puffinus raemdonckii (Beneden): Brodkorb 1962: 707 [New combination.]

Distribution: Middle Oligocene (MP 23-24) of Rupelmonde, Belgium (Beneden 1871, 1872, Miller & Sibley 1941, Brodkorb 1962a).

***Puffinus aquitanicus* (Milne-Edwards)**

Procellaria aquitanica Milne-Edwards, 1874: 6 [Holotype from Léognan: distal end of a humerus; MBo, uncatalogued. Figured by Milne-Edwards 1874, pl. 12, fig. 1, 1a-c.]

Puffinus aquitanicus (Milne-Edwards): Brodkorb 1963: 243 [New combination.]

Distribution: Early Miocene (MN 2-3) of Léognan, France (Milne-Edwards 1874).

***Puffinus antiquus* (Milne-Edwards)**

Procellaria antiqua Milne-Edwards, 1874: 7 [Holotype from Léognan: proximal end of a humerus; MHNB, uncatalogued. Not figured.]

Puffinus antiquus (Milne-Edwards): Brodkorb 1963c: 243 [New combination.]

Distribution: Early Miocene (MN 2-3) of Léognan, France (Milne-Edwards 1874).

***Puffinus nestori* Alcover**

Puffinus nestori Alcover, 1989.

Distribution: Late Pliocene (MN 18) of Ca Na Reia, Eivissa, Spain (Alcover 1989).

Remarks: Not seen.

***Puffinus holeae* Walker et al.**

Puffinus holei Walker, Wragg & Harrison, 1990. [Holotype from Barca.]

Puffinus holeae Walker et al.: Mourer-Chauviré & Antunes 2000: 132 [Spelling emended.]

Distribution: Late Pleistocene (MQ 2C) of Figueira Brava – layer 2, 3 and 4, Portugal (Mourer-Chauviré & Antunes 2000), and early Holocene (MQ 2E) of Figueira Brava – "upper layers" (Mourer-Chauviré & Antunes 2000). Extralimital record is available from the late Pleistocene (MQ 2C-D) of Fuerteventura (incl. type locality), Canary Islands (Walker et al. 1990), and Lanzarote, Canary Islands (McMinn et al. 1990).

Genus *Plotornis* Milne-Edwards

Plotornis Milne-Edwards, 1874: 5 [Type by monotypy: *Plotornis delfortrii* Milne-Edwards, 1874.]

***Plotornis arvernensis* (Milne-Edwards)**

Puffinus arvernensis Milne-Edwards, 1871: 572 [Nomen nudum; no description or indication.]

Puffinus arvernensis Milne-Edwards in Shufeldt, 1896: 510 [Holotype from Saint-Gérard-le-Puy: left tarsometatarsus; MNHN Av-2929. Figured by Shufeldt 1896, pl. 24, fig. 1-2; and Cheneval 1984b, pl. 2, fig. 1a-b.]

Plotornis arvernensis ("Milne-Edwards" = Milne-Edwards in Shufeldt): Cheneval 1984a: 48. [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1874, Cheneval 1984a). The record from the Holocene of Pietro Tampoia cave, Tavolara island near Sardinia, Italy (Shufeldt 1896, Regàlia 1897) is invalid, as the bones belong to the modern *Puffinus yelkouan* (Acerbi, 1827) (Mayaud & Schaub 1950, see also Lambrecht 1933: 269).

Remarks: Wetmore (1930a), Lambrecht (1933: 268) and Brodkorb (1963c: 243) correctly observed, that the binomen *Puffinus arvernensis*, as published by Milne-Edwards (1871: 572), is a nomen nudum, Cheneval's (1984a: 51) arguments to the contrary (under the reference to ICZN 1961, Art. 12 and 16) being incorrect. Wetmore (1930a) attributed the name to Shufeldt (1896), while Brodkorb (1963c) to Milne-Edwards in Shufeldt (1896). The latter interpretation is in accordance with the Code of zoological nomenclature (ICZN 1999, Art. 50.1.).

***Plotornis delfortrii* Milne-Edwards**

Plotornis delfortrii Milne-Edwards, 1874: 5 [Syntypes from Léognan: left tarsometatarsus and distal part of right humerus; MHNB, uncatalogued. Figured by Milne-Edwards 1874, pl. 1, fig. 1, 1a-e (tarsometatarsus), 2, 2a-d (humerus).]

Distribution: Early Miocene (MN 2-3) of Léognan, France (Milne-Edwards 1874), and Saucats, France (Lucazeau 1959).

Genus *Diomedea* Linnaeus

Diomedea Linnaeus, 1758 [Modern genus.]

***Diomedea rumana* Grigorescu & Kessler**

Diomedea rumana Grigorescu & Kessler, 1988: 92 [Lectotype from Credința (here selected): proximal part of right ulna; LPUB 279. Figured by Grigorescu & Kessler 1988, fig. 1a, b, a', b'. Paralectotype from Credința: distal part of right ulna; LPUB 280. Figured by Grigorescu & Kessler 1988, fig. 1b, b', c, c', d, d'. Grigorescu & Kessler (1988) considered both these bone fragments the holotype of the species, but presented no evidence that they originated from the same individual.]

Distribution: Middle Miocene (MN 8) of Credința, Romania (Grigorescu & Kessler 1988).

***Diomedea anglica* Lydekker**

Diomedea anglica Lydekker, 1891a: 189 [Holotype from Foxhall: right tarsometatarsus with associated phalanx I of digit 4; IM, uncatalogued (cast in BMNH A-87). Figured by Lydekker 1891a, fig. 42a-b.]

Distribution: Late Pliocene (MN 16-17) of Foxhall, England (Lydekker 1886, 1891a,d, Harrison & Walker 1978b), and Orford, England (Lydekker 1886, 1891a,d, Harrison & Walker 1978b). A tentative extralimital record is from the early Pliocene of Bone Valley in Florida (Wetmore 1943, Harrison & Walker 1978b). Fisher (1966) attributed to this species an ulna supposedly originating from the late Pleistocene (MQ 2C) of Ilford, England, and formerly described as belonging to the modern *Diomedea exulans* Linnaeus, 1758 by Newton (1891). Harrison & Walker (1977c) showed, that the specimen is at most sub-Recent in age, concluding that "it is plausible that ... [it] did not reach England unaided."

Remarks: Wilkinson (1969: 49) suggested, that *D. anglica* Lydekker is synonymous with the modern *D. albatrus* Pallas, 1769 (see also Miller 1962).

Genus *Calonectris* Mathews & Iredale

Calonectris Mathews & Iredale, 1915 [Modern genus.]

Calonectris diomedea (Scopoli) – Cory's Shearwater

Puffinus diomedea Scopoli, 1769 [Modern species.]

Puffinus eyermani Shufeldt, 1896: 511 [Lectotype from Pietro Tampoia Cave (here selected): right tibiotarsus (USNM; # 11 in Shufeldt 1896). Figured by Shufeldt 1896, pl. 24, fig. 8. Paralectotype from Pietro Tampoia Cave: right tarsometatarsus (USNM; # 10 in Shufeldt 1896).]

Distribution: Holocene (MQ 2E) of Pietro Tampoia Cave, Tavolara island near Sardinia (Shufeldt 1896 sub *P. eyermani*, Regália 1897 sub *P. eyermani*, Lambrecht 1933: 268 sub *P. eyermani*, Mayaud & Schaub 1950).

Remarks: *Puffinus eyermani* Shufeldt was synonymized with the modern *Calonectris diomedea* (Scopoli, 1769) by Mayaud & Schaub (1950).

Family Pelagornithidae Fürbringer

Pelagornithidae Fürbringer, 1888: 1565 [Type genus: *Pelagornis* Lartet, 1857.]

Odontopterygidae Lydekker, 1891a: 57 [Type genus: *Odontopteryx* Owen, 1873.]

Cyphornithidae Wetmore, 1928: 4 [Type genus: *Cyphornis* Cope, 1894.]

Pseudodontornithidae Lambrecht, 1933: 305 [Type genus: *Pseudodontornis* Lambrecht, 1930.]

Dasornithidae Harrison & Walker, 1976b: 24 [Type genus: *Dasornis* Owen, 1870.]

Distribution: Early Eocene (MP 8-9) of England, middle Eocene (MP 11-13) of Belgium (record tentative), late Eocene (MP 19) of France, and early Miocene (MN 2-3) of France. Extralimital record is very rich, ranging from the Paleocene to the Pliocene, and including all continents (Harrison & Walker 1976b, Olson 1985a: 194-201). Extralimital record from the Asian part of the Paratethys is limited to the Paleocene through Oligocene (Aslanova & Burčák-Abramovič 1982, 1999, Aver'janov et al. 1991). Further records from the Atlantic Ocean were reported from the ?Eocene of Nigeria (Andrews 1916, Harrison & Walker 1976b), and South Shetland Islands near Antarctica (Tonni 1980, Tonni & Tambussi 1985), Oligocene of South Carolina (see Hopson 1964, Harrison & Walker 1976b, Olson 1985), and the Miocene of Maryland (Olson 1985a), Virginia (Olson 1984, 1985a), North Carolina (Becker 1987), and Canary Islands (García Talavera 1990).

Remarks: The classification of this family basically follows Harrison & Walker (1976b) in absence of more recent data, although it apparently does not reflect the real phylogeny of the family (Olson 1985a, pers. communication 1997). All the taxa listed below are in need of revision. Subsequent research will certainly result in a marked decrease of pelagornithid taxa recognized from the European Eocene.

Genus *Odontopteryx* Owen

Odontopteryx Owen, 1873: 511 [Type by monotypy: *Odontopteryx toliapica* Owen, 1873.]

Remarks: Extralimital records include *Odontopteryx tschulensis* Aver'janov et al., 1991 from the late Paleocene of Kazakhstan (Aver'janov et al. 1991), *Odontopteryx* sp. from the middle Eocene of Uzbekistan (Aver'janov et al. 1991), and tentatively also from the early Eocene of Virginia (Olson 1999b sub cf. *Odontopteryx* sp.).

Odontopteryx toliapica Owen

Odontopteryx toliapica Owen, 1873: 511 [Holotype from Sheppey: imperfect skull with parts of lower jaws; BMNH 44096. Figured by Owen 1873, pl. 16, fig. 1-8, pl. 17, fig. 1-2 (longitudinal section of a tooth); Krejčí 1877, fig. 869; Woodward 1890. fig. 115; Zittel

1890, fig. 717; Lambrecht 1933, fig. 108; Dement'ev 1964, fig. 691; Harrison & Walker 1976b, text-fig. 1-5, pl. 1, fig. a-f, pl. 2, fig. a-m; and Harrison & Walker 1977a, text-fig. 4-5, pl. 1, fig. k-p.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Owen 1873, Martin 1874, Desor 1876, Harrison & Walker 1976b, 1977a).

Genus *Macrodonopteryx* Harrison & Walker

Macrodonopteryx Harrison & Walker, 1976b: 12 [Type by original designation: *Macrodonopteryx oweni* Harrison & Walker, 1976b.]

***Macrodonopteryx oweni* Harrison & Walker**

Macrodonopteryx oweni Harrison & Walker, 1976b: 12 [Holotype from Sheppey: imperfect skull; BMNH A-1. Figured by Harrison & Walker 1976b, text-fig. 6-10, pl. 3, fig. g,h, pl. 4, fig. a-d, Harrison & Walker 1977a, text-fig. 6-7.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1976b, 1977a).

Genus *Pseudodontornis* Lambrecht

Pseudodontornis Lambrecht, 1930: 1 [Type by monotypy: *Odontopteryx longirostris* Spulski, 1910.]

***Pseudodontornis longirostris* (Spulski)**

Odontopteryx longirostris Spulski, 1910: 507 [Holotype from an unknown locality: partial skull; present location unknown. Figured by Spulski 1910, fig. 1-7.]

Pseudodontornis longirostris (Spulski): Lambrecht 1930: 1 [New combination.]

Distribution: Unknown (Spulski 1910, Lambrecht 1930, Harrison & Walker 1976b). Spulski (1910: 508) reported, that the specimen was purchased for the Institute of Zoology of the Königsberg University in Prussia (now Kaliningrad in Russia) by Professor Braun in 1905 from a local merchant, who said he bought it some five years before that date from a seaman, who reportedly brought it from Brazil. Brodkorb (1963c: 263) conjectured, that the holotype could originate from the Miocene of Germany, but there are no suitable localities of this age in this part of Europe (see Mlíkovský 1996a). If the specimen is from Europe, then it would most probably come from the marine Eocene deposits of the North Sea Basin, which are particularly well represented in northern Germany, Denmark and England (Mlíkovský 1996a).

A mandible, tentatively referred to *Pseudodontornis longirostris* (Spulski), is known from the late Oligocene of Charleston, South Carolina (Hopson 1964, Harrison & Walker 1976b: 15, Olson 1985a: 196-197).

***Pseudodontornis longidentata* Harrison & Walker**

Pseudodontornis longidentata Harrison & Walker, 1976b: 17 [Holotype from Sheppey: portion of rostrum and associated imperfect atlas vertebra; IGS GSM-107876. Figured by Harrison & Walker 1976b, text-fig. 11-13, pl. 5, fig. A-G, 1977, text-fig. 8, pl. 1, fig. Q-S.]

Distribution: Early Eocene (MP 8-9) from Sheppey, England (Harrison & Walker 1976b, 1977a).

Genus *Dasornis* Owen

Dasornis Owen, 1869: 59 [Nomen nudum; no description or indication.]

Dasornis Owen, 1870: 145 [Type by monotypy: *Dasornis londinensis* Owen, 1870.]

Remarks: This genus was included in the Gastornithidae by Lambrecht (1933), and transferred to the Odontopterygiformes by Harrison & Walker (1976b).

***Dasornis londinensis* Owen**

Dasornis londinensis Owen, 1869: 59 [Nomen nudum; no description or indication.]

Dasornis londinensis Owen, 1870: 145 [Holotype from Sheppey: imperfect skull; BMNH 31929. Figured by Owen 1870, pl. 16, Harrison & Walker 1976b, pl. 8, fig. A-C, 1977, pl. 3, fig. A-D.]

Dasornis londiniensis Lydekker, 1891a: 359 [Unsubstantiated emendation. However, it is not a young objective synonym of *Dasornis londinensis*, because this variant spelling is deemed to be identical with the original spelling – ICZN 1999, Art. 58.15.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Owen 1869, 1870, Harrison & Walker 1976b, 1977a).

Genus *Argillornis* Owen

Megalornis Seeley, 1866: 110 [Type by monotypy: *Lithornis emuinus* Bowerbank, 1854. This is a junior homonym of *Megalornis* Gray, 1841, and *Megalornis* Owen, 1843.]

Argillornis Owen, 1878: 124 [Type by monotypy: *Argillornis longipennis* Owen, 1878.]

***Argillornis emuinus* (Bowerbank)**

Lithornis emuinus Bowerbank, 1854: 263 [Holotype from Sheppey: partial shaft of right humerus, originally described as part of a tibiotarsus; BMNH 38941. Figured by Bowerbank 1854, unnumbered fig.; and Harrison & Walker 1977, pl. 4, fig. a.]

Megalornis emuianus (Bowerbank): Seeley, 1866: 110 [New combination, specific name misspelled.]

Argillornis emuinus (Bowerbank): Brodkorb 1963: 248 [New combination.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Bowerbank 1854, Quekett 1855, Seeley 1866, 1869, 1874, Owen 1878, Lydekker 1891a, Lambrecht 1933: 282-284, Harrison & Walker 1976b, 1977a); and middle Eocene (MP 11-13) of Etterbeek, Belgium (Dollo 1909: 111).

Remarks: Lambrecht (1921) attributed *Megalornis emuinus* [sic!] on p. 17 to Seeley (1866) and on p. 36 to Seeley (1874), which is unsubstantiated in both cases.

***Argillornis longipennis* Owen**

Argillornis longipennis Owen, 1878: 124 [Syntypes from Sheppey: proximal end of right humerus (BMNH A-5) and proximal end of left humerus (BMNH A-8). Figured by Owen 1878, pl. 6, fig. 1-3, 7-12, 16, Harrison & Walker 1976b, pl. 6, fig. d-f; and Harrison & Walker 1977a, pl. 4, fig. b-g.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Owen 1878, 1880, Lydekker 1891a, Lambrecht 1933, Harrison & Walker 1976b, 1977), and perhaps also middle Eocene (MP 11-13) of Etterbeek, Belgium (Dollo 1909: 111, Lambrecht 1933: 284).

Genus *Pelagornis* Lartet

Pelagornis Lartet, 1857: 740 [Type by monotypy: *Pelagornis miocaenus* Lartet, 1857. This is a senior homonym of *Pelagornis* Seeley, 1866.]

Remarks: Lambrecht (1933: 282, footnote) mentioned that a humerus labeled "*Pelagornis Delfortrii* 1869" was found in the Museum of Bordeaux. Brodkorb (1963c: 263) attributed this name to Lambrecht (1933) as a nomen nudum, listing it in the synonymy of *Pelagornis miocaenus* Lartet. This is incorrect, because *Pelagornis Delfortrii* is a label name without any standing in zoological nomenclature. It cannot be attributed to Lambrecht.

***Pelagornis miocaenus* Lartet**

Pelagornis miocaenus Lartet, 1857: 740 [Holotype from Armagnac: left humerus; MNHN, uncatalogued. Figured by Milne-Edwards, 1867-1868, pl. 45, fig. 1-7, Harrison & Walker, 1976b, pl. 9, fig. d-e.]

Distribution: Early Miocene (MN 2-3) of Armagnac, France (Lartet 1857, Harrison & Walker 1976b). Harrison & Walker (1976b: 29) tentatively attributed to this species partial humerus from the middle or late Miocene of coastal cliffs near Waipara River mouth, North Canterbury, New Zealand, which was first described by Scarlett (1972).

Order Ciconiiformes Bonaparte

Ciconiae Bonaparte, 1854 [Modern order.]

Family Podicipedidae Bonaparte

Podicipedidae Bonaparte, 1831 [Modern family.]

Genus *Podiceps* Latham

Podiceps Latham, 1787 [Modern genus.]

Thiornis Navás, 1922: 59 [Type by monotypy: *Thiornis sociata* Navás, 1922.]

Miobaptus Švec, 1982: 244 [Type by original designation: *Miobaptus walteri* Švec, 1982.]

Podiceps walteri (Švec)

Miobaptus walteri Švec, 1982: 246 [Holotype from Dolnice: proximal part of right humerus; DP FNSP 4810. Figured by Švec 1982, pl. 1, fig. 2.]

Podiceps walteri (Švec): Mlíkovský 2000f: 97 [New combination.]

Distribution: Early Miocene (MN 3) of Skyřice, Czechia (Mlíkovský 2000f); and early Miocene (MN 4b) of Dolnice, Czechia (Švec 1982, 1984, Mlíkovský 2000f).

Remarks: The genus *Miobaptus* Švec was synonymized with the modern genus *Podiceps* Linnaeus by Mlíkovský (2000f: 97.).

Podiceps miocenicus Kessler

Podiceps miocenicus Kessler, 1984b: 523 [Holotype from Chișinău: right humerus in two parts; LPUI 61-MS. Figured by Kessler 1984b, fig. 13-14.]

Distribution: Late Miocene (MN 9) of Chișinău, Moldova (Kessler 1984b).

Podiceps sociatus (Navás)

Thiornis sociata Navás, 1922: 59 [Holotype from Libros: almost complete skeleton in a slab with feather impressions; MNHN 1930-1. Figured by Navás 1922, pl. 2, and Olson 1995a, fig. 1-4.]

Podiceps sociatus (Navás): Olson 1995: 132 [New combination.]

Distribution: Late Miocene (MN 9-10) of Libros, Spain (Navás 1922, Cracraft 1973: 40, Olson 1995a, Storer 2000).

Remarks: Lambrecht (1933, fig. 144) figured another specimen in slab from Libros (BMNH A-1620), which was later erroneously considered as the holotype of *Thiornis sociata* by Brodkorb (1967: 124) and Cracraft (1973: 40, fig. 40). The latter specimen is indeterminate, but does not belong to the Podicipedidae (Olson 1977: 373, 1995).

Thiornis sociata Navás was long believed to represent a fossil rail, family Rallidae (Lambrecht 1933: 475, Brodkorb 1967: 124, Cracraft 1973: 40), until Olson (1995a) rediscovered its type material and identified it as a grebe, synonymizing *Thiornis* Navás with the modern genus *Podiceps* Linnaeus.

Podiceps cristatus (Linnaeus) – Great Crested Grebe

Colymbus cristatus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub. *P. cf. cristatus*).

Podiceps grisegena (Boddaert) – Red-necked Grebe

Colymbus grisegena Boddaert, 1783 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *P. aff. grisegena*), Stránská skála – Čapek's site 5a, lower layer (Jánossy 1972), and Stránská skála – Musil's talus cone, layer 13 (Mlíkovský 1995b).

***Podiceps auritus* (Linnaeus) – Horned Grebe**

Colymbus auritus Linnaeus, 1758 [Modern species.]

Fulica sp. (*pisana*) Portis, 1889: 13 [Holotype from Orciano Pisano: distal end of right humerus; IGF 14874. Figured by Portis 1889, pl. 1, fig. 24-25, and Regàlia 1902, pl. 27, fig. 21-22.]

Podicipes pisanus (Portis): Regàlia 1902: 233 [New combination, generic name misspelled.]

Distribution: Middle Pliocene (MN 15-16) of Orciano Pisano, Italy (Portis 1889, Regàlia 1902, 1907).

Remarks: *Fulica pisana* Portis was originally described as a coot, but Regàlia (1902) showed that it represents a grebe. After a restudy of a cast of the holotype and a biometrical study (Mlíkovský orig.) I concluded, that *Podiceps pisanus* (Portis) is synonymous with the modern Horned Grebe *Podiceps auritus* (Linnaeus).

***Podiceps nigricollis* Brehm – Black-necked Grebe**

Podiceps nigricollis Brehm, 1831 [Modern species.]

Distribution: Late Miocene (MN 9-10) of Sokolov, Ukraine (Vojinstvens'kyj 1967 sub *Colymbus* cf. *nigricollis*); early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *P. cf. nigricollis*), and Nagyhársánhegy 1-4, Hungary (Jánossy 1980a sub *P. aff. nigricollis*, Jánossy 1980b sub *P. cf. nigricollis*).

***Podiceps ruficollis* Pallas – Little Grebe**

Colymbus ruficollis Pallas, 1764 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Huéscar-1, Spain (Sánchez Marco 1989).

Family Pelecanidae Rafinesque

Pelecanidae Rafinesque, 1815 [Modern family.]

Genus *Miopelecanus* Cheneval

Miopelecanus Cheneval, 1984a: 68 [Type by original designation: *Pelecanus gracilis* Milne-Edwards, 1863.]

***Miopelecanus gracilis* (Milne-Edwards)**

Pelecanus gracilis Milne-Edwards, 1863: 161 [Holotype from Saint-Gérard-le-Puy: proximal end of a tarsometatarsus; present location unknown, probably lost – Cheneval 1984a: 68, 74.

Figured by Milne-Edwards 1867-1868, pl. 38, fig. 1-5.]

Miopelecanus gracilis (Milne-Edwards): Cheneval 1984a: 68 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1867-1868, Cheneval 1984a).

***Miopelecanus intermedius* (Fraas)**

Pelecanus intermedius Fraas, 1870: 281 [Holotype from Hahnenberg: incomplete skull with posterior end of left mandible; SMNS, uncatalogued. Figured by Fraas 1870, pl. 13, fig. 3-4.]

Pelecanus fraasi Lydekker, 1891a: 44 [Holotype from Lierheim: slightly imperfect skull; BMNH 47862. Figured by Lydekker 1891a, fig. 10a; and Heizmann & Hesse 1995, fig. 2a.]

Miopelecanus intermedius (Fraas): Heizmann & Hesse 1995: 175 [New combination.]

Distribution: Middle Miocene (MN 6) of Hahnenberg, Germany (Fraas 1870, Lydekker 1891a), and Lierheim, Germany (Lydekker 1891a, Heizmann & Hesse 1995).

Remarks: *Pelecanus fraasi* Lydekker was synonymized with *Pelecanus intermedius* Fraas by Heizmann & Hesse (1995).

Genus *Pelecanus* Linnaeus

Pelecanus Linnaeus, 1758 [Modern genus.]

Pelecanus odessanus Vidgal'm

Pelecanus odessanus Vidgal'm, 1886: 6 [Lectotype from Novaja Slobodka (here selected): a tarsometatarsus; present location unknown. Figured by Vidgal'm 1886, fig. 1-3.

Paralectotype from Novaja Slobodka: a coracoid lacking cranial end; present location unknown. Figured by Vidgal'm 1886, fig. 4.]

Distribution: Late Miocene (MN 11-13) of Novaja Slobodka, Ukraine (Vidgal'm 1886, Miller 1966: 184, Olson 1999c: 508).

Family Messelornithidae Hesse

Messelornithidae Hesse, 1988a: 84 [Type genus: *Messelornis* Hesse, 1988a.]

Distribution: Middle Eocene (MP 11) of Germany, and perhaps also late Eocene to middle Oligocene (MP 17-23) of France. Extralimital record is limited to the middle Eocene of Wyoming (Hesse 1992).

Genus *Messelornis* Hesse

Messelornis Hesse, 1988a: 90 [Type by original designation: *Messelornis cristata* Hesse, 1988a.]

Distribution: Middle Eocene (MP 11) of Germany, and middle Oligocene (MP 23) of France. Extralimital record includes *Messelornis nearctica* Hesse, 1992 from the middle Eocene (Green River Formation) of Wyoming (Hesse 1992).

Messelornis cristata Hesse

Messelornis cristata Hesse, 1988a: 90 [Holotype from Messel: skeleton in three plates; SMF ME-807a-c. Figured by Hesse 1988a, fig. 15-17, and Hesse 1990, pl. 1, fig. 1-3.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Hesse 1988a,b, 1990, 1991, Hesse & Habersetzer 1993).

Genus *Itardiornis* Mourer-Chauviré

Itardiornis Mourer-Chauviré, 1995a: 96 [Type by original designation: *Itardiornis hessae* Mourer-Chauviré, 1995a.]

Remarks: The taxonomic position of this genus is uncertain.

Itardiornis hessae Mourer-Chauviré

Itardiornis hessae Mourer-Chauviré, 1995a: 97 [Holotype from Itardiès: humeral end of left coracoid; USTL ITD-679. Figured by Mourer-Chauviré 1995a, pl. 1, fig. 6-7.]

Distribution: Late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1995a); early Oligocene (MP 21) of Ravet-Lupovici, France (Mourer-Chauviré 1995a); and middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1995a).

Family Gruidae Vigors

Gruidae Vigors, 1825: 488 [Modern family.]

Genus *Palaeogrus* Portis

Palaeogrus Portis, 1884: 362 [Type by monotypy: *Palaeogrus princeps* Portis, 1884.]

***Palaeogrus princeps* Portis**

Ornitocnemis [sic!] *robustus* Zigno in Portis, 1884: 362 [Nomen nudum; no description or indication.]

Palaeogrus princeps Portis, 1884: 362 [Holotype from Monte Zuello: distal end of left tibiotarsus; MPUP, uncatalogued. Figured by Portis 1884, pl. 1, fig. 1-4.]

Grus princeps (Portis): Lydekker 1891a: 165 [New combination.]

Distribution: Middle Eocene (MP 11-13) of Monte Zuello, Italy (Portis 1884, Cracraft 1973: 78).

Remarks: In describing *Palaeogrus princeps*, Portis (1884: 362) mentioned that the holotype was given to him by Achille de Zigno, and that Zigno provisionally named the fossil *Ornitocnemis robustus*. In the preceding paragraph, Portis (1884) referred to two Zigno's papers, where the latter author described other vertebrates from Monte Zuello. Lambrecht (1921: 63, 1933: 518) erroneously listed the first of the two papers (Zigno 1875) as the source of *Ornitocnemis* [sic!; misspelled] *robustus*, but included it in the synonymy of *Palaeogrus princeps*. Brodkorb (1967: 147) used *Ornitocnemis robustus* as the valid name for the bird, attributing it to Zigno ("1876"), and referring to his paper in vol. 20 of the *Memorie*, p. 445. This is apparently just an erroneous citation of Zigno (1875), because neither Cracraft (1973: 78) nor me were able to find any paper by Zigno, which would correspond to the citation in Brodkorb (1967). Moreover, Zigno (1884) described the bird in a special paper, where he compared it with *Ardea* Linnaeus, not mentioning any name for it. It is thus probable, that Zigno never published the name *Ornitocnemis robustus* in a manner required by the ICZN (1999).

***Palaeogrus hordwelliensis* (Lydekker)**

Grus hordwelliensis Lydekker, 1891a: 165 [Holotype from Hordle: distal end of right tibiotarsus; BMNH 30333. Figured by Lydekker 1891a, fig. 36, Cracraft 1973, fig. 35a-d, Harrison & Walker 1976a, pl. 5, fig. f-j.]

Palaeogrus hordwelliensis (Lydekker): Lambrecht 1933: 519 [New combination.]

Ornitocnemis hordwelliensis (Lydekker): Brodkorb 1967: 148 [New combination.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a, Cracraft 1973, Harrison & Walker 1976a).

Genus *Geranopsis* Lydekker

Geranopsis Lydekker, 1891a: 166 [Type by original designation: *Geranopsis hastingsiae* Lydekker, 1891a. This is a senior homonym of *Geranopsis* Milne-Edwards, 1892.]

***Geranopsis hastingsiae* Lydekker**

Geranopsis hastingsiae Lydekker, 1891a: 166 [Holotype from Hordle: left coracoid; BMNH 30331. Figured by Lydekker 1891a, fig. 37, Cracraft 1973, fig. 39a-b, Harrison & Walker 1976a, pl. 5, fig. k-n.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a, Cracraft 1973, Harrison & Walker 1976a).

Remarks: Generic distinction from *Palaeogrus* Portis has not been documented as yet.

Genus *Balearica* Brisson

Balearica Brisson, 1760 [Modern genus.]

***Balearica excelsa* (Milne-Edwards)**

Grus excelsa Milne-Edwards, 1868: pl. 75 [Lectotype from Saint-Gérand-le-Puy (selected by Fischer & Stephan 1971: 577): distal end of right tibiotarsus; MNHN Av-8556. Figured by Milne-Edwards 1867-1869, pl. 75, fig. 5-6; and Cracraft 1973, fig. 36a-d. Being unaware of the action by Fischer & Stephan (1971), Cracraft (1973: 82) selected as a lectotype the same tibiotarsus. His action is invalid (ICZN 1999, Art. 74.1.1). "Paralectotypes" from Saint-Gérand-le-Puy (in MNHN, one in AMNH): right coracoid (Av-8547), shaft of right humerus (Av-8549a), distal end of left humerus (Av-8549), distal ends of two right humeri (Av-8548, Av-8550), right ulna (Av-8552), proximal end of left ulna (Av-8553), proximal end of right ulna (Av-8551), proximal end of right carpometacarpus (AMNH 10583, cast in FSL 443691), distal end of right carpometacarpus (Av-8554), distal end of left tibiotarsus (Av-8555), proximal end of left tarsometatarsus (Av-8558), proximal end of right tarsometatarsus (Av-8557), shaft of left tarsometatarsus (Av-8559), and two pedal phalanges (Av-8560, Av-8561). Figured by Milne-Edwards 1869-1871, pl. 75, fig. 1-4 (proximal end of left tarsometatarsus), 5-6 (right tibiotarsus), 7 (coracoid), 8-9 (distal end of right humerus), 10-11 (proximal end of ulna), 12-15 (pedal phalanx), 16 (another pedal phalanx), pl. 76, fig. 1-2 (carpometacarpus AMNH 10538); and Cracraft 1973, fig. 37a-b (humerus Av-8548), and 37c-d (tarsometatarsus Av-8557). The ulna Av-8553 seems to be too large for the species – Mlíkovský orig.]

Palaeogrus excelsa (Milne-Edwards): Lambrecht 1933: 519 [New combination.]

Ornitocnemus excelsus (Milne-Edwards): Brodkorb 1967: 148 [New combination.]

Balearica excelsa (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérand-le-Puy, France (Milne-Edwards 1869-1871, Cracraft 1973, Cheneval 2000, Mlíkovský orig.); early Miocene (MN 4) of Artenay, France (Mlíkovský orig., material in MNHN); and middle Miocene (MN 6) of Sansan, France (Cheneval 2000).

Remarks: The lectotypical tibiotarsus of *Grus excelsa* Milne-Edwards differs from the same element of *Grus Pallas* and agrees with that of *Balearica* Brisson in having external condyle slightly convex, not notched. The paralectotypical ulna of *Grus excelsa* Milne-Edwards differs from the same element of *Grus Pallas* and agrees with that of *Balearica* Brisson in having external cotyla broad and rounded in anconal view. In absence of the contrary evidence I transfer here *Grus excelsa* from the genus *Grus Pallas* to the genus *Balearica* Brisson as *Balearica excelsa* (Milne-Edwards), new combination.

Genus *Grus Pallas*

Grus Pallas, 1766 [Modern genus.]

***Grus moldavica* (Kuročkin & Ganea)**

Probalearica moldavica Kuročkin & Ganea, 1972: 64 [Holotype from Golboçica: distal end of left tibiotarsus; GIKM 12173/6. Figured by Kuročkin & Ganea 1972, pl. 2, fig. 9, and text-fig. 11.]

Grus moldavica (Kuročkin & Ganea): Mlíkovský, this paper [New combination.].

Distribution: Late Miocene (MN 9) of Golboçica, Moldova (Kuročkin & Ganea 1972).

Remarks: The holotypical tibiotarsus of *Probalearica moldavica* Kuročkin & Ganea differs from the same element of *Balearica*, and agrees with that of *Grus Pallas*, in being anteroposteriorly less flattened, in having the internal condyle directed anteriorly, not outwardly, and especially in having external trochlea notched. In absence of contrary evidence,

I transfer here *Probalearica moldavica* Kuročkin & Ganea to the modern genus *Grus* Pallas as *Grus moldavica* (Kuročkin & Ganea), new combination. Being smaller than the comparably old *Grus pentelici* Gaudry from the late Miocene (MN 12-13) of Greece and Afghanistan (see below), *Grus moldavica* Kuročkin & Ganea seems to be a valid species of cranes.

***Grus pentelici* Gaudry**

Grus Pentelici Gaudry, 1862a: 634 [Lectotype from Pikermi (selected independently by Fischer & Stephan 1971: 576, and Cracraft 1973: 91): distal end of left tarsometatarsus; MNHN # 39. The lectotype was figured by Gaudry 1862a, pl. 16, fig. 12, Gaudry 1862b, pl. 59, fig. 11, and Cracraft 1973, fig. 41a-b. Paralectotypes from Pikermi: two vertebrae (# 28, 29), anterior end of left coracoid (# 30), right humerus (# 31), proximal end of right humerus (# 40), proximal end of left humerus (# 38), right ulna (# 32), fragmentary pelvis (# 34), distal end of right femur (# 35), proximal end of right tibiotarsus (# 36), proximal end of right tarsometatarsus (# 37), distal end of left tarsometatarsus (# 39), and anterior end of pedal digit 3 (# 33). The syntypes are deposited in MNHN, but are uncatalogued. The numbers given above correspond with numbers I attached to avian bones from Pikermi, when I studied them in 1992. Of the syntypes, the humerus # 40 is not from a crane, belonging instead to a flamingo, and the alleged pedal phalanx (# 33) is an unidentifiable scrap of bone. All other specimens are referable to *Grus pentelici* (the vertebrae # 28-29 tentatively). Paralectotypes were figured by Gaudry 1862a, pl. 16, fig. 8 (humerus), 9 (ulna), 10 (pelvis), and 11 (tarsometatarsus), Gaudry 1862b, pl. 59, fig. 1-2 (vertebrae), 23 (coracoid), 4 (humerus), 5 (ulna), 6 ("phalanx"), 7 (pelvis), 8 (femur), 9 (tibiotarsus), and 10 (tarsometatarsus); and Cracraft 1973, fig. 42b-c (tibiotarsus).]

Pliogrus Pentelici (Gaudry): Lambrecht, 1933: 523 [New combination.]

Grus afghana Mourer-Chauviré, Balouet, Jehenne & Heintz, 1985: 180 [Holotype from Molayan: distal end of left tarsometatarsus; MNHN MOL-428. Figured by Mourer-Chauviré et al. 1985, pl. 1, fig. 1-4.]

Distribution: Late Miocene (MN 11) of Csákvár, Hungary (Kretzoi 1957, see also Jánossy 1987a), and late Miocene (MN 12-13) of Pikermi, Greece (Gaudry 1862a,b, 1862-1867, Cracraft 1973, Mourer-Chauviré et al. 1985, Mlíkovský orig.). An extralimital record is from the late Miocene (MN 12) of Molayan, Afghanistan (Mourer-Chauviré et al. 1985). The tentative record from the middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Depéret 1887 sub ?*Grus pentelici*) should be reevaluated.

Remarks: According to Mourer-Chauviré et al. (1985), the holotypical tarsometatarsus of *Grus afghana* Mourer-Chauviré et al. differs from the same element (lectotype) of *Grus pentelici* Gaudry from similarly aged (MN 12) deposits of Pikermi in Greece in having (1) distal foramen closer to the external intertrochlear notch, (2) channel between distal foramen and external intertrochlear notch deep, and (3) in being smaller. None of these characters is valid. Mourer-Chauviré et al. (1985) apparently did not examine the lectotypical tarsometatarsus of *Grus pentelici* directly, probably deriving its description from fig. 41 in Cracraft (1973), which indeed shows the characters as listed by Mourer-Chauviré et al. (1985). Nevertheless, reflections of lights on the bone in that stereo-figure distort its shape and make exact observation of characters impossible. When the bone is examined directly, it shows exactly the same morphological details as listed by Mourer-Chauviré et al. (1985) for *Grus afghana* Mourer-Chauviré et al. (Mlíkovský, pers. observation in MNHN). The holotype of *Grus afghana* is smaller than the lectotype of *Grus pentelici* only by ca. 10 % (see Tab. 1 in Mourer-Chauviré et al. 1985), i.e. both species fall in the same size class. Hence, I synonymize here *Grus afghana* Mourer-Chauviré et al. with *Grus pentelici* Gaudry.

***Grus primigenia* Milne-Edwards**

Grus primigenia Milne-Edwards, 1869: 33 [Holotype from Eyzies: distal end of right tibiotarsus; MNHN Av-Q.1. Figured by Milne-Edwards 1969-1871, pl. 76, fig. 8-11,

Northcote & Mourer-Chauviré 1985, fig. 1/1-5 (right in all cases), and Northcote & Mourer-Chauviré 1988, fig. 3/4.]

Grus melitensis Lydekker, 1890: 408 [Lectotype from Zebbug (selected by Lydekker 1891a: 162, see also Fischer & Stephan 1971: 576): cranial end of right coracoid; BMNH 49365. Figured by Lydekker 1890, pl. 36, fig. 4. Paralectotypes (from Zebbug): distal end of left tibiotarsus (BMNH 49361), distal end of left tarsometatarsus (BMNH 49358). Figured by Lydekker 1890, pl. 36, fig. 2, 2a (tarsometatarsus), and 5, 5a-b (tibiotarsus), and Lydekker 1891a, fig. 35 (tibiotarsus). Lydekker (1891a: 162-163) attributed to *Grus melitensis* six bones or bone fragments, which were not mentioned in the original description of the species. In the same paper Lydekker (1891a: 162, footnote) declared that the partial coracoid (here termed lectotype) and left fragment of pelvis (BMNH 49322m) are types of the species (this being accepted by Brodkorb 1967: 151). Nevertheless, the pelvis fragment was not part of the syntypical series and cannot be included among the syntypes of *Grus melitensis*.]

Sarcogeranus bohatschevi Serebrovskij, 1940a: 767 [Holotype from Binagady: partial skull; MB R-2. Figured by Serebrovskij 1940a, fig. 1e-f.]

Leucogeranus bohatschevi Serebrovskij, 1941a: 473 [New combination, and spelling of the species name emended. This is junior objective synonym of *Sarcogeranus bohatschevi* Serebrovskij, 1940a.]

Grus leucogeranus bogatschevi (Serebrovskij): Dement'ev 1964: 678 [New combination, new rank, subspecies name misspelled.]

Distribution:); Late Pleistocene (MQ 2C) of Avenc de na Corna, Mallorca, Spain (Mourer-Chauviré et al. 1975 and 1977 sub *G. antigone* – see Northcote & Mourer-Chauviré 1988), and Ilford, England (Harrison & Cowles 1977), Zebbug, Malta (Lydekker 1890 and 1891a sub *G. melitensis*, Northcote 1982a, 1984 and 1992 sub *G. melitensis*), Gnien, Malta (Harrison & Cowles 1977 sub *G. melitensis*, Northcote 1984 and 1992 sub *G. melitensis*), and Mnajdra, Malta (Northcote 1982a, 1984 and 1992 sub *G. melitensis*); late Pleistocene (MQ 2C) of Figueira Brava, Portugal (Mourer-Chauviré & Antunes 2000); late Pleistocene (MQ 2D) of Eyzies, France (Milne-Edwards 1869-1871 and 1875a,b), Harpons – layer B-II, France (Saint-Périer 1920), Madelaine – unknown layer, France (Milne-Edwards 1875b), Morin, France (Delpech 1983), Duruthy – layer 3, France (Delpech 1968 and 1983 sub *G. cf. primigenia*), Gouëris, France (Saint-Périer 1927 sub *G. cf. primigenia*), Gourdan, France (Clot & Mourer-Chauviré 1986), Hohen Vielchen, Germany (Soergel 1961 sub *G. antigone*), Pouàs, Eivissa, Balearics (J. A. Alcover in Tyrberg 1998: 422), and Romanelli – unknown layer, Italy (Cassoli 1992 sub *G. cf. leucogeranus*);

Holocene (MQ 2E; Neolithic) of Ehrenstein, Germany (Soergel 1955 sub *G. antigone*); and Holocene (MQ 2E; Iron Age) of Strathclyde, Isle of Jura, Scotland (Harrison & Cowles 1977), and Somerset, England (Harrison & Cowles 1977).

Extralimital record is available from the late Pleistocene (MQ 2C-D) of Binagady, Azerbaijan (Serebrovskij 1940a,b and 1948 sub *Sarcogeranus cf. leucogeranus*, and *Sarcogeranus bohatschevi* or *Leucoceranus bogatschevi*).

Remarks: Bones of *Grus* cranes, larger than the modern *Grus grus* (Linnaeus), were found in the late Pleistocene (MQ 2C-E) deposits of SW and W Europe. The taxonomic treatment of these bones is uncertain. Large cranes from Malta were traditionally separated at the species level as *Grus melitensis* Lydekker, 1890, while large continental cranes from Europe were referred either to the modern Sarus Crane *Grus antigone* (Linnaeus, 1758) or to the extinct *Grus primigenia* Milne-Edwards, 1869 (see above for references). In addition, similarly sized cranes from the late Pleistocene of Azerbaijan were referred to the extinct *Grus bohatschevi* (Serebrovskij, 1940a) and to the modern Siberian Crane *Grus leucogeranus* Pallas, 1773 (Serebrovskij 1940a,b, 1941a, 1948). Mourer-Chauviré et al. (1975, 1977) synonymized *Grus melitensis* with *Grus antigone*, but Northcote (1982a, 1984, 1985, 1992) resurrected the species. Large continental cranes were separated at the species level on zoogeographical

grounds (Harrison & Cowles 1977) and morphologically (Northcote & Mourer-Chauviré 1985). Alcover et al. (1992) held it for possible that *Grus melitensis* and *Grus primigenia* are conspecific. For the time being, I prefer to treat all these cranes as a single extinct species, to which the name *Grus primigenia* Milne-Edwards, 1869 is applicable. Nevertheless, comparisons with similarly large modern *Grus leucogeranus* Pallas, 1773 from Siberia and *Grus carunculatus* (Gmelin, 1789) from Africa are needed (see Fischer & Stephan 1971 for skeletal measurements of the latter two species). It is also well possible, that different bones listed here under the name *Grus melitensis* Lydekker belong to different fossil and/or modern *Grus* species.

***Grus grus* (Linnaeus) – Common Crane**

Ardea Grus Linnaeus, 1758 [Modern species.]

Grus turfa Portis, 1884: 372 [Syntypes from Peschiera: two vertebrae, left coracoid, left humerus, two ulnae, two radii, two carpometacarpi, two scapulae, partial pelvis, two femora, two tibiotarsi, two fibulae, left tarsometatarsus, all possibly from a single individual; present location unknown. Figured by Portis 1884, pl. 2, fig. 1 (tarsometatarsus), 2 (tibiotarsus), 3 (fibula), 4 (femur), 5 (pelvis), 6 (coracoid), 7 (scapula), 8 (humerus), 9 (ulna), 10 (radius), 11 (carpometacarpus), and 12-13 (vertebrae).]

Distribution: Late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a sub “array of *G. grus*”); early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub *G. cf. grus*); and Holocene (MQ 2E) of Peschiera, Italy (Portis 1884 sub *G. turfa*).

Remarks: Lydekker (1891a: 161) and Paris (1912: 292) indicated, that *Grus turfa* Portis is identical with *Grus primigenia* Milne-Edwards, but Lambrecht (1933: 524) showed that *turfa* is inseparable from the modern *Grus grus* (Linnaeus).

***Grus* sp.**

Distribution: Late Pliocene (MN 17) of Puebla de Valverde, Spain (Adrover et al. 1974).

Family Threskiornithidae Poche

Threskiornithidae Poche, 1904 [Modern family.]

Genus *Rhynchaeites* Wittich

Rhynchaeites Wittich, 1899: 103 [Type by monotypy: *Rhynchaeites messelensis* Wittich, 1899.]

Rhynchaeites Wittich: Lambrecht 1921: 52 [Spelling emended.]

Plumumida Hoch, 1980: 40 [Type by original designation: *Plumumida lutetialis* Hoch, 1980.]

Remarks: *Plumumida* Hoch was synonymized with *Rhynchaeites* Wittich by Peters (1983). The taxonomic position of *Rhynchaeites* within the family was not yet studied.

***Rhynchaeites messelensis* Wittich**

Rhynchaeites messelensis Wittich, 1899: 103 [Holotype from Messel: incomplete skeleton in a slab; lost - Peters 1983: 1-2. Figured by Wittich 1899, pl. 2, fig. 1 (pelvis), 2 (sternum), 3 (tarsometatarsus and phalanges gignitorum pedis), 4 (bill), 5 (proximal end of carpometacarpus), 6a-b (distal end of carpometacarpus), 7 (phalanx proximalis digiti majoris), 8 (phalanx distalis digiti majoris), 9 (proximal end of humerus), 10a-b (distal end of humerus), 11 (phalanx distalis digiti minoris), 12 (ulnare), 13 (proximal end of radius), 14a-b (cranial end of coracoid), 15-16 (quadratum), 17 (caudal vertebra). Neotype (selected by Peters 1983: 4): incomplete skeleton in a slab from Messel; SMF ME-1045. Figured by Peters 1983, fig. 2.]

Plumumida lutetialis Hoch, 1980: 40 [Holotype from Messel: incomplete skeleton in a slab; GIH 2183. Figured by Hoch 1980, fig. 2-6.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Wittich 1899, Hoch 1980, Peters 1983, Mayr 1999b).

Remarks: Both *Rhynchaetes messelensis* Wittich and *Plumumida lutetialis* Hoch were described as charadriiform birds of uncertain taxonomic position. Peters (1983) observed that the latter species is synonymous with the former one, and recognized that *Rhynchaetes messelensis* Wittich was an ibis.

Genus *Plegadis* Kaup

Plegadis Kaup, 1829 [Modern genus.]

Milnea Lydekker, 1891a: 169 [Type by original designation: *Milnea gracilis* Lydekker, 1891a. This is a junior homonym of *Milnea* Reichenbach, 1866.]

Plegadis paganus (Milne-Edwards)

Ibis pagana Milne-Edwards, 1868: 450 [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1984a: 82): right tarsometatarsus; MNHN Av-8693. Figured by Milne-Edwards 1867-1869, pl. 70, fig. 1-5, and Cheneval 1984a, pl. 6, fig. 6. "Paralectotypes" from Saint-Gérard-le-Puy (all in MNHN): incomplete skull (Av-8695), two fragmentary mandibles (Av-4849, 4850), fragmentary sternum (Av-8694), 25 coracoids (Av-2919-2921, 8561-8562), scapula (described by Milne-Edwards 1868: 455, but not mentioned by Cheneval 1983a, 1984a; probably lost), furcula (lost – Cheneval 1984a: 86), 22 humeri (Av-8583-8586, 8588-8604, 8640), 13 ulnae (8605-8616, 8642), four radii (Av-8617, 8618, 8620, 8641), 16 carpometacarpi (Av-8643), fragmentary pelvis (Av-8644), 24 femora (Av-8644-8666, 8691), 24 tibiotarsi (Av-8667-8690, 8692), and 38 tarsometatarsi (Av-2922-2926, 2926a, 8697-8727). Figured by Milne-Edwards 1867-1868, pl. 69 (reconstructed skeleton), pl. 70, fig. 1, 6-8 (tibiotarsus Av-8692), 1, 9-11 (femur Av-8691), 12-14 (pelvis Av-8644), 15-17 (sternum Av-8694), 18-20 (coracoid Av-2920), pl. 71, fig. 1-3 (humerus Av-8640), 1, 4-5 (ulna Av-8642), 1, 6-7 (carpometacarpus Av-8643), 8-9 (furcula), 10-12 (skull Av-8695); and Cheneval 1984a, pl. 6, fig. 1a-b (skull Av-8695), 2a-b (coracoid Av-2920), 3a-b (humerus Av-8640), 4a-b (carpometacarpus Av-8643), 5a-b (tibiotarsus Av-8692). The paralectotypical humerus Av-8691 belongs to *Anas* (= *Mionetta*) *blanchardi* (Milne-Edwards) – see Cheneval (1983b: 88, 1984a: 86). Cheneval (1983b: 88) incorrectly listed this specimen as a paralectotype of *Anas blanchardi* Milne-Edwards.

Milnea gracilis Lydekker, 1891a: 169 [Holotype from Saint-Gérard-le-Puy: left humerus; BMNH 47457. Figured by Lydekker 1891a, fig. 38.]

Eudocimus paganus (Milne-Edwards): Sharpe [New combination; fide Lambrecht 1933: 331.]

Plegadis paganus (Milne-Edwards): Olson 1981: 165 [New combination.]

Actiornis pagana (Milne-Edwards): Harrison 1976: 140 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1967-1869, Lydekker 1891a, Olson 1981, Cheneval 1984a).

Remarks: *Milnea gracilis* Lydekker was described in the Oediceptidae (= Burhinidae). Cracraft (1972: 43) indicated that *Milnea gracilis* Lydekker is synonymous with *Ibis pagana* Milne-Edwards, but did not make the synonymization formally. The latter action was done by Cheneval (1984a: 82).

Genus *Geronticus* Wagler

Geronticus Wagler, 1832 [Modern genus.]

***Geronticus perplexus* (Milne-Edwards)**

Ardea perplexa Milne-Edwards, 1869: 108 [Holotype from Sansan: distal end of right humerus; MNHN Sa-1212. Figured by Milne-Edwards 1869-1871, pl. 96, fig. 1-3; Cheneval 2000, fig. 1.]

Proardea perplexa (Milne-Edwards): Gaillard 1939: 79 [New combination.]

Geronticus perplexus (Milne-Edwards): Cheneval 2000: 324 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871, Cheneval 2000).

***Geronticus eremita* (Linnaeus) – Northern Bald Ibis**

Upupa Eremita Linnaeus, 1758 [Modern species.]

Geronticus balcanicus Boev, 1998c: 49 [Holotype from Slivnica: worn proximal end of left carpometacarpus; NMNHS 14-1996. Figured by Boev 1998c, fig. 1a-d.]

Distribution: Late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 1998c, 2000f sub *G. balcanicus*), and Almenara 1, Spain (Sánchez Marco 1996c); early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001); and middle Pleistocene (MQ 2A) of Spinagallo, Italy (Pavia 1999b).

Remarks: This is the only avian species extinct in Europe, but surviving elsewhere. In historical times, this species was known from Ancient Egypt (Kumerloeve 1983), Ancient Greece (DesFayes 1987), and from Central European Alps (Gesner 1555, see also Fatjo 1906, Killermann 1911, Strohl 1917, Hescheler & Kuhn 1949, Geus 1959, G eroudet 1965, Schenker 1977, Kumerloeve 1978, 1984). The Northern Bald Ibis probably disappeared from Europe in the early 17th century, the last records being available from Switzerland (Bauer & Glutz von Blotzheim 1966, 1987, Smith 1970, Cramp & Simmons 1977, Kumerloeve 1984, Matheu & Del Hoyo 1992). The species is currently on the verge of extinction, with the last free population surviving in Morocco (Matheu & Del Hoyo 1992).

I consider arguments presented by Boev (1998c) for separating *G. balcanicus* Boev from the modern *Geronticus eremita* (Linnaeus) insufficient, and I synonymize here the former with the latter species.

Genus *Platalea* Linnaeus

Platalea Linnaeus, 1758 [Modern genus.]

***Platalea leucorodia* Linnaeus – Eurasian Spoonbill**

Platalea leucorodia Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of P rezletice, Czechia (J anossy 1983 sub *P. cf. leucorodia*).

Family Gastornithidae F urbringer

Gastornithidae F urbringer, 1888: 1178 [Type genus: *Gastornis* H ebert, 1855a.]

Diatrymidae Matthew & Granger, 1917: 321 [Type genus: *Diatryma* Cope, 1876.]

Distribution: Late Paleocene to middle Eocene (MP 6-13) of England, Belgium, France and Germany. Extralimital record is available from the late Paleocene through early Eocene of Wyoming, Colorado and New Mexico (see Andors 1988, 1992), and putatively also from the early Eocene of China (Hou 1980, see also Buffetaut 1997a).

Eggshell fragments from the early Eocene (MP 7) of France, described in the genus *Ornitholithes* by Dughi & Sirugue (1962), were tentatively referred to the Gastornithidae (Michajlov 1991: 223, see also Dughi & Sirugue 1962, 1968).

Remarks: Diatrymidae Matthew & Granger is here synonymized with the Gastornithidae F urbringer (see below under *Gastornis* H ebert).

Genus *Gastornis* Hébert

Gastornis Hébert, 1855a: 579 [Type by monotypy: *Gastornis parisiensis* Hébert, 1855a.]

Diatryma Cope, 1876: 11 [Type by monotypy: *Diatryma gigantea* Cope, 1876.]

Barornis Marsh, 1894: 344 [Type by monotypy: *Barornis regens* Marsh, 1894.]

Omorhamphus Sinclair, 1928: 51 [Type by monotypy: *Omorhamphus storchi* Sinclair, 1928.]

Distribution: Late Paleocene (MP 6) to early Eocene (MP 7-8) of England, Belgium, France and Germany.

Remarks: Coues (1884: 825) and Buffetaut (1997a) suggested that *Diatryma* Cope is synonymous with *Gastornis* Hébert. I concur with them, and formally synonymize here *Diatryma* Cope with *Gastornis* Hébert. Brodkorb (1967: 143-144) recognized two North American species of diatrymas, which should henceforth be known as *Gastornis ajax* (Shufeldt, 1913), new combination, and *Gastornis giganteus* (Cope, 1876), new combination.

For dietary habits of *Gastornis* (incl. *Diatryma*) see Andors (1988, 1991, 1992) and Witmer & Rose (1991).

Gastornis parisiensis Hébert

Gastornis parisiensis Hébert, 1855a: 579 [Holotype from Meudon: left tibiotarsus lacking proximal end; lost – Martin 1992: 98. Casts are in MNHN, uncatalogued, BMNH 32384 and KUVF 25681. Figured by Owen 1856a, pl. 3, fig. 1-1a, Milne-Edwards 1867-1868, pl. 28, fig. 1-4, and Martin 1992, fig. 2a-d.]

Gastornis Edwardsii Lemoine, 1878: 13 [Lectotype from Cernay-les-Reims (selected by Martin 1992: 98): partial left tibiotarsus; MNHN L-2457. Figured by Lemoine 1878, pl. 2, fig. 1-2. Paralectotypes from Cernay-les-Reims: a vertebra (MNHN L-2459), proximal end of right femur (MNHN L-2460), partial tarsometatarsus composed from at least 27 fragments (MNHN L-2458). Figured by Lemoine 1878, pl. 1, fig. 1 above, 2 below (proximal end of femur), fig. 1 below, 2 above (distal end of femur), and pl. 3, fig. 1-3 (tarsometatarsus), 4-6 (vertebra).]

Gastornis Klaasseni Newton, 1885a: 362 [Lectotype from Croydon: proximal end of left femur (IGS GSM-1231, figured by Newton 1886a, pl. 28, fig. 4-6, and Harrison & Walker 1977a, pl. 9, fig. f-h). Paralectotypes from Croydon: distal end of right tibiotarsus (IGS GSM-1228, cast BMNH A-86), distal end of left tibiotarsus (IGS GSM-1229, cast BMNH A-86a). Figured by Newton 1886a, pl. 28, fig. 1-3 (right tibiotarsus), 7-11 (left tibiotarsus), and Harrison & Walker 1977, pl. 9, fig. a-e (right tibiotarsus). The syntypical series included also remains of additional three tibiotarsi, now lost – Harrison & Walker 1977a: 38.]

Distribution: Late Paleocene (MP 6) of Walbeck, Germany (Weigelt 1939, 1942), Mesvin, Belgium (Dollo 1883, 1909, Martin 1992), and Cernay-les-Reims, France (Gervais 1873, 1877, Lemoine 1878, 1880, 1881a,b,c 1884, 1885, 1893, Meunier 1882, Repelin 1924, Andors 1992, Martin 1992, Buffetaut 1997a, see also Lydekker 1893a, Matthew & Granger 1917); early Eocene (MP 7) of Meudon, France (Prévost 1855a,b, Hébert 1855a,b,c,d, Lartet 1855, Valenciennes 1855, Duméril 1855, Owen 1856a,b, Milne-Edwards 1867a,b, 1872, Meunier 1882, Repelin 1924, Martin 1992, see also Bonaparte 1856a,b, Woodward 1886, Lavocat 1899, Russell et al. 1990), Passy, France (Milne-Edwards 1867a,b, 1872), and Croydon, England (Newton 1885a,b, 1886a,b, 1889, Harrison & Walker 1977a, Martin 1992); and early Eocene (MP 8-9) of Mutigny, France (Martin 1992). General reference: Buffetaut 1997b,c.

Remarks: *Gastornis edwardsii* Lemoine and *Gastornis klaasseni* Newton were synonymized with *Gastornis parisiensis* Hébert by Martin (1992).

Lemoine (1881c, pl. 11) reconstructed the skeleton of *Gastornis* using syntypes of *Gastornis parisiensis* and a number of referred material, some of which was later shown to belong to turtles and fishes (Martin 1992, fig. 12; see also Feduccia 1996: 236, Buffetaut 1997c, fig. 1). This reconstruction later entered general monographs (e.g. Meunier 1898,

Lambrecht 1933, fig. 165, Piveteau 1955, fig. 59, Müller 1968, fig. 722), and markedly influenced taxonomic fate of the European *Gastornis* (see Martin 1992, Buffetaut 1997a,b,c).

***Gastornis russeli* Martin**

Gastornis russeli Martin, 1992: 102 [Holotype from Berru: left tarsometatarsus; MNHN R-3560. Figured by Martin 1992, fig. 6a-d, 7a.]

Distribution: Late Paleocene (MP 6) of Berru, France (Martin 1992).

***Gastornis sarasini* (Schaub)**

(?) *Diatryma sarasini* Schaub, 1929b: 588 [Lectotype from Monthelon (here selected): distal end of left tarsometatarsus; BaM TS-94. Figured by Schaub 1929b, fig. 1-3, 6, and Gaillard 1936, fig. 7, 7a. Paralectotype from Monthelon: phalanx I digiti II; NMB TS-8. Figured by Schaub 1929b, fig. 8.]

Diatryma geiselensis Fischer, 1978: 134 [Holotype from Geiseltal XIII: right scapulocoracoid; GM Av-G?/Dia.2. Figured by Fischer 1962, pl. 4, fig. 2.]

Gastornis sarasini (Schaub): Mlíkovský, this paper [New combination.]

Distribution: Early Eocene (MP 10) of Monthelon, France (Schaub 1929a,b, Gaillard 1936); middle Eocene (MP 11) of Messel, Germany (Berg 1965 sub *Diatryma* cf. *steini*), Geiseltal XIII, Germany (Fischer 1962, 1978), and Geiseltal XIV, Germany (Fischer 1962, 1978); and middle Eocene (MP 13) of Geiseltal XXXV, Germany (Fischer 1978), and Geiseltal XXI, Germany (Fischer 1978).

Remarks: There is no evidence that *Diatryma geiselensis* Fischer was different from *Diatryma sarasini* Schaub. They coincide in age, distribution and morphology, so that I synonymize here *Diatryma geiselensis* Fischer with *Diatryma sarasini* Schaub. The species should bear the name *Gastornis sarasini* (Schaub), new combination.

Fischer (1962: 27) hypothesized that *Diatryma sarasini* Schaub could be a subspecies of *Diatryma steini* Matthew & Granger, described from the early Eocene (Willwood Formation) of South Elk Creek, Wyoming (Matthew & Granger 1917), and Berg (1965) tentatively attributed an imprint of a femur from Messel, Germany, to the same species. Until compared directly, I consider the European Eocene gastornithids as separated from the North American ones at the species level.

Family Eogruidae Wetmore

Eogruidae Wetmore, 1934a: 3 [Type genus: *Eogrurus* Wetmore, 1934.]

Ergilornithidae Kozlova, 1960: 329 [Type genus: *Ergilornis* Kozlova, 1960.]

Distribution: Late Miocene (MN 9-13) of Moldova, Ukraine and Greece. Rich extralimital record is limited to the Tertiary of Asia, incl. the middle/late Eocene of China (Wetmore 1934a, Kuročkin 1981), Mongolia (Kuročkin 1981), and Kazakhstan (Bendukidze 1971, 1972a, Kuročkin 1981), early Oligocene of Mongolia (Kozlova 1960, Cracraft 1973, Kuročkin 1981), early Miocene of Kazakhstan (Karchu 1997), late Miocene of Iran (Mecquenem 1908, 1925, Cracraft 1973, Kuročkin 1981), China (Brodkorb 1967) and Krasnodar Province in Russia (Bendukidze 1972b, Kuročkin 1981), and early/middle Pliocene of Kazakhstan (Kuročkin 1981) and Mongolia (Kuročkin 1981, 1985). In addition, Kuročkin (1985: 61) opened the possibility, that large phalanges digitorum pedis from Mioocene Siwalik faunas of Pakistan, described by Davies (1880), belong to this family. However, another alleged record from the Siwalik faunas in the neighboring Pakistan (Harrison & Walker 1982) was rejected by Karchu (1997), so that the former existence of the Eogruidae in the Indian sub-continent remains unproven. Avian footprints described by Lambrecht (1938) from the late Miocene of Hamrin Mountains in Iran as *Urmiornis abeli* do not belong to the Eogruidae (Vjalov 1989). General reference: Kuročkin (1982).

Genus *Amhipelargus* Lydekker

Amhipelargus Lydekker, 1891a: 68 [Type by original designation: *Amhipelargus majori* Lydekker, 1891a. Kuročkin's (1985: 60) statement, that the type of this genus is *Urmionis maraghanus* Mecquenem, 1925, is erroneous.]

Urmionis Mecquenem, 1908: 54 [Type by monotypy *Urmionis maraghanus* Mecquenem, 1908.]

Urmionis Mecquenem, 1925: 27 [Type by monotypy *Urmionis maraghanus* Mecquenem, 1925. Again marked as "nov. gen." This is thus both junior homonym and junior objective synonym of *Urmionis* Mecquenem, 1908.]

Distribution: Late Miocene (MN 9/10 to 11/13) of Moldova, Ukraine and Greece. Extralimital record is known from the early Miocene of Kazakhstan (Karchu 1997), late Miocene of Iran (Mecquenem 1908, 1925, Cracraft 1973, Kuročkin 1981), and early Pliocene of Kazakhstan (Kuročkin 1981) and Mongolia (Kuročkin 1985). The alleged record from the late Miocene of Pakistan (Harrison & Walker 1982) is invalid (Karchu 1997).

Remarks: Kuročkin (1985) and Olson (1985a) followed Harrison's (1981) conjecture and included *Urmionis* Mecquenem in *Amhipelargus* Lydekker. Karchu (1997) separated the genera again, but his data only underscore the specific status of *majori*, being not sufficient for its separation at the generic level.

Amhipelargus ukrainus (Kuročkin)

Urmionis ukrainus Kuročkin, 1981: 78 [Holotype from Hrebnyky: distal end of left tarsometatarsus; IZAN 25-2589. Figured by Burčak-Abramovič 1951, fig. 1/3-4, 2/5-8.]

Amhipelargus ukrainus (Kuročkin): Kuročkin, 1985: 60 [New combination.]

Distribution: Late Miocene (MN 9-10) of Kolkotova Balka, Moldova (Laskarev 1908 and 1912: 41 sub *Urmionis* sp., Burčak-Abramovič 1951 sub *U. maraghanus*, Cracraft 1973 sub *U. maraghanus*, Kuročkin 1981); late Miocene (MN 11) of Hrebnyky, Ukraine (Burčak-Abramovič 1951 sub *U. maraghanus*, Kuročkin 1981); and late Miocene (MN 11-13) of Čebotarevka, Ukraine (Umans'ka 1973 sub *Urmionis* sp., Kuročkin 1981). An extralimital record is available from the late Miocene of Armavir, Krasnodar Province, Russia (Bendukidze 1972b sub *U. maraghanus*, Kuročkin 1981).

Amhipelargus majori Lydekker

Amhipelargus majori Lydekker, 1891a: 69 [Holotype from Samos 1: distal end of left tibiotarsus; BMNH A-123. Figured by Lydekker 1891a, fig. 18, Harrison 1981, fig. 1a-d, 2a.]

Urmionis Maraghanus Mecquenem, 1925: 27 [Lectotype from Maragha (selected by Cracraft 1973: 74): left tarsometatarsus; MNHN uncatalogued. Figured by Mecquenem 1908, fig. 16, Mecquenem 1925, fig. 16; and Cracraft 1973, fig. 33a-d. Paralectotype from Maragha: distal end of right (not left as stated by Mecquenem 1925: 27) tibiotarsus. Figured by Mecquenem 1908, fig. 16; Mecquenem 1925, fig. 16-17; Cracraft 1973, fig. 34a-d). Kuročkin (1981: 78) erroneously called the lectotypical tibiotarsus a holotype.

Amhipelargus maraghanus (Mecquenem): Kuročkin 1985: 60 [New combination.]

Distribution: Late Miocene (MN 11) of Samos 1, Greece (Lydekker 1891a, Harrison 1981, Karchu 1997). An extralimital record is available from the late Miocene of Maragha, Iran (Mecquenem 1908, 1925, Burčak-Abramovič 1951, Cracraft 1973, Harrison 1981, Kuročkin 1981, Karchu 1997). Bones from the late Miocene of Moldova, Ukraine and Krasnodar Province in Russia (Lambrecht 1933: 521, Burčak-Abramovič 1951: 83, Bendukidze 1972b, Cracraft 1973: 74) were later assigned to *Amhipelargus ukrainus* (Kuročkin) by Kuročkin (1981). The alleged extralimital record from the early Pliocene of Kalmakpai, Kazakhstan (Cracraft 1973) became the holotype of *Urmionis orientalis* Kuročkin, 1981.

Remarks: *Amhipelargus majori* Lydekker was transferred from the Ciconiidae to the

Eogruidae by Harrison (1981). The paralectotypical tibiotarsus of *Urmiornis maraghanus* Mecquenem agrees in size and morphology with the holotype of *Amhipelargus majori* Lydekker. Considering the agreement in age and geographic distribution, I synonymize here the former with the latter species (see also Harrison 1981: 112; for an opposite view see Karchu 1997: 105).

Family Otididae Gray

Otididae Gray, 1840 [Modern family.]

Gryzajidae Brodkorb, 1967: 175 [Type genus: *Gryzaja* Zubareva, 1939.]

Remarks: The genus *Gryzaja* Zubareva was described in the Otididae, but Brodkorb (1967: 175) separated it at the family level. I return here the genus to the Otididae, synonymizing thus Gryzajidae Brodkorb with the Otididae Gray. All fossil and modern genera of the Otididae are in need of revision.

Genus *Miootis* Umans'ka

Miootis Umans'ka, 1979a: 40 [Type by original designation: *Miootis compactus* Umans'ka, 1979a.]

Miootis compactus Umans'ka

Miootis compactus Umans'ka, 1979a: 40 [Holotype from Nova Emetivka: left carpometacarpus; IZAN 25-2891. Figured by Umans'ka 1979a, fig. 1a,b,v.]

Distribution: Late Miocene (MN 13) of Nova Emetivka, Ukraine (Umans'ka 1979a).

Remarks: Taxonomic identity of this genus and species is in need of confirmation.

Genus *Chlamydotis* Lesson

Chlamydotis Lesson, 1839 [Modern genus.]

Chlamydotis mesetaria Sánchez Marco

Chlamydotis mesetaria Sánchez Marco, 1990a: 225 [Holotype from Layna: right tibiotarsus; MNCN, uncatalogued. Figured by Sánchez Marco 1990a, text-fig. 1/1-2, 2/1-4, pl. 1, fig. 1-2, pl. 2, fig. 1-4.]

Distribution: Early Pliocene (MN 15) of Layna, Spain (Sánchez Marco 1990a).

Chlamydotis undulata (Jacquin) – Houbara Bustard

Otis undulata Jacquin, 1784 [Modern species.]

Otis khosatzkii Bocheński & Kuročkin, 1987b: 175 [Holotype from Etulia: distal end of right tibiotarsus; PIN 2614-56. Figured by Bocheński & Kuročkin 1987b, pl. 1, fig. 1-3.]

Otis khosatzkii beremendensis Jánossy, 1991: 24 [Holotype from Beremend 15: a "very damaged" tarsometatarsus; NMB V-90-11. Not figured.]

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b and 1990b sub *Tetrax* sp., Jánossy 1991, 1992), and Etulia, Moldova (Bocheński & Kuročkin 1987b); and late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1999e: 70 and 2000a sub *Otis* aff. *khosatzkii*).

Remarks: The available material of *Otis khosatzkii* Bocheński & Kuročkin is not reasonably separable from the modern *Chlamydotis undulata* (Jacquin). Consequently, I synonymize here the former with the latter species. The taxonomic identity of *Otis khosatzkii beremendensis* Jánossy remains uncertain.

Genus *Otis* Linnaeus

Otis Linnaeus, 1758 [Modern genus.]

***Otis affinis* Lydekker**

Otis affinis Lydekker, 1891a: 168 [Holotype from Schnaitheim (= misprint for Steinheim according to Ammon 1918: 55, see also Lambrecht 1933: 528, and Mlíkovský & Hesse 1996: 635): crushed postcranial skeleton in slab; BMNH 36745. Figured by Lambrecht 1933, fig. 157b.]

Chlamydotis affinis (Lydekker): Brodkorb 1967: 174 [New combination.]

Distribution: Middle Miocene (MN 7) of Steinheim, Germany (Lydekker 1891a, Lambrecht 1933, Sánchez Marco 1990a).

Remarks: Mlíkovský (1992b: 441) returned the species to the genus *Otis* Linnaeus.

***Otis bessarabensis* Kessler & Gál**

Otis bessarabensis Kessler & Gál, 1996: 77 [Holotype from Chişinău: distal end of left ulna; LPUI 62-MS. Figured by Kessler & Gál 1996, fig. 3a-b.]

Distribution: Late Miocene (MN 9) of Chişinău, Moldova (Kessler & Gál 1996).

Remarks: Taxonomic identity of this species needs confirmation.

***Otis tetrax* Linnaeus – Little Bustard**

Otis Tetrax Linnaeus, 1758 [Modern species.]

Otis (Tetrax) kalmani Jánossy, 1972: 50 [Holotype from Betfia 2: cranial end of left coracoid; GIB Ob-4845. Not figured.]

Otis paratetrax Bocheński & Kuročkin, 1987b: 179 [Holotype from Etulia: cranial end of left coracoid; PIN 2614-49. Figured by Bocheński & Kuročkin 1987b, pl. 1, fig. 6-7.]

Distribution: Late Pliocene (MN 16) of Etulia, Moldova (Bocheński & Kuročkin 1987b sub *O. paratetrax*); late Pliocene (MN 17) of Kryžanovka 1, Ukraine (Dubrovo & Kapelist 1979 sub *O. cf. tetrax*); early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992), Betfia 2, Romania (Čapek 1917, Jánossy 1972 and 1980); early Pleistocene (MQ 1a) of Nagyarsánhegy 1-4, Hungary (Jánossy 1980a sub *O. kalmani*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5, Czechia (Jánossy 1972), and Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b).

***Otis tarda* Linnaeus – Great Bustard**

Otis Tarda Linnaeus, 1758 [Modern species.]

Otis brevipes Giebel, 1847: 26 [Holotype from Seveckenberg: distal end of right tarsometatarsus; MNHN, uncatalogued. Not figured.]

Otis lambrechtii Kretzoi, 1941: 253 [Holotype from Betfia 5: fragmentary distal end of a tarsometatarsus; GIB, uncatalogued. Not figured.]

Distribution: Early Pliocene (MN 14) of Vojničevo, Ukraine (Vojinstvens'kyj 1967 sub *O. cf. tarda*); late Pliocene (MN 16) of Čișmiki, Moldova (Bocheński & Kuročkin 1987b sub *Otis* sp. - group of *O. tarda*); late Pliocene (MN 18) of Villány 3, Hungary (Jánossy 1980a);

early Pleistocene (MQ 1a) of Osztramos 2, Hungary (Jánossy 1980), Beremend 17, Hungary (Jánossy 1992 sub *O. cf. lambrechtii*), Nagyarsánhegy 1-4, Hungary (Jánossy 1980), and Betfia 5, Romania (Jánossy 1980a sub *O. lambrechtii*); middle Pleistocene (MQ 2A) of Hundsheim, Austria (Jánossy 1974b sub *O. cf. lambrechtii*, Mlíkovský orig.); and late Pleistocene (MQ 2C) of Seveckenberg, Germany (Giebel 1847, Mlíkovský orig.).

Remarks: I restudied the holotype of *Otis brevipes* Giebel and the description of *Otis lambrechtii* Kretzoi, and I synonymize here both of these species with the modern *Otis tarda* Linnaeus (Mlíkovský orig.).

***Otis* sp.**

Distribution: Early Pliocene (MN 15) of Malușteni-Berești, Romania (Kessler 1979).

Genus *Gryzaja* Zubareva

Gryzaja Zubareva, 1939: 607 [Type by monotypy: *Gryzaja odessana* Zubareva, 1939.]

Gryzaja odessana Zubareva

Gryzaja odessana Zubareva, 1939: 607 [Lectotype from Odesa – catacombs (here selected): distal end of right tibiotarsus; IZAN 6617. Figured by Zubareva 1939, 4 unnumbered figs. on p. 607; Zubareva 1948, fig. 7c, 8c; and Dement'ev 1964, fig. 690a,b. Paralectotype: distal end of left tibiotarsus from Odesa–catacombs; IZAN 6618. Not figured.]

Chlamydotis pliodeserti Serebrovs'kyj, 1941a: 474 [Syntypes from Odesa – catacombs: right coracoid and cranial end of left coracoid; present location unknown. Figured by Serebrovs'kyj 1941a; and Dement'ev 1964, fig. 685.]

Otis gryzaja Vojinstvens'kyj, 1967. [New name for *Gryzaja odessana* Zubareva, 1939, hence its junior objective synonym.]

Distribution: Early Pliocene (MN 15) of Odesa – catacombs, Ukraine (Zubareva 1939, 1948, Serebrovs'kyj 1941, Vojinstvens'kyj 1959a, 1967); and late Pliocene (MN 16) of Etulia, Moldova (Chozackij & Kuročkin 1966, Kuročkin & Chozackij 1972, Bocheński & Kuročkin 1987b), and Kotlovina, Ukraine (Kuročkin & Chozackij 1972).

Remarks: *Chlamydotis pliodeserti* Serebrovs'kyj was synonymized with *Gryzaja odessana* Zubareva by Vojinstvens'kyj (1959a: 200).

Family Ciconiidae Sundevall

Ciconiidae Sundevall, 1836 [Modern family.]

Genus *Grallavis* Cheneval

Grallavis Cheneval 1984b: 44 [Type by original designation: *Propelargus edwardsi* Lydekker, 1891c.]

Grallavis edwardsi (Lydekker)

Propelargus (?) *edwardsi* Lydekker, 1891c: 479 [Holotype from Saint-Gérard-le-Puy: right coracoid; BMNH A-427. Figured by Lydekker 1891c, fig. A.]

Grallavis edwardsi (Lydekker): Cheneval 1984b: 44 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Lydekker 1891a sub *Propelargus* (?) sp., Lydekker 1891c, Lambrecht 1933: 319-320, Cheneval 1984b).

Genus *Leptoptilos* Lesson

Leptoptilos Lesson, 1831 [Modern genus.]

Leptoptilos pliocenicus Zubareva

Leptoptilos [sic] *pliocenicus* Zubareva, 1948: 114 [Syntypes from Odesa – catacombs: fragment of right premaxilla (IZAN 8025), two fragments of left mandible from a single individual (IZAN 8026), fragment of right mandible (IZAN 7042), right quadratum (IZAN 7042), shaft of right humerus (IZAN 6939), left carpometacarpus (IZAN 8060), left tarsometatarsus lacking proximal end (IZAN 8024), and phalanx 1 digiti 3 (IZAN 6502). Figured by Zubareva 1948, fig. 1 (quadratum), fig. 2a-c (right mandible), fig. 3 (carpometacarpus), fig. 4a-b (tarsometatarsus), fig. 6a-b (phalanx); and Dement'ev 1964, fig. 695a (carpometacarpus), b,v (tarsometatarsus), g,d (phalanx).]

Distribution: Early Pliocene (MN 15) of Odesa – catacombs, Ukraine (Zubareva 1948).

Genus *Ciconia* Brisson

Ciconia Brisson, 1760 [Modern genus.]

***Ciconia sarmatica* Grigorescu & Kessler**

Ciconia sarmatica Grigorescu & Kessler, 1977: 99 [Holotype from Credința: proximal end of right carpometacarpus; LPUB 259. Figured by Grigorescu & Kessler 1977, pl. 3, fig. 2a-b, 4a, pl. 5, fig. a.]

Distribution: Middle Miocene (MN 8) of Credința, Romania (Grigorescu & Kessler 1977).

Remarks: Taxonomic status of this species is in need of revision.

***Ciconia gaudryi* Lambrecht**

Ciconia gaudryi Lambrecht, 1933: 323 [Holotype from Pikermi: humerus; perhaps in MNHN. I was not able to see the specimen during my visits to MNHN in 1992 and 1999, but other avian bones from the site collected by Gaudry were present there. Figured by Gaudry 1862-1867, pl. 59, fig. 12.]

Distribution: Late Miocene (MN 12-13) of Pikermi, Greece (Gaudry 1862a,b, 1862-1867 sub "Cicogne indéterminée", Lambrecht 1933).

***Ciconia stehlini* Jánossy**

Ciconia stehlini Jánossy, 1992: 16 [Syntypes from Beremend 15: proximal end of tarsometatarsus and distal end of tarsometatarsus; NMB V-91.151. Figured by Jánossy 1992, fig. 4/1-3.]

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b and 1990b sub *Ciconia* sp., Jánossy 1992); and early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992).

Remarks: Taxonomic status of this species is in need of revision.

Genus *incertae sedis*

"*Cygnus*" *bilinicus* Laube

Cygnus bilinicus Laube, 1909: 161 [Holotype from Břešťany: proximal (not distal as stated by Laube 1909) end of an ulna, proximal (not distal as stated by Laube 1909) end of an radius and a carpometacarpus, all apparently from a single individual; RMT 232. Figured by Laube 1909, pl. 1; and Mlíkovský & Švec 1989, pl. 2, fig. b. The bones are too flattened to reveal from which body side they originated.]

Aquilavus bilinicus (Laube): Brodkorb 1964: 264 [New combination; in *Aquilavus* Brodkorb, 1964.]

Distribution: Early Miocene (MN 3) of Břešťany, Czechia (Laube 1909, Mlíkovský & Švec 1989, Mlíkovský 1999c).

Remarks: Described as a swan, and later believed to represent an accipitrid raptor (Lambrecht 1933: 383, Brodkorb 1964: 264, 1978, Howard 1964). Mlíkovský & Švec (1989) showed that it is an indeterminate stork from the family Ciconiidae.

Family Cathartidae Lafresnaye

Cathartidae Lafresnaye, 1839 [Modern family.]

Remarks: In summarizing the evidence for the occurrence of the New World vultures (Cathartidae) in the Old World, Cracraft & Rich (1972) recognized four monotypic genera of Old World cathartids, all from the Paleogene of Europe. Of them, three genera were already

removed from the family, incl. *Eocathartes* Lambrecht (P. Houde in Olson 1985a: 191), *Plesiocathartes* Gaillard (Jollie 1977c: 112), and *Amphiserpentarius* Milne-Edwards (Mourer-Chauviré & Cheneval 1983). The last genus, *Tapinopus* Milne-Edwards, is relegated to the category of Aves incertae sedis in this paper (see below).

The opinion of Cracraft & Rich (1972), that the Cathartidae originated in the Old World, was accepted by Alvarenga (1985) and Emslie (1988), but seems to have no support in the fossil record. There is currently no evidence, that the Cathartidae occurred in the Cenozoic of Europe or elsewhere in the Old World.

Order Anseriformes Wagler

Anseres Wagler, 1831 [Modern order.]

Family Phoenicopteridae Bonaparte

Phoenicopteridae Bonaparte, 1831 [Modern family.]

Palaeolontidae Stejneger, 1885: 154 [Type genus: *Palaelodus* Milne-Edwards, 1863. This is incorrect original spelling of Palaelodidae – see Mlíkovský 2000f.]

Palaelodidae Stejneger: Fürbringer 1888: 1565 [Spelling emended.]

Juncitarsinae Peters, 1987b: 141 [Type genus: *Juncitarsus* Olson & Feduccia, 1980a.]

Remarks: Palaelodidae Stejneger was synonymized with the Phoenicopteridae Bonaparte by Olson & Feduccia (1980a: 45).

Genus *Juncitarsus* Olson & Feduccia

Juncitarsus Olson & Feduccia, 1980a: 48 [Type by original designation: *Juncitarsus gracillimus* Olson & Feduccia, 1980a.]

Distribution: Middle Eocene (MP 11) of Germany. The extralimital record is limited to *Juncitarsus gracillimus* Olson & Feduccia, 1980a from the middle Eocene (early Bridgerian) of Wyoming (Olson & Feduccia 1980a). See also Ericson (1999).

Juncitarsus merkei Peters

Juncitarsus merkei Peters, 1987b: 141 [Holotype from Messel: incomplete skeleton in slab; coll. Kessler, uncatalogued. Cast in SMF, uncatalogued. Figured by Peters 1987b, fig. 1, 3-8.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Peters 1987b). Tentatively referred also from the middle Eocene (early Bridgerian) of Wyoming (Peters 1987b: 142).

Genus *Kashinia* Harrison & Walker

Tenuicrus Harrison & Walker, 1976a: 344 [Type by original designation: *Tenuicrus magnum* Harrison & Walker, 1976a.]

Kashinia Harrison & Walker, 1979c: 110 [New name for *Tenuicrus* Harrison & Walker, 1976a; preoccupied by *Tenuicrus* Womersley, 1940.]

Kashinia magna Harrison & Walker

Tenuicrus magnum Harrison & Walker, 1976a: 344 [Holotype from Hordle: incomplete right coracoid; BMNH 30327. Figured by Harrison & Walker 1976a, pl. 6, fig. h-k.]

Kashinia magna (Harrison & Walker): Harrison & Walker, 1979c: 110 [New combination.]

Distribution: Late Eocene (MP 17) of Hordle, England (Harrison & Walker 1976a).

Remarks: As judged from figures, the holotypical coracoid differs from the same element of the Recurvirostridae (where the species was placed by Harrison & Walker 1976a), and agrees with that of *Juncitarsus* Olson & Feduccia (see Ericson 1999, fig. 2) in having medial angle pointed, shaft shorter and more robust, and head less bent. It is thus possible, that *Kashinia magna* (Harrison & Walker) represents an ancestral flamingo. However, direct study of the specimens is necessary to confirm this conclusion.

Genus *Palaelodus* Milne-Edwards

Palaelodus Milne-Edwards, 1863: 158 [Type by subsequent designation (Milne-Edwards 1869: 59): *Palaelodus ambiguus* Milne-Edwards, 1863. The generic name was often misspelled as *Paloelodus* (see Cheneval 1983c: 181), but I am not aware of any case, where this action could be unequivocally interpreted as a deliberate emendation of *Palaelodus*.]

Probalearica Lambrecht, 1933: 519 [Type by monotypy: *Grus problematica* Milne-Edwards, 1871.]

Megapaloelodus Miller, 1944: 86 [Type by monotypy: *Megapaloelodus connectens* Miller,

1944.]

Distribution: Late Oligocene (MP 25-30) of France and Germany, early Miocene (MN 1-4) of France, Germany and Czechia, and middle Miocene (MN 5-8) of Germany, Czechia and Romania. The genus has a rich extralimital record, which includes early Miocene of South Dakota (A.H. Miller 1944), middle Miocene of California (L.H. Miller 1950, 1952), late Miocene of Oregon (Brodkorb 1961), California (Howard 1971, tentatively), Texas (Becker 1987: 49), and Argentina (Noriega 1995), and late Oligocene through middle Miocene of Australia (Baird & Vickers-Rich 1998). A much younger record from the middle Pleistocene of Australia (Baird & Vickers-Rich 1998) is based on a single bone fragment and should be reevaluated.

Remarks: Olson & Feduccia (1980a: 45) indicated, that *Megapaloelodus* Miller "is very similar to, and may not be separable from *Palaelodus* [Milne-Edwards]". I agree and I synonymize here the former with the latter genus.

***Palaelodus ambiguus* Milne-Edwards**

Palaelodus ambiguus Milne-Edwards, 1863: 159 [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1983c: 181): left tarsometatarsus; MNHN Av-8914. Figured by Cheneval 1983c, pl. 1, fig. 1a-b. Not figured by Milne-Edwards 1869-1871, pl. 83, fig. 1-4, where a side-reversed right tarsometatarsus is shown (contra Cheneval 1983c: 181). "Paralectotypes" from Saint-Gérard-le-Puy (based on the list in Cheneval 1983a: 123-125; all in MNHN): mandible (Av-260 coracoids (Av-1-36, 38-48, 50-90, 92-259, 340-341, 9299, 9302), 287 scapulae (Av-342-628), 80 furculae (Av-260-339), 32 sterna (Av-2768-2805, 2947-2960), 299 humeri (Av-629-639, 641-643, 645-650, 652, 654-664, 666, 668-768, 770-804, 806-849, 851-857, 859-900, 902-945, 6529, 6531-6532), 412 ulnae (Av-1851, 1952-1978, 1981-2018, 2020-2041, 2043-2050, 2052-2069, 2071-2072, 2074-2081, 2083-2107, 2109-2114, 2231-2241, 2244-2294, 2296-2304, 2349-2365, 2367-2412, 2414-2428, 2430-2437), 235 radii (Av-1797, 1799-1800, 1803-1821, 1823-1827, 1829, 1831-1850, 1852-1879, 1881-1883, 2051, 2115-2130, 2132-2152, 2155-2181, 2183-2230, 2305-2316, 2318-2338, 2340-2349), 243 carpometacarpi (Av-1830, 2182, 2439-2440, 2442-2448, 2450-2456, 2458-2476, 2478-2486, 2488, 2490-2541, 2516-2518, 2520-2524, 2526-2538, 2540-2551, 2553-2558, 2560-2572, 2574-2578, 2580-2587, 2589-2615, 2617, 2619-2669, 2701), 185 femora (Av-2976-2989, 2991-3145, 3147, 6503-6510, 6512-6518), 377 tibiotarsi (Av-3221-3225, 3227-3253, 3255-3260, 3262-3361, 3363-3366, 3368-3379, 3414-3462, 3464-3490, 3500-3519, 3521-3539, 3560-3580, 3582-3593, 3595-3613, 3615-3650, 3652-3657, 3659-3661, 6479-6482, 6484, 6487, 6488, 6520-6523), and 281 tarsometatarsi (Av-3662-3757, 3759-3770, 3772-3776, 6528, 8761-8778, 8780-8835, 8837-8893, 8895-8913, 8915-8923, 9027-9254). Figured by Gervais 1859, pl. 51, fig. 9 (tarsometatarsus), Milne-Edwards 1869-1871, pl. 82 (reconstructed skeleton), pl. 83, fig. 1-4 (tarsometatarsus), 5-8 (tibiotarsus), 9-12 (femur), 13-15 (coracoid), pl. 84, fig. 4-5 (humerus), 9-11 (mandible), pl. 85, fig. 4-5 (ulna), 5 (radius), 6-9 (carpometacarpus); Lambrecht 1933, fig. 117b (tarsometatarsus), fig. 118 (reconstructed skeleton); Rothausen 1966, fig. 1 (skeletal reconstruction); and Cheneval 1983a, pl. 11, fig. 1a-b (coracoid), 2a-b (humerus), 3a-b (radius), 4a-b (ulna), 5a-b (carpometacarpus), pl. 12, fig. 1a-b (femur), 2a-b (tibiotarsus).]

Palaelodus gracilipes Milne-Edwards, 1863: 160 [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1983c: 184): right tarsometatarsus; MNHN Av-9255. Figured by Milne-Edwards 1869-1871, pl. 85, fig. 12-16; and Cheneval 1983c, pl. 1, fig. 3a-b. "Paralectotypes" from Saint-Gérard-le-Puy (based on the list in Cheneval 1983a: 132-133; all in MNHN): coracoid (Av-9286), 10 humeri (Av-640, 644, 651, 667, 9282-9285, 9670-9671), 10 ulnae (Av-2019, 2042, 2070, 2073, 2082, 2243, 2295, 2366, 2413, 2429), two radii (Av-2317, 2339), 12 carpometacarpi (Av-2441, 2457, 2487, 2515, 2573, 2579, 2616, 2618, 9287-9290), nine femora (Av-9272-9280), 19 tibiotarsi (Av-3226, 3254, 3261, 3463,

3520, 3594, 3614, 3651, 3658, 6483, 6485-6486, 6489, 6536-6537, 9268-9271), 87 tarsometatarsi (Av-6490-6502, 8779, 8836, 9256-9627). Figured by Milne-Edwards 1869-1871, pl. 86, fig. 1-4 (femur), 5-7 (tibiotarsus), 8-11 (carpometacarpus), 12-14 (coracoid), 15-16 (humerus). Cheneval (1983c: 184) incorrectly listed among the paralectotypes of *Palaelodus gracilipes* also the holotype of *Palaelodus minutus* Milne-Edwards.]

Palaelodus crassipes Milne-Edwards, 1863: 160 [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1983c: 185): proximal end of right tarsometatarsus; MNHN Av-9321. Figured by Cheneval 1983c, pl. 1, fig. 4a-b. Not figured by Milne-Edwards 1869-1871, pl. 88, fig. 4-7, which shows a side-reversed left tarsometatarsus (contra Cheneval 1983c: 185). "Paralectotypes" from Saint-Gérard-le-Puy (based on the list in Cheneval 1983a: 135-136): six coracoids (Av-37, 49, 9296-9298, 9303), eight humeri (Av-653, 665, 805, 850, 901, 6530, 9304-9305), three ulnae (Av-2108, 6533, 9306), eight radii (Av-1822, 1880, 2131, 6534-6535, 9307-9309), 16 carpometacarpi (Av-2438, 2449, 2477, 2489, 2519, 2525, 2539, 2552, 2559, 2588, 3758, 9313-9317), a femur (Av-2990), five tibiotarsi (Av-3367, 3581, 6468, 6519, 6557), and five tarsometatarsi (Av-3367, 3581, 6468, 6519, 6557). Figured by Milne-Edwards 1869-1871, pl. 88, fig. 4-7 (tarsometatarsus), 8-9 (tibiotarsus), 10 (ulna), 11 (radius); and Milne-Edwards 1969-1871, pl. 89, fig. 1-2 (coracoid), 5 (humerus).]

Palaelodus [sic!] *minutus* Milne-Edwards, 1868: pl. 86 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus; MNHN Av-6511. Figured by Milne-Edwards 1869-1871, pl. 86, fig. 17-20; Cheneval 1983a, pl. 10, fig. 2; and Cheneval 1983c, pl. 1, fig. 2a-b.]

Grus problematica Milne-Edwards, 1868: pl. 76 [Holotype from Saint-Gérard-le-Puy: imperfect premaxilla; MNHN Av-8728. Figured by Milne-Edwards 1869-1871, pl. 76, fig. 5-7; and Cracraft 1973, fig. 40a,d.]

Probalearica problematica (Milne-Edwards): Lambrecht 1933: 519 [New combination.]

Grus miocenicus Grigorescu & Kessler, 1977: 102 [Holotype from Credița: distal end of a tarsometatarsus; LPUB 263. Figured by Grigorescu & Kessler 1977.]

Distribution: Middle Oligocene (MP 25) of Chaptuzat, France (Milne-Edwards 1867-1868, 1869-1871); late Oligocene (MP 25-30) of Perignat, France (Lydekker 1891a: 89); late Oligocene (MP 28) of Courmon, France (Milne-Edwards 1869-1871); late Oligocene (MP 29) of Pont-du-Château, France (Milne-Edwards 1869-1871); late Oligocene (MP 30) of Gannat, France (Milne-Edwards 1869-1871);

early Miocene (MN 1) of Weisenau-AS, Germany (Lydekker 1891a: 90, Lambrecht 1933: 340); early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1869-1871, Cheneval 1983c, Cheneval & Escuillié 1992), Wiesbaden-Hessler, Germany (Lambrecht 1933: 340), Wiesbaden 1, Germany (Lambrecht 1933: 342), Budenheim, Germany (Lambrecht 1933: 340, 342); early Miocene (MN 2) of Monsheim, Germany (Lambrecht 1933: 341), Büchelberg, Germany (Kuss 1960), and Ravolzhausen, Germany (Martini 1974); early Miocene (MN 4b) of Dolnice, Czechia (Švec 1981, Mlíkovský orig.);

middle Miocene (MN 5) of Františkovy Lázně, Czechia (Mlíkovský orig.); middle Miocene (MN 6) of Nördlinger Ries, Germany, incl. Steinberg and Hahnenberg (Lambrecht 1933: 339, Heizmann & Hesse 1995); middle Miocene (MN 7) of Lišov, Czechia (Mlíkovský orig. as *P.* cf. *ambiguus*); and middle Miocene (MN 8) of Credița, Romania (Grigorescu & Kessler 1977, Mlíkovský orig.).

The following record is invalid: early Miocene (MN 1) of Pyrimont-Challonges, France (Depéret & Douxami 1902 – see Cheneval 1983c: 184), and Oppenheim, Germany (Rothausen 1966 – see Olson & Feeduccia 1980a, Mlíkovský 1988, Cheneval & Escuillié 1992).

Remarks: *Palaelodus minutus* Milne-Edwards was synonymized with *Palaelodus gracilipes* Milne-Edwards by Cheneval (1983c: 184). The latter species was synonymized with *Palaelodus ambiguus* Milne-Edwards by Lambrecht (1933: 339), and Mlíkovský (orig.). *Grus problematica* Milne-Edwards was synonymized with *Palaelodus ambiguus* Milne-Edwards by Cheneval & Escuillié (1992: 211). The holotypical tarsometatarsus of *Grus miocenicus* Grigorescu & Kessler shows typical morphology of *Palaelodus* flamingos, and agrees in size

with *Palaelodus ambiguus* Milne-Edwards. I synonymize it here with the latter species (Mlíkovský orig.).

The alleged morphological differences between *Palaelodus gracilipes*, *P. ambiguus*, and *P. crassipes* pointed out by Cheneval (1983c) and Heizmann & Hesse (1995) are clearly size-dependent, and can be interpreted as intraspecific variability. *P. gracilipes* is based on smallest individuals of *P. ambiguus*, while *P. crassipes* on its largest individuals.

***Palaelodus goliath* Milne-Edwards**

Palaelodus [sic!] *goliath* Milne-Edwards, 1868: pl. 87 [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1983c: 186): right tarsometatarsus; MNHN Av-6475. Figured by Milne-Edwards 1867-1869, pl. 87, fig. 1-4; and Cheneval 1983c, pl. 1, fig. 5a-b. Paralectotypes from Saint-Gérard-le-Puy (based on the list in Cheneval 1983a: 139-140): two coracoids (Av-6460-6461), sternum (Av-6459), four femora (Av-3146, 6462-6464), nine tibiotarsi (Av-6465-6467, 6469-6474), and five tarsometatarsi (Av-3771, 6476-6477, 8894, 9669). Figured by Milne-Edwards 1867-1868, pl. 87, fig. 5-8 (femur), 13-14 (coracoid); and Milne-Edwards 1867-1868, pl. 88, fig. 1-3 (tibiotarsus).]

Megapalaelodus goliath (Milne-Edwards): Cheneval 1983c: 186 [New combination.]

Distribution: Late Oligocene (MP 25-30) of Perignat, France (Lydekker 1891a: 82); early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1869-1871, Cheneval 1983c), Wiesbaden-Hessler, Germany (Lambrecht 1933: 342), Wiesbaden 1, Germany (Lambrecht 1933: 342), and Budenheim, Germany (Lambrecht 1933: 342); early Miocene (MN 2) of Ravelzhausen, Germany (Martini 1974); early Miocene (MN 4b) of Dolnice, Czechia (Švec 1981, Mlíkovský orig.); middle Miocene (MN 6) of Mátraszőlös 2, Hungary (Gál & Kessler in Gál et al. 2000); and middle Miocene (MN 7) of Steinheim, Germany (Heizmann & Hesse 1995).

Genus *Phoenicopterus* Linnaeus

Phoenicopterus Linnaeus, 1758 [Modern genus.]

Gervaisia Harrison & Walker, 1976d: 308 [Type by original designation: *Phoenicopterus croizeti* Gervais, 1852. This is a junior homonym of *Gervaisia* Bonaparte, 1854: 6, *Gervaisia* Waga, 1858: 829, and *Gervaisia* Robineau-Desvoidy, 1863: 36.]

Harrisonavis Kašín, 1978: 146 [New name for *Gervaisia* Harrison & Walker, 1976d; hence its junior objective synonym.]

Remarks: *Gervaisia* Harrison & Walker and *Harrisonavis* Kašín were synonymized with *Phoenicopterus* Linnaeus by Olson & Feduccia (1980a: 46).

***Phoenicopterus croizeti* Milne-Edwards**

Phoenicopterus croizeti Gervais, 1852: 233 [Syntypes from Gergovie: skull and anterior portion of bill; formerly in coll. Croizet, present location unknown, casts in MNHN Av-6544 (skull; possibly lost – Cheneval 1983a: 146) and Av-6545 (bill). Figured by Gervais 1852, pl. 50, fig. 4 (skull) and 5 (bill). Miller (1963: 292) erroneously stated, that the species was based on a tibiotarsus.]

Gervaisia croizeti [sic!] (Gervais): Harrison & Walker 1976d: 308 [New combination; species name misspelled.]

Harrisonavis croizeti (Gervais): Kašín 1978: 146 [New combination.]

Distribution: Late Oligocene (MP 25) of Chaptuzat, France (Gervais 1848-1852, Lydekker 1891a: 78-79, Milne-Edwards 1867-1868, Cheneval 1984a), Sauvetat, France (Laizer 1828 sub unidentified eggs, Gervais 1848-1852: 234, Lydekker 1891a: 79); late Oligocene (MP 28) of Cournon, France (Gervais 1848-1852: 234, Milne-Edwards 1867-1868); late Oligocene (MP 29) of Pont-du-Château, France (Gervais 1848-1852: 234, eggs only); late Oligocene (MP 30) of Gergovie, France (Gervais 1848-1852, Cheneval 1984a) and Gannat, France (Milne-

Edwards 1867-1868, Cheneval 1984a); late Oligocene (MP 25-30) of Perignat, France (Milne-Edwards 1867-1868, Lydekker 1891a: 78-79), cf. Oligocene (MP 21-30) of Alliers, France (Milne-Edwards 1867-1868), and Aurillac, France (Gervais 1848-1852: 243);

early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1869-1871, Lydekker 1891a: 78-79, Lambrecht 1933: 344, Cheneval 1984a), and Budenheim, Germany (Lambrecht 1933: 345); early Miocene (MN 2) of Ravolzhausen, Germany (Martini 1974); early Miocene (MN 4b) of Dolnice, Czechia (Švec 1981, Mlíkovský orig.); middle Miocene (MN 5) of Františkovy Lázně, Czechia (Mlíkovský, orig.); and middle Miocene (MN 7) of Steinheim, Germany (Fraas 1870, Heizmann & Hesse 1995).

Remarks: After being separated at the generic level by Harrison & Walker (1976d), this species was returned to *Phoenicopterus* Linnaeus by Olson & Feduccia (1980a: 46).

***Phoenicopterus* sp.**

Distribution: Late Miocene (MN 12-13) of Pikermi, Greece (Mlíkovský, orig., material in MNHN).

Family Recurvirostridae Bonaparte

Recurvirostridae Bonaparte, 1831: 59 [Modern family.]

Remarks: Three species from the Paleogene of Europe were assigned to this family, but I had to relegate *Fluviatilavis antunesi* Harrison to the category of Aves incertae sedis, *Tenuicrus magnum* Harrison & Walker may represent an ancient flamingo (see p. 103), and also the taxonomic fate of *Recurvirostra sanctaneboulae* Mourer-Chauviré is uncertain (see below). There is no clear evidence, that the Recurvirostridae existed in Europe before the Holocene.

Genus Recurvirostra Linnaeus

Recurvirostra Linnaeus, 1758 [Modern genus.]

***Recurvirostra sanctaneboulae* Mourer-Chauviré**

Recurvirostra sanctaneboulae Mourer-Chauviré, 1978b: 221 [Holotype from Sainte-Néboule: left ulna; USTL SNB-705. Figured by Mourer-Chauviré 1978b, pl. 1, fig. 7-10.]

Distribution: Late Eocene (MP 18) of Sainte-Néboule, France (Mourer-Chauviré 1978).

Remarks: There is no evidence, that this species belongs to the family Recurvirostridae, or even to the genus *Recurvirostra* Linnaeus. Nevertheless, it seems to belong to one of the following closely related families: Phoenicopteridae, Recurvirostridae and Presbyornithidae. Characters distinguishing proximal end of ulna of these families are not known as yet (Ericson 1999: 248), so that *Recurvirostra sanctaneboulae* Mourer-Chauviré is in danger to become a nomen dubium until more material is found.

Family Anseranatidae Sclater

Anseranatidae Sclater, 1880 [Modern family.]

Anhimidae Stejneger, 1885 [Modern family.]

Romainvillinae Lambrecht, 1933: 351 [Type genus: *Romainvillia* Lebedinsky, 1927. Established as a subfamily of the Anatidae.]

Romainvilliinae Lambrecht: Brodkorb 1964: 208 [Spelling emended.]

Remarks: I synonymize here Anhimidae Stejneger and Romainvilliinae Lambrecht with the Anseranatidae Sclater (see below under *Romainvillia* Lambrecht for a discussion)

Genus Nettapterornis n. g.

Nettapterornis Mlíkovský, this paper [Type by original designation: *Anatalavis oxfordi* Olson,

1999.]

Diagnosis: An anhimid with a flat, duck-like rostrum, ending with an upwards oriented tip. For further details see Olson (1999a: 240-242).

Comparison: Differs from *Anatalavis* Olson & Parris in having brachial depression on the distal end of humerus located in the axis of shaft, external condyle oriented more ventrally (in *Anatalavis* it lays almost parallel with axis of the shaft, with the proximal tip facing the dorsal border of the shaft), internal condyle more protruding distally, and entepicondyle ending more proximally (in *Anatalavis* it almost forms a line with internal and external condyles). Other bones are known for *Anatalavis*.

Included species: Type species only.

Etymology: *Anatalavis* means "duck-winged bird" in Latin (Olson & Parris 1987: 11). *Nettapterornis* means the same in Greek, being formed from Greek words "νεττα" (duck), "πτερον" (wing), and "ορνις" (bird). The name is feminine in gender.

***Nettapterornis oxfordi* (Olson)**

Anatalavis oxfordi Olson, 1999a: 233 [Holotype from Walton-on-the-Naze: associated skeleton lacking hind limbs; BMNH A-5922. Figured by Olson 1999a, fig. 1-2 (skull), 3 (pterygoid), 4 (mandible), 5 (furcula), 61-f (coracoid), 6g (scapula), 7 (sternum and pelvis), 8a-c (humerus), and 9 (ulnare, carpometacarpus and associated phalanges).]

Nettapterornis oxfordi (Olson): Mlíkovský, this paper [New combination.]

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Olson 1999).

Genus *Headonornis* Harrison & Walker

Headonornis Harrison & Walker, 1976a: 335 [Type by original designation: *Agnopterus hantoniensis* Lydekker, 1891a.]

***Headonornis hantoniensis* (Lydekker)**

Agnopterus (?) *hantoniensis* Lydekker, 1891a: 96 [Holotype from Hordle: right coracoid; BMNH 30325. Figured by Lydekker 1891a, fig. 25, Harrison & Walker 1976a, pl. 3, fig. j-o.]

Headonornis hantoniensis (Lydekker): Harrison & Walker, 1976a: 335 [New combination.]

Presbyornis hantoniensis (Lydekker): Bocheński, 1997: 303 [New combination.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a, Harrison & Walker 1976a).

Remarks: Originally not included in a family. Harrison & Walker (1976a) included the species in the *Telmabatidae* (= *Presbyornithidae*). I transfer here this species on the basis of the general morphology of the holotype coracoid.

Genus *Romainvillia* Lebedinsky

Romainvillia Lebedinsky, 1927: 1 [Type by monotypy *Romainvillia stehlini* Lebedinsky, 1927.]

***Romainvillia stehlini* Lebedinsky**

Romainvillia Stehlini Lebedinsky, 1927: 1 [Syntypes from Romainville: furcula in slab (BaM 38), cranial end of left coracoid (BaM 38a), left carpometacarpus (BaM 53) proximal end of right carpometacarpus (BaM 55), and right tarsometatarsus (BaM 25.1); all perhaps from a single individual. Present location unknown: Boxes with respective labels were present in BaM, but bones were absent from them in 1996 (B. Engesser, in litt. 1996). Figured by Lebedinsky 1927, pl. 1, fig. 1-4 (left carpometacarpus), 5-7 (tarsometatarsus), 8-11 (coracoid), and 12 (furcula); and Howard 1964, pl. 1, fig. a-c (left carpometacarpus), d

(tarsometatarsus), e-h (coracoid).]

Distribution: Late Eocene (MP 20) of Romainville, France (Lebedinsky 1927, Brunet 1970).

Remarks: The syntypical carpometacarpus of *Romainvillia* Lebedinsky differs from the same element of the proper Anatidae and agrees with that of *Anseranas* Lesson in having (1) carpal trochlea not notched, (2) processus extensorius rather small, (3) distal end of metacarpal II distorted caudally, which casuses that (4) the facet for digit II is relatively large. In addition, coracoidal fenestra is present as in *Anseranas*, while it is absent in all proper Anatidae. Similarly as Verheyen (1956) and Olson & Feduccia (1980b), I consider *Anseranas* Linnaeus to be closer to the Anhimidae than to the proper Anatidae. Consequently, I transfer here modern anhimids and *Romainvillia* Lebedinsky to the family Anseranatidae, which has nomenclatural priority (see above under the Anseranatidae).

Family Anatidae Leach

Anatidae Leach, 1820 [Modern family.]

Genus *Paracygnopterus* Harrison & Walker

Paracygnopterus Harrison & Walker, 1979b: 35 [Type by original designation: *Paracygnopterus scotti* Harrison & Walker, 1979b.]

Paracygnopterus scotti Harrison & Walker

Paracygnopterus scotti Harrison & Walker, 1979b: 35 [Holotype from Yarmouth: cranial end of right coracoid; BMNH A-4407. Figured by Harrison & Walker 1979b, pl. 2, row A, fig. 1-5.]

Distribution: Early/middle Oligocene (MP 21-23) of Yarmouth, England (Harrison & Walker 1979b).

Remarks: The taxonomic position of this species requires restudy.

Genus *Mionetta* Livezey & Martin

Mionetta Livezey & Martin, 1988: 208 [Type by original designation: *Anas blanchardi* Milne-Edwards, 1863.]

Distribution: Early Miocene (MN 1-4) of France, Germany and Czechia, and middle Miocene (MN 6-8) of France, Germany and Romania.

Mionetta blanchardi Milne-Edwards

Anas Blanchardi Milne-Edwards, 1863: 160 (not 155 as stated by Cheneval 1983b: 88) [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1983b: 88): left tibiotarsus; MNHN Av-8231. Figured by Cheneval 1983a, pl. 14, fig. 1a-b, and Cheneval 1983b, pl. 1, fig. 1a-b. Not figured by Milne-Edwards (1867-1868, pl. 22, fig. 8-14), which shows side-reversed right tibiotarsus (contra Cheneval 1983b: 88). "Paralectotypes" from Saint-Gérard-le-Puy (based on the list in Cheneval 1983a: 157-158, but restricted to limb bones – see Milne-Edwards 1863: 160; all in MNHN): 388 humeri (Av-6966-7047, 7049-7082, 7084-7309, 7311-7317, 7319, 7320, 7322-7339, 7341-7358, 8587), 291 ulnae (Av-2154, 7359-7648), 182 radii (Av-7649-7663, 7665-7735, 7737-7831, 8618), 170 carpometacarpi (Av-7832-7932, 7934, 8029, 8031-8094, 8624, 8626, 8633), 93 femora (Av-8137-8166, 8168-8230), 160 tibiotarsi (Av-6451-6453, 8232-8388), and 156 tarsometatarsi (Av-6456, 6458, 8389-8542). Figured by Milne-Edwards 1867-1868, pl. 21 (reconstructed skeleton), pl. 22, fig. 1 (leg skeleton), 2-7 (tarsometatarsus), 8-14 (tibiotarsus), 17-23 (femur), pl. 23, fig. 1 (wing skeleton), pl. 24, fig. 7-9 (humerus), 10-11, 11a (ulna), 12-13 (radius), 14-17 (carpometacarpus), Howard 1964, pl. 8, fig. a-c (humerus), d-e (tarsometatarsus), f

(tibiotarsus), and Cheneval 1983a, pl. 15, fig. 4a-b (humerus, Av-6966), 5a-b (ulna, Av-7359), and pl. 16, fig. 2a-b (tarsometatarsus, Av-8389).

Cheneval (1983b: 88) erroneously included among the paralectotypes also the tarsometatarsus MNHN Av-6457, upon which Milne-Edwards based *Anas consobrina*, and the femur MNHN Av-8691, attributed by Milne-Edwards to *Ibis pagana*. Howard (1964: 291) erroneously stated, that type of *Anas blanchardi* Milne-Edwards is a "complete skeleton in Paris Museum", possibly referring to the reconstructed skeleton figured by Milne-Edwards 1867-1868, pl. 22, fig. 1.]

Anas consobrina Milne-Edwards, 1867b: 145 [Lectotype from Saint-Gérard-le-Puy (selected by Cheneval 1983b: 91): left tibiotarsus; MNHN Av-6454 = SG-9006. Figured by Cheneval 1983a, pl. 14, fig. 2a-b, and Cheneval 1983b, pl. 1, fig. 2a-b. Not figured by Milne-Edwards 1867-1868, pl. 25, fig. 5-7, which shows a side-reversed right tibiotarsus (contra Cheneval 1983b: 91). "Paralectotypes" from Saint-Gérard-le-Puy (based on the list in Cheneval 1983a: 162-164, restricted to limb bones, as in the case of *Anas blanchardi*; all in MNHN): two carpometacarpi (Av-1055, SG-10054), two femora (Av-6449-6450), right tibiotarsus (Av-6455), and left tarsometatarsus (Av-6457). Figured by Milne-Edwards 1867-1868, pl. 25, fig. 1 (leg skeleton), 2-4 (tarsometatarsus), 5-7 (tibiotarsus), 8-10 (femur), and 11-13 (carpometacarpus).]

Palaeortyx? phasianoides Milne-Edwards, 1869: 237 [Lectotype from Saint-Gérard-le-Puy (here selected): shaft of right humerus; MNHN Av-2896. Figured by Milne-Edwards 1869-1871, pl. 130, fig. 26-27. Paralectotype: cranial end of right scapula from Saint-Gérard-le-Puy; MNHN Av-2895. Figured by Milne-Edwards 1869-1871, pl. 130, fig. 22-25.]

Anas macroptera Milne-Edwards, 1871: 573 [Nomen nudum; no description or indication. Cheneval 1983b: 91 located left humerus from Saint-Gérard-le-Puy (MNHN SG-10053) upon which the species was based, because it was accompanied with Milne-Edwards's handwritten label bearing the name "*Anas macroptera*". *Anas macroptera* Milne-Edwards, 1871 remains nomen nudum, but can be listed in this place.]

Fuligula arvernensis Lydekker, 1891a: 122 [Holotype from Saint-Gérard-le-Puy: associated imperfect right humerus and partial shaft of an ulna in slab; BMNH A-159. Not figured.]

Aythya arvernensis (Lydekker): Brodkorb 1964: 228 [New combination.]

Taoperdix phasianoides (Milne-Edwards): Brodkorb 1964: 301 [New combination.]

Dendrochen blanchardi (Milne-Edwards): Cheneval 1983b: 88 [New combination.]

Dendrochen consobrina (Milne-Edwards): Cheneval 1983b: 91 [New combination.]

Mionetta blanchardi (Milne-Edwards): Livezey & Martin 1988: 208 [New combination.]

Mionetta consobrina (Milne-Edwards): Livezey & Martin 1988: 208 [New combination.]

Distribution: Early Miocene (MN 1) of Weisenau-AS, Germany (Milne-Edwards 1867-1868, Lydekker 1891a); early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1867-1868, Lydekker 1891a, Lambrecht 1933, Cheneval 1983b, Livezey & Martin 1988, Mlíkovský orig.); early Miocene (MN 2) of Ravalzhausen (Martini 1974); early Miocene (MN 4b) of Dolnice, Czechia (Švec 1981, Mlíkovský orig.); middle Miocene (MN 6) of Mátraszőlős 2, Hungary (Gál & Kessler in Gál et al. 2000); middle Miocene (MN 7) of Steinheim, Germany (Fraas 1870, but see Heizmann & Hesse 1995); and middle Miocene (MN 8) of Credința, Romania (Kessler 1992). The record from the early Pliocene (MN 15) of Malușteni-Berești, Romania (Kessler 1992) is doubtful because of its age. The following record is invalid: early Miocene (MN 3) of Skyřice, Czechia (Laube 1910 sub *Anas blanchardi* – reidentified as *Mionetta natator* by Mlíkovský 2000f).

Remarks: Cheneval (1983b), Mlíkovský (1983c) and Livezey & Martin (1988) recognized the dendrocygnine affinities of *blanchardi*. Cheneval (1983b) included the species in the genus *Dendrochen*, created by A.H. Miller (1944) for *Dendrochen robusta* A.H. Miller, 1944 from the early Miocene of South Dakota. Livezey & Martin (1988) separated the species at the generic level, which agrees with my observations. *Anas consobrina* Milne-Edwards is based on large specimens of *Anas blanchardi* (Mlíkovský orig., see also Livezey & Martin 1988).

Palaeortyx phasianoides Milne-Edwards was synonymized with *Mionetta blanchardi* (Milne-Edwards) by Mlíkovský (2000e: 93).

***Mionetta natator* (Milne-Edwards)**

Anas natator Milne-Edwards, 1867b: 148 [Lectotype from Saint-Gérard-le-Puy (selected by Storer 1956: 422): left ulna; MNHN Av-6428. Figured by Milne-Edwards 1867-1868, pl. 25, fig. 21-22; and Cheneval 1983a, pl. 14, fig. 4a-b. "Paralectotypes" from Saint-Gérard-le-Puy (with reference to the data in Cheneval 1983a: 165-166; all reportedly in MNHN): distal end of right humerus (perhaps one of the following: Av-6437-6438, SG-10064-10066), and right tibiotarsus lacking proximal end (perhaps Av-6622; certainly not Av-6436 listed by Cheneval 1983a: 165, 1983b: 93, and figured by Cheneval 1983a, pl. 14, fig. 3a-b, and Cheneval 1983b, pl. 1, fig. 3a-b). Figured by Milne-Edwards 1867-1868, pl. 25, fig. 14-18 (tibiotarsus), and 19-20 (humerus). All syntypes and other material studied by Cheneval (1983a,b) should be in MNHN, but may be lost, because I did not find them there in June 1999.]

Querquedula natator (Milne-Edwards): Brodkorb 1962: 157 [New combination.]

Dendrochen natator (Milne-Edwards): Cheneval 1983b: 93 [New combination.]

Mionetta natator (Milne-Edwards): Livezey & Martin 1988: 208 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1867-1868, Cheneval 1983b, Livezey & Martin 1988); early Miocene (MN 3) of Skyřice, Czechia (Laube 1910 sub *Anas blanchardi*, Mlíkovský 2000f); and early Miocene (MN 4b) of Dolnice, Czechia (Švec 1981, Mlíkovský orig.). The record from the late Miocene (MN ?9) of Chişinău, Moldova (Kessler 1992) is doubtful, because of its age.

Remarks: The taxonomic identity of both paralectotypes of *Mionetta natator* (Milne-Edwards) is doubtful, because both seem to entirely lack characteristic features of respective anatid bones (Mlíkovský orig.). The tibiotarsus perhaps belongs to *Colymboides minutus* Milne-Edwards (Storer 1956: 422).

***Mionetta robusta* (Milne-Edwards)**

Anas robusta Milne-Edwards, 1868: 155 [Holotype from Sansan: distal end of left humerus; MNHN SA-1234. Figured by Milne-Edwards 1867-1868, pl. 25, fig. 23-25, Cheneval 1987, pl. 1, fig. a-b.]

Anserobranta (?) *robusta* (Milne-Edwards): Cheneval 1987: 141 [New combination.]

Mionetta robusta (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Cheneval 1983b sub *Dendrochen consobrina*, Mlíkovský orig.); early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský orig.); middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1867-1868, Cheneval 1987, 2000), and Lierheim, Germany (Lydekker 1891a sub *Anas robusta*); and middle Miocene (MN 7-8) of Dechbetten, Germany (Ammon 1918 sub *Anas* cf. *robusta*).

Remarks: The holotypical humerus of *Anas robusta* Milne-Edwards is characterized by a very narrow space between the external condyle and the facet for the anterior articular ligament, which is typical for the Dendrocygnini (see Woolfenden 1961: 6). In absence of the evidence to the contrary I include here the species in the dendrocygnine genus *Mionetta* Livezey & Martin, which was common in Europe in the older part of the Miocene. Within the latter genus, *Mionetta robusta* (Milne-Edwards), new combination, is larger than both other known species, incl. *Mionetta natator* (Milne-Edwards) and *Mionetta blanchardi* (Milne-Edwards). Cheneval (1983b) correctly recognized from Saint-Gérard-le-Puy a *Mionetta* (his *Dendrochen*) species larger than *blanchardi*, but erroneously applied the name *Anas consobrina* Milne-Edwards to it.

Genus *Cygnavus* Lambrecht

Cygnavus Lambrecht, 1931a: 4 [Type by monotypy: *Cygnavus senckenbergi* Lambrecht 1931a.]

Distribution: Early Miocene (MN 2) of France and Germany. Extralimital record includes *Cygnavus formosus* Kuročkin, 1968 from the middle Oligocene of Kazakhstan (Kuročkin 1968, Mlíkovský & Švec 1986).

***Cygnavus senckenbergi* Lambrecht**

Cygnavus Senckenbergi Lambrecht, 1931a: 3 [Lectotype from Wiesbaden-Hessler (here selected): left femur with abraded ends; SMF Av-132, cast in USNM 215045. Figured by Lambrecht 1931a, pl. 1, fig. 3-4; Lambrecht 1933, fig. 128a-b; Howard 1964, pl. 2, fig. a-b. Paralectotypes from Wiesbaden-Hessler: distal end of left tibiotarsus (SMF Av-131, cast in USNM 215044), and a pedal phalanx (NFS Av-130, cast in USNM 215043). Figured by Lambrecht 1931a, pl. 2, fig. 9-10 (tibiotarsus) and 11-12 (phalanx), Lambrecht 1933, fig. 128c-d (tibiotarsus), and 128e-f (phalanx), and Howard 1964, pl. 2, fig. c-d (phalanx), and e-f (tibiotarsus).]

Cygnopterus alphonsi Cheneval, 1984a: 97 [Holotype from Saint-Gérand-le-Puy: right tarsometatarsus; FSL 91883. Figured by Cheneval 1984a, pl. 9, fig. 3a-c.]

Distribution: Early Miocene (MN 2a) of Saint-Gérand-le-Puy, France (Cheneval 1984a), and Wiesbaden-Hessler, Germany (Lambrecht 1933).

Remarks: The described bones of *Cygnopterus alphonsi* Cheneval agree in size and morphology with the corresponding bones of *Cygnavus senckenbergi* Lambrecht (Mlíkovský orig., after examining casts of all syntypes of the latter species in USNM), both species are contemporaneous, and were described from geographically close localities. Hence, I synonymize here the former with the latter species. This is a large representative of the Dendrocygnini (Mlíkovský orig.).

Genus *Cygnopterus* Lambrecht

Cygnopterus Lambrecht, 1931a: 3 [Type by monotypy: *Sula affinis* Beneden, 1883.]

Distribution: Middle Oligocene (MP 23-24) of Belgium. No extralimital record of this genus is known, because *Cygnopterus lambrechtii* Kuročkin, 1968 from the middle Oligocene of Kazakhstan was shown to be synonymous with the flamingo "*Agnopterus*" *turgaiensis* Tugarinov, 1940a, also described from the middle Oligocene of Kazakhstan (Mlíkovský & Švec 1986).

Remarks: Taxonomic position of this genus is uncertain.

***Cygnopterus affinis* (Beneden)**

Sula affinis Beneden, 1883: 133 [Syntypes from Rupelmonde (said to come from a single individual): cranial end of right coracoid, left humerus, proximal end of right ulna, right femur, and proximal end of left tibiotarsus; IRScNB Ht-Av-2a (femur), Av-2b (tibiotarsus) and Av-2c (humerus). Figured by Lambrecht 1931a, pl. 1, fig. 1-2 (femur), 6-7 (tibiotarsus), pl. 2, fig. 1-2 (humerus), 3-4 (ulna), 5-6 (coracoid); Lambrecht 1933, fig. 127a-b (humerus), 127c-d (ulna), 127e-f (tibiotarsus), 127g-h (femur), 127j-k (coracoid); and Howard 1964, pl. 3, fig. a-b (femur), e-f (humerus), g-h (tibiotarsus), j-k (coracoid). Lambrecht (1931a) described also a scapula, but Beneden (1883) did not mention it, so it could be included among the types only if all the bones were treated as coming from a single individual (which is well possible, but not certain), which would then be termed holotype.

Cygnopterus affinis (Beneden): Lambrecht 1931a: 3 [New combination.]

Palaeopapia hamsteadensis Harrison & Walker, 1979b: 34 [Holotype from Yarmouth: proximal end of right scapula; BMNH A-4412. Figured by Harrison & Walker 1979b, pl. 1, row E, fig. 5-8.]

Distribution: Early/middle Oligocene (MP 21-23) of Yarmouth, England (Harrison & Walker 1979b); and middle Oligocene (MP 23-24) of Rupelmonde, Belgium (Beneden 1883, Lambrecht 1931a, 1933, Cheneval 1984a).

Remarks: The holotypical scapula of *Palaeopapapia hamsteadiensis* Harrison & Walker closely resembles the same element of *Cygnopterus affinis* (Beneden). These species were described from stratigraphically and geographically close localities, so that I synonymize here the former with the latter species.

Genus *Cygnus* Bechstein

Cygnus Bechstein, 1802 [Modern genus]

Olor Wagler, 1832 [Modern genus.]

Palaeocygnus Stejneger, 1882: 180 [Type by original designation: *Cygnus falconeri* Parker, 1865.]

Palaeocygnus Oberholser, 1908. [Unsubstantiated emendation of *Palaeocygnus* Stejneger, 1882; hence an available name. This is a junior objective synonym of *Palaeocygnus* Stejneger, 1882.]

Cygnanser Kretzoi, 1957: 245 [Type by original designation: *Cygnus csakvarensis* Lambrecht, 1931a.]

Remarks: I formerly treated *Olor* Wagler as a separate genus (e.g. Mlíkovský & Švec 1986), but my further studies led me to change the opinion and to consider this genus as a subgenus of *Cygnus* Bechstein. The subgeneric allocation is not known for *Cygnus atavus* Fraas, while all the remaining species seem to belong to the subgenus *Olor* Wagler (but see under *Cygnus equitum* Bate).

Cygnus atavus (Fraas)

Anas atava Fraas, 1870: 275 [Holotype from Steinheim: proximal end of right femur; SMNS, uncatalogued. Figured by Fraas 1870, pl. 13, fig. 1a-c.]

Anas cygniformis Fraas, 1870: 276 [Lectotype from Steinheim (selected by Mlíkovský 1992b: 438): left coracoid; SMNS, uncatalogued. Figured by Fraas 1870, pl. 13, fig. 2a-b. Paralectotypes: proximal ends of two tarsometatarsi and three pedal phalanges; SMNS, uncatalogued. Figured by Fraas 1870, pl. 9, fig. 10-12 (phalanges).

Palaelodus steinheimensis Fraas, 1870: 285 [Holotype from Steinheim: distal end of left tibiotarsus; SMNS, uncatalogued. Figured by Fraas 1870, pl. 7, fig. 13.]

Anser atavus (Fraas): Lambrecht 1933: 369 [New combination.]

Anser cygniformis (Fraas): Lambrecht 1933: 369 [New combination.]

Cygnus atavus (Fraas): Mlíkovský 1992b: 439 [New combination.]

Distribution: Middle Miocene (MN 7) of Steinheim, Germany (Fraas 1870, Lambrecht 1933, Mlíkovský 1992b, Heizmann & Hesse 1995).

Remarks: *Palaelodus steinheimensis* Fraas and *Anas cygniformis* Fraas were synonymized with *Cygnus atavus* (Fraas) by Heizmann & Hesse (1995).

Cygnus (Olor) csakvarensis Lambrecht

Cygnus csákvárens Lambrecht, 1931a: 4 [Nomen nudum; no description or indication.]

Cygnus csákvárens Lambrecht, 1933: 283 [Lectotype from Csákvár (selected by Kretzoi 1957: 240): proximal end of left carpometacarpus; GIB, uncatalogued. Figured by Lambrecht 1933, fig. 128g-h; Kretzoi 1957, fig. 37-39; and Howard 1964, pl. 2, fig. g-h. Paralectotypes: distal end of a carpometacarpus (GIB, uncatalogued), and phalax I digiti majoris (GIB, uncatalogued). Figured by Lambrecht 1933, fig. 128j-k (phalanx); and Kretzoi 1957, fig. 40 (carpometacarpus), fig. 42-43 (phalanx).]

Cygnanser csákvárens (Lambrecht): Kretzoi 1957: 240 [New combination.]

Cygnanser csakvarens Lambrecht: Brodkorb 1964: 210 [Spelling emended.]

Olor csakvarens (Lambrecht): Mlíkovský 1992b: 437 [New combination.]

Distribution: Late Miocene (MN 10) of Csákvár, Hungary (Lambrecht 1933: 283, Kretzoi 1957, Mlíkovský 1992b).

***Cygnus verae* Boev**

Cygnus verae Boev, 2000a: 2 [Nomen nudum; no description or indication. This paper was published before 1 June, while the paper by Boev 2000b on 30 June.]

Cygnus verae Boev, 2000b: 186 [Holotype from Lozenec: proximal end of left humerus; MNHNS 1644. Figured by Boev 2000b, fig. 1a-c.]

Distribution: Early Pliocene (MN 14) of Lozenec, Bulgaria (Boev 1999a, 2000b).

***Cygnus (Olor) falconeri* Parker**

Cygnus falconeri Parker, 1865: 752 [Syntypes from Zebbug (listed according to Parker 1869: 119): two fragmentary skulls, three fragments of ribs, shaft of an ulna, 12 femora, 20 tibiotarsi, 20 tarsometatarsi, and three pedal phalanges; present location unknown, probably lost (Lydekker 1890, 1891a see also Northcote 1982b, 1992). Figured by Parker 1869, pl. 30, fig. 4-5 (ulna), 6-9 (femur), 10-11 (another femur), 12-15 (tibiotarsus), 16-19 (tarsometatarsus), and 20-22 (pedal phalanx). Parker (1865) did not specify the type material of *Cygnus falconeri*, stating that the species was based on "the head and more than half of the long bones [he received from the collectors Falconer and Spratt]", and referring to his forthcoming full paper [= Parker 1869]. Accordingly, the list of syntypes is based on Parker (1869).]

Cygnus melitensis Falconer, 1868: 300 [Types from Zebbug; not specified, but either same as for *Cygnus falconeri* Parker, 1865, or only those collected by Falconer, not those collected by Spratt. Parker (1865, 1869) did not specify which bones he received from which collector. Not figured in the original paper.]

Palaeocygnus falconeri (Parker): Stejneger 1882: 180 [New combination.]

Palaeocygnus falconeri (Parker): Oberholser 1908. [New combination.]

Cygnus (Olor) falconeri Parker: Livezey 1997a: 473 [New combination.]

Distribution: Late Pleistocene (MQ 2C) of Zebbug, Malta (Parker 1865, 1869, Lydekker 1890, 1891a), Ghar Dalam, Malta (Bate 1916), Mnajdra, Malta (Northcote 1982b), and Gnien, Malta (Northcote 1992).

Remarks: Northcote (1982b, 1983) assembled evidence that *falconeri* is related to *Olor*, not to *Cygnus* swans (see also Livezey 1997a: 473). *Cygnus melitensis* Falconer was correctly synonymized with *Cygnus falconeri* Parker by Lambrecht (1933: 385).

***Cygnus equitum* Bate**

Cygnus equitum Bate, 1916: 427 [Holotype from Ghar Dalam: right carpometacarpus; NMM 20 or 21, casts BMNH A-1613 or A-1614 or A-1615 (fide Northcote 1992: 287). Figured by Bate 1916, fig. 1a, 2a-b.]

Anser equitum (Bate): Brodkorb 1964: 213 [New combination.]

Cygnus (Olor) equitum Bate: Northcote 1988a. [New combination.]

Cygnus (Cygnus) equitum Bate: Livezey 1997: 473 [New combination.]

Distribution: Late Pleistocene (MQ 2C) of Ghar Dalam, Malta (Bate 1916), Ta'Kandja, Malta (Lambrecht 1933: 386), Ghaqba, Malta (Lambrecht 1933: 386), Benghisa, Malta (Lambrecht 1933: 386), Kalafrana, Malta (Lambrecht 1933: 386), Gnien, Malta (Northcote 1988, 1992), and Mnajdra, Malta (Northcote 1988, 1992).

Remarks: The taxonomic identity of this species is uncertain. Bate (1916) described it in the genus *Cygnus*, without specifying the subgenus. Brodkorb (1964) transferred it to the genus *Anser* Brisson, while Northcote (1988a, 1992) indicated, that it belongs to the swan subgenus *Olor* Wagler, and Livezey (1997a) placed it in the swan subgenus *Cygnus* Bechstein. The referred carpometacarpus (Lambrecht 1933, fig. 129d) resembles more the same element of *Alopochen* Stejneger than that of *Cygnus* Bechstein or *Anser* Brisson. The referred femur

(Lambrecht 1933, fig. 129e) is too small for an anatid with the referred humerus figured by Lambrecht (1933, fig. 129a) (Mlíkovský, orig.). Although all the bones referred to this species by Bate (1916), Lambrecht (1933) and Northcote (1988) seem to belong to the Anatidae, it is improbable that they belong to a single species.

***Cygnus (Olor) bewickii* Yarrell – Bewick's Swan**

Cygnus Bewickii Yarrell, 1830 [Modern species.]

Anser liskunae Kuročkin, 1976: 59 [Holotype from Dzabchan: distal end of left humerus; PIN 2614-105. Figured by Kuročkin 1976, fig. 6a,b,v, Kuročkin 1985, fig. 12a,b,v.]

Olor liskunae (Kuročkin): Mlíkovský & Švec, 1986: 263 [New combination.]

Anser subanser Jánossy, 1982: 22 [Nomen nudum; no description or indication.]

Anser subanser Jánossy, 1983: 252 [Lectotype from Přezletice (here selected): distal end of right radius; coll. Fejfar, probably destroyed – O. Fejfar, pers. communication, 1999. Not figured. Paralectotype: proximal end of right radius from Přezletice; coll. Fejfar, probably destroyed – O. Fejfar, pers. communication, 1999. Not figured. Jánossy (1983: 252) believed that both these elements belonged to a single bone and termed both of them "holotype". However, there is no support for such an assumption.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1982, 1983, Mlíkovský orig.), East Runton, England (Newton 1882b sub *Anser* sp., Harrison 1979d sub *C. bewickii*, Jánossy 1983: 252 sub *Anser subanser*), West Runton – Upper Freshwater Bed, England (Harrison 1979d: identification uncertain). Extralimital records are available from the middle Pliocene of Dzabchan and Chirgiz-Nur 2, Mongolia (Kuročkin 1976, 1985: 36, Mlíkovský & Švec 1986).

Remarks: Jánossy (1983) attributed to his *Anser subanser* all records he knew about, which were described as *Anser* and which originated from what he called "middle Pleistocene", even when he could not examine the bones. With one exception (retained in *Cygnus bewickii*), these records are listed under *Anser anser* (Linnaeus) in this paper (see below).

The holotypical humerus fragment of *Anser subanser* Jánossy (which I examined in the mid 1980s before it was lost) differs from the same element of *Olor* Wagler in the position of sulcus ligamentosus, shape of the facies articularis carpalis, and in the less pointed processus dorsalis (Mlíkovský orig.). *Anser liskunae* Kuročkin was identified as a small *Olor* swan, but left a valid species in belief that it is even smaller than the modern *Cygnus (Olor) bewickii* Yarrell (Mlíkovský & Švec 1986). The known bones of both of these species agree in morphology with the same elements of the modern *Cygnus bewickii* Yarrell, being comparable in size with the smallest specimens of the latter species. I synonymize here both *Anser subanser* Jánossy and *Anser liskunae* Kuročkin with the modern *Cygnus bewickii* Yarrell.

***Cygnus cygnus* (Linnaeus) – Whooper Swan**

Anas Cygnus Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *C. cf. cygnus*); early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999). The alleged record from Přezletice, Czechia (Jánossy 1983) refers to *Cygnus bewickii* (Mlíkovský orig.).

***Cygnus (Cygnus) olor* (Gmelin) – Mute Swan**

Anas Olor Gmelin, 1789 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Untermassfeld, Germany (Jánossy 1997 sub *C. cf. olor*); early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *C. aff. cygnus* [partim] – Mlíkovský orig.).

***Cygnus* sp.**

Distribution: Late Miocene (MN 9) of Chişinău, Moldova (Macarovici & Oescu 1942); late Miocene (MN 11) of Hrebnyky, Ukraine (Dubrovo & Kapelist 1979), late Miocene (MN 12) of Aljezar B, Spain (Cheneval & Adrover 1993), and early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965).

Genus *Anser* Brisson

Anser Brisson, 1760 [Modern genus]

***Anser thraceiensis* Burčák-Abramovič & Nikolov**

Anser thraceiensis Burčák-Abramovič & Nikolov, 1984: 29 [Syntypes from Trojanovo: left coracoid, distal end of left humerus, and proximal end of right humerus, perhaps from a single individual; NMHNS 2–2/5. Figured by Burčák-Abramovič & Nikolov 1984, pl. 2, fig. 1-1a (right humerus), 3-3a (left humerus), and 5-5a (coracoid).]

Distribution: Late Miocene or early Pliocene (MN 11-15) of Trojanovo, Bulgaria (Burčák-Abramovič & Nikolov 1984).

***Anser anser* (Linnaeus) – Greylag Goose**

Anas Anser Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Untermassfeld, Germany (Jánossy 1997 sub *A. aff. subanser*); early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), West Runton – Upper Freshwater Bed, England (Harrison 1979d), Voigtstedt, Germany (Jánossy 1965 sub *Anser* sp., Jánossy 1983: 252 sub *A. subanser*), and Somssichhégy 2, Hungary (Toula 1910, Jánossy 1983: 252, and 1986: 55 sub *A. subanser*); and middle Pleistocene (MQ 2A) of Várhegy – Fortuna street 25, Hungary (Jánossy 1979: 19 sub *A. aff. anser*, Jánossy 1983: 252, and 1986: 87 sub *A. subanser*), and Ambrona, Spain (Jánossy 1983: 252 sub *A. subanser*, reidentified by Sánchez Marco 1990b as *A. anser*).

***Anser* sp.**

Distribution: Early Miocene (MN 4a) of Dolnice, Czechia (Mlíkovský 1996e); middle Miocene (MN 6-7) of Minişu de Sus, Romania (Kessler & Codrea 1996); early Pliocene (MN 14) of Kamenskoe, Ukraine (Vojinstvens'kyj 1967); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b).

Genus *Branta* Scopoli

Branta Scopoli, 1769 [Modern genus.]

***Branta ruficollis* (Pallas) – Red-breasted Goose**

Anas ruficollis Pallas, 1769 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub *Branta* cf. *ruficollis*).

Remarks: This record is based on a fragmentary cranial end of a coracoid, and should be reevaluated.

Genus *Alopochen* Stejneger

Alopochen Stejneger, 1885 [Modern genus.]

Anserobranta Kuročkin & Ganea, 1972: 54 [Type by original designation: *Anserobranta tarabukini* Kuročkin & Ganea, 1972.]

Proanser Umans'ka, 1979a: 42 [Type by original designation: *Proanser major* Umans'ka,

1979a.]

Remarks: The holotypical carpometacarpi of both *Anserobranta tarabukini* Kuročkin & Ganea and *Proanser major* Umans'ka bear the characters typical for *Alopochen* Stejneger (see Woolfenden 1961). In absence of the contrary evidence I synonymize here the genera *Anserobranta* Kuročkin & Ganea and *Proanser* Umans'ka with *Alopochen* Stejneger.

***Alopochen tarabukini* (Kuročkin & Ganea)**

Anserobranta tarabukini Kuročkin & Ganea, 1972: 54 [Holotype from Chişinău: proximal end of left carpometacarpus; PIN 1713/1722. Figured by Kuročkin & Ganea 1972, text-fig. 4, pl. 1, fig. 5.]

Proanser major Umans'ka, 1979a: 42 [Holotype from Hrebennyky: proximal end of right carpometacarpus; IZAN 25–1682. Figured by Umans'ka 1979a, fig. 2a,b,v.]

Alopochen tarabukini (Kuročkin & Ganea): Mlíkovský, this paper [New combination.]

Distribution: Late Miocene (MN 9) of Chişinău, Moldova (Macarovici & Oescu 1941 sub *Cygnus* sp., Kuročkin & Ganea 1972, Kessler 1984b, 1992); and late Miocene (MN 11) of Hrebennyky, Ukraine (Umans'ka 1979a).

Remarks: The holotype of *Proanser major* Umans'ka agrees in size and morphology with the same element of *Anserobranta tarabukini* Kuročkin & Ganea. Both species were described from stratigraphically and geographically close localities, so that I synonymize here the former with the latter species.

Genus *Tadorna* Oken

Tadorna Oken, 1817 [Modern genus.]

Todarna Reichenbach, 1852: x [Type by monotypy: *Todarna wagneri* Reichenbach, 1852.]

Balcanas Boev, 1998d: 54 [Type by original designation: *Balcanas pliocaenica* Boev, 1998d.]

***Tadorna tadorna* (Linnaeus) – Common Shelduck**

Anas tadorna Linnaeus, 1758 [Modern species.]

Todarna wagneri Reichenbach, 1852: x [Syntypes from Monte Reale: coracoid, phalanx I digiti majoris, femur, and proximal end of tibiotarsus; present location unknown. Figured by Wagner 1832, pl. 2, fig. 49 (femur), 50 (tibiotarsus), 51a-b (coracoid), 52a-b (phalanx).]

Anser anatoides Depéret, 1892: 691 [Holotype from Perpignan: left tibiotarsus; FSL 92881. Figured by Depéret 1897, pl. 13, fig. 1-1a.]

Anser anatoides Depéret: Paris 1912: 290 [Spelling emended.]

Nettapus anatoides (Depéret): Brodkorb 1964: 227 [New combination.]

Anas submajor Jánossy, 1979: 11 [Holotype from Villány 3: left ulna; NMB Vt-83. Figured by Jánossy 1979, pl. 4, fig. 2.]

Balcanas pliocaenica Boev, 1998d: 54 [Holotype from Dorkovo: distal end of left humerus; NHMNS 484. Figured by Boev 1998d, fig. 2a-b.]

Distribution: Early Pliocene (MN 14) of Dorkovo, Bulgaria (Boev 1998d); early Pliocene (MN 15) of Perpignan, France (Depéret 1892, 1897); late Pliocene (MN 17) of Chilhac 2b and Chilhac 3, France (Boeuf & Mourer-Chauviré 1992 sub *T. cf. tadorna*), and Villány 3, Hungary (Jánossy 1979, Mlíkovský 1982a); and early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *Tadorna* sp., Mlíkovský orig.), and Stránská skála – Čapek's site 9a, Absolon's 2nd cave, Czechia (Jánossy 1972, Mlíkovský orig.).

Remarks: *Anas submajor* Jánossy was synonymized with the modern *Tadorna tadorna* (Linnaeus) by Mlíkovský (1982a). *Todarna wagneri* Reichenbach, 1852 was identified with *Tadorna tadorna* (Linnaeus) or *Tadorna ferruginea* (Pallas) by Mlíkovský (1995a; see also C.L. Nitzsch in Wagner 1833). *Anser anatoides* Depéret was compared by its author particularly with *Anser* Brisson and *Nettapus* Brandt, which was probably the reason why

Brodkorb (1964) and Howard (1964) included the species in the latter genus. The holotypical bones of *Anser anatoides* Depéret and *Balcanas pliocaenica* Boev are similar to the same elements of the modern *Tadorna tadorna* (Linnaeus), with which they were not compared formerly. Accordingly, I synonymize here the former two species with the latter one (Mlíkovský, orig.).

Genus *Dendronessa* Wagler

Dendronessa Wagler, 1832 [Modern genus.]

Dendronessa galericulata (Linnaeus)

Anas galericulata Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d sub *Aix galericulata*).

Dendronessa sp.

Distribution: Late Miocene (MN10) of Götzendorf, Austria (Mlíkovský 1992a).

Genus *Anas* Linnaeus

Anas Linnaeus, 1758 [Modern genus.]

Anas velox Milne-Edwards

Anas Meyerii Milne-Edwards, 1868: 129 [Holotype from Öhningen: crushed tarsometatarsus with associated phalanges in slab; BMNH 42805 (= Av-159). Figured by Meyer 1865, pl. 30, fig. 4, and Heer 1865, fig. 367.]

Anas velox Milne-Edwards, 1868: 150 [Lectotype from Sansan (selected by Cheneval 1987: 143, 144): right carpometacarpus; MNHN SA-1230. Figured by Milne-Edwards 1867-1868, pl. 26, fig. 17-18, and Cheneval 1987, pl. 1, fig. 2. Paralectotypes from Sansan: right coracoid (MNHN SA-1232), and proximal end of left ulna (MNHN SA-1231). Figured by Milne-Edwards 1867-1868, pl. 26, fig. 6-8 (coracoid) and 17-18 (ulna), and Cheneval 1987, pl. 1, fig. 3 (coracoid) and 4 (ulna). A tibiotarsus was listed among the syntypes of *Anas velox* Milne-Edwards by Howard (1964: 294) apparently in error.]

Nettion velox (Milne-Edwards): Brodkorb 1964: 224 [New combination.]

Aythya meyerii (Milne-Edwards): Brodkorb 1964: 228 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1867-1868, Cheneval 1987); middle Miocene (MN 7) of Steinheim, Germany (Fraas 1870), Attenfeld, Germany (Schlosser 1916: 33 sub ?*Anas velox*), Öhningen, Germany (Heer 1865, Meyer 1865 sub "Vogelfuss von Oeningen", Milne-Edwards 1867-1868, Mlíkovský 1992b); middle Miocene (MN 8) of Credița, Romania (Grigorescu & Kessler 1977 sub *Anas* cf. *velox*, Kessler 1992); and doubtfully late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1993 sub *Anas* aff. *velox*). The alleged record from the early Miocene (MN 4b) of Dolnice, Czechia (Švec 1981 sub *Nettion* cf. *velox*) was re-identified as *Mionetta natator* (Milne-Edwards) by Mlíkovský (orig.).

Remarks: *Anas meyerii* Milne-Edwards was a tiny duck, even smaller than the modern Holarctic *Anas crecca* Linnaeus. It agrees in size with the better documented *Anas velox* Milne-Edwards, described from the almost contemporaneous deposits of France. I regard these two nominal taxa as conspecific, selecting for them – as the first reviser – the name *Anas velox* Milne-Edwards, 1868. The allocation of this species to the genus *Anas* Linnaeus is uncertain.

Anas sansaniensis Milne-Edwards

Anas sansaniensis Milne-Edwards, 1868: 153 [Lectotype from Sansan (selected by Cheneval

1987: 145); distal end of left tibiotarsus; MNHN SA-1223. Figured by Milne-Edwards 1867-1868, pl. 25, fig. 26-30, and Cheneval 1987, pl. 1, fig. 5a-b. Paralectotype from Sansan: distal end of right humerus; MNHN 1233. Figured by Milne-Edwards 1867-1868, pl. 26, fig. 19-22, and Cheneval 1987, pl. 1, fig. 6a-b.]

Dendrocygna sansaniensis (Milne-Edwards): Mlíkovský 1988: 57 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1867-1868, Cheneval 1987); and middle Miocene (MN 7) of Grive-Saint-Alban, France (Cheneval 1987).

Remarks: Mlíkovský (1988) transferred *Anas sansaniensis* Milne-Edwards in the genus *Dendrocygna* Swainson on the basis of a referred skull from Sansan, but Cheneval (1987) restudied the lectotype and retained the species in *Anas* Linnaeus. I follow here Cheneval's decision, although I am not convinced, that the bird belongs to the genus *Anas* Linnaeus.

***Anas penelope* Linnaeus – Eurasian Wigeon**

Anas Penelope Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1) of Betfia – Aven, Romania (Kessler 1978); and early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), West Runton – Upper Freshwater Bed, England (Harrison 1979d), Voigtstedt, Germany (Jánossy 1965 sub *A. cf. penelope*), Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *A. cf. penelope*), and Nagyharsányhegy 1-4, Hungary (Jánossy 1979).

***Anas strepera* Linnaeus – Gadwall**

Anas strepera Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1) of Betfia – Aven, Romania (Kessler 1978); early Pleistocene (MQ 1a) of Huéscar 1, Spain (Sánchez Marco 1989), Beremend 17, Hungary (Jánossy 1992 sub *Anas cf. strepera*), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *A. aff. strepera*); and early Pleistocene (MQ 1b) of Betfia 7, Romania (Kessler 1975), Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *A. cf. strepera*), and Stránská skála – Musil's talus cone, layers 4, 4a, 5, 6, 8, 13, 14a, 15e, and 16, Czechia (Mlíkovský 1995).

***Anas crecca* Linnaeus – Common Teal**

Anas crecca Linnaeus, 1758 [Modern species]

Anas crecca percrecca Jánossy, 1992: 17 [Holotype from Beremend 16: right ulna; NMB V-91.150. Figured by Jánossy 1992, pl. 4, fig. 8.]

Distribution: Late Miocene (MN 9-10) of Sokolov, Ukraine (Vojinstvens'kyj 1967 sub *A. cf. crecca*); late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b and 1990b sub *Anas cf. querquedula*, Jánossy 1992 sub *Anas crecca percrecca*); early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992 sub *A. crecca percrecca*), and Betfia 2, Romania (Jánossy 1979 sub *A. aff. crecca*); and early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), West Runton – Upper Freshwater Bed, England (Newton 1887 sub *Anas?*, Harrison 1979d), and Přezletice, Czechia (Jánossy 1983 sub *Aythya cf. nyroca* and *Oxyura aff. leucocephala*, Mlíkovský orig.).

***Anas platyrhynchos* Linnaeus – Mallard**

Anas platyrhynchos Linnaeus, 1758 [Modern species.]

Fuligula aretina Portis, 1887: 194 [Lectotype from Montecarlo (here selected): right tarsometatarsus; IGF 930. Figured by Portis 1889, pl. 1, fig. 1-2. Paralectotypes from Montecarlo: axis (IGF 925), 5th (?) cervical vertebra (IGF 929), furcula (IGF 934), shaft of left ulna (IGF 933), proximal end of left carpometacarpus (IGF 935), distal end of left femur (IGF 931), shaft of left femur (IGF 936), and right tarsometatarsus (IGF 932). Figured by Portis 1889, pl. 1, fig. 3 (femur), 4-5 (vertebrae), 6-7 (furcula), 8 (ulna), and 9-10 (carpo-

metacarpus).]

Fuligula sepulta Portis, 1887: 194 [Lectotype from Montecarlo (here selected): left coracoid (IGF 924). Figured by Portis 1889, pl. 1, fig. 14-16. Paralectotypes from Montecarlo: atlas (IGF 922), furcula (IGF 919), proximal end of right rib (IGF 927), left ulna (IGF 920), right ulna (IGF 926), left radius (IGF 921), left carpometacarpus (IGF 928), and right carpometacarpus (IGF 923). Figured by Portis 1889, pl. 1, fig. 11 (vertebra), 12-13 (furcula), 17-19 (ulna), 20-21 (radius), and 22-23 (carpometacarpus).]

Aythya aretina (Portis): Brodkorb 1964: 228 [New combination.]

Aythya sepulta (Portis): Brodkorb 1964: 228 [New combination.]

Anas aretina (Portis): Cheneval 1987: 150 [New combination.]

Anas sepulta (Portis): Cheneval 1987: 150 [New combination.]

Distribution: Late Pliocene or early Pleistocene (MN 16 – MQ 1) of Montecarlo, Italy (Portis 1887 and 1889 sub *F. aretina* and *F. sepulta*), Strette, Italy (Portis 1889 sub *F. aretina*); and early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), East Runton, England (Harrison 1979d, see also Newton 1882b sub *A. boschas*), West Runton – Upper Freshwater Bed, England (Harrison 1979d), Voigtstedt, Germany (Jánossy 1965 sub *A. cf. platyrhynchos*), Přezletice, Czechia (Jánossy 1983 sub *Anas* sp., Mlíkovský orig.), and Stránská skála – Čapek's sites 5a, 5b, and 9a, Czechia (Jánossy 1972 sub *A. cf. platyrhynchos*). Both Pliocene and Pleistocene strata occur at Montecarlo and exact place of excavation of syntypes of *Fuligula aretina* Portis is unknown (see Delle Cave 1996: 677).

Remarks: *Fuligula aretina* Portis was based mainly on forelimb elements, while *Fuligula sepulta* Portis on hindlimb elements. Portis (1887, 1889) believed that *Fuligula sepulta* is smaller than *Fuligula aretina*, but I found in his data no support for this statement. Cheneval (1987) transferred *Fuligula aretina* Portis and *Fuligula sepulta* Portis to the genus *Anas* Linnaeus, without discussing their taxonomic status within the latter genus. The syntypes of both these alleged species agree in size with the same elements of the modern *Anas platyrhynchos* Linnaeus, with which I synonymize here both of them.

***Anas acuta* Linnaeus – Northern Pintail**

Anas acuta Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992 sub *Anas cf. acuta*), and Betfia 2, Romania (Jánossy 1979 sub *A. aff. acuta*); early Pleistocene (MQ 1a) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *A. cf. acuta*), and Nagyarsánygehy 1-4, Hungary (Jánossy 1979 sub *A. aff. acuta*).

***Anas querquedula* Linnaeus – Garganey**

Anas Querquedula Linnaeus, 1758 [Modern species.]

Distribution: Late Miocene (MN 11-13) of Čebotarevka, Ukraine (Vojinstvens'kyj 1967); early Pleistocene (MQ 1) of Betfia – Aven, Romania (Kessler 1978); early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *A. aff. querquedula*); early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999 sub *A. querquedula*), Stránská skála – Čapek's sites 1-3, 4c, 5a, 5b, 9a, Czechia (Jánossy 1972), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972), Stránská skála – Musil's talus cone, layers 4, 4a, 5, 6, 8, 9, 10, 10a, 11, 12, 13, 14a, 15a, 15c, 15e, and 16, Czechia (Mlíkovský 1995b), and Nagyarsánygehy 1-4, Hungary (Jánossy 1979 sub *A. aff. querquedula*).

***Anas clypeata* Linnaeus – Northern Shoveler**

Anas clypeata Linnaeus, 1758 [Modern species.]

Distribution: Late Miocene (MN 11-13) of Čebotarevka, Ukraine (Vojinstvens'kyj 1967 sub *A. cf. clypeata*); early Pleistocene (MQ 1) of Betfia – Aven, Romania (Kessler 1978); early

Pleistocene (MQ 1a) of Huéscar 1, Spain (Sánchez Marco 1989), Beremend 16, Hungary (Jánossy 1992 sub cf. *Spatula clypeata*), Beremend 17, Hungary (Jánossy 1992 sub cf. *Spatula clypeata*), and Betfia 2, Romania (Čapek 1917); early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, and Absolon's 2nd cave, Czechia (Jánossy 1972 sub cf. *Spatula clypeata*), Stránská skála – Musil's talus cone, layers 4a, 5, 6, 7b, 8, 9, 10, 11, 12, 13, 14, 15c, 15e, 16 (Mlíkovský 1995), and Nagyharsányhegy 1-4, Hungary (Jánossy 1979 sub *A. aff. clypeata*). The record from the early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Newton 1887, 1891 sub *Spatula clypeata*) is invalid (Harrison 1979d).

Anas sp.

Distribution: Late Miocene (MN 9) of Chişinău, Moldova (Macarovici & Oescu 1942, Kessler 1984); late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 2000a); and early Pleistocene (MQ 1b) of Sima del Elefante – unknown layer, Spain (Rosas et al. 2001).

Genus *Aythya* Boie

Aythya Boie, 1822 [Modern genus.]

***Aythya chauvirae* Cheneval**

Aythya chauvirae Cheneval, 1987: 148 [Holotype from Sansan: left femur; MNHN SA-1457.

Figured by Cheneval 1987, pl. 1, fig. 7a-b.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Cheneval 1987, 2000); and middle Miocene (MN 8) of Credința, Romania (Grigorescu & Kessler 1977 sub *Aythya* sp., Cheneval 1987, Kessler 1992).

***Aythya ferina* (Linnaeus) – Common Pochard**

Anas ferina Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Huéscar 1, Spain (Sánchez Marco 1989); and early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Lydekker 1891a sub *Fuligula ferina*, Harrison 1979d). An invalid record is from the early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983, reidentified by Mlíkovský, orig., as *Bucephala clangula*).

***Aythya nyroca* (Güldenstädt) – Ferruginous Duck**

Anas nyroca Güldenstädt, 1769 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Huéscar 1, Spain (Sánchez Marco 1989), Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *A. cf. nyroca*); and early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub *A. cf. nyroca*), Přezletice, Czechia (Jánossy 1983, Mlíkovský orig.), and Stránská skála – Čapek's sites 5a, and 9a, Czechia (Jánossy 1972 sub cf. *A. nyroca*).

***Aythya fuligula* (Linnaeus) – Tufted Duck**

Anas fuligula Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), West Runton – Upper Freshwater Bed, England (Harrison 1979d), Voigtstedt, Germany (Jánossy 1965 sub *A. cf. fuligula*), and Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *A. cf. fuligula*). An invalid record is from the early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983, reidentified by Mlíkovský, orig., as *Aythya nyroca* and *Bucephala clangula*).

Genus *Netta* Kaup

Netta Kaup, 1829 [Modern genus.]

***Netta rufina* (Pallas) – Red-crested Pochard**

Anas rufina Pallas, 1773 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Huéscar 1, Spain (Sánchez Marco 1989); and early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d).

Genus *Somateria* Leach

Somateria Leach, 1819 [Modern genus.]

***Somateria mollissima* (Linnaeus) – Common Eider**

Anas mollissima Linnaeus, 1758 [Modern species.]

Somateria gravipes Harrison, 1979: 280 [Holotype from West Runton – Upper Freshwater Bed: left tarsometatarsus; BMNH A-3396. Not figured.]

Distribution: Early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979 sub *S. gravipes*, Mlíkovský 1982c).

Remarks: Mlíkovský (1982c) believed, that the holotypical tarsometatarsus of *Somateria gravipes* Harrison is a pathological specimen, because it differed from the same element of *Somateria mollissima* (Linnaeus) in one dimension. Subsequent research showed, that Harrison (1979) understood this dimension in a different way than me, and that the holotype falls entirely in the size and shape range of the modern *Somateria mollissima* (Mlíkovský orig.).

Genus *Clangula* Leach

Clangula Leach, 1819 [Modern genus.]

***Clangula* sp.**

Distribution: Middle Miocene (MN 6-8) of Mátraszőlös 1, Hungary (Gál & Kessler in Gál et al. 1999).

Remarks: This is a very improbable record, because of its age (Mlíkovský orig.).

Genus *Melanitta* Boie

Melanitta Boie, 1822 [Modern genus.]

***Melanitta nigra* (Linnaeus) – Common Scoter**

Anas nigra Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Lydekker 1891a sub *Fuligula ferina*, Harrison 1979d).

Genus *Bucephala* Baird

Bucephala Baird, 1858 [Modern genus.]

***Bucephala cereti* Boeuf & Mourer-Chauviré**

Bucephala cereti Boeuf & Mourer-Chauviré, 1992: 1093 [Holotype from Chilhac 3: left humerus; MPC CH-3.1971.265. Figured by Boeuf & Mourer-Chauviré 1992, fig. 5-6.]

Distribution: Late Pliocene (MN 17) of Chilhac 3, France (Boeuf & Mourer-Chauviré 1992).

***Bucephala clangula* (Linnaeus) – Common Goldeneye**

Anas clangula Linnaeus, 1758 [Modern species.]

Bucephala angustipes Jánossy, 1965: 345 [Holotype from Voigtstedt: right tarsometatarsus; IQW 2795. Figured by Jánossy 1965, text-fig. 6, pl. 6, fig. 19.]

Distribution: Late Pliocene (MN 16) of Hajnáčka, Slovakia (P. Švec in Fejfar & Heinrich 1985 sub *Mergus* sp., Mlíkovský orig. – tentative identification); late Pliocene (MN 18) of S'Onix, Mallorca (Sondaar et al. 1995 sub *B. cf. clangula*); early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d), Voigtstedt, Germany (Jánossy 1965 sub *Bucephala angustipes*, Mlíkovský 1982c), Přezletice, Czechia (Jánossy 1983 sub *B. cf. angustipes*, *Aythya cf. fuligula*, and *Aythya cf. nyroca*, Mlíkovský orig.), Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *Bucephala angustipes*, Mlíkovský, orig.), and Stránská skála – Musil's talus cone, layers 11, 13, 15c and 15e, Czechia (Mlíkovský 1995b).

Remarks: *Bucephala angustipes* Jánossy was synonymized with *Bucephala clangula* Linnaeus by Mlíkovský (1982c). Early Pleistocene populations of *Bucephala clangula* Linnaeus tend to be smaller than modern ones (Mlíkovský 1995). Pliocene records should be compared with *Bucephala cereti* Boeuf & Mourer-Chauviré.

Genus *Mergellus* Selby

Mergellus Selby, 1840 [Modern genus.]

Mergellus albellus (Linnaeus) – Smew

Mergus Albellus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d).

Genus *Mergus* Linnaeus

Mergus Linnaeus, 1758 [Modern genus.]

Mergus connectens Jánossy

Mergus connectens Jánossy, 1972: 38 [Holotype from Stránská skála – Absolon's 2nd cave: right coracoid; MMB Abs-88. Figured by Jánossy 1972, pl. 1, fig. 5-6.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972, Mlíkovský orig.), Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b), and Betfia 5, Romania (Jánossy 1972: 42).

Mergus serrator Linnaeus – Red-breasted Merganser

Mergus Serrator Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d), Stránská skála – Musil's talus cone, layers 13 and 15e, Czechia (Mlíkovský 1995b).

Mergus merganser Linnaeus – Goosander

Mergus Merganser Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 13 (Mlíkovský 1995b).

Mergus sp.

Distribution: Middle Miocene (MN 6-8) of Mátraszőlös 1, Hungary (Gál & Kessler in Gál et al. 1999).

Remarks: This is an improbable record, because of its age (Mlíkovský orig.).

Genus *Oxyura* Bonaparte

Oxyura Bonaparte, 1828 [Modern genus.]

***Oxyura doksana* n. sp.**

Oxyura doksana Mlíkovský, this paper [Holotype from Dolnice: cranial end of left coracoid; DPFNSP 4828. Figured by Švec & Mlíkovský 1986, pl. 1, fig. a-c.]

Diagnosis: Coracoid of *Oxyura doksana* differs from the coracoids of other *Oxyura* species in having: (1) sulcus musculi procoracoidei deeper, (2) medioventral part of facies articularis less prominent over medioventral margin of canalis triosseus, and (3) in being generally larger and more robust.

Material: Holotype only. For measurements and detailed comparisons see Švec & Mlíkovský 1986.

Etymology: Lacustrine, latinized from *doksa*, Celtic word for a lake, the habitat of *Oxyura doksana*. Both Celts and *Oxyura doksana* were past inhabitants of Czechia, though in different times.

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Švec & Mlíkovský 1986, Mlíkovský orig.).

***Oxyura leucocephala* (Scopoli) – White-headed Duck**

Anas leucocephala Scopoli, 1769 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983, Mlíkovský orig.).

Genus incertae sedis***Anas albae* Jánossy**

Anas albae Jánossy, 1979: 16 [Holotype from Polgárdi: left carpometacarpus; NMB Vt-84. Not figured.]

Distribution: Late Miocene (MN 13) of Polgárdi, Hungary (V. Čapek in Lambrecht 1912a,b sub *Mergus* sp., Jánossy 1979).

***Anas eppelsheimensis* Lambrecht**

Anas eppelsheimensis Lambrecht, 1933: 362 [Holotype from Eppelsheim: fragmentary cranial end of right coracoid; present location unknown. Figured by Lambrecht 1933, fig. 124.]

Distribution: Late Miocene (MN 9) of Eppelsheim, Germany (Lambrecht 1933).

Remarks: The holotypical fragment probably cannot be identified within the family Anatidae (Mlíkovský 1992b).

***Anas isarensis* Lambrecht**

Anas isarensis Lambrecht, 1933: 361 [Holotype: proximal end of right scapula from Aumeister; formerly in BSP 1926-v-12b, but apparently destroyed in the World War II. Figured in Lambrecht 1933, fig. 123a.]

Distribution: Late Miocene (MN 9) of Aumeister, Germany (Lambrecht 1933).

Remarks: Mlíkovský (1992b) transferred this species to the tribe Aythyini, where it was considered indeterminate at the generic level. Unfortunately, in a subsequent test with duck scapulae I tended to misidentify *Anas* scapulae for those of *Aythya*. Taking into account that my re-identification of *Anas isarensis* was based on the restudy of an illustration, I refer here the species to Anatidae incertae sedis. It cannot be regarded as a record of the Aythyini.

***Anser brumeli* Milne-Edwards**

Anser Brumeli Milne-Edwards, 1871: 585 [Holotype from Orleanais: a tarsometatarsus; present location unknown. Not figured.]

Distribution: Middle Miocene (MN 4) of Orleanais, France (Milne-Edwards 1871).

Remarks: This is a valid name, although its description ("un Oie un peu plus petit que la barnache") is highly superficial. Lambrecht (1933: 368) and Brodkorb (1978: 222) erroneously listed the species as a nomen nudum (but see Lambrecht 1921: 12, Howard 1964: 266).

***Anas oeningensis* Meyer**

Anas oeningensis Meyer, 1865: 126 [Holotype from Öhningen: incomplete skeleton in slab; BMNH 42804. Figured by Meyer 1865, pl. 30, fig. 2.]

Anser oeningensis (Meyer): Milne-Edwards 1867b: 127 [New combination.]

Distribution: Middle Miocene (MN 7) of Öhningen, Germany (Meyer 1865, Milne-Edwards 1867b: 127).

Remarks: Taxonomic position of this species is in need of restudy. Lambrecht (1921: 15) listed this species in the synonymy of *Anas meyerii* Milne-Edwards, but the two species markedly differ in size (Mlíkovský orig.). The relation between the lengths of humerus and ulna in *Anas oeningensis* Meyer seems to differ from that of *Anser* Brisson, and to agree with that of the Cairinini (Mlíkovský, orig.). The taxonomic evaluation of this observation requires further study.

***Anas risgoviensis* Ammon**

Anas Risgoviensis Ammon, 1918: 41 [Syntypes from Lierheim: imperfect left coracoid (BMNH 48165) and sternal end of right coracoid (BMNH 48165a). Not figured.]

Distribution: Middle Miocene (MN 6) of Lierheim, Germany (Ammon 1918: 41), and Wallerstein, Germany (Ammon 1918: 7-8, feather imprints).

Remarks: There is no evidence that this species belonged to the genus *Anas* Linnaeus. Lydekker (1891a: 117) mentioned, that the syntypes are "slightly larger than the corresponding bones of *Anas blanchardi*".

***Anser scaldii* Lambrecht**

Anser Scaldii Beneden, 1872: 288 [Nomen nudum; no description or indication.]

Anser Scaldii Beneden, 1873a: 372 [Nomen nudum; no description or indication]

Anser scaldii "Beneden" Lambrecht, 1933: 368 [Holotype from Antwerp: a humerus; IRScNB, uncatalogued. Not figured.]

Distribution: Middle Miocene (MN 7-8) of Antwerp, Belgium (Beneden 1872, 1873a, Shufeldt 1909: 336, Lambrecht 1933: 368).

Remarks: Lambrecht (1933: 368) erroneously stated that the fossil was described by Beneden (1872) as *Anas Scaldii*, and „transferred“ it to the genus *Anser*. This incorrect information was repeated by Gaillard (1939: 78), Brodkorb (1964: 212) and Howard (1964: 267). *Anser Scaldii* is a pure nomen nudum in Beneden (1872, 1873a), but Lambrecht (1933: 368) restudied the holotype and presented a very brief description, thus validating the name.

Order Charadriiformes Huxley

Charadriomorphae Huxley, 1867 [Modern order.]

Family Jacanidae Chenu & des Murs

Jacanidae Chenu & des Murs, 1854 [Modern family.]

Genus *Nupharanassa* Rasmussen et al.

Nupharanassa Rasmussen, Olson & Simons, 1987: 7 [Type by original designation: *Nupharanassa bulotorum* Rasmussen et al., 1987.]

Distribution: Early Miocene (MN 4) of Czechia. Extralimital record is limited to the early Oligocene (MP ?) of Fayum in Egypt (Rasmussen et al. 1987), from where two species were described: *Nupharanassa bulotorum* Rasmussen et al., 1987, and *Nupharanassa tolutaria* Rasmussen et al., 1987.

Nupharanassa bohémica Mlíkovský

Nupharanassa bohémica Mlíkovský 1999a: 121 [Holotype from Dolnice: distal end of right tarsometatarsus; coll. Fejfar, uncatalogued. Figured by Mlíkovský 1999a, unnumbered fig.; and Mourer-Chauviré 1999a, fig. 1.]

Geranopterus bohemicus (Mlíkovský): Mourer-Chauviré 1999a: 151 [New combination.]

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský 1999a, Mourer-Chauviré 1999a).

Remarks: Mourer-Chauviré (1999a) transferred *Nupharanassa bohémica* Mlíkovský to the Coraciidae. She listed characters in which tarsometatarsus of the Jacanidae differ from that of the Coraciidae. Contrary to her opinion (she derived from the figures), the holotypical tarsometatarsus of *Nupharanassa bohémica* Mlíkovský agrees with the same element of the Jacanidae, and differs from that of the Coraciidae in all characters she listed. *Nupharanassa bohémica* Mlíkovský is certainly not a coraciid bird, and no points were presented for its exclusion from the Jacanidae.

Family Rostratulidae Mathews

Rostratulidae Mathews, 1914 [Modern family.]

Genus *Rostratula* Vieillot

Rostratula Vieillot, 1816 [Modern genus.]

Rostratula pulia Mlíkovský

Rostratula pulia Mlíkovský, 1999b: 99 [Holotype from Dolnice: distal part of right tarsometatarsus; DP FNSP 4819. Figured by Švec 1983, pl. 2, fig. a-c.]

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský 1999b).

Remarks: The holotypical tarsometatarsus of *Rostratula pulia* Mlíkovský was originally described by Švec (1983) as a paratype of *Microrallus fejfari* Švec.

Family Scolopacidae Vigors

Scolopacidae Vigors, 1825 [Modern family.]

Remarks: This is an osteologically rather uniform family, which opens considerable space for identification errors. Most of the Tertiary species were described inadequately, none were

properly revised, and the taxonomic status of all of the Tertiary taxa should be reevaluated.

***Calidris* Merrem**

Calidris Merrem, 1804 [Modern genus.]

***Calidris* sp.**

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský orig.); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *Calidris* sp., aff. *alpina* Linnaeus), and Stránská skála – Musil's talus cone, layer 7, Czechia (Mlíkovský 1995b).

Genus *Philomachus* Merrem

Philomachus Merrem, 1804 [Modern genus.]

***Philomachus pugnax* (Linnaeus) – Ruff**

Tringa Pugnax Linnaeus, 1758 [Modern species.]

Philomachus pugnax rhyphaeicus Potapova, 1990: 142 [Holotype from Medvěž'ja cave – layer 3: right humerus; ZIN RO-4003. Figured by Potapova 1990, pl. 3, fig. 6.]

Distribution: Early Pleistocene (MQ 1a) of Tarchankut, Ukraine (Vojinstvens'kyj 1967 sub *P.* cf. *pugnax*); early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub aff. *P. pugnax*); Stránská skála – Musil's talus cone, layers 4, 5, 6, 8, and 11, Czechia (Mlíkovský 1995b); and late Pleistocene (MQ 2D) of Medvěž'ja cave, Russia (Potapova 1990 sub *P. p. rhyphaeicus*).

Genus *Lymnocyptes* Kaup

Lymnocyptes Kaup, 1829 [Modern genus.]

***Lymnocyptes minimus* (Brünnich) – Jack Snipe**

Scolopax minima Brünnich, 1764 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layers 7, 14, and 15e, Czechia (Mlíkovský 1995b).

Genus *Gallinago* Brisson

Gallinago Brisson, 1760 [Modern genus.]

***Gallinago veterior* Jánossy**

Gallinago veterior Jánossy, 1979: 24 [Holotype from Csarnóta 2 – layer 2/B1: nearly complete right coracoid; NMB Vt-81. Figured by Jánossy 1979, fig. 4/9.]

Capella veterior Jánossy: Jánossy 1980b: 13 [New combination.]

Distribution: Early Pliocene (MN 15) of Csarnóta 2, Hungary (Jánossy 1979).

***Gallinago gallinago* (Linnaeus) – Common Snipe**

Scolopax Gallinago Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *G.* cf. *gallinago*), Stránská skála – Čapek site 5a, Czechia (Jánossy 1972 sub *Capella* cf. *gallinago*), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *Capella* cf. *gallinago*), and Stránská skála – Musil's talus cone, layers 4, 4a, 5, 6, 13, and 14, Czechia (Mlíkovský 1995b).

***Gallinago media* (Latham) – Great Snipe**

Scolopax media Latham, 1787 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Jánossy 1979 sub *G. cf. media*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek site 5a, and Absolon's 2nd cave, Czechia (Jánossy 1972 sub *Capella cf. media*), and Stránská skála – Musil's talus cone, layers 5, 6, 13, 14, and 15c, Czechia (Mlíkovský 1995b).

***Gallinago* sp.**

Distribution: Late Miocene (MN 13) of Polgárdi 4 – upper layer, Hungary (Jánossy 1991 sub *Capella* sp.).

Genus *Scolopax* Linnaeus

Scolopax Linnaeus, 1758 [Modern genus.]

***Scolopax carmesinae* Seguí**

Scolopax carmesinae Seguí, 1999b. [Holotype: Humerus.]

Distribution: Late Pliocene (MN 17-18) of the island of Menorca, Spain (Seguí 1999b).

Remarks: Not seen.

***Scolopax rusticola* Linnaeus – Eurasian Woodcock**

Scolopax rusticola Linnaeus, 1758 [Modern species.]

Scolopax rusticola magnus Potapova, 1990: 143 [Holotype from Medvěž'ja cave – layer 3: left humerus; ZIN RO-4029. Figured by Potapova 1990, pl. 3, fig. 7.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 5a, and 9a, Czechia (Jánossy 1972 sub *S. cf. rusticola*), and Stránská skála – Musil's talus cone, layers 4a, 5, 6, 8, 10, 11, 13, and 15e, Czechia (Mlíkovský 1995); and late Pleistocene (MQ 2D) of Medvěž'ja cave, Russia (Potapova 1990 sub *S. rusticola magnus*).

Genus *Limosa* Brisson

Limosa Brisson, 1760 [Modern genus.]

***Limosa limosa* (Linnaeus) – Black-tailed Godwit**

Scolopax Limosa Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992 sub *L. cf. limosa*), Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *L. cf. limosa*); and early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *L. aff. limosa*), Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *L. aff. limosa*), and Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *L. aff. limosa*).

***Limosa* sp.**

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layers 3b, 5, 6, 8, 10, 11, and 13, Czechia (Mlíkovský 1995b: sub *Limosa* spp. Two size classes were recognized, corresponding to *Limosa limosa* (Linnaeus) and *Limosa lapponica* (Linnaeus), respectively, but fragments were usually not identifiable to species).

Genus *Numenius* Brisson

Numenius Brisson, 1760 [Modern genus.]

***Numenius phaeopus* (Linnaeus) – Whimbrel**

Scolopax Phaeopus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layers 5, 6, 8, 11, 13, and 15a, Czechia (Mlíkovský 1995).

***Numenius arquata* (Linnaeus) – Eurasian Curlew**

Scolopax Arquata Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1992 sub *N. cf. arquata*).

Genus *Tringa* Linnaeus

Tringa Linnaeus, 1758 [Modern genus.]

***Tringa erythropus* Linnaeus – Spotted Redshank**

Tringa erythropus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *T. aff. erythropus*).

***Tringa totanus* (Linnaeus) – Common Redshank**

Scolopax totanus Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of Betfia 13, Romania (Kessler 1975 sub *T. aff. totanus*); early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *T. aff. totanus*).

***Tringa ochropus* Linnaeus – Green Sandpiper**

Tringa ochropus [sic!] Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of West Runton – unspecified bed, England (Newton 1882b – bone figured, but not identified, Harrison 1979d), Voigtstedt, Germany (Jánossy 1965 sub *T. cf. ochropus*), and Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *T. aff. ochropus*).

***Tringa glareola* Linnaeus – Wood Sandpiper**

Tringa glareola Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1) of Betfia – Aven, Romania (Kessler 1978).

***Tringa* sp.**

Distribution: Early Miocene (MN 2) of Ravolzhausen, Germany (Martini 1974); late Miocene (MN 13) of Polgárdi 4 – upper layer, Hungary (Jánossy 1991); early Pleistocene (MQ 1) of Betfia 7, Romania (Kessler 1975 sub *T. glareola/ochropus*); early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layers 4, 4a, 5, 6, 8, 9, 13, 14, and 14a, Czechia (Mlíkovský 1995b: three size classes distinguished), and Betfia 5, Romania (Kessler 1975).

Genus *Phalaropus* Brisson

Phalaropus Brisson, 1760 [Modern genus.]

***Phalaropus fulicarius* (Linnaeus) – Red Phalarope**

Tringa fulicaria Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *P. aff.*

fulicarius).

Genus *Incertae sedis*

***Actitis balcanica* Boev**

Actitis balcanica Boev, 1998a: 72 [Holotype from Väršec: distal end of right tarsometatarsus; NMNHS 45. Figured by Boev 1998, fig. 1a-b.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b sub *Actitis* sp., Boev 1998a).

Remarks: Relatively short inner trochlea of the holotypical tarsometatarsus resembles more the condition in the Charadriidae than in the Scolopacidae (Mlíkovský, orig.).

***Elorius paludicola* Milne-Edwards**

Elorius paludicola Milne-Edwards, 1868: 407 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus; MNHN Av-9503. Figured by Milne-Edwards 1867-1868, pl. 63, fig. 23-26. Paralectotypes from Saint-Gérard-le-Puy (according to the material available at MNHN in June 1999 – Mlíkovský orig.): proximal end of right humerus (Av-9501), shaft of right humerus (Av-9495), distal ends of 2 left humeri (Av-9496, Av-9498), distal ends of 2 right humeri (Av-9499, Av-9500), proximal end of left tarsometatarsus (Av-9504), and distal end of right tarsometatarsus (Av-9505). Figured by Milne-Edwards 1867-1868, pl. 63, fig. 27-29 (coracoid), and 30-31 (proximal end of left humerus and distal end of left humerus). I did not find the figured coracoid and proximal end of left humerus and specimens Av-9497 and Av-9502 in MNHN in June 1999. They probably belonged to the syntypical series as well.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1867-1868).

Remarks: *Elorius paludicola* Milne-Edwards, 1868 is type by monotypy of the genus *Elorius* Milne-Edwards, 1868: 407.

***Numenius antiquus* Milne-Edwards**

Numenius antiquus Milne-Edwards, 1868: 415 [Lectotype from Sansan (selected by Cheneval 2000: 361): proximal end of left tarsometatarsus; MNHN Sa-1227. Figured by Milne-Edwards 1867-1868, pl. 64, fig. 27-31; and Cheneval 2000, fig. 18. Paralectotype from Sansan: distal end of right tarsometatarsus; MNHN Sa-1228. Figured by Milne-Edwards 1867-1868, pl. 64, fig. 27; and Cheneval 2000, fig. 19.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1867-1868, Cheneval 2000).

***Scolopax baranensis* Jánossy**

Scolopax baranensis Jánossy, 1979: 23 [Holotype from Csarnóta 2 (layer not given): proximal end of a carpometacarpus (body side not given); NMB Vt-82. Not figured.]

Distribution: Early Pliocene (MN 15) of Csarnóta 2, Hungary (Jánossy 1979).

Remarks: Due to its superficial and irrelevant description by Jánossy (1979) and the absence of any illustration, this species is almost a nomen nudum (see also Mlíkovský 1992b: 442).

***Totanus edwardsi* Gaillard**

Totanus edwardsi Gaillard, 1908: 128 [Holotype from Quercy: distal end of right humerus; BSP 124; apparently destroyed during the World War II (Mlíkovský, orig.). Figured by Gaillard 1908, fig. 37.]

Tringa edwardsi (Gaillard): Bocheński 1997: 315 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1908, Mayr 2000i:

635).

***Totanus lartetianus* Milne-Edwards**

Totanus Lartetianus Milne-Edwards, 1863: 160 [Holotype from Saint-Gérard-le-Puy: right humerus; MNHN. Figured by Milne-Edwards 1867-1868, pl. 63, fig. 17-18. Milne-Edwards (1868: 402-406) described various skeletal elements of *Totanus lartetianus*, which Brodkorb (1967: 186) considered [syn]types of the species. However, Milne-Edwards (1863) described the species only on the basis of a humerus, which has therefore to be regarded as the holotype of the species.]

Tringa lartetiana (Milne-Edwards): Bocheński 1997: 316 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1867-1868).

***Totanus grivensis* Ennouchi**

Totanus grivensis Ennouchi, 1930: 92 [Syntypes from Grive-Saint-Alban – early collections: proximal part of right humerus (ML Lgr-75), and distal part of left humerus (ML LGr-74). Figured by Ennouchi 1930, pl. 2, fig. 1-4 (proximal part) and fig. 5-8 (distal part).]

Tringa grivensis (Ennouchi): Bocheński 1997: 316 [New combination.]

Distribution: Middle Miocene (MN 7-8) of La Grive-Saint-Alban, France (Ennouchi 1930).

***Totanus majori* Lydekker**

Totanus majori Lydekker, 1893: 521 [Holotype from Grive-Saint-Alban: left humerus; BMNH. Figured by Lydekker 1893, pl. 41, fig. 14.]

Tringa majori (Lydekker): Bocheński 1997: 316 [New combination.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Lydekker 1893b, Ennouchi 1930: 92); and middle Miocene (MN 8) of Grive-Saint-Alban – fissure M, France (Ballmann 1969a: 187).

***Totanus minor* Ennouchi**

Totanus minor Ennouchi, 1930: 94 [Syntypes from Grive-Saint-Alban (all in ML): three left humeri (LGr-80, 81, 82), right humerus (LGr-84), and right humerus lacking proximal end (LGr-83). Figured by Ennouchi 1930, pl. 2, fig. 9-12 (one complete humerus); the remaining four syntypes remain not figured.]

Erolia ennouchii Brodkorb, 1967: 189 [New name for *Totanus minor* Ennouchi, 1930, believed to be junior secondary homonym of *Tringa cinclus minor* Schlegel, 1844 = *Erolia alpina* (Linnaeus, 1758) = *Calidris alpina* (Linnaeus, 1758). I was not able to find any paper, in which the species-group name *minor* Schlegel was published in combination with the genus *Totanus* Bechstein, 1803. It is thus probable, that *minor* Ennouchi is not a secondary homonym of *minor* Schlegel.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Ennouchi 1930, Gaillard 1939: 64); and middle Miocene (MN 8) of Grive-Saint-Alban – fissure M, France (Ballmann 1969a: 187).

***Totanus numenioides* Serebrovs'kyj**

Totanus numenioides Serebrovs'kyj, 1941b: 476 [Holotype from Odesa – catacombs: left humerus with damaged distal end; present location unknown. Figured by Serebrovs'kyj 1941b; and Dement'ev 1964, fig. 687a,b.]

Tringa numenioides (Serebrovs'kyj): Dement'ev 1964: 682 [New combination.]

Tringa numenioidea (Serebrovs'kyj): Bocheński 1997: 316 [Marked as a new combination; species name misspelled.]

Distribution: Early Pliocene (MN 15) of Odesa – catacombs, Ukraine (Serebrovs'kyj 1941b).

***Totanus scarabellii* Portis**

Totanus Scarabellii Portis, 1887: 181 [Type from Senigallia: partial hind limb embedded in a slab (body side unknown), incl. distal end of tibiotarsus, tarsometatarsus, two anterior toes, and proximal phalanx of the hind toe; MCI, uncatalogued. Figured by Portis 1887, pl. 1, fig. 1.]

Tringa scarabellii (Portis): Bocheński 1997: 316 [New combination.]

Distribution: Late Miocene (MN 13) of Senigallia, Italy (Massalongo & Scarabelli-Gommi-Flamini 1859: 19 as an unidentified bird remain, Portis 1887).

***Tringa gracilis* Milne-Edwards**

Tringa gracilis Milne-Edwards, 1868: 411 [Syntypes from Saint-Gérard-le-Puy (all in MNHN): large number of bones, incl. at least the following elements: a furcula, proximal end of a scapula, proximal end of a humerus, distal end of a humerus, an ulna, proximal end of a tibiotarsus, distal end of a tibiotarsus, and distal end of a tarsometatarsus. Figured by Milne-Edwards 1867-1868, pl. 58 (not pl. 64 as printed in Milne-Edwards 1867-1868: 411), fig. 2, and pl. 64, fig. 1-4 (tarsometatarsus), 5-10 (tibiotarsus), 11-13 (furcula), 14-16 (scapula), 17-19 (humerus), 20-21 (ulna).]

Erolia gracilis (Milne-Edwards): Brodkorb 1967: 188 [New combination.]

Distribution: Early Miocene (MN 1) of Weisenau-AS, Germany (Milne-Edwards 1867-1868: 413); and early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1867-1868).

***Tringa grigorescui* Kessler & Gál**

Tringa grigorescui Kessler & Gál, 1996: 75 [Holotype from Ciobănița: proximal end of left carpometacarpus; LPUB 283. Figured by Kessler & Gál 1996, fig. 2a-b.]

Distribution: Middle Miocene (MN 8) of Ciobănița, Romania (Kessler & Gál 1996).

Family Glareolidae Brehm

Glareolidae Brehm, 1831 [Modern family.]

Genus *Mioglareola* Ballmann

Mioglareola Ballmann, 1979: 68 [Type by monotypy: *Mioglareola gregaria* Ballmann, 1979.]

Distribution: Early Miocene (MN 4) of Czechia, and middle Miocene (MN 6) of Germany.

***Mioglareola dolnicensis* (Švec)**

Larus dolnicensis Švec, 1980: 380 [Holotype from Dolnice: distal end of left humerus; coll. Fejfar, uncatalogued. Figured by Švec 1980, pl. 1, fig. 2-3.]

Mioglareola dolnicensis (Švec): Mlíkovský 2000c: 94 [New combination.]

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Švec 1980, Mlíkovský 2000c).

***Mioglareola gregaria* Ballmann**

Mioglareola gregaria Ballmann, 1979: 68 [Holotype from Steinberg: imperfect cranium; BSP 1970-XVIII-851. Figured by Ballmann 1979, pl. 1, fig. 3-5, text-fig. 3a-c, 4a-b.]

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský 2000c), and middle Miocene (MN 6) of Steinberg, Germany (Ballmann 1979).

Genus *Glareola* Brisson

Glareola Brisson, 1760 [Modern genus.]

Glareola neogena Ballmann

Glareola neogena Ballmann, 1979: 81 [Holotype from Steinberg: right humerus; BSP 1970-XVIII-852. Figured by Ballmann 1979, pl. 1, fig. 9, pl. 2, fig. 1.]

Distribution: Middle Miocene (MN 6) of Steinberg, Germany (Ballmann 1979), and Goldberg, Germany (Ballmann 1979).

Genus *Cursorius* Latham

Cursorius Latham, 1790 [Modern genus.]

Cursorius sp.

Distribution: Late Miocene (MN 13) of Polgárdi 4 – upper layer, Hungary (Jánossy 1991); and early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992 sub ?*Cursorius* sp.).

Family *Thinocoridae* Sundevall

Thinocoridae Sundevall, 1836 [Modern family.]

Pedionomidae Bonaparte, 1856 [Modern family.]

Turnipacidae Mayr, 2000i: 626 [Type genus: *Turnipax* Mayr, 2000i.]

Remarks: Olson & Steadman (1981) demonstrated, that *Pedionomus* Gould, 1841 is closely related to the Thinocoridae. The discovery of *Turnipax* (Mayr 2000i) allows to understand the Thinocoridae (which have nomenclatural priority) in the broader sense to encompass the presumably monophyletic group of quail-like birds related to the Scolopacidae (see Strauch 1978, Björklund 1994 and Chu 1995 for the latter relationship). Accordingly, I synonymize here Pedionomidae Bonaparte and Turnipacidae Mayr with the Thinocoridae Sundevall.

Mayr (2000i) listed small extremitas omalis of the coracoid as a distinguishing character of his family Turnipacidae. However, the configuration of this element is variable in the Thinocoridae s.l., ranging from open in *Pedionomus* Gould (see Olson & Steadman 1981) over medium small in *Thinocorus* Eschscholtz, 1829 (see Strauch 1976) to small in *Turnipax* Mayr (see Mayr 2000i).

Genus *Turnipax* Mayr

Turnipax Mayr, 2000i: 626 [Type by original designation: *Turnipax dissipata* Mayr, 2000i.]

Turnipax dissipata Mayr

Turnipax dissipata Mayr, 2000i: 627 [Holotype from Lubéron: partial, largely disarticulated skeleton in slab, incl. some vertebrae and ribs, both scapulae, both coracoids, furcula, both humeri, pelvis in two parts, both femora, left tibiotarsus with associated tarsometatarsus and phalanges digitorum pedis, right tibiotarsus with associated tarsometatarsus and phalanges digitorum pedis; SMF Av-427. Figured by Mayr 2000i, fig. 9/1 (left coracoid), fig. 9/2 (right coracoid), 9/3 (right scapula), 9/4 (left humerus), 9/5 (right humerus), 9/6 (one or two pelvises, left femur and right femur), 9/7 (left leg), 9/8 (right leg), 9/9 (all syntypes).

Distribution: Early/middle Oligocene (MP 21-25) of Lubéron, France (Mayr 2000i). If Mayr (2000i) is correct in identifying Lubéron with Céreste, then the known distribution of *Turnipax dissipata* is limited to the middle Oligocene (MP 23) of Céreste, France.

Family Pterocletidae Bonaparte

Pterocletidae Bonaparte, 1831 [Modern family.]

Genus *Archaeoganga* Mourer-Chauviré

Archaeoganga Mourer-Chauviré, 1992a: 231 [Type by original designation: *Archaeoganga pinguis* Mourer-Chauviré, 1992a.]

Distribution: Eocene or Oligocene (MP 16-30) Quercy, France.

Archaeoganga pinguis Mourer-Chauviré

Archaeoganga pinguis Mourer-Chauviré, 1992a: 233 [Holotype from Quercy: right coracoid; MNHN QU-16960. Figured by Mourer-Chauviré 1992a, unnumbered pl., fig. a-b; and Mourer-Chauviré 1993, pl. 1, fig. a-b.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1992a, 1993a).

Archaeoganga larvatus (Milne-Edwards)

Pterocles larvatus Milne-Edwards, 1892: 71 [Holotype from Quercy: left coracoid; MNHN QU-15854. Figured by Mourer-Chauviré 1993, pl. 1, fig. e-f.]

Archaeoganga larvata (Milne-Edwards): Mourer-Chauviré 1992a: 231 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1993a).

Archaeoganga validus (Milne-Edwards)

Pterocles validus Milne-Edwards, 1892: 70 [Holotype from Quercy: left tarsometatarsus; MNHN QU-15853. Figured by Mourer-Chauviré 1993a, pl. 3, fig. c-d.]

Archaeoganga validus (Milne-Edwards): Mourer-Chauviré 1992a: 231 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1993a).

Genus *Leptoganga* Mourer-Chauviré

Leptoganga Mourer-Chauviré, 1993a: 87 [Type by original designation: *Pterocles sepultus* Milne-Edwards, 1869.]

Distribution: Late Oligocene (MP 28) of France, and early Miocene (MN 1-2) of France.

Leptoganga sepultus (Milne-Edwards)

Pterocles sepultus Milne-Edwards, 1869: 294 [Holotype from Saint-Gérard-le-Puy: left tarsometatarsus; MNHN Av-2844. Figured by Milne-Edwards 1869, pl. 141, fig. 1-9.]

Leptoganga sepultus (Milne-Edwards): Mourer-Chauviré 1993a: 88 [New combination.]

Distribution: Late Oligocene (MN 28) of Pech Desse, France (Mourer-Chauviré 1993a), and Pech du Fraysse, France (Mourer-Chauviré 1993a); early Miocene (MN 1) of Paulhiac, France (Mourer-Chauviré 1993); and early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869, Mourer-Chauviré 1993a).

Genus *Gerandia* Lambrecht

Gerandia Lambrecht, 1933: 602 [Type by monotypy: *Columba calcaria* Milne-Edwards, 1869.]

Gerandia calcaria (Milne-Edwards)

Columba calcaria Milne-Edwards, 1869: 292 [Holotype from Saint-Gérard-le-Puy: left humerus; MNHN Av-9672. Figured by Milne-Edwards, 1869-1871, pl. 141, fig. 10-14.]

Gerandia calcaria (Milne-Edwards): Lambrecht 1933: 602 [New combination.]

Distribution: Early Miocene (MN 2a) from Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Lambrecht 1933: 602).

Remarks: Described in the Columbidae (Milne-Edwards 1869). The holotypical humerus differs from the same element of the Columbidae, and agrees with that of the Pterocletidae in having: (1) deep muscular impression on the external side of the head, (2) ectepicondylar process pointed and well separated from the shaft, and (3) distal end of the entepicondyle bent medially. The taxonomic position within the Pterocletidae has to be studied, but the taxon seems to be different from *Archaeoganga* Mourer-Chauviré, *Leptoganga* Mourer-Chauviré, and the modern *Pterocles* Temminck, 1815, with which I compared it.

Family Charadriidae Vigors

Charadriidae Leach, 1820 [Modern family.]

Distribution: Although the earliest identified Charadriidae are known from MN 16, there are two Miocene records of unidentified Charadriidae, incl. those from the early Miocene (MN4b) of Dolnice, Czechia (Mlíkovský 1996e), and the middle Miocene (MN 5) of Vieux-Collonges, France (Ballmann 1972).

Genus *Charadrius* Linnaeus

Charadrius Linnaeus, 1758 [Modern species.]

Charadrius morinellus Linnaeus – Eurasian Dotterel

Charadrius morinellus Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a sub array of *Charadrius morinellus*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *C. aff. morinellus*).

Charadrius sp.

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b and 1990b; not mentioned by Jánossy 1992); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layers 6 and 13, Czechia (Mlíkovský 1995b).

Genus *Pluvialis* Brisson

Pluvialis Brisson, 1760 [Modern genus.]

Pluvialis squatarola (Linnaeus) – Grey Plover

Tringa Squatarola Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub aff. *Squatarola squatarola*), and Stránská skála – Musil's talus cone, layers 3b, 5, 8, and 13, Czechia (Mlíkovský 1995b sub *P. cf. squatarola*).

Pluvialis sp.

Distribution: Early Pleistocene (MQ 1b) of Voigtstedt (Jánossy 1965 sub *Squatarola* sp.).

Genus *Vanellus* Brisson

Vanellus Brisson, 1760 [Modern genus.]

Vanellus vanellus (Linnaeus) – Northern Lapwing

Tringa Vanellus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992 sub *V. cf. vanellus*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 1-3, 5a, and 9a, Czechia (Jánossy 1972 sub (Jánossy 1972 sub *V. cf. vanellus*), and Stránská skála – Musil's talus cone, layers 4, 4a, 5, 6, 7, 8, 10a, 11, and 13, Czechia (Mlíkovský 1995b).

Family Laridae Vigors

Laridae Vigors, 1825 [Modern family.]

Stercorariidae Gray, 1871 [Modern family.]

Remarks: I see no reason why to separate Stercorariidae from the Laridae, because stercorariid genera fit well in the adaptive radiation of the Laridae. Hence, I include here Stercorariidae Gray in the Laridae Vigors.

Genus *Gaviota* Miller & Sibley

Gaviota Miller & Sibley, 1941: 563 [Type by original designation: *Gaviota niobrara* Miller & Sibley, 1941.]

Distribution: The extralimital record is limited to the late Miocene (Clarendonian) of Nebraska, USA (Miller & Sibley 1941, Olson 1985a: 182), from where *Gaviota niobrara* Miller & Sibley, 1941 was described.

Gaviota lipsiensis Fischer

Gaviota lipsiensis Fischer, 1983a: 152 [Holotype from Espenhain: distal end of right humerus; NKMB Av-732. Figured by Fischer 1983a, pl. 12, fig. 1-4.]

Distribution: Middle Oligocene (MP 23-24) of Espenhain, Germany (Fischer 1983a).

Remarks: The taxonomic position of this species needs confirmation.

Genus *Laricola* n. g.

Laricola Mlíkovský, this paper [Type by original designation: *Larus elegans* Milne-Edwards, 1868.]

Diagnosis: A gull with relatively long hindlimbs and short forelimbs (in comparison with the modern *Larus* Linnaeus).

Species included: *Laricola elegans* (Milne-Edwards), and *Laricola totanoides* (Milne-Edwards).

Etymology: “Walking gull”, coined from *Larus*, modern genus of gulls, and *kola*, Greek expression for a runner (Latinized). The name is masculine in gender.

Distribution: Late Oligocene (MP 25-30) of France, and early Miocene (MN 2-4) of France and Czechia.

Laricola elegans (Milne-Edwards)

Larus elegans Milne-Edwards, 1868: 350 [Syntypes from Saint-Gérand-le-Puy: large amount of bones, incl. at least one each of the following elements: skull, coracoid, furcula, scapula, humerus, ulna, radius, carpometacarpus, femur, tibiotarsus, and tarsometatarsus; MNHN. Figured by Milne-Edwards 1867-1868, pl. 56, fig. 11-15 (tarsometatarsus), 16-18 (tibiotarsus), 19-21 (femur), 22 (synsacrum), 23 (sternum), 24-26 (coracoid), 27-29 (scapula), pl. 57, fig. 1 (skull), 2-3 (humerus), 4-5 (ulna), 6-9 (carpometacarpus), 10-11 (furcula), and pl. 58, fig. 1 (reconstructed skeleton).]

Laricola elegans (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Late Oligocene (MP 25) of Antoingt, France (Lydekker 1891a: 178); late Oligocene (MP 30) of Gannat, France (Milne-Edwards 1867b: 350); early Miocene (MN 2a) of Saint-Gérand-le-Puy, France (Milne-Edwards 1867-1868); early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský orig.); and middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Shufeldt 1896: 515). The latter record is uncertain (Brodkorb 1967: 206).

***Laricola totanoides* (Milne-Edwards)**

Larus totanoides Milne-Edwards, 1868: 358 [Syntypes from Saint-Gérard-le-Puy: large amount of bones, incl. at least one each of the following elements: coracoid, humerus, ulna, radius, carpometacarpus, femur, tibiotarsus, tarsometatarsus; MNHN. Figured by Milne-Edwards 1867-1868, pl. 57, fig. 12-15 (tarsometatarsus), and 16-17 (humerus).]

Laricola totanoides (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1867-1868). The record from the middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Shufeldt 1896: 515) is uncertain.

Genus *Larus* Linnaeus

Larus Linnaeus 1758 [Modern genus.]

Rissa Stephen, 1826 [Modern genus.]

***Larus teruelensis* (Villalta)**

Totanus teruelensis Villalta, 1963: 268 [Lectotype from Mansuetos (selected by Olson 1985a: 175): distal end of right humerus; present location unknown – A. Sánchez Marco, in litt. 1995. Figured by Villalta 1963, pl. 3, fig. 1, 1a-b. Paralectotype: distal end of right ulna from Mansuetos; present location unknown – A. Sánchez Marco, in litt. 1995. Figured by Villalta 1963, pl. 2, fig. 5, 5a-b. Note that Villalta (1963: 268) based this species on both listed fragments, which he believed to come probably from a single individual. Olson (1985a: 175) erroneously stated that the species was based solely on the humerus fragment. Nevertheless, the latter action can be interpreted as a selection of the lectotype.]

Tringa teruelensis (Villalta): Bocheński 1997: 316 [New combination.]

Larus teruelensis (Villalta): Mlíkovský, this paper [New combination.]

Distribution: Late Miocene (MN 12) of Mansuetos, Spain (Villalta 1963).

Remarks: Deep longitudinal impression of musculus brachialis antiquus allows to assign the fossil to the Laridae (see also Olson 1985: 175). Within the family the fossil distal end of humerus agrees with the same element of *Larus* Linnaeus and differs from that of *Sterna* Linnaeus and *Chlidonias* Rafinesque especially in being relatively broad. Accordingly, I transfer here *Totanus teruelensis* Villalta to the modern genus *Larus* Linnaeus.

***Larus ridibundus* Linnaeus – Black-headed Gull**

Larus ridibundus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992 sub *Larus* (?) cf. *ridibundus*); and early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), Přezletice, Czechia (Jánossy 1983 sub *L.* cf. *ridibundus*), and Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *L.* cf. *ridibundus*).

***Larus canus* Linnaeus – Mew Gull**

Larus canus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *L.* aff. *canus*), and Stránská skála – Musil's talus cone, layer 5, Czechia (Mlíkovský 1995b).

***Larus tridactylus* Linnaeus – Black-legged Kittiwake**

Larus tridactylus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999 sub cf. *Rissa tridactyla*).

***Larus* sp.**

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský 1996e sub ?*Larus*); middle Miocene (MN 8) of Cretința, Romania (Grigorescu & Kessler 1977); and late Miocene (MN 9) of Chișinău, Moldova (Kessler 1984). The alleged record of *Larus* sp. from the late Miocene (MN 9) of Kalfa, Moldova (Ganea 1965) refers to an unidentified member of the Anatidae (Kuročkin & Ganea 1972).

Genus *Sterna* Linnaeus

Sterna Linnaeus, 1758 [Modern species.]

***Sterna* sp.**

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský 1996e sub ?*Sterna*); and early Pleistocene (MQ 1a) of Primorsk, Ukraine (Vojistvens'kyj 1967).

Genus *Chlidonias* Rafinesque

Chlidonias Rafinesque, 1822 [Modern genus.]

***Chlidonias* sp.**

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub aff. *C. nigra*?).

Genus *Stercorarius* Brisson

Stercorarius Brisson, 1760 [Modern genus.]

***Stercorarius pomarinus* (Temminck) – Pomarine Jaeger**

Lestris pomarinus Temminck, 1815 [Modern species.]

Stercorarius pomarinus philippi Mourer-Chauviré, 1975a: 137 [Holotype from Fage – layer 35: left tarsometatarsus; FSL 41657. Figured by Mourer-Chauviré 1975a, pl. 13, fig. 14.]

Distribution: Middle Pleistocene (MQ 2B) of Fage – layer 35, France (Mourer-Chauviré 1975a sub *S. pomarinus philippi*).

Genus *incertae sedis****Larus desnoyersii* Milne-Edwards**

Larus Desnoyersii Milne-Edwards, 1863: 161 [Holotype from Saint-Gérard-le-Puy: distal end of a humerus; MNHN. Not figured.]

Larus Desnoyersii Milne-Edwards, 1868: 344 [Lectotype from Saint-Gérard-le-Puy: distal end of a humerus; MNHN (same specimen is the holotype of *L. Desnoyersii* Milne-Edwards, 1863); MNHN. Not figured. Paralectotypes (all in MNHN): unknown number of bones, incl. cervical vertebrae, scapula, coracoid, humerus, ulna, radius, phalanx I digiti majoris, fragmentary pelvis, femur, tibiotarsus, and tarsometatarsus. Figured by Milne-Edwards 1867-1868, pl. 54, fig. 15 partim (tibiotarsus), 15-18 (tarsometatarsus), 15, 19-20 (femur), 21-23 (coracoid), 24-26 (humerus), pl. 55 (reconstruction), and pl. 56, fig. 1-2 (pelvis), 3-4 (ulna), 5-7 (carpometacarpus), 8 (phalanx I digiti majoris), 9-10 (scapula). This both junior objective synonym and junior homonym of *Larus Desnoyersii* Milne-Edwards, 1863.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1867-1868, Miller & Sibley 1941, Olson 1985a: 180).

Remarks: Olson (1985a: 180) pointed out that this species possibly should be transferred to the Stercorariidae (see also Miller & Sibley 1941: 566). I agree that it does not belong to the genus *Larus* Linnaeus.

Family Alcidae Vigors

Alcidae Vigors, 1825 [Modern family.]

Petralcinae Mlíkovský, 1987: 136 [Type genus: *Petralca* Mlíkovský, 1987a.]

Genus *Petralca* Mlíkovský

Petralca Mlíkovský, 1987: 136 [Type by original designation: *Petralca austriaca* Mlíkovský, 1987a.]

Petralca austriaca Mlíkovský

Petralca austriaca Mlíkovský, 1987a: 136 [Holotype from Traun-Pucking: incomplete skeleton in slab and counter-slab; NHMW 1980/25. Figured by Mlíkovský 1987a, text-fig. 4-5, pl. 1-2.]

Distribution: Late Oligocene (MP 30) of Traun-Pucking, Austria (Mlíkovský 1987).

Genus *Uria* Brisson

Uria Brisson, 1760 [Modern genus.]

Uria aalge (Pontoppidan) – Common Murre

Colymbus aalge Pontoppidan, 1763 [Modern species.]

Distribution: Late Pliocene (MN 18) of Aldeby, England (Newton 1882b, Harrison 1979d), and Chillesford, England (Harrison 1979d).

Genus *Alca* Linnaeus

Alca Linnaeus, 1758 [Modern genus.]

Plautus Brünnich, 1772 [Modern genus.]

Pinguinus Bonnaterre, 1791 [Modern genus.]

Alca impennis Linnaeus – Great Auk

Alca impennis Linnaeus, 1758: 130 [Modern species.]

Alca major Brisson, 1760: 85 [Unavailable name; book not strictly binomial – ICZN 1955.]

Plautus impennis (Linnaeus): Brünnich, 1772: 78 [New combination.]

Pinguinus impennis (Linnaeus): Bonnaterre, 1791: 29 [New combination.]

Distribution: Early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999); late Pleistocene (MQ 2C) of Devil's Tower, Gibraltar (Bate 1928), Gorham's Cave, Gibraltar (Eastham 1968), Figueira Brava – layer 2, Portugal (Mourer-Chauviré & Antunes 1991, 2000), Archi, Italy (Ascenzi & Segre 1971, Cassoli & Segre 1985); and late Pleistocene (MQ 2D) of Blomvåg, Norway (Lie 1986), Cosquer, France (Clottes et al. 1992, Clottes & Courtin 1993, d'Errico 1994a,b,c, McDonald 1994; see also Cleere 1992), Nerja, Spain (Eastham 1986), Urriaga, Spain (Eastham 1985, 1989), Pendo, Spain (wall engraving, H. Breuil in Alcalde del Río et al. 1912, Eastham 1968, Gonzales Morales 1980, d'Errico 1994a,b,c; but see McDonald 1994), Paglicci, Italy (Mezzena & Palma di Cesnola 1992 sub *A. tordus* or *A. impennis*), Romanelli – layer B, Italy (Regàlia 1904 sub *Uria* sp., Blanc 1927, 1928, Cassoli et al. 1979, Cassoli & Segre 1985), and Arene Candide, Italy (Cassoli 1980: 183). Extralimital records are limited to the late Pleistocene of Porto Santo, Madeira (Pieper 1985), and Norfolk, Virginia (Ray et al. 1968).

The following late Pleistocene records are invalid: Gargas, Jersey, England (Andrews 1920 – see Breuil & Bégouën 1936, d'Errico 1994a,c, McDonald 1994), Raymondén, France (sculpture, Hardy 1891 – see Breuil et al. 1909, H. Breuil in Alcalde del Río et al. 1912: 38, Lambrecht 1933: 790-791, d'Errico 1994a,c, McDonald 1994), and Hermanstorp, Sweden (egg, Munthe 1914, reidentified as that of *Cygnus cygnus* Linnaeus by Löppenthin 1952).

The Holocene record is more abundant, being limited to the northern part of the Atlantic

Ocean. In Europe it includes sites in Iceland (Blasius 1884, Grieve 1885, Bardharson 1911, Petersen 1995), the Hebrides (Mellars 1978, 1987), Scotland (Blasius 1884, Bell 1915, Serjeantson 1993), Ireland (Ussher 1897, 1899a,b, 1902, Knowles 1895, Lambrecht 1933, Woodman 1978), England (Blasius 1884, Grieve 1885, Bell 1915, Fisher 1997, Stewart 1997), Norway (Lambrecht 1933, Olsen 1967, Hufthammer 1982, Montevecchi & Hufthammer 1990), Sweden (Lambrecht 1933, Lageraes 1989), Denmark (Blasius 1884, Winge 1903, Grigson 1989), Holland (Wijngarden-Bakker 1978, Kompanje & Kerkhoff 1991), Baltic Russia (Potapova 1994), Atlantic France (Péquart & Péquart 1926, Péquart et al. 1937), Atlantic Spain (Elorza & Sánchez Marco 1993, Hernández Carrasquilla 1994), and Mediterranean Spain (Hernández Carrasquilla 1994). For the occurrence in Greenland see Meldgaard (1988). In North America, the record is available from the Atlantic coast from Quebec in the north (Bisseling 1938, Jordan & Olson 1982) to Florida in the south (Hay 1902, Bullen & Sleight 1950, Brodkorb 1960b, 1967, Weigel 1988). For maps see Cassoli & Segre (1985), Mourer-Chauviré & Antunes (1991, 1999d), and especially Tyrberg (1999, fig. 1).

The disappearance of the Great Auk from Europe is not mapped as yet. It seems to have disappeared from the Mediterranean Sea at the close of the Pleistocene (being still recorded in Dryas III, but not later – excepting a Holocene record from Nerja in Spain near Gibraltar, Hernández Carrasquilla 1994), and from Norway ca. 1500 years ago (Hufthammer 1982). Its breeding distribution was probably limited to the islands in the northwestern Atlantic Ocean in historical times (see Bengtson 1984), but 18th-century records, probably attributable to vagrants, are available from the British Islands (Grieve 1885) and northwestern France (Scherdlin 1926).

Great Auk became extinct in the mid 19th century. The last generally accepted record dates back to June 1844, when two individuals were killed on the Eldey islet off Iceland (Greenway 1967, Bengtson 1984, Luther 1966, 1986), but an uncertain record is also from 1852 (Newton 1861, Vinokurov 1992). Data on the natural history of the Great Auk were summarized by Steenstrup (1855), Preyer (1862), Blasius (1884, 1903) and Grieve (1885, 1888); see also Violani (1974), Olson et al. 1979, Bengtson (1984), Nettleship & Evans (1985), Livezey (1988), Hobson & Montevecchi (1991), Fuller (1999), and Mourer-Chauviré (1999d). See also Burness & Montevecchi (1992) and Serjeantson (2001)

***Alca torda* Linnaeus – Razorbill**

Alca Torda Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Bacton, England (Harrison 1979d).

Genus *Cephus* Pallas

Cephus Pallas, 1769 [Modern genus.]

***Cephus storeri* Harrison**

Cephus storeri Harrison, 1977: 238 [Holotype from “Red Crag of Suffolk”: left carpometacarpus; BMNH A-4986. Not figured.]

Distribution: Late Pliocene (MN 16-17) of an unknown locality within the Red Crag in Suffolk, England (Harrison 1977).

Genus *incertae sedis*

***Uria ausonia* Portis**

Uria ausonia Portis, 1887: 195 [Nomen nudum; no description or indication.]

Uria ausonia Portis, 1889: 15 [Holotype from Orciano Pisano: distal end of left humerus; IGF 14875. Figured by Portis 1889, pl. 1, fig. 28-30.]

Distribution: Middle Pliocene (MN 15-16) of Orciano Pisano, Italy (Portis 1887, 1889).

Remarks: This species does not belong to the genus *Uria* Brisson, 1760 (S.L. Olson, pers. communication, 1997).

Order Piciformes Meyer & Wolf

Pici Meyer & Wolf, 1810 [Modern order.]

Sandcoleiformes Houde & Olson, 1992 [Type genus: *Sandcoleus* Houde & Olson, 1992.]

Family Zygodactylidae Brodkorb

Zygodactylidae Brodkorb, 1971b: 257 [Type genus: *Zygodactylus* Ballmann, 1969b.]

Primoscenidae Harrison & Walker 1977a: 47 [Type genus: *Primoscens* Harrison & Walker, 1977a.]

Sandcoleidae Houde & Olson, 1992: 139 [Type genus: *Sandcoleus* Houde & Olson, 1992.]

Quercypsittidae Mourer-Chauviré, 1992c: 170 [Type genus: *Quercypsitta* Mourer-Chauviré, 1992c.]

Miopiconidae Mayr 1998c: 50 [Type genus: *Miopico* Mayr, 1998c.]

Pseudasturidae Mayr, 1998b: 200 [Type genus: *Pseudastur* Mayr, 1998b.]

Primoscinidae Harrison & Walker: Mlíkovský 2000d: 81 [Spelling emended.]

Distribution: Early Eocene (MP 7) of Denmark (based on an unnamed form from Fur, figured by Hoch 1975, fig. 6 – see Houde & Olson 1992: 157), early Eocene (MP 8-13) of England (see also Dyke 2001a) and Germany, late Eocene (MP 16-19) of France, early Miocene (MN 3-4) of Germany and Czechia, and middle Miocene (MN 8) of France. Extralimital record includes a number of genera and species from the early to middle Eocene of Wyoming (Houde & Olson 1989, 1992), and *Miopico benimellalensis* Mayr, 1998c from the middle Miocene (MN 6) of Beni Mellal in Morocco (Brunet 1971 sub *Jynx* cf. *torquilla*, Mayr 1998c).

Remarks: I include in this family all ancestral Eocene piciforms, plus their direct Miocene offshoots (*Zygodactylus* Ballmann 1969b, and *Miopico* Mayr, 1998c). Hence, I synonymize here Primoscenidae Harrison & Walker, Sandcoleidae Houde & Olson, Quercypsittidae Mourer-Chauviré, Miopiconidae Mayr, and Pseudasturidae Mayr with the Zygodactylidae Brodkorb. Subsequent research may show, that some of these forms should be assigned to the derived families as representatives of respective ancestral radiations. A comprehensive review of the Zygodactylidae will probably result in a reduction of the number of recognized genera.

Genus *Primoscens* Harrison & Walker

Primoscens Harrison & Walker, 1977a: 47 [Type by original designation: *Primoscens minutus* Harrison & Walker, 1977a.]

Primoscens minutus Harrison & Walker

Primoscens minutus Harrison & Walker, 1977a: 48 [Holotype from Bognor Regis: right carpometacarpus; BMNH A-4681. Figured by Harrison & Walker 1977a, pl. 11, fig. e-j.]

Distribution: Early Eocene (MP 8-9) of Bognor Regis, England (Harrison & Walker 1977a, Olson & Feduccia 1979, Mayr 1998c).

Genus *Primozydactylus* Mayr

Primozydactylus Mayr, 1998c: 39 [Type by original designation: *Primozygodactylus danielsi* Mayr, 1998b.]

Distribution: Early Eocene (MP 8-8/9) of England, and middle Eocene (MN 11) of Germany.

Primozygodactylus danielsi Mayr

Primozygodactylus danielsi Mayr, 1998c: 47 [Holotype from Messel: partial skeleton in plate and counter-plate; SMF ME-2522a,b. Figured by Mayr 1998c, pl. 12, 14 (left).]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998c).

***Primozygodactylus major* Mayr**

Primozygodactylus major Mayr, 1998c: 50 [Holotype from Messel: skeleton in plate and counter-plate; SMF ME-1758a,b. Figured by Mayr 1998c, pl. 16 (below).]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998c).

***Primozygodactylus ballmanni* Mayr**

?*Primozygodactylus ballmanni* Mayr, 1998c: 50 [Holotype from Messel: incomplete skeleton in plate and counter-plate; SMF ME-2108a,b. Figured by Mayr 1998c, pl. 15 (left).]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998c).

***Primozygodactylus* sp.**

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Mayr 1998c: 51); and early Eocene (MP 8-9) of Bognor Regis, England (Mayr 1998c: 52).

Genus *Psittacopes* Mayr & Daniels

Psittacopes Mayr & Daniels, 1998: 158 [Type by original designation: *Psittacopes lepidus* Mayr & Daniels, 1998.]

***Psittacopes lepidus* Mayr & Daniels**

Psittacopes lepidus Mayr & Daniels, 1998: 159 [Holotype from Messel: skeleton lacking right leg; SMF ME-1279. Figured by Mayr & Daniels 1998, pl. 1a-b; and Mayr 2000g, fig. 6.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr & Daniels 1998).

Genus *Pseudastur* Mayr

Pseudastur Mayr, 1998b: 201 [Type by original designation: *Pseudastur macrocephalus* Mayr, 1998b.]

***Pseudastur macrocephalus* Mayr**

Pseudastur macrocephalus Mayr, 1998b: 202 [Holotype from Messel: complete skeleton in slab; WDC C-MG-94. Figured by Mayr 1998b, fig. 1.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Hoch 1988 sub unidentified bird, Mayr 1998b).

Genus *Selmes* Peters

Selmes "Peters" Mayr, 1998c: 64 [Nomen nudum; no definition.]

Selmes "Peters" Mayr & Peters, 1998: 189 [Nomen nudum; no definition.]

Selmes Peters, 1999: 217 [Type by original designation: *Selmes absurdipes* Peters, 1999.]

***Selmes absurdipes* Peters**

Selmes absurdipes "Peters" Mayr, 1998c: 64 [Nomen nudum; no description or indication.]

Selmes absurdipes "Peters" Mayr & Peters, 1998: 189 [Nomen nudum; no description or indication.]

Selmes absurdipes Peters, 1999: 218 [Holotype from Messel: Incomplete skeleton in a slab; SMF ME-2375. Figured by Peters 1999, fig. 1-2; Mayr 2001e, fig. 1.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Peters 1999, Mayr 2001e); and middle Eocene (MP 13) of Geiseltal XXXVI, Germany (Fischer 1987 sub *Eoglaucidium pallas*, Mayr 2001e).

Genus *Masillacolius* Mayr & Peters

Masillacolius Mayr & Peters, 1998: 184 [Type by original designation: *Masillacolius brevidactylus* Mayr & Peters, 1988.]

Masillacolius brevidactylus Mayr & Peters

Masillacolius brevidactylus Mayr & Peters, 1998: 185 [Holotype from Messel: skeleton lacking skull and right wing; SMF ME-10472. Figured by Mayr & Peters 1998, pl. 4, fig. 8; and Mayr 2000g, fig. 4.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr & Peters 1998).

Genus *Messelastur* Peters

Messelastur Peters, 1994: 4 [Type by original designation: *Messelastur gratulator* Peters, 1994.]

Messelastur gratulator Peters

Messelastur gratulator Peters, 1994: 4 [Holotype from Messel: skull and 15 associated cervical vertebrae; SMF ME-2024. Figured by Peters 1994, fig. 1-3 + erratum, fig. 1.]

Distribution: Middle Eocene (MN 11) of Messel, Germany (Peters 1994, Mayr 1998b).

Remarks: Here transferred from the Accipitridae.

Genus *Serudaptus* Mayr

Serudaptus Mayr, 2000h: 208 [Type by original designation: *Serudaptus pohli* Mayr, 2000h.]

Serudaptus pohli Mayr

Serudaptus pohli Mayr, 2000h: 208 [Holotype from Messel: skeleton in slab; WDC C-MG-201a,b. Figured by Mayr 2000h, fig. 1-2.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 2000h).

Remarks: Described in the Pseudasturidae Mayr (Mayr 2000h).

Genus *Eoglaucidium* Fischer

Eoglaucidium Fischer, 1987: 138 [Type by original designation: *Eoglaucidium pallas* Fischer, 1987.]

Eoglaucidium pallas Fischer

Eoglaucidium pallas Fischer, 1987: 138 [Holotype from Geiseltal XX: right humerus; GM XXII/761. Figured by Fischer 1987, fig. 1, 4.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr & Peters 1998) and Geiseltal IL, Germany (Mlíkovský, orig.); and middle Eocene (MP 13) of Geiseltal XX, XXII, XXXV, XXXVII, and XLI, Germany (Fischer 1987, Mlíkovský, orig.). General reference: Mayr & Peters (1998).

Genus *Primocolius* Mourer-Chauviré

Primocolius Mourer-Chauviré, 1988a: 37 [Type by original designation: *Primocolius sigei* Mourer-Chauviré, 1988.]

Primocolius sigei Mourer-Chauviré

Primocolius sigei Mourer-Chauviré, 1988a: 39 [Holotype from Quercy: left humerus; MNHN QU-17092. Figured by Mourer-Chauviré 1988a, pl. 1, fig. 7-8.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1988a); and late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1988a).

***Primocolius minor* Mourer-Chauviré**

Primocolius minor Mourer-Chauviré, 1988a: 39 [Holotype from Quercy: right tarsometatarsus; MNHN QU-15860. Figured by Mourer-Chauviré 1988a, pl. 2, fig. 9, and text-fig. 2c.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1988a); late Eocene (MP 17) of Perrière, France (Mourer-Chauviré 1988a); and late Eocene (MP 19) of Escamps A, France (Mourer-Chauviré 1988a).

***Primocolius* sp.**

Distribution: Late Eocene (MP 19) of Montmartre, France (Mayr 1998c: 205, 2000: 91; tentatively referred).

Genus *Quercypsitta* Mourer-Chauviré

Quercypsitta Mourer-Chauviré, 1992: 170 [Type by original designation: *Quercypsitta sudrei* Mourer-Chauviré, 1992.]

Distribution: Late Eocene (MP 17) of France.

***Quercypsitta sudrei* Mourer-Chauviré**

Quercypsitta sudrei Mourer-Chauviré, 1992: 172 [Holotype from Bouffie: left tarsometatarsus; USTL BFI-1903. Figured by Mourer-Chauviré 1992, pl. 1, fig. 1.]

Distribution: Late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1992).

***Quercypsitta ivani* Mourer-Chauviré**

Quercypsitta ivani Mourer-Chauviré, 1992: 174 [Holotype from Bouffie: left coracoid; USTL BFI-1927. Figured by Mourer-Chauviré 1992, pl. 2, fig. 1-2.]

Distribution: Late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1992).

Genus *Zygodactylus* Ballmann

Zygodactylus Ballmann, 1969a: 196 [Nomen nudum; no definition – see below.]

Zygodactylus Ballmann, 1969b: 52 [Type by monotypy: *Zygodactylus ignotus* Ballmann, 1969b.]

Distribution: Early Miocene (MN 3-4) of Germany and Czechia, and middle Miocene (MN 8) of France.

Remarks: Ballmann's 1969a paper appeared in June, while his 1969b paper was published only on 1 September. In the earlier paper, Ballmann (1969a) cited *Zygodactylus* as originating from Ballmann (1966), which is an unpublished thesis, stating that its type species is *Zygodactylus ignotus*, which was nomen nudum at that point. Hence, *Zygodactylus* Ballmann, 1969a is nomen nudum, and the name must be attributed to Ballmann, 1969b.

***Zygodactylus ignotus* Ballmann**

Zygodactylus ignotus Ballmann, 1969a: 196, 197 [Nomen nudum; no description or indication.]

Zygodactylus ignotus Ballmann, 1969b: 52 [Holotype from Wintershof-West: distal end of right tarsometatarsus; BSP Wi-We-1937-II-18164. Figured by Ballmann 1969b, pl. 2, fig. 13a-b.]

Distribution: Early Miocene (MN 3a) of Wintershof-West, Germany (Ballmann 1969b); and early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský orig.; record uncertain).

***Zygodactylus grivensis* Ballmann**

Zygodactylus grivensis Ballmann, 1969a: 197 [Holotype from Grive-Saint-Alban – fissure L-7: distal end of right tarsometatarsus; FLS, coll. Mein 151. Figured by Ballmann 1969a, pl. 14, fig. 7-9. Note, that the generic name was nomen nudum at this point.]

Distribution: Middle Miocene (MN 8) of Grive-Saint-Alban – fissure L-7, France (Ballmann 1969a).

Family Coliidae Swainson

Coliidae Swainson, 1837 [Modern family.]

Genus *Oligocolius* Mayr

Oligocolius Mayr, 2000a: 86 [Type by original designation: *Oligocolius brevitarsus* Mayr, 2000.]

***Oligocolius brevitarsus* Mayr**

Oligocolius brevitarsus Mayr, 2000a: 87 [Holotype from Frauenweiler: incomplete skeleton in slab; SMNS 80529/2. Figured by Mayr 2000a, fig. 1-2, 3c, 4a, and 4f.]

Distribution: Middle Oligocene (MP 23-24) of Frauenweiler, Germany (Mayr 2000a).

Genus *Urocolius* Bonaparte

Urocolius Bonaparte, 1854 [Modern genus.]

Limnatornis Milne-Edwards, 1871: 392 [Type by monotypy: *Limnatornis paludicola* Milne-Edwards, 1871.]

Palaeopicus Lambrecht, 1933: 629 [Nomen nudum; type not fixed after 1930.]

Palaeopicus Brodkorb, 1952: 175 [Type by original designation: *Picus archiaci* Milne-Edwards, 1871.]

***Urocolius archiaci* (Milne-Edwards)**

Limnatornis paludicola Milne-Edwards, 1871: 392 [Holotype from Saint-Gérard-le-Puy: right humerus; MNHN Av-2858. Figured by Milne-Edwards 1869-1871, pl. 176, fig. 14-18 (not pl. 176, fig. 8-13 as stated by Milne-Edwards 1871: 392). Brodkorb (1971b: 244) erroneously stated that the species was based on three syntypical humeri, NMNH Av-2858, 2859, and 2860.]

Picus archiaci Milne-Edwards, 1871: 396 [Holotype from Saint-Gérard-le-Puy: right tibiotarsus (not left as stated by Brodkorb 1971b: 244), MNHN Av-2852. Figured by Milne-Edwards 1871, pl. 178, fig. 1-5.]

Picus consobrinus Milne-Edwards, 1871: 397 [Holotype from Saint-Gérard-le-Puy: left tibiotarsus; MNHN Av-2854. Figured by Milne-Edwards 1869-1871, pl. 176, fig. 1-7.]

Palaeopicus consobrinus (Milne-Edwards): Lambrecht 1933: 629 [New combination, but the generic name was nomen nudum at this point.]

Colius archiaci (Milne-Edwards): Ballmann 1969a: 195 [New combination.]

Colius paludicola (Milne-Edwards): Ballmann 1969a: 195 [New combination.]

Urocolius archiaci (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Ballmann 1969a, Mlíkovský orig.).

Remarks: *Picus consobrinus* Milne-Edwards was synonymized with *Limnatornis paludicola* Milne-Edwards by Brodkorb (1971b: 244). The holotypical tibiotarsus of *Picus consobrinus* Milne-Edwards is similar in size and morphology with the same element of *Picus archiaci* Milne-Edwards (Mlíkovský orig.). There is no evidence, that more than one coliid species was

found in Saint-Gérard-le-Puy. The configuration of humeral head of *Limnatornis paludicola* Milne-Edwards agrees with that of *Urocolius* Bonaparte, and differs from that of *Colius* Brisson (see also Rich & Haarhoff 1985), so that I synonymize here the genus *Limnatornis* Milne-Edwards, 1871 with *Urocolius* Bonaparte, 1854.

***Urocolius* sp.**

Distribution: Late Miocene (MN 10) of Kohfidisch, Austria (Mlíkovský orig.).

Genus *Colius* Brisson

Colius Brisson, 1760 [Modern genus.]

Necrornis Milne-Edwards, 1871: 388 [Type by monotypy: *Necrornis palustris* Milne-Edwards, 1871.]

***Colius palustris* (Milne-Edwards)**

Necrornis palustris Milne-Edwards, 1871: 388 [Holotype from Sansan: distal end of right tarsometatarsus; MNHN Sa-1243. Figured by Milne-Edwards 1869-1871, pl. 178, fig. 6-10; and Cheneval 2000, fig. 24. Cheneval (2000: 369) incorrectly stated, that the holotype of this species is a lectotype, adding two irrelevant bones as "paralectotypes".]

Colius palustris (Milne-Edwards): Ballmann 1969a: 195 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871, Ballmann 1969a, Rich & Haarhoff 1985, Cheneval 2000); and middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Ballmann 1969a).

Family Psittacidae Illiger

Psittacidae Illiger, 1811 [Modern family.]

Genus *Archaeopsittacus* Lambrecht

Archaeopsittacus Lambrecht, 1933: 609 [Type by monotypy: *Psittacus verreauxi* Milne-Edwards, 1871]

Remarks: This genus belongs to the modern tribe Psittaculini sensu Smith (1975), within which its taxonomic identity has not been evaluated yet (Mlíkovský 1998e). During a preliminary study I found the holotypical tarsometatarsus of *Psittacus verreauxii* Milne-Edwards very similar to the same element of SE Asian parrots of the genera *Alisterus* Mathews, 1911 and *Aprosmictus* Gould, 1843. Both these genera are closely allied, and a further study may well show, that both *Alisterus* Mathews and *Archaeopsittacus* Lambrecht will fall to the synonymy of *Aprosmictus* Gould.

***Archaeopsittacus verreauxi* (Milne-Edwards)**

Psittacus Verreauxii Milne-Edwards, 1870: 558 [Nomen nudum; no description or indication – see Mlíkovský 1998e: 336-337.]

Psittacus Verreauxi Milne-Edwards, 1871: 525 [Holotype from Saint-Gérard-le-Puy: left tarsometatarsus; MNHN Av-2843. Figured by Milne-Edwards 1869-1871, pl. 200, fig. 1-6.]

Archaeopsittacus verreauxi (Milne-Edwards): Lambrecht 1933: 609 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1871, Mlíkovský 1998e).

Genus *Xenopsitta* Mlíkovský

Xenopsitta Mlíkovský, 1998e: 335 [Type by original designation: *Xenopsitta fejfari* Mlíkovský, 1998e.]

***Xenopsitta fejfari* Mlíkovský**

Xenopsitta fejfari Mlíkovský, 1998e: 336 [Holotype from Merkur: distal part of left humerus; coll. Fejfar, uncatalogued. Figured by Mlíkovský 1998, fig. 1a-c.]

Distribution: Early Miocene (MN 3) of Merkur, Czechia (Mlíkovský 1998e).

Remarks: *Xenopsitta fejfari* Mlíkovský belongs to the Psittacini sensu Smith 1975 (Mlíkovský 1998e).

Genus incertae sedis***Psittacus lartetianus* Milne-Edwards**

Psittacus lartetianus Milne-Edwards, 1872: 6. [Lectotype from Sansan (here selected): right tarsometatarsus lacking distal end; MNHN Sa-1668. Figured by Cheneval 2000, fig. 22. Paralectotypes from Sansan (in MNHN): left humerus (Sa-1216), two right humeri (Sa-1274, Sa-1492), distal end of left humerus (Sa-1490), left ulna (Sa-1464), distal ends of two left ulnae (Sa-1312, Sa-1314), distal end of left tibiotarsus (Sa-1606), proximal end of left tarsometatarsus (Sa-1669). Figured by Cheneval 2000, fig. 21 (tibiotarsus). Milne-Edwards (1972) did not state upon which material he created this species, but the catalogue numbers indicate, that it is the bones listed above.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1972, Cracraft 1973, Mlíkovský 1998e).

Remarks: Cheneval (2000: 363) believed, that *Psittacus lartetianus* Milne-Edwards is nomen nudum, but Milne-Edwards (1872: 6) presented a very brief and superficial description of the taxon, which is valid. See also under *Palaeoramides beaumonti* Milne-Edwards.

Family Ramphastidae Vigors

Ramphastidae Vigors, 1825 [Modern family.]

Capitonidae Bonaparte, 1838 [Modern family.]

Capitonididae Prum, 1988: 340 [Type genus: *Capitonides* Ballmann, 1969b.]

Remarks: The family Capitonidae Bonaparte was included in the Ramphastidae Vigors by Prum (1988), and the family Capitonididae Prum was synonymized with the Ramphastidae Vigors by Olson (1991: 224).

Genus *Trachyphonus* Ranzani

Trachyphonus Ranzani, 1821 [Modern genus.]

Capitonides Ballmann, 1969a: 196 [Nomen nudum; no definition.]

Capitonides Ballmann, 1969b: 43 [Type by monotypy: *Capitonides europeus* Ballmann, 1969b.]

Remarks: Ballmann (1983) showed, that *Capitonides* Ballmann is similar to *Trachyphonus* Ranzani (see also Olson 1991). Hence, I synonymize here the former with the latter genus.

***Trachyphonus europeus* (Ballmann)**

Capitonides europeus Ballmann, 1969a: 196 [Nomen nudum; no description or indication.]

Capitonides europeus Ballmann, 1969b: 44 [Holotype from Wintershof-West: right carpometacarpus; BSP Wi-We-1937-II-18162. Figured by Ballmann 1969b, pl. 2, fig. 1a-b.]

Trachyphonus europeus (Ballmann): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 3) of Wintershof-West, Germany (Ballmann 1969b, Olson 1991).

***Trachyphonus protractus* (Ballmann)**

Capitonides protractus Ballmann, 1983: 46 [Holotype from Steinberg: right tarsometatarsus;

BSP 1970-XVIII. Figured by Ballmann 1983, fig. 2: 5-7, 3: a-d.]

Trachyphonus protractus (Ballmann): Olson 1991: 224 [New combination.]

Distribution: Middle Miocene (MN 6) of Steinberg, Germany (Ballmann 1983, Olson 1991).

***Trachyphonus* sp.**

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Ballmann 1969a sub cf. *Capitonides* sp.).

Genus *Pogoniulus* Lafresnaye

Pogoniulus Lafresnaye, 1844 [Modern genus.]

***Pogoniulus* sp.**

Distribution: Late Miocene (MN 10) of Kohfidisch, Austria (Mlíkovský orig.).

Family Picidae Vigors

Picidae Vigors, 1825: 452 [Modern family.]

Genus *Jynx* Linnaeus

Jynx Linnaeus, 1758 [Modern genus.]

***Jynx torquilla* Linnaeus – Eurasian Wryneck**

Jynx Torquilla Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1980a sub *J.* cf. *torquilla*).

Genus *Picus* Linnaeus

Picus Linnaeus, 1758 [Modern genus.]

***Picus peregrinabundus* Umans'ka**

Picus peregrinabundus Umans'ka, 1981: 20 [Holotype from Novoelisavetovka: distal end of left tarsometatarsus; IZAN 43-602. Figured by Umans'ka 1981, fig. 3a,b,v,g.]

Distribution: Late Miocene (MN 11-12) of Novoelisavetovka, Ukraine (Umans'ka 1981).

Remarks: The generic position of this species should be rechecked.

***Picus canus* Gmelin – Grey-headed Woodpecker**

Picus canus Gmelin, 1788 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *P.* cf. *canus*), and Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b).

***Picus viridis* Linnaeus – European Green Woodpecker**

Picus viridis Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Somssichhegy 2, Hungary (Jánossy 1980a sub *P.* aff. *viridis*). The record from the late Oligocene (MP 30) of Aix-en-Provence, France (Bayan 1873, based on a feather imprint) is doubtful because of its age (Mlíkovský, orig.).

Genus *Picoides* Lacépède

Picoides Lacépède, 1799 [Modern genus.]

Dendrocopos Koch, 1816 [Modern genus.]

***Picoides major* (Linnaeus) – Great Spotted Woodpecker**

Picus major Linnaeus, 1758 [Modern species]

Dendrocopos major submajor Jánossy, 1974b: 234 [Lectotype from Hundsheim (here selected); UWPI 1899/66 (partim). Not figured. Paralectotypes: distal end of left tarsometatarsus from Hundsheim (UWPI 1899/66 – partim), and a tarsometatarsus from Tarkő (NMB ?). Not figured.]

Dendrocopos submajor Jánossy: Jánossy 1980a: 24 [New rank.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917); early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 5a, 9a, and 9b, Czechia (Jánossy 1972 sub *Dendrocopos* cf. *major*), and Stránská skála – Musil's talus cone, layers 13 and 15e, Czechia (Mlíkovský 1995b); and middle Pleistocene (MQ 2A) of Hundsheim, Austria (Jánossy 1974b sub *D. major submajor*), and Tarkő – layers 4 and 11, Hungary (Jánossy 1974b and 1976c sub *D. major submajor*, Jánossy 1980a sub *D. submajor*).

Remarks: I restudied the syntypes of *Dendrocopos submajor* Jánossy from Hundsheim (incl. the lectotype), finding no differences between these bones and the same elements of *Picoides major* (Linnaeus). Hence, I synonymize here the former with the latter species.

***Picoides medius* (Linnaeus) – Middle Spotted Woodpecker**

Picus medius Linnaeus, 1758 [Modern species.]

Dendrocopos praemedius Jánossy, 1974b: 237 [Holotype from Villány 3: right carpometacarpus; GIB uncatalogued. Not figured.]

Distribution: Late Pliocene (MN 17) of Obitočnoe, Ukraine (Vojinstvens'kyj 1967 sub *Dendrocopos* cf. *medius*); late Pliocene (MN 18) of Villány 3, Hungary (Jánossy 1974b sub *Dendrocopos praemedius*); and early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1980a sub *Dendrocopos* cf. *medius*), Primorsk, Ukraine (Vojinstvens'kyj 1967 sub *Dendrocopos* cf. *medius*), and Tarchankut, Ukraine (Vojinstvens'kyj 1967 sub *Dendrocopos* aff. *medius*); and middle Pleistocene (MQ 2A) of Tarkő – layers 4 and 11, Hungary (Jánossy 1974b sub *Dendrocopos praemedius*; reidentified by Mlíkovský orig.).

Remarks: Jánossy (1974b) did not present convincing evidence that *Dendrocopos praemedius* Jánossy, 1974b is different from the modern *Picoides medius* (Linnaeus). Hence, I synonymize here the former with the latter species.

Order Galliformes Temminck

Gallinae Temminck, 1820 [Modern order.]

Family Megapodiidae Swainson

Megapodiidae Swainson, 1837 [Modern family.]

Quercymegapodiidae Mourer-Chauviré, 1992b: 77 [Type genus: *Quercymegapodius* Mourer-Chauviré, 1992.]

Quercymegapodiinae Mourer-Chauviré: Mlíkovský, this paper [New rank.]

Remarks: The Quercymegapodiinae are known also from the late Oligocene or early Miocene (Tremembé Formation) of Brazil, incl. *Ameripodius* Alvarenga, 1995 and *Taubacrex* Alvarenga, 1988 (Mourer-Chauviré (1999b, 2000)).

Genus *Quercymegapodius* Mourer-Chauviré

Quercymegapodius Mourer-Chauviré, 1992b: 77 [Type by original designation: *Palaeocryptonyx depereti* Gaillard, 1908.]

Quercymegapodius depereti (Gaillard)

Palaeocryptonyx Depereti Gaillard, 1908: 96 [Holotype from Quercy: left humerus; BSP 127, apparently destroyed in the World War II – Mlíkovský, orig. Cast in FSL PQ-1065. Figured by Gaillard 1908, text-fig. 26, pl. 5, fig. 9-12]

Pirortyx depereti (Gaillard): Brodkorb 1964: 300 [New combination.]

Quercymegapodius depereti (Gaillard): Mourer-Chauviré 1992b: 81 [New combination.]

Quercymegapodius brodkorbi Mourer-Chauviré, 1992b: 82 [Holotype from Rosières 1: left carpometacarpus; USTL ROS-1-202. Figured by Mourer-Chauviré 1992b, fig. 1n-p.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1908, Mourer-Chauviré 1992b); late Eocene (MP 16) of Lavergne, France (Mourer-Chauviré 1992b); late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1992b); late Eocene (MP 18) of Sainte-Néboule, France (Mourer-Chauviré 1992b); and late Eocene (MP 19) of Escamps, France (Mourer-Chauviré 1992b), Rosières 1, and Rosières 2, France (Mourer-Chauviré 1992b).

Remarks: The holotypical carpometacarpus of *Quercymegapodius brodkorbi* was said to be morphologically identical, but slightly smaller than the same element of *Quercymegapodius depereti* (Mourer-Chauviré 1992b: 82-83). The available measurements (Tab. 2 in Mourer-Chauviré 1992b) show that the difference can be attributed to intraspecific variability. Accordingly, I synonymize here *Quercymegapodius brodkorbi* Mourer-Chauviré, 1992 with *Quercymegapodius depereti* (Gaillard, 1908).

Genus *Ameripodius* Alvarenga

Ameripodius Alvarenga, 1995. [Type by original designation: *Ameripodius silvasantosi* Alvarenga, 1995.]

Distribution: The extralimital record includes *Ameripodius silvasantosi* Alvarenga, 1995 from the late Oligocene or early Miocene (Tremembé Formation) of Taubaté, Brazil (Alvarenga 1995, Mourer-Chauviré 1999b, 2000).

Ameripodius alexis Mourer-Chauviré

Ameripodius alexis Mourer-Chauviré, 2000: 482 [Holotype from Saint-Gérard-le-Puy: left humerus, MNHN SG-9333. Figured by Mourer-Chauviré 2000, pl. 1, fig. 1-3.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Mourer-Chauviré 2000).

Family Cracidae Vigors

Cracidae Vigors, 1825 [Modern family.]

Gallinuloididae Lucas, 1900: 84 [Type genus: *Gallinuloides* Eastman, 1900.]

Gallinuloidinae Lucas: Tordoff & Macdonald 1957: 182 [New rank.]

Genus *Taoperdix* Milne-Edwards

Taoperdix Milne-Edwards, 1869: 225 [Type by monotypy: *Tetrao pessieti* Gervais, 1862.]

Distribution: Late Eocene (MP 16) of France (tentative record), middle Oligocene (MP 25) of France, and early Miocene (MN 3-4) of Germany and Czechia.

Taoperdix pessieti (Gervais)

Tetrao? Pessieti Gervais, 1862: 896 [Holotype from Armissan: incomplete skeleton in slab; MNHN 1903-16. Figured by Gervais 1859, pl. 39 (excepting figs. 7bis, and 12bis), Milne-Edwards, 1869-1871, pl. 127, and Eastman 1905, pl. 15-16. Note that the catalogue number in MNHN is the same as for *Sula arvernensis* Milne-Edwards.]

Taoperdix Pessieti (Gervais): Milne-Edwards, 1869: 225 [New combination.]

Distribution: Middle Oligocene (MP 25) of Armissan, France (Gervais 1862, Milne-Edwards 1867-1868, 1869-1871, Eastman 1905).

Taoperdix miocaena Ballmann

Taoperdix miocaena Ballmann, 1969b: 29 [Holotype from Wintershof-West: proximal end of left carpometacarpus; BSP 1937-II-18111. Figured by Ballmann 1969b, pl. 2, fig. 14a-b.]

Distribution: Early Miocene (MN 3) of Wintershof-West, Germany (Ballmann 1969b); and early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský orig.).

Taoperdix sp.

Distribution: Late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1988a, 1992b, tentatively).

Family Phasianidae Vigors

Phasianidae Vigors, 1825 [Modern family.]

Paraortygidae C. Mourer-Chauviré in Fischer, 1990: 133 [Nomen nudum; no definition. Mlíkovský (2000d: 80) erroneously considered this name valid, overlooking Art. 13.6.1. of the ICZN 1999, which invalidated Art. 12.2.4. for names published after 1930.]

Paraortygidae Mourer-Chauviré, 1992b: 70 [Type genus: *Paraortyx* Gaillard, 1908.]

Genus *Paraortyx* Gaillard

Paraortyx Gaillard, 1908: 105 [Type by subsequent designation (Richmond 1917: 612): *Paraortyx lorteti* Gaillard, 1908.]

Paraortyx lorteti Gaillard

Palaeortyx cayluxensis Milne-Edwards, 1892 [Holotype from Quercy: proximal end of right humerus; MNHN 15581. Figured by Gaillard 1908, pl. 6, fig. 1-4. This is a junior primary homonym of *Palaeortyx cayluxensis* Lydekker, 1891a.]

Paraortyx Lorteti Gaillard, 1908: 105 [Holotype from Quercy: left humerus; BSP 65, apparently destroyed during World War II – Mlíkovský, orig. Cast in MGHN PQ-1083. Figured by Gaillard 1908, text-fig. 30, pl. 6, fig. 5-8.]

Palaeortyx gaillardi Lambrecht, 1933: 451 [New name for *Palaeortyx cayluxensis* Milne-

Edwards, 1892.]

Ludiortyx gaillardii (Lambrecht): Brodkorb 1964: 299 [New combination.]

Paraortyx cayluxensis (Milne-Edwards): Cracraft 1973: 19 [New combination.]

Paraortyx gaillardii (Lambrecht): Cracraft 1973: 19 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908, Mourer-Chauviré 1992b); late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1992b) and Perrière, France (Mourer-Chauviré 1992b); late Eocene (MP 19) of Escamps, France (Mourer-Chauviré 1992b), Rosières 1, and Rosières 2, France (Mourer-Chauviré 1992b); early Oligocene (MP 21) of Ravet-Lupovici, France (Mourer-Chauviré 1992b); early Oligocene (MP 22) of Mas de Got B, France (Mourer-Chauviré 1992b); and middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1992b), and Roqueprune 2, France (Mourer-Chauviré 1992b).

Remarks: *Palaeortyx cayluxensis* Milne-Edwards was synonymized with *Paraortyx lorteti* Gaillard by Mourer-Chauviré (1992b).

***Paraortyx brancoi* Gaillard**

Paraortyx Brancoi Gaillard, 1908: 107 [Lectotype from Quercy (selected by Mourer-Chauviré 1992: 76): left humerus; NKMB, uncatalogued. Cast in MGNH PQ-1078. Figured by Gaillard 1908, text-fig. 31, pl. 6, fig. 9-12. Paralectotype from Quercy: left tarsometatarsus with damaged distal end; BSP 119, apparently destroyed during the World War II – Mlikovský, orig. Cast in ML PQ-1078. Figured by Gaillard 1908, pl. 6, fig. 13-16. Gaillard (1908) based this species on two syntypes, one of which was erroneously called by Mourer-Chauviré (1992b: 76) holotype. Nevertheless, her action can be interpreted as a selection of the lectotype.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908, Mourer-Chauviré 1992b); late Eocene (MP 18) of Sainte-Néboule, France (Mourer-Chauviré 1992b); and late Eocene (MP 19) of Escamps, France (Mourer-Chauviré 1992b), and Rosières 2, France (Mourer-Chauviré 1992b).

Genus *Pirortyx* Brodkorb

Pirortyx Brodkorb, 1964: 300 [Type by original designation: *Palaeortyx major* Gaillard, 1939.]

***Pirortyx major* (Gaillard)**

Palaeortyx major Gaillard, 1939: 22 [Holotype from Quercy: right humerus; MNHN QU-16297. Figured by Gaillard 1939, fig. 10, Mourer-Chauviré 1992b, fig. 5e-f.]

Pirortyx major (Gaillard): Brodkorb 1964: 300 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1939, Mourer-Chauviré 1992b); middle Oligocene (MP 23-24) of Espenhain, Germany (Fischer 1990); and late Oligocene (MP 28) of Fraysse (Mourer-Chauviré 1992b).

Genus *Coturnix* Bonnaterre

Coturnix Bonnaterre, 1791 [Modern genus.]

Palaeortyx Milne-Edwards, 1869: 217 [Type by original designation (Milne-Edwards 1869: 230): *Palaeortyx gallica* Milne-Edwards, 1869. Gaillard (1939: 73) and Brodkorb (1964: 298) overlooked the original designation and independently selected *Tringa? hoffmanni* Gervais, 1852 as the type species of *Palaeortyx* Milne-Edwards. Their action is invalid.]

Palaeoperdix Milne-Edwards, 1869: 245 [Type by original designation: *Palaeoperdix longipes* Milne-Edwards, 1869.]

Schaubortyx Brodkorb, 1964: 313 [Type by original designation: *Taoperdix keltica* Eastman, 1905.]

Remarks: The genera *Palaeortyx* Milne-Edwards, *Palaeoperdix* Milne-Edwards, and

Schaubortyx Brodkorb cannot be morphologically separated from the modern quails of the genus *Coturnix* Bonnaterre, with which they were not properly compared formerly.

Within the genus *Coturnix* Bonnaterre I recognize only two size classes and I identify them here with species. The size of both seems to have decreased from the early to the middle Miocene to a similar degree. The larger-bodied lineage (*Coturnix longipes*) became extinct in the middle Miocene, while the smaller-bodied lineage appears to have survived with the modern *Coturnix coturnix* (Linnaeus). Although I would be inclined to name all the forms from this lineage under the latter name, the record from the late Miocene is insufficient and inadequately documented, which opens the possibility, that the lineage was interrupted in that period. Until this problem is solved, I list here deliberately the Oligo-Miocene forms as *Coturnix gallica* Milne-Edwards, and the Plio-Pleistocene forms as *Coturnix coturnix* (Linnaeus).

***Coturnix gallica* (Milne-Edwards)**

Palaeortyx gallica Milne-Edwards, 1869: 230 [Lectotype from Saint-Gérand-le-Puy (selected by Mlíkovský 2000e: 92): right humerus; MNHN Av-2875. Figured by Milne-Edwards 1869-1871, pl. 129, fig. 25-29. Paralectotypes from Saint-Gérand-le-Puy (all in MNHN): left coracoid (Av-2873), proximal end of left humerus (Av-2876), distal end of left humerus (Av-2874), 2 right ulnae (Av-2878, Av-2879), left femur (Av-2886), right femur (Av-2887), right femur (Av-2888), right tibiotarsus (Av-2890), proximal end of right tibiotarsus (Av-2893), distal ends of 3 left tibiotarsi (Av-2889, Av-2891, Av-2892), and right tarsometatarsus (Av-2894). Figured by Milne-Edwards 1869-1871, pl. 128 (reconstructed skeleton), pl. 129, fig. 1-7 (tarsometatarsus Av-2984), 1, 8-11 (tibiotarsus Av-2890), 12-17 (femur Av-2886), 18-22 (femur Av-2888), 23-24 (ulna Av-2878), 25-29 (humerus Av-2875). The syntypical series included also at least one incomplete furcula, one scapula and one radius (see Milne-Edwards 1869-1871, pl. 128). These specimens were absent from the MNHN collections during my study of the remains in June 1999. Assuming consecutive numbering of the bones, the following seven specimens are missing from the collection: Av-2877 and 2880-2885. Of the syntypes, both preserved ulnae and the proximal end of tibiotarsus Av-2893 do not originate from a phasianid (Mlíkovský 2000e).]

Palaeortyx brevipes Milne-Edwards, 1869: 235 [Lectotype from Saint-Gérand-le-Puy (selected by Mlíkovský 2000e: 92): right humerus; MNHN Av-2898. Figured by Milne-Edwards 1869-1871, pl. 130, fig. 12-13. Paralectotypes from Saint-Gérand-le-Puy (all in MNHN): left coracoid (Av-2897), right humerus (Av-2901), proximal end of right humerus (Av-2899), shaft of right humerus (Av-2900), right ulna (Av-2902), another ulna (Av-2903 – absent from the collections in June 1999 – Mlíkovský 2000e), left femur (Av-2908), two right femora (Av-2906, Av-2907), proximal end of right femur (Av-2905), right tibiotarsus (Av-2909), another tibiotarsus (Av-2910 – absent from collections in June 1999 – Mlíkovský 2000e), and right tarsometatarsus (Av-2911). Figured by Milne-Edwards 1869-1871, pl. 130, fig. 1-7 (tarsometatarsus), 7a, 8-11 (femur Av-2906), 13a-b, 14-16 (ulna Av-2902), 17-21 (coracoid), 21a-b (tibiotarsus Av-2911). Assuming consecutive numbering of the bones, specimen Av-2904 was missing from the MNHN collections in June 1999 – Mlíkovský 2000e.]

Palaeortyx ocyptera Milne-Edwards, 1892: 71 [Holotype from Quercy: left humerus; MNHN QU-15592. Figured by Mourer-Chauviré 1992b, fig. 5l-m.]

Rallus dasypus Milne-Edwards, 1892: 73 [Lectotype from Quercy (selected by Cracraft 1973: 24): right femur; MNHN QU-15568. Figured by Cracraft 1973, fig. 10d-e, Mourer-Chauviré 1992b, fig. 10o, 11b. Paralectotype: distal end of right humerus from Quercy, MNHN QU-3071, Figured by Cracraft 1973, fig. 10b-c.]

Palaeortyx grivensis Lydekker, 1893b: 521 [Holotype from Grive-Saint-Alban: right humerus; BMNH. Figured by Lydekker 1893, pl. 41, fig. 12.]

- Quercyrallus dasyypus* (Milne-Edwards): Lambrecht 1933: 461 [New combination.]
- Taoperdix keltica* Eastman, 1905: 134 [Holotype from Armissan: incomplete skeleton in slab and counter-slab; CMP 2023 (slab) and MHN B TF-104 (counter-slab). Figured by Eastman 1905, pl. 13 (slab), and 14 (reconstruction of the skeleton), and Schaub 1945, pl. 23 (counter-slab).]
- Palaeortyx Depereti* Ennouchi, 1930: 72 [Lectotype from Grive-Saint-Alban (here selected): left humerus; ML LGr-50. Figured by Ennouchi 1930, pl. 3, fig 1-4. Paralectotype from Saint-Gérard-le-Puy: left tarsometatarsus; ML LGr-51. Figured by Ennouchi 1930, pl. 3, fig. 5-8.]
- Palaeortyx Joleaudi* Ennouchi, 1930: 76 [Holotype from Grive-Saint-Alban: right humerus; ML LGr-71. Figured by Ennouchi 1930, pl. 3, fig. 9-12.]
- Coturnix (?) miocenica* Villalta, 1963: 271 [Holotype from Mansuetos: proximal end of right carpometacarpus; coll. Villalta 4248. Figured by Villalta 1963, pl. 4, fig. 1, 1a-b.]
- Taoperdix gallica* (Milne-Edwards): Brodkorb 1964: 301 [New combination.]
- Taoperdix brevipes* (Milne-Edwards): Brodkorb 1964: 301 [New combination.]
- Schaubortyx keltica* (Eastman): Brodkorb 1964: 313 [New combination.]
- Plioperdix grivensis* (Lydekker): Brodkorb 1964: 317 [New combination.]
- Plioperdix depereti* (Ennouchi): Brodkorb 1964: 317 [New combination.]
- Plioperdix joleaudi* (Ennouchi): Brodkorb 1964: 317 [New combination.]
- Coturnix gallica* (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908 and 1939 sub *Palaeortyx* aff. *cayluxensis* Lydekker, Cracraft 1973: 24, Olson 1977: 344, Mourer-Chauviré 1992b); early Oligocene (MP 25) of Armissan, France (Eastman 1905, Schaub 1945, Brodkorb 1964: 313), and Belgarric, France (Mourer-Chauviré 1992b); late Oligocene (MP 28) of Desse, France (Mourer-Chauviré 1992b), and Fraysse, France (Mourer-Chauviré 1992b); early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Mourer-Chauviré 1992b, Mlíkovský 2000e); early Miocene (MN 3) of Wintershof-West, Germany (Ballmann 1969b), and Merkur, Czechia (Mlíkovský, orig.); early Miocene (MN 4b) of Dolnice, Czechia (Švec 1980 sub *Palaeortyx* sp., Mlíkovský orig.); middle Miocene (MN 5) of Vieux-Collonges, France (Ballmann 1972); middle Miocene (MN 6) of Petersbuch 39, Germany (Mlíkovský orig.); middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Lydekker 1893, Shufeldt 1896, Ennouchi 1930); middle Miocene (MN 8) of Grive-Saint-Alban – fissures L7 and M, France (Ballmann 1969a); late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1976b and 1993 sub *Palaeortyx* aff. *grivensis*); late Miocene (MN 11-12) of Sümeg, Hungary (Jánossy 1976b sub *P.* aff. *grivensis*); and late Miocene (MN 12) of Mansuetos, Spain (Villalta 1963), Aljezar B, Spain (Cheneval & Adrover 1993), and Tardosbánya, Hungary (Jánossy 1976b sub *Palaeortyx* aff. *grivensis*).

Remarks: *Rallus dasyypus* Milne-Edwards (described in the Rallidae) and *Palaeortyx ocyptera* Milne-Edwards were synonymized with *Palaeortyx gallica* Milne-Edwards by Mourer-Chauviré (1992b: 83). *Palaeortyx brevipes* Milne-Edwards was synonymized with *Palaeortyx gallica* Milne-Edwards by Mlíkovský (2000e). *Taoperdix keltica* Eastman was separated at the genus level by Brodkorb (1964: 313), who compared it with *Palaeortyx hoffmanni* Milne-Edwards, which later turned out to be a rail (Brunet 1970). Judging from the figures and descriptions I see no difference in size or morphology of the types of *Taoperdix keltica* Eastman, *Palaeortyx grivensis* Lydekker, *Palaeortyx depereti* Ennouchi, *Palaeortyx joleaudi* Ennouchi, *Coturnix miocenica* Villalta, and the corresponding elements of *Coturnix gallica* (Milne-Edwards). Hence, I synonymize here the former five species with the latter one.

***Coturnix longipes* (Milne-Edwards)**

Palaeoperdix longipes Milne-Edwards, 1869: 245 [Holotype from Sansan: proximal end of right tarsometatarsus; MNHN Sa-1224. Figured by Milne-Edwards 1869-1871, pl. 130, fig. 28-31; and Cheneval 2000, fig. 7.]

Palaeocryptonyx Gaillardi Ennouchi, 1930: 78 [Holotype from Grive-Saint-Alban: left (not right as stated by Ennouchi 1930: 78) humerus; ML LGr-46. Figured by Ennouchi 1930, pl. 4, fig. 1-4 (not 1-5 as stated by Ennouchi 1930 in captions to pl. 4.).

Proalector gaillardi (Ennouchi): Brodkorb 1964: 316 [New combination.]

Palaeortyx phasianoides grivensis Ballmann, 1969a: 178 [Holotype from Grive-Saint-Alban: left humerus, fissure L7; FSL coll. Mein 29. Figured by Ballmann 1969a, pl. 15, fig. 7. This is a junior primary homonym of *Palaeortyx grivensis* Lydekker, 1893b.]

Palaeortyx? intermedia Ballmann, 1969b: 33 [Holotype from Wintershof-West: left coracoid; BSP 1937-II-18103. Figured by Ballmann 1969b, pl. 1, fig. 1-2.]

Palaeortyx longipes (Milne-Edwards): Mlíkovský 2000e. [New combination.]

Coturnix longipes (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Late Oligocene (MP 28) of Desse, France (Mourer-Chauviré 1992b), and Fraysse, France (Mourer-Chauviré 1992b); early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Mourer-Chauviré 1992b, Mlíkovský 2000e); early Miocene (MN 3a) of Wintershof-West, Germany (Ballmann 1969b sub *Palaeortyx phasianoides*); early Miocene (MN 3) of Skyřice, Czechia (Mlíkovský 2000f); early Miocene (MN 4b) of Dolnice, Czechia (Švec 1980 sub *Palaeortyx* cf. *phasianoides*, Mlíkovský orig.); middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871, Cheneval 2000); middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Lydekker 1893b sub *Palaeortyx depereti*, Ennouchi 1930, Gaillard 1939, Mlíkovský orig.); middle/late Miocene (MN 8-9) of Hostalets de Pierola, Spain (Villalta & Crusafont Pairó 1950, Villalta 1963); and tentatively late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1993 sub *Palaeortyx phasianoides-intermedia*-group).

Remarks: *Palaeortyx intermedia* Ballmann was synonymized with *Palaeortyx longipes* (Milne-Edwards) by Mlíkovský (2000e). Ballmann (1969a: 182) listed *Palaeortyx longipes* Milne-Edwards in the synonymy of *Palaeortyx phasianoides* Milne-Edwards. The lectotype of the latter species belongs to the duck *Mionetta blanchardi* (Milne-Edwards) (Mlíkovský 2000e), but I agree with Ballmann, that early Miocene *Palaeortyx* quails, often reported under the name of *Palaeortyx phasianoides* Milne-Edwards, belong to the same species as those described as *Palaeortyx longipes* Milne-Edwards. Hence, I synonymize here *Palaeortyx phasianoides grivensis* Ballmann with *Coturnix longipes* (Milne-Edwards).

***Coturnix coturnix* (Linnaeus) – Common Quail**

Tetrao Coturnix Linnaeus, 1758 [Modern species.]

Scolopax ghardalamensis Fischer & Stephan, 1974: 516 [Holotype from Ghar Dalam – deer layer: right tarsometatarsus; NMM, uncatalogued. Figured by Fischer & Stephan 1974, fig. 1-2.]

Distribution: Early Pliocene (MN 14-15) of Gargano, Italy, incl. the sites of Biancone 1, Cantatore 3A, Chirò 3, 5A, 6, 11B, 14A, 20A, 24, 27, Fina D, Gervasio 2, Pirro 11A, Rinascita 1, and San Giovannino (Ballmann 1976a,b sub *Palaeortyx grivensis* and *Palaeortyx depereti*); late Pliocene (MN 18) of S'Onix, Mallorca, Spain (Sondaar et al. 1995 sub *C. cf. coturnix*);

early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), and Beremend 17, Hungary (Jánossy 1992); early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 5a, 9a, and 9b, Czechia (Jánossy 1972), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972), Stránská skála – Musil's talus cone, layers 4a, 6, 8, 9, 10a, 11, 13, 14c, 15a, 16e, and 16 (Mlíkovský 1995b), Včeláre 4E, Slovakia (Mlíkovský orig.), and Somssichhegy 2, Hungary (Jánossy 1986 sub *C. cf. coturnix*); and late Pleistocene (MQ 2C) of Ghar Dalam - deer layer, Malta (Fischer & Stephan 1974 sub *Scolopax ghardalamensis*; Olson 1976b).

Remarks: *Scolopax ghardalamensis* Fischer & Stephan was synonymized with *Coturnix coturnix* (Linnaeus) by Olson (1976b: 273).

Genus *Miogallus* Lambrecht

- Miophasianus* Lambrecht, 1933: 439 [Nomen nudum; type species not fixed after 1930.]
Miogallus Lambrecht, 1933: 442 [Type by monotypy: *Gallus longaevus* Ammon, 1918.]
Miophasianus Brodkorb 1952: 175 [Type by original designation: *Phasianus altus* Milne-Edwards, 1869.]

Miogallus altus (Milne-Edwards)

- Phasianus altus* Milne-Edwards, 1869: 239 [Lectotype from Sansan (selected by Cheneval 2000: 351): proximal end of right tarsometatarsus; MNHN Sa-1221. Figured by Milne-Edwards 1869-1871, pl. 131, fig. 27-29; and Cheneval 2000, fig. 9. Paralectotypes from Sansan: distal end of left tibiotarsus (MNHN Sa-1222), and phalanx II digiti majoris (MNHN Sa-1225). Figured by Milne-Edwards 1869-1871, pl. 131, fig. 30-33 (tibiotarsus), 34-36 (phalanx); and Cheneval 2000, fig. 10 (phalanx) and 11 (tibiotarsus).]
- Phasianus medius* Milne-Edwards, 1869: 242 [Holotype from Sansan: distal end of left tarsometatarsus; MNHN Sa-1226. Figured by Milne-Edwards 1869-1871, pl. 131, fig. 24-26; and Cheneval 2000, fig. 8.]
- Phasianus Desnoyersii* Milne-Edwards, 1869: 243 [Holotype from Touraine: left carpometacarpus; MNHN 1906-17. Figured by Milne-Edwards 1869-1871, pl. 131, fig. 37-39.]
- Ardea similis* Fraas, 1870: 49 [Holotype from Steinheim: distal end of a tibiotarsus; SMNS, uncatalogued. Figured by Fraas 1870, pl. 4, fig. 14a-c.]
- Palaeortyx maxima* Lydekker, 1893: 520 [Holotype from Grive-Saint-Alban: right coracoid; BMNH. Figured by Lydekker 1893, pl. 41, fig. 11.]
- Tantalus milne-edwardsii* Shufeldt, 1896: 513 [Holotype from Grive-Saint-Alban: proximal end of right tibiotarsus; USNM 2168. Figured by Shufeldt 1896, fig. 1.]
- Pseudotantalus Milne-Edwardsi* (Shufeldt): Sharpe. [New combination; fide Lambrecht 1921: 25.]
- Gallus longaevus* Ammon, 1918: 41 [Holotype from Dechbetten: cranial end of left coracoid; SMF, uncatalogued. Figured by Ammon 1918, fig. 8.]
- Phasianus augustus* Ammon, 1918: 45 [Holotype from Dechbetten: left femur; SMF, uncatalogued. Figured by Ammon 1918, fig. 9-10.]
- Botaurites similis* (Fraas): Lambrecht 1933: 315 [New combination.]
- Miophasianus altus* (Milne-Edwards): Lambrecht 1933: 439 [New combination.]
- Miophasianus medius* (Milne-Edwards): Lambrecht 1933: 440 [New combination, but the genus name was nomen nudum at this point.]
- Miophasianus augustus* (Ammon): Lambrecht 1933: 440 [New combination, but the genus name was nomen nudum at this point.]
- Miophasianus Desnoyersii* (Milne-Edwards): Lambrecht 1933: 440 [New combination, but the genus name was nomen nudum at this point.]
- Miogallus longaevus* (Ammon): Lambrecht 1933: 442 [New combination.]
- Proardea similis* (Ammon): Gaillard 1939: 79 [New combination.]
- Ibis milne-edwardsi* (Shufeldt): Brodkorb 1963c: 292 [New combination.]
- Miophasianus maximus* (Lydekker): Brodkorb 1964: 315 [New combination.]
- Tantalus milneedwardsii* Shufeldt: Olson 1974: 110 [Spelling emended.]
- Palaeoperdix medius* (Milne-Edwards): Cheneval 2000: 349 [New combination.]
- Miogallus altus* (Milne-Edwards): Mlíkovský, this paper [New combination.]
- Distribution:** Early Miocene (MN 3) of Can Mas, Spain (Villalta 1963); early/middle Miocene (MN 2-5) of Touraine, France (Milne-Edwards 1869); early Miocene (MN 4) of Córcoles, Spain (Alfárez et al. 1982 sub *Miophasianus* sp.); middle Miocene (MN 5) of Vieux-Collonges, France (Ballmann 1972 sub *Miophasianus* sp.); middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869, Gaillard 1939, Cheneval 2000), and Devínská Nová Ves, Slovakia (Švec 1986, Mlíkovský orig.); middle Miocene (MN 7) of Öhningen, Germany

(Lydekker 1891a: 141), Steinheim, Germany (Fraas 1870, Bocheński 1987), and Przeworno 2, Poland (Bocheński 1987); middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Depéret 1887, Lydekker 1893b, Shufeldt 1896, Regàlia 1896, Ennouchi 1930, Gaillard 1939, Olson 1974a, Bocheński 1987), and Dechbetten, Germany (Ammon 1918); middle Miocene (MN 8) of Grive-Saint-Alban – fissures L7 and M, France (Ballmann 1969a); middle/late Miocene (MN 8-9) of Hostalets de Pierola, Spain (Villalta & Crusafont Pairó 1950); and doubtfully late Miocene (MN 9) of Rudabanya, Hungary (Jánossy 1993 sub *Miophasianus* sp.). An uncertain extralimital record is from the middle Miocene (MN 5-8) of Perala, Turkey (Harrison 1985c, but see Cheneval 2000: 349).

Remarks: *Tantalus milneedwardsi* Shufeldt was synonymized with *Miogallus altus* (Milne-Edwards) by Olson (1974a: 112), *Ardea similis* Ammon by Olson (1985a: 166, tentatively), *Phasianus desnoyersi* Milne-Edwards by Švec (1986: 87), and *Phasianus augustus* Ammon by Bocheński (1987: 78, see also Švec 1986: 87). As judged from the figures and descriptions, the holotypes of *Phasianus medius* Milne-Edwards, *Palaeortyx maxima* Lydekker, and *Gallus longaevus* Ammon agree in size and morphology with the corresponding elements of *Miogallus gallus* (Milne-Edwards) (Mlíkovský orig.), and originated from localities similar in age and geographic location. In absence of the evidence to the contrary I synonymize here the former three species with the latter one.

This species seems to be common in the middle Miocene (MN 5-8) of western and central Europe, being most often found in lacustrine deposits. The alleged records from Can Mas and Córcoles are earlier, and need confirmation.

Genus *Alectoris* Kaup

Alectoris Kaup, 1829 [Modern genus.]

Tetraogallus Gray, 1832 [Modern genus.]

Ammoperdix Gould, 1851 [Modern genus.]

Palaeocryptonyx Depéret, 1892: 692 [Type by monotypy: *Palaeocryptonyx donnezani* Depéret, 1892.]

Palaeocryptonyx Depéret, 1897: 131 [Type by monotypy: *Palaeocryptonyx donnezani* Depéret, 1892; again marked as "nov. gen". This is both junior homonym and junior objective synonym of *Palaeocryptonyx* Depéret, 1892.]

Pliogallus Tugarinov, 1940b: 312 [Type by original designation: *Pliogallus coturnoides* Tugarinov, 1940b. This is a junior homonym of *Pliogallus* Gaillard, 1939.]

Plioperdix Kretzoi, 1955b: 367 [New name for *Pliogallus* Tugarinov, 1940b; hence its junior objective synonym.]

Proalector Brodkorb, 1964: 316 [Type by original designation: *Palaeortyx miocaena* Gaillard, 1939.]

Lambrehtia Jánossy, 1974a: 540 [Nomen nudum; type species not fixed after 1930 – see Mlíkovský 1992b: 458.]

Chauvireria Boev, 1997a: 141, 144 [Nomen nudum; not published in combination with an available species-group name after 1930.]

Chauvireria Boev, 1997b: 71 [Type by original designation: *Chauvireria balcanica* Boev, 1997b.]

Remarks: Pliocene and early Pleistocene deposits of Europe are rich in remains of a small phasianid, different from *Perdix* Brisson, which was assigned in SW Europe to the genus *Palaeocryptonyx* Depéret, in Central Europe to *Francolinus* Stephen, in eastern Europe to *Alectoris* Kaup, *Ammoperdix* Gould, and *Pliogallus* Tugarinov (= *Plioperdix* Kretzoi), and in Bulgaria to *Chauvireria* Boev. All these genus-group names (see below for the species-group names) seem to apply to a single taxon, for which the name *Palaeocryptonyx* Depéret is applicable. During a recent osteological revision of the modern Phasianidae I found that three of the extant genera, incl. *Alectoris* Kaup, *Ammoperdix* Gould, and *Tetraogallus* Gray, form a

group of closely related taxa, and that this group is well defined osteologically (Mlíkovský orig.). Taking into account external morphology, habits and distribution of these birds, I believe that all these genera should be united under the name *Alectoris* Kaup, which takes precedence. Understanding *Alectoris* Kaup in this broader sense, there is no reason to separate *Palaeocryptonyx* Depéret at the genus level as well. Accordingly, I synonymize here *Palaeocryptonyx* Depéret, *Pliogallus* Tugarinov, *Plioperdix* Kretzoi, and *Chauvireria* Boev with *Alectoris* Kaup.

Within the genus, the Plio-Pleistocene remains of these partridges from central and eastern Europe seemed to fall into three size classes, which I recognized as species (Mlíkovský 1995b, see also Kuročkin 1985, and Bocheński & Kuročkin 1987a for taxonomic comments). However, my subsequent research convinced me, that the observed range of variability is not sufficient for the recognition of three species-group taxa, and that there is no evidence that more than a single *Alectoris* species inhabited Europe during the Pliocene and early Pleistocene (but see below for a few records of this age attributed to the modern *Alectoris graeca* Meisner).

***Alectoris bavarica* Ballmann**

Alectoris bavarica Ballmann, 1969b: 36 [Holotype from Wintershof-West: left tarsometatarsus; BSP Wi-We-1937-II-18110. Figured by Ballmann 1969b, pl. 1, fig. 4a-b.]

Distribution: Early Miocene (MN 3) of Wintershof-West, Germany (Ballmann 1969b).

Remarks: The taxonomic position of this species requires reexamination, but my brief restudy of its holotype confirmed Ballmann's (1969b) opinion, that the bone is different from the same element of *Palaeortyx* Milne-Edwards (= *Coturnix* Bonnatere).

***Alectoris prisca* (Milne-Edwards)**

Palaeoperdix prisca Milne-Edwards, 1869: 246 [Lectotype Sansan (selected by Cheneval 2000: 345): proximal end of right tarsometatarsus; MNHN Sa-1236. Figured by Milne-Edwards 1869-1871, pl. 131, fig. 1-4; and Cheneval 2000, fig. 4. "Paralectotypes" from Sansan (all in MNHN): furcula (probably lost – Cheneval 2000: 347), two right coracoids (Sa-1239, Sa-1416), right humerus (Sa-1238), proximal end of right humerus (Sa-1297), distal end of left tibiotarsus (Sa-1237). Figured by Milne-Edwards 1869-1871, pl. 131, fig. 5-8 (tibiotarsus), 9-10 (furcula), 11-14 (humerus), 15-17 (coracoid); and Cheneval 2000, fig. 5 (coracoid Sa-1239) and 6 (tibiotarsus). The paralectotypical humerus belongs to the rail *Palaeoaramides beaumonti* (Milne-Edwards) according to Cheneval 2000: 345.]

Palaeoperdix? Sansaniensis Milne-Edwards, 1869: 249 [Holotype from Sansan: distal end of right tibiotarsus; MNHN Sa-1235. Figured by Milne-Edwards 1869-1871, pl. 131, fig. 18-23; and Cheneval 2000, fig. 6.]

Palaeocryptonyx grivensis Ennouchi, 1930: 81 [Holotype from Grive-Saint-Alban: left humerus; ML LGr-55. Figured by Ennouchi 1930, pl. 4, fig. 5-8.]

Alectoris prisca (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871); and middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Ennouchi 1930).

Remarks: The holotypes of both these species agree in morphology with the corresponding elements of *Alectoris* Kaup, they were found in both geographically and stratigraphically close deposits, and belong to the same size class. Hence, I synonymize here *Palaeocryptonyx grivensis* Ennouchi with *Palaeoperdix prisca* Milne-Edwards. *Palaeoperdix sansaniensis* Milne-Edwards was synonymized with *Alectoris prisca* (Milne-Edwards) by Cheneval (2000: 345).

***Alectoris edwardsi* (Depéret)**

Palaeortyx Edwardsi Depéret, 1887: 285 [Syntypes from Grive-Saint-Alban – early collections (all in ML): proximal end of right humerus (LGr-42), three distal ends of right humeri

(uncatalogued), distal end of left ulna (LGr-44), proximal end of right femur (uncatalogued), proximal end of left tibiotarsus (LGr-43), distal end of left tarsometatarsus (uncatalogued). Figured by Depéret 1887, pl. 13, fig. 51 (distal end of humerus), and 52 (proximal end of humerus). The syntypical tarsometatarsus belongs to *Palaeortyx longipes* Lydekker (see also Ballmann 1969a: 182).]

Palaeortyx miocaena Gaillard, 1939: 50 [Lectotype from Grive-Saint-Alban – early collections (here selected): left humerus; ML LGr-6035a. Figured by Gaillard 1939, fig. 26 (left). Paralectotypes from Grive-Saint-Alban – early collections: right humerus (ML LGr-6035b), and left tarsometatarsus (ML LGr-6035c). Figured by Gaillard 1939, fig. 26 (right humerus), and 27 (tarsometatarsus).]

Palaeoperdix edwardsi (Depéret): Brodkorb 1964: 314 [New combination.]

Palaeocryptonyx edwardsi (Depéret): Ballmann 1969a: 182 [New combination.]

Alectoris edwardsi (Depéret): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 3) of Can Mas, Spain (Villalta 1963); middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Depéret 1887, Lydekker 1893, Ennouchi 1930, Gaillard 1939); middle Miocene (MN 8) of Grive-Saint-Alban, fissure L-7, France (Ballmann 1969a); and middle/late Miocene (MN 8-9) of Hostalets de Pierola, Spain (Villalta & Crusafont Pairó 1950).

Remarks: Ballmann (1969a: 182) assigned both syntypical humeri of *Palaeortyx miocaena* Gaillard to *Palaeortyx edwardsii* Ennouchi. I agree, and I synonymize here *Palaeortyx miocaena* Gaillard with *Alectoris edwardsi* (Depéret).

***Alectoris donnezani* (Depéret)**

Palaeocryptonyx Donnezani Depéret, 1892: 131 [Syntypes from Perpignan (all in FSL): "scapula" (= amphibian ilium – Mourer-Chauviré, in litt.), right coracoid (92890), left humerus (92891-1), distal end of left ulna (92891-2), proximal end of left carpometacarpus (92891-5), right femur (92889), proximal end of left tibiotarsus (92891-4), distal end of left tibiotarsus (92892-8), and right tarsometatarsus (92888). Figured by Depéret 1997, pl. 13, fig. 2-2a (tarsometatarsus), 3-3a and 4-4a (tibiotarsus), 5-5a (femur), 6-6a (coracoid), 7-7a (scapula), 8-8a (humerus), 9-9a (ulna), 10-10a (carpometacarpus). The syntypical "scapula" is an amphibian ilium, and the carpometacarpus belongs to the Columbidae (Mourer-Chauviré in litt.).]

Francolinus Čapeki Lambrecht, 1933: 433 [Lectotype from Betfia 2 (here selected): right humerus; GIB uncatalogued. Figured by Lambrecht 1933, fig. 136a, Jurcsák & Kessler 1973, fig. 23a, and Jánossy 1976b, pl. 8, fig. 1. Paralectotypes from Betfia 2: proximal end of left scapula, four left and six right coracoids, incl. two complete ones, fragments of two furculae, fragments of three sterna, nine humeri (one of them complete), incl. three left, two right (one of them juvenile) and four abraded ones, proximal end of ulna, distal ends of three ulnae (incl. a left one), left carpometacarpus, five partial carpometacarpi, three phalanges I digiti 2 alae, and a sacrum. Figured by Lambrecht 1933, fig. 136b (right juvenile humerus, erroneously said by Lambrecht 1933: 434 to be the same bone as in his fig. 136a), 136c-d (coracoid), 136e-f (carpometacarpus), 136g (phalanx I digiti 2 alae), 136h (scapula), 136i (ulna) and 136j (furcula). All syntypes were collected by Tivadar Kormos between 1904 and 1912. The material was studied by Čapek (1917: 27, 68), Lambrecht (1933: 433-436) and Jánossy (1976b: 35-36), each of whom produced a different list of the bones. The above list includes syntypes said to be studied by Lambrecht (1933: 435). Lambrecht (1933: 433) listed among the material of *Francolinus Čapeki* also a humerus from Beremend, but it follows from a comment on p. 691 in the same book, that he did not consider it as a syntype of the species.

Ammoperdix ponticus Tugarinov, 1940b: 311 [Holotype from Odesa – catacombs: right tarsometatarsus with damaged proximal end; IZAN 6489. Figured by Tugarinov 1940b, fig.

- 2; and Bocheński & Kuročkin 1987a, pl. 18, fig. 1-2.]
- Pliogallus coturnoides* Tugarinov, 1940b: 312 [Syntypes from Odesa – catacombs: right coracoid (IZAN 6478), left and right ulna (IZAN 6483 and 6485), right radius (IZAN 6488), and two left carpometacarpi (IZAN 6479 and 6481). Figured by Tugarinov 1940b, fig. 1 (a carpometacarpus); and Dement'ev 1964, fig. 675 (same carpometacarpus).]
- Plioperdix coturnoides* (Tugarinov): Kretzoi 1955b: 305 [New combination.]
- Francolinus capeki* Lambrecht: Brodkorb 1964: 320 [Spelling emended.]
- Francolinus capeki wenzensis* Jánossy, 1974a: 540 [Syntypes from Weże 1: two fragments of premaxillae, two cranial ends of coracoids (1 left, 1 right), four distal ends of humeri (2 left, 2 right), two proximal ends of left ulnae, distal end of left ulna, proximal end of left radius, two proximal ends of carpometacarpi (1 left, 1 right), proximal end of right femur, two distal ends of right femora, proximal end of left tibiotarsus, distal end of right tibiotarsus, three proximal ends of left tarsometatarsi, five distal ends of tarsometatarsi (3 right, 2 left), four phalanges I digiti II alae; ISEZ, uncatalogued. Figured by Jánossy 1974a, pl. 23, fig. 7-8 (premaxilla), and Jánossy 1981, fig. 1-b (premaxilla).]
- Francolinus capeki villányiensis* Jánossy, 1974a: 540 [Syntypes (from Rebiełice Królewskie 1): fragment of premaxilla, fragment of mandibula, anterior fragment of two sterna, cranial ends of 14 coracoids (6 left, 8 right), sternal ends of two right coracoids, proximal ends of 12 scapulae (11 left, 1 right), left humerus, proximal ends of nine humeri (5 left, 4 right), distal ends of 15 humeri (10 left, 5 right), diaphyses of two humeri (1 left, 1 right), two left ulnae, proximal ends of eight ulnae (4 left, 4 right), distal ends of four ulnae (3 left, 1 right), proximal ends of right radius, distal ends of eight radii (5 left, 3 three), an ulnare, left carpometacarpus, proximal ends of eight carpometacarpi (4 left, 4 right), distal ends of 7 carpometacarpi (3 left, 4 right), 12 phalanges I digiti 2 alae, two phalanges II digiti 2 alae, proximal end of left femur, distal ends of two femora (1 left, 1 right), proximal end of right tibiotarsus, distal ends of 15 tibiotarsi (7 left, 7 right, and 1 juvenile), diaphyses of three tibiotarsi (2 left, 1 right), right tarsometatarsus, proximal ends of 14 tarsometatarsi (10 left, 4 right), distal ends of 13 tarsometatarsi (7 left, 6 right), diaphyses of 4 tarsometatarsi (side of body not stated), seven phalanges I digiti 1 pedis, nine phalanges I digiti 2 pedis, two phalanges II digiti 2 pedis, three phalanges I digiti 3 pedis, 10 phalanges I digiti 4 pedis, and 18 phalanges unguis; (from Beremend 1-3): left humerus (said to be from the right side of the body by Jánossy 1976b: 35, but fig. 4 in Lambrecht 1916a shows a left humerus); (from Osztramos 7): cranial ends of three left coracoids, middle trochlea of a tarsometatarsus, phalanx II digiti 2 pedis, and fragmentary phalanx I digiti 3 pedis; (from Villány 3): distal ends of 2 left humeri. Deposited in ISEZ, uncatalogued (syntypes from Rebiełice Królewskie 1) and GIB, uncatalogued (other syntypes). The syntypes from Rebiełice Królewskie 1 are figured in Jánossy 1974a, pl. 23, fig. 1 (sternum), 2 (carpometacarpus), 3 (tarsometatarsus), 4 (humerus), 5 (ulna), and 6 (premaxilla), and Jánossy 1981, fig. 3 (premaxilla). The syntypical humerus from Beremend was figured by Lambrecht 1916b, fig. 4. Jánossy (1974a) based this subspecies on a number of bones from four different localities (see Jánossy 1974a: 547), without even providing list of the material from three of them (Beremend 1-3, Osztramos 7, and Villány 3). Lists of syntypes from the latter three localities were derived from Jánossy (1976b: 35).
- Francolinus (Lambrechtia) capeki* Lambrecht: Jánossy 1974a: 540 [New combination.]
- Francolinus (Lambrechtia) minor* Jánossy, 1974a: 547 [Holotype from Rebiełice Królewskie 1: left carpometacarpus; ISEZ, uncatalogued. Not figured.]
- Francolinus subfrancolinus* Jánossy, 1976b: 36 [Holotype from Villány 3: left carpometacarpus; GIB uncatalogued. Figured by Jánossy 1976b, pl. 8, fig. 3.]
- Alectoris baryosefi* Tchernov, 1980: 25 [Holotype from 'Ubeidyia, layer II-23: left carpometacarpus; HUU 'U-4.5b. Figured by Tchernov 1980, pl. 1, fig. 3.]
- Francolinus wezensis* Jánossy: Jánossy 1981: 380 [New rank; species name misspelled.]
- Plioperdix ponticus* (Tugarinov): Kuročkin 1985: 100 [New combination.]

Plioperdix capeki (Lambrecht): Mlíkovský 1995b: 116 [New combination.]

Plioperdix subfrancolinus (Jánossy): Mlíkovský 1995b: 116 [New combination.]

Chauvireria balcanica Boev, 1997a: 144 [Nomen nudum; no description or indication.]

Chauvireria balcanica Boev, 1997b: 71 [Holotype from Väršec: right carpometacarpus; NMNHS F-47-1996. Figured by Boev 1997b, pl. 1, fig. f.]

Alectoris donnezani (Depéret): Mlíkovský, this paper [New combination.]

Distribution: Early Pliocene (MN 14) of Vojničevo, Ukraine (Vojinstens'kyj 1967 sub *Plioperdix*, Bocheňski & Kuročkin 1987a); early Pliocene (MN 15) of Perpignan, France (Depéret 1892, 1897), Weże 1, Poland (Jánossy 1974a, 1981), Ivanovce 1, Slovakia (Mlíkovský orig.), Csarnóta 2 – layers O, B/1, B/2, 2, 3, 5, 10, 15, and 20, Hungary (Jánossy 1976b), and Odesa – catacombs, Ukraine (Tugarinov 1940, Bocheňski & Kuročkin 1987a); late Pliocene (MN 16) of Rębielice Królewskie 1, Poland (Jánossy 1974a), Rębielice Królewskie 2, Poland (Jánossy 1974a), Osztramos 7, Hungary (Jánossy 1976b), Beremend 15, Hungary (Jánossy 1987b, 1990b, 1992), Etulia, Moldova (Bocheňski & Kuročkin 1987a), Çișmikioi, Moldova (Bocheňski & Kuročkin 1987a), Kotlovina, Ukraine (Bocheňski & Kuročkin 1987a); late Pliocene (MN 16-17) of Tourkobounia 1, Greece (Mourer-Chauviré 1980b, 1993b sub *Palaeocryptonyx* sp.); late Pliocene (MN 17) of Šandalja I – layer 6, Croatia (Malez-Bačić 1979 sub *Francolinus* sp.), Koliňany 2, Slovakia (Mlíkovský orig.), Beremend 11, Hungary (Jánossy 1986: 30), and Kryžanovka 1, Ukraine (Dubrovo & Kapelist 1979), and Väršec, Bulgaria (Boev 1997b); late Pliocene (MN 18) of Montoussé 5, France (Clot et al. 1976a,b), Ca Na Reia, Eivissa, Balearics (Alcover 1989 sub *Palaeocryptonyx* sp.), Kadzielnia 1, Poland (Jánossy 1974a sub *Francolinus* sp.), Villány 3, Hungary (Jánossy 1976b), Betfia 13, Romania (Kessler 1975), and Slivnica, Bulgaria (Boev 2000a); late Pliocene or early Pleistocene of Beremend 1-3, Hungary (Lambrecht 1916b sub "tyúkfél" and "Hühnervogel", Lambrecht 1933, Jánossy 1976b);

early Pleistocene (MQ 1a) of Untermassfeld, Germany (Jánossy 1997), Deutsch-Altenburg 2C, Austria (Jánossy 1981, Mlíkovský 1998d), Deutsch-Altenburg 4B, Austria (Jánossy 1981, Mlíkovský 1998d), Osztramos 2, Hungary (Jánossy 1976b), Osztramos 8, Hungary (Jánossy 1976b), Beremend 16, Hungary (Jánossy 1992), Beremend 17, Hungary (Jánossy 1992), and Betfia 2, Romania (Čapek 1917 sub "egy kihalt *Perdix*-nem" and "eine ausgestorbene *Perdix*-Gattung", Lambrecht 1933, Jánossy 1974b); early Pleistocene (MQ 1b) of Sackdilling, Germany (Heller 1930 sub "ein Hühnerartiger", Lambrecht 1933, K. Lambrecht in Brunner 1933), Chlum 6, Czechia (Mlíkovský 1996t), Koněprusy C-718, Czechia (Mlíkovský 1996t), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972, Mlíkovský orig.), Stránská skála – Musil's talus cone, layers 6, 8, 10, 11, 13 and 16, Czechia (Mlíkovský 1995b), Včeláre 4E, Slovakia (Mlíkovský orig.), Nagyharsányhegy 2, Hungary (Jánossy 1976b), Soave, Italy (Mourer-Chauviré 1980b, 1993b: 54 sub *Palaeocryptonyx* sp.), Podumci, Croatia (Malez 1973, 1986), and Razvode, Croatia (Malez 1973, 1986).

Extralimital record is available from the late Pliocene (MN 16) of Beregovaja, Buryatia (Kuročkin 1985), and early Pleistocene (MQ 1a) of 'Ubeidyia, layers II-23, and II-24, Israel (Tchernov 1980).

Remarks: I deliberately included in this list all relevant records, originally described from the Pliocene and early Pleistocene of Europe and adjacent parts of Asia as belonging to *Palaeocryptonyx* Depéret, *Francolinus* Stephen, and *Plioperdix* Kretzoi. It is possible that some of these bones belong to the modern *Perdix perdix* (Linnaeus), which was recorded from the same time-space unit (see below). No record of *Alectoris donnezani* (Depéret) is known from MQ 2. The record from the late Miocene (MN 9-10) of Sokolov, Ukraine (Vojinstvens'kyj 1967) is doubtful.

***Alectoris graeca* (Meisner) – Rock Partridge**

Perdix graeca Meisner, 1804 [Modern species.]

Alectoris graeca mediterranea Mourer-Chauviré, 1975a: 98 [Holotype from Saint-Estève-

Janson – layer B: left tarsometatarsus; coll. Bonifay CD66-B-4019. Figured by Mourer-Chauviré 1975a, pl. 10, fig. 6.]

Alectoris graeca martelensis Mourer-Chauviré, 1975a: 99 [Holotype from Fage – layer 5: right tarsometatarsus; FSL 41631. Figured by Mourer-Chauviré 1975a, pl. 10, fig. 8.]

Gallus karabachensis Burčák-Abramovič & Aliev, 1989. [Holotype from Azych – layer 5: left tarsometatarsus.]

Alectoris (graeca) mediterranea Mourer-Chauviré: Potapova 2000: 25 [New rank.]

Distribution: Early Pleistocene (MQ 1) of Monte Argentario, Italy (Cassoli 1980: 173 sub *A. g. mediterranea* and *A. g. martelensis*);

middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layers B, C, D and H, France (Mourer-Chauviré 1975a sub *A. g. mediterranea*), and Petralona – layers 2 and 8, Greece (Kretzoi 1977 and Kretzoi & Poulianos 1981 sub *A. g. mediterranea*); middle Pleistocene (MQ 2B) of Lunel-Viel – layers 1-11, France (Mourer-Chauviré 1975a sub *A. g. mediterranea*), Orgnac 3 – layers h, i, j, France (Mourer-Chauviré 1975a sub *A. g. mediterranea*), Lazaret – layers V, V1, V2, 8, 9, 11, 12, 13 and 15, France (Mourer-Chauviré 1975a sub *A. g. mediterranea*), and Fage – layers 2/3, CO, 32, 33 and 36, France (Mourer-Chauviré 1975a sub *A. g. martelensis*).

The extralimital record includes early Pleistocene (MQ 1b) of Treugolnaja Cave, Karachaevo-Cherkessia, Russia (Potapova & Baryšnikov 1993 and Baryšnikov & Potapova 1995 sub *A. g. mediterranea*), and late Pleistocene (MQ 2C) of Azych, Azerbaijan (Burčák-Abramovič & Aliev 1989 sub *Gallus karabachensis*, Baryšnikov & Potapova 1995, Potapova 2000 sub *A. mediterranea*).

Remarks: The early Pleistocene records possibly belong to *Alectoris donnezani* (Depéret). *Gallus karabachensis* Burčák-Abramovič & Aliev was synonymized with the modern *Alectoris graeca* (Meisner) by Baryšnikov & Potapova (1995: 245, see also Potapova 2000: 25).

Genus *Pavo* Linnaeus

Pavo Linnaeus, 1758 [Modern genus.]

Pavo archiaci (Gaudry)

Phasianus Archiaci Gaudry, 1862a: 631 [Syntypes from Pikermi: partial skull with attached posterior part of mandibles (lost, cast # 1), cranial end of right coracoid (# 11), left humerus (lost, cast # 3), right humerus (lost, cast # 2), right ulna (only proximal end preserved; cast of the whole bone available, # 7), proximal end of right ulna (# 8), right radius (only proximal end preserved, # 12), carpometacarpus (lost), right femur with damaged proximal end (#14), proximal end of left femur (# 16), distal end of left tibiotarsus (# 17), distal end of right tibiotarsus (# 26), and a phalanx digiti pedis (# 27); MNHN uncatalogued (the numbers correspond to the labels I left with the bones in 1992). Figured by Gaudry 1862a, pl. 16, fig. 1 (skull), 2+4 (humerus), 3+5 (ulna), and Gaudry 1862-1867, pl. 58, fig. 1-3 (skull), 4-5 (humerus), 6-7 (ulna), 8 (radius), 10 (femur), 11 (tibiotarsus), and 12 (phalanx digiti pedis).]

Gallus Aesculapii Gaudry, 1862a: 632 [Syntypes from Pikermi: associated distal end of right tibiotarsus, right tarsometatarsus and 2 partial phalanges digitorum pedis (only distal end of tibiotarsus and proximal end of tarsometatarsus preserved, cast whole, but damaged toward the distal end, # 18), distal end of right tarsometatarsus (# 19); MNHN uncatalogued (the numbers correspond to the etiquettes I left with the bones in MNHN in 1992). Figured by Gaudry 1862a, pl. 16, fig. 6-7, and Gaudry 1862-1867, pl. 58, fig. 14-15.]

Pavo aesculapi (Gaudry): Jánossy 1991: 17 [New combination.]

Pavo aesculapi phasianoides Jánossy, 1991: 17 [Holotype from Polgárdi 4: right coracoid; GIB Vt-145—V-18.094. Figured by Jánossy 1991, pl. 2, fig. 8.]

Pavo archiaci (Gaudry): Mlíkovský, this paper [New combination.]

Distribution: Late Miocene (MN 9-10) of Kolkotova Balka, Moldova (Laskarev 1908, 1912, Vojinstvens'kyj 1967, Burčak-Abramovič & Bendukidze 1971); late Miocene (MN 11-12) of Novoelisavetovka, Ukraine (Aleksiev 1916 and Vojinstvens'kyj 1967 sub *Gallus* aff. *aesculapii*); late Miocene (MN 11-13) of Belka, Ukraine (Vojinstves'kyj 1967 sub *Gallus aesculapii*), Žovten', Ukraine (Ganea & Kuročkin 1967); late Miocene (MN 12) of Pikermi, Greece (Gaudry 1862a,b, 1862-1867, Weithofer 1888 sub ?*Gallus Aesculapii*, Jánossy 1976b, Mlíkovský orig.); and late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1976b, 1991).

Remarks: I restudied the syntypes of *Phasianus archaici* Gaudry and *Gallus aesculapii* Gaudry, finding that both belong to a single species, and that the species is referable to the genus *Pavo* Linnaeus (Mlíkovský orig., see also Jánossy 1991).

***Pavo bravardi* (Gervais)**

Gallus Bravardi Gervais, 1849: 220 [Holotype from Ardé: spurred shaft of left tarsometatarsus; MNHN, probably lost – Mourer-Chauviré 1989b, 1990. Figured by Gervais 1848-1852, pl. 51, fig. 1-1a, Gervais 1859, pl. 51, fig. 1-1a, and Mourer-Chauviré 1989b, fig. 1-1a.]

Pavo moldavicus Bocheński & Kuročkin, 1987b: 89 [Holotype from Lučești: sternal end of left coracoid; PIN 2614-53. Figured by Bocheński & Kuročkin 1987b, pl. 18, fig. 13-14.]

Phasianus etuliensis Bocheński & Kuročkin, 1987b: 91 [Holotype from Etulia: cranial end of right coracoid; PIN 2614-48. Figured by Bocheński & Kuročkin 1987b, pl. 18, fig. 15-16.]

Pavo bravardi (Gervais): Mourer-Chauviré 1989b: 439 [New combination.]

Distribution: Early Pliocene (MN 14) of Osztramos 1, Hungary (Jánossy 1976b sub *Gallus aesculapii*, Mlíkovský orig.), Vojničevo, Ukraine (Vojinstevs'kyj 1967 sub *Gallus aesculapii*), and Kamenskoe, Ukraine (Vojinstvens'kyj 1967 sub *Gallus* sp.); early Pliocene (MN 15) of Perpignan, France (Depéret 1892, 1897, Mourer-Chauviré 1989b, 1990), Odesa – catacombs, Ukraine (Tugarinov 1940b and Vojinstvens'kyj 1967 sub *Gallus aesculapii*), Muselievo, Bulgaria (Boev 1998e, 1999a), and Megalo Emvolon, Greece (Boev & Koufos 2000); late Pliocene (MN 16) of Ardé, France (Gervais 1844a sub "Gallinacé", Gervais 1849, 1848-1852, 1859, Mourer-Chauviré 1989b, 1990), Etulia, Moldova (Bocheński & Kuročkin 1987b), and Lučești, Moldova (Bocheński & Kuročkin 1987b, Mourer-Chauviré 1989b, 1990); and late Pliocene (MN 17) of Saint-Vallier, France (Viret 1954 sub ?*Gallus Bravardi*, Mourer-Chauviré 1989b, 1990); and late Pliocene (MN 18) of Senèze, France (Stehlin 1923 sub ?*Gallus Bravardi*, Mourer-Chauviré 1989b, 1990).

Remarks: *Pavo moldavicus* Bocheński & Kuročkin was synonymized with *Pavo bravardi* (Gervais) by Mourer-Chauviré (1989b: 445). The holotypical coracoid of *Phasianus etuliensis* Bocheński & Kuročkin differs from the same element of the modern *Phasianus* Linnaeus in characters listed by Bocheński & Kuročkin (1987b), agreeing in them with the coracoids of *Pavo* Linnaeus, with which it was not compared formerly. It belonged to a bird of the same size class as *Pavo bravardi* (Gervais). Accordingly, I synonymize here *Phasianus etuliensis* Bocheński & Kuročkin with *Pavo bravardi* (Gervais).

Genus *Gallus* Brisson

Gallus Brisson, 1760 [Modern genus.]

Pliogallus Gaillard, 1939: 59 [Type by original designation (Gaillard 1939: 81, 93): *Pliogallus crassipes* Gaillard, 1939. This is an senior homonym of *Pliogallus* Tugarinov, 1940b.]

***Gallus gallus* f. *domestica* – Domestic Chicken**

Gallus domesticus fossilis Giebel, 1847: 23 [Nomen nudum; no description or indication.]

Pliogallus crassipes Gaillard, 1939: 59 [Holotype from "Csarnóta": right tarsometatarsus; FSL. Figured by Gaillard 1939, fig. 31.]

Pliogallus kormosi Gaillard, 1939: 61 [Holotype from "Csarnóta": left tarsometatarsus; FSL.]

Figured by Gaillard 1939, fig. 32.]

Gallus gallus levantinus Pichon & Tchernov, 1987: 202 [Holotype from Kebara: left tarso-metatarsus; HUI, uncatalogued. Figured by Tchernov 1962, pl. 4, fig. 24, pl. 5, fig. 6-7; Pichon & Tchernov 1987, text-fig. 2, pl. 1, fig. a-b.]

Distribution: Middle Pleistocene (MQ 2B) of Suard, France (Mourer-Chauviré 1975a sub *G. gallus*); late Pleistocene (MQ 2C) of Crayford, England (Harrison & Walker 1977b sub *G. gallus*), Fontéchevade, France (Mourer-Chauviré 1975a), Grappin, France (Mourer-Chauviré 1975a), Trinka 1 – middle Paleolithic layer, Moldova (Ganea 1972 sub *Gallus* sp.), and Nižnekryvčansk, Ukraine (Marisova & Tatarinov 1962, Marisova 1968 and Tatarinov & Bačinskij 1968 sub *G. gallus*);

late Pleistocene (MQ 2C, pre-Magdalenian 'Würm') of Balauzière – layer III, France (Mourer-Chauviré 1975a sub *G. gallus*, probably mixed – Mourer-Chauviré 1975a: 401), Gonvillard – layer XIV, France (Mourer-Chauviré 1975a sub *G. gallus*), Petit Puymoyen, France (Mourer-Chauviré 1975a), Soulabé – layers C1-C2, France (Mourer-Chauviré 1975a), Dürriloch, Germany (Schlosser 1900 sub *G. gallus*), Zwergloch, Germany (Ranke 1879 sub *G. gallus*), Schusterlucke, Austria (Woldfich 1893 sub *Gallus* [three unnamed forms], Mlíkovský 1997d sub *Gallus* sp.), Vindija – layers D-E, Croatia (Malez 1988 sub *G. gallus*), Sungir, Russia (Gromov 1966 sub *Gallus* sp., Burčák-Abramovič 1972 and 1975 sub *Gallus* sp., see also Sukačev et al. 1966), Bordu Mare, Romania (Jurcsák & Kessler 1988 sub *G. gallus*), Starye Durutory – upper cultural layers, Moldova (Ganea & Ketraru 1964, Ganea 1972), and Kiik-Koba Cave, Ukraine (Tugarinov 1937 sub *Gallus* sp., Vojinstvens'kyj 1967 sub *Gallus* sp., Baryšnikov & Potapova 1992 sub *Gallus* sp.);

late Pleistocene (MQ 2D; Magdalenian) of Lafaye, France (Mourer-Chauviré 1975a sub *G. gallus*), Mège, France (Capitan et al. 1906), Gourdan, France (Clot & Mourer-Chauviré 1986), Veyrier, France (Rütimeyer 1873 sub *G. gallus*?), Colombière, France (Mourer-Chauviré 1975a sub *G. gallus*), Grand Baille, France (Mourer-Chauviré 1975a), Trois Frères, France (Mourer-Chauviré 1975a), and Happurg, Germany (Hörmann 1913 sub *G. gallus*).

Extralimital records are known from the late Pleistocene (MQ 2C; late Mousterian) of Kebara – layer 345-410 cm, Israel (Tchernov 1962 sub *Phasianus hermonis*, Pichon & Tchernov 1987 sub *G. g. levantinus*); and from the late Pleistocene (MQ 2D; late Paleolithic) of Gvardžilas-Kldė, Georgia (Burčák-Abramovič 1966), Kudaro 1, Georgia (Burčák-Abramovič 1980), Mgvimevi, Georgia (Ganea & Burčák-Abramovič 1992), and Bavra, Georgia (Ganea & Burčák-Abramovič 1992); see also Burčák-Abramovič & Lakerbaj (1989), and Burčák-Abramovič (1996).

Remarks: The evidence presented above indicates, that chickens of the genus *Gallus* Brisson were present in Europe during the Riss/Würm interglacial (MQ 2C – lower part) and the lower part of the Würm glacial (MQ 2C – upper part). Nevertheless, the data do not allow to map the distribution of these chickens in that period, because some of the records are based on the material excavated and/or determined long ago, which increases the probability that the bones were assigned to incorrect strata, and/or that the relevant strata were mixed with younger ones, and/or that the bones were incorrectly identified.

On the other hand, there is no clear evidence, that *Gallus* chickens were present in Europe during the last glaciation (i.e. Würm III, which approximately corresponds to the Magdalenian), and subsequent parts of the Holocene. All the data cited above for the Magdalenian were based on the material from early excavations, while more recent excavations did not yield *Gallus* bones from this period. There are also no trustworthy records from the early Holocene (Mesolithic and Neolithic), although some were reported from Ukraine (Liberov 1959, Umans'ka 1972), Romania (Kessler 1975, 1978, 1985, Jurcsák & Kessler 1986), and Greece (see Thesing 1977, Mourer-Chauviré 1981b). It is thus probable, that *Gallus* chickens were absent from Europe during the period of latest Pleistocene (Würm III, i.e. Magdalenian), early Holocene (Preboreal and Boreal, i.e. Mesolithic), and middle Holocene (Atlantic, i.e. Neolithic).

Summarizing this evidence, it seems probable that *Gallus* chickens were present in Europe in the Riss/Würm interglacial and subsequent parts of the Würm glaciations. They probably became extinct with the onset of the last glaciation (Würm III), and were absent from Europe until the Subboreal (i.e. ca. Bronze Age), when they were recorded from sites in Czechia (Benecke 1994), Russia (near Moskva – Karchu 1990), Greece (see West & Zhou 1988), Moldova (Ganea 1972), Ukraine (Liberov 1959, Umans'ka 1972), and Romania (Jurcsák & Kessler 1973, Kessler 1993), and also in adjacent parts of Asia, incl. Armenia (Burčák-Abramovič & Mežlumjan 1986), Turkey (see West & Zhou 1988), and Syria (see West & Zhou 1988).

There is no doubt, that late Holocene chickens of Europe represent domestic forms (see e.g. Petrov 1951, Schweizer 1961, Nogalski 1975, Thesing 1977, Krupka 1978, and Mlíkovský 2002b for their osteology), but the taxonomic status of the late Pleistocene (MQ 2C) form remains unclear. Boule & Villeneuve (1927), Ganea (1965), Burčák-Abramovič & Bendukidze (1971), Burčák-Abramovič 1972 and Ganea & Burčák-Abramovič (1992) argued, that these chickens belonged to a wild, now extinct species of the genus *Gallus* Brisson. If so, the oldest available name for this taxon would be *Gallus gallus levantinus* Pichon & Tchernov. However, the holotypical tarsometatarsus of the latter subspecies differs from the same element of wild *Gallus* spp., and agrees with that of the domestic chicken – as judged from the figures – in having shaft broad and flat, and in being larger (Mlíkovský orig.; see Kuročkin & Anorova 1972 for respective characters).

The existence of a wild *Gallus* species in SE Europe was supported also by Vojinstvens'kyj (1959b), Vojinstvens'kyj & Uman'ska (1959), Marisova & Tatarinov (1965) and Ganea (1965, 1972), who believed that the species survived in this region to the late Holocene, and that it was domesticated here (see also Burčák-Abramovič & Bendukidze 1971, Burčák-Abramovič 1972, Ganea & Burčák-Abramovič 1992). This hypothesis is contradicted by the data presented above (see also Potapova 2000).

Pliogallus crassipes Gaillard and *Pliogallus kormosi* Gaillard were based on the material supposedly originating from the early Pliocene (MN 15) of Csarnóta, Hungary (Gaillard 1939). Jánossy (1976a: 13, 1976b: 38-39) observed that they were based on the remains of modern domestic chicken, and that they were apparently sent to Gaillard as a fake.

Genus *Phasianus* Linnaeus

Phasianus Linnaeus, 1758 [Modern genus.]

***Phasianus* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a).

Genus *Perdix* Brisson

Perdix Brisson, 1760 [Modern genus.]

***Perdix perdix* (Linnaeus) – Grey Partridge**

Tetrao Perdix Linnaeus, 1758 [Modern species.]

Perdix jurcsáki Kretzoi, 1962: 171 [Holotype from Betfia 5: right tarsometatarsus; RMO 1899a/3. Figured by Jurcsák & Kessler 1973, fig. 30 below.]

Perdix jurcsaki Kretzoi: Brodkorb 1964: 320 [Spelling emended.]

Perdix palaeoperdix Mourer-Chauviré, 1975a: 106 [Holotype from Fage – layer 5: right tarsometatarsus; FSL 41639. Figured by Mourer-Chauviré 1975a, pl. 10, fig. 12.]

Perdix perdix jurcsaki Kretzoi: Jánossy 1976b: 37 [New rank.]

Alectoris sutcliffei Harrison, 1980b: 96 [Holotype from Tornewton – glutton stratum: distal end of left tibiotarsus; BMNH A-4165. Figured by Harrison 1980b, fig. 1.]

Distribution: Late Pliocene (MN 17) of Kryžanovka 1, Ukraine (Dubrovo & Kapelist 1979 sub *P. cf. perdix*), and Obitočnoe, Ukraine (Vojinstvens'kyj 1967 sub *P. cf. perdix*);

early Pleistocene (MQ 1) of Betfia 7, Romania (Kessler 1975); early Pleistocene (MQ 1a) of Huéscar 1, Spain (Sánchez Marco 1989), Mas Rambault, France (Mourer-Chauviré 1975a sub *P. palaeoperdix*), Osztramos 5, Hungary (Jánossy 1976b sub *P. perdix jurcsaki*), Beremend 16, Hungary (Jánossy 1992 sub *Perdix* sp.), Beremend 17, Hungary (Jánossy 1992 sub *Perdix* sp.), Betfia 2, Romania (Čapek 1917, Jánossy 1976b sub *P. perdix jurcsaki*), Primorsk, Ukraine (Vojinstvens'kyj 1967), and Ževachova Hill, Ukraine (Dubrovo & Kapelist 1979); early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), Sima del Elefante – unknown layer, Spain (Rosas et al. sub *P. paleoperdix*), Stránská skála – Čapek's sites 1-3, 5a, 9a, and 9b, Czechia (Jánossy 1972 sub *P. aff. perdix*), and Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *P. aff. perdix*), Stránská skála – Musil's talus cone, layers 5, and 6 (Mlíkovský 1995b), and Betfia 5, Romania (Kretzoi 1962, Jurcsák & Kessler 1973, Kessler 1975);

middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layers B, C, D, F, G and H, France (Mourer-Chauviré 1975a sub *P. palaeoperdix*), Vertesszőlős 2 – layer 200-240 cm, Hungary (Jánossy 1976b sub *P. perdix jurcsaki*), Várhegy – Fortuna Street 25, lower layer, Hungary (Jánossy 1976b, tentatively referred, Jánossy 1986 sub *P. cf. jurcsaki*), and Petralona, Greece (Kretzoi 1977 and Kretzoi & Poulianos 1981 sub *P. cf. jurcsaki*); middle Pleistocene (MQ 2B) of Tornewton, England (Harrison 1980b sub *Alectoris sutcliffei*, Stewart 1996, J.R. Stewart in Tyrberg 1998: 479), Atapuerca, Spain (Sánchez Marco 1987a,b), Aridos-1, Spain (Mourer-Chauviré 1980c sub *P. palaeoperdix*), Lunel-Viel – layers 2, 6, 7, 9 and 10, France (Mourer-Chauviré 1975 sub *P. palaeoperdix*), Fage – layers 2/3, CO and 29-40, France (Mourer-Chauviré 1975a,b sub *P. palaeoperdix*), Orgnac 3 – layers d-j, France (Mourer-Chauviré 1975 sub *P. palaeoperdix*), Lazaret 8, France (Mourer-Chauviré 1975a sub *P. palaeoperdix*), Biehle, France (Mourer-Chauviré 1975a,b sub *P. palaeoperdix*), Suard – layers V/VI and VIII, France (Mourer-Chauviré 1975 sub *P. cf. palaeoperdix*), Aldène, France (Mourer-Chauviré 1975a sub *P. palaeoperdix*), Carrière, France (Mourer-Chauviré 1975 sub *P. palaeoperdix*), Combe Grenal, France (Mourer-Chauviré 1975 sub *P. palaeoperdix*), and Azé 2, France (Mourer-Chauviré 1975a sub *P. palaeoperdix*), and Devetaška Cave, Bulgaria (Boev 1999a sub *P. palaeoperdix*); and

late Pleistocene (MQ 2D) of Temnata Cave, Bulgaria (Boev 1999a sub *P. cf. palaeoperdix*), Kozarnika, Bulgaria (Boev 1999a sub *P. palaeoperdix*), Ražiška Cave, Bulgaria (Boev 1999a sub *P. palaeoperdix*, Boev 2000g sub *P. cf. palaeoperdix*), and Cave 16, Bulgaria (Boev 1999a sub *P. palaeoperdix*).

An extralimital record was reported from the early Pleistocene (MQ 1b) of Dursunglu, Turkey (Louchart et al. 1998 sub *P. palaeoperdix*).

Remarks: *Perdix palaeoperdix* Mourer-Chauviré was described as a form ancestral to the modern *Perdix perdix* (Linnaeus). I synonymize it here with the latter species. *Perdix jurcsaki* Kretzoi was synonymized with *Perdix perdix* (Linnaeus) by Jánossy (1976b: 37). *Alectoris sutcliffei* Harrison was synonymized with *Perdix perdix* (Linnaeus) by Stewart (1996, Stewart in Tyrberg 1998: 519).

Genus *Tetrao* Linnaeus

Tetrao Linnaeus, 1758 [Modern genus.]

Tetrao urogallus Linnaeus – Western Capercaillie

Tetrao urogallus Linnaeus, 1758 [Modern species.]

Tetrao praeurogallus Jánossy, 1969: 596 [Holotype from Včeláre 1: right carpometacarpus; GIB V-10.347. Figured by Jánossy 1969, pl. 1, fig. 4, Jánossy 1976a, pl. 1, fig. 1-2.]

Tetrao conjugens Jánossy, 1974a: 537 [Holotype from Węże 1: proximal end of left carpometacarpus; ISEZ, uncatalogued. Figured by Jánossy 1974a, pl. 23, fig. 9.]

Tetrao macropus Jánossy, 1976a: 17 [Holotype from Csarnóta 2: distal end of left tibiotarsus;

GIB V-10.348. Figured by Jánossy 1976a, pl. 1, fig. 6.]

Tetrao rhodopensis Boev, 1998d: 56 [Holotype from Dorkovo: distal end of left humerus; NMNHS 483. Figured by Boev 1998d, fig. 2c-d.]

Distribution: Early Pliocene (MN 14) of Dorkovo, Bulgaria (Boev 1998d); early Pliocene (MN 15) of Weże 1, Poland (Jánossy 1974a), and Csarnóta 2, Hungary (Jánossy 1976a); late Pliocene (MN 16) of Rebiełice Królewskie 1, Poland (Bocheński 1989, 1991, 1993, see also Jánossy 1976a), and Osztramos 7, Hungary (Jánossy 1976a); late Pliocene (MN 18) of Kadzielnia 1, Poland (Bocheński 1989);

early Pleistocene (MQ 1a) of Żabia Cave, Poland (Bosák et al. 1982: 221); early Pleistocene (MQ 1b) of Sackdilling, Germany (Jánossy 1976a), Erpfingen, Germany (Jánossy 1976a), Holštejn 1, Czechia (Musil 1966 sub *T. cf. urogallus*), Stránská skála – Čapek's sites 1-3, 4c, 5a, and 9a, Czechia (Jánossy 1972), Stránská skála – Musil's talus cone, layers 11, and 13, Czechia (Mlíkovský 1995b), Zalesiaki 1A, Poland (Bocheński 1989), Kozi Grzbiet, Poland (Bocheński 1984 sub *T. cf. praeurogallus*), Včeláre 1, Slovakia (Jánossy 1969, 1976a), Nagyarsányhegy 1-4, Hungary (Jánossy 1976a), and Betfia 5, Romania (Kretzoi 1962, Jánossy 1976a); and middle Pleistocene (MQ 2A) of Tarkő – layer 10, Hungary (Jánossy 1976a sub *T. praeurogallus*).

Remarks: *Tetrao praeurogallus* Jánossy, *Tetrao conjugens* Jánossy, *Tetrao macropus* Jánossy, and *Tetrao rhodopensis* Boev are inseparable from the modern *Tetrao urogallus* Linnaeus (Mlíkovský orig.). Hence, I synonymize here the former four species with the latter one.

***Tetrao tetrrix* Linnaeus – Black Grouse**

Tetrao Tetrrix Linnaeus, 1758 [Modern species.]

Lyrurus partium Kretzoi 1962: 171 [Holotype from Betfia 2: right tarsometatarsus, NMB G-50.155/1. Not figured.]

Lyrurus tetrrix longipes Mourer-Chauviré 1975a: 94 [Holotype from Fage – layer 5: left tarsometatarsus, FSL 41617. Figured by Mourer-Chauviré 1975, pl. 10, fig. 3.]

Tetrao partium: Brodkorb 1964: 322 [New combination.]

Lagopus balcanicus Boev 1995a: 263 [Holotype from Väršec: left ulna; MNHNS F-1-1993. Figured by Boev 1995a, fig. 3.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1995a, 1999d); late Pliocene (MN 18) of Senéze, France (Stehlin 1923 sub *?Lyrurus*), Villány 3, Hungary (Jánossy 1976a);

early Pleistocene (MQ 1a) of Deutsch-Altenburg 2C, Austria (Mlíkovský 1998d), Beremend 16, Hungary (Jánossy 1992), Beremend 17, Hungary (Jánossy 1992), Osztramos 2, Hungary (Jánossy 1976a), Osztramos 8, Hungary (Jánossy 1976a), Betfia 1, Romania (Jánossy 1976a), Betfia 2, Romania (Kretzoi 1941 sub *Tetraonidarum* g.ind. n.sp., Kretzoi 1962, Jánossy 1976a); early Pleistocene (MQ 1b) of Sackdilling, Germany (Brunner 1933), Erpfingen, Germany (Jánossy 1976a), Voigtstedt, Germany (Jánossy 1965, 1976a), Chlum 6, Czechia (Mlíkovský 1996t), Stránská skála – Čapek's sites 4a, 4c, 5a, 8, 9a, and 9b, Czechia (Jánossy 1972), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972), Stránská skála – Musil's "new large cave", Czechia (Jánossy 1972), and Stránská skála – Musil's talus cone, layers 4, 4a, 5, 6, 7, 7b, 8, 10, 11, 13, 14, 14c, 15c, and 15e, Czechia (Mlíkovský 1995b), Včeláre 1, Slovakia (Jánossy 1976a), Včeláre 4E, Slovakia (Mlíkovský orig.), Nagyarsányhegy 1-4, Hungary (Jánossy 1976a), and Somssichhegy 2, Hungary (Jánossy 1986);

middle Pleistocene (MQ 2A) of Hundsheim, Austria (Freudenberg 1914 sub *Lyrurus* sp., Jánossy 1974b sub *Lyrurus* cf. *partium*, Mlíkovský orig.), Gombasek, Slovakia (Jánossy 1976a sub *Lyrurus partium*), Tarkő – layers 2, 3, 7, 12, Hungary (Jánossy 1976a sub *Lyrurus partium*), Várhegy – Fortuna Street 25, Hungary (Jánossy 1976a sub *Lyrurus partium*), Vertesszőllős 2, Hungary (Jánossy 1976a and 1990 sub *Lyrurus partium*), and Čertkov 2, Ukraine (Vojinstvens'kyj 1967, Jánossy 1976a sub *Lyrurus partium*); and middle Pleistocene (MQ 2B) of Fage – layers 2/3, CO, 30, 32-40, France (Mourer-Chauviré 1975a sub *T. tetrrix longipes*).

Remarks: *Tetrao partium* (Kretzoi) is a member of the *Tetrao tetrrix* – lineage, and I synonymize it here thus with *Tetrao tetrrix* Linnaeus. The holotypical ulna of *Lagopus balcanicus* Boev differs from the same element of *Lagopus* Brisson, and agrees with that of *Tetrao* Linnaeus (as judged from the figure) in having (1) ventral condyle more protruding distally, and (2) ventral condyle displaced more ventrally, which causes that the dorsal condyle is relatively broad. Within the genus *Tetrao* Linnaeus, *balcanicus* is slightly smaller than the modern *Tetrao tetrrix* Linnaeus, falling thus in the size class of the alleged species *Tetrao partium* (Kretzoi). Hence, I synonymize here *Lagopus balcanicus* Boev with the modern *Tetrao tetrrix* Linnaeus.

The Pliocene and early Pleistocene *Tetrao tetrrix* Linnaeus tended to be smaller than the modern European Black Grouses (Mlíkovský, orig.).

Genus *Bonasa* Stephen

Bonasa Stephen, 1810 [Modern genus.]

***Bonasa bonasia* (Linnaeus) – Hazel Grouse**

Tetrao Bonasia Linnaeus, 1758 [Modern species.]

Lagopus voinstvenskii Ganea, 1972: 29 [Lectotype from Brynzeny 1 – lower cultural layer (here selected. This is the only figured syntype, on the basis of which I determined the identity of the species.): right tarsometatarsus; IZC BRY-408. Figured by Ganea 1969, fig. 2 – center; and Ganea 1972, fig. 1 – center. Paralectotypes: 26 tarsometatarsi from Brynzeny 1 – lower cultural layer, Starye Duruitory – layers II, III-IV, Butești – cultural layer, and Trinka 1, and six other unspecified bones from some or all of these sites (see Ganea 1969: 128-129, 1972: 29-31). Not figured.]

Tetrastes praebonasia Jánossy, 1974b: 218 [Holotype from Tarkő – layer 12: right tarsometatarsus; NMB V-64.789. Not figured.]

Bonasa praebonasia (Jánossy): Bocheński 1991: 574 [New combination.]

Distribution: Early Pleistocene (MQ 1b) of Kozi Grzbiet, Poland (Bocheński 1984), and Stránská skála – Musil's talus cone, layers 8, 9 and 13, Czechia (Mlíkovský 1995b); middle Pleistocene (MQ 2A) of Montoussé 3, France (Clot et al. 1976a,b), Hundsheim, Austria (Freudenberg 1914 sub *?Perdix cinerea*, Jánossy 1974b sub *Tetrastes praebonasia*), and Tarkő – layers 11, and 12, Hungary (Jánossy 1976a sub *Tetrastes praebonasia*). The alleged record from the early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992) is based on a humerus of *Alectoris donnezani* (Mlíkovský, orig.).

Remarks: *Tetrastes praebonasia* Jánossy was described as a chronospecies, directly ancestral to the modern *Bonasa bonasia* (Linnaeus). Hence I synonymize it here with the latter species.

Ganea (1972) named *Lagopus voinstvenskii* on the basis of 33 bones, incl. 27 tarsometatarsi from the late Pleistocene (MQ 2C) of the following four Moldovan localities: Brynzeny 1 – lower cultural layer (Ganea 1969 sub *Lagopus* sp., Ganea 1972), Starye Duruitory – layers II, III-IV (Ganea 1969 sub *Lagopus* sp., Ganea 1972), Butești – cultural layer (Ganea 1972), and Trinka 1 (Ganea 1972). The description of the tarsometatarsi, their measurements and the published photograph of the lectotype (here selected) clearly indicate that this species is identical with the modern *Bonasa bonasia* (Linnaeus), with which I synonymize it here.

Genus *Lagopus* Brisson

Lagopus Brisson, 1760 [Modern genus.]

Remarks: Past distribution of *Lagopus* spp. in Europe was mapped by Nehring (1883), and Tyrberg (1991b, 1995). Osteological variability of fossil populations of *Lagopus* spp. was studied particularly by Bocheński (1974: 138-154, 1985), Mourer-Chauviré (1975a: 85-94), and Stewart (1999a,b). General reference: Potapov 1988.

***Lagopus lagopus* (Linnaeus) – Willow Ptarmigan**

Tetrao Lagopus Linnaeus, 1758 [Modern species.]

Lagopus medius Woldřich, 1893: 621 [Lectotype from Schusterlucke (selected by Mlíkovský 1997d: 109): right tarsometatarsus; NHMW 468. Figured by Woldřich 1893, pl. 6, figs. 30. Paralectotype from Schusterlucke: right tarsometatarsus, NHMW 489. Figured by Woldřich 1893, pl. 6, fig. 31.]

Lagopus lagopus atavus Jánossy 1974a: 534 [Lectotype from Rebiełice Królewskie 1 (selected by Bocheński 1991: 570): proximal end of right humerus, ISEZ AF/26-RK1:22. Figured by Jánossy 1974a, pl. 24, fig. 6. Paralectotypes from Rebiełice Królewskie 1 (listed after Jánossy 1974a: 534, and Bocheński 1991): two partial mandibulae, two cranial ends of left coracoids, three cranial ends of right coracoids, two diaphyses of right coracoids, three proximal ends of left scapulae, two proximal ends of right humeri, distal end of left humerus, proximal end of left ulna, two proximal ends of right ulnae, distal end of left ulna, distal end of right ulna, distal end of right radius, ulnare, left carpometacarpus, right carpometacarpus, proximal end of right carpometacarpus, three phalanges alae, proximal end of left femur, distal end of right tibiotarsus, 2 tarsometatarsi, and phalanx digiti pedis; ISEZ AF/26-RK1. Figured by Jánossy 1974a, pl. 24, fig. 4 (mandible), 5 (femur), 7 (carpometacarpus), and 8 (tarsometatarsus).]

Lagopus lagopus noaillensis Mourer-Chauviré 1975a: 86 [Holotype from Fage – layer 5: right tarsometatarsus, FSL 41594. Figured by Mourer-Chauviré 1975a, pl. 10, fig. 1.]

Lagopus atavus Jánossy: Jánossy 1976a: 33 [New rank.]

Distribution: Early Pliocene (MN 15) of Weże 1, Poland (Jánossy 1974a sub *L. lagopus atavus*), and Muselievo, Bulgaria (Boev 1998e sub *Lagopus* sp.); late Pliocene (MN 16) of Rebiełice Królewskie 1, Poland (Jánossy 1974a sub *L. lagopus atavus*, Bocheński 1991 sub *L. atavus*); late Pliocene (MN 18) of Kielniki 3B, Poland (Jánossy 1974a sub *L. lagopus atavus*, Bocheński 1991 sub *L. atavus*);

early Pleistocene (MQ 1a) of Valerots, France (Mourer-Chauviré 1975a sub *L. lagopus*); early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 1-3, 4a, 4c, 5a, 5b, 9a, Czechia (Jánossy 1972, Mlíkovský orig.), and Stránská skála – Absolon's 2nd cave (Jánossy 1972 sub *L. lagopus*, Mlíkovský orig.); middle Pleistocene (MQ 2B) of Fage – layers CO, 34 and 40, France (Mourer-Chauviré 1975a sub *L. lagopus noaillensis*); and late Pleistocene (MQ 2C) of Schusterlucke, Austria (Woldřich 1893 sub *Lagopus medius*, Mlíkovský 1997d: 109).

Remarks: *Lagopus medius* Woldřich was synonymized with *Lagopus lagopus* (Linnaeus) by Mlíkovský (1997d: 109).

***Lagopus mutus* (Montin) – Rock Ptarmigan**

Tetrao mutus Montin, 1781 [Modern species.]

Lagopus mutus corzezensis Mourer-Chauviré 1975b: 87 [Holotype from Fage – layer 5: right tarsometatarsus, FSL 41601. Figured by Mourer-Chauviré 1975, pl. 10, fig. 2.]

Distribution: Middle Pleistocene (MQ 2B) of Fage – layers 4, CO, 33, 37, 39 and 40, France (Mourer-Chauviré 1975a sub *Lagopus mutus corzezensis*).

Remarks: This species was very abundant in late Pleistocene deposits of Europe, but it has no record earlier than MQ 2B, when it was reported from Westbury-sub-Mendip, England (Andrews 1990), Carrière, France (Mourer-Chauviré 1975a), Fage, France (see under Distribution), Hunas, Germany (Jánossy 1983b), and Hórvölgy, Hungary (Jánossy 1976a).

Genus incertae sedis

***Alectoris pliocaena* Tugarinov**

Alectoris pliocaena Tugarinov, 1940b: 311 [Syntypes from Odesa – catacombs: proximal end of left carpometacarpus (IZAN 6940), and distal end of right ulna (IZAN 6941). Figured by Tugarinov 1940b, fig. 1a (carpometacarpus), 1b (ulna); and Dement'ev 1964, fig. 676a (carpometacarpus), 676b (ulna).]

Distribution: Early Pliocene (MN 15) of Odesa – catacombs, Ukraine (Tugarinov 1940b).

Remarks: Here transferred to the Phasianidae incertae sedis. The syntypes are too large for *Alectoris donnezani* (Depéret).

***Gallus beremendensis* Jánossy**

Gallus beremendensis Jánossy, 1976b: 34 [Holotype from Beremend 5: right humerus; GIB, uncatalogued. Figured by Jánossy 1976b, pl. 8, fig. 5.]

Distribution: Late Pliocene (MN 16a) of Beremend 5, Hungary (Kretzoi 1956 sub "Phasianidae indet.", Jánossy 1976b).

Remarks: Jánossy (1976b) included this species in the genus *Gallus* Brisson without explanation, with the diagnosis "smaller than the hitherto known recent [sic!] or fossil forms of the genus". The taxonomic status of the species requires reexamination, because the holotype seems to differ from the corresponding element of *Gallus* in the shape of shaft, head, and bicipital crest (Mlíkovský orig.).

***Gallus europaeus* Harrison**

Gallus europaeus Harrison, 1978: 375 [Holotype from Ostend: left coracoid; BMNH 18239. Not figured.]

Distribution: Middle Pleistocene (MQ 2A) of Ostend, England (Lydekker 1891a sub *Tetrao urogallus*, Harrison 1978, 1979d, 1985b).

Remarks: The taxonomic status of this species should be reevaluated. There is no evidence, that it belongs to the genus *Gallus* Brisson.

***Gallus moldovicus* Burčák-Abramovič et al.**

Gallus moldovicus Burčák-Abramovič, Ganea & Šušpanov, 1993: 45.

Distribution: Late Pliocene of Moldova (Burčák-Abramovič et al. 1993).

Remarks: Not seen.

***Palaeocryptonyx hungaricus* Jánossy**

Palaeocryptonyx hungaricus Jánossy, 1991: 14 [Holotype from Polgárdi 5: left humerus; GIB Vt-144-V-18.093. Figured by Jánossy 1991, pl. 2, fig. 6.]

Distribution: Late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991), and Polgárdi 5, Hungary (Jánossy 1991).

Remarks: Ratio of the length of humerus and tarsometatarsus of *Palaeocryptonyx hungaricus* Jánossy is different from that of "*Palaeocryptonyx*" partridges, the tarsometatarsi being relatively shorter in *hungaricus* (Mlíkovský orig.).

***Paraortygoides messelensis* Mayr**

Paraortygoides messelensis Mayr, 2000c: 48 [Holotype from Messel: almost complete skeleton in slab; SMF ME-1303a,b. Figured by Mayr 2000c, fig. 1, 2, 4, 7 and 8; and Mayr 2000g, fig. 9.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 2000c).

Remarks: *Paraortygoides messelensis* Mayr is type by original designation of the genus *Paraortygoides* Mayr, 2000c: 47.

Order Accipitriformes Vieillot

Accipitres Vieillot, 1816 [Modern order.]

Family Rallidae Vigors

Rallidae Vigors, 1825 [Modern family.]

Remarks: This list of the Rallidae is based mainly on previous lists by Cracraft (1973) and Olson (1977). It is certain, however, that the list presented below does not reflect true taxonomic structure of the family Rallidae at the genus and species levels, and even the allocation of some Eocene forms to this family is questionable.

Genus *Parvirallus* Harrison & Walker

Parvirallus Harrison & Walker, 1979a: 23 [Type by original designation: *Parvirallus gracilis* Harrison & Walker, 1979a.]

Parvirallus bassetti Harrison

Parvirallus bassetti Harrison, 1984a: 185 [Holotype from Sheppey: distal end of right tibiotarsus; BMNH A-5282. Figured by Harrison 1984a, fig. 3a, 4b,c,e.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison 1984a).

Parvirallus medius Harrison

Parvirallus medius Harrison, 1984a: 185 [Holotype from Warden Point: distal end of right tibiotarsus; BMNH A-3546. Figured by Harrison 1984a, fig. 3b, 4a,d,f.]

Distribution: Early Eocene (MP 8-9) of Warden Point, Isle of Sheppey, England (Harrison 1984a).

Parvirallus gassoni Harrison

Parvirallus gassoni Harrison, 1984a: 185 [Holotype from Sheppey: distal end of right tibiotarsus; BMNH A-5283. Figured by Harrison 1984a, fig. 3c, 4g.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison 1984a).

Parvirallus gracilis Harrison & Walker

Parvirallus gracilis Harrison & Walker, 1979a: 23 [Holotype humerus from Lee-on-Solent: distal end of left; BMNH A-4280. Figured by Harrison & Walker 1979a, pl. 1, row 5 partim.]

Distribution: Middle Eocene (MP 11-13) of Lee-on-Solent, England (Harrison & Walker 1979a).

Genus *Latipons* Harrison & Walker

Latipons Harrison & Walker, 1979a: 24 [Type by original designation: *Latipons gardneri* Harrison & Walker, 1979a.]

Latipons gardneri Harrison & Walker

Latipons gardneri Harrison & Walker, 1979a: 24 [Holotype from Lee-on-Solent: distal end of right tibiotarsus; BMNH A-5009. Figured by Harrison & Walker 1979a, pl. 1, row 3 partim, and row 4 partim.]

Distribution: Middle Eocene (MP 11-13) of Lee-on-Solent, England (Harrison & Walker 1979a).

***Latipons robinsoni* Harrison & Walker**

Latipons robinsoni Harrison & Walker, 1979a: 25 [Holotype from Lee-on-Solent: distal end of left tibiotarsus; BMNH A-5011. Figured by Harrison & Walker 1979a, pl. 1, row 5 partim.]

Distribution: Middle Eocene (MP 11-13) of Lee-on-Solent, England (Harrison & Walker 1979a).

Genus *Ibidopsis* Lydekker

Ibidopsis Lydekker, 1891a: 74 [Type by monotypy: *Ibidopsis hordwelliensis* Lydekker, 1891a.]

Remarks: Lydekker (1891a) described this genus in the Plataleidae (= Threskiornithidae), but Cracraft (1973: 17) and Harrison & Walker (1976a: 342) transferred it to the Rallidae (see also Olson 1977).

***Ibidopsis hordwelliensis* Lydekker**

Ibidopsis hordwelliensis Lydekker, 1891a: 74 [Holotype from Hordle: distal end of right tibiotarsus; BMNH 36793. Figured by Lydekker 1891a, fig. 20, Cracraft 1973, fig. 6a-d, Harrison & Walker 1976a, pl. 6, fig. a-e.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a, Cracraft 1973: 18, Harrison & Walker 1976a: 342).

Genus *Ludiortyx* Brodkorb

Ludiortyx Brodkorb, 1964: 298 [Type by original designation: *Palaeortyx blanchardi* Milne-Edwards, 1869.]

Eortyx Brodkorb, 1967: 111 [Type by original designation: *Tringa? hoffmanni* Gervais, 1852.]

***Ludiortyx hoffmanni* (Gervais)**

Tringa? hoffmanni Gervais, 1852: 229 [Holotype from Montmartre: almost complete skeleton in two slabs; MNHN 7996 (not found in the collections in June 1999 – Mlíkovský, orig.). Figured by Gervais 1848-1852, pl. 49, fig. 4, Milne-Edwards 1869-1871, pl. 124, Zittel 1890, fig. 718, Brunet 1970, pl. D, Cracraft 1973, fig. 7.]

Palaeortyx hoffmanni (Gervais): Milne-Edwards 1869: 217 [New combination.]

Palaeortyx blanchardi Milne-Edwards, 1869: 223 [Holotype from Montmartre: associated skull, cervical vertebrae, scapula, humerus, ulna, radius, carpometacarpus and phalanx 1 digiti II in a slab; MNHN 7994. Figured by Milne-Edwards 1869-1871, pl. 126, fig. 1, Brunet 1970, pl. C below, and Cracraft 1973, fig. 8a-b.]

Ludiortyx blanchardi (Milne-Edwards): Brodkorb 1964: 298 [New combination.]

Eortyx hoffmanni (Gervais): Brodkorb 1967: 111 [New combination.]

Ludiortyx hoffmanni (Gervais): Brunet 1970: 33 [New combination.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Gervais 1848-1852, Bonaparte 1856: 1021 sub "Etourneau", Milne-Edwards 1869-1871, Brunet 1970, Cracraft 1973, Olson 1977).

Remarks: Originally described as a sandpiper (*hoffmanni*) and as a quail (*blanchardi*), the two species are conspecific, representing a rail (Brunet 1970: 33, Cracraft 1973: 19, Olson 1977: 343).

Genus *Montirallus* Mlíkovský

Montirallus Mlíkovský, 1981: 342 [Type by original designation: *Numenius gypsorum* Gervais, 1848. – See Mlíkovský 1995a: 313 for the citation of the type species.]

***Montirallus gypsorum* (Gervais)**

Numenius gypsorum Gervais, 1844a: 18 [Nomen nudum; no description or indication.]

Numenius gypsorum Gervais, 1844b: 294 [Nomen nudum; no description or indication.]

Numenius gypsorum Gervais, 1852: 230 [Lectotype from Montmartre (selected by Mlíkovský 1995a: 313): partial skull imprint showing brain morphology in slab; MNHN 7992 partim. Figured by Gervais 1848-1852, pl. 49, fig. 2, and Dechaseaux 1970b, fig. 1, upper row. Paralectotypes from Montmartre: lateral sternum imprint in slab (MNHN 7920), and associated partial skull imprint and several cervical vertebrae (MNHN 7992 partim). Figured by Gervais 1848-1852, pl. 49, fig. 1 (skull with vertebrae) and 3 (sternum), and Brunet 1970, pl. A, fig. c (sternum).]

Numenius (Limosa) gypsorum (Gervais): Milne-Edwards, 1871: 551 [New combination.]

Limosa gypsorum (Gervais): Brodkorb 1967: 185 [New combination.]

Villetus gypsorum (Gervais): Harrison 1979b: 107 [New combination.]

Montirallus gypsorum (Gervais): Mlíkovský 1981: 342 [New combination.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Gervais 1844a,b, 1848-1852, Brunet 1970, Harrison 1979b, Mlíkovský 1981, 1995a).

Remarks: This species was originally described as a curlew. Brodkorb (1967: 185) and Harrison (1979b: 107) confirmed the identification (though in different genera), but Mlíkovský (1981) moved the species on the basis of the brain morphology to the family Rallidae.

Genus *Quercyrallus* Brodkorb

Quercyrallus Lambrecht, 1933: 461 [Nomen nudum; type not fixed after 1930.]

Quercyrallus Brodkorb, 1952: 175 [Type by original designation: *Rallus arenarius* Milne-Edwards, "1891" = 1892.]

***Quercyrallus arenarius* (Milne-Edwards)**

Rallus arenarius Milne-Edwards, 1892: 74 [Holotype from Quercy: proximal end of left humerus; MNHN QU-3072. Figured by Gaillard 1908, fig. 33, Cracraft 1973, fig. 10a.]

Quercyrallus arenarius (Milne-Edwards): Lambrecht 1933: 461 [New combination, but the genus name was nomen nudum at this point.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908: 110, Cracraft 1973: 22, Olson 1977: 344).

***Quercyrallus quercy* Cracraft**

Quercyrallus quercy Cracraft, 1973: 24 [Holotype from Quercy: distal end of right humerus; MNHN QU-3071. Figured by Cracraft 1973, fig. 10b-c.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Cracraft 1973, Olson 1977: 345).

Genus *Rallicrex* Lambrecht

Rallicrex Lambrecht, 1933: 463 [Type by monotypy: *Rallicrex kolozsvarensis* Lambrecht, 1933.]

***Rallicrex kolozsvarensis* Lambrecht**

Rallicrex kolozsvarensis Lambrecht, 1933: 463 [Holotype from Cetățuie: distal end of left (not right as stated by Olson 1977: 345) tarsometatarsus; GIB, uncatalogued. Figured by Lambrecht 1933, fig. 141a-b, and Jurcsák & Kessler 1973, fig. 9.]

Distribution: Middle Oligocene (MP 24) of Cetățuie, Romania (Lambrecht 1933: 463,

Cracraft 1973: 25, Olson 1977: 345).

***Rallicrex polgardiensis* Jánossy**

Rallicrex polgardiensis Jánossy, 1991: 20 [Holotype from Polgárdi 5: left femur; GIB Vt-147-V-18.096. Figured by Jánossy 1991, pl. 3, fig. 9.]

Distribution: Late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991), and Polgárdi 5, Hungary (Jánossy 1991).

Remarks: The generic position of this species is uncertain.

Genus *Paraortygometra* Lambrecht

Paraortygometra Lambrecht, 1933: 462 [Type by monotypy: *Rallus porzanoides* Milne-Edwards, 1869.]

Microrallus Švec, 1983: 32 [Type by original designation: *Microrallus fejfari* Švec, 1983.]

***Paraortygometra porzanoides* (Milne-Edwards)**

Rallus porzanoides Milne-Edwards, 1869: 150 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus; MNHN Av-2871. Figured by Milne-Edwards 1869-1871, pl. 105, fig. 1-7, Cracraft 1973, fig. 17a-c. Cracraft (1973: 34) called the holotypical tarsometatarsus a lectotype, but Milne-Edwards 1869: 150 clearly based the species on a single tarsometatarsus, tentatively assigning to the species (on his p. 151) also distal end of a right femur (MNHN Av-2870) and a right humerus (MNHN Av-2869).]

Paraortygometra porzanoides (Milne-Edwards): Lambrecht 1933: 463 [New combination.]

Microrallus fejfari Švec, 1983: 37 [Holotype from Dolnice: distal end of left humerus; DP FNSP 4818. Figured by Švec 1983, text-fig. 1a, pl. 1.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Lambrecht 1933: 463, Cracraft 1973: 34, Olson 1977: 345); early Miocene (MN 4b) of Dolnice, Czechia (Švec 1983, Mlíkovský orig.); middle Miocene (MN 8) of Grive-Saint-Alban – fissure H, France (Ballmann 1969a, Cracraft 1973: 34, Olson 1977: 346); and late Miocene (MN 12) of Aljezar B, Spain (Cheneval & Adrover 1993). Extralimital record is available from the early Miocene (MN 3?) of Li Mae Long, Thailand (Cheneval et al. 1984, 1991).

Genus *Palaeoaramides* Lambrecht

Palaeoaramides Lambrecht, 1933: 462 [Type by monotypy: *Rallus christyi* Milne-Edwards, 1869.]

Pararallus Lambrecht, 1933: 466 [Type by monotypy: *Rallus dispar* Milne-Edwards, 1869.]

Tertiaporphyryula Kuročkin & Ganea, 1972: 66 [Type by original designation: *Rallus beaumonti* Milne-Edwards, 1869.]

Remarks: The extralimital record is limited to *Palaeoaramides tugarinovi* Kuročkin, 1980 from the middle Pliocene of Mongolia (Kuročkin 1980). The genus *Tertiaporphyryula* Kuročkin & Ganea was synonymized with *Palaeoaramides* Lambrecht by Cracraft (1973: 30), and the genus *Pararallus* Lambrecht is synonymized with *Palaeoaramides* Lambrecht here (see under *Palaeoaramides beaumontii* Milne-Edwards).

***Palaeoaramides christyi* (Milne-Edwards)**

Rallus christyi Milne-Edwards, 1869: 146 [Lectotype from Saint-Gérard-le-Puy (selected by Cracraft 1973: 27): right tarsometatarsus; MNHN Av-2868. Figured by Milne-Edwards 1869-1871, pl. 104, fig. 1-6 (not pl. 103, fig. 1-5, as stated by Brodkorb 1967: 119). Paralectotype from Saint-Gérard-le-Puy: right tibiotarsus lacking proximal end; MNHN Av-2867. Figured by Milne-Edwards 1869-1871, pl. 104, fig. 1, 7-9.]

Rallus eximius Milne-Edwards, 1869: 149 [Holotype from Saint-Gérard-le-Puy: right tarso-

metatarsus; MNHN Av-2865. Figured by Milne-Edwards 1969-1871, pl. 103, fig. 6-11.]

Palaeoaramides christyi (Milne-Edwards): Lambrecht 1933: 462 [New combination.]

Palaeoaramides eximius (Milne-Edwards): Brodkorb 1967: 119 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871 sub *Rallus christyi*, Cracraft 1973: 27, Olson 1977: 346).

Remarks: *Rallus eximius* Milne-Edwards was synonymized with *Rallus christyi* Milne-Edwards by Lambrecht (1933: 462, see also Cracraft 1973, Olson 1977).

***Palaeoaramides beaumontii* (Milne-Edwards)**

Rallus Beaumontii Milne-Edwards, 1869: 152 [Lectotype from Sansan (selected by Cracraft 1973: 30): distal end of right tibiotarsus; MNHN Sa-1205. Figured by Milne-Edwards 1969-1871, pl. 104, fig. 16-20. Paralectotypes from Saint-Gérard-le-Puy: proximal end of right humerus (MNHN Sa-1204), distal end of right humerus (MNHN Sa-1203), proximal end of right tarsometatarsus (present location unknown), and distal end of right tarsometatarsus (MNHN Sa-1206). Figured by Milne-Edwards 1869-1871, pl. 104, fig. 10-12 (distal end of tarsometatarsus), 13-15 (proximal end of tarsometatarsus), 21-22, 26 (proximal end of humerus), 21-25 (distal end of humerus); and Cracraft 1973, fig. 13a-b (distal end of humerus). The list of paralectotypes presented by Cheneval (2000: 357) incorrectly includes also several syntypes of *Palaeoperdix prisca* Milne-Edwards.]

Rallus dispar Milne-Edwards, 1869: 155 [Lectotype from Sansan (here selected): distal end of left tarsometatarsus; MNHN Sa-1210. Figured by Milne-Edwards 1869-1871, pl. 105, fig. 17-20, 22, and Cracraft 1973, fig. 16a-b. Paralectotypes from Sansan: proximal ends of two right tarsometatarsi (MNHN Sa-1207-1208), and proximal end of a left tarsometatarsus (Sa-1209). Figured by Milne-Edwards 1969-1871, pl. 105, fig. 17-21 (proximal end of right tarsometatarsus). Milne-Edwards (1869: 155) assigned to this species another two elements from Sansan, which led Brodkorb (1967: 120), Cracraft (1973: 33) and Olson (1977: 347) to believe that they form part of the syntypal series, which is not the case (see Mlíkovský 1998e: 338). Cracraft (1973: 33) erroneously selected one of the assigned elements (distal end of left humerus; MNHN Sa-1201) as a lectotype of the species, this action being invalid from the point of view of zoological nomenclature (ICZN 1999). Unfortunately, the "lectotypical" humerus belongs to a parrot (Cheneval 2000). In belief that the latter bone is the name-bearer of *Rallus dispar*, Cheneval (2000) applied the name to the parrot from Sansan, which is unsubstantiated (see also Mlíkovský 1998e).

Pararallus dispar (Milne-Edwards): Lambrecht 1933: 466 [New combination.]

Pararallus beaumonti (Milne-Edwards): Brodkorb 1967: 120 [New combination.]

Tertiariaporphyryula beaumonti (Milne-Edwards): Kuročkin & Ganea 1972: 66 [New combination.]

Palaeoaramides beaumonti (Milne-Edwards): Cracraft 1973: 30 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871, Cracraft 1973, Olson 1977, Mlíkovský 1998e).

Remarks: The lectotype of *Rallus dispar* Milne-Edwards was assigned to *Palaeoaramides beaumonti* (Milne-Edwards) by Cheneval (2000: 357). Accordingly, I synonymize here the former with latter species.

***Palaeoaramides lungi* (Kuročkin & Ganea)**

Tertiariaporphyryula lungi Kuročkin & Ganea, 1972: 67 [Holotype from Bujoru: distal end of right tibiotarsus; TGPI 4/60. Figured by Kuročkin & Ganea 1972, text-fig. 12, pl. 2, fig. 10.]

Palaeoaramides lungi (Kuročkin & Ganea): Olson 1977: 346 [New combination.]

Distribution: Late Miocene (MN 9-11) of Bujoru, Moldova (Kuročkin & Ganea 1972).

Genus *Miorallus* Lambrecht

Miorallus Lambrecht, 1933: 466 [Type by monotypy: *Rallus major* Milne-Edwards, 1869.]

***Miorallus major* (Milne-Edwards)**

Rallus major Milne-Edwards, 1869: 157 [Holotype from Sansan: distal end of left humerus; MNHN Sa-1200. Figured by Milne-Edwards 1869-1871, pl. 103, fig. 12-16; Cracraft 1973, fig. 18a-b; and Cheneval 2000, fig. 17.]

Miorallus major (Milne-Edwards): Lambrecht 1933: 466 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871 sub *Rallus major*, Lambrecht 1933, Cracraft 1973, Olson 1977).

***Miorallus* sp.**

Distribution: Late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1993).

Remarks: This is a doubtful record, based on a proximal end of a humerus, not figured, and said to be "on all sides damaged" (Jánossy 1993: 58).

Genus *Porzana* Vieillot

Porzana Vieillot, 1816 [Modern genus.]

***Porzana estramosi* Jánossy**

Porzana estramosi Jánossy, 1979: 20 [Holotype tarsometatarsus from Osztramos 9: fragmentary distal end of left; GIB V-78.120. Figured by Jánossy 1979, pl. 4, fig. 8.]

Porzana estramosi veterior Jánossy, 1991: 19 [Holotype from Polgárdi 4: right tarsometatarsus; GIB Vt-146-V-18.095. Figured by Jánossy 1991, pl. 3, fig. 7.]

Distribution: Middle Miocene (MN 6-8) of Mátraszőlös 1, Hungary (Gál & Kessler in Gál et al. 1999 sub *Porzana* aff. *estramosi*); late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991); and early Pliocene (MN 14) of Osztramos 9, Hungary (Jánossy 1979).

***Porzana porzana* (Linnaeus) – Spotted Crake**

Rallus Porzana Linnaeus, 1766 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub cf. *P. porzana*); early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992 sub *P.* cf. *porzana*); and early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub *P.* cf. *porzana*), and Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972).

***Porzana parva* (Scopoli) – Little Crake**

Rallus parvus Scopoli, 1769 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992 sub *P.* cf. *parva*).

***Porzana* sp.**

Distribution: Late Pliocene (MN 18) of Väršec, Bulgaria (Boev 1997b: 70, 2000a); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 15e, Czechia (Mlíkovský 1995b).

Genus *Crex* Bechstein

Crex Bechstein, 1803 [Modern genus.]

***Crex crex* (Linnaeus) – Corn Crake**

Rallus Crex Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 17) of Kryžanovka 1, Ukraine (Dubrovo & Kapelist 1979 sub *C. cf. crex*); early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *C. cf. crex*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 4a, and 9a, Czechia (Jánossy 1972 sub *C. cf. crex*), and Stránská skála – Musil's talus cone, layers 8, 11, and 13, Czechia (Mlíkovský 1995b).

***Crex* sp.**

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Mlíkovský orig., material in USNM); late Miocene (MN 10) of Kohfidisch, Austria (Mlíkovský orig., material in NHMW); and early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992).

Genus *Rallus* Linnaeus

Rallus Linnaeus, 1758 [Modern genus.]

***Rallus aquaticus* Linnaeus – Water Rail**

Rallus aquaticus Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 15) of Csarnóta 2, Hungary (Jánossy 1979 sub *Rallus* sp. – *aquaticus* group); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 8, Czechia (Mlíkovský 1995b).

***Rallus* sp.**

Distribution: Middle Miocene (MN 6-8) of Mátraszőlös 1, Hungary (Gál & Kessler in Gál et al. 1999).

Genus *Gallinula* Brisson

Gallinula Brisson, 1760 [Modern genus.]

***Gallinula balcanica* Boev**

Gallinula balcanica Boev, 1997b: 70, 78 [Nomen nudum; no description or indication]

Gallinula balcanica Boev, 1999g: 43 [Holotype from Väršec: left ulna; NMNHS 112. Figured by Boev 1999g, fig. 1.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1999g).

***Gallinula gigantea* Tchernov**

Gallinula gigantea Tchernov, 1980: 30 [Holotype from 'Ubeidiya I: fragmentary proximal end of right humerus; HUI, uncatalogued. Figured by Tchernov 1980, pl. 2, fig. 4.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b). The extralimital record is limited to the early Pleistocene (MQ 1a) of 'Ubeidiya I, Israel (Tchernov 1980).

***Gallinula chloropus* (Linnaeus) – Common Moorhen**

Fulica Chloropus Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene or early Pleistocene (MN 16 - MQ 1) of Bobila Ordis, Spain (Mayr & Gregor 1999 sub *G. cf. chloropus*); early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992 sub *G. cf. chloropus*); and early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d), Boxgrove – quarry 1, England (Harrison & Stewart 1999), and Přezletice, Czechia (Jánossy 1983 sub *G. aff. chloropus*).

***Gallinula* sp.**

Distribution: Early Pliocene (MN 15) of Csarnóta 2, Hungary (Jánossy 1979); and early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub ?*Gallinula*).

Genus *Fulica* Linnaeus

Fulica Linnaeus, 1758 [Modern genus.]

***Fulica atra* Linnaeus – Eurasian Coot**

Fulica atra Linnaeus, 1758 [Modern species.]

Fulica atra pontica Boev & Karajvanova, 1998: 60 [Holotype from Sozopol: right tibiotarsus; NMNHS 6568. Figured by Boev & Karajvanova 1998, fig. 2.]

Distribution: Early Pleistocene (MQ 1b) of Přezletice, Czechia (Jánossy 1983 sub *F. aff. atra*), and Stránská skála – Musil's talus cone – layers 4a, 5, 10b, 143, and 15e, Czechia (Mlíkovský 1995b); and middle Holocene (MQ 2E; ca. 7000-5500 BP) of Sozopol, Bulgaria (Boev & Karajvanova 1998 sub *F. atra pontica*).

Family Cariamidae Bonaparte

Cariamidae Bonaparte, 1850 [Modern family.]

Phorusrhacidae Ameghino, 1889: 659 [Type genus: *Phororhacos* Ameghino, 1889, which is an incorrect subsequent spelling of *Phorusrhacus* Ameghino, 1887 – see Mlíkovský 2000f.]

Brontornithidae Moreno & Mercerat, 1891: 20 [Type genus: *Brontornis* Moreno & Mercerat, 1891.]

Stereornithidae Moreno & Mercerat, 1891: 21 [Type genus: *Stereornis* Moreno & Mercerat, 1891.]

Darwinornithidae Moreno & Mercerat, 1891: 24 [Type genus: *Darwinornis* Moreno & Mercerat, 1891.]

Pelecymnidae Ameghino, 1891: 448 [Type genus: *Pelecymnis* Ameghino, 1891. This is incorrect original spelling. This is a senior objective synonym of Psilopteridae Sáez, 1927 – see Mlíkovský 2000d.]

Patagornithidae Mercerat, 1897: 225 [Type genus: *Patagornis* Moreno & Mercerat, 1891.]

Phororhacidae Ameghino: Lydekker, 1893d: 43 [Spelling emended.]

Orthocnémides Gaillard, 1908: 113 [Vernacular name applied after 1900; not available for nomenclatural purposes – see Brodkorb 1967: 136, Mlíkovský 2000d.]

Hermosiornidae Rovereto, 1914: 110 [Type genus: *Hermosiornis* Rovereto, 1914. This is incorrect original spelling.]

Orthocnemidae Lambrecht, 1921: 70 [Type genus: *Orthocnemus* Milne-Edwards, 1892 = *Idiornis* Oberholser, 1899. Brodkorb (1967: 136) erroneously attributed this family-group name to Lambrecht 1933: 490.]

Bathornithinae Wetmore, 1927: 13 [Type genus: *Bathornis* Wetmore, 1927.]

Psilopteridae Sáez, 1927: 156 [Type genus: *Psilopterus* Moreno & Mercerat, 1891 = *Pelecymnis* Ameghino, 1891. This is a junior objective synonym of Pelecymnidae Ameghino, 1891 – see Mlíkovský 2000d.]

Devincenziidae Kraglievich, 1932: 323 [Type genus: *Devincenzia* Kraglievich, 1932.]

Mesembriornidae Kraglievich, 1932: 332 [Type genus: *Mesembriornis* Moreno & Mercerat, 1891. This is incorrect original spelling.]

Liorninae Kraglievich, 1932: 348 [Type genus: *Liornis* Ameghino, 1895 = *Rostrornis* Moreno & Mercerat, 1891. This is incorrect original spelling.]

Brontorniinae Moreno & Mercerat: Kraglievich 1932: 348 [New rank; spelling incorrect.]

Tolmodinae Kraglievich, 1932: 348 [Type genus: *Tolmodus* Ameghino, 1891.]

Agnopteridae Lambrecht, 1933: 333 [Type genus: *Agnopterus* Milne-Edwards, 1868.]
 Bathornithidae Wetmore: Wetmore 1933: 301 [New rank.]
 Hermosiornithidae Rovereto: Wetmore, 1934b: 7 [Spelling emended.]
 Phororhacoidea Ameghino: Patterson 1941: 49 [New rank.]
 Mesembriornithidae Kraglievich: Kraglievich 1946: 108 [Spelling emended.]
 Hermosiornithinae Rovereto: Patterson & Kraglievich 1960: 17 [New rank.]
 Phorusrhacidae Ameghino: Brodkorb 1963c: 111 [Spelling emended.]
 Phorusrhacoidae Ameghino: Brodkorb 1963c: 111 [Spelling emended.]
 Idiornithidae Brodkorb, 1965: 197 [Type genus: *Idiornis* Oberholser, 1899.]
 Idiornithinae Brodkorb: Mourer-Chauviré 1983: 87 [New rank.]
 Liornithidae Kraglievich: Mlíkovský 2000d: 79 [Spelling emended.]
 Pelecyornithidae Ameghino: Mlíkovský 2000d: 80 [Spelling emended.]
 Phorusrhacinae Mlíkovský this paper [New rank.]

Genus *Gypsornis* Milne-Edwards

Gypsornis Milne-Edwards, 1869: 140 [Type by monotypy: *Gypsornis cuvieri* Milne-Edwards, 1869.]

Percolinus Harrison & Walker, 1977a: 34 [Type by original designation: *Percolinus venablesi* Harrison & Walker, 1977a.]

Distribution: Early Eocene (MP 8-9) of England, and late Eocene (MP 19) of France.

Gypsornis venablesi (Harrison & Walker)

Percolinus venablesi Harrison & Walker, 1977a: 34 [Holotype from Sheppey: proximal end of left tarsometatarsus; BMNH A-3680. Figured by Harrison & Walker 1977a, pl. 8, fig. k-o.]

Gypsornis venablesi (Harrison & Walker): Mlíkovský, this paper [New combination.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Remarks: This species was originally described in the Phasianidae (Harrison & Walker 1977), but its holotypical tarsometatarsus differs from the same element of phasianids and agrees with that of *Gypsornis* Milne-Edwards in the shape of its proximal end (best visible in proximal view), and in the configuration of hypotarsus. Hence, I include here *Percolinus venablesi* in the genus *Gypsornis*. Within the latter genus, *venablesi* is smaller (proximal width of tarsometatarsus = 7.2 mm vs. 11.7 mm) and older than *Gypsornis cuvieri* Milne-Edwards.

Gypsornis cuvieri Milne-Edwards

Gypsornis cuvieri Milne-Edwards, 1869: 140 [Holotype from Montmartre: proximal end of left tarsometatarsus; MNHN 7983. Figured by Cuvier 1822, pl. 74, fig. 7 (not pl. 73 as stated by Cuvier 1822: 324; see also Brunet 1970: 27); Milne-Edwards 1869-1871, pl. 103, fig. 1-5; and Cracraft 1973, fig. 21a-b.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Cuvier 1822 as an unnamed bird, Milne-Edwards 1869-1871, Brunet 1970, Cracraft 1973).

Remarks: This species was originally described as a rail (Milne-Edwards 1869-1871, see also Brunet 1970), but Cracraft (1973: 53) transferred it to the Idiornithidae.

Genus *Talantatos* Reichenbach

Talantatos Reichenbach, 1852: xiv [Type by monotypy: *Tantalus fossilis* Giebel, 1847. Note that this genus name was consistently misspelled as *Tantalatos* by Mlíkovský 1995a.]

Elaphrocnemus Milne-Edwards, 1892: 77 [Type by subsequent designation (Richmond 1902: 680): *Elaphrocnemus phasianus* Milne-Edwards, 1892.]

Filholornis Milne-Edwards, 1892: 67 [Type by subsequent designation (Richmond 1902: 684): *Filholornis paradoxa* Milne-Edwards, 1892.]

Distribution: Middle Eocene (MP 11-13) of England, and late Eocene to late Oligocene (MP 19-28) of France. *Phasianus alfhildae* Shufeldt, 1915 from the late Eocene of Wyoming was included in *Elaphrocnemus* Milne-Edwards by Brodkorb (1967: 138), but Cracraft (1973: 69) excluded it from the Cariamidae.

***Talantatos proudlocki* (Harrison & Walker)**

Percolinus proudlocki Harrison & Walker, 1979a: 21 [Holotype from Lee-on-Solent: proximal end of right tarsometatarsus; BMNH A-4286. Figured by Harrison & Walker 1979a, pl. 1, row 3 partim.]

Talantatos proudlocki (Harrison & Walker): Mlíkovský, this paper [New combination.]

Distribution: Middle Eocene (MP 11-13) of Lee-on-Solent, England (Harrison & Walker 1979a).

Remarks: This species was originally described in the Phasianidae (Harrison & Walker 1979a), but its holotypical tarsometatarsus markedly differs from the same element of phasianids and agrees with that of *Talantatos* Reichenbach in the shape of its proximal end (in proximal view) and in the configuration of the hypotarsus. Accordingly, I transfer here *proudlocki* from the genus *Percolinus* Harrison & Walker to *Talantatos* Reichenbach. Within the latter genus, the holotypical tarsometatarsus of *proudlocki* agrees in size with *Talantatos fossilis* Reichenbach, but its internal hypotarsal ridge seems to be more slender and the species is much older. Hence, I retain here *proudlocki* as a separate species in the genus *Talantatos* Reichenbach until more evidence is obtained.

***Talantatos fossilis* (Giebel)**

T[antalus] fossilis Giebel, 1847: 27 [Holotype from Montmartre: right femur; MNHN 7992 partim. Figured by Cuvier 1822, pl. 73, fig. 14a-c. See Mlíkovský 1995a for the identity of the holotype.]

Talantatos fossilis (Giebel): Reichenbach 1852: xiv [New combination.]

Elaphrocnemus phasianus Milne-Edwards, 1892: 77 [Holotype from Quercy: left tarsometatarsus; MNHN QU-15514. Figured by Cracraft 1973, fig. 27a-b. Milne-Edwards (1892) described this species on the basis of a single tarsometatarsus. Its measurements agree with those of the tarsometatarsus labeled by Cracraft (1973: 61) as a lectotype, this being holotype in fact. Mourer-Chauviré (1983: 92) followed Cracraft in the erroneous assumption that this specimen is a lectotype.]

Filholornis paradoxa Milne-Edwards, 1892: 67 [Holotype from Quercy: left humerus; MNHN QU-16200. Not figured.]

Filholornis debilis Milne-Edwards, 1892: 69 [Holotype from Quercy: left humerus; MNHN QU-16227. Figured by Mourer-Chauviré 1983, pl. 1, fig. 3-4.]

Orthocnemus major Milne-Edwards, 1892: 76 [Holotype from Quercy: distal end of left tarsometatarsus; MNHN QU-15506. Figured by Cracraft 1973, fig. 29c-d. Milne-Edwards (1892: 76-77) clearly based the species on this distal end of tarsometatarsus, referring to it a proximal end of right tarsometatarsus from Quercy (MNHN QU-15505). Cracraft (1973: 57) erroneously selected the referred specimen as a lectotype of the species, being followed by Mourer-Chauviré (1983:110).]

Elaphrocnemus crex Milne-Edwards, 1892: 78 [Lectotype from Quercy: right tarsometatarsus; MNHN QU-15512 (formerly MNHN 3012). Figured by Cracraft 1973, fig. 29a-b. Paralectotypes from Quercy (all in MNHN): distal end of right tibiotarsus (QU-15680), right tarsometatarsus (QU-15518), left tarsometatarsus (QU-15566), distal end of right tarsometatarsus (QU-15513), and distal end of left tarsometatarsus (QU-15506). Figured by Cracraft 1973, fig. 29c-d (distal end of left tarsometatarsus). The list of paralectotypes is based on Cracraft's (1973: 66-67) and Mourer-Chauviré's (1983: 95) lists of bones from the early collections from Quercy from MNHN which could be at disposal to Milne-Edwards, and which were identified as belonging to *Elaphrocnemus crex* by these two modern

authors.]

Idiornis major (Milne-Edwards): Oberholser 1899: 202 [New combination.]

Telecrex peregrinus Mlíkovský, 1989b: 64 [Holotype from Quercy: left femur; NHMW 1988/28/1. Figured by Mlíkovský 1989b, pl. 1, fig. a-f.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908, 1939, Cracraft 1973, Mourer-Chauviré 1983); late Eocene (MP 19) of Montmartre, France (Cuvier 1822 as an unidentified bird, Giebel 1847, Mlíkovský 1995a), Rosières 2, France (Mourer-Chauviré 1983), Escamps C, Escamps III and Escamps IV, France (Mourer-Chauviré 1983); early Oligocene (MP 21) of Aubrelong 1, France (Mourer-Chauviré 1983) and Ravet Lupovici, France (Mourer-Chauviré 1983); early Oligocene (MP 22) of Plante 2, France (Mourer-Chauviré 1983); middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1983), Roqueprune 2, France (Mourer-Chauviré 1983), and Crabit 1, France (Mourer-Chauviré 1983); and late Oligocene (MP 28) of Desse, France (Mourer-Chauviré 1983), and Fraysse, France (Mourer-Chauviré 1983).

Remarks: The nomenclatural history of *Tantalus fossilis* was traced by Mlíkovský (1995a: 312). Gervais (1848-1852: 230) synonymized *Tantalus fossilis* Giebel with *Numenius gypsorum* Gervais. This decision was followed by Brodkorb (1967: 185) and Brunet (1970: 25), who believed that these two forms were based on the same types. This is not true, however (see Mlíkovský 1995a: 312). I examined the holotypical femur of *Tantalus fossilis* in MNHN, finding that it is identical in both size and morphology with the same element of *Elaphrocnemus phasianus* Milne-Edwards. The two species are thus synonymous, with *Talantatos fossilis* (Giebel) being the senior synonym.

Filholornis paradoxa Milne-Edwards and *Filholornis debilis* Milne-Edwards were synonymized with *Elaphrocnemus phasianus* Milne-Edwards by Mourer-Chauviré (1983: 88, 92). Cracraft (1973: 65) and Mourer-Chauviré (1983: 95) recognized *Elaphrocnemus crex* Milne-Edwards as a separate species, size being the distinguishing character. The measurements (Mourer-Chauviré 1983, pl. 1-2) show that *Elaphrocnemus phasianus* Milne-Edwards was based on smaller, while *Elaphrocnemus crex* Milne-Edwards on larger specimens of a single species. Accordingly, I synonymize here the latter with the former species.

Orthocnemus major Milne-Edwards was synonymized with *Orthocnemus cursor* Milne-Edwards by Cracraft (1973: 57; see also Mourer-Chauviré 1983: 88, 110). This synonymization was based on a misidentified holotype (see above). The proper holotype, erroneously considered paralectotype by Cracraft (1973) and Mourer-Chauviré (1983), was referred by both these authors to *Elaphrocnemus crex* Milne-Edwards. Hence, *Orthocnemus major* Milne-Edwards must be removed from the synonymy of *Orthocnemus cursor* Milne-Edwards [= *Agnopterus gallicus* (Milne-Edwards)], and synonymized with *Talantatos fossilis* (Giebel), following the taxonomic treatment of the genus *Talantatos* Reichenbach accepted in the present paper. *Telecrex peregrinus* Mlíkovský, originally described as a guinea-fowl, was synonymized with *Elaphrocnemus phasianus* Milne-Edwards [= *Talantatos fossilis* (Giebel)] by Mourer-Chauviré (1992b: 87).

***Talantatos brodkorbi* (Mourer-Chauviré)**

Elaphrocnemus brodkorbi Mourer-Chauviré, 1983: 97 [Holotype from Quercy: proximal end of left tarsometatarsus; MNHN QU-15500. Figured by Mourer-Chauviré 1983, pl. 2, fig. 3-4.]

Talantatos brodkorbi (Mourer-Chauviré): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1983); and middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1983 sub *Elaphrocnemus crex*, reidentified by Mlíkovský, orig.).

Remarks: Mourer-Chauviré (1983: 95) attributed proximal end of left femur from Itardiès (USTL ITD-597) to *Elaphrocnemus crex* Milne-Edwards [= *Talantatos fossilis* (Giebel)], but the measurements (Mourer-Chauviré 1983, tab. 2) indicate that the specimen is too large for

this species, corresponding in size with the holotype of *Talantatos brodkorbi* (Mourer-Chauviré).

Genus *Agnopterus* Milne-Edwards

Agnopterus Milne-Edwards, 1867: 83 [Type by monotypy: *Agnopterus laurillardi* Milne-Edwards, 1868.]

Orthocnemus Milne-Edwards, 1892: 74 [Type by subsequent designation (Brodkorb 1967: 136): *Orthocnemus gallicus* Milne-Edwards, 1892. The type was designated by original decision according to Brodkorb (1967: 136), but I found nothing in Milne-Edwards (1892) what would indicate which of the three species included in the genus was deemed to be its type. Accordingly, I attribute here the respective nomenclatural act to Brodkorb (1967) himself. This is a junior homonym of *Orthocnemus* Jekel, 1857.]

Idiornis Oberholser, 1899: 202 [New name for *Orthocnemus* Milne-Edwards, 1892; hence its junior objective synonym.]

Distribution: Early Eocene (MP 11) of Germany, and late Eocene to late Oligocene (MP 16-28) of France.

Remarks: My restudy of the holotypical tibiotarsus of *Agnopterus laurillardi* Milne-Edwards showed that it is identical with the same element of the cariamas included by Mourer-Chauviré (1983) in the genus *Idiornis* Oberholser. *Agnopterus* Milne-Edwards has precedence over *Idiornis* Oberholser, which results in several nomenclatural changes (new combinations) that are made below.

Agnopterus tuberculatus (Peters)

Idiornis tuberculata Peters, 1995: 107 [Holotype from Messel: crushed skeleton in slab and counter-slab; IRScNB, uncatalogued (= field-number 160 of the 1986 excavations in Messel). Figured by Peters 1995, fig. 1-6, 9.]

Agnopterus tuberculatus (Peters): Mlíkovský, this paper [New combination.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Peters 1995).

Agnopterus gallicus (Milne-Edwards)

Filholornis gravis Milne-Edwards, 1892: 69 [Holotype from Quercy: left ulna; MNHN QU-15558. Figured by Mourer-Chauviré 1983, pl. 2, fig. 10-11. Milne-Edwards (1892: 69) attributed to *Filholornis gravis* also left carpometacarpus from Quercy (MNHN QU-16206). Mourer-Chauviré (1983: 88, 105) erroneously considered both the specimens syntypes of the species.]

Orthocnemus gallicus Milne-Edwards, 1892: 74 [Lectotype from Quercy (selected by Cracraft 1973: 54): right tarsometatarsus; MNHN QU-15502. Figured by Cracraft 1973, fig. 22a-b, and Mourer-Chauviré 1983, pl. 2, fig. 14-15. Paralectotype from Quercy: distal end of left tibiotarsus, MNHN QU-15605. Figured by Mourer-Chauviré 1983, pl. 3, fig. 10.]

Orthocnemus cursor Milne-Edwards, 1892: 76 [Holotype from Quercy: left tarsometatarsus; MNHN QU-15508. Figured by Cracraft 1973, fig. 24a, and Mourer-Chauviré 1983, pl. 3, fig. 1-2. Cracraft (1973: 57) and Mourer-Chauviré (1983: 110) referred to the holotypical tarsometatarsus as to the lectotype. However, Milne-Edwards (1892: 76) clearly based *Orthocnemus cursor* on a complete tarsometatarsus (only one was available to him), referring to the species several other tarsometatarsi smaller than the holotype, and distal end of a tibiotarsus. The latter specimens thus do not form part of a syntypal series, and the complete tarsometatarsus is to be regarded as the holotype.]

Idiornis gallicus (Milne-Edwards): Oberholser 1899: 202 [New combination.]

Idiornis cursor (Milne-Edwards): Oberholser 1899: 202 [New combination.]

Idiornis gaillardi Cracraft, 1973: 59 [Holotype from Quercy: left tarsometatarsus; MNHN QU-

15534. Figured by Cracraft 1973, fig. 26a-c, and Mourer-Chauviré 1983, pl. 3, fig. 18-19.]
Agnopterus gallicus (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908, 1939, Cracraft & Rich 1972 sub *Diatropornis ellioti* [see Mourer-Chauviré 1983: 111], Cracraft 1973, Mourer-Chauviré 1983); late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1983) and Lavergne, France (Mourer-Chauviré 1983); late Eocene (MP 16-20) of Garrigues, France (Mourer-Chauviré 1983); late Eocene (MP 17) of Perrière, France (Mourer-Chauviré 1983), Salème, France (Mourer-Chauviré 1983), and Bouffie, France (Mourer-Chauviré 1983); late Eocene (MP 19) of Escamps, France (Mourer-Chauviré 1983), and Lostanges, France (Mourer-Chauviré 1983); Oligocene (MP 21-28) of Boussac 2, France (Mourer-Chauviré 1983); middle Oligocene (MP 22) of Mas de Got B, France (Mourer-Chauviré 1983); middle Oligocene (MP 23) of Roqueprune 2, France (Mourer-Chauviré 1983); and late Oligocene (MP 28) of Desse, France (Mourer-Chauviré 1983), and Fraysse, France (Mourer-Chauviré 1983).

Remarks: *Filholornis gravis* Milne-Edwards was synonymized with *Orthocnemus* [= *Agnopterus*] *gallicus* (Milne-Edwards) by Mourer-Chauviré (1983: 88, 104). Both Cracraft (1973) and Mourer-Chauviré (1983) recognized three small-sized species within the genus *Idiornis* Oberholser, incl. *Idiornis gallicus* (Milne-Edwards), *Idiornis cursor* (Milne-Edwards), and *Idiornis gaillardi* Cracraft. Reconsidering the published measurements (Cracraft 1973, tab. 19-21, Mourer-Chauviré 1983, tab. 4-7), I found that the overall coefficient of variability for these three species is less than 10 %, which is well within the standard. Hence, I synonymize here the former two species with *Agnopterus gallicus* (Milne-Edwards).

***Agnopterus minor* (Milne-Edwards)**

Orthocnemus minor Milne-Edwards, 1892: 77 [Holotype from Quercy: distal end of right tarsometatarsus; MNHN QU-15504. Figured by Cracraft 1973, fig. 25a-b.]

Idiornis minor (Milne-Edwards): Oberholser 1899: 202 [New combination.]

Agnopterus minor (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Mourer-Chauviré 1983); late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1983), and Perrière, France (Mourer-Chauviré 1983); and late Eocene (MP 16-20) of Garrigues, France (Mourer-Chauviré 1983).

***Agnopterus laurillardi* Milne-Edwards**

Agnopterus laurillardi Milne-Edwards, 1867: 83 [Holotype from Montmartre: distal end of left tibiotarsus; MNHN 7979. Figured by Milne-Edwards, 1867-1868, pl. 89, fig. 10-15; and Brunet 1970, fig. 3a-d.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Milne-Edwards 1867, Brunet 1970).

Remarks: This species was originally described as a flamingo (see also Brunet 1970). My restudy of the holotypical tibiotarsus showed that it differs in its general shape and in the shape of the internal condyle from the same element of the Phoenicopteridae and agrees with that of *Agnopterus* Milne-Edwards, where it is larger than other species (see Mourer-Chauviré 1983: 108 for measurements). The holotype of *A. laurillardi* Milne-Edwards has the following dimensions: depth of internal condyle = ca. 20 mm, posterior width of distal end = ca. 13 mm (Mlíkovský orig.).

***Agnopterus gracilis* (Milne-Edwards)**

Elaphrocnemus gracilis Milne-Edwards, 1892: 78 [Holotype from Quercy: left tarsometatarsus; MNHN QU-15511. Figured by Cracraft 1973, fig. 31, and Mourer-Chauviré 1983, pl. 4, fig. 1-2.]

Idiornis gracilis (Milne-Edwards): Mourer-Chauviré 1983: 115 [New combination.]

Agnopterus gracilis (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908); middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1983); late Oligocene (MP 25) of Belgarric 1, France (Mourer-Chauviré 1983); and late Oligocene (MP 28) of Desse, France (Mourer-Chauviré 1983).

***Agnopterus itardiensis* (Mourer-Chauviré)**

Idiornis itardiensis Mourer-Chauviré, 1983: 116 [Holotype from Itardiès: distal end of left tarsometatarsus; USTL ITD-595. Figured by Mourer-Chauviré 1983, pl. 4, fig. 12.]

Agnopterus itardiensis (Mourer-Chauviré): Mlíkovský, this paper [New combination.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 2000h sub *I. cf. itardiensis*); Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1983); middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1983), and Roqueprune 2, France (Mourer-Chauviré 1983); late Oligocene (MP 25) of Belgarric 4A, France (Mourer-Chauviré 1983); and late Oligocene (MP 28) of Fraysse, France (Mourer-Chauviré 1983).

Genus *Propelargus* Lydekker

Propelargus Lydekker, 1891a: 65 [Type by original designation: *Propelargus cayluxensis* Lydekker, 1891a.]

Geranopsis Milne-Edwards, 1892: 72 [Type by monotypy: *Geranopsis elatus* Milne-Edwards, 1892. This is a junior homonym of *Geranopsis* Lydekker, 1891a.]

Occitaniavis Mourer-Chauviré, 1983: 121 [Type by original designation: *Geranopsis elatus* Milne-Edwards, 1892.]

Oblitavis Mourer-Chauviré, 1983: 124 [Type by original designation: *Oblitavis insolitus* Mourer-Chauviré, 1983.]

***Propelargus cayluxensis* Lydekker**

Propelargus cayluxensis Lydekker, 1891a: 66 [Holotype from Quercy: distal end of right tarsometatarsus; BMNH A-109. Figured by Lydekker 1891a, fig. 16.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Lydekker 1891a, Mourer-Chauviré 1983).

***Propelargus elatus* (Milne-Edwards)**

Geranopsis elatus Milne-Edwards, 1892: 72 [Holotype from Quercy: distal end of right tibiotarsus; MNHN QU-15601. Figured by Mourer-Chauviré 1983, pl. 5, fig. 7-8.]

Occitaniavis elatus (Milne-Edwards): Mourer-Chauviré 1983: 122 [New combination.]

Propelargus elatus (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Cracraft 1973: 85, Mourer-Chauviré 1983).

Remarks: Milne-Edwards (1892: 72) described this species as being more similar to cranes (family Gruidae) than to other birds. Gaillard (1908: 82) included *Geranopsis elatus* Milne-Edwards in the synonymy of *Geranopsis hastingsiae* Lydekker (see also Lambrecht 1933: 517). Cracraft (1973: 85) resurrected the species, but relegated it to the Aves incertae sedis. Mourer-Chauviré (1983: 122) included this species in the Cariamidae, subfamily Idiornithinae. I found the holotypical tibiotarsus of this species to be strikingly similar to the same element of *Propelargus cayluxensis* Lydekker. Hence, I transfer here the species to the genus *Propelargus* Lydekker, synonymizing at the same time *Geranopsis* Milne-Edwards and *Occitaniavis* Mourer-Chauviré with that genus.

***Propelargus insolitus* (Mourer-Chauviré)**

Oblitavis insolitus Mourer-Chauviré, 1983: 126 [Holotype from Quercy: right humerus; ML PQ-1201. Figured by Mourer-Chauviré 1983, pl. 5, fig. 5-6.]

Propelargus insolitus (Mourer-Chauviré): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1983).

Remarks: Mourer-Chauviré (1983: 126) stated that this species could be a small representative of either *Propelargus* Lydekker, 1891a or *Occitaniavis* Mourer-Chauviré, 1983. Because *Occitaniavis* Mourer-Chauviré is synonymous with *Propelargus* Lydekker (see above), I include *Oblitavis insolitus* Mourer-Chauviré in the latter genus as well. *Oblitavis* Mourer-Chauviré becomes herewith a synonym of *Propelargus* Lydekker.

Genus *Aenigmavis* Peters

Aenigmavis Peters, 1987a: 72 [Type by original designation: *Aenigmavis sapea* Peters, 1987a.]

***Aenigmavis sapea* Peters**

Aenigmavis sapea Peters, 1987a: 73 [Holotype from Messel: incomplete skeleton in two slabs; coll. Maschwitz, uncatalogued. Figured by Peters 1987a, fig. 1-11.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Peters 1987).

Genus *Ameghinornis* Mourer-Chauviré

Ameghinornis Mourer-Chauviré, 1981a: 638 [Type by original designation: *Strigogyps minor* Gaillard, 1939.]

***Ameghinornis minor* (Gaillard)**

Strigogyps minor Gaillard, 1939: 10 [Holotype from Quercy: left humerus; MNHN 16293. Figured by Gaillard 1939, fig. 4, Mourer-Chauviré 1981, pl. 1, fig. 1-2.]

Ameghinornis minor (Gaillard): Mourer-Chauviré 1981a: 639 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1939, Mourer-Chauviré 1981a); and middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1981a). See also Buffetaut & Rage (1982) and Mourer-Chauviré (1982b).

Genus incertae sedis**"*Diatryma*" *cotei* Gaillard**

Diatryma (?) *Côtei* Gaillard, 1937a: 113 [Holotype from Lissieu: distal end of left tarsometatarsus (in a bad state), middle trochlea of left tarsometatarsus (identification of the bone uncertain), two phalanges I digiti, distal end of a phalanx II digiti, and eight ungual phalanges; ML L-71 (tarsometatarsus), L-72 (?trochlea of a tarsometatarsus), L-66 and L-67 (phalanges I), L-76 (phalanx II), and L-69, 70, 73-75, 77, 88 (phalanges III). Figured by Gaillard 1937, fig. 1, 1a, 8, 8a (tarsometatarsus), 2, 2a-c, 9, 9a (trochlea), 3, 3a (phalanx I digiti III), 4, 4a (phalanx I digiti II), 5-6 (ungual phalanges), and 10 (composed skeleton).]

Diatryma cotei Gaillard: Brodkorb 1967: 144 [Spelling emended.]

Distribution: Middle Eocene (MP 14) of Lissieu, France (Gaillard 1937, 1939); and late Eocene (MP 16) of Lavergne, France (Mourer-Chauviré 1983: 128 sub Phorusrhacidae indet.; tentatively referred – Mlíkovský orig.). Brodkorb (1967: 145) attributed to this species also records from the middle Eocene (MP 11) of Messel, Germany (Berg 1965 sub *Diatryma* sp.), and Geiseltal XIII and XIV, Germany (Fischer 1962 sub *Diatryma* sp.). All German records belong to *Diatryma sarasini* Schaub (see above).

Remarks: Andors (1988, 1992) observed that this is not a diatryma, and placed the species in Aves incertae sedis.

Family Cuculidae Wagler

Cuculidae Wagler, 1830 [Modern family.]

Musophagidae Bonaparte, 1831 [Modern family.]

Apopempsidae Brodkorb, 1971b: 201 [Type genus: *Apopempsis* Brodkorb, 1971b.]

Veflintornithidae Kašin, 1976: 231 [Type genus: *Veflintornis* Kašin, 1976.]

Distribution: Unidentified "Musophagidae" were reported also from the late Oligocene (MP 28) of Gaimersheim 1, Germany (Ballmann 1970), and the middle Miocene (MN 5) of Vieux-Collonges, France (Ballmann 1972).

Genus *Dynamopterus* Milne-Edwards

Dynamopterus Milne-Edwards, 1892: 64 [Type by monotypy: *Dynamopterus velox* Milne-Edwards, 1892.]

Dynamopterus velox Milne-Edwards

Dynamopterus velox Milne-Edwards, 1892: 64 [Holotype from Quercy: right humerus; MNHN QU-16225. Not figured.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892).

Dynamopterus boulei Gaillard

Dynamopterus Boulei Gaillard, 1939: 19 [Holotype from Quercy: right humerus; MNHN QU-16296. Figured by Gaillard 1939, fig. 8.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1939).

Genus *Veflintornis* Kašin

Apopempsis Brodkorb, 1971: 202 [Type by original designation: *Musophaga meini* Ballmann, 1969a. This is a junior homonym of *Apopempsis* Schenkling, 1903.]

Veflintornis Kašin, 1976: 231 [New name for *Apopempsis* Brodkorb, 1971; hence its junior objective synonym.]

Remarks: The extralimital record of the genus *Veflintornis* Kašin is limited to "*Apopempsis*" *africanus* Harrison from the early Miocene of Kenya (Harrison 1980d), if the latter species was correctly identified to the genus.

Veflintornis meini (Ballmann)

Musophaga meini Ballmann, 1969a: 188 [Holotype from Grive-Saint-Alban – fissure M: proximal end of right carpometacarpus; FSL 62229 (formerly coll. Mein 118). Figured by Ballmann 1969a, pl. 13, fig. 5-9.]

Apopempsis meini (Ballmann): Brodkorb 1971: 202 [New combination.]

Veflintornis meini (Ballmann): Kašin 1976: 231 [New combination.]

Distribution: Middle Miocene (MN 8) of Grive-Saint-Alban – fissure M, France (Ballmann 1969a).

Genus *Cuculus* Linnaeus

Cuculus Linnaeus, 1758 [Modern genus.]

Cuculus csarnotanus Jánossy

Cuculus csarnotanus Jánossy, 1979: 25 [Holotype: distal end of left humerus from Csarnóta 2 – layer 3; NMB Vt-80. Figured by Jánossy 1979, fig. 4/6.]

Distribution: Early Pliocene (MN 15) of Csarnóta 2 – layer 3, Hungary (Jánossy 1979).

***Cuculus canorus* Linnaeus – Common Cuckoo**

Cuculus canorus Linnseus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001); and (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972), and Stránská skála – Musil's talus cone, layer 6, Czechia (Mlíkovský 1995b).

***Cuculus* sp.**

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b, 1990b, 1992).

Family Accipitridae Vigors

Accipitridae Vigors, 1824 [Modern family.]

Sagittariidae Finsch & Hartlaub, 1870 [Modern family.]

Remarks: All Tertiary forms listed below are in need of revision, excepting the genera *Polemaetus* Heine and *Pelargopappus* Stejneger.

Genus *Palaeocircus* Milne-Edwards

Palaeocircus Milne-Edwards, 1871: 454 [Type by monotypy: *Palaeocircus cuvieri* Milne-Edwards, 1871.]

***Palaeocircus cuvieri* Milne-Edwards**

Palaeocircus Cuvieri Milne-Edwards, 1871: 454 [Holotype from Montmartre: right carpometacarpus; MNHN 7988. Figured by Cuvier 1822, pl. 75, fig. 3, Cuvier 1836, pl. 156, fig. 3, Milne-Edwards 1869-1871, pl. 185, fig. 16, and Brunet 1970, pl. A, fig. a-b.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a: 22, Harrison & Walker 1976a: 339); and late Eocene (MP 19) of Montmartre, France (Cuvier 1822: 317, 326 sub "Oiseau de proie de la grandeur du Balbuzard" = bird of the size of *Pandion haliaeetus*, Milne-Edwards 1869-1871, Brunet 1970: 44, Harrison 1979b).

Genus *Palaeohierax* Milne-Edwards

Palaeohierax Milne-Edwards, 1871: 156 [Type by monotypy: *Palaeohierax gervaisii* Milne-Edwards, 1863.]

***Palaeohierax gervaisii* (Milne-Edwards)**

Aquila gervaisii Milne-Edwards, 1863: 156 [Holotype from Chaptuzat: left tarsometatarsus; MNHN Av-2832. Figured by Gervais 1848-1852, pl. 50, fig. 3-3a, Gervais 1859, pl. 50, fig. 3-3a, and Milne-Edwards 1869-1871, pl. 183, fig. 1-6.]

Palaeohierax gervaisi (Milne-Edwards): Milne-Edwards 1871: 456 [New combination.]

Distribution: Late Oligocene (MP 25) of Chaptuzat, France (Gervais 1844: 22 sub "Oiseau de proie", Gervais 1848-1852: 234 and 1859: 414 sub "*Aquila* ou *Pandion*", Milne-Edwards 1863, 1869-1871); and early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871: 457). Lambrecht (1933: 406) mentioned that Lydekker (1891) attributed two pedal phalanges from the Eocene of Hordwell (= Hordle) to this species. This is obviously an error, because Lydekker's action refers to *Palaeocircus cuvieri* Milne-Edwards, as correctly listed by Lambrecht (1933: 406 above).

Genus *Pernis* Cuvier

Pernis Cuvier, 1817 [Modern genus.]

***Pernis apivorus* (Linnaeus) – European Honey Buzzard**

Falco apivorus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *P.* cf. *apivorus*).

Genus *Milvus* Lacépède

Milvus Lacépède, 1799 [Modern genus.]

***Milvus milvus* (Linnaeus) – Red Kite**

Falco Milvus Linnaeus, 1758 [Modern species.]

Corvus crassipennis Giebel, 1847: 17 [Holotype from Seveckenberg: distal end of left radius; MNHN, uncatalogued. Not figured.]

Distribution: Late Pleistocene (MQ 2C) of Seveckenberg, Germany (Giebel 1847; Mlíkovský orig.).

Remarks: I restudied the holotypical radius of *Corvus crassipennis* Giebel, finding it inseparable from the same element of the modern *Milvus milvus* (Linnaeus). Accordingly, I synonymize here *Corvus crassipennis* Giebel with the latter species.

Genus *Haliaeetus* Savigny

Haliaeetus Savigny, 1809 [Modern genus.]

***Haliaeetus piscator* Milne-Edwards**

Haliaeetus [sic!] *piscator* Milne-Edwards, 1871: 464 [Holotype from Sansan: proximal end of left carpometacarpus; MNHN Sa-14034. Figured by Milne-Edwards 1868-1871, pl. 185, fig. 9-11; and Cheneval 2000, fig. 2.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871, Cheneval 2000).

Remarks: There is no evidence, that this species belongs to the genus *Haliaeetus* Savigny.

***Haliaeetus albicilla* (Linnaeus) – White-tailed Eagle**

Falco albicilla Linnaeus, 1758 [Modern species]

Haliaeetus [sic!] *angustipes* Jánossy, 1983: 257 [Holotype from Přezletice: distal end of left tarsometatarsus; formerly in coll. Fejfar, probably destroyed – O. Fejfar, pers. communication, 1999. Figured by Jánossy 1983, pl. 2, fig. 3-4.]

Distribution: Late Pliocene (MN 17) of Tegelen, Holland (Junge 1953); early Pleistocene (MQ 1a) of Sima del Elefante – layer E-10, Spain (Rosas et al. 2001), and Untermassfeld, Germany (Jánossy 1997 sub *H.* aff. *brevipes* [sic!]); and early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub *Haliaeetus* sp.), Přezletice, Czechia (Jánossy 1983 sub *H.* *angustipes*; Mlíkovský 1997c), and Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *H.* aff. *albicilla*).

Remarks: *Haliaeetus angustipes* Jánossy was synonymized with *Haliaeetus albicilla* (Linnaeus) by Mlíkovský (1997c). Jánossy (1988, 1997) consistently misspelled *angustipes* as *brevipes*. In spite of that, there is no indication in his writings, that he intended to rename the species. Hence, *brevipes* has no standing in zoological nomenclature.

Genus *Gypaetus* Storr

Gypaetus Storr, 1784 [Modern genus.]

Gypaetus barbatus* (Linnaeus) – LammergeierVultur barbatus* Linnaeus, 1758 [Modern species.]**Distribution:** Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001).**Genus *Gyps* Savigny***Gyps* Savigny, 1809 [Modern genus.]***Gyps melitensis* Lydekker**

Gyps melitensis Lydekker, 1890: 404 [Syntypes from Zebbug (as listed by Lydekker 1891a: 29-32): five cervical vertebrae (BMNH 49354), proximal end of right femur (BMNH 49355), distal end of right femur (BMNH 49355a), distal ends of three right tibiotarsi (BMNH 49356, BMNH 49357, BMNH 49359), distal end of left tibiotarsus (BMNH 49360), distal ends of two left tarsometatarsi (BMNH 49363, BMNH 49363a), 3rd trochlea of a tarsometatarsus (BMNH 49363b), and 10 pedal phalanges (BMNH 49364). Figured by Lydekker 1890, pl. 35, fig. 1 (tibiotarsus 49359), 2, 2a (tibiotarsus 39356), 4, 4a (femur 49355a), 6 (tarsometatarsus 49363), 8 (pedal phalanx 49364a), and 9 (pedal phalanx 49364f), pl. 36, fig. 7, 7a-b (vertebra 49354), and Lydekker 1891a, fig. 7 (tibiotarsus 49359).]

Aegyptius melitensis (Lydekker): Tugarinov 1940a: 207 [New combination.]

Distribution: Middle Pleistocene (MQ 2A) of Mosbach, Germany (Mourer-Chauviré 1977, but see Mlíkovský 1998g); late Pleistocene (MQ 2C) of Soulabé – 'galerie profonde', unknown layer, France (Mourer-Chauviré 1975a), Prince – layer B, Monaco (Boule 1919: 307, Mourer-Chauviré 1975a), Coscia – northwestern cave and southern abri, Corsica, France (Bonifay et al. 1998), Zebbug, Malta (Lydekker 1890, 1891a), Ta'Kandja, Malta (Despott 1929 sub *Gyps* sp., Lambrecht 1933: 403 sub *Gyps* sp.), Liko, Crete, Greece (Weesie 1987, 1988), and Simonelli, Crete, Greece (Weesie 1987, 1988); and late Pleistocene (MQ 2D) of Harpons, France (Saint-Périer 1920, Mourer-Chauviré 1975a, Clot & Mourer-Chauviré 1986).

Tyrberg (1998) erroneously listed this species also from Benghisa and Ghaqba, Malta, giving Lambrecht (1933) and Northcote (1992) as a source of this information. However, reports of the latter two authors do not include these data. Similarly, Lambrecht's (1933: 402) statement that Bate (1916) reported on *Gyps melitensis* from Ghar Dalam, Malta, is erroneous (Brodkorb 1964: 278). The following record of *Gyps melitensis* is invalid: middle Pleistocene (MQ 2A) of Hundsheim, Austria (Jánossy 1974b – see Mlíkovský 1998g), and Vértösszölös 2, Hungary (Jánossy 1977, 1986 – see Mlíkovský 1998g), and late Pleistocene (MQ 2C) of Kálmán Lambrecht cave, Hungary (Jánossy 1964; reidentified as *Gyps fulvus* by Jánossy 1977, 1986, see also Mlíkovský 1998g). The record from the late Miocene (MN 11-13) of Novaja Slobodka, Ukraine (Tugarinov 1940a, Burčák-Abramovič 1958) is highly improbable for stratigraphical and zoogeographical reasons (Mlíkovský orig.).

Remarks: The taxonomic status of this species is uncertain. All alleged records from Central Europe were already shown to belong to other species of vultures (Mlíkovský 1998g), while no revision of the syntypes was undertaken thus far. It remains to be shown, whether *Gyps melitensis* Lydekker is different from modern species of vultures.

***Gyps* sp.**

Distribution: Late Pliocene (MN 17) of Váršec, Bulgaria (Boev 1997b, 2000a).

Genus *Aegyptius* Savigny

Aegyptius Savigny, 1809 [Modern genus.]

***Aegyptius mochachus* (Linnaeus) – Eurasian Black Vulture**

Vultur monachus Linnaeus, 1766 [Modern species]

Gyps fulvus spelaeus Friant, 1950c: 1165 [Lectotype from Goyet (selected by Mlíkovský 2000a: 101): incomplete skeleton (= "femelle b" of Friant 1950c, 1951), incl. right coracoid, distal end of left humerus, right radius, shaft of left radius, fragments of an ulna, distal end of left femur, and distal end of left tibiotarsus; IRScNB, uncatalogued. Figured by Friant 1950c, fig. 1 (unnumbered, tibiotarsus), and Friant 1951, fig. 3/i (coracoid), 3/ii (humerus), 3/iii (femur), and 3/iv/1-2 (tibiotarsus), and fig. 4 (radius). Paralectotypes from Goyet (all in IRScNB): fragment of left tibiotarsus of "femelle a", distal end of left humerus of "mâle c", shaft of right tibiotarsus of "mâle d", fragments of an ulna of specimen "e", and fragments of a coracoid, an ulna and ribs of "mâle f". Unfigured.]

Gyps fulvus spelaeus Friant, 1951: 423 [Same type material (and same lectotype – see Mlíkovský 2000a: 101) as for *Gyps fulvus spelaeus* Friant, 1950c. Description of this species was independently published in two journals. The paper by Friant (1950c) was regarded as a preliminary version of Friant (1951), but contains a valid description of the subspecies. This is both a junior homonym and a junior synonym of *Gyps fulvus spelaeus* Friant, 1950c.]

Torgos tracheliotus todei Kleinschmidt, 1953a: 24 [Syntypes from Salzgitter-Lebenstedt: fragmentary sternum, proximal end of left tibiotarsus, fragmentary shaft of left ulna, and fragmentary shaft of left radius; LMB. Figured by Kleinschmidt 1953a, fig. 1 left (sternum), 2 right (sternum), 3 left (tibiotarsus), and 4 (ulna and radius).]

Gyps melitensis aegyptioides Jánossy, 1989: 25 [Lectotype from Repolusthöhle (selected by Mlíkovský 1998g: 25): distal end of left tarsometatarsus; MG 54.625. Figured by Jánossy 1989, pl. 1, fig. 1a-b. Paralectotypes from Repolust Cave: 2 pedal phalanges (MG 54.628 and MG 57.759). Figured by Jánossy 1989, pl. 1, fig 2-3.]

Distribution: Late Pleistocene (MQ 2C) of Goyet, Belgium (Friant 1950c and 1951 sub *G. fulvus spelaeus* – see Mlíkovský 2000a), Salzgitter-Lebenstedt, Germany (Kleinschmidt 1953a,b sub *Torgos tracheliotus todei* – see Mlíkovský 1998g), and Repolusthöhle, Austria (Jánossy 1989 sub *G. melitensis aegyptioides* – see Mlíkovský 1998g).

Remarks: *Torgos tracheliotus todei* Kleinschmidt and *Gyps melitensis aegyptioides* Jánossy were synonymized with the modern *Aegyptius monachus* (Linnaeus) by Mlíkovský (1998g). *Gyps fulvus spelaeus* Friant was synonymized with the modern *Aegyptius monachus* (Linnaeus) by Mlíkovský (2000a).

Genus *Circaetus* Vieillot

Circaetus Vieillot, 1816 [Modern genus.]

***Circaetus* sp.**

Distribution: Late Pliocene (MN 17) of Vărșec, Bulgaria (Boev 1997b: 70, 2000a).

Genus *Circus* Lacépède

Circaetus Lacépède, 1799 [Modern genus.]

***Circus* sp.**

Distribution: Early Pleistocene (MN 1a) of Betfia 2, Romania (Čapek 1917), and Osztramos 2, Hungary (Jánossy 1977).

Genus *Accipiter* Brisson

Accipiter Brisson, 1760 [Modern genus.]

***Accipiter gentilis* (Linnaeus) – Northern Goshawk**

Falco gentilis Linnaeus, 1758 [Modern species.]

Accipiter gentilis brevidactylus Mourer-Chauviré, 1975a: 63 [Holotype from Saint-Estève-Janson – layer B: incomplete, disarticulated skeleton; coll. Bonifay CD66-B-4002, 4004, 4008-4012, 4017, 4026-2029, 4031, 4041, 4045, 4058, 4065-4066, 4071, 4074, 4078, 4091-4092, 4098-4099, 4104, 4108-4109, 4113-4114, 4122, 4129-4130, 4134, 4136-4139. Figured by Mourer-Chauviré 1975a, pl. 22, fig. 6 (carpometacarpus), 7 (coracoid), and 8 (ulna).

Distribution: Early Pliocene (MN 15) of Muselievo, Bulgaria (Boev 1998e: 220 sub *A. cf. gentilis*); early Pleistocene (MQ 1b) of Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972); middle Pleistocene (MQ 2A) of Cimay, France (Mourer-Chauviré 1975a sub *A. g. brevidactylus*), Saint-Estève-Janson – layers B, C, D, E, F and H, France (Mourer-Chauviré 1975a sub *A. g. brevidactylus*); and middle Pleistocene (MQ 2B) of Lunel-Viel – layer 7, France (Mourer-Chauviré 1975a sub *A. g. brevidactylus*), Saint-Sol-Belcastel, France (Philippe et al. 1980 sub *A. g. cf. brevidactylus*).

***Accipiter nisus* (Linnaeus) – Eurasian Sparrowhawk**

Falco Nisus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *A. cf. nisus*).

***Accipiter* sp.**

Distribution: Early Miocene (MN 4b) of Dolnice, Czechia (Mlíkovský, orig., sub *cf. Accipiter*); middle Miocene (MN 6) of Sansan, France (Cheneval 2000 sub aff. *Accipiter*); and late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a).

Genus *Buteo* Lacépède

Buteo Lacépède, 1799 [Modern genus.]

Remarks: Ennouchi (1930: 123) and Ballmann (1969a: 172) mentioned *Buteo arvernensis* Shufeldt, without citing the source of this information. This is probably a misprint for *Bubo arvernensis* Milne-Edwards (a barn owl), but I was not able to find the reason, why Shufeldt was cited as the author of the name.

***Buteo pusillus* Ballmann**

Buteo pusillus Ballmann, 1969a: 173 [Holotype from Grive-Saint-Alban – fissure M: proximal end of right carpometacarpus; FSL, coll. Mein 117. Figured by Ballmann 1969a, pl. 13, fig. 1-4.]

Distribution: Middle Miocene (MN 8) of Grive-Saint-Alban – fissure M, France (Ballmann 1969a).

***Buteo spassovi* Boev**

Buteo spassovi Boev in Boev & Kovačev, 1998: 126 [Holotype from Chadžidimovo: left tibio-tarsus; NMNHS 10190. Figured by Boev & Kovačev 1998, fig. 1, 4.]

Distribution: Late Miocene (MN 11-12) of Chadžidimovo, Bulgaria (Boev & Kovačev 1998).

***Buteo buteo* (Linnaeus) – Common Buzzard**

Falco buteo Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Ževachova Hill, Ukraine (Dubrovo & Kapelist 1979: 88 sub *B. cf. buteo*); and early Pleistocene (MQ 1b) of Sima del Elefante – unknown layer, Spain (Rosas et al. 2001).

***Buteo rufinus* (Cretzschmar) – Long-legged Buzzard**

Falco rufinus Cretzschmar, 1827 [Modern species]

Buteo rufinus jansoni Mourer-Chauviré 1975c: 533 [Holotype from Saint-Estève-Janson – layer H: incomplete skeleton; coll. Bonifay CD66-H base-7381. Figured by Mourer-Chauviré 1975c, pl. 1, fig. 1 (coracoid), 3-4 (humerus), 5 (ulna), 6 (tibiotarsus), 7 (carpometacarpus), 8-9 (femur), 10 tarsometatarsus), 11 (scapula), 12 (pedal phalanx), and 13 (ulnare).]

Distribution: Middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layer H, France (Mourer-Chauviré 1975a, c).

***Buteo* sp.**

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Ennouchi 1930: 57); late Pliocene (MN 17) of Vărșec, Bulgaria (Boev 2000a); late Pliocene (MN 18) of Ca Na Reia, Eivissa, Balearics (Alcover 1989); and early Pleistocene (MQ 1b) of Bacton, England (Harrison 1979d sub *Buteo ?buteo* or *B. buteo/lagopus/rufinus* superspecies).

Genus *Garganoaetus* Ballmann

Garganoaetus Ballmann, 1973: 8 [Type by original designation: *Garganoaetus freudenthali* Ballmann, 1973.]

Distribution: Early Pliocene (MN 14-15) of the island (now peninsula) of Gargano, Italy.

***Garganoaetus freudenthali* Ballmann**

Garganoaetus freudenthali Ballmann, 1973: 9 [Holotype from San Giovannino: right tarsometatarsus; RGM 177924. Figured by Ballmann 1973, text-fig. 1a-g, pl. 1, fig. 1-3.]

Distribution: Early Pliocene (MN 14-15) of Chirò 4, and San Giovannino, Italy (Ballmann 1973, 1976a), and Chirò 20C, Italy (Ballmann 1976a).

***Garganoaetus murivorus* Ballmann**

Garganoaetus murivorus Ballmann, 1973: 17 [Holotype from Chirò 10C: distal end of right tarsometatarsus; RGM 177646. Figured by Ballmann 1973, text-fig. 5a-e, pl. 3, fig. 4-7.]

Distribution: Early Pliocene (MN 14-15) of Chirò 10C, Italy (Ballmann 1973), and Chirò 2S, Chirò 14B, Chirò 20E, and Chirò 24, Italy (Ballmann 1976a).

Genus *Aquila* Brisson

Aquila Brisson, 1760 [Modern genus.]

***Aquila rapax* (Temminck) – Tawny Eagle**

Falco rapax Temminck, 1828 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Vallonnet, France (Mourer-Chauviré 1975a sub *A. cf. rapax*).

***Aquila chrysaetos* (Linnaeus) – Golden Eagle**

Falco Chrysaëtos Linnaeus, 1758 [Modern species]

Aquila chrysaetos bonifaci Mourer-Chauviré, 1975a: 50 [Holotype from Saint-Estève-Janson – layer H below: partial skeleton; coll. Bonifay CD66-H base-4744, 4746, and 4747. Not figured.]

Aquila chrysaetos simurgh Weesie, 1988: 18 [Holotype from Liko cave: left tibiotarsus lacking proximal end; UUT Li-V804. Figured by Weesie 1987, pl. 5, fig. 3a-b., and Weesie 1988, pl. 5, fig. 3a-b.]

Distribution: Early Pleistocene (MQ 1a) of Osztramos 2, Hungary (Jánossy 1977 sub *A. cf. chrysaetos*); middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layers B, C, E/F, F, G, H, France (Mourer-Chauviré 1975a sub *A. chrysaetos bonifacti*); middle Pleistocene (MQ 2B) of Lunel Viel – cave IV (Mourer-Chauviré 1975a sub *A. chrysaetos bonifacti*); and late Pleistocene (MQ 2C) of Liko, Crete, Greece (Weesie 1987 and 1988 sub *A. chrysaetos simurgh*).

***Aquila* sp.**

Distribution: Late Miocene (MN 9-10) of Sokolov, Ukraine (Vojinstvens'kyj 1967); late Miocene (MN 10) of Viladecavalls, Spain (Villalta 1963); late Miocene (MN 11-13) of Kujal'nik, Ukraine (Vojinstvens'kyj 1967); and late Pliocene (MN 17) of Puebla de Valverde, Spain (Adrover et al. 1974), and Väršec, Bulgaria (Boev 1997b: 71, 2000a).

Genus *Hieraaetus* Kaup

Hieraaetus Kaup, 1844 [Modern genus.]

***Hieraaetus edwardsi* (Sharpe)**

Aquila minuta Milne-Edwards, 1871: 463 [Holotype from Sansan: distal end of left tibiotarsus; MNHN Sa-1220. Figured by Milne-Edwards 1869-1871, pl. 185, fig. 5-8. This is a junior primary homonym of *Aquila minuta* Brehm, 1831 = *Hieraaetus pennatus* (Gmelin, 1788).]

Eutolmaetus edwardsi Sharpe, 1899: 262 [New name for *Aquila minuta* Milne-Edwards, 1871.]

Nisaetus edwardsi (Sharpe): Paris 1912: 288 [New combination.]

Hieraaetus [sic!] *edwardsi* (Sharpe): Ammon 1918: 50 [New combination.]

Aquila Edwardsi (Sharpe): Lambrecht 1933: 408 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871, Cheneval 2000); and middle/late Miocene (MN 8-9) of Hostalets de Pierola, Spain (Villalta 1963 sub *A. cf. minuta*).

***Hieraaetus fasciatus* (Vieillot) – Bonelli's Eagle**

Aquila fasciata Vieillot, 1822 [Modern species.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b: 70 sub *Hieraaetus* sp., Boev 2000a sub *H. cf. fasciatus*).

Genus *Polemaetus* Heine

Polemaetus Heine, 1890 [Modern genus.]

***Polemaetus* sp.**

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869, pl. 183, fig. 20-21 as an unnamed raptor, Mlíkovský 1999e, orig.); and early Miocene (MN 3) of Tuchořice, Czechia (Mlíkovský orig.).

Genus *Pelargopappus* Stejneger

Pelargopsis Milne-Edwards, 1868: 460 [Type by monotypy: *Pelargopsis magnus* Milne-Edwards, 1868. This is a junior homonym of *Pelargopsis* Gloger, 1841.]

Pelargopappus Stejneger, 1885: 163 [New name for *Pelargopsis* Milne-Edwards, 1868; hence its junior objective synonym.]

Pelargodes Lydekker, 1891b: 477 [New name for *Pelargopsis* Milne-Edwards, 1868; hence its junior objective synonym.]

Pelargocrex Milne-Edwards, 1893: liv [New name for *Pelargopsis* Milne-Edwards, 1868; hence its junior objective synonym.]

Amphiserpentarius Gaillard, 1908: 44 [Type by monotypy: *Amphiserpentarius schlosseri* Gaillard, 1908.]

Amynoptilon Cracraft & Rich, 1972: 278 [Type by original designation: *Serpentarius robustus* Milne-Edwards, 1868.]

***Pelargopappus magnus* (Milne-Edwards)**

Pelargopsis magnus Milne-Edwards, 1868: 460 [Holotype from Saint-Gérard-le-Puy: distal end of right tarsometatarsus; MNHN Av-8735. Figured by Milne-Edwards, 1867-1868, pl. 72, fig. 4-7, and Mourer-Chauviré & Cheneval 1983, pl. 1, fig. 4a-b. Brodkorb (1963c: 287) and Mourer-Chauviré & Cheneval (1983: 446) erroneously believed, that the holotype is a lectotype designed by Lydekker (1891a: 68), but Milne-Edwards (1868: 460) clearly based the species on the holotypical tarsometatarsus, and Lydekker (1891a: 68) correctly referred to it as to "the type tarso-metatarsus".]

Serpentarius robustus Milne-Edwards, 1870: 557 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus (not left as stated by Mourer-Chauviré & Cheneval 1983: 446); MNHN Av-2816. Figured by Milne-Edwards 1869-1871, pl. 186, fig. 1-6, Lambrecht 1933, fig. 132b, Rich & Cracraft 1972, fig. 10a-c, Jollie 1977b, fig. 127B/a-g, and Mourer-Chauviré & Cheneval 1983, pl. 1, fig. 1, 2a-b, 3a-b.]

Pelargopappus magnus (Milne-Edwards): Stejneger 1885: 163 [New combination.]

Pelargodes magnus (Milne-Edwards): Lydekker 1891b: 477 [New combination.]

Pelargocrex magnus (Milne-Edwards): Milne-Edwards 1893: liv [New combination.]

Amphiserpentarius Schlosseri Gaillard, 1908: 45 [Holotype from Quercy: distal end of left tibiotarsus; BSP 1 - apparently destroyed in the World War II - Mlíkovský, orig. Cast in FSL PQ-1045. Figured by Gaillard 1908, text-fig. 7, pl. 2, fig. 5-8, and Rich & Cracraft 1972, fig. 5a-d.]

Pelargopsis Stehlini Gaillard, 1908: 82 [Holotype from Quercy: distal end of right tarsometatarsus; BaM QH-146. Figured by Gaillard 1908, text-fig. 21, pl. 4, fig. 5-8.]

Pelargopsis Trouessarti Gaillard, 1908: 84 [Holotype from Quercy: distal end of left tarsometatarsus; BaM QH-147. Figured by Gaillard 1908, text-fig. 22, pl. 4, fig. 9-12.]

Pelargodes Stehlini (Gaillard): Lambrecht 1933: 322 [New combination.]

Pelargodes Trouessarti (Gaillard): Lambrecht 1933: 323 [New combination.]

Amphiserpentarius robustus (Milne-Edwards): Lambrecht 1933: 400 [New combination.]

Pelargopappus stehlini (Gaillard): Brodkorb 1963c: 287 [New combination.]

Pelargopappus trouessarti (Gaillard): Brodkorb 1963c: 287 [New combination.]

Amynoptilon robustum (Milne-Edwards): Cracraft & Rich 1972: 278 [New combination.]

Pelargopappus schlosseri (Milne-Edwards): Mourer-Chauviré & Cheneval 1983: 448 [New combination.]

Distribution: Middle Oligocene (MP 23) of Roqueprune 2, France (Mourer-Chauviré & Cheneval 1983); late Oligocene (MP 28) of Desse, France (Mourer-Chauviré & Cheneval 1983) and Fraysse, France (Mourer-Chauviré & Cheneval 1983); and early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1867-1868, 1869-1871, Gaillard 1908, Mourer-Chauviré & Cheneval 1983).

Remarks: The species *magnus*, *stehlini* and *trouessarti* were originally described in the Ciconiidae, while *robustus* and *schlosseri* in the Sagittariidae. Cracraft & Rich (1972) transferred *schlosseri* in the Cathartidae. Mourer-Chauviré & Cheneval (1983) observed that all these species are based on bones of sagittariid birds, recognizing *Pelargopappus schlosseri* (incl. *stehlini* and *trouessarti*) from Saint-Gérard-le-Puy and *Pelargopappus magnus* (incl. *robustus*) from Quercy. The alleged differences in size between the two species are negligible (see Tab. 1 in Mourer-Chauviré & Cheneval 1983: 451), so that I synonymize here *Serpentarius robustus* Milne-Edwards, *Amphiserpentarius schlosseri* Gaillard, *Pelargopsis stehlini* Gaillard and *Pelargopsis trouessarti* Gaillard with *Pelargopappus magnus* (Milne-

Genus incertae sedis

Aquila corroyi Gaillard

Aquila Corroyi Gaillard, 1939: 3 [Holotype from Quercy: right tarsometatarsus; FSUM, uncatalogued. Figured by Gaillard 1939, fig. 1.]

Aquilavus corroyi (Gaillard): Brodkorb 1964: 263 [New combination; in *Aquilavus* Brodkorb, 1952.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1939).

Aquila delphinensis Gaillard

Aquila Delphinensis Gaillard, 1939: 35 [Holotype from Grive-Saint-Alban – early collections: proximal end of left tarsometatarsus; ML LGr-6030. Figured by Gaillard 1939, fig. 16.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Gaillard 1939).

Aquila depredator Milne-Edwards

Aquila depredator Milne-Edwards, 1871: 458 [Holotype: right tarsometatarsus from Saint-Gérard-le-Puy; MNHN Av-2817. Figured by Milne-Edwards 1869-1871, pl. 184, fig. 5-10. Brodkorb's (1964: 263) indication that Lydekker (1891: 25) selected the tarsometatarsus as a lectotype of this species is erroneous.]

Eutolmaëtus depredator (Milne-Edwards): Sharpe 1899. [New combination; fide Lambrecht 1921: 32, 1933: 407.]

Nisaëtus depredator (Milne-Edwards): Paris 1912: 288 [New combination.]

Aquilavus depredator (Milne-Edwards): Lambrecht 1933: 407 [New combination; in *Aquilavus* Lambrecht, 1933.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871).

Remarks: Genus-group name *Aquilavus* was established by Lambrecht (1933: 407) expressly for a collective group, consisting of "alle bisher beschriebenen paläogenen Raubvogelreste des europäischen Paläogens" = "all Paleogene raptor remains from the European Paleogene, described thus far" (the forms from Saint-Gérard-le-Puy, France, were also included in belief that the locality is late Oligocene in age). Accordingly, *Aquilavus* Lambrecht has no type species, and competes with other genus-group names for homonymy, but not for priority (ICZN 1999). Brodkorb (1952: 175) designed *Aquila depredator* Milne-Edwards, 1871 as the type species of *Aquilavus* Lambrecht. Herewith he created a new genus-group name, *Aquilavus* Brodkorb, 1952, which is a junior homonym of *Aquilavus* Lambrecht, 1933.

Aquila fossilis Giebel

Aquila fossilis Giebel, 1847: 9 [Syntypes from Monte Reale: distal end of humerus, ulna, radius, carpometacarpus, costa, and tarsometatarsus; present location unknown. Figured by Wagner 1832, pl. I, fig. 41a-c (tarsometatarsus), 42 (ulna), 43 (radius), 44 (humerus), pl. II, fig. 45 (humerus), 46a-b (carpometacarpus), 46c-d (costa).]

Distribution: Middle/late Pleistocene (MQ 2A-C) of Monte Reale, Sardinia, Italy (Wagner 1832 sub Falke?, Giebel 1847).

Remarks: Lambrecht (1921: 33-34) erroneously attributed *Aquila fossilis* to Portis (1884: 383) and believed that this is a new combination of *Vultur fossilis* Giebel, but both Giebel (1847) and Portis (1884) clearly understood these forms as separate species. Lambrecht (1933) omitted *Aquila fossilis* Giebel, and Brodkorb (1964: 282) listed it in the synonymy of the modern *Buteo buteo* (Linnaeus) without explanation, which is unsubstantiated.

***Aquila hypogaea* Milne-Edwards**

Aquila hypogaea Milne-Edwards, 1892: 60 [Holotype from Quercy: left femur; MNHN QU-15748. Not figured.]

Aquilavus hypogaea (Milne-Edwards): Lambrecht 1933: 407 [New combination; in *Aquilavus* Lambrecht, 1933.]

Aquilavus hypogaeus (Milne-Edwards): Brodkorb 1964: 263 [New combination; in *Aquilavus* Brodkorb, 1964; and ending corrected.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892).

***Aquila pennatoides* Gaillard**

Aquila pennatoïdes Gaillard, 1939: 37 [Holotype from Grive-Saint-Alban – early collections: proximal end of left tarsometatarsus; ML LGr-6029. Figured by Gaillard 1939, fig. 17.]

Aquila pennatoïdes Gaillard: Mlíkovský, this paper [Spelling emended. This spelling appeared already in an upper paragraph on p. 76 in Gaillard 1939. However, Gaillard (1939) consistently used the spelling *pennatoïdes* in his paper, so that I interpret here the “emended” spelling in his paper as a printers error.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Gaillard 1939).

***Aquila prisca* Milne-Edwards**

Aquila prisca Milne-Edwards, 1863: 157 [Syntypes from Saint-Gérard-le-Puy: distal end of left tarsometatarsus and “divers os des ailes” = various wing bones. Should be in MNHN, but I was not able to find them there in June 1999. Figured by Milne-Edwards 1869-1871, pl. 184, fig. 1-4 (tarsometatarsus).]

Etolmaëtus priscus (Milne-Edwards): Sharpe 1899. [New combination; fide Lambrecht 1921: 33, 1933: 408.]

Nisaëtus priscus (Milne-Edwards): Paris 1912: 288 [New combination.]

Aquilavus priscus (Milne-Edwards): Lambrecht 1933: 408 [New combination; in *Aquilavus* Lambrecht, 1933.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1869-1871).

***Buthierax pouliani* Kretzoi**

Buthierax pouliani Kretzoi, 1977: 132 [Holotype from Petralona – section A, layer 2: left tarsometatarsus; MP TA-19. Figured by Kretzoi & Poulianos 1981, fig. 1; and Poulianos 1996, photo 10.7.]

Distribution: Middle Pleistocene (MQ 2A) of Petralona – section A, layer 2, Greece (Kretzoi 1977, Kretzoi & Poulianos 1981).

Remarks: *Buthierax pouliani* Kretzoi, 1977 is type by original designation of the genus *Buthierax* Kretzoi, 1977.

***Milvus brachypterus* Jánossy**

Milvus brachypterus Jánossy, 1977: 26 [Holotype from Nagyharsányhegy 1-4: proximal end of left carpometacarpus; present location unknown. Figured by Jánossy 1977, fig. 5/1-2.]

Distribution: Early Pleistocene (MQ 1b) of Nagyharsányhegy 1-4, Hungary (Lambrecht 1916 sub *Archibuteo lagopus*, Jánossy 1977).

***Milvus deperditus* Milne-Edwards**

Milvus deperditus Milne-Edwards, 1871: 461 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus; MNHN Av-2838. Figured by Milne-Edwards 1869-1871, pl. 185, fig. 1-4.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Lambrecht 1933: 418); and early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Wetmore 1956: 3).

***Milvus incertus* Gaillard**

Milvus incertus Gaillard, 1939: 40 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus; ML SGr-217. Figured by Gaillard 1939, fig. 19.]

Promilio incertus (Gaillard): Brodkorb 1964: 274 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Gaillard 1939); and late Miocene (MN 9 or 11) of Bujoru, Moldova (Kuročkin & Ganea 1972: 63).

Order Columbiformes Latham

Columbae Latham, 1790 [Modern order.]

Family Caprimulgidae Vigors

Caprimulgidae Vigors, 1825 [Modern family.]

Steatornithidae Bonaparte, 1842 [Modern family.]

Nyctibiidae Bonaparte, 1853 [Modern family.]

Archaeotrogonidae Mourer-Chauviré, 1980a: 18 [Type genus: *Archaeotrogon* Milne-Edwards, 1892.]

Eocypselidae Harrison, 1984c: 174 [Type genus: *Eocypselus* Harrison 1984c.]

Preficinae Olson, 1987: 59 [Type genus: *Prefica* Olson, 1987: 59]

Remarks: I consider the modern Steatornithidae and Nyctibiidae survivors of ancient radiation(s) of the Caprimulgidae, and include them in the latter family. The taxonomic position of all Tertiary caprimulgids is uncertain. The genera are thus listed here in the chronological sequence.

Olson (1987: 60) referred partial skeleton of a bird from the middle Eocene (MP 11) of Messel (see Olson 1985a, fig. 4) to *Prefica nivea* Olson, 1987, described from the early Eocene (Wasatchian) of Wyoming as a member of the subfamily Preficinae of the family Steatornithidae. Mayr (1999c: 525) observed, that the Messel specimen does not belong to the latter family (see also Mayr & Daniels 2001). Accordingly, *Prefica nivea* Olson must be deleted from the list of the European fossil birds.

The Caprimulgidae have an interesting fossil record in Europe. They are rather common in the early Eocene through early Miocene deposits, being absent from the middle to late Miocene and Pliocene deposits, to re-appear in the early Pleistocene. The early Miocene record is based on undescribed bones from Merkur, Czechia (MN 3, Mlíkovský orig., bone in coll. Dvořák), and Petersbuch 28, Germany (MN 3-4, Mlíkovský orig., bones in coll. Rummel).

Genus *Eocypselus* Harrison

Eocypselus Harrison, 1984c: 161 [Type by original designation: *Eocypselus vincenti* Harrison, 1984c.]

Eocypselus vincenti Harrison

Eocypselus vincenti Harrison, 1984c: 162 [Holotype from Walton-on-the-Naze: associated left humerus, incomplete right humerus, left coracoid, proximal end of left ulna, left radius; BMNH A-5429. Figured by Harrison 1984c, fig. 1e (humerus), and 3e (wing skeleton).]

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Harrison 1984c).

Remarks: Here transferred from the Apodiformes. *Eocypselus* Harrison may be identical with *Archaeotrogon* Milne-Edwards, but drawings of the holotype do not allow for a closer examination.

Genus *Fluvioviridiavis* Mayr & Daniels

Fluvioviridiavis Mayr & Daniels, 2001: 394 [Type by original designation: *Fluvioviridiavis platyrhamphus* Mayr & Daniels, 2001.]

Distribution: Extralimital record is confined to *Fluvioviridiavis platyrhamphus* Mayr & Daniels, 2001 from the early Eocene (Green River Formation) of Wyoming (Mayr & Daniels 2001).

Remarks: Described in order and family incertae sedis. Here included in the Caprimulgidae in the broad sense accepted here (Mlíkovský, orig.).

***Fluvioviridiavis* sp.**

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Mayr & Daniels 2001; tentatively referred); and middle Eocene (MP 11) of Messel, Germany (Mayr & Daniels; tentatively referred).

Genus *Paraprefica* Mayr

Paraprefica Mayr, 1999c: 525 [Type by original designation: *Paraprefica kelleri* Mayr, 1999c.]

Distribution: Middle Eocene (MP 11) of Germany.

Remarks: Described in the Preficinae (Mayr 1999c), and later transferred in the Nyctibiidae (Mayr 2001b).

***Paraprefica kelleri* Mayr**

Paraprefica kelleri Mayr, 1999c: 526 [Holotype from Messel: partial skeleton in slab; SMF ME-3376. Figured by Mayr 1999c, fig. 6.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1999c).

***Paraprefica major* Mayr**

Paraprefica major Mayr, 1999c: 529 [Holotype from Messel: anterior part of the skeleton in slab; SMNS 81654. Figured by Mayr 1999c, fig. 12; and Mayr 2000g, fig. 11.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1999c, 2000b).

Genus *Ventivorus* Mourer-Chauviré

Ventivorus Mourer-Chauviré, 1988a: 34 [Type by original designation: *Ventivorus ragei* Mourer-Chauviré, 1988a.]

Remarks: Described in the Caprimulgidae s. str. (Mourer-Chauviré 1988a).

***Ventivorus ragei* Mourer-Chauviré**

Ventivorus ragei Mourer-Chauviré, 1988a: 35 [Holotype from Bretou: anterior end of right coracoid; USTL BRT-1015. Figured by Mourer-Chauviré 1988a, pl. 2, fig. 1-3.]

Distribution: Late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1988a).

Genus *Archaeotrogon* Milne-Edwards

Archaeotrogon Milne-Edwards, 1892: 64 [Type by monotypy: *Archaeotrogon venustus* Milne-Edwards, 1892.]

Distribution: Late Eocene (MP 22) through late Oligocene (MP 28) of France.

Remarks: *Archaeotrogon* Milne-Edwards was described as a relative of modern trogons (see also Mourer-Chauviré 1980), but Olson (1985a: 128-120) showed that it has caprimulgid affinities.

***Archaeotrogon cayluxensis* (Lydekker)**

Palaeortyx cayluxensis Lydekker, 1891a: 138 [Syntypes from Quercy: right and left ulna; BMNH A-137a (both specimens). Not figured. This is a senior primary homonym of *Palaeortyx cayluxensis* Milne-Edwards, 1892.]

Archaeotrogon venustus Milne-Edwards, 1892: 64 [Holotype from Quercy: right humerus; MNHN QU-15781. Not figured. The humerus figured by Mourer-Chauviré 1980, fig. 3a-b (MNHN QU-15782) is too small for being the holotype (Mlíkovský orig.).]

Archaeotrogon Zitteli Gaillard, 1908: 70 [Holotype from Quercy: left humerus; formerly in BSP 128, apparently destroyed in the World War II – Mlíkovský, orig.; cast in FSL PQ-

1053. Figured by Gaillard 1908, text-fig. 17, pl. 3, fig. 24-25.]

Ludiortyx cayluxensis (Lydekker): Brodkorb 1964: 299 [New combination.]

Archaeotrogon hoffstetteri Mourer-Chauviré, 1980: 28 [Holotype from Quercy: right humerus; MNHN QU-15796. Figured by Mourer-Chauviré 1980, fig. 3t,w.]

Archaeotrogon cayluxensis (Lydekker): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908, Mourer-Chauviré 1980); late Eocene (MP 16-20) of Garrigues, France (Mourer-Chauviré 1980); late Eocene (MP 17) of Perrière, France (Mourer-Chauviré 1980); late Eocene (MP 19) of Escamps A, and Escamps III, France (Mourer-Chauviré 1980); early Oligocene (MP 22) of Plante 2, France (Mourer-Chauviré 1980), and Mas de Got B, France (Mourer-Chauviré 1980); middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1980), Mounayne, France (Mourer-Chauviré 1980), and Roqueprune 2, France (Mourer-Chauviré 1980); late Oligocene (MP 28) of Desse, France (Mourer-Chauviré 1980), and Fraysse, France (Crochet 1971, Mourer-Chauviré 1980); and Oligocene (MP 21-28) of Belgarite 4A, France (Mourer-Chauviré 1980), Boussac 1, France (Mourer-Chauviré 1980), Boussac 2, France (Mourer-Chauviré 1980), and Fonbonne 1, France (Mourer-Chauviré 1980).

Remarks: These three alleged species of *Archaeotrogon* Milne-Edwards fall in the same size category. Mourer-Chauviré (1992b: 83) reexamined the syntypes of *Palaeortyx cayluxensis* Lydekker, concluding that they belong to an *Archaeotrogon* Milne-Edwards of the size of *Archaeotrogon zitteli* Gaillard. Accordingly, I synonymize here *Archaeotrogon venustus* Milne-Edwards, *Archaeotrogon zitteli* Gaillard and *Archaeotrogon hoffstetteri* Mourer-Chauviré with *Archaeotrogon cayluxensis* (Lydekker), which has priority.

***Archaeotrogon nocturnus* n. n.**

Archaeotrogon Cayluxensis Gaillard, 1908: 67 [Holotype from Quercy: right humerus; FSL PQ-2. Figured by Gaillard 1908, text-fig. 15, pl. 4, fig. 1-4, and Mourer-Chauviré 1980, fig. 4k-l. This is junior secondary homonym of *Palaeortyx* (= *Archaeortogon*) *cayluxensis* Lydekker, 1891a.]

Archaeotrogon nocturnus Mlíkovský, this paper [New name for *Archeotrogon cayluxensis* Gaillard, 1908; hence its junior objective synonym.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1908, 1939, Mourer-Chauviré 1980); early Oligocene (MP 25) of Garouillas, France, and late Oligocene (MP 28) of Fraysse, France (Mourer-Chauviré 1980).

Remarks: *Archaeotrogon cayluxensis* Gaillard, 1908 is preoccupied by *Archaeotrogon cayluxensis* (Lydekker, 1891a). Hence, a new name for the species is proposed here. The species name (lat. nocturnus = nocturnal) is selected in allusion of the nocturnal habits of modern nightjars.

Genus *Euronyctibius* Mourer-Chauviré

Euronyctibius Mourer-Chauviré, 1989a: 2052 [Type by original designation: *Euronyctibius kurochkini* Mourer-Chauviré, 1989a.]

Remarks: Described in the Nyctibiidae.

***Euronyctibius kurochkini* Mourer-Chauviré**

Euronyctibius kurochkini Mourer-Chauviré, 1989a: 2053 [Holotype from Quercy: proximal end of right humerus; MNHN QU-16231. Figured by Mourer-Chauviré 1989a, fig. 2/8-9.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1989a).

Genus *Caprimulgus* Linnaeus

Caprimulgus Linnaeus, 1758 [Modern genus.]

***Caprimulgus europaeus* Linnaeus – European Nightjar**

Caprimulgus europaeus Linnaeus, 1758 [Modern species.]

Caprimulgus europaeus fossilis Čapek, 1917: 29 [Holotype from Betfia 2: left coracoid; GIB Vt-63. Not figured.]

Caprimulgus capeki Jánossy, 1977: 21 [Holotype from Betfia 2: left coracoid; GIB Vt-63. Not figured. This is a junior objective synonym of *Caprimulgus europaeus fossilis* Čapek, 1917.]

Caprimulgus fossilis Čapek: Tyrberg 1998: 552 [New rank.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1977); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub aff. *C. europaeus*, Jánossy 1977 sub *C. capeki*; reidentified as *Caprimulgus europaeus* by Mlíkovský, orig.).

Remarks: Jánossy (1977) regarded Čapek's name for this species as unavailable for nomenclatural purposes, probably because it was proposed conditionally. However, such names are available if published before 1961 (ICZN 1999, Art. 15). In the taxonomic treatment of the form I follow Čapek (1917), and I synonymize accordingly *Caprimulgus capeki* Jánossy with the modern *Caprimulgus europaeus* Linnaeus.

Family Aegialornithidae Lydekker

Aegialornithidae Lydekker, 1891a: 182 [Type genus: *Aegialornis* Lydekker, 1891a: 183.]

Hemiprocnidae Oberholser, 1906 [Modern family.]

Primapinae Harrison, 1984c: 174 [Type genus: *Primapus* Harrison & Walker, 1975.]

Jungornithidae Karchu, 1988: 80 [Type genus: *Jungornis* Karchu, 1988.]

Aegialornithinae Lydekker: Karchu 1992: 384 [New rank.]

Primapodinae Harrison: Mlíkovský 2000d: 81 [Spelling emended.]

Distribution: Early to late Eocene (MP 8/9-19) of England, France and Germany, with a questionable record from the middle Oligocene (MP 23) of France. Extralimital record includes an unnamed form from the early Eocene of Virginia (Olson 1999b), *Jungornis tesselatus* Karchu, 1988 from the early Oligocene of Adygea in Russia, and modern genus *Hemiprocnis* Nitzsch, 1829 from the SE Asia.

Remarks: Although based on a fossil genus, this family includes modern tree-swifts of the genus *Hemiprocnis* Nitzsch, 1829 (Mlíkovský 1982b, 1985, see also Harrison 1975c, 1984c, Mourer-Chauviré 1988b, Karchu 1988, 1992). The family Hemiprocnidae Oberholser was synonymized with the Aegialornithidae Lydekker by Mlíkovský (1982b: 251). See Oberholser (1906), Brodkorb (1971b: 238-240) and Bock (1994: 143, 188) for the nomenclatural status of the Hemiprocnidae.

I see no reason why to separate the genus *Jungornis* Karchu from the Aegialornithidae at the family level, and I synonymize here Jungornithidae Karchu with Aegialornithidae Lydekker.

The taxonomic status of the family Aegialornithidae (incl. Hemiprocnidae) is uncertain. Brodkorb (1977: 240) included the Hemiprocnidae in the Apodidae as a subfamily. It is well possible, that the aegialornithids represent an early radiation of the Apodidae, and that they should be included in the Apodidae. However, typical aegialornithids and typical apodids were coexistent in the Eocene of Europe, which supports the possibility, that the two groups evolved independently within the Columbiformes (in the present sense). Until this question is solved, I leave Aegialornithidae as a separate family.

Genus *Primapus* Harrison & Walker

Primapus Harrison & Walker, 1975: 162 [Type by original designation: *Primapus lacki* Harrison & Walker, 1975.]

Remarks: Peters (1985) synonymized *Primapus* Harrison & Walker with *Aegialornis* Lydekker, but Karchu (1988, 1992) resurrected the genus (see also Mourer-Chauviré 1988b, and Mayr & Peters 1999).

***Primapus lacki* Harrison & Walker**

Primapus lacki Harrison & Walker, 1975: 162 [Holotype from Bognor Regis: right humerus; BMNH A-2166. Figured by Harrison & Walker 1975, pl. 14, fig. a-f.]

Aegialornis lacki (Harrison & Walker): Peters 1985: 158 [New combination.]

Distribution: Early Eocene (MP 8-9) of Bognor Regis, England (Harrison & Walker 1975, Harrison 1984c), and Warden Point on Sheppey, England (Harrison & Walker 1975, Harrison 1984c).

Genus *Aegialornis* Lydekker

Aegialornis Lydekker, 1891a: 183 [Type by original designation: *Aegialornis gallicus* Lydekker, 1891a.]

Tachyornis Milne-Edwards, 1892: 66 [Type by monotypy: *Tachyornis hirundo* Milne-Edwards, 1892.]

Belornis Milne-Edwards, 1893: liv [New name for *Tachyornis* Milne-Edwards, 1892, established in the erroneous belief that the name is preoccupied by *Tachornis* Gosse, 1847. *Belornis* Milne-Edwards is a junior objective synonym of *Tachyornis* Milne-Edwards.]

Distribution: Middle Eocene (MP 12) of Germany and late Eocene (MP 16-17) of France.

***Aegialornis germanicus* n. sp.**

Holotype from Geiseltal L: right humerus with damaged proximal end; GM L-9-1969. Figured by Peters 1998, fig. 1 partim.

Diagnosis: An *Aegialornis*, smaller than *Aegialornis gallicus* Milne-Edwards, the only other species known from the genus.

Measurements: See Peters (1998, for the holotype), and Mourer-Chauviré (1988b, for *Aegialornis gallicus*).

Etymology: After Germany, where the holotype was found.

Distribution: Middle Eocene (MP 12) of Geiseltal L, Germany (Peters 1998 sub *A. breweri*, Mlíkovský orig.).

***Aegialornis gallicus* Lydekker**

Aegialornis gallicus Lydekker, 1891a: 183 [Lectotype from Quercy (selected by Harrison 1984c: 163): right humerus; BMNH A-60. Figured by Lydekker, 1891a, fig. 41a. Paralectotypes from Quercy: 12 humeri from the phosphorites of Quercy; BMNH A-60 (all syntypes). Not figured.]

Tachyornis hirundo Milne-Edwards, 1892: 66 [Holotype from Quercy: right humerus; MNHN QU-15473. Not figured.]

Belornis hirundo (Milne-Edwards): Milne-Edwards 1893a: liv [New combination.]

Aegialornis Leehardti Gaillard, 1908: 60 [Lectotype from Quercy (selected by Collins 1976a: 123): right humerus; MNHM, uncatalogued. Figured by Gaillard 1908, text-fig. 11, pl. 3, fig. 3-6, Collins 1976a, fig. 2, 4a. Paralectotype from Quercy: left tarsometatarsus from Quercy; ML. Figured by Gaillard 1908, pl. 3, fig. 1-2. Gaillard (1908) based this species on two syntypes, one of which was erroneously called holotype by Collins (1976: 123, 125), and Mourer-Chauviré (1988b: 387). Collin's action can be interpreted as a selection of the lectotype.]

Aegialornis leenhardtii Gaillard: Brodkorb 1971b: 234 [Spelling emended.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Lydekker 1891a, Milne-Edwards 1892, 1893a,b, Gaillard 1908, 1939, Lowe 1939, Collins 1976a, Harrison 1975c, 1984c, Mourer-Chauviré 1988b); late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1988b), and Lavergne, France (Mourer-Chauviré 1988b); and late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1988b) and Pradigues, France (Mourer-Chauviré 1988b).

Remarks: *Tachyornis hirundo* Milne-Edwards was synonymized with *Aegialornis gallicus* Lydekker by Milne-Edwards (1892: 80). *Aegialornis leenhardtii* Gaillard seems to be based on large individuals of *Aegialornis gallicus* Lydekker. Accordingly, I synonymize here the former with the latter species.

Genus *Mesogiornis* n. g.

Mesogiornis Mlíkovský, this paper [Type by original designation: *Aegialornis wetmorei* Collins, 1976a.]

Diagnosis: Member of the family Aegialornithidae, differing from *Aegialornis* Lydekker in having on humerus: (1) external tuberosity closer to the top of the head, (2) deltoid crest more bend, (3) deltoid crest narrower and more pointed, and (4) ectepicondylar process located more proximally. It differs from *Cypselavus* Gaillard in having shaft of humerus more robust, and from *Primapus* Harrison in having ectepicondylar process located more proximally.

Etymology: "Inland bird", from Greek *μεσογειος* (inland) and *ορνις* (bird). This an opposite to Lydekker's (1891a) *Aegialornis* (= sea-shore bird, from Greek *αιγιαλος*, sea-shore), coined in the erroneous belief, that the bird was related to gulls.

Included species: Type species only.

Mesogiornis wetmorei (Collins)

Aegialornis wetmorei Collins, 1976a: 124 [Holotype from Quercy: right humerus; MNHN QU-15478. Figured by Collins 1976a, fig. 3a, 4c.]

Aegialornis broweri Collins, 1976a: 124 [Holotype from Quercy: imperfect right humerus; MNHN QU-15481. Figured by Collins 1976a, fig. 3b, 4d.]

Mesogiornis wetmorei (Collins): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Collins 1976a, Mourer-Chauviré 1988b); late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1988b); late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1988b), and Pradigues, France (Mourer-Chauviré 1988b); and late Eocene (MP 18) of Sainte-Néboule, France (Mourer-Chauviré 1978b, 1988b).

Remarks: I examined types of both *Aegialornis wetmorei* Collins and *Aegialornis broweri* Collins in MNHN. *Aegialornis broweri* seems to be based on a small (subadult?) individual of *Aegialornis wetmorei*. Accordingly, I synonymize here the former with the latter species.

Genus *Cypselavus* Gaillard

Cypselavus Gaillard, 1908: 63 [Type by monotypy: *Cypselavus gallicus* Gaillard, 1908.]

Palescyvus Karchu, 1988: 86 [Type by original designation: *Palescyvus escampensis* Karchu, 1988.]

Cypselavus gallicus Gaillard

Cypselavus gallicus Gaillard, 1908: 63 [Lectotype from Quercy (selected by Brodkorb 1971b: 239): right humerus; BSP 29 partim, apparently destroyed in the World War II – Mlíkovský, orig. Figured by Gaillard 1908, text-fig. 13, pl. 3, fig. 12-15. Paralectotypes from Quercy: a humerus (BSP 29 partim, apparently destroyed during World War II – Mlíkovský, orig.),

and another humerus; BaM QD-98.]

Palescyvus escampensis Karchu, 1988: 86 [Holotype from Escamps III: left coracoid; UUT, uncatalogued. Figured by Mourer-Chauviré 1988b, pl. 2, fig. 7-8; and Karchu 1988, fig. 3b.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1908, Mourer-Chauviré 1988b); late Eocene (MP 16) of Lavergne, France (Mourer-Chauviré 1988b); late Eocene (MP 17) of Pradigues, France (Mourer-Chauviré 1988b), and Perrière, France (Mourer-Chauviré 1988b); late Eocene (MP 19) of Escamps III, France (Mourer-Chauviré 1988b, Karchu 1988); and middle Oligocene (MP 23) of Crabit (Mourer-Chauviré 1978).

Remarks: *Palescyvus escampensis* was described by Karchu (1988) on the basis of a coracoid, which was originally identified as belonging to *Cypselavus gallicus* Gaillard (Mourer-Chauviré 1978). Karchu (1988) observed that this coracoid is different from same element of *Jungornis tessellatus* Karchu from the early Oligocene of Adygea, but presented no evidence, why the coracoid should not belong to *Cypselavus gallicus* Gaillard. Accordingly, I synonymize here *Palescyvus escampensis* Karchu with *Cypselavus gallicus* Gaillard, and *Palescyvus* Karchu with *Cypselavus* Gaillard.

Family Apodidae Olphe-Galliard

Apodidae Olphe-Galliard, 1887 [Modern family.]

Genus *Scaniacypselus* Harrison

Scaniacypselus Harrison, 1984c: 166 [Type by original designation: *Scaniacypselus wardi* Harrison 1984c.]

Distribution: Early Eocene (MP 8) of Denmark, and middle Eocene (MP 11) of Germany.

Remarks: General information: Dyke 2001b, Mayr 2001e.

Scaniacypselus wardi Harrison

Scaniacypselus wardi Harrison, 1984c: 166 [Holotype from Røsnaes: associated left humerus, left ulna, left carpometacarpus and alular phalanx; BMNH A-5430. Figured by Harrison 1984c, fig. 2d, 4d, 6a-k.]

Distribution: Early Eocene (MP 8) of Røsnaes, Denmark (Harrison 1984c, Mayr & Peters 1999).

Scaniacypselus szarskii (Peters)

Aegialornis szarskii Peters, 1985: 144 [Holotype from Messel: almost complete skeleton in slab; SMNK Me-301. Figured by Peters 1985, fig. 1, 4a-b.]

Primapus szarskii (Peters): Mourer-Chauviré 1988b: 372 [New combination.]

Scaniacypselus szarskii (Peters): Mayr & Peters 1999: 314 [New combination.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Peters 1985, Karchu 1988, Mayr & Peters 1999).

Genus *Procypseloides* Harrison

Procypseloides Harrison, 1984c: 168 [Type by original designation: *Cypselus ignotus* Milne-Edwards, 1871.]

Distribution: Eocene or Oligocene (MP 16-28) of France, and early Miocene (MN 2-3) of France and Czechia.

Procypseloides mourerchauvireae (Mlíkovský)

Cypseloides mourerchauvireae Mlíkovský, 1989a: 60 [Holotype from Quercy: right tibio-tarsus; NHMW 1988/27. Figured by Mlíkovský 1989a, pl. 1, fig. a-d.]

Procypseloides mourerchauvireae (Mlíkovský): Mlíkovský, this paper [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mlíkovský 1989a). The original expectation, that the specimen originated from the late Eocene deposits of Quercy (Mlíkovský 1989a), is not necessarily correct (C. Mourer-Chauviré, in litt., 1990), and the stratigraphic distribution of *Procypseloides* Harrison makes the Oligocene age of this fossil more probable.

Remarks: While describing the species, I followed the taxonomic treatment of mid-Tertiary swifts proposed by Collins (1976b). My subsequent research showed, that Harrison (1984c) is right in separating these swifts at the generic level. Accordingly, I transfer here *Cypseloides mourerchauvireae* Mlíkovský to the genus *Procypseloides* Harrison.

***Procypseloides ignotus* (Milne-Edwards)**

Cypselus ignotus Milne-Edwards, 1871: 394 [Lectotype from Saint-Gérard-le-Puy (here selected): right carpometacarpus; MNHN 2849. Figured by Milne-Edwards 1869-1871, pl. 159, fig. 18-19, and Collins 1976b, fig. 1b. Paralectotype from Saint-Gérard-le-Puy: left ulna; MNHN Av-2851. Figured by Milne-Edwards 1869-1871, pl. 177, fig. 9-13.]

Collocalia incerta Milne-Edwards, 1871: 394 [Holotype from Saint-Gérard-le-Puy: left tibiotarsus; MNHN Av-2862. Figured by Milne-Edwards 1869-1871, pl. 177, fig. 1-9, Collins 1976b, fig. 1c-d.]

Apus ignotus (Milne-Edwards): Paris 1912: 286 [New combination.]

Cypselavus intermedius Gaillard, 1939: 42 [Holotype from Saint-Gérard-le-Puy: left humerus; ML LGr-218. Figured by Gaillard 1939, fig. 20.]

Cypseloides ignotus (Milne-Edwards): Collins 1976b: 129 [New combination.]

Procypseloides ignotus (Milne-Edwards): Harrison 1984c: 169 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1871, Gaillard 1939, Lowe 1939, Collins 1976b, Harrison 1984c), and early Miocene (MN 3) of Merkur, Czechia (Mlíkovský orig.).

Remarks: *Collocalia incerta* Milne-Edwards and *Cypselus ignotus* Gaillard were synonymized with *Cypselus ignotus* Milne-Edwards by Collins (1976b). As judged from figures and published data, the holotype of *Cypselavus intermedius* Gaillard agrees in both morphology and size with *Cypselus* (= *Procypseloides*) *ignotus* Milne-Edwards (Mlíkovský orig.), so that I synonymize here the former with the latter species.

Genus *Apus* Scopoli

Apus Scopoli, 1777 [Modern genus.]

***Apus gaillardi* (Ennouchi)**

Cypselus Gaillardi Ennouchi, 1930: 98 [Holotype from Grive-Saint-Alban: right humerus; ML LGr-47. Figured by Ennouchi 1930, pl. 4, fig. 9-12.]

Apus gaillardi (Ennouchi): Brodkorb 1971: 240 [New combination.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Ennouchi 1930).

***Apus wetmorei* Ballmann**

Apus wetmorei Ballmann, 1976a: 35 [Holotype from Chirò 24: right humerus; RGM 178975. Figured by Ballmann 1976, text-fig. 15a-c, pl. 3, fig. 11-13.]

Apus baranensis Jánossy, 1977: 24 [Holotype from Beremend 5: left ulna; GIB Vt-65. Figured by Jánossy 1977, fig. 5/12.]

Distribution: Early Pliocene (MN 14-15) of Chirò 24, Italy (Ballmann 1976a); late Pliocene (MN 16a) of Beremend 5, Hungary (Jánossy 1977); and late Pliocene (MN 17) of Väršec,

Bulgaria (Boev 1997b, 1998f, 2000e sub *A. baranensis*).

Remarks: *Apus wetmorei* Ballmann was a very small swift. *Apus baranensis* Jánossy falls in the same size class. Because of general agreement in age and locality, I synonymize here the latter with the former species.

***Apus apus* (Linnaeus) – Common Swift**

Hirundo Apus Linnaeus, 1758 [Modern species.]

Apus apus palapus Jánossy, 1974b: 227 [Lectotype from Hundsheim (here selected): right humerus; UWPI 1899/63-partim). Not figured. Paralectotypes: two left coracoids, seven ulnae (5 left, 2 right), right carpometacarpus, proximal end of right carpometacarpus, and distal end of left tibiotarsus; UWPI 1899/63-partim. Not figured.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 1-3, 4c, 5a, and 9a, Czechia (Jánossy 1972), Stránská skála – Absolon's 2nd cave (Jánossy 1972), Stránská skála – Musil's talus cone, layers 3b, 4a, 6, and 13 (Mlíkovský 1995b), and Kövesvárad, Hungary (Jánossy 1977 sub *A. apus* cf. *palapus*); and middle Pleistocene (MQ 2A) of Hundsheim, Austria (Jánossy 1974b sub *A. apus palapus*, Mlíkovský orig.), and Tarkő – layer 1, Hungary (Jánossy 1977 sub *A. apus* cf. *palapus*, Jánossy 1986 sub *A. apus*).

***Apus melba* (Linnaeus) – Alpine Swift**

Hirundo melba Linnaeus, 1758 [Modern species.]

Apus submelba Jánossy, 1972: 56 [Holotype from Tarkő – layer 4: left ulna; NMB V-64.435. Not figured.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *A. melba*); middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layer B, France (Mourer-Chauviré 1975a sub *A. cf. submelba*), Tarkő – layers 2, 3, and 4, Hungary (Jánossy 1972 sub *A. melba*), and Uppony 1 – layer 7, Hungary (Jánossy 1977 sub *A. submelba*).

Remarks: The alleged metrical differences between *Apus submelba* Jánossy and *Apus melba* Linnaeus (see Jánossy 1972: 57) do not substantiate the specific status of *submelba*. Hence, I synonymize here the former with the latter species.

Genus *Chaetura* Stephen

Chaetura Stephen, 1826 [Modern genus.]

***Chaetura baconica* Jánossy**

Chaetura baconica Jánossy, 1977: 22 [Holotype from Sümeg: right ulna; GIB Vt-64. Figured by Jánossy 1977, fig. 5/13.]

Distribution: Late Miocene (MN 11-12) of Sümeg, Hungary (Jánossy 1977).

Remarks: The generic position of this species should be reevaluated.

Family Podargidae Gray

Podargidae Gray, 1840 [Modern family.]

Remarks: This family is traditionally considered as allied to the Caprimulgidae, but the podargids are much more similar to the Strigidae (Mlíkovský orig., see also Mayr 1999c).

Genus *Masillapodargus* Mayr

Masillapodargus Mayr, 1999c: 521 [Type by original designation: *Masillapodargus* Mayr, 1999c.]

***Masillapodargus longipes* Mayr**

Masillapodargus longipes Mayr, 1999c: 522 [Holotype from Messel: incomplete skeleton in slab; SMNK PAL-1083. Figured by Mayr 1999c, fig. 1, 4.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1999c, 2000b).

Family Sophiornithidae Mourer-Chauviré

Sophiornithidae Mourer-Chauviré, 1987: 117 [Type genus: *Sophiornis* Mourer-Chauviré, 1987.]

Distribution: Late Paleocene (MP 6) of France, and Eocene or Oligocene (MP 16-28) of France. While some of the representatives of this family are certainly younger than MP 15, there is no firm evidence, that the family survived into the Oligocene.

Remarks: The taxonomic position of this family requires restudy.

Genus *Berruornis* Mourer-Chauviré

Berruornis Mourer-Chauviré, 1994: 339 [Type by original designation: *Berruornis orbis-antiqui* Mourer-Chauviré, 1994.]

***Berruornis orbisantiqui* Mourer-Chauviré**

Berruornis orbisantiqui Mourer-Chauviré, 1994: 340 [Holotype from Berru: right tarsometatarsus with damaged distal end; MNHN R-4155. Figured by Mourer-Chauviré 1994, pl. 1, fig. 1-5.]

Distribution: Late Paleocene (MP 6) Cernay-les-Reims (incl. Berru), France (Mourer-Chauviré 1994).

Genus *Palaeobyas* Mourer-Chauviré

Palaeobyas Mourer-Chauviré, 1987: 108 [Type by original designation: *Palaeobyas cracrafti* Mourer-Chauviré, 1987.]

***Palaeobyas cracrafti* Mourer-Chauviré**

Palaeobyas cracrafti Mourer-Chauviré, 1987: 109 [Holotype from Quercy: left tarsometatarsus; MNHN QU-15746. Figured by Mourer-Chauviré 1987, pl. 3, fig. 1-4.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1987).

Remarks: Transferred from the Tytoninae by Mlíkovský (1998c).

Genus *Sophiornis* Mourer-Chauviré

Sophiornis Mourer-Chauviré, 1987: 118 [Type by original designation: *Sophiornis quercynus* Mourer-Chauviré, 1987.]

***Sophiornis quercynus* Mourer-Chauviré**

Sophiornis quercynus Mourer-Chauviré, 1987: 119 [Holotype from Quercy: left tarsometatarsus; ML PQ-1202. Figured by Mourer-Chauviré 1987, pl. 4, fig. 1-4.]

Distribution: Eocene or Oligocene of Quercy (MP 16-28), France (Mourer-Chauviré 1987). Tentatively assigned were bone fragments and pedal phalanges from the late Oligocene (MP 25) of Belgarric, France (Mourer-Chauviré 1987: 119).

Genus *Palaeotyto* Mourer-Chauviré

Palaeotyto Mourer-Chauviré, 1987: 109 [Type by original designation: *Palaeotyto cadurcensis* Mourer-Chauviré, 1987.]

***Palaeotyto cadurcensis* Mourer-Chauviré**

Palaeotyto cadurcensis Mourer-Chauviré, 1987: 111 [Holotype from Quercy: left coracoid; MNHN QU-16953. Figured by Mourer-Chauviré 1987, text-fig. 3a-b, pl. 4, fig. 11-12.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy (MP 16-28), France (Mourer-Chauviré 1987, Mlíkovský 1998c).

Family Strigidae Vigors

Strigidae Vigors, 1825 [Modern family.]

Palaeoglaucinae Mourer-Chauviré, 1987: 116 [Type genus: *Palaeoglaux* Mourer-Chauviré, 1987.]

Necrobyinae Mourer-Chauviré, 1987: 116 [Type genus: *Necrobyas* Milne-Edwards, 1892.]

Selenornithinae Mourer-Chauviré, 1987: 116 [Type genus: *Selenornis* Mourer-Chauviré, 1987.]

Remarks: Three successive radiations can be discerned within this family (Mlíkovský 1998c,f), which are understood here as subfamilies. They include the extinct Palaeoglaucinae Mourer-Chauviré (with *Palaeoglaux* Mourer-Chauviré, and perhaps also *Oligostrix* Fischer), modern Tytoninae Ridgway (with *Necrobyas* Milne-Edwards, *Prosybris* Brodkorb, *Basityto* Mlíkovský, and the modern *Tyto* Billberg and *Phodilus* Geoffroy-Saint-Hilaire), and modern Striginae Vigors (with *Mioglaux* Mlíkovský, *Intulula* Mlíkovský, *Alasio* Mlíkovský, and all modern European genera of owls but *Tyto* Billberg and *Phodilus* Geoffroy-Saint-Hilaire). The remaining fossil genera are of uncertain taxonomic position.

Genus *Palaeoglaux* Mourer-Chauviré

Palaeoglaux Mourer-Chauviré, 1987: 113 [Type by original designation: *Palaeoglaux perrierensis* Mourer-Chauviré, 1987.]

Distribution: Middle to late Eocene (MP 11-17) of France and Germany.

***Palaeoglaux artophoron* Peters**

Palaeoglaux artophoron Peters, 1992: 163 [Holotype from Messel: nearly complete postcranial skeleton in slab and counter-slab; SMF ME-1144a,b. Figured by Peters 1992, fig. 1-8.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Peters 1992).

***Palaeoglaux perrierensis* Mourer-Chauviré**

Palaeoglaux perrierensis Mourer-Chauviré, 1987: 116 [Holotype from Perrière: humeral end of left coracoid; USTL PRR-2585. Figured by Mourer-Chauviré 1987, pl. 3, fig. 7-9.]

Distribution: Late Eocene (MP 17) of Perrière, France (Mourer-Chauviré 1987, Peters 1992), and Salême, France (Mourer-Chauviré 1987, Peters 1992).

Genus *Necrobyas* Milne-Edwards, 1892

Necrobyas Milne-Edwards, 1892: 61 [Type by subsequent designation (Richmond 1902: 699): *Necrobyas harpax* Milne-Edwards 1892.]

Paratyto Brodkorb, 1970: 159 [Type by original designation: *Bubo arvernensis* Milne-Edwards, 1863.]

Distribution: Early Oligocene (MP 21) to early Miocene (MN 2a) of France.

Remarks: *Paratyto* Brodkorb was synonymized with *Necrobyas* Milne-Edwards by Mourer-Chauviré (1987: 91).

***Necrobyas harpax* Milne-Edwards**

Necrobyas harpax Milne-Edwards, 1892: 61 [Holotype from Quercy: imperfect left

tarsometatarsus; MNHN QU-15696. Not figured. In an erroneous belief that Milne-Edwards based this species on a series of syntypes, Mourer-Chauviré (1987: 97) selected a left tarsometatarsus from Quercy, MNHN QU-15695, as a lectotype. The "lectotype" was figured by Mourer-Chauviré 1987, pl. 1, fig. 1-4. Mlíkovský (1998c: 250) attempted to identify another left tarsometatarsus, MNHN QU-15742 as the holotype of the species on the basis of size, but Mourer-Chauviré (1999c: 363) showed that Milne-Edwards's description fits to another specimen, which I list here as the holotype of the species. The actions by Mourer-Chauviré (1987) and Mlíkovský (1998c) have no bearing on the nomenclatural status of *Necrobyas harpax*.]

Necrobyas Rossignoli Milne-Edwards, 1892: 63 [Holotype from Quercy: left tarsometatarsus; MNHN QU-15698. Figured by Mourer-Chauviré 1987, pl. 1, fig. 19-20.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908, 1939, Mourer-Chauviré 1987, 1999c, Mlíkovský 1998c, 1999d); early Oligocene (MP 21-28) of Fonbonne 1, France (Mourer-Chauviré 1987), and Phalip, France (Mourer-Chauviré 1987); early Oligocene (MP 21) of Aubrelong 1, France (Mourer-Chauviré 1987), and Ravet-Lupovici, France (Mourer-Chauviré 1987); early Oligocene (MP 22) of Mas de Got, France (Mourer-Chauviré 1987); and middle Oligocene (MP 23) of Crabit, France (Mourer-Chauviré 1987, Mlíkovský 1998c.). The alleged records from the late Eocene (MP17) of Perrière, France (Mourer-Chauviré 1987), and late Eocene (MP 19) of Rosières, France (Mourer-Chauviré 1987) are doubtful at best (Mlíkovský 1998c: 250). General references: Mlíkovský 1998c, 1999d, Mourer-Chauviré 1999c.

Remarks: *Necrobyas rossignoli* Milne-Edwards was synonymized with *Necrobyas harpax* Milne-Edwards by Mlíkovský (1998c).

***Necrobyas arvernensis* (Milne-Edwards)**

Bubo arvernensis Milne-Edwards, 1863: 158 [Lectotype from Saint-Gérard-le-Puy (selected by Mlíkovský 1998c: 250): right tarsometatarsus; MNHN Av-2834b. Figured by Milne-Edwards 1869-1871, pl. 192, fig. 10-15. Paralectotype from Saint-Gérard-le-Puy: right tibiotarsus; MNHN Av-2834a. Figured by Milne-Edwards 1869-1871, pl. 192, fig. 20-23.]

Necrobyas Edwardsi Gaillard, 1939: 8 [Holotype from Quercy: left tarsometatarsus; MNHN QU-16294. Figured by Gaillard 1939, fig. 3, Mourer-Chauviré 1987, pl. 2, fig. 1-2.]

Paratyto arvernensis (Milne-Edwards): Brodkorb 1970: 159 [New combination.]

Necrobyas arvernensis (Milne-Edwards): Mourer-Chauviré 1987: 91 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1939, Mourer-Chauviré 1987, Mlíkovský 1998c); late Oligocene (MP 28) of Desse, France (Mourer-Chauviré 1987), and Fraysse, France (Mourer-Chauviré 1987); and early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1869-1871, Mourer-Chauviré 1987, Mlíkovský 1998c). The record from the middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Shufeldt 1896) is invalid, as the bones belong to *Tyto sanctialbani* Lydekker (Mlíkovský orig.; bones restudied in USNM).

Remarks: *Necrobyas edwardsi* Gaillard was synonymized with *Necrobyas arvernensis* (Milne-Edwards) by Mlíkovský (1998c).

Genus *Prosybris* Brodkorb

Prosybris Brodkorb, 1970: 159 [Type by original designation: *Strix antiqua* Milne-Edwards, 1863.]

Distribution: Middle Oligocene (MP 23) of France, and early Miocene (MN 3-4) of Austria.

***Prosybris media* (Mourer-Chauviré)**

Necrobyas medius Mourer-Chauviré, 1987: 104 [Holotype from Quercy: distal end of right tarsometatarsus; BaM QH-150. Figured by Mourer-Chauviré 1987, pl. 3, fig. 16-17.]

Prosybris media (Mourer-Chauviré): Mlíkovský 1998c: 252 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1908 sub *Asio henrici*, Mourer-Chauviré 1987, 1999c, Mlíkovský 1998c, 1999d).

***Prosybris antiqua* (Milne-Edwards)**

Strix antiqua Milne-Edwards, 1863: 158 [Nomen nudum; no description or indication.]

Strix antiqua Milne-Edwards, 1869: 498 [Holotype from Saint-Gérard-le-Puy: left tarsometatarsus; MNHN Av-2837. Figured by Milne-Edwards 1869-1871, pl. 192, fig. 3-9.]

Prosybris antiqua (Milne-Edwards): Brodkorb 1970: 159 [New combination.]

Necrobyas minimus Mourer-Chauviré, 1987: 105 [Holotype from Fonbonne 1: distal end of right tarsometatarsus; UUT Fo-1. Figured by Mourer-Chauviré 1987, pl. 2, fig. 19-20.]

Distribution: Oligocene (MP 21-28) of Fonbonne 1, France (Mourer-Chauviré 1987, Mlíkovský 1987c); middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1987, Mlíkovský 1998c); early Miocene (MP 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Brodkorb 1970, Mlíkovský 1998c); and early Miocene (MN 3-4) of Limberg, Austria (Bachmayer 1980 sub *Falco* cf. *naumanni*, Mlíkovský 1998c).

Remarks: *Necrobyas minimus* Mourer-Chauviré was synonymized with *Prosybris antiqua* (Milne-Edwards) by Mlíkovský (1998c: 252, see also Mourer-Chauviré 1999c).

Genus *Basityto* Mlíkovský

Basityto Mlíkovský, 1998c: 248 [Type by original designation: *Basityto rummeli* Mlíkovský, 1998c.]

***Basityto rummeli* Mlíkovský**

Basityto rummeli Mlíkovský, 1998c: 248 [Holotype from Grafenmühle 21: left humerus in two pieces; coll. Rummel, not catalogued. Figured by Mlíkovský 1998c, fig. 1a, 2.]

Distribution: Early Miocene (MN 2-3) of Grafenmühle 21, Germany (Mlíkovský 1998c).

Genus *Tyto* Billberg

Tyto Billberg, 1828 [Modern genus.]

***Tyto sanctialbani* (Lydekker)**

Strix sancti-albani Lydekker, 1893: 518 [Syntypes from Grive-Saint-Alban: "no less than eleven ... bones", BMNH. The following syntypes were illustrated by Lydekker 1893, pl. 41: distal end of right tibiotarsus (fig. 1-2), proximal end of left tarsometatarsus (fig. 3, 3a), and distal end of left tarsometatarsus (fig. 4, 4a).

Strix sancti albani Lydekker: Ennouchi 1930: 66 [Spelling emended.]

Strix Sancti-albani Lydekker: Gaillard 1939: 82 [Spelling emended.]

Tyto sanctialbani (Lydekker): Ballmann 1969a: 191 [New combination and spelling emended.]

Tyto campiterrae Jánossy, 1991: 25 [Holotype from Polgárdi 5: left tarsometatarsus; GIB Vt-148-V-18.097. A tarsometatarsus was figured by Jánossy 1991, pl. 3, fig. 1, but Jánossy (1991) listed two complete tarsometatarsi from the type locality and did not state which of them was figured.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban – early collections, France (Lydekker 1893, Shufeldt 1896 sub *Bubo arvernensis*, Ennouchi 1930, Ballmann 1969a, Mlíkovský 1998c); middle Miocene (MN 8) of Grive-Saint-Alban – fissure M, France (Ballmann 1969a); late Miocene (MN 10) of Kohfidisch, Austria (Mlíkovský 1998c); and late Miocene (MN 13) of Polgárdi 5, Hungary (Jánossy 1991, Mlíkovský 1998c). The alleged record of *Tyto sanctialbani* from the early Pliocene (MN 14-15) of Gargano, Italy (Ballmann

1973, 1976a) is invalid, because the bones belong to *Tyto balearica* Mourer-Chauviré et al. (Mlíkovský 1998c).

Remarks: *Tyto campiterrae* Jánossy was synonymized with *Tyto sanctialbani* (Lydekker) by Mlíkovský (1998c).

***Tyto balearica* Mourer-Chauviré et al.**

Tyto balearica Mourer-Chauviré, Alcover, Moyà & Pons, 1980: 804 [Holotype from Canet: left coracoid; MNM, uncatalogued. Figured by Mourer-Chauviré et al 1980, pl. 1, fig. 3-4.]

Distribution: Late Miocene (MN 12) of Aljezar B, Spain (Cheneval & Adrover 1993); early Pliocene (MN 14-15) of several fissures on Gargano, Italy, incl. Biancone, Cantatore 3A, Chirò 2S, Chirò 3, Chirò 15, Chirò 20A,C,D, Chirò 24, Fina D, Gervaisio 2, Rinascita 1, and San Giovannino (Ballmann 1973, 1976a sub *Tyto sanctialbani*, Mlíkovský 1998c); early Pliocene (MN 15) of Layna, Spain (Mourer-Chauviré & Sánchez Marco 1988), and Sète, France (Mourer-Chauviré & Sánchez Marco 1988); late Pliocene (MN 16) of Moreda, Spain (Sánchez Marco 1996, in litt.), Balaruc 2, France (Mourer-Chauviré & Sánchez Marco 1988); late Pliocene (MN 18) of Canet, Mallorca, Balearics (Mourer-Chauviré et al. 1980, Mourer-Chauviré & Sánchez Marco 1988), S'Onix, Mallorca Balearics (Mourer-Chauviré et al. 1980, Mourer-Chauviré & Sánchez Marco 1988), Binigaus, Minorca, Balearics (Mourer-Chauviré et al. 1980, Mourer-Chauviré & Sánchez Marco 1988), and Almenara 1, Spain (Mourer-Chauviré & Sánchez Marco 1988); middle Pleistocene (MQ 2A-C) of Monte Reale, Sardinia (Mlíkovský 2000b), Son Bauçá, Mallorca, Balearics (Ballmann & Adrover 1970 sub *Strix* sp., Mourer-Chauviré et al. 1980: tentatively referred to *T. balearica*); and middle Pleistocene (MQ 2B) of Castiglione 3, Corsica, France (Salotti et al. 1997). For maps see Mlíkovský (1998c, fig. 6, and 2000b, fig. 1).

***Tyto gigantea* Ballmann**

Tyto robusta Ballmann, 1973: 33 [Holotype from Chirò 12: right femur; RGM 177523. Figured by Ballmann 1973, text-fig. 8a-e, pl. 5, fig. 4-6.]

Tyto gigantea Ballmann, 1973: 37 [Holotype from San Giovannino: right tibiotarsus; RGM 177687. Figured by Ballmann 1973, text-fig. 9a-f, pl. 6, fig. 4-7.]

Distribution: Early Pliocene (MN 14-15) of Gargano, Italy, incl. the following fissures: Chirò 2N, Chirò 2S, Chirò 3, Chirò 4, Chirò 5, Chirò 6, Chirò 7, Chirò 9, Chirò 10, Chirò 11, Chirò 12, Chirò 14A, Chirò 20, Chirò 22, Chirò 24, Chirò 27, Falcone 2A, Fina D, Fina H, Gervaisio 1, Gervaisio 2, Monte Granata, Nazario 2B, Nazario 3, Nazario 4, Pirro 11, Pizzioli 4, Posticchia 1B, and San Giovannino (Ballmann 1973, 1976a).

Remarks: *Tyto robusta* Ballmann was synonymized with *Tyto gigantea* Ballmann by Mlíkovský (1998c).

***Tyto alba* (Scopoli) – Barn Owl**

Strix alba Scopoli, 1769 [Modern species.]

Strix melitensis Lydekker, 1891a: 13 [Holotype from an unknown site in Malta: left femur; BMNH 49322. Not figured.]

Tyto melitensis (Lydekker): Brodkorb 1970a: 231 [New combination.]

Distribution: Late Pleistocene (MQ 2C) of an unknown site in Malta (Lydekker 1891a, Mourer-Chauviré et al. 1980, Mlíkovský 1998c).

Remarks: *Strix melitensis* Lydekker was synonymized with the modern *Tyto alba* (Scopoli) by Mlíkovský (1998c: 255, see also Mourer-Chauviré et al. 1980).

Genus *Mioglaux* Mlíkovský

Mioglaux Mlíkovský, 1998f: 6 [Type by original designation: *Mioglaux debellatrix* Mlíkovský, 1998f.]

Distribution: Early Miocene (MN 2-3) of France and Czechia.

***Mioglaux poirrieri* (Milne-Edwards)**

Bubo Poirrieri Milne-Edwards, 1863: 158 [Holotype from Saint-Gérard-le-Puy: right tarsometatarsus; MNHN Av-2836. Figured by Milne-Edwards 1869-1871, pl. 192, fig. 24-29.]

Mioglaux poirrieri (Milne-Edwards): Mlíkovský 1998f: 11 [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1863, 1869-1871, Mlíkovský 1998f).

***Mioglaux debellatrix* Mlíkovský**

Mioglaux debellatrix Mlíkovský, 1998f: 6 [Holotype from Merkur: distal end of right tibiotarsus; coll. Fejfar, uncatalogued. Figured by Mlíkovský 1998f, fig. 1, 2b.]

Distribution: Early Miocene (MN 3) of Merkur, Czechia (Mlíkovský 1998f).

Genus *Intulula* Mlíkovský

Intulula Mlíkovský, 1998f: 8 [Type by original designation: *Intulula tinnipara* Mlíkovský, 1998f.]

Distribution: Early Miocene (MN 3) of Germany and Czechia.

***Intulula tinnipara* Mlíkovský**

Intulula tinnipara Mlíkovský, 1998f: 9 [Holotype from Merkur: left tarsometatarsus lacking distal end; coll. Fejfar, uncatalogued.]

Distribution: Early Miocene (3a) of Wintershof-West, Germany (Ballmann 1969b sub *Otus wintershofensis*, Mlíkovský orig.); and early Miocene (MN 3) of Merkur, Czechia (Mlíkovský 1998f).

Remarks: Ballmann (1969b) referred to his *Otus wintershofensis* a partial humerus. I restudied the bone in BSP, finding that it has a strongly bent shaft, characteristic for the Surniini. It falls in the same size class with *Intulula tinnipara* Mlíkovský, and I refer here the bone to the latter species. *Otus wintershofensis* Ballmann belongs in the Bubonini (see Mlíkovský 1998f, and below).

***Intulula brevis* (Ballmann)**

Strix brevis Ballmann, 1969b: 38 [Holotype from Wintershof-West: left humerus lacking proximal end; BSP 1937-II-18012. Figured by Ballmann 1969b, pl. 1, fig. 7a-b.]

Intulula brevis (Ballmann): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 3) of Wintershof-West, Germany (Ballmann 1969b, Mlíkovský 1998f); and late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1977 sub *Strix* aff. *brevis*, Jánossy 1980b sub *Strix* cf. *brevis*). The record from Hungary is doubtful because of its age (Mlíkovský, orig.).

Remarks: The holotypical humerus of this species clearly belongs in the Surniini, being indeterminate within the latter tribe (Mlíkovský 1998f, and unpublished reexamination of the specimen in BSP). I restudied in BSP the referred material of this species, of which the tarsometatarsus (BSP 1937-II-18113) was especially diagnostic. I found it to be similar in morphology to the same element (holotype) of *Intulula tinnipara* Mlíkovský from the early Miocene (MN 3) of Czechia. Hence, I transfer here *Strix brevis* Ballmann to the genus *Intulula* Mlíkovský. Within the latter genus, *Intulula brevis* (Ballmann) is larger than *Intulula tinnipara* Mlíkovský (proximal width of the tarsometatarsus of *Intulula tinnipara* from Merkur = ca. 5.5 mm, the same dimension of the tarsometatarsus from Wintershof-West = 8.2 mm; Mlíkovský

Genus *Alasio* Mlíkovský

Alasio Mlíkovský, 1998f: 12 [Type by original designation: *Strix collongensis* Ballmann, 1972.]

Alasio collongensis (Ballmann)

Strix collongensis Ballmann, 1972: 97 [Holotype from Vieux-Collonges: left coracoid; FSL 62226. Figured by Ballmann 1972, text-fig. 1a-b, pl. 1, fig. 1-3.]

Alasio collongensis (Ballmann): Mlíkovský 1998f: 12 [New combination.]

Distribution: Middle Miocene (MN 5) of Vieux-Collonges, France (Ballmann 1972, Mlíkovský 1998f).

Genus *Otus* Pennant

Otus Pennant, 1769 [Modern genus.]

Otus scops (Linnaeus) – European Scops Owl

Strix Scops Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995); and early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), and Betfia 2, Romania (Čapek 1917, Jánossy 1977 sub *?O. scops*)

Genus *Bubo* Duméril

Bubo Duméril, 1806 [Modern genus.]

Nyctea Stephens, 1826 [Modern genus.]

Ophthalmomegas Dehaut, 1911: 55 [Type by monotypy: *Ophthalmomegas lamarmorae* Dehaut, 1911.]

Bubo bubo (Linnaeus) – Great Eagle-Owl

Strix bubo Linnaeus, 1758 [Modern species.]

Ardea lignitum Giebel, 1860: 152 [Holotype from Rippersroda 1: left femur; present location unknown. Figured by Giebel 1860, pl. 1, fig. 2.]

Bubo lignitum (Giebel): Brodkorb 1980: 90 [New combination.]

Bubo bubo davidi Mourer-Chauviré, 1975a: 164 [Holotype from Saint-Estève-Janson – layer B: right tibiotarsus; coll. Bonifay CD66-B-11970. Not figured.]

Distribution: Early Pliocene (MN 14) of Vojničevo, Ukraine (Vojinstvens'kyj 1967 sub *B. cf. bubo*); early Pliocene (MN 15) of Csarnóta 2, Hungary (Jánossy 1977 sub *B. aff. bubo*); late Pliocene (MN 16) of Rębielice Królewskie 1A, Poland (Jánossy 1974a sub *?Bubo* sp.); late Pliocene (MN 16b) of Osztramos 7, Hungary (Jánossy 1977 sub *B. aff. bubo*); late Pliocene (MN 18) of Rippersroda 1, Germany (Giebel 1860 sub *Ardea lignitum*, Brodkorb 1980 sub *B. lignitum*, Mlíkovský 1992b), and Villány 3, Hungary (Jánossy 1977 sub *B. aff. bubo*); early Pleistocene (MQ 1a) of Betfia 2, Romania (Jánossy 1977 sub *B. aff. bubo*); early Pleistocene (MQ 1b) of East Runton, England (Harrison 1979d), and Nagyharsányhegy 1-4, Hungary (Jánossy 1977 sub *B. aff. bubo*); and middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layers B, C, D, F, G and H, France (Mourer-Chauviré 1975a sub *B. bubo davidi*, see also Arredondo & Olson 1994), and Vértesszőlős 2, Hungary (Jánossy 1977 sub *B. aff. bubo*).

Remarks: *Ardea lignitum* Giebel was synonymized with the modern *Bubo bubo* (Linnaeus) by Mlíkovský (1992b: 437).

***Bubo zeylonensis* (Gmelin) – Brown Fish Owl**

Strix zeylonensis Gmelin, 1788 [Modern species.]

Ophthalmomegas lamarmorae Dehaut, 1911: 55 [Lectotype from Figari (selected by Pavia 1999a: 5): fragmentary cranium; MPU PU-29860. Figured by Dehaut 1911, pl. 1, fig. 1, Dehaut 1914, pl. 4, fig. 2, Dehaut 1920, fig. 24, and Pavia 1999a, pl. 1, fig. 1, 5. Paralectotypes from Figari: endocranial cast with associated vertebrae (PU-29861), distal end of a humerus (PU-29862), and imprint of a femur (PU-29863). Figured by Dehaut 1911, pl. 1, fig. 2-5.]

Strix? perpasta Ballmann, 1976a: 27 [Holotype from San Giovannino: left tarsometatarsus; RGM 178978. Figured by Ballmann 1976a, text-fig. 10a-c, pl. 7, fig. 9-11.]

Bubo perpastus (Ballmann): Mlíkovský 1998f: 14 [New combination.]

Bubo insularis Mourer-Chauviré & Weesie, 1986: 198 [Holotype from Rapaci: right ulna; UUT Rp-27. Figured by Mourer-Chauviré & Weesie 1986, pl. 1, fig. 1-2.]

Bubo lamarmorae (Dehaut): Pavia 1999a: 3 [New combination.]

Distribution: Early Pliocene (MN 14-15) of Gargano, Italy, incl. fissures Chirò 5, Chirò 20, Pirro 11, and San Giovannino (Ballmann 1976a sub *Strix? perpasta*); late Pleistocene (MQ 2C) of Coscia – northwestern cave, Corsica, France (Bonifay et al. 1998 sub *B. insularis*), Monte San Giovanni, Sardinia, Italy (Lydekker 1891b sub *B. cf. africanus*, Mourer-Chauviré & Weesie 1986 sub *B. insularis*), Pietro Tampoia Cave, Tavolara near Sardinia, Italy (Lydekker 1891b sub *B. cf. africanus*, Mourer-Chauviré & Weesie 1986 sub *B. insularis*), Figari, Sardinia, Italy (Dehaut 1911, 1914 and 1920 sub *O. lamarmorae*, Comaschi Caria 1969 and 1970 sub *Aquila* sp., Pavia 1999a sub *B. lamarmorae*), Corbeddu – hall I, hall II, hall IV: layer 1, Sardinia, Italy (Mourer-Chauviré & Weesie 1986 and Weesie 1999 sub *B. insularis*), Rapaci, Sardinia, Italy (Mourer-Chauviré & Weesie 1986 sub *B. insularis*), Dorgali, Sardinia, Italy (Kotsakis 1980 sub *Strix uralensis*, Mourer-Chauviré & Weesie 1986 sub *B. insularis*), and Macinaggio, Corsica, France (Mourer-Chauviré & Weesie 1986 sub *B. insularis*), Liko – layer D, Crete, Greece (Weesie 1987, 1988). An extralimital record is available from the early Pleistocene (MQ 1a) of 'Ubeidyia – layer II-23, Israel (Tchernov 1980).

Remarks: *Ophthalmomegas lamarmorae* Dehaut was described as a primate by Dehaut (1911, see also Dehaut 1914, 1920). The avian affinities of the lectotypical skull were recognized by S. Schaub & A. Azzaroli (in Comaschi Caria 1969, 1970), and Pavia (1999a) identified the species with *Bubo insularis* Mourer-Chauviré & Weesie. The paratypes of *Ophthalmomegas lamarmorae* Dehaut belong to the primate *Macaca majori* Azzaroli (S. Schaub & A. Azzaroli in Comaschi Caria 1969, see also Pavia 1999a).

Ophthalmomegas lamarmorae Dehaut, *Strix* (= *Bubo*, see Mlíkovský 1998f: 14) *perpasta* Ballmann, and *Bubo insularis* Mourer-Chauviré & Weesie fall in the same size class as the modern *Bubo zeylonensis* (Gmelin) (Mlíkovský, orig.), with which they were formerly not compared, perhaps because the species is often separated in the genus *Ketupa* Lesson, 1831 (but see Ford 1967 for osteological similarities, and König et al. 1999). Hence I synonymize here the former three species with the latter one.

Bubo zeylonensis (Gmelin) is now extinct in Europe, with nearest existing populations in the Middle East (Cramp 1989, Del Hoyo et al. 1999, König et al. 1999).

***Bubo scandiacus* (Linnaeus) – Snowy Owl**

Strix scandiaca Linnaeus, 1758 [Modern species.]

Nyctea scandiaca gallica Mourer-Chauviré, 1975a: 159 [Holotype from Saint-Estève-Janson – layer H: partial disarticulated skeleton; coll. Bonifay CD66-H-4741. Not figured.]

Distribution: Early Pleistocene (MQ 1b) of Bourgade, France (Mourer-Chauviré 1976); middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layers C? and H, France (Mourer-Chauviré 1975a sub *Nyctea scandiaca gallica*), and Fage – layers CO and 40, France (Mourer-Chauviré 1975a,b sub *Nyctea scandiaca gallica*); and late Pleistocene (MQ 2D) of Morin –

layers A1-2, A3, and A4, France (Mourer-Chauviré 1975a sub *Nyctea scandiaca* cf. *gallica*).

***Bubo* sp.**

Distribution: Late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a, tentatively referred).

Genus *Surnia* Duméril

Surnia Duméril, 1806 [Modern genus.]

***Surnia robusta* Jánossy**

Surnia robusta Jánossy, 1977: 10 [Holotype from Villány 3: left tarsometatarsus; NMB Vt-62. Figured by Jánossy 1977, fig. 2a-d, 5/2.]

Distribution: Late Pliocene (MN 16) of Beremend 4, Hungary (Jánossy 1977), and Osztramos 7, Hungary (Jánossy 1977); late Pliocene (MN 18) of Villány 3, Hungary (Jánossy 1977, Mlíkovský 1998f); and early Pleistocene (MQ 1b) of Nagyharsányhegy 1-4, Hungary (Jánossy 1977).

***Surnia caepki* Jánossy**

Surnia caepki Jánossy, 1972: 53 [Holotype from Stránská skála – Čapek's site 9a: left tarsometatarsus; MMB Ča-807. Figured by Jánossy 1972, pl. 2, fig. 7.]

Surnia caepki Jánossy: Mlíkovský, this paper [Spelling emended.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972).

Genus *Glaucidium* Boie

Glaucidium Boie, 1826 [Modern genus.]

***Glaucidium passerinum* (Linnaeus) – Eurasian Pygmy Owl**

Strix passerina Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1977 sub ?*Otus scops*).

***Glaucidium* sp.**

Distribution: Late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a).

Genus *Athene* Boie

Athene Boie, 1822 [Modern genus.]

Remarks: General reference: Pavia & Mourer-Chauviré 2000.

***Athene angelis* Mourer-Chauviré et al.**

Athene angelis Mourer-Chauviré, Salotti, Pereira, Quinif, Courtois, Dubois & La Milza, 1997: 679 [Holotype from Castiglione 3: incomplete skeleton; MS CAST-3-CG-O-44. Figured by Mourer-Chauviré et al. 1997, unnumbered pl., fig. 1-2 (skull), 3-5 (premaxilla), 6-7 (tibiotarsus), 8-9 (humerus), 11-12 (femur), 13-14 (coracoid), 15-16 (carpometacarpus), and 18-19 (tarsometatarsus).]

Distribution: Middle Pleistocene (MQ 2B) of Castiglione 3, Corsica, France (Mourer-Chauviré et al. 1997); and late Pleistocene (MQ 2C) of Castiglione 1 – layer A, Corsica, France (Mourer-Chauviré et al. 1997).

***Athene cretensis* Weesie**

Athene cretensis Weesie, 1982: 328 [Holotype from Liko: left humerus; RMG Lic-203. Figured by Weesie 1982, fig. 1.]

Distribution: Late Pleistocene (MQ 2C) of Liko – layers A, B, D, a, b, c, and d, Crete, Greece (Weesie 1982, 1987a,b, 1988), and Gumbes B, and Gumbes C, Crete, Greece (Weesie 1982, 1987a,b, 1988).

***Athene noctua* (Scopoli) – Little Owl**

Strix noctua Scopoli, 1769 [Modern species.]

Athene noctua lunellensis Mourer-Chauviré, 1975a: 168 [Holotype from Lunel-Viel, cave I: distal end of right ulna; coll. Bonifay LVI-11-5822. Figured by Mourer-Chauviré 1975a, pl. 22, fig. 11.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), Mas Rambault, France (Mourer-Chauviré 1975a sub *A. noctua lunellensis*), Beremend 16, Hungary (Jánossy 1992 sub *A. veta*; tentatively included here), Beremend 17, Hungary (Jánossy 1992 sub *A. veta*; tentatively included here), and Betfia 2, Romania (Čapek 1917); early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layers 6, 13, and 15e, Czechia (Mlíkovský 1995b); middle Pleistocene (MQ 2A) of Terra Amata, France (Mourer-Chauviré 1975a sub *A. noctua lunellensis*); middle Pleistocene (MQ 2B) of Lunel-Viel – cave 1, layers 1, 2, and 5, France (Mourer-Chauviré 1975a), Orgnac 3 – layers h, i, j, France (Mourer-Chauviré 1975a sub *A. noctua lunellensis*), and Lazaret 8 – layers 8, 12 and 13, France (Mourer-Chauviré 1975a sub *A. noctua lunellensis*).

***Athene* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a).

Remarks: Jánossy (1993: 59) attributed a pedal phalanx from the late Miocene (MN 9) of Rudabánya, Hungary, to *Athene* sp., doubting on the same page its referral to the genus *Athene*, and indicating that the bone is not identifiable even to the genus.

Genus *Strix* Linnaeus

Strix Linnaeus, 1758 [Modern genus.]

***Strix wintershofensis* (Ballmann)**

Otus wintershofensis Ballmann, 1969b: 39 [Holotype from Wintershof-West: distal end of right tibiotarsus; BSP 1937-II-18121. Figured by Ballmann 1969b, pl. 1, fig. 11, 11a, 11b.]

Strix wintershofensis (Ballmann): Mlíkovský 1998f: 12 [New combination.]

Distribution: Early Miocene (MN 3a) of Wintershof-West, Germany (Ballmann 1969b, Mlíkovský 1998f), and Merkur, Czechia (Mlíkovský 1998f).

***Strix edwardsi* (Ennouchi)**

Strix Edwardsi Ennouchi, 1930: 66 [Holotype from Grive-Saint-Alban: distal end of left tibiotarsus; ML LGr-41. Figured by Ennouchi 1930, pl. 5, fig. 9-12.]

Tyto edwardsi (Ennouchi): Brodkorb 1971b: 230 [New combination.]

Strix edwardsi (Ennouchi): Mlíkovský 1998f: 12 [New combination. Note that Ennouchi described the species in the genus *Strix* auct. [= *Tyto* Billberg, 1828], not in *Strix* Linnaeus, 1758.]

Distribution: Middle Miocene (MN 6) of Petersbuch 39, Germany (Mlíkovský 1998f), and middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Ennouchi 1930, Mlíkovský 1998f).

***Strix aluco* Linnaeus – Tawny Owl**

Strix aluco Linnaeus, 1758 [Modern species.]

Strix intermedia Jánossy, 1972: 53 [Holotype from Tarkő – layer 11 (Jánossy 1972: 54) or layer 12 (Jánossy 1977: 21): left coracoid; NMB V-64.746. Not figured.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), Beremend 16, Hungary (Jánossy 1992 sub *S. intermedia*), Beremend 17, Hungary (Jánossy 1992 sub *S. intermedia*); early Pleistocene (MQ 1b) of Boxgrove – quarry 1, England (Harrison & Stewart 1999), Koněprusy C-718, Czechia (Jánossy 1972 and Mlíkovský 1996t sub *S. intermedia*), Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *S. intermedia*, Mlíkovský orig.), and Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *S. intermedia*, Mlíkovský orig.); and middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layer C/D, France (Mourer-Chauviré 1975a sub *S. intermedia*), Hundsheim, Austria (Jánossy 1974b sub *S. intermedia*, Mlíkovský orig.), and Tarkő – layers 10, 11, 12, and 13, Hungary (Jánossy 1972 and 1977 sub *S. intermedia*).

Remarks: Although my first impression was that *Strix intermedia* Jánossy is a valid species (Mlíkovský 1996t), an examination of a large series of bones belonging to modern *Strix aluco* Linnaeus in USNM, and of all fossil material from Czechia and Austria referred by Jánossy (1972, 1974a, 1977) to *Strix intermedia* Jánossy convinced me that *Strix intermedia* Jánossy is not separable from the modern *Strix aluco* Linnaeus. It is questionable, whether all material referred by Jánossy to his *Strix intermedia* belongs to *Strix aluco*.

***Strix* sp.**

Distribution: Late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a sub array of *Strix nebulosa*)

Genus *Aegolius* Kaup

Aegolius Kaup, 1829 [Modern genus.]

***Aegolius funereus* (Linnaeus) – Tengmalm's Owl**

Strix funerea Linnaeus, 1758 [Modern species.]

Athene noctua veta Jánossy, 1974a: 555 [Holotype from Rebielice Królewskie 1: humeral end of right (not left, as stated by Jánossy 1974a) coracoid; ISEZ, uncatalogued. Figured by Jánossy 1974a, pl. 24, fig. 1.]

Athene veta Jánossy: Jánossy 1977: 19 [New rank; see also Jánossy 1981.]

Distribution: Late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a sub *Athene noctua veta*, Jánossy 1977 sub *Athene noctua veta*, Jánossy 1981 sub *Athene veta*, Mlíkovský 1992b), and Osztramos 7, Hungary (Jánossy 1977). Tyrberg (1998: 59) listed *A. cf. funereus* (Linnaeus) from the early Pleistocene (MQ 1b) of "Koněprusy" (= Koněprusy C-718), Czechia, but I was not able to find the species in the papers he cited (i.e. Jánossy 1972, 1976d, 1977, 1979), and I found no bone resembling an owl of this size in my study of the relevant material (Mlíkovský 1996t).

Remarks: *Athene noctua veta* Jánossy was synonymized with the modern *Aegolius funereus* (Linnaeus) by Mlíkovský (1992b: 443).

***Aegolius* sp.**

Distribution: Late Miocene (MN 15) of Csarnóta, Hungary (Jánossy 1980b, exact locality not given); and late Pliocene (MN 18) of Villány 3, Hungary (Jánossy 1980b). However, Jánossy (1986) did not list *Aegolius* from either of the latter two localities.

Genus *Asio* Brisson

Asio Brisson, 1760 [Modern genus.]

***Asio longaevus* (Umans'ka)**

Bubo longaevus Umans'ka, 1979b: 779 [Holotype from Čebotarevka: right tibiotarsus; ZINK 45-3994. Figured by Umans'ka 1979b. fig. 1.]

Asio longaevus (Umans'ka): Mlíkovský 1998f: 13 [New combination.]

Distribution: Late Miocene (MN 11-13) of Čebotarevka, Ukraine (Umans'ka 1979b, Mlíkovský 1998f).

***Asio flammeus* (Pontoppidan) – Short-eared Owl**

Strix flammea Pontoppidan, 1763 [Modern species.]

Distribution: Late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a sub *Asio* aff. *flammeus*); early Pleistocene (MQ 1a) of Osztramos 8, Hungary (Jánossy 1977 sub *A.* aff. *flammeus*), and Betfia 2, Romania (Jánossy 1977 sub *A.* aff. *flammeus*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 1-3, 5a, Czechia (Jánossy 1972 sub *A.* aff. *flammeus* and *Strix intermedia*, Mlíkovský orig.), and Stránská skála – Musil's talus cone, layers 4, 5, 8, 12, and 13, Czechia (Mlíkovský 1995b).

***Asio* sp.**

Distribution: Early Miocene (MN 3a) of Wintershof-West, Germany (Ballmann 1969b); and early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992).

Remarks: Some postcranial bones of the *Asionini* are morphologically very similar to those of the *Surniini* (Mlíkovský orig.). The taxonomic identity of the *Asio* bones from Wintershof-West thus should be treated with caution.

Genus incertae sedis

***Asio pygmaeus* Serebrov's'kyj**

Asio pygmaea Serebrov's'kyj, 1941b: 476 [Holotype from Odesa – catacombs: proximal end of right ulna; present location unknown. Figured by Serebrovskij 1941b, fig.; and Dement'ev 1964, fig. 703a-b.]

Asio pigmaeus Serebrov's'kyj: Brodkorb 1971b: 221 [Ending corrected; species name misspelled.]

Distribution: Early Pliocene (MN 15) of Odesa – catacombs, Ukraine (Serebrov's'kyj 1941b).

Remarks: Indeterminate after the figures (Mlíkovský, orig.).

***Bubo floriana* Kretzoi**

Bubo (?) *floriana* Kretzoi, 1957: 243 [Holotype from Csákvár: phalanx 2 digiti II; GIB, uncatalogued. Figured by Kretzoi 1957, fig. 47-49.]

Distribution: Late Miocene (MN 10) of Csákvár, Hungary (Kretzoi 1957, Jánossy 1977).

Remarks: This phalanx indeed originated from a large owl, but it is indeterminate within the family, being tentatively assigned by Kretzoi (1957) to the genus *Bubo* Duméril on the basis of the size alone (Mlíkovský 1998f).

***Nocturnavis incerta* (Milne-Edwards)**

Bubo incertus Milne-Edwards, 1892: 63 [Holotype from Quercy: right humerus; MNHN QU-16223. Figured by Mourer-Chauviré 1987, text-fig. 2 (left), pl. 2, fig. 15-16.]

Nocturnavis incerta (Milne-Edwards): Mourer-Chauviré 1987: 108 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1982, Mourer-Chauviré 1987); and late Eocene (MP 19) of Escamps, France (Mourer-Chauviré

1987).

Remarks: *Bubo incertus* Milne-Edwards is type by original designation of the genus *Nocturnavis* Mourer-Chauviré, 1987: 106.

***Oligostrix rupelensis* Fischer**

Oligostrix rupeliensis [sic!] Fischer, 1982: 152 [Nomen nudum; no description or indication.]

Oligostrix rupelensis Fischer, 1983b: 483 [Holotype from Espenhain: distal end of left tibiotarsus; NHMB Av-730. Figured by Fischer 1983b.]

Distribution: Middle Oligocene (MP 23-24) of Espenhain, Germany (Fischer 1983b).

Remarks: *Oligostrix rupelensis* Fischer, 1983b is type by original designation of the genus *Oligostrix* Fischer, 1983b: 483. *Oligostrix* Fischer, 1982: 152 is a nomen nudum, because it was not accompanied by an available species name after 1930.

***Selenornis henrici* (Milne-Edwards)**

Otus Henrici Milne-Edwards, 1892: 63 [Holotype from Quercy: distal end of left tibiotarsus; MNHN QU-16222. Figured by Mourer-Chauviré 1987, pl. 3, fig. 5-6.]

Asio Henrici (Milne-Edwards): Gaillard 1908: 36 [New combination.]

Selenornis henrici (Milne-Edwards): Mourer-Chauviré 1987: 113 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Mourer-Chauviré 1987).

Remarks: Taxonomic position of this species is uncertain (Mlíkovský 1998c). *Otus henrici* Milne-Edwards, 1892 is type by original designation of the genus *Selenornis* Mourer-Chauviré, 1987: 112.

***Strix ignota* Milne-Edwards**

Strix ignota Milne-Edwards, 1871: 580 [Holotype from Sansan: distal end of left tarsometatarsus, lacking trochlea of digit IV; MNHN Sa-1254. Figured by Milne-Edwards 1869-1871, pl. 192, fig. 1, 1a, 2, 2a; and Cheneval 2000.]

Tyto ignota (Milne-Edwards): Brodkorb 1971b: 230 [New combination.]

Asio (?) *ignotus* (Milne-Edwards): Cheneval 2000: 365 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871 sub *Strix* sp. (p. 499) and *Strix ignota* (p. 580), Mlíkovský 1998c, Mourer-Chauviré 1999c, Cheneval 2000).

Remarks: Lambrecht (1933: 613) and Brodkorb (1971: 230) erroneously attributed this name to Paris (1912). Paris (1912: 287) and Lambrecht (1921: 97) attributed this name to Milne-Edwards (1871: 499), referring to a page, where the name does not appear. This led Mlíkovský (1998c: 255) to conclude, that the name never did exist, and that Paris (1912) listed it in error. However, Mourer-Chauviré (1999c: 363) and Cheneval (2000: 368) observed, that Milne-Edwards (1871) created the name on p. 580 with a notice, that the species originated from Sansan, and under the reference to p. 499 of the same work. Accordingly, *Strix ignota* is available with Milne-Edwards (1871: 580) as its author.

Cheneval (2000) compared the fossil with only a few modern European owl genera, not considering e.g. the Surniini. He presented no convincing evidence that the species belongs to *Asio* Brisson, and correctly expressed his doubts about this allocation, adding a question-mark to the genus. I leave the species in the Strigidae inc. sedis, but I agree with Cheneval (2000), that it belongs to the Striginae (in the present sense), not to the Tytoninae.

Family Columbidae Illiger

Columbidae Illiger, 1811 [Modern family.]

Genus *Columba* Linnaeus

Columba Linnaeus, 1758 [Modern genus.]

Columba pisana (Portis)

Falco sp. (*Pisanus*) Portis, 1887: 195 [Nomen nudum; no description or indication.]

Falco sp. (*pisanus*) Portis, 1889: 14 [Holotype from Orciano Pisano: distal end of ulna; IGF 14876. Figured by Portis, 1889, pl. 1, fig. 27a-b.]

Columba omnisanctorum Ballmann, 1976a: 30 [Holotype from Biancone 1: distal end of right tarsometatarsus; RGM 178782. Figured by Ballmann 1976, text.-fig. 12, pl. 7, fig. 3-5.]

Columba pisana (Portis): Mlíkovský, this paper [New combination.]

Distribution: Early Pliocene (MN 14-15) of Biancone 1, Italy (Ballmann 1976a), Biancone 2, Italy (Ballmann 1976a), and San Giovannino, Italy (Ballmann 1976a); and middle Pliocene (MN 15-16) of Orciano Pisano, Italy (Portis 1889).

Remarks: C. Mourer-Chauviré (in Olson 1985a: 111) found that the holotypical ulna of *Falco pisanus* is from a columbid bird. In size it agrees with *Columba omnisanctorum* Ballmann, which was described from similarly old deposits near the paleosea shore. In view of the coincidence in age, type of habitat and size I synonymize here *Columba omnisanctorum* Ballmann with *Columba pisana* (Portis). The generic position follows Ballmann (1976a).

Columba melitensis Lydekker

Columba melitensis Lydekker, 1891a: 124 [Holotype from an unknown site in Malta: left coracoid; BMNH A-212. Figured by Lydekker 1891a, fig. 29.]

Distribution: Late Pleistocene (MQ 2C) of an unknown site in Malta (Lydekker 1891a).

Columba livia Gmelin – Rock Dove

Columba livia Gmelin, 1789 [Modern species.]

Columba livia occitanica Mourer-Chauviré, 1975a: 146 [Holotype from Saint-Estève-Janson – layer B/C: left humerus; coll. Bonifay CD66-B = C-11623. Figured by Mourer-Chauviré 1975a, pl. 22, fig. 5.]

Columba livia minuta Mourer-Chauviré, 1975a: 150 [Holotype from Fage – layer 5: left humerus; FSL 41665. Figured by Mourer-Chauviré 1975a, pl. 14, fig. 5. This is a junior primary homonym of *Columba minuta* Linnaeus, 1758, and *Columba minuta* Temminck, 1838-1843.]

Columba livia lazaretensis Mourer-Chauviré, 1975a: 152 [Holotype from Lazaret 8, layer V₂: left humerus; LPLN 23ab. Figured by Mourer-Chauviré 1975a, pl. 14, fig. 11.]

Distribution: Early Pleistocene (MQ 1) of Monte Argentario, Italy (Cassoli 1980: 184); early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), Mas Rambault, France (Mourer-Chauviré 1975a), Beremend 17, Hungary (Jánossy 1992 sub *C. cf. livia*), and Primorsk, Ukraine (Vojinstvens'kyj 1967); early Pleistocene (MQ 1b) of Sima del Elefante – unknown layer, Spain (Rosas et al. 2001); middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layers B, C, D, F, G and H, France (Mourer-Chauviré 1975a sub *C. livia occitanica*), and Cimay, France (Mourer-Chauviré 1975a); and middle Pleistocene (MQ 2B) of Lunel-Viel, France (Mourer-Chauviré 1975a sub *C. livia occitanica*), Fage – layers 2/3, 31 through 40, and CO, France (Mourer-Chauviré 1975a sub *C. livia minuta*), and Lazaret 8, layers I/II through XX, France (Mourer-Chauviré 1975a sub *C. livia lazaretensis*).

The extralimital record includes middle Pleistocene (MQ 2B) of Treugolnaja Cave, Karachaevo-Cherkessia in Russia (Potapova & Baryšnikov 1993 and Baryšnikov & Potapova 1995 sub *C. l. occitanica*).

***Columba oenas* Linnaeus – Stock Dove**

Columba oenas Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Petralona – layers 11, 13, 15, 24, 26, and “mausoleum”, Greece (Kretzoi 1977, Kretzoi & Poulianos 1981, Poulianos 1996).

Remarks: These bones should be compared with those of the modern *Columba livia* Linnaeus.

***Columba palumbus* Linnaeus – Common Wood Pigeon**

Columba palumbus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972), Stránská skála – Musil's talus cone, layer 15e, Czechia (Mlíkovský 1995b), Nagyharsányhegy 1-4, Hungary (Jánossy 1980a sub *C. aff. palumbus*), and Petralona – layers 11 and 15, Greece (Kretzoi 1977, Kretzoi & Poulianos 1981, Poulianos 1996).

***Columba* sp.**

Distribution: Late Pliocene (MN 17) of Šandalja 1 – layer 6, Croatia (Malez-Bačić 1979); and late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b, 2000a).

Genus *Streptopelia* Bonaparte

Streptopelia Bonaparte, 1855 [Modern genus.]

***Streptopelia* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a).

Family Falconidae Leach

Falconidae Leach, 1819 [Modern family.]

Genus *Falco* Linnaeus

Falco Linnaeus, 1758 [Modern genus.]

***Falco medius* Umans'ka**

Falco medius Umans'ka, 1981: 18 [Holotype from Čebotarevka: left carpometacarpus; IZAN 45-4033. Figured by Umans'ka 1981, fig. 2a,b,v.]

Distribution: Late Miocene (MN 11-13) of Čebotarevka, Ukraine (Umans'ka 1981).

Remarks: This fossil is in need of revision.

***Falco bakalovi* Boev**

Falco bakalovi Boev, 1999c: 132 [Holotype from Väršec: postacetabular part of the left part of pelvis; NMNHS 1642. Figured by Boev 1999c, fig. 1a-b, 4 (partim).]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1999c).

Remarks: A dubious species, whose taxonomic position should be reevaluated.

***Falco tinnunculus* Linnaeus – Common Kestrel**

Falco Tinnunculus Linnaeus, 1758 [Modern species]

Falco tinnunculus atavus Jánossy, 1972: 45 [Lectotype (here selected): proximal end of left humerus from Stránská skála – Čapek's site 9a; MMB Ča-622. Not figured. Paralectotypes from Stránská skála – Čapek's site 4c: pedal phalanx (MMB Ča-551); Stránská skála –

Čapek's site 5a: proximal end of left scapula (MMB Ča-194), distal end of left ulna (MMB Ča-387), distal end of left (not right as stated by Jánossy 1972: 45) carpometacarpus (MMB Ča-195), and distal end of right (not left as stated by Jánossy 1972: 45) tarsometatarsus; Stránská skála – Čapek's site 5b: distal end of right tibiotarsus (MMB Ča-483); Stránská skála – Čapek's site 9a: proximal end of right scapula (MMB Ča-630), proximal end of left humerus (MMB Ča-622), distal end of right (not left as stated by Jánossy 1972: 46) radius (MMB Ča-631), left phalanx I digiti majoris (MMB Ča-632), right phalanx I digiti majoris (MMB Ča-630), distal end of left femur (MMB Ča-623), proximal end of left (not right as stated by Jánossy 1972: 46) tarsometatarsus (MMB Ča-627), proximal ends of 2 right tarsometatarsi (MMB Ča-625, Ča-633), distal end of right tarsometatarsus (MMB Ča-628), and 4 pedal phalanges (MMB Ča-634-637; Ča-635 absent from the collection); Stránská skála – Čapek's site 9b: proximal end of right scapula (MMB Ča-794), right carpometacarpus (MMB Ča-793), and distal end of right (not left as stated by Jánossy 1972: 46) tarsometatarsus (MMB Ča-799); Stránská skála – Absolon's 2nd cave: left coracoid (MMB Abs-1063), and right ulna (MMB Abs-64); Stránská skála – unidentified sites: sternal end of left coracoid (MMB Va-6), proximal end of left scapula (MMB SS-2063; absent from the collection), right humerus (MMB Ča-563), proximal end of right humerus (MMB SS-2061), 2 left ulnae (MMB Ča-3561, Ča-3562), right ulna (MMB Va-2), left carpometacarpus (MMB Ča-3564), right carpometacarpus (MMB Ča-3565), proximal end of left femur (MMB Ča-3566), distal ends of 2 left tibiotarsi (MMB Ča-3563, Va-3), and right tarsometatarsus (MMB Va-4); Betfia 2 (after Jánossy 1977: 29; coll. Kormos): two coracoids, proximal end of a humerus, distal ends of three humeri, proximal ends of two carpometacarpi, and a tarsometatarsus; Hundsheim (UWPI 1889/53): distal end of left humerus, distal end of right humerus, distal end of left carpometacarpus, distal end of left tibiotarsus, distal end of right tarsometatarsus, and a pedal phalanx; and "Koněprusy" = Koněprusy C-718 (after Jánossy 1972: 46): a humerus (present location unknown, not found by Mlíkovský 1996t). Not figured. Both syntypes from the Absolon's 2nd cave belong to other species: coracoid Abs-1063 to *Falco vespertinus*, and ulna Abs-64 to *Falco subbuteo* – Mlíkovský, orig.]

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b and 1990b sub *F. tinnunculus* cf. *atavus*, Jánossy 1992 sub *F. tinnunculus atavus*); early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub *F. tinnunculus atavus*), Valerots, France (Mourer-Chauviré 1975a sub *F. tinnunculus atavus*), Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *F. tinnunculus atavus*, Mlíkovský 1998d), Beremend 16, Hungary (Jánossy 1992 sub *F. tinnunculus atavus*), Beremend 17, Hungary (Jánossy 1992 sub *F. tinnunculus atavus*), and Betfia 2, Romania (Čapek 1917, Jánossy 1972 and 1977 sub *F. tinnunculus atavus*); early Pleistocene (MQ 1b) of Kozi Grzbiet, Poland (Bocheński 1984 sub *F. tinnunculus atavus*), Koněprusy C-718, Czechia (Jánossy 1972), Přezletice, Czechia (Jánossy 1983 sub *F. tinnunculus atavus*), Stránská skála – Čapek's sites 4c, 5a, 5b, 9a, 9b and unidentified sites, Czechia (Jánossy 1972 sub *F. tinnunculus atavus*), Stránská skála – Musil's talus cone, layers 4, 5, 6, 13, 15a, 16e, and 16, Czechia (Mlíkovský 1995), Včeláre 1, Slovakia (Jánossy 1977 sub *F. tinnunculus atavus*), Nagyarsányhegy 1-4, Hungary (Jánossy 1977 sub *F. tinnunculus atavus*), Somssichhegy 2, Hungary (Jánossy 1986: 56 sub *F. tinnunculus atavus*), and Betfia 5, Romania (Kretzoi 1962), and Petralona – layer "mausoleum", Greece (Kretzoi 1977, Kretzoi & Poulianos 1981, Poulianos 1996); middle Pleistocene (MQ 2A) of Gombasek, Slovakia (Jánossy 1980b sub *F. tinnunculus atavus*), and Tarkö – layers 2 and 4, Hungary (Jánossy 1986: 74-75 sub *F. tinnunculus atavus*); and middle Pleistocene (MQ 2B) of Fage – layers 2-3, 4, 31, 32, 33, 34, 35, 36, 37, 38, 39, and 41, France (Mourer-Chauviré 1975a sub *F. tinnunculus atavus*), Orgnac 3 – layer h, i, j, France (Mourer-Chauviré 1975a sub *F. tinnunculus atavus*), Biehle, France (Mourer-Chauviré 1975a sub *F. tinnunculus atavus*), and Fournier Quarry, France (Mourer-Chauviré 1975a sub *F. tinnunculus atavus*).

***Falco vespertinus* Linnaeus – Red-footed Falcon**

Falco vespertinus Linnaeus, 1766 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *F. cf. vespertinus*), and Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *F. cf. vespertinus*), and Somssichhegy 2, Hungary (Jánossy 1986 sub *F. cf. vespertinus*).

***Falco subbuteo* Linnaeus – Eurasian Hobby**

Falco subbuteo Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 5, Romania (Kretzoi 1962 sub *F. cf. subbuteo*); and early Pleistocene (MQ 1b) of Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub ?*F. aff. subbuteo*), and Stránská skála – Musil's talus cone, layers 5, 6, and 13, Czechia (Mlíkovský 1995b).

***Falco cherrug* Gray – Saker Falcon**

Falco cherrug Gray, 1844 [Modern species.]

Falco antiquus Mourer-Chauviré, 1975a: 75 [Holotype: right tarsometatarsus from Fage – layer 5; FSL 41584. Figured by Mourer-Chauviré 1975a, pl. 9, fig. 3.]

Distribution: Middle Pleistocene (MQ 2B) of Fage – layer 5, France (Mourer-Chauviré 1975a sub *Falco antiquus*), and Hórvölgy, Hungary (Jánossy 1977: 28 sub *F. aff. atavus*, an apparent misprint for *antiquus*).

Remarks: The differences between *Falco cherrug* Gray and *Falco antiquus* Mourer-Chauviré do not go beyond the intraspecific variability (Mlíkovský, orig.). Hence, I synonymize here the latter with the former species.

***Falco rusticolus* Linnaeus – Gyr Falcon**

Falco rusticolus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *F. aff. rusticolus*).

***Falco peregrinus* Tunstall – Peregrine Falcon**

Falco peregrinus Tunstall, 1771 [Modern species.]

Distribution: Eearly Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *F. aff. cherrug*, Mlíkovský orig.), and Stránská skála – Musil's talus cone, layers 4, 8, 11, 13, and 14a, Czechia (Mlíkovský 1995b).

***Falco* sp.**

Distribution: Early Pliocene (MN 15) of Muselievo, Bulgaria (Boev 1998e, 2000a); late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a); late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 2000a); and early Pleistocene (MQ 1b) of Sima del Elefante – unknown layer, Spain (Rosas et al. 2001), and Stránská skála – Čapek's site 1-3, Czechia (Jánossy 1972 sub *F. aff. cherrug*, Mlíkovský orig.).

Order Coraciiformes Forbes

Coraciiformes Forbes, 1884 [Modern order.]

Family Coraciidae Vigors

Coraciidae Vigors, 1825 [Modern family.]

Eocoraciidae Mayr & Mourer-Chauviré, 2000: 534 [Type genus: *Eocoracias* Mayr & Mourer-Chauviré, 2000.]

Geranopteridae Mayr & Mourer-Chauviré 2000: 538 [Type genus: *Geranopterus* Milne-Edwards, 1892.]

Remarks: Mayr & Mourer-Chauviré (2000) created the families Eocoraciidae and Geranopteridae on the basis of negligible characters, showing at the same time that the taxa are closely allied to the modern Coraciidae. I see no reason why the observed differences should not be interpreted as generic characters of *Eocoracias* Mayr & Mourer-Chauviré, and *Geranopterus* Milne-Edwards, respectively, and I synonymize here thus Eocoraciidae Mayr & Mourer-Chauviré, and Geranopteridae Mayr & Mourer-Chauviré with the modern Coraciidae Vigors.

Genus *Eocoracias* Mayr & Mourer-Chauviré

Eocoracias Mayr & Mourer-Chauviré, 2000: 534 [Type by original designation: *Eocoracias brachyptera* Mayr & Mourer-Chauviré, 2000.]

Eocoracias brachyptera Mayr & Mourer-Chauviré

Eocoracias brachyptera Mayr & Mourer-Chauviré, 2000: 534 [Holotype from Messel: complete articulated skeleton on two slabs; SMNK PAL-2663a,b. Figured by Mayr & Mourer-Chauviré 2000, fig. 1, 6; and Mayr 2000g, fig. 7-8.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr & Mourer-Chauviré 2000).

Genus *Geranopterus* Milne-Edwards

Geranopterus Milne-Edwards, 1892: 66 [Type by monotypy: *Geranopterus alatus* Milne-Edwards, 1892.]

Geranopterus alatus Milne-Edwards

Geranopterus alatus Milne-Edwards, 1892: 66 [Holotype from Quercy: left humerus; MNHN QU-15890, cast in coll. Mlíkovský, uncatalogued. Figured by Mayr & Mourer-Chauviré 2000, fig. 8a-b.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Mourer-Chauviré 1999a, see also Mayr 1998c); late Eocene (MP 18) of Gousnat, France (Mayr & Mourer-Chauviré 2000); and late Eocene (MP 19) of Escamps, France (Mourer-Chauviré 1999a, Mayr & Mourer-Chauviré 2000), Rosières, France (Mayr & Mourer-Chauviré 2000), and Pécarel, France (Mayr & Mourer-Chauviré 2000).

Remarks: Milne-Edwards (1892: 66) compared this species with the Coraciidae and Momotidae. Lambrecht (1933: 627) and Brodkorb (1971: 250) included it in the Coraciidae without comments. Mayr & Mourer-Chauviré (2000) confirmed the coraciid affinities of *Geranopterus* Milne-Edwards.

Geranopterus milneedwardsi Mayr & Mourer-Chauviré

Geranopterus milneedwardsi Mayr & Mourer-Chauviré, 2000: 543 [Holotype from Quercy: proximal end of left humerus; MNHN QU-17037. Figured by Mayr & Mourer-Chauviré

2000, fig. 9a-b.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mayr & Mourer-Chauviré 2000); and late Eocene (MP 17) of Perrière, France (Mayr & Mourer-Chauviré 2000).

Genus *Cryptornis* Milne-Edwards

Cryptornis Milne-Edwards, 1871: 371 [Type by monotypy: *Centropus antiquus* Gervais, 1852.]

Cryptornis antiquus (Gervais)

Centropus? antiquus Gervais, 1852: 229 [Holotype from Montmartre: partial skeleton in slab; MNHN 7991. Figured by Gervais 1852, pl. 49, fig. 1, Laurillard 1849, pl. 2, and Milne-Edwards 1869-1871, pl. 175.]

Cryptornis antiquus (Gervais): Milne-Edwards, 1871: 371 [New combination.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Laurillard 1849 sub *Alcedo*, Gervais 1852, Milne-Edwards 1869-1871).

Remarks: This species was described as a cuckoo (Gervais 1848-1852), and later believed to be related to the Bucerotidae (Milne-Edwards 1869-1871: 374, Murie 1873, Brodkorb 1971: 249), Picidae (Paris 1912: 285, Furon & Soyer 1947, Brunet 1970), and Coraciidae (Harrison 1979b: 107, Mayr 1998c: 29). I follow here Harrison (1979b) in placing provisionally the species in the family Coraciidae. Mayr & Mourer-Chauviré (2000) observed that the holotype agrees in size with the bones of *Geranopterus alatus* Milne-Edwards, concluding that "an evaluation of the possible synonymy of the genera *Cryptornis* Milne-Edwards [...] and *Geranopterus* Milne-Edwards [...] must await the discovery of better preserved specimens of *C. antiquus*" (Mayr & Mourer-Chauviré 2000: 538).

Genus *Coracias* Linnaeus

Coracias Linnaeus, 1758 [Modern genus.]

Coracias garrulus Linnaeus

Coracias garrulus Linnaeus, 1758 [Modern genus.]

Distribution: Early Pleistocene (MQ 1a) of Tarchankut, Ukraine (Vojinstvens'kyj 1967 sub *C. aff. garrulus*).

Family Trogonidae Lesson

Trogonidae Lesson, 1828 [Modern family.]

Genus *Primotrogon* Mayr

Primotrogon Mayr, 1999a: 427 [Type by original designation: *Primotrogon wintersteini* Mayr, 1999a.]

Primotrogon wintersteini Mayr

Primotrogon wintersteini Mayr, 1999a: 428 [Holotype from Céreste: skeleton in a slab; BSP 1997-I-38. Figured by Mayr 1999a, fig. 1-2, 6.]

Distribution: Middle Oligocene (MP 23) of Céreste, France (Mayr 1999a).

Genus *Apaloderma* Swainson

Apaloderma Swainson, 1833 [Modern genus.]

Paratrogon Lambrecht, 1933: 626 [Type by monotypy: *Trogon gallicus* Milne-Edwards,

1871.]

***Apaloderma gallica* (Milne-Edwards)**

Trogon gallicus Milne-Edwards, 1871: 395 [Lectotype from Saint-Gérard-le-Puy (here selected): right humerus; MNHN Av-9675. Figured by Milne-Edwards 1869-1871, pl. 177, fig. 18-22. Milne-Edwards (1871: 395) based this species expressly on two humeri, but I found only one at MNHN in June 1999. The missing humerus should be regarded as the paralectotype of this species.]

Paratrogon gallicus (Milne-Edwards): Lambrecht 1933: 626 [New combination.]

Apaloderma gallica (Milne-Edwards): Mlíkovský, this paper [New combination.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Mourer-Chauviré 1980a, Mlíkovský orig.); and early Miocene (MN 3) of Merkur, Czechia (Mlíkovský orig.).

Remarks: The lectotypical humerus of *Trogon gallicus* Milne-Edwards agrees with the same element of *Apaloderma* Swainson, and differs from that of all other modern trogonid genera (sensu Espinosa de los Monteros 1998), in having entepicondylar prominence large and distinct. In absence to any evidence to the contrary, I transfer here *Trogon gallicus* Milne-Edwards to the genus *Apaloderma* Swainson. Accordingly, I synonymize here the genus *Paratrogon* Lambrecht with *Apaloderma* Swainson.

Family Alcedinidae Rafinesque

Alcedinidae Rafinesque, 1815 [Modern family.]

Genus *Alcedo* Linnaeus

Alcedo Linnaeus, 1758 [Modern genus.]

***Alcedo atthis* (Linnaeus) – Common Kingfisher**

Gracula Atthis Linnaeus, 1758 [Modern species.]

Distribution: Late Oligocene (MP 30) of Aix-en-Provence, France (Bayan 1873). This record is based on a feather imprint, and is highly improbable.

Remarks: Tyrberg (1998) cited three late Pleistocene localities from Europe, where *Alcedo* Linnaeus was found, but none of them contains unmixed material older than the Holocene. There is no evidence, that *Alcedo* Linnaeus existed in Europe before the Holocene.

Family Todidae Vigors

Todidae Vigors, 1825 [Modern family.]

Momotidae Gray, 1840 [Modern family.]

Remarks: A feather fragment found in the late Eocene (MP 14-20) Baltic amber, which originated in Finland (Katinas 1983, Mlíkovský 1996g), was tentatively attributed to the "Momotidae" by Bachofen-Echt (1936b: 346-347, pl. 5, fig. 1-3).

Genus *Palaeotodus* Olson

Palaeotodus Olson, 1976: 112 [Type by original designation and monotypy: *Palaeotodus emryi* Olson, 1976.]

Distribution: Late Eocene (MP 19) and middle Oligocene (MP 23) of France. The extralimital record includes only *Palaeotodus emryi* Olson, 1976a from the middle Oligocene (Orellanian) of Wyoming (Olson 1976a). The European record requires revision.

***Palaeotodus escampsensis* Mourer-Chauviré**

Palaeotodus escampsensis Mourer-Chauviré, 1985b: 408 [Holotype from Escamps: right humerus lacking proximal end; USTL ECA-3215. Figured by Mourer-Chauviré 1985b, pl. 1, fig. 1-3.]

Distribution: Late Eocene (MP 19) of Escamps, France (Mourer-Chauviré 1985b).

***Palaeotodus itardiensis* Mourer-Chauviré**

Palaeotodus itardiensis Mourer-Chauviré, 1985b: 409 [Holotype from Itardies: proximal end of left ulna; USTL ITD-576. Figured by Mourer-Chauviré 1985b, pl. 1, fig. 4-5.]

Distribution: Oligocene (MP 21-28) of Fonbonne 1, France (Mourer-Chauviré 1985b); and middle Oligocene (MP 23) of Itardiès, France (Mourer-Chauviré 1985b).

Genus *Protornis* Meyer

Protornis Meyer, 1844: 338 [Type by monotypy: *Protornis glarniensis* Meyer, 1844.]

***Protornis glarniensis* Meyer**

Protornis glarniensis Meyer, 1844: 338 [Holotype from Matt: Incomplete skeleton in slab and counter-slab; NMZ, uncatalogued. Figured by Meyer 1856, pl. 15, fig. 12, and Peyer 1957, text-fig. 1, 7b-c, 8 (right), 9, 11, 13, 15, 17, 19, 21, pl. 1-5.]

Osteornis scolopacinus Gervais, 1844a: 38 [Holotype: Same as for *Protornis glarniensis* Meyer, 1844. This name was listed as a nomen nudum by Brodkorb 1978: 222, and Mlíkovský 1992b: 444. However, Gervais 1844a: 38 based it on a bird formerly described by Meyer from the “schistes du Canton du Glaris”, with reference to Escher (see Gervais 1844a: 12). This reference can be traced back to the paper by Meyer (1839: 683-684), where the bird is briefly described, though not named. With this reference, *Osteornis scolopacinus* Gervais, 1844a is an available name.]

Protornis glaronensis Meyer, 1856: 92 [Unjustified emendation of *Protornis glarniensis* and consequently its junior objective synonym.]

Distribution: Early/middle Oligocene (MP 21-24) of Matt, Switzerland (Meyer 1839c, 1844, 1856, Studer 1840, Gervais 1844a, Peyer 1949, 1957; see also Baumann 1958, Stüssi 1958, and Olson 1976a).

Remarks: Cracraft (1980: 13) and Mayr (1998c: 38) casted some doubt on the allocation of *Protornis glarniensis* Meyer to the Momotidae (= Todidae). Mlíkovský (1992b: 444) erroneously listed *Protornis glarisiensis* Anonymous, 1844: v as a nomen nudum, this being only an incorrect subsequent spelling.

Family Meropidae Vigors

Meropidae Vigors, 1825 [Modern family.]

Genus *Merops* Linnaeus

Merops Linnaeus, 1758 [Modern genus.]

***Merops radoboyensis* (Meyer)**

Batrachites Radoboyensis H. Freyer, between 1844-1845 [Label name; not available for nomenclatural purposes – see Mlíkovský 1997a.]

Pelophilus Radoboyensis (Freyer) J.J. von Tschudi, between 1845-1849 [New combination; label name; not available for nomenclatural purposes – see Mlíkovský 1997a.]

Fringilla (?) Radoboyensis Meyer, 1865: 125 [Holotype from Radoboj: partial skeleton of hind limbs in a plate; NHMW 1845/XXII/1. Figured by Meyer 1865, pl. 30, fig. 1; Mlíkovský

1997a, fig. 2-3.]

Merops radoboyensis (Meyer): Mlíkovský 1997a: 145 [New combination.]

Distribution: Middle Miocene (MN 7) of Radoboj, Croatia (Meyer 1865, Mlíkovský 1997a).

Order Passeriformes Linnaeus

Passeres Linnaeus, 1758 [Modern order.]

Family Eurylaimidae Lesson

Eurylaiminae Lesson, 1831 [Modern family.]

Distribution: Early Miocene (MN 3) of Wintershof-West, Germany (Ballmann 1969b sub unidentified representative).

Family Hirundinidae Rafinesque

Hirundinidae Rafinesque, 1815 [Modern family.]

Genus *Riparia* Forster

Riparia Forster, 1817 [Modern genus.]

Riparia riparia (Linnaeus) – Sand Martin

Hirundo riparia Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 8, Czechia (Mlíkovský 1995b).

Genus *Hirundo* Linnaeus

Hirundo Linnaeus, 1758 [Modern genus.]

Hirundo rustica Linnaeus – Barn Swallow

Hirundo rustica Linnaeus, 1758 [Modern species.]

Distribution: Early Pliocene (MN 15) of Ivanovce 1, Slovakia (Mlíkovský orig.); early Pleistocene (MQ 1a) of Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *H. cf. rustica*, Mlíkovský 1998d), Deutsch-Altenburg 2C, Austria (Mlíkovský 1998d), Osztramos 8, Hungary (Jánossy 1986 sub *H. cf. rustica*), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *Hirundo* sp.); early Pleistocene (MQ 1b) of Koněprusy C-718, Czechia (Mlíkovský 1996t), Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972), Stránská skála – Musil's talus cone, layers 10a, 13 and 15a, Czechia (Mlíkovský 1995b), Včeláre 4E, Slovakia (Mlíkovský orig.), and Somssichhegy 2, Hungary (Jánossy 1986 sub *H. cf. rustica*).

Hirundo sp.

Distribution: Early Pliocene (MN 15) of Csarnóta 1, Hungary (Kretzoi 1956); late Pliocene (MN 18) of Villány 3, Hungary (Jánossy 1979); early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992); and early Pleistocene (MQ 1b) of Nagyharsányhegy 1-4, Hungary (Jánossy 1979).

Genus *Delichon* Horsfield & Moore

Delichon Horsfield & Moore, 1854 [Modern genus.]

Delichon urbica (Linnaeus) – Common House Martin

Hirundo urbica Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), and Betfia 2, Romania (Čapek 1917); and early Pleistocene (MQ 1b) of Stránská skála – Musil's

talus cone, layers 6, 8, 9, 10a, 13, 14a, and 15e, Czechia (Mlíkovský 1995b).

Family Alaudidae Vigors

Alaudidae Vigors, 1825 [Modern family.]

Remarks: Unidentified Alaudidae were reported also from the middle Miocene (MN 5) of Vieux-Collonges, France (Ballmann 1972).

Genus *Melanocorypha* Boie

Melanocorypha Boie, 1828 [Modern genus.]

Melanocorypha sp.

Distribution: Late Miocene (MN 11-13) of Chrabärsko, Bulgaria (Boev 1996 sub cf. *Galerida*, Boev 2000a,d); late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a); and early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992).

Genus *Calandrella* Kaup

Calandrella Kaup, 1829 [Modern species.]

Calandrella cinerea (Gmelin) – Red-capped Lark

Alauda cinerea Gmelin, 1789 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Tarchankut, Ukraine (Vojinstvens'kyj 1967 sub *C. cf. cinerea*).

Genus *Galerida* Boie

Galerida Boie, 1828 [Modern genus.]

Galerida cristata (Linnaeus) – Crested Lark

Alauda cristata Linnaeus, 1758 [Modern genus.]

Distribution: Early Pleistocene (MQ 1b) of Somssichhegy 2, Hungary (Jánossy 1986 sub *G. cf. cristata*), and Betfia 5, Romania (Kessler 1975).

Galerida sp.

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b, 2000a); and early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992).

Genus *Lullula* Kaup

Lullula Kaup, 1829 [Modern genus.]

Lullula arborea (Linnaeus) – Wood Lark

Alauda arborea Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995).

Lullula sp.

Distribution: Late Miocene (MN 11-13) of Chrabärsko, Bulgaria (Boev 1996 sub cf. *Lullula*); late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b, 2000a), and Slivnica, Bulgaria (Boev 2000a sub aff. *Lullula* sp.).

Genus *Alauda* Linnaeus*Alauda* Linnaeus, 1758 [Modern genus.]***Alauda arvensis* Linnaeus – Eurasian Lark***Alauda arvensis* Linnaeus, 1758 [Modern species.]**Distribution:** Early Pleistocene (MQ 1a) of Valerots, France (Mourer-Chauviré 1975a sub cf. *A. arvensis*), Beremend 16, Hungary (Jánossy 1992 sub *A. cf. arvensis*), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *A. cf. arvensis*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 9a and 9b, Czechia (Jánossy 1972 sub *A. cf. arvensis*).***Alauda* sp.****Distribution:** Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b, 2000a).**Genus *Eremophila* Boie***Eremophila* Boie, 1828 [Modern genus.]***Eremophila alpestris* Linnaeus – Horned Lark***Eremophila alpestris* Linnaeus, 1758 [Modern species.]**Distribution:** Early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub *E. cf. alpestris*).***Eremophila* sp.****Distribution:** Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a).**Family Bombycillidae Swainson**

Bombycillidae Swainson, 1831 [Modern family.]

Genus *Bombycilla* Vieillot*Bombycilla* Vieillot, 1807 [Modern genus.]***Bombycilla garrulus* (Linnaeus) – Bohemian Waxwing***Lanius Garrulus* Linnaeus, 1758 [Modern species.]**Distribution:** Early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992 sub *B. cf. garrulus*).**Family Laniidae Rafinesque**

Laniidae Rafinesque, 1815 [Modern family.]

Genus *Lanius* Linnaeus*Lanius* Linnaeus, 1758 [Modern genus.]***Lanius collurio* Linnaeus – Red-backed Shrike***Lanius collurio* Linnaeus, 1758 [Modern species.]**Distribution:** Early Pleistocene (MQ 1a) of Betfia 7, Romania (Kessler 1975 sub *L. cf. collurio*).***Lanius minor* Gmelin – Lesser Grey Shrike**

Lanius minor Gmelin, 1788 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Jánossy 1979 sub *L. cf. minor*); and early Pleistocene (MQ 1b) of Petralona – layer 24, Greece (Kretzoi 1977, Kretzoi & Poulianos 1981).

***Lanius* sp.**

Distribution: Late Miocene (MN 13) of Polgárdi 2, Hungary (V. Čapek in Lambrecht 1933: 780 sub cf. *L. minor*, Jánossy 1991).

Family Sturnidae Rafinesque

Sturnidae Rafinesque, 1815 [Modern family.]

Genus *Sturnus* Linnaeus

Sturnus Linnaeus, 1758 [Modern genus.]

***Sturnus vulgaris* Linnaeus – Common Starling**

Sturnus vulgaris Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Beremend 16, Hungary (Jánossy 1992 sub *S. cf. vulgaris*), Beremend 17, Hungary (Jánossy 1992 sub *S. cf. vulgaris*), Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *S. cf. vulgaris*); and early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d sub *S. ?vulgaris* or *S. vulgaris/unicolor* superspecies), Boxgrove – quarry 2, England (Harrison & Stewart 1999), and Přezletice, Czechia (Jánossy 1983 sub *S. cf. vulgaris*).

***Sturnus* sp.**

Distribution: Late Pliocene (MN 17) of Váršec, Bulgaria (Boev 2000a).

Family Corvidae Vigors

Corvidae Vigors, 1825 [Modern family.]

Genus *Miocorvus* Lambrecht

"*Miocorax*" = *Miocorvus* Lambrecht, 1933: 636 [Type by monotypy: *Corvus larteti* Milne-Edwards, 1871. *Miocorax* Lambrecht 1933: 636 is a misprint for *Miocorvus* – see Lambrecht 1933: 1024.]

***Miocorvus larteti* (Milne-Edwards)**

Corvus larteti Milne-Edwards, 1871: 381 [Lectotype from Sansan (selected by Cheneval 2000: 374): distal end of right right tarsometatarsus; NMNH Sa-1248. Not figured. Cheneval (2000: 374) believed that this bone is figured by Milne-Edwards 1869-1871, pl. 152, figs. 4, 4a, 5, 5a, 6, 6a, 7, that is was "presque complet" at the time of drawing, and that it subsequently lost its proximal end. However, at least two different tarsometatarsi are shown in those figured (which are apparently side-reversed): an almost complete right tarsometatarsus with damaged distal end (figs. 1-3, and perhaps also 6-8), and a well preserved distal end of a left tarsometatarsus (figs. 4, 5 and 9). Milne-Edwards's statement "de même os" refers to tarsometatarsus as such, not to the specimen.

Paralectotypes from Sansan (all in MNHN; list derived from Cheneval 2000: 374-375): right coracoid (Sa-1412), three cranial ends of left coracoids (Sa-1459, Sa-1607, Sa-1670), sternal end of left coracoid (Sa-1608), sternal end of right coracoid (Sa-1671), two proximal ends of

left scapulae (Sa-1665, Sa-1667), proximal ends of three right scapulae (Sa-1663, Sa-1664, Sa-1666), left humerus (Sa-1246), distal end of left humerus (Sa-1316), distal end of right humerus (Sa-11317), eight proximal ends of left ulnae (Sa-1219, Sa-1309, Sa-1403, Sa-1468, Sa-1652, Sa-1659-1661), seven proximal ends of right ulnae (Sa-1266, Sa-1648, Sa-1653, Sa-1655, Sa-1656, Sa-1658, Sa-1662), seven distal ends of left ulnae (Sa-1389, Sa-1419, Sa-1647, Sa-1649-1651, Sa-1657), five distal ends of right ulnae (Sa-1294, Sa-1315, Sa-1420, Sa-1469, Sa-1654), three right carpometacapi (Sa-1291, Sa-1400, Sa-1615), six proximal ends of left carpometacarpus (Sa-1305, Sa-1332, Sa-1333, Sa-1630, Sa-1631, Sa-1635), nine proximal ends of right carpometacarpus (Sa-1277, Sa-1286, Sa-1287, Sa-1301, Sa-1331, Sa-1609, Sa-1632, Sa-1636, Sa-1637), six distal ends of left carpometacarpus (Sa-1424, Sa-1633, Sa-1634, Sa-1638, Sa-1640, Sa-1643), three distal end of right carpometacarpus (Sa-1639, Sa-1641, Sa-1642), two left phalanges I digiti majoris (Sa-1611, Sa-1644), right phalanx digiti majoris (Sa-1610), two proximal ends of left femora (Sa-1604, Sa-1612), three distal ends of right femora (Sa-1605, Sa-1613, Sa-1614), proximal end of left tibiotarsus (Sa-1437), three proximal ends of right tibiotarsi (Sa-1255, Sa-1433, Sa-1434), 14 distal ends of left tibiotarsi (Sa-1247, Sa-1439-1441, Sa-1444, Sa-1446, Sa-1447, Sa-1449, Sa-1452, Sa-1453, Sa-1455, Sa-1458, Sa-1497, Sa-1646), 12 distal ends of right tibiotarsi (Sa-1313, Sa-1330, Sa-1435, Sa-1436, Sa-1442, Sa-1443, Sa-1445, Sa-1448, Sa-1450, Sa-1451, Sa-1454, Sa-1494), five proximal ends of left tarsometatarsi (Sa-1278, Sa-1289, Sa-1323, Sa-1616, Sa-1621), five proximal ends of right tarsometatarsi (Sa-1256, Sa-1267, Sa-1491, Sa-1617, Sa-1619), 13 distal ends of left tarsometatarsi (Sa-1258, Sa-1275, Sa-1283, Sa-1295, Sa-1322, Sa-1325, Sa-1326, Sa-1329, Sa-1392, Sa-1431, Sa-1624-1626), 13 distal ends of right tarsometatarsi (Sa-1257, Sa-1302, Sa-1311, Sa-1321, Sa-1324, Sa-1327, Sa-1328, Sa-1618, Sa-1620, Sa-1622, Sa-1623, Sa-1627, Sa-1628), and right metatarsus I (Sa-1541). Figured by Milne-Edwards, 1869-1871, pl. 151 (reconstructed skeleton), pl. 52, fig. 1-3 (tarsometatarsus), fig. 4, 4a, 5, 5a, 9 (distal end of tarsometatarsus), fig. 6, 6a, 7, 8 (proximal end of tarsometatarsus), fig. 10, 10a, 11, 11a, 12, 12a, 15 (distal end of tibiotarsus Sa-1247), 13-14 (proximal end of tibiotarsus Sa-1255), 16-18 (proximal end of femur Sa-1604), 16-17, 19 (distal end of femur Sa-1605), 20, 20a, 21, 21a (cranial end of coracoid Sa-1607 and sternal end of coracoid Sa-1608), 22-24 (humerus Sa-1246), 27, 27a, 28, 28a (carpometacarpus Sa-1615), and pl. 153, fig. 21a-c (metatarsus I Sa-1541), and Cheneval 2000, fig. 29 (humerus Sa-1246), 30 (carpometacarpus Sa-1615), 31 (proximal end of tibiotarsus Sa-1255), 32 (distal end of tibiotarsus Sa-1247), and 33 (proximal end of tarsometatarsus Sa-1258).]

Miocorax [sic = *Mocorvus*] *larteti* (Milne-Edwards): Lambrecht 1933: 636 [New combination.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1871, Cheneval 2000).

Remarks: Taxonomic position of this species requires reexamination.

Genus *Corvus* Linnaeus

Corvus Linnaeus, 1758 [Modern genus.]

Remarks: Three size classes of Plio-Pleistocene *Corvus* crows can be distinguished in Europe, which correspond to the modern species *Corvus monedula* Linnaeus, *Corvus corone* Linnaeus plus *Corvus frugilegus* Linnaeus, and *Corvus corax* Linnaeus. For biometrical studies on the bones of modern and fossil corvids of the Holarctic see Mourer-Chauviré (1975a: 228-258), Magish & Harris (1976), Bährmann (1978), Moreno (1986), Kessler & Moldvai (1993), and Tomek & Bocheński (2000).

Corvus corax Linnaeus – Common Raven

Corvus corax Linnaeus, 1758 [Modern species.]

Corvus fossilis Giebel, 1847: 16 [Holotype from Seveckenberg: right tibiotarsus (in two

pieces); MNHN, uncatalogued. Not figured.]

Corvus pliocaenus janossyi Mourer-Chauviré, 1975a: 230 [Holotype from Saint-Estève-Janson – layer D: left humerus; coll. Bonifay CD66-D-11290. Figured by Mourer-Chauviré 1975a, pl. 18, fig. 1-2.]

Corvus antecorax Mourer-Chauviré, 1975a: 239 [Holotype from Fage – layer C5 (Mourer-Chauviré, in litt., 2001): left humerus; FSL 41707. Not figured.]

Corvus corax antecorax Mourer-Chauviré: Brodkorb 1978: 170 [New rank.]

Corvus janossyi Mourer-Chauviré: Jánossy 1979: 31 [New rank.]

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b and 1990b sub *Corvus* sp., Jánossy 1992 sub *C. janossyi*); early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub *C. antecorax*), Untermassfeld, Germany (Jánossy 1997 sub *C. aff. janossyi*), Valerots, France (Mourer-Chauviré 1975a sub *C. antecorax*), Beremend 16, Hungary (Jánossy 1992), Beremend 17, Hungary (Jánossy 1992), and Betfia 5, Romania (Kessler 1975); early Pleistocene (MQ 1b) of Sima del Elefante – layers E-12, E-13, E-14, Spain (Rosas et al. 2001 sub *C. antecorax*), and Stránská skála – Musil's talus cone, layers 13, 15e, Czechia (Mlíkovský 1995 sub *C. pliocaenus*); middle Pleistocene (MQ 2A) of Arago, France (Mourer-Chauviré 1975a sub *C. cf. antecorax*), Saint-Estève-Janson – layers B, C, D, E, E/F, G, H, France (Mourer-Chauviré 1975a sub *C. pliocaenus janossyi*), and Tarkö – layer 2, Hungary (Jánossy 1979 sub *C. janossyi*, Jánossy 1986 sub *C. cf. corax*); and middle Pleistocene (MQ 2B) of Lunel Viel – cave 1, layers 2, 4, 6, France (Mourer-Chauviré 1975a sub *C. antecorax*), Lunel-Viel – cave 1, layer 7, France (Mourer-Chauviré 1975a sub *C. pliocenicus janossyi*), Orgnac 3 – layers f, h, i, j, France (Mourer-Chauviré 1975a sub *C. antecorax*), Fage – layers 2/3, 4, CO, and 29 through 41, France (Mourer-Chauviré 1975a sub *C. antecorax*), Fage – layer 33, France (Mourer-Chauviré 1975a sub *C. pliocenicus janossyi*), Lazaret 8 – layer 17, France (Mourer-Chauviré 1975a sub *C. antecorax*), Biehle, France (Mourer-Chauviré 1975a sub *C. antecorax*), and Azé 2 – layer 6, France (Mourer-Chauviré 1975a sub *C. antecorax*); and late Pleistocene (MQ 2C) of Seveckenberg, Germany (Giebel 1847 sub *C. fossilis*, Mlíkovský orig.).

Remarks: I restudied the holotypical tibiotarsus of *Corvus fossilis* Giebel, finding that it belongs to a large *Corvus* of the size of *Corvus corax* Linnaeus. *Corvus pliocaenus janossyi* Mourer-Chauviré was based on a humerus, which is disproportionately large for most other bones from Saint-Estève-Janson, agreeing in size with the same element of *Corvus corax* Linnaeus (see Mourer-Chauviré 1975a for measurements). Hence, I synonymize here *Corvus fossilis* Giebel and *Corvus pliocaenus janossyi* Mourer-Chauviré with the modern *Corvus corax* Linnaeus.

See Brodkorb (1978: 170) for possible synonyms of *Corvus corax* Linnaeus from the North American Pleistocene.

***Corvus corone* Linnaeus – Carrion Crow**

Corvus corone Linnaeus, 1758 [Modern species.]

Corvus cornix fossilis Giebel, 1847 [Holotype from Monte Reale: right (if the figure is not side-reversed) femur; present location unknown. Figured by Wagner 1832, pl. 2, fig. 53.

This is – by my present decision on the principle of the first reviser – a junior primary homonym of *Corvus fossilis* Giebel.]

Numenius sp. (*pliocaenus*) Portis, 1887: 194 [Nomen nudum; no description or indication.]

Numenius sp. (*pliocaenus*) Portis, 1889: 13 [Holotype from Montecarlo: distal end of right ulna; IGF 937. Figured by Portis 1889, pl. 1, fig. 26a-b, Regàlia 1902, pl. 27, fig. 1.]

Corvus pliocaenus (Portis): Regàlia 1902: 220 [New combination.]

Corvus praecorax Depéret, 1892: 691 [Syntypes from Perpignan: distal end of right humerus, tibiotarsus, and left tarsometatarsus; FSL. Figured by Depéret 1897, pl. 13, fig. 12-13 (tarsometatarsus), 14 + 14a (tibiotarsus), and 17 + 17a (humerus).]

Corvus betfianus Kretzoi, 1962: 172 [Holotype from Betfia 5: distal end of left carpometacarpus; RMO 1899a/4. Figured by Jurcsák & Kessler 1973, fig. 31.]

Corvus simionescui Kessler, 1979: 136 [Holotype from Malușteni-Berești: left coracoid; LPUB 211. Figured by Kessler 1979, fig. 3-4.]

Distribution: Late Miocene (MN 11-13) of Il'inka, Ukraine (Vojinstvens'kyj 1967 sub *C. cf. corone*); early Pliocene (MN 15) of Perpignan, France (Depéret 1892, 1897, and Mourer-Chauviré 1975a: 248 sub *C. praecorax*), and of Malușteni-Berești, Romania (Kessler 1979 sub *C. simionescui*); late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Mourer-Chauviré et al. 1977 and Sondaar et al. 1995 sub *C. pliocaenus*), Montoussé 5, France (Clot et al. 1976a,b sub *C. pliocaenus*), Slivnica, Bulgaria (Boev 1995a and 2000a sub *C. cf. praecorax*), and Tourkobounia 2, Greece (Mourer-Chauviré 1980b sub *C. pliocaenus*); Pliocene or Pleistocene (MN 16 - MQ 1) of Montecarlo, Italy (Portis 1887 and 1889 sub *Numenius pliocaenus*, Regàlià 1902 sub *C. pliocaenus*);

early Pleistocene (MQ 1b) of Soave, France (Mourer-Chauviré 1980b sub *C. pliocaenus*), Stránská skála – Čapek's sites 5a, 9a, Czechia (Jánossy 1972), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972), Stránská skála – Musil's talus cone, layers 5, 13, 15e, Czechia (Mlíkovský 1995b), and Betfia 5, Romania (Kretzoi 1962 sub *C. betfianus*, Kessler 1975 sub *C. corone/betfianus*); middle Pleistocene (MQ 2A) of Arago, France (Mourer-Chauviré 1975a sub *C. corone* and *C. pliocaenus*); and middle Pleistocene (MQ 2B) of Saint-Sol-Belcastel, France (Philippe et al. 1980 sub *C. pliocaenus*).

Remarks: *Corvus cornix fossilis* Giebel was based on a complete femur, which was found by C.L. Nitzsch (in Wagner 1833) to be identical with the same element of the modern *Corvus cornix* Linnaeus (= *Corvus corone* Linnaeus), which is supported by the figure in Wagner (1832). *Numenius* (= *Corvus*) *pliocaenus* Portis, *Corvus praecorax* Depéret, *Corvus betfianus* Kretzoi, and *Corvus simionescui* Kessler fall in the same size category with the modern *Corvus corone* Linnaeus. In absence of the evidence to the contrary, I synonymize here *Corvus cornix fossilis* Giebel, *Numenius pliocaenus* Portis, *Corvus praecorax* Depéret, and *Corvus betfianus* Kessler with *Corvus corone* Linnaeus.

It is symptomatic, that the modern *Corvus frugilegus* Linnaeus, 1758 (Rook) was not identified from pre-middle Pleistocene deposits of Europe. This species is osteologically similar to *Corvus corone* Linnaeus, and the two species cannot be separated from each other on the basis of the postcranial skeleton with certainty, although the bones of *C. corone* Linnaeus tend to be larger and more robust. It is well possible, that some of the records listed here under *Corvus corone* Linnaeus belong to *Corvus frugilegus* Linnaeus.

***Corvus monedula* Linnaeus – Western Jackdaw**

Corvus monedula Linnaeus, 1758 [Modern species.]

Pica pica major Jánossy, 1972: 59 [Lectotype from Stránská skála – Absolon's 2nd cave, whitish fossilization (here selected): distal end of left tarsometatarsus; MMB Abs-146. Not figured. Paralectotypes from Stránská skála – Absolon's 2nd cave (whitish fossilization): right (not left as stated by Jánossy 1972: 59) tibiotarsus (MMB Abs-1003), distal end of left tarsometatarsus (MMB Abs-144); Stránská skála – Absolon's 2nd cave (black fossilization): left humerus (MMB Abs-134), left femur (MMB Abs-131), proximal end of right tibiotarsus (MMB Abs-1084), left (not right as stated by Jánossy 1972: 59) tibiotarsus lacking proximal end (MMB Abs-138), left tarsometatarsus (MMB Abs-132); Stránská skála – unknown site (whitish fossilization): distal end of right humerus (MMB SS-298); and Stránská skála – unknown site (black fossilization): distal end of left humerus (MMB SS-19). Not figured.]

Corvus moravicus Mlíkovský, 1995b: 120 [Holotype from Stránská skála – Musil's talus cone, layer 4: distal end of left femur; MMB 804/2. Figured by Mlíkovský 1995b, fig. 1.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 1-3, 4a, 4c, 5a, 9a, 9b, Czechia (Jánossy 1972 sub

Coloeus cf. monedula), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *Coloeus cf. monedula*), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *Pica pica major*, Mlíkovský orig.), and Stránská skála – Musil's talus cone, layers 3a, 4a, 8, 10, 13, 15a, 15e, Czechia (Mlíkovský 1995b).

Remarks: Black and whitish fossilized bones evidently originated from different sites within the Stránská skála complex. Black bones are older than Quaternary in general appearance and it is possible that Absolon excavated in his cave a fossiliferous pocket of Pliocene or late Miocene age (O. Fejfar, pers. communication). All black bones attributed by Jánossy (1972) to his *Pica pica major* indeed belong to the genus *Pica* Brisson and indeed are larger than corresponding elements of the modern European *Pica pica* (Linnaeus), representing an extinct species of *Pica* Brisson. However, I prefer to leave this species unnamed, because it is of unknown age. Whitish colored bones are apparently of Quaternary origin. One of the whitish syntypes (distal end of tarsometatarsus Abs-146) belongs to the modern *Nucifraga caryocatactes* (Linnaeus), while all other (as listed above, incl. the lectotype) originated from juvenile Jackdaws of the modern species *Corvus monedula* Linnaeus (Mlíkovský orig.). Accordingly, I synonymize here *Pica pica major* Jánossy with the latter species.

According to the data in Kessler & Moldvai (1993), *Corvus moravicus* Mlíkovský is not larger than the modern *C. monedula* Linnaeus (contra Mlíkovský 1995b). Hence, I synonymize here the former with the latter species.

***Corvus* sp.**

Distribution: Late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1993); late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991); late Pliocene (MN 17) of Puebla de Valverde, Spain (Adrover et al. 1974); and late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 2000a).

Genus *Garrulus* Brisson

Garrulus Brisson, 1760 [Modern species.]

***Garrulus glandarius* (Linnaeus) – Eurasian Jay**

Corvus glandarius Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of Montoussé 5, France (Clot et al. 1976a,b); early Pleistocene (MQ 1a) of Untermassfeld, Germany (Jánossy 1997 sub *G. aff. glandarius*), Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *G. cf. glandarius*, Mlíkovský 1998d), Beremend 17, Hungary (Jánossy 1992 sub *G. cf. glandarius*), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *G. aff. glandarius*); and early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d sub *G. ?glandarius* or *Garrulus* superspecies), Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub *G. aff. glandarius*), Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *G. aff. glandarius*), and Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b).

***Garrulus* sp.**

Distribution: Early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965).

Genus *Pica* Brisson

Pica Brisson, 1760 [Modern genus]

***Pica pica* (Linnaeus) – Common Magpie**

Corvus Pica Linnaeus, 1758 [Modern species.]

Distribution: Early Pliocene (MN 15) of Csarnóta 1 and 2, Hungary (Jánossy 1979 sub *P. aff. pica major*); late Pliocene (MN 18) of Villány 3, Hungary (Jánossy 1979 sub *P. aff. pica*

major); early Pleistocene (MQ 1b) of Nagyharsányhegy 1-4, Hungary (Jánossy 1979 sub *P. aff. pica major*); middle Pleistocene (MQ 2A) of Saint-Estève-Janson – layers B, C? and H, France (Mourer-Chauviré 1975a sub *P. pica major*), and Vértesszőllős 2, Hungary (Jánossy 1979 sub *P. aff. pica major*); and middle Pleistocene (MQ 2B) of Lunel-Viel, France (Mourer-Chauviré 1975a). The record from the middle Pleistocene (MQ 2A) of Hundsheim, Austria (Jánossy 1974b) is invalid, because I found that four out of six relevant bones belong to *Pyrrhonorax graculus* (Linnaeus), one to *Garrulus glandarius* (Linnaeus) and one is indeterminate (Mlíkovský orig.).

***Pica* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b, 2000a).

Genus *Nucifraga* Brisson

Nucifraga Brisson, 1760 [Modern species.]

***Nucifraga caryocatactes* (Linnaeus) – Spotted Nutcracker**

Corvus Caryocatactes Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub cf. *N. caryocatactes*); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b).

***Nucifraga* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a sub ?*Nucifraga* sp.).

Genus *Pyrrhonorax* Tunstall

Pyrrhonorax Tunstall, 1771 [Modern genus.]

***Pyrrhonorax graculus* (Linnaeus) – Alpine Chough**

Corvus graculus Linnaeus, 1766 [Modern species]

Pyrrhonorax graculus vetus Kretzoi, 1962: 173 [Nomen nudum; conditionally proposed after 1960.]

Pyrrhonorax graculus vetus "Kretzoi" Mourer-Chauviré, 1975a: 222 [Holotype from Betfia 5: left carpometacarpus; RMO 297/2. Not figured. Mourer-Chauviré (1975a: 222) was first to use this name in a manner required by the ICZN (1999). She thus becomes its author, although she attributed it to Kretzoi 1962.]

Distribution: Early Pliocene (MN 15) of Csarnóta 2 – layers 2 and 23, Hungary (Jánossy 1979 sub *P. aff. graculus*); late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b sub *P. graculus* cf. *vetus*, Boev 2000a sub *P. cf. graculus*); late Pliocene (MN 18) of Meda Gran, Spain (Villalta 1965);

early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *P. aff. graculus*); early Pleistocene (MQ 1b) of Koněprusy C-718, Czechia (Mlíkovský 1996t), Včeláre 1, Slovakia (Jánossy 1979 sub *P. aff. graculus*), Betfia 5, Romania (Kretzoi 1962 and Mourer-Chauviré 1975a sub *P. graculus vetus*), and Petralona – layer 11, Greece (Kretzoi 1977 and Kretzoi & Poulianos 1981 sub *P. graculus vetus*); middle Pleistocene (MQ 2A) of Cimay, France (Mourer-Chauviré 1975a sub *P. graculus vetus*), Saint-Estève-Janson – layers B, C/D, E, F, G and H, France (Mourer-Chauviré 1975a sub *P. graculus vetus*), Arago, France (Mourer-Chauviré 1975a sub *P. graculus vetus*), and Petralona – layer 10, Greece (Kretzoi 1977 and Kretzoi & Poulianos 1981 sub *P. graculus vetus*); and middle Pleistocene (MQ 2B) of Fage – layers 2/3, 4, CO and 29-41, France (Mourer-Chauviré 1975a sub *P. graculus* cf. *vetus*), Orgnac 3 – layers g, h, France

(Mourer-Chauviré 1975a sub *P. graculus* cf. *vetus*), Lazaret 8 – layers iv, v, v1, v2, vi, vii, viii, ix, xi, xii, xiii, xiv, xv, xvi, xvii, xviii, France (Mourer-Chauviré 1975a sub *P. graculus* *vetus*), Suard, France (Mourer-Chauviré 1975a sub *P. graculus* *vetus*), and Aldène – layers v-vi, viii, ix, France (Mourer-Chauviré 1975a sub *P. graculus* *vetus*).

Extralimital records include early Pleistocene (MQ 1b) of Treugol'naja Cave – layers 6, 7a, 7b, Karachaevo-Cherkessia, Russia (Potapova & Baryšnikov 1993, Baryšnikov & Potapova 1995 sub *P. graculus* *vetus*); middle Pleistocene (MQ 2A) of Treugol'naja Cave – layers 5b, 5c, Karachaevo-Cherkessia, Russia (Potapova & Baryšnikov 1993, Baryšnikov & Potapova 1995 sub *P. graculus* *vetus*); and late Pleistocene (MQ 2C-D) of Treugol'naja Cave – layers 4b and 4d, Karachaevo-Cherkessia, Russia (Potapova & Baryšnikov 1993, Baryšnikov & Potapova 1995 sub *P. graculus* *vetus*).

***Pyrrhonorax pyrrhonorax* (Linnaeus) – Red-billed Chough**

Upupa pyrrhonorax Linnaeus, 1758 [Modern species]

Pyrrhonorax primigenius Milne-Edwards, 1875: 487 [Lectotype from Massat (here selected): left tarsometatarsus; MNHN 1906-26 partim. Not figured. Paralectotypes: right tarsometatarsus (MNHN 1906-26 partim), and proximal end of juvenile right tarsometatarsus (MNHN 1906-26 partim). Not figured. Mourer-Chauviré (1975a: 222) selected two tarsometatarsi from Massat deposit at MNHN as syntypes of this species. Her action is invalid according to the ICZN (1999).]

Pyrrhonorax pyrrhonorax primigenius Milne-Edwards: Mourer-Chauviré 1975a: 219 [New rank.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a sub *P. aff. pyrrhonorax*); late Pliocene (MN 18) of Meda Gran, Spain (Villalta 1965 sub *P. cf. pyrrhonorax*), and S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *P. cf. pyrrhonorax*); middle Pleistocene (MQ 2B) of Lazaret, layers iv, v, v1, v2, vi, vii, viii, ix, xiii, xiv, xv, xvi, xviii, France (Mourer-Chauviré 1975a sub *P. pyrrhonorax primigenius*), Fage – layers CO, 30-33, 35, 36, 38 and 40, France (Mourer-Chauviré 1975a sub *P. pyrrhonorax* cf. *primigenius*), and Lazaret, layers i-ii, iii, France (Mourer-Chauviré 1975a sub *P. pyrrhonorax primigenius*); and late Pleistocene (MQ 2D) of Massat, France (Milne-Edwards 1875 and Mourer-Chauviré 1975a sub *P. pyrrhonorax primigenius*), Vieulac, France (Rivière 1887 and 1892 sub *P. primigenius*), and Pageyral, France (Rivière 1887 and 1892 sub *P. primigenius*). Tyrberg (1998) listed *P. pyrrhonorax primigenius* Milne-Edwards from a number of other localities of Europe, referring to Mourer-Chauviré (1975a). However, the latter author listed these sites under the heading "*Pyrrhonorax pyrrhonorax* and *Pyrrhonorax pyrrhonorax primigenius*", and it follows from the subsequent text, that she recognized *primigenius* only from a much more limited number of sites.

Extralimital records include early Pleistocene (MQ 1b) of Treugol'naja Cave – layers 6, 7b, Karachaevo-Cherkessia, Russia (Potapova & Baryšnikov 1993, Baryšnikov & Potapova 1995 sub *P. p. primigenius*); middle Pleistocene (MQ 2A) Treugol'naja Cave – layers 5a, 5b and 5c, Karachaevo-Cherkessia, Russia (Potapova & Baryšnikov 1993, Baryšnikov & Potapova 1995 sub *P. p. primigenius*); and middle Pleistocene (MQ 2B) of Treugol'naja Cave – layers 4b and 4c, Karachaevo-Cherkessia, Russia (Potapova & Baryšnikov 1993 and Baryšnikov & Potapova 1995 sub *P. p. primigenius*).

Remarks: Paris (1912: 284) and Lambrecht (1921: 90) erroneously attributed *Pyrrhonorax primigenius* to Rivière ("1891" = 1892: 372).

***Pyrrhonorax* sp.**

Distribution: Late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 2000a).

Family Troglodytidae Swainson

Troglodytidae Swainson, 1831 [Modern family.]

Genus *Troglodytes* Vieillot*Troglodytes* Vieillot, 1807 [Modern genus.]***Troglodytes troglodytes* (Linnaeus) – Winter Wren***Motacilla Troglodytes* Linnaeus, 1758 [Modern species.]**Distribution:** Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *T. cf. troglodytes*).**Family Turdidae Rafinesque**

Turdidae Rafinesque, 1815 [Modern family.]

Muscicapidae Fleming, 1822 [Modern family.]

Genus *Erithacus* Cuvier*Erithacus* Cuvier, 1800 [Modern genus.]***Erithacus rubecula* (Linnaeus) – European Robin***Motacilla Rubecula* Linnaeus, 1758 [Modern species.]**Distribution:** Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *E. cf. rubecula*); and early Pleistocene (MQ 1b) of Boxgrove – quarry 2, England (Harrison & Stewart 1999 sub cf. *E. rubecula*).***Erithacus* sp.****Distribution:** Early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992).**Genus *Luscinia* Forster***Luscinia* Forster, 1817 [Modern genus.]***Luscinia svecica* (Linnaeus) – Bluethroat***Motacilla svecica* Linnaeus, 1758 [Modern species.]**Distribution:** Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *L. cf. svecica*).***Luscinia* sp.****Distribution:** Late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991); late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a sub array of *L. svecica*); and early Pleistocene (MQ 1a) of Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *Sylvia* sp., Mlíkovský 1998d).**Genus *Phoenicurus* Forster***Phoenicurus* Forster, 1817 [Modern genus.]***Phoenicurus ochruros* (Gmelin)***Motacilla Ochruros* Gmelin, 1774 [Modern species.]**Distribution:** Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001).

Genus *Saxicola* Bechstein

Saxicola Bechstein, 1803 [Modern genus.]

***Saxicola torquata* (Linnaeus) – Common Stonechat**

Muscicapa torquata Linnaeus, 1766 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *S. cf. torquata*); and early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub *S. cf. torquata*).

Genus *Oenanthe* Vieillot

Oenanthe Vieillot, 1816 [Modern genus.]

***Oenanthe oenanthe* (Linnaeus) – Northern Wheatear**

Motacilla Oenanthe Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *O. aff. oenanthe*).

***Oenanthe hispanica* (Linnaeus) – Black-eared Wheatear**

Motacilla hispanica Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001).

***Oenanthe* sp.**

Distribution: Late Pliocene (MN 18) of Montoussé 5, France (Clot et al. 1976a,b).

Genus *Monticola* Boie

Monticola Boie, 1822 [Modern genus.]

***Monticola solitarius* (Linnaeus) – Blue Rock Thrush**

Turdus solitarius Linnaeus, 1758 [Modern species.]

Turdus cyanus Linnaeus, 1758 [Modern species.]

Distribution: Early Pliocene (MN 15) of Perpignan, France (Depéret 1897 sub *Turdus* aff. *cyanus*).

Genus *Turdus* Linnaeus

Turdus Linnaeus, 1758 [Modern species.]

***Turdus merula* Linnaeus – Common Blackbird**

Turdus merula Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b sub *T. cf. merula*); late Pliocene (MN 18) of Montoussé 5, France (Clot et al. 1976a,b sub *T. cf. merula*), and S'Onix, Mallorca (Sondaar et al. 1995 sub *T. cf. merula*); early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub *T. cf. merula*), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *T. cf. merula*); and early Pleistocene (MQ 1b) of West Runton – Upper Freshwater Bed, England (Harrison 1979d sub *T. ?merula* or *T. merula/boulboul* superspecies).

***Turdus pilaris* Linnaeus – Fieldfare**

Turdus pilaris Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 5a, 9a and 9b, Czechia (Jánossy 1972 sub *T. cf. pilaris*).

***Turdus philomelos* Brehm – Song Thrush**

Turdus Ericetorum Turton, 1807 [Modern species.]

Turdus philomelos Brehm, 1831 [Modern species.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b sub *T. cf. philomelos*); late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 2000a sub *T. cf. philomelos*); early Pleistocene (MQ 1a) of Untermassfeld, Germany (Jánossy 1997 sub *T. aff. philomelos*), Beremend 16, Hungary (Jánossy 1992 sub *T. cf. philomelos*), Beremend 17, Hungary (Jánossy 1992 sub *T. cf. philomelos*), Betfia 2, Romania (Jánossy 1979 sub *T. cf. philomelos*), and Primorsk, Ukraine (Vojinstens'kyj 1967 sub *T. cf. ericetorum*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 8, Czechia (Jánossy 1972 sub *T. cf. philomelos*).

***Turdus iliacus* Linnaeus – Redwing**

Turdus iliacus Linnaeus, 1766 [Modern species.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a); late Pliocene (MN 18) of S'Onix, Mallorca (Sondaar et al. 1995); early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992 sub *T. aff. iliacus*); and early Pleistocene (MQ 1b) of Vallonnet, France (Mourer-Chauviré 1975a sub cf. *T. iliacus*).

***Turdus viscivorus* Linnaeus – Mistle Thrush**

Turdus viscivorus Linnaeus, 1758 [Modern species.]

Distribution: Early Pliocene (MN 15) of Csarnóta 2, layer 21, Hungary (Jánossy 1979 sub *T. viscivorus*-group); late Pliocene (MN 18) of Montoussé 5, France (Clot et al. 1976a,b sub *T. cf. viscivorus*); early Pleistocene (MQ 1) of Gènova, Mallorca (Mourer-Chauviré et al. 1977 sub *T. cf. viscivorus*); early Pleistocene (MQ 1a) of Valerots, France (Mourer-Chauviré 1975a sub cf. *T. viscivorus*), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *T. cf. viscivorus*-group); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 5a and 9a, Czechia (Jánossy 1972 sub *T. cf. viscivorus*), and Stránská skála – Absolon's 2nd cave, Czechia (Jánossy 1972 sub *T. cf. viscivorus*).

***Turdus* sp.**

Distribution: Late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991); early Pliocene (MN 15) of Ivanovce 1, Slovakia (Mlíkovský orig.); late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a); late Pliocene (MN 17) of Šandalja 1 – layer 6, Croatia (Malez-Bačić 1979); late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a); late Pliocene (MN 18) of Binigaus, Mallorca, Balearics (Alcover et al. 1981 sub *Turdus iliacus/philomelos*); early Pleistocene (MQ 1a) of Untermassfeld, Germany (Jánossy 1988), and Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *T. cf. viscivorus*, Mlíkovský 1998d sub *Turdus* – large sp.); and early Pleistocene (MQ 1b) of Koněprusy C-718, Czechia (Mlíkovský 1996t), Stránská skála – Musil's talus cone, layer 13, Czechia (Mlíkovský 1995b sub *Turdus* – large and medium spp.), Petralona – layers 15 and 24, Greece (Kretzoi 1977, Kretzoi & Poulianos 1981).

Genus *Muscicapa* Brisson

Muscicapa Brisson, 1760 [Modern genus.]

***Muscicapa striata* (Pallas) – Spotted Flycatcher**

Motacilla striata Pallas, 1764 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *M. cf. striata*).

***Muscicapa* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a).

Genus *Ficedula* Brisson

Ficedula Brisson, 1760 [Modern species.]

***Ficedula hypoleuca* (Pallas) – European Pied Flycatcher**

Motacilla hypoleuca Pallas, 1764 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub cf. *F. hypoleuca*).

***Ficedula* sp.**

Distribution: Late Pliocene (MN 18) of Montoussé 5, France (Clot et al. 1976a,b).

Family Sylviidae Leach

Sylviidae Leach, 1820 [Modern family.]

Genus *Cettia* Bonaparte

Cettia Bonaparte, 1834 [Modern genus.]

***Cettia* sp.**

Distribution: Late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991).

Genus *Locustella* Kaup

Locustella Kaup, 1829 [Modern genus.]

***Locustella* sp.**

Distribution: Late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1993).

Genus *Acrocephalus* Naumann

Acrocephalus Naumann, 1811 [Modern genus.]

***Acrocephalus palustris* (Bechstein) – Marsh Warbler**

Motacilla s. *Sylvia palustris* Bechstein, 1811 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *A. cf. palustris*).

***Acrocephalus arundinaceus* (Linnaeus) – Great Reed Warbler**

Turdus arundinaceus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1) of Betfia 7, Romania (Kessler 1975 sub *A. cf. arundinaceus*).

***Acrocephalus* sp.**

Distribution: Late Miocene (MN 9) of Rudabánya, Hungary (Jánossy 1993); late Miocene

(MN 13) of Polgárdi 4, Hungary (Jánossy 1991), and Polgárdi 5, Hungary (Jánossy 1991); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 5, Czechia (Mlíkovský 1995b).

Genus *Hippolais* Baldenstein

Hippolais Baldenstein, 1827 [Modern genus.]

***Hippolais* sp.**

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Jánossy 1979).

Genus *Sylvia* Scopoli

Sylvia Scopoli, 1769 [Modern species.]

***Sylvia communis* Latham – Common Whitethroat**

Sylvia communis Latham, 1787 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *S. cf. communis*).

***Sylvia atricapilla* (Linnaeus) – Blackcap**

Motacilla atricapilla Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995). The record of *Sylvia* aff. *atricapilla* from the early Pleistocene (MQ 1a) of Deutsch-Altenburg 4B, Austria (Jánossy 1981) was reidentified as belonging to *Luscinia* sp. (Mlíkovský 1998d).

***Sylvia* sp.**

Distribution: Late Miocene (MN 13) of Polgárdi 4, Hungary (Jánossy 1991).

Genus *Phylloscopus* Boie

Phylloscopus Boie, 1826 [Modern genus.]

***Phylloscopus sibilatrix* (Bechstein) – Wood Warbler**

Motacilla sibilatrix Bechstein, 1793 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Čerdževica, Bulgaria (Boev 2000c).

Genus *Regulus* Cuvier

Regulus Cuvier, 1800 [Modern genus.]

***Regulus ignicapillus* (Temminck) – Firecrest**

Sylvia ignicapilla Temminck, 1820 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *R. cf. ignicapillus*).

Family Prunellidae Richmond

Prunellidae Richmond, 1908 [Modern family.]

Genus *Prunella* Vieillot

Prunella Vieillot, 1816 [Modern genus.]

***Prunella modularis* Linnaeus – Dunnock**

Motacilla modularis Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995); and early Pleistocene (MQ 1b) of Boxgrove – quarry 2, England (Harrison & Stewart 1999 sub cf. *P. modularis*).

***Prunella collaris* (Scopoli)**

Sturnus collaris Scopoli, 1769 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 4a, Czechia (Jánossy 1972 sub aff. *P. collaris*).

Family Aegithalidae Reichenbach

Aegithalidae Reichenbach, 1850 [Modern family.]

Genus *Aegithalos* Hermann

Aegithalos Hermann, 1804 [Modern species.]

***Aegithalos caudatus* (Linnaeus) – Long-tailed Tit**

Parus caudatus Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *A. cf. caudatus*); and early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *A. cf. caudatus*).

Family Paridae Vigors

Paridae Vigors, 1825 [Modern family.]

Sittidae Lesson, 1828 [Modern family.]

Genus *Parus* Linnaeus

Parus Linnaeus, 1758 [Modern genus.]

***Parus cristatus* Linnaeus – Crested Tit**

Parus cristatus Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *P. cf. cristatus*).

***Parus ater* Linnaeus – Coal Tit**

Parus ater Linnaeus, 1758

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *P. cf. ater*).

***Parus caeruleus* Linnaeus – Blue Tit**

Parus caeruleus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *P. cf. caeruleus*).

***Parus major* Linnaeus – Great Tit**

Parus major Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *P. cf. major*); and early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *P. cf. major*), and Tarchankut, Ukraine (Vojinstvens'kyj 1967).

***Parus* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1987b, 2000a).

Genus *Sitta* Linnaeus

Sitta Linnaeus, 1758 [Modern genus.]

***Sitta europaea* Linnaeus – Eurasian Nuthatch**

Sitta europaea Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *S. cf. europaea*; Mlíkovský 1998d doubted the record); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *S. cf. europaea*).

***Sitta* sp.**

Distribution: Late Miocene (MN 15) of Csarnóta 2 – layers 17, 19 and 21, Hungary (Jánossy 1974a, 1979); and late Pliocene (MN 16) of Rebielice Królewskie 1, Poland (Jánossy 1974a). The record from early Pleistocene (MQ 1a) of Deutsch-Altenburg 4B, Austria (Jánossy 1981) is invalid (Mlíkovský 1998d).

Family Certhiidae Leach

Certhiidae Leach, 1820 [Modern family.]

Genus *Certhia* Linnaeus

Certhia Linnaeus, 1758 [Modern genus.]

***Certhia familiaris* Linnaeus – Eurasian Treecreeper**

Certhia familiaris Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *C. cf. familiaris*).

Family Motacillidae Horsfield

Motacillidae Horsfield, 1821 [Modern family.]

Genus *Anthus* Bechstein

Anthus Bechstein, 1805 [Modern genus.]

***Anthus campestris* (Linnaeus) – Tawny Pipit**

Alauda campestris Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub *A. cf. campestris*).

***Anthus pratensis* (Linnaeus) – Meadow Pipit**

Alauda pratensis Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001).

***Anthus cervinus* (Pallas) – Red-throated Pipit**

Motacilla cervina Pallas, 1811 [Modern species.]

Distribution: Late Pliocene (MN 16) of Rębielice Królewskie 1, Poland (Jánossy 1974a sub *A. cf. cervinus*).

***Anthus* sp.**

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b, 1990b, 1992); late Pliocene (MN 17) of Vărșec, Bulgaria (Boev 2000a); early Pleistocene (MQ 1a) of Kövesvár, Hungary (Jánossy 1986), and Čerdženica, Bulgaria (Boev 2000c); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972), and Stránská skála – Musil's talus cone, layers 6 and 13, Czechia (Mlíkovský 1995b).

Genus *Motacilla* Linnaeus

Motacilla Linnaeus, 1758 [Modern genus.]

***Motacilla alba* Linnaeus – White Wagtail**

Motacilla alba Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979).

***Motacilla* sp.**

Distribution: Late Miocene (MN 13) of Polgárdi 5, Hungary (Jánossy 1991); late Pliocene (MN 17) of Vărșec, Bulgaria (Boev 2000a); and early Pleistocene (MQ 1b) of Stránská skála – Musil's talus cone, layer 3b, Czechia (Mlíkovský 1995).

Family Passeridae Rafinesque

Passeridae Rafinesque, 1815 [Modern family.]

Genus *Passer* Brisson

Passer Brisson, 1760 [Modern genus.]

***Passer montanus* (Linnaeus) – House Sparrow**

Fringilla montana Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *P. cf. montanus*).

Remarks: This is a doubtful record, based on a single carpometacarpus (Mlíkovský orig.).

***Passer* sp.**

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Mourer-Chauviré 1995a: 342).

Remarks: This record is questionable because of its age (Mlíkovský orig.).

Genus *Petronia* Kaup

Petronia Kaup, 1829 [Modern genus.]

***Petronia petronia* (Linnaeus) – Rock Sparrow**

Fringilla petronia Linnaeus, 1766 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub *P. cf. petronia*).

Family Fringillidae Leach

Fringillidae Leach, 1820 [Modern family.]

Genus *Fringilla* Linnaeus

Fringilla Linnaeus, 1758 [Modern genus.]

***Fringilla coelebs* Linnaeus – Common Chaffinch**

Fringilla coelebs Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b); late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995); and early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *F. cf. coelebs*), and Tarchankut, Ukraine (Vojinstvens'kyj 1967).

***Fringilla* sp.**

Distribution: Middle/late Miocene (MN 8-9) of Hostalets de Pierola, Spain (Villalta 1963); and late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a).

Genus *Serinus* Koch

Serinus Koch, 1816 [Modern genus.]

***Serinus* sp.**

Distribution: Late Pliocene (MN 16) of Beremend 15, Hungary (Jánossy 1987b, 1990b, 1992); and early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992).

Genus *Carduelis* Brisson

Carduelis Brisson, 1760 [Modern genus.]

***Carduelis chloris* (Linnaeus) – European Greefinch**

Loxia chloris Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001), and Mas Rambault, France (Mourer-Chauviré 1975a sub *C. chloris*); and early Pleistocene (MQ 1b) of Sima del Elefante – unknown layer, Spain (Rosas et al. 2001).

***Carduelis carduelis* (Linnaeus) – European Goldfinch**

Fringilla Carduelis Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b sub *C. cf. carduelis*); late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub *C. cf. carduelis*); early Pleistocene (MQ 1a) of Quibas, Spain (Montoya et al. 1999, 2001); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 1-3 and 9a, Czechia (Jánossy 1972 sub *C. aff. carduelis*, both these records are doubtful – Mlíkovský orig.).

***Carduelis spinus* (Linnaeus) – Eurasian Siskin**

Fringilla Spinus Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Mas Rambault, France (Mourer-Chauviré 1975a sub cf. *C. spinus*).

***Carduelis cannabina* (Linnaeus) – Common Linnet**

Fringilla cannabina Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1a) of Čerdženica, Bulgaria (Boev 2000c).

***Carduelis* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 2000a); and early Pleistocene (MQ 1a) of Beremend 17, Hungary (Jánossy 1992).

Genus *Loxia* Linnaeus

Loxia Linnaeus, 1758 [Modern species.]

Remarks: General reference: Tyrberg 1991a.

***Loxia patevi* Boev**

Loxia patevi Boev, 1999h: 53 [Holotype from Väršec: distal end of right humerus; NMNHS 307. Figured by Boev 1999h, fig. 2a.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b sub *Loxia* sp., Boev 1999a,h).

***Loxia curvirostra* Linnaeus – Common Crossbill**

Loxia curvirostra Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 5a, Czechia (Jánossy 1972 sub cf. *L. curvirostra*, the record is doubtful – Mlíkovský, orig.).

***Loxia* sp.**

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Mourer-Chauviré 1995b: 342).

Genus *Pyrrhula* Brisson

Pyrrhula Brisson, 1760 [Modern genus.]

***Pyrrhula pyrrhula* (Linnaeus) – Eurasian Bullfinch**

Loxia Pyrrhula Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's site 9a, Czechia (Jánossy 1972 sub *P.* cf. *pyrrhula*).

***Pyrrhula* sp.**

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b).

Genus *Coccothraustes* Brisson

Coccothraustes Brisson, 1760 [Modern species.]

***Coccothraustes coccothraustes* (Linnaeus) – Hawfinch**

Loxia Coccothraustes Linnaeus, 1758 [Modern species.]

Distribution: Late Pliocene (MN 18) of S'Onix, Mallorca, Balearics (Sondaar et al. 1995 sub

C. cf. coccothraustes); early Pleistocene (MQ 1a) of Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *Pinicola cf. enucleator*, Mlíkovský 1998d), and Betfia 2, Romania (Čapek 1917, Jánossy 1979 sub *C. cf. coccothraustes*); and early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 8 and 9a, Czechia (Jánossy 1972 sub *C. cf. coccothraustes*), and Stránská skála – Musil's talus cone, layer 10b, Czechia (Mlíkovský 1995b).

Family Emberizidae Vigors

Emberizidae Vigors, 1825 [Modern family.]

Genus *Emberiza* Linnaeus

Emberiza Linnaeus, 1758 [Modern genus.]

Emberiza citrinella Linnaeus – Yellowhammer

Emberiza citrinella Linnaeus, 1758 [Modern species.]

Distribution: Early Pleistocene (MQ 1b) of Stránská skála – Čapek's sites 5a and 9a, Czechia (Jánossy 1972 sub *E. cf. citrinella*).

Emberiza sp.

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1997b, 2000a); late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 2000a); early Pleistocene (MQ 1a) of Deutsch-Altenburg 4B, Austria (Jánossy 1981 sub *Serinus* sp., *Serinus cf. canaria* and *Serinus aff. serinus*. All these names refer to a single humerus, reidentified by Mlíkovský 1998d as belonging to *Emberiza* sp.), Beremend 17, Hungary (Jánossy 1992); and early Pleistocene (MQ 1b) of Nagyharsányhegy 1-4, Hungary (Jánossy 1979).

Family incertae sedis

Alauda gypsurum Portis

Alauda gypsurum Portis, 1887: 186 [Holotype from Senigallia: incomplete skeleton in a slab; MCI, uncatalogued. Figured by Portis 1887, fig. 4.]

Distribution: Late Miocene (MN 13) of Senigallia, Italy (Portis 1887).

Alauda major Portis

Alauda major Portis, 1887: 190 [Holotype from Gabbro: incomplete hindlimbs in a slab; present location unknown – Delle Cave 1996: 270].

Distribution: Late Miocene (MN 13) of Gabbro, Italy (Portis 1887).

Anthus bosniaskii Pycraft

Anthus bosniaskii Pycraft, 1909: 370 [Holotype from Gabbro: incomplete hindlimbs in slab and counter-slab; BMNH. Figured by Pycraft 1909, fig. 47.]

Distribution: Late Miocene (MN 13) of Gabbro, Italy (Pycraft 1909).

Coccothraustes balcanicus Boev

Coccothraustes balcanicus Boev, 1998b: 89 [Holotype from Slivnica: imperfect anterior part of a mandible; NMNHS 440. Figured by Boev 1998b, fig. 2a-c, 3b-c (bottom).]

Distribution: Late Pliocene (MN 18) of Slivnica, Bulgaria (Boev 1998b).

Remarks: Posterior end of the symphysis of the holotypical mandible does not resemble

mandibles of *Coccothraustes* Brisson (Mlíkovský, orig.).

***Coccothraustes simeonovi* Boev**

Coccothraustes simeonovi Boev, 1998b: 92 [Holotype from Väršec: distal end of left humerus; NMNHS 120. Figured by Boev 1998b, fig. 4a-b.]

Distribution: Late Pliocene (MN 17) of Väršec, Bulgaria (Boev 1998b).

***Corvus hungaricus* Lambrecht**

Corvus hungaricus Lambrecht, 1916: 164 [Holotype from Nagyharánhegy 1-4: left tarsometatarsus; present location unknown. Figured by Lambrecht 1916, fig. 2.]

Distribution: Late Pliocene (MN 18) of Senèze, France (Stehlin 1923, Lambrecht 1933); early Pleistocene (MQ 1b) of Nagyharsányhegy 1-4, Hungary (Lambrecht 1916, Mourer-Chauviré 1975a: 249, Jánossy 1979: 31).

Remarks: The holotypical tarsometatarsus of this species certainly belongs to a rather large songbird, but no evidence has been presented that it belongs to the family Corvidae or even to the genus *Corvus* Linnaeus, from which it differs in having the tarsometatarsus very slender. I relegate thus *Corvus hungaricus* to the category of Passeriformes incertae sedis until its holotype is located and restudied.

***Fringilla trochanteria* Giebel**

Fringilla trochanteria Giebel, 1847: 15 [Holotype from Seveckenberg: right femur with damaged ends; MNHN uncatalogued. Not figured.]

Distribution: Late Pleistocene (MQ 2C) of Seveckenberg, Germany (Giebel 1847).

Remarks: The holotypical femur is indeterminate within the Passeriformes (Mlíkovský, orig.).

***Hirundo fossilis* Giebel**

Hirundo fossilis Giebel, 1847: 18 [Syntypes from Seveckenberg: furcula, distal end of left ulna and distal end of right ulna (originally referred to as distal ends of two radii), right carpometacarpus, and proximal end of right tarsometatarsus; MNHN uncatalogued. Not figured. I did not find (in 1999) among the syntypes the furcula, but two unidentifiable scraps of tiny bones were present in the box with the remaining syntypes. I am not able to decide whether one or both of them were regarded by Giebel as the syntypical furcula.]

Distribution: Late Pleistocene (MQ 2C) of Seveckenberg, Germany (Giebel 1847).

Remarks: Both ulnae, the carpometacarpus and the tarsometatarsus belong to small song bird(s), but cannot be identified more exactly. Nevertheless, none of the syntypes belongs to any member of the Hirundinidae, which are morphologically distinct (Mlíkovský, orig.).

***Lanius miocaenus* Milne-Edwards**

Lanius miocaenus Milne-Edwards, 1871: 391 [Holotype from Saint-Gérard-le-Puy: left humerus; MNHN Av-2855. Figured by Milne-Edwards 1869-1871, pl. 159, fig. 1, 1a, 2-3.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871).

***Motacilla humata* Milne-Edwards**

Motacilla humata Milne-Edwards, 1871: 390 [Holotype from Saint-Gérard-le-Puy: left tibiotarsus; MNHN Av-2848. Figured by Milne-Edwards, 1869-1871, pl. 159, fig. 10, 10a, 11-12.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871).

***Motacilla major* Milne-Edwards**

Motacilla major Milne-Edwards, 1871: 391 [Syntypes from Saint-Gérard-le-Puy: distal end of tibiotarsus (MNHN Av-2845), and right tarsometatarsus (MNHN Av-2864). Figured by Milne-Edwards 1869-1871, pl. 158, fig. 1-6 (tarsometatarsus), and pl. 159, fig. 13-15 (tibiotarsus).]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871).

Remarks: Lambrecht (1933: 641) observed that the syntypical tarsometatarsus of this species is figured with an open hypotarsus and that this would preclude the species from being a songbird. However, my direct examination of the specimen in MNHN showed that its hypotarsus is broken and that it is correctly assigned to the Passeriformes. The syntypes are indeterminate within the latter order (Mlíkovský, orig.).

***Parus lugubris fossilis* Čapek**

Parus lugubris fossilis Čapek, 1917: 31 [Holotype from Betfia 2: premaxilla; present location unknown. Not figured.]

Distribution: Early Pleistocene (MQ 1a) of Betfia 2, Romania (Čapek 1917).

Remarks: Jánossy (1979) listed this record sub *Parus* cf. *lugubris* Temminck.

***Regulus bulgaricus* Boev**

Regulus bulgaricus Boev, 1999f: 110 [Holotype from Vărșec: left ulna; MNHNS 24. Figured by Boev 1999f, fig. 1, 3 (left), 4 (left).]

Distribution: Late Pliocene (MN 17) of Vărșec, Bulgaria (Boev 1999f).

***Sitta senogalliensis* Portis**

Sitta Senogalliensis Portis, 1887: 184 [Holotype from Senigallia: incomplete skeleton in a slab; MCI, uncatalogued. Figured by Portis 1887, fig. 3.]

Distribution: Late Miocene (MN 13) of Senigallia, Italy (Portis 1887).

***Troglodytes gracilis* Brunner**

Troglodytes gracilis Brunner, 1958: 511 [Holotype from Breitenberghöhle: right carpometacarpus; BSP 1982-X-6651. Figured by Brunner 1958, fig. 10.]

Distribution: Late Pleistocene (MQ 2C-E) of Breitenberghöhle, Germany (Brunner 1958). The age of the locality is uncertain (Tyrberg 1998: 205).

***Turdicus tenuis* Kretzoi**

Turdicus tenuis Kretzoi, 1962: 173 [Holotype from Betfia 5: imperfect left coracoid; RMO 3. Not figured.]

Distribution: Early Pleistocene (MQ 1b) of Betfia 5, Romania (Kretzoi 1962).

Remarks: This species was tentatively placed in the Turdidae by Kretzoi (1962). It is indeterminate within the Passeriformes (Mlíkovský, orig.). *Turdicus tenuis* Kretzoi, 1962 is type by monotypy of the genus *Turdicus* Kretzoi, 1962: 173.

***Turdoides borealis* Jánossy**

Turdoides borealis Jánossy, 1979: 27 [Holotype from Osztramos 1: proximal end of left humerus; GIB V-78119. Figured by Jánossy 1979, fig. 4/7.]

Distribution: Early Pliocene (MN 15) of Osztramos 1, Hungary (Jánossy 1979, see also Jánossy 1972), and Csarnóta 2, Hungary (Jánossy 1979, tentatively referred).

Remarks: The holotype is badly eroded, and the taxonomic position of this species within the Passeres remains unclear (Mlíkovský 1992b: 444). *Turdoides borealis* Jánossy is type by original designation of the genus *Turdoides* Jánossy, 1979: 27.

***Turdus bresciensis* Giebel**

Turdus bresciensis Giebel, 1847: 13 [Holotype from Monte Reale: right (if the figure is not side-reversed) humerus; present location unknown. Figured by Wagner 1832, pl. 2, fig. 57a-c.]

Distribution: Middle Pleistocene (MQ 2A-C) of Monte Reale, Sardinia (Wagner 1832 and 1833 sub *Turdus* sp., Giebel 1847).

Aves incertae sedis

Actiornis anglicus Lydekker

Actiornis anglicus Lydekker, 1891a: 56 [Holotype from Hordle: proximal end of right ulna; BMNH 30328. Figured by Lydekker 1891a, fig. 13, and Harrison & Walker 1976a, pl. 2, fig. k-n.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a, Harrison & Walker 1976a).

Remarks: This fossil was described in the Phalacrocoracidae (which included also the Sulidae at that time). Harrison & Walker (1976a) transferred it to the Threskiornithidae, but Olson (1981) observed that its holotype is neither from a cormorant, nor from an ibis. *Actiornis anglicus* Lydekker, 1891a is type by original designation of the genus *Actiornis* Lydekker, 1891a: 56.

Anas creccoides Beneden

Anas creccoides Beneden, 1871: 260 [Lectotype from Rupelmonde (selected by Brodkorb 1962: 707): a humerus lacking proximal end; IRScNB or ULg. Figured by Beneden 1871, pl. 1 (unnumbered), fig. 3. Paralectotypes from Rupelmonde: shaft of a humerus, proximal end of an ulna, distal end of an ulna, proximal end of a tibiotarsus; IRScNB or ULg. Figured by Beneden 1871, unnumbered pl., fig. 4 (shaft of humerus), 5 (proximal and distal ends of ulnae), and 6 (tibiotarsus). Brodkorb (1962: 707) identified from Beneden's figures from which body side the bones originated. However, drawings of bones were often, though not always, side-reversed in the 19th century publications, so that this information remains unknown. *Anas creccoides* is a junior primary homonym of *Anas creccoides* King, 1828.]

Anas creccoides Beneden: Lambrecht 1921: 15 [Spelling emended.]

Anas benedeni Sharpe 1899: 217 [New name for *Anas creccoides* Beneden, 1871; hence its junior objective synonym.]

Distribution: Middle Oligocene (MP 23-24) of Rupelmonde, Belgium (Beneden 1871).

Remarks: This species was described as a duck (Beneden 1871), but Brodkorb (1962: 707) relegated it to Aves incertae sedis.

Anas lignitifila Portis

Anas lignitifila Portis, 1884: 372 [Holotype from Bamboli: crushed partial skeleton in slab; MGPOT. Figured by Gastaldi 1866b, pl. 6, fig. 7.]

Anas lignitiphila Weithofer, 1888: 367. [Unjustified emendation of *Anas lignitifila* Portis, hence a new name, and a junior objective synonym of *Anas lignitifila* Portis, 1884.]

Distribution: Late Miocene (MN 13) of Bamboli, Italy (Gastaldi 1866a: 43 sub "un magnifico Ornitolite", T. Salvadori in Gastaldi 1866b as an unnamed bird, Portis 1884, Weithofer 1888, Ristori 1891, Howard 1964).

Remarks: This is a meaningless name until the holotype is restudied. Portis (1884: 371-372) attributed the species name to Salvadori, because he created the name on the basis of a description presented by the latter author (T. Salvadori in Gastaldi 1866b: 223-225). Nevertheless, the authorship of the name remains with Portis (1884).

Anas skalicensis Bayer

Anas skalicensis Bayer, 1882: 21 [Holotype from Skalice: scraps of long bones; NMP ČM-1520. Figured by Bayer 1882, fig. 8/2; Bayer 1883, fig. 2; Mlíkovský & Švec 1989, pl. 2, fig. a.]

Anas skalicensis Bayer, 1883: 64 [Same holotype, repeated description. This is both a junior homonym and a junior objective synonym of *Anas skalicensis* Bayer, 1882.]

Distribution: Middle Oligocene (MP 23-24) of Skalice, Czechia (Krejčí 1877: 915 sub "genus indet.", Bayer 1882, 1883, Mlíkovský & Švec 1989).

Remarks: The holotype of this species is indeterminate, even at the ordinal level (Mlíkovský & Švec 1989).

***Ardea rupeliensis* Beneden**

Ardea rupeliensis Beneden, 1873a: 373 [Holotype from Rupelmonde: distal end of a tibiotarsus; present location unknown, perhaps in IRScNB or ULg. – cf. Cheneval 1996a: 538). Not figured.]

Distribution: Middle Oligocene (MP 23-24) of Rupelmonde, Belgium (Beneden 1873a).

Remarks: Superficial description ("un peu petit que le butor" = slightly smaller than *Botaurus stellaris*) and the absence of illustration relegates this species to *Aves incertae sedis* until its holotype is located and restudied (see also Brodkorb 1978: 212).

***Ardea piveteaui* Brunet**

Ardea piveteaui Brunet, 1970: 16 [Holotype from Romainville: associated shaft of right ulna and distal end of right radius; MNHN 1951-1. Figured by Brunet 1970, pl. A, fig. d-e (ulna).]

Distribution: Late Eocene (MP 20) of Romainville, France (Brunet 1970, Harrison 1979a, Olson 1985a: 166).

Remarks: Indeterminate at the present state of our knowledge of the Eocene birds (see Harrison 1979a, Olson 1985a: 166). Nevertheless, the ulna is almost triangular in cross-section and bears distinct papillae remigiales caudales, which opens the possibility, that it will be identified in future (Mlíkovský orig.).

***Ardeacites molassicus* Haushalter**

Ardeacites molassicus Haushalter, 1855: 11 [Holotype from Harbatshofen: right humerus; present location unknown. Figured by Haushalter 1855, pl. 2, fig. 1.]

Distribution: Early Miocene (MN 3) of Harbatshofen, Germany (Haushalter 1855, Brodkorb 1980).

Remarks: The species was originally described as a heron. Brodkorb (1980: 91) relegated *Ardeacites molassicus* Haushalter to *Aves incertae sedis*, being "unable to get it in an order, much less a family." My attempt at identifying the holotype according to its figure in Haushalter (1855) gave the same result (Mlíkovský orig.). *Ardeacites molassicus* Haushalter, 1855 is type by monotypy of the genus *Ardeacites* Haushalter, 1855: 11.

***Argillipes aurorum* Harrison & Walker**

Argillipes aurorum Harrison & Walker, 1977a: 32 [Holotype from the Warden Point on the Isle of Sheppey: proximal end of right tarsometatarsus; BMNH A-3130. Figured by Harrison & Walker 1977, pl. 7, fig. p-t.]

Distribution: Early Eocene (MP 8-9) of Warden Point, Sheppey, England (Harrison & Walker 1977a).

Remarks: Originally described in the Phasianidae (Harrison & Walker 1977a). Here transferred to the *Aves incertae sedis*. *Argillipes aurorum* Harrison & Walker, 1977a is by original designation type of the genus *Argillipes* Harrison & Walker, 1977a: 32.

***Argillipes magnus* Harrison & Walker**

Argillipes magnus Harrison & Walker, 1979b: 37 [Holotype from Yarmouth: proximal end of left tarsometatarsus; BMNH A-4409. Figured by Harrison & Walker 1979b, pl. 2, row B: 3-7.]

Distribution: Middle Oligocene (MP 21-23) of Yarmouth, England (Harrison & Walker 1979b).

Remarks: Described in the Phasianidae. As judged from figures, the holotypical tarsometatarsus resembles the holotype of *Diatropornis ellioti* (Milne-Edwards) (Mlíkovský orig.).

***Argillipes paralectoris* Harrison & Walker**

Argillipes paralectoris Harrison & Walker, 1977a: 33 [Holotype from Sheppey: proximal end of left tarsometatarsus; BMNH A-3604. Figured by Harrison & Walker 1977a, pl. 8, fig. f-j.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Remarks: Described in the Phasianidae (Harrison & Walker 1977a). Here transferred to the Aves incertae sedis.

***Branta minor* Kessler & Gál**

Branta minor Kessler & Gál, 1996: 73 [Holotype from Chişinău: proximal end of right carpometacarpus; LPUI 65-MS. Figured by Kessler & Gál 1996, fig. 1a-b.]

Distribution: Late Miocene (MN 9) of Chişinău, Moldova (Kessler 1984b sub cf. *Phalacrocorax lautus*, Kessler & Gál 1996).

Remarks: The diagnosis is invalid, but the almost useless figures indicate, that this is not an anatid (Mlíkovský orig.).

***Cerestenia pulchrapenna* Mayr**

Cerestenia pulchrapenna Mayr, 2000i: 629 [Holotype from Céreste: nearly complete, articulated postcranial skeleton on a slab; SMNK PAL-3805. Figured by Mayr 2000i, fig. 4-6.]

Distribution: Middle Oligocene (MP 23) of Céreste, France (Mayr 2000i).

Remarks: Mayr (2000i) tentatively included this species in the family Turnipacidae Mayr, 2000i. *Cerestenia pulchrapenna* Mayr, 2000i is type by original designation of the genus *Cerestenia* Mayr, 2000i: 629.

***Chenornis graculoides* Portis**

Chenornis graculoides Portis, 1884: 364 [Holotype from Ceva: partial skeleton in slab; MGPU. Figured by Portis 1884, pl. 1, fig. 5.]

Distribution: Early Miocene (MN 1-2) of Ceva, Italy (Portis 1884).

Remarks: Portis (1884) was uncertain as regards the taxonomic position of this species. Lambrecht (1933: 367) listed it among the Anseriformes incertae sedis. Brodkorb (1964: 211) included it in the subfamily Anserinae of the Anatidae, but Howards (1964) doubted its anatid affinities. The specimen closely resembles cormorants in skeletal proportions and may be attributable to the Phalacrocoracidae (Mlíkovský, orig.). *Chenornis graculoides* Portis, 1884 is type by monotypy of the genus *Chenornis* Portis, 1884: 364.

***Colymbus portisi* Regàlia**

Colymbus Portisi Regàlia, 1902: 231 [Holotype from Orciano Pisano: incomplete cervical vertebra; probably lost – see Delle Cave et al. 1984: 89, Delle Cave 1996: 676. Figured by Regàlia 1902, pl. 27, fig. 19-20.]

Gavia portisi (Regàlia): Brodkorb, 1963: 224 [New combination.]

Distribution: Middle Pliocene (MN 15-16) of Orciano Pisano, Italy (Regàlia 1902, 1907, Delle Cave et al. 1984).

Remarks: This is a meaningless name, based upon a probably indeterminate vertebra. There is no evidence, that the vertebra is referable to a gaviid and, if so, that it represents a valid species.

***Coturnipes cooperi* Harrison & Walker**

Coturnipes cooperi Harrison & Walker, 1977a: 35 [Holotype from Burnham-on-Crouch: distal end of left tarsometatarsus; BMNH A-3706. Figured by Harrison & Walker 1977a, pl. 8, fig. p-t.]

Distribution: Early Eocene (MP 8-9) of Burnham-on-Crouch, England (Harrison & Walker 1977a). A tentative extralimital record is known from the early Eocene (Nanjemoy Formation) of Fisher/Sullivan Site in Virginia (Olson 1999b).

Remarks: Described in the Phasianidae (Harrison & Walker 1977a). Olson (1999b: 127) relegated this species to the category of *Aves incertae sedis*. *Coturnipes cooperi* Harrison & Walker, 1977a is type by original designation of genus *Coturnipes* Harrison & Walker 1977a: 35.

***Cygnus herenthalsi* Lambrecht**

Cygnus Herenthalsii Beneden, 1871: 261 [Nomen nudum; no description or indication.]

Cygnus Herenthalsii Beneden, 1872: 288 [Nomen nudum; no description or indication.]

Cygnus herenthalsii Beneden, 1873a: 372 [Nomen nudum; no description or indication.]

Cygnus herenthalsi [sic!] "Beneden" Lambrecht, 1933: 383 [Holotype from Antwerp: pedal phalanx; IRSeNB. Not figured. Lambrecht (1933: 383) attributed this name to Beneden (see also Stejneger 1882: 181, Lambrecht 1921: 10), giving title of the article as in Beneden 1872 and year and place of publication as in Beneden 1871. Beneden (1871, 1872, 1873a) mentioned this name on three occasions, but it is a pure nomen nudum in all cases. Lambrecht (1933: 383) provided a very brief description of the bird, which makes him the author of its name, although he attributed it to Beneden (see ICZN 1999). Note that Brodkorb (1964: 209) and Howard (1964: 260) just repeated incorrect data given by Lambrecht (1933: 383).]

Distribution: Middle Miocene (MN 7-8) of Antwerp, Belgium (Beneden 1871, 1872, 1873a, Lambrecht 1933).

Remarks: Beneden (1871, 1872, 1873a) believed that the pedal phalanx belongs to a swan, and Lambrecht (1933: 383) confirmed this opinion. Nevertheless, I concur with Howard (1964: 261) that a pedal phalanx is a dubious element upon which to base a new species and I relegate here *Cygnus herenthalsi* Lambrecht, 1933 to the category of *Aves incertae sedis* until its holotype is located and restudied.

***Dolichopterus viator* Milne-Edwards**

Dolicopterus viator Aymard, 1856: 234 [Nomen nudum; no description or indication – see Olson 1978.]

Dolichopterus [sic!] *viator* Milne-Edwards, 1868: 365 [Holotype from Ronzon: associated vertebra, proximal end of radius, carpometacarpus and three unidentified bone fragments in a slab; present location unknown. Figured by Milne-Edwards 1867-1868, pl. 57, fig. 24 (figure caption erroneously refers to fig. 23). Lydekker (1891a: 175) correctly observed that *Dolichopterus* was based solely on this material, but subsequent authors (Lambrecht 1933: 534, Brodkorb 1967: 191, Olson 1978: 448) thought that Milne-Edwards based the species on a syntypical series which included also a partial coracoid in slab (Milne-Edwards 1867-1868, pl. 57, fig. 25; figure caption erroneously refers to fig. 24) and crushed tarsometatarsus in still another slab (Milne-Edwards 1867-1868, pl. 57, fig. 23; figure caption erroneously refers to fig. 25). Referring to the partial skeleton in a slab (pl. 57, fig. 24), Milne-Edwards (1868: 366) clearly stated that this is the specimen upon which the

determination of *Dolichopterus viator* is based ("la pièce sur laquelle est basée cette détermination"). *Dolichopterus* is an incorrect original spelling as evidenced from the etymology and spelling in the caption to pl. 57 in Milne-Edwards 1867-1868, and confirmed by the decision of the first reviser (Lydekker 1891a: 175).]

Distribution: Early Oligocene (MP 21) of Ronzon, France (Aymard 1856, Milne-Edwards 1868-1869, Olson 1978).

Remarks: Aymard (1856) and Milne-Edwards (1968) tentatively included this fossil in their family "Longipennes", which included Wetmorean family Laridae and order Procellariiformes. Lambrecht (1933) and Brodkorb (1967) listed the species in the Charadriidae without explanation. I concur with Olson (1978: 446) that *Dolichopterus viator* Milne-Edwards should be relegated to the category of Aves incertae sedis until its type is located and restudied. *Dolichopterus viator* is type by monotypy of the genus *Dolichopterus* Milne-Edwards, 1868: 365. *Dolichopterus* Aymard, 1856: 234 is a nomen nudum.

***Eleutherornis helveticus* Schaub**

Eleutherornis helveticus Schaub, 1940a: 283 [Holotype from Egerkingen: anterior portion of pelvis; BaM Eh-781. Figured by Schaub 1940a, fig. 1-4.]

Distribution: Middle Eocene (MP 11-13) of Egerkingen, Switzerland (Schaub 1940a,b, Mlíkovský 1992: 444).

Remarks: This species is of uncertain taxonomic position (Mlíkovský 1992b). *Eleutherornis helveticus* Schaub, 1940a is type by original designation of the genus *Eleutherornis* Schaub, 1940a: 283. *Eleutherornis* Schaub, 1940a is type of the family Eleutherornithidae Wetmore, 1951: 3. Cracraft (1974) included this taxon in the Struthionidae as a tribe, Eleutherornithini.

Gaillard (1936) observed, that the pedal phalanx of a giant bird from Egerkingen, described and figured by Schaub (1929: 598, fig. 12-13) resembles the same element of *Diatryma cotei* Gaillard. This opens the possibility, that *Eleutherornis helveticus* Schaub is a phorusrhacid, but it should be compared also with *Gastornis* Hébert.

***Elornis anglicus* Lydekker**

Elornis (?) *anglicus* Lydekker, 1891a: 80 [Holotype from Hordle: left humerus; BMNH 36792. Figured by Lydekker 1891a, fig. 22, and Harrison & Walker 1976a, pl. 2, fig. e-j.]

Helornis anglicus (Lydekker): Lydekker, 1891d: 396 [Unjustified emendation of the genus name, hence a new combination.]

Distribution: Late Eocene (MP 17) of Hordle, England (Lydekker 1891a, Harrison & Walker 1976a).

Remarks: Being originally described in the Phoenicopteridae, this fossil was later transferred to the Threskiornithidae, and synonymized with *Actiornis anglicus* Lydekker (Harrison & Walker 1976a). According to Olson (1981: 168), the holotype was not "convincingly shown to belong in the Threskiornithidae". Accordingly, I relegate here *Elornis anglicus* Lydekker to the Aves incertae sedis until its holotype is properly restudied.

***Elornis grandis* Milne-Edwards**

Elornis grandis Aymard, 1856: 233 [Nomen nudum; no description or indication – see Olson 1978.]

Elornis grandis Milne-Edwards, 1869: 87 [Holotype from Ronzon: proximal end of a humerus; formerly in coll. Aymard, present location unknown. Not figured.]

Distribution: Early Oligocene (MP 21) of Ronzon, France (Aymard 1856, Milne-Edwards 1869, Olson 1978).

Remarks: This is clearly an Avis incertae sedis until its holotype is located and restudied.

***Elornis littoralis* Milne-Edwards**

Elornis (?) *littoralis* Aymard, 1856: 234 [Nomen nudum; no description or indication – see Olson 1978.]

Elorn[is] *antiquus* Aymard, 1856: 234 [Not available; published as a synonym and never used as a valid name – see Olson 1978.]

Elornis littoralis Milne-Edwards, 1868: pl. 90 [Holotype from Ronzon: associated distal end of tibiotarsus and proximal end of tarsometatarsus in a slab; formerly in coll. Aymard, present location unknown. Figured by Milne-Edwards 1967-1868, pl. 90, fig. 1. Several other skeletal elements from Ronzon, figured by Milne-Edwards 1867-1868, pl. 90, fig. 2-7, were included in the type series upon which Milne-Edwards allegedly based this species by Lambrecht (1933: 343), Brodkorb (1963c: 272) and Olson (1978: 447), but Milne-Edwards (1869: 86) attributed the species to Aymard and clearly stated that Aymard based it on the above mentioned slab. The latter specimen should be deemed to be the holotype of *Elornis littoralis* Milne-Edwards.]

Helornis littoralis ("Aymard" = Milne-Edwards): Lydekker 1891d: 396 [New combination, although Lydekker intended just to correct spelling of the generic name.]

Distribution: Early Oligocene (MP 21) of Ronzon, France (Aymard 1856, Milne-Edwards 1867-1868, 1869-1871, Olson 1978).

Remarks: *Elornis littoralis* Milne-Edwards was said to be flamingo-like by Aymard (1856), and included in the Phoenicopteridae by Lydekker (1891a: 80), Lambrecht (1933: 343), Brodkorb (1963c: 272) and Olson (1978: 445, tentatively). As judged from the figure, the distal end of the holotypical; tibiotarsus of *Elornis littoralis* Milne-Edwards markedly differs from the same element of the Phoenicopteridae. Hence, I relegate here this species in the category of Aves incertae sedis until its type is rediscovered and restudied.

Genus *Elornis* Aymard, 1856: 234 is a nomen nudum (see also Olson 1978). *Elornis littoralis* Milne-Edwards, 1868 is type by subsequent designation (Lydekker 1891a: 80) of the genus *Elornis* Milne-Edwards, 1868. *Helornis* Lydekker, 1891d: 396 is an unsubstantiated emendation of *Elornis* "Aymard" Milne-Edwards, 1868, and hence a new name. *Helornis* Lydekker is thus a junior objective synonym of *Elornis* Milne-Edwards.

***Eocathartes robustus* Lambrecht**

Eocathartes robustus Lambrecht, 1935: 362 [Holotype from Cecilie I: partial skeleton in slab; GM 6883. Figured by Lambrecht 1935, pl. 18, fig. 2.]

Distribution: Middle Eocene (MP 13) of Geiseltal – Cecilie I, Germany (Lambrecht 1935).

Remarks: Described as a vulture. This is almost certainly not a vulturid (P. Houde in Olson 1985a: 191). Its affinities await clarification. *Eocathartes robustus* Lambrecht, 1935 is type by monotypy of the genus *Eocathartes* Lambrecht, 1935: 362.

***Eostrix vincenti* Harrison**

Eostrix vincenti Harrison, 1980a: 85 [Holotype from Grange Farm: left phalanx I digiti III; BMNH A-5164. Figured by Harrison 1980a, fig. 1b,d,f, and 2c,d,f.]

Distribution: Early Eocene (MP 7-10) of Grange Farm, England (Harrison 1980a).

Remarks: Here relegated to the category of Aves incertae sedis (see also Houde & Olson 1992: 157). Type species of the genus *Eostrix* Brodkorb, 1971b: 214 is *Protostrix minima* Wetmore, 1938 from the early Eocene (Wasatch Formation) of Wyoming.

***Eupterornis remensis* Lemoine**

Eupterornis remensis Lemoine, 1878: 56 [Lectotype from Cernay-les-Reims (selected by Mourer-Chauviré 1994: 346): distal end of right ulna; MNHN CRL-2480. Figured by Lemoine 1878, pl. 5, fig. 1-3 (not pl. 3 as stated by Lemoine 1878: 56). Paralectotype from

Cernay-les-Reims: phalanx I digiti majori (= ?); MNHN, uncatalogued. Figured by Lemoine 1878, pl. 5, fig. 4-6. Mourer-Chauviré 1994 erroneously called the ulnar fragment a type of the species. Nevertheless, this can be interpreted as a selection of the syntype.]

Distribution: Late Paleocene (MP 6) of Cernay-les-Reims, France (Lemoine 1878, Olson 1985a: 213).

Remarks: This species was described as gull-like (Lemoine 1878). Paris (1912: 214) and Lambrecht (1921: 20) allied it with the Cretaceous Hesperornithidae. Lambrecht (1933: 261) placed it in his order Colymbo-Podicipediformes, and Brodkorb (1963c: 222) included it in the Gaviidae without explanation. Olson (1985a: 213) relegated *Eupterornis remensis* Lemoine to the Aves incertae sedis until its types are restudied. I restudied both syntypes of this species in MNHN and found them indeterminate. The lectotypical ulna is largely reconstructed, its figure in Lemoine being rather fictitious, while the "phalanx" is an indeterminate scrap of bone. *Eupterornis remensis* Lemoine, 1878 is type by monotypy of the genus *Eupterornis* Lemoine, 1878: 56.

***Fluviatilavis antunesi* Harrison**

Fluviatilavis antunesi Harrison, 1983a: 14 [Holotype from Silverinha: right femur; UNL, uncatalogued (C.J. Hazevoet, in litt.). Figured by Harrison 1983a, fig. 1a-e.]

Distribution: Early Eocene (MP 7) of Silveirinha, Portugal (Harrison 1983a).

Remarks: The figure in Harrison (1983a) does not show recurvirostrid features. I relegate this species to the category of Aves incertae sedis until its holotype is restudied. *Fluviatilavis antunesi* Harrison, 1983a is type by original designation of the genus *Fluviatilavis* Harrison, 1983a: 14.

***Fulica dejardinii* Beneden**

Fulica Dejardinii Bedenen, 1871: 261 [Holotype from Antwerp: distal end of left femur; perhaps in IRScNB or ULg. Figured by Beneden 1871, fig. 8.]

Miofulica Dejardini (Beneden): Lambrecht 1933: 480 [New combination.]

Distribution: Middle Miocene (MN 7-8) of Antwerp, Belgium (Beneden 1871).

Remarks: Described as a coot (family Rallidae), but should be relegated to Aves incertae sedis until its holotype is located and restudied (see also Cracraft 1973: 36, Olson 1977: 348). *Fulica dejardinii* Beneden, 1871 is type by monotypy of the genus *Miofulica* Lambrecht, 1933: 480.

***Gastornis minor* Lemoine**

Gastornis minor Lemoine, 1878: 12 [Holotype from Cernay-les-Reims: distal fragment of a tibiotarsus; present location unknown. Figured by Lemoine 1881c, pl. 4, fig. 1-5. Brodkorb (1967: 141) said that the figure in Lemoine shows a left tibiotarsus, but figures in Lemoine's papers were side-reversed, so that the holotypical tibiotarsus originated probably from the right side of the body.]

Remiornis minor (Lemoine): Lydekker 1891a: 360 [New combination.]

Distribution: Late Paleocene (MP 6) of Cernay-les-Reims, France (Lemoine 1878, 1881a,b,c, Martin 1992).

Remarks: Indeterminate from figures, hence a meaningless name until the holotype is found and restudied (Martin 1992).

***Geiseloceros robustus* Lambrecht**

Geiseloceros robustus Lambrecht, 1935: 365 [Holotype from Cecilie I: partial skeleton in a slab; GM 5884. Figured by Lambrecht 1935, pl. 2.]

Distribution: Middle Eocene (MP 13) of Geiseltal – Cecilie I, Germany (Lambrecht 1935).

Remarks: Described as a bucerotid, but certainly not belonging to the Bucerotidae. Taxonomic identity of *Geiseloceros robustus* Lambrecht remains unclear, although P. Houde (in Olson 1985a: 136, 191) conjectured that it may be identical with *Eocathartes robustus* Lambrecht, an alleged vulture from the same locality. *Geiseloceros robustus* Lambrecht, 1935 is type by monotypy of the genus *Geiseloceros* Lambrecht, 1935: 365.

***Gigantibis incognita* Harrison & Walker**

Gigantibis incognita Harrison & Walker, 1976a: 332 [Holotype from Hordle: proximal end of left femur; IGS GSM-1113109. Figured by Harrison & Walker 1976a, pl. 3, fig. c-g.]

Distribution: Late Eocene (MP 17) of Hordle, England (Harrison & Walker 1976a).

Remarks: This fossil was described in the Threskiornithidae, but Olson (1981) relegated it to the category of Aves incertae sedis. *Gigantibis incognita* Harrison & Walker, 1976a is type by original designation of the genus *Gigantibis* Harrison & Walker, 1976a: 331.

***Gracilitarsus mirabilis* Mayr**

Gracilitarsus mirabilis Mayr, 1998c: 56 [Holotype from Messel: incomplete skeleton in a slab; SMNK Me-1085. Figured by Mayr 1998c, pl. 17; Mayr 2001c, fig. 1.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998c, Mayr 2001c).

Remarks: Described in order and family incertae sedis (Mayr 1998c), and later placed in its own family (Mayr 2001d). *Gracilitarsus mirabilis* Mayr, 1998c is type by original designation of the genus *Gracilitarsus* Mayr, 1998c: 55. *Gracilitarsus* Mayr, 1998c is type of the family Gracilitarsidae Mayr, 2001c: 78. Mayr (2001c) indicated that this species can be closely related to *Eutreptodactylus itaboraiensis* Baird & Vickers-Rich, 1997, an alleged cuckoo from the late Paleocene of Brazil (but see Mourer-Chauviré 1999b).

***Hassiavis laticauda* Mayr**

Hassiavis laticauda Mayr, 1998a: 122 [Holotype from Messel: partial skeleton in slab and counter-slab; HLMD Me-9047a,b. Figured by Mayr 1998a, fig. 2-3.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998a).

Remarks: Mayr (1998a) tentatively assigned this species to the Archaeotrogonidae, from which it markedly differs in osteological characters described by Mayr (1998a) and/or those visible in the figures. Until restudied, this species should be relegated to the Aves incertae sedis. *Hassiavis laticauda* Mayr, 1998a is type by original designation of the genus *Hassiavis* Mayr, 1998a: 122.

***Homalopus picoides* Milne-Edwards**

Homalopus picoides Milne-Edwards, 1871: 385 [Holotype from Sansan: distal end of left tarsometatarsus; MNHN Sa-1240. Figured by Milne-Edwards 1869-1871, pl. 178, fig. 15-20; and Cheneval 2000, fig. 27. Cheneval (2000) incorrectly called this specimen a lectotype, without mentioning any paralectotypes.]

Distribution: Middle Miocene (MN 6) of Sansan, France (Milne-Edwards 1869-1871, Cheneval 2000).

Remarks: Milne-Edwards (1869-1871) was not certain about the taxonomic position of this bird. Paris (1912: 285) and Lambrecht (1921: 89, 1933: 631) assigned it to the piciform birds. Brodkorb (1971: 249) placed it in the Bucerotidae, and Cheneval (1996b, 2000: 372) in the Dendrocolaptidae. The latter assignment was doubted on paleozoogeographical grounds by Mlíkovský (1996s: 812). *Homalopus picoides* Milne-Edwards, 1871 is type by original designation of the genus *Homalopus* Milne-Edwards, 1871: 385.

***Horusornis vianeyliaudae* Mourer-Chauviré**

Horusornis vianeyliaudae Mourer-Chauviré, 1991: 190 [Holotype from Bouffie: left tarsometatarsus with damaged proximal end; USTL BFI-1974. Figured by Mourer-Chauviré 1991, pl. 1, fig. 8-9.]

Distribution: Late Eocene (MP 17) of Bouffie, France (Mourer-Chauviré 1991).

Remarks: Described as a raptor of unknown relationships. Until restudied, this species should be relegated to the Aves incertae sedis. *Horusornis vianeyliaudae* Mourer-Chauviré, 1991 is type by original designation of the genus *Horusornis* Mourer-Chauviré, 1991: 190. *Horusornis* Mourer-Chauviré, 1991 is type of the family Horusornithidae Mourer-Chauviré, 1991: 184.

***Howardia eous* Harrison & Walker**

Howardia eous Harrison & Walker, 1976a: 337 [Holotype from Hordle: anterior portion of sternum; BMNH A-4355. Figured by Harrison & Walker 1976a, pl. 4, fig. f.]

Palaeopapia eous (Harrison & Walker): Harrison & Walker 1979c: 110 [New combination.]

Distribution: Late Eocene (MP 17) of Hordle, England (Harrison & Walker 1976a); and early Oligocene (MP 21-23) of Hamstead, England (Harrison & Walker 1979b, tentatively referred specimen).

Remarks: Described in the Anatidae. *Howardia eous* Harrison & Walker, 1976a is type by original designation of the genus *Howardia* Harrison & Walker, 1976a: 337 (a junior homonym of *Howardia* Berlese & Leonardi, 1896). *Palaeopapia* Harrison & Walker, 1979c: 110 is a new name for *Howardia* Harrison & Walker, 1976a, hence its junior objective synonym.

***Ibidopodia palustris* Milne-Edwards**

Ibidopodia palustris Milne-Edwards, 1871: 465 [Holotype from Saint-Gérard-le-Puy: left tarsometatarsus; MNHN, lost? – I was not able to locate it in MNHN in 1999. Figured by Milne-Edwards 1869-1871, pl. 71, fig. 17-21. Milne-Edwards (1869-1871) clearly based this species on the tarsometatarsus, attributing to it tentatively the partial skull MNHN Av-8730 (p. 467: "J'incline à penser qu'un tête d'oiseau ... pourrait bien appartenir à *Ibidopodia palustris* ou à quelque genre voisin, ..." = I am inclined to think that a head of a bird ... could well belong to *Ibidopodia palustris* or to an allied genus, ...) (my translation). The latter skull is figured by Milne-Edwards 1869-1871, pl. 71, fig. 13-16, and Cheneval & Escullié 1992, fig. 5. Brodkorb (1963c: 278) erroneously stated that Lydekker (1891: 74) selected the latter skull as the lectotype of *Ibidopodia palustris*, while the latter author clearly regarded the tarsometatarsus as the holotype of the species. Olson (1981a: 168) erroneously stated that Milne-Edwards based the species on both of these elements.]

Distribution: Early Miocene (MN 2a) of Saint-Gérard-le-Puy, France (Milne-Edwards 1869-1871, Olson 1981).

Remarks: Milne-Edwards (1869-1871) described this species as an ibis, but Olson (1981a: 168) excluded it from the family, relegating it to the category of Aves incertae sedis. The skull mentioned by Milne-Edwards (1869-1871: 467) – and sometimes erroneously regarded as a part of the alleged syntypical series – belongs to *Palaelodus ambiguus* Milne-Edwards (Cheneval & Escullié 1992). *Ibidopodia palustris* Milne-Edwards, 1871 is type by monotypy of the genus *Ibidopodia* Milne-Edwards, 1871: 465.

***Kievornis rogovitshi* Aver'janov et al.**

Kievornis rogovitshi Aver'janov, Potapova & Nesov, 1990: 6 [Holotype from Kyjiv: associated right humerus lacking proximal end, and right ulna; ZIN 3926a-b. The humerus (ZIN 3926a) was called "main specimen of the holotype" and the ulna "additional specimen of the holotype" by Aver'janov et al. in captions to pl. 1-2. Figured by Rogovič 1875a, pl. 2, figs.

3-4; and Aver'janov et al. 1990, pl. 1, fig. 1-5, pl. 2, fig. 1-5.]

Distribution: Late Eocene (MP 17-20) of Kyjiv, Ukraine (Rogovič 1875a: 82 sub *Scolopax* sp., Aver'janov et al. 1990).

Remarks: Aver'janov et al. (1990) described this genus in the family Graculavidae, created by Fürbringer (1888) for the genus *Graculavus* Marsh, 1872 from the late Cretaceous or early Paleocene of New Jersey (see Olson & Parris 1987, Olson 1994). *Kievornis rogovitshi* Aver'janov, Potapova & Nesov, 1990 is type by original designation of the genus *Kievornis* Aver'janov, Potapova & Nesov, 1990: 5.

***Laputa robusta* Dyke**

Laputa robusta Dyke, 2001b.

Laputavis robusta (Dyke): Dyke 2001c: 214 [New combination.]

Distribution: Early Eocene of "London Clay", England (Dyke 2001b,c, Mayr 2001d).

Remarks: Not seen. Described in the Apodiformes. *Laputa* Dyke, 2001b is type by original designation of the genus *Laputa* Dyke, 2001b (preoccupied). *Laputavis* Dyke, 2001c: 214 is new name for *Laputa* Dyke, 2001b.

***Larus toliapicus* König**

Larus toliapicus König, 1825: pl. 16 [Holotype from Sheppey: imperfect cranium; BMNH A-130. Figured by König 1825, pl. 16, fig. 193, Owen 1846, fig. 234a-b, 235, Harrison & Walker 1972, text-fig. 1-5, 6d, 7i, 9a, pl. 1a-c, 2a-c, 3e, Harrison & Walker 1977, pl. 11, fig. a-d.]

Halcyornis toliapicus (König): Owen 1846: 554 [New combination.]

Distribution: Early Eocene (MP 8) of Sheppey, England (König 1825, Owen 1846, Lydekker 1891a, 1896, Harrison & Walker 1972, 1977).

Remarks: This species was described as a gull (König 1825), and later included in the Laridae (Lydekker 1891a, 1896, Lambrecht 1933, Brodkorb 1967: 205), or considered to be allied with the Alcedinidae (Owen 1846, Harrison & Walker 1972, 1977). *Halcyornis toliapicus* Owen, 1846 is type by monotypy of the genus *Halcyornis* Owen, 1846: 554. The latter genus is type of the family Halcyornithidae Harrison & Walker, 1972: 168.

***Litoripes medius* Harrison & Walker**

Litoripes medius Harrison & Walker, 1979a: 22 [Holotype from Yateley: right tarsometatarsus; BMNH A-5014. Figured by Harrison & Walker 1979, pl. 1, row 4 partim.]

Distribution: Middle Eocene (MP 11-13) of Yateley, England (Harrison & Walker 1979a).

Remarks: Described in the Phasianidae. Here transferred to the Aves incertae sedis. *Litoripes medius* Harrison & Walker, 1979a is type by original designation of the genus *Litoripes* Harrison & Walker, 1979a: 22.

***Marinavis longirostris* Harrison & Walker**

Marinavis longirostris Harrison & Walker, 1977a: 6 [Holotype from Abbey Wood: fragmentary right dentary and associated posterior portion of left edge of rostrum; BMNH A-4267. Figured by Harrison & Walker 1977a, fig. 1a-b.]

Distribution: Early Eocene (MP 8-9) of Abbey Wood, England (Harrison & Walker 1977a).

Remarks: This species was based on such a bone splinter, that its taxonomic validity, as well as the taxonomic validity of the genus and the family, remain uncertain. *Marinavis longirostris* Harrison & Walker, 1977a is type by original designation of the genus *Marinavis* Harrison & Walker, 1977a: 6. *Marinavis* Harrison & Walker, 1977a is type of the family Marinavidae Harrison & Walker, 1977a: 6.

***Mergus ronzoni* Gervais**

Mergus Ronzoni Gervais, 1849: 220 [Nomen nudum; no description or indication.]

Mergus Ronzoni Gervais, 1848-1852: 232 [Holotype from Ronzon: imperfect pelvis in slab; MNHN 1909-40. Figured by Milne-Edwards, 1867-1868, pl. 44, fig. 9.]

Sula ronzoni (Gervais): Milne-Edwards 1867: 271 [New combination.]

Prophalacrocorax ronzoni (Gervais): Harrison 1975a: 52 [New combination.]

Distribution: Early Oligocene (MP 21) of Ronzon, France (Gervais 1844a,b sub "Harle", Gervais 1849, 1848-1852, Milne-Edwards 1867-1868, Harrison 1975a).

Remarks: Brodkorb (1963c: 257) and Harrison (1975a: 52) attributed the species name to Milne-Edwards (1867), overlooking that Gervais (1848-1852: 232-233) provided a very brief, but valid description for this species. Harrison (1975a) transferred this species to the Phalacrocoracidae, but Olson (1985a) returned it to the Sulidae until its type is restudied. I briefly studied the holotype in MNHN, concluding that it is indeterminate at the genus, and perhaps even at the family level. *Mergus ronzoni* Gervais, 1848-1852 is type by original designation of the genus *Prophalacrocorax* Harrison, 1975a: 52.

***Messelornis russelli* Mourer-Chauviré**

Messelornis russelli Mourer-Chauviré, 1995a: 96 [Holotype from Cernay-les-Reims: distal end of left humerus; MNHN BR-14033. Figured by Mourer-Chauviré 1995a, text-fig. 1 (left), pl. 1, fig. 2-3.]

Distribution: Late Paleocene (MP 6) of Cernay-les-Reims, France (Mourer-Chauviré 1995a).

Remarks: The holotypical humerus of this species is very different from the same element of the Messelornithidae. Until restudied, I relegate here *Messelornis russelli* Mourer-Chauviré to the category of Aves incertae sedis.

***Microena goodwini* Harrison & Walker**

Microena goodwini Harrison & Walker 1977a: 40 [Holotype from Bognor Regis: left tarsometatarsus; BMNH A-3685. Figured by Harrison & Walker 1977a, pl. 9, fig. n-r.]

Distribution: Early Eocene (MP 8-9) of Bognor Regis, England (Harrison & Walker 1977a).

Remarks: Described in the Columbidae (Harrison & Walker 1977a). Here relegated to the Aves incertae sedis. *Microena goodwini* Harrison & Walker, 1979a is type by original designation of the genus *Microena* Harrison & Walker, 1977a: 39.

***Milvoides kempi* Harrison & Walker**

Milvoides kempi Harrison & Walker, 1979a: 20 [Holotype from Lee-on-Solent: distal end of right tarsometatarsus lacking trochlea for digit II; BMNH A-5005. Figured by Harrison & Walker 1979a, pl. 1, row 1.]

Distribution: Middle Eocene (MP 11-13) of Lee-on-Solent, England (Harrison & Walker 1979a).

Remarks: The holotype of this species is too fragmentary for a certain identification of an Eocene bird. *Milvoides kempi* Harrison & Walker, 1979a is type by original designation of the genus *Milvoides* Harrison & Walker, 1979a: 20.

***Oligocathartes olsoni* Harrison & Walker**

Oligocathartes olsoni Harrison & Walker, 1979b: 36 [Holotype from Yarmouth: incomplete distal end of left tarsometatarsus; BMNH A-4985. Figured by Harrison & Walker 1979b, pl. 2, row A: 6-8, row B: 1-2.]

Distribution: Early/late Oligocene (MP 21-23) of Yarmouth, Isle of Wight, England (Harrison

& Walker 1979b).

Remarks: Described in the Cathartidae, but the holotypical tarsometatarsus seems to be too fragmentary for proper identification (Mlíkovský, orig.). *Oligocathartes olsoni* Harrison & Walker, 1979b is type by original designation of the genus *Oligocathartes* Harrison & Walker, 1979b: 36.

***Ornitholithes bosniaskii* Portis**

Ornitholithes Bosniaskii Portis, 1887: 189 [Syntypes from Ancona: two feather imprints; present location unknown (Delle Cave 1996: 670). Figured by Portis 1887, fig. 9-10.]

Distribution: Late Miocene (MN 11-13) of Ancona, Italy (Portis 1887).

***Ornitholithes faujasi* Omboni**

Ornitholithes Faujasi Omboni, 1885: 772 [Syntypes from Vestena Nova: three feather imprints; MPUP. Figured by Omboni 1885, fig. 1-3.]

Distribution: Late Eocene (MP 17-20) of Vestena Nova, Italy (Omboni 1885).

***Ornitholithes gabbrensis* Portis**

Ornitholithes gabbrensis Portis, 1887: 192 [Syntypes from Gabbro: five feather imprints; MSNUP. Figured by Portis 1887, fig. 13-17.]

Distribution: Late Miocene (MN 13) of Gabbro, Italy (Portis 1887).

***Ornitholithes procaccinii* Portis**

Ornitholithes Procaccinii Portis, 1887: 188 [Syntypes from Senigallia: four feather imprints; MCI. Figured by Portis 1887, fig. 5-8.]

Distribution: Late Miocene (MN 13) of Senigallia, Italy (Procaccini Ricci 1840, Portis 1887).

***Ornitholithes tenuipennis* Omboni**

Ornitholithes tenuipennis Omboni, 1885: 772 [Syntypes from Vestena Nova: three feather imprints; MPUP. Figured by Omboni 1885, fig. 5-7.]

Distribution: Late Eocene (MP 17-20) of Vestena Nova, Italy (Omboni 1885).

***Ornitholithus arcuatus* Dughi & Sirugue**

Ornitholithus arcuatus Dughi & Sirugue, 1962: 75 [Syntypes from Basse Provence: eggshell fragments; MHNA, uncatalogued. Figured by Dughi & Sirugue 1962, pl. 3, fig. 6.]

Ornitholithus mammosus Dughi & Sirugue, 1962: 75 [Syntypes Basse Provence: eggshell fragments from; MHNA, uncatalogued. Figured by Dughi & Sirugue 1962, pl. 3, fig. 3-4.]

Ornitholithus mammillatus Dughi & Sirugue, 1962: 75 [Syntypes from Basse Provence: eggshell fragments; MHNA, uncatalogued. Figured by Dughi & Sirugue 1962, pl. 3, fig. 5.]

Ornitholithus mammeatus Dughi & Sirugue, 1962: 75 [Syntypes from Basse Provence: eggshell fragments; MHNA, uncatalogued. Figured by Dughi & Sirugue 1962, pl. 3, fig. 7.]

Distribution: Early Eocene (MP 7) of Basse Provence, France (Dughi & Sirugue 1959, 1962, 1968, Fabre-Taxy & Touraine 1960, Touraine 1960, 1961, 1969, Rey 1969, Cailleux 1969, Kerourio & Aujard 1987, Dauphin 1991, 1992, 1994); and early/middle Eocene (MP 7-10) of Aude, France (Plaziat 1964, Villatte 1966, Cailleux 1969, Dughi et al. 1969, Kerourio & Aujard 1987).

Remarks: Dughi & Sirugue (1962) understood *arcuatus*, *mammosus*, *mammillatus* and *mammeatus* as a group of species, and they used *arcuatus* in the broader sense elsewhere (Dughi & Sirugue 1968, Dughi et al. 1969, see also Villatte 1966). Accordingly, I list here *mammosus*, *mammillatus* and *mammeatus* in the synonymy of *Ornitholithus arcuatus* Dughi &

Sirugue.

***Ornitholithus biroi* Dughi & Sirugue**

Ornitholithus biroi Dughi & Sirugue, 1962: 74 [Syntypes from Basse Provence: eggshell fragments; MHNA, uncatalogued. Figured by Dughi & Sirugue 1962, pl. 3, fig. 2.]

Distribution: Early Eocene (MN 7) of Basse Provence, France (Dughi & Sirugue 1959, 1962, 1968, Kerourio & Aujard 1987).

Remarks: *Ornitholithus biroi* Dughi & Sirugue, 1962 is type by subsequent designation (Vjalov 1971: 50) of the genus *Ornitholithus* Dughi & Sirugue, 1962: 74.

***Palaeoaramides minutus* Cracraft**

Palaeoaramides minutus Cracraft, 1973: 30 [Holotype from Grive-Saint-Alban: right tarsometatarsus; BMNH A-332. Figured by Cracraft 1973, fig. 14a-b.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Cracraft 1973).

Remarks: The holotypical tarsometatarsus of this species differs in the position and shape of trochlea digiti II (as seen in distal view) from the same element of the Rallidae (Mlíkovský, orig.). Accordingly, I relegate here *Palaeoaramides minutus* Cracraft to the Aves incertae sedis until its holotype is restudied.

***Palaeopsittacus georgei* Harrison**

Palaeopsittacus georgei Harrison, 1982c: 205 [Holotype from Walton-on-the-Naze: 11 associated bones, incl. left scapula lacking the tip, right coracoid, thoracic vertebra, proximal end of right ulna, distal end of right ulna, proximal end of left radius, right cuneiforme, distal end of left carpometacarpus, distal end of left femur, distal end of right tibiotarsus, and proximal end of right tarsometatarsus; BMNH A-5163. Figured by Harrison 1982c, text-fig. 2a (ulna), 3a (scapula), 4c,f (coracoid), 5b (tibiotarsus), and 6a (tarsometatarsus), and pl. 10, fig. a-c (tibiotarsus), h-j (femur), l-m (tarsometatarsus), n-p (proximal end of ulna), q-s (coracoid), t-u (distal end of ulna), v-x (carpometacarpus), y (radius), z (scapula), aa-bb (vertebra).]

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Harrison 1982c, Mayr & Daniels 1998); and middle Eocene (MP 11-13) of Lee-on-Solent, England (Harrison 1982c; specimen tentatively referred).

Remarks: This species was described in the Psittacidae (Harrison 1982c), and later transferred in the Quercypsittidae (Mourer-Chauviré 1992c). Mayr & Daniels (1998) observed that the bird has an anisodactyl foot and relegated it to the category of Aves incertae sedis. The species was called *Poicephalus georgei* in the caption to Fig. 6a by Harrison (1982c: 205). This is certainly a misprint, not a new combination. *Palaeopsittacus georgei* Harrison, 1982a is type by original designation of the genus *Palaeopsittacus* Harrison, 1982a: 204.

***Pararallus hassenkampii* Martini**

Pararallus hassenkampii Martini, 1967a: 8 [Nomen nudum; no description or indication – Mlíkovský 1992b: 441.]

Pararallus hassenkampii Martini, 1967b: 289 [Holotype from Sieblos: associated pedal phalanges with crushed distal fragment of left tarsometatarsus in a slab and a counter-slab; GPIW F-1237 (slab) and F-1238 (counter-slab). Figured by Martini 1967b, text-fig. 1 (reconstruction), pl. 29, fig. 1 (slab), and 2 (counter-slab), and Martini 1988, fig. 1 (counter-slab).]

Distribution: Early/middle Oligocene (MP 21-24) of Sieblos, Germany (Martini 1967a,b, 1988, see also Hassenkamp 1858: 207, Meyer 1858: 111).

Remarks: Probably indeterminate (see also Cracraft 1973: 38).

***Parvicuculus minor* Harrison & Walker**

Parvicuculus minor Harrison & Walker, 1977a: 44 [Holotype from Burnham-on-Crouch: left tarsometatarsus; BMNH A-4919. Figured by Harrison & Walker 1977, pl. 10, fig. a-f, Harrison 1982a, fig. 1a, 2a-e.]

Distribution: Early Eocene (MP 8-9) of Burnham-on-Crouch, England (Harrison & Walker 1977a, Olson & Feduccia 1979, Harrison 1982a).

Remarks: Described in the family Cuculidae. Olson & Feduccia (1979) removed the species to the Primobucconidae, conjecturing that *Parvicuculus* Harrison & Walker "may prove to be a synonym of *Neanis* Brodkorb", but Harrison (1982a: 73) returned it back to cuckoos, placing it together with *Proculcus* Harrison & Walker in his new family Parvicuculidae. Martin (in Martin & Mengel 1984) restudied the holotype of *Parvicuculus minor* Harrison & Walker, concluding that the bird was neither a cuckoo, nor a primobucconid, but that its "assignment to the Coraciiformes would not be unlikely." Baird & Vickers-Rich (1997) restudied a cast of the holotype, and excluded the fossil from both the Cuculidae and Primobucconidae, leaving it in the Aves incertae sedis. *Parvicuculus minor* Harrison & Walker, 1977a is type by original designation of the genus *Parvicuculus* Harrison & Walker, 1977a: 43. *Parvicuculus* Harrison & Walker, 1982a is type of the family Parvicuculidae Harrison, 1982a: 75.

***Parvigyps praecox* Harrison & Walker**

Parvigyps praecox Harrison & Walker, 1977a: 31 [Holotype from Sheppey: distal end of right humerus; BMNH A-71. Figured by Harrison & Walker 1977a, pl. 7, fig. d-h.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Remarks: Described in the Accipitridae. Here relegated to the category of Aves incertae sedis. *Parvigyps praecox* Harrison & Walker, 1977a is type by original designation of the genus *Parvigyps* Harrison & Walker, 1977a: 31.

***Parvulivenator watteli* Harrison**

Parvulivenator watteli Harrison, 1982b: 79 [Holotype from Walton-on-the-Naze: distal end of right tarsometatarsus with associated os metatarsale I, and four pedal phalanges; BMNH A-5172. Figured by Harrison 1982b, fig. 1a-j, 2a-l.]

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Harrison 1982b).

Remarks: Described in the Falconidae. Here transferred to the Aves incertae sedis. *Parvulivenator watteli* Harrison, 1982b is type by original designation of the genus *Parvulivenator* Harrison, 1982b: 79.

***Pelargosteon tothi* Kretzoi**

Pelargosteon tothi Kretzoi, 1962: 169 [Holotype from Betfia 5: incomplete crushed sternum; RMO 1899/1. Figured by Jurcsák & Kessler 1973, fig. 24.]

Pelargosteon tothi Kretzoi: Brodkorb 1963c: 290 [Spelling emended.]

Distribution: Early Pleistocene (MQ 1a) of Köröshegy, Hungary (Kretzoi & Krolopp 1977), and Betfia 2, Romania (Jánossy 1980a); early Pleistocene (MQ 1b) of Voigtstedt, Germany (Jánossy 1965 sub "Ciconiidae indet.", reidentified by Jánossy 1980a: 20), and Betfia 5, Romania (Kretzoi 1962, Jurcsák & Kessler 1973); and middle Pleistocene (MQ 2A) of Várhegy – Fortuna Street 25, Hungary (Jánossy 1980a, Jánossy 1986: 87 sub cf. *Pelargosteon tothi*).

Remarks: Described in the Ciconiidae. The figure of the holotype is too indistinct to allow for a taxonomic assessment of the species. *Pelargosteon tothi* Kretzoi, 1962 is type by monotypy

of the genus *Pelargosteon* Kretzoi, 1962: 169.

***Petropluvialis simplex* Harrison & Walker**

Petropluvialis simplex Harrison & Walker, 1976a: 348 [Holotype from Hordle: cranial end of right coracoid; BMNH 30327. Figured by Harrison & Walker 1976a, pl. 7, fig. f-i.]

Distribution: Late Eocene (MP 17) of Hordle, England (Harrison & Walker 1976a).

Remarks: Described in the Burhinidae. Here relegated to the category of Aves incertae sedis. *Petropluvialis simplex* Harrison & Walker, 1976a is type by original designation of the genus *Petropluvialis* Harrison & Walker, 1976a: 348.

***Picus gaudryi* Depéret**

Picus gaudryi Depéret, 1887: 282 [Holotype from Grive-Saint-Alban: right femur; ML LGr-56. Figured by Depéret 1887, pl. 13, fig. 53, 53a.]

Zygodactylus gaudryi (Depéret): Brodkorb 1971: 258 [New combination.]

Distribution: Middle Miocene (MN 7-8) of Grive-Saint-Alban, France (Depéret 1887).

Remarks: The taxonomic identity of this species is unclear (cf. Brodkorb 1971a,b).

***Piscator tenuirostris* Harrison & Walker**

Piscator tenuirostris Harrison & Walker, 1976a: 328 [Holotype from Hordle: anterior end of premaxilla; BMNH A-146. Figured by Harrison & Walker 1976a, pl. 2, fig. a-d.]

Distribution: Late Eocene (MP 17) of Hordle, England (Harrison & Walker 1976a).

Remarks: Described in the Phalacrocoracidae. Here relegated to the category of Aves incertae sedis. *Piscator tenuirostris* Harrison & Walker, 1976a is type by original designation of the genus *Piscator* Harrison & Walker, 1976a: 328.

***Plesiocathartes europaeus* Gaillard**

Plesiocathartes europaeus Gaillard, 1908: 41 [Holotype from Quercy: distal end of left tarsometatarsus; MNHN QU-1058. Figured by Gaillard 1908, text-fig. 6, pl. 2, fig. 13-16, Cracraft & Rich 1972, fig. 4a-b.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1908, Cracraft & Rich 1972).

Remarks: Jollie (1977c: 112) observed, that "this fossil ... in details shows nothing to link it conclusively with [the Cathartidae]." *Plesiocathartes europaeus* Gaillard, 1908 is type by monotypy of the genus *Plesiocathartes* Gaillard, 1908: 41.

***Plesiocathartes gaillardi* Crusafont & Villalta**

Plesiocathartes (?) *gaillardi* Crusafont Pairó & Villalta, 1955: 236 [Holotype from Fallol: distal end of left tarsometatarsus; formerly in IPS (Crusafont & Villalta 1955: 237), present location unknown (Sánchez Marco 1996: 722). Figured by Crusafont Pairó & Villalta 1955, fig. 55.]

Distribution: Early Miocene (MN 3) of Fallol, Spain (Crusafont Pairó & Villalta 1955, Villalta 1963).

Remarks: Described in the Cathartidae, but important differences from vulturids are mentioned in the description, and the drawing of the holotype is very generalized (Mlíkovský, orig.). Accordingly, I relegate here *Plesiocathartes gaillardi* Crusafont Pairó & Villalta to the Aves incertae sedis until its holotype is located and restudied.

***Pliogrus germanicus* Lambrecht**

Pliogrus germanicus Lambrecht, 1933: 522 [Holotype from Eppelsheim: distal end of left

tibiotarsus; ZGIB. Figured by Lambrecht 1933, fig. 156a-b.]

Distribution: Late Miocene (MN 9) of Eppelsheim, Germany (Lambrecht 1933).

Remarks: Described in the Gruidae. Fischer & Stephan (1971) and Mourer-Chauviré et al. (1985) independently excluded this species from the Gruidae, and Fischer & Stephan (1971) transferred it to the Phoenicopteridae. Figures of the holotype of this species are rather indistinct, but the specimen appears to be neither flamingo-like, nor crane-like in its general shape (Mlíkovský, orig.). *Pliogrus germanicus* Lambrecht thus should be relegated to the category of Aves incertae sedis until its holotype is restudied. *Pliogrus germanicus* Lambrecht, 1933 is type by monotypy of the genus *Pliogrus* Lambrecht, 1933: 522.

***Precursor litorum* Harrison & Walker**

Precursor litorum Harrison & Walker, 1977a: 26 [Holotype from Sheppey: distal end of right humerus; BMNH A-3135. Figured by Harrison & Walker 1977a, pl. 6, fig. k-o.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Remarks: Described in the Glareolidae (tentatively), to which the species bears little resemblance (Mlíkovský orig.).

***Precursor magnus* Harrison & Walker**

Precursor magnus Harrison & Walker, 1977a: 26 [Holotype from Sheppey: distal end of right tarsometatarsus; BMNH A-3683. Figured by Harrison & Walker 1977a, pl. 6, fig. f-j.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1977a).

Remarks: Described in the Glareolidae (tentatively), to which it bears little resemblance (Mlíkovský, orig.).

***Precursor parvus* Harrison & Walker**

Precursor parvus Harrison & Walker, 1977a: 25 [Holotype from Bognor Regis: distal end of left humerus; BMNH A-3684. Figured by Harrison & Walker 1977a, pl. 6, fig. a-e.]

Distribution: Early Eocene (8-9) of Bognor Regis, England (Harrison & Walker 1977a), and Warden Point, Sheppey, England (Harrison & Walker 1977).

Remarks: Described in the Glareolidae (tentatively), to which it bears little resemblance (Mlíkovský, orig.). *Precursor parvus* Harrison & Walker, 1977a is by original designation type of genus *Precursor* Harrison & Walker, 1977: 24.

***Proceriavis martini* Harrison & Walker**

Proceriavis martini Harrison & Walker, 1979b: 30 [Holotype from Yarmouth: imperfect cervical vertebra; BMNH A-4413. Figured by Harrison & Walker 1979b, pl. 1, row A.]

Distribution: Early/middle Oligocene (MP 21-23) of Yarmouth, England (Harrison & Walker 1979b).

Remarks: This species was based on a fragmentary cervical vertebra, unsuitable for exact identification. Harrison & Walker (1979b) tentatively included the species in the Eleutherornithidae. *Proceriavis martini* Harrison & Walker, 1979b is type by original designation of the genus *Proceriavis* Harrison & Walker, 1979b: 30.

***Procuculus minutus* Harrison & Walker**

Procuculus minutus Harrison & Walker, 1977a: 45 [Holotype from Bognor Regis: distal end of right (not left as stated by Harrison & Walker 1977a: 45 – corrected by Harrison 1982a: 73) tarsometatarsus; BMNH A-4680. Figured by Harrison & Walker 1977a, pl. 10, fig. g-k, Harrison 1982a, fig. 3a-e.]

Distribution: Early Eocene (MP 8-9) of Bognor Regis, England (Harrison & Walker 1977a,

Olson & Feduccia 1979, Harrison 1982a).

Remarks: Described as a cuckoo, but subsequently removed from the Cuculidae by Olson & Feduccia (1979). Harrison (1982a: 73) returned the species to the cuckoos, placing it in the separate family Parvicuculidae. Olson (1985a: 135) indicated that *Procuculus minutus* Harrison & Walker may belong in *Primapus* Harrison & Walker from the family Apodidae. Baird & Vickers-Rich (1997: 123) restudied a cast of the holotype, concluding that the species belongs neither to the Cuculidae, nor to the Primobucconidae, and leaving it in the Aves incertae sedis. *Procuculus minutus* Harrison & Walker, 1977a is type by original designation of the genus *Procuculus* Harrison & Walker, 1977a: 44.

***Proherodius oweni* Lydekker**

Proherodius oweni Lydekker, 1891a: 60 [Holotype from Primrose Hill: imperfect sternum; BMNH 43164. Figured by Owen 1846, fig. 236, Harrison & Walker 1977a, pl. 6, fig. p-q, Harrison & Walker 1978a, fig. 1b, 2b.]

Distribution: Early Eocene (MP 7-10) of Primrose Hill, England (Owen 1846: 556 sub "small wading bird", Lydekker 1891a, Harrison & Walker 1977a, 1978a, Harrison 1979a).

Remarks: Described as a heron (Lydekker 1891a). Harrison & Walker (1978a) and Harrison (1979a) transferred the species to the Presbyornithidae, but Olson (1985a: 166) disagreed. Hence, I relegate it to the Aves incertae sedis until its holotype is restudied. *Proherodius oweni* Lydekker, 1891a is type by original designation of the genus *Proherodius* Lydekker, 1891a: 60.

***Proplegadis fisheri* Harrison & Walker**

Proplegadis fisheri Harrison & Walker, 1971: 368 [Holotype from Sheppey: distal end of left humerus; BMNH A-10. Figured by Harrison & Walker 1971, pl. 11, fig. a-e, and Harrison & Walker 1977, pl. 7, fig. a-c.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison & Walker 1971, 1977a).

Remarks: Described in the Threskiornithidae. Olson (1981a: 167) relegated the species to the category of Aves incertae sedis. *Proplegadis fisheri* Harrison & Walker, 1971 is type by original designation of the genus *Proplegadis* Harrison & Walker, 1971: 368.

***Protopelicanus cuvieri* Reichenbach**

Protopelicanus cuvieri Reichenbach, 1852: vii [Holotype from Montmartre: right femur; MNHN 7978. Figured by Cuvier 1822, pl. 73, fig. 13, Cuvier 1835, pl. 139, fig. 13, and Brunet 1970, fig. 2a-d.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Cuvier 1822, 1835, Reichenbach 1852, Brunet 1970, Harrison 1979b, Olson 1985a, Mlíkovský 1995a).

Remarks: This species was originally treated as a pelican (Reichenbach 1852, Brunet 1970). Harrison (1979b) transferred it to the Sulidae, and Olson (1985a: 202) found some similarities to the Pelagornithidae. The taxonomic position and validity of this species is uncertain until its holotype is restudied (see also Mlíkovský 1995a). *Protopelicanus cuvieri* Reichenbach, 1852 is type by monotypy of the genus *Protopelicanus* Reichenbach, 1852: vii.

***Protornis blumeri* Heer**

Protornis blumeri Heer, 1865: 236 [Holotype from Matt: partial skeleton impression in a slab; lost - Peyer 1957. Figured by Heer 1865, fig. 143, Heer 1879, fig. 178, and Peyer 1957, fig. 25.]

Distribution: Early Oligocene (MP 21-24) of Matt, Switzerland (Heer 1865, 1879, Peyer 1957, Olson 1976a).

Remarks: The identity of this species is indeterminate on the basis of the poorly illustrated

hotype. If the specimen belongs to a bird, which is not certain, then *Protornis blumeri* is a meaningless name until its holotype is located and restudied (see also Peyer 1957, Olson 1976a, Mlíkovský 1992b).

***Pumiliornis tessellatus* Mayr**

Pumiliornis tessellatus Mayr, 1999b: 76 [Holotype from Messel: almost complete skeleton in a slab; SMF ME-2092a,b. Figured by Mayr 1999b, fig. 1a,b.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1999b).

Remarks: Described in order and family incertae sedis. *Pumiliornis tessellatus* Mayr, 1999b is type by original designation of the genus *Pumiliornis* Mayr 1999b: 76.

***Quasisyndactylus longibrachis* Mayr**

Quasisyndactylus longibrachis Mayr 1998c: 36 [Holotype from Messel: almost complete skeleton in a slab; SMF ME-1418a-b. Figured by Mayr 1998c, pl. 8-9.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 1998c).

Remarks: Described in the Alcediniformes incertae sedis. *Quasisyndactylus longibrachis* Mayr, 1998c is type by original designation of the genus *Quasisyndactylus* Mayr, 1998c: 30.

***Quercypodargus olsoni* Mourer-Chauviré**

Quercypodargus olsoni Mourer-Chauviré, 1989a: 2051 [Holotype from Quercy: right tarso-metatarsus; MNHN QU-15876. Figured by Mourer-Chauviré 1989a, fig. 2/1-2.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Mourer-Chauviré 1989a); and late Eocene (MP 16) of Lavergne, France (Mourer-Chauviré 1989a).

Remarks: Described in the Podargidae (Mourer-Chauviré 1989a). Mayr (1999c: 524-525) indicated, that the species does not belong to this family. *Quercypodargus olsoni* Mourer-Chauviré, 1989a is type by original designation of the genus *Quercypodargus* Mourer-Chauviré, 1989a: 2048.

***Rallus adelus* Oberholser**

Rallus intermedius Milne-Edwards, 1869: 144 [Holotype from Montmartre: associated skull, cervical vertebrae and crushed pectoral bones in slab; MNHN 7995. Figured by Milne-Edwards 1869-1871, pl. 103, fig. 17, Brunet 1970, pl. C above, Cracraft 1973, fig. 9. This is a junior primary homonym of *Rallus intermedius* Hermann, 1804.]

Rallus adelus Oberholser, 1917. [New name for *Rallus intermedius* Milne-Edwards, 1869; hence its junior objective synonym.]

Quercyrallus intermedius (Milne-Edwards): Lambrecht 1933: 461 [New combination; *Quercyrallus* was nomen nudum at this point.]

Quercyrallus ludianus Brodkorb, 1963a: 542 [New name for *Rallus intermedius* Milne-Edwards, 1869; hence its junior objective synonym.]

Ludiortyx adelus (Oberholser): Olson 1977: 344 [New combination.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Milne-Edwards 1869-1871, Lambrecht 1933, Brunet 1970, Cracraft 1973, Olson 1977).

Remarks: My brief reexamination of the holotype in MNHN showed that the holotypical skull of this species differs in general morphology (particularly in the position of the foramen magnum and in the insertion of the bill) from the Rallidae, and should be removed from the family (Mlíkovský, orig.). It is here relegated to Aves incertae sedis, pending further study.

***Rallus richmondi* Olson**

Rallus dubius Portis, 1887: 183 [Holotype from Senigallia: sternum impression in slab; MCI. Figured by Portis 1887, pl. 1, fig. 2. This is a junior primary homonym of *Rallus dubius* Piller & Mitterpacher, 1783.]

Rallus richmondi Olson, 1977: 349 [New name for *Rallus dubius* Portis, 1887; hence its junior objective synonym.]

Distribution: Late Miocene (MN 13) of Senigallia, Italy (Portis 1887).

Remarks: Described as a rail (family Rallidae). The species is best relegated to Aves incertae sedis until its holotype is located and restudied (see also Olson 1977: 349).

***Remiornis heberti* Lemoine**

Remiornis heberti Lemoine, 1881a: 158 [Lectotype from Cernay-les-Reims (selected by Martin 1992: 102): partial crushed left tarsometatarsus composed from at least 15 fragments (Mlíkovský, orig.); MNHN CRL-2481. Figured by Lemoine 1881a, pl. 9, fig. 13a, and Martin 1992, fig. 7b, 8a-c. Paralectotypes from Cernay-les-Reims: mandibular symphysis (lost?), "left coracoid" = indeterminate scrap of bone (MNHN CRL-2476), thoracic vertebra (MNHN CRL-2482), another thoracic vertebra (lost?), "proximal end of a humerus" = indeterminate scrap of bone (MNHN CRL-2473), distal end of a humerus (lost?), and "proximal end of an ulna" (MNHN CRL-2474). Figured by Lemoine 1881a, pl. 8, fig. 21-22 ("proximal end of humerus"), 23 ("distal end of humerus"), and 24-25 ("ulna"), and pl. 9, fig. 7-8 (mandibular symphysis), 9 (lost vertebra), 10-12 (preserved vertebra), 13 ("coracoid"), and 13a (tarsometatarsus). The mandibular symphysis belongs to a tryonichid turtle, the distal end of the humerus probably also belongs to a turtle, the "coracoid" is probably referable to *Gastornis*, and the preserved vertebra is indeterminate within the Aves according to Martin (1992: 102-103).

Distribution: Late Paleocene (MP 6) of Cernay-les-Reims, France (Lemoine 1881a,b, Martin 1992).

Remarks: *Remiornis heberti* Lemoine was included in the synonymy of *Gastornis minor* Lemoine by Paris (1912: 290) and Lambrecht (1921: 9, 1933: 578). Brodkorb (1967: 141) erroneously believed, that *Remiornis heberti* Lemoine is a new name for *Gastornis minor* Lemoine. Accordingly, all these authors believed that *Remiornis heberti* is a gastornithid. Martin (1992) showed that *Remiornis heberti* Lemoine was based on different type materials than *Gastornis minor* Lemoine, and separated the species at the family and order levels. *Remiornis heberti* Lemoine, 1881a is type by monotypy of the genus *Remiornis* Lemoine, 1881a: 158. *Remiornis* Lemoine, 1881a is type of the family Remiornithidae Martin, 1992: 102, and order Remiornithiformes Martin, 1992: 102.

***Rupelornis definitus* Beneden**

Rupelornis definitus Beneden, 1871: 260 [Holotype from Rupelmonde: distal end of a tibiotarsus; IRScNB or ULg. Figured by Beneden 1871, unnumbered pl., fig. 7.]

Distribution: Middle Oligocene (MP 23-24) of Rupelmonde, Belgium (Beneden 1871), and Espenhain, Germany (Fischer 1983a sub ?*Rupelornis*).

Remarks: Described as a shorebird, but also compared to owls in the original description (Beneden 1871). Lambrecht (1933: 548) and Brodkorb (1967: 205) included the bird in the family Laridae. I relegate here *Rupelornis definitus* Beneden, 1871 to the category of Aves incertae sedis until its holotype is located and restudied. *Rupelornis definitus* Beneden, 1871 is type by monotypy of the genus *Rupelornis* Beneden, 1871: 260.

***Salmila robusta* Mayr**

Salmila robusta Mayr, 2000b: 188 [Holotype from Messel: almost complete skeleton in slab; SMF ME-3014. Figured by Mayr 2000b, fig. 1, 3, 5 and 6.]

Distribution: Middle Eocene (MP 11) of Messel, Germany (Mayr 2000b).

Remarks: Tentatively included in suborder Cariamae by Mayr (2000b). *Salmila robusta* Mayr is type by original designation of the genus *Salmila* Mayr, 2000b: 188.

***Scaniornis lundgreni* Dames**

Scaniornis Lundgreni Dames, 1890: 4 [Holotype from Limnhamn: associated right coracoid, right scapula and right humerus; NRM. Figured by Dames 1890, pl. 1, and Lierl 1993, fig. 4.]

Distribution: Early Paleocene (MP 1-5) of Limnhamn, Sweden (Dames 1890, Lambrecht 1933, Olson & Feduccia 1980a: 44, Tyrberg & Erickson 1996), and Selk, Germany (Lierl 1993 sub *Scaniornis*?).

Remarks: The fossil was originally thought to be related to the Phoenicopteridae. It does not show any characters linking it to flamingos, however, but its taxonomic position within the neornithine birds is not clarified yet (Tyrberg & Erickson 1996). *Scaniornis lundgreni* Dames is type by monotypy of the genus *Scaniornis* Dames, 1890: 4. *Scaniornis* Dames, 1890 is type of the family Scaniornithidae Lambrecht, 1933: 334.

***Sitta cuvieri* Gervais**

Sitta? Cuvieri Gervais, 1848-1852: 228 [Holotype from Montmartre: partial skeleton imprint in slab and conter-slab; MNHN 7975 and 7976. Figured by Gervais, 1848-1852, pl. 50, fig. 2a-b, Milne-Edwards 1869-1871, pl. 161, fig. 2-3, Brunet 1970, pl. A, fig. f-g, and Mayr 1998c, pl. 16 fig. upper right.]

Palaegithalus Cuvieri (Gervais): Milne-Edwards, 1871: 378 [New combination.]

Distribution: Late Eocene (MP 19) of Montmartre, France (Gervais 1848-1852, Milne-Edwards 1869-1871, Brunet 1970, Harrison 1979b, Mayr 1998c).

Remarks: Described as a nuthatch. Lambrecht (1933: 640) transferred the species to the Paridae, while Brunet (1970: 48) to the Motacillidae. Mayr (1998c: 52) observed, that the species is similar to the *Sylviornis* Mourer-Chauviré, and hypothesized that *Sylviornis* Mourer-Chauviré could be synonymous with *Palaegithalus* Milne-Edwards. This is obviously an *Avis incertae sedis* until its holotype is restudied (see also Brodkorb 1978: 217). *Sitta cuvierii* Milne-Edwards, 1871 is type by monotypy of the genus *Palaegithalus* Milne-Edwards, 1871: 378. *Palaegithalus* Milne-Edwards, 1871 is type of the family Palaegithalidae Harrison, 1979: 108.

***Stintonornis mitchelli* Harrison**

Stintonornis mitchelli Harrison, 1984a: 182 [Holotype from Sheppey: distal end of right tarsometatarsus, England; BMNH A-5284. Figured by Harrison 1984a, fig. 1b-c, 2a-f.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (Harrison 1984a).

Remarks: Described in the Falconidae. Here relegated to the category of *Aves incertae sedis*. *Stintonornis mitchelli* Harrison, 1984a is type by original designation of the genus *Stintonornis* Harrison, 1984a: 182.

***Strigogyps dubius* Gaillard**

Strigogyps dubius Gaillard, 1908: 39 [Holotype from Quercy: distal end of left tibiotarsus; BSP 2, apparently destroyed during the World War II – Mlíkovský, orig. Cast in ML PQ-1073. Figured by Gaillard 1908, text-fig. 5, pl. 2, fig. 1-4.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Gaillard 1908, Mourer-Chauviré 1987).

Remarks: Described as an owl. Transferred to *Aves incertae sedis* by Mourer-Chauviré (1987: 119-123). *Strigogyps dubius* Gaillard is type by monotypy of the genus *Strigogyps* Gaillard, 1908: 39.

***Sylphornis bretouensis* Mourer-Chauviré**

Sylphornis bretouensis Mourer-Chauviré, 1988a: 41 [Holotype from Bretou: distal end of right tarsometatarsus; USTL BRT-1025. Figured by Mourer-Chauviré 1988a, pl. 3, fig. 3-4, and text-fig. 3.]

Distribution: Late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1988a). The alleged record of the family Sylphornithidae from the middle Eocene (MP 11) of Messel, Germany (Peters 1991) was reidentified as belonging to *Messelirrisor parvus* Mayr by Mayr (1998c: 25).

Remarks: Described in the Coraciiformes. The holotypical tarsometatarsus of this species markedly differs from the same element of both the Eocene (see Mayr & Mourer-Chauviré 2000 for figures) and modern coraciiforms (Mlíkovský orig.) in the configuration of trochleae. I relegate thus *Sylphornis bretouensis* Mourer-Chauviré to the category of Aves incertae sedis until the holotype is restudied. *Sylphornis bretouensis* Mourer-Chauviré, 1988a is type by original designation of the genus *Sylphornis* Mourer-Chauviré, 1988a: 41. *Sylphornis* Mourer-Chauviré, 1988a as type of the family Sylphornithidae Mourer-Chauviré, 1988a: 41.

Mourer-Chauviré (1999b: 88) indicated that *Eutreptodactylus itaboraiensis* Baird & Vickers-Rich, 1997 from the Paleocene of Itaboraí, Brazil, belongs to the Sylphornithidae (but see Mayr 2001c).

***Tapinopus ellioti* Milne-Edwards**

Tapinopus Ellioti Milne-Edwards, 1892: 79 [Holotype from Quercy: right tarsometatarsus; MNHN QU-3040. Figured by Gaillard 1908, pl. 2, fig. 9-12, Cracraft & Rich 1972, fig. 2a-d.]

Diatropornis ellioti (Milne-Edwards): Oberholser 1899: 203 [New combination.]

Distribution: Eocene or Oligocene (MP 16-28) of Quercy, France (Milne-Edwards 1892, Gaillard 1908, Cracraft & Rich 1972); and late Eocene (MP 16) of Bretou, France (Mourer-Chauviré 1988).

Remarks: As judged from figures, the holotypical tarsometatarsus of *Tapinopus ellioti* Milne-Edwards differs from the same element of the Cathartidae in being slender, somewhat curved antero-posteriorly, and in having much less protruding intercotylar prominence. Hence, I relegate here *Tapinopus* (= *Diatropornis*) *ellioti* Milne-Edwards to the category of Aves incertae sedis. *Diatropornis ellioti* Milne-Edwards, 1892 is type by monotypy of the genus *Tapinopus* Milne-Edwards, 1892: 79, which is a junior homonym of *Tapinopus* Saussure (see Oberholser 1899). *Diatropornis* Oberholser, 1899: 203 is a new name for *Tapinopus* Milne-Edwards, 1892.

***Teracus littoralis* Milne-Edwards**

Teracus littoralis Aymard, 1856: 233 [Nomen nudum; no description or indication.]

Teracus littoralis Aymard: Milne-Edwards, 1863: 154 [Generic name misspelled, but it continued to be nomen nudum, lacking description or indication.]

Teracus littoralis Milne-Edwards, 1871: 453 [Holotype from Ronzon: femur in a slab; formerly in coll. Aymard, present location unknown. Figured by Milne-Edwards 1869-1871, pl. 185 (not pl. 135 as stated by Milne-Edwards 1871: 453), fig. 20-22. Lydekker (1891a: 28) correctly recognized that this specimen is the holotype of the species, while Lambrecht (1933: 405), Brodkorb (1964: 252), Cracraft & Rich (1972: 281) and Olson (1978: 448) included in the alleged syntypical series also a coracoid in another slab, figured by Milne-Edwards (1869-1871) on pl. 185, fig. 23. However, the caption to the latter figure reads "Coracoid ... que M. Aymard considéré comme appartenant au *Teracus littoralis*", i.e. "coracoid, believed by Mr. Aymard to belong to *Teracus littoralis*" (my translation), which clearly disqualifies the specimen as a possible syntype.]

Distribution: Early Oligocene (MP 21) of Ronzon, France (Aymard 1856, Milne-Edwards 1869-1871, Olson 1978).

Remarks: Aymard (1856), Paris (1912), Milne-Edwards (1869-1871) and Lambrecht (1933:

405) considered this species a raptor of some sort. Brodkorb (1964: 252) listed it in the Vulturidae without explanation. Cracraft & Rich (1972: 281) and Olson (1978: 447) excluded *Teracus littoralis* Milne-Edwards from both the Vulturidae (= Cathartidae) and the Falconiformes. The species is best included in the category of the Aves incertae sedis until its holotype is located and restudied. *Teracus littoralis* Milne-Edwards, 1871 is type by monotypy of the genus *Teracus* Milne-Edwards, 1871: 453. *Ieracus* Aymard, 1856: 233 is nomen nudum (no definition).

***Totanus praecursor* Laube**

Totanus praecursor Laube, 1901: 66 [Holotype from Břešřany: pelvic impression in slab and counter-slab; present location unknown – see Mlíkovský 1992b: 442. Figured by Laube 1901, fig. 14.]

Distribution: Early Miocene (MN 3) of Břešřany, Czechia (Laube 1901, Mlíkovský 1992b, 1999c).

Remarks: Described in the Scolopacidae. The figure of the holotype indeed shows a pelvis generally similar to that of the Scolopacidae, but the species was relegated to Aves incertae sedis until its holotype is found and restudied (Mlíkovský 1992b: 442, 1999c).

***Tynskya eocaena* Mayr**

Tynskya eocaena Mayr, 2000d: 60 [Holotype from Tynsky quarry: almost complete skeleton in slab; BSP 1997-1-6. Figured by Mayr 2000d, fig. 1-5.]

Distribution: Early Eocene (MP 8) of Walton-on-the-Naze, England (Mayr 2000d, tentatively referred). Extralimital record is from the early Eocene (Green River Formation) of the Tynsky quarry, Wyoming (Mayr 2000d).

Remarks: Described in family and order incertae sedis, but daid to be raptor-like (Mayr 2000d). *Tynskya eocaena* Mayr, 2000d is type by original designation of the genus *Tynskya* Mayr, 2000d: 60.

***Vanellus Selysii* Beneden**

Vanellus Selysii Beneden, 1871: 259 [Holotype from Rupelmonde: distal end of a humerus; IRScNB. Figured by Beneden 1871, pl. 1, fig. 2.]

Distribution: Middle Oligocene (MP 23-24) of Rupelmonde, Belgium (Beneden 1871, 1872: 287).

Remarks: Described as a lapwing. The name is meaningless until the holotype of the species is located and restudied (see also Olson 1985a: 175).

***Villettus grandis* Harrison & Walker**

Villettus grandis Harrison & Walker, 1976a: 346 [Holotype from Highcliffe: distal end of right tibiotarsus; BMNH A-4354. Figured by Harrison & Walker 1976a, pl. 6, fig. 1-p.]

Distribution: Late Eocene (MP 14-16) of Highcliffe, England (Harrison & Walker 1976a).

Remarks: Described in the Scolopacidae. Here transferred to the Aves incertae sedis. *Villettus grandis* Harrison & Walker, 1976a is type by original designation of the genus *Villettus* Harrison & Walker, 1976a: 346.

***Villettus waltoni* Harrison & Walker**

Villettus waltoni Harrison & Walker, 1976a: 347 [Holotype from Barton: distal end of left tibiotarsus; IGS GSM-113108. Figured by Harrison & Walker 1976a, pl. 7, fig. a-e.]

Distribution: Late Eocene (MP 14-16) of Barton, England (Harrison & Walker 1976a).

Remarks: Described in the Scolopacidae. Here transferred to the Aves incertae sedis.

***Vultur fossilis* Keferstein**

Vultur fossilis Keferstein, 1834: 243 [Holotype from Westeregeln: proximal end of a femur (see Gernar 1826: 612); formerly in coll. Keferstein, present location unknown. Not figured.]

Vultur fossilis Giebel, 1847: 9 [Same holotype as for *Vultur fossilis* Keferstein. This is both a junior homonym and a junior objective synonym of *Vultur fossilis* Keferstein, 1834 (see Mlíkovský 1998g: 24).]

Distribution: Late Pleistocene (MQ 2C) of Westeregeln, Germany (Gernar 1826 sub *Vultur*, Keferstein 1834, Giebel 1847).

Remarks: Both *Vultur fossilis* Keferstein and *Vultur fossilis* Giebel are meaningless names until their holotype is located and restudied. Lydekker (1891a: 32, tentatively) and Brodkorb (1964: 286) synonymized these species (with inaccurate citations – see Mlíkovský 1998g: 24) with the modern *Aegypius monachus* (Linnaeus, 1766), which is not substantiated (Mlíkovský 1998g: 24).

Ichnotaxa

***Anatipeda anas* Panin & Avram**

Anatipeda anas Panin & Avram, 1962: 467 [Syntypes from Putna: series of footprints of several individuals; present location unknown. Figured by Panin & Avram 1962, pl. 10, fig. 31-32; and Panin 1964, fig. 4.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962).

Remarks: *Anatipeda anas* Panin & Avram, 1962 is type by monotypy of the genus *Anatipeda* Panin & Avram, 1962: 467. *Anatipeda* Panin & Avram, 1962 is type of the family Anatipedae Panin & Avram, 1962: 478 (spelling emended to Anatipedidae by Panin 1964: 343 and Mlíkovský 2000d: 78).

***Anatipeda* sp.**

Distribution: Early Miocene (MN 3-4) of Piatra Neamț, Romania (Panin 1964).

***Ardeipeda egretta* Panin & Avram**

Ardeipeda egretta Panin & Avram, 1962: 463 [Holotype from Putna: two footprints (right and left) of a single individual; present location unknown. Figured by Panin & Avram 1962, pl. 5, fig. 21.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962).

Remarks: *Ardeipeda egretta* Panin & Avram, 1962 is type (by present designation) of the genus *Ardeipeda* Panin & Avram, 1962: 363. *Ardeipeda* Panin & Avram, 1962 is type of the family Ardeipedae Panin & Avram, 1962: 478 (spelling corrected to Ardeipedidae by Mlíkovský 2000d: 79).

***Ardeipeda gigantea* Panin & Avram**

Ardeipeda gigantea Panin & Avram, 1962: 463 [Holotype from Putna: a footprint; present location unknown. Figured by Panin & Avram 1962, pl. 5, fig. 22.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962).

***Ardeipeda incerta* Panin & Avram**

Ardeipeda incerta Panin & Avram, 1962: 464 [Syntypes from Putna: series of footprints in a slab, and a footprint in another slab; present location unknown. Figured by Panin & Avram 1962, pl. 6, fig. 23-24.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962).

***Aviadactyla media* Kordos**

Aviadactyla media Kordos, 1983: 276 [Holotype from Ipolytarnóc: two footprints (left and right) of a single individual; GIB V-12729. Figured by Kordos 1983, fig. 6.]

Distribution: Early Miocene (MN 3) of Ipolytarnóc, Hungary (Kordos 1983).

Remarks: *Aviadactyla media* Kordos, 1983 is type by monotypy of the genus *Aviadactyla* Kordos, 1983: 276.

***Avipeda filiportatis* Vjalov**

Avipeda filiportatis Vjalov, 1965: 112 [Holotype from Deljatin: series of three footprints; IGGL. Figured by Vjalov 1965, pl. 15, and Vjalov 1966, pl. 33 and pl. 34, fig. 1.]

Distribution: Early Miocene (MN 3) of Deljatin, Ukraine (Vjalov & Flerov 1952, Vjalov 1965, 1966: 124); and early Miocene (MN 3-4) of Schitu Frumoasa, Romania (Grozescu 1914:

143, pl. 12 sub unidentified avian footprint – tentatively identified by Vjalov 1966: 124-126).

***Avipeda phoenix* Vjalov**

Avipeda phoenix Vjalov, 1965: 112 [Holotype from Deljatin: series of 10 footprints in a slab; IGGL. Figured by Vjalov 1965, pl. 14, fig. 2.]

Distribution: Early Miocene (MN 3) of Deljatin, Ukraine (Vjalov 1965, 1966: 121).

Remarks: *Avipeda phoenix* Vjalov, 1965 is type (by present designation) of the genus *Avipeda* Vjalov, 1965: 112.

***Avipeda sirin* Vjalov**

Avipeda sirin Vjalov, 1965: 112 [Holotype from Nižnij Strutyn': a footprint; IGGL. Figured by Vjalov 1960, fig. 1, and Vjalov 1966, pl. 31, fig. 1.]

Distribution: Early Miocene (MN 3) of Nižnij Strutyn', Ukraine (Vjalov 1960, 1965, 1966: 123).

***Charadriipeda becassi* Panin & Avram**

Charadriipeda becassi Panin & Avram, 1962: 467 [Holotype from Putna: series of footprints of a single individual; present location unknown. Figured by Panin & Avram 1962, pl. 9, fig. 30.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962), and Cvejdi, Romania (Panin 1964).

***Charadriipeda disjuncta* Panin & Avram**

Charadriipeda disjuncta Panin & Avram, 1962: 467 [Holotype from Putna: series of footprints of a single individual; present location unknown. Figured by Panin & Avram 1962, pl. 9, fig. 29.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962), and Călugăr, Romania (Panin 1964).

***Charadriipeda limosa* Rădan & Brustur**

Charadriipeda limosa Rădan & Brustur, 1993: 74. [Holotype from Câmpu lui Neag: footprint; IGGB P-18.592. Figured by Rădan & Brustur 1993, pl. 1, fig. 1c, 2.]

Distribution: Late Oligocene (MP 25-30) of Romania (Rădan & Brustur 1993).

***Charadriipeda minima* Panin & Avram**

Charadriipeda minima Panin & Avram, 1962: 466 [Syntypes from Putna: series of footprints of at least two individuals; present location unknown. Figured by Panin & Avram 1962, pl. 8, fig. 28.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962), and Cvejdi, Romania (Panin 1964).

***Charadriipeda minor* Panin**

Charadriipeda minor Panin, 1964: 346. [Syntypes from Cvejdi: series of footprints in three (?) plates; present location unknown. Figured by Panin 1964, fig. 2, 6b, 7, and 8b.]

Distribution: Early Miocene (MN 3-4) of Cvejdi, Romania (Panin 1964).

Remarks: *Charadriipeda minor* Panin & Avram, 1962 is type (by present designation) of the genus *Charadriipeda* Panin & Avram, 1962: 466. *Charadriipeda* Panin & Avram, 1962 is type of the family Charadriipedae Panin & Avram, 1962: 478 (spelling corrected to Charadriipedidae by Panin 1964: 343 and Mlíkovský 2000d: 79).

***Charadriipeda recurvirostrioidea* Panin & Avram**

Charadriipeda recurvirostrioidea Panin & Avram, 1962: 466 [Syntypes from Putna: series of footprints in three slabs; present location unknown. Figured by Panin & Avram 1962, pl. 7, fig. 26, and pl. 8, fig. 27.]

Charadriipeda recurvirostra Panin, 1964: 343 [New name for *Charadriipeda recurvirostrioidea* Panin & Avram, 1962; hence its junior objective synonym.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962), and Mîngălar, Romania (Panin 1964).

***Gruipeda intermedia* Panin**

Gruipeda intermedia Panin, 1964: 347. [Holotype from Cujești: series of footprints; present location unknown. Figured by Panin 1964, fig. 3.]

Distribution: Early Miocene (MN 3-4) of Cujești, Romania (Panin 1964).

***Gruipeda maxima* Panin & Avram**

Gruipeda maxima Panin & Avram, 1962: 465 [Holotype from Putna: a footprint; present location unknown. Figured by Panin & Avram 1962, pl. 7, fig. 25.]

Distribution: Early Miocene (MN 3-4) of Putna, Romania (Panin & Avram 1962).

Remarks: *Gruipeda maxima* Panin & Avram, 1962 is type by monotypy of the genus *Gruipeda* Panin & Avram, 1962: 465. *Gruipeda* Panin & Avram, 1962 is type of the family Gruipedae Panin & Avram, 1962: 478 (spelling corrected to Gruipedidae by Panin 1964: 343 and Mlíkovský 2000d: 79).

***Ornithotarnocia lambrechtii* Kordos**

Ornithotarnocia lambrechtii Kordos, 1983: 269 [Holotype from Ipolytarnóc: two footprints (left and right) of a single individual; GIB V-12721. Figured by Lambrecht 1912b, pl. 1, Lambrecht 1933, fig. 183, and Kordos 1983, fig. 1.]

Distribution: Early Miocene (MN 3) of Ipolytarnóc, Hungary (Lambrecht 1912b, Kordos 1983).

Remarks: *Ornithotarnocia lambrechtii* Kordos, 1983 is type by monotypy of the genus *Ornithotarnocia* Kordos, 1983: 269.

***Ornitichnites taurinus* Portis**

Ornitichnites [sic!] *taurinus* Portis, 1887: 201 [Holotype from Verrua Savoia: a series of at least 10 footprints in a slab, which was described but not named by Portis 1884: 381; present location unknown. Not figured.]

Distribution: Late Eocene (MP 17-20) of Verrua Savoia, Italy (Portis 1884: 381 sub "gli ornitichniti di Verrua", Portis 1887).

Remarks: Note that citations of relevant Portis's papers in Lambrecht (1933: 659) and Brodkorb (1978: 221) are incorrect.

***Passeripedia ipolyensis* Kordos**

Passeripedia ipolyensis Kordos, 1983: 280 [Holotype from Ipolytarnóc: a footprint (right?); GIB, uncatalogued. Figured by Kordos 1983, fig. 8.]

Distribution: Early Miocene (MN 3) of Ipolytarnóc, Hungary (Kordos 1983).

Remarks: *Passeripedia ipolyensis* Kordos, 1983 is type by monotypy of the genus *Passeripedia* Kordos, 1983: 280.

***Pulchravipes magnificus* Demathieu et al.**

Pulchravipes magnificus Demathieu, Ginsburg, Guérin & Truc, 1984. [Holotype from Saignon: footprints.]

Distribution: Early Oligocene (MP 21) of Saignon, France (Demathieu et al. 1984).

Remarks: Not seen. *Pulchravipes magnificus* Demathieu et al., 1984 is type by monotypy of the genus *Pulchravipes* Demathieu, Ginsburg, Guérin & Truc, 1984.

***Tetraornithopedia tasnadii* Kordos**

Tetraornithopedia tasnadii Kordos, 1983: 279 [Holotype from Ipolytarnóc: footprint (left?); CHI, uncatalogued. Figured by Kordos 1983, fig. 7.]

Distribution: Early Miocene (MN 3) of Ipolytarnóc, Hungary (Kordos 1983).

Remarks: *Tetraornithopedia tasnadii* Kordos is type by monotypy of the genus *Tetraornithopedia* Kordos, 1983: 279.

Taxa non avium

***Bucklandium diluvii* König**

Bucklandium diluvii König, 1825: pl. 8, fig. 91. [Holotype from Sheppey: skull in a slab; BMNH. Figured by König 1825, fig. 91, and Woodward 1889, pl. 22, fig. 1-4.]

Distribution: Early Eocene (MP 8-9) of Sheppey, England (König 1825).

Remarks: Described as a bird. The species was transferred to *Acanthopterygii* by Morris (1843: 193), and Woodward (1889) recognized its affinities to silurid fishes.

***Larus priscus* Giebel**

Larus priscus Giebel, 1847: 31 [Holotype from Seveckenberg: "distal end of right tibia" = distal end of a tarsus; MNHN, uncatalogued. Not figured.]

Distribution: Late Pleistocene (MQ 2C) of Seveckenberg, Germany (Giebel 1847, 1850).

Remarks: This species was based on a partial tarsus of the dipodid rodent *Allactaga* Cuvier, 1836 (Nehring 1880: 436, Mlíkovský orig.).

***Macrornis tanaupus* Seeley**

Macrornis tanaupus Seeley, 1866: 110 [Holotype from Hordle: "proximal end of tibiotarsus" = distal end of a femur; SMC C-20910. Not figured.]

Distribution: Late Eocene (MP 17) of Hordle, England (Seeley 1866).

Remarks: Harrison & Walker (1976a: 344) restudied the holotype of this species, showing that it is not tibiotarsus of a struthious bird, as believed by Seeley (1866), but a reptilian femur. *Macrornis* Seeley, 1866 is type by monotypy of the genus *Macrornis* Seeley, 1866: 110.

***Ornithoidichnites badensis* Böhm**

Ornithoidichnites badensis Böhm, 1896: 238 [Holotype from Bellingen: two footprints (left and right) of a single individual (see Mlíkovský 1992b: 445); present location unknown. Figured by Böhm 1896, text-fig. 1, pl. 1.]

Ichnium badense (Böhm): Böhm 1898: 206 [New combination.]

Distribution: Late Oligocene (MP 25-30) of Bellingen, Germany (Böhm 1896, 1898).

Remarks: These footprints, originally described as avian, belong to a tapir (Böhm 1898, see also Abel 1935). Lambrecht (1921: 99, 1933: 665) and Brodkorb (1978: 221) overlooked Böhm's (1898) correction and continued to list this fossil as avian.

***Ornithichnites argenterae* Portis**

Ornithichnites argenterae Portis, 1879: 221 [Holotype from Argentera: a footprint; MGPUB, cast in MPUT.]

Ornithichnidium argenterae (Portis): Kuhn 1958: 382 [New generic name created for *Ornithichnites* Portis, 1879, but no specific name mentioned.]

Distribution: Late Eocene (MP 17-20) of Argentera, Italy (Portis 1879, 1884).

Remarks: Kuhn (1958, 1963) believed that the genus name *Ornithichnites* was created by Portis (1879), and that it is preoccupied by *Ornithichnites* King, "1844" = 1845. However, *Ornithichnites* was used for the first time by Hitchcock (1836) for alleged avian footprints from the Triassic of Massachusetts. The name is to be understood as a name for all avian footprints (see also Gervais 1844a: 7-9), i.e. for a collective group, which has no type (ICZN 1999, Art. 42.3.1). Kuhn (1958) thought that *Ornithichnites* Portis was based on *Ornithichnites argenterae* Portis, 1879. Although this is not true, *Ornithichnites argenterae* Portis, 1879 constitutes type by monotypy of the genus *Ornithichnidium* Kuhn, 1958: 382. The footprints

were formed by a reptile (Kuhn 1958, 1963).

***Saurornis matthesi* Fischer**

Saurornis matthesi Fischer, 1967: 603 [Holotype from Geiseltal XIV: "proximal end of tarsometatarsus" = distal end of coossified ulna and radius; GM Av-G1956/Dia.13. Figured by Fischer 1967, pl. 1, fig. a-c.]

Distribution: Middle Eocene (MP 11) of Geiseltal XIV, Germany (Fischer 1967).

Remarks: Described in the monotypic genus *Saurornis* Fischer, 1967: 603. Subsequently identified as identical with a perissodactyl mammal of the genus *Lophiodon* Cuvier, 1822 (Fischer 1987).

Nomina nuda

All types of unavailable names are listed under this heading.

Genus-group names

- Aurorornis* Panteleev, 1999: 20 [Eocene of Ukraine; unpublished. The only included species, *Aurorornis taurica* Panteleev, 1999, is nomen nudum.]
- Camaskelus* Aymard, 1856: 233 [Miocene of France; no definition. The only included species, *Camaskelus palustris* Aymard, 1856, is nomen nudum.]
- Palaetus* Milne-Edwards, 1871: 572 [Miocene of France; no definition. The only included species, *Palaetus rapax* Milne-Edwards, 1871, is nomen nudum.]
- Procellornis* Lucazeau, 1959: 21 [Miocene of France; unpublished. Three include species (*P. burdigalensis* Lucazeau, 1959, *P. antiquus* Lucazeau, 1959, and *P. dubius* Lucazeau, 1959) are nomina nuda.]
- Protopelargus* Reichenbach, 1852: xiv [Miocene of Germany; no definition. The only included species, *Protopelargus meyeri* Reichenbach, 1852, is nomen nudum.]
- Suloides* Lucazeau, 1959: 46 [Miocene of France; unpublished. The only included species, *Suloides rozieri* Lucazeau, 1959 is nomen nudum.]
- Tantaleus* Reichenbach, 1852: xiv [Pleistocene of Sardinia, Italy. The only included species, *Tantaleus brecciensis* Giebel, 1847, is nomen nudum.]
- Typhornis miocoenicus* Lucazeau, 1959: 29 [Miocene of France; unpublished. The only included species, *Typhornis miocoenicus* Lucazeau, 1959, is nomen nudum.]

Species-group names

- Anas arcensis* Kretzoi, 1962: 168 [Pleistocene of Romania; no description or indication.]
- Anas crassa* Milne-Edwards, 1871: 573 [Miocene of France; no description or indication.]
- Ardea effosa* Meyer in Lepsius, 1883: 146 [Miocene of Germany; no description or indication.]
- Ardea effosa* Meyer in Lepsius, 1892: 623 [Miocene of Germany; no description or indication.]
- Ardea latipes* Meyer in Lepsius, 1883: 146 [Miocene of Germany; no description or indication.]
- Ardea latipes* Meyer in Lepsius, 1892: 623 [Miocene of Germany; no description or indication.]
- Argala arvernensis* Milne-Edwards, 1871: 572 [Miocene of France; no description or indication.]
- Aurorornis taurica* Panteleev, 1999: 20 [Eocene of Ukraine; unpublished.]
- Camaskelus palustris* Aymard, 1856: 233 [Oligocene of France; no description or indication – see Olson 1978.]
- Carbo risgoviensis* O. Fraas in Engel, 1908: 567 [Miocene of Germany; no description or indication.]
- Gallus steinheimensis* E. Fraas in Engel, 1908: 567 [Miocene of Germany; no description or indication.]
- Himantopus brevipes* Milne-Edwards, 1871: 572 [Miocene of France; no description or indication.]
- Larus burdigalensis* Lucazeau, 1959: 45 [Miocene of France; unpublished.]
- Leptoptilus* [sic!] *arvernensis* "(Milne-Edwards)" Lambrecht, 1921: 26 [New combination for *Argala arvernensis* Milne-Edwards, hence a nomen nudum attributable to Lambrecht (1921). See also Harrison (1974: 42), and Hill & Walker (1979: 217).]
- Ornitholithes ralli* Anonymous: Karg, 1805: 26 [Unpublished label name; specimen thought to have originated from the Miocene of Germany, but a forgery in fact, constructed from fish ribs – Pfannenstiel 1958: 854.]
- Otis agilis* Milne-Edwards, 1871: 572 [Miocene of France; no description or indication.]

- Palaeortyx media* Milne-Edwards, 1871: 571 [Miocene of France; no description or indication.]
- Palaetus rapax* Milne-Edwards, 1871: 571 [Miocene of France; no description or indication.]
- Pelagornis Delfortrii* Anonymous: Lambrecht 1933: 282 [Miocene of France; label name; unpublished.]
- Phalacrocorax risgoviensis* "Fraas" Lambrecht, 1921: 37 [New combination for *Carbo risgoviensis* Fraas in Engel, hence a nomen nudum attributable to Lambrecht (1921).]
- Phasianus Nicheti* "Gaillard" Bastin, 1933: 47, 53 [Pleistocene of France; no description or indication.]
- Procellornis antiquus* Lucazeau, 1959: 31 [Miocene of France; unpublished.]
- Procellornis burdigalensis* Lucazeau, 1959: 21 [Miocene of France; unpublished.]
- Procellornis dubius* Lucazeau, 1959: 42 [Miocene of France; unpublished.]
- Protopelargus meyeri* Reichenbach, 1852: xiv [Miocene of Germany; no description or indication – see Mlíkovský 1995a.]
- Sula rozieri* Lucazeau, 1959: 46 [Miocene of France; unpublished.]
- Suloides miocoenicus* Lucazeau, 1959: 48 [Miocene of France; unpublished.]
- Tantaleus brecciensis* Reichenbach, 1852: xiv [New combination for *Tantalus brecciensis* Giebel, 1847, hence a nomen nudum attributable to Reichenbach (1852) – see Mlíkovský 1995a.]
- Tantalus brecciensis* Giebel, 1847: 27 [Pleistocene of Sardinia; no description or indication – see Mlíkovský 1995a.]
- Typhornis miocoenicus* Lucazeau, 1959: 29 [Miocene of France; unpublished.]

Note: Cheneval (1996a: 538) listed Beneden (1871) as the author of two avian genera and species (*Ardeita gracilis* and *Uriopsis scaldicus*), reportedly described from the middle Oligocene (MP 23-24) of Rupelmonde, Belgium. These forms were not listed in previous catalogues of fossil birds (Lambrecht 1921, 1933, Brodkorb 1963c, 1964, 1967, 1971b, 1978), and I was not able to find them in any of Beneden's relevant papers (Beneden 1871, 1872, 1873a,b, 1883). The existence of these names thus remains unclear until their source is located.

References

The authors are listed alphabetically without respect to diacritical marks. Titles of papers published in languages other than English, French or German are translated to English. The classification of languages follows Prince (1998).

- Abel O. (1935): *Vorzeitliche Lebensspuren*. – Jena: Gustav Fischer Verlag, 644 pp.
- Adrover R. (1975): Principales yacimientos paleomastológicos de la provincia de Teruel y su posición estratigráfica relativa [Main paleomammaliological localities from the Teruel Province, and their relative stratigraphical position.] – *Trabajos del Neógeno-Cuaternario* 4: 31-48. [In Spanish.]
- Adrover R. (1986): Nuevas faunas de roedores en el Mio-Plioceno continental de la región de Teruel (España). Interés bioestratigráfico y paleoecológico [New rodent faunas from the continental Mio-Pliocene of the Teruel region (Spain). Application to biostratigraphy and paleoecology.] – Teruel: Instituto de Estudios Turolenses, 423 pp. [In Spanish.]
- Adrover R., Aguirre E., Heintz E., Moissenet E. & Morales J. (1974): Teruel II [Teruel II.] – In: Aguirre E. & Morales J. (eds.): *Coloquio internacional sobre bioestratigrafía continental del Neógeno Superior y Cuaternario Inferior* [International colloquium on the continental biostratigraphy of the Upper Neogene and Lower Pleistocene]: 69-83. Madrid: Consejo Superior de Investigaciones Científicas. [In Spanish.]
- Adrover R., Mein P. & Moissenet E. (1978): Nuevos datos sobre la edad de las formaciones continentales neógenas de los alrededores de Teruel [New data on the state of the continental Neogene formations in the vicinity of Teruel.] – *Estudios Geológicos* 34: 205-214. [In Spanish.]
- Aguirre E., Hozos M., Mensua S., Morales J., Pérez A., Quirantes J., Sánchez L. & Soria D. (1974): Cuenca del Jalón [The cave of Jalón.] – Aguirre E. & Morales J. (eds.): *Coloquio internacional sobre bioestratigrafía continental del Neógeno Superior y Cuaternario Inferior* [International colloquium on the continental biostratigraphy of the Upper Neogene and Lower Pleistocene]: 11-48. Madrid: Consejo Superior de Investigaciones Científicas. [In Spanish.]
- Agustí J. (1988): El ocells [The birds.] – In: *Història Natural dels Països Catalans* [Natural history of Catalonia]. Vol. 15: *Registre Fòssil* [Catalogue of fossils]: 404-405. Barcelona: Enciclopedia Catalana. [In Catalan.]
- Agustí J., Anadón P., Gibert J., Julià R., Martín-Suárez E., Menéndez E., Moyà-Solà S., Pons-Moyà J., Rivas P. & Toro I. (1984): Estratigrafía y paleontología del Pleistoceno Inferior de Venta Micena (Orce, depresión Guadix-Baza, Granada). Resultados preliminares. [Stratigraphy and paleontology of the Lower Pleistocene of Venta Micena (Orce, Guadix-Baza depression, Granada). Preliminary results.] – *Paleontología y Evolución* 18: 19-38. [In Spanish.]
- Alcalde del Río H., Breuil H. & Sierra L. (1912): *Les cavernes de la région Cantabrique (Espagne)*. – Monaco: Imprimerie Chêne, viii + 265 pp.
- Alcover J.A. (1989): Les aus fòssils de la Cova de Ca Na Reia [Fossil birds from the Ca Na Reia cave.] – *Endins* 14/15: 95-100. [In Catalan.]
- Alcover J.A., Moyà-Solà S. & Pons-Moyà J. (1981): *Les quimeres del passat*. – Ciutat de Mallorca. [In Catalan.]
- Alcover J.A., Florit F., Mourer-Chauviré C. & Weesie P.D.M. (1992): The avifaunas of the isolated Mediterranean islands during the middle and late Pleistocene. – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Pierce Brodkorb*. – Natural History Museum of Los Angeles County, Science Series 36: 273-283.
- Alekseev A.K. (1916): Fauna pozvonočnyh d. Novo-Elisavetovki [Vertebrate fauna of Novo-Elisavetovka village.] – *Zapiski Novorossijskogo Instituta* 9: 273-453. [In

- Russian.]
- Alexander, R. McN. (1983): Allometry of the leg bones of moas (*Dinornithes*) and other birds. – *Journal of the Zoological Society of London* 200: 215-231.
- Alfárez F., Molero G., Brea P. & Santafé J.V. (1982): Precisiones sobre la geología, fauna, cronoestratigrafía y paleoecología del yacimiento mioceno de Córcoles [Details on the geology, fauna, chronostratigraphy and paleoecology of the Miocene site of Córcoles.] – *Real Academia de Ciencias Exactas, Físicas y Naturales* 76: 249-276. [In Spanish.]
- Allen E.G. (1951): A sixteenth century classification of birds. – In: Hörstadius S. (ed.): *Proceedings of the Xth international ornithological congress*: 139-143. Uppsala: Almquist & Wiksell.
- Alvarenga H.M.F. (1985): Notas sobre os Cathartidae (Aves) e descrição de um novo gênero do Cenozóico brasileiro [Comments on the Cathartidae (Aves) and description of a new genus from the Cenozoic of Brazil.] – *Anais da Academia Brasileira de Ciências* 57: 349-357. [In Portuguese.]
- Alvarenga H.M.F. (1988): Ave fóssil (Gruiformes: Rallidae) dos folhelhos da Bacia de Taubaté, Estado de São Paulo, Brasil [A fossil bird (Gruiformes: Rallidae) from the shales of the Taubaté Basin, state of São Paulo, Brazil.] – *Anais da Academia Brasileira de Ciências* 65: 335-345. [In Portuguese.]
- Alvarenga H.M.F. (1995): Um primitivo membro da ordem Galliformes (Aves) do Terciário Médio da Bacia de Taubaté, Estado de São Paulo, Brasil [A primitive member of the order Galliformes (Aves) from the Middle Tertiary of Bacia de Taubaté, São Paulo Province, Brazil.] – *Anais da Academia Brasileira de Ciências* 67: 33-44. [In Portuguese.]
- Amadon, D. (1963) Comparison of fossil and recent species: some difficulties. – *Condor* 65: 407-409.
- Ameghino F. (1889): Contribucion al conocenda de los mamíferos fósiles de la República Argentina [A contribution to the knowledge of fossil mammals of the Republic of Argentina.] – *Actas de la Academia Nacional de Ciencias en Córdoba* 6: 1-1027. [In Spanish.]
- Ameghino F. (1891): Enumeración de las aves fósiles de la República Argentina [List of fossil birds of the Republic of Argentina.] – *Revista Argentina de Historia Natural* 1: 441-453. [In Spanish.]
- Ameghino F. (1895): Sur les oiseaux fossiles de la Patagonia. – *Boletín del Instituto Geográfico Argentino* 15: 501-602.
- Ammon L. von (1918): Tertiäre Vogelreste von Regensburg und die jungmiozäne Vogelwelt. – *Abhandlungen des Naturwissenschaftlichen Vereines in Regensburg* 12: 1-70.
- Anderson J.F., Hall-Martin A. & Russell D.A. (1985): Long-bone circumference and weight in mammals, birds and dinosaurs. – *Journal of Zoology (London) (A)* 207: 53-61.
- Andersson J.G. (1923): Essays on the Cenozoic of northern China. On the occurrence of fossil *Struthionidae* in China. – *Memoirs of the Geological Survey of China, A* 3: 53-77.
- Andors A.V. (1988): Giant groundbirds of North America (Aves: Diatrymidae). – Unpublished thesis, Columbia University, New York, 577 pp.
- Andors A.V. (1991): Paleobiology and relationships of the giant groundbird *Diatryma* (Aves: Gastornithiformes). – In: *Acta XX congressus internationalis ornithologici*: 563-571. Wellington: New Zealand Ornithologists' Congress Trust Board.
- Andors A.V. (1992): Reappraisal of the Eocene groundbird *Diatryma* (Aves: Anserimorphae). – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Pierce Brodkorb*. – *Natural History Museum of Los Angeles County, Science Series* 36: 109-125.

- Andrae K. I. (1855): Beiträge zur Kenntnis der fossilen Flora Siebenburgens und des Banates. – Abhandlungen der Königlich-Kaiserlichen Geologischen Reichsanstalt 2.
- Andrée K. (1937): Der Bernstein und seine Bedeutung in Natur- und Geisteswissenschaften, Kunst und Kunstgewerbe, Technik, Industrie und Handel. – Königsberg [= Kaliningrad]: Gräfe & Unzer, 219 pp.
- Andrée K. (1951): Der Bernstein, das Bernsteinland und sein Leben. – Stuttgart: Kosmos, 96 pp.
- Andrews C.W. (1899): On the remains of a new bird from the London Clay of Sheppey. – Proceedings of the Zoological Society of London 1899: 776-785.
- Andrews C.W. (1916): Note on the sternum of a large carinate bird from the (?) Eocene of southern Nigeria. – Proceedings of the Zoological Society of London 1916: 519-524.
- Andrews C.W. (1920): Remains of the Great Auk and Ptarmigan in the Channel Islands. – Annals and Magazine of Natural History (9) 6: 166.
- Andrews P. (1990): Owls, caves and fossils. – London: Natural History Museum, 231 pp.
- Anonymous (1844): Inhalt. – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1844: iii-xviii.
- Antón M., López G. & Santamaría R. (1993): Estudio preliminar de la icnofauna miocena del yacimiento de Salinas de Añana (provincia de Álava) [A preliminary study of the Miocene ichnofauna from the site of Salinas de Añana (Álava Province).] – Comunicaciones de las IX Jornadas de Paleontología: 23-28. [In Spanish.]
- Anzidei A.P., Bietti A., Cassoli P., Ruffo M. & Segre A.G. (1984): Risultati preliminari dell'lo scavo in un deposito Pleistocenico in località Rebibbia – Casal de'Pazzi (Roma). – Atti XXIV Riunione Scientifica: 131-139. Firenze: Istituto Italiano di Preistoria e Protostoria.
- Archiac d' A., Beaumont E. de, Daubré G.A., Verneuil E. de & Quatrefages A. de (1865): Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France, par Alphonse Milne Edwards. Rapport fait à l'Académie des sciences par la Commission chargée de décerner le grand prix des sciences physiques pour 1865. – Annales des Sciences Naturelles (5) 5: 229-240.
- Arredondo O. & Olson S.L. (1994): A new species of owl of the genus *Bubo* from the Pleistocene of Cuba (Aves: Strigiformes). – Proceedings of the Biological Society of Washington 107: 436-444.
- Ascenzi A. & Segre A. (1971): Il giacimento con mandibola neandertaliana di Archi (Reggio Calabria) [The deposits of Archi (Province of Calabria) with an Neanderthalian mandible.] – Accademia Nazionale dei Lincei 50: 763-771. [In Italian.]
- Aslanova S.M. & Burčak-Abramovič N.I. (1982): Oligocenovaja zubastaja ptica iz s. Pereiškjul' (Apšeroniskij poluostrov) – pervaja i edinstvennaja nachodka v SSSR i na vsem Aziatskom kontinente [An Oligocene toothed bird from Pereiškjul' village (Apšeroniskij Peninsula) – the first and the only record from the USSR and in the whole Asian continent.] – Izvestija Akademii Nauk Gruzinskoj SSR, Biologija 8: 406-412. [In Russian.]
- Aslanova S.M. & Burčak-Abramovič N.I. (1999): A detailed description of *Caspiodontornis kobystanicus* from the oligocene of the Caspian seashore. – Acta Zoologica Cracoviensia 42: 423-433.
- Aver'janov A.O., Potapova O.R. & Nesov L.A. (1990): O pervych otečestvennyh nachodkach kostej drevnich ptic [On the first native records of ancient birds.] – In: Potapov R.L. (ed.): Ornitologičeskie issledovanija v Palearktike [Ornithological research in the Palearctic.] – Trudy Zoologičeskogo Instituta Akademii Nauk SSSR 210: 3-9. [In Russian.]
- Aver'janov A.O., Panteleev A.V., Potapova O.R. & Nesov L.A. (1991): Ložnozubyje pticy (Aves: Pelecaniformes: Odontopterygia) pozdnego paleocena i éocena zapadnoj okrainy drevnej Azii [False-toothed birds (Aves: Pelecaniformes: Odontopterygia) from the late

- Paleocene and Eocene of the western margin of ancient Asia.] – Trudy Zoologičeskogo Instituta Akademii Nauk SSSR 239: 3-12. [In Russian.]
- Aymard A. (1856): [Untitled communication.] – Congrès Scientifique de France 22(1): 227-257, 258-277. [In French.]
- Bacher A. (1967): Vergleichend morphologische Untersuchung an Einzelknochen des postkranialen Skeletts in Mitteleuropa vorkommender Schwäne und Gänse. – Unpubl. Thesis, Ludwig Maximilian University, München, 116 pp.
- Bachmann I. (1878): Fossile Eier aus der obern Süßwassermolasse der Umgebung von Luzern. – Abhandlungen der Schweizerischen Paläontologischen Gesellschaft 5(4): 1-7.
- Bachmayer F. (1964): Fossile Vogelfedern aus den jungtertiären Süßwasserablagerungen von Weingraben (Burgenland, Österreich). – Annalen des Naturhistorischen Museums in Wien 67: 175-180.
- Bachmayer F. (1980): Ein fossiler Vogelrest aus den Diatomeen-Schiefern (Miozän, Ottnangien) von Limberg, Niederösterreich. – Annalen des Naturhistorischen Museums in Wien 83: 25-28.
- Bachmayer F. & Zapfe H. (1962): Reste von *Struthio* aus Pikermi. – Praktika 37: 247-253.
- Bachofen-Echt A. (1929): Leben und Sterben im Bernsteinwald II-III. – Palaeobiologica 2: 15-18, 264-269.
- Bachofen-Echt A. (1936a): Das Vorkommen von Federn in der Braunkohle des Geiseltales. – Nova Acta Leopoldina, Neue Folge 4: 335-340.
- Bachofen-Echt A. (1936b): Das Vorkommen von Federn im Bernstein. – Nova Acta Leopoldina, N.S., 4: 341-347.
- Bachofen-Echt A. (1944): Einschlüsse von Federn und Haaren im Bernstein. – Palaeobiologica 8: 113-119.
- Bachofen-Echt A. (1949): Der Bernstein und seine Einschlüsse. – Wien: Springer-Verlag, 204 pp.
- Bährmann U. (1978): Eine biometrische Analyse zur Morphologie der Nebelkrähe (*Corvus corone cornix* L.) und ihrer intraspezifischer Variation. – Zoologische Abhandlungen (Dresden) 35: 223-252.
- Baird R.F. & Vickers-Rich P. (1997): *Eutreptodactylus itaboriensis* gen. et sp. nov., an early cuckoo (Aves: Cuculidae) from the late Paleocene of Brazil. – Alcheringa 21: 123-127.
- Baird R.F. & Vickers-Rich P. (1998): *Palaelodus* (Aves: Palaelodidae) from the middle to late Cainozoic of Australia. – Alcheringa 22: 135-151.
- Bakulovskij N.N. (1892): Gigantskoe jajco, najdennoe v Rossii [A gigantic egg, found in Russia.] – Nauka i Žizn' 1892 (28): 445. [In Russian.]
- Ballmann P. (1966): Die Vögel aus der altdurdigalen Spaltenfüllung von Wintershof (West) bei Eichstätt in Bayern. – Unpub. Thesis, Universität München, vi + 115 pp. [See Ballmann 1969b for the published version of this thesis.]
- Ballmann P. (1969a): Les oiseaux miocènes de La Grive-Saint-Alban (Isère). – Géobios 2: 157-204.
- Ballmann P. (1969b): Die Vögel aus der altdurdigalen Spaltenfüllung von Wintershof (West) bei Eichstätt in Bayern. – Zitteliana 1: 5-61.
- Ballmann P. (1970): Ein neuer Vertreter der Musophagidae (Aves) aus dem Chattien von Gaimersheim bei Ingolstadt (Bayern). – Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie 10: 271-275.
- Ballmann P. (1972): Les oiseaux miocènes de Vieux-Collonges (Rhône). – Documents du Laboratoire de Géologie de la Faculté des Sciences de Lyon 50: 93-101.
- Ballmann P. (1973): Fossile Vögel aus dem Neogen der Halbinsel Gargano (Italien). – Scripta Geologica 17: 1-75.
- Ballmann P. (1976a): Fossile Vögel aus dem Neogen der Halbinsel Gargano (Italien), zweiter Teil. – Scripta Geologica 38: 1-59.

- Ballmann P. (1976b): The contribution of fossil birds to avian classification. – In: Frith H.J. & Calaby J.H. (eds.): Proceedings of the 16th international ornithological congress: 196-200. Canberra: Australian Academy of Sciences.
- Ballmann, P. (1977): Neuere Erkenntnisse über die zeitliche Entstehung der rezenten Vogelarten. – Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie 17: 169-175.
- Ballmann P. (1979): Fossile Glareolidae aus dem Miozän des Nördlinger Ries (Aves: Charadriiformes). – Bonner Zoologische Beiträge 30: 52-101.
- Ballmann P. (1983): A new species of fossil barbet (Aves: Piciformes) from the middle Miocene of Nördlinger Ries (southern Germany). – Journal of Vertebrate Paleontology 3: 43-48.
- Ballmann P. & Adrover H.R. (1970): Yacimiento paleontológico de la cueva de Son Bauzà (Mallorca) [Paleontological record from the Son Bauzà cave.] – Acta Geológica Hispanica 5: 58-62. [In Spanish.]
- Balme D.M. (1962): Genos and eidos in Aristotle's biology. – Classical Quarterly (new series) 12: 81-98.
- Bardhason, G.G. (1911): [Bones of *Alca impennis* at Hunafloi, North Iceland.] – Videnskabelige Meddelelser Dansk Naturhist. For. København 1910: 73. [In Danish.]
- Baryšnikov G.F. & Potapova O.R. (1992): Paleolithic birds of the Crimean peninsula, USSR. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County, Science Series 36: 293-305.
- Baryšnikov G.F. & Potapova O.R. (1995): Pleistocene birds from the Acheulean site of Treugolnaya Cave in the northern Causasus. – In: Peters D.S. (ed.): Acta palaeornithologica. – Courier Forschungsinstitut Senckenberg 181: 241-248.
- Bastin A. (1933): La faune pléistocène du Département des Ardennes. – Bulletin de la Société d'Histoire Naturelle des Ardennes 28: 43-63.
- Bate D.M.A. (1916): On a small collection of vertebrate remains from the Har Dalam cavern, Malta, with note on a new species of the genus *Cygnus*. – Proceedings of the Zoological Society of London 1916: 421-430.
- Bate D.M.A. (1928): The animal remains. – In: Garrod D.A.E., Buxton L.H.D., Elliot Smith G. & Bate D.M.A.: Excavation of a Mousterian rock shelter at Devil's Tower, Gibraltar. – Journal of the Royal Anthropological Institute of Great Britain and Ireland 58: 92-112.
- Bauer K.M. & Glutz von Blotzheim U.N. (eds., 1966): Handbuch der Vögel Mitteleuropas. Vol. 1. – Frankfurt am Main.
- Bauer K.M. & Glutz von Blotzheim U.N. (eds., 1987): Handbuch der Vögel Mitteleuropas. Vol. 1., 2nd rev. ed. – Wiesbaden: Aula Verlag, 483 pp.
- Baumann, F. (1958): Affine Entzerrung mit einfachen optischen Mitteln. – Schweizerische Paläontologische Abhandlungen 73 (5): 17-21.
- Bayan F. (1873): Sur les plumes d'oiseaux des gypses d'Aix. – Bulletin de la Société Géologique de France (3) 1: 386-387.
- Bayer F. (1882): O nových zbytcích zkamenělých ptáků z českého útvaru třetihorního [On new remains of fossil birds from the Czech Tertiary system.] – Vesmír 12: 20-21. [In Czech.]
- Bayer F. (1883): Über zwei neue Vogelreste aus der böhmischen Tertiärformation. – Sitzungsberichte der königlichen böhmischen Gesellschaft der Wissenschaften Prag 1882: 60-65.
- Beaman M. (1994): Palearctic birds: a check-list of the birds of Europe, North Africa and Asia north of the foothills of the Himalayas. – Stonyhurst: Harrier Publications, 168 pp.
- Bechstein J.M. (1803): Ornithologisches Taschenbuch von und für Deutschland. Vol. 2. –

- Leipzig.
- Becker J.J. (1987): Neogene avian localities of North America. – Washington, D.C.: Smithsonian Institution Press, 171 pp.
- Becker L. (1849): Über Vogel-Eier im Paludinen-Kalke von Weissenau bei Mainz. – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1849: 69-72.
- Beddard F.E. (1902): The structure and classification of birds. – London.
- Bell A. (1915): Pleistocene and later bird fauna of Great Britain and Ireland. – Zoologist (4) 19: 401-412.
- Bendukidze O.G. (1971): Novyj predstavitel' semejstva Geranoididae (Aves, Gruiformes) iz èocenovykh otloženij Zajsana [New representative of the family Geranoididae (Aves, Gruiformes) from the Eocene deposits of Zajsan.] – Soobščeniya Akademii Nauk Gruzinskoj SSR 63: 749-751. [In Russian.]
- Bendukidze O.G. (1972a): Nekotorye čerty istorii tercičnykh žuravlebraznykh podotrjada Grues v Azii [Some features of the history of the Tertiary gruiforms of the suborder Grues in Asia.] – In: (no editor given): Sessija, posvjaščennaja 100-letiju so dnja roždenija akad. A. A. Borisjaka [Symposium, dedicated to the 100th anniversary of the birth of Academician A. A. Borisjak.]: 8-11. Moskva: PIN AN SSSR. [In Russian.]
- Bendukidze O.G. (1972b): O nachodke *Urmioris maraghanus* Mecq. na severnom Kavkaze [On the record of *Urmioris maraghanus* Mecq. in northern Caucasus.] – Soobščeniya Akademii Nauk Gruzinskoj SSR 66: 741-743.
- Benecke N. (1989): Zum Geschlechtsdimorphismus am Skelett mittelalterlicher Haushühner (*Gallus gallus* f. *domestica*) (Aves, Galliformes: Phasianidae). – Zoologische Abhandlungen (Dresden) 45: 23-33.
- Benecke N. (1994): Archäologische Studien zur Entwicklung der Haustierhaltung in Mitteleuropa und Südkandinavien von den Anfängen bis zum ausgehenden Mittelalter. – Schriften zur Ur- und Frühgeschichte 46: 3-451.
- Beneden P.J. van (1871): Les oiseaux de l'argille rupélienne. – Bulletins de l'Académie Royale des Sciences, Lettres et des Beaux-Arts de Belgique (2) 32: 256-261. [Pagination of the offprint is 213-218.]
- Beneden P.J. van (1872): Oiseaux de l'argile rupelienne et du crag d'Anvers. – Journal de Zoologie 1: 284-288.
- Beneden P.J. van (1873a): Paléontologie des vertébrés. In: Bemmels E. van (ed.): Patria Belgica. Vol. 1. Belgique physique: 353-376. Bruxelles: Bruylant-Christophe.
- Beneden P.J. van (1873b): Note sur un oiseau de l'argile rupélienne. – Bulletin de l'Académie Royale des Sciences, Lettres et Beaux-Arts de Belgique (2) 35: 354-357.
- Beneden P.J. van (1883): Sur quelques formes nouvelles des terrains du pays. – Bulletin de l'Académie des Sciences, Lettres et Beaux-Arts de Belgique (3) 6: 132-134.
- Bengtson S.-A. (1984): Breeding ecology and extinction of the Great Auk (*Pinguinus impennis*): anecdotal evidence and conjectures. – Auk 101: 1-12.
- Bensley B.A. (1921): An egg of "*Struthiolithus chersonensis*". – University of Toronto Studies, Biological Series 19: 1-7.
- Berendt G.C. (1845): Die organischen Bernstein-Einschlüsse im Allgemeinen. – In: Goepfert H. & Berendt G.C. (eds.): Der Bernstein und die in ihm befindlichen Pflanzenreste der Vorwelt. – In: Berendt G.C. (ed.): Die im Bernstein befindlichen organischen Reste der Vorwelt. Vol. 1 (1. Abteilung): 40-50. Berlin: Nicolai.
- Berg D.E. (1965): Nachweis des Riesenlaufvogels *Diatryma* im Eozän von Messel bei Darmstadt Hessen. – Notizblatt des Hessischen Landes-Amtes für Bodenforschung 93: 68-72.
- Berger J.-P. (1985): La transgression de la molasse marine supérieure (OMM) en Suisse occidentale. – Müncher Geowissenschaftliche Abhandlungen (A) 5: 1-208.

- Berlese A. & Leonardi G. (1896): Diagnosi di cocciniglie nuove [Diagnoses of new coccinids.] – Rivista di Patologia Vegetale 4: 345-352. [In Italian.]
- Berlin B. (1992): Ethnobiological classification: principles of categorization of plants and animals in traditional societies. – Princeton: Princeton University Press, xvii + 335 pp.
- Bessonat G. & Michaux A. (1973): Découverte d'un squelette complet d'échassier dans le Stampien provencal. – Bulletin du Muséum d'Histoire Naturelle de Marseille 33: 143-145.
- Bisseling G.A.L. (1938): Vondsten van beenderen van de Reuzenalk (*Alca impennis* L.) op Funk Island [Records of bones of the Great Auk (*Alca impennis*) on the Funk Inland.] – Natuur en Mensch 12: 265-269. [In Dutch.]
- Björklund M. (1994): Phylogenetic relationships among Charadriiformes: reanalysis of previous data. – Auk 111: 825-832.
- Blainville H.M.D. (1821): Mémoire sur l'emploi de las forme du sternum et de ses annexes, pour l'établissement ou la confirmation des familles naturelles parmi les oiseaux. – Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts 92: 185-216.
- Blanc G.A. (1927): Sulla presenza di *Alca impennis* Linn. nella formazione pleistocenica superiore di grotta Romanelli in terra d'Otranto [On the presence of *Alca impennis* Linn. in late Pleistocene deposits of the Romanelli cave in the Otranto area.] – Arch. Anthropol. Etnol. 58: 3-39. [In Italian.]
- Blanchard E. (1857): De la détermination de quelques Oiseaux fossiles et des caractères ostéologiques des Gallinacés ou Gallides. – Annales des Sciences Naturelles (4) 7: 91-105.
- Blasius W. (1884): Zur Geschichte der Überreste von *Alca impennis*. – Journal für Ornithologie 32: 58-176.
- Blasius W. (1903): Der Riesenalk, *Alca impennis*. – In: Hennicke C.R. (ed.): Naumann's Naturgeschichte der Vögel Mitteleuropas. Vol 12: 169-208. Gera.
- Blondel J. & Mourer-Chauviré C. (1998): Evolution and history of the western Palearctic avifauna. – Trends in Ecology and Evolution 13: 488-492.
- Bocheński Z. (1974): Ptaki młodszego czwartorzędu Polski [Birds of the late Quaternary of Poland.] – Warszawa: Państwowe Wydawnictwo Naukowe, 212 pp. [In Polish.]
- Bocheński Z. (1984): Middle Pleistocene remains of birds from Kozi Grzbiet in the Świętokrzyskie Mts. (Holy Cross Mts. – Central Poland). – Acta Zoologica Cracoviensia 27: 177-186.
- Bocheński Z. (1985): Osteological differentiation in Willow Grouse. – In: Duncker H.-J. & Fleischer G. (eds.): Vertebrate morphology. – Fortschritte der Zoologie 30: 69-72. Stuttgart: Gustav Fischer Verlag.
- Bocheński Z. (1987): *Miophasianus medius* (Milne-Edwards, 1869) from Przeworno (SW Poland and some general remarks on the genus *Miophasianus*. – Acta Zoologica Cracoviensia 30: 71-80.
- Bocheński Z. (1989): Ptaki - Aves [Birds – Aves.] – In: Kowalski K. (ed.), Historia i ewolucja lądowej fauny Polski [History and evolution of the terrestrial fauna of Poland.] – Folia Quaternaria 59/60: 89-108. [In Polish.]
- Bocheński Z. (1991): Pliocene grouse of the genus *Lagopus* from Poland. – Acta Zoologica Cracoviensia 34: 563-577.
- Bocheński Z. (1992): Historia sów Europy [History of European owls.] – Przegląd Zoologiczny 34: 77-90. [In Polish.]
- Bocheński Z. (1993): Catalogue of fossil and subfossil birds of Poland. – Acta Zoologica Cracoviensia 36: 329-460.
- Bocheński Z. (1995): History of herons of the Western Palearctic. – Acta Zoologica Cracoviensia 38: 343-362.
- Bocheński Z. (1996): Tertiary avian localities of Poland. – Mlíkovský J. (ed.): Tertiary

- avian localities of Europe. – *Acta Universitatis Carolinae, Geologica* 39: 693-697.
- Bocheński Z. (1997): List of European fossil bird species. – *Acta Zoologica Cracoviensia* 40: 293-333.
- Bocheński Z. & Kuročkin E.N. (1987a): New data on Pliocene phasianids (Aves: Phasianidae) of Moldavia and S. Ukraine. – *Acta Zoologica Cracoviensia* 30: 81-95.
- Bocheński Z. & Kuročkin E.N. (1987b): Pliocene bustards (Aves: Otididae and Gryzajidae) of Moldavia and S. Ukraine. – In: Mourer-Chauviré C. (ed.): *L'évolution des oiseaux d'après le témoignage des fossiles*. – *Documents des Laboratoires de Géologie de Lyon* 99: 173-187.
- Bocheński Z. & Szymczyk W. (1979): Szczątki ptaka z oligocenu Karpat Przemyskich [Remains of a bird from the Oligocene of Przemyśl Carpathians.] – In: *Badania paleontologiczne Karpat Przemyskich (Materialy IV Krajowej konferencji paleontologów)* [Paleontological research of Przemyśl Carpathians (Materials of the IVth State conference of paleontologists).] – p. 55. [In Polish.]
- Bocheński Z.M. & Tomek T. (1995): How many comparative skeletons do we need to identify a bird bone? – In: Peters D.S. (ed.): *Acta palaeornithologica*. – *Courier Forschungsinstitut Senckenberg* 181: 357-361.
- Bock W.J. (1994): History and nomenclature of avian family-group names. – *Bulletin of the American Museum of Natural History* 222: 1-281. [For a review see Olson 1995.]
- Boetticher H. von (1955): Der Blaufuß-Tölpel "Camany", *Sula nebouxii* Milne-Edwards. – *Anzeiger der Ornithologischen Gesellschaft in Bayern* 4: 375.
- Boeuf O. & Mourer-Chauviré C. (1992): Les oiseaux d'âge Pliocène de Chilhac, Haute-Loire, France. – *Comptes-Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris)* (2) 314: 1091-1096.
- Boev Z.N. (1988): Paleontologičny izsledovanija na pticite v Bălgarija [Paleontological studies of birds in Bulgaria.] – *Priroda (Sofija)* 5: 63-67. [In Bulgarian.]
- Boev Z.N. (1991): Paleontologičnoto sâkrovište na Vâršec [The paleontological treasure of Vâršec.] – *Priroda i Znanie* 1: 19-20. [In Bulgarian.]
- Boev Z.N. (1992): Paleornithology in Bulgaria. – In: Campbell K.C. (ed.): *Studies in avian paleontology honoring Pierce Brodkorb*. – *Natural History Museum of Los Angeles County, Science Series* 36: 459-463.
- Boev Z.N. (1995a): Middle Villafranchian birds from Varshets (western Balkan range - Bulgaria). – In: Peters D.S. (ed.): *Acta palaeornithologica*. – *Courier Forschungsinstitut Senckenberg* 181: 259-269.
- Boev Z.N. (1995b): On the appearance of the domestic fowl (*Gallus gallus domestica*) in Bulgaria and Balkan Peninsula and the question of domestication of junglefowls [sic!] (genus *Gallus* Brisson, 1760) in southeast Europe. – *Historia Naturalis Bulgarica* 5: 37-49.
- Boev Z.N. (1995c): Eneolithic and early Bronze Age birds from the sunken settlement at the Sozopol bay (Bulgarian Black Sea coast). – *Historia Naturalis Bulgarica* 5: 51-60.
- Boev Z.N. (1996): Tertiary avian localities of Bulgaria. – Mlíkovský J. (ed.): *Tertiary avian localities of Europe*. – *Acta Universitatis Carolinae, Geologica* 39: 541-545.
- Boev Z.N. (1997a): Vidov sâstav i tempove v evoljucijata na pticite (Aves): pregled na sâvremennite schvaščanija [Species composition and evolution rates of birds (Aves): a review of modern concepts.] – *Historia Naturalis Bulgarica* 8: 137-146. [In Bulgarian.]
- Boev Z.N. (1997b): *Chauvireria balcanica* gen. n., sp. n. (Phasianidae – Galliformes) from the Middle Villafranchian of Western Bulgaria. – *Geologica Balcanica* 27: 69-78.
- Boev Z.N. (1998a): *Actitis balcanica* sp. n. – a late Pliocene sandpiper (Aves: Scolopacidae) from Bulgaria. – *Historia Naturalis Bulgarica* 9: 71-77.
- Boev Z.N. (1998b): Late Pliocene Hawfinches (*Coccothraustes* Brisson, 1760) (Aves: Fringillidae) from Bulgaria. – *Historia Naturalis Bulgarica* 9: 87-99.
- Boev Z.N. (1998c): Presence of Bald Ibises (*Geronticus* Wagler, 1832) (Threskiornithidae

- Aves) in the late Pliocene of Bulgaria. – *Geologica Balcanica* 28: 45-52.
- Boev Z.N. (1998d): Fossil birds of Dorkovo – an early Pliocene site in the Rhodope Mts. (Southern Bulgaria). – *Geologica Balcanica* 28: 53-60.
- Boev Z.N. (1998e): Peafowls [sic!] (g. *Pavo* Linnaeus, 1758) and Ptarmigans (g. *Lagopus* Brisson, 1760): an [sic!] unique coexistence in North Bulgaria over 3 m.y. ago. – *Biogeographia* 19: 219-222.
- Boev Z.N. (1998f): Pticitе prez neogen-kvarternata [Birds in the Neogene and Quaternary.] – *Priroda* (Sofija) 1998 (1-2): 57-67. [In Bulgarian.]
- Boev Z.N. (1999a): Neogenski i kvaternerni pticy (Aves) ot Bălgarija [Neogene and Quaternary birds (Aves) of Bulgaria.] – Unpub. thesis, Bulgarian Academy of Sciences, Sofija, 243 + 243 pp. [In Bulgarian.]
- Boev Z.N. (1999b): The late Pleistocene avifauna of Cave No 16 (NW Bulgaria) and its palaeoecological implication – In: 8th international congress on the zoogeography and ecology of Greece and adjacent regions, Abstracts: 21-22. Athene: Hellenic Zoological Society.
- Boev Z.N. (1999c): *Falco bakalovi* sp. n. – a late Pliocene falcon (Falconidae, Aves) from Varshets (W Bulgaria). – *Geologica Balcanica* 29: 131-135.
- Boev Z.N. (1999d): On the presence of *Tetrao partium* (Kretzoi, 1962) (Aves: Tetraonidae) in the late Pliocene of Bulgaria. – *Historia Naturalis Bulgarica* 10: 85-96.
- Boev Z.N. (1999e): Late Pliocene bustards (Aves: Otidae) from western Bulgaria. – *Historia Naturalis Bulgarica* 10: 97-108.
- Boev Z.N. (1999f): *Regulus bulgaricus* sp. n. – the first fossil kinglet (Aves: Sylviidae) from the late Pliocene of Varshets, western Bulgaria. – *Historia Naturalis Bulgarica* 10: 109-115.
- Boev Z.N. (1999g): *Gallinula balcanica* sp. n. (Rallidae: Gruiformes) – a middle Villafranchian moorhen from western Bulgaria. – *Acta Zoologica Bulgarica* 51: 43-47.
- Boev Z.N. (1999h): Earliest finds of crossbills (genus *Loxia*) (Aves: Fringillidae) from Varshets (NW Bulgaria). – *Geologica Balcanica* 29(3-4): 51-57.
- Boev Z.N. (2000a): Neogene avifaunas of Bulgaria. – *Vertebrata Palasiatica* 38 (Supplement): 2-3. [Abstract.]
- Boev Z.N. (2000b): *Cygnus verae* sp. n. (Anseriformes: Anatidae) from the early Pliocene of Sofia (Bulgaria). – *Acta Zoologica Cracoviensia* 43: 185-192.
- Boev Z.N. (2000c): Early Pleistocene and early Holocene avifauna of the Cherdzhenitsa cave, northwestern Bulgaria. – *Historia Naturalis Bulgarica* 11: 107-116.
- Boev Z.N. (2000d): A Lower Miocene gull (Larinae gen. indet.) (Laridae Vigors, 1825 – Charadriiformes Huxley, 1867) from Hrabarsko (W Bulgaria). – *Geologica Balcanica* 30(1-2): 69-75.
- Boev Z.N. (2000e): The presence of *Apus baranensis* Janossy, 1977, (Aves: Apodidae) in the late Pliocene of Bulgaria. – *Acta Zoologica Bulgarica* 52: 43-52.
- Boev Z.N. (2000f): Additional material of *Geronticus balcanicus* Boev, 1998, and precision of the age of the type locality. – *Acta Zoologica Bulgarica* 52: 53-58.
- Boev Z.N. (2000g): Late Pleistocene avifauna of the Razhishkata Cave, western Bulgaria. – *Historia Naturalis Bulgarica* 12: 71-87.
- Boev Z.N. (2001): Early Pliocene avifauna of Muselievo (C Northern Bulgaria). – *Acta Zoologica Cracoviensia* 44: 37-52.
- Boev Z.N. & Karajvanova E. (1998): *Fulica atra pontica* subsp. n. from the Middle Holocene on the south Black Sea coast, Bulgaria. – *Historia Naturalis Bulgarica* 9: 53-69.
- Boev Z.N. & Koufos G. (2000): Presence of *Pavo bravardi* (Gervais, 1849) (Aves, Phasianidae) in the Ruscinian locality of Megalo Emvolon, Macedonia, Greece. – *Geologica Balcanica* 30(1-2): 77-82.
- Boev Z.N. & Kovačev D. (1998): *Buteo spassovi* sp. n. – a late Miocene buzzard

- (Accipitridae, Aves) from SW Bulgaria. – *Geologica Balcanica* 29: 125-129.
- Böhm G. (1896): Tierfährten im Tertiär des Badischen Oberlandes. – In: Freiburger Universitäts-Festprogramm zum 70. Geburtstag des Grossherzogs Friedrich: 227-238. Freiburg.
- Böhm G. (1898): Ueber das fossile Trittpaar im Tertiär des badischen Oberlandes. – *Zeitschrift der Deutschen Geologischen Gesellschaft* 50: 204-206.
- Boles W.E. (1995): The world's oldest songbird. – *Nature (London)* 374: 21-22.
- Boles W.E. (1997): Fossil songbirds (Passeriformes) from the early Eocene of Australia. – *Emu* 97: 43-50.
- Bonaparte C.L. (1850): *Conspectus generum avium*. Vol. 1. – Leyden: E. J. Brill, 543 pp.
- Bonaparte C.L. (1854): Notes ornithologiques sur les collections rapportées en 1853, par M. A. Delattre, de son voyage en Californie et dans le Nicaragua (cont.). – *Comptes Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris)* 38: 1-11, 53-66, 258-266, 378-389, 533-541, 650-665.
- Bonaparte C.L. (1856a): Ornithologie fossile servant d'introduction au Tableau comparatif des Ineptes et des Autruches. – *Comptes-Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris)* 43: 775-783.
- Bonaparte C.L. (1856b): *Conspectus Ineptorum et Struthiorum*. – *Comptes-Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris)* 43: 840-841.
- Bonaparte C.L. (1856c): Additions et corrections aux tableaux paralléliques de la deuxième sous-classe des oiseaux praecoces ou autophages. – *Comptes-Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris)* 43: 1017-1027.
- Bonifay E., Bassiakos Y., Bonifay M.-F., Louchart A., Mourer-Chauviré C., Pereira E., Quinif Y. & Salotti M. (1998): La grotte de la Coscia (Rogliano, Macinaggio): Étude préliminaire d'un nouveau site du Pléistocène supérieur de Corse. – *Paleo* 10: 17-41.
- Bonnaterre J. (1790): *Tableau encyclopédique et méthodique des trois règnes de la Nature. Ornithologie*. Vol. 1. – Paris: Mme veuve Agassi, xcvi + 402 pp.
- Bosák P., Głazek J., Horáček I. & Szyrkiewicz A. (1982): New locality of early Pleistocene vertebrates – Żabia cave at Podlesice, central Poland. – *Acta Geologica Polonica* 32: 217-226.
- Bosniaski S. de (1879): Nuove scoperte paleontologiche [New paleontological discoveries.] – *Atti Società Toscana Scienze Naturali, Processi Verballi*, 9. XI. 1879, p. 13. [In Italian.]
- Boule M. (1919): Les grottes de Grimaldi (Baoussé-Rousse). Vol. 1(4). *Géologie et paléontologie (Fin)*: 237-362. – Monaco. [Aves: pp. 299-331, pls. xxxvi-xli.]
- Boule M. & Villeneuve L. (1927): La grotte de l'Observatoire à Monaco. – *Arch. Inst. Paléontologie Humaine* 1: 1-115.
- Bowerbank J.S. (1852): On the remains of a gigantic bird from the London Clay of Sheppey. – *Reports of the British Association for the Advancement of Science* 21: 55.
- Bowerbank J.S. (1854): On the remains of a gigantic bird (*Lithornis emuinus*) from the London Clay of Sheppey. – *Annals and Magazine of Natural History* (2) 14: 263-264.
- Bowler P. J. (1976): Fossils and progress. *Palaeontology and the idea of progressive evolution in the nineteenth century*. – New York: Neale Watson, viii + 191 pp.
- Bradley F. & Landini W. (1984) I fossili del "tripoli" messiniano di Gabbro (Livorno) [Fossils from the Messinian "tripoli" of Gabbro.] – *Palaeontographica Italica* 73: 5-33.
- Bräm H. (1954): Fährten von Wirbeltieren aus der subalpinen Molasse des Bergsturzgebietes von Goldau. – *Eclogae Geologicae Helvetiae* 85: 399-431.
- Brandt A.F. (1873): Über ein grosses fossiles Vogelei aus der Umgegend von Cherson. – *Bulletin de la Académie Imperiale des Sciences (St. Petersburg)* 18: 158-162.
- Brandt A.F. (1874): On a large fossil egg from neighbourhood of Cherson. – *Ibis* 1874: 4-7. [Extract from Brandt 1873.]
- Brandt A.F. (1885): Über das Schicksaal des Eies von *Struthiolithus chersonensis*. –

- Zoologischer Anzeiger 8: 191-192.
- Bräuhäuser M. (1922): Begleitworte zur geognostischen Spezialkarte von Württemberg. – Atlasblatt Kirchheim. 3rd ed.
- Brehm C.L. (1831): Handbuch der Naturgeschichte aller Vögel Deutschlands. – Ilmenau: F. Voigh, xxiv + 1088 pp.
- Breuil H. & Bégouën H. (1936): Quelques oiseaux inédits ou méconnus de l'art préhistorique. – Congrès Préhistorique de France 12: 475-488.
- Breuil H., Capitan L., Bourrinet P. & Peyrony D. (1909): Observations sur un baton de commandement orné de figures animales et de personnage semi-humains. – Revue de l'École d'Anthropologie 19: 62-76.
- Brisson M.J. (1760): Ornithologia. Vol. 6. – Paris: Bouche, 543 + xlvi + 146 + xxiii pp.
- Brochon E.H. (1880): Un *Pelagornis* dans la molasse fossilifère de Léognan?. – Actes de la Société Linnéenne de Bordeaux 35: xx.
- Brodkorb P. (1952): The types of Lambrecht's fossil bird genera. – Condor 54: 174-175.
- Brodkorb P. (1953): A review of the Pliocene loons. – Condor 55: 211-214.
- Brodkorb P. (1960a): How many species of birds have existed? – Bulletin of the Florida State Museum, Biological Sciences 5: 41-53.
- Brodkorb P. (1960b): Great Auk and Common Murre from a Florida midden. – Auk 77: 342-343.
- Brodkorb P. (1961): Birds from the Pliocene of Juntura, Oregon. – Journal of the Florida Academy of Sciences 24: 169-184.
- Brodkorb P. (1962a): The systematic position of two Oligocene birds from Belgium. – Auk 79: 706-707.
- Brodkorb P. (1962b): A teal from the Lower Pliocene of Kansas. – Quarterly Journal of the Florida Academy of Sciences 25: 157-160.
- Brodkorb P. (1963a): A giant flightless bird from the Pleistocene of Florida. – Auk 80: 111-115.
- Brodkorb P. (1963b): The name of a fossil rail and its date of publication. – Auk 80: 542.
- Brodkorb P. (1963c): Catalogue of fossil birds: Part 1 (Archaeopterygiformes through Ardeiformes). – Bulletin of the Florida State Museum, Biological Sciences 7: 179-293.
- Brodkorb P. (1963d): Miocene birds from the Hawthorne Formation in Florida. – Quarterly Journal of the Florida Academy of Sciences 26: 159-167.
- Brodkorb P. (1964): Catalogue of fossil birds: Part 2 (Anseriformes through Galliformes). – Bulletin of the Florida State Museum, Biological Sciences 8: 195-335.
- Brodkorb P. (1965): New taxa of fossil birds. – Quarterly Journal of the Florida Academy of Sciences 28: 197-198.
- Brodkorb P. (1967): Catalogue of fossil birds: Part 3 (Ralliformes, Ichthyornithiformes, Charadriiformes). – Bulletin of the Florida State Museum, Biological Sciences 11: 99-220.
- Brodkorb P. (1970): Two fossil owls from the Aquitanian of France. – Quarterly Journal of the Florida Academy of Sciences 32: 159-160.
- Brodkorb P. (1971a): The paleospecies of woodpeckers. – Quarterly Journal of the Florida Academy of Sciences 33: 132-136.
- Brodkorb P. (1971b): Catalogue of fossil birds: Part 4 (Columbiformes through Piciformes). – Bulletin of the Florida State Museum, Biological Sciences 15: 163-266.
- Brodkorb P. (1978): Catalogue of fossil birds: Part 5 (Passeriformes). – Bulletin of the Florida State Museum, Biological Sciences 23: 139-228.
- Brodkorb P. (1980): A new fossil heron (Aves: Ardeidae) from the Omo basin of Ethiopia, with remarks on the position of some other species assigned to the Ardeidae. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Hildegard Howard. – Natural History Museum of Los Angeles County, Contributions in Science 330: 87-92.

- Brodkorb P. & Mourer-Chauviré C. (1982): Fossil aningas (Aves: Anhingidae) from early Man sites of Hadar and Omo (Ethiopia) and Olduvai Gorge (Tanzania). – *Géobios* 15: 505-514.
- Bronn H.G. (1834): Ornitholithes. – In: Ersch J.S. & Gruber J.G. (eds.): Allgemeine Encyclopädie der Wissenschaften und Künste. Vol. (3) 5: 463-469. Leipzig.
- Bruijn H. de, Daams R., Daxner-Höck G., Fahlsbusch V., Ginsburg L., Mein P. & Morales J. (1992): Report of the RCMNS working group on fossil mammals, Reisenburg 1990. – *Newsletter on Stratigraphy* 26: 65-118.
- Brunet J. (1961): Oiseaux. – In: Le gisement de Vertébrés miocènes de Beni Mellal (Maroc). – *Notes et Mémoires du Service Géologique (Maroc)* 155: 105-108.
- Brunet J. (1970): Oiseaux de l'Eocène supérieur du bassin de Paris. – *Annales de Paléontologie, Vertébrés*, 56: 1-57.
- Brunner G. (1933): Eine präglaziale Fauna aus dem Windloch bei Sackdilling (Oberpfalz). – *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie (B)* 71: 303-327.
- Brunner G. (1958): Nachtrag zur Breitenberghöhle bei Gössweinstein/Ofr. – *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1958: 500-517.
- Brünnich M.T. (1772): *Zoologiae fundamenta praelectionibus academicis accomodata* [Fundamentals of zoology.] – Copenhagen, 253 pp. [In Latin.]
- Buckland W. (1822): Account of an assemblage of fossil teeth and bones of elephant, rhinoceros, hippopotamus, bear, tiger and hyaena, and sixteen other animals; discovered in a cave at Kirkdale, Yorkshire, in the year 1821; with a comparative view of five similar caverns. – *Philosophical Transactions of the Royal Society of London* 112: 171-236.
- Buckland, W. (1823): *Reliquiae Diluvianae; or observations on the organic Remains contained in Caves, Fissures, and diluvial Gravel, and on other geological Phenomena attesting the action of an universal Deluge.* – London, vii + 303 pp.
- Buffetaut E. (1997a): New remains of the giant bird *Gastornis* from the Upper Paleocene of the eastern Paris Basin and the relationships between *Gastornis* and *Diatryma*. – *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1997 (3): 179-190.
- Buffetaut E. (1997b): The unfinished story of the early Tertiary giant bird *Gastornis*. – *Geological Society of Denmark, Online Series 1: unpaginated.* [<http://home4.inet.tele.dk/dgftth/online/buffetaut.htm>]
- Buffetaut E. (1997c): L'oiseau géant *Gastornis*: interprétation, reconstitution et vulgarisation de fossiles inhabituels dans la France du XIX^e siècle. – *Bulletin de la Société Géologique de France* 168: 805-811.
- Buffetaut E. & Rage J.-C. (1982): Observations à la note de C. Mourer-Chauviré "Première indication de la présence de Phorurhacids, famille d'oiseaux géants d'Amérique du Sud, dans le Tertiaire européen: *Ameghinornis* nov. gen. (Aves, Ralliformes) des Phosphorites du Quercy, France". *Geobios*, 1981, 14-5, p. 637-646. – *Géobios* 15: 267-269.
- Bullen R.P. & Sleight F.W. (1959): Archaeological investigations of the Castle Windy midden, Florida. – William L. Bryant Foundation for American Studies, Report 1: i-vi + 1-32.
- Burčák-Abramovič N.I. (1939): Novye dannye o tretičnyh strausach juga Ukrainy [New data on the Tertiary ostriches of southern Ukraine.] – *Priroda (Moskva)* 28: 94-97. [In Russian.]
- Burčák-Abramovič N.I. (1949): Iskopaemyj straus (*Struthio brachydactylus* sp. nov.) iz meotičeskich otloženij s. Grebennikov na jube Ukrainy [A fossil ostrich (*Struthio brachydactylus* sp. nov.) from the Meotian deposits of Grebenniki village in the south of Ukraine.] – *Doklady Akademii Nauk SSSR* 67: 141-144. [In Russian.]

- Burčak-Abramovič N.I. (1951): *Urmiornis (Urmiornis maraghanus* Mecq.), strausopodobnaja ptica gipparionovoj fauny Zakavkaz'ja i južnoj Ukrainy [Urmiornis (*Urmiornis maraghanus* Mecq.), an ostrich-like bird of the *Hipparion* fauna of Transcaucasia and southern Ukraine.] – *Izvestija Akademii Nauk Azerbajdžanskoj SSR* 1951 (6): 83-94. [In Russian.]
- Burčak-Abramovič N.I. (1953a): Iskopaemye strausy Kavkaza i juga Ukrainy [Fossil ostriches of Caucasus and southern Ukraine.] – *Trudy Estestvenno-Istoričeskogo Muzeja Imeni G. Zardabi* 7: 2-206. [In Russian.]
- Burčak-Abramovič N.I. (1953b): Nachodki iskopaemych strausov [Finds of fossil ostriches.] – *Priroda (Moskva)* 1953 (6): 101-103. [In Russian.]
- Burčak-Abramovič N.I. (1958): Tretičnye pticy SSSR [Tertiary birds of the USSR.] – *Učenyje Zapiski Azerbajdžanskogo Gosudarstvennogo Universiteta Imeni S. M. Kirova, Biologija* 1958 (1): 81-88. [In Russian.]
- Burčak-Abramovič N.I. (1962): [Materials to the knowledge of fossil ostriches of the Old World. Part I. (Review of the literature, history of research, main bibliography).] – *Vertebrata Palasiatica* 6: 299-309. [In Chinese.]
- Burčak-Abramovič N.I. (1966): Pticy verchnepaleolitičeskoj stojanki peščery Gvardžilas-Kldé v Imeretii [Birds of the late Paleolithic site in Gvardžidlas-Kldé cave in Imeretia.] – In: *Kudarskie peščernye paleolitičeskije stojanki v Jugo-Osetii. – Peščery Gruzii* 4: 93-110. Moskva: Kommissija po Izučeniju Četvertičnogo Perioda AN SSSR. [In Russian.]
- Burčak-Abramovič N.I. (1972): O pochodzeniu kur domowych Kaukazu na podstavie danyh paleontologičnyh i archeologičnyh [On the origin of domestic chicken in Caucasus on the basis of paleontological and archaeological data.] – *Przegľad Zoologiczny* 16: 427-431. [In Polish.]
- Burčak-Abramovič N.I. (1975): Die pleistozäne Vogelfauna der UdSSR. – *Quartärpaläontologie* 1: 87-105.
- Burčak-Abramovič N.I. (1980): Ostatki ptic iz peščery Kudaro I [Avian remains from Kudaro I cave.] – In: Ivanova I.K. & Černjachovskij A.G. (eds.): *Kudarskie peščernye paleolitičeskije stojanki v Jugo-Osetii* [Paleolithic sites of the Kudaro caves in South Ossetia.]: 98-110. Moskva: Nauka. [In Russian.]
- Burčak-Abramovič N.I. (1996): [Untitled report.] – *Society of Avian Paleontology and Evolution, Information Letter* 10: 8-10.
- Burčak-Abramovič N.I. & Aliev S.D. (1989): Iskopaemaja ornitofauna paleolitičeskoj stojanki Azychskoj peščery na Malom Kavkaze v Azerbajdžane. I soobščenie [Fossil ornithofauna of the Paleolithic site Azychskaja cave in Little Caucasus in Azerbaidžan. 1st communication.] – In: *Materialy po ekologii životnyh Azerbajdžana* [Materials on the ecology of the animals of Azerbaijan]: 72-80. Baku: Baku Universiteti Nešrijaty. [In Russian.]
- Burčak-Abramovič N.I. & Bendukidze O.G. (1971): O proischoždenii domašnich kur Gruzii [On the origin of domestic chicken in Georgia.] – *Soobščeniya Akademii Nauk Gruzinskoj SSR* 61: 497-500. [In Russian.]
- Burčak-Abramovič N.I. & Kon'kova N.I. (1967): Nachodki iskopaemych strausov v MSSR i drugih punktach SSSR [Finds of fossil ostriches in Moldovan SSR and other sites of the USSR.] – *Paleontologija, Geologija i Poleznye Iskopaemye Moldavii* 2: 146-156. [In Russian.]
- Burčak-Abramovič N.I. & Lakerbaj L.B. (1989): Kurinye (Galliformes (Temminck)) v plejstocenovoj i golocenovoj faune Abchazii, Kavkaza i SSSR. Paleoekologičeskaja obstanovka ich obitanija [Gamefowl (Galliformes (Temminck)) in the Pleistocene and Holocene faunas of Abkhazia, Caucasus and the USSR. Paleoecological conditions of their occurrence.] – *Trudy Abchazskogo Zoologičeskogo Muzeja (Suchumi)* 5: 166-188. [In Russian.]

- Burčák-Abramovič N.I. & Mežlumjan S. (1986): K vidovomu sostavu, rasprostraneniiju i ispol'zovaniju ptic v golocene [On the species composition, distribution and exploitation of birds in the Holocene.] – Zoologičeskij Sbornik (Erevan) 20: 128-149. [In Russian.]
- Burčák-Abramovič N.I. & Nikolov I. (1984): Iskopaemye pticy Phalacrocorax serdicensis sp.n. i Anser thraceiensis sp.n. iz Bolgarii [Fossil birds Phalacrocorax serdicensis sp.n. and Anser thraceiensis sp.n. from Bulgaria.] – Paleontologija, Stratigrafija i Litologija 19: 23-27. [In Russian.]
- Burčák-Abramovič N.I. & Vekua A.K. (1971): The fossil ostrich from the Akchagil layers of Georgia. – Acta Zoologica Cracoviensia 16: 1-26.
- Burčák-Abramovič N.I., Ganea I.M. & Šušpanov K.I. (1993): [New Gallus species from Upper Pliocene in Moldova.] – Buletinul Academiei de Științe a Republicii Moldova 265 (= Științe Biologice și Chimice 4): 45-48, 77. [In Moldovan.]
- Burness G.P. & Montevecchi W.A. (1992): Oceanographic-related variation in the bone sizes of extinct Great Auk. – Polar Biology 11: 545-551.
- Cailleux A. (1969): Observations sur des gastrolites d'Oiseaux holocènes et tertiaires. – Comptes-Rendus de la Société Géologique de France 1969 (3): 73-75.
- Calder, W.A., III. (1984): Size, function and life history. – Cambridge, Massachusetts: Harvard University Press, 431 pp.
- Camp W.H. & Gilly C.L. (1943): The structure and origin of species. – Brittonia 4: 323-385.
- Campbell K.E. & Marcus L. (1992): The relationship of hindlimb bone dimensions to body weight in birds. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County 36: 395-412.
- Camper P. (1786): Conjectures relative to the petrifications found in St. Peter's Mountain near Maestricht. – Philosophical Transactions of the Royal Society of London 76: 443-456. [fide Cuvier 1822]
- Čapek V. (1917): A püspekfürdoi praeglacialis madárfauna [A preglacial avifauna from Püspökfürdő.] – Barlangkutató 5: 25-32. [In Hungarian, with a German translation on pp. 66-77 of the same volume.]
- Čapek V. (1925): Diluvální fauna od Holubic [A Diluvial fauna from the vicinity of Holubice.] – Časopis Moravského Zemského Musea 22-23: 159-178. [In Czech.]
- Capitan L., Breuil H., Bourrinet A. & Peyrony D. (1906): L'abri Mège, une station magdalénienne à Teyjat. – Revue École Antropologie Paris 16: 196-212.
- Casanovas M.L. & Santafé J.V. (1982): Icnofauna oligocena de Agramunt (Lleida, España) [An Oligocene ichnofauna from Argamunt (Lleida, Spain).] – Acta Geológica Hispánica 17: 113-119. [In Spanish.]
- Cassoli P.F. (1980): L'avifauna de Pleistocene superiore delle Arene Candide (Liguria) [Avifauna from the late Pleistocene of Arene Candide (Liguria).] – Memorie Istituto Italiano Paleontologia Umana (n.s.) 3: 155-234. [In Italian.]
- Cassoli P.F. (1992): Avifauna del Pleistocene superiore delle Arene Candide, Praia e Grotta Romanelli (Italia). – Quaternaria Nova 2: 239-246.
- Cassoli P.F. & Segre A.G. (1985): L'Alca impenne *Alca impennis* del Pleistocene d'Italia [The Great Auk *Alca impennis* from the Pleistocene of Italy.] – In: Fasola M. (ed.): Atti III convegno italiano di ornitologia [Proceedings of the IIIrd Italian meeting on ornithology]: 251-254. Pavia: Università di Pavia. [In Italian.]
- Cassoli P.F. & Tagliacozzi A. (1997): Butchering and cooking of birds in the Palaeolithic site of Grotta Romanelli (Italy). – International Journal of Osteoarchaeology 7: 303-320.
- Cassoli P., Segre A.G. & Segre E. (1977): Evolution morphologique et écologique de la côte de Castro (Pouilles) dans le Pléistocène final. – In: Sonnevile-Bordes D. (ed.): La fin des temps glaciaires en Europe. – Colloques Internationaux de CNRS 271: 325-332.
- Cavallo O. & Gaudant J. (1984): Il Messiniano dell'Albese [The Messinian of the Alba

- region.] – *Alba Pompeia* (n. s.) 5 (1): 5-20. [In Italian.]
- Cavallo O., Macagno M. & Pavia G. (1986): Fossili dell'Albese [Fossils of the Alba region.] – *Alba: Famija Albèsia*, 223 pp. [In Italian.]
- Čerkašenko M.I. & Ljaščuk B.F. (1960): Znachidka jajca vykopnogo ptaka na Stanislavščini [Find of an egg of a fossil bird in the Stanislavsk region.] – In: Konferencija po vyvčennju flory i fauny Karpat ta prileglic teritorij, Tezisi Dopovidij [Conference on the study of flora and fauna of the Carpathians and adjacent territories, Proceedings]. L'viv. [In Ukrainian.]
- Chandler R.M. (1990): Fossil birds of the San Diego Formation, late Pliocene, Blacan, San Diego County, California. – In: Olson S.L. (ed.): *Recent advances in the study of Neogene fossil birds.* – *Ornithological Monographs* 44: 73-161. Washington, D.C.: American Ornithologists' Union.
- Cheneval J. (1979): Les oiseaux, genre *Anas*, du gisement aquitain de Saint-Gérard-le-Puy (Allier). – Unpublished Thesis, Université Claude-Bernard, Lyon.
- Cheneval J. (1983a): Révision de l'avifaune du gisement aquitain de Saint-Gérard-le-Puy (Allier, France) (Gaviiformes à Anseriformes). – Unpublished Thesis (Nr. 1380), Université Claude-Bernard, Lyon, 220 pp.
- Cheneval J. (1983b): Les Anatidae (Aves, Anseriformes) du gisement aquitain de Saint-Gérard-le-Puy (Allier, France). – In: Buffeteau E., Mazin J.M. & Salmon E. (eds.): *Actes du symposium paléontologique Georges Cuvier*: 85-98. Montbéliard: Ville de Montbéliard.
- Cheneval J. (1983c): Révision du genre *Palaelodus* Milne-Edwards, 1863 (Aves, Phoenicopteridae) du gisement aquitain de Saint-Gérard-le-Puy (Allier, France). – *Géobios* 16: 179-191.
- Cheneval J. (1984a): Les oiseaux aquatiques (Gaviiformes à Anseriformes) du gisement aquitain de Saint-Gérard-le-Puy (Allier, France): révision systématique. – *Palaeovertebrata* 14: 33-115.
- Cheneval J. (1984b): *Grallavis edwardsi* (Lydekker, 1891), nouveau genre d'oiseau (Ciconiiformes) du gisement aquitain de Saint-Gérard-le-Puy (Allier, France). – *Bulletin de la Société Linnéenne de Lyon* 2: 43-60.
- Cheneval J. (1987): Les Anatidae (Aves, Anseriformes) du Miocène de France. Révision systématique et évolution. – In: Mourer-Chauviré C. (ed.): *L'évolution des oiseaux d'après le témoignage des fossiles.* – *Documents du Laboratoire de Géologie de Lyon* 99: 137-157.
- Cheneval J. (1989): Fossil bird study, and paleoecological consequences: example from the Saint-Gérard-le-Puy deposits (Lower Miocene, Allier, France). – *Palaeogeography, Palaeoclimatology, Palaeoecology* 73: 295-309.
- Cheneval J. (1993): Un oiseau (Aves, Anseriformes, Anatidae) du gisement du Messinien continental de Cherasco (Province de Cuneo, Piémont, Italie). – *Rivista Piemontese di Storia Naturale* 14: 23-32.
- Cheneval J. (1995): A fossil shearwater (Aves: Procellariiformes) from the Upper Oligocene of France and the Lower Miocene of Germany. – In: Peters D.S. (eds.): *Acta palaeornithologica.* – *Courier Forschungsinstitut Senckenberg* 181: 187-198.
- Cheneval J. (1996a): Tertiary avian localities of Belgium. – Mlíkovský J. (ed.): *Tertiary avian localities of Europe.* – *Acta Universitatis Carolinae, Geologica* 39: 535-540.
- Cheneval J. (1996b): Miocene avian localities of France. – Mlíkovský J. (ed.): *Tertiary avian localities of Europe.* – *Acta Universitatis Carolinae, Geologica* 39: 599-611.
- Cheneval J. (2000): L'avifaune de Sansan. In: Ginsburg L. (ed.): *La faune miocène de Sansan (Gers) et son environnement.* – *Mémoires du Muséum National d'Histoire Naturelle (Paris)* 183: 321-388.
- Cheneval J. & Adrover R. (1993): L'avifaune du Miocène supérieur d'Aljezar B (Los Aljezars, Province de Teruel, Espagne). *Systématique et paléoécologie.* –

- Paleontologia i Evolució 26/27: 133-144.
- Cheneval J. & Escuililé F. (1992): New data concerning *Palaelodus ambiguus* (Aves: Phoenicopteriformes: Palaelodidae): ecological and evolutionary interpretations. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County, Science Series 36: 209-224.
- Cheneval J. & Pharisat A. (1995): Contribution à l'avifaune et à la taphonomie et au paleoenvironment du gisement rupelien de Froidefontaine (Territoire de Belfort, France). – In: Gayet M. & Courtinat, B. (eds.): First European paleontological congress. – Géobios, Memoire Special 18: 87-92.
- Chesser R. T. & R. M. Zink, 1994: Modes of speciation in birds: a test of Lynch's method. – Evolution 48: 490-497.
- Chozackij L.I. & Kuročkin E.N. (1966): Novye dannye o pticach pliocena Moldavii [New data on the birds of the Pliocene of Moldova.] – In: Materialy VI Pribaltskoj Ornitologičeskoj Konferencii [Materials of the 6th circum-Baltic ornithological conference]: 153-155. Vilnius. [In Russian.]
- Chozackij L.I. & Kuročkin E.N. (1977): Materialy k poznaniju pliocenovej ornitofauny Moldavii [Materials to the knowledge of the Pliocene avifauna of Moldova.] – In: Sedmaja vsesojuznaja ornitologičeskaja konferencija, Tezisi dokladov [7th all-union ornithological conference, Abstracts.] Vol 1: 31-33. Kyjiv: Naukova Dumka.
- Christol J. de (1835): Comparaison de la population contemporaine des mammifères de deux bassins tertiaires du département de l'Hérault. – Annales des Sciences Naturelles 4 (2): 193-238.
- Chu P.C. (1995): Phylogenetic reanalysis of Strauch's osteological data set for the Charadriiformes. – Condor 97: 174-196.
- Cirić A. (1957): Die Pikermifauna aus der Umgebung von Titov Veles. – Bulletin du Muséum d'Histoire Naturelle du Pays Serbe (A) 8: 1-82.
- Claußen M. (1995): Ein Vogelknochen aus dem miozänen Glimmerton von Gross-Pampau. – Geschiebe-Sammler 28: 11-12. [Undistinct figure does not resemble an avian bone.]
- Cleere H. (1992): [Editorial correction of Clottes et al. 1992.] – Antiquity 66: 828-829. [In English.]
- Clercq S.W. de & Holst H.K.K. (1971): Footprints of birds and sedimentary structures from the subalpine molasse near Flühli (Canton of Luzern). – Eclogae Geologicae Helveticae 64: 63-69.
- Clot A. & Mourer-Chauviré C. (1986): Inventaire systématique des oiseaux quaternaires des Pyrénées Françaises. – Munibe 38: 171-184.
- Clot A., Chaline J., Jammot D., Mourer-Chauviré C. & Rage J.C. (1976a): Les pôches ossifères du Pléistocène moyen et inférieur de Montoussé (Hautes-Pyrénées). – Bulletin de la Société d'Histoire Naturelle de Toulouse 112: 1-15.
- Clot A., Chaline J., Heintz E., Jammot D., Mourer-Chauviré C. & Rage J.C. (1976b): Montoussé 5 (Hautes-Pyrénées), un nouveau remplissage de fissure à faune de vertébrés du Pléistocène inférieur. – Géobios 9: 511-514.
- Clottes J. & Courtin J. (1993): Stone age gallery by the sea. – Archaeology 1993 (May/June): 38-43.
- Clottes J., Beltran A., Courtin J. & Cosquer H. (1992): La Grotte Cosquer (Cap Morgiou, Marseille). – Bulletin de la Société Préhistorique Française 89: 98-128. [Abbreviated English translation: The Cosquer Cave on Cape Morgiou, Marseilles. – Antiquity 66: 583-589, 1992.]
- Coiter V. (1575): De avium sceletis et praecipuis musculis [On the skeleton and main muscles of birds.] – Padova. [In Latin.]
- Collins C.T. (1976a): Two new species of *Aegialornis* from France, with comments on the ordinal affinities of the Aegialornithidae. – In: Olson S.L. (ed.): Collected papers in

- avian paleontology honoring the 90th birthday of Alexander Wetmore. – *Smithsonian Contributions to Paleobiology* 27: 121-127.
- Collins C.T. (1976b): A review of the Lower Miocene swifts (Aves: Apodidae). – In: Olson S.L. (ed.): *Collected papers in avian paleontology honoring the 90th birthday of Alexander Wetmore.* – *Smithsonian Contributions to Paleobiology* 27: 129-132.
- Comaschi Caria I. (1969): Fossili marini e continentali del Quaternario della Sardegna [Marine and continental fossils from the Quaternary of Sardinia.] – *Atti del X *Congr. Intern. Studi Sardi*: 140-229. [In Italian.]
- Comaschi Caria I. (1970): Nuova segnalazione di resti di scimmia nel Quaternario della Sardegna [New data on the remains of monkeys from the Quaternary of Sardinia.] – *Publ. Istituto Geol. Paleont. Univ. Cagliari* 8 (83): 1-7. [In Italian.]
- Cope E.D. (1876): On a gigantic bird from the Eocene of New Mexico. – *Proceedings of the Academy of Natural Sciences of Philadelphia* 28: 10-11.
- Cope E.D. (1894): On *Cyphornis*, an extinct genus of birds. – *Journal of the Academy of Natural Sciences of Philadelphia* (2) 9: 449-452.
- Cordy J.-M. (1993): La grotte de la Belle-Roche (Sprimont, Province de Liège): un gisement paléontologique et archéologique d'exception au Benelux. – *Bull. Classe Sciences* (6) 4: 165-186.
- Coues E. (1884): *Key to North American birds*. 2nd ed. – Boston: Estes and Lauriat.
- Cracraft J. (1972): A new Cretaceous charadriiform family. – *Auk* 89: 36-46.
- Cracraft J. (1973): Systematics and Evolution of the Gruiformes (Class Aves). 3. Phylogeny of the Suborder Grues. – *Bulletin of the American Museum of Natural History* 151: 1-127. [For a review see Olson 1974b.]
- Cracraft J. (1974): Phylogeny and evolution of the Ratite birds. – *Ibis* 116: 494-521.
- Cracraft J. (1980): Phylogenetic theory and methodology in avian paleontology: a critical appraisal. – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Hildegard Howard.* – *Natural History Museum of Los Angeles County, Contributions in Science* 330: 9-16.
- Cracraft J. (1981a): Toward a phylogenetic classification of the Recent birds of the world (class Aves). – *Auk* 98: 681-714. [For comments see Olson 1982.]
- Cracraft J. (1981b): [Review of Feduccia 1980.] – *Systematic Zoology* 30: 219-221.
- Cracraft J. & Rich P.V. (1972): The systematics and evolution of the Cathartidae in the Old World Tertiary. – *Condor* 74: 272-283.
- Cramp S. (ed., 1989): *The birds of the western Palearctic*. Vol. 4. Terns to woodpeckers. – Oxford: Oxford University Press.
- Cramp S. & Simmons K.E.L. (eds., 1977): *The birds of the western Palearctic*. Vol. 1. Ostrich to ducks. – Oxford: Oxford University Press, vii + 722 pp.
- Crochet J.Y. (1971): Les Vertébrés de l'Oligocène supérieur du Pech du Fraysse poche à phosphate du Quercy (commune de Saint-Project, Tarn-et-Garonne). – *Comptes Rendus de la Société Géologique de France* 1971 (6): 316-317.
- Croizet J. B. (1835): [Title?] – *Bulletin de la Société Géologique de France* 7: (104).
- Crusafont Pairó M. & Quintero I. (1970): Noticia preliminar acerca de un nuevo yacimiento de mamíferos fósiles de la provincia de Guadalajara [Preliminary note on a new record of fossil mammals from the Guadalajara Province.] – *Acta Geologica Hispanica* 5: 102-104. [In Spanish.]
- Crusafont Pairó M. & Villalta J.F. de (1955): Parte Paleontológica. Apéndice I. Aves [Section Paleontology. Appendix I. Aves.] – In: Crusafont M., Villalta J.F. de & Truyols J. (eds.): *El Burdigaliense continental de la cuenca del Vallés-Penedés* [Continental Burdigalian of the Vallés-Penedés cave.] – *Memorias y Comunicaciones del Instituto Geológico* 12: 236-237. [In Spanish.]
- Cuvier G. (1800a): Sur un pied d'oiseau fossile incrusté dans le gypse. – *Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts* 51: 128-132.

- Cuvier G. (1800b): Sur le Ornitholithes de Montmartre. – Société Philomatique de Paris, Bulletin des Sciences (II) 41: 129.
- Cuvier G. (1807): Mémoire sur les ossements d'oiseaux qui se trouvent dans les carrières de pierre à platre des environs de Paris. – Annales du Muséum d'Histoire Naturelle 9: 336-356.
- Cuvier G. (1809): Supplément au mémoire sur les ornitholithes de nos carrières à platre. – Annales du Museum d'Histoire Naturelle 14: 43-46.
- Cuvier G. (1812): Discours préliminaire. – In: Cuvier G. (ed.): Recherches sur les ossements de Quadrupèdes, où l'on rétablit les caractères de plusieurs animaux dont les révolutions de globe ont détruit les espèces. Vol. 1: 1-120. Paris: Déterville.
- Cuvier G. (1822): Des ossements d'oiseaux. – In: Cuvier G. (ed.): Recherches sur les ossemens fossiles. Vol. 3: 302-328. Paris: G. Dufour et E. d'Ocagne. [This is the 2nd edition of Cuvier's works, but the article on bird apparently appeared here for the first time.]
- Cuvier G. (1825): Recherches sur les ossemens fossiles. 3rd ed., Vol. 1-5. – Paris: Edmond d'Ocagne, 436 pp.
- Cuvier G. (1836): Des ossemens d'oiseaux. In: Cuvier G. (ed.): Recherches sur les ossemens fossiles. 4th ed, Vol. 5: 549-599, Atlas, pl. 153-156. Paris: Edmond d'Ocagne.
- Dames W. (1890): Über Vogelreste aus dem Saltholmskalk von Limhamn bei Malmö. – Bihang Svenska Vet.-Akad. Handlingar 16, Afd. IV(1): 1-11.
- Daniels M. (1979): A catalogue of the fossil birds from the Eocene London Clay of England contained in the collection of M. C. S. Daniels. – Loughton: M.C.S. Daniels, 10 + 54 pp.
- Daniels M. (1987): The Waltonian avifauna of Britain. – Letter of Information of the Society of Avian Paleontology and Evolution 1: 5-7.
- Daniels M. (1988): [Untitled report.] – Letter of Information of the Society of Avian Paleontology and Evolution 2: 6-9. [In English.]
- Daniels M. (1989): [Untitled report.] – Society of Avian Paleontology and Evolution, Information Letter 2: 6-9. [In English.]
- Daniels M. (1990): [Untitled report.] – Society of Avian Paleontology and Evolution, Information Letter 3: 5-9. [In English.]
- Daniels M. (1991): [Untitled report.] – Society of Avian Paleontology and Evolution, Information Letter 4: 7-13. [In English.]
- Daniels M. (1993): [Untitled report.] – Society of Avian Paleontology and Evolution, Information Letter 7: 8-12. [In English.]
- Daniels M. (1994): [Untitled report.] – Society of Avian Paleontology and Evolution, Information Letter 8: 10-12. [In English.]
- Daniels M. (1997): [Untitled report.] – Society of Avian Paleontology and Evolution, Information Letter 11: 6-9. [In English.]
- Daniels M. (1999): [Untitled report.] – Society of Avian Paleontology and Evolution, Information Letter 13: 11-15. [In English.]
- Darga R., Böhme M., Göhlich U.B. & Rössner G. (1999): Reste höherer Wirbeltiere aus dem Alttertiär des Alpenvorlandes bei Siegsdorf/Oberbayern. – Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie 39: 1-114.
- Darwin C. (1859): On the origin of species by means of natural selection. – London: J. Murray, ix + 502 pp.
- Dauphin C. (1991): Microstructures et composition chimique des coquilles d'oeufs d'oiseaux et de reptiles. IV. Comparaison des coquilles du Sud de la France. – Revue de Paléobiologie 10: 205-216.
- Dauphin Y. (1992): Microstructures et composition chimique des coquilles d'oeufs d'oiseaux et de reptiles. III. Oiseaux fossiles du Sud de la France. – Palaeontographica (A) 224: 55-66.

- Dauphin Y. (1994): Comparaison des microstructures et de la composition chimique d'oeufs fossiles de Sud de la France, de quelques specimens d'Asie et d'oiseaux actuels. – *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 194: 55-71.
- Davies W. (1880): On some fossil bird remains from Siwalik Hills in the British Museum. – *Geological Magazine* (2) 7: 18-27.
- Davitašvili L.Š. (1948) Istorija evoljucionnoj paleontologii ot Darvina do našich dnei [History of the evolutionary paleontology from Darwin till our days]. – Moskva. [In Russian.]
- Dechaseaux C. (1970a): Cérébralisation croissante chez le courlis (*Numenius*) au cours de la période de l'Eocène supérieur à l'époque actuelle. – *Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris) (D)* 270: 771-773.
- Dechaseaux C. (1970b): Moulages endocraniens d'oiseaux de l'Eocène supérieur du Bassin de Paris. – *Annales de Paléontologie, Vertébrés* 56: 67-72.
- Dehaut E.G. (1911): Animaux fossiles du Cap Figari. – *Materiaux pour Servir à l'Histoire Zoologique et Paléontologique des îles de Corse et du Sardaigne* 3: 53-59.
- Dehaut E.G. (1914): Nouvelles recherches sur les Mammifères pléistocènes et récemment éteint de la Sardaigne. 1 partie: Ordre des Singes. – *Materiaux pour Servir à l'Histoire Zoologique et Paléontologique des îles de Corse et du Sardaigne* 5: 71-85.
- Dehaut E. G. (1920): Contributions à l'étude de la vie vertebrée insulaire dans la Région Méditerranéenne occidentale et particulièrement en Sardaigne et en Corse. – Paris: Paul Lechevalier, 95 pp.
- Dehm R. (1935): Über tertiäre Spaltenfüllungen im Fränkischen und Schwäbischen Jura. – *Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Abteilung (Neue Folge)* 29: 1-86.
- Dehm R. (1937): Neue tertiäre Spaltenfüllungen im südlichen Fränkischen Jura (vorläufige Mitteilung). – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 18: 289-313.
- Dehm R. (1962): Altpleistozäne Säuger von Schernfeld bei Eichstätt in Bayern. – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 2: 17-61.
- Dehm R. (1978): Neue tertiäre Spaltenfüllungen im süddeutschen Jura. – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 18: 289-313.
- Del Hoyo J., Elliott A. & Sargatal J. (eds., 1999): *Handbook of the birds of the world*. Vol. 5: Barn owls to hummingbirds. – Barcelona: Lynx Editions, 759 pp.
- Delamétherie J.C. (1800): Sur une empreinte d'oiseau dans un morceau de plâtre de Montmartre. – *Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts* 51: 132.
- Delamétherie J.C. (1802): Note sur un ornitholithe de Montmartre. – *Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts* 54: 59-60.
- Delle Cave L. (1996): Tertiary avian localities of Italy. – Mlíkovský J. (ed.): *Tertiary avian localities of Europe*. – *Acta Universitatis Carolinae, Geologica* 39: 665-681.
- Delle Cave L., Simonetta A. & Azzaroli A. (1984): A skull of a fossil loon (*Gavia*) from the Pliocene of central Italy. – *Palaeontographia Italica* 73: 5-33.
- Delpech F. (1968): Faunes du Magdalénien VI et de l'Azilien du gisement de Duruthy, commune de Sordes-l'Abbaye (Landes). – *Actes de la Société Linnéenne de Bordeaux* 105 (B) 6: 3-25.
- Delpech F. (1983): Les faunes du Paléolithique supérieur dans le Sud-Ouest de la France. – *Cahiers Quaternaires* 6: 1-453.
- Demathieu G., Ginsburg L., Guérin C. & Truc G. (1984): Étude paléontologique, ichnologique et paléoécologique du gisement oligocène de Saignon (Bassin d'Apt, Vaucluse). – *Bulletin du Muséum National d'Histoire Naturelle (Paris) (C)* (4) 6: 153-183.
- Dement'ev G.P. (1964): Klass Aves. Pticy. [Class Aves. Birds]. – In: Roždesvenskij A.K.

- & Tatarinov L.P. (eds.): Osnovy paleontologii: Zemnovodnye, presmykajuščiesja i pticy [Fundamentals of paleontology: Amphibians, reptiles and birds]: 660-699. Moskva: Nauka. [In Russian.]
- Depéret C. (1887): Recherches sur la succession des faunes des vertébrés miocènes de la vallée du Rhône. – Archives du Museum d'Histoire Naturelle de Lyon 4: 45-319.
- Depéret C. (1892): Sur la faune d'oiseaux pliocènes du Roussillon. – Comptes-Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris) 114: 690-692.
- Depéret C. (1897): Les animaux pliocènes du Roussillon. – Mémoires de la Société Géologique de France, Paléontologie 3: 1-198.
- Depéret C. & Douxami H. (1902): Les vertébrés oligocènes de Pyrimont-Challonges (Savoie). – Mémoires de la Société Paléontologique de Suisse 29: 1-90.
- DesFayes M. (1987): Evidence for the ancient presence of Bald Ibis, *Geronticus eremita*, in Greece. – Bulletin of the British Ornithologists' Club 107: 93-94.
- Desnoyers J. (1859a): Sur des empreintes de pas d'animaux dans le gypse des environs de Paris, particulièrement dans la vallée de Montmorency. – Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris) 49: 67-73.
- Desnoyers J. (1859b): Sur des empreintes de pas d'animaux dans le Gypse des environs de Paris, et particulièrement de la vallée de Montmorency. – Bulletin de la Société Géologique de France (2) 16: 936-944.
- Desor E. (1876): Note sur un nouvel oiseau muni de dents de l'argile de Londres. – Bulletin de la Société des Sciences Naturelles Neuchâtel 10: 35-37.
- Despott G. (1929): Annual report on the working of the (Malta) museum department during 1928-1929. – La Valetta: British Museum (Natural History).
- Dieffenbach E. (1853): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1853: 685-686. [In German.]
- Dollo L. (1883): Note sur la présence du *Gastornis edwardsii*, Lemoine, dans l'assise inférieure de l'étage Landénien, à Mesvin, près Mons. – Bulletin du Musée Royal d'Histoire Naturelle de Belgique 2: 297-307.
- Dollo L. (1909): The fossil vertebrates of Belgium. – Annals of the New York Academy of Sciences 19: 99-119.
- Dubois A. (1988): The genus in zoology: a contribution to the theory of evolutionary systematics. – Mémoires du Muséum National d'Histoire Naturelle, Zoologie 140: 1-124.
- Dubrovo I.A. & Kapelist K.V. (1979): Katalog mestonachozdenij tercičnyh pozvonočnyh Ukrainskoj SSR [Catalogue of the localities of Tertiary vertebrates of the Ukrainian SSR.] – Moskva: Nauka, 159 pp. [In Russian.]
- Dughi R. & Sirugue F. (1959): Sur des fragments de coquilles d'oeufs fossiles de l'Eocène de Basse-Provence. – Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris) 249: 959-961.
- Dughi R. & Sirugue F. (1962): Distribution verticale des oeufs d'oiseaux fossiles de l'Eocène de Basse-Provence. – Bulletin de la Société Géologique de France (7) 4: 69-78.
- Dughi R. & Sirugue F. (1968): Marnes à oeufs d'Paléocène de Basse-Provence. – Bulletin de la Société Géologique de France (7) 10: 542-548.
- Dughi R., Plaziat J.C. & Sirugue F. (1969): La répartition stratigraphique des oeufs d'oiseaux du groupe d'*Ornitholithus arcuatus* D. et S. par rapport aux faunes d'eau douce et marines du Sparnacien et du Thanétien de l'Aude. – Comptes Rendus Sommaires de la Société géologique de France 1969: 9-10.
- Duméril C. (1855): Sur une configuration caractéristique du tibia des oiseaux. – Comptes Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris) 40: 619-620.
- Dyke G.J. & Waterhouse D.M. (2000): A mousebird (Aves, Coliiformes) from the London Clay. – Journal für Ornithologie 143.
[Not seen. *Eocolius walkeri* described from the Eocene of England.]

- Dyke G.J. (2001a): Fossil pseudasturid birds (Aves, Pseudasturidae) from the London Clay. – *Bulletin of the Natural History Museum London (Geology)* 57: 1-4.
- Dyke G.J. (2001b): A primitive swift from the London Clay and the relationships of fossil apodiform birds. – *Journal of Vertebrate Paleontology* 21: 195-200.
- Dyke G.J. (2001c): *Laputavis*, a replacement name for *Laputa* Dyke 2001 (preoccupied name). – *Journal of Vertebrate Paleontology* 21: 401.
- Eastham A. (1968): The avifauna of Gorham's cave, Gibraltar. – *Bulletin of the Institute of Archaeology (London)* 7: 37-42.
- Eastham A. (1985): The Magdalenian avifauna at Erralla cave. – *Munibe* 37: 59-80.
- Eastham A. (1986): The birds of the Cueva de Nerja. – In: Jordá Pardo J.F. (ed.): *La prehistoria de la Cueva de Nerja (Málaga)*. Vol. 1(1): 109-130. Málaga: University of Málaga.
- Eastham A. (1989): Cova Negra and Gorham's cave: the place of birds in Mousterian communities. – In: Clutton-Brock J. (ed.): *The walking larder: 350-357*. Southampton: World Archaeological Congress.
- Eastman C.R. (1898a): On the remains of *Struthiolithus chersonensis* from northern China, with remarks on the distribution of struthious birds. – *Bulletin of the Museum of Comparative Zoology* 32: 127-143.
- Eastman C.R. (1898b): Discovery of a second specimen of the fossil egg of *Struthiolithus*. – *Geological Magazine* (4) 5: 434-439.
- Eastman C.R. (1905): Fossil avian remains from Armissan. – *Memoirs of the Carnegie Museum* 2: 131-138.
- Eikamp H. (1979): Fossile Vogeleier aus dem Diluvium und Tertiär. – *Ornithologische Mitteilungen* 31: 286-296.
- Eikamp H. (1980): Fossiler Nachweis eines Fischadlers aus den untermiozänen (aquitänen) Ziegeltonen von Ravolzhausen im Kreis Hanau (Hessen). – *Deutscher Falkenorden, Jahrbuch* 1980: 79-82.
- Eikamp H. (1990): Fossile Vögel aus der Grube Messel. – *Greifvögel und Falkneri* 1989: 66-68.
- Eldredge N. & Gould S.J. (1972): Punctuated equilibria: an alternative to phyletic gradualism. – In: Schopf, T.J.M. (ed.): *Models in paleobiology*: 82-115. San Francisco: Freeman.
- Elorza M. & Sánchez Marco A. (1993): Postglacial fossil Great Auk and associated avian fauna from the Biscay Bay. – *Munibe* 45: 179-185.
- Emslie S.D. (1988): An early condor-like vulture from North America. – *Auk* 105: 529-535.
- Emslie S.D. (1995): An Early Irvingtonian Avifauna from Leisey Shell Pit, Florida. – *Bulletin of the Florida Museum of Natural History* 37: 299-344.
- Engel T. (1908): *Geologischer Wegweiser durch Württemberg*, 3rd rev. ed. – Stuttgart: E. Schweizerbartsche Verlagshandlung, 645 pp.
- Engelhardt H. & Schottler W. (1914): Die tertiäre Kieselgur von Altschlirf im Vogelsburg. – *Abhandlungen der Grossherzoglichen Hessischen Geologischen Landesanstalt zu Darmstadt* 5: 167-192.
- Ennouchi E. (1930): Contribution à l'étude de la faune du Tortonien de La Grive-Saint-Alban (Isère). – Paris: Presses Modernes, 135 pp.
- Ericson P.G.P. (1999): Material of *Juncitarsus* (Phoenicopteriformes), with a guide for differentiating that genus from the Presbyornithidae (Anseriformes). – In: Olson S.L. (ed.): *Avian paleontology at the close of the 20th century*. – *Smithsonian Contributions to Paleobiology* 89: 245-251.
- Errico F. d' (1994a): Birds of the Great Cosquer: the Great Auk and Paleolithic prehistory. – *Antiquity* 68: 39-47.
- Errico F. d' (1994b): Great Auks of the Cosquer Cave again: a reply to McDonald. –

- Antiquity 68: 856-858.
- Errico F. d' (1994c): Birds of Cosquer Cave. The Great Auk (*Pinguinus impennis*) and its significance during the Upper Palaeolithic. – Rock Art Research 11: 45-57.
- Escher A. (1839): [Untitled abstract of a lecture.] – Verhandlungen der Schweizerischen Naturforschenden Gesellschaft 24: 50. [In German.]
- Espinosa de los Monteros A. (1998): Phylogenetic relationships among the trogons. – Auk 115: 937-954.
- Fabre-Taxy S. & Touraine F. (1960): Gisements d'oeufs d'oiseaux de très grande taille dans l'Eocène de Provence. – Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris) 250: 3870-3871.
- Fahlbusch V. (1976): Report on the international symposium on mammalian stratigraphy of the European Tertiary. – Newsletter on Stratigraphy 5: 160-167.
- Fahlbusch V. (1991): The meaning of MN-zonation: consideration for a subdivision of the European continental Tertiary using mammals. – Newsletter on Stratigraphy 24: 159-173.
- Fahlbusch V. & Gall H. (1970): Die obermiozäne Fossil-Lagerstätte Saldelzhausen. I. Entdeckung, Geologie, Faunenübersicht und Grabungsbericht für 1969. – Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie 10: 365-396.
- Falconer H. (1868): [Untitled letter from 2 August 1860 to Capt. Spratt.] – In: Murchinson, C. (ed.): Palaeontological memoirs and notes of the late Hugh Falconer, A.M., M.D., Vol. 2: 300. London: Hardwicke. [In English.]
- Fatio V. (1906): Sur le Waldrapp "Corvus sylvaticus" de Gesner. – Ibis (8) 6: 139-144.
- Faujas-Saint-Fond, B. (1801): Memoire sur quelques fossiles rares de Vestena Nova dans le Veronias, qui n'on pas ete descrit, et que M. de Gazola a donne' au Museum national d'histoire naturelle en l'an 11. – Annales du Museum d'Histoire Naturelle (Paris) 3: 18-24.
- Feduccia A. (1980): The age of birds. – Cambridge, Mass.: Harvard University Press, 196 pp. [For a review see Cracraft 1981b.]
- Feduccia A. (1996): The origin and evolution of birds. – New Haven: Yale University Press, x + 420 pp.
- Fejfar O. & Heinrich W.-D. (1985): Zur Bedeutung der Wirbeltierfundstätten von Ivanovce und Hajnáčka für die Säugetierpaläontologie im Pliozän und frühen Pleistozän in Europa: Kenntnisstand und Probleme. – Věstník Ústředního Ústavu Geologického 60: 213-224.
- Fejfar O. & Heinrich W.-D. (1990a): Muroid rodent biochronology of the Neogene and Quaternary in Europe. – In: Lindsay E.H., Fahlbusch V. & Mein P. (eds.): European Neogene mammal chronology: 91-117. New York: Plenum Press.
- Fejfar O. & Heinrich W.-D. (1990b): Proposed biostratigraphical division of the European continental Neogene and Quaternary based on muroid rodents (Rodentia, Mammalia). In: Fejfar O. & Heinrich W.-D. (eds.): Evolution, phylogeny and biostratigraphy of Arvicolids: 115-124. Praha: Geological Survey, & München: Pfeil Verlag.
- Fejfar O., Heinrich W.-D., Pevzner M.A. & Vangengeim E.A. (1997): Late Cenozoic sequence of mammalian sites in Eurasia: an updated correlation. – Palaeogeography, Palaeoclimatology, Palaeoecology 133: 259-288.
- Fejfar O., Heinrich W.-D. & Lindsay E.H. (1998): Updating the Neogene Rodent biochronology in Europe. – In: Kolfschoten T. van & Gibbard P.L. (eds): The Dawn of the Quaternary. Proceedings of the SEQS-EuroMam symposium 1996. – Mededelingen Nederlands Instituut voor Toegepaste Geowetenschappen TNO 60: 533-553.
- Fischer K. (1962): Der Riesenlaufvogel *Diatryma* aus dem Eozän des Geiseltales. – Hallesches Jahrbuch für Mitteldeutsche Erdgeschichte 4: 26-33.

- Fischer K. (1967): Ein neuer grosser Laufvogel aus dem Eozän des Geiseltales bei Halle (Salle). – Berichte der Deutschen Gesellschaft für Geologische Wissenschaften (A) 12: 601-605.
- Fischer K. (1978): Neue Reste des Riesenlaufvogels *Diatryma* aus dem Eozän des Geiseltales bei Halle (DDR). – Annalen für Ornithologie 2: 133-144.
- Fischer K. (1982): Wirbeltierfunde aus dem marinen Mitteloligozän des Weissensteiner Beckens (Bezirk Leipzig, DDR). – Wissenschaftliche Zeitschrift der Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Reihe 31: 151-153.
- Fischer K. (1983a): Möwenreste (Laridae, Charadriiformes, Aves) aus dem mitteloligozänen Phosphoritknollenhorizont des Weissensteiner Beckens bei Leipzig (DDR). – Annalen für Ornithologie 7: 151-155.
- Fischer K. (1983b): *Oligostrix rupelensis* n.gen., n.sp., eine neue Ureule (Protostrigidae, Strigiformes, Aves) aus dem marinen Mitteloligozän des Weissensteiner Beckens bei Leipzig (DDR). – Zeitschrift für Geologische Wissenschaften 11: 483-487.
- Fischer K. (1985): Ein albatrossartiger Vogel (*Diomedeoides minimus* nov. gen, nov. sp., Diomedeoididae nov. fam., Procellariiformes) aus dem Mitteloligozän bei Leipzig (DDR). – Annalen für Ornithologie 9: 113-118.
- Fischer K. (1987): Eulenreste (*Eoglaucidium pallas* nov. gen, nov. sp., Strigiformes, Aves) aus der mitteloligozänen Braunkohle des Geiseltals bei Halle (DDR). – Annalen für Ornithologie 11: 137-142.
- Fischer K. (1990): Der Hühnervogel *Pirortyx major* (Gaillard, 1939) aus dem marinen Mitteloligozän bei Leipzig (DDR). – Annalen für Ornithologie 14: 133-136.
- Fischer K. & Mauersberger G. (1989): Zu Geschichte und Bedeutung der paläornithologischen Sammlung des Museums für Naturkunde zu Berlin. – Annalen für Ornithologie 13: 139-151.
- Fischer K. & Stephan B. (1971): Ein flugunfähiger Kranich (*Grus cubensis* n.sp.) aus dem Pleistozän von Kuba. – Eine Osteologie der Familie der Kraniche (Gruidae). – Wissenschaftliche Zeitschrift der Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Reihe 20: 541-592.
- Fischer K. & Stephan B. (1974): Eine pleistozäne Avifauna aus der Ghar Dalam Höhle, Malta. – Zeitschrift für Geologische Wissenschaften 2: 515-523.
- Fischer O. & Hably L. (1991): Pliocene flora from the alginite at Gércé. – Annales Historico-Naturales Musei Nationalis Hungarici 83: 25-47.
- Fisher C. T. (1997): Past human exploitation of birds on the Isle of Man. – International Journal of Osteoarchaeology 7: 292-297.
- Flot M. (1892): Description de deux oiseaux nouveaux du gypse parisien. – Mémoires de la Société Géologique de France, Paléontologie 7: 1-10.
- Ford N.L. (1967): A systematic study of the owls based on comparative osteology. – Unpub. Thesis, University of Michigan, Chicago, 128 pp.
- Forsyth Major C.J. (1888): Sur un gisement d'ossements fossiles dans l'île de Samos, contemporains de l'âge de Pikermi. – Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris) 107: 1178-1181.
- Forsyth Major C.J. (1892): Le gisement ossifère de Mytilini. – In: Stefani C. de, Forsyth Major C.J. & Barbey W. (eds): Samos. Étude géologique, paléontologique et botanique: 85-99. Lausanne: Georges Bridel & Cie.
- Forsyth Major C.J. (1894): Le gisement ossifère de Mytilini et Catalogue d'ossements fossiles recueillis à Mytilini, île de Samos, et déposées au Collège Gaillard, à Lausanne. – Lausanne: Georges Bridel & Cie, 51 pp. [Pp. 1-15 is a reprint of Forsyth Major 1892.]
- Fortey R.A. (1985): Gradualism and punctuated equilibria as competing and complementary theories. – Special Papers in Paleontology 33: 17-28.
- Fortey R.A. (1988): Seeing is believing: gradualism and punctuated equilibria in the fossil

- record. – *Science Progress* 72: 1-19.
- Fortis G.B. (1800): Des morceaux de fer et des ornitholithes trouvés dans les carrières de Montmartre. – *Journal de Physique, de Chimie, d'Historie Naturelle et des Arts* 50: 321-341.
- Fortis, G.B. (1801): [Untitled letter to J.-C. Delamétherie.] – *Journal de Physique, de Chimie, d'Historie Naturelle et des Arts* 52: 320-322. [In French.]
- Fraas O. (1870): Die Fauna von Steinheim. Mit Rücksicht auf die miocänen Säugethier- und Vogelreste des Steinheimer Beckens. – *Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg* 26: 145-306. [Also published separately in Stuttgart by E. Schweizerbart'sche Verlagshandlung, 1870. The book has separate pagination and separate numbering of plates.]
- Fraas O. (1879): Brüteplätze von Wasservögeln der jüngsten Tertiärzeit. – *Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde* 1879: 555-556.
- Freudenberg W. (1914): Die Säugetiere des älteren Quartärs von Mitteleuropa mit besonderer Berücksichtigung der Fauna von Hundsheim. – *Geologisch-Paläontologische Abhandlungen* 16: 453-672.
- Friant M. (1950a): Sur le Héron nocturne (*Nycticorax fenensis* nov. spec.) des tourbières anciennes du Cambridgeshire (Angleterre). – *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris)* 230: 2126-2128.
- Friant M. (1950b): Le Héron nocturne des Tourbières anciennes du Cambridgeshire (*Nycticorax fenesis*, nov. spec.). – *Proceedings of the Zoological Society of London* 120: 325-335.
- Friant M. (1950c): Sur le Vautour des cavernes, *Gyps fulvus* Gm. *spelaeus* nov. var. – *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris)* 231: 1164-1165.
- Friant M. (1951): Le "Vautour des Cavernes" (*Gyps fulvus* Gm. ssp. *spelaeus* nov.) de Goyet (Belgique). – *Proceedings of the Zoological Society of London* 121: 417-426.
- Fuller E. (1999): *The Great Auk*. – Southampton: Errol Fuller, 448 pp.
- Fürbringer M. (1888): *Untersuchungen zur Morphologie und Systematik der Vögel*. – Amsterdam: T. J. Van Holkema, & Jena: Gustav Fischer, 1751 pp.
- Fürbringer M. (1902): Zur vergleichenden Anatomie des Brustschulterapparates und der Schultermuskeln. – *Jenaische Zeitschrift für Naturwissenschaften* 36: 289-736.
- Furon R. & Soyer R. (1947): *Catalogue des fossiles tertiaires du bassin de Paris*. – Paris: Paul Lechevalier, 240 pp. + 32 pls.
- Gadow H. (1891): Vögel. Vol. 1. – In: Bronn H.G. (ed.): *Klassen und Ordnungen des Thier-Reichs* Vol. 6. Leipzig: C. F. Winter'sche Verlagshandlung. [With a contribution by E. Selenka.]
- Gadow H. (1892): On the classification of birds. – *Proceedings of the Zoological Society of London* 1892: 229-256.
- Gadow H. (1893): Vögel. Vol. 2. Systematischer Theil. – In: Bronn H.G. (ed.): *Klassen und Ordnungen des Thier-Reichs* Vol. 6 (4): vii + 304 pp. Leipzig: C. F. Winter'sche Verlagshandlung.
- Gaillard C. (1908): Les oiseaux des Phosphorites du Quercy. – *Annales de l'Université de Lyon (Nouvelle Série)* 23: 1-178.
- Gaillard C. (1936): Un oiseau géant dans les dépôts éocènes du Mont-d'Or lyonnais. – *Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris)* 202: 965-967.
- Gaillard C. (1937): Un oiseau géant dans les dépôts éocènes du Mont-d'Or lyonnais. – *Annales de la Société Linnéenne de Lyon (Nouvelle Série)* 80: 111-126.
- Gaillard C. (1939): Contribution à l'étude des oiseaux fossiles. – *Nouvelles Archives du Museum de Lyon* 15(2): 1-100.
- Gál E. (1999): Late Pleistocene bird remains from Grotta Romanelli (Apulia, Italy). – In:

- 2nd International Conference of PhD students (Miskolc, Hungary, 8-14 August 1999), Section Proceedings – Natural Sciences, pp. 121-125.
- Gál E., Hír J., Kessler E., Kókay J., Mészáros L. & Vencel M. (1999): Középső-miocén ősmaradványok, a Mátraszőlős, Rákóczi-kápolna alatti útbevágásból. I. A Mátraszőlős 1. lelőhely [Middle Miocene fossils from the sections at the Rákoczi chapel at Mátraszőlős. I. Locality Mátraszőlős 1.] – *Folia Historico Naturalia Musei Matraensis* 23: 33-78. [In Hungarian.]
- Gál E., Hír J., Kessler E., Kókay J. & Vencel M. (1999): Középső-miocén ősmaradványok a Mátraszőlős, Rákóczi-kápolna alatti útbevágásból. II. A Mátraszőlős 2. lelőhely [Middle Miocene fossils from the sections at the Rákoczi chapel at Mátraszőlős. II. Locality Mátraszőlős 2.] – *Folia Historico Naturalia Musei Matraensis* 24: 39-75. [In Hungarian.]
- Ganea I.M. (1965): Sovremennye dannye ob iskopaemoj ornitofaune Moldavii [New data on the fossil avifauna of Moldova.] – In: *Biologičeskie osnovy rekonstrukcii, racional'nogo ispol'zovanija i ochrany fauny južnoj zony evropejskoj časti SSSR* [Biological bases of the reconstruction, rational exploitation and conservation of the fauna of the southern zone of the European part of the USSR.]: 26-34. Chişinău. [In Russian.]
- Ganea I.M. (1969a): Iskopaemye kurinye pticy Moldavii [Fossil galliforms of Moldova.] – *Ochraňa Prirody Moldavii* 7: 125-134. [In Russian.]
- Ganea I.M. (1969b): Iskopaemye čiščnye pticy Moldavii [Fossil raptors of Moldova.] – *Izvestija Akademii Nauk Respubliki Moldova, Biologičeskie i Chimičeskie Nauki* 1969 (1): 24-28. [In Russian.]
- Ganea I.M. (1972): Istorija ornitofauny Moldavii s pozdnego miocena do našich dnej [History of the Moldovan avifauna since the late Miocene to the present.] – In: *Fauna nazemnyh pozvonočnyh Moldavii i problemy ee rekonstrukcii* [Fauna of terrestrial vertebrates of Moldova and problems of its reconstruction]: 20-43. Chişinău: Ştiinţa. [In Russian.]
- Ganea I.M. & Burčak-Abramovič N.I. (1992): Sostojanie izučennosti iskopaemych ptic roda *Gallus* v Moldove, na Ukraine i Kavkaze [The state of knowledge of fossil birds of the genus *Gallus* in Moldova, Ukraine and the Caucasus.] – *Izvestija Akademii Nauk Respubliki Moldova, Biologičeskie i Chimičeskie Nauki* 1992 (6): 48-50, 71. [In Moldovan.]
- Ganea I.M. & Ketraru N.A. (1964): Nekotorye dannye ob" ornitofaune iz paleolitičeskogo grota Starye Duruitory [Some data on the avifauna from the Paleolithic cave Starye Duruitory.] – *Izvestija Akademii Nauk Moldavskoj SSR, Serija Biologičeskich i Chimičeskich Nauk* 1964(1): 45-48. [In Ukrainian.]
- Ganea I.M. & Ketraru N.A. (1965): Iskopaemaja ornitofauna iz raskopok paleolitičeskoj stojanki Brynzeny 1 [Fossil avifauna from the excavations of the Paleolithic site Brynzeny 1.] – *Izvestija Akademii Nauk Moldavskoj SSR, Serija Biologičeskich i Chimičeskich Nauk* 1965(5): 98-104. [In Russian.]
- Ganea I.M. & Kuročkin E.N. (1967): Kratkie itogi i perspektivy izučenija avifauny neogena jugo-zapadnyh rajonov Evropejskoj časti SSSR [Short results and perspectives of the study of Neogene avifauna of south-western regions of the USSR.] – In: *Mesto i značenie iskopeamych mlekopitajuščich Moldavii v kajnozoe SSSR* [The role and importance of fossil mammals of Moldova in the Cenozoic of the USSR.]: 105-109. Chişinău: AN SSSR. [In Russian.]
- García Talavera I. (1990): Aves gigantes en el Mioceno de Famara (Lanzarote). – *Rev. Acad. Canar. Cienc.* 11: 71-79.
- Gasser U. (1966) Sedimentologische Untersuchungen in der äusseren Zone der subalpinen Molasse des Entlebuches (Kt. Luzern). – *Eclogae Geologicae Helvetiae* 59: 723-772.
- Gastaldi B. (1866a): Intorno ad alcuni fossili della Toscana e del Piemonte [On some

- fossils from Tuscany and Piemonte.] – *Atti della Reale Accademia delle Scienze di Torino* 1: 38-43. [In Italian.]
- Gastaldi B. (1866b): *Intorno ad alcuni fossili del Piemonte e della Toscana* [On some fossils from Piemonte and Tuscany.] – *Memoire della Reale Accademia delle Scienze di Torino* (2) 24: 193-236. [Letter by T. Salvadori on p. 223-225.] [In Italian.]
- Gaudin C.T. (1854): [Untitled abstract of a lecture.] – *Bulletin des Séances de la Société Vaudoise des Sciences Naturelles* 3: 281. [In French.]
- Gaudry A. (1862a): *Note sur les débris d'oiseaux et de reptiles trouvés à Pikermi (Grèce), suivie de quelques remarques de paléontologie générale.* – *Bulletin de la Société Géologique de France* (2) 19: 629-640.
- Gaudry A. (1862b): *Animaux fossiles et géologie de l'Attique d'après les recherches faites en 1855-56 et en 1860 sous les auspices de l'Académie des Sciences.* – Paris: F. Savy, 476 pp. (Aves on pp. 309-316)
- Gaudry A. (1862-1867): *Animaux fossiles et géologie de l'Attique d'après les recherches faites en 1855-56 et en 1860 sous les auspices de l'Académie des Sciences: Atlas.* – Paris: F. Savy, 7 pp. + 76 pls. (Aves on pls. 58-59)
- Gaudry A. (1885): [Untitled comment on the lecture by Lemoine 1885.] – *Bulletin de la Société Géologique de France* (3) 13: 412-413. [In French.]
- Gejvall N.G. (1969): *Lerna. A Preclassical site in the Argolid. Vol. 1. The fauna.* – Princeton: American School of Classical Studies in Athens, 107 pp.
- Geoffroy-Saint-Hilaire I. (1842): [Squelette fossile presque entier d'oiseau, provenant des carrières de Pantin.] – *Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris)* 14: 219.
- George T.N. (1956): *Biospecies, chronospecies, and morphospecies.* – In: Sylvester-Bradley P. C. (ed.): *The species concept in paleontology: 123-137.* London: Systematics Association.
- George W. & Vincent S. (1978): *Notes on the London Clay of the Ockendon Clay plant, South Ockendon, Essex.* – *Tertiary Research* 2: 5-8.
- Germar E.F. (1826): *Bemerkungen über die fossilen Knochen von Westeregeln.* – In: Keferstein C. (ed.): *Teutschland, geognostisch-geologisch dargestellt* 3: 601-612. Weimar.
- Géroutet P. (1965): *Du "Waldrapp" de Gesner aux Ibis chauves du Maroc.* – *Nos Oiseaux* 28: 129-143.
- Gervais P. (1844a): *Remarques sur les oiseaux fossiles.* – Paris: Imprimerie Poussielge, 40 pp. [Thesis defended on 5 August, hence published before that date.]
- Gervais P. (1844b): [Untitled report on Gervais 1844a.] – *L'Institut (Paris)* 12: 293-294. [In French; presented on the meeting of the Société Philomatique de Paris on 17 August, hence published after this date. For other versions of this report see: *Procès-Verbaux de la Société Philomatique de Paris 1844: 67-70*, and *Edinburgh New Philosophical Journal* 38: 175-177 (1845).]
- Gervais P. (1849): *Oiseaux et Reptiles fossiles de France.* – *Mémoires de l'Académie des Sciences de Montpellier* 1: 220-222.
- Gervais P. (1848-52): *Zoologie et Paléontologie françaises (Animaux vertébrés), ou Nouvelles recherches sur les animaux vivants et fossiles de la France.* – Paris: Arthus Bertrand, Vol. I: iv + 271 pp.; Vol. II: Explication des planches et divers mémoires relatifs aux animaux vertébrés; Vol. III: Atlas, 80 pls.
- Gervais P. (1859): *Zoologie et Paléontologie françaises. Nouvelles recherches sur les animaux vertébrés dont trouve les ossements enfouis dans le sol de la France.* 2nd ed. – Paris: Arthus Bertrand, 544 pp. + 84 pls. [Aves on pp. 403-428.]
- Gervais P. (1862): *Examen d'un ornitholithe d'Armissan (Aude).* – *Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris)* 54: 895-896.
- Gervais P. (1867-1869): *Zoologie et paléontologie générales.* – Paris: Arthus Bertrand, vii

- + 263 pp + 50 pls.
- Gervais P. (1873): Enumération de quelques ossements d'animaux vertébrés recueillis aux environs de Reims par M. Lemoine. – *Journal de Zoologie* 2: 351-355.
- Gervais P. (1877): Enumération de quelques ossements d'animaux vertébrés recueillis aux environs de Remis par M. Lemoine. – *Journal de Zoologie* 4: 74-79.
- Gesner C. (1555): *Historia animalium* [History of animals]. Vol. 3. De avibus naturae [On the nature of birds.] – Zürich. [In Latin.]
- Geus A. (1959): Der Waldrapp – in Europa ausgestorben. – *Falke* 6: 133-136.
- Gheorghiu C. et al. (1965): Acumularile de carbonati de la Corund [The carbonate accumulation of Corund.] – *Comunicarile de Geologie* 3: 123-146. [In Romanian.]
- Giebel C.G. (1847): *Fauna der Vorwelt mit steter Berücksichtigung der lebenden Thiere*. Vol. I (2): Vögel und Amphibien. – Leipzig: Brockhaus, 218 pp.
- Giebel C.G. (1850): Über das Vorkommen der diluvialen Knochen in der Provinz Sachsen. – *Jahresberichte des Naturwissenschaftlichen Vereines in Halle* 4: 236-245.
- Giebel C.G. (1860): Zur Fauna der Braunkohlenformation von Rippersroda in Thüringen. – *Zeitschrift für die Gesamten Naturwissenschaften* 16: 147-153.
- Giebel C.G. (1862): Wirbeltier- und Insektenreste im Bernstein. – *Zeitschrift für die Gesamten Naturwissenschaften* 20: 311-321.
- Gingerich P.D. (1979): The stratophenetic approach to phylogeny reconstruction in vertebrate paleontology. – In: Cracraft J. & Eldredge N. (eds.): *Phylogenetic analysis and paleontology*: 40-70. New York: Columbia University Press.
- Gingerich P.D. (1985): Species in the fossil record: concepts, trends, and transitions. – *Paleobiology* 11: 27-41.
- Gisnburg, L., Cheneval J., Janvier P., Pouit D. & Sen, S. (2000): Les Vertébrés des sables continentaux d'âge orléanien inférieur (MN 3) de Mauvières à Marcilly-sur-Maulne (Indre-et-Loire), La Brosse à Meigné-le-Vicomte (Maine-et-Loire) et Chitenay (Loir-et-Cher). – *Geodiversitas* 22: 597-631.
- Glass B. (ed., 1959): *Forerunners of Darwin*. – Baltimore: John Hopkins Press, 471 pp.
- Gloger C.W.L. (1841-1842): *Gemeinnütziges Hand- und Hilfsbuch der Naturgeschichte*. – Breslau [= Wrocław]: Schultz & Co., xlv + 496 pp.
- Göhlich U.B. (2000): The avifauna from the Miocene fossil site of Sandelzhausen (Bavaria, Southern Germany). – *Vertebrata Palasiatica* 38 (Supplement): 14. [Abstract.]
- Göhlich U.B. & Rössner G.E. (1999): Miozäne Wirbeltiere aus der Oberen Süßwassermolasse von Sandelzhausen (Süddeutschland). – *Terra Nostra* (Berlin) 8: 93. [Abstract.]
- Gonzales Morales J.A. (1980): Interpretación ornitológica de los grabados [The ornithological interpretation of engravings.] – In: Gonzales Echegeray J. (ed.): *El yacimiento de la cueva de 'El Pendo' (Excavaciones 1953-57)* [The deposits of El Pendo Cave (Excavations 1953-57).] – *Bibliotheca Praehistorica Hispana* 17: 257-261. [In Spanish.]
- Gosse P.H. (1847): *The birds of Jamaica*. – London: Van Voorst, x + 447 pp.
- Gould S.J. (1966): Allometry and size in ontogeny and phylogeny. – *Biological Reviews* 41: 587-640.
- Gould S.J. & Eldredge N. (1977): Punctuated equilibria: tempo and mode of evolution reconsidered. – *Paleobiology* 3: 115-151.
- Gray G.R. (1841): *A list of the genera of birds, with their synonyms and an indication of the typical species of each genus*. – London: Richard and John E. Taylor, xii + 115 pp.
- Greenway J.C. Jr. (1967): *Extinct and vanishing birds of the world*. 2nd ed. – New York: Dover, xvi + 520 pp.
- Grieve S. (1885): *The Great Auk, or Garefowl (Alca impennis Linn.)*. Its history, archaeology, and remains. – London: Thomas C. Jack, 540 pp.
- Grieve S. (1888): *Recent notes on the Great Auk or Garefowl (Alca impennis Linn.)*. –

- Transactions of the Edinburgh Field Naturalists and Microscopical Society 20: 42-56.
- Grigorescu D. & Kessler E. (1977): The middle Sarmatian avian fauna of South Dobrogea. – *Revue Roumaine de Géologie, Géophysique et Géographie (Géologie)* 21: 93-108.
- Grigorescu D. & Kessler E. (1988): New contributions to the knowledge of the Sarmatian birds from South Dobrogea in the frame of the eastern Paratethyan avifauna. – *Revue Roumaine de Géologie, Géophysique et Géographie (Géologie)* 32: 91-97.
- Grigson C. (1989): Bird-foraging patterns in the Mesolithic. – In: Bonsall C. (ed.): *The Mesolithic in Europe*: 60-72. Edinburgh: John Donald.
- Gromov V.I. (1948): Paleontologičeskoe i archeologičeskoe obosnovanie stratigrafii kontinental'nych otloženij četvertičnogo perioda na territorii SSSR [Paleontological and archaeological foundation of the stratigraphy of the Quaternary continental deposits on the territory of the USSR.] – *Trudy Instituta Geologičeskich Nauk Akademii Nauk SSSR* 64: 1-521. [In Russian.]
- Gromov V.I. (1966): Fauna iz stojanki Sungir' [Fauna from the Sungir' site.] – *Trudy Instituta Geologii Akademii Nauk SSSR* 162: 74-78. [In Russian.]
- Grozescu H.G. (1914): Geologia regiunii subcarpatice din partea de nord a districtului Bacău [Geology of the subcarpathian region in the northern part of the Bacău Province.] – *Analele Institutului de Geologie Romanei* 8: 118-208. [In Romanian.]
- Grycaj, T.G. (1938): Paleontologični rozkopki v karstovych peščerach odes'kich katakomb [Paleontological excavations in karstic caves of the Odesa catacombs.] – *Visti Akademii Nauk Ukrains'koji RSR* 1938 (4). [In Ukrainian.]
- Grycaj, T.G. (1939): Raboty Odesskoj paleontologičeskoj èkspedicii Akademii nauk USSR [Work of the Odesan paleontological expedition of the Academy of Sciences of the Ukrainian SSR.] – *Priroda* 1939 (6): 90-91. [In Russian.]
- Guérin C. (1982): Première biozonation du Pléistocène européen, principal résultats biostratigraphique de l'étude des Rhinocerotidae (Mammalia, Perissodactyla) du Miocène terminal au Pléistocène supérieur d'Europe occidentale. – *Géobios* 15: 593-598.
- Hachler E. (1944): Soustavný přehled recentních i fosilních čeledí ptáčích [A systematic list of extant and fossil families of birds.] – *Ornitholog* 11: 1-8. [In Czech.]
- Haffer J. (1986) Superspecies and species limits in vertebrates. – *Zeitschrift für Zoologische Systematik und Evolutionsforschung* 24: 169-190.
- Haffer J. (1989) Parapatrische Vögelarten der Paläarktischen Region. – *Journal für Ornithologie* 130: 475-512.
- Haffer J. (1995): Species versus phyletic lineages. – In: Peters D.S. (ed.): *Acta palaeornithologica*. – *Courier Forschungsinstitut Senckenberg* 181: 303-309.
- Hardy M. (1891): La station quaternaire de Raymonden à Chancelade (Dordogne), et la sépulture d'un chasseur de rennes. – Paris: Ernest Leroux.
- Harrison C.J.O. (1971): Flamingo (Phoenicopteridae) remains from the British Upper Eocene. – *Bulletin of the British Ornithologists' Club* 91: 36-39.
- Harrison C.J.O. (1974): A re-examination of the extinct marabou stork, *Leptoptilos falconeri*: with descriptions of some new species. – *Bulletin of the British Ornithologists' Club* 94: 42-49.
- Harrison C.J.O. (1975a): The taxonomic status of Milne-Edward's [sic!] fossil Sulids. – *Bulletin of the British Ornithologists' Club*. 95: 51-54.
- Harrison C.J.O. (1975b): *Empheresula*: new name for *Parasula* Harrison 1975. – *Bulletin of the British Ornithologists' Club*. 95: 175.
- Harrison C.J.O. (1975c): Ordinal affinities of the Aegialornithidae. – *Ibis* 117: 164-170.
- Harrison C.J.O. (1976): The wing proportions of the Eocene diver *Colymboides anglicus*. – *Bulletin of the British Ornithologists' Club* 96: 64-65.

- Harrison C.J.O. (1977): A new guillemot, *Cephus* species, from the Lower Pleistocene – Suffolk Natural History 17: 238-239.
- Harrison C.J.O. (1978): A new jungle-fowl from the Pleistocene of Europe. – Journal of Archaeological Science 5: 373-376.
- Harrison C.J.O. (1979a): The herons (Ardeidae) of the Old World Tertiary. – In: Harrison C.J.O. & Walker C.A. (eds.): Studies in Tertiary avian paleontology. – Tertiary Research, Special Paper 5: 11-17.
- Harrison C.J.O. (1979b): The Upper Eocene birds of the Paris Basin: A brief re-appraisal. – Tertiary Research 2: 105-109.
- Harrison C.J.O. (1979c): Small non-passerine birds of the Lower Tertiary as exploiters of ecological niches now occupied by passerines. – Nature 281: 562-563.
- Harrison C.J.O. (1979d): Birds of the Cromer Forest Bed series of the East Anglian Pleistocene. – Transactions of the Norfolk and Norwich Naturalists' Society 24: 277-286.
- Harrison C.J.O. (1979e): The extinct Maltese crane. – Il-Merill 20: 14-15.
- Harrison C.J.O. (1979f): Pleistocene birds of Swanscombe, Kent. – London Naturalist 58: 6-8.
- Harrison C.J.O. (1980a): A small owl from the Lower Eocene of Britain. – Tertiary Research 3: 83-87.
- Harrison C.J.O. (1980b): Pleistocene bird remains from Tornewton Cave and the Brixham Windmill Hill Cave in south Devon. – Bulletin of the British Museum (Natural History), Geology 33: 91-100.
- Harrison C.J.O. (1980c): A re-examination of British Devensian and earlier Holocene bird bones in the British Museum (Natural History). – Journal of Archaeological Science 7: 53-68.
- Harrison C.J.O. (1980d): Fossil birds from Afrotropical Africa in the collection of the British Museum (Natural History). – Ostrich 51: 92-98.
- Harrison C.J.O. (1981): A re-assignment of *Amphipelargus majori* from the Ciconiidae (Ciconiiformes) to Ergilornithidae (Gruiformes). – Tertiary Research 3: 111-112.
- Harrison C.J.O. (1982a): Cuculiform, piciform and passeriform birds in the Lower Eocene of England. – Tertiary Research 4: 71-81.
- Harrison C.J.O. (1982b): A new tiny raptor from the Lower Eocene of England. – Ardea 70: 77-80.
- Harrison C.J.O. (1982c): The earliest parrot: a new species from the British Eocene. – Ibis 124: 203-210.
- Harrison C.J.O. (1982d): An atlas of the birds of the Western Palearctic. – Princeton: Princeton University Press, 332 pp.
- Harrison C.J.O. (1983a): A new wader, *Recurvirostridae* (Charadriiformes), from the early Eocene of Portugal. – Ciências da Terra 7: 9-16.
- Harrison C.J.O. (1983b): Fossil birds from the Lower London Clay of Essex: the W. George and S. Vincent collections. – Tertiary Research 5: 81-82.
- Harrison C.J.O. (1984a): Further additions to the fossil birds of Sheppey: a new falconid and three small rails. – Tertiary Research 5: 179-187.
- Harrison C.J.O. (1984b): Rail-like cursorial birds of the British Lower Eocene, with descriptions of two new species. – London Naturalist 63: 14-23.
- Harrison C.J.O. (1984c): A revision of the fossil swifts (Vertebrata, Aves, suborder Apodi), with descriptions of three new genera and two new species. – Mededelingen van de Werkgroep voor Tertiaire en Kwartaire Geologie 21: 157-177.
- Harrison C.J.O. (1985a): A bony-toothed bird (Odontopterygiformes) from the Paleocene of England. – Tertiary Research 7: 23-25.
- Harrison C.J.O. (1985b): The Pleistocene and prehistoric birds of south-eastern England.

- Bulletin of the Geological Society of Norfolk 35: 53-69.
- Harrison C.J. O. (1985c): A large fossil gamebird, Phasianidae, from the middle Miocene of Turkey. – *Journal of Natural History* 19: 135-137.
- Harrison C.J.O. (1986): A re-examination of some Tertiary Old World ibises, Threskiornithidae. – *Tertiary Research* 7: 139-143.
- Harrison C.J.O. (1987): Pleistocene and prehistoric birds of South-West Britain. – *Proceedings of the University of Bristol Speleological Society* 18: 81-104.
- Harrison C.J.O. & Cowles G.S. (1977): The extinct large cranes of the north-west Palearctic. – *Journal of Anthropological Science* 4: 25-27.
- Harrison C.J.O. & Stewart J.R. (1999): Avifauna. – In: Roberts M.B. & Parfitt S.A. (eds): Boxgrove. A middle Pleistocene hominid site at Earham Quarry, Boxgrove, West Sussex. – *English Heritage Archaeological Report (London)* 17: 187-196.
- Harrison C.J.O. & Walker C.A. (1971): A new ibis from the Lower Eocene of Britain. – *Ibis* 113: 367-368.
- Harrison C.J.O. & Walker C.A. (1972): The affinities of *Halcyornis* from the Lower Eocene. – *Bulletin of the British Museum (Natural History), Geology* 21: 153-170.
- Harrison C.J.O. & Walker C.A. (1975): A new swift from the Lower Eocene of Britain. – *Ibis* 117: 162-164.
- Harrison C.J.O. & Walker C.A. (1976a): Birds of the British Upper Eocene. – *Zoological Journal of the Linnaean Society (London)* 59: 323-351. [For a review see Steadman 1981.]
- Harrison C.J.O. & Walker C.A. (1976b): A review of the bony-toothed birds (Odontopterygiformes): with descriptions of some new species. – *Tertiary Research, Special Paper* 2: 1-62.
- Harrison C.J.O. & Walker C.A. (1976c): A reappraisal of *Prophaethon shrubsolei* Andrews (Aves). – *Bulletin of the British Museum (Natural History), Geology* 27: 1-30.
- Harrison C.J.O. & Walker C.A. (1976d): Cranial material of Oligocene and Miocene flamingos: with a description of a new species from Africa. – *Bulletin of the British Museum (Natural History), Geology* 27: 305-312.
- Harrison C.J.O. & Walker C.A. (1977a): Birds of the British Lower Eocene. – *Tertiary Research, Special Paper* 3: 1-52. [For a review see Steadman 1981.]
- Harrison C.J.O. & Walker C.A. (1977b): A re-examination of the fossil birds from the Upper Pleistocene in the London Basin. – *London Naturalist* 56: 6-9.
- Harrison C.J.O. & Walker C.A. (1977c): The invalidity of Pleistocene records of Wandering Albatross *Diomedea exulans*. – *Ibis* 120: 229-230.
- Harrison C.J.O. & Walker C.A. (1978a): *Proherodius oweni* as a species of Presbyornithidae. – *Tertiary Research* 2: 1-3.
- Harrison C.J.O. & Walker C.A. (1978b): The North Atlantic Albatross *Diomedea anglica*, a Pliocene – Lower Pleistocene species. – *Tertiary Research* 2: 45-46.
- Harrison C.J.O. & Walker C.A. (1979a): Birds of the British Middle Eocene. – In: Harrison C.J.O. & Walker C.A. (eds.): *Studies in Tertiary avian palaeontology*. – *Tertiary Research, Special Paper* 5: 19-27.
- Harrison C.J.O. & Walker C.A. (1979b): Birds of the British Lower Oligocene. – In: Harrison C.J.O. & Walker C.A. (eds.): *Studies in Tertiary avian palaeontology*. – *Tertiary Research, Special Paper* 5: 29-43.
- Harrison C.J.O. & Walker C.A. (1979c): New names for two Upper Eocene bird genera. – *Tertiary Research* 2: 110.
- Harrison C.J.O. & Walker C.A. (1982): Fossil birds from the Upper Miocene of northern Pakistan. – *Tertiary Research* 4: 53-69.
- Hassenkamp E. (1858): Geognostische Beschreibung der Braunkohlenformation in der Rhön. – *Verhandlungen der Physicalisch-Medicinischen Gesellschaft in Würzburg* 8: 185-211.

- Haubold H. & Krumbiegel G. (1984): Typenkatalog der Wirbeltiere aus dem Eozän des Geiseltals. – Halle an der Saale: Martin-Luther-Universität, 67 pp.
- Haushalter C.L. (1855): Merkwürdige fossile Thierüberreste aus der Allgäuer Molasse. – München: Universität München, 16 pp.
- Hay O.P. (1902): On the finding of the bones of the Great Auk (*Plautus impennis*) in Florida. – *Auk* 19: 255-258.
- Hébert E. (1855a): Note sur le tibia du *Gastornis parisiensis*. – *Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris)* 40: 579-582.
- Hébert E. (1855b): Note sur le fémur du *Gastornis parisiensis*. – *Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris)* 40: 1214-1217.
- Hébert E. (1855c): Présentation de plusieurs fossiles remarquables du bassin de Paris. – *Bulletin de la Société Géologique de France* (2) 12: 349-352.
- Hébert E. (1855d): Note sur le fémur de *Gastornis parisiensis*. – *L'Institut (Paris)* 23: 189-190.
- Heer O. (1865): Die Urwelt der Schweiz. – Zürich: Friedrich Schulthess, xxix + 622 pp. [See also Heer 1872, 1876.]
- Heer O. (1872): Le Monde primitif de la Suisse. – Genève, xvi + 801 pp. [Translated from Heer 1865, and edited by I. Demole.]
- Heer O. (1876): The primaeval World of Switzerland. – London. [Translated from Heer 1865 by W. S. Dallas, edited by J. Heywood.]
- Heer O. (1879): Die Urwelt der Schweiz. 2nd rev. ed. – Zürich: Friedrich Schulthess.
- Heinrich W.-D. (1990): Review of fossil arvicolids (Mammalia, Rodentia) from the Pliocene and Quaternary in the German Democratic Republic. – In: Fejfar O. & Heinrich W.-D. (eds.): Evolution, phylogeny and biostratigraphy of arvicolids (Rodentia, Mammalia). – Praha: Geological Survey, & München: Verlag Dr. Friedrich Pfeil.
- Heissig K. (1970): Neue Fundstellen oligozäner Spaltenfaunen im Schwäbisch-Fränkischen Jura. – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 10: 331-350.
- Heissig K. (1973): Oligozäne Vertebraten aus der Spaltenfüllung "Möhren 13" bei Treuchtlingen, Fränkischer Jura. – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 13: 177-182.
- Heissig K. (1978): Fossilführende Spaltenfüllungen Süddeutschlands und die Ökologie ihrer oligozänen Huftiere. – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 18: 237-288.
- Heizmann E.P.J. & Fahlbusch V. (1983): Die mittelmiozäne Wirbeltierfauna von Steinberg (Nördlinger Ries). Eine Übersicht. – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 23: 89-93.
- Heizmann E.P.J. & Hesse A. (1995): Die mittelmiozänen Vogel- und Säugetierreste des Nördlinger Ries (MN 6) und des Steinheimer Beckens (MN 7) – ein Vergleich. – In: Peters D.S. (ed.): *Acta palaeornithologica*. – Courier Forschungsinstitut Senckenberg 181: 171-185.
- Heller F. (1930): Eine Forest-Bed-Fauna aus der Sackdillinger Höhle (Oberpfalz). – *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie (B)* 63: 247-298.
- Heller F. (1932): Fossile Eischalenreste aus der eozänen Braunkohle des Geiseltales. – *Nova Acta Leopoldina (Neue Folge)* 1:55-58.
- Heller F. (1936): Eine oberpliozäne Wirbeltierfauna aus Rheinhessen. – *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie (B)* 76: 99-160.
- Hennig W. (1966): Phylogenetic systematics. – Urbana: University of Illinois Press, 263 pp.
- Hermann J. (1804): Observationes zoologicae quibus novae complures, aliaeque animalium species describuntur et illustrantur [Zoological observations, in which a

- number of new species of different animals are described and illustrated.] – Strasburg: Hammer, 332 pp. [In Latin.]
- Hernández Carrasquilla F. (1994): Addenda al catalogo provisional de yacimientos con aves del cuaternario de la Península ibérica [Addenda to the provisional catalogue of Quaternary avian localities of the Iberian Peninsula.] – *Archaeofauna* 2: 77-92. [In Spanish.]
- Hernández-Pacheco E. (1914a): Los vertebrados terrestres del Mioceno de la Península Ibérica [Terrestrial vertebrates from the Miocene of the Iberian Peninsula.] – *Memorias de la Real Sociedad Española de Historia Natural* 9: 443-488. [In Spanish.]
- Hernández-Pacheco E. (1914b): Mioceno superior de La Puebla de Almuradiel (Toledo) [The Upper Miocene of La Puebla de Almuradiel (Toledo).] – *Boletín de la Real Sociedad Española de Historia Natural* 14: 274-278. [In Spanish.]
- Hernández-Pacheco E. (1929): Pistas de aves fósiles en el Oligoceno de Peralta de la Sal (Lérida) [Footprints of fossil birds in the Oligocene of Peralta de la Sal (Lérida).] – *Memorias de la Real Sociedad Española de Historia Natural* 15: 379-382. [In Spanish.]
- Hernández-Pacheco E. & Dantin J. (1915): Geología y paleontología del Mioceno de Palencia [Geology and paleontology of the Miocene of Palencia.] – *Comisión de Investigaciones Española de Historia Natural* 15: 379-382. [In Spanish.]
- Hescheler K. & Kuhn E. (1949): Die Tierwelt. – In: Tschumi O. (ed.): *Urgeschichte der Schweiz*: 121-368. Frauenfeld: Huber & Co.
- Hesse A. (1988a): Die Messelornithidae - Eine neue Familie der Kranichartigen (Aves: Gruiformes: Rhynocheti) aus dem Tertiär Europas und Nordamerikas. – *Journal für Ornithologie* 129: 83-95.
- Hesse A. (1988b): Taxonomie der Ordnung Gruiformes (Aves) nach osteologischen morphologischen Kriterien unter besonderer Berücksichtigung der Messelornithidae HESSE 1988. – *Courier Forschungsinstitut Senckenberg* 107: 235-247.
- Hesse A. (1990): Die Beschreibung der Messelornithidae (Aves: Gruiformes: Rhynocheti) aus dem Alttertiär Europas und Nordamerikas. – *Courier Forschungsinstitut Senckenberg* 128: 1-176.
- Hesse A. (1991): Die Messelrallen – Rekonstruktion eines Lebensbildes. – *Natur und Museum* 119: 110-112.
- Hesse A. (1992): A new species of *Messelornis* (Aves: Gruiformes: Messelornithidae) from the middle Eocene Green River Formation. – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Pierce Brodkorb*. – Natural History Museum of Los Angeles County, Science Series 36: 171-178.
- Hesse A. & Habersetzer J. (1993): Intraspecific development of various foot- and wing-proportions of *Messelornis cristata* (Aves: Gruiformes: Messelornithidae). – *Kaupia* 3: 41-53.
- Hill A. & Walker A. (1979): A fossil marabout (Aves, Ciconiidae) from the Miocene Ngorora Formation, Baringo District, Kenya. – *Netherlands Journal of Zoology* 29: 215-220.
- Hirsch K.F. & Packard M.J. (1987): Review of fossil eggs and their shell structure. – *S.E.M.* 1: 383-400.
- Hitchcock E. (1836): Ornithichnology. – *American Journal of Science* 29: 307-340.
- Hobson K.A. & Montevecchi W.A. (1991): Stable isotope determination of trophic relationships of Great Auks. – *Oecologia* 87: 528-531.
- Hoch E. (1975): Amniote remnants from the eastern part of the Lower Eocene North Sea basin. – In: *Problèmes actuels de paléontologie (Évolution des vertébrés)*. – *Colloques Internationaux du Centre National de la Recherche Scientifique* 218: 543-562.
- Hoch E. (1980): A new middle Eocene shorebird (Aves: Charadriiformes, Charadrii) with columboid features. – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Hildegard Howard*. – Natural History of Los Angeles County, Contributions in Science

- 330: 33-79.
- Hoch E. (1988): On the ecological role of an Eocene bird from Messel, West Germany. – Courier Forschungs-Institut Senckenberg 107: 249-261.
- Hoch E. (1997): Notes on palaeornithology and on a new bird from the early Tertiary North Sea region. – Geological Society of Denmark, Online Series 1: unpaginated. [<http://home4.inet.tele.dk/dgfth/online/hoch.htm>]
- Hoeninghaus F.W. (1839): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1839: 70-71. [In German.]
- Hoerschelmann H. (1966): Allometrische Untersuchungen an Rumpf und Flügel von Schnepfenvögeln (Charadriidae und Scolopacidae). – Zeitschrift für Zoologische Systematik und Evolutionsforschung 4: 209-317.
- Hoff K.E.A. von (1801): Ueber die Ornitholithen. – Magazin für Gesamte Mineralogie 1: 283-302.
- Hoffmann A. (1982): Punctuated versus gradual mode of evolution: a reconsideration. – Evolutionary Biology 15: 411-436.
- Hondt J.-L. d' (1987): Concepts et critères de famille et superfamille chez les Métazoaires. – Année Biologique 26: 141-175.
- Hooker J.J., Insole A.N., Moody R.T.J., Walker C.A. & Ward D.J. (1980): The distribution of cartilaginous fish, turtles, birds and mammals in the British Paleogene. – Tertiary Research 3: 1-45.
- Hopson J.A. (1964): *Pseudodontornis* and other large marine birds from the Miocene of South Carolina. – Postilla 83: 1-19.
- Horáček I. (1981): Comments on the lithostratigraphic context of the early Pleistocene mammal biozones of Central Europe. – In: Quaternary glaciations in the northern hemisphere (IGCP project 73/1/24), Report 6: 99-117. Praha: IGCP.
- Horáček I. (1985): Survey of the fossil vertebrate localities Včeláre 1-7. – Časopis pro Mineralogii a Geologii 30: 353-366.
- Horáček I. & Ložek V. (1988): Palaeozoology and the mid-European Quaternary past: scope of the approach and selected results. – Rozpravy Československé Akademie Věd, Řada Matematických a Přírodních Věd 98 (4): 1-102.
- Hörmann K. (1913): Der Hohle Fels bei Happurg. – Abhandlungen der Naturhistorischen Gesellschaft in Nürnberg 20: 21-63.
- Hou Lian-Hai (1980): [New form of the Gastornithidae from the Lower Eocene of the Xichuan, Honan.] – Vertebrata Palasiatica 18: 111-115. [In Chinese.]
- Houde P. (1981): Paleognathous carinate birds from the early Tertiary of North America. – Science 214: 1236-1237.
- Houde P. (1986): Ostrich ancestors found in the Northern Hemisphere suggest new hypothesis of ratite origins. – Nature (London) 324: 563-565.
- Houde P. (1988): Paleognathous birds from the early Tertiary of the Northern Hemisphere. – Publications of the Nuttall Ornithological Club 22: 1-148. [For a review see Witmer 1989.]
- Houde P. & Haubold H. (1987): *Palaeotis weigelti* restudied: a small middle Eocene ostrich (Aves: Struthioniformes). – Palaeovertebrata 17: 27-42.
- Houde P. & Olson S.L. (1989): Small arboreal non-passerine birds from the early Tertiary of western North America. – In: Ouellet H. (ed.): Acta XIX congressus internationalis ornithologici: 2030-2036. Ottawa: University of Ottawa Press.
- Houde P. & Olson S.L. (1992): A radiation of coly-like birds from the Eocene of North America (Aves: Sandcoleiformes new order). – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County, Science Series 36: 137-160.
- Howard H. (1949): New avian records for the Pliocene of California. – Carnegie Institute Washington, Publication 584: 177-199.

- Howard H. (1964): Fossil Anseriformes. – In: Delacour J. (ed.): The waterfowl of the world. Vol. 4: 233-326. London: Country Life.
- Howard H. (1971): Pliocene avian remains from Baja California. – Los Angeles County Museum, Contributions in Science 217: 1-17.
- Howgate M.E. (1986): The German «Ostrich» and the molecular clock. – Nature (London) 324: 516.
- Hull D.L. (1970): Morphospecies and biospecies: a reply to Ruse. – British Journal of the Philosophy of Science 21: 280-282.
- Hufthammer A.K. (1982): Geirfuglens utbredelse og morfologiske variasjon i Skandinavia [Distribution and morphological variability of the Great Auk in Scandinavia.] – Unpub. Thesis, University of Bergen, Norway. [In Norwegian.]
- ICZN (1955): Direction 16. – Opinions and Declarations Rendered by the International Commission on Zoological Nomenclature 1: 81-88.
- ICZN (1961): International code of zoological nomenclature. – London: International Trust for Zoological Nomenclature, xviii + 176 pp.
- ICZN (1985): International code of zoological nomenclature. – London: International Trust for Zoological Nomenclature, xx + 338 pp.
- ICZN (1999): International code of zoological nomenclature. 4th ed. – London: International Trust for Zoological Nomenclature, xxix + 306 pp.
- Illies H. (1941): Das insektenführende Alttertiär von Havighorst. – Zeitschrift für Geschiebeforschung und Flachlandgeologie 17: 17-24.
- Illies H. (1942): Über das Eozän von Havighorst bei Hamburg und seine stratigraphische Stellung. – Zeitschrift für Geschiebeforschung und Flachlandgeologie 18: 42-48.
- Imbrie J. (1957): The species problem with fossil animals. – In: Mayr E. (ed.): The species problem: 125-153. New York: American Association for the Advancement of Science.
- Inger R.F. (1958): Comments on the definition of genera. – Evolution 12: 370-384.
- Jan'kova V.S. (1959): Paleontologičeskie nachodki iz Liveneckogo pesčanogo kar'era [Paleontological records from the Liveneck sand quarry.] – Izvestija Rostovskogo Oblastnogo Muzeja Kraevedenija 1959 (1): 41-49. [In Russian.]
- Jánossy D. (1963): Die alt-pleistozäne Wirbeltierfauna von Kövesvár bei Répáshuta (Bükk-Gebirge). Annales Historico-Naturales Musei Nationalis Hungarici 55: 109-141.
- Jánossy D. (1964): Letztinterglaziale Vertebratenfauna aus der Kálmán Lambrecht Höhle (Bükk-Gebirge, Nordost Ungarn). I. – Acta Zoologica Hungarica 9: 139-197.
- Jánossy D. (1965): Vogelreste aus den altpleistozänen Ablagerungen von Voigtstedt in Thüringen. – Paläontologische Abhandlungen (A) 2: 336-361.
- Jánossy D. (1969): Stratigraphische Auswertung der europäischen mittelpleistozänen Wirbeltierfauna. Teil II. – Berichte der Deutschen Gesellschaft für Geologische Wissenschaften (A) 14: 573-643.
- Jánossy D. (1972): Die mittelpleistozäne Vogelfauna der Stránská skála. – In: Musil R. (ed.): Stránská skála I. – Anthropos (Brno) 20: 35-64.
- Jánossy D. (1974a): Upper Pliocene and Lower Pleistocene bird remains from Poland. – Acta Zoologica Cracoviensia 19: 531-566.
- Jánossy D. (1974b): Die mittelpleistozäne Vogelfauna von Hundsheim (Niederösterreich). – Sitzungsberichte der Österreichischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, Abteilung I, 182: 211-257.
- Jánossy D. (1976a): Plio-Pleistocene bird remains from the Carpathian basin. I. Galliformes. 1. Tetraonidae. – Aquila 82: 13-36.
- Jánossy D. (1976b): Plio-Pleistocene bird remains from the Carpathian basin. I. Galliformes. 2. Phasianidae. – Aquila 83: 29-42.
- Jánossy D. (1976c): Die Felsnische Tarkó und die Vertebratenfauna ihrer Ausfüllung. – Karszt- és Barlangkutatás 8: 3-106.

- Jánossy D. (1976d): Die Revision jungmittelpleistozäner Vertebratenfaunen in Ungarn. – *Fragmenta Mineralogica et Palaeontologica* 7: 29-54.
- Jánossy D. (1977): Plio-Pleistocene bird remains from the Carpathian basin. III. Strigiformes, Falconiformes, Caprimulgiformes, Apodiformes. – *Aquila* 84: 9-36.
- Jánossy D. (1979): Plio-Pleistocene bird remains from the Carpathian basin. IV. Anseriformes, Gruiformes, Charadriiformes, Passeriformes. – *Aquila* 85: 11-39. [Includes also a chapter on the Cuculiformes]
- Jánossy D. (1980a): Plio-Pleistocene bird remains from the Carpathian basin. V. Podicipediformes, Ciconiiformes, Otidiformes, Columbiformes, Piciformes. – *Aquila* 86: 19-33.
- Jánossy D. (1980b): Plio-Pleistocene bird remains from the Carpathian basin. VI. Systematical and geographical catalogue. – *Aquila* 87: 9-22.
- Jánossy D. (1981): Die altpleistozänen Vogelfaunen von Deutsch-Altenburg 2 und 4 (Niederösterreich). – *Beiträge zur Paläontologie von Österreich* 8: 375-391.
- Jánossy D. (1982): The extinct ancestor of *Anser anser* in Europe. – *Aquila* 89: 21-22.
- Jánossy D. (1983a): Die mittelpleistozäne Vogelfauna von Přezletice bei Prag (ČSSR). – In: Heinrich W.-D. (ed.): *Wirbeltier-Evolution und Faunenwandel im Känozoikum*. – Schriftenreihe für Geologische Wissenschaften 19-20: 247-269. Berlin: Akademie-Verlag.
- Jánossy D. (1983b): Die jungmittelpleistozäne Vogelfauna von Hunas (Hartmannshof). – In: Heller F. et al. (eds.): *Die Höhlenruine Hunas bei Hartmannshof (Landkreis Nürnberger Land). Eine paläontologische und urgeschichtliche Fundstelle aus dem Spät-Riß*. – Quartär-Bibliothek 4: 265-288. Bonn: Röhrscheid.
- Jánossy D. (1986): Pleistocene vertebrate faunas of Hungary. – Budapest: Akadémiai Kiadó, & Amsterdam: Elsevier, 208 pp.
- Jánossy D. (1987a): Fossil data on cranes in Hungary. – *Aquila* 93-94: 35-37.
- Jánossy D. (1987b): Ältestpleistozäne Vertebratenfauna von Beremend 15 (Süd-Ungarn). – *Fragmenta Mineralogica et Palaeontologica* 13: 89-96.
- Jánossy D. (1988): [Untitled report.]. – *Society of Avian Paleontology and Evolution, Information Letter* 2: 9-10. [In English.]
- Jánossy D. (1989): Geierfunde aus der Repolusthöhle bei Peggau (Steiermark, Österreich). – *Fragmenta Mineralogica et Palaeontologica* 14: 117-120.
- Jánossy D. (1990a): Vertebrate fauna of site II. – In: Kretzoi M. (ed.): *Vértesszőlős: site, man and culture*: 187-229. Budapest: Akadémiai Kiadó.
- Jánossy D. (1990b): Arviculids from the Lower Pleistocene sites at Beremend 15 and Somssich-hegy 2, Hungary. – In: Fejfar O. & Heinrich W.-D. (eds.): *Evolution, phylogeny and biostratigraphy of arviculids (Rodentia, Mammalia)*: 223-230. Praha: Geological Survey, & München: Verlag Dr. Friedrich Pfeil.
- Jánossy D. (1991): Late Miocene bird remains from Polgárdi (W-Hungary). – *Aquila* 98: 13-35.
- Jánossy D. (1992): Lower Pleistocene bird remains from Beremend, (S-Hungary, Loc. 15. and 16.). – *Aquila* 99: 9-25.
- Jánossy D. (1993): Bird remains from the Upper Miocene (MN 9) of Rudabanya (N-Hungary). – *Aquila* 100: 53-70.
- Jánossy D. (1995): A late Miocene avifauna from Polgárdi, western Hungary. – In: Peters D.S. (ed.): *Acta palaeornithologica*. – Courier Forschungsinstitut Senckenberg 181: 203-206.
- Jánossy D. (1996): Lower Pleistocene vertebrate faunas from the localities 16 and 17 of Beremend (southern Hungary). – *Fragmenta Mineralogica et Paleontologica* 18: 91-102.
- Jánossy D. (1997): Die Vogelreste aus dem Unterpleistozän von Untermassfeld. – In: Kahlke R.-D. (ed.): *Das Pleistozän von Untermassfeld bei Meiningen (Thüringen). Teil 1*: 95-100. Bonn: Dr. Rudolf Habelt.

- Jánossy D. & Kordos L. (1977): Az Osztramos gerinces lelöhelyeinek faunisztikai és karszt-morfológiai áttekintése (1974-ig) [Faunistical and karst-morphological review of the paleontological vertebrate localities of Osztramos (till 1974).] – *Fragmenta Mineralogica et Paleontologica* 8: 39-92. [In Hungarian.]
- Jarman M.R. (1976): Early animal husbandry. – *Philosophical Transactions of the Royal Society of London* 275: 85-97.
- Jehl J.R., Jr. (1966): Subspecies of Recent and fossil birds. – *Auk* 83: 306-307.
- Jekel H. (1857): *Fabricia entomologica*. Vol. 1. – Paris.
- Johannsen W. (1909): *Elemente der exakten Erblchkeitslehre*. – Jena: Gustav Fischer, 516 pp.
- Jollie M. (1977a): A contribution to the morphology and phylogeny of the Falconiformes (part II). – *Evolutionary Theory* 2: 115-208.
- Jollie M. (1977b): A contribution to the morphology and phylogeny of the Falconiformes (part III). – *Evolutionary Theory* 2: 209-300.
- Jollie M. (1977c): A contribution to the morphology and phylogeny of the Falconiformes (part IV). – *Evolutionary Theory* 3: 1-141.
- Jordan R.H. & Olson S.L. (1982): First record of the Great Auk (*Pinguinus impennis*) from Labrador. – *Auk* 99: 167-168.
- Jörg E. (1957): Tierwelt und Landschaft am Höwenegg (Hegau) zur Unterpliozänzeit. – *Zeitschrift für Geschichte, Volkskunde und Naturgeschichte* 2(4): 117-125.
- Jullien R., Guérin C., Huguency M. & Mein P. (1979): Découverte d'un gisement de mammifères du Miocène moyen à Collet-Redon, près Saint-Cannat (Bouches-du-Rhône, France): liste faunistique, implications stratigraphiques et paléogéographiques. – *Géobios* 123: 297-301.
- Junge G.C.A. (1953): Fossil bones of a White-tailed Eagle *Haliaeetus albicilla* (L.) from the Tegelen clay. – *Proc. Bulletin van der Koninklijke Akademie de Wetenschappen* 56: 285-290.
- Jurcsák T. & Kessler E. (1973): Cercetari paleornitologice din România [Paleornithological research in Romania.] – *Nymphaea* 1: 263-300. [In Romanian.]
- Jurcsák T. & Kessler E. (1986): Evolutia avifaunei pe teritoriul Romanei. Partea I: Introducere [Evolution of the avifauna in the territory of Romania. Part I: Introduction.] – *Crisia* 16: 577-615. [In Romanian.]
- Jurcsák T. & Kessler E. (1987): Evolutia avifaunei pe teritoriul Romanei. Partea II: Morfologia speciilor fosile [Evolution of the avifauna in the territory of Romania. Part II: Morphology of fossil species.] – *Crisia* 17: 583-609. [In Romanian.]
- Jurcsák T. & Kessler E. (1988): Evolutia avifaunei pe teritoriul Romanei. Partea III: Filogenie și sistematice [Evolution of the avifauna in the territory of Romania. Part III: Phylogeny and systematics.] – *Crisia* 18: 647-688. [In Romanian.]
- Karchu A.A. (1988): Novoe semejstvo strižeobraznyh iz Paleogena Evropy [A new apodiform family from the Paleogene of Europe]. – *Paleontologičeskij Žurnal* 1988 (3): 78-88. [In Russian.]
- Karchu A.A. (1990): Srednegolocenovye pticy iz archeologičeskich pamjatnikov Podmoskov'ja [Middle Holocene birds from the archeological monuments of Podmoskov'e]. – *Ornitologija* 24: 67-71. [In Russian.]
- Karchu A.A. (1992): Morphological divergence within the order Apodiformes as revealed by the structure of the humerus. – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Pierce Brodkorb*. – Natural History Museum of Los Angeles County, Science Series 36: 379-384.
- Karchu A.A. (1997): Novyj vid urmiornisa (Gruiformes: Ergilornithidae: *Urmiornis*) iz rannego miocena Zapadnogo Kazachstana [A new urmiornis species (Gruiformes: Ergilornithidae: *Urmiornis*) from the early Miocene of western Kazakhstan.] – *Paleontologičeskij Žurnal* 1997 (1): 104-110. [In Russian.]

- Karchu A.A. (1999): A new genus and species of the family Jungornithidae (Apodiformes) from the late Eocene of the northern Caucasus, with comments on the ancestry of hummingbirds. – In: Olson S.L. (ed.): Avian paleontology at the close of the 20th century. – Smithsonian Contributions to Paleobiology 89: 207-216.
- Karg J.-M. (1805): Über den Steinbruch zu Oehningen bey Stein am Rhein und dessen Petrefakte. – Denkschrift der Vaterländischen Aerzte und Naturforscher Schwabens 1: 1-74.
- Kašin G.N. (1976): Novoe imja dlja preokkupirovannogo *Apopemopsis* Brodkorb 1971 [New name for the preoccupied *Apopemopsis* Brodkorb 1971.] – Ornitologija 12: 231. [In Russian.]
- Kašin G.N. (1977): Novoe rodovoe nazvanie ptic vzamen preokkupirovannogo. [New genus name of birds in replacing a preoccupied.] – Paleontologičeskij Žurnal 1977 (3): 139. [In Russian.]
- Kašin G.N. (1978): Zamena preokkupirovannogo nazvanija roda v klasse ptic [Replacement of a preoccupied genus name in the class of birds.] – Paleontologičeskij Žurnal 1978 (1): 145-146. [In Russian.]
- Katinas V. (1971): Jantar' i jantar'nye otloženija južnoj Pribaltiki [Amber and amber-bearing deposits of the southern Baltic area.] – Trudy Instituta Geologii (Vilnius) 20: 1-156. [In Russian.]
- Katinas V. (1983): Baltijos gintaras [Baltic amber.] – Vilnius: Mosklas, 112 pp. [In Lithuanian.]
- Keferstein C. (1834): Die Naturgeschichte des Erdkörpers in ihren ersten Grundzügen. – Leipzig: Friedrich Fleischer, x + 896 pp.
- Keller B. (1982): Fossile Vogelfährten aus der Luzerner Molasse. – Mitteilungen der Naturforschenden Gesellschaft Luzern 27: 97-109.
- Kerourio P. & Aujard C. (1987): Les oeufs d'Oiseaux géants du Midi de la France. – Monde et Minéraux 78: 12-16.
- Kessler E. (1975): Contributii noi la studiul avifaunei fosile de Betfia (Jud. Bihor) [Contributions to the study of the fossil avifauna from Betfia (Dept. Bihor).] – Nymphaea 4: 133-138. [In Romanian.]
- Kessler E. (1976): Prezența genului *Gallus* in avifauna fosila a României [Presence of the genus *Gallus* in the fossil avifauna of Romania.] – Nymphaea 4: 133-138. [In Romanian.]
- Kessler E. (1977): Avifauna zonei Crisurilor din perioada Pliocenului pina astazi [Avifauna of the Crișu region from the Pliocene period till today.] – Unpub. Thesis, University of Bucuresti. [In Romanian.]
- Kessler E. (1978): Imbogătirea inventarului paleoavifaunistic al României [Addenda to the list of the paleoavifauna of Romania.] – Nymphaea 6: 165-180. [In Romanian.]
- Kessler E. (1979): Noi date paleornitologice din Pliocenul Romanei [New data on the paleornithology of the Romanian Pliocene.] – Nymphaea 7: 135-138. [In Romanian.]
- Kessler E. (1984a): On some bird remains from the Pliocene of Malusteni in the Laboratory of Paleontology, University of Bucharest. – In: 75 years Laboratory of Paleontology, pp. 287-293. București.
- Kessler E. (1984b): Noi contribuții privind studiul avifaunelor din Paratethys [New contributions to the study of the Paratethys avifauna.] – Crisia 14: 521-532. [In Romanian.]
- Kessler E. (1985): Contribuții noi la studiul avifaunelor cuaternare din România [Contributions to the study of the Quaternary avifauna of Romania.] – Crisia 15: 485-491. [In Romanian.]
- Kessler E. (1986): Fossil bird remains from Romania in context of the avifauna from Eocene. – Studia Universitatis Babeș-Bolyai, Series Geologia-Geographia 31: 75-77. [A list of the "representative genera of the Eocene" is presented in this paper on p. 76,

- where many of the names are misspelled. I do not list these subsequent incorrect spellings in the index, but I list here the European ones with explanation to avoid future misunderstandings: *Boturoides* → *Botaurites*, *Aquilagus* → *Aquilavus*, *Filhornis* → *Filholornis*, *Dyatrma* → *Diatryma*, *Geiselavus* → *Geiseloceros*.]
- Kessler E. (1992): Review of the Neogene waterfowl (Aves: Anatidae) of eastern Paratethys. – *Studia Universitatis Babes-Bolyai, Series Biologia* 37: 47-54.
- Kessler E. (1993): Prezentarea [sic!, should be Prezența] Galinaceelor (Ord. Galliformes, Cl. Aves) printre resturile scheletice colectate din situri arheologice de pe teritoriul României [The presence of galliforms (Ord. Galliformes, Cl. Aves) according to the skeletal remains found in the archeological sites from the territory of Romania.] – *Acta Musei Napocensis* 26-30 (Prehistorie – Istorie Veche – Arheologie 1): 205-219. [In Romanian.]
- Kessler E. (1996): Tertiary avian localities of Romania. – Mlíkovský J. (ed.): Tertiary avian localities of Europe. – *Acta Universitatis Carolinae, Geologica* 39: 703-709.
- Kessler E. & Codrea V. (1996): A bird from the middle Miocene (Badenian) of Zarand Depression (western Romania). – *Studia Universitatis Babes-Bolyai, Series Geologia* 41: 127-130.
- Kessler E. & Gál E. (1996): New taxa in the Neogene bird fauna from eastern Paratethys. – *Studia Universitatis Babes-Bolyai, Series Biologia* 41: 73-79.
- Kessler E. & Moldvai M. (1993): Studii biometrice și morfologice asupra scheletului membrilor la Corvidae (Aves) (Partea I) [Biometrical and morphological studies of the limb skeleton of the Corvidae (Aves) (Part 1).] – *Studia Universitatis Babes-Bolyai, Series Biologia* 38: 69-80. [In Romanian.]
- Kessler E., Codrea V. & Vremir M. (1998): A fossil bird from the Lower Oligocene of Cluj-Napoca (western Romania). – *Studia Universitatis Babes-Bolyai, Series Geologia* 43 (2): 7-12.
- Killermann S. (1911): Der Waldrapp Gesners, neue Zeugnisse für sein ehemaliges Vorkommen in Mitteleuropa. – *Zoologischer Anzeiger* 1911: 271.
- King A.T. (1845): Description of fossil foot marks, supposed to be referable to the classes of birds, Reptilia and Mammalia, found in the Carboniferous series, in Westmoreland County, Pennsylvania. – *Proceedings of the Academy of Natural Sciences of Philadelphia* 2: 175-180.
- King P.P. (1828): On the animals of the Straits of Magellan. – *Zoological Journal (London)* 4: 91-107.
- Kiriakoff S.G. (1967): On the nomenclature of the superspecies. – *Systematic Zoology* 30: 425-458.
- Kitcher P. (1984): Species. – *Philosophy of Science* 51: 308-333.
- Klähn H. (1920): Der Wert der Variationsstatistik für die Paläontologie. – *Berichte der Naturforschenden Gesellschaft in Freiburg i. Br.* 22: 7-224.
- Kleinschmidt A. (1953a): *Torgos tracheliotus todei* forma nova, ein mitteleuropäischer fossiler Ohrengerier aus dem Diluvium von Salzgitter-Lebenstedt. – *Bonner Zoologische Beiträge* 4: 21-30.
- Kleinschmidt A. (1953b): Die zoologischen Funde der Grabung Salzgitter-Lebestedt 1952. – *Eiszeitalter und Gegenwart* 3: 166-188.
- Knowles W.J. (1895): Remains of the Great Auk from White-park Bay, County Antrim. – *Proceedings of the Irish Academy* (3) 3: 650-663.
- Kohring R., Hirsch K.F. (1996): Crocodylian and avian eggshells from the Middle Eocene of the Geiseltal, Eastern Germany. – *Journal of Vertebrate Paleontology* 16: 67-80.
- Kolar-Jurkovšek T. & Jurkovšek B. (1996): Prva najdba terciarne ptice v Sloveniji [First find of a Tertiary bird in Slovenia.] – *Rozprave IV. Razreda SAZU* 37: 23-35. [In Slovenian.]
- Kompanje E.O. & Kerkhoff N.C. (1991): [Find of coracoid of Great Auk on Maasvlakte

- in April 1981.] – Dutch Birding 13: 96-98. [In Dutch.]
- König C., Weick F. & Becking J.-H. (1999): Owls: a guide to the owls of the world. – Mountfield: Pica Press, 462 pp.
- König E. (1820-1825): Icones fossilium sectiles [Collected figures of fossils.] – London, 4 pp. + 19 pls. [In Latin.] [The date of publication of this booklet is sometimes given as 1938, but “no more of the work was published after 1825” according to the on-line catalogue of the Natural History Museum, London (stand: May 2000).]
- Kordos L. (1983): Lábynomok az Ipolytarnóci alsó-miocén korú homokkőben [Footprints from the early Miocene sandstone of Ipolytarnóc.] – Geologia Hungarica, Series Paleontologica 46: 257-415. [In Hungarian.]
- Kormos T. (1911): Die pleistocäne Fauna der Felsnische Puskaporos bei Hámor. – Mitteilungen aus dem Jahrbuch der Kgl. Ungarischen Geologischen Reichsanstalt 19: 148-154.
- Kormos T. (1913): Kleinere Mitteilungen aus dem ungarischen Pleistocän. – Centralblatt für Mineralogie, Geologie und Paläontologie 1913: 13-17.
- Korotkevič E.L. (1961): Do vyvčennja giparionovoj fauny dolyny r. Kujal'nika [On the study of the Hipparion fauna of Kujal'nik river valley.] – Zbirnyk Prac' Zoologičnoho Muzeju 30: 12-129. [In Ukrainian.]
- Korotkevič, E.L. (1972): Èkologičeskaja charakteristika gipparionovych faun juga SSSR [Ecological characteristics of the Hipparion faunas of souther USSR.] – In: David, A. I. (ed.): Pozvonočnye neogena i plejstocena Moldavii [Neogene and Pleistocene vertebrates of Moldova]. Chişinău: Ştiinţa. [In Russian.]
- Kotsakis T. (1980): Osservazioni sui vertebrati quaternari della Sardegna [Observations on the Quaternary vertebrates of Sardinia.] – Bolletino della Societa Geologica Italiana 99: 151-165. [In Italian.]
- Kozlova E.V. (1960): Novye iskopaemye pticy iz jugo-vostočnoj Gobi [New fossil birds from southeastern Gobi.] – Trudy Problemnyh i Tematičeskich Soveščanj Zoologičeskogo Instituta Akademii Nauk SSSR 9: 323-329. [In Russian.]
- Kraft E. (1972): Vergleichend morphologische Untersuchungen an Einzelknochen nord- und mitteleuropäischer kleinerer Hühnervogel. – Unpub. Diss., München, Universität München, 195 pp.
- Kraglievich L. (1932): Una gigantesca ave fósil del Uruguay. *Devincenzia gallinali* n. gen. sp., tipo del una nueva familia Devincenziidae del orden Stereornithes [A gigantic fossil bird from Uruguay. *Devincenzia gallinali* n. gen. sp., type of a new family Devincenziidae of the order Stereornithes.] – Anales del Museo de Historia Natural de Montevideo (2) 3: 323-353. [In Spanish.]
- Kraglievich, L. (1946): Noticia preliminar acerca de un nuevo y gigantesco estereornito de la fauna Chapadmalense [A preliminary note on a new and gigantic stereornithid from the Chapadmalense fauna.] – Anales de la Sociedad des Ciencias de Argentina 142: 104-121. [In Spanish.]
- Krejčí J. (1877): Geologie [Geology.] – Praha: Jan Krejčí, 1042 pp. [In Czech.]
- Kretzoi M. (1941): Ósemlős maradványok Betfiáról – Die unterpleistozäne Säugetierfauna von Betfia bei Nagyvárad. – Földtani Közlöny 71: 308-335. [In Hungarian, with German translation.]
- Kretzoi M. (1954a): Ostrich and camel remains from the central Danube basin. – Acta Geologica 2: 231-242.
- Kretzoi M. (1954b): Befejező jelentés a Csakvári barlang őslénytani feltásáról [Final report on the paleontological findings in the cave of Csákvár.] – A Magyar Állami Földtani Intézet Évi Jelentése az 1952 Évről: 37-69. [In Hungarian.]
- Kretzoi M. (1955a): Strucc-maradványok a Fejér megyei Kislángról [Ostrich remains from Kisláng in the Fejér Province.] – Aquila 59-62: 361-366. [In Hungarian.]
- Kretzoi M. (1955b): *Pliogallus* Gaillard 1939 és *Pliogallus* Tugarinov 1940 [*Pliogallus*

- Gaillard 1939 and *Pliogallus* Tugarinov 1940.] – *Aquila* 59-62: 367. [In Hungarian.]
- Kretzoi M. (1956): A Villányi hegység alsó-pleisztocén gerinces-faunái – Die altpleistozänen Wirbeltierfaunen des Villányer Gebirges. – *Geologica Hungarica* 27: 1-264. [In Hungarian, with German translation.]
- Kretzoi M. (1957): Madármaradványok a Csákvári-faunából [Avian remains in the fauna of Csákvár.] – *Aquila* 63-64: 239-248. [In Hungarian.]
- Kretzoi M. (1962): Madár-maradványok a betfiai alsópleisztocén faunából [Remains of birds in the early Pleistocene fauna of Betfia.] – *Aquila* 67-68: 167-174. [In Hungarian.]
- Kretzoi M. (1975): Fontosabb szórványleletek a máfi gerinces gyűjteményében (2. közlemény) [Important records in the collection of fossil vertebrates of the Hungarian Geological Survey (2nd installment.)] – *A Magyar Állami Földtani Intézet Évi Jelentése az 1975 Évről*: 401-415. [In Hungarian.]
- Kretzoi M. (1977): The fauna of small vertebrates of the Middle Pleistocene at Petralona. – *Anthropos* (Athena) 4: 131-143.
- Kretzoi M. & Krolopp E. (1977): Alsó pleisztocén-végi puhatestű és gerinces fauna a Kőröshegyi téglagyár (Balatonföldvár) feltárásából [An early Pleistocene mollusc and vertebrate fauna from Kőröshegy brickyard (Balatonföldvár).] – *A Magyar Állami Földtani Intézet Évi Jelentése az 1975 Évről*: 369-382. [In Hungarian.]
- Kretzoi M. & Poulianos N.A. (1981): Remarks on the Middle and Lower Pleistocene vertebrate fauna in the Petralona cave. – *Anthropos* (Athena) 8: 57-72.
- Kristoffersen A.V. (1997): New records of perching birds from the latest Paleocene/earliest Eocene Fur Formation of Denmark. – *Geological Society of Denmark, Online Series* 1: unpaginated. [http://home4.inet.tele.dk/dgfh/online/anette.htm]
- Kristoffersen A.V. (1999): Cranial anatomy of an early Paleogene trogon (Aves; Trogoniformes) from northern Europe. – In: IV European workshop on vertebrate paleontology, Abstracts: 55. Albarracín, Spain.
- Krüger J.F. (1823): *Geschichte der Urwelt*. Vol. 2. – Quedlinburg: Gottfried Basse, viii + 967 pp.
- Krumbiegel G. (1959): Die terziären Pflanzen- und Tierwelt der Braunkohle des Geiseltales. – *Die Neue Brehm-Bücherei* 237: 1-156. Wittenberg Lutherstadt: A. Ziemsen Verlag.
- Krumbiegel G. (1962): Die Fossilfundstellen der mitteleozänen Braunkohle des Geiseltales. – *Wissenschaftliche Zeitschrift der Martin-Luther-Universität in Halle* 11: 173-208.
- Krumbiegel G. (1966): Zur Stratigraphie der Fossilfundstellen (Pflanzenfundstellen und Ergänzungen zu den Wirbeltierstellen) im Eozän des Geiseltales. – In: Krumbiegel G. & Schmidt W. (eds.): *Das Geiseltal*: 41-63. Berlin: Deutsche Gesellschaft für Geologische Wissenschaften.
- Krumbiegel G. (1968): Die Fossilfundstellen (Pflanzenfundstellen und Wirbeltierfundstellen) im Eozän des Geiseltales, ihre stratigraphische Bedeutung und ihr Fossilinhalt. – In: Krumbiegel G. & Schmidt W. (eds.): *Das Geiseltal*: 57-85. 2nd rev. ed. Berlin: Deutsche Gesellschaft für Geologische Wissenschaften.
- Krumbiegel G. (1977): Genese, Palökologie und Biostratigraphie der Fossilfundstellen im Eozän des Geiseltales. – In: Matthes H.W. & Thaler B. (eds.): *Eozäne Wirbeltiere des Geiseltales*. – Martin-Luther-Universität Halle-Wittenberg, *Wissenschaftliche Beiträge* 2: 113-138.
- Krumbiegel G., Rufflé L. & Haubold H. (1983): *Das eozäne Geiseltal*. 2nd rev. ed. – *Die Neue Brehm-Bücherei* 237: 1-227. Wittenberg Lutherstadt: A. Ziemsen Verlag.
- Krupska A. (1978): Morfologia kośćca kury domowej na podstawie wczesnośredniowiecznych materiałów archeologicznych z terenu Śląska [Morphology of bones of the domestic chicken on the basis of early Medieval archaeological material

- from the territory of Silesia.] – Unpub. Diss., Wrocław. [In Polish.]
- Kuhn O. (1958): Ein neuer Lacertilier aus dem fränkischen Lithographieschiefer. – Neues Jahrbuch für Geologie und Paläontologie, Monatshefte 1958: 380-382.
- Kuhn O. (1963): Ichnia tetrapodorum. – In: Westphal F. (ed.): Fossilium catalogus. I. Animalia. Pars 101. s'-Gravengahe: Dr. W. Junk, 176 pp.
- Kukla J. (1975): Loess stratigraphy of Central Europe. – In: Butzer K.W. & Isaac G.L. (eds.): After the Australopithecines: 99-188. Haag: Mouton.
- Kukla J. (1977): Pleistocene land-sea correlations: I. Europe. – Earth-Science Reviews 13: 307-374.
- Kukla J. (1978): The classical European glacial stages: correlation with deep-sea sediments. – Transactions of the Nebraska Academy of Sciences 6: 57-93.
- Kumerloeve H. (1978): Waldrapp, *Geronticus eremita* (Linnaeus, 1758) und Glattnackentrapp, *Geronticus calvus* (Boddaert, 1783): zur Geschichte ihrer Erforschung und zur gegenwärtigen Bestandssituation. – Annalen des Naturhistorischen Museums in Wien 81: 319-349.
- Kumerloeve H. (1983): Zur Kenntnis altägyptischer Ibis-Darstellungen, unter besonderer Berücksichtigung des Waldrapps, *Geronticus eremita* (Linnaeus, 1758). – Bonner Zoologische Beiträge 34: 197-234.
- Kumerloeve H. (1984): The Waldrapp, *Geronticus eremita* (Linnaeus, 1758): historical review, taxonomic history, and present status. – Biological Conservation 30: 363-373.
- Kuntz P. (1983): Oeufs fossilisés de Reptiles et d'Oiseaux dans le gisement lutétien de Bouxwiller (Dpt. Bas-Thin/France). – Mitteilungen des Badenschen Landesvereins für Naturkunde und Naturschutz 13: 127-134.
- Kuročkin E.N. (1968): Novye oligocenovye pticy Kazachstana [New Oligocene birds from Kazakhstan.] – Paleontologičeskij Žurnal 1968 (1): 92-101. [In Russian.]
- Kuročkin E.N. (1976): Novye dannye o pticach pliocena zapadnoj Mongolii [New data on the birds of the Pliocene of western Mongolia.] – In: Paleontologija i biostratigrafija Mongolii [Paleontology and biostratigraphy of Mongolia]: 51-67. Moskva: Nauka. [In Russian.]
- Kuročkin E.N. (1980): Middle Pliocene rails from western Mongolia. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Hildegard Howard. – Natural History Museum of Los Angeles County, Contributions in Science 330: 69-73.
- Kuročkin E.N. (1981): Novye predstaviteli i evoljucija dvuch semejstv archaičnych žuravleobraznyh v Evrazii [New representatives and evolution of two families of archaic Gruiformes in Eurasia.] – In: Tatarinov L.P. (ed.): Iskopaemye pozvonočnye Mongolii [Fossil vertebrates of Mongolia]: 59-85. Moskva: Nauka. [In Russian.]
- Kuročkin E.N. (1982): On the evolutionary pathways of didactylous Tertiary gruids under increasing aridization. – In: Novák V.J.A. & Mlíkovský J. (eds.): Evolution and environment: 731-736. Praha: ČSAV.
- Kuročkin E.N. (1985): Pticy central'noj Azii v Pliocene [Birds of Central Asia in the Pliocene.] – Trudy Sovmestnoj Sovetsko-Mongol'skoj Paleontologičeskoj Ekspedicii 26: 1-120. [In Russian.]
- Kuročkin E.N. & Anorova N.S. (1977): K probleme proischoždenija domašnich kur. Svrnvenie skeletov domašnich i bankivskoj kur [On the problem of the origin of domestic chicken. A comparison of skeletons of domestic chicken and the Red Junglefowl.] – Zoologičeskij Žurnal 56: 1511-1521. [In Russian.]
- Kuročkin E.N. & Chozackij L.I. (1972): Gricaja iz rusil'onskoj fauny Moldavii i Ukrainy [Gryzaja from Roussilsonian fauna of Moldova and southern Ukraine.] – Ornitologija 10: 347-349. [In Russian.]
- Kuročkin E.N. & Ganea I.M. (1972): Pticy srednego sarmata Moldavii [Birds of the middle Sarmatian of Moldova.] – In: David A.I. (ed.): Pozvonočnye neogena i plejstocena Moldavii [Vertebrates of the Neogene and Pleistocene of Moldova]: 45-70.

- Chişinău: Ştiinca. [In Russian.]
- Kuročkin E.N. & Lungu A.N. (1970): Novyj straus iz srednego sarmata Moldavii [A new ostrich of the middle Sarmatian of Moldova.] – Paleontologičeskij Žurnal 1970 (1): 118-126. [In Russian.]
- Kuss S.E. (1960): Eine neuentdeckte Säugetierfauna aus dem Aquitan von Büchelberg in der Pfalz. – Notizblatt des Hessischen Landesamts für Bodenforschung 88: 29-62.
- La Barbera M. (1989): Analyzing body size as a factor in ecology and evolution. – Annual Reviews of Ecology and Systematics 20: 97-117.
- Lageræus P. (1989): [What do we know of the Great Auk and its occurrence in Sweden?] – Anser 28: 141-150. [In Swedish.]
- Laizer L. de (1828): Note sur l'existence d'ossements fossiles dans le tuf volcanique ou Pépérino d'Auvergne. – Annales des Sciences Naturelles (Paris) 15: 415-420.
- Lamanon R. de Paul de (1782): Description des divers fossiles trouvés à Montmartre, près Paris, et vues générales sur l'origine des pierres gypseuses. – Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts 19: 173-194.
- Lamanon R. de Paul de (1783): [Untitled letter to the editor.] – Journal de Physique, de Chimie, d'Histoire Naturelle et des Arts 22: 309-313. [In French.]
- Lambrecht K. (1912a): A borsodi Bükk fossilis madarai – Fossile Vögel des Borsoder Bükkgebirges. – Aquila 19: 270-287. [In Hungarian, with German translation.]
- Lambrecht K. (1912b): Magyarország fossilis madarai – Die fossilen Vögel Ungarns. – Aquila 19: 288-320. [In Hungarian, with German translation.]
- Lambrecht K. (1916a): Die Gattung *Plotus* im ungarischen Neogen. – Mitteilungen aus dem Jahrbuche der Königlich Ungarischen Geologischen Reichsanstalt 24: 1-24.
- Lambrecht K. (1916b): Az első magyar praeglacialis madárfauna – Die erste ungarische präglaziale Vogelfauna. – Aquila 22: 160-175. [In Hungarian, with German translation.]
- Lambrecht K. (1917): Az európai madárvilág kialakulása – Die Ausbildung und Geschichte der europäischen Vogelwelt. – Aquila 24:191-221. [In Hungarian, with German translation.]
- Lambrecht K. (1921): Aves. – In: Diener C. (ed.): Fossilium catalogus I: Animalia. – Berlin: W. Junk, 104 pp.
- Lambrecht K. (1928): *Palaeotis weigelti* n.g. n.sp., eine fossile Trappe aus der mitteleozänen Braunkohle des Geiseltales. – Jahrbuch des Halleschen Verbandes für die Erforschung der Mitteldeutschen Bodenschätze (Neue Folge) 7: 20-29.
- Lambrecht K. (1929): Mesozoische und tertiäre Vogelreste aus Siebenburgen. – In: Sciki E. (ed.): X^e congrès international de zoologie: 1262-1275. Budapest: Stephaneum.
- Lambrecht K. (1930) Studien über fossile Riesenvögel. – Geologica Hungarica, Series Palaeontologica 7: 1-37.
- Lambrecht K. (1931a): *Cygnopterus* und *Cygnavus*, zwei fossile Schwäne aus dem Tertiär Europas. – Bulletin du Musée Royal d'Histoire Naturelle de Belgique 7 (31): 1-6.
- Lambrecht K. (1931b): Fortschritte der Palaeo-Ornithologie. – In: Proceedings of the VIIth international ornithological congress: 73-99. Amsterdam.
- Lambrecht K. (1933): Handbuch der Palaeornithologie. – Berlin: Gebrüder Borntraeger, 1024 pp.
- Lambrecht K. (1935): Drei neue Vogelformen aus dem Lutétian des Geiseltales. – Nova Acta Leopoldina (Neue Folge) 3: 361-367.
- Lambrecht K. (1938): *Urmiornis abeli* n. sp., eine pliozäne Vogelfährte aus Persien. – Palaeobiologica 6: 242-245.
- Lartet E. (1837): Sur les débris fossiles trouvés à Sansan, et sur les animaux antédiluviens en général. – Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris) 5: 158-159. [See also L'Institut 1837, p. 126.]
- Lartet E. (1855): Note sur le tibia d'oiseau fossile de Meudon. – Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris) 40: 582-584.

- Lartet E. (1857): Note sur un humérus fossile d'oiseau, attribuée à un très-grand palmipède de la section des Longipennes. – *Compte-Rendu des Séances Hebdomadaires de l'Académie des Sciences (Paris)* 44: 736-741.
- Laskarev V.D. (1908): Geologičeskie nabljudenija v okr. g. Tiraspolja [Geological observations in the vicinity of Tiraspol town.] – *Zapiski Novorossijskogo Obščestva Estesvoispytatelej* 33: 1-22. [In Russian.]
- Laskarev V.D. (1912): Zametki o novych mestonachozhdenijach iskopaemych mlekopitajuščich v tercičnyh otloženijach južnoj Rossii [Comments on new localities of fossil mammals in the Tertiary deposits of southern Russia.] – *Zapiski Novorossijskogo Obščestva Estesvoispytatelej* 38: 36-57. [In Russian.]
- Laskarev V.D. (1936): Miocenska fauna kičmenjaka iz okoline sela Kruševica (Bukulja) [A Miocene fauna of vertebrates from the vicinity of Kruševica village (Bukulja).] – *Geološki Anali Balkanskogo Poluostrva* 13: 14-27. [In Serbo-Croatian.]
- Laube G.C. (1901): Synopsis der Wirbeltierfauna der böhmischen Braunkohlenformation. – *Abhandlungen des Deutschen Naturwissenschaftlichen Vereins in Böhmen "Lotos"* Prag 2(4): 1-76.
- Laube G.C. (1909): Ein neuer Vogelrest aus den Tonen von Preschen bei Bilin. – *Lotos* 57: 159-161.
- Laube G.C. (1910): Vogel- und Reptilienreste aus der Braunkohle von Skiritz bei Brüx. – *Lotos* 58: 115-127.
- Launay L. de (1908): La fourrure d'un écureuil tertiaire. – *Nature (Paris)* 36: 393-395.
- Launay L. de (1923): Études sur le plateau central. V. Notes sur le terrain tertiaire de la Limagne bourbonnaise. – *Bulletin des Services de la Carte Géologique de la France* 36.
- Laurent, G. (1987) *Paléontologie et évolution en France*. Paris: Editions du Comité des Travaux Historiques et Scientifiques, xiv + 554 pp.
- Laurillard C.L. (1849): Atlas (Oiseaux fossiles). – In: d'Orbigny C. (ed.): *Dictionnaire universel d'histoire naturelle*, 2nd ed., Atlas 1, pl. 2. Paris: Abel Pilon et Cie.
- Lavater J.H. (1808): Rhapsodische Anmerkungen über einen bei Oeningen gefundenen Ornitholithen. – *Taschenbuch für Mineralogie* 2: 71-77.
- Lavocat A. (1899): Les oiseaux actuels et anciens. – *Bulletin de l'Académie des Sciences, Inscriptions et Belles-Lettres de Toulouse* 2: 26-33.
- Lebedinsky N.G. (1927): *Romainvillia Stehlini* n.g. n.sp., canard éocène provenant des marnes blanches du Bassin de Paris. – *Mémoires de la Société Paléontologique Suisse* 47(2): 1-8.
- Legendre S. & Lévêque F. (1997): Etalonnage de l'échelle biochronologique mammalienne du Paléogène à l'échelle globale. – In: Aguilar J.-P., Legendre S. & Michaux J. (eds.): *Actes du Congrès BiochroM'97*. – *Mémoires et Travaux de l'Institut de Montpellier* 21: 461-473.
- Leisler B. (1980): Morphological aspects of ecological specialization in bird genera. – *Ökologie der Vögel* 2: 199-220.
- Leisler B. (1985): Ecomorphology. – In: Johnston R.F. (ed.): *Current ornithology* 2: 155-186.
- Lemen C.A. & Freeman P.W. (1984): The genus: a macroevolutionary problem. – *Evolution* 38: 1219-1237.
- Lemoine V. (1878): Recherches sur les oiseaux fossiles des terrains tertiaires inférieurs des environs de Reims. – Reims: Imprimerie Keller, 69 pp.
- Lemoine V. (1880): Communication sur les ossements fossiles des terrains inférieurs des environs de Reims. – Reims: Imprimerie et Lithographie Matot-Braine, 17 pp.
- Lemoine V. (1881a): Sur le *Gastornis Edwardsii* et le *Remiornis Heberti* de l'éocène inférieur des environs de Reims. – *Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris)* 93: 1157-1159.
- Lemoine V. (1881b): Résultats des dernières recherches concernant les mammifères et les

- oiseaux de la faune cernaysienne. – Bulletin de la Société Géologique de France (3) 10: 96.
- Lemoine V. (1881c): Recherches sur les oiseaux fossiles des terrains tertiaires inférieurs des environs de Reims. Deuxième partie. – Reims: Imprimerie et Lithographie Matot-Braine, pp. 73-170.
- Lemoine V. (1884): Présentation de pieces du squelette d'*Eupterornis*, de *Gastornis* et de *Pleuraspidotherium*. – Bulletin de la Société Géologique de France (3) 12: 357.
- Lemoine V. (1885): [Untitled report on a lecture.] – Bulletin de la Société Géologique de France (3) 13: 412. [In French.]
- Lemoine V. (1893): Etude sur les os du pied des mammifères de la faune cernaysienne et sur quelques pieces osseuses nouvelles de cet horizon paléontologique. – Bulletin de la Société Géologique de France (3) 21: 353-368.
- Lepsius G.R. (1883): Das Mainzer Becken. – Darmstadt: A. Bergsträsser, 181 pp.
- Lepsius G.R. (1887-1892): Geologie von Deutschland und den angrenzenden Gebiete. 1. Das westliche und südliche Deutschland. – Stuttgart: Engelhorn, 800 pp. [Appeared in three parts: pp. 1-254 in 1887, pp. 255-458 in 1889, and pp. 459-800 in 1892.]
- Liberov P.D. (1959): K voprosu o vremeni odomašnenija kuricy [On the question of the timing of the domestication of the chicken] – Sovetskaja Archeologija 2: 232-234. [In Russian.]
- Lie R. (1986): Animal bones from the late Weichselian in Norway. – Fauna Norvegica (A) 7: 41-46.
- Lierl H.-J. (1993): Ein Vogelrest in einem Flintgeschiebe von Schleswig. – Geschiebekunde Aktuell 9: 3-6.
- Lindner H. (1979): Zur Frühgeschichte des Haushuhns im Vorderen Orient. – Unpublished Thesis, Ludwig-Maximilian-Universität, München, 65 pp.
- Linnaeus C. (1758): Systema naturae [The system of Nature.] 10th rev. ed. – Holmia [= Stockholm]: Salvius, [iv] + 824 pp. [In Latin.]
- Livezey B.C. (1988): Morphometrics of flightlessness in the Alcidae. – Auk 105: 681-698.
- Livezey B.C. (1997a): A phylogenetic classification of waterfowl (Aves: Anseriformes), including selected fossil species. – Annals of Carnegie Museum 66: 457-496.
- Livezey B.C. (1997b): A phylogenetic analysis of basal Anseriformes, the fossil *Presbyornis*, and the interordinal relationships of waterfowl. – Zoological Journal of the Linnean Society 121: 361-428.
- Livezey B.C. & Martin L.D. (1988): The systematic position of the Miocene anamid *Anas*[?] *blanchardi* Milne-Edwards. – Journal of Vertebrate Paleontology 8: 196-211.
- Livezey B.C. & Zusi R.L. (2001): Higher-order phylogenetics of modern Aves based on comparative anatomy. – Netherlands Journal of Zoology 51: 179-206.
- Ljaščuk B.F. (1962): Ridkiska znachidka v Karpatach [A rare find in the Carpathians.] – Visnik L'vivskogo Deržavnogo Universitetu, Serija Heohrafična 1962 (1): 132-133. [In Ukrainian.]
- Löppenthin B. (1952): Ægget fra Hermanstorp. Et sen-glacialt fund fra Skåne [Eggs from Hermanstorp. A late glacial record from Skåne.] – Dansk Ornithologisk Forenings Tidsskrift 46: 12-31. [In Danish.]
- López, N., Sesé C. & Sanz J. L. (1977): La microfauna (Rodentia, Insectivora, Lagomorpha y Reptilia) de las fisuras del Mioceno medio de Escobosa de Catalañazor (Soria, España) [Microfauna (Rodentia, Insectivora, Lagomorpha and Reptilia) from the middle Miocene fissures of Escobosa de Catalañazor (Soria, Spain). – Trabajos sobre Neógeno-Cuaternario 8: 47-73. [In Spanish.]
- Louchart A., Mourer-Chauviré C., Guleç E., Howell F. C. & White T. D. (1998): L'avifaune de Dursunlu, Turquie, Pléistocène inférieur: climat, environnement et biogéographie. – Comptes Rendus de l'Académie des Sciences, Sciences de la Terre et

- des Planètes 327: 341-346.
- Lowe P.R. (1931): Struthious remains from northern China and Mongolia; with descriptions of *Struthio wimani*, *Struthio anderssoni* and *Struthio mongolicus* spp. nov. – *Palaeontologia Sinica* (C) 6 (4): 1-40.
- Lowe P.R. (1939): On the systematic position of the swifts (suborder Cypseli) and humming-birds (suborder Trochili), with special reference to their relation with the order Passeriformes. – *Transactions of the Zoological Society of London* 24: 307-349.
- Ložek V. & Horáček I. (1992): Slovenský kras ve světle kvartérní paleontologie [Slovak karst in the light of the Quaternary paleontology.] – *Slovenský Kras* 30: 29-56. [In Czech.]
- Lucas F.A. (1900): Characters and relations of *Gallinuloides wyomingensis* Eastman, a fossil gallinaceous bird from the Green River shales of Wyoming. – *Bulletin of the Museum of Comparative Zoölogy* 36: 79-84.
- Lucazeau G. (1959): Observations paléontologiques sur les oiseaux fossiles des faluns de Saucats. – Unpub. thesis, Faculté des Sciences, Université de Bordeaux, Bordeaux, 59 pp.
- Lungu A.N. (1966a): O srednosarmatskoj gipparionovoj faune Moldavii [On the middle Sarmatian Hipparion fauna of Moldova.] – *Izvestija Akademii Nauk Moldavskoj SSR* 1966 (10): 20-25. [In Russian.]
- Lungu A.N. (1966b): Uslovija obitanija i osobennosti sistematičeskogo sostava gipparionovoj fauny srednego sarmata Moldavii [Life conditions and peculiarities of the faunal composition of the Hipparion fauna of the middle Sarmatian of Moldova.] – *Izvestija Akademii Nauk Moldavskoj SSR* 1966 (10): 30-38. [In Russian.]
- Lungu A.N. (1968): Gipparionovaja fauna srednego sarmata Moldavii [Hipparion fauna of the middle Sarmatian of Moldova.] – Unpub. Thesis, University of Tbilisi, Tbilisi. [In Russian.]
- Luther D. (1966): Ausgestorbene und aussterbende Vögel. 1. Der Riesenalk. – *Falke* 13: 112-115.
- Luther D. (1986): Die ausgestorbenen Vögel der Welt. 3rd rev. ed. – *Die Neue Brehm-Bücherei* 424: 1-203. Wittenberg Lutherstadt: A. Ziemsen Verlag.
- Lydekker R. (1886): On some Vertebrata from the Red Crag. – *Quarterly Journal of the Geological Society of London* 42: 364-369.
- Lydekker R. (1890): On the remains of some large extinct birds from the cavern-deposits of Malta. – *Proceedings of the Zoological Society of London* 28: 403-411.
- Lydekker R. (1891a): Catalogue of the fossil birds in the British Museum (Natural History). – London: British Museum (Natural History), 368 pp.
- Lydekker R. (1891b): On Pleistocene bird remains from the Sardinian and Corsican islands. – *Proceedings of the Zoological Society of London* 29: 467-476.
- Lydekker R. (1891c): On remains of a large stork from the Allier Miocene. – *Proceedings of the Zoological Society of London* 29: 476-479.
- Lydekker R. (1891d): On British fossil birds. – *Ibis* (6) 3: 381-410.
- Lydekker R. (1891e): [Untitled report.] – *Nature* 45: 70-71. [In English.]
- Lydekker R. (1892): Recent Researches in Fossil Birds. – *Natural Science* 1: 266-271.
- Lydekker R. (1893a): On the extinct giant birds of Argentina. – *Ibis* (6) 5: 40-47.
- Lydekker R. (1893b): On some bird-bones from the Miocene of Grive-St-Alban, Department of Isere, France. – *Proceedings of the Zoological Society of London* 1893: 517-522.
- Lydekker R. (1896): Recent discoveries in avian palaeontology. – *Science Progress* 5: 398-416.
- Lyell C. (1863): The geological evidences of the antiquity of Man, with remarks on the theories of the origin of species by variation. – London: J. Murray, xii + 520 pp.

- Macarovici N. & Oescu C.V. (1941): Quelques vertébrés fossiles trouvés dans les calcaires recifales de Chişinău. – *Analele Academiei Rômîne, Secţie Ştiinţifică* (3) 17: 351-379.
- Mackie S.J. (1863): A classified and stratigraphical list of fossil birds. – *Geologist* 6: 106-109.
- Magish D.P. & Harris A.H. (1976): Fossil ravens from the Pleistocene of Dry Cave, Eddy County, New Mexico. – *Condor* 78: 399-404.
- Mäklin F.W. (1873): Om ett stort i trakten af Cherson anträffadt fossilt fogelägg [On the history and fate of a fossil bird egg found in Cherson.] – *Öfversigt af Finska Vetenskaps-Societetens Förhandlingar* 15: 68-71. [In Swedish.]
- Malez V. (1973): Fosilne ptice Jugoslavije [Fossil birds of Yugoslavia.] – Unpub. Thesis, Zagreb University, 97 pp. [In Serbocroatian.]
- Malez V. (1984a): Ornitofauna starijeg pleistocena Bosne i Hercegovine [Avifauna of the early Pleistocene of Bosna and Herzegovina.] – *Rad Jugoslavenske Akademije Znanosti i Umjetnosti* 411: 265-297. [In Serbocroatian.]
- Malez V. (1984b): Paleornitološki ostaci iz kvartarnih naslaga nekih spilja Hrvatske i Slovenije [Paleornithological remains from the Quaternary deposits of some caverns of Croatia and Slovenia.] – In: *Deveti jugoslavenski speleoloski kongres, Zbornik predavanja [9th Yugoslavian speleological congress, Proceedings]*: 711-719. [In Serbocroatian.]
- Malez V. (1986): Paleornitološka istrazivanja u našim krajevima [Paleornithological research in our country.] – In: *Mestrov M. & Sušić G. (eds.): Ornitologija u Hrvatskoj [Ornithology in Croatia]*: 205-212. Zagreb. [In Serbocroatian.]
- Malez V. (1988): Pleistocenska ornitofauna iz spilje Vindije u sjeverozapadnoj Hrvatskoj [Pleistocene avifauna from the Vindije cavern in northwestern Croatia.] – *RAD Jugoslavenske Akademije Znanosti i Umjetnosti, Varazdin* 2: 31-203. [In Serbocroatian.]
- Malez-Bačić V. (1979): Pleistocenska ornitofauna iz Šandalje u Istri te njezino stratigrafsko i paleokološko značenje [A pleistocene avifauna from Šandalja in Istria and its stratigraphical and paleoecological significance.] – *Palaeontologia Jugoslavica* 21: 1-46. [In Serbocroatian.]
- Mangin J. F. (1962): Traces de pattes d'oiseaux et flûte-casts associe dans un "facies flysch" du Tertiaire Pyrenéen. – *Sedimentology* 1: 163-166.
- Mankinen E.A. & Dalrymple G.B. (1979): Revised geomagnetic polarity time scale for the interval 0-5 m.y. B.P. – *Journal of Geophysical Research (B)* 84: 615-626.
- Marisova I.V. (1963): Holocenovi ptachy Sereďn'oho Prydnistrov'ja [Holocene birds of Central Prydnistrov'e.] – *Dopovidi Zvitno Naukovej Konferencii Kafedr Kremeneć'koho Pedahohičeskoho Intytutu za 1962 r.*: 45-49. [In Ukrainian.]
- Marisova I.V. (1968): Plejstocenovaja ornitofauna Podolii [The Pleistocene avifauna of Podolie.] – *Ornitologija* 9: 316-323. [In Russian.]
- Marisova I.V. & Tatarinov K.A. (1962): Plejstocenovi ptachy Kryvčanskoji peščery [Pleistocene birds of Kryvčanska cave.] – *Naukovy Zapisky Kremeneć'koho Deržavnogo Pedahohičeskoho Instytutu, Serija Prirodničnych Nauk* 7: 63-76. [In Ukrainian.]
- Marisova I.V. & Tatarinov K.A. (1965): Iskopaemaja ornitofauna zapadnych oblastej Ukrainy i tempy mikroevoljucii nekotorych ptic [Fossil avifauna of the western parts of Ukraine, and tempo of microevolution of some birds.] – In: *Novosti ornitologii [Advances in ornithology]*: 232-233. Alma-Ata: Nauka. [In Russian.]
- Marmora A. de la (1831): Sur quelques coquilles marines dans les breches osseuses, et sur quelques autres particularites d'un terrain supérieur au Calcaire tertiaire subappenin de l'île de Sardaigne et autres localites de bords de la Mediterranee. – *Journal de Géologie* 3: 309-320.

- Marschall A.F. (1864): Auszug aus der Denkschrift des Herrn Alphons Milne-Edwards über die geologische Vertheilung der fossilen Vögel. – Neues Jahrbuch für Mineralogie, Geologie und Paläontologie 1864: 412-425. [Report on, and review of Milne-Edwards 1863.]
- Marsh O.C. (1872): Preliminary description of *Hesperornis regalis*, with notices of four other new species of Cretaceous birds. – American Journal of Science and Arts (3) 3: 360-365.
- Marsh O.C. (1894): A gigantic bird from the Eocene of New Jersey. – American Journal of Science and Arts (3) 48: 344.
- Martin D.S. (1874): On *Odontopteryx toliapicus* Owen. – Proceedings of the Lyceum of Natural History in New York (2) 4: 97-98.
- Martin L.D. (1992): The status of the late Paleocene birds *Gastornis* and *Remiornis*. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County, Science Series 36: 97-108.
- Martin L.D. & Mengel R.M. (1984): A new cuckoo and a chachalaca from the early Miocene of Colorado. – Carnegie Museum of Natural History, Special Publication 9: 171-177.
- Martin R. (1903): Note on some remains of *Struthio karatheodoris* Major on the Island of Samos. – Proceedings of the Zoological Society of London 1903: 203-210.
- Martini E. (1967a): Die oligozäne Fossilfundstelle Sieblos an der Wasserkuppe. – Natur und Museum 97: 1-8.
- Martini E. (1967b): *Pararallus hassenkampfi* n.sp., eine neue Rallen-Art (Aves) aus dem Oligozän von Sieblos/Rhön. – Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 127: 288-292.
- Martini E. (1974): Vogelreste aus dem Miozän von Ravalzhausen (Kr. Hanau, Hessen). – Notizblatt des Hessischen Landes-Amtes für Bodenforschung 102: 136-142.
- Martini E. (1988): Vogelreste aus dem Unter-Oligozän von Sieblos/Rhön. – Beiträge zur Naturkunde Ost Hessens 24: 175-177.
- Martini E. (1998): Reste von Vogel-Eischalen im Schneckenmergel von Sieblos/Rhön (Unter-Oligozän). – In: Martini E. & Rothe P. (eds.): Die alttertiäre Fossilagerstätte Sieblos an der Wasserkuppe/Rhön. – Geologische Abhandlungen Hessen 104: 265-270.
- Martini E. & Tobien H. (1984): Wichtige Wirbeltier-Neufunde der Rhön (Sieblos, Theobaldshof). – Beiträge zur Naturkunde Ost Hessens 20: 133-137.
- Massalongo A. & Scarabelli-Gommi-Flamini G. (1859): Studi della flora e geologia stratigrafica del senigalliese [Studies on the fossil flora and stratigraphic geology of Senigallia.] – Imola, vii + 504 pp. [In Italian.]
- Matheu E. & del Hoyo J. (1992): Family Threskiornithidae (ibises and spoonbills). – In: del Hoyo J., Elliott A. & Sargatal J. (eds.): Handbook of the birds of the world. Vol. 1: Ostrich to ducks: 472-506. Barcelona: Lynx Editions.
- Mathews G.M. (1913): A list of the birds of Australia. – London: Witherby, xxvii + 453 pp.
- Matthes H.W. (1970): Zur Paläogeographie und Stammesgeschichte der eozänen Wirbeltiere des Geiseltales. – Hercynia (Neue Folge) 7: 199-249.
- Matthew W.D. & Granger W. (1917): The skeleton of *Diatryma*, a gigantic bird from the Lower Eocene of Wyoming. – Bulletin of the American Museum of Natural History 37: 307-326.
- Mayaud N. & Schaub S. (1950): Les puffins subfossiles de Sardaigne. – Verhandlungen der Naturforschenden Gesellschaft Basel 61: 19-27.
- Mayr E. (1942): Systematics and the origin of species. – New York: Columbia University Press, 334 pp.
- Mayr E. (1963): Animal species and evolution. – Cambridge: Harvard University Press, 797 pp.

- Mayr E. (1982): The growth of biological thought. – Cambridge: Harvard University Press, 974 pp.
- Mayr E. & Amadon D. (1951): A classification of Recent birds. – American Museum Novitates 1496: 1-42.
- Mayr E. & Bock W.J. (1994): Provisional classifications v standard avian sequences: heuristic and communication in ornithology. – Ibis 136: 12-18.
- Mayr G. (1997): Ein fruchtefressender Spechtvogel. – In: von Koeningswald W. & Storch G. (eds.): Messel – Ein Pompeji der Paläontologie: 128-129. Sigmaringen: Jan Thorbecke.
- Mayr G. (1998a): Ein *Archaeotrogon* (Aves: Archaeotrogonidae) aus dem Mittel-Eozän der Grube Messel (Hessen, Deutschland)? – Journal für Ornithologie 139: 121-129.
- Mayr G. (1998b): A new family of Eocene zygodactyl birds. – Senckenbergiana Lethaea 78: 199-209.
- Mayr G. (1998c): "Coraciiforme" und "piciforme" Kleinvögel aus dem Mittel-Eozän der Grube Messel (Hessen, Deutschland). – Courier Forschungsinstitut Senckenberg 205: 1-101.
- Mayr G. (1998d): Ein eozäner Papagei aus Messel. – Natur und Museum 128: 26-28.
- Mayr G. (1999a): A new trogon from the middle Oligocene of Céreste, France. – Auk 116: 427-434.
- Mayr G. (1999b): *Pumiliornis tessellatus* n. gen. n. sp., a new enigmatic bird from the Middle Eocene of Grube Messel (Hessen, Germany). – Courier Forschungsinstitut Senckenberg 216: 75-83.
- Mayr G. (1999c): Caprimulgiform birds from the Middle Eocene of Messel (Hessen, Germany). – Journal of Vertebrate Paleontology 19: 521-532.
- Mayr G. (2000a): A new mousebird (Coliiformes: Coliidae) from the Oligocene of Germany. – Journal für Ornithologie 140: 85-92.
- Mayr G. (2000b): A remarkable new 'gruiform' bird from the middle Eocene of Messel (Hessen, Germany). – Paläontologische Zeitschrift 74: 187-194.
- Mayr G. (2000c): A new basal galliform bird from the middle Eocene of Messel (Hessen, Germany). – In: Storch G. & Weddige K. (eds.): Advances in vertebrate paleontology – Senckenbergiana Lethaea 80: 45-57.
- Mayr G. (2000d): A new raptor-like bird from the Lower Eocene of North America and Europe. – In: Storch G. & Weddige K. (eds.): Advances in vertebrate paleontology – Senckenbergiana Lethaea 80: 59-65.
- Mayr G. (2000e): Avian remains from the middle Eocene of the Geiseltal (Sachsen-Anhalt, Germany). – Vertebrata PalAsiatica 38, Supplement: 19-20. [Abstract.]
- Mayr G. (2000f): Tiny hoopoe-like birds from the middle Eocene of Messel (Hessen, Germany). – Auk 117: 964-970.
- Mayr G. (2000g): Die Vögel der Grube Messel – ein Einblick in die Vogelwelt Mitteleuropas vor 49 Millionen Jahren. – Natur und Museum 130: 365-378.
- Mayr G. (2000h): New or previously unrecorded avian taxa from the middle Eocene of Messel (Hessen, Germany). – Mitteilungen aus dem Museum für Naturkunde (Geowissenschaftliche Reihe) 3: 207-219.
- Mayr G. (2000i): Charadriiform birds from the early Oligocene of Céreste (France) and the middle Eocene of Messel (Hessen, Germany). – Géobios 33: 625-636.
- Mayr G. (2001a): The earliest fossil record of a modern-type piciform bird from the late Oligocene of Germany. – Journal für Ornithologie 142: 2-6.
- Mayr G. (2001b): Comments on the osteology of *Masillapodius longipes* Mayr 1999 and *Paraprefica major* Mayr 1999, caprimulgiform birds from the middle Eocene of Messel (Hessen, Germany). – Neues Jahrbuch für Geologie und Paläontologie, Monatshefte 2001(2): 65-76.
- Mayr G. (2001c): A new specimen of the tiny middle Eocene bird *Gracilitarsus mirabilis*

- (new family: Gracilitarsidae). – *Condor* 103: 78-84.
- Mayr G. (2001d): The relationships of fossil apodiform birds – a comment on Dyke (2001). – *Senckenbergiana Lethaea* 81: 1-2.
- Mayr G. (2001e): New specimens of the middle Eocene fossil mousebird *Selmes absurdipes* Peters 1999. – *Ibis* 143: 427-434.
- Mayr G. & Daniels M. (1998): Eocene parrots from Messel (Hessen, Germany) and the London Clay of Walton-on-the-Naze (Essex, England). – *Senckenbergiana Lethaea* 78: 157-177.
- Mayr G. & Daniels M. (2001): A new short-legged landbird from the early Eocene of Wyoming and contemporaneous European sites. – *Acta Palaeontologica Polonica* 46: 393-402.
- Mayr G. & Gregor H.-J. (1999): Eine fossile Ralle aus dem Plio-Pleistozän von Bobila Ordis bei Banyols (Gerona, NE-Spanien). – *Documenta Naturae* 127: 1-7.
- Mayr G. & Mourer-Chauviré C. (2000): Rollers (Aves: Coraciiformes s.s.) from the Middle Eocene of Messel (Germany) and the Upper Eocene of the Quercy (France). – *Journal of Vertebrate Paleontology* 20: 533-546.
- Mayr, G. & Gregor, H.-J. (1999): Eine fossile Ralle aus dem Plio-Pleistozän von Bobila Ordis bei Banyols (Gerona, NE-Spanien). – *Documenta Naturae* 127: 1-7.
- Mayr G. & Peters D.S. (1998): The mousebirds (Aves: Coliiformes) from the middle Eocene of Grube Messel (Hessen, Germany). – *Senckenbergiana Lethaea* 78: 179-197.
- Mayr G. & Peters D.S. (1999): On the systematic position of the middle Eocene swift *Aegialornis szarskii* Peters 1985 with description of a new swift-like bird from Messel (Aves, Apodiformes). – *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1999: 312-320.
- McDonald J.F. (1994): Identifying Great Auks and other birds in the Palaeolithic art of western Europe: a reply to d'Errico. – *Antiquity* 68: 850-855.
- McMahon T.A. (1973): Size and shape in biology. – *Science* 173: 349-351.
- McMahon T.A. (1975): Allometry and biomechanics: limb bones in adult ungulates. – *American Naturalist* 109: 547-563.
- McMinn M., Jaume D. & Alcover J.A. (1990): *Puffinus olsoni* n. sp.: nova espècie de baldritja recentment extingida provinent de deposits espeleològics de Fuerteventura i Lanzarote (Illes Canaries, Atlantic Oriental). – *Endins* 16: 63-71. [In Catalan.]
- Mecquenem R. (1908): Contribution à l'étude du gisement des vertébrés de Maragha et ses environs. – *Annales d'Histoire Naturelle, Paléontologie* 1: 27-29.
- Mecquenem R. (1925): Contribution à l'étude des fossiles de Maragha [continuation]. – *Annales de Paléontologie* 24: 26-36. [This article has also a parallel pagination for the whole paper. This part thus bears also page numbers 54-64.]
- Mein P. (1976): Résultats du groupe de travail des vertébrés: biozonation du Neogène méditerranéen à partir des mammifères. – In: Seneš J. (ed.): *Proceedings of the VIth congress of the R.C.M.N.S.* 2: 78-81. Bratislava: IGCP.
- Mein P. (1990): Updating of MN zones. – In: Lindsay E.H., Fahlbusch V. & Mein P. (eds.): *European Neogene mammal chronology*: 73-90. New York: Plenum Press.
- Mein P. & Dauphin Y. (1995): Des coquilles d'oeufs de type *Aepyornis* dans le Bassin de Teruel (Pliocène basal, Espagne). – *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 3: 182-191.
- Meldgaard M. (1988): The Great Auk, *Pinguinus impennis* (L.) in Greenland. – *Hist. Biol.* 1: 145-178.
- Mellars P. (1978): Excavation and economic analysis of Mesolithic shell middens on the Island of Oronsay (Inner Hebrides). – In: Mellars P. (ed.): *The early postglacial settlement of northern Europe*: 371-394. London: Duckworth.
- Mellars P. (1987): Oronsay: prehistoric human ecology on a small island. – Edinburgh: Edinburgh University Press.

- Menzel P. (1906): Über die Flora der Senftenberger Braunkohlenablagerungen. – Abhandlungen der Königlich Preussischen Geologischen Landesanstalt (Neue Folge) 46: 120-129.
- Mercerat A. (1893): Note sur la géologie de la Patagonie. – Buenos Aires: Imprenta Helvetia, 6 pp.
- Mercerat A. (1897): Note sur les oiseaux fossiles de la République Argentine. – Anales de la Sociedad de Ciencias de Argentina 43: 222-240.
- Merrem B. (1816) Tentamen systematis naturalis avium et osteologia principiis [An attempt of the natural system of birds, and fundamentals of osteology.] – Königliche Preussische Akademie der Wissenschaften, Abhandlungen 1812-1813: 237-259.
- Meunier S. (1882): Le *Gastornis*. – Nature (Paris) 466: 353-354.
- Meunier S. (1898): Nos terrains. – Paris: Armand Colin, 192 pp.
- Meunier S. (1906): Remarquables traces de pas sur un banc de Gypse. – Naturaliste 28 (453): 19-21.
- Meyer A.B. (1887): Notiz über im Ostsee-Bernstein eingeschlossene Vogelfedern. – Schriften der Naturforschenden Gesellschaft zu Danzig (Neue Folge) 6: 206-208.
- Meyer H. von (1838): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1838: 413-418. [In German.]
- Meyer H. von (1839a): Die fossilen Säugethiere, Reptilien und Vögel aus den Molasse-Gebilden der Schweiz. – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1839: 1-9.
- Meyer H. von (1839b): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1839: 76-79. [In German.]
- Meyer H. von (1839c): Ein Vogel im Kreideschiefer des Kantons Glaris. – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1839: 683-685.
- Meyer H. von (1843): Summarische Uebersicht der fossilen Wirbelthiere des Mainzer Tertiär-Beckens, mit besonderer Rücksicht auf Weisenau. – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1843: 379-410.
- Meyer H. von (1844): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1844: 329-340. [In German.]
- Meyer H. von (1845): Fossile Säugethiere, Voegel und Reptilien aus dem Molasse-Mergel von Oeningen. – Frankfurt am Main: Verlag der S. Schmerberschen Buchhandlung, 54 pp.
- Meyer H. von (1846): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1846: 462-476. [In German.]
- Meyer H. von (1850): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1850: 195-204. [In German.]
- Meyer H. von (1852): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1852: 301-306. [In German.]
- Meyer H. von (1856): Schildkröte und Vogel aus dem Fischschiefer von Glarus. – Palaeontographica 4: 84-95.
- Meyer H. von (1858): Palaeoniscus obtusus, ein Isopode aus der Braunkohle von Sieblos. – Palaeontographica 5: 111-114. [Includes information on Pararallus hassenkampfi.]
- Meyer H. von (1859): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1859: 723-725. [In German.]
- Meyer H. von (1861): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geologie und Paläontologie 1861: 68. [In German.]
- Meyer H. von (1864): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geologie und Paläontologie 1864: 698-701. [In German.]
- Meyer H. von (1865): Fossile Vögel von Radoboy und Oehningen. – Palaeontographica 14: 125-131.

- Meyer H. von (1867): Ueber fossile Eier und Federn. – *Palaeontographica* 15: 223-252.
- Mezzena F. & Palma di Cesnola A. (1992): Nuove manifestazioni d'arte epigravettiana della Grotta Paglicci nel Gargano [New manifestations of the epi-Gravettian art in the Paglicci Cave at Gargano.] – In: *L'arte in Italia dal Paleolitico all'Età del Bronzo* [The art in Italy from the Paleolithic till the Bronze age.] – Atti della XXVIII Riunione Scientifica: 277-292. Firenze. [In Italian.]
- Michajlov K.E. (1988): Sopostavlenie skorlupy jaic vostočnoevropejskich i aziatskich pliocenovych strausov [A comparison of the eggshell of East European and Asian Pliocene ostriches.] – In: Kuročkin E.N. (ed.): *Iskopaemye reptilii i pticy Mongolii* [Fossil reptiles and birds of Mongolia.] – *Sovmestnaja Sovetsko-Mongol'skaja Paleontologičeskaja Ėkspedicija, Trudy* 34: 65-72. [In Russian.]
- Michajlov K.E. (1991): Classification of fossil eggshells of amniotic vertebrates. – *Acta Palaeontologica Polonica* 36: 193-238.
- Michajlov K.E. & Kuročkin E.N. (1988): Skorlupa jaic iskopaemych Struthioniformes iz Palearktiki i ee mesto v sisteme predstavlenij ob èvoljucii Ratitae [Eggshell of fossil Struthioniformes from the Palearctic and its place in the system of opinions on the evolution of the Ratitae.] – In: Kuročkin E.N. (ed.): *Iskopaemye reptilii i pticy Mongolii* [Fossil reptiles and birds of Mongolia.] – *Sovmestnaja Sovetsko-Mongol'skaja Paleontologičeskaja Ėkspedicija, Trudy* 34: 43-65. [In Russian.]
- Mieg M., Bleicher G. & Fliche P. (1892): Contribution à l'étude des terrains tertiaires d'Alsace. Note complémentaire sur le gisement de Roppentzwiller et le gisement à insectes et à plantes de Kleinkems. – *Bulletin de la Société Géologique de France* (3) 20: 375-385.
- Miller A.H. (1944): An avifauna from the Lower Miocene of South Dakota. – *University of California Publications, Bulletin of the Department of Geological Sciences* 27: 85-100.
- Miller A.H. (1963): The fossil flamingos of Australia. – *Condor* 65: 289-299.
- Miller A.H. (1966): The fossil pelicans of Australia. – *Memoirs of the Queensland Museum* 14: 181-190.
- Miller A.H. & Sibley C.G. (1941): A Miocene gull from Nebraska. – *Auk* 58: 563-566.
- Miller L.H. (1950): A Miocene flamingo from California. – *Condor* 52: 69-73.
- Miller L.H. (1952): The avifauna of the Barstow Miocene of California. – *Condor* 54: 296-301.
- Miller L.H. (1956): A collection of bird remains from the Pliocene of San Diego, California. – *Proceedings of the California Academy of Sciences* 28: 615-621.
- Miller L.H. (1962): A new albatross from the Miocene of California. – *Condor* 64: 471-472.
- Milne-Edwards A. (1863): Mémoire sur la distribution géologique des oiseaux fossiles et description de quelques espèces nouvelles. – *Annales des Sciences Naturelles* (4) 20: 132-176. [Abbreviated versions in *Bulletin de la Société Philomatique* 1863: 61-64, *Comptes Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris)* 56: 1219-1222, and *Revue des Sociétés Savantes* (1) 4: 1-5.]
- Milne-Edwards A. (1867a): Études sur les rapports zoologiques du *Gastornis parisiensis*. – *Annales des Sciences Naturelles* (5) 7: 217-227.
- Milne-Edwards A. (1867-1868): *Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France*. Vol. 1. Paris: Victor Masson et Fils, 472 pp. + 96 pls. [1867b = livraisons 1-17; 1868 = livraisons 18-25; dating after Ronsil 1948: 346.]
- Milne-Edwards A. (1869-1871): *Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France*. Vol. 2. Paris: G. Masson, 627 pp. + pls. 97-200. [1869 = livraisons 26-30; 1871 = livraisons 31 ff.; dating after Ronsil 1948: 346.]

- Milne-Edwards A. (1870): Observations sur la faune ornithologique du Bourbonnais pendant la période tertiaire moyenne. – Annales des Sciences Géologiques (Paris) 2 (2): 1-6. [Same paper in Bibliothèque de l'École des Hautes Études (Paris) 3 (3): 1-6., Comptes Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris) 70: 557-559, and with minor changes in Bulletin Hebdomadaire de l'Association Scientifique de France 7: 280-282.]
- Milne-Edwards A. (1872): Résumé des recherches sur les oiseaux fossiles. – Annales des Sciences Naturelles (Paris) (5) 16 (2): 1-7. [Same paper with minor changes in Bibliothèque de l'École des Hautes Études (Paris) 6: 7 pp., Bulletin Hebdomadaire de l'Association Scientifique de France 10: 99-103, Compte Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris) 74: 1030-1034.]
- Milne-Edwards A. (1874): Observations sur les oiseaux fossiles des Faluns de Saucats et de la Molasse de Léognan. – Bibliothèque de l'École des Hautes Études, Section des Sciences Naturelles 11(3): 3-12. [Reprinted in Annales des Sciences Géologiques 6: 12 pp., 1875]
- Milne-Edwards A. (1875a): Observations on the birds whose bones have been found in the caves of the south-west of France. – Lartet E. & Christy H. (eds.): Reliquiae Aquitanicae: 226-247. London.
- Milne-Edwards A. (1875b): Observations sur les oiseaux dont les ossements ont été retrouvés dans les cavernes du Sud Ouest de la France. – Matériaux pour l'Histoire Primitive et Naturelle de l'Homme (2) 6: 473-503. [French version of Milne-Edwards 1875a.]
- Milne-Edwards A. (1892): Sur les oiseaux fossiles des dépôts éocènes de phosphate de chaux du Sud de la France. – In: Sclater P.L. (ed.): Comptes Rendus du Second Congrès Ornithologique International: 60-80. Budapest.
- Milne-Edwards A. (1893): [Untitled letter.] – Bulletin of the British Ornithologists' Club 1: liii-liv. [In English; see also Ibis (6) 5: 568 (1893).]
- Misonne X. (1958): Faune du Tertiaire et du Pléistocène inférieur de Belgique (oiseaux et mammifères). – Bulletin de l'Institut Royal des Sciences Naturelles de Belgique 46(34): 1-10.
- Mlíkovský J. (1977): Beiträge zur Evolution des Vogelgehirns. – Unpublished thesis, Martin-Luther-Universität, Halle/Saale, 278 pp.
- Mlíkovský J. (1980): Zwei Vogelgehirne aus dem Miozän Böhmens. – Časopis pro Mineralogii a Geologii 25: 409-413.
- Mlíkovský J. (1981a): Relationships of the Eocene bird "*Numenius*" *gyporum* Gervais. – Bulletin du Muséum National d'Histoire Naturelle de Paris (4) (C) 3: 341-343.
- Mlíkovský J. (1981b): Ein fossile [sic!] Vogelgehirn aus dem Oberpliozän Ungarns. – Fragmenta Mineralogica et Palaeontologica 10: 409-413.
- Mlíkovský J. (1982a): Taxonomische Identität der *Anas submajor* Jánossy, 1979 (Aves: Anseriformes) aus dem Oberpliozän Ungarns. – Věstník Československé Společnosti Zoologické 46: 199-202.
- Mlíkovský J. (1982b): Towards a new classification of birds. – In: Il'ičev V.D. & Gavrilov V.M. (eds.): XVIII Congressus internationalis ornithologicus, Abstracts of symposia and poster presentations: 248-251. Moskva: Nauka.
- Mlíkovský J. (1982c): Zur systematischen Stellung von *Bucephala angustipes* Jánossy, 1965 und *Somateria gravipes* Harrison, 1979 (Aves: Anseriformes) aus dem Pleistozän Europas. – Zeitschrift für Geologische Wissenschaften 10: 1463-1475.
- Mlíkovský J. (1982d): Zur Methodologie phylogenetischer Rekonstruktionen in der Paläontologie. – Wissenschaftliche Zeitschrift der Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Reihe 31: 143-146.
- Mlíkovský J. (1983a): On the foundations of biological systematics: a historical approach. – In: Novák V.J.A. & Zemek K. (eds.): General questions of evolution: 305-317. Praha:

ČSAV.

- Mlíkovský J. (1983b): Methodological problems of the contemporary systematics of the family Hominidae. – *Anthropologie* 21: 33-36.
- Mlíkovský J. (1983c): Fossilní doklady evoluce vrubozobých (Aves: Anseriformes) [Fossil evidence for the evolution of the waterfowl (Aves: Anseriformes).] – Unpublished thesis, Czechoslovak Academy of Sciences, Praha, 320 pp. [In Czech.]
- Mlíkovský J. (1985): Towards a new classification of birds. – In: Il'ičev V.D. & Gavrilov V.M. (eds.): *Acta XVIII Congressus internationalis ornithologici*: 1145-1146. Moskva: Nauka. [Reprint of Mlíkovský 1982b.]
- Mlíkovský J. (1987a): Eine neue Alkenart (Aves: Alcidae) aus dem Ober-Oligozän Österreichs. – *Annalen des Naturhistorischen Museums in Wien (A)* 88: 131-147. [With a contribution by J. Kovar.]
- Mlíkovský J. (1987b): Ecological bases of supraspecific evolution. – In: Pesce-Delfino V. (ed.): *Biological evolution*: 253-267. Bari: Adriatica Editrice.
- Mlíkovský J. (1987c): Fundamental units and reductionism in evolutionary biology. – In: Rabinovič V.L. (ed.): *8 International congress of logic, methodology and philosophy of science, Abstracts 2*: 277-279. Moskva: AN SSSR.
- Mlíkovský J. (1988): Notes on the brains of the middle Miocene birds (Aves) of Hahnenberg (F.R.G.). – *Časopis pro Mineralogii a Geologii* 33: 53-61.
- Mlíkovský J. (1989a): A new swift (Aves: Apodidae) from the late Eocene of France. – *Annalen des Naturhistorischen Museums in Wien (A)* 90: 59-62.
- Mlíkovský J. (1989b): A new guineafowl (Aves: Phasianidae) from the late Eocene of France. – *Annalen des Naturhistorischen Museums in Wien (A)* 90: 63-66.
- Mlíkovský J. (1989f): Ptáci staršího a středního pleistocénu Československa: současný stav a perspektivy výzkumu [Birds of the early and middle Pleistocene of Czechoslovakia: current state and perspectives of the research]. – In: Seitl L. (ed.): *Současný stav a perspektivy výzkumu kvartéru v ČSSR* [Current state and perspectives of the Quaternary research in the ČSSR]: 63-67. Brno: Moravské muzeum & Universita J.E. Purkyně. [In Czech.]
- Mlíkovský J. (1992a): Late Miocene birds of Götzendorf/Leitha, Austria. – *Annalen des Naturhistorischen Museums in Wien (A)* 91: 97-100.
- Mlíkovský J. (1992b): The present state of knowledge of the Tertiary birds of Central Europe. – In: Campbell K.E. (ed.): *Studies in avian paleornithology honoring Pierce Brodkorb*. – *Natural History Museum of Los Angeles County, Science Series* 36: 433-458.
- Mlíkovský J. (1995a): Nomenclatural and taxonomic status of fossil birds described by H.G.L. Reichenbach in 1852. – In: Peters D.S. (ed.): *Acta palaeornithologica*. – *Courier Forschungsinstitut Senckenberg* 181: 311-316.
- Mlíkovský J. (1995b): Early Pleistocene birds of Stránská skála: 1. Musil's talus cone. – In: Musil R. (ed.): *Stránská skála Hill: Excavations of open-air sediments 1964-1972*. – *Anthropos (Brno)* 26: 111-126.
- Mlíkovský J. (ed., 1996a): Tertiary avian localities of Europe. – *Acta Universitatis Carolinae, Geologica* 39: 517-848. [For reviews see Mourer-Chauviré 1997 and Olson 1997.]
- Mlíkovský J. (1996b). Tertiary avian localities of Europe: An introduction. In: Mlíkovský J. (ed.): *Tertiary avian localities of Europe*. *Acta Universitatis Carolinae, Geologica* 39: 519-528.
- Mlíkovský J. (1996c): Tertiary avian localities of Austria. – In: Mlíkovský J. (ed.): *Tertiary avian localities of Europe*. – *Acta Universitatis Carolinae, Geologica* 39: 529-533.
- Mlíkovský J. (1996d): Tertiary avian localities of Croatia. – In: Mlíkovský J. (ed.): *Tertiary avian localities of Europe*. – *Acta Universitatis Carolinae, Geologica* 39: 547-

- 549.
- Mlíkovský J. (1996e): Tertiary avian localities of the Czech Republic. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 551-557.
- Mlíkovský J. (1996f): Tertiary avian localities of Denmark. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 559-562.
- Mlíkovský J. (1996g): Tertiary avian localities of Finland. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 563-565.
- Mlíkovský J. (1996h): Tertiary avian localities of Greece. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. Acta Universitatis Carolinae, Geologica 39: 649-653.
- Mlíkovský J. (1996i): Tertiary avian localities of Holland. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 655-656.
- Mlíkovský J. (1996j): Tertiary avian localities of Hungary. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 657-664.
- Mlíkovský J. (1996k): Tertiary avian localities of Macedonia. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 683-684.
- Mlíkovský J. (1996l): Tertiary avian localities of Moldavia. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 685-691.
- Mlíkovský J. (1996m): Tertiary avian localities of Russia (European part). – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 711-713.
- Mlíkovský J. (1996n): Tertiary avian localities of Slovakia. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 715-718.
- Mlíkovský J. (1996o): Tertiary avian localities of Switzerland. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 735-742.
- Mlíkovský J. (1996p): Tertiary avian localities of Ukraine. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 743-757.
- Mlíkovský J. (1996q): Tertiary avian localities of the United Kingdom. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 759-771.
- Mlíkovský J. (1996r): Tertiary avian localities of Yugoslavia. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 773-775.
- Mlíkovský J. (1996s): Tertiary avian faunas of Europe. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 777-818.
- Mlíkovský J. (1996t): Early and middle Pleistocene birds from the Bohemian Karst, Czech Republic. – Acta Societatis Zoologicae Bohemicae 60: 187-190.
- Mlíkovský J. (1997a): Taxonomic identity of *Fringilla radoboyensis* von Meyer 1865 (Aves) from the middle Miocene of Croatia. – Annalen des Naturhistorischen Museums in Wien (A) 98: 143-149.
- Mlíkovský J. (1997b): A new tropicbird (Aves: Phaethontidae) from the late Miocene of Vösendorf, Austria. – Annalen des Naturhistorischen Museums in Wien (A) 98: 151-

154.

- Mlíkovský J. (1997c): Taxonomic identity of *Haliaeetus angustipes* Jánossy, 1983 (Aves: Accipitridae) from the early Pleistocene of the Czech Republic. – *Buteo* 9: 51-56.
- Mlíkovský J. (1997d): Jungpleistozäne Vögel aus der Schusterlucke, Niederösterreich. – *Wissenschaftliche Mitteilungen aus dem Niederösterreichischen Museum* 10: 105-114.
- Mlíkovský J. (1998a): Vertebrates from the Early Miocene lignite deposits of the opencast mine Oberdorf (Western Styrian Basin, Austria): 4. Aves. – *Annalen des Naturhistorischen Museums in Wien (A)* 99: 39-42.
- Mlíkovský J. (1998b): A new loon (Aves: Gaviidae) from the middle Miocene of Austria. – *Annalen des Naturhistorischen Museums in Wien (A)* 99: 331-339.
- Mlíkovský J. (1998c): A new barn owl (Aves: Strigidae) from the early Miocene of Germany, with comments on the fossil history of the Tytoninae. – *Journal für Ornithologie* 139: 247-261.
- Mlíkovský J. (1998d): Early Pleistocene birds of Deutsch-Altenburg, Austria. – *Acta Societatis Zoologicae Bohemiae* 62: 135-141.
- Mlíkovský J. (1998e): A new parrot (Aves: Psittacidae) from the early Miocene of the Czech Republic. – *Acta Societatis Zoologicae Bohemiae* 62: 335-341.
- Mlíkovský J. (1998f): Two new owls (Aves: Strigidae) from the early Miocene of the Czech Republic, with comments on the fossil history of the subfamily Striginae. – *Buteo* 10: 5-21.
- Mlíkovský J. (1998g): Taxonomic comments on the Quaternary vultures (Aves: Accipitridae, Aegypiinae) of Central Europe. – *Buteo* 10: 23-29.
- Mlíkovský J. (1999a): A new jacana (Aves: Jacanidae) from the early Miocene of the Czech Republic. – *Comptes Rendus de l'Académie des Sciences (Paris), Sciences de la Terre et des Planètes* 328: 121-123.
- Mlíkovský J. (1999b): A new painted snipe (Aves: Rostratulidae) from the early Miocene of the Czech Republic. – *Časopis Národního Muzea (Praha), Řada Přírodovědná* 167: 99-101.
- Mlíkovský J. (1999c): Early Miocene birds of Břešňany, Czech Republic. – *Časopis Národního Muzea (Praha), Řada Přírodovědná* 167: 103-109.
- Mlíkovský J. (1999d): Tertiary barn owls of Europe. – *Journal für Ornithologie* 140: 364-366. [Response to Mourer-Chauviré 1999c.]
- Mlíkovský J. (1999e): Note on the osteology and taxonomic position of African Long-tailed Hawk *Urotriorchis macrourus* (Aves: Accipitridae). – *Bulletin of the British Ornithologists' Club* 119: 32-37.
- Mlíkovský J. (2000a): Taxonomic identity of *Gyps fulvus spelaeus* Friant, 1950 (Aves: Accipitridae) from the Pleistocene of Belgium. – *Buteo* 11: 101-102.
- Mlíkovský J. (2000b): First record of *Tyto balearica* (Aves: Strigidae) from the Pleistocene of Sardinia. – *Buteo* 11: 103-105.
- Mlíkovský J. (2000c): Early Miocene pratincoles (Aves: Glareolidae) from Dolnice, Czech Republic. – *Acta Societatis Zoologicae Bohemiae* 64: 93-96.
- Mlíkovský J. (2000d): Family-group names of Cenozoic birds: 1811-1998. – *Časopis Národního Muzea (Praha), Řada Přírodovědná* 169: 75-90.
- Mlíkovský J. (2000e): Early Miocene quails (Aves: Phasianidae) from Saint-Gérand-le-Puy, France. – *Časopis Národního Muzea (Praha), Řada Přírodovědná* 169: 91-96.
- Mlíkovský J. (2000f): Early Miocene birds of Skyřice, Czech Republic. – *Časopis Národního Muzea (Praha), Řada Přírodovědná* 169: 97-100.
- Mlíkovský J. (2001): Late Cenozoic biostratigraphy of Europe: mammal zones and the fossil record of birds. – In: Horáček I. & Mlíkovský J. (eds.): *Papers in paleomammalogy honoring Oldřich Fejfar*. – *Lynx (new series)* 32: in press. Praha.
- Mlíkovský J. (2002): Vögel aus der frühmittelalterlichen Burg Mikulčice, Mähren. – In:

- Poláček L. (ed.): Studien zum Burgwall von Mikulčice 5: in press. Brno: Archeologický ústav AV ČR.
- Mlíkovský J. & Göhlich U.B. (2000): A new wood-hoopoe from the early Miocene of Germany and France. – *Acta Societatis Zoologicae Bohemiae* 64: 419-424.
- Mlíkovský J. & Hesse A. (1996): Tertiary avian localities of Germany. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – *Acta Universitatis Carolinae, Geologica* 39: 619-647.
- Mlíkovský J. & Lukáš J. (1991): Osteological disorders in late Pleistocene birds from the Schusterlucke, Lower Austria. – *Annalen des Naturhistorischen Museums in Wien (A)* 92: 101-103.
- Mlíkovský J. & Švec P. (1986): Review of the Tertiary waterfowl (Aves: Anseridae) of Asia. – *Věstník Československé Společnosti Zoologické* 50: 259-272.
- Mlíkovský J. & Švec P. (1989): Review of the Tertiary waterfowl (Aves: Anseridae) of Czechoslovakia. – *Časopis pro Mineralogii a Geologii* 34: 199-203.
- Mlíkovský J. & Zemek K. (1983): On the history and present state of the species concept. – In: Novák V.J.A. & Zemek K. (eds.): General questions of evolution: 281-304. Praha: ČSAV.
- Mlíkovský J., Bělka L. & Zemek K. (1985): Morphogenesis and the problem of morphospecies. – In: Mlíkovský J. & Novák V.J.A. (eds.): Evolution and morphogenesis: 201-211. Praha: ČSAV.
- Montevocchi W.A. & Hufthammer A.K. (1990): Zooarchaeological implications for prehistoric distributions of seabirds along the Norwegian coast. – *Arctic* 43: 110-114.
- Montoya, P. et al. (1999): La fauna del Pleistoceno inferior de la Sierra de Quibas (Abanilla, Murcia). – *Estudios Geológicos* 55: 127-161.
- Montoya, P. et al. (2001): Une faune très diversifiée du Pléistocène inférieur de la Sierra de Quibas (province de Murcia, Espagne). – *Comptes Rendus de l'Académie des Sciences (Paris), Sciences de la Terre et des Planètes* 332: 387-393.
- Morales J. & Aguirre E. (1976): Carnívoros de Venta del Moro [Carnivores from Venta del Miceno.] – *Trabajos sobre Neógeno-Cuaternario* 5: 31-81. [In Spanish.]
- Moreau R.E. (1954): The main vicissitudes of the European avifauna since the Pliocene. – *Ibis* 96: 411-431.
- Moreau R.E. (1966): On estimates of the past numbers and of the average longevity of avian species. – *Auk* 83: 403-415.
- Moreno E. (1986): Clave osteológica para la identificación de los Passeriformes ibéricos [Osteological key to the identification of the Iberian Passeriformes]. II. Hirundinidae, Prunellidae, Sittidae, Certhiidae, Troglodytidae, Cinclidae, Laniidae, Oriolidae, Corvidae, Sturnidae, Motacillidae. – *Ardeola* 33: 69-129. [In Spanish.]
- Moreno F.P. & Mercerat A. (1891): Catálogo de los pájaros fósiles de la República Argentina conservados en el Museo de La Plata [Catalogue of fossil birds of the Republic of Argentina deposited in the Museum of La Plata.] – *Anales del Museo La Plata* 1: 1-71. [In Spanish.]
- Morris J. (1843): Catalogue of British fossils. London.
- Mourer-Chauviré C. (1975a): Les oiseaux du Pléistocène moyen et supérieur de France. – *Documents des Laboratoires de Géologie de la Faculté des Sciences de Lyon* 64: 1-624.
- Mourer-Chauviré C. (1975b): Les oiseaux (Aves) du gisement pléistocène moyen des Abimes de la Fage à Noailles (Corrèze). – *Nouvelles Archives de la Musée d'Histoire Naturelle de Lyon* 13 (Supplement): 89-112.
- Mourer-Chauviré C. (1975c): Un exemple d'évolution chez les oiseaux au quaternaire: *Buteo rufinus jansoni* nov. subsp. (Aves, Falconida, Accipitridae) du Pléistocène moyen de Saint-Estève-Janson (Bouches-du-Rhône, France). – *Colloque International C.N.R.S.* 218: 533-542.
- Mourer-Chauviré C. (1975d): Faunes d'oiseaux du Pléistocène de France: systématique,

- évolution et adaptation, interprétation paléoclimatique. – *Géobios* 8: 333-352.
- Mourer-Chauviré C. (1976): Les oiseaux. – In: Lumley H. de (ed.): Les civilisations paléolithiques et mésolithiques de la France. Vol. 1: 430-434. – Paris: Éditions du Centre National de la Recherche Scientifique.
- Mourer-Chauviré C. (1977): Die Vogelreste aus den mittelpleistozänen Mosbacher Sanden bei Wiesbaden (Hessen). – *Mainzer Naturwissenschaftliches Archiv* 16: 35-46.
- Mourer-Chauviré C. (1978a): Le bassin lacustre Miocène de Bes-Konak (Anatolie – Turquie): géologie et introduction à la paléontologie des vertébrés. II. Les vertébrés. 5. Oiseaux.. – *Géobios* 11: 52-54.
- Mourer-Chauviré C. (1978b): La poche à phosphate de Sainte-Néboule (Lot) et sa faune de Vertébrés du Ludien supérieur. 6. Oiseaux. – *Palaeovertebrata* 8: 217-229.
- Mourer-Chauviré C. (1980a): The Archaeotrogonidae from the Eocene and Oligocene deposits of "Phosphorites du Quercy", France. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Hildegard Howard. – Natural History Museum of Los Angeles County, Contributions in Science 330: 17-31.
- Mourer-Chauviré C. (1980b): La faune d'oiseaux du Pléistocène en Europe occidentale. – In: Problèmes de stratigraphie quaternaire de France et dans les pays limitrophes. – Bulletin de la Association Française Etude Quaternaire (n.s.), Supplement 1: 353-358.
- Mourer-Chauviré C. (1980c): Las Aves del sitio de ocupación achelense de Aridos-1 (Arganda, Madrid). – In: Santonja M., López N. & Pérez A. (eds.): Ocupaciones achelenses en el valle del Jarama: 145-160. Madrid: Excelentissima Diputación Provincial.
- Mourer-Chauviré C. (1981a): Première indication de la présence de Phorurhacids, famille d'oiseaux géants d'Amérique du Sud, dans le Tertiaire européen: *Ameghinornis* nov. gen. (Aves, Ralliformes) des Phosphorites du Quercy, France. – *Géobios* 14: 637-647.
- Mourer-Chauviré C. (1981b): Les oiseaux de la grotte de Kitsos (Attique, Grèce). – In: Lambert N. (ed.): La grotte préhistorique de Kitsos (Attique). Vol. 2: 595-606. Paris: Editions A.D.P.F, and Athens: Ecole Française.
- Mourer-Chauviré C. (1982a): Les oiseaux fossiles des Phosphorites du Quercy (Eocène supérieur à Oligocène supérieur). Implications paléobiogéographiques. – In: Buffetaut E., Janvier P., Rage J.C. & Tassy P. (eds.): Phylogénie et paleobiogéographie. Livre jubilaire en l'honneur de Robert Hoffstetter. – *Géobios, Mémoire Special* 6: 413-426.
- Mourer-Chauviré C. (1982b): Réponse [to Buffetaut & Rage 1982.] – *Géobios* 15: 268-269.
- Mourer-Chauviré C. (1983): Les Gruiformes (Aves) des Phosphorites du Quercy (France). 1. Sous-ordre Cariamae (Cariamidae et Phorusrhacidae). Systématique et Biostratigraphie. – *Paleovertebrata* 13: 83-143.
- Mourer-Chauviré C. (1985a): Nouvelles découvertes d'oiseaux éocènes et oligocènes dans les Phosphorites du Quercy (France): les Idiornithinae (Ralliformes, Cariamidae). – In: Il'čev V.D. & Gavrilov V.M. (eds.), Acta XVIII congressus internationalis ornithologici: 234-240. Moskva: Nauka.
- Mourer-Chauviré C. (1985b): Les Todidae (Aves, Coraciiformes) des Phosphorites du Quercy (France). – Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen (Amsterdam), B 88: 407-411.
- Mourer-Chauviré C. (1987): Les Strigiformes (Aves) des Phosphorites du Quercy (France): Systématique, biostratigraphie et paléobiogéographie. – In: Mourer-Chauviré C. (ed.): L'évolution des Oiseaux d'après le témoignage des fossiles. – Documents des Laboratoires de Géologie de Lyon 99: 89-135.
- Mourer-Chauviré C. (1988a): Le gisement du Bretou (Phosphorites du Quercy, Tarn-et-Garonne, France) et sa faune de Vertébrés de l'Eocène supérieur. II. Oiseaux. –

- Palaeontographica, A 205: 29-50.
- Mourer-Chauviré C. (1988b): Les Aegialornithidae (Aves: Apodiformes) des Phosphorites du Quercy. Comparaison avec la forme de Messel. – Courier Forschungsinstitut Senckenberg 107: 369-381.
- Mourer-Chauviré C. (1989a): Les Caprimulgiformes et les Coraciiformes de l'Eocène et de l'Oligocène des Phosphorites du Quercy et description de deux genres nouveaux de Podargidae et Nyctibiidae. – Ouellet H. (ed.): Acta XIX Congressus Internationalis Ornithologici: 2047-2055. Ottawa: University of Ottawa Press.
- Mourer-Chauviré C. (1989b): A Peafowl from the Pliocene of Perpignan, France. – Palaeontology 32: 439-446.
- Mourer-Chauviré C. (1990): Les paons du Pliocène et du Pléistocène inférieur d'Europe. Biostratigraphie et paléoécologie. – In: Journée Louis David. – Documents des Laboratoires de Géologie de la Faculté des Sciences Lyon, Hors Série 9: 77-91.
- Mourer-Chauviré C. (1991): Les Horusornithidae nov. fam. (Aves, Accipitriformes) à articulation intertarsienne hyperflexible de l'Eocène du Quercy. – Géobios, Mémoire Spécial 13: 183-192.
- Mourer-Chauviré C. (1992a): Un gânga primitif (Aves, Columbiformes, Pteroclididae) de très grande taille dans le Paléogène des Phosphorites du Quercy (France). – Comptes Rendus Hebdomadaires de l'Académie des Sciences de Paris, Sciences de la Terre et des Planètes 314: 229-235.
- Mourer-Chauviré C. (1992b): Les Galliformes (Aves) of Phosphorites du Quercy (France). Systematics and Biostratigraphy. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County. Science Series 36: 37-95.
- Mourer-Chauviré C. (1992c): Une nouvelle famille de Perroquets (Aves, Psittaciformes) dans l'Eocène supérieur des Phosphorites du Quercy. – Géobios, Mémoire Spécial 14: 169-177.
- Mourer-Chauviré C. (1993a): Les gangas (Aves, Columbiformes, Pteroclididae) du Paleogène et du Miocène inférieur de France. – Palaeovertebrata 22: 73-98.
- Mourer-Chauviré C. (1993b): The Pleistocene avifaunas of Europe. – In: Morales Muniz A. (ed.): Archaeornithology: birds and the archaeological record. – Archaeofauna 2: 53-66.
- Mourer-Chauviré C. (1994): A large owl from the Paleocene of France. – Paleontology 37: 339-348.
- Mourer-Chauviré C. (1995a): The Messelornithidae (Aves: Gruiformes) from the Paleogene of France. – In: Peters D.S. (ed.): Acta palaeornithologica. – Courier Forschungsinstitut Senckenberg 181: 95-105.
- Mourer-Chauviré C. (1995b): Dynamics of the avifauna during the Paleogene and the early Neogene of France. Settling of the recent fauna. – Acta Zoologica Cracoviensia 38: 325-342.
- Mourer-Chauviré C. (1995c): Oiseaux. – In: Le Garouillas et les sites contemporains (Oligocène, MP 25) des Phosphorites du Quercy (Lot, Tarn-et-Garonne, France) et leurs faunes de Vertébrés. – Palaeontographica, A 236: 33-38.
- Mourer-Chauviré C. (1996a): Paleogene avian localities of France. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 567-598.
- Mourer-Chauviré C. (1996b): Pliocene avian localities of France. – In: Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 613-618.
- Mourer-Chauviré C. (1997): [Review of Mlíkovský 1996a.] – Ibis 139: 715. [In English.]
- Mourer-Chauviré C. (1999a): Position systématique de *Nupharanassa bohemica* Mlíkovský, 1999. – Comptes Rendus de l'Académie des Sciences, Sciences de la Terre

- et des Planètes 329: 149-152.
- Mourer-Chauviré C. (1999b): Les relations entre les avifaunes du Tertiaire inférieur d'Europe et d'Amérique du Sud. – Bulletin de la Société Géologique de France 170: 85-90.
- Mourer-Chauviré C. (1999c): Tertiary barn owls of Europe. – Journal für Ornithologie 140: 363-364. [Comments on Mlíkovský 1998c.]
- Mourer-Chauviré C. (1999d): Influence de l'homme sur la répartition de certains oiseaux marins: l'exemple du Grand Pingouin, *Pinguinus impennis*. – Alauda 67: 273-279.
- Mourer-Chauviré C. (1999e): [Untitled response to Rage 1999.] – Bulletin de la Société Géologique de France 170: 951-954. [In French.]
- Mourer-Chauviré C. (2000a): A new species of Ameripodius (Aves: Galliformes: Quercy-megapodiidae) from the Lower Miocene of France. – Palaeontology 43: 481-493.
- Mourer-Chauviré C. (2000b): Influence de l'homme préhistorique sur la répartition de certains oiseaux marins: l'exemple du Grand Pingouin *Pinguinus impennis*. – Alauda 67: 273-279.
- Mourer-Chauviré C. & Antunes M.T. (1991): Présence du grand pingouin, *Pinguinus impennis* (Aves, Charadriiformes) dans le Pléistocène du Portugal. – Géobios 24: 201-205.
- Mourer-Chauviré C. & Antunes M.T. (2000): L'avifaune pléistocène et holocène de Gruta da Figueira Brava (Arrábida, Portugal). – In: Antunes M.T. (ed.): Últimos neandertais em Portugal. Evidência, odontológica e outra [Last neanderthals in Portugal. Odontologic and other evidence.] – Memórias da Academia das Ciências de Lisboa, Classe de Ciências 38: 129-161. [In Catalan.]
- Mourer-Chauviré C. & Cheneval J. (1983): Les Sagittariidae fossiles (Aves, Accipitriiformes) de l'Oligocène des Phosphorites du Quercy et du Miocène inférieur de Saint-Gérard-le-Puy. – Géobios 16: 443-459.
- Mourer-Chauviré C. & Sánchez Marco A. (1988): Présence de *Tyto balearica* (Aves, Strigiformes) dans des gisements continentaux du Pliocène de France et d'Espagne. – Géobios 21: 639-644.
- Mourer-Chauviré C. & Weesie P.D.M. (1986): *Bubo insularis* n. sp., forme endémique insulaire de grand-duc (Aves, Strigiformes) du Pléistocène de Sardaigne et de Corse. – Revue de Paléobiologie 5: 197-205.
- Mourer-Chauviré C., Adrover R. & Pons J. (1975): Présence de *Grus antigone* (L.) dans l'"Avenc de Na Corna" a Majorque (Espagne). – Nouvelles Archives de la Musée d'Histoire Naturelle de Lyon 13 (Supplement): 45-50.
- Mourer-Chauviré C., Moyà S. & Adrover R. (1977): Les oiseaux des gisements quaternaires de Majorque. – Nouvelles Archives de la Musée d'Histoire naturelle de Lyon 15, Supplement: 61-64.
- Mourer-Chauviré C., Alcover J.A., Moyà S. & Pons J. (1980): Une nouvelle forme insulaire d'effraie géante, *Tyto balearica* n.sp. (Aves, Strigiformes) du Plio-Pléistocène des Baléares. – Géobios 13: 803-811.
- Mourer-Chauviré C., Balouet J.-C., Jehenne Y. & Heintz É. (1985): Une nouvelle espèce de grue, *Grus afghana* (Aves, Gruiformes), du Miocène supérieur de Molayan, Afghanistan. – Bulletin du Musée National d'Histoire Naturelle (Paris) (4) 7: 179-187.
- Mourer-Chauviré C., Hugueney M. & Jonet P. (1989): Découverte de Passeriformes dans l'Oligocène supérieur de France. – Comptes Rendus Hebdomadaires de l'Académie des Sciences (Paris), Sciences de la Terre et des Planètes 309: 843-849.
- Mourer-Chauviré C., Salotti M., Pereira É., Quinif Y., Courtois J.-Y., Dubois J.-N. & La Milza J.-C. (1997): *Athene angelis* n. sp. (Aves, Strigiformes), nouvelle espèce endémique insulaire éteinte du Pléistocène moyen et supérieur de Corse (France). – Comptes Rendus de l'Académie des Sciences (Paris), Sciences de la Terre et des

- Planètes 324: 677-684.
- Mourer-Chauviré C., Legendre S., Marandat B. & Sigé B. (1999): Le Quercy tropical. – *Pour la Science* 262: 60-67.
- Mottl M. (1951): Die Repolust-Höhle bei Peggau (Steiermark) und ihre eiszeitlichen Bewohner. – *Archaeologia Austriaca* 8: 1-78.
- Müller A.H. (1955): Der Großablauf der stammesgeschichtlichen Entwicklung. – Jena: Gustav Fischer, 51 pp.
- Müller A.H. (1961): Großabläufe der stammesgeschichtlichen Entwicklung. Erscheinungen und Probleme. – Jena: Gustav Fischer, 116 pp.
- Müller A.H. (1968): Lehrbuch der Paläozoologie. Band III. Vertebraten, Teil 2. Reptilien und Vögel. – Jena: Gustav Fischer Verlag, xv + 656 pp. [Aves on pp. 550-627.]
- Munthe H. (1914): Om några fågelfynd från sen-glaciala lager i Sverige [On some finds of birds from the late Pleistocene deposits in Sweden.] – *Sveriges Geologiska Undersökning, Årbok* 8 (4): 3-26. [In Swedish.]
- Murie J. (1873): On the Upupidae and their relationships. – *Ibis* 15: 181-211.
- Musil R. (1965): Aus der Geschichte der Stránská skála. – *Acta Musei Moraviae* 50: 75-106.
- Musil R. (1966): Holštejn, eine neue altpleistozäne Lokalität in Mähren. – *Acta Musei Moraviae* 51: 13-168.
- Musil R. (ed., 1995a): Stránská skála hill: excavations of open-air sediments 1964-1972. – *Anthropos (Brno)* 26: i-viii, 1-214.
- Musil R. (1995b): The Stránská skála hill – its importance and significance. – In: Musil R. (ed.): *Stránská skála hill: excavations of open-air sediments 1964-1972*. – *Anthropos (Brno)* 26: 177-208.
- Musil R. & Valoch K. (1968): Stránská skála: its meaning for Pleistocene studies. – *Current Anthropology* 9: 534-539.
- Nathusius W. von (1886): Über das fossile Ei von *Struthiolithus chersonensis* Brandt. – *Zoologischer Anzeiger* 9: 47-50.
- Navás L. (1922): Algunos fósiles de Libros (Teruel) [Some fossils from Libros (Teruel).] – *Boletín de la Sociedad Ibérica de Ciencias Naturales* 21(4): 52-61. [In Spanish.]
- Nehring A. (1877): Die quartären Faunen von Thiede und Westeregeln nebst Spuren des vorgeschichtlichen Menschen. – *Archiv für Anthropologie* 10: 359-398.
- Nehring A. (1880): Uebersicht über vierundzwanzig mitteleuropäische Quartär-Faunen. – *Zeitschrift der Deutschen Geologischen Gesellschaft* 32: 468-509.
- Nehring A. (1883): Die ehemalige Verbreitung der Schneehühner in Mitteleuropa. – *Mittheilungen des Ornithologischen Vereines in Wien* 7: 43-45.
- Nehring A. (1904): Diluviale Wirbeltier-Reste aus einer Schlotte des Seveckenberges bei Quedlinburg. – *Zeitschrift der Deutschen Geologischen Gesellschaft* 56: 290-303.
- Nesov L.A. (1992): Mesozoic and Paleogene birds of the USSR and their paleoenvironments. – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Pierce Brodkorb*. – *Natural History Museum of Los Angeles County, Science Series* 36: 465-478.
- Nettleship D.N. & Evans P.G.H. (1985): Distribution and status of the Atlantic Alcidae. – In: Nettleship D.N. & Birkhead T.R. (eds.): *The Atlantic Alcidae*: 53-154. New York: Academic Press.
- Newton A. (1861): Abstract of J. Wolley's researches in Iceland respecting the gare fowl or Great Auk. – *Ibis* 3: 374-399.
- Newton A. (1865): The Gare Fowl and its historians. – *Natural History Review* 1865: 461-488.
- Newton E.T. (1882a): Notes on the vertebrata of the pre-glacial Forest Bed series of the east of England. Part VI – Aves, Reptilia and Amphibia. – *Geological Magazine* 9 (2): 7-9.

- Newton E.T. (1882b): The Vertebrata of the Forest Bed series of Norfolk and Suffolk. – Memoirs of the Geological Survey U.K. [fide Harrison 1979d.]
- Newton E.T. (1885a): *Gastornis klaasseni* Newton, a gigantic bird from the Lower Eocene of Croydon. – Geological Magazine (3) 2: 362-363.
- Newton E.T. (1885b): On the remains of a gigantic species of bird from Lower Eocene beds near Croydon. – Proceedings of the Zoological Society of London 1885: 415-416.
- Newton E.T. (1886a): On the remains of a gigantic species of bird (*Gastornis klaasseni* n. sp.) from the Lower Eocene beds near Croydon. – Transactions of the Zoological Society (London) (12) 5: 143-160.
- Newton E.T. (1886b): Note on the large bird from the Eocene of Croydon, found by Mr. H. M. Klaassen. – Proceedings of the Geological Association of London 9: 349-351.
- Newton E.T. (1887): Note on some recent additions to the vertebrate fauna of the Norfolk pre-glacial bed. – Geological Magazine (3) 4: 145-147.
- Newton E.T. (1889): [Untitled report on a lecture.] – Proceedings of the Zoological Society of London 1889: 220. [In English.]
- Newton E.T. (1891): The Vertebrata of the Pliocene deposits of Britain. – Memoirs of the Geological Survey U.K. [fide Stuart 1974, Harrison & Walker 1977c, and Harrison 1979d.]
- Nitzsch C.L. (1811): Osteographische Beiträge zur Naturgeschichte der Vögel. – Leipzig, x + 122 pp.
- Nogalski S. (1975): Morfologia kośćca wczesnośredniowiecznej kury domowej (*Gallus gallus domesticus*) Pomorza Zachodniego [Morphology of bones of the domestic chicken (*Gallus gallus domesticus*) from the early Medieval Ages of western Pomorze.] – Materiały Zachodniopomorskie 21: 245-320. [In Polish.]
- Nordmann A. (1847): Ueber die Entdeckung reichhaltiger Lagerstätten von fossilen Knochen in Süd-Russland. – In: Jubil. semisaeculare doctor. med. et phil. G. Fischer de Waldheim. Moskva, 11 pp.
- Noriega J.I. (1995): The avifauna from the "Mesopotamian" (Ituzaingó Formation; Upper Miocene) of Entre Rios Province, Argentina. – In: Peters D.S. (ed.): Acta palaeornithologica. – Courier Forschungsinstitut Senckenberg 181: 141-148.
- Northcote M. (1979): Comparative and historical studies of European Quaternary swans and other aquatic birds. – Unpublished PhD. Thesis, Cambridge University, Cambridge, England.
- Northcote M. (1980): Some Cambridgeshire Neolithic to Bronze Age birds and their presence or absence in England in the late-glacial and early Flandrian. – Journal of Archaeological Science 7: 379-383.
- Northcote M. (1981): Size differences between limb bones of Recent and subfossil Mute Swans *Cygnus olor*. – Journal of Archaeological Science 8: 89-98.
- Northcote M. (1982a): The extinct Maltese crane *Grus melitensis*. – Ibis 124: 76-80.
- Northcote M. (1982b): Size, form and habit of the extinct Maltese swan *Cygnus falconeri*. – Ibis 124: 148-158.
- Northcote M. (1982c): Sympatry of Common Cranes *Grus grus* with larger cranes in the last c. 125,000 years. – Bulletin of the British Ornithologists' Club 102: 141-142.
- Northcote M. (1982d): Breeding Bitterns in Cambridgeshire 5,000 years ago. – British Birds 75: 285-286.
- Northcote M. (1983): The giant Maltese swan. – Il-Merill 22: 1-3.
- Northcote M. (1984): Crane *Grus* fossils from the Maltese Pleistocene. – Paleontology 27: 729-735.
- Northcote M. (1985): The giant Maltese crane. – Il-Merill 23: 1-4.
- Northcote M. (1988a): The extinct Pleistocene "swan-goose" *Cygnus (Anser?) equitum* from Malta. – Paleontology 31: 725-740.
- Northcote M. (1988b): The dwarf Maltese swan. – Il-Merill 25: 1-4.

- Northcote M. (1992): Swans (*Cygnus*) and cranes (*Grus*) from the Maltese Pleistocene. – In: Campbell K.E. (ed.): Papers in avian paleontology honoring Pierce Brodkorb. – Natural History Museum of Los Angeles County, Science Series 36: 285-291.
- Northcote M. & Mourer-Chauviré C. (1985): The distinction between the extinct Pleistocene European crane, *Grus primigenia* and the extant Asian Sarus Crane, *G. antigone*. – *Géobios* 18: 877-881.
- Northcote M. & Mourer-Chauviré C. (1988): The extinct crane *Grus primigenia* Milne-Edwards in Mallorca (Spain). – *Géobios* 21: 201-208.
- Novák O. (1877): Fauna der Cyprisschiefer des Egerer Tertiärbeckens. – Sitzungsberichte der Mathematisch-Naturwissenschaftlichen Classe der Kaiserlichen Akademie der Wissenschaften (Wien), I. Abteilung 76: 71-96.
- Nowiński C. & L. Kuźnicki (1965): O rozwoju pojęcia gatunku [On the development of the species notion]. – Warszawa: Państwowe Wydawnictwo Naukowe, 293 pp. [In Polish.]
- Oberholser H.C. (1899): Some untenable names in ornithology. – Proceedings of the Academy of Natural Sciences of Philadelphia 51: 201-216.
- Oberholser H.C. (1905): Notes on the nomenclature of certain genera of birds. – Smithsonian Miscellaneous Collections 48: 59-68.
- Oberholser H.C. (1906): The status of the generic name *Hemiprocne*. – Proceedings of the Biological Society of Washington 19: 67-70.
- Oberholser H.C. (1908): Synopsis of the known species of Swans recent and fossil. – *Emu* 8: 1-11.
- Oberholser H.C. (1917): Mutanda ornithologica I. – Proceedings of the Biological Society of Washington 30: 75-76.
- Olavarría M. de (1898): Huevos fósiles encontrados en Cevico de la Torre (Provincia de Palencia) [Fossil eggs found in Cevico de la Torre (Province of Palencia).] – *Boletín de la Comisión del Mapa Geológico de España* 23: 133-138. [In Spanish.]
- Olsen H. (1967): Varanger-funnene. IV. Osteologisk materiale [Records from Varanger. IV. Osteological material.] – *Tromsø Museum Skrifter* 7 (4). [In Norwegian.]
- Olson S.L. (1974a): *Tantalus milneedwardsii* Shufeldt – a synonym of the Miocene pheasant *Miophasianus altus* (Milne-Edwards). – *Wilson Bulletin* 86: 114-120.
- Olson S.L. (1974b): [Review of Cracraft 1973.] – *Auk* 91: 862-865. [In English.]
- Olson S.L. (1976a): Oligocene fossils bearing on the origins of the Todidae and the Momotidae (Aves: Coraciiformes). – In: Olson S.L. (ed.): Collected papers in avian paleontology honoring the 90th birthday of Alexander Wetmore. *Smithsonian Contributions to Paleobiology* 27: 111-119.
- Olson S.L. (1976b): Fossil woodcocks: an extinct species from Puerto Rico and an invalid species from Malta (Aves: Scolopacidae: Scolopax). – *Proceedings of the Biological Society of Washington* 89: 265-274.
- Olson S.L. (1977): A synopsis of the fossil Rallidae. – In: Ripley S.D. (ed.): *Rails of the world*. Boston: David R. Godine, pp. 339-373.
- Olson S.L. (1978): The nomenclatural status of the taxa of fossil birds attributed to Auguste Aymard. – *Proceedings of the Biological Society of Washington* 91: 444-449.
- Olson S.L. (1981a): The generic allocation of *Ibis pagana* Milne-Edwards, with a review of fossil ibises (Aves: Threskiornithidae). – *Journal of Vertebrate Paleontology* 1: 165-170.
- Olson S.L. (1981b): The museum tradition in ornithology – a response to Ricklefs. – *Auk* 98: 193-195.
- Olson S.L. (1982): A critique of Cracraft's classification of birds. – *Auk* 99: 733-739.
- Olson S.L. (1984): A brief synopsis of the fossil birds from the Pamunkey River and other Tertiary marine deposits in Virginia. – In: Ward L.W. & Krafft K. (eds.): *Stratigraphy*

- and paleontology of the outcropping Tertiary beds in the Pamunkey River region, Central Virginia coastal plain: 217-223. Norfolk: Atlantic Coastal Plain Geological Association.
- Olson S.L. (1985a): The fossil record of birds. – In: Farner D.S., King J.R. & Parkes K.C. (eds.): Avian biology. Vol. 8: 79-256. Orlando: Academic Press.
- Olson S.L. (1985b): A new genus of tropicbird (Pelecaniformes: Phaethontidae) from the middle Miocene Calvert Formation of Maryland. – Proceedings of the Biological Society of Washington 98: 851-855.
- Olson S.L. (1987): An early Eocene oilbird from the Green River Formation of Wyoming (Caprimulgiformes: Steatornithidae). – In: Mourer-Chauviré C. (ed.): L'évolution des oiseaux d'après le témoignage des fossiles. – Documents des Laboratoires de Géologie Lyon 99: 57-69.
- Olson S.L. (1991): Remarks on the fossil record and suprageneric nomenclature of barbets (Aves: Ramphastidae). – Bulletin of the British Ornithologists' Union 111: 222-225.
- Olson S.L. (1992): *Neogaeornis wetzeli* Lambrecht, a Cretaceous loon from Chile (Aves: Gaviidae). – Journal of Vertebrate Paleontology 12: 122-124.
- Olson S.L. (1994): A giant *Presbyornis* (Aves: Anseriformes) and other birds from the Paleocene Aquia Formation of Maryland and Virginia. – Proceedings of the Biological Society of Washington 107: 429-435.
- Olson S.L. (1995a): *Thiornis sociata* Navás, a nearly complete Miocene grebe (Aves: Podicipedidae). – In: Peters D.S. (ed.): Acta palaeornithologica. – Courier Forschungsinstitut Senckenberg 181: 131-140.
- Olson S.L. (1995b): History and nomenclature of avian family-group names. – Auk 112: 539-546. [Review of Bock 1994.]
- Olson S.L. (1997): [Review of Mlíkovský 1996a.]. – Auk 114: 537-538. [In English.]
- Olson S.L. (1999a): The anseriform relationships of *Anatalavis* Olson and Parris (Anseranatidae), with a new species from the Lower Eocene London Clay. – In: Olson S.L. (ed.): Avian paleontology at the close of the 20th century. – Smithsonian Contributions to Paleobiology 89: 231-243.
- Olson S.L. (1999b): Early Eocene birds from eastern North America: a faunule from the Nanjemoy Formation of Virginia. – In: Weems R.E. & Grimsley G.J. (eds.): Early Eocene vertebrates and plants from the Fisher/Sullivan Site (Nanjemoy Formation) Stafford County, Virginia. – Virginia Division of Mineral Resources, Publication 152: 123-132.
- Olson S.L. (1999c): A new species of pelican (Aves: Pelecanidae) from the Lower Pliocene of North Carolina and Florida. – Proceedings of the Biological Society of Washington 112: 503-509.
- Olson S.L. & Daniels M. (1994): [Untitled report.]. – Society of Avian Paleontology and Evolution, Information Letter 8: 10. [In English.]
- Olson S.L. & Feduccia A. (1979): An Old World occurrence of the Eocene avian family Primobucconidae. – Proceedings of the Biological Society of Washington 92: 494-497.
- Olson S.L. & Feduccia A. (1980a): Relationships and evolution of flamingos (Aves: Phoenicopteridae). – Smithsonian Contributions to Zoology 316: i-iii, 1-73.
- Olson S.L. & Feduccia A. (1980b): *Presbyornis* and the origin of the Anseriformes (Aves: Charadriomorphae). – Smithsonian Contributions to Zoology 323: i-iv, 1-24.
- Olson S.L. & Parris D.C. (1987): The Cretaceous birds of New Jersey. – Smithsonian Contributions to Paleobiology 63: i-iii, 1-22.
- Olson S.L. & Steadman D.W. (1981): The relationships of the Pedionomidae (Aves: Charadriiformes). – Smithsonian Contributions to Zoology 337: i-iii, 1-25.
- Olson S.L. & Rasmussen P. (2001): Miocene and Pliocene birds from the Lee Creek Mine, North Carolina. – In: Ray C.E. & Bohaska D.J. (eds.): Geology and paleontology

- of the Lee Creek Mine, North Carolina, III. – *Smithsonian Contributions to Paleobiology* 90: 233-365. [Received too late to be used. Main taxonomic conclusions relevant to European birds are as follows: *Fulica pisana* Portis was synonymized with the modern *Podiceps auritus* (Linnaeus) (p. 248); *Diomedea anglica* Lydekker was transferred to the modern diomedeid genus *Phoebastria* Reichenbach (p. 249); and *Uria ausonia* Portis was transferred to the modern alcid genus *Alca* Linnaeus (p. 272).]
- Olson S.L. & Walker C.A. (1997): A trans-Atlantic record of the fossil tropicbird *Heliadornis ashbyi* (Aves: Phaethontidae) from the Miocene of Belgium. – *Proceedings of the Biological Society of Washington* 110: 624-628.
- Olson S.L., Swift C.C. & Mokhiber C. (1979): An attempt to determine the prey of the Great Auk (*Pinguinus impennis*). – *Auk* 96: 790-792.
- Omboni G. (1885): Penne fossili del Monte Bolca [Fossil feathers from Monte Bolca.] – *Atti Reale Istituto Veneto di Scienze, Lettere e Arti* (6) 3: 1-7. [In Italian.]
- Osborn H.F. (1917): Biocharacters as separable units of organic structure. – *American Naturalist* 51: 449-456.
- Owen R. (1840): Description of the remains of a mammal, a bird, and a serpent from the London Clay. – *Proceedings of the Geological Society of London* 3: 162-166.
- Owen R. (1841): Description of the fossil remains of a mammal (*Hyracotherium leporinum*) and of a bird (*Lithornis vulturinus*) from the London Clay. – *Transactions of the Geological Society of London* 6: 203-208.
- Owen R. (1843): [Untitled report.] – *Proceedings of the Zoological Society of London* 11: 19. [In English.]
- Owen R. (1846): A history of British fossil mammals and birds. London: John Van Voorst, xlvii + 558 pp. [Birds on pp. 545-558.]
- Owen R. (1856a): On the affinities of the large extinct bird (*Gastornis parisiensis*, Hébert), indicated by a fossil femur and tibia discovered in the Lowest Eocene Formation near Paris. – *Quarterly Journal of the Geological Society of London* 12: 204-217.
- Owen R. (1856b): Sur les affinités du *Gastornis parisiensis* Hébert, grand oiseau fossile trouvé dans le terrain éocène inférieur des environs de Paris. – *L'Institut (Paris)* 24: 283.
- Owen R. (1860): Darwin on the origin of species. – *Edinburgh Review* 111: 487-532.
- Owen R. (1869): [Untitled report on a lecture.] – *Proceedings of the Zoological Society of London* 1869: 59. [In English.]
- Owen R. (1870): On *Dinornis* (Part 14), containing contributions to the craniology of the genus, with a description of a fossil cranium of *Dasornis londinensis* Ow., from the London Clay of Sheppey. – *Transactions of the Zoological Society of London* 7: 123-150.
- Owen R. (1873): Description of the skull of a dentigerous bird (*Odontopteryx toliapica*) from the London Clay of Sheppey. – *Quarterly Journal of the Geological Society of London* 29: 511-522.
- Owen R. (1878): On *Argillornis longipennis*, a large bird of flight from the Eocene Clay of Sheppey. – *Quarterly Journal of the Geological Society of London* 34: 124-131.
- Owen R. (1880): On the skull of *Argillornis longipennis*. – *Quarterly Journal of the Geological Society of London* 36: 23-26.
- Palmowski J. & Wachendorf H. (1966): Eine unteroligozäne Wirbeltierfauna aus einer Spaltenfüllung in Herrlingen/Blau (Württ.). – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Historische Geologie* 6: 229-245.
- Panin N. (1961): Asupra unor urme organice și mecanice din miocenul de la confluența Putnei cu Zăbala [On new organical and mechanical imprints in the Miocene of the Putna and Zăbala confluence.] – *Studii și Cercetări de Geologie, Geofizică și Geografie, Seria Geologie* 6: 63-73. [In Romanian.]
- Panin N. (1964): Coexistența urmelor de pași de vertebrate cu mecanoglifele în molasa

- miocenă din Carpații Orientali [Coexistence of vertebrate footprints with mechanoglyphs in the Miocene molasse of East Carpathians.] – Studii și Cercetari de Geologie, Geofizică și Geografie, Seria Geologie 9: 341-363. [In Romanian; see Panin 1965 for the French translation.]
- Panin N. (1965): Coexistence des traces de pas de vertébrés et des mecanoglyphes dans la molasse miocène des Carpates Orientales. – Revue Roumaine de Géologie, Géophysique et de Géographie, Serie de Géologie 9: 141-163. [French version of Panin 1964.]
- Panin N. & Avram E. (1962): Noi urme de vertebrate in miocenul Subcarpaților românești [New vertebrate footprints in the Miocene of the Romanian Subcarpathians.] – Studii și Cercetari de Geologie, Geofizică și Geografie, Seria Geologie 7: 455-484. [In Romanian.]
- Panin N., Lăzărescu V. & Grujinchil C. (1966): Un nou sector ichnofosilifer in terenurile miocene din Carpații Orientali [New ichnofossiliferous section in the Miocene deposits in East Carpathians.] – Buletinul Institutului de Petrol, Gaze și Geologie 16: 9-21. [In Romanian.]
- Pankakoski E., Väisänen R.A. & Nurmi K. (1987): Variability of muskrat skulls: measurement error, environmental modification and size allometry. – Systematic Zoology 36: 35-51.
- Panteleev A.V. (1999): [Untitled communication.] – Society of Avian Paleontology and Evolution, Information Letter 13: 19-20. [In English.]
- Paris P. (1912): Oiseaux fossiles de France. – Revue Française d'Ornithologie 37: 283-298.
- Parker W.K. (1865): Preliminary notes on some fossil birds from the Zebbug Cave, Malta. – Proceedings of the Zoological Society of London 33: 752-753.
- Parker W.K. (1869): On some fossil birds from the Zebbug Cave, Malta. – Transactions of the Zoological Society of London 6: 119-124.
- Parkinson J. (1808-1811) Organic remains of a former world. An examination of the mineralized remains of the vegetables and animals of the antediluvian world, generally termed extraneous fossils. Vol. 1-3. London: Whittingham and Rowland, xii + 461, xvi + 286, and xvi + 455 pp.
- Parry S. (1996): The avifaunal remains. – In: Conway B., McNabb S. & Ashton B. (eds.): The excavations at Barnfield pit, Swanscombe 1968-1972. – British Museum, Occasional Papers 94: 137-143.
- Patterson B. (1941): A new phororhacoid bird from the Deseado Formation of Patagonia. – Field Museum of Natural History, Geology Series 8: 49-54.
- Patterson B. & Kraglievich L.J. (1960): Sistemática y nomenclatura de las aves fororracoideas del Plioceno Argentino. – Publicaciones del Museo Municipal de Ciencias Naturales y Tradicionales Mar del Plata 1: 1-51.
- Paucă M. (1942): Empreintes de pas de palmipèdes dans l'Helvétien carpatique du depar. de Putna. – Bulletin de la Société Roumaine de Géologie 5: 85-88.
- Pavia M. (1999a): Un cranio di *Bubo insularis* Mourer-Chauviré & Weesie, 1986 (Aves, Strigidae) nelle breccie ossifere del Pleistocene di Capo Figari (Sardegna, Italia) [A cranium of *Bubo insularis* Mourer-Chauviré & Weesie, 1986 (Aves, Strigidae) from the ossiferous breccia from the Pleistocene of Capo Figari (Sardinia, Italy).] – Accademia delle Scienze di Torino, Atti Scienze Fisiche 133: 1-10. [In Italian.]
- Pavia M. (1999b): The middle Pleistocene avifauna of Spinagallo Cave (Sicily, Italy): preliminary report. – In: Olson S.L. (ed.): Avian paleontology at the close of the 20th century. – Smithsonian Contributions to Paleobiology 89: 125-127.
- Pavia M. (1999c): Le avifaune pleistoceniche italiane: stato delle conoscenze [The Pleistocene avifauna of Italy: the state of research.] – Avocetta 23: 146. [In Italian.]
- Pavia M. & Mourer-Chauviré C. (2000): An overview on the genus *Athene* in the

- Pleistocene Mediterranean islands, with the description of a new species of *Athene* (Aves, Strigiformes). – *Vertebrata Palasiatica* 38, Supplement: 24-25. [Abstract.]
- Penck A. & Brückner E. (1901-1909): *Die Alpen im Eiszeitalter*. Vol. 1-3. Leipzig: Tauchnitz, 1199 pp.
- Péquart M. & Péquart S.J. (1926): Un kjökkenmödding morbihannais: Er Yoh. Premier outillage en os découvert dans le Morbihan. – *Revue de Anthropologie* 36: 206-211.
- Péquart M., Péquart S.J., Boule M. & Vallois H. (1937): Tévéc, station-nécropole mésolithique du Morbihan. – *Archives de l'Institut de Paléontologie Humaine* 18: 1-227.
- Peters D.S. (1983): Die "Schnepfenralle" *Rhynchaetes messelensis* Wittich 1898 ist ein Ibis. – *Journal für Ornithologie* 124: 1-27.
- Peters D.S. (1985): Ein neuer Segler aus der Grube Messel und seine Bedeutung für den Status der Aegialornithidae (Aves: Apodiformes). – *Senckenbergiana Lethaea* 66: 143-164.
- Peters D.S. (1987a): Ein "Phorusrhacide" aus dem Mittel-Eozän von Messel (Aves: Gruiformes: Cariamae). – In: Mourer-Chauviré C. (ed.): *L'évolution des oiseaux d'après le témoignage des fossiles*. – *Documents de Laboratoires de Géologie Lyon* 99: 71-87.
- Peters D.S. (1987b): *Juncitarsus merkei* n.s. stützt die Ableitung der Flamingos von Regenspfeifervögeln (Aves: Charadriiformes: Phoenicopteridae). – In: Peters D.S. & Wiltschko W. (eds.): *Aspekte der Ornithologie*. – *Courier Forschungsinstitut Senckenberg* 97: 141-155.
- Peters D.S. (1988a): Ein vollständiges Exemplar von *Palaeotis weigelti* (Aves: Palaeognathae). – *Courier Forschungsinstitut Senckenberg* 107: 223-233.
- Peters D.S. (1988b): Die Messelvögel - eine Landvogelfauna. – In: Schaal S. & Ziegler W. (eds.): *Messel - ein Schaufenster in die Geschichte der Erde und des Lebens*: 135-152. Frankfurt am Main: Waldemar Kramer.
- Peters D.S. (1989): Fossil birds from the oil shale of Messel (Lower Middle Eocene, Lutetian). – In: Ouellet H. (ed.): *Acta XIX Congressus internationalis ornithologici*: 2056-2064. Ottawa: University of Ottawa Press.
- Peters D.S. (1991): Zoogeographical relationships of the Eocene avifauna from Messel (Germany). – In: *Acta XX congressus internationalis ornithologici*. Vol. 1: 572-577. Wellington.
- Peters D.S. (1992): A new species of owl (Aves: Strigiformes) from the Middle Eocene Messel oil shale. – In: Campbell K.E. (ed.): *Papers in avian paleontology honoring Pierce Brodkorb*. – *Natural History Museum of Los Angeles County, Science Series* 36: 161-169.
- Peters D.S. (1994): *Messelastur gratulator* n. gen. n. sp., ein Greifvogel aus der Grube Messel (Aves: Accipitridae). – *Courier Forschungsinstitut Senckenberg* 170: 3-9.
- Peters D.S. (1995): *Idiornis tuberculata* n. spec., ein weiterer ungewöhnlicher Vogel aus der Grube Messel. – In: Peters D.S. (ed.): *Acta palaeornithologica*. – *Courier Forschungsinstitut Senckenberg* 181: 107-119.
- Peters D.S. (1997a): Ein verwandter der Mausvögel. – In: Koeningswald W. von & Storch G. (eds.): *Messel – Ein Pompeji der Paläontologie*: 122-123. Sigmaringen: Jan Thorbecke.
- Peters D.S. (1997b): Ein Kranichvogel. – In: Koeningswald W. von & Storch G. (eds.): *Messel – Ein Pompeji der Paläontologie*: 124-125. Sigmaringen: Jan Thorbecke.
- Peters D.S. (1997c): Ein früher Greifvogel. – In: Koeningswald W. von & Storch G. (eds.): *Messel – Ein Pompeji der Paläontologie*: 128-129. Sigmaringen: Jan Thorbecke.
- Peters D.S. (1998): Erstnachweis eines Seglers aus dem Geiseltal (Aves: Apodiformes). – *Senckenbergiana Lethaea* 78: 211-212.
- Peters D.S. (1999): *Selmes absurdipes*, new genus, new species, a sandcoleiform bird from the oil shale of Messel (Germany, middle Eocene). – In: Olson S.L. (ed.): *Avian paleontology at the close of the 20th century*. – *Smithsonian Contributions to*

- Paleobiology 89: 217-222.
- Peters D.S. & Hamedani A. (2000): *Frigidafons babaheydariensis* n. sp., ein Sturmvogel aus dem Oligozän des Irans (Aves: Procellariidae). – *Senckenbergiana Lethaea* 80: 29-37.
- Peters D.S. & Storch G. (1993): South American relationships of Messel birds and mammals. – *Kaupia* 3: 263-269.
- Petersen A. (1995): Some aspects of the history of the Great Auk in Iceland. – *Naturufraedingurinn* 65: 53-66.
- Petersen K.S., Hoch E. & Bonde N. (1973): A new species of Mytilid bivalve, and vertebrate remains from Lower Eocene marine deposits on Røsnaes, Denmark. – *Danmarks Geologiske Undersøgelse, Årbog 1972*: 79-86.
- Petrov S.G. (1951): Proischoždenie i evoljucija domašnich kur [The origin and evolution of domestic chickens.] – Unpub. Diss., Moskva, 185 + 210 pp. [In Russian.]
- Peyer B. (1949): *Protornis glaronensis* H. v. Meyer. Ergebnisse der Präparation. – *Verhandlungen der Schweizerischen Naturforschenden Gesellschaft* 129: 141-142.
- Peyer B. (1950): Geschichte der Tierwelt. – Zürich: Büchergilde Gutenberg, 288 pp.
- Peyer B. (1957): *Protornis glaronensis* H. v. Meyer. Neubeschreibung des Typusexemplars und eines weiteren Fundes. – *Schweizerische Paläontologische Abhandlungen* 73 (1): 1-47.
- Pfannenstiel M. (1958): Fälscher und Fälschungen von Oehninger Fossilien. – *Geologie* 7: 846-860.
- Philippe M., Mourer-Chauviré C. & Evin J. (1980): Les gisements paléontologiques quaternaires des Cuasses de Martel et de Gramat (Corrèze et Lot): faunes et chronologie. – *Nouvelles Archives de la Musée d'Histoire naturelle de Lyon* 18: 57-67.
- Pichon J. & Tchernov E. (1987): Un nouveau coq fossile, *Gallus gallus levantinus* n. subsp. (Aves, Galliformes), dans de Pléistocène d'Israel. – In: Mourer-Chauviré C. (ed.): L'évolution des oiseaux d'après le témoignage des fossiles. – *Documents des Laboratoires de Géologie Lyon* 99: 201-209.
- Pictet F.J. (1844): *Traité élémentaire de paléontologie*. – Paris: J. B. Ballière.
- Pictet F.J. (1853): *Traité élémentaire de paléontologie*. Vol. 2. 2nd ed. – Paris: J. B. Ballière.
- Pidopličko I.G. (1956): Materialy do vyvčennja mynulych faun URSR [Materials on the research of the past faunas of Ukrainian SSR.]. Vol. II. Kyjiv: Naukova Dumka, 235 pp. [In Ukrainian.]
- Pidopličko I.G. & Goldin G.K. (1964): Naibolee severnaja nachodka skorlupy jajca strausa v evropejskoj časti SSSR [The northernmost record of eggshell of ostrich egg in the European part of the USSR.]. – In: Strautman F.I. (ed.): Problemy ornitologii [The problems of ornithology.]: 190-191. L'viv: Izdatel'stvo L'vovskogo Universiteta. [In Russian.]
- Pieper H. (1985): The fossil land birds of Madeira and Porto Santo. – *Bocagiana* 88: 1-6.
- Piller M. & Mitterpacher L. (1783): Iter per poseganum sclavoniae provinciam mensibus junio, et julio anno MDCCLXXXII [An expedition to the provinces of Slavonic peasants in the months June and July of the year 1782.]. – Budapest: Typis Regiae Universitatis, 148 pp. [In Latin.]
- Pitton L.E. (1940): Paléontologie du gisement éocène de Menat (Puy-de-Dôme). Flore et faune. – Paris: Lechevalier, 306 pp.
- Piveteau J. (1955): Oiseaux. – In: Piveteau J. (ed.): *Traité de paléontologie*. Tome 5: Amphibiens – Reptiles – Oiseaux: 994-1091. Paris: Masson et C^{ie}. [With a contribution by C. Dechaseaux.]
- Plaziat J.C. (1964): Pistes d'Oiseaux et remaniements synsédimentaires dans le Lutétien du détroit de Carcassonne (Aude). – *Bulletin de la Société Géologique de France* (7) 6: 289-293.

- Pons J. & Roca Ll. (1974): Restos de grulla fòsil en el »Avenc de Na Corna« (Artù) [Remains of a fossil crane in the »Avenc de Na Corna« (Artù).] – Endins 1: 17-19. [In Catalan.]
- Popov R., Kacarov G. & Gospodinov V. (1921): Predistoričeski i starochristijanski pametnici ot Sofija i okolnostta [Prehistoric and early Christian monuments of Sofija and its surroundings.] – Materialy za Istorijata na Sofija 5: 1-36. [In Bulgarian.]
- Portis A. (1879): Intorno ad alcune impronte eoceniche di Vertebrati recentemente scoperte in Piemonte [On some Eocene vertebrate imprints, recently found in Piemonte.] – Atti Regia Accademia Scienze (Torino) 15: 221-228. [In Italian.]
- Portis A. (1884): Contribuzioni alla ornitolitologia italiana [Contributions to the Italian ornitholithology.] – Memorie Regia Accademia Scienze (Torino) (2) 36: 361-384. [In Italian.]
- Portis A. (1887): Contribuzioni alla ornitolitologia italiana. Parte II [Contributions to the Italian ornitholithology. Part II.] – Memorie Regia Accademia Scienze (Torino) (2) 38: 181-203. [In Italian.]
- Portis A. (1889): Gli ornitoliti del Valdarno superiore e di alcune altre località plioceniche di Toscana [Ornitholiths from the upper Valdarno and some other Pliocene localities of Tuscany.] – Memorie Regio Istituto di Studi Superiori e Practici (Firenze) 1889: 1-20. [In Italian.]
- Portis A. (1900): Di una formazione stagnale presso la Basilica ostiense di Roma e degli avanzi fossili vertebrati in essa rinvenuti [On a lake formation from the Ostian Basilica of Roma, and remains of fossil vertebrates found in it.] – Bolletino della Società Geologica Italiana 19: 179-240. [In Italian.]
- Potapov R.L. (1988): O proischoždenii i èvoljucii belych kuropatok [On the origin and evolution of *Lagopus* grouse.] – In: Andreev A.V. & Kondrat'ev A.Ja. (eds.): Izučenie i ochrana ptic v èkosistemach Severa [Research and conservation of birds in the ecosystems of the North]: 170-176. Vladivostok: Akademija Nauk SSSR. [In Russian.]
- Potapova O.R. (1986): Belye kuropatki (rod *Lagopus*) v plejstocene Severnogo Urala [Grouse (genus *Lagopus*) in the Pleistocene of northern Ural.] – In: Potapov R.L. (ed.): Èkologičeskie i faunističeskie issledovanija ptic [Ecological and faunistic studies of birds.] – Trudy Zoologičeskogo Instituta Akademii Nauk SSSR 147: 46-58. [In Russian.]
- Potapova O.R. (1990): Ostatki ptic iz plejstocenovykh otloženiij Medvež'ej peščery na severnom Urale [Remains of birds from the Pleistocene deposits of the Medvež'ja cave in northern Ural.] – In: Kuz'mina I.E. & Baryšnikov G.F. (eds.): Fauna mlekopitajuščich i ptic pozdnego plejstocena i golocena SSSR [Fauna of mammals and birds of the late Pleistocene and Holocene of the USSR.] – Trudy Zoologičeskogo Instituta Akademii Nauk SSSR 212: 135-153. [In Russian.]
- Potapova O.R. (1994): The first record of the Great Auk *Pinguinus impennis* in Russia. – In: The ornithological notebook of the XXI international congress (Wien, 20-25 August 1994): 723. [Abstract.]
- Potapova O.R. (2000): The Pleistocene records of "wild Junglefowl" in the Caucasus and Trascaucasia: "*Gallus*" remains revision. – Vertebrata Palasiatica 38 (Supplement): 25. [Abstract.]
- Potapova O.R. & Baryšnikov G.F. (1993) Pticy iz Ašel'skoj stojanki v peščere Treugol'naya na severnom Kavkaze [Birds from the Acheulean site in Treugol'naja cave in northern Caucasus.] – Trudy Zoologičeskogo Instituta Russkoj Akademii Nauk 249: 48-65. [In Russian.]
- Poulianos N.A. (1996): Pleistocene mammals of Petralona cave, Chalkidiki, N. Greece. Unpub. Thesis, Charles University, Praha, v + 379 pp.
- Prévost C. (1855a): Annonce de la découverte d'un oiseau fossile de taille gigantesque, trouvé à la partie inférieure de l'argile plastique des terrains parisiens. – Comptes

- Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris) 40: 554-557.
- Prévost C. (1855b): Nouveaux documents sur le gisement du *Gastornis parisiensis* et considérations générales sur les vestiges laissés par des oiseaux dans les terrains des divers âges. – Comptes Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris) 40: 616-619.
- Preyer W. (1862): Über *Plautus impennis*. – Journal für Ornithologie 10: 110-124, 337-356.
- Prince G. (ed., 1998): Encyclopedia of the languages of Europe. – Oxford: Blackwell Publishers, xviii + 499 pp.
- Probst J. (1879): Verzeichniss der Fauna und Flora der Molasse im Württembergischen Oberschwaben. – Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg 35: 221-304.
- Procaccini Ricci V. (1840): Filliti ed altri resti organici trovati nelle gessaie di S. Angelo e S. Gaudenzio presso Sinigaglia [Leaves and other organic remains found in the gypsum deposits of San Angelo and San Gaudenzio near Senigaglia.] – Bolletino della Società Geologica Italiana 19: 179-240. [In Italian.]
- Prum R.O. (1988): Phylogenetic interrelationships of the barbets (Aves: Capitonidae) and toucans (Aves: Ramphastidae) based on morphology with comparisons to DNA-DNA hybridization. – Zoological Journal of the Linnean Society (London) 92: 313-343.
- Prummel W. (1993): Starigard/Oldenburg. Hauptburg der Slawen in Wagrien. IV. Die Tierknochenfunde unter besonderer Berücksichtigung der Beizjagd. – Neumünster: Karl Wachholtz Verlag, 186 pp.
- Pržemysskij K. (1912): Zametka o novom mestonachoždenii "Pikermijskoj fauny" v meotičeskich slojach doliny Bol'sogo Kujal'nika v okr. g. Odessy [Note on a new locality of the "Pikermi fauna" in the Meotian deposits in the vicinity of Odessa city.] – Zapiski Novorossijskogo Obščestva Estesvoispytatelej 38: 25-38. [In Russian.]
- Pržemysskij K. (1914): Fauna pozvonočnych meotičeskich sloev iz okresnostej g. Odessy [Vertebrate fauna of the Meotian deposits from the vicinity of Odessa city.] – Zapiski Novorossijskogo Obščestva Estesvoispytatelej 39: 401-468. [In Russian.]
- Pycraft W.P. (1909): On a fossil bird from the Lower Pliocene. – Proceedings of the Zoological Society of London 1909: 367-370.
- Quekett J.T. (1855): Descriptive and illustrated catalogue of the histological series contained in the Museum [= Hunterian Museum, Royal College of Surgeons]. Vol. 2. Structure of the skeleton of vertebrate animals. London: The College.
- Quenstedt F.A. (1853): [Untitled communication.] – Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg 9: 64-75. [In German.]
- Raaf J.F.M. de, Beets C. & Kortenbout van der Sluijs G. (1965): Lower Oligocene bird-tracks from northern Spain. – Nature (London) 207: 146-148.
- Rădan S. & Brustur T. (1993): Urme de pași de păsări în oligocenul superior din bazinul Petroșani (Carpații Meridionali) [Avian footprints from the late Oligocene of Petroșani Basin (South Carpathians).] – Studii și Cerșetari de Geologie 38: 71-80. [In Romanian.]
- Rage J.-C. (1999): Faunes à affinités sud-américaines dans le Paléogène d'Europe: état de la question. Allons-nous vers une nouvelle interprétation? – Bulletin de la Société Paléontologique de France 170: 951-954.
- Ranke J. (1879): Die natürlichen Höhlen in Bayern. 1. Das Zwergloch und Hasenloch bei Pottenstein in Oberfranken. – Beiträge zur Athropologie und Urgeschichte Bayerns 2: 195-225.
- Rasmussen D.T., Olson S.L. & Simons E.L. (1987): Fossil birds from the Oligocene Jebel Qatrani Formation, Fayum Province, Egypt. – Smithsonian Contributions to Paleobiology 62: i-iii, 1-20.
- Rasmussen P.C. (1998): Early Miocene avifauna from the Pollack Farm Site, Delaware. – In: Benson R.N. (ed.): Geology and paleontology of the lower Miocene Pollack Farm

- Fossil Site, Delaware. – Delaware Geological Survey, Special Publication 21: 149-151.
- Rauscher K. (1984): Ein Fischadlerei aus dem Mittelmiozän von Kärnten (Österreich). – Beiträge zur Paläontologie von Österreich 11: 61-69.
- Ray C.E., Wetmore A. & Dunkle D.H. (1968): Fossil vertebrates from the marine Pleistocene of southeastern Virginia. – Smithsonian Miscellaneous Collection 153(3): 1-25.
- Rayner J.M.V. (1985): Linear relations in biomechanics. – Journal of Zoology (London) (A) 206: 415-439.
- Razoumowsky A. (1789) [Title?] – Mém. de Lausanne 3. [fide Lambrecht 1933]
- Regàlia E. (1897): Fossilli dell'Is. di Tavolara (Sardegna e di Grive St. Alban (Isère) [Fossils from the island of Tavolara (Sardinia) and from Grive St. Alban (Isère).] – Avicula 1: 164-165. [In Italian; parenthesis after Sardegna omitted in the original title.]
- Regàlia E. (1902): Sette uccelli pliocenici del Pisano e de Valdarno superiore [Seven Pliocene birds from Pisano and upper Valdarno.] – Palaeontografia Italica 8: 219-238. [In Italian.]
- Regàlia E. (1904): Sulla fauna di Grotta Romanelli (Castro, Lecce). – Archivio per l'Antropologia e la Etnologia 34:33-85.
- Regàlia E. (1907): Avifaune fossili Italiane [Italian fossil avifauna.] – Avicula 11: 49-54, 79-84. [In Italian.]
- Reichenbach H.G.L. (1852): Das natürliche System der Vögel. – Dresden: Expedition der vollständigen Naturgeschichte, & Berlin: Hofmeister, viii + xxxi + 36 pp.
- Reif W.-E. (1984): Artabgrenzung und das Konzept der evolutionären Art in der Paläontologie. – Zeitschrift für Zoologische Systematik und Evolutionsforschung 22: 263-286.
- Remane J. (1985): Der Artbegriff in Zoologie, Phylogenetik und Biostratigraphie. – Paläontologische Zeitschrift 59: 171-182.
- Rensch B. (1940): Die ganzheitliche Auswirkung der Größenauslese am Vogelskelett. – Journal für Ornithologie 88: 373-388.
- Repelin J. (1924): Les restes des grands oiseaux coureurs d'autrefois au Muséum de Marseille. – Annales du Muséum d'Histoire Naturelle de Marseille 19: 107-112.
- Rey R. (1969): Observations à la note de R. Dughi et F. Sirugue relative à l'âge des marnes à oeufs d'oiseaux en Provence. – Comptes Rendus Hebdomadaires de la Société Géologique de France 1969: 305-306.
- Rhymer J.M. & Simberloff D. (1996): Extinction by hybridization and introgression. – Annual Review of Ecology and Systematics 27: 83-109.
- Rich P.V. (1972): A fossil avifauna from the Upper Miocene Beglia Formation of Tunisia. – Notes du Service Géologique 35: 29-66.
- Rich P.V. (1976): Significance of the Tertiary avifaunas from Africa (with emphasis on a mid to late Miocene avifauna from southern Tunisia. – Annals of the Geological Survey of Egypt 4: 167-210.
- Rich P.V. (1982): Tarsometatarsus of *Protostrix* from the mid-Eocene of Wyoming. – Auk 99: 576-579.
- Rich P.V. & Haarhoff P.J. (1985): Early Pliocene Coliidae (Aves, Coliiformes) from Langebaanweg, South Africa. – Ostrich 56: 20-41.
- Rich P.V. & Walker C.A. (1983): A new genus of Miocene flamingo from East Africa. – Ostrich 54: 95-104.
- Richmond C.W. (1902): List of generic terms proposed for birds during the years 1890-1900 inclusive, to which are added names omitted by Waterhouse in his "Index Generum Avium". – Proceedings of the United States National Museum 24: 663-729.
- Richmond C.W. (1917): Generic names applied to birds during the years 1906 to 1915, inclusive, with additions and corrections to Waterhouse's »Index Generum Avium«. – Proceedings of the United States National Museum 53: 565-636.

- Ristori G. (1891): Ornitoliti di Montebamboli [Ornitholiths of Montebamboli.] – Atti della Società Toscana di Scienze Naturali Residente i Pisa, Processi Verbali 7: 308-309. [In Italian.]
- Rivière E. (1887): Faune des oiseaux, des reptiles et des poissons trouvés des Baoussé-Roussé (Italie) dites Grotte de Menton. – Matériaux pour l'Histoire Primitive et Naturelle de l'Homme (3) 20: 525-535.
- Rivière E. (1892): Nouvelle station quaternaire sur les bords de la Vézère. L'abri sous roche de Pageyral. – Association Française pour l'Avancement des Sciences 20: 372-378.
- Roberton (1834): [Sur un ornithocoprolite trouvé dans un lieu nommé les Meilleries-Saint-Sauveur.] – Bulletin de la Société Géologique de France 4: 415.
- Robineau-Desvoidy A.J.B. (1863): Histoire naturelle des Diptères des environs de Paris. Vol. 2. – Paris.
- Rogovič A.S. (1875a): O pervobytnom mestonachoždenii jantarja okolo Kieva [On a primaeval locality of amber near Kyjiv.] – Trudy S"ezda Russkich Estestvoispytatelej, Otdel Chimii, Mineralogii, Geologii i Paleontologii 4: 81-86. [In Russian.]
- Rogovič A.S. (1875b): Zаметка о местонахождении костей ископаемых животных в юго-западной России [Note on a locality with bones of fossil mammals in southwestern Russia.] – Zapiski Kievskogo Obščestva Estestvoispytatelej 4 (1). [In Russian.]
- Rohlf F.J., Gilmartin A.J. & Hart G. (1983): The Kluge-Kerfoot phenomenon – A statistical artifact. – Evolution 37: 180-202.
- Röhrs M. (1959): Neue Ergebnisse und Probleme der Allometrieforschung. – Zeitschrift für Wissenschaftliche Zoologie 162: 1-96.
- Röhrs M. (1961): Allometrieforschung und biologische Formenanalyse. – Zeitschrift für Morphologie und Anthropologie 51: 289-321.
- Rolle F. (1863): Ch. Darwin's Lehre von der Entstehung der Arten im Pflanzen- und Thierreich in ihrer Anwendung auf die Schöpfungsgeschichte. – Frankfurt am Main, vi + 274 pp.
- Rolle F. (1887): Vögel. – In: Kengott A. (ed.): Handwörterbuch der Mineralogie, Geologie und Palaeontologie. Vol. 3: 392-403. Breslau [= Wrocław]: Verlag von Eduard Trewendt.
- Ronsil R. (1948): Bibliographie ornithologique française. Vol. 1-2. – Paris: Paul Lechevalier, 535 + 100 pp.
- Rosas A., Carbonell E., Cuenca G., Garcia N., Fernandez Jalvo Y., van der Made J., Ollé A., Pares J. M., Perez Gonzalez A., Sánchez Marco A., Sánchez Chillón B. & Vallverdu J. (1998): Cronología, bioestratigrafía y paleoecología del Pleistoceno Medio de Galeria (Sierra de Atapuerca, España) [Chronology, biostratigraphy and paleoecology of the middle Pleistocene of Galeria (Sierra de Atapuerca, Spain).] – Revista Española de Paleontología 13: 71-80. [In Spanish.]
- Rosas, A. et al. (2001): Le gisement Pléistocène de la «Sima del Elefante» (Sierra de Atapuerca, Espagne). L'Anthropologie 105: 301-312.
- Roščin A.D. (1959): Znachidka na Odeščyni jajca vykopnogo strausa [Record of an egg of an fossil ostrich in the Odessa region.] – Dopovidi Akademiji Nauk Ukrajin's'koji RSR 1959: 535-537. [In Ukrainian.]
- Roščin A.D. (1962): Nachodki jaic iskopajemych strausov na jube Ukrainy [Records of eggs of fossil ostriches in southern Ukraine.] – Trudy Odesskogo Gosudarstvennogo Universiteta imeni I. I. Mečnikova 152: 26-30. [In Russian.]
- Roset F. (1974): Nuevos yacimientos de mamíferos fósiles en la cuenca miocénica de Seu d'Urgell (Lérida) (Nota preliminar) [New finds of fossil mammals in the Miocene cavern of Seu d'Urgell (Lérida) (A preliminary note).] – Boletín Informativo del Instituto Provincial de Paleontología de Sabadell 2: 38-42. [In Spanish.]
- Roška V.Ch. (1967): Schema stratigrafii sarmatskich otloženij Moldavii [A scheme of the

- Sarmatian deposits of Moldova.] – *Izvestija Akademii Nauk Moldavskoj SSR* 1967 (4): 72-80. [In Russian.]
- Roth J. & Wagner A. (1855): Die fossilen Knochenüberreste von Pikermi in Griechenland. – *Abhandlungen der Mathematisch-Physicalischen Klasse der Bayerischen Akademie der Wissenschaften* 7: 371-464.
- Rothausen K. (1966): Fossile Flamingos im Tertiär des Mainzer Beckens. – *Zeitschrift der Rheinischen Naturforschenden Gesellschaft in Mainz* 4: 26-31.
- Rothschild W., Hartert E. & Kleinschmidt O. (1897): *Comatibis eremita* (Linn.), a European bird. – *Novitates Zoologicae* 4: 371-377.
- Rovereto, C. (1914): Los Estratos Araucanos y sus fósiles [The Araucano deposits and their fossils.] – *Anales del Museo de Historia Natural de Mendoza* 11: 157-160. [In Spanish.]
- Rudwick M.J.S. (1972) *The meaning of fossils. Episodes in the history of palaeontology.* – London: Macdonald, iii + 287 pp.
- Rühl F. (1896): Beiträge zur Kenntnis der tertiären und quartären Ablagerungen in Bayerisch-Schwaben, von den Alpen bis zum Jura und der Iller bis zum Ammersee. – *Bericht des Naturwissenschaftlichen Vereines für Schwaben und Neuburg (a. V.)* 32: 327-490.
- Ruse M. (1969): Definitions of species in biology. – *British Journal of the Philosophy of Science* 22: 369-371.
- Rusu A. (1989): Problems of correlation and nomenclature concerning the Oligocene formations in NW Transylvanian Basin. – In: Petrescu I. (ed.): *The Oligocene from Transylvanian Basin*: 67-78. Cluj-Napoca.
- Russell D.E. (1964): Les Mammifères paléocènes d'Europe. – *Mémoires du Muséum National d'Histoire Naturelle de Paris (C)* 13: 1-324.
- Russell D.E., Broin F. de, Galoyer A., Gaudant J., Gingerich P.D. & Rage J.C. (1990): Les vertébrés du Sparnacien de Meudon. – *Bulletin d'Information des Géologues du Bassin de Paris* 27(4): 21-31.
- Rüttimeyer L. (1873): Über die Rennthier-Station von Veyrier am Salève. – *Archiv für Anthropologie* 6: 59-73.
- Sáez M.D. de (1927): Las Aves corredoras fósiles del Santacrucense [Fossil cursorial birds of the Santacrucense.] – *Anales de la Sociedad Científica Argentina* 103: 145-160. [In Spanish.]
- Saint-Périer R. de (1920): La grotte des Harpons à Lespugue (Haute-Garonne). – *Anthropologie (Paris)* 30: 209-234.
- Saint-Périer R. de (1927): La grotte de Gouëris à Lespugue. – *Anthropologie (Paris)* 37: 263-276.
- Salotti M., Bailon S., Bonifay M.-F., Courtois J.-Y., Dubois J.-N., Ferrandini J., Ferrandini M., La Milza L.-C., Mourer-Chauviré C., Popelard J.-B., Quinif Y., Réal-Testud A.-M., Miniconi C., Pereira E. & Persiani C. (1997): Castiglione 3, un nouveau remplissage fossilifère d'âge Pléistocène moyen dans le karst de la région d'Oletta (Haute-Corse). – *Comptes Rendus de l'Académie des Sciences Paris, Sciences de la Terre et des Planètes* 324: 67-74.
- Sánchez Marco A. (1987a): Aves fosiles de Atapuerca. – In: Aguirre E., Carbonell E & Bermudez de Castro J.M. (eds.): *El hombre fosil de Ibeas y el Pleistoceno de la Sierra de Atapuerca* 1: 67-73. Junta de Castilla y Leon: Consejería de Cultura y Bienestar Social.
- Sánchez Marco A. (1987b): Saalian s.l. avian findings in the northern Iberian plateau. – In: Mourer-Chauviré C. (ed.): *L'évolution des oiseaux d'après le témoignage des fossiles* – *Documents des Laboratoires de Géologie Lyon* 99: 193-196.
- Sánchez Marco A. (1989): Huéscar-1 (Granada, España): avance de la lista de aves y consiguientes conjeturas sobre paleoambiente y paleogeografía [Huéscar-1 (Granada,

- Spain): list of birds and related ideas on paleoenvironment and paleogeography.] – In: Alberdi M.T. & Bonadonna F.P. (eds.): Geología y paleontología de la cuenca de Guadix-Baza [Geology and paleontology of the Guadix-Baza cavern.] – Trabajos sobre el Neogeno-Cuaternario 11: 175-182. Madrid. [In Spanish.]
- Sánchez Marco A. (1990a): A new bustard (Otídidæ, Aves) from the early Pliocene of Layna (Soria, Spain). – *Paleontologia i Evolució* 23: 223-229.
- Sánchez Marco A. (1990b): Aves de los yacimientos mesopleistocénicos de Torralba y Ambrona (Soria, España) [Birds from middle Pleistocene sites Torralba and Ambrona (Soria, Spain).] – *Actas de Paleontología* 68: 349-356. [In Spanish.]
- Sánchez Marco A. (1995): Las aves de la unidad estratigráfica de la Galería (sierra de Atapuerca, Burgos, España) [Birds from the stratigraphical unit of Galería (Sierra Mountains, Burgos, Spain).] – In: Bermúdez J.M., Arsuaga J.L. & Carbonell E. (eds.): Evolución humana en Europa y los yacimientos de la sierra de Atapuerca [Human evolution in Europe and the record from the Atapuerca Mountains]: 137-146. Valladolid: Junta de Castilla y León.
- Sánchez Marco A. (1996a): Tertiary avian localities of Portugal. – Mlíkovský J. (ed.): Tertiary avian localities of Europe. – *Acta Universitatis Carolinae, Geologica* 39: 699-701.
- Sánchez Marco A. (1996b): Tertiary avian localities of Spain. – Mlíkovský J. (ed.): Tertiary avian localities of Europe. – *Acta Universitatis Carolinae, Geologica* 39: 719-732.
- Sánchez Marco A. (1996c): The presence of the Waldrapp *Geronticus eremita* (Plataleidae) in the Plio-Pleistocene boundary in Spain. – *Ibis* 138: 560-561.
- Sánchez Marco A. (1996d): Aves fósiles del Pleistoceno ibérico: rasgos climáticos, ecológicos y zoogeográficos [Fossil birds of the Iberian Pleistocene: climatic, ecological, and zoogeographic traits.] – *Ardeola* 43: 207-219. [In Spanish.]
- Sánchez Marco A. (1999a): Implications of the avian fauna for paleoecology in the Early Pleistocene of the Iberian Peninsula. – *Journal of Human Evolution* 37: 375-388.
- Sánchez Marco A. (1999b): Ibis eremita. Restos fósiles hallados en yacimientos españoles [The Waldrapp. Fossil remains found in Spanish deposits.] – *Quercus* 157: 48-49. [In Spanish.]
- Sánchez Marco A. (1999c): Nuevas aves fósiles del yacimiento mesopleistocénico de Ambrona (Soria, España) [New fossil birds from the middle Pleistocene site of Ambrona (Soria, Spain).] – *Trabajos de Prehistoria* 56: 115-118. [In Spanish.]
- Sánchez Marco A. (1999d): Aves del yacimiento mesopleistoceno de Galería (sierra de Atapuerca). Patrones ecológicos en el Pleistoceno medio [Birds from the middle Pleistocene of Galería (Atapuerca mountains). Ecological patterns in the middle Pleistocene.] – In: Carbonell E., Rosas A. & Díez J.C. (eds.): *Arqueología en Castilla y León* [Archeology in Castilla and León] 7: 211-224. Valladolid. [In Catalan.]
- Sánchez Marco A. (1999e): Catálogo paleornitológico del Terciario ibérico y balear [Paleornithological catalogue of the Iberian and Balearic Tertiary.] – *Estudios Geológicos* 55: 115-118. [In Spanish.]
- Sánchez Marco A. (2000a): Objeciones al caso de los fósiles gibraltareños [Reply to the case of Gibraltar fossils.] – *Quercus* 168: 52-53. [In Spanish.]
- Sánchez Marco A. (2000b): Vestigios de una lechuza fósil del Mediterráneo occidental [Remains of a fossil barn owl from the western Mediterranean.] – *Quercus* 170: 52-53. [In Spanish.]
- Sánchez Marco A. (2001a): Strigiformes from the Neogene of Spain. – *Ibis* 143: 313-316.
- Sánchez Marco A. & Sastre Páez I. (2001b): Historia de la paleornitología a través de los documentos científicos [History of paleornithology in Spain according to scientific publications.] – *Paleontología* 16: 99-113. [In Spanish.]
- Sánchez Marco A., Pérez B. & Herrero S. (2000): Passeriformes del Neógeno ibérico:

- hallazgos en Layna (Plioceno inferior) [Passeriformes in the Iberian Neogene: records in Layna (lower Pliocene).] – *Revista Española de Paleontología* 15: 101-104. [In Spanish.]
- Scarlett R.J. (1972): Bone of a presumed odontoptyrygian bird from the Miocene of New Zealand. – *New Zealand Journal of Geology and Geophysics* 15: 269-274.
- Schaub S. (1929a): Über einen neuen Riesenvogel aus den Tereginasanden von Monthelon bei Epernay. – *Verhandlungen der Schweizerischen Naturforschenden Gesellschaft* 110: 195.
- Schaub S. (1929b): Über eocäne Ratitenreste in der Osteologischen Sammlung des Basler Museums. – *Verhandlungen der Naturforschenden Gesellschaft in Basel* 40: 588-598.
- Schaub S. (1940a): Ein Ratitenbecken aus dem Bohnerz von Egerkingen. – *Verhandlungen der Schweizerischen Naturforschenden Gesellschaft* 120: 154-155. [Abstract.]
- Schaub S. (1940b): Ein Ratitenbecken aus dem Bohnerz von Egerkingen. – *Eclogae Geologicae Helveticae* 33: 274-284.
- Schaub S. (1945): Bemerkungen zum Typus von *Taoperdix keltica* Eastman. – *Eclogae Geologicae Helveticae* 38: 616-621.
- Schenker A. (1977): [The former distribution of the Bald Ibis *Geronticus eremita* in Europe.] – *Ornithologischer Beobachter* 74: 13-30. [In German.]
- Schenkling S. (1903): Einige neue Cleriden des Stettiner Museums. – *Entomologische Zeitung (Stettin)* 64: 16-21.
- Scherdlin P. (1926): A propos du Grand Pingouin du Musée zoologique de Strasbourg. – *Bulletin de la Société Philomatique d'Alsace et Lorraine* 7: 10-17.
- Scheuchzer J. J. (1708a): *Piscium querelae et vindiciae* [On entitled and accused fishes.] – Zürich, 36 pp. [In Latin.]
- Scheuchzer J.J. (1708b): *Bildnisse verschiedener Fischen und dero Theilen welche in der Sündfluth zu Grund gegangen.* – Zürich: J.J. Scheuchzer, 8 pp. + 5 pls.
- Scheuchzer J.J. (1716): *Museum diluvianum* [Diluvial museum.]. – Zürich.
- Schilder F.A. (1952): Einführung in die Biotaxonomie (Formerkreislehre): Die Entstehung der Arten durch räumliche Sonderung. – Jena: Gustav Fischer, 162 pp.
- Schlegel H. (1844): *Kritische Uebersicht der europäischen Vögel* – *Révue critique des oiseaux d'Europe.* – Leyden: Arnz, cxxxv + 116 pp.
- Schlosser M. (1900): Die Ausgrabungen im Dürrioch bei Schwaighausen, nordwestlich von Regensburg. – *Correspondenzblatt der Deutschen Gesellschaft für Anthropologie, Ethnologie und Urgeschichte* 31: 41-46.
- Schlosser M. (1916): Neue Funde fossiler Säugetiere in der Eichstätter Gegend. – *Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse* 28 (6): 1-78.
- Schlosser M. (1921): Die Hipparionenfaunen von Veles in Mazedonien. – *Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse* 29: 1-55.
- Schlotheim E.F. von (1820): *Die Petrefaktenkunde auf dem jetzigen Standpunkte.* – Gotha: Becker'sche Buchhandlung, 436 pp.
- Schmidt-Kittler N. (ed., 1987): European references levels and correlation tables. – In: Schmidt-Kittler N. (ed.): *International symposium on mammalian biostratigraphy and paleoecology of the European Paleogene.* – *Münchner Geowissenschaftliche Abhandlungen (A)* 10: 13-31.
- Schönwetter M. (1960): *Handbuch der Oologie. Part 1.* – Berlin: Akademie-Verlag, viii + 64 pp.
- Schweizer W. (1961): Zur Frühgeschichte des Haushuhns in Mitteleuropa. – *Studien an Vor- und Frühgeschichtlichen Tierresten Bayerns* 9.

- Seeley H.G. (1866): Notes on some new genera of fossil birds in the Woodwardian Museum. – *Annals and Magazine of Natural History* (3) 18: 109-110.
- Seeley H.G. (1869): Index to the fossil remains of Aves, Ornithosauria, and Reptilia, from the Secondary System of strata arranged in the Woodwardian Museum of the University of Cambridge. – Cambridge: Cambridge University Press, xxiii + 143 pp.
- Seeley H.G. (1874): On the tibia of *Megalornis*, a large struthious bird from the London Clay. – *Quarterly Journal of the Geological Society of London* 30: 708-710.
- Seemann R. (1941): Geologische und paläofaunistische Untersuchungen am Goldberg im Ries. – *Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg* 96: 49-62.
- Seemann R. & Berckhemer F. (1930): Eine Spaltenfüllung mit reicher aquitanen Wirbeltierfauna im Massenkalk des Weissen Jura bei Tomerdingen (Ulmer Alb). – *Palaeontologische Zeitschrift* 12: 14-25.
- Seguí B. (1999a): Els ocells fossils de Mallorca i de Menorca. Successió estratigràfica d'aus en els reblliments carstics de les Gimnesies [Fossil birds of Mallorca and Menorca. Stratigraphic succession of birds in the karstic deposits of the Gymnesic Islands.] – Unpub. Thesis, Universitat de les Illes Balears, 194 pp. [In Catalan.]
- Seguí B. (1999b): A late Tertiary woodcock from Menorca, Balearic Islands, western Mediterranean. – *Condor* 101: 909-915.
- Seguí B. & Alcover J. A. (1999): Comparison of paleoecological patterns in insular bird faunas: a case study from the western Mediterranean and Hawaii. – In: Olson S.L. (ed.): *Avian paleontology at the close of the 20th century*. – *Smithsonian Contributions to Paleobiology* 89: 67-73.
- Selander R.K. (1971): Systematics and speciation in birds. – In: Farner D.S. & King J.R. (eds.): *Avian biology* 1: 57-147. New York: Academic Press.
- Selinka E. (1891): Vögel. Vol. 1. Osteologie. – In: Bronn H.G. (ed.): *Klassen und Ordnungen des Thier-Reichs*. Vol. 6. – Leipzig: C. F. Winter'sche Verlagshandlung.
- Serebrovskij P.V. (1940a): Novye nachodki ptic iz binagadinskih otloženij [New records of birds from the Binagady deposits.] – *Doklady Akademii Nauk SSSR (novaja serija)* 27: 766-768. [In Russian.]
- Serebrovskij P.V. (1940b): O pticah binagadinskih kirovych plastov [On the birds from the tar flots of Binagady.] – *Izvestija Azerbajdžanskogo Filiala Akademii Nauk SSSR* 3: 71-75. [In Russian.]
- Serebrovskij P.V. (1941a): Ostatki plejstocenovych ptic iz binagadinskih otloženij [Remains of Pleistocene birds from the Binagady deposits.] – *Doklady Akademii Nauk SSSR* 33: 473-475. [In Russian.]
- Serebrovskij P.V. (1941b): Pticy iz pliocenovych otloženij Odessy [Birds from the Pliocene deposits of Odessa.] – *Doklady Akademii Nauk SSSR (novaja serija)* 33: 476-479. [In Russian.]
- Serebrovskij P.V. (1948): Pticy binagadinskih kirovych otloženij [Birds of the tar deposits of Binagady.] – *Trudy Estestvenno-Istoričeskogo Muzeja Imeni Zardabi (Baku)* 1: 27-85. [In Russian.]
- Serjeantson D. (1993): Archaeological and ethnographic evidence for seabirds exploitation in Scotland. – In: Morales Muniz A. (ed.): *Archaeornithology: birds and the archaeological record*. – *Archaeozoology* 2: 209-224.
- Serjeantson D. (2001): The Great Auk and the Gannet: a prehistoric perspective on the extinction of the Great Auk. – *International Journal of Osteoarchaeology* 11: 43-55.
- Serres M. de (1830): Observations sur divers ossements de Mammifères et d'Oiseaux découverts dans les Calcaires calcereux quaternaires des environs de Perpignan. – *Annales des Sciences d'Observation* 3: 229-242.
- Serres M. de (1838a): Essai sur les cavernes à ossements et les causes qui les y ont accumulés. – Paris: J. B. Ballière, 412 pp.

- Serres M. de (1838b): Notes sur les animaux des terrains tertiaires marins supérieurs découvert dans le sol immergé des environs de Montpellier. – Annales des Sciences Naturelles 9: 280-292.
- Serres, M. de, Dubreuil J. & Jeanjean B. (1839): Recherches sur les ossements humatiles des cavernes de Lunel-Viel. – Montpellier: Boehm et Cie, i + 275 pp.
- Sesé C. & López N. 1977: El yacimiento de Escoboza de Catalañazor (Soria) y su aportación a la bioestratigrafía de la Meseta Castellana [The locality Escoboza de Catalañazor (Soria) and its importance for the biostratigraphy of Meseta Castellana.] – Trabajos sobre Neógeno-Cuaternario 7: 35-40. [In Spanish.]
- Sharpe R.B. (1899): A hand-list of the genera and species of birds. Vol. 1. London: British Museum (Natural History), xxi + 303 pp.
- Shaw A.B. (1969): Adam, Eve, paleontology, and the non-objective arts. – Journal of Paleontology 43: 1085-1089.
- Shufeldt R.W. (1896): Fossil bones of birds and mammals from Grotto Pietro Tamponi and Grive-St-Alban. – Proceedings of the Academy of Natural Sciences of Philadelphia 1896: 507-516.
- Shufeldt R.W. (1909): Osteology of birds. – New York State Museum of Natural History, Bulletin 130: 1-381, 26 pls.
- Shufeldt R.W. (1913): Further studies of fossil birds with descriptions of new and extinct species. – Bulletin of the American Museum of Natural History 32: 285-306.
- Shufeldt R.W. (1915): Fossil birds in the Marsh collection of Yale University. – Transactions of the Connecticut Academy of Arts and Sciences 19: 1-109.
- Sibley C.G., Ahlquist J.E. & Monroe jr. B.L. (1988): A classification of the living birds of the world, based on DNA-DNA hybridization. – Auk 105: 409-423.
- Sigé B., Hugué M., Crochet J.-Y., Legendre S., Mourer-Chauviré C., Rage J.-C. & Simon-Coinçon R. (1998) Baraval, nouvelle faune de l'Oligocène inférieur (MP 22) des phosphorites du Quercy. Apport à la signification chronologique des remplissages karstiques. – Bulletin de la Société d'Histoire Naturelle de Toulouse 134: 85-90.
- Simpson G.G. (1943): Criteria for genera, species, and subspecies in zoology and paleontology. – Annals of the New York Academy of Sciences 44: 145-178.
- Simpson G.G. (1944): Tempo and mode in evolution. – New York: Columbia University Press, 237 pp.
- Simpson G.G. (1951): The species concept. – Evolution 5: 285-298.
- Simpson G.G. (1953): The major features of evolution. – New York: Columbia University Press, 434 pp.
- Simpson G.G. (1961): Principles of animal taxonomy. – New York: Columbia University Press, 237 pp.
- Sinclair W.J. (1928): *Omorhamphus*, a new flightless bird from the Lower Eocene of Wyoming. – Proceedings of the American Philosophical Society 67: 51-65.
- Skutil J. & Stehlík A. (1939): Moravská diluviální avifauna [Moravian diluvial avifauna.] – Ornitholog 6: 3-27. [In Czech.]
- Skutil J. & Stehlík A. (1944): Čapkovy příspěvky k poznání předhistorického a raně historického ptactva na Moravě [Čapek's contributions to the knowledge of the prehistoric and early historic avifauna in Moravia.] – Ornitholog 11: 37-41. [In Czech.]
- Smirnov E.S. (1960): Taksonomičeskij analiz roda [Taxonomic analysis of the genus]. – Žurnal Obščej Biologii 21: 89-103. [In Russian.]
- Smith G.A. (1975): Systematics of parrots. – Ibis 117: 18-68.
- Smith K.D. (1970): The Waldrapp *Geronticus eremita* (L.). – Bulletin of the British Ornithologists' Club 92: 18-24.
- Smith T. & Smith R. (1995): Synthèse des données actuelles sur les vertébrés de la transition Paleocène-Eocène de Dormaal (Belgique). – Bulletin de la Société Belge de Géologie 104: 119-131.

- Soergel E. (1955): Über einige Vogelreste (Seeadler, Kraniche) aus dem Neolithikum von Ehrenstein bei Ulm. – Jahreshefte des Vereins für Vaterländische Naturkunde von Württemberg 110: 121-124.
- Soergel E. (1961): [In:] Hohen Vielchen. Ein mittelsteinzeitlicher Wohnplatz in Mecklenburg. – Deutsche Akademie der Wissenschaften, Schriften der Sektion für Vor- und Frühgeschichte 10: 64-69. [In German.]
- Solounias N. (1981): The Samos fauna. – Contributions to Vertebrate Paleontology 6: 1-232.
- Sondaar P.Y., McMinn M., Seguí B. & Alcover J.A. (1995): Interès paleontològic dels jaciments càrstic de les Gimnèsies i les Pitiüses [Paleontological finds from the karstic deposits of the Gymnesiens and Pityuses.] – Endins 20: 155-170. [In Catalan.]
- Soulé M.E. (1982): Allometric variation. 1. The theory and some consequences. – American Naturalist 120: 751-764.
- Spasov N. & Iliev N. (1995): Animal remains from the submerged late Eneolithic – early Bronze Age settlements in Sozopol (south Bulgarian Black Sea coast). – In: Proceedings of the International Symposium on Tracia Pontica 6: 287-313. Sozopol.
- Speck J. (1945): Fahrtenfunde aus dem subalpinen Burdigalien und ihre Bedeutung für Fazies und Paläogeographie der oberen Meeresmolasse. – Eclogae Geologicae Helvetiae 38: 411-416.
- Speck J. (1952): Erdgeschichtliches Wesen. – In: Das Buch vom Lande Zug. – Zuger Neujahrsblatt 1952: 17-24.
- Špinar Z. (1980): Paleontologie [Paleontology.] – Praha: SNTL, 361 pp. [In Czech.]
- Spulski B. (1910): *Odontopteryx longirostris* n. sp. – Zeitschrift der Deutschen Geologischen Gesellschaft 62: 507-521.
- Stanley S.M. (1978): Chronospecies' longevities, the origin of genera and the punctuational model of evolution. – Paleobiology 4: 26-40.
- Stadman D.W. (1981): [Review of Harrison & Walker 1976a, 1977a.] – Auk 98: 205-207. [In English.]
- Steenstrup J.J.S. (1855): Et bidrag til geirfuglens, *Alca impennis* Lin., naturhistorie, or saerligt til kundskaben om dens tidligere udbredningskreds. – Videnskabelige Meddelelser fra den Naturhistoriske Forening i Kjøbenhavn 3-7: 33-118. [In Danish.]
- Stehlin H.G. (1923): Die oberpliozäne Fauna von Senèze (Haute-Loire). – Eclogae Geologicae Helvetiae 18: 268-281.
- Steininger F.F. (1999): Chronostratigraphy, geochronology and biochronology of the Miocene “European Land Mammal Mega-Zones (ELMMZ) and the Miocene “Mammal-Zones”. – In: Rössner G.E. & Heissig K. (eds): The Miocene land mammals of Europe: 9-24. München: Dr. F. Pfeil.
- Steininger F.F. & Rögl F. (1984): Paleogeography and palinspastic reconstruction of the Neogene of the Mediterranean and Paratethys. – In: Dixon J.E. & Robertson A.H.F. (eds.): The geological evolution of the eastern Mediterranean: 659-668. Oxford: Blackwell Scientific Publications.
- Steininger F., Rögl F. & Deremitzakis M. (1987): Report on the round table discussion “Mediterranean and Paratethys Correlations”. – In: Déak M. (ed.): VIIIth congress of the regional committee on Mediterranean Neogene stratigraphy. – Annales Instituti Geologici Publici Hungarici 70: 397-421.
- Steininger F.F., Berggren W.A., Kent D.V., Bernor R.L., Sen S. & Agusti J. (1996): Circum-Mediterranean Neogene (Miocene and Pliocene) marine-continental chronologic correlations of European Mammal Units. – In: Bernor R.L., Fahlbusch V. & Mittmann H.-W. (eds.): The evolution of western Eurasian Neogene mammal faunas: 7-46. New York: Columbia University Press.
- Stejneger L. (1882): Outlines of a monograph of the Cygnidae. – Proceedings of the United States National Museum 5: 174-221.

- Stejneger L. (1885): [Articles on individual families.] – In: Kingsley J.S. (ed.): The standard natural history. Vol. 4. Birds. – Boston: S.E. Cassino, vii + 558 pp. [In English.]
- Stemvers-van-Bemmel J. (1984): Een Oligocene fauna in het Bekken van Apt-Manosque-Forcalquier: Zuidfranse vogels en vissen begraven in kalkslib [An Oligocene fauna in the Apt-Manosque-Forcalquier basin: south French birds and fishes buried in limestone.] – *Gea* 17: 134-138. [In Dutch.]
- Stent G.S. (1986): Glass bead game. – *Biology and Philosophy* 1: 227-247.
- Stepanjan L.S. (1970): Konceptija nadvida (superspecies) i predeli ee prilozhenija [The superspecies concept and the limits of its application.] – *Zoologičeskij Žurnal* 49: 653-659. [In Russian.]
- Stepanjan L.S. (1983) Nadvidy i vidy-dvojniki v avifaune SSSR [Superspecies and sibling species in the avifauna of the USSR]. – Moskva: Nauka, 295 pp. [In Russian.]
- Stevanović P.M. (1959): Fosilna jaja u miocenu Brajkovca kod Ljiga i bliži stratigrafski položaj nalazišta [A fossil egg in the Miocene of Brajkovac at Ljig and closer stratigraphical position of the locality.] – *Geološki Anali Balkaskoga Poluostrva* 26: 153-162 + pl. I-II. [In Serbo-Croatian.]
- Stewart J.R. (1996): Quaternary birds from the Torbryan Valley. – In: Charman D.J., Newnham R.M. & Croot D.G. (eds.): Devon and East Cornwall field guide: 190-191. – Quaternary Research Association.
- Stewart J. R. (1997): *Pinguinus impennis*, extinct great auk. – In: Roberts M.B., Parfitt S.A, Pope M.I. & Wenban-Smith F.F. (eds.): Boxgrove, West Sussex: rescue excavations of a Lower Palaeolithic landsurface (Boxgrove Project B, 1989-91). – *Proceedings of the Prehistoric Society* 63: 303-358.
- Stewart J.R. (1999a): Intraspecific variation in modern and Quaternary European *Lagopus*. – In: Olson S.L. (ed.): Avian paleontology at the close of the 20th century. – *Smithsonian Contributions to Paleobiology* 89: 159-168.
- Stewart J. R. (1999b): The evolution of Quaternary birds in the western Palaearctic: aspects of taxonomy and ecomorphology. – Unpub. Thesis, University College, London.
- Stingelin W. (1958a): Vergleichend-morphologische Untersuchungen am Vorderhirn der Vögel auf cytologischer und cytoarchitektonischer Grundlage. – Basel: Helbig & Lichtenhahn, 123 pp.
- Stingelin W. (1958b): Differenzierung und Gestalt des Vorderhirns der Vögel. – *Revue Suisse de Zoologie* 65: 427-434.
- Stingelin W. (1960): Vorderhirn und Anpassungstypus. – In: Bergman G., Donner K.O. & Haartman L. von (eds.): Proceedings of the 12th international ornithological congress: 681-693. Helsinki: Tilgmannin Kirjapaino.
- Storer R.W. (1956): The fossil loon, *Colymboides minutus*. – *Condor* 58: 413-426.
- Storer R.W. (1971): Classification of birds. – In: Farner D.S. & King J.R. (eds.): Avian biology. Vol. 1: 1-18. New York: Academic Press.
- Stresemann E. (1951): Die Entwicklung der Ornithologie von Aristoteles bis zur Gegenwart. – Aachen: Verlag Hans Limberg, xv + 431 pp.
- Strohl J. (1917): Conrad Gesners “Waldrapp”. – *Vierteljahrsschrift der Naturforschenden Gesellschaft Zürich* 62: 501-538.
- Stromer E. von (1928): Wirbeltiere im obermiozänen Flinz Münchens. – *Abhandlungen der Bayerischen Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Abteilung* 32(1): 1-71.
- Storer R.W. (2000): The systematic position of the Miocene grebe, *Thiornis sociata* Navas. – *Annales de Paléontologie* 86: 129-139.
- Stuart A.J. (1974): Pleistocene history of the British vertebrate fauna. – *Biological Reviews* 49: 225-266. Stuart A.J., Wolff R.G., Lister A.M., Singer R. & Egginton J.M.

- (1993): Fossil vertebrates. – In: Singer R., Gladfelter B.G. & Wymer J.J. (eds.): The Lower Paleolithic site at Hoxne, England: 163-206. Chicago.
- Studer B. (1840): [Untitled communication.] – Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde 1840: 211. [In German.]
- Studiati C. (1857): Description des fossiles de la brèche osseuse de Monreale de Bonaria près de Cagliari. – In: Marmora A. de la (ed.): Voyage en Sardaigne III: Suite de la description géologique 2: 651-704. Turin [= Torino]: Les frères Bocca, & Paris: Arthus Bertrand.
- Stüssi F. (1958): Die Entzerrung von Fossilien am Beispiel des *Protornis glaronensis*. – Schweizerische Paläontologische Abhandlungen 73(5): 1-16.
- Sudhaus W. (1995): Analyse der eiszeitbedingten Entstehung von Zwillingarten in der europäischen Vogelfauna. – Sitzungsberichte der Gesellschaft Naturforschender Freunde zur Berlin (Neue Folge) 33: 97-128.
- Sukačev V.N., Gromov V.I. & Bader O.N. (1966): Verchnepaleolitičeskaja stojanka Sungir [The late Paleolithic site Sungir.] – Geologičeskij Institut Akademii Nauk SSSR, Trudy 162: 1-140. [In Russian.]
- Suriano F. (1980): Fossil birds of Simonelli Cave. – Quaderni delle Academia Nazionale dei Lincei 249: 123-126.
- Švec P. (1980): Lower Miocene birds from Dolnice (Cheb basin), western Bohemia. – Časopis pro Mineralogii a Geologii 25: 377-387.
- Švec P. (1981): Lower Miocene birds from Dolnice (Cheb basin), western Bohemia. Part II. – Časopis pro Mineralogii a Geologii 26: 45-56.
- Švec P. (1982): Two new species of diving birds from the Lower Miocene of Czechoslovakia. – Časopis pro Mineralogii a Geologii 27: 243-260.
- Švec P. (1983): Lower Miocene rail from western Bohemia (Ralliformes, Aves). – Věstník Ústředního Ústavu Geologického 58: 31-38.
- Švec P. (1984): Further finds of grebe *Miobaptus walteri* in the Miocene of Bohemia. – Časopis pro Mineralogii a Geologii 29: 167-170.
- Švec P. (1985): New finds of the small fossil loon, *Colymboides minutus* Milne-Edwards, 1867, from the Lower Miocene of western Bohemia. – Acta Universitatis Carolinae, Geologica 4: 447-455.
- Švec P. (1986): The fossil pheasant (Aves, Phasianidae) from the upper Miocene of Děvinská Nová Ves (Slovakia). – Časopis pro Mineralogii a Geologii 31: 83-88.
- Švec P. & Mlíkovský J. (1986): First Tertiary record of the genus *Oxyura* (Aves: Anseridae). – Časopis pro Mineralogii a Geologii 31: 403-407.
- Sylvester-Bradley P.C. (ed., 1956): The species concept in paleontology. – London: Systematists' Association, 145 pp.
- Tatarinov K.A. & Bačinskij G.A. (1968): Peščernye zachoroneniya pliocenovych i antropogenovych pozvonočnych v zapadnykh oblastjach Ukrainy [Cave deposits of Pliocene and Anthropogenic vertebrates in the western parts of Ukraine.] – Bjulleten' Moskovskogo Obščestva Ispytatelej Prirody 13: 114-122. [In Russian.]
- Tchernov E. (1962): Palaeolithic avifauna in Palestine. – Bulletin of the Research Council of Israel (B) 11: 95-131.
- Tchernov E. (1980): The Pleistocene birds of 'Ubeidyia, Jordan Valley. – Jerusalem: The Israel Academy of Sciences and Humanities, 95 pp.
- Temminck C.J. (1838-1843): Histoire naturelle générale des Pigeons. Vol. 2. – Paris, 48 pls.
- Terzea E. & Jurcsák T. (1968): Bemerkungen über die mittelpleistozänen Faunen von Betfia. – Berichte der Deutschen Gesellschaft für Geologische Wissenschaften (A) 13: 381-390.
- Thenius E. (1954): Vösendorf - ein Lebensbild aus dem Pannon des Wiener Beckens. II.

- Systematischer Teil. D. Wirbeltiere. d. Aves. – Mitteilungen der Geologischen Gesellschaft in Wien 46: 49-50.
- Thenius E. (1959): Tertiär. 2. Wirbeltierfaunen. – Stuttgart: Ferdinand Enke, 328 pp.
- Thesing R. (1977): Die Größenentwicklung des Haushuhns in vor- und frühgeschichtlicher Zeit. – Unpub. Diss., Universität München, 62 pp.
- Tintant H. (1980): Problématique de l'espèce en paléozoologie. – In: Bocquet C., Générmont J. & Lamotte M. (eds.): Les problèmes de l'espèce dans le règne animal 3: 321-372. Paris: Société Zoologique de France.
- Tintant H. (1984): L'évolution du concept de genre: de la similitude à la parenté. – Bulletin de la Société Géologique de France 26: 573-582.
- Tkačenko O.F., Pišvanova L.S. & Švareva N.Ja. (1967): Nachodki sledov ptic i chiščnika v slojach "Guka" na r. Rybnice v Predkarpat'e [Records of footprints of birds and a mammalian predator in the deposits of "Guk" at Rybnica river in Predkarpat'e.] – Bjuulletin Moskovskogo Obščestva Ispytatelej Prirody, Otdel Geologii 42(4): 125-128. [In Russian.]
- Tomek T. & Bocheński Z.M. (2000): The comparative osteology of European corvids (Aves: Corvidae), with a key to the identification of their skeletal elements. – Kraków: Institute of Systematics and Evolution of Animals, 102 pp.
- Tonni E.P. (1980): Un pseudornitido (Pelecaniformes, Odontopterygia) de Gran Tamaño, del Terciario temprano de Antartida [A large pseudodontornithid (Pelecaniformes, Odontopterygia) from the early Tertiary of Antarctica.]. – Ameghiniana 17: 273-276. [In Spanish.]
- Tonni E.P. & Tambussi E.P. (1985): Nuevos restos de Odontopterygia (Aves: Pelecaniformes) del Terciario temprano de Antartida [New remains of Odontopterygia (Aves: Pelecaniformes) from the early Tertiary of Antarctica.]. – Ameghiniana 21: 121-124. [In Spanish.]
- Tordoff H.B. & Macdonald J.R. (1957): A new bird (family Cracidae) from the early Oligocene of South Dakota. – Auk 74: 174-184.
- Toula F. (1910): Diluviale Säugetiere vom Gesprengsberg, Kronstadt in Siebenbürgen. – Jahrbuch der K. K. Österreichischen Geologischen Reichsanstalt (Wien) 49: 575-614.
- Touraine F. (1960): Oeufs d'oiseaux de très grande taille dans l'Eocène inférieur de Provence. – Bulletin de la Société Géologique de France (7) 2: 783-789.
- Touraine F. (1961): Les oeufs d'oiseaux éocènes et la stratigraphie du Cengle (Bouches-du-Rhône). – Comptes Rendus de la Société Géologique de France 1961: 90-91.
- Touraine F. (1969): A propos de la note de MM. Dughi, Plaziat et Sirugue traitant de la position stratigraphique des oeufs d'oiseaux découverts dans l'Aude. – Comptes Rendus de la Société Géologique de France 1969: 146.
- Tret'jakov D. K. (1941): Tretičnaja fauna odesskich katakomb [Tertiary fauna of the Odessa catacombs.] – Sovetskaja Nauka 1941 (1): 92-106. [In Russian.]
- Tugarinov A.Ja. (1937): Pticy Kryma vremeni Vjurmского oledenija [Birds of Crimea in the period of the Würmian glaciation.] – Trudy Sovetskoj Sekcii Meždunarodnoj Asociacii po Izučeniju Četvertičnogo Perioda 1: 97-114. [In Russian.]
- Tugarinov A.Ja. (1940a): Novye dannye dlja tretičnoj ornitofauny SSSR [New data on the Tertiary avifauna of the USSR.] – Doklady Akademii Nauk SSSR 26: 205-208, 314-316. [In Russian.]
- Tugarinov A.Ja. (1940b): Novye nachodki pliocenovej ornitofauny Odessy [New records of the Pliocene avifauna of Odessa.] – Doklady Akademii Nauk SSSR 26: 311-313. [In Russian.]
- Tyrberg T. (1985): [*Grus primigenia* – the extinct large crane of Europe.] – Fauna och Flora 80: 169-176. [In Swedish.]
- Tyrberg T. (1991a): Crossbill (Genus *Loxia*) evolution in the West Palearctic – a look at the fossil evidence. – Ornis Svecica 1: 3-10.

- Tyrberg T. (1991b): Arctic, montane and steppe birds as glacial relicts in the West Palearctic. – Ornithologische Verhandlungen 25: 29-49.
- Tyrberg T. (1995): Palaeobiogeography of the genus *Lagopus* in the West Palearctic. – In: Peters D.S. (ed.): Acta palaeornithologica. – Courier Forschungsinstitut Senckenberg 181: 275-291.
- Tyrberg T. (1998): Pleistocene birds of the Palearctic: a catalogue. – Cambridge, Mass.: Nuttall Ornithological Club, ix + 720 pp. [= Publications of the Nuttall Ornithological Club, No. 27.]
- Tyrberg T. (1999): Seabirds and late Pleistocene marine environments in the northeast Atlantic and the Mediterranean. – In: Olson S.L. (ed.): Avian paleontology at the close of the 20th century. – Smithsonian Contributions to Paleobiology 89: 139-157.
- Tyrberg T. (2001): Pleistocene birds of the Palearctic. – <http://w1.115.telia.com/~u11502098/pleistocene.html> [Stand on 7 November 2001.]
- Tyrberg T. & Erickson P.G.P. (1996): Tertiary avian localities of Sweden. – Mlíkovský J. (ed.): Tertiary avian localities of Europe. – Acta Universitatis Carolinae, Geologica 39: 733-734.
- Uhlmann E. (1923): Entwicklungsgedanke und Artbegriff. – Jenaische Zeitschrift für Naturwissenschaften 59: 1-116.
- Umans'ka A.S. (1972): Domašnie pticy iz archeologičeskich pamjatnikov Ukrainy [Domestic birds from the archeological monuments of Ukraine.] – Prirodnaja Obstanovka i Fauny Prošlogo 6: 71-95. [In Russian.]
- Umans'ka A.S. (1973): Novaja nachodka ostatkov urmiornisa iz neogenovych otloženij Ukrainy [New find of Urmiornis remains from the Neogene deposits of Ukraine.] – Vestnik Zoologii (Kyjiv) 9(4): 30-33. [In Russian.]
- Umans'ka A.S. (1979a): Miocenovyte pticy Zapadnogo Pričernomor'ja USSR [Miocene birds of western Pričernomor'e in the Ukrainian SSR.] – Vestnik Zoologii 15(4): 40-45. [In Russian.]
- Umans'ka A.S. (1979b): Novyj vid filina (*Bubo longaevus*) iz pozdneogenovych otloženij severnogo Pričernomor'ja USSR [New species of a horned owl (*Bubo longaevus*) from the late Neogene deposits of northern Pričernomor'e in the Ukrainian SSR.] – Dopovidi Akademiji Nauk Ukrajin's'koji RSR (B) 1979: 779-782. [In Russian.]
- Umans'ka A.S. (1981): Miocenovyte pticy Zapadnogo Pričernomor'ja USSR. Soobščenie II. [Miocene birds of western Pričernomor'e. Communication II.] – Vestnik Zoologii 17(3): 17-21. [In Russian.]
- Unwin D. (1993): Aves. – In: Benton M.J. (ed.): The fossil record 2: 717-737. – London: Chapman & Hall.
- Ussher R.J. (1897a): The discovery of bones of the Great Auk in county Waterford. – Irish Naturalist 6: 208.
- Ussher R.J. (1899a): The Great Auk, once an Irish Bird. – Irish Naturalist 8: 1-3.
- Ussher R.J. (1899b): On remains of *Plautus impennis* from Irish Kitchen middens. – Bulletin of the British Ornithologists' Club 8: 50.
- Ussher R.J. (1902): Great Auk (*Alca impennis*) in Co. Clare. – Irish Naturalist 11: 188.
- Valenciennes A. (1855): Remarques à l'occasion d'une communication de M. Lartet sur le tibia d'oiseau fossile trouvé à Meudon. – Comptes Rendus des Séances Hebdomadaires de l'Académie des Sciences (Paris) 40: 583-584.
- Van Valen L.M. (1988): Species, sets, and the derivative nature of philosophy. – Biology and Philosophy 3: 49-66.
- Verheyen R. (1956): Note sur l'anatomie et la place des kamiches (Anhimiformes) dans le systèmes de classification. – Gerfaut 46: 215-221.
- Verheyen R. (1960): A new classification for the non-passerine birds of the world. – Bulletin de l'Institut Royal des Sciences Naturelles de Belgique 37(27): 1-36.

- Vidgal'm J. (1886): Die fossilen Vogelknochen der Odessaner Steppe-Kalksteinbrüche an der Neuen Slobodka bei Odessa. – Zapiski Novorossijskogo Obščestva Estestvoispytatelej 10, Supplement: 1-10.
- Vilette P. (1994): La paléoavifaune du Pléistocène moyen de la grotte du Lazaret (Nice, Alpes-Maritimes). – Bulletin Mus. Anthropol. Préhist. Monaco 36: 15-29.
- Vilette P. (1995): Les Oiseaux. – In: Defleut A. & Crégut-Bonnoure E. (eds.): Le gisement paléolithique de Pléistocène moyen de la grotte des Cèdres (Le Plan-d'Aups, Var). – Docum. Archéol. Française 49: 154-155.
- Villalta J.F. de (1963): Las aves fósiles del Mioceno español [Fossil birds from the Spanish Miocene.] – Boletín de la Real Sociedad Española de Historia Natural, Geología 61: 263-285. [In Spanish.]
- Villalta J.F. de (1965): Un yacimiento villafranchiense en las Islas Medas (prov. de Gerona) [An Villafranchian locality on the Medas islands.] – In: Actes IV Congrès International d'Etudes Pyrénéennes: 129-131. [In Spanish.]
- Villalta J.F. de & Crusafont Pairó M. (1950): Sobre algunas aves fósiles de Cataluña [On some fossil birds of Catalonia.] – Notas y Comunicaciones del Instituto Geológico y Minero de España 20: 143-156. [In Spanish.]
- Villatte J. (1966): Découverte de fragments de coquille d'oeufs d'Oiseaux dans l'Eocène inférieur de l'Aude. – Comptes Rendus Sommaires de la Société Géologique de France 1966: 345-346.
- Vinokurov A.A. (1992): Redkie i isčezajuščie životnye: pticy [Rare and disappearing animals: birds.] – Moskva: Vysšaja škola, 448 pp. [In Russian.]
- Violani C. (1974): Ecologia di un'estinzione: l'Alca impenne [The ecology of an extinction: *Alca impennis*.] – Boll. Mus. Civ. Venezia 25: 49-60. [In Italian.]
- Viret J. (1938): Sur l'âge des argiles ligniteuses de Nassiet près Amou (Landes). – Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris) 207: 500-501.
- Viret J. (1954): Le loess à bancs de Saint-Vallier (Drôme) et sa faune de mammifères villafranchiens. – Nouvelles Archives du Muséum d'Histoire Naturelle de Lyon 4: 1-200.
- Vjalov O.S. (1951): Schema delenija miocena Prekarpat'ja [A stratigraphical scheme of the Miocene of Predkarpat'e.] – Doklady Akademii Nauk SSSR 135 (5).
- Vjalov O.S. (1960): Novyj iskopaemyj sled pticy v miocene Predkarpat'ja [New excavated footprint of a bird in the Miocene of Predkarpat'e.] – Doklady Akademii Nauk SSSR 135: 1237-1239. [In Russian.]
- Vjalov O.S. (1965): Stratigrafija neogenovych molass predkarpatskogo progiba [Stratigraphy of Neogene molasses of the cis-carpathian step fold.] – Kyjiv: Naukova Dumka, 212 pp. [In Russian.]
- Vjalov O.S. (1966): Sledy žiznedejatel'nosti organizmov i ich paleontologičeskoe značenie [Remains of organismal life activities and their paleontological significance.] – Kyjiv: Naukova Dumka, 220 pp. [In Russian.]
- Vjalov O.S. (1971): Iskopaemye ostatki svidetel'stev razmnoženija i roždenija [Fossil remains of records of reproduction and birth.] – Paleontologičeskij Sbornik (L'viv) 8(2): 42-52. [In Russian.]
- Vjalov O.S. (1989): O sledach ptic iz pliocena Irana, pripisyvaemych rodu *Urmionis*. [On the footprints of birds from the Pliocene of Iran, generally attributed to the genus *Urmionis*.] – Paleontologičeskij Žurnal 1989 (4): 120-121.
- Vjalov, O.S. & Flerov K.K. (1952): Iskopaemye sledy pozvonočnych v tercičnych otloženijach Predkarpat'ja [Fossil footprints of vertebrates in the Tertiary deposits of Predkarpat'je.] – Bžulletin Moskovskogo Obščestva Ispytatelej Prirody, Otdel Geologii 27(5): 80-88.
- Vojinstvens'kyj M.A. (1959a): Novi dani pro systematyčne položennja vykopnoho ptacha

- Gryzaja odessana* Sub. z pliocenovych vidkladiv Odesy [New data on the systematic position of the fossil bird *Gryzaja odessana* Sub. from the Pliocene deposits of Odesa.] – Dopovidi Akademiji Nauk Ukrajin's'koi RSR 1959: 198-202. [In Ukrainian.]
- Vojinstvens'kyj M.A. (1959b): Novye dannye o predkach domašnej kuricy [New data on the ancestry of domestic chicken.] – In: II vsesojuznaja ornitologičeskaja konferencija, Tezisi dokladov [2nd all-union ornithological conference, Abstracts]: 24-25. Moskva: Izdatel'stvo Moskovskogo gosudarstvennogo universiteta. [In Russian.]
- Vojinstvens'kyj M.A. (1963): Iskopaemaja ornitofauna Kryma [The fossil avifauna of Crimea.] – Trudy Kompleksnoj Karstovoj Ėkspedicii Akademii Nauk USSR 1: 106-123. [In Russian.]
- Vojinstvens'kyj M.A. (1967): Iskopaemaja ornitofauna Ukrainy [Fossil avifauna of Ukraine.] – Prirodnaja Obstanovka i Fauny Prošlogo 3: 3-76. [In Russian.]
- Vojinstvens'kyj M.A. & Umans'ka A.S. (1959): Ptachy z sučasnych aljuvial'nych vikladiv nyžn'ogo Dnipra [Birds from the modern alluvial deposits of Dnipr.] – Dopovidi Ukrajin's'koi RSR 1959: 326-330. [In Ukrainian.]
- Voznesens'kyj O.N. (1937): Umovy uložennja meotyčnoji fauny i flory v s. Hrebenyky Tiraspol'skoho rajonu URSS [On a locality of the Meotian fauna and flora in Hrebenyky village in the Tiraspol Province of the Ukrainian SSR.] – Heolohičnyj Žurnal 4 (1). [In Ukrainian.]
- Voznesens'kyj O.N. (1939) Umovy uložennja meotyčnoji fauny chrebetnyh v s. Nova Emetovka Odes'koho rajonu URSS [On a locality of the Meotian vertebrate fauna in Nova Emetovka village in Odesa Province of the Ukrainian SSR.] – Heolohičnyj Žurnal 6. [In Ukrainian.]
- Vuilleumier F. (1976): La notion d'espèce en ornithologie. – *Société Zoologique de France* 38: 29-65.
- Waga M. (1858): Description d'une nouvelle espèce européenne de Crustacés *Philoscia notata* et d'un nouveau Myriapode *Gervaisia costata*. – *Annales de la Société Entomologique de France* (3)5: 829.
- Wagner R. (1828): Ueber die Knochen-Brekzie in Sardinien und die darin gefundenen Thiere, so wie über einige andere hierher gehörige Erscheinungen. – *Archiv für Gesamte Naturgeschichte* 15: 10-31.
- Wagner R. (1832): Ueber die fossilen Insectenfresser, Nager und Vögel der Diluvialzeit, mit besonderer Berücksichtigung der Knochenbrekzien an den Mittelmeerküsten. – *Abhandlungen der Mathematisch-Physicalischen Classe der Bayerischen Akademie* 1: 753-786.
- Wagner R. (1833): [Untitled communication.] – *Neues Jahrbuch für Mineralogie, Geognosie, Geologie und Petrefaktenkunde* 1833: 324. [In German.]
- Wahlert G. von (1965): The role of ecological factors in the origin of higher levels of organization. – *Systematic Zoology* 14: 288-300.
- Wahlert G. von (1973) Phylogenie als ökologischer Prozess. – *Naturwissenschaftliche Rundschau* 26: 247-254.
- Walker C.A. (1980): The distribution of birds in the British Paleogene. – *Tertiary Research* 3: 25-30.
- Walker C.A., Wragg G.M. & Harrison C.J.O. (1990): A new shearwater from the Pleistocene of the Canary Islands and its bearing on the evolution of certain *Puffinus* shearwaters. – *Historical Biology* 3: 203-224.
- Warheit K.I. (1992): The role of morphometrics and cladistics in the taxonomy of fossils: a paleornithological example. – *Systematic Biology* 41: 345-369.
- Warter S.L. (1976): A new Osprey from the Miocene of California (Falconiformes: Pandionidae). – In: Olson S.L. (ed.), *Collected papers in avian paleontology honoring the 90th birthday of Alexander Wetmore*. – *Smithsonian Contributions to Paleobiology* 27: 133-139.

- Weesie P.D.M. (1982): A Pleistocene endemic island form within the genus *Athene*: *Athene cretensis* n. sp. (Aves, Strigiformes) from Crete. – Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen (B) 85: 323-336.
- Weesie P.D.M. (1987a): Preliminary report on the Pleistocene birds from Crete. – In: Mourer-Chauviré C. (ed.): L'évolution des oiseaux d'après le témoignage des fossiles. – Documents des Laboratoires de Géologie Lyon 99: 197-200.
- Weesie P.D.M. (1987b): The Quaternary avifauna of Crete, Greece. – Unpub. PhD. Thesis, State University of Utrecht, Utrecht, x + 90 pp. + 9 pls.
- Weesie P.D.M. (1988): The Quaternary avifauna of Crete, Greece. – *Palaeovertebrata* 18: 1-94. [This is the published version of Weesie 1987b.]
- Weesie P.D.M. (1999): Les oiseaux du Pleistocène supérieur de la grotte de Corbeddu, Sardaigne. – In: Reumer J.W.F. & De Vos J. (eds): Elephants have a snorkel! Papers in honour of Paul Y. Sondaar. – *Deinsea* 7: 401-409.
- Weidmann M. & Reichel M. (1979): Traces de pattes d'oiseaux dans la Molasse suisse. – *Eclogae Geologicae Helvetiae* 72: 953-971.
- Weigel P.H. (1988): Great Auk remains from a Florida shell midden. – *Auk* 75: 215-216.
- Weigelt J. (1933): Die Biostratonomie der 1932 auf der Grube Cecilie im mittleren Geiseltal ausgegrabenen Leichenfelder. – *Nova Acta Leopoldina (Neue Folge)* 1: 157-174.
- Weigelt J. (1939): Die Aufdeckung der bisher ältesten tertiären Säugetierfauna Deutschlands. – *Nova Acta Leopoldina* 7: 515-528.
- Weigelt J. (1942): Die alttertiären Säugetiere Mitteldeutschlands nach den Hallenser Grabungen im Geiseltal und bei Walbeck. – *Preussische Akademie der Wissenschaften, Vorträge und Schriften* 12: 1-48.
- Weithofer K.A. (1888): Alcune osservazione sulla fauna delle ligniti di Casteani e di Montebamboli (Toscana) [Some observations on the fauna of the lignites of Casteani and of Montebamboli (Toscana).] – *Bolletino del Comitato Geologico Italiano* 19: 363-368. [In Italian.]
- West B. & Zhou B.-X. (1988): Did chicken go north? New evidence for domestication. – *Journal of Archaeological Science* 15: 515-533. [A slightly modified version was published in the *World's Poultry Science Journal* 45: 205-218, 1989.]
- Westphal F. (1963): Ein fossilführendes Jungtertiär-Profil aus dem Randecker Maar (Schwäbische Alb). – *Jahresberichte und Mitteilungen des Oberrheinischen Geologischen Vereins (neue Folge)* 45: 27-43.
- Wetmore A. (1927): Fossil birds from the Oligocene of Colorado. – *Proceedings of the Colorado Museum of Natural History* 7: 1-13.
- Wetmore A. (1928): The systematic position of the fossil bird *Cyphornis magnus*. – *Canada Department of Mines, Geological Survey Bulletin* 49: 1-4.
- Wetmore A. (1930a): Fossil bird remains from the Temblor Formation near Bakerfield, California. – *Proceedings of the California Academy of Sciences* (4) 19(8): 85-93.
- Wetmore A. (1930b): A systematic classification for the birds of the world. – *Proceedings of the United States National Museum* 76(24): 1-8.
- Wetmore A. (1933): Bird remains from the Oligocene deposits of Torrington, Wyoming. – *Bulletin of the Museum of Comparative Zoology* 75: 297-311.
- Wetmore A. (1934a): Fossil birds from Mongolia and China. – *American Museum Novitates* 711: 1-16.
- Wetmore A. (1934b): A systematic classification for the birds of the world, revised and amended. – *Smithsonian Miscellaneous Collections* 89(13): 1-11.
- Wetmore A. (1938): Another fossil owl from the Eocene of Wyoming. – *Proceedings of the United States National Museum* 85: 27-29.
- Wetmore A. (1940a): Fossil bird remains from Tertiary deposits in the United States. – *Journal of Morphology* 66: 25-37.

- Wetmore A. (1940b): A systematic classification for the birds of the world. – Smithsonian Miscellaneous Collections 99(7): 1-11.
- Wetmore A. (1943): Fossil birds from the Tertiary deposits of Florida. – Proceedings of the New England Zoological Club 22: 59-68.
- Wetmore A. (1951): A revised classification for the birds of the world. – Smithsonian Miscellaneous Collections 117(4): 1-22.
- Wetmore A. (1956): Miscellaneous notes on fossil birds. – Smithsonian Miscellaneous Collections 138(8): 1-11.
- Wetmore A. (1959): Birds of the Pleistocene in North America. – Smithsonian Miscellaneous Collections 138(4): 1-24.
- Wetmore A. (1960): A classification for the birds of the world. – Smithsonian Miscellaneous Collections 139(11): 1-37.
- Wijngaarden-Bakker L.H. van (1978): A subfossil Great Auk – *Pinguinus impennis* (L.) from the Netherlands. – Ardea 66: 57-61.
- Wiley E.O. (1978): The evolutionary species concept reconsidered. – Systematic Zoology 27: 17-26.
- Wiley E.O. (1981): Phylogenetics. The theory and practice of phylogenetic systematics. – New York: Wiley.
- Wilkinson H.E. (1969): Description of an Upper Miocene albatross from Beaumaris, Victoria, Australia, and a review of fossil Diomedidae. – Memoirs of the National Museum of Victoria 29: 41-51.
- Willmann R. (1985): Die Art im Raum und Zeit. – Hamburg: Paul Parey.
- Wilson L.G. (1970): Sir Charles Lyell's scientific journals on the species question. – New Haven: Yale University Press, lxi + 572 pp.
- Winge H. (1903): Om jordfundne fugle fra Danmark [On the fossil birds from Denmark.] – Videnskabelige Meddelelser fra den Naturhistoriske Forening i Kjøbenhavn 6: 61-109. [In Danish.]
- Witmer L.M. (1989): [Review of Houde 1988.] – Journal of Vertebrate Paleontology 9: 368-369. [In English.]
- Witmer L.M. & Rose K.D. (1991): Biomechanics of the jaw apparatus of the gigantic Eocene bird *Diatryma*: implications for diet and mode of life. – Paleobiology 17: 95-120.
- Wittich E. (1899): Beiträge zur Kenntnis der Messeler Braunkohle und ihrer Fauna. – Abhandlungen der Grossherzoglichen Hessischen Geologischen Landesanstalt zu Darmstadt 3: 77-147.
- Woldrich J.N. (1893): Reste diluvialer Faunen und des Menschen aus dem Waldviertel Niederösterreichs in den Sammlungen des K. K. Naturhistorischen Hofmuseums in Wien. – Denkschriften der K. K. Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse 60: 565-622.
- Wolters H.-E. (1971) Probleme der Gattungsabgrenzung in der Ornithologie. – Bonner Zoologische Beiträge 22: 210-219.
- Womersley H. (1940): Studies on Australian Acarina; Tetranychidae and Trichadenidae. – Transactions of the Royal Society of South Australia 64: 233-265.
- Woodman P.C. (1978): The chronology and economy of the Irish Mesolithic: some working hypotheses. – In: Mellars P. (ed.): The early postglacial settlement of northern Europe: 331-369. London: Duckworth.
- Woodward A.S. (1889): Note on *Bucklandium diluvii* König, a siluroid fish from the London Clay of Sheppey. – Proceedings of the Scientific Meetings of the Zoological Society of London 1889: 208-210.
- Woodward A.S. (1890): A guide to the exhibition galleries of the Department of Geology and Paleontology in the British Museum (Natural History): Fossil mammals and birds. – London: British Museum (Natural History), 96 pp. [Aves on pp. 87-96.] [At least ten,

- only slightly emended editions appeared till 1923, most of them under the title: A guide to the fossil mammals and birds in the Department of Geology and Palaeontology in the British Museum (Natural History).]
- Woodward H. (1886): On "flightless birds" commonly called "wingless birds" fossil and Recent; and a few words on birds as a class. – Proceedings of the Geologists' Association (London) 9: 352-376.
- Woolfenden G.E. (1961): Postcranial osteology of the waterfowl. – Bulletin of the Florida State Museum, Biological Sciences 6: 1-129.
- Zadeh L. (1965): Fuzzy sets. – Information and Control 8: 338-353.
- Zarenkov N.A. (1976): Lekcii po teorii sistematiki [Lessons on the theory of systematics.] – Moskva: Izdatel'stvo Moskovskogo Universiteta, 140 pp. [In Russian.]
- Zbyszewski G. & Ferreira O. da V. (1967): Découverte de vertébrés fossiles dans le Miocène de la région de Leiria. – Comunicações dos Serviços Geológicos de Portugal 52: 5-10.
- Zeuner F. (1931): Die Insektenfauna des Böttinger Marmors. Eine systematische und paläobiologische Studie. – Fortschritte der Geologie und Palaeontologie 9(28): 247-406.
- Zigno A. de (1875): Sirenie fossili trovati nel Veneto [Fossil sirenas discovered near Veneto.] – Memorie Reale Istituto Veneto di Scienze, Lettere e Arti 18: 427-456. [In Italian.]
- Zigno A. de (1884): Comunicazione sopra ossa fossili di uccelli [Report on the fossil bones of birds.] – Rivista Periodica della Accademia di Padova 35: 5-6. [In Italian.]
- Zink R.M. & Slowinski J.B. (1995): Evidence from molecular systematics for decreased avian diversification in the Pleistocene epoch. – Proceedings of the National Academy of Sciences of the U.S.A. 92: 5832-5835.
- Zittel K.A. (1890): Vertebrata (Pisces, Amphibia, Reptilia, Aves). – In: Zittel K.A. (ed.): Handbuch der Palaeontologie, I. Palaeozoologie, vol. 3. – München: R. Oldenbourg, xii + 900 pp. [Aves on pp. 804-863.]
- Zittel K.A. (1899): Geschichte der Geologie und Paläontologie bis Ende des 19. Jahrhunderts. – München: R. Oldenbourg, xi + 868 pp.
- Zubareva V.I. (1939): Novaja forma pticy iz pliocena g. Odessy [New avian form from the Pliocene of Odessa town.] – Doklady Akademii Nauk SSSR 23: 606-608. [In Russian.]
- Zubareva V.I. (1948): Pliocenovi marabu i grycajja [Pliocene marabou and *Gryzaja*.] – Akademija Nauk Ukrajins'koji RSR, Trudy Ynstitutu Zoologiji 1: 114-137. [In Ukrainian.]

Index of authors

The authors are listed alphabetically without respect to diacritical marks. Authors of modern taxa and authors of all taxa cited without the year of publication are not listed here.

- Abel, Othenio (1875-1946) - 35, 277
 Adrover, R. - 37, 44, 47, 48, 49, 54, 92, 115, 154, 174, 193, 210, 234
 Aguirre, E. - 45
 Agustí, J. - 43, 53
 Ahlquist, Jon E. - 19
 Alcalde del Rio, Hermilio (1866-1947) - 57, 139
 Alcover, Josep Antoni - 49, 57, 79, 91, 92, 161, 210, 239
 Alekseev, A. K. - 43, 44, 61, 162
 Aleksiejew → Alekseev
 Alexander, R. McNeil - 17
 Alférez, F. - 156
 Aliev, S. D. - 161, 162
 Allen, E. G. - 25
 Alvarenga, Herculano Marcos Ferraz de - 102, 150
 Amadon, Dean - 19
 Ameghino, Florentino - 178
 Ammon, Ludwig von (1850-1922) - 40, 41, 71, 72, 111, 125, 155, 156, 193
 Anderson, J. F. - 17
 Andersson, Johann Gunnar - 62
 Andors, Allison Victor - 26, 28, 31, 94, 95, 185
 Andrae, Karl [Carl] Justus (1816-1885) - 41, 42
 Andrée, K. - 31
 Andrews, Charles William (1866-1924) - 28, 56, 64, 65, 81, 139
 Andrews, P. - 55, 169
 Anorova, N. S. - 165
 Antunes, Miguel Telles - 55, 57, 79, 91, 139, 140
 Archiac, Adolphe d' (1802-1868) - 25
 Arredondo, Oscar - 212
 Ascenzi, A. - 55, 139
 Aslanova, Sevil Mutalibovna - 81
 Aujard, C. - 26, 27, 261, 262
 Aver'janov, A. O. - 31, 81, 258, 259
 Avram, E. - 39, 273, 274, 275
 Aymard, Auguste - 33, 253, 254, 255, 270, 271, 279
 Azzaroli, A. - 213
 Bacher, Alois (*1940) - 17
 Bachmann, Isidor (1837-1884) - 39
 Bachmayer, Friedrich (*1913) - 39, 40, 45, 62, 209
 Bachofen-Echt, Adolf (1864-1947) - 29, 31, 225
 Bačinskij, G.A. - 164
 Bährmann, Udo - 231
 Baird, Robert F. - 28, 104, 257, 263, 266, 270
 Bakulovskij, N. N. - 44, 61
 Ballmann, Peter - 16, 35, 37, 38, 39, 40, 41, 42, 45, 46, 47, 54, 77, 144, 145, 146, 147, 148, 151, 154, 155, 156, 157, 158, 159, 174, 186, 191, 192, 204, 209, 210, 211, 212, 213, 215, 217, 219, 227, 228
 Balme, D. M. - 15
 Balouet, Jean-Christophe - 90
 Bardharon, G. G. - 139
 Baryshnikov → Baryšnikov
 Baryšnikov, Gennadij F. - 56, 162, 164, 219, 235, 236
 Bastin, A. - 56, 280
 Bate, Dorothea Minola Alice (1880-1951) - 55, 114, 139, 189
 Bauer, K. M. - 94
 Baumann, F. - 33, 226
 Bayan, Félix (1845-1874) - 36, 148, 225
 Bayer, František - 34, 70, 250, 251
 Beaman, Marc - 14
 Becker, Jonathan J. - 81
 Becker, L. - 37
 Beddard, Frank Evers (1858-1925) - 19, 25
 Bégouën, H. - 56, 139
 Bell, A. - 53, 57, 70, 140
 Bendukidze, Oleg Grigor'evič - 43, 96, 97, 162, 165
 Benecke, Norbert - 16, 164
 Beneden, Pierre Joseph van (1809-1894) - 34, 41, 79, 112, 125, 250, 251, 253, 256, 268, 271, 280
 Bengtson, S.-A. - 140
 Bensley, Benjamin Arthur (1875-1934) - 44, 62
 Berckhemer, F. (†1954) - 35
 Berendt, Georg Carl (1790-1850) - 31
 Berg, Dietrich E. - 29, 96, 185
 Berger, J.-P. - 38
 Berlin, Brent - 19
 Bessonat, G. - 34
 Bisseling, G. A. L. - 140
 Björklund, M. - 133
 Blainville, Henri Ducrotay de (1777-1850) - 25
 Blanc, G. A. - 57, 139
 Blanchard, Emile (1819-1900) - 32
 Blasius, August Wilhelm Heinrich (1845-1912) - 140
 Bocheński, Zbigniew M. - 231

- Bocheński, Zygmunt - 6, 18, 37, 41, 42, 47, 48, 49, 50, 51, 98, 99, 100, 108, 131, 132, 137, 156, 157, 158, 159, 161, 163, 166, 167, 168, 169, 221
- Bock, Walter J. - 13, 14, 19, 200
- Boehm → Böhm
- Boeuf, Odile - 49, 117, 122
- Boev, Zlatozar Nikolaev - 40, 44, 45, 47, 50, 51, 55, 56, 57, 94, 98, 113, 114, 117, 121, 130, 157, 160, 161, 163, 165, 166, 167, 169, 176, 177, 178, 189, 190, 191, 192, 193, 204, 215, 220, 222, 228, 229, 230, 233, 234, 235, 236, 238, 239, 241, 244, 245, 246, 247, 249
- Böhm, Georg (1854-1913) - 35, 277
- Boles, Walter E. - 21
- Bonaparte, Charles Lucien (1803-1877) - 32, 39, 95, 172
- Bonifay, E. - 55, 189, 213
- Bosák, Pavel - 51, 167
- Bosniaski, Sigismundo de - 45, 247
- Boule, Marcellin (1861-1942) - 26, 165, 189
- Bowerbank, James Scott (1797-1877) - 28, 83
- Bowler, P. J. - 25
- Bradley, F. - 45
- Bräm, Heinrich - 36
- Brandt, Aleksandr Fedorovič (1844-?1918) - 44, 60, 61
- Bräuhäuser, M. - 42
- Breuil, Henri Edouard Prosper (1877-1961) - 56, 57, 139
- Brjuzgina → Umans'ka
- Brochon, E. H. - 38
- Brodkorb, Pierce (1908-1992) - 6, 13, 16, 19, 31, 32, 34, 37, 38, 41, 45, 47, 49, 50, 60, 62, 63, 64, 66, 68, 71, 72, 73, 76, 79, 80, 82, 83, 85, 87, 89, 95, 96, 98, 99, 101, 104, 107, 110, 111, 113, 114, 117, 118, 120, 124, 125, 131, 140, 145, 150, 152, 154, 155, 156, 157, 159, 160, 165, 172, 173, 174, 175, 178, 179, 181, 182, 185, 186, 189, 194, 195, 199, 200, 201, 202, 204, 207, 208, 209, 210, 212, 215, 217, 218, 223, 224, 226, 232, 250, 251, 252, 253, 255, 257, 258, 259, 260, 263, 264, 267, 268, 269, 270, 272, 275, 277, 280
- Bronn, Heinrich Georg (1800-1862) - 24
- Brückner, Eduard - 8
- Bruijn, H. de - 8
- Brunet, Jean - 32, 33, 76, 108, 141, 154, 172, 173, 179, 181, 187, 224, 251, 266, 267, 269
- Brunner, Georg (1887-1959) - 52, 56, 161, 167, 249
- Brustur, Titus - 35
- Buckland, William (1784-1856) - 24
- Buffeteaut, Eric - 94, 95, 96, 185
- Bullen, R. P. - 140
- Burchak-Abramovich → Burčak-Abramovič
- Burczak-Abramowicz → Burčak-Abramovič
- Burčak-Abramovič, Nikolaj Ivanovič (1900-1997) - 25, 43, 44, 45, 47, 49, 56, 60, 61, 62, 72, 73, 81, 97, 116, 161, 162, 164, 165, 170
- Cailleux, André - 26, 27, 261
- Campbell, Kenneth E. Jr. - 17
- Camper, Pieter (1722-1789) - 24
- Čapek, Václav (1862-1926) - 45, 50, 52, 53, 99, 119, 120, 121, 124, 128, 148, 149, 161, 165, 176, 190, 200, 212, 214, 215, 221, 222, 227, 229, 230, 234, 235, 238, 239, 240, 241, 242, 243, 244, 245, 246, 249
- Capitan, L. - 57, 164
- Casanovas, M. L. - 33, 34
- Cassoli, Pier Franco - 51, 53, 55, 56, 57, 91, 139, 140, 162, 219
- Cavallo, O. - 45
- Čerkašenko, M. I. - 34
- Chandler, Robert M. - 64
- Cheneval, Jacques - 13, 17, 26, 27, 29, 31, 33, 34, 35, 36, 37, 38, 41, 42, 43, 44, 45, 63, 66, 67, 68, 69, 71, 78, 79, 80, 86, 93, 94, 100, 102, 103, 104, 105, 106, 107, 109, 110, 111, 112, 115, 118, 119, 120, 121, 130, 146, 147, 154, 156, 157, 158, 174, 175, 176, 188, 191, 193, 194, 210, 218, 230, 231, 251, 257, 258, 280
- Chesser, R. T. - 18
- Chozackij, Lev Isaakovič - 48, 100
- Christol, J. de - 45
- Chu, P. C. - 133
- Cirić, A. - 45
- Claußen, Marco - 295
- Cleere, H. - 139
- Clerq, S. W. de - 36
- Clot, André - 50, 53, 56, 57, 91, 161, 164, 168, 189, 233, 234, 238, 239, 240
- Clottes, J. - 56, 139
- Codrea, V. - 41
- Coiter, Volcher (1534-1576) - 25
- Collins, Charles T. - 31, 37, 201, 202, 204
- Comaschi Caria, I. - 55, 213
- Cope, Edward Drinker (1840-1897) - 81, 94, 95
- Cordy, J.-M. - 51
- Coues, Elliott (1842-1899) - 95
- Courtin, J. - 56, 139
- Courtois, J.-Y. - 214
- Cowles, Graham S. - 55, 57, 91, 92

- Cracraft, Joel - 19, 28, 30, 31, 32, 33, 34, 37, 41, 43, 44, 45, 58, 59, 85, 87, 89, 90, 93, 96, 97, 101, 102, 147, 152, 153, 154, 171, 172, 173, 174, 175, 176, 179, 180, 181, 182, 183, 184, 193, 194, 226, 254, 256, 262, 264, 270
- Cramp, Stanley (1913-1987) - 94, 213
- Crochet, J.-Y. - 35, 199
- Crusafont, Pairó Miguel - 38, 42, 155, 157, 159, 264
- Cuvier, Georges (1769-1832) - 6, 24, 25, 32, 35, 76, 179, 180, 181, 187, 266
- Dames, Wilhelm (1843-1898) - 26, 268, 269
- Daniels, Michael C. S. - 27, 29, 142, 197, 198, 262
- Dantin, J. - 41
- Darga, Robert - 35
- Darwin, Charles (1809-1882) - 24, 25
- Dauphin, Yannicke - 26, 261
- Davies, William - 96
- Davitašvili, Leo Šiovič (1895-1977) - 25
- De Zigno → see Zigno
- Dechaseaux, Colette - 32, 173
- Dehaut, E. G. - 55, 212, 213
- Dehm, Richard (†1996) - 33, 34, 38, 39, 49
- Del Hoyo, Josep - 94, 213
- Delamétherie, Jean Claude (1743-1817) - 32
- Delle Cave, Laura - 30, 31, 32, 37, 44, 45, 46, 47, 49, 64, 120, 247, 252
- Delpech, F. - 57, 91
- Demathieu, G. - 33, 275, 276
- Dement'ev, Georgij Petrovič (1898-1969) - 82, 91, 100, 131, 159
- Depéret, Charles (1854-1929) - 37, 41, 43, 45, 47, 105, 117, 156, 157, 158, 159, 161, 163, 232, 238, 264
- de Raaf → Raaf
- DesFayes, M. - 94
- Desnoyers, Jules (1800-1887) - 32
- Desor, E. - 82
- Despott, Giuseppe (†1936) - 56, 189
- De Stefani – see Stefani
- Dieffenbach, Ernst (1811-1855) - 41
- Dollo, Louis (1857-1931) - 26, 29, 34, 83, 95
- Douxami, Henri - 37, 105
- Dubois, A. - 14
- Dubrovo, Irina Aleksandrovna - 38, 43, 44, 47, 49, 51, 62, 72, 99, 115, 161, 165, 176, 191
- Dughi, Raymond - 26, 27, 29, 94, 261, 262
- Duméril, Constant (1774-1860) - 26, 95
- Dyke, Paul G. - 141, 203, 259
- Eastham, Anne - 57, 139
- Eastman, Charles Rochester (1868-1918) - 34, 62, 151, 154
- Eikamp, Heinz - 29, 37, 38
- Elorza, Miguel - 140
- Emslie, Steven D. - 64, 102
- Engel, T. - 279, 280
- Engelhardt, H. - 41
- Ennouchi, Emile - 41, 131, 154, 156, 158, 159, 191, 192, 204, 109, 215
- Ericson, Per G. P. - 26, 103, 107, 269
- Errico, Francesco d' - 56, 57, 139
- Escher von der Linth, Arnold (1807-1872) - 33, 226
- Escuillié, François - 37, 105, 258
- Espinosa de los Monteros, A. - 225
- Evans, P. G. H. - 140
- Fabre-Taxy, Suzanne - 26, 261
- Fahlbusch, Volker - 8, 39, 41
- Falconer, Hugh (1808-1865) - 56, 114
- Fatio, V. - 94
- Faujas-de-Saint-Fond, Barthélemy (1741-1819) - 32
- Feduccia, J. Alan - 26, 28, 37, 95, 103, 104, 105, 106, 107, 109, 141, 263, 265, 269
- Fejfar, Oldřich (*1931) - 8, 11, 43, 47, 48, 115, 122, 188, 234
- Ferreira, O. da V. - 39
- Fischer, Karlheinz - 28, 29, 30, 31, 34, 37, 43, 55, 78, 89, 90, 91, 92, 96, 136, 143, 151, 152, 155, 185, 217, 218, 265, 278
- Fischer, O. - 47
- Flerov, K. K. - 38, 273
- Flot, Léon - 32, 75
- Ford, N. L. - 213
- Forsyth Major, Charles Immanuel (1843-1923) - 44, 62
- Fraas, Eberhard (1862-1915) - 279
- Fraas, Oskar (1824-1897) - 40, 41, 86, 87, 107, 110, 113, 118, 156, 279
- Freeman, Patricia W. - 15, 17
- Freudenberg, W. - 53, 167, 168
- Freyer, Heinrich (1802-1866) - 226
- Friant, Madeleine - 57, 69, 70, 189, 190
- Fuller, Errol - 140
- Fürbringer, Max (1846-1920) - 19, 25, 81, 94, 103, 259
- Furon, Raymond - 224
- Gadow, Hans (1855-1928) - 19, 25
- Gaillard, Claude (1861-1946) - 25, 28, 31, 32, 33, 37, 41, 47, 50, 94, 96, 125, 130, 131, 150, 151, 152, 154, 155, 156, 157, 159, 163, 165, 173, 181, 182, 183, 184, 185, 186, 193, 194, 195, 196, 198, 199, 201, 202, 203, 204, 208, 209, 218, 254, 264, 269, 270, 280
- Gál, Erika - 40, 42, 43, 68, 72, 74, 99, 106, 110, 122, 123, 132, 176, 177, 178,

- 252
 Gall, H. - 39
 Ganea, Ivan Michajlovič (1926-1993) - 43, 44, 45, 49, 55, 56, 61, 62, 68, 69, 72, 89, 90, 116, 117, 138, 162, 164, 165, 168, 170, 174, 175, 196
 Ganja → Ganea
 García Talavera, I. - 81
 Gasser, U. - 37
 Gastaldi, Bartolommeo (1818-1879) - 45, 250
 Gaudant, J. - 45
 Gaudin, C. T. - 35
 Gaudry, Albert (1827-1908) - 45, 90, 101, 162
 Geoffroy-Saint-Hilaire, Isidor (1805-1861) - 32
 George, T. N. - 16
 George, W. - 27
 Germar, E. Franz (*1786) - 56, 271, 272
 Géroutet, Paul - 94
 Gervais, Paul (1816-1879) - 6, 25, 26, 32, 33, 34, 35, 36, 45, 47, 95, 106, 107, 151, 152, 163, 172, 173, 181, 187, 224, 226, 260, 269, 277
 Gesner, Konrad (1516-1565) - 94
 Geus, Armin - 94
 Gheorghiu, C. - 44
 Giebel, Christoph Gottfried (1820-1881) - 25, 31, 32, 50, 56, 99, 179, 180, 181, 188, 195, 212, 231, 232, 248, 249, 272, 277, 279, 280
 Gilly, C. L. - 15
 Ginsburg, Léonard - 38, 275, 276
 Glass, B. - 24
 Glutz von Blotzheim, Urs N. - 94
 Göhlich, Ursula B. - 35, 37, 39, 77
 Goldin, G. K. - 47, 62
 Gonzales Morales, J. A. - 57
 Gould, Stephen Jay - 16, 17
 Granger, Walter - 94, 95, 96
 Greenway, James C. Jr. - 140
 Gregor, Hans-Joachim - 40, 177
 Grieve, Symington (1848-1932) - 140
 Grigorescu, Dan - 42, 56, 61, 164
 Grigson, Caroline - 140
 Gromov, V. I. - 49, 56, 62, 164
 Grozescu, H. G. - 39, 273
 Grycaj, T. G. - 47
 Guérin, C. - 8, 11, 275, 276
 Haarhoff, Pippa - 41, 146
 Habersetzer, J. - 29, 87
 Hably, L. - 47
 Hachler, Emil - 19
 Haffer, Jürgen - 18
 Hamedani, H. - 78
 Hardy, M. - 57, 139
 Harris, A. H. - 231
 Harrison, Colin James Oliver - 24, 26, 27, 28, 29, 30, 31, 32, 33, 38, 44, 49, 51, 52, 53, 55, 57, 58, 59, 63, 64, 65, 66, 67, 68, 71, 72, 73, 76, 79, 80, 82, 83, 85, 87, 89, 95, 96, 98, 99, 101, 104, 107, 110, 111, 113, 114, 117, 118, 120, 124, 125, 131, 140, 145, 150, 152, 154, 156, 157, 159, 160, 165, 172, 173, 174, 175, 178, 179, 181, 182, 185, 186, 189, 194, 195, 199, 200, 201, 202, 204, 207, 209, 210, 212, 215, 217, 218, 223, 224, 226, 232, 250, 251, 252, 253, 255, 257, 258, 259, 260, 263, 264, 267, 268, 269, 270, 272, 275, 277, 280
 Hassencamp → Hassenkamp
 Hassenkamp, Ernst (1824-1881) - 33, 262
 Haubold, Hartmut - 29, 30, 60
 Haushalter, Carl Ludwig (1834-1872) - 38, 251
 Hay, Oliver Perry (1846-1930) - 140
 Hazevoet, Cornelis J. - 256
 Hébert, Edmond (1812-1890) - 26, 94, 95
 Heer, Oswald (*1809) - 33, 41, 118, 266
 Heinrich, Wolf-Dieter - 8, 11, 43, 47, 48, 50, 122
 Heintz, Émile - 90
 Heissig, K. - 32, 33, 34, 39
 Heizmann, Elmar P. J. - 17, 41, 64, 86, 87, 105, 106, 107, 110, 113
 Heller, Florian (1905-1978) - 29, 47, 171
 Hernández-Pacheco, Eduardo (*1872) - 34, 37, 41, 48, 140
 Hescheler, K. - 94
 Hesse, Angelika - 17, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 50, 64, 86, 87, 99, 105, 106, 107, 110, 113
 Hill, A. - 279
 Hirsch, Karl F. - 29
 Hitchcock, Edward (1793-1864) - 277
 Hobson, K. A. - 140
 Hoch, Ella - 26, 27, 29, 59, 92, 93, 141, 142
 Hoeninghaus, Friedrich Wilhelm (1770-1854) - 37
 Hoerschelmann, Heinrich - 17
 Hoff, K. E. A. von - 6
 Hoffmann, A. - 16
 Holst, H. K. K. - 36
 Hondt, J.-L. d' - 14
 Hopson, J. A. - 82
 Horáček, Ivan (*1952) - 8, 11, 47, 49, 53
 Hörmann, K. - 57, 164
 Hou, Lian-Hai - 94
 Houde, Peter - 26, 27, 28, 29, 30, 58, 59, 60, 102, 141, 255, 257
 Howard, Hildegard (1901-1998) - 41, 45, 64, 101, 104, 108, 109, 110, 112, 117,

- 118, 124, 125, 250, 253
Hufthammer, A. K. - 140
Hunter, John - 59
ICZN - 6, 8, 13, 14, 68, 80, 85, 88, 89,
151, 175, 195, 200, 235, 236, 253, 277
Illies, H. - 27
Imbrie, J. - 16, 18
Inger, R. F. - 14
Jan'kova, V. S. - 50, 62
Jánossy, Dénes - 17, 25, 43, 44, 45, 47,
48, 49, 50, 51, 52, 53, 54, 55, 56, 62,
69, 73, 76, 85, 86, 90, 92, 94, 98, 99,
101, 115, 116, 117, 118, 119, 120, 121,
122, 123, 124, 127, 128, 129, 130, 133,
135, 137, 138, 148, 149, 154, 155, 157,
159, 160, 161, 162, 163, 165, 166, 167,
168, 169, 170, 174, 176, 177, 178, 186,
187, 188, 189, 190, 191, 192, 196, 200,
204, 205, 209, 211, 212, 213, 214, 216,
217, 219, 220, 221, 222, 227, 228, 229,
230, 232, 233, 234, 235, 237, 238, 239,
240, 241, 242, 243, 244, 245, 246, 247,
249, 263
Jehenne, Yves - 90
Johannsen, W. - 15
Jollie, Malcolm - 31, 102, 194, 264
Jordan, R. H. - 140
Jörg, E. - 43
Jullien, R. - 42
Junge, G. C. A. - 49, 188
Jurcsák, Tiberiu - 51, 55, 66, 159, 164,
173, 232
Jurkovšek, Bogdan - 37
Kapelist, Kira Valerianovna - 38, 43, 44,
47, 49, 51, 62, 72, 99, 115, 161, 165,
176, 191
Karaivanova → Karajvanova
Karajvanova, Elena - 57, 178
Karchu, Aleksandr A. - 31, 32, 44, 96, 97,
98, 164, 200, 201, 202, 203
Karg, J.-M. - 24, 41, 279
Karhu – see Karchu
Karkhu – see Karchu
Kašin, Georgij Nikolaevič - 66, 67, 106,
186
Katinas, V. - 31, 225
Keferstein, Christian (1784-1866) - 56,
271, 272
Kerkhoff, N. C. - 140
Kerourio, P. - 26, 27, 261, 262
Kessler, Eugen - 30, 34, 39, 41, 42, 43,
47, 51, 55, 64, 66, 67, 68, 72, 74, 80,
85, 99, 101, 105, 106, 110, 111, 117,
118, 119, 120, 121, 123, 129, 132, 137,
138, 159, 161, 164, 166, 173, 176, 177,
228, 229, 231, 232, 233, 234, 240, 252
Ketraru, N. A. - 55, 56, 164
Killermann, S. - 94
King, P. P. - 277
Kiriakoff, S. G. - 18
Kitcher, P. - 15
Kleinschmidt, A. - 57, 190
Knowles, W. J. - 139
Koenig → König
Kohring, R. - 29
Kolar-Jurkovšek, Tea - 37
Kompanje, E. O. - 140
König, Carl Dietrich Eberhard (1774-
1851) - 224, 28, 259, 277
König, Claus - 213
Kon'kova, Nadežda Radionovna - 62
Kordos, László - 38, 51, 273, 275, 276
Kormos, Tivadar (1881-1946) - 159
Korotkevič, E. L. - 44, 61
Kotsakis, T. - 55, 213
Koufos, G. - 47, 163
Kovačev, D. - 44, 191
Kozlova, Elizaveta Vladimirovna - 96
Kraft, Elfriede (*1943) - 15
Kraglievich, Lucas (1886-1932) - 178,
179
Krejčí, Jan (1825-1887) - 81, 251
Kretzoi, Miklós - 25, 43, 47, 48, 49, 50,
51, 52, 53, 60, 61, 62, 90, 99, 113, 157,
159, 162, 165, 166, 167, 170, 196, 217,
219, 220, 221, 222, 227, 229, 232, 233,
235, 239, 249, 263, 279
Kristoffersen, Anette Vedding - 26
Krolopp, E. - 51, 263
Krüger, J. F. - 6, 24
Krumbiegel, Günther - 29
Krupska, Anna - 165
Kuhn, E. - 94
Kuhn, O. - 277
Kukla, Jiří [Georg] - 8
Kumerloeve, Hans - 94
Kuntz, P. - 30
Kurochkin → Kuročkin
Kuročkin, Evgenij Nikolaevič - 43, 44,
45, 47, 48, 50, 61, 62, 64, 68, 69, 72,
89, 90, 96, 97, 98, 99, 100, 111, 112,
115, 116, 117, 138, 158, 159, 160, 161,
162, 163, 165, 174, 175, 196
Kuss, S. E. - 37, 105
Kuźnicki, Leszek - 15
La Barbera, M. - 17
La Milza, J.-C. - 214
Lageræus, P. - 140
Laizer, L. de - 35, 106
Lakerbaj, L. B. - 164
Lamanon, Robert Paul de (1752-1787) -
24, 32
Lambrecht, Kálmán (1889-1936) - 6, 8,
11, 13, 19, 24, 25, 26, 28, 29, 30, 31,
32, 33, 34, 35, 36, 37, 38, 39, 40, 41,
42, 43, 44, 45, 47, 49, 50, 55, 56, 57,

- 58, 59, 60, 61, 62, 66, 67, 68, 70, 71, 72, 73, 80, 81, 82, 83, 85, 87, 89, 90, 92, 93, 96, 97, 99, 100, 101, 103, 105, 106, 107, 110, 111, 112, 114, 124, 125, 139, 140, 145, 146, 155, 159, 160, 161, 173, 174, 175, 176, 178, 184, 187, 193, 194, 195, 196, 218, 223, 224, 225, 230, 231, 236, 247, 248, 252, 253, 255, 256, 257, 259, 264, 265, 267, 268, 269, 275, 277, 279, 280
- Landini, W. - 45
- Lartet, Edouard (1801-1871) - 26, 38, 81, 83, 84, 95
- Lascarev → Laskarev
- Laskarev, Vladimir Dmitrievič - 37, 43, 44, 162
- Łaskarew → Laskarev
- Laube, Gustav C. (1839-1923) - 38, 101, 110, 111, 271
- Launay, Louis de (*1860) - 26
- Laurent, Goulven - 24
- Laurillard, Charles Léopold (1783-1853) - 32, 224
- Lavocat, Achille - 26, 95
- Lebedinsky, Naum Gregor (*1888) - 33, 107, 108
- Legendre, S. - 15, 17
- Leisler, Bernd - 15
- Lemen, Cliff A. - 15, 17
- Lemoine, Victor (1837-1897) - 26, 95, 255, 256, 268
- Lepsius, Georg Richard (1851-1915) - 279
- Lévêque, S. - 8
- Liberov, P. D. - 164
- Lie, R. - 56, 139
- Lierl, Hans-Jürgen - 26, 269
- Livezey, Bradley C. - 19, 37, 109, 110, 111, 114, 140
- Ljaščuk, B. F. - 34
- López, N. - 41
- Löppenthin, B. - 57, 139
- Louchart, Antoine - 166
- Lowe, Percy Roycroft (*1870) - 31, 37, 62, 201
- Ložek, Vojen - 8, 47, 53
- Lucas, Frederic Augustus (1852-1929) - 151
- Lucazeau, Ginette - 38, 80, 279, 280
- Lukáš, Jiří - 56
- Lungu, Aleksandr Nikolaevič - 43, 44, 61
- Luther, Dieter - 140
- Lydekker, Richard (1849-1915) - 6, 24, 25, 27, 28, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 44, 49, 53, 55, 56, 57, 59, 62, 63, 67, 70, 71, 80, 81, 83, 86, 87, 91, 92, 93, 95, 97, 99, 100, 105, 106, 107, 108, 110, 111, 114, 125, 131, 136, 153, 155, 156, 159, 170, 172, 178, 184, 187, 189, 193, 194, 198, 200, 201, 209, 210, 217, 219, 250, 253, 254, 258, 259, 266, 270
- Lyell, Charles (1797-1875) - 25
- Macarovici, N. - 43, 115, 117, 121
- Macdonald, J. R. - 151
- Magish, D. P. - 231
- Mäklin, Fredrik Wilhelm - 44, 61
- Malez, V. → Malez-Bačić
- Malez-Bačić, Vesna - 49, 52, 53, 56, 161, 164, 220, 239
- Mangin, J. F. - 33
- Marcus, Leslie - 17
- Marisova, I. V. - 56, 164, 165
- Marmora, Alberto de la (1789-1863) - 24, 54
- Marsh, Othenio Charles (1831-1899) - 95, 259
- Martin, D. S. - 28, 44, 62, 82
- Martin, Larry D. - 26, 28, 37, 95, 96, 109, 110, 111, 256, 263, 268
- Martini, Erlend - 33, 37, 71, 105, 106, 107, 129, 262
- Massalongo, Abramo Bartolomeo (1824-1860) - 45, 132
- Matheu, Eloisa - 94
- Matthes, H. W. - 29
- Matthew, W. D. - 94, 95, 96
- Mauersberger, Gottfried (1931-1994) - 31, 37
- Mayaud, Noel - 57, 80, 81
- Mayr, Ernst (*1904) - 15, 16, 19
- Mayr, Gerald - 27, 29, 31, 32, 33, 34, 36, 49, 75, 76, 93, 130, 133, 141, 142, 143, 144, 145, 170, 177, 184, 197, 198, 201, 203, 205, 223, 224, 226, 252, 257, 259, 262, 267, 268, 269, 270, 271
- McDonald, Joan F. - 56, 57, 139
- McMahon, T. A. - 17
- McMinn, Miquel - 79
- Mecquenem, R. de - 96, 97
- Mein, Pierre - 8, 11, 43
- Meldgaard, M. - 140
- Mellars, Paul A. - 139
- Mengel, R. M. - 28, 263
- Menzel, P. - 38
- Merccerat, A. - 178
- Merrem, Blasius (1761-1824) - 25
- Métherie de la → Delamétherie
- Meunier, Stanislas - 26, 31, 95
- Meyer, Adolf Bernard (1840-1911) - 31
- Meyer, Hermann von (1801-1869) - 25, 31, 33, 37, 38, 41, 118, 124, 125, 226, 262, 279
- Mežlumjan, S. - 165
- Mezzena, F. - 57, 139
- Michajlov, Konstantin E. - 26, 43, 44, 47,

- 48, 50, 61, 62, 94
 Michaux, A. - 34
 Mieg, M. - 34
 Miller, Alden Holmes (1906-1965) - 34, 66, 87, 103, 104, 136, 138
 Miller, Loye H. (1874-1970) - 64, 80, 104
 Milne-Edwards, Alphonse (1835-1900) - 6, 13, 14, 25, 26, 27, 31, 32, 33, 34, 35, 36, 37, 38, 41, 44, 45, 47, 54, 55, 56, 57, 62, 63, 66, 67, 68, 69, 70, 71, 72, 75, 76, 77, 79, 80, 86, 89, 90, 91, 92, 93, 94, 95, 103, 104, 105, 106, 107, 109, 110, 111, 118, 124, 125, 130, 131, 132, 134, 136, 137, 138, 145, 146, 147, 151, 152, 153, 154, 156, 158, 172, 173, 174, 175, 176, 178, 179, 180, 181, 182, 183, 184, 186, 187, 188, 193, 194, 195, 196, 198, 199, 201, 203, 204, 207, 208, 209, 210, 211, 217, 218, 223, 224, 225, 230, 231, 236, 248, 253, 254, 255, 256, 257, 258, 260, 267, 269, 270, 271, 279, 280
 Misonne, Xavier - 26, 27, 29, 33
 Mlíkovský, Jiří (*1954) - 6, 7, 8, 11, 14, 15, 16, 17, 18, 19, 20, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 64, 65, 66, 68, 70, 71, 72, 73, 76, 77, 82, 85, 86, 89, 90, 96, 99, 101, 103, 105, 106, 107, 108, 110, 111, 112, 113, 114, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 132, 135, 136, 137, 138, 139, 141, 143, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 187, 188, 189, 190, 191, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 225, 226, 227, 232, 233, 234, 235, 237, 239, 240, 241, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 254, 257, 262, 265, 266, 267, 269, 270, 271, 272, 273, 274, 275, 277, 280
 Moldvai, Mihai - 231, 234
 Montevecchi, W. A. - 140
 Montoya, P. - 94, 155, 186, 212, 215, 219, 227, 237, 238, 242, 243, 245
 Morales, Arturo - 45
 Moreno, Eulalia - 231
 Moreno, F. P. - 178
 Morris, J. - 277
 Mottl, Maria - 56
 Mourer-Chauviré, Cécile - 16, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 39, 43, 45, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 68, 70, 73, 79, 87, 90, 91, 92, 102, 107, 117, 122, 126, 134, 138, 139, 140, 141, 143, 144, 150, 151, 152, 154, 155, 159, 161, 162, 163, 164, 165, 166, 167, 168, 169, 179, 181, 182, 183, 184, 185, 189, 190, 191, 192, 194, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 212, 213, 214, 215, 217, 218, 219, 221, 222, 223, 224, 225, 226, 229, 230, 231, 232, 233, 234, 235, 236, 238, 239, 240, 243, 244, 245, 246, 247, 255, 258, 260, 262, 265, 267, 269, 270
 Moyà, Salvador - 210
 Müller, Arno Hermann - 14, 96
 Munthe, H. - 57, 139
 Murie, James - 32, 224
 Musil, Rudolf - 51, 52, 53, 167
 Nathusius, Wilhelm von (1828-1899) - 44, 61
 Navás, Longinos (1858-?1936) - 43, 85
 Nehring, Alfred (1845-1904) - 56, 168, 277
 Nesov, Lev Aleksandrovič (1947-1995) - 34, 258, 259
 Nesson → Nesov
 Nettleship, D. N. - 140
 Newton, Alfred (1829-1907) - 140
 Newton, Edwin Tullney - 26, 49, 51, 53, 73, 80, 95, 115, 120, 129, 139
 Nikolov, Ivan - 44, 72, 73, 116
 Nitzsch, Christian Ludwig (1782-1837) - 25, 117
 Nogalski, Stanisław - 165
 Nordmann, Alexander von - 44
 Noriega, Jorge Ignacio - 104
 Northcote, E. Marjorie - 17, 55, 56, 57, 70, 91, 92, 114, 189
 Novák, Otomar (1851-1892) - 39
 Oberholser, Harry Church (*1870) - 32, 63, 113, 178, 180, 182, 183, 200, 267, 270
 Oescu, C. V. - 43, 115, 117, 121
 Ollavaría, M. de - 37
 Olsen, H. - 140
 Olson, Storrs L. - 13, 19, 20, 26, 27, 28, 31, 32, 33, 34, 36, 37, 39, 41, 42, 43, 45, 55, 64, 65, 68, 72, 81, 82, 85, 87, 93, 97, 102, 103, 104, 105, 106, 107, 108, 109, 126, 133, 136, 137, 140, 141, 147, 154, 155, 156, 157, 171, 172, 173, 174, 175, 176, 197, 198, 200, 212, 219, 225, 226, 251, 253, 254, 255, 256, 257, 258, 259, 260, 263, 265, 266, 267, 268, 269, 270
 Omboni, Giovanni (1829-1910) - 32, 261

- Osborn, H. F. - 15
Owen, Richard (1804-1892) - 25, 26, 27, 28, 58, 59, 81, 82, 83, 95, 259, 266
Palma di Cesnola, A. - 57, 139
Palmowski, J. - 34
Panin, N. - 39, 273, 274, 275
Pankakoski, Erkki - 17
Panteleev, Andrej - 279
Paris, Paul (1875-1938) - 76, 92, 117, 193, 195, 196, 204, 218, 224, 236, 256, 257, 268, 270
Parker, William Kitchen (1823-1890) - 56, 113, 114
Parkinson, James (1755-1824) - 24
Parris, David C. - 108, 259
Patterson, Bryan - 178, 179
Pauçã, M. - 39
Pavia, Marco - 53, 55, 94, 212, 213, 214
Penck, Albrecht - 8
Péquart, J. S. - 140
Péquart, Marthe - 140
Pereira, É. - 214
Peters, Dieter Stephan - 29, 30, 60, 92, 93, 103, 142, 143, 182, 185, 201, 203, 207, 270
Petersen, A. - 139
Petrov, S. G. - 165
Peyer, Bernhard - 24, 33, 38, 226, 266
Pfannenstiel, Max (1902-1976) - 279
Pharisat, André - 79
Philippe, M. - 55, 191, 233
Pichon, Joelle - 163, 164
Pictet, François Jules (1809-1872) - 32
Pidopličko, I. G. - 43, 47, 61, 62
Pieper, Harald - 139
Pitton, L. E. - 26
Piveteau, Jean (1899-1991) - 96
Plaziat, Jean-Claude - 27, 261
Pons, Juan - 210
Popov, R. - 44
Portis, Alessandro (1853-1931) - 30, 31, 32, 37, 44, 45, 47, 49, 57, 86, 92, 119, 120, 132, 140, 195, 218, 219, 232, 233, 247, 249, 250, 252, 261, 267, 275, 277
Potapov, R. L. - 168
Potapova, Olga R. - 56, 57, 127, 128, 140, 161, 162, 164, 165, 219, 235, 236, 258, 259
Poulianos, Nikos A. - 52, 53, 162, 166, 196, 219, 220, 221, 229, 235, 239
Prévost, Constant (1787-1856) - 26, 95
Preyer, W. - 140
Probst, Joseph (1823-1905) - 39, 40, 41
Procaccini Ricci, Vico - 37, 45, 261
Prum, R. O. - 147
Prummell, Wietske - 16
Pržemysskij, K. - 44, 61
Pycraft, William Plane (1868-1942) - 45, 247
Quekett, John Thomas (1815-1861) - 83
Quenstedt, Friedrich August (1809-1889) - 33
Quinif, Y. - 214
Raaf, J. F. M. de - 33
Rădan, Silviu - 35, 274
Rabeder, Gernot - 37
Rage, Jean-Claude - 185
Ranke, J. - 56, 164
Rasmussen, D. Tab - 126
Rauscher, Karl L. - 40
Razoumowsky, A. - 24
Ray, C. E. - 139
Rayner, Jeremy M. V. - 17
Regália, Ettore - 41, 47, 57, 71, 73, 80, 81, 86, 139, 156, 232, 233, 252
Reichel, Manfred - 36, 37, 38, 40
Reichenbach, Heinrich Gottlieb Ludwig (1793-1879) - 32, 117, 179, 266, 279, 280
Reif, W.-E. - 18
Remane, J. - 18
Rensch, Bernhard - 17
Repelin, J. - 26, 95
Rey, Roger - 26, 261
Rhymer, J. M. - 18
Rich, Patricia Vickers - 28, 31, 33, 41, 58, 59, 71, 73, 101, 102, 146, 182, 193, 194, 257, 263, 264, 266, 270
Richmond, Charles Wallace (1868-1932) - 151, 179, 207
Ristori, Giuseppe - 45, 250
Rivière, E. - 57, 236
Roberton - 30
Rögl, Fred - 62
Rogovič, Afanasij Semenovič - 31, 258, 259
Rogowich → Rogovič
Rohlf, F. J. - 17
Rolle, Friedrich - 25
Ronsil, René - 330
Rosas, A. - 52, 121, 166, 188, 191, 219, 222, 245
Roščin, A. G. - 43, 44, 61, 62
Rose, K. D. - 95
Roset, F. - 43
Roška, V. Ch. - 43, 61
Rössner, Gertrud E. - 39
Roth, J. - 45
Rothausen, Karlheinz - 37, 38
Rovereto, Cayetano - 178
Rudwick, M. J. S. - 24, 25
Rühl, F. - 39
Russell, Donald E. - 26, 95
Rusu, A. - 34
Rütimeyer, L. - 57, 164
Sáez, Mathilde Dolgopol de - 178

- Saint-Périer, R. de - 57, 91, 189
 Salotti, Michelle - 54, 210, 214
 Salvadori, Tommaso (1825-1923) - 250
 Sánchez, Marco Antonio - 27, 33, 34, 37, 39, 41, 42, 43, 44, 45, 47, 48, 49, 50, 51, 53, 54, 86, 94, 98, 116, 120, 121, 137, 140, 165, 166, 210
 Santafé, J. V. - 33, 34
 Scarabelli-Gommi-Flamini, G. E. - 45, 132
 Scarlett, Ron J. - 83
 Schaub, Samuel - 28, 29, 34, 57, 80, 81, 96, 154, 213, 254
 Schenker, A. - 94
 Scherdlin, P. - 140
 Scheuchzer, Johann Jacob von (1670-1733) - 6, 24, 41
 Schilder, F. A. - 18
 Schlosser, Max - 41, 45, 55, 118, 164
 Schlothheim, Ernst Friedrich von (1764-1832) - 37
 Schmidt-Kittler, Norbert - 8, 11
 Schönwetter, Max - 44, 62
 Schottler, W. - 41
 Schweizer, W. - 165
 Seeley, Harry Govier (1839-1909) - 28, 83, 277
 Seemann, R. - 35, 40, 41
 Segre, A. G. - 55, 57, 139, 140
 Seguí, Bartolomeu - 128
 Seguí Campaner → Seguí
 Selenka, Emil - 25
 Serebrovskij → Serebrovskij
 Serebrovskij, P.V. - 47, 91, 100, 131, 217
 Serjeantson, Dale - 139
 Serres, Marcel de (1786-1862) - 45, 55
 Sesé, C. - 41
 Sharpe, Robert Bowdler (1847-1909) - 93, 156, 193, 195, 196
 Shaw, A. B. - 18
 Shufeldt, Robert Wilson (1850-1934) - 37, 41, 57, 80, 81, 95, 136, 137, 154, 156, 179, 208
 Sibley, Charles Gald (1917-1998) - 19, 34, 136, 138
 Sigé, B. - 31, 34
 Simberloff, D. - 18
 Simmons, K. E. L. - 94
 Simons, Elwyn L. - 126
 Simpson, George Gaylard (1902-1984) - 14, 16, 18
 Sinclair, William John - 95
 Sirugue, François - 26, 29, 94, 261, 262
 Skutil, J. - 52, 53
 Sleight, F. W. - 140
 Slowinski, J. B. - 16
 Smirnov, E. S. - 14
 Smith, G. A. - 146
 Smith, K. D. - 94
 Smith, R. - 26
 Smith, T. - 26
 Soergel, Elisabeth - 57, 91
 Solounias, N. - 44, 62
 Sondaar, Paul Y. - 50, 115, 122, 155, 176, 212, 228, 235, 236, 237, 239, 241, 242, 245, 246
 Soulé, M. E. - 17
 Soyer, Robert - 224
 Speck, Josef - 38
 Špinar, Zdeněk Vlastimil - 34
 Spulski, Boris - 82
 Steadman, David W. - 133
 Steenstrup, Johannes Japetus Smith (1813-1897) - 140
 Stehlík, Alois - 52, 53
 Stehlin, Hans Georg (*1870) - 50, 163, 167, 247
 Steininger, Fritz F. - 8, 62
 Stejneger, Leonhard (1851-1943) - 103, 113, 114, 193, 194, 253
 Stemvers-van-Bemmel, J. - 34
 Stent, G. S. - 15
 Stepanjan, Lev Surenovič - 18
 Stephan, Burkhardt - 43, 55, 89, 90, 91, 92, 155, 265
 Stevanović, Petr M. - 40
 Stewart, John R. - 51, 55, 116, 120, 121, 137, 139, 140, 165, 166, 168, 177, 216, 230, 237, 241
 Stingelin, Werner - 20
 Storch, G. - 29
 Storer, Robert W. - 19, 31, 37, 63, 85, 111
 Strauch, Joseph R. - 133
 Stresemann, Erwin (1889-1972) - 19, 25
 Strohl, J. - 94
 Stromer, E. von (†1952) - 43
 Studer, B. - 226
 Studiati, Césare (†1895) - 54
 Stüssi, F. - 33, 228
 Subarewa → Zubareva
 Sukačev, V. N. - 164
 Suriano, F. - 56
 Šušpanov, K. I. - 170
 Švec, Petr (*1954) - 34, 38, 39, 40, 41, 47, 48, 63, 68, 70, 85, 101, 105, 106, 107, 111, 112, 113, 115, 118, 122, 123, 124, 126, 132, 154, 155, 156, 157, 174, 250, 251
 Sylverster-Bradley, P. C. - 18
 Szymczyk, W. - 37
 Tagliacozzo, A. - 57
 Tambussi, Claudia - 81
 Tatarinov, Konstantin Adrianovič - 56, 164, 165
 Tchernov, Eitan - 160, 161, 163, 164, 177,

- 213
 Terzea, E. - 51
 Thenius, Erich - 29, 43
 Thesing, R. - 165
 Tintant, H. - 14
 Tkačenko, O. F. - 38.
 Tobien, Heinz (†1993) - 33
 Tomek, Tereza - 18, 231
 Tonni, Eduardo P. - 81
 Tordoff, H. B. - 151
 Toula, F. - 52, 116
 Touraine, Fernand - 26, 261
 Tret'jakov, D. K. - 47
 Tschudi, Johann Jacob von - 226
 Tugarinov, Arkadij Jakovlevič - 44, 47, 56, 71, 72, 112, 157, 159, 161, 163, 168, 169, 189
 Tyrberg, Tommy - 6, 7, 19, 26, 49, 50, 51, 52, 53, 54, 55, 56, 57, 91, 140, 166, 189, 200, 216, 225, 236, 246, 249, 269
 Uhlmann, E. - 15
 Umans'ka, Alla Semenovna - 44, 45, 64, 97, 98, 116, 117, 148, 164, 216, 220
 Uman'skaja → Umans'ka
 Ussher, Richard John (1841-1913) - 139
 Valenciennes, Achille - 26, 95
 Valoch, Karel - 52, 53
 Van Beneden → Beneden
 Van Valen, Leigh M. - 15
 Vekua, Abesalom Kapitonovič - 45, 61
 Verheyen, René Carl (1907-1961) - 19, 109
 Vialov → Vjalov
 Vickers-Rich → Rich
 Vidgal'm, Ignatij Martinovič (*1835) - 44, 72, 87
 Vilette, Philippe - 55
 Villalta Comella, José F. de - 38, 42, 43, 44, 49, 72, 137, 154, 155, 156, 159, 193, 235, 236, 245, 264
 Villatte, Juliette - 27, 261
 Villeneuve, L. - 165
 Vincent, Stephen - 27
 Vinokurov, A. A. - 140
 Violani, Carlo - 140
 Viret, J. - 34, 49, 163
 Vjalov, Oleg Stepanovič (1904-1988) - 38, 39, 60, 96, 273, 274
 Vojinstvens'kyj, Michail Anatol'evič - 43, 44, 45, 47, 49, 51, 53, 56, 61, 62, 64, 72, 86, 100, 119, 120, 127, 149, 161, 162, 163, 193, 212, 219, 224, 228, 232, 239, 242, 245
 Voznesens'kyj, O. N. - 44, 45
 Vuilleumier, François - 18
 Vyalov → Vjalov
 Wachendorf, H. - 34
 Wagner, Johann Andreas (1797-1861) - 45
 Wagner, Rudolf (1805-1864) - 24, 54, 117, 195, 232, 233, 249
 Wahlert, G. von - 15
 Walker, A. - 279
 Walker, Cyril Alexander - 26, 27, 28, 29, 30, 31, 33, 37, 38, 41, 49, 55, 58, 59, 65, 63, 64, 65, 73, 78, 80, 81, 82, 83, 84, 87, 95, 96, 97, 103, 106, 107, 109, 112, 141, 163, 171, 172, 180, 200, 201, 250, 251, 252, 253, 254, 257, 258, 259, 261, 263, 264, 265, 266, 271, 277, 279
 Ward, David J.
 Warheit, Kenneth I. - 17, 18
 Warter, Stuart L. - 32
 Weesie, Peter D. M. (*1953) - 54, 55, 56, 189, 192, 213, 214
 Weidmann, Marc - 36, 37, 38, 40
 Weigel, P. H. - 140
 Weigelt, Johannes - 26, 30, 95
 Weithofer, K. Anton - 45, 162, 250
 West, Barbara - 164
 Westphal, F. - 43
 Wetmore, Alexander (1886-1978) - 13, 19, 37, 64, 80, 81, 96, 178, 196
 Widhalm → Vidgal'm
 Wijngaarden-Bakker, Louise H. van - 140
 Wiley, E. O. - 16
 Wilkinson, H.E. - 80
 Willmann, Rainer - 16
 Wilson, L. G. - 25
 Winge, Herluf (1857-1923) - 140
 Witmer, Lawrence M. - 58, 95
 Wittich, Ernst (*1871) - 29, 92, 93
 Woldfich, Jan Nepomuk (1834-1906) - 56, 164, 168
 Wolters, Hans E. - 18, 19
 Woodman, P. C. - 140
 Woodward, Henry (1832-1921) - 26, 32, 33, 34, 36, 41, 81, 95, 277
 Woolfenden, Glen E. - 111, 116
 Zadeh, L. - 15
 Zapfe, Helmut (†1996) - 45, 62
 Zarenkov, N. A. - 16
 Zbyszewski, Georges - 39
 Zemek, Karel (*1946) - 15
 Zeuner, F. - 41
 Zhou, Ben-Xiong - 164
 Zigno, Achille de (1813-1892) - 30, 38
 Zink, R. M. - 16, 18
 Zittel, Karl-Alfred (1838-1904) - 24, 25, 81
 Zubareva, Valentina Ivanovna - 47, 98, 100
 Zusi Richard L. - 19

Index of countries

This list includes names of countries and their selected parts, especially archipelagos and islands

- Afghanistan - 90
 Armenia - 164
 Argentina - 104
 Australia - 104
 Austria - 37, 39, 40, 42, 52, 53, 56, 64, 65, 73, 76, 99, 118, 139, 146, 148, 149, 161, 164, 167, 168, 169, 177, 189, 190, 205, 209, 216, 221, 227, 234, 237
 Azerbaijan - 91, 162
 Balearics - 79, 91, 115, 122, 192, 210, 212, 228, 233, 235, 236, 237, 239, 241, 242, 245, 246
 Belgium - 26, 29, 33, 34, 41, 51, 55, 79, 81, 95, 112, 125, 190, 250, 251, 253, 256, 268, 271
 Brazil - 82, 150, 270
 Bulgaria - 40, 44, 45, 47, 49, 50, 51, 55, 56, 57, 73, 94, 98, 114, 116, 117, 130, 161, 163, 166, 167, 169, 176, 177, 178, 189, 190, 191, 192, 193, 204, 220, 222, 228, 229, 233, 234, 235, 236, 238, 239, 241, 244, 245, 246, 247, 249
 Buryatia - 161
 California - 64, 104
 Canary Islands - 79, 81
 China - 61, 96
 Corsica - 189, 210, 214
 Croatia - 41, 49, 52, 53, 56, 161, 164, 213, 226, 239
 Czechia - 34, 38, 39, 41, 51, 52, 53, 63, 68, 70, 71, 72, 73, 76, 85, 86, 94, 101, 104, 105, 106, 107, 110, 111, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 127, 128, 129, 132, 135, 136, 137, 138, 141, 144, 147, 148, 149, 151, 154, 155, 161, 164, 166, 167, 168, 169, 174, 177, 178, 187, 188, 191, 193, 197, 200, 204, 205, 211, 214, 215, 216, 217, 218, 220, 221, 222, 225, 227, 229, 230, 232, 233, 235, 237, 238, 239, 240, 242, 243, 244, 245, 246, 247, 251, 271
 Delaware - 63
 Denmark - 26, 27, 58, 59, 82, 140, 141, 203
 Egypt - 94, 126
 England - 24, 25, 26, 27, 28, 29, 30, 31, 33, 49, 51, 52, 53, 55, 57, 58, 59, 63, 70, 73, 75, 78, 80, 81, 82, 83, 88, 91, 95, 108, 109, 112, 115, 116, 119, 120, 121, 122, 123, 129, 137, 139, 140, 141, 142, 163, 165, 166, 170, 171, 172, 177, 179, 180, 187, 192, 197, 198, 201, 212, 215, 230, 234, 237, 238, 241, 250, 251, 252, 253, 254, 255, 257, 258, 259, 260, 262, 263, 264, 265, 266, 269, 271, 277
 Finland - 225
 Florida - 64, 80, 140
 France - 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 47, 48, 50, 51, 53, 54, 55, 56, 57, 63, 66, 67, 68, 69, 70, 71, 72, 75, 76, 77, 79, 80, 81, 84, 86, 87, 89, 90, 93, 94, 95, 96, 100, 104, 105, 106, 107, 108, 110, 111, 112, 117, 118, 121, 122, 124, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 143, 144, 145, 146, 147, 148, 150, 151, 152, 154, 155, 156, 157, 158, 159, 161, 162, 163, 164, 165, 166, 167, 169, 172, 173, 174, 175, 176, 179, 181, 182, 183, 184, 185, 186, 187, 188, 189, 191, 192, 193, 194, 195, 196, 198, 199, 201, 202, 203, 204, 206, 207, 208, 209, 210, 211, 212, 213, 215, 216, 217, 218, 219, 221, 222, 223, 224, 225, 226, 228, 229, 231, 232, 233, 234, 235, 236, 238, 239, 240, 243, 244, 245, 246, 247, 248, 251, 252, 254, 255, 256, 257, 258, 260, 261, 262, 264, 266, 267, 268, 269, 270, 276, 279, 280
 Georgia - 164
 Germany - 6, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 50, 51, 52, 53, 55, 56, 57, 60, 64, 65, 69, 70, 71, 72, 75, 77, 78, 79, 82, 87, 91, 92, 93, 95, 96, 99, 103, 104, 105, 106, 107, 110, 111, 112, 113, 115, 116, 118, 119, 120, 121, 122, 124, 125, 129, 132, 133, 136, 141, 142, 143, 144, 145, 147, 151, 152, 154, 155, 156, 158, 161, 164, 167, 170, 176, 177, 182, 184, 185, 186, 188, 189, 190, 197, 198, 201, 203, 205, 207, 209, 211, 212, 215, 217, 218, 223, 227, 232, 234, 238, 239, 248, 249, 251, 255, 256, 257, 262, 263, 264, 267, 268, 269, 270, 272, 277, 278, 279, 280
 Gibraltar → Spain
 Greece - 44, 45, 47, 49, 50, 52, 53, 56, 61, 62, 90, 94, 96, 97, 101, 107, 161, 162, 163, 164, 166, 189, 192, 196, 213, 214, 219, 220, 221, 229, 233, 235, 239
 Greenland - 140
 Holland - 49, 140, 188
 Hungary - 25, 40, 43, 44, 45, 47, 48, 49, 50, 51, 52, 53, 54, 55, 61, 68, 74, 86, 90, 98, 99, 101, 106, 110, 113, 116,

- 117, 118, 119, 120, 122, 123, 124, 127,
128, 129, 130, 133, 135, 137, 148, 149,
154, 155, 157, 161, 162, 163, 165, 166,
167, 168, 170, 174, 176, 177, 186, 187,
189, 190, 192, 196, 204, 205, 209, 211,
212, 214, 215, 216, 217, 219, 220, 221,
222, 227, 228, 229, 230, 232, 234, 235,
237, 239, 240, 241, 243, 244, 245, 247,
249, 263, 273, 275, 276
- Iceland - 139, 140
- India - 62
- Iran - 78, 96, 97
- Ireland - 139
- Israel - 161, 164, 177, 213
- Italy - 30, 31, 32, 37, 44, 45, 46, 47, 49, 51,
52, 53, 54, 55, 56, 57, 64, 73, 86, 88,
91, 92, 94, 120, 132, 139, 140, 155, 161,
162, 192, 204, 209, 210, 213, 219, 233,
247, 249, 250, 252, 261, 267, 275, 277
- Karachaevo-Cherkessia - 162, 219, 235, 236
- Kazakhstan - 81, 96, 97, 111, 112
- Kenya - 186
- Macedonia - 45
- Madagascar - 24
- Madeira - 139
- Malta - 55, 56, 91, 114, 155, 189, 210,
219
- Maryland - 63, 65, 81
- Massachusetts - 277
- Mauritius - 24
- Moldova - 43, 44, 48, 49, 55, 56, 61, 64,
68, 69, 72, 85, 89, 96, 97, 98, 99, 100,
111, 115, 117, 121, 138, 161, 162, 163,
164, 168, 170, 196, 252
- Monaco - 189
- Mongolia - 96, 97, 115, 174
- Montana - 58
- Morocco - 94, 141
- New Zealand - 24, 25, 84
- Nigeria - 81
- North Carolina - 63, 81
- Norway - 56, 139, 140
- Oregon - 104
- Pakistan - 62, 73, 96, 97
- Poland - 37, 41, 47, 49, 50, 51, 92, 135,
161, 166, 167, 168, 169, 212, 213, 214,
216, 217, 221, 237, 239, 243
- Portugal - 27, 37, 39, 43, 57, 79, 91, 139,
256
- Quebeck - 140
- Romania - 30, 34, 35, 37, 39, 41, 42, 47,
49, 50, 51, 55, 66, 67, 73, 80, 99, 101,
104, 105, 110, 116, 118, 119, 120, 121,
128, 129, 132, 137, 148, 149, 161, 164,
165, 166, 167, 173, 176, 190, 200, 212,
214, 215, 217, 221, 222, 227, 228, 229,
230, 232, 233, 234, 235, 238, 239, 240,
241, 242, 243, 244, 245, 246, 249, 263,
272, 274, 275, 279
- Russia (European part) - 31, 47, 49, 50,
56, 57, 61, 82, 127, 128, 140, 164
- Russia (Siberian part) - 96, 97
- Sardinia - 80, 81, 195, 210, 213, 249, 279
- Serbia - 37, 40
- Scotland - 57, 91, 139
- Shetland Islands - 81
- Slovakia - 40, 47, 48, 49, 53, 122, 155,
161, 167, 221, 227, 235, 239
- Slovenia - 37
- South Carolina - 81, 82
- South Dakota - 104, 110
- Spain - 33, 34, 37, 38, 39, 41, 42, 43, 44,
45, 47, 48, 49, 50, 51, 52, 53, 54, 55,
57, 72, 79, 85, 86, 92, 94, 98, 115, 116,
120, 121, 137, 139, 140, 154, 155, 156,
157, 159, 161, 162, 166, 174, 176, 177,
186, 188, 191, 193, 210, 215, 219, 222,
227, 232, 234, 235, 236, 237, 238, 242,
243, 245, 264
- Sweden - 26, 57, 139, 140, 269
- Switzerland - 6, 24, 29, 33, 35, 36, 38, 29,
40, 94, 226, 254, 266
- Syria - 165
- Texas - 104
- Thailand - 174
- Tunisia - 71, 73
- Turkey - 62, 157, 165, 166
- Ukraine - 31, 34, 38, 43, 44, 45, 47, 48,
49, 51, 53, 56, 61, 64, 72, 86, 87, 96,
97, 98, 99, 100, 115, 117, 120, 121, 127,
131, 138, 148, 149, 160, 161, 162, 163,
164, 165, 167, 176, 189, 191, 193, 212,
216, 217, 219, 220, 224, 228, 232, 239,
242, 245, 259, 273, 274, 279
- Uzbekistan - 81
- Virginia - 63, 81, 139, 253
- Wyoming - 58, 103, 141, 179, 271

Index of localities

The localities are listed alphabetically without respect to diacritical marks. Extralimital localities are not included. Pages with type localities are underlined.

- Abbey Wood - 28, 259
 Abri Lafaye → Lafaye
 Abri Suard → Suard
 Aegina - 45
 Agramunt - 33
 Ahnikov - 38
 Aigina → Aegina
 Aix → Aix-en-Provence
 Aix-en-Provence - 36, 148, 225
 Ajnácskő → Hajnáčka
 Aldeby - 49, 51, 139
 Aldène - 54, 166, 235
 Aljezar B - 44, 115, 154, 174, 210
 Allerding - 39
 Alliers - 34, 106
 Almenara 1 - 49, 94, 210
 Alto de Ballester - 39
 Amandoli - 45
 Ambrona - 53, 116
 Amôr 2 - 39
 Ancona - 42, 44, 261
 Andrașiu de Jos - 39
 Andreevka - 44
 Andrianò → Samos 1
 Annetorp → Limhamn
 Antointg - 34, 136
 Antwerp - 40, 41, 65, 125, 253, 256
 Anvers → Antwerp
 Apostolovo - 42, 43, 61
 Arago - 53, 232, 233, 235
 Archi - 55, 139
 Ardé - 47, 48, 163
 Ardes → Ardè
 Arene Candide - 56, 139
 Argentera - 30, 31, 277
 Aridos 1 - 54, 166
 Ariuşd - 37
 Armagnac - 36, 38, 83, 84
 Armaiolo - 49
 Armissan - 34, 35, 151, 153, 154
 Arquillo - 45
 Arquillo 3 - 47
 Arquillo de la Fontana → Arquillo
 Artenay - 39, 89
 Artesilla - 39
 Atapuerca - 54, 166
 Attenfeld - 41
 Aubrelog 1 - 33, 181, 208
 Aude - 27
 Augustine 7 - 38
 Aumeister - 42, 43, 124
 Aurillac - 34, 107
 Auzières → Aude
 Avenc de na Corna - 55, 91
 Azé 2 - 54, 166, 232
 Aveiras de Baixo - 43
 Bachów - 37
 Bacton - 51, 140, 192
 Balaruc 2 - 48, 210
 Balauzière - 55, 164
 Ballestar - 43
 Balša - 50
 Baltic amber → Jantarnyj
 Baltringen - 41
 Bamboli - 42, 45, 250
 Baraque → Basse Provence
 Baraval - 34
 Barranc de Binigaus → Binigaus
 Barros - 33
 Barton - 31, 271
 Basse Provence - 26, 28, 261, 262
 Batallones - 43
 Baume de Gonvillars → Gonvillars
 Bechlejovice - 34
 Bela Stena - 40
 Belgarite 4A - 33, 184, 199
 Belgarric - 34, 154, 183, 184, 206
 Belka - 44, 61, 162
 Belle Roche - 51
 Bellingen - 35, 277
 Benghisa - 55, 114, 189
 Beremend - 48, 50
 Beremend 1-3 - 49, 160, 161
 Beremend 4 - 49, 214
 Beremend 5 - 48, 169, 170, 204, 235
 Beremend 11 - 49, 161
 Beremend 15 - 48, 98, 101, 119, 129, 135, 161, 187, 221, 232, 244, 245
 Beremend 16 - 50, 99, 101, 119, 120, 129, 133, 135, 161, 165, 167, 176, 177, 215, 221, 227, 229, 230, 232, 239
 Beremend 17 - 50, 99, 119, 120, 128, 137, 155, 161, 165, 167, 168, 176, 177, 215, 217, 221, 228, 229, 230, 232, 234, 235, 237, 239, 245, 247
 Berru → Cernay-les-Reims
 Betfia 1 - 50, 167
 Betfia 2 - 50, 99, 119, 120, 121, 128, 148, 149, 159, 161, 165, 167, 176, 190, 200, 212, 214, 215, 217, 221, 222, 227, 229, 230, 234, 235, 238, 239, 240, 241, 242, 243, 244, 245, 246, 249
 Betfia 5 - 50, 51, 99, 129, 165, 166, 167, 221, 222, 228, 232, 233, 235, 249, 263
 Betfia 7 - 51, 119, 129, 165, 229, 240
 Betfia 10 - 51

- Betfia Aven - 51, 119, 120, 129
 Bezymjannaja balka → Malynivcy
 Biancone 1 - 45, 46, 155, 210, 219
 Biancone 2 - 45, 218
 Biehle - 54, 166, 221, 232
 Billy → Saint-Gérand-le-Puy
 Binigaus - 49, 210, 239
 Blomvåg - 56, 139
 Bobila Ordis - 49, 177
 Bognor Regis - 28, 59, 141, 142, 201,
260, 265
 Boom → Rupelmonde
 Bordu Mare - 55, 164
 Böttingen - 41
 Böttinger Marmor → Böttingen
 Bouce → Saint-Gérand-le-Puy
 Bouffie - 30, 31, 87, 144, 150, 152, 183,
 202, 258
 Bourgade - 51, 213
 Boussac 1 - 33, 199
 Boussac 2 - 33, 183, 199
 Bouxviller - 30
 Boxgrove - 51, 115, 116, 119, 120, 137,
 139, 165, 177, 215, 230, 237, 241
 Brajkovac → Peskovita
 Bransat → Saint-Gérand-le-Puy
 Braşov - 41
 Brasso → Braşov
 Bré 8 - 38
 Breitenberghöhle - 54, 56, 249
 Břešňany - 36, 38, 72, 101, 271
 Bretou - 30, 31, 143, 151, 183, 198, 202,
269, 270
 Brunn-Vösendorf → Vösendorf
 Brynseny → Brynzeny
 Brynzeny 1 - 54, 55, 168
 Büchelberg - 37, 105
 Budapest → Várhegy
 Budenheim - 37, 105, 106, 107
 Bujoru - 42, 43, 175
 Burg - 38
 Burggraben → Burg
 Burgmagerbein 3 - 34
 Burnham-on-Crouch - 28, 235, 263
 Burnt Wood - 33
 Burwell - 57, 69, 70
 Buteshty → Buteşti
 Buteşti - 55, 168
 Butešty → Buteşti
 Buzhory → Bujoru
 Bužory → Bujoru
 C-718 → Koněprusy C-718
 Cadillac - 33
 Ca Na Reia - 48, 49, 79, 161, 192
 Cagliari → Monte Reale
 Cala di Luna → Dorgali
 Calafra - 55
 Călugăr - 39
 Câmpu lui Neag - 35, 274
 Can Mas - 38, 156, 159
 Can Ponsic - 43
 Canet - 48, 49, 210
 Cantatore - 45, 155, 210
 Cap de la Biehle → Biehle
 Capins - 54
 Capo Figari → Figari
 Carrière - 54, 166, 169
 Casablanca 1 → Almenara 1
 Casots - 39
 Castelnuovo Berardenga Scalo - 47
 Castiglione 1 - 56, 214
 Castiglione 3 - 52, 53, 210, 214
 Cava Sud de Soave → Soave
 Cave 16 - 56, 166
 Cavillac → Cadillac
 Čebotarevka - 42, 44, 61, 64, 97, 120,
216, 220
 Cecilie I - 29, 30, 60, 255, 256
 Cecilie II - 30, 60
 Cengle → Basse Provence
 Čerdženica - 51, 241, 244, 245
 Céreste - 32, 33, 34, 133, 224, 252
 Čerevičnoe → Čebotarevka
 Čerevičnyj → Čebotarevka
 Cernay-les-Reims - 26, 27, 95, 96, 206,
255, 256, 260, 268
 Cerro Batallones → Batallones
 Cerro de los Espejos → Espejos Hill
 Cerro del Cristo del Otero → Otero
 Cerro del Otero → Otero
 Cerro Pelado → Layna
 Čertkov 2 - 53, 167
 Červený Hill - 51
 Červený vrch → Červený Hill
 Cestats → Léognan
 Cetăţuie - 32, 34, 133
 Cette → Sète
 Ceva - 36, 37, 252
 Cevico de la Torre - 37
 Chadžidimovo - 42, 44, 191
 Chalons-sur-Vesle → Cernay-les-Reims
 Chamblon - 29
 Chantegré → Saint-Gérand-le-Puy
 Chaptuzat - 35, 105, 106, 187
 Chauderon - 35
 Chavroches → Saint-Gérand-le-Puy
 Cherasco - 45
 Cherdzenitsa → Čerdženica
 Chilhac 2 - 49, 117
 Chilhac 3 - 48, 49, 117, 122
 Chillesford - 49, 51
 Chirò 2 - 45, 192, 210
 Chirò 3 - 45, 155, 210
 Chirò 4 - 45, 192, 210
 Chirò 5 - 46, 155, 210, 213
 Chirò 6 - 46, 155, 210

- Chirò 7 - 46, 210
 Chirò 9 - 46, 210
 Chirò 10 - 46, 192, 210
 Chirò 11 - 46, 155, 210
 Chirò 12 - 46, 210
 Chirò 14 - 46, 192, 210
 Chirò 15 - 46, 210
 Chirò 20 - 46, 155, 192, 210, 213
 Chirò 22 - 46, 210
 Chirò 24 - 46, 155, 192, 204, 210
 Chirò 27 - 46, 155, 210
 Chishmikioy → Çiřmikioi
 Chiřinãu - 42, 43, 64, 72, 85, 99, 111, 115, 117, 121, 138, 252
 Chlum 6 - 51, 161, 167
 Chrabãrsko - 42, 44, 72, 73, 228
 Cimay - 53, 191, 219, 235
 Çimiřlia - 44, 61
 Çimiřlija → Çimiřlia
 Ciobanița - 40, 42, 132
 Çiřmikioi - 48, 161
 Çiřmiřkioj → Çiřmikioi
 Ciuperceni - 47
 Climbach - 41
 Cluj–Cetatuie → Cetãtuia
 Cluj–Manastur - 29, 30, 66
 Cluj–Napoca → Cluj–Manastur
 Coderet - 36
 Collet–Redon - 42
 Colombiãre - 56, 164
 Combe Grenal - 55, 166
 Concud - 44
 Corbeddu - 55, 213
 Córcoles - 39, 156
 Corund - 44
 Coscia - 55, 189, 213
 Cosquer - 56, 139
 Cournon - 35, 105, 106
 Cova de Ca Na Reia → Ca Na Reia
 Cova de Canet → Canet
 Cova de Son Bauçã → Son Bauçã
 Crabit 1 - 34, 181, 203, 208
 Crau Coulet - 36
 Crayford - 55, 163
 Crediñta - 40, 42, 67, 80, 101, 105, 110, 118, 121, 137
 Crechy → Saint-Gérard-le-Puy
 Crevillente 2 - 43
 Cristo del Otero → Cerro del Otero
 Croix Rouge - 43
 Croydon - 26, 28, 95
 Csákvár - 42, 43, 90, 113, 217
 Csarnóta 1 - 47, 227, 234
 Csarnóta 2 - 46, 47, 127, 130, 161, 163, 166, 177, 186, 212, 216, 234, 235, 239, 243, 249
 Cuejdi - 36, 39, 274, 275
 Cuesta del Rey - 41
 Dechbetten - 40, 41, 71, 72, 111, 156
 Deljatin - 38, 273, 274
 Der Hohle Fels bei Happburg → Happburg
 Desse - 35, 134, 154, 155, 181, 183, 184, 194, 199, 208
 Deutsch-Altenburg 2C - 51, 161, 167, 227
 Deutsch-Altenburg 4B - 51, 161, 221, 227, 234, 237, 239, 241, 243, 246, 247
 Dévényújfalú → Devínská Nová Ves
 Devetařka Cave - 55, 166
 Devetařka peřtera → Devetařka Cave
 Devil's Tower - 55, 139
 Devínská Nová Ves - 40, 156
 Dolinskoe - 49, 61
 Dolnice - 36, 39, 63, 71, 85, 105, 106, 107, 110, 111, 116, 118, 123, 126, 127, 132, 135, 136, 137, 138, 144, 154, 155, 174, 191
 Dorgali - 55, 213
 Dorkovo - 45, 46, 117, 166
 Dormaal - 26
 Dürrloch - 55, 164
 Duruthy - 91
 Dyckerhoffsche Zementwerke → Wiesbaden 1
 East Runton - 51, 115, 120, 212
 Edeghem → Rupelmonde
 Egerkingen - 29, 254
 Ehrenstein - 57, 91
 El Arquillo → Arquillo
 El Fallol → Fallol
 Els Casots → Casots
 Emmenweid - 39
 Empoli - 46, 64
 Engi → Matt
 Engi-Matt → Matt
 Eppelsheim - 42, 43, 124, 264
 Erkertshofen 1 - 39
 Erpfingen - 51, 167
 Erquelinnes - 27
 Escamps - 30, 32, 144, 150, 152, 183, 217, 223, 225
 Escamps III - 32, 181, 199, 202, 225
 Escamps IV - 32, 181
 Escamps A - 32, 199
 Escamps C - 32, 181
 Escobosa de Calatañazor - 41
 Escorihuela - 48
 Espejos - 48
 Espenhain - 32, 34, 78, 136, 152, 217, 218, 268
 Estacion Imperial - 39
 Esterházy Cave → Csákvár
 Estramos → Osztramos
 Etterbeek - 29, 83
 Etulia - 48, 61, 98, 99, 100, 161, 163
 Etulija → Etulia

- Etulya → Etulia
 Eyzies - 54, 56, 90, 91
 Fage - 52, 55, 138, 161, 162, 165, 166, 167, 169, 213, 219, 221, 222, 231, 232, 235, 236
 Falcone 2 - 46, 210
 Fallol - 36, 38, 264
 Faluns de l'Orleanais → Orleanais
 Faluns de Saucats → Saucats
 Faluns de Suèvres → Orleanais
 Faluns de Touraine → Touraine
 Fellegvár → Cluj-Cetatuie
 Figari - 54, 55, 212, 213
 Figueira Brava - 55, 57, 91, 139
 Fina D - 46, 155, 210
 Fina E - 46
 Fina H - 46, 210
 Fischenbach - 37
 Flon Morand - 38
 Flörsheim - 36
 Fonbonne 1 - 33, 199, 208, 209, 226
 Fontcouverte → Aude
 Fontchévade - 55
 Forcalquier → Vachères
 Fournier Quarry - 55, 221
 Foxhall - 48, 49, 80
 Františkovy Lázně - 39, 105, 107
 Franzensbad → Františkovy Lázně
 Frauenweiler - 32, 33, 145
 Fraysse - 35, 134, 152, 154, 155, 183, 184, 194, 199, 208
 Frohnstetten - 33
 Froidefontaine - 32, 34, 79
 Fronstetten → Frohnstetten
 Fur - 26
 Gabbro - 42, 45, 247, 261
 Gaimersheim 1 - 35, 186
 Galié → Aude
 Gannat - 35, 36, 66, 105, 106, 136
 Gargano - 46, 155, 209, 210, 213
 Gargas - 56, 139
 Garouillas - 35, 199
 Garrigues - 31, 183, 199
 Geisel Valley → Geiseltal
 Geiseltal - 29, 64
 Geiseltal 32 → Geiseltal XXXV
 Geiseltal 36 → Geiseltal XXXVII
 Geiseltal 41 → Geiseltal XLI
 Geiseltal XIII - 28, 96, 185
 Geiseltal XIV - 28, 96, 185, 278
 Geiseltal XX - 30, 143
 Geiseltal XXII - 30, 143
 Geiseltal XXXV - 30, 96, 143
 Geiseltal XXXVI - 30, 142
 Geiseltal XXXVII - 30, 143
 Geiseltal XLI - 30, 96, 143
 Geiseltal L - 30, 201
 Geiseltal II - 28
 Gelinden - 26
 Gênova - 51, 239
 Gérce - 47
 Gergovia → Gergovie
 Gergovie - 35, 36, 106
 Gervaisio 1 - 47, 210
 Gervaisio 2 - 47, 155, 210
 Ghaqba - 55, 114, 189
 Ghar Dalam - 54, 55, 114, 155, 189
 Ghar Dalam cave → Ghar Dalam
 Gnien - 55, 91, 114
 Golbochika → Golboçica
 Golboçica - 42, 43, 68, 72, 89
 Golboçika → Golboçica
 Goldau - 36
 Goldberg - 40
 Gombasek - 53, 167, 221
 Gombaszök → Gombasek
 Gonvillars - 55, 164
 Gorham's Cave - 55, 139
 Gosport → Lee-on-Solent
 Götzendorf - 43, 73, 118
 Gouërris - 55, 91
 Gourdan - 57, 91, 164
 Gousnat - 32, 223
 Goyet - 54, 55, 189, 190
 Grafenmühle - 36, 38, 209
 Grand Baille - 57, 164
 Grange Farm - 27, 28, 255
 Grappin - 55, 163
 Grebeniki → Hrebenyky
 Grebenniki → Hrebenyky
 Grive-Saint-Alban - 40, 41, 42, 118, 131, 136, 137, 144, 145, 146, 148, 153, 154, 155, 156, 157, 158, 159, 174, 177, 192, 195, 204, 208, 209, 215, 262, 264
 Grive-Saint-Alban L-7 - 42
 Grive-Saint-Alban M - 42, 186, 191, 209
 Grotte - 47
 Grotta Romanelli → Romanelli
 Grotte de Dorgali → Dorgali
 Grotte de Gourdan → Gourdan
 Grotte de Macinaggio → Macinaggio
 Grotte des Gouërris → Gouërris
 Grotte des Harpons → Harpons
 Grotte du Lazaret → Lazaret
 Grotte Grappin → Grappin
 Grottes de Veyrier → Veyrier
 Gumbes B - 56, 214
 Gumbes C - 56, 214
 Gundersheim - 47
 Günzburg - 39
 Gurnard Bay - 33
 Gypses de Aix-en-Provence → Aix-en-Provence
 Gypses de Montmartre → Montmartre
 Hahnenberg - 40, 86, 87, 105
 Hajnáčka - 48, 122

- Hamstead - 32, 33, 258
 Happurg - 57, 164
 Harbartshofen - 36, 38, 251
 Harbatzhofen → Harbatshofen
 Harpons - 57, 91, 189
 Havighorst - 27
 Havighorst bei Reinbek → Havighorst
 Heggbach - 39
 Heiligenstadt - 37
 Heitenried 7 - 38
 Hermanstorp - 57, 139
 Herne Bay - 28
 Herrlingen 1 - 34
 Herrlingen 9 - 36
 Herrlingen/Blau → Herrlingen
 Heßler → Wiesbaden-Hessler
 Hessler → Wiesbaden-Hessler
 High Ongar - 27, 28, 58
 Highcliffe - 31, 271
 Higuera - 48
 Hludno - 37
 Hoeleden - 33
 Hohen Vielchen - 57, 91
 Hohler Fels bei Happburg → Happburg
 Holštejn 1 - 51, 167
 Hoogbutsel - 33
 Hordle - 30, 31, 63, 88, 103, 108, 172,
 187, 250, 254, 257, 258, 264, 277
 Hordwell → Hordle
 Hórvölgy - 55, 169, 222
 Hostalets de Pierola - 42, 155, 157, 159,
 193, 245
 Host'ovce 2 - 47
 Höwenegg - 43
 Hrabarsko → Chrãbarsko
 Hrebnyky - 42, 44, 61, 97, 115, 117
 Huéscar 1 - 51, 86, 119, 120, 121
 Hunas - 55, 169
 Hundsheim - 52, 53, 76, 99, 149, 167,
 168, 189, 205, 216, 221, 234
 Ilford - 55, 80, 91
 Il'inka - 44, 61, 232
 Ingelheim - 37
 Ipolytarnóc - 36, 38, 273, 275, 276
 Islas Medas → Meda Gran
 Isle of Sheppey → Sheppey
 Itardiès - 32, 34, 87, 152, 181, 183, 184,
 185, 199, 209, 226
 Ivánháza → Ivanovce
 Ivanócz → Ivanovce
 Ivanovce 1 - 47, 161, 227, 239
 Jantarnyj - 31
 Jaskinia Żabia → Żabia Cave
 Jur'evka → Jur'yvka
 Jur'yvka - 43
 Južen Park → Lozenec
 Juzhen Park → Lozenec
 Kadzielnia 1 - 50
 Kairy - 51, 64
 Kalafrana - 56, 114
 Kalfa - 43, 69, 138
 Kálmán Lambrecht Cave - 56, 189
 Kaltennordheim - 37
 Kamenskoe - 45, 116, 163
 Kardam - 40
 Kastel → Wiesbaden 1
 Kastel-Bruch → Wiesbaden 1
 Katharinenhof - 27
 Kerč' Peninsula 1 - 43
 Kerč' Peninsula 2 - 47, 64
 Kielniki 3B - 50, 169
 Kiev → Kyjiv
 Kiik-Koba - 56, 164
 Kirchberg - 39
 Kishinev → Chişinău
 Kişinev → Chişinău
 Kisláng - 48, 50, 61
 Klausenburg → Klausenburg
 Klein-Kems → Kleinkems
 Klein-Sorheim → Lierheim
 Kleinkems - 34
 Kohfidisch - 42, 43, 148, 177, 209
 Koliňany 2 - 49, 161
 Kolkotova Balka - 43, 97, 162
 Kolkotovskaja Balka → Kolkotova Balka
 Koloszmanostor → Cluj-Manastur
 Koněprusy C-718 - 51, 161, 216, 221,
 227, 235, 239
 Korond → Corund
 Köröshegy - 51, 263
 Kosov - 38
 Kotlovina - 48, 61, 100, 161
 Kövesvárad - 51, 205, 244
 Kozarnika - 57
 Kozi Grzbiet - 51, 167, 168, 221
 Kremnikovci - 45
 Kremnikovtsi → Kremnikovci
 Krivoj Rog → Krivoj Roh
 Krivoj Roh - 43
 Kronstadt → Braşov
 Krottensee → Mokřina
 Kruševica → Milićevo Brdo
 Kryžanovka 1 - 51, 99, 161, 165, 176
 Kryžanovka 2 - 51, 61
 Kryžanovka lower horizon → Kryžanovka
 1
 Kryžanovka upper horizon → Kryžanovka
 2
 Kryzhanovka → Kryžanovka
 Krzywcz → Nižnekryvčansk
 Kubanka - 43
 Kuchurgan → Vojničevo
 Kučurgan → Vojničevo
 Kujal'nickij liman → Kujal'nik
 Kujal'nik - 44, 61, 193
 Kujal'nik harbour → Kujal'nik

- Kunino - 51
 Kyjiv - 30, 31, 258, 259
 La Baraque → Basse Provence
 La Bouffie → Bouffie
 La Colombière → Colombière
 La Croix-Rouge → Croix-Rouse
 La Fontana → Arquillo
 La Grand-Baille → Grand-Baille
 La Grive-de-Saint-Alban → Grive-Saint-Alban
 La Mina → Layna
 La Plante → Plante
 La Puebla de Almoradiel → Puebla de Almoradiel
 La Puebla de Almoradier → Puebla de Almoradiel
 La Puebla de Almuradiel → Puebla de Almoradiel
 La Puebla de Valverde → Puebla de Valverde
 La Retama → Retama
 La Sauvetat → Sauvetat
 Labeur → Saint-Gérard-le-Puy
 Lafaye - 57, 164
 Laina → Layna
 Lammschlucht - 36
 Langenau - 39
 Langy → Saint-Gérard-le-Puy
 Las Higuieruelas → Higuieruelas
 Lausanne - 35
 Lavergne - 31, 150, 183, 185, 202, 203, 267
 Layna - 46, 47, 98, 210
 Lazaret - 52, 55, 162, 166, 215, 219, 232, 235, 236
 Le Bretou → Bretou
 Le Cengle → Basse-Provence
 Le Grotte → Grotte
 Le Lazaret → Lazaret
 Le Pin → Pin
 Le Vallonnet → Vallonnet
 Lee-on-Solent - 29, 171, 172, 180, 260, 262
 Léognan - 36, 38, 67, 79, 80
 Leonhard III - 31, 60
 Les Allets → Alliers
 Les Alletz → Alliers
 Les Alliers → Alliers
 Les Auzières → Aude
 Les Mauquiers → Basse-Provence
 Les Meilleries-Saint-Sauveur → Meilleries-Saint-Sauveur
 Les Montalets → Meudon
 Les Pradigues → Pradigues
 Les Roches → Roches
 Les Ussements → Ussements
 Les Valerots → Valerots
 Libros - 42, 43, 85
 Licata - 44
 Liedena - 33
 Lieffrens 5 - 38
 Lierheim - 40, 86, 87, 111, 125
 Liko - 54, 55, 189, 192, 213, 214
 Limberg - 39, 209
 Limhamn - 26, 27, 268, 269
 Lisboa - 37
 Lišov - 41, 105
 Lissieu - 31, 185
 Liveneck - 50, 61
 Los Barros → Barros
 Los Mansuetos → Mansuetos
 Los Valles de Fuentidueña → Valles de Fuentidueña
 Lostanges - 32, 183
 Lozenec - 46, 47, 114
 Lučești - 48, 163
 Lučešty → Lučești
 Lunel-Viel - 52, 55, 162, 166, 191, 192, 215, 219, 232
 Lupéron - 32, 33, 133
 Maar am Galtenbrunnen - 42
 Macinaggio - 55, 213
 Madelaine - 57, 91
 Mafflon 1 - 38
 Mainz-Weisenau → Weisenau-AS
 Malinivka → Malynivcy
 Malinovka → Malynivcy
 Malinowka → Malynivcy
 Malușteni–Berești - 46, 47, 99, 110, 232, 233
 Malynivcy - 42, 44, 60, 61
 Mansuetos - 42, 44, 137, 154
 Maret - 26
 Mariager Fjord - 27, 59
 Marnes de la Croix-Rousse → Croix-Rouse
 Marnes du Chenay → Cernay-les-Reims
 Mas de Got A - 34
 Mas de Got B - 34, 152, 183, 199
 Mas Rambault - 51, 165, 215, 219, 221, 229, 232, 238, 240, 243, 244, 245
 Massat - 54, 57, 236
 Mátraszőlös 1 - 40, 74, 122, 123, 176, 177
 Mátraszőlös 2 - 40, 68, 106, 110
 Matt - 32, 33, 226, 266
 Matveev Kurgan - 49, 61
 Mauquiers → Basse Provence
 Mauvières - 38
 Meda Gran - 49, 235, 236
 Meda Grande → Meda Gran
 Medvež'ja Cave - 54, 57, 127, 128
 Medvež'ja peščera → Medvež'ja Cave
 Megalo Emvolon - 47, 163
 Mège - 57, 164
 Méhesz → Včeláre 1
 Meilleries-Saint-Sauveur - 30

- Menat - 26
 Merkur - 36, 38, 71, 146, 154, 204, 211,
 215, 225
 Merkur-Nord → Merkur
 Merkur-North → Merkur
 Merkur-sever → Merkur
 Messel - 29, 60, 75, 87, 92, 93, 103, 141,
142, 143, 154, 170, 182, 184, 185, 197,
198, 203, 205, 207, 223, 257, 267, 268,
 270
 Mesvin - 26, 95
 Meudon - 26, 95
 Mihyska → Včeláre 1
 Milford - 30, 31
 Milićevo Brdo - 37
 Míngälar - 39
 Minišu de Sus - 41, 116
 Mnajdra - 55, 91, 114
 Mnajdra Gap → Mnajdra
 Möhren 6 - 32
 Möhren 13 - 34
 Möhren 19 - 33
 Moint Ceindre → Viex Collonges
 Mokřina - 39
 Molasse Coquilliere de l'Armagnac →
 Armagnac
 Mombach - 37
 Mondaino - 45
 Monsheim - 37, 105
 Mont Ceindre → Vieux Collonges
 Mont de Berru → Cernay-les-Reims
 Montaignu-le-Blin → Saint-Gérard-le-Puy
 Montchard → Cernay-les-Reims
 Monte Argentario - 51, 162, 219
 Monte Bamboli → Bamboli
 Monte Bolca → Vestena Nova
 Monte Duello → Monte Zuello
 Monte Granata - 47, 210
 Monte Peglia - 53
 Monte Reale - 52, 53, 54, 117, 195, 210,
232, 249
 Monte San Giovanni - 56, 213
 Monte Zuello - 29, 30, 88
 Montebamboli → Monte Bamboli
 Montecarlo - 48, 49, 119, 120, 232, 233
 Montereale → Monte Reale
 Monthelon - 28, 96
 Montmartre - 24, 30, 32, 75, 76, 144, 172,
173, 179, 180, 181, 183, 187, 224, 266,
267, 269
 Montmorency → Montmartre
 Montoussé 3 - 53, 168
 Montoussé 4 - 53
 Montoussé 5 - 50, 161, 233, 234, 238,
 239, 240
 Montpellier - 45
 Moreda - 48, 210
 Morin - 57, 91, 213
 Morozivka → Čebotarevka
 Morozovka → Čebotarevka
 Mosbach - 53, 189
 Mounayne - 34, 199
 Muntenia - 39
 Muselievo - 46, 47, 163, 169, 191, 222
 Mutigny - 28, 95
 Mytilini → Samos 1
 Nadvornaja - 38
 Nagyarsányhegy 1-4 - 50, 52, 86, 99,
 119, 120, 161, 167, 212, 214, 220, 221,
 227, 234, 247
 Nassiet - 34
 Nástup → Merkur
 Naumovka - 44
 Navarrete del Rio - 37
 Nazario 2 - 47, 210
 Nazario 3 - 47, 210
 Nazario 4 - 47, 210
 Nerja - 57, 139
 Neudorf → Devínská Nová Ves
 Neudorf an der March → Devínská Nová
 Ves
 Nichet - 54, 55
 Niel → Rupelmonde
 Nishnekrivchansk → Nižnekryvčansk
 Nižnekryvčansk - 56, 164
 Nižnij Strutyn' - 36, 38, 274
 Nördlinger Ries - 40, 105
 Nova Emetivka - 42, 45, 61, 98
 Novaja Emetovka → Nova Emetivka
 Novaja Kalfa → Kalfa
 Novaja Slobodka - 42, 44, 72, 87, 189
 Novo-Elisavetovka → Novoelisavetovka
 Novo-Elisavetovka → Novoelisavetovka
 Novo-Pokrovsk → Apostolovo
 Novoelisavetovka - 42, 44, 61, 148, 162
 Novoukrainka - 44, 61
 Oberdorf - 39
 Obitočnoe - 49, 149, 165
 Odesa - 44, 46, 72
 Odesa – catacombs - 47, 61, 72, 100, 131,
159, 161, 163, 217
 Odessa → Odesa
 Oehningen → Öhningen
 Oeningen → Öhningen
 Öhningen - 6, 24, 40, 41, 118, 124, 125,
 156
 Öningen → Öhningen
 Oppenheim - 37, 105
 Oppertshofen 2 - 33
 Orciano Pisano - 46, 47, 48, 73, 86, 140,
218, 219, 252
 Ořechov - 39
 Orford - 49, 80
 Orgnac 3 - 55, 162, 166, 215, 221, 232,
 235
 Orleanais - 36, 39, 69, 71, 72, 124

- Orp-le-Grand - 27
 Orrios 3 - 48
 Orsmaal → Dorsmaal
 Ostend - 52, 53, 170
 Osztramos 1 - 45, 46, 163, 249
 Osztramos 2 - 51, 99, 167, 192
 Osztramos 5 - 51, 165
 Osztramos 7 - 48, 160, 161, 166, 212, 214, 216
 Osztramos 8 - 51, 161, 167, 217
 Osztramos 9 - 45, 176
 Otero - 41
 Otero de Palencia → Otero
 Overstrand - 52, 73
 Pageyral - 57, 236
 Paglicci - 57, 139
 Paracuellos 3 - 42
 Paracuellos 5 - 41
 Passy - 27, 95
 Paulhiac - 37, 134
 Pécarel - 32, 223
 Pech Crabit → Crabit
 Pech Desse → Desse
 Pech du Fraysse → Fraysse
 Pedrera de Gènova → Gènova
 Pendo - 57, 139
 Peninsula de Setubal → Setubal
 Peralta de la Sal - 34
 Perdera de S'Onix → S'Onix
 Perignat - 35, 105, 106
 Perpignan - 46, 47, 117, 159, 161, 163, 232, 238
 Perrière - 30, 31, 144, 152, 183, 199, 203, 207, 208, 224
 Peschiera - 54, 57, 92
 Peshkovita - 40
 Peštera 16 → Cave 16
 Peštera Bordu Mare → Bordu Mare
 Petersbuch 2 - 39
 Petersbuch 4 - 39
 Petersbuch 6 - 42
 Petersbuch 10 - 42
 Petersbuch 18 - 42
 Petersbuch 28 - 39
 Petersbuch 38 - 39, 77
 Petersbuch 39 - 41, 154, 215
 Petit Puymoyen - 56, 164
 Petralona - 52, 53, 162, 166, 219, 220, 221, 229, 235, 239
 Petroverovka → Žovten'
 Peublanc - 37
 Phalip - 33, 208
 Phosphorites du Quercy → Quercy
 Piatra Neamț - 36, 39, 273
 Pietro Tampoia Cave - 54, 57, 80, 81, 213
 Pietro Tamponi Cave → Pietro Tampoia Cave
 Pikermi - 42, 45, 61, 90, 101, 107, 162
 Pin - 31
 Pirro 11 - 47, 155, 210, 213
 Piski → Simeria
 Pizzioli 4 - 47, 210
 Plante 2 - 34, 181, 199
 Podlesnoe → Malynivcy
 Podumci - 52, 161
 Pokrovka → Apostolovo
 Pokrovsk → Apostolovo
 Pokšešti - 43, 61
 Pokšešty → Pokšešti
 Polgárdi - 42, 45, 124
 Polgárdi 2 - 45, 230
 Polgárdi 4 - 129, 133, 162, 170, 174, 176, 234, 237, 239, 240, 241
 Polgárdi 5 - 45, 170, 209, 240, 244
 Pont du Château - 36, 105, 106
 Pontevès → Basse-Provence
 Posticchia 1 - 47, 210
 Pouàs - 57, 91
 Pradigues - 31, 202, 203
 Preschen → Břešťany
 Přezletice - 50, 52, 69, 73, 85, 86, 94, 115, 117, 119, 120, 121, 122, 124, 127, 128, 129, 137, 177, 178, 188, 219, 221, 230
 Primorsk - 51, 138, 149, 165, 239
 Primrose Hill - 27, 28, 266
 Prince - 56, 189
 Prisaca-Putna → Putna
 Przeworno 2 - 41, 156
 Puebla de Almoradiel - 48
 Puebla de Valverde - 49, 193, 234
 Puente Minero - 44
 Putna - 36, 39, 273, 274, 275
 Pymont-Challonges - 37, 105
 Quercy - 30, 31, 32, 68, 130, 134, 143, 144, 150, 151, 152, 153, 154, 173, 180, 181, 182, 183, 184, 185, 186, 194, 195, 198, 199, 201, 202, 203, 206, 207, 208, 217, 218, 223, 224, 264, 267, 269, 270
 Quibas - 51, 94, 155, 186, 188, 212, 215, 219, 227, 237, 238, 242, 243, 245
 Radoboj - 40, 41, 226
 Radoboy → Radoboj
 Raitenbuch 2 - 39
 Rambla de Valdecebro 2 → Arquillo
 Randecker Maar - 43
 Rapaci - 54, 56, 213
 Ravet-Lupovici - 33, 87, 152, 181, 208
 Ravolzhausen - 37, 71, 105, 106, 107, 110, 129
 Raymondén - 57, 139
 Razhishkata Cave → Ražiška Cave
 Ražiška Cave - 57, 166
 Ražiška peštera → Ražiška Cave
 Razvode - 52
 Reach - 54, 57, 69, 70
 Rebiełice Królewskie 1 - 48, 49, 92, 135,

- 160, 161, 166, 168, 169, 212, 239, 243
 Rębielice Królewskie 2 - 49, 161
 Regensburg → Dechbetten
 Reggbach - 38
 Rembielice Królewskie → Rębielice
 Królewskie
 Repolusthöhle - 54, 56, 190
 Retama - 39
 Rilly → Cernay-les-Reims
 Rinascita 1 - 47, 155, 210
 Rippersroda 1 - 50, 212
 Risitobel - 39
 Roches - 36
 Roddi - 45
 Romainville - 30, 33, 108, 251
 Romanelli - 57, 91, 139
 Ronzon - 32, 33, 253, 254, 270
 Roqueprune 2 - 34, 152, 181, 183, 184,
 194, 199, 254, 255, 260
 Rosières - 30, 32, 208, 223
 Rosières 1 - 30, 150, 152
 Rosieres 2 - 30, 150, 152, 181
 Røsnaes - 27, 28, 203
 Rothenstein - 39, 41
 Rott - 37
 Rubí → Fallol
 Rudabanya - 42, 43, 118, 154, 155, 157,
 176, 211, 234, 240
 Rumst → Rupelmonde
 Rungurskaja Sloboda → Sloboda
 Rupelmonde - 32, 34, 79, 112, 250, 251,
 268, 271
 Sables d'Orp-le-Grand → Maret
 Sables de Bruxelles → Etterbeek
 Săcădat - 42
 Sackdilling - 52, 161, 167
 Saignon - 32, 33, 275, 276
 Saint-Antonin → Basse-Provence
 Saint-Estève-Janson - 52, 53, 161, 162,
 166, 190, 191, 192, 205, 212, 213, 216,
 219, 231, 232, 234, 235
 Saint-Gérard-le-Puy - 36, 37, 63, 67, 68,
 70, 71, 77, 80, 86, 89, 93, 100, 104,
 105, 106, 107, 109, 110, 111, 112, 130,
 131, 132, 134, 136, 137, 138, 145, 146,
 150, 153, 154, 155, 174, 175, 187, 193,
 194, 195, 204, 208, 209, 210, 211, 225,
 244, 246, 248, 258
 Saint James Park - 28, 59
 Saint-Julien-le-Montagnié → Basse-
 Provence
 Saint-Maurin → Basse Provence
 Saint-Sol-Belcastel - 55, 191, 233
 Saint-Vallier - 49, 163
 Sainte-Néboule - 30, 32, 107, 150, 152,
 202
 Salême - 31, 183, 207
 Salzgitter-Lebenstedt - 54, 57, 190
 Samos → Samos 1
 Samos 1 - 42, 44, 62, 97
 San Angelo - 36, 37
 San Giovannino - 46, 47, 155, 192, 210,
 213, 219
 Šandalja 1 - 49, 161, 220, 239
 Sandelzhausen - 41
 Sankt Margarethen - 40, 41, 64
 Sansan - 40, 41, 89, 94, 111, 118, 121,
 130, 146, 147, 154, 155, 156, 158, 175,
 176, 188, 191, 193, 218, 230, 257
 Sant'Andrea - 49
 Sarrión → Espejos Hill
 Saucats - 36, 38, 63
 Saulcet → Saint-Gérard-le-Puy
 Sauvetat - 35, 106
 Scaparoni - 45
 Schemmersberg - 40
 Schernfeld - 49
 Schitu Frumneasă - 39, 273
 Schnaitheim → Steinheim
 Schönweg - 40
 Schusterlucke - 54, 56, 164, 168, 169
 Ščigry - 34
 Selk - 26, 269
 Semenovka - 47, 61
 Senèze - 50, 163, 167
 Senftenberg - 38
 Senigallia - 42, 45, 132, 247, 249, 261,
 267
 Serrat-d'en-Vaquer → Perpignan
 Serre de Trigodinna → Aude
 Sète - 47, 210
 Setubal - 37
 Seveckenberg - 54, 56, 99, 188, 231, 232,
 248, 277
 Sheppey - 25, 28, 58, 59, 64, 65, 78, 81,
 82, 83, 171, 179, 201, 251, 252, 259,
 263, 265, 266, 269, 277
 Sieblos - 32, 33, 262
 Sieblos über Fulda → Sieblos
 Sierra de Quibas → Quibas
 Sillans-la-Cascade → Basse-Provence
 Silverinha - 27, 28, 256
 Sima del Elefante - 52, 121, 166, 188,
 191, 219, 222, 232, 245
 Simeria - 37
 Simonelli - 56, 189
 Sinigaglia → Senigallia
 Skalice - 32, 34, 250, 251
 Skalitz → Skalice
 Skiritz → Skřice
 Škodova Gora → Škodova Hill
 Škodova Hill - 44, 72
 Skyřice - 36, 38, 85, 110, 111, 155
 Slivnica - 48, 50, 94, 121, 161, 222, 228,
 233, 234, 236, 239, 247
 Slivnitsa → Slivnica

- Sloboda - 38
 Slobodka → Novaja Slobodka
 Snigirevka - 44, 61
 Soave - 52, 161, 233
 Sodbach-Heitenried → Heitenried
 Sofia → Sofija
 Sofija 1 - 44
 Sofija 2 - 47
 Sokolov (Czechia) - 39
 Sokolov (Ukraine) - 43, 86, 119, 193
 Sokolov sand pit → Sokolov (Ukraine)
 Sokolovskij peščannyj kar'er → Sokolov (Ukraine)
 Solnhofen - 38
 Somerset - 57, 91
 Somssich Hill 2 → Somssichhegy 2
 Somssichhegy 2 - 52, 116, 148, 155, 167, 221, 227, 228
 Son Bauçà - 54, 210
 Son Bauzà → Son Bauçà
 S'Onix - 50, 115, 122, 155, 176, 210, 212, 228, 233, 235, 236, 237, 239, 241, 242, 245, 246
 Soulabé - 56, 57, 164, 189
 Sozopol - 54, 57, 178
 Sozopol Bay → Sozopol
 Spinagallo - 53, 94
 Spitzberg → Steinberg
 St. Gerand-le-Puy → Saint-Gérard-le-Puy
 St. James Park → Saint James Park
 St. Margarethen → Sankt Margarethen
 Starye Duruitory - 56, 164, 168
 Staryje Duruitory → Starye Duruitory
 Steendorp → Rupelmonde
 Stein → Öhningen
 Stein am Rhein → Öhningen
 Steinberg - 40, 41, 99, 105, 115, 132, 133, 147
 Steinheim - 40, 41, 64, 106, 107, 110, 118, 156
 Steinheim am Albuch → Steinheim
 Steyn → Öhningen
 Storzigen - 41
 Stránská skála - 50, 52, 53
 Stránská skála – Absolon's 2nd cave - 50, 52, 117, 120, 123, 127, 128, 137, 155, 161, 166, 167, 169, 191, 216, 220, 221, 222, 233, 234, 239
 Stránská skála – Čapek's sites 1-3 - 53, 120, 135, 166, 167, 169, 205, 217, 222, 233, 245
 Stránská skála – Čapek's site 4 - 52, 120, 167, 169, 177, 205, 220, 221
 Stránská skála – Čapek's site 5 - 52, 86, 99, 119, 120, 121, 122, 127, 128, 129, 135, 138, 149, 155, 166, 167, 169, 176, 187, 191, 205, 217, 220, 221, 233, 234, 238, 239, 246, 247
 Stránská skála – Čapek's site 8 - 53, 167, 239, 246
 Stránská skála – Čapek's site 9 - 53, 86, 99, 116, 119, 120, 122, 148, 149, 155, 166, 167, 169, 177, 187, 188, 200, 205, 216, 220, 221, 222, 227, 229, 233, 237, 238, 239, 242, 243, 244, 245, 246, 247
 Stránská skála – Musil's talus cone - 50, 53, 86, 99, 116, 119, 120, 122, 123, 127, 128, 129, 135, 137, 148, 149, 155, 161, 166, 167, 168, 176, 177, 178, 187, 205, 215, 217, 220, 221, 222, 227, 232, 233, 234, 235, 239, 240, 244, 246
 Strathclyde - 57, 91
 Strette - 49, 120
 Stshigry → Sčigry
 Stubersheim - 38
 Suard - 55, 163, 166
 Suèvres → Orleanais
 Suchomasty 3 - 43
 Sulzigobel - 40
 Sümeg - 42, 44, 154, 205
 Sundby - 26
 Sungir - 56, 164
 Suvorovskoe - 49, 61
 Szakadát → Săcădat
 Ta'Kandja - 56, 114, 189
 Tal Ghaqba → Ghaqba
 Tal Gnien → Gnien
 Tal Herba → Herba
 Tanne → Kaltennordheim
 Tarchankut - 51, 127, 149, 224, 228, 242, 245
 Tardosbanya - 44
 Tarkö - 52, 53, 149, 167, 168, 205, 215, 216, 221, 232
 Tatarești - 49, 61
 Tatarești → Tatarești
 Tataros → Tataruș-Brusturi
 Tataruș-Brusturi - 42, 43, 73
 Tatinja Draga - 53
 Tegelen - 49, 188
 Tegelen Clay → Tegelen
 Temnata Cave - 57, 166
 Temnata doubka → Temnata Cave
 Temnata dupka → Temnata Cave
 Templomhegy → Villány 3
 Terhaegen → Rupelmode
 Terra Amata - 215
 Thalberggraben - 35
 Thalberg-Graben → Thalberggraben
 Thalheim → Brașov
 Titov Veles → Veles
 Tjaginka - 43
 Tomerdingen - 35
 Toril 3 - 42
 Tornewton - 52, 55, 165, 166
 Touraine - 36, 38, 156

- Tourkobounia 1 - 49, 161
 Tourkobounia 2 - 50, 233
 Tourkobounia 5 - 53
 Traun/Pucking - 35, 37, 139
 Trebovlje - 37
 Treteau-Verneau → Saint-Gérard-le-Puy
 Treuchtlingen 2 - 38
 Trinka 1 - 54, 56, 164, 168
 Trinki → Trinka
 Trois Frères - 57, 164
 Trojanovo - 42, 44, 116
 Tschischmikioi → Çiřmikioi
 Tuchořice - 38, 193
 Ulm–Westtangente - 38
 Untermassfeld - 51, 115, 116, 161, 188,
 232, 234, 239
 Uppony 1 - 53, 205
 Üromhegy - 53
 Urtiaga - 57
 Ussemets - 38
 Vachères - 34
 Val d'Arno → Montecarlo
 Val Mala - 45
 Valdarno → Montecarlo
 Valdecebro 5 - 44
 Valerots - 51, 169, 221, 229, 232, 239
 Valles de Fuentidueña - 43, 72
 Vallonnet - 51, 192, 239
 Várgegy – Fortuna Street 25 - 52, 53, 116,
 166, 167, 263
 Várhegy – Uri Street 72 - 52, 54
 Varnica → Varnița
 Varnița - 42, 43, 61
 Varnitsa → Varnița
 Varnsdorf - 32, 34, 70
 Väršec - 48, 49, 98, 130, 160, 161, 165,
 167, 176, 177, 189, 190, 191, 192, 193,
 204, 215, 220, 222, 228, 229, 230, 233,
 235, 236, 238, 239, 242, 244, 245, 246,
247, 249
 Varshets → Väršec
 Vaumas → Saint-Gérard-le-Puy
 Včeláre 1 - 50, 53, 166, 167, 221, 235
 Včeláre 3 - 49
 Včeláre 4E - 53, 155, 161, 167, 227
 Veles - 45
 Venta del Moro - 45
 Venta Micena - 53
 Verrua Savoia - 30, 31, 275
 Vértesszöllös 2 - 54, 166, 167, 189, 212,
 234
 Vestena Nova - 30, 32, 261
 Vestenanova → Vestena Nova
 Veyrier - 57, 164
 Vieulac - 57, 236
 Vieux de Collonges → Vieux Collonges
 Vieux-Collonges → Vieux Collonges
 Vieux Collonges - 39, 40, 135, 154, 156,
 186, 212, 228
 Viladecavalls - 43, 193
 Villány - 48
 Villány 3 - 50, 99, 117, 149, 160, 161,
 167, 210, 214, 216, 227, 234
 Villány 6 - 53
 Villány 8 - 53
 Villány-Kalkberg → Villány 3
 Villánu-Nord → Villány 3
 Vinalmont - 27
 Vindija - 56, 164
 Voigtstedt - 53, 69, 92, 115, 116, 119,
 120, 121, 122, 129, 167, 176, 177, 188,
 234, 238, 263
 Voiničevo → Vojničevo
 Vojničevo - 45, 61, 99, 160, 163, 212
 Vorochta - 34
 Vösendorf - 42, 43, 65
 Vřivo → Apostolovo
 Vyřiva → Apostolovo
 Waes → Rupelmonde
 Walbeck - 26, 95
 Waldhalde - 38
 Wallerstein - 41, 125
 Walton-on-the-Naze - 27, 28, 59, 108,
 142, 197, 198, 262, 263, 271
 Warden Point → Sheppey
 Warnsdorf → Varnsdorf
 Weingraben - 40
 Weisnau - 36, 77
 Weisnau-AS - 36, 37, 71, 79, 105, 110,
 132
 Weissenau → Weissenau-AS
 Weissenburg 6 - 38
 Weissenburg 9 - 33
 Weissenburg in Bayern → Weissenburg
 West Ronton – Upper Forest Bed - 52, 53,
 73, 115, 116, 118, 119, 120, 121, 122,
 123, 129
 Westbury-sub-Mendip - 55, 169
 Westeregeln - 54, 56, 271, 272
 Weže 1 - 47, 160, 161, 166, 177, 230,
 234, 238
 Wiesbaden 1 - 36, 38, 105, 106, 169
 Wiesbaden-Heßler → Wiesbaden-Hessler
 Wiesbaden-Hessler - 36, 38, 105, 106,
112
 Wiesbaden-Kastel → Wiesbaden 1
 Wintershof (West) → Wintershof-West
 Wintershof-West - 36, 38, 77, 144, 147,
151, 154, 155, 158, 211, 215, 217, 227
 Yarmouth - 32, 33, 109, 112, 251, 252,
260, 265
 Yateley - 29, 30, 259
 Žabia Cave - 51, 167
 Žabia skala → Žabia Cave
 Zalesiaki 1A - 51, 167
 Zebbug - 54, 56, 91, 144, 189

Zebbug cave → Zebbug
Zelená - 39
Zell - 37
Ževachova Hill - 51, 165, 191

Zhevakhova gora → Ževachova Hill
Žovten' - 44, 61, 162
Zwergloch - 56, 164

Index of collections

For the explanation of abbreviations see pp. 21-24.

- Aix-en-Provence (MHNA) - 261, 262
 Baku (MB) - 91
 Bâle → Basel
 Basel (BaM) - 96, 108, 194, 208, 254
 Berlin (NKMB) - 78, 136, 152, 154, 217
 Berlin (ZGIB) - 264
 Bologna (MGPUB) - 277
 Bordeaux (MHNB) - 79, 80, 154, 194
 Braunschweig (LMB) - 190
 Brno (MMB) - 123, 214, 220, 221, 233
 Brünn → Brno
 Bruxelles (IRScNB) - 79, 112, 125, 182, 189, 250, 251, 253, 256, 268, 271
 Bucharest → București
 București (IGGB)
 București (LPUB) - 64, 67, 80, 101, 105, 132, 232
 Budapest (GIB) - 61, 73, 99, 113, 149, 159, 160, 166, 169, 170, 173, 174, 176, 200, 204, 205, 209, 217, 249, 273, 275
 Budapest (NMB) - 101, 117, 124, 127, 130, 167, 168, 186, 205, 214, 215
 Cambridge (SMC) - 69, 277
 Chilhac (MPC)
 Chișinău (GIKM) - 68, 72, 89
 Chișinău (IZC) - 168
 Darmstadt (HLMD) - 257
 Firenze (IGF) - 86, 119, 140, 232
 Florence → Firenze
 Frankfurt am Main (SMF) - 71, 75, 87, 92, 103, 112, 133, 141, 142, 143, 156, 170, 198, 207, 267, 268
 Graz (MG) - 190
 Halle an der Saale (GM) - 60, 143, 201, 255, 256, 278
 Hamburg (GIH) - 93
 Iași (LPUI) - 85, 99, 252
 Imola (MCI) - 132, 247, 249, 261, 267
 Ipolytarnóc (CHI) - 276
 Ipswich (IM) - 80
 Jerusalem (HUJ) - 160, 163, 177
 Karlsruhe (SMNK) - 75, 203, 205, 223, 252, 257
 Kiev → Kyjiv
 Kišinev → Chișinău
 Kraków (ISEZ) - 160, 166, 168, 216
 Kyjiv (IZAN) - 64, 97, 98, 100, 117, 148, 159, 169, 220
 Kyjiv (GMK) - 61
 Leningrad → Sankt Peterburg
 Leyden (RGM) - 192, 204, 213, 219
 Liège (ULg) - 250, 251, 256, 268
 Lisboa (UNL) - 256
 London (BMNH) - 24, 58, 59, 63, 64, 68, 78, 81, 82, 83, 86, 88, 91, 93, 97, 99, 100, 103, 108, 109, 110, 112, 118, 122, 124, 125, 131, 140, 141, 153, 165, 170, 171, 172, 179, 184, 189, 197, 201, 203, 209, 219, 247, 250, 251, 252, 253, 254, 255, 258, 259, 260, 261, 263, 264, 265, 266, 269, 271, 277
 London (IGS) - 82, 95, 257, 271
 L'vyv (IGGL) - 273, 274
 Lyon (FSL) - 112, 124, 130, 132, 133, 144, 147, 150, 155, 159, 219, 222, 231, 232
 Lyon (ML) - 131, 152, 154, 158, 159, 184, 185, 195, 196, 204, 206, 215, 264, 269
 Lyon (coll. Bonifay): 161, 190, 191, 192, 212, 213, 215, 219, 231
 Madrid (MNCN) - 98
 Marseille (FSUM) - 194
 Montauban (MNHM)
 Montpellier (USTL) - 87, 107, 144, 184, 198, 207, 225, 226, 258, 261, 269
 Moscow → Moskva
 Moskva (PIN) - 72, 98, 99, 115, 117
 Munich → München
 München (BSP) - 77, 124, 130, 132, 133, 144, 147, 150, 151, 155, 158, 194, 198, 202, 211, 215, 224, 249, 269, 271
 Novorosijsk (GUN) - 61
 Oradea (RMO) - 165, 232, 235, 249, 263
 Padova (MPUP) - 88, 261
 Padua → Padova
 Paris (MNHN) - 63, 66, 68, 70, 71, 75, 80, 83, 85, 89, 90, 93, 94, 95, 96, 97, 99, 101, 104, 105, 106, 109, 110, 111, 118, 121, 130, 131, 132, 134, 136, 143, 144, 145, 146, 147, 150, 151, 152, 153, 154, 156, 158, 162, 163, 172, 173, 174, 175, 176, 180, 181, 182, 183, 185, 186, 187, 188, 193, 194, 195, 196, 198, 199, 201, 202, 204, 206, 208, 209, 217, 218, 223, 224, 225, 230, 231, 236, 248, 251, 255, 256, 257, 258, 260, 264, 266, 267, 268, 269, 270, 277
 Petralona (MP) - 196
 Pisa (MSNUP) - 261
 Pittsburg (CMP) - 154
 Prague → Praha
 Praha (DPFNSP) - 63, 85, 123, 126, 174
 Praha (NMP) - 70, 250
 Praha (coll. Fejfar) - 115, 126, 132, 146, 188, 211
 Sabadell (IPS) - 264
 Sankt Peterburg (ZIN) - 127, 128, 258

- Sartène (MS) - 214
Sofia → Sofija
Sofija (MNHNS) - 72, 94, 114, 116, 117,
130, 160, 166, 167, 177, 178, 191, 220,
246, 247, 249
Stockholm (NRM) - 269
Stuttgart (SMNS) - 86, 113, 145, 156, 198
Teplice (RMT) - 101
Thermopolis (WDC) - 142
Tiraspol (TGPI) - 61, 175
Torino (MGPUT) - 212, 250, 252
Turin → Torino
Utrecht (UUT) - 192, 202, 209, 213
Valletta (NMM) - 114, 155
Vienna → Wien
Washington (USNM) - 65, 81, 112, 156
- Weimar (IQW) - 122
Weißenburg (coll. Rummel): 41, 209
Wien (UWPI) - 76, 149, 205, 221
Wien (NHMW) - 64, 65, 66, 139, 168,
180, 203, 226
Würzburg (GPIW) - 262
Zürich (NMZ) - 24, 226
- coll. Croizet - 106
coll. Keferstein - 271
coll. Kessler - 103
coll. Maschwitz - 185
coll. Michel - 79
coll. Paupe - 79
coll. Villalta - 72, 154

Index of stratigraphical units

Only European localities are considered here. The 'Quercy', spanning MP 16-28, is not included here (see p. 382 for its occurrence). This index is arranged stratigraphically.

- MQ 2E (Holocene) - 54, 57, 70, 79, 80, 81, 91, 94, 139, 140, 164, 178, 249
- MQ 2C-D (late Pleistocene) - 54, 55, 56, 57, 79, 91, 114, 139, 155, 163, 164, 166, 168, 169, 188, 189, 190, 192, 195, 210, 213, 214, 219, 232, 236, 248, 249, 272, 277
- MQ 2A-B (middle Pleistocene) - 52, 53, 54, 55, 76, 99, 116, 138, 149, 162, 163, 166, 167, 168, 169, 189, 191, 192, 195, 196, 205, 210, 212, 213, 214, 215, 216, 222, 232, 236, 249, 263
- MQ 1 (early Pleistocene) - 50, 51, 52, 53, 64, 69, 73, 85, 86, 92, 94, 99, 101, 115, 116, 117, 119, 120, 121, 122, 123, 127, 128, 129, 133, 135, 137, 139, 140, 148, 149, 155, 161, 165, 166, 167, 168, 169, 176, 177, 178, 186, 187, 188, 190, 191, 192, 196, 100, 205, 212, 213, 214, 215, 216, 217, 219, 220, 221, 222, 224, 227, 228, 229, 230, 232, 233, 234, 235, 237, 238, 239, 240, 241, 242, 244, 245, 246, 247, 249, 263
- MN 16-18 (late Pliocene) - 47, 48, 49, 50, 61, 73, 80, 86, 92, 94, 98, 99, 100, 115, 117, 119, 120, 121, 122, 129, 130, 135, 139, 140, 149, 155, 161, 163, 165, 166, 167, 169, 170, 176, 177, 187, 188, 191, 192, 193, 204, 210, 212, 213, 214, 215, 216, 217, 219, 220, 221, 222, 228, 229, 230, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 249, 252
- MN 14-15 (early Pliocene) - 44, 45, 46, 47, 61, 64, 72, 73, 86, 98, 99, 100, 101, 108, 114, 116, 117, 127, 130, 131, 140, 155, 161, 163, 165, 166, 169, 176, 177, 186, 191, 192, 204, 209, 210, 212, 213, 216, 217, 219, 222, 227, 232, 233, 235, 238, 239, 243, 249, 252
- MN 11-13 (late Miocene, Turolian) - 43, 44, 45, 61, 62, 64, 72, 73, 87, 90, 97, 98, 101, 107, 115, 116, 117, 120, 124, 128, 129, 132, 133, 148, 154, 162, 170, 174, 189, 191, 193, 196, 205, 209, 210, 216, 220, 228, 230, 232, 234, 237, 239, 240, 241, 244, 247, 249, 250, 261, 267
- MN 09-10 (late Miocene, Vallesian) - 40, 41, 42, 43, 61, 64, 65, 68, 71, 72, 73, 85, 86, 89, 97, 99, 111, 113, 115, 117, 118, 119, 121, 124, 138, 146, 148, 154, 155, 157, 159, 162, 175, 176, 193, 196, 209, 211, 217, 234, 240, 245, 264
- MN 05-08 (middle Miocene) - 24, 39, 40, 64, 65, 67, 68, 72, 80, 99, 101, 105, 106, 107, 110, 111, 113, 114, 118, 121, 122, 123, 124, 130, 131, 132, 133, 136, 137, 145, 146, 147, 148, 154, 155, 156, 158, 159, 174, 175, 176, 177, 186, 188, 191, 192, 193, 195, 196, 208, 209, 212, 215, 218, 226, 228, 231, 245, 253, 256, 257, 262, 264
- MN 01-04 (early Miocene) - 37, 38, 39, 40, 63, 67, 68, 69, 71, 72, 76, 77, 79, 80, 84, 89, 93, 100, 101, 105, 106, 107, 110, 111, 112, 116, 118, 124, 126, 127, 129, 130, 131, 132, 134, 136, 137, 138, 144, 145, 146, 147, 150, 151, 154, 155, 156, 158, 159, 174, 175, 187, 191, 193, 194, 195, 196, 204, 208, 209, 211, 215, 217, 225, 246, 248, 251, 252, 258, 264, 271, 273, 274, 275, 276
- MP 25-30 (late Oligocene) - 34, 35, 36, 37, 66, 105, 106, 134, 136, 139, 151, 154, 155, 184, 186, 187, 194, 199, 206, 208, 225, 277
- MP 23-24 (middle Oligocene) - 32, 34, 70, 79, 87, 109, 112, 133, 136, 145, 152, 173, 181, 183, 184, 185, 194, 199, 203, 208, 209, 218, 224, 226, 250, 251, 252, 258, 260, 262, 265, 266, 268, 271
- MP 21-22 (early Oligocene) - 32, 33, 34, 87, 109, 112, 133, 152, 181, 183, 190, 208, 226, 252, 254, 255, 258, 260, 262, 265, 266, 270, 276
- MP 17-20 (late Eocene, Priabonian) - 31, 32, 33, 63, 76, 88, 103, 107, 108, 144, 150, 152, 172, 173, 179, 181, 183, 187, 199, 202, 203, 207, 217, 223, 224, 250, 251, 254, 257, 258, 259, 261, 264, 266, 267, 269, 275, 277
- MP 14-16 (late Eocene, Bartonian) - 31, 143, 150, 183, 185, 198, 202, 203, 267, 270, 271
- MP 11-13 (middle Eocene) - 28, 29, 30, 31, 60, 64, 66, 75, 83, 87, 88, 93, 96, 103, 141, 142, 143, 170, 171, 172, 180, 182, 184, 185, 190, 201, 203, 205, 207, 223, 254, 255, 256, 257, 259, 267, 268, 277
- MP 07-10 (early Eocene) - 26, 27, 28, 58, 59, 65, 78, 82, 83, 95, 96, 108, 141, 142, 171, 179, 197, 198, 201, 203, 251, 252, 253, 255, 256, 259, 260, 261, 262, 263, 266, 269, 271, 277
- MP 01-06 (Paleocene) - 26, 27, 95, 96, 206, 256, 260, 268, 269

Index of family-group names

All names are listed with the ending -idae, without respect to taxonomic rank. Incorrect spellings are not considered. Only home-pages are listed.

- | | | |
|--------------------------|--------------------------|---------------------------|
| Accipitridae - 187 | Gastornithidae - 94 | Phasianidae - 151 |
| Aegialornithidae - 200 | Gaviidae - 63 | Phoenicopteridae - 103 |
| Aegithalidae - 242 | Geranopteridae - 223 | Phoeniculidae - 77 |
| Agnopteridae - 178 | Glareolidae - 132 | Phorusrhacidae - 178 |
| Alaudidae - 228 | Gracilatarsidae - 257 | Picidae - 148 |
| Alcedinidae - 225 | Gruidae - 88 | Podargidae - 205 |
| Alcidae - 138 | Gruipedidae - 275 | Podicipedidae - 85 |
| Anatidae - 109 | Gryzajidae - 98 | Preficidae - 197 |
| Anatipedidae - 273 | Hemiprocnidae - 200 | Primapidae - 200 |
| Anhimidae - 107 | Hermosornithidae - 178 | Primosciniidae - 141 |
| Anhingidae - 73 | Hirundinidae - 227 | Procellariidae - 78 |
| Anseranatidae - 107 | Horusornithidae - 258 | Prophaethontidae - 64 |
| Apodidae - 203 | Idiornithidae - 179 | Prunellidae - 241 |
| Apopenpsidae - 186 | Jacanidae - 126 | Pseudasturidae - 141 |
| Ardeidae - 67 | Juncitarsidae - 103 | Pseudodontornithidae - 81 |
| Ardeipedidae - 273 | Jungornithidae - 200 | Pseudosulidae - 66 |
| Archaeotrogonidae - 197 | Laniidae - 229 | Psilopteridae - 178 |
| Bathornithinae - 178 | Laridae - 136 | Psittacidae - 146 |
| Bombycillidae - 229 | Laurillardidae - 75 | Pterocletidae - 133 |
| Brontornithidae - 178 | Lithornithidae - 178 | Quercymegapodiidae - 150 |
| Capitonidae - 147 | Marinavidae - 259 | Quercypsittidae - 141 |
| Capitonididae - 147 | Megapodiidae - 150 | Rallidae - 171 |
| Caprimulgidae - 197 | Meropidae - 226 | Ramphastidae - 147 |
| Cariamidae - 178 | Mesembriornithidae - 178 | Recurvirostridae - 107 |
| Cathartidae - 101 | Messelirrisoridae - 75 | Remiornithidae - 268 |
| Certhiidae - 243 | Messelornithidae - 87 | Romainvilliidae - 107 |
| Charadriidae - 135 | Miopiconidae - 141 | Rostratulidae - 126 |
| Charadriipedidae - 274 | Momotidae - 225 | Sagittariidae - 187 |
| Ciconiidae - 100 | Motacillidae - 243 | Sandcoleidae - 141 |
| Coliidae - 145 | Muscicapidae - 237 | Scaniornithidae - 269 |
| Columbidae - 218 | Musophagidae - 185 | Scolopacidae - 126 |
| Colymboididae - 63 | Necrobyidae - 207 | Selenornithidae - 207 |
| Coraciidae - 223 | Nyctibiidae - 197 | Sittidae - 242 |
| Corvidae - 230 | Odontopterygidae - 81 | Sophiornithidae - 206 |
| Cracidae - 151 | Orthocnemidae - 178 | Steatornithidae - 197 |
| Cuculidae - 185 | Orthocnémides - 178 | Stercorariidae - 197 |
| Cyphornithidae - 81 | Otididae - 98 | Stereornithidae - 178 |
| Darwinornithidae - 178 | Palaeogithalidae - 269 | Strigidae - 207 |
| Dasornithidae - 81 | Palaelodidae - 101 | Struthiolithidae - 60 |
| Devincenziidae - 178 | Palaeoglaucidae - 207 | Struthionidae - 60 |
| Diatrymidae - 94 | Palaeotididae - 60 | Sturnidae - 230 |
| Diomedeoididae - 78 | Paraortygidae - 151 | Sulidae - 66 |
| Eleutherornithidae - 254 | Paridae - 242 | Sylphornithidae - 270 |
| Emberizidae - 246 | Parvicuculidae - 263 | Sylviidae - 240 |
| Enkurosulidae - 66 | Passeridae - 244 | Thinocoridae - 133 |
| Eocoraciidae - 223 | Patagornithidae - 178 | Threskiornithidae - 92 |
| Eocypselidae - 197 | Pedionomidae - 133 | Tinamidae - 58 |
| Eogruidae - 96 | Pelagornithidae - 178 | Todidae - 225 |
| Ergilornithidae - 96 | Pelecanidae - 86 | Tolmodidae - 178 |
| Eurylaimidae - 227 | Pelecyornithidae - 178 | Troglodytidae - 236 |
| Falconidae - 220 | Petralcidae - 138 | Trogonidae - 224 |
| Fringillidae - 244 | Phaethontidae - 64 | Turdidae - 237 |
| Gallinuloididae - 151 | Phalacrocoracidae - 70 | |

Turnipacidae - 133
Upupidae - 76

Veflintornithidae - 186
Zygodactylidae - 141

Index of genus-group namesHome-pages of genera are underlined.

- Accipiter - 190, 191
 Acrocephalus - 240
 Actiornis - 93, 250
 Actitis - 130
 Aegialornis - 200, 201, 202, 203
 Aegithalos - 242
 Aegolius - 200, 201, 202, 203
 Aegyptius - 189, 190, 272
 Aenigmavis - 185
 Agnopterus - 108, 112, 178, 181, 182, 183, 184
 Alasio - 207, 211, 212
 Alauda - 228, 229, 243, 247
 Alca - 139, 140, 341
 Alcedo - 224, 225
 Alektoris - 157, 159, 160, 161, 162, 165, 166, 169
 Alisterus - 146
 Alopochen - 116, 117
 Ameghinornis - 185
 Ameripodius - 150
 Ammoperdix - 157, 159
 Amphipelargus - 97, 98
 Amphiserpentarius - 102, 193, 194
 Amynoptilon - 193, 194
 Anas - 70, 109, 110, 111, 113, 115, 116, 117, 118, 119, 120, 121, 122, 124, 125, 250, 279
 Anatalavis - 107, 108
 Anatipeda - 273
 Anhinga - 73, 74
 Anser - 113, 115, 116, 117, 124, 125
 Anseranas - 109
 Anserobranta - 111, 116, 117
 Anthus - 243, 244, 245
 Apaloderma - 224, 225
 Apopempsis - 186
 Aprosmictus - 146
 Apus - 204, 205
 Aquila - 187, 192, 193, 194, 195, 196
 Aquilavus - 101, 194, 195, 196
 Archaeoganga - 134, 135
 Archaeopsittacus - 146
 Archaeotrogon - 197, 198, 199
 Ardea - 68, 69, 71, 88, 92, 94, 156, 157, 212, 251, 279
 Ardeacites - 251
 Ardeagranda - 68
 Ardeipeda - 273
 Ardeita - 280
 Ardeola - 68, 69
 Argala - 279
 Argillipes - 251, 252
 Argillornis - 83
 Asio - 208, 216, 217, 218
 Athene - 214, 215, 216
 Aurornis - 279
 Aviadactyla - 273
 Avipeda - 273, 274
 Aythya - 110, 118, 119, 120, 121, 122, 124
 Balcanas - 117
 Balearica - 89
 Barornis - 95
 Basityto - 207, 209
 Bathornis - 178
 Batrachites - 226
 Belornis - 201
 Berruornis - 206
 Bombycilla - 229
 Bonasa - 168
 Botaurites - 71, 72, 156
 Botaurus - 69, 70
 Branta - 72, 116, 252
 Brontornis - 178
 Bubo - 208, 209, 212, 213, 216, 217
 Bucephala - 121, 122, 123
 Bucklandium - 24, 277
 Buteo - 191, 192, 195
 Buthierax - 196
 Calandrella - 228
 Calidris - 127, 131
 Calonectris - 81
 Camaskelus - 279
 Capella - 127, 128
 Capitonides - 147
 Caprimulgus - 199, 200
 Carbo - 280
 Carduelis - 245
 Centropus - 224
 Cepphus - 140
 Cerestenia - 252
 Certhia - 243
 Cettia - 240
 Chaetura - 205
 Charadriipeda - 274
 Charadrius - 135
 Chauvireria - 157, 158, 160
 Chenornis - 252
 Chlamydotis - 98, 99, 100
 Chlidonias - 137, 138
 Ciconia - 101
 Circaetus - 190
 Circus - 190
 Clangula - 122
 Coccythraustes - 246, 247
 Colius - 145, 146
 Collocalia - 204
 Columba - 134, 218, 219, 220
 Colymboides - 63, 111
 Colymbus - 85, 86, 139, 252
 Coracias - 224
 Corvus - 188, 230, 231, 232, 233, 234, 235, 247, 248
 Coturnipes - 253
 Coturnix - 152, 153, 154, 155, 158
 Crex - 176, 177
 Cryptornis - 224
 Cuculus - 186, 187
 Cursorius - 133
 Cygnanser - 113
 Cygnavus - 111, 112
 Cygnopterus - 112
 Cygnus - 101, 113, 114, 115, 253
 Cyphornis - 81
 Cypselavus - 202, 203, 204
 Cypseloides - 203, 204
 Cypselus - 203, 204
 Darwinornis - 178
 Dasornis - 81, 82, 83
 Delichon - 227
 Dendrochen - 110, 111
 Dendrocopos - 148, 149
 Dendrocygna - 118, 119
 Dendronessa - 118
 Devincenzia - 178
 Diatropornis - 252, 270
 Diatryma - 94, 95, 96, 185, 254
 Diomedea - 80, 341
 Diomedeoides - 78
 Dolichopterus - 253, 254
 Dolicopterus - 253, 254
 Dynamopterus - 186
 Dyspetornis - 63
 Egretta - 68, 69
 Elaphrocnemus - 179, 180, 181, 183

- Eleutherornis - 254
 Elorius - 130
 Elornis - 254, 255
 Emberiza - 246, 247
 Empheresula - 66
 Enkurosula - 66, 67
 Eocathartes - 102, 255
 Eocolius - 300
 Eocoracias - 223
 Eocypselus - 197
 Eoglaucidium - 142, 143
 Eogrus - 96
 Eortyx - 172
 Eostega - 66
 Eostrix - 255
 Eremophila - 229
 Ergilornis - 96
 Erithacus - 237
 Erolia - 131, 132
 Eudocimus - 93
 Eupterornis - 255, 256
 Euronyctibius - 199
 Eutolmaetus - 193, 195, 196
 Eutreptodactylus - 257, 270
 Falco - 187, 188, 190, 191, 192, 209, 218, 219, 220, 221, 222
 Ficedula - 240
 Filholornis - 179, 180, 181, 182, 183
 Fluviatilavis - 107, 256
 Fluvioviridiavis - 197, 198
 Francolinus - 157, 159, 160, 161
 Frigidafons - 78, 79
 Fringilla - 226, 244, 245, 248
 Fulica - 86, 177, 178, 256, 341
 Fuligula - 110, 119, 120, 121
 Galerida - 228
 Gallinago - 127, 128
 Gallinula - 177
 Gallinuloides - 151
 Gallus - 155, 156, 161, 163, 164, 165, 169, 170, 279
 Garganoaetus - 192
 Garrulus - 234, 235
 Gastornis - 94, 95, 96, 256, 268
 Gavia - 63, 64, 252
 Gaviota - 136
 Geiseloceros - 256, 257
 Gerandia - 134
 Geranopsis - 88, 184
 Geranopterus - 126, 223
 Geronticus - 93, 94
 Gervaisia - 106
 Gigantibis - 257
 Glareola - 132, 133
 Glaucidium - 214
 Gracilitarsus - 257
 Gracula - 225
 Graculavus - 259
 Graculus - 70, 71
 Grallavis - 100
 Gruipeda - 275
 Grus - 88, 89, 90, 91, 92, 105
 Gryzaja - 98, 100
 Gypaetus - 188
 Gyps - 188, 189, 190
 Gypsornis - 179
 Halcyornis - 259
 Haliaeetus - 188
 Haliaeus - 72
 Harrisonavis - 106
 Hassiavis - 257
 Headornis - 108
 Heliadornis - 65
 Helornis - 254, 255
 Hemiprocne - 200
 Hermosiornis - 178
 Hieraaetus - 193
 Himantopus - 279
 Hippolais - 240, 241
 Hirundo - 205, 227, 248
 Homalopus - 257
 Horusornis - 258
 Howardia - 258
 Hydromis - 63
 Ibis - 93, 156
 Ichnium - 277
 Idiornis - 178, 179, 180, 182, 183, 184
 Ieracus - 270, 271
 Intulula - 207, 211
 Itardiornis - 87
 Ixobrychus - 69
 Jungornis - 200
 Juncitarsus - 103
 Jynx - 148
 Kashinia - 103
 Kievornis - 258, 259
 Lagopus - 15, 167, 168, 169
 Lambrechtia - 157, 160
 Lanius - 229, 230, 248
 Laputa - 259
 Laputavis - 259
 Laricola - 136
 Larus - 24, 79, 132, 136, 137, 138, 259, 277, 279
 Latipons - 171, 172
 Laurillardia - 75, 76
 Leptoganga - 134, 135
 Leptoptilos - 100, 279
 Lestris - 138
 Leucogeranus - 91
 Limnatornis - 77, 145, 146
 Limosa - 128, 173
 Liornis - 178
 Lithornis - 58, 59, 83
 Litoripes - 259
 Locustella - 240
 Loxia - 245, 246
 Ludiortyx - 152, 172, 199, 267
 Lullula - 228
 Luscinia - 237, 241
 Lymnocryptes - 127
 Lyrurus - 167
 Macrodonopteryx - 82
 Macrornis - 277
 Marinavis - 259
 Masillacoliis - 143
 Masillapodargus - 205
 Megalornis - 83
 Megapaloelodus - 103, 104, 106
 Melanitta - 122
 Melanocorypha - 228
 Mergellus - 123
 Mergus - 123, 124, 260
 Merops - 226
 Mesembriornis - 178
 Mesogiornis - 202
 Messelastur - 143
 Messelirrisor - 75, 270
 Messelornis - 87, 260
 Microena - 260
 Microrallus - 126, 174
 Microsula - 67
 Milnea - 93
 Milvoidea - 260
 Milvus - 188, 196
 Miobaptus - 85
 Miocorax - 72, 230
 Miocorvus - 230
 Miofulica - 256
 Miogallus - 155, 156, 157
 Mioglareola - 132
 Mioglaux - 207, 210, 211
 Mionetta - 109, 111, 118, 155
 Miootis - 98
 Miopelecanus - 86, 87
 Miophasianus - 155, 156
 Miopico - 141
 Miorallus - 175, 176
 Monticola - 238
 Montirallus - 172, 173

- Morus - 67
 Motacilla - 237, 238, 239,
 240, 241, 243, 244, 248
 Muscicapa - 237, 239
 Musophaga - 186
 Neanis - 263
 Necrobyas - 207, 208, 209
 Necornis - 146
 Nectornis - 70, 71
 Neptuniavis - 59, 78
 Netta - 121
 Nettapterornis - 107, 108
 Nettapus - 117
 Nettion - 118
 Nisaetus - 193, 195, 196
 Nocturnavis - 217
 Nucifraga - 235
 Numenius - 128, 129, 130,
 172, 173, 232, 233
 Nupharanassa - 126
 Nyctea - 212, 213
 Nycticorax - 69, 70
 Oblitavis - 184, 185
 Occitaniavis - 184, 185
 Odontopteryx - 81, 82
 Oenanthe - 238
 Oligocathartes - 260, 261
 Oligocolius - 145
 Oligocorax - 70, 71
 Oligostrix - 207, 217, 218
 Olor - 113, 114, 115
 Omorhamphus - 95
 Ophthalmomegas - 212, 213
 Ornithichnidium - 277
 Ornithichnites - 275, 277
 Ornithoidichnites - 277
 Ornitholithes - 94, 261, 279
 Ornitholithus - 261, 262
 Ornithotarnocia - 275
 Ornithichnites - 275, 277
 Ornitocnemis - 88
 Orthocnemus - 60, 88, 89,
 178, 180, 182, 183
 Osteornis - 226
 Otis - 98, 99, 100, 280
 Otus - 210, 212, 215, 218
 Oxyura - 119, 123, 124
 Pachystruthio - 60, 61
 Palaegithalus - 269
 Palaelodus - 103, 104, 105,
 106, 113, 258
 Palaeoaramides - 147, 158,
174, 175, 262
 Palaeobyas - 206
 Palaeocircus - 187
 Palaeocryptonyx - 150, 154,
157, 158, 159, 161, 170
 Palaeocycnus - 113, 114
 Palaeocycgnus - 113, 114
 Palaeoglaux - 207
 Palaeogrus - 60, 88, 89
 Palaeohierax - 187
 Palaeopapia - 112, 258
 Palaeoperdix - 152, 156,
 158, 159, 175
 Palaeopicus - 145
 Palaeopsittacus - 262
 Palaeortyx - 110, 151, 152,
 153, 154, 155, 156, 157,
 158, 159, 172, 198, 280
 Palaeostruthio - 60, 61
 Palaeotis - 60
 Palaeotodus - 225, 226
 Palaeotyto - 206
 Palaetus - 279, 280
 Palescyvus - 202, 203
 Pandion - 187
 Paracorax - 71, 73
 Paracygnopterus - 109
 Paraortygoides - 170
 Paraortygometra - 174
 Paraortyx - 151, 152
 Paraprefica - 198
 Pararallus - 174, 175, 262
 Parasula - 66
 Paratrogon - 224, 225
 Paratyto - 207, 208
 Parus - 242, 249
 Parvicuculus - 263, 266
 Parvigyps - 59, 263
 Parvirallus - 171
 Parvulivinator - 263
 Passer - 244
 Passeripedia - 275
 Patagornis - 178
 Pavo - 16, 162, 163
 Pedionomus - 133
 Pediorallus - 58, 59
 Pelagornis - 81, 83, 280
 Pelargocrex - 193, 194
 Pelargodes - 193, 194
 Pelargopappus - 187, 193,
 194
 Pelargopsis - 193, 194
 Pelargosteon - 263
 Pelecanus - 73, 86, 87
 Pelecyornis - 178
 Pelophilus - 226
 Percolinus - 179, 180
 Perdix - 157, 161, 165, 166
 Pernis - 187
 Petralca - 138, 139
 Petronia - 244
 Petropluvialis - 264
 Phalacrocorax - 70, 71, 72,
 73, 252, 280
 Phalaropus - 129
 Phasianus - 156, 157, 162,
 163, 164, 165, 179, 280
 Philomachus - 127
 Phirriculus - 77
 Phodilus - 207
 Phoebastria - 341
 Phoenicopterus - 106, 107
 Phoeniculus - 76
 Phoenicurus - 237
 Phororhacos → Phorusrhacus
 Phorusrhacus - 178
 Phylloscopus - 241
 Pica - 233, 234, 235
 Picoides - 148
 Picus - 145, 148, 264
 Pinguinus - 139
 Pinicola - 246
 Pirortyx - 150, 152
 Piscator - 264
 Platalea - 94
 Plautus - 139
 Plegadis - 93
 Plesiocathartes - 102, 264
 Pliocarbo - 71, 72
 Pliogallus - 157, 158, 159,
163, 165
 Pliogrus - 90, 264, 265
 Plioperdix - 154, 157, 158,
 159, 160, 161
 Plotornis - 79, 80
 Plotus - 73
 Plumumida - 92, 93
 Pluvialis - 135
 Podiceps - 85, 86, 341
 Pogoniulus - 149
 Poicephalus - 262
 Polemaetus - 187, 193
 Porzana - 176
 Precursor - 265
 Prefica - 197
 Presbyornis - 108
 Primapus - 200, 201, 203,
 266
 Primocolius - 143, 144
 Primodroma - 78
 Primoscens - 141
 Primotrogon - 224
 Primozygodactylus - 141,
 142
 Proalector - 155, 157
 Proanser - 116, 117
 Proardea - 68, 94, 156
 Proardeola - 67, 68, 69
 Probalearica - 89, 90, 103,
 105
 Procellaria - 79
 Procellornis - 279, 280

- Proceriavis - 265
 Proculus - 263, 265, 266
 Procypseloides - 203, 204
 Proherodius - 59, 266
 Promilio - 196
 Promusophaga - 58, 59
 Propelargus - 100, 184, 185
 Prophaethon - 64
 Prophalacrocorax - 260
 Proplegadis - 266
 Prosybris - 207, 208, 209
 Prototelargus - 279, 280
 Protopelicanus - 266
 Protornis - 226, 266
 Protostrix - 255
 Prunella - 241, 242
 Pseudastur - 141, 142
 Pseudodontornis - 81, 82
 Pseudosula - 66, 67
 Pseudotantalus - 156
 Psilopterus - 178
 Psittacopes - 142
 Psittacus - 146, 147
 Pterocles - 134, 135
 Puffinus - 79, 80, 81
 Pulchravipes - 275, 276
 Pumiliornis - 267
 Pyrrhocorax - 234, 235, 236
 Pyrrhula - 246
 Quasisyndactylus - 267
 Quercymegapodius - 150
 Quercypodargus - 267
 Quercypsitta - 141, 144
 Quercyrallus - 153, 173, 267
 Querquedula - 111
 Rallicrex - 173, 174
 Rallus - 153, 154, 173, 175,
 176, 177, 267
 Recurvirostra - 107
 Regulus - 241, 249
 Remiornis - 256, 268
 Rhynchaetes - 92
 Riparia - 227
 Rissa - 137
 Romainvillia - 107, 108,
 109
 Rostratula - 126
 Rostrornis - 178
 Rupelornis - 268
 Salmila - 268
 Sandcoleus - 141
 Sarcogeranus - 91
 Sarmatosula - 67
 Saurornis - 278
 Saxicola - 237
 Scaniacypselus - 203
 Scanionis - 268, 269
 Schaubortyx - 152, 154
 Scolopax - 127, 128, 129,
 130, 155
 Selenornis - 207, 218
 Selmes - 142
 Serinus - 245, 247
 Serpentarius - 194
 Serudaptus - 143
 Sitta - 243, 249, 269
 Somateria - 121, 122
 Sophiornis - 206
 Spatula - 120
 Stercorarius - 138
 Stereornis - 178
 Sterna - 137, 138
 Stintonornis - 269
 Streptopelia - 220
 Strigogyps - 185, 269
 Strix - 208, 209, 210, 211,
 212, 213, 214, 215, 216,
 217, 218
 Struthio - 60, 61
 Struthiolithus - 60, 61
 Sturnus - 230, 242
 Sula - 66, 67, 112, 260, 280
 Suloides - 279, 280
 Surnia - 214
 Sylphornis - 269, 270
 Sylvia - 237, 240, 241
 Tachornis - 201
 Tachyornis - 201, 202
 Tadorna - 117
 Talantatos - 179, 180, 181
 Tantaleus - 279
 Tantalus - 156, 157, 179,
 180, 280
 Taoperdix - 110, 151, 152,
 153, 154
 Tapinopus - 102, 270
 Taubacrex - 150
 Telecrex - 180
 Teracus - 270, 271
 Tertiariaporphyryla - 174,
 175
 Tetrao - 151, 155, 165, 166,
 167, 168, 169
 Tetraogallus - 157
 Tetraornithopedia - 276
 Tetrastes - 168
 Tetrax - 98, 99
 Thinocorus - 133
 Thiornis - 85
 Todarna - 117
 Tolmodus - 178
 Torgos - 190
 Totanus - 130, 131, 132,
 137, 271
 Trachyphonus - 147, 148
 Tringa - 127, 129, 130, 131,
 132, 135, 137, 152, 172
 Troglodytes - 236, 237, 249
 Trogon - 224, 225
 Turdicus - 249
 Turdoides - 249
 Turdus - 238, 239, 240, 249
 Turnipax - 133
 Tynskya - 271
 Typhornis - 279, 280
 Tyto - 207, 209, 210, 215,
 218
 Upupa - 76, 94, 236
 Uria - 139, 140, 341
 Uriopsis - 280
 Urmionis - 96, 97, 98
 Urocolius - 145, 146
 Vanellus - 135, 271
 Veflintornis - 186
 Ventivorus - 198
 Villetus - 173, 271
 Vultur - 188, 189, 195, 271,
 272
 Xenopsitta - 146, 147
 Zygodactylus - 141, 144,
 264

Index of species-group names

Trinominals are indexed as binominals. Diacritical marks are omitted. Home-pages of fossil species and subspecies are underlined.

- aalge, Colymbus - 139
 Uria - 139
 abeli, Urmionis - 96
 absurdipes, Selmes - 142
 acuta, Anas - 120
 adelus, Ludiortyx - 267
 Rallus - 267
 adzalycensis, Struthiolithus - 61
 aegyptioides, Gyps - 190
 aesculapi, Gallus - 162, 163
 Pavo - 162
 affinis, Chlamydotis - 99
 Cygnopterus - 112
 Otis - 99
 Sula - 112
 afghana, Grus - 90
 africanus, Bubo - 213
 agilis, Otis - 279
 ajax, Diatryma - 95
 Gastornis - 95
 alatus, Geranopterus - 223
 alba, Motacilla - 244
 Tyto - 210
 albae, Anas - 124
 albellus, Mergellus - 123
 Mergus - 123
 albicilla, Falco - 188
 Haliaeetus - 188
 alexjevi, Struthiolithus - 61
 alexis, Ameripodius - 150
 alfhildae, Elaphrocnemus - 179
 Phasianus - 179
 alpestris, Eremophila - 229
 alphonsi, Cygnopterus - 112
 alpina, Calidris - 127, 131
 Erolia - 131
 altus, Miogallus - 156, 157
 Miophasianus - 156
 Phasianus - 156
 aluco, Strix - 215
 ambiguus, Palaelodus - 103, 104, 105,
 106, 258
 amissa, Ardea - 68
 Egretta - 68
 Proardea - 68, 69
 anas, Anatipeda - 273
 anatoides, Anser - 117
 Nettapus - 117
 anatolicus, Nectornis - 70
 angelis, Athene - 214
 anglica, Diomedea - 80
 Phoebastria - 341
 anglicus, Actiornis - 250, 254
 Colymboides - 63
 Elornis - 254
 angustipes, Bucephala - 122, 123
 Haliaeetus - 188
 anser, Anas - 116
 Anser - 116
 antecorax, Corvus - 231, 232
 antigone, Grus - 91
 antiqua, Procellaria - 79
 Prosybris - 208, 209
 Strix - 208, 209
 antiquus, Centropus - 224
 Cryptornis - 224
 Elornis - 255
 Falco - 222
 Numenius - 130
 Procellornis - 279, 280
 Puffinus - 79
 antunesi, Fluviatilavis - 107, 256
 apivorus, Pernis - 187
 apus, Apus - 205
 aquaticus, Rallus - 177
 aquitanica, Procellaria - 79
 aquitanicus, Puffinus - 79
 arborea, Ardeagrandsis - 68
 Lullula - 228
 arcensis, Anas - 279
 archiaci, Colius - 145
 Pavo - 162
 Phasianus - 162, 163
 Picus - 145
 Urocolius - 145
 arctica, Gavia - 64
 arcuatus, Ornitholithus - 261
 arenarius, Quercyrallus - 173
 Rallus - 173
 aretina, Anas - 120
 Aythya - 120
 Fuligula - 119, 120
 argenterae, Ormithichnidium - 277
 Ornitichnites - 277
 aristotelis, Pelecanus - 73
 Phalacrocorax - 73
 arquata, Numenius - 129
 Scolopax - 129
 artophoron, Palaeoglaux - 207
 arundinaceus, Acrocephalus - 240
 Turdus - 240
 arvensis, Alauda - 229
 arvernensis, Argala - 279
 Aythya - 110
 Bubo - 191, 207, 208, 209
 Buteo - 191

- Empheresula - 66
 Fuligula - 110
 Leptoptilos - 279
 Necrobyas - 208
 Parasula - 66
 Paratyto - 208
 Plotornis - 80
 Puffinus - 80
 Sula - 66
 ashbyi, Heliadornis - 65
 asiaticus, Struthio - 62
 atava, Anas - 113
 atavus, Anser - 113
 Cygnus - 113
 Falco - 220, 221, 222
 Lagopus - 168, 169
 ater, Parus - 242
 atra, Fulica - 178
 atricapilla, Motacilla - 241
 Sylvia - 241
 atthis, Alcedo - 225
 Gracula - 225
 augustus, Miophasianus - 156
 Phasianus - 156, 157
 aurelianensis, Ardea - 69
 auritus, Podiceps - 86, 341
 aurosum, Argillipes - 251
 ausonia, Alca - 341
 Uria - 140, 341
 austriaca, Petralca - 138, 139
 avitus, Botaurites - 71, 72
 babeheydariensis, Frigidafons - 78
 baconica, Chaetura - 205
 badense, Ichnium - 277
 badensis, Ornithoidichnites - 277
 bakalovi, Falco - 220
 balcanica, Actitis - 130
 Chauvireria - 157, 160
 Gallinula - 177
 balcanicus, Coccythraustes - 247
 Geronticus - 94
 Lagopus - 167
 balearica, Tyto - 209, 210
 ballmanni, Primozygodactylus - 142
 baranensis, Apus - 204
 Scolopax - 130
 barbarae, Pediorallus - 58, 59
 barbatus, Gypaetus - 188
 Vultur - 188
 baryosefi, Alectoris - 160
 basaltica, Anas - 70
 bassetti, Parvirallus - 171
 bavarica, Alectoris - 158
 beaumonti, Palaeoaramides - 147, 175
 Pararallus - 175
 Rallus - 174, 175
 Tertiariaporphyryula - 175
 becassi, Charadriipeda - 274
 benedeni, Anas - 250
 benimellalensis, Miopico - 141
 beremendensis, Gallus - 169
 Otis - 98
 bessarabensis, Otis - 99
 betfianus, Corvus - 232, 233
 bewickii, Cygnus - 115
 bilinicus, Aquilavus - 101
 Cygnus - 101
 biroi, Ornitholithus - 262
 blanchardi, Anas - 109, 110, 111, 125
 Dendrochen - 110
 Ludiortyx - 172
 Mionetta - 110
 Palaeortyx - 172
 blumeri, Protornis - 266
 bohatschevi, Leucogeranus - 91
 bohatshevi, Grus - 91
 Leucogeranus - 91
 Sarcogeranus - 91
 bohemica, Nupharanassa - 126
 bohemicus, Geranopterus - 126
 bonasia, Bonasa - 168
 Tetrao - 168
 bonifacti, Aquila - 192
 borealis, Turdoides - 249
 bosniaskii, Anthus - 247
 Ornitholithes - 261
 boulei, Dynamopterus - 186
 bournei, Primodroma - 78
 brachydactylus, Struthio - 61
 brachyptera, Eocoracias - 223
 Milvus - 196
 brancoi, Paraortyx - 152
 bravardi, Gallus - 163
 Pavo - 163
 brecciensis, Tantalus - 280
 Tantalus - 280
 bresciensis, Turdus - 249
 bretouensis, Sylphornis - 269, 270
 brevidactylus, Accipiter - 190, 191
 Masillacolius - 143
 brevipes, Haliaeetus - 188
 Himantopus - 279
 Otis - 99
 Palaeortyx - 153, 154, 155
 Taoperdix - 154
 brevis, Intulula - 211
 Strix - 211
 brevitarsus, Oligocolius - 145
 brodkorbi, Elaphrocnemus - 181
 Frigidafons - 78, 79
 Quercymegapodius - 250
 Tantalatos - 181
 broweri, Aegialornis - 202
 brumeli, Anser - 124
 brunhuberi, Ardea - 71
 Phalacrocorax - 72

- bubo, Bubo - 212
 bulgaricus, Regulus - 249
 bulotorum, Nupharanassa - 126
 burdigalensis, Larus - 279
 Procellornis - 279, 280
 buteo, Buteo - 191, 192, 195
 Falco - 191
 cadurensis, Palaeotyto - 206
 caeruleus, Parus - 242
 calcaria, Columba - 134
 Gerandia - 134
 campestris, Alauda - 243
 Anthus - 243
 campiterrae, Tyto - 209
 canaria, Serinus - 247
 cannabina, Carduelis - 245
 Fringilla - 245
 canorus, Cuculus - 186
 canus, Larus - 137
 Picus - 148
 capeki, Caprimulgus 200
 Francolinus - 159, 160
 Lambrechtia - 160
 Plioperdix - 160
 Surnia - 214
 carbo, Pelecanus - 73
 Phalacrocorax - 72, 73
 carduelis, Carduelis - 245
 carmesinae, Scolopax - 128
 carunculatus, Grus - 92
 caryocatactes, Nucifraga - 234, 235
 caudatus, Aegithalos - 242
 Parus - 242
 cayluxensis, Archaeotrogon - 199
 Palaeortyx - 151, 154, 198, 199
 Ludiortyx - 199
 Propelargus - 184
 cereti, Bucephala - 122, 123
 cervinus, Anthus - 243
 Motacilla - 243
 chauvirae, Aythya - 121
 cherrug, Falco - 222
 chersonensis, Struthio - 60, 62
 Struthiolithus - 60
 chloris, Carduelis - 245
 Loxia - 245
 chloropus, Fulica - 177
 Gallinula - 177
 christyi, Palaeoaramides - 174
 Rallus - 174, 175
 chrysaetos, Aquila - 192
 Falco - 192
 cinclus, Tringa - 131
 cinerea, Alauda - 228
 Ardea - 69
 Calandrella - 228
 citrinella, Emberiza - 247
 clangula, Bucephala - 121, 122, 123
 clypeata, Anas - 120
 Spatula - 120
 coccothraustes, Coccothraustes - 246
 Loxia - 246
 coelebs, Fringilla - 245
 collaris, Prunella - 242
 Sturnus - 242
 collurio, Lanius - 229
 collongensis, Alasio - 211, 212
 Strix - 212
 communis, Sylvia - 241
 compactus, Miotitis - 98
 concinna, Gavia - 64
 conjugens, Tetrao - 166, 167
 connectens, Megapaloelodus - 103
 Mergus - 123
 consobrina, Anas - 110, 111
 Dendrochen - 110, 111
 Mionetta - 110
 consobrinus, Palaeopicus - 145
 Picus - 145
 cooperi, Coturnipes - 253
 corax, Corvus - 231, 232
 corone, Corvus - 231, 232, 233
 correzensis, Lagopus - 169
 corroyi, Aquila - 194
 Aquilavus - 194
 cotei, Diatryma - 185, 254
 coturnix, Coturnix - 153, 155
 Tetrao - 155
 coturnoides, Pliogallus - 157, 159
 Plioperdix - 159
 cracrafti, Palaeobyas - 206
 crassa, Anas - 279
 crassipennis, Corvus - 188
 crassipes, Palaelodus - 105, 106
 Pliogallus - 163, 165
 crecca, Anas - 118, 119
 crex, Crex - 176, 177
 Rallus - 176
 creccoides, Anas - 250
 cretensis, Athene - 214
 crex, Elaphrocnemus - 180, 181
 cristata, Alauda - 228
 Galerida - 228
 Messelornis - 87
 cristatus, Parus - 242
 Pavo - 16
 Podiceps - 85
 croizeti, Gervaisia - 106
 Harrisonavis - 106
 Phoenicopterus - 106
 csakvarensis, Cygnanser - 113
 Cygnus - 113
 Olor - 113
 csarnotanus, Cuculus - 186
 cursor, Idiornis - 182, 183
 Orthocnemus - 181, 182

- curvirostra, Loxia - 246
 cuvieri, Gypsornis - 179
 Palaegithalus - 269
 Palaeocircus - 187
 Protopelicanus - 266
 Sitta - 269
 cyanus, Turdus - 238
 cygniformis, Anas - 113
 Anser - 113
 cygnus, Anas - 115
 Cygnus - 115, 139
 danielsi, Primozygodactylus - 141
 dasypus, Rallus - 153, 154
 Quercyrallus - 153
 davidi, Bubo - 212
 debellatrix, Mioglaux - 211
 debilis, Filholornis - 180, 181
 definitus, Rupelornis - 268
 dejardinii, Fulica - 256
 Miofulica - 256
 delfortrii, Pelagornis - 83, 279
 Plotornis - 79, 80
 delphinensis, Aquila - 195
 deperditus, Milvus - 196
 depereti, Palaeocryptonyx - 150
 Palaeortyx - 154, 155
 Pirortyx - 150
 Quercymegapodius - 150
 depredator, Aquila - 195
 Aquilavus - 195
 Eutolmaetus - 195
 Nisaetus - 195
 desnoyersii, Larus - 138
 Miophasianus - 156
 Phasianus - 156, 157
 destefanii, Paracorax - 73
 Phalacrocorax - 71, 73
 diluvii, Bucklandium - 277
 diomedea, Calonectris - 81
 Puffinus - 81
 disjuncta, Charadriipeda - 274
 dispar, Pararallus - 175
 Rallus - 175
 dissipata, Turnipax - 133
 dobrogensis, Sarmatosula - 67
 doksana, Oxyura - 123, 124
 dolnicensis, Larus - 132
 Mioglareola - 132
 domesticus, Gallus - 163
 donnezani, Alectoris - 160, 162, 169
 Palaeocryptonyx - 157, 159
 dubius, Procellornis - 279, 280
 Rallus - 267
 Strigogyps - 269
 edwardsi, Alectoris - 159
 Aquila - 193
 Eutolmaetus - 193
 Hieraaetus - 193
 Necrobyas - 208
 Nisaetus - 193
 Gastornis - 95
 Grallavis - 100
 Palaeocryptonyx - 159
 Palaeoperdix - 159
 Palaeortyx - 158
 Propelargus - 100
 Strix - 215
 Totanus - 130
 Tringa - 130
 Tyto - 215
 effosa, Ardea - 279
 egeriana, Gavia - 63
 egretta, Ardeipeda - 273
 elatus, Geranopsis - 184
 Occitaniavis - 184
 Propelargus - 184
 elegans, Laricola - 136
 Larus - 136
 elliotti, Diatropornis - 182, 252, 270
 Tapinopus - 270
 emryi, Palaeotodus - 225
 emuinus, Argillornis - 83
 Lithornis - 83
 Megalornis - 83
 ennouchii, Erolia - 131
 enucleator, Pinicola - 246
 eocaena, Tynskya - 271
 eous, Howardia - 258
 Palaeopapapia - 258
 epops, Upupa - 76
 eppelsheimensis, Anas - 124
 equitum, Anser - 114
 Cygnus - 114
 Olor - 114
 eremita, Geronticus - 94
 Upupa - 94
 ericetorum, Turdus - 238, 239
 erythropus, Tringa - 129
 escampensis, Palescyvus - 202, 203
 escampsiensis, Palaeotodus - 225
 estramosi, Porzana - 176
 etuliensis, Phasianus - 163
 europaea, Sitta - 243
 europaeus, Caprimulgus - 200
 Gallus - 170
 Plesiocathartes - 264
 europeus, Capitonides - 147
 Trachyphonus - 147
 excelsa, Balearica - 89
 Grus - 89
 Palaeogrus - 89
 excelsus, Ornithocemus - 89
 eximius, Palaeoaramides - 175
 Rallus - 174, 175
 exulans, Diomedea - 80
 eyermani, Puffinus - 81

- falconeri, Cygnus - 113, 114
 Olor - 114
 Palaeocycnus - 114
 Palaeocygnus - 114
 familiaris, Certhia - 243
 fasciata, Aquila - 193
 fasciatus, Hieraaetus - 193
 faujasi, Ornitholithes - 261
 fejfari, Microrallus - 126, 174
 Xenopsitta - 146, 147
 fenensis, Nycticorax - 69, 70
 ferina, Anas - 121
 Aythya - 121
 Fuligula - 122
 ferruginea, Tadorna - 111
 filiportatis, Avipeda - 273
 fisheri, Proplegadis - 266
 flammea, Strix - 217
 flammeus, Asio - 217
 floriana, Bubo - 217
 formosa, Ardea - 68
 formosus, Cygnavus - 111
 fossilis, Aquila - 195
 Caprimulgus - 200
 Corvus - 231, 232, 233
 Gallus - 163, 231, 232, 233
 Haliaeetus - 72
 Hirundo - 248
 Parus - 249
 Talantatos - 180, 181
 Tantalus - 179, 180, 181
 Vultur - 195, 271, 272
 fraasi, Pelecanus - 86, 87
 freudenthali, Garganoaetus - 192
 frugilegus, Corvus - 231, 233
 fulicaria, Tringa - 129
 fulicarius, Phalaropus - 129
 fuligula, Anas - 121
 Aythya - 121
 fulvus, Gyps - 189, 190
 funerea, Strix - 216
 funereus, Aegolius - 216
 gabbrensis, Ornitholithes - 261
 gaillardii, Apus - 204
 Cypselus - 204
 Idiornis - 182, 183
 Ludiortyx - 152
 Palaeocryptonyx - 154
 Palaeortyx - 151
 Paraortyx - 152
 Plesiocathartes - 264
 Proalector - 154, 155
 galericulata, Aix - 118
 Anas - 118
 Dendronessa - 118
 gallica, Apaloderma - 225
 Coturnix - 154
 Nyctea - 213
 Palaeortyx - 152, 153, 154
 Taoperdix - 154
 gallicus, Aegialornis - 201, 202
 Agnopterus - 181
 Cypselavus - 202, 203
 Idiornis - 182, 183
 Orthocnemus - 182, 183
 Paratrogon - 225
 Trogon - 224, 225
 gallinago, Capella - 127
 Gallinago - 127
 Scolopax - 127
 gallus, Gallus - 163, 164, 165
 gardneri, Latipons - 171
 garrulus, Bombycilla - 229
 Coracias - 224
 Lanius - 229
 garzetta, Ardea - 69
 Egretta - 69
 gassoni, Parvirallus - 171
 gaudryi, Ciconia - 101
 Picus - 264
 Zygodactylus - 264
 geiselensis, Diatryma - 96
 geiseltalensis, Ornithocnemus - 60
 Palaeogrus - 60
 gentilis, Accipiter - 190, 191
 Falco - 190
 georgei, Palaeopsittacus - 262
 Poicephalus - 262
 germanicus, Aegialornis - 201
 Pliogrus - 264, 265
 gervaisii, Aquila - 187
 Palaeohierax - 187
 ghardalensis, Scolopax - 155
 gigantea, Ardeipeda - 273
 Diatryma - 95
 Gallinula - 177
 Tyto - 210
 giganteus, Gastornis - 95
 glandarius, Corvus - 234
 Garrulus - 234, 235
 glareola, Tringa - 129
 glarisiensis, Protornis - 226
 glarniensis, Protornis - 226
 glaronensis, Protornis - 226
 goliath, Megapaloelodus - 106
 Palaelodus - 106
 goodwini, Microena - 260
 gracilipes, Palaelodus - 104, 105, 106
 gracilis, Agnopterus - 183
 Ardeita - 280
 Elaphrocnemus - 183
 Erolia - 132
 Idiornis - 183
 Milnea - 93
 Miopelecanus - 86
 Parvirallus - 171

- Pelecanus - 86
 Tringa - 132
 Troglodytes - 249
 gracillimus, Juncitarsus - 103
 graculoides, Chenornis - 252
 graculus, Corvus - 235
 Pyrrhocorax - 235
 graeca, Alectoris - 158, 161, 162
 Perdix - 161
 grandis, Elornis - 254
 Messelirrisor - 75
 Villetus - 271
 gratulator, Messelastur - 143
 gravipes, Somateria - 122
 gravis, Filholornis - 182, 183
 gregaria, Mioglareola - 132
 grigorescui, Tringa - 132
 grisegena, Podiceps - 85, 86
 grivensis, Palaeocryptonyx - 158
 Palaeortyx - 153, 154, 155
 Plioperdix - 154
 Totanus - 131
 Tringa - 131
 Zygodactylus - 144
 grus, Ardea - 92
 Grus - 92
 gryzaja, Otis - 100
 gypсорum, Alauda - 247
 Limosa - 173
 Montirallus - 173
 Numenius - 172, 173, 181
 Villetus - 173
 halcyrostris, Messelirrisor - 75
 haliaetus, Pandion - 187
 hamstediensis, Palaeopapua - 112
 hantoniensis, Agnopterus - 108
 Headonornis - 108
 harpax, Necrobyas - 207, 208
 hassenkampi, Pararallus - 262
 hastingsiae, Geranopsis - 88, 184
 heberti, Remiornis - 268
 helveticus, Eleutherornis - 254
 henrici, Asio - 208, 218
 Otus - 218
 Selenornis - 218
 herenthalsi, Cygnus - 253
 hermonis, Phasianus - 164
 herrenthalsi, Cygnus - 253
 hessae, Itardiornis - 87
 hirundo, Belornis - 201
 Tachyornis - 201, 202
 hispanica, Motacilla - 238
 Oenanthe - 238
 hoffmanni, Eortyx - 172
 Ludiortyx - 172
 Palaeortyx - 154, 172
 Tringa - 152, 172
 hoffstetteri, Archaeotrogon - 199
 holeae, Puffinus - 79
 holei, Puffinus - 79
 hookeri, Lithornis - 59
 Pediorallus - 59
 hordwelliensis, Grus - 88
 Ornitocnemus - 88
 Palaeogrus - 88
 Ibdopsis - 172
 humata, Motacilla - 248
 hungaricus, Corvus - 247, 248
 Palaeocryptonyx - 170
 hypogaea, Aquila - 195
 Aquilavus - 195
 hypoleuca, Ficedula - 240
 Motacilla - 240
 ibericum, Phalacrocorax - 72
 ibericus, Phalacrocorax - 72
 ignicapilla, Sylvia - 241
 ignicapillus, Regulus - 241
 ignota, Strix - 218
 Tyto - 218
 ignotus, Apus - 204
 Asio - 218
 Cypseloides - 204
 Cypselavus - 204
 Procypseloides - 204
 Zygodactylus - 144
 iliacus, Turdus - 239
 impennis, Alca - 139
 Pinguinus - 139
 Plautus - 139
 incerta, Ardeipeda - 273
 Collocalia - 204
 Nocturnavis - 217
 incertus, Bubo - 217
 Milvus - 196
 Promilio - 196
 incognita, Gigantibis - 257
 insolitus, Oblitavis - 184, 185
 Propelargus - 184
 insularis, Bubo - 213
 intermedia, Gruipeda - 275
 Palaeortyx - 155
 Strix - 215, 216
 intermedius, Cypselavus - 204
 Graculus - 71
 Miocorax - 71
 Miopelecanus - 87
 Pelecanus - 86, 87
 Phalacrocorax - 71
 Quercyrallus - 267
 Rallus - 267
 ipolyensis, Passeripedia - 275
 isarensis, Anas - 124
 itaboraiensis, Eutreptodactylus - 257, 270
 itardiensis, Agnopterus - 184
 Idiornis - 184
 Palaeotodus - 226

- ivani, Quercypsitta - 144
 janossyi, Corvus - 231, 232
 jansoni, Buteo - 191
 joleaudi, Palaeortyx - 154
 Plioperdix - 154
 juresaki, Perdix - 161, 162
 kalmani, Otis - 99
 karabachensis, Gallus - 161, 162
 karatheodoris, Struthio - 62
 kelleri, Paraprefica - 198
 keltica, Schaubortyx - 154
 Taoperdix - 152, 153
 kempi, Milvoidea - 260
 khosatzkii, Otis - 98
 klaasseni, Gastornis - 95
 kolozsvarensis, Rallicrox - 173
 kormosi, Pliogallus - 163, 165
 kurochkini, Euronyctibius - 199
 lacki, Aegialornis - 201
 Primapus - 200, 201
 lagopus, Archibuteo - 196
 Buteo - 192
 Lagopus - 15, 168
 lamarmorae, Bubo - 213
 Ophthalmomegas - 212, 213
 lambrechtii, Cygnopterus - 112
 Ornithotarnocia - 275
 Otis - 99
 lapponica, Limosa - 128
 larteti, Corvus - 230
 Miocorax - 231
 Miocorvus - 231
 lartetiana, Tringa - 131
 lartetianus, Psittacus - 147
 Totanus - 131
 larvatus, Archaeoganga - 134
 Pterocles - 134
 laticauda, Hassiavis - 257
 latipes, Ardea - 279
 laurillardi, Agnopterus - 182, 183
 lautus, Phalacrocorax - 72, 252
 lazaretensis, Columba - 219
 lebedinskyi, Eostega - 66
 leehardti → leenhardtii
 leenhardtii, Aegialornis - 201, 202
 lepidus, Psittacops - 142
 leucocephala, Anas - 124
 Oxyura - 119, 124
 leucogeranus, Grus - 91, 92
 leucorodia, Platalea - 94
 levantinus, Gallus - 163, 165
 lignitifila, Anas - 250
 lignitiphila, Anas - 250
 lignitum, Ardea - 212
 Bubo - 212
 limosa, Charadriipeda - 274
 Limosa - 128
 Scolopax - 128
 lipsiensis, Gaviota - 136
 liskunae, Anser - 115
 Olor - 115
 litorum, Precursor - 265
 littoralis, Elornis - 255
 Graculus - 71
 Helornis - 255
 Ieracus - 270
 Oligocorax - 71
 Phalacrocorax - 71
 Teracus - 270, 271
 livia, Columba - 219
 londinensis, Dasornis - 82, 83
 londiniensis, Dasornis - 83
 longaevus, Asio - 216
 Bubo - 216
 Gallus - 155, 156, 157
 longibrachis, Quasisyndactylus - 267
 longidentata, Pseudodontornis - 82
 longipennis, Argillornis - 83
 longipes, Coturnix - 153, 155
 Lyrurus - 167
 Masillapodargus - 205
 Palaeoperdix - 152, 154
 Palaeortyx - 155
 Phalacrocorax - 72
 Pliocarbo - 71, 72
 longirostris, Laurillardia - 75, 76
 Marinavis - 259
 lorteti, Paraortyx - 151, 152
 ludianus, Quercyrallus - 267
 lugubris, Parus - 249
 lundgreni, Scaniornis - 268, 269
 lunellensis, Athene - 215
 lungi, Palaeoaramides - 175
 Tertiariaporphyryla - 175
 lutetialis, Plumumida - 92, 93
 macrocephalus, Pseudastur - 142
 macroptera, Anas - 110
 macropus, Tetrao - 166, 167
 magna, Kashinia - 103
 magnifica, Promusophaga - 58, 59
 magnificus, Pulchravipes - 275, 276
 magnum, Tenuicrus - 103, 107
 magnus, Argillipes - 251
 Pelargocrex - 193
 Pelargodes - 193
 Pelargopappus - 193
 Pelargopsis - 193
 Precursor - 265
 Scolopax - 128
 major, Alauda - 247
 Alca - 139
 Dendrocopos - 149
 Motacilla - 248
 Idiornis - 180
 Miorallus - 176
 Orthocnemus - 180, 181

- Palaeortyx - 152
 Paraprefica - 198
 Parus - 242
 Pica - 233, 234
 Picoides - 148, 149
 Pirortyx - 152
 Primozygodactylus - 142
 Proanser - 116, 117
 Rallus - 175, 176
 majori, Amphipelargus - 97, 98
 Totanus - 131
 Tringa - 130
 mammeatus, Ornitholithus - 261
 mammillatus, Ornitholithus - 261
 mammosus, Ornitholithus - 261
 maraghanus, Amphipelargus - 97
 Urmiornis - 97, 98
 martelensis, Alectoris - 161, 162
 martini, Proceriavis - 265
 matthesi, Saurornis - 278
 maxima, Gruipeda - 275
 Palaeortyx - 280
 maximus, Miophasianus - 156
 media, Aviadactyla - 273
 Capella - 128
 Gallinago - 127, 128
 Palaeortyx - 280
 Prosybris - 208
 Scolopax - 127
 mediterranea, Alectoris - 161, 162
 medius, Dendrocopos - 149
 Falco - 220
 Lagopus - 168, 169
 Litoripes - 259
 Miophasianus - 156
 Necrobyas - 208
 Palaeoperdix - 156
 Parvirallus - 171
 Phasianus - 156, 157
 Picoides - 149
 Picus - 149
 meini, Apopemopsis - 185
 Musophaga - 185
 Veflintornis - 185
 melba, Apus - 205
 Hirundo - 205
 melitensis, Aegyptius - 189
 Columba - 219
 Cygnus - 114
 Grus - 91, 92
 Gyps - 189, 190
 Strix - 210
 Tyto - 210
 merganser, Mergus - 123
 merkeli, Juncitarsus - 103
 merula, Turdus - 238
 mesetaria, Chlamydotis - 98
 messelensis, Paraortygoides - 170
 Rhynchaetes - 92, 93
 meyerii, Anas - 118, 125
 Aythya - 118
 Protopelargus - 279, 280
 milneedwardsi, Geranopterus - 233
 Ibis - 156
 Pseudotantalus - 156
 Tantalus - 156, 157
 milvus, Falco - 188
 Milvus - 188
 minima, Charadriipeda - 274
 Protostrix - 255
 Scolopax - 127
 minimus, Diomedeoides - 78
 Lymnocryptes - 127
 Necrobyas - 209
 minor, Agnopterus - 183
 Ameghinornis - 185
 Branta - 252
 Francolinus - 160
 Charadriipeda - 274
 Gastornis - 256, 258
 Idiornis - 183
 Lanius - 229, 230
 Neptuniavis - 59, 78
 Orthocnemus - 183
 Parvicuculus - 263
 Remiornis - 256
 Primocolius - 144
 Strigogyps - 185
 Totanus - 131
 Tringa - 131
 minuta, Aquila - 193
 Columba - 219
 minutus, Colymboides - 63, 111
 Ixobrychus - 69
 Palaelodus - 105
 Palaeoaramides - 262
 Primoscens - 141
 Procuculus - 265
 miocaena, Palaeortyx - 157, 159
 Taoperdix - 151
 miocaenus, Graculus - 70
 Lanius - 248
 Nectornis - 70, 71
 Oligocorax - 70
 Pelagornis - 83
 Phalacrocorax - 70
 miocenica, Coturnix - 154
 miocenicus, Grus - 105
 Podiceps - 85
 miocoenicus, Suloides - 279, 280,
 Typhornis - 279, 280
 mirabilis, Gracilitarsus - 257
 miranda, Neptuniavis - 78
 mitchelli, Stintonornis - 269
 modularis, Motacilla - 241
 Prunella - 241

- molassicus, Ardeacites - 251
 moldavica, Gavia - 64
 Grus - 89, 90
 Probalearica - 89, 90
 moldavicus, Pavo - 64
 moldovicus, Gallus - 170
 mollissima, Anas - 122
 Somateria - 122
 monachus, Aegyptius - 189, 272
 Vultur - 189
 monedula, Coloeus - 233
 Corvus - 231, 233, 234
 montana, Fringilla - 244
 montanus, Passer - 244
 moravicus, Corvus - 233, 234
 morinellus, Charadrius - 135
 mourerchauvireae, Cypseloides - 203, 204
 Procypseloides - 203
 munieri, Laurillardia - 76
 murivorus, Garganoetus - 192
 muticus, Pavo - 16
 mutus, Lagopus - 15, 169
 Tetrao - 169
 nasi, Lithornis - 58, 59
 Pediorallus - 58, 59
 natator, Anas - 111
 Dyspetornis - 63
 Hydromis - 63
 Mionetta - 111, 118
 Querquedula - 111
 nearctica, Messelornis - 87
 nebulosa, Strix - 216
 neogena, Glareola - 133
 nestori, Puffinus - 79
 nicheti, Phasianus - 280
 nigra, Anas - 122
 Melanitta - 122
 nigricollis, Podiceps - 86
 niobrara, Gaviota - 136
 nisus, Accipiter - 191
 Falco - 191
 nivea, Prefica - 197
 noaillensis, Lagopus - 169
 noctua, Athene - 215
 Strix - 215
 nocturnus, Archaeotrogon - 199
 novorossicus, Struthio - 61
 numenioides, Totanus - 131
 Tringa - 131
 nyroca, Anas - 121
 Aythya - 119, 121
 occitanica, Columba - 219
 ochropus, Tringa - 129
 ochruros, Motacilla - 237
 Phoenicurus - 237
 ocyptera, Palaeortyx - 153, 154
 odessana, Gryzaja - 100
 odessanus, Pelecanus - 87
 oenanthe, Motacilla - 238
 Oenanthe - 238
 oenas, Columba - 219
 oeningensis, Anas - 124
 Anser - 125
 olor, Anas - 115
 Cygnus - 115
 olsoni, Morus - 67
 Oligocathartes - 260, 261
 Quercypodargus - 267
 omnisacrorum, Columba - 219
 orbisantiqvi, Berruornis - 206
 orientalis, Urmiornis - 97
 orlovi, Struthio - 61
 oxfordi, Anatalavis - 107, 108
 oxfordi, Nettapterornis - 107, 108
 oweni, Macrodonpteryx - 82
 Proherodius - 59, 266
 pagana, Actiornis - 93
 Ibis - 93
 paganus, Eudocimus - 93
 Plegadis - 93
 palaeoperdix, Perdix - 165, 166
 palapus, Apus - 205
 pallas, Eoglaucidium - 142, 143
 paludicola, Colius - 145
 Elorius - 130
 Limnatornis - 77, 145
 palumbus, Columba - 219
 palustris, Acrocephalus - 240
 Camaskelus - 279
 Colius - 146
 Ibidopodia - 258
 Motacilla - 240
 Necornis - 146
 Sylvia - 240
 pannonica, Anhinga - 73
 pannonicus, Pachystruthio - 61
 Plotus - 73
 Struthio - 60, 61
 paradoxa, Filholornis - 179, 180, 181
 Gavia - 64
 paralectoris, Argillipes - 252
 paratethydicus, Heliadornis - 65
 paratetrax, Otis - 99
 parisiensis, Gastornis - 95
 Laurillardia - 75, 76
 partium, Lyrurus - 167
 Tetrao - 167
 parva, Porzana - 176
 parvus, Messelirrisor - 75, 270
 Precursor - 265
 passerinum, Glaucidium - 214
 Strix - 214
 patevi, Loxia - 246
 penelope, Anas - 119
 pennatoides, Aquila - 196
 pennatus, Hieraaetus - 193

- pentelici, Grus - 90
 Pliogrus - 90
 percreca, Anas - 119
 peregrinabundus, Picus - 148
 perdix, Perdix - 161, 165
 Tetrao - 165
 peregrinus, Falco - 222
 Telecrex - 180, 181
 perpasta, Strix - 213
 perpastus, Bubo - 213
 perplexa, Ardea - 94
 Proardea - 94
 perplexus, Geronticus - 94
 perrierensis, Palaeoglaux - 207
 pessieti, Taoperdix - 151
 Tetrao - 66, 151
 petronia, Fringilla - 244
 Petronia - 244
 phaeopus, Numenius - 128
 Scolopax - 128
 phasianoides, Palaeortyx - 110, 155
 Pavo - 162
 Taoperdix - 110
 phasianus, Elaphrocnemus - 179, 180, 181
 philippi, Stercorarius - 138
 philomelos, Turdus - 238, 239
 phoeniculides, Upupa - 76
 phoenix, Avipeda - 274
 pica, Corvus - 233
 Pica - 233, 234
 picoides, Homalopus - 257
 pilaris, Turdus - 238
 pinguis, Archaeoganga - 134
 pinicola, Phirriculus - 77
 pisana, Columba - 218
 Fulica - 86, 341
 pisanus, Falco - 218
 Podiceps - 86
 piscator, Haliaeetus - 188
 piveteaui, Ardea - 251
 platyrhamphus, Fluvioviridiavis - 197
 platyrhynchos, Anas - 119, 120
 plioaena, Alectoris - 169
 plioaenica, Balcanas - 117
 plioaenus, Corvus - 231, 232, 233
 Numenius - 232, 233
 pliocenicus, Leptoptilos - 100
 pliodeserti, Chlamydotis - 100
 pohli, Serudaptus - 143
 poirrieri, Bubo - 210
 Mioglaux - 210, 211
 polgardiensis, Rallicrex - 174
 pomarinus, Lestris - 138
 Stercorarius - 138
 pontica, Fulica - 178
 ponticus, Ammoperdix - 178
 Plioperdix - 160
 portisi, Colymbus - 252
 Gavia - 252
 porzana, Porzana - 176
 porzanoides, Paraortygometra - 174
 Rallus - 174
 pouliani, Buthierax - 196
 praebonasia, Bonasa - 168
 Tetrastes - 168
 praecarbo, Phalacrocorax - 71, 72
 praecorax, Corvus - 232, 233
 praecox, Parvigyps - 59, 263
 praecursor, Totanus - 271
 praemedius, Dendrocopos - 149
 praeurogallus, Tetrao - 166, 167
 pratensis, Alauda - 243
 Anthus - 243
 primigenia, Grus - 90, 91, 92
 primigenius, Pyrrhacorax - 236
 princeps, Grus - 88
 Palaeogrus - 88
 prisca, Alectoris - 158
 Aquila - 196
 Palaeoperdix - 158
 priscus, Aquilavus - 196
 Larus - 277
 Eutolmaetus - 196
 Nisaetus - 196
 problematica, Grus - 103, 105
 Probalearica - 105
 procaccinii, Ornitholithes - 261
 protractus, Capitonides - 147
 Trachyphonus - 147
 proudlocki, Percolinus - 180
 Talantatos - 180
 pugnax, Philomachus - 127
 Tringa - 127
 pulchrapenna, Cerestenia - 252
 pulia, Rostratula - 126
 pusillus, Buteo - 191
 pygmaea,asio - 217
 Enkurosula - 67
 Microsula - 67
 Pseudosula - 67
 Sula - 67
 pygmaeus,asio - 217
 pyrrhacorax, Pyrrhacorax - 236
 Upupa - 236
 pyrrhula, Loxia - 246
 Pyrrhula - 246
 quercy, Quercyrallus - 173
 quercynus, Sophiornis - 206
 querquedula, Anas - 120
 radoboyensis, Batrachites - 226
 Fringilla - 226
 Merops - 226
 Pelophilus - 226
 raemdonckii, Larus - 79
 Puffinus - 79
 ragei, Ventivorus - 198

- ralli, Ornitholites - 179
 ralloides, Ardeola - 69
 rapax, Aquila - 192
 Falco - 192
 Palaetus - 279, 280
 recurvirostra, Charadriipeda - 275
 recurvirostrioides, Charadriipeda - 274
 regens, Barornis - 95
 remensis, Eupternis - 255, 256
 rhodopensis, Tetrao - 166, 167
 rhyphaeicus, Philomachus - 127
 richmondi, Rallus - 267
 ridibundus, Larus - 137
 riparia, Hirundo - 227
 Riparia - 227
 risgoviensis, Anas - 125
 Carbo - 279
 robinsoni, Latipons - 172
 robusta, Anas - 111
 Anserobranta - 111
 Laputa - 259
 Laputavis - 259
 Mionetta - 111
 Salmila - 268
 Surnia - 214
 Tyto - 210
 robustum, Amynoptilon - 194
 robustus, Amphiserpentarius - 194
 Eocathartes - 255, 257
 Geiseloceros - 256, 257
 Ornitocnemis - 88
 Serpentarius - 193, 194
 rogovitshi, Kievornis - 258, 259
 ronzoni, Mergus - 260
 Prophalacrocorax - 260
 Sula - 260
 rossignoli, Necrobyas - 208
 rozieri, Sula - 280
 rubecula, Erithacus - 237
 Motacilla - 237
 rufficollis, Anas - 116
 Branta - 116
 Podiceps - 86
 rufina, Anas - 121
 Netta - 121
 rufinus, Buteo - 191, 192
 Falco - 191
 rumana, Diomedea - 80
 rummeli, Basityto - 209
 rupelensis, Oligostrix - 217, 218
 rupeliensis, Ardea - 251
 Oligostrix - 217
 russelli, Gastornis - 96
 Messelornis - 260
 rustica, Hirundo - 227
 rusticola, Scolopax - 128
 rusticolus, Falco - 222
 sanctaneboulae, Recurvirostra - 107
 sanctialbani, Strix - 209
 Tyto - 208, 209, 210
 sansaniensis, Anas - 118, 119
 Palaeoperdix - 118, 158
 sapea, Aenigmavis - 185
 sarmatica, Ciconia - 101
 sarasini, Diatryma - 96, 185
 Gastornis - 96
 scaldicus, Uriopsis - 280
 scaldii, Anas - 125
 Anser - 125
 scandiaca, Nyctea - 213
 Strix - 213
 scandiacus, Bubo - 213
 scarabellii, Totanus - 132
 Tringa - 132
 schlosseri, Amphiserpentarius - 193
 Pelargopappus - 194
 schultzi, Gavia - 64
 scolopacinus, Osteornis - 226
 scops, Otus - 212, 214
 Strix - 212
 scotti, Paracygnopterus - 109
 selysii, Vanellus - 271
 senckenbergi, Cygnavus - 111, 112
 senogalliensis, Sitta - 249
 sepulta, Anas - 120
 Aythya - 120
 Fuligula - 119, 120
 sepultus, Leptoganga - 134
 Pterocles - 134
 serdicensis, Phalacrocorax - 72
 serinus, Serinus - 247
 serrator, Mergus - 123
 shrubsolei, Prophaethon - 64
 sibilatrix, Motacilla - 241
 Phylloscopus - 241
 sigei, Primocolius - 143
 silvasantosi, Ameripodius - 150
 simeonovi, Coccythraustes - 247
 similis, Ardea - 156, 157
 Botaurites - 156
 Proardea - 156
 simionescui, Corvus - 232, 233
 simplex, Petropluvialis - 264
 simurgh, Aquila - 192
 sirin, Avipeda - 274
 skalicensis, Anas - 250
 sociata, Podiceps - 85
 sociatus, Thiornis - 85
 solitarius, Monticola - 238
 Turdus - 238
 spassovi, Buteo - 191
 spelaeus, Gyps - 189, 190
 spinus, Carduelis - 245
 Fringilla - 245
 squatarola, Pluvialis - 135
 Squatarola - 135

- Tringa - 135
 stehlini, Ciconia - 101
 Pelargodes - 194
 Pelargopappus - 194
 Pelargopsis - 194
 Romainivillia - 108
 steini, Diatryma - 96
 steinheimensis, Gallus - 279
 Palaelodus - 113
 stellaris, Ardea - 69
 Botaurus - 69, 70, 251
 sternatus, Palaeostruthio - 60, 61
 storchi, Omorhamphus - 95
 storeri, Cepphus - 140
 strepera, Anas - 119
 striata, Motacilla - 239
 Muscicapa - 239
 subanser, Anser - 115
 subbuteo, Falco - 221, 222
 subfrancolinus, Francolinus - 160
 Plioperdix - 160
 submajor, Anas - 117
 Dendrocopos - 149
 submelba, Apus - 205
 sudrei, Quercypsitta - 144
 sutcliffei, Alectoris - 165, 166
 svecica, Luscinia - 237
 Motacilla - 237
 szarskii, Aegialornis - 203
 Primapus - 203
 Scaniacypselus - 203
 tadorna, Anas - 117
 Tadorna - 117
 tanaupus, Macromis - 277
 tarabukini, Alopochen
 Anserobranta - 116, 117
 tarda, Otis - 99
 tasnadii, Tetraornithopedia - 236
 taurica, Aurorornis - 279
 taurinus, Ornithichnites - 275
 tenuipennis, Ornitholithes - 261
 tenuirostris, Piscator - 264
 tenuis, Turdicus - 249
 teruelensis, Larus - 137
 Totanus - 137
 Tringa - 139
 tessellatus, Jungornis - 200
 Pumiliornis - 267
 tetrax, Otis - 99
 tetrax, Tetrao - 167, 168
 thraceiensis, Anser - 116
 tinnipara, Intulula - 211
 tinnunculus, Falco - 220, 221
 topei, Torgos - 190
 toliapica, Odontopteryx - 259
 toliapicus, Halcyornis - 259
 Larus - 24, 259
 tolutaria, Nupharanassa - 126
 torda, Alca - 140
 tordus, Alca - 139
 torquata, Muscicapa - 237
 Saxicola - 237, 238
 torquilla, Jynx - 141, 148
 totanoides, Laricola - 137
 Larus - 136
 totanus, Tringa - 129
 tothi, Pelargosteon - 263
 tracheliotus, Torgos - 190
 tridactyla, Rissa - 137
 tridactylus, Larus - 137
 trochanteria, Fringilla - 240
 troglodytes, Motacilla - 237
 Troglodytes - 237
 trouessarti, Prelargodes - 194
 Pelargopappus - 194
 Pelargopsis - 194
 tschulensis, Odontopteryx - 81
 tuberculata, Idiornis - 182
 tuberculatus, Agnopterus - 182
 turfa, Grus - 92
 turgaiensis, Agnopterus - 112
 ukrainus, Amphipelargus - 97
 Urmiornis - 97
 undulata, Chlamydotis - 98
 unicolor, Sturnus - 230
 uralensis, Strix - 213
 urbica, Delichon - 227
 Hirundo - 227
 urogallus, Tetrao - 166, 167, 170
 validus, Archaeoganga - 134
 Pterocles - 134
 vanellus, Tringa - 135
 Vanellus - 135
 velox, Anas - 118
 Nettion - 118
 Dynamopterus - 186
 venablesi, Gypsornis - 179
 Percolinus - 179
 venustus, Archaeotrogon - 198, 199
 verae, Cygnus - 113, 114
 verreauxii, Archaeopsittacus - 146
 Psittacus - 146
 vesperinus, Falco - 221
 veta, Athene - 216
 veterior, Capella - 127
 Gallinago - 127
 Porzana - 176
 vetus, Pyrrhocorax - 235, 236
 vianeyliaudae, Horusornis - 258
 viator, Dolichopterus - 253, 254
 Dolichopterus - 253
 villanziensis, Francolinus - 160
 vincenti, Eocypselus - 197
 Eostrix - 255
 viridis, Picus - 148
 viscivorus, Turdus - 239

- voinstvenskii, Lagopus - 168
- vulgaris, Sturnus - 230
- vulturinus, Lithornis - 58, 59
- wagneri, Todarna - 117
- walkeri, Ardeola - 68
- Eocolius - 300
- Proardeola - 67, 68, 69
 - walteri, Miobaptus - 85
- Podiceps - 85
 - waltoni, Villetus - 271
 - wardi, Scaniacypselus - 203
 - watteli, Parvulivenator - 263
 - weigelti, Palaeotis - 60
 - wenzensis, Francolinus - 160
 - wetmorei, Aegialornis - 202
- Apus - 204
- Mesogiornis - 202
 - wintershofensis, Strix - 215
- Otus - 211, 215
 - wintersteini, Primotrogon - 224
 - yelkouan, Puffinus - 80
 - zeylonensis, Bubo - 212, 213
- Strix - 212
 - zitteli, Archaeotrogon - 198, 199

