

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 401 "M" St., S.W.

Washington D.C. 20460

NOTICE OF PESTICIDE:

x Registration Reregistration

6218-78

EPA Rea

Number

Date of issuance

APR 2 7 2004

Term of Issuance

Conditional

Name of Pesticide Product

Ready to Use Year Round Spray Oil

Name and Address of Registrant (include ZIP Code):

Summit 7657Canton Center Drive Baltimore, MD 21224-2079

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide. Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
  - 2. Make the following label changes before you release the product for shipment:

## SEE PAGE 2 FOR LABEL CORRECTIONS.

3. Submit three copies of the final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "accepted with comments" is enclosed for your records.

Signature of Approving Official:

Akiva Abramovitch, Ph.D.

Chemist

Insecticide-Rodenticide Branch

Registration Division (7505C)

APR 2 7 2004

EPA Form 8570-6

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 6218-78".
- b. The name of the active ingredient must be changed to petroleum oil from paraffinic oil.
- c. You must add under the Directions for Use "Not to be applied to plants being grown for commercial use".
- d. The Storage and disposal statements must be revised. "Keep container tightly closed (misspelled) in storage". Delete "to prevent entry of contaminants". The Pesticide disposal is incorrect and it should be revised per PR Notice 2001-6.

If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions (toll free number). Never put unused product down any indoor or outdoor drain.

e. Delete "Leaf polish for Hardy Plants" on page 4 since it is not a pesticidal claim.

Enclosure

ACCEPTED with COMMENTS In EPA Letter Dateds

SHAKE WELL BEFORE USE

APR 2 7 2004

# READY-TO-USE Pungicide, and Rodent as amended, for the registered under EPA

Under the Federal Insecticide, Pungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

KILLS INSECTS FOUND ON FRUIT & ORNAMENTAL TREES
VEGETABLES, HOUSEPLANTS & ORNAMENTALS
ENVIRONMENTALLY RESPONSIBLE INSECTICIDE

## KEEP OUT OF THE REACH OF CHILDREN CAUTION

See back panel for additional precautions.
EPA Reg.# 6218- EPA Est.# 6218-MD-2

ACTIVE INGREDIENT	Paraffinic Oil*	1.00%
INERT INGREDIENTS		99.00%
*Unsulfonated Residue of Paraffinic Oil	92.0% Min.	

NET CONTENTS\_\_\_\_\_

Summit Responsible Solutions
7657 Canton Center Dr.Baltimore, MD 21224

## CAUTION KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS

## **FIRST AID**

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

If swallowed \*Call a poison control center or doctor immediately for treatment advice.

\*Do not induce vomiting unless told to by a poison control center or doctor.

If in eyes \*Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue

rinsing.

\*Call a poison control center or doctor for treatment advise.

If on skin \*Take off contaminated clothing.

\*Rinse skin immediately with plenty of water for 15-20 minutes.

\*Call a poison control center or doctor for treatment advise.

If inhaled \*Move person to fresh air.

\*If person is not breathingcall 911 or an ambulance, then give artificial

respiration, preferably mouth-to-mouth if possible.

\*Call a poison control center or doctor for further treatment advise.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Keep container tightly closeed in storage to prevent entry of contaminants. Store in cool, dry, locked area out of the reach of children. Do not store this product in direct sunlight for prolonged periods of time. This product should be stored in its original container only.

**Pesticide Disposal**: Securely wrap original container in several layers of newspaper and discard in the trash.

Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.

#### USE PRECAUTIONS

Do not spray plants suffering from drought (mositure stress). Do not apply during periods of prolonged high temperature combined with high relative humidity. Do not use this product with dimetholate, herbicides or fungicides such as Captan, Folpet, Dyrene, Karathene, Morestan or any product containing sulfur. Do not use with Sevin 50W formulation (carbaryl) on deciduous fruit trees.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

## APPLICATION INSTRUCTIONS

Shake the container vigoriously prior to use.

- The best times to spray are the early evening and early morning.
- The target pest and plant should be completely covered with the spray solution to be effective.
- Certain plants may be sensitive to oil. A small number of plants should be test sprayed prior to general spraying.
- Use caution on open blooms-bleaching and spotting have been observed on the open bloom of certain plants.
- May be used up to the day of harvest.

## FRUIT & NUT TREES

TREE	PEST	APPLICATION NOTES
Apple ,Pear Almonds,Apricots, Cherries,Peaches, Nectarines,Plum Prune,Pecans	Aphids(eggs) Apple Red Bug/ Mites (eggs) including European Red Mite/Scales(hard, soft)/Sourly Scale/ FruitTree Leaf Roller(eggs),Pear Leal, Blister Mite (eggs),Pear Psylla,Two Spotted Spider Mite, Pear Rust Mite, Tent Caterpillans, San Jose Scale, Peach Twig Borer (hibernacules),Cottony Peach Sc	Do not apply over fruit after waxy bloom forms. Application should not be made on oil sensistive varities-Do not apply to treeslacking adequate moisture. ale,

#### **CITRUS TREES**

Lemon & Grapefruit Naval & Valencia Oranges, other Citrus	Mites including: Rust Mite, Spider Mite Scales including: Black Scale, California Red Scale, Whitefly & Blackfly, Sooty Mold
Varities including Non-	
bearing	

For best results spray during fall, early Winter & Spring months when Red Scale are mpore easily controlled and trees are generally less reactive to oil. For all Citrus trees ensure adequate moisture.

## SHADE TREES, SHRUBS, ORNAMENTALS, FLOWERS & FOLIAGE PLANTS, CONIFERS

#### TREE OR PLANT

#### **PEST**

#### APPLICATION NOTES

Shade Trees dormant, and shrubs Conifers, Deciduous, Broadleaf Evergreens & Woody Ornamentals Ornamental Trees Shrubs including Conifers, Deciduous and Broadleaf Evergreens

Aphids, Adelgids, Eriophyid Mites, Gall Mits, Lace Bugs Leaf Beetle Larve, Leafminer Mealybugs, Psylids (Immature) Spider Mites, Whitefly

Oils remove the glaucus (blue) bloom from such evergreens as Colorado Blue Spruce and Koster Spruce. Always test spray sensitive Sawfly Larvae, Scale(immature) plants such as Cryptomeria, Smoke Tree, chamaecypris, Juniper, Japanese Holly & Spruce. Do not spray Walnut foliage. Most effective on Whitefly during immature stages.

Flowering & Foliage Plants including: Roses and other Flowering Shrubs, Bedding Plants, Foliage Ornamentals.

CAUTION: Up to 4 applications during the growing season are typically sufficient to control pests. Use a 2 week application interval. Do not spray when buds are fully open and elongation is occurring.

## SMALL FRUITS

Blueberries, Strawberries, Grapes

Mites, Sawfly(eggs), Scales Aphids, Mealybugs

## VEGETABLES

Beans, Corn, Peppers, Squash Tomatoes, Asparagus, Beets Cabbage, Cauliflower Cucurbits, Eggplants, Lettuce Melons, Potatoes, Radishes

Aphids, Mites, Beetle Larvae Leafminers, Thrips, Leafhopper Whitefly

## **HOBBY GREENHOUSE**

Azaleas, Begonia, Camellia Chrysanthemum, Oleffenbacfia Easter Lily, Fern, Gardenia Geranium, Hibiscus Foliage, Jade Plant, New Guinea Impatiens, Palms, Philodendron, Poinsettia Portulaca, Reiger Begonia, Zinnias Dieffenbachia, Poinsettias,

Aphids, Fungus Gnats, Leafminers, Mealybugs, Scales (soft & hardy), Spider Mites, Thirps Whitefly(immature)

Do not use on Coconut Palms or Maidenhair Fern. Chrysanthemum and Geranium Blooms have shown phytotoxicity at high rates

Leaf Polish for Hardy Plants

FREQUENCY OF APPLICATION: For the greenhouse pests listed, use once a week initially. As the pest is controlled, decrease the frequency to 2-3 weeks as needed.

Application safety during bloom period should be determined for each individual plant to be treated by conducting a small test.

## **INTERIORSCAPES & HOUSEHOLDS**

## **PLANTS**

13 July 2

## PEST

## APPLICATION NOTES

Chrysanthemum, Dieffenbachia, Aphids, Leaf Miner Dracaena, Fern, Ficus, Gardenia Meal Bugs, Scales Jade Plant, Palm, Philodendron, Schefflera

Spider Mites, Whitefly

Do not apply to plants in direct sunlight. Behind glass. Do not use on Coconut Palms, & Maidenhair Ferns. Chrysanthemum blooms have shown phytotoxic symptoms at higher usage.

Applicator should conduct a test for phytotoxicity by treating a few specimens before making a large scale application. Protect floor and floor coverings and furnishings from over spray.

CAUTION: Do not exceed 3 consecutive applications without allowing a 2-week interval without application.