

Certificate of Analysis

pGL4.30[*luc2P*/NFAT-RE/Hygro] Vector:

Part No. Size
E848A 20µg



Instructions for use of this product can be found in the pGL4 Luciferase Reporter Vectors Technical Manual #TM259, available online at: www.promega.com/protocols

Description: The pGL4.30[*luc2P*/NFAT-RE/Hygro] Vector^(a-f) contains an NFAT response element (NFAT-RE) that drives the transcription of the luciferase reporter gene *luc2P* (*Photinus pyralis*). *luc2P* is a synthetically-derived luciferase sequence with humanized codon optimization that is designed for high expression and reduced anomalous transcription. The *luc2P* gene contains hPEST, a protein destabilization sequence. The protein encoded by *luc2P* responds more quickly than the protein encoded by the *luc2* gene upon induction. The vector backbone contains an ampicillin resistance gene to allow selection in *E. coli* and a mammalian selectable marker for hygromycin resistance.

See the pGL4 Luciferase Reporter Vectors Technical Manual #TM259 for more information.

Concentration: 1µg/µl.

GenBank® Accession Number: DQ904462.

Storage Buffer: The pGL4.30[*luc2P*/NFAT-RE/Hygro] Vector is supplied in 10mM Tris-HCl (pH 7.4), 1mM EDTA.

Storage Conditions: See the product information label for storage temperature recommendations. Avoid multiple freeze-thaw cycles and exposure to frequent temperature changes. These fluctuations can greatly alter product stability. See the expiration date on the product information label.

Usage Note: Concentration gradients may form in frozen products and should be dispersed upon thawing. Mix well prior to use.

Quality Control Assays

Nuclease Assay: Following incubation of 1µg of the vector in restriction digest buffer B at 37°C for 16 hours, no evidence of nuclease activity is detected by agarose gel electrophoresis.

Physical Purity: $A_{260}/A_{280} \geq 1.80$, $A_{260}/A_{250} \geq 1.05$ at pH 7.4.

Sequence: The pGL4.30[*luc2P*/NFAT-RE/Hygro] Vector has been completely sequenced and has 100% identity with the published sequence, available at: www.promega.com/vectors

Signed by:

J. Stevens, Quality Assurance

^(a)BY USE OF THIS PRODUCT, RESEARCHER AGREES TO BE BOUND BY THE TERMS OF THIS LIMITED USE LABEL LICENSE. If the researcher is not willing to accept the terms of this label license, and the product is unused, Promega will accept return of the unused product and provide the researcher with a full refund.

Researchers may use this product for research use only, no commercial use is allowed. "Commercial use" means any and all uses of this product and derivatives by a party for money or other consideration and may include but is not limited to use in: (1) product manufacture; and (2) to provide a service, information or data; and/or resale of the product or its derivatives, whether or not such product or derivatives are resold for use in research. Researchers shall have no right to modify or otherwise create variations of the nucleotide sequence of the luciferase gene except that researchers may: (1) create fused gene sequences provided that the coding sequence of the resulting luciferase gene has no more than four deoxynucleotides missing at the affected terminus compared to the intact luciferase gene sequence, and (2) insert and remove nucleic acid sequences in splicing research predicated on the inactivation or reconstitution of the luminescence of the encoded luciferase. No other use or transfer of this product or derivatives is authorized without the prior express written consent of Promega. In addition, researchers must either: (1) use luminescent assay reagents purchased from Promega for all determinations of luminescence activity of this product and its derivatives; or (2) contact Promega to obtain a license for use of the product and its derivatives. Researchers may transfer derivatives to others for research use provided that at the time of transfer a copy of this label license is given to the recipients and recipients agree to be bound by the terms of this label license. With respect to any uses outside this label license, including any diagnostic, therapeutic or prophylactic uses, please contact Promega for supply and licensing information. PROMEGA MAKES NO REPRESENTATIONS OR WARRANTIES OF ANY KIND, EITHER EXPRESSED OR IMPLIED, INCLUDING FOR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE WITH REGARDS TO THE PRODUCT. The terms of this label license shall be governed under the laws of the State of Wisconsin, USA. This label license relates to Promega patents and/or patent applications on improvements to the luciferase gene.

^(b)U.S. Pat. No. 5,670,356.

^(c)U.S. Pat. No. 8,008,006 and European Pat. No. 1341808.

^(d)Patent Pending.

^(e)U.S. Pat. No. 7,728,118.

^(f)The method of recombinant expression of *Coleoptera* luciferase is covered by U.S. Pat. Nos. 5,583,024, 5,674,713 and 5,700,673. A license (from Promega for research reagent products and from The Regents of the University of California for all other fields) is needed for any commercial sale of nucleic acid contained within or derived from this product, and from The Regents of the University of California for all other fields) is needed for any commercial sale of nucleic acid contained within or derived from this product.

Part# 9PIE848
Revised 6/13



Promega

Promega Corporation

2800 Woods Hollow Road	
Madison, WI 53711-5399	USA
Telephone	608-274-4330
Toll Free	800-356-9526
Fax	608-277-2516
Internet	www.promega.com

PRODUCT USE LIMITATIONS, WARRANTY, DISCLAIMER

Promega manufactures products for a number of intended uses. Please refer to the product label for the intended use statements for specific products. Promega products contain chemicals which may be harmful if misused. Due care should be exercised with all Promega products to prevent direct human contact.

Each Promega product is shipped with documentation stating specifications and other technical information. Promega products are warranted to meet or exceed the stated specifications. Promega's sole obligation and the customer's sole remedy is limited to replacement of products free of charge in the event products fail to perform as warranted. Promega makes no other warranty of any kind whatsoever, and SPECIFICALLY DISCLAIMS AND EXCLUDES ALL OTHER WARRANTIES OF ANY KIND OR NATURE WHATSOEVER, DIRECTLY OR INDIRECTLY, EXPRESS OR IMPLIED, INCLUDING, WITHOUT LIMITATION, AS TO THE SUITABILITY, PRODUCTIVITY, DURABILITY, FITNESS FOR A PARTICULAR PURPOSE OR USE, MERCHANTABILITY, CONDITION, OR ANY OTHER MATTER WITH RESPECT TO PROMEGA PRODUCTS. In no event shall Promega be liable for claims for any other damages, whether direct, incidental, foreseeable, consequential, or special (including but not limited to loss of use, revenue or profit), whether based upon warranty, contract, tort (including negligence) or strict liability arising in connection with the sale or the failure of Promega products to perform in accordance with the stated specifications.

© 2006, 2008, 2009, 2013 Promega Corporation. All Rights Reserved.

Bright-Glo is a trademark of Promega Corporation.

GenBank is a registered trademark of the U.S. Department of Health and Human Services. *TransIT* is a registered trademark of Mirus Corporation.

Products may be covered by pending or issued patents or may have certain limitations. Please visit our Web site for more information.

All specifications are subject to change without prior notice.

Product claims are subject to change. Please contact Promega Technical Services or access the Promega online catalog for the most up-to-date information on Promega products.

Part# 9PIE848
Printed in USA. Revised 6/13.



pGL4.30[*luc2P*/NFAT-RE/Hygro] Vector Features List and Map:

NFAT response element	33–122
Minimal promoter	136–166
<i>luc2P</i> reporter gene	228–2003
SV40 late poly(A) signal	2043–2264
SV40 early enhancer/promoter	2312–2730
Synthetic hygromycin (Hyg ^r) coding region	2755–3792
Synthetic poly(A) signal	3816–3864
Reporter Vector primer 4 (RVprimer4) binding region	3931–3950
<i>ColE1</i> -derived plasmid replication origin	4188
Synthetic β-lactamase (Amp ^r) coding region	4979–5839
Synthetic poly(A) signal/transcriptional pause site	5944–6097
Reporter Vector primer 3 (RVprimer3) binding region	6046–6065




Figure 1. Vector map.

Sample Protocol to Determine Luciferase Induction by Ionomycin/PMA in HEK 293 Cells Transfected with pGL4.30[*luc2P*/NFAT-RE/Hygro] Vector

Materials to be Supplied by User

- Dulbecco's PBS (DPBS)
- 0.05% (w/v) trypsin in DPBS
- DMEM supplemented with 10% fetal bovine serum (DMEM/FBS)
- *TransIT*[®]-LT1 (Mirus Part# MIR2304)
- DMSO
- 1mM ionomycin in DMSO
- 1mg/ml phorbol 12-myristate 13-acetate (PMA) in DMSO
- Bright-Glo™ Luciferase Assay System (Cat.# E2610)
- HEK 293 cells

Day 1: Plate Cells

1. Grow HEK 293 cells in DMEM/FBS to approximately 75% confluency.
2. Harvest the cells via trypsinization. Remove the DMEM/FBS, wash the cells with DPBS and add the trypsin/DPBS (1X volume). After 2 minutes, add 4X volume of DMEM/FBS, collect the cell suspension and pellet the cells by centrifugation. Aspirate supernatant and resuspend in DMEM/FBS at a concentration of 20,000 viable cells/90µl DMEM/FBS.
3. Dispense 90µl of the cell suspension to the wells of a 96-well plate. Plate enough wells to perform each test condition in triplicate.
4. Cover the plate and place it in a tissue culture incubator at 37°C overnight (or for 24 hours).

Day 2: Transfect Cells

1. Prepare the DNA transfection master mix. Each well of 96-well plate to be transfected requires 10µl DMEM, 0.3µl *TransIT*[®]-LT1 and 0.1µg pGL4.30[*luc2P*/NFAT-RE/Hygro] plasmid DNA. To prepare the master mix, calculate the total number of wells that will be transfected and prepare 110% of this amount. It is recommended that at least 10 wells of master mix be prepared.
 - For each well, mix 10µl DMEM and 0.3µl *TransIT*[®]-LT1 in a microcentrifuge tube, briefly vortex at maximum setting and incubate at room temperature for 15 minutes.
 - For each well to be transfected, add 0.1µg of pGL4.30[*luc2P*/NFAT-RE/Hygro] Vector to the DMEM/*TransIT*[®]-LT1, vortex briefly and incubate at room temperature for 15 minutes.
2. Add 10µl of master mix to each well that is to be transfected.
3. Cover the plate and place it in a tissue culture incubator at 37°C overnight (or for 24 hours).

Day 3: Induce Transfected Cells and Measure Luciferase Activity

1. Prepare 1X induction and 1X control solutions. Calculate the volume of 1X induction and 1X control solution by multiplying the number of wells needed for each solution by 100µl and prepare 110% of this amount.
 - 1X induction solution: Dilute 1mM stock ionomycin solution to 1µM (1:1,000) and 1mg/ml stock solution of PMA to 10ng/ml (1:100,000) in DMEM/FBS.
 - 1X control solution: Dilute DMSO 1:1,000 in DMEM/FBS.
2. Remove medium from wells that will be treated with either 1X induction or 1X control solutions.
3. Add 100µl of 1X induction solution to the cells to be induced and 100µl of 1X control solution to the control noninduced cells.
4. Return the plate to the tissue culture incubator and induce for 17 hours.

Day 4: Analyze Luciferase Activity

1. Analyze luciferase activity using the Bright-Glo™ Luciferase Assay System as described in Technical Manual #TM052.
2. Calculate the fold induction as follows:

$$\text{Fold Induction} = \frac{\text{Average relative light units of induced cells}}{\text{Average relative light units of control cells}}$$