



NIAGARA SPRINGS FISH HATCHERY

2005 Steelhead Brood Year Report



By:
Jarrett L. Page, Fish Hatchery Assistant Manager
Jerry Chapman, Fish Hatchery Manager II
Doug S. Young, Fish Culturist
Kevin Yelton, Fish Culturist

IDFG 06-45
November 2006

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT	1
INTRODUCTION	2
OBJECTIVES	2
IDAHO DEPARTMENT OF FISH AND GAME GOALS	2
FACILITY DESCRIPTION.....	3
WATER SUPPLY	4
STAFFING	4
FISH PRODUCTION.....	4
Egg Shipments and Early Rearing	4
Final Production Rearing.....	6
Fish Distribution	7
FISH HEALTH	8
FISH MARKING	9
Fin Clipping, CWT, and PIT Tags	9
SPAWN TIMING MANIPULATIONS.....	10
RECOMMENDATIONS.....	11
Completed Improvements	11
Needed Improvements.....	12
Early Rearing and Incubation	12
Final Rearing	12
Employee Safety	12
Water Source	12
Building Improvements.....	13
LITERATURE CITED.....	14

TABLE OF CONTENTS (continued)

	<u>Page</u>
APPENDICES	15
Appendix 1. NSFH monthly water allocations	16
Appendix 2. Volume of water discharged from NSFH to Rim View Trout Company by month from November 2005 to May 20062006	16
Appendix 3. Results of annual analysis of NSFH source water, 1004 through 2005	17
Appendix 4. NSFH brood year 2005 steelhead survival from egg to smolt.....	17
Appendix 5. NSFH BY2005 feed usage	18
Appendix 6. NSFH brood year 2005 steelhead smolt distribution.....	19
Appendix 7. NSFH production costs for brood year 2005	19
Appendix 8. Fin lengths of NSFH steelhead; March 2006.....	19
Appendix 9. Fork length frequencies at release for four Coded Wire-tagged raceways; March 2005.....	20
Appendix 10. CWT summary for brood year 2005 steelhead at NSFH.....	21
Appendix 11. PIT tag summary for brood year 2005 steelhead at NSFH	21
Appendix 12. Comparison of mortality in vaccinated and non-vaccinated raceways at NSFH for brood year 2005	22
Appendix 13. NSFH history, BY66 to present	23
Appendix 14. Oxbow and Pahsimeroi stock spawn timing manipulations at NSFH for brood year 2005.....	24

ABSTRACT

Niagara Springs Fish Hatchery (NSFH) received 2,021,450 steelhead (*Oncorhynchus mykiss*) eggs and fry during the 2005 brood year. A total of 1,129,690 Pahsimeroi stock eggs and fry (555,023 eggs and 574,667 swim-up fry) were received from Oxbow Hatchery. After spawning, Pahsimeroi stock eggs were shipped green to Oxbow Hatchery for incubation on chilled well water and then transferred to NSFH as eyed eggs or swim-up fry. Just over half (50.87%) of the Pahsimeroi egg lots were shipped to NSFH as first feeding fry. This is a sixteen percent (16.29%) decrease from the previous year. A total of 891,760 Hells Canyon stock eggs and fry (443,987 eggs and 447,773 swim-up fry) were received from Oxbow Hatchery. Approximately 50% of the earliest Hells Canyon egg lots were shipped to NSFH as first feeding fry.

Total production for the 2005 brood year at NSFH was 1,812,874 steelhead (402,306 lbs) for anadromous smolt releases. No excess fish were available to be stocked as fall releases.

A total of 1,812,874 steelhead smolts (402,306 lbs at 4.51 fish/lb) were released into the Snake and Salmon rivers from March 20 to April 27, 2006. A total of 828,883 smolts (200,350 lbs at 4.14 fish/lb) of Pahsimeroi stock were released in the Pahsimeroi River at the weir, and 222,419 smolts (51,980 lbs at 4.28 fish/lb) of Pahsimeroi stock were released in the Little Salmon River near Hazard Creek. A total of 520,458 smolts (97,596 lbs at 5.33 fish/lb) of Hells Canyon stock were released in the Snake River at Hells Canyon Dam, and 241,114 smolts (52,380 lbs at 4.60 fish/lb) of Hells Canyon stock were stocked in the Little Salmon River near Hazard Creek.

Mortalities from pathogens increased this year. For this brood year, fifty percent (50%) of the total steelhead production was vaccinated for furunculosis (*Aeromonas salmonicida*). No steelhead at NSFH were vaccinated for enteric redmouth disease (ERM-*Yersinia ruckerii*) during this brood year. Furunculosis and ERM were not isolated during the 2005 brood year. Clinical bacterial coldwater disease, caused by *Flavobacterium psychrophilum*, was detected in July 2005. *F. psychrophilum* was detected again during preliberation sampling. Infectious Hematopoietic Necrosis Virus (IHNV) caused acute losses to one raceway during October.

Comment [d1]: Were?

A total of 381,758 lbs of fish feed were fed (378,722 lbs of Rangen and 3,036 lbs of Skretting (formerly Moore-Clark)) at a cost of \$150,541.83 to produce 402,306 lbs of steelhead for a conversion rate of 0.948:1.

Comment [d2]: Were?

Authors:

Jarrett L. Page, Fish Hatchery Assistant Manager

Jerry Chapman, Fish Hatchery Manager II

INTRODUCTION

The NSFH is owned and financed by Idaho Power Company (IPC), and operated and staffed by the Idaho Department of Fish and Game (Department). It is located in the Snake River Canyon, ten miles south of Wendell, Idaho. The NSFH is one of four hatcheries IPC owns and which the Department staffs and operates that fulfill IPC's mitigation requirement under the Federal Energy Regulatory Commission license #1971. The goal of NSFH is to rear 400,000 pounds (lbs) of steelhead smolts annually. Originally, these smolts were used to relocate a portion of the Snake River steelhead run into the Salmon River. Since 1980, 200,000 lbs of production are used to sustain a steelhead run below Hells Canyon Dam in the Snake River, and 200,000 lbs are stocked in the Salmon River drainage.

OBJECTIVES

The two major mitigation requirements that must be met at IPC's NSFH are to produce quality steelhead smolts to sustain steelhead trout runs in the Snake River below Hells Canyon Dam and in the Salmon River and its tributaries by successfully meeting these objectives:

1. Rear 200,000 lbs of quality steelhead smolts to be released in the Salmon River and its tributaries. The steelhead are to return as adults in sufficient numbers to provide quality sport fisheries in these waters and to supply sufficient broodstock (1,000 adults) to the Pahsimeroi Fish Hatchery for the collection of spawn for the next production cycle.
2. Rear 200,000 lbs of quality steelhead smolts to be released in the Snake River below Hells Canyon Dam. These are to return as adults in sufficient numbers to provide a quality sport fishery in the Snake River and to supply sufficient broodstock (1,000 adults) to the Hells Canyon Trap for the collection of spawn for the next production cycle.

IDAHO DEPARTMENT OF FISH AND GAME GOALS

1. Provide quality steelhead smolts to the Snake and Salmon rivers that will survive downstream migration and return as adults in sufficient numbers to provide a quality sport fishery in these waters and their tributaries.
2. Provide quality hatchery steelhead for supplementation where wild stocks of steelhead have diminished below desired levels and where managers feel quality hatchery steelhead would enhance the fisheries resource.
3. Enhance the genetic quality of hatchery stocks through management and hatchery practices that favor genetic variability.

FACILITY DESCRIPTION

Fish culture facilities at NSFH consists of an indoor nursery area, outdoor rearing raceways, and two flow-through settling ponds. Spring water supplies 42 upwelling incubators and 21 rectangular vats for the hatching and early rearing of fry. The nursery tanks provide 749.61 ft³ of hatching and early rearing space.

The outdoor rearing space consists of nineteen raceways (300-ft x 10-ft), (142,500 ft³), which are supplied by constant temperature, gravity flow, spring water. This allows for the production of 400,000 lbs of steelhead at a density index of less than 0.35 lbs/ft³/in as recommended by Piper (1982).

Two flow-through settling ponds (150-ft x 60-ft) are provided to remove settleable solids from the NSFH effluent. The settling ponds handle all the flow from the raceways and meet Environmental Protection Agency (EPA) guidelines for aquaculture discharge.

Dissolved nutrients are monitored on a biweekly basis to comply with the terms of a consent order between IPC and the Idaho Division of Environmental Quality. Samples of inflow, raceway effluent, and settling pond discharges to Niagara Springs Creek and Rimview Trout Company are collected using Sigma automated water samplers and sent to Rangen Aquaculture Research Center for analysis. Additional water analysis is performed monthly in accordance with the EPA National Pollutant Discharge Elimination System permit. Samples are collected only on inflow and discharge water. The Rangen Aquaculture Research Center conducts nutrient analysis, while the Department's Eagle Fish Health Laboratory completes analysis for total suspended solids.

Swim-up fry are hand fed in all nursery areas with some supplementation from the use of Ziegler belt feeders. Once the fish reach approximately 75 fish per pound (fpp) they are transitioned to the automated feed delivery system. Two moveable bridges span the outdoor rearing area. A total of 19 Nielsen automatic feeders are mounted on the bridges. The fish are fed by moving the bridges down the length of the rearing area and energizing the individual feeders on the control panels. Bulk feed is dispensed to the feeders by a conveyor supplied by two 20,000-lb storage bins with an associated fines separator.

Raceway cleaning is accomplished with an automatic air blower system. Three blower motors supply approximately 10 psi of air to weighted, perforated, airlines on the bottom side corner of each raceway. The resulting bubble screen creates a vortex of water currents that keep waste material suspended along the length of the raceways. While this system saves many hours of labor sweeping raceways, it is not completely efficient. Raceways are broomed daily to augment the air blower system and a power scrubber is also employed on a regular basis to remove excessive algal growth.

Buildings on the NSFH grounds include five residences. Three are wood-frame houses, one is a doublewide modular home, and one is a 16-ft wide mobile home. A 32-ft x 80-ft metal building contains an office, two incubator rooms, garage, shop, and feed storage room. Additional buildings include: two screen storage buildings (14-ft x 24-ft and 10-ft x 12ft), an open front shed (10ft x 30ft), and a masonry block chiller building (70-ft x 45-ft) which contains the chiller and blower-electrical room, a heated shop, and garage.

The NSFH staff is also responsible for care of the IPC-owned two-acre park adjacent to Niagara Springs Creek. It has a public, handicapped-accessible restroom, picnic tables, BBQ grills, and refuse containers.

WATER SUPPLY

In addition to NSFH, Niagara Springs supplies water to Rimview Trout Company, Niagara Springs Wildlife Management Area, and Idaho State's Pugmire Park. Niagara Springs' total flow is 220 cubic feet per second (cfs), which is divided into water rights by the four users.

The IPC has entered into an agreement with the four other users of Niagara Springs water whereby NSFH will receive water according to a stepped flow chart (Appendix 1). The NSFH has a water right of 132 cfs. A diversion canal was built by IPC in 2001, allowing up to 75 cfs of water to be discharged to Rimview Trout Company from November through April. This canal crosses onto Rimview property and attaches at their second-use head-ditch. Three slide gates located in the east and west settling basins regulate the volume of water discharged to Rimview. The volume of water discharged to Rimview this season is summarized by month in Appendix 2.

Water from Niagara Springs is a constant 59°F and flows by gravity to the incubators, nursery vats, outdoor raceways, fire hydrants, and irrigation system. Water quality is checked annually during the spring at the NSFH for herbicides, pesticides, heavy metal contaminants and normal water chemistry parameters (Appendix 3).

STAFFING

Four permanent personnel staff the NSFH. Jerry Chapman, Fish Hatchery Manager II, Jarrett Page, Fish Hatchery Assistant Manager, Doug Young, and Kevin Yelton, Fish Culturists, handle most operational duties. During peak work activities there are several Bio Aides throughout the year: Mike Anderson, Glen Owsley, and Lisa Brittan who assist the permanent staff with fish culture, maintenance, and other assignments.

FISH PRODUCTION

Egg Shipments and Early Rearing

Eggs and fry received at NSFH originate from broodstock trapping and spawning operations at IPC's Oxbow and Pahsimeroi hatcheries. To retard embryonic development, steelhead eggs spawned at Pahsimeroi Hatchery were transported by aircraft to Oxbow Hatchery for incubation in chilled wellwater (43°F) prior to delivery to NSFH. This procedure was done to control smolt size while minimizing the need to take fish off feed during the rearing cycle at NSFH. At the Oxbow Hatchery, a 70 horsepower chiller unit was utilized to chill 52°F wellwater to 43°F for incubation. Pahsimeroi Hatchery does not have a chiller unit for this purpose. The NSFH received both eggs and fry for the 2005 brood year (Appendix 4). Of the

Pahsimeroi stock eggs incubated at Oxbow Hatchery and destined for NSFH, approximately one-half (49.13%) were delivered to NSFH as eyed eggs while the remaining (50.87%) were delivered as first feeding fry. This brood year saw an 16.29% increase in eyed eggs and subsequent decrease of fry received at NSFH. This was created due to concerns over water quality and amounts of water delivered to the incubation building. Eyed eggs were transported in conventional coolers while fry were transported in specially designed, perforated fry transport tubes; and placed inside a 500-gallon fish-stocking tank acquired by NSFH from Hagerman State Fish Hatchery (HSFH).

Comment [13]: Here you have two numbers in the same sentence which should add to 100%. One number has 2 decimal places and the other has none. There should be a consistent number of decimals in both cases. I suggest eliminating the .03.

The 500-gallon tank is mounted to a trailer and towed by a 3/4-ton hatchery vehicle. The tank is equipped with Point-4 ceramic air stones and bottled oxygen. This tank and trailer design was created in 2002 to help prevent the possible transmission of disease pathogens between hatcheries. This tank/trailer configuration also eliminated the transportation bottleneck created when NSFH wanted to use HSFH transport trucks during their busy "catchable" stocking season.

The NSFH received 555,023 eyed-eggs of Pahsimeroi stock between June 7 and June 9, 2005. This is one week earlier than the prior brood year and came from lots 6-9. A total of 574,667 Pahsimeroi stock swim-up fry from egg lots 1-5, were received from July 5 through July 14. The last of the fry were received 9 days earlier than the prior brood year. A total of 1,129,690 eggs and fry of the Pahsimeroi stock were received at NSFH for the 2005 brood year.

A total of 443,987 eyed-eggs of Hells Canyon stock (lots 10-15) were shipped to NSFH from May 27 through June 7, 2005. Egg lots 1-9 were transported to NSFH as swim-up fry (447,773) between June 16 and July 7, 2005, and placed directly into outdoor nursery raceways. The timing of the fry shipments coincided with the swim-up timing of fish from the eyed-eggs that were received earlier at NSFH, and was approximately the same as the prior brood year. A total of 891,760 eggs and fry of Hells Canyon stock were received at NSFH for the 2005 brood year. NSFH's steelhead egg to smolt survival for brood year 2005 is summarized in Appendix 4.

Upon arrival at NSFH, all eggs were tempered and disinfected with iodine at 100-ppm for 30 minutes and placed in upwelling incubators (25,250 per incubator average) inside the vats. All fry shipping containers were disinfected prior to shipping. Upon arrival at the hatchery, fry were tempered in their shipping containers before ponding.

Fry were not inventoried from the nursery vats to the nursery raceways this brood year. Consequently, hatching success and mortality could only be estimated. The NSFH staff observed normal survival to eyed eggs during the early rearing cycle. Survival of fry to fingerling was 99.34% in Pahsimeroi steelhead and 95.76% in Hells Canyon steelhead. Overall, fry to fingerling survival was 97.78%.

As was done last year, nursery sections were screened at both ends and remained expanded from 20-ft to 35-ft in length. This effectively prevented fry and fingerlings from getting into the headrace, and allowed the fry to be reared at lower starting densities. As densities increased, fry were given more rearing volume by relocating screens to 50-ft, 75-ft, and finally, to the end of the first section at 100-ft. In 2001, additional screen brackets were placed at 75-ft and 150-ft allowing for greater flexibility than the standard keyway distances of 100, 200, and 300-ft. This methodology reduces wasted feed and reduces cleaning times created by using the standard keyway distances, yet still allows densities to be lowered and greater management of inventories. This practice was expanded even more, during the 2003 brood year, by creating

screen brackets at 250-ft for the Coded-Wire-Tag (CWT) raceways. These CWT raceways have approximately 30,000 to 78,000 untagged steelhead per raceway, depending on the stock and tagging requests. In the production cycle, these 30,000 to 78,000 fish require less or more than the normal 100-ft key-way sections depending on the number. This allows for better cleaning, feeding and density control. Fish are given more room before they attain a density index of 0.30. All of these fish culture changes will continue in future brood years.

Comment [14]: Can you provide some information on the target density index that was used or a maximum not-to-exceed density figure that might help the reader understand how you manage your inventories. In other words at what DI do you move the screens and give the fish more room?

Throughout the entire early rearing period, steelhead at NSFH were fed Rangen dry feeds. Feed was dispensed by hand feeding and supplemented with Ziegler belt feeders in the indoor and outdoor nursery areas. When the fingerlings reached approximately 275 fish/lb 50% of the population was fed Skretting's "Clarks Fry" (ProActive) dry feeds for 14 days. The purpose of feeding Skretting's ProActive feed was to stimulate the fry's immune system prior to the vaccination program. When they reached 75 fish/lb, all NSFH fish were switched to a Rangen extruded diet. The switch to Rangen bulk extruded feed allowed NSFH staff to utilize the bulk tanks, feed conveyor system, fines separator, and bridge feeders.

Final Production Rearing

Adipose fin-clipping operations are used to split the fish into even-numbered and odd-numbered raceway sections. During this program, fish are crowded to the lower 100-ft section of each odd-numbered raceway. Half the fish are clipped and put into the upper two-thirds of the raceway, while the other half are clipped into the adjacent even-numbered raceway. Fin-clipping operations started on September 19 and were completed by October 12, 2005.

Fish were given the final 100-ft of rearing space in early January. Hells Canyon fish were placed in raceways 1 through 8, while Pahsimeroi fish were placed in raceways 9 through 19. Normal fish culture techniques during this time include: feeding fish with the bridge, sweeping raceways, conducting sample counts, cleaning screens and air lines, removing mortalities, equipment maintenance, record keeping, nutrient sampling, pond scrubbing, supervising and running the adipose-marking trailer, length frequency and fin quality collection and reporting, assisting with CWT and passive integrated transponder (PIT) tagging operations, and conducting tag and mark retention checks.

Pahsimeroi steelheads were held off feed for 6 days to slow their growth, and 14 days for other fish culture reasons. Hells Canyon steelhead were off feed for 4 days to slow their growth and 12 days for other fish culture reasons. These reasons include; off feed prior to handling for vaccination, adipose fin clipping, CWT and PIT-tagging programs, and off feed prior to shipping. This is a dramatic decrease from prior years, since it was not uncommon to have fish off feed for up to 45 days. Although early growth rates exceeded 0.033 inches per day, growth rates were slowed to 0.021 inches per day. Slowing the growth rates has been accomplished through feeding practices and reducing the amount of feed fed per raceway per day, but still continually feeding the fish. Past practices reduced growth rates by holding fish off feed for 5 days at a time.

A total of 378,722 lbs of Rangen and 3,036 lbs of Skretting were fed over the course of the brood year (Appendix 5). The Rangen feed total includes 29,750 lbs of Oxytetracycline (OTC) medicated feed used for two medicated feed treatment during this brood year. OTC was fed allowing for a 21-day withdrawal time prior to stocking, meeting Food and Drug Administration (FDA) requirements.

The total cost of the OTC feed was \$10,605.55. The total cost of regular feed was \$139,831.92. A total of 402,306 lbs of fish were produced on lbs of feed for a conversion rate of 0.948:1. Total NSFH production costs incurred by IPC during the 2005 brood year were \$892,728.72 which includes IPC overhead, smolt hauling, and shop expenditures, but does not include capital outlay expenditures. The cost/lb of fish produced was \$2.22 (Appendix 7).

Fin quality was assessed in March using methods developed by the NSFH personnel. NSFH personnel had previously performed fin quality measurements in March and April. However, alterations to the stocking schedule this year dictated that all fin quality measurements be taken in March to allow NSFH staff time to meet the FDA required guidelines for withdrawal periods on fish treated with MS-222 anesthetic.

Fins of steelhead reared at NSFH were compared to fins of wild, outmigrating steelhead collected from the Salmon and Pahsimeroi Rivers over a two-year period. The resulting fin index (0.1185) will be used for all fin quality measurements at NSFH.

A total of 80 steelhead, from all four of the CWT-tagged raceways, were used for this comparison. After measuring the lengths of the dorsal and pectoral fins, a fork length was taken from each fish. By comparing the average fin length to the average fork length, a fin quality index was calculated. This index was then compared to that of wild steelhead. Results indicate that the fin quality index from fish raised at NSFH was 70.00% of that of wild fish (Appendix 8). This was a 3.42% decrease from the 2004 releases using the fin index based on wild steelhead.

A target smolt size of 170 to 250 mm fork length has been established by NOAA Fisheries to maximize smolt out-migration and minimize the potential for predation by hatchery steelhead on wild salmon. To demonstrate compliance with these criteria, length frequency data were collected prior to shipping to determine fish size at the time of release (Appendix 9). The average length of CWT tagged fish sampled from four raceways in March was 207.89 mm (8.18 inches).

Fish Distribution

The IPC contracted with Neil Ring Trucking of Buhl, Idaho, to transport steelhead smolts to release sites using three IPC tank trailers. Transport of steelhead from NSFH began on March 20 and ended on April 27, 2006. Eighty loads of steelhead (402,306 lbs at 4.51 fish/lb) were transported to the Snake and Salmon rivers (Appendix 6). The first fish were transported to Hells Canyon (Hells Canyon stock), then to the Little Salmon River near Hazard Creek (Hells Canyon stock and then Pahsimeroi stock), then to the Pahsimeroi River below the weir, and finally back to the Little Salmon River to meet requests. The Hazard Creek release site was utilized because of reduced space and high snow levels at Stinky Springs. Department biologists feel that Pahsimeroi fish do better if stocked after the second week in April (Kent Ball, personal communication).

Steelhead smolt release figures are as follows; Snake River at Hells Canyon Dam (Hells Canyon stock): 520,458 fish (97,596 lbs at 5.33 fish/lb); Little Salmon near Hazard Creek (Hells Canyon stock): 241,114 fish (52,380 lbs at 4.60 fish/lb); Pahsimeroi River below the weir (Pahsimeroi stock): 828,883 fish (200,350 lbs at 4.14 fish/lb); and the Little Salmon near Hazard Creek (Pahsimeroi stock): 222,419 fish (51,980 lbs at 4.28 fish/lb) (Appendix 6). Total

Comment [15]: Can you explain why the 9.23% increase occurred? What did you do different this year?

Comment [16]: Jerry, Paul, Unless you are doing some additional statistical analysis that I am not aware of I don't think this sentence is accurate. There is no statistical test that can be done to determine whether Henrys Fork rainbow or wild steelhead smolts are a better point of comparison. As I recall we determined that the new fin quality index using wild steelhead was a better point of comparison for 3 reasons: it compares steelhead to steelhead rather than steelhead to resident trout (apples to apples instead of apples to oranges); you have a larger sample size of wild smolts; and the wild steelhead are closer in size to the hatchery steelhead than the resident trout. This was not really determined statistically. Rather it was just a objective assessment that one is a better point of comparison than the other.

Comment [d7]: Total lbs. of fish released is not mentioned in Appendix 4 (NSFH Steelhead Survival from Egg to Smolt). Only the total # of fish released is mentioned. Should this reference be to appendix 6?

Comment [d8]: This is listed as 4.94 fish/lb in the abstract. Please keep decimal places consistent?

Pahsimeroi production was 252,330 lbs, or 1,051,302 steelhead smolts at 4.17 fish/lb, and total Hells Canyon production was 149,976 lbs, or 761,572 steelhead smolts at 5.07 fish/lb. Total NSFH production for the year was 402,306 lbs, or 1,812,874 fish.

Total survival to release was 93.06% for Pahsimeroi steelhead, while total survival to release for Hells Canyon steelhead was 85.40%. Overall, combined survival to release for NSFH steelhead smolts was 89.68% (Appendix 4).

Comment [d9]: Is this meant to be an arithmetic mean or a weighted mean? If you just add the 2 percentages and divide by 2 the number I get is 90.72%. It does not appear that you used weighted means elsewhere so I suspect an error here. Please check you math.

FISH HEALTH

Fish health is always a concern at NSFH. The location of NSFH, in the heart of the commercial trout industry, makes it vulnerable to the horizontal transmission of many etiologic agents. Disease problems from IHNV, bacterial furunculosis, ERM, and bacterial coldwater disease (CWD) have caused significant losses in years past (Munson, 1996). In addition, the NSFH and its spring-water source are located directly below agricultural land, exposing both to toxic drift and runoff from chemical application to fields above NSFH. Stringent sanitation programs and fish culture practices are implemented to facilitate disease control.

Fifty percent of the brood year 2005 fish were vaccinated with an autogenous *Aeromonas salmonicida* bacterin obtained from Aqua Health Limited. Fish were dipped in an oxygenated solution of 18 liters of water to 2 liters of vaccine with a one-percent (1%) salt solution incorporated into the vaccination solution. The salt solution was introduced to the vaccination protocol to reduce stress brought about by physical handling and to increase the uptake of vaccine by the fish. Vaccine was applied at a rate of 220 lbs of fish per liter of vaccine, for 40 seconds. The vaccination program started on August 24 and ended on September 13, 2005. Average fish size at the time of vaccination was 112.83 fish/lb. Because Furunculosis has not been a problem in recent years (the last clinical diagnosis was brood year 1998), NSFH decided to vaccinate only 50% of the steelhead population again during the 2005 brood year. Fish were chosen for vaccination that represented both egg and fry transfers to NSFH, from both stocks of fish. No conclusions were made about the non-vaccinated vs. vaccinated fish health since there was no epizootic during this brood year. Mortalities were recorded on all the raceways each month after vaccination until shipping (Appendix 12). Burton (2005) suggests that hatchery staff reinstate 100% vaccination to protect NSFH due to Rim View Hatchery discontinuing the use of all antibiotics to market their product as organic.

Mortality for the year was above normal. Acute losses due to infectious agents were experienced in one raceway at NSFH. Acute losses to IHN were limited to raceway 6 during October. Clinical bacterial coldwater disease was detected in July 2005 in one raceway of Hells Canyon steelhead. Treatment with OTC-medicated feed was administered with good results. Minor losses were attributed to Motile Aeromonad Septicemia (MAS), (*Aeromonas sp.*). An application of OTC - medicated feed was administered prior to release to reduce mortality due to CWD being detected during preliberation necropsy. Fish were treated for 10 days with 4% OTC incorporated into the feed in accordance with FDA Investigational New Animal Drug #9332 requirements.

The organosomatic index showed normal values in all categories for both Pahsimeroi and Hells Canyon stocks. Blood work was also taken on both stocks of steelhead at NSFH. Parameter levels for leukocrit and serum protein were normal for the Hells Canyon stock. Mean

hematocrit levels and serum protein for the Hells Canyon raceway sampled were 44.3% and 4.76% respectively (Burton 2006).

Mean hematocrit levels and serum protein for the Pahsimeroi raceway sampled were 41.4% and 4.66% respectively (Burton 2006). CTL and KTL (metric and English condition factors) were not assayed for this brood year. The condition of fish from both Hells Canyon and Pahsimeroi stocks at liberation was good.

Comment [d10]: What do these stand for?

Furunculosis, and ERM were not isolated again at this facility during the 2005 brood year. A continuing aggressive disease management program at this facility has been effective in controlling mortality due to these etiological agents. During the first 31 years of production at NSFH, (1966-1997), the yearly mortality averaged 30.70%. During the last 5 years of production at NSFH, the average yearly mortality has declined to 10.37%, a decrease of nearly 3 times from the first thirty-one years of production (Appendix 13).

Comment [d11]: Does this stand for Enteric Red Mouth? If so, it is not mentioned at all before this. Should it be added to the first paragraph that talks about the diseases that can be problems at NSFH?

As NSFH reduces egg request numbers and the stated management practices of the inventory continue, a reduction in losses due to MAS and CWD should occur. Hatchery personnel will need to investigate some of the new feeds breaking into the market. These feeds might help fin quality, survival, or any number of fish health parameters while balancing gains verses costs (Munson 2003). Hatchery staff should also decide whether to continue current vaccination protocol or to resume 100% vaccination (Burton 2006).

Comment [d12]: Appendix 12 only refers to BY03 mortality comparisons between vaccinated and unvaccinated raceways, not to the last 31 years of mortalities. This is not an appropriate reference?

FISH MARKING

Fin Clipping, CWT, and PIT Tags

All hatchery-reared steelhead in the state are marked with an adipose fin clip. Adipose fin clipping is done so that anglers can differentiate between hatchery and wild steelhead. Steelhead were adipose-fin clipped at NSFH between September 19 and October 12, 2005.

Brood year 2005 steelhead were implanted with CWTs from September 28 to October 1, 2005. A total of 164,341 steelhead received CWTs (65,458 Hells Canyon stock, and 98,883 Pahsimeroi stock). Each tag group was held in an individual raceway section so that separate mortality information could be gathered. The CWT groups of 30,000 fish were given a 100-ft section and the one raceway with 60,000 (RCWY 12) was given 200 ft as densities increased. The fish with CWTs (Hells Canyon and Pahsimeroi stock) that were destined for the Little Salmon River were again moved into raceways closer together so that they could be shipped within days of each group. In prior years, the Pahsimeroi stock of fish that were destined for the Little Salmon River were shipped approximately a month later than the Hells Canyon stock fish. This continued practice should allow better evaluation of stock performance, as well as some possible insight into travel times for each stock, and comparisons into prior years' downstream migration times. The first fish that could be evaluated would be the returning adults in 2006, and then every succeeding year.

Comment [d13]: If you take the total number of fish that received CWTs (156,124 fish) and subtract 30,000 multiple times, you end up with 66,124 for the final raceway, which is a difference of 3,945 fish from the number provided (62,179 fish). If the 30,000 fish put into the other raceways is approximate, then maybe that word should be added to the text. It seems odd though that one number would be exact (62,179) and the others (30,000) would be approximate.

A total of 158,051 CWT-tagged fish were released at three release sites (Appendix 10). A total of 32,254 CWT-tagged fish were released in the Snake River at Hells Canyon Dam from March 27 through March 28, 2006, while 64,035 CWT-tagged fish were released at the Pahsimeroi weir (Pahsimeroi River) between April 12 and April 13, 2006. Both Hells Canyon stock, and Pahsimeroi stock, were released in the Little Salmon River near Hazard Creek for the 2005 brood year. A total of 30,422 CWT-tagged Hells Canyon stock were released at Hazard

Creek between April 3, and April 5. The Pahsimeroi stock were released at Hazard Creek between April 5, and April 7, 2006 and totaled 31,340 CWT- tagged fish.

In addition to the CWT-tagged fish, 1,196 fish were tagged with PIT tags on February 27 and 28 2006 (raceways 4, 8, 9, and 13). These computer chips are injected into the body cavities of the fish and information can be accessed as to hatchery origin, length, weight, release watershed, date of release, downstream migration, timing, and travel rates. In this manner, an individual fish can be tracked on its seaward migration without sacrificing the fish.

Comment [PEA14]:

All mortalities were scanned for PIT-tag detection after tagging had occurred, prior to release and during release. Only two (2) mortalities of PIT-tagged fish occurred after the tagging was completed and prior to release. Consequently, a total of 1,194 PIT-tagged fish were released from NSFH for the 2004 brood year. Of these, a total of 299 PIT-tagged fish were released below Hells Canyon Dam (Hells Canyon stock), while 299 PIT-tagged fish (Hells Canyon stock) and 300 (Pahsimeroi stock) were released in the Little Salmon River near Hazard Creek. In addition, 298 PIT-tagged fish were released at the Pahsimeroi weir in the Pahsimeroi River (Pahsimeroi stock) (Appendix 11).

SPAWN TIMING MANIPULATIONS

Several years ago, the Department consulted University of Idaho geneticist Dr. Madison Powell for recommendations on the proper methods to move the spawn timing back to historical spawning times. Dr. Powell suggested that 10% of early-spawning steelhead and 100% of late-spawning steelhead be spawned if Department personnel desire to move the spawn timing back without impacting the existing program. Department personnel at Oxbow and Pahsimeroi hatcheries attempted to employ these recommendations again this year and will continue to do so in future years. While a high percentage of the early spawn takes were kept, the numbers were small compared to the larger lots kept later (Appendix 14).

Comment [d15]: Matt's full name is actually Madison.

Comment [d16]: Appendix 13 is a summary of NSFH production history from BY66 to present. Please check your numbering on this and all references to appendices.

Approximately 21% of the eggs shipped to NSFH from early spawning adults (3/31-4/13) were utilized for smolt production destined for the Pahsimeroi River, while 17% of eggs from the middle of the spawn (4/18-4/28) and 62% of eggs from late spawning fish (5/2-5/12) were utilized for smolt production at the Pahsimeroi River. At Oxbow Hatchery, 48% of early spawning fish (3/14-4/9) were utilized for smolt production back to Hells Canyon Dam, while 35% of the middle of the spawn (4/18-4/25) and 17% of eggs from late spawning fish (4/28-5/5) were utilized. Five lots of fish were spawned and retained at Oxbow Hatchery during March, while only one lot from March (31) was retained for NSFH at the Pahsimeroi Hatchery.

Comment [d17]: Utilized where? If they were utilized for smolt production at the Pahsimeroi River, then shouldn't "for smolt production at the Pahsimeroi River" come at the end of the sentence?

Comment [d18]: I did not see any reference to Appendix 14 in the text. This seems to be the appropriate spot in my opinion.

Comment [d19]: Again, utilized where? If they were utilized for smolt production back to Hells Canyon Dam, then shouldn't "for smolt production back to Hells Canyon Dam" come at the end of the sentence?

Spawn timing may also be directly correlated to winter river water temperatures. Colder winter river temperatures may delay spawning by delaying egg development in the female based on the temperature unit philosophy. A correlation could exist to associate winter river water temperature with ensuing spawn timing in conjunction with artificially moving the spawn timing back by choosing eggs from later females. In the future, hatchery managers may be able to predict when the bulk of spawning will occur based on winter river water temperatures and spawn timing manipulations from preceding years.

RECOMMENDATIONS

Completed Improvements

Several hatchery improvement projects were completed this past year. A layer of gravel was added to the road from NSFH to Crystal Springs Fish Hatchery as part of a long-term maintenance plan. IPC entered a cost-share agreement with State Parks and Clear Springs Feed Company for this work. Dust-Gard solution was applied again this year to the main springs road to minimize dust and reduce road "washboards". A locksmith fixed all the door locks around the hatchery and replaced the doorknobs on the chiller building main doors. New latches were installed on all the fry tubes. The birdnetting was repaired several times. The interior of the park restroom was repainted to cover numerous vandalism inscriptions. The interiors of residence one and three were painted prior to new employees moving into them. The carpets were also professionally cleaned. Rocks were placed around the well casing as instructed by the safety inspector. Two new miniature gram scales were purchased for hatchery use, and new air lines were purchased to replace one third of the old air lines in the raceways. The feed room air conditioning unit was moved to the exterior of the building to work more efficiently.

A new smolt hauling tanker was purchased by IPC, and a used forklift was received from the Malad IPC shop for use at the hatchery. All personnel received professional training to operate the forklift. The forklift clutch and other transmission problems were repaired. Idaho Power Company also purchased a new leaf collection system for the lawn tractor and installed a safety gate on the feed separator as required by the safety inspector. New central air systems were installed in all three wood frame houses, and a new cooling system was installed in the blower room of the chiller building. IPC shop personnel also repaired the bridge feeder electrical problems and tightened the bridge cable reel. An electrician was hired to rewire the clipping trailer according to electrical specifications. New outlet covers were installed on all the outlets while the trailer and steps were properly grounded to the birdnetting fence posts. Overhead indoor outlets were rendered useless to prevent electrical malfunctions. New floor mats and dip nets were purchased for the clipping trailer.

Several landscaping projects were also completed this past year. All trees along the entrance and spring roads were trimmed for better visibility. Trees were trimmed along Rimview's intake canal on IPC property at Rimview's request. Both hatchery and professional crews removed dead trees around the hatchery. Numerous live elm trees were removed around the intake to reduce the leaves and seeds that plug our raceway intake screens. Numerous sprinkler heads were replaced for better lawn coverage. Kimberly Nurseries repaired broken buried sprinkler lines and changed broken solenoids. Redwood stain was applied to all the parking barrier logs in the park, and the yellow bollards around the property were repainted. Landscaping rock was applied to new areas around the hatchery to make mowing easier and added to existing rock areas. Soil-pep was added around all the trees and bushes on the property, and wildflowers were planted on the border with Rimview Trout Company. Grass was replanted in many areas. Weeds were sprayed in the spring and fall, and fertilizer was applied to all the grounds and park in the spring. New trees were sprayed to prevent coddling moth infestations.

Needed Improvements

Early Rearing and Incubation

An expansion of the present nursery facility to at least twelve times its present size is needed. The number of vats should be based on a desired density index of 0.30 at a fish size of 200 fish/lb or 2.5 inches in length. Using these criteria, there should be at least 15,120 ft³ of rearing space to ensure adequate rearing space for fry. This system would protect fry from bird predation and provide them with shade from the sun.

Comment [d20]: Does he mean vats? They are referred to in the facility description as vats.

Final Rearing

Concrete repair work needs to be completed at 300' on all the raceways, with additional repairs on some raceway walls. Galvanized keyways should be installed to replace the severely rusted steel keyways in the raceways. The birdnetting will have to be replaced in the near future because it requires more maintenance and is starting to deteriorate. The large bridge needs to be professionally sandblasted and painted, and a diesel-powered compressor needs to be purchased so hatchery personnel don't have to borrow the IPC shop compressor or rent one twice per month.

A fish counter is needed for better inventory accuracy and management. Other adipose fin clipping equipment needed includes a hydraulic pump unit, and 4 inch pump, 170 feet of 4 inch flexible line with cam locks on both ends, and a new fresh-flow pump for backup water to the trailer. A Kawasaki "Mule" or Cushman is needed for quick trips around the hatchery to haul equipment and save wear and tear on the trucks and mowers.

Employee Safety

A trash rack needs to be installed in front of the intake gate at the upper pool to prevent access to the spring and injury to the public. A trash rack should also be installed at the entrance to the discharge canal to Rimview Hatchery.

The bulk tank, conveyor line, and entrance gates to the outdoor raceways need to be raised. This is a safety issue as the low height of the conveyor line and gate doorways have caused numerous bumps and bruises.

Water Source

An intake-traveling screen is needed to remove leaves and macrophytes from the water at our intake. Entire raceways are in jeopardy every fall when leaves plug head screens during the night. Weeds also break loose from the springs all year long that could plug our head screens.

Building Improvements

A new hatchery and incubation building with functional nursery vats is badly needed. The building should also include public restrooms that are handicapped accessible, an office, shop, meeting room, and an adequate feed storage space. A three-stall garage for the trucks and mowers would be beneficial to protect these items from vandalism and weather.

Comment [d21]: Does he mean vats?

Sliding glass doors should be installed in the living rooms of the three wood frame houses to allow access to decks and improved access to the outside for fire safety. Bathroom remodeling in all the wood-frame houses is needed, along with some kind of awning or roof over the outside decks for the two residences with the most sun exposure.

LITERATURE CITED

- Burton, D. 2005. Niagara Springs Fish Inspection Report. Department Eagle Fish Health Laboratory. 1800 Trout Road, Eagle, Idaho.
- Munson, A. D. 1996. Niagara Springs Fish Inspection Report. Department Eagle Fish Health Laboratory. 1800 Trout Road, Eagle, Idaho.
- Munson, A. D. 2001. Niagara Springs Fish Inspection Report. Department Eagle Fish Health Laboratory. 1800 Trout Road, Eagle, Idaho.
- Munson, A. D. 2003. Niagara Springs Fish Inspection Report. Department Eagle Fish Health Laboratory. 1800 Trout Road, Eagle, Idaho.
- Munson, A. D. 2004. Niagara Springs Fish Inspection Report. Department Eagle Fish Health Laboratory. 1800 Trout Road, Eagle, Idaho.
- Piper, R.G., I.B. McElwain, L..E. Orme, J.P. McCraren, L.G. Fowler, and J.R. Leanard. 1982. Fish Hatchery Management. U.S. Fish and Wildlife Service, Washington, D.C.

APPENDICES

Comment [122]: I notice that the captions for all of the appendices are capitalized as if they were titles. Example: Appendix 1. NSFH Monthly Water Use Allocations. General writing style manuals say that these captions should read just like a sentence with only the first word and proper names being capitalized. So the proper way to write the caption for Appendix 1 would be... Appendix 1. NSFH monthly water use allocations. Please correct all appendices to follow this format.

Appendix 1. NSFH monthly water use allocations.

Month	Max. Flow	Month	Max. Flow
May	50 cfs	November	70 cfs
June	50 cfs	December	90 cfs
July	50 cfs	January	100 cfs
August	50 cfs	February	110 cfs
September	50 cfs	March	120 cfs
October	60 cfs	April	120 cfs

Appendix 2. Volume of water discharged from NSFH to Rimview Trout Company by month from November 2005 to May 2006.

Month	Hatchery Inflow	Flow to Rimview	Discharge Flow to Niagara Springs Crk.
November	70 cfs	0 cfs	70 cfs
December	90 cfs	25 cfs	65cfs
January	90 cfs	25 cfs	65 cfs
February	110 cfs	60 cfs	50 cfs
March	120 cfs	70 cfs	50 cfs
April	80 cfs	45 cfs	35 cfs
May	30 cfs	0 cfs	30 cfs

Comment [123]: There is not set allocation for the volume of water sent to RimView. It will vary annually based on spring flows. Therefore this caption should be changed as noted.

Appendix 3. Results of annual analysis of NSFH source water, 1994 through 2005.

Analysis	Yearly Results							Maximum Allowable Contaminant Levels
	1994 (mg/l)	2000 (mg/l)	2001 (mg/l)	2002 (mg/l)	2003 (mg/l)	2004 (mg/l)	2005 (mg/l)	
Alkalinity	166	170	180	170	170	160	170	10.0
Antimony	0.002	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.006
Arsenic	0.005	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.05
Barium	0.180	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	2
Beryllium	0.0002	N/T*	N/T*	N/T*	N/T*	N/T*	N/T*	0.004
Cadmium	0.00034	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.005
Chromium	0.002	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.1
Chloride	N/T	46	44	45	41	46	44	250
Copper	0.010	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	1.3
Cyanide	0.005	N/T*	N/T*	N/T*	N/T*	N/T*	N/T*	0.200
Fluoride	0.570	0.6	0.7	N/D*	0.6	N/D*	N/D*	4
Hardness	234	230	230	240	220	230	230	100
Iron	0.010	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.3
Lead	0.002	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.015
Manganese	N/T	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.05
Mercury	0.0002	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.002
Nickel	0.003	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.1
Nitrate as N	1.630	1.6	1.7	1.8	1.4	1.8	2.0	10
Nitrite as N	0.01	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	1
PH	8.00	8.4	8.2	8.2	7.9	8.1	8.0	6.5 - 8.5
Selenium	0.005	N/D*	N/D*	N/D*	N/D*	N/D*	N/D*	0.05

*N/D = Not detected

*N/T = Not tested

Appendix 4. NSFH brood year 2005 steelhead survival from egg to smolt .

Source	Eggs Received	Fry Received	Total Received	Fingerlings Released	% Survival Fingerlings	Smolts Released	Total Release	% Survival To Release
Pahsimeroi	555,023	574,667	1,129,690	0	*99.34%	1,051,302	1,051,302	93.06%
Oxbow	443,987	447,773	891,760	0	*95.76 %	761,572	761,572	85.40%
Totals	999,010	1,022,440	2,021,450	0	*97.78%	1,812,874	1,812,874	89.68%

*Estimated percentages

Appendix 5. NSFH brood year 2005 feed usage.

Manufacturer	Type and Size	Dates Received	Total Pounds Received	Total Pounds Used	Total Feed cost (\$)
			2,550		
Rangen	Trout and Salmon Starter Swimup	6/15/05-8/4/05		2,550	1,364.01
Rangen	Trout and Salmon Starter #1	6/15/05-8/18/05	2,000	2,000	1,087.18
Rangen	Trout and Salmon Starter #2	7/5/05-8/29/05	3,650	3,650	1,975.33
Rangen	Trout and Salmon Starter #3	7/26/05-9/23/05	4,550	4,550	2,452.81
Moore Clark	Clark's Fry 1.5mm ProActive	8/04/05	3,036	3,036	1,757.05
Rangen	Bulk 470 exsl. 2.0mm slow-sink	8/08/05-10/24/05	43,600	43,600	17,574.14
Rangen	Bulk 470 exsl. 3/32 slow-sink	10/28/05-1/05/06	100,860	100,860	40,666.70
Rangen	Bulk 470 exsl. 1/8 slow-sink	1/16/06-4/03/06	194,820	194,820	
Rangen	Sack Trout Production TM (8000g) 1/8 pellet	2/15/06-3/24/06	29,600	29,600	10,406.20
Rangen	TM Medicated Feed 4000g #2	7/22/06	100	100	62.12
Rangen	TM Medicated Feed 8000g #2	7/20/05	50	50	137.23
Rangen	Fines credit "exsl" bulk / sack	7/22/05, 5/5/06	-2,780	-2,780	1,142.51
Rangen	Fines credit "exsl" Bulk	7/13/06-3/17/06	-578	-578	209.43
Grand Total			381,758	381,758	\$150,541.83

Appendix 6. NSFH brood year 2005 steelhead smolt distribution.

Destination	Stock	Weight	Dates	Number Per Pound	Number Released
Hells Canyon (Snake R.)	H.C.	97,596	3/20-3/30	5.33	520,458
Hazard Creek (Little Salmon R.)	H.C.	52,380	3/30-4/5	4.60	241,114
Pahsimeroi (Pahsimeroi R.)	Pah.	200,350	4/7-4/25	4.14	828,883
Hazard Creek (Little Salmon R.)	Pah.	51,980	4/5-4/7,4/26-4/27	4.28	222,419
Total		402,306		4.51	1,812,874

Appendix 7. NSFH production costs for brood year 2005.

Number of Fish	Lbs of Feed	Cost of Feed	Pounds of Fish	Feed Conversion	Total Cost	Cost per 1,000	Cost per Pound
1,812,874	381,758	\$150,541.83	402,306	.948	\$892,728.72	\$492.44	\$2.22


*Cost includes IPC cost for overhead, smolt hauling and shop expenditures and does not include capital outlay expenditures.

Appendix 8. Fin lengths of NSFH steelhead; March 2005.

Raceway	Fork Length	Right Pectoral	Left Pectoral	Dorsal	Ave. Fin Length	Fin Quality Index	Wild Fin Quality Index	Percent of Wild FQI
4	196.3	19	19.8	14.4	17.7	.090	0.1185	76%
8	198	20	19.4	15.8	18.35	.093	0.1185	78%
9	203.1	16.7	18.6	11.4	15.53	.077	0.1185	65%
12	221.4	18.8	18.0	11.6	16.14	.073	0.1185	62%
Average						.083	0.1185	70%

Appendix 9. Fork length frequencies at release for four Coded Wire-tagged raceways;
March 2005.

Raceway #	H.C. 4	H.C. 8	Pah. 9	Pah. 12
Sample Size	100	100	100	100
Ave. Frk. Length	197.46	203.14	203.3	224.52
Lower Range (mm)	115.00	98.00	94.00	94.00
Upper Range (mm)	243.00	261.00	258.00	274.00
	(mm)	(Inches)		
Hells Canyon Average Length	201.87	7.95		
Pahsimeroi Average Length	213.91	8.42		
Overall Average Length	207.89	8.18		



Comment [124]: In this graph several corrections are needed. 1. you forgot a % sign on the 0.5>250mm. 2. There is only one set of data therefore the legend which explains # of fish are presented in blue is not necessary there is no other color to confuse it with. 3. The title should be deleted and a caption inserted listing this graph as appendix 10.

Appendix 10. CWT summary for brood year 2005 steelhead at NSFH.


Raceway	Release Site	CWT Number	Number Tag	Mortality to Release	Number Shed	CWT Number Released	Untagged	Total Tagged Group Release	Total Site Release
Snake River									
4	Hells Canyon Dam	10-80-77	33,005	421	330	32,254	78,829	32,254	
Total			33,005	421	330	32,254	78,829	32,254	111,083
Salmon River									
8	Little Salmon River	10-81-77	32,453	279	1,752	30,422	53,000	30,422	
Total			32,453	279	1,752	30,422	53,000	30,422	83,883
Pahsimeroi River									
12	Pahsimeroi Trap	10-92-77	66,279	587	1,657	64,035	32,925	64,035	
Total			66,279	587	1,657	64,035	32,925	64,035	96,960
Salmon River									
9	Little Salmon River	10-05-75	10,631	93	319	10,217	68,038	31,340	99,378
9	Little Salmon River	10-23-71	21,973	193	659	21,123			
Total			32,604	286	978	31,340	68,038	31,340	99,378

Total CWT Release: 158,051
Total Site Releases: 1,812,874
Total Smolt Releases: 1,812,874

Appendix 11. PIT tag summary for brood year 2005 steelhead at NSFH.

Raceway	Release Site	Number Tagged	Number Released	Mortality
4	Hells Canyon Dam Snake River	299	298	1
8	Hazard Creek Little Salmon River	299	298	1
9	Hazard Creek Little Salmon River	300	300	0
13	Pahsimeroi Weir Pahsimeroi River	298	298	0
Totals		1,196	1,194	2

Appendix 12. Comparison of mortality in vaccinated and non-vaccinated raceways at NSFH for brood year 2005.



Appendix 13. NSFH production history, BY66 to present.

NIAGARA SPRINGS HATCHERY HATCHERY HISTORY BY66-PRESENT													
YEAR	PAHSIM. Eggs/fry Received	OXBOW Eggs/fry Received	TOTAL Eggs/fry Received	TOTAL Yearly MORT.	% MORT	FALL Releases	Salmon R. SMOLT Release	Hells C. SMOLT Release	SPRING Releases	TOTAL LBS Released	Feed fed Total	Conv.	Fish/lb
1965-66	0	3,085,194	3,085,194	---	---	---	---	---	---	---	---	---	---
1966-67	0	2,605,288	2,605,288	623,533	23.93	29,400	1,364,842	587,513	1,952,355	153,552	305,890	1.99	12.7
1967-68	0	3,215,652	3,215,652	1,209,183	37.60	0	1,664,325	342,144	2,006,469	204,251	298,450	1.46	9.82
1968-69	0	2,469,536	2,469,536	695,219	28.15	0	1,665,117	109,200	1,774,317	184,186	280,430	1.52	9.63
1969-70	1,477,695	1,927,727	3,405,422	654,022	19.21	757,500	1,608,000	385,900	1,993,900	299,235	502,410	1.68	6.66
1970-71	1,330,494	1,480,150	2,810,644	305,176	10.86	670,960	1,630,002	0	2,444,860	202,025	384,040	1.90	12.1
1971-72	1,439,842	700,061	2,139,903	153,603	7.18	215,625	1,555,050	0	1,770,675	235,375	376,080	1.60	7.52
1972-73	8,850,764	1,819,721	10,670,485	3,105,637	29.10	3,008,664	1,543,349	0	4,556,184	163,839	266,800	1.63	27.8
1973-74	3,663,990	1,264,384	4,928,374	2,953,847	59.94	0	1,960,378	0	1,974,527	187,494	319,130	1.70	10.5
1974-75	3,160,144	280,098	3,440,242	2,108,426	61.29	0	1,331,280	0	1,331,816	166,640	352,890	2.12	7.99
1975-76	2,234,978	51,559	2,286,537	513,688	22.47	40,977	1,690,390	0	1,731,872	248,708	437,600	1.76	6.96
1976-77	2,487,824	730,862	3,218,686	1,642,383	51.03	0	1,433,675	141,005	1,576,303	251,835	454,762	1.81	6.26
1977-78	2,540,728	517,250	3,057,978	1,229,537	40.21	281,208	1,266,025	0	1,547,233	154,829	370,080	2.39	9.99
1978-79	2,048,350	441,069	2,489,419	426,977	17.15	344,944	1,372,454	0	1,717,498	244,887	643,680	2.63	7.01
1979-80	2,622,425	124,814	2,747,239	203,985	7.43	548,987	1,097,060	348,220	1,994,267	314,100	629,580	2.00	6.35
1980-81	1,697,010	498,416	2,195,426	720,172	32.80	0	862,494	612,760	1,475,254	316,330	622,930	1.97	4.66
1981-82	2,003,418	298,952	2,302,370	953,015	41.39	0	995,205	354,150	1,349,355	374,350	663,850	1.77	3.60
1982-83	2,313,339	253,776	2,567,115	1,431,975	55.78	500,000	542,390	92,750	635,140	181,150	448,860	2.48	3.51
1983-84	2,749,292	709,716	3,459,008	1,849,313	53.46	449,070	752,195	408,430	1,160,625	310,000	632,400	2.04	3.74
1984-85	2,333,760	598,404	2,932,164	613,771	20.93	630,500	1,273,181	414,712	1,687,893	314,650	541,198	1.72	5.36
1985-86	1,332,152	1,582,340	2,914,492	903,999	31.02	330,640	860,358	819,495	1,679,853	339,885	580,850	1.71	4.94
1986-87	1,339,176	935,195	2,274,371	422,476	18.58	39,995	1,011,900	800,000	1,811,900	419,000	557,960	1.33	4.32
1987-88	1,640,040	1,289,029	2,929,069	775,569	26.48	404,000	872,100	877,400	1,749,500	405,515	584,290	1.44	4.31
1988-89	1,256,289	1,213,399	2,469,688	803,488	32.53	0	930,700	735,500	1,666,200	406,800	574,770	1.41	4.10
1989-90	1,925,795	833,397	2,759,192	252,892	9.17	603,000	956,100	947,200	1,903,300	465,400	597,310	1.25	4.09
1990-91	1,966,434	113,190	2,079,624	311,624	14.98	0	856,000	912,000	1,768,000	484,025	632,030	1.28	3.65
1991-92	650,400	691,500	1,341,900	311,400	23.21	0	786,600	243,900	1,030,500	232,500	283,000	1.22	4.43
		812,000	812,000	394,936	48.64	0		417,064	417,064	72,786			5.73
1992-93	Wallowa												
1992-93	1,131,951	1,013,846	2,145,797				761,800	353,600		235,075			
1992-93	Babington		*Babington	Release	In Little	Salmon	*222,560	306,907	**47,089	131,090			
			**Brownlee	Reservoir									
1993-94	954,294	1,509,596	2,463,890	1,263,820	54.89	0	928,981	609,115	1,538,096	350,151	440,143	1.26	4.40
1994-95	1,042,728	1,099,915	2,142,643	281,034	13	160,000	741,180	960,429	1,701,609	376,060	489,960	1.29	4.52
1995-96	1,400,000	1,397,103	2,797,103	906,008	32.4	157,600	890,135	843,360	1,733,495	352,750	429,528	1.22	5.00
1996-97	1,297,250	1,303,599	2,600,849	698,156	26.84	149,040	1,093,002	660,651	1,753,653	370,520	421,144	1.14	4.79
1997-98	1,434,497	1,211,977	2,646,474	992,649	37.5	0	942,430	711,395	1,653,825	361,745	412,624	1.14	4.57
1998-99	1,412,000	1,393,383	2,805,383	759,809	27.08	60,634	1,185,535	657,665	1,843,200	444,455	484,110	1.09	4.63
1999-00	1,712,675	1,133,871	2,846,546	281,131	9.87	364,923	1,011,633	792,902	2,295,605	457,626	469,043	1.02	4.30
2000-01	1,416,442	1,045,825	2,462,267	100,330	4.07	431,133	1,351,337	579,467	1,930,804	459,580	473,540	1.03	4.29
2001-02	1,502,313	950,907	2,453,220	137,481	5.60	478,586	1,310,985	526,168	1,837,153	454,430	442,864	0.98	4.11
2002-03	1,161,547	919,416	2,080,963	224,277	10.78	0	1,330,802	525,884	1,856,686	417,275	415,155	0.99	4.45
2003-04	1,151,911	921,800	2,073,711	185,403	8.94	0	1,355,364	532,944	1,888,308	409,050	388,744	0.95	4.61
2004-05	1,133,186	924,927	2,058,113	353,035	17.15	0	1,179,054	526,024	1,705,078	369,600	368,040	0.99	4.61
2005-06	1,129,690	891,760	2,021,450	208,576	10.32	0	1,292,416	520,458	1,812,874	402,306	364,435	0.95	4.51

Appendix 14. Oxbow and Pahsimeroi stock spawn timing manipulations at NSFH for brood year 2004.

Hells Canyon Stock Eggs Used for Production Purposes				Pahsimeroi Stock Eggs Used for Production Purposes			
Lot Number	Spawn Date	Percent of Eggs Available Utilized for Smolt Production	Percent of Total Smolts Utilized for Production Back to Rack	Lot Number	Spawn Date	Percent of Eggs Available Utilized for Smolt Production	Percent of Total Smolts Utilized for Production Back to Rack
1	03/14/05	0%	0.0%	1	03/31/05	74%	13%
2	03/17/05	73%	1.2%	2	04/13/05	32%	7.8%
3	03/21/05	61%	3.5%	3	04/18/05	87%	6.3%
4	03/28/05	62%	5.4%	4	04/25/05	8%	1.0%
5	03/31/05	87%	14.9%	5	04/28/05	97%	9.6%
6	04/04/05	57%	12.8%	6	05/02/05	71%	20.7%
7	04/07/05	57%	3.4%	7	05/05/05	96%	26.4%
8	04/11/05	52%	2.0%	8	05/09/05	91%	10.8%
9	04/14/05	86%	4.6%	9	05/12/05	100%	4.4%
10	04/18/05	76%	7.7%				
11	04/21/05	66%	6.3%				
12	04/25/05	82%	21.2%				
13	04/28/05	94%	11.4%				
14	05/02/05	94%	4.6%				
15	05/05/05	100%	1.0%				

Submitted by:

Approved by:

Jarrett L. Page
Fish Hatchery Assistant Manager

Steve Yundt, Chief
Fisheries Bureau

Jerry Chapman
Fish Hatchery Manager II

Tom Rogers
Anadromous Fish Hatcheries Supervisor

Doug S. Young
Fish Culturist

Kevin Yelton
Fish Culturist