

Homeostasis or channelopathy? Acquired cell type-specific ion channel changes in temporal lobe epilepsy and their antiepileptic potential

OPEN ACCESS

Jakob Wolfart* and Debora Laker

Oscar Langendorff Institute of Physiology, University of Rostock, Rostock, Germany

Edited by:

Maria Cristina D'Adamo,
Charité - Universitätsmedizin Berlin,
Germany

Reviewed by:

Imre Vida,
Charité - Universitätsmedizin Berlin,
Germany

Giuseppe Di Giovanni,
University of Malta, Malta

*Correspondence:

Jakob Wolfart,
Oscar Langendorff Institute of
Physiology, University of Rostock,
Gertrudenstr. 9,
18057 Rostock, Germany
jakob.wolfart@uni-rostock.de

Specialty section:

This article was submitted to
Membrane Physiology and Membrane
Biophysics,
a section of the journal
Frontiers in Physiology

Received: 04 March 2015

Accepted: 19 May 2015

Published: 15 June 2015

Citation:

Wolfart J and Laker D (2015)
Homeostasis or channelopathy?
Acquired cell type-specific ion channel
changes in temporal lobe epilepsy and
their antiepileptic potential.
Front. Physiol. 6:168.
doi: 10.3389/fphys.2015.00168

Neurons continuously adapt the expression and functionality of their ion channels. For example, exposed to chronic excitotoxicity, neurons homeostatically downscale their intrinsic excitability. In contrast, the “acquired channelopathy” hypothesis suggests that proepileptic channel characteristics develop during epilepsy. We review cell type-specific channel alterations under different epileptic conditions and discuss the potential of channels that undergo homeostatic adaptations, as targets for antiepileptic drugs (AEDs). Most of the relevant studies have been performed on temporal lobe epilepsy (TLE), a widespread AED-refractory, focal epilepsy. The TLE patients, who undergo epilepsy surgery, frequently display hippocampal sclerosis (HS), which is associated with degeneration of *cornu ammonis* subfield 1 pyramidal cells (CA1 PCs). Although the resected human tissue offers insights, controlled data largely stem from animal models simulating different aspects of TLE and other epilepsies. Most of the cell type-specific information is available for CA1 PCs and dentate gyrus granule cells (DG GCs). Between these two cell types, a dichotomy can be observed: while DG GCs acquire properties decreasing the intrinsic excitability (in TLE models and patients with HS), CA1 PCs develop channel characteristics increasing intrinsic excitability (in TLE models without HS only). However, thorough examination of data on these and other cell types reveals the coexistence of protective and permissive intrinsic plasticity within neurons. These mechanisms appear differentially regulated, depending on the cell type and seizure condition. Interestingly, the same channel molecules that are upregulated in DG GCs during HS-related TLE, appear as promising targets for future AEDs and gene therapies. Hence, GCs provide an example of homeostatic ion channel adaptation which can serve as a primer when designing novel anti-epileptic strategies.

Keywords: hippocampus, homeostasis, kainic acid, pilocarpine, channelacoids

Introduction

The relationship between epileptic seizures and ion channels is typically focused on the proepileptic (meaning seizure-supporting) nature of ion channel abnormalities. This perspective, embodied in the terms “channelopathy” and “channelepsy” (Hoffman, 1995; Ptacek, 1997; George, 2004; Kullmann and Waxman, 2010; D’adamo et al., 2013), is fueled by the increasing number of ion channel mutations discovered in epilepsy patients (Biervert et al., 1998; Charlier et al., 1998; Singh et al., 1998, 2008; Zuberi et al., 1999; Heilstedt et al., 2001; Chioza et al., 2002; Schulte et al., 2006; Cavalleri et al., 2007; Tomlinson et al., 2010; Lachance-Touchette et al., 2011; Weckhuysen et al., 2013), and the seizure phenotypes of corresponding engineered channel mutants (Signorini et al., 1997; Schroeder et al., 1998; Smart et al., 1998; Spigelman et al., 2002; Ludwig et al., 2003; Peters et al., 2005; Huang et al., 2009; Ishii et al., 2009; Riazanski et al., 2011; Hedrich et al., 2014). In contrast to genetic channelopathies, an “acquired channelopathy” is declared when ion channel abnormalities develop independently of the genetic background (Waxman, 2001; Bernard et al., 2004; Poolos and Johnston, 2012). Thus, the prevailing view of progressive acquisition of proepileptic channel properties during epilepsy is conceptually similar to the old “seizures beget seizures” hypothesis (Gower, 1881; Hauser and Lee, 2002; Sills, 2007; Ben-Ari, 2008).

For those studying homeostasis, the acquired channelopathy scenario may appear puzzling; how is it possible that neurons exposed to an environment already hyperexcitable, further enhance their excitability in a potentially self-destructive manner? Usually, biological cells are well equipped to counteract detrimental excitation and especially neurons respond to extrinsic hyperexcitation with intrinsic hypoexcitability on all time scales (Davis and Bezprozvanny, 2001; Turrigiano and Nelson, 2004; Marder and Goaillard, 2006; Meier et al., 2015). Are these mechanisms of protective homeostasis disabled in epilepsy? And if so, is it the rule or rather the exception? It is important to answer these questions, because it will help to understand the mechanisms of seizures and antiepileptic drugs (AEDs). The prevailing AED strategy is to inhibit excitatory channels such as sodium (Na) and calcium (Ca) channels (Löscher et al., 2013). A more recent approach is to support a specific potassium (K) channel (further discussion below) (Tatulian et al., 2001; Faulkner and Burke, 2013). Generally, AED strategies imply relatively fixed ion channel repertoires, but channels are dynamically adjusted all the time and these changes are not well understood in particular with respect to epilepsy.

Most of the discussion on channel-related AED mechanisms of action takes place without mentioning cell types even though AED channel targets can have behaviorally opposed effects depending on which cell types are affected (Prakriya and Mennerick, 2000; He et al., 2002). Hence, to discuss antiepileptic strategies in a meaningful manner, the first step is to obtain an overview on available information about channel molecule alterations, cell types, and epilepsy model methods. The present review focuses on temporal lobe epilepsy (TLE), because the majority of the available data concern this most common form

of partial epilepsies, which in turn account for 60% of all adult epilepsy cases (Tellez-Zenteno and Hernandez-Ronquillo, 2012). Some of the discussed channel mechanisms might also be relevant for other epilepsies as many TLE animal models display generalized seizures. Studies on the principal neurons of the hippocampal formation, especially *cornu ammonis* 1 (CA1) and dentate gyrus (DG), outnumber studies on other areas by far. This fact is mirrored in our list of acquired ion channel alterations (Table 1). Other cell types are expected to gain importance in the future and are also discussed in respective sections. We especially highlight channels with net inhibitory effects and relate these to existing or promising AED mechanisms of action (Table 2), asking whether certain cell types can signpost effective molecule target combinations for future AED and antiepileptic gene therapies.

Temporal Lobe Epilepsy

TLE seizures start in the temporal lobe and impair the consciousness, among other symptoms (Blumenfeld and Meador, 2014). TLE is often refractory to AEDs; therefore, anterior temporal lobe resection is a standard treatment (De Tisi et al., 2011). In its most common form, TLE affects the medial temporal/limbic network with involvement of the entorhinal cortex, the amygdala, and the hippocampus with the DG and the CA regions 1-3 (hippocampus *sensu strictu*) (Spencer, 2002). The neuropathological correlate of TLE is hippocampal sclerosis (HS) (Margerison and Corsellis, 1966; Curia et al., 2014) and quite often is HS interpreted to be the cause of TLE (Blümcke et al., 2012). However, as human TLE is heterogeneous, it appears difficult to find even simple correlations between the severity of HS and epileptic seizures, let alone the certainty that HS causes TLE or indeed vice versa (King et al., 1997; Jefferys, 1999; Blümcke et al., 2002; Mathern et al., 2002; Sutula et al., 2003; De Lanerolle and Lee, 2005; Briellmann et al., 2007; Mueller et al., 2007). A widespread criterion for HS diagnosis is loss of more than half of the CA1 PCs (Wyler et al., 1992). Other structural changes associated to HS, are sprouting of the mossy fiber axons of DG granule cells (GCs) (Sutula et al., 1989), changed GC morphology (Isokawa and Levesque, 1991), and GC dispersion (GCD) (Houser, 1990). Severe HS is correlated with severe GCD (Thom et al., 2002), although this relationship is not always strict (Blümcke et al., 2013). Occasionally, studies counting cells per area confuse GCD with degeneration of GCs. However, GCs are only lost in extreme HS (Wyler grade IV), i.e., when most of the CA1 PCs have already degenerated (Wyler et al., 1992). More than 90% of the hippocampi resected during epilepsy surgery exhibit HS (Blümcke et al., 2012) but it should be kept in mind that the decision on surgery itself depends on HS because the success rates of epilepsy surgery are higher with diagnosis of a lesion (Jobst and Cascino, 2015). Unbiased postmortem studies reveal HS in about half of TLE patients (Thom et al., 2010).

Epilepsy surgery offers direct experimental access to living human hippocampal tissue but lack of proper control tissue makes animal models indispensable. Different approaches were used to simulate a chronic epileptic state with anatomical changes resembling TLE. Since human TLE is heterogeneous and has

unknown causes, animal models can only reproduce partial aspects of the disease (Coulter et al., 2002; Morimoto et al., 2004). One hypothesis for TLE is an initial precipitating injury (Mathern et al., 2002). Therefore, most animal models generate a chronic epileptic state via one status epilepticus (SE). Widespread is SE induction via i.p. (systemic) injection of kainate (sKA) (Nadler et al., 1978; Ben-Ari, 1985) or pilocarpine (sPilo) (Turski et al., 1983, 1989). The sKA and sPilo models produce generalized seizures and bilateral brain damage, the extent of which depends on the SE-termination protocol (Schwob et al., 1980; Turski et al., 1983). Often, no HS or only mild forms of it (noHS) occur in sKA and sPilo models (Okazaki et al., 1999; Scharfman et al., 2000; Dietrich et al., 2005; Curia et al., 2014). In contrast, development of HS and GCD, as well as focal spontaneous TLE seizures, can be induced by intracranial (intrahippocampal) kainate injection (iKA) (Suzuki et al., 1995; Fritschy, 2004). Another TLE model, which reproduces HS and chronic (bilateral) seizures, is perforant path stimulation-induced non-convulsive- (Kienzler et al., 2009) or convulsive SE (Bumanglag and Sloviter, 2008). In addition, there are different forms of electrical stimulation (“kindling”) of amygdala or hippocampus (aKindl, hKindl) which mostly evoke seizures only during the stimulation (i.e., no chronic epilepsy), (Goddard et al., 1969; Mcnamara, 1984; Morimoto et al., 2004).

Dentate Gyrus

The DG is often viewed as a strategic “gate keeper” of the hippocampus and failure in its filter function was hypothesized to be a potential cause for TLE seizures (Heinemann et al., 1992; Lothman et al., 1992; Hsu, 2007; Krook-Magnuson et al., 2015). An alternative hypothesis is that the DG does not actively contribute to hippocampal seizures (Sloviter, 1994; Liu et al., 2000; Harvey and Sloviter, 2005). The following changes were considered responsible for a proepileptic role of the DG (De Lanerolle et al., 1992; Mody et al., 1992b): (i) mossy fiber sprouting (Tauck and Nadler, 1985; Sutula et al., 1989), (ii) loss of specific interneurons (Sloviter, 1987; Magloczky and Freund, 2005), and (iii) intrinsic hyperexcitability of the principal DG neurons, the GCs (Magloczky et al., 1997; Beck et al., 1998; Dietrich et al., 1999; Coulter, 2000; De Lanerolle et al., 2003; Selke et al., 2006; Mehranfarid et al., 2014a). In contrast to the last point, many studies concluded that TLE does not change DG GCs intrinsically (Mody et al., 1992a; Beck et al., 1996; Isokawa, 1996; Molnar and Nadler, 1999; Okazaki et al., 1999; Scharfman et al., 2003; Dietrich et al., 2005; Beck and Yaari, 2008). Contrary to both prior hypotheses, we found a *decrease* of the intrinsic excitability of GCs which was due to a reduction in input resistance (R_{in}); it occurred in samples of TLE patients with HS vs. mild/no HS as well as in iKA vs. control mice (Stegen et al., 2009, 2012; Young et al., 2009; Kirchheim et al., 2013). A reasonable question is: why are there so many disparate results on the same cell type (Vida, 2009)? With rare exceptions (Isokawa and Mello, 1991; Mehranfarid et al., 2014b), most studies reporting unchanged GCs were those employing TLE models without HS. In TLE patients and the iKA TLE model, the R_{in} of DG GCs correlates with the degree of HS (Stegen et al., 2009, 2012; Young et al., 2009). Although it cannot be ruled out that

some of the studies missed R_{in} differences due to methodological procedures, such as applying minimum R_{in} as cell selection criterion, low seal resistance sharp electrodes, or by dissociating GC somata from their dendritic conductances (Mehranfarid et al., 2014b), the conservative conclusion currently is: in TLE with HS, the ion channel expression of GCs is more drastically changed than in TLE without HS. It is important to note that GC channel adaptations only occur in the HS area, i.e., where GCD and neurodegeneration are clearly visible, but not outside of this HS focus. Ventral parts of the ipsilateral DG of iKA mice as well as the contralateral DG may even harbor hyperexcitable GCs (Le Duigou et al., 2008; Young et al., 2009; Häussler et al., 2012).

The molecular mechanism behind the reduced excitability of GCs is mainly transcriptional upregulation of K leak channels, i.e., channels that are open at resting membrane potential (V_{rest}) (Stegen et al., 2009, 2012; Young et al., 2009). Specifically, these are inwardly rectifying K (K_{ir}) channels of classic leak subtype $K_{ir}2.1-4$ and two pore domain K leak channels of subtype $K_{2p}1.1$ and $K_{2p}6.1$. In addition, elevated tonic chloride (Cl) leak conductances mediated by gamma amino butyric receptors type A ($GABA_A$ Rs) were detected in GCs of iKA mice (Young et al., 2009). Such tonic $GABA_A$ currents are likely mediated by extrasynaptic $GABA_A$ Rs composed of $\alpha 4-6$ plus β , and $\gamma 2$ or δ subunits (Peng et al., 2002; Zhang et al., 2007; Glykys et al., 2008) which could underlie epilepsy-related changes (Peng et al., 2004). In the adult stage, the Cl equilibrium potential (E_{Cl}) of GCs is between V_{rest} and action potential (AP) threshold, although E_{Cl} can change during TLE (Palma et al., 2006; Huberfeld et al., 2007; Pathak et al., 2007; Khirug et al., 2010; Barmashenko et al., 2011). Therefore, the functional influence of TLE-related $GABA_A$ leak elevation is an enhancement of shunting inhibition. To explain the latter: if a large conductance ($1/R$) is added, its influence via Ohm’s law ($U = R \times I$) will minimize the voltage impact (U) of any further input currents (I). All conductances have this influence, but the counterintuitive effect of “inhibitory depolarization” occurs specifically when E is between V_{rest} and AP threshold, as with E_{Cl} (Staley and Mody, 1992; Wolfart et al., 2005; Meier et al., 2015). There are few but notable differences between the iKA mouse model and the human condition. Human GCs lack the $GABA_A$ leak increase and instead show an HS-related enhancement of ZD7288-sensitive, hyperpolarization-activated cation conductance, most likely mediated by HCN1 channels (Stegen et al., 2012). Thus, contrary to prior reports (Stabel et al., 1992), a functional h-current (I_H) exists in rodent and human GCs (Young et al., 2009; Stegen et al., 2012) and this I_H , as well as respective HCN1 subunits are enhanced in HS-related TLE (Bender et al., 2003; Stegen et al., 2012). The functional effect of I_H in GCs is similar to the $GABA_A$ leak because E_H is also between V_{rest} and AP threshold, again contributing to enhanced shunting inhibition (Stegen et al., 2012) (see CA Section for more discussion on HCN channels). Another interesting difference between the GCs of TLE patients and iKA mice is that human GCs almost never display a pronounced delay of AP responses as their iKA counterparts (Stegen et al., 2009, 2012; Young et al., 2009). These “ramp” delays of GCs are mediated by *shaker*-related, voltage-gated K (K_v) channels containing $K_v1.1$, $K_v1.2$, or $K_v1.6$ subunits,

which are sensitive to dendrotoxin and μM concentrations of 4-aminopyridine (4-AP) (Kirchheim et al., 2013). The molecular mechanism for the 3-fold delayed AP responses of iKA GCs is transcriptional upregulation of $\text{K}_v1.1$ subunits, which, consistent with a homeostatic response, is reversible upon interruption of chronic hyperexcitation (Kirchheim et al., 2013). Without these K_v1 currents, GCs are much more vulnerable during excitotoxic insults (Kirchheim et al., 2013). The dissimilarities between human and mouse GCs could be due to species differences or (more likely) due to the disease etiology. However, for the present perspective on AED strategies it is notable that in both human and mouse TLE GCs, a depolarizing but shunting conductance is co-upregulated with K_{ir} channels such that V_{rest} is almost unchanged (Stegen et al., 2009, 2012; Young et al., 2009). This downscaling is not only suitable to maintain basic metabolic functions dependent on V_{rest} ; it is also a native example of static shunt, enforcing a subtractive gain shift of the neuronal input-output curve (Wolfart et al., 2005). A recent network simulation study has demonstrated that the experimentally observed channel scaling of GCs could also restore spatiotemporal pattern separation under epileptic conditions, i.e., maintain the proposed function of the DG network (Yim et al., 2015).

In addition to the discussed leak channel modifications, other epilepsy-related changes occur in excitatory ion channels and ionotropic receptors of DG cell types (Table 1). For example, voltage-gated Ca (Ca_v) channels of P/Q-type ($\text{Ca}_v2.2$) were found increased while the L-type subunit $\text{Ca}_v1.2$ was diminished in the DG molecular layer of TLE patient vs. autopsy samples (Djamshidian et al., 2002). No such changes were observed in the TLE model of ventricular KA injection (Westenbroek et al., 1998). Many immunohistochemistry studies exist on GABA_A R and GABA_B R changes during TLE: the GABA_A R $\alpha 1$ -3, $\beta 2$ -3, and $\gamma 2$ subunits were all found elevated in GCs of TLE patients (Loup et al., 2000) and similar results (except $\alpha 2$) were obtained in the iKA model (Bouilleret et al., 2000; Knuesel et al., 2001). In sPilo, the DG immunostaining of GABA_A R subunits is heterogeneous (Brooks-Kayal et al., 1998). Some of the confusion, created by various tissue-level studies was clarified by electron microscopy and functional analysis. For example, Sun et al. (2007) revealed that although $\text{GABA}_A\alpha 4$ subunits are reduced extrasynaptically in GCs of the hKindl model, they are in fact increased in synaptic locations which was interpreted as proepileptic. Similarly, $\text{GABA}_A\gamma 2$ subunits disappear from synaptic locations, reducing the phasic inhibition in sPilo but they reappear in extrasynaptic locations, apparently replacing lost $\text{GABA}_A\delta$ subunits because functionally, tonic inhibition is maintained (Zhang et al., 2007). Measuring surface-coupled protein during sPilo yielded reduction of $\text{GABA}_A\beta 2/3\gamma 2$ subunits on GC membranes; however, GC recordings revealed that tonic GABA_A currents were, if anything, increased (Goodkin et al., 2008). In contrast to GABA_A Rs, the GABA_B R1a-b immunosignal of GCs was found reduced in HS-TLE patients (Munoz et al., 2002).

Interneurons of the DG degenerate in the hilus and the molecular layer but somatic inhibition of GCs is apparently spared (Magloczky and Freund, 2005). Here we refrain from

reviewing TLE-related interneuron numbers, as respective cell counting studies mostly rely on Ca binding proteins which themselves change in TLE as shown by Magloczky et al. (1997); for review see Magloczky and Freund (2005). With respect to ion channels of interneurons in the hilus, a decrease of voltage-gated Na type 1 (Na_v1) channels (Qiao et al., 2013) and an increase of Ca_v1 channels (Xu et al., 2007) had been reported. Interestingly, in basket cell interneurons of the DG, tonic GABA_A currents are homeostatically adjusted in sPilo rats (Yu et al., 2013).

Astrocytes may also play an important role in epilepsy, in particular via their (potentially impaired) capacity to buffer extracellular K ions (Bordey and Sontheimer, 1998; Jabs et al., 2008; Boison, 2012). For example, in the DG, less glial K_{ir} current was measured in HS- vs. noHS TLE tissue (Hinterkeuser et al., 2000) and less $\text{K}_{2p3.1}$ -positive glia was noted in the molecular layer (Kim et al., 2008b). Concerning the K_{ir} channels, it is likely that downregulation of the $\text{K}_{ir4.1}$ subunit is responsible (Buono et al., 2004; Heuser et al., 2012). However, in another sPilo study, no changes of hippocampal $\text{K}_{ir4.1}$ protein were detected or even upregulation was noted in cortical and subcortical regions (Nagao et al., 2013). In a different model (seizure-sensitive gerbils), upregulation of $\text{K}_{2p3.1}$ channels was reported (Kim et al., 2007a). In two human cases, gain of function mutation in the glial $\text{K}_{ir4.1}$ channel was associated with infantile epileptic spasms (Sicca et al., 2011).

In summary, pro- and anticonvulsive channel changes have been described in the DG of different TLE models. For noHS models, it is difficult to draw a final conclusion on channel changes in GCs as these are heterogeneous. In TLE tissue with HS, the ion channel expression of GCs is clearly changed to decreased excitability.

Cornu Ammonis

A prominent example of acquired channelopathy is the reduced influence of A-type (rapidly inactivating) K channels in dendrites of CA1 PCs of sPilo rats (Bernard et al., 2004). The respective identification as A-type current was performed in current-clamp experiments via 5 mM 4-AP (Bernard et al., 2004), which blocks delayed rectifier (not rapidly inactivating) K_v1 channels as well as A-type K_v4 and K_v1 channels (Pongs, 1992). In addition, a reduction of $\text{K}_v4.2$ and $\text{K}_v1.4$ mRNA was detected via RT-PCR from microdissected CA1 tissue with mixed cell types (Bernard et al., 2004). The conclusion that $\text{K}_v4.2$ channel downregulation mediated the dendritic channelopathy in sPilo rats (Bernard et al., 2004) was confirmed later by immunocytochemistry, although quantification on the tissue level yielded no difference in the chronic phase (Monaghan et al., 2008). In another sPilo study, $\text{K}_v4.2$ protein of the CA1 region was increased during the first week after SE but decreased in the chronic phase (Su et al., 2008). In the sKA model, $\text{K}_v4.2$ mRNA was reported unchanged in CA1 PCs in the first 24 h but appeared upregulated 12 h later (Francis et al., 1997). In the same model, A-currents of CA1 PCs were decreased but their sensitivity to 50 μM 4-AP and the immunohistochemistry suggested that these currents were rather mediated by K_v1 than K_v4 channels (Sosanya et al., 2014).

TABLE 1 | Ion channel changes during temporal lobe epilepsy.

Type of cells and TLE/model	Ion channel/current	Regulation/methods	References
DG GRANULE CELLS			
HS/noHS human	I _{Kir2.x}	↑ funct (vc, 0.1 mM Ba ²⁺)	Stegen et al., 2009
HS/no HS human	K _{ir} 2.2	↑ prot IP	Stegen et al., 2012
HS/noHS human	GABA _A α1/α2 GABA _A α3 GABA _A β2/β3 GABA _A γ2	↑ prot IP ± prot IP ↑ prot IP ↑ prot IP	Loup et al., 2000
HS/noHS human	GABA _B R1a-b	↓ prot IP	Munoz et al., 2002
HS/noHS human	HCN1	↑ RNA _{ISH} , prot IP	Bender et al., 2003
HS/noHS human	I _H HCN1	↑ funct (vc, ZD7288) ↑ prot IP	Stegen et al., 2012
TLE/autopsy human	K _v 7.5	± prot IP	Yus-Najera et al., 2003
TLE/autopsy human	Na _v 1.1/1.2/1.3	± RNA _{ISH}	Whitaker et al., 2001
TLE/autopsy human	Ca _v 1.2 Ca _v 1.3 Ca _v 2.2/2.3 Ca _v 2.1	↓ prot IP ± prot IP ± prot IP ↑ prot IP	Djamshidian et al., 2002
HS-iKA mouse	I _{Kir2.x} K _{ir} 2.1/2.2/2.3/2.4	↑ funct (vc, 40 μM Ba ²⁺) ↑ prot IP	Young et al., 2009
HS-iKA mouse	K _{2P} 1.1 K _{2P} 6.1	↑ prot IP ↑ prot IP	Young et al., 2009
HS-iKA mouse	I _{Kv1.x} K _v 1.1 K _{ir} 2.1	↑ funct (vc, DTX) ↑ RNA _{SC-PCR} , prot IF ↑ RNA _{SC-PCR}	Kirchheim et al., 2013
HS-iKA mouse	GABA _A α1/α5 GABA _A α2 GABA _A α3 GABA _A γ2	↑ prot IP ↓ prot IP ± prot IP ↑ prot IP	Bouilleret et al., 2000
HS-iKA mouse	GABA _A α2	↑ prot IF	Knuesel et al., 2001
HS-iKA mouse	I _{tGABA-A}	↑ funct	Young et al., 2009
hKindl rat	GABA _A α4	↓ _{syn} ↑ _{ex} prot EM	Sun et al., 2007
hKindl rat	I _{Na}	± funct	Ketelaars et al., 2001
aKindl rat	Na _v 1.1/1.2/1.6 Na _v 1.6	± prot IF ± RNA _{ISH}	Blumenfeld et al., 2009
aKindl mouse	Na _v 1.6	± prot IF	Blumenfeld et al., 2009
sPilo rat	K _v 1.4 K _v 4.2/KChIP2	± prot IP ± prot IP	Monaghan et al., 2008
sPilo rat	I _{GABAeff} I _{GABApot}	↑ funct ± funct	Gibbs et al., 1997
sPilo rat	I _{GABAeff} I _{GABApot} GABA _A α1 GABA _A α2/α3 GABA _A α4 GABA _A β1 GABA _A β2 GABA _A β3 /δ /ε GABA _γ 1/γ2/γ3	↑ funct ± funct ↓ RNA _{SC-PCR} ± RNA _{SC-PCR} ↑ RNA _{SC-PCR} ↓ RNA _{SC-PCR} ± RNA _{SC-PCR} ↑ RNA _{SC-PCR} ± RNA _{SC-PCR}	Brooks-Kayal et al., 1998
sPilo mouse	GABA _A α4 GABA _A δ GABA _A γ2	↓ _{1-4days} ↑ _{30days} prot IP ↓ prot IP ± _{1-4days} ↑ _{60days} prot IP	Peng et al., 2004
sPilo rat	GABA _A β2 GABA _A β3 GABA _A δ GABA _A γ2	↓ surface prot W ↓ surface prot W ± prot _{surface} (↑), funct ↓ surface prot W	Goodkin et al., 2008
sPilo rat	HCN1	↑ RNA _{ISH}	Bender et al., 2003

(Continued)

TABLE 1 | Continued

Type of cells and TLE/model	Ion channel/current	Regulation/methods	References
sPilo rat	TRP3 TRP6	↓ 1days–1week prot IP ↑ 1days–1week prot IP	Kim et al., 2013
sPilo rat, mouse	I _{Cav} Ca _v 3.2	± funct ± prot IF	Becker et al., 2008
sPilo mouse	I _t GABA–A I _p GABA–A GABA _A α GABA _A δ GABA _A γ2	± funct ↓ funct ↓ _{syn/ex} prot EM ↓ _{syn/ex} port EM ↓ _{syn} ↑ _{ex} prot EM	Zhang et al., 2007
P20 sPilo rat	GABA _A α1	↑ RNA _{SC} –PCR	Raol et al., 2006b
P20 sKA rat	GABA _A α1	↑ RNA _{SC} –PCR	Raol et al., 2006b
sKA rat	K _v 4.2	↓ 3h,6h ↑ 24h RNA _{ISH}	Francis et al., 1997
sKA rat	Na _v 1.1 Na _v 1.2 Na _v 1.3	± 3–24h RNA _{ISH} ↑ 3–6h ± 24h RNA _{ISH} ↑ 3–6h ↑ 24h RNA _{ISH}	Bartolomei et al., 1997
DG INTERNEURONS			
sPilo rat	KChIP1 (hilus)	↓ prot	Monaghan et al., 2008
sPilo rat	tGABA _A (BCs) GABA _A δ(BCs)	↑ 6–8days funct ↑ 1week prot IF	Yu et al., 2013
sPilo mouse	GABA _A δ(BCs)	↑ prot IP	Peng et al., 2004
DG GLIAL CELLS			
HS/noHS human	I _{Kir}	↓ funct	Hinterkeuser et al., 2000
HS/noHS human	K _{ir} 4.1 (hilus)	↓ prot	Heuser et al., 2012
sPilo rat	K _{2p} 3.1	↓ prot	Kim et al., 2008b
CA1 PYRAMIDAL CELLS			
TLE/autopsy human	K _v 7.5	± prot IP	Yus-Najera et al., 2003
TLE/autopsy human	Na _v 1.1/1.3 Na _v 1.2	± RNA _{ISH} ↓ RNA _{ISH}	Whitaker et al., 2001
TLE/autopsy human	Ca _v 1.2/1.3 Ca _v 2.1/2.2/2.3	± prot IP ± prot IP	Djamshidian et al., 2002
HS-iKA mouse	GABA _A α2	↓ prot IF	Knuesel et al., 2001
HS-iKA mouse	GABA _A α1/α3/α5 GABA _A α2 GABA _A γ2	↓ prot IP ± prot IP ↓ prot IP	Bouillere et al., 2000
hKindl rat	I _{Na} –window	↑ funct	Ketelaars et al., 2001
aKindl rat	Na _v 1.1/1.2/1.6 Na _v 1.6	± prot IF ± RNA _{ISH}	Blumenfeld et al., 2009
aKindl mouse	Na _v 1.6	± prot IF	Blumenfeld et al., 2009
sPilo rat	I _A –type/dend	↓ funct (cc, 5 mM 4-AP)	Bernard et al., 2004
sPilo rat	K _v 4.2/KChIP2	↓ prot IP	Monaghan et al., 2008
sPilo rat	I _{SK}	↓ funct (vc, UCL1684)	Schulz et al., 2012
sPilo rat	I _{GABAeff} I _{GABApot}	↓ funct ↑ funct	Gibbs et al., 1997
sPilo rat	I _{Cl} CLC2	↑ funct ↑ prot IF, W	Ge et al., 2011
sPilo rat	I _H /dend I _H /soma	↓ 1h–5weeks funct ± 3days–5weeks funct	Jung et al., 2007, 2011
sPilo rat	I _H HCN2	↓ funct (vc, ZD7288) ↓ prot (surface)	Marcelin et al., 2009
sPilo rat	I _{Cav}	↑ funct (cc, 1 mM Ni ²⁺)	Sanabria et al., 2001
sPilo rat	TRP3 TRP6	↓ 1days–1week prot IP ↑ 1days–1week prot IP	Kim et al., 2013
sPilo rat	I _{Cav} 3	↑ funct (vc, 0.1 mM Ni ²⁺)	Su et al., 2002

(Continued)

TABLE 1 | Continued

Type of cells and TLE/model	Ion channel/current	Regulation/methods	References
sKA rat	K _v 4.2	± 3h,6h,24h RNA _{ISH}	Francis et al., 1997
sKA rat	I _{Kv1.x} K _v 1.1	↑ 14days ↓ 30days funct (cc, 50 μM 4-AP) ↑ 14days ↓ 30days prot IF/W	Sosanya et al., 2014
sKA rat	HCN1 HCN2	↓ RNA _{ISH} ↑ RNA _{ISH}	Brewster et al., 2002
sKA rat	I _H	↑ 1–2days ↓ 28–30days funct	Shin et al., 2008
sKA rat	Na _v 1.1 Na _v 1.2 Na _v 1.3	± 3–24h RNA _{ISH} ± 3–24h RNA _{ISH} ± 3h ↑ 6h ± 24h RNA _{ISH}	Bartolomei et al., 1997
CA1 GLIAL CELLS			
TLE/autopsy human	K _{ir} 4.1	↓ prot IP	Heuser et al., 2012
sPilo rat	K _{2p} 3.1	± prot	Kim et al., 2008b
sKA rat	K _v 1.3	↑ funct	Menteyne et al., 2009
CA2 PYRAMIDAL CELLS			
HS/noHS human	GABA _A α2 GABA _A α1/α3	↑ prot IP ↓ prot IP	Loup et al., 2000
TLE/autopsy human	Na _v 1.1/1.3 Na _v 1.2	± RNA _{ISH} ↓ RNA _{ISH}	Whitaker et al., 2001
TLE/autopsy human	Ca _v 1.2/1.3 Ca _v 2.1/2.2/2.3	± prot IP ± prot IP	Djamshidian et al., 2002
noHS-aKindl rat	Na _v 1.1/1.2/1.6 Na _v 1.6	± prot IF ± RNA _{ISH}	Blumenfeld et al., 2009
CA3 PYRAMIDAL CELLS			
HS/noHS human	Na _v 1.1/1.3 Na _v 1.2	± RNA _{ISH} ↓ RNA _{ISH}	Whitaker et al., 2001
TLE/autopsy human	Ca _v 1.3, Ca _v 2.2/2.3 Ca _v 2.1 Ca _v 1.2	± prot IP ± prot IP ↑ prot IP ↓ prot IP	Djamshidian et al., 2002
HS-iKA mouse	GABA _A α1/α2 GABA _A α5 GABA _A γ2	± prot IP ↓ prot IP ↓ prot IP	Bouilleret et al., 2000
aKindl rat	Na _v 1.1/1.2 Na _v 1.6	± prot IF ↑ prot IF, RNA _{ISH}	Blumenfeld et al., 2009
aKindl mouse	I _{persist inward} Na _v 1.6	↑ funct ↑ prot IF, RNA _{ISH}	Blumenfeld et al., 2009
sKA rat	K _v 4.2	± 3h,6h ↓ 24h RNA _{ISH}	Francis et al., 1997
sKA rat	I _{Kv1.x} K _v 1.1	↑ 14days ↓ 30days funct ↑ 14days ↓ 30days prot IF/W	Sosanya et al., 2014
sKA rat	HCN1 HCN2	↓ RNA _{ISH} ↑ RNA _{ISH}	Brewster et al., 2002
sKA rat	Na _v 1.1 Na _v 1.2 Na _v 1.3	± 3–24h RNA _{ISH} ± 3–24h RNA _{ISH} ± 3h ↑ 6h ± 24h RNA _{ISH}	Bartolomei et al., 1997
sPilo rat	TRP3 TRP6	↓ 1days–1week prot IP ↑ 1days–1week prot IP	Kim et al., 2013
CA INTERNEURONS			
ventral of iKA	I _H (CA3 O-LM)	↓ funct	Dugladze et al., 2007
sPilo rat	KChIP1 (CA3 SP)	↓ prot IP	Monaghan et al., 2008
SUBICULAR NEURONS			
sPilo rat	I _{Ca} (PCs)	↑ funct (cc, 1 mM Ni ²⁺)	Wellmer et al., 2002
AMYGDALA NEURONS			
a/hKindl rat	K _v 7.2	↑ prot IP	Penschuck et al., 2005

(Continued)

TABLE 1 | Continued

Type of cells and TLE/model	Ion channel/current	Regulation/methods	References
CORTICAL NEURONS			
hKindl rat EC	I _{Nav} (layer II stellate cells) Na _v 1.2/6 (layer II stellate cells) Na _v 1.1/3 (layer II stellate cells)	↑ funct ↑ prot IF ± prot IF	Hargus et al., 2013
hKindl rat PC	K _v 1.6 (interneurons)	↓ funct, prot IF	Gavrilovici et al., 2012
sKA rat EC	I _H (layer III PCs)	↓ 24h ↓ 1week funct (dendritic)	Shah et al., 2004
THALAMIC NEURONS			
sPilo mouse	I _A (VM relay cells) K _v 4.2 (surf)	↑ funct ↑ prot IF	Smith et al., 2012
sPilo mouse	I _{T-type} (relay cells) Ca _v 3.1 (thalamus) Ca _v 3.2 (thalamus) Ca _v 3.3 (thalamus)	± 4h ↑ 10days ↑ 31days funct ± 4h/10days/31days RNA _{PCR} ± 4h ↑ 10days ↑ 31days RNA _{PCR} ± 4h ↑ 10days ± 31days RNA _{PCR}	Graef et al., 2009
TISSUE LEVEL ANALYSIS			
HS/noHS human	K _v 4.2 (hipp)	↑ prot W	Aronica et al., 2009
TLE/autopsy human	K _v 7.5 (temporal cortex)	± prot IP	Yus-Najera et al., 2003
TLE/autopsy human	CLC2 (temporal lobe)	↓ RNA _{PCR}	Bertelli et al., 2007
HS-iKA mouse	K _v 3.2 (DG ML)	↓ prot IP	Young et al., 2009
sKA rat	HCN1/2 (EC)	↓ 24h = 1week prot W	Shah et al., 2004
sKA rat	HCN1 (CA1) HCN2 (CA1)	↑ 1–2 ↓ 28–30days prot W ± 1–2days ↓ 28–30days prot W	Shin et al., 2008
sKA rat	HCN1 (CA1/DG) HCN1 (CA3) HCN1 (EC) HCN2 (CA1) HCN2 (CA3) HCN2 (DG) HCN2 (EC)	± 24h ↓ 7days/6weeks RNA _{PCR} ± 24h ↓ 7days ± 6weeks RNA _{PCR} ± 24h/7days/6weeks RNA _{PCR} ↓ 24h/7days/6weeks RNA _{PCR} ↓ 24h ± 7days/6weeks RNA _{PCR} ↓ 24h ± 7days ↓ 6weeks RNA _{PCR} ↓ 24h ± 7days/6weeks RNA _{PCR}	Powell et al., 2008
sPilo rat	K _v 4.2 (CA1)	↓ 30days RNA _{PCR}	Bernard et al., 2004
sPilo rat	K _v 1.4/3.3/3.4/4.2/4.3 (DG)	± RNA _{PCR}	Rüschenschmidt et al., 2006
sPilo rat	KChIP2 (CA1) KChIP2 (DG) K _v 4.2 (CA1) K _v 4.2 (DG)	↓ 1week ↓ 4weeks prot IF ↓ 1week ± 4weeks prot IF ↓ 1week ± 4weeks prot IF ↓ 1week ± 4weeks prot IF	Monaghan et al., 2008
sPilo rat	K _v 4.2 (CA1/2) K _v 4.2 (CA3) K _v 4.2 (DG)	↑ 2days ± 50days prot IF ↑ 2days ↓ 50days prot IF ± prot IF	Su et al., 2008
sPilo rat	K _{2p} 5.1 (CA1) K _{2p} 5.1 (CA3) K _{2p} 5.1 (DG)	↓ 3days–5weeks prot IF (cell loss) ↑ 3days–5weeks prot IF ↑ 3days–5weeks prot IF	Kim et al., 2009
sPilo rat	SK1/2 (hipp) SK3 (hipp) SK1 (hipp) SK2/3 (hipp)	↓ 10days ± chronic prot W ↓ prot W ± RNA _{PCR} ↓ RNA _{PCR}	Oliveira et al., 2010
sPilo rat	SK1/2/3 (CA1) SK2 (CA1) SK1/3 (CA1)	↓ RNA _{PCR} ↓ prot W ± prot W	Schulz et al., 2012
sPilo rat	BK (hilus/CA3)	↓ prot IF/W	Pacheco Otalora et al., 2008
sPilo rat	HCN1 (CA1) HCN2 (CA1)	↓ 3,6days ↓ 3–5weeks prot W ↓ 3,6days ± 3–5weeks prot W	Jung et al., 2007
sPilo rat	HCN1 (CA1)	↓ RNA _{PCR}	Marcelin et al., 2009
sPilo rat	HCN1 (CA1-3/DG) HCN2 (CA1) HCN2 (CA2/3) HCN2 (DG) HCN4 (CA1-3/DG)	↑ 12h ± 2weeks ↑ 11weeks prot IP ↓ 12h ± 7days ↓ 5weeks prot IP ± 12h–5weeks prot IP ↓ 12h–5weeks prot IP ± 12h–5weeks prot IP	Oh et al., 2012

(Continued)

TABLE 1 | Continued

Type of cells and TLE/model	Ion channel/current	Regulation/methods	References
sPilo rat	K _{ir} 2.1 (amyg, hipp, ctx) K _{ir} 4.1 (hipp) K _{ir} 4.1 (amyg, striat, ctx) K _{ir} 5.1 (amyg, hipp, ctx)	± prot W ± prot W IP ↑ prot W IP ± prot W	Nagao et al., 2013
sPilo rat /mouse	Ca _v 3.2 (CA1)	↑ 5days± chronic RNA _{PCR} , prot W	Becker et al., 2008
sPilo mouse	Ca _v 3.1 (hippocampus) Ca _v 3.2 (hippocampus) Ca _v 3.3 (hippocampus)	± 10days/31days RNA _{PCR} ± 10days ↓ 31days RNA _{PCR} ± 10days/31days RNA _{PCR}	Graef et al., 2009
noHS-aKindl rat	HCN1 (CA1/3/DG/EC) HCN2 (CA1/DG/EC) HCN2 (CA3)	± partial/full RNA _{PCR} ± partial/full RNA _{PCR} ± partial ↓ full RNA _{PCR}	Powell et al., 2008
vKA rat	Ca _v 1.2/1.3 (CA1-3/DG) Ca _v 2.1/2.2 (CA1-3/DG) Ca _v α2 (CA3 astrocyt)	± prot IP ± prot IP ↑ prot IP	Westenbroek et al., 1998

1st column: aKindl, hKindl, Epilepsy susceptibility model via repeated electrical stimulation (kindling) of amygdala or hippocampal areas; BCs, Basket cell interneurons; CLC, Chloride channel; dend, Dendritic recording; EC, Entorhinal cortex; HS/noHS human, Tissue from TLE patients with hippocampal sclerosis (HS) compared to samples from TLE patients with mild or no HS (noHS); HS-iKA, intrahippocampal unilateral kainic acid (KA) injection status epilepticus (SE) TLE animal model with severe HS; KChIP, K channel interacting protein (enhances K_v4 function); O-LM, Interneurons projecting from stratum oriens to stratum lacunosum moleculare; PC, Piriform cortex sKA, systemic (intraperitoneal) KA injection SE TLE model; sPilo, systemic pilocarpine injection SE TLE model using adult rats; SP, Interneurons scattered in stratum pyramidale; ventral of iKA, Same iKA model as above but data collected in ventral (noHS) areas; vKA, intraventricular KA injection SE TLE model shows sclerosis in CA3. 2nd column: For ion channel nomenclature see text; amyg, Amygdala; ctx, Cortex; hipp, Hippocampus; I_x, Current x measured (t/pGABA_A, tonic/phasic current via GABA_ARs; GABA_{eff/pot}, efficacy or potency of GABA application; striat, Striatum TRP, Transient receptor potential cation channels; 3rd column: ↑ / ↓, Up-/downregulation (interpreted as antiepileptic/stabilizing (↑ ↓) or as proepileptic/excitatory (↑ ↓) effect; ±, No difference; subscripts, time period in which the changes took place (otherwise chronic) and/or method (full/partial refers to kindling procedure) and/or subcellular location like synaptic (syn), extra - and/or peri synaptic (ex), and surface (surf); funct, Functionally (electrophysiologically) verified via, voltage-clamp (vc) or current-clamp (cc), or with pharmacology (drug); prot, Protein detection via immunofluorescence (IF) or immunoperoxidase (IP) or western blot (W) labeling, or electron microscopy (EM); ISH, RNA detection via *in situ* hybridization; P10/20 sPilo, systemic pilocarpine injection SE TLE model using postnatal day 10 or 20 rats; PCR, mRNA detection via reverse transcriptase (RT)PCR on the tissue level; SC-PCR, Single cell (cell type specific) RT-PCR.

In contrast to the sPilo rats, tissue from TLE patients with HS displays similar hippocampal K_v4.2 mRNA levels compared to patients without HS, and on the protein level K_v4.2 channels are even elevated, despite the cell loss in the CA1 region (Aronica et al., 2009). In a model of cortical heterotopia, K_v4.2 channels are also elevated in CA1 PCs (Castro et al., 2001). In other epilepsy models, K_v4.2 changes were transient (Tsaour et al., 1992) or increased in the chronic phase (Pei et al., 1997) and also K_v1.2 levels renormalize after the acute phase (Tsaour et al., 1992; Pei et al., 1997). Genetic deletion of K_v4.2 alone is not sufficient to create epilepsy (Hu et al., 2006) which may be due to compensatory mechanisms.

A second prominent channelopathy scenario for TLE also takes place in the distal dendrites of CA1 PCs: downregulation of HCN channels (Bender and Baram, 2007; Dyhrfeld-Johnsen et al., 2009; Baruscotti et al., 2010; Noam et al., 2011). Opposite to the described increase in DG GCs, HCN1 channels are decreased in CA1 PCs of both sPilo and sKA rats (Jung et al., 2007; Marcelin et al., 2009; Jung et al., 2011). On the tissue level, upregulation of HCN1 was observed in the CA1-3 regions of sPilo, but it was speculated that this staining could have been due to interneurons (Oh et al., 2012). The HCN-mediated ZD7288-sensitive h-current has a shunting effect not only in DG GCs (Stegen et al., 2012) but also in CA1 PCs (Gasparini and Difrancesco, 1997; Magee, 1999; Berger et al., 2001). Consequently, loss of HCN is usually interpreted as proepileptic (Brewster et al., 2002; Poolos et al., 2002; Jung et al., 2007, 2011; Marcelin et al., 2009). However, there are situations, e.g., in febrile seizure models, where an increase of I_H in CA1 PCs

was interpreted as proepileptic (Chen et al., 2001; Poolos, 2004, 2009; Dyhrfeld-Johnsen et al., 2008, 2009; Noam et al., 2011). Similar to the A-type channelopathy, the HCN channelopathy of noHS models could so far not be confirmed with human data (Bender et al., 2003). In patients with mild HS, abundant HCN1 protein decorates the distal dendrites of surviving CA1 PCs (Stegen et al., 2012). Similar to K_v4 channels, HCN1 loss in CA1 PCs alone does not produce epilepsy, but it can enhance the susceptibility for certain seizure induction protocols (Huang et al., 2009; Poolos, 2009; Santoro et al., 2010).

Other ion channel modifications in CA1 PCs include an enhanced Ca_v channel function in the sPilo, determined with 0.1 mM nickel (Sanabria et al., 2001; Su et al., 2002). This concentration inhibits Ca_v2.3 (R-type) and Ca_v3 (T-type) channels. The Ca_v3.2 RNA from homogenized tissue was indeed found elevated in sPilo, but only in the acute phase and not in the chronic phase (Becker et al., 2008). Another sPilo study detected no changes or even downregulation of Ca_v3 RNA, but also only on the tissue level (Graef et al., 2009). Concerning L-type channels (Ca_v1.2, Ca_v1.3), P-type (Ca_v2.1), N-type channels (Ca_v2.2), and R-type Ca_v channels, no immunocytochemistry differences were detected in CA1 PCs of TLE vs. autopsy samples (Djamshidian et al., 2002). Kindled rats displayed an increased window current in CA1 PCs mediated by Na_v channels at V_{rest} (Ketelaars et al., 2001). These differences appeared unlikely to arise from Na_v1.1, Na_v1.2, and Na_v1.6 channels, as their immuno signal was not changed under similar conditions (Blumenfeld et al., 2009). In contrast, a lowered amount of Na_v1.2 RNA was found via *in situ* hybridization in human TLE

vs. autopsy samples (Whitaker et al., 2001). Small conductance, Ca-activated K (SK) channels mediating the medium duration afterhyperpolarization (AHP) appeared reduced in CA1 PCs of sPilo rats and this phenomenon corresponded to a reduction of the SK2 RNA amount on the tissue level (Schulz et al., 2012). Another sPilo study found transient changes in SK1 and SK2 protein and permanent downregulation of SK3 via western blotting of homogenized hippocampus (Oliveira et al., 2010). In a maximal electroshock convulsions (MES) model, no changes were detected in $K_v1.1/2/4$, $K_v4.2$, and $K_{ir}3.1/2$ channels of CA1 PCs (Pei et al., 1997). In seizure-sensitive gerbils, CA1 PCs displayed elevated $K_v3.1b$ and $K_v3.2$ channel immuno signals (Lee et al., 2009).

The PCs of CA2 and CA3 (in particular CA3a/b) are notably less affected by HS-related cell death than CA1 PCs (Wyler et al., 1992; Blümcke et al., 2012). In samples from TLE patients vs. autopsy, CA3 PCs exhibited increased immunoreactivity for $Ca_v2.1$ subunits (Djamshidian et al., 2002). In the same work, the $Ca_v1.2$ was diminished in CA3 PCs, but enhanced in astrocytes. The $Na_v1.2$ RNA signal was reduced (Whitaker et al., 2001) while the $Ca_v2.1$ immuno signal was elevated in CA3 PCs of HS-TLE patients vs. autopsy (Djamshidian et al., 2002). In noHS models, the RNA and immuno intensities of $Na_v1.6$ channels were found raised in CA3 PCs (Blumenfeld et al., 2009) while those of $K_v4.2$ and $K_v1.1$ channels were decreased, respectively (Francis et al., 1997; Sosanya et al., 2014). Because chronic Na_v upregulation could lead to depolarization block, functional verification is warranted (Auvin et al., 2008; Cestele et al., 2008). The HCN1 RNA was diminished in CA3 PCs of sKA rats, but not after febrile seizures; the HCN2 appeared elevated in both seizure forms, although some of these signals are transient and may be due to interneurons (Brewster et al., 2002). In the MES model, CA cells show little changes in the tested K_v1 , K_v4 , and $K_{ir}3$ channels (Pei et al., 1997) while in an absence epilepsy model CA3 PCs displayed upregulation of some Na_v channels (Xu et al., 2013b). With respect to HS-related iKA seizures, CA3a/b PCs appear similar to DG GCs, i.e., high $K_{ir}2$ levels are found in surviving cells (Young et al., 2009). Thus, more functional characterizations of CA3 cell subtypes in relation to different seizure phenotypes are needed. Also, CA interneurons display ion channel changes in TLE. For example, the oriens-lacunosum moleculare interneurons ventral of the HS area in iKA mice possess reduced I_H and show increased oscillatory activity in the gamma frequency (Dugladze et al., 2007).

In summary, also in the CA regions, pro- and antiepileptic channel changes have been described in TLE. In comparison with DG GCs, particularly the CA1 PCs stand out with proepileptic changes, as here the two most prominent examples of acquired channelopathy have been described. However, in resected tissue of TLE patients no hyperexcitability was detected in CA1 (Cohen et al., 2002). Thus, the CA1 PC-based acquired channelopathy hypothesis either has to be limited for TLE models without HS or it has to be demonstrated that CA1 PCs in non-sclerotic areas of TLE patients are intrinsically hyperexcitable and/or that surviving hyperexcitable CA1 PCs within the sclerotic hippocampus are connected in a hub-like manner (Morgan and Soltesz, 2008). In its current form, the CA1 PC channelopathy

hypothesis collides with the simple principle “dead cells do not seize” (Delorenzo et al., 2005).

Beyond the Hippocampus

Although the entorhinal cortex is a likely source of TLE seizures (Spencer and Spencer, 1994; Spencer, 2002), it received less attention compared to the hippocampus. Nevertheless, there is evidence for ion channel alterations in this region. For example, in layer III PCs of sKA rats, the I_H was found decreased in the chronic phase although western blot signals of HCN1 and HCN2 channel subunits were at control level at the same time (Shah et al., 2004). In kindled rats, the neocortical layer II stellate cells display elevated Na_v currents and $Na_v1.2/1.6$ immunostaining (Hargus et al., 2013). In contrast, a sPilo study found no changes in the intrinsic excitability of layer II PCs and concluded that loss of interneurons was responsible for the elevated perforant path output (Kobayashi et al., 2003). Similarly, $K_v1.1$ reduction in cortical interneurons could play a role in some forms of TLE (Li et al., 2012). In frontal lobe epilepsy, layer II/III PCs displayed h-current downregulation (Wierschke et al., 2010). In some forms of cortical epilepsies, the opposite of the above described hippocampal $K_v4.2$ channelopathy occurs; here these channels are upregulated suggesting homeostatic adaptation of cortical neurons (Aronica et al., 2009).

Other brain areas connected to the hippocampus also display epilepsy-related ion channel adaptations or pathologies. For example, in the subiculum, increased Ca currents were detected in sPilo (Wellmer et al., 2002) while in kindling models, the amygdala showed elevated levels of KCNQ2 ($K_v7.2$) channel protein (Penschuck et al., 2005). Two brain areas rarely implicated in TLE, are the substantia nigra (Depaulis et al., 1994; Ma et al., 2007) and the thalamus (Bertram et al., 2001). One sPilo study found elevated T-type currents in thalamic relay cells and consistent Ca_v3 channel RNA upregulation on the tissue level (Graef et al., 2009). In contrast, the enhanced $K_v4.2$ surface protein expression in ventromedial thalamic relay neurons of sPilo argued for an adaptive response (Smith et al., 2012). We use the thalamus to illustrate that similar ion channels can have opposite functions in different cell types and epilepsies: when sufficiently hyperpolarized, thalamic neurons are capable of rebound burst firing which is mediated by T-type (Ca_v3) channels and thought to underlie absence epilepsy (Budde et al., 2005; Cope et al., 2009; Kanyshkova et al., 2012) although other mechanisms may also play a role (Crunelli and Leresche, 2002; Strauss et al., 2004; Kole et al., 2007). Therefore, it makes sense that absence epilepsy is treated with Ca_v3 channel inhibitors, while hyperpolarizing AEDs can even aggravate absence seizures (Rogawski and Löscher, 2004; Powell et al., 2014). Bursting of “epileptic neurons” has long been suspected as the cellular correlate of epileptic seizures in general (Sypert and Ward, 1967). Also for TLE, pathological Ca_v3 -supported burst firing has been suggested as a cellular mechanism for seizures (Yaari and Beck, 2002). However, the same T-type channel responsible for epileptic bursting in thalamic and cortical neurons could actually prevent burst firing in other cell types (Wolfart and Roeper,

2002). Hence, it is important to determine the role of AED-targeted ion channels in a cell type-specific manner, in particular since current AEDs are applied systemically.

Mechanisms Underlying Epilepsy-Related Ion Channel Alterations

The molecular upstream links of many of the above described ion channel modifications are unclear. For example, the correlation between HS and the leak channel upregulation in DG GCs currently only allows the statement that some part of the sequence, which leads to HS, must also be responsible for the observed changes in channel transcription. One possibility would be that morphological alterations, e.g., related to cytoskeletal changes as they occur during GCD, directly also cause the channel changes (O'Malley and Harvey, 2007). As GCD is caused by reelin deficiency (Haas et al., 2002), reelin is a candidate molecule and indeed it can affect transcription factors (Feng et al., 1999; Chen et al., 2007). However, the hypothesis requires that reelin deficiency alone (without epilepsy) must also trigger the respective channel changes, which does not appear to be the case (Kowalski et al., 2010). Another possible mechanism for transcriptional channelopathies is that the access to neurotrophic factors is interrupted in areas of injury (Waxman, 2001).

We currently favor the hypothesis that the seizures themselves partially cause HS (Mathern et al., 2002) and that the same seizure-induced mechanisms which cause HS, also cause the ion channel changes. The reasoning is as follows. In animal models, SE-related excitotoxic neurodegeneration has two phases: an acute glutamate receptor overstimulation cell swelling phase, and a late Ca-related phase, which gradually fades into the chronic phase of spontaneous seizures, the latter being associated with permanently disturbed intracellular Ca levels (Olney and Sharpe, 1969; Choi, 1992; Magloczky and Freund, 1995; Delorenzo et al., 2005). Also for the iKA model two phases of HS have been described; in the first phase, i.e., hours after KA injection, CA1 PCs and hilar neurons degenerate; in the second phase, about 2 weeks after KA injection, full HS develops with GCD and neurodegeneration affecting CA1, CA3c PCs and eventually also CA3a/b, CA2 PCs, and DG GCs (Bouillieret et al., 1999). While the initial injury has its own channel changes (see below), the second phase of HS-related neurodegeneration in iKA mice correlates well with the described downscaling GC excitability in the same model (Young et al., 2009). Another argument for the hypothesis "chronic seizures cause chronic channel adaptations" is the evidence showing that activity-dependent Ca signals directly couple the degree of excitation to the regulation of gene expression (Marder and Goaillard, 2006). Furthermore, hippocampal seizures are associated with extensive NMDA receptor activation and Ca influx inducing cell stress and neurodegeneration (Isokawa and Levesque, 1991; Magloczky and Freund, 1995; Golowasch et al., 1999; Limbrick et al., 2003; Raza et al., 2004; Ayala and Tapia, 2005; Suzuki et al., 2005). From what we know about the mechanisms of long term synaptic potentiation (LTP), the NMDA-mediated Ca-influx during seizures is likely to activate Ca-dependent kinases (Varga et al., 2004; Nassirpour et al., 2010) and transcription

factors, which in turn regulate the transcription of specific ion channel genes (Scharfman, 2002; Fan et al., 2005; Blair et al., 2008; Mucha et al., 2010). This view is also consistent with many homeostasis studies demonstrating how increased neuronal activity can directly influence ion channel expression (Turriano et al., 1995; Desai et al., 1999; Van Welie et al., 2004; Misonou et al., 2006). Even the subunit-specific heteromerization can be influenced by seizure-like activity (Zha et al., 2008). Another aspect of epilepsy-related channel regulation is the subject of age. Generally, the immature brain appears more sensitive to seizures than the adult brain (Jensen and Baram, 2000), but chronic epilepsy may develop easier in adult animals (Brooks-Kayal et al., 1998; Zhang et al., 2004; Raol et al., 2006b).

Many mechanistic questions remain. For example: if TLE models without HS show similar elevated input to GCs as HS models (Kobayashi et al., 2003; Kumar and Buckmaster, 2006), why do they not display the same channel changes as HS-TLE models? We would like to know what kind of patterns evoke which type of homeostatic ion channel regulation. There are many molecules modified in HS-related TLE (Becker et al., 2003; Elliott et al., 2003; Lukasiuk and Pitkanen, 2004; Motti et al., 2010; Okamoto et al., 2010). Which of these molecules relates to ion channel transformations? A major question is: if the hypothesis of activity- and Ca-dependent homeostatic channel regulation is true, why are there so few adaptive changes in CA1 PCs (Whitaker et al., 2001; Ge et al., 2011)? One possibility would be that CA PCs have a network task that does not allow such homeostatic adaptations. For example, if activity-dependent LTP strengthens synapses on CA1 PCs, it could be counterproductive if the same activity would weaken synaptic impact. In this context, it makes sense that LTP-inducing mechanisms are accompanied by permissive K_v channel internalization (Kim et al., 2007b; Hyun et al., 2013) or HCN1 downregulation (McClelland et al., 2011). On the other hand, it is unlikely that LTP of CA1 PCs is a positive feedback mechanism that exists without homeostatic ion channel control (Abbott and Nelson, 2000). Astonishingly, the survival rate of CA1 PC cells is even higher in the noHS TLE models where exactly these detrimental mechanisms have been discovered. If noHS TLE would be an early stage of HS TLE, the LTP-permissive channel regulation of CA1 PCs could be even stronger in HS-related TLE and in fact be responsible for the CA1 PC degeneration. Back to the initial question: CA1 PCs are indeed also capable of homeostatic ion channel regulation counteracting chronic hyperexcitability (Van Welie et al., 2004, 2006; Otto et al., 2006). Thus, similar to the interaction of LTP and synaptic depression, there must be mechanisms to separate homeostatic and LTP-permissive ion channel regulation within the same neuron.

In addition to the control of gene transcription, epilepsy-related changes in ion channel function can be fine-tuned via post-transcriptional mechanisms like splicing and RNA editing, as well as oxidation or phosphorylation. For example, glycine receptors of hippocampi resected from TLE patients show altered RNA editing, which is particularly relevant when combined with abnormal expression of Cl cotransporter 2 (KCC2) and proconvulsive shift of E_{Cl} (Eichler et al., 2008; Meier et al., 2014). Another example is the increased $K_v1.1$ RNA editing found in

sKA rats (Streit et al., 2011). The intracellular redox state is also known to influence ion channel function via post-translational modulation (Ruppertsberg et al., 1991) and redox-sensitivity of A-type channels appears to be modified in DG GCs from sPilo rats (Rüschenschmidt et al., 2006). Hypoxia is another stimulus for adaptive ion channel modification. For example, in CA1 PCs, reduced oxygen levels lead to I_H downregulation within less than an hour (Zhang et al., 2006). Seizure-related pH changes can also affect channel function and vice versa (Ziemann et al., 2008). Furthermore, phosphorylation is a cellular mechanism, not only to regulate protein trafficking but also to dynamically control ion channel gating (Levitan, 1994). For example, the above discussed (sKA K_v4 and sPilo HCN) channelopathies had both been linked to the intracellular phosphorylation status (Lugo et al., 2008; Jung et al., 2010). Finally, reciprocal changes in phosphorylation and methylation of Na_v channels had been observed with sKA-related seizures (Baek et al., 2014).

Ion channel function depends on the subcellular location to which the channels are targeted and this targeting can be altered in epilepsy (Chung et al., 2006). One example is the subcellular redistribution of dendritic $K_v4.2$ channels in sPilo rats from the inner to the outer molecular layer of the DG (Monaghan et al., 2008). Another example is the disturbed HCN channel trafficking into dendrites of CA1 PCs in epilepsy models (Shin et al., 2008). Furthermore, the axon initial segment and presynaptic terminals have been specifically implicated in TLE (Wimmer et al., 2010; Meier et al., 2014). Thus, ion channels are constantly adapted on pre- and post-translational levels and epilepsy interferes with both levels. Nevertheless, it may be possible to separate the channel changes into those which are part of a cure, i.e., homeostatic in nature, and those which are part of the disease, i.e., either straightforward channelopathy or failure of homeostasis. In any case, AEDs must act on the background of these channel changes. In the next chapter we ask which AEDs have mechanisms of actions comparable to the homeostatic channel regulations observed in neurons during TLE.

Ion Channel-Related Antiepileptic Drug and Gene Therapy Mechanisms

The available AEDs are hypothesized to work via reducing the impact of excitatory ion channels and/or increasing the effect of net inhibitory channels (Löscher et al., 2013). For example, Phenytoin, Carbamazepine, Lamotrigine, Oxcarbazepine, Zonisamide, Rufinamide, Lacosamide, and Eslicarbazepine are thought to act via Na channels, whereas Ethosuximide, Gabapentin, Pregabalin are Ca channel antagonists. Phenobarbital, Primidone, Diazepam, Clonazepam, Clobazam, Progabide, Vigabatrin, and Tiagabine support inhibition via $GABA_A$ channels. Other AEDs like Valproate, Felbamate, and Topiramate are effective at multiple of the above targets. There is only one K channel enhancer AED on the market: the recently approved, first-in-class AED Retigabine (Faulkner and Burke, 2013). Thus, current AEDs mainly target excitatory channels while in contrast, the neurons frequently upregulate K channels when faced with epileptic hyperexcitability

(Table 1). Despite the dissimilar routes of AED mechanisms and cell type-specific homeostasis, some of the cellular strategies are indeed comparable to (potential) AED mechanisms (Table 2). More than 100 K channel subunits are currently known (Coetzee et al., 1999; Goldstein et al., 2005; Gutman et al., 2005; Kubo et al., 2005; Wei et al., 2005) and interaction of native subunits enlarges the number of K currents that can be considered as potential AED targets in preclinical research considerably. Therefore, we propose that cell type-specific approaches based on endogenous homeostasis mechanisms, could guide target-driven development of AEDs.

Early studies investigated ATP-sensitive K_{ir} (K_{ir6}) channel openers such as cromakalim and diazoxide as potential AEDs (Alzheimer and Ten Bruggencate, 1988; Gandolfo et al., 1989a,b). These K_{ir6} channel enhancers were also found protective in anoxia-induced seizures (Mattia et al., 1994; Yamada et al., 2001). However, because K_{ir6} channels are also expressed in the periphery and because the substances were ineffective in standard AED testing models, K_{ir6} channel activators may be of limited utility for epilepsy therapy (Wickenden, 2002; Meldrum and Rogawski, 2007). Systemic administration of SK channel enhancer EBIO reduces seizures in certain seizure models but also produced severe adverse effects (Anderson et al., 2006). Also, Cl channels were considered in the context of TLE (Stogmann et al., 2006; Bertelli et al., 2007; Rinke et al., 2010). Retigabine was initially thought to exert its anticonvulsive action only via GABARs (Rostock et al., 1996; Otto et al., 2002). Later it was shown that Retigabine also activates K_v7 channels, fortunately those of the brain ($K_v7.2-5$) and not those of the heart ($K_v7.1$) raising hopes on the new AED class (Main et al., 2000; Rundfeldt and Netzer, 2000; Wickenden et al., 2000; Tatulian et al., 2001; Dost et al., 2004). However, to establish how retigabine and similar related compounds are best used in epilepsy therapy, still has to be determined (Splinter, 2013).

The $K_v1.1$ channel subunit has often been implicated in epilepsy (Smart et al., 1998; Wenzel et al., 2007; Robbins and Tempel, 2012). A gene therapy approach showed that viral $K_v1.1$ overexpression in neocortical PCs of mice with neocortical epilepsy, effectively reduces the respective seizures (Wykes et al., 2012). A $K_v1.1$ reduction in interneurons was also suggested to play a role in some forms of TLE (Li et al., 2012). Interestingly, precisely the $K_v1.1$ subunits are enhanced endogenously in DG GCs of iKA mice with severe HS (Kirchheim et al., 2013) but apparently not in TLE patients with HS (Stegen et al., 2009, 2012). Thus, $K_v1.1$ enhancement in specific hippocampal neurons could be an excellent antiepileptic and neuroprotective strategy (Kirchheim et al., 2013; Sosanya et al., 2014).

A number of drugs, originally approved for a different action, were later found to enhance K_{ir} channel function; e.g., $K_{ir1.1}$ (Pregabalin) and $K_{ir2.3}$ (Tenidap; Liu et al., 2002; Lee and Liou, 2014). Furthermore, supporting K_{ir3} channel (Kaufmann et al., 2013) and K_{ir2} channel (Xu et al., 2013a) function *in vivo* is effective against seizures of certain epilepsy models (Table 2). Finally, $K_{2p2.1}$ upregulation via adenoviral gene therapy reduces EEG seizures in the sPilo model (Dey et al., 2014). Hence, leak K channels could be an attractive AED target. Again, especially $K_{ir2.1-2.4}$ and $K_{2p2.1}$ leak channels, which are endogenously

TABLE 2 | Inhibitory ion channel changes with antiepileptic potential.

Ion channel	AED, Seizure model	Cell type/Method	References
POTASSIUM CHANNELS			
K _{2P} 2.1 ↑	GenTher (hip, EC, pre SE)/sPilo	Hip neur recs, CellCult, EEG, behavior	Dey et al., 2014
K _{ir} 1.1 ↑	Pregabalin/A	Thal neur recs, HetEx	Lee and Liou, 2014
K _{ir} 2.3 ↑	Tenidap/A*	HetEx recs	Liu et al., 2002
K _{ir} 2.3 ↑	Tenidap/A*/sPilo	EEG, behavior	Xu et al., 2013a
K _{ir} 3.1 ↑	ML297/MES/PTZ	HetEx recs, behavior	Kaufmann et al., 2013
K _{ir} 3.4 ↑			
K _{ir} 6.2 ↑	Pregabalin/A H19-7	CellCult hip neur recs	Huang et al., 2006
K _{ir} 6.x ↑	Cromakalim/diazoxide anoxia seizure model	Field recs	Mattia et al., 1994
K _v 1.1 ↑	PTZ	Behavior	Lu et al., 2008
K _v 1.1 ↑	GenTher (neo, post SE)/NCTX iTX	NCTX PC recs, EEG	Wykes et al., 2012
K _v 1.1 ↑	Rapamycin/sKA	Behavior	Sosanya et al., 2014
K _v 7.2/7.3 ↑	ICA-27243/MES/PTZ/aKindl	Behavior, HetEx recs, Hip neur recs	Roeloffs et al., 2008; Wickenden et al., 2008
K _v 7.x? ↑	Somatostatin/Gen sPent/sKA	CA1 PCs, field recs, behavior	Qiu et al., 2008
K _v 7.2/7.3 ↑	Flupirtine/A*/sKA/FNeo	EEG, behavior	Raol et al., 2009
K _v 7.2/7.3 ↑	Retigabine/A	CA3 PCs recs	Kim et al., 2012
K _v 7.2/7.3 ↑	Retigabine/A/case	EEG, behavior	Walleigh et al., 2013
K _v 7.2/7.3 ↑	Retigabine/A	HetEx recs	Schenzer et al., 2005; Zhou et al., 2013
K _v 7.2–7.5 ↑	ICA-105665 i.p./P ₂	EEG, behavior	Kasteleijn-Nolst Trenite et al., 2013
K _v Del ↓	Lamotrigine/A	CA1 PC recs	Grunze et al., 1998
K _v Del ↓	Levetiracetam/A	Hip neur recs, CellCult	Madeja et al., 2003
K _v A-type ↓	Lamotrigine/A	Hip neur recs, CellCult	Huang et al., 2004
hERG ↓	Lamotrigine/A/Phenytoin/A	HetEx recs	Danielsson et al., 2003, 2005
KCa (SK) ↑	EBIO/MES, PTZ	Behavior	Anderson et al., 2006
KCa (BK) ↑	Zonisamide/A H19-7	Hip neur recs, CellCult	Huang et al., 2007
KCa2 ↑	SKA-19 i.p./aKindl/in vitro PTX/4-AP	CA1 PC recs	Coleman et al., 2015
Na _v ↓			
GABA_A RECEPTORS AND OTHER CHLORIDE CHANNELS			
GABA _{Aα1} ↑	GenTher (DG, pre SE)/sPilo	DG, EG, behavior	Raol et al., 2006a
I _{GABAA} ↑	Retigabine/A	Cort neur recs, CellCult	Otto et al., 2002
Cl NpHR ↑	Optogenetic act/NCTX iTX	EEG	Wykes et al., 2012
GABA _{Aδ} ↑	GenTher/CTZ/sKA	CA1 PC recs, behavior	Sun et al., 2013
HCN CHANNELS			
I _H ↑	Lamotrigine/A	CA1 PCs, dendritic recs	Poolos et al., 2002
I _H ↑	Lamotrigine/A	Interneurons (CA1 _{O-LM} ?)	Peng et al., 2010
I _H ↑	Gabapentin/A	CA1 PCs	Surges et al., 2003

1st column: ↑↓, Manipulated up (↑) and down (↓) regulation of respective ion channel interpreted as pro- or anticonvulsive (red and green, respectively). 2nd column: /A, approved drug as AED or /A* for another indication; /case, case report of centroparietal focal seizures stopped by Retigabine; /P₂, phase 2 clinical study; /GenTher, in vivo gene therapy via stereotaxic intracranial injection of viral vectors in hippocampus (hip), entorhinal cortex (EC) or neocortex (neo). Tested in animal models: /aKindl, amygdala kindling seizure model; /FNeo, rat model of neonatal seizures via convulsant gas flurothyl; MES, maximal electroshock model of epilepsy; NCTX iTX, motorcortex tetanus toxin injection model of neocortical epilepsy; PTZ, systemic pentylenetetrazol injection model of epilepsy; hKindl, hippocampal kindling to SE seizure model; iKA, intrahippocampal unilateral kainic acid injection SE TLE model; Neo sKA, systemic KA injection SE model of neonatal seizures; sPilo, systemic pilocarpine injection SE TLE model; 3rd column: CellCult, cell culture, e.g., of primary fetal cortical neurons; Cor neur, cortical neurons; field recs, extracellular field recordings; HetEx, heterologous expression and pharmacological testing of cloned channels; Hip neur, hippocampal neurons, Cl NpHR, chloride pump halorhodopsin from *Natronomonas pharaonis*, NpHR; recs, electrophysiological recordings; Thal neur, thalamic neurons.

upregulated in DG GCs during epileptic hyperexcitability, appear to be good candidates (Young et al., 2009).

Although enhancing K channel function is a plausible antiepileptic strategy, there are drawbacks. For example, too much silencing via K channels can also be detrimental (Du et al., 2005; Taverna et al., 2005; Coulson et al., 2008).

Furthermore, as discussed for the thalamus, de-inactivation of Na_v and Ca_v channels necessary for AP activation results in the counterintuitive effect that enhancing K currents can increase seizure susceptibility. One example is the big conductance, Ca-activated K (BK) channel. These channels contribute to the fast AHP, enabling high AP frequencies. Consequently, enhancing BK

currents has proepileptic effects (Jin et al., 2000; Brenner et al., 2005; Shruti et al., 2008). If in turn the fast AHP is impaired in interneurons, e.g., via loss of $K_v3.2$ channels, seizures become more likely (Lau et al., 2000). Hence, the usability of K channel enhancing AEDs has to be carefully evaluated.

As explained, certain Cl and cation conductances, mediated by $GABA_A$ Rs and HCN channels, confer shunting inhibition which may also be used as an AED strategy. Indeed, enhancing tonic $GABA_A$ currents by overexpression of $\alpha5/\beta3/\gamma2$ and $\alpha6/\beta3/\delta$ GABARs, reduced epileptiform activity in hippocampal cell culture and elevation of the δ GABARs *in vivo* lowered cyclothiazide (CTZ)-induced seizures (Sun et al., 2013). Also HCN channels are an AED target which requires further investigation (Poolos et al., 2002; Shah et al., 2013): certain AEDs, initially approved as Na_v and Ca_v channel blockers, later turned out to be enhancers of I_H in CA1 PCs as well (Poolos et al., 2002; Surges et al., 2003). However, since one of them (Lamotrigine) was also effective in interneurons (Peng et al., 2010), further experiments must clarify to which extend I_H is involved in the mechanisms of action of these AEDs. As explained above, the combination of K_{ir} and HCN/Cl channel upregulation achieves a homeostatic shunt in GCs during TLE (Young et al., 2009; Stegen et al., 2012). It is tempting to speculate that support of such shunting channel combinations indicated by GCs (which could be called “channelacoids”) is a particularly promising AED strategy.

Impaired interneuron activity, e.g., due to Na_v channel mutations, often increases seizure susceptibility (Lau et al., 2000; Chen et al., 2002; Ogiwara et al., 2007; Martin et al., 2010; Mashimo et al., 2010; Rossignol et al., 2013; De Kovel et al., 2014; Hedrich et al., 2014). Therefore, it makes sense that supporting GABAergic transmission is a successful AED treatment (see above). However, more cell type-specific data is needed to explain how systemically administered AEDs, which block Na_v channels also expressed in interneurons, actually work. Apparently these AEDs preferentially target excitatory neurons (Prakriya and

Mennerick, 2000; He et al., 2002; Pothmann et al., 2014). On the other hand, functional enhancement of interneurons can be an effective AED strategy (Jensen et al., 2014). The next question would be how interneurons inhibiting interneurons fit into these scenarios (Kim et al., 2008a). Generally, the specific role of interneuron subtypes in epilepsy is far from clear, in particular when considering that many of the synchronous AP rhythms, generated by interneurons, are suspiciously akin to epileptic seizures (Cobb et al., 1995; Cohen et al., 2002; D'antuono et al., 2004; Vida et al., 2006). Last but not least, it should be kept in mind that all ion channel abnormalities acquired during epilepsy, can affect their sensitivity for AEDs. Indeed, changes in the channel subunit composition are among the mechanisms proposed to underlie acquired pharmacoresistance (Sun et al., 2007; Zhang et al., 2007; Streit et al., 2011). One example would be a decrease of the $K_v7.2/7.3$ ratio, as suggested by (Otto et al., 2006), which is expected to increase the Retigabine sensitivity of K_v7 channels (Schenzer et al., 2005).

In summary, cell type-specific information on epilepsy-related ion channel modifications can explain and support AED strategies. Precisely those inhibitory ion channels which appear to be effective AED targets in preclinical tests are the ones upregulated in DG GCs during TLE. These data indicate that cell-endogenous ion channel homeostasis mechanisms could be used as “channelacoid” archetypes in the search of antiepileptic strategies. In particular, the enhancement of static shunt via combined K/Cl/cation leak channel support appears to be a promising strategy.

Acknowledgments

JW and DL were/are funded via the grants SFB780/C2 and WO1563/1-1 of the Deutsche Forschungsgemeinschaft (DFG), respectively.

References

- Abbott, L. F., and Nelson, S. B. (2000). Synaptic plasticity: taming the beast. *Nat. Neurosci.* 3(Suppl.), 1178–1183. doi: 10.1038/81453
- Alzheimer, C., and Ten Bruggencate, G. (1988). Actions of BRL 34915 (Cromakalim) upon convulsive discharges in guinea pig hippocampal slices. *Naunyn Schmiedeberg's Arch. Pharmacol.* 337, 429–434. doi: 10.1007/BF00169535
- Anderson, N. J., Slough, S., and Watson, W. P. (2006). *In vivo* characterisation of the small-conductance KCa (SK) channel activator 1-ethyl-2-benzimidazolone (1-EBIO) as a potential anticonvulsant. *Eur. J. Pharmacol.* 546, 48–53. doi: 10.1016/j.ejphar.2006.07.007
- Aronica, E., Boer, K., Doorn, K. J., Zurolo, E., Spliet, W. G., Van Rijen, P. C., et al. (2009). Expression and localization of voltage dependent potassium channel $K_v4.2$ in epilepsy associated focal lesions. *Neurobiol. Dis.* 36, 81–95. doi: 10.1016/j.nbd.2009.06.016
- Auvin, S., Dulac, O., and Vallee, L. (2008). Do SCN1A mutations protect from hippocampal sclerosis? *Epilepsia* 49, 1107–1108. doi: 10.1111/j.1528-1167.2008.01549_3.x
- Ayala, G. X., and Tapia, R. (2005). Late N-methyl-D-aspartate receptor blockade rescues hippocampal neurons from excitotoxic stress and death after 4-aminopyridine-induced epilepsy. *Eur. J. Neurosci.* 22, 3067–3076. doi: 10.1111/j.1460-9568.2005.04509.x
- Baek, J. H., Rubinstein, M., Scheuer, T., and Trimmer, J. S. (2014). Reciprocal changes in phosphorylation and methylation of mammalian brain sodium channels in response to seizures. *J. Biol. Chem.* 289, 15363–15373. doi: 10.1074/jbc.M114.562785
- Barmashenko, G., Hefft, S., Aertsen, A., Kirschstein, T., and Köhling, R. (2011). Positive shifts of the GABA(A) receptor reversal potential due to altered chloride homeostasis is widespread after status epilepticus. *Epilepsia* 52, 1570–1578. doi: 10.1111/j.1528-1167.2011.03247.x
- Bartolomei, F., Gastaldi, M., Massacrier, A., Planells, R., Nicolas, S., and Cau, P. (1997). Changes in the mRNAs encoding subtypes I, II and III sodium channel alpha subunits following kainate-induced seizures in rat brain. *J. Neurocytol.* 26, 667–678. doi: 10.1023/A:1018549928277
- Baruscotti, M., Bottelli, G., Milanese, R., Difrancesco, J. C., and Difrancesco, D. (2010). HCN-related channelopathies. *Pflugers Arch.* 460, 405–415. doi: 10.1007/s00424-010-0810-8
- Beck, H., Blumcke, I., Kral, T., Clusmann, H., Schramm, J., Wiestler, O. D., et al. (1996). Properties of a delayed rectifier potassium current in dentate granule cells isolated from the hippocampus of patients with chronic

- temporal lobe epilepsy. *Epilepsia* 37, 892–901. doi: 10.1111/j.1528-1157.1996.tb00043.x
- Beck, H., Steffens, R., Elger, C. E., and Heinemann, U. (1998). Voltage-dependent Ca^{2+} currents in epilepsy. *Epilepsy Res.* 32, 321–332. doi: 10.1016/S0920-1211(98)00062-X
- Beck, H., and Yaari, Y. (2008). Plasticity of intrinsic neuronal properties in CNS disorders. *Nat. Rev. Neurosci.* 9, 357–369. doi: 10.1038/nrn2371
- Becker, A. J., Chen, J., Zien, A., Sochivko, D., Normann, S., Schramm, J., et al. (2003). Correlated stage- and subfield-associated hippocampal gene expression patterns in experimental and human temporal lobe epilepsy. *Eur. J. Neurosci.* 18, 2792–2802. doi: 10.1111/j.1460-9568.2003.02993.x
- Becker, A. J., Pitsch, J., Sochivko, D., Opitz, T., Staniek, M., Chen, C. C., et al. (2008). Transcriptional upregulation of Cav3.2 mediates epileptogenesis in the pilocarpine model of epilepsy. *J. Neurosci.* 28, 13341–13353. doi: 10.1523/JNEUROSCI.1421-08.2008
- Ben-Ari, Y. (1985). Limbic seizure and brain damage produced by kainic acid: mechanisms and relevance to human temporal lobe epilepsy. *Neuroscience* 14, 375–403. doi: 10.1016/0306-4522(85)90299-4
- Ben-Ari, Y. (2008). Epilepsies and neuronal plasticity: for better or for worse? *Dialogues Clin. Neurosci.* 10, 17–27.
- Bender, R. A., and Baram, T. Z. (2007). Epileptogenesis in the developing brain: what can we learn from animal models? *Epilepsia* 48(Suppl. 5), 2–6. doi: 10.1111/j.1528-1167.2007.01281.x
- Bender, R. A., Soleymani, S. V., Brewster, A. L., Nguyen, S. T., Beck, H., Mathern, G. W., et al. (2003). Enhanced expression of a specific hyperpolarization-activated cyclic nucleotide-gated cation channel (HCN) in surviving dentate gyrus granule cells of human and experimental epileptic hippocampus. *J. Neurosci.* 23, 6826–6836.
- Berger, T., Larkum, M. E., and Luscher, H. R. (2001). High I(h) channel density in the distal apical dendrite of layer V pyramidal cells increases bidirectional attenuation of EPSPs. *J. Neurophysiol.* 85, 855–868.
- Bernard, C., Anderson, A., Becker, A., Poolos, N. P., Beck, H., and Johnston, D. (2004). Acquired dendritic channelopathy in temporal lobe epilepsy. *Science* 305, 532–535. doi: 10.1126/science.1097065
- Bertelli, M., Cecchin, S., Lapucci, C., De Gemmis, P., Danieli, D., D'Amore, E. S., et al. (2007). Quantification of chloride channel 2 (CLCN2) gene isoforms in normal versus lesion- and epilepsy-associated brain tissue. *Biochim. Biophys. Acta* 1772, 15–20. doi: 10.1016/j.bbadis.2006.10.015
- Bertram, E. H., Mangan, P. S., Zhang, D., Scott, C. A., and Williamson, J. M. (2001). The midline thalamus: alterations and a potential role in limbic epilepsy. *Epilepsia* 42, 967–978. doi: 10.1046/j.1528-1157.2001.04200.8967.x
- Biervert, C., Schroeder, B. C., Kubisch, C., Berkovic, S. F., Propping, P., Jentsch, T. J., et al. (1998). A potassium channel mutation in neonatal human epilepsy. *Science* 279, 403–406. doi: 10.1126/science.279.5349.403
- Blair, R. E., Sombati, S., Churn, S. B., and Delorenzo, R. J. (2008). Epileptogenesis causes an N-methyl-D-aspartate receptor/ Ca^{2+} -dependent decrease in Ca^{2+} /calmodulin-dependent protein kinase II activity in a hippocampal neuronal culture model of spontaneous recurrent epileptiform discharges. *Eur. J. Pharmacol.* 588, 64–71. doi: 10.1016/j.ejphar.2008.04.021
- Blümcke, I., Coras, R., Miyata, H., and Ozkara, C. (2012). Defining clinico-neuropathological subtypes of mesial temporal lobe epilepsy with hippocampal sclerosis. *Brain Pathol.* 22, 402–411. doi: 10.1111/j.1750-3639.2012.00583.x
- Blümcke, I., Thom, M., Aronica, E., Armstrong, D. D., Bartolomei, F., Bernasconi, A., et al. (2013). International consensus classification of hippocampal sclerosis in temporal lobe epilepsy: a Task Force report from the ILAE Commission on Diagnostic Methods. *Epilepsia* 54, 1315–1329. doi: 10.1111/epi.12220
- Blümcke, I., Thom, M., and Wiestler, O. D. (2002). Ammon's horn sclerosis: a maldevelopmental disorder associated with temporal lobe epilepsy. *Brain Pathol.* 12, 199–211.
- Blumenfeld, H., Lampert, A., Klein, J. P., Mission, J., Chen, M. C., Rivera, M., et al. (2009). Role of hippocampal sodium channel Nav1.6 in kindling epileptogenesis. *Epilepsia* 50, 44–55. doi: 10.1111/j.1528-1167.2008.01710.x
- Blumenfeld, H., and Meador, K. J. (2014). Consciousness as a useful concept in epilepsy classification. *Epilepsia* 55, 1145–1150. doi: 10.1111/epi.12588
- Boison, D. (2012). Adenosine dysfunction in epilepsy. *Glia* 60, 1234–1243. doi: 10.1002/glia.22285
- Bordey, A., and Sontheimer, H. (1998). Properties of human glial cells associated with epileptic seizure foci. *Epilepsy Res.* 32, 286–303. doi: 10.1016/S0920-1211(98)00059-X
- Bouillere, V., Loup, F., Kiener, T., Marescaux, C., and Fritschy, J. M. (2000). Early loss of interneurons and delayed subunit-specific changes in GABA(A)-receptor expression in a mouse model of mesial temporal lobe epilepsy. *Hippocampus* 10, 305–324. doi: 10.1002/1098-1063(2000)10:3<305::AID-HIPO11>3.0.CO;2-I
- Bouillere, V., Ridoux, V., Depaulis, A., Marescaux, C., Nehlig, A., and Le Gal La Salle, G. (1999). Recurrent seizures and hippocampal sclerosis following intrahippocampal kainate injection in adult mice: electroencephalography, histopathology and synaptic reorganization similar to mesial temporal lobe epilepsy. *Neuroscience* 89, 717–729. doi: 10.1016/S0306-4522(98)00401-1
- Brenner, R., Chen, Q. H., Vilaythong, A., Toney, G. M., Noebels, J. L., and Aldrich, R. W. (2005). BK channel beta4 subunit reduces dentate gyrus excitability and protects against temporal lobe seizures. *Nat. Neurosci.* 8, 1752–1759. doi: 10.1038/nn1573
- Brewster, A., Bender, R. A., Chen, Y., Dube, C., Eghbal-Ahmadi, M., and Baram, T. Z. (2002). Developmental febrile seizures modulate hippocampal gene expression of hyperpolarization-activated channels in an isoform- and cell-specific manner. *J. Neurosci.* 22, 4591–4599.
- Briellmann, R. S., Mark Wellard, R., Masterton, R. A., Abbott, D. F., Berkovic, S. F., and Jackson, G. D. (2007). Hippocampal sclerosis: MR prediction of seizure intractability. *Epilepsia* 48, 315–323. doi: 10.1111/j.1528-1167.2006.00932.x
- Brooks-Kayal, A. R., Shumate, M. D., Jin, H., Rikhter, T. Y., and Coulter, D. A. (1998). Selective changes in single cell GABA(A) receptor subunit expression and function in temporal lobe epilepsy. *Nat. Med.* 4, 1166–1172. doi: 10.1038/2661
- Budde, T., Caputi, L., Kanyshkova, T., Staak, R., Abrahamczik, C., Munsch, T., et al. (2005). Impaired regulation of thalamic pacemaker channels through an imbalance of subunit expression in absence epilepsy. *J. Neurosci.* 25, 9871–9882. doi: 10.1523/JNEUROSCI.2590-05.2005
- Bumanglag, A. V., and Sloviter, R. S. (2008). Minimal latency to hippocampal epileptogenesis and clinical epilepsy after perforant pathway stimulation-induced status epilepticus in awake rats. *J. Comp. Neurol.* 510, 561–580. doi: 10.1002/cne.21801
- Buono, R. J., Lohoff, F. W., Sander, T., Sperling, M. R., O'Connor, M. J., Dlugos, D. J., et al. (2004). Association between variation in the human KCNJ10 potassium ion channel gene and seizure susceptibility. *Epilepsy Res.* 58, 175–183. doi: 10.1016/j.epilepsyres.2004.02.003
- Castro, P. A., Cooper, E. C., Lowenstein, D. H., and Baraban, S. C. (2001). Hippocampal heterotopia lack functional Kv4.2 potassium channels in the methylazoxymethanol model of cortical malformations and epilepsy. *J. Neurosci.* 21, 6626–6634.
- Cavalleri, G. L., Weale, M. E., Shianna, K. V., Singh, R., Lynch, J. M., Grinton, B., et al. (2007). Multicentre search for genetic susceptibility loci in sporadic epilepsy syndrome and seizure types: a case-control study. *Lancet Neurol.* 6, 970–980. doi: 10.1016/S1474-4422(07)70247-8
- Cestele, S., Scalmani, P., Rusconi, R., Terragni, B., Franceschetti, S., and Mantegazza, M. (2008). Self-limited hyperexcitability: functional effect of a familial hemiplegic migraine mutation of the Nav1.1 (SCN1A) Na^{+} channel. *J. Neurosci.* 28, 7273–7283. doi: 10.1523/JNEUROSCI.4453-07.2008
- Charlier, C., Singh, N. A., Ryan, S. G., Lewis, T. B., Reus, B. E., Leach, R. J., et al. (1998). A pore mutation in a novel KQT-like potassium channel gene in an idiopathic epilepsy family. *Nat. Genet.* 18, 53–55. doi: 10.1038/ng098-53
- Chen, C., Bharucha, V., Chen, Y., Westenbroek, R. E., Brown, A., Malhotra, J. D., et al. (2002). Reduced sodium channel density, altered voltage dependence of inactivation, and increased susceptibility to seizures in mice lacking sodium channel beta 2-subunits. *Proc. Natl. Acad. Sci. U.S.A.* 99, 17072–17077. doi: 10.1073/pnas.212638099
- Chen, K., Aradi, I., Thon, N., Eghbal-Ahmadi, M., Baram, T. Z., and Soltesz, I. (2001). Persistently modified h-channels after complex febrile seizures convert the seizure-induced enhancement of inhibition to hyperexcitability. *Nat. Med.* 7, 331–337. doi: 10.1038/85480
- Chen, Y., Kundakovic, M., Agis-Balboa, R. C., Pinna, G., and Grayson, D. R. (2007). Induction of the reelin promoter by retinoic acid is mediated by Sp1. *J. Neurochem.* 103, 650–665. doi: 10.1111/j.1471-4159.2007.04797.x

- Chioza, B., Osei-Lah, A., Wilkie, H., Nashef, L., McCormick, D., Asherson, P., et al. (2002). Suggestive evidence for association of two potassium channel genes with different idiopathic generalised epilepsy syndromes. *Epilepsy Res.* 52, 107–116. doi: 10.1016/S0920-1211(02)00195-X
- Choi, D. W. (1992). Excitotoxic cell death. *J. Neurobiol.* 23, 1261–1276. doi: 10.1002/neu.480230915
- Chung, H. J., Jan, Y. N., and Jan, L. Y. (2006). Polarized axonal surface expression of neuronal KCNQ channels is mediated by multiple signals in the KCNQ2 and KCNQ3 C-terminal domains. *Proc. Natl. Acad. Sci. U.S.A.* 103, 8870–8875. doi: 10.1073/pnas.0603376103
- Cobb, S. R., Buhl, E. H., Halasy, K., Paulsen, O., and Somogyi, P. (1995). Synchronization of neuronal activity in hippocampus by individual GABAergic interneurons. *Nature* 378, 75–78. doi: 10.1038/378075a0
- Coetzee, W. A., Amarillo, Y., Chiu, J., Chow, A., Lau, D., McCormack, T., et al. (1999). Molecular diversity of K⁺ channels. *Ann. N.Y. Acad. Sci.* 868, 233–285. doi: 10.1111/j.1749-6632.1999.tb11293.x
- Cohen, I., Navarro, V., Clemenceau, S., Baulac, M., and Miles, R. (2002). On the origin of interictal activity in human temporal lobe epilepsy *in vitro*. *Science* 298, 1418–1421. doi: 10.1126/science.1076510
- Coleman, N., Nguyen, H. M., Cao, Z., Brown, B. M., Jenkins, D. P., Zolkowska, D., et al. (2015). The riluzole derivative 2-amino-6-trifluoromethylthio-benzothiazole (SKA-19), a mixed K₂ Activator and na blocker, is a potent novel anticonvulsant. *Neurotherapeutics* 12, 234–249. doi: 10.1007/s13311-014-0305-y
- Cope, D. W., Di Giovanni, G., Fyson, S. J., Orban, G., Errington, A. C., Lorincz, M. L., et al. (2009). Enhanced tonic GABAA inhibition in typical absence epilepsy. *Nat. Med.* 15, 1392–1398. doi: 10.1038/nm.2058
- Coulson, E. J., May, L. M., Osborne, S. L., Reid, K., Underwood, C. K., Meunier, F. A., et al. (2008). p75 neurotrophin receptor mediates neuronal cell death by activating GIRK channels through phosphatidylinositol 4,5-bisphosphate. *J. Neurosci.* 28, 315–324. doi: 10.1523/JNEUROSCI.2699-07.2008
- Coulter, D. A. (2000). Mossy fiber zinc and temporal lobe epilepsy: pathological association with altered “epileptic” gamma-aminobutyric acid A receptors in dentate granule cells. *Epilepsia* 41(Suppl. 6), S96–S99. doi: 10.1111/j.1528-1157.2000.tb01565.x
- Coulter, D. A., McIntyre, D. C., and Löscher, W. (2002). Animal models of limbic epilepsies: what can they tell us? *Brain Pathol.* 12, 240–256. doi: 10.1111/j.1750-3639.2002.tb00439.x
- Crunelli, V., and Leresche, N. (2002). Block of thalamic T-Type Ca(2+) channels by ethosuximide is not the whole story. *Epilepsy Curr.* 2, 53–56. doi: 10.1046/j.1535-7597.2002.00024.x
- Curia, G., Lucchi, C., Vinet, J., Gualtieri, F., Marinelli, C., Torsello, A., et al. (2014). Pathophysiology of mesial temporal lobe epilepsy: is prevention of damage antiepileptogenic? *Curr. Med. Chem.* 21, 663–688. doi: 10.2174/0929867320666131119152201
- D’adamo, M. C., Catacuzzeno, L., Di Giovanni, G., Franciolini, F., and Pessia, M. (2013). K(+) channelopathy: progress in the neurobiology of potassium channels and epilepsy. *Front. Cell. Neurosci.* 7:134. doi: 10.3389/fncel.2013.00134
- D’antuono, M., Louvel, J., Köhling, R., Mattia, D., Bernasconi, A., Olivier, A., et al. (2004). GABAA receptor-dependent synchronization leads to ictogenesis in the human dysplastic cortex. *Brain* 127, 1626–1640. doi: 10.1093/brain/awh181
- Danielsson, B. R., Lansdell, K., Patmore, L., and Tomson, T. (2003). Phenytoin and phenobarbital inhibit human HERG potassium channels. *Epilepsy Res.* 55, 147–157. doi: 10.1016/S0920-1211(03)00119-0
- Danielsson, B. R., Lansdell, K., Patmore, L., and Tomson, T. (2005). Effects of the antiepileptic drugs lamotrigine, topiramate and gabapentin on HERG potassium currents. *Epilepsy Res.* 63, 17–25. doi: 10.1016/j.eplepsyres.2004.10.002
- Davis, G. W., and Bezprozvany, I. (2001). Maintaining the stability of neural function: a homeostatic hypothesis. *Annu. Rev. Physiol.* 63, 847–869. doi: 10.1146/annurev.physiol.63.1.847
- De Kovel, C. G., Meisler, M. H., Brilstra, E. H., Van Berkestijn, F. M., Van ’T Slot, R., Van Lieshout, S., et al. (2014). Characterization of a de novo SCN8A mutation in a patient with epileptic encephalopathy. *Epilepsy Res.* 108, 1511–1518. doi: 10.1016/j.eplepsyres.2014.08.020
- De Lanerolle, N. C., Brines, M., Williamson, A., Kim, J. H., and Spencer, D. D. (1992). Neurotransmitters and their receptors in human temporal lobe epilepsy. *Epilepsy Res. Suppl.* 7, 235–250.
- De Lanerolle, N. C., Kim, J. H., Williamson, A., Spencer, S. S., Zaveri, H. P., Eid, T., et al. (2003). A retrospective analysis of hippocampal pathology in human temporal lobe epilepsy: evidence for distinctive patient subcategories. *Epilepsia* 44, 677–687. doi: 10.1046/j.1528-1157.2003.32701.x
- De Lanerolle, N. C., and Lee, T. S. (2005). New facets of the neuropathology and molecular profile of human temporal lobe epilepsy. *Epilepsy Behav.* 7, 190–203. doi: 10.1016/j.yebeh.2005.06.003
- De Tisi, J., Bell, G. S., Peacock, J. L., Mcevoy, A. W., Harkness, W. F., Sander, J. W., et al. (2011). The long-term outcome of adult epilepsy surgery, patterns of seizure remission, and relapse: a cohort study. *Lancet* 378, 1388–1395. doi: 10.1016/S0140-6736(11)60890-8
- Delorenzo, R. J., Sun, D. A., and Deshpande, L. S. (2005). Cellular mechanisms underlying acquired epilepsy: the calcium hypothesis of the induction and maintenance of epilepsy. *Pharmacol. Ther.* 105, 229–266. doi: 10.1016/j.pharmthera.2004.10.004
- Depaulis, A., Vergnes, M., and Marescaux, C. (1994). Endogenous control of epilepsy: the nigral inhibitory system. *Prog. Neurobiol.* 42, 33–52. doi: 10.1016/0301-0082(94)90020-5
- Desai, N. S., Rutherford, L. C., and Turrigiano, G. G. (1999). Plasticity in the intrinsic excitability of cortical pyramidal neurons. *Nat. Neurosci.* 2, 515–520. doi: 10.1038/9165
- Dey, D., Eckle, V. S., Vitko, I., Sullivan, K. A., Lasiecka, Z. M., Winckler, B., et al. (2014). A potassium leak channel silences hyperactive neurons and ameliorates status epilepticus. *Epilepsia* 55, 203–213. doi: 10.1111/epi.12472
- Dietrich, D., Clusmann, H., Kral, T., Steinhauser, C., Blumcke, I., Heinemann, U., et al. (1999). Two electrophysiologically distinct types of granule cells in epileptic human hippocampus. *Neuroscience* 90, 1197–1206. doi: 10.1016/S0306-4522(98)00574-0
- Dietrich, D., Podlogar, M., Ortman, G., Clusmann, H., and Kral, T. (2005). Calbindin-D28k content and firing pattern of hippocampal granule cells in amygdala-kindled rats: a perforated patch-clamp study. *Brain Res.* 1032, 123–130. doi: 10.1016/j.brainres.2004.10.060
- Djamshidian, A., Grassl, R., Seltenthaler, M., Czech, T., Baumgartner, C., Schmidbauer, M., et al. (2002). Altered expression of voltage-dependent calcium channel alpha(1) subunits in temporal lobe epilepsy with Ammon’s horn sclerosis. *Neuroscience* 111, 57–69. doi: 10.1016/S0306-4522(01)00528-0
- Dost, R., Rostock, A., and Rundfeldt, C. (2004). The anti-hyperalgesic activity of retigabine is mediated by KCNQ potassium channel activation. *Naunyn Schmiedeberg’s Arch. Pharmacol.* 369, 382–390. doi: 10.1007/s00210-004-0881-1
- Du, W., Bautista, J. F., Yang, H., Diez-Sampedro, A., You, S. A., Wang, L., et al. (2005). Calcium-sensitive potassium channelopathy in human epilepsy and paroxysmal movement disorder. *Nat. Genet.* 37, 733–738. doi: 10.1038/ng1585
- Dugladze, T., Vida, I., Tort, A. B., Gross, A., Otahal, J., Heinemann, U., et al. (2007). Impaired hippocampal rhythmicity in a mouse model of mesial temporal lobe epilepsy. *Proc. Natl. Acad. Sci. U.S.A.* 104, 17530–17535. doi: 10.1073/pnas.0708301104
- Dyhrfeld-Johnsen, J., Morgan, R. J., Foldy, C., and Soltesz, I. (2008). Upregulated H-Current in Hyperexcitable CA1 Dendrites after Febrile Seizures. *Front. Cell. Neurosci.* 2:2. doi: 10.3389/neuro.03.002.2008
- Dyhrfeld-Johnsen, J., Morgan, R. J., and Soltesz, I. (2009). Double trouble? potential for hyperexcitability following both channelopathic up- and downregulation of I(h) in epilepsy. *Front. Neurosci.* 3, 25–33. doi: 10.3389/neuro.01.005.2009
- Eichler, S. A., Kirischuk, S., Jüttner, R., Schaefermeier, P. K., Legendre, P., Lehmann, T. N., et al. (2008). Glycinergic tonic inhibition of hippocampal neurons with depolarizing GABAergic transmission elicits histopathological signs of temporal lobe epilepsy. *J. Cell. Mol. Med.* 12, 2848–2866. doi: 10.1111/j.1582-4934.2008.00357.x
- Elliott, R. C., Miles, M. F., and Lowenstein, D. H. (2003). Overlapping microarray profiles of dentate gyrus gene expression during development- and epilepsy-associated neurogenesis and axon outgrowth. *J. Neurosci.* 23, 2218–2227.
- Fan, Y., Fricker, D., Brager, D. H., Chen, X., Lu, H. C., Chitwood, R. A., et al. (2005). Activity-dependent decrease of excitability in rat hippocampal neurons through increases in I(h). *Nat. Neurosci.* 8, 1542–1551. doi: 10.1038/nn1568
- Faulkner, M. A., and Burke, R. A. (2013). Safety profile of two novel antiepileptic agents approved for the treatment of refractory partial seizures: ezogabine

- (retigabine) and perampanel. *Expert Opin. Drug Saf.* 12, 847–855. doi: 10.1517/14740338.2013.823399
- Feng, Z., Chang, R. C., Bing, G., Hudson, P., Tiao, N., Jin, L., et al. (1999). Long-term increase of Sp-1 transcription factors in the hippocampus after kainic acid treatment. *Brain Res. Mol. Brain Res.* 69, 144–148. doi: 10.1016/S0169-328X(99)00099-6
- Francis, J., Jugloff, D. G., Mingo, N. S., Wallace, M. C., Jones, O. T., Burnham, W. M., et al. (1997). Kainic acid-induced generalized seizures alter the regional hippocampal expression of the rat Kv4.2 potassium channel gene. *Neurosci. Lett.* 232, 91–94. doi: 10.1016/S0304-3940(97)00593-4
- Fritschy, J. M. (2004). A new animal model of temporal lobe epilepsy. *Epileptologie*, 21–28.
- Gandolfo, G., Gottesmann, C., Bidard, J. N., and Lazdunski, M. (1989a). K⁺ channels openers prevent epilepsy induced by the bee venom peptide MCD. *Eur. J. Pharmacol.* 159, 329–330.
- Gandolfo, G., Romettino, S., Gottesmann, C., Van Luijtelar, G., Coenen, A., Bidard, J. N., et al. (1989b). K⁺ channel openers decrease seizures in genetically epileptic rats. *Eur. J. Pharmacol.* 167, 181–183.
- Gasparini, S., and Difrancesco, D. (1997). Action of the hyperpolarization-activated current (I_h) blocker ZD 7288 in hippocampal CA1 neurons. *Pflugers Arch.* 435, 99–106. doi: 10.1007/s004240050488
- Gavrilovici, C., Pollock, E., Everest, M., and Poulter, M. O. (2012). The loss of interneuron functional diversity in the piriform cortex after induction of experimental epilepsy. *Neurobiol. Dis.* 48, 317–328. doi: 10.1016/j.nbd.2012.07.002
- Ge, Y. X., Liu, Y., Tang, H. Y., Liu, X. G., and Wang, X. (2011). CLC-2 contributes to tonic inhibition mediated by alpha5 subunit-containing GABA(A) receptor in experimental temporal lobe epilepsy. *Neuroscience* 186, 120–127. doi: 10.1016/j.neuroscience.2011.04.029
- George, A. L. Jr. (2004). Inherited channelopathies associated with epilepsy. *Epilepsy Curr.* 4, 65–70. doi: 10.1111/j.1535-7597.2004.42010.x
- Gibbs, J. W. III, Shumate, M. D., and Coulter, D. A. (1997). Differential epilepsy-associated alterations in postsynaptic GABA(A) receptor function in dentate granule and CA1 neurons. *J. Neurophysiol.* 77, 1924–1938.
- Glykys, J., Mann, E. O., and Mody, I. (2008). Which GABA(A) receptor subunits are necessary for tonic inhibition in the hippocampus? *J. Neurosci.* 28, 1421–1426. doi: 10.1523/JNEUROSCI.4751-07.2008
- Goddard, G. V., McIntyre, D. C., and Leech, C. K. (1969). A permanent change in brain function resulting from daily electrical stimulation. *Exp. Neurol.* 25, 295–330. doi: 10.1016/0014-4886(69)90128-9
- Goldstein, S. A., Bayliss, D. A., Kim, D., Lesage, F., Plant, L. D., and Rajan, S. (2005). International union of pharmacology. LV. Nomenclature and molecular relationships of two-P potassium channels. *Pharmacol. Rev.* 57, 527–540. doi: 10.1124/pr.57.4.12
- Golowasch, J., Abbott, L. F., and Marder, E. (1999). Activity-dependent regulation of potassium currents in an identified neuron of the stomatogastric ganglion of the crab *Cancer borealis*. *J. Neurosci.* 19:RC33.
- Goodkin, H. P., Joshi, S., Mtchedlishvili, Z., Brar, J., and Kapur, J. (2008). Subunit-specific trafficking of GABA(A) receptors during status epilepticus. *J. Neurosci.* 28, 2527–2538. doi: 10.1523/JNEUROSCI.3426-07.2008
- Gower, W. R. (1881). *Epilepsy and Other Chronic Convulsive Disorders, their Causes, Symptoms and Treatment*. London: J & A Churchill.
- Graef, J. D., Nordskog, B. K., Wiggins, W. F., and Godwin, D. W. (2009). An acquired channelopathy involving thalamic T-type Ca²⁺ channels after status epilepticus. *J. Neurosci.* 29, 4430–4441. doi: 10.1523/JNEUROSCI.0198-09.2009
- Grunze, H., Von Wegerer, J., Greene, R. W., and Walden, J. (1998). Modulation of calcium and potassium currents by lamotrigine. *Neuropsychobiology* 38, 131–138. doi: 10.1159/000026528
- Gutman, G. A., Chandy, K. G., Grissmer, S., Lazdunski, M., McKinnon, D., Pardo, L. A., et al. (2005). International Union of Pharmacology. LIII. Nomenclature and molecular relationships of voltage-gated potassium channels. *Pharmacol. Rev.* 57, 473–508. doi: 10.1124/pr.57.4.10
- Haas, C. A., Dudeck, O., Kirsch, M., Huszka, C., Kann, G., Pollak, S., et al. (2002). Role for reelin in the development of granule cell dispersion in temporal lobe epilepsy. *J. Neurosci.* 22, 5797–5802.
- Hargus, N. J., Nigam, A., Bertram, E. H. III, and Patel, M. K. (2013). Evidence for a role of Nav1.6 in facilitating increases in neuronal hyperexcitability during epileptogenesis. *J. Neurophysiol.* 110, 1144–1157. doi: 10.1152/jn.00383.2013
- Harvey, B. D., and Sloviter, R. S. (2005). Hippocampal granule cell activity and c-Fos expression during spontaneous seizures in awake, chronically epileptic, pilocarpine-treated rats: implications for hippocampal epileptogenesis. *J. Comp. Neurol.* 488, 442–463. doi: 10.1002/cne.20594
- Hauser, W. A., and Lee, J. R. (2002). Do seizures beget seizures? *Prog. Brain Res.* 135, 215–219.
- Häussler, U., Bielefeld, L., Frieppel, U. P., Wolffart, J., and Haas, C. A. (2012). Septotemporal position in the hippocampal formation determines epileptic and neurogenic activity in temporal lobe epilepsy. *Cereb. Cortex* 22, 26–36. doi: 10.1093/cercor/bhr054
- He, Y., Zorumski, C. F., and Mennerick, S. (2002). Contribution of presynaptic Na⁺ channel inactivation to paired-pulse synaptic depression in cultured hippocampal neurons. *J. Neurophysiol.* 87, 925–936.
- Hedrich, U. B., Liautard, C., Kirschenbaum, D., Pofahl, M., Lavigne, J., Liu, Y., et al. (2014). Impaired Action Potential Initiation in GABAergic Interneurons Causes Hyperexcitable Networks in an Epileptic Mouse Model Carrying a Human NaV1.1 Mutation. *J. Neurosci.* 34, 14874–14889. doi: 10.1523/JNEUROSCI.0721-14.2014
- Heilstedt, H. A., Burgess, D. L., Anderson, A. E., Chedrawi, A., Tharp, B., Lee, O., et al. (2001). Loss of the potassium channel beta-subunit gene, KCNAB2, is associated with epilepsy in patients with 1p36 deletion syndrome. *Epilepsia* 42, 1103–1111. doi: 10.1046/j.1528-1157.2001.08801.x
- Heinemann, U., Beck, H., Dreier, J. P., Ficker, E., Stabel, J., and Zhang, C. L. (1992). The dentate gyrus as a regulated gate for the propagation of epileptiform activity. *Epilepsy Res. Suppl.* 7, 273–280.
- Heuser, K., Eid, T., Lauritzen, F., Thoren, A. E., Vindedal, G. F., Tauboll, E., et al. (2012). Loss of perivascular Kir4.1 potassium channels in the sclerotic hippocampus of patients with mesial temporal lobe epilepsy. *J. Neuropathol. Exp. Neurol.* 71, 814–825. doi: 10.1097/NEN.0b013e318267b5af
- Hinterkeuser, S., Schroder, W., Hager, G., Seifert, G., Blumcke, I., Elger, C. E., et al. (2000). Astrocytes in the hippocampus of patients with temporal lobe epilepsy display changes in potassium conductances. *Eur. J. Neurosci.* 12, 2087–2096. doi: 10.1046/j.1460-9568.2000.00104.x
- Hoffman, E. P. (1995). Voltage-gated ion channelopathies: inherited disorders caused by abnormal sodium, chloride, and calcium regulation in skeletal muscle. *Annu. Rev. Med.* 46, 431–441. doi: 10.1146/annurev.med.46.1.431
- Houser, C. R. (1990). Granule cell dispersion in the dentate gyrus of humans with temporal lobe epilepsy. *Brain Res.* 535, 195–204. doi: 10.1016/0006-8993(90)91601-C
- Hsu, D. (2007). The dentate gyrus as a filter or gate: a look back and a look ahead. *Prog. Brain Res.* 163, 601–613. doi: 10.1016/S0079-6123(07)63032-5
- Hu, H. J., Carrasquillo, Y., Karim, F., Jung, W. E., Nerbonne, J. M., Schwarz, T. L., et al. (2006). The kv4.2 potassium channel subunit is required for pain plasticity. *Neuron* 50, 89–100. doi: 10.1016/j.neuron.2006.03.010
- Huang, C. W., Huang, C. C., Liu, Y. C., and Wu, S. N. (2004). Inhibitory effect of lamotrigine on A-type potassium current in hippocampal neuron-derived H19-7 cells. *Epilepsia* 45, 729–736. doi: 10.1111/j.0013-9580.2004.58403.x
- Huang, C. W., Huang, C. C., and Wu, S. N. (2006). The opening effect of pregabalin on ATP-sensitive potassium channels in differentiated hippocampal neuron-derived H19-7 cells. *Epilepsia* 47, 720–726. doi: 10.1111/j.1528-1167.2006.00498.x
- Huang, C. W., Huang, C. C., and Wu, S. N. (2007). Activation by zonisamide, a newer antiepileptic drug, of large-conductance calcium-activated potassium channel in differentiated hippocampal neuron-derived H19-7 cells. *J. Pharmacol. Exp. Ther.* 321, 98–106. doi: 10.1124/jpet.106.116954
- Huang, Z., Walker, M. C., and Shah, M. M. (2009). Loss of dendritic HCN1 subunits enhances cortical excitability and epileptogenesis. *J. Neurosci.* 29, 10979–10988. doi: 10.1523/JNEUROSCI.1531-09.2009
- Huberfeld, G., Wittner, L., Clemenceau, S., Baulac, M., Kaila, K., Miles, R., et al. (2007). Perturbed chloride homeostasis and GABAergic signaling in human temporal lobe epilepsy. *J. Neurosci.* 27, 9866–9873. doi: 10.1523/JNEUROSCI.2761-07.2007
- Hyun, J. H., Eom, K., Lee, K. H., Ho, W. K., and Lee, S. H. (2013). Activity-dependent downregulation of D-type K⁺ channel subunit Kv1.2 in rat hippocampal CA3 pyramidal neurons. *J. Physiol. (Lond)*.

- Ishii, A., Fukuma, G., Uehara, A., Miyajima, T., Makita, Y., Hamachi, A., et al. (2009). A de novo KCNQ2 mutation detected in non-familial benign neonatal convulsions. *Brain Dev.* 31, 27–33. doi: 10.1016/j.braindev.2008.05.010
- Isokawa, M. (1996). Decrement of GABAA receptor-mediated inhibitory postsynaptic currents in dentate granule cells in epileptic hippocampus. *J. Neurophysiol.* 75, 1901–1908.
- Isokawa, M., and Levesque, M. F. (1991). Increased NMDA responses and dendritic degeneration in human epileptic hippocampal neurons in slices. *Neurosci. Lett.* 132, 212–216. doi: 10.1016/0304-3940(91)90304-C
- Isokawa, M., and Mello, L. E. (1991). NMDA receptor-mediated excitability in dendritically deformed dentate granule cells in pilocarpine-treated rats. *Neurosci. Lett.* 129, 69–73. doi: 10.1016/0304-3940(91)90722-6
- Jabs, R., Seifert, G., and Steinhauser, C. (2008). Astrocytic function and its alteration in the epileptic brain. *Epilepsia* 49(Suppl. 2), 3–12. doi: 10.1111/j.1528-1167.2008.01488.x
- Jefferys, J. G. (1999). Hippocampal sclerosis and temporal lobe epilepsy: cause or consequence? *Brain* 122(Pt 6), 1007–1008.
- Jensen, F. E., and Baram, T. Z. (2000). Developmental seizures induced by common early-life insults: short- and long-term effects on seizure susceptibility. *Ment. Retard. Dev. Disabil. Res. Rev.* 6, 253–257. doi: 10.1002/1098-2779(2000)6:4<253::AID-MRDD4>3.0.CO;2-P
- Jensen, H. S., Grunnet, M., and Bastlund, J. F. (2014). Therapeutic potential of Na(V)1.1 activators. *Trends Pharmacol. Sci.* 35, 113–118. doi: 10.1016/j.tips.2013.12.007
- Jin, W., Sugaya, A., Tsuda, T., Ohguchi, H., and Sugaya, E. (2000). Relationship between large conductance calcium-activated potassium channel and bursting activity. *Brain Res.* 860, 21–28. doi: 10.1016/S0006-8993(00)01943-0
- Jobst, B. C., and Cascino, G. D. (2015). Resective epilepsy surgery for drug-resistant focal epilepsy: a review. *JAMA* 313, 285–293. doi: 10.1001/jama.2014.17426
- Jung, S., Bullis, J. B., Lau, I. H., Jones, T. D., Warner, L. N., and Poolos, N. P. (2010). Downregulation of dendritic HCN channel gating in epilepsy is mediated by altered phosphorylation signaling. *J. Neurosci.* 30, 6678–6688. doi: 10.1523/JNEUROSCI.1290-10.2010
- Jung, S., Jones, T. D., Lugo, J. N. Jr., Sheerin, A. H., Miller, J. W., D'ambrosio, R., et al. (2007). Progressive dendritic HCN channelopathy during epileptogenesis in the rat pilocarpine model of epilepsy. *J. Neurosci.* 27, 13012–13021. doi: 10.1523/JNEUROSCI.3605-07.2007
- Jung, S., Warner, L. N., Pitsch, J., Becker, A. J., and Poolos, N. P. (2011). Rapid loss of dendritic HCN channel expression in hippocampal pyramidal neurons following status epilepticus. *J. Neurosci.* 31, 14291–14295. doi: 10.1523/JNEUROSCI.1148-11.2011
- Kanyshkova, T., Meuth, P., Bista, P., Liu, Z., Ehling, P., Caputi, L., et al. (2012). Differential regulation of HCN channel isoform expression in thalamic neurons of epileptic and non-epileptic rat strains. *Neurobiol. Dis.* 45, 450–461. doi: 10.1016/j.nbd.2011.08.032
- Kasteleijn-Nolst Trenite, D. G., Biton, V., French, J. A., Abou-Khalil, B., Rosenfeld, W. E., Diventura, B., et al. (2013). Kv7 potassium channel activation with ICA-105665 reduces photoparoxysmal EEG responses in patients with epilepsy. *Epilepsia* 54, 1437–1443. doi: 10.1111/epi.12224
- Kaufmann, K., Romaine, I., Days, E., Pascual, C., Malik, A., Yang, L., et al. (2013). ML297 (VU0456810), the first potent and selective activator of the GIRK potassium channel, displays antiepileptic properties in mice. *ACS Chem. Neurosci.* 4, 1278–1286. doi: 10.1021/cn400062a
- Ketelaars, S. O., Gorter, J. A., van Vliet, E. A., Lopes da Silva, F. H., and Wadman, W. J. (2001). Sodium currents in isolated rat CA1 pyramidal and dentate granule neurons in the post-status epilepticus model of epilepsy. *Neuroscience* 105, 109–120. doi: 10.1016/S0306-4522(01)00176-2
- Khirug, S., Ahmad, F., Puskarjov, M., Afzalov, R., Kaila, K., and Blaesse, P. (2010). A Single Seizure episode leads to rapid functional activation of KCC2 in the neonatal rat hippocampus. *J. Neurosci.* 30, 12028–12035. doi: 10.1523/JNEUROSCI.3154-10.2010
- Kienzler, F., Norwood, B. A., and Sloviter, R. S. (2009). Hippocampal injury, atrophy, synaptic reorganization, and epileptogenesis after perforant pathway stimulation-induced status epilepticus in the mouse. *J. Comp. Neurol.* 515, 181–196. doi: 10.1002/cne.22059
- Kim, D. S., Kim, J. E., Kwak, S. E., Choi, H. C., Song, H. K., Kim, Y. I., et al. (2007a). Up-regulated astroglial TWIK-related acid-sensitive K⁺ channel-1 (TASK-1) in the hippocampus of seizure-sensitive gerbils: a target of anti-epileptic drugs. *Brain Res.* 1185, 346–358. doi: 10.1016/j.brainres.2007.09.043
- Kim, D. S., Ryu, H. J., Kim, J. E., and Kang, T. C. (2013). The reverse roles of transient receptor potential canonical channel-3 and -6 in neuronal death following pilocarpine-induced status epilepticus. *Cell. Mol. Neurobiol.* 33, 99–109. doi: 10.1007/s10571-012-9875-6
- Kim, J., Jung, S. C., Clemens, A. M., Petralia, R. S., and Hoffman, D. A. (2007b). Regulation of dendritic excitability by activity-dependent trafficking of the A-type K⁺ channel subunit Kv4.2 in hippocampal neurons. *Neuron* 54, 933–947. doi: 10.1016/j.neuron.2007.05.026
- Kim, J. E., Kwak, S. E., Choi, H. C., Song, H. K., Kim, Y. I., Jo, S. M., et al. (2008a). Voltage-gated Na⁺ channel II immunoreactivity is selectively up-regulated in hippocampal interneurons of seizure sensitive gerbils. *Neurosci. Lett.* 438, 295–299. doi: 10.1016/j.neulet.2008.04.079
- Kim, J. E., Kwak, S. E., Choi, S. Y., and Kang, T. C. (2008b). Region-specific alterations in astroglial TWIK-related acid-sensitive K⁺-1 channel immunoreactivity in the rat hippocampal complex following pilocarpine-induced status epilepticus. *J. Comp. Neurol.* 510, 463–474. doi: 10.1002/cne.21767
- Kim, J. E., Kwak, S. E., and Kang, T. C. (2009). Upregulated TWIK-related acid-sensitive K⁺ channel-2 in neurons and perivascular astrocytes in the hippocampus of experimental temporal lobe epilepsy. *Epilepsia* 50, 654–663. doi: 10.1111/j.1528-1167.2008.01957.x
- Kim, K. S., Kobayashi, M., Takamatsu, K., and Tzingounis, A. V. (2012). Hippocampal calcin and KCNQ channels contribute to the kinetics of the slow afterhyperpolarization. *Biophys. J.* 103, 2446–2454. doi: 10.1016/j.bpj.2012.11.002
- King, D., Bronen, R. A., Spencer, D. D., and Spencer, S. S. (1997). Topographic distribution of seizure onset and hippocampal atrophy: relationship between MRI and depth EEG. *Electroencephalogr. Clin. Neurophysiol.* 103, 692–697. doi: 10.1016/S0013-4694(97)00090-4
- Kirchheim, F., Tinnes, S., Haas, C. A., Stegen, M., and Wolfart, J. (2013). Regulation of action potential delays via voltage-gated potassium Kv1.1 channels in dentate granule cells during hippocampal epilepsy. *Front. Cell. Neurosci.* 7:248. doi: 10.3389/fncel.2013.00248
- Knuesel, I., Zuellig, R. A., Schaub, M. C., and Fritschy, J. M. (2001). Alterations in dystrophin and utrophin expression parallel the reorganization of GABAergic synapses in a mouse model of temporal lobe epilepsy. *Eur. J. Neurosci.* 13, 1113–1124. doi: 10.1046/j.0953-816x.2001.01476.x
- Kobayashi, M., Wen, X., and Buckmaster, P. S. (2003). Reduced inhibition and increased output of layer II neurons in the medial entorhinal cortex in a model of temporal lobe epilepsy. *J. Neurosci.* 23, 8471–8479.
- Kole, M. H., Brauer, A. U., and Stuart, G. J. (2007). Inherited cortical HCN1 channel loss amplifies dendritic calcium electrogenesis and burst firing in a rat absence epilepsy model. *J. Physiol.* 578, 507–525. doi: 10.1113/jphysiol.2006.122028
- Kowalski, J., Geuting, M., Paul, S., Dieni, S., Laurens, J., Zhao, S., et al. (2010). Proper layering is important for precisely timed activation of hippocampal mossy cells. *Cereb. Cortex* 20, 2043–2054. doi: 10.1093/cercor/bhp267
- Krook-Magnuson, E., Armstrong, C., Bui, A., Lew, S., Ojiala, M., and Soltesz, I. (2015). *In vivo* evaluation of the dentate gate theory in epilepsy. *J. Physiol.* doi: 10.1113/JP270056
- Kubo, Y., Adelman, J. P., Clapham, D. E., Jan, L. Y., Karschin, A., Kurachi, Y., et al. (2005). International Union of Pharmacology. LIV. Nomenclature and molecular relationships of inwardly rectifying potassium channels. *Pharmacol. Rev.* 57, 509–526. doi: 10.1124/pr.57.4.11
- Kullmann, D. M., and Waxman, S. G. (2010). Neurological channelopathies: new insights into disease mechanisms and ion channel function. *J. Physiol.* 588, 1823–1827. doi: 10.1113/jphysiol.2010.190652
- Kumar, S. S., and Buckmaster, P. S. (2006). Hyperexcitability, interneurons, and loss of GABAergic synapses in entorhinal cortex in a model of temporal lobe epilepsy. *J. Neurosci.* 26, 4613–4623. doi: 10.1523/JNEUROSCI.0064-06.2006
- Lachance-Touchette, P., Brown, P., Meloche, C., Kinirons, P., Lapointe, L., Lacasse, H., et al. (2011). Novel alpha1 and gamma2 GABAA receptor subunit mutations in families with idiopathic generalized epilepsy. *Eur. J. Neurosci.* 34, 237–249. doi: 10.1111/j.1460-9568.2011.07767.x
- Lau, D., Vega-Saenz De Miera, E. C., Contreras, D., Ozaita, A., Harvey, M., Chow, A., et al. (2000). Impaired fast-spiking, suppressed cortical inhibition, and

- increased susceptibility to seizures in mice lacking Kv3.2 K⁺ channel proteins. *J. Neurosci.* 20, 9071–9085.
- Le Duigou, C., Bouillere, V., and Miles, R. (2008). Epileptiform activities in slices of hippocampus from mice after intra-hippocampal injection of kainic acid. *J. Physiol.* 586, 4891–4904. doi: 10.1113/jphysiol.2008.156281
- Lee, C. H., and Liou, H. H. (2014). Pregabalin activates ROMK1 channels via cAMP-dependent protein kinase and protein kinase C. *Eur. J. Pharmacol.* 740, 35–44. doi: 10.1016/j.ejphar.2014.06.049
- Lee, S. M., Kim, J. E., Sohn, J. H., Choi, H. C., Lee, J. S., Kim, S. H., et al. (2009). Down-regulation of delayed rectifier K⁺ channels in the hippocampus of seizure sensitive gerbils. *Brain Res. Bull.* 80, 433–442. doi: 10.1016/j.brainresbull.2009.07.016
- Levitan, I. B. (1994). Modulation of ion channels by protein phosphorylation and dephosphorylation. *Annu. Rev. Physiol.* 56, 193–212. doi: 10.1146/annurev.ph.56.030194.001205
- Li, K. X., Lu, Y. M., Xu, Z. H., Zhang, J., Zhu, J. M., Zhang, J. M., et al. (2012). Neuregulin 1 regulates excitability of fast-spiking neurons through Kv1.1 and acts in epilepsy. *Nat. Neurosci.* 15, 267–273. doi: 10.1038/nn.3006
- Limbrick, D. D. Jr., Sombati, S., and Delorenzo, R. J. (2003). Calcium influx constitutes the ionic basis for the maintenance of glutamate-induced extended neuronal depolarization associated with hippocampal neuronal death. *Cell Calcium* 33, 69–81. doi: 10.1016/S0143-4160(02)00054-4
- Liu, M., Pleasure, S. J., Collins, A. E., Noebels, J. L., Naya, F. J., Tsai, M. J., et al. (2000). Loss of BETA2/NeuroD leads to malformation of the dentate gyrus and epilepsy. *Proc. Natl. Acad. Sci. U.S.A.* 97, 865–870. doi: 10.1073/pnas.97.2.865
- Liu, Y., Liu, D., Printzenhoff, D., Coghlan, M. J., Harris, R., and Krafte, D. S. (2002). Tenidap, a novel anti-inflammatory agent, is an opener of the inwardly rectifying K⁺ channel hKir2.3. *Eur. J. Pharmacol.* 435, 153–160. doi: 10.1016/S0014-2999(01)01590-4
- Löscher, W., Klitgaard, H., Twyman, R. E., and Schmidt, D. (2013). New avenues for anti-epileptic drug discovery and development. *Nat. Rev. Drug Discov.* 12, 757–776. doi: 10.1038/nrd4126
- Lothman, E. W., Stringer, J. L., and Bertram, E. H. (1992). The dentate gyrus as a control point for seizures in the hippocampus and beyond. *Epilepsy Res. Suppl.* 7, 301–313.
- Loup, F., Wieser, H. G., Yonekawa, Y., Aguzzi, A., and Fritschy, J. M. (2000). Selective alterations in GABA_A receptor subtypes in human temporal lobe epilepsy. *J. Neurosci.* 20, 5401–5419.
- Lu, Q., Peevey, J., Jow, F., Monaghan, M. M., Mendoza, G., Zhang, H., et al. (2008). Disruption of Kv1.1 N-type inactivation by novel small molecule inhibitors (disinactivators). *Bioorg. Med. Chem.* 16, 3067–3075. doi: 10.1016/j.bmc.2007.12.031
- Ludwig, A., Budde, T., Stieber, J., Moosmang, S., Wahl, C., Holthoff, K., et al. (2003). Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. *EMBO J.* 22, 216–224. doi: 10.1093/emboj/cdg032
- Lugo, J. N., Barnwell, L. F., Ren, Y., Lee, W. L., Johnston, L. D., Kim, R., et al. (2008). Altered phosphorylation and localization of the A-type channel, Kv4.2 in status epilepticus. *J. Neurochem.* 106, 1929–1940. doi: 10.1111/j.1471-4159.2008.05508.x
- Lukasiuk, K., and Pitkanen, A. (2004). Large-scale analysis of gene expression in epilepsy research: is synthesis already possible? *Neurochem. Res.* 29, 1169–1178. doi: 10.1023/B:NERE.0000023604.91584.6c
- Ma, W., Berg, J., and Yellen, G. (2007). Ketogenic diet metabolites reduce firing in central neurons by opening K(ATP) channels. *J. Neurosci.* 27, 3618–3625. doi: 10.1523/JNEUROSCI.0132-07.2007
- Madeja, M., Margineanu, D. G., Gorji, A., Siep, E., Boerrigter, P., Klitgaard, H., et al. (2003). Reduction of voltage-operated potassium currents by levetiracetam: a novel antiepileptic mechanism of action? *Neuropharmacology* 45, 661–671. doi: 10.1016/S0028-3908(03)00248-X
- Magee, J. C. (1999). Dendritic Ih normalizes temporal summation in hippocampal CA1 neurons. *Nat. Neurosci.* 2, 508–514. doi: 10.1038/9158
- Magloczky, Z., and Freund, T. F. (1995). Delayed cell death in the contralateral hippocampus following kainate injection into the CA3 subfield. *Neuroscience* 66, 847–860. doi: 10.1016/0306-4522(94)00613-A
- Magloczky, Z., and Freund, T. F. (2005). Impaired and repaired inhibitory circuits in the epileptic human hippocampus. *Trends Neurosci.* 28, 334–340. doi: 10.1016/j.tins.2005.04.002
- Magloczky, Z., Halasz, P., Vajda, J., Czirkaj, S., and Freund, T. F. (1997). Loss of Calbindin-D28K immunoreactivity from dentate granule cells in human temporal lobe epilepsy. *Neuroscience* 76, 377–385. doi: 10.1016/S0306-4522(96)00440-X
- Main, M. J., Cryan, J. E., Dupere, J. R., Cox, B., Clare, J. J., and Burbidge, S. A. (2000). Modulation of KCNQ2/3 potassium channels by the novel anticonvulsant retigabine. *Mol. Pharmacol.* 58, 253–262.
- Marcelin, B., Chauviere, L., Becker, A., Migliore, M., Esclapez, M., and Bernard, C. (2009). h channel-dependent deficit of theta oscillation resonance and phase shift in temporal lobe epilepsy. *Neurobiol. Dis.* 33, 436–447. doi: 10.1016/j.nbd.2008.11.019
- Marder, E., and Goaillard, J. M. (2006). Variability, compensation and homeostasis in neuron and network function. *Nat. Rev. Neurosci.* 7, 563–574. doi: 10.1038/nrn1949
- Margerison, J. H., and Corsellis, J. A. (1966). Epilepsy and the temporal lobes. A clinical, electroencephalographic and neuropathological study of the brain in epilepsy, with particular reference to the temporal lobes. *Brain* 89, 499–530. doi: 10.1093/brain/89.3.499
- Martin, M. S., Dutt, K., Papale, L. A., Dube, C. M., Dutton, S. B., De Haan, G., et al. (2010). Altered function of the SCN1A voltage-gated sodium channel leads to gamma-aminobutyric acid-ergic (GABAergic) interneuron abnormalities. *J. Biol. Chem.* 285, 9823–9834. doi: 10.1074/jbc.M109.078568
- Mashimo, T., Ohmori, I., Ouchida, M., Ohno, Y., Tsurumi, T., Miki, T., et al. (2010). A missense mutation of the gene encoding voltage-dependent sodium channel (Nav1.1) confers susceptibility to febrile seizures in rats. *J. Neurosci.* 30, 5744–5753. doi: 10.1523/JNEUROSCI.3360-09.2010
- Mathern, G. W., Adelson, P. D., Cahan, L. D., and Leite, J. P. (2002). Hippocampal neuron damage in human epilepsy: Meyer's hypothesis revisited. *Prog. Brain Res.* 135, 237–251. doi: 10.1016/S0079-6123(02)35023-4
- Mattia, D., Nagao, T., Rogawski, M. A., and Avoli, M. (1994). Potassium channel activators counteract anoxic hyperexcitability but not 4-aminopyridine-induced epileptiform activity in the rat hippocampal slice. *Neuropharmacology* 33, 1515–1522. doi: 10.1016/0028-3908(94)90124-4
- Mcclelland, S., Flynn, C., Dube, C., Richichi, C., Zha, Q., Ghestem, A., et al. (2011). Neuron-restrictive silencer factor-mediated hyperpolarization-activated cyclic nucleotide gated channelopathy in experimental temporal lobe epilepsy. *Ann. Neurol.* 70, 454–464. doi: 10.1002/ana.22479
- Mcnamara, J. O. (1984). Kindling: an animal model of complex partial epilepsy. *Ann. Neurol.* 16(Suppl.), S72–S76. doi: 10.1002/ana.410160712
- Mehranfar, N., Gholamipour-Badie, H., Motamedi, F., Janahmadi, M., and Naderi, N. (2014a). The effect of paxilline on early alterations of electrophysiological properties of dentate gyrus granule cells in pilocarpine-treated rats. *Iran. J. Pharm. Res.* 13, 125–132. doi: 10.9734/ARRB/2014/11778
- Mehranfar, N., Gholamipour-Badie, H., Motamedi, F., Janahmadi, M., and Naderi, N. (2014b). Occurrence of two types of granule cells with different excitability in rat dentate gyrus granule cell layer following pilocarpine-induced status epilepticus. *Annu. Res. Rev. Biol.* 4, 3707–3715. doi: 10.9734/ARRB/2014/11778
- Meier, J., Semtner, M., Winkelmann, A., and Wolfart, J. (2014). Presynaptic mechanisms of neuronal plasticity and their role in epilepsy. *Front. Cell. Neurosci.* 8:164. doi: 10.3389/fncel.2014.00164
- Meier, J., Semtner, M., and Wolfart, J. (2015). “Homeostasis of neuronal excitability via synaptic and intrinsic inhibitory mechanisms,” in *Homeostatic Control of Brain Function*, eds. D. Boison and S. A. Masino (Oxford: Oxford University Press).
- Meldrum, B. S., and Rogawski, M. A. (2007). Molecular targets for antiepileptic drug development. *Neurotherapeutics* 4, 18–61. doi: 10.1016/j.nurt.2006.11.010
- Menteyne, A., Levavasseur, F., Audinat, E., and Avignone, E. (2009). Predominant functional expression of Kv1.3 by activated microglia of the hippocampus after status epilepticus. *PLoS One* 4:e6770. doi: 10.1371/journal.pone.0006770
- Misonou, H., Menegola, M., Mohapatra, D. P., Guy, L. K., Park, K. S., and Trimmer, J. S. (2006). Bidirectional activity-dependent regulation of neuronal ion channel phosphorylation. *J. Neurosci.* 26, 13505–13514. doi: 10.1523/JNEUROSCI.3970-06.2006
- Mody, I., Kohr, G., Otis, T. S., and Staley, K. J. (1992a). The electrophysiology of dentate gyrus granule cells in whole-cell recordings. *Epilepsy Res. Suppl.* 7, 159–168.

- Mody, I., Otis, T. S., Staley, K. J., and Kohr, G. (1992b). The balance between excitation and inhibition in dentate granule cells and its role in epilepsy. *Epilepsy Res. Suppl.* 9, 331–339.
- Molnar, P., and Nadler, J. V. (1999). Mossy fiber-granule cell synapses in the normal and epileptic rat dentate gyrus studied with minimal laser photostimulation. *J. Neurophysiol.* 82, 1883–1894.
- Monaghan, M. M., Menegola, M., Vacher, H., Rhodes, K. J., and Trimmer, J. S. (2008). Altered expression and localization of hippocampal A-type potassium channel subunits in the pilocarpine-induced model of temporal lobe epilepsy. *Neuroscience* 156, 550–562. doi: 10.1016/j.neuroscience.2008.07.057
- Morgan, R. J., and Soltesz, I. (2008). Nonrandom connectivity of the epileptic dentate gyrus predicts a major role for neuronal hubs in seizures. *Proc. Natl. Acad. Sci. U.S.A.* 105, 6179–6184. doi: 10.1073/pnas.0801372105
- Morimoto, K., Fahnestock, M., and Racine, R. J. (2004). Kindling and status epilepticus models of epilepsy: rewiring the brain. *Prog. Neurobiol.* 73, 1–60. doi: 10.1016/j.pneurobio.2004.03.009
- Motti, D., Le Duigou, C., Eugene, E., Chemaly, N., Wittner, L., Lazarevic, D., et al. (2010). Gene expression analysis of the emergence of epileptiform activity after focal injection of kainic acid into mouse hippocampus. *Eur. J. Neurosci.* 32, 1364–1379. doi: 10.1111/j.1460-9568.2010.07403.x
- Mucha, M., Ooi, L., Linley, J. E., Mordaka, P., Dalle, C., Robertson, B., et al. (2010). Transcriptional control of KCNQ channel genes and the regulation of neuronal excitability. *J. Neurosci.* 30, 13235–13245. doi: 10.1523/JNEUROSCI.1981-10.2010
- Mueller, S. G., Laxer, K. D., Schuff, N., and Weiner, M. W. (2007). Voxel-based T2 relaxation rate measurements in temporal lobe epilepsy (TLE) with and without mesial temporal sclerosis. *Epilepsia* 48, 220–228. doi: 10.1111/j.1528-1167.2006.00916.x
- Munoz, A., Arellano, J. I., and Defelipe, J. (2002). GABABR1 receptor protein expression in human mesial temporal cortex: changes in temporal lobe epilepsy. *J. Comp. Neurol.* 449, 166–179. doi: 10.1002/cne.10287
- Nadler, J. V., Perry, B. W., and Cotman, C. W. (1978). Intraventricular kainic acid preferentially destroys hippocampal pyramidal cells. *Nature* 271, 676–677. doi: 10.1038/271676a0
- Nagao, Y., Harada, Y., Mukai, T., Shimizu, S., Okuda, A., Fujimoto, M., et al. (2013). Expressional analysis of the astrocytic Kir4.1 channel in a pilocarpine-induced temporal lobe epilepsy model. *Front. Cell. Neurosci.* 7:104. doi: 10.3389/fncel.2013.00104
- Nassirpour, R., Bahima, L., Lalive, A. L., Lüscher, C., Lujan, R., and Slesinger, P. A. (2010). Morphine- and CaMKII-dependent enhancement of GIRK channel signaling in hippocampal neurons. *J. Neurosci.* 30, 13419–13430. doi: 10.1523/JNEUROSCI.2966-10.2010
- Noam, Y., Bernard, C., and Baram, T. Z. (2011). Towards an integrated view of HCN channel role in epilepsy. *Curr. Opin. Neurobiol.* 21, 873–879. doi: 10.1016/j.conb.2011.06.013
- O'Malley, D., and Harvey, J. (2007). MAPK-dependent actin cytoskeletal reorganization underlies BK channel activation by insulin. *Eur. J. Neurosci.* 25, 673–682. doi: 10.1111/j.1460-9568.2007.05347.x
- Ogiwara, I., Miyamoto, H., Morita, N., Atapour, N., Mazaki, E., Inoue, I., et al. (2007). Nav1.1 localizes to axons of parvalbumin-positive inhibitory interneurons: a circuit basis for epileptic seizures in mice carrying a Scn1a gene mutation. *J. Neurosci.* 27, 5903–5914. doi: 10.1523/JNEUROSCI.5270-06.2007
- Oh, Y. J., Na, J., Jeong, J. H., Park, D. K., Park, K. H., Ko, J. S., et al. (2012). Alterations in hyperpolarization-activated cyclic nucleotide-gated cation channel (HCN) expression in the hippocampus following pilocarpine-induced status epilepticus. *BMB Rep.* 45, 635–640. doi: 10.5483/BMBRep.2012.45.11.091
- Okamoto, O. K., Janjoppi, L., Bonone, F. M., Pansani, A. P., Da Silva, A. V., Scorza, F. A., et al. (2010). Whole transcriptome analysis of the hippocampus: toward a molecular portrait of epileptogenesis. *BMC Genomics* 11:230. doi: 10.1186/1471-2164-11-230
- Okazaki, M. M., Molnar, P., and Nadler, J. V. (1999). Recurrent mossy fiber pathway in rat dentate gyrus: synaptic currents evoked in presence and absence of seizure-induced growth. *J. Neurophysiol.* 81, 1645–1660.
- Oliveira, M. S., Skinner, F., Arshadmansab, M. F., Garcia, I., Mello, C. F., Knaus, H. G., et al. (2010). Altered expression and function of small-conductance (SK) Ca(2+)-activated K+ channels in pilocarpine-treated epileptic rats. *Brain Res.* 1348, 187–199. doi: 10.1016/j.brainres.2010.05.095
- Olney, J. W., and Sharpe, L. G. (1969). Brain lesions in an infant rhesus monkey treated with monosodium glutamate. *Science* 166, 386–388. doi: 10.1126/science.166.3903.386
- Otto, J. F., Kimball, M. M., and Wilcox, K. S. (2002). Effects of the anticonvulsant retigabine on cultured cortical neurons: changes in electroresponsive properties and synaptic transmission. *Mol. Pharmacol.* 61, 921–927. doi: 10.1124/mol.61.4.921
- Otto, J. F., Yang, Y., Frankel, W. N., White, H. S., and Wilcox, K. S. (2006). A spontaneous mutation involving Kcnq2 (Kv7.2) reduces M-current density and spike frequency adaptation in mouse CA1 neurons. *J. Neurosci.* 26, 2053–2059. doi: 10.1523/JNEUROSCI.1575-05.2006
- Pacheco Otorlora, L. F., Hernandez, E. F., Arshadmansab, M. F., Francisco, S., Willis, M., Ermolinsky, B., et al. (2008). Down-regulation of BK channel expression in the pilocarpine model of temporal lobe epilepsy. *Brain Res.* 1200, 116–131. doi: 10.1016/j.brainres.2008.01.017
- Palma, E., Amici, M., Sobrero, F., Spinelli, G., Di Angelantonio, S., Ragozzino, D., et al. (2006). Anomalous levels of Cl⁻ transporters in the hippocampal subiculum from temporal lobe epilepsy patients make GABA excitatory. *Proc. Natl. Acad. Sci. U.S.A.* 103, 8465–8468. doi: 10.1073/pnas.0602979103
- Pathak, H. R., Weissinger, F., Terunuma, M., Carlson, G. C., Hsu, F. C., Moss, S. J., et al. (2007). Disrupted dentate granule cell chloride regulation enhances synaptic excitability during development of temporal lobe epilepsy. *J. Neurosci.* 27, 14012–14022. doi: 10.1523/JNEUROSCI.4390-07.2007
- Pei, Q., Burnet, P. W., Grahame-Smith, D. G., and Zetterstrom, T. S. (1997). Differential effects of acute and chronic electroconvulsive shock on the abundance of messenger RNAs for voltage-dependent potassium channel subunits in the rat brain. *Neuroscience* 78, 343–350. doi: 10.1016/S0306-4522(96)00574-X
- Peng, B. W., Justice, J. A., Zhang, K., He, X. H., and Sanchez, R. M. (2010). Increased basal synaptic inhibition of hippocampal area CA1 pyramidal neurons by an antiepileptic drug that enhances I(H). *Neuropsychopharmacology* 35, 464–472. doi: 10.1038/npp.2009.150
- Peng, Z., Hauer, B., Mihalek, R. M., Homanics, G. E., Sieghart, W., Olsen, R. W., et al. (2002). GABA(A) receptor changes in delta subunit-deficient mice: altered expression of alpha4 and gamma2 subunits in the forebrain. *J. Comp. Neurol.* 446, 179–197. doi: 10.1002/cne.10210
- Peng, Z., Huang, C. S., Stell, B. M., Mody, I., and Houser, C. R. (2004). Altered expression of the delta subunit of the GABA(A) receptor in a mouse model of temporal lobe epilepsy. *J. Neurosci.* 24, 8629–8639. doi: 10.1523/JNEUROSCI.2877-04.2004
- Penschuck, S., Bastlund, J. F., Jensen, H. S., Stensbol, T. B., Egebjerg, J., and Watson, W. P. (2005). Changes in KCNQ2 immunoreactivity in the amygdala in two rat models of temporal lobe epilepsy. *Brain Res. Mol. Brain Res.* 141, 66–73. doi: 10.1016/j.molbrainres.2005.08.004
- Peters, H. C., Hu, H., Pongs, O., Storm, J. F., and Isbrandt, D. (2005). Conditional transgenic suppression of M channels in mouse brain reveals functions in neuronal excitability, resonance and behavior. *Nat. Neurosci.* 8, 51–60. doi: 10.1038/nn1375
- Pongs, O. (1992). Structural basis of voltage-gated K⁺ channel pharmacology. *Trends Pharmacol. Sci.* 13, 359–365. doi: 10.1016/0165-6147(92)90109-J
- Poolos, N. P. (2004). The Yin and Yang of the H-Channel and Its role in epilepsy. *Epilepsy Curr.* 4, 3–6. doi: 10.1111/j.1535-7597.2004.04101.x
- Poolos, N. P. (2009). Genetic loss of HCN1 channels is exciting, but is it epileptic? *Epilepsy Curr.* 10, 51–52. doi: 10.1111/j.1535-7511.2009.01352.x
- Poolos, N. P., and Johnston, D. (2012). Dendritic ion channelopathy in acquired epilepsy. *Epilepsia* 53 (Suppl. 9), 32–40. doi: 10.1111/epi.12033
- Poolos, N. P., Migliore, M., and Johnston, D. (2002). Pharmacological upregulation of h-channels reduces the excitability of pyramidal neuron dendrites. *Nat. Neurosci.* 5, 767–774. doi: 10.1038/nn891
- Pothmann, L., Muller, C., Averkin, R. G., Bellistri, E., Miklitz, C., Uebachs, M., et al. (2014). Function of inhibitory micronetworks is spared by Na⁺ channel-acting anticonvulsant drugs. *J. Neurosci.* 34, 9720–9735. doi: 10.1523/JNEUROSCI.2395-13.2014
- Powell, K. L., Lukasiuk, K., O'Brien, T. J., and Pitkanen, A. (2014). Are alterations in transmitter receptor and ion channel expression responsible for epilepsies? *Adv. Exp. Med. Biol.* 813, 211–229. doi: 10.1007/978-94-017-8914-1_17
- Powell, K. L., Ng, C., O'Brien, T. J., Xu, S. H., Williams, D. A., Foote, S. J., et al. (2008). Decreases in HCN mRNA expression in the hippocampus after

- kindling and status epilepticus in adult rats. *Epilepsia* 49, 1686–1695. doi: 10.1111/j.1528-1167.2008.01593.x
- Prakriya, M., and Mennerick, S. (2000). Selective depression of low-release probability excitatory synapses by sodium channel blockers. *Neuron* 26, 671–682. doi: 10.1016/S0896-6273(00)81203-9
- Ptacek, L. J. (1997). Channelopathies: ion channel disorders of muscle as a paradigm for paroxysmal disorders of the nervous system. *Neuromuscul. Disord.* 7, 250–255. doi: 10.1016/S0960-8966(97)00046-1
- Qiao, X., Werkman, T. R., Gorter, J. A., Wadman, W. J., and Van Vliet, E. A. (2013). Expression of sodium channel alpha subunits 1.1, 1.2 and 1.6 in rat hippocampus after kainic acid-induced epilepsy. *Epilepsy Res.* 106, 17–28. doi: 10.1016/j.eplepsyres.2013.06.006
- Qiu, C., Zeyda, T., Johnson, B., Hochgeschwender, U., De Lecea, L., and Tallent, M. K. (2008). Somatostatin receptor subtype 4 couples to the M-current to regulate seizures. *J. Neurosci.* 28, 3567–3576. doi: 10.1523/JNEUROSCI.4679-07.2008
- Raol, Y. H., Lapedes, D. A., Keating, J. G., Brooks-Kayal, A. R., and Cooper, E. C. (2009). A KCNQ channel opener for experimental neonatal seizures and status epilepticus. *Ann. Neurol.* 65, 326–336. doi: 10.1002/ana.21593
- Raol, Y. H., Lund, I. V., Bandyopadhyay, S., Zhang, G., Roberts, D. S., Wolfe, J. H., et al. (2006a). Enhancing GABA(A) receptor alpha 1 subunit levels in hippocampal dentate gyrus inhibits epilepsy development in an animal model of temporal lobe epilepsy. *J. Neurosci.* 26, 11342–11346. doi: 10.1523/JNEUROSCI.3329-06.2006
- Raol, Y. H., Zhang, G., Lund, I. V., Porter, B. E., Maronski, M. A., and Brooks-Kayal, A. R. (2006b). Increased GABA(A)-receptor alpha1-subunit expression in hippocampal dentate gyrus after early-life status epilepticus. *Epilepsia* 47, 1665–1673. doi: 10.1111/j.1528-1167.2006.00640.x
- Raza, M., Blair, R. E., Sombati, S., Carter, D. S., Deshpande, L. S., and Delorenzo, R. J. (2004). Evidence that injury-induced changes in hippocampal neuronal calcium dynamics during epileptogenesis cause acquired epilepsy. *Proc. Natl. Acad. Sci. U.S.A.* 101, 17522–17527. doi: 10.1073/pnas.0408155101
- Riazanski, V., Deriy, L. V., Shevchenko, P. D., Le, B., Gomez, E. A., and Nelson, D. J. (2011). Presynaptic CLC-3 determines quantal size of inhibitory transmission in the hippocampus. *Nat. Neurosci.* 14, 487–494. doi: 10.1038/nn.2775
- Rinke, I., Artmann, J., and Stein, V. (2010). ClC-2 voltage-gated channels constitute part of the background conductance and assist chloride extrusion. *J. Neurosci.* 30, 4776–4786. doi: 10.1523/JNEUROSCI.6299-09.2010
- Robbins, C. A., and Tempel, B. L. (2012). Kv1.1 and Kv1.2: similar channels, different seizure models. *Epilepsia* 53(Suppl. 1), 134–141. doi: 10.1111/j.1528-1167.2012.03484.x
- Roeloffs, R., Wickenden, A. D., Crean, C., Werness, S., McNaughton-Smith, G., Stables, J., et al. (2008). In vivo profile of ICA-27243 [N-(6-chloro-pyridin-3-yl)-3,4-difluoro-benzamide], a potent and selective KCNQ2/Q3 (Kv7.2/Kv7.3) activator in rodent anticonvulsant models. *J. Pharmacol. Exp. Ther.* 326, 818–828. doi: 10.1124/jpet.108.137794
- Rogawski, M. A., and Löscher, W. (2004). The neurobiology of antiepileptic drugs. *Nat. Rev. Neurosci.* 5, 553–564. doi: 10.1038/nrn1430
- Rossignol, E., Kruglikov, I., Van Den Maagdenberg, A. M., Rudy, B., and Fishell, G. (2013). CaV 2.1 ablation in cortical interneurons selectively impairs fast-spiking basket cells and causes generalized seizures. *Ann. Neurol.* 74, 209–222. doi: 10.1002/ana.23913
- Rostock, A., Tober, C., Rundfeldt, C., Bartsch, R., Engel, J., Polymeropoulos, E. E., et al. (1996). D-23129: a new anticonvulsant with a broad spectrum activity in animal models of epileptic seizures. *Epilepsy Res.* 23, 211–223. doi: 10.1016/0920-1211(95)00101-8
- Rundfeldt, C., and Netzer, R. (2000). The novel anticonvulsant retigabine activates M-currents in Chinese hamster ovary-cells transfected with human KCNQ2/3 subunits. *Neurosci. Lett.* 282, 73–76. doi: 10.1016/S0304-3940(00)00866-1
- Ruppersberg, J. P., Stocker, M., Pongs, O., Heinemann, S. H., Frank, R., and Koenen, M. (1991). Regulation of fast inactivation of cloned mammalian IK(A) channels by cysteine oxidation. *Nature* 352, 711–714. doi: 10.1038/352711a0
- Rüschenschmidt, C., Chen, J., Becker, A., Riazanski, V., and Beck, H. (2006). Functional properties and oxidative modulation of A-type K currents in hippocampal granule cells of control and chronically epileptic rats. *Eur. J. Neurosci.* 23, 675–685. doi: 10.1111/j.1460-9568.2006.04608.x
- Sanabria, E. R., Su, H., and Yaari, Y. (2001). Initiation of network bursts by Ca²⁺-dependent intrinsic bursting in the rat pilocarpine model of temporal lobe epilepsy. *J. Physiol.* 532, 205–216. doi: 10.1111/j.1469-7793.2001.0205g.x
- Santoro, B., Lee, J. Y., Englot, D. J., Gildersleeve, S., Piskorowski, R. A., Siegelbaum, S. A., et al. (2010). Increased seizure severity and seizure-related death in mice lacking HCN1 channels. *Epilepsia* 51, 1624–1627. doi: 10.1111/j.1528-1167.2010.02554.x
- Scharfman, H. E. (2002). Epilepsy as an example of neural plasticity. *Neuroscientist* 8, 154–173. doi: 10.1177/107385840200800211
- Scharfman, H. E., Goodman, J. H., and Sollas, A. L. (2000). Granule-like neurons at the hilar/CA3 border after status epilepticus and their synchrony with area CA3 pyramidal cells: functional implications of seizure-induced neurogenesis. *J. Neurosci.* 20, 6144–6158.
- Scharfman, H. E., Sollas, A. L., Berger, R. E., and Goodman, J. H. (2003). Electrophysiological evidence of monosynaptic excitatory transmission between granule cells after seizure-induced mossy fiber sprouting. *J. Neurophysiol.* 90, 2536–2547. doi: 10.1152/jn.00251.2003
- Schenzer, A., Friedrich, T., Pusch, M., Saftig, P., Jentsch, T. J., Grotzinger, J., et al. (2005). Molecular determinants of KCNQ (Kv7) K⁺ channel sensitivity to the anticonvulsant retigabine. *J. Neurosci.* 25, 5051–5060. doi: 10.1523/JNEUROSCI.0128-05.2005
- Schroeder, B. C., Kubisch, C., Stein, V., and Jentsch, T. J. (1998). Moderate loss of function of cyclic-AMP-modulated KCNQ2/KCNQ3 K⁺ channels causes epilepsy. *Nature* 396, 687–690. doi: 10.1038/25367
- Schulte, U., Thumfart, J. O., Klocker, N., Sailer, C. A., Bildl, W., Biniossek, M., et al. (2006). The epilepsy-linked Lgi1 protein assembles into presynaptic Kv1 channels and inhibits inactivation by Kvbeta1. *Neuron* 49, 697–706. doi: 10.1016/j.neuron.2006.01.033
- Schulz, R., Kirschstein, T., Brehme, H., Porath, K., Mikkat, U., and Köhling, R. (2012). Network excitability in a model of chronic temporal lobe epilepsy critically depends on SK channel-mediated AHP currents. *Neurobiol. Dis.* 45, 337–347. doi: 10.1016/j.nbd.2011.08.019
- Schwob, J. E., Fuller, T., Price, J. L., and Olney, J. W. (1980). Widespread patterns of neuronal damage following systemic or intracerebral injections of kainic acid: a histological study. *Neuroscience* 5, 991–1014. doi: 10.1016/0306-4522(80)90181-5
- Selke, K., Muller, A., Kukley, M., Schramm, J., and Dietrich, D. (2006). Firing pattern and calbindin-D28k content of human epileptic granule cells. *Brain Res.* 1120, 191–201. doi: 10.1016/j.brainres.2006.08.072
- Shah, M. M., Anderson, A. E., Leung, V., Lin, X., and Johnston, D. (2004). Seizure-induced plasticity of h channels in entorhinal cortical layer III pyramidal neurons. *Neuron* 44, 495–508. doi: 10.1016/j.neuron.2004.10.011
- Shah, M. M., Huang, Z., and Martinello, K. (2013). HCN and KV7 (M-) channels as targets for epilepsy treatment. *Neuropharmacology* 69, 75–81. doi: 10.1016/j.neuropharm.2012.03.005
- Shin, M., Brager, D., Jaramillo, T. C., Johnston, D., and Chetkovich, D. M. (2008). Mislocalization of h channel subunits underlies h channelopathy in temporal lobe epilepsy. *Neurobiol. Dis.* 32, 26–36. doi: 10.1016/j.nbd.2008.06.013
- Shruti, S., Clem, R. L., and Barth, A. L. (2008). A seizure-induced gain-of-function in BK channels is associated with elevated firing activity in neocortical pyramidal neurons. *Neurobiol. Dis.* 30, 323–330. doi: 10.1016/j.nbd.2008.02.002
- Sicca, F., Imbrici, P., D'adamo, M. C., Moro, F., Bonatti, F., Brovedani, P., et al. (2011). Autism with seizures and intellectual disability: possible causative role of gain-of-function of the inwardly-rectifying K⁺ channel Kir4.1. *Neurobiol. Dis.* 43, 239–247. doi: 10.1016/j.nbd.2011.03.016
- Signorini, S., Liao, Y. J., Duncan, S. A., Jan, L. Y., and Stoffel, M. (1997). Normal cerebellar development but susceptibility to seizures in mice labievcking G protein-coupled, inwardly rectifying K⁺ channel GIRK2. *Proc. Natl. Acad. Sci. U.S.A.* 94, 923–927. doi: 10.1073/pnas.94.3.923
- Sills, G. J. (2007). Seizures beget seizures: a lack of experimental evidence and clinical relevance fails to dampen enthusiasm. *Epilepsy Curr.* 7, 103–104. doi: 10.1111/j.1535-7511.2007.00189.x
- Singh, N. A., Charlier, C., Stauffer, D., Dupont, B. R., Leach, R. J., Melis, R., et al. (1998). A novel potassium channel gene, KCNQ2, is mutated in an inherited epilepsy of newborns. *Nat. Genet.* 18, 25–29. doi: 10.1038/ng0198-25
- Singh, N. A., Otto, J. F., Dahle, E. J., Pappas, C., Leslie, J. D., Vilaythong, A., et al. (2008). Mouse models of human KCNQ2 and KCNQ3 mutations for benign familial neonatal convulsions show seizures and neuronal plasticity without synaptic reorganization. *J. Physiol. (Lond)*. 586, 3405–3423. doi: 10.1113/jphysiol.2008.154971

- Sloviter, R. S. (1987). Decreased hippocampal inhibition and a selective loss of interneurons in experimental epilepsy. *Science* 235, 73–76. doi: 10.1126/science.2879352
- Sloviter, R. S. (1994). The functional organization of the hippocampal dentate gyrus and its relevance to the pathogenesis of temporal lobe epilepsy. *Ann. Neurol.* 35, 640–654. doi: 10.1002/ana.410350604
- Smart, S. L., Lopantsev, V., Zhang, C. L., Robbins, C. A., Wang, H., Chiu, S. Y., et al. (1998). Deletion of the K(V)1.1 potassium channel causes epilepsy in mice. *Neuron* 20, 809–819. doi: 10.1016/S0896-6273(00)81018-1
- Smith, S. E., Xu, L., Kasten, M. R., and Anderson, M. P. (2012). Mutant LGI1 inhibits seizure-induced trafficking of Kv4.2 potassium channels. *J. Neurochem.* 120, 611–621. doi: 10.1111/j.1471-4159.2011.07605.x
- Sosanya, N. M., Brager, D. H., Wolfe, S., Niere, F., and Raab-Graham, K. F. (2014). Rapamycin reveals an mTOR-independent repression of Kv1.1 expression during epileptogenesis. *Neurobiol. Dis.* 73C, 96–105. doi: 10.1016/j.nbd.2014.09.011
- Spencer, S. S. (2002). Neural networks in human epilepsy: evidence of and implications for treatment. *Epilepsia* 43, 219–227. doi: 10.1046/j.1528-1157.2002.26901.x
- Spencer, S. S., and Spencer, D. D. (1994). Entorhinal-hippocampal interactions in medial temporal lobe epilepsy. *Epilepsia* 35, 721–727. doi: 10.1111/j.1528-1157.1994.tb02502.x
- Spigelman, I., Li, Z., Banerjee, P. K., Mihalek, R. M., Homanics, G. E., and Olsen, R. W. (2002). Behavior and physiology of mice lacking the GABAA-receptor delta subunit. *Epilepsia* 43(Suppl. 5), 3–8. doi: 10.1046/j.1528-1157.43.s.5.8.x
- Splinter, M. Y. (2013). Efficacy of retigabine in adjunctive treatment of partial onset seizures in adults. *J. Cent. Nerv. Syst. Dis.* 5, 31–41. doi: 10.4137/JCNSD.S9299
- Stabel, J., Ficker, E., and Heinemann, U. (1992). Young CA1 pyramidal cells of rats, but not dentate gyrus granule cells, express a delayed inward rectifying current with properties of IQ. *Neurosci. Lett.* 135, 231–234. doi: 10.1016/0304-3940(92)90443-B
- Staley, K. J., and Mody, I. (1992). Shunting of excitatory input to dentate gyrus granule cells by a depolarizing GABAA receptor-mediated postsynaptic conductance. *J. Neurophysiol.* 68, 197–212.
- Stegen, M., Kirchheim, F., Hanuschkin, A., Staszewski, O., Veh, R. W., and Wolfart, J. (2012). Adaptive intrinsic plasticity in human dentate gyrus granule cells during temporal lobe epilepsy. *Cereb. Cortex* 22, 2087–2101. doi: 10.1093/cercor/bhr294
- Stegen, M., Young, C. C., Haas, C. A., Zentner, J., and Wolfart, J. (2009). Increased leak conductance in dentate gyrus granule cells of temporal lobe epilepsy patients with Ammon's horn sclerosis. *Epilepsia* 50, 646–653. doi: 10.1111/j.1528-1167.2009.02025.x
- Stogmann, E., Lichtner, P., Baumgartner, C., Schmied, M., Hotzy, C., Asmus, F., et al. (2006). Mutations in the CLCN2 gene are a rare cause of idiopathic generalized epilepsy syndromes. *Neurogenetics* 7, 265–268. doi: 10.1007/s10048-006-0057-x
- Strauss, U., Kole, M. H., Brauer, A. U., Pahnke, J., Bajorat, R., Rolfs, A., et al. (2004). An impaired neocortical Ih is associated with enhanced excitability and absence epilepsy. *Eur. J. Neurosci.* 19, 3048–3058. doi: 10.1111/j.0953-816X.2004.03392.x
- Streit, A. K., Derst, C., Wegner, S., Heinemann, U., Zahn, R. K., and Decher, N. (2011). RNA editing of Kv1.1 channels may account for reduced ictogenic potential of 4-aminopyridine in chronic epileptic rats. *Epilepsia* 52, 645–648. doi: 10.1111/j.1528-1167.2011.02986.x
- Su, H., Sochivko, D., Becker, A., Chen, J., Jiang, Y., Yaari, Y., et al. (2002). Upregulation of a T-type Ca²⁺ channel causes a long-lasting modification of neuronal firing mode after status epilepticus. *J. Neurosci.* 22, 3645–3655.
- Su, T., Cong, W. D., Long, Y. S., Luo, A. H., Sun, W. W., Deng, W. Y., et al. (2008). Altered expression of voltage-gated potassium channel 4.2 and voltage-gated potassium channel 4-interacting protein, and changes in intracellular calcium levels following lithium-pilocarpine-induced status epilepticus. *Neuroscience* 157, 566–576. doi: 10.1016/j.neuroscience.2008.09.027
- Sun, C., Mchedlishvili, Z., Erisir, A., and Kapur, J. (2007). Diminished neurosteroid sensitivity of synaptic inhibition and altered location of the alpha4 subunit of GABA(A) receptors in an animal model of epilepsy. *J. Neurosci.* 27, 12641–12650. doi: 10.1523/JNEUROSCI.4141-07.2007
- Sun, Y., Wu, Z., Kong, S., Jiang, D., Pitre, A., Wang, Y., et al. (2013). Regulation of epileptiform activity by two distinct subtypes of extrasynaptic GABAA receptors. *Mol. Brain* 6:21. doi: 10.1186/1756-6606-6-21
- Surges, R., Freiman, T. M., and Feuerstein, T. J. (2003). Gabapentin increases the hyperpolarization-activated cation current Ih in rat CA1 pyramidal cells. *Epilepsia* 44, 150–156. doi: 10.1046/j.1528-1157.2003.36802.x
- Sutula, T., Cascino, G., Cavazos, J., Parada, I., and Ramirez, L. (1989). Mossy fiber synaptic reorganization in the epileptic human temporal lobe. *Ann. Neurol.* 26, 321–330. doi: 10.1002/ana.410260303
- Sutula, T. P., Hagen, J., and Pitkanen, A. (2003). Do epileptic seizures damage the brain? *Curr. Opin. Neurol.* 16, 189–195. doi: 10.1097/01.wco.0000063770.15877.bc
- Suzuki, F., Heinrich, C., Boehrer, A., Mitsuya, K., Kurokawa, K., Matsuda, M., et al. (2005). Glutamate receptor antagonists and benzodiazepine inhibit the progression of granule cell dispersion in a mouse model of mesial temporal lobe epilepsy. *Epilepsia* 46, 193–202. doi: 10.1111/j.0013-9580.2005.35504.x
- Suzuki, F., Junier, M. P., Guilhem, D., Sorensen, J. C., and Onteniente, B. (1995). Morphogenetic effect of kainate on adult hippocampal neurons associated with a prolonged expression of brain-derived neurotrophic factor. *Neuroscience* 64, 665–674. doi: 10.1016/0306-4522(94)00463-F
- Sypert, G. W., and Ward, A. A. Jr. (1967). The hyperexcitable neuron: microelectrode studies of the chronic epileptic focus in the intact, awake monkey. *Exp. Neurol.* 19, 104–114. doi: 10.1016/0014-4886(67)90010-6
- Tatulian, L., Delmas, P., Abogadie, F. C., and Brown, D. A. (2001). Activation of expressed KCNQ potassium currents and native neuronal M-type potassium currents by the anti-convulsant drug retigabine. *J. Neurosci.* 21, 5535–5545.
- Tauk, D. L., and Nadler, J. V. (1985). Evidence of functional mossy fiber sprouting in hippocampal formation of kainic acid-treated rats. *J. Neurosci.* 5, 1016–1022.
- Taverna, S., Tkatch, T., Metz, A. E., and Martina, M. (2005). Differential expression of TASK channels between horizontal interneurons and pyramidal cells of rat hippocampus. *J. Neurosci.* 25, 9162–9170. doi: 10.1523/JNEUROSCI.2454-05.2005
- Tellez-Zenteno, J. F., and Hernandez-Ronquillo, L. (2012). A review of the epidemiology of temporal lobe epilepsy. *Epilepsy Res. Treat.* 2012:630853. doi: 10.1155/2012/630853
- Thom, M., Mathern, G. W., Cross, J. H., and Bertram, E. H. (2010). Mesial temporal lobe epilepsy: How do we improve surgical outcome? *Ann. Neurol.* 68, 424–434. doi: 10.1002/ana.22142
- Thom, M., Sisodiya, S. M., Beckett, A., Martinian, L., Lin, W. R., Harkness, W., et al. (2002). Cytoarchitectural abnormalities in hippocampal sclerosis. *J. Neuropathol. Exp. Neurol.* 61, 510–519.
- Tomlinson, S. E., Tan, S. V., Kullmann, D. M., Griggs, R. C., Burke, D., Hanna, M. G., et al. (2010). Nerve excitability studies characterize Kv1.1 fast potassium channel dysfunction in patients with episodic ataxia type 1. *Brain* 133, 3530–3540. doi: 10.1093/brain/awq318
- Tsaur, M. L., Sheng, M., Lowenstein, D. H., Jan, Y. N., and Jan, L. Y. (1992). Differential expression of K⁺ channel mRNAs in the rat brain and down-regulation in the hippocampus following seizures. *Neuron* 8, 1055–1067. doi: 10.1016/0896-6273(92)90127-Y
- Turrigiano, G., Lemasson, G., and Marder, E. (1995). Selective regulation of current densities underlies spontaneous changes in the activity of cultured neurons. *J. Neurosci.* 15, 3640–3652.
- Turrigiano, G. G., and Nelson, S. B. (2004). Homeostatic plasticity in the developing nervous system. *Nat. Rev. Neurosci.* 5, 97–107. doi: 10.1038/nrn1327
- Turski, L., Ikonomidou, C., Turski, W. A., Bortolotto, Z. A., and Cavalheiro, E. A. (1989). Review: cholinergic mechanisms and epileptogenesis. The seizures induced by pilocarpine: a novel experimental model of intractable epilepsy. *Synapse* 3, 154–171. doi: 10.1002/syn.890030207
- Turski, W. A., Cavalheiro, E. A., Schwarz, M., Czuczwar, S. J., Kleinrok, Z., and Turski, L. (1983). Limbic seizures produced by pilocarpine in rats: behavioural, electroencephalographic and neuropathological study. *Behav. Brain Res.* 9, 315–335. doi: 10.1016/0166-4328(83)90136-5
- Van Welie, I., Van Hooft, J. A., and Wadman, W. J. (2004). Homeostatic scaling of neuronal excitability by synaptic modulation of somatic hyperpolarization-activated Ih channels. *Proc. Natl. Acad. Sci. U.S.A.* 101, 5123–5128. doi: 10.1073/pnas.0307711101
- Van Welie, I., Van Hooft, J. A., and Wadman, W. J. (2006). Background activity regulates excitability of rat hippocampal CA1 pyramidal neurons by adaptation of a K⁺ conductance. *J. Neurophysiol.* 95, 2007–2012. doi: 10.1152/jn.00220.2005
- Varga, A. W., Yuan, L. L., Anderson, A. E., Schrader, L. A., Wu, G. Y., Gatchel, J. R., et al. (2004). Calcium-calmodulin-dependent kinase II modulates Kv4.2

- channel expression and upregulates neuronal A-type potassium currents. *J. Neurosci.* 24, 3643–3654. doi: 10.1523/JNEUROSCI.0154-04.2004
- Vida, I. (2009). 'Leaky' neurons in the epileptic hippocampus: should we get excited? *J. Physiol.* 587, 4127–4128. doi: 10.1113/jphysiol.2009
- Vida, I., Bartos, M., and Jonas, P. (2006). Shunting inhibition improves robustness of gamma oscillations in hippocampal interneuron networks by homogenizing firing rates. *Neuron* 49, 107–117. doi: 10.1016/j.neuron.2005.11.036
- Walleigh, D. J., Legido, A., and Valencia, I. (2013). Ring chromosome 20: a pediatric potassium channelopathy responsive to treatment with ezogabine. *Pediatr. Neurol.* 49, 368–369. doi: 10.1016/j.pediatrneurol.2013.06.005
- Waxman, S. G. (2001). Transcriptional channelopathies: an emerging class of disorders. *Nat. Rev. Neurosci.* 2, 652–659. doi: 10.1038/35090026
- Weckhuysen, S., Ivanovic, V., Hendrickx, R., Van Coster, R., Hjalgrim, H., Moller, R. S., et al. (2013). Extending the KCNQ2 encephalopathy spectrum: clinical and neuroimaging findings in 17 patients. *Neurology* 81, 1697–1703. doi: 10.1212/01.wnl.0000435296.72400.a1
- Weï, A. D., Gutman, G. A., Aldrich, R., Chandy, K. G., Grissmer, S., and Wulff, H. (2005). International Union of Pharmacology. LII. Nomenclature and molecular relationships of calcium-activated potassium channels. *Pharmacol. Rev.* 57, 463–472. doi: 10.1124/pr.57.4.9
- Wellmer, J., Su, H., Beck, H., and Yaari, Y. (2002). Long-lasting modification of intrinsic discharge properties in subicular neurons following status epilepticus. *Eur. J. Neurosci.* 16, 259–266. doi: 10.1046/j.1460-9568.2002.02086.x
- Wenzel, H. J., Vacher, H., Clark, E., Trimmer, J. S., Lee, A. L., Sapolsky, R. M., et al. (2007). Structural consequences of Kcna1 gene deletion and transfer in the mouse hippocampus. *Epilepsia* 48, 2023–2046. doi: 10.1111/j.1528-1167.2007.01189.x
- Westenbroek, R. E., Bausch, S. B., Lin, R. C., Franck, J. E., Noebels, J. L., and Caterall, W. A. (1998). Upregulation of L-type Ca²⁺ channels in reactive astrocytes after brain injury, hypomyelination, and ischemia. *J. Neurosci.* 18, 2321–2334.
- Whitaker, W. R., Faull, R. L., Dragunow, M., Mee, E. W., Emson, P. C., and Clare, J. J. (2001). Changes in the mRNAs encoding voltage-gated sodium channel types II and III in human epileptic hippocampus. *Neuroscience* 106, 275–285. doi: 10.1016/S0306-4522(01)00212-3
- Wickenden, A. D. (2002). Potassium channels as anti-epileptic drug targets. *Neuropharmacology* 43, 1055–1060. doi: 10.1016/S0028-3908(02)00237-X
- Wickenden, A. D., Krajewski, J. L., London, B., Wagoner, P. K., Wilson, W. A., Clark, S., et al. (2008). N-(6-chloro-pyridin-3-yl)-3,4-difluoro-benzamide (ICA-27243): a novel, selective KCNQ2/Q3 potassium channel activator. *Mol. Pharmacol.* 73, 977–986. doi: 10.1124/mol.107.043216
- Wickenden, A. D., Yu, W., Zou, A., Jegla, T., and Wagoner, P. K. (2000). Retigabine, a novel anti-convulsant, enhances activation of KCNQ2/Q3 potassium channels. *Mol. Pharmacol.* 58, 591–600.
- Wierschke, S., Lehmann, T. N., Dehnicke, C., Horn, P., Nitsch, R., and Deisz, R. A. (2010). Hyperpolarization-activated cation currents in human epileptogenic neocortex. *Epilepsia* 51, 404–414. doi: 10.1111/j.1528-1167.2009.02275.x
- Wimmer, V. C., Reid, C. A., So, E. Y., Berkovic, S. F., and Petrou, S. (2010). Axon initial segment dysfunction in epilepsy. *J. Physiol.* 588, 1829–1840. doi: 10.1113/jphysiol.2010.188417
- Wolfart, J., Debay, D., Le Masson, G., Destexhe, A., and Bal, T. (2005). Synaptic background activity controls spike transfer from thalamus to cortex. *Nat. Neurosci.* 8, 1760–1767. doi: 10.1038/nn1591
- Wolfart, J., and Roeper, J. (2002). Selective coupling of T-type calcium channels to SK potassium channels prevents intrinsic bursting in dopaminergic midbrain neurons. *J. Neurosci.* 22, 3404–3413.
- Wykes, R. C., Heeroma, J. H., Mantoan, L., Zheng, K., Macdonald, D. C., Deisseroth, K., et al. (2012). Optogenetic and potassium channel gene therapy in a rodent model of focal neocortical epilepsy. *Sci. Transl. Med.* 4, 161–152. doi: 10.1126/scitranslmed.3004190
- Wyler, A. R., Dohan, F. C., Schweitzer, J. B., and Berry, A. D. (1992). A grading system for hippocampal sclerosis. *J. Epilepsy* 5, 220–225. doi: 10.1016/S0896-6974(05)80120-3
- Xu, J. H., Long, L., Tang, Y. C., Hu, H. T., and Tang, F. R. (2007). Ca(v)1.2, Ca(v)1.3, and Ca(v)2.1 in the mouse hippocampus during and after pilocarpine-induced status epilepticus. *Hippocampus* 17, 235–251. doi: 10.1002/hipo.20263
- Xu, L., Hao, Y., Wu, X., Yu, P., Zhu, G., and Hong, Z. (2013a). Tenidap, an agonist of the inwardly rectifying K⁺ channel Kir2.3, delays the onset of cortical epileptiform activity in a model of chronic temporal lobe epilepsy. *Neurol. Res.* 35, 561–567. doi: 10.1179/1743132813Y.0000000157
- Xu, X., Guo, F., Lv, X., Feng, R., Min, D., Ma, L., et al. (2013b). Abnormal changes in voltage-gated sodium channels Na(V)1.1, Na(V)1.2, Na(V)1.3, Na(V)1.6 and in calmodulin/calmodulin-dependent protein kinase II, within the brains of spontaneously epileptic rats and tremor rats. *Brain Res. Bull.* 96, 1–9. doi: 10.1016/j.brainresbull.2013.04.003
- Yaari, Y., and Beck, H. (2002). "Epileptic neurons" in temporal lobe epilepsy. *Brain Pathol.* 12, 234–239. doi: 10.1111/j.1750-3639.2002.tb00438.x
- Yamada, K., Ji, J. J., Yuan, H., Miki, T., Sato, S., Horimoto, N., et al. (2001). Protective role of ATP-sensitive potassium channels in hypoxia-induced generalized seizure. *Science* 292, 1543–1546. doi: 10.1126/science.1059829
- Yim, M. Y., Hanuschkin, A., and Wolfart, J. (2015). Intrinsic rescaling of granule cells restores pattern separation ability of a dentate gyrus network model during epileptic hyperexcitability. *Hippocampus* 25, 297–308. doi: 10.1002/hipo.22373
- Young, C. C., Stegen, M., Bernard, R., Muller, M., Bischofberger, J., Veh, R. W., et al. (2009). Upregulation of inward rectifier K⁺ (Kir2) channels in dentate gyrus granule cells in temporal lobe epilepsy. *J. Physiol.* 587, 4213–4233. doi: 10.1113/jphysiol.2009.170746
- Yu, J., Proddutur, A., Elgammal, F. S., Ito, T., and Santhakumar, V. (2013). Status epilepticus enhances tonic GABA currents and depolarizes GABA reversal potential in dentate fast-spiking basket cells. *J. Neurophysiol.* 109, 1746–1763. doi: 10.1152/jn.00891.2012
- Yus-Najera, E., Munoz, A., Salvador, N., Jensen, B. S., Rasmussen, H. B., Defelipe, J., et al. (2003). Localization of KCNQ5 in the normal and epileptic human temporal neocortex and hippocampal formation. *Neuroscience* 120, 353–364. doi: 10.1016/S0306-4522(03)00321-X
- Zha, Q., Brewster, A. L., Richichi, C., Bender, R. A., and Baram, T. Z. (2008). Activity-dependent heteromerization of the hyperpolarization-activated, cyclic-nucleotide gated (HCN) channels: role of N-linked glycosylation. *J. Neurochem.* 105, 68–77. doi: 10.1111/j.1471-4159.2007.05110.x
- Zhang, G., Raol, Y. H., Hsu, F. C., Coulter, D. A., and Brooks-Kayal, A. R. (2004). Effects of status epilepticus on hippocampal GABA_A receptors are age-dependent. *Neuroscience* 125, 299–303. doi: 10.1016/j.neuroscience.2004.01.040
- Zhang, K., Peng, B. W., and Sanchez, R. M. (2006). Decreased IH in hippocampal area CA1 pyramidal neurons after perinatal seizure-inducing hypoxia. *Epilepsia* 47, 1023–1028. doi: 10.1111/j.1528-1167.2006.00574.x
- Zhang, N., Wei, W., Mody, I., and Houser, C. R. (2007). Altered localization of GABA(A) receptor subunits on dentate granule cell dendrites influences tonic and phasic inhibition in a mouse model of epilepsy. *J. Neurosci.* 27, 7520–7531. doi: 10.1523/JNEUROSCI.1555-07.2007
- Zhou, P., Yu, H., Gu, M., Nan, F. J., Gao, Z., and Li, M. (2013). Phosphatidylinositol 4,5-bisphosphate alters pharmacological selectivity for epilepsy-causing KCNQ potassium channels. *Proc. Natl. Acad. Sci. U.S.A.* 110, 8726–8731. doi: 10.1073/pnas.1302167110
- Ziemann, A. E., Schnizler, M. K., Albert, G. W., Severson, M. A., Howard, M. A. 3rd, Welsh, M. J., et al. (2008). Seizure termination by acidosis depends on ASIC1a. *Nat. Neurosci.* 11, 816–822. doi: 10.1038/nn.2132
- Zuberi, S. M., Eunson, L. H., Spauschus, A., De Silva, R., Tolmie, J., Wood, N. W., et al. (1999). A novel mutation in the human voltage-gated potassium channel gene (Kv1.1) associates with episodic ataxia type 1 and sometimes with partial epilepsy. *Brain* 122(Pt 5), 817–825.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Wolfart and Laker. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.