



# 1 Hydroclimatic processes as the primary drivers of the Early 2 Khvalynian transgression of the Caspian Sea: new developments

3  
4 Alexander Gelfan<sup>1,2</sup>, Andrey Panin<sup>1,3</sup>, Andrey Kalugin<sup>1</sup>, Polina Morozova<sup>3</sup>,  
5 Vladimir Semenov<sup>1,3,4</sup>, Alexey Sidorchuk<sup>2</sup>, Vadim Ukraintsev<sup>1,3</sup>, Konstantin Ushakov<sup>1,5</sup>

6 <sup>1</sup> Water Problems Institute, Russian Academy of Sciences, Moscow, 119333, Russia

7 <sup>2</sup> Lomonosov Moscow State University, Faculty of Geography, Moscow, 119991, Russia

8 <sup>3</sup> Institute of Geography, Russian Academy of Sciences, Moscow, 119017, Russia

9 <sup>4</sup> Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences, 119017 Moscow, Russia

10 <sup>5</sup> Shirshov Institute of Oceanology, Russian Academy of Sciences, 117997 Moscow, Russia

11 *Correspondence to:* Alexander Gelfan (hydrowpi@mail.ru)

12 **Abstract.** It has been well established that during the late Quaternary, the Khvalynian transgression of the Caspian  
13 Sea occurred, when the sea level rose tens of meters above the present one. Here, we evaluate the physical feasibility  
14 of the hypothesis that the maximum phase of this extraordinary event (known as the “Early Khvalynian transgression”)  
15 could be initiated and maintained for several thousand years solely by hydroclimatic factors. The hypothesis is based  
16 on recent studies dating the highest sea level stage (well above +10 m a.s.l.) to the final period of deglaciation, 17-13  
17 kyr BP, and studies estimating the contribution of the glacial waters in the sea level rise for this period as negligible.  
18 To evaluate the hypothesis put forward, we first applied the coupled ocean and sea-ice general circulation model  
19 driven by the climate model and estimated the equilibrium water inflow (irrespective of its origin) sufficient to  
20 maintain the sea level at the well-dated marks of the Early Khvalynian transgression as 400-470 km<sup>3</sup>/year. Secondly,  
21 we conducted an extensive 14C-dating of the large paleochannels (signs of high flow of atmospheric origin) located  
22 in the Volga basin and found that the period of their origin (17.5-14 ka BP) is almost identical to the recent dating of  
23 the main phase of the Early Khvalynian transgression. Water flow that could form these palaeochannels was earlier  
24 estimated for the ancient Volga River as 420 km<sup>3</sup>/year, i.e. close to the equilibrium runoff we determined. Thirdly, we  
25 applied a hydrological model forced by paleoclimate data to reveal physically consistent mechanisms of an  
26 extraordinarily high water inflow into the Caspian Sea in the absence of visible glacial meltwater effect. We found  
27 that the inflow could be caused by the spread of post-glacial permafrost in the Volga paleo-catchment. The numerical  
28 experiments demonstrated that the permafrost resulted in a sharp drop in infiltration into the frozen ground and reduced  
29 evaporation, which all together generated the Volga runoff during the Oldest Dryas, 17-14.8 kyr BP, up to 360  
30 km<sup>3</sup>/year (i.e. the total inflow into the Caspian Sea could reach 450 km<sup>3</sup>/year). The closeness of the estimates of river  
31 inflow into the sea, obtained by three independent methods, in combination with the previously obtained results, gave  
32 us reason to conclude that the hypothesis put forward is physically consistent.

## 33 1 Introduction

34 Paleogeographical data give grounds to assert that during the late Quaternary the largest highstand in the Quaternary  
35 history of the Caspian Sea took place, which was called the “Great” Khvalynian transgression. The boundaries of the  
36 Khvalynian Sea are well-detected in the relief of the Northern Caspian lowland (e.g. Leontiev, 1968, 1977; Rychagov,  
37 1974, 1997), and confirmed by stratigraphic and biostratigraphic analysis of Quaternary deposits (Fedorov, 1957,  
38 1978; Svitoch and Yanina 1997; Svitoch, 2009, 2014; Yanina 2012; Makshaev and Svitoch 2016; Yanina et al., 2018;



39 Kurbanov et al., 2021). The accumulated data show that in the early, maximum stage of the Khvalynian transgression,  
40 the sea level rose up to +48 m a.s.l., i.e. almost 80 meters above the current Caspian Sea level (CSL), while the sea  
41 surface area was 940,000 km<sup>2</sup>, which is 2.5 times larger than its current area (Yanko-Hombach and Kislov, 2018).

42 Although the very fact of the Early Khvalynian transgression and the assessment of the maximum sea level are not  
43 questioned by most researchers, there are significant disagreements regarding the dating of this extraordinary  
44 hydrological phenomenon and the views on its genesis.

45 In the 1970s-90s, it was assumed that the maximum phase of the Khvalynian transgression was synchronous to the  
46 Early Valdai (Early Weichselian, MIS 4) glaciation of the Russian Plain and occurred 50-70 ka BP (see reviews by  
47 Kislov et al., 2014; Arslanov et al., 2016 and references there). Accumulation of geochronometric, mostly radiocarbon  
48 (14C) data has allowed a reassessment of this viewpoint and proposal for a younger age of the Early Khvalynian  
49 transgression, corresponding to the second half of the last glaciation (Late Valdai, Late Weichselian, MIS 2) (Svitoch  
50 et al., 1994, 1998; Svitoch and Yanina, 1997). A number of compilations of the accumulated geochronological data  
51 have been published in recent years that enable a more detailed interpretation of the transgression. Arslanov et al.  
52 (2016) summarized the 14C and 230Th/234U dates of the Lower Khvalynian deposits performed at St. Petersburg  
53 University and proposed to date the +35 and +22 m a.s.l. transgressive stages at 16 and 14 ka BP, respectively, while  
54 the period 14-12 ka BP was attributed stages 0 and -12 m a.s.l. of the subsequent Late Khvalynian transgression.  
55 Krijgsman et al. (2019), based on a review of available dates, assigned the entire Khvalynian epoch to the 35-10 ka  
56 BP interval, with the Yenotaevian regression separating the Early and Late Khvalynian phases, about 15 ka BP.  
57 Koriche et al. (2022) attributed the Early Khvalynian stage to 35-25 ka BP and the Late Khvalynian stage to 17-12 ka  
58 BP. The latter, according to (Koriche et al., 2022), reached +35 m a.s.l. during 14.5-16.5 ka BP. Makshaev and Tkach  
59 (2023), based on generalization of more than 180 14C dates, attributed the Early Khvalynian stage of the Caspian Sea  
60 to the period 46-12.5 ka BP. In their opinion, sea level exceeded the contemporary level at the beginning of MIS 2  
61 (28-25 ka BP). This was followed by two transgressive events of 25-18 ka BP (level reached +10+15 m a.s.l.) and 17-  
62 13.5 ka BP (+20+22 m a.s.l.), separated by a regressive phase between 18 and 17 ka BP. These authors attribute the  
63 Yenotaevian regression and the subsequent Late Khvalynian transgression to 12.5-8.5 ka BP.

64 Recently, a series of papers have been published where sections containing the Khvalynian sediments were first dated  
65 by optically stimulated luminescence (OSL) (Kurbanov et al., 2021, 2022, 2023; Butuzova et al., 2022; Taratunina et  
66 al., 2022). These results were summarized in Kurbanov et al. (2023), who identified the following transgression stages:  
67 1) sea level rise to about +5 m a.s.l (32 m above the present CSL) between 30-35 and 27 ka BP; 2) sea level stabilization  
68 with a slight (about 2 m) rise within the interval of 27-20 ka BP; 3) a sharp rise in the sea level beginning from 18-17  
69 ka BP; 4) maximum stage of the sea level during the period around 16-15 ka BP; 5) rapid fall of the sea level during  
70 the period 15-14 ka BP from its maximum values to less than +11 m a.s.l.

71 Thus, the Khvalynian stage in the development of the Caspian Sea can currently be referred to the period from the end  
72 of MIS 3 (about 35 ka BP) to the Early Holocene (8.5 ka BP). At the beginning of that period, the sea level was lower  
73 than it is now, but no later than 27 ka BP it was already much higher. It should be emphasized that no direct dates for  
74 the maximum stage of +48+50 m a.s.l. have been obtained in any study. The recently published OSL data on the  
75 Raygorod section in the Northern Caspian Lowland at +13.5 m a.s.l. (Taratunina et al., 2022) show that from at least  
76 90 ka BP up to 18 ka BP, subaerial deposits (alluvium, loess) were accumulating there, i.e., the maximum phase of  
77 transgression could not have occurred before the Last Glacial Maximum (LGM). The age of the maximum stage is  
78 best justified by (Kurbanov et al., 2023), where the maximum stage is sandwiched between the rise and fall phases



79 and is assigned to the interval of 15-16 ka BP. Therefore, taking into account the reliable recent dating reviewed above,  
80 we will limit our attempt to explain the genesis of the Early Khvalynian transgression to the final period of  
81 deglaciation, (18)17-13 kyr BP.

82 Another widely debated question is: what are the causes of the Early Khvalynian transgression? The discussed  
83 hypotheses are reduced to the consideration of the sources of a huge water influx into the sea, which, under the climatic  
84 conditions of the Late Pleistocene, could provide the sea level rise of tens of meters above the present CSL. Other  
85 causes, such as tectonic factors or natural, internal fluctuations of the water body, are considered unlikely (Rychagov,  
86 1997; Yanko-Hombach and Kislov, 2018, respectively). According to paleoclimatic modeling experiments (e.g.  
87 Kislov and Toropov, 2007; Morozova, 2014; Yanko-Hombach and Kislov, 2018; Morozova et al., 2021), the LGM  
88 and post-LGM climate is characterized by low air temperatures and low precipitation with a reduced, relative to the  
89 modern, climatic runoff, that is, the difference between precipitation and evaporation in the catchment area of the  
90 Caspian Sea. To explain the increased river inflow into the Caspian Sea as a factor of the Early Khvalynian  
91 transgression, hypotheses are put forward about additional, in comparison with atmospheric precipitation, sources of  
92 water. The most discussed hypothesis is the recharge of glacial meltwater from the south-eastern flank of the  
93 Scandinavian ice sheet (SIS) via the Volga River during the LGM and deglaciation (Kvasov, 1979; Varuschenko et  
94 al., 1987; Toropov and Morozova, 2011; Tudryn et al., 2016; Koriche et al., 2022). Hypotheses are also put forward  
95 about the overflow of glacially dammed lakes and water discharge from outside the drainage basin of the Caspian Sea  
96 - from the upper Dnieper catchment and from the Sukhona and Vychegda Rivers that belong to the Arctic Ocean  
97 catchment (Kvasov, 1979; Larsen et al., 2006; Lyså et al., 2011), from the Aral Sea basin through a hypothetical  
98 hydrological system connecting it with both the ice-dammed lakes of the West Siberian ice-sheet and the Caspian Sea  
99 (see Grosswald and Kotlyakov, 1989; Chepalyga, 2007, as well as a critique of this hypothesis by Svitoch (2009) and  
100 Panin et al. (2020)). Kvasov (1979) estimated the contribution of the SIS meltwater and proglacial lakes as 46% and  
101 input from the Aral Sea as 21% of the total water inflow into the Early Khvalynian Caspian Sea, which was estimated  
102 by this author as 560 km<sup>3</sup>/year. Based on the PMIP2 (Paleoclimate Modelling Intercomparison Project, Phase 2)  
103 climate simulation data, Toropov and Morozova (2011) estimated that the SIS meltwater could have made the main  
104 contribution to the Khvalynian transgressions: 83% of the ancient Volga River inflow assessed as 462 km<sup>3</sup>/year. The  
105 coupled atmosphere-ocean-vegetation HadCM3 climate model experiments allowed Koriche et al. (2022) to conclude  
106 that meltwater combined with the changes (due to isostatic adjustment) in the drainage system leading to an increase  
107 in the Caspian Sea catchment area by 60-70% of its modern size, had the most substantial influence on the sea level  
108 rise during the last deglaciation from 20 kyr BP to 14 kyr BP. Note that all the above estimates of the SIS meltwater  
109 contribution were obtained solely from modelling results, which were not confirmed by geological and/or  
110 geomorphological evidence.

111 The validity of the above hypotheses considering glacial meltwater as a substantial source of water inflow into the  
112 Caspian Sea and confidence in the corresponding estimates of meltwater contribution to the Early Khvalynian  
113 transgression, are directly related to the assessed age of the transgression. According to the present-day state of  
114 geochronological studies described above, the stages well above +10 m a.s.l. are dated to the period of (18)17-13 kyr  
115 BP. Tudryn et al. (2016) proposed that glacial meltwater entered the Caspian Sea during the entire deglaciation epoch  
116 up to 13.8 kyr BP. However, Panin et al. (2021) showed that the inflow of meltwater into the Volga basin occurred  
117 only from its upper part directly covered by the Scandinavian ice-sheet, and was limited to a period from 21 to 16.5  
118 kyr BP, i.e. the transgression was developing towards its highest stage, while the input of glacial waters ceased. The  
119 authors estimated the possible glacial meltwater input to the upper Volga River in the range of 15-70 km<sup>3</sup>/year, or



120 only 5–25% of the present-day Volga runoff into the Caspian Sea, which is far from enough to support the Khvalynian  
121 highstand. The insignificant role of glacial meltwater in the genesis of the Early Khvalynian transgression during the  
122 deglaciation period is also argued in earlier works (Kalinin et al, 1966; Panin et al., 2005; Sidorchuk et al., 2009).  
123 Also, a number of recent studies (Panin et al., 2020, 2022; Borisova et al., 2022) showed that neither the proglacial  
124 lakes in the upper Volga region proposed by Kvasov (1979), nor the overflow to the Volga River from the Arctic basin  
125 occurred in MIS 2.

126 The hypothesis of hydroclimatic initiation of the Early Khvalynian transgression, in the absence of a noticeable  
127 contribution from glacial meltwater, is supported by the ubiquitous presence in the southern half of the Eastern  
128 European Plain, including the Volga basin, of signs of high flow of atmospheric origin - river palaeochannels that are  
129 many times greater in size than the contemporary rivers (Sidorchuk et al., 2009, 2011, 2021; Ukraintsev, 2022). On  
130 the basis of the developed morphometric analysis of palaeochannels, Sidorchuk et al. (2009, 2021) estimated the  
131 meteoric (formed due to atmospheric precipitation) runoff of the ancient Volga River, which was capable of forming  
132 the palaeochannels, as 420 km<sup>3</sup>/year, i.e. 65% higher than the modern annual runoff. At physically reasonable ratios  
133 of precipitation and evaporation in the Caspian Sea, this is quite sufficient to maintain levels of the Early Khvalynian  
134 transgression (Sidorchuk et al., 2009; Kislov et al., 2014).

135 The age of large palaeochannels in the Dnieper, Don, and Volga basins obtained by the 14C method falls within the  
136 interval of 18-13 kyr BP (Borisova et al., 2006; Sidorchuk et al., 2009; Panin et al., 2013, 2017; Panin and Matlakhova,  
137 2015), that is, exactly at the time when the CSL rose above +10 m a.s.l. However, it should be noted that in the Volga  
138 basin itself, only two large palaeochannels have been dated so far on the Moskva River, a tributary of the Oka River,  
139 and on the Samara River, a tributary of the lower Volga (Sidorchuk et al., 2009). This is insufficient for such a large  
140 basin encompassing several natural zones with significant differences in the present climate. In this study, we clarified  
141 the period of activity of large palaeochannels in the Volga basin.

142 Thus, according to the above review there is a knowledge gap, which drives the main motivation for our study. On  
143 the one hand, the well-founded modern datings show that in the final period of deglaciation, 18(17)-13 kyr BP, the  
144 CSL rose well above +10 m a.s.l. (likely, up to +22 ÷ +35 m a.s.l.), but, on the other hand, it has been proved that the  
145 meltwater runoff – due to the Scandinavian ice-sheet melting and outbursts of ice-dammed proglacial lakes - was  
146 either absent or contributed insignificantly to the transgression of the sea during this period. A research question arises:  
147 could the Early Khvalynian transgression of the Caspian Sea have been initiated and maintained solely by  
148 hydroclimatic factors in the cryoarid climate of the deglaciation period and in the absence of an inflow of glacial  
149 meltwater?

150 Kislov and Toropov (2007), Sidorchuk et al. (2009) hypothesized that during the decline in the glacier melt, river flow  
151 into the sea could significantly exceed the current one due to the spread of post-glacial permafrost in the river  
152 catchments of the East European Plain. Permafrost could reduce evaporation for the sea catchment territory owing to  
153 a drastic decrease in the infiltration capacity of frozen ground. Gelfan and Kalugin (2021) applied a physically based  
154 hydrological model to assess the sensitivity of the Volga River runoff to the hypothetical spread of permafrost in the  
155 river basin. The authors demonstrated that under the modern climatic conditions mean annual runoff may increase by  
156 85% due to modeled "freezing" of the basin. They concluded that river inflow into the Caspian Sea is markedly  
157 sensitive to presence of permafrost over the sea catchment area, thus further verification of the hypothesis is advisable  
158 in the cryoarid climatic conditions of the late Pleistocene. One of the objectives of our study is to verify this hypothesis  
159 explaining the maintenance of the CSL at +22 ÷ +35 m a.s.l. reliably dated to the period of 18(17)-13 kyr BP in the  
160 absence of significant glacial meltwater runoff during this period.



161 The logic of our study was as follows. Using a full ocean model coupled with a model of sea-ice dynamics INMIO  
162 COMPASS – CICE (Ibrayev et al., 2012; Hunke et al., 2015), we simulated the Caspian Sea water balance  
163 components under the climate conditions of the Late Pleistocene – Middle Holocene, which were re-constructed with  
164 the help of the climate model INMCM4.8 (Volodin et al., 2018). On the basis of the simulation data, we estimated the  
165 equilibrium river water inflow into the sea maintaining its level at the well-dated marks of the Early Khvalynian  
166 transgression. To verify the model-based estimations, the river runoff assessments derived from the morphometry of  
167 palaeochannels formed in the period 18-13 kyr BP (Sidorchuk et al., 2021) were used. Also, we made an attempt to  
168 improve the knowledge on the chronology of widespread geomorphological evidence of high river runoff in the Late  
169 Pleniglacial – Late Glacial in the Volga basin. To achieve this, additional dating of large palaeochannels in different  
170 parts of the basin was carried out. Then, the hydrological model was forced by the paleoclimate data, and numerical  
171 experiments were conducted to assess the water inflow to the Caspian Sea from the ancient Volga catchment with  
172 underlying permafrost. Comparison of estimates of water inflow into the Caspian Sea obtained using three independent  
173 approaches (1 – estimating equilibrium inflow into the sea via an ocean model coupled with a climate model; 2 -  
174 paleogeographic reconstructions of water flow through palaeochannels, and 3 – hydrological modeling river runoff  
175 generation in the sea catchment area under the paleoclimatic conditions) provided us with grounds for answering the  
176 above research question.

177 The remaining part of this paper is organized as follows. General information about the Caspian Sea is given in the  
178 next section. Section 3 contains methodology of our study including brief description of the models used and the  
179 numerical experiments designed. The results are presented and discussed in Section 4. The overall conclusions are  
180 given in Section 5.

## 181 **2. General information on the Caspian Sea**

182 The Caspian Sea (36°33'–47°07' N, 46°43'–54°50' E) is the world's largest inland water body located within an  
183 endorheic (no outflow) basin. The sea surface area at the current sea level is equal to 365,000 km<sup>2</sup>. The coastline  
184 length is 5970 km. The greatest length of the sea (along the meridian 50°00'E) is 1030 km. The greatest width along  
185 the parallel 45°30' N reaches 435 km. The large meridional extent results in climate variations over the basin: from  
186 sub-tropical in the southwest to desertic in the east and northeast.

187 Owing to the endorheic nature of the Caspian Sea, its level widely fluctuated in the past. During the late Cenozoic,  
188 the CSL variations exceeded, probably, several hundreds of meters (Forte and Cowgill 2013) and at least 100 m,  
189 during the last 500,000–700,000 years (Water balance..., 2016), during the Holocene the CSL changes were from 15  
190 m (Water balance..., 2016) to several tens of meters (Kakroodi et al. 2012), during the last millennium the CSL  
191 changed by 10 m (Naderi Beni et al. 2013) and during the period of instrumental observations (beginning from 1830)  
192 within the range of 4 m: from -25.1 m a.s.l. at the beginning of 1880s to -29.0 m a.s.l. in the middle of 1970s (Frolov,  
193 2003). The present (December of 2022) CSL is -28.6 m a.s.l.

194 The CSL variations are controlled mainly by water inflow from rivers and precipitation on the sea, as well as by water  
195 outflow through evaporation from the sea surface (Ratkovich, 1993; Golitsyn et al., 1998; Kroonenberg et al., 2000;  
196 Arpe and Leroy, 2007; Arpe et al., 2012; Naderi Beni et al., 2013; Panin and Dianskii, 2014; Chen et al., 2017), i.e.  
197 they are strongly dependent on climatic variations (Kroonenberg et al., 2000; Arpe and Leroy, 2007; ), at least as long  
198 as no significant changes are occurring in the sea catchment area. Groundwater inflow contribution is estimated to be  
199 small (Zektser, 1996) and expected to partly compensate for the impact from the outflow to the Kara-Bogaz-Gol Bay  
200 (Chen et al., 2017) accounting for the uncertainty of both estimates.



201 The Caspian Sea is fed by more than 130 large and small rivers with the total annual flow of about 300 km<sup>3</sup> (average  
202 value for 1880-2001 (Frolov, 2003)). The total catchment area of the sea is 3,050,000 km<sup>2</sup>, which is 8 times the area  
203 of its water area (386,400 km<sup>2</sup> at the sea level of -27.50 m a.s.l.). The largest of the tributaries is the Volga River,  
204 whose catchment area is 1,360,000 km<sup>2</sup>. For the period of instrumental observations (1881-2012), the mean annual  
205 flow of the Volga in the river outlet (Volgograd city) is about 250 km<sup>3</sup> (e.g. Arpe et al., 2019). Taking into account  
206 water losses due to evaporation in the Volga delta, the Volga water inflow into the Caspian Sea is about 233 km<sup>3</sup> of  
207 water per year (Frolov, 2003) or about 80% of the total inflow of river water into the sea. According to (Kislov and  
208 Toropov, 2007), the relative contribution of the Volga runoff has changed insignificantly over the past 20 thousand  
209 years and accounts for 75 to 90% of the total inflow into the Caspian Sea. According to various estimates, the long-  
210 term mean precipitation on the Caspian Sea surface in the 20th century was about 200 mm/year (about 77 km<sup>3</sup>/year),  
211 evaporation from the sea surface was 960 mm/year (about 371 km<sup>3</sup>/year), and effective evaporation (the difference  
212 between evaporation and precipitation) was 760 mm/year (about 294 km<sup>3</sup>/year), respectively (e.g. Frolov, 2003; Water  
213 Balance..., 2016).

214 The relationship between water input to and output from the Caspian Sea controls the sea level. The CSL response to  
215 changes in the main water balance components of the sea depends on the peculiarities of the sea bathymetry, namely,  
216 a significant fraction of shallow water areas. The northern part of the sea is shallow, in the southern and central parts  
217 of the sea there are deep depressions that are intersected by an underwater ridge. The average depth of the sea is 208  
218 m, the maximum depth is 1025 m. About 69% of the total sea area is at depths less than 200 meters, and a shallow  
219 zone with depths less than 10 m occupies 28% of the sea area. In the range of the CSL fluctuations from -28.0 to -  
220 24.0 m a.s.l., a one-meter change in the CSL results in a 1500 km<sup>2</sup> change in the area of the deep-water part of the sea,  
221 and a 12500 km<sup>2</sup> change in the area of the shallow-water North Caspian part (Frolov, 2021). The predominant increase  
222 in the water area due to the shallow waters of the Northern Caspian with a rise in the sea level creates a non-linear  
223 dependence of evaporation from sea level fluctuations (Frolov, 2003).

224

### 225 **3 Research Methods**

#### 226 **3.1 Hydro- thermodynamics model of the Caspian Sea**

227 To simulate the Caspian Sea water balance components, we used a regional configuration of the coupled ocean and  
228 sea-ice general circulation model INMIO COMPASS – CICE. This approach involves a detailed description of marine  
229 dynamic processes with a high spatiotemporal resolution, taking into account ice drift and energy-mass transfer in the  
230 water-ice-atmosphere system. Thus, it is possible to obtain more reasonable values of evaporation from the sea surface  
231 compared to global climate models, in which a coarser resolution is typically used and the sea level is set constant,  
232 allowing no change in the surface area when the water balance of the sea is different from zero. The importance of  
233 using a full ocean model for the Caspian Sea was demonstrated by (Arpe et al., 2019).

234 The coupled model built from INMIO COMPASS (Ibrayev et al., 2012) and CICE (Hunke et al., 2015) codes in the  
235 CMF2.0 software environment (Kalmykov et al., 2018) was used earlier for weather forecasting and climate research  
236 (Fadeev et al., 2018; Kalnitskii et al., 2020; Ushakov and Ibrayev, 2018, and references therein). The model solves  
237 the equations of three-dimensional dynamics and thermodynamics of the ocean and sea ice cover, explicitly  
238 reproducing a wide range of processes responsible for the main energy-carrying elements of the circulation. The






239 calculations were performed using a model configuration tuned for the Caspian Sea region with a spatial resolution of  
240 about 22 km and a time step of 20 minutes, which was described in (Morozova et al., 2021).

### 241 3.2 Assessing equilibrium river inflow into the paleo-Caspian Sea under the transgressive levels of the sea

242 To assess an equilibrium river inflow into the paleo-Caspian Sea, the paleo-climate data simulated by the INMCM4.8  
243 climate model (Volodin et al., 2018) were set as atmospheric boundary conditions for the coupled ocean-ice model  
244 according to the protocols of PMIP4 (Paleoclimate Modelling Intercomparison Project, Phase 4) and CMIP6 (Coupled  
245 Model Intercomparison Project, Phase 6). The paleo-climate data represent two periods: the Last Glacial Maximum  
246 (experiment LGM, 21 kyr BP, Kageyama et al., 2021) and the mid-Holocene (experiment midHolocene, 6 kyr BP,  
247 Brierley et al., 2020). The data included near-surface air temperature and specific humidity, precipitation, wind  
248 velocity vector, fluxes of incoming longwave and shortwave radiation. The time resolution of the boundary fields was  
249 6 hours, which made it possible to explicitly consider a wide range of variability, from synoptic to interannual scales.

250 Since the Caspian Sea in the experiments of the climate model was specified in the modern coastline, the isolines of  
251 some boundary fields (air temperature and humidity, incoming longwave radiation) showed a tendency to follow this  
252 coastline. For these fields, an extrapolation was made from the sea area domain adopted by the climate model to the  
253 area of transgression. Since the sea level rise affects mainly the northern coastal regions, the extrapolation was  
254 performed from south to north using the meridional gradients calculated for each field by the least square method over  
255 the central part of the climate model water area (Fig. 1a).



256 a) The Caspian Sea area representation in the climate model INMCM4.8 (blue cells), red shading - cells used to  
257 calculate meridional gradients, green and red cells - extrapolation areas for transgressive stages (green cells – meridional  
258 extrapolation, red cells – extrapolation by the nearest neighbor method); b) The model representation of the Caspian Sea  
259 coastline for the sea levels assigned in the numerical experiments. The grey fill shows modern boundaries of the sea.  
260  
261



262 Further, for several transgressive cells, where this meridional procedure is not applicable, a simple extrapolation by  
263 the nearest neighbor method was performed. Precipitation, wind velocity components, and incoming shortwave  
264 radiation were used directly without extrapolation.

265 Calculations of the water balance in the LGM and mid-Holocene were carried out for a range of the CSL: from the  
266 near-modern one (-25 m a.s.l.) to the maximum level of the Early Khvalynian transgression (+50 m a.s.l.), with a step  
267 of 15 meters, a total of six experiments. The corresponding model domains are shown in Fig. 1b.

268 For each of the two paleo-periods and each sea level, the experiment was performed for 50 model years and was  
269 organized as follows (Table 1). First, a rough initial approximation for the annual mean river runoff was specified as  
270 a linear function of the sea area (Morozova et al., 2021). After that, a model spin-up was performed for five years, and  
271 then during the next 15 years of model integration the average water imbalance was calculated. At the end of the 20<sup>th</sup>  
272 year, the obtained average imbalance was subtracted from the river runoff, and the average anomaly was subtracted  
273 from the sea level field. This resulted in the equilibrium runoff value and reinitialized sea level, which were used to  
274 further proceed with the calculations. Another spin-up was performed for 10 years, and finally, the last 20 years of the  
275 experiment were used to analyze the fields of evaporation and precipitation over the sea.

276

277 **Table 1 – Stages of numerical experiments with the coupled ocean-ice model**

Years	Experiment stage
1 – 5	Initial approximation for the runoff. Spin-up.
6 – 20	Initial approximation for the runoff. Calculating water imbalance.
end of year 20	Applying corrections to runoff and sea level
21 – 30	Corrected runoff. Spin-up.
31 – 50	Corrected runoff. Analyzing the Caspian Sea water balance components

### 278 3.3 Investigating the chronology of large palaeochannels

279 Dating was carried out by the radiocarbon (<sup>14</sup>C) method in the laboratories of the Institute of Earth Sciences, St.  
280 Petersburg University (index LU) and the Institute of Geography, Russian Academy of Sciences, Moscow (index  
281 IGRAN). Plant remains and dispersed organic matter in gyttja were used for dating. Fresh water mollusk shells, which  
282 are frequently met in drill cores, were not used because of the high probability of date distortion due to the hard water  
283 effect. Boring for organics sampling was carried out by a mechanical corer, usually in the centre of the palaeochannel  
284 (depending on its accessibility for the machine). The geological structure of the palaeochannels usually distinguishes  
285 3-4 sedimentary units, from top to bottom: (1) overbank alluvia - silty loam, sandy loam, or peat in place of the filled  
286 up oxbow lake; (2) oxbow lake sediments - clayey loam; (3) sediments of the intermediated stage of the palaeochannel  
287 abandonment, when it was not yet completely isolated from the river and flow still continued; usually silty sand or  
288 sandy silts; (4) channel alluvium - sands, sands with gravel and pebbles. Below the bed of channel alluvium  
289 corresponding to the studied palaeochannel, there were often older alluvial deposits, which could be of diverse  
290 composition - sands, loams, gyttja (unit 5).

291 Samples from channel alluvium (unit 4) are preferred for dating as they correspond to the time of active palaeochannel  
292 development. However, the channel alluvium is well-washed and organic inclusions are rare. They are much more  
293 commonly found in unit 3 sediments. The process of gradual abandonment of channel meanders usually takes a few  
294 years, at the most a few decades. This is less than the usual interval of uncertainty of <sup>14</sup>C dates and from the point of





295 view of geological time can be considered as a moment. Therefore, we considered that the samples from unit 4 also  
296 belong to the time of active development of the palaeochannels, its very end. Unfortunately, in unit 4, as well as in  
297 unit 5, organic materials suitable for dating were found only in a small number of boreholes. They were much more  
298 common in unit 2. Because the existence of an oxbow lake in the palaeochannels could be very long (millennia),  
299 samples were taken only from the very bottom of unit 3 and when interpreting the dates obtained, it was taken into  
300 consideration that they refer to the time when the active development of the palaeochannels ceased. In addition, in  
301 some cases, it was possible to sample at 14C from unit 5, the ancient alluvium underlying the channel alluvium of the  
302 palaeochannel under study. Such dates were interpreted as predating the time of activity of the studied palaeochannel.

303 Thus, in terms of the stratigraphic position, the dates have been divided into three groups:

- 304 • dates from units 3, 4, giving the time of activity of large palaeochannels - activity dates;
- 305 • dates from unit 2, referring to the time when the studied palaeochannels had already been abandoned - post-  
306 dates;
- 307 • dates from unit 5, indicating the time when the large palaeochannels were not yet active - pre-dates.

308 In order to determine the total activity interval of large palaeochannels in the Volga basin within each of the groups,  
309 the dates were summarised. For this purpose, the OxCal 4.4 software Sum module (Bronk Ramsey, 2009) was used.

#### 310 **3.4 Modeling water inflow into the Caspian Sea from the ancient Volga catchment covered by permafrost**

311 Numerical experiments were carried out with a physically based model of runoff generation in the Volga River basin  
312 (Motovilov, 2016; Kalugin, 2022) developed on the basis of the ECOMAG hydrological modeling platform  
313 (Motovilov et al., 1999). Earlier, Gelfan and Kalugin (2021) applied the ECOMAG-based model of the Volga basin  
314 for assessing the river runoff sensitivity to the hypothetical permafrost distribution over the basin area.

315 The model describes spatially variable processes of snow accumulation and snowmelt, heat and water transfer within  
316 the vegetation-soil system, evapotranspiration, infiltration into frozen and unfrozen soil, soil freezing and thawing,  
317 surface, subsurface and groundwater flow into the river network, and river channel flow with a daily time-step. The  
318 model inputs include spatially distributed daily precipitation, air temperature and air humidity data. The Volga River  
319 basin was schematized onto grid cells with a mean area of 1750 km<sup>2</sup>.

320 A detailed description of the ECOMAG-based Volga River model, methods for setting the parameters and model  
321 verification results for the modern climate were presented by Gelfan and Kalugin (2021). In particular, it was shown  
322 that the developed model is robust against climate changes, i.e. it allows one to obtain stable (in statistical sense)  
323 results of hydrological simulations within the Volga River basin for years with contrasting climatic conditions. We  
324 consider the robustness of the hydrological model as a necessary condition for its applicability for paleohydrological  
325 reconstructions.

326 As the boundary conditions in our experiments, we used climate data simulated by the MPI-ESM-CR global climate  
327 model, which reproduced climate conditions of the deglaciation period (26-0 kyr BP) with prescribed ice sheets and  
328 surface topographies from ICE-6G reconstruction (Peltier et al., 2015) within the framework of PMIP4 experiment  
329 (Kapsch et al., 2021). The used climate data included monthly series of the near ground meteorological data obtained  
330 within a transit experiment Ice6G\_P2 (Kapsch et al., 2021) for the last 26,000 years with a hundred-year averaging  
331 period. The MPI-ESM-CR model has a spatial resolution of 3.75° in longitude and 3.7° in latitude on average.



332 For hydrological modeling, we applied climate simulation data for the four following periods: the post-LGM (18-17.1  
333 kyr BP), the Oldest Dryas (17-14.8 kyr BP), the Bølling (14.7-14.1 kyr BP) and the Allerød (14-12.8 kyr BP). Since  
334 a hydrological model requires daily data, the monthly MPI-ESM-CR-simulated data were transformed into the series  
335 of the corresponding daily values by the delta-change temporary downscaling method (Gelfan et al., 2017). For the  
336 transformation, we used daily data of the meteorological observations for the period of 1985-2014 at 306  
337 meteorological stations located within the Volga River basin. As a result, we constructed 30-year artificial time-series  
338 of daily precipitation, air temperature and air humidity, so that their mean values were equal to the corresponding  
339 long-term means calculated from monthly series for each of the four considered paleo-periods. The constructed series  
340 were assigned as the boundary conditions for the hydrological model.

341 Taking into account that the climatic boundaries of permafrost follow approximately with an isotherm of the mean  
342 annual air temperature below  $-5^{\circ}\text{C}$  (Smith, Riseborough, 2002), in our experiments, the presence of permafrost was  
343 assumed if the climatic data demonstrated a drop in the mean annual air temperature in the Volga basin below  $-5^{\circ}\text{C}$ ,  
344 i.e. by about  $10^{\circ}\text{C}$  less than the mean air temperature in the modern climate ( $+4.5^{\circ}\text{C}$ ). For all elements of the  
345 computational domain underlain by permafrost, the initial temperature of soils was set as negative from the ground  
346 surface to the depth of 3 meter (the depth of attenuation of the seasonal temperature fluctuations).

347 The hydrological model also took into account the features of the vegetation cover in the considered paleoperiods.  
348 Simakova (2008) and Makshaev (2019) showed that during the post-LGM and the Oldest Dryas, periglacial tundra  
349 landscapes were common in the ancient Volga basin. The model parameters corresponding to these landscapes were  
350 set using the Global Land Cover Characterization database (Loveland et al., 2000).

## 351 **4. Results and Discussion**

### 352 **4.1 Estimates of equilibrium river runoff to the Caspian Sea at the Early Khvalynyan transgression levels**


353 The numerical simulations with the INMIO COMPASS - CICE model (Sec. 3.2) provided estimates of the Caspian  
354 Sea water balance components for a wide range of possible CSLs under climatic conditions of the Last Glacial  
355 Maximum and the Holocene Climatic Optimum. Fig. 2 shows the average simulated values of evaporation and  
356 precipitation (mm/year) over the Caspian Sea surface area, as well as the river runoff volume ( $\text{km}^3/\text{year}$ ) required to  
357 maintain different prescribed values of the Caspian Sea at equilibrium conditions. As can be seen from Fig. 2, the  
358 average evaporation decreases when the CSL rises. This is related to the peculiarities of the Caspian Sea morphology:  
359 under the CSL rise, the coastline expands predominantly in the northern direction, where temperatures are lower, and  
360 the sea ice cover period is longer. Precipitation, on the contrary, slightly increases, but this growth does not compensate  
361 for the decrease in evaporation, so the average values of effective evaporation for the entire Caspian Sea surface area  
362 also decrease with the rising sea level above  $-25\text{ m a.s.l.}$  In general, the change in the equilibrium runoff is proportional  
363 to the change in the Caspian Sea surface area, but this dependence is not linear. For the CSL above  $-25\text{ m a.s.l.}$ , the  
364 Caspian Sea expands to the northern flat shore and the increase in the sea area accelerates.

365 This is accompanied by a decrease in the river discharge increment per unit area increase. For the level range of  $-25$   
366  $\div -10\text{ m a.s.l.}$ , this increment is  $0.55\text{ km}^3/\text{year}$  per  $10^3\text{ km}^2$  for mid-Holocene conditions, and  $0.40\text{ km}^3/\text{year}$  per  $10^3$   
367  $\text{km}^2$  for LGM. For the transgressive  $+35\div +50\text{ m a.s.l.}$  range, however, it becomes  $0.25\text{ km}^3/\text{year}$  per  $10^3\text{ km}^2$  for both  
368 mid-Holocene and LGM. Under LGM conditions, both evaporation and precipitation over the sea surface area are  
369 much lower than the corresponding values during mid-Holocene. Simulated evaporation is on average 180-200




370 mm/year lower, and precipitation is 70-90 mm/year lower, which results in 15-20% lower values of the equilibrium  
371 runoff in LGM compared to mid-Holocene conditions for the CSLs above -25 m a.s.l.


372



373



374



375

376 **Figure 2: Simulated Caspian Sea water balance components for different transgressive states under climatic conditions of**  
377 **the Last Glacial Maximum and the Holocene Climatic Optimum: averaged over the sea area evaporation (a), precipitation**  
378 **(b), and equilibrium river runoff (c) as a function of the sea level.**

379

380 Given lower air temperatures during LGM and a large shallow water area in the north at transgressive states of the  
381 Caspian Sea, the sea ice cover extent and duration play a major role in the decrease in evaporation from the sea surface.  
382 Model simulations suggest that the evaporation changes are affected by sea ice export to the warmer southern part of  
383 the Sea driven by sea circulation and surface winds. This effect is important not only during the spring melting season,  
384 but also in winter on the marginal freezing part of the water area, where the sea ice is thin.


385 The chosen LGM and mid-Holocene periods presumably represent the most contrasting climatic conditions during  
386 the late Pleistocene-early Holocene, so we interpreted the simulated values of the equilibrium river runoff as a possible  
387 range of changes during the deglaciation period under consideration. According to our results, the river runoff values  
388 required to sustain the CSL at the highest dated transgressive state at +35 m a.s.l. (17-13 kyr BP) belong to the range  
389 of 400-470 km<sup>3</sup>/year. Assuming that the contribution of the Volga River runoff to the total river discharge in that  
390 period was close to the modern one (about 80%), we estimated the river runoff from in the Volga watershed during



391 the period of the Early Khvalynian transgression ((18)17-13 kyr BP) as 320-375 km<sup>3</sup>/year, i.e. 1.3-1.5 times larger  
392 than the present day's values.


#### 393 4.2 Results of dating large palaeochannels in the Volga basin

394 Drilling of large palaeochannels in different parts of the Volga basin was carried out and 14C dates were obtained for  
395 a part of the boreholes (Fig. 3). A total of 57 dates suitable for statistical analysis of the palaeochannel activity time  
396 were obtained. Dates were received from the valleys of 18 rivers: Dubna, Medveditsa, Ustyia (upper Volga basin),  
397 Moskva, Protva, Moksha (Oka basin), upper Kama, Izh, Kilmez, Lolog, Yazva, Dema (Kama basin), Samara, Sok,  
398 Buzuluk, B. Cheremshan, B. Kinel (lower Volga basin), B. Uzen (Northern Pre-Caspian).



399  
400 **Figure 3: Map of cores made in large palaeochannels over the Volga basin (1 – all cores, 2 – dated cores; type of dates: 3 –**  
401 **post-dates, 4 – activity dates, 5 – pre-dates. Numbers are central points of 14C calibrated dates.**  
402

403 All dates are divided into three groups - 19 activity dates, 21 post-dates and 17 pre-dates (see the Methods section)  
404 and for each group the summation was done in OxCal 4.4 (Fig. 4). The resulting distributions suggest the following.  
405 The direct dates in the channel alluvium of the large palaeochannels form two clusters, the main one between 13.8-  
406 17.3 ka BP and a small complementary one between 18.2-18.8 ka BP. The latter overlaps with the youngest part of  
407 the distribution of dates in the underlying sediments (pre-dates), from which we can conclude that, with a high  
408 probability, there is no generation of palaeochannels of the corresponding age. This cluster of dates may be related to  
409 the dating of redeposited ancient organics.



410

411

412

**Figure 4: Summed distributions of radiocarbon dates from large palaeochannels in river valleys of the Volga basin.**

413

414

415

416

417

418

419

420

421

422

On the right, the distribution of dates in the fluvial alluvium is clearly limited to dates in the overlying sediments (post-dates). It should be noted that in the interval of 12.5-13.8 ka BP, the dates for the overlying sediments are derived from the bottoms of the palaeochannel fills (those cases where there was no material suitable for dating in the channel alluvium). However, at present one can only say with certainty that the stage of large palaeochannel formation and the corresponding epoch of high river runoff in the Volga basin lasted from at least 17.5 to 14 ka BP. A visual analysis of the map (Fig. 3) shows no regional differences in dates, i.e. the epoch started and ended geologically simultaneously in the whole Volga basin. Attention is drawn to the gap in the dates in the interval from 12.5 to 11.5 ka BP corresponding to the Younger Dryas epoch and the very beginning of the Holocene. This may be a result of a shortage of organic material due to scarcity of vegetation during this harsh epoch, but more likely reflects low fluvial activity and a significant drop in river flow in general.

423

424

425

426

427

428

429

430

431

432

433

434

435

The determined interval of activity of big palaeochannels shows that from at least 17.5 to 14 ka BP the Volga River runoff considerably exceeded the modern one. This corresponds generally to the palaeoclimate estimates from paleofloristic data by Borisova (2021) who established a significant increase in atmospheric precipitation in the central East European Plain in the second half of MIS 2 during the warming events 17–19 ka BP (the Late Pleniglacial) and 13–14.5 ka BP (the Bølling and Allerød interstadials). The Oldest Dryas cooling at 14.5–17 ka BP was characterized by a decrease in precipitation below the present-day values, but the high runoff coefficients due to the existence of permafrost could have favored still high runoff values. These estimates point that during the aforementioned period of big palaeochannel activity, the flow hardly remained constant, but it cannot be determined by geomorphological methods: among large palaeochannels there are no distinctive age generations that would differ consistently in size. All large palaeochannels make up a single set of forms, clearly differing in size and position in the valley floor topography from younger palaeochannels, the sizes of which correspond to modern rivers. The distribution of dates for the large palaeochannels also does not reveal clear periodicity or discontinuity on the basis of which the internal periodicity of the high flow epoch could be judged. Perhaps the available number of dates is not yet sufficient for this.



436 At this stage we can only mark the time frames of the epoch of high river discharge, which began no later than 17.5  
437 ka BP and ended no earlier than 14 ka BP, and relate the estimate of the annual Volga runoff magnitude obtained from  
438 the size of the palaeochannels ( $420 \text{ km}^3$  (Sidorchuk et al., 2021)) to this epoch as a whole. Probably the drop of activity  
439 dates at around 16 ka (Fig. 4) marks the Oldest Dryas pause in high river flow and big channel formation, but to  
440 establish it reliably a much larger massif of dates is necessary.

441 The interval of increased inflow of river water into the Caspian Sea from 17.5 to 14 ka BP corresponds exactly to the  
442 main phase of the Early Khvalynian transgression dated by marine sediments in the Northern Caspian Lowland from  
443 18-17 to 14-13 ka BP (see the review in the Introduction). It was shown in section 4.1 that such amount of the Volga  
444 runoff was more than enough to keep the Caspian level at +35 m a.s.l. - the highest dated shoreline of the Khvalynian  
445 transgression (remember that the considered maximum level of  $+48 \div +50$  m a.s.l. has not yet been characterized by  
446 any direct date - see the review in the Introduction).

447 What could be the reasons for such a significant increase in river runoff? The involvement of glacial meltwater is  
448 excluded because large palaeochannels are present in various parts of the Volga basin, including those completely  
449 isolated not only from the last, but also from all Quaternary glaciations in general (for example, basins of the lower  
450 Volga or right tributaries of the Oka). It is easy to show that possible increase in river runoff due to thawing of  
451 permafrost, which undoubtedly took place after the LGM, was also negligible. Let us assume that water exchange  
452 between groundwater and river water covered the upper 100 m of the Earth's crust. Let us also assume that during the  
453 last glacial epoch, this entire stratum had a deliberately overestimated ice content of 50%, and the deliberately  
454 unfeasible condition that all meltwater entered the river network when the permafrost melted. It is not difficult to  
455 calculate that if this 100-meter layer of permafrost had melted during the above 3,000-year period, it would have  
456 increased the annual river runoff from the modern basin area by less than  $23 \text{ km}^3$ , which is less than 10% of the  
457 average modern flow volume in the Volga basin. It should be emphasized that this estimate is repeatedly  
458 overestimated. In reality, the additional inflow of water due to melting permafrost could be an order of magnitude  
459 less.

460 Thus, huge water flowing into the Caspian Sea from the Volga basin during the period from 17 to 13 ka BP could only  
461 be of atmospheric origin (except for possible minor glacial meltwater runoff from the sources of the Volga itself at  
462 the very beginning of this period as demonstrated by Panin et al. (2021)). As mentioned in the Introduction, Gelfan  
463 and Kalugin (2021) quantified a significant decrease in runoff losses due to the hypothetical spread of permanently  
464 frozen soils over the Volga catchment and the resulting increase in the runoff coefficient, i.e. proportion of  
465 precipitation involved in the river runoff formation. But the question arises: is the amount of precipitation  
466 corresponding to the cryoarid climate of the deglaciation epoch enough to form an extraordinary river runoff even  
467 with the spread of permafrost over the catchment area of the Caspian Sea? To answer this question, we carried out  
468 numerical experiments with a hydrological model that reproduce the formation of river inflow into the Caspian Sea in  
469 the climatic conditions of the period from 17 to 13 ka BP and under the assumption of frozen catchment area of the  
470 sea. The results are presented in the next section.


#### 471 **4.3 Modeling the Volga River runoff in the climate conditions from the post-LGM to the Allerød (18-13 kyr** 472 **BP)**

473 Fig. 5 illustrates changes in the mean annual precipitation, air temperature and air humidity deficit assessed from the  
474 MPI-ESM-CR-simulated monthly data and averaged over the Volga basin for four periods: the post-LGM (18-17.1  
475 kyr BP), the Oldest Dryas (17-14.8 kyr BP), the Bølling (14.7-14.1 kyr BP) and the Allerød (14-12.8 kyr BP), covering





476 the epoch of the Early Khvalynian transgression. According to these data, all considered periods were colder than the  
477 modern climate in the Volga River basin, herewith each subsequent period was warmer than the previous one. Mean  
478 annual precipitation values assessed for different periods were 18-34% less than the modern value. Due to the cold  
479 climate, all the periods are characterized by an increase in the mean annual solid precipitation from 7% in the post-  
480 LGM and the Bølling to 41% in the Allerød (relative to the modern values). On the contrary, the mean annual liquid  
481 precipitation sum decreased from 45% in the Oldest Dryas to 54% in the Bølling. The mean annual air humidity  
482 deficit, which affects evaporation from the catchment surface, turned out to be lower than the modern one by an  
483 average of 40-50% in different periods.



484  
485 **Figure 5: MPI-ESM-CR-simulated data of the mean annual air temperature, total precipitation and air humidity deficit,**  
486 **averaged over the Volga basin, during the considered periods of paleo-time and under the modern climate.**  
487

488 Taking into account the cold climate in the post-LGM period, when the average annual temperature was 12.6°C lower  
489 than the present one (see Fig. 5), the Oldest Dryas (10.9°C lower) and the Bølling (9.4°C lower), we assumed that the  
490 whole catchment area was covered by continuous permafrost during these three periods. Generally, this assumption  
491 corresponds to the paleogeographic findings of Sidorchuk et al. (2008) and Borisova (2021). An algorithm that allows



492 taking into account the hypothetical presence of permanently frozen ground in the Volga River catchment and  
493 modeling the hydrological effect of permafrost was described by Gelfan and Kalugin (2021).


494 Numerical experiments with the hydrological model, which was forced by the temporary downscaled paleo-climate  
495 data, demonstrated that the mean annual runoff of the ancient Volga during the post-LGM period and the Oldest Dryas  
496 increased in comparison with the modern one for the period of 1985-2014 (259 km<sup>3</sup>) by 24% and 38%, respectively  
497 (Fig. 6). The runoff rising during the Oldest Dryas was larger due to larger mean precipitation. The permafrost led to  
498 a decrease in the infiltration capacity of the soils by more than an order of magnitude in comparison with the unfrozen  
499 soil over the river catchment. Decreased soil infiltration resulted in an increase in the mean runoff coefficient to as  
500 much as 0.67, i.e. 2/3 of precipitation falling on the catchment was not lost and reached the river channels and then  
501 the Caspian Sea (note that the mean annual runoff coefficient in the modern climate for the Volga basin is 0.35, i.e.  
502 almost twice as low). As a result, the assessed permafrost-induced changes in the runoff coefficient could themselves  
503 lead to an increase in the mean runoff even with a decrease in the mean precipitation comparing with the modern one.  
504 And this growth became especially noticeable due to the reduced evaporation from the catchment area caused by the  
505 decrease in the air humidity deficit during the post-LGM period and the Oldest Dryas (see Fig. 5). At the same time,  
506 the mean runoff visibly dropped during the Bølling period in spite of the permafrost presence that can be explained  
507 by a 5-15% decrease in precipitation with a simultaneous 40-45% increase in evaporation (owing to the rise in air  
508 humidity deficit) during this period comparing with the previous ones. During the Allerød, the mean runoff was also  
509 less than during the post-LGM or the Oldest Dryas, but the difference is not as significant as for the Bølling, owing to  
510 the rising precipitation and decreasing evaporation. The response of different parts of the Volga River basin to climate  
511 impacts differed from the response of the entire basin as a whole (see Fig. 6).

512 During the high-flow post-LGM and Oldest Dryas periods, the river runoff was mostly formed in the right-bank sub-  
513 catchments of the middle Volga: e.g. within the boundary of the modern Oka River basin, the runoff was 70% more  
514 than the spatially averaged one for the Volga basin. This result is confirmed by the data of a paleogeographic  
515 reconstruction of the runoff of ancient channels, most of the traces of which are located on the right bank of the middle  
516 Volga. On the contrary, on the catchment areas of the Upper Volga and the left-bank part of the middle Volga (Kama  
517 basin), the river runoff is estimated to be 30-40% less than the average value for the basin.


518 According to the simulation results, significant changes occurred in the intra-annual flow regime of the Volga in  
519 comparison with the modern regime. In the modern climate, the high flow season runs from April to June and makes  
520 up 54% of the annual runoff. In the considered paleo-periods, the high-flow season was a month later (from May to  
521 July), and the share of the annual runoff for these months varied from 75% to 85% with the largest value in the Oldest  
522 Dryas (Fig. 7). The simulated runoff from the sub-basins of the Oka and Kama Rivers, as well as from the Upper  
523 Volga was generally characterized by the same tendencies as for the runoff from the whole Volga. The most notable  
524 difference was a significant increase in the Oka freshet during the post-LGM and the Oldest Dryas, which we  
525 explained by a larger influence of permafrost together with the increased snow water equivalent due to an increased  
526 sum of the solid precipitation as mentioned above. The long-term mean of the annual peak discharge at the outlet of  
527 the Volga River during the post-LGM and the Oldest Dryas turned out to be 3 times higher than the corresponding  
528 mean simulated under the modern climate, and reached the values of 100,000 m<sup>3</sup>/s. The mean maximum discharge of  
529 the Oka River was as much as 4 times higher than the modern value, reaching 21,000 m<sup>3</sup>/s during the Oldest Dryas.  
530 A significant increase in the mean peak discharge of snowmelt flood compared to the current one was also obtained  
531 for the Upper Volga (3.7 times) and for the Kama (2.5 times). Peak flow makes the greatest contribution to the re-  
532 shaping of river channels, activates sediment flow and processes of transformation of channel forms. According to the



533 hypothesis of Sidorchuk et al. (2021), it was the snowmelt floods that turned out to be the main driver of fluvial  
534 activity and the formation of the palaeochannels occurring in the modern Volga basin and formed between 17.5 ka  
535 BP and 14 ka BP.  
536



537  
538 **Figure 6: The mean annual runoff simulated for different periods of deglaciation and for the modern climate. Top to down:**  
539 **the entire Volga basin, the Oka River Basin (right-bank part of the middle Volga), upper part of the basin, the Kama River**  
540 **Basin (left-bank part of the middle Volga).**  
541



542

543 **Figure 7: The mean monthly flow simulated for the different periods of deglaciation and for the modern climate. Top to**  
544 **down: the entire Volga basin, the Oka River Basin (right-bank part of the middle Volga), upper part of the basin, the Kama**  
545 **River Basin (left-bank part of the middle Volga).**

546

547 Our results do not contradict this hypothesis, since the largest increase in the simulated mean peak flow occurred in  
548 the Volga basin during the Oldest Dryas period (17-14.8 kyr BP).

549 Thus, we summarized that, according to the data of paleoclimate modeling, the climate of the Volga basin in the period  
550 from 18 kyr BP to the end of the Oldest Dryas (14.8 kyr BP) was characterized by low air temperature (11-13°C less  
551 than in the modern climate) and low precipitation (24-32% less than in the modern climate). At the same time,  
552 according to our experiments with the hydrological model, the mean annual Volga runoff during the Oldest Dryas  
553 (17-14.8 kyr BP) could reach up to 360 km<sup>3</sup>, which is almost 40% higher than the modern runoff, and the mean annual  
554 peak flow could increase 3 times. The main factors of the increased runoff were a decrease in evaporation from the



555 Volga paleo-catchment as well as the spread of permafrost reducing runoff losses due to infiltration into soils, which  
556 all together compensated, over and above, for the decrease in precipitation.

557 Note that the significant hydrological role of permafrost in the considered paleoperiod could be significantly less in  
558 the process of its degradation in later periods. This can be evidenced, in particular, by the end of increased flow shortly  
559 after 14 ka BP, i.e. in the Allerød, which can hypothetically be associated with thawing of the permafrost by that time.  
560 However, the permafrost completely recovered during the Younger Dryas stadial (12.8-11.8 ka BP), but the formation  
561 of large palaeochannels did not resume during this period. On the contrary, it was noted above that there is a dip in  
562 dates for the 12.5-11.5 ka BP interval, which may indicate a decrease in fluvial activity. This is also supported by the  
563 coincidence of this period with a drop in the sea level, the Yenotaevkian regression (Makshaev and Tkach, 2023).

## 564 **5 Conclusions**

565 Our study was aimed at verifying the physical consistency of the hypothesis asserting the hydroclimatic origin of the  
566 Early Khvalynian transgression of the Caspian Sea. When *a priori* formulating the hypothesis, we firstly relied on the  
567 up-to-date and well-founded OSL-datings (Kurbanov et al., 2021, 2022, 2023; Butuzova et al., 2022; Taratunina et  
568 al., 2022), which referred the sea level stage well above +10 m a.s.l. (likely up to +22 ÷ +35 m a.s.l.) to the final period  
569 of deglaciation, 17-13 kyr BP. Nowadays, this is the highest dated sea level rise in the Quaternary history of the  
570 Caspian Sea, since the maximum stage of the Early Khvalynian transgression (+48+50 m a.s.l.) has still not been dated  
571 in any geochronological study. Secondly, we relied on the results of recent (Panin et al., 2020, 2021; Borisova et al.,  
572 2021) and earlier (Kalinin et al., 1966; Panin et al., 2005; Sidorchuk et al., 2009) publications, which argued a  
573 negligible contribution of meltwater runoff (due to the Scandinavian ice-sheet melting and outflows of ice-dammed  
574 proglacial lakes) to the transgression of the sea during the considered, 17-13 kyr BP, period. Thirdly, our hypothesis  
575 was based on the ubiquitous presence of large river palaeochannels, whose age was estimated within the close interval,  
576 18-13 kyr BP, in the Caspian Sea catchment and adjacent river basins (Borisova et al., 2006; Sidorchuk et al., 2009;  
577 Panin et al., 2013, 2017; Panin and Matlakhova, 2015). Herewith, the palaeochannels are located in various parts of  
578 the Volga basin, including those completely isolated not only from the last, but also from all Quaternary glaciations,  
579 so the glacial meltwater was unlikely to contribute to their formation (Sidorchuk et al., 2009; 2021).

580 Thus, previous studies have given us the reasons to believe that the hypothesis put forward does not contradict the  
581 present knowledge on the nature of the Early Khvalynian transgression. That is why we reduced the hypothesis  
582 verification to evaluation of its physical feasibility, i.e. the physical feasibility of the CSL rise above +10 m a.s.l. under  
583 the climate of the deglaciation period, 17-13 kyr BP, in the absence of visible glacial meltwater effect. We carried out  
584 a comprehensive study of the physical consistency of the proposed hypothesis and obtained the following new results:

585 1. Using the coupled ocean and sea-ice general circulation model INMIO COMPASS – CICE driven by the climate  
586 model INMCM4.8 in accordance with the PMIP4 and CMIP6 modelling protocols, we estimated the equilibrium water  
587 runoff (irrespective of its origin), which could be sufficient to maintain the considered sea level under the modelled  
588 effective evaporation from the entire sea surface area. We found that the mean equilibrium runoff into the Caspian  
589 Sea for its highest dated transgressive state at +35 m a.s.l. (17-13 kyr BP) should fall within the range of 400-470  
590 km<sup>3</sup>/year. Assuming that the contribution of the Volga River runoff to the total river discharge in that period was close  
591 to the modern one (about 80%), we estimated the river runoff from the Volga River basin during the aforementioned  
592 period as 320-375 km<sup>3</sup>/year, i.e. 1.3-1.5 times larger than the present day's annual runoff.



593 2. An extensive <sup>14</sup>C-dating of the activity of palaeochannels located in the valleys of 18 rivers in the Volga basin we  
594 conducted, allowed us to narrow down the time frames of the epoch of high river discharge to 17.5-14 ka BP and  
595 relate the estimate of the annual Volga runoff magnitude derived earlier from the size of the palaeochannels (420  
596 km<sup>3</sup>/year (Sidorchuk et al., 2021)) to this epoch. Again, the updated time frames are almost identical to the  
597 aforementioned modern dating of the main phase of the Early Khvalynian transgression (17-13 ka BP), i.e. the  
598 estimates obtained by the independent methods turned out to be very close. Importantly, the estimate of the runoff that  
599 formed the studied palaeochannels occurred not far from and higher than the above maximum estimate of the  
600 equilibrium runoff: 420 km<sup>3</sup>/year and 375 km<sup>3</sup>/year, respectively. That is, the river flow passing through the ancient  
601 palaeochannels could maintain the sea level above +10 m a.s.l. under the climate of the considered epoch. As a result,  
602 we argued that 17.5-14 ka BP were thousands of years with a huge water inflow capable of maintaining the Caspian  
603 Sea level at the maximum dated marks of the Early Khvalynian transgression, and this inflow was not of glacial origin.

604 3. Using an ECOMAG-based hydrological model of the Volga runoff generation forced by paleoclimate data, we  
605 analyzed physically consistent mechanisms of an extraordinary high water inflow into the Caspian Sea both in the  
606 absence of visible glacial meltwater effect and under the drier and colder climate than the modern one (e.g., during  
607 the Oldest Dryas, 17-14.8 kyr BP, the air temperature was 10.9°C and precipitation was 24% less than in the modern  
608 climate). Nevertheless, our numerical experiments demonstrated that the mean annual Volga runoff during the Oldest  
609 Dryas could reach up to 360 km<sup>3</sup>, which is almost 40% higher than the modern runoff, and the mean annual peak flow  
610 could increase 3 times. The main factors of the increased runoff were the spread of permafrost which resulted in a  
611 sharp drop in infiltration into the frozen ground and reduced evaporation from the Volga paleo-catchment, which all  
612 together compensated, over and above, for the decrease in precipitation. A huge growth of peak flow during the Oldest  
613 Dryas, 17-14.8 kyr BP, greatly contributed to the processes of river channel transformation and could have formed  
614 the giant channels over the ancient Volga catchment.

615 Thus, our results do not contradict the hypothesis put forward, that the Early Khvalynian transgression of the Caspian  
616 Sea could be initiated and maintained solely by hydroclimatic factors within the deglaciation period, 17-13 ka BP.  
617 Also, the hypothesis has proven to be physically feasible, since we found a possible cause of the huge inflow into the  
618 Caspian Sea in the absence of visible glacial meltwater contribution.

#### 619 **Code/Data availability**

620 Paleoclimate Simulation Datasets related to this paper can be found  
621 at [https://pure.mpg.de/pubman/faces/ViewItemOverviewPage.jsp?itemId=item\\_3187396\\_4](https://pure.mpg.de/pubman/faces/ViewItemOverviewPage.jsp?itemId=item_3187396_4), an open-source online  
622 data repository hosted at MPG PuRe (Kageyama et al., 2021).

#### 623 **Author contribution**

624 **Alexander Gelfan:** Conceptualization of the study, Methodology of paleo-hydrological study, Writing, Reviewing  
625 and Editing; **Andrey Panin:** Methodology of palaeochannels dating, Field works; Writing, Reviewing and  
626 Editing; **Andrey Kalugin:** Paleo-hydrological simulations, Writing and Editing; **Polina Morozova:** Paleo-climate  
627 simulations, Writing; **Vladimir Semenov:** Methodology of assessing equilibrium river inflow into the sea, Writing;  
628 **Alexey Sidorchuk:** Methodology of assessing palaeochannel flow; **Vadim Ukraintsev:** Palaeochannels dating, Field  
629 works; **Konstantin Ushakov:** coupled ocean and sea-ice simulations.





630 **Competing interests**

631 The authors declare that they have no known competing financial interests or personal relationships that could have  
632 appeared to influence the work reported in this paper.

633 **Acknowledgements**

634 Radiocarbon dating of alluvial deposits and the numerical experiments with the ocean model were financially  
635 supported by the Russian Science Foundation (Grant 19-17-00215). Hydrological simulations were designed within  
636 the framework of the State Assignment theme № FMWZ-2022-0001. Geomorphological investigations in river  
637 floodplain contribute to the State Assignment theme № FMGE-2019-0005.

638 The present work was carried out within the framework of the Panta Rhei Research Initiative of the International  
639 Association of Hydrological Sciences (IAHS).

640 **References**

641 Arpe, K., and Leroy, S. A.: The Caspian Sea Level forced by the atmospheric circulation, as observed and modelled,  
642 *Quaternary international*, 173, 144–152, <https://doi.org/10.1016/j.quaint.2007.03.008>, 2007.

643 Arpe, K., Leroy, S. A. G., Lahijani, H., and Khan, V.: Impact of the European Russia drought in 2010 on the Caspian  
644 Sea level, *Hydrol. Earth Syst. Sci.*, 16, 19–27, <https://doi.org/10.5194/hess-16-19-2012>, 2012.

645 Arpe, K., Tsuang, B. J., Tseng, Y. H., Liu, X. Y., and Leroy, S. A.: Quantification of climatic feedbacks on the Caspian  
646 Sea level variability and impacts from the Caspian Sea on the large-scale atmospheric circulation, *Theoretical  
647 and Applied Climatology*, 136, 475–488, <https://doi.org/10.1007/s00704-018-2481-x>, 2019.

648 Arslanov, K. A., Yanina, T. A., Chepalyga, A. L., Svitoch, A. A., Makshaev, R. R., Maksimov, F. E., Chernov, S. B.,  
649 Tertychniy, N. I., and Starikova, A. A.: On the age of the Khvalynian deposits of the Caspian Sea coast according  
650 to  $^{14}\text{C}$  and  $^{230}\text{Th}/^{234}\text{U}$  methods, *Quaternary International*, 409, 81–87,  
651 <https://doi.org/10.1016/j.quaint.2015.05.067>, 2016.

652 Borisova, O., Konstantinov, E., Utkina, A., Baranov, D., and Panin, A.: On the existence of a large proglacial lake in  
653 the Rostov-Kostroma lowland, north-central European Russia, *J. Quat. Sci.*, 37 (8), 1442–1459,  
654 <https://doi.org/10.1002/jqs.3454>, 2022.

655 Borisova, O., Sidorchuk, A., and Panin, A.: Palaeohydrology of the Seim River basin, Mid-Russian Upland, based on  
656 palaeochannel morphology and palynological data, *Catena*, 66 (1), 53–73,  
657 <https://doi.org/10.1016/j.catena.2005.07.010>, 2006.

658 Borisova, O. K.: Landscape and Climatic Conditions in the Central East European Plain in the last 22 thousand Years:  
659 Reconstruction based on Paleobotanical Data, *Water Resour.*, 48, 886–896,  
660 <https://doi.org/10.1134/S0097807821060038>, 2021.

661 Brierley, C. M., Zhao, A., Harrison, S. P., Braconnot, P., Williams, C. J. R., Thornalley, D. J. R., Shi, X., Peterschmitt,  
662 J.-Y., Ohgaito, R., Kaufman, D. S., Kageyama, M., Hargreaves, J. C., Erb, M. P., Emile-Geay, J., D'Agostino,  
663 R., Chandan, D., Carré, M., Bartlein, P. J., Zheng, W., Zhang, Z., Zhang, Q., Yang, H., Volodin, E. M., Tomas,  
664 R. A., Routson, C., Peltier, W. R., Otto-Bliesner, B., Morozova, P. A., McKay, N. P., Lohmann, G., Legrande,



- 665 A. N., Guo, C., Cao, J., Brady, E., Annan, J. D., and Abe-Ouchi, A.: Large-scale features and evaluation of the  
666 PMIP4-CMIP6 midHolocene simulations, *Clim. Past.*, 16, 1847–1872, [https://doi.org/10.5194/cp-16-1847-](https://doi.org/10.5194/cp-16-1847-2020)  
667 2020, 2020.
- 668 Bronk Ramsey, C.: Bayesian analysis of radiocarbon dates, *Radiocarbon*, 51 (1), 337–360,  
669 [https://doi.org/10.2458/azu\\_js\\_rc.51.3494](https://doi.org/10.2458/azu_js_rc.51.3494), 2009.
- 670 Butuzova, E. A., Kurbanov, R. N., Taratunina, N. A., Makeev, A. O., Rusakov, A. V., Lebedeva, M. P., Murray, A.  
671 S., and Yanina, T. A.: Shedding light on the timing of the largest Late Quaternary transgression of the Caspian  
672 Sea, *Quaternary Geochronology*, 73, 101378, <https://doi.org/10.1016/j.quageo.2022.101378>, 2022.
- 673 Chen, J. L., Pekker, T., Wilson, C. R., Tapley, B. D., Kostianoy, A. G., Cretaux, J. F., and Safarov, E. S.: Long-term  
674 Caspian Sea level change, *Geophys. Res. Lett.*, 44 (13), 6993–7001, <https://doi.org/10.1002/2017GL073958>,  
675 2017.
- 676 Chepalyga, A. L.: Late glacial great flood in the Ponto-Caspian basin, in: *The Black Sea Flood Question: Changes in*  
677 *Coastline, Climate, and Human Settlement*, edited by: Yanko-Hombach, V., Gilbert, A.S., Panin, A., and  
678 Dolukhanov, P.M., Springer, Dordrecht, 119–148, [https://doi.org/10.1007/978-1-4020-5302-3\\_6](https://doi.org/10.1007/978-1-4020-5302-3_6), 2007.
- 679 Fadeev, R., Ushakov, K., Tolstykh, M., and Ibrayev, R.: Design and development of the SLAV-INMIO-CICE coupled  
680 model for seasonal prediction and climate research, *Russian J. Numerical Analysis and Mathematical Modelling*,  
681 33 (6), 333–340, <https://doi.org/10.1515/rnam-2018-0028>, 2018.
- 682 Fedorov, P. V.: Stratigraphy of Quaternary sediments and the history of the development of the Caspian Sea, *Proc. of*  
683 *the Geological Institute of the Academy of Science of the USSR*, 2 (10), 1–308, 1957 (in Russian).
- 684 Fedorov, P. V. (Ed.): *Pleistocene of the Ponto-Caspian*. Nauka Press, Moscow, 165 pp., 1978 (in Russian).
- 685 Forte A. M., Cowgill E.: Late Cenozoic base-level variations of the Caspian Sea: a review of its history and proposed  
686 driving mechanisms, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 386, 392–407,  
687 <https://doi.org/10.1016/j.palaeo.2013.05.035>, 2013.
- 688 Frolov A. V. (Ed.): *Modeling of long-term fluctuations of the Caspian Sea level: theory and applications*, GEOS Publ.,  
689 Moscow, 174 pp., ISBN 5-89118-298-X, 2003 (in Russian).
- 690 Frolov, A. V.: Dynamic-Stochastic Modeling of the Paleo-Caspian Sea Long-Term Level Variations (14–4 Thousand  
691 Years BC), *Water Resour.*, 48, 854–863, <https://doi.org/10.1134/S0097807821060051>, 2021.
- 692 Gelfan, A., Gustafsson, D., Motovilov, Y., Arheimer, B., Kalugin, A., Krylenko, I., and Lavrenov, A.: Climate change  
693 impact on the water regime of two great Arctic rivers: Modeling and uncertainty issues, *Clim. Chang.*, 141, 499–  
694 515, <https://doi.org/10.1007/s10584-016-1710-5>, 2017.
- 695 Gelfan, A. N., and Kalugin, A. S.: Permafrost in the Caspian Basin as a Possible Trigger of the Late Khvalynian  
696 Transgression: Testing Hypothesis Using a Hydrological Model, *Water Resour.*, 48, 831–843,  
697 <https://doi.org/10.1134/S0097807821060063>, 2021.
- 698 Golitsyn, G. S., Ratkovich, D. Ya., Fortus, M. I., and Frolov, A. V.: On the present-day rise in the Caspian Sea level,  
699 *Water Resour.*, 25 (2), 117–122, 1998.



- 700 Grosswald, M. G., and Kotlyakov, V. M.: The great proglacial drainage system in Northern Eurasia and its significance  
701 for inter-regional correlations, in: Quaternary Period: Paleogeography and Lithology, edited by: Yanshin, A.L.,  
702 Kishinev, Stiintsya, 5–13, 1989 (in Russian).
- 703 Hunke E. C., Lipscomb W. H., Turner A. K., Jeffery N., and Elliott, S.: CICE: the Los Alamos Sea Ice Model  
704 Documentation and Software User's Manual Version 5.1, Los Alamos National Laboratory, 2015.
- 705 Ibrayev, R. A., Khabeev, R. N., and Ushakov, K. V.: Eddy-resolving 1/10° model of the World Ocean, *Izv. Atmos.*  
706 *Ocean Phys.*, 48, 37–46. <https://doi.org/10.1134/S0001433812010045>, 2012.
- 707 Kageyama, M., Harrison, S. P., Kapsch, M.-L., Lofverstrom, M., Lora, J. M., Mikolajewicz, U., Sherriff-Tadano, S.,  
708 Vadsaria, T., Abe-Ouchi, A., Bouttes, N., Chandan, D., Gregoire, L. J., Ivanovic, R. F., Izumi, K., LeGrande, A.  
709 N., Lhardy, F., Lohmann, G., Morozova, P. A., Ohgaito, R., Paul, A., Peltier, W. R., Poulsen, C. J., Quiquet, A.,  
710 Roche, D. M., Shi, X., Tierney, J. E., Valdes, P. J., Volodin, E., and Zhu, J.: The PMIP4 Last Glacial Maximum  
711 experiments: preliminary results and comparison with the PMIP3 simulations, *Clim. Past.*, 17, 1065–1089,  
712 <https://doi.org/10.5194/cp-17-1065-2021>, 2021.
- 713 Kakroodi, A. A., Kroonenberg, S. B., Hoogendoorn, R. M., Mohammadkhani, H., Yamani, M., Ghassemi, M. R., and  
714 Lahijani, H. A. K.: Rapid Holocene sea-level changes along the Iranian Caspian coast, *Quaternary International*,  
715 263, 93–103, <https://doi.org/10.1016/j.quaint.2011.12.021>, 2012.
- 716 Kalinin, G. P., Markov, K. K., and Suetova, I. A.: Fluctuations in the level of the Earth's water bodies in the geological  
717 past. Part I, *Oceanology*, 6 (5), 737–746, 1966 (in Russian).
- 718 Kalmykov, V. V., Ibrayev, R. A., Kaurkin, M. N., and Ushakov, K. V.: Compact Modeling Framework v3.0 for high-  
719 resolution global ocean–ice–atmosphere models, *Geosci. Model Dev.*, 11 (10), 3983–3997.  
720 <https://doi.org/10.5194/gmd-11-3983-2018>, 2018.
- 721 Kalnitskii, L. Y., Kaurkin, M. N., Ushakov, K. V., and Ibrayev, R. A.: Seasonal Variability of Water and Sea-Ice  
722 Circulation in the Arctic Ocean in a High-Resolution Model, *Izv. Atmos. and Ocean. Physics*, 56 (5), 522–533,  
723 <https://10.1134/S0001433820050060>, 2020.
- 724 Kalugin, A.: Hydrological and meteorological variability in the Volga River basin under global warming by 1.5 and  
725 2 degrees, *Climate*, 10 (7), 107, <https://doi.org/10.3390/cli10070107>, 2022.
- 726 Kapsch, M.-L., Mikolajewicz, U., Ziemer, F., and Schannwell, C.: Ocean response in transient simulations of the last  
727 deglaciation dominated by underlying ice-sheet reconstruction and method of meltwater distribution,  
728 *Geophysical Research Letters*, 49, e2021GL096767, <https://doi.org/10.1029/2021GL096767>, 2022.
- 729 Kislov, A., and Toropov, P.: East European River runoff and Black Sea and Caspian Sea level changes as simulated  
730 within the Paleoclimate Modeling Intercomparison Project, *Quaternary International*, 167, 40–48,  
731 <https://doi.org/10.1016/j.quaint.2006.10.005>, 2007.
- 732 Kislov, A. V., Panin, A. V., and Toropov, P.: Current status and palaeostages of the Caspian Sea as a potential  
733 evaluation tool for climate model simulations, *Quaternary International*, 345, 48–55,  
734 <https://doi.org/10.1016/j.quaint.2014.05.014>, 2014.
- 735 Koriche, S. A., Singarayer, J. S., Cloke, H. L., Valdes, P. J., Wesselingh, F. P., Kroonenberg, S. B., Wickert, A. D.,  
736 and Yanina, T. A.: What are the drivers of Caspian Sea level variation during the late Quaternary? *Quaternary*  
737 *Science Reviews*, 283, 107457, <https://doi.org/10.1016/j.quascirev.2022.107457>, 2022.



- 738 Krijgsman, W., Tesakov, A., Yanina, T., Lazarev, S., Danukalova, G., Van Baak, C. G. C., Agustí, J., Alçiçek, M. C.,  
739 Aliyeva, E., Bista, D., Bruch, A., Büyükeriç, Y., Bukhsianidze, M., Flecker, R., Frolov, P., Hoyle, T. M.,  
740 Jorissen, E. L., Kirscher, U., Koriche, S. A., Kroonenberg, S. B., Lordkipanidze, D., Oms, O., Rausch, L.,  
741 Singarayer, J., Stoica, M., van de Velde, S., Titov, V. V., and Wesselingh, F. P.: Quaternary time scales for the  
742 Pontocaspian domain: interbasinal connectivity and faunal evolution., *Earth-Sci. Rev.*, 188, 1–40,  
743 <https://doi.org/10.1016/j.earscirev.2018.10.013>, 2019.
- 744 Kroonenberg, S. B., Badyukova, E. N., Storms, J. E. A., Ignatov, E. I., and Kasimov, N. S.: A full sea level cycle in  
745 65 years: barrier dynamics along Caspian shores, *Sediment. Geol.*, 134, 257–274, [https://doi.org/10.1016/S0037-](https://doi.org/10.1016/S0037-0738(00)00048-8)  
746 [0738\(00\)00048-8](https://doi.org/10.1016/S0037-0738(00)00048-8), 2000.
- 747 Kurbanov, R., Murray, A., Thompson, W., Svistunov, M., Taratunina, N., and Yanina, T.: First reliable chronology  
748 for the Early Khvalynian Caspian Sea transgression in the Lower Volga River valley, *Boreas*, 50 (1), 134–146,  
749 <https://doi.org/10.1111/bor.12478>, 2021.
- 750 Kurbanov, R. N., Buylaert, J.-P., Stevens, T., Taratunina, N. A., Belyaev, V. R., Makeev, A. O., Lebedeva, M. P.,  
751 Rusakov, A. V., Solodovnikov, D., Költringer, C., Rogov, V. V., Streletskay, I. D., Murray, A. S., and Yanina,  
752 T. A.: A detailed luminescence chronology of the Lower Volga loess-palaeosol sequence at Leninsk, *Quaternary*  
753 *Geochronology*, 73, 101376, <https://doi.org/10.1016/j.quageo.2022.101376>, 2022.
- 754 Kurbanov, R. N., Belyaev, V. R., Svistunov, M. I., Butuzova, E. A., Solodovnikov, D. A., Taratunina, N. A., and  
755 Yanina, T. A.: New data on the age of the Early Khvalynian transgression of the Caspian Sea, *Izvestiya*  
756 *Rossiiskoi Akademii Nauk. Seriya Geograficheskaya*, 87(3), 1, 2023 (in Russian).
- 757 Kvasov, D. D. (Ed.): *The Late Quaternary history of large lakes and inland seas of Eastern Europe*, Suomalainen  
758 tiedekad., Helsinki, 71 pp, 1979.
- 759 Larsen, E., Kjar, K. H., Demidov, I., Funder, S., Grosfjeld, K., Houmark-Nielsen, M., Jensen, M., Linge, H., and Lysa,  
760 A.: Late Pleistocene glacial and lake history of northwestern Russia, *Boreas*, 35, 394–424,  
761 <https://doi.org/10.1080/03009480600781958>, 2006.
- 762 Leontiev, O. K.: Evolution of the Caspian shores in the Upper Pliocene and Quaternary period, in: *Geomorphological*  
763 *analysis during geological research in the Caspian lowland*, edited by: Aristarchova L. B., MSU Press, Moscow,  
764 106–140, 1968 (in Russian).
- 765 Leontiev, O. K., Rychagov, G. I., Kaplin, P. A., Svitoch, A. A., Parunin, O. B., and Shlyukov, A. I.: Chronology and  
766 palaeogeography of Ponto-Caspian (based on result of radiocarbon dating), *Pleistocene Palaeogeography and*  
767 *Sediments of Southern Seas of the USSR*, 26–38, 1977 (in Russian).
- 768 Loveland, T. R., Reed, B. C., Brown, J. F., Ohlen, D. O., Zhu, Z., Yang, L., and Merchant, J. W.: Development of a  
769 global land cover characteristics database and IGBP DISCover from 1 km AVHRR data, *International Journal*  
770 *of Remote Sensing*, 21 (6–7), 1303–1330, 2000.
- 771 Lyså, A., Jensen, M. A., Larsen, E., Fredin, O. L. A., and Demidov I. N.: Ice-distal landscape and sediment signatures  
772 evidencing damming and drainage of large proglacial lakes, NW Russia, *Boreas*, 40 (3), 481–497,  
773 <https://doi.org/10.1111/j.1502-3885.2010.00197.x>, 2011.
- 774 Makshaev, R. R.: Paleogeography of the Middle and Lower Volga Region during the Early Khvalynian Transgression  
775 of the Caspian Sea, Ph.D. thesis, Lomonosov Moscow State University, Moscow, 160 pp., 2019 (in Russian).



- 776 Makshaev, R. R., and Svitoch, A. A.: Chocolate Clays of the northern Caspian Sea region: distribution, structure, and  
777 origin, *Quaternary International*, 409, 44–49, <https://doi.org/10.15356/0435-4281-2015-1-101-112>, 2016.
- 778 Makshaev, R. R., and Tkach, N. T.: Chronology of Khvalynian stage of the Caspian Sea according to radiocarbon  
779 dating, *Geomorfologiya i Paleogeografiya*, 54 (1), 37–54 <https://doi.org/10.31857/S0435428123010108>, 2023  
780 (in Russian).
- 781 Morozova P. A.: Influence of the Scandinavian Ice Sheet on the climate conditions of the East European Plain  
782 according to the numerical modeling data of the project PMIP II, *Ice and Snow*, 54 (1), 113–124,  
783 <https://doi.org/10.15356/2076-6734-2014-1-113-124>, 2014 (in Russian).
- 784 Morozova P. A., Ushakov K. V., Semenov V. A., and Volodin E. M.: Water budget of the Caspian Sea in the Last  
785 Glacial Maximum by data of experiments with mathematical models, *Water Resour.*, 48 (6), 823–830,  
786 <https://doi.org/10.1134/S0097807821060130>, 2021.
- 787 Motovilov, Y.: Hydrological simulation of river basins at different spatial scales: 1. Generalization and averaging  
788 algorithms, *Water Resour.*, 43, 429–437, <https://doi.org/10.1134/S0097807816030118>, 2016.
- 789 Motovilov, Y., Gottschalk, L., Engeland, K., and Rodhe, A.: Validation of a distributed hydrological model against  
790 spatial observations, *Agric. For. Meteorol.*, 98–99, 257–277, [https://doi.org/10.1016/S0168-1923\(99\)00102-1](https://doi.org/10.1016/S0168-1923(99)00102-1),  
791 1999.
- 792 Naderi Beni, A., Lahijani, H., Mousavi Harami, R., Arpe, K., Leroy, S. A. G., Marriner, N., Berberian, M., Andrieu-  
793 Ponel, V., Djamali, M., Mahboubi, A., and Reimer, P.J.: Caspian Sea-level changes during the last millennium:  
794 historical and geological evidence from the south Caspian Sea, *Climate of the Past*, 9 (4), 1645–1665,  
795 <https://doi.org/10.5194/cp-9-1645-2013>, 2013.
- 796 Panin, A., Adamiec, G., Buylaert, J.-P., Matlakhova, E., Moska, P., and Novenko, E.: Two Late Pleistocene climate-  
797 driven incision/aggradation rhythms in the middle Dnieper River basin, west-central Russian Plain, *Quaternary  
798 Science Reviews*, 166, 266–288, <https://doi.org/10.1016/j.quascirev.2016.12.002>, 2017.
- 799 Panin, A. V., Astakhov, V. I., Lotsari, E., Komatsu, G., Lang, J., and Winsemann, J.: Middle and Late Quaternary  
800 glacial lakeoutburst floods, drainage diversions and reorganization of fluvial systems in northwestern Eurasia,  
801 *Earth-Science Reviews*, 201, 103069, <https://doi.org/10.1016/j.earscirev.2019.103069>, 2020.
- 802 Panin, A. V., and Matlakhova, E. Yu.: Fluvial chronology in the East European plain over the last 20 ka and its  
803 palaeohydrological implications, *Catena*, 130, 46–61, <https://doi.org/10.1016/j.catena.2014.08.016>, 2015.
- 804 Panin, A. V., and Sidorchuk, A. Ju., Borisova, O. K.: Fluvial processes and river runoff in the Russian Plain in the  
805 end of the Late Valdai epoch, in: *Geography Perspectives: to the 100th anniversary of K.K. Markov*, Geogr.  
806 Dep. MSU, Moscow, 114–127, 2005 (in Russian).
- 807 Panin, A. V., Sidorchuk, A. Y., and Ukraintsev, V. Y.: The Contribution of Glacial Melt Water to Annual Runoff of  
808 River Volga in the Last Glacial Epoch, *Water Resour.*, 48, 877–885,  
809 <https://doi.org/10.1134/S0097807821060142>, 2021.
- 810 Panin, A. V., Sidorchuk, A. Yu, and Vlasov, M. V.: High Late Valdai (Vistulian) runoff in the Don River basin,  
811 *Izvestiya Rossiiskoi Akademii Nauk. Seriya Geograficheskaya*, 1, 118–129, 2013 (in Russian).



- 812 Panin, A. V., Sorokin, A. N., Bricheva, S. S., Matasov, V. M., Morozov, V. V., Smirnov, A. L., Solodkov, N. N., and  
813 Uspenskaia, O. N.: Landscape development history of the Zabolotsky peat bog in the context of initial settlement  
814 of the Dubna River lowland (Upper Volga basin), *Vestnik Archeologii, Antropologii i Etnografii*, 2, 85–100.  
815 <https://doi.org/10.20874/2071-0437-2022-57-2-7>, 2022.
- 816 Panin, G. N. and Dianskii, N. A.: On the correlation between oscillations of the Caspian Sea level and the North  
817 Atlantic climate, *Izvestiya, Atmospheric and Oceanic Physics*, 50 (3), 266–278, [https://doi.org/](https://doi.org/10.1134/S000143381402008X)  
818 [10.1134/S000143381402008X](https://doi.org/10.1134/S000143381402008X), 2014.
- 819 Peltier, W. R., Argus, D. F., and Drummond, R.: Space geodesy constrains ice age terminal deglaciation: The global  
820 ICE-6G\_C (VM5a) model, *J. Geophys. Res. Solid Earth*, 120, 450–487, <https://doi.org/10.1002/2014JB011176>,  
821 2015.
- 822 Ratkovich, D. Ya.: Modern variations of the Caspian Sea level, *Water Resour.*, 20 (2), 160–171, 1993.
- 823 Rychagov, G.I.: Late Pleistocene history of the Caspian Sea, in: *Studies of the Caspian Sea*, edited by: Leontiev, O.K.,  
824 Maev, E.G., MSU Press., Moscow, 18–29, 1974 (in Russian).
- 825 Rychagov, G. I. (Ed.): *Pleistocene History of the Caspian Sea*, MSU Press, Moscow, 267 pp., ISBN 5-211-03828-2,  
826 1997 (in Russian).
- 827 Sidorchuk, A. Yu., Panin, A. V., and Borisova, O.K.: Climate-induced changes in surface runoff on the North-Eurasian  
828 plains during the late glacial and Holocene, *Water Resour.*, 35, 386–396.  
829 <https://doi.org/10.1134/S0097807808040027>, 2008.
- 830 Sidorchuk, A. Y., Panin, A. V., and Borisova, O.K.: Morphology of river channels and surface runoff in the Volga  
831 River basin (East European Plain) during the Late Glacial period, *Geomorphology*, 113 (3–4), 137–157,  
832 <https://doi.org/10.1016/j.geomorph.2009.03.007>, 2009.
- 833 Sidorchuk, A., Panin, A., and Borisova, O.: Surface runoff to the Black Sea from the East European Plain during Last  
834 Glacial Maximum–Late Glacial time, in: *Geology and Geoarchaeology of the Black Sea Region: Beyond the*  
835 *Flood Hypothesis*, edited by: Buynevich, I., Yanko–Hombach, V., Gilbert, A.S., and Martin, R.E., Geological  
836 Society of America Special Paper 473, pp. 1–25, [https://doi.org/10.1130/2011.2473\(01\)](https://doi.org/10.1130/2011.2473(01)), 2011.
- 837 Sidorchuk, A. Y., Ukraintsev, V. Y., and Panin, A. V.: Estimating Annual Volga Runoff in the Late Glacial Epoch  
838 from the Size of River Paleochannels, *Water Resour.*, 48, 864–876,  
839 <https://doi.org/10.1134/S0097807821060178>, 2021.
- 840 Simakova, A. N.: Evolution of vegetation of the Russian Plain and Western Europe in the Late Neopleistocene–Middle  
841 Holocene (33–4.8 thousand years BP) (from palynological data), Ph.D. thesis, Geological Institute of the Russian  
842 Academy of Sciences, Moscow, 34 pp., 2008 (in Russian).
- 843 Smith, M., and Riseborough, D.: Climate and the limits of permafrost: A zonal analysis, *Permafrost and Periglacial*  
844 *Processes*, 13 (1), 1–15, <https://doi.org/10.1002/ppp.410>, 2002.
- 845 Svitoch, A. A.: Khvalynian transgression of the Caspian Sea was not a result of water overflow from the Siberian  
846 proglacial lakes, nor a prototype of the Noachian flood, *Quaternary International*, 197, 115–125,  
847 <https://doi.org/10.1016/j.quaint.2008.02.006>, 2009.





- 848 Svitoch, A. A. (Ed.): The Big Caspian: Structure and History of Development, MSU Press, Moscow, 270 pp., ISBN  
849 978-5-19-010904-7, 2014 (in Russian).
- 850 Svitoch, A. A., Parunin, O. B., and Yanina, T. A.: Radiocarbon chronology of the deposits and events of late  
851 Pleistocene of the Ponto-Caspian region, in: Quaternary Geochronology, edited by: Murzaev, V.E., Puning Ya-  
852 M.K., Chichagova, O.A., Nauka, Moscow, pp. 75–82, 1994 (in Russian).
- 853 Svitoch, A. A., Selivanov, A. O., and Yanina, T. A.: Paleogeographic Events of the Ponto-Caspian and Mediterranean  
854 Pleistocene, RASHN, Moscow, pp. 289, 1988 (in Russian).
- 855 Svitoch, A. A., and Yanina, T. A. (Eds.): Quaternary Deposits of the Caspian Sea Coasts, MSU Press, Moscow, 267  
856 pp., 1997 (in Russian).
- 857 Taratunina, N. A., Buylaert, J. P., Kurbanov, R. N., Yanina, T. A., Makeev, A. O., Lebedeva, M. P., Utkina, A. O.,  
858 and Murray, A. S.: Late Quaternary evolution of lower reaches of the Volga River (Raygorod section) based on  
859 luminescence dating, Quaternary Geochronology, 72, 101369, <https://doi.org/10.1016/j.quageo.2022.101369>,  
860 2022.
- 861 Toropov, P. A., and Morozova, P. A.: Evaluation of the Caspian Sea level fluctuations during the Late Pleistocene  
862 cryochrome epoch based on the results of the numerical climate modeling, Vestn. Mosc. Univ. Ser. 5. Geogr. 2,  
863 55–61, 2011 (in Russian).
- 864 Tudryn, A., Leroy, S. A. G., Toucanne, S., Gibert-Brunet, E., Tucholka, P., Lavrushin, Y. A., Dufaure, O., Miska, S.,  
865 and Bayon, G.: The Ponto-Caspian basin as a final trap for southeastern Scandinavian Ice-Sheet meltwater,  
866 Quaternary Science Reviews, 148, 29–43, <https://doi.org/10.1016/j.quascirev.2016.06.019>, 2016.
- 867 Ukraintsev, V. Yu.: Evidences of the high river runoff in the river valleys of the Volga basin during the Late Glacial,  
868 Geomorfologiya, 53 (1), <https://doi.org/10.31857/S0435428122010126>, 2022.
- 869 Ushakov, K. V. and Ibrayev, R. A.: Assessment of mean world ocean meridional heat transport characteristics by a  
870 high-resolution model, Russ. J. Earth. Sci., 18, ES1004, <https://doi.org/10.2205/2018ES000616>, 2018.
- 871 Varuschenko, S. I., Varuschenko, A. N., and Klige, R. K. (Eds.): Changes in the Regime of the Caspian Sea and  
872 Closed Basins in Paleotime, Nauka, Moscow, 239 pp., 1987 (in Russian).
- 873 Volodin, E. M., Mortikov, E. V., Kostykin, S. V., Galin, V. Y., Lykossov, V. N., Gritsun, A. S., Diansky, N. A.,  
874 Gusev, A. V., Iakovlev, N. G., Shestakova, A. A., and Emelina, S. V.: Simulation of the modern climate using  
875 the INMCM48 climate model, Russ. J. Numer. Anal. Math. Modelling, 33 (6), 367–374,  
876 <https://doi.org/10.1515/rnam-2018-0032>, 2018.
- 877 Water balance and level fluctuations of the Caspian Sea. Modeling and prediction, (Ed. V. Gruzinov). Moscow,  
878 Rosgidromet. 375 pp, ISBN 978-5-9908623-0-2, 2016 (in Russian).
- 879 Yanina, T. A.: Correlation of the Late Pleistocene paleogeographical events of the Caspian Sea and Russian plain,  
880 Quaternary International, 271, 120–129, <https://doi.org/10.1016/j.quaint.2012.06.003>, 2012.
- 881 Yanina, T., Sorokin, V., Bezrodnykh, Y., and Romanyuk, B.: Late Pleistocene climatic events reflected in the Caspian  
882 Sea geological history (based on drilling data), Quaternary International, 465 (4), 130–141,  
883 <https://doi.org/10.1016/j.quaint.2017.08.003>, 2018.



- 884 Yanko-Hombach, V., and Kislov, A.: Late Pleistocene and Holocene sea-level dynamics in the Caspian and Black  
885 seas: Data synthesis and paradoxical interpretations, *Quaternary International*, 465, 63–71,  
886 <https://doi.org/10.1016/j.quaint.2017.11.030>, 2018.
- 887 Zekster, I. S.: Groundwater discharge into lakes: a review of recent studies with particular regard to large saline lakes  
888 in central Asia, *Int. J. Salt Lake Res.*, 4, 233–249, <https://doi.org/10.1007/BF02001493>, 1995