

INFECTIOUS DISEASES OF UGANDA



Stephen Berger, MD

GIDEON
E-BOOK SERIES

Infectious Diseases of Uganda - 2013 edition

Stephen Berger, MD

Copyright © 2013 by GIDEON Informatics, Inc. All rights reserved.

Published by GIDEON Informatics, Inc, Los Angeles, California, USA. www.gideononline.com

Cover design by GIDEON Informatics, Inc

No part of this book may be reproduced or transmitted in any form or by any means without written permission from the publisher. Contact GIDEON Informatics at ebook@gideononline.com.

ISBN-13: 978-1-61755-633-3

ISBN-10: 1-61755-633-5

Visit <http://www.gideononline.com/ebooks/> for the up to date list of GIDEON ebooks.

DISCLAIMER: Publisher assumes no liability to patients with respect to the actions of physicians, health care facilities and other users, and is not responsible for any injury, death or damage resulting from the use, misuse or interpretation of information obtained through this book. Therapeutic options listed are limited to published studies and reviews. Therapy should not be undertaken without a thorough assessment of the indications, contraindications and side effects of any prospective drug or intervention. Furthermore, the data for the book are largely derived from incidence and prevalence statistics whose accuracy will vary widely for individual diseases and countries. Changes in endemicity, incidence, and drugs of choice may occur. The list of drugs, infectious diseases and even country names will vary with time.

Scope of Content:

Disease designations may reflect a specific pathogen (ie, Adenovirus infection), generic pathology (Pneumonia - bacterial) or etiologic grouping (Coltivirus - Old world). Such classification reflects the clinical approach to disease allocation in the Infectious Diseases Module of the GIDEON web application. Similarly, a number of diseases which are generally diagnosed and treated outside of the field of Infectious Diseases are not included, despite the fact that a clear infectious etiology exists. Examples include Peptic ulcer, Tropical spastic paraparesis, Hairy-cell leukemia, Creutzfeldt-Jakob disease, Human papilloma virus infections, etc. In contrast, a number of other entities of unknown etiology which do present to Infectious Diseases specialists have been included: Kawasaki's disease, Chronic fatigue syndrome, Kikuchi and Kimura diseases. Several minor infections having minimal relevance to the field of Geographic Medicine are not covered: Paronychia, Otitis externa, Molluscum contagiosum, etc.

Introduction: The GIDEON e-book series

Infectious Diseases of Uganda is one in a series of GIDEON [ebooks](#) which summarize the status of individual infectious diseases, in every country of the world. Data are based on the GIDEON web application (www.gideononline.com) which relies on standard text books, peer-review journals, Health Ministry reports and ProMED, supplemented by an ongoing search of the medical literature.

Chapters are arranged alphabetically, by disease name. Each section is divided into four sub-sections:

1. Descriptive epidemiology
2. Summary of clinical features
3. Status of the disease in Uganda
4. References

The initial items in the first section, Descriptive epidemiology, are defined as follows:

- Agent** Classification (e.g., virus, parasite) and taxonomic designation.
- Reservoir** Any animal, arthropod, plant, soil or substance in which an infectious agent normally lives and multiplies, on which it depends primarily for survival, and where it reproduces itself in such a manner that it can be transmitted to a susceptible host.
- Vector** An arthropod or other living carrier which transports an infectious agent from an infected organism or reservoir to a susceptible individual or immediate surroundings.
- Vehicle** The mode of transmission for an infectious agent. This generally implies a passive and inanimate (i.e., non-vector) mode.

There are 348 generic infectious diseases in the world today. 236 of these are endemic, or potentially endemic, to Uganda. A number of other diseases are not relevant to Uganda and have not been included in this book.

In addition to endemic diseases, we have included all published data regarding imported diseases and infection among expatriates from Uganda.

The availability and quality of literature regarding specific infectious diseases vary from country to country. As such, you may find that many of the sections in this book are limited to a general discussion of the disease itself - with no data regarding Uganda.

This is a book about the geography and epidemiology of Infection. Comprehensive and up-to-date information regarding the causes, diagnosis and treatment of each disease is available in the [GIDEON web application](#). Many of the diseases are generic. For example, such designations as Pneumonia bacterial and Urinary tract infection include a number of individual diseases. These appear under the subheading, Synonyms, listed under each disease.

We welcome feedback, and will be pleased to add any relevant, sourced material. Email us at ebook@gideononline.com

For more information about GIDEON see the section [About GIDEON](#) and visit www.gideononline.com

Last updated: February 19, 2013

Table of Contents

Introduction: The GIDEON e-book series	2	Dengue*	88
Actinomycosis	6	Dermatophytosis	90
Adenovirus infection ⁺	8	Dientamoeba fragilis infection	91
Aeromonas & marine Vibrio infx. ⁺	10	Diphtheria ⁺	92
AIDS ⁺	12	Diphyllobothriasis	95
Amoeba - free living	18	Dipylidiasis	96
Amoebic abscess ⁺	20	Dirofilariasis	97
Amoebic colitis ⁺	21	Dracunculiasis ⁺	98
Animal bite-associated infection	23	Ebola ⁺	101
Anisakiasis	24	Echinococcosis - unilocular ⁺	103
Anthrax ⁺	25	Endocarditis - infectious	105
Ascariasis ⁺	29	Enteritis necroticans	106
Aspergillosis	31	Enterobiasis ⁺	107
Bacillary angiomatosis	32	Enterovirus infection ⁺	108
Bacillus cereus food poisoning	33	Entomophthoromycosis	110
Bacterial vaginosis ⁺	34	Epidural abscess	111
Balantidiasis	36	Erysipelas or cellulitis	112
Bartonellosis - cat borne	37	Erysipeloid	113
Bartonellosis - other systemic	39	Erythrasma	114
Bertiella and Inermicapsifer*	40	Escherichia coli diarrhea	115
Blastocystis hominis infection	41	Fascioliasis ⁺	117
Blastomycosis ⁺	42	Filariasis - Bancroftian ⁺	119
Botulism ⁺	44	Fungal infection - invasive	122
Brain abscess	46	Gastroenteritis - viral	124
Brucellosis ⁺	47	Gianotti-Crosti syndrome	125
Bunyaviridae infections - misc. ⁺	51	Giardiasis ⁺	126
Campylobacteriosis ⁺	53	Gnathostomiasis*	128
Candidiasis ⁺	55	Gonococcal infection ⁺	129
Chancroid ⁺	56	Granuloma inguinale	131
Chikungunya ⁺	57	Hantavirus infection - Old World	132
Chlamydia infections, misc. ⁺	59	Hepatitis A	134
Chlamydia pneumoniae infection	61	Hepatitis B ⁺	136
Cholecystitis & cholangitis	62	Hepatitis C ⁺	139
Cholera ⁺	63	Hepatitis D ⁺	142
Chromomycosis	67	Hepatitis E ⁺	144
Chronic meningococemia	68	Hepatitis G ⁺	146
Clostridial food poisoning	69	Herpes B infection	147
Clostridial myonecrosis	70	Herpes simplex encephalitis	148
Clostridium difficile colitis	71	Herpes simplex infection ⁺	149
Coenurosis ⁺	72	Herpes zoster	151
Common cold	73	Histoplasmosis*	153
Conjunctivitis - inclusion	74	Histoplasmosis - African ⁺	155
Conjunctivitis - viral ⁺	75	HIV infection - initial illness ⁺	157
Crimean-Congo hemorrhagic fever ⁺	76	Hookworm ⁺	158
Cryptococcosis ⁺	78	Human herpesvirus 6 infection	160
Cryptosporidiosis ⁺	80	Hymenolepis diminuta infection	161
Cutaneous larva migrans	82	Hymenolepis nana infection ⁺	162
Cyclosporiasis	83	Infection of wound, puncture, IV line, etc	163
Cysticercosis ⁺	84	Infectious mononucleosis or EBV infection	164
Cytomegalovirus infection	86	Influenza ⁺	166
		Intestinal spirochetosis	178

Intra-abdominal abscess	179	Pediculosis	268
Intracranial venous thrombosis	180	Pentastomiasis - Linguatula	269
Isosporiasis ⁺	181	Pericarditis - bacterial	270
Kawasaki disease	182	Perinephric abscess	271
Kikuchi's disease and Kimura disease	184	Perirectal abscess.....	272
Kingella infection.....	186	Peritonitis - bacterial.....	273
Laryngotracheobronchitis.....	187	Pertussis ⁺	274
Lassa fever.....	188	Pharyngeal & cervical space infx.....	277
Legionellosis	190	Pharyngitis - bacterial	278
Leishmaniasis - cutaneous ⁺	192	Pityriasis rosea.....	279
Leishmaniasis - visceral ⁺	195	Plague ⁺	280
Leprosy ⁺	197	Plesiomonas infection.....	283
Leptospirosis ⁺	201	Pleurodynia	284
Listeriosis ⁺	203	Pneumocystis pneumonia	285
Liver abscess - bacterial	205	Pneumonia - bacterial	286
Loiasis ⁺	206	Poliomyelitis ⁺	287
Lymphocytic choriomeningitis.....	207	Protothecosis and chlorellosis.....	291
Lymphogranuloma venereum	208	Pseudocowpox	292
Malaria ⁺	210	Pyodermas (impetigo, abscess, etc).....	293
Malignant otitis externa.....	215	Pyomyositis	295
Mansonelliasis - M. perstans ⁺	216	Q-fever.....	296
Mansonelliasis - M. streptocerca ⁺	218	Rabies ⁺	298
Marburg virus disease ⁺	219	Rat bite fever - spirillary.....	302
Measles ⁺	221	Rat bite fever - streptobacillary	303
Melioidosis.....	224	Relapsing fever ⁺	304
Meningitis - aseptic (viral)	226	Respiratory syncytial virus infection	306
Meningitis - bacterial ⁺	227	Respiratory viruses - miscellaneous	308
Microsporidiosis ⁺	232	Reye's syndrome.....	310
Moniliformis and Macracanthorhynchus.....	233	Rheumatic fever.....	311
Mumps.....	234	Rhinocleroma and ozena ⁺	312
Myalgic encephalomyelitis.....	236	Rhinospordiosis ⁺	313
Mycetoma	238	Rhodococcus equi infection	314
Mycobacteriosis - M. marinum	240	Rift Valley fever	315
Mycobacteriosis - M. scrofulaceum	241	Rotavirus infection ⁺	316
Mycobacteriosis - M. ulcerans ⁺	242	Rubella ⁺	318
Mycobacteriosis - miscellaneous nontuberculous ⁺	244	Salmonellosis ⁺	320
Mycoplasma (miscellaneous) infections ⁺	246	Sarcocystosis.....	322
Mycoplasma pneumoniae infection	248	Scabies.....	323
Myiasis.....	250	Scarlet fever.....	324
Necrotizing skin/soft tissue infx.	251	Schistosomiasis - haematobium ⁺	325
Neutropenic typhlitis	253	Schistosomiasis - intercalatum ⁺	328
Nocardiosis.....	254	Schistosomiasis - mansoni ⁺	330
O'nyong nyong ⁺	255	Septic arthritis	333
Oesophagostomiasis ⁺	256	Septicemia - bacterial	334
Onchocerciasis ⁺	257	Shigellosis ⁺	335
Orbital and eye infections	260	Sindbis ⁺	337
Orf ⁺	261	Sinusitis	338
Ornithosis.....	262	Smallpox [*]	339
Osteomyelitis ⁺	263	Sparganosis.....	341
Otitis media.....	264	Spondweni ⁺	342
Parainfluenza virus infection.....	265	Sporotrichosis.....	343
Parvovirus B19 infection	266	Spotted fevers - Old World ⁺	345

Staphylococcal food poisoning	347	Tropical pulmonary eosinophilia.....	378
Staphylococcal scalded skin syndrome.....	348	Trypanosomiasis - African ⁺	379
Streptococcus suis infection	349	Tuberculosis ⁺	383
Strongyloidiasis ⁺	350	Tungiasis ⁺	387
Subdural empyema.....	352	Typhoid and enteric fever ⁺	389
Suppurative parotitis.....	353	Typhus - endemic.....	392
Syphilis ⁺	354	Typhus - epidemic ⁺	393
Taeniasis ⁺	356	Urinary tract infection	395
Tetanus ⁺	357	Varicella ⁺	397
Thelaziasis.....	362	Vibrio parahaemolyticus infection	399
Thogoto ⁺	363	Wesselsbron ⁺	400
Toxic shock syndrome	364	West Nile fever ⁺	401
Toxocariasis.....	365	Whipple's disease	403
Toxoplasmosis ⁺	367	Yaws ⁺	405
Trachoma ⁺	369	Yellow fever ⁺	407
Trichinosis ⁺	370	Yersiniosis	410
Trichomoniasis ⁺	372	Zika ⁺	412
Trichostrongyliasis [*]	374	Zygomycosis	413
Trichuriasis ⁺	375	About GIDEON	415
Tropical phagedenic ulcer	377		

* Not endemic. Imported, expatriate or other context reported.

⁺ Country specific note exists for disease

Actinomycosis

Agent	BACTERIUM. Actinomycetes, Actinomyces spp. An anaerobic gram-positive bacillus
Reservoir	Human - oral, fecal, vaginal flora
Vector	None
Vehicle	Endogenous
Incubation Period	Unknown
Diagnostic Tests	Gram stain and bacteriological culture using strict anaerobic technique. Growth is apparent in 3-7 days.
Typical Adult Therapy	Ampicillin 50 mg/kg/day IV X 4 to 6 weeks - then Amoxicillin 1.5 g/d PO X 6 months. OR Penicillin G 10 to 20 million units/day X 4 to 6w; then Penicillin V X 6 to 12m. Alternatives: Doxycycline , ceftriaxone , Erythromycin Excision/drainage
Typical Pediatric Therapy	Ampicillin 50 mg/kg/day IV X 4 to 6 weeks - then Amoxicillin 20 mg/kg/day PO X 6 months. Penicillin G 100,000 units/kg/day X 4 to 6w; then Penicillin V 25,000 units/day X 6 to 12m. Excision/drainage
Clinical Hints	Mandibular osteomyelitis with fistulae (sulfur granules) in the setting of poor dental hygiene [oral actinomycosis]; intrauterine device and pelvic abscesses [pelvic actinomycosis]; fever, right lower quadrant mass and fistulae [abdominal actinomycosis].
Synonyms	Actinomyces, Aktinomykose, Lumpy jaw. ICD9: 039. ICD10: A42

Clinical

Anatomic variants of Actinomycosis

Oral-cervical actinomycosis accounts for 55% of actinomycosis, and may be manifested as soft tissue swelling, an abscess, or a mass lesion. ¹

- Lesions may be multiple, and relapse following short courses of therapy.
- The disease often spreads to adjacent structures (masseter muscle, carotid artery, cranium, cervical spine, trachea, or thorax) without regard for normal tissue planes.
- Lymphatic spread and lymphadenopathy are rare.
- Infection is associated with pain, fever, and leukocytosis.

Periapical actinomycosis ² is common and responds to dental care and antibiotics.

- The most common location for actinomycosis is the perimandibular region.
- Periapical infection often precedes infection, which is usually seen at the angle of the jaw; however, the cheek, submental space, retromandibular space, and temporomandibular joint may be affected.
- The overlying skin is often blue to red-purple in color, and sinuses may appear.
- An abscess may ensue, with trismus.
- Mandibular periostitis and osteomyelitis are rarely encountered.
- Maxillary or ethmoid disease, with or without osteomyelitis, is uncommon; but maxillary sinusitis and associated cutaneous fistulas can occur.
- Masses of the hard palate, tongue, nasal septum, head and neck, salivary glands, thyroglossal ducts, thyroid, branchial cleft cysts, lacrimal ducts, orbital structures and larynx have also been reported.
- The tonsils are rarely, if ever, involved; however, infection of the external or middle ear, temporal bone and mastoid may occur following spread of facial disease.

Thoracic actinomycosis ³ accounts for 15% of actinomycosis cases, and represents aspiration of organisms from the pharynx (rarely direct extension from the head and neck or abdominal cavity).

- Most cases present as an indolent, slowly progressive process involving the lung parenchyma and pleura.
- Chest pain, fever, and weight loss are common; occasionally with hemoptysis and a productive cough.
- X-ray findings are non-specific.
- The usual appearance is either a mass lesion or pneumonitis with or without pleural involvement.
- An air bronchogram within a mass lesion is suggestive when present, pleural thickening, effusion, or empyema is seen in more than 50% of cases.
- An isolated pleural effusion may drain spontaneously through the chest wall or produce a soft tissue or breast mass; or posteriorly, to involve the vertebrae or paraspinal structures or spinal cord
- Pulmonary disease may extend across fissures or pleura, and involve the mediastinum, pericardium (rarely endocardium)

or contiguous bone.

Abdominal actinomycosis ⁴ accounts for 20% of actinomycosis and represents ingestion of bacteria, hematogenous infection or extension from the female pelvis.

- Associated fever, weight loss, abdominal pain or fullness and changing bowel habits may be present for months before the diagnosis is suspected.
- Physical findings include mass lesions and sinus tracts of the abdominal wall.
- Lymphadenopathy is uncommon.
- 65% of cases are associated with appendicitis, and 65% of lesions present in the right iliac fossa.
- Associated tuboovarian infection, hepatic abscesses ⁵, diverticulitis or foreign body perforation in the transverse or sigmoid colon may also be encountered.
- Other associated factors include previous gastric or bowel surgery, typhoid fever, amebic dysentery, trauma, and pancreatitis.
- Abdominal infection may extend to the liver hematogenously; and perirectal or perianal infection is occasionally encountered, resulting in chronic fistulae, sinuses and strictures.

Pelvic actinomycosis ⁶ may represent spread from intra-abdominal infection; but is most often a complication of intra-uterine device (IUD) placement.

- Any type of IUD can cause infection; and on average, the device has been in place for eight years prior to the appearance of actinomycosis.
- Infection may even occur months following removal of the device.
- Infection is manifest as endometritis or a mass/abscess of the tubes or ovaries. ⁷
- Presenting features consist of chronic fever, weight loss, abdominal pain, and vaginal bleeding.
- A "frozen pelvis" suggestive of malignancy or endometriosis is often encountered; and the infection may involve the ureters, bladder, rectum, small or large bowel or peritoneum.
- The diagnostic value of smears and cultures for Actinomyces among asymptomatic women with IUD's is controversial.

Other forms of actinomycosis include:

- brain abscess
- chronic meningitis
- urogenital infection
- musculoskeletal infection
- isolated skin ⁸ and muscle disease (including mycetoma)
- infected orthopedic prostheses
- thyroiditis
- disseminated hematogenous infection of multiple organs

This disease is endemic or potentially endemic to all countries.

References

1. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 1998 May ;85(5):496-508.
 2. Quintessence Int 2005 Feb ;36(2):149-53.
 3. Chest 1993 Aug ;104(2):366-70.
 4. Clin Microbiol Infect 2003 Aug ;9(8):881-5.

5. Orv Hetil 2011 Feb 13;152(7):268-72.
 6. Am J Obstet Gynecol 1999 Feb ;180(2 Pt 1):265-9.
 7. Ned Tijdschr Geneesk 2012 ;156(16):A4511.
 8. Int J Dermatol 2008 Dec ;47(12):1271-3.

Adenovirus infection

Agent	VIRUS - DNA. Adenoviridae, Adenovirus Enteric strains classified in genus Mastadenovirus
Reservoir	Human Non-human primates
Vector	None
Vehicle	Droplet Water
Incubation Period	4d - 12d
Diagnostic Tests	Viral culture/serology or antigen assay. Direct fluorescence of secretions. Nucleic acid amplification.
Typical Adult Therapy	Enteric/secretion precautions. Cidofovir has been used in some cases. Symptomatic therapy
Typical Pediatric Therapy	As for adult
Vaccine	Adenovirus
Clinical Hints	Atypical pneumonia, upper respiratory infection, tracheitis, bronchiolitis or keratoconjunctivitis with preauricular adenopathy; uncomplicated illness usually lasts 3 to 5 days; this agent may also cause hemorrhagic cystitis.
Synonyms	Adenovirus gastroenteritis, Epidemic keratoconjunctivitis, Pharyngoconjunctival fever. ICD9: 047.9,077.1,077.2,008.62,480.0 ICD10: A08.2,B30.1,B34.0,J12.0

Clinical

Only 50% of Adenovirus infections are clinically apparent.

- Infection in children usually presents as mild pharyngitis or tracheitis.
- Adenovirus type 7 can cause fulminant bronchiolitis and pneumonia in infants.
- Severe respiratory infection is associated with serotype 14 ¹
- Adenoviruses have been isolated more often than any other nonbacterial pathogen from patients with the whooping cough syndrome; however, a causal relation has not been established.

Cough, fever, sore throat, tonsillitis ² and rhinorrhea are the most common findings ³, and usually last 3 to 5 days. ^{4 5}

- Rales and rhonchi may be present.
- X-ray studies in patients with pneumonias reveal patchy ground-glass infiltrates primarily in the lower lung fields.
- Outbreaks among military personnel are characterized by tracheobronchitis, with 20% requiring hospitalization.
- The disease is usually self-limited, superinfection and death are rare.
- Severe infections are increasingly reported among immunocompromised patients. ⁶⁻⁸
- There are also case reports of severe Adenovirus pneumonia in immunocompetent adults. ⁹
- Rare instances of fatal Adenovirus myocarditis have been reported. ^{10 11}
- In one series of 3,298 adenoviral infections, pneumonia was found in only 2.4%. ¹²
- Adenoviral pneumonia is often followed by bronchiolitis obliterans in children. ^{13 14}

Pharyngoconjunctival fever:

Pharyngoconjunctival fever often occurs in the setting of small outbreaks.

- Illness is characterized by conjunctivitis, pharyngitis, rhinitis, cervical lymphadenitis, and fever to 38 C.
- The onset is acute, and symptoms last 3 to 5 days.
- Bulbar and palpebral conjunctivitis, usually bilateral, may be the only finding.
- The palpebral conjunctivae have a granular appearance.
- Bacterial superinfection and permanent residua are unusual.
- Respiratory involvement usually does not progress to the bronchi or lungs.
- Contaminated swimming pools and ponds have been implicated as sources of spread.

Epidemic keratoconjunctivitis:

Epidemic keratoconjunctivitis has an incubation period of 4 to 24 days, and lasts for 1 to 4 weeks.

- The conjunctivitis is often bilateral, and preauricular adenopathy is common. ¹⁵⁻¹⁷
- Multiple subepithelial corneal infiltrates are often present. ¹⁸
- Visual disturbance may persist for several months.
- Secondary spread to household contacts occurs in 10% of the cases.

Hemorrhagic cystitis:

Hemorrhagic cystitis is two to three times more common in boys than girls (unlike bacterial cystitis which is predominantly seen in girls). ¹⁹

- Hematuria usually persists for approximately three days.
- There was no seasonal preponderance.
- Adenoviral urethritis is also reported. ²⁰

Infantile adenoviral enteritis:

Infantile adenoviral enteritis is characterized by watery diarrhea is watery with fever, and may last for 1 to 2 weeks.

- Adenoviruses have also been implicated in the etiology of intussusception, encephalitis and meningoencephalitis. ²¹
- Rare instances of intestinal intussusception have been associated with adenoviral gastroenteritis. ²²

Other forms of infection:

Adenoviruses have emerged as important pathogens in immunosuppressed patients, particularly those undergoing bone marrow or solid organ transplantation.

- Syndromes include infection of the transplanted organ, or disseminated infection involving the lung, colon (ie, chronic diarrhea ²³), and central nervous system.
- Infection, notably of the urinary ²⁴ and gastrointestinal tracts, is also a common complication of AIDS.
- Adenoviral parotitis and encephalitis are also reported in AIDS patients.

This disease is endemic or potentially endemic to all countries.

Adenovirus infection in Uganda**Prevalence surveys:**

4% of respiratory disease among young children (1977 publication) ²⁵

References

1. Clin Infect Dis 2008 Feb 1;46(3):421-5.
2. Pediatr Infect Dis J 2005 Aug ;24(8):733-4.
3. Rev Med Virol 2008 Nov-Dec;18(6):357-74.
4. Pediatrics 2004 Jan ;113(1 Pt 1):e51-6.
5. Br Med Bull 2002 ;61:247-62.
6. Br J Haematol 2005 Jan ;128(2):135-44.
7. Pediatr Blood Cancer 2008 Mar ;50(3):647-9.
8. Curr Opin Organ Transplant 2009 Dec ;14(6):625-33.
9. Eur J Clin Microbiol Infect Dis 2008 Feb ;27(2):153-8.
10. J Med Virol 2008 Oct ;80(10):1756-61.
11. J Clin Microbiol 2010 Feb ;48(2):642-5.
12. J Clin Virol 2012 Sep 25;
13. Zhonghua Er Ke Za Zhi 2008 Oct ;46(10):732-8.
14. Pediatr Pulmonol 2009 May ;44(5):450-6.
15. Prog Retin Eye Res 2000 Jan ;19(1):69-85.
16. Rev Med Virol 1998 Oct ;8(4):187-201.
17. Postgrad Med 1997 May ;101(5):185-6, 189-92, 195-6.
18. Jpn J Ophthalmol 2011 Jan ;55(1):11-5.
19. Arch Dis Child 2005 Mar ;90(3):305-6.
20. Sex Health 2007 Mar ;4(1):41-4.
21. J Neurovirol 2006 Jun ;12(3):235-40.
22. Infez Med 2010 Dec ;18(4):256-8.
23. Pediatr Infect Dis J 2008 Apr ;27(4):360-2.
24. Am J Kidney Dis 2008 Jan ;51(1):121-6.
25. Bull World Health Organ 1977 ;55(5):625-31.

Aeromonas & marine Vibrio infx.

Agent	BACTERIUM. <i>Aeromonas hydrophila</i> & <i>Vibrio vulnificus</i> , et al Facultative gram-negative bacilli
Reservoir	Salt or brackish water Fish
Vector	None
Vehicle	Water/shellfish - contact or ingestion
Incubation Period	Range 2d - 7d
Diagnostic Tests	Culture. Notify laboratory if these organisms are suspected in stool.
Typical Adult Therapy	Fluoroquinolone or <i>Sulfamethoxazole/trimethoprim</i> . Other antimicrobial agent as determined by susceptibility testing
Typical Pediatric Therapy	<i>Sulfamethoxazole/trimethoprim</i> . Or other antimicrobial agent as determined by susceptibility testing
Clinical Hints	Diarrhea, fever, vomiting or sepsis after marine injury or ingestion of raw oysters/contaminated fresh or brackish water; fecal leukocytes present; severe or fatal in immunosuppressed or alcoholic patients.
Synonyms	<i>Aeromonas</i> , <i>Aeromonas hydrophila</i> , <i>Vibrio mimicus</i> , <i>Vibrio vulnificus</i> . ICD9: 005.81,027.9 ICD10: A48.8

Clinical

***Aeromonas hydrophila* gastroenteritis:**

There is controversy as to whether *Aeromonas hydrophila* can cause gastroenteritis.

- Volunteer feeding studies using as many as 1 billion cells have failed to elicit illness.
- The presence of this species in the stools of individuals with diarrhea, in the absence of other known enteric pathogens, suggests that it has some role in disease. ¹
- *Aeromonas* species are often implicated in traumatic and surgical wound sepsis ^{2 3} and a variety of localized infections. ⁴⁻⁸
- *Aeromonas caviae* and *A. sobria* are considered by many as "putative pathogens," in diarrheal disease.

Two types of gastroenteritis have been associated with *A. hydrophila* ⁹ :

- a cholera-like illness with a watery diarrhea
- a dysenteric illness characterized by loose stools containing blood and mucus.
- cases of hemolytic uremic syndrome have followed *Aeromonas* infection ¹⁰

Generalized systemic infection has been observed in individuals with underlying illness.

***Vibrio vulnificus*:**

Vibrio vulnificus causes septicemia in persons with chronic liver disease, alcoholism or hemochromatosis, and immunosuppressed patients. ^{11 12}

- The disease appears 12 hours to 3 days after eating raw or undercooked seafood, especially oysters.
- One third of the patients are in shock within 12 hours after hospital admission.
- Three quarters have distinctive, bullous skin lesions which may be mistaken for pemphigus or pemphigoid.
- Thrombocytopenia is common and there is often evidence of disseminated intravascular coagulation.
- Over 50 percent of patients with septicemia die; and the mortality rate exceeds 90 percent among those with hypotension.

Relatively high mortality rates are associated with necrotizing fasciitis caused by *Aeromonas* or *Vibrio* species. ¹³

V. vulnificus can also infect wounds sustained in coastal or estuarine waters.

- Infections range from mild self limited lesions to rapidly progressive cellulitis or myositis that can mimic clostridial myonecrosis clinically.

Additional species of *Aeromonas* and *Vibrio* are described in the Microbiology module.

This disease is endemic or potentially endemic to all countries.

Aeromonas & marine Vibrio infx. in Uganda

Prevalence surveys:

7% of HIV-positive patients with diarrhea in rural Uganda (*Aeromonas* species, 2010 publication)

References

1. Infection 2007 Apr ;35(2):59-64.
2. Scand J Infect Dis 2009 ;41(3):164-70.
3. Surg Infect (Larchmt) 2011 Jun ;12(3):241-5.
4. Clin Nephrol 2011 Feb ;75 Suppl 1:65-8.
5. Mil Med 2011 Dec ;176(12):1444-6.
6. Neurocirugia (Astur) 2012 Sep ;23(5):200-2.
7. Eur J Clin Microbiol Infect Dis 2012 Aug 24;
8. Am J Trop Med Hyg 2012 Nov ;87(5):933-5.
9. Crit Rev Microbiol 2002 ;28(4):371-409.
10. Diagn Microbiol Infect Dis 2007 Jun ;58(2):231-4.
11. South Med J 2004 Feb ;97(2):163-8.
12. Eur J Clin Microbiol Infect Dis 2007 Nov ;26(11):785-92.
13. Am J Emerg Med 2008 Feb ;26(2):170-5.

AIDS

Agent	VIRUS - RNA. Retroviridae, Lentivirinae: Human Immunodeficiency Virus, HIV
Reservoir	Human
Vector	None
Vehicle	Blood Semen Sexual Transplacental Breast-feeding
Incubation Period	2m - 10y (50% within 10y)
Diagnostic Tests	HIV antibody (ELISA, Western blot). Nucleic acid amplification. Tests for HIV antigen & viral load as indicated.
Typical Adult Therapy	Two nucleosides + 1 protease inhibitor; or two nucleosides + 1 non-nucleoside; or 2 nucleosides + Ritonavir (alone or with lopinavir) + (indinavir, amprenavir, saquinavir or nelfinavir)
Typical Pediatric Therapy	As for adult
Clinical Hints	Most often associated with drug abuse, blood products, men who have sex with men, hemophilia. Hints: severe herpes simplex or moniliasis, chronic cough, diarrhea, weight loss, lymphadenopathy, retinitis, encephalitis or Kaposi's sarcoma.
Synonyms	ARC, Gay cancer, GRID, HIV-1, HIV-2, HIV-AIDS, SIDA, Slim disease. ICD9: 042 ICD10: B20,B21,B22,B23,B24

Clinical

CDC case surveillance definition:

As of 1993, the CDC (The United States Centers for Disease Control) surveillance case definition for AIDS includes all HIV-infected persons age 13 or over who have either. ¹

- a) a <200 CD4+ T-lymphocytes
- b) a CD4+ T-lymphocyte percentage of total lymphocytes of <14%
- or c) any of the following: pulmonary tuberculosis, recurrent pneumonia, or invasive cervical cancer; or any of the 23 clinical conditions defined in the case definition published in 1987. ²
- For WHO case definition (1994) see reference ³

The clinical features of AIDS are protean and often characterized by multisystem illness, evidence of immune suppression and the presence of one or more superinfections (tuberculosis ⁴, Cytomegalovirus infection, cerebral toxoplasmosis ⁵, pneumocystosis ^{6 7}, penicilliosis ^{8 9}, severe or recalcitrant candidiasis, disseminated *Acanthamoeba* infection ¹⁰, etc).

Acute HIV infection is characterized by fever, generalized lymphadenopathy, headache, fatigue, myalgia, rash, nausea, vomiting, night sweats, sore throat, diarrhea or weight loss. ¹¹

- 40% to 90% of persons have symptoms suggestive of an acute viral infection.
- Symptoms tend to subside within two weeks; however, some patients continue to be ill for as long as ten weeks.
- In most cases, a history of likely acquisition within the past several weeks can be established: unprotected sex, extra-medical injection, transfusion, etc.

HIV infection and opportunistic pathogens:

HIV infection increases the incidence and severity of a wide variety of infectious diseases ¹² caused by viruses, mycobacteria, actinomycetes, treponemes, fungi ¹³⁻¹⁷, protozoa and helminths.


- HIV infection increases the incidence and severity of clinical malaria; however, in severe malaria the level of parasitemia is similar in HIV-positive and HIV-negative patients. ¹⁸⁻²⁵
- During pregnancy, HIV infection increases the incidence of clinical malaria, maternal morbidity, and fetal and neonatal morbi-mortality.
- HIV infection increases severity of malaria, the risk of malaria treatment failure, and for cerebral malaria in children. ^{26 27}
- Some antimalarial drugs may inhibit HIV, while certain anti-retroviral drugs are effective against *Plasmodium* species. ²⁸
- Reactivation of Chagas disease encephalopathy has been reported among infected HIV-positive patients. ²⁹
- Acquired syphilis in patients with HIV infection is characterized by severe and accelerated infection, often with overt meningitis, hepatitis ³⁰ and other forms of systemic involvement. ³¹⁻³⁸ The presence of concurrent syphilis does not affect the progression of AIDS. ³⁹
- *Haemophilus ducreyi* has been associated with esophageal ulceration in HIV-positive patients. ⁴⁰

- Hepatitis G infection appears to improve survival among persons with concurrent HIV infection. ⁴¹ 41% of infants born to mothers with HIV-HGB-C coinfection acquired HGB-C infection (Thailand, 2009 publication) ⁴²
- Concurrent HIV infection increases the incidence of cirrhosis and HCC among Hepatitis B carriers ^{43 44} ; and shortens the time to development of chronic liver disease in patients with Hepatitis C. ⁴⁵
- HIV-HCV coinfection is characterized by more rapid progression to cirrhosis and diminished response to peginterferon/ribavirin therapy. ⁴⁶⁻⁵⁰
- Hepatitis D is associated with relatively aggressive disease among patients with HIV-HBV coinfection. ⁵¹
- Concurrent HIV infection may prolong the duration of viremia in patients with hepatitis A. ⁵²
- Lesions of Herpes simplex in HIV-positive patients may be vegetative, hypertrophic, condyloma-like, nodular, ulcerative, or tumor-like nodules or plaques. ⁵³

This disease is endemic or potentially endemic to all countries.

AIDS in Uganda

The first cases of AIDS were reported in 1983.




Graph: Uganda. AIDS, cases

Notes:

1. See reference ⁵⁴

Individual years:


2005 - The true incidence was estimated at 135,000 new HIV infections, 88,100 AIDS cases and 76,400 AIDS deaths. ⁵⁵



Graph: Uganda. AIDS, cumulative cases

Notes:

1. 47% of cases to December 1997 were in males.
2. The true number of AIDS cases to December 1997 is estimated at 1,900,000, with 1,800,000 AIDS deaths.




Graph: Uganda. AIDS, estimated deaths

Notes:

1. In 1994, over 80% of deaths for the age group 25 to 44 were attributable to HIV infection.
2. 1,700,000 AIDS orphans were estimated to December 1999; 880,000 in 2001.
3. An estimated 1,000,000 AIDS orphans were alive in 2003; 1,000,000 in 2005.

Seroprevalence surveys:

- 12% (rural) to 33% (urban) of the general population
 - 4.1% of the general population in 2003
 - 8.2% of the population (20.9% of females ages 20 to 24) in Masaka (rural southwest, 1989 to 1990)
 - 7.8% of persons in a rural parish (2011) ⁵⁶
 - 6.9% of the general population (13.8% of females ages 20 to 24) during 1996 to 1997
 - 2.3% of males and 4.8% of females ages 15 to 24 (2006) ⁵⁷
 - 0.7% of children ages 2 to 12 years (rural southwest, 2004 to 2005) ⁵⁸
 - 5.9% of persons in a national serosurvey, and 8.8% of persons undergoing voluntary counseling and testing, ages 15 to 49 (2004 to 2005) ⁵⁹
 - 25% of hospital patients tested for the first time (Mulago and Mbarara, 2004 to 2006) ⁶⁰
 - 14% of adults attending a health program in Rakai, southwestern Uganda (2011 publication) ⁶¹
 - 28.8% of persons in fishing communities on Lake Victoria (2005) ⁶²
 - 50% of emergency medical patients (Kampala, 2004) ⁶³
 - 42% of inpatients
 - 16% of blood donors
 - 2.2% of blood donors in central Uganda (1999) ⁶⁴
 - 8.7% of pregnant women, nationwide (2003 to 2004) ⁶⁵
 - 12.6% of pregnant women and 10.8% of their male partners (Entebbe, 2002 to 2006) ⁶⁶
 - 11.9% of pregnant women in Entebbe (2004 to 2005) ⁶⁷
 - 15.3% of pregnant women attending antenatal care clinics in western Uganda (1991 to 2004) ⁶⁸
 - 29.5% of urban pregnant women in 1992; 14.7% in 1997; 13.8% in 1998; 11.25% in 2000; 5.2% in 2005
 - 8.75% of rural pregnant women in 1997; 7.7% in 1998
 - 12.1% of pregnant women in Gulu district in 2000, 11.3% in 2003 ⁶⁹
 - 11.9% of otherwise healthy pregnant women at enrolment to a trial of deworming. (Entebbe, 2007 publication) ⁷⁰
 - 23.4% of adults in Rakai in 1990, and 20.9% in 1992
 - 26.5% of military personnel (1996)
 - 33.9% of urban STD patients in 1995; 29.4% in 1998 (males)
 - 7.9% of adults in Tanga region (2005 publication) ⁷¹
 - 4.9% of persons age 15 to 54 in the rural southwest (1999 publication) ⁷²
 - 20% of patients with primary hepatocellular carcinoma (Kampala, 2008) ⁷³
 - 34.4% of malnourished children hospitalized with diarrhea (Kampala, 2002 to 2003)
 - 30.0% of hospitalized children suffering from severe acute malnutrition (2006) ⁷⁴
 - 9% of children with cerebral malaria, vs. 2.3% of children with uncomplicated malaria (2011 publication) ⁷⁵
- The rate of HIV acquisition among young women was 2.7 per 100 woman-years during 1999 to 2004 ⁷⁶ ; and 4.12 per 100 woman-years in a 2008 publication. ⁷⁷
- The rate of HIV infection among men in Rakai is 1.09 per 100 person-years (2009 publication) ⁷⁸



Graph: Uganda. AIDS - estimated living with HIV/AIDS, cases

Notes:

- Figure for 1997 represented 9.51% of all adults; 5.0% in 2001; 4.1% in 2003; 6.7% in 2005.

Associated infections:

- The incidence of tuberculosis among persons living with HIV/AIDS is 3.4% per year (1997). 58% of tuberculosis patients are HIV-positive (1989 publication)⁷⁹; 50% in 1997. 4.6% of newly-diagnosed HIV-positive individuals have tuberculosis (Kampala, 2006 publication)⁸⁰. *M. tuberculosis* bacteremia was identified in 12.3% of febrile HIV-positive adults, and *Salmonella* bacteremia in 5.7% (Kampala, 2007)⁸¹.
- Cryptococcosis occurred in 77 of 1,372 HIV-positive patients followed in a cohort study in Entebbe during 1995 to 1999⁸². The rate of cryptococcosis among HIV-positive adults was 40.4/1,000 person-years, and the fungus accounted for 17% of all HIV-associated death. Pulmonary cryptococcosis was identified in 2.7% of hospitalized HIV-positive patients (Kampala, 2007 to 2008)⁸³.
- A case of African histoplasmosis was confirmed in Japan in a Ugandan national with AIDS.⁸⁴
- 24.5% (HIV-negative) to 64.0% (HIV-positive) persons in Kitgum are seropositive for syphilis (1986).⁸⁵
- 34% of HIV-positive patients, and 27% of controls are seropositive toward *Toxoplasma gondii*.⁸⁶
- In a separate study, 54% of HIV-positive patients were found to be seropositive.⁸⁷
- 9% of children with cerebral malaria were found to be HIV-positive, vs. 2.3% of children with uncomplicated malaria (2011 publication)⁸⁸. Asymptomatic *Plasmodium falciparum* antigenemia was found in 4% of urban AIDS patients receiving HAART (2004 to 2005)⁸⁹.
- 9.7% of HIV-positive patients were found to be co-infected by gonorrhea, 7.1% *Chlamydia trachomatis*, 3.3% *Trichomonas vaginalis* and 64.3% syphilis (2012 publication)⁹⁰.
- 4.9% of HIV-positive pregnant women were HBsAg-positive and 0.6% seropositive toward Hepatitis C (Kampala, 2007 publication)⁹¹. 9% of HIV-positive patients were HBsAg-positive (2010 publication)⁹². 6.1% of HIV-positive adolescents and young adults were HBsAg-positive (Kampala, 2011 to 2012)⁹³. 15% of HIV-positive patients were HBsAg-positive and 3% seropositive toward Hepatitis C (2008 publication)⁹⁴. 3.3% of HIV-positive patients were found to be seropositive toward HCV (Kampala, 2003 to 2004)⁹⁵. 17.3% of HIV-positive patients are infected by Hepatitis GB virus-C. (2007 publication)⁹⁶. 3.1% of HIV-positive persons above age 18 are seropositive toward Hepatitis A (IgM), 18.1% Hepatitis C; and 17.3% carry HBsAg (2006)⁹⁷.
- 87.5% of HIV-positive and 29.5% of HIV-negative persons ages 15 to 44 years in Kilimanjaro were found to be seropositive toward HSV-2 (2005)⁹⁸.
- *Aeromonas* was found in 7% of HIV-positive patients with diarrhea in rural Uganda, *Strongyloides stercoralis* 8% and *Cryptosporidium parvum* 5%. (2010 publication)

References

1. MMWR Recomm Rep 1992 Dec 18;41(RR-17):1-19.
2. MMWR Morb Mortal Wkly Rep 1987 Aug 14;36 Suppl 1:1S-15S.
3. Wkly Epidemiol Rec 1994 Sep 16;69(37):273-5.
4. N Engl J Med 1991 Jun 6;324(23):1644-50.
5. CNS Drugs 2003 ;17(12):869-87.
6. N Engl J Med 1990 Jan 18;322(3):161-5.
7. Curr Opin Pulm Med 2008 May ;14(3):228-34.
8. Curr Opin Infect Dis 2008 Feb ;21(1):31-6.
9. AIDS Alert 1999 Nov ;14(11):suppl 4.
10. Diagn Microbiol Infect Dis 2007 Mar ;57(3):289-94.
11. J Microbiol Immunol Infect 2005 Feb ;38(1):65-8.
12. Int J STD AIDS 2009 Jun ;20(6):369-72.
13. AIDS 2007 Oct 18;21(16):2119-29.
14. Ann N Y Acad Sci 2007 Sep ;1111:336-42.
15. AIDS 2008 May 31;22(9):1047-53.
16. Clin Infect Dis 1995 Aug ;21 Suppl 1:S108-10.
17. Clin Infect Dis 2000 Jun ;30(6):877-81.
18. Med Mal Infect 2007 Oct ;37(10):629-36.
19. Malar J 2007 ;6:143.
20. Clin Infect Dis 2007 Nov 1;45(9):1208-13.
21. Malar J 2007 ;6:143.
22. Lancet Infect Dis 2011 Jul ;11(7):541-56.
23. Sante 2011 Jul 1;21(3):174-177.
24. Future Virol 2012 ;7(7):699-708.
25. Parasit Vectors 2013 Jan 17;6(1):18.
26. BMC Pediatr 2011 ;11:5.
27. Mediterr J Hematol Infect Dis 2012 ;4(1):e2012032.
28. Trends Parasitol 2008 Jun ;24(6):264-71.
29. Int J Infect Dis 2008 Nov ;12(6):587-92.
30. Int J STD AIDS 2012 Aug ;23(8):e4-6.
31. AIDS Rev 2008 Apr-Jun;10(2):85-92.
32. Mayo Clin Proc 2007 Sep ;82(9):1091-102.
33. MMWR Morb Mortal Wkly Rep 2007 Jun 29;56(25):625-8.
34. Clin Infect Dis 2007 May 1;44(9):1222-8.
35. Dermatol Clin 2006 Oct ;24(4):497-507, vi.
36. Int J STD AIDS 2009 Apr ;20(4):278-84.
37. Eur J Intern Med 2009 Jan ;20(1):9-13.
38. Int J STD AIDS 2012 Aug ;23(8):599-600.
39. Int J STD AIDS 2010 Jan ;21(1):57-9.
40. Int J STD AIDS 2009 Apr ;20(4):238-40.
41. Trans R Soc Trop Med Hyg 2008 Dec ;102(12):1176-80.
42. J Infect Dis 2009 Jul 15;200(2):227-35.
43. J Antimicrob Chemother 2010 Jan ;65(1):10-7.
44. Semin Liver Dis 2012 May ;32(2):114-9.
45. Lancet Infect Dis 2009 Dec ;9(12):775-83.
46. Semin Liver Dis 2012 May ;32(2):138-46.
47. Eur Rev Med Pharmacol Sci 2012 Nov ;16(11):1473-83.
48. J Infect Dis 2013 Mar ;207 Suppl 1:S40-4.
49. J Infect Dis 2013 Mar ;207 Suppl 1:S26-32.
50. J Infect Dis 2013 Mar ;207 Suppl 1:S1-6.
51. Semin Liver Dis 2012 May ;32(2):120-9.
52. Clin Infect Dis 2002 Feb 1;34(3):379-85.
53. Int J STD AIDS 2011 Apr ;22(4):181-6.
54. Lancet Infect Dis 2008 Aug ;8(8):471.
55. AIDS 2008 Feb 19;22(4):503-10.
56. PLoS One 2012 ;7(8):e43400.
57. Trop Med Int Health 2012 Jul ;17(7):796-807.
58. Trop Med Int Health 2010 Apr ;15(4):414-22.
59. J Acquir Immune Defic Syndr 2008 Oct 1;49(2):183-9.
60. Bull World Health Organ 2008 Apr ;86(4):302-9.
61. J Med Virol 2011 May ;83(5):796-800.
62. Sex Transm Infect 2011 Oct ;87(6):511-5.
63. Afr Health Sci 2007 Sep ;7(3):124-8.
64. Trop Med Int Health 2006 Jun ;11(6):951-4.
65. BMC Public Health 2006 ;6:91.
66. Trop Med Int Health 2008 May ;13(5):680-2.
67. PLoS Negl Trop Dis 2009 ;3(6):e473.
68. Int J STD AIDS 2009 Jul ;20(7):499-502.
69. Trans R Soc Trop Med Hyg 2006 Jun ;100(6):586-93.
70. Trans R Soc Trop Med Hyg 2007 Sep ;101(9):899-907.
71. Trans R Soc Trop Med Hyg 2006 Jun ;100(6):543-50.
72. Sex Transm Infect 1999 Apr ;75(2):98-102.
73. Afr Health Sci 2011 Aug ;11 Suppl 1:S20-3.
74. Trans R Soc Trop Med Hyg 2009 Jun ;103(6):541-8.
75. BMC Pediatr 2011 ;11:5.
76. AIDS 2007 May 11;21(8):1027-34.
77. J Acquir Immune Defic Syndr 2008 Jun 1;48(2):203-10.
78. AIDS 2009 Jul 31;23(12):1589-94.
79. WHO AIDS Tech Bull 1989 Mar ;2:37.
80. Int J Tuberc Lung Dis 2006 Jul ;10(7):761-7.
81. J Acquir Immune Defic Syndr Hum Retrovirol 1998 Dec 15;19(5):484-9.
82. AIDS 2002 May 3;16(7):1031-8.
83. J Acquir Immune Defic Syndr 2010 Jul ;54(3):269-74.
84. Nihon Ishinkin Gakkai Zasshi 2003 ;44(4):299-306.
85. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):144-7.
86. Trans R Soc Trop Med Hyg 1991 Mar-Apr;85(2):227-9.
87. Acta Trop 2006 Dec ;100(3):218-22.
88. BMC Pediatr 2011 ;11:5.
89. Malar J 2011 ;10:66.
90. J Infect Dev Ctries 2012 ;6(3):276-82.
91. J Med Virol 2007 Dec ;79(12):1797-801.
92. Int J STD AIDS 2010 Aug ;21(8):553-7.
93. J Int AIDS Soc 2012 ;15(6):18433.
94. Afr Health Sci 2008 Mar ;8(1):8-12.
95. Afr Health Sci 2009 Sep ;9(3):143-6.
96. Int J STD AIDS 2007 Apr ;18(4):244-9.
97. BMC Public Health 2008 ;8:416.
98. East Afr J Public Health 2011 Mar ;8(1):28-32.

Amoeba - free living

Agent	PARASITE - Protozoa. Centramoebida, Acanthamoebidae: Acanthamoeba and Balamuthia Schizopyrenida, Vahkampfiidae: Naegleria
Reservoir	Water Soil
Vector	None
Vehicle	Water (diving, swimming)
Incubation Period	5d - 6d (range 2d - 14d)
Diagnostic Tests	Wet preparation. Specialized cultures. Serology available in reference centers.
Typical Adult Therapy	CNS Naegleria: Amphotericin B to 1 mg/kg/d IV + 1.5 mg intrathecal X 8 days; plus Miconazole 350 mg/sq m/d IV + 10 mg intrathecal qod X 8d Acanthamoeba: Sulfonamides + Flucytosine
Typical Pediatric Therapy	CNS Naegleria: Amphotericin B to 1 mg/kg/d IV + 1.5 mg intrathecal X 8 days; plus Miconazole 350 mg/sq m/d IV + 10 mg intrathecal qod X 8d Acanthamoeba: Sulfonamides + Flucytosine
Clinical Hints	Severe, rapidly-progressing meningoencephalitis (<i>Naegleria</i> , <i>Acanthamoeba</i> or <i>Balamuthia</i>) following swimming or diving in fresh water; or keratitis (<i>Acanthamoeba</i>), often following use of contaminated solutions to clean contact lenses.
Synonyms	Acanthamoben, Acanthamoeba, Amebic keratitis, Balamuthia, Balmuthia, Dictyostelium, Free-living ameba, Leptomyxid ameba, Naegleria, Paravahlkampfia, Primary amebic meningoencephalitis, Sappinia, Vahlkampfia. ICD9: 136.2 ICD10: B60.1,B60.2

Clinical

Primary amebic meningoencephalitis usually occurs in children and young adults who have been swimming in warm fresh water. ¹

Infection is heralded by abnormal sensations of taste or smell followed by abrupt onset of fever, nausea, and vomiting.

- The majority of patients have headache, meningitis and disorders of mental status changes.
- Coma and death may ensue within one week
- Only three nonfatal infections had been reported to 2003.

Acanthamoeba encephalitis:

Granulomatous amebic encephalitis due to *Acanthamoeba* occurs in immunocompromised and debilitated patients.

- Infection has a gradual onset characterized focal neurological deficits, mental status abnormalities, seizures, fever, headache, hemiparesis and meningismus.
- Visual disturbances and ataxia are often encountered.
- Death may ensue within 7 to as long as 120 days.
- Secondary infection of a cerebral ependymal cyst has been reported. ²
- Disseminated *Acanthamoeba* infection has been reported in an HIV-positive patient. ³

Balamuthia encephalitis:

Balamuthia mandrillaris encephalitis may be associated with headache, low-grade fever, vomiting, ataxia, photophobia, cranial nerve palsy, speech disturbances, cerebellar nystagmus, seizures, and altered mental status. ^{4 5}

- The case-fatality rate for *Balamuthia* encephalitis is over 90%.

Acanthamoeba keratitis:

Acanthamoeba keratitis is clinically similar to herpetic infection, and presents with a foreign-body sensation followed by severe pain, photophobia, tearing, blepharospasm, conjunctivitis, iritis, anterior uveitis, dendriform keratitis, radial keratoneuritis, ptosis and blurred vision. ⁶⁻¹¹

- In rare instances, the infection is painless. ^{12 13}
- Rupture of Descemet's membrane may occur. ¹⁴
- Bilateral infection is common. ¹⁵
- Dacryoadenitis may be present in some cases. ¹⁶
- Ocular discharge and endophthalmitis are very rare. ¹⁷
- Sympathetic ophthalmia of the un-infected eye has been reported. ¹⁸

- Atypical presentations have been described in patients with keratoconus. ¹⁹

Acanthamoeba infection has also been associated with skin ulcers ²⁰, pneumonia, adenitis, vasculitis, osteomyelitis, and sinusitis.

- Cutaneous acanthamoebiasis has been associated with ulceronecrotic lesions, an infiltrative bluish plaque, or periorbital tumor. ²¹
- Fatal disseminated *Acanthamoeba lenticulata* infection has been reported in a heart transplant patient.
- Four cases of disseminated *Acanthamoeba* infection in stem-cell transplant recipients had been reported as of 2008. ²²

This disease is endemic or potentially endemic to all countries.

References

1. Int J Parasitol 2004 Aug ;34(9):1001-27.
2. Surg Neurol 2009 Aug ;72(2):185-9.
3. Diagn Microbiol Infect Dis 2007 Mar ;57(3):289-94.
4. MMWR Morb Mortal Wkly Rep 2008 Jul 18;57(28):768-71.
5. Clin Microbiol Rev 2008 Jul ;21(3):435-48.
6. Eye (Lond) 2003 Nov ;17(8):893-905.
7. Ophthalmology 2006 Mar ;113(3):412-6.
8. Curr Opin Ophthalmol 2006 Aug ;17(4):327-31.
9. Eye Contact Lens 2008 Sep ;34(5):247-53.
10. Nihon Ganka Gakkai Zasshi 2010 Dec ;114(12):1030-5.
11. Middle East Afr J Ophthalmol 2011 Jul ;18(3):252-5.
12. Coll Antropol 2009 Sep ;33(3):951-4.
13. Ophthalmologie 2007 May ;104(5):415-7.
14. Eye Contact Lens 2009 Nov ;35(6):338-40.
15. Am J Ophthalmol 2008 Feb ;145(2):193-197.
16. Arch Ophthalmol 2006 Sep ;124(9):1239-42.
17. Trends Parasitol 2006 Apr ;22(4):175-80.
18. Eye Contact Lens 2011 Nov ;37(6):374-6.
19. Eye Contact Lens 2009 Jan ;35(1):38-40.
20. Transpl Infect Dis 2007 Mar ;9(1):51-4.
21. Int J Dermatol 2009 Dec ;48(12):1324-9.
22. Transpl Infect Dis 2008 Dec ;10(6):437-41.

Amoebic abscess

Agent	PARASITE - Protozoa. Sarcocystid, Entamoebidae: <i>Entamoeba histolytica</i> (must be distinguished from non-invasive, <i>Entamoeba dispar</i>)
Reservoir	Human
Vector	Fly (<i>Musca</i>) - occasionally
Vehicle	Food Water Sexual contact Fly
Incubation Period	2w - 6m (rarely years; 95% within 6m)
Diagnostic Tests	Imaging. Serology. Nucleic acid amplification. Note: Amoebae are usually not present in stool at this stage.
Typical Adult Therapy	Metronidazole 750 mg TID X 10d OR Tinidazole 800 mg TID X 5d
Typical Pediatric Therapy	Metronidazole 15 mg/kg TID X 10d OR Tinidazole 15 to 20 mg/kg TID X 5d
Clinical Hints	Fever, local pain, weight loss. Remember that liver abscess may be bacterial or amoebic - latter most often single and in right hepatic lobe.
Synonyms	Absceso amebiano, Amebic liver abscess. ICD9: 006.3,006.4,006.5,006.6,006.8 ICD10: A06.4,106.5,A06.7,106.8

Clinical

The clinical presentation may be acute or subacute in onset.

- Fever than 50% of patients have fever, hepatomegaly or abdominal pain.
- 30% to 40% have concurrent diarrhea.
- Other findings may include shoulder pain, cough, chest pain, pleural or pericardial effusion. ^{1 2}
- The findings of ameboma may mimic those of malignancy. ³
- A case of Budd-Chiari syndrome complicating amebic abscess has been reported. ⁴

Laboratory findings include leukocytosis without eosinophilia in 80%, anemia in over 50%, elevated serum alkaline phosphatase levels in 80%.

Pleuropulmonary amebiasis is the most common complication of amebic liver abscess, usually representing rupture of a superior right lobe abscess through the diaphragm.

- Symptoms include cough, pleuritic pain, and dyspnea.
- Empyema, hepatobronchial fistula or pericarditis (from left lobe abscesses) may follow.
- Although most cases involve the liver, abscesses may occur in virtually any organ. ^{5 6}
- *Entamoeba histolytica* encephalitis has been reported. ⁷

This disease is endemic or potentially endemic to all countries.

Amoebic abscess in Uganda

Epidemiological data regarding Amebic abscess are included in the notes for Amebic colitis

References

1. South Med J 2004 Jul ;97(7):673-82.
2. Curr Gastroenterol Rep 2004 Aug ;6(4):273-9.
3. Indian J Pathol Microbiol 2009 Apr-Jun;52(2):228-30.
4. Am J Trop Med Hyg 2009 Nov ;81(5):768-9.
5. No Shinkei Geka 2007 Sep ;35(9):919-25.
6. Parasitol Res 2012 Mar ;110(3):1291-6.
7. Trans R Soc Trop Med Hyg 2007 Mar ;101(3):311-3.

Amoebic colitis

Agent	PARASITE - Protozoa. Sarcomastigota, Entamoebidea: Entamoeba histolytica (must be distinguished from non-invasive, Entamoeba dispar)
Reservoir	Human
Vector	Fly (Musca) - occasionally
Vehicle	Food Water Sexual contact Fly
Incubation Period	1w - 3w (range 3d - 90d)
Diagnostic Tests	Fresh stool/aspirate for microscopy. Stool antigen assay. Stool PCR. Note: serological tests usually negative.
Typical Adult Therapy	Metronidazole 750 mg TID X 10d OR Tinidazole 2 g as single dose daily X 5d
Typical Pediatric Therapy	Metronidazole 15 mg/kg TID X 10d OR Tinidazole 50 mg/kg as single dose daily X 5d
Clinical Hints	Dysentery, abdominal pain, tenesmus - without hyperemia of rectal mucosa or fecal pus (i.e., unlike shigellosis); liver abscess and dysentery rarely coexist in a given patient.
Synonyms	Amebiasis, Amebiasis intestinal, Amebic dysentery, Amoebenruhr, Entamoeba bangladeshi, Entamoeba gingivalis, Entamoeba moshkovskii. ICD9: 006.0,006.1,006.2 ICD10: A06.0,A06.1,A06.2

Clinical

Patients with noninvasive infection may present with nonspecific gastrointestinal complaints such as chronic intermittent diarrhea, mucus, abdominal pain, flatulence and weight loss ^{1 2}

Infection has been documented in children as young as two weeks of age. ^{3 4}

A review of amebiasis among men who have sex with men • see reference ⁵

Cases of cutaneous amebiasis of the penis have been acquired through insertive anal intercourse. ⁶

Invasive amebiasis:

The onset of invasive infection is usually gradual (over 1 to 3 weeks) and characterized by abdominal pain, tenderness, and bloody stools.

- Fever is present in one third of cases, and the may be enlarged and tender.
- Signs of fluid loss and electrolyte loss may be seen in severe infections.
- In children, colitis can present as rectal bleeding alone without diarrhea.
- Fecal leukocytes may not be present, and are not as numerous as in shigellosis.
- Charcot-Leyden crystals are often seen in the stool.

Fulminant colitis:

Fulminant colitis is rare and carries a very high mortality.

- Predisposing factors include malnourishment, pregnancy and corticosteroid treatment.
- Such patients are severely ill with fever, leukocytosis, profuse bloody and mucoid diarrhea, generalized abdominal pain.
- Hypotension and peritonitis may be evident.
- Intestinal perforation and necrosis, or hepatic abscess may ensue.
- The clinical features of Cytomegalovirus colitis in AIDS patients may mimic those of amebic colitis. ⁷

Additional complications:

Additional complications include toxic megacolon (complicates 0.5% of amebic colitis cases); annular ameboma of the colon, which may mimic carcinoma.

- Chronic, irritative bowel syndromes, ulcerative post-dysenteric colitis or perianal amebiasis may also follow acute amebic colitis.
- Extraintestinal amebiasis may involve a wide variety of organs.
- Other forms of amebiasis include colocolocutaneous fistula ⁸ or amebiasis cutis ⁹, brain abscess, cervicitis ¹⁰, rectovaginal fistulae and penile infection

Liver abscess is discussed separately in this module.

This disease is endemic or potentially endemic to all countries.

Amoebic colitis in Uganda

Notable outbreaks:

2010 - An outbreak (15 fatal cases) of amebic dysentery was reported in the eastern region. ¹¹

References

1. Curr Opin Infect Dis 2003 Oct ;16(5):479-85.
2. N Engl J Med 2003 Apr 17;348(16):1565-73.
3. Indian J Pediatr 2010 Aug ;77(8):903-4.
4. Case Rep Infect Dis 2012 ;2012:614398.
5. Lancet Infect Dis 2012 Sep ;12(9):729-36.
6. Sex Transm Infect 2012 Aug 28;
7. Am J Med Sci 2008 Oct ;336(4):362-4.
8. Int J Surg Case Rep 2011 ;2(3):40-3.
9. Australas J Dermatol 2010 Feb ;51(1):52-5.
10. J Microbiol Immunol Infect 2012 Jun 26;
11. ProMED <promedmail.org> archive: 20101130.4309

Animal bite-associated infection

Agent	BACTERIUM. <i>Pasteurella multocida</i> , and other zoonotic bite pathogens
Reservoir	Cat Dog Marsupial (Tasmanian devil) Other mammal Rarely bird
Vector	None
Vehicle	Cat (60%), dog (30%) or other bite. No obvious source in 10%
Incubation Period	3h - 3d
Diagnostic Tests	Gram stain/culture. Hold specimen for 2 weeks to discount Capnocytophaga & other genera.
Typical Adult Therapy	Penicillin, a Tetracycline or Cefuroxime . Dosage and duration appropriate for nature and severity of infection
Typical Pediatric Therapy	Penicillin or Cefuroxime . Dosage and duration appropriate for nature and severity of infection
Clinical Hints	Infection of cat, dog or other bite wound - acquired during the preceding 3 to 72 hours (no history of bite in 10%); systemic infection (meninges, bone, lungs, joints, etc) may occur.
Synonyms	<i>Bacteroides pyogenes</i> , <i>Bacteroides tectus</i> , <i>Bergeyella zoohelcum</i> , Bisgaard's taxon 16, <i>Capnocytophaga canimorsus</i> , <i>Capnocytophaga cynodegmi</i> , CDC EF-4, CDC NO-1, <i>Corynebacterium kutscheri</i> , <i>Corynebacterium canis</i> , <i>Corynebacterium freiburgense</i> , <i>Fusobacterium canifelinum</i> , <i>Halomonas venusta</i> , <i>Kingella potus</i> , <i>Moraxella canis</i> , <i>Neisseria animaloris</i> , <i>Neisseria canis</i> , <i>Neisseria weaveri</i> , <i>Neisseria zoodegmatis</i> , <i>Pasteurella caballi</i> , <i>Pasteurella canis</i> , <i>Pasteurella dagmatis</i> , <i>Pasteurella multocida</i> , <i>Pasteurella stomatis</i> , <i>Psychrobacter immobilis</i> , <i>Staphylococcus intermedius</i> , <i>Vibrio harveyi</i> . ICD9: 027.2 ICD10: A28.0

Clinical

These are typically skin and soft infections which follow the bites of cats, dogs or other animals • usually during the preceding 3 to 72 hours. ¹

- There is no history of bite in ten percent of cases.
- Systemic infection (meninges ², bone, lungs ³, joints, etc) may occur, with rare instance of severe septicemia. ⁴⁻⁷

See the Microbiology module (Bacteria • Characterize) for a comprehensive discussion of bacterial species associated with bite wound infection in humans.

This disease is endemic or potentially endemic to all countries.

References

1. *J Am Acad Dermatol* 1995 Dec ;33(6):1019-29.
2. *Scand J Infect Dis* 2002 ;34(3):213-7.
3. *Semin Respir Infect* 1997 Mar ;12(1):54-6.
4. *Am J Emerg Med* 2008 Mar ;26(3):380.e1-3.
5. *Kansenshogaku Zasshi* 2009 Sep ;83(5):557-60.
6. *Indian J Pathol Microbiol* 2011 Apr-Jun;54(2):368-70.
7. *Conn Med* 2011 Nov-Dec;75(10):603-5.

Anisakiasis

Agent	PARASITE - Nematoda. Phasmidea: <i>Anisakis simplex</i> and <i>Pseudoterranova decipiens</i>
Reservoir	Marine mammals Fish
Vector	None
Vehicle	Undercooked fish
Incubation Period	Hours - 14d
Diagnostic Tests	Endoscopic identification of larvae.
Typical Adult Therapy	Endoscopic removal of larvae; surgery for complications
Typical Pediatric Therapy	As for adult
Clinical Hints	Allergic reactions; or acute and chronic abdominal pain, often with 'peritoneal signs' or hematemesis; follows ingestion of undercooked fish (e.g., sushi), squid or octopus.
Synonyms	Anasakis, Bolbosoma, Cod worm disease, Contraecaecum, Eustrongylides, Herring worm disease, Pseudoterranova, Whaleworm. ICD9: 127.1 ICD10: B81.0

Clinical

The location of the worms and presenting features depend somewhat on the genus.

- *Phocanema* more commonly associated with infection of the stomach.
- *Anisakis* is usually associated with intestinal disease. ¹

Invasive anisakiasis:

Symptoms occur within 48 hours after ingestion.

- Gastric anisakiasis is characterized by intense abdominal pain, nausea, and vomiting. ²
- Small intestinal involvement results in lower abdominal pain and signs of obstruction ³, and may cause ⁴ or mimic appendicitis. ^{5 6}
- Rare instances of duodenal ulcer have been reported. ⁷
- Symptoms may last for months, rarely for years.
- The disease may also suggest tumor, regional enteritis or diverticulitis. ⁸
- Rare instances of overt hemorrhage ⁹ and intussusception are reported. ^{10 11}
- Patients may also experience a pharyngeal "tickling sensation", cough or a foreign body in the mouth or throat. ¹²

Allergic anisakiasis:

Ingestion of *Anisakis* larvae with seafood is often responsible for acute allergic manifestations such as urticaria and anaphylaxis ¹³, with or without accompanying gastrointestinal symptomatology. ¹⁴

- Eosinophilia is usually not present in either gastric or intestinal anisakiasis; however, leukocytosis is noted in two thirds of patients with intestinal involvement.
- Urticaria is present in 20% of cases ¹⁵

This disease is endemic or potentially endemic to all countries.

References

1. Trends Parasitol 2002 Jan ;18(1):20-5.
2. Gastroenterol Hepatol 2003 Jun-Jul;26(6):341-6.
3. Korean J Gastroenterol 2010 Sep ;56(3):192-5.
4. J Med Case Reports 2012 Apr 23;6(1):114.
5. Clin Microbiol Infect 2003 Jul ;9(7):734-7.
6. Ann Chir 2005 Jul-Aug;130(6-7):407-10.
7. Chonnam Med J 2012 Apr ;48(1):73-5.
8. Rev Esp Enferm Dig 2002 Aug ;94(8):463-72.
9. Updates Surg 2011 Sep ;63(3):213-7.
10. World J Gastroenterol 2010 Apr 14;16(14):1804-7.
11. Intern Med 2013 ;52(2):223-6.
12. J Parasitol 2007 Apr ;93(2):440-3.
13. J Infect Chemother 2011 Aug ;17(4):544-6.
14. Clin Microbiol Rev 2008 Apr ;21(2):360-79, table of contents.
15. Bull Acad Natl Med 2007 Jan ;191(1):53-65; discussion 65-6.

Anthrax

Agent	BACTERIUM. <i>Bacillus anthracis</i> An aerobic gram positive bacillus
Reservoir	Soil Goat Cattle Sheep Water Horse
Vector	Fly (rare)
Vehicle	Hair Wool Hides Bone products Air Meat
Incubation Period	1d-7d; 1-12 cutaneous, 1-7 GI; 1-43 pulmonary
Diagnostic Tests	Bacteriological culture. Alert laboratory that organism may be present. Serology and rapid tests by Ref. Centers.
Typical Adult Therapy	Isolation (secretions). Ciprofloxacin ; alt. Doxycycline , Penicillin G . Add Clindamycin + Rifampin for pulmonary infection. Dosage/route/duration as per severity
Typical Pediatric Therapy	Isolation (secretions). Ciprofloxacin (Doxycycline if age \geq 8y). Add Clindamycin + Rifampin for pulmonary infection. Dosage/route/duration as per severity
Vaccine	Anthrax
Clinical Hints	Edematous skin ulcer covered by black eschar - satellite vesicles may be present; fulminant gastroenteritis or pneumonia; necrotizing stomatitis; hemorrhagic meningitis. Acquired from contact with large mammals or their products (meat, wool, hides, bone).
Synonyms	Antrace, Antrax, Antraz, Carhunco, Carbunculo, Malcharbon, Malignant pustule, Miltbrann, Miltvuur, Milzbrand, Mjaltbrand, Siberian plague, Siberian ulcer, Splenic fever, Wool-sorter's disease. ICD9: 022 ICD10: A22

Clinical

Most cases of anthrax occur in one of four forms: cutaneous, gastrointestinal, oropharyngeal and inhalational. ¹

CDC case definition for reporting:

As of 1996, the CDC (The United States Centers for Disease Control) case definition for reporting purposes consists of any illness with acute onset characterized by one or more of the following:

- cutaneous (a skin lesion evolving during a period of 2-6 days from a papule, through a vesicle to a depressed black eschar)
- pulmonary (hypoxia, dyspnea and mediastinal widening following a brief 'viral-type' prodrome)
- intestinal (severe abdominal distress followed by fever or signs of septicemia)
- oropharyngeal (mucosal lesion, cervical adenopathy and edema, and fever)
- demonstration of *Bacillus anthracis* by culture, immunofluorescence or serological response.

WHO case definition for surveillance:

The WHO Case definition for surveillance is as follows:

Clinical description:

An illness with acute onset characterized by several clinical forms. These are:

(a) localized form:

- cutaneous: skin lesion evolving over 1 to 6 days from a papular through a vesicular stage, to a depressed black eschar invariably accompanied by edema that may be mild to extensive

• systemic forms:

- gastro-intestinal: abdominal distress characterized by nausea, vomiting, anorexia and followed by fever
- pulmonary (inhalation): brief prodrome resembling acute viral respiratory illness, followed by rapid onset of hypoxia, dyspnea and high temperature, with X-ray evidence of mediastinal widening
- meningeal: acute onset of high fever possibly with convulsions, loss of consciousness, meningeal signs and symptoms; commonly noted in all systemic infections

Laboratory criteria for diagnosis

- isolation of *Bacillus anthracis* from a clinical specimen (e.g., blood, lesions, discharges)
- demonstration of *B. anthracis* in a clinical specimen by microscopic examination of stained smears (vesicular fluid, blood, cerebrospinal fluid, pleural fluid, stools)
- positive serology (ELISA, Western blot, toxin detection, chromatographic assay, fluorescent antibody test (FAT))
- Note: It may not be possible to demonstrate *B. anthracis* in clinical specimens if the patient has been treated with antimicrobial agents.

Case classification

- Suspected: A case that is compatible with the clinical description and has an epidemiological link to confirmed or suspected

animal cases or contaminated animal products.

- Probable: A suspected case that has a positive reaction to allergic skin test (in non-vaccinated individuals).
- Confirmed: A suspected case that is laboratory-confirmed.

Cutaneous anthrax:

- 95% of anthrax cases (worldwide) are cutaneous.
- The incubation period for cutaneous anthrax ranges from 12 hours to 12 days.
- Cutaneous anthrax begins with pruritus at the affected site, typically followed by a small, painless papule that progresses to a vesicle in 1 to 2 days.²
- The lesion erodes, leaving a necrotic ulcer with a characteristic black center.
- Secondary vesicles are sometimes observed.
- Lymphadenopathy may occur, and local edema may be extensive.
- Patients may have fever, malaise, and headache.
- The most common sites of cutaneous anthrax are the hands, forearms, and head.
- Anthrax related to illicit drug injection may present as subcutaneous infection rather than overt skin lesions.³
- Rarely infection may involve the genital area⁴, eyelid⁵⁻⁷ or other areas.
- Cutaneous anthrax is fatal in approximately 20% of cases if left untreated.

Inhalational anthrax: ^{8 9}

- Infection may progress to respiratory failure and shock within 1 to 2 days following onset of symptoms.
- The case-fatality rate exceeds 80%, even with appropriate antibiotic therapy.¹⁰
- Symptoms include pharyngeal pain, cough, fever and myalgia • followed by respiratory distress, cervical edema and venous engorgement suggestive of mediastinitis.^{11 12}

Gastrointestinal anthrax: ¹³

- Infection is characterized by pharyngeal pain, nausea, vomiting, and bloody diarrhea.
- Intestinal gangrene, obstruction and perforation may ensue.¹⁴
- The case-fatality rate for intestinal infection ranges from 25% to 60%.
- Ulcerative lesions, usually multiple and superficial, may occur in the stomach, sometimes in association with similar lesions of the esophagus and jejunum.
- Ulcers may bleed, and in severe cases the hemorrhage may be massive and fatal.
- Ascites may be present.
- Lesions in the mid-jejunum, terminal ileum, or cecum tend to develop around a single site or a few sites of ulceration and edema, similar to cutaneous anthrax.

Oropharyngeal anthrax:

- Infection is characterized by painful neck swelling and fever.
- The other common symptoms are sore throat, dysphagia, and hoarseness, enlargement of cervical lymph nodes and soft tissue edema.
- Oral lesions are located on the tonsils, posterior pharyngeal wall, or the hard palate.¹⁵
- In severe cases, the tonsillar lesions extended to involve the anterior and posterior pillars of fauces, as well as the soft palate and uvula.
- Early lesions are edematous and congested.
- By the end of the first week, central necrosis and ulceration produce a whitish patch, which evolves to a pseudomembrane which covers the ulcer after an additional week.


Meningeal anthrax: ¹⁶

- Infection is characterized by fever, malaise, meningeal signs, hyperreflexia, and delirium, stupor, or coma.¹⁷
- CSF analyses demonstrated hemorrhagic meningitis, with positive Gram's stains and CSF cultures.
- 75% of patients die within 24 hours of presentation; mortality rates of 100% are reported in some series.^{18 19}
- Pathologic findings include hemorrhagic meningitis, multifocal subarachnoid and intraparenchymal hemorrhages, vasculitis, and cerebral edema.²⁰

Published case-fatality rates are as follows: cutaneous <1%, gastrointestinal 25% to 60%, inhalational 46% and injectional 33%.²¹

This disease is endemic or potentially endemic to 147 countries.

Anthrax in Uganda




Graph: Uganda. Anthrax, cases

Notes:

Individual years:


1969 - Included 2 fatal cases. ²²



Graph: Uganda. Anthrax - bovine, outbreaks

Notes:

1. 4 to 5 cattle died of anthrax in 1974 and 1975.



Graph: Uganda. Anthrax - bovine

One outbreak (64 cases) of buffalo anthrax was reported in 2004.

Notable outbreaks:

- 1976 - An outbreak of gastrointestinal infection was traced to an infected zebu.
 - 1984 - An outbreak (143 cases, 9 fatal) of gastrointestinal anthrax was reported. ²³
 - 2003 - An outbreak (23 fatal cases) was reported among cattle in Hoima district.
 - 2003 - An outbreak (23 cases) was reported among cattle in Hoima. ²⁴
 - 2004 to 2005 - An outbreak (200 fatal cases) was ascribed to ingestion of infected corpses by other hippopotami. ^{25 26}
- Ten fatal cases were also reported among humans who ate meat from affected animals. ²⁷
- 2005 - An outbreak (50 fatal cases, approximate) was reported among zebra in Lake Mburo National Park (western Uganda). ²⁸
 - 2009 - An outbreak (6 fatal cases) in Bushenyi district was associated with ingestion of meat from an antelope. ^{29 30}
 - 2010 (publication year) - An outbreak (4 cases) of cutaneous anthrax in Kasese district was associated with the skinning of a goat. ³¹
 - 2010 - An outbreak (101 fatal cases) of anthrax was reported among hippopotami in Kalangala District ³²⁻³⁴
 - 2011 - An outbreak was reported among humans and cattle in western Uganda. ³⁵⁻³⁷

References

1. Curr Top Microbiol Immunol 2002 ;271:1-19.
2. Dermatol Clin 2004 Jul ;22(3):247-56, v.
3. J Bone Joint Surg Br 2011 Mar ;93(3):414-7.
4. Ann Trop Paediatr 2007 Dec ;27(4):307-9.
5. Turk J Pediatr 2009 Jan-Feb;51(1):67-8.
6. Bull Soc Belge Ophtalmol 2009 ;(312):29-36.
7. Clin Ophthalmol 2010 ;4:713-6.
8. Curr Infect Dis Rep 2002 Jun ;4(3):238-243.
9. Clin Infect Dis 2003 May 15;36(10):1275-83.
10. Chest 1999 Nov ;116(5):1369-76.
11. J Am Acad Nurse Pract 2001 Apr ;13(4):164-8; quiz 169-70.
12. Mod Pathol 2001 May ;14(5):482-95.
13. Emerg Infect Dis 2002 Jul ;8(7):649-51.
14. Arch Intern Med 2003 Nov 10;163(20):2527-31.
15. Emerg Radiol 2010 Mar ;17(2):161-4.
16. Lancet Infect Dis 2005 May ;5(5):287-95.
17. Scand J Infect Dis 2002 ;34(1):66-7.
18. J Neurol Sci 2009 Jun 15;281(1-2):41-5.
19. Mikrobiyol Bul 2009 Oct ;43(4):671-6.
20. Neurology 2002 Aug 13;59(3):327-34.
21. Am J Respir Crit Care Med 2011 Dec 15;184(12):1333-41.
22. Bull World Health Organ 1973 ;49(2):143-8.
23. Ann Ist Super Sanita 1984 ;20(2-3):205-8.
24. ProMED <promedmail.org> archive: 20030218.0422
25. ProMED <promedmail.org> archive: 20050221.0559
26. ProMED <promedmail.org> archive: 20091021.3624
27. ProMED <promedmail.org> archive: 20041109.3027
28. ProMED <promedmail.org> archive: 20050427.1176
29. ProMED <promedmail.org> archive: 20090219.0690
30. ProMED <promedmail.org> archive: 20090223.0758
31. East Afr J Public Health 2009 Dec ;6(3):235-9.
32. ProMED <promedmail.org> archive: 20100719.2418
33. ProMED <promedmail.org> archive: 20101012.3696
34. ProMED <promedmail.org> archive: 20101018.3779
35. ProMED <promedmail.org> archive: 20111107.3316
36. ProMED <promedmail.org> archive: 20111108.3322
37. ProMED <promedmail.org> archive: 20111111.3343

Ascariasis

Agent	PARASITE - Nematoda. Phasmidea: <i>Ascaris lumbricoides</i>
Reservoir	Human ? Dog
Vector	None
Vehicle	Vegetables Fly
Incubation Period	10d - 14d (range 7d - >200d)
Diagnostic Tests	Stool microscopy.
Typical Adult Therapy	Albendazole 400 mg X 1 dose OR Mebendazole 100 mg BID X 3d
Typical Pediatric Therapy	Mebendazole 100 mg BID X 3 d (> age 2).
Clinical Hints	An acute illness characterized by cough, wheezing and eosinophilia; adult worms are associated with abdominal pain (occasionally obstruction), pancreatic or biliary disease; highest rates among children and in areas of crowding and poor sanitation.
Synonyms	<i>Ascaris</i> , <i>Ascaris lumbricoides</i> , Askariasis. ICD9: 127.0 ICD10: B77

Clinical

The pulmonary manifestations of ascariasis occur during the stage of larval migration through the lungs and resemble Löffler's syndrome: cough, wheezing, pulmonary infiltration and eosinophilia. ^{1 2}

- Children with heavy *Ascaris* infection experience impaired digestion and absorption of proteins, often with moderate steatorrhea.
- A mass of worms may block the lumen of the small bowel, resulting in acute intestinal obstruction, with vomiting, abdominal distention, cramps ³⁻⁶ and occasionally hemorrhage ⁷, gangrene or perforation. ⁸
- Ileal volvulus and intussusception are also reported. ⁹

Worms may also invade and obstruct the biliary duct (pancreatic-biliary ascariasis) ¹⁰⁻¹⁹, producing abdominal pain, which may be associated with ascending cholangitis, acute or recurrent pancreatitis ^{20 21}, pancreatic pseudotumor ²² or obstructive jaundice. ²³⁻²⁶

- The majority of patients with hepatobiliary and pancreatic ascariasis present with biliary colic. ²⁷
- Choledocholithiasis, hepatolithiasis, liver abscess and cirrhosis are associated with the presence of dead, rather than viable worms. ²⁸
- Aberrant worms may appear at umbilical and hernial fistulas, the fallopian tubes, ovaries ²⁹, lower esophagus ³⁰, urinary bladder, pleural space ³¹, trans-nasal ³² or trans-ostomy ³³ feeding tubes, lungs, nose ³⁴, paranasal sinuses ³⁵ and other sites.

Ascaris suum has been reported to cause rare cases of myelitis, eosinophilic pneumonia and focal liver lesions in humans, and is discussed under 'Toxocariasis.' ³⁶⁻³⁹

This disease is endemic or potentially endemic to all countries.

Ascariasis in Uganda

Prevalence surveys:

- 6.3% nationwide (1998 to 2005) ⁴⁰
- 17.4% of persons in Kabale District (2003) ⁴¹
- 2.3% of otherwise healthy pregnant women at enrolment to a trial of deworming (Entebbe, 2007 publication) ⁴²
- 2% of pregnant women (Entebbe and Katabi, 2003 to 2005) ⁴³
- 0.5% of pregnant women in Masindi District (2003 to 2004) ⁴⁴
- 47.6% of mothers and 33.3% of their children in Habutobere; 36.8% of mothers and 27.3% of their children in Musezero (southwestern Uganda, 2010) ⁴⁵

- 17.5% of children below age 20 in Southern Uganda (1998) ⁴⁶
- 17% of primary school children in Kampala (1997 publication) ⁴⁷
- 2.4% of school children in northwest and southeast Uganda (2002 to 2003) ⁴⁸
- 9.3% of school children in proximity to Lake Victoria (2009 publication) ⁴⁹
- 2.8% of school children in 2003; 1.6% in 2004; 0.6% in 2005 (nationwide)
- 6.8% of school children in the Great Lakes region of Uganda, Tanzania, Kenya and Burundi (2010 publication) ⁵⁰

References

1. Adv Parasitol 2001 ;48:285-375.
2. J Clin Pathol 1965 Nov ;18(6):737-42.
3. Lancet 2006 May 6;367(9521):1521-32.
4. Indian J Pediatr 2007 Dec ;74(12):1085-7.
5. Pediatr Surg Int 2009 Dec ;25(12):1099-102.
6. World J Surg 2010 May ;34(5):963-8.
7. Malays J Med Sci 2012 Apr ;19(2):92-5.
8. Pathol Res Pract 2010 May 15;206(5):292-4.
9. J Indian Assoc Pediatr Surg 2012 Jul ;17(3):116-9.
10. AJR Am J Roentgenol 2007 Jun ;188(6):1596-603.
11. Chir Ital 2008 Sep-Oct;60(5):733-8.
12. Saudi J Gastroenterol 2007 Jan-Mar;13(1):25-32.
13. Ultrasound Q 2009 Dec ;25(4):207-9.
14. Eur J Pediatr Surg 2010 May ;20(3):187-90.
15. Trop Doct 2010 Oct ;40(4):227-9.
16. Clin Radiol 2011 Mar ;66(3):275-7.
17. Turk J Gastroenterol 2011 ;22(2):178-82.
18. Trop Gastroenterol 2011 Jul-Sep;32(3):210-3.
19. JOP 2013 ;14(1):88-91.
20. Singapore Med J 2009 Jun ;50(6):e218-9.
21. Rev Med Interne 2011 Jun ;32(6):e84-7.
22. Gastroenterol Hepatol 2011 Aug-Sep;34(7):464-7.
23. Indian J Gastroenterol 2001 Mar ;20 Suppl 1:C28-32.
24. World J Surg 2006 Aug ;30(8):1500-6.
25. Saudi J Gastroenterol 2009 Apr ;15(2):121-4.
26. BMJ Case Rep 2012 ;2012
27. Southeast Asian J Trop Med Public Health 2007 Jul ;38(4):631-5.
28. Saudi J Gastroenterol 2010 Jul-Sep;16(3):203-6.
29. Int J Gynecol Pathol 2011 Nov ;30(6):549-52.
30. World J Gastroenterol 2012 Apr 7;18(13):1552-4.
31. Ulus Travma Acil Cerrahi Derg 2010 Mar ;16(2):183-4.
32. Saudi J Gastroenterol 2009 Oct-Dec;15(4):288.
33. Turk J Gastroenterol 2011 ;22(2):203-4.
34. Br J Surg 1972 Jun ;59(6):437-42.
35. Trans R Soc Trop Med Hyg 1997 Jan-Feb;91(1):37.
36. Abdom Imaging 2004 Sep-Oct;29(5):598-602.
37. Rinsho Shinkeigaku 2004 Mar ;44(3):198-202.
38. Nihon Kokyuki Gakkai Zasshi 1998 Feb ;36(2):208-12.
39. J Helminthol 2012 Mar 19;:1-8.
40. Trop Med Int Health 2005 Nov ;10(11):1187-9.
41. Trans R Soc Trop Med Hyg 2005 Jan ;99(1):18-24.
42. Trans R Soc Trop Med Hyg 2007 Sep ;101(9):899-907.
43. J Infect Dis 2008 Sep 15;198(6):920-7.
44. Trans R Soc Trop Med Hyg 2008 Feb ;102(2):130-6.
45. Trans R Soc Trop Med Hyg 2012 Feb ;106(2):75-83.
46. East Afr Med J 2001 Jun ;78(6):283-6.
47. East Afr Med J 1997 May ;74(5):311-4.
48. Trans R Soc Trop Med Hyg 2006 Nov ;100(11):1039-48.
49. Geospat Health 2009 Nov ;4(1):39-53.
50. Trop Med Int Health 2010 Feb ;15(2):198-207.

Aspergillosis

Agent	FUNGUS. Ascomycota, Euascomycetes, Eurotiales: <i>Aspergillus</i> . A hyaline hyphomycete
Reservoir	Compost Hay Cereal Soil
Vector	None
Vehicle	Air
Incubation Period	3d - 21d
Diagnostic Tests	Fungal culture. Biopsy. Nasal culture or serologic testing may be useful in select cases.
Typical Adult Therapy	Voriconazole 6 mg/kg IV Q12h, day 1; follow with 4 mg/kg IV OR Amphotericin B - if invasive, rapidly increase to max dose 0.6 mg/kg/d and to total 2.5g. OR Itraconazole
Typical Pediatric Therapy	Voriconazole 3 to 9 mg/kg IV Q12h OR Amphotericin B - if invasive, rapidly increase to max dose 0.6 mg/kg/d X 6w. OR Itraconazole
Clinical Hints	Pulmonary "fungus ball"; adult-onset asthma; consolidation or infected "pulmonary infarct" in setting of immune suppression (e.g., AIDS, leukemia, etc) leads to widespread hematogenous dissemination if not treated promptly.
Synonyms	Aspergillose, Aspergillus. ICD9: 117.3 ICD10: B44

Clinical

Clinical forms of aspergillosis include: ^{1 2}

- allergy (allergic bronchopulmonary aspergillosis)
- colonization of air spaces (otomycosis, fungus ball or mycetoma of the paranasal sinuses ³ or lungs)
- non-pulmonary invasive (eye, sinuses, cardiac valve, skin, gastrointestinal tract) ^{4 5}
- pulmonary-invasive

Invasion of the ears and sinuses can cause extensive necrosis in immunocompromised hosts.

- The most common central nervous system manifestations include brain abscess or cerebral infarction
- Meningitis is rare
- Endophthalmitis and keratitis usually occur following injury
- Wound infections and infection of vascular access sites has also been reported. ⁶
- Sporadic instances of Isolated invasive *Aspergillus* tracheobronchitis ⁷ and chronic necrotizing pulmonary aspergillosis are encountered. ⁸

Case-fatality rates range from 10% to 90%.

- One series of 289 cases cited a mortality rate of 40.2% (2008 publication) ⁹

This disease is endemic or potentially endemic to all countries.

References

1. *J Infect Chemother* 2004 Jun ;10(3):138-45.
2. *Rev Pneumol Clin* 2004 Apr ;60(2):73-7.
3. *Quintessence Int* 2012 Feb ;43(2):143-6.
4. *Infection* 2006 Dec ;34(6):333-8.
5. *J Burn Care Res* 2007 Nov-Dec;28(6):918-21.
6. *Clin Microbiol Infect* 2004 Mar ;10 Suppl 1:24-30.
7. *Clin Microbiol Infect* 2010 Jun ;16(6):689-95.
8. *Int J Infect Dis* 2010 Jun ;14(6):e479-82.
9. *Clin Infect Dis* 2008 Nov 1;47(9):1176-84.

Bacillary angiomatosis

Agent	BACTERIUM. Bartonella henselae or Bartonella quintana . Rickettsia-like bacteria
Reservoir	Human ? Tick ? Cat
Vector	Cat flea Tick (ixodid) - rare
Vehicle	None
Incubation Period	Unknown
Diagnostic Tests	Histology with special stains. Specialized culture techniques. Serology. Nucleic acid amplification.
Typical Adult Therapy	Clarithromycin 500 mg BID X 8 weeks Alternatives Azithromycin 250 mg QD or Ciprofloxacin 500 mg BID
Typical Pediatric Therapy	Clarithromycin 7.5 mg/kg PO BID X 8 months. OR Gentamicin 2 mg/kg IMq12h
Clinical Hints	Hemangiomas papules and nodules of skin, spleen, liver (peliosis hepatis), bone or other tissues; virtually all in the setting of AIDS or other immune deficiency; rare instances following tick bite in immune-competent individuals.
Synonyms	Bacillary peliosis, Peliosis hepatis. ICD9: 757.32,083.8 ICD10: K76.4,A44.0

Clinical

Bacillary angiomatosis was originally described as involving skin and regional lymph nodes of HIV-infected persons. ¹

- Subsequent infections have involved patients with other forms of immune suppression, and presented in a variety of organs including liver, spleen, bone, brain, lung, bowel, and uterine cervix.

Cutaneous lesions often arise in crops and resemble the lesions of verruga peruana.

- Lesions may present as fixed or mobile subcutaneous or dermal nodules.
- Single or multiple dome-shaped, skin-colored, red or purple papules are also described, which may ulcerate and discharge serosanguinous fluid. ^{2 3}
- Lesions can range in diameter from millimeters to centimeters, and may mimic pyogenic granuloma. ⁴
- Regional lymph nodes are frequently enlarged in a variety of distributions.
- Involved organs contain multiple blood-filled cystic structures that range from microscopic to several millimeters in size.
- Bone disease may present as multiple osteolytic lesions.

This disease is endemic or potentially endemic to all countries.

References

1. Clin Infect Dis 2003 Aug 15;37(4):559-66.
2. Dermatology 2000 ;201(4):326-31.

3. Ophthal Plast Reconstr Surg 2010 Sep-Oct;26(5):371-2.
4. Ann Plast Surg 2012 Oct 3;

Bacillus cereus food poisoning

Agent	BACTERIUM. <i>Bacillus cereus</i> (toxin). An aerobic gram-positive bacillus
Reservoir	Soil Processed & dried foods
Vector	None
Vehicle	Food
Incubation Period	2h - 9h (range 1h - 24h)
Diagnostic Tests	No practical test available. Isolation of organism from suspect food.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Usually follows ingestion of rice or other vegetables; vomiting within 1 to 6 hours and/or diarrhea within 6 to 24 hours; no fecal leucocytes.
Synonyms	Bacillus cytotoxicus. ICD9: 005.89 ICD10: A05.4

Clinical

Two types of illness are caused by two distinct metabolites. ¹

- Diarrhea is caused by a large molecular weight protein.
- Vomiting is caused by a low molecular weight, heat-stable peptide. ²

Symptoms of *B. cereus* diarrheal food poisoning mimic those of *Clostridium perfringens* food poisoning.

- Symptoms of the emetic form mimic *S. aureus* food poisoning. ³

Diarrheal form:

The onset of watery diarrhea, abdominal cramps, and pain occurs 6 to 15 hours after consumption of contaminated food. ⁴

- Nausea may accompany diarrhea, but vomiting (emesis) rarely occurs.
- Symptoms persist for 24 hours in most instances.

Emetic form:

The emetic type of food poisoning is characterized by nausea and vomiting within 0.5 to 6 h after consumption of contaminated foods.

- Occasionally, abdominal cramps and/or diarrhea may also occur.
- Duration of symptoms is generally less than 24 h.

Only two fatal cases had been reported to 2005. ^{5 6} Illness was characterized by rhabdomyolysis and renal failure.

- A case of encephalopathy and hepatic failure • similar to Reye's syndrome • was related to *Bacillus cereus* food poisoning. ⁷
- A case report of fatal *Bacillus cereus* food poisoning was published from Belgium in 2011. ⁸

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 1993 Oct ;6(4):324-38.
2. J Food Prot 2005 Mar ;68(3):636-48.
3. FEMS Microbiol Lett 1997 Dec 15;157(2):223-8.
4. ProMED <promedmail.org> archive: 20071207.3948

5. J Clin Microbiol 2005 Aug ;43(8):4277-9.
6. N Engl J Med 1997 Apr 17;336(16):1142-8.
7. Brain Dev 2010 Sep ;32(8):688-90.
8. J Clin Microbiol 2011 Dec ;49(12):4379-81.

Bacterial vaginosis

Agent	BACTERIUM. <i>Gardnerella vaginalis</i> (facultative gram-negative bacillus), <i>Mobiluncus curtisii</i> , <i>Mobiluncus mulieris</i> , Prevotella, et al
Reservoir	Human
Vector	None
Vehicle	Sexual contact - normal flora in 14% (girls) to 70% (women)
Incubation Period	Unknown
Diagnostic Tests	Identification of "clue cells" or positive KOH test in vaginal discharge. Culture.
Typical Adult Therapy	Metronidazole 500 mg BID X 7d OR Tinidazole 2 g PO daily X 3d OR Clindamycin 300 mg BID X 7d + intravaginal Clindamycin or Metronidazole ? Also treat sexual partner
Typical Pediatric Therapy	Metronidazole 7.5 mg/kg BID X 7d
Clinical Hints	Thin vaginal discharge - "fishy" odor when mixed with KOH; mild to moderate pruritus; occasionally urethritis in sexual partner.
Synonyms	Gardnerella, Gardnerella vaginalis, Mobiluncus. ICD9: 041.89,616,10,099.8 ICD10: N76.1

Clinical

The diagnosis of bacterial vaginosis required three of the following: ¹⁻³

1. A white, noninflammatory vaginal discharge or coating
2. The presence of clue cells ⁴
3. A vaginal pH above 4.5
4. A fishy odor following addition of 10% KOH to the vaginal discharge (presumably due to liberated trimethylamine).

Note that routine culture is unnecessary.

Associated conditions:

Sequelae of bacterial vaginosis include preterm birth ⁵⁻⁷ and neonatal distress ⁸, low birth weight ⁹, chorioamnionitis, cervicitis ¹⁰, scalp abscess of the newborn, an increased risk of late miscarriage ¹¹ and maternal infection. ¹²

- Some studies have suggested a correlation between bacterial vaginosis and infertility. ¹³⁻¹⁹
- Bacterial vaginosis may increase the risk for acquisition of HIV infection.
- Bacterial vaginosis may predispose to urinary tract infection ²⁰ and endometritis. ²¹

Gardnerella vaginalis has rarely been associated with balanitis, urethritis, urinary tract infections, asymptomatic bacteremia and infectious endocarditis in adult males. ²²

Cases of osteomyelitis, discitis and septic arthritis due to *Gardnerella vaginalis* have been reported. ²³⁻²⁶

This disease is endemic or potentially endemic to all countries.

Bacterial vaginosis in Uganda

95% of sexually-experienced women followed for two years experienced at least one episode of bacterial vaginosis when followed for two years (Rakai, 2011 publication) ²⁷

Prevalence surveys:

- 36.1% of women with genital disease (southwestern Uganda, 1999 to 2001) ²⁸
- 38.3% of women in Rakai (2008 publication) ²⁹

References

1. Am Fam Physician 2004 Dec 1;70(11):2125-32.
2. J Reprod Med 2004 Oct ;49(10):781-6.
3. Infect Dis Clin North Am 2005 Jun ;19(2):387-406.
4. BMJ 2004 May 29;328(7451):1306-8.
5. J Perinat Med 2009 ;37(2):130-4.
6. Mymensingh Med J 2011 Jan ;20(1):115-20.
7. Gynecol Obstet Fertil 2012 Jan ;40(1):48-54.
8. J Matern Fetal Neonatal Med 2012 Jan ;25(1):64-7.
9. Bull World Health Organ 2007 Jan ;85(1):9-18.
10. J Infect Dis 2006 Mar 1;193(5):617-24.
11. Fertil Steril 2007 Nov ;88(5):1396-403.
12. Best Pract Res Clin Obstet Gynaecol 2007 Jun ;21(3):375-90.
13. Symp Soc Exp Biol 1990 ;44:225-40.
14. Eur J Obstet Gynecol Reprod Biol 2012 Nov 27;
15. Int J STD AIDS 2009 Nov ;20(11):778-81.
16. Eur J Obstet Gynecol Reprod Biol 2008 Sep ;140(1):3-11.
17. BJOG 2002 Jun ;109(6):714-7.
18. J Reprod Med 2001 Sep ;46(9):806-10.
19. Lancet 1999 Aug 7;354(9177):511.
20. J Obstet Gynaecol 2007 Apr ;27(3):252-4.
21. Infect Dis Obstet Gynecol 2006 ;2006:84140.
22. Int J STD AIDS 2010 Sep ;21(9):653-7.
23. J Clin Microbiol 2012 Dec ;50(12):4154-6.
24. J Med Microbiol 2009 Oct ;58(Pt 10):1382-4.
25. Clin Infect Dis 1995 Aug ;21(2):443-5.
26. J Clin Microbiol 2009 Jan ;47(1):264-5.
27. Sex Transm Dis 2011 Nov ;38(11):1040-5.
28. Sex Transm Infect 2005 Dec ;81(6):488-93.
29. Am J Obstet Gynecol 2009 Jan ;200(1):42.e1-7.

Balantidiasis

Agent	PARASITE - Protozoa. Ciliate (Ciliophora), Litostomatea: <i>Balantidium coli</i>
Reservoir	Pig Non-human primate Rodent
Vector	None
Vehicle	Water Food
Incubation Period	1d - 7d (range 1d - 60d)
Diagnostic Tests	Microscopy of stool or colonic aspirates.
Typical Adult Therapy	Tetracycline 500 mg QID X 10d. OR Metronidazole 750 mg TID X 5d. OR Iodoquinol 650 mg TID X 20d
Typical Pediatric Therapy	Age >= 8 years: Tetracycline 10 mg/kg QID (max 2g/d) X 10d. Age <8 yrs, Metronidazole 15 mg/kg TID X 5d; or Iodoquinol 13 mg/kg TID X 20d
Clinical Hints	Dysentery, often with vomiting; mimics intestinal amebiasis. The disease is most common in pig-raising areas. Symptoms last for one to four weeks, and may recur.
Synonyms	Balantidiose, Balantidiosis, <i>Balantidium coli</i> , Balantidosis, Balindosis, Ciliary dysentery. ICD9: 007.0 ICD10: A07.0

Clinical

Most cases are asymptomatic.

- Clinical manifestations, when present, include persistent diarrhea, occasionally dysentery, abdominal pain, and weight loss. ¹

Symptoms can be severe in debilitated individuals.

- *Balantidium* pneumonia has been reported in immune-compromised patients ² and persons with occupational exposure. ³

Diagnosis is based on detection of trophozoites in stool specimens or in tissue collected during endoscopy.

- Cysts are less frequently encountered.
- *Balantidium coli* is passed intermittently and once outside the colon is rapidly destroyed. Thus stool specimens should be collected repeatedly, and immediately examined or preserved.
- Rare cases of pulmonary infection ⁴ and osteomyelitis have been reported. ⁵
- *Balantidium coli* has been identified in the urine. ⁶

This disease is endemic or potentially endemic to 109 countries.

References

1. Gastroenterol Hepatol 2000 Mar ;23(3):129-31.
2. Am J Hematol 2003 Jul ;73(3):180-3.
3. Can J Infect Dis 2003 May ;14(3):163-6.
4. S Afr Med J 2010 Aug ;100(8):534-6.
5. J Neurosurg Spine 2012 Dec 21;
6. J Nephrol 2010 Nov-Dec;23(6):732-7.

Bartonellosis - cat borne

Agent	BACTERIUM. <i>Afipia felis</i> , <i>Bartonella henselae</i> , <i>Bartonella clarridgeiae</i> , et al. A facultative gram-negative coccobacillus
Reservoir	Cat Possibly tick
Vector	Flea (cat flea = Ctenocephalides)
Vehicle	Cat scratch Plant matter (thorn, etc)
Incubation Period	3d - 14d
Diagnostic Tests	Visualization of organisms on Warthin Starry stain. Culture. Serology. Nucleic acid amplification.
Typical Adult Therapy	Aspiration of nodes as necessary. Azithromycin 500 mg day 1, then 250 daily X 4 days Alternatives: Clarithromycin , Ciprofloxacin , Sulfamethoxazole/trimethoprim
Typical Pediatric Therapy	Aspiration of nodes as necessary. Azithromycin 10 mg/kg day 1, then 5 mg/kg daily X 4 days
Clinical Hints	Tender suppurative regional adenopathy following cat scratch (usually kitten); fever present in 25%. systemic infection (liver, brain, endocardium, bone, etc) occasionally encountered; most cases resolve within 6 weeks.
Synonyms	<i>Afipia felis</i> , <i>Bartonella clarridgeiae</i> , <i>Bartonella henselae</i> , <i>Bartonella koehlerae</i> , Cat scratch disease, Debre's syndrome, Foshay-Mollaret cat-scratch fever, Katszenkrat-Krankheit, Petzetakis' syndrome, SENLAT. ICD9: 078.3 ICD10: A28.1

Clinical

Clinical history:

Approximately 90% of patients have a history of exposure to a cat.

- The disease has also been reported after exposure to squirrels, dogs, goats, thorns and barbed wire. ¹
- 75% of patients report a bite or scratch to the head, neck or upper limbs.
- Subclinical bacteremia is common among immuno-competent persons with animal and arthropod contact.

Symptoms:

Following an incubation period of 3 to 10 days, a small skin lesion appears consisting of a macule, papule, pustule or vesicle.

- Within 1 to 2 weeks, edema and tenderness of the regional lymph nodes appear.
- In some cases, the patient may present with Parinaud oculoglandular syndrome (conjunctival granuloma with suppurative preauricular adenitis), encephalopathy, erythema nodosum, thrombocytopenic purpura, arthritis, synovitis or pneumonia.

Signs:

Physical examination reveals involvement of a single node in 50% of cases.

- 30% have involvement of multiple sites, and 20% involvement of several nodes in the same region.
- Lymph nodes typically measure 1 to 5 cm.
- The majority of lesions regress over 2 to 6 months, but may last for as long as 2 years.
- Suppuration occurs in 10% of cases, and cellulitis is rare.
- Inguinal lymphadenopathy in cat-scratch disease may suggest a diagnosis of lymphogranuloma venereum. ²

Additional findings:

One third of patients manifest fever, lasting 1 to 7 days; and some cases may present as Fever of Unknown Origin. ³

- Malaise, fatigue, anorexia, vomiting, weight loss, headache, splenomegaly and pharyngitis are occasionally observed.
- 10.5% of patients have musculoskeletal manifestations ⁴, including osteitis ⁵ and osteomyelitis ⁶⁻⁸
- Rare features include a transient truncal maculopapular rash, encephalopathy ⁹ or encephalitis ¹⁰ with seizures, lethargy, coma, parotitis ¹¹, cranial or peripheral nerve involvement, facial nerve paresis, myelitis ¹², uveitis or neuroretinitis ¹³⁻²⁵, optic neuritis ²⁶ with transient blindness, vitreal hemorrhage ²⁷, polyneuritis, radiculitis, Guillain-Barre syndrome ²⁸, disseminated visceral infection ^{29 30}, osteomyelitis ^{31 32}, endocarditis of native or prosthetic valves ³³⁻³⁸ or vascular prostheses ³⁹, hepatosplenomegaly with hepatic granulomata ⁴⁰, renal microabscesses ⁴¹, erythema marginatum, erythema multiforme, erythema nodosum ⁴² and thrombocytopenic purpura. ⁴³
- Scalp eschar with neck lymphadenopathy (SENLAT) has been reported in some cases ⁴⁴, and could be confused with tularemia or infection by *Rickettsia slovaca* or *Rickettsia raoultii*. ⁴⁵

- *B. henselae* accounts for 6.1% of bacterial species causing uveitis (2001 to 2007) ⁴⁶

29 cases of *Bartonella henselae* infection of solid-organ transplant recipients were reported to 2011 • many with disseminated disease. ⁴⁷

In one case, *Bartonella koehlerae* infection was associated with depression, anxiety, mood swings, severe headaches, muscle spasms, interphalangeal joint stiffness, decreased peripheral vision, diminished tactile sensation and hallucinations. ⁴⁸

This disease is endemic or potentially endemic to all countries.

References

1. Am J Clin Pathol 2004 Jun ;121 Suppl:S71-80.
2. Int J STD AIDS 2009 Aug ;20(8):585-6.
3. Case Report Med 2011 ;2011:183937.
4. Clin Infect Dis 2007 Dec 15;45(12):1535-40.
5. J Infect 2007 May ;54(5):417-21.
6. Pediatr Infect Dis J 2006 Dec ;25(12):1177-81.
7. Rev Med Interne 2009 Jul ;30(7):602-8.
8. Pediatr Radiol 2012 Jan ;42(1):116-9.
9. Emerg Med J 2008 Oct ;25(10):703-4.
10. Arch Pediatr 2012 Aug ;19(8):823-6.
11. Rev Stomatol Chir Maxillofac 2008 Jun ;109(3):183-6.
12. Clin Infect Dis 2007 Aug 15;45(4):e42-5.
13. Ocul Immunol Inflamm 2008 Jan-Feb;16(1):45-9.
14. Eur J Ophthalmol 2009 Mar-Apr;19(2):307-9.
15. Hong Kong Med J 2009 Oct ;15(5):391-3.
16. J AAPOS 2009 Dec ;13(6):602-4.
17. Clin Microbiol Infect 2009 Dec ;15 Suppl 2:132-3.
18. Int J Pediatr 2010 ;2010:763105.
19. Rev Med Interne 2011 Apr ;32(4):e46-8.
20. Int Ophthalmol 2010 Oct ;30(5):553-8.
21. Cornea 2011 Apr ;30(4):468-71.
22. Rev Chilena Infectol 2010 Oct ;27(5):417-22.
23. Clin Ophthalmol 2011 ;5:817-29.
24. Infection 2011 Aug 9;
25. Arch Pediatr 2012 Aug ;19(8):823-6.
26. Vojnosanit Pregl 2006 Nov ;63(11):971-4.
27. Int Ophthalmol 2011 Apr ;31(2):125-8.
28. Pediatr Infect Dis J 2006 Jan ;25(1):90-1.
29. Neth J Med 2008 Apr ;66(4):160-2.
30. J Heart Lung Transplant 2009 Jul ;28(7):736-9.
31. Semin Arthritis Rheum 2011 Dec ;41(3):511-6.
32. Clin Nucl Med 2012 Aug ;37(8):772-4.
33. Pediatr Infect Dis J 2009 Oct ;28(10):922-5.
34. J Heart Valve Dis 2011 Jan ;20(1):94-7.
35. Vector Borne Zoonotic Dis 2011 Nov ;11(11):1503-5.
36. J Card Surg 2011 Sep ;26(5):483-5.
37. J Card Surg 2012 Mar 1;
38. J Heart Valve Dis 2012 Sep ;21(5):682-5.
39. Ann Thorac Surg 2012 Apr ;93(4):e93-5.
40. Wien Klin Wochenschr 2006 Oct ;118(19-20):615-8.
41. Pediatr Infect Dis J 2010 May ;29(5):472-3.
42. Rev Med Interne 2011 Mar ;32(3):e34-6.
43. Infez Med 2008 Jun ;16(2):99-102.
44. Clin Infect Dis 2010 Feb 15;50(4):549-51.
45. J Med Case Reports 2011 ;5:108.
46. Medicine (Baltimore) 2008 May ;87(3):167-76.
47. Medicine (Baltimore) 2012 Mar ;91(2):111-21.
48. J Clin Microbiol 2011 Sep ;49(9):3415-7.

Bartonellosis - other systemic

Agent	BACTERIUM. <i>Bartonella quintana</i> , <i>B. koehlerae</i> , <i>B. elizabethae</i> , <i>B. tamiae</i> , <i>B. washoensis</i> , etc A fastidious gram-negative coccobacillus
Reservoir	Human Louse Rat Cat Dog Sheep
Vector	Louse (Pediculus) Flea - rare (Ctenocephalides, Pulex) Mite - rare (Dermanyssus)
Vehicle	Wound or eye contact with secretions/louse feces
Incubation Period	9d - 25d (range 4d - 35d)
Diagnostic Tests	Serology. Culture. Nucleic acid amplification.
Typical Adult Therapy	Doxycycline 100 mg PO BID X 3 to 5 days (if endocarditis, add Gentamicin 3 mg/kg daily X 28 days) Alternatives: Clarithromycin , Azithromycin , Gentamicin , Fluoroquinolone (Levofloxacin , Trovafoxacin , Pefloxacin , Sparfloxacin or Moxifloxacin)
Typical Pediatric Therapy	Erythromycin 10 mg/kg PO QID X 3 to 5 days. OR Gentamicin 2 mg/kg IM q12h. Alternatives: Clarithromycin , Azithromycin
Clinical Hints	Headache, myalgias, shin pain, macular rash, splenomegaly; endocarditis & bacteremia seen; relapse common; often associated with poor hygiene & crowding.
Synonyms	<i>Bartonella alsatica</i> , <i>Bartonella bovis</i> , <i>Bartonella capreoli</i> , <i>Bartonella doshiae</i> , <i>Bartonella elizabethae</i> , <i>Bartonella grahamii</i> , <i>Bartonella quintana</i> , <i>Bartonella rochalimae</i> , <i>Bartonella schoenbuchensis</i> , <i>Bartonella tamiae</i> , <i>Bartonella vinsonii</i> , <i>Bartonella vinsonii berkhoffii</i> , <i>Bartonella volans</i> , <i>Bartonella washoensis</i> , <i>Candidatus Bartonella mayotimonensis</i> , <i>Candidatus Bartonella melophagi</i> , <i>Candidatus Bartonella merieuxii</i> , <i>Candidatus Bartonella rochalimae</i> , Five day fever, His-Werner disease, Meuse fever, Quintan fever, Quintana fever, Shank fever, Shin fever, Shinbone fever, Trench fever, Volhynian fever. ICD9: 083.1 ICD10: A44.0,A44.8,A79.0

Clinical

Infection is characterized by abrupt onset of headache, postorbital pain, conjunctivitis, leg and back pain, relapsing fevers, splenomegaly and an erythematous maculopapular rash on the chest, back and abdomen. ¹

- In 50% of cases, as many as 3 to 8 relapses occur.

Subclinical bacteremia is common among immuno-competent persons with animal and arthropod contact.

No fatalities have been reported in classic trench fever.

Bartonella quintana (formerly *Rochalimaea quintana*) and related bacteria may also produce bacillary angiomatosis (discussed separately in this module), bacteremia, endocarditis ²⁻⁵, myocarditis ⁶, uveitis ^{7 8}, neuroretinitis ⁹ or chronic lymphadenopathy.

- *Bartonella* species other than *B. henselae* account for 8.1% of bacterial uveitis (France, 2008 publication) ¹⁰
- A single reported case of *Bartonella rochalimae* infection was characterized by fever, myalgia, headache and splenomegaly. ¹¹
- *Bartonella vinsonii* subsp *berkhoffii* genotype has been implicated in a case of epithelioid hemangioendothelioma. ¹²

This disease is endemic or potentially endemic to all countries.

References

1. *Vector Borne Zoonotic Dis* 2001 ;1(2):91-118.
2. *Emerg Infect Dis* 2002 Feb ;8(2):202-3.
3. *Curr Opin Infect Dis* 1998 Apr ;11(2):189-93.
4. *Ann N Y Acad Sci* 2009 May ;1166:120-6.
5. *Thorac Cardiovasc Surg* 2012 Jul ;60(5):363-5.
6. *J Med Case Rep* 2009 ;3:7325.
7. *Clin Microbiol Infect* 2009 Dec ;15 Suppl 2:132-3.
8. *Clin Ophthalmol* 2011 ;5:817-29.
9. *Med Glas Ljek komore Zenicko-doboj kantona* 2012 Aug ;9(2):435-7.
10. *Medicine (Baltimore)* 2008 May ;87(3):167-76.
11. *N Engl J Med* 2007 Jun 7;356(23):2381-7.
12. *J Clin Microbiol* 2009 Jun ;47(6):1957-60.

Bertiella and Inermicapsifer

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Anoplocephalidae: Bertiella spp. and Inermicapsifer spp.
Reservoir	Rodent Non-human primate
Vector	None
Vehicle	Mite (ingestion)
Incubation Period	Unknown
Diagnostic Tests	Identification of ova or proglottids in stool.
Typical Adult Therapy	Not established
Typical Pediatric Therapy	As for adult
Clinical Hints	Abdominal pain, vomiting, diarrhea or constipation following contact with primates.
Synonyms	Bertiella, Bertiella, Bertiella mucronata, Bertiella studeri, Bertielliasis, Inermicapsifer. ICD9: 123.8 ICD10: B71.8

Clinical

The few cases reported have ranged from asymptomatic infection to moderate abdominal pain, vomiting and diarrhea. ¹

- Symptoms may be intermittent or continuous. ²
- Adult tapeworms are known to live for at least two years.
- Diagnosis is based on finding worms, worm segments or ova in stool.

This disease is endemic or potentially endemic to 29 countries. Although Bertiella and Inermicapsifer is not endemic to Uganda, imported, expatriate or other presentations of the disease have been associated with this country.

Bertiella and Inermicapsifer in Uganda

Bertiella spp. have been identified in chimpanzees (*Pan troglodytes*) ³, guenons ⁴ and colobus monkeys in Uganda. ^{5 6}

References

1. Folia Parasitol (Praha) 1998 ;45(1):1-8.
2. Rev Inst Med Trop Sao Paulo 1997 Mar-Apr;39(2):123-7.
3. Am J Primatol 2012 Jul ;74(7):642-50.
4. J Parasitol 2004 Dec ;90(6):1356-60.
5. J Parasitol 2005 Jun ;91(3):569-73.
6. Am J Primatol 2008 Mar ;70(3):222-30.

Blastocystis hominis infection

Agent	PARASITE - Protozoa. Chromista, Bigyra, Blastocystea: Blastocystis hominis. [taxonomic status remains uncertain]
Reservoir	Human
Vector	None
Vehicle	Fecal-oral Water
Incubation Period	Unknown
Diagnostic Tests	Stool microscopy. Nucleic acid amplification.
Typical Adult Therapy	Nitazoxanide 500 mg BID X 3 d. OR Metronidazole 750 mg TID X 10d. OR Iodoquinol 650 mg TID X 20 d. or Sulfamethoxazole/trimethoprim
Typical Pediatric Therapy	Nitazoxanide - Age 1 to 3 years: 5 ml (100 mg) PO Q12h X 3 days - Age 4 to 11 years: 10 mg (200 mg) PO Q12h X 3 days; OR Metronidazole 15 mg/kg/d X 10d. Sulfamethoxazole/trimethoprim
Clinical Hints	Diarrhea and flatulence; usually no fever; illness similar to giardiasis; increased risk among immune-suppressed patients; the exact role of this organism in disease is controversial.
Synonyms	Apoi, Blastocystiose, Blastocystis hominis, Zierdt-Garavelli disease. ICD9: 007.8 ICD10: A07.8

Clinical

¹ Symptoms ascribed to blastocystosis include leucocyte-negative diarrhea, nausea, pain ², flatulence and abdominal distention. ^{3 4}

- Some reports suggest an association between urticaria and *Blastocystis* infection. ⁵⁻¹¹
- Symptoms usually last for 3 to 10 days, but may persist for weeks or months.
- *Blastocystis hominis* has also been implicated in the etiology of irritable bowel syndrome and urticaria ¹²⁻¹⁴, and may contribute to the development of anemia among infected pregnant women. ¹⁵

A search for alternative etiologies (including other infectious agents) should always be made in such patients. ^{16 17}

This disease is endemic or potentially endemic to all countries.

References

1. Clin Exp Dermatol 2011 Dec ;36(8):908-10.
2. J Pediatr Surg 2006 Aug ;41(8):1489-91.
3. Gastroenterol Clin North Am 2001 Sep ;30(3):797-815, x.
4. J Trop Med Hyg 1991 Apr ;94(2):118-22.
5. Am J Med Sci 2013 Jan 28;
6. Parasitol Res 2011 Mar ;108(3):553-60.
7. Acta Derm Venereol 2008 ;88(1):80-1.
8. Australas J Dermatol 2006 May ;47(2):117-9.
9. Acta Derm Venereol 2005 ;85(4):357-8.
10. Eur Rev Med Pharmacol Sci 2004 May-Jun;8(3):117-20.
11. Allergol Immunopathol (Madr) 1993 Jul-Aug;21(4):149-51.
12. Clin Microbiol Rev 2008 Oct ;21(4):639-65.
13. Parasitol Res 2011 Mar ;108(3):553-60.
14. Parasitol Res 2012 Mar ;110(3):1269-75.
15. Parasitol Res 2012 Jun ;110(6):2167-74.
16. Clin Microbiol Rev 1996 Oct ;9(4):563-84.
17. J Microbiol Immunol Infect 2008 Jun ;41(3):222-6.

Blastomycosis

Agent	FUNGUS. Ascomycota, Euascomycetes, Onygenales. <i>Blastomyces dermatitidis</i> . A dimorphic fungus
Reservoir	Soil Beaver Dog Rodent
Vector	None
Vehicle	Air
Incubation Period	14d - 44d (range 7d - 100d)
Diagnostic Tests	Microscopy and culture. Skin tests and serology not useful. Nucleic acid amplification.
Typical Adult Therapy	Itraconazole 200 to 400 mg PO daily X 6 months. OR Ketoconazole 400 mg/d X 6 months. OR Amphotericin B - total dose 2.0g. Excision as required
Typical Pediatric Therapy	Ketoconazole (if age >2) 5 mg/kg/d X 6 months. OR Amphotericin B - total cumulative dose 30 mg/kg
Clinical Hints	Acute or chronic lung infection, often complicated by hematogenous involvement of skin (verrucous or ulcerated skin/subcutaneous lesions), osteomyelitis of vertebrae or long bones, meningitis, prostatitis, etc.
Synonyms	<i>Blastomyces dermatitidis</i> , Blastomykose, Chicago disease, Gilchrist's disease, North American blastomycosis. ICD9: 116.0 ICD10: B40

Clinical

Blastomycosis typically presents as a flu-like illness and is often diagnosed as a pneumonia.

- Symptoms include sudden onset of fever, cough, chest pain, weight loss, hemoptysis, shortness of breath and fatigue. ¹
- Hematogenous, lymphatic, or macrophage-borne dissemination occur.
- Pulmonary involvement occurs in approximately 60%, and is manifest as airspace consolidation, focal masses, intermediate-sized nodules, interstitial disease, miliary disease, or cavitary lesions. ^{2 3}
- Cavities favor the upper lobes.
- 35% have involvement of both lung and skin; and 19% have infection of skin only.

Dissemination may involve the genitourinary tract, skin, liver, CNS ^{4 5}, spleen ⁶, bone, lymph nodes, heart, adrenals, GI tract, skeletal muscles ^{7 8} and pancreas.

- Central nervous system infection may present as epidural abscess, intracranial mass lesions or meningitis. ⁹
- A case of hypopituitarism associated with blastomycosis has been reported. ¹⁰

In chronic cutaneous blastomycosis the initial skin lesion presents as one or more verrucous or pustular nodules which eventually ulcerate. ^{11 12}

- Lesions are most common on exposed skin such as the face ¹³, hands, wrists, and lower legs.
- If untreated, elevated granulomatous lesions with advancing borders develop.
- Skeletal involvement has been reported in 33% of patients • therefore, an extensive radiographic examination is recommended for all patients with blastomycosis.

This disease is endemic or potentially endemic to 28 countries.

Blastomycosis in Uganda

A single case of blastomycosis had been reported in Uganda to 1988. ¹⁴⁻¹⁶

References

1. *Infect Dis Clin North Am* 2003 Mar ;17(1):21-40, vii.
2. *Radiographics* 2007 May-Jun;27(3):641-55.
3. *Mycopathologia* 2009 Mar ;167(3):115-24.
4. *Med Mycol* 2006 Nov ;44(7):659-64.
5. *Clin Neurol Neurosurg* 2011 Oct ;113(8):665-7.
6. *Can J Infect Dis Med Microbiol* 2010 ;21(1):53-6.
7. *Am J Med Sci* 2009 Apr ;337(4):293-4.
8. *Mycoses* 2010 Jul ;53(4):356-9.
9. *Clin Infect Dis* 2010 Mar 15;50(6):797-804.
10. *J Pediatr Endocrinol Metab* 2011 ;24(9-10):775-7.

11. *Int J Dermatol* 2008 Aug ;47(8):824-30.
12. *J Am Acad Dermatol* 2009 Aug ;61(2):355-8.
13. *Ophthal Plast Reconstr Surg* 2009 May-Jun;25(3):230-2.
14. *Mycopathologia* 1989 Jul ;107(1):25-32.
15. *S Afr Med J* 1989 Jul 1;76(1):13-6.
16. *Sabouraudia* 1964 Oct ;3(4):306-11.

Botulism

Agent	BACTERIUM. Clostridium botulinum . An anaerobic gram-positive bacillus
Reservoir	Soil Animal Fish
Vector	None
Vehicle	Food Occasionally soil (wound contamination)
Incubation Period	1d - 2d
Diagnostic Tests	Electrophysiologic (EMG) pattern. Isolation of organism from food (occ. from infant stomach). Mouse toxin assay
Typical Adult Therapy	Heptavalent (types A-G) or trivalent (types A, B, E) antitoxin [following test dose] 10 ml in 100 ml saline over 30 min Additional 10 ml at 2 and 4 hours if necessary. Respiratory support
Typical Pediatric Therapy	As for adult
Vaccine	Botulism antitoxin
Clinical Hints	Clinical manifestations similar to those of atropine poisoning: dysarthria, diplopia, dilated pupils, dry mouth, constipation, flaccid paralysis, etc); onset approximately 36 hrs after ingestion of poorly-preserved food.
Synonyms	Botulisme, Botulismo, Botulismus, Kerner's disease. ICD9: 005.1 ICD10: A05.1

Clinical

For reporting purposes, the CDC (The United States Centers for Disease Control) case definitions for Foodborne, Infant and Wound Botulism are as follows:

- 1) Neurological syndrome (diplopia, blurred vision, bulbar weakness, symmetric paralysis); or
- 2) Infant exhibiting constipation, poor feeding and failure to thrive, followed by progressive weakness, impaired respiration and death. ¹

Symptoms and signs of botulism reflect characteristic electrophysiological abnormalities ² and include diplopia ^{3 4}, blurred vision, ptosis, slurred speech, difficulty swallowing, dry mouth ⁵, and muscle weakness. Infants are lethargic, 'floppy,' constipated and feed poorly• exhibiting a weak cry and poor muscle tone. ^{6 7}

- In food-borne botulism, symptoms generally begin 18 to 36 hours after ingestion (range 6 hours to 10 days). ⁸
- Type F botulism is characterized by the appearance of respiratory failure within 24 hours, quadriplegia by the fifth day and rapid recovery beginning on the eighth day. ^{9 10}
- A case of asymmetric cranial nerve demyelination due to type F botulism has been reported. ¹¹
- If untreated, these symptoms progress to paralysis of the arms, legs, trunk and respiratory muscles.
- Patients who experience nausea and vomiting, cranial neuropathy or urinary retention are most likely to develop respiratory failure. ¹²
- Botulinum toxin may persist in the serum of patients for as long as 12 days. ¹³

Infant botulism should be suspected if a previously healthy infant (age <12 months) develops constipation and weakness in sucking, swallowing, or crying; hypotonia; and progressive bulbar and extremity muscle weakness. ¹⁴

- Approximately 50% of patients require mechanical ventilation.
- Lumbar puncture and brain imaging studies are usually normal, in contrast to other causes of flaccid weakness.
- The findings of infant botulism may mimic those of Hirschprung's disease ¹⁵ or acute abdomen. ¹⁶

This disease is endemic or potentially endemic to all countries.

Botulism in Uganda

Notable outbreaks:

2008 - An outbreak (3 cases, 1 fatal) was reported among high school students in Kampala. ^{17 18}

References

1. J Perinatol 2007 Mar ;27(3):175-80.
2. Muscle Nerve 2009 Aug ;40(2):271-8.
3. JAMA 1979 Feb 2;241(5):475-7.
4. Eye (Lond) 1994 ;8 (Pt 6):646-8.
5. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2011 Apr ;111(4):e15-8.
6. Semin Neurol 2004 Jun ;24(2):155-63.
7. Muscle Nerve 1998 Jun ;21(6):701-10.
8. Clin Infect Dis 2005 Oct 15;41(8):1167-73.
9. Neurology 2005 Dec 13;65(11):1694-700.
10. Emerg Infect Dis 2009 Jun ;15(6):969-71.
11. Emerg Infect Dis 2012 Jan ;18(1):102-4.
12. Am J Trop Med Hyg 2007 Aug ;77(2):386-9.
13. J Infect Dis 2009 Apr 1;199(7):1029-31.
14. ProMED <promedmail.org> archive: 20070420.1295
15. J Pediatr Surg 2009 Oct ;44(10):e5-7.
16. Infez Med 2009 Dec ;17(4):254-6.
17. ProMED <promedmail.org> archive: 20081113.3574
18. ProMED <promedmail.org> archive: 20081118.3639

Brain abscess

Agent	BACTERIUM OR FUNGUS. Mixed oral anaerobes / streptococci, <i>Staphylococcus aureus</i> (from endocarditis), etc.
Reservoir	Human
Vector	None
Vehicle	None
Incubation Period	Variable
Diagnostic Tests	Imaging techniques (CT, scan, etc).
Typical Adult Therapy	Antibiotic(s) appropriate to likely pathogens + drainage
Typical Pediatric Therapy	As for adult
Clinical Hints	Headache, vomiting and focal neurological signs; often associated with chronic sinusitis or otitis media, pleural or heart valve infection; patients are often afebrile.
Synonyms	Ascesso cerebrale, Cerebral abscess. ICD9: 324.0 ICD10: G06.0

Clinical

The clinical presentation of brain abscess may range from indolent to fulminant. ¹

- Most manifestations are due to the size and location of this space-occupying lesion within the brain and the virulence of the infecting microorganism, and not to infection per se.
- Headache is observed in approximately 70% of patients and may be moderate to severe and unilateral or generalized.
- Sudden worsening of the headache, accompanied by meningismus, may herald rupture of the abscess into the ventricular space.
- Less than 50% of patients present with a classic triad of fever, headache, and focal neurological deficit.
- Mental status changes are seen in 70% of cases, fever in 45 to 50%, seizures in 25 to 35%, vomiting in 25 to 50%, nuchal rigidity in 25% and papilledema in 25%.

Metastatic infections are most often associated with endocarditis, and may present with multiple abscesses.

- Although the distribution of the middle cerebral artery is most often involved, any part of the brain may be infected.
- Common pathogens in this setting reflect the usual flora of endocarditis and bacteremia.

Etiological associations:

- Congenital heart disease: viridans streptococci, *Haemophilus* spp.
- Endocarditis: *Staphylococcus aureus*, streptococci
- Immunodeficiency: Toxoplasmosis, *Nocardia*, fungi
- Otitis: Peptostreptococci, streptococci, Enterobacteriaceae
- Pleuropulmonary infection: anaerobes, *Nocardia*
- Sinusitis: Streptococci, Enterobacteriaceae, *Bacteroides*, *Haemophilus influenzae*
- Traumatic or post-surgical: *Staphylococcus aureus*, streptococci, Enterobacteriaceae

This disease is endemic or potentially endemic to all countries.

References

1. *Curr Neurol Neurosci Rep* 2004 Nov ;4(6):448-56.

Brucellosis

Agent	BACTERIUM. <i>Brucella abortus</i> , <i>Brucella melitensis</i> , <i>Brucella suis</i> , <i>Brucella canis</i> An aerobic gram-negative bacillus
Reservoir	Pig Cattle Sheep Goat Dog Coyote Caribou
Vector	None
Vehicle	Food Air Dairy products Animal excretions
Incubation Period	10d - 14d (range 5d - 60d)
Diagnostic Tests	Culture of blood or bone marrow. Serology. Note: Alert laboratory to possibility of Brucella.
Typical Adult Therapy	Doxycycline 100 mg BID + Rifampin 600 mg BID X 6 weeks. Alternatives Tetracycline + Gentamicin
Typical Pediatric Therapy	Rifampin 20 mg/kg/day (maximum 600 mg) plus: >age 8 years: Doxycycline 2 mg/kg BID PO X 6w age < 8 years Sulfamethoxazole/trimethoprim 4/20 mg/kg BID X 4 to 6w Add Gentamicin if severe
Clinical Hints	Prolonged fever, hepatosplenomegaly, lymphadenopathy, arthritis, osteomyelitis or chronic multisystem infection following ingestion of unpasteurized dairy products, contact with farm animals or meat processing.
Synonyms	Bang's disease, Bangsche Krankheit, Brucella, Brucellemia, Brucelliasis, Brucellose, Brucellosen, Brucellosi, Brucelose, Brucelosis, Cyprus fever, Febris melitensis, Febris sudoralis, Febris undulans, Fievre caprine, Gibraltar fever, Goat fever, Malta fever, Maltafieber, Melitococcosis, Neapolitan fever, Rock fever, Typhomalarial fever, Undulant fever. ICD9: 023 ICD10: A23

Clinical

For surveillance purposes the CDC (The United States Centers for Disease Control) case definition of brucellosis consists of "an illness characterized by acute or insidious onset of fever, night sweats, undue fatigue, weight loss, headache and arthralgia" associated with epidemiological or laboratory evidence for infection.

WHO Case definition for surveillance:

The WHO Case definition for surveillance is as follows:

Clinical description

- An illness characterized by acute or insidious onset, with continued, intermittent or irregular fever of variable duration, profuse sweating particularly at night, fatigue, anorexia, weight loss, headache, arthralgia and generalized aching. Local infection of various organs may occur

Laboratory criteria for diagnosis

- Isolation of *Brucella* spp. from clinical specimen or
- *Brucella* agglutination titer (e.g., standard tube agglutination tests: SAT>160) in one or more serum specimens obtained after onset of symptoms or
- ELISA (IgA, IgG, IgM), 2-mercaptoethanol test, complement fixation test, Coombs, fluorescent antibody test (FAT), and radioimmunoassay for detecting antilipopopolysaccharide antibodies; and counterimmunoelectrophoresis (CIEP)

Case classification

- Suspected: A case that is compatible with the clinical description and is epidemiologically linked to suspected or confirmed animal cases or contaminated animal products.
- Probable: A suspected case that has a positive Rose Bengal test.
- Confirmed: A suspected or probable case that is laboratory-confirmed.

Clinical manifestations: ¹

The clinical picture of brucellosis is nonspecific, and most often consists of fever, sweats, malaise, anorexia, headache, depression and back pain. ^{2 3}

- The fever of brucellosis may mimic that of enteric fever ⁴ ; and an undulant fever pattern is seen in chronic infections.
- Fever may be absent among patients with end-stage renal disease who acquire brucellosis. ⁵
- Mild lymphadenopathy is seen in 10 to 20% of patients; and splenomegaly or hepatomegaly in 20 to 30%. Rare instances of splenic rupture have been reported. ⁶
- Bone and joint infections are common ⁷⁻¹⁰ , including a high rate of vertebral osteomyelitis. ¹¹⁻¹⁴ Rare instances of acute or granulomatous myositis ¹⁵ , bursitis ¹⁶ and soft tissue or muscular abscesses have also been reported. ¹⁷⁻²⁰ Most cases of brucellar monoarthritis represent reactive rather than septic disease. ²¹ Infection of natural ²² or prosthetic joints ^{23 24}

and soft tissue has been reported. ²⁵

- Vertebral osteomyelitis is characterized by osteolysis, often associated with paravertebral masses, spondylodiscitis ^{26 27}, epidural abscess ²⁸⁻³⁰, or psoas abscesses. ³¹⁻³³
- Epididymo-orchitis is found in 7.6% to 12.7% of male patients with brucellosis. ³⁴⁻⁴⁰ Prostatitis has also been reported. ⁴¹
- Endocarditis is well documented ⁴²⁻⁵¹, including isolated case reports of *Brucella* infection of prosthetic valves ^{52 53} and devices such as implantable defibrillators ⁵⁴ and pacemaker leads. ⁵⁵ Rare instances of aortitis ^{56 57}, arterial thrombosis ⁵⁸, myocarditis ⁵⁹ and pericarditis are also reported. ⁶⁰⁻⁶⁴
- Pulmonary infiltrates ⁶⁵⁻⁶⁷, pleural effusion ⁶⁸, ileitis ⁶⁹, chest wall infection ⁷⁰, cholestatic jaundice ⁷¹, acalculous cholecystitis ⁷², peritonitis associated with dialysis ⁷³, and liver abscess have been reported. ^{74 75}
- Ocular manifestations include uveitis, visual loss due to suprasellar mass ⁷⁶, keratitis, conjunctivitis, papillitis, retinal hemorrhages and third-nerve palsy. ^{77 78}
- Neurological manifestations may include encephalitis ⁷⁹, meningitis ⁸⁰⁻⁸⁴, cranial ⁸⁵ or peripheral neuropathy ⁸⁶, progressive paraparesis ⁸⁷ or Guillain-Barre syndrome ⁸⁸, cerebral vasculitis with infarct ⁸⁹, intracranial hypertension or hydrocephalus ^{90 91}, psychosis ⁹², and parenchymal granulomata ⁹³ or abscesses. ⁹⁴⁻¹⁰¹
- Renal infection may present at hematuria, proteinuria, pyuria, overt nephritis or renal failure. ¹⁰² Rare instances of glomerulonephritis have also been reported. ¹⁰³⁻¹⁰⁵
- Persons working with animals may present with severe pharyngitis as an initial feature of brucellosis. ¹⁰⁶
- Abscesses involving a variety of body areas and solid organs may occur ¹⁰⁷⁻¹¹¹
- Various forms of rash occur in 6% to 13% of patients including generalized or localized papules or macules ¹¹², ulcers, purpura, vasculitis / leukocytoclastic vasculitis ¹¹³, panniculitis ¹¹⁴ and erythema nodosum ^{115 116}
- Brucellosis has been implicated in cases of human abortion. ^{117 118}


Virtually any organ or body system may be infected during the course of illness ¹¹⁹⁻¹³⁰

- Chronic brucellosis generally represents persistence of local infection in bone, joints, liver, spleen or kidneys.
- Relapses are common, especially following inadequate therapy.
- Pancytopenia is reported in 15% of cases ^{131 132}
- Brucellosis has been reported to cause myelofibrosis ¹³³, and to trigger hemolytic anemia in patients with Glucose-6-Phosphate Dehydrogenase deficiency. ¹³⁴
- Isolated thrombocytopenia mimicking ITP is reported in 6% of cases ¹³⁵⁻¹⁴⁰; hepatic dysfunction ¹⁴¹, colitis ¹⁴² Coombs-positive hemolytic anemia ^{143 144}, reactive hemophagocytic ^{145 146} or myelodysplastic syndrome ¹⁴⁷, pancytopenia ¹⁴⁸, disseminated intravascular coagulation ¹⁴⁹, TTP ^{150 151}, Guillain-Barre syndrome ¹⁵² and syndrome of inappropriate secretion of antidiuretic hormone (SIADH) have also been documented. ¹⁵³

This disease is endemic or potentially endemic to 179 countries.

Brucellosis in Uganda

Human disease in this country is due to *Brucella abortus* and *B. melitensis*. ¹⁵⁴



Graph: Uganda. Brucellosis, cases

Notes:

1. The rate of brucellosis in urban Kampala was estimated at 5.8 per 100,000 per year (2010 publication). ¹⁵⁵

652 cases of brucellosis were treated at a referral hospital in Kampala during 2004 to 2006 - most associated with consumption of raw milk transported from peri-urban and rural areas of Kampala and/or dairy production areas outside Kampala. ¹⁵⁶

- There is a female predominance among hospitalized brucellosis cases in Kampala (2011 publication) ¹⁵⁷

Prevalence surveys:

6.5% of cattle herds in Kampala (2011 publication) ¹⁵⁸

Seroprevalence surveys:

55.6% of cattle herds in the Mbarra milk basin (2002) ¹⁵⁹

21.5% of cattle in Kumi and Mbale districts (western region) and 3.4% in Kiboga, Mpigi and Kiruhura districts (eastern region) (2008 to 2009) ¹⁶⁰

100% of herds and 34.0% of cattle in the pastoral system; 5.5% of herds and 3.3% of cattle in the zerograzing system (2006 to 2007) ¹⁶¹

38% of livestock in Makerere, 32% in Entebbe and 7% in Tororo (2010 publication) ¹⁶²

12.6% of milk marketed in urban Kampala (2010 publication) ¹⁶³

References

1. PLoS Negl Trop Dis 2012 Dec ;6(12):e1929.
2. J Commun Dis 2002 Dec ;34(4):287-301.
3. Int J Infect Dis 2002 Sep ;6(3):182-6.
4. J Infect Dev Ctries 2009 ;3(3):239-40.
5. Nephrol Dial Transplant 2008 Jul ;23(7):2344-9.
6. Chang Gung Med J 2011 Nov ;34(6 Suppl):52-5.
7. J Clin Rheumatol 2004 Dec ;10(6):300-7.
8. Rheumatol Int 2011 Jun ;31(6):721-4.
9. Intern Med 2011 ;50(5):421-8.
10. Semin Musculoskelet Radiol 2011 Nov ;15(5):470-9.
11. Clin Infect Dis 2008 Feb 1;46(3):426-33.
12. Mikrobiyol Bul 2009 Jan ;43(1):141-5.
13. Radiol Med 2010 Aug ;115(5):794-803.
14. Rheumatol Int 2012 Jul 29;
15. Muscle Nerve 2012 Feb ;45(2):290-3.
16. J Med Microbiol 2010 Dec ;59(Pt 12):1514-8.
17. Intern Med 2008 ;47(23):2091-3.
18. Int J Infect Dis 2009 Nov ;13(6):e485-7.
19. Ann Saudi Med 2011 May-Jun;31(3):311-3.
20. J Med Case Reports 2011 ;5:125.
21. Rheumatol Int 2012 May ;32(5):1465-8.
22. Chir Organi Mov 2009 Sep ;93(2):75-8.
23. Diagn Microbiol Infect Dis 2007 Aug ;58(4):481-5.
24. Rev Med Suisse 2007 Apr 18;3(107):1007-9.
25. Intern Med 2008 ;47(3):171-2.
26. Neurol Neurochir Pol 2010 Sep-Oct;44(5):516-9.
27. Rheumatol Int 2012 Nov 4;
28. Cases J 2009 ;2:7614.
29. J Infect Dev Ctries 2011 May ;5(5):403-5.
30. BMJ Case Rep 2012 ;2012
31. Clin Infect Dis 2008 Feb 1;46(3):426-33.
32. Trop Doct 2009 Apr ;39(2):124-7.
33. J Back Musculoskelet Rehabil 2009 ;22(2):121-3.
34. Int J Infect Dis 2006 Mar ;10(2):171-7.
35. Int Urol Nephrol 2006 ;38(3-4):637-9.
36. Diagn Microbiol Infect Dis 2007 Apr ;57(4):367-72.
37. Urol Int 2009 ;82(4):481-3.
38. Urologe A 2011 Jan ;50(1):71-3.

39. BMC Res Notes 2011 ;4:286.
40. J Infect Chemother 2012 Oct ;18(5):760-3.
41. Mikrobiyol Bul 2009 Jul ;43(3):493-7.
42. Eur J Clin Microbiol Infect Dis 2003 Nov ;22(11):647-50.
43. Circ J 2008 Dec ;72(12):2096-7.
44. Int J Cardiol 2009 Jan 24;131(3):e87-9.
45. Can J Cardiol 2006 Sep ;22(11):971-4.
46. Turk Kardiyol Dern Ars 2008 Jul ;36(5):329-31.
47. Rev Port Cardiol 2008 Oct ;27(10):1309-15.
48. Trop Doct 2009 Apr ;39(2):85-8.
49. Clin Cardiol 2010 Feb ;33(2):E20-6.
50. Turk J Gastroenterol 2009 Jun ;20(2):135-7.
51. Int J Antimicrob Agents 2012 Aug ;40(2):145-50.
52. Int Urol Nephrol 2011 Jan 15;
53. J Cardiovasc Med (Hagerstown) 2009 Mar ;10(3):257-8.
54. Clin Infect Dis 2007 Feb 15;44(4):e37-9.
55. Saudi Med J 2010 Apr ;31(4):448-50.
56. Vasa 2011 Mar ;40(2):150-6.
57. Vector Borne Zoonotic Dis 2012 Oct ;12(10):827-40.
58. Case Rep Infect Dis 2012 ;2012:581489.
59. Intern Med 2012 ;51(8):901-4.
60. Intern Med 2009 ;48(19):1773-4.
61. J Med Case Reports 2011 ;5(1):22.
62. Ren Fail 2011 ;33(3):367-70.
63. Case Rep Infect Dis 2013 ;2013:796437.
64. Int J Infect Dis 2013 Feb 11;
65. Vector Borne Zoonotic Dis 2008 Apr ;8(2):245-8.
66. Respir Care Clin N Am 2004 Mar ;10(1):99-109.
67. Afr Health Sci 2011 Aug ;11 Suppl 1:S112-6.
68. Acta Med Iran 2011 ;49(5):325-6.
69. Am J Gastroenterol 1988 Jan ;83(1):80-2.
70. J Glob Infect Dis 2011 Jan ;3(1):86-8.
71. Int J Infect Dis 2010 Sep ;14 Suppl 3:e322-4.
72. J Infect Dev Ctries 2010 Jul ;4(7):464-7.
73. Perit Dial Int 2012 Mar ;32(2):126-30.
74. Eur J Pediatr 2008 Jun ;167(6):699-700.
75. Asian J Surg 2007 Oct ;30(4):283-5.
76. Pediatr Neurol 2009 May ;40(5):401-3.
77. Clin Infect Dis 2008 May 1;46(9):1338-45.
78. Can J Ophthalmol 2009 Oct ;44(5):598-601.
79. Saudi Med J 2006 Apr ;27(4):539-41.
80. Arch Iran Med 2008 Jan ;11(1):21-5.
81. Eur J Pediatr 2006 Oct ;165(10):726-7.
82. Trop Doct 2009 Oct ;39(4):233-5.
83. Turk J Pediatr 2010 Jul-Aug;52(4):426-9.
84. Arch Iran Med 2010 Nov ;13(6):486-91.
85. South Med J 2009 Aug ;102(8):855-7.
86. Eur Neurol 2009 ;61(1):33-8.
87. Rev Neurol (Paris) 2011 Feb ;167(2):181-4.
88. J Glob Infect Dis 2011 Oct ;3(4):390-2.
89. Int J Infect Dis 2010 Sep ;14 Suppl 3:e202-4.
90. J Pediatr Neurosci 2010 Jul ;5(2):144-6.
91. J Child Neurol 2011 Oct ;26(10):1316-8.
92. Arch Iran Med 2012 Nov ;15(11):723-5.
93. Eur Spine J 2007 Dec ;16 Suppl 3:255-9.
94. Med Sci Monit 2006 Dec ;12(12):CS119-122.
95. J Clin Neurosci 2006 May ;13(4):485-7.
96. Int J Infect Dis 2004 Nov ;8(6):379-81.
97. J Korean Neurosurg Soc 2008 Jan ;43(1):37-40.
98. Int J Infect Dis 2009 Nov ;13(6):e339-43.
99. Cases J 2009 ;2:6698.
100. Mikrobiyol Bul 2011 Jul ;45(3):401-10.
101. J Infect Chemother 2012 Oct ;18(5):767-70.
102. Urology 2009 Jun ;73(6):1179-83.
103. Int Urol Nephrol 2011 Jan 15;
104. Ren Fail 2011 ;33(3):367-70.
105. J Infect Dev Ctries 2011 Dec ;5(12):893-5.
106. Occup Med (Lond) 2008 Jun ;58(4):305-7.
107. Eur J Gastroenterol Hepatol 2008 Apr ;20(4):349-52.
108. J Med Microbiol 2009 Feb ;58(Pt 2):267-9.
109. Mikrobiyol Bul 2009 Jan ;43(1):141-5.
110. Gastroenterol Hepatol 2009 Apr ;32(4):291-3.
111. Cases J 2009 ;2:7143.
112. Am J Dermatopathol 2009 Oct ;31(7):687-90.
113. Indian J Dermatol 2011 May ;56(3):339-40.
114. Int J Dermatol 2011 Jul 26;
115. Am J Dermatopathol 2008 Apr ;30(2):169-71.
116. J Coll Physicians Surg Pak 2009 Dec ;19(12):794-5.
117. J Obstet Gynaecol Res 2010 Apr ;36(2):418-23.
118. Recent Pat Antiinfect Drug Discov 2012 Jul 17;
119. Respir Care Clin N Am 2004 Mar ;10(1):99-109.
120. Eur J Pediatr 2006 Oct ;165(10):726-7.
121. Breast J 2006 Jul-Aug;12(4):375-6.
122. Indian J Med Microbiol 2006 Oct ;24(4):286-8.
123. Virchows Arch 2008 Jan ;452(1):97-101.
124. Neurol Sci 2008 Dec ;29(6):481-3.
125. Eur Neurol 2009 ;61(1):33-8.
126. Tuberk Toraks 2008 ;56(4):443-7.
127. Mikrobiyol Bul 2009 Jan ;43(1):141-5.
128. Annu Rev Biochem 1998 ;67:181-98.
129. Ann Thorac Surg 2010 Jun ;89(6):2038-40.
130. J Med Case Reports 2011 ;5:125.
131. Am J Hematol 2008 Apr ;83(4):334-9.
132. Arch Med Sci 2011 Feb ;7(1):173-5.
133. Int J Infect Dis 2010 Feb ;14(2):e158-60.
134. Med Princ Pract 2009 ;18(4):329-31.
135. Int J Lab Hematol 2007 Dec ;29(6):442-5.
136. Infez Med 2008 Sep ;16(3):158-61.
137. Clin Appl Thromb Hemost 2011 Nov-Dec;17(6):E36-8.
138. J Med Liban 2010 Oct-Dec;58(4):241-3.
139. Arch Iran Med 2012 May ;15(5):303-6.
140. Intern Med 2012 ;51(23):3291-3.
141. Infez Med 2008 Sep ;16(3):148-53.
142. Mil Med 2008 Nov ;173(11):1145-7.
143. Intern Med 2008 ;47(11):1043-5.
144. Ann Pharmacother 2010 Oct ;44(10):1677-80.
145. J Microbiol Immunol Infect 2010 Apr ;43(2):159-62.
146. Indian J Pediatr 2010 Dec ;77(12):1434-6.
147. Med Princ Pract 2012 ;21(2):183-5.
148. Pediatr Hematol Oncol 2011 Apr ;28(3):203-8.
149. Clin Appl Thromb Hemost 2011 Nov-Dec;17(6):E10-2.
150. Am J Med Sci 2008 Mar ;335(3):230-2.
151. Clin Appl Thromb Hemost 2011 Jun ;17(3):245-7.
152. Rev Peru Med Exp Salud Publica 2010 Jun ;27(2):292-5.
153. Case Report Med 2010 ;2010
154. Prev Vet Med 2001 Dec 3;52(2):91-108.
155. PLoS One 2010 ;5(12):e14188.
156. Ann N Y Acad Sci 2008 Dec ;1149:309-11.
157. Int J Health Geogr 2011 ;10:52.
158. BMC Vet Res 2011 ;7:60.
159. Prev Vet Med 2005 Mar 15;67(4):267-81.
160. J Vet Med Sci 2012 Jun 18;
161. Trop Anim Health Prod 2009 Dec ;41(8):1765-71.
162. Trop Anim Health Prod 2011 Mar ;43(3):603-8.
163. PLoS One 2010 ;5(12):e14188.

Bunyaviridae infections - misc.

Agent	VIRUS - RNA. Bunyaviridae, Orthobunyavirus. Over 30 strains have been associated with human disease (see Synonyms)
Reservoir	Rat Bird Marsupial Chipmunk Cattle Sheep Horse Bat
Vector	Mosquito (exceptions: Shuni is transmitted by culicoid flies; Bhanja, Tamdy, Wanowrie and Zirqa by ticks)
Vehicle	None
Incubation Period	3d - 12d
Diagnostic Tests	Serology and virus isolation. Nucleic acid amplification. Biosafety level 2 or 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Abrupt onset of fever, chills, headache; photophobia, rash arthralgia, myalgia, vomiting, diarrhea or cough may be present; meningitis or myocarditis may occur with Bwamba virus; usual course 2 to 7 days.
Synonyms	Bangui, Batai, Bhanja, Bunyamwera, Bwamba, Cache Valley, Calovo, Catu, Fort Sherman, Garissa, Germiston, Guama, Hartland virus, HNF virus, Huaiyangshan virus, Ilesha, Ingwavuma, Kairi, Lumbo, Ngari, Northway, Nyando, Pongola, Severe fever with thrombocytopenia syndrome, SFTSV, Shokwe, Shuni, Tacaiuma, Tamdy, Tataguine, Tensaw, Wanowrie, Wyeomyia, Zirqa. ICD9: 066.3 ICD10: A93.8

Clinical

As a group, these diseases are characterized by acute febrile illness occurring in persons exposed to wild or forest environments.

- Additional features may include headache, myalgia, arthralgia, rash or aseptic meningitis.

Avalon virus has been implicated in isolated cases of conjunctivitis ¹ and polyradiculitis. ^{2 3}

Ilesha virus infection may be associated with fever or rash, or hemorrhagic fever.

Huaiyangshan virus (Severe fever with thrombocytopenia syndrome bunyavirus, SFTSV) appears to infect primarily farmers, above age 50. ⁴⁻⁷

- Contact with the blood of patients appears to facilitate spread of the disease. ⁸
- Clinical features include fever, conjunctivitis, diarrhea, abdominal pain, thrombocytopenia, leukopenia, proteinuria and hematuria. ⁹
- In one series, 8 of 49 hospitalized infections (16.3%) were fatal. ¹⁰

A novel Bunyavirus ("heartland virus") has been associated with cases of fever, fatigue, diarrhea, thrombocytopenia, and leukopenia • following tick bites. ^{11 12}

This disease is endemic or potentially endemic to 90 countries.

Bunyaviridae infections - misc. in Uganda

Bunyamwera ¹³ , Bwamba and Pongola ¹⁴ viruses are encountered.

Seroprevalence surveys:

20.3% of the general population (Bunyamwera virus, 1967 to 1969, eight areas) ¹⁵

37% to 44% of the general population (Bwamba virus, vector *Anopheles funestus*) ¹⁶⁻¹⁸

References

1. J Fr Ophtalmol 1984 ;7(6-7):457-62.
2. Presse Med 1985 Sep 14;14(30):1616.
3. Presse Med 1985 Sep 14;14(30):1616.
4. N Engl J Med 2011 Apr 21;364(16):1523-32.
5. Emerg Infect Dis 2012 Jun ;18(6):963-5.
6. ProMED <promedmail.org> archive: 20110317.0853
7. ProMED <promedmail.org> archive: 20110619.1880
8. J Infect Dis 2012 Dec 6;
9. J Virol 2012 May 23;
10. Clin Infect Dis 2012 Feb 15;54(4):527-33.
11. N Engl J Med 2012 Aug 30;367(9):834-41.
12. ProMED <promedmail.org> archive: 20120830.1274007
13. Trans R Soc Trop Med Hyg 1953 Jan ;47(1):13-48.
14. Trans R Soc Trop Med Hyg 1985 ;79(4):567.
15. Bull World Health Organ 1970 ;42(5):797-805.
16. Am J Trop Med Hyg 1999 Jul ;61(1):158-62.
17. Ann Trop Med Parasitol 1952 May ;46(1):61-7.
18. Afr Health Sci 2002 Apr ;2(1):24-8.

Campylobacteriosis

Agent	BACTERIUM. <i>Campylobacter jejuni</i> subsp <i>jejuni</i> , et al A microaerophilic gram-negative bacillus
Reservoir	Human Mammal Bird
Vector	None
Vehicle	Water Food
Incubation Period	2d - 4d (range 1d - 10d)
Diagnostic Tests	Stool (rarely blood, CSF) culture. Nucleic acid amplification. Alert laboratory when these organisms are suspected.
Typical Adult Therapy	Stool precautions. Erythromycin 500 mg QID X 7d. Alternatives Azithromycin , Fluoroquinolone (Levofloxacin , Trovaflaxacin , Pefloxacin , Sparflaxacin or Moxifloxacin), Gentamicin
Typical Pediatric Therapy	Stool precautions. Erythromycin 12.5 mg/kg PO QID X 7d. Alternatives - Azithromycin , Gentamicin
Clinical Hints	Febrile diarrhea or dysentery; vomiting or bloody stool often noted; severe abdominal pain may mimic appendicitis; disease is most common among children and lasts one to four days.
Synonyms	Campylobacter. ICD9: 008.43 ICD10: A04.5

Clinical

Following an incubation period of 1 to 10 days, patients develop diarrhea (often bloody) and abdominal pain.

- Initial symptoms of malaise, dizziness, fever, headache and myalgia are common.
- Vomiting is unusual.
- Leucocytes are usually seen on stool smears.

Infection may be complicated by cholecystitis ¹, pseudoappendicitis, peritonitis ² (including peritonitis associated with dialysis ^{3 4}), hemolytic-uremic syndrome, bacteremia ⁵⁻⁷, myocarditis ⁸⁻¹¹, endocarditis ¹², pericarditis ^{13 14}, pleurisy ^{15 16}, mycotic iliac ¹⁷ and aortic aneurysms ¹⁸, meningitis ¹⁹, encephalopathy ²⁰, epidural abscess ^{21 22}, septic arthritis ²³, cellulitis ²⁴, Sweet's syndrome ²⁵, spontaneous abortion, reactive arthritis or Guillain-Barre syndrome.

- Reactive arthritis has been reported in 1% to 13% of cases ^{26 27}
- The risk for reactive arthritis following *Campylobacter* infection was 2.1/100,000 cases (United States, 2002 to 2004) ²⁸
- Elderly patients are at risk for complicated or fatal infection. ²⁹

Guillain Barre syndrome (GBS) has been estimated to complicate 0.1% of *Campylobacter* infections. ³⁰⁻³⁴

- *Campylobacter* infection is implicated in 14% to 40% of GBS episodes. ³⁵⁻⁴⁰
- Risk for GBS continues for up to 2 months following an episode of Campylobacteriosis.
- The rate of GBS is 19.2 per 100,000 episodes of Campylobacteriosis. ⁴¹
- There have been case reports of brain stem encephalitis ⁴², cranial neuropathy ⁴³, acute transverse myelitis ⁴⁴ and demyelization of the central nervous system or spinal cord following *C. jejuni* infection. ⁴⁵
- There is evidence that campylobacteriosis may increase the risk for later development of inflammatory bowel disease. ⁴⁶

This disease is endemic or potentially endemic to all countries.

Campylobacteriosis in Uganda

Prevalence surveys:

- 9.3% of children with acute diarrhea (Kampala, 2005) ⁴⁷
- 9.3% of children with acute diarrhea (Kampala, 2010 publication) ⁴⁸

References

1. Int J Med Sci 2009 ;6(6):374-5.
2. J Clin Microbiol 2010 Jan ;48(1):336-7.
3. Perit Dial Int 2010 Jan-Feb;30(1):99-104.
4. Perit Dial Int 2012 Oct 2;

5. Clin Microbiol Infect 2010 Jan ;16(1):57-61.
6. Medicine (Baltimore) 2010 Sep ;89(5):319-30.
7. Clin Infect Dis 2011 Oct ;53(8):e99-e106.
8. Resuscitation 2008 Oct ;79(1):165-7.
9. Eur J Pediatr 2010 Jan ;169(1):63-5.
10. Scand J Infect Dis 2009 ;41(6-7):528-31.
11. Wien Klin Wochenschr 2010 May ;122(9-10):315-9.
12. Tex Heart Inst J 2011 ;38(5):584-7.
13. Int J Cardiol 2010 Sep 24;144(1):e14-6.
14. Ir J Med Sci 2011 Sep ;180(3):753-5.
15. J Clin Microbiol 2007 Jul ;45(7):2334-6.
16. Intern Med 2010 ;49(22):2481-6.
17. Vascular 2009 Jul-Aug;17(4):226-9.
18. Surg Today 2009 ;39(2):137-40.
19. Scand J Infect Dis 1996 ;28(3):269-70.
20. Neurol Sci 2012 Feb ;33(1):155-8.
21. J Clin Microbiol 2009 Mar ;47(3):857-8.
22. Muscle Nerve 2009 Nov ;40(5):875-9.
23. J Clin Microbiol 2009 Oct ;47(10):3370-1.
24. Acta Clin Belg 2009 Jul-Aug;64(4):346-8.
25. J Med Microbiol 2012 Oct ;61(Pt 10):1473-5.
26. J Rheumatol 2008 Mar ;35(3):480-7.
27. J Rheumatol 1983 Feb ;10(1):107-8.
28. Ann Rheum Dis 2008 Dec ;67(12):1689-96.
29. J Infect 2008 Sep ;57(3):214-22.
30. Clin Microbiol Rev 1998 Jul ;11(3):555-67.
31. J Peripher Nerv Syst 2009 Jun ;14(2):72-4.
32. Emerg Infect Dis 2009 Aug ;15(8):1315-7.
33. Curr Allergy Asthma Rep 2011 Jun ;11(3):197-204.
34. Curr Gastroenterol Rep 2012 Oct ;14(5):395-405.
35. J Health Popul Nutr 2010 Dec ;28(6):545-52.
36. Emerg Infect Dis 2006 Jun ;12(6):990-3.
37. N Engl J Med 1995 Nov 23;333(21):1374-9.
38. Clin Microbiol Rev 1998 Jul ;11(3):555-67.
39. J Child Neurol 2009 Jun ;24(6):664-8.
40. J Neurol Neurosurg Psychiatry 2011 Mar ;82(3):300-5.
41. J Infect Dis 2006 Jul 1;194(1):95-7.
42. J Clin Pathol 2007 Oct ;60(10):1161-2.
43. Rinsho Shinkeigaku 2007 Jan ;47(1):53-5.
44. J Clin Neurosci 2012 Feb ;19(2):316-8.
45. Spinal Cord 2007 Oct ;45(10):690-4.
46. Gastroenterology 2009 Aug ;137(2):495-501.
47. Afr Health Sci 2009 Sep ;9(3):201-5.
48. Tanzan J Health Res 2010 Jan ;12(1):100-3.

Candidiasis

Agent	FUNGUS - Yeast. Ascomycota, Hemiascomycetes, Saccharomycetales. Candida albicans , and other species.
Reservoir	Human
Vector	None
Vehicle	Contact Catheter
Incubation Period	Variable
Diagnostic Tests	Culture. Serology and assays for cell-specific antigens are performed in some centers,
Typical Adult Therapy	Topical, oral, systemic antifungal agent depending on clinical presentation and species [in Therapy module, scroll through upper left box]
Typical Pediatric Therapy	As for adult
Clinical Hints	Dermal erythema with satellite pustules; "cheesy" mucosal discharge; severe, widespread or intractable disease should suggest the possibility of underlying diabetes, AIDS or other form of immune suppression.
Synonyms	Candida, Candida-Mykosen, Candidiase, Candidiasi, Candidose, Monilia, Moniliasis, Salmonella, Thrush. ICD9: 112 ICD10: B37

Clinical

The clinical features of candidiasis range from localized mucosal or skin inflammation to multi-organ candidal sepsis.

Often infection represents overgrowth of *Candida* species following use of antimicrobial agents, or in the presence of the high mucosal glucose concentrations found in diabetics.

- Other predisposing factors include chronic intertrigo, oral contraceptive use, and cellular immune deficiency.
- Candidiasis is a common initial event in HIV-infected individuals.
- White exudative plaques may occur on the tongue or buccal mucosa (thrush), vaginal or rectal mucosa.
- Fissured, macerated lesions at the corners of the mouth (perleche) are common among individuals with poorly-fitting dentures. In fact, candidal infections have a predilection for sites that are chronically wet and macerated.
- Intertriginous lesions are edematous, erythematous, and scaly; and associated with scattered "satellite pustules." ¹
- The glans penis and scrotum as inner aspect of the thighs are often involved.

Systemic *Candida* infections may involve virtually any organ or organ system, and mimic bacterial sepsis. ^{2 3}

- Case fatality rates for infected vascular catheters range from 26% to 38%; 33% for infected prosthetic cardiac valves; 20% to 40% for urinary catheters.

This disease is endemic or potentially endemic to all countries.

Candidiasis in Uganda

Prevalence surveys:

18.6% of women with genital disease (southwestern Uganda, 1999 to 2001) ⁴

References

1. *Infect Dis Clin North Am* 2002 Dec ;16(4):793-820, v.
2. *Infect Dis Clin North Am* 2002 Dec ;16(4):821-35.

3. *Case Report Ophthalmol* 2012 Sep ;3(3):277-82.
4. *Sex Transm Infect* 2005 Dec ;81(6):488-93.

Chancroid

Agent	BACTERIUM. <i>Haemophilus ducreyi</i> . A facultative gram-negative bacillus
Reservoir	Human
Vector	None
Vehicle	Sexual contact
Incubation Period	3d - 10d (2d - 21d)
Diagnostic Tests	Culture (inform laboratory when this diagnosis is suspected). Fluorescent staining under development
Typical Adult Therapy	Azithromycin 1.0 g PO X 1 dose. OR Ceftriaxone 250 mg IM X 1 dose. OR Ciprofloxacin 500 mg PO BID X 3 days OR Erythromycin 500 mg PO TID X 7d.
Typical Pediatric Therapy	Azithromycin 12 mg/kg PO X 1 dose OR Erythromycin 10 mg/kg PO TID X 7d. OR Ceftriaxone 10 mg/kg IM X 1
Clinical Hints	Soft, painful and tender chancre on erythematous base, with regional lymphadenopathy (generally unilateral and painful); onset 3 to 10 days following sexual exposure.
Synonyms	Blot sjanker, Chancre mou, Chancro blando, <i>Haemophilus ducreyi</i> , Nkumunye, Soft chancre, Ulcera mole, Ulcus molle, Weeke sjanker, Weicher Schanker. ICD9: 099.0 ICD10: A57

Clinical

For surveillance the CDC (The United States Centers for Disease Control) case definition consist of a sexually-transmitted disease characterized by painful genital ulceration and inflammatory inguinal adenopathy; but without evidence for *Treponema pallidum* by dark field and serological examination (after at least 7 days) and without clinical or laboratory evidence for herpes simplex infection.

Infection begins with a papule or pustule which ulcerates and enlarges over a period of 1 to 2 days. ¹

- The lesion is soft, painful and bleeds easily; and the ulcer edges are undermined and irregular. ²
- Two thirds of patients present with more than one ulcer
- Painful unilateral or bilateral lymphadenopathy is present in 40% of cases.
- Systemic signs are unusual.
- Extragenital skin ulcers are occasionally encountered. ^{3 4}
- *Haemophilus ducreyi* has been associated with esophageal ulceration in HIV-positive patients. ⁵

This disease is endemic or potentially endemic to all countries.

Chancroid in Uganda

Prevalence surveys:

9.8% of men and 7.3% of women ages 15 to 54 in the rural southwest (1999 publication) ⁶

2% of patients with genital ulcer disease (rural Rakai District, 2008 publication) ⁷

References

1. *Sex Transm Infect* 2003 Feb ;79(1):68-71.
2. *Curr Opin Infect Dis* 2002 Feb ;15(1):43-7.
3. *Clin Infect Dis* 2007 May 15;44(10):e85-7.
4. *Med J Aust* 2010 Mar 15;192(6):348-50.
5. *Int J STD AIDS* 2009 Apr ;20(4):238-40.
6. *Sex Transm Infect* 1999 Apr ;75(2):98-102.
7. *Sex Transm Infect* 2009 Apr ;85(2):97-101.

Chikungunya

Agent	VIRUS - RNA. Togaviridae, Alphavirus: Chikungunya virus. Related Semliki Forest and Me Tri viruses are found in Africa & Asia
Reservoir	Non-human primate
Vector	Mosquito (Aedes spp. ; Ae. fuscifer-taylori group in Africa)
Vehicle	None
Incubation Period	2d - 12d
Diagnostic Tests	Viral culture (blood). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Abrupt fever, leukopenia, myalgia and prominent bilateral joint pain; maculopapular rash appears on 2nd to 5th days in greater than 50% of cases; fever resolves within 7 days, but joint pain may persist for months.
Synonyms	Buggy Creek, Getah, Knuckle fever, Me Tri, Semliki Forest. ICD9: 062.8,066.3 ICD10: A92.1

Clinical

The fever of Chikungunya is characterized by a rapid rise in temperature to as high as 40 C, often accompanied by rigors, myalgia, headache, photophobia, retro-orbital pain, sore throat with objective signs of pharyngitis, nausea, and vomiting. ¹

- Fever may abate after a few days, only to recrudescence ("saddle-back" fever curve").
- Polyarthralgia occurs in 70% of cases, favors small joints and sites of previous injury, and is most intense on arising.
- Joints may swell, but without significant fluid accumulation. ^{2 3}
- Joint pain is most severe in adults.
- Symptoms may last for from 1 week to several months. ⁴
- Joint involvement may progress to residual chronic pain ^{5 6} or destructive arthritis. ⁷
- Arthralgia may persist for as long as 24 months. ⁸⁻¹⁴ In one series, 57% of patients continued to experience rheumatological symptoms for 15 months or more ^{15 16} ; and in another, 66.5% reported myalgia, asthenia or arthralgia for more than 12 months. ¹⁷
- Imaging studies may reveal joint effusion, bony erosion, marrow edema, synovial thickening, tendonitis and tenosynovitis. ¹⁸
- Laboratory tests reveal mild leukopenia and relative lymphocytosis; persistent mixed cryoglobulinemia is present in most cases. ¹⁹ Cases of hepatic dysfunction have also been reported. ²⁰

Dermatological manifestations:

A rash characteristically appears on the first day of illness, but may be delayed.

- The patient exhibits erythema of the face and neck, which evolves to a macular or maculopapular exanthem of the trunk, limbs, face, palms, and soles in 50% of cases. ²¹
- Common findings also include hyperpigmentation, xerosis, excoriated papules, aphthous-like ulcers, vesiculobullous and lichenoid eruptions, and exacerbation of pre-existing or quiescent dermatoses. ^{22 23}
- Pigmentary changes are seen in 42% of cases, intertriginous aphthous-like ulcers in 21.37% and a vesiculobullous eruption in 2.75% (only in infants). ²⁴
- Morbilliform eruptions are most common, followed by scaling, macular erythema, intertrigo, hypermelanosis, xerosis, excoriated papules, urticaria and petechiae. ²⁵
- Vesiculo-bullous lesions are most common in children ²⁶ ; and extensive bullous lesions have been reported in infected infants. ²⁷
- Pruritus is common, and petechiae have been seen in some patients.
- Purpuric macules ²⁸ , genital ulcers, desquamation of the facial skin ²⁹ , erythema multiforme and erythema nodosum have also been reported in patients with Chikungunya. ³⁰

Complications:

Complications include hemorrhagic syndrome, myopericarditis ^{31 32} , hemodynamic disorders ³³ and rare instances of renal

failure. ³⁴

- Fatal infection ^{35 36} and transplacental infections have been reported. ³⁷⁻³⁹
- Peritonitis, encephalitis and secondary bacterial infections have been reported among immunocompromised patients with Chikungunya. ⁴⁰
- The case fatality rate may be as high as 1 per 1,000 cases. ^{41 42}
- Children occasionally present with seizures or convulsions.
- Sudden sensorineural hearing loss has been reported ⁴³
- Eye involvement may present as transient granulomatous and nongranulomatous anterior uveitis, optic neuritis ⁴⁴ , retinitis ^{45 46} , retrobulbar neuritis ⁴⁷ , Fuchs' heterochromic iridocyclitis. ⁴⁸ and dendritic lesions. ⁴⁹
- Chikungunya has no observable effect on the outcome of pregnancy ⁵⁰ ; however, infection of infants during the perinatal period is characterized by fever, rash, peripheral edema, thrombocytopenia, lymphopenia, decreased prothrombin value, and elevation of aspartate aminotransferase levels.
- Neurological complications include altered mental function, seizures ⁵¹ , encephalitis ^{52 53} , myelopathy ^{54 55} , myeloradiculopathy ⁵⁶ , acute flaccid paralysis ⁵⁷ , focal neurological deficit ⁵⁸ with abnormal CT scan of head, Guillain-Barre syndrome ⁵⁹ , urinary retention ⁶⁰ and altered CSF biochemistry. ⁶¹⁻⁶³

In some cases Chikungunya may mimic Kawasaki disease. ⁶⁴

- Although the clinical features of dengue and chikungunya are similar, chikungunya patients are more likely to exhibit early myalgia or arthralgia; while sore throat, cough, nausea, vomiting, diarrhea, abdominal pain, anorexia, tachycardia and thrombocytopenia will favor a diagnosis of dengue. ⁶⁵

Infection by a related agent, **Semliki Forest virus**, is characterized by fever, myalgia, arthralgia and persistent headache. ⁶⁶

This disease is endemic or potentially endemic to 48 countries.

Chikungunya in Uganda

Rare instances of human infection are documented. ^{67 68}

Seroprevalence surveys:

47% of adults in Karamoja district (1984) ⁶⁹

References

1. Clin Infect Dis 2007 Jun 1;44(11):1401-7.
2. Baillieres Clin Rheumatol 1995 Feb ;9(1):145-50.
3. J Rheumatol 1980 Mar-Apr;7(2):231-6.
4. J Rheumatol 1980 Mar-Apr;7(2):231-6.
5. BMC Infect Dis 2010 ;10:31.
6. Int J Clin Pract 2011 Dec ;65(12):1306-12.
7. BMC Infect Dis 2009 ;9:200.
8. Clin Infect Dis 2008 Aug 15;47(4):469-75.
9. Trans R Soc Trop Med Hyg 2010 Jun ;104(6):392-9.
10. J Assoc Physicians India 2011 Feb ;59:83-6.
11. Epidemiol Infect 2011 Jul 18;:1-9.
12. Best Pract Res Clin Rheumatol 2011 Jun ;25(3):337-46.
13. Medicine (Baltimore) 2012 Jul ;91(4):212-9.
14. ProMED <promedmail.org> archive: 20110115.0178
15. PLoS Negl Trop Dis 2009 ;3(3):e389.
16. Rheumatology (Oxford) 2012 Mar 16;
17. J Infect 2012 Apr 17;
18. Trans R Soc Trop Med Hyg 2010 Jun ;104(6):392-9.
19. PLoS Negl Trop Dis 2009 ;3(2):e374.
20. Med J Malaysia 2010 Mar ;65(1):83-4.
21. Med Trop (Mars) 2007 Apr ;67(2):167-73.
22. Indian J Dermatol 2010 ;55(1):64-7.
23. Indian J Dermatol Venereol Leprol 2010 Nov-Dec;76(6):671-6.
24. Int J Dermatol 2008 Feb ;47(2):154-9.
25. Int J Dermatol 2008 Nov ;47(11):1148-52.
26. Indian Pediatr 2012 Jan ;49(1):51-3.
27. Eur J Pediatr 2010 Jan ;169(1):67-72.
28. Int J Dermatol 2011 Jan ;50(1):61-9.
29. Indian J Dermatol 2011 May ;56(3):290-4.
30. Indian J Dermatol 2009 ;54(2):128-31.
31. J Indian Med Assoc 1978 Jun 1;70(11):256-8.
32. Am J Trop Med Hyg 2008 Feb ;78(2):212-3.
33. Pediatr Infect Dis J 2007 Sep ;26(9):811-5.
34. Trans R Soc Trop Med Hyg 2010 Feb ;104(2):89-96.
35. Emerg Infect Dis 2008 Aug ;14(8):1327.
36. Med J Malaysia 2010 Mar ;65(1):83-4.
37. ProMED <promedmail.org> archive: 20061006.2873
38. ProMED <promedmail.org> archive: 20070524.1669
39. ProMED <promedmail.org> archive: 20070718.2305
40. Emerg Infect Dis 2010 Jun ;16(6):1038-40.
41. Arch Pediatr 2008 Mar ;15(3):253-62.
42. ProMED <promedmail.org> archive: 20080304.0895
43. Int J Pediatr Otorhinolaryngol 2008 Feb ;72(2):257-9.
44. Int J Infect Dis 2011 Feb ;15(2):e147-50.
45. Indian J Ophthalmol 2008 Jul-Aug;56(4):329-31.
46. Indian J Ophthalmol 2009 Mar-Apr;57(2):148-50.
47. Indian J Ophthalmol 2009 Mar-Apr;57(2):148-50.
48. Indian J Ophthalmol 2010 Nov-Dec;58(6):545-7.
49. Am J Ophthalmol 2007 Oct ;144(4):552-6.
50. Emerg Infect Dis 2010 Mar ;16(3):418-25.
51. Rev Neurol (Paris) 2009 Jan ;165(1):48-51.
52. Scand J Infect Dis 2008 ;40(11-12):995-6.
53. J Assoc Physicians India 2012 Apr ;60:68-70.
54. J Clin Virol 2009 Oct ;46(2):145-9.
55. Am J Trop Med Hyg 2011 Aug ;85(2):386-9.
56. J Glob Infect Dis 2012 Oct ;4(4):207-8.
57. Epidemiol Infect 2008 Sep ;136(9):1277-80.
58. Neurol India 2009 Mar-Apr;57(2):177-80.
59. Emerg Infect Dis 2009 Mar ;15(3):495-6.
60. Urol Ann 2010 Sep ;2(3):110-3.
61. J Assoc Physicians India 2007 Nov ;55:765-9.
62. J Child Neurol 2008 Sep ;23(9):1028-35.
63. Crit Care Med 2008 Sep ;36(9):2536-41.
64. Pediatr Infect Dis J 2010 Mar ;29(3):275-7.
65. PLoS Negl Trop Dis 2012 Sep ;6(9):e1786.
66. Am J Trop Med Hyg 1990 Apr ;42(4):386-93.
67. Trans R Soc Trop Med Hyg 1985 ;79(4):567.
68. Trans R Soc Trop Med Hyg 1971 ;65(2):152-68.
69. Trans R Soc Trop Med Hyg 1989 Nov-Dec;83(6):851-4.

Chlamydia infections, misc.

Agent	BACTERIUM. Chlamydiaceae, Chlamydiae , Chlamydia trachomatis; Simkania negevensis; Waddlia chondrophila
Reservoir	Human
Vector	None
Vehicle	Sexual contact
Incubation Period	5d - 10d
Diagnostic Tests	Microscopy and immunomicroscopy of secretions. Serology. Tissue culture. Nucleic acid amplification.
Typical Adult Therapy	Doxycycline 100 mg BID X 7d. OR Azithromycin 1g as single dose OR Levofloxacin 500 mg daily X 7 days OR Ofloxacin 300 mg BID X 7 days
Typical Pediatric Therapy	Weight <45 kg: Erythromycin 10 mg/kg QID X 14d Weight ≥45 kg, but age <8 years: Azithromycin 1 g as single dose Age ≥ 8 years: Azithromycin 1 g as single dose OR Doxycycline 100 mg BID X 7 d
Clinical Hints	Thin, scant penile discharge; cervicitis; conjunctivitis; neonatal pneumonia; pelvic inflammatory disease; concurrent gonorrhea may be present.
Synonyms	Bedsonia, Chlamydia trachomatis, Chlamyden-Urethritis, Chlamyden-Zervizitis, Chlamydophila, Inclusion blenorrhea, Non-gonococcal urethritis, Nonspecific urethritis, Parachlamydia, Parachlamydia acanthamoebae, Prachlamydia, Protochlamydia, Protochlamydia naegleriophila, Rhabdochlamydia, Simkania negevensis, Waddlia chondrophila. ICD9: 099.41,099.5 ICD10: A56,A55

Clinical

Genito-urinary Infection with *Chlamydia trachomatis* may result in urethritis, epididymitis ¹, obstructive uropathy ², cervicitis, Fitz-Hugh-Curtis syndrome ³⁻⁵, acute salpingitis, tubal scarring, reduced conception rates (even in the absence of scarring) ⁶, ectopic pregnancy ⁷⁻¹⁰, miscarriage ^{11 12}, low birth weight or pre-term delivery. ¹³⁻¹⁶

- The rates of orchitis/epididymitis, prostatitis, infertility, and urethral stricture following genital infection in males are 4.28%, 1.41%, 1.27%, and 0.13% • respectively. ¹⁷
- The extent to which *Chlamydia* infection contributes to male and female infertility is unclear. ¹⁸⁻²²
- Levels of serum Prostate-specific Antigen (PSA) may be elevated in patients with *Chlamydia trachomatis* infection. ²³
- Perinatal infections may result in inclusion conjunctivitis or pneumonia in the newborn. ²⁴
- Asymptomatic pharyngeal infection or acute chlamydial tonsillopharyngitis may follow oro-genital contact. ²⁵

Chlamydia trachomatis infection is implicated in the etiology of reactive arthritis. ²⁶⁻⁴⁵

Parachlamydiaceae (including *Parachlamydia acanthamoebae*) have been associated with human respiratory infections, conjunctivitis, keratitis and uveitis. ^{46 47}

- The signs and symptoms of infection are similar to those of genital *Mycoplasma* infection. ⁴⁸
- Recurrent infection may represent either reinfection or treatment failure. ⁴⁹

For surveillance purposes, the CDC (The United States Centers for Disease Control) case definition of nongonococcal urethritis requires that gonorrhea has been discounted in the setting of:

- a visible abnormal urethral discharge
- or, a positive leukocyte esterase test from a male aged <60 who does not have a history of kidney disease or bladder infection, prostatic enlargement, anatomical abnormality of the urogenital tract, or recent urinary tract instrumentation
- or microscopic evidence of urethritis (over 5 leukocytes per high-power field) on stain of a urethral smear.

This disease is endemic or potentially endemic to all countries.

Chlamydia infections, misc. in Uganda

Prevalence surveys:

- 3.2% of women in Rakai (2003 publication) ⁵⁰
- 9.0% of female and 5.7% of male adolescents in Kampala (2006 publication) ⁵¹
- 6.8% of urban and 7.8% of rural women below age 20 (*Chlamydia trachomatis* and/or *Neisseria gonorrhoeae*, 2010 publication) ⁵²
- 7.1% of sexually experienced women below age 20 years (*Chlamydia trachomatis* and/or *Neisseria gonorrhoeae*, 2011 publication) ⁵³
- 7.1% of HIV-positive patients (2012 publication) ⁵⁴

References

1. Sex Transm Dis 2008 Sep ;35(9):827-33.
2. Hinyokika Kiyo 2008 Apr ;54(4):301-4.
3. Korean J Lab Med 2008 Aug ;28(4):293-8.
4. Korean J Gastroenterol 2010 Mar ;55(3):203-7.
5. Arch Pediatr 2013 Jan 29;
6. Hum Reprod 2011 Nov ;26(11):3061-7.
7. Sex Transm Dis 2007 Oct ;34(10):739-43.
8. J Obstet Gynaecol Res 2009 Aug ;35(4):775-81.
9. Fertil Steril 2012 Nov ;98(5):1175-85.
10. Ginekol Pol 2012 Nov ;83(11):819-21.
11. Emerg Infect Dis 2011 Sep ;17(9):1630-5.
12. Ugeskr Laeger 2013 Feb 4;175(6):354-357.
13. Eur J Epidemiol 2011 Jun ;26(6):493-502.
14. Clin Exp Obstet Gynecol 2004 ;31(3):175-8.
15. Braz J Infect Dis 2011 Nov-Dec;15(6):533-9.
16. Gynecol Obstet Invest 2012 Mar 29;
17. Sex Transm Dis 2008 Sep ;35(9):827-33.
18. Sex Transm Infect 2008 Jun ;84(3):171-5.
19. Am J Trop Med Hyg 2008 Feb ;78(2):323-7.
20. Fertil Steril 2009 Apr ;91(4 Suppl):1448-50.
21. Sex Transm Infect 2008 Jun ;84(3):171-5.
22. Hum Reprod Update 2010 Mar-Apr;16(2):189-204.
23. Br J Cancer 2011 Aug 23;105(5):602-5.
24. Semin Pediatr Infect Dis 2005 Oct ;16(4):235-44.
25. Case Report Otolaryngol 2012 ;2012:736107.
26. Curr Opin Rheumatol 2010 Jul ;22(4):424-30.
27. Curr Opin Rheumatol 2010 Jul ;22(4):363-7.
28. Int J Rheum Dis 2010 Feb 1;13(1):27-38.
29. Curr Opin Rheumatol 2010 Jan ;22(1):72-7.
30. Arthritis Rheum 2009 May ;60(5):1311-6.
31. Scand J Rheumatol 2009 ;38(5):353-6.
32. Curr Rheumatol Rep 2007 Apr ;9(1):4-5.
33. Scand J Rheumatol 2006 Nov-Dec;35(6):459-62.
34. Ann Rheum Dis 2006 Mar ;65(3):281-4.
35. Rheumatol Int 2006 Aug ;26(10):879-85.
36. Clin Dermatol 2004 Nov-Dec;22(6):469-75.
37. Curr Opin Rheumatol 2004 Jul ;16(4):380-92.
38. Rheum Dis Clin North Am 2003 Aug ;29(3):613-29.
39. Rheum Dis Clin North Am 2003 Feb ;29(1):21-36, v-vi.
40. Arthritis Res 2002 ;4(1):5-9.
41. Curr Rheumatol Rep 2001 Oct ;3(5):412-8.
42. Ann Rheum Dis 2001 Apr ;60(4):337-43.
43. Ann Med 2011 Aug 24;
44. Reumatol Clin 2012 Mar 13;
45. Am J Med Sci 2013 Jan 16;
46. Clin Microbiol Rev 2006 Apr ;19(2):283-97.
47. Medicine (Baltimore) 2008 May ;87(3):167-76.
48. Clin Infect Dis 2009 Jan 1;48(1):41-7.
49. J Infect Dis 2010 Jan 1;201(1):42-51.
50. Sex Transm Infect 2003 Apr ;79(2):98-105.
51. J Adolesc Health 2006 Mar ;38(3):213-21.
52. Sex Reprod Healthc 2010 Aug ;1(3):111-5.
53. Afr Health Sci 2011 Sep ;11(3):383-9.
54. J Infect Dev Ctries 2012 ;6(3):276-82.

Chlamydia pneumoniae infection

Agent	BACTERIUM. Chlamydiaceae, Chlamydiae , Chlamydia [Chlamydia] pneumoniae
Reservoir	Human
Vector	None
Vehicle	Droplet
Incubation Period	7d - 28d
Diagnostic Tests	Direct fluorescence of sputum. Serology and culture in specialized laboratories. Nucleic acid amplification.
Typical Adult Therapy	Respiratory isolation. Doxycycline 50 mg BID X 10d. Alternatives: Erythromycin 500 mg QID X 10d. Azithromycin 1 g, then 0.5 g daily. Clarithromycin 0.5 g BID
Typical Pediatric Therapy	Respiratory isolation; Erythromycin 10 mg/kg QID X 10d
Clinical Hints	Atypical pneumonia, often associated with pharyngitis and myalgia; consider when Mycoplasma, Legionella and influenza are discounted.
Synonyms	Chlamydia pneumoniae, Chlamydia TWAR, Chlamydia pneumoniae, TWAR. ICD9: 078.88 ICD10: J16.0

Clinical

Asymptomatic infection is common.

- Pneumonia and bronchitis are the most common clinical syndromes associated with *C. pneumoniae*.¹
- Sinusitis and pharyngitis may also occur, even in the absence of lower respiratory tract infection.
- Initial symptoms may consist of rhinitis, sore throat, or hoarseness; followed after several days or weeks prominent cough.
- Fever is often absent.
- Cough and malaise may persist for months; and reinfection may occur.

A single, subsegmental, patchy infiltrate may be seen on chest X ray.

- Other findings described include, lobar pulmonary consolidation, interstitial infiltrates, bilateral pneumonia, pleural effusion, acute respiratory distress syndrome², hilar adenopathy³ and myo-pericarditis.⁴
- The appearance of a miliary infiltrate may suggest a diagnosis of tuberculosis.⁵
- *Chlamydia pneumoniae* has been identified as an agent of otitis media.⁶
- Rare cases of intra-hepatic cholestasis have been reported.⁷
- The peripheral white blood cell count is usually not elevated.

C. pneumoniae has been identified as a cause of acute respiratory exacerbations in patients with cystic fibrosis and acute respiratory infection in children with sickle cell disease.

- *C. pneumoniae* infection is implicated in the etiology of recurrent tonsillitis.⁸
- The organism has also been implicated in development of asthma⁹⁻¹², chronic rhinosinusitis¹³, otitis media, migraine¹⁴, endocarditis, lumbosacral meningoradiculitis, erythema nodosum, erythema exsudativum multiforme¹⁵, nodular vasculitis¹⁶, Guillain-Barre syndrome, keratoconjunctivitis sicca¹⁷, hemophagocytic lymphohistiocytosis^{18 19}, reactive arthritis and atherosclerosis.²⁰

This disease is endemic or potentially endemic to all countries.

References

1. Expert Rev Anti Infect Ther 2003 Oct ;1(3):493-503.
2. J Med Case Rep 2012 ;6(1):20.
3. Pediatr Pulmonol 2011 Oct ;46(10):1038-40.
4. Pediatr Cardiol 2009 Apr ;30(3):336-9.
5. Pediatr Emerg Care 2009 Sep ;25(9):597-8.
6. Scand J Infect Dis 1998 ;30(4):377-80.
7. Med Mol Morphol 2011 Mar ;44(1):52-7.
8. Eur J Clin Microbiol Infect Dis 2008 Dec ;27(12):1233-7.
9. Curr Allergy Asthma Rep 2010 Jan ;10(1):67-73.
10. Immunol Allergy Clin North Am 2010 Nov ;30(4):575-85, vii-viii.
11. Immunol Allergy Clin North Am 2010 Nov ;30(4):565-74, vii.
12. Allergy 2011 Apr ;66(4):458-68.
13. Acta Otolaryngol 2006 Sep ;126(9):952-7.
14. J Headache Pain 2009 Apr ;10(2):121-4.
15. J Dermatol 2012 Mar ;39(3):306-8.
16. Case Rep Dermatol 2011 Sep ;3(3):263-7.
17. Cesk Slov Oftalmol 2011 Apr ;67(2):42, 44-8, 50.
18. PMID 21058284
19. Pediatr Blood Cancer 2011 May ;56(5):853-5.
20. Clin Infect Dis 2005 Apr 15;40(8):1131-2.

Cholecystitis & cholangitis

Agent	BACTERIUM. Escherichia coli , Klebsiella pneumoniae, enterococci, et al.
Reservoir	Human
Vector	None
Vehicle	Endogenous bacteria
Incubation Period	Variable
Diagnostic Tests	Roentgenograms/imaging (cholecystogram, ultrasound, CT, etc).
Typical Adult Therapy	Antibiotics and surgical intervention as required
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, chills and right upper quadrant abdominal pain; often "female, fat and 40"; may be associated with gallstones or pancreatitis, or present as 'fever of unknown origin'.
Synonyms	Acute cholecystitis, Angiocholite, Ascending cholangitis, Cholangitis, Cholecystite, Cholecystitis, Cholezystitis, Colangite, Colangitis, Colectite, Gall bladder. ICD9: 575.0,576.1 ICD10: K81,K83.0

Clinical

Cholangitis is caused by obstruction of the common bile duct, which subsequently becomes infected. ¹

- Strictures, stenosis, tumors, or endoscopic manipulation of the CBD cause bile stasis.
- The resultant infection ascends into the hepatic ducts, while increased biliary pressure spreads infection into the biliary canaliculi, hepatic veins and perihepatic lymphatics, leading to bacteremia.

Charcot's triad (fever, right upper quadrant pain, and jaundice) is found in 70% of patients.

- Additional findings include right upper quadrant pain, mild hepatomegaly, tachycardia, altered mental status, rigors, fever, hypotension, jaundice, pruritis, acholic stools.
- The case-fatality rate is 7% to 40%, and is highest in patients with hypotension, renal failure, liver abscess, cirrhosis, inflammatory bowel disease, malignant strictures and advanced age, or delays in diagnosis or surgery.

This disease is endemic or potentially endemic to all countries.

References

1. [Mayo Clin Proc 1998 May ;73\(5\):473-8.](#)

Cholera

Agent	BACTERIUM. <i>Vibrio cholerae</i> A facultative gram-negative bacillus
Reservoir	Human
Vector	None
Vehicle	Water Fecal-oral Seafood (oyster, ceviche) Vegetables Fly
Incubation Period	1d - 5d (range 9h - 6d)
Diagnostic Tests	Stool culture. Advise laboratory when this organism is suspected.
Typical Adult Therapy	Stool precautions. Doxycycline 100 mg BID X 5d, or Fluoroquinolone (Levofloxacin , Trovafoxacin , Pefloxacin , Sparfloxacin or Moxifloxacin), or Azithromycin Fluids (g/l): NaCl 3.5, NaHCO ₃ 2.5, KCl 1.5, glucose 20
Typical Pediatric Therapy	Stool precautions. Age >=8 years: Doxycycline 2 mg/kg BID X 5d. Age <8 years: Sulfamethoxazole/trimethoprim Fluids (g/l): NaCl 3.5, NaHCO ₃ 2.5, KCl 1.5, glucose 20
Vaccines	Cholera - injectable Cholera - oral
Clinical Hints	Massive, painless diarrhea and dehydration; occasionally vomiting; apathy or altered consciousness common; rapid progression to acidosis, electrolyte imbalance and shock; fever is uncommon.
Synonyms	Colera, Kolera. ICD9: 001 ICD10: A00

Clinical

WHO Case definition for surveillance:

The WHO Case definition for surveillance is as follows:

Clinical case definition

- In an area where the disease is not known to be present: severe dehydration or death from acute watery diarrhea in a patient aged 5 years or more or
- In an area where there is a cholera epidemic: acute watery diarrhea, with or without vomiting in a patient aged 5 years or more

Laboratory criteria for diagnosis

- Isolation of *Vibrio cholerae* O1 or O139 from stools in any patient with diarrhea.

Case classification

- Suspected: A case that meets the clinical case definition.
- Probable: Not applicable.
- Confirmed: A suspected case that is laboratory-confirmed.

Note: In a cholera-threatened area, when the number of confirmed cases rises, shift should be made to using primarily the suspected case classification.

- Cholera does appear in children under 5 years; however, the inclusion of all cases of acute watery diarrhea in the 2-4 year age group in the reporting of cholera greatly reduces the specificity of reporting.
- For management of cases of acute watery diarrhea in an area where there is a cholera epidemic, cholera should be suspected in all patients.

Symptoms and signs of cholera reflect the degree of fluid loss: thirst, postural hypotension, tachycardia, weakness, fatigue and dryness of the mucous membranes.

- Following an incubation period of 12 hours to 5 days¹, the patient experiences sudden onset of painless, watery diarrhea, which may later be accompanied by vomiting.²
- Abdominal cramps may occur.
- Fever is typically absent in adults, but present in children.
- The diarrhea has a "rice water" appearance and fishy odor.
- In patients with severe disease, stool volume can exceed 250 ml per /kg during the first 24 hours (17.5 liters in a 70 kg adult!).
- Severe cases exhibit sunken eyes (depressed fontanelles in infants), thready pulse, somnolence or coma.
- Without replacement of fluids and electrolytes, hypovolemic shock and death ensue.
- The clinical features of cholera due to *Vibrio cholerae* O139 are indistinguishable from disease due to other strains.³
- Rare cases of acalculous⁴⁻⁶ and infectious cholecystitis have been ascribed to *Vibrio cholerae*.⁷


This disease is endemic or potentially endemic to 119 countries.

Cholera in Uganda

Cholera is currently or recently reported from:

- Central Region
- Eastern Region
- Western Region

In recent years, the disease had been limited to Kasase District.




Graph: Uganda. Cholera, cases

Notes:

1. During December 1997, 370 cases (3 fatal) were confirmed in Kampala City and 26 cholera deaths were reported in Bugiri District (Busoga Province). ^{8 9}

Individual years:

- 1999 - Activity in Eastern Region and Western Region (Kasese).
- 2003 - Primarily in Bundibugyo district.
- 2007 - 100 cases (0 fatal) were reported during January.



Graph: Uganda. Cholera, deaths

Notable outbreaks:

1997 to 1998 - An outbreak (6,228 cases) of cholera was reported in Kampala. ¹⁰

2006 to 2007 - An outbreak (1,099 cases, 11 fatal) was reported in the Kampala region. ¹¹

2007 to 2008 - Outbreaks were reported in Arua, Bulisa, Butaleja, Kampala, Kibaale, Ksoro, Manafwa, Mbele, Mpigi, Nebbi, Ntungano, Pallisa and Yumbi. ¹², ¹³⁻³³ Cases were also reported among refugees fleeing Dem. Rep. Congo to Uganda. ³⁴ ³⁵

2009 - Outbreaks were reported in Buigiri (23 fatal cases) ³⁶, Kasese (584 cases, 10 fatal) ³⁷ and Kampala (35 cases, 5 fatal). ³⁸⁻⁴⁰

2010 - Outbreaks were reported in Manafwa (42 cases, 4 fatal) ⁴¹ ⁴², Butaleja (3 fatal cases) ⁴³, Tororo (50 cases, 6 fatal) and Moroto (220 cases, 5 fatal). ⁴⁴⁻⁴⁶

2011 - Outbreaks were reported in Kasese District (54 cases, 4 fatal) ⁴⁷ and Hoima (8 suspect cases). ⁴⁸

2012 - Outbreaks were reported. ⁴⁹⁻⁵⁶

2013 - An outbreak (28 cases) was reported in Nebbiu District. ⁵⁷

References

1. J Infect 2012 Nov 29;
2. Clin Infect Dis 2003 Aug 1;37(3):398-405.
3. Dtsch Tierarztl Wochenschr 1993 Jul ;100(7):255-8.
4. Diagn Microbiol Infect Dis 1998 Mar ;30(3):187-91.
5. Surg Endosc 1995 Jun ;9(6):730-2.
6. Lancet 1994 May 7;343(8906):1156-7.
7. Am J Gastroenterol 1996 Oct ;91(10):2241-2.
8. East Afr Med J 2000 Jul ;77(7):347-9.
9. Wkly Epidemiol Rec 1998 Jul 3;73(27):201-8.
10. East Afr Med J 2000 Jul ;77(7):347-9.
11. ProMED <promedmail.org> archive: 20080128.0359
12. ProMED <promedmail.org> archive: 20070105.0047
13. ProMED <promedmail.org> archive: 20070112.0147
14. ProMED <promedmail.org> archive: 20080108.0104
15. ProMED <promedmail.org> archive: 20071219.4087
16. ProMED <promedmail.org> archive: 20071213.4015
17. ProMED <promedmail.org> archive: 20071207.3944
18. ProMED <promedmail.org> archive: 20071128.3845
19. ProMED <promedmail.org> archive: 20070924.3164
20. ProMED <promedmail.org> archive: 20070921.3141
21. ProMED <promedmail.org> archive: 20070323.1019
22. ProMED <promedmail.org> archive: 20070205.0457
23. ProMED <promedmail.org> archive: 20070126.0350
24. ProMED <promedmail.org> archive: 20080201.0411
25. ProMED <promedmail.org> archive: 20080212.0563
26. ProMED <promedmail.org> archive: 20080223.0742
27. ProMED <promedmail.org> archive: 20080314.1020
28. ProMED <promedmail.org> archive: 20080602.1764
29. ProMED <promedmail.org> archive: 20080617.1896
30. ProMED <promedmail.org> archive: 20080627.1983
31. ProMED <promedmail.org> archive: 20081001.3099
32. ProMED <promedmail.org> archive: 20081014.3253
33. ProMED <promedmail.org> archive: 20081218.3988
34. ProMED <promedmail.org> archive: 20081118.3641
35. ProMED <promedmail.org> archive: 20081212.3906
36. ProMED <promedmail.org> archive: 20090930.3411
37. ProMED <promedmail.org> archive: 20091125.4044
38. ProMED <promedmail.org> archive: 20091009.3500
39. ProMED <promedmail.org> archive: 20091023.3653
40. ProMED <promedmail.org> archive: 20091103.3795
41. ProMED <promedmail.org> archive: 20100216.0550
42. ProMED <promedmail.org> archive: 20100319.0881
43. ProMED <promedmail.org> archive: 20100825.2993
44. Epidemiol Infect 2011 Sep 27;:1-10.
45. ProMED <promedmail.org> archive: 20100525.1737
46. ProMED <promedmail.org> archive: 20101025.3869
47. ProMED <promedmail.org> archive: 20111203.3523
48. ProMED <promedmail.org> archive: 20110617.1849
49. ProMED <promedmail.org> archive: 20120302.1057887
50. ProMED <promedmail.org> archive: 20120310.1065940
51. ProMED <promedmail.org> archive: 20120402.1087425
52. ProMED <promedmail.org> archive: 20120405.1091348
53. ProMED <promedmail.org> archive: 20120419.1106799
54. ProMED <promedmail.org> archive: 20120709.1191289

- 55. [ProMED <promedmail.org>](mailto:promedmail.org) archive: 20120724.1211703
- 56. [ProMED <promedmail.org>](mailto:promedmail.org) archive: 20121126.1424886
- 57. [ProMED <promedmail.org>](mailto:promedmail.org) archive: 20130131.1522070

Chromomycosis

Agent	FUNGUS. Ascomycota, Euascomycetes, Chaetothyriales. Dematiaceous molds: Phialophora, Cladophialophora, Fonsecaea, Rhinocladiella
Reservoir	Wood Soil Vegetation
Vector	None
Vehicle	Minor trauma
Incubation Period	14d - 90d
Diagnostic Tests	Biopsy and fungal culture.
Typical Adult Therapy	Itraconazole 100 mg PO QID X (up to) 18 m. OR (for late disease) Flucytosine 25 mg/kg QID X 4m. Terbinafine has been used in some cases. Local heat; excision as necessary
Typical Pediatric Therapy	Itraconazole 1 mg/kg PO BID X (up to) 18 m. OR Ketoconazole (if age >2) 5 mg/kg/d X 3 to 6m. Local heat; excision as necessary
Clinical Hints	Violaceous, verrucous, slowly-growing papule(s) or nodules, most commonly on lower extremities; usually follows direct contact with plant matter in tropical regions.
Synonyms	Chromoblastomycosis, Chromomykose, Verrucous dermatitis. ICD9: 117.2 ICD10: B43.0

Clinical

The lesions of chromomycosis typically progress from a papule to cicatricial fibrosis: nodules, tumors, plaques, warty lesions, and scarring lesions. ^{1 2}

- The verrucous form appears at the site of inoculation.
- The primary lesion, a small pink scaly papule, may be pruritic but rarely painful. ³
- Over time (often months to years), new crops of lesions appear in the same or adjacent areas as warty, purplish, scaly nodules or smooth, firm tumors. ⁴
- Peripheral spread may occur with healing in the center, as lesions enlarge and become grouped.
- Older lesions resemble cauliflower, with small ulcerations or "black dots" of hemopurulent material on the surface. ⁵
- These lesions can be pruritic and are rarely painful.
- Satellite lesions may develop through autoinoculation or lymphatic spread.
- Coalesced lesions form a large verrucous mass.
- Occasionally, an annular, flattened, papular lesion having a raised border is encountered.
- Keloid formation, fibrosis, lymphostasis and marked edema may follow.
- Fistulae are not seen.
- Malignant transformation has been reported in long-lasting lesions. ⁶

Signs of mucosal infection may mimic those of rhinosporidiosis ⁷, while those of cutaneous infection may mimic dermal leishmaniasis. ⁸

Rare cases of postoperative eye infection have been reported. ⁹

Rare cases of hematogenous spread to the brain, lymph nodes, liver, lungs, bones and joints ¹⁰, soft tissues and other organs have been reported. ¹¹

This disease is endemic or potentially endemic to all countries.

References

1. Curr Opin Infect Dis 2006 Apr ;19(2):148-52.
2. Clin Dermatol 2012 Jul ;30(4):403-8.
3. Clin Dermatol 2007 Mar-Apr;25(2):188-94.
4. Infect Dis Clin North Am 2003 Mar ;17(1):59-85, viii.
5. Med Mycol 2009 Feb ;47(1):3-15.
6. An Bras Dermatol 2010 Mar-Apr;85(2):267-70.
7. J Clin Pathol 1960 Jul ;13:287-90.
8. Dermatol Online J 2012 ;18(10):3.
9. J Cataract Refract Surg 2011 May ;37(5):963-6.
10. Am J Orthop (Belle Mead NJ) 2012 Jul ;41(7):328-31.
11. Skeletal Radiol 2009 Feb ;38(2):177-80.

Chronic meningococemia

Agent	BACTERIUM. Neisseria meningitidis An aerobic gram-negative coccus
Reservoir	Human
Vector	None
Vehicle	Air Infected secretions
Incubation Period	Unknown
Diagnostic Tests	Blood culture. Test patient for complement component deficiency.
Typical Adult Therapy	Intravenous Penicillin G 20 million units daily X 7 days
Typical Pediatric Therapy	Intravenous Penicillin G 200,000 units daily X 7 days
Clinical Hints	Recurrent episodes of low-grade fever, rash, arthralgia and arthritis - may persist for months; rash is distal, prominent near joints and may be maculopapular, petechial or pustular; may be associated with complement component deficiency.
Synonyms	Meningococemia, chronic. ICD9: 036.2 ICD10: A39.3

Clinical

Chronic meningococemia is characterized by persistent meningococcal bacteremia associated with low-grade fever, rash and arthritis.

- The rash is similar to that of gonococemia. ^{1 2}
- The illness may recur over a period of weeks to months.
- Patients (or their contacts) may ultimately present with acute bacterial meningitis or septicemia.

Non-bacteremic cases occur, and may be diagnosed through demonstration of meningococci in skin lesions. ³

This disease is endemic or potentially endemic to all countries.

References

1. Schweiz Med Wochenschr 1998 Dec 12;128(50):1988-93.
2. Pediatr Dermatol 1996 Nov-Dec;13(6):483-7.
3. Arch Dermatol 2008 Jun ;144(6):770-3.

Clostridial food poisoning

Agent	BACTERIUM. Clostridium perfringens An anaerobic gram-positive bacillus
Reservoir	Soil Human Pig Cattle Fish Poultry
Vector	None
Vehicle	Food
Incubation Period	8h - 14h (range 5h - 24h)
Diagnostic Tests	Laboratory diagnosis is usually not practical. Attempt culture of food for <i>C. perfringens</i> .
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Abdominal pain; watery diarrhea (usually no fever or vomiting) onset 8 to 14 hours after ingestion of meat, fish or gravy; no fecal leucocytes; usually resolves within 24 hours.
Synonyms	

Clinical

Seven to 15 hours after ingestion of toxin (range 6 to 24), the patient develops watery diarrhea (90%), abdominal cramps (80%); and occasionally nausea (25%), vomiting (9%) or fever (24%).¹

- Symptoms may persist for 8 to 72 hours (usually one day)
- Fatal cases are rare^{2 3}

This disease is endemic or potentially endemic to all countries.

References

1. [Int J Food Microbiol 2002 Apr 5;74\(3\):195-202.](#)
2. [Rev Physiol Biochem Pharmacol 2004 ;152:183-204.](#)
3. [Anesthesiol Clin North America 2004 Sep ;22\(3\):509-32, vii.](#)

Clostridial myonecrosis

Agent	BACTERIUM. Clostridium perfringens An anaerobic gram-positive bacillus
Reservoir	Soil Human
Vector	None
Vehicle	Soil Trauma
Incubation Period	6h - 3d
Diagnostic Tests	Gram stain of exudate. Wound and blood cultures. Presence of gas in tissue (not specific).
Typical Adult Therapy	Prompt, aggressive debridement. Penicillin G 3 million units IV Q3h + Clindamycin 900 mg IV Q8h. Hyperbaric oxygen
Typical Pediatric Therapy	Prompt, aggressive debridement. Penicillin G 50,000 units/kg IV Q3h + Clindamycin 10 mg/kg IV Q6h. Hyperbaric oxygen
Vaccine	Gas gangrene antitoxin
Clinical Hints	Gas gangrene is heralded by rapidly progressive tender and foul smelling infection of muscle associated with local gas (crepitus or seen on X-ray), hypotension, intravascular hemolysis and obtundation.
Synonyms	Anaerobic myonecrosis, Clostridial gangrene, Gas gangrene. ICD9: 040.0 ICD10: A48.0

Clinical

Gas gangrene is a fulminant infection with prominent findings at the infection site and severe systemic disease. ¹

The process may follow trauma (usually of an extremity), surgery (notably intestinal or biliary), septic abortion or delivery, vascular insufficiency or burns, underlying colorectal or pelvic cancer, or neutropenia complicating leukemia or cytotoxic therapy.

Following an incubation period of 1 to 4 days (range 6 hours to 3 weeks) the patient develops severe local pain, heaviness or pressure.

- The infection then progresses within minutes to hours, with localized edema, pallor and tenderness.
- Gas may be noted in the soft tissues by palpation, x-ray or scans, but crepitance is a late finding .
- The skin initially appears pale, and progresses to a magenta or bronze discoloration with hemorrhagic bullae and subcutaneous emphysema.
- A thin, brown, serosanguinous discharge may be present, associated with an offensive odor described as sweetish or "mousey."
- Gram's stain of the discharge shows a large number of gram-positive or gram-variable rods, with few or no white blood cells.

Profound systemic toxicity is also present, diaphoresis, anxiety, and tachycardia disproportionate to fever.

- In fact, fever may be low or absent in the early stages.
- Other complications include intravascular hemolysis, hemoglobinuria, hypotension, renal failure, and metabolic acidosis.
- Central nervous system manifestations are rare and most frequently comprise meningitis with or without pneumocephalon, encephalitis, plexitis, cerebral abscess, or subdural empyema. ²
- Coma and generalized 'bronze' edema are seen preterminally.

This disease is endemic or potentially endemic to all countries.

References

1. [Int Orthop 2004 Oct ;28\(5\):257-60.](#)
2. [Infection 2007 Dec ;35\(6\):396-405.](#)

Clostridium difficile colitis

Agent	BACTERIUM. Clostridium difficile An anaerobic gram-positive bacillus
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Assay of stool for <i>C. difficile</i> toxin.
Typical Adult Therapy	Metronidazole 500 mg PO TID X 10d. OR Vancomycin 125 mg [oral preparation] QID X 10d
Typical Pediatric Therapy	Vancomycin 2 mg/kg [oral preparation] QID X 10d
Clinical Hints	Fever, leukocytosis, abdominal pain; mucoid or bloody diarrhea during / following antibiotic therapy; fecal leucocytes present; suspect even when mild diarrhea follows antibiotic intake.
Synonyms	<i>Klebsiella oxytoca colitis</i> , Pseudomembranous colitis. ICD9: 008.45 ICD10: A04.7

Clinical

Symptoms may appear as early as the first or second day of antimicrobial therapy; or as late as 10 weeks after cessation. ¹

- Occasionally, a single dose of an antimicrobial or antineoplastic agent has been implicated. ²

The frequency of diarrhea ranges from three to as many as 20 stools per day.

- Stools may be soft or watery, but rarely demonstrate overt blood.
- Occult blood in the stool is found in approximately 25% of patients. ³
- Abdominal pain is present in 22% of patients, fever in 28% and leukocytosis in 50%.
- Reactive polyarthritis and hemolytic-uremic syndrome ⁴ have been reported in some cases.
- Rare instances of *Clostridium difficile* bacteremia are reported. ^{5 6}
- Disease caused by *C. difficile* 027 is relatively severe and carries a higher mortality rate than infection by other strains. ^{7 8}

This disease is endemic or potentially endemic to all countries.

References

- Can Fam Physician 2004 Nov ;50:1536-40, 1543-5.
- BMJ 2005 Sep 3;331(7515):498-501.
- Clin Microbiol Infect 2005 Jul ;11 Suppl 4:57-64.
- Clin Nephrol 2013 Jan 15;
- Emerg Infect Dis 2010 Aug ;16(8):1204-10.
- J Med Microbiol 2011 Mar ;60(Pt 3):378-80.
- Postgrad Med J 2007 May ;83(979):291-5.
- Curr Opin Infect Dis 2007 Aug ;20(4):376-83.

Coenurosis

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Taeniidae: Taenia multiceps (Multiceps spp.)
Reservoir	Sheep Wild carnivore, Horse Dog
Vector	None
Vehicle	Water Food Soil (contaminated by dog)
Incubation Period	Unknown
Diagnostic Tests	Identification of parasite in biopsy material.
Typical Adult Therapy	Excision
Typical Pediatric Therapy	As for adult
Clinical Hints	Mass in brain, eye, muscle or subcutaneous tissue; may present months to years after exposure in sheep-raising areas; basilar arachnoiditis with internal hydrocephalus is common.
Synonyms	Multiceps, Taenia multiceps. ICD9: 123.8 ICD10: B71.8

Clinical

Human infection has a predilection for the cysterna magna, and presents as basal arachnoiditis and hydrocephalus. ¹

- Subcutaneous tissue, muscle and eye infections are also reported, and present as a cystic masses (often containing daughter cysts) which may attain the size of a hen's egg.
- The clinical features of coenurosis may mimic those of echinococcosis. ²

This disease is endemic or potentially endemic to 25 countries.

Coenurosis in Uganda

Most cases are reported from the western mountain regions or near the Kenyan border. ³

References

1. Clin Infect Dis 1998 Sep ;27(3):519-23.
2. Clin Neuropathol 2011 Jan-Feb;30(1):28-32.
3. Trans R Soc Trop Med Hyg 1968 ;62(2):251-5.

Common cold

Agent	VIRUS - RNA. Picornaviridae. Rhinoviruses, Coronavirus, et al.
Reservoir	Human
Vector	None
Vehicle	Droplet Contact
Incubation Period	1d - 3d
Diagnostic Tests	Viral culture and serology are available, but not practical.
Typical Adult Therapy	Supportive; Pleconaril under investigation
Typical Pediatric Therapy	As for adult
Clinical Hints	Nasal obstruction or discharge, cough and sore throat are common; fever >38 C unusual in adults; illness usually lasts one week, occasionally two.
Synonyms	Acute coryza, Raffreddore. ICD9: 079,460 ICD10: J00

Clinical

In young adults, the common cold runs its course in an average of 7 days.

Fever is uncommon, and in most cases, rhinorrhea and nasal obstruction predominate. ¹

- Sore throat, cough and hoarseness are often present.
- The nasal tip is often red, and mucoid secretions and a glistening nasal mucosa are evident.
- The pharynx may be mildly edematous and erythematous, but without exudate.

Complications include bacterial sinusitis, otitis media, exacerbation of chronic bronchitis and precipitation of asthma. ²

- Rare instances of pneumonia have been attributed to infection by Coronavirus strains OC43 and 229E.
- Severe symptoms, including bronchiolitis are associated with Coronavirus HCoV-NL63 infection in young children.

This disease is endemic or potentially endemic to all countries.

References

1. *Pediatr Infect Dis J* 2004 Nov ;23(11):1049-50.
2. *Allergy* 2011 Apr ;66(4):458-68.

Conjunctivitis - inclusion

Agent	BACTERIUM. Chlamydiae , Chlamydia trachomatis
Reservoir	Human
Vector	None
Vehicle	Infected secretions Sexual contact Water (swimming pools)
Incubation Period	5d - 12d
Diagnostic Tests	Demonstration of chlamydiae on direct fluorescence or culture of exudate.
Typical Adult Therapy	Secretion precautions. Topical Erythromycin . Erythromycin 250 mg PO QID. X 14 days OR Doxycycline 100 mg PO BID X 14 days
Typical Pediatric Therapy	Secretion precautions. Topical Erythromycin . Azithromycin 1 g PO as single dose. Alternative If age >8 years, Doxycycline 100 mg PO BID X 7 days.
Clinical Hints	Ocular foreign body sensation, photophobia and discharge which may persist for months to as long as 2 years; keratitis and conjunctival follicles may be evident.
Synonyms	Inclusion conjunctivitis, Paratrachoma. ICD9: 077.0 ICD10: P39.1,A74.0

Clinical

Ophthalmia neonatorum caused by *Chlamydia* is characterized by conjunctival injection without follicles. ¹

Follicular conjunctivitis in adults is most prominent on the lower lid, and the presence of bulbar follicles is highly suggestive of a *Chlamydia* etiology. ²

- The infection is usually bilateral and accompanied by profuse discharge.

Parachlamydiaceae (including *Parachlamydia acanthamoebae*) have been associated with conjunctivitis, keratitis and uveitis. ³

Trachoma may be differentiated from inclusion conjunctivitis by the presence of corneal scarring and a preference of the latter for the upper tarsal conjunctivae.

This disease is endemic or potentially endemic to all countries.

References

1. Arch Pediatr 1999 Mar ;6(3):317-20.
2. J Fr Ophtalmol 1999 May ;22(5):577-80.
3. Clin Microbiol Rev 2006 Apr ;19(2):283-97.

Conjunctivitis - viral

Agent	VIRUS. Picornavirus, Adenovirus
Reservoir	Human
Vector	None
Vehicle	Contact
Incubation Period	1d - 3d
Diagnostic Tests	Viral isolation is available but rarely practical.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Watery discharge, generalized conjunctival injection and mild pruritus; may be associated with an upper respiratory infection.
Synonyms	Apollo conjunctivitis, Apollo eye, Congiuntivite virale, Hemorrhagic conjunctivitis, Viral conjunctivitis. ICD9: 077.1,077.2,077.3,077.4,077.8,372.0 ICD10: B30,B30.3,H10

Clinical

The symptoms of viral conjunctivitis include erythema, itching and lacrimation.

- The presence of large quantities of pus may suggest a bacterial etiology. ^{1 2}

Hemorrhagic conjunctivitis is characterized by sudden onset of painful, swollen, red eyes with subconjunctival hemorrhaging, palpebral follicles, photophobia, foreign body sensation, eyelid edema, punctate keratitis, and excessive tearing. ^{3 4}

- Symptoms usually persist for 3 to 5 days.

This disease is endemic or potentially endemic to all countries.

Conjunctivitis - viral in Uganda

Notable outbreaks:

2010 - An outbreak (397 cases) of viral conjunctivitis was reported in Kampala. ⁵

2010 - An outbreak (6,818 cases) of acute hemorrhagic conjunctivitis was reported. ^{6 7}

References

1. BMJ 2003 Oct 4;327(7418):789.
2. Postgrad Med 1997 May ;101(5):185-6, 189-92, 195-6.
3. Prog Med Virol 1984 ;29:23-44.
4. ProMED <promedmail.org> archive: 20071006.3302
5. ProMED <promedmail.org> archive: 20100514.1574
6. MMWR Morb Mortal Wkly Rep 2010 Aug 20;59(32):1024.
7. ProMED <promedmail.org> archive: 20100819.2890

Crimean-Congo hemorrhagic fever

Agent	VIRUS - RNA. Bunyaviridae, Nairovirus: CCHF virus. Infections also ascribed to related agents (Nairobi sheep and Dugbe viruses)
Reservoir	Hare Bird Tick Cattle Sheep Goat
Vector	Tick (Hyalomma - over 30 potential vectors in this genus)
Vehicle	Infected secretions from patient or livestock
Incubation Period	1d - 6d (range 2d - 53d)
Diagnostic Tests	Viral culture (blood, CSF, tissue. Serology. Nucleic acid amplification. Biosafety level 4.
Typical Adult Therapy	Isolation. Supportive therapy. Ribavirin : 1g PO QID X 4d, then 0.5g QID X 6d
Typical Pediatric Therapy	Isolation. Supportive therapy Ribavirin (dosage not established)
Clinical Hints	Headache, chills, myalgia, abdominal pain, photophobia, petechiae, thrombocytopenia and leukopenia; conjunctivitis and pharyngitis are often present; onset 3 to 7 days following a tick bite. Case-fatality rate approximately 30%.
Synonyms	Acute infectious capillary toxinosis, CCHF, Crimea Congo hemorrhagic fever, Dugbe, Erve, Ganjam, Nairobi sheep, Tribec, Xinjiang hemorrhagic fever. ICD9: 065.0 ICD10: A98.0

Clinical

The incubation period following tick bite is usually one to three days; however, incubation periods as long as 53 days have been reported. ^{1 2}

- The incubation period following contact with infected blood or tissues is usually five to six days, with a maximum of 13 days.

Onset of illness is sudden, with fever, myalgia, vertigo, neck pain and stiffness, backache, headache and photophobia. ³

- There may be initial nausea, vomiting and sore throat accompanied by diarrhea and generalized abdominal pain. ⁴
- Later, the patient may experience sharp mood swings, and may become confused and aggressive.
- After two to four days, agitation is replaced by somnolence, depression and lassitude, and the abdominal pain may localize to the right upper quadrant, with detectable hepatomegaly. ⁵
- Other clinical signs at this stage include tachycardia, lymphadenopathy and a petechial rash which progresses to ecchymoses and other bleeding diatheses.
- There is usually evidence of hepatitis.
- Increased serum ferritin levels may suggest severe infection ⁶
- Acalculous cholecystitis ⁷, myocardial hypokinesia ⁸, hemorrhagic pleural effusion ⁹, parotitis ¹⁰ and diffuse alveolar hemorrhage with / without overt hemoptysis have been reported. ^{11 12}
- Sub-conjunctival and retinal hemorrhages are present in some cases. ¹³

The severely ill may develop hepatorenal and pulmonary failure after the fifth day of illness. ^{14 15}

- Severe endo-myocardial dysfunction has been reported. ^{16 17}
- The mortality rate may vary from 10% to 30%, with death occurring in the second week of illness. ^{18 19}
- In those patients who recover, improvement generally begins on the ninth or tenth day after onset of illness.

Sub-clinical infection is common. ²⁰

- Crimean-Congo hemorrhagic fever may be mis-diagnosed as dengue in regions where the two diseases co-exist. ²¹
- Q-fever may mimic Crimean-Congo hemorrhagic fever. ²²

Diagnosis of suspected CCHF is performed in specially-equipped, high biosafety level laboratories.

- IgG and IgM antibodies may be detected in serum by enzyme-linked immunoassay from day six of illness.
- IgM remains detectable for up to four months, and IgG levels decline but remain detectable for up to five years.

Although an inactivated, mouse brain-derived vaccine against CCHF has been developed and used on a small scale in Eastern Europe, there is no safe and effective vaccine widely available for human use.

This disease is endemic or potentially endemic to 54 countries.

Crimean-Congo hemorrhagic fever in Uganda

Seropositive humans have been identified in Karamoja district (1984). ²³

References

1. Int J Infect Dis 2011 Jul ;15(7):e449-52.
2. Mikrobiyol Bul 2012 Jan ;46(1):129-33.
3. Rev Infect Dis 1989 May-Jun;11 Suppl 4:S777-82.
4. ProMED <promedmail.org> archive: 20080622.1935
5. Rev Infect Dis 1989 May-Jun;11 Suppl 4:S777-82.
6. Int J Infect Dis 2010 Jan ;14(1):e50-4.
7. J Clin Virol 2011 Feb ;50(2):162-3.
8. Med Sci Monit 2011 Aug ;17(8):CR457-460.
9. Jpn J Infect Dis 2009 Jan ;62(1):70-2.
10. J Clin Virol 2012 Feb ;53(2):159-61.
11. Trop Doct 2008 Oct ;38(4):252-4.
12. Southeast Asian J Trop Med Public Health 2011 Sep ;42(5):1100-5.
13. Am J Ophthalmol 2009 Apr ;147(4):634-638.e1.
14. Int J Infect Dis 2008 Jul ;12(4):374-9.
15. Lancet Infect Dis 2006 Apr ;6(4):203-14.
16. Arch Biochem Biophys 1991 Oct ;290(1):143-52.
17. J Vector Borne Dis 2011 Sep ;48(3):150-4.
18. Curr Opin Infect Dis 2007 Oct ;20(5):495-500.
19. ProMED <promedmail.org> archive: 20110530.1653
20. Emerg Infect Dis 2012 Apr ;18(4):640-2.
21. J Infect Dev Ctries 2010 Jul ;4(7):459-63.
22. Indian J Med Microbiol 2011 Oct-Dec;29(4):418-9.
23. Trans R Soc Trop Med Hyg 1989 Nov-Dec;83(6):851-4.

Cryptococcosis

Agent	FUNGUS - Yeast. Basidiomycota, Hymenomycetes, Sporidiales: Cryptococcus neoformans
Reservoir	Pigeon Soil
Vector	None
Vehicle	Air
Incubation Period	Variable
Diagnostic Tests	Fungal culture and stains. Latex test for fungal antigen in CSF and serum. Nucleic acid amplification.
Typical Adult Therapy	Amphotericin B 0.3 mg/kg/d X 6w (+/- Flucytosine); then 0.8 mg/kg qod X 8w. OR Fluconazole 200 mg/d
Typical Pediatric Therapy	Amphotericin B 0.3 mg/kg/d X 6w (+/- Flucytosine); then 0.8 mg/kg qod X 8w. OR Fluconazole 3 mg/kg/d
Clinical Hints	Chronic lymphocytic meningitis or pneumonia in an immune-suppressed patient; meningitis may be subclinical, or "wax and wane" - nuchal rigidity absent or minimal; bone, skin, adrenals, liver, prostate and other sites may be infected.
Synonyms	Busse-Buschke disease, Cryptococcus, European blastomycosis, Torulosis. ICD9: 117.5,321.0 ICD10: B45

Clinical

Central nervous system infection:

Central nervous system infection may be acute or gradual in onset, with acute manifestations most common in immunosuppressed patients (eg, with AIDS). ¹

- Often, the onset is characterized by waxing and waning manifestations over weeks to months, interspersed by asymptomatic periods.
- Complaints may be mild and nonspecific, and consist of headache, nausea, dizziness, irritability, somnolence, confusion, or obtundation. ²
- Decreased visual acuity, diplopia, and facial weakness may be evident.
- Fever is often absent, and patients have minimal or no nuchal rigidity.
- Papilledema is noted as many as one third of cases, and cranial nerve palsies in 20%. Bilateral amaurosis has been reported as a sequela of infection. ^{3 4}
- Hyperreflexia, choreoathetoid movements or myoclonic jerks may be present.
- Elevated CSF protein concentrations are present in 50%, hypoglycorrhachia in 33% and pleocytosis above 20 cells per cu. Mm. In 20%.
- Peripheral blood eosinophilia may be present. ^{5 6}

Respiratory tract infection:

Respiratory tract cryptococcosis may be asymptomatic, or limited to a mild productive cough with blood-streaked sputum and minor ache in the chest. ^{7 8}

- Pulmonary infection may present as a single rounded lesion, lobar pneumonia, bronchiolitis obliterans ⁹ or miliary disease.
- Rales or pleural friction rub are unusual, and pleural effusions are uncommon.
- Pulmonary infection in immunocompetent patients may progress or regress spontaneously over long periods.
- Cryptococcosis among patients with AIDS often presents as a solitary cavitory pulmonary nodule. ¹⁰
- Concurrent CNS infection may be evident in some cases.

One-half of AIDS patients with cryptococcal meningitis have concurrent pulmonary involvement, and two-thirds are fungemic. ¹¹

- Initial cough and dyspnea are found in 5 to 25% of HIV-positive patients with cryptococcosis.
- Cryptococcal immune reconstitution inflammatory syndrome may present as a clinical worsening of cryptococcal disease after initiation of antiretroviral therapy. ¹²
- Case-fatality rates for treated cryptococcosis in AIDS patients are 10% to 25%.

The clinical features of *Cryptococcus neoformans* var. *gattii* infection are similar to those of *C. neoformans* infection. ¹³

- *C. neoformans* var. *gattii* infections usually involve the lungs (75 percent), although neurological (8 percent) and combined (9 percent) infections are seen. ¹⁴

- Blindness due to high cerebrospinal fluid pressure, optic neuropathy or endophthalmitis, is relatively common among immunocompetent individuals infected with *C. gattii*.¹⁵

Cryptococcosis may involve a variety of other sites including skin¹⁶⁻²⁶ and soft tissues²⁷⁻³⁰, blood stream³¹⁻³², mucosa, colon or intestine³³⁻³⁵, gall bladder and bile ducts³⁶, liver, peritoneum³⁷⁻³⁹, lymph nodes⁴⁰⁻⁴², bones and joints⁴³⁻⁴⁹, breasts, pericardium, genital tract⁵⁰⁻⁵², prostate⁵³, placenta (without neonatal involvement)⁵⁴, adrenals⁵⁵, eyes⁵⁶⁻⁵⁷, parotid glands⁵⁸, tongue⁵⁹⁻⁶⁰, retropharyngeal space⁶¹, etc.

The cutaneous features of cryptococcosis include papules, pustules, nodules, subcutaneous swelling, abscesses, molluscum contagiosum-like or tumor-like lesions, cellulitis, blisters, ulcers and very rarely, necrotizing fasciitis⁶²

- Primary cutaneous cryptococcosis may occur in persons working with birds.⁶³

Note: *Cryptococcus neoformans* is one of at least a dozen *Cryptococcus* species. See the Microbiology • Yeasts module.

This disease is endemic or potentially endemic to all countries.

Cryptococcosis in Uganda

Cryptococcosis occurred in 77 of 1,372 HIV-positive patients followed in a cohort study in Entebbe during 1995 to 1999⁶⁴

The rate of cryptococcosis among HIV-positive adults is 40.4 per 1,000 person-years⁶⁵

Prevalence surveys:

17% of all HIV-associated deaths.

2.7% of hospitalized HIV-positive patients and 11% of BAL specimens from HIV-positive patients (pulmonary cryptococcosis, Kampala, 2007 to 2008)⁶⁶

References

1. CNS Drugs 2003 ;17(12):869-87.
2. Infect Dis Clin North Am 2002 Dec ;16(4):837-74, v-vi.
3. Arq Bras Oftalmol 2008 Jan-Feb;71(1):101-3.
4. Rev Iberoam Micol 2010 Oct-Dec;27(4):207-9.
5. J Infect Chemother 2008 Aug ;14(4):319-24.
6. Kansenshogaku Zasshi 2010 Sep ;84(5):597-601.
7. Semin Respir Crit Care Med 2008 Apr ;29(2):141-50.
8. Curr Opin Pulm Med 2009 May ;15(3):254-60.
9. J Infect Chemother 2010 Jun ;16(3):206-9.
10. Scand J Infect Dis 2012 Dec 17;
11. AIDS 2007 Oct 18;21(16):2119-29.
12. Lancet Infect Dis 2010 Nov ;10(11):791-802.
13. Can J Infect Dis Med Microbiol 2009 ;20(1):23-8.
14. ProMED <promedmail.org> archive: 20100426.1341
15. ProMED <promedmail.org> archive: 20110427.1305
16. Indian J Med Microbiol 2006 Jul ;24(3):228-30.
17. Postepy Hig Med Dosw (Online) 2008 ;62:1-3.
18. J Drugs Dermatol 2008 Jan ;7(1):53-4.
19. J Cutan Pathol 2008 Nov ;35(11):1007-13.
20. Dermatol Online J 2008 ;14(7):9.
21. Transpl Infect Dis 2009 Feb ;11(1):68-71.
22. Trop Doct 2009 Apr ;39(2):114-5.
23. Mycoses 2010 May ;53(3):256-8.
24. BMC Infect Dis 2010 ;10:239.
25. Int J Dermatol 2012 Jul ;51(7):780-4.
26. J Infect Dev Ctries 2013 ;7(1):60-3.
27. Med Mycol 2008 May ;46(3):269-73.
28. Cutis 2010 Jun ;85(6):303-6.
29. Acta Cytol 2010 Nov-Dec;54(6):1130-2.
30. Sex Transm Dis 2012 Oct ;39(10):792-3.
31. Clin Nephrol 2009 Jan ;71(1):88-91.
32. Arerugi 2009 Nov ;58(11):1536-43.
33. Singapore Med J 2008 Nov ;49(11):e305-7.
34. Korean J Gastroenterol 2008 Oct ;52(4):255-60.
35. Med Mycol 2011 Oct ;49(7):775-8.
36. Int J Med Sci 2012 ;9(3):200-6.
37. PMID 21046602
38. Diagn Cytopathol 2011 May ;39(5):365-7.
39. Mycopathologia 2012 Jul ;174(1):81-5.
40. Br J Radiol 2008 Feb ;81(962):e53-6.
41. AIDS Res Hum Retroviruses 2011 Apr ;27(4):373-6.
42. Rev Med Chil 2010 Dec ;138(12):1535-8.
43. Med Mycol 2011 Aug ;49(6):667-71.
44. Hip Int 2011 Jul-Aug;21(4):495-7.
45. J Infect Dev Ctries 2011 Sep ;5(9):669-73.
46. J Indian Med Assoc 2011 Aug ;109(8):592, 594.
47. Med Mycol 2012 Mar 21;
48. Hand (N Y) 2011 Dec ;6(4):450-3.
49. Mycopathologia 2012 Dec 29;
50. Prostate Cancer Prostatic Dis 2008 ;11(2):203-6.
51. Int J Gynecol Pathol 2008 Jan ;27(1):37-40.
52. AIDS Patient Care STDS 2009 Feb ;23(2):71-3.
53. J Assoc Physicians India 2012 May ;60:57-9.
54. Pediatr Dev Pathol 2009 May-Jun;12(3):249-52.
55. BMC Infect Dis 2011 ;11:340.
56. Arq Bras Oftalmol 2006 Mar-Apr;69(2):265-7.
57. Ocul Immunol Inflamm 2008 Jul-Aug;16(4):191-3.
58. Med Mycol 2006 May ;44(3):279-83.
59. Southeast Asian J Trop Med Public Health 2010 Sep ;41(5):1188-91.
60. Case Report Otolaryngol 2012 ;2012:517415.
61. Travel Med Infect Dis 2010 Sep ;8(5):322-5.
62. Clin Exp Dermatol 2009 Dec ;34(8):e751-3.
63. Ugeskr Laeger 2011 Sep 26;173(39):2423-4.
64. AIDS 2002 May 3;16(7):1031-8.
65. Clin Infect Dis 2008 Jun 1;46(11):1694-701.
66. J Acquir Immune Defic Syndr 2010 Jul ;54(3):269-74.

Cryptosporidiosis

Agent	PARASITE - Protozoa. Sporozoa, Coccidea, Eimeriida: <i>Cryptosporidium hominis</i> and <i>C. parvum</i> (rarely <i>C. muris</i> , <i>felis</i> , <i>meleagridis</i> , et al).
Reservoir	Mammal (over 150 species)
Vector	None
Vehicle	Water Feces Oysters Fly
Incubation Period	5d - 10d (range 2d - 14d)
Diagnostic Tests	Stool/duodenal aspirate for acid-fast, direct fluorescence staining, or antigen assay. Nucleic acid amplification
Typical Adult Therapy	Stool precautions. Nitazoxanide 500 mg PO BID X 3 days
Typical Pediatric Therapy	Stool precautions. Nitazoxanide : 1 to 3 years: 100 mg PO BID X 3 days 4 to 11 years: 200 mg PO BID X 3 days >12 years: 500 mg PO BID X 3 days
Clinical Hints	Watery diarrhea, vomiting, abdominal pain; although self-limited in healthy subjects, this is a chronic and wasting illness and may be associated with pulmonary disease among immunosuppressed (e.g., AIDS) patients.
Synonyms	<i>Cryptosporidium</i> , <i>Cryptosporidium andersoni</i> , <i>Cryptosporidium fayeri</i> , <i>Cryptosporidium felis</i> , <i>Cryptosporidium hominis</i> , <i>Cryptosporidium parvum</i> , <i>Cryptosporidium tyzzeri</i> , <i>Cryptosporidium ubiquitum</i> , <i>Cryptosporidium viatorum</i> , Kryptosporidiose. ICD9: 007.4 ICD10: A07.2

Clinical

Cryptosporidiosis affects the gastrointestinal tract and may be asymptomatic or associated with watery diarrhea and abdominal cramps.

- Fever and anorexia are uncommon, and fecal leukocytes are not seen.
- Although vomiting is not common among adults, it is often encountered in children. ¹

Rare instances of pulmonary infection have been reported. ^{2 3}

There is some evidence that *Cryptosporidium hominis* infection in children is associated with diarrhea, nausea, vomiting, general malaise, and increased oocyst shedding intensity and duration.

- In contrast, infections caused by *C. parvum*, *C. meleagridis*, *C. canis*, and *C. felis* are associated with diarrhea only.

Illness persists for 1 to 20 days (mean 10) in immunocompetent individuals

- Protracted, severe diarrhea leading to malabsorption, dehydration, extraintestinal (ie, biliary or pulmonary ⁴⁻⁶) and fatal infection may develop in immunocompromised individuals. ^{7 8}

This disease is endemic or potentially endemic to all countries.

Cryptosporidiosis in Uganda

Prevalence surveys:

- 25.0% of children (ages 0 to 5) hospitalized for diarrhea in Kampala (1999 to 2000) ⁹
- 12.5% of children ages 9 to 36 months with diarrhea (Kampala, 2007 to 2009) ¹⁰
- 28.1% of malnourished children hospitalized with diarrhea (Kampala, 2002 to 2003)
- 73.6% of HIV-positive and 5.9% of HIV-negative children with persistent diarrhea (Kampala, 2002 to 2003) ¹¹
- 5% of HIV-positive patients with diarrhea in rural Uganda (2010 publication)
- 32.4% of humans, 11.1% of non-human primates and 2.2% of livestock (Kibale National Park, 2012 publication) ¹²
- 20% red colobus (*Ptilocolobus tephrosceles*), 5% of red-tailed guenons (*Cercopithecus ascanius*) and 0% of black-and-white colobus (*Colobus guereza*) from forest adjacent to a national park (*Cryptosporidium* sp. or *Giardia* sp.) (Kibale National Park, 2005) ¹³

Infection has been linked to contact with free mountain gorillas (*Gorilla gorilla*). ¹⁴

References

1. Semin Pediatr Infect Dis 2004 Oct ;15(4):253-9.
2. Indian J Pathol Microbiol 2009 Apr-Jun;52(2):267-8.
3. Ann Biol Clin (Paris) 2011 Sep-Oct;69(5):605-8.
4. Rev Argent Microbiol 2008 Apr-Jun;40(2):106-8.
5. Emerg Infect Dis 2007 Mar ;13(3):462-4.
6. N Engl J Med 1996 Jan 4;334(1):19-23.
7. Curr Opin Infect Dis 2002 Oct ;15(5):523-7.
8. Parasitol Today 1998 Apr ;14(4):150-6.
9. Am J Trop Med Hyg 2005 Nov ;73(5):921-5.
10. Clin Infect Dis 2010 May 15;50(10):1366-72.
11. Am J Trop Med Hyg 2005 Nov ;73(5):921-5.
12. PLoS Negl Trop Dis 2012 Apr ;6(4):e1597.
13. J Parasitol 2007 Apr ;93(2):439-40.
14. Am J Trop Med Hyg 2002 Apr ;66(4):442-4.

Cutaneous larva migrans

Agent	PARASITE - Nematoda. Phasmidea: Ancylostoma braziliense, A. caninum, Bunostomum phlebotomum, Strongyloides myopotami
Reservoir	Cat Dog Cattle
Vector	None
Vehicle	Soil Contact
Incubation Period	2d - 3d (range 1d - 30d)
Diagnostic Tests	Biopsy is usually not helpful.
Typical Adult Therapy	Albendazole 200 mg BID X 3d OR Ivermectin 200 micrograms/kg as single dose. OR Thiabendazole topical, and oral 25 mg/kg BID X 5d (max 3g).
Typical Pediatric Therapy	Albendazole 2.5 mg/kg BID X 3d OR Ivermectin 200 micrograms/kg once OR Thiabendazole topical, and oral 25 mg/kg BID X 5d (max 3g).
Clinical Hints	Erythematous, serpiginous, pruritic advancing lesion(s) or bullae - usually on feet; follows contact with moist sand or beach front; may recur or persist for months.
Synonyms	Creeping eruption, Pelodera, Plumber's itch. ICD9: 126.2,126.8,126.9 ICD10: B76.9

Clinical

Cutaneous larva migrans is characterized by one or more erythematous linear, vesicular or bullous ¹ lesions which tend to be raised and palpable. ²⁻⁴

- The lesions are intensely pruritic and extend in length from day to day. ⁵
- The site of the lesions reflects contact with sand / soil, as from walking barefoot or lying on a beach. ⁶
- Infection may persist for over one year. ⁷

This disease is endemic or potentially endemic to all countries.

References

1. Acta Dermatovenerol Croat 2011 ;19(2):120-1.
2. Clin Dermatol 2003 Sep-Oct;21(5):407-16.
3. Cutis 2003 Aug ;72(2):111-5.
4. Lancet Infect Dis 2008 May ;8(5):302-9.
5. J Travel Med 2007 Sep-Oct;14(5):326-33.
6. Quintessence Int 2006 Oct ;37(9):721-3.
7. Int J Infect Dis 2012 Dec 4;

Cyclosporiasis

Agent	PARASITE - Protozoa. Sporozoa, Coccidea, Eimeriida: <i>Cyclospora cayetanensis</i>
Reservoir	Human ? Non-human primate
Vector	None
Vehicle	Water Vegetables
Incubation Period	1d - 11d
Diagnostic Tests	Identification of organism in stool smear. Cold acid fast stains and ultraviolet microscopy may be helpful.
Typical Adult Therapy	Sulfamethoxazole/trimethoprim 800/160 mg BID X 7d Ciprofloxacin 500 mg PO BID X 7 d (followed by 200 mg TIW X 2 w) has been used in sulfa-allergic patients
Typical Pediatric Therapy	Sulfamethoxazole/trimethoprim 10/2 mg/kg BID X 7d
Clinical Hints	Watery diarrhea (average 6 stools daily), abdominal pain, nausea, anorexia and fatigue lasting up to 6 weeks (longer in AIDS patients); most cases follow ingestion of contaminated water in underdeveloped countries.
Synonyms	<i>Cryptosporidium muris</i> , Cyanobacterium-like agent, <i>Cyclospora</i> . ICD9: 007.5 ICD10: A07.8

Clinical

Symptoms appear abruptly in 68% of cases

- Patients usually present with intermittent watery diarrhea, with up to eight or more stools per day. ^{1 2}
- Other symptoms may include anorexia, nausea, abdominal cramps, bloating, flatulence, mild to moderate weight loss, fatigue, and myalgia.
- Fever is rare.

In the immunocompetent patient, the diarrhea may last from a few days to up to three months, with the organism detectable in the stool for up to two months.

- In immune compromised individual, particularly AIDS patients, the disease can persist for weeks to several months.

Reactive arthritis syndrome (Reiter's syndrome) has been associated with progression of the disease. ³

Acalculous *Cyclospora* cholecystitis has been demonstrated in a patient with AIDS.

This disease is endemic or potentially endemic to all countries.

References

1. *Int J Parasitol* 2003 Apr ;33(4):371-91.
2. *Curr Opin Infect Dis* 2002 Oct ;15(5):519-22.
3. *Afr Health Sci* 2007 Jun ;7(2):62-7.

Cysticercosis

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Taeniidae: <i>Taenia solium</i>
Reservoir	Pig Human
Vector	None
Vehicle	Soil (contaminated by pigs) Fecal-oral Fly
Incubation Period	3m - 3y
Diagnostic Tests	Serology (blood or CSF) and identification of parasite in biopsy material.
Typical Adult Therapy	Albendazole 400 mg PO BID X 30d. OR Praziquantel 30 mg/kg TID X 14d (15 to 30d for neurocysticercosis). Surgery as indicated Add corticosteroids if brain involved.
Typical Pediatric Therapy	Albendazole 15 mg/kg PO BID X 30d. OR Praziquantel 30 mg/kg TID X 14d (15 to 30d for neurocysticercosis). Surgery as indicated Add corticosteroids if brain involved.
Clinical Hints	Cerebral, ocular or subcutaneous mass; usually no eosinophilia; calcifications noted on X-ray examination; lives in area where pork is eaten; 25% to 50% of patients have concurrent <i>Taenia</i> infestation.
Synonyms	<i>Taenia crassiceps</i> . ICD9: 123.1 ICD10: B69

Clinical

Cysticercosis is manifest as painless, rubbery (average 2 cm) nodules in skin and soft tissues, or other body sites. ¹⁻³

- "Rice grain" calcifications are often visible on routine roentgenograms of soft tissue, notably the pelvis and upper legs.
- Virtually any area of the body may be affected. ⁴⁻²⁴
- Cysticercosis involving the subcutaneous tissues may mimic malignancy or tuberculous lymphadenitis. ^{25 26}
- Rare instances of cysticercosis are reported in infants and young children. ^{27 28}

Central nervous system infection may present as seizures, increased intracranial pressure or hydrocephalus ²⁹⁻³¹, altered mental status, eosinophilic meningitis ³², ventriculitis ³³, focal neurological defects, stroke ³⁴, intramedullary ³⁵⁻³⁸ or extramedullary spinal mass ³⁹⁻⁴², quadriplegia ⁴³, spinal subarachnoid infection ⁴⁴ or encephalitis. ⁴⁵⁻⁴⁷

- In humans, cysticerci are more frequently located in the ventricles and subarachnoid space at the base of the brain, while in pigs, cysticerci are more frequently found in the parenchyma. ⁴⁸
- Parenchymal infestation and epilepsy are most common among children, while ventricular cysts with blockage of cerebrospinal fluid predominates among adults. ⁴⁹

The eyes are infested in 15% to 45% of patients ^{50 51}, usually presenting as a cyst in the vitreous cavity. ^{52 53}, less commonly the anterior chamber. ⁵⁴

- The first ophthalmologic signs of cysticercosis are papilledema, pupillary abnormalities, or nystagmus. ⁵⁵
- Cysticercosis of the extraocular muscles is associated with limitation of eye movement, ptosis, proptosis and local mass. ⁵⁶⁻⁶¹

This disease is endemic or potentially endemic to all countries.

Cysticercosis in Uganda

Prevalence surveys:

- 0.12% to 1.2% of pigs in Kampala slaughterhouses (*Taeniasis*, 1998 to 1999) ⁶²
- 33.7% of pigs in rural Lara (*Taeniasis*, northern region)

References

1. *J Laryngol Otol* 2008 Sep ;122(9):1005-7.
2. *Dentomaxillofac Radiol* 2008 Feb ;37(2):113-6.
3. *J Med Case Rep* 2008 ;2:196.
4. *Am J Trop Med Hyg* 2008 Dec ;79(6):864-5.

5. Trans R Soc Trop Med Hyg 2009 Feb ;103(2):206-8.
6. Indian Heart J 2008 May-Jun;60(3):260-2.
7. Ear Nose Throat J 2009 Nov ;88(11):1218-20.
8. Acta Cytol 2010 Sep-Oct;54(5 Suppl):853-6.
9. Indian J Plast Surg 2010 Jul ;43(2):210-2.
10. Indian Pediatr 2011 Feb ;48(2):141-3.
11. J Korean Neurosurg Soc 2011 Mar ;49(3):190-3.
12. J Glob Infect Dis 2011 Jul ;3(3):306-8.
13. Skeletal Radiol 2012 Sep ;41(9):1061-6.
14. Indian J Dent Res 2011 Jul-Aug;22(4):617.
15. J Oral Maxillofac Surg 2012 Feb 22;
16. J Oral Maxillofac Pathol 2011 May ;15(2):219-22.
17. Asian Pac J Trop Med 2012 Jul ;5(7):582-6.
18. Indian J Otolaryngol Head Neck Surg 2011 Jul ;63(Suppl 1):127-30.
19. J Lab Physicians 2012 Jan ;4(1):56-8.
20. Niger J Clin Pract 2012 Jul-Sep;15(3):361-3.
21. Urol Int 2012 Sep 5;
22. Indian J Dent Res 2012 May ;23(3):436.
23. J Clin Diagn Res 2012 Nov ;6(9):1555-6.
24. J Clin Diagn Res 2012 Dec ;6(10):1669-71.
25. J Clin Pathol 2010 Oct ;63(10):926-9.
26. Kathmandu Univ Med J (KUMJ) 2010 Apr-Jun;8(30):257-60.
27. Am J Trop Med Hyg 2009 Sep ;81(3):449-51.
28. Pathog Glob Health 2012 May ;106(2):122-3.
29. J Clin Neurosci 2011 Jun ;18(6):867-9.
30. Childs Nerv Syst 2011 Oct ;27(10):1709-21.
31. Br J Neurosurg 2012 Jun ;26(3):305-9.
32. Rev Inst Med Trop Sao Paulo 2007 Sep-Oct;49(5):331-4.
33. Rev Neurol (Paris) 2011 Aug-Sep;167(8-9):632-4.
34. Neurologist 2012 Jan ;18(1):17-22.
35. Acta Biomed 2008 Apr ;79(1):39-41.
36. Int J Med Sci 2011 ;8(5):420-3.
37. Asian J Neurosurg 2012 Apr ;7(2):90-2.
38. West J Emerg Med 2012 Nov ;13(5):434-6.
39. Acta Neurol Taiwan 2009 Sep ;18(3):187-92.
40. J Korean Neurosurg Soc 2010 Dec ;48(6):547-50.
41. Spine J 2011 Apr ;11(4):e1-5.
42. J Travel Med 2011 Jul-Aug;18(4):284-7.
43. BMJ Case Rep 2012 ;2012
44. Spine (Phila Pa 1976) 2012 May 11;
45. Neurol Res 2010 Apr ;32(3):229-37.
46. Surg Neurol 2005 Feb ;63(2):123-32; discussion 132.
47. Clin Neurol Neurosurg 2013 Jan 22;
48. Trop Med Int Health 2008 May ;13(5):697-702.
49. Childs Nerv Syst 2011 Oct ;27(10):1709-21.
50. Trop Doct 2003 Jul ;33(3):185-8.
51. Ophthalmol Clin North Am 2002 Sep ;15(3):351-6.
52. Ocul Immunol Inflamm 2011 Aug ;19(4):240-5.
53. Orbit 2011 Oct ;30(5):230-5.
54. Parasitol Int 2012 Jun ;61(2):378-80.
55. J AAPOS 2007 Oct ;11(5):495-6.
56. Strabismus 2008 Jul-Sep;16(3):97-106.
57. Eur J Ophthalmol 2010 Jan-Feb;20(1):240-2.
58. Cases J 2009 ;2:7025.
59. Ophthal Plast Reconstr Surg 2009 Nov-Dec;25(6):499-501.
60. Ophthalmology 2010 Mar ;117(3):600-5, 605.e1.
61. J Fr Ophtalmol 2012 Oct 31;
62. Acta Trop 2003 Jun ;87(1):13-23.

Cytomegalovirus infection

Agent	VIRUS - DNA. Herpesviridae, Betaherpesvirinae: Human herpesvirus 5 (Cytomegalovirus)
Reservoir	Human
Vector	None
Vehicle	Droplet (respiratory) Urine Dairy products Tears Stool Sexual contact (rare) Transplacental
Incubation Period	3w - 5w (range 2w - 12w)
Diagnostic Tests	Viral culture (blood, CSF, urine, tissue). Serology. Direct viral microscopy. Nucleic acid amplification
Typical Adult Therapy	[Most cases self-limited]. Ganciclovir 5 mg/kg q12h IV X 2 to 3w. OR Foscarnet 90 mg/kg Q12h IV
Typical Pediatric Therapy	[Most cases self-limited] Ganciclovir 5 mg/kg q12h IV X 2 to 3w
Vaccine	Cytomegalovirus immunoglobulin
Clinical Hints	Heterophile-negative "mononucleosis"; mild pharyngitis (without exudate); variable lymphadenopathy and splenomegaly; retinitis in AIDS patients; pneumonia in setting of immune suppression.
Synonyms	Cytomegalovirus, Zytomegalie. ICD9: 078.5 ICD10: B25

Clinical

Acute Cytomegalovirus infection is clinically similar to infectious mononucleosis (IM), and characterized by fever, generalized lymphadenopathy and hepatosplenomegaly. ¹

- In contrast to IM, pharyngitis is uncommon in Cytomegalovirus infection.
- Cytomegalovirus infection is often identified in cases of fatal myocarditis in immunocompetent patients. ²
- Primary CMV infection may be associated with uveitis ³, retinitis or pneumonia ⁴ • even in immunocompetent patients ⁵
⁶
- Additional manifestations of CMV infection may include prostatitis ⁷, adrenal failure ⁸, protracted diarrhea ⁹, esophagitis ¹⁰, gastritis ¹¹, duodenitis ¹², colitis with megacolon ^{13 14}, appendicitis ¹⁵, colonic pseudotumor ¹⁶ or colonic polyposis ¹⁷, pancreatitis ¹⁸, myocarditis ¹⁹ and protein-losing gastropathy (Menterier's disease). ²⁰
- Sexually-acquired Cytomegalovirus proctitis is characterized by rectal bleeding associated with a mononucleosis-like syndrome. ²¹
- The clinical features of Cytomegalovirus colitis in AIDS patients may mimic those of amebic colitis ^{22 23} or Crohn's disease. ²⁴
- Cases of pruritic maculo-papular exanthem due to CMV infection are reported among patients with AIDS. ²⁵
- Evidence for primary CMV infection is often present among infants hospitalized for wheezing. ²⁶
- Ocular infection may present as inflammatory ocular hypertensive syndrome (IOHS), corneal endothelitis, or retinitis with retinal necrosis. ^{27 28}
- Rare instances of erythema multiforme complicating Cytomegalovirus infection have been reported. ^{29 30}
- CMV / EBV co-infection may be associated with prolonged illness. ³¹

Severe or fatal multisystem disease occurs is encountered in congenital infection ³²⁻³⁷ and infection of immune-suppressed individuals. ³⁸⁻⁴¹

- Instances of pure red-cell aplasia ⁴² and hemophagocytic syndrome have been reported. ⁴³
- Sensorineural hearing loss detected in 21% of asymptomatic and 33% of symptomatic congenital infections ⁴⁴⁻⁴⁹
- Residual neurological damage including epilepsy is common among infants with congenital infection. ⁵⁰
- Rare instances of persistent pulmonary hypertension have been reported in infants with congenital infection. ⁵¹

Immunocompetent persons may also develop major complications ⁵², including cerebral sinus thrombosis ⁵³; peripheral venous ⁵⁴⁻⁶², mesenteric ⁶³⁻⁶⁷ or portal vein thrombosis ⁶⁸⁻⁷⁸, colitis ⁷⁹, transverse myelitis ⁸⁰, hemolytic anemia ⁸¹, rhabdomyolysis ^{82 83}, prostatitis ⁸⁴ and cholecystitis. ⁸⁵

This disease is endemic or potentially endemic to all countries.

References

1. Lancet Infect Dis 2004 Dec ;4(12):725-38.
2. Clin Infect Dis 2005 Mar 1;40(5):683-8.
3. Medicine (Baltimore) 2008 May ;87(3):167-76.
4. Infect Dis Clin North Am 2010 Mar ;24(1):147-58.
5. J Neuroophthalmol 2008 Jun ;28(2):126-7.
6. Eur J Ophthalmol 2008 Sep-Oct;18(5):813-5.
7. Int J Surg Pathol 2010 Feb ;18(1):55-9.
8. Eur J Pediatr 2010 Jul ;169(7):887-9.
9. An Pediatr (Barc) 2009 Jun ;70(6):582-5.
10. Dis Esophagus 2010 Aug ;23(6):E32-4.
11. World J Gastrointest Endosc 2010 Nov 16;2(11):379-80.
12. Scand J Gastroenterol 2011 Oct ;46(10):1228-35.
13. J Med Virol 2010 Apr ;82(4):638-41.
14. BMC Pediatr 2011 ;11:52.
15. Transpl Infect Dis 2012 Oct 17;
16. Case Rep Gastroenterol 2012 May ;6(2):266-72.
17. Pathology 2011 Aug ;43(5):440-6.
18. JOP 2011 Mar ;12(2):158-61.
19. Intern Med 2010 ;49(2):131-3.
20. Acta Gastroenterol Belg 2007 Jul-Sep;70(3):296-9.
21. Sex Transm Dis 2011 Sep ;38(9):876-8.
22. Am J Med Sci 2008 Oct ;336(4):362-4.
23. Gastroenterol Hepatol 1998 Jan ;21(1):26-7.
24. World J Gastroenterol 2009 Sep 14;15(34):4327-30.
25. Int J Dermatol 2008 Sep ;47(9):944-6.
26. Pediatr Int 2008 Oct ;50(5):654-7.
27. Graefes Arch Clin Exp Ophthalmol 2011 Jan ;249(1):103-10.
28. Eye (Lond) 2012 Feb ;26(2):237-40.
29. J Cutan Med Surg 2011 Mar-Apr;15(2):115-7.
30. Pediatr Dermatol 2012 May 29;
31. J Med Virol 2009 Aug ;81(8):1399-402.
32. Curr Opin Obstet Gynecol 2011 Apr ;23(2):123-8.
33. Pediatr Clin North Am 2004 Aug ;51(4):889-908, viii.
34. Curr Opin Infect Dis 2005 Jun ;18(3):241-5.
35. J Infect Dis 2007 Mar 15;195(6):782-8.
36. Semin Fetal Neonatal Med 2007 Jun ;12(3):154-9.
37. Indian J Pathol Microbiol 2008 Jul-Sep;51(3):402-4.
38. Transplantation 2005 Feb 27;79(4):381-6.
39. Curr Infect Dis Rep 2006 May ;8(3):222-30.
40. World J Gastroenterol 2008 Aug 21;14(31):4849-60.
41. J Med Case Rep 2008 ;2:289.
42. J Pediatr Hematol Oncol 2010 May ;32(4):315-6.
43. Braz J Infect Dis 2009 Feb ;13(1):72-3.
44. J Pediatr 2008 Jul ;153(1):84-8.
45. Int J Pediatr Otorhinolaryngol 2011 Mar ;75(3):376-82.
46. Otolaryngol Head Neck Surg 2011 May ;144(5):662-75.
47. Georgian Med News 2011 Jun ;(195):61-5.
48. Int J Pediatr Otorhinolaryngol 2012 Mar 1;
49. Acta Paediatr 2012 Apr 20;
50. Brain Dev 2008 Jun ;30(6):420-4.
51. Pulm Med 2011 ;2011:293285.
52. Virol J 2008 ;5:47.
53. J Med Virol 2012 Dec ;84(12):1934-6.
54. Clin Microbiol Infect 2010 Jul ;16(7):851-4.
55. J Thromb Thrombolysis 2009 Nov ;28(4):496-9.
56. Isr Med Assoc J 2007 Oct ;9(10):757-8.
57. Am J Med Sci 2007 Aug ;334(2):111-4.
58. Ann Clin Microbiol Antimicrob 2005 ;4:11.
59. Clin Infect Dis 2003 Jun 1;36(11):E134-9.
60. J Infect 2003 Feb ;46(2):141-2.
61. Thromb Res 2010 Dec ;126(6):e439-43.
62. Ann Hematol 2012 Apr ;91(4):597-604.
63. Blood Coagul Fibrinolysis 2007 Jul ;18(5):509-11.
64. Eur J Gastroenterol Hepatol 2006 Apr ;18(4):443-5.
65. Enferm Infecc Microbiol Clin 2002 Feb ;20(2):96-7.
66. Clin Infect Dis 2001 Mar 15;32(6):983-6.
67. J Med Case Reports 2012 Apr 24;6(1):118.
68. J Thromb Thrombolysis 2009 Nov ;28(4):496-9.
69. Clin Infect Dis 2007 Jan 15;44(2):e13-6.
70. Thromb Haemost 2006 Jan ;95(1):199-201.
71. BMC Gastroenterol 2006 ;6:10.
72. Rev Med Interne 2006 May ;27(5):426-8.
73. Rev Med Interne 2006 Jan ;27(1):54-8.
74. Presse Med 2001 Dec 15;30(38):1876-8.
75. Eur J Intern Med 2011 Apr ;22(2):195-9.
76. Recent Prog Med 2011 Jul-Aug;102(7-8):294-5.
77. J Med Virol 2012 Jan ;84(1):116-8.
78. J Thromb Thrombolysis 2012 Feb ;33(2):160-72.
79. Singapore Med J 2011 Sep ;52(9):e170-2.
80. BMC Res Notes 2012 ;5:364.
81. J Med Case Rep 2010 ;4:334.
82. BMJ Case Rep 2013 ;2013
83. Med Clin (Barc) 1991 Feb 16;96(6):220-1.
84. Am J Transplant 2011 Jun ;11(6):1330-3.
85. Am J Transplant 2009 May ;9(5):1249-52.

Dengue

Agent	VIRUS - RNA. Flaviviridae, Flavivirus: Dengue virus
Reservoir	Human Mosquito ? Monkey (in Malaysia and Africa)
Vector	Mosquito - Stegomyia (Aedes) aegypti, S. albopictus, S. polynesiensis, S. scutellaris
Vehicle	Blood (rare)
Incubation Period	5d - 8d (range 2d - 15d)
Diagnostic Tests	Viral isolation (blood). Serology. Nucleic acid amplification. Biosafety level 2.
Typical Adult Therapy	Supportive; IV fluids to maintain blood pressure and reverse hemoconcentration
Typical Pediatric Therapy	As for adult
Clinical Hints	Headache, myalgia, arthralgia, relative bradycardia, leukopenia and macular rash; dengue hemorrhagic (DHF) = dengue + thrombocytopenia and hemoconcentration; dengue shock = DHF + hypotension.
Synonyms	Bouquet fever, Break-bone fever, Dandy fever, Date fever, Dengue Fieber, Duengero, Giraffe fever, Petechial fever, Polka fever. ICD9: 061 ICD10: A90,A91

Clinical

WHO Case definitions for surveillance:

1. DENGUE FEVER:

Clinical description

• An acute febrile illness of 2-7 days duration with 2 or more of the following: headache, retro-orbital pain, myalgia, arthralgia (as many as 41% of cases ¹ , rash, hemorrhagic manifestations, leucopenia.

Laboratory criteria for diagnosis • One or more of the following:

- Isolation of the dengue virus from serum, plasma, leukocytes, or autopsy samples
- Demonstration of a fourfold or greater change in reciprocal IgG or IgM antibody titers to one or more dengue virus antigens in paired serum samples
- Demonstration of dengue virus antigen in autopsy tissue by immunohistochemistry or immunofluorescence or in serum samples by EIA
- Detection of viral genomic sequences in autopsy tissue, serum or CSF samples by polymerase chain reaction (PCR)

Case classification

- Suspected: A case compatible with the clinical description.
- Probable: A case compatible with the clinical description with one or more of the following:
 - Supportive serology (reciprocal hemagglutination-inhibition antibody titer >1280, comparable IgG EIA titer or positive IgM antibody test in late acute or convalescent-phase serum specimen).
 - Occurrence at same location and time as other confirmed cases of dengue fever.
- Confirmed: A case compatible with the clinical description, laboratory confirmed.

2. DENGUE HEMORRHAGIC FEVER:

A probable or confirmed case of dengue and hemorrhagic tendencies evidenced by one or more of the following:

- Positive tourniquet test (sensitivity questioned • see reference ²)
- Petechiae, ecchymoses or purpura
- Bleeding: mucosa, gastrointestinal tract, injection sites or other
- Hematemesis or melena
- And thrombocytopenia (100 000 cells or less per mm³)
- And evidence of plasma leakage due to increased vascular permeability, manifested by one or more of the following:
 - 20% rise in average hematocrit for age and sex
 - 20% drop in hematocrit following volume replacement treatment compared to baseline
 - signs of plasma leakage (pleural effusion, ascites, hypoproteinemia)

3. DENGUE SHOCK SYNDROME:

All the above criteria, plus evidence of circulatory failure manifested by rapid and weak pulse, and narrow pulse pressure (<=20 mm Hg) or hypotension for age, cold, clammy skin and altered mental status.

CDC case definition:

For surveillance purposes, the U.S. Centers for Disease Control (CDC) case definition of dengue fever consists of "acute febrile illness characterized by frontal headache, retro-ocular pain, muscle and joint pain, and rash."

- The initial fever rises rapidly and lasts for two to seven days.
- Occasionally "saddleback" fever pattern is evident, with a drop after a few days and rebound within 24 hours.³ Relative bradycardia is common.⁴
- Conjunctival injection and pharyngeal inflammation may occur as well as lymphadenopathy.
- Rash occurs in up to 50 percent of patients, either early in the illness with flushing or mottling, or between the 2nd to the 6th day as a scarlatiniform or maculopapular rash that usually spreads centrifugally.
- The later rash usually lasts for two to three days.
- Diffuse erythema and late desquamation of hands and feet may be confused with toxic shock syndrome.
- As fever drops, petechiae may be seen.
- Additional manifestations of dengue may include post-dengue depression, acalculous cholecystitis, uveitis, retinitis and psychological depression.

Additional clinical features:

- The likelihood of encountering classic clinical findings of dengue fever increases with patient age.⁵
- The rash of dengue may be mistaken for measles or rubella.⁶
- A long time interval between attacks of dengue may actually increase the risk of dengue hemorrhagic fever.⁷
- Rare instances of encephalopathy^{8 9}, seizures, splenic rupture^{10 11}, pancreatitis¹² myocarditis, pericarditis¹³ and aplastic anemia complicating dengue are reported.¹⁴
- Hepatic dysfunction is often encountered¹⁵⁻¹⁷; however, overt hepatitis is less common.¹⁸
- Retinal involvement may manifest as foveolitis, which can be diagnosed by funduscopy and optical coherence tomography¹⁹
- Prolonged post-dengue fatigue is common.²⁰
- Renal failure is associated with increased mortality rates in dengue.²¹
- Risk factors for fatal dengue hemorrhagic fever among elderly patients in include male sex, chronic obstructive pulmonary disease, dengue shock syndrome and acute renal failure.²²

The diagnosis of Dengue Hemorrhagic Fever (DHF) is defined by²³

- thrombocytopenia (<100,000/mm³)
- evidence of plasma leakage (hematocrit increased by at least 20%) or other objective evidence of increased capillary permeability
- Dengue Shock Syndrome (DSS) consists of DHF in addition to hypotension or narrow pulse pressure (less than 21 mm Hg).²⁴

Note that Leptospirosis²⁵, Zika, Crimean-Congo hemorrhagic fever²⁶ and dengue are clinically similar, and may coexist in a given country.

- Although the clinical features of dengue and chikungunya are similar, chikungunya patients are more likely to exhibit early myalgia or arthralgia; while sore throat, cough, nausea, vomiting, diarrhea, abdominal pain, anorexia, tachycardia and thrombocytopenia will favor a diagnosis of dengue.²⁷

This disease is endemic or potentially endemic to 144 countries. Although Dengue is not endemic to Uganda, imported, expatriate or other presentations of the disease have been associated with this country.

Dengue in Uganda

Notable outbreaks:

2011 - An outbreak (122 cases) of dengue was reported among Ugandan military personnel serving in Somalia.²⁸

References

1. Clin Rheumatol 2009 Sep ;28(9):1067-71.
2. J Assoc Physicians India 1999 Feb ;47(2):203-4.
3. Postgrad Med J 2004 Oct ;80(948):588-601.
4. Emerg Infect Dis 2008 Feb ;14(2):350-1.
5. Emerg Infect Dis 2007 Jun ;13(6):924-5.
6. Rev Soc Bras Med Trop 2008 Jul-Aug;41(4):338-44.
7. Trans R Soc Trop Med Hyg 2008 Jun ;102(6):522-3.
8. Am J Trop Med Hyg 1996 Mar ;54(3):253-5.
9. BMJ Case Rep 2013 ;2013
10. Trop Med Parasitol 1993 Dec ;44(4):327-8.
11. Braz J Infect Dis 2003 Dec ;7(6):423-5.
12. Acta Med Indones 2012 Jan ;44(1):57-61.
13. BMC Res Notes 2011 ;4:268.
14. Travel Med Infect Dis 2009 Mar ;7(2):118-20.
15. Am J Trop Med Hyg 1992 Sep ;47(3):265-70.
16. Am J Trop Med Hyg 2010 Oct ;83(4):774-80.
17. Iran J Pediatr 2012 Jun ;22(2):231-6.
18. Asian Pac J Trop Med 2011 Apr ;4(4):323-4.
19. Ophthalmologica 2008 ;222(5):317-20.
20. J Clin Virol 2007 Jan ;38(1):1-6.
21. Clin J Am Soc Nephrol 2008 Sep ;3(5):1350-6.
22. Am J Trop Med Hyg 2008 Aug ;79(2):149-53.
23. Lancet 2007 Nov 10;370(9599):1644-52.
24. Clin Microbiol Rev 1998 Jul ;11(3):480-96.
25. Trop Doct 2010 Apr ;40(2):92-4.
26. J Infect Dev Ctries 2010 Jul ;4(7):459-63.
27. PLoS Negl Trop Dis 2012 Sep ;6(9):e1786.
28. ProMED <promedmail.org> archive: 20110919.2845

Dermatophytosis

Agent	FUNGUS. Ascomycota, Euascomyces, Onygenales: Epidermophyton, Microsporum, Trichophyton, Trichosporon spp., Arthroderma, et al
Reservoir	Human Dog Cat Rabbit Marsupial Other mammal
Vector	None
Vehicle	Contaminated soil/flooring Animal contact
Incubation Period	2w - 38w
Diagnostic Tests	Fungal culture and microscopy of skin, hair or nails. Nucleic acid amplification.
Typical Adult Therapy	Skin - topical Clotrimazole, Miconazole, etc. Hair/nails - Terbinafine, Griseofulvin, Itraconazole or Fluconazole PO
Typical Pediatric Therapy	As for adult
Clinical Hints	Erythematous, circinate, scaling or dyschromic lesions of skin, hair or nails; pruritus, secondary infection and regional lymphadenopathy may be present.
Synonyms	Arthroderma, DermatOMICOSE, DermatomyCose, DermatomyCosis, DermatomyKose, DermatomyKosen, Emericella, Favus, Granuloma trichophyticum, Gruby's disease, Kodamaea, Leukonychia trichophytica, Microsporum, Natrassia, Onychocola, Onychomycosis, Pityriasis versicolor, Ringworm, Saint Aignan's disease, Scopulariopsis, Scytalidium, Tinea, Tinea barbae, Tinea capitis, Tinea corporis, Tinea cruris, Tinea favosa, Tinea imbricata, Tinea manum, Tinea pedis, Tinea unguinum, Tokelau ringworm, Triadelphia pulvinata, Trichomycosis, Trichophytosis, Trichophytosis gladiatorum. ICD9: 110,111 ICD10: B35,B36

Clinical

Dermatophytosis is characterized by indolent infection of skin, hair or nails. ^{1 2}

Common findings include scaling, pruritis and discoloration • usually without overt signs of inflammation.

Tinea imbricata, a superficial mycosis caused by *Trichophyton concentricum*, an anthropophilic dermatophyte.

- The skin lesions are characteristically concentric and lamellar (imbricata: in Latin, tiled) plaques of scale. ³
- Predisposing conditions include humidity, inheritance, and immunologic factors. ⁴

Rare instances of mycetoma of the scalp due to *Microsporum canis* have been reported. ⁵

This disease is endemic or potentially endemic to all countries.

References

1. Dermatol Ther 2004 ;17(6):517-22.
2. Lancet 2004 Sep 25-Oct 1;364(9440):1173-82.
3. Trans R Soc Trop Med Hyg 2008 Apr ;102(4):389-93.
4. Int J Dermatol 2004 Jul ;43(7):506-10.
5. Diagn Microbiol Infect Dis 2011 May ;70(1):145-9.

Dientamoeba fragilis infection

Agent	PARASITE - Protozoa. Archezoa, Parabasala, Trichomonadea. Flagellate: Dientamoeba fragilis
Reservoir	Human Gorilla Pig
Vector	None
Vehicle	Fecal-oral (? on pinworm ova)
Incubation Period	8d - 25d
Diagnostic Tests	Identification of trophozoites in stool. Nucleic acid amplification. Alert laboratory if this diagnosis is suspected.
Typical Adult Therapy	Stool precautions. Iodoquinol 650 mg PO TID X 20d. OR Tetracycline 500 mg QID X 10d. OR Paromomycin 10 mg/kg TID X 7d OR Metronidazole 750 mg PO TID X 10d
Typical Pediatric Therapy	Stool precautions. Iodoquinol 13 mg/kg PO TID X 20d. OR (age >8) Tetracycline 10 mg/kg QID X 10d OR Paromomycin 10 mg/kg TID X 7d OR Metronidazole 15 mg/kg PO TID X 10d
Clinical Hints	Abdominal pain with watery or mucous diarrhea; eosinophilia may be present; infestation may persist for more than one year.
Synonyms	

Clinical

Most infections are asymptomatic.

- Symptoms may include diarrhea, flatulence, abdominal pain, fatigue and anorexia; and may rarely mimic acute appendicitis. ¹⁻³
- Clinical features are similar to those of giardiasis; however, vomiting, anorexia and weight loss are less common in *Dientamoeba* infection. ⁴
- The presence of abdominal pain or diarrhea in a patient with enterobiasis should suggest the diagnosis of concurrent *Dientamoeba* infection. ⁵

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 2004 Jul ;17(3):553-70, table of contents.
2. Am J Trop Med Hyg 2010 Apr ;82(4):614-9.
3. Parasitology 2011 Apr 27;:1-5.
4. Int J Infect Dis 2006 May ;10(3):255-61.
5. Trends Parasitol 2006 Feb ;22(2):92-6.

Diphtheria

Agent	BACTERIUM. Corynebacterium diphtheriae A facultative gram-positive bacillus
Reservoir	Human
Vector	None
Vehicle	Droplet Contact Dairy products Clothing
Incubation Period	2d - 5d (range 1d - 10d)
Diagnostic Tests	Culture on special media. Advise laboratory when this diagnosis is suspected.
Typical Adult Therapy	Respiratory isolation. Equine antitoxin 20,000 to 80,000 units IM. Erythromycin 500 mg QID X 10d
Typical Pediatric Therapy	Respiratory isolation. Equine antitoxin 1,000 units/kg IM. Erythromycin 10 mg/kg QID X 10d
Vaccines	Diphtheria antitoxin Diphtheria DTP DT DTaP Td
Clinical Hints	Pharyngeal membrane with cervical edema and lymphadenopathy; or punched out skin ulcers with membrane; myocarditis or neuropathy (foot/wrist drop) appears weeks later.
Synonyms	<i>Corynebacterium diphtheriae</i> , Difteri, Difteria, Diferie, Diferite, Diphterie. ICD9: 032 ICD10: A36

Clinical

WHO Case definition for surveillance:

Clinical description

- An illness of the upper respiratory tract characterized by laryngitis or pharyngitis or tonsillitis, and adherent membranes of tonsils, pharynx and/or nose

Laboratory criteria for diagnosis

- Isolation of *Corynebacterium diphtheriae* from a clinical specimen.
- Note: A rise in serum antibody (fourfold or greater) is of interest only if both serum samples were obtained before administration of diphtheria toxoid or antitoxin. This is not usually the case in surveillance, where serological diagnosis of diphtheria is thus unlikely to be an issue.

Case classification

- Suspected: Not applicable.
- Probable: A case that meets the clinical description.
- Confirmed: A probable case that is laboratory confirmed or linked epidemiologically to a laboratory confirmed case.

Note: Persons with positive *C. diphtheriae* cultures who do not meet the clinical description (i.e. asymptomatic carriers) should not be reported as probable or confirmed diphtheria cases.

Faucal diphtheria:

Following an incubation period of 2 to 5 days (7 days after primary skin infection for cutaneous diphtheria), the patient presents with nonspecific symptom which may include fever and chills, malaise, sore throat, hoarseness or dysphagia, cervical edema and lymphadenopathy, rhinorrhea (mucopurulent or blood-tinged), cough, stridor, wheezing, nausea and vomiting and headache. ¹

- Respiratory diphtheria may progress rapidly to respiratory arrest from airway obstruction by a tracheobronchial pseudomembrane.
- Tachycardia, pallor, and foul breath may be present.
- The pseudomembrane is generally firm, adherent, thick, fibrinous and of a gray-brown color.
- It may occur over the palate, pharynx, epiglottis, larynx, or trachea • occasionally extending into the tracheobronchial tree.
- The area may bleed if disturbed.
- Marked edema of the tonsils, uvula, submandibular region and anterior neck ("bull neck) may be observed and may be associated with thick speech, stridor, anterior cervical lymphadenopathy, and petechial hemorrhages.

Cutaneous diphtheria:

Cutaneous diphtheria is associated with a history of a break in the skin, followed by pain, tenderness, erythema, or exudate.

- Lesions appear as punched-out ulcers with dirty gray membranes at their margins.
- Genital ulcers may be misdiagnosed as venereal disease. ²

Cardiac complications:

Cardiovascular signs ensue 1 to 2 weeks following the initial illness.

- Myocarditis occurs in as many as two thirds of patients, and approximately 20% develop cardiac dysfunction.
- Circulatory collapse, heart failure, atrioventricular blocks and arrhythmias may occur.
- Endocarditis and mycotic aneurysms also have been reported, typically in intravenous drug users.

Neurological complications:

Approximately 70% of patients with severe infection develop neuropathy, neuritis or motor paralysis 2 to 8 weeks following initial illness.

- Clinical and cerebrospinal fluid findings at this stage are indistinguishable from those Guillain-Barre syndrome.
- Potentially fatal paralysis of the diaphragm may ensue.
- Paralysis typically resolves completely with resolution of infection.

The neurological manifestations of diphtheria include:

- hypesthesia and paralysis of the soft palate
- weakness of the posterior pharyngeal, laryngeal, and facial nerves, resulting in a "nasal tone" to the voice, difficulty in swallowing, and occasionally aspiration
- cranial neuropathies, typically during the fifth week, leading to oculomotor and ciliary paralysis (strabismus, blurred vision, and loss of accommodation)
- symmetric polyneuropathy beginning within 10 days to 3 months after infection, and manifest as motor deficit with diminished deep tendon reflexes
- proximal muscle weakness of the extremities progressing distally (or distal weakness progressing proximally).

Other forms of diphtheria:

Other less common manifestations include infection of the genitourinary tract, gastrointestinal tract, vagina, external ear, and conjunctiva.

- Hemorrhagic conjunctivitis and dissolution of the cornea may occur. ³
- Focal necrosis of the kidneys ⁴, liver, and adrenal glands may be observed.
- Cases of septic arthritis, osteomyelitis, splenic abscesses, and bacteremia have been reported.

A rare case of diphtherial urethritis was acquired through orogenital contact. ⁵

This disease is endemic or potentially endemic to all countries.

Diphtheria in Uganda

Vaccine Schedule:

BCG - birth

DTwP - 6, 10, 14 weeks


DTwPHibHep - 6, 10, 14 weeks

Measles (monovalent) - 9 months


OPV - birth; 6, 10, 14 weeks

TT - Women of child-bearing age; +1, +6 months; +1, +1 year

Vitamin A - 6, 12, 18, 24, 30, 36, 42 months



Graph: Uganda. Diphtheria - WHO-UNICEF est. vaccine (DTP3 %) coverage



Graph: Uganda. Diphtheria, cases

References

1. J Infect Dis 2000 Feb ;181 Suppl 1:S110-5.
2. Indian J Dermatol Venereol Leprol 2008 Mar-Apr;74(2):187.
3. Arch Ophthalmol 1978 Jan ;96(1):53-6.
4. Lik Sprava 2010 Oct-Dec;(7-8):44-53.
5. Sex Transm Infect 2012 May 23;

Diphyllobothriasis

Agent	PARASITE - Platyhelminthes, Cestoda. Pseudophyllidea, Diphylobothriidae: <i>Diphyllobothrium latum</i> , et al
Reservoir	Human Dog Bear Fish-eating mammal
Vector	None
Vehicle	Fresh-water fish - notably (for <i>D. latum</i>) perch, burbot and pike
Incubation Period	4w - 6w (range 2w - 2y)
Diagnostic Tests	Identification of ova or proglottids in feces.
Typical Adult Therapy	Praziquantel 10 mg/kg PO as single dose OR Niclosamide 2 g PO once
Typical Pediatric Therapy	Praziquantel 10 mg/kg PO as single dose OR Niclosamide 50 mg/kg PO once
Clinical Hints	Abdominal pain, diarrhea and flatulence; vitamin B12 deficiency is noted in 0.02% of patients; rare instances of intestinal obstruction have been described; worm may survive for decades in human intestine.
Synonyms	Bandwormer [<i>Diphyllobothrium</i>], <i>Bothriocephalus latus</i> , Broad fish tapeworm, <i>Diphyllobothrium latum</i> , Diplogonoporiasis, Fish tapeworm. ICD9: 123.4 ICD10: B70.0

Clinical

Patients may experience abdominal pain, diarrhea, weight loss, asthenia or vertigo. ¹

- Vitamin B-12 deficiency is described in cases of prolonged infestation by *Diphyllobothrium latum* ²⁻¹¹ (but not other *Diphyllobothrium* species ^{12 13}).

This disease is endemic or potentially endemic to all countries.

References

1. Parasitol Res 2003 Nov ;91(5):412-38.
2. Acta Med Scand 1947 Oct 15;129(1):59-76.
3. Blood 1948 Jan ;3(1):91-102.
4. Acta Med Scand Suppl 1950 ;259:112-22.
5. Br Med J 1950 Jul 22;2(4672):188-92.
6. Acta Med Scand Suppl 1952 ;271:1-68.
7. Exp Parasitol 1956 Mar ;5(2):207-30.
8. Nature 1956 Oct 27;178(4539):934-5.
9. Am J Clin Nutr 1961 Sep-Oct;9:606-12.
10. World Med J 1964 May ;11:170-2.
11. Br Med J 1976 Oct 30;2(6043):1028.
12. Am J Trop Med Hyg 2012 Nov ;87(5):897-901.
13. Arctic Med Res 1991 Jan ;50(1):18-24.

Dipylidiasis

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Dipylidiidae: <i>Dipylidium caninum</i>
Reservoir	Dog Cat
Vector	None
Vehicle	Flea = <i>Ctenocephalides</i> spp. (by ingestion)
Incubation Period	21d - 28d
Diagnostic Tests	Identification of proglottids in feces.
Typical Adult Therapy	Praziquantel 10 mg/kg PO as single dose OR Niclosamide 2 g PO once
Typical Pediatric Therapy	Praziquantel 10 mg/kg PO as single dose OR Niclosamide 50 mg/kg PO once
Clinical Hints	Diarrhea, abdominal distention and restlessness (in children); eosinophilia may be observed; proglottids may migrate out of anus.
Synonyms	Cucumber tapeworm, <i>Dipylidium caninum</i> , Dog tapeworm, Double-pored dog tapeworm. ICD9: 123.8 ICD10: B71.1

Clinical

Most infections with *Dipylidium caninum* are asymptomatic.

- Severe diarrhea, urticaria, fever and eosinophilia are occasionally encountered. ¹
- The principal sign (in animals and children) consists of the passage of proglottids on the perianal region, feces, diapers, or occasionally on floor covering and furniture.
- Infection has been reported in patients as young as four months ² to two years. ³
- Proglottids are motile when freshly passed and may be mistaken for maggots or fly larvae.

This disease is endemic or potentially endemic to all countries.

References

1. *Parasitol Res* 2003 Nov ;91(5):412-38.
2. *Clin Lab Sci* 2011 ;24(4):212-4.
3. *Rev Chilena Infectol* 2008 Dec ;25(6):465-71.

Dirofilariasis

Agent	PARASITE - Nematoda. Phasmidea, Filariae: <i>Dirofilaria</i> (<i>Nochtiella</i>) <i>immitis</i> (pulmonary); <i>D. tenuis</i> & <i>D. repens</i> (subcutaneous infection) & <i>D. ursi</i>
Reservoir	Mammal Dog Wild carnivore (<i>D. tenuis</i> in raccoons; <i>D. ursi</i> in Bears)
Vector	Mosquito
Vehicle	None
Incubation Period	60d - 90d
Diagnostic Tests	Identification of parasite in tissue (ie, lung biopsy). Serologic tests available in some centers.
Typical Adult Therapy	Not available; excision is often diagnostic and curative
Typical Pediatric Therapy	As for adult
Clinical Hints	Most patients are asymptomatic; occasional instances of cough and chest pain, with solitary pulmonary coin lesion; or multiple tender subcutaneous nodules; eosinophilia usually not present.
Synonyms	Candidatus <i>Dirofilaria hongkongensis</i> , <i>Dirofilariosis</i> , <i>Dirofilaria</i> , Dog heartworm, <i>Filaria conjunctivae</i> , <i>Loaina</i> , <i>Pelecitus</i> . ICD9: 125.6 ICD10: B74.8

Clinical

Pulmonary infections usually present as a well-circumscribed coin lesion. ¹

- Occasionally the lesions are transient or multiple. ²
- Symptoms such as chest pain, dyspnea, fever, cough and eosinophilia are present in only 50% of cases.
- Isolated infections have been reported in the mesentery, spermatic cord, epididymis ³, peritoneal cavity ⁴, anterior chamber of the eye ⁵, orbital muscles ⁶ and liver.
- Lesions may suggest malignancy ⁷, and coexistence of dirofilariasis and lung cancer has been reported. ⁸
- In rare cases pulmonary cavitation may occur ⁹

Skin and subcutaneous infections are caused by *D. tenuis*, *D. repens* ¹⁰, *D. ursi*, *D. immitis* and *D. striata*.

- Clinical manifestations are limited to a small (0.5 to 1.5 cm) discrete nodule which may appear on any area of the body. ¹¹⁻¹³
- Local pain, inflammation, eosinophilia and a sensation of motion may be present in some cases.
- Rare instances of scrotal pseudotumor are reported. ¹⁴

A novel *Dirofilaria* species ("*Candidatus Dirofilaria hongkongensis*") has been identified as a cause of human (cervical lymphadenopathy, abdominal subcutaneous mass and subconjunctival nodule) and canine infection in Hong Kong. ¹⁵

This disease is endemic or potentially endemic to 228 countries.

References

1. South Med J 1999 Mar ;92(3):276-9.
2. Vet Parasitol 2005 Oct 24;133(2-3):157-80.
3. Urology 2009 Jan ;73(1):209.e1-3.
4. Magy Seb 2008 Oct ;61(5):281-4.
5. Middle East Afr J Ophthalmol 2012 Jul-Sep;19(3):349-51.
6. Korean J Parasitol 2009 Dec ;47(4):397-9.
7. Pediatr Blood Cancer 2009 Sep ;53(3):485-7.
8. J Infect 2008 Apr ;56(4):241-3.
9. Nihon Kokyuki Gakkai Zasshi 2009 May ;47(5):372-5.
10. Emerg Infect Dis 2007 Jan ;13(1):150-2.
11. Int J Surg Pathol 2008 Jan ;16(1):101-3.
12. Eur J Ophthalmol 2009 May-Jun;19(3):475-7.
13. Chir Main 2011 Feb ;30(1):66-8.
14. Pathog Glob Health 2012 Oct ;106(6):370-2.
15. J Clin Microbiol 2012 Nov ;50(11):3534-41.

Dracunculiasis

Agent	PARASITE - Nematoda. Phasmidea, Filariae: <i>Dracunculus medinensis</i>
Reservoir	Human
Vector	None
Vehicle	Copepod (<i>Mesocyclops</i> and <i>Thermocyclops</i>) in drinking water
Incubation Period	12m - 18m
Diagnostic Tests	Identification of adult worm in situ; or identification of discharged larvae from wound.
Typical Adult Therapy	Worm removal Metronidazole 500 mg PO TID X 10d. OR Thiabendazole 30 mg/kg PO BID X 3d have been used to facilitate worm removal.
Typical Pediatric Therapy	Worm removal Metronidazole 8 mg/kg PO TID X 10d. OR Thiabendazole 30 mg/kg BID X 3d have been used to facilitate worm removal.
Clinical Hints	Nausea and urticaria followed by the appearance of a papule or bulla (usually lower leg) which ruptures; calcified worm on x-ray; occasional eosinophilia; worm may survive for 18 months in human.
Synonyms	Dracunculose, <i>Dracunculus medinensis</i> , Dracontiasis, <i>Filaria medinensis</i> , Guinea worm, Medina worm. ICD9: 125.7 ICD10: B72

Clinical

WHO Case definition for surveillance:

Clinical case definition

- A case of dracunculiasis is defined as an individual exhibiting or having a history of a skin lesion with the emergence of a Guinea worm.
- A recent history (within one year) of a skin lesion with emergence of a Guinea worm (*Dracunculus medinensis*) is the only time-frame which must be used in surveillance programs.

As the adult female migrates, a blister develops on the skin where the worm will emerge. ¹

- Symptoms consist of fever, urticaria and other allergic phenomena, swelling, and local pain and burning. ²
- The blister will eventually rupture, and the patient seeks relief through immersing the affected skin in water.
- The resulting temperature change causes the blister to erupt, exposing the worm, which then releases a milky white liquid containing millions of larvae into the water.
- The process of larval shedding continues for several days after it has emerged from the ulcer.

More than 90% of the worms appear on the legs and feet, but may occur anywhere on the body. ³

- Ulcers may take many weeks (8 weeks average) to heal; and are secondarily infected with bacteria in approximately 50% of cases.
- Permanent disabling scars and crippling may result.
- Each time a worm emerges, the patient may be unable to work and resume daily activities for an average of 3 months.
- Overtly symptomatic infection become apparent during planting or harvesting season, resulting in heavy crop losses.
- Dermal onchocerciasis may mimic dracunculiasis. ⁴
- Calcified worms representing prior infection are occasionally noted on x-ray. ⁵⁻⁷


This disease is endemic or potentially endemic to 16 countries.

Dracunculiasis in Uganda

Time and Place:

Dracunculiasis was first reported in Uganda in 1934, in West Nile District. ⁸

- In recent years, 94.9% of cases have been reported from the northeast (Kitgum, Moroto and Kotido Districts). ^{9 10}
- Peak transmission occurs during April to July.



Graph: Uganda. Dracunculiasis, cases

Notes:

Individual years:


1992 - Uganda reported the second highest incidence, after Nigeria. ¹¹1994 - 98% of these from the 5 northernmost districts: Arus, Gulu, Kitgum, Kotido and Moroto. ¹²1997 - Over 90% from Kotido and Moroto Districts. ¹³1998 - Included 162 cases imported from Sudan. ¹⁴

1999 - 58% from Kitido district, and including 5 imported cases.

2000 - From the northeast (36 cases in Kitgum district, 29 in Kotido district and 27 in Moroto district), and including 4 imported cases. ¹⁵2001 - 84% from two villages in Kotido district), and including 4 imported cases. ¹⁶

2002 - Included 18 imported cases.

2003 - All cases reported from Nawopoet village, Kotido district), and including 13 imported cases. ¹⁷2004 - All imported. ¹⁸2005 - All imported. ¹⁹2006 - Both cases imported from Sudan. ²⁰2007 - All imported from Sudan. ²¹



Graph: Uganda. Dracunculiasis, endemic villages

Notes:

1. 2,677 villages were found to be infected during 1991 to 1992, with a point prevalence of 52,816 cases.

References

1. Bull World Health Organ 1979 ;57(5):683-9.
2. J Trop Med Hyg 1991 Feb ;94(1):35-41.
3. J Radiol Case Rep 2012 Nov ;6(11):29-33.
4. Am J Trop Med Hyg 2010 Dec ;83(6):1348-51.
5. J Cutan Aesthet Surg 2012 Oct ;5(4):281-3.
6. J Radiol Case Rep 2012 Nov ;6(11):29-33.
7. J Assoc Physicians India 2006 Mar ;54:201.
8. Am J Trop Med Hyg 2006 Jul ;75(1):3-8.
9. Int J Epidemiol 1988 Jun ;17(2):434-40.
10. East Afr Med J 1987 Feb ;64(2):102-7.
11. Wkly Epidemiol Rec 1993 Apr 30;68(18):125-31.
12. Wkly Epidemiol Rec 1995 May 5;70(18):125-31.
13. Wkly Epidemiol Rec 1998 May 1;73(18):129-35.
14. Wkly Epidemiol Rec 1999 May 14;74(19):146-52.
15. Wkly Epidemiol Rec 2001 May 4;76(18):133-9.
16. Wkly Epidemiol Rec 2002 Jun 28;77(26):216-20.
17. MMWR Morb Mortal Wkly Rep 2003 Sep 19;52(37):881-3.
18. Wkly Epidemiol Rec 2005 May 13;80(19):165-76.
19. Wkly Epidemiol Rec 2006 May 5;81(18):173-82.
20. Wkly Epidemiol Rec 2007 Apr 20;82(16):133-40.
21. Wkly Epidemiol Rec 2008 May 2;83(18):159-67.

Ebola

Agent	VIRUS - RNA. Mononegavirales, Filoviridae, Filovirus: Ebola virus
Reservoir	? Primate ? Guinea pig ? Bat
Vector	None
Vehicle	Infected secretions Contact Needle Syringe
Incubation Period	5d - 12d (range 2d - 21d)
Diagnostic Tests	Viral culture (blood, liver, spleen). Serology. Nucleic acid amplification. Biosafety level 4.
Typical Adult Therapy	Strict isolation; supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, myalgia, arthralgia, sore throat, vomiting, diarrhea, conjunctivitis, hepatic dysfunction and maculopapular rash; hemorrhagic diatheses from 3rd to 4th day of illness; case-fatality rate 50% to 90%.
Synonyms	Lloviu virus, Tai Forest. ICD9: 065.8 ICD10: A98.4

Clinical

WHO Case definition for surveillance: ¹

Clinical description

- Ebola hemorrhagic fever begins with acute fever, diarrhea that can be bloody (referred to as "diarrhee rouge" in francophone Africa), and vomiting.
- Headache, nausea, and abdominal pain are common.
- Conjunctival injection, dysphagia, and hemorrhagic symptoms such as nosebleeds, bleeding gums, vomiting of blood, blood in stools, purpura may further develop.
- Some patients may also show a maculopapular rash on the trunk.
- Dehydration and significant wasting occur as the disease progresses.
- At a later stage, there is frequent involvement of the central nervous system, manifested by somnolence, delirium, or coma.
- The case-fatality rate ranges from 50% to 90%.

Laboratory criteria for diagnosis

- Supportive: Positive serology (ELISA for IgG and/or IgM), or
- Confirmatory: Positive virus isolation (only in a laboratory of biosafety level 4) or
- Positive skin biopsy (immunohistochemistry) or
- Positive PCR

Case classification

Suspected: A case that is compatible with the clinical description.

Probable: in epidemic situation:

- Any person having had contact with a clinical case and presenting with acute fever, or
- Any person presenting with acute fever and 3 of the following symptoms: headache, vomiting / nausea, loss of appetite, diarrhea, intense fatigue, abdominal pain, general or articular pain, difficulty in swallowing, difficulty in breathing, hiccoughs, or
- Any unexplained death

Confirmed: Any suspected or probable case that is laboratory-confirmed.

Contact: in epidemic situation:

- An asymptomatic person having had physical contact within the past 21 days with a confirmed or probable case or his/her body fluids (e.g., care for patient, participation in burial ceremony, handling of potentially infected laboratory specimens).
- In epidemic situations and after laboratory confirmation of a few initial cases, there is no need for individual laboratory confirmation and the use of .suspected or probable. case classifications is sufficient for surveillance and control purposes.

The symptoms and signs of Marburg and Ebola virus infections are similar. ²

As in smallpox ³, transmission of Ebola ⁴ has occurred during funerals.

Following an incubation period of 4 to 16 days, onset is sudden, marked by anorexia, fever, chills, headache and myalgia. ⁵

- Later, the patient develops nausea, vomiting, sore throat, abdominal pain and diarrhea. ⁶

- Patients are dehydrated, apathetic and disoriented and exhibit pharyngeal and conjunctival injection.
- A non-specific rash may be visible. ⁷
- Most develop severe hemorrhagic manifestations between days 5 and 7.
- Bleeding is often from multiple sites, most commonly from the gastrointestinal tract, lungs and gingiva. ^{8 9}
- Clinical features of infection by Bundibugyo ebolavirus include non-bloody diarrhea (81%), severe headache (81%), and asthenia (77%). ¹⁰
- Hemorrhage and oropharyngeal lesions carry a particularly poor prognosis.
- Death occurs between days 7 and 16.
- Infection by Ebola virus without development of symptoms has been reported. ¹¹

This disease is endemic or potentially endemic to 14 countries.

Ebola in Uganda

Seroprevalence surveys:

1.5% of persons in West Nile Province (1990 to 1992)

Notable outbreaks:

2000 to 2001 - An outbreak (425 cases, 224 fatal) of Ebola was reported, including 393 cases (203 fatal) in Gulu town (northern Uganda); 27 cases (17 fatal) in Masindi; 5 cases (4 fatal) in Mbarara district. ¹²⁻²⁸ Cases were reported in Kitgum, Rwot-Obillo in Aswa county, northern Gulu, Kasubi and Kabedo Opong in Gulu municipality. 16 health workers died during the outbreak. One additional case (fatal) was confirmed in Gulu in January 2001.

2007 to 2008 - An outbreak (116 cases, 39 fatal) was reported in Bundibugyo District, western Uganda. ²⁹⁻⁴⁴ The infecting strain was identified as Bundibugyo Ebolavirus. ⁴⁵⁻⁴⁷

2009 - A case of suspected Ebola was reported in Mbale. ⁴⁸

2011 - An outbreak (1 case, fatal) of Sudan virus infection was reported in Luwero District. ⁴⁹⁻⁶⁰

2012 - An outbreak (24 probable and confirmed, 17 fatal) of Sudan virus infection was reported in Kibaale. ⁶¹⁻⁸⁶ A second outbreak (7 cases, 4 fatal) was subsequently reported in Luwero district. ⁸⁷⁻⁹²

References

1. New Microbiol 2009 Oct ;32(4):359-67.
2. Curr Opin Infect Dis 2001 Aug ;14(4):467-80.
3. Am J Epidemiol 1971 Oct ;94(4):341-7.
4. Bull Soc Pathol Exot 2005 Sep ;98(3):218-23.
5. J Infect 2000 Jan ;40(1):16-20.
6. J Infect Dis 1999 Feb ;179 Suppl 1:ix-xvi.
7. Int J Dermatol 2012 Sep ;51(9):1037-43.
8. Lancet Infect Dis 2004 Aug ;4(8):487-98.
9. Rev Infect Dis 1989 May-Jun;11 Suppl 4:S790-3.
10. PLoS One 2012 ;7(12):e52986.
11. Lancet 2000 Jun 24;355(9222):2210-5.
12. MMWR Morb Mortal Wkly Rep 2001 Feb 9;50(5):73-7.
13. Int J Infect Dis 2004 Jan ;8(1):27-37.
14. US News World Rep 2000 Oct 30;129(17):39.
15. Wkly Epidemiol Rec 2000 Oct 20;75(42):337-8.
16. Bull World Health Organ 2000 ;78(12):1476-7.
17. BMJ 2000 Oct 28;321(7268):1037.
18. Afr Health Sci 2001 Dec ;1(2):60-5.
19. Can Commun Dis Rep 2001 Mar 15;27(6):49-53.
20. JAMA 2001 Feb 28;285(8):1010-2.
21. Wkly Epidemiol Rec 2001 Feb 9;76(6):41-6.
22. Trop Med Int Health 2002 Dec ;7(12):1068-75.
23. Trop Doct 2002 Jan ;32(1):10-5.
24. Emerg Infect Dis 2003 Nov ;9(11):1430-7.
25. Emerg Infect Dis 2003 Oct ;9(10):1242-8.
26. Virus Res 2005 Oct ;113(1):16-25.
27. BMC Infect Dis 2011 ;11:357.
28. Global Health 2012 ;8:15.
29. Wkly Epidemiol Rec 2008 Mar 7;83(10):89-90.
30. PLoS Pathog 2008 Nov ;4(11):e1000212.
31. Emerg Infect Dis 2010 Dec ;16(12):1969-72.
32. J Infect Dis 2011 Nov ;204 Suppl 3:S761-7.
33. PLoS One 2012 ;7(12):e52986.
34. ProMED <promedmail.org> archive: 20071130.3869
35. ProMED <promedmail.org> archive: 20071202.3885
36. ProMED <promedmail.org> archive: 20071203.3903
37. ProMED <promedmail.org> archive: 20071204.3917
38. ProMED <promedmail.org> archive: 20071206.3933
39. ProMED <promedmail.org> archive: 20071208.3965
40. ProMED <promedmail.org> archive: 20071212.3996
41. ProMED <promedmail.org> archive: 20071215.4031
42. ProMED <promedmail.org> archive: 20080104.0050
43. ProMED <promedmail.org> archive: 20080221.0704
44. ProMED <promedmail.org> archive: 20081121.3675
45. Emerg Infect Dis 2010 Jul ;16(7):1087-92.
46. Virology 2012 Feb 20;423(2):119-24.
47. ProMED <promedmail.org> archive: 20101030.3934
48. ProMED <promedmail.org> archive: 20090821.2962
49. Wkly Epidemiol Rec 2011 May 27;86(22):221.
50. Pan Afr Med J 2011 ;9:43.
51. Emerg Infect Dis 2012 Sep ;18(9):1480-3.
52. ProMED <promedmail.org> archive: 20110514.1476
53. ProMED <promedmail.org> archive: 20110516.1489
54. ProMED <promedmail.org> archive: 20110519.1511
55. ProMED <promedmail.org> archive: 20110520.1532
56. ProMED <promedmail.org> archive: 20110525.1593
57. ProMED <promedmail.org> archive: 20110526.1609
58. ProMED <promedmail.org> archive: 20110526.1613
59. ProMED <promedmail.org> archive: 20110604.1711
60. ProMED <promedmail.org> archive: 20110617.1856
61. Pediatr Infect Dis J 2012 Sep ;31(9):998, 1002.
62. Wkly Epidemiol Rec 2012 Sep 7;87(36):339.
63. Wkly Epidemiol Rec 2012 Dec 7;87(49/50):493.
64. ProMED <promedmail.org> archive: 20120728.1218683
65. ProMED <promedmail.org> archive: 20120728.1218652
66. ProMED <promedmail.org> archive: 20120725.1214822
67. ProMED <promedmail.org> archive: 20120729.1219160
68. ProMED <promedmail.org> archive: 20120730.1220418
69. ProMED <promedmail.org> archive: 20120730.1221657
70. ProMED <promedmail.org> archive: 20120731.1223266
71. ProMED <promedmail.org> archive: 20120801.1224404
72. ProMED <promedmail.org> archive: 20120802.1226036
73. ProMED <promedmail.org> archive: 20120802.1226459
74. ProMED <promedmail.org> archive: 20120803.1227357
75. ProMED <promedmail.org> archive: 20120804.1228594
76. ProMED <promedmail.org> archive: 20120804.1228678
77. ProMED <promedmail.org> archive: 20120805.1229338
78. ProMED <promedmail.org> archive: 20120806.1230691
79. ProMED <promedmail.org> archive: 20120807.1233213
80. ProMED <promedmail.org> archive: 20120810.1238765
81. ProMED <promedmail.org> archive: 20120813.1241762
82. ProMED <promedmail.org> archive: 20120815.1245722
83. ProMED <promedmail.org> archive: 20120817.1249794
84. ProMED <promedmail.org> archive: 20120825.1263284
85. ProMED <promedmail.org> archive: 20120903.1278832
86. ProMED <promedmail.org> archive: 20121005.1326565
87. ProMED <promedmail.org> archive: 20121114.1409832
88. ProMED <promedmail.org> archive: 20121116.1411532
89. ProMED <promedmail.org> archive: 20121118.1414575
90. ProMED <promedmail.org> archive: 20121130.1432644
91. ProMED <promedmail.org> archive: 20130110.1491237
92. ProMED <promedmail.org> archive: 20130117.1502950

Echinococcosis - unilocular

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Taeniidae: <i>Echinococcus granulosus</i> , <i>Echinococcus canadensis</i>
Reservoir	Dog Wolf Dingo Sheep Horse Pig
Vector	None
Vehicle	Soil Dog Feces Fly
Incubation Period	1y - 20y
Diagnostic Tests	Serology. Identification of parasite in surgical specimens.
Typical Adult Therapy	Albendazole 400 mg BID X 28d. Repeat X 3, with 2 week hiatus between cycles. Praziquantel has been used preoperatively to sterilize cyst. Follow by surgery as indicated. PAIR (puncture-aspiration-injection-reaspiration) is also used
Typical Pediatric Therapy	Albendazole 10 mg/kg/day X 28d. Repeat X 3, with 2 week hiatus between cycles. Praziquantel has been used preoperatively to sterilize cyst. Follow by surgery as indicated. PAIR (puncture-aspiration-injection-reaspiration) also used
Clinical Hints	Calcified hepatic cyst or mass lesions in lungs and other organs; brain and lung involvement are common in pediatric cases.
Synonyms	<i>Echinococcus canadensis</i> , <i>Echinococcus granulosus</i> , <i>Echinococcus ortleppi</i> , Hydatid cyst, Unilocular echinococcosis. ICD9: 122.0,122.1,122.2,122.3,122.4 ICD10: B67.0,B67.1,B67.2,B67.3,B67.4

Clinical

Symptoms are often absent, even when large cysts are present; and cysts are often discovered incidentally on a routine x-ray or ultrasound study. ¹

Hepatic echinococcosis often presents as abdominal pain with or without a palpable mass in the right upper quadrant. ²

- Biliary compression or rupture of the cysts into a bile duct may mimic cholecystitis or cholelithiasis.
- Ductal compression may also result in pancreatitis. ³
- Leakage from a cyst may produce fever, pruritis, urticaria, eosinophilia or even anaphylactic shock. ⁴

Extra-hepatic echinococcosis presents as space-occupying lesions of brain ⁵, lung ⁶ and thorax ⁷, bone ⁸⁻¹⁸, muscles ¹⁹⁻²⁷, joints ²⁸, parapharyngeal spaces ²⁹ or paranasal sinuses ³⁰, heart ³¹⁻³⁶ and heart valves ^{37 38}, pericardium ³⁹ ⁴⁰, breast ⁴¹⁻⁴³, subcutaneous tissue ⁴⁴⁻⁴⁶, abdominal wall ^{47 48}, supraclavicular region ⁴⁹, peripheral nerves ⁵⁰, ovary, thyroid ⁵¹⁻⁵³, orbits ⁵⁴⁻⁵⁹, parotid gland ⁶⁰, spleen ⁶¹⁻⁶⁶, adrenal ⁶⁷, kidneys ⁶⁸⁻⁷², urinary bladder ⁷³⁻⁷⁵, peritoneum / mesentery / omentum ⁷⁶⁻⁷⁸ or virtually any other organ. ⁷⁹⁻⁸⁵

• In contrast to hepatic echinococcosis, extrahepatic cysts are often non-calcified and may at times be mistaken for malignancy. ^{86 87}

- The brain is involved in 1 to 2% of all *Echinococcus granulosus* infections. ⁸⁸
- The spleen is involved in 0.5% to 6.0% of abdominal infections. ⁸⁹
- The clinical features of cerebral coenurosis may mimic those of echinococcosis. ⁹⁰
- Primary spinal hydatidosis occurs in 1% of cases and may be confused with space-occupying non-infectious disorders ⁹¹⁻⁹⁵

Pulmonary cysts ⁹⁶ may rupture into the bronchial tree and produce cough, hemoptysis and chest pain. ⁹⁷

- Rupture of cysts may disseminate protoscolices to contiguous organs or into the vascular system, resulting in the formation of additional cysts.
- Late intrathoracic complications include intrapulmonary or pleural rupture, infection of the ruptured cysts, reactions of the adjacent tissues, thoracic wall invasion and iatrogenic involvement of pleura. ⁹⁸
- Rupture can occur spontaneously or as a result of trauma or surgery. ⁹⁹
- Anaphylaxis may follow cyst rupture ^{100 101}, but has also reported in patients with intact cysts. ¹⁰² In rare cases, anaphylactic shock (eg, following blunt trauma) may be the initial presenting feature of echinococcosis. ¹⁰³
- Secondary colonization of hydatid cysts by *Aspergillus* has been reported. ¹⁰⁴

Primary superinfection of cysts by bacteria or fungi occurs in approximately 7.3% of cases. ¹⁰⁵

This disease is endemic or potentially endemic to 155 countries.

Echinococcosis - unilocular in Uganda

Prevalence surveys:

66.3% of dogs in Moroto District (2007 to 2008) ¹⁰⁶

Echinococcus granulosus infection has been identified in warthogs in Queen Elizabeth National Park. ¹⁰⁷

References

1. J Gastroenterol Hepatol 2005 Mar ;20(3):352-9.
2. Scand J Gastroenterol Suppl 2004 ;(241):50-5.
3. Cases J 2009 ;2:7374.
4. Surg Today 2004 ;34(12):987-96.
5. J Child Neurol 2008 May ;23(5):585-8.
6. Curr Opin Pulm Med 2010 May ;16(3):257-61.
7. Pan Afr Med J 2012 ;13:7.
8. Trans R Soc Trop Med Hyg 2008 Mar ;102(3):233-8.
9. Ann Trop Med Parasitol 2007 Sep ;101(6):551-3.
10. Postgrad Med J 2007 Aug ;83(982):536-42.
11. JNMA J Nepal Med Assoc 2008 Jul-Sep;47(171):139-41.
12. Thorac Cardiovasc Surg 2007 Dec ;55(8):525-7.
13. Orthopedics 2008 Jul ;31(7):712.
14. J Pediatr Surg 2010 Nov ;45(11):2247-9.
15. Rev Mal Respir 2011 Mar ;28(3):306-11.
16. J Hand Surg Am 2012 Apr 4;
17. Rev Fac Cien Med Univ Nac Cordoba 2012 Mar ;69(1):51-5.
18. Int J Prev Med 2012 Sep ;3(9):660-3.
19. Clin Invest Med 2008 ;31(5):E296-9.
20. Kathmandu Univ Med J (KUMJ) 2008 Oct-Dec;6(24):511-3.
21. Arch Pediatr 2010 Mar ;17(3):263-5.
22. Srp Arh Celok Lek 2010 Jul-Aug;138(7-8):502-5.
23. BMC Infect Dis 2011 ;11:103.
24. Surg Infect (Larchmt) 2011 Oct ;12(5):401-3.
25. Neurol Neurochir Pol 2011 Jul-Aug;45(4):387-90.
26. Diagn Cytopathol 2012 Sep 25;
27. Ekleml Hastalik Cerrahisi 2012 Dec ;23(3):173-6.
28. Am J Trop Med Hyg 2009 Sep ;81(3):371-2.
29. Dysphagia 2011 Mar ;26(1):75-7.
30. Gen Dent 2008 Jul-Aug;56(5):444-6.
31. Acta Med Okayama 2008 Oct ;62(5):341-4.
32. Gen Thorac Cardiovasc Surg 2010 May ;58(5):248-50.
33. Bull Soc Pathol Exot 2010 Dec ;103(5):305-8.
34. J Cardiothorac Surg 2010 ;5:124.
35. Tex Heart Inst J 2011 ;38(6):719-22.
36. Case Report Med 2012 ;2012:603087.
37. Eur J Echocardiogr 2008 Mar ;9(2):342-3.
38. J Clin Ultrasound 2011 Sep ;39(7):431-3.
39. J Emerg Med 2010 Jun ;38(5):582-6.
40. Intern Med 2012 ;51(4):391-3.
41. Acta Chir Belg 2007 Sep-Oct;107(5):570-1.
42. J Pak Med Assoc 2010 Mar ;60(3):232-4.
43. Singapore Med J 2010 Apr ;51(4):e72-5.
44. Parasitol Int 2008 Jun ;57(2):236-8.
45. Rev Stomatol Chir Maxillofac 2012 Jun 13;
46. J Med Case Rep 2012 Nov 26;6(1):404.
47. J Med Case Rep 2011 ;5:270.
48. Chirurgia (Bucur) 2012 Sep-Oct;107(5):655-8.
49. Case Report Med 2012 ;2012:484638.
50. J Neurosurg Spine 2008 Apr ;8(4):394-7.
51. J Infect Dev Ctries 2009 ;3(9):732-4.
52. Acta Med Iran 2011 ;49(4):262-4.
53. Rev Stomatol Chir Maxillofac 2012 Apr ;113(2):124-6.
54. J Clin Neurosci 2012 Feb 11;
55. Childs Nerv Syst 2011 May ;27(5):693-5.
56. Orbit 2010 Feb ;29(1):51-6.
57. J Craniomaxillofac Surg 2010 Jun ;38(4):274-8.
58. Minim Invasive Neurosurg 2007 Dec ;50(6):367-9.
59. Int Ophthalmol 2004 Jul ;25(4):193-200.
60. Rev Stomatol Chir Maxillofac 2011 Jun ;112(3):190-2.
61. J Coll Physicians Surg Pak 2009 Jun ;19(6):380-2.
62. Trop Doct 2009 Oct ;39(4):248-9.
63. Klin Mikrobiol Infekc Lek 2009 Oct ;15(5):185-8.
64. Rev Gastroenterol Peru 2010 Jul-Sep;30(3):224-7.
65. Korean J Parasitol 2012 Jun ;50(2):147-50.
66. Vector Borne Zoonotic Dis 2013 Feb 12;
67. Saudi Med J 2008 Jul ;29(7):1004-8.
68. Urology 2009 May ;73(5):999-1001.
69. Can Urol Assoc J 2011 May 1 ;1:1-6.
70. Cell Biol Int Rep 1990 Feb ;14(2):179.
71. PLoS One 2012 ;7(11):e47667.
72. Urologia 2012 ;79 Suppl 19:e67-71.
73. J Postgrad Med 2008 Oct-Dec;54(4):313-5.
74. Arch Gynecol Obstet 2010 Jul ;282(1):29-32.
75. Can Urol Assoc J 2012 Oct ;6(5):E192-4.
76. World J Emerg Surg 2009 ;4:13.
77. BMJ Case Rep 2012 ;2012:
78. Case Rep Surg 2012 ;2012:654282.
79. Tunis Med 2009 Feb ;87(2):123-6.
80. Int Urogynecol J 2010 Dec ;21(12):1577-9.
81. J Clin Med Res 2011 Feb 12;3(1):52-4.
82. Auris Nasus Larynx 2012 Oct ;39(5):537-9.
83. J Infect Dev Ctries 2011 Nov ;5(11):825-7.
84. Bull Soc Pathol Exot 2012 Oct ;105(4):256-8.
85. Case Report Med 2012 ;2012:362610.
86. Med Oncol 2009 Dec ;26(4):424-8.
87. J Chin Med Assoc 2011 May ;74(5):237-9.
88. J Clin Neurosci 2007 Apr ;14(4):394-6.
89. Korean J Parasitol 2012 Jun ;50(2):147-50.
90. Clin Neuropathol 2011 Jan-Feb;30(1):28-32.
91. J Spinal Cord Med 2008 ;31(1):106-8.
92. Eur Spine J 2009 Jul ;18 Suppl 2:179-82.
93. Spine (Phila Pa 1976) 2010 Apr 20;35(9):E356-8.
94. Eur Spine J 2010 Sep ;19(9):1415-22.
95. Infez Med 2011 Mar ;19(1):39-41.
96. Curr Opin Pulm Med 2010 May ;16(3):257-61.
97. Ann Thorac Surg 2004 Apr ;77(4):1200-4.
98. Eur J Radiol 2009 Apr ;70(1):49-56.
99. Saudi Med J 2010 Jan ;31(1):37-42.
100. Respir Care 2011 Jun ;56(6):863-5.
101. Am J Trop Med Hyg 2011 Sep ;85(3):452-5.
102. J Gastrointest Surg 2008 Dec ;12(12):2243-5.
103. Acta Gastroenterol Belg 2011 Sep ;74(3):462-4.
104. Clin Med Insights Case Rep 2011 ;4:63-8.
105. Am J Trop Med Hyg 2010 Mar ;82(3):376-8.
106. Trop Anim Health Prod 2010 Oct ;42(7):1451-7.
107. Int J Parasitol 2009 Sep ;39(11):1269-76.

Endocarditis - infectious

Agent	BACTERIUM OR FUNGUS. viridans streptococci, <i>Staphylococcus aureus</i> , enterococci, <i>Candida albicans</i> , et al.
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Blood culture, clinical findings, ultrasonography of heart valves.
Typical Adult Therapy	Bactericidal antibiotic appropriate to species
Typical Pediatric Therapy	As for adult
Clinical Hints	Consider in any patient with fever, multisystem disease (i.e., skin lesions, hematuria, neurological symptoms, single or multiple abscesses or bone, brain, lung, etc) and a preexisting cardiac valvular lesion.
Synonyms	Bacterial endocarditis, Endocardite, Endocarditis, Endokarditis, Fungal endocarditis, Infectious endocarditis, S.B.E.. ICD9: 421 ICD10: I33

Clinical

The definitive diagnosis of infective endocarditis requires: [1](#) [2](#)

- 1) Demonstration of microorganisms; and/or histological lesions in the heart or heart valves; or
- 2) Presence of two major criteria; or 1 major and 3 minor criteria; or 5 minor criteria, as follows:

Major Criteria:

A. Culture:

- 1. Typical microorganisms (*HACEK*, *Streptococcus viridans*, *Streptococcus bovis*) in 2 separate blood cultures; or community acquired *Staphylococcus aureus* or enterococcus without obvious focus.
- 2. Persistently positive blood cultures (drawn more than 12 hours apart; or three positive cultures at least one hour apart).

B. Evidence of endocardial or valvular involvement (echocardiogram, abscess, new valvular regurgitant lesion)

Minor Criteria:

A. Predisposition (heart condition, drug abuse)

B. Fever

C. Embolic phenomena, mycotic aneurysm, Janeway lesion, or intracranial hemorrhage.

D. Immunological phenomena (Osler nodes, positive rheumatoid factor)

E. Echocardiogram with suggestive, but not specific findings.

F. Positive blood culture, but not meeting Major criteria.

Etiological associations:

- Injecting drug user: *Staphylococcus aureus*, enterococci, Enterobacteriaceae, *Pseudomonas aeruginosa*, *Candida*
- Prosthetic valve: *Staphylococcus epidermidis* Enterobacteriaceae, *Candida*, *Aspergillus*
- Rheumatic or other valvular disease: viridans Streptococci, enterococci
- "Culture negative" endocarditis: *Coxiella burnetii*, *Bartonella* spp., *Tropheryma whipplei*, et al.

This disease is endemic or potentially endemic to all countries.

References

1. *Am J Med* 1994 Mar ;96(3):200-9.
2. *Clin Infect Dis* 2000 Apr ;30(4):633-8.

Enteritis necroticans

Agent	BACTERIUM. Clostridium perfringens An anaerobic gram positive bacillus
Reservoir	Soil Feces Mammal
Vector	None
Vehicle	Food
Incubation Period	1d - 4d
Diagnostic Tests	Recovery of type C Clostridium perfringens from stool in association with suggestive findings.
Typical Adult Therapy	Supportive, with surgery as indicated. Oral Vancomycin not proven but may be helpful (125 mg QID X 10d)
Typical Pediatric Therapy	Supportive, with surgery as indicated. Oral Vancomycin not proven but may be helpful (2 mg/kg QID X 10d)
Clinical Hints	Bloody diarrhea, fever, vomiting and abdominal distention - may progress to ileus or bowel perforation and shock; onset one day after ingestion of pork.
Synonyms	Darmbrand, Necrotizing jejunitis, Pigbel, Segmental necrotizing enteritis. ICD9: 569.89,008,49 ICD10: A05.2

Clinical

This condition is characterized by abdominal distention, pain and hypotension.

- Dilated, thickened loops of bowel exhibiting segmental necrosis are found at laparotomy. ¹
- Published case-fatality rates vary from 12% (treated) to 58%.

This disease is endemic or potentially endemic to 13 countries.

References

1. [P N G Med J 1979 Mar ;22\(1\):18-23.](#)

Enterobiasis

Agent	PARASITE - Nematoda. Phasmidea: Enterobius vermicularis
Reservoir	Human
Vector	None
Vehicle	Fecal-oral Air Clothing Sexual contact (rare)
Incubation Period	14d - 42d
Diagnostic Tests	Apply scotch tape to anal verge in a.m. & paste onto glass slide for microscopy.
Typical Adult Therapy	Albendazole 400 mg PO as single dose - repeat in 2w. OR Mebendazole 100 mg PO as single dose - repeat in 2w. OR Pyrantel pamoate 11 mg/kg (max 1g) PO as single dose; or
Typical Pediatric Therapy	Mebendazole 100 mg PO as single dose (>age 2) - repeat in 2w. OR Pyrantel pamoate 11 mg/kg (max 1g) PO X 1
Clinical Hints	Nocturnal anal pruritus; occasionally vaginitis or abdominal pain; eosinophilia is rarely, if ever, encountered.
Synonyms	Enterobio, Enterobius vermicularis, Oxyuriasis, Oxyuris, Pinworm, Seatworm. ICD9: 127.4 ICD10: B80

Clinical

The typical manifestation of enterobiasis is nocturnal pruritus and related to hypersensitivity to worm antigens.

- Local dermal "tingling" is also encountered. ¹
- Migration of adult females to the vulva may result in vaginal pain ² and vulvovaginitis ³, or predispose to urinary tract infection.
- Eosinophilia is occasionally present.

Complications are rare, and include salpingitis ⁴, cystitis ⁵, peritonitis ^{6 7}, hepatitis granuloma ⁸ and urethritis. ⁹

- Although abdominal symptoms may mimic those of appendicitis, *Enterobius* is at least as common in normal as in inflamed appendices. ¹⁰⁻¹³
- Cases of *Enterobius* prostatitis ¹⁴ and peritonitis have been reported. ¹⁵
- Adults and ova of *Enterobius* have been identified in the kidneys ^{16 17} and eyes ¹⁸ of infested patients.

The presence of diarrhea or abdominal pain suggests coinfection with *Dientamoeba fragilis*.

This disease is endemic or potentially endemic to all countries.

Enterobiasis in Uganda

Prevalence surveys:

0.05% of pregnant women in Entebbe (2004 to 2005) ¹⁹

References

1. Gastroenterol Clin North Am 1996 Sep ;25(3):579-97.
2. J Paediatr Child Health 2013 Jan 18;
3. Int J Immunopathol Pharmacol 2008 Oct-Dec;21(4):1031-3.
4. Pathol Res Pract 2010 Jun 15;206(6):405-7.
5. Braz J Infect Dis 2008 Aug ;12(4):352.
6. Surg Infect (Larchmt) 2009 Dec ;10(6):545-7.
7. Clin Exp Obstet Gynecol 2012 ;39(3):379-81.
8. Int J Surg Case Rep 2012 ;3(1):6-9.
9. J Infect 1992 Jan ;24(1):87-90.
10. Trop Doct 2006 Jul ;36(3):160-2.
11. Cases J 2008 ;1(1):376.
12. Scand J Gastroenterol 2009 ;44(4):457-61.
13. Pediatr Surg Int 2004 May ;20(5):372-5.
14. J Med Case Rep 2007 ;1:137.
15. Eur J Gynaecol Oncol 2007 ;28(6):513-5.
16. J Med Microbiol 2010 Jul ;59(Pt 7):860-1.
17. Scand J Urol Nephrol 2012 Feb ;46(1):70-2.
18. J Clin Microbiol 2011 Dec ;49(12):4369-70.
19. PLoS Negl Trop Dis 2009 ;3(6):e473.

Enterovirus infection

Agent	VIRUS - RNA. Picornaviridae: Coxsackievirus, ECHO virus, Enterovirus, Parechovirus
Reservoir	Human
Vector	None
Vehicle	Droplet Fecal-oral
Incubation Period	2d-7d
Diagnostic Tests	Viral culture (stool, pharynx, CSF). Serology. Nucleic acid amplification.
Typical Adult Therapy	Supportive. Pleconaril 200 to 400 mg PO TID X 7d has been used for severe infections
Typical Pediatric Therapy	Supportive. Pleconaril 5 mg/kg PO BID has been used for severe infections
Clinical Hints	Summer-to-autumn sore throat; occasionally chest pain, macular or vesicular rash, meningitis, myopericarditis, etc.
Synonyms	Boston exanthem [Coxsackie. A 16], Coxsackie, Coxsackievirus, ECHO, Echovirus, Enteroviruses, Hand, foot and mouth disease, Hand-foot-and-mouth disease, Herpangina [Coxsackievirus A], HEV 68, HPeVs, Human Enterovirus 68, Human Parechovirus, Ljungan virus, Myocarditis, enteroviral, Parechovirus, Pericarditis, enteroviral. ICD9: 049,079.2,008.67,074.0,074.8,074.3,070.4,078.89 ICD10: A88.0,A87.0,B08.4,B08.5,B08.8,B30.3,B34.1

Clinical

The various enteroviruses are associated with fever and pharyngitis, which may be followed by appearance of: ^{1 2}

- rash
- aseptic meningitis
- encephalitis ³
- acute disseminated encephalomyelitis ⁴
- epidemic conjunctivitis
- herpangina
- hand-foot-and-mouth disease
- myocarditis
- pericarditis
- pleurodynia
- pneumonia
- acute flaccid paralysis ⁵⁻⁷
- conjunctivitis, etc

Hand, foot and mouth disease (HFMD) is characterized by a prodrome of fever and sore throat, followed by the appearance of vesicles on the palmar and plantar regions, and oral mucosa.

- Vesicles in the mouth are often pleomorphic, with rectangular and triangular shapes.
- Hand foot and mouth disease has been associated with onychomadesis • complete nail shedding from the proximal portion, affecting both fingernails and toenails. ⁸⁻¹⁶
- HFMD due to Enterovirus 71 is often complicated by central nervous system disease and sequelae. ¹⁷⁻⁴²

The clinical features of Enterovirus infection among neonates and infants are similar to those of Parechovirus infection. ⁴³

Human Enterovirus 68 infection is associated with respiratory illness ranging from relatively mild illness that did not require hospitalization to severe illness requiring intensive care and mechanical ventilation. Some infections have been fatal. ^{44 45}

Echoviruses 22 and 23 have been reclassified as human parechovirus (HPeV) 1 and 2, respectively. ⁴⁶

- Parechovirus infections have been associated with respiratory and gastrointestinal disease ^{47 48}, epidemic myalgia ⁴⁹ or rarely encephalitis. ⁵⁰
- HPeV2 is usually associated with gastrointestinal illness.
- HPeV3 has been associated with transient paralysis, sepsis-like syndromes, or myalgia with muscle weakness.
- HPeV4 has been associated with fever in a neonate ⁵¹
- HPeV6 (NII561-2000) has been associated with infectious gastroenteritis, fever with rash, upper respiratory infection and

Reye's syndrome

This disease is endemic or potentially endemic to all countries.

Enterovirus infection in Uganda

Notable outbreaks:

2010 - An outbreak (6,818 cases) of acute hemorrhagic conjunctivitis was reported. [52](#) [53](#)

References

1. Dermatol Clin 2002 Apr ;20(2):217-23.
2. Semin Pediatr Infect Dis 2002 Jan ;13(1):40-7.
3. N Engl J Med 1999 Sep 23;341(13):936-42.
4. Neurosciences (Riyadh) 2010 Jan ;15(1):46-8.
5. Rev Infect Dis 1984 May-Jun;6 Suppl 2:S387-90.
6. AJNR Am J Neuroradiol 2001 Jan ;22(1):200-5.
7. Int J Neurosci 2012 Jul ;122(7):338-44.
8. Eur J Pediatr 2001 Nov ;160(11):649-51.
9. Pediatr Dermatol 2000 Jan-Feb;17(1):7-11.
10. Euro Surveill 2008 Jul 3;13(27)
11. Epidemiol Infect 2010 Dec ;138(12):1775-8.
12. Euro Surveill 2010 Sep 16;15(37)
13. Euro Surveill 2010 Sep 16;15(37)
14. ProMED <promedmail.org> archive: 20100916.3356
15. ProMED <promedmail.org> archive: 20100921.3401
16. ProMED <promedmail.org> archive: 20100922.3421
17. Korean J Pediatr 2011 Jan ;54(1):11-6.
18. Clin Infect Dis 2000 Sep ;31(3):678-83.
19. N Engl J Med 2000 Feb 3;342(5):355-6.
20. Wkly Epidemiol Rec 1997 Jul 11;72(28):211-2.
21. Jpn J Infect Dis 1999 Feb ;52(1):12-5.
22. Scand J Infect Dis 1999 ;31(4):331-5.
23. Virus Res 1999 May ;61(1):1-9.
24. Med J Malaysia 2002 Mar ;57(1):88-91.
25. Med J Malaysia 2005 Aug ;60(3):297-304.
26. N Engl J Med 1999 Sep 23;341(13):929-35.
27. Emerg Infect Dis 2003 Mar ;9(3):291-3.
28. Pediatrics 2002 Jun ;109(6):e88.
29. Clin Infect Dis 2002 May 1;34 Suppl 2:S52-7.
30. J Clin Virol 2000 Jun ;17(1):23-30.
31. Emerg Infect Dis 2001 Jan-Feb;7(1):146-8.
32. Scand J Infect Dis 2010 Aug ;42(8):609-12.
33. ProMED <promedmail.org> archive: 20060305.0712
34. ProMED <promedmail.org> archive: 20060313.0792
35. ProMED <promedmail.org> archive: 20060319.0854
36. ProMED <promedmail.org> archive: 20060406.1035
37. ProMED <promedmail.org> archive: 20100329.0985
38. ProMED <promedmail.org> archive: 20100412.1184
39. ProMED <promedmail.org> archive: 20100416.1235
40. ProMED <promedmail.org> archive: 20100507.1489
41. ProMED <promedmail.org> archive: 20100625.2121
42. Zhonghua Yi Xue Za Zhi 2012 Jul 3;92(25):1742-6.
43. Pediatr Infect Dis J 2008 Mar ;27(3):241-5.
44. MMWR Morb Mortal Wkly Rep 2011 Sep 30;60(38):1301-4.
45. ProMED <promedmail.org> archive: 20110929.2945
46. Clin Infect Dis 2006 Jan 15;42(2):204-10.
47. Curr Opin Infect Dis 2010 Jun ;23(3):224-30.
48. J Clin Virol 2009 May ;45(1):1-9.
49. Emerg Infect Dis 2012 Nov ;18(11):1787-93.
50. Brain Dev 2013 Jan 21;
51. Emerg Infect Dis 2006 Oct ;12(10):1572-5.
52. MMWR Morb Mortal Wkly Rep 2010 Aug 20;59(32):1024.
53. ProMED <promedmail.org> archive: 20100819.2890

Entomophthoromycosis

Agent	FUNGUS. Zygomycota, Zygomycetes, Entomophthorales: Basidiobolus or Conidiobolus
Reservoir	Vegetation Soil Amphibian Reptile
Vector	None
Vehicle	Air (inhalation) Direct inoculation
Incubation Period	Unknown
Diagnostic Tests	Biopsy and fungal culture.
Typical Adult Therapy	Antifungal agents and excision as indicated. Oral potassium iodide may be helpful
Typical Pediatric Therapy	As for adult
Clinical Hints	Slowly-spreading subcutaneous nodule involving nose, upper face, pharynx; no skin ulceration or systemic signs.
Synonyms	Basidiobolomycosis, Basidiobolus, Conidiobolosis, Conidiobolus, Rhinoentomophthoromycosis, Rhinophycomycosis entomophthorae, Subcutaneous phycomycosis, Subcutaneous zygomycosis. ICD9: 117.9 ICD10: B48.8

Clinical

Basidiobolomycosis is caused by *Basidiobolus ranarum* (*B. haptosporus*, *B. meristosporus*) and involves the subcutaneous tissues and rarely the intestine. ^{1 2}

- The typical infection is characterized by a single, painless, sharply circumscribed, hard subcutaneous mass • most commonly on the thigh or buttock.
- Infection of the nose and other facial structures is also reported. ³
- The overlying skin remains intact, and bone involvement is not seen.
- Abdominal disease is characterized by abdominal pain and a palpable mass, and may mimic an intestinal malignancy or acute appendicitis. ⁴⁻⁶
- Additional findings may include hepatosplenomegaly, hepatic mass lesion, jaundice, gastrointestinal hemorrhage ⁷ or eosinophilia. ⁸

Conidiobolomycosis begins in the nasal submucosa and spreads to the skin, glabella, cheek, lips, paranasal sinuses and pharynx. ^{9 10}

- Central facial swelling may progress to overt elephantiasis of the face. ¹¹
- Histological examination reveals hyphae surrounded by an eosinophilic halo (Spendore-Hoeppli phenomenon) ¹²
- Rare cases of disseminated conidiobolomycosis have been reported. ¹³

This disease is endemic or potentially endemic to 32 countries.

References

1. Clin Microbiol Rev 2000 Apr ;13(2):236-301.
2. Clin Microbiol Infect 2004 Mar ;10 Suppl 1:31-47.
3. Mycopathologia 2010 Sep ;170(3):165-8.
4. BMC Infect Dis 2006 ;6:140.
5. J Med Microbiol 2011 Sep ;60(Pt 9):1395-402.
6. Clin Infect Dis 2012 Mar 22;
7. J Clin Microbiol 2001 Jun ;39(6):2360-3.
8. J Med Microbiol 2011 Jul ;60(Pt 7):871-80.
9. Clin Microbiol Infect 2005 Mar ;11(3):249-50.
10. Infection 2008 Dec ;36(6):594-6.
11. Am J Dermatopathol 2012 Jul ;34(5):511-22.
12. Clin Dermatol 2012 Jul ;30(4):409-12.
13. J Clin Microbiol 2011 Feb ;49(2):752-6.

Epidural abscess

Agent	BACTERIUM. Staphylococcus aureus , facultative gram negative bacilli, etc
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Imaging (CT scan, MRI). Gram-stain and culture of blood or pus.
Typical Adult Therapy	Intravenous antibiotic(s) appropriate to identified or suspected pathogens. Drainage as indicated
Typical Pediatric Therapy	Intravenous antibiotic(s) appropriate to identified or suspected pathogen. Drainage as indicated
Clinical Hints	Frontal bone abscess; or spinal cord compression with signs of infection - often in setting of injecting drug abuse or preexisting staphylococcal infection.
Synonyms	

Clinical

Intracranial epidural abscesses:

Intracranial epidural abscesses may appear gradually, with initial findings suggestive of the underlying sinusitis or otitis. ¹

- Early findings include local pain followed by generalized headache, often with alteration of mental status.
- Focal neurological signs and focal or generalized seizures appear, which reflect the local anatomy of the lesion, for example:
 - abscess near the petrous bone may involve cranial nerves V and VI, with unilateral facial pain and lateral rectus weakness (Gradenigo's syndrome)
 - an occipital epidural abscess may obstruct the superior sagittal sinus

Eventually, papilledema and other signs of elevated intracranial pressure develop.

- Extension into the subdural space is accompanied by rapid neurological deterioration.

Spinal epidural abscess:

Spinal epidural abscess is more common in men than in women and may occur at any age.

- The presentation may be acute or gradual, over several months. ²
- Most begin with focal vertebral pain, which begins to radiate along the course of involved nerve roots.
- Signs of spinal cord compression (long-tract findings), later progress to paralysis below the level of the lesion.
- Hematogenous infection of the epidural space produces rapid progression with prominent systemic signs, and severe local pain.
- Chronic abscesses may mimic epidural neoplasia, often without systemic signs of infection.
- Cervical abscesses may compromise respiration, and produce rapid evolving flaccid hyporeflexia, suggestive of Guillain-Barre syndrome.
- Epidural abscess has occasionally been reported as a complication of pyomyositis. ³

This disease is endemic or potentially endemic to all countries.

References

1. South Med J 2004 Mar ;97(3):279-82; quiz 283.
2. J Am Acad Orthop Surg 2004 May-Jun;12(3):155-63.
3. J Neurosurg Pediatr 2010 Jul ;6(1):33-7.

Erysipelas or cellulitis

Agent	BACTERIUM. Erysipelas: <i>Streptococcus pyogenes</i> Cellulitis: <i>Staphylococcus aureus</i> , <i>Streptococcus pyogenes</i> , occasionally others
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	1d - 7d
Diagnostic Tests	Clinical diagnosis is usually sufficient. Aspiration of lesion for smear and culture may be helpful in some cases.
Typical Adult Therapy	Antibiotic directed at likely pathogens (Group A <i>Streptococcus</i> and <i>Staphylococcus aureus</i>)
Typical Pediatric Therapy	As for adult
Clinical Hints	Erysipelas is well-circumscribed, tender, edematous (peau d'orange), warm and painful; cellulitis is less painful, flat and without a distinct border.
Synonyms	Cellulite, Cellulitis, Celulite, Celulitis, Erisipela, Erysipelas, St. Anthony's fire (erysipelas), St. Francis' fire (erysipelas), Zellulitis. ICD9: 035,681,682 ICD10: A46,L03

Clinical

Erysipelas:

Erysipelas is characterized by abrupt onset of "fiery-red" superficial swelling of the face or extremities. ¹

- The lesion is typically recognized by the presence of well-defined indurated margins, particularly along the nasolabial fold; rapid progression; and intense pain. ²
- Flaccid bullae may develop on the second or third day of illness; but extension to deeper soft tissues is rare.
- Desquamation occurs between the fifth and tenth days of illness.

Cellulitis:

Cellulitis is characterized by local pain, erythema, swelling, and heat. ^{3 4}

- Cellulitis may be caused by any of a wide variety of bacteria or yeasts; however, *S. aureus* or *S. pyogenes* are most often implicated.
- A history of preceding trauma, insect bite, needle insertion or surgery is often present.
- Cultures of biopsy specimens or aspirates are positive in only 20% of cases.
- Infection by *S. aureus* often spreads out from a localized infection (abscess, folliculitis) or foreign body
- Streptococcal cellulitis tends to be more diffuse and rapid in onset, and associated with lymphangitis and fever.
- Streptococci also cause recurrent cellulitis in the setting of lymphedema resulting from elephantiasis or lymph node damage.

Recurrent staphylococcal cutaneous infections are encountered in patients with "Job's syndrome" (eosinophilia and elevated serum levels of IgE); and nasal carriers of staphylococci.

This disease is endemic or potentially endemic to all countries.

References

1. Clin Evid 2003 Dec ;(10):1878-83.
2. Am J Clin Dermatol 2003 ;4(3):157-63.

3. Am Fam Physician 2002 Jul 1;66(1):119-24.
4. Curr Opin Infect Dis 2007 Apr ;20(2):118-23.

Erysipeloid

Agent	BACTERIUM. <i>Erysipelothrix rhusiopathiae</i> A facultative gram-positive bacillus
Reservoir	Mammal Bird Fish
Vector	None
Vehicle	Contact with meat, mammal, poultry or fish
Incubation Period	1d - 4d
Diagnostic Tests	Culture.
Typical Adult Therapy	Penicillin V, Ampicillin, third-generation cephalosporin, Fluoroquinolone (<i>Levofloxacin</i> , <i>Trovafloxacin</i> , <i>Pefloxacin</i> , <i>Sparfloxacin</i> or <i>Moxifloxacin</i>), <i>Erythromycin</i> or <i>Tetracycline</i> generally adequate
Typical Pediatric Therapy	Penicillin V, Ampicillin, third-generation cephalosporin or <i>Erythromycin</i> generally adequate
Clinical Hints	Annular erythema or 'target lesion' on hand following contact with raw animal or fish products; local pain and swelling; no discharge is noted and fever is present in only 10% of cases.
Synonyms	<i>Erysipelothrix rhusiopathiae</i> , Rutlauf. ICD9: 027.1 ICD10: A26

Clinical

Erysipeloid is generally limited to the skin (mainly hands and fingers)

Infection is characterized by pain, edema and purplish erythema with sharp irregular margins which extends peripherally but clears centrally. ^{1 2}

- Relapses and extensions of the lesions to distant areas are common, but there is no fever.
- There is no permanent immunity following an attack.
- Lesions of cutaneous leishmaniasis may mimic those of erysiploid. ³

Complications:

- 31 cases of endocarditis due to *Erysipelothrix rhusiopathiae* had been reported to 1976 ^{4 5} ; and approximately 50 to 1988. ⁶
- Rarely-reported complications have included chronic granulomatous cheilitis ⁷ , peritonitis associated with peritoneal dialysis ⁸ , pneumonia ⁹ and spinal infection with epidural empyema. ¹⁰

This disease is endemic or potentially endemic to all countries.

References

1. Prim Care 2000 Jun ;27(2):459-73.
2. J Med Microbiol 1999 Sep ;48(9):789-99.
3. Bull Soc Pathol Exot 2008 Dec ;101(5):395-7.
4. Pathol Biol (Paris) 1977 May ;25(5):345-52.
5. Dtsch Med Wochenschr 1976 Nov 12;101(46):1672-4.
6. Clin Microbiol Rev 1989 Oct ;2(4):354-9.
7. Ann Dermatol Venereol 2010 Feb ;137(2):124-7.
8. J Korean Med Sci 2010 Aug ;25(8):1234-6.
9. J Med Microbiol 2012 Mar ;61(Pt 3):450-1.
10. Eur Spine J 2012 Apr 17;

Erythrasma

Agent	BACTERIUM. <i>Corynebacterium minutissimum</i> A facultative gram-positive bacillus
Reservoir	Human
Vector	None
Vehicle	Indigenous flora
Incubation Period	Unknown
Diagnostic Tests	Coral fluorescence of skin lesion under Wood's lamp. Culture (alert lab regarding diagnosis).
Typical Adult Therapy	Erythromycin 250 mg PO QID X 14d. Topical Clindamycin 2% and topical Fusidic acid have also been used
Typical Pediatric Therapy	Erythromycin 10 mg/kg PO QID X 14d. Topical Clindamycin 2% and topical Fusidic acid have also been used
Clinical Hints	Pruritic, scaling, slowly-progressive red-brown patch; usually in groin - occasionally in toe webs; common in obese or diabetic males; coral fluorescence with Wood's light.
Synonyms	<i>Corynebacterium minutissimum</i> , Eritrasma. ICD9: 039.0 ICD10: L08.1

Clinical

Erythrasma is characterized by slowly spreading, reddish-brown, pruritic patches • usually in the groin and axillae. ¹

- Other areas include the interdigital regions of the feet ² , the vulva ³ and intergluteal and crural folds.
- Most patients are obese, male diabetics. ⁴⁻⁶
- The lesions fluoresce red when exposed to Wood's lamp. ⁷⁻¹⁰
- The differential diagnosis of erythrasma includes psoriasis, dermatophytosis, candidiasis and intertrigo.

The etiologic agent of erythrasma, *Corynebacterium minutissimum*, has also been associated with bacteremia ¹¹⁻¹⁴ , meningitis ¹⁵ , breast abscess ¹⁶ , eye infection ¹⁷ , endocarditis ^{18 19} , peritonitis ²⁰ , cutaneous granulomas ²¹ , costochondral abscess ²² , puerperal infection ²³ and pyelonephritis. ^{24 25}

This disease is endemic or potentially endemic to all countries.

References

1. N Engl J Med 2004 Oct 14;351(16):1666.
2. J Am Acad Dermatol 1990 Apr ;22(4):578-82.
3. Obstet Gynecol 1993 May ;81(5 (Pt 2)):862-4.
4. Clin Dermatol 2006 Jul-Aug;24(4):237-46.
5. Br J Dermatol 1974 Oct ;91(4):481-4.
6. Arch Dermatol 1969 Jun ;99(6):674-80.
7. AMA Arch Derm Syphilol 1952 May ;65(5):614-5.
8. JAMA 1967 Mar 13;199(11):841.
9. N Engl J Med 2004 Oct 14;351(16):1666.
10. Br J Dermatol 1972 Aug ;87(2):130-7.
11. Clin Infect Dis 2002 Aug 15;35(4):e40-2.
12. Infect Control Hosp Epidemiol 1998 Oct ;19(10):786-9.
13. Clin Infect Dis 1994 Jul ;19(1):204-5.
14. Diagn Microbiol Infect Dis 1986 Nov ;5(4):327-30.
15. J Infect 2008 Jan ;56(1):77-9.
16. J Clin Microbiol 1984 Dec ;20(6):1219-20.
17. Int Ophthalmol 1995-1996;19(5):313-6.
18. J Infect 2007 Feb ;54(2):e79-81.
19. Br J Ophthalmol 1985 Jan ;69(1):29-31.
20. Perit Dial Int 1998 May-Jun;18(3):345-6.
21. J Eur Acad Dermatol Venereol 2002 Nov ;16(6):643-5.
22. J Infect 2000 Jul ;41(1):103-5.
23. Klin Lab Diagn 1995 Jul-Aug;(4):45-8.
24. J Infect 2005 Dec ;51(5):e299-303.
25. Pediatr Infect Dis J 1994 Dec ;13(12):1151-2.

Escherichia coli diarrhea

Agent	BACTERIUM. <i>Escherichia coli</i> A facultative gram-negative bacillus
Reservoir	Human Mammal
Vector	None
Vehicle	Food Water Fecal-oral
Incubation Period	1d - 3d (range 12h - 10d)
Diagnostic Tests	Stool culture. Request characterization of <i>E. coli</i> isolates.
Typical Adult Therapy	Supportive therapy. Avoid anti-motility drugs and antimicrobial agents. Plasma exchange may be effective in HUS Note that antimicrobial agents may increase risk for hemolytic-uremic syndrome when used in cases of <i>E. coli</i> O157:H7 infection
Typical Pediatric Therapy	Supportive therapy. Avoid anti-motility drugs and antimicrobial agents. Plasma exchange may be effective in HUS Note that antimicrobial agents may increase risk for hemolytic-uremic syndrome when used in cases of <i>E. coli</i> O157:H7 infection
Clinical Hints	Watery diarrhea or dysentery - common among travelers and infants; hemorrhagic colitis and hemolytic uremic syndrome are associated with type O157:H7 (& occasionally other types).
Synonyms	DAEC (Diffusely Adherent <i>E. coli</i>), <i>E. coli</i> diarrhea, EAEC (Enteroadherent <i>E. coli</i>), EAggEC (Enteraggregative <i>E. coli</i>), EHEC (Enterohemorrhagic <i>E. coli</i>), EIEC (Enteroinvasive <i>E. coli</i>), EPEC (Enteropathogenic <i>E. coli</i>), <i>Escherichia albertii</i> , ETEC (Enterotoxigenic <i>E. coli</i>), Hemolytic-uremic syndrome, Hemolytic Uremic Syndrome, HUS. ICD9: 008.0 ICD10: A04.0,A04.1,A04.2,A04.3,A04.4

Clinical

Enterotoxigenic *Escherichia coli* (ETEC) infection is characterized by a short incubation period, and watery diarrhea without blood or mucus.

- Fever and vomiting occur in a minority of patients. ¹
- The disease may be life-threatening in infants.

Enteropathogenic *E. coli* (EPEC) causes watery diarrhea with fever and vomiting, primarily among children under age 2 years.

Enteroinvasive *E. coli* (EIEC) causes watery diarrhea; only a minority of patients experience dysentery.

Enterohemorrhagic *E. coli* (EHEC) causes diarrhea without fever, often with blood and cramps at all ages. ²

- Rare instances of toxic megacolon have been reported ³
- One strain of EHEC, O157:H7 is an important cause of hemolytic-uremic syndrome (HUS). ⁴
- Approximately 6% to 10% of patients infected by this strain develop HUS • with an overall mortality rate of 0.6% for STEC O157 infections and 4.6% for HUS. ⁵
- Nearly 40% of patients with STEC-HUS require at least temporary renal replacement therapy and up to 20% will have permanent residual kidney dysfunction. ⁶
- Hemolytic-uremic syndrome can also follow infection by *Clostridium difficile* ⁷ and by non-O157 strains of *E. coli*. ⁸
- Reactive arthritis is reported in 10% of cases ⁹

Enteraggregative *E. coli* (EAggEC) causes watery, persistent diarrhea (over 2 weeks) without vomiting. ¹⁰

- Low-grade fever may be observed, and gross blood may occasionally be present in stools. ¹¹

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 2005 Jul ;18(3):465-83.
2. Curr Opin Infect Dis 2003 Jun ;16(3):259-63.
3. J Infect 2006 Apr ;52(4):e103-6.
4. Curr Gastroenterol Rep 2004 Aug ;6(4):297-301.
5. Clin Infect Dis 2009 Nov 15;49(10):1480-5.
6. Nat Rev Nephrol 2012 Sep 18;8(11):658-69.
7. Clin Nephrol 2013 Jan 15;
8. Emerg Infect Dis 2011 Feb ;17(2):180-5.
9. J Rheumatol 2008 Mar ;35(3):480-7.

10. [Emerg Infect Dis 1998 Apr-Jun;4\(2\):251-61.](#)
11. [Curr Opin Gastroenterol 2005 Jan ;21\(1\):4-8.](#)

Fascioliasis

Agent	PARASITE - Platyhelminthes, Trematoda. Echinostomatida, Fasciolidae: Fasciola hepatica or Fasciola gigantica
Reservoir	Sheep Cattle Snail (Lymnaea, Fossaria)
Vector	None
Vehicle	Food Aquatic plants Watercress (Nasturtium officinale)
Incubation Period	2w - 3m
Diagnostic Tests	Identification of ova in stool or duodenal aspirates (adult parasite in tissue). Serology. PCR. CT scan.
Typical Adult Therapy	Triclabendazole 10 mg/kg PO X 2 doses. OR Bithionol 50 mg/kg every other day X 10 doses OR Nitazoxanide 500 mg PO BID X 7d
Typical Pediatric Therapy	Triclabendazole 10 mg/kg PO X 2 doses. OR Bithionol 50 mg/kg every other day X 10 doses OR Nitazoxanide: Age 1 to 3y 100 mg BID X 7 d Age 4 to 11y 200 mg BID X 7d
Clinical Hints	Fever, hepatomegaly, cholangitis, jaundice and eosinophilia; urticaria occasionally observed during the acute illness; parasite may survive more than 10 years in the biliary tract.
Synonyms	Eurytrema, Fasciola gigantica, Fasciola hepatica, Hepatic distomiasis, Lederegelbefall, Sheep liver fluke. ICD9: 121.3 ICD10: B663.

Clinical

The presence and severity of disease depend on the intensity of infection and the host.

Symptoms may appear a few days after ingestion of larvae, when the immature worms reach the abdominal cavity and begin migrating across or within the liver. ¹

- Typical early symptoms include fever, abdominal pain, gastrointestinal disturbances and urticaria. ²
- Hepatomegaly, anemia and jaundice may also be present.
- Creeping eruption has been reported. ³
- Rare instance of ectopic adult worms are reported ⁴⁻⁸

A latent phase follows during which the only finding is prominent eosinophilia.

- Eventually, the patient enters a chronic phase characterized by biliary colic ⁹, epigastric pain, jaundice, hepatomegaly and abdominal tenderness. ^{10 11}
- Sporadic cases of systemic vasculitis ¹² and pancreatitis are encountered. ¹³⁻¹⁸

This disease is endemic or potentially endemic to 99 countries.

Fascioliasis in Uganda

Prevalence surveys:

- 43.7% of cattle at low altitudes and 1.1% at high altitudes (*Fasciola gigantica*, Mount Eglon, 2012 publication) ¹⁹
- 28.6% of cattle in Tororo and Soroti Districts (2000) ²⁰

Reservoirs:

- Fascioliasis has been identified in Zebu cattle (1960 publication) ²¹
- *Lymnaea natalensis* is prevalent at low altitudes, and *L. trunculata* at elevations >1,800 meters. ²²

References

1. Gastroenterol Clin North Am 1996 Sep ;25(3):627-36.
2. Curr Opin Infect Dis 2008 Oct ;21(5):523-30.
3. Am J Trop Med Hyg 2005 May ;72(5):508-9.
4. Neurosurgery 2006 Sep ;59(3):E706-7; discussion E706-7.
5. Clin Infect Dis 2007 Nov 1;45(9):1207, 1238-9.
6. Trans R Soc Trop Med Hyg 2009 Mar ;103(3):318-20.
7. Trop Doct 2010 Oct ;40(4):253-4.
8. Turk J Gastroenterol 2011 Jun ;22(3):347-50.

9. AJR Am J Roentgenol 2007 Jun ;188(6):1596-603.
10. Bull World Health Organ 1999 ;77(4):340-6.
11. Mayo Clin Proc 1998 May ;73(5):473-8.
12. Rev Peru Med Exp Salud Publica 2012 Sep ;29(3):386-9.
13. Turk J Gastroenterol 2010 Jun ;21(2):183-7.
14. Gastrointest Endosc 2009 Aug ;70(2):386-7; discussion 387.
15. Z Gastroenterol 2007 Apr ;45(4):313-6.
16. JOP 2005 Jan ;6(1):36-9.
17. Gastrointest Endosc 1991 Jul-Aug;37(4):473-5.
18. Ann Gastroenterol Hepatol (Paris) 1987 Mar-Apr;23(2):67-70.
19. Parasit Vectors 2012 ;5:196.
20. Onderstepoort J Vet Res 2002 Jun ;69(2):133-40.
21. J Comp Pathol 1960 Apr ;70:176-81.
22. Parasit Vectors 2012 ;5:196.

Filariasis - Bancroftian

Agent	PARASITE - Nematoda. Phasmidea, Filariae: Wuchereria bancrofti
Reservoir	Human
Vector	Mosquito (Anopheles, Aedes, Culex)
Vehicle	None
Incubation Period	5m - 18m (range 1m - 2y)
Diagnostic Tests	Identification of microfilariae in nocturnal blood specimen. Nucleic acid amplification. Serology may be helpful.
Typical Adult Therapy	Diethylcarbamazine : 50 mg day 1 50 mg TID day 2 100 mg TID day 3 Then 2 mg/kg TID X 18 days. OR Ivermectin 200ug/kg PO as single dose. Doxycycline 200 mg daily X 8 w is also effective.
Typical Pediatric Therapy	As for adult
Clinical Hints	Lymphangitis, lymphadenitis, eosinophilia, epididymitis, orchitis, hydrocoele or progressive edema; episodes of fever and lymphangitis may recur over several years; chyluria occasionally encountered.
Synonyms	Bancroftian filariasis, Rosetta leg, Wuchereria bancrofti. ICD9: 125.0 ICD10: B74.0

Clinical

WHO Case definition for surveillance:

Clinical case definition

- Hydrocoele or lymphedema in a resident of an endemic area for which other causes of these findings have been excluded.

Laboratory criteria for diagnosis

- Microfilaria positive, antigen positive or biopsy positive.

Case classification

Suspected: Not applicable.

Probable: A case that meets the clinical case definition.

Confirmed: A person with laboratory confirmation even if he/she does not meet the clinical case definition.

Clinical manifestations reflect either acute inflammation or lymphatic obstruction. ¹⁻³

- Repeated episodes of lymphangitis, lymphadenitis, fever, headache, backache and nausea may occur; and arthritis ⁴, funiculitis, epididymitis, or orchitis are common.
- In long-standing cases lymphedema or persistent adenopathy may develop.
- Hydrocoele ⁵ is the most common clinical manifestation of lymphatic filariasis, and causes sexual disability.
- Hydrocoelelectomy accounts for 25% of all surgical procedures performed in endemic areas of Ghana and Kenya.
- Lower limb involvement is characterized by initial pretibial pitting edema, which eventually becomes nonpitting and involves the entire leg.
- The skin of the leg or scrotum becomes thick, fissured, and warty; and ulceration and secondary infection may occur.
- Rare instances of pleural effusion ⁶, multiple subcutaneous nodules ⁷ and intra-abdominal cysts are reported ⁸
- Chyluria reflects rupture of swollen lymphatics into the urinary tract. ⁹⁻¹⁴ Microscopic (occasionally gross) hematuria is reported in some cases. ^{15 16}
- Filarial granuloma may mimic testicular cancer. ¹⁷

Microfilariae may be found in properly timed blood specimens, hydrocoele fluid, chylous urine and organ aspirates. ^{18 19}

- Adult worms are identified in biopsy material.
- Eosinophilia usually appears only during acute episodes of inflammation.

There is extensive evidence that endosymbiont bacteria (*Wolbachia* spp.) are necessary for the development of filarial larvae, and fertility of adult parasites. ²⁰⁻²³


- Doxycycline has proven effective in therapy, presumably through inhibition of *Wolbachia* spp. ²⁴⁻²⁷

This disease is endemic or potentially endemic to 117 countries.

Filariasis - Bancroftian in Uganda

Time and Place:

- High prevalence occurs north of Lake Victoria.
- 90.6% of the population were at risk for Bancroftian filariasis as of 2000.



Graph: Uganda. Filariasis - at risk, cases

Notes:

1. Number of persons targeted for mass treatment.
2. Additional references: 2008 ²⁸ 2010 ²⁹

Disease in Uganda is nocturnally periodic.


Prevalence surveys:

20.3% of school children in Moyo (20.3), 30.7% in Yumbe, 26.6% in Kotido, 24.9% in Katakwi and 21.5% in Lira districts (microfilaremia, 2000 to 2003). ³⁰

3.7% of school children in an endemic region in 2007; 0.4% in 2010 (microfilaremia) ³¹

18% to 30% of persons in the high altitude districts of Lira, Soroti and Katakwi

2% to 4% of persons in Masindi district (2011 publication) ³²



Graph: Uganda. Filariasis - mass treatment, cases

Notes:

1. Additional references: 2005 ³³ 2007 ³⁴ 2008 ³⁵ 2010 ³⁶

Vectors:

- The local vector is *Anopheles bwambae*.
- *Culex quinquefasciatus* is also implicated.
- *Anopheles gambiae* and *An. funestus* serve as vectors in high-altitude areas.

References

1. Ann N Y Acad Sci 2002 Dec ;979:131-42; discussion 188-96.
2. Int J Parasitol 2002 Jul ;32(8):947-60.
3. Parasite Immunol 2009 Nov ;31(11):664-72.
4. Mymensingh Med J 2007 Jul ;16(2 Suppl):S7-11.
5. PLoS Negl Trop Dis 2010 ;4(6):e695.
6. Trop Doct 2011 Oct ;41(4):238-9.
7. Acta Med Indones 2011 Oct ;43(4):249-51.
8. Indian J Radiol Imaging 2011 Jan ;21(1):18-20.
9. Can Med Assoc J 1928 Oct ;19(4):458-9.
10. BMJ Case Rep 2012 ;2012
11. Arch Ital Urol 1961 Aug ;34:305-17.
12. Dtsch Med Wochenschr 1969 May 16;94(20):1074-6.
13. Clin Radiol 1975 Apr ;26(2):237-42.
14. J Urol 1977 Mar ;117(3):393-4.
15. Trans R Soc Trop Med Hyg 2008 May ;102(5):506-7.
16. Am J Trop Med Hyg 1992 Jun ;46(6):745-51.
17. J Med Case Rep 2008 ;2:321.
18. Diagn Cytopathol 2008 Jan ;36(1):40-1.
19. Diagn Cytopathol 2011 Nov ;39(11):847-8.
20. Am J Trop Med Hyg 2005 Aug ;73(2):354-8.
21. Microbes Infect 2004 Jan ;6(1):113-28.
22. Cell Microbiol 2004 Feb ;6(2):97-104.
23. Cell Microbiol 2012 Dec 4;
24. J Infect Dis 2005 Oct 15;192(8):1483-93.
25. Lancet 2005 Jun 18-24;365(9477):2067-8.
26. Microbes Infect 2003 Apr ;5(4):261-73.
27. Am J Trop Med Hyg 2008 Jun ;78(6):854-5.
28. Wkly Epidemiol Rec 2009 Oct 9;84(42):437-44.
29. Wkly Epidemiol Rec 2011 Aug 26;86(35):377-88.
30. Trans R Soc Trop Med Hyg 2001 Mar-Apr;95(2):161-7.
31. Parasit Vectors 2011 ;4:134.
32. ProMED <promedmail.org> archive: 20110512.1458
33. Wkly Epidemiol Rec 2006 Jun 2;81(22):221-32.
34. Wkly Epidemiol Rec 2008 Sep 12;83(37):333-41.
35. Wkly Epidemiol Rec 2009 Oct 9;84(42):437-44.
36. Wkly Epidemiol Rec 2011 Aug 26;86(35):377-88.

Fungal infection - invasive

Agent	FUNGUS. Various (major syndromes such as Candidiasis, Blastomycosis, etc are discussed separately in this module)
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Culture of blood, urine, biopsy material. Serum antigen or antibody assay in some cases.
Typical Adult Therapy	Antifungal agent(s) directed at known or likely pathogen
Typical Pediatric Therapy	As for adult
Clinical Hints	This diagnosis should be suspected in any patient with evidence of severe local or multisystem infection, particularly in the setting of immune suppression.
Synonyms	Acremonium, Adiaspiromycosis, Allescheriasis, Alternaria, Arthrographis kalrae, Athopsis, Aureobasidium, Bipolaris, Blastobotrys proliferans, Chaetomium, Chrysosporium, Cladophialophora, Cladosporium, Curvularia, Cyphellophora, Dactylaria, Debaryomyces, Dreschslera, Emmonsia, Exophiala, Exserohilum, Fonsecaea, Fungal meningitis, Fungal sepsis, Fusarium, Geosmithia, Geotrichosis, Graphium, Hansenula, Haplomycosis, Hendersonula, Hyalophycomycosis, Kluyveromyces, Lasiodiplodia, Lasiodiplodia, Lecythophora, Malassezia furfur, Monascus, Monosporiosis, Mycocentrospora, Neocosmospora vasinfesta, Neosartorya hiratsukae, Neosartorya udagawae, Ochroconis, Oidiodendron, Paecilomyces, Paraconiothyrium, Phaeoacremonium, Phaeohyphomycosis, Phialophora, Phoma, Pichia, Pseudallescheria, Pseudallescheriasis, Pyrenochaeta, Ramichloridium, Rhinocladiella, Sarcopodium, Scedosporium, Septicemia - fungal, Taeniolella, Thielavia, Trichoderma, Ulocladium, Veronacea, Wallemia. ICD9: 117.6,117.8,117.9,118 ICD10: B43.1,B43.2,B43.8,B48.2,B48.3,B48.7,B48.8

Clinical

Major syndromes (Aspergillosis, Candidiasis, Coccidioidomycosis, Cryptococcosis, Penicilliosis, etc) are discussed elsewhere in this module.

Clinical syndromes associated with systemic fungal infection (in alphabetical order):

Adiaspiromycosis (Haplomycosis) is a pulmonary infection due to *Emmonsia* (*Chrysosporium*) species.

- Most cases have been described in Latin America and Central Europe, with additional reports from Israel and the United States.
- Three forms are recognized: solitary granuloma, localized granulomatous disease and diffuse, disseminated granulomatous disease. ¹

Arthrographis kalrae has been reported as a cause of sinusitis and meningitis in patient with AIDS.

Blastobotrys proliferans is an ascomyctous yeast that has been reported to cause peritonitis in a dialysis patient. ²

Curvularia inaequalis has been associated with several cases of peritonitis complicating peritoneal dialysis. ³

Exophiala jaenselmei and **Rhinocladiella** species have been implicated in cases of nosocomial fungemia.

- An outbreak of *Exophiala* infection in the United States was associated with contamination of injectable steroids.

Exserohilum is a dematiaceous fungus that has been associated with skin infections, keratitis, systemic infections and sinusitis. ⁴

Fusarium often infects the cornea ⁵, but may occasionally cause subcutaneous infection, fungemia, pneumonia, arthritis, bursitis, brain abscess and a variety of other systemic infection. ⁶

- Pathogenic members of the *Fusarium solani* complex are common in the environment. ⁷

Geotrichosis is a rare form of pneumonia and systemic mycosis caused by *Geotrichum candidum*.

- The organism is ubiquitous in nature and often found in the stool of healthy humans.
- Pulmonary disease simulates tuberculosis; and mucosal infection is similar to moniliasis.

Graphium basitruncatum has been associated with fungemia in a patient with leukemia. ⁸

Hansenula species have been implicated in nosocomial infections, endocarditis, fungemia and urinary tract infection

Lasiodiplodia theobromae has been reported to cause keratomycoses. ⁹

Neocosmospora vasinfecta, a plant pathogen, has caused at least 3 cases of soft tissue infection (lower extremities, in Senegal) or fatal disseminated infection in immunocompromised humans. ¹⁰

Neosartorya hiratsukae has been implicated in a case of brain abscess.

Penicillium • 31 cases of invasive infection by *Penicillium* species other than *P. marneffeii* were reported during 1951 to 2001 • including 12 of pulmonary disease, and 4 prosthetic valve endocarditis.

Phaeoophomycosis (infection by dematiaceous fungi) is manifested as:

- brain abscess (typically *Cladosporium trichoides*; also *Exophiala dermatitidis* ¹¹, *Fonsecaea pedrosoi*, *Ramichloridium obovoideum*, *Ochroconis gallopavum*, *Chaetomium atrobruneum*, et al),
- sinusitis (*Drechslera*, *Bipolaris*, *Exserohilum*, *Curvularia*, *Alternaria*, *Cladosporium*)
- subcutaneous infection (typically due to *Exophiala* and *Phialophora* species • occasionally *Fonsecaea*, *Cladosporidium*, *Alternaria*, *Dactylaria*, *Mycocentrospora*, *Phaeoacremonium* ¹², *Veronaea*, *Cyphellophora pluriseptata*, etc)
- endocarditis.

Pseudoallescheriasis (Petriellidiosis) is caused by *Scedosporium apiospermum* (*Pseudoallescheria boydii*) and may present as mycetoma; or infection of the brain, bone and joints, orbits and other tissues. ^{13 14}

Ramichloridium mackenziei has been reported to cause brain abscess in the Middle East.

Sarcopodium oculorum has been implicated as a cause of corneal ulcer in Brazil.

Trichoderma spp. are associated with peritonitis among dialysis patients, and disseminated infection in the immune-suppressed.

Fungal eye infection:

- Fungal endophthalmitis may be exogenous or endogenous.
- Clinically, onset is delayed and more gradual than infection due to bacteria.
- Hyaline fungi:
 - Fusarium* species are implicated in keratitis, scleritis and intraocular infections
 - Aspergillus* in keratitis following industrial trauma or surgery, orbital infection, dacryocystitis, scleritis and endophthalmitis
 - Scedosporium* in keratitis, scleritis, endophthalmitis, orbital infection
 - Paecilomyces* in keratitis, endophthalmitis and intralenticular infections
 - Acremonium* in keratitis and endophthalmitis.
- Dematiaceous fungi
 - Bipolaris*, *Curvularia*, *Exophiala*, *Exserohilum*, *Lecytophora* and *Phialophora* are implicated in keratitis and intraocular infections
 - Lasiodiplodia* in keratitis and endophthalmitis.
- Other fungal agents (*Candida*, *Cryptococcus*, *Coccidioides*, *Paracoccidioides*, *Blastomyces*, *Histoplasma*, *Sporothrix*) which may cause ocular infection are discussed separately in this module.

This disease is endemic or potentially endemic to all countries.

References

1. J Clin Microbiol 2005 Apr ;43(4):1495-504.
2. J Clin Microbiol 2007 Oct ;45(10):3453-5.
3. J Clin Microbiol 2005 Aug ;43(8):4288-92.
4. Eur J Clin Microbiol Infect Dis 2006 Apr ;25(4):247-53.
5. Am J Ophthalmol 2007 Feb ;143(2):356-8.
6. Clin Microbiol Rev 2007 Oct ;20(4):695-704.
7. J Clin Microbiol 2006 Jun ;44(6):2186-90.
8. J Clin Microbiol 2007 May ;45(5):1644-7.
9. Sabouraudia 1976 Jul ;14(2):155-70.
10. Emerg Infect Dis 2001 Jan-Feb;7(1):149-52.
11. J Child Neurol 2009 Mar ;24(3):342-5.
12. J Clin Microbiol 2006 Jun ;44(6):2207-11.
13. Expert Rev Anti Infect Ther 2005 Oct ;3(5):765-73.
14. Med Mycol 2006 Jun ;44(4):295-327.

Gastroenteritis - viral

Agent	VIRUS - RNA Calicivirus (Norwalk, Hawaii, Sapporo, Snow Mountain, Norovirus); Torovirus; or Astrovirus
Reservoir	Human
Vector	None
Vehicle	Food Water Shellfish Vegetables
Incubation Period	Norwalk 1d - 2d; astrovirus 3d - 4d
Diagnostic Tests	Demonstration of virus (electron microscopy or stool antigen analysis). Serology. Nucleic acid amplification.
Typical Adult Therapy	Stool precautions; supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Vomiting (less common with Astrovirus), abdominal pain; loose, watery diarrhea lasting 1 to 3 days; no fecal leucocytes; fever in 50% - headache and myalgia in some cases.
Synonyms	Aichi, Astroviridae, Astrovirus, Calicivirus gastroenteritis, Chiba, Cosavirus, Diarrhea, Gastroenterite virale, Hawaii agent gastroenteritis, Klassevirus, Mexico virus, Mini-reovirus, Minireovirus, Norovirus gastroenteritis, Norwalk agent gastroenteritis, Norwalk-like, Parkville virus gastroenteritis, Picobirnavirus, Recovirus, Roskilde disease, Saffold Cardiovirus, Salivirus, Sapovirus, Sapporo, Sapporo-like, Snow Mountain, SRSV gastroenteritis, Toronto virus, Torovirus, Vinterkraksjuka, Viral gastroenteritis, Winter vomiting disease. ICD9: 008.8,008.69,008.62,008.63,008.64,008.65,008.66,008.67 ICD10: A08.1,A08.2,A08.3,A08.4

Clinical

The onset of infection due to the Norwalk virus group may be gradual or abrupt, and is heralded by abdominal cramps with or without nausea.

- In most cases, both vomiting and diarrhea occur. ¹
- Four to eight non-bloody stools are passed per day; and fecal leucocytes are absent.
- 87% of patients with NLV infection develop diarrhea within 5 days; and only 60% of patients with Sapporo-like virus [SLV] infection.
- 59% of children below age 1 year develop vomiting with NLV, and 44% with SLV.
- Myalgias, malaise, headaches and benign febrile seizures ^{2 3} may also be present.
- A low-grade fever occurs in 50% of cases.
- Original publications stated that symptoms remit in 48 to 72 hours without sequelae; however, recent studies suggest that illness usually persists for 5 to 6 days.
- The duration of illness has been correlated with fecal concentration of virus.
- Residual dyspepsia, constipation or gastroesophageal reflux disease may persist following Norovirus infection. ⁴
- Cases of Guillain-Barre syndrome ⁵ and necrotizing enterocolitis in newborn infants have been ascribed to Norovirus infection. ^{6 7}
- Review of the clinical features of fatal Norovirus infection • see reference ⁸

Astrovirus diarrhea is similar to NLV infection; however, the former is characterized by a milder illness and lower incidence of vomiting. ⁹

Rare instances of meningitis have been associated with Saffold virus infection. ¹⁰

This disease is endemic or potentially endemic to all countries.

References

1. Gastroenterol Clin North Am 2001 Sep ;30(3):779-95.
2. Clin Infect Dis 2009 Apr 1;48(7):849-55.
3. J Paediatr Child Health 2011 Jun ;47(6):373-7.
4. Clin Infect Dis 2012 Oct ;55(7):915-22.
5. BMJ Case Rep 2012 ;2012
6. J Pediatr 2008 Sep ;153(3):339-44.
7. Pediatr Infect Dis J 2010 Jul ;29(7):644-7.
8. Am J Infect Control 2012 Dec 21;
9. Korean J Pediatr 2012 Mar ;55(3):77-82.
10. Emerg Infect Dis 2012 Jan ;18(1):7-12.

Gianotti-Crosti syndrome

Agent	UNKNOWN
Reservoir	Unknown
Vector	None
Vehicle	Unknown
Incubation Period	Unknown
Diagnostic Tests	Clinical features and skin biopsy findings.
Typical Adult Therapy	None
Typical Pediatric Therapy	None
Clinical Hints	Generalized skin eruption involving the extremities, face and buttocks; lymphadenopathy of the axillae and inguinal region; anicteric hepatitis; resolves in 15 to 42 days. Rare outbreaks have been reported.
Synonyms	Acrodermatitis papulosa infantilis, Papular acrodermatitis of childhood, Papulovesicular acrolocated syndrome. ICD9: 693.0 ICD10: L27.8

Clinical

Most patients are in the age group 2 to 6 years; however, the disease has occasionally been reported in infants and young adults. ^{1 2}

Clinical features are largely limited to discrete flat-topped papules on the face, extensor surfaces of the extremities and buttocks. ³

- The eruption is symmetrical, occasionally pruritic, either skin-colored or erythematous, and evolves over a period of two to three days.
- The skin lesions measure 2 to 4 mm in diameter, with a tendency for larger lesions among young children. ⁴
- Koebner phenomenon has been described.
- In most cases, the exanthem resolves after 15 to 20 days, but may persist for as long as 5 weeks.
- Hemorrhagic skin lesions and petechiae have been described in some cases. ⁵
- Prominent lymphadenopathy is noted, primarily in the inguinal and axillary regions.
- Hepatomegaly and anicteric hepatitis are common.

Gianotti-Crosti syndrome may be the only presenting manifestation of Epstein-Barr virus infection. ⁶

The features of Gianotti-Crosti syndrome may mimic those of atopic dermatitis. ⁷

The diagnosis is confirmed by skin biopsy, which reveals spongiosis of the upper epidermis and upper dermis, with perivascular lymphocytic and histiocytic infiltrates. ⁸

This disease is endemic or potentially endemic to all countries.

References

1. J Cutan Med Surg 2008 May-Jun;12(3):121-5.
2. Cutis 2012 Apr ;89(4):169-72.
3. Cutis 2001 Sep ;68(3):207-13.
4. Pediatr Dermatol 1991 Sep ;8(3):224-7.
5. J Am Acad Dermatol 1992 Feb ;26(2 Pt 1):207-10.
6. Turk J Pediatr 2008 May-Jun;50(3):302-4.
7. Asia Pac Allergy 2012 Jul ;2(3):223-6.
8. J Am Acad Dermatol 2000 Dec ;43(6):1076-9.

Giardiasis

Agent	PARASITE - Protozoa. Archezoa, Metamonada, Treponadea. Flagellate: <i>Giardia lamblia</i> [<i>G. intestinalis</i> , <i>G. duodenalis</i>]
Reservoir	Human Beaver Muskrat Dog Cat Carnivores Sheep Goat Horse Cattle
Vector	None
Vehicle	Food Water Fecal-oral Fly
Incubation Period	1w - 3w (range 3d - 6w)
Diagnostic Tests	String test (gelatin capsule containing string). Stool microscopy or antigen assay. Nucleic acid amplification.
Typical Adult Therapy	Metronidazole 250 mg PO TID X 5d. OR Nitazoxanide 500 mg PO BID X 3d OR Tinidazole 2 g PO X1. OR Furazolidone 100 mg PO QID X 7d. OR Paromomycin 10 mg/kg PO TID X 7d OR Quinacrine 100 mg PO TID X 5d
Typical Pediatric Therapy	Metronidazole 5 mg/kg PO TID X 5d. OR Tinidazole 50 mg PO X 1 (maximum 2g). OR Furazolidone 1.5 mg/kg QID X 7d OR Nitazoxanide : Age 1 to 3y 100 mg BID X 7 d Age 4 to 11y 200 mg BID X 7d
Clinical Hints	Foul smelling, bulky diarrhea, nausea and flatulence; may "wax and wane"; weight loss and low-grade fever are common.
Synonyms	Beaver fever, <i>Giardia duodenalis</i> , <i>Giardia intestinalis</i> , <i>Giardia lamblia</i> , Lambliasis. ICD9: 007.1 ICD10: A07.1

Clinical

The usual interval between infection and the onset of acute symptoms ranges from one to two weeks.

In most instances, the individual will experience sudden explosive, watery, foul-smelling diarrhea; excessive gas; abdominal pain; bloating; nausea; asthenia; and anorexia. ¹

- Symptoms consistent with irritable bowel syndrome and functional dyspepsia are reported in 80.5% and 24.5% of patients, respectively ²
- Upper gastrointestinal symptoms such as vomiting may predominate. ³
- Fever is unusual, and asymptomatic infection is common.
- Blood or mucus in the stool is rare, and there is neither leucocytosis nor eosinophilia.

Occasionally, the illness may last for months, or even years, causing recurrent episodes of impaired digestion, lactose intolerance, diarrhea, depression, asthenia and weight loss. ⁴⁻⁸

- Recurrence of symptoms is also common following effective treatment. ⁹
- Severe and prolonged infections are reported among patients with IgA deficiency and malnutrition.
- Infection in children may result in stunted growth, delayed development ^{10 11} and vitamin A deficiency. ¹²

Sequelae:

- Reactive arthritis may occasionally follow infection by *Giardia intestinalis*. ¹³
- Giardiasis has been implicated in the etiology of irritable bowel syndrome and chronic fatigue syndrome. ¹⁴

This disease is endemic or potentially endemic to all countries.

Giardiasis in Uganda

Prevalence surveys:

- 40.7% of persons near Kibale National Park, 12.4% of livestock and 11.1% of primates (2010 publication) ¹⁵
- 20.1% of asymptomatic children ages 1 to <5 years, in urban Kampala (2012 publication) ¹⁶
- 20% red colobus (*Ptilocolobus tephrosceles*), 5% of red-tailed guenons (*Cercopithecus ascanius*) and 0% of black-and-white colobus (*Colobus guereza*) from forest adjacent to a national park (*Cryptosporidium* sp. Or *Giardia* sp.) (Kibale National Park, 2005) ¹⁷

References

1. Clin Microbiol Rev 2001 Jan ;14(1):114-28.
2. BMC Gastroenterol 2009 ;9:27.
3. Curr Opin Infect Dis 2003 Oct ;16(5):453-60.
4. BMC Infect Dis 2009 ;9:206.
5. Trends Parasitol 2010 Feb ;26(2):75-82.
6. Fam Pract 2010 Jun ;27(3):255-9.
7. BMC Gastroenterol 2012 Feb 8;12(1):13.
8. BMC Gastroenterol 2013 Feb 12;13(1):28.
9. Scand J Prim Health Care 2009 ;27(1):12-7.
10. Lancet 2002 Feb 16;359(9306):564-71.
11. J Trop Pediatr 2004 Apr ;50(2):90-3.
12. Ann Nutr Metab 2010 ;57(3-4):228-33.
13. J Clin Rheumatol 2004 Apr ;10(2):86-8.
14. Gut 2012 Feb ;61(2):214-9.
15. PLoS Negl Trop Dis 2010 ;4(5):e683.
16. PLoS Negl Trop Dis 2012 Aug ;6(8):e1780.
17. J Parasitol 2007 Apr ;93(2):439-40.

Gnathostomiasis

Agent	PARASITE - Nematoda. Phasmidea: Gnathostoma spinigerum (rarely G. hispidum, G. doloresi and G. nipponicum)
Reservoir	Cat Dog Poultry Frog Fish
Vector	None
Vehicle	Food Fish Amphibian Reptile
Incubation Period	3w - 4w (range 2d - 1y)
Diagnostic Tests	Identification of larva in tissue. Serological testing in specialized laboratories.
Typical Adult Therapy	Albendazole 400 mg daily for 21 days has been recommended as an adjunct to surgical excision Ivermectin , 200 ug/kg PO as a single dose has also been advocated.
Typical Pediatric Therapy	As for adult
Clinical Hints	Follows ingestion of raw meat, poultry, fish or frog; migratory nodules of skin, soft tissues, brain or eye; eosinophilia; parasite may survive for more than 10 years in human tissue.
Synonyms	Gnathostoma, Gongylnematiasis, Larva migrans profundus, Nodular migratory eosinophilic panniculitis, Physaloptera, Spiruroid larva migrans, Wandering swelling, Yangtze edema. ICD9: 128.1 ICD10: B83.1

Clinical

Initial symptoms may include nausea, abdominal pain or urticaria.

- The presence of worms in skin or soft tissue results in migratory, pruritic or painful swellings which may be erythematous and attain a size of several centimeters. ¹
- Swellings may last for 1 to 4 weeks in a given area, and then reappear in a new location • a pattern which can continue for months or years.
- Findings of central nervous system infection (less than 1% of patients with subcutaneous gnathostomiasis) have included radiculopathy, meningitis, encephalitis, subarachnoid or intercerebral hemorrhage or paralysis. ²⁻⁴
- Other syndromes include eye infestation ⁵⁻¹⁰, persistent abdominal pain with hepatomegaly, or pneumonitis. ¹¹
- 74 cases of intra-ocular gnathostomiasis had been reported as of 2012 • including 14 from India. ¹²
- Eosinophilia is prominent. Eosinophiles may also be found in CSF and pleural effusions. ¹³

This disease is endemic or potentially endemic to 22 countries. Although Gnathostomiasis is not endemic to Uganda, imported, expatriate or other presentations of the disease have been associated with this country.

Gnathostomiasis in Uganda

A related species, *Physaloptera caucasica* has been reported to produce similar infection in Malawi, Mozambique and Uganda.

References

1. Clin Infect Dis 1993 Jan ;16(1):33-50.
2. Southeast Asian J Trop Med Public Health 2008 Sep ;39(5):800-3.
3. Clin Infect Dis 2009 Feb 1;48(3):322-7.
4. Emerg Infect Dis 2011 Jul ;17(7):1174-80.
5. Rev Infect Dis 1986 May-Jun;8(3):350-6.
6. J Neuroophthalmol 2006 Sep ;26(3):184-6.
7. Retina 2007 Jan ;27(1):67-73.
8. Eur J Ophthalmol 2007 Jan-Feb;17(1):130-2.
9. Indian J Med Microbiol 2007 Jul ;25(3):276-8.
10. Cases J 2009 ;2:9370.
11. Southeast Asian J Trop Med Public Health 2008 Sep ;39(5):804-7.
12. Am J Trop Med Hyg 2012 Apr ;86(4):620-3.
13. Emerg Infect Dis 2004 Sep ;10(9):1690-91.

Gonococcal infection

Agent	BACTERIUM. Neisseria gonorrhoeae An aerobic gram-negative coccus
Reservoir	Human
Vector	None
Vehicle	Sexual contact Childbirth Exudates
Incubation Period	2d - 7d
Diagnostic Tests	Smear (male), culture. Consult laboratory for proper acquisition & transport. Nucleic acid amplification.
Typical Adult Therapy	Ceftriaxone 250 mg IM X 1. PLUS Azithromycin 1 g PO as single dose.
Typical Pediatric Therapy	Weight <=45 kg: Ceftriaxone 125 mg IM X 1 Weight >45 kg: as for adult. PLUS Azithromycin
Clinical Hints	Copious urethral discharge (male) or cervicitis beginning 2 to 7 days after sexual exposure; PID; fever, painful pustules and suppurative arthritis (primarily encountered in postmenstrual females).
Synonyms	Blennorrhagie, Blenorrhagia, Gonococemia, Gonore, Gonorre, Gonorrea, Gonorrhea, Gonorrhoe, Gonorrhho, Gonorrhoe, Infeccion gonococica, Infeccoes gonococicas, Neisseria gonorrhoeae. ICD9: 098 ICD10: A54

Clinical

Gonorrhea:

Gonorrhea in males typically presents as urethral discomfort, dysuria, and discharge.

- The degree of discomfort and discharge are variable.
- Asymptomatic infection is common among females, but may also occur in males ¹
- Gonococcal epididymitis presents with unilateral pain and swelling localized posteriorly within the scrotum.
- Gonorrhea in the female are usually manifest as vaginal discharge and endocervicitis.
- The discharge is thin, purulent and mildly odorous.
- Dysuria or a scant urethral discharge may be present.
- Non-gonococcal urethritis, including infection by *Chlamydia trachomatis* and other *Neisseria* species ² may mimic gonococcal infection.
- Infection can be passed to the male urethra from the pharynx through fellatio. ³
- Levels of serum Prostate-specific Antigen (PSA) may be elevated in patients with gonorrhea. ⁴

Gonococcal PID:

Pelvic or lower abdominal pain suggests infection of the endometrium, fallopian tubes, ovaries or peritoneum.

- Pain may be midline, unilateral, or bilateral.
- Fever and vomiting may be present.
- Right upper quadrant pain from perihepatitis (Fitz-Hugh-Curtis syndrome) may occur following the spread of organisms upward along peritoneal planes to the hepatic capsule ⁵⁻⁷ (The syndrome is also reported as a complication of gonorrhea in males) ⁸

Other clinical forms: ⁹

Gonococcal proctitis is often asymptomatic, but rectal pain, pruritus, tenesmus, bloody diarrhea and rectal discharge may be present.

Gonococcal pharyngitis may be asymptomatic, or associated with severe inflammation. *Neisseria gonorrhoeae* is often present in throat specimens from patients with urethritis. ¹⁰

Gonococcal conjunctivitis is usually unilateral in adults; however, neonatal infection (ophthalmia neonatorum) involves both eyes.

- Symptoms include pain, redness, and a purulent discharge and may result in blindness.
- Rare instances of corneal perforation are reported. ¹¹

Disseminated gonococcal infection is characterized by joint or tendon pain, of single or multiple joints. ¹²

- Severe pain, swelling, and decreased mobility in a single joint (usually the knee) suggest purulent arthritis.
- Tenosynovitis is common, usually affecting the small joints of the hands.
- A rash is present in 25% of patients with gonococemia.
- Additional complications include meningitis, endocarditis, septic shock with ARDS ¹³, subcutaneous abscess, Fournier's

gangrene ¹⁴ , pyomyositis ¹⁵ and other localized infections. ¹⁶

This disease is endemic or potentially endemic to all countries.

Gonococcal infection in Uganda

Prevalence surveys:

- 1.1% of men and 1.9% of women in Rakai (2003 publication) ¹⁷
- 4.5% of female and 4.7% of male adolescents in Kampala (2006 publication) ¹⁸
- 40.0% of pregnant women (1973 publication) ¹⁹
- 6.8% of urban and 7.8% of rural women below age 20 (*Chlamydia trachomatis* and/or *Neisseria gonorrhoeae*, 2010 publication) ²⁰
- 7.1% of sexually experienced women below age 20 years (*Chlamydia trachomatis* and/or *Neisseria gonorrhoeae*, 2011 publication) ²¹
- 9.7% of HIV-positive patients (2012 publication) ²²

References

1. Sex Transm Dis 2006 May ;33(5):314-9.
2. Sex Transm Dis 2011 May ;38(5):439-41.
3. Sex Transm Dis 2011 May ;38(5):372-3.
4. Br J Cancer 2011 Aug 23;105(5):602-5.
5. Cleve Clin J Med 2004 Mar ;71(3):233-9.
6. Korean J Hepatol 2008 Jun ;14(2):178-84.
7. World J Gastroenterol 2008 Dec 7;14(45):6975-80.
8. Korean J Gastroenterol 2010 Mar ;55(3):203-7.
9. Curr Infect Dis Rep 2006 Mar ;8(2):132-8.
10. J Infect Chemother 2008 Dec ;14(6):442-4.
11. Sex Transm Infect 2010 Nov ;86(6):447-8.
12. Curr Infect Dis Rep 2006 Mar ;8(2):132-8.
13. Int J Infect Dis 2010 Sep ;14 Suppl 3:e239-41.
14. Case Rep Urol 2012 ;2012:312365.
15. Case Rep Infect Dis 2012 ;2012:790478.
16. AIDS Patient Care STDS 2007 Jan ;21(1):4-8.
17. Sex Transm Infect 2003 Apr ;79(2):98-105.
18. J Adolesc Health 2006 Mar ;38(3):213-21.
19. Bull World Health Organ 1990 ;68(5):639-54.
20. Sex Reprod Healthc 2010 Aug ;1(3):111-5.
21. Afr Health Sci 2011 Sep ;11(3):383-9.
22. J Infect Dev Ctries 2012 ;6(3):276-82.

Granuloma inguinale

Agent	BACTERIUM. Klebsiella granulomatis (formerly <i>Calymmatobacterium granulomatis</i>) An gram-negative bacillus
Reservoir	Human
Vector	None
Vehicle	Sexual contact Direct contact
Incubation Period	7d - 30d (range 3d - 1 year)
Diagnostic Tests	Identification of organism in stained smears. Culture in specialized laboratories (HEp-2 cells).
Typical Adult Therapy	Doxycycline 100 mg BID PO X 3w. Alternatives: Azithromycin 1 g daily X 3 w. Sulfamethoxazole/trimethoprim 800/160 mg BID X 3w Erythromycin 500 mg QID X 3w.
Typical Pediatric Therapy	Doxycycline 2 mg/kg BID X 2 to 3w (above age 8). Alternatives: Sulfamethoxazole/trimethoprim , Erythromycin or Azithromycin
Clinical Hints	Slowly expanding, ulcerating skin nodule with friable base; usually painless; may be complicated by edema or secondary infection - rarely spreads to bone or joints.
Synonyms	<i>Calymmatobacterium granulomatis</i> , Donovanosis, Granuloma genitoinguinale, Granuloma inguinale tropicum, Granuloma venereum, Sixth venereal disease. ICD9: 099.2 ICD10: A58

Clinical

The primary lesion of granuloma inguinale appears on the perineum or genitals in 80% to 90% of cases.

- Infection begins as a small painless papule or indurated nodule which progresses to a painless beefy-red ulcer with rolled edges and a friable surface.
- Multiple ulcers may coalesce, and new lesions may also form through autoinoculation. ¹
- Scar formation, deformity, keloids and lymphedema may develop. ²
- The most common sites of infection are the prepuce, coronal sulcus, and penile shaft; the labia and the fourchette .
- Rectal lesions may follow anal intercourse.
- Systemic disease of bones, joints, liver and lymphatics is rare, and may follow infection of the uterine cervix.
- Granuloma inguinale may present as mass lesions which mimic malignancy ³ or elephantiasis ⁴ ; and cutaneous metastases from mucinous carcinoma may mimic granuloma inguinale. ⁵

This disease is endemic or potentially endemic to all countries.

References

1. *Sex Transm Infect* 2002 Dec ;78(6):452-7.
2. *Int J STD AIDS* 2001 Jul ;12(7):423-7.
3. *South Med J* 2009 Jan ;102(1):104-5.
4. *Int J STD AIDS* 2012 Nov ;23(11):835-6.
5. *Intern Med* 2012 ;51(17):2479-81.

Hantavirus infection - Old World

Agent	VIRUS - RNA. Bunyaviridae, Hantavirus - Old world : Hantaan, Puumala, Dobrava/Belgrade, Saaremaa & Seoul viruses
Reservoir	Field mouse (Apodemus agrarius-Hantaan) Vole (Myodes glareolus-Puumala) Rat (Rattus norvegicus-Seoul) ? Bat ? Bird
Vector	None
Vehicle	Animal excreta
Incubation Period	12d - 21d (range 4d - 42d)
Diagnostic Tests	Serology. Viral culture. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive. Suggest Ribavirin : 1g IV q6h X 4d, then 0.5g q6h X 6d
Typical Pediatric Therapy	Supportive. Suggest Ribavirin
Vaccine	Hantavirus [old world]
Clinical Hints	Headache, backache, myalgia, diarrhea, vomiting, conjunctivitis, hemorrhage and azotemia; proteinuria and thrombocytopenia common; history of local rodent infestation may be elicited; case-fatality rates 0.1% (Puumala) to 15% (Belgrade).
Synonyms	Acute epidemic hemorrhagic fever, Bosnian hemorrhagic fever, Churilov disease, Dobrava/Belgrade, Endemic benign nephropathy, Epidemic hemorrhagic fever, Far eastern hemorrhagic fever, Haemorrhagic nephrosonephritis, Hantaan, Hemorrhagic fever & renal syndrome, Infectious hemorrhagic fever, Khabarovsk, Korean hemorrhagic fever, Mouse fever, Muju, Muroid virus nephropathy, Nephropathia epidemica, Puumala, Rodent-borne viral nephropathy, Saaremaa, Sandinavian epidemic nephropathy, Sangassou, Seoul, Sochi virus, Songo fever, Sorkfeber, Thailand virus, Thottapalayam, Topografov, Tula, Viral hemorrhagic fever, Viral hemorrhagic fevers. ICD9: 078.6 ICD10: A98.5

Clinical

The course of severe Hemorrhagic Fever and Renal Syndrome (HFRS • Nephropathia epidemica ¹) involves five overlapping stages: ²

- febrile
- hypotensive
- oliguric
- diuretic
- convalescent

Febrile stage:

It is not uncommon for one or more of these stages to be inapparent or absent. ^{3 4}

- The onset of the disease is sudden, with intense headache, backache, fever, and chills.
- Hemorrhage is manifested during the febrile phase as a flushing of the face or injection of the conjunctiva and mucous membranes.
- A petechial rash may appear on the palate and axillary skin folds.
- Extreme albuminuria, typically appearing on the fourth day, is characteristic of severe HFRS.
- Severe renal disease ⁵ is more likely in patients with thrombocytopenia. ⁶

Hypotensive stage:

As the febrile stage ends, hypotension may develop and last for hours to days, accompanied by nausea and vomiting.

- One-third of deaths occur during this phase, related to vascular leakage and shock.
- The severity and mortality rate for Seoul virus infection are reported to be lower than those for Hantaan virus infection. ⁷

The final (convalescent) stage can last weeks to months.

Additional findings:

- Electrocardiographic abnormalities are present in 57% of patients with Puumala virus infection ^{8 9} and patients who recover from the infection are at increased risk for death from cardiovascular disease. ¹⁰
- Approximately 50% of deaths occur during the subsequent (oliguric) phase.

- Both interstitial nephritis and glomerulonephritis are encountered. ¹¹
- Patients who survive and progress to the diuretic phase show improved renal function but may still die of shock or pulmonary complications.
- Overt lung disease suggestive of Hantavirus pulmonary syndrome has been reported in patients infected with Puumala virus. ¹²
- Residual hypopituitarism may follow severe Dobrava ¹³ or Puumala virus infection. ¹⁴
- Visual impairment is common ¹⁵ , and acute angle-closure glaucoma has been reported in some cases. ¹⁶
- Additional findings have included pancreatitis, cholangitis, myocarditis, pericarditis, orchitis ¹⁷ , meningitis, cerebritis and gastrointestinal hemorrhage. ¹⁸
- Severe central nervous system manifestations of Puumala virus infection appear to be most common among young male patients. ¹⁹

Case-fatality rates range from less than 0.1% for Hemorrhagic Fever Renal Syndrome [HFRS] caused by Puumala [PUU] virus to approximately 5% to 10% for HFRS caused by Hantaan [HTN] virus.

Scrub typhus and hemorrhagic fever with renal syndrome (HFRS) often coexist in areas of Asia. ²⁰

- Retro-orbital, lumbar, or flank tenderness; proteinuria; and microscopic hematuria, and hemorrhagic manifestations are most common in HFRS.
- Dermal eschar, regional lymphadenopathy, and maculopapular rash should suggest scrub typhus.
- Gastrointestinal symptoms of Hantavirus infection may mimic those of acute appendicitis. ²¹

This disease is endemic or potentially endemic to 93 countries.

References

1. JAMA 1977 Aug 22;238(8):874-7.
2. Scand J Infect Dis 2000 ;32(2):125-32.
3. Curr Top Microbiol Immunol 2001 ;256:135-51.
4. Eur J Emerg Med 2001 Mar ;8(1):17-20.
5. Virchows Arch A Pathol Anat Histol 1978 Feb 10;377(2):129-44.
6. Emerg Infect Dis 2004 Aug ;10(8):1420-5.
7. Scand J Infect Dis 2011 Aug ;43(8):632-41.
8. Scand J Infect Dis 2009 ;41(1):57-62.
9. Scand J Infect Dis 2005 ;37(8):594-598.
10. Emerg Infect Dis 2013 Jan ;19(1):126-8.
11. Clin Nephrol 2011 Jun ;75(6):550-6.
12. Eur J Clin Microbiol Infect Dis 2011 May ;30(5):685-90.
13. Emerg Infect Dis 2012 Jul ;18(7):1180-3.
14. Eur J Clin Microbiol Infect Dis 2010 Jun ;29(6):705-13.
15. Scand J Infect Dis 2012 Dec ;44(12):956-62.
16. Ophthalmologie 2011 Aug ;108(8):753-8.
17. Virol Sin 2011 Aug ;26(4):285-8.
18. Am J Trop Med Hyg 2011 Feb ;84(2):229-33.
19. BMC Infect Dis 2011 ;11:217.
20. Am J Trop Med Hyg 2007 May ;76(5):801-5.
21. J Clin Virol 2011 Feb ;50(2):164-6.

Hepatitis A

Agent	VIRUS - RNA. Picornaviridae, Hepatovirus: Hepatitis A virus
Reservoir	Human Non-human primate
Vector	None
Vehicle	Fecal-oral Food Water Fly
Incubation Period	21d - 30d (range 14d - 60d)
Diagnostic Tests	Serology. Nucleic acid amplification.
Typical Adult Therapy	Stool precautions; supportive
Typical Pediatric Therapy	As for adult
Vaccines	Hepatitis A Hepatitis A + Hepatitis B Immune globulin
Clinical Hints	Vomiting, anorexia, dark urine, light stools and jaundice; rash and arthritis occasionally encountered; fulminant disease, encephalopathy and fatal infections are rare (case-fatality rate 0.15% to 2.7%, depending on age).
Synonyms	Botkin's disease, Epatite A, HAV, Hepatite per virus A, Infectious hepatitis. ICD9: 070.0 ICD10: B15.0, B15.9

Clinical

WHO Case definition for surveillance of acute viral hepatitis (all types):

Clinical description

- Acute illness typically including acute jaundice, dark urine, anorexia, malaise, extreme fatigue, and right upper quadrant tenderness.
- Biological signs include increased urine urobilinogen and >2.5 times the upper limit of serum alanine aminotransferase.
- Note: Most infections occur in early childhood. A variable proportion of adult infections is asymptomatic.

Laboratory criteria for diagnosis

- Hepatitis A: IgM anti-HAV positive
- Hepatitis B: positive for Hepatitis B surface antigen (HBsAg) or IgM anti-HBc-positive
- Non-A, non-B: IgM anti-HAV and IgM anti-HBc (or HBsAg) negative

Note 1: The anti-HBc IgM test, specific for acute infection, is not available in most countries.

- HBsAg, often available, cannot distinguish between acute new infections and exacerbations of chronic hepatitis B, although continued HBsAg seropositivity (>6 months) is an indicator of chronic infection.

Note 2: For patients negative for hepatitis A or B, further testing for a diagnosis of acute hepatitis C, D, or E is recommended:

Hepatitis C: anti-HCV positive

Hepatitis D: HBsAg positive or IgM anti-HBc positive plus anti-HDV positive (only as co-infection or super-infection of hepatitis B)

Hepatitis E: IgM anti-HEV positive

Case classification

- Suspected: A case that is compatible with the clinical description.
- Probable: Not applicable.
- Confirmed: A suspected case that is laboratory confirmed or, for hepatitis A only, a case compatible with the clinical description, in a person who has an epidemiological link with a laboratory-confirmed case of hepatitis A (i.e. household or sexual contact with an infected person during the 15-50 days before the onset of symptoms).

Clinical features of Hepatitis A:

The prodrome is characterized by anorexia, asthenia, headache, myalgia and moderate fever.

- Patients develop nausea, vomiting and right upper abdominal pain • and later overt jaundice. ¹
- Symptoms persist for 4 to 8 weeks, and the patient may remain asthenic and anorectic for several months thereafter.
- As many as 90% of cases in children less than 5 years of age are asymptomatic; fewer 50% among adults.
- Relapses may occur for up to 6 months following the initial infection.
- Rare instances of acute disseminated encephalomyelitis ², myelitis ³, meningoencephalitis ⁴, acute cholestatic syndrome ⁵, acalculous cholecystitis ^{6 7}, urticaria ⁸, pancreatitis ^{9 10}, pleural effusion or ascites ^{11 12}, acute

glomerulonephritis or renal failure ¹³⁻²² , pure red-cell aplasia ²³ , cerebral venous thrombosis ²⁴ and rhabdomyolysis have been reported. ²⁵

- Concurrent HIV infection may prolong the duration of viremia in patients with hepatitis A. ²⁶

Hepatitis A accounts for 3.1% of acute hepatic failure cases (United States, 1998 to 2005) ²⁷

- The case-fatality rate is 0.1% among children below age 4 years; 0.4% ages 5 to 29 years; and 1% above age 40.
- 55% of hepatitis A patients with acute hepatic failure recover • the remainder either die of the disease or require transplantation. ²⁸

A false positive serological reaction toward Epstein-Barr virus has been associated with Hepatitis A. ²⁹

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 2001 Jan ;14(1):38-58.
2. Rev Neurol (Paris) 2008 Oct ;164(10):852-4.
3. Med Trop (Mars) 2010 Feb ;70(1):7-8.
4. Trop Doct 2010 Jul ;40(3):176-7.
5. Folia Med (Plovdiv) 2012 Jan-Mar;54(1):30-5.
6. Ugeskr Laeger 1991 Jul 15;153(29):2076.
7. Pediatr Emerg Care 2012 Jun ;28(6):560-1.
8. Am J Gastroenterol 1993 Feb ;88(2):277-8.
9. Pancreas 2008 May ;36(4):424-7.
10. JNMA J Nepal Med Assoc 2011 Jan-Mar;51(181):7-10.
11. Ann Trop Paediatr 2009 Dec ;29(4):317-9.
12. Australas Med J 2012 ;5(7):369-72.
13. Trop Doct 2009 Jul ;39(3):186-7.
14. Ren Fail 2009 ;31(8):756-64.
15. J Viral Hepat 2010 Sep ;17(9):611-7.
16. Korean J Gastroenterol 2010 Dec ;56(6):391-3.
17. Acta Paediatr 2011 Sep ;100(9):e132-4.
18. Diagn Microbiol Infect Dis 2011 Apr ;69(4):400-4.
19. Scand J Infect Dis 2012 Feb ;44(2):144-8.
20. J Clin Virol 2011 Nov ;52(3):192-7.
21. Korean J Gastroenterol 2007 Aug ;50(2):116-20.
22. PMID 19685373
23. Chonnam Med J 2011 Apr ;47(1):51-3.
24. Case Report Neurol Med 2012 ;2012:120423.
25. Korean J Hepatol 2009 Mar ;15(1):85-9.
26. Clin Infect Dis 2002 Feb 1;34(3):379-85.
27. Hepatology 2006 Dec ;44(6):1589-97.
28. Transplant Proc 2010 Dec ;42(10):4658-60.
29. Pediatr Infect Dis J 1994 May ;13(5):413-4.

Hepatitis B

Agent	VIRUS - DNA. Hepadnaviridae, Orthohepadnavirus: Hepatitis B virus
Reservoir	Human Non-human primate
Vector	None
Vehicle	Blood Infected secretions Sexual contact Transplacental
Incubation Period	2m - 3m (range 1m - 13m)
Diagnostic Tests	Serology. Nucleic acid amplification.
Typical Adult Therapy	Needle precautions; supportive. For post-exposure or chronic infection: Peginterferon alfa-2a or Peginterferon alfa-2b ; OR Lamivudine ; OR Adefovir
Typical Pediatric Therapy	As for adult
Vaccines	Hepatitis A + Hepatitis B Hepatitis B + Haemoph. influenzae Hepatitis B immune globulin Hepatitis B
Clinical Hints	Vomiting and jaundice; rash or arthritis occasionally noted; risk group (drug abuse, blood products, sexual transmission); cirrhosis or hepatoma may follow years after acute illness; fulminant and fatal infections are encountered.
Synonyms	Epatite B, HBV, Hepatite per virus B, Serum hepatitis. ICD9: 070.1 ICD10: B16.2,B16.9, B16.1

Clinical

WHO Case definition for surveillance of acute viral hepatitis (all types):

Clinical description

- Acute illness typically including acute jaundice, dark urine, anorexia, malaise, extreme fatigue, and right upper quadrant tenderness.
- Biological signs include increased urine urobilinogen and >2.5 times the upper limit of serum alanine aminotransferase.
- Note: Most infections occur in early childhood. A variable proportion of adult infections is asymptomatic.

Laboratory criteria for diagnosis

- Hepatitis A: IgM anti-HAV positive
- Hepatitis B: positive for Hepatitis B surface antigen (HBsAg) or IgM anti-HBc positive
- Non-A, non-B: IgM anti-HAV and IgM anti-HBc (or HBsAg) negative

Note 1: The anti-HBc IgM test, specific for acute infection, is not available in most countries.

- HBsAg, often available, cannot distinguish between acute new infections and exacerbations of chronic hepatitis B, although continued HBsAg seropositivity (>6 months) is an indicator of chronic infection.

Note 2: For patients negative for hepatitis A or B, further testing for a diagnosis of acute hepatitis C, D, or E is recommended:

Hepatitis C: anti-HCV positive

Hepatitis D: HBsAg positive or IgM anti-HBc positive plus anti-HDV positive (only as co-infection or super-infection of hepatitis B)

Hepatitis E: IgM anti-HEV positive

Case classification

- Suspected: A case that is compatible with the clinical description.
- Probable: Not applicable.
- Confirmed: A suspected case that is laboratory confirmed or, for hepatitis A only, a case compatible with the clinical description, in a person who has an epidemiological link with a laboratory-confirmed case of hepatitis A (i.e. household or sexual contact with an infected person during the 15-50 days before the onset of symptoms).

Clinical features of Hepatitis B:

Infection can be asymptomatic (particularly in young children) or quite mild, with only fatigue, anorexia, and malaise.

- Clinical disease with jaundice occurs in 50% of adults and 10% of young children.
- Extrahepatic manifestations include arthralgia, arthritis, rash ¹, inflammatory myopathy ², focal segmental glomerulosclerosis ³ and acute glomerulonephritis. ⁴⁻⁸
- Rare instances of pure red cell aplasia ⁹, symmetric sensorimotor polyneuropathy ¹⁰, and pancreatitis have been

reported. ¹¹

- Chronic infection occurs in most young children and in 5% to 10% of adults, and can lead to persistent hepatitis, retarded growth in children ¹², active hepatitis, cirrhosis, or hepatocellular carcinoma. ¹³
- Acute exacerbation of chronic Hepatitis B may occur. ¹⁴
- Patients with HBV-HDV coinfection appear to have more severe acute disease and a higher risk of fulminant hepatitis (2% to 20%) compared with those infected with HBV alone ¹⁵; however, chronic HBV infection appears to occur less frequently in persons with HBV-HDV coinfection.
- Concurrent HIV infection increases the incidence of cirrhosis and HCC among Hepatitis B carriers. ^{16 17}

One to two million deaths are attributed to hepatitis B annually. 25% of chronic carriers died of primary liver cancer or cirrhosis as adults.

- Hepatitis B is responsible for 60% to 80% of the world's primary liver cancer.
- Primary liver cancer is one of the three leading causes of cancer death in East Asia, Southeast Asia, the Pacific Basin and sub-Saharan Africa.
- Hepatitis B predominates among patients with hepatocellular carcinoma in most Asian, African and Latin American countries; while hepatitis C predominates in Japan, Pakistan, Mongolia, Egypt, Europe and the United States. ¹⁸

This disease is endemic or potentially endemic to all countries.

Hepatitis B in Uganda

Vaccine Schedule:

BCG - birth

DTwP - 6, 10, 14 weeks


DTwPHibHep - 6, 10, 14 weeks

Measles (monovalent) - 9 months

OPV - birth; 6, 10, 14 weeks

TT - Women of child-bearing age; +1, +6 months; +1, +1 year

Vitamin A - 6, 12, 18, 24, 30, 36, 42 months



Graph: Uganda. Hepatitis B - WHO-UNICEF est. % (HepB3) vaccine coverage

HBsAg-positivity surveys:

10.3% of adults (2005) ¹⁹

10% of persons in Kitgum - of which 30.6% are seropositive toward hepatitis D (1986) ²⁰

5% of adults attending a health program in Rakai, southwestern Uganda (2011 publication) ²¹

4.9% of HIV-positive pregnant women (Kampala, 2007 publication) ²²

- 15% of HIV-positive patients (2008 publication) ²³
- 9% of HIV-positive patients (2010 publication) ²⁴
- 6.1% of HIV-positive adolescents and young adults (Kampala, 2011 to 2012) ²⁵
- 11.0% of university students in Kampala (2000 to 2001) ²⁶
- 3.0% of blood donors in central Uganda (1999) ²⁷
- 8.7% of health workers (2006 publication) ²⁸
- 8% of hospital patients (RSA, 2010 publication) ²⁹
- 8.1% of health care workers in Mulgo (2003) ³⁰
- 87% of patients with primary hepatocellular carcinoma (Kampala, 2008) ³¹

Notable outbreaks:

- 2010 to 2011 - An outbreak (74 cases) of blood-borne Hepatitis B was reported in Moyo District. ³²
- 2011 - An outbreak (12 cases) of Hepatitis B was reported in Kitgum and Gulu. ³³
- 2013 - An outbreak (29 fatal cases) was reported in Moyo and Adjumani. ³⁴

References

1. J Travel Med 2011 May-Jun;18(3):224-5.
2. J Clin Neuromuscul Dis 2011 Sep ;13(1):26-37.
3. Nephrol Dial Transplant 2011 Jan ;26(1):371-3.
4. Curr Opin Gastroenterol 2004 May ;20(3):241-7.
5. Clin Liver Dis 2004 May ;8(2):403-18.
6. Postgrad Med J 2010 Aug ;86(1018):486-92.
7. Eur J Intern Med 2011 Apr ;22(2):161-6.
8. Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi 2010 Dec ;24(6):464-7.
9. Korean J Gastroenterol 2012 Sep 25;60(3):177-81.
10. J Clin Neurosci 2012 Nov 16;
11. Pancreas 2008 May ;36(4):424-7.
12. Adv Virol 2012 ;2012:670316.
13. Clin Liver Dis 2002 May ;6(2):317-34, v.
14. J Gastroenterol Hepatol 2009 Jul ;24(7):1179-86.
15. J Gen Virol 2009 Nov ;90(Pt 11):2638-43.
16. J Antimicrob Chemother 2010 Jan ;65(1):10-7.
17. Semin Liver Dis 2012 May ;32(2):114-9.
18. Br J Cancer 2007 Apr 10;96(7):1127-34.
19. Afr Health Sci 2009 Jun ;9(2):98-108.
20. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):144-7.
21. J Med Virol 2011 May ;83(5):796-800.
22. J Med Virol 2007 Dec ;79(12):1797-801.
23. Afr Health Sci 2008 Mar ;8(1):8-12.
24. Int J STD AIDS 2010 Aug ;21(8):553-7.
25. J Int AIDS Soc 2012 ;15(6):18433.
26. Afr Health Sci 2005 Jun ;5(2):93-8.
27. Trop Med Int Health 2006 Jun ;11(6):951-4.
28. Vaccine 2006 Nov 17;24(47-48):6930-7.
29. J Med Virol 2010 Aug ;82(8):1334-40.
30. BMC Infect Dis 2010 ;10:191.
31. Afr Health Sci 2011 Aug ;11 Suppl 1:S20-3.
32. ProMED <promedmail.org> archive: 20110304.0714
33. ProMED <promedmail.org> archive: 20110916.2826
34. ProMED <promedmail.org> archive: 20130108.1487843

Hepatitis C

Agent	VIRUS - RNA. Flaviviridae, Hepacivirus: Hepatitis C virus
Reservoir	Human
Vector	None
Vehicle	Blood Sexual contact Transplacental
Incubation Period	5w - 10w (range 3w - 16w)
Diagnostic Tests	Serology. Nucleic acid amplification.
Typical Adult Therapy	Needle precautions; supportive. If hepatocellular disease: Weekly Peginterferon alfa-2a 180 mcg SC or Peginterferon alfa-2b 1.5 mcg SC AND Ribavirin 400 mg in AM & 600 mg in PM daily AND Telepravir OR Bocepravir Duration per viral genotype
Typical Pediatric Therapy	Peginterferon alfa-2b 3 MU/m ² SC x1 weekly AND Ribavirin 15mg/kg
Clinical Hints	Vomiting and jaundice; may be history of transfusion within preceding 1 to 4 months; chronic hepatitis and fulminant infections are encountered.
Synonyms	Epatite C, HCV, Hepatite per virus C, Non-A, non-B parenteral hepatitis. ICD9: 070.2,070.3,070.44,070.51,070.54,070.7 ICD10: B17.1

Clinical

WHO Case definition for surveillance of acute viral hepatitis (all types):

Clinical description

- Acute illness typically including acute jaundice, dark urine, anorexia, malaise, extreme fatigue, and right upper quadrant tenderness.
- Biological signs include increased urine urobilinogen and >2.5 times the upper limit of serum alanine aminotransferase.
- Note: Most infections occur in early childhood. A variable proportion of adult infections is asymptomatic.

Laboratory criteria for diagnosis

- Hepatitis A: IgM anti-HAV positive
 - Hepatitis B: positive for Hepatitis B surface antigen (HBsAg) or IgM anti-HBc positive
 - Non-A, non-B: IgM anti-HAV and IgM anti-HBc (or HBsAg) negative
- Note 1: The anti-HBc IgM test, specific for acute infection, is not available in most countries.
- HBsAg, often available, cannot distinguish between acute new infections and exacerbations of chronic hepatitis B, although continued HBsAg seropositivity (>6 months) is an indicator of chronic infection.

Note 2: For patients negative for hepatitis A or B, further testing for a diagnosis of acute hepatitis C, D, or E is recommended:

Hepatitis C: anti-HCV positive ¹

Hepatitis D: HBsAg positive or IgM anti-HBc positive plus anti-HDV positive (only as co-infection or super-infection of hepatitis B)

Hepatitis E: IgM anti-HEV positive

Case classification

- Suspected: A case that is compatible with the clinical description.
- Probable: Not applicable.
- Confirmed: A suspected case that is laboratory confirmed or, for hepatitis A only, a case compatible with the clinical description, in a person who has an epidemiological link with a laboratory-confirmed case of hepatitis A (i.e. household or sexual contact with an infected person during the 15-50 days before the onset of symptoms).

Clinical features of Hepatitis C:

Patients with acute infection typically are either asymptomatic or have a mild clinical illness. ²

- 60% to 70% of patients have no symptoms
- 20% to 30% of patients have jaundice
- 10% to 20% of patients have non-specific symptoms, such as anorexia, malaise, or abdominal pain.

Clinical illness in patients with acute hepatitis C who seek medical care is similar to that of other types of viral hepatitis.

- The average time period from exposure to symptom onset is 6-7 weeks, whereas the average time period from exposure to seroconversion is 8-9 weeks.
- Anti-HCV can be detected in 80% of patients within 15 weeks after exposure, in >90% within 5 months after exposure, and in >97% by 6 months after exposure.

- Rarely, seroconversion is delayed for as long as 9 months after exposure.
- Rare instances of optic neuritis have been reported. ³

The clinical course is variable; and fluctuating elevations in serum ALT levels, are the most characteristic feature. ^{4 5}

- Fulminant hepatic failure following acute infection is rare.
- 15% to 25% of infections resolve without sequelae.
- Chronic HCV infection develops 75% to 85% of patients who exhibit persistent or fluctuating ALT elevations.
- 75% to 85% of patients with acute hepatitis C infection progress to chronic disease, and 20% to cirrhosis within 20 to 25 years. ⁶
- No clinical or epidemiological features among patients with acute infection are predictive of persistent infection or chronic liver disease.
- Chronic liver disease is usually insidious, progressing without symptoms or physical signs in the majority of patients during 20 or more years following acute infection.
- Cirrhosis develops in 10% to 20% of persons with chronic hepatitis C over a period of 20 to 30 years; and hepatic cell carcinoma in 1% to 5%.
- HCV infection appears to have little short-term impact on survival after bone marrow transplantation, but is a risk factor for veno-occlusive disease and graft-versus-host disease. ⁷
- Concurrent HIV infection shortens the time to development of chronic liver disease in patients with Hepatitis C. ⁸⁻¹³

Hepatitis B predominates among patients with hepatocellular carcinoma in most Asian, African and Latin American countries; while hepatitis C predominates in Japan, Pakistan, Mongolia, Egypt, Europe and the United States. ¹⁴

Additional manifestations seen in patients with chronic hepatitis C infection ¹⁵ may include mixed cryoglobulinemia with systemic vasculitis of the skin, erythema induratum ¹⁶, retarded growth in children ¹⁷, renal disease ¹⁸⁻²³; CNS vasculitis ²⁴, acute disseminated encephalomyelitis ²⁵, dorsal root ganglionopathy ²⁶ and other nervous system disorders ²⁷⁻³⁰; thrombocytopenia ³¹⁻³³; non-Hodgkin lymphoma; porphyria cutanea tarda and lichen planus ³⁴; hypothyroidism ³⁵; lymphocytic sialoadenitis (similar to that of Sjogren's syndrome) ³⁶; autoimmune and other rheumatological disorders ³⁷⁻⁴⁰, nectolytic acral erythema ⁴¹; scleritis ⁴²; and orbital plasmacytoma. ⁴³

This disease is endemic or potentially endemic to all countries.

Hepatitis C in Uganda

The nationwide carriage rate in 1997 was estimated at 1.17%.

Seroprevalence surveys:

- 4.1% of blood donors in central Uganda (1999) ⁴⁴
- 4% of children and 12% of their mothers (Kampala, 2006 publication) ⁴⁵
- 0.6% of HIV-positive pregnant women (Kampala, 2007 publication) ⁴⁶
- 3% of HIV-positive patients (2008 publication) ⁴⁷
- 3.3% of HIV-positive patients (Kampala, 2003 to 2004) ⁴⁸
- 13% of hospital patients (2010 publication) ⁴⁹
- 2.6% of hospital patients (2011 publication) ⁵⁰

References

1. J Hepatol 2012 Dec ;57(6):1349-60.
2. Am J Gastroenterol 2008 May ;103(5):1283-97; quiz 1298.
3. J Neuroophthalmol 2009 Jun ;29(2):128-33.
4. Clin Liver Dis 1997 Nov ;1(3):569-85.
5. Clin Liver Dis 2002 May ;6(2):317-34, v.
6. J Gastroenterol 2007 Jul ;42(7):513-21.
7. Hematology Am Soc Hematol Educ Program 2006 ;:375-80.
8. Lancet Infect Dis 2009 Dec ;9(12):775-83.
9. Semin Liver Dis 2012 May ;32(2):1127-34.
10. Eur Rev Med Pharmacol Sci 2012 Nov ;16(11):1473-83.
11. J Infect Dis 2013 Mar ;207 Suppl 1:S40-4.
12. J Infect Dis 2013 Mar ;207 Suppl 1:S26-32.
13. J Infect Dis 2013 Mar ;207 Suppl 1:S1-6.
14. Br J Cancer 2007 Apr 10;96(7):1127-34.
15. Clin Liver Dis 2008 Aug ;12(3):611-36, ix.
16. J Clin Virol 2009 Apr ;44(4):333-6.
17. Adv Virol 2012 ;2012:670316.
18. Postgrad Med J 2010 Aug ;86(1018):486-92.
19. Hepat Res Treat 2010 ;2010:534327.
20. Hepatol Int 2011 Jun 23;
21. Liver Int 2011 Sep ;31(8):1071-80.
22. Contrib Nephrol 2012 ;176:10-23.
23. Gastroenterol Hepatol (N Y) 2012 Jul ;8(7):434-45.
24. World J Gastroenterol 2012 Jan 14;18(2):188-91.
25. Yonsei Med J 2012 Jul 1;53(4):856-8.
26. Ann Hepatol 2012 Nov ;11(6):958-60.
27. Neurologist 2008 May ;14(3):151-6.
28. J Clin Neuromuscul Dis 2011 Sep ;13(1):14-25.
29. Clin Dev Immunol 2012 ;2012:236148.
30. Curr Neurol Neurosci Rep 2012 Dec ;12(6):642-54.
31. J Gastrointestin Liver Dis 2010 Dec ;19(4):381-5.
32. BMC Res Notes 2012 Mar 13;5(1):142.
33. Int J Lab Hematol 2012 Oct ;34(5):525-32.
34. Oral Dis 2010 Oct ;16(7):601-12.
35. Presse Med 2012 Feb ;41(2):190-5.
36. Rheum Dis Clin North Am 2009 Feb ;35(1):111-23.
37. Autoimmun Rev 2008 Oct ;8(1):48-51.
38. J Gastrointestin Liver Dis 2007 Mar ;16(1):65-73.
39. Rheumatol Int 2010 Feb ;30(4):515-7.
40. Clin Rheumatol 2010 Dec ;29(12):1373-80.
41. J Gastroenterol Hepatol 2006 Jul ;21(7):1200-6.
42. Ann Ophthalmol (Skokie) 2008 Fall-Winter;40(3-4):197-9.
43. Orbit 2009 ;28(1):71-3.
44. Trop Med Int Health 2006 Jun ;11(6):951-4.
45. Emerg Infect Dis 2006 Sep ;12(9):1440-3.
46. J Med Virol 2007 Dec ;79(12):1797-801.

47. Afr Health Sci 2008 Mar ;8(1):8-12.
48. Afr Health Sci 2009 Sep ;9(3):143-6.

49. J Med Virol 2010 Aug ;82(8):1371-8.
50. J Trop Med 2011 ;2011:598341.

Hepatitis D

Agent	VIRUS - RNA. Deltavirus: Hepatitis D virus - a 'satellite' virus which is encountered as infection with a co-virus (Hepatitis B)
Reservoir	Human
Vector	None
Vehicle	Infected secretions Blood Sexual contact
Incubation Period	4w - 8w (range 2w - 20w)
Diagnostic Tests	Serology. Nucleic acid amplification.
Typical Adult Therapy	Needle precautions; supportive Interferon alfa 2-a has been used.
Typical Pediatric Therapy	As for adult
Clinical Hints	Vomiting and jaundice - biphasic course often noted; occurs as a coinfection or superinfection of hepatitis B; may be chronic or fulminant (combined hepatitis B and delta carries a worse prognosis than seen with hepatitis B alone).
Synonyms	Epatite D, Hepatitis delta. ICD9: 070.41,070.52 ICD10: B17.0

Clinical

WHO Case definition for surveillance of acute viral hepatitis (all types):

Clinical description

- Acute illness typically including acute jaundice, dark urine, anorexia, malaise, extreme fatigue, and right upper quadrant tenderness. ¹
- Biological signs include increased urine urobilinogen and >2.5 times the upper limit of serum alanine aminotransferase.
- Note: Most infections occur in early childhood. A variable proportion of adult infections is asymptomatic.

Laboratory criteria for diagnosis

- Hepatitis A: IgM anti-HAV positive
- Hepatitis B: Positive for Hepatitis B surface antigen (HBsAg) or IgM anti-HBc positive
- Non-A, non-B: IgM anti-HAV and IgM anti-HBc (or HBsAg) negative

Note 1: The anti-HBc IgM test, specific for acute infection, is not available in most countries.

- HBsAg, often available, cannot distinguish between acute new infections and exacerbations of chronic hepatitis B, although continued HBsAg seropositivity (>6 months) is an indicator of chronic infection.

Note 2: For patients negative for hepatitis A or B, further testing for a diagnosis of acute hepatitis C, D, or E is recommended:

Hepatitis C: anti-HCV positive

Hepatitis D: HBsAg positive or IgM anti-HBc positive plus anti-HDV positive (only as co-infection or super-infection of hepatitis B)

Hepatitis E: IgM anti-HEV positive

Case classification

- Suspected: A case that is compatible with the clinical description.
- Probable: Not applicable.
- Confirmed: A suspected case that is laboratory confirmed or, for hepatitis A only, a case compatible with the clinical description, in a person who has an epidemiological link with a laboratory-confirmed case of hepatitis A (i.e. household or sexual contact with an infected person during the 15-50 days before the onset of symptoms).

Clinical features of Hepatitis D:

Hepatitis D is characterized by gradual onset of abdominal pain and vomiting, followed by development of jaundice.

- A biphasic course often noted.
- When coinfection by hepatitis B is often present, the course may be chronic or fulminant. ²
- 80% of patients with chronic hepatitis D infection progress to cirrhosis within 5 to 10 years. ³

This disease is endemic or potentially endemic to all countries.

Hepatitis D in Uganda

10% of persons in Kitgum are HBsAg-positive.
- 30.6% of the latter are seropositive toward hepatitis D (1986).⁴

References

1. Lancet 2008 Jul 26;372(9635):321-32.
2. J Gastroenterol Hepatol 1997 Apr ;12(4):289-98.
3. Curr Treat Options Gastroenterol 2007 Dec ;10(6):456-63.
4. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):144-7.

Hepatitis E

Agent	VIRUS - RNA. Caliciviridae: Hepatitis E virus
Reservoir	Human Rodent Pig
Vector	None
Vehicle	Fecal-oral Water Shellfish Blood (rare) Meat (rare)
Incubation Period	30d - 40d (range 10d - 70d)
Diagnostic Tests	Identification of virus by immune electron microscopy (stool). Serology. Nucleic acid amplification.
Typical Adult Therapy	Stool precautions; supportive
Typical Pediatric Therapy	As for adult
Vaccine	Hepatitis E
Clinical Hints	Clinically similar to hepatitis A - no chronic residua; severe or fatal if acquired during pregnancy (10% to 24% case-fatality rate).
Synonyms	Epatite E, Non-A, non-B enteric hepatitis. ICD9: 070.43,070.53 ICD10: B17.2

Clinical

WHO Case definition for surveillance of acute viral hepatitis (all types):

Clinical description

- Acute illness typically including acute jaundice, dark urine, anorexia, malaise, extreme fatigue, and right upper quadrant tenderness.
- Biological signs include increased urine urobilinogen and >2.5 times the upper limit of serum alanine aminotransferase.
- Note: Most infections occur in early childhood. A variable proportion of adult infections is asymptomatic.

Laboratory criteria for diagnosis

- Hepatitis A: IgM anti-HAV positive
- Hepatitis B: positive for Hepatitis B surface antigen (HBsAg) or IgM anti-HBc positive
- Non-A, non-B: IgM anti-HAV and IgM anti-HBc (or HBsAg) negative

Note 1: The anti-HBc IgM test, specific for acute infection, is not available in most countries.

- HBsAg, often available, cannot distinguish between acute new infections and exacerbations of chronic hepatitis B, although continued HBsAg seropositivity (>6 months) is an indicator of chronic infection.

Note 2: For patients negative for hepatitis A or B, further testing for a diagnosis of acute hepatitis C, D, or E is recommended:

Hepatitis C: anti-HCV positive

Hepatitis D: HBsAg positive or IgM anti-HBc positive plus anti-HDV positive (only as co-infection or super-infection of hepatitis B)

Hepatitis E: IgM anti-HEV positive

Case classification

- Suspected: A case that is compatible with the clinical description.
- Probable: Not applicable.
- Confirmed: A suspected case that is laboratory confirmed or, for hepatitis A only, a case compatible with the clinical description, in a person who has an epidemiological link with a laboratory-confirmed case of hepatitis A (i.e. household or sexual contact with an infected person during the 15-50 days before the onset of symptoms).

Clinical features of Hepatitis E:

In contrast to hepatitis A, hepatitis E is characterized by:

- relatively long incubation period
- prolonged clinical course
- severe and often fatal illness among pregnant women ¹⁻⁴, patients with pre-existing hepatic cirrhosis ⁵, hemodialysis patients ⁶ and possibly women taking oral contraceptive medication. ⁷
- poor protective value of immune serum globulin. ⁸

In most hepatitis E outbreaks, the highest rates of clinically evident disease have been among young to middle-age adults.

- Lower disease rates in younger age groups may be the result of anicteric and/or subclinical HEV infection.
- Clinical disease in western countries and Japan is most common among males and persons above age 60 years. ⁹

Clinical signs and symptoms are similar to those of other types of viral hepatitis and include abdominal pain, anorexia, dark urine, fever, hepatomegaly, jaundice, malaise, nausea, and vomiting.¹⁰

- Less common findings include arthralgia, arthritis¹¹, diarrhea, acute pancreatitis¹²⁻¹⁴, pruritus, an urticarial rash, severe thrombocytopenia¹⁵, photophobia, Guillain-Barre syndrome¹⁶⁻¹⁸, inflammatory polyradiculopathy¹⁹, encephalitis^{20 21}, aplastic anemia²², pregnancy associated with fetal ascites²³, and hemophagocytic syndrome.²⁴
- A false positive serological reaction toward Epstein-Barr virus has been reported in Hepatitis E virus infection.²⁵
- The case fatality rate for young adults is 0.5% to 3%; 15% to 20% for pregnant women.²⁶
- A subsequent publication estimated the CFR for all cases at 0.019 among non-pregnant patients vs. 0.198 among pregnant women.²⁷

The period of infectivity following acute infection is not known; however, virus excretion in stools has been demonstrated up to 14 days after illness onset.

- The period of viral excretion appears to be prolonged among patients with hematological malignancy.²⁸

Sporadic cases of chronic Hepatitis E virus infection are reported, notably among immunosuppressed patients.²⁹⁻³⁹

- Rare cases of chronic hepatitis E infection have been reported in immuno-competent individuals.^{40 41}

This disease is endemic or potentially endemic to all countries.

Hepatitis E in Uganda

Notable outbreaks:

2007 to 2009 - Outbreaks (10,300 cases, 172 fatal) was reported in Kaabong, Kitgum, Pader and West Nile districts and Bundibugyo region.⁴²⁻⁶²

2011 - Outbreaks (908 cases) were reported in Karamoja.⁶³

References

1. J Hepatol 2011 Jun ;54(6):1107-13.
2. Indian J Med Res 2001 Feb ;113:35-9.
3. Epidemiol Infect 2012 May ;140(5):767-87.
4. Emerg Infect Dis 2012 Sep ;18(9):1401-4.
5. J Hepatol 2007 Mar ;46(3):387-94.
6. Am J Nephrol 2010 ;31(5):398-407.
7. Am J Trop Med Hyg 2010 Jan ;82(1):12-5.
8. World J Gastroenterol 2004 Aug 1;10(15):2157-62.
9. Curr Opin Infect Dis 2006 Oct ;19(5):460-6.
10. Infect Dis Clin North Am 2000 Sep ;14(3):669-87.
11. Clin Rheumatol 2007 Nov ;26(11):1973-5.
12. Pancreas 2008 May ;36(4):424-7.
13. J Clin Virol 2011 Jul ;51(3):202-4.
14. JNMA J Nepal Med Assoc 2011 Jan-Mar;51(181):7-10.
15. J Clin Microbiol 2008 Jul ;46(7):2450-2.
16. World J Gastroenterol 2009 Apr 7;15(13):1645-7.
17. Infection 2011 Aug 30;
18. J Clin Neurosci 2012 Apr ;19(4):607-8.
19. Emerg Infect Dis 2011 Aug ;17(8):1510-2.
20. Emerg Infect Dis 2011 Feb ;17(2):173-9.
21. Ann Hepatol 2012 Sep-Oct;11(5):618-22.
22. J Clin Virol 2012 Mar 20;
23. Fetal Diagn Ther 2012 Jun 1;
24. Nihon Shokakibyō Gakkai Zasshi 2008 Jun ;105(6):841-6.
25. Eur J Gastroenterol Hepatol 2009 Dec ;21(12):1433-5.
26. J Med Virol 2008 Apr ;80(4):646-58.
27. Hepatology 2012 Apr ;55(4):988-97.
28. J Clin Virol 2010 Oct ;49(2):141-4.
29. Ned Tijdschr Geneesk 2010 ;154:A1790.
30. Ann Intern Med 2010 Jul 20;153(2):135-6.
31. Minerva Gastroenterol Dietol 2010 Jun ;56(2):121-8.
32. J Viral Hepat 2011 Mar ;18(3):227-8.
33. N Engl J Med 2009 Sep 3;361(10):1025-7.
34. Ann Intern Med 2009 Mar 17;150(6):430-1.
35. J Hepatol 2009 Mar ;50(3):622-4.
36. J Hepatol 2009 Feb ;50(2):435-7.
37. Hepatology 2008 Oct ;48(4):1328-30.
38. Neth J Med 2012 Aug ;70(6):261-6.
39. Med Clin (Barc) 2012 Feb 4;138(2):69-72.
40. Gastroenterol Hepatol 2011 Jun-Jul;34(6):398-400.
41. Hepatogastroenterology 2011 Mar-Apr;58(106):324-5.
42. Emerg Infect Dis 2010 Jan ;16(1):126-9.
43. Clin Infect Dis 2010 Apr 1;50(7):1006-10.
44. Clin Infect Dis 2010 Jul 1;51(1):117-8; author reply 118-9.
45. Am J Trop Med Hyg 2010 Nov ;83(5):1170-3.
46. PLoS One 2012 ;7(7):e41135.
47. ProMED <promedmail.org> archive: 20080304.0894
48. ProMED <promedmail.org> archive: 20080512.1612
49. ProMED <promedmail.org> archive: 20080529.1737
50. ProMED <promedmail.org> archive: 20080620.1919
51. ProMED <promedmail.org> archive: 20080716.21
52. ProMED <promedmail.org> archive: 20080804.2385
53. ProMED <promedmail.org> archive: 20080806.2413
54. ProMED <promedmail.org> archive: 20080813.2511
55. ProMED <promedmail.org> archive: 20080905.2777
56. ProMED <promedmail.org> archive: 20081014.3256
57. ProMED <promedmail.org> archive: 20081017.3288
58. ProMED <promedmail.org> archive: 20090309.0982
59. ProMED <promedmail.org> archive: 20090507.1705
60. ProMED <promedmail.org> archive: 20090530.2009
61. ProMED <promedmail.org> archive: 20090925.3366
62. ProMED <promedmail.org> archive: 20091211.4215
63. ProMED <promedmail.org> archive: 20110613.1803

Hepatitis G

Agent	VIRUS - RNA. Flaviviridae, Hepacivirus: Hepatitis G virus. HGBV-A, B and C appear to be related
Reservoir	Human
Vector	None
Vehicle	Blood Vertical transmission has also been documented Sexual transmission suspected
Incubation Period	Unknown
Diagnostic Tests	Serology. Nucleic acid amplification.
Typical Adult Therapy	Supportive. Alpha interferon has been shown to ? transiently eliminate the carrier state
Typical Pediatric Therapy	As for adult
Clinical Hints	Acute or chronic hepatitis acquired from blood (needles, etc); clinically milder than hepatitis C - most cases limited to anicteric elevation of hepatic enzyme levels; viremia documented for as long as 10 years.
Synonyms	Epatite G, Hepatitis GB. ICD9: 070,59 ICD10: B17.8

Clinical

Hepatitis G is characterized by acute or chronic hepatitis acquired from blood (needles, etc). ¹

- The disease is milder than hepatitis C, with most cases limited to anicteric elevation of hepatic enzyme levels. ²
- Viremia has been documented for as long as 10 years.
- A case of aplastic anemia complicating Hepatitis G infection has been reported. ³

This disease is endemic or potentially endemic to all countries.

Hepatitis G in Uganda

Prevalence surveys:

17.3% of HIV-positive patients (2007 publication) ⁴

References

1. World J Gastroenterol 2008 Aug 14;14(30):4725-34.
2. Curr Opin Infect Dis 2002 Oct ;15(5):529-34.

3. Virol J 2011 ;8:30.
4. Int J STD AIDS 2007 Apr ;18(4):244-9.

Herpes B infection

Agent	VIRUS - DNA. Herpesviridae, Alphaherpesviridae, Simplexvirus: Cercopithecine herpesvirus 1 (Herpes B virus)
Reservoir	Monkey (usually <i>Macaca</i> species and <i>cynomolgus</i>)
Vector	None
Vehicle	Contact or bite
Incubation Period	10d - 20d (range 2d - 60d)
Diagnostic Tests	Viral culture (skin exudates). Nucleic acid amplification. Biosafety level 4.
Typical Adult Therapy	Therapy: Acyclovir 12 mg/kg IV q8h. OR Ganciclovir 5 mg/kg IV q12h. Follow with prolonged Acyclovir 800 mg PO 5X daily. Postexposure prophylaxis: Valacyclovir 1g PO q8h X 14 days. OR Acyclovir 800 mg PO X 5 X 14 days
Typical Pediatric Therapy	Acyclovir or Ganciclovir as for adult.
Clinical Hints	Vesicles, lymphadenopathy, myalgia, singultus, major neurological signs; usually within one month following contact with monkey; case-fatality rates exceed 80%. permanent neurological residua are common.
Synonyms	Cercopithecine herpesvirus 1, Herpes B, Herpesvirus simiae, Macacine herpesvirus 1. ICD9: 078.89 ICD10: B00.4

Clinical

Most human infections have been fatal, consisting of myelitis and hemorrhagic encephalitis with concomitant multiorgan involvement. ¹

The illness begins with fever, malaise, diffuse myalgia, nausea, abdominal pain and headache.

- Lymphadenitis is seen proximal to the site of inoculation.
- Dermal vesicles may be present.
- Abdominal pain and nausea may occur.
- Neurological findings then predominate, with dysesthesia, ataxia, diplopia, seizures, and ascending flaccid paralysis. ²
- A lymphocytic CSF pleocytosis and elevated protein levels are noted, often with numerous erythrocytes.
- In contrast to herpes simplex infection, the encephalitis is multifocal.
- Rarely, isolated skin infection and even an isolated meningitis may be encountered.

This disease is endemic or potentially endemic to all countries.

References

1. [Emerg Infect Dis 2003 Feb ;9\(2\):246-50.](#)
2. [Clin Infect Dis 2002 Nov 15;35\(10\):1191-203.](#)

Herpes simplex encephalitis

Agent	VIRUS - DNA. Herpesviridae, Alphaherpesvirinae, Simplexvirus: Human herpesvirus (usually type I)
Reservoir	Human
Vector	None
Vehicle	Infected secretions, including Sexual contact
Incubation Period	Unknown
Diagnostic Tests	Viral culture CSF usually negative. CT brain. Compare CSF/blood antibody levels. Nucleic acid amplification.
Typical Adult Therapy	Acyclovir 10 mg/kg IV Q8h
Typical Pediatric Therapy	Acyclovir 10 mg/kg IV Q8h
Clinical Hints	Rapidly-progressive severe encephalitis, usually without exanthem; often unilateral, temporal and parietal lobe predominance; permanent residua and high case-fatality rate in untreated cases.
Synonyms	

Clinical

Although fever, headache, behavioral changes, confusion, focal neurological findings, and abnormal cerebrospinal fluid are suggestive of herpetic encephalitis, signs are not pathognomonic.

- Typical findings include fever, headache, psychiatric symptoms, altered consciousness, dysphagia, seizures¹ and vomiting.²
- Relatively severe and atypical presentations of encephalitis may occur in immunosuppressed patients.³
- Focal weakness, ataxia, hemiparesis, and memory loss are common.
- In some cases, patients exhibit memory loss, psychiatric disorders⁴⁻¹¹, photophobia, cranial nerve deficits, papilledema, loss of visual fields, olfactory disturbance¹², choreoathetosis¹³ or other movement disorders.¹⁴
- Concurrent herpetic encephalitis and cutaneous herpes simplex are uncommon.
- Infection is usually frontotemporal and unilateral, and characterized by severe, often fatal disease.¹⁵
- Unilateral involvement of the temporoparietal region is typical, and helps distinguish herpetic infection from other forms of viral encephalitis • which tend to be bilateral and symmetrical.¹⁶
- Cases of overt cerebral hemorrhage¹⁷, acute disseminated encephalomyelitis¹⁸, Charles Bonnet syndrome (complex visual hallucinations)¹⁹ and symmetric brain stem encephalitis have been reported.²⁰
- West Nile viral encephalitis may mimic herpes simplex encephalitis.²¹

Herpes encephalitis is a risk factor for acute retinal necrosis.²²

Relapse of encephalitis occurs in 12% of treated patients.^{23 24}

This disease is endemic or potentially endemic to all countries.

References

1. J Neurol 2012 Apr 18;
2. J Neurol 2005 Mar ;252(3):268-72.
3. Neurology 2012 Nov 20;79(21):2125-32.
4. Case Report Med 2012 ;2012:241710.
5. Am J Psychiatry 1976 Feb ;133(2):165-70.
6. Postgrad Med J 1978 Nov ;54(637):763-7.
7. Am J Med 1982 Sep ;73(3):445-8.
8. Fortschr Neurol Psychiatr 1982 Dec ;50(12):387-95.
9. South Med J 1985 Nov ;78(11):1347-50.
10. Biol Psychiatry 1992 Jul 15;32(2):211-2.
11. No To Hattatsu 2002 Jan ;34(1):61-5.
12. J Neurol 2010 Mar ;257(3):439-43.
13. Indian J Pediatr 2010 Aug ;77(8):901-2.
14. Eur J Neurol 2005 May ;12(5):331-43.
15. Eur J Neurol 2005 May ;12(5):331-43.
16. Pediatr Radiol 2007 Oct ;37(10):949-63.
17. AIDS Read 2009 Apr ;19(4):153-5.
18. Acta Neurol Belg 2010 Dec ;110(4):340-4.
19. Pediatr Neurol 2012 Apr ;46(4):250-2.
20. J Clin Neurosci 2009 Apr ;16(4):589-90.
21. Pediatr Neurol 2006 Jul ;35(1):62-4.
22. Neurology 2008 Oct 14;71(16):1268-74.
23. J Neurol 2006 Feb ;253(2):163-70.
24. J Child Neurol 2011 Mar ;26(3):369-72.

Herpes simplex infection

Agent	VIRUS - DNA. Herpesviridae, Alphaherpesvirinae, Simplexvirus: Human herpesvirus I and II
Reservoir	Human
Vector	None
Vehicle	Infected secretions, including Sexual contact
Incubation Period	1d - 14d
Diagnostic Tests	Viral culture or microscopy of lesions. Serology. Nucleic acid amplification.
Typical Adult Therapy	Acyclovir 400 mg PO TID X 7d. OR Famciclovir 250 mg PO TID X 7d. OR Valacyclovir 1 g PO BID X 7d
Typical Pediatric Therapy	Acyclovir 10 mg/kg PO QID X 7 d
Clinical Hints	Recurring localized crops of painful vesicles on a red base; regional adenopathy often present; may follow a prodrome of neuropathy or hyperesthesia.
Synonyms	Herpes gladiatorum, Herpes rugbiorum, Herpes simplex, Scrum pox. ICD9: 054.0,054.1,054.2,054.4,054.5,054.6,054.7,054.8,054.9 ICD10: A60,B00

Clinical

The initial attack of herpes simplex is generally more overt than recurrent episodes; however, primary infections are often asymptomatic. ¹

- Symptoms will also vary depending on the site of infection (eye ^{2 3} , esophagus ^{4 5} , anal region, etc).

Signs and symptoms:

Following a prodrome of local discomfort, tender papular, vesicular or ulcerative lesions on an erythematous base appear. ⁶

- Anorexia, malaise and fever may accompany individual episodes.
- The lesions coalesce, and tender bilateral lymphadenopathy develops.
- Skin lesions usually heal over the next several days to weeks.
- Patients may give a history of occupational exposure (ie, herpetic whitlow, found in medical or dental personnel; herpes gladiatorum among wrestlers).
- Vesicular skin lesions of tularemia may mimic those of herpes simplex ⁷ ; and herpetic infection may present as folliculitis. ⁸

Complications:

Immunosuppressed patients and neonates are at particular risk for disseminated and severe infections. ⁹⁻¹³

- Lesions of the tongue may present as herpetic geometric glossitis. ¹⁴
- Mucosal herpetic lesions may serve as a portal for bacterial invasion. ¹⁵
- Ocular complications include conjunctivitis, scleritis ¹⁶ , severe keratitis and retinal necrosis. ^{17 18}
- Corneal infection may present as epithelial keratitis (dendritica/geographica), stromal keratitis (necrotizing vs. non-necrotizing or "interstitial keratitis"), endotheliitis (disciform keratitis), neurotrophic keratopathy (metaherpetic keratitis) or vascularized corneal scars. ¹⁹⁻²¹
- Over 10% of keratouveitis cases are complicated by secondary glaucoma ²² Herpetic keratitis may complicated ocular steroid injection ²³
- Herpes simplex infection has been etiologically linked to facial (Bell's) palsy. ^{24 25}
- Pancreatitis ²⁶ , esophagitis ²⁷ , cardiomyopathy ²⁸ and rhabdomyolysis with renal failure have been reported to complicate herpes simplex infection. ²⁹
- Herpes simplex hepatitis is most common in the setting of pregnancy or immune suppression ³⁰ ; however, rare instances of hepatitis and fulminant hepatic failure due to HSV infection have been reported in immunocompetent persons. ³¹⁻³⁵
- HSV-related erythema multiforme ^{36 37} has been reported in stem-cell transplant recipients ³⁸
- Disseminated infection among patients with eczema (Eczema herpeticum) may resemble smallpox ^{39 40} or present as atopic dermo-respiratory syndrome. ⁴¹
- Chronic (>1 month) mucocutaneous infections may occur in HIV-positive patients, in the absence of disseminated disease. ⁴²
- Herpetic lesions in HIV-positive patients may be vegetative, hypertrophic, condyloma-like, nodular, ulcerative, or tumor-like

nodules or plaques. ^{43 44}

- Herpes simplex may contribute to the pathology of periodontitis. ⁴⁵
- Herpes simplex has been reported to cause Gerhardt syndrome (inspiratory dyspnea without dysphonia) from vocal cord paralysis ⁴⁶

Anterior uveitis • differential diagnosis:

Anterior uveitis due to Rubella virus is characterized by younger age at onset and a chronic course, typically associated with cataract at presentation. ⁴⁷

- Rubella virus has been implicated in the etiology of Fuchs heterochromic iridocyclitis. ⁴⁸
- Anterior uveitis due to Herpes simplex and Varicella-Zoster viruses is more common in adults, and often follows an acute course.
- Herpes simplex anterior uveitis presents with conjunctival redness, corneal edema, a history of keratitis, and the presence of posterior synechiae. Anterior chamber inflammation is common with Herpes simplex virus, while vitritis is more common with Rubella and Varicella-Zoster virus.
- Rubella, Herpes simplex and Varicella-zoster viruses are associated with intraocular pressure of more than 30 mmHg and development of glaucoma (18%-30%; P = 0.686).
- Focal chorioretinal scars were present in 22% of Rubella cases, 0% of HSV and in 11% of VZV uveitis cases.

Acquisition of Herpes simplex by the newborn at the time of delivery is associated with severe illness and results in death in approximately 50% of cases. ⁴⁹

- Neonatal herpes simplex infection is characterized by vesicular rash, hypothermia, lethargy, seizures, respiratory distress, hepatosplenomegaly, thrombocytopenia, hepatic dysfunction and cerebrospinal fluid pleocytosis. ^{50 51}

Herpes simplex virus is an important cause of encephalitis (discussed separately in this module) and keratitis. ⁵²

This disease is endemic or potentially endemic to all countries.

Herpes simplex infection in Uganda

Prevalence surveys:

- 62% of patients with genital ulcer disease (HSV-2, rural Rakai District, 2008 publication) ⁵³
- 43.8% of patients with genital ulcer disease (HSV-2, 1999 to 2001) ⁵⁴
- 96.3% of female genital ulcer disease with an identified cause (2009 publication) ⁵⁵

Seroprevalence surveys:

- 32.2% of males and 58.2% of females (ages 15 to 54) in Rakai district (HSV-2, 2000 publication) ⁵⁶
- 33.76% of men ages 15 to 49 in Rakai (HSV-2, 2009 publication) ⁵⁷
- 30.6% of men and 71.5% of women ages 15 to 54 in the rural southwest (HSV-2, 1999 publication) ⁵⁸
- 52% of sexually-active women ages 18 to 35 (HSV-2, 2007 publication) ⁵⁹

References

1. N Engl J Med 2004 May 6;350(19):1970-7.
2. Prog Retin Eye Res 2006 Jul ;25(4):355-80.
3. Medicine (Baltimore) 2008 May ;87(3):167-76.
4. Medicine (Baltimore) 2010 Jul ;89(4):204-10.
5. Pediatr Infect Dis J 2011 Oct ;30(10):911-2.
6. Dent Clin North Am 2005 Jan ;49(1):15-29, vii.
7. Clin Infect Dis 2008 Jul 1;47(1):e4-6.
8. Ned Tijdschr Geneesk 2009 ;153:A285.
9. Semin Pediatr Infect Dis 2005 Oct ;16(4):271-81.
10. Semin Perinatol 2007 Feb ;31(1):19-25.
11. Curr Opin Infect Dis 2007 Feb ;20(1):73-6.
12. Cancer 2009 Jan 1;115(1):199-206.
13. Infection 2010 Oct ;38(5):423-6.
14. Indian J Pathol Microbiol 2010 Jan-Mar;53(1):133-4.
15. J Periodontol 2008 Feb ;79(2):376-8.
16. Am J Ophthalmol 2009 Nov ;148(5):779-789.e2.
17. Pediatr Infect Dis J 2009 Feb ;28(2):163-4.
18. Arch Ophthalmol 2011 Apr ;129(4):403-8.
19. Ophthalmologie 2011 Apr ;108(4):385-95; quiz 396-7.
20. Prog Retin Eye Res 2012 Aug 7;
21. Prog Retin Eye Res 2012 Aug 27;
22. Int Ophthalmol 2010 Apr ;30(2):191-4.
23. Cornea 2009 May ;28(4):463-4.
24. Pediatr Infect Dis J 2008 May ;27(5):468-9.
25. J Med Virol 2010 Sep ;82(9):1582-5.
26. Eur J Gastroenterol Hepatol 2009 Jan ;21(1):114-6.
27. Medicine (Baltimore) 2010 Jul ;89(4):204-10.
28. Wien Klin Wochenschr 2010 Oct ;122(19-20):592-5.
29. South Med J 2008 Dec ;101(12):1271-2.
30. J Gastrointest Liver Dis 2011 Mar ;20(1):93-6.
31. Transpl Infect Dis 2007 Dec ;9(4):323-6.
32. Liver Transpl 2007 Oct ;13(10):1428-34.
33. Ned Tijdschr Geneesk 2009 ;153:A55.
34. J Clin Pathol 1969 Jan ;22(1):60-6.
35. Saudi J Kidney Dis Transpl 2011 Jan ;22(1):107-11.
36. Hua Xi Kou Qiang Yi Xue Za Zhi 2008 Aug ;26(4):452-3.
37. Contemp Clin Dent 2011 Oct ;2(4):372-5.
38. Arch Dermatol 2008 Jul ;144(7):902-7.
39. Emerg Infect Dis 2009 Jul ;15(7):1102-4.
40. Chem Immunol Allergy 2012 ;96:89-95.
41. Allergy 2011 Jul ;66(7):925-33.
42. J Am Acad Dermatol 2012 Jun ;66(6):e217-27.
43. Int J STD AIDS 2011 Apr ;22(4):181-6.
44. Int J Gynecol Pathol 2012 Jan ;31(1):33-7.
45. J Calif Dent Assoc 2011 Jun ;39(6):393-9.
46. Eur Ann Otorhinolaryngol Head Neck Dis 2012 May 11;
47. Ophthalmology 2011 Oct ;118(10):1905-10.
48. Graefes Arch Clin Exp Ophthalmol 2010 Oct ;248(10):1487-91.
49. Obstet Gynecol Surv 2011 Oct ;66(10):629-38.
50. Pediatr Infect Dis J 2008 May ;27(5):425-30.
51. Clin Obstet Gynecol 2012 Dec ;55(4):938-44.
52. J Infect Dis 2008 Sep 1;198(5):659-63.
53. Sex Transm Infect 2009 Apr ;85(2):97-101.
54. Sex Transm Infect 2005 Dec ;81(6):488-93.
55. Int J STD AIDS 2009 Sep ;20(9):650-1.
56. East Afr Med J 2000 Aug ;77(8):428-30.
57. J Infect Dis 2009 Apr 1;199(7):945-9.
58. Sex Transm Infect 1999 Apr ;75(2):98-102.
59. AIDS 2007 Jul 31;21(12):1515-23.

Herpes zoster

Agent	VIRUS - DNA. Herpesviridae, Alphaherpesvirinae: Varicella-zoster virus
Reservoir	Human
Vector	None
Vehicle	Air Direct contact
Incubation Period	Unknown
Diagnostic Tests	Viral culture (vesicles). Serology. Nucleic acid amplification.
Typical Adult Therapy	Acyclovir 800 mg PO X 5 daily X 7 to 10d. OR Famciclovir 500 PO TID. OR Valacyclovir 1 g PO TID
Typical Pediatric Therapy	Acyclovir 20 mg/kg PO QID X 7 to 10d
Vaccine	Herpes zoster
Clinical Hints	Unilateral dermatomal pain, tenderness and paresthesia followed in 3 to 5 days by macular, erythematous rash evolving to vesicles; trunk and chest most common, but other areas possible; patients usually above age 50.
Synonyms	Fuocodi Saint'Antonio, Shingles, Zona, Zoster. ICD9: 053 ICD10: B02

Clinical

The condition represents reactivation of dormant Varicella-Zoster virus in dorsal root ganglia.

Disease is characterized by grouped vesicular lesions distributed along one to three sensory dermatomes, usually unilateral and on the trunk or face. ¹

- Mild pruritis or excruciating pain may be present, and persis after the disappearance of the rash.
- Although pain typically presents for 1 to 3 days prior to the appearance of a rash, the pre-eruptive prodromal period may persist for as long as 18 days. ²
- In immunocompromised individuals, herpes zoster may become disseminated.
- A chronic verrucous form of herpes zoster seen in HIV-positive patients is associated with antiviral drug-resistance. ³

Most healthy persons recover without complications; however, individuals above age 50 years are at increased risk of postherpetic neuralgia which may persist for months to years after the rash has healed.

- The possible effect of antiviral drugs in prevention of pos-herpetic neuralgia is controversial. ⁴
- Immunocompromised patients are risk for chronic herpes zoster; or infection of the central nervous system ⁵, liver, lungs or pancreas.
- Chronic (>1 month) mucocutaneous infections may occur in HIV-positive patients, in the absence of disseminated disease. ⁶
- Visual impairment or scleral damage may follow zoster ophthalmia. ⁷⁻⁹ Over 10% of keratouveitis cases are complicated by secondary glaucoma ¹⁰ Rare instances of orbital apex syndrome ¹¹ and optic neuritis are also reported. ¹²
- VZ virus infection may be associated with myotomal paresis ¹³, facial nerve palsy ¹⁴ or Ramsay-Hunt syndrome (Bell palsy unilateral or bilateral, vesicular eruptions on the ears, ear pain, dizziness, preauricular swelling, tingling, tearing, loss of taste sensation, and nystagmus) ¹⁵
- VZ virus infection can be a presenting symptom of hyperparathyroidism and occurs twice as often in persons with hypercalcemia than age-matched controls. ¹⁶
- In some cases, reactivation of VZ virus may present as radiculitis, cranial nerve palsy or other features of herpes zoster • but without rash (zoster sine herpete). ¹⁷

This disease is endemic or potentially endemic to all countries.

References

1. N Engl J Med 2002 Aug 1;347(5):340-6.
2. Am J Med 2013 Jan 28;
3. Clin Exp Dermatol 1999 Sep ;24(5):346-53.
4. Med Mal Infect 2012 Feb ;42(2):53-8.
5. Lancet Neurol 2007 Nov ;6(11):1015-28.
6. J Am Acad Dermatol 2012 Jun ;66(6):e217-27.

7. Curr Opin Ophthalmol 2004 Dec ;15(6):531-6.
8. Ophthalmologie 2008 May ;105(5):480-4.
9. Curr Treat Options Neurol 2011 Feb ;13(1):79-91.
10. Int Ophthalmol 2010 Apr ;30(2):191-4.
11. Clin Ophthalmol 2011 ;5:1603-8.
12. Int Ophthalmol 2011 Jun ;31(3):233-6.
13. Am J Orthop (Belle Mead NJ) 2012 May ;41(5):220-2.
14. Pediatr Int 2006 Jun ;48(3):245-9.
15. J Craniofac Surg 2011 Sep ;22(5):1961-3.
16. Clin Infect Dis 2008 May 1;46(9):1452-4.
17. Curr Top Microbiol Immunol 2010 ;342:243-53.

Histoplasmosis

Agent	FUNGUS. Ascomycota, Euascomycetes, Onygenales: <i>Histoplasma capsulatum</i> var. <i>capsulatum</i> A dimorphic fungus
Reservoir	Soil Caves Chicken roosts Bat
Vector	None
Vehicle	Air
Incubation Period	10d - 14d (range 5d - 25d)
Diagnostic Tests	Fungal culture. Serologic tests less helpful. Antigen tests currently under study. Nucleic acid amplification.
Typical Adult Therapy	Itraconazole 200 mg daily X 9m For severe or immunocompromized patients: Amphotericin B 0.4 mg/kg/d X 6w, then 0.8 mg/kg qod X 8w
Typical Pediatric Therapy	Itraconazole 2 mg/kg daily X 9 m. For severe or immunocompromized patients: Amphotericin B 0.4 mg/kg/d X 6w, then 0.8 mg/kg qod X 8w
Clinical Hints	Fever, cough, myalgia, pulmonary infiltrates and calcifying hilar lymphadenopathy; chronic multisystem infection often encountered.
Synonyms	Darling's disease, <i>Histoplasma capsulatum</i> , Histoplasmosis, Ohio River Valley Fever, Ohio Valley disease, Reticuloendothelial cytomycosis. ICD9: 115.0 ICD10: B39.0,B39.1,B39.2,B39.3,B39.4

Clinical

Asymptomatic infection is common, and may be found as an incidental finding on chest X-ray, or through serological or skin tests. ¹

Pulmonary histoplasmosis:

Acute benign respiratory infection is characterized by weakness, fever, chest pains, and cough. ²

- The severity of illness is related to the magnitude of the exposure.
- Chronic pulmonary infection occurs in persons with pre-existing lung diseases such as emphysema.
- The infection is most common in males over the age of 40.
- Chronic pulmonary lesions are characterized by extensive cavitation, but may resemble those of tuberculosis. ³

Disseminated histoplasmosis:

Disseminated infection is seen in immunocompromised patients (AIDS ⁴⁻⁶, leukemia, corticosteroid therapy, anti-TNF therapy ⁷, etc) and may be characterized by fever, anemia, hepatitis, pneumonia, pleuritis, pericarditis ⁸, acalculous cholecystitis ⁹, meningitis, atypical skin lesions (10% of cases) ^{10 11} and ulcers of the mouth ¹², tongue ¹³, nose ¹⁴, paranasal sinuses ¹⁵, esophagus ^{16 17}, colon ¹⁸⁻²⁰ and larynx. ^{21 22}

- Associated findings include upper lobe cavitation with fibrosis (similar to tuberculosis); sclerosing mediastinitis with obstruction of the superior vena cava, pulmonary arteries and veins; esophagus; and constrictive pericarditis. ²³
- Fungemia is most common in patients with immunosuppression or neutropenia (<3,000 per cu mm). ²⁴
- Central nervous system infection can present as chronic meningitis, focal parenchymal lesions of the brain or spinal cord, stroke due to infected emboli, and diffuse encephalitis. ²⁵
- Spinal infection may mimic tuberculosis spondylodiscitis. ²⁶
- Adrenal masses ²⁷⁻³¹ and renal infection are occasionally reported ^{32 33} and may mimic carcinoma. ³⁴
- Peritoneal histoplasmosis has been reported as a complication of peritoneal dialysis. ³⁵
- Epididymo-orchitis and prostatitis are occasionally reported. ³⁶⁻³⁸
- Gastrointestinal infection may mimic colonic carcinoma ^{39 40} or abdominal tuberculosis. ⁴¹
- Dermatological manifestations include erythema nodosum ⁴², erythema multiforme ⁴³, purpuric lesions, or the appearance of ulcerating verrucous plaques ^{44 45}. Primary infection may present as a dermal nodule with regional adenopathy. ⁴⁶ Skin lesions may mimic secondary syphilis. ⁴⁷
- Hypercalcemia has been reported in some cases. ⁴⁸⁻⁵²

"Ocular histoplasmosis syndrome" is characterized by peripapillary atrophy, punched out lesions, a macular disciform lesion or scar in one eye without vitritis.

- The role of *Histoplasma capsulatum* in this condition is unclear. ^{53 54}
- Overt *Histoplasma* keratitis has been reported ⁵⁵

Acute disseminated infection is also seen in infants and young children and is marked by fever, cough, exhaustion and hepatosplenomegaly. ⁵⁶

- Roentgenographic findings include multiple nodules (3 to 4 mm) changing into punctate calcifications; histoplasmosis (non-calcifying nodules <3 mm); a "target lesion" (ie, central calcification); or hilar/mediastinal adenopathy ("popcorn" calcification).

Primary histoplasmosis of the mouth has been reported. ⁵⁷

This disease is endemic or potentially endemic to 93 countries. Although Histoplasmosis is not endemic to Uganda, imported, expatriate or other presentations of the disease have been associated with this country.

Histoplasmosis in Uganda

1997 (publication year) - A British resident acquired "histoplasmosis" in Uganda. ⁵⁸

2013 (publication year) - A Swiss traveler acquired "histoplasmosis" in Uganda. ⁵⁹

References

1. Trends Microbiol 2003 Oct ;11(10):488-94.
2. Am Fam Physician 2002 Dec 15;66(12):2247-52.
3. Mycoses 2006 Jul ;49(4):274-82.
4. AIDS 2008 May 31;22(9):1047-53.
5. Clin Infect Dis 1995 Aug ;21 Suppl 1:S108-10.
6. Clin Infect Dis 2000 Jun ;30(6):877-81.
7. Scand J Rheumatol 2009 ;38(4):311-6.
8. Medicine (Baltimore) 1983 Mar ;62(2):110-9.
9. J Infect Dev Ctries 2011 Mar ;5(3):235-8.
10. Mycopathologia 2007 Dec ;164(6):295-9.
11. Clin Dermatol 2012 Nov ;30(6):592-8.
12. Rev Med Chil 2010 May ;138(5):586-9.
13. Indian J Dermatol Venereol Leprol 2009 Mar-Apr;75(2):173-6.
14. Sao Paulo Med J 2010 Jul ;128(4):236-8.
15. Mycopathologia 2011 Jan ;171(1):57-9.
16. Am J Trop Med Hyg 2009 Mar ;80(3):347-50.
17. Ear Nose Throat J 2001 Oct ;80(10):702.
18. Rev Gastroenterol Peru 2010 Apr-Jun;30(2):163-6.
19. Tenn Med 2011 May ;104(5):49-50.
20. J Glob Infect Dis 2011 Apr ;3(2):195-8.
21. Semin Respir Infect 2002 Jun ;17(2):158-81.
22. Mycoses 2009 Nov ;52(6):539-40.
23. Clin Microbiol Rev 2007 Jan ;20(1):115-32.
24. Med Mycol 2010 Feb ;48(1):85-9.
25. Curr Treat Options Neurol 2008 May ;10(3):161-7.
26. Acta Reumatol Port 2008 Jul-Sep;33(3):360-3.
27. Diagn Cytopathol 2010 May ;38(5):357-9.
28. Diagn Cytopathol 2011 Apr ;39(4):294-6.
29. Diagn Cytopathol 2011 Mar 8;
30. Southeast Asian J Trop Med Public Health 2011 Jul ;42(4):920-5.
31. Med J Malaysia 2011 Dec ;66(5):504-6.
32. Mycopathologia 2009 Jun ;167(6):315-23.
33. Am J Kidney Dis 2011 Dec ;58(6):1018-21.
34. Virchows Arch 2009 Feb ;454(2):229-32.
35. Nat Rev Nephrol 2010 Jul ;6(7):435-9.
36. Am J Med Sci 2009 Sep ;338(3):238-40.
37. Transpl Infect Dis 2011 Oct ;13(5):489-91.
38. Indian J Urol 2012 Jul ;28(3):359-61.
39. Int J STD AIDS 2009 Jun ;20(6):429-30.
40. J Glob Infect Dis 2011 Apr ;3(2):195-8.
41. J Assoc Physicians India 2009 Jan ;57:76-8.
42. Arch Dermatol 1981 Nov ;117(11):709-12.
43. N Engl J Med 1966 Feb 24;274(8):415-20.
44. Dermatol Online J 2008 ;14(2):19.
45. Clin Dermatol 2012 Nov ;30(6):592-8.
46. Rev Soc Bras Med Trop 2008 Nov-Dec;41(6):680-2.
47. Dermatol Online J 2011 ;17(11):10.
48. J Med Liban 2012 Jul-Sep;60(3):165-8.
49. JAMA 1977 Mar 28;237(13):1350-2.
50. Am J Med 1985 May ;78(5):881-4.
51. Pediatr Infect Dis J 1994 May ;13(5):421-2.
52. J Infect 1999 Jul ;39(1):88-90.
53. Br J Ophthalmol 1999 May ;83(5):535-9.
54. Med Mycol 2012 Feb ;50(2):202-6.
55. Braz J Infect Dis 2007 Dec ;11(6):595-7.
56. Infect Dis Clin North Am 2003 Mar ;17(1):1-19, vii.
57. Int J Infect Dis 2010 Sep ;14 Suppl 3:e325-8.
58. J Oral Pathol Med 1997 Jul ;26(6):294-6.
59. Dtsch Med Wochenschr 2013 Feb ;138(7):313-6.

Histoplasmosis - African

Agent	FUNGUS. Ascomycota, Euascomycetes, Onygenales: <i>Histoplasma capsulatum</i> var. <i>duboisii</i> A dimorphic fungus
Reservoir	? Bat ? Chicken Baboon Bird
Vector	None
Vehicle	Air
Incubation Period	Unknown
Diagnostic Tests	Fungal culture or biopsy.
Typical Adult Therapy	Amphotericin B; or Ketoconazole
Typical Pediatric Therapy	As for adult
Clinical Hints	Skin nodule, abscess or ulcer; underlying bone lesion (commonly skull, femur and ribs) - may be multiple and associated with lymphadenopathy and weight loss.
Synonyms	African histoplasmosis. ICD9: 115.1 ICD10: B39.5

Clinical

Although *H. capsulatum* var. *duboisii* is probably acquired through inhalation, primary pulmonary infection has not been demonstrated.

- Infection is characterized by suppurative granulomata in the skin, bones and subcutaneous tissues • rarely in the abdominal viscera, lungs and other organs. ¹
- There is neither pain nor systemic illness in local infection; however, disseminated and even fatal infections are described.

Skin and bone are most frequently affected.

- Typical findings consist of ulcers, nodules or psoriaform lesions which may resolve spontaneously. ²⁻⁴
- Wide dissemination is occasionally reported. ⁵
- Cutaneous lesions may mimic molluscum contagiosum. ⁶
- Tender, subcutaneous cold abscesses may also be present.
- Osteolytic bone occur in up to 50% of cases • primarily of the skull and ribs, and often involving the vertebrae as well. ⁷
- Local, draining lymphadenitis may also be present.
- Immunosuppressed patients may develop miliary infection; or multiorgan involvement of liver, spleen, kidney and lung.
- Addison's disease resulting from adrenal gland involvement has been reported. ⁸

This disease is endemic or potentially endemic to 26 countries.

Histoplasmosis - African in Uganda

1997 (publication year) - A British resident acquired "histoplasmosis" in Uganda. ⁹

2003 (publication year) - A case of African histoplasmosis was confirmed in Japan in a Ugandan national with AIDS. ¹⁰

2013 (publication year) - A Swiss traveler acquired "histoplasmosis" in Uganda. ¹¹

Notable outbreaks:

2011 - An outbreak (9 cases) was reported among researchers studying in Uganda - from Austria, Madagascar, the Netherlands, Poland, South Africa, Sweden, Switzerland and the United Kingdom (2 cases). ¹²

References

1. Indian J Chest Dis Allied Sci 2000 Oct-Dec;42(4):271-7.
2. Rev Iberoam Micol 1997 Dec ;14(4):155-9.
3. Indian J Chest Dis Allied Sci 2000 Oct-Dec;42(4):271-7.
4. Am J Trop Med Hyg 2012 Feb ;86(2):261-3.
5. Eur J Clin Microbiol Infect Dis 2009 Jun ;28(6):697-9.
6. Bull Soc Pathol Exot 2008 Feb ;101(1):5-7.
7. Afr Health Sci 2001 Aug ;1(1):28-9.
8. Saudi Med J 2008 Jun ;29(6):904-6.
9. J Oral Pathol Med 1997 Jul ;26(6):294-6.
10. Nihon Ishinkin Gakkai Zasshi 2003 ;44(4):299-306.

11. [Dtsch Med Wochenschr 2013 Feb ;138\(7\):313-6.](#)
12. [ProMED <promedmail.org> archive: 20110908.2738](#)

HIV infection - initial illness

Agent	VIRUS - RNA. Retroviridae, Lentivirinae: Human Immunodeficiency Virus
Reservoir	Human
Vector	None
Vehicle	Blood Semen Sexual Transplacental Breast-feeding
Incubation Period	1w - 6w
Diagnostic Tests	HIV antibody (ELISA, Western blot). HIV or HIV antigen assays. Nucleic acid amplification.
Typical Adult Therapy	Supportive; 'prophylactic' Zidovudine + additional drugs (DDI, 3TC, etc) should be considered - particularly during pregnancy
Typical Pediatric Therapy	Supportive; role for 'prophylactic' Zidovudine + additional drugs (DDI, 3TC, etc) should be considered
Clinical Hints	Fever, diarrhea, sore throat and a mononucleosis-like illness in a 'high risk' patient (eg, men who have sex with men, drug abuser, etc).
Synonyms	HIV, HIV infection. ICD9: 042 ICD10: B20,B21,B22,B23,B24

Clinical

The clinical features of acute HIV infection are protean and often characterized by fever, generalized lymphadenopathy, headache, fatigue, myalgia, rash, nausea, vomiting, night sweats, sore throat, diarrhea or weight loss. ¹

- 40% to 90% of persons have symptoms suggestive of an acute viral infection.
- Symptoms tend to subside within two weeks; however, some patients continue to be ill for as long as ten weeks.
- In most cases, a history of likely acquisition within the past several weeks can be established: unprotected sex, extra-medical injection, transfusion, etc.

This disease is endemic or potentially endemic to all countries.

HIV infection - initial illness in Uganda

Data and background information regarding HIV infection are included in the note for AIDS

References

1. J Microbiol Immunol Infect 2005 Feb ;38(1):65-8.
2. Lancet Infect Dis 2008 Aug ;8(8):471.
3. AIDS 2008 Feb 19;22(4):503-10.
4. J Acquir Immune Defic Syndr 2008 Oct 1;49(2):183-9.
5. Bull World Health Organ 2008 Apr ;86(4):302-9.
6. Afr Health Sci 2007 Sep ;7(3):124-8.
7. Trop Med Int Health 2006 Jun ;11(6):951-4.
8. BMC Public Health 2006 ;6:91.
9. Trop Med Int Health 2008 May ;13(5):680-2.
10. PLoS Negl Trop Dis 2009 ;3(6):e473.
11. Int J STD AIDS 2009 Jul ;20(7):499-502.
12. Trans R Soc Trop Med Hyg 2006 Jun ;100(6):586-93.
13. Trans R Soc Trop Med Hyg 2007 Sep ;101(9):899-907.
14. Trans R Soc Trop Med Hyg 2006 Jun ;100(6):543-50.
15. Sex Transm Infect 1999 Apr ;75(2):98-102.
16. Trans R Soc Trop Med Hyg 2009 Jun ;103(6):541-8.
17. AIDS 2007 May 11;21(8):1027-34.
18. J Acquir Immune Defic Syndr 2008 Jun 1;48(2):203-10.
19. AIDS 2009 Jul 31;23(12):1589-94.
20. WHO AIDS Tech Bull 1989 Mar ;2:37.
21. Int J Tuberc Lung Dis 2006 Jul ;10(7):761-7.
22. J Acquir Immune Defic Syndr Hum Retrovirol 1998 Dec 15;19(5):484-9.
23. AIDS 2002 May 3;16(7):1031-8.
24. Nihon Ishinkin Gakkai Zasshi 2003 ;44(4):299-306.
25. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):144-7.
26. Trans R Soc Trop Med Hyg 1991 Mar-Apr;85(2):227-9.
27. Acta Trop 2006 Dec ;100(3):218-22.
28. J Med Virol 2007 Dec ;79(12):1797-801.
29. Afr Health Sci 2008 Mar ;8(1):8-12.
30. Int J STD AIDS 2007 Apr ;18(4):244-9.
31. BMC Public Health 2008 ;8:416.

Hookworm

Agent	PARASITE - Nematoda. Phasmidea: <i>Necator americanus</i> , <i>Ancylostoma duodenale</i> , <i>A. ceylonicum</i> (in Calcutta and the Philippines)
Reservoir	Human
Vector	None
Vehicle	Soil Contact
Incubation Period	7d - 2y
Diagnostic Tests	Examination of stool for ova.
Typical Adult Therapy	Albendazole 400 mg X 1 dose. OR Mebendazole 100 mg BID X 3d. OR Pyrantel pamoate 11 mg/kg (max 3g) X 3d; or
Typical Pediatric Therapy	As for adult
Clinical Hints	Pruritic papules (usually of feet) - later cough and wheezing; abdominal pain and progressive iron-deficiency anemia; eosinophilia common; dyspnea and peripheral edema in heavy infections; <i>Ancylostoma caninum</i> implicated in eosinophilic enteritis.
Synonyms	<i>Anchilostoma</i> , <i>Ancylostoma ceylanicum</i> , <i>Ancylostoma duodenale</i> , <i>Ancylostomiasis</i> , <i>Anquilostomiasis</i> , <i>Cyclodontostomum</i> , <i>Eosinophilis enteritis</i> , <i>Hakenwurmer-Befall</i> , <i>Miner's anemia</i> , <i>Necator americanus</i> , <i>Necatoriasis</i> , <i>Uncinariasis</i> . ICD9: 126.0,126.1 ICD10: B76.0,B76.1,B76.8

Clinical

Initial manifestations of hookworm consist of pruritus, erythema, and a papular, or vesicular rash at the site of larval penetration ("ground itch"). ¹

- Migration of larvae through the lungs may result in a Loeffler-like syndrome with transitory cough, wheezing, diffuse opacities on x-ray and eosinophilia in sputum and blood. ^{2 3}
- Migration of *A. duodenale* larvae to the breast, with infection of nursing infants ('hypobiosis') has been described. ⁴
- The major finding in overt infection is iron-deficiency anemia. ⁵
- Heavy intestinal infection may also produce local symptoms of abdominal pain, diarrhea, and occasionally malabsorption with weight loss (most commonly in children).
- Rare instances of overt bleeding ⁶ or melena have been reported. ⁷

This disease is endemic or potentially endemic to all countries.

Hookworm in Uganda

Prevalence surveys:

- 43.5% nationwide (1998 to 2005) ⁸
- 39.9% of persons in rural Uganda (2010 publication) ⁹
- 32.1% of persons in Kabale District (2003) ¹⁰
- 44.5% of children below age 20 in Southern Uganda (1998) ¹¹
- 12.9% of primary school children in Kampala (1997 publication) ¹²
- 51.8% of school children in northwest and southeast Uganda (2002 to 2003) ¹³
- 50.9% of school children in 2003; 24.1% in 2004; 10.7% in 2005 (nationwide)
- 50.0% of school children in the Great Lakes region of Uganda, Tanzania, Kenya and Burundi (2010 publication) ¹⁴
- 16% of children below age 3 years living on the Ugandan shore of Lake Victoria (2006 publication) ¹⁵
- 44.5% of healthy pregnant women at enrolment to a trial of deworming (Entebbe, 2007 publication) ¹⁶
- 2.4% of school children in proximity to Lake Victoria (2009 publication) ¹⁷
- 45% of pregnant women (Entebbe and Katabi, 2003 to 2005) ¹⁸
- 66.6% of pregnant women in Masindi District (2003 to 2004) ¹⁹

44.5% of pregnant women in Entebbe (2004 to 2005) ²⁰
45% of pregnant women in Entebbe (2010 publication) ²¹

References

1. Dermatol Clin 1989 Apr ;7(2):275-90.
2. Gastroenterol Clin North Am 1996 Sep ;25(3):579-97.
3. Semin Respir Infect 1988 Jun ;3(2):172-8.
4. Trans R Soc Trop Med Hyg 1995 Jul-Aug;89(4):379.
5. Lancet 2006 May 6;367(9521):1521-32.
6. Asian Pac J Trop Med 2012 Apr ;5(4):331-2.
7. Ann Trop Paediatr 2008 Dec ;28(4):293-6.
8. Trop Med Int Health 2005 Nov ;10(11):1187-9.
9. PLoS Negl Trop Dis 2010 ;4(6):e713.
10. Trans R Soc Trop Med Hyg 2005 Jan ;99(1):18-24.
11. East Afr Med J 2001 Jun ;78(6):283-6.
12. East Afr Med J 1997 May ;74(5):311-4.
13. Trans R Soc Trop Med Hyg 2006 Nov ;100(11):1039-48.
14. Trop Med Int Health 2010 Feb ;15(2):198-207.
15. Ann Trop Med Parasitol 2006 Jun ;100(4):315-26.
16. Trans R Soc Trop Med Hyg 2007 Sep ;101(9):899-907.
17. Geospat Health 2009 Nov ;4(1):39-53.
18. J Infect Dis 2008 Sep 15;198(6):920-7.
19. Trans R Soc Trop Med Hyg 2008 Feb ;102(2):130-6.
20. PLoS Negl Trop Dis 2009 ;3(6):e473.
21. Clin Infect Dis 2010 Feb 15;50(4):531-40.

Human herpesvirus 6 infection

Agent	VIRUS - DNA. Herpesviridae, Betaherpesvirinae, Roseolovirus: Herpesvirus 6 (Herpesvirus 7 is also implicated)
Reservoir	Human
Vector	None
Vehicle	Droplet Contact
Incubation Period	10d - 15d
Diagnostic Tests	Viral isolation and serologic tests rarely indicated. Nucleic acid amplification has been used
Typical Adult Therapy	Supportive Gancyclovir has been used in unusual and severe cases.
Typical Pediatric Therapy	As for adult
Clinical Hints	High fever followed by sudden defervescence and fleeting rash; most patients are below the age of 2 years; only 10% to 20% of herpesvirus 6 infections are associated with a rash.
Synonyms	Dreitagefieber, Exanthem criticum, Exanthem subitum, Herpesvirus 6, HHV-6, Pseudorubella, Roseola, Roseola infantilis, Roseola subitum, Sixth disease, Zahorsky's disease. ICD9: 057.8 ICD10: B08.2

Clinical

Roseola typically is characterized by high fever (often to 40 C) lasting from three to seven days, followed by rapid defervescence and a characteristic pink rash. ^{1 2}

- The rash is maculopapular or erythematous, beginning on the trunk and spreading to the neck and extremities. ³
- Skin lesions are discrete, not pruritic, blanch on pressure and fade within 3 to 48 hours.

Diarrhea, cough and irritability are common, and seizures may rarely occur in individual cases. ⁴

- HHV-6 infection accounts for 10% to 20% of febrile seizures in children below the age of two years. ^{5 6}
- Other findings may include bulging anterior fontanel, Nagayama spots (erythematous papules on the soft palate and uvula), periorbital edema, inflamed tympanic membranes, cervical, post auricular, and post occipital lymphadenopathy, splenomegaly, meningitis with radiculitis ⁷, encephalopathy or encephalitis ⁸⁻¹⁶, arthropathy (4.3% of cases) ¹⁷, rhabdomyolysis ¹⁸, uveitis ^{19 20}, optic neuritis ²¹, corneal inflammation ²² and conjunctival injection.
- Rare instances of acute hepatic failure ²³ and purpura fulminans have been reported. ²⁴

Reactivation and severe disease have been encountered in bone-marrow, solid organ transplant and other immune-deficient patients. ²⁵⁻²⁸

- HHV-6-associated pleurisy has been reported following stem-cell transplantation (2007 publication) ²⁹
- Fatal hepatitis and myocarditis has been reported in immunocompetent adults. ^{30 31}

This disease is endemic or potentially endemic to all countries.

References

1. J Med Microbiol 2003 Jan ;52(Pt 1):5-18.
2. Herpes 2006 May ;13(1):20-4.
3. Curr Opin Infect Dis 2001 Jun ;14(3):343-56.
4. N Engl J Med 2005 Feb 24;352(8):768-76.
5. Pediatr Neurol 2010 Jan ;42(1):28-31.
6. Epilepsy Res 2011 Sep ;96(1-2):89-95.
7. J Neurovirol 2009 Jan ;15(1):108-9.
8. Curr Infect Dis Rep 2008 Jul ;10(4):292-9.
9. Arch Pediatr 2007 May ;14(5):472-5.
10. Pediatr Neurol 2006 Feb ;34(2):160-3.
11. Emerg Infect Dis 2004 Sep ;10(9):1700-2.
12. Pediatr Neurol 2009 Nov ;41(5):353-8.
13. Pediatr Neurol 2010 Jan ;42(1):32-9.
14. Curr Infect Dis Rep 2004 Aug ;6(4):316-321.
15. AJR Am J Roentgenol 2010 Mar ;194(3):754-60.
16. Brain Nerve 2010 Aug ;62(8):869-75.
17. Clin Rheumatol 2009 Sep ;28(9):1067-71.
18. Pediatr Infect Dis J 2012 Nov ;31(11):1202-3.
19. J Infect 2007 Apr ;54(4):e237-40.
20. Invest Ophthalmol Vis Sci 2012 Jul ;53(8):4692-8.
21. Jpn J Ophthalmol 2011 Sep ;55(5):502-5.
22. Cornea 2011 Feb ;30(2):204-7.
23. Pediatr Med Chir 2012 Sep-Oct;34(5):229-33.
24. Br J Dermatol 2009 Jul ;161(1):181-3.
25. Arch Pediatr 2006 Dec ;13(12):1518-20.
26. Herpes 2007 Sep ;14(2):41-4.
27. Liver Transpl 2009 Oct ;15(10):1242-6.
28. Br J Radiol 2010 Dec ;83(996):e255-8.
29. J Pediatr Hematol Oncol 2007 Oct ;29(10):709-12.
30. Hum Pathol 2009 May ;40(5):740-5.
31. J Clin Virol 2011 Oct ;52(2):142-5.

Hymenolepis diminuta infection

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Hymenolepididae: Hymenolepis diminuta
Reservoir	Rodent Various insects
Vector	None
Vehicle	Arthropod - ingestion
Incubation Period	2w - 4w
Diagnostic Tests	Identification of ova in stool
Typical Adult Therapy	Praziquantel 25 mg/kg as single dose. OR Niclosamide 2g, then 1g/d X 6d
Typical Pediatric Therapy	Praziquantel 25 mg/kg as single dose. OR Niclosamide 1g, then 0.5g/d X 6d (1.5g, then 1g for weight >34kg)
Clinical Hints	Nausea, abdominal pain and diarrhea; eosinophilia may be present; primarily a pediatric disease, in rodent-infested areas; infestation resolves spontaneously within 2 months.
Synonyms	Hymenolepis diminuta, Mathevotaenia, Rat tapeworm. ICD9: 123.6 ICD10: B71.0

Clinical

Patients, usually children, may develop mild abdominal pain, nausea diarrhea and eosinophilia. ¹

This disease is endemic or potentially endemic to all countries.

References

1. [Pediatr Infect Dis J 1990 Mar ;9\(3\):216-9.](#)

Hymenolepis nana infection

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Hymenolepididae: Hymenolepis (Rodentolepis) nana
Reservoir	Human Rodent (especially hamster)
Vector	None
Vehicle	Food Water Fecal-oral
Incubation Period	2w - 4w
Diagnostic Tests	Identification of ova in stool
Typical Adult Therapy	Praziquantel 25 mg/kg once. OR Nitazoxanide 500 mg daily for 3 days OR Niclosamide 2g/d X 1, then 1g/d X 6d
Typical Pediatric Therapy	Praziquantel 25 mg/kg once. OR Nitazoxanide 100 mg (age 1 to 3 years) to 200 mg (age 4 to 11 years) BID X 3d OR Niclosamide 1g/d X 1, then 0.5g/d X 6d (1.5g, then 1g for weight >34kg)
Clinical Hints	Nausea, abdominal pain, diarrhea, irritability and weight loss; eosinophilia may be present; infection is maintained by autoinfection (worm reproduces within the intestinal lumen).
Synonyms	Dwarf tapeworm, Hymenolepis nana, Rodentolepis (Hymenolepis) microstoma, Rodentolepsiasis, Vampirolepis nana. ICD9: 123.6 ICD10: B71.0

Clinical

Infestation by *Hymenolepis nana* is largely asymptomatic. ¹

- Children are most likely to exhibit symptoms consisting of abdominal pain and diarrhea. ²
- Pruritis ani and behavioral and sleep disturbances are occasionally encountered. ³
- Most patients have eosinophilia (5% to 10% of total leucocyte count).

This disease is endemic or potentially endemic to all countries.

Hymenolepis nana infection in Uganda

Prevalence surveys:

- 11% of children below age 3 years living on the Ugandan shore of Lake Victoria (2006 publication) ⁴
- 0.16% of pregnant women in Entebbe (2004 to 2005) ⁵

References

1. [Trans R Soc Trop Med Hyg 2007 Feb ;101\(2\):203-5.](#)
2. [J Med Assoc Thai 2000 Sep ;83\(9\):1035-8.](#)
3. [Parasitol Res 2003 Nov ;91\(5\):412-38.](#)
4. [Ann Trop Med Parasitol 2006 Jun ;100\(4\):315-26.](#)
5. [PLoS Negl Trop Dis 2009 ;3\(6\):e473.](#)

Infection of wound, puncture, IV line, etc

Agent	BACTERIUM. <i>Staphylococcus aureus</i> , streptococci, facultative or aerobic gram negative bacilli, anaerobes, et al
Reservoir	Human Soil Water Air (spores) Various animals and plants
Vector	None
Vehicle	Trauma Water Medications Bandages Autoinoculation
Incubation Period	Variable
Diagnostic Tests	Smear and culture of catheter, material from wound.
Typical Adult Therapy	Drainage, remove catheter, debridement and antibiotics appropriate to infecting species
Typical Pediatric Therapy	As for adult
Clinical Hints	Source (ie, venous line, postoperative, marine, animal bite) may suggest species; onset less than 24 hrs = group A Strep. or <i>Cl. perfringens</i> ; 2 to 7 days <i>S. aureus</i> ; over 7 days gram negative bacilli; foul odor anaerobes.
Synonyms	Intravenous catheter infection, Line infection, Surgical wound infection, Wound infection. ICD9: 686.9,451 ICD10: T79.3,I80.0, Y95

Clinical

Wound infection is a self-defined illness.

The features and severity of infection are largely determined by the health status of the patient, and the nature of the wound and infecting organism.

Signs of infection which develop in a patient with an intravenous catheter should be assumed to be related to the catheter until proven otherwise.

This disease is endemic or potentially endemic to all countries.

Infectious mononucleosis or EBV infection

Agent	VIRUS - DNA. Herpesviridae. Gammaherpesvirinae, Lymphocryptovirus: Human herpesvirus 4 (Epstein Barr virus)
Reservoir	Human
Vector	None
Vehicle	Saliva Blood transfusion
Incubation Period	28d - 42d
Diagnostic Tests	Serology. Nucleic acid amplification.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Exudative pharyngitis, symmetrical cervical lymphadenopathy, splenomegaly and hepatic dysfunction; atypical lymphocytes and positive serology appear after 10 to 14 days; acute illness resolves in 2 to 3 weeks, but malaise and weakness may persist for months.
Synonyms	EBV, EBV, Epstein-Barr, Febbre ghiandolare, Glandular fever, Infectious mononucleosis, Monocytic angina, Mononucleose, Mononucleosi, Mononucleosis - infectious, Mononukleose, Pfeiffer's disease. ICD9: 075 ICD10: B27.0

Clinical

Symptoms of Infectious Mononucleosis (IM) usually consist of fever, pharyngitis, and lymphadenopathy. ¹

- Patients usually do not recall a history of possible exposure.
- A prodrome consisting of 1 to 2 weeks of fatigue, malaise, and myalgia is common; however, abrupt presentations may occur.
- A low-grade fever is usually present and lasts for 1 to 2 weeks, occasionally up to 5 weeks.
- CMV / EBV co-infection may be associated with prolonged illness. ²

Pharyngitis may be severe, particularly during the first week of illness. ³

- Tonsillitis may be present, and lymphadenopathy is almost universal, lasting for 1 to 2 weeks.
- Posterior cervical nodes are often affected, and generalized adenopathy may occur.
- Periorbital edema and palatal petechiae are often present.
- Splenomegaly is found in most cases, and hepatomegaly in 25%.
- Asymptomatic pericardial effusions are common. ⁴
- Patients often complain of headache.
- A morbilliform or papular erythematous eruption of the upper extremities or trunk is noted in 5% of cases.
- Lemmiere's syndrome has been reported as a complication of infectious mononucleosis. ⁵
- Guillain-Barre syndrome and membranous glomerulonephritis have been reported following primary EBV infection. ⁶

It is of note that a macular erythematous rash may occur in patients treated with ampicillin, usually appearing 5 to 9 days following the first dose.

- This phenomenon should not be misinterpreted as a penicillin allergy.
- Erythema nodosum and erythema multiforme have been associated with IM, as have petechiae and jaundice.
- The presence of severe abdominal pain may herald splenic rupture.

Other diseases ascribed to Epstein-Barr virus include nasopharyngeal carcinoma, Burkitt's lymphoma (African type) ⁷, post-transfusion lymphoproliferative disorder (PTLD) ⁸, hemophagocytic lymphohistiocytosis ⁹ and hemolytic anemia. ¹⁰

- Epstein-Barr virus infection, like many other infectious diseases, is occasionally followed by Guillain-Barre syndrome.
- Gianotti-Crosti syndrome may be the only presenting manifestation of Epstein-Barr virus infection. ¹¹

A false positive serological reaction toward Epstein-Barr virus has been associated with a variety of conditions, including rheumatoid arthritis ^{12 13}, Hepatitis E ¹⁴, Hepatitis A ¹⁵ and Parvovirus B19 infection. ¹⁶

This disease is endemic or potentially endemic to all countries.

References

1. Postgrad Med 2000 Jun ;107(7):175-9, 183-4, 186.
2. J Med Virol 2009 Aug ;81(8):1399-402.
3. Scand J Infect Dis Suppl 1991 ;80:94-104.
4. Indian Pediatr 2012 Mar 8;49(3):195-8.
5. Wien Klin Wochenschr 2008 ;120(5-6):181-3.
6. Arch Pediatr 2010 Nov ;17(11):1535-9.
7. Am J Trop Med Hyg 2011 Mar ;84(3):397-401.
8. Crit Rev Oncol Hematol 1989 ;9(2):149-95.
9. J Pediatr Hematol Oncol 2012 Mar ;34(2):e45-8.
10. Acta Haematol 1992 ;88(2-3):142-6.
11. Turk J Pediatr 2008 May-Jun;50(3):302-4.
12. Ann Rheum Dis 1985 Nov ;44(11):742-6.
13. Clin Exp Immunol 1979 Jun ;36(3):415-22.
14. Eur J Gastroenterol Hepatol 2009 Dec ;21(12):1433-5.
15. Pediatr Infect Dis J 1994 May ;13(5):413-4.
16. Clin Vaccine Immunol 2009 Mar ;16(3):372-5.

Influenza

Agent	VIRUS - RNA. Orthomyxoviridae, Orthomyxovirus: Influenza virus
Reservoir	Human Occasionally Ferret Bird Pig
Vector	None
Vehicle	Droplet
Incubation Period	1d - 3d
Diagnostic Tests	Viral culture (respiratory secretions). Serology. Nucleic acid amplification techniques are available.
Typical Adult Therapy	Respiratory precautions. Influenza A or B: Oseltamivir 75 mg PO BID X 5d OR Zanamavir 10 mg BID X 5 days
Typical Pediatric Therapy	Respiratory precautions. Influenza A or B: Oseltamivir 2 mg/kg (max 75 mg) PO BID X 5d OR Zanamavir (age > 5 years) 10 mg BID X 5 days
Vaccines	Influenza - inactivated Influenza - live
Clinical Hints	Myalgia, headache, cough, fever; pharyngitis and conjunctivitis often present; usually encountered in the setting of an outbreak; leucocytosis, chest pain and lobar infiltrate herald bacterial (pneumococcal or staphylococcal) pneumonia.
Synonyms	Asian flu, Aviaire influenza, Avian flu, Avian influenza, Bird flu, Epidemic catarrh, Grippe, H1N1, H2N2, H3N2, H5N1, Hong Kong flu, LPAI, Spanish influenza, Swine flu, Swine influenza. ICD9: 487 ICD10: J09,J10,J11

Clinical

Influenza is characterized by acute onset of fever, headache, myalgia, nonproductive cough, sore throat, and rhinitis. ¹

- The illness usually resolves in 2 to 7 days; however, symptoms often persist for up to two weeks.
- Severe illness or death may complicate the acute infection, notably in pregnant women ², the elderly and patients with underlying medical conditions. ³
- Complications include primary viral pneumonia or bacterial pneumonia (most commonly pneumococcal) ⁴; myocarditis, myositis, Guillain-Barre syndrome ⁵, encephalitis ⁶, Gianotti-Crosti syndrome ⁷ and transverse myelitis. ⁸⁻¹⁰

WHO Case definition for surveillance • Influenza:

Clinical case definition

A person with sudden onset of fever of >38°C and cough or sore throat in the absence of other diagnoses.

Laboratory criteria for diagnosis

- Virus isolation: Swab or aspirate from the suspected individual, or
- Direct detection of influenza viral antigen.
- Serology: Fourfold rise in antibody titer between early and late serum.

Case classification

- Suspected: A case that meets the clinical case definition.
- Confirmed: A case that meets the clinical case definition and is laboratory-confirmed (used mainly in epidemiological investigation rather than surveillance).

WHO definition for surveillance • Swine influenza (H1H1):

confirmed case • person with swine influenza A (H1N1) virus infection laboratory confirmed by

- real-time RT-PCR and/or
- viral culture and/or
- 4-fold rise in swine influenza A(H1N1) virus specific neutralizing antibodies

probable case • either

- person with influenza test positive for influenza A, but unsubtypeable by reagents used to detect seasonal influenza virus infection, or
- person with clinically compatible illness or who died of unexplained acute respiratory illness who is considered to be epidemiologically linked to probable or confirmed case

CDC definition for surveillance • Swine influenza (H1H1):

confirmed case • person with acute respiratory illness with swine influenza A (H1N1) virus infection laboratory confirmed at

CDC by

- real-time reverse transcriptase polymerase chain reaction (RT-PCR) and/or
- viral culture

probable case • person with acute febrile respiratory illness who is

- positive for influenza A, but negative for H1 and H3 by influenza RT-PCR
- positive for influenza A by influenza rapid test or influenza immunofluorescence assay (IFA) plus meets criteria for suspected case

suspected case • person with acute respiratory illness (defined as recent onset of ≥ 2 of rhinorrhea or nasal congestion, sore throat, or cough) plus

- close contact to confirmed case of swine influenza A (H1N1) virus infection during case's infectious period, or
- close contact defined as within about 6 feet of ill person
- infectious period defined as 1 day prior to illness onset to 7 days after onset
- travel to or residence in area with confirmed cases of swine influenza A (H1N1) virus infection

Avian influenza H5N1 infection:

Avian influenza H5N1 infection is characterized by fever greater than 38 C, shortness of breath and cough. ^{11 12}

- The incubation period is 2 to 4 days.
- All patients reported to date have presented with significant lymphopenia and marked chest radiograph abnormalities consisting of diffuse, multifocal or patchy infiltrates.
- Some cases showed segmental or lobular consolidation with air bronchograms.
- Crackles were frequently heard on auscultation.
- Some of the patients reported sore throat, conjunctivitis, myalgia, rash or rhinorrhea.
- Watery diarrhea or loose stools was noted in approximately 50% of the cases.
- Myocardial dysfunction and hepatic dysfunction are also reported.
- Reactive hemophagocytic syndrome is the most characteristic pathological finding and may contribute to the lymphopenia, liver dysfunction, and abnormal clotting profiles observed among patients with severe infection.
- Approximately 90% of patients with H5N1 infection have been below age 40. ¹³
- Approximately 60% of patients have died, on an average of 10 days after onset of symptoms.

Influenza virus H1N1 infection

- During the "Spanish flu" H1N1 pandemic of 1918 to 1919, illness was characterized by unusual severity, tendency to affect young healthy adults, rapid progression and overwhelming pneumonia.
- During the outbreak of A(H1N1)pdm09 virus outbreak of 2009 to 2010 ^{14 15}, children ^{16 17} and young adults accounted for a large proportion of cases. ^{18 19} Severe cases were not necessarily associated with underlying disease. Obesity ²⁰⁻²⁹, immune-compromise ³⁰ (but not necessarily AIDS ³¹), pregnancy ³²⁻⁵³, infection while hospitalized ⁵⁴, preexisting neurological disorders ⁵⁵, sickle cell disease ⁵⁶ and asthma were identified as risk factors for complications. ⁵⁷⁻⁶³ Children below age 5 years, particularly those with neuro-developmental disorders, were also found to be at risk. ⁶⁴⁻⁶⁶
- Most deaths were caused by primary viral pneumonia ⁶⁷⁻⁷⁷, and bacterial co-infection was identified in as many as 29% of fatal cases. ⁷⁸⁻⁸³
- Vomiting and diarrhea were reported in up to 25% of patients ⁸⁴, and as many as 6% were afebrile. ⁸⁵ Case-fatality rates were not necessarily higher than those reported for other strains of Influenza virus. ^{86 87}
- Additional complications included myopathy or rhabdomyolysis ⁸⁸⁻⁹⁴, encephalitis or encephalopathy ⁹⁵⁻¹²⁰, ischemic stroke ¹²¹, aseptic meningitis ¹²², acute disseminated or hemorrhagic leukoencephalitis ¹²³⁻¹³², deafness ¹³³, cerebellitis ¹³⁴, acute myelopathy ¹³⁵, Guillain-Barre syndrome ¹³⁶⁻¹³⁹, parkinsonism ¹⁴⁰, narcolepsy ^{141 142}, quadriplegia ¹⁴³, glomerulonephritis ^{144 145}, tubulointerstitial nephritis ¹⁴⁶, renal failure ¹⁴⁷⁻¹⁵⁶, hemolytic-uremic syndrome ¹⁵⁷⁻¹⁶¹, reactive thrombocytosis ¹⁶², hemophagocytic lymphohistiocytosis ¹⁶³⁻¹⁶⁸, myopathy ¹⁶⁹, cold agglutinin syndrome ¹⁷⁰, autoimmune ¹⁷¹ and thrombotic thrombocytopenic purpura ¹⁷², myocarditis ¹⁷³⁻¹⁹³ or reversible myocardial dysfunction ¹⁹⁴⁻¹⁹⁶, pericarditis ^{197 198}, subacute thyroiditis ¹⁹⁹, rash ²⁰⁰, pancreatitis ²⁰¹, vascular thrombosis ²⁰², plastic bronchitis ²⁰³, hemorrhagic pneumonia ²⁰⁴ and Acute Respiratory Distress Syndrome (ARDS). ²⁰⁵⁻²¹⁹
- In some cases, the clinical features of leptospirosis suggested a diagnosis of H1N1 influenza. ²²⁰

This disease is endemic or potentially endemic to all countries.

Influenza in Uganda

GIDEON does not follow routine country reports on human Influenza, since the scope and nature of these data are often diffuse, sporadic or inconsistent. See the "Worldwide" note for material regarding pandemic influenza, influenza vaccine, avian influenza in humans and other relevant subjects.

Notable outbreaks:

2009 to 2010 - An outbreak (263 cases) was reported. **Context:** A pandemic of H1N1 Influenza virus [A (H1N1) pdm09 ²²¹ infection occurred. ²²²⁻³¹⁹ Over 600,000 cases had been officially-reported worldwide as of March, 2010. ³²⁰⁻³²² 18,449 fatal cases were reported to August 1, 2010 (true number for first 12 months estimated at 293,500 ^{323 324}

). ³²⁵⁻³⁴⁹ Indigenous populations from Australia, Canada, the United States and New Zealand were found to have at least a 3-fold greater death rate than others in their countries. ³⁵⁰⁻³⁶⁹ Reporting of case-number summaries was suspended by WHO as of July 6 ³⁷⁰; and on August 10, the pandemic was declared to have ended. ^{371 372} The pandemic began in Mexico, spreading rapidly to the United States and Canada. Swine were not implicated in the transmission of disease. ³⁷³⁻³⁷⁵ Human-to-swine transmission was confirmed in Argentina ³⁷⁶, Cambodia ^{377 378}, Vietnam ^{379 380}, Italy ³⁸¹ and Canada during the outbreak ³⁸²⁻³⁹³; and infected swine were identified in Argentina ³⁹⁴⁻³⁹⁶, Australia ³⁹⁷⁻³⁹⁹, Brazil ⁴⁰⁰, Cameroon ^{401 402}, China ⁴⁰³⁻⁴⁰⁶, Denmark ⁴⁰⁷, Finland ⁴⁰⁸, Germany ^{409 410}, Iceland ⁴¹¹, India ⁴¹², Indonesia ⁴¹³, Ireland ⁴¹⁴, Italy ^{415 416}, Japan ^{417 418}, England ⁴¹⁹, Mexico ⁴²⁰, Northern Ireland ⁴²¹, Norway ⁴²², Republic of Korea ^{424 425}, Reunion Island ⁴²⁶, Russian Federation ⁴²⁷, Scotland ⁴²⁸, Taiwan ⁴²⁹, Thailand ⁴³⁰, ⁴³¹, the United Kingdom ^{432 433} and the United States. ⁴³⁴⁻⁴⁴² Infected turkeys were subsequently identified in Canada ⁴⁴³⁻⁴⁴⁵, Chile ⁴⁴⁶⁻⁴⁵⁰, France ⁴⁵¹, the United Kingdom ^{452 453} and the United States. ^{454 455} Infection was reported in cats ⁴⁵⁶⁻⁴⁶⁸, ferrets ⁴⁶⁹⁻⁴⁷⁴, a dog ⁴⁷⁵, a badger (*Taxidea taxus*), a captive Bornean binturong (*Arctictis binturong penicillata*) ^{476 477} and a cheetah ^{478 479} in the United States ⁴⁸⁰⁻⁴⁸⁶; skunks in Canada ⁴⁸⁷; dogs in Italy ⁴⁸⁸ and China ⁴⁸⁹; farmed American mink (*Neovison vison*) in the Netherlands ⁴⁹⁰; and in dogs and swine in Hong Kong. ⁴⁹¹

- Reporting dates vary by country. The following updates include incidence data as of December 31, 2010. ^{492 493}:

Afghanistan (17 fatal) ^{494 495}, Albania (6 fatal), Algeria (57 fatal cases), American Samoa (94 - 0 fatal), Andorra (1), Angola (37) ⁴⁹⁶, Anguilla (14), Antigua and Barbuda (0 fatal), Argentina (626 fatal) ⁴⁹⁷⁻⁵¹⁸, Armenia (3 fatal), Aruba (13), Australia (51,170 - 195 fatal) ⁵¹⁹⁻⁵⁶⁵, Austria (24 fatal) ⁵⁶⁶⁻⁵⁶⁸, Azerbaijan (2), Bahamas (4 fatal), Bahrain (7 fatal), Bangladesh (7 fatal) ⁵⁶⁹, Barbados (157 - 3 fatal) ⁵⁷⁰, Belarus (20 fatal) ⁵⁷¹, Belgium (17 fatal) ⁵⁷²⁻⁵⁷⁵, Belize (60), Bermuda (1 fatal), Bhutan (487) ⁵⁷⁶, Bolivia (59 fatal) ⁵⁷⁷, Bosnia and Herzegovina (10 fatal), Botswana (23), Brazil (2,125 fatal) ⁵⁷⁸⁻⁵⁸⁸, British Virgin Islands (25), Brunei (850 - 1 fatal), Bulgaria (40 fatal), Burundi (7), Cambodia (6 fatal) ⁵⁸⁹⁻⁵⁹¹, Cameroon (4) ⁵⁹², Canada (429 fatal) ⁵⁹³⁻⁶²⁴, Cape Verde (118), Cayman Islands (130 - 1 fatal), Chad (1), Chile (156 fatal) ⁶²⁵⁻⁶⁴⁰, Central African Republic ⁶⁴¹, China (724 fatal - including 56 in Hong Kong ⁶⁴²⁻⁶⁶⁶ and 2 in Macao) ⁶⁶⁷⁻⁷²², Colombia (254 fatal) ⁷²³, Comoros (2 fatal in Mayotte) ⁷²⁴, Congo (21) ⁷²⁵, Cook Islands (106 - 1 fatal), Costa Rica (65 fatal), Croatia (25 fatal), Cuba (1,805 - 83 fatal) ⁷²⁶, Cyprus (6 fatal) ⁷²⁷, Czech Republic (98 fatal), Democratic Republic of Congo (222) ^{728 729}, Democratic Republic of Korea (9) ⁷³⁰, Denmark (32 fatal) ⁷³¹⁻⁷³⁵, Dominica (51), Dominican Republic (464 - 24 fatal), Ecuador (130 fatal) ⁷³⁶, Egypt (281 fatal) ⁷³⁷⁻⁷⁴⁰, El Salvador (34 fatal) ⁷⁴¹, Estonia (19 fatal), Ethiopia (12), Falkland Islands (7), Fiji (268 - 0 fatal), Finland (43 fatal) ⁷⁴²⁻⁷⁴⁵, France (349 fatal) ⁷⁴⁶⁻⁷⁷⁸, French Guiana (29 - 1 fatal) ^{779 780}, French Polynesia (185 - 7 fatal) ^{781 782}, Gabon (4), Georgia (20 fatal), Germany (253 fatal) ⁷⁸³⁻⁸⁰¹, Ghana (1 fatal) ⁸⁰², Gibraltar (16), Greece (141 fatal) ⁸⁰³⁻⁸¹², Grenada (28), Guadeloupe (5 fatal) ⁸¹³, Guam (341 - 2 fatal) ⁸¹⁴, Guatemala (26 fatal) ^{815 816}, Guyana (30), Haiti (95) ^{817 818}, Honduras (18 fatal), Hong Kong (56 fatal) ⁸¹⁹⁻⁸⁴³, Hungary (133 fatal) ⁸⁴⁴, Iceland (2 fatal) ^{845 846}, India (44,958 - 2,703 fatal) ⁸⁴⁷⁻⁸⁶⁴, Indonesia (691 - 10 fatal) ^{865 866}, Iran (147 fatal) ⁸⁶⁷⁻⁸⁷⁵, Iraq (42 fatal) ⁸⁷⁶, Ireland (24 fatal) ⁸⁷⁷⁻⁸⁷⁹, Israel (113 fatal, including 28 in Gaza and the West Bank) ⁸⁸⁰⁻⁸⁹³, Italy (256 fatal) ⁸⁹⁴⁻⁹⁰⁶, Ivory Coast (5) ⁹⁰⁷, Jamaica (7 fatal), Japan (198 fatal) ⁹⁰⁸⁻⁹³⁶, Jordan (19 fatal) ⁹³⁷, Kazakhstan (17) ⁹³⁸, Kenya (417) ⁹³⁹⁻⁹⁴², Kiribati (4 - 0 fatal), Kuwait (30 fatal), Laos (156 - 1 fatal) ⁹⁴³⁻⁹⁴⁵, Latvia (34 fatal), Lebanon (5 fatal) ^{946 947}, Lesotho (65), Libya (1 fatal), Liechtenstein (5), Lithuania (23 fatal) ⁹⁴⁸, Luxembourg (3 fatal) ⁹⁴⁹, Macao (2 fatal) ⁹⁵⁰, Macedonia (23 fatal), Madagascar (3 fatal) ⁹⁵¹⁻⁹⁵⁶, Malaysia (1,780 - 77 fatal) ⁹⁵⁷⁻⁹⁶⁴, Malawi (4), Maldives (1 fatal), Mali (12) ⁹⁶⁵, Malta (5 fatal), Marshall Islands (115 - 1 fatal), Martinique (44 - 1 fatal) ⁹⁶⁶, Mauritania (15), Mauritius (8 fatal), Mexico (1,316 fatal) ⁹⁶⁷⁻¹⁰⁰⁹, Micronesia (82 - 0 fatal), Moldova (35 fatal) ¹⁰¹⁰, Monaco (1), Mongolia (29 fatal) ^{1011 1012}, Montenegro (7 fatal), Montserrat (21), Morocco (64 fatal) ¹⁰¹³⁻¹⁰¹⁵, Mozambique (2 fatal), Myanmar (137) ¹⁰¹⁶⁻¹⁰²¹, Namibia (1 fatal), Nauru (8 - 0 fatal), Nepal (172 - 3 fatal) ¹⁰²²⁻¹⁰²⁵, The Netherlands (62 fatal) ¹⁰²⁶⁻¹⁰³⁹, Netherlands Antilles (128 cases - 59 in Curacao, including 3 on a cruise ship; 29 in St. Maarten and 38 on Bonaire), New Caledonia (508 - 7 fatal), New Zealand (4,974 - 51 fatal) ¹⁰⁴⁰⁻¹⁰⁶⁶, Nicaragua (2,152 cases - 11 fatal) ¹⁰⁶⁷, Niger (12), Nigeria (2 fatal) ^{1068 1069}, Niue (0), Northern Marianas (6 - 0 fatal), Norway (12,513 cases, 29 fatal) ¹⁰⁷⁰⁻¹⁰⁷³, Oman (31 fatal) ^{1074 1075}, Pakistan (14 fatal) ¹⁰⁷⁶⁻¹⁰⁷⁹, Palau (47 - 0 fatal), Panama (12 fatal) ^{1080 1081}, Papua New Guinea (29 - 0 fatal), Paraguay (47 fatal), Peru (238 fatal) ¹⁰⁸²⁻¹⁰⁸⁶, Philippines (3,207 - 30 fatal), Pitcairn Island (0), Poland (148 fatal) ^{1087 1088}, Portugal (83 fatal) ¹⁰⁸⁹⁻¹⁰⁹², Puerto Rico (20), Qatar (8 fatal) ¹⁰⁹³, Republic of Korea (171 fatal) ¹⁰⁹⁴⁻¹¹⁰⁷, Reunion (12 fatal) ¹¹⁰⁸⁻¹¹²², Romania (122 fatal) ¹¹²³, Russian Federation (19 fatal) ¹¹²⁴⁻¹¹³³, Rwanda (494 - 0 fatal) ¹¹³⁴⁻¹¹³⁶, Saint Kitts and Nevis (2 fatal), Saint Lucia (1 fatal), Saint Vincent and the Grenadines (19), Samoa (173 - 2 fatal), Sao Tome and Principe (2 fatal), Saudi Arabia (124 fatal) ¹¹³⁷⁻¹¹⁵², Scotland (38 fatal) ¹¹⁵³⁻¹¹⁶⁶, Senegal (184), Serbia (71 fatal) ^{1167 1168}, Seychelles (33), Singapore (19 fatal) ¹¹⁶⁹⁻¹¹⁸⁹, Slovakia (53 fatal), Slovenia (19 fatal) ¹¹⁹⁰, Solomon Islands (4 - 0 fatal), South Africa (93 fatal) ¹¹⁹¹⁻¹¹⁹⁵, Spain (271 fatal) ¹¹⁹⁶⁻¹²³³, Sri Lanka (48 fatal), Sudan (5 fatal), Suriname (144 - 2 fatal) ¹²³⁴, Swaziland (2), Sweden (25 fatal) ¹²³⁵⁻¹²³⁸, Switzerland (18 fatal) ¹²³⁹⁻¹²⁴², Syria (127 fatal) ^{1243 1244}, Taiwan (44 fatal) ¹²⁴⁵⁻¹²⁵⁸, Tanzania (1 fatal), Thailand (212 fatal) ¹²⁵⁹⁻¹²⁷⁷, Tokelau (0), Tonga (20 - 1 fatal), Trinidad and Tobago (5 fatal), Tunisia (21 fatal), Turkey (627 fatal) ¹²⁷⁸⁻¹²⁸⁷, Turks and Caicos Islands (45), Tuvalu (23 - 0 confirmed) ¹²⁸⁸, Uganda (263), Ukraine (282 fatal) ¹²⁸⁹⁻¹²⁹⁸, United Arab Emirates (6 fatal) ¹²⁹⁹⁻¹³⁰², United Kingdom (474 fatal: at least 142 in England, 38 in Scotland - including the first fatal case in Europe ¹³⁰³⁻¹³⁰⁵, 21 in Wales and 13 in Northern Ireland) ¹³⁰⁶⁻¹³⁵⁵, United

States (2,718 fatal) ¹³⁵⁶⁻¹⁴³², Uruguay (20 fatal), Vanuatu (3 - 0 fatal), Venezuela (137 fatal) ^{1433 1434}, Vietnam (53 fatal) ¹⁴³⁵⁻¹⁴⁴¹, Virgin Islands, U.S. (49), Wallis and Futuna (55 - 0 fatal) ¹⁴⁴², Yemen (28 fatal), Zambia (90) ¹⁴⁴³ and Zimbabwe (41). ¹⁴⁴⁴⁻¹⁵⁰²

References

1. JAMA 2005 Feb 23;293(8):987-97.
2. Emerg Infect Dis 2008 Jan ;14(1):95-100.
3. Lancet 2003 Nov 22;362(9397):1733-45.
4. Int J Infect Dis 2012 Mar 1;
5. Clin Infect Dis 2009 Jan 1;48(1):48-56.
6. Curr Opin Neurol 2010 Jun ;23(3):305-11.
7. Pediatr Dermatol 2011 Nov-Dec;28(6):733-5.
8. Travel Med Infect Dis 2008 May ;6(3):114-24.
9. Pediatr Infect Dis J 2008 May ;27(5):384-9.
10. Pediatr Infect Dis J 2008 May ;27(5):390-5.
11. N Engl J Med 2005 Sep 29;353(13):1374-85.
12. Lancet 2008 Apr 26;371(9622):1464-75.
13. ProMED <promedmail.org> archive: 20070211.0522
14. Infect Dis Clin North Am 2010 Mar ;24(1):203-28.
15. Int J Infect Dis 2012 Oct ;16(10):e714-23.
16. Pediatr Infect Dis J 2010 Aug ;29(8):694-8.
17. Clin Pediatr (Phila) 2010 Aug ;49(8):777-81.
18. N Engl J Med 2009 Nov 12;361(20):2000-1.
19. Infect Control Hosp Epidemiol 2010 Jul ;31(7):676-82.
20. Clin Infect Dis 2011 Feb 1;52(3):301-12.
21. Lancet Infect Dis 2011 Jan ;11(1):14-5.
22. PLoS One 2010 ;5(3):e9694.
23. Chest 2011 Feb ;139(2):382-6.
24. Obes Rev 2011 Aug ;12(8):653-9.
25. ProMED <promedmail.org> archive: 20090711.2482
26. Clin Infect Dis 2011 Sep ;53(5):413-21.
27. Rev Pneumol Clin 2011 Sep ;67(4):244-9.
28. Influenza Other Respi Viruses 2012 Feb 15;
29. Pulm Pharmacol Ther 2012 May 23;
30. Crit Care 2010 ;14(2):139.
31. Clin Infect Dis 2011 Jan 15;52(2):219-27.
32. Rev Med Virol 2012 Mar 13;
33. J Obstet Gynaecol 2010 Feb ;30(2):97-100.
34. BMJ 2010 ;340:c1279.
35. BMJ 2010 ;340:c1235.
36. Obstet Gynecol 2010 Apr ;115(4):717-26.
37. Crit Care Med 2010 Apr ;38(4 Suppl):e52-7.
38. Obstet Gynecol 2010 Feb ;115(2 Pt 2):412-4.
39. Obstet Gynecol 2010 Feb ;115(2 Pt 2):409-12.
40. Expert Rev Respir Med 2010 Jun ;4(3):321-8.
41. Int J Obstet Anesth 2010 Oct ;19(4):465-6.
42. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S13-20.
43. Am J Obstet Gynecol 2011 Jul ;205(1):10-8.
44. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S38-45.
45. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S1-3.
46. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S38-45.
47. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S89-93.
48. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S13-20.
49. BMJ 2011 ;342:d3237.
50. BMJ 2011 ;342:d3214.
51. Obstet Gynecol 2011 Sep ;118(3):593-600.
52. Influenza Other Respi Viruses 2012 Feb 15;
53. J Infect Dis 2012 Oct ;206(8):1260-8.
54. J Infect Dis 2011 Jun 15;203(12):1739-47.
55. Acta Paediatr 2011 Jul ;100(7):e12-6.
56. Pediatr Blood Cancer 2011 Jan ;56(1):95-8.
57. Wkly Epidemiol Rec 2009 Jul 24;84(30):305-8.
58. MMWR Morb Mortal Wkly Rep 2009 Jul 24;58(28):773-8.
59. Euro Surveill 2009 ;14(33)
60. CMAJ 2010 Jan 12;182(1):39-44.
61. Clin Infect Dis 2010 Mar 1;50(5):686-90.
62. Hosp Pract (Minneap) 2010 Apr ;38(2):74-81.
63. ProMED <promedmail.org> archive: 20090619.2260
64. MMWR Morb Mortal Wkly Rep 2009 Sep 4;58(34):941-7.
65. Radiology 2010 Mar ;254(3):934-41.
66. Pediatr Crit Care Med 2010 Mar ;11(2):173-8.
67. N Engl J Med 2010 May 6;362(18):1708-19.
68. BMJ 2009 ;339:b4313.
69. Am J Respir Crit Care Med 2010 Jan 1;181(1):72-9.
70. N Engl J Med 2009 Nov 12;361(20):2001-3.
71. Crit Care Med 2010 Apr ;38(4 Suppl):e91-7.
72. AJR Am J Roentgenol 2009 Dec ;193(6):1494-9.
73. AJR Am J Roentgenol 2009 Dec ;193(6):1488-93.
74. J Clin Virol 2010 Feb ;47(2):199-200.
75. Diagn Interv Radiol 2011 Sep ;17(3):216-22.
76. Eur Respir J 2011 Jul ;38(1):106-11.
77. ProMED <promedmail.org> archive: 20091208.4188
78. MMWR Morb Mortal Wkly Rep 2009 Oct 2;58(38):1071-4.
79. PLoS One 2009 ;4(12):e8540.
80. Infection 2010 Jun ;38(3):237-9.
81. Thorax 2011 Mar ;66(3):247-52.
82. ProMED <promedmail.org> archive: 20090930.3410
83. ProMED <promedmail.org> archive: 20091017.3577
84. Gut 2009 Nov ;58(11):1567-8.
85. Wkly Epidemiol Rec 2009 May 22;84(21):185-9.
86. ProMED <promedmail.org> archive: 20090425.1552
87. ProMED <promedmail.org> archive: 20090425.1557
88. Pediatr Infect Dis J 2009 Dec ;28(12):1138-9.
89. Gac Med Mex 2009 Nov-Dec;145(6):519-21.
90. Am J Kidney Dis 2010 Mar ;55(3):615.
91. Emerg Infect Dis 2010 Mar ;16(3):565.
92. Neth J Med 2010 Aug ;68(1):317-9.
93. Arq Neuropsiquiatr 2012 May ;70(5):316-8.
94. Pediatr Int 2012 Oct ;54(5):703-5.
95. Pediatr Radiol 2010 Feb ;40(2):200-5.
96. Emerg Infect Dis 2010 Nov ;16(11):1811-3.
97. Emerg Infect Dis 2010 Jan ;16(1):139-42.
98. Lancet Neurol 2010 Feb ;9(2):142-3.
99. AJNR Am J Neuroradiol 2010 Mar ;31(3):396-400.
100. Neurology 2010 Mar 30;74(13):1020-1.
101. Curr Opin Neurol 2010 Jun ;23(3):305-11.
102. Neurology 2010 Aug 17;75(7):666-7.
103. Ann Neurol 2010 Nov ;68(5):762-6.
104. J Neurol Sci 2010 Nov 15;298(1-2):110-3.
105. Arch Neurol 2011 Apr ;68(4):455-62.
106. Rev Chilena Infectol 2010 Oct ;27(5):413-6.
107. Eur J Pediatr 2011 Aug ;170(8):1007-15.
108. J Child Neurol 2011 Apr ;26(4):501-6.
109. Acta Neuropathol 2011 Sep ;122(3):371-3.
110. J Neuroophthalmol 2012 Mar ;32(1):48-50.
111. Influenza Other Respi Viruses 2010 May 1;4(3):117-20.
112. J Clin Microbiol 2010 Oct ;48(10):3803-5.
113. Neurocrit Care 2010 Dec ;13(3):396-406.
114. Neurology 2012 Oct 2;79(14):1474-1481.
115. Neurology 2012 Oct 2;79(14):1479.
116. Pediatr Emerg Care 2012 Oct ;28(10):998-1002.
117. J Clin Virol 2012 Oct 26;
118. ProMED <promedmail.org> archive: 20091129.4089
119. Pediatrics 2012 Mar 12;
120. Scand J Infect Dis 2012 Dec ;44(12):941-947.
121. Acta Anaesthesiol Taiwan 2012 Mar ;50(1):41-5.
122. Turk J Pediatr 2011 Jan-Feb;53(1):91-3.
123. Arch Neurol 2010 Jun ;67(6):756-8.
124. Pediatr Infect Dis J 2011 Feb ;30(2):165-7.
125. J Neurol 2011 Mar ;258(3):513-4.
126. Biomed Environ Sci 2010 Aug ;23(4):323-6.
127. Eur J Pediatr 2011 Jun ;170(6):779-88.
128. Neuro Sci 2011 Oct ;32(5):907-9.
129. J Pediatr Neurosci 2010 Jul ;5(2):157-9.
130. J Neuropsychiatry Clin Neurosci 2011 ;23(2):E7-8.
131. Rev Clin Esp 2012 Apr ;212(4):e28-31.
132. Pediatr Neurol 2012 Jul ;47(1):62-4.
133. J Laryngol Otol 2012 Dec ;126(12):1274-5.
134. PMID 20972135
135. J Neurol 2011 Jan ;258(1):34-6.
136. Neurocrit Care 2010 Aug ;13(1):98-100.
137. Intensive Care Med 2010 Jul ;36(7):1275.
138. Curr Neurol Neurosci Rep 2010 Nov ;10(6):476-83.
139. Nervenarzt 2012 Apr 14;
140. J Neuroinflammation 2010 ;7:47.
141. ProMED <promedmail.org> archive: 20110823.2561
142. Ann Neurol 2011 Sep ;70(3):410-7.
143. Emerg Infect Dis 2010 Mar ;16(3):569-70.
144. Pediatr Nephrol 2011 Jan ;26(1):151-2.
145. Pediatr Nephrol 2011 Jan ;26(1):153-4.
146. Pediatr Nephrol 2012 Oct ;27(10):1985-7.
147. Clin Microbiol Infect 2010 Apr ;16(4):330-2.
148. Contrib Nephrol 2010 ;165:310-4.
149. Clin J Am Soc Nephrol 2010 Nov ;5(11):1916-21.
150. Intensive Care Med 2011 May ;37(5):763-7.
151. Intensive Care Med 2011 May ;37(5):768-74.
152. Intensive Care Med 2011 May ;37(5):880-1.
153. Intensive Care Med 2011 May ;37(5):729-31.
154. J Crit Care 2011 Dec ;26(6):577-85.
155. J Ren Care 2011 Sep ;37(3):128-33.
156. Eur J Pediatr 2012 Oct 13;
157. Pediatr Nephrol 2011 Jan ;26(1):147-8.
158. Pediatr Nephrol 2011 Jan ;26(1):149-50.
159. Pediatr Nephrol 2011 Jan ;26(1):143-4.
160. Pediatr Nephrol 2011 Jan ;26(1):3-5.
161. Clin Exp Nephrol 2011 Dec ;15(6):948-52.
162. J Lab Physicians 2011 Jul ;3(2):131-2.
163. J Pediatr Hematol Oncol 2011 Mar ;33(2):135-7.
164. Case Report Med 2011 ;2011:951910.
165. J Med Case Rep 2011 ;5:280.
166. Ann Diagn Pathol 2012 Dec ;16(6):477-84.
167. Ann Saudi Med 2012 Jan-Feb;32(1):86-9.
168. J Infect Chemother 2012 Oct ;18(5):771-4.

169. Trop Doct 2010 Oct ;40(4):242-3.
170. Am J Med 2011 Feb ;124(2):e1-2.
171. J Chin Med Assoc 2011 Sep ;74(9):425-7.
172. J Clin Virol 2011 Jun ;51(2):146-7.
173. J Am Coll Cardiol 2010 Mar 2;55(9):928-9.
174. J Clin Microbiol 2010 Jun ;48(6):2298-300.
175. Cardiol Young 2010 Jun ;20(3):351-2.
176. Clin Microbiol Infect 2011 Apr ;17(4):572-4.
177. Hellenic J Cardiol 2010 Jul-Aug;51(4):379-80.
178. Circ J 2010 Oct ;74(10):2193-9.
179. Virchows Arch 2011 Mar ;458(3):371-6.
180. Rev Assoc Med Bras 2010 Jul-Aug;56(4):394.
181. Heart Lung 2010 Sep-Oct;39(5):453-8.
182. Eur Heart J 2011 Feb ;32(3):283.
183. Circ J 2010 Nov ;74(12):2758-9.
184. Rev Esp Cardiol 2010 Nov ;63(11):1386-7.
185. J Formos Med Assoc 2010 Nov ;109(11):857-8.
186. Heart Lung 2011 Nov-Dec;40(6):566-8.
187. J Chin Med Assoc 2011 Mar ;74(3):130-3.
188. ProMED <promedmail.org> archive: 20100215.0526
189. Chest 2011 Jun ;139(6):1545-6; author reply 1546.
190. Can J Cardiol 2011 Jul-Aug;27(4):514-22.
191. Heart 2011 Oct ;97(19):1630.
192. Cardiol J 2011 ;18(5):552-5.
193. Rev Port Cardiol 2012 Jul-Aug;31(7-8):517-20.
194. Chest 2010 May ;137(5):1195-7.
195. Eur Respir J 2011 Jan ;37(1):112-8.
196. Am J Cardiol 2010 Nov 15;106(10):1517-9.
197. Korean Circ J 2010 Oct ;40(10):539-42.
198. Scand J Infect Dis 2012 Jun ;44(6):470-2.
199. Endocrine 2010 Jun ;37(3):440-1.
200. Influenza Other Respi Viruses 2011 Jul ;5(4):223-4.
201. Isr Med Assoc J 2010 Oct ;12(10):640-1.
202. Clin Infect Dis 2011 Jan 15;52(2):e14-7.
203. Chest 2010 Dec ;138(6):1486-8.
204. Influenza Other Respi Viruses 2012 Dec 20;
205. Curr Opin Crit Care Med 2011 Feb ;17(1):64-71.
206. Am J Respir Crit Care Med 2010 Sep 15;182(6):857; author reply 857-8.
207. Int J Obstet Anesth 2010 Oct ;19(4):465-6.
208. Am J Med Sci 2010 Dec ;340(6):499-504.
209. Minerva Anestesiol 2009 Dec ;75(12):741-5.
210. Ann Fr Anesth Reanim 2010 Feb ;29(2):165-6.
211. Am J Med 2010 Mar ;123(3):282-285.e2.
212. Acad Radiol 2010 Jun ;17(6):681-5.
213. J Bras Pneumol 2010 May-Jun;36(3):386-9.
214. South Med J 2010 Aug ;103(8):786-90; quiz 791-2.
215. Ned Tijdschr Geneesk 2010 ;154:A1640.
216. Eur J Intern Med 2011 Dec ;22(6):e119-24.
217. J Crit Care 2012 Jun 12;
218. PMID 23027398
219. ProMED <promedmail.org> archive: 20091001.3419
220. Emerg Infect Dis 2011 Jun ;17(6):1145-6.
221. Wkly Epidemiol Rec 2011 Oct 21;86(43):480.
222. Ann Thorac Surg 1991 Nov ;52(5):1122-6.
223. BMJ 2009 ;338:b2425.
224. Nature 2009 Aug 20;460(7258):1021-5.
225. Science 2009 Oct 30;326(5953):729-33.
226. Wkly Epidemiol Rec 2009 Aug 21;84(34):341-8.
227. Infection 2009 Oct ;37(5):381-9.
228. Wkly Epidemiol Rec 2009 Sep 4;84(36):361-5.
229. Postgrad Med 2009 Sep ;121(5):43-7.
230. J Infect Dis 2009 Oct 1;200(7):1018-21.
231. Euro Surveill 2009 ;14(42)
232. Lancet Infect Dis 2009 Oct ;9(10):583.
233. Rev Med Virol 2009 Sep ;19(5):253-6.
234. N Engl J Med 2009 Jul 9;361(2):115-9.
235. Nature 2009 May 21;459(7245):324-5.
236. Nature 2009 Apr 30;458(7242):1082-3.
237. Public Health 2009 Jun ;123(6):405-6.
238. Science 2009 Jun 19;324(5934):1496-7.
239. Science 2009 May 29;324(5931):1127.
240. Science 2009 May 22;324(5930):987.
241. Science 2009 May 15;324(5929):871.
242. Science 2009 May 8;324(5928):700-2.
243. Wkly Epidemiol Rec 2009 Nov 13;84(46):481-4.
244. Euro Surveill 2009 ;14(46)
245. Emerg Infect Dis 2009 Dec ;15(12):1963-9.
246. Mayo Clin Proc 2010 Jan ;85(1):64-76.
247. PLoS Curr 2009 ;1:RRN1133.
248. PLoS Curr 2009 ;1:RRN1123.
249. PLoS Curr 2009 ;1:RRN1047.
250. Crit Care Med 2010 Apr ;38(4 Suppl):e1-9.
251. BMJ 2009 ;339:b5681.
252. Mt Sinai J Med 2010 Jan-Feb;77(1):103-13.
253. Am J Respir Crit Care Med 2010 Jul 1;182(1):41-8.
254. Science 2010 Mar 26;327(5973):1563-4.
255. Vaccine 2010 Jul 12;28(31):4895-902.
256. N Engl J Med 2010 Jun 10;362(23):2221-3.
257. N Engl J Med 2010 Jun 10;362(23):2175-84.
258. Curr Opin Infect Dis 2010 Aug ;23(4):293-9.
259. Curr Opin Pediatr 2010 Aug ;22(4):530-5.
260. Eur J Clin Microbiol Infect Dis 2010 Nov ;29(11):1327-47.
261. Influenza Other Respi Viruses 2010 Jul ;4(4):187-97.
262. Curr Opin Infect Dis 2010 Oct ;23(5):415-20.
263. Infect Dis Clin North Am 2010 Sep ;24(3):603-17.
264. Virol J 2010 ;7:196.
265. Am J Med Sci 2010 Sep ;340(3):202-8.
266. Epidemics 2010 Sep ;2(3):132-8.
267. Epidemiol Infect 2011 Jan ;139(1):27-40.
268. PLoS Med 2010 Oct ;7(10)
269. Euro Surveill 2010 Dec 9;15(49)
270. Vaccine 2011 Feb 17;29(9):1836-43.
271. Wkly Epidemiol Rec 2011 Feb 18;86(8):61-5.
272. 2010 ;
273. Influenza Other Respi Viruses 2011 May ;5(3):157-66.
274. Influenza Other Respi Viruses 2011 May ;5(3):148-56.
275. PLoS One 2011 ;6(3):e17823.
276. Public Health 2011 May ;125(5):247-56.
277. PLoS One 2011 ;6(5):e19432.
278. Lancet Infect Dis 2011 Jun ;11(6):423-4.
279. Respirology 2011 Aug ;16(6):876-82.
280. Influenza Other Respi Viruses 2011 Sep ;5(5):306-16.
281. Influenza Other Respi Viruses 2011 Nov ;5(6):e487-98.
282. Influenza Other Respi Viruses 2011 Nov ;5(6):375-9.
283. Euro Surveill 2011 ;16(26)
284. PLoS Med 2011 Jul ;8(7):e1001053.
285. Vaccine 2011 Sep 2;29(38):6472-84.
286. PLoS One 2011 ;6(8):e21828.
287. Semin Respir Crit Care Med 2011 Aug ;32(4):400-8.
288. Semin Respir Crit Care Med 2011 Aug ;32(4):393-9.
289. Semin Respir Crit Care Med 2011 Aug ;32(4):373-92.
290. Vaccine 2011 Jul 22;29 Suppl 2:B56-62.
291. PLoS Pathog 2011 Sep ;7(9):e1002225.
292. Wkly Epidemiol Rec 2011 Oct 21;86(43):480.
293. Curr Opin Virol 2011 Oct ;1(4):254-62.
294. Vaccine 2012 Feb 8;30(7):1255-64.
295. East Mediterr Health J 2011 Apr ;17(4):342-8.
296. BMC Infect Dis 2012 Feb 10;12(1):36.
297. Euro Surveill 2012 ;17(4)
298. Vaccine 2012 Mar 1;
299. Clin Microbiol Rev 2012 Apr ;25(2):344-61.
300. Clin Microbiol Rev 2012 Apr ;25(2):223-63.
301. Bull World Health Organ 2012 Apr 1;90(4):306-10.
302. Curr Top Microbiol Immunol 2012 May 26;
303. Health Place 2012 Jul ;18(4):726-36.
304. J Infect Public Health 2012 Aug ;5(4):286-96.
305. PLoS One 2012 ;7(10):e45450.
306. Epidemiol Infect 2012 Nov 27;:1-15.
307. J Infect Dis 2012 Dec ;206 Suppl 1:S22-8.
308. J Infect Dis 2012 Dec ;206 Suppl 1:S101-7.
309. BMC Med 2012 Dec 13;10(1):165.
310. Influenza Other Respi Viruses 2012 Dec 24;
311. BMC Infect Dis 2013 Jan 7;13(1):6.
312. Influenza Other Respi Viruses 2013 Jan 21;
313. Crit Care Med 2013 Feb 8;
314. ProMED <promedmail.org> archive: 20091022.3637
315. ProMED <promedmail.org> archive: 20091204.4138
316. ProMED <promedmail.org> archive: 20091209.4193
317. ProMED <promedmail.org> archive: 20100126.0289
318. ProMED <promedmail.org> archive: 20090706.2430
319. ProMED <promedmail.org> archive: 20100603.1841
320. ProMED <promedmail.org> archive: 20091010.3510
321. ProMED <promedmail.org> archive: 20091017.3568
322. ProMED <promedmail.org> archive: 20091116.3961
323. Lancet Infect Dis 2012 Sep ;12(9):687-95.
324. Lancet Infect Dis 2012 Sep ;12(9):651-3.
325. Euro Surveill 2009 Jul 2;14(26)
326. Euro Surveill 2009 ;14(33)
327. Eur J Public Health 2012 Feb ;22(1):7-8.
328. PLoS One 2012 ;7(9):e43491.
329. ProMED <promedmail.org> archive: 20100313.0805
330. ProMED <promedmail.org> archive: 20100216.0546
331. ProMED <promedmail.org> archive: 20091231.4404
332. ProMED <promedmail.org> archive: 20091222.4313
333. ProMED <promedmail.org> archive: 20090703.2391
334. ProMED <promedmail.org> archive: 20100107.3568
335. ProMED <promedmail.org> archive: 20100327.0965
336. ProMED <promedmail.org> archive: 20091116.3961
337. ProMED <promedmail.org> archive: 20100501.1418
338. ProMED <promedmail.org> archive: 20100417.1250
339. ProMED <promedmail.org> archive: 20100522.1700
340. ProMED <promedmail.org> archive: 20100530.1798
341. ProMED <promedmail.org> archive: 20100605.1867
342. ProMED <promedmail.org> archive: 20100612.1970
343. ProMED <promedmail.org> archive: 20100619.2059
344. ProMED <promedmail.org> archive: 20100704.2223
345. ProMED <promedmail.org> archive: 20100710.2311
346. ProMED <promedmail.org> archive: 20100716.2381
347. ProMED <promedmail.org> archive: 20100724.2479
348. ProMED <promedmail.org> archive: 20100731.2565
349. ProMED <promedmail.org> archive: 20100807.2680
350. MMWR Morb Mortal Wkly Rep 2009 Dec 11;58(48):1341-4.
351. Euro Surveill 2009 ;14(50)
352. Int J Tuberc Lung Dis 2010 Feb ;14(2):130.
353. Med J Aust 2010 May 17;192(10):623.
354. J Physiol 1978 Feb ;275:60P-61P.
355. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S189-97.
356. Ann Epidemiol 2011 Aug ;21(8):623-30.

357. Am J Public Health 2011 Sep ;101(9):1776-84.
358. Rural Remote Health 2011 Jul-Sep;11(3):1781.
359. Emerg Infect Dis 2011 Sep ;17(9):1615-23.
360. Commun Dis Intell 2011 Jun ;35(2):172-6.
361. Emerg Infect Dis 2012 Jan ;18(1):71-7.
362. Emerg Infect Dis 2012 May ;18(5):866-8.
363. Can J Public Health 2012 Mar-Apr;103(2):90-3.
364. Am J Public Health 2012 Dec 13;
365. BMC Public Health 2012 Dec 20;12(1):1098.
366. ProMED <promedmail.org> archive: 20101203.4341
367. ProMED <promedmail.org> archive: 20091213.4236
368. ProMED <promedmail.org> archive: 20100114.0160
369. ProMED <promedmail.org> archive: 20091214.4246
370. ProMED <promedmail.org> archive: 20090717.2553
371. BMJ 2010 ;341:c4393.
372. ProMED <promedmail.org> archive: 20100811.2753
373. Nature 2009 Aug 6;460(7256):683.
374. PLoS One 2009 ;4(12):e8367.
375. Curr Top Microbiol Immunol 2012 Dec 21;
376. Emerg Infect Dis 2010 Feb ;16(2):304-7.
377. Influenza Other Respi Viruses 2012 May 30;
378. ProMED <promedmail.org> archive: 20120601.1152800
379. Influenza Other Respi Viruses 2012 Sep ;6(5):348-57.
380. ProMED <promedmail.org> archive: 20120106.1001398
381. J Environ Biol 2012 Mar ;33(2):155-7.
382. J Virol 2010 Mar ;84(5):2245-56.
383. Emerg Infect Dis 2010 Apr ;16(4):706-8.
384. J Virol 2011 Sep ;85(17):8667-79.
385. J Gen Virol 2012 Oct ;93(Pt 10):2195-203.
386. Curr Top Microbiol Immunol 2012 Dec 21;
387. ProMED <promedmail.org> archive: 20090505.1680
388. ProMED <promedmail.org> archive: 20090505.1683
389. ProMED <promedmail.org> archive: 20090506.1691
390. ProMED <promedmail.org> archive: 20090507.1709
391. ProMED <promedmail.org> archive: 20090513.1790
392. ProMED <promedmail.org> archive: 20090615.2215
393. ProMED <promedmail.org> archive: 20090828.3027
394. ProMED <promedmail.org> archive: 20090626.2322
395. ProMED <promedmail.org> archive: 20090701.2376
396. ProMED <promedmail.org> archive: 20090703.2401
397. Aust Vet J 2011 Nov ;89(11):427-31.
398. Influenza Other Respi Viruses 2012 Feb 15;
399. ProMED <promedmail.org> archive: 20090801.2698
400. Influenza Other Respi Viruses 2012 Dec 28;
401. Vet Microbiol 2012 Apr 23;156(1-2):189-92.
402. ProMED <promedmail.org> archive: 20110924.2897
403. Res Vet Sci 2012 Aug ;93(1):125-32.
404. J Virol 2011 Oct ;85(19):10279-85.
405. Emerg Infect Dis 2011 Sep ;17(9):1757-9.
406. Emerg Infect Dis 2012 Feb ;18(2):357-9.
407. ProMED <promedmail.org> archive: 20100111.0128
408. ProMED <promedmail.org> archive: 20100901.3114
409. J Gen Virol 2011 May ;92(Pt 5):1184-8.
410. ProMED <promedmail.org> archive: 20091211.4220
411. ProMED <promedmail.org> archive: 20091028.3737
412. Indian J Med Res 2010 Aug ;132:160-7.
413. ProMED <promedmail.org> archive: 20091127.4071
414. ProMED <promedmail.org> archive: 20091002.3427
415. Open Virol J 2010 ;4:52-6.
416. ProMED <promedmail.org> archive: 20091205.4144
417. Microbiol Immunol 2012 Nov ;56(11):792-803.
418. ProMED <promedmail.org> archive: 20091022.3635
419. ProMED <promedmail.org> archive: 20091204.4141
420. ProMED <promedmail.org> archive: 20091211.4214
421. ProMED <promedmail.org> archive: 20090918.3280
422. Euro Surveill 2009 ;14(45)
423. ProMED <promedmail.org> archive: 20091019.3589
424. Vet Rec 2011 Aug 6;169(6):155.
425. ProMED <promedmail.org> archive: 20100422.1296
426. Emerg Infect Dis 2012 Oct ;18(10):1665-8.
427. ProMED <promedmail.org> archive: 20091226.4353
428. Vet Rec 2010 May 1;166(18):548-51.
429. ProMED <promedmail.org> archive: 20091106.3840
430. Emerg Infect Dis 2010 Oct ;16(10):1587-90.
431. Virus Genes 2011 Aug ;43(1):1-5.
432. PLoS Curr 2011 ;3:RRN1209.
433. Vet Rec 2012 Sep 15;171(11):271.
434. Science 2010 Jun 18;328(5985):1529.
435. Ecohealth 2011 Sep ;8(3):376-80.
436. Emerg Infect Dis 2012 Sep ;18(9):1519-21.
437. ProMED <promedmail.org> archive: 20091019.3592
438. ProMED <promedmail.org> archive: 20091202.4111
439. ProMED <promedmail.org> archive: 20091020.3600
440. ProMED <promedmail.org> archive: 20091106.3834
441. ProMED <promedmail.org> archive: 20091228.4372
442. ProMED <promedmail.org> archive: 20091107.3857
443. ProMED <promedmail.org> archive: 20091020.3602
444. ProMED <promedmail.org> archive: 20091022.3629
445. ProMED <promedmail.org> archive: 20091027.3719
446. Emerg Infect Dis 2010 Apr ;16(4):709-11.
447. Rev Chilena Infectol 2012 Dec ;29(6):695.
448. ProMED <promedmail.org> archive: 20090821.2961
449. ProMED <promedmail.org> archive: 20090823.2978
450. ProMED <promedmail.org> archive: 20090829.3036
451. ProMED <promedmail.org> archive: 20100131.0337
452. Avian Dis 2012 Dec ;56(4 Suppl):1062-7.
453. ProMED <promedmail.org> archive: 20110113.0150
454. ProMED <promedmail.org> archive: 20100108.0087
455. ProMED <promedmail.org> archive: 20091202.4111
456. Emerg Infect Dis 2010 Mar ;16(3):534-7.
457. Zoonoses Public Health 2012 Dec ;59(8):549-52.
458. Vet Pathol 2010 May ;47(3):378-86.
459. Arch Virol 2011 Jan ;156(1):117-20.
460. J Am Vet Med Assoc 2011 Apr 15;238(8):964.
461. Zoonoses Public Health 2011 Dec ;58(8):573-81.
462. J Clin Microbiol 2011 Dec ;49(12):4101-5.
463. Clin Vaccine Immunol 2012 Oct 31;
464. ProMED <promedmail.org> archive: 20091105.3816
465. ProMED <promedmail.org> archive: 20091121.4008
466. Emerg Infect Dis 2012 Feb ;18(2):315-7.
467. ProMED <promedmail.org> archive: 20091211.4213
468. ProMED <promedmail.org> archive: 20091209.4192
469. J Infect Dis 2009 Dec 15;200(12):1884-92.
470. PLoS Pathog 2010 ;6(7):e1001022.
471. Virol J 2010 ;7:149.
472. J Vet Diagn Invest 2010 Sep ;22(5):784-8.
473. Zoonoses Public Health 2012 Jun 15;
474. PLoS One 2012 ;7(8):e42343.
475. ProMED <promedmail.org> archive: 20091222.4305
476. Emerg Infect Dis 2011 Apr ;17(4):747-9.
477. ProMED <promedmail.org> archive: 20110324.0925
478. Emerg Infect Dis 2012 Feb ;18(2):315-7.
479. ProMED <promedmail.org> archive: 20091211.4213
480. Science 2009 Jul 3;325(5936):17.
481. J Infect Dis 2009 Dec 15;200(12):1884-92.
482. ProMED <promedmail.org> archive: 20091101.3777
483. ProMED <promedmail.org> archive: 20091021.3618
484. ProMED <promedmail.org> archive: 20091114.3936
485. ProMED <promedmail.org> archive: 20091202.4111
486. ProMED <promedmail.org> archive: 20091115.3947
487. Emerg Infect Dis 2010 Jun ;16(6):1043-5.
488. Emerg Infect Dis 2010 Dec ;16(12):2019-21.
489. J Gen Virol 2012 Jan ;93(Pt 1):119-23.
490. Vet Rec 2012 Mar 23;
491. ProMED <promedmail.org> archive: 20100101.0014
492. ProMED <promedmail.org> archive: 20091104.3813
493. ProMED <promedmail.org> archive: 20091118.3981
494. Presse Med 2010 Jun ;39(6):730-4.
495. ProMED <promedmail.org> archive: 20091115.3945
496. J Infect Dis 2012 Dec ;206 Suppl 1:S136-9.
497. Medicina (B Aires) 2009 ;69(4):393-423.
498. Medicina (B Aires) 2009 ;69(4):478-82.
499. An Pediatr (Barc) 2010 Jan ;72(1):62-6.
500. Clin Infect Dis 2009 Nov 1;49(9):1458-60.
501. Nature 2009 Jul 16;460(7253):311.
502. Emerg Infect Dis 2009 Dec ;15(12):2060-1.
503. N Engl J Med 2010 Jan 7;362(1):45-55.
504. Emerg Infect Dis 2010 Feb ;16(2):311-3.
505. Intensive Care Med 2010 Jun ;36(6):1015-22.
506. Rev Panam Salud Publica 2010 Mar ;27(3):226-9.
507. Mem Inst Oswaldo Cruz 2010 Mar ;105(2):179-83.
508. Scand J Infect Dis 2011 Jan ;43(1):70-4.
509. J Am Geriatr Soc 2010 Sep ;58(9):1813-5.
510. Medicina (B Aires) 2010 ;70(6):518-23.
511. Infect Control Hosp Epidemiol 2011 Jan ;32(1):87-90.
512. Public Health Rep 2011 Jan-Feb;126(1):9-12.
513. Arch Argent Pediatr 2011 Jun ;109(3):198-203.
514. PLoS One 2012 ;7(4):e33670.
515. PLoS One 2012 ;7(10):e47540.
516. J Infect Dev Ctries 2013 ;7(1):36-40.
517. ProMED <promedmail.org> archive: 20091227.4359
518. ProMED <promedmail.org> archive: 20090626.2322
519. BMJ 2009 ;339:b2858.
520. Euro Surveill 2009 Aug 6;14(31)
521. Med J Aust 2009 Sep 7;191(5):267-9.
522. Med J Aust 2009 Oct 19;191(8):454-8.
523. Euro Surveill 2009 ;14(42)
524. Euro Surveill 2009 ;14(42)
525. Euro Surveill 2009 ;14(42)
526. Med J Aust 2009 Nov 2;191(9):502-6.
527. Aust Fam Physician 2009 Aug ;38(8):567.
528. Crit Care Resusc 2009 Sep ;11(3):170-2.
529. Med J Aust 2009 Nov 16;191(10):573-4.
530. N Engl J Med 2009 Dec 31;361(27):2591-4.
531. Med J Aust 2010 Jan 4;192(1):33-6.
532. Disaster Med Public Health Prep 2009 Dec ;3 Suppl 2:S154-9.
533. Euro Surveill 2009 ;14(50)
534. Med J Aust 2010 Jan 18;192(2):94-7.
535. Med J Aust 2010 Jan 18;192(2):84-6.
536. Am J Respir Crit Care Med 2010 Feb 15;181(4):300-6.
537. Respiriology 2010 Jan ;15(1):51-6.
538. PLoS One 2010 ;5(3):e9880.
539. Med J Aust 2010 Apr 5;192(7):364-5.
540. N S W Public Health Bull 2010 Jan-Feb;21(1-2):1-3.
541. Med J Aust 2010 May 17;192(10):623.
542. J Physiol 1978 Feb ;275:60P-61P.
543. Crit Care Resusc 2010 Jun ;12(2):121-30.
544. PLoS One 2010 ;5(6):e11341.

545. Intensive Crit Care Nurs 2010 Aug ;26(4):207-14.
546. Aust N Z J Public Health 2010 Jun ;34(3):228-31.
547. Emerg Infect Dis 2010 Aug ;16(8):1211-6.
548. Commun Dis Intell 2010 Jun ;34(2):102-9.
549. J Paediatr Child Health 2010 Nov ;46(11):673-9.
550. Euro Surveill 2010 ;15(31)
551. Emerg Infect Dis 2010 Sep ;16(9):1396-402.
552. Emerg Infect Dis 2010 Sep ;16(9):1388-95.
553. PLoS One 2010 ;5(9):e12562.
554. Med J Aust 2010 Oct 4;193(7):401-4.
555. Euro Surveill 2010 ;15(40)
556. Aust Health Rev 2010 Nov ;34(4):477-86.
557. Pediatrics 2011 Jan ;127(1):e156-63.
558. Soc Sci Med 2011 Mar ;72(6):912-8.
559. Med J Aust 2011 Feb 21;194(4):169-74.
560. Crit Care 2011 ;15(3):R143.
561. Anaesth Intensive Care 2011 May ;39(3):384-91.
562. BMC Public Health 2012 Oct 12;12(1):869.
563. Emerg Infect Dis 2013 Jan ;19(1):92-101.
564. ProMED <promedmail.org> archive: 20101021.3812
565. ProMED <promedmail.org> archive: 20090806.2784
566. Infection 2011 Aug ;39(4):341-52.
567. Eur J Epidemiol 2012 Jul ;27(7):567-75.
568. J Med Virol 2012 Sep ;84(9):1331-4.
569. Emerg Infect Dis 2012 Jan ;18(1):146-9.
570. BMC Public Health 2010 ;10 Suppl 1:S10.
571. ProMED <promedmail.org> archive: 20091110.3893
572. Euro Surveill 2009 Jul 16;14(28)
573. Euro Surveill 2009 Aug 6;14(31)
574. J Hosp Infect 2011 Feb ;77(2):118-22.
575. Influenza Other Respi Viruses 2011 Sep ;5(5):351-6.
576. Influenza Other Respi Viruses 2012 Jul 20;
577. Euro Surveill 2009 ;14(35)
578. Clinics (Sao Paulo) 2009 ;64(10):1025-30.
579. Rev Saude Publica 2009 Oct ;43(5):900-4.
580. Mem Inst Oswaldo Cruz 2010 Mar ;105(2):179-83.
581. Euro Surveill 2009 ;14(42)
582. Braz J Infect Dis 2011 May-Jun;15(3):220-4.
583. Diagn Microbiol Infect Dis 2011 Sep ;71(1):98-9.
584. J Bras Pneumol 2012 Feb ;38(1):57-65.
585. Vaccine 2012 May 17;
586. Cien Saude Colet 2012 Jun ;17(6):1629-34.
587. Cad Saude Publica 2012 Jul ;28(7):1325-36.
588. Cad Saude Publica 2013 Jan ;29(1):189-94.
589. BMC Public Health 2010 ;10:322.
590. Southeast Asian J Trop Med Public Health 2010 Sep ;41(5):1104-15.
591. Southeast Asian J Trop Med Public Health 2012 Jan ;43(1):68-77.
592. BMC Infect Dis 2010 ;10:56.
593. BMJ 2009 ;339:b2746.
594. CMAJ 2009 Aug 4;181(3-4):159-63.
595. CMAJ 2009 Sep 15;181(6-7):E102-5.
596. JAMA 2009 Nov 4;302(17):1872-9.
597. Wkly Epidemiol Rec 2009 Nov 20;84(47):485-91.
598. CMAJ 2010 Feb 9;182(2):131-6.
599. Emerg Infect Dis 2009 Dec ;15(12):2001-3.
600. Can J Public Health 2009 Sep-Oct;100(5):337-9.
601. PLoS Curr 2009 ;1:RRN1137.
602. Int J Tuberc Lung Dis 2010 Feb ;14(2):130.
603. CMAJ 2010 Feb 23;182(3):257-64.
604. Vaccine 2010 Apr 19;28(18):3180-4.
605. Pediatr Crit Care Med 2010 Sep ;11(5):603-9.
606. PLoS Med 2010 Apr ;7(4):e1000258.
607. Vaccine 2010 Aug 31;28(38):6210-20.
608. CMAJ 2010 Nov 23;182(17):1851-6.
609. BMC Res Notes 2010 ;3:283.
610. J Popul Ther Clin Pharmacol 2010 ;17(3):e358-62.
611. CMAJ 2010 Dec 14;182(18):1981-7.
612. Influenza Other Respi Viruses 2011 Mar ;5(2):83-8.
613. BMC Infect Dis 2011 ;11:90.
614. Epidemiol Infect 2011 Sep 16;:1-12.
615. Can J Public Health 2011 Sep-Oct;102(5):345-8.
616. PLoS One 2011 ;6(11):e26427.
617. Am J Infect Control 2012 May 8;
618. Pediatrics 2012 Sep ;130(3):397-406.
619. BMJ Open 2012 ;2(5)
620. Infect Control Hosp Epidemiol 2012 Oct ;33(10):1043-6.
621. Influenza Other Respi Viruses 2012 Nov 5;
622. J Health Commun 2013 Jan 10;
623. ProMED <promedmail.org> archive: 20091204.4138
624. ProMED <promedmail.org> archive: 20091203.4127
625. Rev Chilena Infectol 2009 Aug ;26(4):307-10.
626. Gut 2009 Nov ;58(11):1567-8.
627. Crit Care Med 2010 Apr ;38(4 Suppl):e133-7.
628. Euro Surveill 2010 Jan 7;15(1)
629. Clin Infect Dis 2010 Mar 15;50(6):860-8.
630. Eur Respir J 2010 Oct ;36(4):864-9.
631. Crit Care Med 2010 Apr ;38(4 Suppl):e133-7.
632. Mem Inst Oswaldo Cruz 2010 Mar ;105(2):179-83.
633. Can Assoc Radiol J 2010 Oct ;61(4):233-40.
634. Rev Chilena Infectol 2010 Apr ;27(2):144-7.
635. Rev Med Chil 2010 Sep ;138(9):1186-96.
636. Emerg Infect Dis 2011 Jul ;17(7):1256-8.
637. Rev Med Chil 2011 Mar ;139(3):321-6.
638. Rev Med Chil 2011 Jul ;139(7):833-40.
639. BMC Infect Dis 2012 Nov 13;12(1):298.
640. ProMED <promedmail.org> archive: 20090618.2253
641. Virol J 2013 Feb 8;10(1):49.
642. J Clin Microbiol 2009 Jul ;47(7):2344-6.
643. Hong Kong Med J 2009 Oct ;15(5):381-4.
644. BMJ 2009 ;339:b4164.
645. J Clin Virol 2009 Nov ;46(3):298-9.
646. Emerg Infect Dis 2010 Mar ;16(3):538-41.
647. Vaccine 2010 Jun 23;28(29):4632-7.
648. BMC Infect Dis 2010 ;10:139.
649. Am J Infect Control 2010 Jun ;38(5):374-80.
650. PLoS One 2010 ;5(6):e10911.
651. J Infect Dis 2010 Sep 15;202(6):867-76.
652. Epidemiology 2010 Nov ;21(6):842-6.
653. J Med Virol 2010 Nov ;82(11):1809-15.
654. Clin Infect Dis 2010 Nov 15;51(10):1184-91.
655. PLoS One 2011 ;6(3):e17713.
656. J Hosp Infect 2011 Aug ;78(4):308-11.
657. J Infect 2011 Oct ;63(4):274-80.
658. Clin Infect Dis 2011 Jul 1;53(1):100-1; author reply 101, 103.
659. Influenza Other Respi Viruses 2011 May ;5 Suppl 1:190-4.
660. Epidemiol Infect 2012 Sep ;140(9):1542-50.
661. Matern Child Health J 2012 Feb 2;
662. Hong Kong Med J 2012 Aug ;18(4):310-7.
663. Influenza Other Respi Viruses 2012 Aug 9;
664. Euro Surveill 2012 ;17(45)
665. ProMED <promedmail.org> archive: 20090616.2221
666. ProMED <promedmail.org> archive: 20121108.1400408
667. J Clin Microbiol 2009 Jul ;47(7):2344-6.
668. Science 2009 Sep 18;325(5947):1482-3.
669. Emerg Infect Dis 2009 Sep ;15(9):1418-22.
670. Hong Kong Med J 2009 Oct ;15(5):381-4.
671. Emerg Infect Dis 2009 Nov ;15(11):1849-50.
672. Zhonghua Jie He He Hu Xi Za Zhi 2009 Jul ;32(7):482-4.
673. Zhonghua Liu Xing Bing Xue Za Zhi 2009 Jul ;30(7):684-6.
674. Clin Infect Dis 2010 Feb 15;50(4):622-3.
675. Zhonghua Liu Xing Bing Xue Za Zhi 2009 Jul ;30(7):653-5.
676. N Engl J Med 2009 Dec 24;361(26):2507-17.
677. Biosci Trends 2009 Aug ;3(4):127-30.
678. Zhonghua Yu Fang Yi Xue Za Zhi 2009 Oct ;43(10):856-60.
679. QJM 2010 May ;103(5):311-7.
680. Zhonghua Liu Xing Bing Xue Za Zhi 2009 Nov ;30(11):1121-4.
681. Zhonghua Liu Xing Bing Xue Za Zhi 2009 Nov ;30(11):1106-10.
682. Zhonghua Liu Xing Bing Xue Za Zhi 2009 Nov ;30(11):1102-5.
683. Chin Med J (Engl) 2010 Feb 20;123(4):401-5.
684. Emerg Infect Dis 2010 Apr ;16(4):725-6.
685. Zhonghua Jie He He Hu Xi Za Zhi 2009 Jul ;33(2):81-5.
686. Emerg Infect Dis 2010 Jun ;16(6):1011-3.
687. Clin Infect Dis 2010 Jul 15;51(2):251-2.
688. Infect Control Hosp Epidemiol 2010 Sep ;31(9):961-3.
689. J Infect 2010 Oct ;61(4):277-83.
690. Asia Pac J Public Health 2012 Mar ;24(2):352-60.
691. Epidemiol Infect 2011 Jan ;139(1):52-8.
692. Hum Vaccin 2010 Sep 8;6(9)
693. Infect Genet Evol 2011 Jan ;11(1):222-6.
694. Zhonghua Jie He He Hu Xi Za Zhi 2010 Jun ;33(6):403-5.
695. Emerg Infect Dis 2010 Nov ;16(11):1809-11.
696. Intervirology 2011 ;54(3):164-70.
697. Prev Med 2011 Jan ;52(1):71-4.
698. Chin Med J (Engl) 2010 Oct ;123(19):2651-4.
699. N Engl J Med 2010 Dec 16;363(25):2416-23.
700. Zhonghua Liu Xing Bing Xue Za Zhi 2010 May ;31(5):497-9.
701. Zhonghua Liu Xing Bing Xue Za Zhi 2010 May ;31(5):494-6.
702. Zhonghua Liu Xing Bing Xue Za Zhi 2010 May ;31(5):485-8.
703. Zhonghua Er Ke Za Zhi 2010 Oct ;48(10):733-8.
704. Clin Infect Dis 2011 Feb 15;52(4):457-65.
705. Intervirology 2011 ;54(5):233-45.
706. PLoS One 2011 ;6(4):e17919.
707. Asia Pac J Public Health 2011 May 5;
708. BMC Infect Dis 2011 ;11:128.
709. PLoS One 2011 ;6(10):e25934.
710. Pediatr Infect Dis J 2012 Apr 4;
711. Am J Epidemiol 2012 Apr 5;
712. Public Health 2012 Apr 17;
713. Emerg Infect Dis 2012 May ;18(5):758-66.
714. Zhonghua Liu Xing Bing Xue Za Zhi 2012 Jan ;33(1):62-6.
715. Acta Virol 2012 ;56(4):329-35.
716. ProMED <promedmail.org> archive: 20100303.0702
717. ProMED <promedmail.org> archive: 20100115.0180
718. ProMED <promedmail.org> archive: 20100105.0040
719. ProMED <promedmail.org> archive: 20091203.4116
720. ProMED <promedmail.org> archive: 20090616.2221
721. ProMED <promedmail.org> archive: 20121108.1400408
722. ProMED <promedmail.org> archive: 20091103.3787
723. Euro Surveill 2009 Jul 30;14(30):19284.
724. Bull Soc Pathol Exot 2011 May ;104(2):114-8.
725. ProMED <promedmail.org> archive: 20091112.3920
726. Emerg Infect Dis 2012 Feb ;18(2):336-7.
727. Euro Surveill 2009 ;14(33)
728. J Infect Dis 2012 Dec ;206 Suppl 1:S36-40.
729. ProMED <promedmail.org> archive: 20091110.3893
730. ProMED <promedmail.org> archive: 20091231.4403
731. Ugeskr Laeger 2009 Jun 8;171(24):1996.
732. Euro Surveill 2010 Dec 9;15(49)

733. Euro Surveill 2011 ;16(3)
734. Vaccine 2011 Jul 22;29 Suppl 2:B63-9.
735. Influenza Other Respi Viruses 2013 Feb 9;
736. Med Intensiva 2010 Jun-Jul;34(5):310-7.
737. ProMED <promedmail.org> archive: 20090609.2128
738. ProMED <promedmail.org> archive: 20091215.4255
739. ProMED <promedmail.org> archive: 20090611.2150
740. ProMED <promedmail.org> archive: 20091101.3776
741. ProMED <promedmail.org> archive: 20091025.3695
742. Euro Surveill 2010 Feb 4;15(5)
743. Euro Surveill 2010 ;15(45)
744. Euro Surveill 2011 ;16(27)
745. Euro Surveill 2012 ;17(38)
746. Euro Surveill 2009 May 28;14(21)
747. Euro Surveill 2009 Jul 9;14(27)
748. Med Mal Infect 2010 Jan ;40(1):48-50.
749. Euro Surveill 2009 Jul 23;14(29)
750. Euro Surveill 2010 Jan 14;15(2)
751. Clin Microbiol Infect 2010 Apr ;16(4):322-5.
752. Clin Microbiol Infect 2010 Apr ;16(4):304-8.
753. Clin Microbiol Infect 2010 Apr ;16(4):303.
754. Clin Microbiol Infect 2010 Apr ;16(4):393-6.
755. Euro Surveill 2010 Feb 11;15(6)
756. PLoS One 2010 ;5(2):e9214.
757. Euro Surveill 2009 ;14(39)
758. PLoS One 2010 ;5(4):e10199.
759. BMC Infect Dis 2010 ;10:162.
760. Euro Surveill 2010 Jun 24;15(25)
761. Med Mal Infect 2010 Jul ;40(7):404-11.
762. PLoS Curr 2010 ;2:RRN1188.
763. Bull Soc Pathol Exot 2011 May ;104(2):119-24.
764. Vaccine 2010 Nov 29;28(51):8157-61.
765. Prev Med 2011 Feb ;52(2):178-81.
766. Arch Pediatr 2011 Mar 31;
767. Vaccine 2011 Jun 20;29(28):4632-7.
768. PLoS One 2011 ;6(5):e19621.
769. Bull Acad Natl Med 2010 Apr-May;194(4-5):719-32; discussion 732.
770. Clin Microbiol Infect 2012 Feb ;18(2):177-83.
771. Vaccine 2011 Sep 16;29(40):7075-9.
772. Vaccine 2012 Feb 1;30(6):995-7.
773. Influenza Other Respi Viruses 2012 Mar 22;
774. Presse Med 2012 May 3;
775. J Med Virol 2012 Jul ;84(7):1071-9.
776. Drug Saf 2012 Oct 1;35(10):845-54.
777. PLoS One 2012 ;7(9):e45051.
778. ProMED <promedmail.org> archive: 20110111.0128
779. Bull Soc Pathol Exot 2011 May ;104(2):119-24.
780. Rev Panam Salud Publica 2012 Aug ;32(2):124-30.
781. Clin Microbiol Infect 2010 Apr ;16(4):303.
782. Clin Microbiol Infect 2010 Apr ;16(4):304-8.
783. Euro Surveill 2009 May 7;14(18)
784. Euro Surveill 2009 Aug 6;14(31)
785. Euro Surveill 2009 ;14(32)
786. Euro Surveill 2009 ;14(34)
787. MMW Fortschr Med 2009 Oct 1;151(40):35-7.
788. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2010 May ;53(5):510-9.
789. Euro Surveill 2010 May 6;15(18)
790. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2010 Dec ;53(12):1277-82.
791. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2010 Dec ;53(12):1257-66.
792. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2010 Dec ;53(12):1223-30.
793. J Clin Virol 2009 Nov ;46(3):295-7.
794. Euro Surveill 2010 Dec 9;15(49)
795. BMC Infect Dis 2010 ;10:155.
796. Vaccine 2011 May 23;29(23):4008-12.
797. PLoS One 2011 ;6(6):e21340.
798. PLoS One 2011 ;6(9):e23955.
799. Gesundheitswesen 2011 Nov ;73(11):722-9.
800. PMID 22278764
801. Euro Surveill 2012 ;17(13)
802. J Infect Dis 2012 Dec ;206 Suppl 1:S108-13.
803. Euro Surveill 2009 May 28;14(21)
804. Euro Surveill 2010 Feb 11;15(6)
805. Acta Paediatr 2010 Aug ;99(8):1114-5.
806. MMWR Morb Mortal Wkly Rep 2010 Jun 11;59(22):682-6.
807. Euro Surveill 2009 Jul 23;14(29)
808. Int J Adolesc Med Health 2010 Apr-Jun;22(2):339-40.
809. PLoS Curr 2010 ;2:RRN1194.
810. Eur J Public Health 2011 Jun ;21(3):329-32.
811. PLoS One 2011 ;6(6):e20593.
812. Vaccine 2011 Sep 29;29(38):6664-9.
813. Bull Soc Pathol Exot 2011 May ;104(2):119-24.
814. Hawaii Med J 2010 Jun ;69(6 Suppl 3):50-1.
815. Influenza Other Respi Viruses 2010 May 1;4(3):129-40.
816. PLoS One 2010 ;5(12):e15826.
817. Influenza Other Respi Viruses 2012 Nov 30;
818. ProMED <promedmail.org> archive: 20091203.4127
819. J Clin Microbiol 2009 Jul ;47(7):2344-6.
820. Hong Kong Med J 2009 Oct ;15(5):381-4.
821. BMJ 2009 ;339:b4164.
822. J Clin Virol 2009 Nov ;46(3):298-9.
823. Emerg Infect Dis 2010 Mar ;16(3):538-41.
824. Vaccine 2010 Jun 23;28(29):4632-7.
825. BMC Infect Dis 2010 ;10:139.
826. Am J Infect Control 2010 Jun ;38(5):374-80.
827. PLoS One 2010 ;5(6):e10911.
828. J Infect Dis 2010 Sep 15;202(6):867-76.
829. Epidemiology 2010 Nov ;21(6):842-6.
830. J Med Virol 2010 Nov ;82(11):1809-15.
831. Clin Infect Dis 2010 Nov 15;51(10):1184-91.
832. PLoS One 2011 ;6(3):e17713.
833. J Hosp Infect 2011 Aug ;78(4):308-11.
834. J Infect 2011 Oct ;63(4):274-80.
835. Clin Infect Dis 2011 Jul 1;53(1):100-1; author reply 101, 103.
836. Influenza Other Respi Viruses 2011 May ;5 Suppl 1:190-4.
837. Epidemiol Infect 2012 Sep ;140(9):1542-50.
838. Matern Child Health J 2012 Feb 2;
839. Hong Kong Med J 2012 Aug ;18(4):310-7.
840. Influenza Other Respi Viruses 2012 Aug 9;
841. Euro Surveill 2012 ;17(45)
842. ProMED <promedmail.org> archive: 20090616.2221
843. ProMED <promedmail.org> archive: 20121108.1400408
844. Lege Artis Med 2011 Feb ;21(2):89-95.
845. Laeknabladid 2010 Feb ;96(2):83-90.
846. Euro Surveill 2010 Dec 9;15(49)
847. Laeknabladid 2010 Feb ;96(2):83-90.
848. Indian J Med Res 2009 May ;129(5):465-7.
849. Indian J Public Health 2009 Jul-Sep;53(3):190-1.
850. J Infect Dev Ctries 2010 Jan ;4(1):7-14.
851. Indian Pediatr 2010 Jan ;47(1):25-31.
852. Ann Trop Paediatr 2010 ;30(1):51-5.
853. PLoS One 2010 ;5(3):e9693.
854. PLoS One 2010 ;5(5):e10540.
855. Indian J Pediatr 2010 Sep ;77(9):981-5.
856. BMC Infect Dis 2010 ;10:255.
857. Arch Virol 2011 Feb ;156(2):207-17.
858. Bioinformation 2011 ;5(10):416-21.
859. Indian J Med Res 2012 Apr ;135(4):534-7.
860. ProMED <promedmail.org> archive: 20100726.2506
861. ProMED <promedmail.org> archive: 20100720.2436
862. ProMED <promedmail.org> archive: 20100630.2177
863. ProMED <promedmail.org> archive: 20091103.3796
864. ProMED <promedmail.org> archive: 20100706.2249
865. BMC Public Health 2010 ;10:322.
866. Microbiol Immunol 2011 Jul ;55(7):514-7.
867. Arch Iran Med 2009 Nov ;12(6):533-41.
868. Arch Iran Med 2010 Mar ;13(2):91-8.
869. Influenza Other Respi Viruses 2010 Jul ;4(4):179-86.
870. Nihon Rinsho 2010 Sep ;68(9):1605-10.
871. Influenza Other Respi Viruses 2012 Apr 6;
872. Iran Red Crescent Med J 2011 Oct ;13(10):698-701.
873. Int J Prev Med 2012 Dec ;3(12):860-6.
874. ProMED <promedmail.org> archive: 20090906.3140
875. ProMED <promedmail.org> archive: 20091115.3946
876. ProMED <promedmail.org> archive: 20091102.3780
877. Euro Surveill 2009 Sep 24;14(38)
878. Euro Surveill 2009 ;14(44)
879. Ir Med J 2009 Sep ;102(8):267-8.
880. Biosecur Bioterror 2009 Dec ;7(4):399-404.
881. Harefuah 2009 Nov ;148(11):738-42, 795.
882. Harefuah 2009 Sep ;148(9):577-82, 659.
883. Int J Tuberc Lung Dis 2010 Feb ;14(2):130.
884. Harefuah 2009 Dec ;148(12):799-803, 857.
885. Emerg Infect Dis 2010 Apr ;16(4):720-1.
886. J Public Health Policy 2010 Jul ;31(2):256-69.
887. Arch Pediatr Adolesc Med 2010 Nov ;164(11):1015-22.
888. BMC Infect Dis 2011 ;11:92.
889. Infection 2011 Oct ;39(5):399-404.
890. Emerg Infect Dis 2011 Sep ;17(9):1740-3.
891. Clin Infect Dis 2011 Dec ;53(12):1199-207.
892. Influenza Other Respi Viruses 2012 Dec 22;
893. ProMED <promedmail.org> archive: 20091215.4255
894. Euro Surveill 2009 Jun 18;14(24)
895. Euro Surveill 2009 ;14(49)
896. Clin Microbiol Infect 2011 Feb ;17(2):247-50.
897. Risk Anal 2011 Apr ;31(4):645-56.
898. Euro Surveill 2010 Dec 9;15(49)
899. Euro Surveill 2009 Jul 9;14(27)
900. Eur J Epidemiol 2011 Mar ;26(3):211-9.
901. Antiviral Res 2011 Jun ;90(3):205-12.
902. Intensive Care Med 2011 Nov ;37(11):1746-55.
903. Hum Vaccin 2011 Jan-Feb;7 Suppl:217-25.
904. Int J Immunopathol Pharmacol 2011 Jul-Sep;24(3):651-9.
905. J Med Virol 2011 Dec ;83(12):2057-65.
906. Epidemics 2012 Mar ;4(1):9-21.
907. Influenza Other Respi Viruses 2012 Aug 2;
908. Euro Surveill 2009 Jun 4;14(22)
909. Wkly Epidemiol Rec 2009 Jun 12;84(24):237-44.
910. Euro Surveill 2009 Jun 18;14(24)
911. Euro Surveill 2009 Jul 23;14(29)
912. Euro Surveill 2009 ;14(35)
913. Am J Disaster Med 2009 May-Jun;4(3):133-4.
914. Euro Surveill 2009 ;14(44)
915. J Infect Chemother 2011 Jun ;17(3):401-6.
916. PLoS Curr 2009 ;1:RRN1139.

917. Nihon Koshu Eisei Zasshi 2010 Mar ;57(3):157-64.
918. PLoS One 2010 ;5(6):e11057.
919. Emerg Infect Dis 2009 Oct ;15(10):1685.
920. J Infect Chemother 2011 Aug ;17(4):468-72.
921. Emerg Infect Dis 2011 Mar ;17(3):470-9.
922. Influenza Other Respi Viruses 2010 Jul ;4(4):179-86.
923. Nihon Rinsho 2010 Sep ;68(9):1605-10.
924. Environ Health Prev Med 2011 Sep ;16(5):320-6.
925. Emerg Infect Dis 2011 Apr ;17(4):746-7.
926. PLoS One 2011 ;6(4):e19409.
927. J Infect Chemother 2011 Oct ;17(5):595-601.
928. Emerg Infect Dis 2011 Sep ;17(9):1763-5.
929. Emerg Infect Dis 2011 Sep ;17(9):1737-9.
930. Emerg Infect Dis 2011 Nov ;17(11):1993-2000.
931. Jpn J Infect Dis 2011 ;64(6):473-81.
932. Pediatr Infect Dis J 2012 Apr ;31(4):368-72.
933. PLoS One 2012 ;7(2):e31289.
934. J Obstet Gynaecol Res 2012 Apr 9;
935. PLoS One 2012 ;7(6):e36455.
936. PLoS One 2013 ;8(1):e54786.
937. Biosecur Bioterror 2009 Dec ;7(4):399-404.
938. Virol Sin 2011 Oct ;26(5):306-14.
939. MMWR Morb Mortal Wkly Rep 2009 Oct 23;58(41):1143-6.
940. Emerg Infect Dis 2011 Sep ;17(9):1744-6.
941. Vaccine 2011 Apr 27;29(19):3617-22.
942. J Infect Dis 2012 Dec ;206 Suppl 1:S68-73.
943. BMC Public Health 2010 ;10:322.
944. Southeast Asian J Trop Med Public Health 2010 Sep ;41(5):1104-15.
945. Southeast Asian J Trop Med Public Health 2012 Jan ;43(1):68-77.
946. J Med Liban 2012 Apr-Jun;60(2):70-6.
947. ProMED <promedmail.org> archive: 20090611.2150
948. Medicina (Kaunas) 2011 ;47(1):11-8.
949. Emerg Infect Dis 2011 Mar ;17(3):403-11.
950. Chin Med J (Engl) 2010 Oct ;123(19):2651-4.
951. PLoS One 2012 ;7(5):e37067.
952. Epidemiol Infect 2012 Jul 20;:1-6.
953. J Infect Dis 2012 Dec ;206 Suppl 1:S5-S13.
954. J Infect Dis 2012 Dec ;206 Suppl 1:S140-7.
955. Epidemiol Infect 2013 Jan 18;:1-2.
956. ProMED <promedmail.org> archive: 20091113.3933
957. Med J Malaysia 2009 Jun ;64(2):105-7.
958. Prev Med 2010 Jul ;51(1):92-3.
959. Vaccine 2010 Jun 17;28(28):4499-505.
960. J Behav Med 2011 Feb ;34(1):23-31.
961. J Community Health 2010 Dec ;35(6):676-82.
962. J Infect 2010 Nov ;61(5):440-2.
963. Int J Behav Med 2011 Jun ;18(2):112-21.
964. Emerg Infect Dis 2011 Apr ;17(4):708-10.
965. Clin Microbiol Infect 2012 Oct ;18(10):976-81.
966. Bull Soc Pathol Exot 2011 May ;104(2):119-24.
967. MMWR Morb Mortal Wkly Rep 2009 May 8;58(17):467-70.
968. MMWR Morb Mortal Wkly Rep 2009 Jun 5;58(21):585-9.
969. Wkly Epidemiol Rec 2009 Jun 5;84(23):213-9.
970. Nature 2009 Jun 25;459(7250):1122-5.
971. Nature 2009 Jun 18;459(7249):931-9.
972. N Engl J Med 2009 Aug 13;361(7):674-9.
973. N Engl J Med 2009 Aug 13;361(7):680-9.
974. Euro Surveill 2009 Jul 2;14(26)
975. Influenza Other Respi Viruses 2009 Sep ;3(5):215-22.
976. PLoS One 2009 ;4(9):e6895.
977. J Infect Dev Ctries 2009 ;3(5):327-30.
978. Health Res Policy Syst 2009 ;7:21.
979. JAMA 2009 Nov 4;302(17):1880-7.
980. Lancet 2009 Dec 19;374(9707):2032-3.
981. Lancet 2009 Dec 19;374(9707):2072-9.
982. Salud Publica Mex 2009 Sep-Oct;51(5):361-71.
983. PLoS Curr 2009 ;1:RRN1129.
984. Emerg Infect Dis 2010 Jan ;16(1):27-34.
985. Cell 2009 Dec 24;139(7):1203-5.
986. Arch Med Res 2009 Nov ;40(8):669-72.
987. Arch Med Res 2009 Nov ;40(8):681-6.
988. Arch Med Res 2009 Nov ;40(8):705-11.
989. PLoS One 2010 ;5(5):e10658.
990. Thorax 2010 Jun ;65(6):505-9.
991. Salud Publica Mex 2010 Jul-Aug;52(4):288-9.
992. Emerg Infect Dis 2010 Aug ;16(8):1292-5.
993. PLoS One 2010 ;5(10):e13256.
994. Am J Prev Med 2010 Nov ;39(5):395-402.
995. Rev Invest Clin 2010 Jul-Aug;62(4):289-98.
996. Math Biosci Eng 2011 Jan ;8(1):223-38.
997. Salud Publica Mex 2011 Mar-Apr;53(2):105-6.
998. PLoS Med 2011 May ;8(5):e1000436.
999. Int J Infect Dis 2011 Nov ;15(11):e781-6.
1000. PLoS One 2011 ;6(8):e23853.
1001. Clin Infect Dis 2011 Nov ;53(10):985-93.
1002. Arch Med Res 2011 Oct ;42(7):627-32.
1003. PLoS Curr 2012 ;4:RRN1306.
1004. J Infect Dev Ctries 2012 ;6(4):302-10.
1005. Rev Panam Salud Publica 2012 Apr ;31(4):269-74.
1006. PLoS One 2012 ;7(7):e41069.
1007. Politics Life Sci 2012 ;31(1-2):52-67.
1008. ProMED <promedmail.org> archive: 20090425.1557
1009. ProMED <promedmail.org> archive: 20090703.2391
1010. ProMED <promedmail.org> archive: 20091110.3893
1011. Influenza Other Respi Viruses 2012 Feb 13;
1012. ProMED <promedmail.org> archive: 20100117.0194
1013. Int Arch Med 2010 ;3:26.
1014. Pathol Biol (Paris) 2012 Jun 22;
1015. J Infect Dis 2012 Dec ;206 Suppl 1:S94-S100.
1016. Emerg Infect Dis 2010 Sep ;16(9):1366-72.
1017. J Health Popul Nutr 2010 Dec ;28(6):537-44.
1018. Southeast Asian J Trop Med Public Health 2012 Jul ;43(4):871-6.
1019. Jpn J Infect Dis 2011 ;64(5):377-81.
1020. Emerg Infect Dis 2012 Jul ;18(7):1058-64.
1021. Influenza Other Respi Viruses 2012 Nov 5;
1022. Virol J 2011 ;8:133.
1023. J Nepal Health Res Counc 2010 Oct ;8(2):75-7.
1024. Jpn J Infect Dis 2011 ;64(5):444-5.
1025. ProMED <promedmail.org> archive: 20091016.3563
1026. J Clin Virol 2009 Jul ;45(3):179-84.
1027. Euro Surveill 2010 Jan 14;15(2)
1028. Eur J Public Health 2012 Feb ;22(1):150-7.
1029. Euro Surveill 2009 Jul 9;14(27)
1030. BMC Public Health 2011 ;11:2.
1031. Clin Vaccine Immunol 2011 Mar ;18(3):469-76.
1032. Euro Surveill 2011 ;16(7)
1033. Eur J Epidemiol 2011 Mar ;26(3):195-201.
1034. Influenza Other Respi Viruses 2011 Nov ;5(6):e513-20.
1035. BMC Public Health 2011 ;11:758.
1036. PLoS One 2012 ;7(2):e31197.
1037. Influenza Other Respi Viruses 2012 Feb 28;
1038. Vaccine 2012 Dec 13;
1039. Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz 2013 Jan ;56(1):67-75.
1040. N Z Med J 2009 Jul 3;122(1298):11-6.
1041. Euro Surveill 2009 ;14(34)
1042. MMWR Morb Mortal Wkly Rep 2009 Aug 28;58(33):918-21.
1043. N Z Med J 2009 Aug 21;122(1301):66-9.
1044. Crit Care Resusc 2009 Sep ;11(3):170-2.
1045. N Z Med J 2009 Jul 24;122(1299):73-7.
1046. Euro Surveill 2009 ;14(44)
1047. Emerg Infect Dis 2010 Jan ;16(1):100-2.
1048. Am J Respir Crit Care Med 2010 Feb 15;181(4):300-6.
1049. N Z Med J 2010 Apr 9;123(1312):45-53.
1050. Crit Care Resusc 2010 Jun ;12(2):121-30.
1051. Pediatrics 2011 Jan ;127(1):e156-63.
1052. Euro Surveill 2010 ;15(24)
1053. PLoS One 2010 ;5(10):e13211.
1054. Euro Surveill 2011 ;16(6)
1055. Crit Care 2011 ;15(3):R143.
1056. Anaesth Intensive Care 2011 May ;39(3):384-91.
1057. Emerg Infect Dis 2012 Jan ;18(1):71-7.
1058. Emerg Infect Dis 2012 May ;18(5):866-8.
1059. N Z Med J 2012 ;125(1365):54-66.
1060. N Z Med J 2012 ;125(1365):16-20.
1061. Emerg Infect Dis 2013 Jan ;19(1):92-101.
1062. ProMED <promedmail.org> archive: 20101021.3812
1063. ProMED <promedmail.org> archive: 20100804.2632
1064. ProMED <promedmail.org> archive: 20100618.2042
1065. ProMED <promedmail.org> archive: 20090827.3022
1066. ProMED <promedmail.org> archive: 20090828.3032
1067. Clin Infect Dis 2010 Jun 1;50(11):1462-7.
1068. J Infect Dis 2012 Dec ;206 Suppl 1:S121-8.
1069. ProMED <promedmail.org> archive: 20091112.3920
1070. Euro Surveill 2010 Mar 4;15(9)
1071. Euro Surveill 2010 ;15(31)
1072. PLoS One 2012 ;7(1):e30018.
1073. BMC Infect Dis 2012 Mar 20;12(1):63.
1074. Int J Infect Dis 2012 Apr 19;
1075. Oman Med J 2012 May ;27(3):201-6.
1076. J Pak Med Assoc 2010 Apr ;60(4):250-2.
1077. J Pak Med Assoc 2010 Apr ;60(4):329-30.
1078. PLoS One 2012 ;7(8):e41866.
1079. ProMED <promedmail.org> archive: 20091115.3945
1080. J Infect Dev Ctries 2009 ;3(11):811-6.
1081. J Infect Dev Ctries 2011 Sep ;5(9):664-8.
1082. Euro Surveill 2009 ;14(32)
1083. Euro Surveill 2009 ;14(42)
1084. PLoS One 2010 ;5(7):e11719.
1085. Influenza Other Respi Viruses 2009 Nov ;3(6):253-6.
1086. PLoS One 2011 ;6(6):e21287.
1087. Przegl Epidemiol 2011 ;65(2):199-203.
1088. Adv Exp Med Biol 2013 ;756:271-83.
1089. Rev Port Pneumol 2010 Nov-Dec;16(6):880-6.
1090. Rev Port Pneumol 2010 Nov-Dec;16(6):870-9.
1091. J Infect Dev Ctries 2011 Sep ;5(9):658-63.
1092. Euro Surveill 2012 ;17(27)
1093. Vaccine 2011 Mar 3;29(11):2206-11.
1094. Scand J Infect Dis 2010 Apr ;42(4):311-4.
1095. J Prev Med Public Health 2010 Mar ;43(2):109-16.
1096. J Prev Med Public Health 2010 May ;43(3):274-8.
1097. J Prev Med Public Health 2010 Mar ;43(2):99-104.
1098. Pediatr Pulmonol 2010 Oct ;45(10):1014-20.
1099. Korean J Radiol 2010 Jul-Aug;11(4):417-24.
1100. Vaccine 2011 Feb 4;29(7):1395-8.
1101. J Korean Med Sci 2011 Jan ;26(1):22-7.
1102. Int J Tuberc Lung Dis 2011 Feb ;15(2):270-5, i.
1103. Am J Infect Control 2011 Aug 24;

1104. BMC Res Notes 2011 ;4:351.
 1105. J Korean Med Sci 2012 Apr ;27(4):408-15.
 1106. Epidemiol Infect 2012 Aug 1 ;1-10.
 1107. Scand J Infect Dis 2012 Dec 14;
 1108. Euro Surveill 2009 ;14(42)
 1109. Clin Microbiol Infect 2010 Apr ;16(4):303.
 1110. Clin Microbiol Infect 2010 Apr ;16(4):309-16.
 1111. Clin Microbiol Infect 2010 Apr ;16(4):304-8.
 1112. Presse Med 2010 Jul-Aug;39(7-8):e147-57.
 1113. PLoS One 2010 ;5(5):e10896.
 1114. Ann Fr Anesth Reanim 2010 Dec ;29(12):902-8.
 1115. Bull Soc Pathol Exot 2011 May ;104(2):108-13.
 1116. PLoS Curr 2010 ;2:RRN1145.
 1117. Bull Soc Pathol Exot 2011 May ;104(2):105-7.
 1118. Bull Soc Pathol Exot 2011 May ;104(2):125-34.
 1119. Emerg Infect Dis 2011 Jan ;17(1):140-1.
 1120. Bull Soc Pathol Exot 2011 May ;104(2):97-104.
 1121. PLoS One 2012 ;7(9):e44755.
 1122. Epidemiol Health 2012 ;34:e2012009.
 1123. Vector Borne Zoonotic Dis 2010 Nov ;10(9):935-8.
 1124. Zh Mikrobiol Epidemiol Immunobiol 2010 Jan-Feb;(1):3-9.
 1125. Vopr Virusol 2011 Jan-Feb;56(1):44-9.
 1126. Zh Mikrobiol Epidemiol Immunobiol 2011 Jan-Feb;(1):26-34.
 1127. Vopr Virusol 2011 Mar-Apr;56(2):4-9.
 1128. Infect Genet Evol 2011 Dec ;11(8):2107-12.
 1129. Zh Mikrobiol Epidemiol Immunobiol 2011 May-Jun;(3):14-20.
 1130. Vestn Ross Akad Med Nauk 2011 ;(7):30-6.
 1131. Zh Mikrobiol Epidemiol Immunobiol 2011 Jul-Aug;(4):24-7.
 1132. Virol Sin 2011 Oct ;26(5):306-14.
 1133. ProMED <promedmail.org> archive: 20091110.3893
 1134. PLoS One 2012 ;7(6):e31572.
 1135. J Infect Dis 2012 Dec ;206 Suppl 1:S74-9.
 1136. ProMED <promedmail.org> archive: 20091112.3920
 1137. Lancet 2009 Nov 21;374(9703):1724.
 1138. Lancet 2009 Nov 21;374(9703):1786-91.
 1139. Saudi Med J 2009 Dec ;30(12):1532-6.
 1140. Science 2009 Nov 13;326(5955):938-40.
 1141. Ann Saudi Med 2010 Jan-Feb;30(1):59-62.
 1142. Ann Saudi Med 2010 Jan-Feb;30(1):11-4.
 1143. Lancet 2010 Jan 16;375(9710):199-200.
 1144. BMC Infect Dis 2010 ;10:42.
 1145. J Travel Med 2010 Mar-Apr;17(2):75-81.
 1146. J R Soc Med 2010 Oct ;103(10):386.
 1147. Saudi Med J 2011 Jun ;32(6):598-602.
 1148. J Travel Med 2012 Jan-Feb;19(1):15-21.
 1149. Saudi Med J 2012 Jan ;33(1):98.
 1150. J Travel Med 2012 May ;19(3):163-168.
 1151. ProMED <promedmail.org> archive: 20091120.3997
 1152. ProMED <promedmail.org> archive: 20091122.4013
 1153. Euro Surveill 2009 Apr 30;14(17)
 1154. Euro Surveill 2009 May 28;14(21)
 1155. BMJ 2009 ;338:b2470.
 1156. Euro Surveill 2010 ;15(24)
 1157. Rural Remote Health 2010 Jul-Sep;10(3):1413.
 1158. Euro Surveill 2011 ;16(2)
 1159. Euro Surveill 2011 ;16(20):19871.
 1160. PLoS One 2011 ;6(6):e20358.
 1161. BMC Infect Dis 2011 ;11:192.
 1162. J Clin Virol 2011 Dec ;52(4):300-3.
 1163. J Environ Public Health 2011 ;2011:407505.
 1164. J Gen Virol 2012 Feb 15;
 1165. Lancet Infect Dis 2012 Sep ;12(9):696-702.
 1166. Lancet Infect Dis 2012 Sep ;12(9):653-4.
 1167. Croat Med J 2011 Apr 15;52(2):141-50.
 1168. Srp Arh Celok Lek 2012 Nov-Dec;140(11-12):751-5.
 1169. Emerg Infect Dis 2010 Jan ;16(1):21-6.
 1170. JAMA 2010 Apr 14;303(14):1383-91.
 1171. Ann Acad Med Singapore 2010 Apr ;39(4):328-5.
 1172. Ann Acad Med Singapore 2010 Apr ;39(4):325-3.
 1173. Ann Acad Med Singapore 2010 Apr ;39(4):313-12.
 1174. Ann Acad Med Singapore 2010 Apr ;39(4):307-6.
 1175. Ann Acad Med Singapore 2010 Apr ;39(4):303-4.
 1176. Ann Acad Med Singapore 2010 Apr ;39(4):291-4.
 1177. Ann Acad Med Singapore 2010 Apr ;39(4):283-8.
 1178. Ann Acad Med Singapore 2010 Apr ;39(4):273-10.
 1179. Ann Acad Med Singapore 2010 Apr ;39(4):265-2.
 1180. Singapore Med J 2009 Jun ;50(6):581-3.
 1181. Br J Sports Med 2010 Jun ;44(7):528-32.
 1182. Emerg Infect Dis 2010 Oct ;16(10):1554-61.
 1183. PLoS One 2011 ;6(3):e17468.
 1184. Emerg Infect Dis 2011 Aug ;17(8):1455-62.
 1185. Influenza Other Respi Viruses 2011 Nov ;5(6):e563-7.
 1186. Expert Rev Anti Infect Ther 2012 Jul ;10(7):751-60.
 1187. Emerg Infect Dis 2013 Jan ;19(1):92-101.
 1188. ProMED <promedmail.org> archive: 20101021.3812
 1189. ProMED <promedmail.org> archive: 20100530.1795
 1190. Wien Klin Wochenschr 2012 Feb 15;
 1191. Euro Surveill 2009 ;14(42)
 1192. QJM 2010 May ;103(5):319-25.
 1193. PLoS One 2012 ;7(11):e49482.
 1194. J Infect Dis 2012 Dec ;206 Suppl 1:S166-72.
 1195. J Infect Dis 2012 Dec ;206 Suppl 1:S148-53.
 1196. Euro Surveill 2009 Apr 30;14(17)
 1197. Euro Surveill 2009 May 14;14(19)
 1198. Euro Surveill 2009 ;14(49)
 1199. Med Intensiva 2010 Mar ;34(2):87-94.
 1200. Euro Surveill 2010 Feb 4;15(5)
 1201. Med Clin (Barc) 2010 Jun 5;135(1):21-2.
 1202. Med Clin (Barc) 2010 Jun 5;135(1):1-7.
 1203. Arch Bronconeumol 2010 Mar ;46 Suppl 2:32-8.
 1204. Arch Bronconeumol 2010 Mar ;46 Suppl 2:3-12.
 1205. Vaccine 2010 Jul 5;28(30):4751-7.
 1206. Med Clin (Barc) 2010 Oct 16;135(12):543-5.
 1207. Clin Microbiol Infect 2011 Jun ;17(6):845-50.
 1208. Crit Care 2009 ;13(5):196.
 1209. AIDS 2010 Oct 23;24(16):2461-7.
 1210. Clin Microbiol Infect 2011 May ;17(5):738-46.
 1211. Vet Microbiol 2011 Apr 21;149(1-2):56-63.
 1212. Chest 2011 Mar ;139(3):555-62.
 1213. Vaccine 2011 Feb 1;29(6):1332-8.
 1214. Respirology 2011 Jan ;16(1):78-85.
 1215. Rev Esp Salud Publica 2010 Sep-Oct;84(5):657-63.
 1216. Rev Esp Salud Publica 2010 Sep-Oct;84(5):569-88.
 1217. Crit Care Med 2011 May ;39(5):945-51.
 1218. Gac Sanit 2011 Jan-Feb;25(1):23-8.
 1219. Gac Sanit 2011 Jul-Aug;25(4):296-302.
 1220. Rev Esp Salud Publica 2011 Jan-Feb;85(1):73-80.
 1221. Rev Esp Salud Publica 2011 Jan-Feb;85(1):1-2.
 1222. Influenza Other Respi Viruses 2011 Nov ;5(6):e544-51.
 1223. Medicine (Baltimore) 2011 Sep ;90(5):328-36.
 1224. Enferm Infecc Microbiol Clin 2012 Feb ;30(2):60-3.
 1225. J Clin Virol 2012 Jan ;53(1):16-21.
 1226. Eur J Public Health 2012 Dec ;22(6):792-6.
 1227. Clin Microbiol Infect 2012 Mar ;18(3):E55-62.
 1228. PLoS One 2012 ;7(3):e33139.
 1229. Eur J Pediatr 2012 Mar 21;
 1230. J Antimicrob Chemother 2012 Mar 30;
 1231. J Am Geriatr Soc 2012 Mar 29;
 1232. Enferm Infecc Microbiol Clin 2012 Oct ;30S4:2-9.
 1233. Pediatr Emerg Care 2013 Jan ;29(1):49-52.
 1234. Influenza Other Respi Viruses 2012 Nov 8;
 1235. Euro Surveill 2009 Jun 18;14(24)
 1236. Euro Surveill 2009 Sep 17;14(37)
 1237. Euro Surveill 2011 ;16(18)
 1238. Health Policy 2012 Oct 5;
 1239. Swiss Med Wkly 2009 Dec 26;139(51-52):730.
 1240. Swiss Med Wkly 2011 ;141:w13177.
 1241. Eur J Epidemiol 2011 Mar ;26(3):203-10.
 1242. PLoS One 2012 ;7(11):e49806.
 1243. Avicenna J Med 2012 Apr ;2(2):34-7.
 1244. Avicenna J Med 2012 Apr ;2(2):38-9.
 1245. J Infect 2010 Feb ;60(2):168-74.
 1246. Pediatr Neonatol 2010 Apr ;51(2):83-8.
 1247. Euro Surveill 2010 May 27;15(21):19575.
 1248. BMC Public Health 2010 ;10:322.
 1249. Emerg Infect Dis 2010 Aug ;16(8):1309-11.
 1250. Clin Vaccine Immunol 2010 Dec ;17(12):1958-62.
 1251. Clin Infect Dis 2010 Dec 15;51(12):1465-7.
 1252. Emerg Infect Dis 2011 Jan ;17(1):76-8.
 1253. J Chin Med Assoc 2011 Jul ;74(7):298-304.
 1254. PLoS One 2011 ;6(9):e24440.
 1255. BMC Infect Dis 2011 ;11:332.
 1256. PLoS One 2012 ;7(4):e36120.
 1257. ProMED <promedmail.org> archive: 20120426.1115182
 1258. ProMED <promedmail.org> archive: 20091001.3421
 1259. Euro Surveill 2009 Aug 6;14(31)
 1260. J Infect 2010 May ;60(5):402-3.
 1261. PLoS One 2010 ;5(3):e9717.
 1262. BMC Public Health 2010 ;10:322.
 1263. Asian Pac J Allergy Immunol 2010 Mar ;28(1):67-75.
 1264. Vaccine 2010 Jul 26;28(33):5437-44.
 1265. Infect Control Hosp Epidemiol 2010 Aug ;31(8):854-6.
 1266. Infect Control Hosp Epidemiol 2009 Dec ;30(12):1236-7.
 1267. Jpn J Infect Dis 2010 Jul ;63(4):251-6.
 1268. Clin Infect Dis 2010 Aug 1;51(3):368-9.
 1269. Emerg Infect Dis 2010 Sep ;16(9):1366-72.
 1270. PLoS One 2011 ;6(1):e16164.
 1271. J Health Popul Nutr 2010 Dec ;28(6):537-44.
 1272. Southeast Asian J Trop Med Public Health 2012 Jul ;43(4):871-6.
 1273. Jpn J Infect Dis 2011 ;64(5):377-81.
 1274. Emerg Infect Dis 2012 Jul ;18(7):1058-64.
 1275. Influenza Other Respi Viruses 2012 Nov 5;
 1276. PLoS One 2012 ;7(12):e51275.
 1277. PLoS One 2013 ;8(2):e54946.
 1278. Euro Surveill 2009 ;14(32)
 1279. Jpn J Infect Dis 2010 Jul ;63(4):239-45.
 1280. BMC Infect Dis 2010 ;10:281.
 1281. Vaccine 2010 Dec 16;29(2):329-33.
 1282. Pediatr Crit Care Med 2012 Jan ;13(1):e11-7.
 1283. BMC Infect Dis 2011 ;11:87.
 1284. BJOG 2011 Sep ;118(10):1216-22.
 1285. Scand J Infect Dis 2011 Dec ;43(11-12):923-9.
 1286. Jpn J Infect Dis 2012 ;65(1):13-8.
 1287. Mikrobiyol Bul 2012 Oct ;46(4):575-93.
 1288. BMC Infect Dis 2013 Jan 7;13(1):6.
 1289. ProMED <promedmail.org> archive: 20091031.3764
 1290. ProMED <promedmail.org> archive: 20091101.3771
 1291. ProMED <promedmail.org> archive: 20091102.3781

1292. ProMED <promedmail.org> archive: 20091103.3799
 1293. ProMED <promedmail.org> archive: 20091105.3827
 1294. ProMED <promedmail.org> archive: 20091107.3858
 1295. ProMED <promedmail.org> archive: 20091110.3893
 1296. ProMED <promedmail.org> archive: 20091113.3931
 1297. ProMED <promedmail.org> archive: 20091116.3959
 1298. ProMED <promedmail.org> archive: 20091117.3970
 1299. Asia Pac J Public Health 2010 Jul ;22(3 Suppl):195-245.
 1300. Emerg Infect Dis 2011 Feb ;17(2):292-5.
 1301. Med Princ Pract 2011 ;20(1):97-9.
 1302. East Mediterr Health J 2012 Jan ;18(1):31-6.
 1303. BMJ 2009 ;338:b2470.
 1304. Euro Surveill 2009 Apr 30;14(17)
 1305. Euro Surveill 2009 May 28;14(21)
 1306. Euro Surveill 2009 Jun 4;14(22)
 1307. Euro Surveill 2009 Jul 9;14(27)
 1308. BMJ 2009 ;338:b2094.
 1309. BMJ 2009 ;339:b2829.
 1310. Euro Surveill 2009 Jul 9;14(27)
 1311. BMJ 2009 ;339:b2897.
 1312. Euro Surveill 2009 ;14(33)
 1313. BMJ 2009 ;339:b3403.
 1314. BMJ 2009 ;339:b2829.
 1315. Epidemiol Infect 2010 Feb ;138(2):183-91.
 1316. Crit Care 2009 ;13(6):426; author reply 426.
 1317. Lancet 2010 Mar 27;375(9720):1100-8.
 1318. BMJ 2009 ;339:b5213.
 1319. PLoS Curr 2009 ;1:RRN1130.
 1320. Clin Med 2009 Dec ;9(6):534-8.
 1321. Euro Surveill 2010 Jan 21;15(3)
 1322. Euro Surveill 2010 May 20;15(20)
 1323. Epidemiol Infect 2010 Nov ;138(11):1531-41.
 1324. Arch Dis Child 2011 Sep ;96(9):857-9.
 1325. Epidemiol Infect 2011 Oct ;139(10):1560-9.
 1326. J Clin Virol 2011 Feb ;50(2):183.
 1327. J Epidemiol Community Health 2011 Oct ;65(10):941-6.
 1328. Health Technol Assess 2010 Dec ;14(55):83-114.
 1329. Euro Surveill 2011 ;16(2)
 1330. Euro Surveill 2011 ;16(20):19871.
 1331. PLoS One 2011 ;6(6):e20358.
 1332. BMC Infect Dis 2011 ;11:192.
 1333. J Clin Virol 2011 Dec ;52(4):300-3.
 1334. J Environ Public Health 2011 ;2011:407505.
 1335. J Gen Virol 2012 Feb 15;
 1336. Lancet Infect Dis 2012 Sep ;12(9):696-702.
 1337. Lancet Infect Dis 2012 Sep ;12(9):653-4.
 1338. Euro Surveill 2011 ;16(3)
 1339. Euro Surveill 2011 ;16(3)
 1340. Vaccine 2011 Mar 21;29(14):2613-8.
 1341. Euro Surveill 2011 ;16(5)
 1342. PLoS One 2011 ;6(2):e17074.
 1343. Epidemiol Infect 2011 Sep ;139(9):1431-9.
 1344. Emerg Infect Dis 2011 Apr ;17(4):592-8.
 1345. Epidemiol Infect 2012 Jan ;140(1):100-5.
 1346. Influenza Other Respi Viruses 2011 Nov ;5(6):e504-12.
 1347. PLoS One 2011 ;6(8):e23779.
 1348. BMJ 2011 ;343:d5408.
 1349. Epidemiol Infect 2011 Nov 11;:1-9.
 1350. Vaccine 2012 Feb 8;30(7):1371-8.
 1351. Int J Public Health 2012 Feb 7;
 1352. Euro Surveill 2012 ;17(14)
 1353. J Paediatr Child Health 2012 Dec 16;
 1354. ProMED <promedmail.org> archive: 20091210.4208
 1355. ProMED <promedmail.org> archive: 20090806.2784
 1356. MMWR Morb Mortal Wkly Rep 2009 May 1;58(16):435-7.
 1357. MMWR Morb Mortal Wkly Rep 2009 May 1;58(16):431-3.
 1358. MMWR Morb Mortal Wkly Rep 2009 Sep 18;58(36):1009-12.
 1359. MMWR Morb Mortal Wkly Rep 2009 Oct 2;58(38):1071-4.
 1360. N Engl J Med 2009 Nov 12;361(20):1935-44.
 1361. Ann Emerg Med 2009 Nov ;54(5):732-6.
 1362. MMWR Morb Mortal Wkly Rep 2009 Nov 13;58(44):1236-41.
 1363. Emerg Infect Dis 2009 Dec ;15(12):2004-7.
 1364. PLoS Med 2009 Dec ;6(12):e1000207.
 1365. PLoS Curr 2009 ;1:RRN1127.
 1366. PLoS Curr 2009 ;1:RRN1042.
 1367. N Engl J Med 2009 Dec 31;361(27):2619-27.
 1368. MMWR Morb Mortal Wkly Rep 2010 Jan 8;58(51):1436-40.
 1369. Clin Infect Dis 2010 Feb 15;50(4):528-30.
 1370. Clin Infect Dis 2010 Feb 15;50(4):523-7.
 1371. MMWR Morb Mortal Wkly Rep 2010 Jan 22;59(2):44-8.
 1372. MMWR Morb Mortal Wkly Rep 2010 Jan 22;59(2):38-43.
 1373. MMWR Morb Mortal Wkly Rep 2010 Jan 29;59(3):74-7.
 1374. PLoS Curr 2010 ;2:RRN1148.
 1375. Emerg Infect Dis 2010 Mar ;16(3):504-6.
 1376. PLoS Curr 2010 ;2:RRN1153.
 1377. MMWR Morb Mortal Wkly Rep 2010 Apr 16;59(14):423-30.
 1378. Mem Inst Oswaldo Cruz 2010 Mar ;105(2):179-83.
 1379. Am J Pathol 2010 Jul ;177(1):166-75.
 1380. PLoS One 2010 ;5(5):e10722.
 1381. Am J Pathol 2010 Jul ;177(1):166-75.
 1382. BMC Infect Dis 2010 ;10:162.
 1383. PLoS One 2010 ;5(7):e11601.
 1384. PLoS One 2010 ;5(7):e11677.
 1385. MMWR Morb Mortal Wkly Rep 2010 Jul 30;59(29):901-8.
 1386. Stud Health Technol Inform 2010 ;160(Pt 1):447-51.
 1387. MMWR Morb Mortal Wkly Rep 2010 Dec 3;59(47):1541-5.
 1388. J Occup Environ Med 2010 Dec ;52(12):1212-6.
 1389. Am J Public Health 2011 Feb ;101(2):285-93.
 1390. Am J Emerg Med 2012 Feb ;30(2):275-82.
 1391. MMWR Morb Mortal Wkly Rep 2011 Jan 21;60(2):37-41.
 1392. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S90-3.
 1393. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S83-9.
 1394. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S75-82.
 1395. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S69-74.
 1396. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S60-8.
 1397. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S27-35.
 1398. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S198-204.
 1399. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S13-26.
 1400. Clin Infect Dis 2011 Jan 1;52 Suppl 1:S1-3.
 1401. JAMA 2011 Mar 16;305(11):1080, 1082-3.
 1402. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S7-12.
 1403. Am J Disaster Med 2011 Jan-Feb;6(1):23-30.
 1404. Am J Public Health 2011 Jul ;101(7):1252-5.
 1405. Vaccine 2011 Jul 18;29(32):5284-9.
 1406. Am J Obstet Gynecol 2011 Jun ;204(6 Suppl 1):S7-12.
 1407. Influenza Other Respi Viruses 2011 Sep ;5(5):321-7.
 1408. Ann Epidemiol 2011 Aug ;21(8):623-30.
 1409. Emerg Infect Dis 2011 Sep ;17(9):1685-91.
 1410. Influenza Other Respi Viruses 2012 Sep ;6(5):305-8.
 1411. J Health Care Poor Underserved 2011 ;22(4 Suppl):39-60.
 1412. Value Health 2012 Jan ;15(1):158-66.
 1413. Emerg Infect Dis 2012 Feb ;18(2):308-311.
 1414. Influenza Other Respi Viruses 2012 Feb 21;
 1415. MMWR Morb Mortal Wkly Rep 2012 Feb 24;61:113-8.
 1416. Clin Infect Dis 2012 Mar 20;
 1417. Emerg Infect Dis 2012 Apr ;18(4):556-62.
 1418. Clin Obstet Gynecol 2012 Jun ;55(2):487-97.
 1419. Am J Epidemiol 2012 May 11;
 1420. Influenza Other Respi Viruses 2012 Nov ;6(6):e134-e142.
 1421. Influenza Other Respi Viruses 2012 Nov ;6(6):e129-33.
 1422. Influenza Other Respi Viruses 2012 Nov ;6(6):e169-e177.
 1423. PLoS One 2012 ;7(8):e40984.
 1424. J Infect Dis 2012 Nov ;206(9):1350-1358.
 1425. PLoS One 2012 ;7(10):e48187.
 1426. Med Care 2013 Jan 3;
 1427. ProMED <promedmail.org> archive: 20090501.1646
 1428. ProMED <promedmail.org> archive: 20090422.1516
 1429. ProMED <promedmail.org> archive: 20090424.1541
 1430. ProMED <promedmail.org> archive: 20090425.1552
 1431. ProMED <promedmail.org> archive: 20090426.1566
 1432. ProMED <promedmail.org> archive: 20090507.1709
 1433. Invest Clin 2009 Sep ;50(3):279-81.
 1434. ProMED <promedmail.org> archive: 20091105.3820
 1435. BMC Med 2009 ;7:43.
 1436. PLoS Med 2010 May ;7(5):e1000277.
 1437. BMC Public Health 2010 ;10:322.
 1438. MMWR Morb Mortal Wkly Rep 2011 Jun 17;60(23):781-5.
 1439. ProMED <promedmail.org> archive: 20090708.2450
 1440. ProMED <promedmail.org> archive: 20090809.2819
 1441. ProMED <promedmail.org> archive: 20090728.2655
 1442. Presse Med 2011 Mar ;40(3):e145-51.
 1443. J Infect Dis 2012 Dec ;206 Suppl 1:S173-7.
 1444. Euro Surveill 2009 May 14;14(19)
 1445. MMWR Morb Mortal Wkly Rep 2009 May 1;58(16):431-3.
 1446. Euro Surveill 2009 May 14;14(19)
 1447. Euro Surveill 2009 May 7;14(18)
 1448. MMWR Morb Mortal Wkly Rep 2009 May 8;58(17):453-8.
 1449. Euro Surveill 2009 May 7;14(18)
 1450. N Engl J Med 2009 Jun 18;360(25):2605-15.
 1451. Euro Surveill 2009 May 21;14(20)
 1452. Ann Intern Med 2009 Jul 7;151(1):59-62.
 1453. Wkly Epidemiol Rec 2009 May 15;84(20):173-9.
 1454. Wkly Epidemiol Rec 2009 May 15;84(20):173-9.
 1455. Euro Surveill 2009 Jun 11;14(23):19238.
 1456. J Clin Virol 2009 Jul ;45(3):174-8.
 1457. Euro Surveill 2009 Jun 18;14(24)
 1458. Wkly Epidemiol Rec 2009 Jun 19;84(25):249-57.
 1459. BMJ 2009 ;338:b2470.
 1460. Euro Surveill 2009 Jul 23;14(29)
 1461. Wkly Epidemiol Rec 2009 Jul 24;84(30):305-8.
 1462. BMC Infect Dis 2009 ;9:129.
 1463. Virology 2012 Jan 5;422(1):151-60.
 1464. ProMED <promedmail.org> archive: 20090426.1577
 1465. ProMED <promedmail.org> archive: 20090427.1583
 1466. ProMED <promedmail.org> archive: 20090427.1586
 1467. ProMED <promedmail.org> archive: 20090428.1609
 1468. ProMED <promedmail.org> archive: 20090429.1614
 1469. ProMED <promedmail.org> archive: 20090428.1600
 1470. ProMED <promedmail.org> archive: 20090429.1622
 1471. ProMED <promedmail.org> archive: 20090430.1638
 1472. ProMED <promedmail.org> archive: 20090428.1646
 1473. ProMED <promedmail.org> archive: 20090502.1654
 1474. ProMED <promedmail.org> archive: 20090503.1660
 1475. ProMED <promedmail.org> archive: 20090504.1675
 1476. ProMED <promedmail.org> archive: 20090505.1681
 1477. ProMED <promedmail.org> archive: 20090507.1715
 1478. ProMED <promedmail.org> archive: 20090508.1722
 1479. ProMED <promedmail.org> archive: 20090520.1895

1480. ProMED <promedmail.org> archive: 20090521.1906
1481. ProMED <promedmail.org> archive: 20090522.1921
1482. ProMED <promedmail.org> archive: 20090523.1931
1483. ProMED <promedmail.org> archive: 20090525.1945
1484. ProMED <promedmail.org> archive: 20090526.1960
1485. ProMED <promedmail.org> archive: 20090527.1972
1486. ProMED <promedmail.org> archive: 20090531.2025
1487. ProMED <promedmail.org> archive: 20090605.2089
1488. ProMED <promedmail.org> archive: 20090608.2117
1489. ProMED <promedmail.org> archive: 20090611.2166
1490. ProMED <promedmail.org> archive: 20090616.2221
1491. ProMED <promedmail.org> archive: 20121108.1400408
1492. ProMED <promedmail.org> archive: 20090619.2261
1493. ProMED <promedmail.org> archive: 20090622.2288
1494. ProMED <promedmail.org> archive: 20090625.2306
1495. ProMED <promedmail.org> archive: 20090627.2338
1496. ProMED <promedmail.org> archive: 20090701.2372
1497. ProMED <promedmail.org> archive: 20090722.2599
1498. ProMED <promedmail.org> archive: 20090820.2955
1499. ProMED <promedmail.org> archive: 20090906.3138
1500. ProMED <promedmail.org> archive: 20090911.3209
1501. ProMED <promedmail.org> archive: 20090918.3272
1502. ProMED <promedmail.org> archive: 20100108.0093

Intestinal spirochetosis

Agent	BACTERIUM. Brachyspira pilosicoli and <i>B. aalborgi</i> Anaerobic gram-negative spirochetes
Reservoir	Human Fowl Pigs
Vector	None
Vehicle	Endogenous
Incubation Period	Unknown
Diagnostic Tests	Spirochetes resemble 'brush border' on bowel biopsy; identification of <i>Brachyspira</i> by PCR
Typical Adult Therapy	Metronidazole appears to be effective in some cases.
Typical Pediatric Therapy	As for adult.
Clinical Hints	Chronic diarrhea and abdominal pain in the absence of other identifiable etiology
Synonyms	Human intestinal spirochetosis. ICD9: 009.1 ICD10: A04.8

Clinical

This diagnosis should be suspected in patients with persistent or chronic diarrhea lasting more than several weeks, in whom alternative etiologies are not identified.

- Abdominal pain, hematochezia, flatulence and intermittent constipation are also reported in some cases. ¹⁻³
- *Brachyspira* has been identified in the blood in some cases ⁴, even in the absence of intestinal disease. ⁵
- Asymptomatic infection is common. ⁶
- Intestinal spirochetosis in children may mimic inflammatory bowel disease. ⁷
- Although some patients improve following administration of Metronidazole, other cases resolve without specific therapy. ⁸

Roentgenographic studies may reveal colonic mucosal edema and luminal narrowing. ⁹

Standard H & E staining of colonic biopsies reveals a 'pseudo-brush border' consisting of tiny spirochetes ¹⁰⁻¹²; or free-floating spirochetes in the intestinal mucus. ¹³

- Similar findings are often present in asymptomatic individuals. ¹⁴
- The organism can be identified using specialized culture ¹⁵ or molecular methods. ¹⁶⁻¹⁸

This disease is endemic or potentially endemic to all countries.

References

1. Scand J Gastroenterol 2007 Dec ;42(12):1422-7.
2. Rev Soc Bras Med Trop 2005 Jan-Feb;38(1):56-7.
3. Pediatr Dev Pathol 2010 Nov-Dec;13(6):471-5.
4. J Pak Med Assoc 2009 Oct ;59(10):723-4.
5. J Clin Microbiol 2011 Oct ;49(10):3697-9.
6. J Med Microbiol 2010 Jul ;59(Pt 7):791-6.
7. BMC Pediatr 2012 ;12:163.
8. Ned Tijdschr Geneeskd 2005 Dec 17;149(51):2873-6.
9. J Gastroenterol 2007 Mar ;42(3):253-6.
10. Cesk Patol 2004 Jul ;40(3):117-20.
11. Klin Mikrobiol Infekc Lek 2004 Apr ;10(2):61-7.
12. Colorectal Dis 2002 Mar ;4(2):97-100.
13. Hum Pathol 2010 Feb ;41(2):249-54.
14. Int J Surg Pathol 2010 Apr ;18(2):144-8.
15. J Med Microbiol 2003 Jun ;52(Pt 6):509-13.
16. FEMS Microbiol Lett 2001 Apr 13;197(2):167-70.
17. J Clin Microbiol 2000 Oct ;38(10):3555-60.
18. J Clin Microbiol 1999 Jun ;37(6):2093-8.

Intra-abdominal abscess

Agent	BACTERIUM. Mixed anaerobic / aerobic, staphylococci, Neisseria gonorrhoeae , Chlamydia trachomatis, etc
Reservoir	Human
Vector	None
Vehicle	None
Incubation Period	Variable
Diagnostic Tests	Various imaging techniques (CT, Gallium scan, ultrasound, etc).
Typical Adult Therapy	Percutaneous or open drainage + antibiotics directed at known or suspected pathogen(s)
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, chills and localizing pain (e.g., chest pain in subphrenic abscess) - setting of prior surgery, biliary or colonic disease, appendicitis, vaginal discharge (PID); FUO, subdiaphragmatic gas or limited diaphragmatic motion may be present.
Synonyms	Abscess - Abdominal, Acute appendicitis, Appendicitis, Intraabdominal abscess, Intraperitoneal abscess, P.I.D., Pancreatic abscess, Pelvic abscess, Pelvic inflammatory disease, Pylephlebitis, Subhepatic abscess, Subphrenic abscess, Suppurative pancreatitis, Tuboovarian abscess. ICD9: 614,577.0 ICD10: K35,N73,K75.1,K85

Clinical

Intraabdominal abscesses often occur in the setting of prior abdominal trauma, surgery or infection.

Signs and symptoms may include fever, pain, tenderness and leucocytosis.

- In many cases, the sole presenting feature is prolonged fever, which may be accompanied by weight loss, lethargy and anemia.
- One or more localized masses may be detectable on palpation or through the use of imaging techniques.

Comprehensive reviews of clinical presentation:

- Pelvic Inflammatory Disease ¹⁻⁶
- Splenic Abscess ^{7 8}
- Pancreatic Abscess ^{9 10}
- Pylephlebitis. ¹¹

This disease is endemic or potentially endemic to all countries.

References

1. Int J STD AIDS 2005 Nov ;16(11):715-20; quiz 721.
2. Clin Evid 2004 Jun ;(11):2121-7.
3. Rev Infect Dis 1990 Jul-Aug;12 Suppl 6:S656-64.
4. Rev Infect Dis 1986 Jan-Feb;8(1):86-116.
5. Int J STD AIDS 2005 Nov ;16(11):715-20; quiz 721.
6. Expert Rev Anti Infect Ther 2006 Apr ;4(2):235-47.
7. Am J Surg 1987 Jul ;154(1):27-34.
8. Am Surg 2001 Jan ;67(1):80-5.
9. World J Surg 1990 Jul-Aug;14(4):505-11; discussion 511-2.
10. Br J Surg 1984 Feb ;71(2):141-3.
11. Am J Gastroenterol 1996 Jun ;91(6):1251-3.

Intracranial venous thrombosis

Agent	BACTERIUM. Oral anaerobes, streptococci, et al
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Culture (blood, CSF if indicated). Ophthalmoscopy. Roentgenographic studies of skull & sinuses.
Typical Adult Therapy	Antibiotic(s) directed at known or suspected pathogens
Typical Pediatric Therapy	As for adult
Clinical Hints	Headache, seizures and fever; cranial nerve dysfunction may be present; usually occurs in the setting of facial, otic or sinus infection.
Synonyms	Cavernous sinus thrombosis, Cerebral sinus thrombosis, Cortical vein thrombosis, Internal cerebral vein thrombosis, Straight sinus thrombosis, Superior sinus thrombosis, Transverse sinus thrombosis. ICD9: 325 ICD10: G08

Clinical

Cortical vein thrombosis may occasionally be clinically silent, or produce only transient neurological findings. ¹

- Septic cortical vein or venous sinus occlusion may progress to subdural empyema, meningitis, brain abscess, systemic infection or pulmonary embolism.
- Severe headache is present in 90% of cases, and cerebral lesions with neurological signs in 50%. ²
- If collateral flow is compromised, the resulting neurological may mimic brain abscess, with impairment of consciousness, focal or generalized seizures, and increased intracranial pressure.
- Depending on the site of the lesion, one may encounter hemiparesis, which involves the face and hand if veins; unilateral or bilateral leg weakness; aphasia; etc. ³

Cavernous sinus thrombosis is characterized by diplopia, photophobia, orbital edema, and progressive exophthalmos.

- Involvement of cranial nerves III, IV, V, and VI is reflected by ophthalmoplegia, fixed pupil, a loss of the corneal reflex and diminished upper facial.
- Papilledema, retinal hemorrhages, and visual loss may also occur.

Anterior superior sagittal sinus thrombosis may produce intracranial hypertension without other signs.

- More extensive blockage of this sinus is associated with bilateral leg weakness followed by arm weakness and clouding of consciousness.

Lateral sinus thrombosis causes pain over the ear and mastoid, occasionally with edema over the mastoid itself (Griesinger's sign); or ipsilateral facial pain and lateral rectus weakness (Gradenigo's syndrome).

This disease is endemic or potentially endemic to all countries.

References

1. *Neurol Sci* 2005 Feb ;25(6):311-5.
2. *N Engl J Med* 2005 Apr 28;352(17):1791-8.
3. *J Neurol* 2004 Jan ;251(1):11-23.

Isosporiasis

Agent	PARASITE - Protozoa. Sporozoa, Coccidea, Eimeriida: Isospora [Cystoisospora] belli
Reservoir	Human
Vector	None
Vehicle	Food Liquids Fecal-oral Sexual (homosexual) contact
Incubation Period	7d - 10d
Diagnostic Tests	Microscopy of stool or duodenal contents. Advise laboratory when this organism is suspected.
Typical Adult Therapy	Sulfamethoxazole/trimethoprim 800/160 mg BID X 10 days - Then BID X 3 weeks (may be indefinite in AIDS patient) Increase dosage / duration in immune-suppressed patients Pyrimethamine 50 to 75 mg per day + leucovorin if allergic to sulfa
Typical Pediatric Therapy	Sulfamethoxazole/trimethoprim 25/5 mg/kg BID X 10 days - Then BID X 3 weeks
Clinical Hints	Myalgia, watery diarrhea, nausea and leukocytosis; eosinophilia may be present; prolonged and severe in AIDS patients.
Synonyms	Isospora belli. ICD9: 007.2 ICD10: A07.3

Clinical

Isosporiasis is characterized by abdominal cramps, watery diarrhea, headache, weight loss and myalgias. ¹

- Fever and vomiting may also be present.
- A low-grade eosinophilia is present in 50% of patients
- Fecal leucocytes are not seen.

Infection in AIDS patients may cause significant weight loss and dehydration, requiring hospitalization. ²

- Disease is also more severe among patients with lymphoma and leukemia. ³
- Chronic and severe infection may occasionally affect immunocompetent patients as well, and infants and young children are most likely to suffer severe disease. ⁴
- Paralysis related to severe potassium depletion has been reported in an AIDS patient with isosporiasis. ⁵
- Biliary disease similar to primary sclerosing cholangitis has been reported. ⁶
- Disseminated extraintestinal infection has rarely been reported.

This disease is endemic or potentially endemic to all countries.

Isosporiasis in Uganda

Prevalence surveys:

13% of AIDS-related infection in this country

References

1. *Parasitology* 1998 ;117 Suppl:S143-59.
2. *AIDS Res* 1983-1984;1(5):327-38.
3. *Case Rep Infect Dis* 2012 ;2012:640104.
4. *Semin Gastrointest Dis* 1997 Jan ;8(1):33-44.
5. *J Emerg Med* 2011 Dec ;41(6):e129-32.
6. *Hum Pathol* 2009 Sep ;40(9):1342-6.

Kawasaki disease

Agent	UNKNOWN
Reservoir	Unknown
Vector	None
Vehicle	Unknown
Incubation Period	Unknown
Diagnostic Tests	Diagnosis is based on clinical criteria only.
Typical Adult Therapy	Intravenous gamma globulin 2.0 g/kg over 10 to 12h X 1 dose. Plus aspirin 100 mg/kg/day X 14d (or until defervescence) - then 5 to 10 mg/kg/day until normal ESR. Infliximab 5 mg/kg has been successful in some studies.
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, conjunctivitis, stomatitis, erythematous rash which desquamates; occasional coronary artery occlusion; the disease is most common among children; case-fatality rates of 1% to 4% are reported.
Synonyms	Kawasaki's disease, Mucocutaneous lymph node syndrome. ICD9: 446.1 ICD10: M30.3

Clinical

Diagnostic criteria: ^{1 2}

Fever for at least five days in addition to at least 4 of the following:

1. Changes in the oral mucosa (erythema, strawberry tongue, etc)
2. Changes in hands and feet (erythema, swelling, periungual desquamation, rarely gangrene ³)
3. Rash, primarily on trunk (maculopapular, scarlatiniform, erythema multiforme).
4. Cervical lymphadenopathy ⁴
5. Absence of other etiology.

Kawasaki disease is encountered among adults ⁵⁻⁷ as well as children.

- The incidence of specific diagnostic criteria are roughly similar in both groups
- Cheilitis, meningitis, and thrombocytosis are more common in children. Rare instances of thrombocytopenia are also reported ⁸
- Arthralgia is common, and may involve one or multiple joints ⁹
- Arthralgia, adenopathy, and liver function abnormality ^{10 11} are more common in adults. ¹²
- Older children may have a more marked inflammatory response and worse outcome, as compared to young children. ¹³
- Absence of fever ¹⁴ , acute hepatitis ¹⁵ , pleural effusion, disseminated intravascular coagulopathy ¹⁶ , pancreatitis ^{17 18} and cholestasis have been reported in some cases. ¹⁹
- Rare instances of recurrent Kawasaki disease have been reported. ^{20 21}

There is no diagnostic test for Kawasaki disease.

Atypical or Incomplete Kawasaki Disease:

As many as 23% of patients may present with "incomplete (atypical) Kawasaki disease" characterized by fever ≥ 5 days and the presence of < 4 "classic signs." ²²⁻²⁵

- The clinical picture in atypical Kawasaki disease may be dominated by one unusual finding: seizure, bloody diarrhea, nephrotic syndrome, hyponatremia or compressive cervical lymphadenopathy. ²⁶
- Of 232,263 cases reported in Japan during 2007 to 2008, 80% had classic clinical findings and 20% had "incomplete" Kawasaki disease. ²⁷
- Occasionally, the initial presentation of Kawasaki disease may be limited to erythema multiforme ²⁸ or fever with cervical lymphadenopathy. ²⁹
- Patients with incomplete and atypical KS are more likely than those with other febrile diseases to present with mucosal changes, conjunctivitis, extremity abnormalities, perineal desquamation ³⁰ , and later development of coronary artery abnormalities. ³¹

The appearance of redness or crusting at a BCG inoculation site is a valuable predictive sign for Kawasaki disease. ³²

- Orange-brown discoloration of nails (chromonychia) is a common finding in some series. ³³

Additional findings:

Infants below age 1 year have a relatively high incidence of cardiac involvement. ³⁴

- Cardiac involvement is present in 13.6% of cases (Japan, 2003 to 2004) ³⁵
- Coronary arteritis is common, and coronary artery aneurysms may rupture ^{36 37} or persist into adulthood. ³⁸⁻⁴¹
- Meningoencephalitis, often with seizures, has been reported as a presenting feature of Kawasaki disease. ^{42 43}
- Additional complications include oculomotor ⁴⁴ or facial palsy ⁴⁵, parotitis ⁴⁶, large pleural effusions ⁴⁷, retropharyngeal mass ⁴⁸, cholestatic jaundice ⁴⁹⁻⁵¹, colitis ⁵², nephrotic syndrome ⁵³, sterile pyuria ⁵⁴, sensorineural hearing loss ^{55 56} and peripheral vascular gangrene ⁵⁷ and necrotic lesions on the face. ⁵⁸
- 7% of affected children develop Kawasaki disease shock syndrome, with decreased systolic blood pressure or evidence of hypoperfusion. The shock syndrome is characterized by an increased rate of echocardiographic abnormalities and is less likely to respond to IVIG therapy ^{59 60}
- Neutrophilia, anemia, thrombocytosis, hemophagocytic lymphohistiocytosis ^{61 62}, hepatic dysfunction ⁶³ and sterile pyuria ⁶⁴ are common. Syndrome of inappropriate ADH secretion has been reported. ⁶⁵

Diseases which may mimic Kawasaki disease include Chikungunya ⁶⁶, meningococcal septicemia ⁶⁷, Takayasu's arteritis ⁶⁸, systemic onset juvenile idiopathic arthritis ⁶⁹ and Q fever. ⁷⁰

This disease is endemic or potentially endemic to all countries.

References

1. J Paediatr Child Health 2005 Mar ;41(3):87-93.
2. Pediatr Int 2005 Apr ;47(2):232-4.
3. J Pediatr 2006 Jul ;149(1):131-3.
4. J Comput Assist Tomogr 2012 Jan-Feb;36(1):138-42.
5. Ann Thorac Surg 2008 Mar ;85(3):1081-3.
6. Ugeskr Laeger 2009 Feb 2;171(6):430-3.
7. J Clin Rheumatol 2012 Feb 12;
8. Rheumatol Int 2009 Dec ;30(2):245-8.
9. J Pediatr 2006 Jun ;148(6):800-5.
10. Arch Mal Coeur Vaiss 2007 May ;100(5):439-47.
11. Ann Trop Paediatr 2007 Dec ;27(4):303-6.
12. Semin Arthritis Rheum 2005 Jun ;34(6):785-92.
13. J Paediatr Child Health 2006 Jul-Aug;42(7-8):423-7.
14. Pediatr Infect Dis J 2009 Oct ;28(10):927-8.
15. Ann Trop Paediatr 2007 Dec ;27(4):303-6.
16. Clin Pediatr (Phila) 2010 Jun ;49(6):598-600.
17. Pediatr Rheumatol Online J 2010 ;8:8.
18. Pediatr Infect Dis J 2010 Jun ;29(6):571-3.
19. Dig Liver Dis 2008 Jul ;40(7):582-4.
20. J Clin Rheumatol 2012 Feb 12;
21. Pediatrics 2011 Feb ;127(2):e489-93.
22. Eur J Pediatr 2012 Apr ;171(4):657-62.
23. Pediatr Infect Dis J 2012 Apr ;31(4):417-8.
24. Pediatr Cardiol 2012 Feb 10;
25. Pediatr Cardiol 2012 Feb 15;
26. Arch Pediatr 2012 Nov ;19(11):1264-8.
27. Eur J Pediatr 2012 Apr ;171(4):651-6.
28. Ital J Pediatr 2013 Feb 13;39(1):11.
29. J Pediatr 2010 May ;156(5):786-91.
30. Clin Exp Rheumatol 2012 Sep-Oct;30(5):799-804.
31. Eur J Pediatr 2012 Nov 16;
32. Pediatr Infect Dis J 2010 May ;29(5):430-3.
33. Rheumatol Int 2012 Sep 15;
34. J Microbiol Immunol Infect 2006 Oct ;39(5):387-91.
35. Pediatr Int 2008 Jun ;50(3):287-90.
36. Pediatr Cardiol 2008 Nov ;29(6):1115-9.
37. Interact Cardiovasc Thorac Surg 2010 Feb ;10(2):317-8.
38. Obstet Gynecol 2007 Feb ;109(2 Pt2):517-9.
39. J Cardiol 2008 Feb ;51(1):65-9.
40. World J Pediatr 2010 Feb ;6(1):38-42.
41. Cardiol Young 2011 Feb ;21(1):74-82.
42. J Child Neurol 2006 Dec ;21(12):1080-1.
43. No To Hattatsu 2008 Jul ;40(4):289-94.
44. Rheumatol Int 2011 Jan ;31(1):97-9.
45. Acta Paediatr Taiwan 2008 Jan-Feb;49(1):24-7.
46. Korean Circ J 2009 Nov ;39(11):502-4.
47. J Paediatr Child Health 2007 Mar ;43(3):191-2.
48. Yonsei Med J 2010 Sep ;51(5):784-6.
49. Pediatr Infect Dis J 2012 Mar 14;
50. J Pediatr Gastroenterol Nutr 2012 Mar 20;
51. Indian J Pediatr 2012 Mar 24;
52. BMJ Case Rep 2013 ;2013
53. Pediatr Nephrol 2012 Apr 24;
54. Korean J Pediatr 2013 Jan ;56(1):13-8.
55. J Clin Rheumatol 2010 Oct ;16(7):322-5.
56. Eur J Pediatr 2012 May ;171(5):851-4.
57. Heart Surg Forum 2007 ;10(1):E70-2.
58. Arch Pediatr 2010 Dec ;17(12):1667-9.
59. Pediatrics 2009 May ;123(5):e783-9.
60. Eur J Pediatr 2011 Jul ;170(7):941-3.
61. Pediatr Hematol Oncol 2011 Apr ;28(3):230-6.
62. Arch Pediatr 2012 May 29;
63. Pediatr Infect Dis J 2011 Feb ;30(2):141-4.
64. Pediatr Nephrol 2007 Jul ;22(7):987-91.
65. Pediatr Int 2011 Jun ;53(3):354-7.
66. Pediatr Infect Dis J 2010 Mar ;29(3):275-7.
67. Pediatr Emerg Care 2009 Mar ;25(3):190-2.
68. Rheumatol Int 2012 Nov ;32(11):3655-9.
69. Rheumatol Int 2010 Dec 5;
70. Kansenshogaku Zasshi 2009 May ;83(3):245-50.

Kikuchi's disease and Kimura disease

Agent	UNKNOWN
Reservoir	Unknown
Vector	None
Vehicle	Unknown
Incubation Period	Unknown
Diagnostic Tests	Biopsy.
Typical Adult Therapy	Supportive Hydroxychloroquine and corticosteroids have been successful for Kikuchi's disease in some cases.
Typical Pediatric Therapy	As for adult
Clinical Hints	Most patients of Asian origin. Kikuchi disease: prolonged (1 to 12 months) cervical lymphadenopathy (rubbery, non-matted - may be tender), fever (40%), weight loss, 'sweats', leukopenia. Salivary gland involvement, glomerulitis, painless subcutaneous masses and eosinophilia suggest Kimura disease.
Synonyms	Angiolymphoid hyperplasia, Angiolymphoid hyperplasia-eosinophilia, Eosinophilic follicular lymphadenitis, Histiocytic necrotizing lymphadenitis, Kikuchi and Fujimoto's disease, Kikuchi's disease, Kimura disease. ICD9: 289.3 ICD10: I89.8

Clinical

Kikuchi's disease:

Kikuchi's disease (histiocytic necrotizing lymphadenitis) is characterized by histiocytic necrotizing lymphadenitis, usually of the cervical region ^{1 2} ; however, other anatomic regions may be involved. ³

- Generalized lymphadenopathy is occasionally encountered ⁴
- The disease is primarily seen in young Japanese women or women of Oriental descent in the third decade of life. ⁵
- Pediatric ⁶ , male and elderly patients are occasionally encountered. ⁷
- Leukopenia is present in 50% of cases, and atypical lymphocytes may be seen in the peripheral blood smear.
- Additional features may include aseptic meningitis ^{8 9} , maculopapular or urticarial rash ¹⁰ , arthralgia, myalgia, hepatosplenomegaly, hepatic dysfunction, neuropathy, venous thrombosis ¹¹ , orbital pseudotumor ¹² , pericarditis, pulmonary infiltrates with pleural effusion ¹³ and pulmonary hemorrhage.
- Biopsy material reveals paracortical hyperplasia without granulocytic infiltration and a typical 'starry sky' pattern. ^{14 15}
- Clinical features may mimic those of lupus erythematosus ¹⁶ , tuberculous meningitis ¹⁷ , or lymphoma. ¹⁸⁻²¹
- The prognosis is good, and patients recover after a mean of 3 months.
- Relapse occurs in 20% of cases ²² and recurrence in 3% to 4%. ²³
- Hydroxychloroquine and corticosteroids have been advocated by some authorities.

Kimura disease:

Kimura disease (angiolymphoid hyperplasia with eosinophiles (eosinophilic follicular lymphadenitis) is also most common among Oriental males. ²⁴

- Most present as painless subcutaneous masses and lymphadenopathy of the cervical region.
- Cases of isolated Kimura disease of the earlobe ²⁵ and eyelid have been reported ^{26 27}
- In contrast to Kikuchi's disease, salivary gland involvement ^{28 29} , glomerulitis, nephrotic syndrome ³⁰ , elevated IgE and eosinophilia are often encountered.
- Hypercoagulability has been reported in some cases ³¹

Angiolymphoid hyperplasia with eosinophilia is clinically similar to Kimura disease, but is histologically distinct from the latter. ³²⁻³⁸

- The condition is characterized by characterized by reddish-brown nodules and plaques in the dermis and the subcutaneous tissues, typically occurring on the neck and head. ³⁹

This disease is endemic or potentially endemic to all countries.

References

1. Pediatrics 2004 Dec ;114(6):e752-6.
2. Am J Hematol 2003 Sep ;74(1):60-3.
3. Intern Med 2011 ;50(6):649-52.
4. Joint Bone Spine 2006 May ;73(3):311-3.
5. Clin Rheumatol 2007 Jan ;26(1):50-4.
6. Ear Nose Throat J 2008 Jun ;87(6):350-3.
7. Hawaii Med J 2006 Nov ;65(11):315-7.
8. Clin Infect Dis 2005 Oct 15;41(8):e80-2.
9. Neurol Sci 2012 Nov 3;
10. Eur J Pediatr 2004 Apr ;163(4-5):210-3.
11. Vasa 2012 Sep ;41(5):371-4.
12. Zhonghua Yan Ke Za Zhi 2011 May ;47(5):427-30.
13. BMC Pulm Med 2010 ;10:54.
14. Zhonghua Nei Ke Za Zhi 2006 Feb ;45(2):127-9.
15. Hum Pathol 2010 Sep ;41(9):1245-54.
16. Lupus 2006 ;15(6):384-7.
17. Neurol Sci 2012 Nov 3;
18. Heart Lung 2009 Sep-Oct;38(5):450-6.
19. Breast Cancer 2009 Nov 27;
20. Eur J Radiol 2012 Aug ;81(8):1817-20.
21. Exp Oncol 2011 Dec ;33(4):242-4.
22. Int J Infect Dis 2009 May ;13(3):322-6.
23. Case Report Otolaryngol 2013 ;2013:364795.
24. Eur J Radiol 2011 Nov ;80(2):489-97.
25. Clin Exp Dermatol 2010 Jun ;35(4):e97-9.
26. Ophthal Plast Reconstr Surg 2006 Nov-Dec;22(6):495-8.
27. Nihon Ganka Gakkai Zasshi 2011 Aug ;115(8):699-705.
28. J Craniofac Surg 2011 Jan ;22(1):337-8.
29. Eur Ann Otorhinolaryngol Head Neck Dis 2012 Sep 25;
30. Eur J Dermatol 2009 Nov-Dec;19(6):626-8.
31. J Thromb Thrombolysis 2010 Apr ;29(3):354-7.
32. Rev Med Interne 2007 May ;28(5):346-8.
33. Zhonghua Bing Li Xue Za Zhi 2005 Jun ;34(6):353-7.
34. Br J Oral Maxillofac Surg 2005 Jun ;43(3):249-52.
35. Cesk Slov Oftalmol 2003 Sep ;59(5):319-24.
36. J Laryngol Otol 2003 Jul ;117(7):570-3.
37. J Dermatol 2010 Apr ;37(4):355-9.
38. Clin Colorectal Cancer 2010 Jul ;9(3):179-82.
39. Int J Surg Case Rep 2011 ;2(8):258-60.

Kingella infection

Agent	BACTERIUM. <i>Kingella kingae</i> , et al A facultative gram-negative coccobacillus
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Unknown
Diagnostic Tests	Culture of blood, joint fluid, CSF, etc. Alert laboratory if these organisms are suspected.
Typical Adult Therapy	Penicillin G or Penicillin V usually effective - dosage per severity/site
Typical Pediatric Therapy	As for adult
Clinical Hints	A relatively rare cause of septic arthritis, endocarditis, meningitis and other infections; most infections have been in young children.
Synonyms	

Clinical

Kingella kingae, *K. (Suttonella) indologenes*, *K. denitrificans* and *K. oralis* are found in the normal respiratory tract, and occasionally associated with bacteremia, bone and joint infection (notably in young children) ^{1 2} and endocarditis (the 'K' in the HACEK group). ³

- *Kingella potus* has been isolated from a kinkajou wound in a zookeeper. ⁴

This disease is endemic or potentially endemic to all countries.

References

1. Lancet Infect Dis 2004 Jun ;4(6):358-67.
2. Expert Rev Anti Infect Ther 2004 Oct ;2(5):787-94.
3. Clin Microbiol Rev 2001 Jan ;14(1):177-207.
4. J Clin Microbiol 2005 Jul ;43(7):3526-9.

Laryngotracheobronchitis

Agent	VIRUS OR BACTERIUM. Parainfluenza virus, Influenza virus, Mycoplasma, et al
Reservoir	Human
Vector	None
Vehicle	Droplet
Incubation Period	3d - 8d
Diagnostic Tests	Viral culture (respiratory secretions). Serology. Nucleic acid amplification.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Usually encountered in the setting of bronchiolitis, laryngitis or croup following a minor upper respiratory infection in young children.
Synonyms	Bronchitis, Croup, Laringitis, Laryngite, Laryngitis, Laryngotracheitis. ICD9: 464,466 ICD10: J04,J05,J20,J21

Clinical

Laryngotracheobronchitis is a self-defined syndrome consisting of hacking cough, often with an 'itching' or 'foreign body' sensation in the airways, and hoarseness. ¹

- Viral croup and epiglottitis are two major inflammatory causes of airway obstruction in children.
- Spasmodic croup and membranous laryngotracheobronchitis may be associated with obstruction. ²

Bacterial tracheitis is an uncommon (>200 cases reported worldwide) severe condition usually affecting children that manifests as cough, stridor, mucopurulent tracheal secretions and lack of response to therapeutic modalities used for treating viral croup. ³

- Fever may be low-grade or even absent.
- 75% of patients require intubation and mechanical ventilation.
- The case/fatality rate is approximately 2%.
- Causative pathogens include *Staphylococcus aureus* (50% of cases) and *S. pneumoniae*, *H. influenzae*, *M. catarrhalis* and *S. pyogenes*. Gram-negative bacilli are also reported in some cases.
- Occasionally, co-infection with viral croup agents is found.

This disease is endemic or potentially endemic to all countries.

References

1. *Pediatr Clin North Am* 1994 Apr ;41(2):265-76.
2. *Pediatr Emerg Care* 1991 Dec ;7(6):337-42.
3. *Scand J Infect Dis* 2009 ;41(8):548-57.

Lassa fever

Agent	VIRUS - RNA. Arenaviridae, Arenavirus: Lassa virus
Reservoir	Multimammate rat (<i>Mastomys natalensis</i>)
Vector	None
Vehicle	Rodent secretions Dust Food Patient secretions
Incubation Period	8d - 14d (range 3d - 21d)
Diagnostic Tests	Viral culture (blood, urine, throat, liver, spleen). Serology. Nucleic acid amplification. Biosafety level 4.
Typical Adult Therapy	Strict isolation. Ribavirin 2g IV Then 1g q6h IV X 4d Then 0.5g IV q8h X 6d
Typical Pediatric Therapy	Strict isolation. Ribavirin 30 mg/kg IV Then 15 mg/kg IV q6h X 4d Then 7.5 mg/kg IV q8h X 6d
Clinical Hints	Gastrointestinal symptoms, cough, pharyngitis, conjunctivitis and retrosternal pain; leukopenia, proteinuria and hepatic dysfunction may be present; case-fatality rates of 15% to 25% are reported.
Synonyms	Lassa-Fieber, Lujo, Luna virus. ICD9: 078.89 ICD10: A96.2

Clinical

WHO Case definition for surveillance:

Clinical description

An illness of gradual onset with one or more of the following:

- malaise, fever, headache, sore throat, cough, nausea, vomiting, diarrhea, myalgia, chest pain, hearing loss, and
- A history of contact with excreta of rodents or with a probable or confirmed case of Lassa fever.

Laboratory criteria for diagnosis

- Isolation of virus (only in laboratory of biosafety level 4) from blood, urine or throat washings or
- Positive IgM serology or seroconversion (IgG antibody) in paired serum specimens or
- Demonstration of Lassa virus antigen in autopsy tissues by immunohistochemistry or in serum by ELISA
- Positive PCR from serum or autopsy tissues

Case classification

- Suspected: A case compatible with the clinical description.
- Probable: A suspected case that is epidemiologically linked to a confirmed case.
- Confirmed: A suspected case that is laboratory-confirmed.
- Contact: A person having close personal contact with the patient (living with, caring for) or a person testing the laboratory specimens of a patient in the 3 weeks after the onset of the illness.

Acute illness:

The onset of symptoms is gradual, with fever, malaise, headache, sore throat, cough, nausea, vomiting, diarrhea, myalgia and chest and abdominal pain.

- The fever may be either constant or intermittent with spikes. ¹
- Inflammation of the throat and eyes is commonly observed.
- In severe cases, hypotension or shock, pleural effusion, hemorrhage, seizures, encephalopathy and swelling of the face and neck are commonly encountered ²
- Case-fatality rates of 15% to 28% are reported. ³
- The disease is more severe in pregnancy, and fetal loss occurs in greater than 80% of cases. ⁴

Additional signs and symptoms:

Hair loss and loss of coordination may occur in convalescence.

- Sensorineural deafness occurs in 29%, making this the most common cause of deafness in West Africa. ⁵⁻⁷

The clinical syndrome of Lassa fever is difficult to distinguish from severe malaria, septicemia, yellow fever and other viral hemorrhagic fevers (e.g., Ebola).

- Inflammation of the throat with white tonsillar patches is an important distinguishing feature.

Definitive diagnosis requires testing that is available only in highly-specialized laboratories.

- Laboratory specimens may be hazardous and must be handled with extreme care at the highest level of biosafety containment.

This disease is endemic or potentially endemic to 13 countries.

References

1. BMJ 2003 Nov 29;327(7426):1271-5.
2. Rev Infect Dis 1989 May-Jun;11 Suppl 4:S783-9.
3. Trans R Soc Trop Med Hyg 2010 Jan ;104(1):51-4.
4. Virol J 2011 ;8:404.
5. JAMA 1990 Oct 24-31;264(16):2093-6.
6. J Med Case Rep 2009 ;3:36.
7. Eur Arch Otorhinolaryngol 2011 Feb ;268(2):197-201.

Legionellosis

Agent	BACTERIUM. <i>Legionella pneumophila</i> , et al An aerobic gram-negative bacillus
Reservoir	Water
Vector	None
Vehicle	Water Aerosols
Incubation Period	5- 6d (range 2-12d); Pontiac fever = 1-2d
Diagnostic Tests	Serology. Culture. Urine antigen (certain types). Nucleic acid amplification. Alert lab if organism suspected.
Typical Adult Therapy	Fluoroquinolone (<i>Levofloxacin</i> , <i>Trovafoxacin</i> , <i>Pefloxacin</i> , <i>Sparfloxacin</i> or <i>Moxifloxacin</i>). OR <i>Azithromycin</i> . OR <i>Erythromycin</i> + <i>Rifampin</i>
Typical Pediatric Therapy	<i>Azithromycin</i> . OR <i>Erythromycin</i> + <i>Rifampin</i>
Clinical Hints	Respiratory illness with extrapulmonary manifestations (diarrhea, confusion, renal or hepatic dysfunction, relative bradycardia, etc.); most cases reported during summer in temperate areas; case-fatality rates of 5% to 25% are reported.
Synonyms	Doença dos legionarios, Legionarsjuka, Legionarssjuka, Legionella, Legionellose, Legionellosi, Legionnaire's disease, Pontiac fever. ICD9: 482.84 ICD10: A48.1,A48.2

Clinical

WHO Case definition for surveillance:

Clinical description

An illness characterized by an acute lower respiratory infection with focal signs of pneumonia on clinical examination and/or radiological evidence of pneumonia.

Laboratory criteria for diagnosis

Presumptive: one or more of the following:

- Detection of specific *Legionella* antigen in respiratory secretions or urine
- Direct fluorescent antibody (DFA) staining of the organism in respiratory secretions or lung tissue, using evaluated monoclonal reagents
- A fourfold or greater rise in specific serum antibody titer to *Legionella* species other than *Legionella pneumophila* serogroup 1, using a locally validated serological test

Confirmative: one or more of the following:

- Isolation of *Legionella* from respiratory secretions, lung tissue, pleural fluid, or blood
- A fourfold or greater rise in specific serum antibody titer to *L. pneumophila* serogroup 1 by indirect immunofluorescence antibody test or microagglutination
- Most European countries and others such as the United States now include the detection of *L. pneumophila* serogroup 1 antigen in urine as a confirmatory test.

Case classification

- Suspected: Not applicable.
- Probable: A case compatible with the clinical description, with presumptive laboratory results.
- Confirmed: A case compatible with the clinical description, with confirmative laboratory results.

Pneumonia associated with extrapulmonary findings should suggest the possibility of Legionnaire's disease.

- Q-fever may be mistaken for Legionnaires' disease ¹
- The most common clinical manifestation is pneumonia, ranging from mild • to severe, with respiratory failure and death.
- Risk factors for overt disease include advanced age, smoking, chronic obstructive pulmonary disease, immunosuppression, and recent surgery.
- Person-to-person transmission has not been demonstrated.

Legionnaire's disease vs. Pontiac fever:

There are 2 currently recognized distinct clinicoepidemiological manifestations of legionellosis:

- Both forms are characterized initially by anorexia, vomiting, myalgia and headache, followed within a day by rising fevers and chills.
- Legionnaires. disease. (pneumonic form) and
- Pontiac fever (non-pneumonic Legionnaires disease)

Legionnaires disease ^{2 3}

- In the pneumonic form, non-productive cough, abdominal pain / diarrhea, confusion / delirium are common.
- It is not possible, clinically, to distinguish *Legionella* pneumonia from other pneumonias ⁴ ; suspicion should be raised in any pneumonia connected with epidemiological information (e.g., recent traveling, hospitalization, gatherings, immunosuppression).
- In addition, age (>50), sex (M), smoking, alcohol consumption have been shown to be risk factors.

Pontiac fever:

Pontiac fever is a self-limited, influenza-like illness lasting 2 to 5 days, often in healthy persons following exposure to contaminated whirlpools or spas. ⁵⁻⁷

- Pontiac fever is not associated with pneumonia. It is thought to represent a reaction to inhaled antigen, rather than to bacteria.
- Proposed case definition for Pontiac fever include occurrence of at least one symptom among headache, myalgia, fever and rigors, beginning 2 to 8 days following exposure. ⁸

Complications:

Reported complications of legionellosis have included empyema, pleural effusion, lung abscess, renal failure (in 10% to 50% of cases), endocarditis ^{9 10} , peritonitis, cerebellar ataxia ¹¹ , cutaneous and visceral abscesses ¹² , arteriovenous fistula infection, pericarditis and myocarditis.

- Case-fatality rates may approach 40%, particularly among patients with underlying disease or immunosuppression.
- Additional risk factors for fatal infection include heart disease, malignancy, alcoholism and renal disease. ¹³

This disease is endemic or potentially endemic to all countries.

References

1. Heart Lung 2009 Jul-Aug;38(4):354-62.
2. Ann Intern Med 1979 Apr ;90(4):509-17.
3. Ann Intern Med 1979 Apr ;90(4):522-6.
4. J Thorac Imaging 1991 Jul ;6(3):6-13.
5. Am J Epidemiol 1978 Feb ;107(2):149-60.
6. Lancet 1980 Nov 1;2(8201):969.
7. Medicine (Baltimore) 1983 Mar ;62(2):120-8.
8. BMC Public Health 2006 ;6:112.
9. Heart 2003 May ;89(5):e16.
10. Emerg Infect Dis 2012 Jan ;18(1):95-7.
11. Acta Med Port 2011 Dec ;24 Suppl 3:667-70.
12. Emerg Infect Dis 2011 Jan ;17(1):145-6.
13. Med Mal Infect 2007 Jun ;37(6):325-30.

Leishmaniasis - cutaneous

Agent	PARASITE - Protozoa. Neozoa, Euglenozoa, Kinetoplastea. Flagellate: <i>Leishmania tropica</i> , et al
Reservoir	Human Hyrax Rodent Marsupial Dog Sloth Anteater Armadillo
Vector	Fly (sandfly = <i>Phlebotomus</i> for old world; <i>Lutzomyia</i> or <i>Psychodopygus</i> for new world)
Vehicle	None
Incubation Period	2w - 8w (range 1w - months)
Diagnostic Tests	Identification of organism on smear or specialized culture. Nucleic acid amplification
Typical Adult Therapy	Pentavalent antimonials 20 mg/kg/d IV or IM X 21d & / or topical paromomycin . Alternatives: <i>L. major</i> - Fluconazole or Azithromycin , PO <i>L. mexicana</i> or <i>L. panamensis</i> - Ketoconazole , PO <i>L. braziliensis</i> - Azithromycin , PO
Typical Pediatric Therapy	As for adult
Clinical Hints	Chronic ulcerating skin nodule; painless (<i>Leishmania tropica</i>) or painful (<i>L. major</i>); diffuse infection or regional lymphadenopathy occasionally encountered.
Synonyms	Aleppo button, Antep boil, Baghdad boil, Bay sore, Bejuco, Biskra button, Bolho, Bosjaws, Bush yaws, Busi-yasi, Chiclero ulcer, Cutaneous leishmaniasis, Delhi ulcer, Domal, El-Mohtafura, Forest yaws, Gafsa boil, Granuloma endemicum, Hashara, Jericho boil, Kaal Daana, Kandahar sore, <i>Leishmania major</i> , <i>Leishmania tropica</i> , Leishmaniasis, Leishmaniose: Kutane, Leishmaniosi cutanea, Lepra de montana, Liana, Okhet, One-year boil, Oriental sore, Pendjeh sore, Pian bois, Saldana, Ulcera de Bejuco, Urfa boil, Uta, Yatevi, Year boil. ICD9: 085.1,085.2,085.3,085.4 ICD10: B55.1

Clinical

WHO Case definition for surveillance:

Clinical description

- Appearance of one or more lesions, typically on uncovered parts of the body.
- The face, neck, arms and legs are the most common sites.
- At the site of inoculation a nodule appears, and may enlarge to become an indolent ulcer.
- The sore remains in this stage for a variable time before healing, and typically leaves a depressed scar.
- Other atypical forms may occur.
- In some individuals, certain strains can disseminate and cause mucosal lesions. These sequelae involve nasopharyngeal tissues and can be very disfiguring.

Laboratory criteria for diagnosis

- positive parasitology (stained smear or culture from the lesion)
- mucocutaneous leishmaniasis only: positive serology (IFA, ELISA)

Case classification

WHO operational definition:

- A case of cutaneous leishmaniasis is a person showing clinical signs (skin or mucosal lesions) with parasitological confirmation of the diagnosis (positive smear or culture) and/or, for mucocutaneous leishmaniasis only, serological diagnosis.

Typically, a nodule develops at the site of a sandfly bite following a few days to several months. ^{1 2}

- The lesion may be erythematous, or covered by a thin yellow crust.
- The nodule reaches a diameter of 1 to 5 cm over a period of weeks or months, and is not painful. ³
- The crust may thicken, and even replace the nodule; or may fall away to reveal an ulcer with a raised edge.
- Panniculitis ⁴ and satellite papules are common.
- The lesion may heal over a period of months or even years, leaving a depressed scar.
- Secondary infection is not prominent, and the major residua are scarring and disability.
- Rare instances of late scar carcinoma have been reported. ⁵

Lesions caused by *Leishmania major* evolve and heal most rapidly, and are often inflamed or exudative ("wet sore" or "rural sore").

- Lesions caused by *L. tropica* are less inflamed ("dry sore" or "urban sore").
- Lesions due to *L. infantum* appear only after several months, and are small, nodular, and persist for years.

Lesions of *L. aethiopia* are typically single, and often involve the face.

- Satellite papules evolve to produce a slowly growing, shiny tumor or plaque that may not crust nor ulcerate.
- If the site borders an area of mucosa, mucocutaneous leishmaniasis may develop, with swelling of the lips and enlargement of the nose over many years.

Leishmania brasiliensis produces deep, usually single, ulcers with a granulomatous base.

- 15 per cent of patients will relapse after spontaneous recovery or therapeutic improvement.

The lesions of **L. guyanensis** are multiple, fleshy and protuberant, and involve the limbs.

- Unlike other *Leishmania* species, **L. braziliensis** and **L. panamensis** are commonly associated with metastatic lesions along the path of draining lymphatics.
- Nodular lymphadenitis occurs, and may mimic nocardiosis. ⁶

The lesions of **L. mexicana** ('chiclero ulcer') are commonly located on the side of the face or behind the ears.

- The lesion consists of a single ulcerative lesion, most commonly involving the ear pinna, without a tendency for cutaneous metastasis, lymphatic or mucosal involvement. ⁷
- Destruction of the pinna is common.

Skin lesions with regional adenopathy may also occur in visceral leishmaniasis, and suggest a diagnosis of cutaneous leishmaniasis. ⁸

- Facial lesions of chromomycosis may be misdiagnosed as cutaneous leishmaniasis. ⁹

Other clinical forms:

Three forms of cutaneous leishmaniasis do not heal spontaneously: **Disseminated cutaneous leishmaniasis**, **Leishmaniasis recidivans** and **American mucosal leishmaniasis**.

- Diffuse cutaneous leishmaniasis is often seen with *L. amazoensis* infections, and also occurs in about 0.01% of *L. aethiopica* infections.
- The nodule spreads locally without ulceration, while secondary hematogenous lesions appear on other body sites.
- These are often symmetrical, and involve the face and extensor surfaces of the limbs.
- The external genitalia may also be affected, but the eye, mucosae and peripheral nerves are not infected (in contrast to lepromatous leprosy).
- The infection evolves gradually over many years.

Cases of erysipeloid, recidiva cutis (LRC), and disseminated leishmaniasis (DL) have been ascribed to *L. panamensis* infection. ¹⁰

Leishmaniasis recidivans (lupoid leishmaniasis) is a rare complication of *L. tropica* infection.

- The lesion presents as a spreading nodule, leading to a plaque formation simulating discoid lupus erythematosus. ^{11 12}
- After initial healing, papules reappear in the edge of the scar and the lesion spreads slowly over many years.
- The condition most commonly involves the face, and may be quite disfiguring.

Sporotrichoid cutaneous leishmaniasis may mimic cutaneous sporotrichosis. ¹³

- Lesions of cutaneous leishmaniasis may mimic those of erysipeloid ¹⁴ or carcinoma. ¹⁵
- Diffuse cutaneous leishmaniasis may mimic lepromatous leprosy ¹⁶ The lesions of both cutaneous and mucocutaneous leishmaniasis could be mistaken for those of borderline tuberculoid leprosy. ¹⁷

Atypical, non-ulcerating nodular granulomatous lesions caused by *L. mexicana* and *L. chagasi* have been described in Central America.

- Most cases have involved exposed areas on the body, and most patients have been children.

In rare cases, leishmaniasis of the nose may present as rhinophyma ¹⁸ or mimic erysipelas. ¹⁹

This disease is endemic or potentially endemic to 85 countries.

Leishmaniasis - cutaneous in Uganda

An endemic focus exists on the Kenyan border.

The local vectors are *Phlebotomus (Larroussius) pedifer*, *Ph. (Synphlebotomus) martini*, *Ph. longipes* and *Ph. orientalis*.

References

1. *Comp Immunol Microbiol Infect Dis* 2004 Sep ;27(5):305-18.
2. *Curr Opin Infect Dis* 2003 Apr ;16(2):97-101.
3. *J Postgrad Med* 2003 Jan-Mar;49(1):50-4.
4. *Case Rep Dermatol Med* 2012 ;2012:612434.
5. *J Craniofac Surg* 2007 May ;18(3):708-10.
6. *Curr Infect Dis Rep* 2008 Sep ;10(5):404-10.
7. *Travel Med Infect Dis* 2012 Nov 9;
8. *Clin Exp Dermatol* 2009 Jul ;34(5):e196-8.
9. *Dermatol Online J* 2012 ;18(10):3.
10. *Am J Trop Med Hyg* 2005 Aug ;73(2):281-4.

11. Trop Doct 2012 Feb 15;
12. Dermatol Ther (Heidelb) 2011 Dec ;1(2):36-41.
13. Ann Dermatol Venereol 2008 Jan ;135(1):63-7.
14. Bull Soc Pathol Exot 2008 Dec ;101(5):395-7.
15. Am J Otolaryngol 2009 Jul-Aug;30(4):285-7.
16. Dermatol Online J 2009 ;15(4):9.
17. Acta Derm Venereol 2012 Mar 20;
18. Dermatol Online J 2009 ;15(3):10.
19. Eur Ann Otorhinolaryngol Head Neck Dis 2011 Apr ;128(2):95-7.

Leishmaniasis - visceral

Agent	PARASITE - Protozoa. Neozoa, Euglenozoa, Kinetoplastea. Flagellate: <i>Leishmania donovani</i> , <i>L. infantum</i> , <i>L. cruzi</i> ; rarely, <i>L. tropica</i>
Reservoir	Human Rodent Dog Fox Hares
Vector	Fly (sandfly = <i>Phlebotomus</i> for old world; <i>Lutzomyia</i> for new world)
Vehicle	Blood
Incubation Period	2m - 6m (10d - 12m)
Diagnostic Tests	Smear / culture of bone marrow, splenic aspirate, lymph nodes. Serology. Nucleic acid amplification.
Typical Adult Therapy	Pentavalent antimonials (Stibogluconate) 20 mg/kg/d X 28d. OR Amphotericin B 1 mg/kg/QOD X 8w (or lipid complex 3 mg/kg/d X 5d) OR Paromomycin 11 mg/kg IM QD X 21 days OR Miltefosine 50 to 150 mg PO daily X 4 to 6 weeks.
Typical Pediatric Therapy	Pentavalent antimonials (Stibogluconate) 20 mg/kg/d X 28d. OR Amphotericin B 1 mg/kg/QOD X 8w (or lipid complex 3 mg/kg/d X 5d) OR Paromomycin 11 mg/kg IM QD X 21 days OR Miltefosine 2.5 mg/kg daily (maximum 150 mg) X 28d
Clinical Hints	Chronic fever, weight loss, diaphoresis, hepatosplenomegaly, lymphadenopathy and pancytopenia; grey pigmentation (Kala Azar = 'black disease') may appear late in severe illness; case-fatality rate = 5% (treated) to 90% (untreated).
Synonyms	Burdwan fever, Cachectic fever, Dum Dum fever, Kala azar, <i>Leishmania donovani</i> , <i>Leishmania infantum</i> , <i>Leishmania siamensis</i> , Leishmaniose: Viszerale, Leishmaniosi viscerale, Ponos, Visceral leishmaniasis. ICD9: 085.0 ICD10: B55.0

Clinical

WHO Case definition for surveillance:

Clinical description

- An illness with prolonged irregular fever, splenomegaly and weight loss as its main symptoms.

Laboratory criteria for diagnosis

- positive parasitology (stained smears from bone marrow, spleen, liver, lymph node, blood or culture of the organism from a biopsy or aspirated material)
- positive serology (IFA, ELISA)

Case classification

WHO operational definition:

- A case of visceral leishmaniasis is a person showing clinical signs (mainly prolonged irregular fever, splenomegaly and weight loss) with serological (at geographical area level) and/or parasitological confirmation (when feasible at central level) of the diagnosis.
- In endemic malarious areas, visceral leishmaniasis should be suspected when fever lasts for more than two weeks and no response has been achieved with anti-malaria drugs (assuming drug resistant malaria has also been considered).

Following an incubation period of two to eight months, the patient develops chronic fever, abdominal pain (from an enlarged spleen) and swelling, weight loss, cough and occasionally, diarrhea.

- The classical fever rises twice daily, without rigors; however, single 'spikes,' irregular or undulant fevers are common.
- Caucasians may experience an abrupt onset of high fever, with rapid progression of illness, toxemia, weakness, dyspnea, and anemia.
- Visceral leishmaniasis in HIV-positive patients is characterized by short incubation period, high incidence of multi-organ disease, and tendency to relapse. ¹

Physical signs may be limited to splenomegaly; but chronically-ill patients are typically pale and cachectic.

- Hyperpigmentation of face, extremities and abdomen ('Kala azar') may be present in advanced cases.
- The spleen is non-tender, and may be massively enlarged, reaching the left or even right iliac fossa.
- Moderate hepatomegaly is present in one-third of cases.
- Rare instances of granulomatous hepatitis are reported. ²
- Generalized lymphadenopathy is found in 50% of African patients, and a smaller percentage of Indian and European cases. In some cases, localized lymphadenopathy may be the only sign of infection. ³
- Jaundice, mucosal and retinal hemorrhage, laryngeal lesions ⁴, uveitis, chronic diarrhea with malabsorption ⁵, interstitial

nephritis ⁶ , pericardial effusion ⁷ and panniculitis ⁸ are occasionally encountered.

- Skin lesions with regional adenopathy may suggest a diagnosis of cutaneous leishmaniasis. ⁹
- A chronic rash (Post kala-azar dermal leishmaniasis = PKDL) resembling leprosy, and involving primarily the extremities and face often appears months to years following infection. ¹⁰
- Other rare manifestations include the hemophagocytic syndrome ¹¹ , laryngeal infection ¹² , leukemoid changes or myelodysplasia or pyothorax ¹³
- The clinical features of visceral leishmaniasis may mimic those of autoimmune hepatitis, primary biliary cirrhosis, and systemic lupus erythematosus. ¹⁴

Laboratory studies reveal pancytopenia, hypoalbuminemia, hyperglobulinemia, and only mild hepatic dysfunction.

- In endemic countries, leishmaniasis is the principal cause of febrile pancytopenia among children without hematologic malignancy. ¹⁵
- Intercurrent infections are common notably pneumococcal disease (otitis, pneumonia or septicemia), tuberculosis and measles.
- Evidence of primary adrenal insufficiency, hypoparathyroidism and hypothyroidism are common. ¹⁶

The case/fatality rate without treatment is 80% to 90%.

This disease is endemic or potentially endemic to 108 countries.

Leishmaniasis - visceral in Uganda

Kala Azar was first reported in Uganda until the 1950's. ¹⁷

Kala azar in Uganda appears to be limited to Pokot Country, Nakapiripirit District (eastern border with Kenya). ¹⁸

2,088 cases were confirmed during 2000 to 2006.

- Incidence increased from 175 cases in 2000, to 690 cases in 2005.

Malaria was identified in 19% of patients with visceral leishmaniasis (2000 to 2006) ¹⁹

The local vector is *Phlebotomus martini*.

References

1. Ann Trop Med Parasitol 2003 Oct ;97 Suppl 1:99-105.
2. Turkiye Parazitoloj Derg 2008 ;32(1):12-5.
3. Vector Borne Zoonotic Dis 2011 Aug ;11(8):1213-5.
4. Indian J Pathol Microbiol 2009 Jan-Mar;52(1):62-4.
5. Indian J Gastroenterol 2006 Nov-Dec;25(6):309-10.
6. Am J Transplant 2010 Jun ;10(6):1486-9.
7. Trans R Soc Trop Med Hyg 2007 May ;101(5):429-32.
8. Acta Med Port 2011 Jul ;24(4):649-52.
9. Clin Exp Dermatol 2009 Jul ;34(5):e196-8.
10. Lancet Infect Dis 2003 Feb ;3(2):87-98.
11. Zhonghua Er Ke Za Zhi 2011 Jul ;49(7):550-3.
12. Head Neck 2012 Sep 13;
13. Indian J Pediatr 2009 Aug ;76(8):843-5.
14. Korean J Parasitol 2012 Jun ;50(2):133-6.
15. Pediatr Emerg Care 2012 May 29;
16. Am J Trop Med Hyg 2011 May ;84(5):668-73.
17. East Afr Med J 1969 Apr ;46(4):204-7.
18. Emerg Infect Dis 2007 Mar ;13(3):507-9.
19. PLoS Negl Trop Dis 2012 Apr ;6(4):e1617.

Leprosy

Agent	BACTERIUM. <i>Mycobacterium leprae</i> An acid-fast bacillus
Reservoir	Human ? Armadillo
Vector	None
Vehicle	Patient secretions
Incubation Period	3y - 5y (range 3m - 40y)
Diagnostic Tests	Visualization of organisms in exudate, scrapings or biopsy. Nucleic acid amplification.
Typical Adult Therapy	Multibacillary: One year therapy Dapsone 100 mg + Clofazimine 50 mg daily; and, Rifampin 600 mg + Clofazimine 300 mg once monthly Paucibacillary: Six month therapy Dapsone 100 mg daily; and Rifampin 600 mg once monthly
Typical Pediatric Therapy	Multibacillary: One year therapy Dapsone 1 to 2 mg/kg + Clofazimine 1 mg/kg daily; and, Rifampin 10 mg/kg + Clofazimine 1 mg/kg once monthly Paucibacillary: Six month therapy Dapsone 1 to 2 mg/kg daily; and Rifampin 10 mg/kg once monthly
Clinical Hints	Anesthetic, circinate hypopigmented skin lesions and thickened peripheral nerves (tuberculoid leprosy); or diffuse, destructive papulonodular infection (lepromatous leprosy); or combined/intermediate forms.
Synonyms	Aussatz, Doence de Hansen, Hansen's disease, Lebbra, Lepra, <i>Mycobacterium leprae</i> , <i>Mycobacterium lepromatosis</i> . ICD9: 030 ICD10: A30

Clinical

WHO Case definition for surveillance:

Clinical description

- The clinical manifestations of the disease vary in a continuous spectrum between the two polar forms, lepromatous and tuberculoid leprosy:
- In lepromatous (multibacillary) leprosy, nodules, papules, macules and diffuse infiltrations are bilateral symmetrical and usually numerous and extensive; involvement of the nasal mucosa may lead to crusting, obstructed breathing and epistaxis; ocular involvement leads to iritis and keratitis
- In tuberculoid (paucibacillary) leprosy, skin lesions are single or few, sharply demarcated, anesthetic or hypoesthetic, and bilateral asymmetrical, involvement of peripheral nerves tends to be severe
- Borderline leprosy has features of both polar forms and is more labile
- Indeterminate leprosy is characterized by hypopigmented maculae with ill-defined borders; if untreated, it may progress to tuberculoid, borderline or lepromatous disease

Laboratory criteria for confirmation

- Alcohol-acid-fast bacilli in skin smears (made by the scrape-incision method).
- In the paucibacillary form the bacilli may be so few that they are not demonstrable.
- In view of the increasing prevalence of HIV and hepatitis B infection in many countries where leprosy remains endemic, the number of skin smear sites and the frequency of smear collection should be limited to the minimum necessary.

Case classification:

WHO operational definition:

A case of leprosy is defined as a person showing one or more of the following features, and who as yet has to complete a full course of treatment:

- hypopigmented or reddish skin lesions with definite loss of sensation
- involvement of the peripheral nerves, as demonstrated by definite thickening with loss of sensation
- skin smear positive for acid-fast bacilli

Classification (microbiological):

Paucibacillary (PB): includes all smear-negative cases

Multibacillary (MB): includes all smear-positive cases.

Classification (clinical):

Paucibacillary single lesion leprosy: 1 skin lesion.

Paucibacillary leprosy: 2 to 5 patches or lesions on the skin.

Multibacillary leprosy: >5 patches or lesions on the skin.

The major forms of leprosy are as follows: ^{1 2}

Tuberculoid • one or a few well-demarcated, hypopigmented, and anesthetic skin lesions, frequently with active, spreading edges and a clearing center; peripheral nerve swelling or thickening also may occur.

Lepromatous • a number of erythematous papules and nodules or an infiltration of the face (including oral mucosa ³ , hands, and feet with lesions in a bilateral and symmetrical distribution that progress to thickening of the skin.

Borderline (dimorphous) • skin lesions characteristic of both the tuberculoid and lepromatous forms.

Indeterminate • early lesions, usually hypopigmented macules, without developed tuberculoid or lepromatous features.

Relapsing disease may manifest as lymphadenopathy mimicking tuberculosis ⁴

- Relapses may follow effective antimicrobial therapy. ⁵⁻⁷

The skin lesions of paracoccidioidomycosis may mimic those of tuberculoid leprosy. ⁸

- Lepromatous leprosy may mimic sarcoidosis. ^{9 10}
- Lupus vulgaris may mimic actinomycosis or mycetoma. ¹¹
- Diffuse cutaneous leishmaniasis may mimic lepromatous leprosy ¹² The lesions of both cutaneous and mucocutaneous leishmaniasis could be mistaken for those of borderline tuberculoid leprosy. ¹³
- Leprosy may be initially misdiagnosed as adult stills disease or an auto-immune disorder. ¹⁴

Leprosy may be associated with endocrine dysfunction including hypogonadism, sterility and osteoporosis ^{15 16}

Six percent of leprosy patients exhibit rheumatological manifestations, most commonly resembling rheumatoid arthritis ^{17 18}

- Rare instances of spondylodiscitis have been reported. ¹⁹

Lucio's phenomenon is a rare and aggressive necrotizing variant of erythema nodosum leprosum that classically occur in patients with undiagnosed, diffuse non-nodular lepromatous leprosy. ²⁰

Erythema multiforme is occasionally encountered among patients with leprosy. ²¹

Chronic skin lesions may undergo malignant transformation. ²²


Cranial nerve involvement, most often trigeminal and facial, is not uncommon. ²³

- Segmental necrotizing granulomatous neuritis is reported in some cases. ²⁴

This disease is endemic or potentially endemic to all countries.

Leprosy in Uganda


The disease is most common in the eastern and central regions.



Graph: Uganda. Leprosy - registered prevalence, cases

Notes:

- Individual years:
- 1975 - True number estimated at 100,000 cases.
- 1991 - True number estimated at 30,000 cases.



Graph: Uganda. Leprosy, cases

MDT coverage is 100% (1998).

References

1. Lancet 2004 Apr 10;363(9416):1209-19.
2. Int J Infect Dis 2012 Nov 14;
3. Med Oral Patol Oral Cir Bucal 2008 Aug ;13(8):E479-82.
4. Indian J Dermatol Venereol Leprol 2009 Mar-Apr;75(2):177-9.
5. Lepr Rev 2008 Sep ;79(3):331-4.
6. Lepr Rev 2008 Sep ;79(3):320-4.
7. Lepr Rev 2008 Sep ;79(3):303-10.
8. Rev Inst Med Trop Sao Paulo 2008 Jan-Feb;50(1):47-50.
9. J Clin Rheumatol 2011 Dec ;17(8):432-5.
10. Minn Med 2008 Nov ;91(11):30-3.
11. Int J Dermatol 2009 Feb ;48(2):150-3.
12. Dermatol Online J 2009 ;15(4):9.
13. Acta Derm Venereol 2012 Mar 20;
14. J Clin Rheumatol 2011 Dec ;17(8):432-5.
15. Eur J Clin Microbiol Infect Dis 2009 Jan ;28(1):1-7.
16. Biomedica 2012 Mar ;32(1):13-22.
17. J Indian Med Assoc 2008 Mar ;106(3):165-6.
18. Rheumatology (Oxford) 2010 Dec ;49(12):2237-42.
19. Eur Spine J 2010 Jul ;19 Suppl 2:S211-5.
20. Int J Dermatol 2009 Sep ;48(9):984-8.
21. Mem Inst Oswaldo Cruz 2012 Dec ;107 Suppl 1:34-42.
22. Bull Soc Pathol Exot 2011 Feb ;104(1):6-9.
23. Indian J Lepr 2011 Apr-Jun;83(2):101-2.
24. Case Rep Dermatol Med 2012 ;2012:758093.

Leptospirosis

Agent	BACTERIUM. <i>Leptospira interrogans</i> An aerobic non-gram staining spirochete
Reservoir	Cattle Dog Horse Deer Rodent Fox Marine mammal Cat Marsupial Frog
Vector	None
Vehicle	Water Soil urine contact
Incubation Period	7d - 12d (range 2d - 26d)
Diagnostic Tests	Culture on specialized media. Dark field microscopy of urine, CSF. Serology.
Typical Adult Therapy	Doxycycline 100 mg BID X 5 to 7d
Typical Pediatric Therapy	Age >= 8y: Doxycycline 2.2 mg/kg BID X 5 to 7d. Age < 8y: IV Penicillin G 50,000u/kg q6h X 5 to 7d
Clinical Hints	"Sterile" meningitis, nephritis, hepatitis, myositis and conjunctivitis; often follows recent skin contact with fresh water in rural or rodent-infested areas; case-fatality rates of 5% to 40% are reported.
Synonyms	Andaman hemorrhagic fever, Canefield fever, Canicola fever, Field fever, Fish handler's disease, Fort Bragg fever, Japanese autumnal fever, <i>Leptospira</i> , Leptospirose, Leptospirosen, Leptospirosi, Mud fever, Pre-tibial fever, Rat fever, Rice field fever, Swamp fever, Swineherd disease, Weil's disease. ICD9: 100 ICD10: A27

Clinical

WHO Case definition for surveillance:

Clinical description

Acute febrile illness with headache, myalgia and prostration associated with any of the following symptoms:

- conjunctival suffusion
- meningeal irritation
- anuria or oliguria and/or proteinuria
- jaundice
- hemorrhages (from the intestines; lung bleeding is notorious in some areas)
- cardiac arrhythmia or failure
- skin rash

and a history of exposure to infected animals or an environment contaminated with animal urine.

Other common symptoms include nausea, vomiting, abdominal pain, diarrhea, arthralgia.

Laboratory criteria for diagnosis

- Isolation (and typing) from blood or other clinical materials through culture of pathogenic leptospire
- Positive serology, preferably Microscopic Agglutination Test (MAT), using a range of *Leptospira* strains for antigens that should be representative of local strains

Case classification

- Suspected: A case that is compatible with the clinical description.
- Probable: Not applicable.
- Confirmed: A suspect case that is confirmed in a competent laboratory.

Note: Leptospirosis is difficult to diagnose clinically in areas where diseases with symptoms similar to those of leptospirosis occur frequently.

SPECIAL ASPECTS

- Serology by Microscopic Agglutination Test (MAT) may provide presumptive information on causative serogroups.
- Attempts should be made to isolate leptospire, and isolates should be typed to assess locally circulating serovars.
- Questioning the patient may provide clues to infection source and transmission conditions.
- Animal serology may give presumptive information on serogroup status of the infection Isolation followed by typing gives definite information on serovar.

Disease due to *Leptospira interrogans* serovar. *icterohaemorrhagiae* is usually overt, and often manifest as hepatitis, meningitis and nephritis. ¹

- Canicola fever is due to serovar. *canicola* (occasionally *L. interrogans* serovar. *pomona*) and characterized by a milder lymphocytic meningitis, without hepatic or renal involvement.
- Serovar. *autumnalis* (occasionally *L. interrogans* serovar. *pomona*) produces Fort Bragg fever, a febrile illness associated with raised, erythematous, and mildly tender pretibial skin lesions.

Acute phase

Subclinical infection is common.

- Overt leptospirosis (90% of cases) is characterized by a self-limited, systemic illness.
- Patients are at risk for severe and potentially fatal illness which may present with renal failure, liver failure, pneumonia²³ or hemorrhagic diathesis.⁴
- Illness begins abruptly with such symptoms as fever (38 to 40 C), headache (over 95% of cases), rigors, myalgia (over 80%), conjunctivitis (30 to 40%), abdominal pain (30%), vomiting (30 to 60%), diarrhea (15 to 30%), cough, muscular (calf) tenderness, pharyngitis (20%) and a pretibial maculopapular rash (fewer than 10%).
- Additional findings may include lymphadenopathy, splenomegaly, atypical lymphocytosis⁵, thrombocytopenia, transitory paraparesis⁶, hepatomegaly or pancreatitis.⁷
- During the acute illness, bacteria can be recovered from or seen in blood, CSF, or tissue using specialized techniques.
- Organisms are demonstrated in urine after the 5th to 7th days. Pyuria, hematuria and proteinuria may be evident as well.
- Severe hypomagnesemia has been reported during the acute phase of infection.⁸⁻¹⁰

Latency and relapse:

The acute phase is followed by an asymptomatic period of 4 to 30 days.

- At this point, illness reappears, with conjunctival suffusion, photophobia, eye pain, myalgia, lymphadenopathy and hepatosplenomegaly.
- Additional findings may lymphocytic meningitis (70 to 80% of patients) with normal glucose levels; pretibial purpura, uveitis¹¹¹², iridocyclitis or chorioretinitis, facial nerve palsy¹³, thrombocytopenia, hypotension, myopericarditis¹⁴ and pancreatitis.¹⁵
- Weil's disease is characterized by hepatic and renal function which may progress to severe and even fatal hepatorenal failure which carries a case-fatality rate of 5 to 40%.
- Renal involvement, principally interstitial nephritis and tubular necrosis¹⁶ may be severe, even in the absence of jaundice.¹⁷
- Pulmonary infiltrates¹⁸, severe hemorrhagic pneumonia¹⁹²⁰ and acute pulmonary distress syndrome may be encountered, even in the absence of hepatic and renal failure.²¹⁻²³
- Congestive heart failure is rare; however, cardiac arrhythmias may occur and result in sudden deaths.
- Acute disseminated encephalomyelitis has been reported as a complication of leptospirosis.²⁴
- Relatively severe infection is reported among pregnant women.²⁵

Persistent, asymptomatic renal colonization by *Leptospirae* may follow infection in humans.²⁶

The clinical features of dengue²⁷, influenza pneumonia²⁸ and pyomyositis may mimic those of leptospirosis.²⁹

This disease is endemic or potentially endemic to all countries.

Leptospirosis in Uganda

Leptospirosis, cases: None reported between 1999 and 2003

References

1. *Curr Opin Infect Dis* 2001 Oct ;14(5):527-38.
2. *Emerg Infect Dis* 2002 Sep ;8(9):1004-5.
3. *Emerg Infect Dis* 2009 May ;15(5):834-5.
4. *Lancet Infect Dis* 2003 Dec ;3(12):757-71.
5. *Rev Soc Bras Med Trop* 2011 Oct ;44(5):611-5.
6. *J Microbiol Immunol Infect* 2012 Feb ;45(1):75-8.
7. *Am J Trop Med Hyg* 2007 May ;76(5):886-7.
8. *Am J Trop Med Hyg* 2008 Dec ;79(6):915-7.
9. *Am J Trop Med Hyg* 2009 Jun ;80(6):1067.
10. *Nat Clin Pract Nephrol* 2008 Feb ;4(2):91-101.
11. *Medicine (Baltimore)* 2008 May ;87(3):167-76.
12. *Am J Ophthalmol* 1997 Jul ;124(1):71-9.
13. *Braz J Infect Dis* 2009 Aug ;13(4):319-21.
14. *Arch Cardiol Mex* 2009 Oct-Dec;79(4):268-73.
15. *BMC Infect Dis* 2013 Jan 24;13(1):38.
16. *J Bras Nefrol* 2010 Dec ;32(4):400-7.
17. *Clin Nephrol* 2009 Sep ;72(3):186-92.
18. *Br J Radiol* 1981 Nov ;54(647):939-43.
19. *Emerg Infect Dis* 2011 Dec ;17(12):2375-7.
20. *BMC Infect Dis* 2013 Jan 24;13(1):38.
21. *Clin Infect Dis* 2005 Feb 1;40(3):343-51.
22. *W V Med J* 2010 Nov-Dec;106(7):20-2.
23. *Neth J Med* 2012 Jun ;70(5):215-21.
24. *Pediatr Neurol* 2009 Jun ;40(6):471-3.
25. *Eur J Clin Microbiol Infect Dis* 2012 May 2;
26. *PLoS Negl Trop Dis* 2010 ;4(2):e612.
27. *Trop Doct* 2010 Apr ;40(2):92-4.
28. *Emerg Infect Dis* 2011 Jun ;17(6):1145-6.
29. *Trop Doct* 2008 Oct ;38(4):254-6.

Listeriosis

Agent	BACTERIUM. <i>Listeria monocytogenes</i> A facultative gram-positive bacillus
Reservoir	Mammal Human Bird Soil Water
Vector	None
Vehicle	Transplacental Dairy products (eg, soft cheeses), Infected secretions Vegetables Poultry Water
Incubation Period	3d - 21d (-60d post-ingestion)
Diagnostic Tests	Culture of blood or CSF.
Typical Adult Therapy	Ampicillin 2g IV q6h X 2w (higher dosage in meningitis) + Gentamicin . Sulfamethoxazole/trimethoprim recommended for Penicillin-allergic patients
Typical Pediatric Therapy	Ampicillin 50 mg/kg IV Q6h X 2w (higher dosage in meningitis). Sulfamethoxazole/trimethoprim recommended for Penicillin-allergic patients
Clinical Hints	Meningitis or sepsis, often immune-suppressed patients (lymphoma, AIDS, etc); gastroenteritis - may follow ingestion of 'over-the-counter' foods; neonatal septicemia occasionally encountered.
Synonyms	<i>Listeria monocytogenes</i> , Listeriose, Listeriosi. ICD9: 027.0 ICD10: A32

Clinical

Major risk factors for invasive Listeriosis reflect T-cell mediated immune compromise, including old age, pregnancy, hematological malignancy, chemotherapy, corticosteroid therapy and anti-TNF-alpha agents

Signs of *Listeria* meningitis are often atypical: ¹

- brain stem and cerebellar involvement (rhombencephalitis) occurs in 11% of cases ^{2 3}
- nuchal rigidity in only 80% to 85%
- movement disorders (ataxia, myoclonus) in 15% to 20% ⁴
- seizures in 25%. ⁵

The blood culture is positive in 75% of meningitis cases; and the cerebrospinal fluid gram stain is positive in only 40%.

Symptoms of food-borne listeriosis develop between one day and three months after ingestion the bacteria in food. ⁶

- Most cases are characterized by diarrhea and fever ^{7 8}
- Headache, myalgia and arthralgia are common. ⁹
- The bacteria may be excreted in stool for several months.

Other forms of listeriosis:

- Hepatic listeriosis may present as single or multiple abscesses, or diffuse granulomatous hepatitis. ¹⁰
- Numerous cases of *Listeria* endocarditis of both native and prosthetic valves have been reported. ¹¹⁻²⁵ Instances of pericarditis ²⁶, cardiac pseudotumor ²⁷, and aortitis with aortic dissection have also been reported. ²⁸
- Sporadic cases of prosthetic joint infection ²⁹⁻³², renal failure, brain abscess ³³, cutaneous infection ³⁴, mycotic aortic aneurysm ^{35 36}, pericarditis ³⁷, uveitis ³⁸, cholecystitis ^{39 40} and rhabdomyolysis have been reported. ⁴¹
- *Listeria* peritonitis has been reported in a patient undergoing peritoneal dialysis ⁴² and in a patient with biliary cirrhosis. ⁴³ Spontaneous bacterial peritonitis due to *Listeria monocytogenes* ⁴⁴⁻⁴⁸ and infection of a ventriculo-peritoneal shunt have also been reported. ⁴⁹

This disease is endemic or potentially endemic to all countries.

Listeriosis in Uganda

Listeriosis, cases: None reported between 1999 and 2003

References

1. *Curr Infect Dis Rep* 2008 Jul ;10(4):300-6.
2. *Scand J Infect Dis* 2005 ;37(3):190-4.
3. *Medicine (Baltimore)* 2011 Jul ;90(4):256-61.
4. *Rev Med Chil* 2009 Dec ;137(12):1602-6.
5. *FEMS Immunol Med Microbiol* 2003 Apr 1;35(3):173-5.
6. *FEMS Immunol Med Microbiol* 2003 Apr 1;35(3):173-5.
7. *N Engl J Med* 1997 Jan 9;336(2):130-2.
8. *N Engl J Med* 1997 Jan 9;336(2):100-5.
9. *Clin Infect Dis* 2005 May 1;40(9):1327-32.
10. *Infection* 2007 Jun ;35(4):212-8.
11. *Eur J Intern Med* 2008 Jun ;19(4):295-6.
12. *Am J Emerg Med* 2007 Nov ;25(9):1086.e3-5.
13. *Intern Med* 2007 ;46(15):1209-12.
14. *Int J Cardiol* 2007 Jun 12;118(3):e106-7.
15. *Int J Antimicrob Agents* 2006 Nov ;28(5):480-1.
16. *Nihon Naika Gakkai Zasshi* 2005 Dec 10;94(12):2609-12.
17. *Scand J Infect Dis* 2004 ;36(10):709-11.
18. *Echocardiography* 2004 Jul ;21(5):423-7.
19. *Int J Infect Dis* 2004 Mar ;8(2):97-102.
20. *Scand J Infect Dis* 2002 ;34(5):383-4.
21. *Transpl Infect Dis* 1999 Dec ;1(4):284-7.
22. *Rev Esp Cardiol* 2000 Feb ;53(2):300.
23. *Acta Clin Belg* 1994 ;49(2):95-8.
24. *Rev Infect Dis* 1988 May-Jun;10(3):616-23.
25. *Scand J Infect Dis* 1988 ;20(4):359-68.
26. *Jpn J Infect Dis* 2012 Jul ;65(4):312-4.
27. *J Infect* 2009 Feb ;58(2):161-3.
28. *Gen Thorac Cardiovasc Surg* 2008 Aug ;56(8):417-20.
29. *Enferm Infecc Microbiol Clin* 2009 Aug-Sep;27(7):389-93.
30. *Acta Ortop Mex* 2009 Sep-Oct;23(5):302-5.
31. *Klin Mikrobiol Infekc Lek* 2011 Apr ;17(2):62-6.
32. *Clin Infect Dis* 2012 Jan 15;54(2):240-8.
33. *Enferm Infecc Microbiol Clin* 2010 Feb ;28(2):87-94.
34. *Br J Hosp Med (Lond)* 2009 Nov ;70(11):659.
35. *J Vasc Surg* 2010 Aug ;52(2):456-9.
36. *Vasc Endovascular Surg* 2011 Nov ;45(8):773-4.
37. *Acta Cardiol* 2011 Aug ;66(4):537-8.
38. *Case Report Ophthalmol* 2010 ;1(2):63-65.
39. *Acta Clin Belg* 2012 Jul-Aug;67(4):295-7.
40. *Am J Med Sci* 2012 Dec 5;
41. *Neurocrit Care* 2009 ;10(1):70-2.
42. *Int Urol Nephrol* 2008 ;40(3):815-9.
43. *J Clin Pathol* 2010 Sep ;63(9):835-6.
44. *Case Rep Gastroenterol* 2008 ;2(3):321-5.
45. *Z Gastroenterol* 2011 Jul ;49(7):832-5.
46. *J Infect Public Health* 2011 Sep ;4(4):211-6.
47. *Scand J Gastroenterol* 2012 Oct ;47(10):1129-40.
48. *Ann Hepatol* 2012 Nov ;11(6):955-7.
49. *J Clin Microbiol* 2011 Nov ;49(11):3924-7.

Liver abscess - bacterial

Agent	BACTERIUM. Various species from portal (Bacteroides, mixed aerobe-anaerobe) or biliary (<i>Escherichia coli</i> , etc) source
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Ultrasonography, CT or radionucleotide scan. If amoebic abscess suspected, perform Entamoeba serology
Typical Adult Therapy	Intravenous antibiotic(s) directed at likely or suspected pathogens. Percutaneous or open drainage
Typical Pediatric Therapy	As for adult
Clinical Hints	Tender liver, and prolonged fever in a patient with history of diverticulosis, cholecystitis, appendicitis, etc; clinically similar to amoebic abscess, but often multiple.
Synonyms	Ascesso fegato, Bacterial liver abscess, Hepatic abscess - bacterial, Liver abscess. ICD9: 572.0 ICD10: K75.0

Clinical

Symptoms of pyogenic hepatic abscess include fever and rigors of several days' to several weeks' duration.

- Dull right upper quadrant pain may be associated with cough and pleuritic pain with radiation to the right shoulder and an associated pleural rub. ¹
- Tender hepatomegaly is present in 50 to 70% of the patients.
- Jaundice is uncommon, unless the abscess is extensive or associated with ascending.
- In some cases, the sole presentation may be fever of unknown origin.

Serological studies, a history of diarrhea, edema of the right chest wall, and limitation to a single abscess in the posterior, superior right hepatic lobe may be suggestive of amoebic abscess. ^{2 3}

Alkaline phosphatase is the most consistently elevated serum enzyme in patients with liver abscess.

- Blood cultures are positive in 50% of cases.

This disease is endemic or potentially endemic to all countries.

References

1. Infect Dis Clin North Am 2000 Sep ;14(3):547-63, vii.
2. Curr Gastroenterol Rep 2004 Aug ;6(4):273-9.
3. Trop Med Int Health 2004 Jun ;9(6):718-23.

Loiasis

Agent	PARASITE - Nematoda. Phasmidea, Filariae: Loa loa
Reservoir	Human
Vector	Fly (deer fly = Chrysops)
Vehicle	None
Incubation Period	4m - 3y
Diagnostic Tests	Microfilariae in blood (take during daylight hours). Adult worm recovered. Serology. Nucleic acid amplification.
Typical Adult Therapy	Diethylcarbamazine : 50 mg PO day 1 50 mg PO TID day 2 100 mg PO TID day 3 3 mg/kg PO TID days 4 to 21. Notes: Ivermectin may cause encephalopathy if dual infection with <i>Onchocerca</i> . Albendazole = alternative to DEC in some cases
Typical Pediatric Therapy	As for adult
Clinical Hints	Migrating pruritic or painful subcutaneous nodules (Calabar swellings), fever and eosinophilia; adult worms may exit to subconjunctival space.
Synonyms	Calabar swellings, Filaria lacrimalis, Filaria loa, Filaria oculi humani, Filaria subconjunctivalis, Fugitive swellings, Loa loa, Microfilaria diurna. ICD9: 125.2 ICD10: B74.3

Clinical

In many cases, infected patients are asymptomatic, with eosinophilia as their sole clinical finding. ¹

- Transient areas of localized subcutaneous edema (Calabar swellings) are preceded by overlying urticaria, pain and itching for several hours. ²
- The lesions are nonerythematous, measure 10 to 20 cm, and last for days to weeks.
- Calabar swellings are most common around joints and may be painful (74%) or painless (26%). ³
- Often, an adult worm will pass beneath the conjunctiva, resulting in local swelling, pain and inflammation. ⁴

Complications include endomyocardial fibrosis, retinopathy, encephalopathy ⁵, peripheral neuropathy ⁶, arthritis, pleural or peritoneal effusions ⁷, hypoechogenic splenic lesions and breast calcification.

This disease is endemic or potentially endemic to 17 countries.

Loiasis in Uganda

Controversy regarding the existence of loiasis in this country are based on the local occurrence of 'Kampala eye worm', variously ascribed to *Loa Loa*, *Mansonella perstans* and a 'Wuchereria bancrofti-like worm.'

Microfilaremia prevalence approaching 7% was demonstrated in Toro district in 1961.

Loiasis was identified in 0.05% of pregnant women in Entebbe (2004 to 2005) ⁸

Some reports may represent importations from southern Sudan.

2011 (publication year) - A Brazilian traveler acquired loiasis in Uganda. ⁹

References

1. Ann Trop Med Parasitol 2006 Dec ;100(8):715-31.
2. Dermatol Clin 1989 Apr ;7(2):313-21.
3. Ann Trop Med Parasitol 2007 Mar ;101(2):151-60.
4. Am J Trop Med Hyg 1989 Jan ;40(1):40-5.
5. West Afr J Med 2007 Apr-Jun;26(2):156-9.
6. Am J Trop Med Hyg 2011 May ;84(5):733-7.
7. J Travel Med 2012 May ;19(3):186-8.
8. PLoS Negl Trop Dis 2009 ;3(6):e473.
9. Rev Inst Med Trop Sao Paulo 2011 Sep-Oct;53(5):295-7.

Lymphocytic choriomeningitis

Agent	VIRUS - RNA. Arenaviridae, Arenavirus: Lymphocytic choriomeningitis virus
Reservoir	House mouse Guinea pig Hamster Monkey
Vector	None
Vehicle	Urine Saliva Feces Food Dust
Incubation Period	8d - 12d (range 6d - 14d)
Diagnostic Tests	Viral culture (blood, throat, CSF). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Headache, myalgia, meningitis and encephalitis; photophobia or pharyngitis may be present; prior exposure to rodents; infection resolves within 2 weeks, however convalescence may require an additional 2 months.
Synonyms	

Clinical

Acute infection:

35% of Lymphocytic choriomeningitis virus infections are asymptomatic and 50% are characterized by a nonspecific flu-like illness.

- Overt infections are characterized by fever, headache, nausea and systemic symptoms, leukopenia and thrombocytopenia. ^{1 2}
- Patients may also exhibit lymphadenopathy and a maculopapular rash (12% to 15% of patients have rash and/or meningitis or encephalitis).
- Relapses characterized by a more severe headache with meningitis may occur after initial improvement.
- Papilledema may be noted

The CSF protein concentration ranges from 50 to 300 mg/dl.

- A pleocytosis of several hundred lymphocytes/mm³ is commonly observed.
- Decreases in CSF glucose concentration are documented in over 20% of cases.

Complications:

Complications include encephalitis, psychosis, paraplegia, transitory aqueductal stenosis, and disturbances of cranial, sensory, or autonomic nervous function.

- Occasionally, orchitis, myocarditis, arthritis, or alopecia are encountered.
- Lymphocytic choriomeningitis is increasingly recognized as a cause of hydrocephalus, psychomotor retardation, congenital chorioretinitis and blindness, most often when acquired during the first or second trimesters of pregnancy. ^{3 4}
- Congenital infection is also associated with microencephaly, periventricular calcifications, ventriculomegaly, pachygyria, cerebellar hypoplasia, porencephalic and periventricular cysts. ⁵

The case-fatality rate for Lymphocytic choriomeningitis is less than one percent; however, patients with sustained viremia lacking an inflammatory response seem to be at risk for fatal outcome. ⁶

This disease is endemic or potentially endemic to all countries.

References

1. Semin Pediatr Infect Dis 2003 Apr ;14(2):72-82.
2. ProMED <promedmail.org> archive: 20050804.2273
3. MMWR Morb Mortal Wkly Rep 2005 Jun 3;54(21):537-9.
4. Semin Pediatr Neurol 2012 Sep ;19(3):89-95.
5. Ann Neurol 2007 Oct ;62(4):347-55.
6. N Engl J Med 2006 May 25;354(21):2208-11.

Lymphogranuloma venereum

Agent	BACTERIUM. Chlamydiaceae, Chlamydiae , Chlamydia trachomatis, types L1, L2, L3
Reservoir	Human
Vector	None
Vehicle	Sexual contact
Incubation Period	7d - 12d (range 3d - 30d)
Diagnostic Tests	Serology. Culture of pus performed in specialized laboratories.
Typical Adult Therapy	Doxycycline 100 mg PO BID X 3w. OR Erythromycin 500 mg QID X 3w
Typical Pediatric Therapy	Age < 8 years: Erythromycin 10 mg/kg PO QID X 2 to 4w. Age >= 8 years: Doxycycline 2 mg/kg PO BID X 2 to 4w
Clinical Hints	Genital nodule or vesicle with large, suppurating regional nodes; generalized lymphadenopathy or proctitis may be present; late complications include genital edema, rectal strictures and perianal abscesses.
Synonyms	Bubonulus, Durand-Nicolas-Favre disease, Linfogranuloma venereo, Lymphogranuloma inguinale, Lymphopathia venereum, Maladie de Nicolas et Favre, Tropical bubo, Venereal bubo, Venerisk lymphogranulom. ICD9: 099.1 ICD10: A55

Clinical

Acute illness:

The first stage of Lymphogranuloma venereum (LGV) is characterized by a papule or ulcer on the genital or anal mucosa, or of the adjacent skin. ¹⁻⁴

- Occasionally, the lesion is intraurethral or cervical, producing urethritis or cervicitis.
- The secondary stage occurs days to weeks after the primary lesion and is characterized by lymphadenopathy and systemic illness.
- Cervical lymphadenopathy may occur if infection is acquired through oro-genital contact. ⁵

Lymphadenitis:

The inguinal lymph nodes are most often affected, and are unilateral in two thirds of patients.

- The obturator and iliac nodes are occasionally affected in women.
- Initially the lymph nodes are discrete and tender with overlying erythema.
- A characteristic "groove" may be evident between the femoral and inguinal lymph nodes.
- In some cases, patients may present with a "bubonulus": penile adenopathy and secondary local acute lymphedema. ⁶
- Later, the nodes may suppurate and coalesce, forming a bubo that may rupture spontaneously (30% of cases) to produce fistulae ⁷ or sinus tracts which may drain for months.

Inguinal lymphadenopathy in cat-scratch disease may suggest a diagnosis of lymphogranuloma venereum. ⁸

- Rectal involvement may suggest a diagnosis of inflammatory bowel disease. ^{9 10}

Systemic manifestations at this stage include fever, headache, and myalgia.

- Meningitis may also occur.
- LGV is increasingly recognized as a cause of hemorrhagic proctitis in men who have sex. ¹¹
- Reactive arthritis has been reported following LGV proctitis ^{12 13}

Relapse occurs in 20% of untreated patients.

Only 25% of women present with inguinal lymphadenopathy.

- Women and homosexual men may present with proctitis or pain in the lower abdomen and back pain related to involvement of pelvic and lumbar lymph nodes.
- Late complications include esthiomene (chronic hypertrophic and ulceration of the vulva, scrotum or penis), and elephantiasis of the male or female genitalia.
- Major lower leg involvement may suggest a diagnosis of deep-vein thrombosis. ¹⁴

This disease is endemic or potentially endemic to all countries.

References

1. Sex Transm Infect 2002 Apr ;78(2):90-2.
2. J Am Acad Dermatol 1999 Oct ;41(4):511-32.
3. Med Clin North Am 1998 Sep ;82(5):1081-104, vi.
4. Sex Transm Infect 2011 Mar ;87(2):123-4.
5. Ear Nose Throat J 2008 Aug ;87(8):478-80.
6. Sex Transm Infect 2007 Jul ;83(4):337-8.
7. Infect Dis Obstet Gynecol 1999 ;7(4):199-201.
8. Int J STD AIDS 2009 Aug ;20(8):585-6.
9. Scand J Gastroenterol 2011 Apr ;46(4):503-10.
10. World J Gastroenterol 2012 Jul 7;18(25):3317-21.
11. Curr Infect Dis Rep 2007 Mar ;9(2):143-50.
12. Sex Transm Infect 2009 Jun ;85(3):180-1.
13. Int J STD AIDS 2011 Jan ;22(1):59-60.
14. Int J STD AIDS 2012 Mar ;23(3):219-20.

Malaria

Agent	PARASITE - Protozoa. Sporozoa, Coccidea, Haemosporida: Plasmodium spp.
Reservoir	Human Primate (Plasmodium knowlesi)
Vector	Mosquito (Anopheles)
Vehicle	Blood
Incubation Period	12d -30d
Diagnostic Tests	Examination of blood smear. Serology, antigen & microscopic techniques. Nucleic acid amplification.
Typical Adult Therapy	Resistant falcip: Lumefantrine/Artemether OR Quinine + Doxycycline or Clindamycin OR Atovaquone/proguanil OR Artesunate IV if severe malaria If sens., Chloroquine 1g, then 500 mg at 6, 24 & 48 hrs. If P. ovale or P. vivax - follow with Primaquine
Typical Pediatric Therapy	Resistant falcip: Lumefantrine/Artemether OR Quinine + Clindamycin OR Atovaquone/proguanil OR Artesunate (>age 8) IV (severe malaria) If sens, Chloroquine 10 mg/kg, then 5 mg/kg at 6, 24, & 48 hrs. If P. ovale or P. vivax - follow with Primaquine
Clinical Hints	Fever, headache, rigors ("shaking chills"), vomiting, myalgia, diaphoresis and hemolytic anemia; fever pattern (every other or every third day) and splenomegaly may be present; clinical disease may relapse after 7 (ovale and vivax) to 40 (malariae) years.
Synonyms	Ague, Bilious remittent fever, Chagres fever, Estiautumnal fever, March fever, Marsh fever, Paludism, Paludismo, Plasmodium falciparum, Plasmodium knowlesi, Plasmodium malariae, Plasmodium ovale, Plasmodium vivax. ICD9: 084 ICD10: B50,B51,B52,B53,B54

Clinical

WHO Case definition for surveillance (For use in endemic areas and people exposed to malaria, e.g., a history of visit to endemic area).

- Malaria must be defined in association with clinical disease symptoms.
- The case definition for malaria cannot be uniform: it will vary according to how malaria is perceived in a given country, local patterns of transmission, and disease consequences.
- The suggested definitions are deliberately broad.
- Each national malaria control program must adapt the definition and introduce additional indicators to make it more applicable to local epidemiology and control targets.

Clinical description

- Signs and symptoms vary; most patients experience fever.
- Splenomegaly and anemia are commonly associated signs.
- Common but non-specific symptoms include otherwise unexplained headache, back pain, chills, sweating, myalgia, nausea, vomiting.
- Untreated *Plasmodium falciparum* infection can lead to coma, generalized convulsions, hyperparasitemia, normocytic anemia, disturbances of fluid, electrolyte, and acid-base balance, renal failure, hypoglycemia, hyperpyrexia, hemoglobinuria, circulatory collapse / shock, spontaneous bleeding (disseminated intravascular coagulation), pulmonary edema, and death.

Laboratory criteria for diagnosis

Demonstration of malaria parasites in blood films (mainly asexual forms).

Case classification

In areas without access to laboratory-based diagnosis.

- Probable uncomplicated malaria: A person with symptoms and/or signs of malaria who receives anti-malarial treatment.
- Probable severe malaria: A patient who requires hospitalization for symptoms and signs of severe malaria and receives anti-malarial treatment.
- Probable malaria death: death of a patient diagnosed with probable severe malaria.

In areas with access to laboratory-based diagnosis.

- Asymptomatic malaria: A person with no recent history of symptoms and/or signs of malaria who shows laboratory confirmation of parasitemia.
- Confirmed uncomplicated malaria: A patient with symptoms and/or signs of malaria who received anti-malarial treatment, with laboratory confirmation of diagnosis.

Acute infection:

Most cases present with non-specific signs suggestive of 'sepsis,' such as fever, rigors, headache and myalgia.

- Clinical findings include cough, fatigue, malaise, arthralgia, myalgia, headache, and diaphoresis.

- In Africa, tickborne relapsing fever ¹ and rabies are often mis-diagnosed as malaria. ²

The typical malarial paroxysm begins with rigors lasting 1 to 2 hours, followed by high fever.

- This is followed by marked diaphoresis and a fall in temperature.
- Tertian (fever every other day) fever may occur in infection by *P. falciparum*, *P. vivax* and *P. ovale*; quartan (every third day) fever with *P. malariae* infection; and daily fever with *P. knowlesi* infection. ^{3 4}
- *P. knowlesi* malaria appears to be more severe than *P. malariae* malaria, with higher rates of parasitemia and fatality. ⁵⁻⁹
- 'Classical' fever patterns are rarely helpful, and anemia and splenomegaly develop only after several attacks.
- Less common findings include anorexia, vomiting, diarrhea and hypotension.
- In some cases, malaria may present as fever accompanied by an urticarial rash. ¹⁰

Complications:

Complications include pulmonary disease (ARDS) ^{11 12}, encephalopathy (cerebral malaria) ¹³⁻¹⁵, nephropathy, retinopathy ¹⁶⁻¹⁹, cranial nerve palsy, cerebellar ataxia ²⁰, acute disseminated encephalomyelitis ²¹, hypocalcemia with tetany ²², shock ('algid malaria'), purpura fulminans ²³, massive diarrhea, pancreatitis ²⁴, myocarditis ²⁵ and dysfunction of other organs. ^{26 27}

- Patients with *falciparum* malaria are at increased risk for bacteremia. ²⁸
- Occasionally, patients experience Post-malaria Neurological Syndrome: acute confusion, cerebellar ataxia, diffuse cerebral demyelination, seizures, hearing loss ²⁹, cognitive dysfunction ³⁰ or other neuropsychiatric findings several days to weeks following successful treatment of falciparum malaria. ³¹⁻³⁶
- *Plasmodium falciparum* infection accounts for most complications and deaths from malaria ³⁷; however, severe disease may occasionally complicate infection by other species. ³⁸⁻⁴³
- The presence of malarial retinopathy is associated with a poor prognosis. ⁴⁴
- *P. falciparum* is also responsible for most malarial drug resistance.
- Maternal infection is associated with vertical transmission to the newborn ⁴⁵, fetal loss and low birth weight in infants. ⁴⁶⁻⁵⁰
- 5% of African children with severe malaria were found to have concomitant bacteremia ⁵¹
- Severe and fatal disease associated with *Plasmodium vivax* infection is increasingly reported in recent years. ⁵²⁻⁶⁵ Instances of acute glomerulonephritis ⁶⁶⁻⁶⁸, IgA nephropathy ⁶⁹, renal cortical necrosis ⁷⁰, acalculous cholecystitis ⁷¹, jaundice, pancreatitis ⁷², thrombocytopenia ^{73 74}, shock, peripheral gangrene ⁷⁵, cerebral malaria ⁷⁶, hypoglycemia and acute respiratory distress syndrome have been reported with *Plasmodium vivax* infections. ⁷⁷⁻⁸⁰
- *Plasmodium malariae* infection is rarely associated with severe illness ⁸¹; and may lead to renal glomerular damage and nephrotic syndrome. ⁸²⁻⁸⁴
- Pericarditis ⁸⁵ and acute respiratory distress syndrome have been reported in *Plasmodium ovale* infection. ⁸⁶

Malaria and HIV infection:

HIV infection increases the incidence and severity of clinical malaria; however, in severe malaria the level of parasitemia is similar in HIV-positive and HIV-negative patients. ⁸⁷⁻⁹⁵

- During pregnancy, HIV infection increases the incidence of clinical malaria, maternal morbidity, and fetal and neonatal morbi-mortality.
- HIV infection increases the risk for malaria treatment failure, and for cerebral malaria in children. ⁹⁶
- Some antimalarial drugs may inhibit HIV, while certain anti-retroviral drugs are effective against *Plasmodium* species. ⁹⁷

Relapse:

Relapse may occur months to years following the initial episode.

- Relapses of *P. vivax* and *P. ovale* infection result from release of parasites which had remained dormant in the liver.
- As such, treatment of infection by either of these two species should include a drug (eg, primaquine) active against intrahepatic parasites.
- Although infections caused by *P. falciparum* and *P. knowlesi* do not relapse, reinfection may occur. ⁹⁸

Plasmodium malariae persists without symptoms in the blood, rather than the liver.

- Relapse has been reported as long as 40 to 50 years following exit from an endemic area. ⁹⁹

This disease is endemic or potentially endemic to 186 countries. Chloroquine resistant falciparum malaria endemic to 81 countries. Chloroquine-sensitive malaria endemic to 29 countries.

Malaria in Uganda

Time and Place:

Malaria is reported throughout the country during all seasons. ¹⁰⁰

- The potential exists for highland transmission in the southwest region.
- 36,112 cases were reported during an epidemic of highland malaria in the southwestern highlands between January to

August 1998. ¹⁰¹ ¹⁰²


- Status report on malaria in Uganda as of 2011 - see reference ¹⁰³
- Analysis of the economic impact of malaria on Uganda (1997 to 2003) - see reference ¹⁰⁴

Infecting species:

- Chloroquine-resistant *P. falciparum* **IS** reported. ¹⁰⁵
- *Plasmodium falciparum* accounts for over 85% of cases.
- *Plasmodium ovale* accounts for 1% to 6% of malaria cases in lake-side and inland districts. (2011 publication) ¹⁰⁶
- Sporadic infection by *Plasmodium vivax* is reported. ¹⁰⁷

Prevalence surveys:


- 35.2% of pregnant women in Masindi District (2003 to 2004) ¹⁰⁸
 - 10.9% of pregnant women in Entebbe (2004 to 2005) ¹⁰⁹
 - 11% of rural women in the third trimester of pregnancy (2006 to 2008) ¹¹⁰
 - 10.9% of otherwise healthy pregnant women at enrolment to a trial of deworming. (Entebbe, 2007 publication) ¹¹¹
 - 9.5% of mothers and 16.7% of their children in Habutobere; 5.6% of mothers and 3.1% of their children in Musezero (southwestern Uganda, 2010) ¹¹²
 - 9.5% of febrile episodes among children in an urban area of low endemicity (Kampala, 2010 publication) ¹¹³
 - 46.4% of school-age children in the Lake Victoria region (2011 publication) ¹¹⁴
 - 71.9% of children below age 5 years (PCR, Apac, northern region, 2011 publication) ¹¹⁵
 - 43% of children below age 5 years in rural areas, in 2004; 23% in 2010 ¹¹⁶
 - 10% of children in a rural parish (2011) ¹¹⁷
 - 20% of infants in a high-transmission area (2012 publication) ¹¹⁸
 - 42.6% in primigravida and 33.3% in multigravida women attending an antenatal clinic in an area of stable malaria transmission. (2007 publication) ¹¹⁹
 - 4% of urban AIDS patients receiving HAART (asymptomatic *P. falciparum* antigenemia, 2004 to 2005) ¹²⁰
 - 19% of patients with visceral leishmaniasis (2000 to 2006) ¹²¹
- 9% of children with cerebral malaria were found to be HIV-positive, vs. 2.3% of children with uncomplicated malaria (2011 publication) ¹²²



Graph: Uganda. Malaria, cases

Notes:

1. Incidence for 2000 to 2005 includes data from reference ¹²³
2. Malaria accounted for 20% of hospital admissions in 1999; 38% in 2000; 38% in 2001. ¹²⁴



Graph: Uganda. Malaria, deaths

Notes:

1. Figures for 1980, 1990, 2000 and 2010 are based on estimates of true mortality.¹²⁵ Since these estimates are significantly higher than official Health Ministry reports for other years during this period, resultant graphs will suggest unusual fluctuation in trends.

2. Malaria accounted for 23% of hospital deaths in 1998; 16% in 1999; 35% in 2001.

Individual years:

2003 - True number estimated at 8,450.

2004 - 490 fatal cases were reported during an outbreak in Kasese (southern region).

Vectors:

42 *Anopheles* species have been identified in Uganda.

- The principal vectors are *Anopheles gambiae*¹²⁶ and *An. funestus*.^{127 128}

- *An. marshalli* is reported to be a secondary vector.

- *An. christyi* was previously (1940's) reported as a vector in the Keigzi Highlands.

- Additional vectors include *An. nili*, *An. arabiensis*¹²⁹, *An. coustani*, *An. paludis*, *An. brohieri*, *An. hancocki*, *An. bwanbae*, *An. quadriannulatus*, and *An. hargreavesi*.

Notable outbreaks:

1998 - An outbreak of highland malaria was reported in the southwestern region.^{130 131}

2013 - An outbreak (30 cases, 5 fatal) was reported in Mubende.^{132 133}

References

- Emerg Infect Dis 2007 Jan ;13(1):117-23.
- Emerg Infect Dis 2007 Jan ;13(1):136-9.
- Trends Parasitol 2008 Sep ;24(9):406-10.
- Trends Parasitol 2011 Oct ;27(10):442-9.
- Clin Infect Dis 2008 Jan 15;46(2):165-71.
- Emerg Infect Dis 2011 Jul ;17(7):1248-55.
- Malar J 2012 ;11:284.
- Clin Infect Dis 2012 Oct 19;
- ProMED <promedmail.org> archive: 20080105.0060
- J Infect Dev Ctries 2012 ;6(12):895-6.
- J Natl Med Assoc 2011 Jan ;103(1):64-7.
- Chest 2012 Aug ;142(2):492-505.
- Brain 2011 May ;134(Pt 5):1519-28.
- Future Microbiol 2012 Feb ;7:291-302.
- Rev Neurol (Paris) 2012 Mar 1;
- Trans R Soc Trop Med Hyg 2009 Jul ;103(7):661-4.
- Trans R Soc Trop Med Hyg 2009 Jul ;103(7):665-71.
- Ocul Immunol Inflamm 2008 Sep-Oct;16(5):239-41.
- Neurology 2012 Sep 18;79(12):1196-7.
- J Infect Dev Ctries 2012 ;6(3):290-4.
- Indian J Pediatr 2012 Jun 15;
- J Assoc Physicians India 2012 Jul ;60:57-8.
- Trans R Soc Trop Med Hyg 2007 Oct ;101(10):1045-7.
- J Assoc Physicians India 2011 Nov ;59:731-3.
- Case Report Med 2011 ;2011:202083.
- Crit Care 2003 Aug ;7(4):315-23.
- Leg Med (Tokyo) 2012 Feb 25;
- Lancet 2011 Oct 8;378(9799):1316-23.
- Ann Trop Paediatr 2011 ;31(1):1-10.
- Malar J 2010 ;9:366.
- J Travel Med 2009 Nov-Dec;16(6):424-30.
- Lancet 1996 Oct 5;348(9032):917-21.

33. *Ann Trop Med Parasitol* 2001 Mar ;95(2):215-7.
34. *Neurol Sci* 2006 Dec ;27(6):442-4.
35. *J Vector Borne Dis* 2007 Sep ;44(3):227-9.
36. *Am J Trop Med Hyg* 2008 May ;78(5):699-701.
37. *Leg Med (Tokyo)* 2012 Feb 25;
38. *Emerg Infect Dis* 2005 Jan ;11(1):132-4.
39. *Trans R Soc Trop Med Hyg* 2007 Jul ;101(7):655-9.
40. *J Travel Med* 2009 Mar-Apr;16(2):138-40.
41. *J Vector Borne Dis* 2009 Jun ;46(2):141-4.
42. *J Clin Neurol* 2010 Jun ;6(2):102-3.
43. *Parasitol Res* 2012 Jun ;110(6):2573-6.
44. *Am J Trop Med Hyg* 2011 Jan ;84(1):141-7.
45. *Malar J* 2011 ;10:239.
46. *Bull World Health Organ* 2007 Jan ;85(1):9-18.
47. *Infect Dis Obstet Gynecol* 2005 Dec ;13(4):229-36.
48. *J Infect Dis* 2011 Mar 1;203(5):691-9.
49. *Trends Parasitol* 2011 Apr ;27(4):168-75.
50. *Lancet Infect Dis* 2012 May ;12(5):388-96.
51. *Trop Med Int Health* 2009 Sep ;14(9):1011-9.
52. *Curr Opin Infect Dis* 2009 Oct ;22(5):430-5.
53. *PLoS Negl Trop Dis* 2011 Jun ;5(6):e1032.
54. *J Infect Public Health* 2011 Jun ;4(2):91-5.
55. *Trop Doct* 2011 Jul ;41(3):168-9.
56. *Ann Trop Paediatr* 2011 ;31(4):351-6.
57. *Lancet Infect Dis* 2012 May ;12(5):388-96.
58. *Parasitol Res* 2012 Jun ;110(6):2573-6.
59. *Malar J* 2012 ;11:12.
60. *Trop Doct* 2012 Apr ;42(2):92-3.
61. *Malar J* 2012 May 2;11(1):144.
62. *Clin Infect Dis* 2012 Oct ;55(8):e67-74.
63. *Parasit Vectors* 2012 Jul 30;5(1):154.
64. *Trop Doct* 2012 Aug 2;
65. *Clin Microbiol Rev* 2013 Jan ;26(1):36-57.
66. *Ann Trop Paediatr* 2011 ;31(2):181-4.
67. *Trop Doct* 2012 Jan ;42(1):63-4.
68. *Parasitol Res* 2012 Jul 18;
69. *J Korean Med Sci* 2012 Apr ;27(4):446-9.
70. *Parasitol Res* 2012 Nov ;111(5):2213-6.
71. *Am J Trop Med Hyg* 2011 Jul ;85(1):42-9.
72. *JOP* 2012 ;13(2):215-6.
73. *Ann Trop Med Parasitol* 2011 Dec ;105(8):593-8.
74. *Trop Doct* 2012 Aug 2;
75. *Paediatr Int Child Health* 2012 ;32(3):164-6.
76. *Asian Pac J Trop Med* 2012 Aug ;5(8):665-6.
77. *Am J Trop Med Hyg* 2009 Nov ;81(5):758-62.
78. *Harefuah* 2010 Sep ;149(9):580-2, 620.
79. *Emerg Infect Dis* 2012 May ;18(5):842-5.
80. *Chest* 2012 Aug ;142(2):492-505.
81. *Emerg Infect Dis* 2009 May ;15(5):832-4.
82. *Lancet* 1960 Apr 9;1(7128):806-7.
83. *J Travel Med* 2011 Jul-Aug;18(4):288-91.
84. *Ann Pathol* 2012 Feb ;32(1):40-52.
85. *Med Trop (Mars)* 2011 Feb ;71(1):79-80.
86. *Am J Trop Med Hyg* 2008 Sep ;79(3):391-3.
87. *Med Mal Infect* 2007 Oct ;37(10):629-36.
88. *Malar J* 2007 ;6:143.
89. *Clin Infect Dis* 2007 Nov 1;45(9):1208-13.
90. *Malar J* 2007 ;6:143.
91. *Lancet Infect Dis* 2011 Jul ;11(7):541-56.
92. *Sante* 2011 Jul 1;21(3):174-177.
93. *Mediterr J Hematol Infect Dis* 2012 ;4(1):e2012032.
94. *Future Virol* 2012 ;7(7):699-708.
95. *Parasit Vectors* 2013 Jan 17;6(1):18.
96. *BMC Pediatr* 2011 ;11:5.
97. *Trends Parasitol* 2008 Jun ;24(6):264-71.
98. *Emerg Infect Dis* 2011 Jul ;17(7):1314-5.
99. *Clin Microbiol Rev* 2007 Oct ;20(4):579-92.
100. *Acta Trop* 2012 Mar ;121(3):184-95.
101. *Trans R Soc Trop Med Hyg* 1999 Sep-Oct;93(5):480-7.
102. *Trop Med Int Health* 2000 Dec ;5(12):865-75.
103. *Acta Trop* 2012 Mar ;121(3):196-201.
104. *Int Arch Med* 2012 Mar 22;5(1):12.
105. *Acta Trop* 2005 Sep ;95(3):172-82.
106. *Int J Parasitol* 2011 May ;41(6):677-83.
107. *PLoS One* 2011 ;6(5):e19801.
108. *Trans R Soc Trop Med Hyg* 2008 Feb ;102(2):130-6.
109. *PLoS Negl Trop Dis* 2009 ;3(6):e473.
110. *PLoS One* 2011 ;6(7):e21862.
111. *Trans R Soc Trop Med Hyg* 2007 Sep ;101(9):899-907.
112. *Trans R Soc Trop Med Hyg* 2012 Feb ;106(2):75-83.
113. *J Clin Microbiol* 2010 Oct ;48(10):3539-43.
114. *Parasit Vectors* 2011 ;4:232.
115. *Am J Trop Med Hyg* 2011 May ;84(5):830-7.
116. *Malar J* 2011 ;10:132.
117. *PLoS One* 2012 ;7(8):e43400.
118. *Malar J* 2012 Dec 27;11(1):432.
119. *Trans R Soc Trop Med Hyg* 2008 Feb ;102(2):130-6.
120. *Malar J* 2011 ;10:66.
121. *PLoS Negl Trop Dis* 2012 Apr ;6(4):e1617.
122. *BMC Pediatr* 2011 ;11:5.
123. *Am J Trop Med Hyg* 2007 Dec ;77(6 Suppl):36-47.
124. *BMC Med* 2011 ;9:37.
125. *Lancet* 2012 Feb 4;379(9814):413-31.
126. *Malar J* 2005 ;4:59.
127. *Mol Ecol* 2005 Dec ;14(14):4235-48.
128. *Am J Trop Med Hyg* 2006 Aug ;75(2):219-25.
129. *Malar J* 2006 ;5:16.
130. *Trans R Soc Trop Med Hyg* 1999 Sep-Oct;93(5):480-7.
131. *Trop Med Int Health* 2000 Dec ;5(12):865-75.
132. *ProMED <promedmail.org> archive: 20130131.1523957*
133. *ProMED <promedmail.org> archive: 20130126.1515447*

Malignant otitis externa

Agent	BACTERIUM. Pseudomonas aeruginosa : aerobic gram-negative bacillus (virtually all cases)
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Culture of otic exudate and biopsy material. Careful roentgenographic and neurological examinations.
Typical Adult Therapy	Early debridement complemented by at least 2 parenteral antibiotics active against <i>Pseudomonas aeruginosa</i>
Typical Pediatric Therapy	As for adult
Clinical Hints	Otic pain, swelling and discharge; infection of bony and cartilaginous ear canal; over 80% of patients are diabetics over age 50; cranial nerve (usually VII) signs in 50%. case-fatality rate > 55%.
Synonyms	

Clinical

The case definition of Malignant Otitis Externa consists of pain, edema, exudate, granulations, microabscesses (when explored), positive bone scan or failure of local treatment often more than 1 week. ¹

- Additional criteria may include cranial nerve involvement, positive radiograph, debilitating condition and old age.

Severe pain and tenderness in the mastoid area are accompanied by drainage of pus from the external canal. ²

- Involvement of the temporal bone, meninges, venous sinuses, cranial nerves (IX, X, XII) and brain may follow.

This disease is endemic or potentially endemic to all countries.

References

1. *J Laryngol Otol* 1987 Mar ;101(3):216-21.
2. *Lancet Infect Dis* 2004 Jan ;4(1):34-9.

Mansonelliasis - *M. perstans*

Agent	PARASITE - Nematoda. Phasmidea, Filariae: <i>Mansonella</i> (<i>Esslingeria</i>) <i>perstans</i>
Reservoir	Human
Vector	Midge (<i>Culicoides</i> spp.)
Vehicle	None
Incubation Period	5m - 18m (range 1m - 2y)
Diagnostic Tests	Identification of microfilariae in blood. Nucleic acid amplification.
Typical Adult Therapy	Albendazole 400 mg PO BID X 10 d OR Mebendazole 100 mg PO BID X 30 d. Recent data suggest that addition of doxycycline may be of benefit.
Typical Pediatric Therapy	Age >2 years: As for adult. OR Albendazole 10 mg/kg/day PO X 10d
Clinical Hints	Recurrent pruritic subcutaneous lesions, arthralgia and eosinophilia; headache, fever or abdominal pain may also be present.
Synonyms	<i>Acanthocheilonema perstans</i> , Bung eye disease, <i>Dipetalonema berghei</i> , <i>Dipetalonema perstans</i> , <i>Dipetalonema semiclarum</i> , <i>Esslingeria perstans</i> , <i>Filaria perstans</i> , <i>Mansonella perstans</i> , <i>Mansonella rhodhaini</i> , <i>Mansonella semiclarum</i> , <i>Meningonema peruzzii</i> , <i>Tetrapetalonema berghei</i> , <i>Tetrapetalonema perstans</i> . ICD9: 125.4 ICD10: B74.4

Clinical

Patients develop recurrent pruritic subcutaneous swellings, fever, headache, joint pain, abdominal or chest pain and eosinophilia.

- Hepatosplenomegaly and intraocular lesions are occasionally seen.
- Asymptomatic microfilaremia from transfusion of infected blood has been reported.
- "Bung eye," characterized by the formation of yellowish nodules on the bulbar conjunctivae with proptosis and lid edema, has been reported in Uganda and neighboring countries. ¹
- Microfilariae of *Mansonella perstans* have been identified in vaginal secretions. ^{2 3}

Human cerebral infection by *Meningonema peruzzii* has been reported. ⁴

This disease is endemic or potentially endemic to 49 countries.

Mansonelliasis - *M. perstans* in Uganda

Time and Place:

- M. perstans* infection in Uganda was first reported in 1901. ⁵
- The disease is most common between Lake Victoria and Lake Kioga. ⁶

Prevalence surveys:

- 39% to 81% of adults in the western region (1996 publication) ⁷
- 0.4% to 72.8% of children (61.8% of school districts, 2000 to 2003) ⁸
- 21.3% of otherwise healthy pregnant women at enrolment to a trial of deworming. (Entebbe, 2007 publication) ⁹
- 21% of pregnant women (Entebbe and Katabi, 2003 to 2005) ¹⁰
- 76.5% of persons in Mukono District and 57.7% in Luwero District (2005 to 2006) ¹¹

"**Bung eye**" characterized by the formation of yellowish nodules on the bulbar conjunctivae with proptosis and lid edema, has been reported in Uganda and neighboring countries. ¹²

- The causative agent appears to be a zoonotic *Mansonella* species, possibly *M. vanhoofi*, *M. rodhaini* or *M. leopoldi*.

References

1. Am J Trop Med Hyg 1988 May ;38(3):553-7.
2. Pan Afr Med J 2012 ;12:47.
3. Med Sante Trop 2013 Feb 7;
4. Parasite 1995 Jun ;2(2):173-6.
5. Ann Trop Med Parasitol 2005 Jun ;99(4):383-93.
6. Trans R Soc Trop Med Hyg 2001 Mar-Apr;95(2):161-7.
7. Appl Parasitol 1996 Jan ;37(1):32-7.
8. Ann Trop Med Parasitol 2005 Jun ;99(4):383-93.
9. Trans R Soc Trop Med Hyg 2007 Sep ;101(9):899-907.
10. J Infect Dis 2008 Sep 15;198(6):920-7.
11. Trans R Soc Trop Med Hyg 2009 Mar ;103(3):266-73.
12. Am J Trop Med Hyg 1988 May ;38(3):553-7.

Mansonelliasis - M. streptocerca

Agent	PARASITE - Nematoda. Phasmidea, Filariae: Mansonella (Esslingeria) streptocerca
Reservoir	Non-human primate
Vector	Midge (Culicoides grahami and Cu. milnei)
Vehicle	None
Incubation Period	5m - 18m (speculative)
Diagnostic Tests	Identification of microfilariae in skin snips.
Typical Adult Therapy	Ivermectin 150 ug/kg PO (single dose). OR Diethylcarbamazine : 50 mg PO day 1 50 mg PO TID day 2 100 mg PO TID day 3 Then 2 mg/kg PO TID X 18 days
Typical Pediatric Therapy	As for adult
Clinical Hints	Pruritic dermatitis (usually over chest wall and shoulder area), hypopigmented macules, urticaria and eosinophilia.
Synonyms	Acanthocheilonema streptocerca, Agamofilaria streptocerca, Dipetalonema streptocerca, Esslingeria streptocerca, Mansonella streptocerca. ICD9: 125.6 ICD10: B74.4

Clinical

Infection is rare, and limited to pruritic and hypopigmented macules and papules, principally over the thorax and shoulders.

- Regional edema may occur in some cases.
- Recurrent pruritic subcutaneous lesions, arthralgia and eosinophilia; headache, fever or abdominal pain may also be present.

This disease is endemic or potentially endemic to 16 countries.

Mansonelliasis - M. streptocerca in Uganda

Prevalence surveys:

5% to 89% of adults and 35% of children in the north (Ruwenzori Mountains, 1997 publication) ¹

References

1. [Acta Trop 1997 Jan ;63\(1\):43-55.](#)

Marburg virus disease

Agent	VIRUS - RNA. Mononegavirales, Filoviridae, Filovirus: Marburg virus
Reservoir	? African green monkey ? Bat
Vector	None
Vehicle	Infected secretions Contact Syringe Needle
Incubation Period	5d - 7d (range 3d - 13d)
Diagnostic Tests	Viral culture (blood, body fluids, liver, spleen). Serology. Nucleic acid amplification. Biosafety level 4.
Typical Adult Therapy	Strict isolation; supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, myalgia, arthralgia, sore throat, vomiting, diarrhea, conjunctivitis and maculopapular rash; hemorrhagic diatheses begins on 3rd to 4th day of illness; hepatic dysfunction common; may follow contact with monkey.
Synonyms	Durba syndrome, Green monkey disease, Marburg. ICD9: 078.89 ICD10: A98.3

Clinical

WHO Case definition for surveillance: ¹

Clinical description

- Ebola hemorrhagic fever and Marburg virus disease both begin with acute fever, diarrhea that can be bloody (referred to as "diarrhee rouge" in francophone Africa), and vomiting.
- Headache, nausea, and abdominal pain are common.
- Conjunctival injection, dysphagia, and hemorrhagic symptoms such as nosebleeds, bleeding gums, vomiting of blood, blood in stools, purpura may further develop.
- Some patients may also show a maculopapular rash on the trunk.
- Dehydration and significant wasting occur as the disease progresses.
- At a later stage, there is frequent involvement of the central nervous system, manifested by somnolence, delirium, or coma.
- The case-fatality rate ranges from 50% to 90%.

Laboratory criteria for diagnosis

- Supportive: Positive serology (ELISA for IgG and/or IgM), or
- Confirmatory: Positive virus isolation (only in a laboratory of biosafety level 4) or
- Positive skin biopsy (immunohistochemistry) or
- Positive PCR

Case classification

Suspected: A case that is compatible with the clinical description.

Probable: in epidemic situation:

- Any person having had contact with a clinical case and presenting with acute fever, or
- Any person presenting with acute fever and 3 of the following symptoms: headache, vomiting / nausea, loss of appetite, diarrhea, intense fatigue, abdominal pain, general or articular pain, difficulty in swallowing, difficulty in breathing, hiccoughs, or
- Any unexplained death

Confirmed: Any suspected or probable case that is laboratory-confirmed.

Contact: in epidemic situation:

- An asymptomatic person having had physical contact within the past 21 days with a confirmed or probable case or his/her body fluids (e.g., care for patient, participation in burial ceremony, handling of potentially infected laboratory specimens).
- In epidemic situations and after laboratory confirmation of a few initial cases, there is no need for individual laboratory confirmation and the use of .suspected or probable. case classifications is sufficient for surveillance and control purposes.

The symptoms and signs of Marburg and Ebola virus infections are similar. ^{2 3}

Following an incubation period of 4 to 16 days, onset is sudden, marked by anorexia, fever, chills, headache and myalgia.

- After approximately three days, the patient develops nausea, vomiting, sore throat, abdominal pain and diarrhea.
- Patients are dehydrated, apathetic and disoriented and exhibit pharyngeal and conjunctival injection.
- A non-specific rash may be visible. ⁴

- Most develop severe hemorrhagic manifestations between days 5 and 7.
- Bleeding is often from multiple sites, most commonly from the gastrointestinal tract, lungs and gingivae.
- Hemorrhage and oropharyngeal lesions carry a particularly poor prognosis.
- Death occurs between days 7 and 16.

This disease is endemic or potentially endemic to 11 countries.

Marburg virus disease in Uganda

Prevalence surveys:

5.1% of Egyptian fruit bats (*Rousettus aegyptiacus*) in a cave associated with human infection (2007) ⁵

In 2008, a Dutch woman acquired fatal Marburg virus infection while visiting a cave in the Maramagambo Forest (between Queen Elisabeth Park and Kabale). ⁶⁻⁹

- In 2008, an American woman acquired Marburg infection in the same cave. ¹⁰⁻¹⁴

Prevalence surveys:

2.5% of *Rousettus aegyptiacus* bats in Python Cave (2007 to 2008) ^{15 16}

Notable outbreaks:

2007 - An outbreak (5 cases suspected, 4 confirmed, 1 fatal) was reported among gold miners in Kamwenge. ¹⁷⁻²⁶

2012 - An outbreak (20 cases, 9 fatal) was reported in Kabale District, Kampala, Ibanda, Mbarara, and Kabarole. ²⁷⁻³⁹

References

1. *New Microbiol* 2009 Oct ;32(4):359-67.
2. *Clin Infect Dis* 2001 Nov 15;33(10):1707-12.
3. *Rev Infect Dis* 1989 May-Jun;11 Suppl 4:S777-82.
4. *Int J Dermatol* 2012 Sep ;51(9):1037-43.
5. ProMED <promedmail.org> archive: 20090801.2700
6. *Emerg Infect Dis* 2009 Aug ;15(8):1171-5.
7. *Travel Med Infect Dis* 2011 May ;9(3):126-34.
8. ProMED <promedmail.org> archive: 20080710.2107
9. ProMED <promedmail.org> archive: 20080711.2115
10. *MMWR Morb Mortal Wkly Rep* 2009 Dec 18;58(49):1377-81.
11. *Travel Med Infect Dis* 2011 May ;9(3):126-34.
12. *J Infect Dis* 2011 Nov ;204 Suppl 3:S796-9.
13. ProMED <promedmail.org> archive: 20090131.0423
14. ProMED <promedmail.org> archive: 20091218.4276
15. *PLoS Pathog* 2012 Oct ;8(10):e1002877.
16. ProMED <promedmail.org> archive: 20121029.1374103
17. *Wkly Epidemiol Rec* 2007 Oct 26;82(43):381-4.
18. *Wkly Epidemiol Rec* 2007 Aug 17;82(33):297-8.
19. *Wkly Epidemiol Rec* 2007 Sep 7;82(36):317.
20. ProMED <promedmail.org> archive: 20070801.2490
21. ProMED <promedmail.org> archive: 20070801.2475
22. ProMED <promedmail.org> archive: 20070803.2514
23. ProMED <promedmail.org> archive: 20070804.2533
24. ProMED <promedmail.org> archive: 20070810.2609
25. ProMED <promedmail.org> archive: 20070814.2656
26. ProMED <promedmail.org> archive: 20071002.3257
27. *Wkly Epidemiol Rec* 2012 Oct 26;87(43):414.
28. *Wkly Epidemiol Rec* 2012 Nov 9;87(45):437-8.
29. *Lancet* 2012 Nov 17;380(9855):1726.
30. ProMED <promedmail.org> archive: 20121021.1356423
31. ProMED <promedmail.org> archive: 20121020.1355141
32. ProMED <promedmail.org> archive: 20121019.1352925
33. ProMED <promedmail.org> archive: 20121023.1359519
34. ProMED <promedmail.org> archive: 20121026.1366477
35. ProMED <promedmail.org> archive: 20121029.1372569
36. ProMED <promedmail.org> archive: 20121031.1382300
37. ProMED <promedmail.org> archive: 20121105.1392660
38. ProMED <promedmail.org> archive: 20121110.1403820
39. ProMED <promedmail.org> archive: 20121123.1422035

Measles

Agent	VIRUS - RNA. Paramyxoviridae, Paramyxovirinae, Morbillivirus: Measles virus
Reservoir	Human
Vector	None
Vehicle	Droplet
Incubation Period	8d - 14d
Diagnostic Tests	Viral culture (difficult and rarely indicated). Serology. Nucleic acid amplification.
Typical Adult Therapy	Respiratory isolation; supportive. Ribavirin 20 to 35 mg/kg/day X 7 days has been used for severe adult infection
Typical Pediatric Therapy	As for adult
Vaccines	Measles Measles-Mumps-Rubella Measles-Rubella
Clinical Hints	Coryza, fever, headache, conjunctivitis, photophobia and a maculopapular rash after 3 to 5 days; Koplik's spots (bluish-grey lesions on buccal mucosa, opposite second molars) often precede rash; encephalitis or viral pneumonia occasionally encountered.
Synonyms	Masern, Massling, Mazelen, Meslinger, Morbilli, Morbillo, Rubeola, Rugeole, Sarampion, Sarampo. ICD9: 055 ICD10: B05

Clinical

WHO Case definition for surveillance:

Any person with:

- fever, and
- maculopapular (i.e. non-vesicular) rash, and
- cough, coryza (i.e. runny nose) or conjunctivitis (i.e. red eyes).

or

Any person in whom a clinician suspects measles infection.

Laboratory criteria for diagnosis

- At least a fourfold increase in antibody titer or
- Isolation of measles virus or
- Presence of measles-specific IgM antibodies

Case classification

- Clinically confirmed: A case that meets the clinical case definition.
- Probable: Not applicable.
- Laboratory-confirmed: only for outbreak confirmation and during elimination phase A case that meets the clinical case definition and that is laboratory-confirmed or linked epidemiologically to a laboratory-confirmed case.

Acute illness:

Symptoms begin to appear about 10 to 12 days after exposure to the virus, with fever followed by cough, rhinorrhea, and/or conjunctivitis. ¹

- The rash appears approximately 14 days after exposure and lasts 5 to 6 days.
- The rash begins at the hairline, spreading to the face and neck.
- Over the next three days, the rash gradually extends, eventually reaching the hands and feet. ²

Complications:

Complications of measles include diarrhea, otitis media (10%), pneumonia (5%), encephalitis (0.1%) ^{3 4}, arthropathy (28%) ⁵, seizures, and death. ⁶

- Twenty percent of patients experience one or more complications, most often children below five years of age and adults over 20.
- Measles in pregnancy may be associated with maternal pneumonia, abortion, low birth weight ^{7 8} or congenital infection of the newborn. ^{9 10}
- In developing countries, measles has been known to kill as many as one out of four people.

- Measles is the leading cause of blindness among African children, as a result of concomitant vitamin A deficiency.
- Measles pneumonia accounts for approximately 17% of bronchiolitis obliterans in children (Beijing, 2001 to 2007) ¹¹
- Rare instances of thyroiditis, pancreatitis and sialoadenitis have been reported. ¹²

This disease is endemic or potentially endemic to all countries.

Measles in Uganda

Vaccine Schedule:

BCG - birth

DTwP - 6, 10, 14 weeks


DTwPHibHep - 6, 10, 14 weeks

Measles (monovalent) - 9 months


OPV - birth; 6, 10, 14 weeks

TT - Women of child-bearing age; +1, +6 months; +1, +1 year

Vitamin A - 6, 12, 18, 24, 30, 36, 42 months



Graph: Uganda. Measles - WHO-UNICEF est. % vaccine coverage



Graph: Uganda. Measles, cases

Notes:

1. Measles transmission in Uganda was interrupted following a vigorous vaccination campaign undertaken during 2002 to 2006. ¹³

Individual years:

1997 - 12,562 cases reported to WHO - and true incidence estimated at 110,909 cases.

1998 - 42,632 cases reported to WHO - and true incidence estimated at 74,458 cases.

1999 - 42,737 cases reported to WHO.

Notable outbreaks:

1990 to 1991 - An outbreak of measles was reported in Kampala. ^{14 15}

2011 - Outbreaks (3,312 cases) were reported. ¹⁶⁻¹⁸

2012 - Outbreaks (3,270 cases during January to March) were reported in multiple districts. ¹⁹⁻²¹

2013 - An outbreak (141 cases, 5 fatal) was reported in Western Uganda. ^{22 23}

References

1. J Infect Dis 2004 May 1;189 Suppl 1:S4-16.
2. Dermatol Clin 2002 Apr ;20(2):209-15, v.
3. Pediatr Neurol 2003 Apr ;28(4):281-4.
4. Semin Pediatr Neurol 2012 Sep ;19(3):107-14.
5. Clin Rheumatol 2009 Sep ;28(9):1067-71.
6. Lancet 2003 Mar 1;361(9359):763-73.
7. N Engl J Med 1966 Apr 7;274(14):768-71.
8. J Infect 2003 Jul ;47(1):40-4.
9. Ann Trop Paediatr 2011 ;31(2):185-8.
10. J Gynecol Obstet Biol Reprod (Paris) 2012 Mar 7;
11. Zhonghua Er Ke Za Zhi 2008 Oct ;46(10):732-8.
12. Histopathology 2000 Aug ;37(2):141-6.
13. Emerg Infect Dis 2011 Jan ;17(1):110-3.
14. Wkly Epidemiol Rec 1991 Dec 6;66(49):364-7.
15. Bull World Health Organ 1992 ;70(3):317-21.
16. ProMED <promedmail.org> archive: 20111211.3574
17. ProMED <promedmail.org> archive: 20111218.3627
18. ProMED <promedmail.org> archive: 20130119.1506143
19. ProMED <promedmail.org> archive: 20120402.1087489
20. ProMED <promedmail.org> archive: 20120422.1110279
21. ProMED <promedmail.org> archive: 20120527.1146773
22. ProMED <promedmail.org> archive: 20130112.1494531
23. ProMED <promedmail.org> archive: 20130216.1545849

Melioidosis

Agent	BACTERIUM. <i>Burkholderia pseudomallei</i> An aerobic gram-negative bacillus
Reservoir	Soil Water Sheep Goat Horse Pig Rodent Monkey Marsupial
Vector	None
Vehicle	Water: Contact, ingestion, aerosol Breast milk (rare)
Incubation Period	3d - 21d (range 2d - 1y)
Diagnostic Tests	Culture of blood, sputum, tissue. Serology. Nucleic acid amplification.
Typical Adult Therapy	Ceftazidime or Meropenem or Imipenem IV X at least 14 days May be combined with Sulfamethoxazole/trimethoprim PO Follow with Sulfamethoxazole/trimethoprim +/- Doxycycline X at least 3 months.
Typical Pediatric Therapy	Ceftazidime or Meropenem or Imipenem IV X at least 14 days May be combined with Sulfamethoxazole/trimethoprim PO Follow with Sulfamethoxazole/trimethoprim X at least 3 months.
Clinical Hints	May present as: lymphangitis with septicemia; or fever, cough and chest pain; or diarrhea; bone, central nervous system, liver and parotid infection are occasionally encountered; case-fatality rate 10% to over 50% (septicemic form).
Synonyms	<i>Burkholderia pseudomallei</i> , <i>Burkholderia thailandensis</i> , Melioidose, Nightcliff Gardeners' Disease, Whitmore disease. ICD9: 025 ICD10: A24.1,A24.2,A24.3,A24.4

Clinical

The clinical features of melioidosis are similar to those of tuberculosis: prolonged fever, weight loss, latency with reactivation, upper-lobe infiltrates, etc. ¹⁻⁵

- As in tuberculosis, long latent periods may precede appearance of the disease; in some reports 29 years ⁶, or even 69 years. ⁷
- Disease rates are highest among diabetics. ⁸⁻¹⁰ Other predisposing conditions include collagen-vascular disease, alcoholism, malnutrition, chronic renal or hepatic disease, corticosteroid therapy, splenectomy, pregnancy, chronic granulomatous disease, leukemia and lymphoma.

Acute melioidosis can be divided into five clinical forms:

- septicemia without abscess formation
- septicemia with disseminated foci
- localized infection
- transitory bacteremia
- "fever of unknown origin"

Most patients with overt infection present with pneumonia which may include pulmonary nodules, consolidation, necrotizing lesions, pleural effusion, pleural thickening and mediastinal abscesses. ¹¹⁻¹³

- Occasionally, the only lesion may be a pleural mass.
- Although confluent upper lobe infiltrates are common, the apices are generally spared in non-septicemic cases.
- Rapid progression and early cavitation are common.
- Pleural effusion is seen in 21% of patients with acute disease, and 13% of patients with chronic melioidosis
- Pericarditis occurs in six to ten percent of all patients.
- Patients with cystic fibrosis (ie, traveling to endemic countries) appear to be at high risk for pulmonary infection.
- The pattern of organ involvement in recurrent or relapsing melioidosis is similar to that of primary infection. ¹⁴

45% of cases present as septicemia with infection of multiple organs.

- Pericarditis ^{15 16} may complicate the pulmonary infection, and necessitate surgical drainage for tamponade.
- Visceral abscesses may involve the spleen ^{17 18}, liver ¹⁹⁻²¹, kidneys ²², pancreas ²³, prostate ²⁴ or other organs. ²⁵
- Osteomyelitis is common. ²⁶⁻³⁰
- Generalized or local suppurative lymphadenitis is occasionally encountered. ³¹
- Primary cutaneous diseases occurs in 12% of cases, and secondary cutaneous dissemination in 2% ^{32 33}
- Complications of melioidosis include nasopharyngitis, brain abscess ³⁴, septic arthritis ^{35 36}, dural sinus thrombosis ³⁷, brainstem dysfunction ³⁸, orbital infection ^{39 40}, meningitis, transverse myelitis ⁴¹, urinary tract infection ⁴²,

epididymo-orchitis, prostatitis ^{43 44}, suppurative parotitis ⁴⁵, mycotic aneurysm ⁴⁶, parapharyngeal abscess, corneal ulcer, necrotizing fasciitis ⁴⁷⁻⁴⁹, septic arthritis ^{50 51}, psoas and other muscular abscesses. ⁵²⁻⁵⁴

- Persistent bacteremia beyond one week under therapy is associated with poor prognosis. ⁵⁵
- Melioidosis is the most common cause of mycotic aneurysm in some areas of Thailand. ⁵⁶

Renal failure occurs in up to one-third of hospitalized patients with melioidosis, and carries a poor prognosis.

In nonendemic regions, patients present with reactivated disease occurring months to years after initial exposure to the organism.

- Typical symptoms include fever, cough, weight loss and apical changes on chest x-ray • all suggestive of tuberculosis.
- The presence of a mediastinal mass may mistakenly suggest malignancy. ⁵⁷
- The clinical features of melioidosis may also mimic those of enteric fever. ⁵⁸
- It is not uncommon for the two diseases to coexist.

This disease is endemic or potentially endemic to 73 countries.

References

1. Curr Opin Infect Dis 2004 Apr ;17(2):131-6.
2. Eur Respir J 2003 Sep ;22(3):542-50.
3. Int J Tuberc Lung Dis 2008 Oct ;12(10):1209-15.
4. Br J Radiol 2009 Jun ;82(978):514-21.
5. J Thorac Imaging 2012 Jul 9;
6. Hosp Pract (Minneapolis) 1997 May 15;32(5):219-21.
7. J Clin Microbiol 2005 Feb ;43(2):970-2.
8. Infect Immun 2012 Apr 2;
9. Clin Infect Dis 2012 Feb 1;54(3):362-9.
10. PLoS Negl Trop Dis 2010 ;4(11):e900.
11. Clin Microbiol Rev 2005 Apr ;18(2):383-416.
12. Clin Infect Dis 2012 Feb 1;54(3):362-9.
13. Curr Probl Diagn Radiol 2012 Nov-Dec;41(6):199-209.
14. Am J Trop Med Hyg 2009 Aug ;81(2):335-7.
15. Scand J Infect Dis 2007 ;39(4):357-9.
16. Clin Infect Dis 2010 Sep 1;51(5):e46-9.
17. Ann Acad Med Singapore 2008 Sep ;37(9):749-52.
18. J Med Assoc Thai 2009 Nov ;92(11):1476-84.
19. Indian J Med Microbiol 2007 Apr ;25(2):150-1.
20. J Med Assoc Thai 2010 Jul ;93(7):838-48.
21. J Med Imaging Radiat Oncol 2011 Apr ;55(2):176-82.
22. Nephrology (Carlton) 2012 Dec 21;
23. JOP 2010 ;11(4):365-8.
24. Singapore Med J 2009 Apr ;50(4):385-9.
25. Trans R Soc Trop Med Hyg 2012 Oct ;106(10):629-31.
26. Emerg Infect Dis 2007 Aug ;13(8):1257-9.
27. Indian J Orthop 2010 Apr ;44(2):216-20.
28. Semin Musculoskelet Radiol 2011 Nov ;15(5):480-8.
29. Australas Med J 2012 ;5(2):141-3.
30. J Foot Ankle Surg 2013 Feb 7;
31. Trans R Soc Trop Med Hyg 2006 Aug ;100(8):798-801.
32. Clin Infect Dis 2008 Sep 1;47(5):603-9.
33. Emerg Infect Dis 2012 Feb ;18(2):359-60.
34. J Neurosurg 2008 Feb ;108(2):243-7.
35. J Clin Rheumatol 2001 Aug ;7(4):242-7.
36. J Assoc Physicians India 2012 Jun ;60:44-5.
37. J Med Assoc Thai 2006 Feb ;89(2):242-7.
38. J Emerg Med 2012 Apr 9;
39. J Clin Microbiol 2007 Dec ;45(12):4073-4.
40. Ophthal Plast Reconstr Surg 2013 Jan 8;
41. BMC Infect Dis 2012 Sep 28;12(1):232.
42. J Postgrad Med 2007 Apr-Jun;53(2):108-10.
43. Clin Rheumatol 2008 Dec ;27 Suppl 2:S59-61.
44. Nat Clin Pract Urol 2007 Feb ;4(2):111-4.
45. Indian Pediatr 2010 Sep ;47(9):799-801.
46. Clin Microbiol Infect 2011 May ;17(5):719-21.
47. Emerg Infect Dis 2003 Nov ;9(11):1484-5.
48. Jpn J Infect Dis 2008 Mar ;61(2):151-3.
49. Southeast Asian J Trop Med Public Health 2008 Jul ;39(4):656-8.
50. J Microbiol Immunol Infect 2007 Apr ;40(2):178-82.
51. Clin Rheumatol 2008 Dec ;27 Suppl 2:S59-61.
52. Lancet 2003 May 17;361(9370):1715-22.
53. Southeast Asian J Trop Med Public Health 2008 Jul ;39(4):649-55.
54. J Trop Pediatr 2011 Jun ;57(3):185-91.
55. Am J Trop Med Hyg 2011 Jun ;84(6):858-61.
56. Clin Infect Dis 2008 Dec 1;47(11):1436-9.
57. J Med Case Rep 2012 ;6(1):28.
58. Trans R Soc Trop Med Hyg 2008 Dec ;102 Suppl 1:S117-8.

Meningitis - aseptic (viral)

Agent	VIRUS - RNA. Picornaviridae, enteroviruses
Reservoir	Human
Vector	None
Vehicle	Fecal-oral Droplet
Incubation Period	Variable
Diagnostic Tests	Viral isolation (stool, CSF, throat). Serology.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Lymphocytic meningitis (normal CSF glucose); often follows sore throat; typically occurs during late summer and early autumn in temperate regions.
Synonyms	Aseptic meningitis, Encephalitis - viral, Meningite virale, Meningitis, viral, Meningo-encefalite virale, Viral encephalitis, Viral meningitis. ICD9: 047,048,049,320.2 ICD10: A87,G03.0

Clinical

WHO Case definition for surveillance:

Clinical case definition

A case with fever 38.5°C and one or more of the following:

- neck stiffness
 - severe unexplained headache
 - neck pain and 2 or more of the following: photophobia, nausea, vomiting, abdominal pain, pharyngitis with exudates
- For children <2 years of age a case is defined as
- A case with fever 38.5°C and one or more of the following: irritability, bulging fontanelle

Laboratory criteria for confirmation

- The specific virus confirmed on cell culture.

Case classification

Suspected: A case that meets the clinical case definition and one or more of the following:

- normal CSF glucose and normal or mild increase in CSF protein (>50 mg/dl), moderate increase CSF cells (<500/mm³) and lymphocyte predominance (>50%)
- CSF Positive for viral genomic sequences using PCR (Polymerase Chain Reaction)
- Epidemiological link to a confirmed case

Confirmed: A suspected or probable case with laboratory confirmation.

As a group, the viral meningitides are characterized by fever, headache, meningismus and lymphocytic pleocytosis. ^{1 2}

- Major complications and sequelae are unusual. ^{3 4}
- The cerebrospinal fluid glucose level is normal, and a transitory neutrophilic pleocytosis is occasionally encountered.
- CSF pleocytosis is often absent among children with enteroviral meningitis. ^{5 6}

This disease is endemic or potentially endemic to all countries.

References

1. Semin Pediatr Infect Dis 2002 Jan ;13(1):40-7.
2. Semin Neurol 2000 ;20(3):277-92.
3. CMAJ 2003 May 27;168(11):1421-3.
4. J Clin Microbiol 2003 May ;41(5):2230-2.
5. Pediatr Emerg Care 2010 Feb ;26(2):77-81.
6. Arch Dis Child 2012 Oct ;97(10):874-8.

Meningitis - bacterial

Agent	BACTERIUM. Neisseria meningitidis , Streptococcus pneumoniae , Haemophilus influenzae , et al
Reservoir	Human
Vector	None
Vehicle	Air Infected secretions
Incubation Period	Variable
Diagnostic Tests	CSF microscopy and culture. Blood culture. Note: Antigen detection is non-specific and rarely useful.
Typical Adult Therapy	Bactericidal agent(s) appropriate to known or suspected pathogen + dexamethasone
Typical Pediatric Therapy	As for adult
Vaccines	H. influenzae (HbOC-DTP or -DTaP) Haemophilus influenzae (HbOC) Haemophilus influenzae (PRP-D) Haemophilus influenzae (PRP-OMP) Haemophilus influenzae (PRP-T) Meningococcal Hepatitis B + Haemoph. influenzae
Clinical Hints	Headache, stiff neck, obtundation, high fever and leukocytosis; macular or petechial rash and preceding sore throat suggest meningococcal infection.
Synonyms	Bacterial meningitis, Enfermedad Meningococica, Haemophilus influenzae, Haemophilus influenzaes, HIB meningitis, HIBs, Infections a meningocoque, Meningite batterica, Meningite meningococcica, Meningococcal, Meningokokken Erkr., Meningokokkose. ICD9: 036.0,320 ICD10: A39,G00,G01,G02

Clinical

WHO Case definition for surveillance of Meningococcal infection:

Clinical case definition

- An illness with sudden onset of fever (>38.5°C rectal or >38.0°C axillary) and one or more of the following:
- Neck stiffness
- Altered consciousness
- Other meningeal sign or petechial or purpuric rash
- In patients <1 year, suspect meningitis when fever accompanied by bulging fontanelle.

Laboratory criteria for diagnosis

- Positive CSF antigen detection or
- Positive culture

Case classification

- Suspected: A case that meets the clinical case definition.
- Probable: A suspected case as defined above and turbid CSF (with or without positive Gram stain) or ongoing epidemic and epidemiological link to a confirmed case
- Confirmed: A suspected or probable case with laboratory confirmation.

WHO Case definition for surveillance of Haemophilus influenzae type b (Hib disease):

Clinical description

- Bacterial meningitis is characterized by fever of acute onset, headache and stiff neck.
- Meningitis is not a specific sign for Hib disease, and Hib disease cannot be diagnosed on clinical grounds.

Laboratory criteria for diagnosis

- Culture: isolation of Hib from a normally sterile clinical specimen, such as cerebrospinal fluid (CSF) or blood.
- Culture of Hib from non-sterile sites such as the throat, where bacteria can grow without causing disease, does not define Hib disease.
- Antigen detection: identification of Hib antigen in normally sterile fluids, by methods such as latex agglutination or counter-immunoelectrophoresis (CIE).

Case classification

- Potential: (bacterial meningitis case): a child with a clinical syndrome consistent with bacterial meningitis.
- Probable: Not applicable.
- Confirmed: A case that is laboratory-confirmed (growth or identification of Hib in CSF or blood).


Note: Any person with Hib isolated from CSF or blood may be reported as a confirmed case, regardless of whether their clinical syndrome was meningitis.

As a group, the bacterial meningitides are characterized by signs of sepsis, fever, headache, meningismus and neutrophilic pleocytosis. ^{1 2}


- 33% to 69% of patients with meningococcal infection have hyperglycemia on admission ^{3 4}
- 7.5% of patients with meningococcal infection present with arthritis. ⁵
- Major complications and sequelae are common.
- Delayed cerebral thrombosis is encountered in 1.1% of cases. ⁶

This disease is endemic or potentially endemic to all countries.

Meningitis - bacterial in Uganda



Graph: Uganda. Meningitis - meningococcal, cases




Graph: Uganda. Meningococcal infection, cases

Notes:

Individual years:

2006 - 728 cases (37 fatal) were reported during weeks 1 to 15. ⁷




Graph: Uganda. Meningococcal infection, deaths

In one hospital series, *Streptococcus pneumoniae* accounted for 41% of meningitis cases among children below age 2 years, *Salmonella* species 31% and *H. influenzae* 9% (2003 to 2008) ⁸


Vaccine Schedule:

BCG - birth
 DTwP - 6, 10, 14 weeks
 DTwPHibHep - 6, 10, 14 weeks
 Measles (monovalent) - 9 months
 OPV - birth; 6, 10, 14 weeks
 TT - Women of child-bearing age; +1, +6 months; +1, +1 year
 Vitamin A - 6, 12, 18, 24, 30, 36, 42 months

Routine use of *Haemophilus influenzae* vaccine was introduced in 2002.⁹



Graph: Uganda. Hib3 - WHO-UNICEF est. % vaccine coverage



© 2013 - GIDEON Informatics Inc - www.gideononline.com

Graph: Uganda. *Haemophilus influenzae* meningitis, cases**Notable outbreaks:**

1994 to 1995 - An outbreak (291 cases, 13.3% fatal) of group A meningococcal infection was reported in a refugee camp in northern Uganda. ¹⁰

2006 - An outbreak (515 cases, 44 fatal) of group A meningococcal infection was reported in Nakapiripirit and Moroto. ¹¹⁻¹⁶

2007 - An outbreak (2,961 cases, 105 fatal) of meningococcal infection was reported in the northwestern region (Karamoja and West Nile States). ¹⁷⁻²⁴ An additional 147 cases were reported in the northeast. ²⁵

2008 to 2009 - Outbreaks (336 cases, 42 fatal) were reported in Masindi (14 cases, 4 fatal) ²⁶, Hoima (47 cases, 13 fatal) and Arua (150 cases, 18 fatal). ²⁷⁻²⁹

References

1. Semin Neurol 2000 ;20(3):293-306.
2. Infect Dis Clin North Am 2001 Dec ;15(4):1047-71.
3. BMC Infect Dis 2009 ;9:57.
4. Crit Care 2011 ;15(1):R44.
5. Eur J Clin Microbiol Infect Dis 2012 Apr 3;
6. Intensive Care Med 2012 Dec 18;
7. Wkly Epidemiol Rec 2007 Feb 2;82(5):34-40.
8. J Paediatr Child Health 2012 Apr ;48(4):324-8.
9. MMWR Morb Mortal Wkly Rep 2008 Feb 15;57(6):148-51.
10. Epidemiol Infect 2000 Feb ;124(1):75-81.
11. Wkly Epidemiol Rec 2006 Mar 31;81(13):119-20.
12. Wkly Epidemiol Rec 2006 Feb 17;81(7):62.
13. Wkly Epidemiol Rec 2006 Feb 3;81(5):43.
14. ProMED <promedmail.org> archive: 20060121.0207
15. ProMED <promedmail.org> archive: 20060215.0500
16. ProMED <promedmail.org> archive: 20060321.0880
17. Wkly Epidemiol Rec 2007 Feb 2;82(5):33.
18. ProMED <promedmail.org> archive: 20070123.0300
19. ProMED <promedmail.org> archive: 20070125.0339
20. ProMED <promedmail.org> archive: 20070203.0438
21. ProMED <promedmail.org> archive: 20070207.0484
22. ProMED <promedmail.org> archive: 20070213.0538
23. ProMED <promedmail.org> archive: 20070222.0654
24. ProMED <promedmail.org> archive: 20070316.0934
25. ProMED <promedmail.org> archive: 20070302.0740
26. ProMED <promedmail.org> archive: 20090120.0249
27. ProMED <promedmail.org> archive: 20090115.0170
28. ProMED <promedmail.org> archive: 20090123.0304
29. ProMED <promedmail.org> archive: 20090220.0716

Microsporidiosis

Agent	PARASITE - Protozoa. Microspora: Enterocytozoon, Encephalitozoon (Septata), Vittaforma (Nosema), Pleistophora, Trachipleistophora, et al.
Reservoir	Rabbit Rodent Carnivore Non-human primate Fish Dog Bird
Vector	None
Vehicle	? Fecal-oral
Incubation Period	Unknown
Diagnostic Tests	Microscopy of duodenal aspirates. Inform laboratory if this organism is suspected. Nucleic acid amplification.
Typical Adult Therapy	Albendazole 400 mg PO BID X 3 weeks. Add Fumagillin for ocular <i>S. intestinalis</i> may respond to Albendazole and Fumagillin Nitazoxanide has been used for <i>E. bienersi</i> .
Typical Pediatric Therapy	Albendazole 200 mg PO BID X 3 weeks. Add Fumagillin for ocular <i>S. intestinalis</i> may respond to Albendazole and Fumagillin Nitazoxanide has been used for <i>E. bienersi</i> .
Clinical Hints	In AIDS patients, infection is characterized by chronic diarrhea, wasting and bilateral keratoconjunctivitis; hepatitis and myositis may be present.
Synonyms	Brachiola, Encephalitozoon, Enterocytozoon, Microsporidium, Nosema, Pleistophora, Trachipleistophora, Tubulinosema, Vittaforma. ICD9: 136.8 ICD10: A07.8

Clinical

Intestinal disease in immunocompetent patients is characterized by self-limited diarrhea, traveler's diarrhea or asymptomatic carriage. ¹

- Immunocompromised patients present with diarrhea, cholangitis, cholecystitis, sinusitis or pneumonia. ^{2 3}

Ocular microsporidiosis is associated with keratoconjunctivitis.

Other syndromes include sinusitis, nephritis, myositis and prostatitis. ⁴

This disease is endemic or potentially endemic to all countries.

Microsporidiosis in Uganda

Prevalence surveys:

17.4% of children ages 0 to 60 months with diarrhea, and 16.8% of asymptomatic controls (*Enterocytozoon bienersi*, Kampala, 1999 to 2001) ⁵

30.4% of malnourished children hospitalized with diarrhea (Kampala, 2002 to 2003)

76.9% of HIV-positive and 6.6% of HIV-negative children with persistent diarrhea (Kampala, 2002 to 2003) ^{6 7}

Encephalitozoon intestinalis has been identified in gorillas and persons sharing their habitats. ⁸

References

1. Curr Opin Infect Dis 2006 Oct ;19(5):485-92.
2. Clin Infect Dis 1994 May ;18(5):819-25.
3. Clin Microbiol Rev 1994 Oct ;7(4):426-61.
4. Clin Microbiol Rev 2005 Jul ;18(3):423-45.

5. Am J Trop Med Hyg 2005 Nov ;73(5):921-5.
6. Am J Trop Med Hyg 2005 Nov ;73(5):921-5.
7. Emerg Infect Dis 2009 Jan ;15(1):49-52.
8. Parasitol Res 2002 Oct ;88(10):926-31.

Moniliformis and Macracanthorhynchus

Agent	PARASITE - Archiacanthocephala. Moniliformida: Moniliformis moniliformis, Oligocanthorhynchida: Maracanthorhynchus hirudinaceus.
Reservoir	Pig (Maracanthorhynchus), rat and fox (Moniliformis),
Vector	None
Vehicle	Insect (ingestion)
Incubation Period	Unknown - presumed 15 to 40 days
Diagnostic Tests	Identification of worm in stool.
Typical Adult Therapy	Infection is usually self-limited. Pyrantel pamoate has been used against Moniliformis moniliformis - 11 mg/kg PO - repeat once in 2 weeks
Typical Pediatric Therapy	Infection is usually self-limited. Pyrantel pamoate has been used against Moniliformis moniliformis - 11 mg/kg PO - repeat once in 2 weeks
Clinical Hints	Most infections are characterized by asymptomatic passage of a worm; however, vague complaints such as 'periumbilical discomfort' and 'giddiness' have been described.
Synonyms	Acanthocephalan worms, Macracanthorhynchus, Moniliform acanthocephalan, Moniliformis moniliformis. ICD9: 128.9 ICD10: B83.8

Clinical

Most infections are characterized by asymptomatic passage of a worm; however, vague complaints such as 'periumbilical discomfort' and 'giddiness' have been described. ¹

- In one instance, a man developed marked abdominal pain following experimental self-infection.
- In another case, intestinal perforation was associated with *Macracanthorhynchus hirudinaceus* infestation. ²

This disease is endemic or potentially endemic to all countries.

References

1. J Helminthol 1992 Sep ;66(3):241-2.
2. Trop Med Parasitol 1989 Dec ;40(4):476-7.

Mumps

Agent	VIRUS - RNA. Paramyxoviridae, Paramyxovirinae, Rubulavirus: Mumps virus
Reservoir	Human
Vector	None
Vehicle	Aerosol
Incubation Period	14d - 24d (range 12d - 24d)
Diagnostic Tests	Viral culture (saliva, urine, CSF) indicated only in complicated cases. Serology. Nucleic acid amplification.
Typical Adult Therapy	Respiratory isolation; supportive
Typical Pediatric Therapy	As for adult
Vaccines	Measles-Mumps-Rubella Mumps Rubella - Mumps
Clinical Hints	Fever, parotitis, orchitis (20% of post-pubertal males), meningitis (clinically apparent in 1% to 10%), oophoritis, or encephalitis (0.1%); most cases resolve within 1 to 2 weeks.
Synonyms	Bof, Epidemic parotitis, Fiebre urliana, Infectious parotitis, Kusma, Oreillons, Paperas, Parotidite epidemica, Parotiditis, Parotite epidemica, Passjuka. ICD9: 072 ICD10: B26

Clinical

One third of Mumps virus infections are asymptomatic.

Acute illness:

The prodrome of mumps consists of low-grade fever, anorexia, malaise, and headache.

- Usually within one day, the patient complains "earache" and tenderness is noted over the parotid gland. ¹
- The gland is soon visibly enlarged and progresses to maximum size over the next 2 to 3 days, often with lifting of the ear lobe upward and outward.
- The orifice of Stensen's duct is edematous and erythematous, and trismus and pain on chewing may be present.
- It is important to remember that the enlarged gland obscures the angle of the mandible, while cervical adenopathy does not.
- Parotid involvement is unilateral in 25% of cases.
- As the disease progresses, fever may reach 40C.
- Subsequently pain, fever, and tenderness resolve, and the parotid gland returns to normal size within a week.
- Involvement of the other salivary glands occurs in 10% of cases, but are rare in the absence of parotid involvement.
- Presternal edema develops in 6% of patients, most often in those who have submandibular adenitis.

8% to 15% of patients will continue shedding Mumps virus 5 days after the onset of symptoms. ²

Neurological manifestations:

Central nervous system involvement is the most common extrasalivary gland manifestation of this disease.

- Cerebrospinal fluid pleocytosis has been documented in 51% patients with mumps, without other evidence of meningitis.
- Clinical meningitis occurs in 1 to 10% of persons with mumps parotitis; while parotitis is documented in less than 50% of patients with mumps.
- Meningitis may occur before, during or after salivary gland involvement.
- The features of mumps meningitis are similar to those of other viruses, and the clinical course is benign; however, polymorphonuclear CSF pleocytosis and reduced glucose levels are not unusual.

Encephalitis occurs in less than 0.1% of cases, and may be accompanied by altered consciousness, seizures, paresis, aphasia, involuntary movements; and sequelae such as psychomotor retardation, deafness (1 per 1,000 to 20,000 cases ³) ⁴ and convulsive disorders.

- Other neurological complications of mumps include cerebellar ataxia ⁵, facial nerve palsy, transverse myelitis, Guillain-Barre syndrome, and aqueductal stenosis.

Epididymo-orchitis:

Epididymo-orchitis is the most common extra-salivary gland manifestation in adults, developing in 20 to 30% of infected postpubertal males.

- This complication is bilateral in 15% of cases, and appears during the first week of mumps in 70% of cases.
- Rarely, this is the only manifestation of mumps.
- Onset is abrupt, with elevation of fever, chills, headache, vomiting, and testicular pain.
- The testis is warm, swollen (to as much as four times normal size), and tender, with erythema of the scrotum.
- Epididymitis is present in 85%, and usually precedes the orchitis.
- Tenderness may persist for more than 2 weeks in 20% of cases; and some degree of atrophy is noted in 50% of the patients, even after 2 years.
- Impotence is not encountered, and sterility is rare.

Additional manifestations of mumps:

Other features of mumps include oophoritis, fetal wastage ⁶, migratory polyarthritits, monoarticular arthritis and arthralgia, electrocardiographic changes (with or without overt myocarditis), nephritis, thyroiditis, mastitis, prostatitis, hepatitis, cholecystitis and thrombocytopenia.

This disease is endemic or potentially endemic to all countries.

References

1. Lancet 2008 Mar 15;371(9616):932-44.
2. Clin Infect Dis 2008 May 1;46(9):1447-9.
3. Pediatr Infect Dis J 2009 Mar ;28(3):173-5.
4. Laryngol Rhinol Otol (Stuttg) 1977 Apr ;56(4):342-5.
5. Am J Dis Child 1992 Aug ;146(8):930-1.
6. N Engl J Med 1966 Apr 7;274(14):768-71.

Myalgic encephalomyelitis

Agent	UNKNOWN
Reservoir	Unknown
Vector	None
Vehicle	Unknown
Incubation Period	Unknown
Diagnostic Tests	Clinical diagnosis; ie, discount other diseases.
Typical Adult Therapy	Supportive; ? immune modulators (experimental)
Typical Pediatric Therapy	As for adult
Clinical Hints	Unexplained depression, fatigue, cognitive disorders, sleep disturbance, recurrent bouts of pharyngitis and adenopathy, rheumatological symptoms and fever lasting more than six months.
Synonyms	Chronic fatigue syndrome. ICD9: 780.71 ICD10: G93.3

Clinical

The CDC (The United States Centers for Disease Control) consensus definition of Chronic Fatigue Syndrome (currently Myalgic encephalomyelitis ¹) requires the presence of two major criteria, in addition to at least six symptom criteria and at least two physical criteria (or the presence of eight symptom criteria, without need for physical criteria) as follows: ²⁻⁶

Major criteria:

- A. New onset of persistent or relapsing, debilitating fatigue or fatigability without a history of similar illness. Fatigue does not resolve with bed rest, and reduces daily activity by at least 50% for at least 6 months.
- B. Exclusion of other disorders through history, physical examination and laboratory studies.

Minor criteria:

A. Symptoms.

1. Mild fever or chills
2. Sore throat
3. Painful cervical or axillary adenopathy
4. Myalgia
5. Muscle weakness
6. Migratory arthralgia
7. Prolonged fatigue not meeting major criteria
8. Generalized headaches
9. Neuropsychological complaints (photophobia), scotomata, forgetfulness, irritability, confusion, problems in thinking or concentration ⁷⁻⁹ , depression)
10. Sleep disturbances
11. Description of the initial symptom complex as developing over a period of hours to days.

B. Physical criteria.

1. Low grade fever
2. Nonexudative pharyngitis
3. Cervical or axillary lymphadenopathy (nodes may be tender, and are usually no larger than 2 cm).

Affected children present with low levels of school attendance, fatigue, anxiety, functional disability and pain. ¹⁰

- Three phenotypes of Chronic Fatigue Syndrome are described in children: musculoskeletal, migraine and "sore throat." ¹¹

Patients with disease onset above age 50 years present with relatively high rates of fatigue, depression and autonomic dysfunction. ¹²

Additional findings described in Chronic fatigue syndrome have included generalized hyperalgesia ^{13 14} , impaired cardiac function ¹⁵ , migraine headache ¹⁶ and postural orthostatic tachycardia. ^{17 18}

In one series, 33% of patients referred to an Infectious Diseases clinic for suspected Lyme disease were found to have Chronic Fatigue Syndrome, and only 23% Lyme disease. ¹⁹

This disease is endemic or potentially endemic to all countries.

References

1. J Intern Med 2011 Oct ;270(4):327-38.
2. Occup Med (Lond) 2005 Jan ;55(1):13-9.
3. Clin Evid 2003 Dec ;(10):1289-303.
4. Lancet 2006 Jan 28;367(9507):346-55.
5. BMC Med 2009 ;7:57.
6. J Eval Clin Pract 2012 Feb ;18(1):25-31.
7. Psychol Med 2010 Aug ;40(8):1253-67.
8. J Psychosom Res 2010 May ;68(5):489-94.
9. Clin Neurol Neurosurg 2011 May ;113(4):295-302.
10. Arch Dis Child 2008 May ;93(5):419-21.
11. Arch Dis Child 2010 Apr ;95(4):245-9.
12. Eur J Clin Invest 2013 Mar ;43(3):302-8.
13. Clin Rheumatol 2010 Apr ;29(4):393-8.
14. Eur J Clin Invest 2012 Feb ;42(2):203-12.
15. J Intern Med 2011 Jul 27;
16. BMC Neurol 2011 ;11:30.
17. QJM 2008 Dec ;101(12):961-5.
18. J Intern Med 2012 Dec 4;
19. QJM 2012 Feb 1;

Mycetoma

Agent	BACTERIUM OR FUNGUS. <i>Nocardia</i> spp, <i>Madurella mycetomatis</i> , <i>Actinomadura pelletieri</i> , <i>Streptomyces somaliensis</i> , et al
Reservoir	Soil Vegetation
Vector	None
Vehicle	Contact Wound Soil
Incubation Period	2w - 2y
Diagnostic Tests	Bacterial and fungal culture of material from lesion.
Typical Adult Therapy	Antimicrobial or antifungal agent as determined by culture. Excision as indicated
Typical Pediatric Therapy	As for adult
Clinical Hints	Painless, chronic, draining, fistulous subcutaneous nodule - usually involving lower extremity; osteolytic lesions may be noted on x-ray; usually no fever; most patients are males age 20 to 40 (ie, occupational exposure).
Synonyms	<i>Leptosphaeria tompkinsii</i> , Madura foot, Madura-Fuss, <i>Madurella</i> , Mycetom, Pleurostomophora, White grain eumycetoma. ICD9: 039.4,117.4 ICD10: B47

Clinical

Mycetoma is typically characterized by a painless nodule or thickening, which involve the feet in 80% of cases. ¹

- The lesions slowly enlarge and form sinus tracts which drain bloody, serous or purulent fluid containing granules of various colors. ²
- Systemic findings are absent.
- Lesional hyperhidrosis is common, and tendons and nerves are usually spared until late stages of the infection. ³
- Regional lymphadenopathy is encountered in 1% to 3% of cases.
- Lupus vulgaris may mimic mycetoma. ⁴

Hematogenous spread of infection is extremely rare. ⁵

- Mycetoma may spread to involve contiguous bone or regional lymph nodes.
- In Actinomycotic infections, the course is more rapid and aggressive, with prominent inflammation and early destruction of bone.

Dark granules characterize *Madurella* infection, while pale colored granules are seen in *Acremonium* infection.

- *Actinomadura madurae*, *Nocardia brasiliensis*, and *Streptomyces somaliensis* produce smaller white, yellow, or brownish granules.

Rare instances of mycetoma of the scalp due to *Microsporum canis* have been reported. ⁶

- Perianal actinomycetoma may mimic other chronic diseases of the anal region.
- Ocular mycetoma has been reported as a complication of a trauma ⁷ or sub-tenon injection ⁸
- Rare cases of palatal ⁹, lingual ¹⁰, paranasal and cavernous sinus infection have been reported. ^{11 12}
- The clinical features of mycetoma may mimic those of soft tissue tumors. ¹³

Diagnosis is based on radiological and ultrasonic imaging, histology, culture and serology.

- A characteristic "dot in circle" sign may be seen on magnetic resonance imaging (MRI) studies. ¹⁴
- Although Actinomycotic lesions may be amenable to antibiotic therapy, eumycetoma requires aggressive surgical excision.

This disease is endemic or potentially endemic to all countries.

References

1. Am J Clin Dermatol 2006 ;7(5):315-21.
2. J Foot Ankle Surg 2004 Sep-Oct;43(5):327-31.
3. Infect Dis Clin North Am 2003 Mar ;17(1):59-85, viii.
4. Int J Dermatol 2009 Feb ;48(2):150-3.
5. J Neurosurg Pediatr 2008 Jun ;1(6):493-5.
6. Diagn Microbiol Infect Dis 2011 May ;70(1):145-9.
7. J Fr Ophtalmol 2013 Jan 25;
8. Cornea 2009 Sep ;28(8):933-5.

9. Rev Inst Med Trop Sao Paulo 2011 May-Jun;53(3):165-8.
10. J Oral Maxillofac Surg 2012 Nov ;70(11):e622-4.
11. PMID 19818480
12. World Neurosurg 2010 Jan ;73(1):69-71.
13. J Foot Ankle Surg 2011 Sep-Oct;50(5):569-76.
14. J Clin Imaging Sci 2012 ;2:66.

Mycobacteriosis - *M. marinum*

Agent	BACTERIUM. Actinomycetes, <i>Mycobacterium marinum</i> An aerobic acid-fast bacillus
Reservoir	Fresh and salt water (eg, swimming pools, aquaria) Fish (ornamental, salmon, sturgeon, bass)
Vector	None
Vehicle	Water per areas of minor skin trauma
Incubation Period	5d - 270d (median 21d)
Diagnostic Tests	Mycobacterial culture from lesion. Alert laboratory when this organism is suspected.
Typical Adult Therapy	Rifampicin 600 mg/day + Ethambutol 20 mg/kg/day X 6w. Alternative: Minocycline
Typical Pediatric Therapy	Sulfamethoxazole/trimethoprim 5 mg-25 mg/kg BID X 6w. Alternative Minocycline (Age >= 8)
Clinical Hints	Violaceous papule, ulcer, plaque, psoriaform lesion; onset weeks after exposure (swimming pool, aquarium); commonly involves the elbow, knee, hand or foot.
Synonyms	Aquarium granuloma, Fish fanciers' finger syndrome, Fish tank granuloma, Mariner's TB, <i>Mycobacterium balnei</i> , <i>Mycobacterium marinum</i> , <i>Mycobacterium scrofulaceum</i> , Spam, Swimming pool granuloma. ICD9: 031.1 ICD10: A31.1

Clinical

The incubation period varies from 5 to 170 days (median 21 days); with 35% of cases exceeding 30 days.

- Characteristic painful, slowly-growing blue papules usually involve the extremities, and may ulcerate. ¹
- The upper extremities are involved in 75% to 95%, and spread to deeper structures (tendons, bones, joints) occurs in 29%. ²⁻⁵
- Dissemination is rare, but has been described in AIDS patients. ⁶
- Multiple sporotrichoid subcutaneous nodules have been reported. ^{7 8}
- Extensive verrucous dermal plaques have been reported among Pacific Islanders infected by *Mycobacterium marinum*. ^{9 10}
- Tenosynovitis ("fish-tank finger") is occasionally encountered. ¹¹⁻¹⁴
- A rare case of nasal infection presenting as epistaxis has been reported. ¹⁵
- Scarring may occur, but is less pronounced than that which follows *M. ulcerans* infection.

This disease is endemic or potentially endemic to all countries.

References

1. Travel Med Infect Dis 2008 Jul ;6(4):240-4.
2. Arch Intern Med 2002 Aug 12-26;162(15):1746-52.
3. Md Med J 1995 Dec ;44(12):1043-6.
4. Semin Arthritis Rheum 1995 Jun ;24(6):382-90.
5. ProMED <promedmail.org> archive: 20110704.2026
6. Emerg Infect Dis 2003 Nov ;9(11):1496-8.
7. J Clin Rheumatol 2008 Jun ;14(3):156-60.
8. Curr Infect Dis Rep 2008 Sep ;10(5):404-10.
9. Australas J Dermatol 1998 Aug ;39(3):173-6.
10. Am J Trop Med Hyg 2008 Aug ;79(2):166-7.
11. Acta Orthop Belg 2004 Jun ;70(3):279-82.
12. N Engl J Med 2004 Aug 26;351(9):911.
13. J Infect 2007 Jun ;54(6):584-8.
14. Jpn J Infect Dis 2006 Oct ;59(5):337-40.
15. Am J Trop Med Hyg 2011 Aug ;85(2):195-6.

Mycobacteriosis - M. scrofulaceum

Agent	BACTERIUM. Actinomycetes, <i>Mycobacterium scrofulaceum</i> An aerobic acid-fast bacillus
Reservoir	Water (lakes, rivers) Soil Raw milk Plant material
Vector	None
Vehicle	Water Soil ? Through areas of minor trauma
Incubation Period	Unknown
Diagnostic Tests	Culture of tissue or aspirates.
Typical Adult Therapy	Excision. Drugs (Isoniazid - Rifampin - streptomycin - Cycloserine) are rarely indicated
Typical Pediatric Therapy	As for adult
Clinical Hints	Painless lymphadenopathy, most commonly unilateral and submandibular (true tuberculosis involves the lower neck and produces a strongly positive tuberculin reaction and/or suggestive chest X ray). The disease is most common during early childhood.
Synonyms	

Clinical

Mycobacterium scrofulaceum is a common cause of lymphadenitis, most commonly among children ages 1 to 3 years.

- Most infections involve the submandibular region, however involvement of other lymph node groups or body organs may occur. ¹
- Rare instances of dissemination are reported. ^{2 3}

This disease is endemic or potentially endemic to all countries.

References

1. Clin Dermatol 1995 May-Jun;13(3):277-80.
2. Clin Infect Dis 1996 Jan ;22(1):159-61.
3. Clin Infect Dis 1995 Mar ;20(3):549.

Mycobacteriosis - M. ulcerans

Agent	BACTERIUM. Actinomycetes, <i>Mycobacterium ulcerans</i> An aerobic acid-fast bacillus
Reservoir	Vegetation
Vector	Mosquitoes (probable)
Vehicle	Contact
Incubation Period	3w - 3m
Diagnostic Tests	Mycobacterial culture from lesion. Alert laboratory that this organism is suspected.
Typical Adult Therapy	Rifampin + amikacin. OR Ethambutol + Sulfamethoxazole/trimethoprim X 6w. OR Rifampin + a fluoroquinolone. Excision as indicated
Typical Pediatric Therapy	Rifampin 20 mg/kg/day + amikacin 7.5 mg/kg q12h X 6w. Excision as indicated
Clinical Hints	Slowly growing, painless ulcerative nodule with undermined edges - lymphadenopathy usually not present; in most cases, a single leg lesion involving the extensor surface (face and trunk may be involved in children).
Synonyms	Bairnsdale ulcer, Buruli ulcer, Kakerifu ulcer, Kasongo ulcer, Kisikro, Kumusi ulcer, Mexican ulcer, <i>Mycobacterium buruli</i> , <i>Mycobacterium ulcerans</i> , Searl's ulcer, Tora ulcer, Ulcerans disease. ICD9: 031.1 ICD10: A31.1

Clinical

Buruli ulcer is a chronic, indolent, necrotizing disease of the skin. ¹

- Lesions usually involve the limbs (the face ² and trunk in children); 60% involve the lower limbs, 30% upper limbs and 10% other body regions. ³⁻⁵
- The process begins as firm, nontender subcutaneous nodules 1 to 2 cm in diameter at sites of penetrating skin trauma.
- Within the next 1 to 2 months, these areas become fluctuant, followed by the formation of a painless, undermined ulceration. ⁶

Ulcerations may involve up to 15% of the patient's skin surface, and may destroy nerves and blood vessels, or even invade bone.

- Some cases may present as dermal plaques or cellulitis, without ulcer.
- Most lesions eventually heal spontaneously, but frequently result in chronic lymphedema and disfiguring scarring. ^{7 8}
- Skin lesions may re-emerge as long as one year following therapy, possibly as a result of immune reconstitution. ⁹
- Severe infection may induce contracture and atrophy of contiguous skeletal muscles. ¹⁰
- Systemic symptoms are rare, and healing may take 4 to 6 months, with extensive scar formation and limb deformity.
- Rare instances of systemic spread ¹¹ and multifocal osteomyelitis have been reported. ¹²
- Hemoglobinopathy appears to be a risk factor for *Mycobacterium ulcerans* osteomyelitis ¹³


This disease is endemic or potentially endemic to 37 countries.

Mycobacteriosis - M. ulcerans in Uganda

The term 'Buruli ulcer' dates from reports of lesions among Rwandan refugees living in Buruli County, Mengo, Uganda.
- Highest rates previously occurred in swampy regions along the Nile River.

Cases were reported on the east bank of the Victoria Nile during 1962 to 1964, following heavy rains.

As of 2002, the disease appeared to have disappeared from Uganda ¹⁴ ; however, 117 suspect cases were reported in the Nakasongola district (formerly Buruli district) in 2003. ¹⁵
- Additional cases may exist in Adjumani district (formerly Madi district).



Graph: Uganda. Buruli ulcer, cases

References

1. Expert Rev Anti Infect Ther 2003 Aug ;1(2):217-22.
2. Am J Trop Med Hyg 2011 Dec ;85(6):1100-5.
3. Wkly Epidemiol Rec 2008 Apr 25;83(17):145-54.
4. Med Trop (Mars) 2008 Dec ;68(6):643-4.
5. Ann Dermatol Venereol 2013 Feb ;140(2):125-8.
6. Bull World Health Organ 2005 Oct ;83(10):785-91.
7. Trans R Soc Trop Med Hyg 1999 Jul-Aug;93(4):337-40.
8. Lancet 1999 Sep 18;354(9183):1013-8.
9. PLoS Negl Trop Dis 2011 Aug ;5(8):e1252.
10. Med Sci (Paris) 2011 Feb ;27(2):187-92.
11. World J Gastroenterol 2008 Feb 7;14(5):808-11.
12. Am J Trop Med Hyg 2010 Aug ;83(2):307-13.
13. Trop Med Int Health 2007 Apr ;12(4):511-8.
14. Wkly Epidemiol Rec 2002 May 17;77(20):165-6.
15. Wkly Epidemiol Rec 2004 May 14;79(20):194-9.

Mycobacteriosis - miscellaneous nontuberculous

Agent	BACTERIUM. Actinomycetes, Mycobacterium spp. An aerobic acid-fast bacillus
Reservoir	Water Soil Fish Mammal Bird
Vector	None
Vehicle	Air Water Contact Ingestion Trauma
Incubation Period	Variable
Diagnostic Tests	Microscopy & culture of tissue, secretions, blood. Nucleic acid amplification. Inform laboratory if suspected
Typical Adult Therapy	Drug, route and duration appropriate to clinical setting and species [in Therapy module, scroll through upper left box]
Typical Pediatric Therapy	As for adult
Clinical Hints	Pneumonia, or chronic granulomatous infection of various tissues; systemic disease may complicate immune suppression; <i>M. avium-intracellulare</i> characterized by aggressive course and resistance to most antimycobacterial drugs.
Synonyms	<i>Mycobacterium abscessus</i> , <i>Mycobacterium avium</i> , <i>Mycobacterium avium-intracellulare</i> , <i>Mycobacterium franklinii</i> , <i>Mycobacterium immunogenum</i> , <i>Mycobacterium jaccussii</i> , <i>Mycobacterium xenopi</i> , <i>Segniliparus</i> . ICD9: 031.9,031.2 ICD10: A31.0,A31.1,A31.8

Clinical

The clinical features of systemic mycobacterial infection are protean, and can involve disease of virtually any organ or tissue. ¹⁻⁵

- Specific syndromes reflect the immune status of the patient and the specific fungal species involved (see Worldwide note)

Mycobacterium avium-intracellulare infection is clinically similar to tuberculosis, producing localized pulmonary disease ⁶ or disseminated lesions of virtually any organ. ^{7 8}

- Bacteremia is common, and can be detected using specialized blood culture systems.

Mycobacterium kansasii infection is characterized by productive cough, dyspnea, and chest pain.

- 16% of patients are asymptomatic.
- A right sided, apical or subapical, thin walled cavitary infiltrate is characteristic. ⁹

Mycobacterium malmoense infection is usually characterized by pulmonary disease suggestive of tuberculosis, or pediatric cervical lymphadenopathy. ¹⁰

Note: Over 110 species of *Mycobacterium* have been associated with human infection.

- See Microbiology • Mycobacteria module

This disease is endemic or potentially endemic to all countries.

Mycobacteriosis - miscellaneous nontuberculous in Uganda

Organisms associated with cervical lymphadenitis among pastoral communities in the Karamoja region include *Mycobacterium avium*, *M. tuberculosis*, *M. bovis* and *M. intracellulare*. (2008 publication). ¹¹

References

1. Infection 2004 Oct ;32(5):257-70.
2. Clin Chest Med 2002 Sep ;23(3):553-67.
3. Scand J Infect Dis 2006 ;38(4):246-55.
4. Emerg Infect Dis 2010 Jan ;16(1):166-8.
5. Clin Microbiol Infect 2004 Nov ;10(11):951-3.
6. Int J Tuberc Lung Dis 2007 Feb ;11(2):215-21.
7. Am J Med Sci 2009 Mar ;337(3):218-20.
8. J Med Microbiol 2010 Nov ;59(Pt 11):1365-7.

9. [Postgrad Med J 2005 Aug ;81\(958\):530-3.](#)
10. [Int J Tuberc Lung Dis 2008 Sep ;12\(9\):987-93.](#)
11. [Epidemiol Infect 2008 May ;136\(5\):636-43.](#)

Mycoplasma (miscellaneous) infections

Agent	BACTERIUM. Mycoplasmatales <i>Mycoplasma genitalium</i> , <i>Mycoplasma hominis</i> , <i>Mycoplasma fermentans</i> , <i>Mycoplasma penetrans</i> , <i>Ureaplasma urealyticum</i>
Reservoir	Human
Vector	None
Vehicle	Secretion, Sexual transmission
Incubation Period	Unknown
Diagnostic Tests	Culture (urine, pharynx). Serology. Nucleic acid amplification.
Typical Adult Therapy	<i>Azithromycin</i> 1 g orally as single dose OR <i>Doxycycline</i> 100 mg PO BID X 7 days OR <i>Levofloxacin</i> 500 mg daily X 7 days OR <i>Ofloxacin</i> 300 mg BID X 7 days
Typical Pediatric Therapy	<i>Erythromycin</i> 10 mg/kg PO QID X 2w
Clinical Hints	Urethritis, vaginitis, neonatal pneumonia; rarely stillbirth, prematurity or infertility
Synonyms	<i>Acholeplasma laidlawii</i> , <i>Epirythrozoon</i> , Hemotrophic <i>Mycoplasma</i> , <i>Mycoplasma amphoriforme</i> , <i>Mycoplasma buccale</i> , <i>Mycoplasma faucium</i> , <i>Mycoplasma felis</i> , <i>Mycoplasma fermentans</i> , <i>Mycoplasma genitalium</i> , <i>Mycoplasma hominis</i> , <i>Mycoplasma lipophilum</i> , <i>Mycoplasma orale</i> , <i>Mycoplasma penetrans</i> , <i>Mycoplasma pirum</i> , <i>Mycoplasma primum</i> , <i>Mycoplasma salivarium</i> , <i>Mycoplasma spermatophilum</i> , T <i>Mycoplasmas</i> , T strains, <i>Ureaplasma parvum</i> , <i>Ureaplasma urealyticum</i> . ICD9: 041.81 ICD10: A49.3

Clinical

Asymptomatic pharyngeal and vaginal carriage of *Mycoplasma* species and *Ureaplasma* is common.

- As many as 70% of sexually-active persons are colonized.

The signs and symptoms of infection are similar to those of *Chlamydia* infection.¹

- Urogenital infection may present as vaginitis, cervicitis, non-gonococcal urethritis, epididymitis², prostatitis³ or urethral discharge.
- Less common findings may include pelvic inflammatory disease⁴⁻⁶, post-partum fever^{7 8}, chorioamnionitis, infertility⁹, prematurity¹⁰ and stillbirth.¹¹⁻¹⁴
- Bronchitis, arthritis^{15 16}, neonatal meningitis and encephalitis¹⁷⁻¹⁹, osteitis²⁰, endocarditis^{21 22}, brain abscess²³, soft tissue infections²⁴, genital under disease²⁵, bacteremia²⁶, respiratory distress in the newborn²⁷ and pneumonia have been reported.²⁸⁻³¹

Infection by hemotrophic *Mycoplasma* species (formerly *Epirythrozoon*) is characterized by fever, anemia and hemolytic jaundice • notably among pregnant women and newborns.³²

This disease is endemic or potentially endemic to all countries.

Mycoplasma (miscellaneous) infections in Uganda

Prevalence surveys:

Mycoplasma genitalium infection was found in 14% of CSW (Kampala, 2011 publication)³³

References

1. Clin Infect Dis 2009 Jan 1;48(1):41-7.
2. Genitourin Med 1988 Dec ;64(6):367-8.
3. Scand J Urol Nephrol 2005 ;39(6):479-82.
4. Genitourin Med 1985 Jun ;61(3):185-9.
5. J Med Microbiol 2005 Dec ;54(Pt 12):1249-50.
6. Curr Opin Infect Dis 2008 Feb ;21(1):65-9.
7. Lancet 1980 Dec 6;2(8206):1217-21.
8. Pediatr Infect Dis 1986 Nov-Dec;5(6 Suppl):S258-61.
9. Pediatr Infect Dis 1986 Nov-Dec;5(6 Suppl):S262-5.
10. Trans Assoc Am Physicians 1981 ;94:261-6.
11. Am J Obstet Gynecol 1983 Jan 15;145(2):245-50.
12. Pediatr Infect Dis 1986 Nov-Dec;5(6 Suppl):S282-7.
13. Scand J Infect Dis 2001 ;33(8):604-10.
14. Infect Dis Obstet Gynecol 2011 ;2011:959816.

15. *J Infect* 2007 Nov ;55(5):e135-7.
16. *Am J Transplant* 2005 Jan ;5(1):183-8.
17. *Arch Dis Child* 1979 Mar ;54(3):231-3.
18. *Sex Transm Dis* 1983 Oct-Dec;10(4 Suppl):331-4.
19. *Eur J Pediatr* 2003 Jul ;162(7-8):514-6.
20. *Eur J Clin Microbiol Infect Dis* 2006 Nov ;25(11):715-7.
21. *Pediatr Infect Dis J* 2006 Sep ;25(9):851-2.
22. *Clin Infect Dis* 2004 Feb 1;38(3):e21-4.
23. *Pediatr Infect Dis J* 2002 Nov ;21(11):1083-5.
24. *Pediatr Infect Dis J* 2002 Dec ;21(12):1171-3.
25. *Sex Transm Dis* 1983 Oct-Dec;10(4 Suppl):285-8.
26. *Eur J Clin Microbiol Infect Dis* 2003 Oct ;22(10):608-11.
27. *BMC Infect Dis* 2006 ;6:166.
28. *Pediatr Infect Dis* 1986 Nov-Dec;5(6 Suppl):S288-91.
29. *Clin Microbiol Rev* 1993 Jan ;6(1):69-87.
30. *Intensive Care Med* 2007 Jan ;33(1):143-7.
31. *Scand J Infect Dis* 2003 ;35(4):282-4.
32. *Emerg Infect Dis* 2009 Jul ;15(7):1139-40.
33. *J Infect Dis* 2012 Jan 15;205(2):289-96.

Mycoplasma pneumoniae infection

Agent	BACTERIUM. Mollicutes. Mycoplasma pneumoniae
Reservoir	Human
Vector	None
Vehicle	Droplet
Incubation Period	6d - 23d
Diagnostic Tests	Culture (sputum, throat). Serology. Nucleic acid amplification.
Typical Adult Therapy	Erythromycin 500 mg PO BID X 2w. OR Azithromycin 1 g, followed by 500 mg PO daily X 5 days. OR Doxycycline 100 mg PO BID
Typical Pediatric Therapy	Erythromycin 10 mg/kg PO QID X 2w
Clinical Hints	Coryza, "hacking" cough; subsegmental infiltrate; bullous otitis media is often present; most patients below age 30; cold agglutinins are neither sensitive nor specific for infection, and appear only during second week.
Synonyms	<i>Mycoplasma pneumoniae</i> , Primary atypical pneumonia. ICD9: 041.81,483.0 ICD10: B96.0

Clinical

Acute infection:

Onset is insidious and gradual, and characterized by fever, malaise, a dry cough, headache, 'scratchy' throat and chest wall (ie, muscular) pain. ¹

- Pleuritic pain, productive cough and rigors are unusual and should suggest infection by other bacterial species.
- A lymphocytic pleural effusion may be present. ²
- The pharynx and tympanic membranes are often erythematous, without adenopathy; and the lungs are usually normal to auscultation.
- A macular, urticarial or vesicular rash is occasionally present; and erythema multiforme / mucositis ³ (including Stevens-Johnson syndrome) is reported in some cases. ⁴⁻¹¹

Atypical manifestations: ¹²

Atypical and severe disease is encountered among older adults.

- Rare instances of acute hepatitis ^{13 14}, glomerulonephritis ^{15 16}, rhabdomyolysis ¹⁷⁻¹⁹, septic shock ²⁰, endocarditis ²¹, ARDS ²², multi-organ failure ²³, acute respiratory distress syndrome ²⁴, pericarditis and empyema have been reported. ²⁵
- Neurological findings may include encephalitis ²⁶⁻³⁶, brainstem / striatal encephalopathy ³⁷, post-encephalitic seizures ³⁸, aseptic meningitis ³⁹⁻⁴¹, acute transverse myelitis ^{42 43}, stroke ^{44 45}, or polyradiculopathy. ⁴⁶⁻⁴⁸
- Obsessive-compulsive disorder has been ascribed to *Mycoplasma pneumoniae* infection ⁴⁹
- Extrapulmonary manifestations also include hematologic (including autoimmune hemolytic anemia ^{50 51}), pancytopenia ⁵², acute thrombocytosis ⁵³; renal; gastrointestinal; genitourinary ⁵⁴; hepatic ^{55 56}; osteoarticular ⁵⁷; cutaneous (rash, angioedema with eosinophilia ⁵⁸), papular purpuric gloves and socks syndrome (PPGSS) ⁵⁹, leukocytoclastic vasculitis ⁶⁰, urticarial vasculitis suggestive of adult Still's disease ⁶¹, toxic epidermal necrolysis ⁶², mucositis ⁶³; possible splenic infarction ⁶⁴; and ocular involvement (including vasculitis ⁶⁵ and optic neuritis ⁶⁶ / papillitis). ⁶⁷
- *Mycoplasma pneumoniae* infection is implicated in the etiology of Guillain-Barre syndrome ⁶⁸⁻⁷⁰, recurrent tonsillitis ⁷¹ and asthma. ⁷²⁻⁷⁸
- *Mycoplasma pneumoniae* infection is independently associated with risk of subsequent development of ischemic stroke ⁷⁹ and may play a role in the development of atherosclerosis. ⁸⁰

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 2004 Oct ;17(4):697-728, table of contents.
2. Scand J Infect Dis 2012 Oct ;44(10):793-7.
3. Acta Paediatr 2011 Nov ;100(11):e238-40.
4. Pediatr Dermatol 2006 Nov-Dec;23(6):546-55.
5. Minerva Pediatr 2008 Feb ;60(1):135-9.
6. J Dtsch Dermatol Ges 2009 May ;7(5):445-8.
7. Rev Chilena Infectol 2009 Oct ;26(5):457-63.
8. Pediatrics 2011 Jun ;127(6):e1605-9.
9. Indian J Pediatr 2011 Oct 20;
10. Case Report Med 2012 ;2012:430490.
11. Case Rep Infect Dis 2013 ;2013:457161.
12. J Infect Chemother 2010 Jun ;16(3):162-9.
13. Yonsei Med J 2008 Dec 31;49(6):1055-9.
14. Eur J Gastroenterol Hepatol 2009 Feb ;21(2):220-1.
15. Pediatr Nephrol 2010 Sep ;25(9):1765-9.
16. Medicina (Kaunas) 2010 ;46(5):360-3.
17. Pediatr Neurol 2009 Feb ;40(2):128-30.
18. Emerg Infect Dis 2012 May ;18(5):849-51.
19. Hong Kong Med J 2012 Jun ;18(3):247-9.
20. Clin Pediatr (Phila) 2009 Apr ;48(3):320-2.
21. Emerg Infect Dis 2008 Oct ;14(10):1664-5.
22. Indian J Pathol Microbiol 2010 Jul-Sep;53(3):555-7.
23. Allergy Asthma Immunol Res 2012 Jan ;4(1):55-7.
24. Rev Mal Respir 2013 Jan ;30(1):77-80.
25. BMC Infect Dis 2006 ;6:18.
26. Tunis Med 2010 Feb ;88(2):125-8.
27. Pediatr Neurol 2010 Jul ;43(1):41-5.
28. Curr Infect Dis Rep 2010 Jul ;12(4):282-90.
29. Klin Padiatr 2011 Jul ;223(4):209-13.
30. Neuropathology 2012 Feb ;32(1):91-9.
31. Case Rep Neurol 2011 May ;3(2):109-12.
32. Clin Infect Dis 1998 Feb ;26(2):398-409.
33. Eur J Clin Microbiol Infect Dis 2009 Jan ;28(1):91-4.
34. J Neurol Sci 2011 Oct 15;309(1-2):18-25.
35. Rev Neurol (Paris) 2012 Jan ;168(1):49-52.
36. Acta Neuropathol 2012 Feb 28;
37. No To Hattatsu 2011 Nov ;43(6):471-5.
38. Epilepsia 2011 Nov ;52(11):1979-85.
39. Acta Paediatr 2009 Aug ;98(8):1300-6.
40. Med Mal Infect 2007 Nov ;37(11):738-45.
41. Einstein (Sao Paulo) 2012 Jan-Mar;10(1):100-2.
42. Pediatr Neurol 2009 Feb ;40(2):128-30.
43. Eur J Neurol 2006 Feb ;13(2):112-24.
44. Pediatr Pulmonol 2012 Mar 29;
45. Zhongguo Dang Dai Er Ke Za Zhi 2012 Nov ;14(11):823-6.
46. J Infect 2005 Dec ;51(5):343-54.
47. Curr Opin Neurol 2006 Aug ;19(4):374-8.
48. Rev Med Brux 2008 Mar-Apr;29(2):103-6.
49. J Child Neurol 2008 Mar ;23(3):338-40.
50. Braz J Infect Dis 2009 Feb ;13(1):77-9.
51. Case Report Med 2012 ;2012:649850.
52. J Pediatr Hematol Oncol 2009 Oct ;31(10):760-2.
53. Heart Lung 2009 Sep-Oct;38(5):444-9.
54. Pediatr Infect Dis J 2008 Mar ;27(3):280-2.
55. Nihon Kyobu Shikkan Gakkai Zasshi 1991 Jun ;29(6):693-7.
56. Digestion 2012 Oct 23;86(4):302-308.
57. Pediatr Int 2008 Aug ;50(4):511-3.
58. Clin Rheumatol 2008 Dec ;27(12):1573-6.
59. Clin Pediatr (Phila) 2011 Dec ;50(12):1140-3.
60. Acta Dermatovenerol Croat 2012 ;20(2):119-22.
61. Rheumatol Int 2011 Sep 15;
62. Arch Argent Pediatr 2013 Feb ;111(1):e24-7.
63. J Dtsch Dermatol Ges 2012 Oct ;10(10):740-6.
64. J Infect Chemother 2012 Feb 22;
65. Clin Pediatr (Phila) 2007 Jun ;46(5):451-3.
66. Case Rep Neurol 2011 May ;3(2):109-12.
67. Neurol Sci 2012 Apr ;33(2):355-8.
68. J Infect Dev Ctries 2011 Jun ;5(6):459-64.
69. Clin Microbiol Infect 2007 Mar ;13(3):334-7.
70. J Neurol 2011 Nov ;258(11):1958-64.
71. Eur J Clin Microbiol Infect Dis 2008 Dec ;27(12):1233-7.
72. FEMS Immunol Med Microbiol 2009 Jun ;56(1):25-31.
73. Curr Allergy Asthma Rep 2010 Jan ;10(1):67-73.
74. Rev Mal Respir 2010 Oct ;27(8):890-7.
75. Immunol Allergy Clin North Am 2010 Nov ;30(4):575-85, vii-viii.
76. Immunol Allergy Clin North Am 2010 Nov ;30(4):565-74, vii.
77. Allergy 2011 Apr ;66(4):458-68.
78. Allergol Immunopathol (Madr) 2007 Jan-Feb;35(1):4-9.
79. Stroke 2011 Oct ;42(10):2940-3.
80. GMS Krankenhhyg Interdiszip 2011 ;6(1):Doc04.

Myiasis

Agent	PARASITE - Insecta (Diptera) larvae
Reservoir	Mammal
Vector	Biting arthropod
Vehicle	Fly eggs deposited by biting arthropod
Incubation Period	1w - 3m
Diagnostic Tests	Identification of extracted maggot.
Typical Adult Therapy	Removal of maggot
Typical Pediatric Therapy	As for adult
Clinical Hints	Pruritic or painful draining nodule; fever and eosinophilia may be present; instances of brain, eye, middle ear and other deep infestations are described.
Synonyms	Calliphora, Chrysomya, Chrysomyia, Cochliomyia, Cordylobia, Cuterebrosis, Dermatobia, Eristalis, Furuncular myiasis, Gasterophilus, Hypoderma, Lucilia, Lund's fly, Maggot infestation, Megaselia, Musca, Muscina, Oedemagena, Oestrus larvae, Ophthalmomyiasis, Psychoda, Rectal myiasis, Sarcophaga, Screw worm, Telmatoscopus, Urinary myiasis, Vaginal myiasis, Wohlfarthia. ICD9: 134.0 ICD10: B87

Clinical

Myiasis may be primary (active invasion) or secondary (colonization of wound). ¹

- Primary furuncular myiasis is usually characterized by one or more erythematous, painful "pustules" having a central perforation. ²
- Eosinophilia may be present. ³
- Other clinical forms include ophthalmomyiasis (migrating larvae in the conjunctival sac), pharyngeal, nasal ⁴, urinary, vaginal, tracheopulmonary and rectal infestation.
- Larvae may rarely invade the paranasal sinuses and even cause eosinophilic meningitis. ⁵
- Penile myiasis may mimic a sexually transmitted disease ⁶

This disease is endemic or potentially endemic to all countries.

References

1. J Am Acad Dermatol 2008 Jun ;58(6):907-26; quiz 927-9.
2. J Am Acad Dermatol 2004 Feb ;50(2 Suppl):S26-30.
3. Turk J Pediatr 2009 May-Jun;51(3):279-81.
4. Braz J Otorhinolaryngol 2009 May-Jun;75(3):356-61.
5. Clin Microbiol Rev 2009 Apr ;22(2):322-48, Table of Contents.
6. Braz J Infect Dis 2008 Apr ;12(2):155-7.

Necrotizing skin/soft tissue infx.

Agent	BACTERIUM. <i>Streptococcus pyogenes</i> , <i>Clostridium perfringens</i> , mixed anaerobic and/or gram-negative bacilli
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Clinical features. Smear and culture (including anaerobic culture) of exudate.
Typical Adult Therapy	Debridement and parenteral antibiotics directed by smear and culture results. Hyperbaric oxygen in more severe infections
Typical Pediatric Therapy	As for adult
Clinical Hints	At least 7 syndromes in this category: most characterized by local pain and swelling, skin discoloration or edema, gas formation, foul odor and variable degrees of systemic toxicity.
Synonyms	Anaerobic cellulitis, Chancrum oris, Clostridial cellulitis, <i>Clostridium novyi</i> , Fasciitis, Fournier's gangrene, Gangrenous cellulitis, Gangrenous stomatitis, Invasive group A strep. Infections, Meleney's synergistic gangrene, Necrotizing fasciitis, Noma, Streptococcal fasciitis, Synergistic necrotizing cellulitis. ICD9: 686.8,528.1 ICD10: M72.6,A69.0

Clinical

Infections often begin in areas of minor trauma or loss of dermal integrity (as in varicella), and may spread within hours to involve large areas and endanger life. ¹⁻⁵

Clinical forms of necrotizing skin and soft tissue infection (in alphabetical order):

Clostridial cellulitis usually follows local trauma or surgery, and has a gradual onset following an incubation period of 3 or more days.

- There is minimal pain and discoloration, with moderate swelling.
- A thin, occasionally foul and dark colored exudate is noted and copious gas is present.
- Systemic signs are minimal.

Clostridial myonecrosis is discussed elsewhere in this module • but is distinguishable from the above syndromes by its severity, prominent systemic toxicity and the presence of overt muscle involvement.

Fournier's gangrene is a form of necrotizing fasciitis which involves the scrotum and penis. ⁶⁻⁸

- Most patients are over the age of 50 • diabetic, alcoholic or suffering from rectal cancer.
- The lesion is markedly destructive and mutilating, and typically due to a mixed flora of anaerobic and facultative or aerobic gram negative bacilli.
- Fournier's gangrene may occasionally complicate varicella ⁹
- The case fatality rate for Fournier's gangrene is over 20% ¹⁰

Gangrenous stomatitis (chancrum oris, Noma) is a mutilating condition of the skin and soft tissues of the face which affects primarily immune-suppressed ¹¹⁻¹³ and malnourished children. ¹⁴⁻¹⁹

- Most patients are under the age of 6 years.
- The disease usually begins as a painful red or purple intraoral lesion, which rapidly spreads to destroy surrounding bone and soft tissues of the mouth and face.
- The case-fatality rate is 70% to 90%.

Infected vascular gangrene is a complication of peripheral vascular insufficiency and has a gradual onset beginning 5 or more days after the initiating event.

- Onset is gradual, and pain may vary from absent to prominent.
- The area is discolored and painful, and associated with foul malodorous gas and involvement of underlying muscle.
- Systemic signs are minimal.

Meleney's gangrene (progressive bacterial synergistic gangrene) usually involves sites of fistulae, retention sutures or draining empyema. ^{20 21}

- The infection begins 1 to 2 weeks following surgery, and is characterized by erythema and moderate swelling, with minimal crepitus.

Necrotizing fasciitis is typically associated with diabetes mellitus or recent abdominal surgery. ^{22 23}

- Following an incubation period of 1 to 4 days, the patient becomes increasingly ill, with moderate local pain and gas formation, and a foul seropurulent discharge.
- Pain may be severe, and areas of erythema and necrosis are evident.
- Relatively high mortality rates are associated with necrotizing fasciitis caused by *Aeromonas* or *Vibrio* species. ²⁴

Non-clostridial anaerobic cellulitis is usually associated with diabetes mellitus or a preexisting local infection.

- Onset may be gradual or rapid, with moderate swelling, dark pus, minimal discoloration and copious foul-smelling gas.
- Pain is minimal, and the patient is moderately ill.

Synergistic necrotizing cellulitis is associated with diabetes, renal disease, obesity or preexisting perirectal infection.

- The incubation period varies from 3 to 14 days, and onset is acute.
- Swelling may be marked, and associated with intense local pain, foul 'dishwater' pus and small amounts of gas.
- Moderate muscle involvement and marked systemic disease are present.

This disease is endemic or potentially endemic to all countries.

References

1. Clin Infect Dis 2007 Mar ;44(5):705-10.
2. Am Fam Physician 2003 Jul 15;68(2):323-8.
3. Infect Dis Clin North Am 1996 Dec ;10(4):835-55.
4. J Antimicrob Chemother 2004 Jun ;53 Suppl 2:ii37-50.
5. Lancet Infect Dis 2005 Aug ;5(8):501-13.
6. Urology 2004 Aug ;64(2):218-22.
7. Ann Ital Chir 2004 Jan-Feb;75(1):97-106; discussion 106.
8. Surg Clin North Am 2002 Dec ;82(6):1213-24.
9. Pediatr Emerg Care 2007 Oct ;23(10):719-20.
10. J Microbiol Immunol Infect 2007 Dec ;40(6):500-6.
11. Ann Dermatol Venereol 2003 Feb ;130(2 Pt 1):199-201.
12. Odontostomatol Trop 2001 Dec ;24(96):26-9.
13. Lancet 1989 Jul 8;2(8654):108.
14. Br J Oral Maxillofac Surg 2004 Jun ;42(3):267-9.
15. Br J Plast Surg 2003 Sep ;56(6):524-7.
16. Trop Med Int Health 2003 May ;8(5):402-7.
17. Plast Reconstr Surg 2003 Apr 15;111(5):1702-7.
18. Dakar Med 1998 ;43(1):45-8.
19. Quintessence Int 1997 Apr ;28(4):277-81.
20. Ann Med Interne (Paris) 1989 ;140(3):182-5.
21. Br J Surg 1990 Mar ;77(3):260-4.
22. Curr Opin Infect Dis 2005 Apr ;18(2):101-6.
23. Curr Opin Infect Dis 2007 Apr ;20(2):118-23.
24. Am J Emerg Med 2008 Feb ;26(2):170-5.

Neutropenic typhlitis

Agent	BACTERIUM. <i>Clostridium septicum</i> (occasionally <i>Clostridium tertium</i> , <i>Clostridium sporogenes</i> , <i>Clostridium sordellii</i> or <i>Clostridium tertium</i>)
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Unknown
Diagnostic Tests	Typical findings in the setting of neutropenia. Ultrasonography may be helpful.
Typical Adult Therapy	Broad spectrum antimicrobial coverage, which should include clostridia and <i>Pseudomonas aeruginosa</i> . Role of surgery is controversial
Typical Pediatric Therapy	Broad spectrum antimicrobial coverage, which should include clostridia and <i>Pseudomonas aeruginosa</i> . Role of surgery is controversial
Clinical Hints	Fever, abdominal pain, diarrhea (occasionally bloody) and right lower quadrant signs in a neutropenic (leukemic, etc) patient; may spread hematogenously to extremities; case-fatality rate 50% to 75%.
Synonyms	Neutropenic enterocolitis. ICD9: 540.0 ICD10: A04.8

Clinical

Neutropenic typhlitis is clinically similar to acute appendicitis, but limited to patients with severe neutropenia. ¹⁻³

This disease is endemic or potentially endemic to all countries.

References

1. Clin Infect Dis 1993 Sep ;17(3):484-90.
2. Ann Surg 1987 Dec ;206(6):711-6.
3. Curr Opin Gastroenterol 2006 Jan ;22(1):44-7.

Nocardiosis

Agent	BACTERIUM. Actinomycetes, Nocardia spp. An aerobic gram positive bacillus (acid-fast using special technique)
Reservoir	Soil
Vector	None
Vehicle	Air Dust Wound Contact
Incubation Period	? days to weeks
Diagnostic Tests	Culture and gram stain of exudates, sputa, tissue specimens. Advise laboratory when Nocardia suspected.
Typical Adult Therapy	Sulfamethoxazole/trimethoprim - dosage and duration of therapy appropriate to clinical severity
Typical Pediatric Therapy	As for adult
Clinical Hints	Pneumonia, lung abscess, brain abscess, or other chronic suppurative infection; often in the setting of immune suppression.
Synonyms	Nocardia, Nocardiose. ICD9: 039 ICD10: A43

Clinical

Nocardiosis may present as an acute or chronic suppurative infection with a tendency to remission and exacerbation. ¹

- Infections are most common among immunocompromised patients. ^{2 3}
- The most common presentation is pneumonia.
- Brain abscesses account for 33% of cases.
- Infection of virtually any other organ may occur. ⁴⁻⁷

Nocardiosis may mimic tuberculosis, particularly in the setting of HIV infection. ⁸

- Nodular lymphadenitis, seen with *Nocardia brasiliensis* infection, may mimic nocardiosis. ⁹

The ecology and phenotypic characteristics of *Nocardia* species ¹⁰ are discussed in the Microbiology module.

This disease is endemic or potentially endemic to all countries.

References

1. *Medicine (Baltimore)* 2004 Sep ;83(5):300-13.
2. *Am J Transplant* 2004 Nov ;4 Suppl 10:47-50.
3. *Medicine (Baltimore)* 2004 Sep ;83(5):300-13.
4. *Intern Med* 2013 ;52(1):129-33.
5. *J Microbiol Immunol Infect* 2011 Jun ;44(3):238-40.
6. *BMC Infect Dis* 2009 ;9:194.
7. *Breast J* 2012 Nov 27;
8. *Trans R Soc Trop Med Hyg* 2008 Mar ;102(3):219-24.
9. *Curr Infect Dis Rep* 2008 Sep ;10(5):404-10.
10. *J Clin Microbiol* 2005 Jun ;43(6):2624-8.

O'nyong nyong

Agent	VIRUS - RNA. Togaviridae, Alphavirus: O'nyong nyong virus
Reservoir	Unknown
Vector	Mosquito (<i>Anopheles funestus</i> and <i>An. gambiae</i>)
Vehicle	None
Incubation Period	3d - 12d
Diagnostic Tests	Viral culture (blood). Serology. Nucleic acid amplification. Biosafety level 2.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Myalgia and severe arthralgia; maculopapular rash (often pruritic) and leukopenia; conjunctivitis and cervical lymphadenopathy; fever resolves within 7 days, however arthralgia may persist.
Synonyms	Igbo Ora. ICD9: 066.3 ICD10: A92.8

Clinical

O'nyong nyong is characterized by fever, arthralgia, headache, conjunctivitis, myalgia and lymphadenopathy. ¹

- Knees and ankles are most commonly involved, and lymphadenopathy affects primarily the cervical region.
- Most patients develop a generalized rash which may be pruritic.

Infection by a related virus, Igbo Ora, is characterized by fever, headache, rash and arthralgia.

This disease is endemic or potentially endemic to 16 countries.

O'nyong nyong in Uganda

The etiologic agent of O'nyong nyong was first isolated in Acholi (O'nyong nyong means 'weakening of the joints' in the Acholi dialect). ²

The virus has been recovered from *Anopheles gambiae*, *An. funestus* and *Mansonia (Mansonioides) uniformis*. ³

Rates as high as 38% were recorded in affected areas.

Notable outbreaks:

1959 to 1962 - An outbreak of O'nyong nyong was reported, with *Anopheles gambiae* and *An. funestus* implicated as vectors. ⁴

1996 to 1997 - An outbreak of O'nyong nyong was reported in southwest Uganda (Mbarara, Masaka, Kabarole and Rakai districts), later spreading north-westward into the Sembabule area. ⁵⁻⁸ The implicated vectors were *An. funestus* and *Mansonia uniformis*. ⁹

References

1. Clin Infect Dis 1999 Nov ;29(5):1243-50.
2. Trop Geogr Med 1963 Dec ;15:411-21.
3. Am J Trop Med Hyg 1999 Jul ;61(1):158-62.
4. Trans R Soc Trop Med Hyg 1965 May ;59:300-6.
5. J Infect Dis 1999 Nov ;180(5):1436-43.
6. Emerg Infect Dis 1997 Jan-Mar;3(1):77.
7. Am J Trop Med Hyg 1999 Jul ;61(1):158-62.
8. ProMED <promedmail.org> archive: 19970311.0552
9. Emerg Infect Dis 1997 Jan-Mar;3(1):77.

Oesophagostomiasis

Agent	PARASITE - Nematoda. Phasmidea: Oesophagostomum bifurcum (O. apiostomum, O. stephanostomum)
Reservoir	Non-human primate Soil
Vector	None
Vehicle	Feces Water Soil
Incubation Period	2w - 2m
Diagnostic Tests	Demonstration of parasite in tissue.
Typical Adult Therapy	Albendazole (400 mg as single dose), or Pyrantel pamoate may be effective. Excision as necessary
Typical Pediatric Therapy	Albendazole or Pyrantel pamoate may be effective. Excision as necessary
Clinical Hints	Right lower quadrant abdominal pain and tenderness, often with intraabdominal mass or peritoneal signs.
Synonyms	Dapaong tumor, Oesophagostomum apiostomum, Oesophagostomum bifurcum, Oesophagostomum stephanostomum, Ternidens. ICD9: 127.7 ICD10: B81.8

Clinical

Oesophagostomiasis is contracted through ingestion of soil-contaminated food or water, and is characterized by development of an inflammatory mass in the ileum or colon.

- Approximately 15% of patients present with multinodular disease, characterized by abdominal pain, fever, vomiting and mucous diarrhea. ^{1 2}
- 85% of patients develop an intestinal mass adherent to the overlying abdominal wall (helminthoma = Dapaong tumor), often associated with pain and fever. ^{3 4}

This disease is endemic or potentially endemic to 35 countries.

Oesophagostomiasis in Uganda

Sporadic cases of *Oesophagostomum bifurcum* infection have been reported. ⁵

Oesophagostomum bifurcum, *O. stephanostomum* and *Oesophagostomum* spp. have been identified in wild chimpanzees ⁶, mountain gorillas (*Gorilla beringei beringei*), colobus monkeys, guenons and baboons in Uganda. ⁷⁻¹²

References

1. Trans R Soc Trop Med Hyg 2001 May-Jun;95(3):295-9.
2. Trans R Soc Trop Med Hyg 2000 Mar-Apr;94(2):177-82.
3. Clin Infect Dis 2001 Jul 15;33(2):166-70.
4. Gut 1972 Jan ;13(1):8-16.
5. Virchows Arch 2001 Jul ;439(1):21-6.
6. Am J Primatol 2012 Jul ;74(7):642-50.
7. PLoS Negl Trop Dis 2010 ;4(3):e630.
8. J Wildl Dis 2008 Oct ;44(4):878-87.
9. Am J Primatol 2008 Mar ;70(3):222-30.
10. J Parasitol 2005 Jun ;91(3):569-73.
11. J Parasitol 2004 Dec ;90(6):1356-60.
12. Primates 2005 Jan ;46(1):59-63.

Onchocerciasis

Agent	PARASITE - Nematoda. Phasmidea, Filariae: <i>Onchocerca volvulus</i>
Reservoir	Human
Vector	Fly (black fly = <i>Simulium</i>)
Vehicle	None
Incubation Period	12m - 18m
Diagnostic Tests	Identification of microfilariae in skin snips or on ophthalmoscopy. Nucleic acid amplification.
Typical Adult Therapy	Excision of nodules. Ivermectin 150ug/kg PO once. Repeat every 6 months Doxycycline 100 mg PO daily for 6 weeks prior to Ivermectin improves cure rate If eye involved, administer corticosteroid for several days prior to ivermectin .
Typical Pediatric Therapy	Excision of nodules. Ivermectin 150ug/kg PO once. Repeat every 6 months Age > 8 years: Doxycycline , as for adult
Clinical Hints	Macular, papular or dyschromic skin lesions; pruritus; lymphadenopathy; keratitis or uveitis; eosinophilia; firm nodules over bony prominences; adult worms may survive for 15 years in the human host.
Synonyms	Aswad, Craw-craw, Dipetalonema arbuta, Dipetalonema sprenti, Erysipelas de la Costa, Flussblindheit, Jur blindness, Lichenified onchodermatitis, Nakalanga syndrome, <i>Onchocerca cervicalis</i> , <i>Onchocerca dewitiei</i> , <i>Onchocerca guttarosa</i> , <i>Onchocerca jakutensis</i> , <i>Onchocerca lupi</i> , <i>Onchocerca reticulata</i> , <i>Onchocerca volvulus</i> , Onchozerkose, River blindness, Robles' disease, Sowda. ICD9: 125.3 ICD10: B73

Clinical

WHO Case definition for surveillance:

- In an endemic area, a person with fibrous nodules in subcutaneous tissues.
- Laboratory criteria for confirmation • one or more of the following
- Presence of microfilariae in skin snips taken from the iliac crest
- Presence of adult worms in excised nodules
- Presence of typical ocular manifestations, such as slit-lamp observations of microfilariae in the cornea, the anterior chamber, or the vitreous body

Case classification

Suspected: A case that meets the clinical case definition.

Probable: Not applicable.

Confirmed: A suspected case that is laboratory-confirmed.

W.H.O. recognizes five forms of skin disease for purposes of survey and control:

- acute papular onchodermatitis
- chronic papular onchodermatitis
- lichenified onchodermatitis
- atrophy
- depigmentation

The distribution of dermal nodules is related to body regions which are exposed to vector bites (ie, local clothing practices). ¹

The microfilariae of *Onchocerca* migrate throughout the body and give rise to visual impairment (punctate keratitis) ², rashes, intense pruritis and depigmentation of the skin ³; lymphadenitis; "hanging groin" and elephantiasis of the genitals. ⁴

- Rare instance of eosinophilic meningitis have been reported. ⁵
- Some studies have suggested a possible association between onchocerciasis "nodding syndrome" and epilepsy; however data are inconclusive. ⁶⁻²¹

Onchocerciasis has been implicated in the etiology of Nakalanga syndrome (hyposexual dwarfism) in Sudan; and sowda (a form of endemic filarial limb dermatosis with adenopathy) on the Arabian Peninsula. ²²

- It has been suggested that sowda may be caused by a zoonotic species rather than *Onchocerca volvulus*.

Dermal onchocerciasis may mimic dracunculiasis. ²³

There is extensive evidence that endosymbiont bacteria (*Wolbachia* spp.) are necessary for the development of filarial larvae, and fertility of adult parasites. ²⁴⁻²⁷

- Doxycycline has proven effective in therapy, presumably through inhibition of *Wolbachia* spp. ²⁸⁻³⁰

This disease is endemic or potentially endemic to 36 countries.

Onchocerciasis in Uganda

Time and Place:

Onchocerciasis is considered the second most common parasitic disease in Uganda (ie, after malaria).

- The disease is endemic to 17 of the country's 39 districts: Apac, Arua, Bushenyi ³¹, Gulu, Hoima ³², Kabale, Kabarole ³³, Kasese, Kibale, Kitgum, Kisoro, Masindi, Mbale, Mbarara, Moyo, Nebbi ³⁴ and Rukungiri.
- 1.8 million persons live in areas of risk. Highest rates are registered at Mt. Eglon.


30,000 were infested (and 800 blind) in 1987; 1,200,000 in 1993 (this figure does not include the estimated 100,000 infested refugees residing in the country).

Prevalence surveys:

19% in Nebbi (onchocercal skin disease, 1994)

33.89% of persons in Kasese, Kisoro, Kanangu, Moyo and Nebbi in 1993; 3.1% in 2006 (microfilaremia) ³⁵


Onchocerciasis has been implicated in the etiology of Nakalanga syndrome (hyposexual dwarfism) ³⁶ and epilepsy in this country. ^{37 38}



Graph: Uganda. Onchocerciasis - treatments administered, cases

Notes:

1. Additional references: 2007 ³⁹ 2008 ⁴⁰
2. Community-directed treatment with ivermectin was launched in Uganda in 1997. ⁴¹
3. Greater than 70% of Ugandans have received an annual dose of ivermectin for at least 10 years (2007 publication) ⁴²



Graph: Uganda. Onchocerciasis, treatments administered (% coverage)

Vectors:

- The local vectors are *Simulium damnosum* (along the lower Victoria Nile and Ruwenzori mountains) and *S. naevi*⁴³ - the latter accounting for over 85% of transmission.

- *S. kilibanum* may also act as a vector south of the Ruwenzori focus.⁴⁴

Notable outbreaks:

2009 (publication year) - An outbreak of onchocerciasis was reported in Northern Uganda.⁴⁵

References

1. J Vector Borne Dis 2012 Sept;49(3):140-142.
2. Ophthalmol Clin North Am 2002 Sep ;15(3):351-6.
3. Ann Trop Med Parasitol 2006 Dec ;100(8):733-46.
4. BMJ 2003 Jan 25;326(7382):207-10.
5. Clin Microbiol Rev 2009 Apr ;22(2):322-48, Table of Contents.
6. East Afr Med J 1992 Oct ;69(10):554-6.
7. Bull World Health Organ 1996 ;74(4):361-7.
8. Trans R Soc Trop Med Hyg 1997 Sep-Oct;91(5):525-7.
9. Am J Epidemiol 1999 Mar 15;149(6):565-70.
10. Trans R Soc Trop Med Hyg 2002 Sep-Oct;96(5):537-41.
11. Neuroepidemiology 2004 May-Jun;23(3):144-9.
12. Trends Parasitol 2006 Jan ;22(1):17-20.
13. Trans R Soc Trop Med Hyg 2007 Jan ;101(1):48-55.
14. Am J Trop Med Hyg 2008 Mar ;78(3):400-1.
15. Epilepsia 2008 Dec ;49(12):2008-15.
16. Epilepsia 2009 Oct ;50(10):2325-6.
17. Parasite Immunol 2010 Jan ;32(1):79-80.
18. Parasitology 2010 Sep ;137(10):1559-68.
19. Am J Trop Med Hyg 2011 Aug ;85(2):225-8.
20. Int J Infect Dis 2012 Nov 5;
21. Afr Health Sci 2012 Sep ;12(3):242-8.
22. Trop Geogr Med 1987 Jan ;39(1):73-6.
23. Am J Trop Med Hyg 2010 Dec ;83(6):1348-51.
24. Am J Trop Med Hyg 2005 Aug ;73(2):354-8.
25. Microbes Infect 2004 Jan ;6(1):113-28.
26. Cell Microbiol 2004 Feb ;6(2):97-104.
27. Cell Microbiol 2012 Dec 4;
28. J Infect Dis 2005 Oct 15;192(8):1483-93.
29. Lancet 2005 Jun 18-24;365(9477):2067-8.
30. Microbes Infect 2003 Apr ;5(4):261-73.
31. East Afr Med J 2004 Feb ;81(2):92-6.
32. Ann Trop Med Parasitol 2005 Mar ;99(2):165-72.
33. Trop Med Parasitol 1992 Jun ;43(2):80-2.
34. East Afr Med J 1995 May ;72(5):295-8.
35. Trop Med Int Health 2008 Sep ;13(9):1196-203.
36. Am J Trop Med Hyg 1996 Jan ;54(1):80-3.
37. Bull Soc Pathol Exot 2000 Nov ;93(4):255-9.
38. Epilepsy Res 1998 May ;30(3):247-51.
39. Wkly Epidemiol Rec 2008 Aug 22;83(34):307-12.
40. Wkly Epidemiol Rec 2010 Jan 22;85(4):23-8.
41. Ann Trop Med Parasitol 2005 Mar ;99(2):165-72.
42. Ann Trop Med Parasitol 2007 Jun ;101(4):323-33.
43. East Afr Med J 2004 May ;81(5):244-7.
44. Trop Med Int Health 1999 Dec ;4(12):819-26.
45. CMAJ 2009 Jul 7;181(1-2):E4.

Orbital and eye infections

Agent	BACTERIUM OR FUNGUS. Streptococcus pyogenes , oral anaerobes, <i>Aspergillus</i> spp., facultative gram-negative bacilli, et al
Reservoir	Endogenous Introduced flora (trauma, surgery)
Vector	None
Vehicle	Trauma Surgery Contiguous (sinusitis) Hematogenous
Incubation Period	Variable
Diagnostic Tests	Imaging techniques (CT or MRI). Culture of aspirates or surgical material.
Typical Adult Therapy	Local and systemic antimicrobial agents appropriate for species and severity
Typical Pediatric Therapy	As for adult
Clinical Hints	Proptosis, chemosis, extraocular palsy, or hypopyon associated with sinusitis, bacteremia, eye trauma or surgery. Involves the eye (endophthalmitis); periosteum (periorbital infection); orbit (orbital cellulitis); orbit + eye (panophthalmitis).
Synonyms	Bacterial keratitis, Ceratite, Cheratite, Endophthalmitis, Eye infection, Keratite, Keratitis, Orbital infection, Panophthalmitis, Queratitis. ICD9: 360.0 ICD10: H05.0

Clinical

Endophthalmitis involves the ocular cavity and adjacent structures. ^{1 2}

- Infection may occur in the setting of endocarditis or other bacteremic infections, or follow surgery or penetrating trauma.
- The onset of fungal endophthalmitis is more gradual than infection due to bacteria.
- Several species of parasites (ie, *Toxoplasma*, *Toxocara*, *Onchocerca*, etc) and viruses (CMV, Herpes simplex, measles) may also infect a variety of orbital structures, and are discussed elsewhere in this module.

Panophthalmitis involves all ocular tissue layers, including the episclera. ^{3 4}

- Pain on eye movement is prominent.

Orbital cellulitis is an acute infection of the orbital contents.

- Infection can easily spread to the cavernous sinuses.
- The most common sources for infection are the paranasal sinuses (most commonly ethmoid in children).
- Fever, lid edema, orbital pain, proptosis and limited motion of the globe are important symptoms.

Keratitis can be caused by viruses (Herpes simplex, zoster, smallpox), bacteria, fungi, protozoa (*Acanthamoeba*) or helminths (*Onchocerca volvulus*)

- Microbial keratitis complicating orthokeratology is mainly caused by *P. aeruginosa* or *Acanthamoeba* ⁵

This disease is endemic or potentially endemic to all countries.

References

1. Clin Infect Dis 1995 Sep ;21(3):479-86; quiz 487-8.
2. N Engl J Med 1978 Jul 6;299(1):28-31.
3. Clin Infect Dis 1995 Sep ;21(3):479-86; quiz 487-8.
4. N Engl J Med 1978 Jul 6;299(1):28-31.
5. Eye Contact Lens 2007 Nov ;33(6 Pt 2):373-7; discussion 382.

Orf

Agent	VIRUS - DNA. Poxviridae, Parapoxvirus: Orf virus
Reservoir	Sheep Goat Reindeer Musk ox
Vector	None
Vehicle	Contact Infected secretions Fomite Cat-scratch (rare)
Incubation Period	3d - 6d (range 2d - 7d)
Diagnostic Tests	Viral culture (skin lesion or exudate). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Skin pustule or ulcer following contact with sheep or goats; most lesions limited to finger or hand; heals without scarring within 6 weeks.
Synonyms	Contagious pustular dermatitis, Ecthyma contagiosum, Ovine pustular dermatitis. ICD9: 078.89 ICD10: B08.0

Clinical

Human infection is milder than that of sheep, and usually limited to indolent vesicles and pustules on the hands. ^{1 2}

- Pustules may attain a size of 1 to 2 cm, and are often associated with low-grade fever and regional lymphadenitis.
- Lesions heal over a period of 2 to 6 weeks, without scarring.
- Bullous lesions ³, secondary bacterial infection, disseminated orf, Guillain-Barre syndrome ⁴ and erythema multiforme ⁵⁻¹⁰ have been described in some cases.

This disease is endemic or potentially endemic to all countries.

Orf in Uganda

Notable outbreaks:

2011 - An outbreak (200 cases) of contagious ecthyma was reported among goats in Nwoya District. ¹¹

References

1. Br J Plast Surg 1993 Sep ;46(6):532-4.
2. Cleve Clin J Med 1991 Nov-Dec;58(6):531-4.
3. J Am Acad Dermatol 2008 Jan ;58(1):49-55.
4. Australas J Dermatol 2011 Feb ;52(1):62-5.
5. Int J Dermatol 2008 Dec ;47(12):1333-4.
6. J Eur Acad Dermatol Venereol 2006 May ;20(5):612-3.
7. Ann Dermatol Venereol 2000 Apr ;127(4):397-9.
8. Pediatr Dermatol 1997 Mar-Apr;14(2):154-5.
9. Tidsskr Nor Laegeforen 1984 May 20;104(14):978-9.
10. Cutis 1983 Mar ;31(3):334-8.
11. ProMED <promedmail.org> archive: 20111220.3641

Ornithosis

Agent	BACTERIUM. Chlamydiaceae, Chlamydiae , Chlamydophila [Chlamydia] psittaci
Reservoir	Parakeet Parrot Pigeon Turkey Duck Cat Sheep Goat Cattle ? Dog
Vector	None
Vehicle	Bird droppings Dust Air Aerosol from cat [rare]
Incubation Period	7d - 14d (range 4d - 28d)
Diagnostic Tests	Serology. Culture (available in special laboratories) rarely indicated.
Typical Adult Therapy	Doxycycline 100 mg PO BID X 10d. Alternatives: Erythromycin 500 mg PO QID X 10d. Azithromycin 1 g, then 0.5 g daily. Clarithromycin 0.5 g BID
Typical Pediatric Therapy	Age < 8 years: Erythromycin 10 mg/kg QID X 10d Age >=8 years: Doxycycline 100 mg PO BID X 10d.
Clinical Hints	Headache, myalgia and pneumonia, often with relative bradycardia, hepatomegaly or splenomegaly; onset 1 to 4 weeks following contact with pigeons, psittacine birds or domestic fowl; case-fatality rate without treatment = 20%.
Synonyms	Chlamydophila abortus, Chlamydophila psittaci, Ornitose, Papegojsjuka, Parrot fever, Psitacosis, Psittacosis, Psittakose. ICD9: 073 ICD10: A70

Clinical

Onset may be insidious or abrupt, and the illness may be subclinical, or take the form of nonspecific fever and malaise, pharyngitis, hepatosplenomegaly, and adenopathy. ¹

- Bradycardia and splenomegaly may suggest typhoid at this stage.

A more common presentation consists of atypical pneumonia, with nonproductive cough, fever, headache and pulmonary infiltrates. ^{2 3}

- Additional findings may include photophobia, tinnitus, ataxia, deafness, anorexia, vomiting, abdominal pain ⁴, diarrhea, constipation, hemoptysis, epistaxis, arthralgia, and rash (Horder's spots) reminiscent of the rose spots of typhoid. ⁵
- Fever, pharyngitis, rales and hepatomegaly are noted in over 50% of cases.

Complications include pericarditis, myocarditis, and "culture-negative" endocarditis, ARDS ⁶, overt hepatitis, hemolytic anemia, DIC, reactive arthritis, cranial nerve palsy, cerebellar dysfunction, transverse myelitis, meningitis, encephalitis and seizures, thrombophlebitis, pancreatitis and thyroiditis.

- Subclinical infection by *Chlamydophila psittaci* has been implicated in the etiology of chronic polyarthritis. ⁷
- Rare instances of abortion have been reported.

Chlamydophila abortus, a related species which affects goats, cattle and sheep, had been associated with rare instances of abortion, stillbirth and even maternal death in humans.

This disease is endemic or potentially endemic to all countries.

References

1. Aust Fam Physician 2001 Aug ;30(8):739-41.
2. Semin Respir Infect 1997 Mar ;12(1):7-11.
3. Acta Clin Belg 2010 May-Jun;65(3):192-6.
4. J Med Microbiol 2011 Apr ;60(Pt 4):547-9.

5. J Infect 1990 Nov ;21(3):251-9.
6. Nihon Kokyuki Gakkai Zasshi 2007 May ;45(5):419-23.
7. Clin Exp Rheumatol 2011 Nov-Dec;29(6):977-82.

Osteomyelitis

Agent	BACTERIUM OR FUNGUS. <i>Staphylococcus aureus</i> , facultative gram-negative bacilli, <i>Candida albicans</i> , etc
Reservoir	Endogenous Introduced flora (trauma, surgery)
Vector	None
Vehicle	Trauma Hematogenous Extension from other focus
Incubation Period	Variable
Diagnostic Tests	Radiography, including bone scan. Culture of biopsy material.
Typical Adult Therapy	Systemic antimicrobial agent(s) appropriate to known or suspected pathogen. Surgery as indicated
Typical Pediatric Therapy	As for adult
Clinical Hints	Limb pain or gait disturbance; obscure fever; prior skin infection; may be hematogenous, or arise from contiguous (soft tissue, joint) infection; X-ray changes are not apparent for at least 10 days in acute infection.
Synonyms	Osteomyelitis, Osteomyelitis, Osteomyelitis, Paravertebral abscess. ICD9: 015,730.9 ICD10: M86

Clinical

Osteomyelitis is a self-defined condition characterized by infection of one or more bones.

- Signs and symptoms vary widely, and reflect associated underlying conditions, infecting species and location of the infection. ¹⁻³

Etiological associations:

- Animal bite: *Pasteurella multocida*
- Diabetes and vascular insufficiency: Usually mixed infection (*Staphylococcus aureus*, *Staphylococcus epidermidis*, Gram-negative bacilli, Anaerobes)
- Hematogenous: Usually single organism (*Staphylococcus aureus*, Enterobacteriaceae)
- Injecting drug user: staphylococci, Gram-negative bacilli, *Candida* spp.
- Secondary to contiguous infection: Often mixed infection (*Staphylococcus aureus*, Gram-negative bacilli)
- Sick cell anemia: *Staphylococcus aureus*, *Salmonella* spp.

This disease is endemic or potentially endemic to all countries.

Osteomyelitis in Uganda

Osteomyelitis cases accounted for 10% of outpatients (5 hospitals, 2006 to 2007) ⁴

References

1. J Paediatr Child Health 2005 Jan-Feb;41(1-2):59-62.
2. Skull Base 2009 Jul ;19(4):247-54.

3. Spine (Phila Pa 1976) 2010 Nov 1;35(23):E1317-21.
4. Trans R Soc Trop Med Hyg 2010 Feb ;104(2):139-42.

Otitis media

Agent	BACTERIUM OR VIRUS. Haemophilus influenzae & Streptococcus pneumoniae in most acute cases; RSV, Parainfluenza, et al
Reservoir	Human
Vector	None
Vehicle	None
Incubation Period	Variable
Diagnostic Tests	Clinical findings. Culture of middle ear fluid if available.
Typical Adult Therapy	Antimicrobial agent directed at likely pathogens
Typical Pediatric Therapy	As for adult
Vaccine	Pneumococcal conjugate
Clinical Hints	Acute bacterial otitis media often represents the final stage in a complex of anatomic, allergic or viral disorders of the upper airways; recurrent or resistant infections may require surgical intervention.
Synonyms	Otitis media aguda. ICD9: 382.0 ICD10: H65,H66

Clinical

Signs and symptoms of otitis media consist of local pain and tenderness, with or without fever and signs of sepsis. ¹ ²

This disease is endemic or potentially endemic to all countries.

References

1. Laryngoscope 2004 Nov ;114(11 Pt 3 Suppl 105):1-26.
2. JAMA 2003 Sep 24;290(12):1633-40.

Parainfluenza virus infection

Agent	VIRUS - RNA. Paramyxoviridae: Respirovirus - Human Parainfluenza virus 1 and 3. Rubulavirus - Human Parainfluenza virus 2 and 4.
Reservoir	Human
Vector	None
Vehicle	Droplet
Incubation Period	3d - 8d
Diagnostic Tests	Viral culture (respiratory secretions). Serology. Nucleic acid amplification.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Upper respiratory infection - often croup or laryngitis. The disease is most common during infancy; older children develop a 'cold-like' illness; the infection is complicated by pneumonia in 7% to 17% of cases.
Synonyms	Parainfluenza, Sendai. ICD9: 078.89,480.2 ICD10: J12.2

Clinical

Clinical forms of Parainfluenza virus infection include 'the common cold,' otitis media, croup (acute laryngotracheobronchitis) ¹, 'flu-like illness' ², bronchiolitis ³ and pneumonia.

This disease is endemic or potentially endemic to all countries.

References

1. J Pediatr Health Care 2004 Nov-Dec;18(6):297-301.
2. J Med Virol 2009 Dec ;81(12):2066-71.
3. Curr Opin Pulm Med 2002 Mar ;8(2):112-6.

Parvovirus B19 infection

Agent	VIRUS - DNA. Parvoviridae, Parvovirinae: Erythrovirus B19
Reservoir	Human
Vector	None
Vehicle	Droplet
Incubation Period	4d - 14d (range 3d - 21d)
Diagnostic Tests	Serology. Nucleic acid amplification (testing should be reserved for the rare instance of complicated infection).
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Erythema infectiosum (erythema of cheeks; lacelike or morbilliform rash on extremities); febrile polyarthralgia, or bone marrow aplasia/hypoplasia may be present.
Synonyms	Duke's disease, Erythema infantum febrile, Erythema infectiosum, Erythema simplex marginatum, Erythrovirus B19, Fifth disease, Fourth disease, Funfte Krankheit, Parascarlantina, Parvovirus 4, Parvovirus B19, Sticker's disease. ICD9: 057.0 ICD10: B08.3

Clinical

Acute infection:

Erythema infectiosum is a mild childhood illness characterized by a facial rash ("slapped cheek" appearance), and a reticulated or lacelike rash on the trunk and extremities. ¹

- Papular-purpuric gloves-and-socks syndrome ², or localized and generalized petechial rashes may occur in some cases. ³⁻⁸
- Reappearance of the rash may occur for several weeks following nonspecific stimuli such as change in temperature, sunlight, and emotional stress.
- The patient is otherwise well at rash onset but often gives a history of a systemic prodrome lasting 1 to 4 days.
- In some outbreaks, pruritis has been a common clinical feature. ⁹
- Rubella-like, morbilliform ¹⁰, vesicular and purpuric ¹¹ rashes have also been reported.
- Asymptomatic infection has been reported in approximately 20% of children and adults.
- Rare instances of hepatosplenomegaly ¹² and heart failure have been reported. ¹³
- Co-infection with parvovirus and other hepatitis viruses may result in fulminant hepatic failure ¹⁴

Joint manifestations:

- In some outbreaks, arthralgias and arthritis have been commonly reported. ¹⁵
- Infection may produce a symmetrical peripheral polyarthropathy.
 - The hands are most frequently affected, followed by the knees and wrists.
 - Symptoms are usually self-limited but may persist for several months.
 - Joint symptoms, more common in adults, are encountered in approximately 20% of cases ¹⁶ and may occur as the sole manifestation of infection.

Instances of seizure ¹⁷, coma, encephalitic ataxia or chorea ^{18 19}, meningoencephalitis ²⁰, autonomic or sensory neuropathy ²¹, cranial nerve palsy ²², myocarditis ^{23 24}, severe endothelialitis (Degos-like syndrome) ²⁵, myositis ²⁶, and acute hepatitis have been reported. ²⁷⁻³⁰

- Sequelae remain in 22% of patients with neurological involvement ³¹
- A distinct form of Parvovirus infection known as "papular-purpuric gloves and socks syndrome" is characterized by fever and edematous rash, often associated with conjunctivitis and arthritis ³²
- Additional complications may include glomerulonephritis ^{33 34}, Melkersson-Rosenthal syndrome and hemophagocytic lymphohistiocytosis ^{35 36}
- Hepatic dysfunction may be present in some cases. ³⁷

Parvovirus B19 infection and hematological disease:

Parvovirus B19 is the primary etiologic agent causing Transient Aplastic Crisis (TAC) in patients with chronic hemolytic

anemias (e.g., sickle cell disease, hemoglobin SC disease, hereditary spherocytosis, alpha-thalassemia, and autoimmune hemolytic anemia) and occasionally follows anemia due to blood loss. ³⁸

- Patients with TAC typically present with pallor, weakness, and lethargy and may report a nonspecific prodromal illness during the preceding 1 to 7 days.
- Few patients with TAC report a rash.
- In the acute phase, patients usually have a moderate to severe anemia with absence of reticulocytes; and bone marrow examination shows a hypoplastic or an aplastic erythroid series with a normal myeloid series.
- Recovery is indicated by a return of reticulocytes in the peripheral smear approximately 7 to 10 days after their disappearance.
- TAC may require transfusion and hospitalization and can be fatal if not treated promptly.

A false positive serological reaction toward Epstein-Barr virus has been reported in Parvovirus B19 infection. ³⁹

A Parvovirus B19-related severe chronic anemia associated with red cell aplasia has been described in transplant recipients ⁴⁰, patients on maintenance chemotherapy for acute lymphocytic leukemia, patients with congenital immunodeficiencies, and patients with human immunodeficiency virus (HIV)-related immunodeficiency. ⁴¹

Infection of the intestinal mucosa may produce symptoms of inflammatory bowel disease. ⁴²

Intrapartum infections:

Intrauterine infections can lead to specific or permanent organ defects in the fetus (e.g. heart anomalies, eye diseases, micrognathia, chronic anemia, myocarditis, hepatitis, meconium peritonitis and central nervous system anomalies). ⁴³⁻⁴⁵

- Thrombocytopenia is reported in 46% of cases ⁴⁶
- Rare cases of transient neonatal leukoerythroblastosis have been reported ⁴⁷
- In most reported B19 infections occurring during pregnancy, the fetus has not been adversely affected; however, in some cases B19 infection has been associated with fetal death. ⁴⁸⁻⁵⁰
- The risk of fetal death attributable to maternal parvovirus infection is estimated at less than 10%.
- Fetal death most commonly occurs from the 10th through the 20th weeks of pregnancy.
- Although maternal infection appears to be common in late pregnancy, hydrops is relatively rare. ⁵¹

A related member of the family Parvovirinae, Human Bocavirus, is discussed under 'Respiratory viruses • miscellaneous'

This disease is endemic or potentially endemic to all countries.

References

1. Int J Dermatol 2004 Oct ;43(10):747-9.
2. J Clin Virol 2011 Nov ;52(3):269-71.
3. Pediatrics 2010 Apr ;125(4):e787-92.
4. Int J Dermatol 2008 Jul ;47(7):760-2.
5. Clin Pediatr (Phila) 2006 Apr ;45(3):275-80.
6. New Microbiol 2006 Jan ;29(1):45-8.
7. J Am Acad Dermatol 2005 May ;52(5 Suppl 1):S109-13.
8. Pediatr Dermatol 1998 Jan-Feb;15(1):35-7.
9. J R Coll Gen Pract 1987 May ;37(298):210-1.
10. Rev Soc Bras Med Trop 2008 Jul-Aug;41(4):338-44.
11. Ann Dermatol Venereol 2010 Nov ;137(11):709-12.
12. J Trop Med 2011 ;2011:287914.
13. Clin Exp Dermatol 2008 Aug ;33(5):588-90.
14. Pediatr Infect Dis J 2009 Jul ;28(7):649-50.
15. Clin Perinatol 2005 Sep ;32(3):697-704.
16. Clin Rheumatol 2009 Sep ;28(9):1067-71.
17. J Trop Med 2011 ;2011:287914.
18. J Child Neurol 2008 Sep ;23(9):1078-80.
19. Ann Trop Paediatr 2010 ;30(4):339-44.
20. Clin Infect Dis 2008 Aug 1;47(3):385-7.
21. Brain Dev 2011 Feb ;33(2):161-5.
22. Eur J Ophthalmol 2010 Jul-Aug;20(4):802-4.
23. N Engl J Med 2010 Apr 1;362(13):1248-9.
24. J Clin Virol 2011 Jan ;50(1):61-4.
25. J Cutan Pathol 2008 Oct ;35 Suppl 1:20-5.
26. Case Report Rheumatol 2012 ;2012:250537.
27. Ugeskr Laeger 2007 Nov 19;169(47):4075-7.
28. BMC Infect Dis 2010 ;10:246.
29. Scand J Infect Dis 2011 Jul ;43(6-7):547-9.
30. J Clin Microbiol 2011 Sep ;49(9):3422-4.
31. Clin Infect Dis 2009 Jun 15;48(12):1713-23.
32. J Am Acad Dermatol 2009 Apr ;60(4):691-5.
33. PMID 19735054
34. Ren Fail 2012 Nov 1;
35. Clin Exp Dermatol 2009 Dec ;34(8):e623-5.
36. APMIS 2009 Oct ;117(10):773-7.
37. Ugeskr Laeger 1998 Oct 26;160(44):6355-6.
38. Rev Med Virol 2003 Nov-Dec;13(6):347-59.
39. Clin Vaccine Immunol 2009 Mar ;16(3):372-5.
40. Clin Infect Dis 2006 Jul 1;43(1):40-8.
41. Arch Pathol Lab Med 2007 May ;131(5):799-804.
42. J Clin Microbiol 2009 May ;47(5):1591-5.
43. Z Geburtshilfe Neonatol 2007 Apr ;211(2):60-8.
44. Infect Dis Obstet Gynecol 2008 ;2008:524601.
45. N Engl J Med 1987 Jan 22;316(4):183-6.
46. BJOG 2008 Jan ;115(1):76-81.
47. Int J Infect Dis 2009 Nov ;13(6):e473-5.
48. Infect Dis Obstet Gynecol 2003 ;11(3):175-9.
49. J Clin Virol 2006 May ;36(1):1-7.
50. Reprod Toxicol 2006 May ;21(4):421-35.
51. Fetal Diagn Ther 2011 ;30(1):41-7.

Pediculosis

Agent	PARASITE - Insecta. Anoplura: <i>Pediculus humanus</i> , <i>Phthirus pubis</i> .
Reservoir	Human
Vector	Louse
Vehicle	Contact
Incubation Period	7d
Diagnostic Tests	Identification of adults and "nits."
Typical Adult Therapy	Permethrin 1%; or malathion 0.5%; or lindane OR Ivermectin 200 mcg/kg PO
Typical Pediatric Therapy	Permethrin 1%; or malathion 0.5%; or lindane OR Ivermectin 200 mcg/kg PO (> 15 kg body weight)
Clinical Hints	Pruritus in the setting of poor personal hygiene; adults or nits may be visible; note that the body louse (<i>Pediculus humanus</i> var. corporis; not the head louse) transmits diseases such as epidemic typhus, trench fever and relapsing fever.
Synonyms	Crab louse, Lousebefall, Pediculose, <i>Pediculus capitus</i> , <i>Pediculus corporis</i> , Pedikulose, <i>Phthirus pubis</i> , Pidocci. ICD9: 132 ICD10: B85

Clinical

Most louse infestations are asymptomatic, with only 15% to 36% of patients complaining of pruritus.

- The principal clinical finding consists of presence of the lice themselves, and their eggs ('nits'). ^{1 2}

This disease is endemic or potentially endemic to all countries.

References

1. *J Am Acad Dermatol* 2004 Jun ;50(6):819-42, quiz 842-4.
2. *N Engl J Med* 2002 May 23;346(21):1645-50.

Pentastomiasis - Linguatula

Agent	PARASITE - Pentastomid worm. <i>Linguatula serrata</i>
Reservoir	Herbivore
Vector	None
Vehicle	Meat (liver or lymph nodes of sheep/goat)
Incubation Period	Unknown
Diagnostic Tests	Identification of larvae in nasal discharge.
Typical Adult Therapy	No specific therapy available
Typical Pediatric Therapy	As for adult
Clinical Hints	Pharyngeal or otic itching, cough, rhinitis or nasopharyngitis which follows ingestion of undercooked liver.
Synonyms	Linguatula, Marrara syndrome. ICD9: 128.8 ICD10: B83.8

Clinical

Infestation ("halzoun" or "marrara syndrome") is associated with pain and itching in the throat or ear, lacrimation, cough, hemoptysis, rhinorrhea or hoarseness. ^{1 2} (Halzoun is also associated with infection by *Dicrocoelium dendriticum*) ³

- Complications include respiratory obstruction, epistaxis, facial paralysis or involvement of the eye.

This disease is endemic or potentially endemic to 184 countries.

References

1. Rev Infect Dis 1987 Nov-Dec;9(6):1087-94.
2. Acta Trop 1996 Dec 16;62(3):127-34.
3. Acta Trop 2013 Jan ;125(1):115-8.

Pericarditis - bacterial

Agent	BACTERIUM. Streptococcus pneumoniae , Staphylococcus aureus , et al
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Ultrasonography and cardiac imaging techniques. Culture of pericardial fluid (include mycobacterial culture).
Typical Adult Therapy	Antimicrobial agent(s) appropriate to known or anticipated pathogen. Drainage as indicated
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, chest pain and dyspnea; patients are acutely ill and have overt signs such as venous distention, and an enlarged cardiac 'shadow'; concurrent pneumonia or upper respiratory infection may be present; case-fatality rate = 20%.
Synonyms	Bacterial pericarditis, Pericardite. ICD9: 074.23,074.2,115.03,420 ICD10: I30

Clinical

Viral pericarditis often follows a prodrome of upper respiratory infection.

- Typical findings include fever and chest pain. ^{1 2}
- The pain may be pleuritic or positional (ie, exacerbated by bending forward) and associated with signs and symptoms of congestive heart failure.
- Concurrent myocarditis, pneumonia or pleuritis are often present.

This disease is endemic or potentially endemic to all countries.

References

1. [N Engl J Med 2004 Nov 18;351\(21\):2195-202.](#)
2. [Lancet 2004 Feb 28;363\(9410\):717-27.](#)

Perinephric abscess

Agent	BACTERIUM OR FUNGUS. Escherichia coli , other facultative gram negative bacilli, Candida albicans , et al
Reservoir	Human
Vector	None
Vehicle	None
Incubation Period	Variable
Diagnostic Tests	Urine and blood culture. Renal imaging (CT, etc).
Typical Adult Therapy	Antimicrobial agent(s) appropriate to known or anticipated pathogen. Surgery as indicated
Typical Pediatric Therapy	As for adult
Clinical Hints	Unexplained fever, leukocytosis and flank pain; patients are typically over age 50, often diabetic; consider in the patient with nonresponsive 'pyelonephritis' or a renal mass (by examination or x-ray).
Synonyms	

Clinical

Symptoms may be overt or subtle, and limited to unexplained fever; indeed, 33% of such lesions are first diagnosed at autopsy.

- Typical patients are female and over the age of 50. [1-3](#)
- Diabetes and evidence for preceding or current urinary tract infection or bacteremia (including endocarditis) may be present.

This disease is endemic or potentially endemic to all countries.

References

1. [Med Clin North Am 1988 Sep ;72\(5\):993-1014.](#)
2. [Infect Dis Clin North Am 1987 Dec ;1\(4\):907-26.](#)
3. [Infect Dis Clin North Am 1997 Sep ;11\(3\):663-80.](#)

Perirectal abscess

Agent	BACTERIUM. Various (often mixed anaerobic and aerobic flora)
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Culture of drainage material.
Typical Adult Therapy	Surgical drainage and antibiotics effective against fecal flora
Typical Pediatric Therapy	As for adult
Clinical Hints	Anal or perianal pain with fever and a tender mass suggest this diagnosis; granulocytopenic patients commonly develop small, soft and less overt abscesses - often due to <i>Pseudomonas aeruginosa</i> .
Synonyms	

Clinical

Perirectal abscess is a self-defined illness usually associated with overt local pain, swelling, tenderness and fluctuance. ¹

- Abscesses in neutropenic patients are often more subtle, and may present as unexplained fever without marked local findings.

This disease is endemic or potentially endemic to all countries.

References

1. [Ann Emerg Med 1995 May ;25\(5\):597-603.](#)

Peritonitis - bacterial

Agent	BACTERIUM. Various (often mixed anaerobic and aerobic flora)
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Culture of blood and peritoneal fluid. Peritoneal fluid cell count may also be useful.
Typical Adult Therapy	Antimicrobial agent(s) appropriate to known or anticipated pathogens. Surgery as indicated
Typical Pediatric Therapy	As for adult
Clinical Hints	Abdominal pain and tenderness, vomiting, absent bowel sounds, guarding and rebound; diarrhea may be present in children; search for cause: visceral infection or perforation, trauma, underlying cirrhosis (spontaneous peritonitis) etc.
Synonyms	Acute peritonitis, Bacterial peritonitis, Peritonite. ICD9: 567 ICD10: K65

Clinical

Bacterial peritonitis following trauma, infection or perforation of an abdominal viscus is usually overt clinically. ¹

Spontaneous bacterial peritonitis is somewhat more subtle, and should be suspected when unexplained deterioration occurs in a patient with ascites or chronic liver disease. ^{2 3}

- As many as 30% of patients are asymptomatic, and the remainder present with fever, chills, abdominal pain, diarrhea, increasing ascites, encephalopathy or renal dysfunction.
- Abdominal tenderness, guarding and hypotension may be present.
- Bacteremia is a poor prognostic factor in these patients. ⁴

This disease is endemic or potentially endemic to all countries.

References

1. Am J Surg 2003 Nov 28;186(5A):15S-22S; discussion 31S-34S.
2. Eur J Clin Microbiol Infect Dis 1998 Aug ;17(8):542-50.
3. Semin Liver Dis 1997 ;17(3):203-17.
4. Scand J Infect Dis 2007 ;39(8):697-702.

Pertussis

Agent	BACTERIUM. Bordetella pertussis An aerobic gram-negative coccobacillus
Reservoir	Human
Vector	None
Vehicle	Air Infected secretions
Incubation Period	7d - 10d (range 5d - 21d)
Diagnostic Tests	Culture & direct fluorescence (nasopharynx). Alert laboratory when suspected. Serology.
Typical Adult Therapy	Respiratory precautions. Erythromycin 500 mg QID X 10d. Alternatives: Azithromycin , Clarithromycin
Typical Pediatric Therapy	Respiratory precautions: Erythromycin 10 mg/kg QID X 10d. Alternatives: Azithromycin , Clarithromycin
Vaccines	DTaP DTP
Clinical Hints	Coryza, paroxysmal cough, occasional pneumonia or otitis; lymphocytosis; most often diagnosed in young children; epistaxis and subconjunctival hemorrhage often noted; seizures (below age 2); case-fatality rate = 0.5%.
Synonyms	Bordetella holmesii , Bordetella parapertussis , Bordetella pertussis , Coqueluche, Keuchhusten, Kikhosta, Kikhoste, Kinkhoest, Parapertussis, Pertosse, Syndrome coqueluchoide, Tos convulsa, Tos farina, Tosse convulsa, Tussis convulsa, Whooping cough. ICD9: 033 ICD10: A37

Clinical

WHO Case definition for surveillance: ¹⁻³

Clinical case definition

A person with a cough lasting at least 2 weeks with at least one of the following:

- paroxysms (i.e. fits) of coughing
- inspiratory .whooping.
- post-tussive vomiting (i.e. vomiting immediately after coughing)
- without other apparent cause

Laboratory criteria for diagnosis

- Isolation of *Bordetella pertussis*, or
- Detection of genomic sequences by polymerase chain reaction (PCR)

Case classification

- Suspected: A case that meets the clinical case definition.
- Confirmed: A person with a cough that is laboratory-confirmed.

Acute illness:

Following an incubation period of 7 to 10 days (range 6 to 20) the patient develops coryza and cough (the catarrhal stage).

- After one to two weeks, the cough progresses into the paroxysmal stage. ^{4 5}
- Post-tussive vomiting is common, and young children and older infants may exhibit an inspiratory "whoop."
- Among infants younger than six months, apnea is common and the whoop may be absent. ⁶
- The paroxysmal stage lasts three to four weeks (range one to six).
- The convalescent stage lasts for two to four weeks.

Complications:

Infants are at increased risk of complications from pertussis, while pertussis among adolescents and adults tends to be milder and may be limited to a persistent cough. ⁷

- Over 70% of infants younger than 6 months require hospitalization.
- Complications of pertussis can include secondary bacterial pneumonia (the most common cause of death in pertussis), seizures and encephalopathy. ⁸
- Other, less serious complications include otitis media and dehydration.
- Severe coughing can lead to pneumothorax, epistaxis, subdural hematoma, acute carotid dissection with stroke ⁹, hernia, and rectal prolapse.
- Pertussis in adults is often characterized by unexplained prolonged cough. ^{10 11}

- Pertussis-RSV infection is common. ¹²
- Rare cases of acute disseminated encephalomyelitis ¹³ and hemolytic-uremic syndrome have been ascribed to pertussis ¹⁴
- Human Bocavirus infection may mimic the symptoms of pertussis ¹⁶

Parapertussis is caused by *Bordetella parapertussis*, and shares many of the clinical features of pertussis.

- 70% of infections are asymptomatic.

This disease is endemic or potentially endemic to all countries.

Pertussis in Uganda

Vaccine Schedule:

BCG - birth

DTwP - 6, 10, 14 weeks


DTwPHibHep - 6, 10, 14 weeks

Measles (monovalent) - 9 months


OPV - birth; 6, 10, 14 weeks

TT - Women of child-bearing age; +1, +6 months; +1, +1 year

Vitamin A - 6, 12, 18, 24, 30, 36, 42 months



Graph: Uganda. Pertussis - WHO-UNICEF est. vaccine (DTP3 %) coverage



Graph: Uganda. Pertussis, cases

References

1. J Public Health Manag Pract 2009 Nov-Dec;15(6):479-84.
2. Int J Infect Dis 2010 Dec ;14(12):e1072-5.
3. Clin Infect Dis 2012 Mar 19;
4. Aust Fam Physician 2004 May ;33(5):317-9.
5. Lancet Infect Dis 2002 Dec ;2(12):744-50.
6. CMAJ 2005 Feb 15;172(4):509-15.
7. Paediatr Respir Rev 2008 Sep ;9(3):201-11; quiz 211-2.
8. J Neurol Sci 2012 Sep 15;320(1-2):145-8.
9. J Stroke Cerebrovasc Dis 2013 Feb 11;
10. N Engl J Med 2005 Mar 24;352(12):1215-22.
11. J Emerg Med 2012 Dec 31;
12. Pediatr Infect Dis J 2007 Apr ;26(4):316-8.
13. Ann Trop Paediatr 2011 ;31(3):269-72.
14. Eur J Clin Microbiol Infect Dis 2006 Aug ;25(8):515-7.
15. Pediatr Nephrol 2010 Jul ;25(7):1361-4.
16. Pediatrics 2008 Mar ;121(3):e631-7.

Pharyngeal & cervical space infx.

Agent	BACTERIUM. <i>Streptococcus pyogenes</i> , mixed oral anaerobes, etc.
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Careful examination of region and X-ray (or CT scan). Smear and culture of pus if available.
Typical Adult Therapy	Surgical drainage and parenteral antibiotics effective against oral flora
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, painful swelling and displacement of the tongue, fauces and other intraoral structures; dysphagia, dyspnea or jugular phlebitis may ensue in more virulent infections.
Synonyms	Cervical space infection, descending necrotizing mediastinitis, Lemmier's syndrome, Ludwig's angina, Post-anginal septicemia, Quinsy. ICD9: 682.0,682.1 ICD10: J36,J39.0,J39.1

Clinical

Signs and symptoms reflect the site of infection: ¹

- masticator, buccal, canine or parotid spaces
- submandibular, submaxillary and submandibular spaces (Ludwig's angina)
- lateral pharyngeal, retropharyngeal or paratracheal spaces
- peritonsillar tissues (quinsy)
- jugular vein (post-anginal septicemia = Lemmiere's syndrome) ^{2 3}

Lemmiere's syndrome is a potentially fatal infection caused by *Fusobacterium necrophorum*.

- The condition is most common among young healthy persons and typically begins with pharyngotonsillitis which spreads to the parapharyngeal spaces to produce septic phlebitis of the internal jugular vein. ⁴⁻⁶
- Submandibular edema and tenderness along the sternocleidomastoid muscle are noted, usually unilaterally.
- After one to two weeks, the patient develops multiple metastatic abscesses of the lungs, muscles ⁷, bones, joints • or rarely, brain.
- Hyperbilirubinemia and mild disseminated intravascular coagulation may be present.
- The case-fatality rate is 4% to 33%, even with appropriate antimicrobial therapy.

This disease is endemic or potentially endemic to all countries.

References

1. Eur Arch Otorhinolaryngol 2009 Feb ;266(2):273-7.
2. South Med J 2003 Sep ;96(9):928-32.
3. Am J Otolaryngol 2003 Mar-Apr;24(2):111-7.
4. Clin Microbiol Rev 2007 Oct ;20(4):622-59.
5. ORL J Otorhinolaryngol Relat Spec 2003 Mar-Apr;65(2):117-20.
6. Medicine (Baltimore) 2002 Nov ;81(6):458-65.
7. Am J Med Sci 2008 Jun ;335(6):499-501.

Pharyngitis - bacterial

Agent	BACTERIUM. Most often Streptococcus pyogenes ; Str. groups B, C, F and G are occasionally isolated
Reservoir	Human
Vector	None
Vehicle	Droplet Rarely food
Incubation Period	1d - 5d
Diagnostic Tests	Throat swab for culture or antigen detection (group A Streptococcus) ASLO titer may not indicate current infection
Typical Adult Therapy	Penicillin G or Penicillin V or other antistreptococcal antibiotic to maintain serum level for 10 days
Typical Pediatric Therapy	As for adult
Clinical Hints	Purulent pharyngitis and cervical lymphadenopathy usually indicate streptococcal etiology; however, viruses (mononucleosis, enteroviruses) and other bacteria (gonorrhea, diphtheria) should also be considered.
Synonyms	Acute pharyngitis, Bacterial pharyngitis, Mal di gola batterica, Oral thrush, Streptococcal pharyngitis, Tonsillitis - bacterial, Vincent's angina. ICD9: 034.0,462 ICD10: J02,J03

Clinical

This is a self-defined condition characterized by erythema and pain in the pharynx, often associated with fever, dysphagia and upper respiratory tract infection. ¹

This disease is endemic or potentially endemic to all countries.

References

1. [Paediatr Drugs 2003 ;5 Suppl 1:13-23.](#)

Pityriasis rosea

Agent	UNKNOWN. Human herpesvirus 7 has been implicated
Reservoir	Unknown
Vector	Unknown
Vehicle	Unknown
Incubation Period	Unknown
Diagnostic Tests	Clinical features.
Typical Adult Therapy	Supportive; ultraviolet B exposure is suggested
Typical Pediatric Therapy	As for adult
Clinical Hints	3 to 8 week illness; herald patch followed by crops of salmon-colored macules and papules; pruritus; systemic symptoms rare.
Synonyms	

Clinical

Pityriasis rosea is a mild exanthem characterized by oval or round macules or papules which evolve following the appearance of a "herald patch" (80% of cases). ¹⁻⁵

- Fine desquamation and pruritus are common.
- Rarely, the condition may recur. ⁶
- In Black patients, Pityriasis rosea may present with facial and scalp involvement, post-inflammatory disorders of pigmentation and papular lesions. ⁷
- Acral lesions ⁸ or dermal follicles may predominate in some cases. ⁹
- The disease should be distinguished from secondary syphilis • the latter characterized by prominent lymphadenopathy; lack of pruritus and herald patch; and accompanying fever and systemic signs. ¹⁰

This disease is endemic or potentially endemic to all countries.

References

1. Australas J Dermatol 2012 Aug ;53(3):e64-5.
2. Curr Opin Pediatr 2009 Aug ;21(4):481-5.
3. Clin Exp Dermatol 2009 Mar ;34(2):269-70.
4. Indian J Dermatol Venereol Leprol 1998 Jul-Aug;64(4):185-6.
5. Dermatologica 1981 ;162(1):64-5.
6. Clin Exp Dermatol 2009 Jul ;34(5):e114-6.
7. Cases J 2009 ;2:6796.
8. Indian Dermatol Online J 2010 Jul ;1(1):21-3.
9. J Dermatol Case Rep 2012 Jun 30;6(2):36-9.
10. J Am Acad Dermatol 1985 Apr ;12(4):597-624.

Plague

Agent	BACTERIUM. <i>Yersinia pestis</i> A facultative gram-negative bacillus
Reservoir	Rodent Rabbit Cat Wild carnivore
Vector	Flea (Pulex; Xenopsylla)
Vehicle	Air Contact
Incubation Period	2d - 7d (range 1d - 14d)
Diagnostic Tests	Culture (blood, sputum, pus). Fluorescent (DFA) staining of pus. Nucleic acid amplification.
Typical Adult Therapy	Strict isolation. Gentamicin 2 mg/kg IV loading dose, then 1.7 mg/kg Q8h. OR Streptomycin 15 mg/kg q12h X 10d. OR Doxycycline 100 mg PO BID X 10d. OR Chloramphenicol 20 mg/kg PO QID
Typical Pediatric Therapy	Gentamicin 2 mg/kg IV loading dose, then 1.7 mg/kg Q8h OR Streptomycin 10 mg/kg q8h X 10d. OR Chloramphenicol 15 mg/kg PO QID X 10d
Vaccine	Plague
Clinical Hints	Suppurative lymphadenitis; septicemia; hemorrhagic pneumonia; history of animal contact in many cases; case-fatality rates for bubonic plague without therapy are 50% to 60%.
Synonyms	Black death, Black plague, Bubonic plague, Glandular plague, Hemorrhagic plague, Peste, Pneumonic plague, Saint Roch's disease, <i>Yersinia pestis</i> . ICD9: 020 ICD10: A20

Clinical

WHO Case definition for surveillance:

Disease characterized by rapid onset of fever, chills, headache, severe malaise, prostration, with

- bubonic form: extreme painful swelling of lymph nodes (buboes)
- pneumonic form: cough with blood-stained sputum, chest pain, difficult breathing
- Note: Both forms can progress to a septicemic form with toxemia: sepsis without evident buboes rarely occurs.

Laboratory criteria for diagnosis

- Isolation of *Yersinia pestis* in cultures from buboes, blood, CSF or sputum or
- Passive hemagglutination (PHA) test, demonstrating an at least fourfold change in antibody titer, specific for F1 antigen of *Y. pestis*, as determined by the hemagglutination inhibition test (HI) in paired sera.

Case classification

- Suspected: A case compatible with the clinical description. May or may not be supported by laboratory finding of Gram stain negative bipolar coccobacilli in clinical material (bubo aspirate, sputum, tissue, blood).
- Probable: A suspected case with Positive direct fluorescent antibody (FA) test for *Y. pestis* in clinical specimen; or passive hemagglutination test, with antibody titer of at least 1:10, specific for the F1 antigen of *Y. pestis* as determined by the hemagglutination inhibition test (HI); or epidemiological link with a confirmed case.
- Confirmed: A suspected or probable case that is laboratory-confirmed.

Symptoms:

The initial features of plague are nonspecific and include fever, chills, myalgias, pharyngitis, headache.

- Regional lymph nodes are enlarged, painful and extremely tender. ¹
- Additional features, notably in patients with septicemic or pneumonic plague include nausea, vomiting, diarrhea, hematemesis, hematochezia, cough with hemoptysis, dyspnea and signs of meningitis.

Signs:

The physical examination reveals fever, tachycardia, tachypnea, and hypotension.

- Buboes are usually inguinal (60% to 90%), axillary (30%), cervical (10%), or epitrochlear (10%).
- Femoral nodes are involved more frequently than inguinal nodes. ²
- Nodes are typically no larger than 5 cm, extremely tender, erythematous, and surrounded by a boggy hemorrhagic area.
- A maculopapular lesion may be found at the site of the flea bite.
- Acral cyanosis, ecchymosis, petechiae, and digital gangrene are seen in patients with septicemic plague.
- Signs of septic shock or DIC may also be present.

Plague pneumonia: ³

Primary plague pneumonia follows an incubation period of 1 to 3 days, with sudden onset of fever, chills, headache and

malaise. ⁴

- Cough is prominent, with copious sputum production, chest pain and dyspnea.
- Profuse hemoptysis is common.
- Physical examination reveals rales and diffuse areas of dullness to percussion. ⁵
- Untreated plague pneumonia is virtually always fatal.

Rare instances of gastrointestinal plague have been associated with ingestion of contaminated meat. ⁶

This disease is endemic or potentially endemic to 41 countries.

Plague in Uganda

Plague is currently or recently endemic to: ⁷


Western Region:

Nebbi District

Northern Region:

Arua District


The earliest record of plague in Africa was reported by missionaries in Uganda in 1877. ⁸



Graph: Uganda. Plague, cases

Notes:

1. 5,086 cases were reported during 1935 to 1949. ⁹



Graph: Uganda. Plague, deaths

Notes:

1. 31,305 plague deaths were reported during 1910 to 1919; 17,410 during 1920 to 1929; 11,387 during 1930 to 1939. ¹⁰

Presumed local reservoirs include the Nile rat, *Arvicanthus niloticus*. ¹¹
 - Presumed local vectors include *Xenopsylla cheopis* and *X. brasiliensis*.

Notable outbreaks:

- 1982 - An outbreak (152 cases, 3 fatal) was reported.
- 1986 - An outbreak (340 cases, 27 fatal) was reported.
- 1993 - An outbreak (167 cases, 18 fatal) was reported in Western Region (Nebbi District). ¹²
- 1998 - An outbreak was reported in Arua district. ^{13 14}
- 2001 - An outbreak was reported in Nebbi District. ^{15 16}
- 2004 - An outbreak (4 cases, 1 confirmed, 3 fatal) of pneumonic plague was reported in Kango Subcounty.
- 2006 - An outbreak (127 cases, 11 fatal) was reported in Arua and Nebbi Districts - including 12 cases of pneumonic plague. ¹⁷⁻¹⁹
- 2007 - Outbreaks (179 cases, total) were reported in Masindi District (19 cases, 9 fatal) ²⁰⁻²², and in Arua (121 cases, 10 fatal) and Nebbi (39 cases, 9 fatal). ^{23 24}
- 2008 - An outbreak (68 deaths) was reported in Arua and Nebbi. ²⁵⁻²⁷
- 2012 - An outbreak (2 fatal cases) was reported in Arua. ²⁸

References

1. Infect Dis Clin North Am 1991 Mar ;5(1):165-75.
2. ProMED <promedmail.org> archive: 20060528.1500
3. Semin Respir Infect 1997 Mar ;12(1):12-23.
4. ProMED <promedmail.org> archive: 20060614.1650
5. Semin Respir Infect 2003 Sep ;18(3):159-67.
6. Epidemiol Infect 2011 May ;139(5):728-35.
7. Am J Trop Med Hyg 2009 Jun ;80(6):1014-22.
8. Trans R Soc Trop Med Hyg 2010 Feb ;104(2):97-103.
9. Bull World Health Organ 1953 ;9(5):665-700.
10. Bull World Health Organ 1951 ;4(4):475-533.
11. Am J Trop Med Hyg 2009 Oct ;81(4):718-24.
12. Wkly Epidemiol Rec 1995 Feb 17;70(7):45-8.
13. Wkly Epidemiol Rec 2000 Oct 20;75(42):338-43.
14. ProMED <promedmail.org> archive: 19981108.2169
15. Wkly Epidemiol Rec 2003 Apr 18;78(16):130-5.
16. ProMED <promedmail.org> archive: 20011031.2689
17. MMWR Morb Mortal Wkly Rep 2009 Jul 24;58(28):778-81.
18. ProMED <promedmail.org> archive: 20061020.3022
19. ProMED <promedmail.org> archive: 20090724.2617
20. ProMED <promedmail.org> archive: 20070301.0725
21. ProMED <promedmail.org> archive: 20070302.0736
22. ProMED <promedmail.org> archive: 20070309.0835
23. ProMED <promedmail.org> archive: 20070914.3044
24. ProMED <promedmail.org> archive: 20071206.3932
25. ProMED <promedmail.org> archive: 20081104.3474
26. ProMED <promedmail.org> archive: 20081223.4037
27. ProMED <promedmail.org> archive: 20090714.2509
28. ProMED <promedmail.org> archive: 20120213.1039922

Plesiomonas infection

Agent	BACTERIUM. <i>Plesiomonas shigelloides</i> A facultative gram-negative bacillus
Reservoir	Fish Animal Soil Reptile Bird
Vector	None
Vehicle	Water Food
Incubation Period	1d - 2d
Diagnostic Tests	Stool culture - alert laboratory when this organism is suspected. Nucleic acid amplification.
Typical Adult Therapy	Stool precautions. Antimicrobial agent per in-vitro susceptibility (<i>Ciprofloxacin</i> considered 'drug of choice')
Typical Pediatric Therapy	Stool precautions. Antimicrobial agent per in-vitro susceptibility. Fluid replacement
Clinical Hints	Fever, abdominal pain, vomiting and severe diarrhea; symptoms often persist for 2 to 4 weeks; follows ingestion of shellfish or recent travel to developing countries in many cases.
Synonyms	<i>Plesiomonas shigelloides</i> . ICD9: 008.8 ICD10: A04.8

Clinical

The infection is characterized by a self-limited diarrhea, often with blood or mucus in stool. ¹

- Watery diarrhea is most common; however, a cholera-like illness with as many 30 bowel movements per day may occur.
- Associated abdominal pain may mimic that of appendicitis, including enlargement of peritoneal lymph nodes. ²
- Fecal leucocytes are present.
- As many as 30% of cases continue for over four weeks, and symptoms may persist for as long as 3 months.
- *Plesiomonas* has been rarely associated with fatal meningitis and septicemia ³⁻¹³, proctitis ¹⁴, cellulitis and dermal abscesses ¹⁵, pneumonia ¹⁶, pleural effusion ¹⁷, osteomyelitis ¹⁸, cholecystitis ¹⁹, peritonitis ^{20 21}, salpingitis ²², epididymo-orchitis ²³, pancreatitis ²⁴, splenic abscess ²⁵, keratitis ²⁶ and endophthalmitis. ²⁷
- 21 cases of *Plesiomonas* septicemia had been reported as of 1996. ²⁸

This disease is endemic or potentially endemic to all countries.

References

1. Rev Infect Dis 1988 Mar-Apr;10(2):303-16.
2. J Clin Microbiol 1988 Dec ;26(12):2675-7.
3. Pediatr Infect Dis J 1988 Dec ;7(12):877-9.
4. Rev Invest Clin 1988 Oct-Dec;40(4):353-7.
5. Kansenshogaku Zasshi 1985 Nov ;59(11):1154-8.
6. Pediatrics 1983 Mar ;71(3):389-91.
7. J Med Assoc Ga 1982 Nov ;71(11):775-6.
8. Dtsch Med Wochenschr 1982 Aug 20;107(33):1238-9.
9. J Singapore Paediatr Soc 1981 ;23(3-4):156-8.
10. South Med J 1980 Mar ;73(3):393-4.
11. Rev Cubana Med Trop 2000 Jan-Apr;52(1):10-4.
12. Heart Lung 2010 Jul-Aug;39(4):335-9.
13. J Microbiol Immunol Infect 2010 Aug ;43(4):344-6.
14. J Clin Microbiol 1988 Feb ;26(2):388-91.
15. Med Mal Infect 2007 Dec ;37(12):840.
16. Med Mal Infect 2009 Jun ;39(6):397-400.
17. Postgrad Med J 1986 Jul ;62(729):663-4.
18. J Clin Microbiol 1987 Sep ;25(9):1791-3.
19. J Clin Microbiol 1984 Nov ;20(5):985-7.
20. J Clin Microbiol 1988 Dec ;26(12):2675-7.
21. Am J Gastroenterol 1995 Sep ;90(9):1529-30.
22. Clin Microbiol Infect 2002 Dec ;8(12):803-5.
23. AIDS Read 2001 Dec ;11(12):617-9.
24. Rev Infect Dis 1990 Sep-Oct;12(5):813-6.
25. Pediatr Infect Dis J 2001 Dec ;20(12):1178-9.
26. Pediatr Infect Dis J 2012 Nov ;31(11):1200-1.
27. Am J Ophthalmol 1983 Sep ;96(3):403-4.
28. Pediatr Hematol Oncol 1996 May-Jun;13(3):265-9.

Pleurodynia

Agent	VIRUS - RNA. Picornaviridae: Coxsackievirus
Reservoir	Human
Vector	None
Vehicle	Air Fecal-oral Fomite
Incubation Period	3d - 5d
Diagnostic Tests	Viral culture (throat, stool). Serology. Nucleic acid amplification.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Sore throat followed by pleuritic chest pain - a late summer illness in temperate regions; pain is often recurrent and appears in 'waves' - local pressure on affected area may elicit identical pain; usually resolves within one week.
Synonyms	Balme disease, Bamie disease, Bornholm disease, Devil's grip, Drangedal disease, Epidemic benign dry pleurisy, Epidemic myalgia, Sylvest's disease. ICD9: 074.1 ICD10: B33.0

Clinical

Pleurodynia is characterized by a prodrome of upper respiratory tract infection, followed by abrupt onset of pleuritic chest pain. ¹

- The pain may be severe and lead to a misdiagnosis of myocardial infarction.
- Some patients present with abdominal pain suggestive of peritonitis.
- Important diagnostic features include appearance of cases in clusters (often in late summer to autumn) and lack of leucocytosis or other findings suggestive of pneumonia or peritonitis.

This disease is endemic or potentially endemic to all countries.

References

1. Trop Geogr Med 1975 Jun ;27(2):151-9.

Pneumocystis pneumonia

Agent	FUNGUS. Ascomycota ?, Archiascomycetes, Pneumocystidales: <i>Pneumocystis jiroveci</i> (now separate from <i>Pneumocystis carinii</i>)
Reservoir	Human
Vector	None
Vehicle	? Air
Incubation Period	4d - 8w
Diagnostic Tests	Identification of organisms in induced sputum, bronchial washings, tissue. Serology. Nucleic acid amplification.
Typical Adult Therapy	Therapy: Sulfamethoxazole/trimethoprim 25 mg/5 mg/kg QID X 14d. OR Pentamidine 4 mg/kg/d X 14d. OR Dapsone + Trimethoprim . OR Atovaquone OR Primaquine + Clindamycin Prophylaxis - similar, but at altered dosage. Dapsone also used.
Typical Pediatric Therapy	Therapy: Sulfamethoxazole/trimethoprim 25 mg/5 mg/kg QID X 14d. OR Pentamidine 4 mg/kg/d X 14d. OR Dapsone + Trimethoprim . OR Atovaquone OR Primaquine + Clindamycin Prophylaxis - similar, but at altered dosage.
Clinical Hints	Dyspnea, hypoxia and interstitial pneumonia; usually encountered in the setting of severe immune suppression (AIDS, leukemia, etc); roentgenographic findings (typically bilateral alveolar pattern) may follow symptoms only after several days.
Synonyms	PCP, <i>Pneumocystis carinii</i> , <i>Pneumocystis jiroveci</i> . ICD9: 136.3 ICD10: B59

Clinical

P. jiroveci infection often presents as a self-limiting upper respiratory tract infection in infants, predominantly in the age group 1.5 to 4 months of age.

The major presenting symptoms are shortness of breath, fever, and a nonproductive cough. ¹

- Sputum production, hemoptysis and chest pain are rarely encountered. ²
- Tachypnea and tachycardia are usually prominent
- Children may demonstrate cyanosis, flaring of the nasal alae, and intercostal retractions.

Lung auscultation is usually not helpful, with rales present in only 1/3 of adults with this disease.

- The x-ray usually shows bilateral diffuse infiltrates extending from the perihilar region. ³
- Other findings can unilateral infiltrates, nodules, cavities, pneumatoceles, hilar lymphadenopathy and pleural effusion.
- Patients receiving aerosolized pentamidine as prophylaxis have an increased incidence of apical infiltrates and pneumothorax.
- Impaired oxygenation is common.

Extrapulmonary infection by *P. jiroveci* may occur in as many as 3% of infected patients and is reported as an unexpected finding at autopsy.

- The main sites of involvement are lymph nodes, spleen, liver, bone marrow, gastrointestinal tract, eyes ⁴, thyroid, adrenal glands, and kidneys.
- The clinical correlate of these findings is rapidly progressive multisystem disease, an enlarging thyroid mass, pancytopenia, retinal infiltrates, pleural effusion, splenic lesions, and calcifications in the spleen, liver, adrenal, or kidney.
- Rare instances of intestinal pseudotumor ⁵ and cutaneous infection have been reported. ⁶

This disease is endemic or potentially endemic to all countries.

References

1. *Curr Opin Pulm Med* 2005 May ;11(3):203-7.
2. *Curr Opin Infect Dis* 2005 Apr ;18(2):165-70.
3. *Curr Opin Pulm Med* 2008 May ;14(3):228-34.

4. *Ophthalmology* 1997 Nov ;104(11):1853-6.
5. *Arch Pathol Lab Med* 2012 Sep ;136(9):1001-3.
6. *Am J Dermatopathol* 2012 Aug 11;

Pneumonia - bacterial

Agent	BACTERIUM. Streptococcus pneumoniae , <i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i> , other aerobic and facultative gram negative bacilli, etc.
Reservoir	Human
Vector	None
Vehicle	Droplet Endogenous infection
Incubation Period	1d - 3d
Diagnostic Tests	Culture of sputum, blood. Analyze ("grade") sputum cytology to assess significance of culture.
Typical Adult Therapy	Antimicrobial agent(s) appropriate to known or suspected pathogen
Typical Pediatric Therapy	As for adult
Vaccine	Pneumococcal
Clinical Hints	Rigors ("shaking chills"), pleuritic pain, hemoptysis, lobar infiltrate and leukocytosis; empyema and lung abscess suggest etiology other than pneumococcus; foul sputum with mixed flora may herald anaerobic (aspiration) pneumonia.
Synonyms	Bacterial pneumonia, Empiema, Empeem, Empyem, Empyema, Empyeme, Lung abscess, Neumonia, Pleurisy, Pneumococcal infection - invasive, Pneumococcal pneumonia, Polmonite batterica, <i>Streptococcus pneumoniae</i> , <i>Streptococcus pneumoniae</i> - invasive. ICD9: 481,482,483,484 ICD10: J13,J14,J15,J17,J18,J85,J86

Clinical

The designation "Pneumonia • bacterial" in this module is generic, and includes a large variety of etiological agents and anatomical presentations (ie, empyema, lung abscess, lobar• vs. broncho-pneumonia, etc.)

- The clinical features of bacterial pneumonia are largely determined by the infecting species and clinical setting. [1-4](#)
- All forms are characterized by fever, chest pain, productive cough, and physical or roentgenographic evidence for pulmonary consolidation.

Etiological associations:

- AIDS: *Pneumocystis jiroveci*, Mycobacteria (non-tuberculous), Tuberculosis, Nocardiosis, Cryptococcosis, Cytomegalovirus
- Animal contact: Q-fever, Ornithosis
- Aspiration: Oral Anaerobes; if nosocomial, Enterobacteriaceae, *Acinetobacter*, *Pseudomonas*
- Cystic fibrosis (Fibrocystic disease) • *Burkholderia cepacia*
- Drowning ("near-drowning"): *Pseudoallescheria boydii*
- Endocarditis: *Staphylococcus aureus*
- Immunosuppression: *Aspergillosis*, Cryptococcosis, Nocardiosis, *Pneumocystis jiroveci*, Cytomegalovirus
- Infant: see Respiratory syncytial virus, Parainfluenza virus, Respiratory viruses • misc.
- Influenza: Influenza virus, *Streptococcus pneumoniae*, *Staphylococcus aureus*
- Myeloma: *Streptococcus pneumoniae*
- Nosocomial pneumonia: Enterobacteriaceae, *Acinetobacter*, *Pseudomonas*, *Staphylococcus aureus*
- Pulmonary alveolar proteinosis: *Nocardia*
- Traveler or tourist: Histoplasmosis, Legionellosis, Melioidosis

This disease is endemic or potentially endemic to all countries.

References

1. Curr Opin Pulm Med 2004 May ;10(3):171-5.
2. Curr Opin Pulm Med 2005 May ;11(3):218-25.
3. Clin Infect Dis 2004 Dec 1;39(11):1642-50.
4. Infect Dis Clin North Am 2004 Dec ;18(4):791-807; viii.

Poliomyelitis

Agent	VIRUS - RNA. Picornaviridae, Picornavirus: Polio virus
Reservoir	Human
Vector	None
Vehicle	Fecal-oral Dairy products Food Water Fly
Incubation Period	7d - 14d (range 3d - 35d)
Diagnostic Tests	Viral culture (pharynx, stool). Serology. Nucleic acid amplification.
Typical Adult Therapy	Stool precautions; supportive
Typical Pediatric Therapy	As for adult
Vaccines	Poliomyelitis - injectable Poliomyelitis - oral
Clinical Hints	Sore throat, headache, vomiting and myalgia followed by flaccid paralysis; meningeal involvement in 1% of cases - paralysis in only 0.1%. paralysis tends to be more extensive in adult patients.
Synonyms	Acute flaccid paralysis, Heine-Medin disease, Infantile paralysis, Kinderlahmung, Kinderverlamming, Paralisi infantile, Paralysis flaccida, Paralysis flacida aguda, PFA (Paralysis Flacidas Agudas), Polio, Poliomyelite, Poliomyelitt. ICD9: 045 ICD10: A80

Clinical

CDC (The United States Centers for Disease Control) case definition for surveillance:

For surveillance purposes, the CDC (The United States Centers for Disease Control) case definition of paralytic poliomyelitis requires, "Acute onset of a flaccid paralysis ¹ of one or more limbs with decreased or absent tendon reflexes in the affected limbs, without other apparent cause, and without sensory or cognitive loss."

• A 'confirmed case' requires persistence of the neurological deficit 60 days after onset of initial symptoms, fatal illness or unknown follow-up status.

The WHO Case definition for surveillance includes any child under fifteen years of age with acute, flaccid paralysis or any person with paralytic illness at any age when poliomyelitis is suspected. ²

Poliomyelitis is typically a late summer illness in temperate climates, and often begins as a mild upper respiratory tract infection.

- In some cases, the disease follows vaccination (live vaccine) or recent contact with a vaccinee.
- Patients have been known to excrete virus for as long as ten years following an episode of poliomyelitis ³
- Antecedent injection in a given site may precipitate paralytic poliomyelitis in the same limb. ⁴⁻⁹

90% to 95% of poliomyelitis infections are asymptomatic.

- Symptoms include fever, sore throat, headache, vomiting and stiff neck.
- Paralysis is typically asymmetrical, and most often involves the lower extremities.
- Bulbar paralysis or encephalitis may occur in patients in the absence of limb paralysis.
- 4% to 8% experience minor symptoms, and 1% to 2% develop paralysis.
- Paralysis is most common in the very young and very old, following minor blunt trauma to a limb, and among persons who had undergone tonsillectomy.
- The case/fatality rate for paralytic poliomyelitis is 2% to 10%.

This disease is endemic or potentially endemic to 87 countries.


Poliomyelitis in Uganda

Vaccine Schedule:


BCG - birth

DTwP - 6, 10, 14 weeks

DTwPHibHep - 6, 10, 14 weeks
 Measles (monovalent) - 9 months
 OPV - birth; 6, 10, 14 weeks
 TT - Women of child-bearing age; +1, +6 months; +1, +1 year
 Vitamin A - 6, 12, 18, 24, 30, 36, 42 months



Graph: Uganda. Poliomyelitis - WHO-UNICEF est. % vaccine (POL3) coverage




Graph: Uganda. Poliomyelitis, cases


Notes:

1. Historical data from references [10](#) [11](#)
 Individual years:


- 1997 - None due to wild virus.
- 1998 - None due to wild virus.
- 2008 - All due to wild virus. ¹²
- 2009 - Imported infection, from Sudan ¹³⁻¹⁵
- 2010 - Case in Bugiri district. ¹⁶



Graph: Uganda. Poliomyelitis, deaths



Graph: Uganda. Acute flaccid paralysis (AFP), cases



Graph: Uganda. AFP, rate per 100,000 below age 15

References

1. Curr Opin Infect Dis 2003 Oct ;16(5):375-81.
2. Bull World Health Organ 1992 ;70(1):79-84.
3. Biologicals 2006 Jun ;34(2):113-6.
4. Trans R Soc Trop Med Hyg 1985 ;79(3):355-8.
5. J Virol 1998 Jun ;72(6):5056-60.
6. N Engl J Med 1995 Jul 6;333(1):64.
7. N Engl J Med 1995 Jul 6;333(1):63; author reply 64.
8. N Engl J Med 1995 Jul 6;333(1):62; author reply 64.
9. N Engl J Med 1995 Jul 6;333(1):63; author reply 64.
10. Bull World Health Organ 1955 ;12(4):595-649.
11. Bull World Health Organ 1956 ;15(1-2):43-121.
12. Wkly Epidemiol Rec 2009 Jun 19;84(25):257-60.
13. MMWR Morb Mortal Wkly Rep 2010 Nov 5;59(43):1393-9.
14. ProMED <promedmail.org> archive: 20090226.0808
15. ProMED <promedmail.org> archive: 20090305.0900
16. ProMED <promedmail.org> archive: 20101027.3900

Protothecosis and chlorellosis

Agent	ALGA. <i>Prototheca wickerhamii</i> ; rarely <i>Pr. zopfii</i> , <i>Pr. cutis</i> Achloric algae <i>Chlorella</i> spp. contain chloroplasts
Reservoir	A rare animal pathogen (cat, dog, cattle). <i>Chlorella</i> spp. are reported to infect domestic and wild mammals.
Vector	None
Vehicle	Water Sewage Food Local trauma
Incubation Period	Unknown
Diagnostic Tests	Culture on fungal media. Biopsy. Nucleic acid amplification.
Typical Adult Therapy	Surgical excision. There are anecdotal reports of successful therapy with <i>Amphotericin B</i> , <i>Ketoconazole</i> and <i>Itraconazole</i> (latter 200 mg/day X 2 months) or <i>voriconazole</i>
Typical Pediatric Therapy	As for adult (<i>Itraconazole</i> 2 mg/kg/day X 2 months)
Clinical Hints	May follow immune suppression or skin trauma; dermal papules, plaques, eczematoid or ulcerated lesions; olecranon bursitis; systemic infection also reported.
Synonyms	Chlorellosis, <i>Prototheca</i> , Protothecosis. ICD9: 136.8 ICD10: B99

Clinical

Four forms of disease are reported:

- cutaneous infection
- olecranon bursitis
- disseminated
- onychomycosis ¹⁻³

The incubation period of protothecosis is unknown; however, infections which have followed trauma have appeared after approximately two weeks. ⁴

- Most cases presented as a single painless, slowly progressive, well-circumscribed plaque or papulonodular skin lesion that may become eczematoid or ulcerated. ⁵⁻⁷
- Soft tissue lesions favor the olecranon bursa; sites of minor trauma or corticosteroid injection; or surgical wounds which have been exposed to soil or water. ^{8 9}
- Lesions enlarge gradually over weeks to months, with no tendency for healing.
- Other presentations have included tenosynovitis ¹⁰ ; algemia complicating immune-suppression ¹¹ ; nasopharyngeal ulcerated lesion followed prolonged intubation, and infection of ambulatory peritoneal catheters.
- Skin lesions in HIV-infected patients are similar to those of healthy patients
- Peritonitis due to *P. wickerhamii* has been reported in peritoneal dialysis patients. ¹²
- A case of subacute endocarditis due to *Prototheca wickerhamii* has been reported. ¹³

A single case of *Chlorella* wound infection has been reported. ¹⁴

This disease is endemic or potentially endemic to all countries.

References

1. Int J Syst Evol Microbiol 2006 Jun ;56(Pt 6):1419-25.
2. Int J Dermatol 2006 Sep ;45(9):1071-3.
3. Mycopathologia 2011 Sep ;172(3):207-13.
4. Cutis 1999 Mar ;63(3):185-8.
5. Med Mycol 2004 Apr ;42(2):95-106.
6. Dermatol Clin 2003 Apr ;21(2):249-55.
7. Clin Dermatol 2012 Jul ;30(4):432-6.
8. Clin Microbiol Rev 2007 Apr ;20(2):230-42.
9. J Cutan Med Surg 2009 Sep-Oct;13(5):273-5.
10. Clin Orthop Relat Res 2008 Dec ;466(12):3143-6.
11. Emerg Infect Dis 2009 Jul ;15(7):1129-30.
12. Nefrologia 2007 ;27(1):81-2.
13. Klin Med (Mosk) 2011 ;89(4):69-76.
14. Am J Clin Pathol 1983 Jul ;80(1):102-7.

Pseudocowpox

Agent	VIRUS - DNA. Poxviridae, Parapoxvirus: Pseudocowpox virus
Reservoir	Cattle
Vector	None
Vehicle	Contact
Incubation Period	5d - 14d
Diagnostic Tests	Viral culture (skin lesion or exudate). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Umbilicated nodule on the hand following contact with cattle; mild regional lymphadenopathy.
Synonyms	Bovine papular stomatitis, Farmyard pox, Milker's nodule, Noduli mulgentinum, Sealpox. ICD9: 051.1 ICD10: B08.0

Clinical

Pseudocowpox is mild and self-limited and characterized by a red-to-blue dermal nodule associated with minimal lymphadenopathy. ¹

This disease is endemic or potentially endemic to all countries.

References

1. [J Am Acad Dermatol 2001 Jan ;44\(1\):1-16.](#)

Pyodermas (impetigo, abscess, etc)

Agent	BACTERIUM. Various (<i>Staphylococcus aureus</i> & <i>Streptococcus pyogenes</i> predominate)
Reservoir	Human
Vector	None
Vehicle	Endogenous & contact with infected secretions
Incubation Period	Variable
Diagnostic Tests	Clinical diagnosis usually sufficient. Aspiration of lesion for smear and culture may be helpful in some cases.
Typical Adult Therapy	Antibiotic directed at likely pathogens (Group A Streptococcus and <i>Staphylococcus aureus</i>)
Typical Pediatric Therapy	As for adult
Clinical Hints	Impetigo characterized by vesicles which progress to pustules ('honey-colored pus'); highly contagious; may be complicated by acute glomerulonephritis.
Synonyms	Acne vulgaris, Carbonchio, Carbuncle, Follicolite, Follicolite, Folliculite, Folliculitis, Follikulitis, Foroncolosi, Forunculose, Forunculosi, Furunculosis, Furunkulose, Furunulose, Hydradenitis, Impetigine, Impetigo, Paronychia, Pyoderma. ICD9: 680,684,686 ICD10: L01,L02,L08.0,L73.2

Clinical

Impetigo is characterized by multiple superficial lesions caused by group A-hemolytic streptococci and/or *Staphylococcus aureus*.¹

- The lesions consist of pustules that rupture and form a characteristic honey-colored crust.
- Lesions caused by staphylococci are associated with tense, clear bullae (bullous impetigo.).
- Ecthyma is a variant of impetigo that usually presents as punched-out ulcers on the lower extremities.
- Streptococcal impetigo is most common among children 2 to 5 years of age, and epidemics may occur in settings of poor hygiene, lower socioeconomic status or tropical climates.
- The most important complication of impetigo is poststreptococcal glomerulonephritis.

Folliculitis is most often caused by *Staphylococcus aureus*.²

- Blockage of sebaceous glands may result in sebaceous cysts, which may present as extensive abscesses or become secondarily infected.
- Infection of specialized sweat glands (hidradenitis suppurativa) occur in the axillae.
- Chronic folliculitis is a hallmark of acne vulgaris, in which normal flora (e.g., *Propionibacterium acnes*) may play a role.
- Diffuse folliculitis may herald infection by *Pseudomonas aeruginosa* or *Aeromonas hydrophila*³, in waters that are insufficiently chlorinated and maintained at temperatures above 37 C. Although such Infection is usually self-limited, bacteremia and septic shock have been reported.

Erysipelas is caused by *Streptococcus pyogenes* and is characterized by abrupt onset of "fiery-red" superficial swelling of the face or extremities.

- The lesion is typically recognized by the presence of well-defined indurated margins, particularly along the nasolabial fold; rapid progression; and intense pain.
- Flaccid bullae may develop on the second or third day of illness; but extension to deeper soft tissues is rare.
- Desquamation occurs between the fifth and tenth days of illness.

Cellulitis is characterized by local pain, erythema, swelling, and heat.

- Cellulitis may be caused by any of a wide variety of bacteria or yeasts; however, *S. aureus* or *S. pyogenes* are most often implicated.
- A history of preceding trauma, insect bite, needle insertion or surgery is often present.
- Cultures of biopsy specimens or aspirates are positive in only 20% of cases.
- Infection by *S. aureus* often spreads out from a localized infection (abscess, folliculitis) or foreign body.
- Streptococcal cellulitis tends to be more diffuse and rapid in onset, and associated with lymphangitis and fever.
- Streptococci also cause recurrent cellulitis in the setting of lymphedema resulting from elephantiasis or lymph node damage.
- Recurrent staphylococcal cutaneous infections are encountered in patients with "Job's syndrome" (eosinophilia and elevated serum levels of IgE); and nasal carriers of staphylococci.

Cellulitis associated with animal bites is commonly caused by *Pasteurella multocida*, *Staphylococcus intermedius* and *Capnocytophaga canimorsus* (formerly DF-2) and is discussed separately in this module under 'Pasteurellosis, etc.'

- Human bites contain a variety of anaerobic organisms (*Fusobacterium*, *Bacteroides*), aerobic and anaerobic streptococci, and *Eikenella corrodens*.
- *Aeromonas hydrophila* causes an aggressive form of cellulitis following minor trauma in marine environments.
- *P. aeruginosa* is the most common cause of ecthyma gangrenosum and infection following penetrating injuries to the foot.
- Gram-negative bacillary cellulitis, (including *P. aeruginosa* infection) is common among hospitalized, immunocompromised patients.

This disease is endemic or potentially endemic to all countries.

References

1. [Semin Cutan Med Surg 2004 Mar ;23\(1\):29-38.](#)
2. [Am Fam Physician 2002 Jul 1;66\(1\):119-24.](#)
3. [Australas J Dermatol 2008 Feb ;49\(1\):39-41.](#)

Pyomyositis

Agent	BACTERIUM. Usually Staphylococcus aureus
Reservoir	Human
Vector	None
Vehicle	Hematogenous
Incubation Period	Variable
Diagnostic Tests	Ultrasonography or CT scan.
Typical Adult Therapy	Antibiotic directed at confirmed or suspected pathogen (usually <i>Staphylococcus aureus</i>); drainage
Typical Pediatric Therapy	As for adult
Clinical Hints	Pain, swelling and "woody" induration of a large muscle (usually lower limb or trunk) associated with fever and leukocytosis; often follows trauma to the involved region; lymphadenopathy uncommon; leucocytosis in most cases.
Synonyms	Tropical pyomyositis. ICD9: 040.81 ICD10: M60.0

Clinical

The initiating lesion may be overt blunt or penetrating trauma; however, some cases may represent complications of viral or parasitic myositis. ¹

- An increasing percentage of reported patients have been HIV-positive. ²
- 20 to 50% of patients with pyomyositis recall recent blunt trauma or vigorous exercise involving the area of infection; and most infections involve a single muscle or muscle group.
- Rare cases of pyomyositis have been associated with spinal epidural abscess ³ or Lemmiere's syndrome ⁴
- The major muscles of the lower extremities and trunk muscles are most often infected ⁵ ; however, virtually any muscle can be involved. ⁶⁻¹¹

Onset is often subacute with fever, swelling with or without erythema, mild pain and minimal tenderness. ¹²

- The involved area is indurated or has a wooden consistency.
- 10 to 21 or more days later, the patient complains of fever, with muscle tenderness and swelling.
- Overlying skin is intact and warm, usually without erythema.
- There is no regional lymphadenitis.
- At this point, pus can be aspirated from the involved muscle.
- Eventually, manifestations of sepsis appear, with local erythema, tenderness and fluctuance. ¹³
- Additional symptoms may reflect compression of contiguous structures. ^{14 15}
- Septicemia, ARDS and rapidly progressive or fatal infections are also encountered. ^{16 17}

Leukocytosis is present.

- Eosinophilia suggests a diagnosis of 'tropical myositis' but is thought to represent the presence of concurrent parasitic infection.

The clinical features of pyomyositis may mimic those of leptospirosis. ¹⁸ \

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 2008 Jul ;21(3):473-94.
2. J Natl Med Assoc 1996 Sep ;88(9):565-9.
3. J Neurosurg Pediatr 2010 Jul ;6(1):33-7.
4. Am J Med Sci 2008 Jun ;335(6):499-501.
5. J Rheumatol 1997 Sep ;24(9):1734-8.
6. Orthopedics 2008 Nov ;31(11):1146.
7. Br J Oral Maxillofac Surg 2010 Apr ;48(3):216-7.
8. Indian J Ophthalmol 2010 Nov-Dec;58(6):532-5.
9. J Med Case Rep 2011 ;5:195.
10. J Pediatr Orthop B 2012 Apr 8;
11. BMJ Case Rep 2012 ;2012
12. Semin Arthritis Rheum 1994 Jun ;23(6):396-405.
13. Pediatr Radiol 2006 Apr ;36(4):338-43.
14. J Med Case Rep 2008 ;2:204.
15. QJM 2008 Dec ;101(12):983-4.
16. Am J Forensic Med Pathol 2008 Jun ;29(2):131-5.
17. Asian Pac J Trop Med 2011 Apr ;4(4):325-7.
18. Trop Doct 2008 Oct ;38(4):254-6.

Q-fever

Agent	BACTERIUM. <i>Coxiella burnetii</i> Intracellular organism related to Rickettsiae
Reservoir	Cattle Sheep Goat Bird Fish Rodent Rabbit Tick Bandicoot Marsupial Dog Cat
Vector	None
Vehicle	Air Dust Infected secretions Dairy products
Incubation Period	18d - 21d (range 4d - 40d)
Diagnostic Tests	Serology. Culture possible in specialized laboratories. Nucleic acid amplification.
Typical Adult Therapy	Doxycycline 100 mg BID X 2w OR Fluoroquinolone Add Hydroxychloroquine 600 mg per day if endocarditis
Typical Pediatric Therapy	Age < 8 years: Erythromycin 10 mg/kg QID X 2 weeks Age >= 8 years: Doxycycline 100 mg BID X 2 weeks
Vaccine	Q fever
Clinical Hints	Headache, myalgia, cough and hepatic dysfunction; hepatosplenomegaly, 'F.U.O.' and endocarditis encountered; proximity to farming or animals during 2 to 4 weeks preceding illness; most infections resolve in 1 to 2 weeks; case-fatality rate = 1.5%.
Synonyms	Balkan grippe, <i>Coxiella burnetii</i> , Febbre australiana, Febre Q, Nine Mile fever, Q-Fieber, Q-koorts, Query fever, Red River fever. ICD9: 083.0 ICD10: A78

Clinical

The typical clinical presentation of Q-fever (pneumonia vs. hepatitis) seems to vary from region to region. ^{1 2}

Q-fever is often asymptomatic or mistaken for an acute viral illness.

- Q-fever may be mistaken for Legionnaires' disease ³
- After an incubation period of 2 to 3 weeks, the patients develops fever, headache, and myalgias. ⁴
- Cough is present in 25% to 70%, and hepatosplenomegaly in 30% to 50%.
- An evanescent rash may appear in 5% of cases.
- The blood CRP is elevated; however leukocytosis is present in only 20% of cases. ⁵
- Acute thrombocytosis may also be encountered. ⁶
- False-positive tests toward a variety of non-related agents and conditions may be encountered: anti-nuclear antibody (ANA), smooth muscle antibody, rheumatoid factor, Epstein-Barr Virus, Cytomegalovirus, *Mycoplasma pneumoniae*, Parvovirus, *Bordetella pertussis*, *Rickettsia conorii* and *Rickettsia typhi*. ⁷

The frequency of pneumonitis is highly variable (10% to 60%) ^{8 9} ; and clinical and radiological features are non-specific. ¹⁰⁻¹²

- Neurological complications may include encephalitis, brachial plexopathy ¹³ , Guillain-Barre syndrome ¹⁴ , status epilepticus and pseudotumor cerebri ¹⁵
- Several cases of Q-fever uveitis have been reported. ¹⁶ In one case, a patient developed anterior uveitis accompanied by exudative bilateral inferior retinal detachment and optic disk edema. ¹⁷

Occasionally, the illness may be prolonged, with severe pneumonitis ^{18 19} and hepatic involvement. ²⁰⁻²²

- Independent risk factors for development of chronic Q fever include valvular surgery, vascular prosthesis, aneurysm, renal insufficiency, and older age. ²³
- Chronic fatigue is common following Q-fever, and in some cases may actually represent persistent infection. ²⁴⁻²⁸
- Although the acute disease is usually self-limited, Q-fever endocarditis may occasionally develop 3 to 20 years following the acute infection and is often fatal. ^{29 30}
- Pericarditis ³¹⁻³⁴ , myocarditis ^{35 36} , aortitis ³⁷ , optic neuritis ³⁸ , uveitis ³⁹⁻⁴¹ , disseminated intravascular coagulation ⁴² , hemophagocytic syndrome ⁴³⁻⁴⁵ , bleeding phenomena (melena, epistaxis, petechiae) ⁴⁶ , autoimmune hemolytic anemia ⁴⁷ , osteomyelitis ^{48 49} , recurrent subcutaneous abscesses and nodules ⁵⁰ , spontaneous abortion ^{51 52} , prosthetic joint infection ⁵³ , cholecystitis ⁵⁴ and tubulointerstitial nephritis ⁵⁵ have been reported as complications of Q-

fever. ⁵⁶⁻⁵⁸

- Over 80% of patients with Q-fever endocarditis have a history of underlying valvular disease.
- Additional vascular complications of Q-fever include aneurysm rupture, aorto-enteric fistulae and lower-limb embolisation. ^{59 60}
- Q fever may mimic Kawasaki disease ⁶¹ , lupus erythematosus ⁶² or Crimean-Congo hemorrhagic fever. ⁶³

This disease is endemic or potentially endemic to all countries.

References

1. Mayo Clin Proc 2008 May ;83(5):574-9.
2. Infect Dis Clin North Am 2008 Sep ;22(3):505-14, ix.
3. Heart Lung 2009 Jul-Aug;38(4):354-62.
4. Eur J Clin Microbiol Infect Dis 2003 Feb ;22(2):108-10.
5. Clin Chem Lab Med 2009 ;47(11):1407-9.
6. Heart Lung 2009 Sep-Oct;38(5):444-9.
7. Eur J Clin Microbiol Infect Dis 2011 Dec ;30(12):1527-30.
8. Respir Care Clin N Am 2004 Mar ;10(1):99-109.
9. Curr Opin Infect Dis 2004 Apr ;17(2):137-42.
10. Nihon Kokyuki Gakkai Zasshi 2008 Dec ;46(12):967-71.
11. Br J Radiol 1991 Dec ;64(768):1101-8.
12. Respir Care Clin N Am 2004 Mar ;10(1):99-109.
13. Muscle Nerve 2008 Dec ;38(6):1644-8.
14. Rev Neurol (Paris) 2013 Feb 7;
15. Pediatr Neurol 2008 Jan ;38(1):44-6.
16. Medicine (Baltimore) 2008 May ;87(3):167-76.
17. Clin Ophthalmol 2011 ;5:1273-5.
18. Chest 1998 Sep ;114(3):808-13.
19. Respir Care Clin N Am 2004 Mar ;10(1):99-109.
20. J Med Case Rep 2007 ;1:154.
21. Gastroenterol Hepatol 2010 Jan ;33(1):21-4.
22. J Infect 2012 Mar ;64(3):247-59.
23. Emerg Infect Dis 2012 Apr ;18(4):563-70.
24. Lancet 1996 Apr 6;347(9006):978-9.
25. BMC Infect Dis 2011 ;11:97.
26. J Psychosom Res 2012 Apr ;72(4):300-4.
27. Ned Tijdschr Geneesk 2012 ;156(48):A5474.
28. Ned Tijdschr Geneesk 2012 ;156(48):A5258.
29. Scand J Infect Dis 1997 ;29(1):41-9.
30. Can J Cardiol 2006 Jul ;22(9):781-5.
31. Clin Infect Dis 1999 Aug ;29(2):393-7.
32. Clin Microbiol Infect 2009 Dec ;15 Suppl 2:173-5.
33. Ann N Y Acad Sci 2006 Oct ;1078:248-51.
34. Ann Thorac Surg 2004 Jul ;78(1):326-8.
35. Ann N Y Acad Sci 2006 Oct ;1078:248-51.
36. Hippokratia 2008 Jan ;12(1):46-9.
37. Acta Cardiol 2012 Apr ;67(2):261-4.
38. Int J Infect Dis 2010 Sep ;14 Suppl 3:e269-73.
39. Bull Soc Ophtalmol Fr 1959 Sep-Oct;7:599-605.
40. Clin Microbiol Infect 2009 Dec ;15 Suppl 2:176-7.
41. Clin Experiment Ophthalmol 2008 Nov ;36(8):797-8.
42. Diagn Microbiol Infect Dis 2011 Feb ;69(2):210-2.
43. Infection 2011 Dec ;39(6):579-82.
44. Scand J Infect Dis 2006 ;38(11-12):1119-22.
45. Int J Infect Dis 2013 Feb 8;
46. Ter Arkh 2011 ;83(11):51-5.
47. Ann Clin Microbiol Antimicrob 2012 May 18;11(1):14.
48. Infection 2011 Apr ;39(2):167-9.
49. Comp Immunol Microbiol Infect Dis 2012 Mar ;35(2):169-72.
50. Pediatr Infect Dis J 2012 May ;31(5):525-7.
51. Clin Microbiol Infect 2011 Apr 25;
52. Eur J Clin Microbiol Infect Dis 2012 Sep 28;
53. J Clin Microbiol 2012 Oct 17;
54. Pediatr Infect Dis J 2012 Dec 26;
55. Ann Clin Microbiol Antimicrob 2012 May 18;11(1):14.
56. Lancet Infect Dis 2005 Dec ;5(12):734-5.
57. Vector Borne Zoonotic Dis 2010 May ;10(4):421-3.
58. Med Clin (Barc) 2008 Dec 6;131(20):798-9.
59. Eur J Vasc Endovasc Surg 2011 Sep ;42(3):384-92.
60. J Med Case Rep 2011 ;5:565.
61. Kansenshogaku Zasshi 2009 May ;83(3):245-50.
62. Reumatol Clin 2012 May-Jun;8(3):143-4.
63. Indian J Med Microbiol 2011 Oct-Dec;29(4):418-9.

Rabies

Agent	VIRUS - RNA. Rhabdoviridae, Mononegavirales, Lyssavirus: Rabies virus. Other human Lyssaviruses = Mokola, Duvenhage, European Bat (EBL)
Reservoir	Dog Fox Skunk Jackal Wolf Cat Raccoon Mongoose Bat Rarely rodent or Rabbit
Vector	None
Vehicle	Saliva Bite Transplants Air (bat aerosol)
Incubation Period	1m - 3m (range 4d to 19 years !)
Diagnostic Tests	Viral culture & direct immunofluorescence of saliva, CSF, corneal smears. Serology. Nucleic acid amplification.
Typical Adult Therapy	Strict isolation; supportive. The Milwaukee protocol (prolonged deep sedation and support) has been successful in some cases. See Vaccines module for pre- and post-exposure schedules
Typical Pediatric Therapy	As for adult
Vaccines	Rabies Rabies immune globulin
Clinical Hints	Follows animal bite (rarely lick) - often after months: agitation, confusion, seizures, painful spasms of respiratory muscles, progressive paralysis, coma and death; case-fatality rate > 99%.
Synonyms	Aravan, Australian bat lyssavirus, Ballina, BBLV, Bokeloh bat lyssavirus, Duvenhage, EBL, European bat Lyssavirus, Hondsdolheid, Hydrophobia, Ikoma lyssavirus, Irkut, Khujand, Lyssa, Mokola, Pteropus lyssavirus, Rabia, Rage, Raiva, Saint Hubert's disease, Shimoni bat virus, Tollwut, West Caucasian bat, Wutkrankheit. ICD9: 071 ICD10: A82

Clinical

WHO Case definition for surveillance:

- An acute neurological syndrome (encephalitis) dominated by forms of hyperactivity (furious rabies) or paralytic syndromes (dumb rabies) that progresses towards coma and death, usually by respiratory failure, within 7 to 10 days after the first symptom if no intensive care is instituted.
- Bites or scratches from a suspected animal can usually be traced back in the patient medical history.
- The incubation period may vary from days to years ¹ but usually falls between 30 and 90 days.

Laboratory criteria for diagnosis

One or more of the following

- Detection of rabies viral antigens by direct fluorescent antibody (FA) in clinical specimens, preferably brain tissue (collected post mortem)
- Detection by FA on skin or corneal smear (collected ante mortem)
- FA positive after inoculation of brain tissue, saliva or CSF in cell culture, in mice or in suckling mice
- Detectable rabies-neutralizing antibody titer in the CSF of an unvaccinated person
- Identification of viral antigens by PCR on fixed tissue collected post mortem or in a clinical specimen (brain tissue or skin, cornea or saliva)
- Isolation of rabies virus from clinical specimens and confirmation of rabies viral antigens by direct fluorescent antibody testing

Case classification

Rabies:

- Suspected: A case that is compatible with the clinical description.
- Probable: A suspected case plus history of contact with suspected rabid animal.
- Confirmed: A suspected case that is laboratory-confirmed.

Rabies exposure:

- Possibly exposed: A person who had close contact (usually a bite or scratch) with a rabies-susceptible animal in (or originating from) a rabies-infected area.
- Exposed: A person who had a close contact (usually a bite or scratch) with a laboratory-confirmed rabid animal.


The initial symptoms of rabies are often limited to low grade fever and pain or paresthesia at the site of inoculation.

- Progressive encephalitis then ensues. ²
- "Furious rabies" is characterized by hyperactivity, fluctuating level of consciousness, aerophobia and hydrophobia.
- Bizarre behavior and lack of focal neurological signs are typical.

- Hydrophobia may manifest as 'jerky' inspiratory spasms progressing to opisthotonus, generalized seizures or cardiorespiratory arrest.
- Similar reactions may be elicited by fanning the patient ("aerophobia).
- Paralytic ("dumb") rabies is characterized by progressive flaccid paralysis, with fasciculation and pain in the affected muscles.
- Minor sensory disturbances may be present. Such patients may survive for as long as one month, ultimately dying of bulbar and respiratory paralysis.
- Rare instances of survival have been documented. [3-16](#)
- In Africa, rabies is often mis-diagnosed as cerebral malaria.

This disease is endemic or potentially endemic to 151 countries.

Rabies in Uganda




Graph: Uganda. Rabies, cases

Notes:


Individual years:

- 1993 - All from dogs.
- 1994 - All from dogs.
- 1995 - All from dogs.
- 1996 - 14 from dogs, and one from a fox.
- 1997 - All from dogs.
- 1999 - All from dogs.
- 2009 - 59 cases of suspected rabies were reported in Kampala and Wakiso during a single week. [17](#)

An estimated 6,601 dog bites occur each year (39.6 per 100,000), 40% to children below age 10 years. [18](#)




Graph: Uganda. Rabies - postexposure prophylaxis, cases



Graph: Uganda. Rabies, animal

Notes:

- Individual years:
- 1970 - One cat and three farm animals.
- 1971 - Four ruminants.
- 2000, 2001, 2002 - All bovines.
- 2002 - No rabid bats were identified.



Graph: Uganda. Rabies, dog

References

1. Emerg Infect Dis 2008 Dec ;14(12):1950-1.
2. Mayo Clin Proc 2004 May ;79(5):671-6.
3. N Engl J Med 2005 Jun 16;352(24):2508-14.
4. MMWR Morb Mortal Wkly Rep 2007 Apr 20;56(15):361-5.
5. Trans R Soc Trop Med Hyg 2008 Oct ;102(10):979-82.
6. MMWR Morb Mortal Wkly Rep 2010 Feb 26;59(7):185-90.
7. MMWR Morb Mortal Wkly Rep 2012 Feb 3;61:61-5.
8. ProMED <promedmail.org> archive: 20081114.3599
9. ProMED <promedmail.org> archive: 20081122.3689
10. ProMED <promedmail.org> archive: 20090214.0638
11. ProMED <promedmail.org> archive: 20090919.3292
12. MMWR Morb Mortal Wkly Rep 2012 Feb 3;61:61-5.
13. ProMED <promedmail.org> archive: 20110527.1619
14. ProMED <promedmail.org> archive: 20110606.1728
15. ProMED <promedmail.org> archive: 20110613.1802
16. ProMED <promedmail.org> archive: 20120106.1002196
17. ProMED <promedmail.org> archive: 20090424.1548
18. Trop Med Int Health 2005 Aug ;10(8):790-8.

Rat bite fever - spirillary

Agent	BACTERIUM. <i>Spirillum minus</i> An aerobic gram-negative spirochete
Reservoir	Rat Mouse Cat
Vector	None
Vehicle	Bite
Incubation Period	7d - 21d (range 5d - 40d)
Diagnostic Tests	Dark-field exam of wound. Animal inoculation.
Typical Adult Therapy	Amoxicillin/clavulanate 875/125 mg PO BID X 7d. OR Procaine Penicillin G 600,000u IM q12h X 7d. OR Doxycycline 200 mg BID X 7d
Typical Pediatric Therapy	Amoxicillin/clavulanate 10 mg/kg PO BID X 7d OR Procaine Penicillin G 25,000u/kg IM q12h X 7d
Clinical Hints	Lymphadenopathy, myalgia, maculopapular rash and recurrent fever beginning 1 to 3 weeks after rat bite; infection resolves after 3 to 6 days; case-fatality rate = 6%.
Synonyms	Sodoku, Spirillosis, Spirillum minor, Spirillum minus. ICD9: 026.0 ICD10: A25.0

Clinical

Most patients present with a recent rat bite wound, which may later form an ulcer with local swelling, pain and skin changes.

- Regional lymphatics and lymph nodes are enlarged and tender.
- Fever rises to as high as 40 C, with accompanying rigors.
- After 3 days, fever ends in 'crisis,' followed by a quiescent interval of 5 to 10 days.
- One or more relapses follow, and are associated with a purple papular exanthem on the chest and arms.
- Additional findings include generalized hyperreflexia, arthralgia, myalgia and hyperesthesia.
- The fatality rate without treatment is 10%.

Features which may distinguish spirillary [S] from streptobacillary [B] rat bite fever include the following: ^{1 2}

- | | |
|------------|------------------------------------|
| incubation | - S up to 30 days |
| | - B up to 10 days |
| bite wound | - S may produce a chancre |
| | - B heals promptly |
| relapses | - S regular |
| | - B intermittent |
| rash | - S generalized macular |
| | - B macular, pustular or petechial |
| arthritis | - S rare |
| | - B common ³ |

This disease is endemic or potentially endemic to all countries.

References

1. Scand J Infect Dis 2002 ;34(6):474-7.
2. Lancet Infect Dis 2001 Sep ;1(2):91.
3. Clin Orthop Relat Res 2000 Nov ;(380):173-6.

Rat bite fever - streptobacillary

Agent	BACTERIUM. <i>Streptobacillus moniliformis</i> A facultative gram-negative bacillus
Reservoir	Rat Squirrel Weasel Turkey
Vector	None
Vehicle	Infected secretions Bite Dairy products
Incubation Period	3d - 10d (range 1d - 22d)
Diagnostic Tests	Culture of blood or joint fluid. Nucleic acid amplification.
Typical Adult Therapy	<i>Amoxicillin/clavulanate</i> 875/125 mg PO BID X 7d. OR <i>Doxycycline</i> 100 mg PO BID X 7d
Typical Pediatric Therapy	<i>Amoxicillin/clavulanate</i> 10 mg/kg TID X 7d. OR (if age>8 years) <i>Doxycycline</i> 2 mg/kg PO BID X 7 days (maximum 200 mg/day)
Clinical Hints	Headache, myalgia, maculopapular rash and arthralgia or arthritis; history of a rat bite during the preceding 1 to 3 weeks in most cases; case-fatality rate = 10%.
Synonyms	Haverhill fever, Streptobacillosis, <i>Streptobacillus moniliformis</i> . ICD9: 026.1 ICD10: A25.1

Clinical

Most patients present with a recent rat bite wound, which may later form an ulcer with local swelling, pain and skin changes.¹

- Symptoms include fever, prostration, marked myalgia and muscle tenderness, headache and a generalized morbilliform rash • most marked on the hands and feet.²
- Generalized lymphadenopathy is present, and migratory arthropathy is often present.
- Fever resolves in 5 to 10 days, but may relapse repeatedly over a period of weeks to months.

One or more relapses follow, and are associated with a purple papular exanthem on the chest and arms.

- Additional findings include generalized hyperreflexia, migratory polyarthralgia (over 50% of cases), myalgia and hyperesthesia.
- Arthritis affects more than one joint in 83.3% of patients, involving the knee in most.³
- Rare instances of endocarditis^{4 5}, psoas abscess, epidural abscess⁶ and spondylodiscitis have been reported.⁷

The fatality rate without treatment is 10%, and results from endocarditis or multiple visceral abscesses.

Features which may distinguish spirillary [S] from streptobacillary [B] rat bite fever include the following:^{8 9}

- | | |
|------------|------------------------------------|
| incubation | - S up to 30 days |
| | - B up to 10 days |
| bite wound | - S may produce a chancre |
| | - B heals promptly |
| relapses | - S regular |
| | - B intermittent |
| rash | - S generalized macular |
| | - B macular, pustular or petechial |
| arthritis | - S rare |
| | - B common ¹⁰ |

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 2007 Jan ;20(1):13-22.
2. Clin Microbiol Rev 2007 Jan ;20(1):13-22.
3. BMC Infect Dis 2007 ;7:56.
4. Emerg Infect Dis 2006 Jun ;12(6):1037-8.
5. J Clin Microbiol 2007 Sep ;45(9):3125-6.
6. J Clin Microbiol 2012 Sep ;50(9):3122-4.
7. J Clin Microbiol 2008 Aug ;46(8):2820-1.
8. Scand J Infect Dis 2002 ;34(6):474-7.
9. Lancet Infect Dis 2001 Sep ;1(2):91.
10. Clin Orthop Relat Res 2000 Nov ;(380):173-6.

Relapsing fever

Agent	BACTERIUM. <i>Borrelia</i> spp. A microaerophilic spirochete
Reservoir	Human Tick Rodent
Vector	Tick (<i>Ornithodoros</i>), louse (<i>Pediculus</i>)
Vehicle	Blood products
Incubation Period	7d - 8d (range 2d - 18d)
Diagnostic Tests	Examination of blood smears (thick and thin smears). Some species (<i>B. hermsii</i>) may grow in BSK II medium.
Typical Adult Therapy	Doxycycline 100 mg PO BID X 7d. OR Erythromycin 500 mg QID X 7d
Typical Pediatric Therapy	Chloramphenicol 12.5 mg/kg PO QID X 7d. OR Erythromycin 10 mg/kg QID X 7d
Clinical Hints	Headache, myalgia, hepatosplenomegaly, rash and relapsing illness; louse-borne (vs. tick borne) characterized by higher case fatality rate, fewer relapses and higher incidence of hepatosplenomegaly, jaundice and neurological complications.
Synonyms	Bilious typhoid, <i>Borrelia microti</i> , <i>Borrelia miyamotoi</i> , <i>Borrelia turicatae</i> , Borreliosis, Famine fever, Febbre recidiva, Febbre ricorrente, Febris recurrens, Fiebre recurrente, Lauseruckfallfieber, Mianeh fever, Ruckfall fieber, Tilbakefallsfever, Vagabond fever, Yellow famine fever, Yellow plague. ICD9: 087.9,087.0,087.1 ICD10: A68

Clinical

The clinical manifestations of louse-borne and tick-borne ¹ relapsing fevers are similar. ^{2 3}

- Louse-borne disease is characterized by a longer incubation period, longer febrile periods and afebrile intervals, and fewer relapses.
- Both types have an acute onset of high fever with rigors, headache, myalgia, arthralgia, photophobia and cough.
- In Africa, tickborne relapsing fever is often mis-diagnosed as malaria. ⁴

Physical findings often include conjunctivitis, petechiae, and abdominal tenderness with hepatomegaly and splenomegaly.


- Nuchal rigidity, pulmonary rales, lymphadenopathy, jaundice and ARDS ⁵ are occasionally encountered.
- Hemorrhagic phenomena are common but rarely severe.
- Iritis and iridocyclitis may lead to permanent impairment of vision. Uveitis is also described. ⁶
- A petechial, macular, or papular rash over the trunk may be noted toward the end of the illness.
- As many as 30% of patients develop neurological findings such as coma, cranial nerve palsies, hemiplegia, meningitis, and seizures. ⁷
- Rare instances of acute respiratory distress syndrome ^{8 9} and dermal eschar ¹⁰ have been associated with tick-borne relapsing fever.
- Deaths are ascribed to myocarditis with associated arrhythmias, cerebral hemorrhage or hepatic failure.
- "Tropical thrombophlebitis" has been associated with outbreaks of relapsing fever in South Africa. ¹¹

Borrelia miyamotoi infection is characterized by fever, headache, myalgia and multiple relapses. ¹²

- *B. miyamotoi* meningoenzephalitis was reported in an immunocompromised patient. ¹³

This disease is endemic or potentially endemic to 118 countries.

Relapsing fever in Uganda



Graph: Uganda. Relapsing fever, cases

Notes:

1. Historical data from reference ¹⁴

Both louse-borne and tick-borne disease are reported.

Borrelia duttoni (vector *Ornithodoros erraticus*) has been identified in Uganda. ¹⁵

References

1. Infect Dis Clin North Am 2008 Sep ;22(3):449-68, viii.
2. Tex Med 1995 May ;91(5):56-9.
3. Trop Geogr Med 1995 ;47(2):49-52.
4. Emerg Infect Dis 2007 Jan ;13(1):117-23.
5. MMWR Morb Mortal Wkly Rep 2007 Oct 19;56(41):1073-6.
6. Am J Ophthalmol 2006 Aug ;142(2):348-9.
7. Emerg Infect Dis 2013 Feb ;19(2):301-4.
8. Wilderness Environ Med 2008 ;19(4):280-6.
9. Infection 2012 Dec ;40(6):695-7.
10. J Travel Med 2012 Jul ;19(4):261-3.
11. S Afr Med J 1975 Nov 15;49(49):2057-8.
12. ProMED <promedmail.org> archive: 20130118.1504740
13. N Engl J Med 2013 Jan 17;368(3):240-5.
14. Bull World Health Organ 1958 ;19(4):673-710.
15. East Afr Med J 1952 Nov ;29(11):477-9.

Respiratory syncytial virus infection

Agent	VIRUS - RNA. Paramyxoviridae, Pneumovirinae: Human respiratory syncytial virus
Reservoir	Human
Vector	None
Vehicle	Droplet Infected secretions (hands)
Incubation Period	2d - 8d
Diagnostic Tests	Viral culture or DFA (nasal and other respiratory secretions). Serology. Nucleic acid amplification.
Typical Adult Therapy	Ribavirin aerosol 20 mg/ml for 12h/d X 3 to 5d [severe infections]. Effectiveness not proven
Typical Pediatric Therapy	As for adult
Vaccine	RSV immune globulin
Clinical Hints	Rhinorrhea, cough, wheezing, bronchiolitis and respiratory distress; encountered primarily in infancy.
Synonyms	Chimpanzee coryza agent, Respiratory syncytial virus, RSV. ICD9: 079.6,480.1 ICD10: B97.4,J12.1

Clinical

RSV infections are manifested as:

- lower respiratory tract disease (pneumonia, bronchiolitis, tracheobronchitis)
- upper respiratory tract illness, often accompanied by fever and otitis media. ¹

Asymptomatic infection is rare.

- Pneumonia or bronchiolitis occurs in 30% to 71% of patients (89% among closed populations of infants).
- Croup accounts for only 5% to 10% of cases.
- Wheezing ²⁻⁴, rhonchi, rales, and pulmonary infiltrates are encountered with bronchiolitis as well as pneumonia. ⁵
- Bronchiolitis is characterized by wheezing and hyperaeration of the lung.
- RSV infection in adults is usually mild; however severe disease may develop. ⁶⁻⁸

Lower respiratory tract infection is heralded by nasal congestion and often pharyngitis.

- Fever occurs in young children, with temperatures ranging from 38 to 40C.
- Fever is present for 2 to 4 days; however, the extent and duration of the fever does not correlate with the severity of the disease.
- Fever is frequently absent at the time of admission to the hospital.
- Cough is often a predominant sign.
- The cough may be paroxysmal and associated with vomiting, but without the "whoop" typical of pertussis.
- Laryngitis and hoarseness are not common.

Dyspnea, increased respiratory rate, and retractions of the intercostal muscles are common.

- In bronchiolitis, expiration is prolonged, and the respiratory rate may be remarkably elevated. ⁹
- Intercostal retractions are also prominent in bronchiolitis.
- On auscultation, the infant may have crackles and wheezing, which may be present intermittently and may fluctuate in intensity.
- Cyanosis is rare, despite hypoxemia. In most infants, the duration of illness is 7 to 21 days, and hospitalization, if required, averages 3 to 7 days.
- Thrombocytosis is common among children hospitalized with RSV bronchiolitis. ¹⁰
- The severity and / or duration of RSV bronchiolitis is exacerbated by concomitant human metapneumovirus infection. ¹¹⁻¹⁴
- RSV infection accounts for approximately 5% of bronchiolitis obliterans in children (Beijing, 2001 to 2007) ¹⁵
- Infection in premature infants may result in long term effects on airway function. ¹⁶⁻¹⁸

Otitis media is a common complication of RSV infection in young children. ¹⁹⁻²²

- Viral meningitis ²³, encephalopathy and seizures are also encountered. ²⁴⁻²⁶
- Repeated or secondary infections occurring after the first 3 years of life are most commonly manifested as an upper respiratory tract illness or tracheobronchitis.

- Young adults may present with flu-like illness, pneumonia, chronic cough suggestive of tracheobronchitis or bronchitis, and occasionally with otitis. ²⁷
- Infection among the elderly is often nosocomially acquired, and may result in pneumonia in 5% to 50% of the cases, with a fatal outcome in up to 20%.
- Additional extrapulmonary manifestations of RSV infection include myocarditis ²⁸, supraventricular tachycardia, ventricular tachycardias, pericardial effusion ²⁹, seizures, focal neurological abnormalities, hyponatremia and hepatitis ³⁰

Signs and symptoms of Human Metapneumovirus (hMPV) infection are similar to those of Respiratory syncytial virus infection ^{31 32}, and coinfection by these two agents may be particularly severe. ³³⁻³⁷ Children with hMPV infection are likely to be older than those with RSV, and more likely to present with pneumonia and less likely to present with bronchiolitis. ³⁸

- Clinical signs of Rhinovirus infection ³⁹ and of Human Bocavirus infection are also similar to those of Respiratory syncytial virus infection; however, hypoxia, and neutrophilia may be more common in Human Bocavirus infection. ⁴⁰
- Superinfection of RSV by *Staphylococcus aureus* ⁴¹, *Bordetella pertussis* ⁴² and other bacteria is not unusual. ⁴³

This disease is endemic or potentially endemic to all countries.

References

1. Paediatr Respir Rev 2004 ;5 Suppl A:S119-26.
2. Pediatr Int 2008 Oct ;50(5):654-7.
3. Early Hum Dev 2011 Mar ;87 Suppl 1:S51-4.
4. Early Hum Dev 2011 Mar ;87 Suppl 1:S51-4.
5. Pediatr Infect Dis J 2003 Feb ;22(2 Suppl):S94-9.
6. Semin Respir Crit Care Med 2011 Aug ;32(4):423-32.
7. Infect Disord Drug Targets 2012 Feb 15;
8. J Infect Dis 2012 Apr 23;
9. J Pediatr 2003 Nov ;143(5 Suppl):S112-7.
10. Isr Med Assoc J 2010 Jan ;12(1):39-41.
11. J Infect Dis 2005 Feb 1;191(3):382-6.
12. Emerg Infect Dis 2003 Mar ;9(3):372-5.
13. Emerg Infect Dis 2004 Jul ;10(7):1318-20.
14. Pediatr Pulmonol 2007 Aug ;42(8):740-3.
15. Zhonghua Er Ke Za Zhi 2008 Oct ;46(10):732-8.
16. Thorax 2009 Jun ;64(6):490-5.
17. Clin Ther 2010 Dec ;32(14):2422-32.
18. J Allergy Clin Immunol 2012 Mar 21;
19. Vaccine 2007 Feb 19;25(9):1683-9.
20. Acta Paediatr 2010 Jun ;99(6):867-70.
21. Acta Paediatr 1995 Apr ;84(4):419-23.
22. J Clin Microbiol 2011 Nov ;49(11):3750-5.
23. Kansenshogaku Zasshi 2011 Nov ;85(6):682-5.
24. Minerva Pediatr 2005 Jun ;57(3):137-42.
25. J Child Neurol 2009 Dec ;24(12):1499-503.
26. J Infect Chemother 2011 Dec ;17(6):776-81.
27. Semin Respir Crit Care Med 2007 Apr ;28(2):171-81.
28. Fetal Pediatr Pathol 2011 ;30(1):64-8.
29. Cardiol Young 2012 Jun 14;:1-2.
30. Crit Care 2006 ;10(4):R107.
31. Pediatr Infect Dis J 2004 Jan ;23(1 Suppl):S25-32.
32. Clin Microbiol Rev 2006 Jul ;19(3):546-57.
33. Curr Opin Infect Dis 2005 Apr ;18(2):141-6.
34. J Infect Dis 2005 Feb 1;191(3):382-6.
35. Emerg Infect Dis 2003 Mar ;9(3):372-5.
36. Emerg Infect Dis 2004 Jul ;10(7):1318-20.
37. ProMED <promedmail.org> archive: 20120223.1050554
38. J Paediatr Child Health 2011 Oct ;47(10):737-41.
39. Pediatr Infect Dis J 2012 Jan ;31(1):84-6.
40. Eur J Pediatr 2010 Sep ;169(9):1087-92.
41. Pediatr Infect Dis J 2010 Nov ;29(11):1048-50.
42. Pediatr Infect Dis J 2007 Apr ;26(4):316-8.
43. J Infect Chemother 2011 Feb ;17(1):87-90.

Respiratory viruses - miscellaneous

Agent	VIRUS - RNA and DNA Pneumovirinae: Human Metapneumovirus Coronaviridae: New Haven Coronavirus, HKU1 Parvovirinae: Human Bocavirus
Reservoir	Human
Vector	None
Vehicle	Droplet Infected secretions (on hands)
Incubation Period	Unknown
Diagnostic Tests	Viral culture. Serology. Nucleic acid amplification.
Typical Adult Therapy	NA
Typical Pediatric Therapy	NA
Clinical Hints	Rhinorrhea, cough, wheezing, bronchiolitis and respiratory distress; encountered primarily in infancy.
Synonyms	Acanthamoeba polyphaga mimivirus, Bat reovirus, Bocavirus, Bradford coccus, Cardiovirus, Coronavirus HKU1, Coronavirus NL63, Encephalomyocarditis Virus, HCoV-HKU1, HCoV-NL63, HK23629/07, HKU1, HRV-A, HRV-B, HRV-C, Human Bocavirus, Human Coronavirus NL63, Human CoV 229E, Human CoV OC43, Human metapneumovirus, Human rhinovirus, Kampar, Karolinska Institutet virus, KI virus, Melaka, Metapneumovirus, Mimivirus, New Haven coronavirus, Pulau, Rhinovirus, Small Anellovirus, Tioman virus, Torque tenovirus, Torquetenovirus, Washington University virus, WU polyomavirus, WU virus. ICD9: 079.89 ICD10: B34.2,J12.8

Clinical

For a comprehensive review of newer respiratory viral infections, see ¹

Human Metapneumovirus:

Signs and symptoms of Human Metapneumovirus (hMPV) infection are similar to those of Respiratory syncytial virus infection ^{2 3}, and coinfection by these two agents may be relatively severe and / or prolonged. ⁴⁻⁹ Children with hMPV infection are likely to be older than those with RSV, and more likely to present with pneumonia and less likely to present with bronchiolitis. ¹⁰

- Findings include either lower respiratory tract disease (pneumonia, bronchiolitis, tracheobronchitis) or upper respiratory tract illness, often accompanied by fever and otitis media. ^{11 12}
- Asymptomatic infection is reported. ^{13 14}
- Wheezing, rhonchi, rales, and pulmonary infiltrates are encountered with bronchiolitis, hyperaeration and pneumonia. ¹⁵ Severe and potentially-fatal infections are reported. ¹⁶
- Apnea has been reported in newborn infants. ¹⁷
- hMPV has been recovered from the middle ear in patients with otitis media. ¹⁸ and is associated with 6% of otitis media cases in children. ¹⁹
- Central nervous system disease has been reported, ranging from febrile seizures to fatal encephalitis. ²⁰⁻²²
- Reinfection is common. ²³⁻²⁵
- Although infection in adults is usually mild or asymptomatic ²⁶, severe disease is reported in elderly adults with underlying disease. ²⁷⁻²⁹

New Haven coronavirus:

New Haven coronavirus infection is characterized by fever, cough and rhinorrhea. ^{30 31}

- Tachypnea, hypoxia and pulmonary infiltrates may be present.
- The agent has also been identified as a common cause for croup. ³²

Coronavirus infections:

HKU1 (HCoV-HKU1), a human coronavirus, was isolated in Hong Kong in 2005, from two adult patients with pneumonia. ³³

- An additional 6 cases in Hong Kong were characterized by gastroenteritis, fever, otitis and febrile seizures.
- Human Coronavirus OC43 infection is associated with fever, rhinitis, pharyngitis, laryngitis, otitis, bronchitis, bronchiolitis or pneumonia. ³⁴

Human Bocavirus:

Human Bocavirus is a common cause of lower respiratory tract infection in children. ^{35 36}

- Bocavirus infections, including cases of severe pneumonia, have also been reported in adults. ³⁷
- Patients are often co-infected by Respiratory syncytial virus, Adenovirus, Influenza virus, Human metapneumovirus or other pathogens. ³⁸
- Clinical presentation may include fever, cough, rhinorrhea, conjunctivitis, wheezing, respiratory distress, pneumonia or pleural effusion. ³⁹
- Rarely, severe and life-threatening infection is encountered. ⁴⁰
- Human Bocavirus infection may mimic the symptoms of pertussis ⁴¹
- Rare instances of Human Bocavirus encephalitis have been reported. ⁴²
- Clinical signs are also similar to those of Respiratory syncytial virus infection; however, hypoxia, and neutrophilia may be more common in Human Bocavirus infection. ⁴³
- Disseminated Bocavirus infection, including diarrhea and viremia, has been reported in a stem cell transplant patient. ⁴⁴

Other viruses:

Although Rhinovirus infection is usually associated with the common cold, infection may be associated with severe lower respiratory tract infections ⁴⁵, and outbreaks of major and even fatal disease have been reported in chronic care facilities. ⁴⁶⁻⁴⁹

Melaka virus, a bat-associated Reovirus, has been identified as a cause of fever and acute respiratory tract infection in Malaysia. ⁵⁰

Saffold Cardiovirus, a member of the Picornaviridae, has been associated with cases of upper respiratory tract infection in children. ^{51 52}

- Human infection by an additional Cardiovirus, Encephalomyocarditis Virus, have been characterized by fever, headache, nausea and dyspnea. (2009 publication) ⁵³ One such patient also experienced weight loss, arthralgia, photophobia, myalgia, chills, vomiting, and abdominal pain.

This disease is endemic or potentially endemic to all countries.

References

1. Clin Microbiol Rev 2008 Apr ;21(2):274-90, table of contents.
2. Pediatr Infect Dis J 2004 Jan ;23(1 Suppl):S25-32.
3. Clin Microbiol Rev 2006 Jul ;19(3):546-57.
4. Curr Opin Infect Dis 2005 Apr ;18(2):141-6.
5. J Infect Dis 2005 Feb 1;191(3):382-6.
6. Emerg Infect Dis 2003 Mar ;9(3):372-5.
7. Emerg Infect Dis 2004 Jul ;10(7):1318-20.
8. Pediatr Pulmonol 2007 Aug ;42(8):740-3.
9. ProMED <promedmail.org> archive: 20120223.1050554
10. J Paediatr Child Health 2011 Oct ;47(10):737-41.
11. Semin Respir Crit Care Med 2007 Apr ;28(2):213-21.
12. Emerg Infect Dis 2004 Apr ;10(4):700-5.
13. J Infect Dis 2006 Aug 15;194(4):474-8.
14. New Microbiol 2009 Jul ;32(3):297-301.
15. Infect Dis Clin North Am 2005 Sep ;19(3):569-84.
16. Case Rep Pediatr 2012 ;2012:268074.
17. Rev Chilena Infectol 2007 Aug ;24(4):313-8.
18. Pediatr Infect Dis J 2005 Jul ;24(7):655-7.
19. Int J Pediatr Otorhinolaryngol 2006 Jul ;70(7):1189-93.
20. Emerg Infect Dis 2005 Mar ;11(3):467-70.
21. Pediatr Infect Dis J 2009 Dec ;28(12):1057-60.
22. Arch Neurol 2012 May ;69(5):649-52.
23. Uirusu 2006 Dec ;56(2):173-81.
24. J Infect Dis 2008 Sep 15;198(6):836-42.
25. Semin Respir Crit Care Med 2007 Apr ;28(2):213-21.
26. Arch Intern Med 2008 Dec 8;168(22):2489-96.
27. Pediatr Infect Dis J 2008 Oct ;27(10 Suppl):S80-3.
28. Rev Chilena Infectol 2011 Apr ;28(2):174-8.
29. Viruses 2013 ;5(1):87-110.
30. Curr Opin Infect Dis 2005 Apr ;18(2):141-6.
31. Curr Opin Pediatr 2006 Feb ;18(1):42-7.
32. ProMED <promedmail.org> archive: 20050825.2512
33. Clin Infect Dis 2006 Mar 1;42(5):634-9.
34. J Clin Virol 2008 Jul ;42(3):233-43.
35. Pediatr Infect Dis J 2007 Aug ;26(8):745-6.
36. ProMED <promedmail.org> archive: 20050824.2494
37. Emerg Infect Dis 2006 Oct ;12(10):1614-6.
38. Clin Microbiol Rev 2008 Apr ;21(2):291-304, table of contents.
39. J Clin Microbiol 2011 Mar ;49(3):1179-81.
40. Emerg Infect Dis 2011 Dec ;17(12):2303-5.
41. Pediatrics 2008 Mar ;121(3):e631-7.
42. Clin Infect Dis 2012 Apr ;54(7):964-7.
43. Eur J Pediatr 2010 Sep ;169(9):1087-92.
44. Emerg Infect Dis 2007 Sep ;13(9):1425-7.
45. Pediatr Infect Dis J 2009 Apr ;28(4):337-9.
46. Ann Intern Med 1995 Oct 15;123(8):588-93.
47. J Clin Virol 2007 Mar ;38(3):227-37.
48. Clin Infect Dis 2005 Jul 15;41(2):262-5.
49. J Am Geriatr Soc 2006 Feb ;54(2):284-9.
50. ProMED <promedmail.org> archive: 20070626.2063
51. Emerg Infect Dis 2008 May ;14(5):834-6.
52. Emerg Infect Dis 2008 Sep ;14(9):1398-405.
53. Emerg Infect Dis 2009 Apr ;15(4):640-6.

Reye's syndrome

Agent	UNKNOWN
Reservoir	Unknown
Vector	None
Vehicle	Unknown
Incubation Period	Unknown
Diagnostic Tests	Clinical diagnosis.
Typical Adult Therapy	Electrolyte & glucose management, ? enemas, ? dialysis
Typical Pediatric Therapy	As for adult
Clinical Hints	Vomiting, lethargy, coma, seizures, hepatomegaly, hypoglycemia and elevated blood ammonia concentration; usually anicteric; follows viral infection; aspirin ingestion is often implicated.
Synonyms	Reye syndrome. ICD9: 331.81 ICD10: G93.7

Clinical

Signs and symptoms of Reye's syndrome include protracted vomiting and encephalopathy, in the absence of fever or jaundice. ^{1 2}

- Hepatomegaly is present in 50% of cases.
- Twelve hours to 3 weeks following an antecedent viral illness, the patient develops vomiting and lethargy, followed by restlessness, irritability, combativeness, disorientation, delirium, tachycardia, hyperventilation, dilated pupils with sluggish response, hyperreflexia, positive Babinski sign, and appropriate response to noxious stimuli.

Diarrhea and hyperventilation are often the first signs in children below age 2 years.

- Later, obtundation, coma and decorticate rigidity are associated with inappropriate response to noxious stimuli.
- Coma deepens, and the patient is found to have fixed and dilated pupils, loss of oculovestibular reflexes and dysconjugate gaze with caloric stimulation.
- Seizures ensue, with flaccid paralysis, absent deep tendon reflexes, lack of pupillary response and respiratory arrest.

Similar disease (Reye-like syndrome) is caused by inborn errors of metabolism, hypoglycemia, hypoketonemia, elevated ammonia, and organic aciduria. ³

- A case of encephalopathy and hepatic failure • similar to Reye's syndrome • was related to *Bacillus cereus* food poisoning. ⁴

This disease is endemic or potentially endemic to all countries.

References

1. Eur J Pediatr 2000 Sep ;159(9):641-8.
2. N Engl J Med 1999 Sep 9;341(11):846-7.
3. Pediatr Neurol 2008 Sep ;39(3):198-200.
4. Brain Dev 2010 Sep ;32(8):688-90.

Rheumatic fever

Agent	BACTERIUM. Streptococcus pyogenes A facultative gram-positive coccus
Reservoir	Human
Vector	None
Vehicle	Droplet
Incubation Period	1w - 5w
Diagnostic Tests	Clinical diagnosis.
Typical Adult Therapy	Supportive; salicylates
Typical Pediatric Therapy	As for adult
Clinical Hints	Migratory arthritis, fever, carditis, chorea, subcutaneous nodules, erythema marginatum and leukocytosis; follows overt pharyngitis after 1 to 5 weeks in most cases; acute attack persists for approximately 3 months.
Synonyms	Febbre reumatica. ICD9: 390,391 ICD10: I00,I01,I02

Clinical

Case definition for surveillance:

The CDC (The United States Centers for Disease Control) case definition for surveillance requires evidence for preceding group A streptococcal infection (culture, serology) in addition to two major clinical criteria; or one major and two minor criteria, as follows:

Major clinical criteria:

- carditis
- polyarthritis
- chorea ^{1 2}
- subcutaneous nodules
- erythema marginatum. ³

Minor criteria:

- previous rheumatic fever or rheumatic heart disease
- arthralgia
- fever
- elevation of erythrocyte sedimentation rate [ESR]
- positive C-reactive protein
- leucocytosis
- prolongation of the P-R interval on electrocardiogram.

This disease is endemic or potentially endemic to all countries.

References

1. S Afr Med J 1997 Jun ;87 Suppl 3:C157-60.
2. Rev Neurol 2004 Nov 1-15;39(9):810-5.
3. Rheum Dis Clin North Am 1997 Aug ;23(3):545-68.

Rhinoscleroma and ozena

Agent	BACTERIUM. Klebsiella pneumoniae ssp ozaenae and Klebsiella pneumoniae ssp rhinoscleromatis Facultative gram-negative bacilli
Reservoir	Human
Vector	None
Vehicle	Infected secretions
Incubation Period	Unknown
Diagnostic Tests	Culture. Biopsy. Nucleic acid amplification. Advise laboratory when this diagnosis is suspected.
Typical Adult Therapy	Rhinoscleroma: Streptomycin , often with systemic or topical Rifampin - for 3 to 6 weeks; fluoroquinolones also appear to be effective. Ozena: Ciprofloxacin or Sulfamethoxazole/trimethoprim for 3 months
Typical Pediatric Therapy	As for adult
Clinical Hints	Rhinorrhea associated with a painless intranasal mass; may extend to sinuses or ears.
Synonyms	Klebsiella pneumoniae ssp ozaenae , Ozena, Rhinoscleroma. ICD9: 040.1 ICD10: J31.0

Clinical

The nose is involved in over 90% of cases of rhinoscleroma.

- Findings include fetid discharge, a crusting granulomatous mass and cicatrization. ^{1 2}
- The pharynx is involved in 15% to 40%, the larynx in 2% to 2%, the tracheobronchial tree in 15% ³ and the paranasal sinuses in 2% to 25%. ⁴
- Rare instances of laryngeal stenosis resulting from rhinoscleroma are reported. ⁵
- Standard therapy consists of streptomycin in combination with topical or systemic rifampicin, for at least 3 to 6 weeks.
- Recent studies suggest that fluoroquinolones are also effective.

Ozena (primary atrophic rhinitis) is characterized by progressive atrophy of the nasal mucosa and underlying bone.

- Findings include foul-smelling, thick, dry crusts and greatly enlarged nasal cavities. ⁶
- Laryngeal involvement has been reported. ⁷
- Ozena may be associated with tracheobronchopathia osteochondroplastica ⁸
- Rare instances of disseminated systemic infection are reported. ⁹

This disease is endemic or potentially endemic to all countries.

Rhinoscleroma and ozena in Uganda

Highest rates of rhinoscleroma are found in Itesot and among tribes in the southwestern region. ¹⁰

85 cases were documented during 1972 to 1981.

The disease is most common among females and middle aged adults.

References

- South Med J 1988 Dec ;81(12):1580-2.
- Laryngoscope 1982 Oct ;92(10 Pt 1):1149-53.
- Ann Otol Rhinol Laryngol 1996 May ;105(5):336-40.
- Rhinology 2008 Dec ;46(4):338-41.
- Acta Otorrinolaringol Esp 2010 May-Jun;61(3):241-3.
- J Otolaryngol 1990 Oct ;19(5):345-9.
- J Clin Microbiol 2005 Nov ;43(11):5811-3.
- Rev Mal Respir 2007 Sep ;24(7):883-7.
- Malays J Pathol 2009 Dec ;31(2):147-50.
- East Afr Med J 1993 Mar ;70(3):186-8.

Rhinosporidiosis

Agent	PROTOCTISTA Rhinosporidium seeberi [may in fact be Microcystis, a cyanobacterium]
Reservoir	Water Soil Vegetation
Vector	None
Vehicle	Aerosol from soil or water
Incubation Period	2w - 6m
Diagnostic Tests	Histology of resected material (organism does not grow in-vitro).
Typical Adult Therapy	Excision
Typical Pediatric Therapy	As for adult
Clinical Hints	Friable, painless vascular masses of nose, conjunctivae and larynx; recurrence is common.
Synonyms	Rhinosporidium seeberi. ICD9: 117.0 ICD10: B48.1

Clinical

Clinical forms include:

- nasal (chronic, painless unilateral obstruction, mucoid discharge) ¹
- conjunctival (usually palpebral) ² or lacrimal lesion. ^{3 4}
- ENT (mucous membrane mass of the epiglottis, tongue, palate, tonsil, uvula, larynx ^{5 6}, paranasal sinuses. ⁷
- urethral (predominantly male), presenting as a painless, friable polyp of the fossa navicularis. ⁸

Multiple painless dermal or subcutaneous nodules may be present. ⁹⁻¹⁴

- Rarely, skin lesions may be polymorphic ¹⁵
- Additional manifestations have included primary cutaneous lesions ¹⁶, osteomyelitis ¹⁷⁻¹⁹, bronchial mass lesion ²⁰, obstructive tracheitis ²¹ and infection of the parotid duct. ²²⁻²⁵

Relapse occurs in approximately 10% of cases following excision.

Signs of mucosal chromomycosis may mimic those of rhinosporidiosis. ²⁶

This disease is endemic or potentially endemic to 65 countries.

Rhinosporidiosis in Uganda

A review of 51 cases was published in 1978. ²⁷

References

1. Indian J Pathol Microbiol 2001 Jan ;44(1):17-21.
2. J Indian Med Assoc 2003 Nov ;101(11):667-8, 670.
3. Ophthal Plast Reconstr Surg 2009 May-Jun;25(3):234-5.
4. Indian J Otolaryngol Head Neck Surg 2011 Jul ;63(3):243-6.
5. Ear Nose Throat J 2004 Aug ;83(8):568, 570.
6. Indian J Otolaryngol Head Neck Surg 2011 Jul ;63(3):243-6.
7. J Laryngol Otol 2010 Oct ;124(10):1139-41.
8. J Coll Physicians Surg Pak 2008 May ;18(5):314-5.
9. J Dermatol 1998 Aug ;25(8):527-32.
10. J Eur Acad Dermatol Venereol 2006 Jan ;20(1):88-9.
11. Indian J Dermatol Venereol Leprol 2007 May-Jun;73(3):185-7.
12. Indian J Dermatol Venereol Leprol 2007 May-Jun;73(3):179-81.
13. Indian J Dermatol Venereol Leprol 2007 Sep-Oct;73(5):343-5.
14. Australas J Dermatol 2011 May ;52(2):e4-6.
15. Indian J Dermatol Venereol Leprol 2008 May-Jun;74(3):298.
16. Diagn Cytopathol 2009 Feb ;37(2):125-7.
17. Indian J Pathol Microbiol 2005 Apr ;48(2):215-7.
18. J Orthop Surg (Hong Kong) 2008 Apr ;16(1):99-101.
19. Skeletal Radiol 2011 Feb ;40(2):225-8.
20. Lung India 2012 Apr ;29(2):173-5.
21. J Laryngol Otol 2008 Apr ;122(4):e13.
22. Indian J Pathol Microbiol 2007 Apr ;50(2):320-2.
23. Indian J Dent Res 2009 Jul-Sep;20(3):388-9.
24. Indian J Med Microbiol 2012 Jan ;30(1):108-11.
25. J Investig Clin Dent 2013 Feb 1;
26. J Clin Pathol 1960 Jul ;13:287-90.
27. East Afr Med J 1978 Dec ;55(12):582-6.

Rhodococcus equi infection

Agent	BACTERIUM. Rhodococcus equi An aerobic gram-positive coccobacillus
Reservoir	Farm animal Farm soil
Vector	None
Vehicle	? Inhalation Contact Ingestion
Incubation Period	Unknown
Diagnostic Tests	Culture of blood, body fluids and secretions. Advise laboratory when these organisms are suspected.
Typical Adult Therapy	Vancomycin 500 mg q8h. Alternatives: Erythromycin , Gentamicin , Rifampin
Typical Pediatric Therapy	Vancomycin 10 mg/kg q6h. Alternatives: Erythromycin , Gentamicin , Rifampin
Clinical Hints	Most often encountered as pleuropulmonary infection in an immune-suppressed patient; history of contact with farm or farm animals in 40% of cases.
Synonyms	Rhodococcus. ICD9: 027.9 ICD10: A92.8

Clinical

The clinical features of *Rhodococcus equi* disease are largely determined by the site of infection and clinical substrate in which it occurs. ^{1 2}

- 49% of patients are HIV-positive.
- Pulmonary infection predominates among HIV-positive patients ³
- Extrapulmonary disease (abscesses, septicemia, eye or wound infection, etc) is most common in immunocompetent individuals. ⁴

This disease is endemic or potentially endemic to all countries.

References

1. Clin Transplant 2004 Dec ;18(6):748-52.
2. Emerg Infect Dis 1997 Apr-Jun;3(2):145-53.
3. Clin Microbiol Infect 1997 Feb ;3(1):12-18.
4. J Med Case Reports 2011 ;5:358.

Rift Valley fever

Agent	VIRUS - RNA. Bunyaviridae, Phlebovirus: Rift Valley fever virus
Reservoir	Sheep Ruminant
Vector	Mosquito (Culex, Aedes, Anopheles, Eretmapodites, Mansonia, Culicoides, Coquillettia spp.)
Vehicle	None
Incubation Period	3d - 5d (range 2d - 7d)
Diagnostic Tests	Viral culture (blood, CSF). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive. Animal studies suggest a possible role for Ribavirin .
Typical Pediatric Therapy	As for adult
Vaccine	Rift Valley fever
Clinical Hints	Headache, myalgia, photophobia, arthralgia and a maculopapular rash; occasional jaundice and retinitis; history of contact with sheep or cattle during the preceding week may be elicited; case fatality rate = 0.1%.
Synonyms	Arumowot, Enzootic hepatitis, Gabek Forest, Gordil, Riftvalleykoorts, Zinga. ICD9: 066.3 ICD10: A92.4

Clinical

Disease is heralded by a 'flu-like' illness with sudden onset of fever, headache, myalgia and back pain. ^{1 2}

- Following an incubation period of 2 to 6 days, the patient may develop a mild, flu-like illness which may mimic dengue fever or viral meningitis.
- A characteristic syndrome consists of fever, large-joint arthralgia, and gastrointestinal complaints followed by jaundice, right upper-quadrant pain, and delirium, often coinciding with hemorrhagic manifestations. ³
- Nuchal rigidity, arthralgia, myalgia and photophobia may be present.
- Retinitis occurs in 15% of patients, and is characterized by macular, paramacular, and/or extramacular lesions, often occurring bilaterally. Hemorrhage and edema are often present, and vasculitis, vascular occlusion and optic atrophy are also observed. ⁴⁻⁶

Complications include hemorrhagic fever ⁷ on the second to fourth day of illness; or retinal hemorrhage or meningoencephalitis appearing after the first week. ⁸⁻¹⁰

- Hemorrhagic phenomena and fatal encephalitis have been observed in approximately 1% to 2% of patients during epidemics and account for much of the mortality.
- Renal dysfunction is encountered in 60% of cases. ¹¹
- The case-fatality rate in epidemics is usually below 1%.

This disease is endemic or potentially endemic to 34 countries.

References

1. Rev Infect Dis 1989 May-Jun;11 Suppl 4:S777-82.
2. Wkly Epidemiol Rec 2008 Jan 11;83(2):17-22.
3. Am J Trop Med Hyg 2010 Mar ;82(3):371-5.
4. Br J Ophthalmol 1980 May ;64(5):366-74.
5. Ann Saudi Med 1997 May ;17(3):377-80.
6. Ophthalmology 2005 Feb ;112(2):313-8.
7. Ugeskr Laeger 2007 Jun 25;169(26):2537-8.
8. Bull Soc Pathol Exot Filiales 1989 ;82(5):620-7.
9. Bull Soc Pathol Exot Filiales 1989 ;82(5):611-9.
10. Bull Soc Pathol Exot Filiales 1989 ;82(5):605-10.
11. Saudi J Kidney Dis Transpl 2009 Nov ;20(6):1047-52.

Rotavirus infection

Agent	VIRUS - RNA. Reoviridae: Rotavirus
Reservoir	Human
Vector	None
Vehicle	Fecal-oral Water
Incubation Period	12h - 3d
Diagnostic Tests	Stool assay for viral antigen. Serology. Nucleic acid amplification.
Typical Adult Therapy	Stool precautions; supportive
Typical Pediatric Therapy	As for adult
Vaccine	Rotavirus
Clinical Hints	Vomiting, diarrhea and mild fever: the illness lasts approximately 1 week, and is most severe in infancy; fatal cases are associated with dehydration and electrolyte imbalance.
Synonyms	Rotavirus. ICD9: 008.61 ICD10: A08.0

Clinical

Infants and young children present with fever, vomiting, diarrhea, and occasionally dehydration. ¹

- Most hospitalized patients had experienced fever and vomiting for 2 to 3 days, and diarrhea for 4 to 5 days.
- The diarrhea is watery without blood or mucus.
- Leukocytes are detected in the stool in a small percentage of patients.
- Approximately 36% of episodes are characterized by 'dehydrating diarrhea.'
- Viremia is present in over 50% of patients with Rotavirus diarrhea. ^{2 3}
- Asymptomatic infection is common. ⁴

Infection in immunodeficient children may persist for weeks to months.

Rotavirus infection is not unusual in adults. ⁵

Complications:

- Rotavirus infection increases the risk of bacteremia in children with nontyphoid *Salmonella* gastroenteritis ⁶
- Rare instances of toxic megacolon have been reported. ⁷
- Although intestinal intussusception may occur in some cases ⁸, a causal role for Rotavirus infection (ie, as opposed to Rotavirus vaccine ⁹) is not established. ¹⁰
- Central nervous system dysfunction may complicate Rotavirus infection, in the form of seizures ¹¹⁻¹⁴, cerebellitis ¹⁵⁻¹⁸, encephalopathy ^{19 20} and death. ²¹
- Some reports have linked Rotavirus infections with instances of aseptic meningitis ^{22 23}, necrotizing enterocolitis, myositis, liver abscess, pancreatitis ^{24 25}, pneumonia, Kawasaki's disease, acute hemorrhagic edema ²⁶, sudden infant death syndrome and Crohn's disease.

This disease is endemic or potentially endemic to all countries.

Rotavirus infection in Uganda

Prevalence surveys:

45.4% of children ages 3 to 59 months with diarrhea (2010 publication) ²⁷

References

1. Curr Opin Gastroenterol 2005 Jan ;21(1):26-31.
2. J Med Virol 2008 Dec ;80(12):2169-76.
3. Pediatr Infect Dis J 2010 Sep ;29(9):836-9.
4. Am J Epidemiol 2010 May 1;171(9):1023-30.
5. Lancet Infect Dis 2004 Feb ;4(2):91-9.
6. Eur J Clin Microbiol Infect Dis 2009 Apr ;28(4):425-8.
7. Acta Paediatr 2009 Nov ;98(11):1850-2.
8. Turk J Gastroenterol 2009 Sep ;20(3):209-13.
9. J Infect Dis 2009 Nov 1;200 Suppl 1:S264-70.
10. J Infect Dis 2009 Nov 1;200 Suppl 1:S277-81.
11. An Pediatr (Barc) 2008 Sep ;69(3):263-6.
12. Pediatr Neurol 2010 Jun ;42(6):404-8.
13. An Pediatr (Barc) 2010 Aug ;73(2):70-3.
14. Pediatr Emerg Care 2011 Nov ;27(11):1062-4.
15. AJNR Am J Neuroradiol 2010 Oct ;31(9):1591-5.
16. Brain Dev 2011 Jan ;33(1):21-7.
17. World J Pediatr 2011 Aug 27;
18. Pediatr Neurol 2012 Jan ;46(1):48-50.
19. Pediatr Infect Dis J 2009 Apr ;28(4):318-21.
20. Eur J Pediatr 2010 Oct ;169(10):1287-91.
21. J Child Neurol 2007 Dec ;22(12):1367-70.
22. Jpn J Infect Dis 2009 Jul ;62(4):279-83.
23. J Med Virol 2011 Sep ;83(9):1637-40.
24. Indian Pediatr 2009 Dec ;46(12):1099-101.
25. Pediatr Emerg Care 2010 Aug ;26(8):592-3.
26. Pediatr Dermatol 2004 Sep-Oct;21(5):548-50.
27. BMC Pediatr 2010 ;10:69.

Rubella

Agent	VIRUS - RNA. Togaviridae: Rubella virus
Reservoir	Human
Vector	None
Vehicle	Contact Air Transplacental
Incubation Period	16d - 18d (range 14d - 23d)
Diagnostic Tests	Viral culture (throat, urine). Serology. Nucleic acid amplification.
Typical Adult Therapy	Respiratory precautions. Supportive
Typical Pediatric Therapy	As for adult
Vaccines	Rubella Rubella - Mumps Measles-Mumps-Rubella Measles-Rubella
Clinical Hints	Maculopapular rash following a one-day prodrome of coryza and headache; post auricular lymphadenopathy; arthralgia and arthritis encountered in adults; severe thrombocytopenia or encephalitis may follow acute infection.
Synonyms	Epidemic roseola, German measles, Roda hund, Rode hond, Rode hunder, Rodehond, Rosolia, Roteln, Rubeola [Spanish], Three-day measles. ICD9: 056 ICD10: B06

Clinical

CDC (The United States Centers for Disease Control) case definition for surveillance:

For surveillance purposes, the CDC (The United States Centers for Disease Control) case definition of rubella requires, "An illness that has all of the following characteristics:

- acute onset of generalized maculopapular rash
- temperature >37.2 C if measured
- arthralgia/arthritis, lymphadenopathy, or conjunctivitis" ¹ Arthropathy may occur in as many as 41% of cases ²

A "confirmed" case requires either laboratory confirmation or epidemiological link to a laboratory-confirmed case.

- Atypical features may be seen in adults with rubella; ie, hepatitis, conjunctival hemorrhage ³, uveitis ⁴, retinitis ⁵ and a high incidence of polyarthritis.
- Rare instances of acute hepatic failure ⁶ and hemophagocytic syndrome ⁷ are reported.

Congenital rubella should be suspected if any of the following is present in a newborn infant ^{8 9}:

- cataracts (45% of cases), congenital glaucoma, pigmentary retinopathy
- congenital heart disease (70%, most commonly patent ductus arteriosus or pulmonary artery stenosis) Both anomalies may appear concurrently in up to 50% of cases ¹⁰
- hearing loss (35% to 60%)
- purpura
- splenomegaly
- jaundice
- microcephaly, mental retardation ¹¹, meningoencephalitis
- radiolucent bone disease
- duodenal stenosis ¹²

The chance of fetal defects from a viremic mother is 40% to 90% during the first trimester. ¹³


- Infection also increases the risk for spontaneous abortion and miscarriage by 50%. ¹⁴
- The rate of congenital rubella syndrome during epidemics is 0.5 to 2.2 per 1,000 live births.
- 60% of children with CRS have hearing impairment, 45% congenital heart disease, 27% microcephaly, 25% cataracts, 23% low birth weight (< 2,500 grams), 17% purpura, 19% hepatosplenomegaly, 13% mental retardation and 10% meningoencephalitis.

Anterior uveitis • differential diagnosis:

Anterior uveitis due to Rubella virus is characterized by younger age at onset and a chronic course, typically associated with cataract at presentation. ¹⁵

- Rubella virus has been implicated in the etiology of Fuchs heterochromic iridocyclitis. ¹⁶
- Anterior uveitis due to Herpes simplex and Varicella-Zoster viruses is more common in adults, and often follows an acute course.
- Herpes simplex anterior uveitis presents with conjunctival redness, corneal edema, a history of keratitis, and the presence of posterior synechiae. Anterior chamber inflammation is common with Herpes simplex virus, while vitritis is more common with Rubella and Varicella-Zoster virus.
- Rubella, Herpes simplex and Varicella-zoster viruses are associated with intraocular pressure of more than 30 mmHg and development of glaucoma (18%-30%; P = 0.686).
- Focal chorioretinal scars were present in 22% of Rubella cases, 0% of HSV and in 11% of VZV uveitis cases.

This disease is endemic or potentially endemic to all countries.

Rubella in Uganda

Graph: Uganda. Rubella, cases

References

1. Lancet 2004 Apr 3;363(9415):1127-37.
2. Clin Rheumatol 2009 Sep ;28(9):1067-71.
3. Med J Malaysia 2006 Jun ;61(2):242-4.
4. Am J Ophthalmol 2008 Aug ;146(2):292-7.
5. Clin Ophthalmol 2010 ;5:3-4.
6. Pediatr Infect Dis J 2010 Jun ;29(6):573-4.
7. Mediterr J Hematol Infect Dis 2012 ;4(1):e2012050.
8. Pediatr Infect Dis J 2004 Dec ;23(12):1116-22.
9. Reprod Toxicol 2006 May ;21(4):390-8.
10. Birth Defects Res A Clin Mol Teratol 2010 Jan ;88(1):1-8.
11. Semin Pediatr Neurol 1994 Sep ;1(1):26-35.
12. J Infect 2006 Nov ;53(5):e207-10.
13. Semin Fetal Neonatal Med 2007 Jun ;12(3):182-92.
14. N Engl J Med 1966 Apr 7;274(14):768-71.
15. Ophthalmology 2011 Oct ;118(10):1905-10.
16. Graefes Arch Clin Exp Ophthalmol 2010 Oct ;248(10):1487-91.

Salmonellosis

Agent	BACTERIUM. <i>Salmonella</i> A facultative gram-negative bacillus
Reservoir	Mammal Bird Reptile
Vector	None
Vehicle	Food Milk Eggs Poultry Shellfish Meat Vegetables Fruit Fecal-oral Fly
Incubation Period	12h - 36h (range 6h - 5d)
Diagnostic Tests	Culture (stool, blood, infected tissue). Serology.
Typical Adult Therapy	Stool precautions. Therapy not indicated for uncomplicated diarrhea; if necessary, treat per antibiogram
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, chills & watery diarrhea 12 to 24 hours after ingestion of eggs, meat, poultry; fecal leucocytes present; fever resolves in 2 days; but diarrhea persists for up to 7 days (occasionally weeks).
Synonyms	Salmonellosen, Salmonellosi. ICD9: 003 ICD10: A02

Clinical

WHO Case definition for surveillance:

- An illness with the following symptoms: diarrhea, abdominal cramps, fever, vomiting and malaise.

Laboratory criteria for confirmation

- Isolation of *Salmonella* spp. from the stool or blood of a patient.

Case classification

- Suspected: An individual showing one or more of the clinical features.
- Confirmed: A suspected case with laboratory confirmation.

Acute infection:

Salmonella gastroenteritis is usually indistinguishable from that caused by other bacterial and viral pathogens. ¹

- Nausea, vomiting, and diarrhea begin 6 to 48 hours following ingestion of contaminated food or water.
- Incubation periods as long as 8 days have been reported. ²
- Abdominal cramps and fever as high as 39 C are common.
- The diarrhea is usually characterized as loose, non-bloody stools of moderate volume.
- Voluminous diarrhea, bloody stools, and tenesmus may also occur.

The infection is usually self-limited.

- Fever resolves within 3 days, and diarrhea resolves within 3 to 7 days.
- Stool cultures may remain positive for 4 to 5 weeks after infection, and carriage may persist for as long as one year in fewer than 1% of cases. ³
- Antibiotic treatment is reserved for unusual and complicated infections: septicemia, neonates, immunosuppressed patients, etc.

Complications:

The spectrum of extraintestinal salmonellosis is similar to that of other gram-negative bacterial infections: osteomyelitis ⁴⁻⁸, meningitis ⁹⁻¹¹, endocarditis ¹²⁻¹⁴, etc.

- Endovascular infections are particularly common, and may result in aneurysms of the aorta and other large vessels. ^{15 16}
- *Salmonella* osteomyelitis is common in children with underlying hemoglobinopathies. Pyomyositis has also been reported in such cases. ¹⁷
- Septicemia is often described in patients with schistosomiasis ¹⁸⁻²², lymphoma, lupus erythematosus ^{23 24}, bartonellosis, malaria and hepatic cirrhosis.
- Rotavirus infection increases the risk of bacteremia in children with nontyphoid *Salmonella* gastroenteritis ²⁵
- Elderly patients are at risk for complicated or fatal infection. ²⁶
- Reactive arthritis has been reported in as many as 16.8% of cases ²⁷⁻²⁹
- The risk for reactive arthritis following *Salmonella* infection ³⁰ was 1.4/100,000 cases (United States, 2002 to 2004) ³¹
- There is evidence that salmonellosis may increase the risk for later development of inflammatory bowel disease. ³²

This disease is endemic or potentially endemic to all countries.

Salmonellosis in Uganda

Prevalence surveys:

39.4% of bacteremia episodes among severely malnourished children (Kampala, 2006 publication) ³³

5.7% of febrile HIV-positive adults (bacteremia, Kampala, 2007) ³⁴

References

1. Clin Infect Dis 2003 Aug 15;37(4):e47-52.
2. Foodborne Pathog Dis 2012 Mar ;9(3):245-8.
3. Rev Infect Dis 1984 May-Jun;6(3):345-56.
4. Spine (Phila Pa 1976) 2010 Nov 1;35(23):E1317-21.
5. Orthopedics 2009 Sep ;32(9)
6. Spine J 2009 Nov ;9(11):e1-4.
7. Acta Obstet Gynecol Scand 2009 ;88(10):1171-3.
8. Ann Saudi Med 2003 Nov-Dec;23(6):358-62.
9. Euro Surveill 2010 Feb 18;15(7)
10. Ann Trop Paediatr 2009 Mar ;29(1):13-22.
11. Int J Infect Dis 2003 Mar ;7(1):53-60.
12. Can J Cardiol 2010 Oct ;26(8):323-5.
13. J Heart Valve Dis 2009 Jul ;18(4):401-10.
14. J Am Soc Echocardiogr 2009 Feb ;22(2):210.e1-3.
15. Clin Infect Dis 1999 Oct ;29(4):862-8.
16. Hong Kong Med J 2007 Jun ;13(3):234-7.
17. J Med Case Rep 2010 ;4:198.
18. J Trop Med Hyg 1977 Jan ;80(1):14-8.
19. Mem Inst Oswaldo Cruz 1998 ;93 Suppl 1:135-9.
20. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):121.
21. Arch Intern Med 1971 Aug ;128(2):254-7.
22. Rev Soc Bras Med Trop 2009 Jul-Aug;42(4):436-45.
23. Int J Rheum Dis 2009 Dec ;12(4):319-23.
24. Lupus 2012 Oct ;21(12):1356-8.
25. Eur J Clin Microbiol Infect Dis 2009 Apr ;28(4):425-8.
26. J Infect 2008 Sep ;57(3):214-22.
27. J Rheumatol 2008 Mar ;35(3):480-7.
28. Rev Rhum Engl Ed 1999 Jan 30;66(1 Suppl):14S-18S; discussion 19S.
29. J Rheumatol 2002 Apr ;29(4):767-71.
30. Arthritis Rheum 1988 Nov ;31(11):1377-83.
31. Ann Rheum Dis 2008 Dec ;67(12):1689-96.
32. Gastroenterology 2009 Aug ;137(2):495-501.
33. BMC Infect Dis 2006 ;6:160.
34. J Acquir Immune Defic Syndr Hum Retrovirol 1998 Dec 15;19(5):484-9.

Sarcocystosis

Agent	PARASITE - Protozoa. Sporozoa, Coccidea, Eimeriida: <i>Sarcocystis bovis</i> or <i>S. suis</i>
Reservoir	Cattle Pig
Vector	None
Vehicle	Meat Water
Incubation Period	9d - 39d
Diagnostic Tests	Identification of cysts in stool.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Diarrhea and abdominal pain of varying severity; muscle pain and eosinophilia occasionally encountered.
Synonyms	<i>Isospora hominis</i> , Kudoa, Sarcocystiasis, Sarcocystis, Sarcosporidiosis. ICD9: 136.5 ICD10: A07.8

Clinical

Human infection follows ingestion of undercooked beef or pork.

- Clinical features are limited to abdominal pain, vomiting, moderate diarrhea or asymptomatic infection of muscle. ^{1 2}
- Rare instances of myositis ³ with eosinophilia have also been reported.

This disease is endemic or potentially endemic to all countries.

References

1. Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi 1999 ;17(1):25-7.
2. Clin Microbiol Rev 2004 Oct ;17(4):894-902, table of contents.
3. Arq Neuropsiquiatr 1985 Sep ;43(3):296-302.

Scabies

Agent	PARASITE - Arthropod. Arachnid, Acarina (Mite), Sarcoptidae: Sarcoptes [Acarus] scabiei
Reservoir	Human
Vector	mite
Vehicle	Contact, including Sexual contact
Incubation Period	3d - 42d
Diagnostic Tests	Identification of mites in skin scrapings.
Typical Adult Therapy	Permethrin 5%. OR Lindane. OR Crothamiton 10% OR Ivermectin 150 to 200 ug/kg PO as single dose
Typical Pediatric Therapy	Permethrin 5%. OR Lindane. OR Crothamiton 10% OR Ivermectin 200 mcg/kg PO (> 15 kg body weight)
Clinical Hints	Intensely pruritic papules, vesicles and burrows - interdigital webs, wrists, elbows, axillae, perineal region, buttocks, penis; pruritus most intense at night; severe psoriaform infestation (Norwegian scabies) noted in debilitated patients.
Synonyms	Cheyletiella, Cheyletiella infestation, Escabiose, Escabiosis, Histiostomatid mites, Krätze, Mange, Ornithonyssus, Pyemotes, Sarcoptes scabiei, Sarna, Scabbia, Skabies, Tropical rat mite. ICD9: 133 ICD10: B86

Clinical

The lesions of scabies are usually symmetrical.

- Typical sites include the interdigital webs, buttocks, penis, scrotum, breasts and nipples, axillae and flexor surfaces of the wrists. ¹
- Pruritus is often worse at night.
- Skin lesions consist of burrows, papules or vesicles. ²
- Exaggerated eczematous patches ('crusted', or Norwegian scabies) ^{3 4} may be encountered • notably in institutions for Down's syndrome and leprosy. ⁵
- Lesions in children are atypical and tend to involve the buttocks and perineum. ⁶
- Complications include secondary infection and acute glomerulonephritis.

Otoacariasis due to Histiostomatid mites has been reported in Saudi Arabia. ⁷

This disease is endemic or potentially endemic to all countries.

References

1. J Am Acad Dermatol 2004 Jun ;50(6):819-42, quiz 842-4.
2. Cutis 1995 Jun ;55(6):370-1.
3. Lancet Infect Dis 2006 Dec ;6(12):769-79.
4. Arch Dermatol 1976 Feb ;112(2):179-81.
5. Adv Parasitol 2004 ;57:309-76.
6. Cutis 2003 Mar ;71(3):193-6.
7. Am J Trop Med Hyg 2007 May ;76(5):967-71.

Scarlet fever

Agent	BACTERIUM. Streptococcus pyogenes A facultative gram-positive coccus
Reservoir	Human
Vector	None
Vehicle	Infected secretions Occasionally food
Incubation Period	1d - 4d
Diagnostic Tests	Typical clinical features associated with group A streptococcal pharyngitis.
Typical Adult Therapy	Benzathine Penicillin G 1.2 million units IM as single dose
Typical Pediatric Therapy	Benzathine Penicillin G : Weight <14kg: 300,000 units IM Weight 14 to 28kg: 600,000 units IM Weight >28kg: 1.2 million units IM
Clinical Hints	Overt pharyngitis followed within 24 to 48 hrs by florid erythematous rash.
Synonyms	Escarlatina, Lanhousha, Scarlattina, Scharlach. ICD9: 034.1 ICD10: A38

Clinical

Signs of streptococcal pharyngitis (fever, pharyngeal exudate and pain) are followed by the appearance of a rash within 12 to 24 hours.

- The exanthem appears initially on the trunk and spreads rapidly over the body to finally involve the extremities. ¹
- The exanthem has the texture of sandpaper, and blanches with pressure.
- Pruritis may be present.
- Facial flushing and circumoral pallor are characteristic.

The patient appears ill, with fever, tachycardia, pharyngitis, tender adenopathy and palatal petechiae.

- Within a few days, the rash becomes more intense along skin folds, producing lines of confluent petechiae (Pastia sign).
- The rash begins to fade within 3 to 4 days, with desquamation evident over the face, palms and fingers.
- Skin peeling may persist for as long as a month.

During the first 2 days of illness, the tongue has a white coat through which the red and edematous papillae project ('white strawberry tongue').

- The tongue later desquamates and becomes markedly reddened ('red strawberry tongue').

Complications are those associated with the streptococcal infection itself • spread to regional, retropharyngeal tissues, middle ears, and sinuses; acute rheumatic fever or post-streptococcal glomerulonephritis.

- Septic complications such as meningitis, pyogenic arthritis, and endocarditis, are occasionally encountered.

This disease is endemic or potentially endemic to all countries.

References

1. [J Am Acad Dermatol 1989 Nov ;21\(5 Pt 1\):891-903.](#)

Schistosomiasis - haematobium

Agent	PARASITE - Platyhelminthes, Trematoda. Strigeida, Schistosomatidae: <i>Schistosoma haematobium</i>
Reservoir	Snail (<i>Bulinus</i> , <i>Planorbarius</i> , <i>Ferrissia</i>) Rarely baboon or monkey
Vector	None
Vehicle	Water (skin contact)
Incubation Period	2w - 6w
Diagnostic Tests	Identification of ova in urine or stool. Serology. Antigen detection.
Typical Adult Therapy	Praziquantel 20 mg/kg PO BID X 1 day
Typical Pediatric Therapy	As for adult
Clinical Hints	Early urticaria, fever and eosinophilia; later, dysuria, hematuria and obstructive nephropathy; often complicated by bladder cancer in advanced cases; parasite may survive for decades in human host.
Synonyms	Bilharziasis, urinary, Egyptian hematuria, Katayama fever [1], <i>Schistosoma guineensis</i> , <i>Schistosoma haematobium</i> , Schistosomal hematuria, Schistosomiasis, Vesicle bilharziasis. ICD9: 120.0 ICD10: B65.0

Clinical

WHO Case definition for surveillance:

Endemic areas (moderate or high prevalence)

- Suspected: Not applicable.
- Probable: Not applicable.
- Confirmed: A person with visible hematuria or with positive reagent strip for hematuria or with eggs of *S. haematobium* in urine (microscope).

Non-endemic areas and areas of low prevalence

- Suspected: A person with visible hematuria or with positive reagent strip for hematuria.
- Probable: Not applicable.
- Confirmed: A person with eggs of *S. haematobium* in urine (microscope).

The clinical features caused by *Schistosoma* species infecting man are similar ¹, and will be discussed together.

Acute infection:

Within 24 hours of penetration by cercariae, the patient develops a pruritic papular skin rash known as swimmer's itch. [The more overt form of Cercarial dermatitis associated with avian schistosomes is discussed elsewhere in this module.]

- One to two months after exposure, an overt systemic illness known as Katayama fever (named for Katayama district, Hiroshima, Japan) begins, heralded by acute onset of fever, chills, diaphoresis, headache, and cough. ²
- The liver, spleen, and lymph nodes are enlarged, and eosinophilia is present.
- Rare instances of myocarditis have been reported during acute schistosomiasis. ^{3 4}
- Although deaths have been described at this point (notably in *S. japonicum* infection) these findings subside within a few weeks in most cases.

Chronic schistosomiasis:

The likelihood of progression to chronic schistosomiasis is related to the extent of infestation.

- Chronic schistosomiasis caused by *S. mansoni*, *S. japonicum*, or *S. mekongi* is characterized by fatigue, abdominal pain and intermittent diarrhea or dysentery.
- Blood loss from intestinal ulcerations may lead to moderate anemia.
- In *S. mansoni*, *S. japonicum*, and *S. mekongi* infections, ova remain in the venous portal circulation and are carried to the liver where they produce granulomata and fibrosis ⁵, and block portal blood flow.
- Colonic polyposis has been associated with infection by *S. mansoni*, *S. japonicum*, and *S. intercalatum*.
- Retroperitoneal fibrosis has been reported with *S. japonicum* infection. ⁷
- Portal hypertension and portosystemic collateral circulation result.
- Although liver function tests remain normal for a long time, hepatosplenomegaly and variceal hemorrhage develop.
- The spleen is firm and may reach massive size.
- Fatal hematemesis is unusual.
- Laboratory tests reveal moderate eosinophilia and anemia related to blood loss and hypersplenism.
- Eventually, hepatic function deteriorates, with late ascites and jaundice.

In *S. haematobium* infection, ova are located in the bladder and ureters, leading to granuloma formation, inflammation, hematuria, ureteral obstruction, secondary infection and often carcinoma of the bladder.⁸⁻¹⁰ Ova are also commonly present in the seminal vesicles and prostate.¹¹

- Areas of chronic inflammation, fibrous tissue and calcifications ("sandy patches") in the genital mucosa and bladder contain ova, and are considered pathognomonic for *S. haematobium* infection.¹²
- Genital lesions may present a risk factor for acquisition of HIV infection¹³; and schistosomal co-infection may accelerate HIV disease progression and facilitate viral transmission to sexual partners.¹⁴
- Terminal hematuria and dysuria are common symptoms.
- Although best known for damage to the urinary bladder and ureters, the female genitalia are involved in 50% to 70% of women with *S. haematobium* infection • resulting in vaginal deformities and fistulae¹⁵, hypogonadism, ectopic pregnancy, miscarriage and malignancy.¹⁶⁻¹⁹ *Schistosoma mansoni* is implicated in the etiology of appendicitis²⁰, and membranoproliferative glomerulonephritis and amyloidosis²¹; and may also involve the fallopian tubes²²⁻²³ or cause ovarian²⁴ or testicular pseudotumor²⁵ and acute abdomen associated with granulomatous peritonitis²⁶ or panniculitis.²⁷
- Reinfection or inadequately treated infection may lead to extra-anogenital bilharziasis cutanea tarda. Lesion may typically complicate pre-existing skin conditions.²⁸
- Proctitis is occasionally encountered.²⁹

S. intercalatum infection is characterized by abdominal pain and bloody diarrhea.

S. mekongi is an important cause of hepatomegaly in endemic areas.

Complications:

The following are some of the many complications described in chronic schistosomiasis.

- Pulmonary schistosomiasis is manifested by symptoms and signs of right ventricular congestion related to blockage of pulmonary capillaries by ova in the course of hepatosplenic schistosomiasis.³⁰⁻³²
- Central nervous system schistosomiasis is manifested as delirium, coma, seizures, dysphasia, visual impairment, ataxia, a cerebral mass, generalized encephalopathy, cerebral vasculitis with stroke, or focal epilepsy (notably in *S. japonicum* infection).³³⁻⁴⁵
- Granulomata of *S. haematobium* and *S. mansoni* may involve the spinal cord (most commonly the cauda equina or conus medularis), producing transverse myelitis.⁴⁶⁻⁵⁹ Rare instances of cerebral infection by *S. haematobium* have been reported.⁶⁰ *Schistosoma mansoni* infection may occasionally involve the bladder, mimicking *S. haematobium* infection or malignancy.⁶¹ *S. mansoni* infection has been implicated in cases of colo-rectal cancer.⁶²
- *Salmonella* bacteremia is often reported among persons with hepato-splenic schistosomiasis.⁶³⁻⁶⁷

This disease is endemic or potentially endemic to 58 countries.

Schistosomiasis - haematobium in Uganda

The disease is endemic to the north, East (Mbale area), northern shore of Lake Victoria and area north of Lake Kyoga. - *Schistosoma mansoni* is more widely distributed than *S. haematobium* in this country.⁶⁸

17% of persons seroconverted toward *Schistosoma mansoni* and / or *Schistosoma haematobium* following recreational exposure to the Upper Nile River (2007)⁶⁹

Prevalence surveys:

40.8% of school-age children in the Lake Victoria region (*Schistosoma* sp, 2011 publication)⁷⁰

The local snail reservoir is *Bulinus tropicus*.⁷¹⁻⁷²

References

1. Lancet 2006 Sep 23;368(9541):1106-18.
2. Lancet Infect Dis 2007 Mar ;7(3):218-24.
3. Am J Trop Med Hyg 2010 Mar ;82(3):365-7.
4. Med Trop (Mars) 1980 May-Jun;40(3):271-9.
5. Parasite Immunol 2009 Nov ;31(11):656-63.
6. Trans R Soc Trop Med Hyg 2010 Jun ;104(6):443-5.
7. Nihon Hinyokika Gakkai Zasshi 2010 Jul ;101(5):694-7.
8. Parasite Immunol 2009 Nov ;31(11):686-96.
9. Cancer Lett 2011 Jun 28;305(2):239-49.
10. J Infect Dev Ctries 2010 May ;4(5):267-81.
11. Am J Trop Med Hyg 1970 Sep ;19(5):779-84.
12. Trans R Soc Trop Med Hyg 2006 Aug ;100(8):740-52.
13. AIDS 2006 Feb 28;20(4):593-600.
14. PLoS Negl Trop Dis 2011 Dec ;5(12):e1396.
15. Afr J Reprod Health 2009 Sep ;13(3):137-40.
16. Acta Trop 2001 Jun 22;79(3):193-210.
17. Parasitol Today 1999 Sep ;15(9):378-81.
18. Am J Trop Med Hyg 2009 Oct ;81(4):549-50.
19. Trends Parasitol 2012 Feb ;28(2):58-65.
20. Trop Gastroenterol 2009 Oct-Dec;30(4):230-2.
21. Ann Pathol 2012 Feb ;32(1):40-52.
22. Braz J Infect Dis 2010 May-Jun;14(3):288-90.
23. Braz J Infect Dis 2011 Mar-Apr;15(2):174-7.
24. Arch Gynecol Obstet 2010 Jan ;281(1):141-3.
25. Fertil Steril 2011 May ;95(6):2124.e1-4.
26. Trans R Soc Trop Med Hyg 2009 Oct ;103(10):1068-70.
27. Ann Dermatol Venereol 2012 Feb ;139(2):132-6.
28. J Cutan Pathol 2009 Jul ;36(7):766-71.

29. Am J Trop Med Hyg 2009 Feb ;80(2):179-81.
30. Ann Trop Med Parasitol 2009 Mar ;103(2):129-43.
31. Chest 2010 Jun ;137(6 Suppl):20S-29S.
32. Travel Med Infect Dis 2012 Sep 11;
33. J Neurol 2012 Jan ;259(1):22-32.
34. Pediatr Infect Dis J 2011 Nov ;30(11):1006-8.
35. Ann Indian Acad Neurol 2011 Apr ;14(2):107-10.
36. Lancet Neurol 2011 Sep ;10(9):853-64.
37. BMC Infect Dis 2012 ;12:220.
38. Neurologist 2012 Nov ;18(6):333-42.
39. Neurochirurgie 2012 May 29;
40. Rev Med Interne 2012 Oct ;33(10):580-2.
41. Curr Neurol Neurosci Rep 2012 Dec ;12(6):666-74.
42. Trans R Soc Trop Med Hyg 2008 Feb ;102(2):107-16.
43. Lancet Neurol 2011 Sep ;10(9):853-64.
44. BMC Infect Dis 2012 ;12:220.
45. Neurologist 2012 Nov ;18(6):333-42.
46. Q J Med 1986 Dec ;61(236):1131-9.
47. Acta Trop 2008 Nov-Dec;108(2-3):89-97.
48. J Infect 2010 Mar ;60(3):244-7.
49. Neurol Res 2010 Apr ;32(3):252-62.
50. Am J Trop Med Hyg 2009 Oct ;81(4):551-4.
51. J Neurol 2012 Jan ;259(1):22-32.
52. Pediatr Infect Dis J 2011 Nov ;30(11):1006-8.
53. Ann Indian Acad Neurol 2011 Apr ;14(2):107-10.
54. Lancet Neurol 2011 Sep ;10(9):853-64.
55. BMC Infect Dis 2012 ;12:220.
56. Neurologist 2012 Nov ;18(6):333-42.
57. Neurochirurgie 2012 May 29;
58. Rev Med Interne 2012 Oct ;33(10):580-2.
59. Curr Neurol Neurosci Rep 2012 Dec ;12(6):666-74.
60. J Clin Microbiol 2011 Oct ;49(10):3703-6.
61. Rev Soc Bras Med Trop 2009 Sep-Oct;42(5):581-2.
62. World J Surg Oncol 2010 ;8:68.
63. J Trop Med Hyg 1977 Jan ;80(1):14-8.
64. Mem Inst Oswaldo Cruz 1998 ;93 Suppl 1:135-9.
65. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):121.
66. Arch Intern Med 1971 Aug ;128(2):254-7.
67. Rev Soc Bras Med Trop 2009 Jul-Aug;42(4):436-45.
68. Afr Health Sci 2008 Dec ;8(4):239-43.
69. Emerg Infect Dis 2010 May ;16(5):866-8.
70. Parasit Vectors 2011 ;4:232.
71. East Afr Med J 1996 Aug ;73(8):495-8.
72. East Afr Med J 1992 Jun ;69(6):316-8.

Schistosomiasis - intercalatum

Agent	PARASITE - Platyhelminthes, Trematoda. Strigeida, Schistosomatidae: <i>Schistosoma intercalatum</i>
Reservoir	Snail (<i>Bulinus forskalii</i> and <i>Bulinus africanus</i> group)
Vector	None
Vehicle	Water (skin contact)
Incubation Period	2w - 6w
Diagnostic Tests	Identification of ova in stool or biopsy specimens. Serology. Antigen detection.
Typical Adult Therapy	Praziquantel 20 mg/kg PO BID X 1 day
Typical Pediatric Therapy	As for adult
Clinical Hints	Diarrhea (often bloody), abdominal pain, eosinophilia and hepatosplenomegaly; significant disease of liver, lung or central nervous system is rare.
Synonyms	<i>Schistosoma intercalatum</i> . ICD9: 120.8 ICD10: B65.8

Clinical

WHO Case definition for surveillance (all forms of intestinal schistosomiasis):

Endemic areas (moderate or high prevalence)

- Suspected: A person with chronic or recurrent intestinal symptoms (blood in stool, bloody diarrhea, diarrhea, abdominal pains) or, at a later stage, hepatosplenomegaly.
- Probable: A person who meets the criteria for presumptive treatment, according to the locally applicable diagnostic algorithms.
- Confirmed: A person with eggs of *S. mansoni*, or *S. japonicum/mekongi* in stools (microscope).

Non-endemic areas and areas of low prevalence

- Suspected: A person with chronic or recurrent intestinal symptoms (blood in stool, bloody diarrhea, diarrhea, abdominal pains) or, at a later stage, hepatosplenomegaly.
- Probable: Not applicable.
- Confirmed: A person with eggs of *S. mansoni* or *S. japonicum* in stools (microscope). A person with positive reaction to immunoblot test.

The clinical features caused by *Schistosoma* species infecting man are similar ¹, will be discussed together.

Acute infection:

Within 24 hours of penetration by cercariae, the patient develops a pruritic papular skin rash known as swimmer's itch. [The more overt form of Cercarial dermatitis associated with avian schistosomes is discussed elsewhere in this module.]

- One to two months after exposure, an overt systemic illness known as Katayama fever (named for Katayama district, Hiroshima, Japan) begins, heralded by acute onset of fever, chills, diaphoresis, headache, and cough. ²
- The liver, spleen, and lymph nodes are enlarged, and eosinophilia is present.
- Although deaths have been described at this point (notably in *S. japonicum* infection) these findings subside within a few weeks in most cases.

Chronic schistosomiasis:

The likelihood of progression to chronic schistosomiasis is related to the extent of infestation.

- Chronic schistosomiasis caused by *S. mansoni*, *S. japonicum*, or *S. mekongi* is characterized by fatigue, abdominal pain and intermittent diarrhea or dysentery.
- Colonic polyposis has been associated with infection by *S. mansoni*, *S. japonicum*, and *S. intercalatum*.
- ³ Retroperitoneal fibrosis has been reported with *S. japonicum* infection. ⁴
- Blood loss from intestinal ulcerations may lead to moderate anemia.
- In *S. mansoni*, *S. japonicum*, and *S. mekongi* infections, ova remain in the venous portal circulation and are carried to the liver where they produce granulomata and fibrosis ⁵, and block portal blood flow.
- Portal hypertension and portosystemic collateral circulation result.
- Although liver function tests remain normal for a long time, hepatosplenomegaly and variceal hemorrhage develop.
- The spleen is firm and may reach massive size.
- Fatal hematemesis is unusual.
- Laboratory tests reveal moderate eosinophilia and anemia related to blood loss and hypersplenism.

- Eventually, hepatic function deteriorates, with late ascites and jaundice.

In *S. haematobium* infection, ova are located in the bladder and ureters, leading to granuloma formation, inflammation, hematuria, ureteral obstruction, secondary infection and often carcinoma of the bladder.⁶⁻⁸ Ova are also commonly present in the seminal vesicles and prostate.⁹

- Genital lesions may present a risk factor for acquisition of HIV infection¹⁰; and schistosomal co-infection may accelerate HIV disease progression and facilitate viral transmission to sexual partners.¹¹
- Terminal hematuria and dysuria are common symptoms.

S. intercalatum infection is characterized by abdominal pain and bloody diarrhea.

S. mekongi is an important cause of hepatomegaly in endemic areas.

Complications:

The following are some of the many complications described in chronic schistosomiasis.

- Pulmonary schistosomiasis is manifested by symptoms and signs of right ventricular congestion related to blockage of pulmonary capillaries by ova in the course of hepatosplenic schistosomiasis.¹²⁻¹⁴
- Central nervous system schistosomiasis is manifested as delirium, coma, seizures, dysphasia, visual impairment, ataxia, a cerebral mass, generalized encephalopathy, cerebral vasculitis with stroke, or focal epilepsy (notably in *S. japonicum* infection).¹⁵⁻²⁷
- Granulomata of *S. haematobium* and *S. mansoni* may involve the spinal cord (most commonly the cauda equina or conus medularis), producing transverse myelitis.²⁸⁻⁴¹ Rare instances of cerebral infection by *S. haematobium* have been reported.⁴² *Schistosoma mansoni* infection may occasionally involve the bladder, mimicking *S. haematobium* infection or malignancy.⁴³ *S. mansoni* infection has been implicated in cases of colo-rectal cancer.⁴⁴
- Although best known for damage to the urinary bladder and ureters, the female genitalia are involved in 50% to 70% of women with *S. haematobium* infection • resulting in vaginal deformities and fistulae⁴⁵, hypogonadism, ectopic pregnancy, miscarriage and malignancy.⁴⁶⁻⁴⁹ *Schistosoma mansoni* is implicated in the etiology of appendicitis⁵⁰, and membranoproliferative glomerulonephritis and amyloidosis⁵¹; and may also involve the fallopian tubes^{52 53} or cause ovarian⁵⁴ or testicular pseudotumor⁵⁵ and acute abdomen associated with granulomatous peritonitis⁵⁶ or panniculitis.⁵⁷
- *Salmonella* bacteremia is often reported among persons with hepato-splenic schistosomiasis.⁵⁸⁻⁶²

This disease is endemic or potentially endemic to 15 countries.

Schistosomiasis - intercalatum in Uganda

The disease is identified in Arua, Gulu, Maracha and the Albert Nile region.⁶³

References

1. Lancet 2006 Sep 23;368(9541):1106-18.
2. Lancet Infect Dis 2007 Mar ;7(3):218-24.
3. Trans R Soc Trop Med Hyg 2010 Jun ;104(6):443-5.
4. Nihon Hinyokika Gakkai Zasshi 2010 Jul ;101(5):694-7.
5. Parasite Immunol 2009 Nov ;31(11):656-63.
6. Parasite Immunol 2009 Nov ;31(11):686-96.
7. Cancer Lett 2011 Jun 28;305(2):239-49.
8. J Infect Dev Ctries 2010 May ;4(5):267-81.
9. Am J Trop Med Hyg 1970 Sep ;19(5):779-84.
10. AIDS 2006 Feb 28;20(4):593-600.
11. PLoS Negl Trop Dis 2011 Dec ;5(12):e1396.
12. Ann Trop Med Parasitol 2009 Mar ;103(2):129-43.
13. Chest 2010 Jun ;137(6 Suppl):20S-29S.
14. Travel Med Infect Dis 2012 Sep 11;
15. J Neurol 2012 Jan ;259(1):22-32.
16. Pediatr Infect Dis J 2011 Nov ;30(11):1006-8.
17. Ann Indian Acad Neurol 2011 Apr ;14(2):107-10.
18. Lancet Neurol 2011 Sep ;10(9):853-64.
19. BMC Infect Dis 2012 ;12:220.
20. Neurologist 2012 Nov ;18(6):333-42.
21. Neurochirurgie 2012 May 29;
22. Rev Med Interne 2012 Oct ;33(10):580-2.
23. Curr Neurol Neurosci Rep 2012 Dec ;12(6):666-74.
24. Trans R Soc Trop Med Hyg 2008 Feb ;102(2):107-16.
25. Lancet Neurol 2011 Sep ;10(9):853-64.
26. BMC Infect Dis 2012 ;12:220.
27. Neurologist 2012 Nov ;18(6):333-42.
28. Q J Med 1986 Dec ;61(236):1131-9.
29. Acta Trop 2008 Nov-Dec;108(2-3):89-97.
30. J Infect 2010 Mar ;60(3):244-7.
31. Neurol Res 2010 Apr ;32(3):252-62.
32. Am J Trop Med Hyg 2009 Oct ;81(4):551-4.
33. J Neurol 2012 Jan ;259(1):22-32.
34. Pediatr Infect Dis J 2011 Nov ;30(11):1006-8.
35. Ann Indian Acad Neurol 2011 Apr ;14(2):107-10.
36. Lancet Neurol 2011 Sep ;10(9):853-64.
37. BMC Infect Dis 2012 ;12:220.
38. Neurologist 2012 Nov ;18(6):333-42.
39. Neurochirurgie 2012 May 29;
40. Rev Med Interne 2012 Oct ;33(10):580-2.
41. Curr Neurol Neurosci Rep 2012 Dec ;12(6):666-74.
42. J Clin Microbiol 2011 Oct ;49(10):3703-6.
43. Rev Soc Bras Med Trop 2009 Sep-Oct;42(5):581-2.
44. World J Surg Oncol 2010 ;8:68.
45. Afr J Reprod Health 2009 Sep ;13(3):137-40.
46. Acta Trop 2001 Jun 22;79(3):193-210.
47. Parasitol Today 1999 Sep ;15(9):378-81.
48. Am J Trop Med Hyg 2009 Oct ;81(4):549-50.
49. Trends Parasitol 2012 Feb ;28(2):58-65.
50. Trop Gastroenterol 2009 Oct-Dec;30(4):230-2.
51. Ann Pathol 2012 Feb ;32(1):40-52.
52. Braz J Infect Dis 2010 May-Jun;14(3):288-90.
53. Braz J Infect Dis 2011 Mar-Apr;15(2):174-7.
54. Arch Gynecol Obstet 2010 Jan ;281(1):141-3.
55. Fertil Steril 2011 May ;95(6):2124.e1-4.
56. Trans R Soc Trop Med Hyg 2009 Oct ;103(10):1068-70.
57. Ann Dermatol Venereol 2012 Feb ;139(2):132-6.
58. J Trop Med Hyg 1977 Jan ;80(1):14-8.
59. Mem Inst Oswaldo Cruz 1998 ;93 Suppl 1:135-9.
60. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):121.
61. Arch Intern Med 1971 Aug ;128(2):254-7.
62. Rev Soc Bras Med Trop 2009 Jul-Aug;42(4):436-45.
63. Am J Trop Med Hyg 1994 Jun ;50(6):723-6.

Schistosomiasis - mansoni

Agent	PARASITE - Platyhelminthes, Trematoda. Strigeida, Schistosomatidae: Schistosoma mansoni
Reservoir	Snail (Biomphalaria) Dog Cat Pig Cattle Rodent Horse Non-human primate
Vector	None
Vehicle	Water (skin contact)
Incubation Period	2w - 6w
Diagnostic Tests	Identification of ova in stool or biopsy specimens. Serology. Antigen detection.
Typical Adult Therapy	Praziquantel 20 mg/kg PO BID X one day OR Oxamniquine 15 mg PO X one dose
Typical Pediatric Therapy	Praziquantel 20 mg/kg PO BID X one day OR Oxamniquine 10 mg PO BID X one day
Clinical Hints	Early urticaria, fever and eosinophilia; later, hepatosplenomegaly and portal hypertension; parasite may survive for decades in human host.
Synonyms	Bilharziasis, intestinal, Katayama fever [3], Schistosoma mansoni. ICD9: 120.1 ICD10: B65.1

Clinical

WHO Case definition for surveillance (all forms of intestinal schistosomiasis):

Endemic areas (moderate or high prevalence)

- Suspected: A person with chronic or recurrent intestinal symptoms (blood in stool, bloody diarrhea, diarrhea, abdominal pains) or, at a later stage, hepatosplenomegaly.
- Probable: A person who meets the criteria for presumptive treatment, according to the locally applicable diagnostic algorithms.
- Confirmed: A person with eggs of *S. mansoni*, or *S. japonicum/mekongi* in stools (microscope).

Non-endemic areas and areas of low prevalence

- Suspected: A person with chronic or recurrent intestinal symptoms (blood in stool, bloody diarrhea, diarrhea, abdominal pains) or, at a later stage, hepatosplenomegaly.
- Probable: Not applicable.
- Confirmed: A person with eggs of *S. mansoni* or *S. japonicum* in stools (microscope). A person with positive reaction to immunoblot test.

The clinical features caused by *Schistosoma* species infecting man are similar ¹, will be discussed together.

Acute infection:

Within 24 hours of penetration by cercariae, the patient develops a pruritic papular skin rash known as swimmer's itch. [The more overt form of Cercarial dermatitis associated with avian schistosomes is discussed elsewhere in this module.]

- One to two months after exposure, an overt systemic illness known as Katayama fever (named for Katayama district, Hiroshima, Japan) begins, heralded by acute onset of fever, chills, diaphoresis, headache, and cough. ²
- The liver, spleen, and lymph nodes are enlarged, and eosinophilia is present.
- Although deaths have been described at this point (notably in *S. japonicum* infection) these findings subside within a few weeks in most cases.

Chronic schistosomiasis:

The likelihood of progression to chronic schistosomiasis is related to the extent of infestation.

- Chronic schistosomiasis caused by *S. mansoni*, *S. japonicum*, or *S. mekongi* is characterized by fatigue, abdominal pain and intermittent diarrhea or dysentery.
- Colonic polyposis is has been associated with infection by *S. mansoni*, *S. japonicum*, and *S. intercalatum*.
- ³ Retroperitoneal fibrosis has been reported with *S. japonicum* infection. ⁴
- Blood loss from intestinal ulcerations may lead to moderate anemia.
- In *S. mansoni*, *S. japonicum*, and *S. mekongi* infections, ova remain in the venous portal circulation and are carried to the liver where they produce granulomata and fibrosis ⁵, and block portal blood flow.
- Portal hypertension and portosystemic collateral circulation result.
- Although liver function tests remain normal for a long time, hepatosplenomegaly and variceal hemorrhage develop.
- The spleen is firm and may reach massive size.
- Fatal hematemesis is unusual.
- Laboratory tests reveal moderate eosinophilia and anemia related to blood loss and hypersplenism.

- Eventually, hepatic function deteriorates, with late ascites and jaundice.

In *S. haematobium* infection, ova are located in the bladder and ureters, leading to granuloma formation, inflammation, hematuria, ureteral obstruction, secondary infection and often carcinoma of the bladder.⁶⁻⁸ Ova are also commonly present in the seminal vesicles and prostate.⁹

- Genital lesions may present a risk factor for acquisition of HIV infection¹⁰; and schistosomal co-infection may accelerate HIV disease progression and facilitate viral transmission to sexual partners.¹¹
- Terminal hematuria and dysuria are common symptoms.

S. intercalatum infection is characterized by abdominal pain and bloody diarrhea.

S. mekongi is an important cause of hepatomegaly in endemic areas.

Complications:

The following are some of the many complications described in chronic schistosomiasis.

- Pulmonary schistosomiasis is manifested by symptoms and signs of right ventricular congestion related to blockage of pulmonary capillaries by ova in the course of hepatosplenic schistosomiasis.¹²⁻¹⁴
- Central nervous system schistosomiasis is manifested as delirium, coma, seizures, dysphasia, visual impairment, ataxia, a cerebral mass, generalized encephalopathy, cerebral vasculitis with stroke, or focal epilepsy (notably in *S. japonicum* infection).¹⁵⁻²⁷
- Granulomata of *S. haematobium* and *S. mansoni* may involve the spinal cord (most commonly the cauda equina or conus medularis), producing transverse myelitis.²⁸⁻⁴¹ Rare instances of cerebral infection by *S. haematobium* have been reported.⁴² *Schistosoma mansoni* infection may occasionally involve the bladder, mimicking *S. haematobium* infection or malignancy.⁴³ *S. mansoni* infection has been implicated in cases of colo-rectal cancer.⁴⁴
- Although best known for damage to the urinary bladder and ureters, the female genitalia are involved in 50% to 70% of women with *S. haematobium* infection • resulting in vaginal deformities and fistulae⁴⁵, hypogonadism, ectopic pregnancy, miscarriage and malignancy.⁴⁶⁻⁴⁹ *Schistosoma mansoni* is implicated in the etiology of appendicitis⁵⁰, and membranoproliferative glomerulonephritis and amyloidosis⁵¹; and may also involve the fallopian tubes^{52 53} or cause ovarian⁵⁴ or testicular pseudotumor⁵⁵ and acute abdomen associated with granulomatous peritonitis⁵⁶ or panniculitis.⁵⁷
- *Salmonella* bacteremia is often reported among persons with hepato-splenic schistosomiasis.⁵⁸⁻⁶²

This disease is endemic or potentially endemic to 59 countries.

Schistosomiasis - mansoni in Uganda

Time and Place:

Schistosoma mansoni infestation was first reported in Uganda in 1902, in Arua district.

- The disease is endemic to the north, East (Mbale area), northern shore of Lake Victoria and area north of Lake Kyoga.^{63 64}

- *Schistosoma mansoni* is more widely distributed than *S. haematobium* in this country.⁶⁵

- Transmission is limited to areas with rainfall >900 mm per year.

- Disease prevalence rates and intensity increase with age, peaking in the age group 10 to 20 years (1998 to 2002).

Prevalence surveys:

- 4.1% of children ages 10 to 15 (Kampala, 1997 publication)⁶⁶
- 92.6% of schools in proximity to Lake Victoria (at least one infected child, 2009 publication)⁶⁷
- Over 50% of school children living in proximity to Lake Victoria (1998 to 2002)
- 20.4% of school children nationwide, with a male/female ratio of 1.22/1 (1998 to 2002)
- 42.4% of school children in 2003; 26.8% in 2004; 17.9% in 2005 (nationwide)
- 44.3% of school children in northwest and southeast Uganda (2002 to 2003)⁶⁸
- 27.8% of school children near high altitude (>1,400 meters) crater lakes in western Uganda (2008 publication)⁶⁹
- 18.1% of school children in the Great Lakes region of Uganda, Tanzania, Kenya and Burundi (2010 publication)⁷⁰
- 40.8% of school-age children in the Lake Victoria region (*Schistosoma* sp, 2011 publication)⁷¹
- 44.3% of preschool children and 60.0% of their mothers near Lake Albert, 16.0% of preschool children and 29.2% of their mothers near Lake Victoria (2010 publication)
- 47.5% of preschool children in Bugoigo, Lake Albert region (2011 publication)⁷²
- 27.2% of children and 47.6% of their mothers (2010 publication)⁷³
- 47% of children below age 3 years living on the Ugandan shore of Lake Victoria (2006 publication)⁷⁴
- 72% of the population of Butiaba, Lake Albert (1996 to 1997)⁷⁵
- 65% of persons ages 10 to 20 years in Walukuba Division, Jinja District (2009 publication)^{76 77}
- 86% in Buboigo and Walakuba, 92% in Booma and 91% in Piida (Lake Albert region, 2006 publication)⁷⁸

- 18.3% of otherwise healthy pregnant women at enrolment to a trial of deworming (Entebbe, 2007 publication) ⁷⁹
- 18% of pregnant women (Entebbe and Katabi, 2003 to 2005) ⁸⁰
- 18% of pregnant women (Entebbe, 2010 publication) ⁸¹
- 3.7% of pregnant women in Masindi District (2003 to 2004) ⁸²

17% of persons seroconverted toward *Schistosoma mansoni* and / or *Schistosoma haematobium* following recreational exposure to the Upper Nile River (2007) ⁸³

Seroprevalence surveys:

- 90% of semi-captive wild-born chimpanzees on an island sanctuary in Lake Victoria (2011 publication) ^{84 85}

Schistosoma mansoni infection is found among chimpanzees on Ngamba Island. ⁸⁶⁻⁸⁹

Reservoirs:

- Local snail reservoirs include *Biomphalaria pfeifferi*, *Biomphalaria sudanica* ⁹⁰, *Biomphalaria stanleyi* ^{91 92} and *Biomphalaria choanophala*. ^{93 94}

References

1. Lancet 2006 Sep 23;368(9541):1106-18.
2. Lancet Infect Dis 2007 Mar ;7(3):218-24.
3. Trans R Soc Trop Med Hyg 2010 Jun ;104(6):443-5.
4. Nihon Hinyokika Gakkai Zasshi 2010 Jul ;101(5):694-7.
5. Parasite Immunol 2009 Nov ;31(11):656-63.
6. Parasite Immunol 2009 Nov ;31(11):686-96.
7. Cancer Lett 2011 Jun 28;305(2):239-49.
8. J Infect Dev Ctries 2010 May ;4(5):267-81.
9. Am J Trop Med Hyg 1970 Sep ;19(5):779-84.
10. AIDS 2006 Feb 28;20(4):593-600.
11. PLoS Negl Trop Dis 2011 Dec ;5(12):e1396.
12. Ann Trop Med Parasitol 2009 Mar ;103(2):129-43.
13. Chest 2010 Jun ;137(6 Suppl):20S-29S.
14. Travel Med Infect Dis 2012 Sep 11;
15. J Neurol 2012 Jan ;259(1):22-32.
16. Pediatr Infect Dis J 2011 Nov ;30(11):1006-8.
17. Ann Indian Acad Neurol 2011 Apr ;14(2):107-10.
18. Lancet Neurol 2011 Sep ;10(9):853-64.
19. BMC Infect Dis 2012 ;12:220.
20. Neurologist 2012 Nov ;18(6):333-42.
21. Neurochirurgie 2012 May 29;
22. Rev Med Interne 2012 Oct ;33(10):580-2.
23. Curr Neurol Neurosci Rep 2012 Dec ;12(6):666-74.
24. Trans R Soc Trop Med Hyg 2008 Feb ;102(2):107-16.
25. Lancet Neurol 2011 Sep ;10(9):853-64.
26. BMC Infect Dis 2012 ;12:220.
27. Neurologist 2012 Nov ;18(6):333-42.
28. Q J Med 1986 Dec ;61(236):1131-9.
29. Acta Trop 2008 Nov-Dec;108(2-3):89-97.
30. J Infect 2010 Mar ;60(3):244-7.
31. Neurol Res 2010 Apr ;32(3):252-62.
32. Am J Trop Med Hyg 2009 Oct ;81(4):551-4.
33. J Neurol 2012 Jan ;259(1):22-32.
34. Pediatr Infect Dis J 2011 Nov ;30(11):1006-8.
35. Ann Indian Acad Neurol 2011 Apr ;14(2):107-10.
36. Lancet Neurol 2011 Sep ;10(9):853-64.
37. BMC Infect Dis 2012 ;12:220.
38. Neurologist 2012 Nov ;18(6):333-42.
39. Neurochirurgie 2012 May 29;
40. Rev Med Interne 2012 Oct ;33(10):580-2.
41. Curr Neurol Neurosci Rep 2012 Dec ;12(6):666-74.
42. J Clin Microbiol 2011 Oct ;49(10):3703-6.
43. Rev Soc Bras Med Trop 2009 Sep-Oct;42(5):581-2.
44. World J Surg Oncol 2010 ;8:68.
45. Afr J Reprod Health 2009 Sep ;13(3):137-40.
46. Acta Trop 2001 Jun 22;79(3):193-210.
47. Parasitol Today 1999 Sep ;15(9):378-81.
48. Am J Trop Med Hyg 2009 Oct ;81(4):549-50.
49. Trends Parasitol 2012 Feb ;28(2):58-65.
50. Trop Gastroenterol 2009 Oct-Dec;30(4):230-2.
51. Ann Pathol 2012 Feb ;32(1):40-52.
52. Braz J Infect Dis 2010 May-Jun;14(3):288-90.
53. Braz J Infect Dis 2011 Mar-Apr;15(2):174-7.
54. Arch Gynecol Obstet 2010 Jan ;281(1):141-3.
55. Fertil Steril 2011 May ;95(6):2124.e1-4.
56. Trans R Soc Trop Med Hyg 2009 Oct ;103(10):1068-70.
57. Ann Dermatol Venereol 2012 Feb ;139(2):132-6.
58. J Trop Med Hyg 1977 Jan ;80(1):14-8.
59. Mem Inst Oswaldo Cruz 1998 ;93 Suppl 1:135-9.
60. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):121.
61. Arch Intern Med 1971 Aug ;128(2):254-7.
62. Rev Soc Bras Med Trop 2009 Jul-Aug;42(4):436-45.
63. Trop Med Int Health 2004 Mar ;9(3):372-80.
64. Acta Trop 2012 Feb 22;
65. Afr Health Sci 2008 Dec ;8(4):239-43.
66. East Afr Med J 1997 May ;74(5):311-4.
67. Geospat Health 2009 Nov ;4(1):39-53.
68. Trans R Soc Trop Med Hyg 2006 Nov ;100(11):1039-48.
69. BMC Infect Dis 2008 ;8:110.
70. Trop Med Int Health 2010 Feb ;15(2):198-207.
71. Parasit Vectors 2011 ;4:232.
72. PLoS Negl Trop Dis 2011 ;5(1):e938.
73. Am J Trop Med Hyg 2010 Nov ;83(5):1048-55.
74. Ann Trop Med Parasitol 2006 Jun ;100(4):315-26.
75. Trans R Soc Trop Med Hyg 2004 Dec ;98(12):711-8.
76. Trop Med Int Health 2009 Jun ;14(6):612-9.
77. ProMED <promedmail.org> archive: 20110413.1162
78. Trans R Soc Trop Med Hyg 2006 Mar ;100(3):216-23.
79. Trans R Soc Trop Med Hyg 2007 Sep ;101(9):899-907.
80. J Infect Dis 2008 Sep 15;198(6):920-7.
81. Clin Infect Dis 2010 Feb 15;50(4):531-40.
82. Trans R Soc Trop Med Hyg 2008 Feb ;102(2):130-6.
83. Emerg Infect Dis 2010 May ;16(5):866-8.
84. Vector Borne Zoonotic Dis 2011 Feb ;11(2):169-76.
85. Trends Parasitol 2012 Aug ;28(8):320-6.
86. Parasitology 2012 Oct 25;144:1-11.
87. Trends Parasitol 2012 Aug ;28(8):320-6.
88. J Parasitol 2012 Oct ;98(5):1049-51.
89. Vector Borne Zoonotic Dis 2011 Feb ;11(2):169-76.
90. Parasitologia 2005 Mar ;47(1):115-25.
91. Trop Med Parasitol 1992 Sep ;43(3):162-6.
92. Parasitol Int 2010 Mar ;59(1):49-53.
93. East Afr Med J 1996 Aug ;73(8):495-8.
94. East Afr Med J 1992 Jun ;69(6):316-8.

Septic arthritis

Agent	BACTERIUM or FUNGUS. Gram positive cocci most common; gram negative bacilli, gonococci, mycobacteria, fungi, et al
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Smear and culture of joint fluid. Cytological and chemical analysis of joint fluid also useful.
Typical Adult Therapy	Antimicrobial agent(s) directed at known or likely pathogen
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever (60% to 80%) associated with swelling, erythema and tenderness (usually single joint, most commonly a knee; elbow or ankle in child); mean fluid leukocyte count in acute bacterial forms = 50,000 / cu mm.
Synonyms	

Clinical

Most cases present with fever, malaise and local findings of warmth, swelling and decreased range of motion. ^{1 2}

- Lack of erythema and local warmth are not uncommon.
- The most commonly involved joints are the knee and hip, followed by the shoulder and ankle. ³
- Non-gonococcal arthritis is mono-articular in 80% to 90% of cases.
- Infection of the costochondral, sternoclavicular and sacroiliac joints is common in intravenous drug users.

Synovial fluid demonstrates low viscosity and turbidity.

- Leucocyte counts usually exceed 50,000 per cu mm.
- Note that leucocytosis, low glucose and high lactate levels are also encountered in some non-infectious forms of arthritis.
- Gram stains are positive in 50% of cases, and cultures in 90%.

Etiological associations:

- Adult below age 30: *Neisseria gonorrhoeae* (often monoarticular involving knee)
- Associated rash: Lyme disease, gonococcemia (often monoarticular, involving knee)
- Child below age 5 years: *Haemophilus influenzae*, *Staphylococcus aureus*, *Streptococcus* spp.
- Chronic arthritis: Tuberculosis, Mycobacteria • nontuberculous, Sporotrichosis and other fungi
- Hematogenous infection: *Staphylococcus aureus*, *Streptococcus pyogenes*
- Injecting drug user: *Pseudomonas aeruginosa* (often sternoclavicular or sacroiliac)
- Traumatic injury to joint: *Staphylococcus aureus*, Enterobacteriaceae, *Pseudomonas aeruginosa*

This disease is endemic or potentially endemic to all countries.

References

1. J Paediatr Child Health 2005 Jan-Feb;41(1-2):59-62.
2. Clin Microbiol Infect 2006 Apr ;12(4):309-14.
3. Clin Microbiol Rev 2002 Oct ;15(4):527-44.

Septicemia - bacterial

Agent	BACTERIUM. Escherichia coli , Staphylococcus aureus , facultative gram negative bacilli, et al
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Culture of blood and sepsis source.
Typical Adult Therapy	Antimicrobial agent(s) directed at known or likely pathogen
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, rigors, leukocytosis, tachypnea, mental changes; hypotension, acidosis and bleeding diathesis herald septic shock; further signs (eg, urinary infection, phlebitis, etc) may point to the source of infection .
Synonyms	Sepsis, Septicaemia, Septicemia, Septicemie, Septikemie, Setticeimia. ICD9: 036.2,036.3,038 ICD10: A40,A41

Clinical

Bacterial septicemia is defined as the presence of signs and symptoms related to bacteremia. ¹

- The clinical spectrum and severity of disease are largely determined by the infecting species, underlying diseases and source of infection.
- Most patients present with fever, tachycardia and leucocytosis, in addition to signs and symptoms referable to a primary infectious focus (eg, urinary tract, abdominal infection, endocarditis, etc).

This disease is endemic or potentially endemic to all countries.

References

1. [Scand J Infect Dis 2003 ;35\(9\):529-34.](#)

Shigellosis

Agent	BACTERIUM. <i>Shigella sonnei</i> , <i>Shigella flexneri</i> , <i>Shigella boydii</i> or <i>Shigella dysenteriae</i> A facultative gram-negative bacillus
Reservoir	Human Non-human primate
Vector	None
Vehicle	Fecal-oral Water Dairy products Fomite Fly Vegetables
Incubation Period	48h - 72h (range 7h - 1w)
Diagnostic Tests	Stool culture.
Typical Adult Therapy	Stool precautions. Choice of antimicrobial agent based on regional susceptibility patterns. Continue treatment for five days
Typical Pediatric Therapy	As for adult
Clinical Hints	Watery or bloody diarrhea, tenesmus, abdominal pain and headache; colonic hyperemia and abundant fecal leucocytes are present; usually resolves in 3 days (may persist for up to 14); case fatality rate = 1%.
Synonyms	Bacillaire dysenterie, Bacillary dysentery, Dissenteria batterica, Dysentery bacillaris, Leptospiroserkrankung, Ruhr, Shigella, Shigellose, Shigelose, Ubertragbare Ruhr. ICD9: 004 ICD10: A03

Clinical

Acute infection:

Approximately 50% of infections are limited to transient fever or self-limited diarrhea.

- 50% of patients progress to bloody diarrhea and dysentery. ¹
- Fever may rise rapidly to 40 C, and febrile seizures are common in children.
- Seizures rarely recur or result in neurological sequelae.
- Dysentery is characterized by passage of 10 to 30 small-volume stools consisting of blood, mucus, and pus.
- Abdominal cramps and tenesmus are noted, and straining may lead to rectal prolapse, notably in young children. ²
- On endoscopy, the colonic mucosa is hemorrhagic, with mucous discharge and focal ulcerations. Most lesions are in the distal colon.

Complications:

Patients with mild disease generally recover without specific therapy in two to seven days.

- Severe shigellosis can progress to toxic dilatation or perforation of the colon, which may be fatal.
- Mild dehydration is common, and protein-losing enteropathy can occur with severe disease.
- Complications are most commonly described in developing countries and are related both to the relative prevalence of *S. dysenteriae* type 1 and *S. flexneri*, and the poor nutritional state of the local populations.
- *Shigella* bacteremia is not uncommon, and is associated with increased mortality, particularly among infants below one year of age and persons with protein-energy malnutrition.
- Hemolytic-uremic syndrome (HUS) may complicate shigellosis due to *S. dysenteriae* type 1, and usually develops toward the end of the first week of shigellosis. ³⁻⁵ The case-fatality rate in these cases is 36%. ⁶
- Profound hyponatremia and hypoglycemia may occur.
- Other complications include encephalopathy ⁷, seizures, altered consciousness, and bizarre posturing, pneumonia ⁸, meningitis, vaginitis, keratoconjunctivitis ⁹, pneumonia and "rose spots."
- Reiter's syndrome is seen in patients having histocompatibility antigen HLA-B27. ^{10 11}
- Reactive arthritis follows 7% to 10% of *Shigella* infections. ¹²⁻¹⁵

This disease is endemic or potentially endemic to all countries.

Shigellosis in Uganda

56,734 cases of dysentery were reported in 1999.

References

1. Rev Infect Dis 1991 Mar-Apr;13 Suppl 4:S220-5.
2. Semin Pediatr Infect Dis 2004 Oct ;15(4):246-52.
3. J Med Assoc Thai 1990 Jul ;73(7):401-5.
4. Cent Afr J Med 1995 Sep ;41(9):267-74.
5. J Health Popul Nutr 2012 Sep ;30(3):257-61.
6. Trans R Soc Trop Med Hyg 2012 May 9;
7. Pediatr Infect Dis J 2010 May ;29(5):444-7.
8. Emerg Infect Dis 2009 Nov ;15(11):1874-6.
9. J Clin Microbiol 2006 Jun ;44(6):2291-4.
10. Ann Rheum Dis 1979 ;38 Suppl 1:suppl 119-22.
11. J Rheumatol 1981 Nov-Dec;8(6):969-73.
12. Ann Rheum Dis 2005 Apr ;64(4):594-8.
13. J Rheumatol 2008 Mar ;35(3):480-7.
14. Rev Rhum Engl Ed 1999 Jan 30;66(1 Suppl):14S-18S; discussion 19S.
15. J Clin Rheumatol 2012 Aug ;18(5):257-8.

Sindbis

Agent	VIRUS - RNA. Togaviridae, Alphavirus: Sindbis virus
Reservoir	Wild bird
Vector	Mosquito (Culex univittatus and Cx. tritaeniorhynchus)
Vehicle	None
Incubation Period	3d - 6d
Diagnostic Tests	Viral culture (blood, vesicle fluid). Serology. Nucleic acid amplification. Biosafety level 2.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, myalgia, arthritis and a papular-to-vesicular rash; arthralgias may persist for more than three years; fatality not reported.
Synonyms	Babanki, Whataroa. ICD9: 078.89 ICD10: A92.8

Clinical

Sindbis virus infection is a self-limited febrile illness associated with myalgia and diffuse papular or vesicular rash (typically of the plantar region). ¹

- Arthralgia is common, and may be severe.
- Fever and rash persist for 2 to 3 weeks; and arthralgias for months, or even years.
- No deaths have been reported.

Infection by a related agent, Babanki virus, is characterized by fever, rash and arthralgia.

This disease is endemic or potentially endemic to 33 countries.

Sindbis in Uganda

Seroprevalence surveys:

12.6% of the general population (1967 to 1969, eight areas) ²

References

1. J Intern Med 2004 Dec ;256(6):457-71.
2. Bull World Health Organ 1970 ;42(5):797-805.

Sinusitis

Agent	BACTERIUM. Various (Haemophilus influenzae & Streptococcus pneumoniae in most acute cases)
Reservoir	Human
Vector	None
Vehicle	None
Incubation Period	Variable
Diagnostic Tests	Imaging techniques. Culture of sinus drainage.
Typical Adult Therapy	Antimicrobial agent(s) directed at likely pathogens. Drainage as indicated
Typical Pediatric Therapy	As for adult
Clinical Hints	Sinusitis often follows upper respiration infections; headache, fever and local tenderness are common, however the precise presentation varies with patient age and anatomic localization.
Synonyms	Acute sinusitis, Mastoidite, Mastoiditis, Rhinosinusitis, Sinusite. ICD9: 473.9,383.0,461 ICD10: H70,J01

Clinical

Acute community-acquired bacterial sinusitis is usually superimposed on preexisting viral sinusitis.

- In most cases, it is not possible to distinguish between viral and bacterial infections.
- Sneezing, nasal discharge and obstruction, facial pressure and headache are common in both conditions. ¹
- Fever of 38C or more, facial pain, and erythema occur may occasionally herald bacterial infections.
- The nasal discharge may be colored in both viral and bacterial sinusitis.
- Cough and hyposmia may also be present.

Sinusitis following dental infection is associated with molar pain and a foul breath odor.

- Sphenoid sinusitis is associated with severe frontal, temporal, or retroorbital headache that radiates to the occipital region; and hypesthesia or hyperesthesia of the ophthalmic or maxillary dermatomes of the fifth cranial nerve.
- Lethargy and findings suggestive of cavernous sinus or cortical vein thrombosis, orbital cellulitis or orbital abscess may also be present.
- In severe cases of frontal sinusitis, pus may collect under the periosteum of the frontal bone resulting in a 'Pott puffy tumor.'

Rare instances of toxic shock syndrome have followed sinusitis. ²

This disease is endemic or potentially endemic to all countries.

References

1. *N Engl J Med* 2004 Aug 26;351(9):902-10.
2. *Arch Otolaryngol Head Neck Surg* 2009 Jun ;135(6):538-42.

Smallpox

Agent	VIRUS - DNA. Poxviridae, Orthopoxvirus: Variola virus
Reservoir	Human
Vector	None
Vehicle	Contact Infected secretions Fomite
Incubation Period	7d - 17d
Diagnostic Tests	Culture and electron microscopy of skin lesions. Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Isolation; supportive. Cidofovir is effective in vitro
Typical Pediatric Therapy	As for adult
Vaccine	Smallpox
Clinical Hints	Fever, myalgia, headache, pustular or hemorrhagic rash; disease resolves in 2 to 3 weeks; case-fatality rate = 25% for severe form (variola major) and 1% for minor form; last naturally-acquired case reported in Somalia in 1977.
Synonyms	Alastrim, Eczema vaccinatum, Kopper, Smallpox, Vailo, Variola, Variola minor, Varioloid. ICD9: 050 ICD10: B03

Clinical

Acute infection: ^{1 2}

12 to 14 days after exposure (range 7 to 17 days), the patient experiences a 2 to 3 day prodrome of high fever, malaise, prostration and severe headache and backache.

- This "preeruptive stage" is followed by the appearance of a maculopapular rash (i.e., eruptive stage) that progresses to papules within one to two days.
- Vesicles appear on the fourth or fifth day; pustules by the seventh day; and scab lesions on the fourteenth day.
- The rash first appears on the oral mucosa, face, and forearms; and then spreads to the trunk and legs. ³
- The palms and soles may also be involved.
- Skin lesions are deeply embedded in the dermis and feel like firm round objects in the skin.
- As the lesions heal, the scabs separate and pitted scarring gradually develops.
- Patients are most infectious during the first week of the rash when the oral mucosa lesions ulcerate and release large amounts of virus into the saliva.
- A patient is no longer infectious after all scabs have separated (3 to 4 weeks after the onset of the rash).
- Rare instances of bone involvement (osteomyelitis variolosa) are described. ⁴⁻⁶
- During the smallpox era, overall mortality rates were approximately 30%.

Other less common but more severe forms of smallpox include

- a) flat-type smallpox (mortality rate over 96%) characterized by severe toxemia and flat, velvety, confluent lesions that did not progress to the pustular stage or scarring
- b) hemorrhagic smallpox, characterized by severe prodromal symptoms, toxemia, and a hemorrhagic rash.

Hemorrhagic smallpox is uniformly fatal and occur among all ages and in both sexes, but pregnant women appear to be unusually susceptible. ^{7 8}

- Illness usually begins with a somewhat shorter incubation period and is characterized by high fever and pain in the head, back, and abdomen.
- Soon thereafter, a dusky erythema develops, followed by petechiae and frank hemorrhages into the skin and mucous membranes.
- Death usually occurs by the fifth or sixth day after onset of rash.


Variola minor is generally less severe, with fewer constitutional symptoms and a more sparse rash.

- A milder form of disease is also seen among those who have residual immunity from previous vaccination.
- In partially immune persons, the rash tends to be atypical and more scant and the evolution of the lesions more rapid.

Disseminated herpes simplex in patients with eczema (Eczema herpeticum) may resemble smallpox. ⁹

This disease is not currently endemic to any country. Although Smallpox is not endemic to Uganda, imported, expatriate or other presentations of the disease have been associated with this country.

Smallpox in Uganda



Graph: Uganda. Smallpox, cases

Four fatal cases were reported during 1961 to 1964; 63 in 1965; 25 in 1966; 24 in 1967; 5 in 1968; 0 in 1969; 0 in 1970.

1,045 cases of variola major (54 fatal) were reported during 1966 to 1970.

Indigenous transmission ended in 1968.

Importations from Sudan were reported in 1969; from Rwanda, Zaire and Sudan in 1970; from Sudan in 1971; from Sudan in 1972.

References

1. Emerg Infect Dis 1999 Jul-Aug;5(4):537-9.
2. Wkly Epidemiol Rec 2001 Nov 2;76(44):337-44.
3. JAMA 1997 Aug 6;278(5):399-411.
4. J Orthop Surg (Hong Kong) 2008 Dec ;16(3):355-8.
5. Rheumatol Int 2011 Sep ;31(9):1231-3.
6. J Orthop Surg (Hong Kong) 2011 Apr ;19(1):120-2.
7. Bull World Health Organ 1965 ;33(5):607-13.
8. Bull World Health Organ 1965 ;33(6):773-82.
9. Emerg Infect Dis 2009 Jul ;15(7):1102-4.

Sparganosis

Agent	PARASITE - Platyhelminthes, Cestoda. Pseudophyllidea, Diphyllbothriidae: Spirometra spp.
Reservoir	Copepod - to bird, amphibian or reptile
Vector	None
Vehicle	Water Undercooked reptile or amphibian meat
Incubation Period	20d - 3y
Diagnostic Tests	Identification of parasite in tissue.
Typical Adult Therapy	Excision
Typical Pediatric Therapy	As for adult
Clinical Hints	Painful or pruritic nodules and eosinophilia; worm present in skin, eye, brain or other foci and may survive for over five years.
Synonyms	

Clinical

The worm usually lodges in subcutaneous tissue or muscle in the chest, abdominal wall, extremities, or scrotum.

- Infection typically presents as a nodular mass, swelling and painful edema.
- Orbital sparganosis is also common, and is manifest by erythema of the lids and conjunctivae, proptosis and the presence of fixed or migrating subconjunctival masses. ¹
- 35 cases of sparganosis of the breast were identified in the literature as of 2010. ²
- Less common sites have included the urinary tract, pleura, pericardium ³, brain ⁴⁻⁶, spinal canal ^{7 8}, scrotum ^{9 10}, articular bursa ¹¹, liver ¹² and other abdominal viscera.
- Spinal disease may present as a lumbar radiculopathy ¹³⁻¹⁵ or mimic intramedullary tumor. ¹⁶
- Pulmonary ¹⁷ or cerebral sparganosis may mimic tuberculosis ¹⁸
- Genital sparganosis presents as a palpable subcutaneous nodule in the groin, labia, or scrotum and may mimic a tumor in the epididymis and testis or hydrocoele. ¹⁹⁻²¹
- The patient may notice "lumps" which appear and then spontaneously disappears, over a period of weeks to months.
- The overlying skin is redness and pruritic, and local bleeding or necrosis may occur.

This disease is endemic or potentially endemic to 49 countries.

References

1. Can J Ophthalmol 2012 Oct ;47(5):453-7.
2. World J Surg 2011 Mar ;35(3):573-9.
3. Korean Circ J 2011 Jan ;41(1):38-42.
4. Rev Infect Dis 1991 Jan-Feb;13(1):155-9.
5. J R Army Med Corps 2007 Sep ;153(3):189-90.
6. AJNR Am J Neuroradiol 2012 Aug 2;
7. J Korean Neurosurg Soc 2011 Apr ;49(4):241-4.
8. J Korean Neurosurg Soc 2012 Mar ;51(3):170-2.
9. Urology 2008 Feb ;71(2):351.e11-2.
10. Korean J Parasitol 2010 Mar ;48(1):57-9.
11. Clin Orthop Relat Res 2011 Jul ;469(7):2072-4.
12. BMJ Case Rep 2012 ;2012
13. Clin Neurol Neurosurg 2008 Sep ;110(8):843-6.
14. J Korean Neurosurg Soc 2011 Apr ;49(4):241-4.
15. J Korean Neurosurg Soc 2012 Mar ;51(3):170-2.
16. J Clin Neurosci 2011 Aug ;18(8):1128-9.
17. Trop Biomed 2012 Jun ;29(2):220-3.
18. Br J Neurosurg 2008 Dec ;22(6):784-6.
19. Urology 2007 Dec ;70(6):1223.e1-2.
20. J Ultrasound Med 2010 Nov ;29(11):1627-33.
21. Korean J Parasitol 2012 Dec ;50(4):353-5.

Spondweni

Agent	VIRUS - RNA. Flaviviridae, Flavivirus: Spondweni virus
Reservoir	Unknown
Vector	Mosquito (Aedes circumluteolus, Armigeres, Culex, Eretmapodites, Mansonia)
Vehicle	None
Incubation Period	Unknown
Diagnostic Tests	Viral culture (blood). Serology. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, headache, myalgia and a pruritic maculopapular rash; fatality and sequelae have not been reported.
Synonyms	Banzi, Koutango, Sepik. ICD9: 078.89 ICD10: A92.8

Clinical

Spondweni is a mild illness, limited to fever, myalgia and a maculopapular rash. ¹

- No fatal cases have been reported.

Infection by a related agent, Koutango virus, is characterized by fever, headache, eye pain, rash and arthralgia.

This disease is endemic or potentially endemic to 10 countries.

Spondweni in Uganda

Banzi virus infection has been documented in this country.

Seroprevalence surveys:

8.9% of the general population (Banzi virus, 1967 to 1969, eight areas) ²

References

1. Lancet 1982 Dec 11;2(8311):1306-8.
2. Bull World Health Organ 1970 ;42(5):797-805.

Sporotrichosis

Agent	FUNGUS. Ascomycota, Euascomycetes, Ophiostomatales: <i>Sporothrix schenckii</i> , <i>S. brasiliensis</i> and <i>S. globosa</i> A dimorphic dematiaceous fungus
Reservoir	Soil Vegetation Wood
Vector	None
Vehicle	Trauma Contact Air (rare)
Incubation Period	1w - 3m
Diagnostic Tests	Fungal culture. Serologic tests available in some centers.
Typical Adult Therapy	Itraconazole 100 to 200 mg PO daily X 3 to 6 months. OR Fluconazole 400 mg PO daily X 6 months. OR Potassium iodide 1 to 5 ml PO TID X 3 to 6 months
Typical Pediatric Therapy	Itraconazole 2 mg/kg PO daily X 3 to 6 months. OR Fluconazole 3 mg/kg PO daily X 6 months.
Clinical Hints	Draining nodules which follow lymphatics; acquired from contact with flowers, thorns, trees or other plant material; eye, brain, testis, bone and other tissues may be involved.
Synonyms	Schenck's disease, <i>Sporothrix brasiliensis</i> , <i>Sporothrix globosa</i> , <i>Sporothrix mexicana</i> , <i>Sporothrix schenckii</i> , Sporotrichose. ICD9: 117.1 ICD10: B42

Clinical

Clinical forms of sporotrichosis:

Cutaneous sporotrichosis begins as a painless erythematous papule which enlarges and suppurates, without systemic symptoms. ^{1 2}

- Multiple lesions may spread along lymphatic channels. ³
- Occasionally only a single lesion appears, and may persist for decades.
- Bilateral infection may occur. ⁴
- Hematogenous infection of multiple skin sites has also been described. ^{5 6}
- In some cases, ulcers appear on multiple body sites. ⁷

Nodular lymphadenitis is also seen in *Nocardia brasiliensis* infection, tularemia, *Mycobacterium marinum* infection, and infections caused by *Leishmania panamensis/guyanensis* ^{8 9}

- Lesions of sporotrichosis may rarely mimic those of pyoderma gangrenosum ¹⁰ or keratoacanthoma. ¹¹⁻¹³

Pulmonary sporotrichosis characteristically presents as a single upper lobe cavity associated with cough and low-grade fever.

- Multifocal lung lesions have also been reported. ¹⁴
- 86 cases of pulmonary sporotrichosis were reported in the world's literature during 1960 to 2010. ¹⁵ Extrapulmonary multifocal disease involved the joints in 45.4%.

Osteoarticular sporotrichosis is characterized by infection of a single bone or large peripheral joint ¹⁶ • hip and shoulder involvement is not encountered. ¹⁷⁻²⁰

- Most patients are afebrile when first seen.
- Occasionally, the infection presents as tenosynovitis, usually of the wrist or ankle.

Other forms include conjunctival infection ^{21 22} , hematogenous endophthalmitis, brain abscess, soft tissue mass ²³ , meningitis ²⁴ , orchitis, etc. ²⁵

This disease is endemic or potentially endemic to all countries.

References

1. Infect Dis Clin North Am 2003 Mar ;17(1):59-85, viii.
2. Clin Dermatol 2012 Jul ;30(4):437-43.

3. Dermatol Clin 1996 Jan ;14(1):69-76.
4. Rev Iberoam Micol 2009 Dec 31;26(4):247-9.

5. Med Mycol 2012 Feb ;50(2):197-201.
6. Int J Infect Dis 2011 Oct ;15(10):e727-9.
7. Clin Exp Dermatol 2008 Mar ;33(2):135-8.
8. Curr Infect Dis Rep 2008 Sep ;10(5):404-10.
9. Ann Dermatol Venereol 2008 Jan ;135(1):63-7.
10. Mycopathologia 1991 Dec ;116(3):165-8.
11. Am J Trop Med Hyg 2012 May ;86(5):741.
12. Cutis 1998 Jul ;62(1):37-9.
13. J Dermatol Surg 1976 Jun ;2(3):209-14.
14. Rev Port Pneumol 2008 May-Jun;14(3):443-9.
15. Med Mycol 2013 Jan 4;
16. Rev Iberoam Micol 2008 Mar ;25(1):54-6.
17. Acta Derm Venereol 2012 Aug 1;
18. J Oral Surg 1981 Jun ;39(6):468-72.
19. Skeletal Radiol 1984 ;12(1):23-8.
20. J Infect 1989 Nov ;19(3):273-6.
21. Cornea 2005 May ;24(4):491-3.
22. Cornea 2010 May ;29(5):573-6.
23. Malays J Pathol 2009 Dec ;31(2):143-5.
24. J Neurol Neurosurg Psychiatry 2010 Jun ;81(6):696-9.
25. Cutis 2006 Oct ;78(4):253-6.

Spotted fevers - Old World

Agent	BACTERIUM. <i>Rickettsia conorii</i> subsp. <i>Conorii</i> R. <i>aeschlimannii</i> , R. <i>helvetica</i> , R. <i>massiliae</i> , R. <i>monacensis</i> , R. <i>slovaca</i>
Reservoir	Dog Rodent Tick
Vector	Tick (<i>Rhipicephalus sanguineus</i>)
Vehicle	None
Incubation Period	6d - 7d (range 3d - 18d)
Diagnostic Tests	Serology. Demonstration of rickettsiae by immunofluorescence or culture. Nucleic acid amplification.
Typical Adult Therapy	Doxycycline 100 mg PO BID X 3 to 5d. OR Chloramphenicol 500 mg PO QID X 3 to 5d
Typical Pediatric Therapy	Doxycycline 2 mg/kg PO BID X 3 to 5d (maximum 200 mg/day). OR Chloramphenicol 10 mg/kg PO QID X 3 to 5d
Clinical Hints	Headache, myalgia, maculopapular rash; an eschar may be identifiable; patient may recall tick bite or dog contact during the preceding 1 to 3 weeks; untreated disease resolves within two weeks; case-fatality rates of 2% to 3% are reported.
Synonyms	Boutonneuse fever, Candidatus <i>Rickettsia kellyi</i> , DEBONEL, Febre escaro-nodular, Febre escaronodular, Indian tick typhus, Kenya tick typhus, Marseilles fever, Mediterranean spotted fever, R. <i>aeschlimannii</i> , <i>Rickettsia aeschlimannii</i> , <i>Rickettsia conorii</i> subsp <i>conorii</i> , <i>Rickettsia conorii</i> subsp <i>indica</i> , <i>Rickettsia helvetica</i> , <i>Rickettsia massiliae</i> , <i>Rickettsia monacensis</i> , <i>Rickettsia raoultii</i> , <i>Rickettsia slovaca</i> , Thai spotted fever, TIBOLA, Tick-borne lymphadenopathy. ICD9: 082.1 ICD10: A77.1

Clinical

The clinical features of Mediterranean spotted fever (MSM) are similar to those of Rocky Mountain spotted fever (q.v.); however, an eschar ("tache noire") and diffuse distribution of the rash characterize MSM. ^{1 2}

- Hepatomegaly, elevation of serum transaminase levels and splenomegaly are common. ³
- Complications may include meningitis with CSF pleocytosis (either lymphocytic or polymorphonuclear), encephalitis ⁴⁻⁶, renal failure ⁷, myocarditis ⁸, coronary artery ectasia ⁹, acute pancreatitis ¹⁰, bleeding diatheses, splenic rupture ¹¹, hemophagocytic syndrome ^{12 13} and retinitis. ¹⁴
- There is evidence that Israeli spotted fever is more virulent than Mediterranean spotted fever. ¹⁵

Spotted fever in India differs from the Mediterranean form in that the rash is often purpuric, and an inoculation eschar at the bite site is rarely found. ¹⁶

- The clinical course is mild to moderately severe.

A syndrome of Dermacentor-borne necrosis with erythema and painful lymphadenopathy (DEBONEL) described in Spain has been ascribed to possible infection by *Rickettsia slovaca*. ^{17 18}

- This syndrome appears to be identical to Tick-borne lymphadenopathy (TIBOLA), reported in Hungary. ¹⁹
- Clinical features may include fever, dermal eschar, lymphadenopathy, facial edema, rash, headache, asthenia and alopecia.
- Rarely, tularemia presenting with scalp eschar and cervical lymphadenopathy may suggest infection by *Rickettsia slovaca*, *Rickettsia raoultii* or *Bartonella henselae* ²⁰

Rickettsia helvetica has been implicated in cases of mild flu-like illness (myalgia, arthralgia, headache, conjunctivitis) without rash, in Denmark, Italy, France and Thailand; and in myocarditis reported from Sweden. ²¹

Rickettsia monacensis infection has been associated with headache, joint pain, a nonpruritic, disseminated maculopapular rash of the trunk and lower extremities, including palms and soles.

- An inoculation site eschar is not reported.

This disease is endemic or potentially endemic to 104 countries.

Spotted fevers - Old World in Uganda

Rickettsia conorii was identified in a tick (*Haemaphysalis punctataechei*) in 2003. ²²

References

1. Rev Infect Dis 1985 Sep-Oct;7(5):635-42.
2. BMC Infect Dis 2006 ;6:60.
3. Infez Med 2007 Jun ;15(2):105-10.
4. J Med Microbiol 2009 Apr ;58(Pt 4):521-5.
5. Rev Neurol (Paris) 2011 Feb ;167(2):173-6.
6. J Infect Chemother 2012 Feb ;18(1):105-8.
7. Nephrol Ther 2011 Jul ;7(4):245-7.
8. Ann Cardiol Angeiol (Paris) 2011 May 25;
9. New Microbiol 2011 Oct ;34(4):421-4.
10. Acta Gastroenterol Belg 2011 Mar ;74(1):91-2.
11. Emerg Infect Dis 2008 Jun ;14(6):995-7.
12. J Med Microbiol 2011 Apr ;60(Pt 4):537-42.
13. Int J Infect Dis 2013 Feb 8;
14. Infection 2008 Aug ;36(4):384-6.
15. Ann N Y Acad Sci 2003 Jun ;990:285-94.
16. Int J Dermatol 1991 Nov ;30(11):790-4.
17. Clin Microbiol Rev 2005 Oct ;18(4):719-56.
18. Ann N Y Acad Sci 2006 Oct ;1078:206-14.
19. Wien Klin Wochenschr 2002 Jul 31;114(13-14):648-54.
20. J Med Case Reports 2011 ;5:108.
21. Ann Trop Med Parasitol 2005 Apr ;99(3):325-30.
22. Emerg Infect Dis 2007 Oct ;13(10):1508-10.

Staphylococcal food poisoning

Agent	BACTERIUM. <i>Staphylococcus aureus</i> exotoxins
Reservoir	Human (nares, hands) Occasionally cattle (udder), dog/cat (nasopharyngeal)
Vector	None
Vehicle	Food (creams, gravies, sauces)
Incubation Period	2h - 4h (range 30 min - 9h)
Diagnostic Tests	Identification of bacterium in food.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	'Explosive' diarrhea and vomiting; usually no fever; no fecal leucocytes; onset 1 to 6 hours after food; resolves within 1 to 2 days; fatality is rare.
Synonyms	Staphylococcus aureus food poisoning. ICD9: 005.0 ICD10: A05.0

Clinical

Usually symptoms start within several hours of ingestion of potentially contaminated foods

- Illness is heralded by nausea, vomiting and intestinal cramping, followed by urgency and profuse watery non-bloody diarrhea.
- Symptoms resolve within 12 to 24 hours.
- Multiple family members or patrons of the same eating establishment may be affected.
- The presence of both explosive diarrhea and vomiting, lack of fever and short incubation period are helpful in distinguishing this entity from other forms of food poisoning.

This disease is endemic or potentially endemic to all countries.

Staphylococcal scalded skin syndrome

Agent	BACTERIUM. Staphylococcus aureus phage group 2 A facultative gram-positive coccus
Reservoir	Human
Vector	None
Vehicle	Direct contact; infected secretions
Incubation Period	1d - 4d
Diagnostic Tests	Typical clinical features; Recovery of <i>S. aureus</i> from localized wound or blood ; skin biopsy may be helpful
Typical Adult Therapy	Fluid replacement (as for burn) ; Intravenous Nafcillin or Oxacillin , in addition to application of anti-staphylococcal drug to local source infection; Vancomycin if MRSA
Typical Pediatric Therapy	Fluid replacement (as for thermal burn) ; Intravenous Nafcillin or Oxacillin , in addition to application of anti-staphylococcal drug to local source infection; Vancomycin if MRSA
Clinical Hints	Acute, generalized exfoliative dermatitis which occurs primarily in infants and young children; a pre-existing localized skin infection is present in most - but not all - cases.
Synonyms	Lyell disease, Ritter disease, Ritter von Ritterschein disease, Scalded skin syndrome, SSSS. ICD9: 695.81 ICD10: L00

Clinical

Staphylococcal scalded skin syndrome (SSSS) is characterized by diffuse erythematous cellulitis followed by extensive skin exfoliation. ^{1 2}

- Generalized erythema and then bulla formation with separation of the skin at the granular cell layer. ^{3 4}
- A warm, 'sandpaper' erythema with accentuation in the flexor creases may mimic scarlet fever; while the presence of flaccid bullae and Nikolsky sign may suggest pemphigus. ⁵
- Skin biopsy can be used to differential SSSS from Toxic epidermal necrolysis. ⁶
- Facial edema and perioral crusting are often present.

Dehydration may indicate fluid loss (as in thermal burns)

- Complete recovery occurs in most cases, within one to two weeks. ⁷
- The case-fatality rate in uncomplicated SSSS is less than 2%.
- Rare instances of recurrence have been reported ⁸
- Staphylococcal septicemia complicates SSSS in a minority of cases.

This disease is endemic or potentially endemic to all countries.

References

1. N Engl J Med 2006 Oct 26;355(17):1800-10.
2. Arch Dis Child 1998 Jan ;78(1):85-8.
3. Clin Microbiol Infect 2001 Jun ;7(6):301-7.
4. Semin Dermatol 1992 Mar ;11(1):11-8.
5. N Engl J Med 2006 Oct 26;355(17):1800-10.
6. J Dermatolog Treat 2005 ;16(5-6):278-86.
7. Pediatrics 1980 Aug ;66(2):285-90.
8. East Afr Med J 1997 Sep ;74(9):603-4.

Streptococcus suis infection

Agent	BACTERIUM. <i>Streptococcus suis</i> I and <i>Streptococcus suis</i> II A facultative gram-positive coccus
Reservoir	Pig
Vector	None
Vehicle	Air Secretions Meat Local wounds
Incubation Period	Unknown. Probably hours to few days
Diagnostic Tests	Culture of blood, tissue, body fluids
Typical Adult Therapy	Systemic antibiotic. Usually susceptible in vitro to Penicillin, Amoxicillin , Chloramphenicol and Gentamicin
Typical Pediatric Therapy	Systemic antibiotic
Clinical Hints	Severe multisystem disease, hemorrhagic diatheses, deafness or meningitis appearing hours to a few days after contact with pigs or pig products.
Synonyms	<i>Streptococcus suis</i> . ICD9: 027.8 ICD10: A48.8

Clinical

Demography:

Virtually all patients have been farmers and butchers, of whom 80 percent were men.

- Most had been involved in butchering sick pigs or selling the pork.
- Over 40 percent of the patients were in the age group 50 to 60 years, and none were children. ¹

Signs and symptoms:

- Clinical features of *Streptococcus suis* II infection include high fever, malaise, nausea and vomiting • followed by meningitis, subcutaneous hemorrhage, multi-organ failure (hepatic, renal, pulmonary, cardiac) and coma in severe cases. ^{2 3}
- Toxic shock syndrome is common. ⁴
- Sensorineural hearing loss is often present. ^{5 6}
- Peritonitis ⁷, endocarditis ⁸⁻¹⁰, mycotic aortic aneurysm ¹¹, rhabdomyolysis ¹², spondylodiscitis ¹³, salcroiliitis ¹⁴, monoarthritis ¹⁵⁻¹⁷, endophthalmitis ¹⁸ and cranial nerve palsy ¹⁹ have been reported.
- Persons with occupational exposure may exhibit asymptomatic seropositivity toward *S. suis*. ²⁰
- Relapses of meningitis may occur. ²¹

This disease is endemic or potentially endemic to 227 countries.

References

1. ProMED <promedmail.org> archive: 20050816.2399
2. Lancet Infect Dis 2007 Mar ;7(3):201-9.
3. ProMED <promedmail.org> archive: 20050804.2271
4. PLoS Med 2006 May ;3(5):e151.
5. J Med Assoc Thai 2008 May ;91(5):654-8.
6. Singapore Med J 2010 Feb ;51(2):e30-3.
7. J Med Assoc Thai 2000 Oct ;83(10):1274-7.
8. Eur J Clin Microbiol Infect Dis 1996 Sep ;15(9):765-6.
9. Kansenshogaku Zasshi 2009 Sep ;83(5):544-8.
10. Clin Med Insights Cardiol 2012 ;6:119-23.
11. Surg Infect (Larchmt) 2010 Apr ;11(2):179-81.
12. Clin Infect Dis 1997 Apr ;24(4):710-2.
13. Presse Med 1996 Oct 5;25(29):1348.
14. An Med Interna 1994 Jun ;11(6):309.
15. Ann Rheum Dis 1988 Jul ;47(7):598-9.
16. J Infect 1987 May ;14(3):237-41.
17. Korean J Lab Med 2011 Apr ;31(2):115-7.
18. Br J Ophthalmol 1978 Oct ;62(10):729-31.
19. J Med Assoc Thai 1999 Sep ;82(9):922-4.
20. ProMED <promedmail.org> archive: 20070823.2756
21. Singapore Med J 2010 Feb ;51(2):e30-3.

Strongyloidiasis

Agent	PARASITE - Nematoda. Phasmidea: <i>Strongyloides stercoralis</i> (<i>Strongyloides fulleborni</i> is occasionally implicated in systemic disease)
Reservoir	Human ? Dog Monkey (for <i>Strongyloides fulleborni</i>)
Vector	None
Vehicle	Skin contact Soil Feces Autoinfection Sexual contact (rare)
Incubation Period	14d - 30d
Diagnostic Tests	Identification of larvae (or ova, for <i>Strongyloides fulleborni</i>) in stool or duodenal aspirate. Serology.
Typical Adult Therapy	Ivermectin 200 micrograms/kg/d PO daily X 2d OR Thiabendazole 25 mg/kg BID (max 3g) X 2d OR Albendazole 400 mg/d X 3d (7 days for hyperinfection syndrome)
Typical Pediatric Therapy	Ivermectin 200 micrograms/kg/d PO daily X 2d OR Thiabendazole 25 mg/kg BID (max 3g) X 2d. OR Albendazole 200 mg/d X 3d (7 days for hyperinfection syndrome)
Clinical Hints	Diarrhea, gluteal or perineal pruritus and rash; eosinophilia often present; widespread dissemination encountered among immune-suppressed patients because of uncontrolled autoinfection (case-fatality rate for this complication = 80%).
Synonyms	Anguilluliasis, Anguillulosis, Cochin China gastroenteritis, Diploscapter, Halicephalobus, Halicephalobus, Larva currens, Leptodera intestinals, Leptodera stercoralis, Micronema, Pseudo-rhabdis stercoralis, Rhabditis stercoralis, Rhabdonema intestinale, Rhabdonema stercoralis, <i>Strongyloides fulleborni</i> , <i>Strongyloides stercoralis</i> , Strongyloidose, Threadworm, Turbatrix. ICD9: 127.2 ICD10: B78

Clinical

Strongyloidiasis may present as long as 75 years following initial acquisition of the parasite. ¹

Gastrointestinal strongyloidiasis:

The symptoms of strongyloidiasis reflect invasion of the skin, larval migration of larvae intestinal penetration.

- Approximately one third of patients are asymptomatic.
- Dermal and pulmonary symptoms resemble those of hookworm ², pruritic papular or linear urticarial rash (larva currens ³ ⁴) and a Loeffler-like syndrome.
- Intestinal penetration is characterized by abdominal pain, mucous diarrhea and eosinophilia. ⁵
- Vomiting, weight loss, protein-losing enteropathy and inappropriate ADH excretion ⁶ are occasionally encountered.
- Intestinal obstruction has been reported. ^{7 8}
- Yellowish mucosal nodules are seen on colonoscopy, predominantly in the ascending colon. ⁹
- Findings in colonic infection may mimic those of ulcerative colitis. ¹⁰

Generalized strongyloidiasis:

5 to 22% of patients develop a generalized or localized urticarial rash beginning in the anal region and extending to the buttocks, abdomen, and thighs.

- Extraintestinal infection may involve a wide variety of organs. ¹¹⁻¹⁶
- Autoinfection is characterized by massive larval invasion of the lungs and other organs.
- Massive systemic strongyloidiasis occurs in patients with lymphoma, leukemia and AIDS; and during high-dose therapy with corticosteroids. ¹⁷
- Findings include generalized abdominal pain, concurrent gram-negative bacillary septicemia (55% of cases) ¹⁸, bilateral diffuse pulmonary infiltrates and ileus.
- Hyperinfection may mimic acute exacerbation of COPD ¹⁹
- Eosinophilia may be present or absent at this stage; and rare instances of eosinophilic meningitis have been reported. ²⁰
- An outbreak of hyperinfection strongyloidiasis has been reported among immune-suppressed renal transplant recipients. ²¹
- *Strongyloides stercoralis* is the only helminth responsible for disseminated infection in immunocompromised patients. ²²

Strongyloides fulleborni infection is usually asymptomatic.

Strongyloides fulleborni kellyi infection ²³ is most common among infants, and consist of abdominal distention, mild

diarrhea and protein-losing enteropathy.

- Respiratory distress may occur, and is associated with a characteristic high-pitched cry.

This disease is endemic or potentially endemic to all countries.

Strongyloidiasis in Uganda

Prevalence surveys:

- 12.3% of otherwise healthy pregnant women at enrolment to a trial of deworming (Entebbe, 2007 publication) ²⁴
- 12% of pregnant women (Entebbe and Katabi, 2003 to 2005) ²⁵
- 5.4% of mothers and children in eastern Uganda (2010 publication) ²⁶
- 8% of HIV-positive patients with diarrhea in rural Uganda (2010 publication)

Seroprevalence surveys:

- 4% of mothers and 2% of their preschool children (Western Uganda, 2008 publication) ²⁷

Strongyloides fulleborni has been identified in primates. ^{28 29}

References

1. Emerg Infect Dis 2011 May ;17(5):931-2.
2. Semin Respir Infect 1997 Jun ;12(2):122-9.
3. Cutis 2008 May ;81(5):409-12.
4. Hautarzt 2011 May ;62(5):380-3.
5. J Clin Gastroenterol 2005 Mar ;39(3):203-11.
6. Southeast Asian J Trop Med Public Health 2007 Mar ;38(2):239-46.
7. Trop Gastroenterol 2005 Oct-Dec;26(4):201-2.
8. World J Emerg Surg 2010 ;5:23.
9. Digestion 2011 ;83(3):210-4.
10. Hum Pathol 2009 Apr ;40(4):572-7.
11. J Pediatr Adolesc Gynecol 2006 Oct ;19(5):329-32.
12. Trans R Soc Trop Med Hyg 2009 Jan ;103(1):106-7.
13. J Gastrointest Liver Dis 2009 Sep ;18(3):367-9.
14. J Intensive Care Med 2010 May-Jun;25(3):172-4.
15. Head Neck 2011 Mar 7;
16. Trop Biomed 2011 Apr ;28(1):64-7.
17. Clin Microbiol Rev 2004 Jan ;17(1):208-17.
18. Am J Clin Pathol 2007 Oct ;128(4):622-7.
19. J Chin Med Assoc 2009 Aug ;72(8):442-5.
20. Clin Microbiol Rev 2009 Apr ;22(2):322-48, Table of Contents.
21. Transplant Proc 2007 May ;39(4):1014-5.
22. Rev Prat 2007 Jan 31;57(2):167-73.
23. Parasitol Today 1992 Sep ;8(9):314-8.
24. Trans R Soc Trop Med Hyg 2007 Sep ;101(9):899-907.
25. J Infect Dis 2008 Sep 15;198(6):920-7.
26. J Helminthol 2011 Sep ;85(3):325-33.
27. J Helminthol 2008 Sep ;82(3):263-9.
28. J Parasitol 2005 Jun ;91(3):569-73.
29. PMID 18576800

Subdural empyema

Agent	BACTERIUM. Haemophilus influenzae , oral anaerobes, streptococci, et al
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Imaging techniques (CT scan, etc).
Typical Adult Therapy	Antimicrobial agent(s) directed at known or likely pathogen
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, severe headache, vomiting, and signs of meningeal irritation and increased cerebrospinal fluid pressure; may follow head trauma, meningitis, otitis or sinusitis; case-fatality rate 15% (alert) to 60% (comatose).
Synonyms	

Clinical

Most patients present with headache, meningismus, decreased mental status and hemiparesis. ¹

- 32 cases of suppurative parotitis in neonates were reported during 1970 to 2004. ²
- In 60 to 90% of cases, sinusitis or otitis is present.
- Extension of the infection into the subdural space is heralded by fever, focal and later generalized headache, vomiting, and meningismus. ³
- 50% of patients exhibit altered mental function.
- Focal neurological signs appear within 24 to 48 hours, and rapidly progress to hemispheric dysfunction with hemiparesis and hemisensory deficit.
- Seizures, usually focal, occur in 50% of cases, and papilledema in less than 50%.
- Signs of increased intracranial pressure appear, leading to cerebral herniation and death.
- Chronic and even sterile subdural collections are also encountered, often following antibiotic therapy.

This disease is endemic or potentially endemic to all countries.

References

1. *Curr Neurol Neurosci Rep* 2004 Nov ;4(6):448-56.
2. *Pediatr Int* 2012 Oct 8;
3. *Curr Neurol Neurosci Rep* 2004 Nov ;4(6):448-56.

Suppurative parotitis

Agent	BACTERIUM. Most commonly Staphylococcus aureus
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Unknown
Diagnostic Tests	Clinical features (local swelling and purulent discharge from salivary ducts). Stain and culture of discharge.
Typical Adult Therapy	Surgical drainage and aggressive parenteral antistaphylococcal therapy
Typical Pediatric Therapy	As for adult
Clinical Hints	Consider when confronted by unexplained fever in the setting of malnutrition, dehydration and obtundation; local swelling and discharge of pus from salivary duct are diagnostic.
Synonyms	Parotitis, bacterial. ICD9: 527.2 ICD10: K11.3

Clinical

Suppurative parotitis is characterized by the sudden onset of firm, erythematous swelling of the pre- and post auricular areas, extending to the angle of the mandible. ¹

- Marked pain and tenderness is accompanied by high fever, chills and marked toxicity.
- Pus may be seen exiting from the parotid duct.
- Progression of the disease can result in massive swelling of the neck, respiratory obstruction, septicemia, fistula formation ² and osteomyelitis of the adjacent facial bones.
- The condition should be suspected in any patient with unexplained or prolonged fever.

This disease is endemic or potentially endemic to all countries.

References

1. [J Craniofac Surg 2003 Jan ;14\(1\):37-40.](#)
2. [J Med Case Rep 2010 ;4:249.](#)

Syphilis

Agent	BACTERIUM. <i>Treponema pallidum</i> subsp. pallidum A microaerophilic gram-negative spirochete
Reservoir	Human
Vector	None
Vehicle	Sexual contact Infected secretions
Incubation Period	2w - 4w (range 10d - >8w)
Diagnostic Tests	Dark field microscopy (chancre). VDRL confirmed by antitreponemal test (FTA, MHTP). Nucleic acid amplification.
Typical Adult Therapy	Primary, secondary or early (< 1 year) latent: Benzathine Penicillin G 2.4 million units IM Other stages: Repeat dosage at one and two weeks Alternatives: Tetracycline , Ceftriaxone
Typical Pediatric Therapy	Primary, secondary or early (< 1 year) latent: Benzathine Penicillin G : Weight <14 kg: 600,000u IM Weight 14 to 28 kg: 1,200,000u IM Other stages: Repeat dosage at one and two weeks
Clinical Hints	Firm, painless chancre (primary syphilis); later fever, papulosquamous rash and multisystem infection (secondary syphilis); late lesions of brain, aorta, bone or other organs (tertiary syphilis).
Synonyms	Canton rash, Chinese ulcer, Christian disease, French disease, German sickness, Harde sjanker, Lues, Neopolitan itch, Polish sickness, Sifilide, Sifilis, Spanish pockes, Syphilis, <i>Treponema pallidum</i> . ICD9: 090,091,092,093,094,095,096,097 ICD10: A50,A51,A52,A53

Clinical

WHO Case definition for surveillance:

The signs and symptoms of syphilis are multiple.

- The primary stage usually, but not necessarily, involves ulceration of the external genital organs and local lymphadenopathy; secondary and tertiary syphilis show mainly dermatological and systemic manifestations. For surveillance purposes, only confirmed cases will be considered.

Confirmed case

- A person with a confirmed positive serology for syphilis (Rapid Plasma Reagin (RPR) or VDRL confirmed by TPHA (*Treponema pallidum* hemagglutination antibodies) or FTA (fluorescent treponemal antibody absorption)).

Case classification

- Congenital syphilis: An infant with a positive serology, whether or not the mother had a positive serology during pregnancy.
- Acquired syphilis: All others.

Additional notes:

- The prevalence rate among pregnant women in developing countries varies between 3% and 19%. Maternal syphilis is associated with congenital syphilis (one third of births from such pregnancies), and with spontaneous abortion and stillbirth.
- Because the primary lesion is often painless and secondary syphilis is usually not diagnosed, women are mainly identified through serological screening.

Syphilis is a chronic disease with a waxing and waning course; and is reported from all countries.

- Transmission is mainly by sexual contact.
- Primary, secondary, and early latent syphilis are potentially infectious.

Stages of syphilis:

- Primary syphilis is characterized by a painless chancre at the site of inoculation. ¹ Penile swelling without an overt chancre has also been reported. ²
- The secondary stage is characterized by a generalized (rarely localized ³ non-pruritic polymorphic ⁴⁻⁶ or papulonecrotic ⁷ rash, lymphadenopathy, and systemic manifestations. ⁸⁻¹³ Moist flat genital or mucosal lesions (condyloma lata) ¹⁴ or granulomatous dermatitis ¹⁵ may be evident.
- An asymptomatic latent period follows, which for epidemiological purposes is divided into early (<1 year) and late (>1 year) stages.
- The tertiary stage is the most destructive and is marked by cardiovascular ¹⁶ and neurological sequelae ¹⁷⁻²², and gummatous involvement of any organ system. ²³⁻²⁶
- As of 2009, the world's literature contained 165 reports of cerebral syphilitic gummata • 64% in men and 66% located on the cerebral convexities. ²⁷

- Syphilitic uveitis may present in the absence of other clinical manifestations of syphilis.²⁸ Acute posterior placoid chorioretinitis is also encountered.^{29 30} 143 cases of syphilitic uveitis were reported in the English Language literature during 1984 to 2008.³¹

The clinical features of congenital infection are similar to those of secondary syphilis, and may be associated with deformation of teeth, bones and other structures.³²

Acquired syphilis in patients with HIV infection is characterized by severe and accelerated infection, often with overt meningitis, hepatitis³³, lues maligna (a florid papulopustular rash)³⁴ and other forms of systemic involvement.³⁵⁻⁴⁴

- The presence of concurrent syphilis does not affect the progression of AIDS.⁴⁵

This disease is endemic or potentially endemic to all countries.

Syphilis in Uganda

Prevalence surveys:

5% of patients with genital ulcer disease (rural Rakai District, 2008 publication)⁴⁶

Seroprevalence surveys:

- 6.1% of CSW (VDRL, 1988 publication)⁴⁷
- 24.5% (HIV-negative) to 64.0% (HIV-positive) persons in Kitgum (1986)⁴⁸
- 2.3% of men and 1.4% of women in Rakai (2003 publication)⁴⁹
- 4.3% of persons in fishing communities on Lake Victoria (2005)⁵⁰
- 12.9% of men and 12.6% of women ages 15 to 54 in the rural southwest (1999 publication)⁵¹
- 4.0% of female and 2.8% of male adolescents in Kampala (2006 publication)⁵²
- 4.0% of pregnant women and 6.2% of their male partners (Entebbe, 2002 to 2006)⁵³
- 1.1% of pregnant women in Entebbe (2004 to 2005)⁵⁴
- 5.3% of intervention-program pregnant women (2012 publication)⁵⁵
- 0.2% of blood donors in central Uganda (1999)⁵⁶
- 64.3% of HIV-positive patients (2012 publication)⁵⁷

References

1. Clin Dermatol 2004 Nov-Dec;22(6):461-8.
2. Int J STD AIDS 2008 Sep ;19(9):640-1.
3. J Coll Physicians Surg Pak 2008 May ;18(5):303-4.
4. Clin Microbiol Rev 2005 Jan ;18(1):205-16.
5. J Cutan Med Surg 2008 May-Jun;12(3):114-6.
6. Int J Dermatol 2010 Nov ;49(11):1321-4.
7. South Med J 2007 Dec ;100(12):1221-2.
8. Australas J Dermatol 2013 Feb ;54(1):e19-21.
9. J Clin Microbiol 2011 Dec ;49(12):4394-6.
10. J Korean Med Sci 2010 Nov ;25(11):1661-4.
11. Int J STD AIDS 2010 Mar ;21(3):215-6.
12. Clin Nephrol 2008 Dec ;70(6):532-6.
13. Immune Netw 2012 Dec ;12(6):261-8.
14. Int J Dermatol 2008 Jan ;47(1):56-8.
15. Ned Tijdschr Geneesk 2012 ;155(33):A5130.
16. Am J Cardiol 2009 Dec 1;104(11):1578-87.
17. J Neurol Neurosurg Psychiatry 2004 Dec ;75(12):1727-30.
18. Curr Infect Dis Rep 2005 Jul ;7(4):277-284.
19. Curr Treat Options Neurol 2006 May ;8(3):185-92.
20. J Emerg Med 2011 Dec ;41(6):613-5.
21. Int J Stroke 2011 Apr ;6(2):136-43.
22. BMJ Case Rep 2013 ;2013
23. World J Hepatol 2010 Sep 27;2(9):362-6.
24. Int J STD AIDS 2011 Sep ;22(9):531-3.
25. Neurol Sci 2012 Oct ;33(5):1179-81.
26. Clin Nucl Med 2012 Sep ;37(9):e231-3.
27. Neurosurgery 2009 Mar ;64(3):568-75; discussion 575-6.
28. J Chin Med Assoc 2007 Jul ;70(7):274-80.
29. Clin Ophthalmol 2008 Sep ;2(3):669-73.
30. Retina 2012 Oct ;32(9):1915-41.
31. Clin Experiment Ophthalmol 2010 Jan ;38(1):68-74.
32. Infect Dis Obstet Gynecol 2012 ;2012:430585.
33. Int J STD AIDS 2012 Aug ;23(8):e4-6.
34. Sex Transm Dis 2009 Aug ;36(8):512-4.
35. AIDS Rev 2008 Apr-Jun;10(2):85-92.
36. Mayo Clin Proc 2007 Sep ;82(9):1091-102.
37. MMWR Morb Mortal Wkly Rep 2007 Jun 29;56(25):625-8.
38. Clin Infect Dis 2007 May 1;44(9):1222-8.
39. Dermatol Clin 2006 Oct ;24(4):497-507, vi.
40. Int J STD AIDS 2009 Apr ;20(4):278-84.
41. Eur J Intern Med 2009 Jan ;20(1):9-13.
42. J Clin Microbiol 2010 Jul ;48(7):2640-2.
43. Clin Rheumatol 2011 May ;30(5):729-33.
44. Int J STD AIDS 2012 Aug ;23(8):599-600.
45. Int J STD AIDS 2010 Jan ;21(1):57-9.
46. Sex Transm Infect 2009 Apr ;85(2):97-101.
47. Bull World Health Organ 1990 ;68(5):639-54.
48. Trans R Soc Trop Med Hyg 1990 Jan-Feb;84(1):144-7.
49. Sex Transm Infect 2003 Apr ;79(2):98-105.
50. Sex Transm Infect 2011 Oct ;87(6):511-5.
51. Sex Transm Infect 1999 Apr ;75(2):98-102.
52. J Adolesc Health 2006 Mar ;38(3):213-21.
53. Trop Med Int Health 2008 May ;13(5):680-2.
54. PLoS Negl Trop Dis 2009 ;3(6):e473.
55. J Acquir Immune Defic Syndr 2012 Nov 1;61(3):e40-e46.
56. Trop Med Int Health 2006 Jun ;11(6):951-4.
57. J Infect Dev Ctries 2012 ;6(3):276-82.

Taeniasis

Agent	PARASITE - Platyhelminthes, Cestoda. Cyclophyllidea, Taeniidae: <i>Taenia solium</i> & <i>T. saginata</i> (other species occasionally encountered)
Reservoir	Cattle Pig
Vector	None
Vehicle	Meat
Incubation Period	6w - 14w
Diagnostic Tests	Identification of ova or proglottids in feces.
Typical Adult Therapy	Praziquantel 10 mg/kg PO as single dose OR Niclosamide 2 g PO once
Typical Pediatric Therapy	Praziquantel 10 mg/kg PO as single dose OR Niclosamide 50 mg/kg PO once
Clinical Hints	Vomiting and weight loss; often symptomatic or first appreciated due to passage of proglottids or 'tape' segments; parasite may survive for over 25 years in the human intestine.
Synonyms	Bandwurm [Taenia], Drepanidotaenia, Gordiid worm, Hair snake, Mesocestoides, Raillietina, <i>Taenia longihamatus</i> , <i>Taenia saginata</i> , <i>Taenia solium</i> , <i>Taenia taeniaformis</i> , Taeniarhynchiasis, Tapeworm (pork or beef), Tenia. ICD9: 123.0,123.2 ICD10: B68

Clinical

Most cases of *Taenia* infestation are subclinical.

Symptomatic taeniasis may be associated with nausea, vomiting, epigastric fullness, weight loss or diarrhea. ¹

- *Taenia saginata* often becomes apparent when motile proglottids are passed through the anus; however, this is uncommon with *T. solium* infestations.
- Eosinophilia is not a prominent finding.
- Rare complications include appendicitis, cholangitis ², cholecystitis ³, pancreatitis or intestinal obstruction. ⁴
- The major complication of *T. solium* infection, Cysticercosis, is discussed separately in this module.

This disease is endemic or potentially endemic to all countries.

Taeniasis in Uganda

Prevalence surveys:

- 0.12% to 1.2% of pigs in Kampala slaughterhouses are infested (1998 to 1999) ⁵
- 33.7% of pigs in rural Lara (northern region)

References

1. Parasitol Res 2003 Nov ;91(5):412-38.
2. Acta Clin Belg 2012 Nov-Dec;67(6):436-7.
3. Case Rep Surg 2012 ;2012:572484.
4. Trans R Soc Trop Med Hyg 2007 May ;101(5):527-8.
5. Acta Trop 2003 Jun ;87(1):13-23.

Tetanus

Agent	BACTERIUM. <i>Clostridium tetani</i> An anaerobic gram-positive bacillus
Reservoir	Animal feces Soil
Vector	None
Vehicle	Injury
Incubation Period	6d - 8d (range 1d - 90d)
Diagnostic Tests	Isolation of <i>C. tetani</i> from wound is rarely helpful. Serology (specimen taken before administration of antitoxin).
Typical Adult Therapy	Human antitoxin (see Vaccine module). Metronidazole (2 g daily) or Penicillin G (24 million u daily) or Doxycycline (200 mg daily). Diazepam (30 to 240 mg daily). Tracheostomy, hyperalimentation
Typical Pediatric Therapy	Human antitoxin (see Vaccine module). Metronidazole (30 mg/kg daily); or Penicillin G (300,000 units/kilo daily). Diazepam. Tracheostomy, hyperalimentation
Vaccines	DT DTaP DTP Td Tetanus immune globulin Tetanus
Clinical Hints	Trismus, facial spasm, opisthotonus, tachycardia and recurrent tonic spasms of skeletal muscle; sensorium is clear; disease may persist for 4 to 6 weeks; case fatality rate = 10% to 40%.
Synonyms	Lockjaw, Starrkrampf, Stelkrampf, Tetano, Tetanos. ICD9: 037,771.3 ICD10: A33,A34,A35

Clinical

Tetanus may present in any of four clinical forms: generalized, localized, cephalic, and neonatal. ¹

- In general, shorter incubation periods are associated with a worse prognosis.
- Certain portals of entry (compound fractures) and underlying conditions (heroin addiction) are also associated with poorer prognoses.
- A series of 11 cases of tetanus related to tungiasis (25% of all tetanus cases) was reported by a single hospital in Brazzaville over an 11-month period (1989 publication). ²
- An outbreak of 12 cases of tetanus in Argentina was reported among elderly women treated with sheep cell therapy (1996). ³
- In one case, tetanus was associated with chronic otitis media. ⁴
- Tetanus has been reported in a child who bit her own tongue during a convulsion ⁵ and following a snake bite (2007 publication) ⁶
- An attack of tetanus does not result in immunity. Therefore, recurrent tetanus is possible, unless the patient is given a series of toxoid following recovery. ⁷⁻¹⁴

Generalized tetanus, the most common form, begins with trismus ("lockjaw") and risus sardonicus (increased tone in the orbicularis oris).

- Abdominal wall rigidity may be present.
- The generalized spasm consists of opisthotonic posturing with flexion of the arms and extension of the legs. ¹⁵
- The patient does not lose consciousness, and experiences severe pain during these spasms.
- Spasms often are triggered by sensory stimuli.
- Respiration may be compromised by upper airway obstruction, or by participation of the diaphragm in the general muscular contraction.
- Autonomic dysfunction, usually occurring after several days of symptoms, is currently the leading cause of death in tetanus.
- Complications of tetanus include rhabdomyolysis and renal failure ^{16 17}
- The illness can progress for two weeks, while the severity of illness may be decreased by partial immunity.
- Recovery takes an additional month, but is complete unless complications supervene.
- Lower motor neuron dysfunction may appear after the spasms remit, and persist for several additional weeks.
- A case of *Clostridium tetani* bacteremia has been reported. ¹⁸
- Case-fatality rates of 10% to 50% are reported, but may be as high as 70% in Africa. ¹⁹

- The differential diagnosis of tetanus includes strychnine poisoning and neuromyotonia (Isaac's syndrome). ²⁰

Localized tetanus presents as rigidity of the muscles associated with the site of inoculation.

- Initial symptomatology may be limited to back pain ²¹
- The illness may be mild and persistent, and tends to resolve spontaneously.
- Weakness and diminished muscle tone are often present in the most involved muscle.
- Localized tetanus is often a prodrome of generalized tetanus.

Cephalic tetanus is a form of localized disease affecting the cranial nerve musculature

- Facial nerve weakness, is often apparent, and extraocular muscle involvement is occasionally noted.

Neonatal tetanus follows infection of the umbilical stump, most commonly as a result of a failure of aseptic technique following delivery of non-immune mothers.

- The condition usually manifests with generalized weakness and failure to nurse; followed by rigidity and spasms.
- The mortality rate exceeds 90%, and psychomotor retardation is common among survivors.
- Poor prognostic factors include age younger than 10 days, symptoms present for fewer than 5 days before presentation to hospital, fever, and the presence of risus sardonicus or fever.
- Apnea is the leading cause of death in the first week of disease, and sepsis in the second week.
- Bacterial infection of the umbilical stump leads to sepsis in almost half of babies with neonatal tetanus.

The WHO Case definition for surveillance of neonatal tetanus is as follows:

- Suspected case: Any neonatal death between 3-28 days of age in which the cause of death is unknown; or any neonate reported as having suffered from neonatal tetanus between 3-28 days of age and not investigated.
- Confirmed case: Any neonate with a normal ability to suck and cry during the first two days of life, and who between 3 and 28 days of age cannot suck normally, and becomes stiff or has convulsions (i.e. jerking of the muscles) or both.
- Hospital-reported cases of neonatal tetanus are considered confirmed.
- The diagnosis is purely clinical and does not depend upon laboratory or bacteriological confirmation.

This disease is endemic or potentially endemic to all countries.

Tetanus in Uganda

Vaccine Schedule:

BCG - birth

DTwP - 6, 10, 14 weeks


DTwPHibHep - 6, 10, 14 weeks

Measles (monovalent) - 9 months


OPV - birth; 6, 10, 14 weeks

TT - Women of child-bearing age; +1, +6 months; +1, +1 year


Vitamin A - 6, 12, 18, 24, 30, 36, 42 months




Graph: Uganda. Tetanus - WHO-UNICEF est. vaccine (DTP3 %) coverage



Graph: Uganda. Tetanus - WHO-UNICEF est. vaccine (TT2+ %) coverage




Graph: Uganda. Tetanus, cases



Graph: Uganda. Tetanus - neonatal, cases

Notes:

1. Neonatal tetanus in Uganda was certified by W.H.O. as "eliminated" as of 2011. ²²



Graph: Uganda. Tetanus - neonatal, estimated deaths

Notes:

1. 1992 - The highest rate in the world (13.0 per 1,000 live births).

References

1. Neurol India 2002 Dec ;50(4):398-407.
2. Dakar Med 1989 ;34(1-4):44-8.
3. Clin Infect Dis 1997 Apr ;24(4):738.
4. BMJ Case Rep 2012 ;2012
5. Trop Doct 2012 Jul ;42(3):180-1.
6. Bull Soc Pathol Exot 2007 Aug ;100(3):184-5.
7. Lancet 2004 Jun 19;363(9426):2048.
8. Trop Doct 2011 Apr ;41(2):127-8.
9. Rev Inst Med Trop Sao Paulo 1982 Mar-Apr;24(2):107-15.
10. South Med J 1971 Jul ;64(7):859 passim.
11. Br Med J 1968 Jun 29;2(5608):829.
12. Presse Med 1967 May 20;75(24):1223-6.
13. Calif Med 1962 Jul ;97:31-3.
14. Ann Intern Med 1954 Jul ;41(1):159-63.
15. J Neurol Neurosurg Psychiatry 2000 Sep ;69(3):292-301.
16. Clin Nephrol 2010 Jan ;73(1):64-7.
17. Arch Pediatr 2012 May 29;
18. J Med Microbiol 2012 Sep 13;
19. Trop Doct 2009 Jan ;39(1):39-40.
20. J Clin Neurosci 2010 Jun ;17(6):814-5.
21. Emerg Med J 2007 Jan ;24(1):e5.
22. Wkly Epidemiol Rec 2011 Dec 9;86(50):565-75.

Thelaziasis

Agent	PARASITE - Nematoda. Phasmidea: <i>Thelazia callipaeda</i> [rarely <i>T. californiensis</i>]
Reservoir	Dog Rabbit Deer Cat
Vector	Fly (? <i>Musca</i> and <i>Fannia</i> species)
Vehicle	None
Incubation Period	not known
Diagnostic Tests	Identification of parasite.
Typical Adult Therapy	Extraction of parasite
Typical Pediatric Therapy	As for adult
Clinical Hints	Conjunctivitis and lacrimation associated with the sensation of an ocular foreign body.
Synonyms	Conjunctival spirurosis, Oriental eye worm, Rictularia, <i>Thelazia californiensis</i> , <i>Thelazia callipaeda</i> . ICD9: 372.15 ICD10: B83.8

Clinical

The signs and symptoms of Thelaziasis are related to the presence of a worm in the conjunctival sac, and consist of pain, lacrimation and a foreign body sensation. ^{1 2}

This disease is endemic or potentially endemic to all countries.

References

1. Trends Parasitol 2005 Jan ;21(1):1-4.
2. J Parasitol 2006 Aug ;92(4):872-5.

Thogoto

Agent	VIRUS - RNA. Orthomyxoviridae, Thogotovirus: Thogoto virus
Reservoir	Sheep Bird
Vector	Tick
Vehicle	None
Incubation Period	4d - 5d
Diagnostic Tests	Viral culture (blood, CSF, brain tissue). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Encephalitis or optic neuritis which follows a tick bite or contact with livestock; occasional reports of hepatitis and fatal infection.
Synonyms	Dhori. ICD9: 078.89 ICD10: A85.2

Clinical

Reported clinical features in humans have included encephalitis, hepatitis, optic neuritis and lymphadenopathy. ¹

This disease is endemic or potentially endemic to 12 countries.

Thogoto in Uganda

Thogoto virus has been isolated from a Ugandan banded mongoose (*Mungos mungo*) - the first instance of isolation from a wild vertebrate. ²

References

1. Arch Virol Suppl 1996 ;11:33-40.
2. Am J Trop Med Hyg 1999 Mar ;60(3):439-40.

Toxic shock syndrome

Agent	BACTERIUM. <i>Staphylococcus aureus</i> , <i>Streptococcus pyogenes</i> , et al - (toxins) Facultative gram-positive cocci
Reservoir	Human
Vector	None
Vehicle	Tampon (occasionally bandage, etc) which induces toxinosis
Incubation Period	Unknown
Diagnostic Tests	Isolation of toxigenic <i>Staphylococcus aureus</i> . Toxin assay available in specialized laboratories.
Typical Adult Therapy	The role of topical (eg, vaginal) and systemic antistaphylococcal antibiotics is unclear
Typical Pediatric Therapy	The role of topical (eg, vaginal) and systemic antistaphylococcal antibiotics is unclear
Clinical Hints	Fever (>38.9), hypotension (<90 mm Hg) and dermal erythema with desquamation; respiratory, cardiac or other disease present; most cases associated with 'super absorbent' tampon use or staphylococcal wound infection; case-fatality rate = 5% to 10%.
Synonyms	Streptococcal toxic shock syndrome, TSS. ICD9: 040.82 ICD10: A48.3

Clinical

CDC (The United States Centers for Disease Control) case definition for surveillance:

For surveillance purposes, the CDC (The United States Centers for Disease Control) case definition of toxic shock syndrome ¹ requires an illness with the following clinical manifestations:

1. fever at least 38.9 C
2. diffuse macular erythema ²
3. desquamation 1 to 2 weeks after onset of illness (particularly of the palms and soles)
4. hypotension (less than 90 mm Hg for adults, or less than fifth percentile if below age 16 years • or orthostatic hypotension)
5. multisystem involvement, consisting of three or more of the following: acute vomiting or diarrhea; myalgia and elevation of creatine phosphokinase levels; vaginal, oropharyngeal or conjunctival hyperemia; elevation of blood urea nitrogen or creatine to at least twice normal, or sterile pyuria; elevation of serum bilirubin or aminotransferase levels to at least twice normal; platelet count < 100,000/ cu mm; disorientation or alteration in consciousness unrelated to fever and hypotension
6. laboratory examination
 - negative cultures of blood, throat or cerebrospinal fluid (however, *S. aureus* may be present in blood)
 - negative tests for measles, leptospirosis or rickettsiosis

A probable case requires at least five of the above clinical findings. A confirmed case requires all six clinical findings (unless the patient dies before desquamation can occur).

The case definition for Streptococcal toxic shock syndrome ^{3 4} includes isolation of *Streptococcus pyogenes* in addition to:

1. hypotension as above
2. multiorgan involvement characterized by at least two of the following (defined above)
 - renal impairment
 - coagulopathy
 - hepatic dysfunction
 - acute respiratory distress syndrome
 - a generalized erythematous macular rash which may desquamate ⁵
 - soft tissue necrosis (fasciitis, myositis, gangrene).

This disease is endemic or potentially endemic to all countries.

References

1. Infect Dis Clin North Am 1996 Dec ;10(4):727-46.
2. J Am Acad Dermatol 1998 Sep ;39(3):383-98; quiz 399-400.
3. J Emerg Med 2002 May ;22(4):357-66.
4. Emerg Infect Dis 1995 Jul-Sep;1(3):69-78.
5. J Am Acad Dermatol 1998 Sep ;39(3):383-98; quiz 399-400.

Toxocariasis

Agent	PARASITE - Nematoda. Phasmidea: <i>Toxocara cati</i> and <i>canis</i>
Reservoir	Cat Dog Mouse
Vector	None
Vehicle	Soil ingestion
Incubation Period	1w - 2y
Diagnostic Tests	Identification of larvae in tissue. Serology.
Typical Adult Therapy	Albendazole 400 mg BID X 5d. OR Mebendazole 100 to 200 mg PO bid X 5 days Add corticosteroids if eye, brain, heart or lung involvement is present.
Typical Pediatric Therapy	As for adult
Clinical Hints	Cough, myalgia, seizures, urticaria, hepatomegaly, pulmonary infiltrates or retrobulbar lesion; marked eosinophilia often present; symptoms resolve after several weeks, but eosinophilia may persist for years.
Synonyms	<i>Ascaris suum</i> , <i>Toxocara canis</i> , <i>Toxocara cati</i> , Toxocarose, Visceral larva migrans. ICD9: 128.0 ICD10: B83.0

Clinical

Most infections present in children below the age of 5 years, and are asymptomatic or mild; however rare instances of infection are reported in adults. ¹

Overt disease is characterized by fever, cough, wheezing, eosinophilia, myalgia, tender hepatomegaly and abdominal pain. ²

- A tender nodular rash may be present on the trunk and legs.
- Chronic urticaria, chronic pruritus, relapsing eosinophilic cellulitis ³ and eczema are also reported. ⁴
- Myocarditis ^{5 6}, pericarditis ⁷, pulmonary infiltrates, acute respiratory distress syndrome ⁸, seizures, nephritis, encephalopathy ⁹, spinal involvement (usually cervical or thoracic) including transverse myelitis ^{10 11}, encephalomyelitis ¹², cerebral vasculitis ¹³, Bell's palsy ¹⁴, eosinophilic meningitis ¹⁵⁻¹⁷, eosinophilic pneumonia ¹⁸ or pleural effusion ^{19 20}, eosinophilic ascites ^{21 22}, eosinophilic abscesses of the liver ^{23 24}, cystitis ²⁵ and renal dysfunction have been described in heavy infections.
- Eye disease is rare in toxocariasis. ²⁶ Ocular toxocariasis usually presents in children ages 5 to 10 years, and is characterized by formation of a retinal granuloma at or near the macula, resulting in strabismus, iridocyclitis, glaucoma, papillitis or visual loss. ²⁷⁻³²
- Toxocariasis has been identified as a cause of chronic cough in childhood ³³ and of diminished lung function (FEV-1) at any age. ³⁴
- In some cases, pulmonary and hepatic nodular lesions could be mistaken for malignancy. ^{35 36}
- Toxocariasis has been implicated in the etiology of epilepsy ³⁷ and decreased cognitive function among children. ³⁸

Ascaris suum, a parasite of pigs, has been reported to cause rare cases of myelitis ³⁹, encephalopathy ⁴⁰, eosinophilic pneumonia ⁴¹⁻⁴³ and focal liver lesions in humans. ⁴⁴⁻⁴⁸

- *A. suum* has been implicated in cases of eosinophilic colitis ⁴⁹ and intestinal obstruction. ⁵⁰

This disease is endemic or potentially endemic to all countries.

References

1. *Vojnosanit Pregl* 2011 Oct ;68(10):881-5.
2. *Clin Microbiol Rev* 2003 Apr ;16(2):265-72.
3. *Cases J* 2008 ;1(1):356.
4. *J Am Acad Dermatol* 2008 Dec ;59(6):1031-42.
5. *Circ J* 2009 Jul ;73(7):1344-8.
6. *J Cardiovasc Ultrasound* 2012 Sep ;20(3):150-3.
7. *Acta Med Port* 1997 Feb-Mar;10(2-3):157-60.
8. *Rev Mal Respir* 2010 May ;27(5):505-8.
9. *Am J Trop Med Hyg* 2003 Sep ;69(3):341-3.
10. *Eur J Radiol* 2010 Aug ;75(2):236-40.
11. *Medicine (Baltimore)* 2011 Sep ;90(5):337-43.
12. *Case Report Med* 2011 ;2011:503913.
13. *Rev Neurol (Paris)* 2012 May 22;
14. *Rev Chilena Infectol* 2012 Oct ;29(5):551-3.
15. *Clin Neurol Neurosurg* 2005 Aug ;107(5):432-8.
16. *Pediatr Infect Dis J* 2008 Sep ;27(9):849-50.

17. Clin Microbiol Rev 2009 Apr ;22(2):322-48, Table of Contents.
18. Turkiye Parazitol Derg 2012 ;36(4):258-259.
19. Am J Trop Med Hyg 2004 Dec ;71(6):764.
20. Kansenshogaku Zasshi 2006 Nov ;80(6):716-20.
21. Pediatr Infect Dis J 2008 Jun ;27(6):563-4.
22. Pol Merkur Lekarski 2008 Jun ;24(144):533-5.
23. Bratisl Lek Listy 2011 ;112(11):644-7.
24. Abdom Imaging 2012 Jul 17;
25. Iran J Parasitol 2011 Mar ;6(1):92-4.
26. Rev Chilena Infectol 2011 Oct ;28(5):431-4.
27. Ophthalmol Clin North Am 2002 Sep ;15(3):351-6.
28. Klin Oczna 2008 ;110(10-12):364-6.
29. J Pediatr Ophthalmol Strabismus 2011 Jul-Aug;48(4):204-12.
30. Eur J Ophthalmol 2011 Jul-Aug;21(4):490-4.
31. Jpn J Ophthalmol 2011 Mar ;55(2):143-7.
32. J Pediatr Ophthalmol Strabismus 2012 Aug 28;:1-11.
33. J Helminthol 2008 Dec ;82(4):357-63.
34. Int J Parasitol 2011 Feb ;41(2):243-7.
35. Dig Dis Sci 2012 Jan ;57(1):155-60.
36. Clin Gastroenterol Hepatol 2013 Feb 5;
37. PLoS Negl Trop Dis 2012 Aug ;6(8):e1775.
38. Int J Parasitol 2012 Nov 1;
39. Rinsho Shinkeigaku 2004 Mar ;44(3):198-202.
40. J Neurol Sci 1999 Apr 1;164(2):195-9.
41. Kansenshogaku Zasshi 2004 Dec ;78(12):1036-40.
42. Nihon Kokyuki Gakkai Zasshi 2001 Sep ;39(9):716-20.
43. Nihon Kokyuki Gakkai Zasshi 1998 Feb ;36(2):208-12.
44. Intern Med 2002 Jul ;41(7):574-9.
45. Abdom Imaging 2004 Sep-Oct;29(5):598-602.
46. Rinsho Shinkeigaku 2004 Mar ;44(3):198-202.
47. Nihon Kokyuki Gakkai Zasshi 1998 Feb ;36(2):208-12.
48. J Helminthol 2012 Mar 19;:1-8.
49. J Gastroenterol Hepatol 1997 Mar ;12(3):204-6.
50. Trans R Soc Trop Med Hyg 1978 ;72(1):107.

Toxoplasmosis

Agent	PARASITE - Protozoa. Sporozoa, Coccidea, Eimeriida: <i>Toxoplasma gondii</i>
Reservoir	Rodent Pig Cattle Sheep Chicken Bird Cat Marsupial (kangaroo)
Vector	None
Vehicle	Transplacental Meat ingestion Soil ingestion Water or milk (rare) Fly
Incubation Period	1w - 3w (range 5d - 21d)
Diagnostic Tests	Serology. Cultivation or identification of organisms per specialized laboratories. Nucleic acid amplification.
Typical Adult Therapy	Pyrimethamine 25 mg/d + Sulfonamides 100 mg/kg (max 6g)/d X 4w - give with folinic acid. Alternatives: Clindamycin , Azithromycin , Dapsone . Spiramycin (in pregnancy) 4g/d X 4w
Typical Pediatric Therapy	Pyrimethamine 2 mg/kg/d X 3d, then 1 mg/kg/d + Sulfonamides 100 mg/kg/d X 4w - give with folinic acid. Alternatives: Clindamycin , Azithromycin , Dapsone .
Clinical Hints	Fever, lymphadenopathy and hepatic dysfunction; chorioretinitis; cerebral cysts (patients with AIDS); congenital hydrocephalus, mental retardation or blindness.
Synonyms	Toxoplasma, Toxoplasmosis, Toxoplasmosi. ICD9: 130 ICD10: B58

Clinical

Acquired toxoplasmosis:

The clinical features of acquired toxoplasmosis can range from subclinical infection to lymphadenopathy (the most common presentation) to fatal, fulminant disease.

- In healthy adults, infection is usually subclinical, or mimics infectious mononucleosis; however, pharyngitis, posterior and posterior cervical lymphadenopathy are unusual in toxoplasmosis.
- Most patients with acute *Toxoplasma* lymphadenitis experience fatigue, headache, difficulty concentrating and muscle aches. ¹
- In immunocompromised hosts, toxoplasmosis may mimic other opportunistic infections, such as tuberculosis or infection with *P. jiroveci* (formerly *P. carinii*) ², or extensive varicella. ³
- In patients with AIDS, CNS involvement is the most common manifestation, followed by pulmonary disease. ⁴

Congenital toxoplasmosis:

The rate and severity of congenital toxoplasmosis are largely related to gestational age at the time of infection. ^{5 6}

- Overt clinical signs appear to be more common among American infants vs. European infants with congenital toxoplasmosis. ^{7 8}
- The brain and eyes are often affected, presenting as chorioretinitis, hydrocephalus, intracranial calcifications, and seizures. ⁹
- 97% of children infected during the first trimester of pregnancy and having normal antenatal ultrasounds are asymptomatic or only slightly affected. ¹⁰
- Rare instances of nephrotic syndrome complicating congenital toxoplasmosis have been reported. ¹¹

Ocular toxoplasmosis: ¹²

Ocular toxoplasmosis occurs from reactivation of cysts in the retina.

- Focal necrotizing retinitis is characteristic lesion, and approximately 35% of all cases of retinochoroiditis can be attributed to toxoplasmosis. ¹³
- Risk factors for early (first two years of life) retinochoroiditis include a delay of >8 weeks between maternal seroconversion and the beginning of treatment, female gender, and the presence of cerebral calcifications. ¹⁴
- The incidence and severity of ocular toxoplasmosis varies from country to country. ¹⁵

CNS toxoplasmosis:

The manifestations of CNS toxoplasmosis in the immunocompromised patient range from an insidious process evolving over several weeks to acute onset of a confusional state.

- Signs may be focal or symmetrical.
- *T. gondii* has a predilection to localize in the basal ganglia and brain stem, producing extrapyramidal symptoms resembling those of Parkinson's disease.

- A normal CT scan does not rule out cerebral toxoplasmosis. MRI is the imaging modality of choice ¹⁶
- Nonfocal evidence of neurological dysfunction may include generalized weakness, headache, confusion, lethargy, alteration of mental status, personality changes, and coma.
- Infection in transplant recipients is often diffuse and disseminated.
- In patients with underlying malignancy (e.g. Hodgkin's disease), the presentation is evenly distributed between focal and nonfocal forms of encephalitis.

Toxoplasmosis and AIDS:

Patients with AIDS tend to present subacutely with nonspecific symptoms such as neuropsychiatric complaints, headache, fever, weight loss, disorientation, confusion, and lethargy evolving over 2 to 8 weeks.

- Later findings include evidence of focal CNS mass lesions, ataxia, aphasia, hemiparesis, visual field loss, vomiting, confusion, dementia, stupor and seizures. ¹⁷
- Toxoplasmosis presenting as subcutaneous mass has been reported among HIV-positive patients. ^{18 19}
- Primary cerebral lymphoma in AIDS patients may be mistaken for Toxoplasmosis. ²⁰

This disease is endemic or potentially endemic to all countries.

Toxoplasmosis in Uganda

Seroprevalence surveys:

- 34% of HIV-positive patients, and 27% of controls (1991 publication) ²¹
- 54% of HIV-positive patients (2006 publication) ²²
- 31% of domestic goats (2000 publication) ²³
- 47% of chickens in Kampala (2007 publication) ²⁴

Toxoplasmosis has been reported in a sand fox (*Vulpes rueppelli*). ²⁵

References

1. Scand J Infect Dis 2012 Dec 4;
2. J Am Acad Dermatol 2008 Nov ;59(5):781-4.
3. J Clin Microbiol 2013 Feb 6;
4. Parasitol Today 1998 Apr ;14(4):150-6.
5. Am J Med 2005 Mar ;118(3):212-6.
6. Reprod Toxicol 2006 May ;21(4):458-72.
7. Pediatr Infect Dis J 2011 Dec ;30(12):1056-61.
8. Expert Rev Anti Infect Ther 2012 Jul ;10(7):815-28.
9. Neonatal Netw 2001 Jun ;20(4):23-30.
10. Eur J Obstet Gynecol Reprod Biol 2007 Nov ;135(1):53-7.
11. Rev Inst Med Trop Sao Paulo 2012 Apr ;54(2):61-4.
12. Clin Experiment Ophthalmol 2012 Jun 19;
13. Am J Ophthalmol 2004 Jan ;137(1):1-17.
14. Pediatr Infect Dis J 2008 Jan ;27(1):27-32.
15. PLoS Negl Trop Dis 2008 ;2(8):e277.
16. Neth J Med 2009 Apr ;67(4):150-2.
17. CNS Drugs 2003 ;17(12):869-87.
18. Trop Biomed 2010 Dec ;27(3):447-50.
19. Diagn Cytopathol 2010 Oct ;38(10):716-20.
20. Brain Tumor Pathol 2011 Feb ;28(1):83-7.
21. Trans R Soc Trop Med Hyg 1991 Mar-Apr;85(2):227-9.
22. Acta Trop 2006 Dec ;100(3):218-22.
23. Acta Trop 2000 Jul 21;76(1):33-8.
24. Parasitology 2008 Jan ;135(Pt 1):39-45.
25. J Parasitol 2008 Aug ;94(4):976-7.

Trachoma

Agent	BACTERIUM. Chlamydia trachomatis, type A
Reservoir	Human
Vector	Fly
Vehicle	Infected secretions Fly Fomite
Incubation Period	5d - 12d
Diagnostic Tests	Culture or direct immunofluorescence of secretions. Serology. Nucleic acid amplification.
Typical Adult Therapy	Azithromycin 20 mg/kg as single dose. OR Doxycycline 100 mg/day PO X 14 days. Also administer topical Tetracycline
Typical Pediatric Therapy	Erythromycin 10 mg/kg PO QID X 4w. Also administer topical Tetracycline
Clinical Hints	Keratoconjunctivitis with palpebral scarring and pannus formation; 0.5% of infections result in blindness.
Synonyms	Egyptian ophthalmia, Granular conjunctivitis, Kornerkrankheit, Trachom, Tracoma. ICD9: 076 ICD10: A71

Clinical

Early symptoms include erythema and swelling of both bulbar and palpebral conjunctivae, associated with a watery or purulent discharge. ^{1 2}

- Additional findings may include preauricular lymphadenopathy and rhinitis.
- Examination reveals follicular hypertrophy and conjunctival scarring.
- Corneal scars (Herbert's pits), punctate keratitis and pannus formation may also be present. ³
- As scarring progresses, the eyelashes deviate (entropion) and may produce additional trauma and ulceration of the conjunctivae. ⁴
- Reinfection and bacterial superinfection are common ⁵ and may contribute to the progression of follicular trachoma. ⁶

Trachoma may be differentiated from inclusion conjunctivitis by the presence of corneal scarring and a preference of the latter for the upper tarsal conjunctivae

This disease is endemic or potentially endemic to all countries.

Trachoma in Uganda

1,050,797 cases of active trachoma were estimated in 2003.

Prevalence surveys:

21% of blindness among Sudanese refugees in Uganda (2000 publication) ⁷

Notable outbreaks:

2012 - An outbreak (50,000 cases) of trachoma was reported in Karamoja subregion. ⁸

References

1. Clin Microbiol Rev 2004 Oct ;17(4):982-1011, table of contents.
2. Lancet 2003 Jul 19;362(9379):223-9.
3. Lancet 2008 Jun 7;371(9628):1945-54.
4. Br Med Bull 2007 ;84:99-116.
5. Invest Ophthalmol Vis Sci 2011 Apr ;52(5):2181-6.
6. Invest Ophthalmol Vis Sci 2011 Jul ;52(8):6012-7.
7. East Afr Med J 2000 Nov ;77(11):580-2.
8. ProMED <promedmail.org> archive: 20121107.1396915

Trichinosis

Agent	PARASITE - Nematoda. Adenophorea: <i>Trichinella spiralis</i> (occasionally <i>T. nativa</i> , <i>T. britovi</i> , <i>T. pseudospiralis</i> , <i>T. nelsoni</i> , et al)
Reservoir	Wild carnivore Omnivore Marine mammal
Vector	None
Vehicle	Meat ingestion
Incubation Period	10d - 20d (range 1w - 10w)
Diagnostic Tests	Identification of larvae in tissue. Serology.
Typical Adult Therapy	Albendazole 400 mg PO BID X 14d. OR Mebendazole 200 to 400 mg PO tid X 3 days, then 400 to 500 mg PO. tid X 10 days. Give with prednisone 50 mg PO daily X 3 to 5 days (then 'taper' dosage)
Typical Pediatric Therapy	Albendazole 7 mg/kg BID X 14 d. OR Mebendazole 200 to 400 mg PO tid X 3 days, then 400 to 500 mg PO. tid X 10 days. Give with prednisone 50 mg PO daily X 3 to 5 days (then 'taper' dosage)
Clinical Hints	Early diarrhea and vomiting; subsequent myalgia, facial edema and eosinophilia; onset 1 to 4 weeks following ingestion of undercooked meat (usually pork); symptoms may persist for two months; case-fatality rate for symptomatic infection = 2%.
Synonyms	Trichinellose, Trichinellosis, Trichinose, Trikinose, Triquiniasis, Triquonosis. ICD9: 124 ICD10: B75

Clinical

The great majority of infections are subclinical.

- The development of symptoms depends on the number of larvae ingested.

Signs and symptoms:

During the first week of illness, the patient may diarrhea, abdominal pain and vomiting. ¹⁻³

- Symptoms associated with larval invasion appear during the second week and include fever, periorbital edema, subconjunctival hemorrhages and chemosis. ⁴
- Myositis is also common, and often appears in the extraocular muscles, progressing to involve the masseters, neck muscles, limb and lumbar muscles.
- Additional symptoms may include headache, cough, dyspnea, hoarseness and dysphagia.
- Occasionally, a macular or petechial rash, or retinal or subungual splinter hemorrhages are seen.
- Hepatomegaly is common. ⁵
- Laboratory studies may reveal marked eosinophilia, hypoalbuminemia, decreased erythrocyte sedimentation rate, proteinuria or hematuria.
- Rare instances of renal dysfunction ⁶, encephalitis ⁷ and eosinophilic meningitis have been reported. ⁸

Clinical course:

- Systemic symptoms usually peak 2 to 3 weeks after infection and then slowly subside; however, weakness may persist for weeks.
- A number of clinical findings may persist for several months: hypocalcemia, hypomagnesemia, fatigue, myalgia (notably in the legs), cardiovascular disorders, neurological, psychiatric, and allergic illnesses. ⁹
- Deaths are ascribed to myocarditis ^{10 11}, encephalitis or pneumonia.

This disease is endemic or potentially endemic to all countries.

Trichinosis in Uganda

No cases were reported in 1999.

References

1. *Expert Rev Anti Infect Ther* 2003 Oct ;1(3):471-82.
2. *Postgrad Med J* 2002 Jan ;78(915):15-22.

3. *Clin Microbiol Rev* 1996 Jan ;9(1):47-54.
4. *Clin Microbiol Rev* 2009 Jan ;22(1):127-45, Table of Contents.

5. Foodborne Pathog Dis 2011 Sep ;8(9):943-8.
6. Foodborne Pathog Dis 2011 Feb ;8(2):179-88.
7. Foodborne Pathog Dis 2011 May ;8(5):579-85.
8. Clin Microbiol Rev 2009 Apr ;22(2):322-48, Table of Contents.
9. Vet Parasitol 2009 Feb 23;159(3-4):320-3.
10. Foodborne Pathog Dis 2010 Oct ;7(10):1235-8.
11. Foodborne Pathog Dis 2011 Aug ;8(8):853-60.

Trichomoniasis

Agent	PARASITE - Protozoa. Archezoa, Parabasala, Trichomonadea. Flagellate: <i>Trichomonas vaginalis</i>
Reservoir	Human
Vector	None
Vehicle	Sexual contact
Incubation Period	4d - 28d
Diagnostic Tests	Microscopy of vaginal discharge. ELISA, culture, antigen detection tests available. Nucleic acid amplification.
Typical Adult Therapy	Metronidazole or Tinidazole 2g PO as single dose to both sexual partners
Typical Pediatric Therapy	Metronidazole 5 mg/kg PO TID X 7d. OR Tinidazole 50 mg/kg PO X 1 (maximum 2 grams)
Clinical Hints	Vaginal pruritus, erythema and thin or frothy discharge; mild urethritis may be present in male or female.
Synonyms	Pentatrichomonas, Tetratrichomonas, Trichomonaden, Trichomonas, Trichomonas vaginalis, Trichomoniasis, Tritrichomonas. ICD9: 131 ICD10: A59

Clinical

10% to 50% of infections are asymptomatic.

- Symptoms often begin or worsen during the menstrual period.
- Infection is usually characterized by vaginal discharge and vulvovaginal irritation. ¹
- Dysuria may be present, and dyspareunia is common.
- As many as two thirds of infected women complain of a disagreeable odor.
- Abdominal discomfort is present in 5% to 12%.

Examination reveals a copious loose discharge that pools in the posterior vaginal fornix. ²

- The discharge is yellow or green in 5% to 40%, and bubbles are observed in the discharge in 10% to 33%. ³
- The material has a pH above 4.5 in 66% to 91% of cases.
- Endocervical disease is not caused by *T. vaginalis*.
- Punctate hemorrhages (colpitis macularis or "strawberry cervix") are seen on colposcopically in 45% of infected women, but in only 2% by visual inspection alone.
- Parasites can be recovered from the urethra and paraurethral glands in more than 95% of the women, and may explain the association of the infection with urinary frequency and dysuria.

Reported complications of trichomonal vaginitis include vulvar ulceration ⁴, and vaginitis emphysematosa • the presence gas-filled blebs in the vaginal wall. ⁵

- Gestational trichomoniasis may be associated with premature labor and low birth weight, postabortal infection or premature rupture of the membranes.
- Spread of trichomonads beyond the lower urogenital tract is extremely rare.
- Sporadic cases of neonatal pneumonia due to *Trichomonas vaginalis* are reported. ⁶

Trichomoniasis has been associated with endometritis, adnexitis, pyosalpinx, infertility, preterm birth, low birth weight, bacterial vaginosis, and increased risk of cervical cancer, HPV, and HIV infection. ⁷

- In men, its complications include urethritis, prostatitis, epididymitis, and infertility through interference with sperm function. ⁸

Most men carrying trichomonads are asymptomatic; however, the organism is implicated in 5% to 15% of patients with nongonococcal urethritis.

- The discharge from trichomonal urethritis is usually milder than that seen with other infections.
- Epididymitis, superficial penile ulcerations (often beneath the prepuce) and prostatitis are also described.

Tritrichomonas foetus pneumonia ⁹ and peritonitis have been reported in immunosuppressed patients. ¹⁰

- *Trichomonas tenax* has been reported to cause pneumonia in an immunosuppressed patient. ¹¹

This disease is endemic or potentially endemic to all countries.

Trichomoniasis in Uganda

Prevalence surveys:

- 30.8% of women in Rakai (2003 publication) ¹²
- 18.9% of women with genital disease (southwestern Uganda, 1999 to 2001) ¹³
- 8.0% of female and 0% of male adolescents in Kampala (2006 publication) ¹⁴
- 3.3% of HIV-positive patients (2012 publication) ¹⁵

References

1. J Reprod Med 2004 Oct ;49(10):781-6.
2. Clin Microbiol Rev 2004 Oct ;17(4):794-803, table of contents.
3. J Infect Dis 1980 Feb ;141(2):137-143.
4. Int J STD AIDS 2010 Sep ;21(9):664-5.
5. Infect Dis Clin North Am 2005 Jun ;19(2):387-406.
6. Am J Trop Med Hyg 2008 Jan ;78(1):17-9.
7. J Reprod Immunol 2009 Dec ;83(1-2):185-9.
8. Andrologia 2011 Aug ;43(4):283-5.
9. J Clin Microbiol 2006 Mar ;44(3):1165-8.
10. Emerg Infect Dis 2011 Jul ;17(7):1312-3.
11. New Microbiol 2012 Jan ;35(1):83-7.
12. Sex Transm Infect 2003 Apr ;79(2):98-105.
13. Sex Transm Infect 2005 Dec ;81(6):488-93.
14. J Adolesc Health 2006 Mar ;38(3):213-21.
15. J Infect Dev Ctries 2012 ;6(3):276-82.

Trichostrongyliasis

Agent	PARASITE - Nematoda. Phasmidea: Trichostrongylus colubriformis, T. orientalis, T. probolurus
Reservoir	Herbivore
Vector	None
Vehicle	Water Food Vegetation
Incubation Period	21d
Diagnostic Tests	Identification of ova in stool or duodenal aspirate.
Typical Adult Therapy	Albendazole 400 mg PO X 1. OR Pyrantel pamoate 11 mg/kg (max 1g) PO once. OR Mebendazole 100 mg PO BID X 7d
Typical Pediatric Therapy	As for adult
Clinical Hints	Diarrhea, abdominal pain and weight loss; eosinophilia is often present; infestation may persist for years; fatality and sequelae are not reported.
Synonyms	Haemonchus, Marshallagia, Ostertagia, Trichostrongylus. ICD9: 127.6 ICD10: B81.2

Clinical

Most infections are asymptomatic, or characterized by mild nonspecific abdominal symptoms.

- Heavy infections may result in episodic diarrhea, abdominal pain and weight loss. ¹
- Rare instances of cholecystitis are reported.

This disease is endemic or potentially endemic to 38 countries. Although Trichostrongyliasis is not endemic to Uganda, imported, expatriate or other presentations of the disease have been associated with this country.

Trichostrongyliasis in Uganda

Prevalence surveys:

1% of pregnant women in Entebbe (2004 to 2005) ²

References

1. Rev Ig Bacteriol Virusol Parazitol Epidemiol Pneumoftiziol Bacteriol Virusol Parazitol Epidemiol 1978 Oct-Dec;23(4):201-6.
2. PLoS Negl Trop Dis 2009 ;3(6):e473.

Trichuriasis

Agent	PARASITE - Nematoda. Adenophorea: Trichuris trichiura
Reservoir	Human
Vector	None
Vehicle	Soil ingestion Sexual contact (rare) Fly
Incubation Period	2m - 2y
Diagnostic Tests	Stool microscopy or visualization of adult worms (adults are approximately 3 cm long).
Typical Adult Therapy	Mebendazole 100 mg PO BID X 3d. OR Albendazole 400 mg PO daily X 3 to 7 days OR Ivermectin 200 mg/kg PO daily X 3 days
Typical Pediatric Therapy	Mebendazole 100 mg PO BID X 3d (>age 2). OR Albendazole 400 mg PO X 3 to 7 days OR Ivermectin 200 mg/kg PO daily X 3 days
Clinical Hints	Abdominal pain, bloody diarrhea, rectal prolapse or intestinal obstruction are occasionally encountered; the parasite may survive for as long as five years in the human host.
Synonyms	Trichocephaliasis, Trichuris trichiura, Tricuriasis, Whipworm. ICD9: 127.3 ICD10: B79

Clinical

The vast majority of infections are asymptomatic. ¹

- Symptoms are aggravated by concurrent shigellosis, balantidiasis or amebiasis.
- Heavy infestations are characterized by dysentery or rectal prolapse. ^{2 3}
- Infants may develop hypoproteinemia, anemia, mental retardation and digital clubbing. ⁴⁻⁸
- In some cases, chronic infection may result in edema of the ileocecal valve, suggestive of "malignancy" ⁹

This disease is endemic or potentially endemic to all countries.

Trichuriasis in Uganda

Prevalence surveys:

- 5.0% nationwide (1998 to 2005) ¹⁰
- 8.1% of persons in Kabale District (2003) ¹¹
- 9.1% of otherwise healthy pregnant women at enrolment to a trial of deworming. (Entebbe, 2007 publication) ¹²
- 9% of pregnant women (Entebbe and Katabi, 2003 to 2005) ¹³
- 4.6% of pregnant women in Masindi District (2003 to 2004) ¹⁴
- 33.3% of mothers and 52.8% of their children in Habutobere; 47.4% of mothers and 54.6% of their children in Musezero (southwestern Uganda, 2010) ¹⁵
- 7.3% of children below age 20 in Southern Uganda (1998) ¹⁶
- 28% of primary school children in Kampala (1997 publication) ¹⁷
- 2.4% of school children in northwest and southeast Uganda (2002 to 2003) ¹⁸
- 2.2% of school children in 2003; 2.5% in 2004; 1.6% in 2005 (nationwide)
- 5% of children below age 3 years living on the Ugandan shore of Lake Victoria (2006 publication) ¹⁹
- 12.9% of school children in proximity to Lake Victoria (2009 publication) ²⁰
- 6.8% of school children in the Great Lakes region of Uganda, Tanzania, Kenya and Burundi (2010 publication) ²¹

References

1. Lancet 2006 May 6;367(9521):1521-32.
2. Gastroenterol Clin North Am 1996 Sep ;25(3):579-97.
3. J Natl Med Assoc 2004 Jan ;96(1):93-6.
4. Gastrointest Endosc 2010 Jan ;71(1):200-4.
5. Trans R Soc Trop Med Hyg 1976 ;70(4):313-6.
6. Arch Dis Child 1978 May ;53(5):411-3.
7. Parasitology 2000 ;121 Suppl:S73-95.
8. Trop Biomed 2012 Dec ;29(4):626-31.
9. ISRN Gastroenterol 2011 ;2011:105178.
10. Trop Med Int Health 2005 Nov ;10(11):1187-9.

11. [Trans R Soc Trop Med Hyg 2005 Jan ;99\(1\):18-24.](#)
12. [Trans R Soc Trop Med Hyg 2007 Sep ;101\(9\):899-907.](#)
13. [J Infect Dis 2008 Sep 15;198\(6\):920-7.](#)
14. [Trans R Soc Trop Med Hyg 2008 Feb ;102\(2\):130-6.](#)
15. [Trans R Soc Trop Med Hyg 2012 Feb ;106\(2\):75-83.](#)
16. [East Afr Med J 2001 Jun ;78\(6\):283-6.](#)
17. [East Afr Med J 1997 May ;74\(5\):311-4.](#)
18. [Trans R Soc Trop Med Hyg 2006 Nov ;100\(11\):1039-48.](#)
19. [Ann Trop Med Parasitol 2006 Jun ;100\(4\):315-26.](#)
20. [Geospat Health 2009 Nov ;4\(1\):39-53.](#)
21. [Trop Med Int Health 2010 Feb ;15\(2\):198-207.](#)

Tropical phagedenic ulcer

Agent	BACTERIUM Mixed infection by ? Fusobacterium species and Borrelia
Reservoir	Human
Vector	None
Vehicle	Direct inoculation ? via minor trauma
Incubation Period	Unknown
Diagnostic Tests	Wound smear suggestive of fusobacterial infection.
Typical Adult Therapy	Systemic Penicillin G . Excision/debridement as necessary
Typical Pediatric Therapy	As for adult
Clinical Hints	A deep, painful, foul-smelling ulcer (usually of the leg) with undermined edges; may be complicated by secondary infection.
Synonyms	Acute phagedenic ulcer. ICD9: 682.7 ICD10: A69.8,L97

Clinical

95% of ulcers involve the ankle or lower third of the leg.

- Minor trauma is followed by a tender indurated area which evolves into a round or oval skin ulcer. ^{1 2}
- Ulcers favor the lower extremities, and tend to be single, painful and foul-smelling.
- Ulcers spread rapidly, and result in exposure of underlying muscles and tendons.
- Fever and restlessness are common.
- After 4 or more weeks, ulcers may become painless and chronic, and persist for decades.
- Scar carcinomas develop in 2% of cases, and constitute a common form of malignancy in parts of Africa.

This disease is endemic or potentially endemic to 69 countries.

References

1. [Int J Dermatol 1988 Jan-Feb;27\(1\):49-53.](#)
2. [J Int Coll Surg 1962 Aug ;38:120-8.](#)

Tropical pulmonary eosinophilia

Agent	UNKNOWN Possibly related to filarial infection
Reservoir	Unknown
Vector	Unknown
Vehicle	Unknown
Incubation Period	Unknown
Diagnostic Tests	Antifilarial antibodies may be present. Response to therapeutic trial.
Typical Adult Therapy	Diethylcarbamazine 2 mg/kg PO TID X 10d
Typical Pediatric Therapy	As for adult
Clinical Hints	Chronic cough, wheezing, dyspnea, reticular-nodular pulmonary infiltrates and eosinophilia (over 3,000/cu. mm.) acquired in countries known to be endemic for filariasis.
Synonyms	

Clinical

Tropical pulmonary eosinophilia is characterized by recurrent episodes of paroxysmal, dry cough, wheezing, and dyspnea. ¹⁻³

- Malaise, anorexia, and weight loss are common.
- Symptoms are worse at night.
- Physical examination reveals scattered wheezes and crackles.
- Some patients have fever, hepatomegaly and lymphadenopathy.
- Symptoms fluctuate in severity over many months.

Eosinophilia is present in the majority of patients, often at very high levels (as high as 60,000/cu mm) • however, the level of eosinophilia is not related to the severity of symptoms.

- Chest radiographs reveal scattered reticulonodular opacities ⁴ which may be mistaken for miliary tuberculosis. ⁵
- Serum antibodies to filaria are present.
- A presumptive clinical diagnosis can usually be made through successful response to antifilarial therapy.
- A second course may be necessary in some cases.

This disease is endemic or potentially endemic to 109 countries.

References

1. Clin Infect Dis 2005 Feb 15;40(4):635-6.
2. Clin Chest Med 2002 Jun ;23(2):377-96, ix.
3. Curr Opin Pulm Med 2007 Sep ;13(5):428-33.
4. Semin Respir Crit Care Med 2006 Apr ;27(2):171-84.
5. Parasitol Int 2012 Jun ;61(2):381-4.

Trypanosomiasis - African

Agent	PARASITE - Protozoa. Neozoa, Euglenozoa, Kinetoplastidea. Flagellate: <i>Trypanosoma</i> [Trypanozoon] <i>brucei gambiense</i> and <i>T. b. rhodesiense</i>
Reservoir	Human Deer Wild carnivore Cattle
Vector	Fly (<i>Glossina</i> = tsetse fly)
Vehicle	None
Incubation Period	3d - 21d (acute illness)
Diagnostic Tests	Identification of protozoa in CSF, blood, lymph node aspirate. Serology. Nucleic acid amplification.
Typical Adult Therapy	Early: Pentamidine 4 mg/kg IM qod X 10 doses. OR Suramin 1g IV days 1, 3, 7, 14, 21 (after test dose 100 mg) OR Eflornithine (<i>gambiense</i> only) 100 mg q6h IV X 14 d; then 75 mg/kg PO X 21-30 d. Late + CNS disease: Melarsoprol
Typical Pediatric Therapy	Early: Pentamidine 4 mg/kg IM qod X 10 doses. OR Suramin 20 mg/kg IV days 1, 3, 7, 14, 21 (after test dose 20 mg) Late + CNS: Melarsoprol
Clinical Hints	Chancre, myalgia, arthralgia, lymphadenopathy and recurrent fever; later mental changes, sensory disorders and heart failure; disease due to <i>Trypanosoma brucei rhodesiense</i> is more rapid and virulent than that due to <i>T. b. gambiense</i> .
Synonyms	African sleeping sickness, African trypanosomiasis, Gambian fever, Schlafkrankheit, <i>Trypanosoma brucei</i> , <i>Trypanosoma congolense</i> , <i>Trypanosoma evansi</i> , <i>Trypanosoma lewisi</i> , Trypanosomiasis, afrikanische, U.T.I., UTI. ICD9: 086.3,086.4,086.5 ICD10: B56

Clinical

WHO Case definition for surveillance:

- In the early stages, a painful chancre, which originates as a papule and evolves into a nodule may be found at the primary site of tsetse fly bite.
- There may be fever, intense headache, insomnia, painless lymphadenopathy, anemia, local edema and rash. In the later stage, there is cachexia, somnolence and signs of central nervous system involvement.
- The disease may run a protracted course of several years in the case of *Trypanosoma brucei gambiense*. In case of *T. b. rhodesiense*, the disease has a rapid and acute evolution.
- Both diseases are always fatal without treatment.
- The painful chancre is very rare in *T. b. gambiense* infection.

Laboratory criteria for diagnosis

- Presumptive: serological: card agglutination trypanosomiasis test (CATT) for *T. b. gambiense* only or immunofluorescent assay (IFA) for *T. b. rhodesiense* mainly and possibly for *T. b. gambiense*.
- Confirmative: parasitological: detection (microscopy) of trypanosomes in blood, lymph nodes aspirates or CSF.

Case classification

- Suspected: A case that is compatible with the clinical description and/or a history of exposure.
- Probable: A case with a positive serology with or without clinical symptoms in persons without previous history of trypanosomiasis diagnosis or treatment.
- Confirmed: A case with positive parasitology, with or without clinical symptoms.

Notes:

- In the early stage or even early in the late stage of the disease there are often no clinical signs or symptoms which can be associated with the disease.
- Suspicion is then based on local risk of contracting the disease and local disease historical background.
- Confirmed positive healthy carriers are a major public health risk. As a reservoir of parasites, they disseminate the disease, and must be treated as soon as possible.

Acute trypanosomiasis:

The initial sign of African trypanosomiasis is a chancre which develops at the site of inoculation, 1 to 2 weeks following the bite of a tsetse fly.

- The chancre may reach a diameter of several centimeters, and be associated with regional adenopathy, but resolves over several weeks.
- In most cases, the chancre is noted by neither the patient nor the clinician.
- Fever appears weeks to months following inoculation, and is characteristically intermittent.
- Lymphadenopathy is a fairly constant feature of west African trypanosomiasis.

- The nodes are discrete, movable, rubbery, and nontender.
- Supraclavicular and cervical nodes are often visibly discernible, and enlargement of the nodes of the posterior cervical triangle ("Winterbottom's sign") is common in the west African form.
- Additional findings at this point may include hepatosplenomegaly; edema of the face, hands and feet; pruritis; an irregular circinate, 5 to 10 cm rash on the trunk, shoulders, buttocks and thighs ¹ ; headache, asthenia, weight loss, arthralgias, and tachycardia.
- In some cases, trypanosomiasis has been associated with cryoglobulinemic membranoproliferative glomerulonephritis. ²

***Trypanosoma brucei gambiense* infection:**

In the West African form, the meningoencephalitic stage may develop months or even years after the initial infection. ³

- Findings include irritability, personality changes, indifference, apathy, daytime somnolence (often with insomnia at night), slurred speech, choreiform movements of the trunk, neck, and extremities, tremors of the tongue and fingers, ataxia, and muscular fasciculations. ⁴
- CSF cell counts above 5 per cu mm are considered indicative of brain involvement. ^{5 6}
- The final phase of the CNS disease is progression to coma and death ⁷ ; however, survival without therapy is reported. ⁸

***Trypanosoma brucei rhodesiense* infection:**

The East African form tends to follow a more acute course, with an incubation of a few weeks to several weeks.

- Intermittent fever, headache, myalgia, and rash develop early; while lymphadenitis is not a prominent feature.
- Persistent tachycardia is common, and some patients die of arrhythmias, congestive heart failure or pericarditis before the onset of neurological disease.
- If untreated, the East African form is fatal within weeks to months.

This disease is endemic or potentially endemic to 36 countries.


Trypanosomiasis - African in Uganda

Time and Place:

Trypanosomiasis is most common in the area of the White Nile, Sudan border, north shore of Lake Victoria, Iganga ⁹ , Kambuli, Mokono and Tinja.

- Epidemics of sleeping sickness occurred during 1901 to 1915, 1940 to 1946, and 1976 to 1989. ¹⁰
- The disease focus has moved from lakeshore Buganda (1905 to 1920) to the Busoga and south-east districts during recent years. ^{11 12}
- In recent years, outbreaks of trypanosomiasis have been introduced into Soroti District by imported cattle. ^{13 14}
- Outbreaks were reported in Luuka Country (western Iganga district) in 1976, and Soroti District in 1998. ¹⁵
- An epidemic of *Trypanosoma brucei rhodesiense* infection in eastern Uganda began in 1998 as a result of livestock movements, and continued unabated as of 2005. ^{16 17}
- As of 2004, foci of *T. rhodesiense* infection are found in Busia, Busiri, Iganga, Jinja, Kamule, Kamuli, Nayuge, Mukono, Soroti and Toro districts.
- Recent spread of *T.b. rhodesiense* into the midwestern part of Uganda (Masindi district) ^{18 19} and southeastern region (Mayuge District) has been reported. ²⁰ Northward extension of the disease may be related to movement of livestock. ²¹
- An outbreak of Ngana (bovine trypanosomiasis) was reported in Tororo in 2008. ²²

Two million persons are considered at risk.



Graph: Uganda. Trypanosomiasis - African, cases

Notes:

1. Rates range from 8 to 50 per 100,000 per year.
2. 140 fatal cases were officially reported during 2000 to 2009 - the true number was estimated at 1,700. ²³

Individual years:

1995 - True number estimated at 5,000 cases.

Over 250,000 fatal cases were estimated in the Lake Victoria region during 1900 to 1920 - most in the Busoga region. ²⁴
 - 19,974 cases were identified in Busoga during 1976 to 1983. ²⁵

Exported cases:

- 2006 - A South African tourist acquired trypanosomiasis in Uganda. ²⁶
 2008 - An Australian who had immigrated from Sudan acquired trypanosomiasis (nonfatal) in Uganda. ²⁷
 2009 - A Polish tourist acquired trypanosomiasis in Queen Elizabeth National Park. ²⁸

Prevalence surveys:

- 0.75% of cattle in south-western Uganda (*Trypanosoma brucei*, 2003).
 5% of cattle, 13.9% of pigs, and 0.4% of small ruminants (southwestern Uganda, 2000)
 17.3% of cattle in Tororo and Soroti Districts (*Trypanosoma brucei*, 2000) ²⁹
 0.2% of goats, 3.5% of dogs, 5.0% of sheep, 7.5% of cattle, and 15.5% of pigs (northwestern Uganda, 2011 publication) ³⁰

Infecting species:

The endemic species in Uganda are *Trypanosoma brucei rhodesiense* (in the southeast and center) and *T.b. gambiense* (in the northwest and south). ³¹

- *T.b. rhodesiense* accounted for over 80% of cases reported during 1978 to 1989.
- *T.b. gambiense* accounted for 45% of cases reported during 1990 to 1994; 57.8% during 2000 to 2004.
- As of 2004, foci of *T.b. gambiense* infection are found in Adjumani, Arua, Moyo and Yumbe districts.

Vectors:

- The local vectors are *Glossina fuscipes* (ssp. *quanzensis*, *martinii* and *fuscipes*) ^{32 33} and *G. pallidipes*. ³⁴

Notable outbreaks:

- 1957 - An outbreak of trypanosomiasis was reported. ³⁵
 1981 (publication year) - An outbreak was reported in Busoga. ³⁶⁻⁴¹
 2001 (publication year) - An outbreak was reported in Soroti District. ⁴²
 2009 - An outbreak (29 cases, 18 fatal) was reported in Dokola District. ⁴³

References

1. J Travel Med 2007 May-Jun;14(3):192-6.
2. Ann Pathol 2012 Feb ;32(1):40-52.
3. J Clin Invest 2004 Feb ;113(4):496-504.
4. J Neurol 2006 Apr ;253(4):411-6.
5. Trop Med Int Health 2005 May ;10(5):395-403.
6. PLoS Negl Trop Dis 2011 ;5(3):e968.
7. Lancet Infect Dis 2003 Dec ;3(12):804-8.
8. PLoS Negl Trop Dis 2012 ;6(6):e1691.
9. Afr Anal 1991 Jun 14;:14.
10. Emerg Infect Dis 2006 May ;12(5):813-20.
11. Afr Health Sci 2006 Dec ;6(4):223-31.
12. Ann Trop Med Parasitol 1997 Oct ;91(7):851-6.
13. Lancet Infect Dis 2003 Jan ;3(1):42-5.
14. Lancet 2001 Aug 25;358(9282):625-8.
15. Lancet 2001 Aug 25;358(9282):625-8.
16. Lancet 2005 Aug 27-Sep 2;366(9487):745-7.
17. ProMED <promedmail.org> archive: 20050829.2552
18. Curr Opin Infect Dis 2002 Oct ;15(5):477-84.
19. Ann Trop Med Parasitol 1999 Dec ;93(8):817-22.
20. ProMED <promedmail.org> archive: 20071208.3962
21. ProMED <promedmail.org> archive: 20091217.4272
22. ProMED <promedmail.org> archive: 20080410.1316
23. Emerg Infect Dis 2012 Oct ;18(10):1686-7.
24. Emerg Infect Dis 2004 Apr ;10(4):567-73.
25. Trop Med Parasitol 1985 Jun ;36(2):72-6.
26. J Travel Med 2012 Jan-Feb;19(1):44-53.
27. Med J Aust 2010 Apr 5;192(7):417-9.
28. ProMED <promedmail.org> archive: 20090810.2844
29. Onderstepoort J Vet Res 2002 Jun ;69(2):133-40.
30. Vet Parasitol 2012 Jun 8;187(1-2):157-67.
31. BMJ 2005 Nov 26;331(7527):1238-41.
32. East Afr Med J 1983 Jun ;60(6):397-401.
33. Ann Soc Belg Med Trop 1991 ;71 Suppl 1:35-47.
34. Schweiz Med Wochenschr 1999 Aug 10;129(31-32):1091-8.
35. Trans R Soc Trop Med Hyg 1959 Sep ;53:384-93.
36. Trop Doct 1981 Oct ;11(4):175-8.
37. Bull Anim Health Prod Afr 1981 Mar ;29(1):55-8.
38. Trans R Soc Trop Med Hyg 1983 ;77(2):277-8.
39. Trop Med Parasitol 1985 Jun ;36(2):72-6.
40. Ann Soc Belg Med Trop 1989 ;69 Suppl 1:173-9; discussion 212-4.
41. Ann Soc Belg Med Trop 1991 ;71 Suppl 1:35-47.
42. Lancet 2001 Aug 25;358(9282):625-8.
43. ProMED <promedmail.org> archive: 20090315.1057

Tuberculosis

Agent	BACTERIUM. Actinomycetes, <i>Mycobacterium tuberculosis</i> An aerobic acid-fast bacillus
Reservoir	Human Cattle
Vector	None
Vehicle	Air Dairy products
Incubation Period	4w - 12w (primary infection)
Diagnostic Tests	Microscopy. Culture. Nucleic acid amplification. Inform laboratory when this diagnosis is suspected.
Typical Adult Therapy	Respiratory isolation. Typical pulmonary infection is treated with 6 months of Isoniazid , Rifampin & Pyrazinamide
Typical Pediatric Therapy	As for adult
Vaccine	BCG
Clinical Hints	Cough, "night sweats" and weight loss; often presents as prolonged fever (FUO) or infection of bone, meninges, kidneys or other organs; most infections represent reactivation of old foci in lungs, brain, bone, kidneys etc.
Synonyms	Consumption, <i>Mycobacterium africanum</i> , <i>Mycobacterium bovis</i> , <i>Mycobacterium caprae</i> , <i>Mycobacterium orygis</i> , <i>Mycobacterium tuberculosis</i> , <i>Oryx bacillus</i> , Phthisis, TB, TB meningitis, Tuberculose, Tuberculose miliar, Tuberculosi, Tuberculous meningitis, Tuberkulose, White plague. ICD9: 010,012,013,014,015,016,017,018 ICD10: A15,A16,A17,A18,A19

Clinical

WHO Case definition for surveillance:

Pulmonary tuberculosis, sputum smear positive (PTB+)

- Tuberculosis in a patient with at least two initial sputum smear examinations (direct smear microscopy) positive for Acid-Fast Bacilli (AFB), or
- Tuberculosis in a patient with one sputum examination positive for acid fast bacilli and radiographic abnormalities consistent with active pulmonary tuberculosis as determined by the treating medical officer, or
- Tuberculosis in a patient with one sputum specimen positive for acid-fast bacilli and at least one sputum that is culture positive for acid-fast bacilli.

Pulmonary tuberculosis, sputum smear negative (PTB-)

Tuberculosis in a patient with symptoms suggestive of tuberculosis and having one of the following:

- Three sputum specimens negative for acid-fast bacilli
- Radiographic abnormalities consistent with pulmonary tuberculosis and a lack of clinical response to one week of a broad-spectrum antibiotic
- Decision by a physician to treat with a full curative course of antituberculous chemotherapy

Pulmonary tuberculosis, sputum smear negative, culture positive

- Tuberculosis in a patient with symptoms suggestive of tuberculosis and having sputum smear negative for acid-fast bacilli and at least one sputum that is culture positive for *M. tuberculosis* complex

Extra-pulmonary tuberculosis

- Tuberculosis of organs other than lungs: pleura, lymph nodes, abdomen, genito-urinary tract, skin, joints and bones, tuberculous meningitis, etc.
- Diagnosis should be based on one culture positive specimen from an extra-pulmonary site, or histological or strong clinical evidence consistent with active extra-pulmonary tuberculosis, followed by a decision by a medical officer to treat with a full course of anti-tuberculous therapy
- Any patient diagnosed with both pulmonary and extra-pulmonary tuberculosis should be classified as a case of pulmonary tuberculosis

The clinical features of tuberculosis are protean, and largely determined by the site of infection and clinical substrate.

- Most infections represent reactivation of a dormant focus in a lung, with resultant chronic fever, weight loss, nocturnal diaphoresis, productive cough and typical roentgenographic findings. ¹
- Reactivation of an extrapulmonary focus (kidney, bone, central nervous system ^{2 3}, skin ⁴, gastrointestinal ⁵⁻¹⁰ and hepatobiliary system ¹¹, eyes ^{12 13}, skeletal muscle ¹⁴⁻¹⁸, reproductive tract ¹⁹, breast ^{20 21}, etc) will result in signs referable to the infected organ.
- The extent and severity of disease are influenced by patient age, nutrition, immune function ^{22 23}, and many other

factors which are beyond the scope of this module.

- Nocardiosis may mimic tuberculosis, particularly in the setting of HIV infection. ²⁴
- The appearance of a miliary infiltrates in tropical pulmonary eosinophilia ²⁵ or *Chlamydomphila pneumoniae* infection may suggest a diagnosis of tuberculosis. ²⁶
- Spinal histoplasmosis may mimic tuberculosis spondylodiscitis ²⁷ ; and gastrointestinal histoplasmosis may mimic abdominal tuberculosis. ²⁸
- The clinical features of melioidosis are similar to those of tuberculosis: prolonged fever, weight loss, latency with reactivation, upper-lobe infiltrates, etc. ²⁹⁻³¹
- Tularemia ³² and leprosy may manifest as lymphadenopathy mimicking tuberculosis ³³

This disease is endemic or potentially endemic to all countries.

Tuberculosis in Uganda

Vaccine Schedule:

BCG - birth

DTWP - 6, 10, 14 weeks


DTwPHibHep - 6, 10, 14 weeks

Measles (monovalent) - 9 months


OPV - birth; 6, 10, 14 weeks

TT - Women of child-bearing age; +1, +6 months; +1, +1 year

Vitamin A - 6, 12, 18, 24, 30, 36, 42 months



Graph: Uganda. Tuberculosis - WHO-UNICEF est. % BCG coverage



Graph: Uganda. Tuberculosis, cases


Notes:

Individual years:

1997 - True number estimated at 66,000 = 320 per 100,000

The prevalence of tuberculosis in Kampala is 1,400 per 100,000 - and 440 per 100,000 have smear-positive tuberculosis (2001 to 2002).

57% of health care workers in Kampala are tuberculin-positive (2001). ³⁴



Graph: Uganda. Tuberculosis, estimated (WHO) deaths

Tuberculosis and HIV infection:

The incidence of tuberculosis among persons living with HIV infection is 3.4% per year (1997).

- 4.6% of newly-diagnosed HIV-positive individuals have tuberculosis (Kampala, 2006 publication) ³⁵
- 58% of tuberculosis patients are HIV-positive (1989 publication) ³⁶ ; 50% in 1997.
- *M. tuberculosis* bacteremia was identified in 12.3% of febrile HIV-positive adults (Kampala, 2007) ³⁷

During 1996 to 1997, primary resistance to INH was present in 6.7% of isolates; multidrug resistance (MDR) in 0.5%.

- During 2003 to 2006, 24.7% of isolates in Kampala were resistant to INH and 12.7% were MDR. ³⁸
- In 2008, INH resistance / MDR-TB was identified in 5.8% / 1.1% of new cases, and 20.0% / 11.7% of previously-treated cases (Kampala). ³⁹

The most common cause of tuberculosis in Kampala is *Mycobacterium africanum* subtype II.

Bovine tuberculosis is found in 74.1% of herds in the Mbara milk basin (2002). ⁴⁰

- Organisms associated with cervical lymphadenitis among pastoral communities in the Karamoja region include *Mycobacterium avium*, *M. tuberculosis*, *M. bovis* and *M. intracellulare* (2008 publication). ⁴¹

References

1. Acad Emerg Med 2000 Sep ;7(9):1056-60.
2. Clin Microbiol Rev 2008 Apr ;21(2):243-61, table of contents.
3. Lancet Neurol 2005 Mar ;4(3):160-70.
4. Trop Med Int Health 2006 Oct ;11(10):1521-8.
5. Southeast Asian J Trop Med Public Health 2009 May ;40(3):505-10.
6. Trop Gastroenterol 2009 Jan-Mar;30(1):35-9.
7. Singapore Med J 2009 Jun ;50(6):638-45; quiz 646.
8. Curr Opin Infect Dis 2009 Oct ;22(5):490-6.
9. J Infect Dev Ctries 2010 Oct ;4(10):650-4.
10. J Gastrointest Surg 2011 Oct ;15(10):1837-41.
11. South Med J 2008 Apr ;101(4):356-61.
12. J Bras Pneumol 2008 Feb ;34(2):98-102.
13. J Fr Ophtalmol 2009 Nov ;32(9):673-8.
14. Hand (N Y) 2009 Mar ;4(1):88-91.
15. Int J Rheum Dis 2010 Feb 1;13(1):82-5.
16. Indian J Tuberc 2010 Jan ;57(1):34-40.
17. Med Ultrason 2011 Sep ;13(3):245-8.
18. Acta Orthop Traumatol Turc 2011 ;45(4):276-9.
19. J Clin Microbiol 2008 Dec ;46(12):4068-70.
20. Trans R Soc Trop Med Hyg 2009 Jun ;103(6):559-63.
21. Int J Tuberc Lung Dis 2010 Jun ;14(6):758-63.
22. Curr Opin Pulm Med 2005 May ;11(3):203-7.
23. N Engl J Med 1991 Jun 6;324(23):1644-50.
24. Trans R Soc Trop Med Hyg 2008 Mar ;102(3):219-24.
25. Parasitol Int 2012 Jun ;61(2):381-4.
26. Pediatr Emerg Care 2009 Sep ;25(9):597-8.
27. Acta Reumatol Port 2008 Jul-Sep;33(3):360-3.
28. J Assoc Physicians India 2009 Jan ;57:76-8.
29. Curr Opin Infect Dis 2004 Apr ;17(2):131-6.
30. Eur Respir J 2003 Sep ;22(3):542-50.
31. Int J Tuberc Lung Dis 2008 Oct ;12(10):1209-15.
32. Ann Dermatol Venereol 2009 Oct ;136(10):718-22.
33. Indian J Dermatol Venereol Leprol 2009 Mar-Apr;75(2):177-9.
34. Int J Tuberc Lung Dis 2005 Jun ;9(6):686-8.
35. Int J Tuberc Lung Dis 2006 Jul ;10(7):761-7.
36. WHO AIDS Tech Bull 1989 Mar ;2:37.
37. J Acquir Immune Defic Syndr Hum Retrovirol 1998 Dec 15;19(5):484-9.
38. Clin Infect Dis 2008 Nov 1;47(9):1126-34.
39. ProMED <promedmail.org> archive: 20120206.1034331
40. Prev Vet Med 2005 Mar 15;67(4):267-81.
41. Epidemiol Infect 2008 May ;136(5):636-43.

Tungiasis

Agent	PARASITE - Insecta Siphonaptera (Flea), Tungidae: Tunga penetrans and T. trimamillata ("sand fleas")
Reservoir	Pig Dog ? Various other mammals
Vector	None
Vehicle	Contact
Incubation Period	8d - 12d
Diagnostic Tests	Identification of parasite.
Typical Adult Therapy	Extraction of parasite Ivermectin has been advocated in some publications.
Typical Pediatric Therapy	As for adult
Clinical Hints	Painful papule or nodule, usually on the feet - may be multiple; begins 1 to 2 weeks after walking on dry soil; secondary infections and tetanus are described.
Synonyms	Bicho de pe, Chica, Chigger, Chigoe flea, Jigger, Nigua, Puce-chique, Tu, Tunga penetrans, Tunga trimamillata, Tungosis. ICD9: 134.1 ICD10: B88.1

Clinical

Virtually all infestations are limited to the foot, notably the interdigital and periungual regions. ^{1 2}

- Ectopic infections are occasionally noted on the hands, elbows, thighs or gluteal region • and even the eyelids. ³
- Irritation begins 8 to 12 days following infection, and is manifested as a small 'pit' which evolves into a circular ulcer associated with pain, edema, erythema and pruritis.
- On dermoscopy, circumferential rings may be evident surrounding a central black lesion • the 'radial crown' sign. ⁴
- Secondary bacterial infection, thrombophlebitis or even tetanus may follow.
- Most infestations are characterized by 2 to 3 fleas, although hundreds may occasionally be present. ^{5 6}
- Severe disease may be characterized by deep ulcerations, necrosis leading to denudation of underlying bone, and auto-amputation of digits.
- Ectopic infection (hands, elbows, knees, neck ⁷, anus and genitals) is encountered, often in small children.
- Studies in an endemic region of Brazil revealed 17 lesions (maximum 98) per patient, and almost all had nail deformation and edema.
- Nail loss (46%), pain and fissures (70%), digit deformation (25%), abscesses (42%), and walking difficulty (59%) were common. (Brazil, 2007 publication) ⁸

A series of 11 cases of tetanus related to tungiasis (25% of all tetanus cases) was reported by a single hospital in Brazzaville over an 11-month period (1989 publication). ⁹

- Tungiasis is implicated in the etiology of 10% of tetanus cases in Sao Paulo, Brazil (2001 publication). ¹⁰

This disease is endemic or potentially endemic to 89 countries.

Tungiasis in Uganda

2011 (publication year) - A Finnish traveler acquired tungiasis in Uganda - the first case of tungiasis reported in the Nordic region. ¹¹

Notable outbreaks:

2010 - An outbreak (20,000 clinical cases, 20 fatal) of tungiasis was reported. ^{12 13}

References

1. *Parasitol Res* 2004 Oct ;94(4):275-282.
2. *J Am Acad Dermatol* 1989 May ;20(5 Pt 2):941-4.
3. *Ophthalmologica* 2007 ;221(6):439-42.
4. *Arch Dermatol* 2009 Mar ;145(3):348-9.
5. *Parasitol Res* 2007 Jan ;100(2):413-21.
6. *An Bras Dermatol* 2011 Sep-Oct;86(5):1027-8.

7. [Am J Trop Med Hyg 2010 Jun ;82\(6\):1076-8.](#)
8. [Rev Soc Bras Med Trop 2007 Jan-Feb;40\(1\):63-7.](#)
9. [Dakar Med 1989 ;34\(1-4\):44-8.](#)
10. [Braz J Infect Dis 2001 Dec ;5\(6\):319-23.](#)
11. [APMIS 2008 Mar ;116\(3\):219-22.](#)
12. [ProMED <promedmail.org> archive: 20101106.4023](#)
13. [ProMED <promedmail.org> archive: 20101107.4035](#)

Typhoid and enteric fever

Agent	BACTERIUM. <i>Salmonella serotype Typhi</i> (other <i>Salmonella</i> species cause 'paratyphoid' fever) A facultative gram-negative bacillus
Reservoir	Human
Vector	None
Vehicle	Fecal-oral Food, Fly Water
Incubation Period	15d - 21d (range 5d - 34d)
Diagnostic Tests	Culture (blood, urine, sputum culture). Stool usually negative unless late untreated infection). Serology.
Typical Adult Therapy	Ceftriaxone 2 g IV q12h to q 24h X 5 to 7d. OR Ciprofloxacin 750 mg PO (400 mg IV) Q12h X 2w. OR Azithromycin 1 gram PO on day 1; then 500 mg days 2 to 7. Add corticosteroids if evidence of shock or decreased mental status.
Typical Pediatric Therapy	Ceftriaxone 50 to 80 mg/kg IV daily X 5 to 7d. OR Azithromycin 15 mg/kg PO on day 1; then 7.5 mg/kg on days 2 to 7.
Vaccines	Typhoid - injectable Typhoid - oral
Clinical Hints	Transient diarrhea followed by fever, splenomegaly, obtundation, rose spots (during second week of illness); leukopenia and relative bradycardia often observed; case fatality rate = 0.8% (treated) to 15% (untreated).
Synonyms	Abdominal typhus, Abdominaltyphus, Buiktyphus, Enteric fever, Febbre tifoide, Febbre tifoidea, Fiebre tifoidea, Paratifoidea, Paratyfus, Paratyphoid, <i>Salmonella serotype Typhi</i> , Tyfoïd, Typhoid, Typhoïde. ICD9: 002 ICD10: A01

Clinical

Enteric fever is a defined syndrome of systemic illness associated with *Salmonella* infection.

- Enteric fever caused by *S. typhi* is referred to as "typhoid fever," and that caused by *S. paratyphi*, is referred to as "paratyphoid fever."
- Symptoms are often nonspecific and insidious in onset. ^{1 2}
- The differential diagnosis of fever, abdominal pain with hepatosplenomegaly also includes malaria, amebic liver abscess, brucellosis ³, visceral leishmaniasis, and dengue fever.
- The clinical features of scrub typhus ⁴ and melioidosis may also mimic those of enteric fever. ⁵

Acute illness:

Following an incubation period of 5 to 21 days, an initial enterocolitis may develop without associated fever.

- Constipation is present in 10 to 40% of patients; abdominal pain 20 to 40%; hepatosplenomegaly in 50%.
- Such symptoms as chills, diaphoresis, headache, anorexia, cough, sore throat, vertigo and myalgia often precede the onset of fever.
- Psychosis or confusion ("muttering delirium") occur in 5 to 10%, encephalopathy in 21% ⁶, and seizures and coma in less than 1%.
- Patients appear acutely ill.
- Cervical lymphadenopathy develops in some patients, and pulmonary disease is rare at this stage.
- 3% have signs and symptoms of cholecystitis, and jaundice is reported in as many as 12% of cases. ⁷
- Instances of "typhoid hepatitis" appear to represent super-infection by hepatitis virus, rather than a complication of typhoid fever. ⁸

Course of illness and complications:

Symptoms resolve by the fourth week of infection without antimicrobial therapy.

- Weight loss, and debilitation may persist for months, and 10% of patients will experience a relapse.
- Relapse is more common among antibiotic-treated than non-treated patients.
- Intestinal perforation is characterized by recurrent fever, abdominal pain, intestinal hemorrhage and tachycardia occurring in the 3rd to 4th week of illness. 65.7% of perforations are solitary and involved the anti-mesenteric border of the terminal ileum ⁹⁻¹². There is a male predominance among patients with typhoidal perforation. ¹³ During a typhoid fever outbreak in

Uganda, 43% of patients presented with intestinal perforation. ¹⁴

- 70% of pregnancies will end in miscarriage when complicated by untreated typhoid.
- Rare instances of acalculous cholecystitis ¹⁵⁻¹⁸, gall-bladder perforation ¹⁹, pancreatitis, intestinal intussusception ²⁰, rhabdomyositis, renal failure ²¹, genital ulceration ²², spondylitis/spondylodiscitis ²³, transverse myelitis ²⁴, cranial nerve palsy ²⁵, cerebral venous sinus thrombosis ²⁶, endophthalmitis ²⁷ and ectopic abscesses ^{28 29} have been reported in typhoid patients.
- The case-fatality rate is 10% to 15%

Carrier state:

The carrier state is defined as persistent shedding of *Salmonella typhi* in stool and/or urine for ≥ 12 months. ³⁰

- Approximately 5% of people who contract typhoid continue to carry the disease after they recover.


- Long-term carriage is associated with an increased incidence of cancers of the gallbladder ^{31 32}, pancreas, colo-rectum and lung. ^{33 34}

Laboratory findings include leukopenia (albeit an initial leucocytosis is common), thrombocytopenia, coagulopathy and hepatic dysfunction.

- The most sensitive laboratory test for enteric fever is blood culture.
- Serum transaminase elevations appear to reflect myopathy rather than hepatic disease in most cases. ³⁵

This disease is endemic or potentially endemic to all countries.

Typhoid and enteric fever in Uganda




Graph: Uganda. Typhoid and paratyphoid, cases

Notes:

1. During 1947 to 1949, the mean annual incidence was 359 cases ³⁶

5,871 cases of typhoid were reported in 1999.



Graph: Uganda. Typhoid, deaths

Prevalence surveys:

2.3% of febrile illness in the principal hospital in Kampala during 1992, 0.3% in 1995. ³⁷

Notable outbreaks:

2007 to 2009 - An outbreak (577 cases, 43% with intestinal perforation - 47 fatal) of typhoid fever was reported in Kasese. ³⁸⁻⁴⁰

References

1. Curr Gastroenterol Rep 2003 Aug ;5(4):279-86.
2. N Engl J Med 2002 Nov 28;347(22):1770-82.
3. J Infect Dev Ctries 2009 ;3(3):239-40.
4. Trans R Soc Trop Med Hyg 2010 Apr ;104(4):309-10.
5. Trans R Soc Trop Med Hyg 2008 Dec ;102 Suppl 1:S117-8.
6. Am J Trop Med Hyg 2012 Apr ;86(4):698-702.
7. Ann Afr Med 2010 Jul-Sep;9(3):135-40.
8. Trans R Soc Trop Med Hyg 1994 Jul-Aug;88(4):437-8.
9. Kathmandu Univ Med J (KUMJ) 2006 Jan-Mar;4(1):22-4.
10. J Infect Dev Ctries 2010 Oct ;4(10):650-4.
11. J Gastrointest Surg 2011 Oct ;15(10):1837-41.
12. Ann Afr Med 2011 Oct-Dec;10(4):259-65.
13. Clin Exp Gastroenterol 2012 ;5:213-7.
14. Clin Infect Dis 2012 Feb 22;
15. Singapore Med J 2006 Apr ;47(4):327-8.
16. J Natl Med Assoc 2009 Jul ;101(7):717-9.
17. Travel Med Infect Dis 2009 Jul ;7(4):203-6.
18. Scand J Infect Dis 2006 ;38(3):196-200.
19. Niger J Med 2011 Jan-Mar;20(1):181-3.
20. Acta Chir Belg 2011 May-Jun;111(3):174-5.
21. Int J Infect Dis 2009 Sep ;13(5):e282-5.
22. Am J Obstet Gynecol 2009 May ;200(5):e6-7.
23. J Med Assoc Thai 2010 Jan ;93(1):137-41.
24. Paediatr Int Child Health 2012 ;32(3):174-6.
25. Ann Trop Paediatr 2011 ;31(3):255-8.
26. Neurologist 2012 Jul ;18(4):202-3.
27. Semin Ophthalmol 2012 May-Jul;27(3-4):94-8.
28. J Gynecol Obstet Biol Reprod (Paris) 2007 Nov ;36(7):709-12.
29. J Glob Infect Dis 2011 Oct ;3(4):402-4.
30. Trends Microbiol 2012 May 14;
31. World J Gastroenterol 2010 Nov 21;16(43):5395-404.
32. Nat Rev Microbiol 2011 Jan ;9(1):9-14.
33. Lancet 1994 Jan 8;343(8889):83-4.
34. J Infect Dev Ctries 2010 May ;4(5):267-81.
35. Indian J Med Microbiol 2007 Oct ;25(4):351-3.
36. Bull World Health Organ 1955 ;13(1):173-91.
37. East Afr Med J 1998 Feb ;75(2):68-72.
38. Clin Infect Dis 2012 Feb 22;
39. ProMED <promedmail.org> archive: 20080814.2524
40. ProMED <promedmail.org> archive: 20081118.3640

Typhus - endemic

Agent	BACTERIUM. Rickettsia typhi
Reservoir	Rat
Vector	Flea (Xenopsylla or Nosopsyllus spp.)
Vehicle	None
Incubation Period	10d - 12d (range 4d - 18d)
Diagnostic Tests	Serology. Identification of rickettsiae in smear or culture of skin lesions. Nucleic acid amplification.
Typical Adult Therapy	Doxycycline 100 mg BID X 3 to 5d
Typical Pediatric Therapy	Doxycycline 2 mg/kg BID X 3 to 5d (maximum 200 mg/day); or Chloramphenicol 12.5 mg/kg QID X 3 to 5d
Clinical Hints	Fever, headache and myalgia; truncal maculopapular rash (present in 60%) appears on days 3 to 5 and persists for 4 to 8 days; fever resolves after 12 to 16 days; case fatality rate (untreated) = 2%.
Synonyms	Endemic typhus, Murine typhus, Rickettsia typhi, Ship typhus, Tifo murino, Tifus pulgas, Vlektyphus. ICD9: 081.0 ICD10: A75.2

Clinical

The features of endemic typhus are similar to those of epidemic typhus, but less severe. ¹

- Headache and myalgia predominate.
- The rash is nonspecific and may be lacking in 50% of patients. ²
- Major complications are rare.
- The severity of infection has been associated with old age, delayed diagnosis, hepatic and renal dysfunction, central nervous system abnormalities, and pulmonary compromise.
- Ocular complications include uveitis, retinal hemorrhage, choroidal dots, papilledema and optic neuritis ^{3 4}
- Rare instances of splenic infarction ⁵ and hemophagocytic syndrome have been reported. ⁶
- As many as 4% of hospitalized patients die.

This disease is endemic or potentially endemic to all countries.

References

1. Postgrad Med J 2000 May ;76(895):269-72.
2. Clin Dermatol 1996 May-Jun;14(3):271-8.
3. Br J Ophthalmol 2009 Jul ;93(7):938-42.
4. J Neuroophthalmol 2011 Dec ;31(4):342-3.
5. Am J Med Sci 2008 Jun ;335(6):502-3.
6. Am J Trop Med Hyg 2012 Jun ;86(6):1049-53.

Typhus - epidemic

Agent	BACTERIUM. Rickettsia prowazekii
Reservoir	Human ? Flying squirrel
Vector	Louse (Pediculus) ? Squirrel flea
Vehicle	None
Incubation Period	10d - 14d (range 5d - 23d)
Diagnostic Tests	Serology. Identification of rickettsiae in smear or culture of skin lesions. Nucleic acid amplification.
Typical Adult Therapy	Doxycycline 100 mg PO BID X 3 to 5d. OR Chloramphenicol 500 mg QID X 3 to 5d
Typical Pediatric Therapy	Doxycycline 2 mg/kg PO BID X 3 to 5d (maximum 200 mg/day). OR Chloramphenicol 10 mg/kg PO QID X 3 to 5d
Clinical Hints	Fever, headache and myalgia; truncal maculopapular rash appears on days 4 to 7; encephalopathy or myocarditis may ensue; fever resolves after 2 weeks, but convalescence is prolonged; case-fatality rate (untreated) = 10% to 20%.
Synonyms	Camp fever, Epidemic typhus, Jail fever, Red louse disease, Rickettsia prowazekii, Ship fever, Shop typhus, Sutama, Sylvatic epidemic typhus, Tifus piojos, Tobardillo. ICD9: 080 ICD10: A75.0

Clinical

Typhus is characterized by the sudden onset of headache, chills, prostration, vomiting, high fever, coughing, severe myalgia and shin pain. ¹

- Conjunctival suffusion, splenomegaly and cerebral obtundation are common.
- A macular eruption appears on the second to sixth day, initially on the upper trunk, with spread to the entire body except the face, palms and soles. ²
- The rash may later become petechial or necrotic.
- No rash is noted in as many as 30% of cases.

Meningoencephalitis occurs in as many as 50% of severe cases.


- Other complications include secondary bacterial infection, peripheral gangrene and myocarditis.
- The case-fatality rate among untreated patients is 10% to 15%.

Important clinical features which distinguish epidemic typhus from malaria and typhoid are lack of splenomegaly, rigors and diarrhea.

Recrudescence typhus (Brill-Zinsser disease) may present years after the initial episode, and is characterized by classic features of epidemic typhus with a milder clinical course.

This disease is endemic or potentially endemic to 63 countries.

Typhus - epidemic in Uganda



Graph: Uganda. Typhus - epidemic, cases

Notes:

1. Several cases of typhus were reported in 1959; 0 during 1960 to 1968.

Individual years:

1977 - Cases reported in East Cases and West Ankole. Two fatal cases reported.

1980 - None fatal.

1982 - 47 cases were reported during January to March.

Notable outbreaks:

1969 - An outbreak (46 cases) was reported in Masaka District. ³

1975 to 1976 - An outbreak was reported in Ankole. ⁴

References

1. Postgrad Med J 2000 May ;76(895):269-72.
2. Clin Dermatol 1996 May-Jun;14(3):271-8.

3. Trans R Soc Trop Med Hyg 1972 ;66(5):783-8.
4. East Afr Med J 1983 Jan ;60(1):55-9.

Urinary tract infection

Agent	BACTERIUM OR FUNGUS. Escherichia coli , other facultative gram negative bacilli, enterococci, et al
Reservoir	Human
Vector	None
Vehicle	Endogenous
Incubation Period	Variable
Diagnostic Tests	Urine culture and leucocyte count.
Typical Adult Therapy	Antimicrobial agent(s) directed at known or likely pathogen
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, dysuria, frequency, flank pain and vomiting; infection in children or men and infection which relapses in women may warrant radiological studies to rule out underlying obstruction or calculus.
Synonyms	Cistite, Cistitis, Cystite, Cystitis, Pielite, Pielitis, Pielonefrite, Pielonefritis, Prostatite, Pyelitis, Pyelonephrite, Pyelonephritis, Trigonitis, Tubulointerstitial nephritis, Urethritis, Uretrite, Zystitis. ICD9: 791.9,136.9,599.0,590,601.0 ICD10: N10,N30,N41

Clinical

Young children often exhibit nonspecific signs such as fever, poor feeding and vomiting.

- Abdominal pain may be present.
- After early childhood, dysuria, urgency, and frequency are generally present in UTI.
- Adult women with cystitis have frequent and urgency, often with lower abdominal or lower back pain.
- The urine may be foul smelling or turbid and is often bloody.
- Onset of symptoms is usually abrupt.
- Some infections progress to upper tract involvement, with fever, rigors, nausea, vomiting, abdominal and flank pain.
- Classical signs of 'upper' vs. 'lower' UTI are often misleading and do not necessarily point to the location of infection.

In the elderly, UTIs are often asymptomatic or manifest by nonspecific signs.

- Frequency, urgency, nocturia, and incontinence in this age group may also mimic other disorders in this age group.
- Infection associated with neurogenic bladders and indwelling catheters may not necessarily present with localizing symptoms.

Acute uncomplicated cystitis is most common in young women but may also be seen in men, children or the elderly. ¹

- Typical symptoms include dysuria, frequency, urgency, and suprapubic or pelvic pain. ²
- Suprapubic tenderness is present in 10 to 20 percent, and gross hematuria in 20 to 30 percent.
- Approximately ten percent of patients with symptoms of acute cystitis will be found to have occult infection of the upper urinary tract.
- Bacterial vaginosis may predispose to urinary tract infection ³

Acute pyelonephritis presents with flank, low back, or abdominal pain, in addition to fever, rigors, sweats, headache, nausea, vomiting, malaise, and prostration. ⁴

- Antecedent or concomitant symptoms of cystitis may or may not be present.
- Fever and flank pain are relatively specific indicators of renal infection.
- A minority of patients with pyelonephritis develop septicemia, or necrotizing renal or perinephric abscesses.
- The latter are often associated with urinary tract obstruction or diabetes [see Perinephric abscess].

All urinary infections in males should be considered complicated until proven otherwise, and prompt a careful search for anatomical or functional abnormality of the urinary tract.

Comprehensive reviews of prostatitis. ^{5 6}

This disease is endemic or potentially endemic to all countries.

References

1. Minerva Urol Nefrol 2004 Mar ;56(1):15-31.
2. N Engl J Med 2003 Jul 17;349(3):259-66.
3. J Obstet Gynaecol 2007 Apr ;27(3):252-4.
4. J Urol 2002 Dec ;168(6):2351-8.
5. Clin Microbiol Rev 1998 Oct ;11(4):604-13.
6. BMC Infect Dis 2008 ;8:12.

Varicella

Agent	VIRUS - DNA. Herpesviridae, Alphaherpesvirinae: Human Herpesvirus 3 (Varicella-zoster virus)
Reservoir	Human
Vector	None
Vehicle	Air Direct contact
Incubation Period	2w - 3w
Diagnostic Tests	Viral culture (vesicles). Serology. Nucleic acid amplification.
Typical Adult Therapy	Respiratory isolation. Severe/complicated cases: Acyclovir 10 to 12 mg/kg IV q8h X 7d Adolescent / young adult: 800 mg PO X 5 per day X 7 d. Alternatives: Valacyclovir 1 g PO TID; or Famciclovir 500 mg PO TID
Typical Pediatric Therapy	Respiratory isolation. Acyclovir [severe/complicated cases] 150 mg/sq m IV q8h X 7d
Vaccines	Varicella Varicella-Zoster immune globulin
Clinical Hints	Cough and fever followed by a pruritic papulovesicular rash after 1 to 2 days; pneumonia is often encountered; case fatality rate = 4.3 per 100,000 cases (7% in immune-suppressed patients).
Synonyms	Chickenpox, Lechina, Skoldkopper, Vannkopper, Varicela, Varizellen, Vattenkoppor, Waterpokken, Windpocken. ICD9: 052 ICD10: B01

Clinical

Acute infection:

The predominant features of varicella are fever, cough, malaise, lymphadenopathy and a generalized pruritic vesicular rash typically consisting of 250 to 500 lesions.

- The rash generally begins on the scalp and proceeds to the trunk and extremities, with most lesions on the trunk.
- Skin lesions are initially maculopapular, progressing to vesicles on an erythematous base. ¹
- Atypical varicella, including lesions on palms and soles, may mimic monkeypox in endemic areas. ²

Complications:

Complications include hepatitis ^{3 4}, encephalitis (notably involving the cerebellum) ⁵⁻⁷, myelitis ⁸, arthritis ⁹, secondary bacterial infections, Reye's syndrome, disorders of the facial ¹⁰⁻¹² and other cranial nerves ¹³, meningitis ¹⁴, cerebral venous thrombosis ¹⁵, transverse myelitis ¹⁶, acute urinary retention ¹⁷, pancreatitis ¹⁸, appendicitis ¹⁹, pneumonia ²⁰⁻²², empyema ²³, acute respiratory distress syndrome (ARDS) ²⁴⁻²⁷, spontaneous pneumothorax ²⁸, myocarditis ²⁹, atrioventricular block ³⁰, hemorrhagic pericarditis ^{31 32}, optic neuritis ³³⁻³⁵, uveitis ³⁶, acute retinal necrosis ^{37 38}, necrotizing scleritis ^{39 40}, deep venous thrombosis or thromboembolism ⁴¹, purpura fulminans ^{42 43}, idiopathic thrombocytopenic purpura ⁴⁴ and hemophagocytic lymphohistiocytosis. ⁴⁵

- Pyomyositis ⁴⁶, osteomyelitis ⁴⁷, necrotizing fasciitis or Fournier's gangrene may occasionally complicate varicella ⁴⁸
- Post varicella cerebral infarction has been described in young, previously healthy children within a few months of VZV infection and is characterized by middle cerebral artery territory infarction and proximal MCA disease. ⁴⁹ A similar condition has been reported in immunocompromised patients following herpes zoster involving the ophthalmic branch of the trigeminal nerve as well as in the context of primary varicella complicated by granulomatous angiitis ⁵⁰ Extra-cranial vascular thrombosis of large or small vessels has also been reported ⁵¹
- VZ virus infection may be associated with facial nerve palsy ⁵² or Ramsay-Hunt syndrome (Bell palsy unilateral or bilateral, vesicular eruptions on the ears, ear pain, dizziness, preauricular swelling, tingling, tearing, loss of taste sensation, and nystagmus) ⁵³
- Immunocompromised individuals, neonates, infants, adolescents and adults are at risk of severe illness and complications. ⁵⁴⁻⁵⁶
- VZ virus infection can be a presenting symptom of hyperparathyroidism and occurs twice as often in persons with hypercalcemia than age-matched controls. ⁵⁷
- Use of nonsteroidal anti-inflammatory drugs during primary varicella, has been implicated as a risk factor for subsequent occurrence of streptococcal necrotizing fasciitis.

Anterior uveitis • differential diagnosis:

Anterior uveitis due to Rubella virus is characterized by younger age at onset and a chronic course, typically associated with cataract at presentation. ⁵⁸

- Rubella virus has been implicated in the etiology of Fuchs heterochromic iridocyclitis. ⁵⁹
- Anterior uveitis due to Herpes simplex and Varicella-Zoster viruses is more common in adults, and often follows an acute course.
- Herpes simplex anterior uveitis presents with conjunctival redness, corneal edema, a history of keratitis, and the presence of posterior synechiae. Anterior chamber inflammation is common with Herpes simplex virus, while vitritis is more common with Rubella and Varicella-Zoster virus.
- Rubella, Herpes simplex and Varicella-zoster viruses are associated with intraocular pressure of more than 30 mmHg and development of glaucoma (18%-30%; P = 0.686).
- Focal chorioretinal scars were present in 22% of Rubella cases, 0% of HSV and in 11% of VZV uveitis cases.

Perinatal infection: ⁶⁰

Newborn infants whose mothers had onset of varicella within 5 days before delivery or within the 48 hours after delivery are at risk for neonatal varicella. ⁶¹⁻⁶⁶

- Neonatal varicella carries a case-fatality rate as high as 30%.
- Maternal infection ⁶⁷ during the first 20 weeks of pregnancy carries a risk (0.4% to 2.0%) of congenital varicella, characterized by low birth weight, hypoplasia of extremities, dermal scarring, focal muscular atrophy, encephalitis, cortical atrophy, chorioretinitis and microcephaly. ⁶⁸⁻⁷⁰

This disease is endemic or potentially endemic to all countries.

Varicella in Uganda**Notable outbreaks:**

2010 - An outbreak of varicella was reported in Bukuda, eastern region. ^{71 72}

References

1. Dermatol Clin 2002 Apr ;20(2):267-82.
2. Clin Infect Dis 2009 Jan 1;48(1):e6-8.
3. Liver Transpl 2008 Sep ;14(9):1309-12.
4. Bone Marrow Transplant 2009 Oct ;44(7):441-7.
5. Lancet Neurol 2007 Nov ;6(11):1015-28.
6. Curr Neurol Neurosci Rep 2009 Nov ;9(6):430-4.
7. Vaccine 2012 Aug 24;30(39):5785-90.
8. J Neurol Sci 2012 May 5;
9. Pediatr Infect Dis J 2011 Nov ;30(11):980-2.
10. Pediatr Int 2006 Jun ;48(3):245-9.
11. Enferm Infecc Microbiol Clin 2010 Oct ;28(8):504-8.
12. J Med Virol 2010 Sep ;82(9):1582-5.
13. Laryngoscope 2011 Aug ;121(8):1627-30.
14. Pediatr Infect Dis J 2008 Apr ;27(4):362-3.
15. J Stroke Cerebrovasc Dis 2012 Nov ;21(8):917.e1-4.
16. Acta Med Iran 2010 Nov-Dec;48(6):417-8.
17. S Afr Med J 2012 Apr ;102(4):196.
18. Saudi J Gastroenterol 2007 Jul-Sep;13(3):138-40.
19. Acta Medica (Hradec Kralove) 2012 ;55(3):150-2.
20. Eur Respir J 2003 May ;21(5):886-91.
21. Scand J Infect Dis 2010 Mar ;42(3):215-21.
22. Eur J Clin Microbiol Infect Dis 2011 Mar ;30(3):435-7.
23. Arch Pediatr 2008 Nov ;15(11):1643-7.
24. Med Sante Trop 2013 Jan 30;
25. Intern Med 2004 Dec ;43(12):1205-9.
26. J Formos Med Assoc 1999 Nov ;98(11):778-82.
27. Intensive Care Med 1985 ;11(6):319-22.
28. Acta Dermatovenerol Alp Panonica Adriat 2009 Jun ;18(2):71-2.
29. Pediatr Cardiol 2011 Dec ;32(8):1241-3.
30. Pediatr Infect Dis J 2011 May ;30(5):445-6.
31. Indian J Pathol Microbiol 2009 Apr-Jun;52(2):237-9.
32. Pediatr Cardiol 2010 Jul ;31(5):703-6.
33. Neurol Sci 2006 Sep ;27(4):278-80.
34. Pediatr Infect Dis J 2010 Dec ;29(12):1150-2.
35. Case Rep Ophthalmol Med 2012 ;2012:371584.
36. Medicine (Baltimore) 2008 May ;87(3):167-76.
37. J Fr Ophtalmol 2009 Jan ;32(1):60.e1-6.
38. Case Report Ophthalmol 2012 May ;3(2):180-4.
39. Ocul Immunol Inflamm 2006 Oct ;14(5):317-9.
40. Ophthalmologie 2008 May ;105(5):480-4.
41. Pediatr Infect Dis J 2012 Sep ;31(9):985-7.
42. J Nippon Med Sch 2009 Jun ;76(3):165-8.
43. Arch Pediatr 2011 Jul ;18(7):783-6.
44. Acta Paediatr 2010 Sep ;99(9):1385-8.
45. Pediatr Blood Cancer 2009 Aug ;53(2):226-8.
46. J Pediatr Orthop B 2011 Jul ;20(4):264-9.
47. Musculoskelet Surg 2012 Jun 17;
48. Pediatr Emerg Care 2007 Oct ;23(10):719-20.
49. Dev Med Child Neurol 2007 Jun ;49(6):417-22.
50. J Clin Neurosci 1998 Apr ;5(2):228-30.
51. Int J Infect Dis 2009 Nov ;13(6):e498-500.
52. Pediatr Int 2006 Jun ;48(3):245-9.
53. J Craniofac Surg 2011 Sep ;22(5):1961-3.
54. Lancet 2006 Oct 14;368(9544):1365-76.
55. Pediatr Infect Dis J 2008 Oct ;27(10):946-8.
56. Bone Marrow Transplant 2009 Oct ;44(7):441-7.
57. Clin Infect Dis 2008 May 1;46(9):1452-4.
58. Ophthalmology 2011 Oct ;118(10):1905-10.
59. Graefes Arch Clin Exp Ophthalmol 2010 Oct ;248(10):1487-91.
60. J Obstet Gynaecol Can 2012 Mar ;34(3):287-92.
61. Australas Med J 2011 ;4(6):291-3.
62. BMJ Case Rep 2012 ;2012
63. Arch Dis Child 2011 May ;96(5):453-6.
64. Indian Pediatr 2010 Feb ;47(2):181.
65. Arch Pediatr Adolesc Med 2009 May ;163(5):481-2.
66. Semin Fetal Neonatal Med 2009 Aug ;14(4):209-17.
67. N Engl J Med 1966 Apr 7;274(14):768-71.
68. Semin Fetal Neonatal Med 2009 Aug ;14(4):209-17.
69. Indian J Dermatol Venereol Leprol 2010 Nov-Dec;76(6):724.
70. BJOG 2011 Sep ;118(10):1155-62.
71. ProMED <promedmail.org> archive: 20100325.0953
72. ProMED <promedmail.org> archive: 20100324.0947

Vibrio parahaemolyticus infection

Agent	BACTERIUM Vibrio parahaemolyticus A facultative gram-negative bacillus
Reservoir	Marine water Seafood Fish
Vector	None
Vehicle	Seafood
Incubation Period	10h - 20h (range 2h - 4d)
Diagnostic Tests	Stool culture - alert laboratory when this organism is suspected.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Vomiting and explosive diarrhea, 4 to 24 hours following ingestion of seafood (often steamed crabs); diarrhea may persist for 7 to 10 days; case fatality rate = 0.1%.
Synonyms	Vibrio parahaemolyticus. ICD9: 005.4 ICD10: A05.3

Clinical

Symptoms usually begin within 10 to 20 hours after ingestion of seafood, and persist for 2 to 10 days.

- Illness is characterized by vomiting (50%), abdominal pain and watery or explosive diarrhea.
- Fever is noted in 25% of patients.
- Dysentery has been described in some cases. ¹

Rare instances of bacteremia and extra-intestinal infection are reported. ²⁻⁴

This disease is endemic or potentially endemic to all countries.

References

1. J Infect Dis 2002 Dec 1;186(11):1615-20.
2. N Z Med J 2008 Oct 3;121(1283):99-101.
3. Rev Chilena Infectol 2009 Aug ;26(4):360-2.
4. J Med Microbiol 2010 Feb ;59(Pt 2):235-8.

Wesselsbron

Agent	VIRUS - RNA. Flaviridae, Flavivirus: Wesselsbron virus
Reservoir	Sheep Cattle
Vector	Mosquito (Aedes sp., Anopheles gambiae, An. pharoensis, Culex telesilla, Cu. univittatus, Mansonia uniformis)
Vehicle	None
Incubation Period	2d - 4d
Diagnostic Tests	Viral culture (blood, throat). Serology. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Fever, myalgia, arthralgia, dermal hyperesthesia, maculopapular rash and leukopenia; illness lasts up to 10 days; no fatality reported.
Synonyms	

Clinical

Wesselsbron is characterized by abrupt onset of fever, myalgias, arthralgias, dermal hyperesthesia, leucopenia.

- A maculopapular rash which appears after 3 to 4 days. ¹
- Meningoencephalitis may follow.
- Although human infection may be severe, no fatalities have been reported.

This disease is endemic or potentially endemic to 31 countries.

Wesselsbron in Uganda

Seropositive humans have been identified in West Nile A, West Nile B, Madi and Bwamba (1967 to 1969). ²

References

1. S Afr Med J 1958 May 24;32(21):543-5.
2. Bull World Health Organ 1970 ;42(5):797-805.

West Nile fever

Agent	VIRUS - RNA. Flaviridae, Flavivirus: West Nile virus
Reservoir	Bird Horse Bat ? Tick
Vector	Mosquito (Culex univittatus. Cu. pipiens, Cu. vishnui, Cu. neavei, Coquillettidia, Aedes and Anopheles spp.)
Vehicle	Blood transmission [rare]
Incubation Period	3d - 6d (range 1d - 7d)
Diagnostic Tests	Viral culture (blood, CSF). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Myalgia, arthralgia, lymphadenopathy, headache, conjunctivitis and a macular rash; sporadic instances of encephalitis, meningitis and myocarditis are reported; illness resolves within one week in most cases.
Synonyms	Bagaza, Fiebre del Oeste del Nilo, Lourdige, Near Eastern equine encephalitis, Ntaya, Usutu, WNF. ICD9: 066.4 ICD10: A92.3

Clinical

Acute infection:

West Nile fever in humans usually is a minor influenza-like illness, characterized by an abrupt onset of moderate to high fever lasting 3 to 5 days.

- The fever is occasionally biphasic, and may be accompanied by rigors.
- Additional findings include frontal headache, sore throat, backache, myalgia, arthralgia, fatigue, conjunctivitis and retrobulbar pain. ¹
- A maculopapular or roseolar rash ^{2 3} appears in approximately 50% of cases, spreading from the trunk to the extremities and head.
- Lymphadenopathy, anorexia, nausea, abdominal pain, diarrhea, and respiratory symptoms are also encountered.

Neuroinvasive disease:

Occasionally (<15% of cases), acute aseptic meningitis or encephalitis occurs, associated with neck stiffness, vomiting, confusion, disturbed consciousness, somnolence, tremor of extremities, abnormal reflexes, convulsions, pareses, and coma. ^{4 5}

- Such patients may then develop anterior myelitis and acute asymmetric flaccid paralysis, reminiscent of poliomyelitis or Guillain-Barre syndrome. ⁶⁻⁹ Rare instances of progressive brachial diplegia are reported. ¹⁰
- Focal encephalitis with seizures may mimic herpes simplex encephalitis. ^{11 12}
- A case of stuttering (? lingual myoclonus) during the course of West Nile fever has been reported. ¹³
- Risk factors for neuroinvasive disease include age >45 years, male sex, hypertension and diabetes mellitus. ¹⁴
- Multifocal chorioretinitis is common among patients with neuroinvasive disease. ¹⁵⁻²⁰
- Most fatal cases occur in patients older than 50 years. ²¹

Hepatosplenomegaly, hepatitis, pancreatitis ²², myocarditis ²³ and hemorrhagic fever have been reported. ²⁴

Prolonged convalescence (up to one year) may follow recovery from encephalitis; and myalgia, confusion and lightheadedness may persist beyond this period. ²⁵⁻²⁹

- Recovery is complete (less rapid in adults than in children, often accompanied by long-term myalgias and weakness), and permanent sequelae have not been reported.
- Prolonged depression persists in as many as 31% of patients following recovery. ^{30 31}
- Residual kidney disease is common following West Nile fever. ³²

Laboratory findings:

Laboratory findings consist of a slightly increased sedimentation rate and mild leukocytosis.

- Profound and prolonged lymphocytopenia is reported in some cases. ³³
- Cerebrospinal fluid in patients with central nervous system involvement is clear, with moderate pleocytosis and elevated

protein.

- A distinctive CSF plasmacytosis may be present.
- The virus can be recovered from the blood for up to 10 days in immunocompetent febrile patients, and as late as 22 to 28 days after infection in immunocompromised patients.
- Peak viremia occurs 4 to 8 days postinfection.

Usutu virus infection was reported in two immune-compromised humans in 2009. ³⁴⁻³⁶

- Clinical features included fever of 39.5 °C, headache and neurological findings. One patient developed fulminant hepatitis.

Ntaya virus, a related flavivirus, has been associated with febrile illness and neurological findings. ³⁷

This disease is endemic or potentially endemic to 86 countries.

West Nile fever in Uganda

West Nile fever is named after the West Nile district of Uganda, where the agent was first described (1937). ³⁸

- The virus has been found in local mosquitoes (*Coquillettidia metallica*). ³⁹

Zoonotic infection by Usutu virus (a related flavivirus) has been identified in this country. ⁴⁰

A third flavivirus, Ntaya virus, was first isolated from mosquitoes in Uganda. ^{41 42}

Seroprevalence surveys:

11.0% of the general population (1967 to 1969, eight areas) ⁴³

References

1. Mayo Clin Proc 2003 Sep ;78(9):1137-43; quiz 1144.
2. Clin Infect Dis 2005 Oct 15;41(8):1204-7.
3. Dermatology 2005 ;211(4):348-50.
4. Curr Opin Infect Dis 2004 Oct ;17(5):413-20.
5. Curr Opin Neurol 2004 Jun ;17(3):343-6.
6. Curr Opin Infect Dis 2004 Oct ;17(5):413-20.
7. Emerg Infect Dis 2005 Jul ;11(7):1021-7.
8. Neurology 2000 Jul 12;55(1):144-6.
9. Front Neurol 2012 ;3:37.
10. Muscle Nerve 2012 Jun ;45(6):900-4.
11. Pediatr Neurol 2006 Jul ;35(1):62-4.
12. Clin Infect Dis 2003 Dec 1;37(11):1573-8.
13. Emerg Infect Dis 2011 Aug ;17(8):1567-8.
14. Emerg Infect Dis 2007 Dec ;13(12):1918-20.
15. Mayo Clin Proc 2006 Jan ;81(1):12-6.
16. Ocul Immunol Inflamm 2007 Nov-Dec;15(6):435-9.
17. Ophthalmic Surg Lasers Imaging 2006 May-Jun;37(3):240-2.
18. Eye (Lond) 2007 Jul ;21(7):952-5.
19. Ophthalmology 2004 Nov ;111(11):2065-70.
20. Ophthalmology 2003 Sep ;110(9):1732-6.
21. J Am Geriatr Soc 2002 Nov ;50(11):1844-6.
22. Travel Med Infect Dis 2008 Nov ;6(6):373-5.
23. Am J Trop Med Hyg 2006 Dec ;75(6):1232-3.
24. Clin Infect Dis 2006 Jun 1;42(11):1527-35.
25. Emerg Infect Dis 2006 Aug ;12(8):1260-2.
26. Ann Intern Med 2008 Aug 19;149(4):232-41.
27. J Neuropsychol 2008 Sep ;2(Pt 2):477-99.
28. Emerg Infect Dis 2004 Aug ;10(8):1405-11.
29. Front Neurol 2012 ;3:37.
30. Emerg Infect Dis 2007 Mar ;13(3):479-81.
31. S D Med 2010 Apr ;63(4):127-9, 131-3.
32. ProMED <promedmail.org> archive: 20120714.1202043
33. Clin Infect Dis 2000 Oct ;31(4):1116-7.
34. Euro Surveill 2009 ;14(50)
35. Euro Surveill 2009 ;14(50)
36. ProMED <promedmail.org> archive: 20091217.4273
37. Br Med J 1978 Apr 15;1(6118):956-8.
38. Curr Infect Dis Rep 2005 Jul ;7(4):292-296.
39. Ann Trop Med Parasitol 1961 Dec ;55:398-402.
40. Ann Trop Med Parasitol 1964 Sep ;58:367-74.
41. Proc Soc Exp Biol Med 1951 May ;77(1):130-3.
42. Trans R Soc Trop Med Hyg 1953 Jan ;47(1):13-48.
43. Bull World Health Organ 1970 ;42(5):797-805.

Whipple's disease

Agent	BACTERIUM. Actinomycetes, <i>Tropheryma whipplei</i> A gram positive bacillus
Reservoir	Unknown
Vector	None
Vehicle	None
Incubation Period	Unknown
Diagnostic Tests	Identification of inclusions in lamina propria (other tissues). Tissue culture. Nucleic acid amplification.
Typical Adult Therapy	Ceftriaxone 2.0 g IV daily X 14 days. OR Penicillin G 6 to 324 million units daily + + streptomycin 1 g daily X 14d. Then: Sulfamethoxazole/trimethoprim 800/160 mg PO BID X 1 year. OR Doxycycline 100 mg PO BID X 1 year
Typical Pediatric Therapy	This disease is not described in children
Clinical Hints	A chronic multisystem disorder characterized by weight loss, diarrhea, abdominal and joint pain; dermal hyperpigmentation, fever and lymphadenopathy often present; PAS-positive macrophages present in intestinal biopsy material.
Synonyms	Intestinal lipodystrophy, Lipophagic granulomatosis, Mesenteric chyladenectasis, Steatorrhea arthropericarditica, <i>Tropheryma whipplei</i> . ICD9: 040.2 ICD10: K90.8

Clinical

The typical patient with Whipple's disease has a history of recurrent arthralgia or arthritis involving multiple joints for several years.¹

- Joint complaints precede systemic and gastrointestinal disease in approximately one-third of patients², and may persist for years in the absence of diarrhea.³
- Infection of prosthetic joints has been reported.⁴
- Diarrhea, low-grade fever and weight loss are characteristic, and hyperpigmentation is present in 50% of patients.
- Generalized lymphadenopathy is common.
- A syndrome of dementia and obesity or ataxia linked associated with *T. whipplei* infection has been recently described.⁵

As many as one third of the patients develop cardiac involvement characterized by the presence of systolic murmurs, a pericardial friction rub, congestive heart failure, and nonspecific electrocardiographic changes.⁶

- The most common pathological changes are endocarditis⁷⁻⁹ with negative blood cultures, presenting with thickened and deformed mitral or aortic valves.¹⁰
- In one series, Whipple's disease was identified in 6.3% of patients with culture-negative endocarditis (2011 publication)¹¹
- 30 to 40% of patients develop pleuritic chest pain, chronic nonproductive cough, and dyspnea.
- The chest X-ray may show a pleural effusion or pulmonary infiltrates.
- Isolated *T. whipplei* endocarditis may occur without other systemic features of Whipple's disease.^{12 13}

Relapse of Whipple's disease has been reported following therapy.¹⁴

- Recurrence of symptoms following therapy may represent an immune reconstitution syndrome.¹⁵

Tropheryma whipplei was isolated from 6.4% of blood specimens from febrile patients with cough (Senegal, 2008 to 2009)¹⁶

Other features of Whipple's disease include personality changes or dementia¹⁷⁻¹⁹, hypersomnia²⁰, amnesic syndrome²¹, peripheral or cranial nerve neuropathy²², encephalitis^{23 24}, cerebral pseudotumor^{25 26}, chronic headache²⁷, endocarditis²⁸⁻³¹, pericarditis³², pneumonia^{33 34}, pulmonary hypertension^{35 36}, subcutaneous nodules, anemia, myoclonus, ataxia³⁷, chorioretinitis³⁸, vitritis³⁹, uveitis^{40 41}, Parinaud syndrome⁴², sacroiliitis⁴³ and spondylitis⁴⁴, hypoalbuminemia and hypokalemia.⁴⁵

The features of Whipple's disease may resemble those of lymphoma, celiac disease, Crohn's vasculitis, sepsis, an inflammatory process, liposarcoma, rheumatoid arthritis, seizure disorder, cerebrovascular accident, xanthoma, or central

nervous system neoplasm. ⁴⁶

This disease is endemic or potentially endemic to all countries.

References

1. Lancet Infect Dis 2008 Mar ;8(3):179-90.
2. Clin Gastroenterol Hepatol 2004 Oct ;2(10):849-60.
3. J Clin Microbiol 2009 Feb ;47(2):492-5.
4. J Clin Microbiol 2008 Apr ;46(4):1556-7.
5. BMC Infect Dis 2011 ;11:171.
6. Medicine (Baltimore) 2010 Sep ;89(5):337-45.
7. Rev Med Interne 2005 Oct ;26(10):784-90.
8. Ann Cardiol Angeiol (Paris) 2012 Feb ;61(1):61-3.
9. BMC Res Notes 2012 ;5:600.
10. Eur J Echocardiogr 2008 May ;9(3):426-7.
11. J Clin Microbiol 2012 Feb ;50(2):216-22.
12. Int J Infect Dis 2011 Nov ;15(11):e804-6.
13. Interact Cardiovasc Thorac Surg 2012 Apr 11;
14. Intern Med 2012 ;51(15):2045-50.
15. Ann Intern Med 2010 Dec 7;153(11):710-7.
16. Clin Infect Dis 2010 Sep 1;51(5):515-21.
17. Clin Neurol Neurosurg 2008 Jul ;110(7):747-9.
18. Pract Neurol 2008 Oct ;8(5):311-7.
19. Dtsch Med Wochenschr 2011 Jun ;136(24):1312-5.
20. Arq Neuropsiquiatr 2006 Sep ;64(3B):865-8.
21. Cogn Behav Neurol 2010 Mar ;23(1):49-51.
22. Curr Infect Dis Rep 2006 Jun ;8(4):301-6.
23. Rev Med Interne 2011 Aug ;32(8):513-6.
24. Arch Neurol 2011 Nov ;68(11):1471-3.
25. Acta Neurochir (Wien) 2009 Feb ;151(2):173-5.
26. J Neurol Sci 2011 Sep 15;308(1-2):1-8.
27. J Gen Intern Med 2008 Dec ;23(12):2131-3.
28. J Clin Microbiol 2007 Jun ;45(6):2078-81.
29. J Med Case Rep 2010 ;4:245.
30. Am J Med 2010 Oct ;123(10):962.e1-4.
31. Clin Microbiol Infect 2010 Aug ;16(8):1213-22.
32. Can J Cardiol 2009 Mar ;25(3):e89-91.
33. Emerg Infect Dis 2010 Feb ;16(2):258-63.
34. Chest 2012 Jun ;141(6):1595-8.
35. Can Respir J 2011 Sep-Oct;18(5):e70-2.
36. Case Rep Pulmonol 2012 ;2012:382460.
37. J Neurol Sci 2010 Oct 15;297(1-2):97-100.
38. Semin Arthritis Rheum 2009 Apr ;38(5):403-6.
39. Transpl Infect Dis 2008 Dec ;10(6):413-8.
40. Ophthalmologe 2008 Nov ;105(11):1046, 1048-51.
41. Medicine (Baltimore) 2008 May ;87(3):167-76.
42. Ophthalmology 2012 Mar 13;
43. Acta Clin Belg 2008 Mar-Apr;63(2):107-11.
44. Orphanet J Rare Dis 2009 ;4:13.
45. Postgrad Med J 2000 Dec ;76(902):760-6.
46. Am J Surg Pathol 2012 Jul ;36(7):1066-73.

Yaws

Agent	BACTERIUM. <i>Treponema pallidum</i> subsp. pertenue: microaerophilic gram-negative spirochete
Reservoir	Human ? Non-human primate
Vector	None
Vehicle	Contact ? Insect bite ? Fomite
Incubation Period	3w - 5w (range 10d - 12w)
Diagnostic Tests	VDRL and antitreponemal tests (FTA, MHTP) positive as in syphilis.
Typical Adult Therapy	Benzathine <i>Penicillin G</i> 1.2 million units IM as single dose. A single oral dose of <i>Azithromycin</i> is also effective.
Typical Pediatric Therapy	Benzathine <i>Penicillin G</i> : Weight <14kg: 300,000u IM Weight 14 to 28kg: 600,000u IM Weight >28kg - 1.2 million u IM A single oral dose (30 mg/kg) of <i>Azithromycin</i> is also effective.
Clinical Hints	Dermal papillomata, periostitis and soft tissue suppuration; regional lymphadenopathy common; relapses often seen during initial 5 years of illness; gummas and hyperkeratotic plaques in later stages.
Synonyms	Anakhre, Bouba, Breda's disease, Charlouis' Disease, Frambesia, Gangosa, Goundou, Granuloma tropicum, Gundo, Henpue, Henpuye, Ogo Mutilans, Parangi, Patek, Pian, <i>Treponema pallidum</i> subsp. pertenue. ICD9: 102 ICD10: A66

Clinical

Yaws has three clinical stages. ^{1 2}

- Stage 1 is characterized by the a variety of flat and/or raised skin lesions.
- Stage 2 (Gangosa Syndrome, Ogo, or Rhinopharyngitis Mutilans) may involve the bones, joints, and/or skin.
- Stage 3 (Goundou Syndrome, Henpue, Henpuye, Gundo, or Anakhre) may also involve the bones, joints, and/or skin.


After an incubation period of approximately 3 weeks, a primary painless 2 to 5 cm pruritic papule ("mother yaw") appears at the site of inoculation. ³

- The lesions may ulcerate, but generally heal completely after 3 to 6 months.
- Secondary lesions appear in crops from weeks to months later, measure 1 to 5 cm and tend to ulcerate or take the shape of raspberries (frambesoids), round or discoidal papillomas. ⁴
- Osteoperiostitis may be evident at this stage ⁵ ; however systemic symptoms are usually not present
- The secondary stage may persist for up to 6 months, and relapse over periods as long as 10 years. ⁶
- The third stage is characterized by destructive necrotic and gummatous lesions of skin, bone, nasopharynx and contiguous structures.

This disease is endemic or potentially endemic to 67 countries.

Yaws in Uganda

Yaws occurs in all areas of the country, with highest incidence in the northwestern and southwestern regions.



Graph: Uganda. Yaws, cases

Notes:

1. Historical data from reference ⁷

References

<ol style="list-style-type: none"> 1. J Am Acad Dermatol 1993 Oct ;29(4):519-35; quiz 536-8. 2. Bull World Health Organ 1951 ;4(2):201-46. 3. Microbes Infect 2002 Jan ;4(1):83-94. 4. Clin Dermatol 2006 May-Jun;24(3):181-90. 	<ol style="list-style-type: none"> 5. Clin Infect Dis 2011 Mar 15;52(6):771-4. 6. Bull World Health Organ 1956 ;15(6):869-96. 7. Bull World Health Organ 1953 ;8(1-3):129-82; discussion 205-10.
---	---

Yellow fever

Agent	VIRUS - RNA. Flaviridae, Flavivirus: Yellow fever virus
Reservoir	Human Mosquito Monkey ? Marsupial
Vector	Mosquito - Stegomyia (Aedes), Haemagogus, Sabethes
Vehicle	None
Incubation Period	3d - 6d (range 2.5d - 14d)
Diagnostic Tests	Viral culture (blood, liver). Serology. Nucleic acid amplification. Biosafety level 3.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Vaccine	Yellow fever
Clinical Hints	Headache, backache, vomiting, myalgias, jaundice, hemorrhagic diathesis, relative bradycardia and leukopenia; illness is often biphasic; 10% to 60% die within 7 days of onset.
Synonyms	Bulan fever, Febbre gialla, Febre amarela, Fever of Fernando Po, Fever of the blight of Benin, Fiebre amarilla, Fievre jaune, Gelbfieber, Gele koorts, Gul feber, Gula febern, Inflammatory fever, Kendal's disease, Magdalena fever, Maladie de Siam, Pest of Havana, Stranger's fever. ICD9: 060 ICD10: A95

Clinical

WHO Case definition for surveillance: ¹

Clinical description

- Characterized by acute onset of fever followed by jaundice within 2 weeks of onset of first symptoms.
- Hemorrhagic manifestations and signs of renal failure may occur.

Laboratory criteria for diagnosis

- Isolation of yellow fever virus, or
- Presence of yellow fever specific IgM or a four-fold or greater rise in serum IgG levels in paired sera (acute and convalescent) or
- Positive post-mortem liver histopathology or detection of yellow fever antigen in tissues by immunohistochemistry or
- Detection of yellow fever virus genomic sequences in blood or organs by PCR

Case classification

- Suspected: A case that is compatible with the clinical description.
- Probable: A suspected case with presence of yellow fever IgM antibody (in the absence of vaccination within 30 days); or positive postmortem liver histopathology; or an epidemiological link to a confirmed case or outbreak.
- Confirmed: A probable case; and a fourfold or greater increased in antibody titers; or presence of yellow fever neutralization antibody; or detection of yellow fever virus, viral genome or antigen in blood or tissues.

The clinical presentation of yellow fever can range from a self-limited flu-like illness to overwhelming hemorrhagic fever, with a case-fatality rate of 50%. ²

- As many as 50% of infections may be clinically inapparent.


Infection is heralded by abrupt onset of fever, headache, and myalgias associated with conjunctival injection, facial flushing, relative bradycardia (Faget's sign) and leukopenia. ³

- Although most cases do not progress beyond this stage, a remission of fever for a few hours to several days may be followed by high fever, headache, lumbosacral pain, nausea, vomiting, abdominal pain, and somnolence.
- At this stage, the patient exhibits icteric hepatitis and a hemorrhagic diathesis with prominent bleeding from the gastrointestinal tract, epistaxis, bleeding gums, and petechial and purpuric hemorrhages.
- Weakness, prostration, protracted vomiting and albuminuria are prominent.
- Deepening jaundice and elevations in serum transaminase levels continue for several days, accompanied by azotemia and progressive oliguria.
- Direct bilirubin levels rise to 5 to 10 mg/dl, while alkaline phosphatase levels are only slightly raised.
- Eventually, hypotension, shock, and metabolic acidosis develop, compounded by myocardial dysfunction and arrhythmias.
- Additional findings may include acute tubular necrosis, confusion, seizures, and coma.
- CSF examination reveals an elevated protein level without pleocytosis.
- Death usually occurs within 7 to 10 days after onset.

This disease is endemic or potentially endemic to 47 countries.

Yellow fever in Uganda

Uganda is designated a COUNTRY WITH RISK OF YELLOW FEVER VIRUS TRANSMISSION.




Graph: Uganda. Yellow fever, cases

Notes:

1. Historical data from reference ⁴

Individual years:

1964 - See references ⁵ ⁶



Graph: Uganda. Yellow fever, deaths

Proof of vaccination **IS** required if traveling from a country with risk of YFV transmission and ≥ 1 year of age - CDC (The United States Centers for Disease Control) recommends vaccination for all travelers ≥ 9 months of age (arriving from any country).

Notable outbreaks

- 1949 (publication year) - An outbreak of sylvan yellow fever (vector *Aedes (Stegomyia) africanus*) was reported. ⁷
- 2010 - An outbreak (226 cases, 53 fatal) of yellow fever was reported in the northern region. ⁸⁻²⁷

References

1. Wkly Epidemiol Rec 2010 Nov 19;85(47):465-72.
2. Clin Lab Med 2002 Dec ;22(4):981-1020, viii.
3. Lancet Infect Dis 2001 Aug ;1(1):11-20.
4. Bull World Health Organ 1954 ;11(3):325-89.
5. Trans R Soc Trop Med Hyg 1965 Jul ;59:436-40.
6. Trans R Soc Trop Med Hyg 1965 Jul ;59:441-3.
7. Ann Trop Med Parasitol 1949 Apr ;43(1):74-89.
8. Wkly Epidemiol Rec 2011 Jan 28;86(5):37-8.
9. ProMED <promedmail.org> archive: 20101129.4297
10. Virology 2012 Jan 5;422(1):1-5.
11. Int J Infect Dis 2012 May 8;
12. ProMED <promedmail.org> archive: 20101130.4302
13. ProMED <promedmail.org> archive: 20101202.4334
14. ProMED <promedmail.org> archive: 20101203.4340
15. ProMED <promedmail.org> archive: 20101205.4354
16. ProMED <promedmail.org> archive: 20101208.4382
17. ProMED <promedmail.org> archive: 20101224.4531
18. ProMED <promedmail.org> archive: 20101228.4572
19. ProMED <promedmail.org> archive: 20101229.4588
20. ProMED <promedmail.org> archive: 20101230.4595
21. ProMED <promedmail.org> archive: 20110102.0017
22. ProMED <promedmail.org> archive: 20110103.0027
23. ProMED <promedmail.org> archive: 20110106.0074
24. ProMED <promedmail.org> archive: 20110111.0123
25. ProMED <promedmail.org> archive: 20110118.0214
26. ProMED <promedmail.org> archive: 20110119.0233
27. ProMED <promedmail.org> archive: 20110202.0387

Yersiniosis

Agent	BACTERIUM. <i>Yersinia enterocolitica</i> and <i>Yersinia pseudotuberculosis</i> A facultative gram-negative bacillus
Reservoir	Pig Rodent Rabbit Sheep Goat Cattle Horse Dog Cat Bat
Vector	None
Vehicle	Food Water Meat Dairy products Vegetables Fecal-oral Blood
Incubation Period	4d - 7d (range 1d - 11d)
Diagnostic Tests	Culture stool, blood. Alert laboratory when these organisms are suspected.
Typical Adult Therapy	Stool precautions; diarrhea is self-limited. If severe disease - Ciprofloxacin 500 mg BID X 5 to 7d. OR Sulfamethoxazole/trimethoprim
Typical Pediatric Therapy	Stool precautions; diarrhea is self-limited. If severe disease - Sulfamethoxazole/trimethoprim 20 mg-4 mg/kg BID X 5 to 7d
Clinical Hints	Fever, diarrhea, right lower quadrant pain; fecal leucocytes present; may be associated with rheumatologic manifestations such as erythema multiforme, Reiter's syndrome and chronic arthritis.
Synonyms	<i>Yersinia enterocolitica</i> , <i>Yersinia pseudotuberculosis</i> , Yersiniose. ICD9: 008.44 ICD10: A04.6,A28.2

Clinical

Yersinia enterocolitica infection typically presents as febrile diarrhea, and occasionally bloody diarrhea.

- Lower abdominal pain without diarrhea occurs in over 15% of cases, and may mimic acute appendicitis. ¹⁻⁵ Several instances of intestinal intussusception have been associated with *Yersinia enterocolitica* ⁶⁻¹⁵ and *Y. pseudotuberculosis* infections. ¹⁶
- Pharyngitis ¹⁷ and tonsillitis ¹⁸ are encountered ; and metastatic infection of bone, soft tissues, spleen, meninges ¹⁹ or other organs may occur. ²⁰⁻²³
- Chronic arthritis, erythema nodosum, Reiter's syndrome ²⁴ , Sweet's syndrome ²⁵ , glomerulonephritis ²⁶⁻³² , pneumonia ³³ and endocarditis ³⁴⁻⁴⁷ are also reported.
- Reactive arthritis has been reported in over 20% of cases ^{48 49}
- *Yersinia enterocolitica* septicemia (associated with transfusion of contaminated red blood cell products) is fatal in over 50% of cases. ⁵⁰

Yersinia enterocolitica is one of at least a dozen *Yersinia* species encountered in humans. See the Microbiology module for further details.

This disease is endemic or potentially endemic to all countries.

References

1. *Pediatr Infect Dis J* 2000 Oct ;19(10):954-8.
2. *Infect Dis Clin North Am* 1988 Sep ;2(3):625-41.
3. *Emerg Radiol* 2008 Mar ;15(2):123-6.
4. *Pediatr Surg Int* 1998 Jan ;13(1):2-5.
5. *Acta Paediatr* 2011 Jul ;100(7):1028-32.
6. *Intern Med* 2012 ;51(18):2545-9.
7. *Med Welt* 1976 May 21;27(21):1063-8.
8. *Am J Dis Child* 1983 Aug ;137(8):803-4.
9. *Pediatr Med Chir* 1984 Sep-Oct;6(5):667-8.
10. *J Pediatr Surg* 1992 Dec ;27(12):1591-2.
11. *Presse Med* 1996 Apr 20;25(14):668-70.
12. *J Pediatr Gastroenterol Nutr* 1996 Jul ;23(1):77-80.
13. *Klin Padiatr* 1998 Mar-Apr;210(2):61-4.
14. *Arch Pathol Lab Med* 2001 Nov ;125(11):1486-8.
15. *Clin Med Insights Pathol* 2010 ;3:7-11.
16. *Acta Paediatr* 1996 Oct ;85(10):1253-5.
17. *Am J Med* 1987 Mar 23;82(3 Spec No):636-7.
18. *Acta Otorrinolaringol Esp* 2011 Sep-Oct;62(5):381-4.
19. *Fortschr Neurol Psychiatr* 2012 Sep ;80(9):527-9.
20. *J Clin Microbiol* 2010 Sep ;48(9):3438-9.
21. *J Clin Microbiol* 2010 Sep ;48(9):3438-9.
22. *Am J Med* 2008 Mar ;121(3):e1.
23. *Case Report Med* 2012 ;2012:259475.
24. *Scand J Infect Dis* 1971 ;3(1):83-5.
25. *J Med Microbiol* 2012 Oct ;61(Pt 10):1473-5.
26. *Contrib Microbiol Immunol* 1991 ;12:301-6.
27. *Acta Med Scand* 1986 ;220(5):471-6.
28. *Lakartidningen* 1983 Apr 6;80(14):1461-4.
29. *Acta Med Scand* 1981 ;209(1-2):97-101.
30. *Acta Med Scand* 1981 ;209(1-2):103-10.
31. *Lancet* 1978 Mar 4;1(8062):498-9.
32. *Scand J Infect Dis* 1977 ;9(3):253-6.
33. *J Med Microbiol* 2012 Dec 14;
34. *Scand J Infect Dis* 2012 Oct 31;
35. *Infection* 2007 Jun ;35(3):203-5.
36. *J Infect Dis* 1983 Nov ;148(5):940.
37. *J Infect* 1983 Nov ;7(3):267-9.
38. *Ugeskr Laeger* 1985 Jul 8;147(28):2236-7.
39. *Scand J Infect Dis* 1986 ;18(4):269-79.
40. *Presse Med* 1986 Mar 8;15(10):487.
41. *Am J Med* 1989 Jun ;86(6 Pt 1):723-4.
42. *Postgrad Med J* 1992 Sep ;68(803):762-3.

43. [Eur J Clin Microbiol Infect Dis 1995 Feb ;14\(2\):126-30.](#)
44. [Infection 1998 Sep-Oct;26\(5\):320-1.](#)
45. [Clin Microbiol Rev 2001 Jan ;14\(1\):177-207.](#)
46. [Scand J Infect Dis 2001 ;33\(5\):397.](#)
47. [Chemotherapy 2002 Jul ;48\(3\):158-9.](#)
48. [J Rheumatol 2008 Mar ;35\(3\):480-7.](#)
49. [Rev Rhum Engl Ed 1999 Jan 30;66\(1 Suppl\):14S-18S; discussion 19S.](#)
50. [Clin Infect Dis 2011 Sep ;53\(6\):583-91.](#)

Zika

Agent	VIRUS - RNA. Flaviviridae, Flavivirus: Zika virus
Reservoir	Human Mosquito Monkey
Vector	Mosquito - <i>Stegomyia (Aedes)</i> species
Vehicle	None
Incubation Period	5d - 8d (range 2d - 15d)
Diagnostic Tests	Viral isolation (blood). Serology. Nucleic acid amplification.
Typical Adult Therapy	Supportive
Typical Pediatric Therapy	As for adult
Clinical Hints	Zika is characterized by a mild dengue-like illness with conjunctivitis and a maculopapular rash that starts on the face and spreads to the rest of the body; joint pain is common; myalgia, retroorbital pain and leg edema may occur.
Synonyms	Zika fever. ICD9: 078.89 ICD10: A92.8

Clinical

Zika virus infection is mild illness which lasts for 4 to 7 days. ¹

- The female/male ratio of patients in 2/1.
- Signs and symptoms include fever, headache (40%) conjunctivitis (65%) and a maculopapular rash (80%) that starts on the face and spreads to the rest of the body.
- Pruritis is common
- Patients may experience arthralgia (35%) or arthritis (25%), mostly affecting the smaller joints of the hands and feet. Larger joints may also be involved
- Additional findings have included myalgias (25%) retro-orbital pain (30%) , lower extremity edema (20%), cough, diarrhea, ² , lymphadenopathy, and abdominal pain (8%).
- Clinical features tend to be relatively mild in children.
- Symptoms resolve within 4 to 7 days.

Note that Zika and Dengue are clinically similar, and may coexist in a given country.

This disease is endemic or potentially endemic to 13 countries.

Zika in Uganda

Zika virus was first described in 1947 in rhesus monkey serum in Uganda. ³

- Epizootics of Zika virus infection were documented in the Zika Forest during 1969 to 1970. ⁴

Seroprevalence surveys:

5.6% of the general population (1967 to 1969, eight areas) ⁵

The local vector appears to be *Stegomyia (Aedes) africanus*. ⁶

References

1. ProMED <promedmail.org> archive: 20070706.2157
2. Emerg Infect Dis 2012 Feb ;18(2):349-51.
3. Emerg Infect Dis 2008 Aug ;14(8):1232-9.

4. Trans R Soc Trop Med Hyg 1982 ;76(4):552-62.
5. Bull World Health Organ 1970 ;42(5):797-805.
6. Bull World Health Organ 1964 ;31:57-69.

Zygomycosis

Agent	FUNGUS. Zygomycota, Zygomycetes, Mucorales: Mucor spp., Rhizopus spp., Lichtheimia (formerly Absidia) spp, Saksenaea spp, et al
Reservoir	Saprophytes
Vector	None
Vehicle	Air Bandages Contact
Incubation Period	Variable
Diagnostic Tests	Fungal smear and culture.
Typical Adult Therapy	Amphotericin B to maximum dose 0.8 mg/kg/d; and to total dose of 3g. Excision as indicated
Typical Pediatric Therapy	Amphotericin B max dose 0.8 mg/kg/d; and to total dose of 40 mg/kg. Excision as indicated
Clinical Hints	Periorbital pain, sinusitis, and palatal, nasal or cerebral infarcts; occurs in the setting of preexisting acidosis (diabetes, uremia); pulmonary infection may complicate leukemia.
Synonyms	Absidia, Actinomucor, Apophysomyces, Cokeromyces, Cunninghamella, Hormographiella, Lichtheimia, Lichtheimia, Mucor, Mucormycosis, Mycocladus, Phycomycosis, Rhizomucor, Rhizopus, Saksenaea, Syncephalastrum. ICD9: 117.7 ICD10: B46

Clinical

Infection is most commonly associated with hyperglycemia, metabolic (diabetic, uremic) acidosis, corticosteroid therapy and neutropenia ¹, transplantation, heroin injection or administration of desferoxamine. ²

- Major risk factors identified in children are neutropenia, diabetes mellitus, and prematurity. ³
- Virtually any organ can be involved ⁴⁻⁷; however, most infections involve the paranasal sinuses and contiguous structures (orbit, cavernous sinus, cranial nerves, cerebral arteries), lungs, skin ^{8 9} and gastrointestinal tract. ¹⁰

Disease manifestations reflect the mode of transmission, with rhinocerebral and pulmonary diseases being most common. ¹¹

- Cutaneous ¹², gastrointestinal, and allergic diseases are also seen.
- The Mucorales are associated with blood vessel invasion, often leading to thrombosis, infarction and tissue destruction.
- Rare cases of sinusitis have been ascribed to *Actinomucor elegans*.
- Dissemination is common.
- Therapy must be started early and consists of antifungal drugs, surgical intervention, and reversal

Rhinocerebral zygomycosis initially manifests with headache (often unilateral), fever, facial pain, diplopia, lacrimation, and nasal stuffiness.

- As the infection spreads, necrotic lesions appear in the turbinates, nose, paranasal skin or hard palate. ¹³
- Rare cases of mycotic mandibular osteomyelitis have been reported.
- Chemosis, proptosis, and external ophthalmoplegia may occur.
- Cranial nerve abnormalities are common (nerves II through VII, IX, and X) ¹⁴, and blindness may ensue following invasion of the cavernous sinus, ophthalmic artery, and orbit.
- Hemiparesis, seizures, or monocular blindness suggest advanced disease.
- Invasion of the internal carotid artery in the cavernous sinus can occur, with metastatic lesions in the frontoparietal cortex and deepening coma.

Pulmonary zygomycosis presents with nonspecific symptoms such as fever, cough and dyspnea. ¹⁵

- Hemoptysis may occur with vascular invasion.
- Radiological findings include segmental consolidation which progresses to contiguous areas of the lung and may cavitate.
- In 74% of pulmonary zygomycosis cases, the infection is limited to the lung. ¹⁶

Gastrointestinal zygomycosis usually affects patients with severe malnutrition, and may involve the stomach, ileum, colon ¹⁷ or peritoneum. ¹⁸

- Clinical findings mimic intra-abdominal abscess.
- The diagnosis is often made at autopsy.

Cutaneous zygomycosis may present as primary infection, characterized by necrotic lesions following trauma; or

secondary extension from a focus of rhinocerebral infection. ^{19 20}

Renal zygomycosis may mimic malignancy ²¹⁻²³

59 case reports (38 fatal) of neonatal zygomycosis had been published to July 2007 • 77% premature infants, 54% gastrointestinal and 36% dermal. ²⁴

Zygomycosis has a poor prognosis, with a mortality rate of 44%. ²⁵

This disease is endemic or potentially endemic to all countries.

References

1. Haematologica 2004 Feb ;89(2):207-14.
2. Curr Opin Infect Dis 2004 Dec ;17(6):517-25.
3. Pediatr Infect Dis J 2007 Aug ;26(8):723-7.
4. Cir Cir 2007 Nov-Dec;75(6):465-9.
5. Intern Med 2008 ;47(9):839-42.
6. Transpl Infect Dis 2008 Dec ;10(6):419-25.
7. Int J Surg Pathol 2011 Feb ;19(1):75-9.
8. Ann Plast Surg 2008 Apr ;60(4):433-6.
9. Indian J Dermatol Venereol Leprol 2008 Jul-Aug;74(4):367-70.
10. Clin Microbiol Rev 2005 Jul ;18(3):556-69.
11. Clin Infect Dis 2012 Feb ;54 Suppl 1:S55-60.
12. Clin Microbiol Infect 2009 Oct ;15 Suppl 5:41-5.
13. Clin Microbiol Infect 2004 Mar ;10 Suppl 1:31-47.
14. Case Report Med 2011 ;2011:216404.
15. Semin Respir Crit Care Med 2008 Apr ;29(2):111-20.
16. Am J Transplant 2009 Sep ;9(9):2166-71.
17. Med Mycol 2006 Nov ;44(7):683-7.
18. Singapore Med J 2012 May ;53(5):e106-9.
19. Clin Dermatol 2012 Jul ;30(4):413-9.
20. Clin Dermatol 2012 Nov ;30(6):628-32.
21. Am J Med Sci 2009 Oct ;338(4):330-3.
22. Urol Int 2012 Feb 17;
23. Med Mycol 2012 May 15;
24. Am J Perinatol 2009 Sep ;26(8):565-73.
25. Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2007 Nov ;104(5):e28-34.

About GIDEON

GIDEON Informatics produces the GIDEON [web application](#) and the GIDEON [ebooks](#) series.

GIDEON online

GIDEON online is the world's premier global infectious disease knowledge management tool. GIDEON (Global Infectious Diseases and Epidemiology Online Network) is an easy to use, interactive and comprehensive web based tool that helps overcome information overload, save time and access a vast knowledge database. GIDEON is used for diagnosis and reference in the fields of Tropical and Infectious Diseases, Epidemiology, Microbiology, Antimicrobial Therapy and Occupational Toxicology.

Content

GIDEON is made up of three modules, which are updated continually: Infectious Diseases, Microbiology and Toxicology. The Infectious Diseases module encompasses 347 diseases, 231 countries, and 500+ anti-infective drugs and vaccines. Microbiology includes over 1,500 microbial taxa; and Toxicology, over 3,000 agents and 205 diseases. GIDEON's worldwide data sources access the entire world's literature and adhere to the standards of Evidence Based Medicine. Over 18,000 notes outline the status of specific infections within each country. Also featured are over 33,000 images, graphs, and interactive maps and more than 150,000 linked references.

Users

GIDEON is used in hospitals, universities (colleges and medical schools), private practice, Public Health departments and Military installations - by physicians (emergency room, infectious diseases, pediatrics and hospitalists), teachers, clinical microbiologists and health professionals. It is an ideal teaching tool for health care students, residents and fellows.

Accuracy

The Infectious Diseases Diagnosis module has been tested in a blinded multi-center field trial of 495 patients. The correct diagnosis was displayed in over 94% of cases, and was listed first in over 75%. GIDEON has been reviewed in numerous journals and is continually updated daily to maintain content and accuracy.

GIDEON ebooks

GIDEON ebooks complement the GIDEON web application by expanding easy access to the GIDEON's vast content without a subscription or continual internet access. Ebooks can be downloaded to a variety of devices and can be read anywhere. These ebooks summarize the status of individual infectious diseases, in every country of the world.

To learn more about GIDEON online, visit www.gideononline.com and follow our blog at www.gideononline.com/blog. For the latest list of GIDEON ebooks, visit www.gideononline.com/ebooks