

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2014287009 B2**

(54) Title
Compositions comprising synthetic polynucleotides encoding CRISPR related proteins and synthetic sgRNAs and methods of use

(51) International Patent Classification(s)
C12N 15/85 (2006.01) **C12N 15/63** (2006.01)

(21) Application No: **2014287009** (22) Date of Filing: **2014.07.11**

(87) WIPO No: **WO15/006747**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/844,890	2013.07.11	US
61/886,545	2013.10.03	US

(43) Publication Date: **2015.01.15**

(44) Accepted Journal Date: **2020.10.29**

(71) Applicant(s)
Moderna Therapeutics, Inc.

(72) Inventor(s)
Hoge, Stephen G.;Huang, Eric Yi-Chun;Chakraborty, Tirtha

(74) Agent / Attorney
Griffith Hack, GPO Box 4164, Sydney, NSW, 2001, AU

(56) Related Art
WANG HAOYI ET AL, CELL, CELL PRESS, US, (2013-05-02), vol. 153, no. 4, doi:10.1016/J.CELL.2013.04.025, ISSN 0092-8674, pages 910 - 918
WO 2012/031043 A1
WO 2012135805 A2



- (51) **International Patent Classification:**
C12N 15/85 (2006.01) C12N 15/63 (2006.01)
- (21) **International Application Number:**
PCT/US2014/046434
- (22) **International Filing Date:**
11 July 2014 (11.07.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/844,890 11 July 2013 (11.07.2013) US
61/886,545 3 October 2013 (03.10.2013) US
- (71) **Applicant: MODERNA THERAPEUTICS, INC.**
[US/US]; 200 Technology Square, Cambridge, MA 02139 (US).
- (72) **Inventors: HOGE, Stephen, G.;** c/o Moderna Therapeutics, Inc., 200 Technology Square, Cambridge, MA 02139 (US). **HUANG, Eric, Yi-chun;** c/o Moderna Therapeutics, Inc., 200 Technology Square, Cambridge, MA 02139 (US). **CHAKRABORTY, Tirtha;** c/o Moderna Therapeutics, Inc., 200 Technology Square, Cambridge, MA 02139 (US).
- (74) **Agents: BELLIVEAU, Michael J. et al.;** Clark & Elbing LLP, 101 Federal Street, 15th Floor, Boston, MA 02110 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- of inventorship (Rule 4.17(iv))


Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

- (88) **Date of publication of the international search report:**
23 April 2015

(54) **Title:** COMPOSITIONS COMPRISING SYNTHETIC POLYNUCLEOTIDES ENCODING CRISPR RELATED PROTEINS AND SYNTHETIC SGRNAS AND METHODS OF USE

Figure 1



(57) **Abstract:** The present invention relates to compositions comprising and methods of using synthetic polynucleotides, e.g., modified mRNA, encoding CRISPR related proteins including dCAS9 and synthetic sgRNAs targeting a gene of interest.

WO 2015/006747 A3

TITLE

[0001] Compositions comprising synthetic polynucleotides encoding CRISPR related proteins and synthetic sgRNAs and methods of use.

CROSS-REFERENCE TO RELATED APPLICATION

[0002] This application claims the benefit of U.S. Provisional Patent Application No. 61/844,890, filed July 11, 2013, and US Provisional Patent Application No. 61/886,545, filed October 3, 2013, the disclosures of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0003] The present invention is directed to compositions comprising and methods of using synthetic polynucleotides and short guide RNAs (sgRNAs). The synthetic polynucleotides are, e.g., synthetic modified RNA encoding CRISPR related proteins, e.g., dCAS9 and dCAS9-effector domain fusion proteins. The synthetic sgRNAs target a gene of interest. The synthetic polynucleotides and synthetic sgRNAs can be used to modulate transcription, e.g., in therapeutics and/or clinical and research settings.

BACKGROUND OF THE INVENTION

[0004] Bacterial and archaeal Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) systems rely on CRISPR RNAs (crRNAs) in complex with CRISPR-associated (CAS) proteins to direct degradation of complementary sequences present within invading viral and plasmid DNA (Mali et al. *Science*. 2013. 339:823-826; herein incorporated by reference in its entirety).

[0005] In type II CRISPR systems, the pre-crRNAs are processed by trans-activating crRNA (tracrRNA) complementary to the repeat sequences in pre-crRNA which triggers processing by the double-stranded (ds) RNA-specific ribonuclease RNase III in the presence of the CAS9 protein. CAS9 (formerly CSN1) is the key protein in the type II CRISPR systems and is thought to be the sole protein responsible for crRNA-guided silencing of foreign DNA (Jinek et al. *Science*. 2012. 337:816-821; herein incorporated by reference in its entirety). The CAS9 protein is also hypothesized to be involved in both crRNA maturation and crRNA-guided DNA interference (Jinek et al. *Science*. 2012. 337:816-821; herein incorporated by reference in its entirety).

[0006] The CAS9 endonuclease family can be programmed with single RNA molecules to cleave specific DNA sites which may be used to develop a versatile RNA-directed system to generate dsDNA breaks for genome targeting and editing (Jinek et al.

Science. 2012. 337:816-821; herein incorporated by reference in its entirety). This use of CAS9 could enhance the ease of genome engineering.

[0007] The Church Lab engineered the bacterial and archaeal type II CRISPR system to function with custom guide RNA (gRNA) in human cells (Mali et al. Science. 2013. 339:823-826; herein incorporated by reference in its entirety).

[0008] It has been shown that fusion of CRISPR-associated catalytically inactive dCas9 protein to distinct effector domains (e.g. VP64, p65AD, KRAB, and Mxi1) enables repression or activation of transcription in human and yeast cells, with the site of delivery determined solely by a co-expressed short guide RNA (sgRNA). Coupling of dCas9 to a transcriptional repressor domain can silence expression of multiple endogenous genes, with no detectable off-targets as verified by RNA-seq analysis. (Qi et al. Cell. 2013. 152:1173-1183)

[0009] One example of a catalytically inactive Cas9 protein is the mutant (D10A, H840A) *Streptococcus pyogenes* Cas9 protein, typically referred to as dCAS9 (Qi et al. Cell. 2013. 152:1173-1183; Gilbert et al. Cell. 2013. 154:1-10). A *Streptococcus thermophiles* Cas9 with 2 endonuclease domains and single amino acid substitution (D31A, H858A) has been described which also resulted in catalytically inactive Cas9 (Sapranaukas et al. Nucleic Acids Research. 2011. 39:9275-9282).

[0010] Other publications describing the CRISPR systems, Cas9, and dCas9 include the following Cong et al (2013); Jinek et al (2012); Lei et al (2013); Gilbert et al (2013); Perez-Pinela et al (2013); Maider et al (2013).

[0011] It has been shown that certain modified mRNA sequences have the potential as therapeutics with benefits beyond just evading, avoiding or diminishing the immune response. Such studies are detailed in published co-pending applications International Publication No WO2012019168 filed August 5, 2011 and International Publication No WO2012045075 filed October 3, 2011, International Publication No WO2013052523 filed October 3, 2011, the contents of which are incorporated herein by reference in their entirety.

SUMMARY

The present invention as claimed herein is described in the following items 1 to 19:

1. A composition comprising:

- (a) an mRNA comprising:
 - (i) at least one 5'-cap structure;
 - (ii) a 5'-UTR;
 - (iii) an open reading frame encoding a polypeptide,
wherein the polypeptide has an amino acid sequence that is at least 80% identical to SEQ ID NO: 7, 8, 9, 61, 62, 63, 64, or 65, and
wherein the open reading frame consists of nucleotides selected from 1-methyl-pseudouridine, cytidine, adenosine, and guanosine; and
 - (iv) a 3'-UTR;
 - (b) an sgRNA comprising:
 - (i) a first region of 20-25 linked nucleosides complementary to either strand of a 5'-UTR of the gene of interest;
 - (ii) a second region located at the 3'-terminus of the first region comprising a guide RNA scaffold sequence;
 - (iii) at least one phosphorothioate linkage; and
 - (iv) at least one 2'-methoxy-nucleotide;
- and
- (c) a plurality of lipid nanoparticles comprising:
 - (i) a biodegradable cationic lipid;
 - (ii) a non-cationic lipid;
 - (iii) a sterol; and
 - (iv) a PEGylated lipid,
- wherein the lipid nanoparticles encapsulate the mRNA and/or the sgRNA.

2. The composition of item 1, wherein the mRNA further comprises a poly-A tail.
3. The composition of item 2, wherein the poly-A tail is from 80 to 140 nucleotides in length.
4. The composition of item 3, wherein the poly-A tail is 100 nucleotides in length.

5. The composition of any one of items 1 to 4, wherein the biodegradable cationic lipid comprises at least one ester linkage.
6. The composition of item 5, wherein the biodegradable cationic lipid comprises an internal ester.
7. The composition of item 5, wherein the biodegradable cationic lipid comprises a terminal ester.
8. The composition of any one of items 1 to 7, wherein the polypeptide comprises a sequence having at least 80% identity to SEQ ID NO: 61.
9. The composition of item 8, wherein the polypeptide comprises a sequence that is 100% identical to SEQ ID NO: 61.
10. The composition of any one of items 1 to 9, wherein the open reading frame further encodes an effector domain.
11. The composition of item 10, wherein the effector domain is selected from the group consisting of KRAB, VP64, p65AD, and Mxi.
12. The composition of item 11, wherein the effector domain is KRAB or VP64.
13. The composition of any one of items 1 to 12, wherein the 3'-UTR comprises at least one miR binding site or one miR binding site seed.
14. The composition of item 13, wherein the 3'-UTR comprises a miR-122 or miR-142 or miR-146 binding site.

15. The composition of any one of items 1 to 14, wherein the guide RNA scaffold sequence comprises SEQ ID NO: 82.

16. The composition of any one of items 1 to 15, wherein the mRNA is produced using an in vitro transcription method or using a chemical synthesis method.

17. The composition of any one of items 1 to 16, wherein the mRNA is substantially purified.

18. The composition of any one of items 1 to 17, further comprising a pharmaceutically acceptable excipient.

19. The composition of item 18, wherein the pharmaceutically acceptable excipient is selected from the group consisting of a solvent, aqueous solvent, non-aqueous solvent, dispersion media, diluent, dispersion, suspension aid, surface active agent, isotonic agent, thickening or emulsifying agent, preservative, polymer, peptide, protein, cell, hyaluronidase, and mixtures thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012] The foregoing and other objects, features and advantages will be apparent from the following description of particular embodiments of the invention, as illustrated in the accompanying drawings in which like reference characters refer to the same parts throughout the different views. The drawings are not necessarily to scale, emphasis

instead being placed upon illustrating the principles of various embodiments of the invention.

[0013] FIG. 1 is a schematic of a synthetic polynucleotide of the present invention.

[0014] FIG. 2 shows the results of an anti-FLAG Western and Immunoprecipitation performed on the lysates of HeLa cells transfected with FLAG-tagged Cas9 constructs or controls.

[0015] FIG. 3 shows the results of a western analysis of HeLa cells that were transfected with HA-tagged Gilbert Cas9 mRNA using Trilink-HA-tagged Cas9 mRNA as a positive control.

[0016] FIG. 4 shows the results of a LC-PRM Cas9 peptide quantification in cell lysates of HeLa cells transfected with Maeder and Gilbert Cas9 constructs versus untreated HeLa cells. Each chart shows the quantification of a peptide fragment of the Cas9 construct. Figure 4A represents quantification of GNELALPSK peptide (SEQ ID NO: 120). Figure 4B represents quantification of the YFDTTIDR peptide (SEQ ID NO: 121). Figure 4C represents quantification of the IPYYVGPLAR peptide (SEQ ID NO: 122).

[0017] FIG. 5 shows the results of an anti-FLAG Western and Immunoprecipitation performed on the lysates of mouse livers from animals dosed with 14ug Cas9 LNP (MC3) or mock. WCE = whole cell extract. IP = immunoprecipitation with M2-agarose beads (Sigma). Ladder = See Blue Plus 2™ prestained protein marker (LifeTech).

DETAILED DESCRIPTION

[0018] The present invention is based, at least in part, on the surprising discovery that mRNAs can be expressed in the cytoplasm and translocated to the nucleus. The mRNA technology of the invention is surprisingly suited for delivery of intracellular proteins. The mRNA technology of the invention is also desirable suited for delivery of multiple mRNAs. The methodologies of the invention avoid or circumvent problems associated with conventional gene therapy approaches, such as toxicity, integration of DNA into host genome, and the like. The methodologies of the invention are particularly suited to the delivery of a plurality of mRNAs, without inducing an innate immune and without undesirable side effects associated with conventional gene therapy approaches. In particular, the mRNA delivery technology of the invention facilitates the expression both the mRNA encoding CRISPR-related proteins, *e.g.*, Cas9, dCas9 and variants thereof, in combination with one or more guide RNAs (sgRNAs) to enable gene regulation *in vitro*

and *in vivo*. Using mRNA-based pharmaceutical compositions, the technology provides for therapeutic applications of the CRISPR technologies that were previously unattainable

[0019] The data presented herein evidence that it is possible to achieve intracellular expression of CRISPR-related proteins, including proteins engineered to include regulatory sequence facilitating regulation of protein expression, and the corresponding mRNA targeting sequences (*i.e.*, sgRNAs). The data presented herein demonstrate that it is possible to achieve intracellular protein expression of CRISPR-related proteins, in particular, *via in vivo* administration of pharmaceutical compositions containing modified mRNAs encoding such proteins.

[0020] The details of various embodiments of the invention are set forth in the description below. Other features, objects, and advantages of the invention will be apparent from the description and the drawings, and from the claims.

[0021] Described herein are compositions (including pharmaceutical compositions) and methods for the design, preparation, manufacture and/or formulation of synthetic polynucleotides encoding one or more one or more CRISPR related proteins, e.g., dCAS9 and/or dCAS9-effector fusion proteins. Also described herein are compositions (including pharmaceutical compositions) and methods for the design, preparation, manufacture and/or formulation of one or more synthetic small guide RNAs (“sgRNAs”) targeting a gene of interest. Both the synthetic polynucleotides and sgRNAs are typically manufactured using *in vitro* transcription. In one embodiment the synthetic polynucleotide and sgRNA are modified, e.g., comprise at least one modified nucleotide. Also described herein are method of modulation of expression of a gene of interest using a synthetic polynucleotide encoding a CRISPR related protein and an sgRNA targeting the gene of interest. Included are *in vitro* and *in vivo* methods, e.g., methods of treatment of a disease state correlated with expression of the gene of interest. Also provided are systems, processes, devices and kits for the selection, design and/or utilization of the synthetic polynucleotides and synthetic sgRNAs described herein.

[0022] A key limitation of the CRISPR/Inhibition/Activation (CRISPR/I/A) systems described to date is their dependence on delivery of DNA by viral vectors to drive intracellular transcription (of the sgRNA) and transcription and translation of Cas9, dCas9 or dCas9-fusion proteins such as dCas9-VP64, and dCas9-KRAB. There are several implications of this limitation for *in vivo* systems using transcriptional activation/repression as this will require repeat dosing. First, delivery of DNA by pseudo virus will induce measurable innate immune activation (e.g., thru TLRs) and these will

impact transfection efficiency and have undesirable side effects. Second, the repeat dosing of viral capsids could and likely will induce an adaptive immune response on repeat dosing. Third, the use of DNA – and the risk of chronically activated dCas9 represent a significant safety and toxicity risk for in vivo therapeutics.

[0023] Modified synthetic polynucleotide (e.g., modified mRNA) presents an ideal platform for the clinical translation of this prokaryotic platform. Technologies are disclosed herein to safely delivery modified mRNA in vivo that encodes for CRISPR related protein, e.g., dCA9, in a time limited and – potentially cell-type - targeted fashion. Moreover, the use of modified mRNA will avoid the risk of significant innate immune activation, and diminish the risk of adaptive responses.

I. Synthetic polynucleotides encoding CRISPR related proteins

[0024] The present invention provides synthetic polynucleotides which encode one or more CRISPR related proteins, e.g., dCAS9 and dCAS9 fusion proteins. The term “polynucleotide,” in its broadest sense, includes any compound and/or substance that comprise a polymer of nucleotides. Exemplary polynucleotides of the invention include, but are not limited to, ribonucleic acids (RNAs), deoxyribonucleic acids (DNAs), threose nucleic acids (TNAs), glycol nucleic acids (GNAs), peptide nucleic acids (PNAs), locked nucleic acids (LNAs, including LNA having a β - D-ribo configuration, α -LNA having an α -L-ribo configuration (a diastereomer of LNA), 2'-amino-LNA having a 2'-amino functionalization, and 2'-amino- α -LNA having a 2'-amino functionalization) or hybrids thereof.

[0025] In one embodiment, the synthetic polynucleotide is a synthetic messenger RNA (mRNA). As used herein, the term “messenger RNA” (mRNA) refers to any polynucleotide, which may be synthetic, which encodes a polypeptide, e.g., a CRISPR related protein, and which is capable of being translated to produce the encoded polypeptide *in vitro*, *in vivo*, *in situ* or *ex vivo*. In the present invention, the mRNA encodes a CRISPR related protein, e.g., dCAS9 and dCA9-effector (activator or inhibitor) fusion proteins. Additional description of and sequences for CRISPR related proteins are below.

[0026] The present invention expands the scope of functionality of traditional mRNA molecules by providing synthetic polynucleotides which comprise one or more structural and/or chemical modifications or alterations which impart useful properties to the polynucleotides including, in some embodiments, the lack of a substantial induction of

the innate immune response of a cell into which the polynucleotide is introduced. As used herein, a “structural” feature or modification is one in which two or more linked nucleotides are inserted, deleted, duplicated, inverted or randomized in a synthetic polynucleotide, primary construct or mmRNA without significant chemical modification to the nucleotides themselves. Because chemical bonds will necessarily be broken and reformed to effect a structural modification, structural modifications are of a chemical nature and hence are chemical modifications. However, structural modifications will result in a different sequence of nucleotides. For example, the polynucleotide “ATCG” (SEQ ID NO: 123) may be chemically modified to “AT-5meC-G” (SEQ ID NO: 124). The same polynucleotide may be structurally modified from “ATCG” (SEQ ID NO: 123) to “ATCCCG” (SEQ ID NO: 125). Here, the dinucleotide “CC” has been inserted, resulting in a structural modification to the polynucleotide.

CRISPR related proteins

[0027] The synthetic polynucleotides of the invention encode a CRISPR related protein. The term “CRISPR related protein” includes but is not limited to CAS9, CSY4, dCAS9, and dCAS9-effector domain (activator and/or inhibitor domain) fusion proteins. Examples of CRISPR related protein polypeptide sequences and polynucleotide sequences are found in the Tables below.

[0028] The CRISPR related protein can be from any number of species including but not limited to *Streptococcus pyogenes*, *Listeria innocua*, and *Streptococcus thermophilus*.

[0029] The synthetic polynucleotide can encode the wild-type sequence of the CRISPR related protein or a variant CRISPR related protein. Variants are described in further details herein. As described in more detail herein, the synthetic polynucleotide can include either the wild-type codon usage or codon usage optimized for a particular application, e.g., human codon optimization for human therapeutics.

[0030] According to the present invention, the synthetic polynucleotides comprise at least a first region of linked nucleosides encoding at least one CRISPR related proteins. Example of CRISPR related proteins are listed in the Tables below. For any particular CRISPR related protein there may exist one or more variants or isoforms. It will be appreciated by those of skill in the art that disclosed in the Tables are potential flanking regions. These are encoded in each transcript either to the 5' (upstream) or 3' (downstream) of the ORF or coding region. The coding region is definitively and specifically disclosed by teaching the protein sequence. Consequently, the sequences

taught flanking that encoding the protein are considered flanking regions. It is also possible to further characterize the 5' and 3' flanking regions by utilizing one or more available databases or algorithms. Databases have annotated the features contained in the flanking regions of the transcripts and these are available in the art.

[0031] In one embodiment the CRISPR related protein is dCAS9. Typical sequences are included in the Tables below.

[0032] In another embodiment the CRISPR related protein is a dCAS9-effector domain fusion protein. The effector domain can be an activation domain or an inhibition domain, depending on the application. Examples of effector domains are well known to one of skill in the art and include, e.g., KRAB, CSD, WRPW, VP64, or p65AD.

[0033] In some embodiments, the CRISPR related protein is dCAS9-KRAB or dCAS9-VP64 fusion proteins. Examples of sequences are provided in the Tables below and the Sequence Listing.

[0034] Further constructs useful in various aspects of the invention include those set forth as SEQ ID NOs: 1-9. These include the nucleotide sequence encoding Cas9, serotype M1 (SEQ ID NO:1); nucleotide sequence encoding FLAG-tagged (*i.e.*, trimer-FLAG) CAS9 having a nuclear localization sequence (NLS) (SEQ ID NO:2); nucleotide sequence encoding HA-tagged Cas9 having a NLS (Cas9-HAtag/2xNLS) conjugated to GFP (SEQ ID NO:3); nucleotide sequence encoding Cas9 C-terminal fragment having a NLS (SEQ ID NO:4); nucleotide sequence encoding Cas9 N-terminal fragment (SEQ ID NO:5); nucleotide sequence encoding FLAG-tagged Cas9 having NLSs (FLAG-NLS-Cas9-NLS) (SEQ ID NO:6); as well as Cas9 protein sequences (Cas9 - SEQ ID NO:8, Cas9 serotype M1 - SEQ ID NO:9) and FLAG-tagged (*i.e.*, trimer-FLAG) CAS9 having a nuclear localization sequence (NLS) (SEQ ID NO:9.)

Synthetic polynucleotide Architecture

[0035] The modified synthetic polynucleotides or of the present invention are distinguished from wild type mRNA in their functional and/or structural design features which serve to, as evidenced herein, overcome existing problems of effective polypeptide production using nucleic acid-based therapeutics.

[0036] Figure 1 shows a representative polynucleotide primary construct **100** of the present invention. As used herein, the term "primary construct" or "primary mRNA construct" refers to a polynucleotide transcript which encodes one or more polypeptides of interest and which retains sufficient structural and/or chemical features to allow the

polypeptide of interest encoded therein to be translated. Primary constructs may be polynucleotides of the invention. When structurally or chemically modified, the primary construct may be referred to as an modified synthetic polynucleotides or mmRNA.

[0037] Returning to FIG. 1, the primary construct **100** here contains a first region of linked nucleotides **102** that is flanked by a first flanking region **104** and a second flanking region **106**. As used herein, the “first region” may be referred to as a “coding region” or “region encoding” or simply the “first region.” This first region may include, but is not limited to, the encoded polypeptide of interest, i.e., the CRISPR related protein. The polypeptide of interest may comprise at its 5' terminus one or more signal sequences encoded by a signal sequence region **103**. The first flanking region **104** may comprise a region of linked nucleotides comprising one or more complete or incomplete 5' UTRs sequences. The first flanking region **104** may also comprise a 5' terminal capping region **108**. The second flanking region **106** may comprise a region of linked nucleotides comprising one or more complete or incomplete 3' UTRs. The second flanking region **106** may also comprise a 3' tailing sequence **110**.

[0038] Bridging the 5' terminus of the first region **102** and the first flanking region **104** is a first operational region **105**. Traditionally this first operational region comprises a Start codon. The operational region may alternatively comprise any translation initiation sequence or signal including a Start codon.

[0039] Bridging the 3' terminus of the first region **102** and the second flanking region **106** is a second operational region **107**. Traditionally this second operational region comprises a Stop codon. The second operational region may alternatively comprise any translation initiation sequence or signal including a Stop codon. According to the present invention, multiple serial stop codons may also be used.

[0040] In some embodiments, the synthetic polynucleotide, primary construct, or mmRNA includes from about 30 to about 100,000 nucleotides (e.g., from 30 to 50, from 30 to 100, from 30 to 250, from 30 to 500, from 30 to 1,000, from 30 to 1,500, from 30 to 3,000, from 30 to 5,000, from 30 to 7,000, from 30 to 10,000, from 30 to 25,000, from 30 to 50,000, from 30 to 70,000, from 100 to 250, from 100 to 500, from 100 to 1,000, from 100 to 1,500, from 100 to 3,000, from 100 to 5,000, from 100 to 7,000, from 100 to 10,000, from 100 to 25,000, from 100 to 50,000, from 100 to 70,000, from 100 to 100,000, from 500 to 1,000, from 500 to 1,500, from 500 to 2,000, from 500 to 3,000, from 500 to 5,000, from 500 to 7,000, from 500 to 10,000, from 500 to 25,000, from 500 to 50,000, from 500 to 70,000, from 500 to 100,000, from 1,000 to 1,500, from 1,000 to

2,000, from 1,000 to 3,000, from 1,000 to 5,000, from 1,000 to 7,000, from 1,000 to 10,000, from 1,000 to 25,000, from 1,000 to 50,000, from 1,000 to 70,000, from 1,000 to 100,000, from 1,500 to 3,000, from 1,500 to 5,000, from 1,500 to 7,000, from 1,500 to 10,000, from 1,500 to 25,000, from 1,500 to 50,000, from 1,500 to 70,000, from 1,500 to 100,000, from 2,000 to 3,000, from 2,000 to 5,000, from 2,000 to 7,000, from 2,000 to 10,000, from 2,000 to 25,000, from 2,000 to 50,000, from 2,000 to 70,000, and from 2,000 to 100,000).

[0041] Generally, the length of the first region encoding the polypeptide of interest, e.g., CRISPR related protein of the present invention is greater than about 30 nucleotides in length (e.g., at least or greater than about 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, 2,000, 2,500, and 3,000, 4,000, 5,000, 6,000, 7,000, 8,000, 9,000, 10,000, 20,000, 30,000, 40,000, 50,000, 60,000, 70,000, 80,000, 90,000 or up to and including 100,000 nucleotides).

[0042] According to the present invention, the first and second flanking regions may range independently from 15-1,000 nucleotides in length (e.g., greater than 30, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, and 900 nucleotides or at least 30, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, and 1,000 nucleotides).

[0043] According to the present invention, the 3' tailing sequence may range from absent to 500 nucleotides in length (e.g., at least 60, 70, 80, 90, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, or 500 nucleotides). Where the tailing sequence is a polyA tail, the length may be determined in units of or as a function of polyA Binding Protein binding. In this embodiment, the polyA tail is long enough to bind at least 4 monomers of PolyA Binding Protein. PolyA Binding Protein monomers bind to stretches of approximately 38 nucleotides. As such, it has been observed that polyA tails of about 80 nucleotides and 160 nucleotides are functional.

[0044] According to the present invention, the 5' terminal capping region may comprise a single cap or a series of nucleotides forming the cap. In this embodiment the capping region may be from 1 to 10, e.g. 2-9, 3-8, 4-7, 1-5, 5-10, or at least 2, or 10 or fewer nucleotides in length. In some embodiments, the capping region is absent.

[0045] According to the present invention, the first and second operational regions may range from 3 to 40, e.g., 5-30, 10-20, 15, or at least 4, or 30 or fewer nucleotides in

length and may comprise, in addition to a Start and/or Stop codon, one or more signal and/or restriction sequences.

Cyclic Synthetic Polynucleotides

[0046] According to the present invention, a synthetic polynucleotide may be cyclized, or concatemerized, to generate a translation competent molecule to assist interactions between poly-A binding proteins and 5'-end binding proteins. The mechanism of cyclization or concatemerization may occur through at least 3 different routes: 1) chemical, 2) enzymatic, and 3) ribozyme catalyzed. The newly formed 5'-/3'-linkage may be intramolecular or intermolecular.

[0047] In the first route, the 5'-end and the 3'-end of the nucleic acid contain chemically reactive groups that, when close together, form a new covalent linkage between the 5'-end and the 3'-end of the molecule. The 5'-end may contain an NHS-ester reactive group and the 3'-end may contain a 3'-amino-terminated nucleotide such that in an organic solvent the 3'-amino-terminated nucleotide on the 3'-end of a synthetic mRNA molecule will undergo a nucleophilic attack on the 5'-NHS-ester moiety forming a new 5'-/3'-amide bond.

[0048] In the second route, T4 RNA ligase may be used to enzymatically link a 5'-phosphorylated nucleic acid molecule to the 3'-hydroxyl group of a nucleic acid forming a new phosphodiester linkage. In an example reaction, 1µg of a nucleic acid molecule is incubated at 37°C for 1 hour with 1-10 units of T4 RNA ligase (New England Biolabs, Ipswich, MA) according to the manufacturer's protocol. The ligation reaction may occur in the presence of a split oligonucleotide capable of base-pairing with both the 5'- and 3'-region in juxtaposition to assist the enzymatic ligation reaction.

[0049] In the third route, either the 5'-or 3'-end of the cDNA template encodes a ligase ribozyme sequence such that during *in vitro* transcription, the resultant nucleic acid molecule can contain an active ribozyme sequence capable of ligating the 5'-end of a nucleic acid molecule to the 3'-end of a nucleic acid molecule. The ligase ribozyme may be derived from the Group I Intron, Group I Intron, Hepatitis Delta Virus, Hairpin ribozyme or may be selected by SELEX (systematic evolution of ligands by exponential enrichment). The ribozyme ligase reaction may take 1 to 24 hours at temperatures between 0 and 37°C.

Synthetic Polynucleotide Multimers

[0050] According to the present invention, multiple distinct synthetic polynucleotides may be linked together through the 3'-end using nucleotides which are modified at the 3'-terminus. Chemical conjugation may be used to control the stoichiometry of delivery into cells. For example, the glyoxylate cycle enzymes, isocitrate lyase and malate synthase, may be supplied into HepG2 cells at a 1:1 ratio to alter cellular fatty acid metabolism. This ratio may be controlled by chemically linking polynucleotides, primary constructs or mmRNA using a 3'-azido terminated nucleotide on one polynucleotide, primary construct or mmRNA species and a C5-ethynyl or alkynyl-containing nucleotide on the opposite polynucleotide, primary construct or mmRNA species. The modified nucleotide is added post-transcriptionally using terminal transferase (New England Biolabs, Ipswich, MA) according to the manufacturer's protocol. After the addition of the 3'-modified nucleotide, the two polynucleotide, primary construct or mmRNA species may be combined in an aqueous solution, in the presence or absence of copper, to form a new covalent linkage via a click chemistry mechanism as described in the literature.

[0051] In another example, more than two polynucleotides may be linked together using a functionalized linker molecule. For example, a functionalized saccharide molecule may be chemically modified to contain multiple chemical reactive groups (SH-, NH₂-, N₃, etc...) to react with the cognate moiety on a 3'-functionalized mRNA molecule (i.e., a 3'-maleimide ester, 3'-NHS-ester, alkynyl). The number of reactive groups on the modified saccharide can be controlled in a stoichiometric fashion to directly control the stoichiometric ratio of conjugated polynucleotide, primary construct or mmRNA.

Synthetic Polynucleotide Conjugates and Combinations

[0052] In order to further enhance protein production, synthetic polynucleotides of the present invention can be designed to be conjugated to other polynucleotides, dyes, intercalating agents (*e.g.* acridines), cross-linkers (*e.g.* psoralene, mitomycin C), porphyrins (TPPC4, texaphyrin, Sapphyrin), polycyclic aromatic hydrocarbons (*e.g.*, phenazine, dihydrophenazine), artificial endonucleases (*e.g.* EDTA), alkylating agents, phosphate, amino, mercapto, PEG (*e.g.*, PEG-40K), MPEG, [MPEG]₂, polyamino, alkyl, substituted alkyl, radiolabeled markers, enzymes, haptens (*e.g.* biotin), transport/absorption facilitators (*e.g.*, aspirin, vitamin E, folic acid), synthetic ribonucleases, proteins, *e.g.*, glycoproteins, or peptides, *e.g.*, molecules having a specific affinity for a co-ligand, or antibodies *e.g.*, an antibody, that binds to a specified cell type

such as a cancer cell, endothelial cell, or bone cell, hormones and hormone receptors, non-peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, or a drug.

[0053] Conjugation may result in increased stability and/or half life and may be particularly useful in targeting the polynucleotides, primary constructs or mmRNA to specific sites in the cell, tissue or organism.

[0054] According to the present invention, the mmRNA or primary constructs may be administered with, or further encode one or more of RNAi agents, siRNAs, shRNAs, miRNAs, miRNA binding sites, antisense RNAs, ribozymes, catalytic DNA, tRNA, RNAs that induce triple helix formation, aptamers or vectors, and the like.

Bifunctional Synthetic Polynucleotides

[0055] In one embodiment of the invention are bifunctional polynucleotides (e.g., bifunctional primary constructs or bifunctional mmRNA). As the name implies, bifunctional polynucleotides are those having or capable of at least two functions. These molecules may also by convention be referred to as multi-functional.

[0056] The multiple functionalities of bifunctional polynucleotides may be encoded by the RNA (the function may not manifest until the encoded product is translated) or may be a property of the polynucleotide itself. It may be structural or chemical. Bifunctional modified polynucleotides may comprise a function that is covalently or electrostatically associated with the polynucleotides. Further, the two functions may be provided in the context of a complex of a mmRNA and another molecule.

[0057] Bifunctional polynucleotides may encode peptides which are anti-proliferative. These peptides may be linear, cyclic, constrained or random coil. They may function as aptamers, signaling molecules, ligands or mimics or mimetics thereof. Anti-proliferative peptides may, as translated, be from 3 to 50 amino acids in length. They may be 5-40, 10-30, or approximately 15 amino acids long. They may be single chain, multichain or branched and may form complexes, aggregates or any multi-unit structure once translated.

Noncoding Polynucleotides and Primary Constructs

[0058] As described herein, provided are polynucleotides and primary constructs having sequences that are partially or substantially not translatable, e.g., having a noncoding region. Such noncoding region may be the "first flanking region" of the primary construct. Alternatively, the noncoding region may be a region other than the first region. Such molecules are generally not translated, but can exert an effect on protein

production by one or more of binding to and sequestering one or more translational machinery components such as a ribosomal protein or a transfer RNA (tRNA), thereby effectively reducing protein expression in the cell or modulating one or more pathways or cascades in a cell which in turn alters protein levels. The polynucleotide or primary construct may contain or encode one or more long noncoding RNA (lncRNA, or lincRNA) or portion thereof, a small nucleolar RNA (sno-RNA), micro RNA (miRNA), small interfering RNA (siRNA) or Piwi-interacting RNA (piRNA).

Polypeptide of interest (CRISPR related proteins)

[0059] According to the present invention, the synthetic polynucleotide is designed to encode one or more polypeptides of interest or fragments thereof. The polypeptides of interest include CRISPR related proteins, e.g., CAS9, dCAS9, and dCAS9-effector domain (activator and /or inhibitor domain) fusion proteins. CRISPR related proteins are described in more detail herein.

[0060] A polypeptide of interest may include, but is not limited to, whole polypeptides, a plurality of polypeptides or fragments of polypeptides, which independently may be encoded by one or more nucleic acids, a plurality of nucleic acids, fragments of nucleic acids or variants of any of the aforementioned. As used herein, the term “polypeptides of interest” refers to any polypeptides which are selected to be encoded in the primary construct of the present invention. As used herein, “polypeptide” means a polymer of amino acid residues (natural or unnatural) linked together most often by peptide bonds. The term, as used herein, refers to proteins, polypeptides, and peptides of any size, structure, or function. In some instances the polypeptide encoded is smaller than about 50 amino acids and the polypeptide is then termed a peptide. If the polypeptide is a peptide, it will be at least about 2, 3, 4, or at least 5 amino acid residues long. Thus, polypeptides include gene products, naturally occurring polypeptides, synthetic polypeptides, homologs, orthologs, paralogs, fragments and other equivalents, variants, and analogs of the foregoing. A polypeptide may be a single molecule or may be a multi-molecular complex such as a dimer, trimer or tetramer. They may also comprise single chain or multichain polypeptides such as antibodies or insulin and may be associated or linked. Most commonly disulfide linkages are found in multichain polypeptides. The term polypeptide may also apply to amino acid polymers in which one or more amino acid residues are an artificial chemical analogue of a corresponding naturally occurring amino acid.

[0061] The term “polypeptide variant” refers to molecules which differ in their amino acid sequence from a native or reference sequence. The amino acid sequence variants may possess substitutions, deletions, and/or insertions at certain positions within the amino acid sequence, as compared to a native or reference sequence. Ordinarily, variants will possess at least about 50% identity (homology) to a native or reference sequence, and preferably, they will be at least about 80%, more preferably at least about 90% identical (homologous) to a native or reference sequence.

[0062] In some embodiments “variant mimics” are provided. As used herein, the term “variant mimic” is one which contains one or more amino acids which would mimic an activated sequence. For example, glutamate may serve as a mimic for phospho-threonine and/or phospho-serine. Alternatively, variant mimics may result in deactivation or in an inactivated product containing the mimic, e.g., phenylalanine may act as an inactivating substitution for tyrosine; or alanine may act as an inactivating substitution for serine.

[0063] “Homology” as it applies to amino acid sequences is defined as the percentage of residues in the candidate amino acid sequence that are identical with the residues in the amino acid sequence of a second sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent homology. Methods and computer programs for the alignment are well known in the art. It is understood that homology depends on a calculation of percent identity but may differ in value due to gaps and penalties introduced in the calculation.

[0064] By “homologs” as it applies to polypeptide sequences means the corresponding sequence of other species having substantial identity to a second sequence of a second species.

[0065] “Analog” is meant to include polypeptide variants which differ by one or more amino acid alterations, e.g., substitutions, additions or deletions of amino acid residues that still maintain one or more of the properties of the parent or starting polypeptide.

[0066] The present invention contemplates several types of compositions which are polypeptide based including variants and derivatives. These include substitutional, insertional, deletion and covalent variants and derivatives. The term “derivative” is used synonymously with the term “variant” but generally refers to a molecule that has been modified and/or changed in any way relative to a reference molecule or starting molecule.

[0067] As such, mRNA encoding polypeptides containing substitutions, insertions and/or additions, deletions and covalent modifications with respect to reference sequences, in particular the polypeptide sequences disclosed herein, are included within the scope of this invention. For example, sequence tags or amino acids, such as one or more lysines, can be added to the peptide sequences of the invention (e.g., at the N-terminal or C-terminal ends). Sequence tags can be used for peptide purification or localization. Lysines can be used to increase peptide solubility or to allow for biotinylation. Alternatively, amino acid residues located at the carboxy and amino terminal regions of the amino acid sequence of a peptide or protein may optionally be deleted providing for truncated sequences. Certain amino acids (e.g., C-terminal or N-terminal residues) may alternatively be deleted depending on the use of the sequence, as for example, expression of the sequence as part of a larger sequence which is soluble, or linked to a solid support.

[0068] “Substitutional variants” when referring to polypeptides are those that have at least one amino acid residue in a native or starting sequence removed and a different amino acid inserted in its place at the same position. The substitutions may be single, where only one amino acid in the molecule has been substituted, or they may be multiple, where two or more amino acids have been substituted in the same molecule.

[0069] As used herein the term “conservative amino acid substitution” refers to the substitution of an amino acid that is normally present in the sequence with a different amino acid of similar size, charge, or polarity. Examples of conservative substitutions include the substitution of a non-polar (hydrophobic) residue such as isoleucine, valine and leucine for another non-polar residue. Likewise, examples of conservative substitutions include the substitution of one polar (hydrophilic) residue for another such as between arginine and lysine, between glutamine and asparagine, and between glycine and serine. Additionally, the substitution of a basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue such as aspartic acid or glutamic acid for another acidic residue are additional examples of conservative substitutions. Examples of non-conservative substitutions include the substitution of a non-polar (hydrophobic) amino acid residue such as isoleucine, valine, leucine, alanine, methionine for a polar (hydrophilic) residue such as cysteine, glutamine, glutamic acid or lysine and/or a polar residue for a non-polar residue.

[0070] “Insertional variants” when referring to polypeptides are those with one or more amino acids inserted immediately adjacent to an amino acid at a particular position

in a native or starting sequence. “Immediately adjacent” to an amino acid means connected to either the alpha-carboxy or alpha-amino functional group of the amino acid.

[0071] “Deletional variants” when referring to polypeptides are those with one or more amino acids in the native or starting amino acid sequence removed. Ordinarily, deletional variants will have one or more amino acids deleted in a particular region of the molecule.

[0072] “Covalent derivatives” when referring to polypeptides include modifications of a native or starting protein with an organic proteinaceous or non-proteinaceous derivatizing agent, and/or post-translational modifications. Covalent modifications are traditionally introduced by reacting targeted amino acid residues of the protein with an organic derivatizing agent that is capable of reacting with selected side-chains or terminal residues, or by harnessing mechanisms of post-translational modifications that function in selected recombinant host cells. The resultant covalent derivatives are useful in programs directed at identifying residues important for biological activity, for immunoassays, or for the preparation of anti-protein antibodies for immunoaffinity purification of the recombinant glycoprotein. Such modifications are within the ordinary skill in the art and are performed without undue experimentation.

[0073] Certain post-translational modifications are the result of the action of recombinant host cells on the expressed polypeptide. Glutaminy and asparaginy residues are frequently post-translationally deamidated to the corresponding glutamyl and aspartyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues may be present in the polypeptides produced in accordance with the present invention.

[0074] Other post-translational modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the alpha-amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)).

[0075] “Features” when referring to polypeptides are defined as distinct amino acid sequence-based components of a molecule. Features of the polypeptides encoded by the mmRNA of the present invention include surface manifestations, local conformational shape, folds, loops, half-loops, domains, half-domains, sites, termini or any combination thereof.

[0076] As used herein when referring to polypeptides the term “surface manifestation” refers to a polypeptide based component of a protein appearing on an outermost surface.

[0077] As used herein when referring to polypeptides the term “local conformational shape” means a polypeptide based structural manifestation of a protein which is located within a definable space of the protein.

[0078] As used herein when referring to polypeptides the term “fold” refers to the resultant conformation of an amino acid sequence upon energy minimization. A fold may occur at the secondary or tertiary level of the folding process. Examples of secondary level folds include beta sheets and alpha helices. Examples of tertiary folds include domains and regions formed due to aggregation or separation of energetic forces. Regions formed in this way include hydrophobic and hydrophilic pockets, and the like.

[0079] As used herein the term “turn” as it relates to protein conformation means a bend which alters the direction of the backbone of a peptide or polypeptide and may involve one, two, three or more amino acid residues.

[0080] As used herein when referring to polypeptides the term “loop” refers to a structural feature of a polypeptide which may serve to reverse the direction of the backbone of a peptide or polypeptide. Where the loop is found in a polypeptide and only alters the direction of the backbone, it may comprise four or more amino acid residues. Oliva et al. have identified at least 5 classes of protein loops (J. Mol Biol 266 (4): 814-830; 1997). Loops may be open or closed. Closed loops or “cyclic” loops may comprise 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acids between the bridging moieties. Such bridging moieties may comprise a cysteine-cysteine bridge (Cys-Cys) typical in polypeptides having disulfide bridges or alternatively bridging moieties may be non-protein based such as the dibromozylyl agents used herein.

[0081] As used herein when referring to polypeptides the term “half-loop” refers to a portion of an identified loop having at least half the number of amino acid residues as the loop from which it is derived. It is understood that loops may not always contain an even number of amino acid residues. Therefore, in those cases where a loop contains or is identified to comprise an odd number of amino acids, a half-loop of the odd-numbered loop will comprise the whole number portion or next whole number portion of the loop (number of amino acids of the loop/2+/-0.5 amino acids). For example, a loop identified as a 7 amino acid loop could produce half-loops of 3 amino acids or 4 amino acids ($7/2=3.5\pm 0.5$ being 3 or 4).

[0082] As used herein when referring to polypeptides the term “domain” refers to a motif of a polypeptide having one or more identifiable structural or functional characteristics or properties (e.g., binding capacity, serving as a site for protein-protein interactions).

[0083] As used herein when referring to polypeptides the term “half-domain” means a portion of an identified domain having at least half the number of amino acid residues as the domain from which it is derived. It is understood that domains may not always contain an even number of amino acid residues. Therefore, in those cases where a domain contains or is identified to comprise an odd number of amino acids, a half-domain of the odd-numbered domain will comprise the whole number portion or next whole number portion of the domain (number of amino acids of the domain/2+/-0.5 amino acids). For example, a domain identified as a 7 amino acid domain could produce half-domains of 3 amino acids or 4 amino acids ($7/2=3.5\pm 0.5$ being 3 or 4). It is also understood that sub-domains may be identified within domains or half-domains, these subdomains possessing less than all of the structural or functional properties identified in the domains or half domains from which they were derived. It is also understood that the amino acids that comprise any of the domain types herein need not be contiguous along the backbone of the polypeptide (i.e., nonadjacent amino acids may fold structurally to produce a domain, half-domain or subdomain).

[0084] As used herein when referring to polypeptides the terms “site” as it pertains to amino acid based embodiments is used synonymously with “amino acid residue” and “amino acid side chain.” A site represents a position within a peptide or polypeptide that may be modified, manipulated, altered, derivatized or varied within the polypeptide based molecules of the present invention.

[0085] As used herein the terms “termini” or “terminus” when referring to polypeptides refers to an extremity of a peptide or polypeptide. Such extremity is not limited only to the first or final site of the peptide or polypeptide but may include additional amino acids in the terminal regions. The polypeptide based molecules of the present invention may be characterized as having both an N-terminus (terminated by an amino acid with a free amino group (NH₂)) and a C-terminus (terminated by an amino acid with a free carboxyl group (COOH)). Proteins of the invention are in some cases made up of multiple polypeptide chains brought together by disulfide bonds or by non-covalent forces (multimers, oligomers). These sorts of proteins will have multiple N- and C-termini. Alternatively, the termini of the polypeptides may be modified such that they

begin or end, as the case may be, with a non-polypeptide based moiety such as an organic conjugate.

[0086] Once any of the features have been identified or defined as a desired component of a polypeptide to be encoded by the primary construct or mmRNA of the invention, any of several manipulations and/or modifications of these features may be performed by moving, swapping, inverting, deleting, randomizing or duplicating. Furthermore, it is understood that manipulation of features may result in the same outcome as a modification to the molecules of the invention. For example, a manipulation which involved deleting a domain would result in the alteration of the length of a molecule just as modification of a nucleic acid to encode less than a full length molecule would.

[0087] Modifications and manipulations can be accomplished by methods known in the art such as, but not limited to, site directed mutagenesis. The resulting modified molecules may then be tested for activity using *in vitro* or *in vivo* assays such as those described herein or any other suitable screening assay known in the art.

[0088] According to the present invention, the polypeptides may comprise a consensus sequence which is discovered through rounds of experimentation. As used herein a "consensus" sequence is a single sequence which represents a collective population of sequences allowing for variability at one or more sites.

[0089] As recognized by those skilled in the art, protein fragments, functional protein domains, and homologous proteins are also considered to be within the scope of polypeptides of interest of this invention. For example, provided herein is any protein fragment (meaning a polypeptide sequence at least one amino acid residue shorter than a reference polypeptide sequence but otherwise identical) of a reference protein 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or greater than 100 amino acids in length. In another example, any protein that includes a stretch of about 20, about 30, about 40, about 50, or about 100 amino acids which are about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 95%, or about 100% identical to any of the sequences described herein can be utilized in accordance with the invention. In certain embodiments, a polypeptide to be utilized in accordance with the invention includes 2, 3, 4, 5, 6, 7, 8, 9, 10, or more mutations as shown in any of the sequences provided or referenced herein.

Variant Polypeptides of Interest

[0090] In one embodiment, synthetic polypeptides of the invention may encode variant polypeptides, e.g., variant CRISPR related proteins which have a certain identity with a reference polypeptide sequence. As used herein, a “reference polypeptide sequence” refers to a starting polypeptide sequence. Reference sequences may be wild type sequences or any sequence to which reference is made in the design of another sequence. A “reference polypeptide sequence” may be, e.g., any encoding CAS9, dCAS9, a dCAS9-activator domain fusion protein, a dCAS9-inhibitor domain fusion protein, and/or variants thereof.

[0091] The term “identity” as known in the art, refers to a relationship between the sequences of two or more peptides, as determined by comparing the sequences. In the art, identity also means the degree of sequence relatedness between peptides, as determined by the number of matches between strings of two or more amino acid residues. Identity measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., “algorithms”). Identity of related peptides can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; and Carillo et al., SIAM J. Applied Math. 48, 1073 (1988).

[0092] In some embodiments, the polypeptide variant may have the same or a similar activity as the reference polypeptide. Alternatively, the variant may have an altered activity (e.g., increased or decreased) relative to a reference polypeptide. Generally, variants of a particular polynucleotide or polypeptide of the invention will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% but less than 100% sequence identity to that particular reference polynucleotide or polypeptide as determined by sequence alignment programs and parameters described herein and known to those skilled in the art. Such tools for alignment include those of the BLAST suite (Stephen F. Altschul, Thomas L. Madden, Alejandro A. Schäffer, Jinghui Zhang, Zheng Zhang, Webb Miller, and David J. Lipman

(1997), "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", Nucleic Acids Res. 25:3389-3402.) Other tools are described herein, specifically in the definition of "Identity."

[0093] Default parameters in the BLAST algorithm include, for example, an expect threshold of 10, Word size of 28, Match/Mismatch Scores 1, -2, Gap costs Linear. Any filter can be applied as well as a selection for species specific repeats, e.g., Homo sapiens.

Flanking Regions: Untranslated Regions (UTRs)

[0094] As described herein, the synthetic polynucleotides of the invention include untranslated regions. Untranslated regions (UTRs) of a gene are transcribed but not translated. The 5'UTR starts at the transcription start site and continues to the start codon but does not include the start codon; whereas, the 3'UTR starts immediately following the stop codon and continues until the transcriptional termination signal. There is growing body of evidence about the regulatory roles played by the UTRs in terms of stability of the nucleic acid molecule and translation. The regulatory features of a UTR can be incorporated into the polynucleotides, primary constructs and/or mmRNA of the present invention to enhance the stability of the molecule. The specific features can also be incorporated to ensure controlled down-regulation of the transcript in case they are misdirected to undesired organs sites.

[0095] Examples of UTRs include but are not limited to those found in the Tables below.

[0096] Tables 2 and 3 of co-pending U.S. Provisional Patent Application No 61/737,130 filed December 14, 2012 provide a listing of exemplary UTRs which may be utilized in the primary construct of the present invention as flanking regions. Variants of 5' or 3'UTRs may be utilized wherein one or more nucleotides are added or removed to the termini, including A, T, C or G.

[0097] It should be understood that those listed are examples and that any UTR from any gene may be incorporated into the respective first or second flanking region of the primary construct. Furthermore, multiple wild-type UTRs of any known gene may be utilized. It is also within the scope of the present invention to provide artificial UTRs which are not variants of wild type genes. These UTRs or portions thereof may be placed in the same orientation as in the transcript from which they were selected or may be altered in orientation or location. Hence a 5' or 3' UTR may be inverted, shortened, lengthened, made chimeric with one or more other 5' UTRs or 3' UTRs. As used herein,

the term “altered” as it relates to a UTR sequence, means that the UTR has been changed in some way in relation to a reference sequence. For example, a 3' or 5' UTR may be altered relative to a wild type or native UTR by the change in orientation or location as taught above or may be altered by the inclusion of additional nucleotides, deletion of nucleotides, swapping or transposition of nucleotides. Any of these changes producing an “altered” UTR (whether 3' or 5') comprise a variant UTR.

[0098] In one embodiment, the UTR may be selected from the UTRs described in Lengthy Table 21 in co-pending US Provisional Application No. US 61/775,509, filed March 9, 2013, entitled Heterologous Untranslated Regions for mRNA and in Lengthy Table 21 and in Table 22 in co-pending US Provisional Application No. US 61/829,372, filed May 31, 2013, entitled Heterologous Untranslated Regions for mRNA; each of which is herein incorporated by reference in its entirety.

[0099] In one embodiment, a double, triple or quadruple UTR such as a 5' or 3' UTR may be used. As used herein, a “double” UTR is one in which two copies of the same UTR are encoded either in series or substantially in series. For example, a double beta-globin 3' UTR may be used as described in US Patent publication 20100129877, the contents of which are incorporated herein by reference in its entirety.

[00100] It is also within the scope of the present invention to have patterned UTRs. As used herein “patterned UTRs” are those UTRs which reflect a repeating or alternating pattern, such as ABABAB or AABBAABBAABB or ABCABCABC or variants thereof repeated once, twice, or more than 3 times. In these patterns, each letter, A, B, or C represent a different UTR at the nucleotide level.

[00101] In one embodiment, flanking regions are selected from a family of transcripts whose proteins share a common function, structure, feature of property. For example, polypeptides of interest may belong to a family of proteins which are expressed in a particular cell, tissue or at some time during development. The UTRs from any of these genes may be swapped for any other UTR of the same or different family of proteins to create a new chimeric primary transcript. As used herein, a “family of proteins” is used in the broadest sense to refer to a group of two or more polypeptides of interest which share at least one function, structure, feature, localization, origin, or expression pattern.

[00102] In one embodiment, the synthetic polynucleotides of the invention may comprise an untranslated region that is not heterologous to the encoded protein of interest. As a non-limiting example, the untranslated region may comprise all or a portion or fragment thereof of one or more of the untranslated regions described in co-pending US

Provisional Application No. US 61/775,509, filed March 9, 2013, entitled Heterologous Untranslated Regions for mRNA and US Provisional Application No. US 61/829,372, filed March 15, 2013, entitled Heterologous Untranslated Regions for mRNA, each of which is herein incorporated by reference in its entirety.

5' UTR and Translation Initiation

[00103] The synthetic polynucleotides of the invention typically include a 5' UTR. Natural 5'UTRs bear features which play roles in for translation initiation. They harbor signatures like Kozak sequences which are commonly known to be involved in the process by which the ribosome initiates translation of many genes. Kozak sequences have the consensus CCR(A/G)CCAUGG (SEQ ID NO: 126), where R is a purine (adenine or guanine) three bases upstream of the start codon (AUG), which is followed by another 'G'. 5'UTR also have been known to form secondary structures which are involved in elongation factor binding.

[00104] By engineering the features typically found in abundantly expressed genes of specific target organs, one can enhance the stability and protein production of the synthetic polynucleotides, primary constructs or mmRNA of the invention. For example, introduction of 5' UTR of liver-expressed mRNA, such as albumin, serum amyloid A, Apolipoprotein A/B/E, transferrin, alpha fetoprotein, erythropoietin, or Factor VIII, could be used to enhance expression of a nucleic acid molecule, such as a mmRNA, in hepatic cell lines or liver. Likewise, use of 5' UTR from other tissue-specific mRNA to improve expression in that tissue is possible – for muscle (MyoD, Myosin, Myoglobin, Myogenin, Herculin), for endothelial cells (Tie-1, CD36), for myeloid cells (C/EBP, AML1, G-CSF, GM-CSF, CD11b, MSR, Fr-1, i-NOS), for leukocytes (CD45, CD18), for adipose tissue (CD36, GLUT4, ACRP30, adiponectin) and for lung epithelial cells (SP-A/B/C/D).

[00105] Other non-UTR sequences may be incorporated into the 5' (or 3' UTR) UTRs. For example, introns or portions of introns sequences may be incorporated into the flanking regions of the polynucleotides, primary constructs or mmRNA of the invention. Incorporation of intronic sequences may increase protein production as well as mRNA levels.

[00106] The 5'UTR may selected for use in the present invention may be a structured UTR such as, but not limited to, 5'UTRs to control translation. As a non-limiting example, a structured 5'UTR may be beneficial when using any of the terminal modifications described in copending U.S. Provisional Application No. 61/758,921 filed

January 31, 2013, entitled Differential Targeting Using RNA Constructs; U.S. Provisional Application No. 61/781,139 filed March 14, 2013, entitled Differential Targeting Using RNA Constructs; U.S. Provisional Application No. 61/729,933, filed November 26, 2012 entitled Terminally Optimized RNAs; U.S. Provisional Application No 61/737,224 filed December 14, 2012 entitled Terminally Optimized RNAs and U.S. Provisional Application No 61/829,359 filed May 31, 2013 entitled Terminally Optimized RNAs, each of which is herein incorporated by reference in their entirety.

3' UTR and the AU Rich Elements

[00107] The synthetic polynucleotides of the invention typically include a 3' UTR. 3'UTRs are known to have stretches of Adenosines and Uridines embedded in them. These AU rich signatures are particularly prevalent in genes with high rates of turnover. Based on their sequence features and functional properties, the AU rich elements (AREs) can be separated into three classes (Chen et al, 1995): Class I AREs contain several dispersed copies of an AUUUA motif within U-rich regions. C-Myc and MyoD contain class I AREs. Class II AREs possess two or more overlapping UUAUUUA(U/A)(U/A) nonamers. Molecules containing this type of AREs include GM-CSF and TNF- α . Class III AREs are less well defined. These U rich regions do not contain an AUUUA motif. c-Jun and Myogenin are two well-studied examples of this class. Most proteins binding to the AREs are known to destabilize the messenger, whereas members of the ELAV family, most notably HuR, have been documented to increase the stability of mRNA. HuR binds to AREs of all the three classes. Engineering the HuR specific binding sites into the 3' UTR of nucleic acid molecules will lead to HuR binding and thus, stabilization of the message *in vivo*.

[00108] Introduction, removal or modification of 3' UTR AU rich elements (AREs) can be used to modulate the stability of polynucleotides, primary constructs or mmRNA of the invention. When engineering specific polynucleotides, primary constructs or mmRNA, one or more copies of an ARE can be introduced to make polynucleotides, primary constructs or mmRNA of the invention less stable and thereby curtail translation and decrease production of the resultant protein. Likewise, AREs can be identified and removed or mutated to increase the intracellular stability and thus increase translation and production of the resultant protein. Transfection experiments can be conducted in relevant cell lines, using polynucleotides, primary constructs or mmRNA of the invention and protein production can be assayed at various time points post-transfection. For

example, cells can be transfected with different ARE-engineering molecules and by using an ELISA kit to the relevant protein and assaying protein produced at 6 hr, 12 hr, 24 hr, 48 hr, and 7 days post-transfection.

3' UTR and microRNA Binding Sites

[00109] In some embodiments, the synthetic polynucleotides of the invention include a miRNA binding site in the 3'UTR. A microRNA (or miRNA) is a 19-25 nucleotide long noncoding RNA that binds to the 3'UTR of nucleic acid molecules and down-regulate gene expression either by reducing nucleic acid molecule stability or by inhibiting translation. The polynucleotides, primary constructs or mmRNA of the invention may comprise one or more microRNA target sequences, microRNA sequences, microRNA binding sites, or microRNA seeds. Such sequences may correspond to any known microRNA such as those taught in US Publication US2005/0261218 and US Publication US2005/0059005, or those listed in Table 7 of co-pending application USSN 61/758,921 filed January 31, 2013 (Attorney Docket Number 2030.1039), the contents of which are incorporated herein by reference in their entirety.

[00110] Examples of 3'UTRs containing miRNA binding sites include but are not limited to those found in the Tables below.

[00111] A microRNA sequence comprises a "seed" region, i.e., a sequence in the region of positions 2-8 of the mature microRNA, which sequence has perfect Watson-Crick complementarity to the miRNA target sequence. A microRNA seed may comprise positions 2-8 or 2-7 of the mature microRNA. In some embodiments, a microRNA seed may comprise 7 nucleotides (e.g., nucleotides 2-8 of the mature microRNA), wherein the seed-complementary site in the corresponding miRNA target is flanked by an adenine (A) opposed to microRNA position 1. In some embodiments, a microRNA seed may comprise 6 nucleotides (e.g., nucleotides 2-7 of the mature microRNA), wherein the seed-complementary site in the corresponding miRNA target is flanked by an adenine (A) opposed to microRNA position 1. See for example, Grimson A, Farh KK, Johnston WK, Garrett-Engele P, Lim LP, Bartel DP; Mol Cell. 2007 Jul 6;27(1):91-105. The bases of the microRNA seed have complete complementarity with the target sequence. By engineering microRNA target sequences into the 3'UTR of polynucleotides, primary constructs or mmRNA of the invention one can target the molecule for degradation or reduced translation, provided the microRNA in question is available. This process will reduce the hazard of off target effects upon nucleic acid molecule delivery.

Identification of microRNA, microRNA target regions, and their expression patterns and role in biology have been reported (Bonauer et al., *Curr Drug Targets* 2010 11:943-949; Anand and Cheresch *Curr Opin Hematol* 2011 18:171-176; Contreras and Rao *Leukemia* 2012 26:404-413 (2011 Dec 20. doi: 10.1038/leu.2011.356); Bartel *Cell* 2009 136:215-233; Landgraf et al, *Cell*, 2007 129:1401-1414).

[00112] For example, if the nucleic acid molecule is an mRNA and is not intended to be delivered to the liver but ends up there, then miR-122, a microRNA abundant in liver, can inhibit the expression of the gene of interest if one or multiple target sites of miR-122 are engineered into the 3'UTR of the polynucleotides, primary constructs or mmRNA.

Introduction of one or multiple binding sites for different microRNA can be engineered to further decrease the longevity, stability, and protein translation of a polynucleotides, primary constructs or mmRNA.

[00113] As used herein, the term "microRNA site" refers to a microRNA target site or a microRNA recognition site, or any nucleotide sequence to which a microRNA binds or associates. It should be understood that "binding" may follow traditional Watson-Crick hybridization rules or may reflect any stable association of the microRNA with the target sequence at or adjacent to the microRNA site.

[00114] Conversely, for the purposes of the polynucleotides, primary constructs or mmRNA of the present invention, microRNA binding sites can be engineered out of (i.e. removed from) sequences in which they naturally occur in order to increase protein expression in specific tissues. For example, miR-122 binding sites may be removed to improve protein expression in the liver. Regulation of expression in multiple tissues can be accomplished through introduction or removal of one or several microRNA binding sites.

[00115] Examples of tissues where microRNA are known to regulate mRNA, and thereby protein expression, include, but are not limited to, liver (miR-122), muscle (miR-133, miR-206, miR-208), endothelial cells (miR-17-92, miR-126), nervous system (miR-124a, miR-9), pluripotent cells (miR-302, miR-367, miR-290, miR-371, miR-373), pancreatic islet cells (miR-375), myeloid cells (miR-142-3p, miR-142-5p, miR-16, miR-21, miR-223, miR-24, miR-27), adipose tissue (let-7, miR-30c), heart (miR-1d, miR-149), kidney (miR-192, miR-194, miR-204), and lung epithelial cells (let-7, miR-133, miR-126).

[00116] MicroRNA can also regulate complex biological processes such as angiogenesis (miR-132) (Anand and Cheresch *Curr Opin Hematol* 2011 18:171-176). In

the polynucleotides, primary constructs or mmRNA of the invention, binding sites for microRNAs that are involved in such processes may be removed or introduced, in order to tailor the expression of the polynucleotides, primary constructs or mmRNA expression to biologically relevant cell types or to the context of relevant biological processes.

[00117] MicroRNAs are also known to be expressed in the immune cells (also called hematopoietic cells), for example, the antigen presenting cells (e.g. dendritic cells and macrophage). Immune cell specific microRNAs are involved in immunogenicity, autoimmunity, the immune -response to infection, inflammation, as well as unwanted immune response after gene therapy and tissue/organ transplantation. For example, miR-142 and miR-146 are exclusively expressed in the immune cells, particularly abundant in myeloid dendritic cells. Introducing the miR-142 binding site into the 3'-UTR of a polynucleotide or a gene delivery construct can selectively suppress the gene expression in the antigen presenting cells through miR-142 mediated mRNA degradation, limiting antigen presentation in professional APCs (e.g. dendritic cells) and thereby preventing antigen-mediated immune response after gene delivery (see, Annoni A et al., blood, 2009, 114, 5152-5161, the content of which is herein incorporated by reference in its entirety.). In the polynucleotides, primary constructs or mmRNA of the present invention, binding sites for microRNAs that are involved in such processes may be introduced, in order to reduce the expression of the polynucleotides, primary constructs or mmRNA of the present invention in APCs and to subdue the antigen mediated immune response.

[00118] Lastly, through an understanding of the expression patterns of microRNA in different cell types, polynucleotides, primary constructs or mmRNA can be engineered for more targeted expression in specific cell types or only under specific biological conditions. Through introduction of tissue-specific microRNA binding sites, polynucleotides, primary constructs or mmRNA could be designed that would be optimal for protein expression in a tissue or in the context of a biological condition.

[00119] Transfection experiments can be conducted in relevant cell lines, using engineered polynucleotides, primary constructs or mmRNA and protein production can be assayed at various time points post-transfection. For example, cells can be transfected with different microRNA binding site-engineering polynucleotides, primary constructs or mmRNA and by using an ELISA kit to the relevant protein and assaying protein produced at 6 hr, 12 hr, 24 hr, 48 hr, 72 hr and 7 days post-transfection. *In vivo* experiments can also be conducted using microRNA-binding site-engineered molecules to examine

changes in tissue-specific expression of formulated polynucleotides, primary constructs or mmRNA.

Viral Sequences

[00120] Additional viral sequences such as, but not limited to, the translation enhancer sequence of the barley yellow dwarf virus (BYDV-PAV) can be engineered and inserted in the 3' UTR of the synthetic polynucleotides, polynucleotides, primary constructs or mmRNA of the invention and can stimulate the translation of the construct *in vitro* and *in vivo*. Transfection experiments can be conducted in relevant cell lines at and protein production can be assayed by ELISA at 12hr, 24hr, 48hr, 72 hr and day 7 post-transfection.

[00121]

5' Capping

[00122] The synthetic polynucleotides of the invention can include a 5' capping region or 5' cap. The 5' cap structure of an mRNA is involved in nuclear export, increasing mRNA stability and binds the mRNA Cap Binding Protein (CBP), which is responsible for mRNA stability in the cell and translation competency through the association of CBP with poly(A) binding protein to form the mature cyclic mRNA species. The cap further assists the removal of 5' proximal introns removal during mRNA splicing.

[00123] Endogenous mRNA molecules may be 5'-end capped generating a 5'-ppp-5'-triphosphate linkage between a terminal guanosine cap residue and the 5'-terminal transcribed sense nucleotide of the mRNA molecule. This 5'-guanylate cap may then be methylated to generate an N7-methyl-guanylate residue. The ribose sugars of the terminal and/or antiterminal transcribed nucleotides of the 5' end of the mRNA may optionally also be 2'-O-methylated. 5'-decapping through hydrolysis and cleavage of the guanylate cap structure may target a nucleic acid molecule, such as an mRNA molecule, for degradation.

[00124] Modifications to the polynucleotides, primary constructs, and mmRNA of the present invention may generate a non-hydrolyzable cap structure preventing decapping and thus increasing mRNA half-life. Because cap structure hydrolysis requires cleavage of 5'-ppp-5' phosphodiester linkages, modified nucleotides may be used during the capping reaction. For example, a Vaccinia Capping Enzyme from New England Biolabs (Ipswich, MA) may be used with α -thio-guanosine nucleotides according to the manufacturer's instructions to create a phosphorothioate linkage in the 5'-ppp-5' cap.

Additional modified guanosine nucleotides may be used such as α -methyl-phosphonate and seleno-phosphate nucleotides.

[00125] Additional modifications include, but are not limited to, 2'-O-methylation of the ribose sugars of 5'-terminal and/or 5'-antiterminal nucleotides of the mRNA (as mentioned above) on the 2'-hydroxyl group of the sugar ring. Multiple distinct 5'-cap structures can be used to generate the 5'-cap of a nucleic acid molecule, such as an mRNA molecule.

[00126] Cap analogs, which herein are also referred to as synthetic cap analogs, chemical caps, chemical cap analogs, or structural or functional cap analogs, differ from natural (i.e. endogenous, wild-type or physiological) 5'-caps in their chemical structure, while retaining cap function. Cap analogs may be chemically (i.e. non-enzymatically) or enzymatically synthesized and/linked to a nucleic acid molecule.

[00127] For example, the Anti-Reverse Cap Analog (ARCA) cap contains two guanines linked by a 5'-5'-triphosphate group, wherein one guanine contains an N7 methyl group as well as a 3'-O-methyl group (i.e., N7,3'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine (m^7G -3'mppp-G; which may equivalently be designated 3' O-Me- $m^7G(5')$ ppp(5')G). The 3'-O atom of the other, unmodified, guanine becomes linked to the 5'-terminal nucleotide of the capped nucleic acid molecule (e.g. an mRNA or mmRNA). The N7- and 3'-O-methylated guanine provides the terminal moiety of the capped nucleic acid molecule (e.g. mRNA or mmRNA).

[00128] Another exemplary cap is mCAP, which is similar to ARCA but has a 2'-O-methyl group on guanosine (i.e., N7,2'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine, m^7Gm -ppp-G).

[00129] While cap analogs allow for the concomitant capping of a nucleic acid molecule in an in vitro transcription reaction, up to 20% of transcripts remain uncapped. This, as well as the structural differences of a cap analog from an endogenous 5'-cap structures of nucleic acids produced by the endogenous, cellular transcription machinery, may lead to reduced translational competency and reduced cellular stability.

[00130] Polynucleotides, primary constructs and mmRNA of the invention may also be capped post-transcriptionally, using enzymes, in order to generate more authentic 5'-cap structures. As used herein, the phrase "more authentic" refers to a feature that closely mirrors or mimics, either structurally or functionally, an endogenous or wild type feature. That is, a "more authentic" feature is better representative of an endogenous, wild-type, natural or physiological cellular function and/or structure as compared to synthetic

features or analogs, etc., of the prior art, or which outperforms the corresponding endogenous, wild-type, natural or physiological feature in one or more respects. Non-limiting examples of more authentic 5' cap structures of the present invention are those which, among other things, have enhanced binding of cap binding proteins, increased half life, reduced susceptibility to 5' endonucleases and/or reduced 5' decapping, as compared to synthetic 5' cap structures known in the art (or to a wild-type, natural or physiological 5' cap structure). For example, recombinant Vaccinia Virus Capping Enzyme and recombinant 2'-O-methyltransferase enzyme can create a canonical 5'-5'-triphosphate linkage between the 5'-terminal nucleotide of an mRNA and a guanine cap nucleotide wherein the cap guanine contains an N7 methylation and the 5'-terminal nucleotide of the mRNA contains a 2'-O-methyl. Such a structure is termed the Cap1 structure. This cap results in a higher translational-competency and cellular stability and a reduced activation of cellular pro-inflammatory cytokines, as compared, e.g., to other 5' cap analog structures known in the art. Cap structures include 7mG(5')ppp(5')N,pN2p (cap 0), 7mG(5')ppp(5')NlmpNp (cap 1), and 7mG(5')-ppp(5')NlmpN2mp (cap 2).

[00131] Because the polynucleotides, primary constructs or mmRNA may be capped post-transcriptionally, and because this process is more efficient, nearly 100% of the polynucleotides, primary constructs or mmRNA may be capped. This is in contrast to ~80% when a cap analog is linked to an mRNA in the course of an in vitro transcription reaction.

[00132] According to the present invention, 5' terminal caps may include endogenous caps or cap analogs. According to the present invention, a 5' terminal cap may comprise a guanine analog. Useful guanine analogs include inosine, N1-methyl-guanosine, 2'-fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, and 2-azido-guanosine.

IRES Sequences

[00133] Further, provided are synthetic polynucleotides, primary constructs or mmRNA which may contain an internal ribosome entry site (IRES). First identified as a feature Picorna virus RNA, IRES plays an important role in initiating protein synthesis in absence of the 5' cap structure. An IRES may act as the sole ribosome binding site, or may serve as one of multiple ribosome binding sites of an mRNA. Synthetic polynucleotides, primary constructs or mmRNA containing more than one functional ribosome binding site may encode several peptides or polypeptides that are translated

independently by the ribosomes (“multicistronic nucleic acid molecules”). When polynucleotides, primary constructs or mmRNA are provided with an IRES, further optionally provided is a second translatable region. Examples of IRES sequences that can be used according to the invention include without limitation, those from picornaviruses (e.g. FMDV), pest viruses (CFFV), polio viruses (PV), encephalomyocarditis viruses (ECMV), foot-and-mouth disease viruses (FMDV), hepatitis C viruses (HCV), classical swine fever viruses (CSFV), murine leukemia virus (MLV), simian immune deficiency viruses (SIV) or cricket paralysis viruses (CrPV).

Stop Codons

[00134] In one embodiment, the primary constructs of the present invention may include at least two stop codons before the 3' untranslated region (UTR). The stop codon may be selected from TGA, TAA and TAG. In one embodiment, the primary constructs of the present invention include the stop codon TGA and one additional stop codon. In a further embodiment the additional stop codon may be TAA.

[00135] In another embodiment, the primary constructs of the present invention may include three stop codons before the 3' untranslated region (UTR).

Signal Sequences

[00136] The primary constructs or mmRNA may also encode additional features which facilitate trafficking of the polypeptides to therapeutically relevant sites. One such feature which aids in protein trafficking is the signal sequence. As used herein, a “signal sequence” or “signal peptide” is a polynucleotide or polypeptide, respectively, which is from about 9 to 200 nucleotides (3-60 amino acids) in length which is incorporated at the 5' (or N-terminus) of the coding region or polypeptide encoded, respectively. Addition of these sequences result in trafficking of the encoded polypeptide to the endoplasmic reticulum through one or more secretory pathways. Some signal peptides are cleaved from the protein by signal peptidase after the proteins are transported.

[00137] Signal sequences may be selected from any of those listed in co-pending patent applications: U.S. Provisional Patent Application No 61/618,862, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Biologics; U.S. Provisional Patent Application No 61/681,645, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Biologics; U.S. Provisional Patent Application No 61/737,130, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Biologics; U.S. Provisional Patent Application No 61/618,866, filed April

2, 2012, entitled Modified Polynucleotides for the Production of Antibodies; U.S. Provisional Patent Application No 61/681,647, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Antibodies; U.S. Provisional Patent Application No 61/737,134, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Antibodies; U.S. Provisional Patent Application No 61/618,868, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Vaccines; U.S. Provisional Patent Application No 61/681,648, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Vaccines; U.S. Provisional Patent Application No 61/737,135, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Vaccines; U.S. Provisional Patent Application No 61/618,870, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Therapeutic Proteins and Peptides; U.S. Provisional Patent Application No 61/681,649, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Therapeutic Proteins and Peptides; U.S. Provisional Patent Application No 61/737,139, filed December 14, 2012, Modified Polynucleotides for the Production of Therapeutic Proteins and Peptides; U.S. Provisional Patent Application No 61/618,873, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Secreted Proteins; U.S. Provisional Patent Application No 61/681,650, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Secreted Proteins; U.S. Provisional Patent Application No 61/737,147, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Secreted Proteins; U.S. Provisional Patent Application No 61/618,878, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Plasma Membrane Proteins; U.S. Provisional Patent Application No 61/681,654, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Plasma Membrane Proteins; U.S. Provisional Patent Application No 61/737,152, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Plasma Membrane Proteins; U.S. Provisional Patent Application No 61/618,885, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Cytoplasmic and Cytoskeletal Proteins; U.S. Provisional Patent Application No 61/681,658, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Cytoplasmic and Cytoskeletal Proteins; U.S. Provisional Patent Application No 61/737,155, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Cytoplasmic and Cytoskeletal Proteins; U.S. Provisional Patent Application No 61/618,896, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Intracellular Membrane Bound Proteins; U.S.

Provisional Patent Application No 61/668,157, filed July 5, 2012, entitled Modified Polynucleotides for the Production of Intracellular Membrane Bound Proteins; U.S. Provisional Patent Application No 61/681,661, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Intracellular Membrane Bound Proteins; U.S. Provisional Patent Application No 61/737,160, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Intracellular Membrane Bound Proteins; U.S. Provisional Patent Application No 61/618,911, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Nuclear Proteins; U.S. Provisional Patent Application No 61/681,667, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Nuclear Proteins; U.S. Provisional Patent Application No 61/737,168, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Nuclear Proteins; U.S. Provisional Patent Application No 61/618,922, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Proteins; U.S. Provisional Patent Application No 61/681,675, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Proteins; U.S. Provisional Patent Application No 61/737,174, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Proteins; U.S. Provisional Patent Application No 61/618,935, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/681,687, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/737,184, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/618,945, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/681,696, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/737,191, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/618,953, filed April 2, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/681,704, filed August 10, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; U.S. Provisional Patent Application No 61/737,203, filed December 14, 2012, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease,

International Application No PCT/US2013/030062, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Biologics and Proteins Associated with Human Disease; International Application No PCT/US2013/030063, filed March 9, 2013, entitled Modified Polynucleotides; International Application No. PCT/US2013/030064, entitled Modified Polynucleotides for the Production of Secreted Proteins; International Application No PCT/US2013/030059, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Membrane Proteins; International Application No. PCT/US2013/030066, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Cytoplasmic and Cytoskeletal Proteins; International Application No. PCT/US2013/030067, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Nuclear Proteins; International Application No. PCT/US2013/030060, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Proteins; International Application No. PCT/US2013/030061, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Proteins Associated with Human Disease; International Application No. PCT/US2013/030068, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Cosmetic Proteins and Peptides; and International Application No. PCT/US2013/030070, filed March 9, 2013, entitled Modified Polynucleotides for the Production of Oncology-Related Proteins and Peptides; International Patent Application No. PCT/US2013/031821, filed March 15, 2013, entitled In Vivo Production of Proteins, the contents of which are incorporated herein by reference. Protein signal sequences which may be incorporated for encoding by the polynucleotides, primary constructs or mmRNA of the invention include signal sequences from α -1-antitrypsin, G-CSF, Factor IX, Prolactin, Albumin, HMMSP38, ornithine carbamoyltransferase, Cytochrome C Oxidase subunit 8A, Type III, bacterial, viral, secretion signals, Vrg-6, PhoA, OmpA, STI, STII, Amylase, Alpha Factor, Endoglucanase V, Secretion signal, fungal, fibronectin and interleukins (e.g., IL12).

[00138] In the table of the co-pending patent applications, SS is secretion signal and MLS is mitochondrial leader signal. The primary constructs or mmRNA of the present invention may be designed to encode any of the signal sequences or fragments or variants thereof. These sequences may be included at the beginning of the polypeptide coding region, in the middle or at the terminus or alternatively into a flanking region.

[00139] Additional signal sequences which may be utilized in the present invention include those taught in, for example, databases such as those found at <http://www.signalpeptide.de/> or <http://proline.bic.nus.edu.sg/spdb/>. Those described in

US Patents 8,124,379; 7,413,875 and 7,385,034 are also within the scope of the invention and the contents of each are incorporated herein by reference in their entirety.

Protein Cleavage Signals and Sites

[00140] In one embodiment, the polypeptides of the present invention may include at least one protein cleavage signal containing at least one protein cleavage site. The protein cleavage site may be located at the N-terminus, the C-terminus, at any space between the N- and the C- termini such as, but not limited to, half-way between the N- and C-termini, between the N-terminus and the half way point, between the half way point and the C-terminus, and combinations thereof.

[00141] The polypeptides of the present invention may include, but is not limited to, a proprotein convertase (or prohormone convertase), thrombin or Factor Xa protein cleavage signal. Proprotein convertases are a family of nine proteinases, comprising seven basic amino acid-specific subtilisin-like serine proteinases related to yeast kexin, known as prohormone convertase 1/3 (PC1/3), PC2, furin, PC4, PC5/6, paired basic amino-acid cleaving enzyme 4 (PACE4) and PC7, and two other subtilases that cleave at non-basic residues, called subtilisin kexin isozyme 1 (SKI-1) and proproteinconvertase subtilisin kexin 9 (PCSK9).

[00142] In one embodiment, the primary constructs and mmRNA of the present invention may be engineered such that the primary construct or mmRNA contains at least one encoded protein cleavage signal. The encoded protein cleavage signal may be located before the start codon, after the start codon, before the coding region, within the coding region such as, but not limited to, half way in the coding region, between the start codon and the half way point, between the half way point and the stop codon, after the coding region, before the stop codon, between two stop codons, after the stop codon and combinations thereof.

[00143] In one embodiment, the primary constructs or mmRNA of the present invention may include at least one encoded protein cleavage signal containing at least one protein cleavage site. The encoded protein cleavage signal may include, but is not limited to, a proprotein convertase (or prohormone convertase), thrombin and/or Factor Xa protein cleavage signal. One of skill in the art may use Table 1 below or other known methods to determine the appropriate encoded protein cleavage signal to include in the primary constructs or mmRNA of the present invention. For example, starting with a

signal sequence and considering the codons of Table 1 one can design a signal for the primary construct which can produce a protein signal in the resulting polypeptide.

[00144] In one embodiment, the polypeptides of the present invention include at least one protein cleavage signal and/or site.

[00145] As a non-limiting example, U.S. Pat. No. 7,374,930 and U.S. Pub. No. 20090227660, herein incorporated by reference in their entireties, use a furin cleavage site to cleave the N-terminal methionine of GLP-1 in the expression product from the Golgi apparatus of the cells. In one embodiment, the polypeptides of the present invention include at least one protein cleavage signal and/or site with the proviso that the polypeptide is not GLP-1.

[00146] In one embodiment, the primary constructs or mmRNA of the present invention includes at least one encoded protein cleavage signal and/or site.

[00147] In one embodiment, the primary constructs or mmRNA of the present invention includes at least one encoded protein cleavage signal and/or site with the proviso that the primary construct or mmRNA does not encode GLP-1.

[00148] In one embodiment, the primary constructs or mmRNA of the present invention may include more than one coding region. Where multiple coding regions are present in the primary construct or mmRNA of the present invention, the multiple coding regions may be separated by encoded protein cleavage sites. As a non-limiting example, the primary construct or mmRNA may be signed in an ordered pattern. One such pattern follows AXBY form where A and B are coding regions which may be the same or different coding regions and/or may encode the same or different polypeptides, and X and Y are encoded protein cleavage signals which may encode the same or different protein cleavage signals. A second such pattern follows the form AXYBZ where A and B are coding regions which may be the same or different coding regions and/or may encode the same or different polypeptides, and X, Y and Z are encoded protein cleavage signals which may encode the same or different protein cleavage signals. A third pattern follows the form ABXCY where A, B and C are coding regions which may be the same or different coding regions and/or may encode the same or different polypeptides, and X and Y are encoded protein cleavage signals which may encode the same or different protein cleavage signals.

[00149] In one embodiment, the polypeptides, primary constructs and mmRNA can also contain sequences that encode protein cleavage sites so that the polypeptides,

primary constructs and mmRNA can be released from a carrier region or a fusion partner by treatment with a specific protease for said protein cleavage site.

Poly-A tails

[00150] The synthetic polynucleotides of the invention typically include a 3' tailing sequences, e.g., a poly-A tail. During RNA processing, a long chain of adenine nucleotides (poly-A tail) may be added to a polynucleotide such as an mRNA molecules in order to increase stability. Immediately after transcription, the 3' end of the transcript may be cleaved to free a 3' hydroxyl. Then poly-A polymerase adds a chain of adenine nucleotides to the RNA. The process, called polyadenylation, adds a poly-A tail that can be between 100 and 250 residues long.

[00151] It has been discovered that unique poly-A tail lengths provide certain advantages to the polynucleotides, primary constructs or mmRNA of the present invention.

[00152] Generally, the length of a poly-A tail of the present invention is greater than 30 nucleotides in length. In another embodiment, the poly-A tail is greater than 35 nucleotides in length (e.g., at least or greater than about 35, 40, 45, 50, 55, 60, 70, 80, 90, 100, 120, 140, 160, 180, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, 1,700, 1,800, 1,900, 2,000, 2,500, and 3,000 nucleotides). In some embodiments, the polynucleotide, primary construct, or mmRNA includes from about 30 to about 3,000 nucleotides (e.g., from 30 to 50, from 30 to 100, from 30 to 250, from 30 to 500, from 30 to 750, from 30 to 1,000, from 30 to 1,500, from 30 to 2,000, from 30 to 2,500, from 50 to 100, from 50 to 250, from 50 to 500, from 50 to 750, from 50 to 1,000, from 50 to 1,500, from 50 to 2,000, from 50 to 2,500, from 50 to 3,000, from 100 to 500, from 100 to 750, from 100 to 1,000, from 100 to 1,500, from 100 to 2,000, from 100 to 2,500, from 100 to 3,000, from 500 to 750, from 500 to 1,000, from 500 to 1,500, from 500 to 2,000, from 500 to 2,500, from 500 to 3,000, from 1,000 to 1,500, from 1,000 to 2,000, from 1,000 to 2,500, from 1,000 to 3,000, from 1,500 to 2,000, from 1,500 to 2,500, from 1,500 to 3,000, from 2,000 to 3,000, from 2,000 to 2,500, and from 2,500 to 3,000).

[00153] In one embodiment, the poly-A tail is designed relative to the length of the overall polynucleotides, primary constructs or mmRNA. This design may be based on the length of the coding region, the length of a particular feature or region (such as the first or

flanking regions), or based on the length of the ultimate product expressed from the polynucleotides, primary constructs or mmRNA.

[00154] In this context the poly-A tail may be 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100% greater in length than the polynucleotides, primary constructs or mmRNA or feature thereof. The poly-A tail may also be designed as a fraction of polynucleotides, primary constructs or mmRNA to which it belongs. In this context, the poly-A tail may be 10, 20, 30, 40, 50, 60, 70, 80, or 90% or more of the total length of the construct or the total length of the construct minus the poly-A tail. Further, engineered binding sites and conjugation of polynucleotides, primary constructs or mmRNA for Poly-A binding protein may enhance expression.

[00155] Additionally, multiple distinct polynucleotides, primary constructs or mmRNA may be linked together to the PABP (Poly-A binding protein) through the 3'-end using modified nucleotides at the 3'-terminus of the poly-A tail. Transfection experiments can be conducted in relevant cell lines and protein production can be assayed by ELISA at 12hr, 24hr, 48hr, 72 hr and day 7 post-transfection.

[00156] In one embodiment, the polynucleotide primary constructs of the present invention are designed to include a polyA-G Quartet. The G-quartet is a cyclic hydrogen bonded array of four guanine nucleotides that can be formed by G-rich sequences in both DNA and RNA. In this embodiment, the G-quartet is incorporated at the end of the poly-A tail. The resultant mmRNA construct is assayed for stability, protein production and other parameters including half-life at various time points. It has been discovered that the polyA-G quartet results in protein production equivalent to at least 75% of that seen using a poly-A tail of 120 nucleotides alone.

II. Synthetic small guide RNAs (sgRNA)

[00157] The invention also includes synthetic small guide RNAs or sgRNAs. A synthetic sgRNA targets a gene of interest, e.g., a gene where modulation of transcription is desired. A synthetic sgRNA includes a sequence, typically 20-25 nucleotides long, that is complementary to one strand of the 5'UTR of the gene of interest upstream of the transcription start site. The synthetic sgRNA also includes a guide scaffold sequence. A typical guide scaffold sequence is as follows:

```
GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCC
GTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGGTGC
(SEQ ID NO:100)
```

[00158] A description of sgRNA design can be found at, e.g., Mali et al. Science. 2013. 339:823-826.

[00159] Examples of sgRNA sequences can be found in the Tables below.

Genes of interest

[00160] The sgRNA target a gene of interest, directing the CRISPR related protein encoded by the synthetic polynucleotide to interact with the gene of interest. The gene of interest is selected depending on the application. Examples genes of interest include VEGF, TPO, and/or genes apoptosis or senescence genes.

III. Design and synthesis of synthetic polynucleotides and sgRNAs

[00161] Synthetic polynucleotides and sgRNAs of the invention may be prepared according to any available technique including, but not limited to chemical synthesis; enzymatic synthesis, which is generally termed *in vitro* transcription (IVT); or enzymatic or chemical cleavage of a longer precursor, etc. Methods of synthesizing RNAs are known in the art (see, e.g., Gait, M.J. (ed.) *Oligonucleotide synthesis: a practical approach*, Oxford [Oxfordshire], Washington, DC: IRL Press, 1984; and Herdewijn, P. (ed.) *Oligonucleotide synthesis: methods and applications*, Methods in Molecular Biology, v. 288 (Clifton, N.J.) Totowa, N.J.: Humana Press, 2005; both of which are incorporated herein by reference).

[00162] The process of design and synthesis of the primary constructs, e.g. synthetic polynucleotides encoding a CRISPR related protein or synthetic sgRNA, of the invention generally includes the steps of gene construction, synthetic mRNA production (either with or without modifications) and purification. In the enzymatic synthesis method, a polynucleotide sequence encoding the CRISPR related protein or sgRNA is first selected for incorporation into a vector which will be amplified to produce a cDNA template. Optionally, the CRISPR related protein polynucleotide sequence and/or any flanking sequences may be codon optimized. The cDNA template is then used to produce mRNA or sgRNA through *in vitro* transcription (IVT). After production, the mRNA or sgRNA may undergo purification and clean-up processes. The steps of which are provided in more detail below.

Gene Construction of the Synthetic Polynucleotide encoding a CRISPR related protein

[00163] The step of gene construction may include, but is not limited to gene synthesis, vector amplification, plasmid purification, plasmid linearization and clean-up, and cDNA template synthesis and clean-up.

[00164] Once a polypeptide of interest, e.g., a CRISPR related protein, e.g., dCAS9 or a dCAS9-effector domain fusion protein, is selected for production, a primary construct is designed. Within the primary construct, a first region of linked nucleosides encoding the polypeptide of interest may be constructed using an open reading frame (ORF) of a selected nucleic acid (DNA or RNA) transcript. The ORF may comprise the wild type ORF, an isoform, variant or a fragment thereof. As used herein, an “open reading frame” or “ORF” is meant to refer to a nucleic acid sequence (DNA or RNA) which is capable of encoding a polypeptide of interest. ORFs often begin with the start codon, ATG and end with a nonsense or termination codon or signal.

[00165] Further, the nucleotide sequence of the first region may be codon optimized. Codon optimization methods are known in the art and may be useful in efforts to achieve one or more of several goals. These goals include to match codon frequencies in target and host organisms to ensure proper folding, bias GC content to increase mRNA stability or reduce secondary structures, minimize tandem repeat codons or base runs that may impair gene construction or expression, customize transcriptional and translational control regions, insert or remove protein trafficking sequences, remove/add post translation modification sites in encoded protein (e.g. glycosylation sites), add, remove or shuffle protein domains, insert or delete restriction sites, modify ribosome binding sites and mRNA degradation sites, to adjust translational rates to allow the various domains of the protein to fold properly, or to reduce or eliminate problem secondary structures within the mRNA. Codon optimization tools, algorithms and services are known in the art, non-limiting examples include services from GeneArt (Life Technologies) and/or DNA2.0 (Menlo Park CA). In one embodiment, the ORF sequence is optimized using optimization algorithms. Codon options for each amino acid are given in Table 1.

Table 1. Codon Options

Amino Acid	Single Letter Code	Codon Options
Isoleucine	I	ATT, ATC, ATA
Leucine	L	CTT, CTC, CTA, CTG, TTA, TTG
Valine	V	GTT, GTC, GTA, GTG
Phenylalanine	F	TTT, TTC

Methionine	M	ATG
Cysteine	C	TGT, TGC
Alanine	A	GCT, GCC, GCA, GCG
Glycine	G	GGT, GGC, GGA, GGG
Proline	P	CCT, CCC, CCA, CCG
Threonine	T	ACT, ACC, ACA, ACG
Serine	S	TCT, TCC, TCA, TCG, AGT, AGC
Tyrosine	Y	TAT, TAC
Tryptophan	W	TGG
Glutamine	Q	CAA, CAG
Asparagine	N	AAT, AAC
Histidine	H	CAT, CAC
Glutamic acid	E	GAA, GAG
Aspartic acid	D	GAT, GAC
Lysine	K	AAA, AAG
Arginine	R	CGT, CGC, CGA, CGG, AGA, AGG
Selenocysteine	Sec	UGA in mRNA in presence of Selenocystein insertion element (SECIS)
Stop codons	Stop	TAA, TAG, TGA

[00166] After a sequence has been codon optimized it may be further evaluated for regions containing restriction sites. At least one nucleotide within the restriction site regions may be replaced with another nucleotide in order to remove the restriction site from the sequence but the replacement of nucleotides does alter the amino acid sequence which is encoded by the codon optimized nucleotide sequence.

[00167] In some embodiments and as described in more detail herein, a 5' UTR and/or a 3' UTR may be provided as flanking regions. Multiple 5' or 3' UTRs may be included in the flanking regions and may be the same or of different sequences. Any portion of the flanking regions, including none, may be codon optimized and any may independently contain one or more different structural or chemical modifications, before and/or after codon optimization. Combinations of features may be included in the first and second flanking regions and may be contained within other features. For example, the ORF may be flanked by a 5' UTR which may contain a strong Kozak translational initiation signal and/or a 3' UTR which may include an oligo(dT) sequence for templated addition of a poly-A tail.

[00168] After optimization (if desired), the primary construct components are reconstituted and transformed into a vector such as, but not limited to, plasmids, viruses, cosmids, and artificial chromosomes. For example, the optimized construct may be reconstituted and transformed into chemically competent *E. coli*, yeast, neurospora, maize, drosophila, etc. where high copy plasmid-like or chromosome structures occur by methods described herein.

Vector Amplification

[00169] The vector containing the primary construct, encoding either the synthetic polynucleotide encoding a CRISPR related protein or the sgRNA, is then amplified and the plasmid isolated and purified using methods known in the art such as, but not limited to, a maxi prep using the Invitrogen PURELINK™ HiPure Maxiprep Kit (Carlsbad, CA).

Plasmid Linearization

[00170] The plasmid may then be linearized using methods known in the art such as, but not limited to, the use of restriction enzymes and buffers. The linearization reaction may be purified using methods including, for example Invitrogen's PURELINK™ PCR Micro Kit (Carlsbad, CA), and HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC) and Invitrogen's standard PURELINK™ PCR Kit (Carlsbad, CA). The purification method may be modified depending on the size of the linearization reaction which was conducted. The linearized plasmid is then used to generate cDNA for *in vitro* transcription (IVT) reactions.

cDNA Template Synthesis

[00171] A cDNA template may be synthesized by having a linearized plasmid undergo polymerase chain reaction (PCR). Table 4 of U.S. Patent Application No 13/791,922 filed March 9, 2013 provides a listing of primers and probes that may be usefully in the PCR reactions of the present invention. It should be understood that the listing is not exhaustive and that primer-probe design for any amplification is within the skill of those in the art. Probes may also contain chemically modified bases to increase base-pairing fidelity to the target molecule and base-pairing strength.

[00172] In one embodiment, the cDNA may be submitted for sequencing analysis before undergoing transcription.

mRNA Production

[00173] The process of synthetic polynucleotide or sgRNA production may include, but is not limited to, *in vitro* transcription, cDNA template removal and RNA clean-up, and RNA capping and/or tailing reactions. Alternatively the synthetic polynucleotide or sgRNA can be chemically synthesized.

In Vitro Transcription

[00174] The cDNA produced in the previous step may be transcribed using an *in vitro* transcription (IVT) system. The system typically comprises a transcription buffer,

nucleotide triphosphates (NTPs), an RNase inhibitor and a polymerase. The NTPs may be manufactured in house, may be selected from a supplier, or may be synthesized as described herein. The NTPs may be selected from, but are not limited to, those described herein including natural and unnatural (modified) NTPs. The polymerase may be selected from, but is not limited to, T7 RNA polymerase, T3 RNA polymerase and mutant polymerases such as, but not limited to, polymerases able to incorporate modified nucleic acids. Inorganic pyrophosphatase can be included in the transcription system.

RNA Polymerases

[00175] Any number of RNA polymerases or variants may be used in the design of the primary constructs of the present invention.

[00176] RNA polymerases may be modified by inserting or deleting amino acids of the RNA polymerase sequence. As a non-limiting example, the RNA polymerase may be modified to exhibit an increased ability to incorporate a 2'-modified nucleotide triphosphate compared to an unmodified RNA polymerase (see International Publication WO2008078180 and U.S. Patent 8,101,385; herein incorporated by reference in their entireties).

[00177] Variants may be obtained by evolving an RNA polymerase, optimizing the RNA polymerase amino acid and/or nucleic acid sequence and/or by using other methods known in the art. As a non-limiting example, T7 RNA polymerase variants may be evolved using the continuous directed evolution system set out by Esvelt *et al.* (Nature (2011) 472(7344):499-503; herein incorporated by reference in its entirety) where clones of T7 RNA polymerase may encode at least one mutation such as, but not limited to, lysine at position 93 substituted for threonine (K93T), I4M, A7T, E63V, V64D, A65E, D66Y, T76N, C125R, S128R, A136T, N165S, G175R, H176L, Y178H, F182L, L196F, G198V, D208Y, E222K, S228A, Q239R, T243N, G259D, M267I, G280C, H300R, D351A, A354S, E356D, L360P, A383V, Y385C, D388Y, S397R, M401T, N410S, K450R, P451T, G452V, E484A, H523L, H524N, G542V, E565K, K577E, K577M, N601S, S684Y, L699I, K713E, N748D, Q754R, E775K, A827V, D851N or L864F. As another non-limiting example, T7 RNA polymerase variants may encode at least mutation as described in U.S. Pub. Nos. 20100120024 and 20070117112; herein incorporated by reference in their entireties. Variants of RNA polymerase may also include, but are not limited to, substitutional variants, conservative amino acid substitution, insertional variants, deletional variants and/or covalent derivatives.

[00178] In one embodiment, the primary construct may be designed to be recognized by the wild type or variant RNA polymerases. In doing so, the primary construct may be modified to contain sites or regions of sequence changes from the wild type or parent primary construct.

[00179] In one embodiment, the primary construct may be designed to include at least one substitution and/or insertion upstream of an RNA polymerase binding or recognition site, downstream of the RNA polymerase binding or recognition site, upstream of the TATA box sequence, downstream of the TATA box sequence of the primary construct but upstream of the coding region of the primary construct, within the 5'UTR, before the 5'UTR and/or after the 5'UTR.

[00180] In one embodiment, the 5'UTR of the primary construct may be replaced by the insertion of at least one region and/or string of nucleotides of the same base. The region and/or string of nucleotides may include, but is not limited to, at least 3, at least 4, at least 5, at least 6, at least 7 or at least 8 nucleotides and the nucleotides may be natural and/or unnatural. As a non-limiting example, the group of nucleotides may include 5-8 adenine, cytosine, thymine, a string of any of the other nucleotides disclosed herein and/or combinations thereof.

[00181] In one embodiment, the 5'UTR of the primary construct may be replaced by the insertion of at least two regions and/or strings of nucleotides of two different bases such as, but not limited to, adenine, cytosine, thymine, any of the other nucleotides disclosed herein and/or combinations thereof. For example, the 5'UTR may be replaced by inserting 5-8 adenine bases followed by the insertion of 5-8 cytosine bases. In another example, the 5'UTR may be replaced by inserting 5-8 cytosine bases followed by the insertion of 5-8 adenine bases.

[00182] In one embodiment, the primary construct may include at least one substitution and/or insertion downstream of the transcription start site which may be recognized by an RNA polymerase. As a non-limiting example, at least one substitution and/or insertion may occur downstream the transcription start site by substituting at least one nucleic acid in the region just downstream of the transcription start site (such as, but not limited to, +1 to +6). Changes to region of nucleotides just downstream of the transcription start site may affect initiation rates, increase apparent nucleotide triphosphate (NTP) reaction constant values, and increase the dissociation of short transcripts from the transcription complex curing initial transcription (Briebe et al, *Biochemistry* (2002) 41: 5144-5149; herein incorporated by reference in its entirety). The modification, substitution and/or

insertion of at least one nucleic acid may cause a silent mutation of the nucleic acid sequence or may cause a mutation in the amino acid sequence.

[00183] In one embodiment, the primary construct may include the substitution of at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12 or at least 13 guanine bases downstream of the transcription start site.

[00184] In one embodiment, the primary construct may include the substitution of at least 1, at least 2, at least 3, at least 4, at least 5 or at least 6 guanine bases in the region just downstream of the transcription start site. As a non-limiting example, if the nucleotides in the region are GGGAGA (SEQ ID NO: 127) the guanine bases may be substituted by at least 1, at least 2, at least 3 or at least 4 adenine nucleotides. In another non-limiting example, if the nucleotides in the region are GGGAGA (SEQ ID NO: 127) the guanine bases may be substituted by at least 1, at least 2, at least 3 or at least 4 cytosine bases. In another non-limiting example, if the nucleotides in the region are GGGAGA (SEQ ID NO: 127) the guanine bases may be substituted by at least 1, at least 2, at least 3 or at least 4 thymine, and/or any of the nucleotides described herein.

[00185] In one embodiment, the primary construct may include at least one substitution and/or insertion upstream of the start codon. For the purpose of clarity, one of skill in the art would appreciate that the start codon is the first codon of the protein coding region whereas the transcription start site is the site where transcription begins. The primary construct may include, but is not limited to, at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7 or at least 8 substitutions and/or insertions of nucleotide bases. The nucleotide bases may be inserted or substituted at 1, at least 1, at least 2, at least 3, at least 4 or at least 5 locations upstream of the start codon. The nucleotides inserted and/or substituted may be the same base (e.g., all A or all C or all T or all G), two different bases (e.g., A and C, A and T, or C and T), three different bases (e.g., A, C and T or A, C and T) or at least four different bases. As a non-limiting example, the guanine base upstream of the coding region in the primary construct may be substituted with adenine, cytosine, thymine, or any of the nucleotides described herein. In another non-limiting example the substitution of guanine bases in the primary construct may be designed so as to leave one guanine base in the region downstream of the transcription start site and before the start codon (see Esvelt *et al.* Nature (2011) 472(7344):499-503; herein incorporated by reference in its entirety). As a non-limiting example, at least 5 nucleotides may be

inserted at 1 location downstream of the transcription start site but upstream of the start codon and the at least 5 nucleotides may be the same base type.

cDNA Template Removal and Clean-Up

[00186] The cDNA template may be removed using methods known in the art such as, but not limited to, treatment with Deoxyribonuclease I (DNase I). RNA clean-up may also include a purification method such as, but not limited to, AGENCOURT® CLEANSEQ® system from Beckman Coulter (Danvers, MA), HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC).

Capping and/or Tailing Reactions

[00187] The primary construct or mmRNA may also undergo capping and/or tailing reactions. A capping reaction may be performed by methods known in the art to add a 5' cap to the 5' end of the primary construct. Methods for capping include, but are not limited to, using a Vaccinia Capping enzyme (New England Biolabs, Ipswich, MA).

[00188] A poly-A tailing reaction may be performed by methods known in the art, such as, but not limited to, 2' O-methyltransferase and by methods as described herein. If the primary construct generated from cDNA does not include a poly-T, it may be beneficial to perform the poly-A-tailing reaction before the primary construct is cleaned.

[00189] Samples undergoing capping reactions may have varying amounts of 5' capped structures ranging from 0 to 100%. In some embodiments, the sample comprises 0, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 95, 99, or 100% 5' capped RNA.

Synthetic polynucleotide and sgRNA Purification

[00190] Purification of the synthetic polynucleotide or sgRNA may include, but is not limited to, RNA clean-up, quality assurance and quality control. RNA clean-up may be performed by methods known in the arts such as, but not limited to, AGENCOURT® beads (Beckman Coulter Genomics, Danvers, MA), poly-T beads, LNATM oligo-T capture probes (EXIQON® Inc, Vedbaek, Denmark), HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC) or RNase III treatment (a non-limiting example of treating mRNA with RNase III is described by Meis et al. in International Publication No. WO2013102203, herein incorporated by reference in its entirety). The term "purified" when used in relation to a

polynucleotide such as a “purified RNA“ refers to one that is separated from at least one contaminant. As used herein, a “contaminant” is any substance which makes another unfit, impure or inferior. Thus, a purified polynucleotide (e.g., DNA and RNA) is present in a form or setting different from that in which it is found in nature, or a form or setting different from that which existed prior to subjecting it to a treatment or purification method.

[00191] A quality assurance and/or quality control check may be conducted using methods such as, but not limited to, gel electrophoresis, UV absorbance, or analytical HPLC.

[00192] In another embodiment, the RNA may be sequenced by methods including, but not limited to reverse-transcriptase-PCR.

[00193] In one embodiment, the mRNA or mmRNA may be quantified using methods such as, but not limited to, ultraviolet visible spectroscopy (UV/Vis). A non-limiting example of a UV/Vis spectrometer is a NANODROP® spectrometer (ThermoFisher, Waltham, MA). The quantified mRNA or mmRNA may be analyzed in order to determine if the mRNA or mmRNA may be of proper size, check that no degradation of the mRNA or mmRNA has occurred. Degradation of the mRNA and/or mmRNA may be checked by methods such as, but not limited to, agarose gel electrophoresis, HPLC based purification methods such as, but not limited to, strong anion exchange HPLC, weak anion exchange HPLC, reverse phase HPLC (RP-HPLC), and hydrophobic interaction HPLC (HIC-HPLC), liquid chromatography-mass spectrometry (LCMS), capillary electrophoresis (CE) and capillary gel electrophoresis (CGE).

IV. Modified synthetic polynucleotides and sgRNAs

[00194] The synthetic polynucleotides and sgRNAs of the invention are typically modified. Herein, in a polynucleotide (such as synthetic polynucleotide or sgRNA), the terms “modification” or, as appropriate, “modified” refer to modification with respect to A, G, U or C ribonucleotides. Generally, herein, these terms are not intended to refer to the ribonucleotide modifications in naturally occurring 5'-terminal mRNA cap moieties. Examples of modifications can be found in US patent application no. 13/644,072, filed 03-Oct-2012 and published as US20130115272, .e.g, the contents of which are incorporated by reference for all purposes.

[00195] The modifications may be various distinct modifications. In some embodiments, the coding region, the flanking regions and/or the terminal regions may

contain one, two, or more (optionally different) nucleoside or nucleotide modifications. In some embodiments, a modified polynucleotide, primary construct, or mmRNA introduced to a cell may exhibit reduced degradation in the cell, as compared to an unmodified polynucleotide, primary construct, or mmRNA.

[00196] The polynucleotides, primary constructs, and mmRNA can include any useful modification, such as to the sugar, the nucleobase, or the internucleoside linkage (*e.g.* to a linking phosphate / to a phosphodiester linkage / to the phosphodiester backbone). One or more atoms of a pyrimidine nucleobase may be replaced or substituted with optionally substituted amino, optionally substituted thiol, optionally substituted alkyl (*e.g.*, methyl or ethyl), or halo (*e.g.*, chloro or fluoro). In certain embodiments, modifications (*e.g.*, one or more modifications) are present in each of the sugar and the internucleoside linkage. Modifications according to the present invention may be modifications of ribonucleic acids (RNAs) to deoxyribonucleic acids (DNAs), threose nucleic acids (TNAs), glycol nucleic acids (GNAs), peptide nucleic acids (PNAs), locked nucleic acids (LNAs) or hybrids thereof). Additional modifications are described herein.

[00197] As described herein, the polynucleotides, primary constructs, and mmRNA of the invention do not substantially induce an innate immune response of a cell into which the mRNA is introduced. Features of an induced innate immune response include 1) increased expression of pro-inflammatory cytokines, 2) activation of intracellular PRRs (RIG-I, MDA5, etc, and/or 3) termination or reduction in protein translation.

[00198] In certain embodiments, it may be desirable to intracellularly degrade a modified nucleic acid molecule introduced into the cell. For example, degradation of a modified nucleic acid molecule may be preferable if precise timing of protein production is desired. Thus, in some embodiments, the invention provides a modified nucleic acid molecule containing a degradation domain, which is capable of being acted on in a directed manner within a cell.

[00199] The polynucleotides, primary constructs, and mmRNA can optionally include other agents (*e.g.*, RNAi-inducing agents, RNAi agents, siRNAs, shRNAs, miRNAs, antisense RNAs, ribozymes, catalytic DNA, tRNA, RNAs that induce triple helix formation, aptamers, vectors, etc.).

[00200] In some embodiments, the polynucleotide, primary construct, or mmRNA (*e.g.*, the first region, first flanking region, or second flanking region) includes *n* number of linked nucleosides having any base, sugar, backbone, building block or other structure or formula, including but not limited to those of Formulas I through IX or any substructures

thereof as described in International Publication No WO2013052523 filed October 3, 2012, entitled Modified Nucleosides, Nucleotides, and Nucleic Acids, and Uses Thereof, the contents of which are incorporated herein by reference in their entirety. Such structures include modifications to the sugar, nucleobase, internucleoside linkage, or combinations thereof.

[00201] Combinations of chemical modifications include those taught in including but not limited to those described in International Publication No WO2013052523 filed October 3, 2012, entitled Modified Nucleosides, Nucleotides, and Nucleic Acids, and Uses Thereof, the contents of which are incorporated herein by reference in their entirety.

[00202] The synthesis of polynucleotides, primary constructs or mmRNA of the present invention may be according to the methods described in International Publication No WO2013052523 filed October 3, 2012, entitled Modified Nucleosides, Nucleotides, and Nucleic Acids, and Uses Thereof, the contents of which are incorporated herein by reference in their entirety.

[00203] In some embodiments, the nucleobase selected from the group consisting of cytosine, guanine, adenine, and uracil.

[00204] In some embodiments, the modified nucleobase is a modified uracil. Exemplary nucleobases and nucleosides having a modified uracil include pseudouridine (ψ), pyridin-4-one ribonucleoside, 5-aza-uridine, 6-aza-uridine, 2-thio-5-aza-uridine, 2-thio-uridine (s^2U), 4-thio-uridine (s^4U), 4-thio-pseudouridine, 2-thio-pseudouridine, 5-hydroxy-uridine (ho^5U), 5-aminoallyl-uridine, 5-halo-uridine (e.g., 5-iodo-uridine or 5-bromo-uridine), 3-methyl-uridine (m^3U), 5-carboxymethyl-uridine (cm^5U), 1-carboxymethyl-pseudouridine, 5-carboxyhydroxymethyl-uridine (chm^5U), 5-carboxyhydroxymethyl-uridine methyl ester ($mchm^5U$), 5-methoxycarbonylmethyl-uridine (mcm^5U), 5-methoxycarbonylmethyl-2-thio-uridine (mcm^5s^2U), 5-aminomethyl-2-thio-uridine (nm^5s^2U), 5-methylaminomethyl-uridine (mnm^5U), 5-methylaminomethyl-2-thio-uridine (mnm^5s^2U), 5-methylaminomethyl-2-seleno-uridine (mnm^5se^2U), 5-carbamoylmethyl-uridine (nem^5U), 5-carboxymethylaminomethyl-uridine ($cmnm^5U$), 5-carboxymethylaminomethyl-2-thio-uridine ($cmnm^5s^2U$), 5-propynyl-uridine, 1-propynyl-pseudouridine, 5-taurinomethyl-uridine (τm^5U), 1-taurinomethyl-pseudouridine, 5-taurinomethyl-2-thio-uridine (τm^5s^2U), 1-taurinomethyl-4-thio-pseudouridine, 5-methyl-uridine (m^5U , i.e., having the nucleobase deoxythymine), 1-methyl-pseudouridine ($m^1\psi$), 5-methyl-2-thio-uridine (m^5s^2U), 1-methyl-4-thio-pseudouridine ($m^1s^4\psi$), 4-thio-1-methyl-pseudouridine, 3-methyl-pseudouridine ($m^3\psi$), 2-thio-1-methyl-pseudouridine, 1-

methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydrouridine (D), dihydropseudouridine, 5,6-dihyrouridine, 5-methyl-dihyrouridine (m^5D), 2-thio-dihyrouridine, 2-thio-dihydropseudouridine, 2-methoxy-uridine, 2-methoxy-4-thio-uridine, 4-methoxy-pseudouridine, 4-methoxy-2-thio-pseudouridine, N1-methyl-pseudouridine, 3-(3-amino-3-carboxypropyl)uridine (acp^3U), 1-methyl-3-(3-amino-3-carboxypropyl)pseudouridine ($acp^3\psi$), 5-(isopentenylaminomethyl)uridine (inm^5U), 5-(isopentenylaminomethyl)-2-thio-uridine (inm^5s^2U), α -thio-uridine, 2'-O-methyl-uridine (Um), 5,2'-O-dimethyl-uridine (m^5Um), 2'-O-methyl-pseudouridine (ψm), 2-thio-2'-O-methyl-uridine (s^2Um), 5-methoxycarbonylmethyl-2'-O-methyl-uridine (mcm^5Um), 5-carbamoylmethyl-2'-O-methyl-uridine (ncm^5Um), 5-carboxymethylaminomethyl-2'-O-methyl-uridine ($cmnm^5Um$), 3,2'-O-dimethyl-uridine (m^3Um), 5-(isopentenylaminomethyl)-2'-O-methyl-uridine (inm^5Um), 1-thio-uridine, deoxythymidine, 2'-F-ara-uridine, 2'-F-uridine, 2'-OH-ara-uridine, 5-(2-carbomethoxyvinyl) uridine, and 5-[3-(1-E-propenylamino)uridine.

[00205] In some embodiments, the modified nucleobase is a modified cytosine.

Exemplary nucleobases and nucleosides having a modified cytosine include 5-aza-cytidine, 6-aza-cytidine, pseudoisocytidine, 3-methyl-cytidine (m^3C), N4-acetyl-cytidine (ac^4C), 5-formyl-cytidine (f^5C), N4-methyl-cytidine (m^4C), 5-methyl-cytidine (m^5C), 5-halo-cytidine (e.g., 5-iodo-cytidine), 5-hydroxymethyl-cytidine (hm^5C), 1-methyl-pseudoisocytidine, pyrrolo-cytidine, pyrrolo-pseudoisocytidine, 2-thio-cytidine (s^2C), 2-thio-5-methyl-cytidine, 4-thio-pseudoisocytidine, 4-thio-1-methyl-pseudoisocytidine, 4-thio-1-methyl-1-deaza-pseudoisocytidine, 1-methyl-1-deaza-pseudoisocytidine, zebularine, 5-aza-zebularine, 5-methyl-zebularine, 5-aza-2-thio-zebularine, 2-thio-zebularine, 2-methoxy-cytidine, 2-methoxy-5-methyl-cytidine, 4-methoxy-pseudoisocytidine, 4-methoxy-1-methyl-pseudoisocytidine, lysidine (k_2C), α -thio-cytidine, 2'-O-methyl-cytidine (Cm), 5,2'-O-dimethyl-cytidine (m^5Cm), N4-acetyl-2'-O-methyl-cytidine (ac^4Cm), N4,2'-O-dimethyl-cytidine (m^4Cm), 5-formyl-2'-O-methyl-cytidine (f^5Cm), N4,N4,2'-O-trimethyl-cytidine (m^4_2Cm), 1-thio-cytidine, 2'-F-ara-cytidine, 2'-F-cytidine, and 2'-OH-ara-cytidine.

[00206] In exemplary embodiments, the modified nucleobase is a modified uracil selected from pseudouridine (ψ) and 1-methylpseudouridine. In some embodiments, the modified nucleobase is a modified uracil in combination with a modified cytosine, e.g., 5-methylcytosine.

[00207] In some embodiments, a polynucleotide of the invention is fully modified with pseudouridine (ψ), optionally in combination with 5-methylcytosine. In some embodiments, a polynucleotide of the invention is fully modified with 1-methylpseudouridine ($m^1\psi$), optionally in combination with 5-methylcytosine.

[00208] In some embodiments, the mRNA molecules are codon optimized.

[00209] In some embodiments, the modified nucleobase is a modified adenine.

Exemplary nucleobases and nucleosides having a modified adenine include 2-amino-purine, 2, 6-diaminopurine, 2-amino-6-halo-purine (e.g., 2-amino-6-chloro-purine), 6-halo-purine (e.g., 6-chloro-purine), 2-amino-6-methyl-purine, 8-azido-adenosine, 7-deaza-adenine, 7-deaza-8-aza-adenine, 7-deaza-2-amino-purine, 7-deaza-8-aza-2-amino-purine, 7-deaza-2,6-diaminopurine, 7-deaza-8-aza-2,6-diaminopurine, 1-methyl-adenosine (m^1A), 2-methyl-adenine (m^2A), N6-methyl-adenosine (m^6A), 2-methylthio-N6-methyl-adenosine (ms^2m^6A), N6-isopentenyl-adenosine (i^6A), 2-methylthio-N6-isopentenyl-adenosine (ms^2i^6A), N6-(cis-hydroxyisopentenyl)adenosine (io^6A), 2-methylthio-N6-(cis-hydroxyisopentenyl)adenosine (ms^2io^6A), N6-glycinylocarbamoyl-adenosine (g^6A), N6-threonylocarbamoyl-adenosine (t^6A), N6-methyl-N6-threonylocarbamoyl-adenosine (m^6t^6A), 2-methylthio-N6-threonylocarbamoyl-adenosine (ms^2g^6A), N6,N6-dimethyl-adenosine (m^6_2A), N6-hydroxynorvalylcarbonyl-adenosine (hn^6A), 2-methylthio-N6-hydroxynorvalylcarbonyl-adenosine (ms^2hn^6A), N6-acetyl-adenosine (ac^6A), 7-methyl-adenine, 2-methylthio-adenine, 2-methoxy-adenine, α -thio-adenosine, 2'-O-methyl-adenosine (Am), N6,2'-O-dimethyl-adenosine (m^6Am), N6,N6,2'-O-trimethyl-adenosine (m^6_2Am), 1,2'-O-dimethyl-adenosine (m^1Am), 2'-O-ribosyladenosine (phosphate) ($Ar(p)$), 2-amino-N6-methyl-purine, 1-thio-adenosine, 8-azido-adenosine, 2'-F-ara-adenosine, 2'-F-adenosine, 2'-OH-ara-adenosine, and N6-(19-amino-pentaoxonadecyl)-adenosine.

[00210] In some embodiments, the modified nucleobase is a modified guanine.

Exemplary nucleobases and nucleosides having a modified guanine include inosine (I), 1-methyl-inosine (m^1I), wyosine (imG), methylwyosine (mimG), 4-demethyl-wyosine (imG-14), isowyosine (imG2), wybutosine (yW), peroxywybutosine (o_2yW), hydroxywybutosine (OHyW), undermodified hydroxywybutosine (OHyW*), 7-deaza-guanosine, queuosine (Q), epoxyqueuosine (oQ), galactosyl-queuosine (galQ), mannosyl-queuosine (manQ), 7-cyano-7-deaza-guanosine ($preQ_0$), 7-aminomethyl-7-deaza-guanosine ($preQ_1$), archaeosine (G^+), 7-deaza-8-aza-guanosine, 6-thio-guanosine, 6-thio-7-deaza-guanosine, 6-thio-7-deaza-8-aza-guanosine, 7-methyl-guanosine (m^7G), 6-thio-7-

methyl-guanosine, 7-methyl-inosine, 6-methoxy-guanosine, 1-methyl-guanosine (m^1G), N2-methyl-guanosine (m^2G), N2,N2-dimethyl-guanosine (m^2_2G), N2,7-dimethyl-guanosine ($m^{2,7}G$), N2, N2,7-dimethyl-guanosine ($m^{2,2,7}G$), 8-oxo-guanosine, 7-methyl-8-oxo-guanosine, 1-methyl-6-thio-guanosine, N2-methyl-6-thio-guanosine, N2,N2-dimethyl-6-thio-guanosine, α -thio-guanosine, 2'-O-methyl-guanosine (Gm), N2-methyl-2'-O-methyl-guanosine (m^2Gm), N2,N2-dimethyl-2'-O-methyl-guanosine (m^2_2Gm), 1-methyl-2'-O-methyl-guanosine (m^1Gm), N2,7-dimethyl-2'-O-methyl-guanosine ($m^{2,7}Gm$), 2'-O-methyl-inosine (Im), 1,2'-O-dimethyl-inosine (m^1Im), and 2'-O-riboseylguanosine (phosphate) (Gr(p)).

[00211] The nucleobase of the nucleotide can be independently selected from a purine, a pyrimidine, a purine or pyrimidine analog. For example, the nucleobase can each be independently selected from adenine, cytosine, guanine, uracil, or hypoxanthine. In another embodiment, the nucleobase can also include, for example, naturally-occurring and synthetic derivatives of a base, including pyrazolo[3,4-d]pyrimidines, 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo (*e.g.*, 8-bromo), 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, deazaguanine, 7-deazaguanine, 3-deazaguanine, deazaadenine, 7-deazaadenine, 3-deazaadenine, pyrazolo[3,4-d]pyrimidine, imidazo[1,5-a]1,3,5 triazinones, 9-deazapurines, imidazo[4,5-d]pyrazines, thiazolo[4,5-d]pyrimidines, pyrazin-2-ones, 1,2,4-triazine, pyridazine; and 1,3,5 triazine. When the nucleotides are depicted using the shorthand A, G, C, T or U, each letter refers to the representative base and/or derivatives thereof, *e.g.*, A includes adenine or adenine analogs, *e.g.*, 7-deaza adenine).

[00212] Modified nucleosides and nucleotides (*e.g.*, building block molecules) can be prepared according to the synthetic methods described in Ogata et al., *J. Org. Chem.* 74:2585-2588 (2009); Purmal et al., *Nucl. Acids Res.* 22(1): 72-78, (1994); Fukuhara et al., *Biochemistry*, 1(4): 563-568 (1962); and Xu et al., *Tetrahedron*, 48(9): 1729-1740 (1992), each of which are incorporated by reference in their entirety.

[00213] The polypeptides, primary constructs, and mmRNA of the invention may or may not be uniformly modified along the entire length of the molecule. For example, one or more or all types of nucleotide (*e.g.*, purine or pyrimidine, or any one or more or all of A, G, U, C) may or may not be uniformly modified in a polynucleotide of the invention, or in a given predetermined sequence region thereof (*e.g.* one or more of the sequence regions represented in Figure 1). In some embodiments, all nucleotides X in a polynucleotide of the invention (or in a given sequence region thereof) are modified, wherein X may any one of nucleotides A, G, U, C, or any one of the combinations A+G, A+U, A+C, G+U, G+C, U+C, A+G+U, A+G+C, G+U+C or A+G+C.

[00214] Different sugar modifications, nucleotide modifications, and/or internucleoside linkages (*e.g.*, backbone structures) may exist at various positions in the polynucleotide, primary construct, or mmRNA. One of ordinary skill in the art will appreciate that the nucleotide analogs or other modification(s) may be located at any position(s) of a polynucleotide, primary construct, or mmRNA such that the function of the polynucleotide, primary construct, or mmRNA is not substantially decreased. A modification may also be a 5' or 3' terminal modification. The polynucleotide, primary construct, or mmRNA may contain from about 1% to about 100% modified nucleotides (either in relation to overall nucleotide content, or in relation to one or more types of nucleotide, *i.e.* any one or more of A, G, U or C) or any intervening percentage (*e.g.*, from 1% to 20%, from 1% to 25%, from 1% to 50%, from 1% to 60%, from 1% to 70%, from 1% to 80%, from 1% to 90%, from 1% to 95%, from 10% to 20%, from 10% to 25%, from 10% to 50%, from 10% to 60%, from 10% to 70%, from 10% to 80%, from 10% to 90%, from 10% to 95%, from 10% to 100%, from 20% to 25%, from 20% to 50%, from 20% to 60%, from 20% to 70%, from 20% to 80%, from 20% to 90%, from 20% to 95%, from 20% to 100%, from 50% to 60%, from 50% to 70%, from 50% to 80%, from 50% to 90%, from 50% to 95%, from 50% to 100%, from 70% to 80%, from 70% to 90%, from 70% to 95%, from 70% to 100%, from 80% to 90%, from 80% to 95%, from 80% to 100%, from 90% to 95%, from 90% to 100%, and from 95% to 100%).

[00215] In some embodiments, the polynucleotide, primary construct, or mmRNA includes a modified pyrimidine (*e.g.*, a modified uracil/uridine/U or modified cytosine/cytidine/C). In some embodiments, the uracil or uridine (generally: U) in the polynucleotide, primary construct, or mmRNA molecule may be replaced with from about 1% to about 100% of a modified uracil or modified uridine (*e.g.*, from 1% to 20%,

from 1% to 25%, from 1% to 50%, from 1% to 60%, from 1% to 70%, from 1% to 80%, from 1% to 90%, from 1% to 95%, from 10% to 20%, from 10% to 25%, from 10% to 50%, from 10% to 60%, from 10% to 70%, from 10% to 80%, from 10% to 90%, from 10% to 95%, from 10% to 100%, from 20% to 25%, from 20% to 50%, from 20% to 60%, from 20% to 70%, from 20% to 80%, from 20% to 90%, from 20% to 95%, from 20% to 100%, from 50% to 60%, from 50% to 70%, from 50% to 80%, from 50% to 90%, from 50% to 95%, from 50% to 100%, from 70% to 80%, from 70% to 90%, from 70% to 95%, from 70% to 100%, from 80% to 90%, from 80% to 95%, from 80% to 100%, from 90% to 95%, from 90% to 100%, and from 95% to 100% of a modified uracil or modified uridine). The modified uracil or uridine can be replaced by a compound having a single unique structure or by a plurality of compounds having different structures (e.g., 2, 3, 4 or more unique structures, as described herein).

[00216] In some embodiments, the cytosine or cytidine (generally: C) in the polynucleotide, primary construct, or mmRNA molecule may be replaced with from about 1% to about 100% of a modified cytosine or modified cytidine (e.g., from 1% to 20%, from 1% to 25%, from 1% to 50%, from 1% to 60%, from 1% to 70%, from 1% to 80%, from 1% to 90%, from 1% to 95%, from 10% to 20%, from 10% to 25%, from 10% to 50%, from 10% to 60%, from 10% to 70%, from 10% to 80%, from 10% to 90%, from 10% to 95%, from 10% to 100%, from 20% to 25%, from 20% to 50%, from 20% to 60%, from 20% to 70%, from 20% to 80%, from 20% to 90%, from 20% to 95%, from 20% to 100%, from 50% to 60%, from 50% to 70%, from 50% to 80%, from 50% to 90%, from 50% to 95%, from 50% to 100%, from 70% to 80%, from 70% to 90%, from 70% to 95%, from 70% to 100%, from 80% to 90%, from 80% to 95%, from 80% to 100%, from 90% to 95%, from 90% to 100%, and from 95% to 100% of a modified cytosine or modified cytidine). The modified cytosine or cytidine can be replaced by a compound having a single unique structure or by a plurality of compounds having different structures (e.g., 2, 3, 4 or more unique structures, as described herein).

[00217] In some embodiments, at least 25% of the cytosines are replaced by a compound of Formula (b10)-(b14) (e.g., at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%).

[00218] In some embodiments, at least 25% of the uracils are replaced by a compound of Formula (b1)-(b9) (e.g., at least about 30%, at least about 35%, at least about 40%, at

least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%).

[00219] In some embodiments, at least 25% of the cytosines are replaced by a compound of Formula (b10)-(b14), and at least 25% of the uracils are replaced by a compound of Formula (b1)-(b9) (e.g., at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%).

[00220] In one embodiment, the synthetic polynucleotide and/or sgRNA includes a terminal modification. The terminal modification may be a modification described in US Provisional Application No. US 61/729,933, filed November 26, 2012, entitled Terminally Optimized RNAs; US Provisional Application No. US 61/737,224, filed December 14, 2012, entitled Terminally Optimized RNAs; US Provisional Application No. US61/758,921, filed January 31, 2013, entitled Differential Targeting Using RNA Constructs; US Provisional Application No. US 61/781,139, filed March 14, 2013, entitled Differential Targeting Using RNA Constructs; US Provisional Application No. US 61/829,359, filed May 31, 2013, entitled Differential Targeting Using RNA Constructs; US Provisional Application No. US 61/839,903, filed June 27, 2013, entitled Differential Targeting Using RNA Constructs; US Provisional Application No. 61/842,709, filed July 3, 2013, entitled Differential Targeting Using RNA Constructs; each of which is herein incorporated by reference in its entirety.

[00221] In another embodiment, the chemical modification may be a modification described in International Publication No. WO2013052523, herein incorporated by reference in its entirety.

V. Pharmaceutical Compositions: Formulation, Administration, Delivery and Dosing

[00222] In one embodiment, the invention includes synthetic polynucleotide and/or sgRNA compositions and complexes in combination with one or more pharmaceutically acceptable excipients. Pharmaceutical compositions may optionally comprise one or more additional active substances, e.g. therapeutically and/or prophylactically active substances. General considerations in the formulation and/or manufacture of pharmaceutical agents may be found, for example, in *Remington: The Science and*

Practice of Pharmacy 21st ed., Lippincott Williams & Wilkins, 2005 (incorporated herein by reference in its entirety). synthetic polynucleotides and/or synthetic sgRNAs

[00223] In one embodiment, the synthetic polynucleotide and/or sgRNA are formulated. As a non-limiting example the synthetic polynucleotides and/or synthetic sgRNAs may be formulated by the methods described in International Publication No. WO2013090648 and/or co-pending US Provisional Application No. US 61/821,406, filed March 14, 2013, entitled Formulation and Delivery of Modified Nucleoside, Nucleotide, and Nucleic Acid Compositions, US Provisional Application No. US 61/821,406, filed May 9, 2013, entitled Formulation and Delivery of Modified Nucleoside, Nucleotide, and Nucleic Acid Compositions and US Provisional Application No. US 61/840,510, filed June 28, 2013, entitled Formulation and Delivery of Modified Nucleoside, Nucleotide, and Nucleic Acid Compositions, each of which is herein incorporated by reference in its entirety.

[00224] Although the descriptions of pharmaceutical compositions provided herein are principally directed to pharmaceutical compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to any other animal, *e.g.*, to non-human animals, *e.g.* non-human mammals. Modification of pharmaceutical compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with merely ordinary, if any, experimentation. Subjects to which administration of the pharmaceutical compositions is contemplated include, but are not limited to, humans and/or other primates; mammals, including commercially relevant mammals such as cattle, pigs, horses, sheep, cats, dogs, mice, and/or rats; and/or birds, including commercially relevant birds such as poultry, chickens, ducks, geese, and/or turkeys.

[00225] Formulations of the pharmaceutical compositions described herein may be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include the step of bringing the active ingredient into association with an excipient and/or one or more other accessory ingredients, and then, if necessary and/or desirable, dividing, shaping and/or packaging the product into a desired single- or multi-dose unit.

[00226] A pharmaceutical composition in accordance with the invention may be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of

single unit doses. As used herein, a “unit dose” is discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

[00227] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition in accordance with the invention will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of example, the composition may comprise between 0.1% and 100%, e.g., between .5 and 50%, between 1-30%, between 5-80%, at least 80% (w/w) active ingredient

Formulations

[00228] The synthetic polynucleotides and/or synthetic sgRNAs of the invention can be formulated using one or more excipients to: (1) increase stability; (2) increase cell transfection; (3) permit the sustained or delayed release (e.g., from a depot formulation of the polynucleotide, primary construct, or mmRNA); (4) alter the biodistribution (e.g., target the polynucleotide, primary construct, or mmRNA to specific tissues or cell types); (5) increase the translation of encoded protein *in vivo*; and/or (6) alter the release profile of encoded protein *in vivo*. In addition to traditional excipients such as any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, excipients of the present invention can include, without limitation, lipidoids, liposomes, lipid nanoparticles, polymers, lipoplexes, core-shell nanoparticles, peptides, proteins, cells transfected with polynucleotide, primary construct, or mmRNA (e.g., for transplantation into a subject), hyaluronidase, nanoparticle mimics and combinations thereof. Accordingly, the formulations of the invention can include one or more excipients, each in an amount that together increases the stability of the polynucleotide, primary construct, or mmRNA, increases cell transfection by the polynucleotide, primary construct, or mmRNA, increases the expression of polynucleotide, primary construct, or mmRNA encoded protein, and/or alters the release profile of polynucleotide, primary construct, or mmRNA encoded proteins. Further, the

primary construct and mmRNA of the present invention may be formulated using self-assembled nucleic acid nanoparticles.

[00229] Formulations of the pharmaceutical compositions described herein may be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include the step of associating the active ingredient with an excipient and/or one or more other accessory ingredients.

[00230] A pharmaceutical composition in accordance with the present disclosure may be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. As used herein, a “unit dose” refers to a discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient. The amount of the active ingredient may generally be equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage including, but not limited to, one-half or one-third of such a dosage.

[00231] Relative amounts of the active ingredient, the pharmaceutically acceptable excipient, and/or any additional ingredients in a pharmaceutical composition in accordance with the present disclosure may vary, depending upon the identity, size, and/or condition of the subject being treated and further depending upon the route by which the composition is to be administered. For example, the composition may comprise between 0.1% and 99% (w/w) of the active ingredient.

[00232] In some embodiments, the formulations described herein may contain at least one synthetic polynucleotide. As a non-limiting example, the formulations may contain 1, 2, 3, 4 or 5 mmRNA. In one embodiment the formulation may contain modified mRNA encoding proteins selected from categories such as, but not limited to, human proteins, veterinary proteins, bacterial proteins, biological proteins, antibodies, immunogenic proteins, therapeutic peptides and proteins, secreted proteins, plasma membrane proteins, cytoplasmic and cytoskeletal proteins, intracellular membrane bound proteins, nuclear proteins, proteins associated with human disease and/or proteins associated with non-human diseases. In one embodiment, the formulation contains at least three modified mRNA encoding proteins. In one embodiment, the formulation contains at least five modified mRNA encoding proteins.

[00233] Pharmaceutical formulations may additionally comprise a pharmaceutically acceptable excipient, which, as used herein, includes, but is not limited to, any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents,

preservatives, and the like, as suited to the particular dosage form desired. Various excipients for formulating pharmaceutical compositions and techniques for preparing the composition are known in the art (see Remington: The Science and Practice of Pharmacy, 21st Edition, A. R. Gennaro, Lippincott, Williams & Wilkins, Baltimore, MD, 2006; incorporated herein by reference). The use of a conventional excipient medium may be contemplated within the scope of the present disclosure, except insofar as any conventional excipient medium may be incompatible with a substance or its derivatives, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition.

[00234] In some embodiments, the particle size of the lipid nanoparticle may be increased and/or decreased. The change in particle size may be able to help counter biological reaction such as, but not limited to, inflammation or may increase the biological effect of the modified mRNA delivered to mammals.

[00235] Pharmaceutically acceptable excipients used in the manufacture of pharmaceutical compositions include, but are not limited to, inert diluents, surface active agents and/or emulsifiers, preservatives, buffering agents, lubricating agents, and/or oils. Such excipients may optionally be included in the pharmaceutical formulations of the invention.

Lipidoids

[00236] The synthesis of lipidoids has been extensively described and formulations containing these compounds are particularly suited for delivery of polynucleotides, primary constructs or mmRNA (see Mahon et al., *Bioconjug Chem.* 2010 21:1448-1454; Schroeder et al., *J Intern Med.* 2010 267:9-21; Akinc et al., *Nat Biotechnol.* 2008 26:561-569; Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869; Siegwart et al., *Proc Natl Acad Sci U S A.* 2011 108:12996-3001; all of which are incorporated herein in their entirety).

[00237] While these lipidoids have been used to effectively deliver double stranded small interfering RNA molecules in rodents and non-human primates (see Akinc et al., *Nat Biotechnol.* 2008 26:561-569; Frank-Kamenetsky et al., *Proc Natl Acad Sci U S A.* 2008 105:11915-11920; Akinc et al., *Mol Ther.* 2009 17:872-879; Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869; Leuschner et al., *Nat Biotechnol.* 2011 29:1005-1010; all of which is incorporated herein in their entirety), the present disclosure describes their formulation and use in delivering single stranded polynucleotides, primary

constructs, or mmRNA. Complexes, micelles, liposomes or particles can be prepared containing these lipidoids and therefore, can result in an effective delivery of the polynucleotide, primary construct, or mmRNA, as judged by the production of an encoded protein, following the injection of a lipidoid formulation via localized and/or systemic routes of administration. Lipidoid complexes of polynucleotides, primary constructs, or mmRNA can be administered by various means including, but not limited to, intravenous, intramuscular, or subcutaneous routes.

[00238] *In vivo* delivery of nucleic acids may be affected by many parameters, including, but not limited to, the formulation composition, nature of particle PEGylation, degree of loading, oligonucleotide to lipid ratio, and biophysical parameters such as particle size (Akinc et al., *Mol Ther.* 2009 17:872-879; herein incorporated by reference in its entirety). As an example, small changes in the anchor chain length of poly(ethylene glycol) (PEG) lipids may result in significant effects on *in vivo* efficacy. Formulations with the different lipidoids, including, but not limited to penta[3-(1-laurylamino propionyl)]-triethylenetetramine hydrochloride (TETA-5LAP; aka 98N12-5, see Murugaiah et al., *Analytical Biochemistry*, 401:61 (2010)), C12-200 (including derivatives and variants), and MD1, can be tested for *in vivo* activity.

[00239] The lipidoid referred to herein as “98N12-5” is disclosed by Akinc et al., *Mol Ther.* 2009 17:872-879 and is incorporated by reference in its entirety. (See Figure 2)

[00240] The lipidoid referred to herein as “C12-200” is disclosed by Love et al., *Proc Natl Acad Sci U S A.* 2010 107:1864-1869 (see Figure 2) and Liu and Huang, *Molecular Therapy.* 2010 669-670 (see Figure 2); both of which are herein incorporated by reference in their entirety. The lipidoid formulations can include particles comprising either 3 or 4 or more components in addition to polynucleotide, primary construct, or mmRNA. As an example, formulations with certain lipidoids, include, but are not limited to, 98N12-5 and may contain 42% lipidoid, 48% cholesterol and 10% PEG (C14 alkyl chain length). As another example, formulations with certain lipidoids, include, but are not limited to, C12-200 and may contain 50% lipidoid, 10% distearylphosphatidyl choline, 38.5% cholesterol, and 1.5% PEG-DMG.

[00241] In one embodiment, a polynucleotide, primary construct, or mmRNA formulated with a lipidoid for systemic intravenous administration can target the liver. For example, a final optimized intravenous formulation using polynucleotide, primary construct, or mmRNA, and comprising a lipid molar composition of 42% 98N12-5, 48% cholesterol, and 10% PEG-lipid with a final weight ratio of about 7.5 to 1 total lipid to

polynucleotide, primary construct, or mmRNA, and a C14 alkyl chain length on the PEG lipid, with a mean particle size of roughly 50–60 nm, can result in the distribution of the formulation to be greater than 90% to the liver (see, Akinc et al., Mol Ther. 2009 17:872-879; herein incorporated in its entirety). In another example, an intravenous formulation using a C12-200 (see US provisional application 61/175,770 and published international application WO2010129709, each of which is herein incorporated by reference in their entirety) lipidoid may have a molar ratio of 50/10/38.5/1.5 of C12-200/disteroylphosphatidyl choline/cholesterol/PEG-DMG, with a weight ratio of 7 to 1 total lipid to polynucleotide, primary construct, or mmRNA, and a mean particle size of 80 nm may be effective to deliver polynucleotide, primary construct, or mmRNA to hepatocytes (see, Love et al., Proc Natl Acad Sci U S A. 2010 107:1864-1869 herein incorporated by reference). In another embodiment, an MD1 lipidoid-containing formulation may be used to effectively deliver polynucleotide, primary construct, or mmRNA to hepatocytes *in vivo*. The characteristics of optimized lipidoid formulations for intramuscular or subcutaneous routes may vary significantly depending on the target cell type and the ability of formulations to diffuse through the extracellular matrix into the blood stream. While a particle size of less than 150 nm may be desired for effective hepatocyte delivery due to the size of the endothelial fenestrae (see, Akinc et al., Mol Ther. 2009 17:872-879 herein incorporated by reference), use of a lipidoid-formulated polynucleotide, primary construct, or mmRNA to deliver the formulation to other cells types including, but not limited to, endothelial cells, myeloid cells, and muscle cells may not be similarly size-limited. Use of lipidoid formulations to deliver siRNA *in vivo* to other non-hepatocyte cells such as myeloid cells and endothelium has been reported (see Akinc et al., Nat Biotechnol. 2008 26:561-569; Leuschner et al., Nat Biotechnol. 2011 29:1005-1010; Cho et al. Adv. Funct. Mater. 2009 19:3112-3118; 8th International Judah Folkman Conference, Cambridge, MA October 8-9, 2010 herein incorporated by reference in its entirety). Effective delivery to myeloid cells, such as monocytes, lipidoid formulations may have a similar component molar ratio. Different ratios of lipidoids and other components including, but not limited to, disteroylphosphatidyl choline, cholesterol and PEG-DMG, may be used to optimize the formulation of the polynucleotide, primary construct, or mmRNA for delivery to different cell types including, but not limited to, hepatocytes, myeloid cells, muscle cells, etc. For example, the component molar ratio may include, but is not limited to, 50% C12-200, 10% disteroylphosphatidyl choline, 38.5% cholesterol, and %1.5 PEG-DMG (see Leuschner et al., Nat Biotechnol 2011

29:1005-1010; herein incorporated by reference in its entirety). The use of lipidoid formulations for the localized delivery of nucleic acids to cells (such as, but not limited to, adipose cells and muscle cells) via either subcutaneous or intramuscular delivery, may not require all of the formulation components desired for systemic delivery, and as such may comprise only the lipidoid and the polynucleotide, primary construct, or mmRNA.

[00242] Combinations of different lipidoids may be used to improve the efficacy of polynucleotide, primary construct, or mmRNA directed protein production as the lipidoids may be able to increase cell transfection by the polynucleotide, primary construct, or mmRNA; and/or increase the translation of encoded protein (see Whitehead et al., Mol. Ther. 2011, 19:1688-1694, herein incorporated by reference in its entirety).

Liposomes, Lipoplexes, and Lipid Nanoparticles

[00243] The polynucleotide, primary construct, and mmRNA of the invention can be formulated using one or more liposomes, lipoplexes, or lipid nanoparticles. In one embodiment, pharmaceutical compositions of polynucleotide, primary construct, or mmRNA include liposomes. Liposomes are artificially-prepared vesicles which may primarily be composed of a lipid bilayer and may be used as a delivery vehicle for the administration of nutrients and pharmaceutical formulations. Liposomes can be of different sizes such as, but not limited to, a multilamellar vesicle (MLV) which may be hundreds of nanometers in diameter and may contain a series of concentric bilayers separated by narrow aqueous compartments, a small unicellular vesicle (SUV) which may be smaller than 50 nm in diameter, and a large unilamellar vesicle (LUV) which may be between 50 and 500 nm in diameter. Liposome design may include, but is not limited to, opsonins or ligands in order to improve the attachment of liposomes to unhealthy tissue or to activate events such as, but not limited to, endocytosis. Liposomes may contain a low or a high pH in order to improve the delivery of the pharmaceutical formulations.

[00244] The formation of liposomes may depend on the physicochemical characteristics such as, but not limited to, the pharmaceutical formulation entrapped and the liposomal ingredients, the nature of the medium in which the lipid vesicles are dispersed, the effective concentration of the entrapped substance and its potential toxicity, any additional processes involved during the application and/or delivery of the vesicles, the optimization size, polydispersity and the shelf-life of the vesicles for the intended application, and the batch-to-batch reproducibility and possibility of large-scale production of safe and efficient liposomal products.

[00245] In one embodiment, pharmaceutical compositions described herein may include, without limitation, liposomes such as those formed from 1,2-dioleoyloxy-*N,N*-dimethylaminopropane (DODMA) liposomes, DiLa2 liposomes from Marina Biotech (Bothell, WA), 1,2-dilinoleoyloxy-3-dimethylaminopropane (DLin-DMA), 2,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), and MC3 (US20100324120; herein incorporated by reference in its entirety) and liposomes which may deliver small molecule drugs such as, but not limited to, DOXIL® from Janssen Biotech, Inc. (Horsham, PA).

[00246] In one embodiment, pharmaceutical compositions described herein may include, without limitation, liposomes such as those formed from the synthesis of stabilized plasmid-lipid particles (SPLP) or stabilized nucleic acid lipid particle (SNALP) that have been previously described and shown to be suitable for oligonucleotide delivery in vitro and in vivo (see Wheeler et al. *Gene Therapy*. 1999 6:271-281; Zhang et al. *Gene Therapy*. 1999 6:1438-1447; Jeffs et al. *Pharm Res*. 2005 22:362-372; Morrissey et al., *Nat Biotechnol*. 2005 2:1002-1007; Zimmermann et al., *Nature*. 2006 441:111-114; Heyes et al. *J Contr Rel*. 2005 107:276-287; Semple et al. *Nature Biotech*. 2010 28:172-176; Judge et al. *J Clin Invest*. 2009 119:661-673; deFougerolles *Hum Gene Ther*. 2008 19:125-132; all of which are incorporated herein in their entireties). The original manufacture method by Wheeler et al. was a detergent dialysis method, which was later improved by Jeffs et al. and is referred to as the spontaneous vesicle formation method. The liposome formulations are composed of 3 to 4 lipid components in addition to the polynucleotide, primary construct, or mmRNA. As an example a liposome can contain, but is not limited to, 55% cholesterol, 20% distearylphosphatidyl choline (DSPC), 10% PEG-S-DSG, and 15% 1,2-dioleoyloxy-*N,N*-dimethylaminopropane (DODMA), as described by Jeffs et al. As another example, certain liposome formulations may contain, but are not limited to, 48% cholesterol, 20% DSPC, 2% PEG-c-DMA, and 30% cationic lipid, where the cationic lipid can be 1,2-distearloxy-*N,N*-dimethylaminopropane (DSDMA), DODMA, DLin-DMA, or 1,2-dilinolenyloxy-3-dimethylaminopropane (DLenDMA), as described by Heyes et al.

[00247] In one embodiment, the polynucleotides, primary constructs and/or mmRNA may be formulated in a lipid vesicle which may have crosslinks between functionalized lipid bilayers.

[00248] In one embodiment, the polynucleotides, primary constructs and/or mmRNA may be formulated in a lipid-polycation complex. The formation of the lipid-polycation

complex may be accomplished by methods known in the art and/or as described in U.S. Pub. No. 20120178702, herein incorporated by reference in its entirety. As a non-limiting example, the polycation may include a cationic peptide or a polypeptide such as, but not limited to, polylysine, polyornithine and/or polyarginine. In another embodiment, the polynucleotides, primary constructs and/or mmRNA may be formulated in a lipid-polycation complex which may further include a neutral lipid such as, but not limited to, cholesterol or dioleoyl phosphatidylethanolamine (DOPE).

[00249] The liposome formulation may be influenced by, but not limited to, the selection of the cationic lipid component, the degree of cationic lipid saturation, the nature of the PEGylation, ratio of all components and biophysical parameters such as size. In one example by Semple et al. (Semple et al. *Nature Biotech.* 2010 28:172-176), the liposome formulation was composed of 57.1 % cationic lipid, 7.1% dipalmitoylphosphatidylcholine, 34.3 % cholesterol, and 1.4% PEG-c-DMA. As another example, changing the composition of the cationic lipid could more effectively deliver siRNA to various antigen presenting cells (Basha et al. *Mol Ther.* 2011 19:2186-2200; herein incorporated by reference in its entirety).

[00250] In some embodiments, the ratio of PEG in the LNP formulations may be increased or decreased and/or the carbon chain length of the PEG lipid may be modified from C14 to C18 to alter the pharmacokinetics and/or biodistribution of the LNP formulations. As a non-limiting example, LNP formulations may contain 1-5% of the lipid molar ratio of PEG-c-DOMG as compared to the cationic lipid, DSPC and cholesterol. In another embodiment the PEG-c-DOMG may be replaced with a PEG lipid such as, but not limited to, PEG-DSG (1,2-Distearoyl-sn-glycerol, methoxypolyethylene glycol) or PEG-DPG (1,2-Dipalmitoyl-sn-glycerol, methoxypolyethylene glycol). The cationic lipid may be selected from any lipid known in the art such as, but not limited to, DLin-MC3-DMA, DLin-DMA, C12-200 and DLin-KC2-DMA.

[00251] In one embodiment, the cationic lipid may be selected from, but not limited to, a cationic lipid described in International Publication Nos. WO2012040184, WO2011153120, WO2011149733, WO2011090965, WO2011043913, WO2011022460, WO2012061259, WO2012054365, WO2012044638, WO2010080724, WO201021865 and WO2008103276, US Patent Nos. 7,893,302 and 7,404,969 and US Patent Publication No. US20100036115; each of which is herein incorporated by reference in their entirety. In another embodiment, the cationic lipid may be selected from, but not limited to, formula A described in International Publication Nos. WO2012040184, WO2011153120,

WO2011149733, WO2011090965, WO2011043913, WO2011022460, WO2012061259, WO2012054365 and WO2012044638; each of which is herein incorporated by reference in their entirety. In yet another embodiment, the cationic lipid may be selected from, but not limited to, formula CLI-CLXXIX of International Publication No. WO2008103276, formula CLI-CLXXIX of US Patent No. 7,893,302, formula CLI-CLXXXII of US Patent No. 7,404,969 and formula I-VI of US Patent Publication No. US20100036115; each of which is herein incorporated by reference in their entirety. As a non-limiting example, the cationic lipid may be selected from (20Z,23Z)-N,N-dimethylnonacosan-20,23-dien-10-amine, (17Z,20Z)-N,N-dimethylhexacosan-17,20-dien-9-amine, (1Z,19Z)-N,N-dimethylpentacosan-16,19-dien-8-amine, (13Z,16Z)-N,N-dimethyldocosan-13,16-dien-5-amine, (12Z,15Z)-N,N-dimethylhenicosan-12,15-dien-4-amine, (14Z,17Z)-N,N-dimethyltricosan-14,17-dien-6-amine, (15Z,18Z)-N,N-dimethyltetracosan-15,18-dien-7-amine, (18Z,21Z)-N,N-dimethylheptacosan-18,21-dien-10-amine, (15Z,18Z)-N,N-dimethyltetracosan-15,18-dien-5-amine, (14Z,17Z)-N,N-dimethyltricosan-14,17-dien-4-amine, (19Z,22Z)-N,N-dimethyloctacosan-19,22-dien-9-amine, (18Z,21Z)-N,N-dimethylheptacosan-18,21-dien-8-amine, (17Z,20Z)-N,N-dimethylhexacosan-17,20-dien-7-amine, (16Z,19Z)-N,N-dimethylpentacosan-16,19-dien-6-amine, (22Z,25Z)-N,N-dimethylhentriacontan-22,25-dien-10-amine, (21Z,24Z)-N,N-dimethyltriacontan-21,24-dien-9-amine, (18Z)-N,N-dimethylheptacosan-18-en-10-amine, (17Z)-N,N-dimethylhexacosan-17-en-9-amine, (19Z,22Z)-N,N-dimethyloctacosan-19,22-dien-7-amine, N,N-dimethylheptacosan-10-amine, (20Z,23Z)-N-ethyl-N-methylnonacosan-20,23-dien-10-amine, 1-[(1Z,14Z)-1-nonylicosan-1,14-dien-1-yl]pyrrolidine, (20Z)-N,N-dimethylheptacosan-20-en-10-amine, (15Z)-N,N-dimethylheptacosan-15-en-10-amine, (14Z)-N,N-dimethylnonacosan-14-en-10-amine, (17Z)-N,N-dimethylnonacosan-17-en-10-amine, (24Z)-N,N-dimethyltriacontan-24-en-10-amine, (20Z)-N,N-dimethylnonacosan-20-en-10-amine, (22Z)-N,N-dimethylhentriacontan-22-en-10-amine, (16Z)-N,N-dimethylpentacosan-16-en-8-amine, (12Z,15Z)-N,N-dimethyl-2-nonylhenicosan-12,15-dien-1-amine, (13Z,16Z)-N,N-dimethyl-3-nonyldocosan-13,16-dien-1-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]heptadecan-8-amine, 1-[(1S,2R)-2-hexylcyclopropyl]-N,N-dimethylnonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]nonadecan-10-amine, N,N-dimethyl-21-[(1S,2R)-2-octylcyclopropyl]henicosan-10-amine, N,N-dimethyl-1-[(1S,2S)-2-[(1R,2R)-2-pentylcyclopropyl]methyl]cyclopropyl]nonadecan-10-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]hexadecan-8-amine, N,N-dimethyl-1-[(1R,2S)-2-

undecylcyclopropyl]tetradecan-5-amine, N,N-dimethyl-3- {7-[(1S,2R)-2-octylcyclopropyl]heptyl} dodecan-1-amine, 1-[(1R,2S)-2-heptylcyclopropyl]-N,N-dimethyloctadecan-9-amine, 1-[(1S,2R)-2-decylcyclopropyl]-N,N-dimethylpentadecan-6-amine, N,N-dimethyl-1-[(1S,2R)-2-octylcyclopropyl]pentadecan-8-amine, R-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, S-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-(octyloxy)propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}pyrrolidine, (2S)-N,N-dimethyl-1-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-3-[(5Z)-oct-5-en-1-yloxy]propan-2-amine, 1-{2-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]-1-[(octyloxy)methyl]ethyl}azetidine, (2S)-1-(hexyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2S)-1-(heptyloxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(nonyloxy)-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-[(9Z)-octadec-9-en-1-yloxy]-3-(octyloxy)propan-2-amine (Compound 9); (2S)-N,N-dimethyl-1-[(6Z,9Z,12Z)-octadeca-6,9,12-trien-1-yloxy]-3-(octyloxy)propan-2-amine, (2S)-1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(pentyloxy)propan-2-amine, (2S)-1-(hexyloxy)-3-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethylpropan-2-amine, 1-[(11Z,14Z)-icosa-11,14-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2S)-1-[(13Z,16Z)-docosa-13,16-dien-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, (2S)-1-[(13Z)-docos-13-en-1-yloxy]-3-(hexyloxy)-N,N-dimethylpropan-2-amine, 1-[(13Z)-docos-13-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, 1-[(9Z)-hexadec-9-en-1-yloxy]-N,N-dimethyl-3-(octyloxy)propan-2-amine, (2R)-N,N-dimethyl-H(1-metoyloctyl)oxy]-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, (2R)-1-[(3,7-dimethyloctyl)oxy]-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-2-amine, N,N-dimethyl-1-(octyloxy)-3-({8-[(1S,2S)-2-[(1R,2R)-2-pentylcyclopropyl]methyl}cyclopropyl]octyl}oxy)propan-2-amine, N,N-dimethyl-1-{{8-(2-octylcyclopropyl)octyl}oxy}-3-(octyloxy)propan-2-amine and (11E,20Z,23Z)-N,N-dimethylnonacos-11,20,2-trien-10-amine or a pharmaceutically acceptable salt or stereoisomer thereof.

[00252] In one embodiment, the cationic lipid may be synthesized by methods known in the art and/or as described in International Publication Nos. WO2012040184, WO2011153120, WO2011149733, WO2011090965, WO2011043913, WO2011022460,

WO2012061259, WO2012054365, WO2012044638, WO2010080724 and WO201021865; each of which is herein incorporated by reference in their entirety.

[00253] In one embodiment, the LNP formulations of the polynucleotides, primary constructs and/or mmRNA may contain PEG-c-DOMG 3% lipid molar ratio. In another embodiment, the LNP formulations of the polynucleotides, primary constructs and/or mmRNA may contain PEG-c-DOMG 1.5% lipid molar ratio.

[00254] In one embodiment, the pharmaceutical compositions of the polynucleotides, primary constructs and/or mmRNA may include at least one of the PEGylated lipids described in International Publication No. 2012099755, herein incorporated by reference.

[00255] In one embodiment, the LNP formulation may contain PEG-DMG 2000 (1,2-dimyristoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000). In one embodiment, the LNP formulation may contain PEG-DMG 2000, a cationic lipid known in the art and at least one other component. In another embodiment, the LNP formulation may contain PEG-DMG 2000, a cationic lipid known in the art, DSPC and cholesterol. As a non-limiting example, the LNP formulation may contain PEG-DMG 2000, DLin-DMA, DSPC and cholesterol. As another non-limiting example the LNP formulation may contain PEG-DMG 2000, DLin-DMA, DSPC and cholesterol in a molar ratio of 2:40:10:48 (see Geall et al., Nonviral delivery of self-amplifying RNA vaccines, PNAS 2012; PMID: 22908294).

[00256] In one embodiment, the LNP formulation may be formulated by the methods described in International Publication Nos. WO2011127255 or WO2008103276, each of which is herein incorporated by reference in their entirety. As a non-limiting example, modified RNA described herein may be encapsulated in LNP formulations as described in WO2011127255 and/or WO2008103276; each of which is herein incorporated by reference in their entirety.

[00257] In one embodiment, LNP formulations described herein may comprise a polycationic composition. As a non-limiting example, the polycationic composition may be selected from formula 1-60 of US Patent Publication No. US20050222064; herein incorporated by reference in its entirety. In another embodiment, the LNP formulations comprising a polycationic composition may be used for the delivery of the modified RNA described herein *in vivo* and/or *in vitro*.


[00258] In one embodiment, the LNP formulations described herein may additionally comprise a permeability enhancer molecule. Non-limiting permeability enhancer

molecules are described in US Patent Publication No. US20050222064; herein incorporated by reference in its entirety.


[00259] In one embodiment, the pharmaceutical compositions may be formulated in liposomes such as, but not limited to, DiLa2 liposomes (Marina Biotech, Bothell, WA), SMARTICLES® (Marina Biotech, Bothell, WA), neutral DOPC (1,2-dioleoyl-sn-glycero-3-phosphocholine) based liposomes (e.g., siRNA delivery for ovarian cancer (Landen et al. Cancer Biology & Therapy 2006 5(12)1708-1713)) and hyaluronan-coated liposomes (Quiet Therapeutics, Israel).

[00260] Lipid nanoparticle formulations may be improved by replacing the cationic lipid with a biodegradable cationic lipid which is known as a rapidly eliminated lipid nanoparticle (reLNP). Ionizable cationic lipids, such as, but not limited to, DLinDMA, DLin-KC2-DMA, and DLin-MC3-DMA, have been shown to accumulate in plasma and tissues over time and may be a potential source of toxicity. The rapid metabolism of the rapidly eliminated lipids can improve the tolerability and therapeutic index of the lipid nanoparticles by an order of magnitude from a 1 mg/kg dose to a 10 mg/kg dose in rat. Inclusion of an enzymatically degraded ester linkage can improve the degradation and metabolism profile of the cationic component, while still maintaining the activity of the reLNP formulation. The ester linkage can be internally located within the lipid chain or it may be terminally located at the terminal end of the lipid chain. The internal ester linkage may replace any carbon in the lipid chain.

[00261] In one embodiment, the internal ester linkage may be located on either side of the saturated carbon. Non-limiting examples of reLNPs include,



And



[00262] In one embodiment, an immune response may be elicited by delivering a lipid nanoparticle which may include a nanospecies, a polymer and an immunogen. (U.S. Publication No. 20120189700 and International Publication No. WO2012099805; each of which is herein incorporated by reference in their entirety). The polymer may encapsulate the nanospecies or partially encapsulate the nanospecies.

[00263] Lipid nanoparticles may be engineered to alter the surface properties of particles so the lipid nanoparticles may penetrate the mucosal barrier. Mucus is located on mucosal tissue such as, but not limited to, oral (e.g., the buccal and esophageal membranes and tonsil tissue), ophthalmic, gastrointestinal (e.g., stomach, small intestine, large intestine, colon, rectum), nasal, respiratory (e.g., nasal, pharyngeal, tracheal and bronchial membranes), genital (e.g., vaginal, cervical and urethral membranes). Nanoparticles larger than 10-200 nm which are preferred for higher drug encapsulation efficiency and the ability to provide the sustained delivery of a wide array of drugs have been thought to be too large to rapidly diffuse through mucosal barriers. Mucus is continuously secreted, shed, discarded or digested and recycled so most of the trapped particles may be removed from the mucosal tissue within seconds or within a few hours. Large polymeric nanoparticles (200nm -500nm in diameter) which have been coated densely with a low molecular weight polyethylene glycol (PEG) diffused through mucus only 4 to 6-fold lower than the same particles diffusing in water (Lai et al. PNAS 2007 104(5):1482-487; Lai et al. Adv Drug Deliv Rev. 2009 61(2): 158-171; each of which is herein incorporated by reference in their entirety). The transport of nanoparticles may be determined using rates of permeation and/or fluorescent microscopy techniques including, but not limited to, fluorescence recovery after photobleaching (FRAP) and high resolution multiple particle tracking (MPT).

[00264] The lipid nanoparticle engineered to penetrate mucus may comprise a polymeric material (i.e. a polymeric core) and/or a polymer-vitamin conjugate and/or a tri-block co-polymer. The polymeric material may include, but is not limited to, polyamines, polyethers, polyamides, polyesters, polycarbamates, polyureas, polycarbonates, poly(styrenes), polyimides, polysulfones, polyurethanes, polyacetylenes, polyethylenes, polyethyleneimines, polyisocyanates, polyacrylates, polymethacrylates, polyacrylonitriles, and polyarylates. The polymeric material may be biodegradable and/or biocompatible. Non-limiting examples of specific polymers include poly(caprolactone) (PCL), ethylene vinyl acetate polymer (EVA), poly(lactic acid) (PLA), poly(L-lactic acid) (PLLA), poly(glycolic acid) (PGA), poly(lactic acid-co-glycolic acid) (PLGA), poly(L-

lactic acid-co-glycolic acid) (PLLGA), poly(D,L-lactide) (PDLA), poly(L-lactide) (PLLA), poly(D,L-lactide-co-caprolactone), poly(D,L-lactide-co-caprolactone-co-glycolide), poly(D,L-lactide-co-PEO-co-D,L-lactide), poly(D,L-lactide-co-PPO-co-D,L-lactide), polyalkyl cyanoacrylate, polyurethane, poly-L-lysine (PLL), hydroxypropyl methacrylate (HPMA), polyethyleneglycol, poly-L-glutamic acid, poly(hydroxy acids), polyanhydrides, polyorthoesters, poly(ester amides), polyamides, poly(ester ethers), polycarbonates, polyalkylenes such as polyethylene and polypropylene, polyalkylene glycols such as poly(ethylene glycol) (PEG), polyalkylene oxides (PEO), polyalkylene terephthalates such as poly(ethylene terephthalate), polyvinyl alcohols (PVA), polyvinyl ethers, polyvinyl esters such as poly(vinyl acetate), polyvinyl halides such as poly(vinyl chloride) (PVC), polyvinylpyrrolidone, polysiloxanes, polystyrene (PS), polyurethanes, derivatized celluloses such as alkyl celluloses, hydroxyalkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, hydroxypropylcellulose, carboxymethylcellulose, polymers of acrylic acids, such as poly(methyl(meth)acrylate) (PMMA), poly(ethyl(meth)acrylate), poly(butyl(meth)acrylate), poly(isobutyl(meth)acrylate), poly(hexyl(meth)acrylate), poly(isodecyl(meth)acrylate), poly(lauryl(meth)acrylate), poly(phenyl(meth)acrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate) and copolymers and mixtures thereof, polydioxanone and its copolymers, polyhydroxyalkanoates, polypropylene fumarate, polyoxymethylene, poloxamers, poly(ortho)esters, poly(butyric acid), poly(valeric acid), poly(lactide-co-caprolactone), and trimethylene carbonate, polyvinylpyrrolidone. The lipid nanoparticle may be coated or associated with a co-polymer such as, but not limited to, a block copolymer, and (poly(ethylene glycol))-(poly(propylene oxide))-(poly(ethylene glycol)) triblock copolymer (see US Publication 20120121718 and US Publication 20100003337; each of which is herein incorporated by reference in their entirety). The co-polymer may be a polymer that is generally regarded as safe (GRAS) and the formation of the lipid nanoparticle may be in such a way that no new chemical entities are created. For example, the lipid nanoparticle may comprise poloxamers coating PLGA nanoparticles without forming new chemical entities which are still able to rapidly penetrate human mucus (Yang et al. *Angew. Chem. Int. Ed.* 2011 50:2597-2600; herein incorporated by reference in its entirety).

[00265] The vitamin of the polymer-vitamin conjugate may be vitamin E. The vitamin portion of the conjugate may be substituted with other suitable components such as, but not limited to, vitamin A, vitamin E, other vitamins, cholesterol, a hydrophobic moiety, or

a hydrophobic component of other surfactants (e.g., sterol chains, fatty acids, hydrocarbon chains and alkylene oxide chains).

[00266] The lipid nanoparticle engineered to penetrate mucus may include surface altering agents such as, but not limited to, mmRNA, anionic protein (e.g., bovine serum albumin), surfactants (e.g., cationic surfactants such as for example dimethyldioctadecylammonium bromide), sugars or sugar derivatives (e.g., cyclodextrin), nucleic acids, polymers (e.g., heparin, polyethylene glycol and poloxamer), mucolytic agents (e.g., N-acetylcysteine, mugwort, bromelain, papain, clerodendrum, acetylcysteine, bromhexine, carbocysteine, eprazinone, mesna, ambroxol, sobrerol, domiodol, letosteine, stepronin, tiopronin, gelsolin, thymosin β 4 dornase alfa, neltexine, erdosteine) and various DNases including rhDNase. The surface altering agent may be embedded or enmeshed in the particle's surface or disposed (e.g., by coating, adsorption, covalent linkage, or other process) on the surface of the lipid nanoparticle. (see US Publication 20100215580 and US Publication 20080166414; each of which is herein incorporated by reference in their entirety).

[00267] The mucus penetrating lipid nanoparticles may comprise at least one mmRNA described herein. The mmRNA may be encapsulated in the lipid nanoparticle and/or disposed on the surface of the particle. The mmRNA may be covalently coupled to the lipid nanoparticle. Formulations of mucus penetrating lipid nanoparticles may comprise a plurality of nanoparticles. Further, the formulations may contain particles which may interact with the mucus and alter the structural and/or adhesive properties of the surrounding mucus to decrease mucoadhesion which may increase the delivery of the mucus penetrating lipid nanoparticles to the mucosal tissue.

[00268] In one embodiment, the polynucleotide, primary construct, or mmRNA is formulated as a lipoplex, such as, without limitation, the ATUPLEXTM system, the DACC system, the DBTC system and other siRNA-lipoplex technology from Silence Therapeutics (London, United Kingdom), STEMFECTTM from STEMGENT® (Cambridge, MA), and polyethylenimine (PEI) or protamine-based targeted and non-targeted delivery of nucleic acids (Aleku et al. *Cancer Res.* 2008 68:9788-9798; Strumberg et al. *Int J Clin Pharmacol Ther* 2012 50:76-78; Santel et al., *Gene Ther* 2006 13:1222-1234; Santel et al., *Gene Ther* 2006 13:1360-1370; Gutbier et al., *Pulm Pharmacol. Ther.* 2010 23:334-344; Kaufmann et al. *Microvasc Res* 2010 80:286-293; Weide et al. *J Immunother.* 2009 32:498-507; Weide et al. *J Immunother.* 2008 31:180-188; Pascolo *Expert Opin. Biol. Ther.* 4:1285-1294; Fotin-Mleczek et al., 2011 *J.*

Immunother. 34:1-15; Song et al., Nature Biotechnol. 2005, 23:709-717; Peer et al., Proc Natl Acad Sci U S A. 2007 6;104:4095-4100; deFougerolles *Hum Gene Ther.* 2008 19:125-132; all of which are incorporated herein by reference in its entirety).

[00269] In one embodiment such formulations may also be constructed or compositions altered such that they passively or actively are directed to different cell types *in vivo*, including but not limited to hepatocytes, immune cells, tumor cells, endothelial cells, antigen presenting cells, and leukocytes (Akinc et al. Mol Ther. 2010 18:1357-1364; Song et al., Nat Biotechnol. 2005 23:709-717; Judge et al., J Clin Invest. 2009 119:661-673; Kaufmann et al., Microvasc Res 2010 80:286-293; Santel et al., Gene Ther 2006 13:1222-1234; Santel et al., Gene Ther 2006 13:1360-1370; Gutbier et al., Pulm Pharmacol. Ther. 2010 23:334-344; Basha et al., Mol. Ther. 2011 19:2186-2200; Fenske and Cullis, Expert Opin Drug Deliv. 2008 5:25-44; Peer et al., Science. 2008 319:627-630; Peer and Lieberman, Gene Ther. 2011 18:1127-1133; all of which are incorporated herein by reference in its entirety). One example of passive targeting of formulations to liver cells includes the DLin-DMA, DLin-KC2-DMA and MC3-based lipid nanoparticle formulations which have been shown to bind to apolipoprotein E and promote binding and uptake of these formulations into hepatocytes *in vivo* (Akinc et al. Mol Ther. 2010 18:1357-1364; herein incorporated by reference in its entirety). Formulations can also be selectively targeted through expression of different ligands on their surface as exemplified by, but not limited by, folate, transferrin, N-acetylgalactosamine (GalNAc), and antibody targeted approaches (Kolhatkar et al., Curr Drug Discov Technol. 2011 8:197-206; Musacchio and Torchilin, *Front Biosci.* 2011 16:1388-1412; Yu et al., Mol Membr Biol. 2010 27:286-298; Patil et al., Crit Rev Ther Drug Carrier Syst. 2008 25:1-61; Benoit et al., Biomacromolecules. 2011 12:2708-2714; Zhao et al., Expert Opin Drug Deliv. 2008 5:309-319; Akinc et al., Mol Ther. 2010 18:1357-1364; Srinivasan et al., Methods Mol Biol. 2012 820:105-116; Ben-Arie et al., Methods Mol Biol. 2012 757:497-507; Peer 2010 J Control Release. 20:63-68; Peer et al., Proc Natl Acad Sci U S A. 2007 104:4095-4100; Kim et al., Methods Mol Biol. 2011 721:339-353; Subramanya et al., Mol Ther. 2010 18:2028-2037; Song et al., Nat Biotechnol. 2005 23:709-717; Peer et al., Science. 2008 319:627-630; Peer and Lieberman, Gene Ther. 2011 18:1127-1133; all of which are incorporated herein by reference in its entirety)..

[00270] In one embodiment, the polynucleotide, primary construct, or mmRNA is formulated as a solid lipid nanoparticle. A solid lipid nanoparticle (SLN) may be spherical with an average diameter between 10 to 1000 nm. SLN possess a solid lipid

core matrix that can solubilize lipophilic molecules and may be stabilized with surfactants and/or emulsifiers. In a further embodiment, the lipid nanoparticle may be a self-assembly lipid-polymer nanoparticle (see Zhang et al., ACS Nano, 2008, 2 (8), pp 1696–1702; herein incorporated by reference in its entirety).

[00271] Liposomes, lipoplexes, or lipid nanoparticles may be used to improve the efficacy of polynucleotide, primary construct, or mmRNA directed protein production as these formulations may be able to increase cell transfection by the polynucleotide, primary construct, or mmRNA; and/or increase the translation of encoded protein. One such example involves the use of lipid encapsulation to enable the effective systemic delivery of polyplex plasmid DNA (Heyes et al., Mol Ther. 2007 15:713-720; herein incorporated by reference in its entirety). The liposomes, lipoplexes, or lipid nanoparticles may also be used to increase the stability of the polynucleotide, primary construct, or mmRNA.

[00272] In one embodiment, the polynucleotides, primary constructs, and/or the mmRNA of the present invention can be formulated for controlled release and/or targeted delivery. As used herein, “controlled release” refers to a pharmaceutical composition or compound release profile that conforms to a particular pattern of release to effect a therapeutic outcome. In one embodiment, the polynucleotides, primary constructs or the mmRNA may be encapsulated into a delivery agent described herein and/or known in the art for controlled release and/or targeted delivery. As used herein, the term “encapsulate” means to enclose, surround or encase. As it relates to the formulation of the compounds of the invention, encapsulation may be substantial, complete or partial. The term “substantially encapsulated” means that at least greater than 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.9 or greater than 99.999% of the pharmaceutical composition or compound of the invention may be enclosed, surrounded or encased within the delivery agent. “Partially encapsulation” means that less than 10, 10, 20, 30, 40 50 or less of the pharmaceutical composition or compound of the invention may be enclosed, surrounded or encased within the delivery agent. Advantageously, encapsulation may be determined by measuring the escape or the activity of the pharmaceutical composition or compound of the invention using fluorescence and/or electron micrograph. For example, at least 1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the pharmaceutical composition or compound of the invention are encapsulated in the delivery agent.

[00273] In another embodiment, the polynucleotides, primary constructs, or the mmRNA may be encapsulated into a lipid nanoparticle or a rapidly eliminating lipid nanoparticle and the lipid nanoparticles or a rapidly eliminating lipid nanoparticle may then be encapsulated into a polymer, hydrogel and/or surgical sealant described herein and/or known in the art. As a non-limiting example, the polymer, hydrogel or surgical sealant may be PLGA, ethylene vinyl acetate (EVAc), poloxamer, GELSITE® (Nanotherapeutics, Inc. Alachua, FL), HYLENEX® (Halozyme Therapeutics, San Diego CA), surgical sealants such as fibrinogen polymers (Ethicon Inc. Cornelia, GA), TISSELL® (Baxter International, Inc Deerfield, IL), PEG-based sealants, and COSEAL® (Baxter International, Inc Deerfield, IL).

[00274] In one embodiment, the lipid nanoparticle may be encapsulated into any polymer or hydrogel known in the art which may form a gel when injected into a subject. As another non-limiting example, the lipid nanoparticle may be encapsulated into a polymer matrix which may be biodegradable.

[00275] In one embodiment, the polynucleotide, primary construct, or mmRNA formulation for controlled release and/or targeted delivery may also include at least one controlled release coating. Controlled release coatings include, but are not limited to, OPADRY®, polyvinylpyrrolidone/vinyl acetate copolymer, polyvinylpyrrolidone, hydroxypropyl methylcellulose, hydroxypropyl cellulose, hydroxyethyl cellulose, EUDRAGIT RL®, EUDRAGIT RS® and cellulose derivatives such as ethylcellulose aqueous dispersions (AQUACOAT® and SURELEASE®).

[00276] In one embodiment, the controlled release and/or targeted delivery formulation may comprise at least one degradable polyester which may contain polycationic side chains. Degradable polyesters include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In another embodiment, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

[00277] In one embodiment, the polynucleotides, primary constructs, and/or the mmRNA of the present invention may be encapsulated in a therapeutic nanoparticle. Therapeutic nanoparticles may be formulated by methods described herein and known in the art such as, but not limited to, International Pub Nos. WO2010005740, WO2010030763, WO2010005721, WO2010005723, WO2012054923, US Pub. Nos. US20110262491, US20100104645, US20100087337, US20100068285, US20110274759, US20100068286, and US Pat No. 8,206,747; each of which is herein

incorporated by reference in their entirety. In another embodiment, therapeutic polymer nanoparticles may be identified by the methods described in US Pub No.

US20120140790, herein incorporated by reference in its entirety.

[00278] In one embodiment, the therapeutic nanoparticle of may be formulated for sustained release. As used herein, “sustained release” refers to a pharmaceutical composition or compound that conforms to a release rate over a specific period of time. The period of time may include, but is not limited to, hours, days, weeks, months and years. As a non-limiting example, the sustained release nanoparticle may comprise a polymer and a therapeutic agent such as, but not limited to, the polynucleotides, primary constructs, and mmRNA of the present invention (see International Pub No. 2010075072 and US Pub No. US20100216804 and US20110217377, each of which is herein incorporated by reference in their entirety).

[00279] In one embodiment, the therapeutic nanoparticles may be formulated to be target specific. As a non-limiting example, the therapeutic nanoparticles may include a corticosteroid (see International Pub. No. WO2011084518). In one embodiment, the therapeutic nanoparticles may be formulated to be cancer specific. As a non-limiting example, the therapeutic nanoparticles may be formulated in nanoparticles described in International Pub No. WO2008121949, WO2010005726, WO2010005725, WO2011084521 and US Pub No. US20100069426, US20120004293 and US20100104655, each of which is herein incorporated by reference in their entirety.

[00280] In one embodiment, the nanoparticles of the present invention may comprise a polymeric matrix. As a non-limiting example, the nanoparticle may comprise two or more polymers such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester) or combinations thereof.

[00281] In one embodiment, the diblock copolymer may include PEG in combination with a polymer such as, but not limited to, polyethylenes, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates,

polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester) or combinations thereof.

[00282] In one embodiment, the therapeutic nanoparticle comprises a diblock copolymer. As a non-limiting example the therapeutic nanoparticle comprises a PLGA-PEG block copolymer (see US Pub. No. US20120004293 and US Pat No. 8,236,330, herein incorporated by reference in their entireties). In another non-limiting example, the therapeutic nanoparticle is a stealth nanoparticle comprising a diblock copolymer of PEG and PLA or PEG and PLGA (see US Pat No 8,246,968, each of which is herein incorporated by reference in its entirety).

[00283] In one embodiment, the therapeutic nanoparticle may comprise at least one acrylic polymer. Acrylic polymers include but are not limited to, acrylic acid, methacrylic acid, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, amino alkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), polycyanoacrylates and combinations thereof.

[00284] In one embodiment, the therapeutic nanoparticles may comprise at least one cationic polymer described herein and/or known in the art.

[00285] In one embodiment, the therapeutic nanoparticles may comprise at least one amine-containing polymer such as, but not limited to polylysine, polyethylene imine, poly(amidoamine) dendrimers and combinations thereof.

[00286] In one embodiment, the therapeutic nanoparticles may comprise at least one degradable polyester which may contain polycationic side chains. Degradable polyesters include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In another embodiment, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

[00287] In another embodiment, the therapeutic nanoparticle may include a conjugation of at least one targeting ligand.

[00288] In one embodiment, the therapeutic nanoparticle may be formulated in an aqueous solution which may be used to target cancer (see International Pub No. WO2011084513 and US Pub No. US20110294717, each of which is herein incorporated by reference in their entirety).

[00289] In one embodiment, the polynucleotides, primary constructs, or mmRNA may be encapsulated in, linked to and/or associated with synthetic nanocarriers. The synthetic

nanocarriers may be formulated using methods known in the art and/or described herein. As a non-limiting example, the synthetic nanocarriers may be formulated by the methods described in International Pub Nos. WO2010005740, WO2010030763 and US Pub. Nos. US20110262491, US20100104645 and US20100087337, each of which is herein incorporated by reference in their entirety. In another embodiment, the synthetic nanocarrier formulations may be lyophilized by methods described in International Pub. No. WO2011072218 and US Pat No. 8,211,473; each of which is herein incorporated by reference in their entireties.

[00290] In one embodiment, the synthetic nanocarriers may contain reactive groups to release the polynucleotides, primary constructs and/or mmRNA described herein (see International Pub. No. WO20120952552 and US Pub No. US20120171229, each of which is herein incorporated by reference in their entirety).

[00291] In one embodiment, the synthetic nanocarriers may contain an immunostimulatory agent to enhance the immune response from delivery of the synthetic nanocarrier. As a non-limiting example, the synthetic nanocarrier may comprise a Th1 immunostimulatory agent which may enhance a Th1-based response of the immune system (see International Pub No. WO2010123569 and US Pub. No. US20110223201, each of which is herein incorporated by reference in its entirety).

[00292] In one embodiment, the synthetic nanocarriers may be formulated for targeted release. In one embodiment, the synthetic nanocarrier is formulated to release the polynucleotides, primary constructs and/or mmRNA at a specified pH and/or after a desired time interval. As a non-limiting example, the synthetic nanoparticle may be formulated to release the polynucleotides, primary constructs and/or mmRNA after 24 hours and/or at a pH of 4.5 (see International Pub. Nos. WO2010138193 and WO2010138194 and US Pub Nos. US20110020388 and US20110027217, each of which is herein incorporated by reference in their entireties).

[00293] In one embodiment, the synthetic nanocarriers may be formulated for controlled and/or sustained release of the polynucleotides, primary constructs and/or mmRNA described herein. As a non-limiting example, the synthetic nanocarriers for sustained release may be formulated by methods known in the art, described herein and/or as described in International Pub No. WO2010138192 and US Pub No. 20100303850, each of which is herein incorporated by reference in their entireties.

[00294] Polymers, Biodegradable Nanoparticles, and Core-Shell Nanoparticles

[00295] The polynucleotide, primary construct, and mmRNA of the invention can be formulated using natural and/or synthetic polymers. Non-limiting examples of polymers which may be used for delivery include, but are not limited to, Dynamic POLYCONJUGATE™ formulations from MIRUS® Bio (Madison, WI) and Roche Madison (Madison, WI), PHASERX™ polymer formulations such as, without limitation, SMARTT POLYMER TECHNOLOGY™ (Seattle, WA), DMRI/DOPE, poloxamer, VAXFECTIN® adjuvant from Vical (San Diego, CA), chitosan, cyclodextrin from Calando Pharmaceuticals (Pasadena, CA), dendrimers and poly(lactic-co-glycolic acid) (PLGA) polymers. RONDEL™ (RNAi/Oligonucleotide Nanoparticle Delivery) polymers (Arrowhead Research Corporation, Pasadena, CA) and pH responsive co-block polymers such as, but not limited to, PHASERX™ (Seattle, WA).

[00296] A non-limiting example of PLGA formulations include, but are not limited to, PLGA injectable depots (e.g., ELIGARD® which is formed by dissolving PLGA in 66% N-methyl-2-pyrrolidone (NMP) and the remainder being aqueous solvent and leuprolide. Once injected, the PLGA and leuprolide peptide precipitates into the subcutaneous space).

[00297] Many of these polymer approaches have demonstrated efficacy in delivering oligonucleotides *in vivo* into the cell cytoplasm (reviewed in deFougerolles *Hum Gene Ther.* 2008 19:125-132; herein incorporated by reference in its entirety). Two polymer approaches that have yielded robust *in vivo* delivery of nucleic acids, in this case with small interfering RNA (siRNA), are dynamic polyconjugates and cyclodextrin-based nanoparticles. The first of these delivery approaches uses dynamic polyconjugates and has been shown *in vivo* in mice to effectively deliver siRNA and silence endogenous target mRNA in hepatocytes (Rozema et al., *Proc Natl Acad Sci U S A.* 2007 104:12982-12887). This particular approach is a multicomponent polymer system whose key features include a membrane-active polymer to which nucleic acid, in this case siRNA, is covalently coupled via a disulfide bond and where both PEG (for charge masking) and *N*-acetylgalactosamine (for hepatocyte targeting) groups are linked via pH-sensitive bonds (Rozema et al., *Proc Natl Acad Sci U S A.* 2007 104:12982-12887). On binding to the hepatocyte and entry into the endosome, the polymer complex disassembles in the low-pH environment, with the polymer exposing its positive charge, leading to endosomal escape and cytoplasmic release of the siRNA from the polymer. Through replacement of the *N*-acetylgalactosamine group with a mannose group, it was shown one could alter targeting from asialoglycoprotein receptor-expressing hepatocytes to sinusoidal

endothelium and Kupffer cells. Another polymer approach involves using transferrin-targeted cyclodextrin-containing polycation nanoparticles. These nanoparticles have demonstrated targeted silencing of the *EWS-FLI1* gene product in transferrin receptor-expressing Ewing's sarcoma tumor cells (Hu-Lieskovan *et al.*, *Cancer Res.* 2005 65: 8984-8982) and siRNA formulated in these nanoparticles was well tolerated in non-human primates (Heidel *et al.*, *Proc Natl Acad Sci USA* 2007 104:5715-21). Both of these delivery strategies incorporate rational approaches using both targeted delivery and endosomal escape mechanisms.

[00298] The polymer formulation can permit the sustained or delayed release of the polynucleotide, primary construct, or mmRNA (e.g., following intramuscular or subcutaneous injection). The altered release profile for the polynucleotide, primary construct, or mmRNA can result in, for example, translation of an encoded protein over an extended period of time. The polymer formulation may also be used to increase the stability of the polynucleotide, primary construct, or mmRNA. Biodegradable polymers have been previously used to protect nucleic acids other than mmRNA from degradation and been shown to result in sustained release of payloads *in vivo* (Rozema *et al.*, *Proc Natl Acad Sci U S A.* 2007 104:12982-12887; Sullivan *et al.*, *Expert Opin Drug Deliv.* 2010 7:1433-1446; Convertine *et al.*, *Biomacromolecules.* 2010 Oct 1; Chu *et al.*, *Acc Chem Res.* 2012 Jan 13; Manganiello *et al.*, *Biomaterials.* 2012 33:2301-2309; Benoit *et al.*, *Biomacromolecules.* 2011 12:2708-2714; Singha *et al.*, *Nucleic Acid Ther.* 2011 2:133-147; deFougerolles *Hum Gene Ther.* 2008 19:125-132; Schaffert and Wagner, *Gene Ther.* 2008 16:1131-1138; Chaturvedi *et al.*, *Expert Opin Drug Deliv.* 2011 8:1455-1468; Davis, *Mol Pharm.* 2009 6:659-668; Davis, *Nature* 2010 464:1067-1070; each of which is herein incorporated by reference in its entirety).

[00299] In one embodiment, the pharmaceutical compositions may be sustained release formulations. In a further embodiment, the sustained release formulations may be for subcutaneous delivery. Sustained release formulations may include, but are not limited to, PLGA microspheres, ethylene vinyl acetate (EVAc), poloxamer, GELSITE® (Nanotherapeutics, Inc. Alachua, FL), HYLENEX® (Halozyme Therapeutics, San Diego CA), surgical sealants such as fibrinogen polymers (Ethicon Inc. Cornelia, GA), TISSELL® (Baxter International, Inc Deerfield, IL), PEG-based sealants, and COSEAL® (Baxter International, Inc Deerfield, IL).

[00300] As a non-limiting example modified mRNA may be formulated in PLGA microspheres by preparing the PLGA microspheres with tunable release rates (e.g., days

and weeks) and encapsulating the modified mRNA in the PLGA microspheres while maintaining the integrity of the modified mRNA during the encapsulation process. EVAc are non-biodegradable, biocompatible polymers which are used extensively in pre-clinical sustained release implant applications (e.g., extended release products Ocusert a pilocarpine ophthalmic insert for glaucoma or progestasert a sustained release progesterone intrauterine device; transdermal delivery systems Testoderm, Duragesic and Selegiline; catheters). Poloxamer F-407 NF is a hydrophilic, non-ionic surfactant triblock copolymer of polyoxyethylene-polyoxypropylene-polyoxyethylene having a low viscosity at temperatures less than 5°C and forms a solid gel at temperatures greater than 15°C. PEG-based surgical sealants comprise two synthetic PEG components mixed in a delivery device which can be prepared in one minute, seals in 3 minutes and is reabsorbed within 30 days. GELSITE® and natural polymers are capable of in-situ gelation at the site of administration. They have been shown to interact with protein and peptide therapeutic candidates through ionic interaction to provide a stabilizing effect.

[00301] Polymer formulations can also be selectively targeted through expression of different ligands as exemplified by, but not limited by, folate, transferrin, and N-acetylgalactosamine (GalNAc) (Benoit et al., *Biomacromolecules*. 2011 12:2708-2714; Rozema et al., *Proc Natl Acad Sci U S A*. 2007 104:12982-12887; Davis, *Mol Pharm*. 2009 6:659-668; Davis, *Nature* 2010 464:1067-1070; each of which is herein incorporated by reference in its entirety).

[00302] The polynucleotides, primary constructs and/or mmRNA of the invention may be formulated with or in a polymeric compound. The polymer may include at least one polymer such as, but not limited to, polyethenes, polyethylene glycol (PEG), poly(L-lysine)(PLL), PEG grafted to PLL, cationic lipopolymer, biodegradable cationic lipopolymer, polyethyleneimine (PEI), cross-linked branched poly(alkylene imines), a polyamine derivative, a modified poloxamer, a biodegradable polymer, biodegradable block copolymer, biodegradable random copolymer, biodegradable polyester copolymer, biodegradable polyester block copolymer, biodegradable polyester block random copolymer, linear biodegradable copolymer, poly[α -(4-aminobutyl)-L-glycolic acid] (PAGA), biodegradable cross-linked cationic multi-block copolymers, polycarbonates, polyanhydrides, polyhydroxyacids, polypropylfumerates, polycaprolactones, polyamides, polyacetals, polyethers, polyesters, poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polycyanoacrylates, polyureas, polystyrenes, polyamines, polylysine, poly(ethylene

imine), poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), acrylic polymers, amine-containing polymers or combinations thereof.

[00303] As a non-limiting example, the polynucleotides, primary constructs and/or mmRNA of the invention may be formulated with the polymeric compound of PEG grafted with PLL as described in U.S. Pat. No. 6,177,274 herein incorporated by reference in its entirety. The formulation may be used for transfecting cells *in vitro* or for *in vivo* delivery of the polynucleotides, primary constructs and/or mmRNA. In another example, the polynucleotides, primary constructs and/or mmRNA may be suspended in a solution or medium with a cationic polymer, in a dry pharmaceutical composition or in a solution that is capable of being dried as described in U.S. Pub. Nos. 20090042829 and 20090042825 each of which are herein incorporated by reference in their entireties.

[00304] As another non-limiting example the polynucleotides, primary constructs or mmRNA of the invention may be formulated with a PLGA-PEG block copolymer (see US Pub. No. US20120004293 and US Pat No. 8,236,330, each of which is herein incorporated by reference in their entireties). As a non-limiting example, the polynucleotides, primary constructs or mmRNA of the invention may be formulated with a diblock copolymer of PEG and PLA or PEG and PLGA (see US Pat No 8,246,968, herein incorporated by reference in its entirety).

[00305] A polyamine derivative may be used to deliver nucleic acids or to treat and/or prevent a disease or to be included in an implantable or injectable device (U.S. Pub. No. 20100260817 herein incorporated by reference in its entirety). As a non-limiting example, a pharmaceutical composition may include the modified nucleic acids and mmRNA and the polyamine derivative described in U.S. Pub. No. 20100260817 (the contents of which are incorporated herein by reference in its entirety).

[00306] The polynucleotides, primary constructs or mmRNA of the invention may be formulated with at least one acrylic polymer. Acrylic polymers include but are not limited to, acrylic acid, methacrylic acid, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, amino alkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), polycyanoacrylates and combinations thereof.

[00307] In one embodiment, the polynucleotides, primary constructs or mmRNA of the present invention may be formulated with at least one polymer described in International Publication Nos. WO2011115862, WO2012082574 and WO2012068187, each of which is herein incorporated by reference in their entireties. In another embodiment the

polynucleotides, primary constructs or mmRNA of the present invention may be formulated with a polymer of formula Z as described in WO2011115862, herein incorporated by reference in its entirety. In yet another embodiment, the polynucleotides, primary constructs or mmRNA may be formulated with a polymer of formula Z, Z' or Z'' as described in WO2012082574 or WO2012068187, each of which are herein incorporated by reference in their entireties. The polymers formulated with the modified RNA of the present invention may be synthesized by the methods described in WO2012082574 or WO2012068187, each of which is herein incorporated by reference in their entireties.

[00308] Formulations of polynucleotides, primary constructs or mmRNA of the invention may include at least one amine-containing polymer such as, but not limited to polylysine, polyethylene imine, poly(amidoamine) dendrimers or combinations thereof.

[00309] For example, the polynucleotides, primary constructs and/or mmRNA of the invention may be formulated in a pharmaceutical compound including a poly(alkylene imine), a biodegradable cationic lipopolymer, a biodegradable block copolymer, a biodegradable polymer, or a biodegradable random copolymer, a biodegradable polyester block copolymer, a biodegradable polyester polymer, a biodegradable polyester random copolymer, a linear biodegradable copolymer, PAGA, a biodegradable cross-linked cationic multi-block copolymer or combinations thereof. The biodegradable cationic lipopolymer may be made by methods known in the art and/or described in U.S. Pat. No. 6,696,038, U.S. App. Nos. 20030073619 and 20040142474 each of which is herein incorporated by reference in their entireties. The poly(alkylene imine) may be made using methods known in the art and/or as described in U.S. Pub. No. 20100004315, herein incorporated by reference in its entirety. The biodegradable polymer, biodegradable block copolymer, the biodegradable random copolymer, biodegradable polyester block copolymer, biodegradable polyester polymer, or biodegradable polyester random copolymer may be made using methods known in the art and/or as described in U.S. Pat. Nos. 6,517,869 and 6,267,987, the contents of which are each incorporated herein by reference in its entirety. The linear biodegradable copolymer may be made using methods known in the art and/or as described in U.S. Pat. No. 6,652,886. The PAGA polymer may be made using methods known in the art and/or as described in U.S. Pat. Nos. 6,217,912 herein incorporated by reference in its entirety. The PAGA polymer may be copolymerized to form a copolymer or block copolymer with polymers such as but not limited to, poly-L-lysine, polyarginine, polyornithine, histones, avidin, protamines,

polylactides and poly(lactide-co-glycolides). The biodegradable cross-linked cationic multi-block copolymers may be made by methods known in the art and/or as described in U.S. Pat. No. 8,057,821 or U.S. Pub. No. 2012009145 each of which is herein incorporated by reference in their entireties. For example, the multi-block copolymers may be synthesized using linear polyethyleneimine (LPEI) blocks which have distinct patterns as compared to branched polyethyleneimines. Further, the composition or pharmaceutical composition may be made by the methods known in the art, described herein, or as described in U.S. Pub. No. 20100004315 or U.S. Pat. Nos. 6,267,987 and 6,217,912 each of which is herein incorporated by reference in their entireties.

[00310] The polynucleotides, primary constructs, and mmRNA of the invention may be formulated with at least one degradable polyester which may contain polycationic side chains. Degradable polyesters include, but are not limited to, poly(serine ester), poly(L-lactide-co-L-lysine), poly(4-hydroxy-L-proline ester), and combinations thereof. In another embodiment, the degradable polyesters may include a PEG conjugation to form a PEGylated polymer.

[00311] In one embodiment, the polymers described herein may be conjugated to a lipid-terminating PEG. As a non-limiting example, PLGA may be conjugated to a lipid-terminating PEG forming PLGA-DSPE-PEG. As another non-limiting example, PEG conjugates for use with the present invention are described in International Publication No. WO2008103276, herein incorporated by reference in its entirety.

[00312] In one embodiment, the polynucleotides, primary constructs and/or mmRNA described herein may be conjugated with another compound. Non-limiting examples of conjugates are described in US Patent Nos. 7,964,578 and 7,833,992, each of which are herein incorporated by reference in their entireties. In another embodiment, the polynucleotides, primary constructs and/or mmRNA of the present invention may be conjugated with conjugates of formula 1-122 as described in US Patent Nos. 7,964,578 and 7,833,992, each of which are herein incorporated by reference in their entireties.

[00313] As described in U.S. Pub. No. 20100004313, herein incorporated by reference in its entirety, a gene delivery composition may include a nucleotide sequence and a poloxamer. For example, the polynucleotide, primary construct and/or mmRNA of the present invention may be used in a gene delivery composition with the poloxamer described in U.S. Pub. No. 20100004313.

[00314] In one embodiment, the polymer formulation of the present invention may be stabilized by contacting the polymer formulation, which may include a cationic carrier,

with a cationic lipopolymer which may be covalently linked to cholesterol and polyethylene glycol groups. The polymer formulation may be contacted with a cationic lipopolymer using the methods described in U.S. Pub. No. 20090042829 herein incorporated by reference in its entirety. The cationic carrier may include, but is not limited to, polyethylenimine, poly(trimethylenimine), poly(tetramethylenimine), polypropylenimine, aminoglycoside-polyamine, dideoxy-diamino- β -cyclodextrin, spermine, spermidine, poly(2-dimethylamino)ethyl methacrylate, poly(lysine), poly(histidine), poly(arginine), cationized gelatin, dendrimers, chitosan, 1,2-Dioleoyl-3-Trimethylammonium-Propane(DOTAP), N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride (DOTMA), 1-[2-(oleoyloxy)ethyl]-2-oleyl-3-(2-hydroxyethyl)imidazolium chloride (DOTIM), 2,3-dioleoyloxy-N-[2(sperminecarboxamido)ethyl]-N,N-dimethyl-1-propanaminium trifluoroacetate (DOSPA), 3B-[N—(N',N'-Dimethylaminoethane)-carbonyl]Cholesterol Hydrochloride (DC-Cholesterol HCl) diheptadecylamidoglycyl spermidine (DOGS), N,N-distearyl-N,N-dimethylammonium bromide (DDAB), N-(1,2-dimyristyloxyprop-3-yl)-N,N-dimethyl-N-hydroxyethyl ammonium bromide (DMRIE), N,N-dioleoyl-N,N-dimethylammonium chloride DODAC) and combinations thereof.

[00315] The polynucleotide, primary construct, and mmRNA of the invention can also be formulated as a nanoparticle using a combination of polymers, lipids, and/or other biodegradable agents, such as, but not limited to, calcium phosphate. Components may be combined in a core-shell, hybrid, and/or layer-by-layer architecture, to allow for fine-tuning of the nanoparticle so to delivery of the polynucleotide, primary construct and mmRNA may be enhanced (Wang et al., *Nat Mater.* 2006 5:791-796; Fuller et al., *Biomaterials.* 2008 29:1526-1532; DeKoker et al., *Adv Drug Deliv Rev.* 2011 63:748-761; Endres et al., *Biomaterials.* 2011 32:7721-7731; Su et al., *Mol Pharm.* 2011 Jun 6;8(3):774-87; herein incorporated by reference in its entirety).

[00316] Biodegradable calcium phosphate nanoparticles in combination with lipids and/or polymers have been shown to deliver polynucleotides, primary constructs and mmRNA *in vivo*. In one embodiment, a lipid coated calcium phosphate nanoparticle, which may also contain a targeting ligand such as anisamide, may be used to deliver the polynucleotide, primary construct and mmRNA of the present invention. For example, to effectively deliver siRNA in a mouse metastatic lung model a lipid coated calcium phosphate nanoparticle was used (Li et al., *J Contr Rel.* 2010 142: 416-421; Li et al., *J Contr Rel.* 2012 158:108-114; Yang et al., *Mol Ther.* 2012 20:609-615). This delivery

system combines both a targeted nanoparticle and a component to enhance the endosomal escape, calcium phosphate, in order to improve delivery of the siRNA.

[00317] In one embodiment, calcium phosphate with a PEG-polyanion block copolymer may be used to delivery polynucleotides, primary constructs and mmRNA (Kazikawa et al., J Contr Rel. 2004 97:345-356; Kazikawa et al., J Contr Rel. 2006 111:368-370).

[00318] In one embodiment, a PEG-charge-conversional polymer (Pitella et al., Biomaterials. 2011 32:3106-3114) may be used to form a nanoparticle to deliver the polynucleotides, primary constructs and mmRNA of the present invention. The PEG-charge-conversional polymer may improve upon the PEG-polyanion block copolymers by being cleaved into a polycation at acidic pH, thus enhancing endosomal escape.

[00319] The use of core-shell nanoparticles has additionally focused on a high-throughput approach to synthesize cationic cross-linked nanogel cores and various shells (Siegwart et al., Proc Natl Acad Sci U S A. 2011 108:12996-13001). The complexation, delivery, and internalization of the polymeric nanoparticles can be precisely controlled by altering the chemical composition in both the core and shell components of the nanoparticle. For example, the core-shell nanoparticles may efficiently deliver siRNA to mouse hepatocytes after they covalently attach cholesterol to the nanoparticle.

[00320] In one embodiment, a hollow lipid core comprising a middle PLGA layer and an outer neutral lipid layer containing PEG may be used to delivery of the polynucleotide, primary construct and mmRNA of the present invention. As a non-limiting example, in mice bearing a luciferase-expressing tumor, it was determined that the lipid-polymer-lipid hybrid nanoparticle significantly suppressed luciferase expression, as compared to a conventional lipoplex (Shi et al, Angew Chem Int Ed. 2011 50:7027-7031).

Peptides and Proteins

[00321] The synthetic polynucleotides and/or synthetic sgRNAs of the invention can be formulated with peptides and/or proteins in order to increase transfection of cells by the polynucleotide, primary construct, or mmRNA. In one embodiment, peptides such as, but not limited to, cell penetrating peptides and proteins and peptides that enable intracellular delivery may be used to deliver pharmaceutical formulations. A non-limiting example of a cell penetrating peptide which may be used with the pharmaceutical formulations of the present invention includes a cell-penetrating peptide sequence attached to polycations that facilitates delivery to the intracellular space, e.g., HIV-derived TAT peptide,

penetratins, transportans, or hCT derived cell-penetrating peptides (see, e.g., Caron et al., Mol. Ther. 3(3):310-8 (2001); Langel, Cell-Penetrating Peptides: Processes and Applications (CRC Press, Boca Raton FL, 2002); El-Andaloussi et al., Curr. Pharm. Des. 11(28):3597-611 (2003); and Deshayes et al., Cell. Mol. Life Sci. 62(16):1839-49 (2005), all of which are incorporated herein by reference). The compositions can also be formulated to include a cell penetrating agent, e.g., liposomes, which enhance delivery of the compositions to the intracellular space. polynucleotides, primary constructs, and mmRNA of the invention may be complexed to peptides and/or proteins such as, but not limited to, peptides and/or proteins from Aileron Therapeutics (Cambridge, MA) and Permeon Biologics (Cambridge, MA) in order to enable intracellular delivery (Cronican et al., ACS Chem. Biol. 2010 5:747-752; McNaughton et al., Proc. Natl. Acad. Sci. USA 2009 106:6111-6116; Sawyer, Chem Biol Drug Des. 2009 73:3-6; Verdine and Hilinski, Methods Enzymol. 2012;503:3-33; all of which are herein incorporated by reference in its entirety).

[00322] In one embodiment, the cell-penetrating polypeptide may comprise a first domain and a second domain. The first domain may comprise a supercharged polypeptide. The second domain may comprise a protein-binding partner. As used herein, "protein-binding partner" includes, but are not limited to, antibodies and functional fragments thereof, scaffold proteins, or peptides. The cell-penetrating polypeptide may further comprise an intracellular binding partner for the protein-binding partner. The cell-penetrating polypeptide may be capable of being secreted from a cell where the polynucleotide, primary construct, or mmRNA may be introduced.

[00323] Formulations of the including peptides or proteins may be used to increase cell transfection by the polynucleotide, primary construct, or mmRNA, alter the biodistribution of the polynucleotide, primary construct, or mmRNA (e.g., by targeting specific tissues or cell types), and/or increase the translation of encoded protein.

Cells

[00324] The polynucleotide, primary construct, and mmRNA of the invention can be transfected *ex vivo* into cells, which are subsequently transplanted into a subject. As non-limiting examples, the pharmaceutical compositions may include red blood cells to deliver modified RNA to liver and myeloid cells, virosomes to deliver modified RNA in virus-like particles (VLPs), and electroporated cells such as, but not limited to, from MAXCYTE® (Gaithersburg, MD) and from ERYTECH® (Lyon, France) to deliver

modified RNA. Examples of use of red blood cells, viral particles and electroporated cells to deliver payloads other than mmRNA have been documented (Godfrin et al., *Expert Opin Biol Ther.* 2012 12:127-133; Fang et al., *Expert Opin Biol Ther.* 2012 12:385-389; Hu et al., *Proc Natl Acad Sci U S A.* 2011 108:10980-10985; Lund et al., *Pharm Res.* 2010 27:400-420; Huckriede et al., *J Liposome Res.* 2007;17:39-47; Cusi, *Hum Vaccin.* 2006 2:1-7; de Jonge et al., *Gene Ther.* 2006 13:400-411; all of which are herein incorporated by reference in its entirety).

[00325] The polynucleotides, primary constructs and mmRNA may be delivered in synthetic VLPs synthesized by the methods described in International Pub No. WO2011085231 and US Pub No. 20110171248, each of which is herein incorporated by reference in their entirety.

[00326] Cell-based formulations of the polynucleotide, primary construct, and mmRNA of the invention may be used to ensure cell transfection (e.g., in the cellular carrier), alter the biodistribution of the polynucleotide, primary construct, or mmRNA (e.g., by targeting the cell carrier to specific tissues or cell types), and/or increase the translation of encoded protein.

[00327] A variety of methods are known in the art and suitable for introduction of nucleic acid into a cell, including viral and non-viral mediated techniques. Examples of typical non-viral mediated techniques include, but are not limited to, electroporation, calcium phosphate mediated transfer, nucleofection, sonoporation, heat shock, magnetofection, liposome mediated transfer, microinjection, microprojectile mediated transfer (nanoparticles), cationic polymer mediated transfer (DEAE-dextran, polyethylenimine, polyethylene glycol (PEG) and the like) or cell fusion.

[00328] The technique of sonoporation, or cellular sonication, is the use of sound (e.g., ultrasonic frequencies) for modifying the permeability of the cell plasma membrane. Sonoporation methods are known to those in the art and are used to deliver nucleic acids *in vivo* (Yoon and Park, *Expert Opin Drug Deliv.* 2010 7:321-330; Postema and Gilja, *Curr Pharm Biotechnol.* 2007 8:355-361; Newman and Bettinger, *Gene Ther.* 2007 14:465-475; all herein incorporated by reference in their entirety). Sonoporation methods are known in the art and are also taught for example as it relates to bacteria in US Patent Publication 20100196983 and as it relates to other cell types in, for example, US Patent Publication 20100009424, each of which are incorporated herein by reference in their entirety.

[00329] Electroporation techniques are also well known in the art and are used to deliver nucleic acids *in vivo* and clinically (Andre et al., Curr Gene Ther. 2010 10:267-280; Chiarella et al., Curr Gene Ther. 2010 10:281-286; Hojman, Curr Gene Ther. 2010 10:128-138; all herein incorporated by reference in their entirety). In one embodiment, polynucleotides, primary constructs or mmRNA may be delivered by electroporation as described in Example 26.

Hyaluronidase

[00330] The intramuscular or subcutaneous localized injection of polynucleotide, primary construct, or mmRNA of the invention can include hyaluronidase, which catalyzes the hydrolysis of hyaluronan. By catalyzing the hydrolysis of hyaluronan, a constituent of the interstitial barrier, hyaluronidase lowers the viscosity of hyaluronan, thereby increasing tissue permeability (Frost, Expert Opin. Drug Deliv. (2007) 4:427-440; herein incorporated by reference in its entirety). It is useful to speed their dispersion and systemic distribution of encoded proteins produced by transfected cells. Alternatively, the hyaluronidase can be used to increase the number of cells exposed to a polynucleotide, primary construct, or mmRNA of the invention administered intramuscularly or subcutaneously.

Nanoparticle Mimics

[00331] The polynucleotide, primary construct or mmRNA of the invention may be encapsulated within and/or absorbed to a nanoparticle mimic. A nanoparticle mimic can mimic the delivery function organisms or particles such as, but not limited to, pathogens, viruses, bacteria, fungus, parasites, prions and cells. As a non-limiting example the polynucleotide, primary construct or mmRNA of the invention may be encapsulated in a non-viron particle which can mimic the delivery function of a virus (see International Pub. No. WO2012006376 herein incorporated by reference in its entirety).

Nanotubes

[00332] The polynucleotides, primary constructs or mmRNA of the invention can be attached or otherwise bound to at least one nanotube such as, but not limited to, rosette nanotubes, rosette nanotubes having twin bases with a linker, carbon nanotubes and/or single-walled carbon nanotubes, The polynucleotides, primary constructs or mmRNA may be bound to the nanotubes through forces such as, but not limited to, steric, ionic, covalent and/or other forces.

[00333] In one embodiment, the nanotube can release one or more polynucleotides, primary constructs or mmRNA into cells. The size and/or the surface structure of at least one nanotube may be altered so as to govern the interaction of the nanotubes within the body and/or to attach or bind to the polynucleotides, primary constructs or mmRNA disclosed herein. In one embodiment, the building block and/or the functional groups attached to the building block of the at least one nanotube may be altered to adjust the dimensions and/or properties of the nanotube. As a non-limiting example, the length of the nanotubes may be altered to hinder the nanotubes from passing through the holes in the walls of normal blood vessels but still small enough to pass through the larger holes in the blood vessels of tumor tissue.

[00334] In one embodiment, at least one nanotube may also be coated with delivery enhancing compounds including polymers, such as, but not limited to, polyethylene glycol. In another embodiment, at least one nanotube and/or the polynucleotides, primary constructs or mmRNA may be mixed with pharmaceutically acceptable excipients and/or delivery vehicles.

[00335] In one embodiment, the polynucleotides, primary constructs or mmRNA are attached and/or otherwise bound to at least one rosette nanotube. The rosette nanotubes may be formed by a process known in the art and/or by the process described in International Publication No. WO2012094304, herein incorporated by reference in its entirety. At least one polynucleotide, primary construct and/or mmRNA may be attached and/or otherwise bound to at least one rosette nanotube by a process as described in International Publication No. WO2012094304, herein incorporated by reference in its entirety, where rosette nanotubes or modules forming rosette nanotubes are mixed in aqueous media with at least one polynucleotide, primary construct and/or mmRNA under conditions which may cause at least one polynucleotide, primary construct or mmRNA to attach or otherwise bind to the rosette nanotubes.

Conjugates

[00336] The polynucleotides, primary constructs, and mmRNA of the invention include conjugates, such as a polynucleotide, primary construct, or mmRNA covalently linked to a carrier or targeting group, or including two encoding regions that together produce a fusion protein (e.g., bearing a targeting group and therapeutic protein or peptide).

[00337] The conjugates of the invention include a naturally occurring substance, such as a protein (e.g., human serum albumin (HSA), low-density lipoprotein (LDL), high-

density lipoprotein (HDL), or globulin); an carbohydrate (e.g., a dextran, pullulan, chitin, chitosan, inulin, cyclodextrin or hyaluronic acid); or a lipid. The ligand may also be a recombinant or synthetic molecule, such as a synthetic polymer, e.g., a synthetic polyamino acid, an oligonucleotide (e.g. an aptamer). Examples of polyamino acids include polyamino acid is a polylysine (PLL), poly L-aspartic acid, poly L-glutamic acid, styrene-maleic acid anhydride copolymer, poly(L-lactide-co-glycolid) copolymer, divinyl ether-maleic anhydride copolymer, N-(2-hydroxypropyl)methacrylamide copolymer (HMPA), polyethylene glycol (PEG), polyvinyl alcohol (PVA), polyurethane, poly(2-ethylacrylic acid), N-isopropylacrylamide polymers, or polyphosphazine. Example of polyamines include: polyethylenimine, polylysine (PLL), spermine, spermidine, polyamine, pseudopeptide-polyamine, peptidomimetic polyamine, dendrimer polyamine, arginine, amidine, protamine, cationic lipid, cationic porphyrin, quaternary salt of a polyamine, or an alpha helical peptide.

[00338] Representative U.S. patents that teach the preparation of polynucleotide conjugates, particularly to RNA, include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941; 6,294,664; 6,320,017; 6,576,752; 6,783,931; 6,900,297; 7,037,646; each of which is herein incorporated by reference in their entireties.

[00339] In one embodiment, the conjugate of the present invention may function as a carrier for the polynucleotides, primary constructs and/or mmRNA of the present invention. The conjugate may comprise a cationic polymer such as, but not limited to, polyamine, polylysine, polyalkylenimine, and polyethylenimine which may be grafted to with poly(ethylene glycol). As a non-limiting example, the conjugate may be similar to the polymeric conjugate and the method of synthesizing the polymeric conjugate described in U.S. Pat. No. 6,586,524 herein incorporated by reference in its entirety.

[00340] The conjugates can also include targeting groups, e.g., a cell or tissue targeting agent, e.g., a lectin, glycoprotein, lipid or protein, e.g., an antibody, that binds to a specified cell type such as a kidney cell. A targeting group can be a thyrotropin,

melanotropin, lectin, glycoprotein, surfactant protein A, Mucin carbohydrate, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-gulucosamine multivalent mannose, multivalent fucose, glycosylated polyaminoacids, multivalent galactose, transferrin, bisphosphonate, polyglutamate, polyaspartate, a lipid, cholesterol, a steroid, bile acid, folate, vitamin B12, biotin, an RGD peptide, an RGD peptide mimetic or an aptamer.

[00341] Targeting groups can be proteins, e.g., glycoproteins, or peptides, e.g., molecules having a specific affinity for a co-ligand, or antibodies e.g., an antibody, that binds to a specified cell type such as a cancer cell, endothelial cell, or bone cell. Targeting groups may also include hormones and hormone receptors. They can also include non-peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-gulucosamine multivalent mannose, multivalent fucose, or aptamers. The ligand can be, for example, a lipopolysaccharide, or an activator of p38 MAP kinase.

[00342] The targeting group can be any ligand that is capable of targeting a specific receptor. Examples include, without limitation, folate, GalNAc, galactose, mannose, mannose-6P, aptamers, integrin receptor ligands, chemokine receptor ligands, transferrin, biotin, serotonin receptor ligands, PSMA, endothelin, GCPII, somatostatin, LDL, and HDL ligands. In particular embodiments, the targeting group is an aptamer. The aptamer can be unmodified or have any combination of modifications disclosed herein.

[00343] In one embodiment, pharmaceutical compositions of the present invention may include chemical modifications such as, but not limited to, modifications similar to locked nucleic acids.

[00344] Representative U.S. Patents that teach the preparation of locked nucleic acid (LNA) such as those from Santaris, include, but are not limited to, the following: U.S. Pat. Nos. 6,268,490; 6,670,461; 6,794,499; 6,998,484; 7,053,207; 7,084,125; and 7,399,845, each of which is herein incorporated by reference in its entirety.

[00345] Representative U.S. patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found, for example, in Nielsen *et al.*, Science, 1991, 254, 1497-1500.

[00346] Some embodiments featured in the invention include polynucleotides, primary constructs or mmRNA with phosphorothioate backbones and oligonucleosides with other

modified backbones, and in particular --CH₂--NH--CH₂--, --CH₂--N(CH₃)--O--CH₂-- [known as a methylene (methylimino) or MMI backbone], --CH₂--O--N(CH₃)--CH₂--, --CH₂--N(CH₃)--N(CH₃)--CH₂-- and --N(CH₃)--CH₂--CH₂-- [wherein the native phosphodiester backbone is represented as --O—P(O)₂--O--CH₂--] of the above-referenced U.S. Pat. No. 5,489,677, and the amide backbones of the above-referenced U.S. Pat. No. 5,602,240. In some embodiments, the polynucleotides featured herein have morpholino backbone structures of the above-referenced U.S. Pat. No. 5,034,506.

[00347] Modifications at the 2' position may also aid in delivery. Preferably, modifications at the 2' position are not located in a polypeptide-coding sequence, i.e., not in a translatable region. Modifications at the 2' position may be located in a 5'UTR, a 3'UTR and/or a tailing region. Modifications at the 2' position can include one of the following at the 2' position: H (i.e., 2'-deoxy); F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Exemplary suitable modifications include O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. In other embodiments, the polynucleotides, primary constructs or mmRNA include one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties, or a group for improving the pharmacodynamic properties, and other substituents having similar properties. In some embodiments, the modification includes a 2'-methoxyethoxy (2'-O--CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78:486-504) i.e., an alkoxy-alkoxy group. Another exemplary modification is 2'-dimethylaminoethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples herein below, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O--CH₂--O--CH₂--N(CH₂)₂, also described in examples herein below. Other modifications include 2'-methoxy (2'-OCH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide. Polynucleotides of the

invention may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative U.S. patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Pat. Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920 and each of which is herein incorporated by reference.

[00348] In still other embodiments, the polynucleotide, primary construct, or mmRNA is covalently conjugated to a cell penetrating polypeptide. The cell-penetrating peptide may also include a signal sequence. The conjugates of the invention can be designed to have increased stability; increased cell transfection; and/or altered the biodistribution (e.g., targeted to specific tissues or cell types).

Self-Assembled Nucleic Acid Nanoparticles

[00349] Self-assembled nanoparticles have a well-defined size which may be precisely controlled as the nucleic acid strands may be easily reprogrammable. For example, the optimal particle size for a cancer-targeting nanodelivery carrier is 20-100 nm as a diameter greater than 20 nm avoids renal clearance and enhances delivery to certain tumors through enhanced permeability and retention effect. Using self-assembled nucleic acid nanoparticles a single uniform population in size and shape having a precisely controlled spatial orientation and density of cancer-targeting ligands for enhanced delivery. As a non-limiting example, oligonucleotide nanoparticles are prepared using programmable self-assembly of short DNA fragments and therapeutic siRNAs. These nanoparticles are molecularly identical with controllable particle size and target ligand location and density. The DNA fragments and siRNAs self-assembled into a one-step reaction to generate DNA/siRNA tetrahedral nanoparticles for targeted *in vivo* delivery. (Lee et al., Nature Nanotechnology 2012 7:389-393).

Excipients

[00350] Pharmaceutical formulations may additionally comprise a pharmaceutically acceptable excipient, which, as used herein, includes any and all solvents, dispersion media, diluents, or other liquid vehicles, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's *The Science and Practice of Pharmacy*, 21st Edition, A. R. Gennaro (Lippincott, Williams &

Wilkins, Baltimore, MD, 2006; incorporated herein by reference) discloses various excipients used in formulating pharmaceutical compositions and known techniques for the preparation thereof. Except insofar as any conventional excipient medium is incompatible with a substance or its derivatives, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutical composition, its use is contemplated to be within the scope of this invention.

[00351] In some embodiments, a pharmaceutically acceptable excipient is at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% pure. In some embodiments, an excipient is approved for use in humans and for veterinary use. In some embodiments, an excipient is approved by United States Food and Drug Administration. In some embodiments, an excipient is pharmaceutical grade. In some embodiments, an excipient meets the standards of the United States Pharmacopoeia (USP), the European Pharmacopoeia (EP), the British Pharmacopoeia, and/or the International Pharmacopoeia.

[00352] Pharmaceutically acceptable excipients used in the manufacture of pharmaceutical compositions include, but are not limited to, inert diluents, dispersing and/or granulating agents, surface active agents and/or emulsifiers, disintegrating agents, binding agents, preservatives, buffering agents, lubricating agents, and/or oils. Such excipients may optionally be included in pharmaceutical compositions.

[00353] Exemplary diluents include, but are not limited to, calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, *etc.*, and/or combinations thereof.

[00354] Exemplary granulating and/or dispersing agents include, but are not limited to, potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinylpyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (VEEGUM[®]), sodium lauryl sulfate, quaternary ammonium compounds, *etc.*, and/or combinations thereof.

[00355] Exemplary surface active agents and/or emulsifiers include, but are not limited to, natural emulsifiers (e.g. acacia, agar, alginic acid, sodium alginate, tragacanth, chondrux, cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin), colloidal clays (e.g. bentonite [aluminum silicate] and VEEGUM[®] [magnesium aluminum silicate]), long chain amino acid derivatives, high molecular weight alcohols (e.g. stearyl alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (e.g. carboxy polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan, cellulosic derivatives (e.g. carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan fatty acid esters (e.g. polyoxyethylene sorbitan monolaurate [TWEEN[®]20], polyoxyethylene sorbitan [TWEENn[®]60], polyoxyethylene sorbitan monooleate [TWEEN[®]80], sorbitan monopalmitate [SPAN[®]40], sorbitan monostearate [Span[®]60], sorbitan tristearate [Span[®]65], glyceryl monooleate, sorbitan monooleate [SPAN[®]80]), polyoxyethylene esters (e.g. polyoxyethylene monostearate [MYRJ[®]45], polyoxyethylene hydrogenated castor oil, polyethoxylated castor oil, polyoxymethylene stearate, and SOLUTOL[®]), sucrose fatty acid esters, polyethylene glycol fatty acid esters (e.g. CREMOPHOR[®]), polyoxyethylene ethers, (e.g. polyoxyethylene lauryl ether [BRIJ[®]30]), poly(vinyl-pyrrolidone), diethylene glycol monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid, ethyl laurate, sodium lauryl sulfate, PLUORINC[®]F 68, POLOXAMER[®]188, cetrimonium bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, etc. and/or combinations thereof.

[00356] Exemplary binding agents include, but are not limited to, starch (e.g. cornstarch and starch paste); gelatin; sugars (e.g. sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol,); natural and synthetic gums (e.g. acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum silicate (Veegum[®]), and larch arabogalactan); alginates; polyethylene oxide; polyethylene glycol; inorganic calcium salts; silicic acid; polymethacrylates; waxes; water; alcohol; *etc.*; and combinations thereof.

[00357] Exemplary preservatives may include, but are not limited to, antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, alcohol preservatives, acidic preservatives, and/or other preservatives. Exemplary antioxidants include, but are not limited to, alpha tocopherol, ascorbic acid, acorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and/or sodium sulfite. Exemplary chelating agents include ethylenediaminetetraacetic acid (EDTA), citric acid monohydrate, disodium edetate, dipotassium edetate, edetic acid, fumaric acid, malic acid, phosphoric acid, sodium edetate, tartaric acid, and/or trisodium edetate. Exemplary antimicrobial preservatives include, but are not limited to, benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and/or thimerosal. Exemplary antifungal preservatives include, but are not limited to, butyl paraben, methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and/or sorbic acid. Exemplary alcohol preservatives include, but are not limited to, ethanol, polyethylene glycol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and/or phenylethyl alcohol. Exemplary acidic preservatives include, but are not limited to, vitamin A, vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroacetic acid, ascorbic acid, sorbic acid, and/or phytic acid. Other preservatives include, but are not limited to, tocopherol, tocopherol acetate, deteroxime mesylate, cetrimide, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium metabisulfite, GLYDANT PLUS[®], PHENONIP[®], methylparaben, GERMALL[®] 115, GERMABEN[®] II, NEOLONE[™], KATHON[™], and/or EUXYL[®].

[00358] Exemplary buffering agents include, but are not limited to, citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium gluconate, calcium gluceptate, calcium gluconate, D-gluconic acid, calcium glycerophosphate, calcium lactate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium

acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, Ringer's solution, ethyl alcohol, *etc.*, and/or combinations thereof.

[00359] Exemplary lubricating agents include, but are not limited to, magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behanate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, *etc.*, and combinations thereof.

[00360] Exemplary oils include, but are not limited to, almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macademia nut, mallow, mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughly, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soybean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils. Exemplary oils include, but are not limited to, butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 360, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and/or combinations thereof.

[00361] Excipients such as cocoa butter and suppository waxes, coloring agents, coating agents, sweetening, flavoring, and/or perfuming agents can be present in the composition, according to the judgment of the formulator.

Delivery

[00362] The present disclosure encompasses the delivery of polynucleotides, primary constructs or mmRNA for any of therapeutic, pharmaceutical, diagnostic or imaging by any appropriate route taking into consideration likely advances in the sciences of drug delivery. Delivery may be naked or formulated.

Naked Delivery

[00363] The polynucleotides, primary constructs or mmRNA of the present invention may be delivered to a cell naked. As used herein in, “naked” refers to delivering polynucleotides, primary constructs or mmRNA free from agents which promote transfection. For example, the polynucleotides, primary constructs or mmRNA delivered to the cell may contain no modifications. The naked polynucleotides, primary constructs or mmRNA may be delivered to the cell using routes of administration known in the art and described herein.

Formulated Delivery

[00364] The polynucleotides, primary constructs or mmRNA of the present invention may be formulated, using the methods described herein. The formulations may contain polynucleotides, primary constructs or mmRNA which may be modified and/or unmodified. The formulations may further include, but are not limited to, cell penetration agents, a pharmaceutically acceptable carrier, a delivery agent, a bioerodible or biocompatible polymer, a solvent, and a sustained-release delivery depot. The formulated polynucleotides, primary constructs or mmRNA may be delivered to the cell using routes of administration known in the art and described herein.

[00365] The compositions may also be formulated for direct delivery to an organ or tissue in any of several ways in the art including, but not limited to, direct soaking or bathing, via a catheter, by gels, powder, ointments, creams, gels, lotions, and/or drops, by using substrates such as fabric or biodegradable materials coated or impregnated with the compositions, and the like.

Administration

[00366] The polynucleotides, primary constructs or mmRNA of the present invention may be administered by any route which results in a therapeutically effective outcome. These include, but are not limited to enteral, gastroenteral, epidural, oral, transdermal, epidural (peridural), intracerebral (into the cerebrum), intracerebroventricular (into the cerebral ventricles), epicutaneous (application onto the skin), intradermal, (into the skin itself), subcutaneous (under the skin), nasal administration (through the nose), intravenous (into a vein), intraarterial (into an artery), intramuscular (into a muscle), intracardiac (into the heart), intraosseous infusion (into the bone marrow), intrathecal (into the spinal canal), intraperitoneal, (infusion or injection into the peritoneum), intravesical infusion, intravitreal, (through the eye), intracavernous injection, (into the

base of the penis), intravaginal administration, intrauterine, extra-amniotic administration, transdermal (diffusion through the intact skin for systemic distribution), transmucosal (diffusion through a mucous membrane), insufflation (snorting), sublingual, sublabial, enema, eye drops (onto the conjunctiva), or in ear drops. In specific embodiments, compositions may be administered in a way which allows them cross the blood-brain barrier, vascular barrier, or other epithelial barrier. Non-limiting routes of administration for the polynucleotides, primary constructs or mmRNA of the present invention are described below.

Parenteral and Injectable Administration

[00367] Liquid dosage forms for oral and parenteral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and/or elixirs. In addition to active ingredients, liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, oral compositions can include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and/or perfuming agents. In certain embodiments for parenteral administration, compositions are mixed with solubilizing agents such as CREMOPHOR[®], alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and/or combinations thereof.

[00368] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations may be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally acceptable diluents and/or solvents, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables.

[00369] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00370] In order to prolong the effect of an active ingredient, it is often desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle. Injectable depot forms are made by forming microcapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

Rectal and Vaginal Administration

[00371] Compositions for rectal or vaginal administration are typically suppositories which can be prepared by mixing compositions with suitable non-irritating excipients such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active ingredient.

Oral Administration

[00372] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, an active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient such as sodium citrate or dicalcium phosphate and/or fillers or extenders (*e.g.* starches, lactose, sucrose, glucose, mannitol, and silicic acid), binders (*e.g.* carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia), humectants (*e.g.* glycerol), disintegrating agents (*e.g.* agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate), solution retarding agents (*e.g.* paraffin), absorption accelerators (*e.g.* quaternary ammonium compounds), wetting agents (*e.g.* cetyl alcohol

and glycerol monostearate), absorbents (*e.g.* kaolin and bentonite clay), and lubricants (*e.g.* talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate), and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may comprise buffering agents.

Topical or Transdermal Administration

[00373] As described herein, compositions containing the polynucleotides, primary constructs or mmRNA of the invention may be formulated for administration topically. The skin may be an ideal target site for delivery as it is readily accessible. Gene expression may be restricted not only to the skin, potentially avoiding nonspecific toxicity, but also to specific layers and cell types within the skin.

[00374] The site of cutaneous expression of the delivered compositions will depend on the route of nucleic acid delivery. Three routes are commonly considered to deliver polynucleotides, primary constructs or mmRNA to the skin: (i) topical application (*e.g.* for local/regional treatment and/or cosmetic applications); (ii) intradermal injection (*e.g.* for local/regional treatment and/or cosmetic applications); and (iii) systemic delivery (*e.g.* for treatment of dermatologic diseases that affect both cutaneous and extracutaneous regions). polynucleotides, primary constructs or mmRNA can be delivered to the skin by several different approaches known in the art. Most topical delivery approaches have been shown to work for delivery of DNA, such as but not limited to, topical application of non-cationic liposome–DNA complex, cationic liposome–DNA complex, particle-mediated (gene gun), puncture-mediated gene transfections, and viral delivery approaches. After delivery of the nucleic acid, gene products have been detected in a number of different skin cell types, including, but not limited to, basal keratinocytes, sebaceous gland cells, dermal fibroblasts and dermal macrophages.

[00375] In one embodiment, the invention provides for a variety of dressings (*e.g.*, wound dressings) or bandages (*e.g.*, adhesive bandages) for conveniently and/or effectively carrying out methods of the present invention. Typically dressing or bandages may comprise sufficient amounts of pharmaceutical compositions and/or polynucleotides, primary constructs or mmRNA described herein to allow a user to perform multiple treatments of a subject(s).

[00376] In one embodiment, the invention provides for the polynucleotides, primary constructs or mmRNA compositions to be delivered in more than one injection.

[00377] In one embodiment, before topical and/or transdermal administration at least one area of tissue, such as skin, may be subjected to a device and/or solution which may increase permeability. In one embodiment, the tissue may be subjected to an abrasion device to increase the permeability of the skin (see U.S. Patent Publication No. 20080275468, herein incorporated by reference in its entirety). In another embodiment, the tissue may be subjected to an ultrasound enhancement device. An ultrasound enhancement device may include, but is not limited to, the devices described in U.S. Patent Publication No. 20040236268 and U.S. Patent Nos. 6,491,657 and 6,234,990; each of which is herein incorporated by reference in their entireties. Methods of enhancing the permeability of tissue are described in U.S. Patent Publication Nos. 20040171980 and 20040236268 and U.S. Patent No. 6,190,315; each of which are herein incorporated by reference in their entireties.

[00378] In one embodiment, a device may be used to increase permeability of tissue before delivering formulations of the polynucleotides, primary constructs and mmRNA described herein. The permeability of skin may be measured by methods known in the art and/or described in U.S. Patent No. 6,190,315, herein incorporated by reference in its entirety. As a non-limiting example, a modified mRNA formulation may be delivered by the drug delivery methods described in U.S. Patent No. 6,190,315, herein incorporated by reference in its entirety.

[00379] In another non-limiting example tissue may be treated with a eutectic mixture of local anesthetics (EMLA) cream before, during and/or after the tissue may be subjected to a device which may increase permeability. Katz et al. (*Anesth Analg* (2004); 98:371-76; herein incorporated by reference in its entirety) showed that using the EMLA cream in combination with a low energy, an onset of superficial cutaneous analgesia was seen as fast as 5 minutes after a pretreatment with a low energy ultrasound.

[00380] In one embodiment, enhancers may be applied to the tissue before, during, and/or after the tissue has been treated to increase permeability. Enhancers include, but are not limited to, transport enhancers, physical enhancers, and cavitation enhancers. Non-limiting examples of enhancers are described in U.S. Patent No. 6,190,315, herein incorporated by reference in its entirety.

[00381] In one embodiment, a device may be used to increase permeability of tissue before delivering formulations of polynucleotides, primary constructs and/or mmRNA described herein, which may further contain a substance that invokes an immune response. In another non-limiting example, a formulation containing a substance to

invoke an immune response may be delivered by the methods described in U.S. Publication Nos. 20040171980 and 20040236268; each of which is herein incorporated by reference in their entirety.

[00382] Dosage forms for topical and/or transdermal administration of a composition may include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants and/or patches. Generally, an active ingredient is admixed under sterile conditions with a pharmaceutically acceptable excipient and/or any needed preservatives and/or buffers as may be required.

[00383] Additionally, the present invention contemplates the use of transdermal patches, which often have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms may be prepared, for example, by dissolving and/or dispensing the compound in the proper medium. Alternatively or additionally, rate may be controlled by either providing a rate controlling membrane and/or by dispersing the compound in a polymer matrix and/or gel.

[00384] Formulations suitable for topical administration include, but are not limited to, liquid and/or semi liquid preparations such as liniments, lotions, oil in water and/or water in oil emulsions such as creams, ointments and/or pastes, and/or solutions and/or suspensions. Topically-administrable formulations may, for example, comprise from about 0.1% to about 10% (w/w) active ingredient, although the concentration of active ingredient may be as high as the solubility limit of the active ingredient in the solvent. Formulations for topical administration may further comprise one or more of the additional ingredients described herein.

Depot Administration

[00385] As described herein, in some embodiments, the composition is formulated in depots for extended release. Generally, a specific organ or tissue (a "target tissue") is targeted for administration.

[00386] In some aspects of the invention, the polynucleotides, primary constructs or mmRNA are spatially retained within or proximal to a target tissue. Provided are method of providing a composition to a target tissue of a mammalian subject by contacting the target tissue (which contains one or more target cells) with the composition under conditions such that the composition, in particular the nucleic acid component(s) of the composition, is substantially retained in the target tissue, meaning that at least 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the

composition is retained in the target tissue. Advantageously, retention is determined by measuring the amount of the nucleic acid present in the composition that enters one or more target cells. For example, at least 1, 5, 10, 20, 30, 40, 50, 60, 70, 80, 85, 90, 95, 96, 97, 98, 99, 99.9, 99.99 or greater than 99.99% of the nucleic acids administered to the subject are present intracellularly at a period of time following administration. For example, intramuscular injection to a mammalian subject is performed using an aqueous composition containing a ribonucleic acid and a transfection reagent, and retention of the composition is determined by measuring the amount of the ribonucleic acid present in the muscle cells.

[00387] Aspects of the invention are directed to methods of providing a composition to a target tissue of a mammalian subject, by contacting the target tissue (containing one or more target cells) with the composition under conditions such that the composition is substantially retained in the target tissue. The composition contains an effective amount of a polynucleotide, primary construct or mmRNA such that the polypeptide of interest is produced in at least one target cell. The compositions generally contain a cell penetration agent, although “naked” nucleic acid (such as nucleic acids without a cell penetration agent or other agent) is also contemplated, and a pharmaceutically acceptable carrier.

[00388] In some circumstances, the amount of a protein produced by cells in a tissue is desirably increased. Preferably, this increase in protein production is spatially restricted to cells within the target tissue. Thus, provided are methods of increasing production of a protein of interest in a tissue of a mammalian subject. A composition is provided that contains polynucleotides, primary constructs or mmRNA characterized in that a unit quantity of composition has been determined to produce the polypeptide of interest in a substantial percentage of cells contained within a predetermined volume of the target tissue.

[00389] In some embodiments, the composition includes a plurality of different polynucleotides, primary constructs or mmRNA, where one or more than one of the polynucleotides, primary constructs or mmRNA encodes a polypeptide of interest. Optionally, the composition also contains a cell penetration agent to assist in the intracellular delivery of the composition. A determination is made of the dose of the composition required to produce the polypeptide of interest in a substantial percentage of cells contained within the predetermined volume of the target tissue (generally, without inducing significant production of the polypeptide of interest in tissue adjacent to the

predetermined volume, or distally to the target tissue). Subsequent to this determination, the determined dose is introduced directly into the tissue of the mammalian subject.

[00390] In one embodiment, the invention provides for the polynucleotides, primary constructs or mmRNA to be delivered in more than one injection or by split dose injections.

[00391] In one embodiment, the invention may be retained near target tissue using a small disposable drug reservoir or patch pump. Non-limiting examples of patch pumps include those manufactured and/or sold by BD® (Franklin Lakes, NJ), Insulet Corporation (Bedford, MA), SteadyMed Therapeutics (San Francisco, CA), Medtronic (Minneapolis, MN), UniLife (York, PA), Valeritas (Bridgewater, NJ), and SpringLeaf Therapeutics (Boston, MA).

Pulmonary Administration

[00392] A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for pulmonary administration via the buccal cavity. Such a formulation may comprise dry particles which comprise the active ingredient and which have a diameter in the range from about 0.5 nm to about 7 nm or from about 1 nm to about 6 nm. Such compositions are suitably in the form of dry powders for administration using a device comprising a dry powder reservoir to which a stream of propellant may be directed to disperse the powder and/or using a self propelling solvent/powder dispensing container such as a device comprising the active ingredient dissolved and/or suspended in a low-boiling propellant in a sealed container. Such powders comprise particles wherein at least 98% of the particles by weight have a diameter greater than 0.5 nm and at least 95% of the particles by number have a diameter less than 7 nm. Alternatively, at least 95% of the particles by weight have a diameter greater than 1 nm and at least 90% of the particles by number have a diameter less than 6 nm. Dry powder compositions may include a solid fine powder diluent such as sugar and are conveniently provided in a unit dose form.

[00393] Low boiling propellants generally include liquid propellants having a boiling point of below 65 °F at atmospheric pressure. Generally the propellant may constitute 50% to 99.9% (w/w) of the composition, and active ingredient may constitute 0.1% to 20% (w/w) of the composition. A propellant may further comprise additional ingredients such as a liquid non-ionic and/or solid anionic surfactant and/or a solid diluent (which may have a particle size of the same order as particles comprising the active ingredient).

[00394] Pharmaceutical compositions formulated for pulmonary delivery may provide an active ingredient in the form of droplets of a solution and/or suspension. Such formulations may be prepared, packaged, and/or sold as aqueous and/or dilute alcoholic solutions and/or suspensions, optionally sterile, comprising active ingredient, and may conveniently be administered using any nebulization and/or atomization device. Such formulations may further comprise one or more additional ingredients including, but not limited to, a flavoring agent such as saccharin sodium, a volatile oil, a buffering agent, a surface active agent, and/or a preservative such as methylhydroxybenzoate. Droplets provided by this route of administration may have an average diameter in the range from about 0.1 nm to about 200 nm.

Intranasal, nasal and buccal Administration

[00395] Formulations described herein as being useful for pulmonary delivery are useful for intranasal delivery of a pharmaceutical composition. Another formulation suitable for intranasal administration is a coarse powder comprising the active ingredient and having an average particle from about 0.2 μm to 500 μm . Such a formulation is administered in the manner in which snuff is taken, *i.e.* by rapid inhalation through the nasal passage from a container of the powder held close to the nose.

[00396] Formulations suitable for nasal administration may, for example, comprise from about as little as 0.1% (w/w) and as much as 100% (w/w) of active ingredient, and may comprise one or more of the additional ingredients described herein. A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for buccal administration. Such formulations may, for example, be in the form of tablets and/or lozenges made using conventional methods, and may, for example, 0.1% to 20% (w/w) active ingredient, the balance comprising an orally dissolvable and/or degradable composition and, optionally, one or more of the additional ingredients described herein. Alternately, formulations suitable for buccal administration may comprise a powder and/or an aerosolized and/or atomized solution and/or suspension comprising active ingredient. Such powdered, aerosolized, and/or aerosolized formulations, when dispersed, may have an average particle and/or droplet size in the range from about 0.1 nm to about 200 nm, and may further comprise one or more of any additional ingredients described herein.

Ophthalmic Administration

[00397] A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for ophthalmic administration. Such formulations may, for example, be in the form of eye drops including, for example, a 0.1/1.0% (w/w) solution and/or suspension of the active ingredient in an aqueous or oily liquid excipient. Such drops may further comprise buffering agents, salts, and/or one or more other of any additional ingredients described herein. Other ophthalmically-administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form and/or in a liposomal preparation. Ear drops and/or eye drops are contemplated as being within the scope of this invention.

Payload Administration: Detectable Agents and Therapeutic Agents

[00398] The polynucleotides, primary constructs or mmRNA described herein can be used in a number of different scenarios in which delivery of a substance (the "payload") to a biological target is desired, for example delivery of detectable substances for detection of the target, or delivery of a therapeutic agent. Detection methods can include, but are not limited to, both imaging *in vitro* and *in vivo* imaging methods, *e.g.*, immunohistochemistry, bioluminescence imaging (BLI), Magnetic Resonance Imaging (MRI), positron emission tomography (PET), electron microscopy, X-ray computed tomography, Raman imaging, optical coherence tomography, absorption imaging, thermal imaging, fluorescence reflectance imaging, fluorescence microscopy, fluorescence molecular tomographic imaging, nuclear magnetic resonance imaging, X-ray imaging, ultrasound imaging, photoacoustic imaging, lab assays, or in any situation where tagging/staining/imaging is required.

[00399] The polynucleotides, primary constructs or mmRNA can be designed to include both a linker and a payload in any useful orientation. For example, a linker having two ends is used to attach one end to the payload and the other end to the nucleobase, such as at the C-7 or C-8 positions of the deaza-adenosine or deaza-guanosine or to the N-3 or C-5 positions of cytosine or uracil. The polynucleotide of the invention can include more than one payload (*e.g.*, a label and a transcription inhibitor), as well as a cleavable linker. In one embodiment, the modified nucleotide is a modified 7-deaza-adenosine triphosphate, where one end of a cleavable linker is attached to the C7 position of 7-deaza-adenine, the other end of the linker is attached to an inhibitor (*e.g.*, to the C5 position of the nucleobase on a cytidine), and a label (*e.g.*, Cy5) is attached to the

center of the linker (see, e.g., compound 1 of A*pCp C5 Parg Capless in Fig. 5 and columns 9 and 10 of U.S. Pat. No. 7,994,304, incorporated herein by reference). Upon incorporation of the modified 7-deaza-adenosine triphosphate to an encoding region, the resulting polynucleotide will have a cleavable linker attached to a label and an inhibitor (e.g., a polymerase inhibitor). Upon cleavage of the linker (e.g., with reductive conditions to reduce a linker having a cleavable disulfide moiety), the label and inhibitor are released. Additional linkers and payloads (e.g., therapeutic agents, detectable labels, and cell penetrating payloads) are described herein.

[00400] For example, the polynucleotides, primary constructs or mmRNA described herein can be used in induced pluripotent stem cells (iPS cells), which can directly track cells that are transfected compared to total cells in the cluster. In another example, a drug that may be attached to the polynucleotides, primary constructs or mmRNA via a linker and may be fluorescently labeled can be used to track the drug *in vivo*, e.g. intracellularly. Other examples include, but are not limited to, the use of a polynucleotide, primary construct or mmRNA in reversible drug delivery into cells.

[00401] The polynucleotides, primary constructs or mmRNA described herein can be used in intracellular targeting of a payload, e.g., detectable or therapeutic agent, to specific organelle. Exemplary intracellular targets can include, but are not limited to, the nuclear localization for advanced mRNA processing, or a nuclear localization sequence (NLS) linked to the mRNA containing an inhibitor.

[00402] In addition, the polynucleotides, primary constructs or mmRNA described herein can be used to deliver therapeutic agents to cells or tissues, e.g., in living animals. For example, the polynucleotides, primary constructs or mmRNA attached to the therapeutic agent through a linker can facilitate member permeation allowing the therapeutic agent to travel into a cell to reach an intracellular target.

[00403] In some embodiments, the payload may be a therapeutic agent such as a cytotoxin, radioactive ion, chemotherapeutic, or other therapeutic agent. A cytotoxin or cytotoxic agent includes any agent that may be detrimental to cells. Examples include, but are not limited to, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxyanthracenedione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, maytansinoids, e.g., maytansinol (see U.S. Pat. No. 5,208,020 incorporated herein in its entirety), rachelmycin (CC-1065, see U.S. Pat. Nos. 5,475,092, 5,585,499,

and 5,846,545, all of which are incorporated herein by reference), and analogs or homologs thereof. Radioactive ions include, but are not limited to iodine (*e.g.*, iodine 125 or iodine 131), strontium 89, phosphorous, palladium, cesium, iridium, phosphate, cobalt, yttrium 90, samarium 153, and praseodymium. Other therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thiotepa chlorambucil, rachelmycin (CC-1065), melphalan, carmustine (BSNU), lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (*e.g.*, vincristine, vinblastine, taxol and maytansinoids).

[00404] In some embodiments, the payload may be a detectable agent, such as various organic small molecules, inorganic compounds, nanoparticles, enzymes or enzyme substrates, fluorescent materials, luminescent materials (*e.g.*, luminol), bioluminescent materials (*e.g.*, luciferase, luciferin, and aequorin), chemiluminescent materials, radioactive materials (*e.g.*, ^{18}F , ^{67}Ga , $^{81\text{m}}\text{Kr}$, ^{82}Rb , ^{111}In , ^{123}I , ^{133}Xe , ^{201}Tl , ^{125}I , ^{35}S , ^{14}C , ^3H , or $^{99\text{m}}\text{Tc}$ (*e.g.*, as pertechnetate (technetate(VII), TcO_4^-)), and contrast agents (*e.g.*, gold (*e.g.*, gold nanoparticles), gadolinium (*e.g.*, chelated Gd), iron oxides (*e.g.*, superparamagnetic iron oxide (SPIO), monocrystalline iron oxide nanoparticles (MIONs), and ultrasmall superparamagnetic iron oxide (USPIO)), manganese chelates (*e.g.*, Mn-DPDP), barium sulfate, iodinated contrast media (iohexol), microbubbles, or perfluorocarbons). Such optically-detectable labels include for example, without limitation, 4-acetamido-4'-isothiocyanatostilbene-2,2'-disulfonic acid; acridine and derivatives (*e.g.*, acridine and acridine isothiocyanate); 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS); 4-amino-N-[3-vinylsulfonyl]phenyl]naphthalimide-3,5 disulfonate; N-(4-anilino-1-naphthyl)maleimide; anthranilamide; BODIPY; Brilliant Yellow; coumarin and derivatives (*e.g.*, coumarin, 7-amino-4-methylcoumarin (AMC, Coumarin 120), and 7-amino-4-trifluoromethylcoumarin (Coumarin 151)); cyanine dyes; cyanosine; 4',6-diaminidino-2-phenylindole (DAPI); 5' 5"-dibromopyrogallol-sulfonaphthalein (Bromopyrogallol Red); 7-diethylamino-3-(4'-isothiocyanatophenyl)-4-methylcoumarin; diethylenetriamine pentaacetate; 4,4'-diisothiocyanatodihydro-stilbene-2,2'-disulfonic acid; 4,4'-diisothiocyanatostilbene-2,2'-disulfonic acid; 5-[dimethylamino]-naphthalene-1-sulfonyl

chloride (DNS, dansylchloride); 4-dimethylaminophenylazophenyl-4'-isothiocyanate (DABITC); eosin and derivatives (e.g., eosin and eosin isothiocyanate); erythrosin and derivatives (e.g., erythrosin B and erythrosin isothiocyanate); ethidium; fluorescein and derivatives (e.g., 5-carboxyfluorescein (FAM), 5-(4,6-dichlorotriazin-2-yl)aminofluorescein (DTAF), 2',7'-dimethoxy-4'5'-dichloro-6-carboxyfluorescein, fluorescein, fluorescein isothiocyanate, X-rhodamine-5-(and-6)-isothiocyanate (QFITC or XRITC), and fluorescamine); 2-[2-[3-[[1,3-dihydro-1,1-dimethyl-3-(3-sulfopropyl)-2H-benz[e]indol-2-ylidene]ethylidene]-2-[4-(ethoxycarbonyl)-1-piperazinyl]-1-cyclopenten-1-yl]ethenyl]-1,1-dimethyl-3-(3-sulfopropyl)-1H-benz[e]indolium hydroxide, inner salt, compound with n,n-diethylethanamine(1:1) (IR144); 5-chloro-2-[2-[3-[(5-chloro-3-ethyl-2(3H)-benzothiazol-ylidene)ethylidene]-2-(diphenylamino)-1-cyclopenten-1-yl]ethenyl]-3-ethyl benzothiazolium perchlorate (IR140); Malachite Green isothiocyanate; 4-methylumbelliferone orthocresolphthalein; nitrotyrosine; pararosaniline; Phenol Red; B-phycoerythrin; o-phthaldialdehyde; pyrene and derivatives(e.g., pyrene, pyrene butyrate, and succinimidyl 1-pyrene); butyrate quantum dots; Reactive Red 4 (CIBACRONTM Brilliant Red 3B-A); rhodamine and derivatives (e.g., 6-carboxy-X-rhodamine (ROX), 6-carboxyrhodamine (R6G), lissamine rhodamine B sulfonyle chloride rhodamine (Rhod), rhodamine B, rhodamine 123, rhodamine X isothiocyanate, sulforhodamine B, sulforhodamine 101, sulfonyl chloride derivative of sulforhodamine 101 (Texas Red), N,N,N',N' tetramethyl-6-carboxyrhodamine (TAMRA) tetramethyl rhodamine, and tetramethyl rhodamine isothiocyanate (TRITC)); riboflavin; rosolic acid; terbium chelate derivatives; Cyanine-3 (Cy3); Cyanine-5 (Cy5); cyanine-5.5 (Cy5.5), Cyanine-7 (Cy7); IRD 700; IRD 800; Alexa 647; La Jolla Blue; phthalocyanine; and naphthalocyanine.

[00405] In some embodiments, the detectable agent may be a non-detectable pre-cursor that becomes detectable upon activation (e.g., fluorogenic tetrazine-fluorophore constructs (e.g., tetrazine-BODIPY FL, tetrazine-Oregon Green 488, or tetrazine-BODIPY TMR-X) or enzyme activatable fluorogenic agents (e.g., PROSENSE® (VisEn Medical))). *In vitro* assays in which the enzyme labeled compositions can be used include, but are not limited to, enzyme linked immunosorbent assays (ELISAs), immunoprecipitation assays, immunofluorescence, enzyme immunoassays (EIA), radioimmunoassays (RIA), and Western blot analysis.

Combinations

[00406] The polynucleotides, primary constructs or mmRNA may be used in combination with one or more other therapeutic, prophylactic, diagnostic, or imaging agents. By “in combination with,” it is not intended to imply that the agents must be administered at the same time and/or formulated for delivery together, although these methods of delivery are within the scope of the present disclosure. Compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. In some embodiments, the present disclosure encompasses the delivery of pharmaceutical, prophylactic, diagnostic, or imaging compositions in combination with agents that may improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body. As a non-limiting example, the polynucleotides, primary constructs and/or mmRNA may be used in combination with a pharmaceutical agent for the treatment of cancer or to control hyperproliferative cells. In U.S. Pat. No. 7,964,571, herein incorporated by reference in its entirety, a combination therapy for the treatment of solid primary or metastasized tumor is described using a pharmaceutical composition including a DNA plasmid encoding for interleukin-12 with a lipopolymer and also administering at least one anticancer agent or chemotherapeutic. Further, the polynucleotides, primary constructs and/or mmRNA of the present invention that encodes anti-proliferative molecules may be in a pharmaceutical composition with a lipopolymer (see e.g., U.S. Pub. No. 20110218231, herein incorporated by reference in its entirety, claiming a pharmaceutical composition comprising a DNA plasmid encoding an anti-proliferative molecule and a lipopolymer) which may be administered with at least one chemotherapeutic or anticancer agent.

Dosing

[00407] The present invention provides methods comprising administering polynucleotides, primary constructs and/or mmRNA and their encoded proteins or complexes in accordance with the invention to a subject in need thereof. nucleic acids, proteins or complexes, or pharmaceutical, imaging, diagnostic, or prophylactic compositions thereof, may be administered to a subject using any amount and any route of administration effective for preventing, treating, diagnosing, or imaging a disease, disorder, and/or condition (e.g., a disease, disorder, and/or condition relating to working

memory deficits). The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease, the particular composition, its mode of administration, its mode of activity, and the like. Compositions in accordance with the invention are typically formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of the compositions of the present invention may be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective, prophylactically effective, or appropriate imaging dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

[00408] In certain embodiments, compositions in accordance with the present invention may be administered at dosage levels sufficient to deliver from about 0.0001 mg/kg to about 100 mg/kg, from about 0.001 mg/kg to about 0.05 mg/kg, from about 0.005 mg/kg to about 0.05 mg/kg, from about 0.001 mg/kg to about 0.005 mg/kg, from about 0.05 mg/kg to about 0.5 mg/kg, from about 0.01 mg/kg to about 50 mg/kg, from about 0.1 mg/kg to about 40 mg/kg, from about 0.5 mg/kg to about 30 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.1 mg/kg to about 10 mg/kg, or from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic, diagnostic, prophylactic, or imaging effect. The desired dosage may be delivered three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, or every four weeks. In certain embodiments, the desired dosage may be delivered using multiple administrations (*e.g.*, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations).

[00409] According to the present invention, it has been discovered that administration of mmRNA in split-dose regimens produce higher levels of proteins in mammalian subjects. As used herein, a “split dose” is the division of single unit dose or total daily dose into two or more doses, *e.g.*, two or more administrations of the single unit dose. As used herein, a “single unit dose” is a dose of any therapeutic administered in one dose/at

one time/single route/single point of contact, i.e., single administration event. As used herein, a “total daily dose” is an amount given or prescribed in 24 hr period. It may be administered as a single unit dose. In one embodiment, the mmRNA of the present invention are administered to a subject in split doses. The mmRNA may be formulated in buffer only or in a formulation described herein.

Dosage Forms

[00410] A pharmaceutical composition described herein can be formulated into a dosage form described herein, such as a topical, intranasal, intratracheal, or injectable (e.g., intravenous, intraocular, intravitreal, intramuscular, intracardiac, intraperitoneal, subcutaneous).

Liquid dosage forms

[00411] Liquid dosage forms for parenteral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups, and/or elixirs. In addition to active ingredients, liquid dosage forms may comprise inert diluents commonly used in the art including, but not limited to, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. In certain embodiments for parenteral administration, compositions may be mixed with solubilizing agents such as CREMOPHOR[®], alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and/or combinations thereof.

Injectable

[00412] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art and may include suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations may be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally acceptable diluents and/or solvents, for example, a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed include, but are not limited to, are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono-

or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables.

[00413] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00414] In order to prolong the effect of an active ingredient, it may be desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the polynucleotide, primary construct or mmRNA then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered polynucleotide, primary construct or mmRNA may be accomplished by dissolving or suspending the polynucleotide, primary construct or mmRNA in an oil vehicle. Injectable depot forms are made by forming microcapsule matrices of the polynucleotide, primary construct or mmRNA in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of polynucleotide, primary construct or mmRNA to polymer and the nature of the particular polymer employed, the rate of polynucleotide, primary construct or mmRNA release can be controlled. Examples of other biodegradable polymers include, but are not limited to, poly(orthoesters) and poly(anhydrides). Depot injectable formulations may be prepared by entrapping the polynucleotide, primary construct or mmRNA in liposomes or microemulsions which are compatible with body tissues.

Pulmonary

[00415] Formulations described herein as being useful for pulmonary delivery may also be use for intranasal delivery of a pharmaceutical composition. Another formulation suitable for intranasal administration may be a coarse powder comprising the active ingredient and having an average particle from about 0.2 μm to 500 μm . Such a formulation may be administered in the manner in which snuff is taken, i.e. by rapid inhalation through the nasal passage from a container of the powder held close to the nose.

[00416] Formulations suitable for nasal administration may, for example, comprise from about as little as 0.1% (w/w) and as much as 100% (w/w) of active ingredient, and

may comprise one or more of the additional ingredients described herein. A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for buccal administration. Such formulations may, for example, be in the form of tablets and/or lozenges made using conventional methods, and may, for example, contain about 0.1% to 20% (w/w) active ingredient, where the balance may comprise an orally dissolvable and/or degradable composition and, optionally, one or more of the additional ingredients described herein. Alternately, formulations suitable for buccal administration may comprise a powder and/or an aerosolized and/or atomized solution and/or suspension comprising active ingredient. Such powdered, aerosolized, and/or aerosolized formulations, when dispersed, may have an average particle and/or droplet size in the range from about 0.1 nm to about 200 nm, and may further comprise one or more of any additional ingredients described herein.

[00417] General considerations in the formulation and/or manufacture of pharmaceutical agents may be found, for example, in Remington: The Science and Practice of Pharmacy 21st ed., Lippincott Williams & Wilkins, 2005 (incorporated herein by reference).

Coatings or Shells

[00418] Solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally comprise opacifying agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type may be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

Properties of Pharmaceutical Compositions

[00419] The pharmaceutical compositions described herein can be characterized by one or more of bioavailability, therapeutic window and/or volume of distribution.

Bioavailability

[00420] The polynucleotides, primary constructs or mmRNA, when formulated into a composition with a delivery agent as described herein, can exhibit an increase in bioavailability as compared to a composition lacking a delivery agent as described herein.

As used herein, the term “bioavailability” refers to the systemic availability of a given amount of polynucleotides, primary constructs or mmRNA administered to a mammal. Bioavailability can be assessed by measuring the area under the curve (AUC) or the maximum serum or plasma concentration (C_{\max}) of the unchanged form of a compound following administration of the compound to a mammal. AUC is a determination of the area under the curve plotting the serum or plasma concentration of a compound along the ordinate (Y-axis) against time along the abscissa (X-axis). Generally, the AUC for a particular compound can be calculated using methods known to those of ordinary skill in the art and as described in G. S. Banker, *Modern Pharmaceutics, Drugs and the Pharmaceutical Sciences*, v. 72, Marcel Dekker, New York, Inc., 1996, herein incorporated by reference.

[00421] The C_{\max} value is the maximum concentration of the compound achieved in the serum or plasma of a mammal following administration of the compound to the mammal. The C_{\max} value of a particular compound can be measured using methods known to those of ordinary skill in the art. The phrases “increasing bioavailability” or “improving the pharmacokinetics,” as used herein mean that the systemic availability of a first polynucleotide, primary construct or mmRNA, measured as AUC, C_{\max} , or C_{\min} in a mammal is greater, when co-administered with a delivery agent as described herein, than when such co-administration does not take place. In some embodiments, the bioavailability of the polynucleotide, primary construct or mmRNA can increase by at least about 2%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.

Therapeutic Window

[00422] The polynucleotides, primary constructs or mmRNA, when formulated into a composition with a delivery agent as described herein, can exhibit an increase in the therapeutic window of the administered polynucleotide, primary construct or mmRNA composition as compared to the therapeutic window of the administered polynucleotide, primary construct or mmRNA composition lacking a delivery agent as described herein. As used herein “therapeutic window” refers to the range of plasma concentrations, or the range of levels of therapeutically active substance at the site of action, with a high

probability of eliciting a therapeutic effect. In some embodiments, the therapeutic window of the polynucleotide, primary construct or mmRNA when co-administered with a delivery agent as described herein can increase by at least about 2%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or about 100%.

Volume of Distribution

[00423] The polynucleotides, primary constructs or mmRNA, when formulated into a composition with a delivery agent as described herein, can exhibit an improved volume of distribution (V_{dist}), e.g., reduced or targeted, relative to a composition lacking a delivery agent as described herein. The volume of distribution (V_{dist}) relates the amount of the drug in the body to the concentration of the drug in the blood or plasma. As used herein, the term “volume of distribution” refers to the fluid volume that would be required to contain the total amount of the drug in the body at the same concentration as in the blood or plasma: V_{dist} equals the amount of drug in the body/concentration of drug in blood or plasma. For example, for a 10 mg dose and a plasma concentration of 10 mg/L, the volume of distribution would be 1 liter. The volume of distribution reflects the extent to which the drug is present in the extravascular tissue. A large volume of distribution reflects the tendency of a compound to bind to the tissue components compared with plasma protein binding. In a clinical setting, V_{dist} can be used to determine a loading dose to achieve a steady state concentration. In some embodiments, the volume of distribution of the polynucleotide, primary construct or mmRNA when co-administered with a delivery agent as described herein can decrease at least about 2%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%.

Biological Effect

[00424] In one embodiment, the biological effect of the modified mRNA delivered to the animals may be categorized by analyzing the protein expression in the animals. The reprogrammed protein expression may be determined from analyzing a biological sample collected from a mammal administered the modified mRNA of the present invention. In

one embodiment, the expression protein encoded by the modified mRNA administered to the mammal of at least 50 pg/ml may be preferred. For example, a protein expression of 50-200 pg/ml for the protein encoded by the modified mRNA delivered to the mammal may be seen as a therapeutically effective amount of protein in the mammal.

Quantification

[00425] In one embodiment, the polynucleotides, primary constructs or mmRNA of the present invention may be quantified in exosomes derived from one or more bodily fluid. As used herein “bodily fluids” include peripheral blood, serum, plasma, ascites, urine, cerebrospinal fluid (CSF), sputum, saliva, bone marrow, synovial fluid, aqueous humor, amniotic fluid, cerumen, breast milk, bronchoalveolar lavage fluid, semen, prostatic fluid, cowper's fluid or pre-ejaculatory fluid, sweat, fecal matter, hair, tears, cyst fluid, pleural and peritoneal fluid, pericardial fluid, lymph, chyme, chyle, bile, interstitial fluid, menses, pus, sebum, vomit, vaginal secretions, mucosal secretion, stool water, pancreatic juice, lavage fluids from sinus cavities, bronchopulmonary aspirates, blastocyl cavity fluid, and umbilical cord blood. Alternatively, exosomes may be retrieved from an organ selected from the group consisting of lung, heart, pancreas, stomach, intestine, bladder, kidney, ovary, testis, skin, colon, breast, prostate, brain, esophagus, liver, and placenta.

[00426] In the quantification method, a sample of not more than 2mL is obtained from the subject and the exosomes isolated by size exclusion chromatography, density gradient centrifugation, differential centrifugation, nanomembrane ultrafiltration, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof. In the analysis, the level or concentration of a polynucleotide, primary construct or mmRNA may be an expression level, presence, absence, truncation or alteration of the administered construct. It is advantageous to correlate the level with one or more clinical phenotypes or with an assay for a human disease biomarker. The assay may be performed using construct specific probes, cytometry, qRT-PCR, real-time PCR, PCR, flow cytometry, electrophoresis, mass spectrometry, or combinations thereof while the exosomes may be isolated using immunohistochemical methods such as enzyme linked immunosorbent assay (ELISA) methods. Exosomes may also be isolated by size exclusion chromatography, density gradient centrifugation, differential centrifugation, nanomembrane ultrafiltration, immunoabsorbent capture, affinity purification, microfluidic separation, or combinations thereof.

[00427] These methods afford the investigator the ability to monitor, in real time, the level of polynucleotides, primary constructs or mmRNA remaining or delivered. This is possible because the polynucleotides, primary constructs or mmRNA of the present invention differ from the endogenous forms due to the structural or chemical modifications.

[00428]

Detection of Modified Polynucleotides by Mass Spectrometry

[00429] Mass spectrometry (MS) is an analytical technique that can provide structural and molecular mass/concentration information on molecules after their conversion to ions. The molecules are first ionized to acquire positive or negative charges and then they travel through the mass analyzer to arrive at different areas of the detector according to their mass/charge (m/z) ratio.

[00430] Mass spectrometry is performed using a mass spectrometer which includes an ion source for ionizing the fractionated sample and creating charged molecules for further analysis. For example ionization of the sample may be performed by electrospray ionization (ESI), atmospheric pressure chemical ionization (APCI), photoionization, electron ionization, fast atom bombardment (FAB)/liquid secondary ionization (LSIMS), matrix assisted laser desorption/ionization (MALDI), field ionization, field desorption, thermospray/plasmaspray ionization, and particle beam ionization. The skilled artisan will understand that the choice of ionization method can be determined based on the analyte to be measured, type of sample, the type of detector, the choice of positive versus negative mode, etc.

[00431] After the sample has been ionized, the positively charged or negatively charged ions thereby created may be analyzed to determine a mass-to-charge ratio (i.e., m/z). Suitable analyzers for determining mass-to-charge ratios include quadropole analyzers, ion traps analyzers, and time-of-flight analyzers. The ions may be detected using several detection modes. For example, selected ions may be detected (i.e., using a selective ion monitoring mode (SIM)), or alternatively, ions may be detected using a scanning mode, e.g., multiple reaction monitoring (MRM) or selected reaction monitoring (SRM).

[00432] Liquid chromatography-multiple reaction monitoring (LC-MS/MRM) coupled with stable isotope labeled dilution of peptide standards has been shown to be an effective method for protein verification (e.g., Keshishian et al., Mol Cell Proteomics 2009 8: 2339-2349; Kuhn et al., Clin Chem 2009 55:1108-1117; Lopez et al., Clin Chem 2010

56:281-290). Unlike untargeted mass spectrometry frequently used in biomarker discovery studies, targeted MS methods are peptide sequence-based modes of MS that focus the full analytical capacity of the instrument on tens to hundreds of selected peptides in a complex mixture. By restricting detection and fragmentation to only those peptides derived from proteins of interest, sensitivity and reproducibility are improved dramatically compared to discovery-mode MS methods. This method of mass spectrometry-based multiple reaction monitoring (MRM) quantitation of proteins can dramatically impact the discovery and quantitation of biomarkers via rapid, targeted, multiplexed protein expression profiling of clinical samples.

[00433] In one embodiment, a biological sample which may contain at least one protein encoded by at least one modified mRNA of the present invention may be analyzed by the method of MRM-MS. The quantification of the biological sample may further include, but is not limited to, isotopically labeled peptides or proteins as internal standards.

[00434] According to the present invention, the biological sample, once obtained from the subject, may be subjected to enzyme digestion. As used herein, the term “digest” means to break apart into shorter peptides. As used herein, the phrase “treating a sample to digest proteins” means manipulating a sample in such a way as to break down proteins in a sample. These enzymes include, but are not limited to, trypsin, endoproteinase Glu-C and chymotrypsin. In one embodiment, a biological sample which may contain at least one protein encoded by at least one modified mRNA of the present invention may be digested using enzymes.

[00435] In one embodiment, a biological sample which may contain protein encoded by modified mRNA of the present invention may be analyzed for protein using electrospray ionization. Electrospray ionization (ESI) mass spectrometry (ESIMS) uses electrical energy to aid in the transfer of ions from the solution to the gaseous phase before they are analyzed by mass spectrometry. Samples may be analyzed using methods known in the art (e.g., Ho et al., *Clin Biochem Rev.* 2003 24(1):3-12). The ionic species contained in solution may be transferred into the gas phase by dispersing a fine spray of charge droplets, evaporating the solvent and ejecting the ions from the charged droplets to generate a mist of highly charged droplets. The mist of highly charged droplets may be analyzed using at least 1, at least 2, at least 3 or at least 4 mass analyzers such as, but not limited to, a quadrupole mass analyzer. Further, the mass spectrometry method may include a purification step. As a non-limiting example, the first quadrupole may be set to select a single m/z ratio so it may filter out other molecular ions having a different m/z

ratio which may eliminate complicated and time-consuming sample purification procedures prior to MS analysis.

[00436] In one embodiment, a biological sample which may contain protein encoded by modified mRNA of the present invention may be analyzed for protein in a tandem ESIMS system (e.g., MS/MS). As non-limiting examples, the droplets may be analyzed using a product scan (or daughter scan) a precursor scan (parent scan) a neutral loss or a multiple reaction monitoring.

[00437] In one embodiment, a biological sample which may contain protein encoded by modified mRNA of the present invention may be analyzed using matrix-assisted laser desorption/ionization (MALDI) mass spectrometry (MALDIMS). MALDI provides for the nondestructive vaporization and ionization of both large and small molecules, such as proteins. In MALDI analysis, the analyte is first co-crystallized with a large molar excess of a matrix compound, which may also include, but is not limited to, an ultraviolet absorbing weak organic acid. Non-limiting examples of matrices used in MALDI are α -cyano-4-hydroxycinnamic acid, 3,5-dimethoxy-4-hydroxycinnamic acid and 2,5-dihydroxybenzoic acid. Laser radiation of the analyte-matrix mixture may result in the vaporization of the matrix and the analyte. The laser induced desorption provides high ion yields of the intact analyte and allows for measurement of compounds with high accuracy. Samples may be analyzed using methods known in the art (e.g., Lewis, Wei and Siuzdak, Encyclopedia of Analytical Chemistry 2000:5880-5894). As non-limiting examples, mass analyzers used in the MALDI analysis may include a linear time-of-flight (TOF), a TOF reflectron or a Fourier transform mass analyzer.

[00438] In one embodiment, the analyte-matrix mixture may be formed using the dried-droplet method. A biologic sample is mixed with a matrix to create a saturated matrix solution where the matrix-to-sample ratio is approximately 5000:1. An aliquot (approximately 0.5-2.0 uL) of the saturated matrix solution is then allowed to dry to form the analyte-matrix mixture.

[00439] In one embodiment, the analyte-matrix mixture may be formed using the thin-layer method. A matrix homogeneous film is first formed and then the sample is then applied and may be absorbed by the matrix to form the analyte-matrix mixture.

[00440] In one embodiment, the analyte-matrix mixture may be formed using the thick-layer method. A matrix homogeneous film is formed with a nitro-cellulose matrix additive. Once the uniform nitro-cellulose matrix layer is obtained the sample is applied and absorbed into the matrix to form the analyte-matrix mixture.

[00441] In one embodiment, the analyte-matrix mixture may be formed using the sandwich method. A thin layer of matrix crystals is prepared as in the thin-layer method followed by the addition of droplets of aqueous trifluoroacetic acid, the sample and matrix. The sample is then absorbed into the matrix to form the analyte-matrix mixture.

VI. Uses of synthetic polynucleotides and/or synthetic sgRNAs

[00442] The synthetic polynucleotides and/or synthetic sgRNAs of the present invention may be used to alter the phenotype of cells. The polynucleotides, primary constructs and mmRNA of the invention may encode peptides, polypeptides or multiple proteins to produce polypeptides of interest. The polypeptides of interest may be used in therapeutics and/or clinical and research settings. As a non-limiting example, the polypeptides of interest may include reprogramming factors, differentiation factors and de-differentiation factors.

Therapeutic Agents

[00443] The polynucleotides, primary constructs or mmRNA of the present invention, such as modified nucleic acids and modified RNAs, and the proteins translated from them described herein can be used as therapeutic or prophylactic agents. They are provided for use in medicine, therapy and preventative treatments. For example, a polynucleotide, primary construct or mmRNA described herein (*e.g.*, a modified mRNA encoding a CRISPR-related polypeptide or protein) can be administered to a subject, wherein the polynucleotide, primary construct or mmRNA is translated *in vivo* to produce a therapeutic or prophylactic polypeptide in the subject. Likewise, and optionally in combination with the above, a polynucleotide, primary construct or mmRNA described herein (*e.g.*, a modified sgRNA or plurality of same) can be administered to a subject, wherein the polynucleotide, primary construct or mmRNA guides gene-editing by a CRISPR-related polypeptide or protein.) Provided are compositions, methods, kits, and reagents for diagnosis, treatment or prevention of a disease or condition in humans and other mammals. The active therapeutic agents of the invention include polynucleotides, primary constructs or mmRNA, cells containing the polynucleotides, primary constructs or mmRNA or polypeptides translated from the polynucleotides, primary constructs or mmRNA.

[00444] In certain embodiments, provided herein are combination therapeutics containing one or more polynucleotide, primary construct or mmRNA containing translatable regions that encode for a protein or proteins. In certain embodiments,

provided herein are combination therapeutics containing one or more polynucleotide, primary construct or mmRNA containing non-translatable regions, for example, regions complementary to target gene sequences, *e.g.*, sgRNAs.

[00445] Provided herein are methods of inducing translation of a recombinant polypeptide in a cell population using the polynucleotide, primary construct or mmRNA described herein. Such translation can be *in vivo*, *ex vivo*, *in culture*, or *in vitro*. The cell population is contacted with an effective amount of a composition containing a nucleic acid that has at least one nucleoside modification, and a translatable region encoding the recombinant polypeptide. The cell population is optionally also contacted with an effective amount of a composition containing a nucleic acid that has at least one nucleoside modification, and has complementarity to a target gene sequence, *e.g.*, a sgRNA. The population is contacted under conditions such that the nucleic acid is localized into one or more cells of the cell population and the recombinant polypeptide is translated in the cell from the nucleic acid and or the sgRNA (or plurality thereof) is also localized into one or more cells of the cell population.

[00446] An “effective amount” of the composition is provided based, at least in part, on the target tissue, target cell type, means of administration, physical characteristics of the nucleic acid (*e.g.*, size, and extent of modified nucleosides), and other determinants. In general, an effective amount of the composition provides efficient protein production in the cell, preferably more efficient than a composition containing a corresponding unmodified nucleic acid. Increased efficiency may be demonstrated by increased cell transfection (*i.e.*, the percentage of cells transfected with the nucleic acid), increased protein translation from the nucleic acid, decreased nucleic acid degradation (as demonstrated, *e.g.*, by increased duration of protein translation from a modified nucleic acid), or reduced innate immune response of the host cell.

[00447] Aspects of the invention are directed to methods of inducing *in vivo* translation of a recombinant polypeptide in a mammalian subject in need thereof, *e.g.*, a recombinant CRISPR-related protein. Further aspects of the invention are directed to methods of inducing gene editing *in vivo* in a mammalian subject in need thereof. Therein, an effective amount of a composition containing a nucleic acid that has at least one structural or chemical modification and a translatable region encoding the recombinant polypeptide, optionally in combination with an effective amount of a composition containing one or more nucleic acids having at least one structural or chemical modification and/or having sequence complementarity to a target gene, *e.g.*, one or more sgRNAs, is administered to

the subject using the delivery methods described herein. The nucleic acid is provided in an amount and under other conditions such that the nucleic acid is localized into a cell of the subject and (1) the recombinant polypeptide is translated in the cell from the nucleic acid and/or (2) the sgRNA(s) bind complementary sequences within target genes. The cell in which a nucleic acid of the invention is localized, or the tissue in which the cell is present, may be targeted with one or more than one rounds of nucleic acid administration.

[00448] In certain embodiments, the administered polynucleotide, primary construct or mmRNA directs production of one or more recombinant polypeptides and/or regulatory RNA, *e.g.*, sgRNA, that provides a functional activity which is substantially absent in the cell, tissue or organism in which the recombinant polypeptide is translated, *e.g.*, a gene editing activity. In certain embodiments, an administered polynucleotide, primary construct or mmRNA directs production of one or more recombinant polypeptides and/or regulatory RNA, *e.g.*, sgRNA, involved in gene editing, and is administered in combination with one or more additional polynucleotide, primary construct or mmRNA that directs production of one or more additional recombinant polypeptides having therapeutic utility, *e.g.*, providing a functional activity which is substantially absent in the cell, tissue or organism in which the recombinant polypeptide is translated. For example, the missing functional activity may be enzymatic, structural, or gene regulatory in nature. In related embodiments, the administered polynucleotide, primary construct or mmRNA directs production of one or more recombinant polypeptides that increases (*e.g.*, synergistically) a functional activity which is present but substantially deficient in the cell in which the recombinant polypeptide is translated.

[00449] In other embodiments, the administered polynucleotide, primary construct or mmRNA directs production of one or more recombinant polypeptides that replace a polypeptide (or multiple polypeptides) that is substantially absent in the cell in which the recombinant polypeptide is translated. Such absence may be due to genetic mutation of the encoding gene or regulatory pathway thereof. In some embodiments, the recombinant polypeptide increases the level of an endogenous protein in the cell to a desirable level; such an increase may bring the level of the endogenous protein from a subnormal level to a normal level or from a normal level to a super-normal level.

[00450] Alternatively, the recombinant polypeptide functions to antagonize the activity of an endogenous protein present in, on the surface of, or secreted from the cell. Usually, the activity of the endogenous protein is deleterious to the subject; for example, due to mutation of the endogenous protein resulting in altered activity or localization.

Additionally, the recombinant polypeptide antagonizes, directly or indirectly, the activity of a biological moiety present in, on the surface of, or secreted from the cell. Examples of antagonized biological moieties include lipids (e.g., cholesterol), a lipoprotein (e.g., low density lipoprotein), a nucleic acid, a carbohydrate, a metabolite (e.g. bilirubin), a protein toxin such as shiga and tetanus toxins, or a small molecule toxin such as botulinum, cholera, and diphtheria toxins. Additionally, the antagonized biological molecule may be an endogenous protein that exhibits an undesirable activity, such as a cytotoxic or cytostatic activity.

[00451] Without being bound in theory, it is proposed that the CRISPR gene-editing process can reverse disease symptoms *in vivo*. As such, this methodology has potential for treating many genetic disorders.

In certain embodiments, a mmRNA encoding a CRISPR-related polypeptide or protein is administered in combination with one or more sgRNAs, optionally in combination with a corrected DNA template. The DNA template includes a correct sequence of corresponding to a mutated gene associated with a disease. Exemplary DNA templates correspond to single-mutation disease causing genes. A cell into which the requisite components are introduced repairs the damaged gene using the CRISPER-related protein. In doing so, it copies from the template, introducing new genetic material into the genome.

[00452] The data presented herein demonstrate effective liver delivery of components of the invention. As such, the methodology described herein is particularly suited to treatment of metabolic disease, *e.g.*, metabolic disease caused by inborn genetic errors, *e.g.*, ornithine transcarbamylase deficiency-related diseases. The methods of the invention can also serve to attenuate disease progression. It is also proposed that intramuscular delivery, *e.g.*, via multiple muscle injection, can serve to treat muscle diseases, for example, muscular dystrophy, by inclusion of appropriate corrected DNA templates. In yet other embodiments, *ex vivo* approaches are envisioned involving, for example, administration of agents to hematopoietic cells *ex vivo* (*e.g.*, via electroporation), followed optionally by selection of cells expressing the proper components, followed by administration of treated cells, where recolonization can subsequently occur.

[00453] The recombinant proteins, mmRNAs encoding same, and/or sgRNAs, described herein may be engineered for localization within the cell, potentially within a specific compartment such as the nucleus, or are engineered for secretion from the cell or translocation to the plasma membrane of the cell. In exemplary embodiments, the

recombinant proteins, mmRNAs encoding same, and/or sgRNAs, are engineered for nuclear localization.

[00454] Other aspects of the present disclosure relate to transplantation of cells containing polynucleotide, primary construct, or mmRNA to a mammalian subject. Administration of cells to mammalian subjects is known to those of ordinary skill in the art, and include, but is not limited to, local implantation (e.g., topical or subcutaneous administration), organ delivery or systemic injection (e.g., intravenous injection or inhalation), and the formulation of cells in pharmaceutically acceptable carrier. Such compositions containing polynucleotide, primary construct, or mmRNA can be formulated for administration intramuscularly, transarterially, intraperitoneally, intravenously, intranasally, subcutaneously, endoscopically, transdermally, or intrathecally. In some embodiments, the composition may be formulated for extended release.

[00455] The subject to whom the therapeutic agent may be administered suffers from or may be at risk of developing a disease, disorder, or deleterious condition. Provided are methods of identifying, diagnosing, and classifying subjects on these bases, which may include clinical diagnosis, biomarker levels, genome-wide association studies (GWAS), and other methods known in the art.

VII. Kits and Devices

[00456] The invention provides a variety of kits for conveniently and/or effectively carrying out methods of the present invention. Typically kits will comprise sufficient amounts and/or numbers of components to allow a user to perform multiple treatments of a subject(s) and/or to perform multiple experiments, and contact cells and/or a population of cells at least once.

[00457] In one aspect, the present invention provides kits comprising the molecules (polynucleotides, primary constructs or mmRNA) of the invention. In one embodiment, the kit comprises one or more functional antibodies or function fragments thereof.

[00458] Kits and devices useful in combination with the polynucleotides, primary constructs or mmRNA) of the invention include those disclosed in co-pending U.S. Provisional Patent Application No 61/737,130 filed December 14, 2012, the contents of which are incorporated herein by reference in their entirety.

VIII. Definitions

[00459] At various places in the present specification, substituents of compounds of the present disclosure are disclosed in groups or in ranges. It is specifically intended that the present disclosure include each and every individual subcombination of the members of such groups and ranges. For example, the term “C₁₋₆ alkyl” is specifically intended to individually disclose methyl, ethyl, C₃ alkyl, C₄ alkyl, C₅ alkyl, and C₆ alkyl.

[00460] *About*: As used herein, the term “about” means +/- 10% of the recited value.

[00461] *Administered in combination*: As used herein, the term “administered in combination” or “combined administration” means that two or more agents are administered to a subject at the same time or within an interval such that there may be an overlap of an effect of each agent on the patient. In some embodiments, they are administered within about 60, 30, 15, 10, 5, or 1 minute of one another. In some embodiments, the administrations of the agents are spaced sufficiently closely together such that a combinatorial (*e.g.*, a synergistic) effect is achieved.

[00462] *Animal*: As used herein, the term “animal” refers to any member of the animal kingdom. In some embodiments, “animal” refers to humans at any stage of development. In some embodiments, “animal” refers to non-human animals at any stage of development. In certain embodiments, the non-human animal is a mammal (*e.g.*, a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, and worms. In some embodiments, the animal is a transgenic animal, genetically-engineered animal, or a clone.

[00463] *Approximately*: As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[00464] *Associated with*: As used herein, the terms “associated with,” “conjugated,” “linked,” “attached,” and “tethered,” when used with respect to two or more moieties, means that the moieties are physically associated or connected with one another, either directly or via one or more additional moieties that serves as a linking agent, to form a

structure that is sufficiently stable so that the moieties remain physically associated under the conditions in which the structure is used, *e.g.*, physiological conditions. An “association” need not be strictly through direct covalent chemical bonding. It may also suggest ionic or hydrogen bonding or a hybridization based connectivity sufficiently stable such that the “associated” entities remain physically associated.

[00465] *Bifunctional*: As used herein, the term “bifunctional” refers to any substance, molecule or moiety which is capable of or maintains at least two functions. The functions may affect the same outcome or a different outcome. The structure that produces the function may be the same or different. For example, bifunctional modified RNAs of the present invention may encode a cytotoxic peptide (a first function) while those nucleosides which comprise the encoding RNA are, in and of themselves, cytotoxic (second function). In this example, delivery of the bifunctional modified RNA to a cancer cell would produce not only a peptide or protein molecule which may ameliorate or treat the cancer but would also deliver a cytotoxic payload of nucleosides to the cell should degradation, instead of translation of the modified RNA, occur.

[00466] *Biocompatible*: As used herein, the term “biocompatible” means compatible with living cells, tissues, organs or systems posing little to no risk of injury, toxicity or rejection by the immune system.

[00467] *Biodegradable*: As used herein, the term “biodegradable” means capable of being broken down into innocuous products by the action of living things.

[00468] *Biologically active*: As used herein, the phrase “biologically active” refers to a characteristic of any substance that has activity in a biological system and/or organism. For instance, a substance that, when administered to an organism, has a biological effect on that organism, is considered to be biologically active. In particular embodiments, a polynucleotide, primary construct or mmRNA of the present invention may be considered biologically active if even a portion of the polynucleotide, primary construct or mmRNA is biologically active or mimics an activity considered biologically relevant.

[00469] *Cancer stem cells*: As used herein, “cancer stem cells” are cells that can undergo self-renewal and/or abnormal proliferation and differentiation to form a tumor.

[00470] *Chemical terms*: Chemical terms not otherwise defined herein, will conform to the chemical term definitions provided in co-pending U.S. Provisional Patent Application No 61/737,130 filed December 14, 2012, the contents of which are incorporated herein by reference in their entirety.

[00471] The term “diastereomer,” as used herein means stereoisomers that are not mirror images of one another and are non-superimposable on one another.

[00472] The term “effective amount” of an agent, as used herein, is that amount sufficient to effect beneficial or desired results, for example, clinical results, and, as such, an “effective amount” depends upon the context in which it is being applied. For example, in the context of administering an agent that treats cancer, an effective amount of an agent is, for example, an amount sufficient to achieve treatment, as defined herein, of cancer, as compared to the response obtained without administration of the agent.

[00473] The term “enantiomer,” as used herein, means each individual optically active form of a compound of the invention, having an optical purity or enantiomeric excess (as determined by methods standard in the art) of at least 80% (i.e., at least 90% of one enantiomer and at most 10% of the other enantiomer), preferably at least 90% and more preferably at least 98%.

[00474] The term “isomer,” as used herein, means any tautomer, stereoisomer, enantiomer, or diastereomer of any compound of the invention. It is recognized that the compounds of the invention can have one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (i.e., geometric E/Z isomers) or diastereomers (e.g., enantiomers (i.e., (+) or (-)) or cis/trans isomers). According to the invention, the chemical structures depicted herein, and therefore the compounds of the invention, encompass all of the corresponding stereoisomers, that is, both the stereomerically pure form (e.g., geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric and stereoisomeric mixtures, e.g., racemates. Enantiomeric and stereoisomeric mixtures of compounds of the invention can typically be resolved into their component enantiomers or stereoisomers by well-known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and stereoisomers can also be obtained from stereomerically or enantiomerically pure intermediates, reagents, and catalysts by well-known asymmetric synthetic methods.

[00475] The term “stereoisomer,” as used herein, refers to all possible different isomeric as well as conformational forms which a compound may possess (e.g., a compound of any formula described herein), in particular all possible stereochemically and conformationally isomeric forms, all diastereomers, enantiomers and/or conformers of the basic molecular structure. Some compounds of the present invention may exist in

different tautomeric forms, all of the latter being included within the scope of the present invention.

[00476] *Compound:* As used herein, the term “compound,” is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted.

[00477] The compounds described herein can be asymmetric (*e.g.*, having one or more stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present disclosure that contain asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically active starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present disclosure. Cis and trans geometric isomers of the compounds of the present disclosure are described and may be isolated as a mixture of isomers or as separated isomeric forms.

[00478] Compounds of the present disclosure also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond and the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Examples prototropic tautomers include ketone – enol pairs, amide – imidic acid pairs, lactam – lactim pairs, amide – imidic acid pairs, enamine – imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, such as, 1H- and 3H-imidazole, 1H-, 2H- and 4H- 1,2,4-triazole, 1H- and 2H- isoindole, and 1H- and 2H-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution.

[00479] Compounds of the present disclosure also include all of the isotopes of the atoms occurring in the intermediate or final compounds. “Isotopes” refers to atoms having the same atomic number but different mass numbers resulting from a different number of neutrons in the nuclei. For example, isotopes of hydrogen include tritium and deuterium.

[00480] The compounds and salts of the present disclosure can be prepared in combination with solvent or water molecules to form solvates and hydrates by routine methods.

[00481] *Committed*: As used herein, the term “committed” means, when referring to a cell, when the cell is far enough into the differentiation pathway where, under normal circumstances, it will continue to differentiate into a specific cell type or subset of cell type instead of into a different cell type or reverting to a lesser differentiated cell type.

[00482] *Conserved*: As used herein, the term “conserved” refers to nucleotides or amino acid residues of a polynucleotide sequence or polypeptide sequence, respectively, that are those that occur unaltered in the same position of two or more sequences being compared. Nucleotides or amino acids that are relatively conserved are those that are conserved amongst more related sequences than nucleotides or amino acids appearing elsewhere in the sequences.

[00483] In some embodiments, two or more sequences are said to be “completely conserved” if they are 100% identical to one another. In some embodiments, two or more sequences are said to be “highly conserved” if they are at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some embodiments, two or more sequences are said to be “highly conserved” if they are about 70% identical, about 80% identical, about 90% identical, about 95%, about 98%, or about 99% identical to one another. In some embodiments, two or more sequences are said to be “conserved” if they are at least 30% identical, at least 40% identical, at least 50% identical, at least 60% identical, at least 70% identical, at least 80% identical, at least 90% identical, or at least 95% identical to one another. In some embodiments, two or more sequences are said to be “conserved” if they are about 30% identical, about 40% identical, about 50% identical, about 60% identical, about 70% identical, about 80% identical, about 90% identical, about 95% identical, about 98% identical, or about 99% identical to one another. Conservation of sequence may apply to the entire length of an oligonucleotide or polypeptide or may apply to a portion, region or feature thereof.

[00484] *Controlled Release*: As used herein, the term “controlled release” refers to a pharmaceutical composition or compound release profile that conforms to a particular pattern of release to effect a therapeutic outcome.

[00485] *Cyclic or Cyclized*: As used herein, the term “cyclic” refers to the presence of a continuous loop. Cyclic molecules need not be circular, only joined to form an unbroken chain of subunits. Cyclic molecules such as the engineered RNA or mRNA of the present invention may be single units or multimers or comprise one or more components of a complex or higher order structure.

[00486] *Cytostatic*: As used herein, “cytostatic” refers to inhibiting, reducing, suppressing the growth, division, or multiplication of a cell (*e.g.*, a mammalian cell (*e.g.*, a human cell)), bacterium, virus, fungus, protozoan, parasite, prion, or a combination thereof.

[00487] *Cytotoxic*: As used herein, “cytotoxic” refers to killing or causing injurious, toxic, or deadly effect on a cell (*e.g.*, a mammalian cell (*e.g.*, a human cell)), bacterium, virus, fungus, protozoan, parasite, prion, or a combination thereof.

[00488] *Delivery*: As used herein, “delivery” refers to the act or manner of delivering a compound, substance, entity, moiety, cargo or payload.

[00489] *Delivery Agent*: As used herein, “delivery agent” refers to any substance which facilitates, at least in part, the *in vivo* delivery of a polynucleotide, primary construct or mmRNA to targeted cells.

[00490] *Destabilized*: As used herein, the term “destable,” “destabilize,” or “destabilizing region” means a region or molecule that is less stable than a starting, wild-type or native form of the same region or molecule.

[00491] *Detectable label*: As used herein, “detectable label” refers to one or more markers, signals, or moieties which are attached, incorporated or associated with another entity that is readily detected by methods known in the art including radiography, fluorescence, chemiluminescence, enzymatic activity, absorbance and the like. Detectable labels include radioisotopes, fluorophores, chromophores, enzymes, dyes, metal ions, ligands such as biotin, avidin, streptavidin and haptens, quantum dots, and the like. Detectable labels may be located at any position in the peptides or proteins disclosed herein. They may be within the amino acids, the peptides, or proteins, or located at the N- or C- termini.

[00492] *Developmental Potential*: As used herein, “developmental potential” or “developmental potency” refers to the total of all developmental cell fates or cell types that can be achieved by a cell upon differentiation.

[00493] *Developmental Potential Altering Factor*: As used herein, “developmental potential altering factor” refers to a protein or RNA which can alter the developmental potential of a cell.

[00494] *Digest*: As used herein, the term “digest” means to break apart into smaller pieces or components. When referring to polypeptides or proteins, digestion results in the production of peptides.

[00495] *Differentiated cell*: As used herein, the term “differentiated cell” refers to any somatic cell that is not, in its native form, pluripotent. Differentiated cell also encompasses cells that are partially differentiated.

[00496] *Differentiation*: As used herein, the term “differentiation factor” refers to a developmental potential altering factor such as a protein, RNA or small molecule that can induce a cell to differentiate to a desired cell-type.

[00497] *Differentiate*: As used herein, “differentiate” refers to the process where an uncommitted or less committed cell acquires the features of a committed cell.

[00498] *Disease*: As used herein, the term “disease” refers to an abnormal condition affecting the body of an organism often showing specific bodily symptoms.

[00499] *Disorder*: As used herein, the term “disorder,” refers to a disruption of or an interference with normal functions or established systems of the body.

[00500] *Distal*: As used herein, the term “distal” means situated away from the center or away from a point or region of interest.

[00501] *Dose splitting factor (DSF)*-ratio of PUD of dose split treatment divided by PUD of total daily dose or single unit dose. The value is derived from comparison of dosing regimens groups.

[00502] *Embryonic stem cell*: As used herein, the term “embryonic stem cell” refers to naturally occurring pluripotent stem cells of the inner cell mass of the embryonic blastocyst.

[00503] *Encapsulate*: As used herein, the term “encapsulate” means to enclose, surround or encase.

[00504] *Encoded protein cleavage signal*: As used herein, “encoded protein cleavage signal” refers to the nucleotide sequence which encodes a protein cleavage signal.

[00505] *Engineered*: As used herein, embodiments of the invention are “engineered” when they are designed to have a feature or property, whether structural or chemical, that varies from a starting point, wild type or native molecule.

[00506] *Exosome*: As used herein, “exosome” is a vesicle secreted by mammalian cells or a complex involved in RNA degradation.

[00507] *Expression*: As used herein, “expression” of a nucleic acid sequence refers to one or more of the following events: (1) production of an RNA template from a DNA sequence (*e.g.*, by transcription); (2) processing of an RNA transcript (*e.g.*, by splicing, editing, 5' cap formation, and/or 3' end processing); (3) translation of an RNA into a polypeptide or protein; and (4) post-translational modification of a polypeptide or protein.

[00508] *Feature*: As used herein, a “feature” refers to a characteristic, a property, or a distinctive element.

[00509] *Formulation*: As used herein, a “formulation” includes at least a polynucleotide, primary construct or mmRNA and a delivery agent.

[00510] *Fragment*: A “fragment,” as used herein, refers to a portion. For example, fragments of proteins may comprise polypeptides obtained by digesting full-length protein isolated from cultured cells.

[00511] *Functional*: As used herein, a “functional” biological molecule is a biological molecule in a form in which it exhibits a property and/or activity by which it is characterized.

[00512] *Homology*: As used herein, the term “homology” refers to the overall relatedness between polymeric molecules, *e.g.* between nucleic acid molecules (*e.g.* DNA molecules and/or RNA molecules) and/or between polypeptide molecules. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identical or similar. The term “homologous” necessarily refers to a comparison between at least two sequences (polynucleotide or polypeptide sequences). In accordance with the invention, two polynucleotide sequences are considered to be homologous if the polypeptides they encode are at least about 50%, 60%, 70%, 80%, 90%, 95%, or even 99% for at least one stretch of at least about 20 amino acids. In some embodiments, homologous polynucleotide sequences are characterized by the ability to encode a stretch of at least 4–5 uniquely specified amino acids. For polynucleotide sequences less than 60 nucleotides in length, homology is determined by the ability to encode a stretch of at least 4–5 uniquely specified amino acids. In accordance with the invention, two protein sequences are considered to be homologous if the proteins are at least about 50%, 60%, 70%, 80%, or 90% identical for at least one stretch of at least about 20 amino acids.

[00513] *Identity*: As used herein, the term “identity” refers to the overall relatedness between polymeric molecules, *e.g.*, between oligonucleotide molecules (*e.g.* DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Calculation of the percent identity of two polynucleotide sequences, for example, can be performed by aligning the two sequences for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second nucleic acid sequences for optimal alignment and non-identical sequences can be disregarded for comparison purposes). In certain

embodiments, the length of a sequence aligned for comparison purposes is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or 100% of the length of the reference sequence. The nucleotides at corresponding nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which needs to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. For example, the percent identity between two nucleotide sequences can be determined using methods such as those described in *Computational Molecular Biology*, Lesk, A. M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D. W., ed., Academic Press, New York, 1993; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; *Computer Analysis of Sequence Data, Part I*, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; each of which is incorporated herein by reference. For example, the percent identity between two nucleotide sequences can be determined using the algorithm of Meyers and Miller (CABIOS, 1989, 4:11-17), which has been incorporated into the ALIGN program (version 2.0) using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. The percent identity between two nucleotide sequences can, alternatively, be determined using the GAP program in the GCG software package using an NWSgapdna.CMP matrix. Methods commonly employed to determine percent identity between sequences include, but are not limited to those disclosed in Carillo, H., and Lipman, D., *SIAM J Applied Math.*, 48:1073 (1988); incorporated herein by reference. Techniques for determining identity are codified in publicly available computer programs. Exemplary computer software to determine homology between two sequences include, but are not limited to, GCG program package, Devereux, J., *et al.*, *Nucleic Acids Research*, 12(1), 387 (1984)), BLASTP, BLASTN, and FASTA Altschul, S. F. *et al.*, *J. Molec. Biol.*, 215, 403 (1990)).

[00514] *Inhibit expression of a gene:* As used herein, the phrase “inhibit expression of a gene” means to cause a reduction in the amount of an expression product of the gene. The expression product can be an RNA transcribed from the gene (*e.g.*, an mRNA) or a

polypeptide translated from an mRNA transcribed from the gene. Typically a reduction in the level of an mRNA results in a reduction in the level of a polypeptide translated therefrom. The level of expression may be determined using standard techniques for measuring mRNA or protein.

[00515] *In vitro*: As used herein, the term “*in vitro*” refers to events that occur in an artificial environment, *e.g.*, in a test tube or reaction vessel, in cell culture, in a Petri dish, *etc.*, rather than within an organism (*e.g.*, animal, plant, or microbe).

[00516] *In vivo*: As used herein, the term “*in vivo*” refers to events that occur within an organism (*e.g.*, animal, plant, or microbe or cell or tissue thereof).

[00517] *Isolated*: As used herein, the term “isolated” refers to a substance or entity that has been separated from at least some of the components with which it was associated (whether in nature or in an experimental setting). Isolated substances may have varying levels of purity in reference to the substances from which they have been associated. Isolated substances and/or entities may be separated from at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or more of the other components with which they were initially associated. In some embodiments, isolated agents are more than about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components.

[00518] *Substantially isolated*: By “substantially isolated” is meant that the compound is substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compound of the present disclosure. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compound of the present disclosure, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

[00519] *Linker*: As used herein, a linker refers to a group of atoms, *e.g.*, 10-1,000 atoms, and can be comprised of the atoms or groups such as, but not limited to, carbon, amino, alkylamino, oxygen, sulfur, sulfoxide, sulfonyl, carbonyl, and imine. The linker can be attached to a modified nucleoside or nucleotide on the nucleobase or sugar moiety at a first end, and to a payload, *e.g.*, a detectable or therapeutic agent, at a second end. The linker may be of sufficient length as to not interfere with incorporation into a nucleic

acid sequence. The linker can be used for any useful purpose, such as to form mmRNA multimers (e.g., through linkage of two or more polynucleotides, primary constructs, or mmRNA molecules) or mmRNA conjugates, as well as to administer a payload, as described herein. Examples of chemical groups that can be incorporated into the linker include, but are not limited to, alkyl, alkenyl, alkynyl, amido, amino, ether, thioether, ester, alkylene, heteroalkylene, aryl, or heterocyclyl, each of which can be optionally substituted, as described herein. Examples of linkers include, but are not limited to, unsaturated alkanes, polyethylene glycols (e.g., ethylene or propylene glycol monomeric units, e.g., diethylene glycol, dipropylene glycol, triethylene glycol, tripropylene glycol, tetraethylene glycol, or tetraethylene glycol), and dextran polymers, Other examples include, but are not limited to, cleavable moieties within the linker, such as, for example, a disulfide bond (-S-S-) or an azo bond (-N=N-), which can be cleaved using a reducing agent or photolysis. Non-limiting examples of a selectively cleavable bond include an amido bond can be cleaved for example by the use of tris(2-carboxyethyl)phosphine (TCEP), or other reducing agents, and/or photolysis, as well as an ester bond can be cleaved for example by acidic or basic hydrolysis.

[00520] *MicroRNA (miRNA) binding site:* As used herein, a microRNA (miRNA) binding site represents a nucleotide location or region of a nucleic acid transcript to which at least the “seed” region of a miRNA binds.

[00521] *Modified:* As used herein “modified” refers to a changed state or structure of a molecule of the invention. Molecules may be modified in many ways including chemically, structurally, and functionally. In one embodiment, the mRNA molecules of the present invention are modified by the introduction of non-natural nucleosides and/or nucleotides, e.g., as it relates to the natural ribonucleotides A, U, G, and C. Noncanonical nucleotides such as the cap structures are not considered “modified” although they differ from the chemical structure of the A, C, G, U ribonucleotides.

[00522] *Mucus:* As used herein, “mucus” refers to the natural substance that is viscous and comprises mucin glycoproteins.

[00523] *Multipotent:* As used herein, “multipotent” or “partially differentiated cell” when referring to a cell refers to a cell that has a developmental potential to differentiate into cells of one or more germ layers, but not all three germ layers.

[00524] *Naturally occurring:* As used herein, “naturally occurring” means existing in nature without artificial aid.

[00525] *Non-human vertebrate*: As used herein, a “non human vertebrate” includes all vertebrates except *Homo sapiens*, including wild and domesticated species. Examples of non-human vertebrates include, but are not limited to, mammals, such as alpaca, banteng, bison, camel, cat, cattle, deer, dog, donkey, gayal, goat, guinea pig, horse, llama, mule, pig, rabbit, reindeer, sheep water buffalo, and yak.

[00526] *Off-target*: As used herein, “off target” refers to any unintended effect on any one or more target, gene, or cellular transcript.

[00527] *Oligopotent*: As used herein, “oligopotent” when referring to a cell means to give rise to a more restricted subset of cell lineages than multipotent stem cells..

[00528] *Open reading frame*: As used herein, “open reading frame” or “ORF” refers to a sequence which does not contain a stop codon in a given reading frame.

[00529] *Operably linked*: As used herein, the phrase “operably linked” refers to a functional connection between two or more molecules, constructs, transcripts, entities, moieties or the like.

[00530] *Optionally substituted*: Herein a phrase of the form “optionally substituted X” (e.g., optionally substituted alkyl) is intended to be equivalent to “X, wherein X is optionally substituted” (e.g., “alkyl, wherein said alkyl is optionally substituted”). It is not intended to mean that the feature “X” (e.g. alkyl) *per se* is optional.

[00531] *Peptide*: As used herein, “peptide” is less than or equal to 50 amino acids long, e.g., about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 amino acids long.

[00532] *Paratope*: As used herein, a “paratope” refers to the antigen-binding site of an antibody.

[00533] *Patient*: As used herein, “patient” refers to a subject who may seek or be in need of treatment, requires treatment, is receiving treatment, will receive treatment, or a subject who is under care by a trained professional for a particular disease or condition.

[00534] *Pharmaceutically acceptable*: The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[00535] *Pharmaceutically acceptable excipients*: The phrase “pharmaceutically acceptable excipient,” as used herein, refers any ingredient other than the compounds described herein (for example, a vehicle capable of suspending or dissolving the active

compound) and having the properties of being substantially nontoxic and non-inflammatory in a patient. Excipients may include, for example: antiadherents, antioxidants, binders, coatings, compression aids, disintegrants, dyes (colors), emollients, emulsifiers, fillers (diluent), film formers or coatings, flavors, fragrances, glidants (flow enhancers), lubricants, preservatives, printing inks, sorbents, suspending or dispersing agents, sweeteners, and waters of hydration. Exemplary excipients include, but are not limited to: butylated hydroxytoluene (BHT), calcium carbonate, calcium phosphate (dibasic), calcium stearate, croscarmellose, crosslinked polyvinyl pyrrolidone, citric acid, crospovidone, cysteine, ethylcellulose, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, maltitol, mannitol, methionine, methylcellulose, methyl paraben, microcrystalline cellulose, polyethylene glycol, polyvinyl pyrrolidone, povidone, pregelatinized starch, propyl paraben, retinyl palmitate, shellac, silicon dioxide, sodium carboxymethyl cellulose, sodium citrate, sodium starch glycolate, sorbitol, starch (corn), stearic acid, sucrose, talc, titanium dioxide, vitamin A, vitamin E, vitamin C, and xylitol.

[00536] *Pharmaceutically acceptable salts:* The present disclosure also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, “pharmaceutically acceptable salts” refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form (e.g., by reacting the free base group with a suitable organic acid). Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, quaternary ammonium, and amine cations,

including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. The pharmaceutically acceptable salts of the present disclosure include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present disclosure can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418, *Pharmaceutical Salts: Properties, Selection, and Use*, P.H. Stahl and C.G. Wermuth (eds.), Wiley-VCH, 2008, and Berge et al., *Journal of Pharmaceutical Science*, 66, 1-19 (1977), each of which is incorporated herein by reference in its entirety.

[00537] *Pharmacokinetic*: As used herein, "pharmacokinetic" refers to any one or more properties of a molecule or compound as it relates to the determination of the fate of substances administered to a living organism. Pharmacokinetics is divided into several areas including the extent and rate of absorption, distribution, metabolism and excretion. This is commonly referred to as ADME where: (A) Absorption is the process of a substance entering the blood circulation; (D) Distribution is the dispersion or dissemination of substances throughout the fluids and tissues of the body; (M) Metabolism (or Biotransformation) is the irreversible transformation of parent compounds into daughter metabolites; and (E) Excretion (or Elimination) refers to the elimination of the substances from the body. In rare cases, some drugs irreversibly accumulate in body tissue.

[00538] *Pharmaceutically acceptable solvate*: The term "pharmaceutically acceptable solvate," as used herein, means a compound of the invention wherein molecules of a suitable solvent are incorporated in the crystal lattice. A suitable solvent is physiologically tolerable at the dosage administered. For example, solvates may be prepared by crystallization, recrystallization, or precipitation from a solution that includes organic solvents, water, or a mixture thereof. Examples of suitable solvents are ethanol, water (for example, mono-, di-, and tri-hydrates), *N*-methylpyrrolidinone (NMP), dimethyl sulfoxide (DMSO), *N,N'*-dimethylformamide (DMF), *N,N'*-dimethylacetamide

(DMAC), 1,3-dimethyl-2-imidazolidinone (DMEU), 1,3-dimethyl-3,4,5,6-tetrahydro-2-(1H)-pyrimidinone (DMPU), acetonitrile (ACN), propylene glycol, ethyl acetate, benzyl alcohol, 2-pyrrolidone, benzyl benzoate, and the like. When water is the solvent, the solvate is referred to as a “hydrate.”

[00539] *Physicochemical*: As used herein, “physicochemical” means of or relating to a physical and/or chemical property.

[00540] *Pluripotent*: As used herein, “pluripotent” refers to a cell with the developmental potential, under different conditions, to differentiate to cell types characteristic of all three germ layers.

[00541] *Pluripotency*: As used herein, “pluripotency” or “pluripotent state” refers to the developmental potential of a cell where the cell has the ability to differentiate into all three embryonic germ layers (endoderm, mesoderm and ectoderm).

[00542] *Preventing*: As used herein, the term “preventing” refers to partially or completely delaying onset of an infection, disease, disorder and/or condition; partially or completely delaying onset of one or more symptoms, features, or clinical manifestations of a particular infection, disease, disorder, and/or condition; partially or completely delaying onset of one or more symptoms, features, or manifestations of a particular infection, disease, disorder, and/or condition; partially or completely delaying progression from an infection, a particular disease, disorder and/or condition; and/or decreasing the risk of developing pathology associated with the infection, the disease, disorder, and/or condition.

[00543] *Prodrug*: The present disclosure also includes prodrugs of the compounds described herein. As used herein, “prodrugs” refer to any substance, molecule or entity which is in a form predicate for that substance, molecule or entity to act as a therapeutic upon chemical or physical alteration. Prodrugs may be covalently bonded or sequestered in some way and which release or are converted into the active drug moiety prior to, upon or after administered to a mammalian subject. Prodrugs can be prepared by modifying functional groups present in the compounds in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compounds. Prodrugs include compounds wherein hydroxyl, amino, sulfhydryl, or carboxyl groups are bonded to any group that, when administered to a mammalian subject, cleaves to form a free hydroxyl, amino, sulfhydryl, or carboxyl group respectively. Preparation and use of prodrugs is discussed in T. Higuchi and V. Stella, “Pro-drugs as Novel Delivery Systems,” Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in*

Drug Design, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987, both of which are hereby incorporated by reference in their entirety.

[00544] *Proliferate*: As used herein, the term “proliferate” means to grow, expand or increase or cause to grow, expand or increase rapidly. “Proliferative” means having the ability to proliferate. “Anti-proliferative” means having properties counter to or inapposite to proliferative properties.

[00545] *Progenitor cell*: As used herein, the term “progenitor cell” refers to cells that have greater developmental potential relative to a cell which it can give rise to by differentiation.

[00546] *Protein cleavage site*: As used herein, “protein cleavage site” refers to a site where controlled cleavage of the amino acid chain can be accomplished by chemical, enzymatic or photochemical means.

[00547] *Protein cleavage signal*: As used herein “protein cleavage signal” refers to at least one amino acid that flags or marks a polypeptide for cleavage.

[00548] *Protein of interest*: As used herein, the terms “proteins of interest” or “desired proteins” include those provided herein and fragments, mutants, variants, and alterations thereof.

[00549] *Proximal*: As used herein, the term “proximal” means situated nearer to the center or to a point or region of interest.

[00550] *Purified*: As used herein, “purify,” “purified,” “purification” means to make substantially pure or clear from unwanted components, material defilement, admixture or imperfection.

[00551] *Repeated transfection*: As used herein, the term “repeated transfection” refers to transfection of the same cell culture with a polynucleotide, primary construct or mmRNA a plurality of times. The cell culture can be transfected at least twice, at least 3 times, at least 4 times, at least 5 times, at least 6 times, at least 7 times, at least 8 times, at least 9 times, at least 10 times, at least 11 times, at least 12 times, at least 13 times, at least 14 times, at least 15 times, at least 16 times, at least 17 times at least 18 times, at least 19 times, at least 20 times, at least 25 times, at least 30 times, at least 35 times, at least 40 times, at least 45 times, at least 50 times or more.

[00552] *Reprogramming*: As used herein, “reprogramming” refers to a process that reverses the developmental potential of a cell or population of cells.

[00553] *Reprogramming factor*: As used herein, the term “reprogramming factor” refers to a developmental potential altering factor such as a protein, RNA or small

molecule, the expression of which contributes to the reprogramming of a cell to a less differentiated or undifferentiated state.

[00554] *Sample*: As used herein, the term “sample” or “biological sample” refers to a subset of its tissues, cells or component parts (e.g. body fluids, including but not limited to blood, mucus, lymphatic fluid, synovial fluid, cerebrospinal fluid, saliva, amniotic fluid, amniotic cord blood, urine, vaginal fluid and semen). A sample further may include a homogenate, lysate or extract prepared from a whole organism or a subset of its tissues, cells or component parts, or a fraction or portion thereof, including but not limited to, for example, plasma, serum, spinal fluid, lymph fluid, the external sections of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, tumors, organs. A sample further refers to a medium, such as a nutrient broth or gel, which may contain cellular components, such as proteins or nucleic acid molecule.

[00555] *Signal Sequences*: As used herein, the phrase “signal sequences” refers to a sequence which can direct the transport or localization of a protein.

[00556] *Single unit dose*: As used herein, a “single unit dose” is a dose of any therapeutic administered in one dose/at one time/single route/single point of contact, i.e., single administration event.

[00557] *Similarity*: As used herein, the term “similarity” refers to the overall relatedness between polymeric molecules, e.g. between polynucleotide molecules (e.g. DNA molecules and/or RNA molecules) and/or between polypeptide molecules. Calculation of percent similarity of polymeric molecules to one another can be performed in the same manner as a calculation of percent identity, except that calculation of percent similarity takes into account conservative substitutions as is understood in the art.

[00558] *Somatic cell*: As used herein, “somatic cells” refers to any cell other than a germ cell, a cell present in or obtained from a pre-implantation embryo, or a cell resulting from proliferation of such a cell *in vitro*.

[00559] *Somatic stem cell*: As used herein, a “somatic stem cell” refers to any pluripotent or multipotent stem cell derived from non-embryonic tissue including fetal, juvenile and adult tissue.

[00560] *Somatic pluripotent cell*: As used herein, a “somatic pluripotent cell” refers to a somatic cell that has had its developmental potential altered to that of a pluripotent state.

[00561] *Split dose*: As used herein, a “split dose” is the division of single unit dose or total daily dose into two or more doses.

[00562] *Stable*: As used herein “stable” refers to a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and preferably capable of formulation into an efficacious therapeutic agent.

[00563] *Stabilized*: As used herein, the term “stabilize”, “stabilized,” “stabilized region” means to make or become stable.

[00564] *Stem cell*: As used herein, the term “stem cell” refers to a cell in an undifferentiated or partially differentiated state that has the property of self-renewal and has the developmental potential to differentiate into multiple cell types, without a specific developmental potential. A stem cell may be able capable of proliferation and giving rise to more such stem cells while maintaining its developmental potential.

[00565] *Subject*: As used herein, the term “subject” or “patient” refers to any organism to which a composition in accordance with the invention may be administered, *e.g.*, for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include animals (*e.g.*, mammals such as mice, rats, rabbits, non-human primates, and humans) and/or plants.

[00566] *Substantially*: As used herein, the term “substantially” refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of interest. One of ordinary skill in the biological arts will understand that biological and chemical phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term “substantially” is therefore used herein to capture the potential lack of completeness inherent in many biological and chemical phenomena.

[00567] *Substantially equal*: As used herein as it relates to time differences between doses, the term means plus/minus 2%.

[00568] *Substantially simultaneously*: As used herein and as it relates to plurality of doses, the term means within 2 seconds.

[00569] *Suffering from*: An individual who is “suffering from” a disease, disorder, and/or condition has been diagnosed with or displays one or more symptoms of a disease, disorder, and/or condition.

[00570] *Susceptible to*: An individual who is “susceptible to” a disease, disorder, and/or condition has not been diagnosed with and/or may not exhibit symptoms of the disease, disorder, and/or condition but harbors a propensity to develop a disease or its symptoms. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition (for example, cancer) may be characterized by one or more of the

following: (1) a genetic mutation associated with development of the disease, disorder, and/or condition; (2) a genetic polymorphism associated with development of the disease, disorder, and/or condition; (3) increased and/or decreased expression and/or activity of a protein and/or nucleic acid associated with the disease, disorder, and/or condition; (4) habits and/or lifestyles associated with development of the disease, disorder, and/or condition; (5) a family history of the disease, disorder, and/or condition; and (6) exposure to and/or infection with a microbe associated with development of the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition will develop the disease, disorder, and/or condition. In some embodiments, an individual who is susceptible to a disease, disorder, and/or condition will not develop the disease, disorder, and/or condition.

[00571] *Sustained release*: As used herein, the term “sustained release” refers to a pharmaceutical composition or compound release profile that conforms to a release rate over a specific period of time.

[00572] *Synthetic*: The term “synthetic” means produced, prepared, and/or manufactured by the hand of man. Synthesis of polynucleotides or polypeptides or other molecules of the present invention may be chemical or enzymatic.

[00573] *Targeted Cells*: As used herein, “targeted cells” refers to any one or more cells of interest. The cells may be found *in vitro*, *in vivo*, *in situ* or in the tissue or organ of an organism. The organism may be an animal, preferably a mammal, more preferably a human and most preferably a patient.

[00574] *Therapeutic Agent*: The term “therapeutic agent” refers to any agent that, when administered to a subject, has a therapeutic, diagnostic, and/or prophylactic effect and/or elicits a desired biological and/or pharmacological effect.

[00575] *Therapeutically effective amount*: As used herein, the term “therapeutically effective amount” means an amount of an agent to be delivered (*e.g.*, nucleic acid, drug, therapeutic agent, diagnostic agent, prophylactic agent, *etc.*) that is sufficient, when administered to a subject suffering from or susceptible to an infection, disease, disorder, and/or condition, to treat, improve symptoms of, diagnose, prevent, and/or delay the onset of the infection, disease, disorder, and/or condition.

[00576] *Therapeutically effective outcome*: As used herein, the term “therapeutically effective outcome” means an outcome that is sufficient in a subject suffering from or susceptible to an infection, disease, disorder, and/or condition, to treat, improve

symptoms of, diagnose, prevent, and/or delay the onset of the infection, disease, disorder, and/or condition.

[00577] *Total daily dose*: As used herein, a “total daily dose” is an amount given or prescribed in 24 hr period. It may be administered as a single unit dose.

[00578] *Totipotency*: As used herein, “totipotency” refers to a cell with a developmental potential to make all of the cells found in the adult body as well as the extra-embryonic tissues, including the placenta.

[00579] *Transcription factor*: As used herein, the term “transcription factor” refers to a DNA-binding protein that regulates transcription of DNA into RNA, for example, by activation or repression of transcription. Some transcription factors effect regulation of transcription alone, while others act in concert with other proteins. Some transcription factor can both activate and repress transcription under certain conditions. In general, transcription factors bind a specific target sequence or sequences highly similar to a specific consensus sequence in a regulatory region of a target gene. Transcription factors may regulate transcription of a target gene alone or in a complex with other molecules.

[00580] *Transcription*: As used herein, the term “transcription” refers to methods to introduce exogenous nucleic acids into a cell. Methods of transfection include, but are not limited to, chemical methods, physical treatments and cationic lipids or mixtures.

[00581] *Transdifferentiation*: As used herein, “transdifferentiation” refers to the capacity of differentiated cells of one type to lose identifying characteristics and to change their phenotype to that of other fully differentiated cells.

[00582] *Treating*: As used herein, the term “treating” refers to partially or completely alleviating, ameliorating, improving, relieving, delaying onset of, inhibiting progression of, reducing severity of, and/or reducing incidence of one or more symptoms or features of a particular infection, disease, disorder, and/or condition. For example, “treating” cancer may refer to inhibiting survival, growth, and/or spread of a tumor. Treatment may be administered to a subject who does not exhibit signs of a disease, disorder, and/or condition and/or to a subject who exhibits only early signs of a disease, disorder, and/or condition for the purpose of decreasing the risk of developing pathology associated with the disease, disorder, and/or condition.

[00583] *Unmodified*: As used herein, “unmodified” refers to any substance, compound or molecule prior to being changed in any way. Unmodified may, but does not always, refer to the wild type or native form of a biomolecule. Molecules may undergo a series of

modifications whereby each modified molecule may serve as the “unmodified” starting molecule for a subsequent modification.

Equivalents and Scope

[00584] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments in accordance with the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the appended claims.

[00585] In the claims, articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[00586] It is also noted that the term “comprising” is intended to be open and permits but does not require the inclusion of additional elements or steps. When the term “comprising” is used herein, the term “consisting of” is thus also encompassed and disclosed.

[00587] Where ranges are given, endpoints are included. Furthermore, it is to be understood that unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or subrange within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

[00588] In addition, it is to be understood that any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Since such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the compositions of the invention (*e.g.*, any nucleic acid or

protein encoded thereby; any method of production; any method of use; *etc.*) can be excluded from any one or more claims, for any reason, whether or not related to the existence of prior art.

[00589] All cited sources, for example, references, publications, databases, database entries, and art cited herein, are incorporated into this application by reference, even if not expressly stated in the citation. In case of conflicting statements of a cited source and the instant application, the statement in the instant application shall control.

[00590] Section and table headings are not intended to be limiting.

EXAMPLES

Example 1. DNA constructs for Synthetic Polynucleotide and sgRNA Production

[00591] Synthetic polynucleotides and sgRNA of the invention are produced via creation of a DNA construct; PCR amplification of a cDNA template; in-vitro transcription of the cDNA template (using, in some embodiments, modified nucleotides); if necessary, post-in vitro transcription enzymatic poly-A tailing and/or 5' capping; and purification and analysis.

[00592] The DNA construct for the synthetic polynucleotide includes an open reading frame. The open reading frame (ORF) coding for the CRISPR related protein, e.g., dCAS9 or dCAS9-activator fusion protein, may be flanked by a 5' untranslated region (UTR) which may contain a strong Kozak translational initiation signal and/or an alpha-globin 3' UTR which may include an oligo(dT) sequence for templated addition of a poly-A tail.

[00593] The ORF may also include various upstream or downstream additions (such as, but not limited to, β -globin, tags, etc.) may be ordered from an optimization service such as, but limited to, DNA2.0 (Menlo Park, CA) and may contain multiple cloning sites which may have XbaI recognition.

[00594] The DNA construct for the sgRNA of the invention is made using similar methods. The plasmid construct includes an insert with a sgRNA sequence including the N(20-25) guides sequence and the guide scaffold sequence flanked on the 5' end with a RNA polymerase specific 5'UTR, e.g., a T7 polymerase 5' UTR.

[00595] The DNA construct for the synthetic polynucleotide and/or the sgRNA is transformed in *E. coli*. For the present invention, NEB DH5-alpha Competent *E. coli* are used. Transformations are performed according to NEB instructions using 100 ng of plasmid. The protocol is as follows:

1. Thaw a tube of NEB 5-alpha Competent *E. coli* cells on ice for 10 minutes.
2. Add 1-5 µl containing 1 pg-100 ng of plasmid DNA to the cell mixture. Carefully flick the tube 4-5 times to mix cells and DNA. Do not vortex.
3. Place the mixture on ice for 30 minutes. Do not mix.
4. Heat shock at 42°C for exactly 30 seconds. Do not mix.
5. Place on ice for 5 minutes. Do not mix.
6. Pipette 950 µl of room temperature SOC into the mixture.
7. Place at 37°C for 60 minutes. Shake vigorously (250 rpm) or rotate.
8. Warm selection plates to 37°C.
9. Mix the cells thoroughly by flicking the tube and inverting.

[00596] Alternatively, incubate at 30°C for 24-36 hours or 25°C for 48 hours.

[00597] A single colony is then used to inoculate 5 ml of LB growth media using the appropriate antibiotic and then allowed to grow (250 RPM, 37° C) for 5 hours. This is then used to inoculate a 200 ml culture medium and allowed to grow overnight under the same conditions.

[00598] To isolate the plasmid (up to 850 µg), a maxi prep is performed using the Invitrogen PURELINK™ HiPure Maxiprep Kit (Carlsbad, CA), following the manufacturer's instructions.

[00599] In order to generate cDNA for *In Vitro* Transcription (IVT), the plasmid is first linearized using a restriction enzyme such as XbaI. A typical restriction digest with XbaI will comprise the following: Plasmid 1.0 µg; 10x Buffer 1.0 µl; XbaI 1.5 µl; dH₂O up to 10 µl; incubated at 37° C for 1 hr. If performing at lab scale (< 5µg), the reaction is cleaned up using Invitrogen's PURELINK™ PCR Micro Kit (Carlsbad, CA) per manufacturer's instructions. Larger scale purifications may need to be done with a product that has a larger load capacity such as Invitrogen's standard PURELINK™ PCR Kit (Carlsbad, CA). Following the cleanup, the linearized vector is quantified using the NanoDrop and analyzed to confirm linearization using agarose gel electrophoresis.

Example 2. PCR for cDNA Production

[00600] The linearized plasmid (encoding the synthetic polynucleotide encoding the CRISPR related protein or encoding the sgRNA) is amplified using PCR for preparation of cDNA, e.g., a transcription template. PCR procedures for the preparation of cDNA are performed using 2x KAPA HIFI™ HotStart ReadyMix by Kapa Biosystems (Woburn, MA). This system includes 2x KAPA ReadyMix 12.5 µl; Forward Primer (10 µM); 0.75

μl; Reverse Primer (10 uM) 0.75 μl; Template cDNA ; 100 ng; and dH2O diluted to 25.0 μl. The reaction conditions are at 95° C for 5 min. and 25 cycles of 98° C for 20 sec, then 58° C for 15 sec, then 72° C for 45 sec, then 72° C for 5 min. then 4° C to termination.

[00601] In some embodiments, the reverse primer of the instant invention incorporates a poly-T₁₂₀ for a poly-A₁₂₀ in the mRNA. Other reverse primers with longer or shorter poly(T) tracts can be used to adjust the length of the poly(A) tail in the mRNA.

[00602] The reaction is cleaned up using Invitrogen's PURELINK™ PCR Micro Kit (Carlsbad, CA) per manufacturer's instructions (up to 5 μg). Larger reactions will require a cleanup using a product with a larger capacity. Following the cleanup, the cDNA is quantified using the NanoDrop and analyzed by agarose gel electrophoresis to confirm the cDNA is the expected size. The cDNA is then submitted for sequencing analysis before proceeding to the *in vitro* transcription reaction.

Example 3. In vitro Transcription (IVT)

[00603] The amplified product cDNA is used as a template for in vitro transcription to generate synthetic polynucleotides and/or synthetic sgRNAs. The input nucleotide triphosphate (NTP) mix is made in-house using natural and un-natural NTPs. The *in vitro* transcription reaction generates modified synthetic polynucleotides and/or sgRNA.

[00604] A typical *in vitro* transcription reaction includes the following:

Template cDNA	1.0 μg
10x transcription buffer (400 mM Tris-HCl pH 8.0, 190 mM MgCl ₂ , 50 mM DTT, 10 mM Spermidine)	2.0 μl
Custom NTPs (25mM each)	7.2 μl
RNase Inhibitor	20 U
T7 RNA polymerase	3000 U
dH2O	Up to 20.0 μl
Incubation at 37° C for 3 hr-5 hrs.	

[00605] The synthetic polynucleotide and/or synthetic sgRNA may be modified to reduce the cellular innate immune response. The modifications to reduce the cellular response may include those described herein, e.g., pseudouridine (ψ) and 5-methylcytidine (5meC, 5mc or m⁵C). (See, Kariko K et al. Immunity 23:165-75 (2005), Kariko K et al. Mol Ther 16:1833-40 (2008), Anderson BR et al. NAR (2010); each of which is herein incorporated by reference in their entirety).

[00606] The crude IVT mix may be stored at 4° C overnight for cleanup the next day. 1 U of RNase-free DNase is then used to digest the original template. After 15 minutes of

incubation at 37° C, the mRNA is purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. This kit can purify up to 500 µg of RNA. Following the cleanup, the RNA is quantified using the NanoDrop and analyzed by agarose gel electrophoresis to confirm the RNA is the proper size and that no degradation of the RNA has occurred.

Example 4. PolyA Tailing Reaction

[00607] In some embodiments, the synthetic polynucleotide is synthesized without a poly-A tail, i.e., the cDNA construct is amplified without a poly-T. Without a poly-T in the cDNA, a poly-A tailing reaction must be performed before cleaning the final product. This is done by mixing Capped IVT RNA (100 µl); RNase Inhibitor (20 U); 10x Tailing Buffer (0.5 M Tris-HCl (pH 8.0), 2.5 M NaCl, 100 mM MgCl₂)(12.0 µl); 20 mM ATP (6.0 µl); Poly-A Polymerase (20 U); dH₂O up to 123.5 µl and incubation at 37° C for 30 min. If the poly-A tail is already in the transcript, then the tailing reaction may be skipped and proceed directly to cleanup with Ambion's MEGACLEAR™ kit (Austin, TX) (up to 500 µg). Poly-A Polymerase is preferably a recombinant enzyme expressed in yeast.

[00608] For studies performed and described herein, the poly-A tail is encoded in the IVT template to comprise 160 nucleotides in length. However, it should be understood that the processivity or integrity of the polyA tailing reaction may not always result in exactly 160 nucleotides. Hence polyA tails of approximately 160 nucleotides, e.g., about 150-165, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164 or 165 are within the scope of the invention.

Example 5. 5' capping of the synthetic polynucleotide

[00609] 5'-capping of the synthetic polynucleotide (modified RNA) may be completed concomitantly during the *in vitro*-transcription reaction or alternatively enzymatically capped post-transcription.

[00610] 5'-capping of the synthetic polynucleotide(modified RNA) may be completed concomitantly during the *in vitro*-transcription reaction using the following chemical RNA cap analogs to generate the 5'-guanosine cap structure according to manufacturer protocols: 3'-O-Me-m⁷G(5')ppp(5') G [the ARCA cap]; G(5')ppp(5')A; G(5')ppp(5')G; m⁷G(5')ppp(5')A; m⁷G(5')ppp(5')G (New England BioLabs, Ipswich, MA).

[00611] 5'-capping of modified RNA may be completed post-transcriptionally using a Vaccinia Virus Capping Enzyme to generate the "Cap 0" structure: m⁷G(5')ppp(5')G

(New England BioLabs, Ipswich, MA). Cap 1 structure may be generated using both Vaccinia Virus Capping Enzyme and a 2'-O methyl-transferase to generate: m7G(5')ppp(5')G-2'-O-methyl. Cap 2 structure may be generated from the Cap 1 structure followed by the 2'-O-methylation of the 5'-antepenultimate nucleotide using a 2'-O methyl-transferase. Cap 3 structure may be generated from the Cap 2 structure followed by the 2'-O-methylation of the 5'-preantepenultimate nucleotide using a 2'-O methyl-transferase. Enzymes are preferably derived from a recombinant source.

[00612] Post-transcription capping of the synthetic polynucleotide is performed as follows. The reaction mixture includes: IVT RNA 60 µg-180µg and dH₂O up to 72 µl. The mixture is incubated at 65° C for 5 minutes to denature RNA, and then is transferred immediately to ice.

[00613] The protocol then involves the mixing of 10x Capping Buffer (0.5 M Tris-HCl (pH 8.0), 60 mM KCl, 12.5 mM MgCl₂) (10.0 µl); 20 mM GTP (5.0 µl); 20 mM S-Adenosyl Methionine (2.5 µl); RNase Inhibitor (100 U); 2'-O-Methyltransferase (400U); Vaccinia capping enzyme (Guanylyl transferase) (40 U); dH₂O (Up to 28 µl); and incubation at 37° C for 30 minutes for 60 µg RNA or up to 2 hours for 180 µg of RNA.

[00614] The capped synthetic polynucleotide is then purified using Ambion's MEGACLEAR™ Kit (Austin, TX) following the manufacturer's instructions. Following the cleanup, the capped synthetic polynucleotide is quantified using the NANODROP™ (ThermoFisher, Waltham, MA) and analyzed by agarose gel electrophoresis to confirm the RNA is the proper size and that no degradation of the RNA has occurred. The RNA product may also be sequenced by running a reverse-transcription-PCR to generate the cDNA for sequencing.

[00615] The capping efficiency of the synthetic polynucleotide is assayed using a number of experimental techniques. The capped synthetic polynucleotide is transfected into cells and the amount of CRISPR related protein is assayed using, e.g., ELISA. Higher levels of protein expression correlates with higher capping efficiency. The capped synthetic polynucleotide is run on an agarose-urea gel-electrophoresis; a synthetic polynucleotide with a single, consolidated band by electrophoresis corresponds to a higher purity product compared to a synthetic polynucleotide with multiple bands or streaking bands. The capped synthetic polynucleotide is run on an HPLC column; synthetic polynucleotide with a single HPLC peak corresponds to a higher purity product. The capped synthetic polynucleotide is transfected into human primary keratinocytes at multiple concentrations. At 6, 12, 24 and 36 hours post-transfection the amount of pro-

inflammatory cytokines such as TNF-alpha and IFN-beta secreted into the culture medium are assayed by ELISA. Synthetic mRNAs that secrete higher levels of pro-inflammatory cytokines into the medium correspond to a synthetic mRNA containing an immune-activating cap structure. The capped synthetic polynucleotide is analyzed for capping reaction efficiency by LC-MS after capped mRNA nuclease treatment. Nuclease treatment of capped mRNAs yields a mixture of free nucleotides and the capped 5'-5'-triphosphate cap structure detectable by LC-MS. The amount of capped product on the LC-MS spectra is expressed as a percent of total mRNA from the reaction and would correspond to capping reaction efficiency. The cap structure with higher capping reaction efficiency would have a higher amount of capped product by LC-MS.

Example 6. Methods for Analysis of synthetic polynucleotides and synthetic sgRNA

[00616] Agarose Gel Electrophoresis of Modified RNA or RT PCR Products:

Individual modified RNAs (200-400 ng in a 20 µl volume) or reverse transcribed PCR products (200-400 ng) are loaded into a well on a non-denaturing 1.2% Agarose E-Gel (Invitrogen, Carlsbad, CA) and run for 12-15 minutes according to the manufacturer protocol.

[00617] Nanodrop Modified RNA Quantification and UV Spectral Data: Modified RNAs in TE buffer (1 µl) are used for Nanodrop UV absorbance readings to quantitate the yield of each modified RNA from an *in vitro* transcription reaction.

Example 7. Method of Screening for Protein Expression

[00618] After transfection of cells with synthetic polynucleotide encoding a CRISPR related protein, protein expression is analyzed.

A. Electrospray Ionization

[00619] A biological sample which may contain proteins encoded by synthetic polynucleotides administered to the cells or subject is prepared and analyzed according to the manufacturer protocol for electrospray ionization (ESI) using 1, 2, 3 or 4 mass analyzers. A biologic sample may also be analyzed using a tandem ESI mass spectrometry system.

[00620] Patterns of protein fragments, or whole proteins, are compared to known controls for a given protein and identity is determined by comparison.

B. Matrix-Assisted Laser Desorption/Ionization

[00621] A biological sample which may contain proteins encoded by synthetic polynucleotides administered to the cells or subject is prepared and analyzed according to the manufacturer protocol for matrix-assisted laser desorption/ionization (MALDI).

[00622] Patterns of protein fragments, or whole proteins, are compared to known controls for a given protein and identity is determined by comparison.

C. Liquid Chromatography-Mass spectrometry-Mass spectrometry

[00623] A biological sample, which may contain proteins encoded by synthetic polynucleotides, may be treated with a trypsin enzyme to digest the proteins contained within. The resulting peptides are analyzed by liquid chromatography-mass spectrometry-mass spectrometry (LC/MS/MS). The peptides are fragmented in the mass spectrometer to yield diagnostic patterns that can be matched to protein sequence databases via computer algorithms. The digested sample may be diluted to achieve 1 ng or less starting material for a given protein. Biological samples containing a simple buffer background (e.g. water or volatile salts) are amenable to direct in-solution digest; more complex backgrounds (e.g. detergent, non-volatile salts, glycerol) require an additional clean-up step to facilitate the sample analysis.

[00624] Patterns of protein fragments, or whole proteins, are compared to known controls for a given protein and identity is determined by comparison.

Example 8. Microphysiological Systems

[00625] Synthetic polynucleotides encoding CRISPR related proteins are formulated using one of the methods described herein such as in buffer, lipid nanoparticles and PLGA. These formulations are then administered to or contacted with microphysiological systems created from organ chips as described in International Publication Nos. WO2013086502, WO2013086486 and WO2013086505, the contents of each of which are herein incorporated by reference in its entirety.

Example 9: Design of DNA templates for synthetic polynucleotides encoding Cas9 and dCas9-effector domain fusion proteins

[00626] Plasmids were designed for use as in vitro transcription templates to manufacture chemically synthetic polynucleotides as described herein. Two plasmids were designed to contain codon optimized Cas9 as described by Cong and Mali, respectively (Science (2013) 339:819-23 and Science (2013) 339:823-6.) The first plasmid encoded FLAG-tagged Cas9 and the second plasmid encoded untagged Cas9. Two plasmids were designed to contain codon optimized dcas9 that is fused to different

effector domains (see Cell (2013) 154:442-51 and Nature Methods (2013) 10:977-979). Coding sequences were placed under the control of T7 polymerase promoter for in vitro transcription of modified mRNA.

[00627] Sequences of the Cas9, dCAS9 gene and effector domains are found in the Tables below.

[00628] The following 5'UTR and 3'UTR sequences were used:

5' UTR

TCAAGCTTTTGGACCCCTCGTACAGAAGCTAATACGACTCACTATAGGGAAATAAGAGAGAAA
AGAAGAGTAAGAAGAAATATAAGAGCCACC (SEQ ID NO:71)

3' UTR

TGATAATAGgctg gagcctcgggtggccatgctt cttgcccctt gggcctcccc
ccagcccctc ctccccttc tgcaccogtaccoccggtgtt ctttgaataa
agtctgagtg ggcggc TCTAGA (SEQ ID NO:81)

[00629] The following constructs were created: (3) dCas9- HA tag-2xSV40NLS-KRAB and (4) dCas9-NLS-FLAG-VP64.

[00630] In other embodiments, the sequence-encoding codon-optimized *Streptococcus pyogenes* dCas9 gene can be fused with an HA-tag, one or more NLSs (*e.g.*, SV40 NLSs, for example, one or two NLSs, optionally at the N- and/or C-terminus) and/or a detection tag (*e.g.*, HA tag, FLAG tag, tagBFP, or the like). Any of the above can be featured alone or in combination with an effector domain (KRAB, CSD, WRPW, VP64, or p65AD domains) in the Cas9 constructs of the invention, for example, to facilitate up or down-regulation). Sequences of various Cas9, dCAS9 gene and effector domains are found in the Tables below.

[00631] Cas9 or dCas9 expression can be regulated further by engineering of miR regulatory sequences into UTR of the various constructs. Two miR-122 regulatory sequence-containing constructs are made by replacing the standard 3' UTR with a miR-122 regulatory site-containing 3' UTR. miR-122 is highly expressed in the hepatocytes and not in other tissues; therefore, translation of mRNA containing miR-122 regulatory (binding) sites in its 3' UTR is selectively down-regulated in hepatocytes but not in other cells.

[00632] The following constructs were designed: (5) dCas9-KRAB-miR-122 and (6) dCas9-NLS-FLAG-VP64-miR-122).

[00633] Constructs 1-4 (DNA cassettes) were synthesized and inserted into a vector (pJ364 from DNA 2.0, Inc., Menlo Park, CA). The plasmid was transfected into E.coli as

described above, plasmid DNA was purified and linearized using a restriction enzyme (XbaI). The linearized plasmid was PCR amplified and used as a template for in vitro transcription to produce the synthetic polynucleotides of the invention.

[00634]

Example 10: Design of DNA templates for synthetic sgRNAs targeting VEGF.

[00635] Four constructs each encoding a sgRNA targeting a gene of interest, i.e., a VEGF gene, were created for use as templates for in vitro transcription.

[00636] The constructs each included a 5'UTR (T7), a GGG, a 20-25 nucleotide sequence complementary to a sequence upstream of the VEGF gene transcriptional start, and a guide RNA scaffold sequence. The VEGF sgRNA sequences were based on Maeder et al.

[00637] The following sequence was used as the 5'UTR sequence:

TCAAGCTTTTGGACCCCTCGTACAGAAGCTAATACGACTCACTATAGGG (SEQ ID NO:72)

[00638] The following four sgRNA constructs were created; the sequences in bold italic are those complementary to the VEGF gene.

VEGF V1 sgRNA construct sequence

TCAAGCTTTTGGACCCCTCGTACAGAAGCTAATACGACTCACTATAGGG ***TGTGCAGACGGCAGTCACTAGG***GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTGAAAAGTGGCACCCGAGTCGGTGC (SEQ IN NO:101)

VEGF V2 sgRNA construct sequence

TCAAGCTTTTGGACCCCTCGTACAGAAGCTAATACGACTCACTATAGGG ***AGCAGCGTCTTCGAGAGTGAGG***GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTGAAAAGTGGCACCCGAGTCGGTGC (SEQ ID NO:102)

VEGF V3 sgRNA construct sequence

TCAAGCTTTTGGACCCCTCGTACAGAAGCTAATACGACTCACTATAGGG ***TGAGTGAGTGTGTGCGTGTGG***GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTGAAAAGTGGCACCCGAGTCGGTGC (SEQ ID NO:103)

VEGF V4 sgRNA construct sequence

TCAAGCTTTTGGACCCCTCGTACAGAAGCTAATACGACTCACTATAGGG ***TTGGAGCGGGGAGAGGCCAGG***GTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTGAAAAGTGGCACCCGAGTCGGTGC (SEQ ID NO:104)

[00639] The four constructs (DNA cassettes) were synthesized and inserted into a vector (pJ224 from DNA 2.0, Inc., Menlo Park, CA). The plasmid was transfected into E.coli as described above and plasmid DNA was purified and linearized using a

restriction enzyme (XbaI). The linearized plasmid was PCR amplified as described herein and used as a template for in vitro transcription to produce the synthetic sgRNA.

Example 11: Production of synthetic polynucleotides and sgRNAs via in vitro transcription.

[00640] Using the methods described herein the two synthetic polynucleotides encoding Cas9, the two synthetic polynucleotides encoding the dCAS9-effector gene fusion proteins, and the four synthetic sgRNAs targeting VEGF were synthesized using in vitro transcription of the constructs described above. Synthetic polynucleotides and synthetic sgRNAs with the following sequences were synthesized.

mRNA sequence of dCas9- HA tag-2xSV40NLS- KRAB

GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAAUAUAAGAGCCACCAUGGACAAGAAAUAC
UCAAUCGGACUUGCUAUCGGAACUAACAGCGUGGGAUGGGCCGUCAUUACUGACGAAUACAA
GGUGCCCUCAAAGAAGUUAAGGUCCUGGGAAAUACCGAUAGACACUCCAUCAAAAAGAACC
UGAUCGGGGCACUGUUGUUCGACUCGGGAGAGACUGCAGAAGCAACCAGGCUCAAGCGCACU
GCGAGGCGCCGGUACACCCGGAGGAAGAACC GCAUCUGCUACCUCCAAGAGAUUUUCAGCAA
CGAGAUGGCAAAGGUCGAUGAUUCGUUCUUCACCCGCCUUGAGGAGUCGUUCCUUGUCGAGG
AGGACAAAAAGCAUGAAAGACAUCCGAUCUUCGGAACAUCGUGGACGAAGUCGCAUACCAU
GAAAAGUACCCUACCAUCUACCAUCUCAGAAAGAAACUCGUCGAUUAACUGAUAAAGGCCGA
CUUGC GGCGUAUCUACCUGGCUCUGGGCGACAUGAUCAAGUUUCGGGGUCACUUUCUCAUCG
AGGGUGAUCUCAACCCGGACAAUUCGACGUUGACAAACUCUUAUCAACUGGUCCAGACG
UACAACCAGCUGUUCGAAGAAAUCCGAUCAACGCAAGCGGAGUGGACGCCAAAGCCAUUCU
GUCGGCCCGCCUCUCGAAGUCGCGUCGCCUGGAAAUCUGAUUGCUCAGCUC CCGGGCGAAA
AGAAGAAUGGCCUGUUUGGAAACCUCUAUCGCACUGUCCUCGGGCUGACUCCCAACUUCAAA
UCGAACUUUGACUUGGCUGAGGAUGCAAAGCUGCAACUCUCCAAAGACACUUACGAUGAUGA
CCUGGACAAUCUCCUGGCGCAGAUCCGGGGAUCAGUAUGCUGACCUGUUCUGGGCGCCAAGA
ACCUGUCUGAUGCCAUCUGCUCUCGGAUAUCCUGAGAGUGAACACUGAGAUACCAAGGCG
CCUCUGAGCGCCUCGAUGAUCAAACGCUACGAUGAACACCAUCAGGACCUCACUCUUCUGAA
GGCUUUGGUGCGGCAGCAGCUUCGGAAAAGUACAAAGAGAUUCUUCUUCGACCAGUCGAAAA
ACGGCUACGCCGGAUACAUUGAUGGCGGCGCAAGCCAGGAGGAAUUCUAUAAGUUUAUACAA
CCGAUCCUGGAGAAGAUGGACGGCACUGAAGAACUUCUGGUCAAGCUGAAUCGAGAGGAUCU
GCUCCGGAAGCAGCGGACCUUCGACAAUGGGUCUAUCCUCACCAAUCCAUCUCGGCGAGC
UGCAUGCGAUUCUGAGGCGCCAGGAGGACUUCUACCCAUUCUGAAAGACAAUCGGGAGAAA
AUCGAAAAGAUUCUGACGUUCGCAUUCCAUACUACGUCGGGCCACUUGC GCGGGGUAAUUC
GAGAUUCGCCUGGAUGACGCGGAAGUCCGAAGAAACCAUCACGCCGUGGAAUUCGAAGAAG
UGGUCGACAAGGGAGCCAGCGCACAGUCCUUCAUUGAGCGCAUGACCAAUUUCGACAAAAAU
CUGCCGAACGAGAAGGUCCUGCCGAAGCAUUCACUGCUGUACGAAUACUUUACCGUGUACAA
CGAACUGACCAAGGUGAAGUACGUCACCGAGGGAAUGAGAAAGCCUGCUUUCUGAGCGGAG
AACAGAAGAAGGCCAUUGUUGACCUCUCUUCUUAAGACUAAUCGCAAAGUGACCGUGAAGCAG
CUUAAAGAGGAUUACUUCAAAAAGAUCGAAUGUUUCGACUCCGUGGAAAUCAGCGGGCUGGA
GGAUAGAUUCAACGCGUCCUUGGGACUUAACCACGACCUCUUAAGAUCAUCAAGGAUAAGG
AUUUCUCGACAAUGAGGAAAACGAAGAUUCCUGGAGGACAUCGUUCUGACUCUGACCCUC
UUUGAGGACCGGGAGAUGAUCGAGGAGAGACUCAAGACCUACGCGCACCUGUUUGACGACAA
AGUGAUGAAGCAACUUAACGCAGGCGCUACCCGGCUGGGGCAGACUGUCACGCAAGUUGA
UCAACGGAAUAGAGAUAAACAGUCCGGAAAGACCAUCCUGGACUCCUGAAGUCCGAUGGA

UUCGCCAACCGGAAUUUCAUGCAGCUCAUCCAUGACGACUCAUUGACUUUCAAGGAGGAUUA
 CAAAAGGCCCAAGUGAGCGGCCAAGGGGACUCCCUUCACGAACACAUCGCAAUUUUGGCCG
 GAUACCAGCGAUUAAGAAGGGAAUCCUGCAGACCGUGAAGGUGGUGGACGAGCUGGUGAAA
 GUGAUGGGACGGCACAAGCCGGAAAACAUCGUGAUCGAGAUGGCCAGAGAGAACCAGACGAC
 UCAAAGGGCCAGAAGAACUCGCGCGAACGCAUGAAGAGAAUAGAAGAGGGAAUUAAGGAAC
 UGGGAUCGCAGAUUCUUGAAGGAGCACCCUGUCGAAAAUACUCAACUCCAGAACGAGAAGCUG
 UACCUGUACUAUCUUCAAAACGGCAGGGACAUGUAUGUCGACCAAGAGCUCGACAUUAACCG
 CCUGUCCGAUUAUGACGUGGACGCCAUCGUGCCGCAGAGCUUUCUCAAGGACGAUUCCAUCG
 ACAACAAAGUGCUCACCCGCAGCGACAAGAAUAGAGGGGAAGUCGGAUAACGUCCCUUCGAA
 GAGGUGGUGAAAAAGAUGAAGAAUUCUGGCGGCAGCUCUGAAUGCAAAGCUCUACACCCA
 ACGGAAGUUUGACAACCUCACCAAGGCAGAAAGAGGAGGACUGUCGGAUUUGGAUAAGGCCG
 GUUUCAUCAAGCGACA AUUGGUGGAAACUCGGCAAUUAACCAAGCAUGUGGCACAGAUUCUG
 GACUCCCGUAUGAACACCAAGUACGACGAGAACGAUAAGCUGAUCCGCGAGGUCAAGGUGAU
 CACCCUCAAAAGCAAACUUGUGUCAGACUUCGGAAGGACUUCCAAUUCUACAAGGUCCGCG
 AAUUCAACAACUACCACCACGCUCAUGACGCAUACCUGAACGCGUGUGGUCGGGACUGCCCUC
 AUCAAGAAGUACCCUAAACUCGAAAGCGAAUUGUGUACGGCGACUACAAAGUGUACGAUGU
 CCGGAAGAUGAUCGCGAAAUCGAGCAGGAGAUCGGAAGGGCGACUGCUAAGUACUUUUUCU
 ACUCGAACAUCAUGAACUUCUUCAAAACCGAAUACCCUGGCUAAUGGGCGAGAUACAGAAAG
 CGCCCGCUGAUCGAAACCAACGGCGAAACCGGUGAAAUCGUGUGGGACAAGGGCCGCGAUUU
 CGCUACUGUGAGAAAGGUCCUUCUCCAUUGCCGCAAGUGAAUUCGUCAAAAAGACUGAGGUGC
 AGACUGGCGGAUUUUCCAAGGAAUCGAUCCUCCCAAAGAGGAACUCAGAUAAAGCUCUACGCG
 CGGAAAAGGAUUGGGACCCUAGAAGUACGGAGGAUUUGAUAGCCCAACUGUGGCCUACUC
 UGUGCUCGUGGUGGCCAAAGUCGAGAAAGGAAAGUCGAAGAAGUUGAAAUCGUGAAAAGAAC
 UCUUGGGAAUCACUAUCAUGGAGCGGUCGUCAUUUGAAAAGAACCCAAUCGACUUCUGGAA
 GCCAAGGGAUACAAGAAGUCAAGAAGACCUGAUCUACAAGCUCUUAAGUACAGCCUGUU
 CGAACUGGAGAACGGAAGGAAACGGAUGCUGGCUUCCGCCGGCGAACUGCAAAGGGCAAUG
 AGCUGGCCUCCCAUCGAAUACGUGAACUUCUUCUACCUUGCCUCCAUUACGAAAAGUUG
 AAGGGCUCACCCGAGGACAAUGAGCAGAAACAGCUCUUGUUGAACAAACACAAACACUACCU
 GGACGAAAUCAUCGAACAAAUCAGCGAGUUCAGCAAGCGCGUCAUUCUGGCGGACGCGAACC
 UGGAUAAAAGUGCUGUCCGCGUACAACAAGCACCCGCGAUAAAGCCGAUACGGGAACAGGCUGAG
 AACAUCAUUCACCUUCUACUCUCUAAUCUGGGAGCCCCCGCCGCCUUCAGUACUUUGA
 UACUACCAUCGACCCGAAGAGAUACACGAGCACCAAGGAAGUGCUCGAUGCCACCCUGAUCC
 ACCAGUCCAUUCUGGUCUGUACGAAACCGCAAUCGAUCUGUCACAGCUCGGAGGAGAUGCG
 UACCCCUACGAUGUCCCCGACUACGCGUCACUCGGUAGCGGCAGCCCGAAGAAGAAAAGAAA
 GGUGGAGGACCCGAAGAAAAGAGGAAGGUUGACGGGAUCGGAAGCGGAUCGAAUGGAUCGU
 CAGGGGGAGGUGGCGGAGGUUUGGACGCAAAUACUUCGCGCCUGGUCACGGACCUUGGUG
 ACCUUAAAAGACGUGUUCGUGGAUUUACACAGGGAAAGAAUGGAAACUGUUGGACACCGCCCA
 GCAGAUUCGUGUACCGGAAUGUGAUGCUGGAGAACUACAAAACUUGGUGUCCUGGGGUUUC
 AACUCACUAAGCCAGAUGUCAUUCUUGACUGGAAAAGGGAGAAGAACCUGUAUAAUAGGCU
 GGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCU
 GCACCCGUACCCCCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC+A(140nt) tail
 (SEQ ID NO:51)

mRNA sequence of dCas9-NLS-FLAG-VP64

GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAUAUAAGAGCCACCAUGGACAAGAAAUA
 UCAAUCGGACUUGCUAUCGGAACUAACAGCGUGGGAUGGGCCGUCAUUACUGACGAAUACAA
 GGUGCCCUCAAAGAAGUUAAGGUCCUGGAAAUACCGAUAGACACUCCAUCAAAAAGAACC
 UGAUCGGGGCACUGUUGUUCGACUCGGGAGAGACUGCAGAAGCAACCAGGCUCAAGCGCACU
 GCGAGGCGCCGGUACACCCGGAGGAAGAACCGAUCUGCUACCUCCAAGAGAUUUUCAGCAA

CGAGAUGGCAAAGGUCGAUGAUUCGUUCUUUCACCGCCUUGAGGAGUCGUUCCUUGUCGAGG
AGGACAAAAGCAUGAAAAGACAUCCGAUCUUCGGAACAUCGUGGACGAAGUCGCAUACCAU
GAAAAGUACCCUACCAUCUACCAUCUCAGAAAAGAAACUCGUCGAUUAACUGAUAAAGGCCGA
CUUGC GGUCGAUCUACCUGGCUCUGGCGCACAUGAUC AAGUUUCGGGGUCACUUUCUCAUCG
AGGGUGAUCUCAACCCGGACAAUUCGACGUUGACAAACUCUUAUCCAACUGGUCCAGACG
UACAACCAGCUGUUCGAAGAAAAUCCGAUCAACGCAAGCGGAGUGGACGCCAAAGCCAUUCU
GUCGGCCCGCCUCUCGAAGUCGCGUCGCCUGGAAAAUCUGAUUGCUCAGCUC CCGGGCGAAA
AGAAGAAUGGCCUGUUUGGAAACCUCAUCGCACUGUCCUCGGGCUGACUCCCAACUUCAAA
UCGAACUUUGACUUUGGCUGAGGAUGCAAAGCUGCAACUCUCCAAAGACACUUACGAUGAUGA
CCUGGACAAUCUCCUGGCGCAGAUCGGGGAUCAGUAUGCUGACCUGUUCUGGGCGCCAAGA
ACCUGUCUGAUGCCAUCCUGCUCUCCGAUAUCCUGAGAGUGAACACUGAGAUACCAAGGCG
CCUCUGAGCGCCUCGAUGAUCAAACGCUACGAUGAACACCAUCAGGACCUCACUCUUCUGAA
GGCUUUGGUGCGGCAGCAGCUUCGGAAAAGUACAAAGAGAUUCUUCUUGACCAGUCGAAAA
ACGGCUCACGCCGGAUACAUGAUGGCGGGCGCAAGCCAGGAGGAAUUCUAUAAGUUUAUCAA
CCGAUCCUGGAGAAGAUGGACGGCACUGAAGAACUUCUGGUCAAGCUGAAUCGAGAGGAUCU
GCUCCGGAAAGCAGCGGACCUUCGACAAUGGGUCUAUCCUCACCAAUCCAUCUCGGCGAGC
UGCAUGCGAUUCUGAGGCGCCAGGAGGACUUCUACCCAUUCUGAAAGACAAUCGGGAGAAA
AUCGAAAAGAUUCUGACGUUCCGCAUUCCAUACUACGUCGGGCCACUUGC GCGGGGUAAUUC
GAGAUUCGCCUGGAUGACGCGGAAGUCCGAAGAAACCAUCACGCCGUGGAAUUCGAAGAAG
UGGUCGACAAGGGAGCCAGCGCACAGUCCUUCAUUGAGCGCAUGACCAAUUCGACAAAAAU
CUGCCGAACGAGAAGGUCCUGCCGAAGCAUUCACUGCUGUACGAAUACUUUACCGUGUACAA
CGAACUGACCAAGGUGAAGUACGUCACCGAGGGAAUGAGAAAGCCUGCUUUCUGAGCGGAG
AACAGAAGAAGGCCAUUGUUGACCUCUCUUC AAGACUAAUCGCAAAGUGACCGUGAAGCAG
CUUAAAAGAGGAUUACUUCAAAAAGAU CGAAUGUUUCGACUCCGUGGAAAUCAGCGGCGUGGA
GGAUAGAUUCAACGCGUCCUUGGGACUUAACCACGACCUCUUAAGAUCAUCAAGGAUAAGG
AUUUCUCGACAAUGAGGAAAACGAAGAUUCCUGGAGGACAUCGUUCUGACUCUGACCCUC
UUUGAGGACCGGGAGAUGAUCGAGGAGAGACUCAAGACCUACGCGCACCUGUUUGACGACAA
AGUGAUGAAGCAACUUAACCGCAGGCGCUACACCGGCUGGGGCAGACUGUCACGCAAGUUGA
UCAACGGAAUUGAGAUAAACAGUCCGGAAAGACCAUCCUGGACUUCUGAAGUCCGAUGGA
UUCGCCAACC GGAAUUCAUUGCAGCUCAUCCAUGACGACUCAUUGACUUUCAAGGAGGAUUA
CCAAAAGGCCAAGUGAGCGGCCAAGGGGACUCCUUCACGAACACAUCGCAAUUUGGCCG
GAUCACCAGCGAUUAAGAAGGGAAUCCUGCAGACCGUGAAGGUGGUGGACGAGCUGGUGAAA
GUGAUGGGACGGCACAAGCCGGAACAUCGUGAUCGAGAUGGCCAGAGAGAACCAGACGAC
UCAAAAAGGGCCAGAAGAACUCGCGCGAACGCAUGAAGAGAAUAGAAGAGGGAAUUAAGGAAC
UGGGAUCGCAGAUUCUUGAAGGAGCACCUCUGCGAAAAUACUCAACUCCAGAACGAGAAGCUG
UACCUGUACUACUUCAAAACGGCAGGGACAUGUAUGUCGACCAAGAGCUCGACAUUAACCG
CCUGUCCGAUUAUGACGUGGACGCCAUCGUGCCGCAGAGCUUUCU CAAGGACGAUUC CAUCG
ACAACAAAGUUCUCACCCGCGAGCGACAAGAAUAGAGGGAAAGUCGGAUAAACGUCCCUUCGGAA
GAGGUGGUGAAAAAGAUGAAGAAUACUGGCGGCAGCUCCUGAAUGCAAAGCUCAUACCCCA
ACGGAAGUUUGACAACCUCACCAAGGCAGAAAGAGGAGGACUGUCGGAAUUGGAUAAGGCCG
GUUUCAUCAAGCGACAAUUGGUGGAAACUCGGCAAUUAACCAAGCAUGUGGCACAGAUUCUG
GACUCCCGUAUGAACACCAAGUACGACGAGAACGAUAAGCUGAUCCGCGAGGUCAAGGUGAU
CACCCUCAAAAGCAAACUUGUGUCAGACUUCGGAAAGGACUUCCAAUUCUACAAGGUCCGCG
AAUUCAACAACUACCACCACGCUCAUGACGCAUACCUGAACGCUGUGGUCGGGACUGCCCUC
AUC AAGAAGUACCCUAAACUCGAAAGCGAAUUGUGUACGGCGACUACAAAGUGUACGAUGU
CCGGAAGAUGAUCGCGAAAUCGAGCAGGAGAUCCGAAAGGCGACUGCUAAGUACUUUUUCU
ACUCGAACAUCAUGAACUUCUUCAAAACCGAAAUCACCCUGGCUAAUGGCGAGAUACAGAAAG
CGCCCGCUGAUCGAAACCAACGGCGAAACCGGUGAAAUCGUGUGGGACAAGGGCCGCGAUUU
CGCUACUGUGAGAAAGGUCCUUUCAUGCCGCAAGUGAAUAUCGUCAAAAAGACUGAGGUGC
AGACUGGCGGAUUUCCAAGGAAUCGAUCCUCCCAAAGAGGAACUCAGAUAAAGCUCAUUCGCG
CGGAAAAGGAUUGGGACCCUAGAAGUACGGAGGAUUUGAUAGCCCAACUGUGGCCUACUC

UGUGCUCGUGGUGGCCAAAGUCGAGAAAGGAAAGUCGAAGAAGUUGAAAUCCGUGAAAAGAAC
 UCUUGGGAAUCACUAUCAUGGAGCGGUCGUCAUUUGAAAAGAACCCAAUCGACUUCUGGAA
 GCCAAGGGAAUACAAAGAAGUCAAGAAAGACCUGAUCUAAGCUCCCUAAGUACAGCCUGUU
 CGAACUGGAGAACGGAAGGAAACGGAUGCUGGCUUCCGCCGGCGAACUGCAAAGGGCAAUG
 AGCUGGCCUCCCAUCGAAAUACGUGAACUUCUCUACCUUGCCUCCAUUACGAAAAGUUG
 AAGGGCUCACCCGAGGACAAUGAGCAGAAACAGCUCUUGUUGAACAAACACAAACACUACCU
 GGACGAAAUCAUCGAACAAUCAGCGAGUUCAGCAAGCGCGUCAUUCUGGCGGACGCGAACC
 UGGAUAAAAGUGCUGCCGCGUACAACAAGCACCGCGAUAAAGCCGAUACGGGAACAGGCUGAG
 AACAUCAUUCACCUUCUACUCUCACUAAUCUGGGAGCCCCCGCCGCUUCAAGUACUUUGA
 UACUACCAUCGACCGCAAGAGAUACACGAGCACCAAGGAAGUGCUCGAUGCCACCCUGAUCC
 ACCAGUCCAUUACUGGUCUGUACGAAACGCGAAUCGAUCUGUCACAGCUCGGAGGAGAUGGG
 UCACCGAAAAAGAAACGGAAAGUCAGCUCGGAUUAACAAGGAUCACGACGGAGACUACAAGGA
 CCAUGACAUCGACUAUAAGGACGACGACGACAAGGCCGUGGAGGCGGUGGAUCGGGACGCG
 CGGACGCCUUGGAUGACUUCGACCUUGACAUGCUGGGAUCCGACGCACUUGAUGAUUUUGAU
 CUCGAUAUGCUUGGCAGCGACGCACUGGACGAUUUCGACCUCGACAUGCUCGGAUCGGAUGC
 GCUCGACGACUUCGAUCUGGAUAUGCUGUAUAUAGGCUGGAGCCUCGGUGGCCAUGCUUC
 UUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCUUCCUGCACCCGUACCCCGUGGUCUU
 UGAAUAAAGUCUGAGUGGGCGGC+A(140nt) tail (SEQ ID NO:52)

mRNA sequence of dCas9-KRAB-miR122

GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAUAUAAGAGCCACCAUGGACAAGAAGUAC
 UCAAUCGGACUCGCAAUCGGAACCAAUUCAGUCGGCUGGGCAGUCAUUACCGAUGAAUACAA
 GGUGCCGUCGAAGAAGUUCAAAGUCCUGGGUAAACACUGACAGACAUUCGAUCAAAAAGAACC
 UGAUCGGAGCCUUGCUGUUUGAUUCAGGGCAAACCGCCGAAGCUACCCGGUUGAAACGAACU
 GCUAGACGCCGCUACACGCGCCGCAAGAACCAGGUAUCGCUACCUCCAAGAAAUCUUCUCGAA
 CGAAUUGGCUAAGGUGGACGACUCGUUCUUCACCGGCUCGAGGAGUCAUUCUUGUGGAGG
 AAGAUAGAAGCACGAAAGACACCCGAUCUUCGGCAACAUCGUGGACGAAGUCGCGUACCAC
 GAAAAGUACCCGACUAUCUACCAUCUCCGGAAGAAGCUCGUGGAUAGCACCGAUAAAGGCCGA
 UCUGCGACUGAUCUACCUCGCGCUGGCCCAUAUGAUUAAGUUCGCGGGCACUUCUUCUACUG
 AAGGGGACCUGAAUCCAGACAACUCGGACGUGGAUAAGCUGUUUAUCCAGCUGGUGCAGACU
 UACAAUCAAUUGUUUGAAGAAAACCCUAUCAACGCGUCUGGGGUGGACGCAAAGGCCAUCCU
 GAGCGCGCGGUGUCAAAAUCCAGACGGCUGGAAAUCUGAUAGCCCAACUGCCGGGCGAGA
 AGAAAACGGCCUGUUUGGAAAUCUUAUCGCCUGUCCUGGGACUGACCCCAACUUCUACAG
 UCGAACUUCGACUUGGCCGAGGAUGCGAAGCUCAGCUCAGCAAAGACACCUACGACGAUGA
 CCUCGAUAACCUGUUGGCCAGAUCCGUGACCAGUAUGCUGAUCUUCUUGGCGGCCAAGA
 ACCUGUCAGACGCAAUUCUGCUCUCCGACAUCUCCGCGGUGAAUACUGAGAUACUAAAAGCC
 CCAUUGAGCGCGUCGAUGAUCAAAGAUACGACGAGCACCACCAGGAUCUGACUCUCCUCAA
 GGCACUGGUCCGCCAACAGCUCCCGGAAAAGUACAAAGAGAUUCUUCUUGACCAAUCCAAAA
 ACGGAUACGCUUGGUUACAUAAGACGGCGGAGCGUCACAAGAAGAGUUCUACAAGUUCUACAAG
 CCUAUCCUGGAAAAGAUGGACGGGACCGAGGAACUCCUGGUUAAGCUCAAUAGGGAGGAUCU
 GCUGCGCAAGCAACGCACGUUCGACAAUGGAAGCAUCCCAUCAGAUCCACUGGGGGAGC
 UCCACGCGAUCCUGAGGCGCCAGGAAGAUUUCUACCAUUCUGAAGGACAAUAGAGAGAAA
 AUCGAAAAGAUCCUGACUUCGGAUCCCGUACUACGUGGGCCCGCUCGCACGGGAAACUC
 ACGGUUUGCCUGGAUGACUCGCAAUCCGAAGAAACCAUUAACCCUUGGAAUUCGAGGAGG
 UGGUCGAUAAAGGCGCCUCAGCCAGUCGUUCAUCGAAAGAAUGACCAACUUCGACAAGAAC
 CUCCAAAUGAGAAGGUGCUGCCAAAACAUAAGCCUGCUGUACGAGUACUUUACUGUGUAUAA
 CGAACUCACCAAGGUGAAAUACGUGACCGAGGGAAUGCGCAAGCCGGCAUUUCUGUCGGGCG
 AACAGAAGAAGGCAAUUGUGGACUUGCUGUUCAAAACCAACCAGGAGGACCGUGAAACAG

CUCAAGGAAGAUUACUUUAAGAAGAUCGAGUGUUUCGAUAGCGUCGAAAUUUCGGGGGUGGA
AGAUCGCUUCAUGCAAGCCUUGGGACGUACCACGAUCUGCUUAAGAUCAUUAAGGACAAGG
AUUUCUUGACAACGAAGAGAACGAGGAUAUUCUCGAGGAUAUCGUCCUGACCCUGACUCUG
UUUGAGGAUAGAGAAAUGAUCGAGGAGAGAUUGAAAACUUACGCACACCUCUUCGACGAUAA
GGUGAUGAAACAGCUGAAAAGGCGUAGAUACACUGGUUUGGGGAAGGCUGUCGAGAAAGCUGA
UCAACGGAAUUAAGGACAAGCAGUCCGGAAAAACCAUCCUGGAUUUCCUCAAGUCCGACGGU
UUCGCCAACCGCAACUUCUUGCAGCUGAUCCACGAUGAUUCCUGACGUUCAAGAGGAUUAU
CCAGAAGGCACAAGUGUCCGGACAAGGAGACUCACUCCACGAGCAUAUCGCUAAUUCGCGG
GAUCGCCAGCUAUCAGAAGGGAAUUCUUGCAGACUGUCAAGGUGGUGGACGAACUGGUGAAA
GUGAUGGGAAAGGCAUAAGCCGGAGAAUAUCGUGAUCGAGAUUGGCGAGGGAAAACCAGACGAC
CCAGAAAGGACAGAAAAACAGCCGGGAACGCAUGAAGCGCAUCGAAGAGGGAAUCAAAGAGC
UUGGGAGCCAAAUCUCAAAGAACACCCUGUGGAAAAUACCCAACUGCAGAAUGAGAAGCUU
UACCUGUAUUACCUCAAAACGGGCGCGACAUGUACGUUGACCAGGAAUUGGACAUUAACCG
GCUUUCGACUACGAUGUGGACGCUAUCGUCCCGCAGUCCUUCUGAAAGACGAUUCGAUCG
ACAAUAAGGUCCUGACUAGAUACAGACAAGAAUCCGGGAAAGUCAGACAACGUGCCUAGCGAA
GAGGUCGUUAAGAAGAUGAAGAAUACUGGCGCCAGCUGCUGAACGCGAAGCUUAUCACUCA
GCGCAAGUUCGACAACCUCACCAAGGCAGAAAGAGGCGGAUUGUCGGAGCUCGACAAAGCUG
GCUUCAUCAAGCGCCAGCUCGUCGAAACUCGCCAGAUUACUAAGCAUGUGGCGCAGAUCCUG
GACAGCCGCAUGAAUACUAAGUAUGAUGAGAAUGACAAGCUCAUCCGGGAGGUGAAGGUCAU
CACCCUGAAGUCCAAGCUGGUGUCCGACUUCGGAAGGACUUCCAAUUCUACAAAGUCAGAG
AAUACAACAAUUAACCAUCACGCGCAUGACGCCUACUUGAAUGCAGUGGUGGGUACUGCCCUC
AUCAAGAAAUACCCAAAGCUUGAAAGCGAGUUUGUCUACGGAGACUACAAGGUGUACGACGU
CCGGAAGAUGAUCGCCAAAUCGGAACAGGAAAUUGGGAAAGGCGACCUCUAAAGUACUUCUUCU
ACUCGAAUAUCAUGAAUUUCUUAAGACCAGGAUCACGCUUGCAAUUGGCGAAAUCGGAAG
CGGCCCCUCAUCGAAACCAACGGAGAAACCGGAGAAAUCGUGUGGGACAAGGGUCGCGAUUU
UGCGACCGUCCGAAAGGUUCUUAAGCAUGCCUCAAGUGAACAUUCGUCAAGAAAACGGAAGUGC
AGACUGGAGGCUUCAGCAAGGAGUCCAUUCUCCGAAACGCAACUCCGACAAACUGAUCGCA
CGCAAGAAAGACUGGGACCCGAAGAAAUACGGAGGCUUCGAUUCGCCGACUGUGGCUUACUC
GGUCCUGGUUGUGGGCAAGGUGGAAAAGGGAAAGUCCAAGAAGCUGAAGUCCGUCAAGGAGC
UCCUCGGAAUCACCAUCAUGGAACGGUCAAGCUUCGAGAAAAACCCAAUUGACUUCUGGAG
GCAAAGGGGUACAAGGAGGUGAAGAAGGAUCUGAUCUCAAACUGCCGAAGUACAGCCUCU
UGAGCUCGAAAACGGACGCAAAAAGGAUGCUGGCCUCCGCCGGAGAGCUGCAAAGGGGAAACG
AGCUUGCCUUGCCUUCCAAGUACGUGAACUUCUGUACCUGGCAUCCACUACGAAAACUG
AAGGGAUCGCCGGAGGACAACGAACAGAAGCAGCUGUUUGUGGAACAACACAAGCAUUAUCU
GGAUGAAAUCAUCGAACAAAUCAGCGAAUUCUCAAAGGGUGAUCUUGGCCGACGCCAACC
UGGAUAAAAGUGCUUUCGCCCUACAACAAACUUCGCGACAAGCCGAUCCGGGAGCAGGCCGAA
AACAUCAUUCACCUGUUUACCCUGACUAAUCUGGGUGCGCCCGCGGCUUUCAAAUAUCUUCGA
UACCACGAUCGACCCGGAAGAGAUACACCAGCACCAAAGAGGUGUUGGACGCGACCCUCAUCC
ACCAAUCUAUUAACGGCCUCUAUGAAACUAGGAUCGACCUCAGCCAGCUGGGAGGCGAUGCC
UACCCUUCGAUGUCCCGGACUACGCCUCGCUUGGGAUCCGGAUUCGGAAGAAGAAGCGGAA
GGUCGAGGACCCAAAGAAAAAGCGCAAAGUGGAUUGGGAUCGGUAGCGGUUCCAACGGUUCU
CGGGUGGCGGCGGAGGCGGCAUGGAUGCUAAGUCACUUAACCGCCUGGUCGCGGACGCUGGUG
ACUUUCAAGAUGUGUUCGUGGAUUUCACUCGUGAGGAAUGGAAAUUGCUGGACACUGCCCA
ACAGAUCGUCUACCGCAACGUCAUGCUUGAAAACUACAAAAACCUUGUCGUCGUGGGAUUC
AGCUGACCAAGCCCAGCUGAUUCUGAGACUGGAGAAGGGCGAAGAACCUUGAUAAUAGGCU
GGAGCCUCGGUGGCCAUGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCU
GCACCCGUACCCCCAAACACCAUUGUCACACUCCAGUGGUCUUUGAAUAAAGUCUGAGUGG
GCGGC+A(140nt) tail (SEQ ID NO:53)

mRNA sequence of dCas9-VP64-miR122

GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAUAUAAGAGCCACCAUGGACAAGAAGUAC
UCAAUCGGACUCGCAAUCGGAACCAAUUCAGUCGGCUGGGCAGUCAUUACCGAUGAAUACAA
GGUGCCGUCGAAGAAGUUCAAAGUCCUGGGUAACACUGACAGACAUUCGAUCAAAAAGAACC
UGAUCGGAGCCUUGCUGUUUGAUUCAGGCGAAACCGCCGAAGCUACCCGGUUGAAACGAACU
GCUAGACGCCGCUACACGCGCCGCAAGAACCGGAUCUGCUACCUCCAAGAAAUCUUCUCGAA
CGAAAUGGCUAAGGUGGACGACUCGUUCUUCACCCGGCUCGAGGAGUCAUUCUUGUGGAGG
AAGAUAGAAGCACGAAAGACACCCGAUCUUCGGCAACAUCGUGGACGAAGUCGCGUACCAC
GAAAAGUACCCGACUAUCUACCAUCUCCGGAAGAAGCUCGUGGAUAGCACCGAUAAAGGCCGA
UCUGCGACUGAUUCACCUCGCGCUGGCCCAUAUGAUUAAGUUCGCGGGCACUUCUCCUCAUCG
AAGGGGACCUGAAUCCAGACAACUCGGACGUGGAUAAGCUGUUUAUCCAGCUGGUGCAGACU
UACAAUCAAUUGUUUGAAGAAAACCCUAUCAACGCGUCUGGGGUGGACGCAAAGGCCAUCCU
GAGCGCGCGCUGUCAAAAUCCAGACGGCUGGAAAUCUGAUAGCCCAACUGCCGGGCGAGA
AGAAAACGGCCUGUUUGGAAAUCUUAUCGCCUCUGCCUGGGACUGACCCCAACUUCAAG
UCGAACUUCGACUUGGCCGAGGAUGCGAAGCUCAGCUCAGCAAAGACACCUACGACGAUGA
CCUCGAUAACCUGUUGGCCAGAUCCGUGACCAGUAUGCUGAUCUCUUCUUGGCCGCCAAGA
ACCUGUCAGACGCAAUUCUGCUCUCCGACAUCUCCGCGGGUGAAUACUGAGAUACUAAAAGCC
CCAUUGAGCGCGUCGAUGAUCAAAAGAUACGACGAGCACCACCAGGAUCUGACUCUCCUCA
GGCACUGGUCCGCCAACAGCUCCCGGAAGUACAAAGAGAUUCUUCUUGACCAAUCCAAAA
ACGGAUACGCUUGGUUACAUAGACGGCGGAGCGUCACAAGAAGAGUUCUACAAGUUCAUCAA
CCUAUCCUGGAAAAGAUGGACGGGACCGAGGAACUCCUGGUUAAGCUCAAUAGGGAGGAUCU
GUCGCGCAAGCAACGCACGUUCGACAAUGGAAGCAUCCCAUCAGAUCCACUGGGGGAGC
UCCACGCGAUCCUGAGGCGCCAGGAAGAUUUCUACCAUUCUGAAGGACAAUAGAGAGAAA
AUCGAAAAGAUCCUGACUUCUCCGAAUCCGUACUACGUGGGCCCGCUCGCACGGGAAAACUC
ACGGUUUGCCUGGAUGACUCGCAAUCCGAAGAAACCAUUAACCCCGGAAUUCGAGGAGG
UGGUCGAUAAAGGCGCCUCAGCCCAGUCGUUCAUCGAAAGAAUGACCAACUUCGACAAGAAC
CUCCCAAUAGAGAAGGUGCUGCCAAAACAUGCCUGCUGUACGAGUACUUUACUGUGUAUAA
CGAACUCACCAAGGUGAAAUACGUGACCGAGGGAAUGCGCAAGCCGGCAUUUCUGUCGGGCG
AACAGAAGAAGGCAAUUGUGGACUUGCUGUUCAAAACCAACCGGAAGGUGACCGUGAAACAG
CUCAAGGAAGAUUACUUAAGAAGAUUCGAGUGUUUCGAUAGCGUCGAAAUUUCGGGGGUGGA
AGAUUCGUUCAUUGCAAGCCUUGGGACGUACCACGAUCUGCUUAAGAUCAUUAAGGACAAGG
AUUUCUUGACAACGAAGAGAACGAGGAUUAUCUGAGGAUAUCGUCCUGACCCUGACUCUG
UUUGAGGAUAGAGAAAUGAUCGAGGAGAGAUUGAAAACUUAACGCACACCUCUUCGACGAUAA
GGUGAUGAAACAGCUGAAAAGGCGUAGAUACACUGGUUGGGGAAGGCUGUCGAGAAAGCUGA
UCAACGGAAUUGGGACAAGCAGUCCGGAACCAUCCUGGAUUUCCUCAAGUCCGACGGU
UUCGCCAACCGCAACUUCAUUGCAGCUGAUCCACGAUGAUUCCUGACGUUCAAGAGGAUUA
CCAGAAGGCACAAGUGUCCGGACAAGGAGACUCACUCCACGAGCAUAUCGCUAAUCUCGCCG
GAUCGCCAGCUAUCAAGAAGGGAAUCUUGCAGACUGUCAAGGUGGUGGACGAACUGGUGAAA
GUGAUGGGAAGGCAUAAGCCGGAGAAUAUCGUGAUCGAGAUUGGCGAGGGAAAACCGACGAC
CCAGAAAGGACAGAAAACAGCCGGGAACGCAUGAAGCGCAUCGAAGAGGGAAUCAAAGAGC
UUGGGAGCCAAAUCUCAAAGAACACCCUGUGGAAAUAACCAACUGCAGAAUGAGAAGCUU
UACCUGUAUUACCUCCAAAACGGGCGCGACAUGUACGUUGACCAGGAAUUGGACAUUAACCG
GCUUUCGACUACGAUGUGGACGCUAUCGUCCCGCAGUCCUCCUGAAAGACGAUUCGAUCG
ACAAUAAGGUCCUGACUAGAUCAGACAAGAAUCGGGGAAAGUCAGACAACGUGCCUAGCGAA
GAGGUCGUUAAGAAGAUGAAGAAUACUGGGCGCCAGCUGCUGAACGCGAAGCUUAUCACUCA
GCGCAAGUUCGACAACCUCACCAAGGCAGAAAGAGGCGGAUUGUCGGAGCUCGACAAAGCUG
GCUUCAUCAAGCGCCAGCUCGUCGAAACUCGCCAGAUUACUAAGCAUGUGGGCGCAGAUCCUG
GACAGCCGCAUGAAUACUAAGUAUGAUGAGAAUGACAAGCUCAUCCGGGAGGUGAAGGUCAU
CACCCUGAAGUCCAAGCUGGUGUCCGACUUCGGAAGGACUUCCAAUUCUACAAGUCAGAG
AAAUCAACAUAUACCAUCACGCGCAUGACGCCUACUUGAAUGCAGUGGUGGGUACUGCCCUC
AUCAAGAAAUAACCAAAGCUUGAAAGCGAGUUUGUCUACGGAGACUACAAGGUGUACGACGU
CCGGAAGAUGAUCGCCAAAUCGGAACAGGAAAUUGGGAAGGCGACCUCUAAGUACUUCUUCU

ACUCGAAUAUCAUGAAUUUCUUCAAGACCGAGAUCACGCUUGCAAUUGGCGAAAUCCGGAAG
 CGGCCCCUCAUCGAAACCAACGGAGAAAACCGGAGAAAUCGUGUGGGACAAGGGUCGCGAUUU
 UGCGACCGUCCGAAAGGUUCUUAGCAUGCCUCAAGUGAACAU CGUCAAGAAAACGGAAGUGC
 AGACUGGAGGCUUCAGCAAGGAGUCCAUUCUCCGAAACGCAACUCCGACAAACUGAUCGCA
 CGCAAGAAAAGACUGGGACCCGAAGAAAUACGGAGGCUUCGAUUCGCCGACUGUGGCUUACUC
 GGUCCUGGUUGUGGGCCAAGGUGGAAAAGGGAAAGUCCAAGAAGCUGAAGUCCGUCAAGGAGC
 UCCUCGGAAUCACCAUCAUGGAAACGGUCAAGCUUCGAGAAAAACCCAAUUGACUUCUGGAG
 GCAAAGGGGUACAAGGAGGUGAAGAAGGAUCUGAUCAUCAAACUGCCGAAGUACAGCCUCUU
 UGAGCUCGAAAACGGACGCAAAAAGGAUGCUGGCCUCCGCCGGAGAGCUGCAAAGGGGAAACG
 AGCUUGCCUUGCCUUCCAAGUACGUGAACUUCUGUACCUGGCAUCCACUACGAAAAACUG
 AAGGGAUCGCCGGAGGACAACGAACAGAAGCAGCUGUUUGUGGAACAACACAAGCAUUAUCU
 GGAUGAAAUCAUCGAACAAAUCAGCGAAUUCUCAA AAAAGGGUGAUCUUGGCCGACGCCAACC
 UGGAUAAAAGUGCUUUCGCCUACAACAAACAU CGCGACAAGCCGAUCCGGGAGCAGGCCGAA
 ACAUCAUUCACCUGUUUACCCUGACUAAUCUGGGUGCGCCCGCGGCUUUCAAUACUUCGA
 UACCACGAUCGACCCGGAAGAGAUACACCAGCACCAAAGAGGUGUUGGACGCGACCCUCAUCC
 ACCAAUCUAUUAACCGCCUCUAUGAAACUAGGAUCGACCUCAGCCAGCUGGGAGGCGAUGGA
 UCCCCAAAGAAGAAGAGAAAAGUGUCCUCCGACUACAAGGAUCAUGAUGGGGACUAUAAAGA
 UCAUGAUUUGAUUACAAGGACGACGACGACAAGGCCGUGGAGGAGGAGGUUCCGGCCGCG
 CCGAUGCUCUCGACGACUUCGACCUCGACAUGCUGGGAUCCGACGCCUGGACGACUUGAU
 CUGGAUAUGCUGGGCUCGGACGCCCUUGAUGACUUCGAUCUGGACAUGCUGGGGUCGGAUGC
 ACUGGACGACUUCGACCUUGAUUUGCUGUGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCUCUC
 UUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUUCUGCACCCGUACCCCCCAAACACC
 AUUGUCACACUCCAGUGGUCUUUGAAUAAAAGUCUGAGUGGGCGGC+A(140nt) tail
 (SEQ ID NO:54)

mRNA sequence of Cas9 (Mali)

GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAAUAUAAGAGCCACCAUGGACAAGAAAUAC
 AGCAUCGGCCUGGAUUAUGGAACUAACAGCGUUGGAUGGGCAGUGAUCACCGACGAGUACAA
 GGUGCCGAGCAAGAAGUUCAAGGUGCUGGGGAACACUGACCGCCAUUCAAUUAGAAAAACC
 UCAUUGGAGCACUGCUUUUUGACUCGGGUGAGACUGCCGAAGCUACCAGGCUCAAACGCACC
 GCACGCAGACGGUACACCCGCCGCAAGAAUCGCAUCUGCUAUCUGCAAGAGAUCUUUCCAA
 CGAGAUGGCGAAGGUUGACGACAGCUUUUCCACCGGCUGGAAGAGAGCUUCCUCUGGGAAG
 AGGACAAAAGCACGAAAGGCAUCCAAUCUUCGGUAACAUCGUGGACGAAGUGGCGUAUCAC
 GAAAAGUACCCUACCAUCUACCAUCUGCGGAAGAAGCUGGUCGAUUCACGGAUAAGGCAGA
 CCUGAGACUGAUCUACCUGGCUUUGGCCCAUAUGAUCAAUUCGCGGCCAUUCCUGAUCG
 AGGGGGACC UUAACCCGGAUAACUCGGAUGUCGACAAGCUGUUAUCCAGCUGGUCCAAACG
 UAUAACCAACUGUUUGAGGAAAUCCCAUCAACGCUUCGGGGUGGACGCCAAAGCAAUCCU
 CUCCGCGCGCCUGAGCAAGUCACGGCGGCUCGAAAACCUGAUCGCGCAGCUGCCGGGAGAAA
 AGAAAAAUGGACUGUUUGGGAAUCUGAUCGCGCUGUCGCUCCGCCUGACUCCAACUUUAAG
 UCAAUUUCGAUUUGGCCGAAGAUGCCAAGCUGCAGCUGUCAAGGACACUACGACGACGA
 CCUGGACAAUCUGCUGGCCAGAUUGGGGACCAUACGCAGACCUGUUCUUGGCCGCGAAGA
 ACCUGAGCGACGCCAUUCUUCUGUCCGAUUAUCUGAGAGUCAAUACCGAAAUCACUAAGGCU
 CCGCUGUCCGCUCAAUGAUCAAGCGCUACGAUGAACACCACCAGGAUCUCACUCUGUCUCAA
 AGCCCUCGUGAGACAACAAUUGCCUGAAAAGUACAAGGAGAUUCUUCGACCAGAGCAAAA
 ACGGCUACGCAGGCUACAUCGAUGGAGGAGCGUCACAAGAAGAGUUCUACAAGUUAUCAAG
 CCAAUCUUGGAGAAGAUGGACGGUACUGAAGAACUCCUUGUGAAGCUGAAUAGGGAGGAUUU
 GCUCAGAAAGCAGCGGACUUUUGACAACGGCUCGAUCCUCAUCAGAUUCACCUCGGUGAGC
 UGCAUGCCAUCUUUCGGCGCAAGAGGAUUUUUACCCUUCUGAAGGAUAAUCGCGAGAAA
 AUCGAAAAGAUCUGACGUUCAGAAUCCCUACUACGUGGGACCGCUGGCGCGCGGUAACUC
 GCGGUUUGCAUGGAUGACUCGCAAGUCAGAGGAAACUAUCACUCCUUGGAAUUUUGAGGAGG

UCGUCGAUAAGGGAGCCUCCGCCAGUCAUUCAUCGAACGCAUGACCAACUUCGACAAGAAU
 CUUCCGAACGAGAAGGUCCUCCAAAGCACUCCUGUUGUACGAAUACUUCACCGUGUACAA
 UGAGCUGACCAAAGUUAAGUAUGUCACCGAGGGCAUGAGAAAGCCGGCCUCCUCAGCGGCG
 AACAAAAGAAGGCCAUCGUCGACCUCUCUUAAGACCAACCGGAAGGUGACCGUCAAGCAA
 CUCAAGGAGGACUACUUAAGAAGAU CGAAUGCUUUGACUCGGUCGAAAUCAGCGGAGUGGA
 GGACC GGUUAACGCGUCACUGGGUACCUACCAUGAUCUCCUGAAAAUCAUCAAGACAAGG
 ACUUCUGGACAACGAAGAAAACGAGGACAUCUGGAAGAUUUGUGUUGACCCUGACGCUG
 UUCGAGGACC GGAAAUGAUCGAGGAAAGGCUUAAGACCUACGCACACCUCUUCGAUGACAA
 AGUGAUGAAGCAACUGAAGCGGCGGAGAUUACUGGCUGGGGGAGGCUCUCCCGGAAGCUCA
 UUA AUGAUCAGAGACAAACAGUCGGGUAAAACUAUCCUCGACUCCUCAAGUCGGAUGGG
 UUCGCCAACCGGAACUUC AUGCAGCUGAUCCACGAUGAUUCCUUGACCUUCAAGGAAGAUAU
 CCAGAAGGCGCAAGUGAGCGGACAGGGAGAUUCGUUGCACGAACAUUUCGCUAAUCUCGCCG
 GAUCCCCAGCCAUCAAGAAAGGAAUCCUGCAGACCGUGAAGGUGGUGGAUGAACUGGUGAAA
 GUGAUGGGGGCGCCACAAACCAGAGAACAUCGUAUUGAGAUGGCCCGGAGAAUCAGACCAC
 UCAGAAGGGACAAAAGAUCUCCAGAGAGCGGAUGAAAACGCAUCGAGGAAGGCAUCAAGAGC
 UUGGUAGCCAAAUCUGAAGGAACACCCGGUCGAGAACACCCAGCUC CAGAACGAAAAGCUU
 UACCUGUACUACCUC AAAAUGGACGGGACAUGUACGUCGACCAGGAAUUGGACAUCAACAG
 ACUCAGCGACUACGAUGUGGACCAUAUUGUGCCACAGUCCUUCUUAAGGACGACAGCAUCG
 AUAACAAAGUGCUCACUAGAUCAGACAAAAUCGCGGAAAUCAGACAAUGUGCCAUCGGAA
 GAGGUUGUCAAGAAGAUGAAAAACUACUGGAGACAGCUGCUCAAUGCCAAACUUAUCACCCA
 GCGGAAGUUCGACAACCUUACCAAGGCCGAGCGCGGAGGAUUGUCCGAACUCGACAAGGCCG
 GCUUCAUCAAAAGGCAGCUGGUGGAAACCCGGCAGAUACUAAAACGUGGCCCAGAUCCUC
 GAUUCGCGCAUGAACACUAAAUACGAUGAGAAUGACAAGCUGAUUAGGGAAGUCAAGGUCAU
 CACUCUGAAGUCGAAACUGGUGUCGGACUUAAGAAAAGGAUUUCCAGUUCUACAAAGUCCGCG
 AGAUUAACAACUACCACCACGCUCAUGACGCCUACCUGAAUGCAGUUGUGGGCACCCGCGCUG
 AUCAAGAAGUAUCCGAAGCUGGAAUCCGAGUUCGUGUACGGAGAUUACAAAGUGUACGACGU
 GCGCAAGAU GAUCGCCAAGUCGGAACAGGAAAUCGGAAAGGCUACUGCAAAGUACUUCUUCU
 ACUCAACAUC AUGAACUUCUUCAAAACGGAGAUACGCUCGCGAACGGCGAAAUCCGGAAA
 AGGCCGCUCAUUGAAACCAACGGAGAAACCGGGGAGAU CGUGUGGGACAAGGGAAAGGGAUUU
 UGCGACUGUGAGGAAGGUGUUGUCCAU GCCGCAAGUCAUAUUGUGAAAAAGACGGAAGUGC
 AAACCGGAGGAUUCAGCAAAGAAUCCAUCUCCAAAAGCGCAACUCGGACAAACUCAUCGCG
 CGCAAGAAGGAUUGGGACCCCAAGAAUACGGUGGCUUUGACAGCCCUACUGUGGCUUACUC
 CGUCCUCGUCGUGGCUAAAAGUGGAAAAGGGUAAAUCCAAAAGCUCAAAUCGGUGAAGGAGC
 UCCUGGGAAUCACGAUCAUGGAGCGGUCGAGCUUCGAAAAGAAUCCUAUUGAUUUCUGGAG
 GCGAAGGGCUACAAGGAAGUCAAGAAAGACCUGAUCAUCAAGCUCCCGAAGUACAGCCUCU
 CGAGCUCGAAAACGGCAGAAAGAGGAUGCUGGCAUCAGCGGGAGAAUUGCAGAAGGGAAACG
 AACUGGCACUGCCGUCCAAGUACGUGAAUUUUCUCAUCUGGCUAGCCACUACGAAAAGCUG
 AAGGGAUCGCCCCGAGGACAACGAGCAAAAAACAUCGUUCGUGGAGCAGCACAAAGCACUACCU
 GGAUGAGAUCAUCGAGCAGAUUCCGAAUUCGAAAACGCGUGAUCCUUGCCGAUGCCAAUC
 UGGAUAAAAGUGUUGUCGGCUUACAACAAGCAUCGGGAUAAACCGAUCCGCGAACAGGCAGAA
 AACAUCAUUAUCUGUUCACUUGACCAAUCUGGGAGCGCCUGCCGCGUUUAAGUACUUCGA
 CACCACUAUUGAUAGAAAGCGCUACACCUCGACCAAGGAAGUGCUGGACGCUACCCUGAUCC
 ACCAGUCCAUCACCGGACUCUACGAAACUCGCAUUGACCUGUCCAGCUUGGAGGAGAUUCA
 CGGGCCGAUCCAAAGAAAAAGCGCAAGGUCUGAUAAUAGGCUGGAGCCUCGGUGGCCAUGCU
 UCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCCUCCUGCACCCGUACCCCGUGGUC
 UUUGAAUAAAAGUCUGAGUGGGCGGC+A(140nt) tail (SEQ ID NO:55)

mRNA sequence of Cas9 (Cong)

GGGAAUAAGAGAGAAAAGAAGAGUAAGAAGAAUAUAAGAGCCACCAUGGACUACAAGGAC
 CACGACGGAGACUACAAAGACCAUGACAUCGAUUAACAAGGAUGACGAUGACAAAUGGCACC

GAAGAAGAAGAGAAAAGGUCGGAAUUCACGGGGUGCCGGCCCGGACAAAGAAGUACUCAAUUCG
 GACUGGAUAUCGGCACGAACAGCGUGGGUUGGGCAGUGAUCACCGACGAAUACAAGGUGCCG
 AGCAAGAAGUUCAAAGUGCUGGGAAAUAACCGAUCGCCAUUCGAUCAAGAAAAAUCUGAUUGG
 CGCGCUCUUGUUCGACUCGGGAGAGACUGCCGAGGCCACUAGACUGAAGAGGACCGCUAGGC
 GCCGCUACACGAGGGCGCAAAAACCGCAUCUGCUAUCUUCUUCUUCUUCUUCUUCUUCUUCUUC
 GCCAAGGUGGACGACUCCUUUUUCCAUCGGCUGGAAGAUAUUUCUGGUGGAGGAGGACAA
 GAAGCACGAACGCCAUCCCAUUUUCGGCAACAUUGUCGACGAAGUGGCCUAUCAUGAGAAGU
 AUCCGACUAUCUACCACUUGAGAAAAGAAGCUGGUGGACUCCACUGACAAGGCAGAUUCGCGG
 UUGAUCUACCUCGCACUGGCCCAUAUGAUCAAAUUCGGGGACACUUCUUCUUCGAGGGCGA
 CCUUAUACCCGACAAUUCGGAUGUGGAUAAGCUUUUCAUCCAGCUGGUCCAGACCUACAACC
 AACUGUUUGAAGAAAAUCCAAUCAUUGCGAGCGGUGUCGAUGCAAAGGCCAUCCUGAGCGCC
 CGCCUCUCGAAAAGCAGAAGGCUCGAAAACCGAUCGCACAGUUGCCUGGAGAGAAGAAGAA
 CGGCCUCUUCGGCAAUCUACUGCAUUGUCCUGGGACUGACUCCAAACUUCAAAUCCAACU
 UCGACUUGGCCGAGGACGCCAAACUGCAACUGAGCAAAGAUACCUACGAUGAUGACUUGGAC
 AAUCUUCUGGCUCAGAUCGGCGACCAGUACGCCGACCUGUCCUUGCGGCUAAGAACCUGUC
 GGACGCCAUCCUGCUGUCCGACAUCUCCUGCGCGUCAAUACCGAAAUAUCUAAAAGCACCACUCU
 CGGCAUCCAUGAUCAAAGAGAUACGAUGAACACCACCAGGAUCUCACCCUCCUGAAAGCACUG
 GUGCGGCAGCAGCUCUCCUGAGAAAUAACAAGGAAUUCUUCUUGAUCAGUCCAAGAACGGUAU
 CGCCGGAUACAUCGACGGCGGCGGAGCCAAGAGGAAUUCUACAAGUUCAUCAAGCCGAUCC
 UGGAAAAGAUUGGAUGGCACGGAAGAACUCCUGGUCAAACUGAAUAGAGAGGAUCUGCUCCGC
 AAACAACGGACCUUCGAUAACGGAUUCGUAUCCCGCACCAAGAUCCACCUCGGCGAACUGCAUC
 CAUCCUGCGGGCGCAGGAGGACUUUUACCCGUUCCUCAAGACAACAGAGAAAAGAUUCGAGA
 AGAUCUUGACCUUCGCAUCCCGUACUACGUGGGCCCCGUCGCGAGAGGUAACUCCCGCUUU
 GCUUGGAUGACUAGAAAGUCAGAGGAAACGAUACCCCAUGGAACUUCGAAGAGGUGGUUGA
 CAAAGGAGCGAGCGCCCAAUCGUUCAUCGAACGGGAUGACUAAAUUCGAUAAGAUAUCGCCGA
 AUGAGAAGGUCCUGCCUAAGCACUCACUUCUGUAUGAAUACUUAACUGUGUAUAACGAACUC
 ACCAAAGUCAAAUACGUGACUGAGGGAAUGCGCAAGCCUGCGUUUUUGUCCGGCGAGCAGAA
 AAAGGCCAUUCGUGGACUUGCUGUUCAAAACCAACCGCAAGGUGACUGUUAAGCAACUCAAG
 AGGACUACUUAAGAAGAUUCGAAUGCUUUGACUCGGUCGAGAUUUCGGGGUUGAAGAUAGA
 UUCAACGCGUCGCGUGGGAACCUACCAUGAUCUCCUCAAGAUUAUCAAGGACAAAGACUCCU
 GGUAACGAGGAGAAUGAGGACAUCUCCGAAGAUUAUGUGCUUACCCUGACCCUUUUCGAAG
 AUCGCGAAAUGAUCGAAGAACGCCUGAAAACCUACGCUCACCUGUUCGACGAUAAGGUGAUG
 AAACAGUUGAAACGCCGGCGGUACACGGGUUGGGGGCGGCUGUCGCGCAAGCUGAUC AACGG
 AAUUCGGGACAAACAGAGCGGAAAGACCAUCCUCGAUUUUCUGAAGUCCGAUGGUUUUGCCA
 ACCGCAACUUCAUUGCAGCUCAUCCAUGACGAUUCGCUUACCUUUAAGGAGGAUAUCCAGAAG
 GCACAAGUGUCGGGACAAGGGGAUUCGCUCCACGAACACAUCGCCAAUCUGGGCGGGUCGCC
 GGCAAUUAAGAAGGGAAUCCUCCAGACUGUUAAGGUGGUCGACGAGCUGGUGAAGGUGAUGG
 GGAGACAUAAGCCUGAAAACAUAUGAUCGAGAUUGGCGAGAGAAAUAACAACUACUCAGAAG
 GGACAGAAGAAUUCGGGGAGCGGAUGAAGCGCAUCGAGGAGGGAAUCAAGGAACUGGGCUC
 CAAAUCCUGAAAGAGCAUCCCGUGGAAAUAUCUACGUCGAGAACGAGAAGCUUUACCGUGU
 ACUAUCUUCAAAUUGGACGGGACAUGUACGUCGACCAAGAACUGGAUAUCAUUCGGCUCUCC
 GAUUACGACGUCGAUCACAUCGUCCCCCAAUCAUUCUGAAGGAUGAUAGCAUCGAUAACAA
 GGUGUCACUAGAUCAGACAAAACCGGGGAAAGUCAGAUAAACGUCGCCAGCGAAGAAGUCG
 UGAAGAAGAUGAAGAAUUCUGGAGGCAACUUCUGAACGCCAAACUCAUCACUCAGCGCAAG
 UUCGACAACCUGACCAAAGCAGAAAGGGGAGGACUCAGCGAGCUGGACAAGGCUGGUUUCAU
 CAAACGGCAGCUGGUGGAGACUCGCCAAAUCACGAAGCAUGUGGCCCAGAUUCUCGACUCGC
 GCAUGAAUACUAAAGUACGACGAAAACGAUAAGCUGAUCGGGAGGUGAAGGUGAUCACCCUC
 AAGAGCAAGCUCGUGUCCGAUUUCGGAAAGACUUCAGUUCUACAAGGUGCGGGGAGAUUAA
 CAACUACCAUCACGCUCACGACGCUUACCUCAAUGCUGUGGUGGGGACGGCGUUGAUUAAGA
 AGUACCCAAAACUGGAGUCCGAAUUCGUCUACGGAGAUUAACAAGGUCUACGACGUGCGCAAG
 AUGAUUGCCAAGUCGGAGCAGGAAAUUGGGAAAGCGACUGCUAAGUACUUCUUCUACUCGAA

UAUCAUGAACUUCUUCUUCAGACCGAAAUCACCCUGGCUAACGGCGAGAUUCAGGAAACGGCCGC
 UGAUCGAAACUAAUUGGUGAGACUGGGUGAAAUCGUGUGGGUAUAGGGACGGGACUUCGCCACG
 GUCCGCAAGGUCCUCAGCAUGCCGCAAGUGAAUUAUUGUUAAGAAAACCGAAGUGCAGACCGG
 UGGGUUCUCGAAGGAAUCCAUCUGCCAAAGCGCAACUCGGAUAAGCUUAUUGCCCCGCAAGA
 AGGAUUGGGACCCGAAAAAGUACGGUGGGUUCGACUCCCUACCGUGGCGUACUCGGUGUUG
 GUGGUGGCCAAAGUGGAAAAGGGCAAUCAAAGAAGCUCAAGAGCGUCAAGGAGCUGCUGGG
 AAUCACCAUCAUGGAGAGGUCCAGCUUUGAGAAAAACCCGAUCGACUUCUUGGAAGCCAAGG
 GAUACAAAAGAGGUGAAGAAAGACCUGAUCAUCAACUUCCAAAGUACUCCCUGUUCGAACUC
 GAAAACGGGAGGAAGCGCAUGCUCGCCUCAGCCGGGGAACUGCAAAGGGCAACGAACUGGC
 CCUCCCGUCAAAAUACGUCAACUUCUGUACUUGGCGUCACACUACGAAAAGCUGAAAGGAU
 CCCCAGAGGACAACGAACAGAAACAGCUGUUCGUCGAGCAGCACAAGCACUACCUGGACGAG
 AUCAUCGAACAGAUUCGGAUUUCAGCAAGAGAGUGAUUCUUGGCAGACGCUAACCUUGACAA
 AGUCCUCUCGGCAUACAAUAAAGCAUCGCGACAAGCCGAUCAGAGAACAGGCGGAGAACAUCA
 UCCACCUGUUCACUCUCACCAACCUGGGCGGCCAGCGGCUUUUAAGUACUUGAUACCACC
 AUUGACCGCAAGAGAUACACCUCAAUAAAAGAAGUGCUGGACGCAACCCUGAUCCAUCAAG
 CAUACCCGGACUUUAUGAAACUCGGAUCGAUCUCUCACAGCUCGGAGGAGACAAAAGACCGG
 CUGCCACCAAGAAGGCCGGACAGGCAAAGAAGAAGAAAUGAUAAUAGGCUGGAGCCUCGGUG
 GCAUGCUUCUUGCCCCUUGGGCCUCCCCCAGCCCCUCCUCCCUUCCUGCACCCGUACCC
 CCGUGGUCUUUGAAUAAAGUCUGAGUGGGCGGC+A(140nt) tail (SEQ ID NO:56)

VEGF V1 sgRNA

GGGTGTGCAGACGGCAGTCACTAGGGTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTA
 GTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC (SEQ ID NO:91)

VEGF V2 sgRNA

GGGTGAGTGAGTGTGTGCGTGTGGGTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAG
 TCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC (SEQ ID NO:92)

VEGF V3 sgRNA

GGGTGAGTGAGTGTGTGCGTGTGGGTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAG
 TCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC (SEQ ID NO:93)

VEGF V4 sgRNA

GGGTTGGAGCGGGGAGAAGGCCAGGGTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTA
 GTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC (SEQ ID NO:94)

In vitro expression of CRISPR-related proteins from mRNA constructs

[00641] HeLa cells were transfected with various tagged dCas9 mRNAs and cellular extracts were subjected to Western, Immunoprecipitation, and LC/MS analysis to detect expression of protein expressed by said RNAs.

[00642] A FLAG tagged Cas9 mRNA constructs containing Cong Cas9 (SEQ ID NO: 55), an untagged Cas9 mRNA construct containing Mali Cas9 (SEQ ID NO: 56), a HA-

tagged dCas9 construct containing Gilbert dCas9 (SEQ ID NO: 55), a FLAG-tagged dCas9 construct containing Maeder dCas9 (SEQ ID NO: 52), and and a HA-tagged Trilink Cas9 construct (Trilink) were transfected into HeLa cells according to the following protocol:

[00643] 3×10^6 HeLa cells were plated on 10 cm plates on the day prior to transfection. 30 μ L of Lipofectamine 2000 was diluted into a tube containing 200 μ L Opti-MEM for each sample to be transfected and was held at room temperature for 2-5 minutes. 7.5 ug Cas9 mRNA was diluted into another tube containing 200 μ L Opti-MEM. Then the two dilutions were combined for each sample and mixed. The final solution was allowed to stand at room temperature for 20-25 minutes before the transfection solution was overlaid onto the plated HeLa cells.

[00644] After seven hours post-transfection, cells were lysed with lysis buffer in the presence of 1x HALT protease inhibitors (Pierce). Protein from cell lysates was isolated and reserved for western, immunoprecipitation, or LC-MS assays.

[00645] For immunoprecipitation, lysates normalized for protein concentration were added to Sigma M2 agarose beads and washed. Immunoprecipitation was performed at 4°C overnight. Beads were then washed and resuspended in 1X LDS protein sample buffer. Samples were then run on a 4-12% Bis-Tris gel with MES biffer (Life Technologies) using See BluePlus 2™ prestained protein standards (Life Technologies). The nitrocellulose was blocked with 5% milk in PBS for one hour, then blotted with 1:2000 anti-FLAG (Cell Signaling Technologies) (Figure 2) or 1:1000 anti-HA (Covance HA.11 ascites) (Figure 3) in PBS + 0.05% Tween for 4 hours overnight at 4°C. Membranes were then wahed and incubated with anti-rabbit (Sigma) or anti-mouse (Sigma) conjugated to HRP for 45 minutes at room temperature and detected with Pierce Super Signal West Femto (Pierce) according to the manufacturer's instructions.

[00646] Cong Cas9-FLAG protein was detected in HeLa cells transfected with Cong Cas9-FLAG mRNA (SEQ ID NO: 55) *via* both direct anti-FLAG western and immunoprecipitation anti-FLAG western (Figure 2). Maeder Cas9-FLAG protein was detected in HeLA cells transfected with Maeder Cas9-FLAG mRNA (SEQ ID NO: 52) *via* immunoprecipitation anti-FLAG western (Figure 2). Both Mali Cas9 and Gilbert Cas9-HA did not have a FLAG domain and thus did not show up on the anti-FLAG western or immunoprecipitation (Figure 2).

[00647] Gilbert dCas9-HA protein was detected in HeLa cells transfected with Gilbert dCas9-HA mRNA (SEQ ID NO: 51) *via* direct anti-HA western (Figure 3). Trilink Cas9-HA was used as a positive control for dCas9 expression.

[00648] Protein extracts of HeLa cells transfected with Gilbert dCas9-HA mRNA (SEQ ID NO: 51) and Maeder dCas9-FLAG (SEQ ID NO: 52) were analyzed for protein expression of the Cas9 constructs via LC-PRM peptide quantification and compared to an untreated control HeLa cell protein extracts.

[00649] All solvents were HPLC-grade from Sigma-Aldrich and all chemicals where not stated otherwise were obtained from Sigma-Aldrich.

[00650] Samples were prepared in cell lysis buffer containing 8M urea and 0.1M ammonium bicarbonate. BCA assays to determine total protein content were carried out. Samples with 30 µg of protein were reduced using and digested overnight with trypsin. C18 cleanup for mass spectrometry was carried out using MICROSpin™ columns (The Nest Group) according to the manufacturer's instructions. Peptides were dried down redissolved in 30 µl to the final concentration of 1 µg/ul in LC solvent A (1% acetonitrile in water with 0.1% formic acid (FA)) containing a controlpeptide mix for retention time calibration.

[00651] Peptides (1 µg per sample) were analyzed on C18 chromatography columns on a Thermo Scientific Easy nLC nano-liquid chromatography system for all mass spectrometric analysis. LC solvents were A: 1% acetonitrile in water with 0.1% FA; B: 3% water in acetonitrile with 0.1% FA (gradient elution).

[00652] LC-MS/MS runs for peptide discovery were carried out on a Thermo Scientific Q Exactive™ mass spectrometer equipped with a standard nano-electrospray source.

[00653] LC-PRM for was carried out using a predetermined inclusion list containing target peptides precursor masses on a Thermo Scientific Q Exactive™ mass spectrometer equipped with a standard nano-electrospray source as described before (gradient elution).

[00654] The LC gradient for LC-PRM was 5-35% solvent B in 50 minutes followed by 35-100%B in 2 minutes and 100% B for 8 minutes (total gradient length was 60 minutes).

[00655] LC-MS/MS datasets were analyzed using the MaxQuant software package and searches were performed against the UniProt HUMAN database and other databases. For all peptides of target proteins that were successfully detected in LC-MS/MS, LC-PRM assays were generated using the most intense fragment ions from LC-MS/MS analysis. Top 10 most intense assays were selected.

[00656] Data were processed using SpectroDive™ (Biognosys) software for the analysis of LC-MRM and LC-PRM data based on mProphet (Reiter, Rinner et al., Nature Methods 8 (2011), 430–435). 10 peptides representing housekeeping proteins were quantified in every sample to assess sample loading on the column.

[00657] Relative quantification of peptide segments of Cas9 was carried out for each of the protein extracts via LC-PRM. Quantification of the Cas9 peptide segments: “GNELALPSK” (SEQ ID NO: 120), “YFDTTIDR” (SEQ ID NO: 121), and “IPYYVGPLAR” (SEQ ID NO: 122) is shown in Figures 4A, 4B, and 4C, respectively.

Table 2 shows relative quantification for each:

Table 2– LC-PRM Cas9 Peptide Quantification in Cell Lysate Samples

Peptides	Untreated	Raw Data		Normalization to housekeepers	
		Maeder	Gilbert	Maeder	Gilbert
GNELALPSK (SEQ ID NO: 120)	None	28079	10848	24712	12559
YFDTTIDR (SEQ ID NO: 121)	None	18405	8277	16198	9583
IPYYVGPLAR (SEQ ID NO: 122)	3071	49031	21150	43152	24486

In vivo

[00658] Mice (n=1) (8 wk old female BalbC) were administered 350 ul intravenously of 0.04 mg/mL (14µg) FLAG-Cas9 Cong mRNA as described as set forth in SEQ ID NO:55_ (polyA tail of approximately 140 nucleotides not shown in sequence; 5' cap, Cap1; fully modified with 1-methylpseudouridine) formulated in a lipid nanoparticle comprising the lipid Dlin-MC3-DMA described in Table 3 or a control of PBS (mock).

Table 3. Lipid Nanoparticle Formulation

Ionizable Amino Lipid (Lipid/Mol%)	Non-Cationic Lipid (Lipid/Mol%)	Cholesterol (Mol%)	PEG Lipid (Lipid/Mol %)
Dlin-MC3-DMA 50	DSPC 10	Cholesterol 38.5	PEG-DOMG 1.5

[00659] 6 hours after administration the mice were humanely euthanized. Livers were collected from the mice and Cas9 expression was determined by immunoprecipitation as described in detail below.

[00660] Liver tissue was lysed in a Tissue lyser formulation in the presence of protease inhibitors. Protein was extracted and quantitated using a BCA protein assay (Pierce). Samples were normalized by total protein quantification. Approximately 88 mg of protein extract was added to M2 agarose beads (Sigma) for immunoprecipitation. After gently mixing for 3-4 hours at 4°C, beads were aspirated and washed with RIPA buffer to remove supernatant. RIPA buffer supplemented with 200µg / mL 3X FLAG peptide (Sigma) was added to the beads and incubated at 4°C for 30 minutes with shaking at 1100 rpm in a thermomixer. Peptide elution was then performed by spinning down beads and removing supernatant. Samples were incubated at 72°C for 12 minutes on 4-12% Bis Tris gel using MES running buffer (LifeTech) along with prestained protein standards (LifeTech). The gel was transferred onto nitrocellulose using an iBlot™ apparatus for 6 minutes according to manufacturer's instructions. Membrane was removed and placed in a vessel suitable for blotting and washed 2x with ddH₂O. Nitrocellulose was then blocked for 45 minutes with 5% nonfat dry milk in PBS. Nitrocellulose was then blotted with anti-M2 HRP at 1:15,000 in 1% milk PBS 0.05% Tween for 30 minutes at room temperature. The primary antibody solution was then removed and the nitrocellulose was washed with PBS 0.05% Tween. FLAG-Cas9 peptides were detected with Pierce Super Signal Femto™. The nitrocellulose membrane was then exposed for imaging.

[00661] Results of the immunoprecipitation are shown in Figure 5. As shown in Figure 5, a band is present at 150kD only in the Cas9 LNP sample and not in the PBS control treated sample. These data provide evidence of Cas9 protein expression from LNP formulated synthetic mRNA *in vivo* in mice dosed with Cas9-LNP.

Example 12: In vitro modulation of transcription with synthetic polynucleotides encoding dCas9-effector fusion proteins and synthetic sgRNA

[00662] VEGF (vascular endothelial growth factor), one of the most potent angiogenic factors, has recently been identified as an inducer of neoangiogenesis in many tumors including gliomas. U-87 MG and A-172 are two established glioblastoma cell lines. Basal VEGF expression was an order of magnitude higher in U-87 MG compared to A-172. (Int J Dev Neurosci. 1999 Aug-Oct;17(5-6):565-77). This provides a system to test

gene regulation by dCas9-KRAB and dCas9-VP64 with sgRNA directed to 5' flanking genomic sequence of VEGF.

Inhibition of transcription with dCas9-KRAB

[00663] Gilbert 2013 described gene down-regulation using dCas9-kRAB fusion protein. KRAB is a category of transcriptional repression domains present in approximately 400 human zinc finger protein-based transcription factors (KRAB zinc finger proteins) (Huntley 2006). KRAB domains have been shown to be effective repressors of transcription.

[00664] The U-87MG cell line produces endogenous VEGF. The dCas9-kRAB synthetic polynucleotide together with the 4 synthetic sgRNA targeting VEGF described above are used to inhibit transcription of a VEGF gene in U-87MG cells. The following experimental protocol is used.

1. U-87MG are maintained in Dulbecco's modified Eagle medium (DMEM) in 10% FBS, 2 mM glutamine, 100 units/ml streptomycin, and 100 mg/ml penicillin.
2. All transfections are performed in triplicate using Lipofectamine and using transfection protocol described in U.S. patent publication no. US 2013/0259924 (U.S. patent application no. 13/791,922, filed on March 9, 2013) incorporated by reference.
3. 160,000 cells are seeded into 24-well plates the day before transfection.
4. Transfect U-87MG cells with 100ng of dCas9-KRAB mRNA and 100ng of VEGF V1-V4 sgRNAs. For synergy experiments, transfect 100ng of dCas9-KRAB mRNA with 25ng each of VEGF V1-V4 sgRNAs (a total of 100ng of four different VEGF sgRNA)
5. Incubate cells for 8 hours and wash cells to remove transfection agent.
6. Continue to incubate cells in DMEM
7. Culture medium of U-87MG transfected with mRNA encoding VEGFA-targeted sgRNA and dCas9-KRAB was collected 0h, 12h, 24h, 48h after transfection
8. VEGFA protein expression is measured by ELISA.

[00665] Introduction of synthetic polynucleotides encoding dCas9-KRAB with sgRNA targeting VEGF genomic sequence inhibits VEGF transcription compared to untreated cells.

Activation of transcription with dCas9-VP64

[00666] Maeder 2013 and Perez-Pinera 2013 describe an RNA-guided transcriptional activator, created by fusing dCas9 (D10A, H840A) to the VP64 transactivation domain. dCas9-VP64 recognizes genomic target sites through the hybridization of a sgRNA to a target sequence.

[00667] The HEK293 cell line does not produce endogenous VEGF. The dCas9-VP64 synthetic polynucleotide together with the 4 synthetic sgRNA targeting VEGF described above are used to activate transcription of a VEGF gene in HEK293 cells. The following experimental protocol is used.

1. HEK293 are maintained in Dulbecco's modified Eagle medium (DMEM) in 10% FBS, 2 mM glutamine, 100 units/ml streptomycin, and 100 mg/ml penicillin.
2. All transfections are performed in triplicate using Lipofectamine and using transfection protocol described in U.S. patent publication no. US 2013/0259924 (U.S. patent application no. 13/791,922, filed on March 9, 2013).
3. 160,000 cells were seeded into 24-well plates the day before transfection.
4. HEK293 cells are transfected with 100ng of dCas9-VP64 mRNA and 100ng of VEGF V1-V4 sgRNAs. For synergy experiments, transfect 100ng of dCas9-VP64 mRNA with 25ng each of VEGF V1-V4 sgRNAs (a total of 100ng of four different VEGF sgRNA)
5. Incubate cells for 8 hours and wash cells to remove transfection agent.
6. Continue to incubate cells in DMEM.
7. Culture medium of HEK293 transfected with mRNA encoding VEGFA-targeted sgRNA and dCas9-VP64 is collected 0h, 12h, 24h, 48h after transfection
8. VEGFA protein expression is measured by ELISA.
9. VEGFA mRNA is measured by RT-PCR or Taqman or Q-PCR

[00668] Introduction of synthetic polynucleotides encoding dCas9-VP64 with sgRNA targeting VEGF genomic sequence activates VEGF transcription in HEK293 cells compared to untreated cells.

Example 13: Tissue specific modulation of in vitro protein expression

[00669] miR-122 is present in primary human hepatocytes and is not present in hepatocellular carcinoma cell lines such as HepG2, Hep3B, and SK-Hep-1 cells. It has

been shown that miR-122 can down-regulate protein production encoded by an mRNA containing miR-122 binding sites in the 3' UTR in primary hepatocytes, but not in HepG2 and Hep3B

[00670] Primary hepatocytes and HepG2 cells are transfected with dCas9-KRAB, dCas9-KRAB-miR122, dCas9-VP64, dCas9-VP64-miR122 synthetic polynucleotides along with a combination of the four synthetic sgRNA targeting VEGF.

[00671] Transfection of primary hepatocytes is performed as described herein. Table 4 and 5 provide the results.

Table 4: Results of primary hepatocyte transfection.

Synthetic polynucleotide	dCas9-KRAB	dCas9-KRAB-miR122	dCas9-VP64	dCas9-VP64-miR122	Untreated
Synthetic sgRNA					
VEGF V1 sgRNA	Potential decrease in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-KRAB-miR122 mRNA	Potential increase in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-VP64-miR122 mRNA	
VEGF V2 sgRNA	Potential decrease in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-KRAB-miR122 mRNA	Potential increase in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-VP64-miR122 mRNA	
VEGF V3 sgRNA	Potential decrease in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-KRAB-miR122 mRNA	Potential increase in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-VP64-miR122 mRNA	
VEGF V4 sgRNA	Potential decrease in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-KRAB-miR122 mRNA	Potential increase in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-VP64-miR122 mRNA	
VEGF V1-V4 sgRNA	Potential decrease in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-KRAB-miR122 mRNA	Potential increase in protein expression and mRNA transcript	Potentially no change because miR-122 down-regulate the translation of dCas9-VP64-miR122 mRNA	

[00672] Transfection of HEpG2 cells is performed as described herein. The following table provides the results.

Table 5: Results of HEpG2 transfection.

	dCas9-KRAB	dCas9-KRAB-miR122	dCas9-VP64	dCas9-VP64-miR122	Untreated
VEGF V1 sgRNA	Potential decrease in protein expression and mRNA transcript	Potential decrease in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	
VEGF V2 sgRNA	Potential decrease in protein expression and mRNA transcript	Potential decrease in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	
VEGF V3 sgRNA	Potential decrease in protein expression and mRNA transcript	Potential decrease in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	
VEGF V4 sgRNA	Potential decrease in protein expression and mRNA transcript	Potential decrease in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	
VEGF V1-V4 sgRNA	Potential decrease in protein expression and mRNA transcript	Potential decrease in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	Potential increase in protein expression and mRNA transcript	

Example 14: Modulation of *in vivo* protein expression with dCas9

[00673] Delivery of synthetic polynucleotides encoding dCAS9-effector fusion proteins and synthetic sgRNA targeting a gene of interest to the liver using lipid-nucleic acid particle formulation is performed following procedures described U.S. patent publication no. US 2013/0259924 (U.S. patent application no. 13/791,922, filed on March 9, 2013). Delivery of synthetic polynucleotides encoding dCAS9-effector fusion proteins and synthetic sgRNA targeting a gene of interest to muscle cells via intramuscular injection is performed as described in U.S. patent publication no. US 2013/0259924 (U.S. patent application no. 13/791,922, filed on March 9, 2013).

a. *In vivo* inhibition of TPO gene transcription with dCas9-KRAB

[00674] Thrombopoietin (TPO) is produced mainly by the liver. Synthetic polynucleotides encoding dCas9-KRAB or dCas9-KRAB-miR122 along with sgRNA targeting the 5' flanking sequence of thrombopoietin are formulated in an LNP formulation. 0.005mg/kg to 0.5mg/kg of LNP formulated dCas9-KRAB and TPO

sgRNA is administered intravenously to mice. Blood and liver samples are collected at 8h, 24hr, 48hr, 72hr post administration. TPO protein level is measured by ELISA and TPO mRNA is measured by RT-PCR or q-PCR.

[00675] With dCas9-KRAB mediated suppression with TPO sgRNA, TPO protein and mRNA expression will decrease. However, because of the presence of miR-122, groups treated with dCas9-KRAB-miR122 TPO sgRNA will not show significant down regulation of TPO protein production and mRNA expression.

b. *In vivo* inhibition of reporter gene transcription with dCas9-KRAB

[00676] An exogenous DNA plasmid comprising a reporter molecule, such as luciferase, GFP, or human VEGF, and a specific 5' flanking sequence containing VEGF V1-V4 sgRNA sequences and a constitutively active promoter to drive protein expression is co-administered with the synthetic polynucleotide encoding dCas9-KRAB/dCas9-KRAB-miR122 and with the 4 synthetic sgRNA targeting VEGF. Polynucleotides are LNP formulated and administered as described herein.

[00677] There is a decrease in expression of the reporter molecule in groups treated with dCas9-KRAB/VEGF sgRNA. There is not a significant decrease in expression of the reporter molecule in groups treated with dCas9-KRAB-mir122/VEGF sgRNA.

c. *In vivo* activation of LDHC gene transcription with dCas9-VP64

[00678] To demonstrate activation of transcription and protein expression using synthetic polynucleotide encoding dCas9-VP64, proteins that are not typically expressed by the liver are targeted. Lactate dehydrogenase C (LDHC) is a testis specific enzyme. Administration of LNP formulated synthetic polynucleotide encoding dCas9-VP64 along with synthetic sgRNA targeting 5' genomic sequence of mouse LDHC induces the expression of LDHC in the liver. The LDHC mRNA transcript is measured by RT-PCR or qPCR and the protein expression is measured by ELISA or western blot. Administration of LNP formulated synthetic polynucleotide encoding dCas9-VP64-miR122 and sgRNA targeting the 5' genomic sequence of the mouse LDHC does not activate LDHC expression because of the presence of miR-122 in the liver cells.

d. *In vivo* activation of reporter gene transcription with dCas9-VP64

[00679] An exogenous DNA plasmid comprising a reporter molecule, such as luciferase, GFP, or human VEGF, and a specific 5' flanking sequence containing VEGF V1-V4 sgRNA sequences and a constitutively inactive promoter is co-administered with the synthetic polynucleotide encoding dCas9-VP64/dCas9-VP64-miR122 with and the

synthetic sgRNA targeting VEGF. There is an increase in expression of the reporter molecule in groups treated with dCas9-VP64/VEGF sgRNA. There is not a significant increase in expression of the reporter molecule in groups treated with dCas9-VP64-mir122/VEGF sgRNA.

e. Tissue specific *in vivo* modulation of LDHC gene transcription

[00680] As described herein, microRNA can provide tissue specificity. To demonstrate tissue selectivity of dCas9 system using microRNA 122, one group of mice is dosed intravenously and another group is dosed intramuscularly with LNP formulated synthetic polynucleotide encoding dCas9-VP64-mir122 and synthetic sgRNA targeting LDHC. Due to the presence of miR-122 in the liver cells, the synthetic polynucleotide encoding dCas9-VP64 will not be expressed; thus, no dCas9-VP64 mediated activation of LDHC expression. Since muscle cells do not contain miR-122, the group receiving intramuscular injection of dCas9-VP64-mir122 and LDHC sgRNA will show expression of a testis-specific protein.

f. Tissue specific *in vivo* modulation of reporter gene transcription

[00681] Another example is to co-administer a DNA plasmid described in Gilbert et al as a target for gene suppression of dCas9-KRAB. A formulated LNP carrying 1) SV40-GFP DNA plasmid (described in Gilbert et al), 2) sgGFP-NT1 sgRNA (described in Gilbert et al), and 3) mRNA encoding either dCas9-KRAB or dCas9-KRAB-miR122 is administered via IV to the liver, which contain miR-122. The group receiving dCas9-KRAB will show a reduced GFP expression compared to the control group that received no dCas9-KRAB mRNA as measure by flow cytometry or immunohistochemistry because the translation of dCas9-KRAB will not be reduced by miR122. The group receiving dCas9-KRAB-miR122 will show GFP expression similar to the control group receiving no dCas9-KRAB-miR122 mRNA because of the presence of miR-122 in the hepatocytes which inhibits the translation of the dCas9-KRAB-miR122 mRNA. To compared IM delivery to the muscle cells which do not contain miR-122, a formulated LNP carrying 1) SV40-GFP DNA plasmid (described in Gilbert et al), 2) sgGFP-NT1 sgRNA (described in Gilbert et al), and 3) mRNA encoding either dCas9-KRAB or dCas9-KRAB-miR122 is administered via IM injection. We expect to see both the dCas9-KRAB and the dCas9-KRAB-miR122 groups to show GFP expression in the muscle cells as measure by flow cytometry or immunohistochemistry.

[00682] Another example is to co-administer a DNA plasmid described in Gilbert et al as a target for gene activation of dCas9-VP64. A formulated LNP carrying 1) GAL4 UAS GFP DNA plasmid (described in Gilbert et al), 2) sgGAL4-1 sgRNA (described in Gilbert et al), and 3) mRNA encoding either dCas9-VP64 or dCas9-VP64-miR122 is administered via IV to the liver. The group receiving dCas9-VP64 mRNA will show GFP expression as measure by flow cytometry or immunohistochemistry whereas the group receiving dCas9-VP64-miR122 will not show GFP expression because of the presence of miR-122 in the hepatocytes suppressing the translation of dCas9-VP64-miR122 mRNA. To compared IM delivery to the muscle cells which do not contain miR-122, A formulated LNP carrying 1) GAL4 UAS GFP DNA plasmid (described in Gilbert et al), 2) sgGAL4-1 sgRNA (described in Gilbert et al), and 3) mRNA encoding either dCas9-VP64 or dCas9-VP64-miR12 is administered via IM injection. We expect to see both the dCas9-VP64 and the dCas9-VP64-miR122 groups to activate GFP expression in the muscle cells as measure by flow cytometry or immunohistochemistry.

Example 15: Use of 3 different delivery systems

[00683] A combination of targeting mechanism is used to allow for a significant safety window when administering synthetic polynucleotides encoding a CRISPR related protein and sgRNA.

[00684] First, a synthetic polynucleotide encoding a dCAS9-effector domain fusion protein is designed using a miR-142.3p targeted 3'UTR to prevent expression in APCs.

[00685] Second, the synthetic polynucleotide is formulated in an ionizable LNP that targets LDL-R mediated uptake into hepatocytes. This leads to expression of the dCAS9-effector domain fusion protein in the liver.

[00686] Third, the synthetic sgRNA targeting the hepatic gene of interest is conjugated with, e.g., a GalNac conjugate or similar targeting moiety that is preferentially be taken up by hepatocytes. The synthetic sgRNA can be administered separately and after a 2-3 hour delay.

[00687] The combination of targeting mechanisms (in this case 3: mir-3'UTR, LNP, GalNac) and the absence of DNA would allow for significant safety window.

[00688] The result could be transcriptional activation or inhibition of any protein target (our first in vivo experiments will focus on IHC to confirm localization of dCas9 fusion into the liver and functional read outs using reporters like luciferase and/or GFP).

OTHER EMBODIMENTS

[00689] It is to be understood that the words which have been used are words of description rather than limitation, and that changes may be made within the purview of the appended claims without departing from the true scope and spirit of the invention in its broader aspects.

[00690] While the present invention has been described at some length and with some particularity with respect to the several described embodiments, it is not intended that it should be limited to any such particulars or embodiments or any particular embodiment, but it is to be construed with references to the appended claims so as to provide the broadest possible interpretation of such claims in view of the prior art and, therefore, to effectively encompass the intended scope of the invention.

[00691] All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, section headings, the materials, methods, and examples are illustrative only and not intended to be limiting.

Sequence Tables

Table 5: CRISPR related proteins

Description	Sequence	SEQ ID NO
CRISPR-associated endonuclease CAS9/Csn1 protein	MDKKYSIGLDIGTNSVGVAVITDEYKVPKFKVLTGNTDRHSIKKNLIGALLFDSGETAE ATRLKRTARRRYTRRNKRICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFG NIVDEVAYHEKYPTIYHLRKKLVDSADKADRLIYLALAHMIKFRGHFLIEGDLNPDNSD VDKLFIQLVQTYNQFFENPINASVDKAILLSARLSKSRLENLIAQLPEGKKNGLFGN LIALSLGLTPNFKSNFDLAEDAKLQLSKDTYDDDDLNLLAQIGDYADLFLAAKNLSDAI LLSDLRVNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLPEKYKEIFFDQSKNGYA GYIDGGASQEEFYKFKPILEKMDGTEELLVKNREDLLRQKQRTFDNGSIPHQIHLGELH AILRRQEDFYFFLKDNPEKIEKILTFRIPYVYVGLARGNSRFAMTRKSEETITPWNFEE VVDKGASAQSFIERMTNFDKNLPNEKVLPKHSLLYEYFTVYNELTKYVYTEGMRKPAFL SGEQKKAIVDLLFKNKVVTKQKEDYFKKIECFDSVEISGVEDRFNASLGTYYHDLKIKI IKDKDFLDNEENEDILEIVLTLTFEDREMIERLKYAHLFDDKVMKQKRRRYTGWG RLSRKLINGIRDKQSGKTIIDFLKSDGFANRFNQLIHDDSLTFKEDIQKAQVSGQDLSL HEHIANLAGSPAIIKKGILQTVKVVDELVKVGRHKPENIVIEARENQFTQKGGKNSRER MKRIEIGIKELSGOILKEHPVENTQLQNEKLYLYLQNGRDMYVDQELINRLSDYDVDH IVPQSFLLKDDSIDNKVILTRSDKNRGKSDNVPSEVVKKMKNYRQLLNAKLI TQRKFDNL TKAERGLSELDKAGFIKRQLVETRQITKHVAQILDSRMNTKYDENDKLIREVKVI TLKS KLVSDFRKDFQFYKVREINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGDYKVDVVRK MIAKSEQEI GKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDF ATVRKVLMPQVNI VKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFDSPTVA YSVLIVAKVEKSKKLSVKELIGITIMERSSEKPNIDFLEAKGYEKKDLIIKLPK YSLFELENGRKRMLASAGELQKGNELALPSKYVNFYLAHYEKLKSGSPEDNEQKQLFVE QHKHYLDEII EQISEFSKRVILADANLKDVLKSAYNKHRDKPIREQAENIIHLFTLTNLGA PAAFYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGD	7
CRISPR-associated endonuclease CAS9/Csn1 mRNA	auggacaaga aguaucuccau ugggucugau aucggcacaac acagcugcugc cugggcccugc auuacggacg aguaacaaggu gccgagcaaaa aaaaaucaaa uucugggcaa uaccgaucgc cacagcauaa agaagaaccu cauuggcgcc cuccugucg acuccgggga gacggccgaa gccacgcgcg ucaaaaagaac agcacggcgc agauuaacc gcagaaagaa ucggaucugc uaccugcagg agaucuuuag uaaugagau gcuuaaggug augacuuuu cuuccauagg cuggaggagu ccuuuuuggu ggaggaggau aaaaagcacg agcggcacc aauuuuggc aauaucugg acgaguggc guaccaugaa aaguuaccac ccauaauca ucugaggaaag aagcuuugua acaguaucuga uaaggcugac uugcgguuga ucuauucugc gcuggcgcau	60 120 180 240 300 360 420 480

Description	Sequence	SEQ ID NO
	augaucaaa uucgggggaca cuuccucauc gaggggggacc ugaaccacaga caacagcgau	540
	gucgacaaac ucuuuaucca acuggnucag acuuacaauc agcuuuucga agagaaccccg	600
	aucaacgcau cgggaguuga cgccaaagca auccugagcg cuaggcuguc caaaucccgg	660
	cggcucgaaa accucaucgc acagcucccu ggggagaaga agaagggccu guuuguaau	720
	cuuauccccc ugucacucgg gcugaccccc aacuuuaau cuacuucga ccuggccgaa	780
	gaugccaagc uucaacugag caagacacc uacguugaug aucucgacaa ucugcuggcc	840
	cagaucggcg acccagucgc agaccuuuuu uugggggcaa agaaccucug agacgccaau	900
	cugcugagug auuuucugcg agugaacag gagaucacca aagcucgcu gagcgcuaqu	960
	augaucaagc gcuuugaug gacaccacaa gacuuuacuu ugcugaaggc ccuugucaga	1020
	cagcaacugc cugagaagua caaggaaau uuucucgac agucuaaaaa uggcuacgcc	1080
	ggauacaau gggcgggagc agccaggag gaauuuaca auuuuuuaa gcccaucuu	1140
	gaaaaauug accggaccga ggagcugcug guaaagcuaa acagagaaga ucuguuugcg	1200
	aaacagcga cuuucgacaa uggaagcauc cccaccaga uucaccuggg cgaacugcac	1260
	gcuaucuca ggcggcaaga ggauuucuc cccuuuuuga aagauaacag ggaaaagau	1320
	ggaaaaauc ucacauucg gauaccuac uauguagggc ccucgcccg gggaaaauuc	1380
	agaauccgu ggaugacuc caaaucaaga gagaccauca cuccucggaa cuucgaggaa	1440
	gucguggaa agggggccc ugcccagucc uucaucgaaa ggaugacuaa cuuugauaaa	1500
	aaucugcua acgaaaaggu gcuucuaaa cacucucgc uguacgagua cuucacaguu	1560
	uaaaacgagc ucaccaaggu caauacguc acagaagggg ugagaaagcc agcauuccug	1620
	ucuggagagc agaaagaaagc uaucguggac cuccucuca agacgaaccg gaaaguuacc	1680
	gugaaacagc ucaaaagaaagg aggaucgaa aagaauugaa guuucgacuc uguuugaaauc	1740
	agcggagugg aggalucguu caacgcauc cugggaaacgu aucacgauc ccugaaaauc	1800
	auaaagaca aggacuuccu ggacaauagf gagaacgagf acuuucuuu ggcacaauugc	1860
	cucacccua cguuuguuga agauaggagf augauugaaf aacgcuugaa acuuacgcu	1920
	caucucuucg acgacaaaagu caugaaacag cucaagaggg gccgaauac aggauggggg	1980
	cggcugucua gaaaacugau caaugggaur cgagacaagc agaguggaaa gacaauccug	2040
	gauuuuuuu aguccgaurf auuugccaac cggaaacuuca ugcaguuuau ccaugaugac	2100
	ucucucaccu uuaagggagg cauccagaaa gcacaaguuu cuggccagggf ggacagucuu	2160
	cacgagcaca ucgcuuauc ugcagguagc ccagcuauca aaaaaggaa acugcagacc	2220
	guuaaggucg uggauugaacu cguaaaaua augggaaagg auaaagccga gaaauucguu	2280
	aucgagaugf cccgagagaa ccaaacuacc cagaaggagc agaagaaacag uagggaaaagg	2340
	augaaagagg uugagagggf uuuaaaagaa cuggggucucc aaauccuuua ggaacacca	2400
	guuuaaaca cccagcuuca gaaugagaaf cucuaccugu acuaucugca gaacggcagf	2460
	gacaugucagf uggaucaagg acuggacauc auucggcucu ccgacuaaga cguggaucau	2520
	aucgugcccc agucuuuuu caaagaugau uuuaaugua auaaaguuu gacaagauc	2580
	gaaaaaaau gagggaagag ugauaacguc cccucagaaag aaguuuguaa gaaaaugaaa	2640

Description	Sequence	SEQ ID NO
	<p>aauuuuuggc ggagcugcu gaacgccaaa cugaucacac aacggaaguu cgauaaucug 2700</p> <p>acuaaggcug aacgagugug cugucugag uuggauaaag ccggcuucau caaaaggcag 2760</p> <p>cuugugaga cagccagau caccaagcac guggcccaaa uucucgauc acgcaugaac 2820</p> <p>accaaguacg augaaaauga caaacugahu cgaagagguga aghuuuuac ucugaagucu 2880</p> <p>aagcugucuc cagauuucag aaaggacuuu caguuuuuaa agugagagaga gaucaacaaa 2940</p> <p>uaccaccaug cgaugaugc cuaccugaau cagugugug gacugcacu uaucaaaaaa 3000</p> <p>uaucccaagc uuuaaucuga auuuuuuac ggagacuaa aaguguaaga uuuuaggaaa 3060</p> <p>augaucgcaa agucugagca ggaauaggc aaggccaccg cuaaguacu uuuuacagc 3120</p> <p>auuuuuuga auuuuuucaa gaccgagahu acacuggcca auggagagau ucggaagcga 3180</p> <p>ccauuuacg aaacaaacgg agaaacagga gaaaucugug gggacaaggg uagggaauuc 3240</p> <p>gcgacagucc ggaagguccu guccaugccg caggugaaca ucguuuuuuu gaccgaagua 3300</p> <p>cagaccggag gcuuucccaa ggaauuauuc cucccgaaaa ggaacagcga caagcugauc 3360</p> <p>gcacgcaaaa aagauuugga ccccaagaaa uacggcggau ucgauucucc uacagucgcu 3420</p> <p>uacaguuac ugguuguggc caaaguggag aaagggaagu cuaaaaaacu caaaagcugc 3480</p> <p>aaggaaucg ugggcaucac aucauggag cgaucuaagc ucgaaaaaaa ccccaucgac 3540</p> <p>uuucucgag cgaaggauga uaaagagguc aaaaagacc ucaucauuua ccuucccaag 3600</p> <p>uacucucucu uugagcuuga aaacggccgg aaacgaaugc ucgcuagugc gggcggagcug 3660</p> <p>cagaagguu acgagcuggc acugccucu aaauacguu auuuuuugua ucuggccagc 3720</p> <p>cacuaugaaa agcucaaaag gucucccga gaaauagagc agaaagcagcu guucguggaa 3780</p> <p>caacacaaac acuaacuuuga ugagaucauc gagcaaaa gcaaaucuc caaaagagug 3840</p> <p>auccucgccc agcuuaaccu cgauaaggug cuuucugcuu acaauaagca cagggaauaag 3900</p> <p>cccaucaggg agcaggcaga aaacaauuac cacuuuuua cucugacca cuugggcccg 3960</p> <p>ccugcagccu ucaaguacuu cgacaccacc auagacagaa agcguuacac cucuacaaa 4020</p> <p>gaggucccug acgccaacac gauucaucag ucauuuacgg ggcucuuga acaagaauc 4080</p> <p>gaccucuc agcucgugg agacagcagg gcugacccca agaagaagag gaagguguga 4140</p>	
<p>CRISPR-associated endonuclease CAS9/Csn1 protein</p>	<p>MDKYSIGLDIGTNSVGWAVITDEYKVPKFKVLGNTDRHSIKKNLIGALLFDSGETAE</p> <p>ATRLKRTARRRYTRKNRI CYLQEIFSNEMAKVDDSFHRLSEFLVEEDKKHERHPFG</p> <p>NIVDEVAHYHEKYPTI YHLRKKLVSDTKADLRLLI YALAHMIKFRGHFLIEGDLNPDNSD</p> <p>VDKLFIQLVQTVNQLFEENPINASGVDAKAILSARLSKSRRLLENLIAQLPGEKKNGLFGN</p> <p>LIALSLGLTPNPKSNFDLAEDAKLQSKDLYDDLDNLLAQIGDQYADLFLAAKNLSDAI</p> <p>LLSDIILRVNFEITKAPLSASMIKRYDEHHQDLTLKALVRQQLPEKYKEIFFDQSKNGYA</p> <p>GYIDGGASQEEFYKFIKPI LEKMDGTEELLVKLNRELLRKRQTFDNGSI PHQIHLGELH</p> <p>AILRRQEDFYFLKDNREKIEKILTFRIYVYVGLARGNSRFAMTRKSEETITPWNFEE</p> <p>VVDKGSASQSFIERMTNFDKNLPNEKVLPKHSLLEYEYFTVYNELTKVYVTEGMRKPAFL</p> <p>SGEQKKAIVDLLLFTNRKVTVKQLKEDYFKKIECFDSVEISGVEDRNFNASLGTYHDLLEKI</p>	8

Description	Sequence	SEQ ID NO
<p>CRISPR-associated endonuclease CAS9/Csn1 mRNA</p>	<p>I KDKDFLDNEENEDI LEDIVLTLTFEDREMIERLKT YAHLFDDKVMQKLRRTYTGWG RLSRKLINGTRDKQSGKTI LDFLKSDGFANRNFMLI HDDSLTFKEDI QKAQVSGQDLSL HEHIANLAGSPAIKKGI LQTVKVVDELVKVMGRHKPENIV IEMARENETTQKQKNSRER MKRIEEGIKELGSOI LKEHPVENTQLEKLYLYLQNGRDMVYDQELDINRLSDYDVH IVPQSFLLKDDSIDNKVLT RSDKNRGSNDVPSEEVKKMKNYWFQLLNAKLITQRKFDNL TKAERGLSELDKAGFI KRQLVETRQITKHVAQI LDSRMNTKYDENDKLIREVKVIITLKS KLVSDFRKFQFYKVI RINNYHHADAYLNAVVG TALI KKYPKLESEFVYGDYKVIDVRK MIAKSEQEI GKATAKYFFYSNIMNFKTEI TLANGEIRKRPLIETNGETGEIVWDKGRDF ATVRKVLSPQVNI VKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFSDSPIVA YSVLVAKVEKSKKLSVKELLGITIMERSSEFKNPIDFLEAKGYEVKDDLI IKLPK YSLFELENGRKRMLASAGELQKGNELALPSKYVNFLYLASHYERLKGSPEDNEQKQLFVE QHKHYLDEII EQISEFSKRVI LADANLDKVL SAYNKHRDKPIRQAENI IHLFTLTNLGA PAAFKYFDTTIDRKRYTSTKEVLDATLIHQSI TGLYETRIDLSQLGGDSRADPKKKRKY</p>	<p>2</p>
	<p>auggacuaua aggacacaga cggagacuac aaggaucaug auuugauua caaagacgau 60 gacgauaaga uggcccaaaa gaagaagcgg aagguccgua uccacggagu cccagcagcc 120 gacaagaagu acagcaucgg ccuggacauc ggcaccaacu cugugggucg gcccugauc 180 accgacgagu acaagugcc cagcaagaaa uucaaggugc ugggaacac cgaccggcac 240 agcaucaaga agaaccugau cggagcccg cuguucgaca gcgggaaac agccgagcc 300 accggcuga agagaaccgc cagaagaaga uacaccagac ggaagaaccg gaucugcuau 360 cugcaagaga ucuuacgcaa cgagauggcc aagguggacg acagcuuuu ccacagacug 420 gaagaguccu uccuggugga agaggauaag aagcacgagc ggcaccccau cuucggcaac 480 aucguggacg aggucccuu ccacgagaag uacccacca ucuaccacu gagaaagaaa 540 cugguggaca gcaccgacaa ggcggaccug cggcugaucu aucuggcccu ggcaccacau 600 aucaaguucc ggggccacuu ccugaucgag ggcgaccuga accccgacaa cagcagcug 660 gacaagcugu ucauccagcu ggugcagacc uacaaccagc ugnucgagga aaccccauc 720 aacgccagcg gcgugfacg caaggccauc cugucugcca gacugagcaa gagcagacgg 780 cuggaaauc ugaucgcca gcugccggc gagaagaaga augccuguu cggcaaccu 840 auugcccu ga gcccacac gaccccaac uucaagagca acucgaccu ggcggagga 900 gccaaacugc agcugagcaa ggacaccuac gacgacgacc uggacaaccu gcuggccca 960 aucggcgacc aguaaccga ccuguuucug gccgccaaga accuguccga cgccaucucc 1020 cugagcgaca uccugagagu gaacaccgag aucaccaagg ccccccugag cgcucuaug 1080 aucaagagau acgacgagca ccaccaggac cugaccucgc ugaaagcucu cgugcgccag 1140 cagcugccug agaaauacaa agagaauuuc uucgaccaga gcaagaacgg cuaccggcg 1200 uacaauagc gcggagccag ccaggaaagag uucaucaagu ucaucaagcc cauccuggaa 1260 aagaugcag gcaccgagga acugcucug aagcugaaca gagaggaccu cugcgggaa 1320</p>	

Description	Sequence	SEQ ID NO
	cagcggaccu ucgacaacgg cagcauuccc caccagauc accugggaga gcugcacgcc	1380
	auucugcggc ggcaggaaga uuuuuaccca uuccugaagg acaaccggga aaagaucgag	1440
	aagaucuccga ccuucggcau cccuacua cccuacua cccuacua uggcagggg aaacagcaga	1500
	uucgcccuga ugaccagaaa gacgagaaa accauacccc ccuggaacuu cgaggaaug	1560
	guggacaagg gcgcuuccgc ccagaguu cccagucgga ugaccaacuu cgauaagaac	1620
	cugcccaacg agaagugcu gcccaagcac agccugcugu acgaguacuu caccguguan	1680
	aacgagcuga ccaaaugua auacugacc gagggaauga gaaagcccgc cuuccugagc	1740
	ggcggagcaga aaaaggccau cggggaccug cuguucaaga ccaaccggaa agugaccgug	1800
	aagcagcuga aagaggacua cuucaagaaa aucgagugcu ucgacuccgu ggaauucc	1860
	ggcgggaaag aucgguuca cgcuccccug ggcacauacc acgacucgcu gaaaauuac	1920
	aaggacaagg acuuuccuga caaugaggaa aacgaggaca uuucggaaaga uaucgugcug	1980
	accugacac uguuuagaga cagagagaug aucgaggaac ggcugaaaa cuaugcccac	2040
	cuguucgacg acaaaugau gaagcagcug aagcggcga gaaaccccg cuggggcagg	2100
	cugagccgga agcugaucua cggcauccgg gacaagcagu ccggcaagac auuccuggau	2160
	uuccugaagu ccgacggcuu cgcacaacaga acuucaugc acgugaacca cgacgacagc	2220
	cugaccuuua aagaggacau ccagaaaggc cagguguccg gccaggcga uagccugcac	2280
	gagcacauug ccauucggc cggcagccc gccauuaaga agggcauccu gcagacagug	2340
	aaggugugug acgagcucgu gaaagugaug gcccggcaca agcccagaa caucgugauc	2400
	gaauggcc aagagaacca gaccaccag aagggacaga agaacagccg cgagagaug	2460
	aagcggaucg aagaggcgu caaagagcug ggcagccaga uccugaaaga acaccggug	2520
	gaaaacacc agcugcagaa cgagaagcug uaccugua cuccugcaga ugggcccggau	2580
	auguacgugg accaggaa cuccagcua cggcuguccg acuaagcugu ggaccuuac	2640
	gugccucaga gcuuucugaa ggacgacuc aucgacaaca aggugcugac cagaagcgc	2700
	aagaaccggg gcaagagcga caacgugccc uccgaagagg ucgugaagaa gaugaagaac	2760
	uacugcggc agcugcugaa cgcgaagcug auuaccaga gaaaguuuca caaucgacc	2820
	aaggccgaga gaggcggccu gagcgaacug gaaagggccg gcuucauca gagacagcug	2880
	guggaaacc ggcaagacac aagcagcug gcacagaucc uggacuccc gaugaacacu	2940
	aaguacgacg agaaugacaa gcugaucggg gaagugaaag ugaucocccu gaaguccaa	3000
	cugguguccg auuucgggaa ggaauccag uuuuacaaa ugccggagau caacaacuac	3060
	caccacgcc acgacgccua ccugaacgcc gucgugggaa ccgcccugau caaaaauac	3120
	ccuaagcugg aaagcaguu cguuacggc gacuaacaag uguacfacgu gcggaagau	3180
	aucgccaaga gcgagcagga auccggcaag gcuaccgcca aguaucuu cuacagcaac	3240
	aucaugaacu uuuucaagac cgagauuacc cuggccaacg gcgagauccg gaagcggccu	3300
	cugaucgaga caaacggcga accggggag aucguguggg auaggggccg ggaauuucc	3360
	accgucgga aagucugag caugcccaac gugaauauc ugaaaaagac cgaggucgag	3420
	acaggcggcu ucagcaaga gcuauuccug ccaagaggga acagcagaua gcugaucgcc	3480

Description	Sequence	SEQ ID NO
<p>CRISPR-associated endonuclease CAS9/Csn1 protein</p>	<p>agaaagaagg acugggacc uaagaaguac ggcggcuucg acagcccccac cguggccuau 3540 ucugugcugg ugguggccaa aguggaaaa ggaagucca agaaacugaa gagugugaaa 3600 gagcugcugg ggaucaccu cauggaaaga agcagcuucg agaagaaucc caucgacuuu 3660 cuugaagcca agggcuacaa agaagugaaa aaggaccuga ucaucaagcu gccuaaguac 3720 ucccuguucg agcuggaaaa cggccggaag agaaugcugg ccucugccgg cgaacugcag 3780 aaggaaaacg aacugcccu gccuccaaa uaugugaacu uccuugaccu gcccagcccac 3840 uaugagaagc ugaaggcuc ccccgaggau aaugagcaga aacagcuguu uguggaaacag 3900 cacaagcacu accuggacga gaucaucgag cagaucagcg aguuuccaa gagagugauc 3960 cuggccgacg cuaaucugga caaugugcug uccgcuaca acaagcaccg ggaauagccc 4020 aucagagagc agcccgagaa naucauccac cuguuuacc ugaccaauu cggagccccu 4080 gccgcuuca aguacuuuga caccaccauc gaccggaaga gguacaccag caccaaaag 4140 gugcuggacg ccaccugau ccaccagagc aucaccggcc uguacgagac acggauccgac 4200 cugucucagc ugggagcga caagcuccu gcugcuacua agaaagcugg ucaagcuaag 4260 aaaaaagaaa</p>	<p>3540 3600 3660 3720 3780 3840 3900 3960 4020 4080 4140 4200 4260 4269</p>
<p>MDYKDHGDYKDHDI DYKDDDDKMA PKKRKVG I HGVPAADKKYSIGLIDIGTNSVGVAVI TDEYKVP SKFKVLGNTDRHS IKKNLIGALLFDSGETAEATRLKFTARRRYTRRKNRICY LQEIFSNEMAKVDDSFHRLSEEFV EEDKKHERHP IFGNIVDEVA YHEKYPTIYHLRKK LVDSTDKADLR LIY LALAHMIKFRGHFI EGDLPDNDSDVKLF IQLVQTYNQLFEENPI NASGVDAKA ILSARLSKRRLENLI AQLPGEKKNGLFGNLIALSLGLTPNFKSNFDLAE AKLQLSKDTYDDDLNLLAQIGDQYADLFLAAKNLSDA ILLSDILRVNTEITKAPLSASM IKRYDEHHODLTLKALVRQQLPEKYKEIFFDQSKNGYAGYIDGGASQEEFYKFIKPILE KMDGTEELLVKLNREDLLRKQRTFDNGS IPHQIHLGELHAILRRQEDFY PFLKDNREKIE KILTFRIPIYVGPLARGNSRFAMTRKSEETI TPWNFEVVDKGASAQS FIERMTNFDKN LPNEKVLPHSLLYEYFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVD LLLFKTNRKVTV KQLKEDYFKKIECFDSVEISGVEDRFNASLGT YHDL LKIIKDKDFLDNEENEDILEDIVL TLTLFEDREMI EERLKT YAHLFDDKVMKQLKRRRYTGWRLSRKRLINGIRDKQSGKTILD FLKSDGFANRFMQLIHDDSLTFKEDIQAQVSGQDLSHEHIANLAGSPA IKGILQTV KVVDLVKVMGRHKPENIV IEMARENQTQKGOKNSRERMKRIEIGIKELGSQLKEHPV ENTQLQNEKLYLYLQNGRDMYVDQELDINRLSDYVDHI VPOSF LKDDSDI DNKVLTRSD KNRGKSDNVPSEEVVKKMKNYWRQLNAKLI TQRKFDNLTKAERGLSELDKAGFIKRQL VETROI TKHVAQI LLSRMTKYDENDKL IREVKVITL KSKLVDFRKFDFYK VREINNY HHADAYLNAVGTALIKKYPKLESEFVYGDYKVDYDRKMI AKSEQEI GKATAKYFFYSN IMNFFKTEITLANGEIRKRPLIETNGETGEI VWDKGRDFA TVRKVLMPQVNI VVKTEVQ TGGFSKESILPKRNSDKLIARKKDWDPKYG GFDSP TVAYSVLVVAKVEK GKSKKLSVK ELLGITIMERSSEKNPIDFLEAKGYKEVKKDLI IKLPKYSLFELENGRKRMLASAGELQ</p>	<p>9</p>	

Description	Sequence	SEQ ID NO
CRISPR-associated endonuclease CAS9/Csn1 mRNA	<p>KGNELALPSKYVNFVLYLASHYEKLGKSPEDNEQKQLFVEQHKHYLDEIIEQISEFSKRVI LADANLDKVLISAYNKHRDKPIREQAENIHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKE VLDATLIHQSTGLYETRIDLSQLGGDKRPAATKKAGQAKKKK</p> <p>auggacaaga aguaacagcau cggccuggac aucggcacca acucuguggg cugggcccug 60 aucaccgacg aguaacaagg gcccagcaag aaauucaagg ugcugggcaa caccgaccgg 120 cacagcauca agaagaaccu gaucggcgc cugcuguucg acagcgagga acagccggag 180 gccaccggc ugaagagaac cgccagaaga agauacacca gacggaagaa cgggaucugc 240 uaucugcaag agaucuucag caacgagau gccaaggugg acgacagcuu cuuccacaga 300 cuggaagagu ccuuccuggu ggaagaggu aagaagcacg agcggcacc caucuucggc 360 aacaucgug acgaguggc cuaccacgag aaguaaccca ccaucuacca ccugagaaaag 420 aaacuggug acagcaccga caaggccgac cugcggcuga ucuaucuggc ccuggccccac 480 augaucaagu uccggggcca cuuccugauc gaggcggacc ugaaccccga caacagcgac 540 guggacaagc uhuucaucca gcuggugcag accuacaacc agcuguucga ggaaaacccc 600 aucaacgcca cggcggugga gcccaggcc auccugucg ccagacugag caagagcaga 660 cggcuggaaa aucugaucg ccagcugccc ggcgagaaga agaauggccu uuucggcaac 720 cugauugccc ugagccuggg ccugaccccc aacuucaaga gcaacuucga ccuggcccag 780 gaugccaaac ugcagcugag caaggacacc uacgacgacg accuggacaa ccugcuggcc 840 cagaucggcg accagucgc gcaccuguuu cuggccgcca agaaccuguc cgacgccauc 900 cugcugagcg acauccugag aguaacacc gagaucacca aggcccccuc gagcggccuc 960 augaucaaga gauacgacga gccaccacg gaccugacc ucugaaagc ucucgugcgg 1020 cagcagcugc cugagaagua caaagagau uucuuacgacc agagcaagaa cggcuacgcc 1080 ggcuaaucg auggcggagc cagccaggaa gaguucaaca aguucauca gcccauccug 1140 gaaaagaug acggcacccga ggaacugcuc gugaagcuga acagagagga ccugcugcgg 1200 aagcagcgg ccuucgacaa cggcagcauc cccaccacga uccaccuggg agagcugcac 1260 gccauucug gccggcagga agauuuuac ccauuucuga aggacaaccg ggaaaagauc 1320 gagaagauc ugaccuuccg caucccuac uacgugggcc cucuggccag gggaaacagc 1380 agauucgcu ggaugaccag aaagagcag gaaaccauca ccccuuggaa cuucgaggaa 1440 guggugaca agggcgccag cgcccagagc uucaucgagc ggaugaccaa cuucgaaag 1500 aaccugcca acgagaaggu gcugcccaag cacagccugc uguacgagua cuucaccgug 1560 uacaacgagc ugaccaaaagu gaaauacgug accgagggaa ugagaaaagcc cgccuuccug 1620 agcggcgagc agaaaaagc caucguggac cugcuguuca agaccaccg gaaagugacc 1680 gugaagcagc ugaagagga cuacucaa gaaaucgagu gcuucgacuc cguggaauc 1740 uccggcgugg aagauccgguu caacgccucc cugggcacau accagcauc gcugaaaau 1800 aucaagaca aggacuuccu ggacaauag gaaaacgagg acauucugga agauaucgug 1860 cugaccucca cacuguuuga ggacagagag augaucgagg aaccgucgaa aaccuaugcc 1920</p>	3

Description	Sequence	SEQ ID NO
	caccugnuccg acgacaaaagu gaugaagcag cugaagcggc ggagauaacac cggcuggggc	1980
	aggcugagcc ggaagcugau caacggcauc cgggacaagc agucccgcaa gacaauccug	2040
	gauuuccuga aguccgacgg cuucgccaac agaaacuua ugcagcugau ccacgacgac	2100
	agccugaccu uuaaagagga cauccagaaa gcccaggugu ccggccaggg cgaugccug	2160
	cacgagcaca uugccaauu ggccggcagc cccgccaaua agaagggcau ccugcagaca	2220
	gugaagggug uggacagacu cgugaaaug augggcccgc acaagcccga gaacaucug	2280
	aucgaaaugg ccagagagaa ccagaccacc cagaaggagc agaagaacag ccgagagaga	2340
	augaaaggga ucgaaagagg caucaaag cugggcagcc agauccugaa agaaccacc	2400
	guggaaaaca cccagcugca gaacgagaag cuhuaccugu acuaaccugca gauggggcg	2460
	gauauguacg uggaccagga acuggacauc aaccggcugu ccgacuacga uugggaccuu	2520
	aucgugccuc agagcuuucu gaaggacgac uccaucgaa acaaaugucu gacucggagc	2580
	gacaagaacc ggggcaagag cgacaacgug ccuccggaag aggucgugaa gaagaugaa	2640
	aacuacuggc gccagcugcu gaugccaag cugauuacc agaggaguu cgacaauug	2700
	accaagggcg agagggcgg ccugagcga cuggauaagg ccggcuuucuu caagagacag	2760
	cugugugaaa cccgacagau caaaagcac guggcacaga uccuggacuc ccggaugaac	2820
	acuaaguacg ccgagaacga caacugauc cgggaaguga aagugaucac ccugaagucc	2880
	aagcuguguu ccgahuuucc gaaggauuuc caguuuuaca aagugcgga gaucaacaac	2940
	uaccaccag cccacgacgc cuaccugaac gccugcugug gaaccgccc gaucaaaaag	3000
	uaccuuagc uggaaagcga guucgugac ggcgacuaca agguhuacga cugcgggaag	3060
	augaucgcca agagcgagca ggaauccggc aaggcuaccg ccaaguauuu cuucuacagc	3120
	aaacauaga acuuuuucaa gcccgaguu acccuggcca accggcgagau ccggaagcgg	3180
	ccucugaucg agacaacgg cgaaacaggc gagaucgugu gggauaaggf ccgggacuuu	3240
	gccaccgugc ggaagugcu gucuauccc caagugaaua ucgugaaaaa gaccgaggug	3300
	cagacagggc gcuuacagcaa agagucuauc cugcccaga ggaacagcga caagcugauc	3360
	gccagaaaag aggacuggga ccuaaagaag uacggcgccu ucgacagccc caccguggcc	3420
	uuuucuguc uggugggugc caaaguggaa aagggcaagu ccaagaaaacu gaagagugug	3480
	aaagagcugc uggggaucau caucauggaa agaagcagcu ucgagaagaa ucccuaugc	3540
	uuucuggaag ccaagggcua caaagaagug aaaaaggacc ugaucuucaa gcugccuaag	3600
	uacucccugu ucgagcugga aaacggccgg aagagaaugc uggccucugc cggcgaacug	3660
	cagaaggga acgaaacugc ccugccucc aaauauguga acuuuccugua ccuggcccgc	3720
	cacuaugaga agcugaaggg cucccccgag gaaauaugagc agaaacagcu guuugggaa	3780
	cagcacaac acuaaccugga cgagaucalc gagcagauc gcgaguucuc caagagagug	3840
	auccugggcg agcuaaucu ggacaagug cuhagcgcuu caacaagca cagagacaag	3900
	ccuaucagag agcagggcga gaaaucauc caccuguuuu ccugagccaa ucuggggacc	3960
	ccugccgccc ucaaguacuu ugacaccacc accugaccga agagguaacac cagcaccaaa	4020
	gaggugcugc acgccaaccu gauccaccag agcauaccgg gccuuguaacg gacacggauc	4080

Description	Sequence	SEQ ID NO
	gaccuguc agcugggagg cgagccuau ccuauagacg ugcccgaaua ugccagccug ggcagcggcu ccccacaaga aaacgcaag guggaagauc cuaagaaaaa gcggaaaagug gacggcauug guagugggag caacggcagc agcggauccg ugagcaaggg cgaggagcug uacaccgggg ugugcccdu ccugucgag cuggacggcg acguaaacgg ccacaaguuc agcugcgcg ggcagggcga gggcgaugcc accaacggca agcugaccuu gaaguucauc ugcaaccacc gcaagcugcc cgugcccug cccaccucg ugaccauccu gaccuacggc gugcagugcu ucagccgcu cccgaccac augaagcagc acgacuucuu caaguccgcc augcccgaag gcuacgucca ggagcgcacc aucucuuca aggacgacgg caccuacaag acccgcgcc aggugaaguu cgagggcag acccuguga accgcaucga gcugaaagggc aucgacuua aggagjacgg caacauccug gggcacaagc uggaguacaa cuucaacagc cacaaagucu auaucaaggc cgacaagcag aagaacggca ucaaggcgaa cuucaagauc cgccacaacg ucgaggjacgg cagcgugcag cucgcccacc acuaaccagc gaacaccccc aucggcgacg gcccgcugcu gcugcccagc aaccacuacc ugagcaccga guccaagcug agcaaaagacc ccaacgagaa gcgcgaucac augguccugc uggaguucgu gaccgcccgc gggaucacuc ucggcaugga cgagcugnac aagua	4140 4200 4260 4320 4380 4440 4500 4560 4620 4680 4740 4800 4860 4920 4956
CRISPR-associated endonuclease CAS9/Csn1 mRNA	ugaucgcaa gucggaacag gaaucggaa aggcuaucgc aaaguacuuc uucaucuaa acaucauga cuucuuaaa acggagauc gcucgcgaa cggcgaauuc cggaaaagc cgcucauga aaccacgga gaaaccggg agaucgugug ggacaagggg agggauuuug cgacugugag gaaguguuug uccaugccgc agucaauuu uguaaaaaag acggaaguc aaaccggagg auucaagcaaa gaaucuauc ucccagaagc caacucggac aaacuaucg cgcgcaagaa ggaugggac cccaagaaau acggugggcu ugacagcccu acuguggcu acuccgucu cgucgugcu aaaguggaa aggguaaauc caaaaagcuc aaucgguga aggagcuccu gggaucacg aucauggagc gducgagcuu gaaaaagaa ccuaugahu uccuggagg gaaggcuac aggaaguc aaaaagaccu gaucaucag cucccgaagu acagccuuu cgagcugaa acggcagaa agaggauugc ggcaucagcg ggagaauugc agaagggaaa cgaacuggca cugccgucca aguacgugaa uuuuucuaug cuggcuagcc acuaagaaa gcugaagga ucgcccggg caacagagca aaaaacug uucguggagc agcacaagca cuaccugau gagaucucg agcagaucuc cgaaucucg aaacgcguga uccuugccga ugccaucug gaaaguguu ugucggcuua caacaagcau cgggaaaaac cgaucgcga acagcgagaa acaucauuc aucuguauc uuugaccaau cugggagcgc cugccgcguu uaaguacuuc gacaccua uugauagaaa gcgcuacacc ucgaccaagg aagugcugga cgcuaaccug auccaccagu ccaucaccgg acucacgaa acucgcaug accugucca gcuhggagga guuacaggg ccgauccaaa gaaaaagcgc aagucugau aaagggcugg agccucggug gccaugcuuc uuugccuuug ggcucccccc cagcccccc uccccuuucc gaccccguac ccccgguguc uuugaauaaa gucugagugg cgggcuucag	60 120 180 240 300 360 420 480 540 600 660 720 780 840 900 960 1020 1080 1140 1200

Description	Sequence	SEQ ID NO
	<p>a</p> <p>1201</p>	
CRISPR-associated endonuclease CAS9/Csn1 mRNA	<p>ucaaagcuuuu ggaccucug acagaagcua auacgacuca cuuaagggaa auuagagaga 60</p> <p>aaagaagagu aagaagaaau aaagagcca ccauggacaa gaaauacagc auaggccug 120</p> <p>auauuggaac uaacagcguu ggauggcgag ugaucaccga cgaguaacaag gugccgagca 180</p> <p>agaaguuaaa ggugcugggg aacacugacc gccauaaau uaagaaaaa cucauuggag 240</p> <p>cacugcuuuu ugacucgggu gagacugccg agcuaaccag gcuaaaaaa accgcacgca 300</p> <p>gacgguacac ccgccgcaag aaucgcauc gcuauucgca agagauuuu uccaacgaga 360</p> <p>uggcgaaggu ugacgacagc uuuuuacc ccugugaa gaagauucg guggaagagg 420</p> <p>acaaaaagca cgaagggcau ccaaucuucg guaacaucgu ggacgaagug gcguaucacg 480</p> <p>aaaaauacc uaccaucuac caucugcga aagaugcugu cgaauccacg gaaaggcag 540</p> <p>accugagacu gaucuaccug gcuuuggccc auaugaucaa auuccgggc cauuuccuga 600</p> <p>ucgaggggga ccuaaaccg gaaacucgg auucgacaa gcuguucauc cagcuggucc 660</p> <p>aaacguauaa ccaacuguuu gaggaauuuc ccaucaaccg uucgggggug gacgcaaaag 720</p> <p>caaucucuc cgcgcggcug agcaagucac ggcggcucga aaaccugauc gcgcagcugc 780</p> <p>cgggagaaa gaaaaaugga cuguuuggga aucugaucg ccugucguc gcucugacuc 840</p> <p>caaacuuuaa gucaaaauuc guuuuggccg aagaugcaca gcugcagcug ucaaaaggaca 900</p> <p>cuaacgacga cgaucuggac aaucugcug cccagaauug ggaccaauc gcagaccug 960</p> <p>ucuuuggccg gaagaaccug agcgacgca uucucuguc cgaauucug agagucaaua 1020</p> <p>ccgaaauac naagcucug cuguccgcu caaugaucaa gcgcuacgcu gaacaccacc 1080</p> <p>aggauucac ucugcucaaa gccucugga gacaacaau gccugaaaa uacaaggaga 1140</p> <p>ucuuucuga ccagagcaaa accggcuacg caggcuacau cgauggagga gcgucacaag 1200</p> <p>aagaguuaa caaguucauc aagccaauc uggagaagau ggacgguacu gaagaacuc 1260</p> <p>uugugaagcu gaauagggag gaaugcuca gaaagcagcg gachuugac accggcuga 1320</p> <p>ucccucaua gaaucaccuc ggugagcug augccaucuu ucggcgccaa gaggauuuu 1380</p> <p>acccuuuccu gaaggauaa cgcgagaaa ucgaaaagau ccugacguc agaauuccu 1440</p> <p>acuaacgugg accgcuggcg cgcgguaacu cgcgguuugc auggauacu cgcaagucag 1500</p> <p>aggaaacua cacuccuug cauuugagg aggucgucga uaaggagcc uccgcccagu 1560</p> <p>cauucaucga acgcaugacc acuucgaca agaaucuucc gaacgagaag guccuuccaa 1620</p> <p>agcacuccu guuuaacgaa uacuuaccg uguacaaua gcugaccaaa guuaaguau 1680</p> <p>ucaccgagg caugdaaag ccggccuucc ucagcggcga acaaaagaag gccaucgucg 1740</p> <p>accuccuuu caagaccaac cggaaagguga ccgucaagca acucaaggag gacuacuuca 1800</p> <p>agaagaucga augcuugac ucggucgaaa uacagcggagu ggaggaccgg uuuaacgcu 1860</p> <p>cacuggguac cuaccaugau cuccugaaaa ucaucaaaaga caaggauuc cuggacaacg 1920</p> <p>aaagaaaacga ggacaucucg gaagauuug uguugaccuu gacgcuugc gaggaccggg 1980</p> <p>aaauauguca ggaaggcuu agaccuucg cacaccucu cgaugacaaa gugaugagc 2040</p>	5

Description	Sequence	SEQ ID NO
	aacugaagcg gcggaauuu acuggcuggg ggaggcucuc ccggaagcuc auuauggaa 2100 ucagagacaa acagucgggu aaacuauc ucgacuuccu caagucggau gggucgcca 2160 accggaacuu caugcagcug uccacgaug auuccuugac cuucaaggaa gauuuccaga 2220 aggcgcaagu gagcgacag ggagauucgu ugcacgaaca uaucguuuu cugcccgga 2280 cccagccau caagaaagga uccucgaga cgguaaggu ggugaugaa cuggugaaa 2340 ugauggggc ccacaaacca gagaaacag ucauugagau gcccgcgag aucagacca 2400 cucagaagg acaaaagaac uccagagagc ggaugaaagc caucgagaa ggcaucaaag 2460 agcuugguag ccaauuccu aggaacacc cggucgagaa caccaagc cagaacgaa 2520 agcuuuaccu guacuaccu caaauggac gggacaugua cguccgacc gaauggaca 2580 ucaacagacu cagcgaucac gaugggacc auuuuggcc acaguccuu cuuaaggac 2640 acagcaucga uacaaagug cucacuagau cagacaaaa ucgcgggaaa ucagacaaug 2700 ugccaucgga agagguuugc aagaagaua aaacuacug gagacagcug cucaaugcca 2760 aacuuuacac ccagcggag uucgacaacc uuaccaaggc cgagcggga ggaauugcc 2820 aacucgacaa ggcggcuuc ucaaaaggc agcuguguga aaccggcag aucacuaaac 2880 acguggccca gaucucgau ucgcgcauga acacuaaaua cgaugagaa gacaagcuga 2940 uuagggaaug caaggucauc acucugaagu cgaaacuggu ucgugacuuu agaaaggau 3000 uccaguucua caaaguccgc gaguuuaaca acuaccacca cgcucaugac gccuaccuga 3060 augcaguugu gggcaccgc cugaucaaga aguaucgaa gcuggaaucc gaguucgugu 3120 acggagaaua caaaguuac gacgucgca aga 3153	
CRISPR-associated endonuclease CAS9/Csn1 mRNA	ucaagcuuu ggaccucgu acagaagcua auacgacua cuauaggaa auaaagaga 60 aaagaagagu aagaagaaau auaagagcca ccauggacua caaggaccac gacggagacu 120 acaagacca ugacuagau uacaaggaug acgaugacaa auuggaccg agaagaaga 180 gaaagucgg auuacacggg gugccggccg cggacaagaa guacucauc ggacuggaa 240 ucggcacgaa cagcuggggu ugggcaguga uaccgacga auacaaggug ccgagcaaga 300 aguucaaaug ucugggaaau accgaucgc auucgacaa gaaaaucug auuggcgc 360 uccuguucga cucgggagag acugccgagg ccacuagacu gaagagacc gcuaggcgc 420 gcuaacagag gcgcaaaac cgaucugcu auuucaaga auuuuca aacgagau 480 ccaaggugga cgacuucuu uucaucggc uggaagaauc auuuugug gaggagaca 540 agaagcacga acgcauucc auuuucgca acauugcga cgaagugcc uaucauga 600 aguaucgac uaucuaccac uugagaaaga agcugguga cuccacugac aaggcagau 660 ugcgguugau cuaccugca cuggcccau ugaucuuuu cgggggac uuccucauc 720 agggcgaccu uauuccgac auuuucgug uggauaagcu uucauccag cugguccaga 780 ccuaacacca acuguuugaa gaaaaucca ucaauvcgag cggugucgau gcaagacca 840 uccugagcg ccgcuucg aaaagcaga ggcucgaaa ccugaucgca caguugccug 900 gagagaaga gaacggcuc uucggcauuc ucauvcgaa uucccugga cugacuccaa 960	6

Description	Sequence	SEQ ID NO
	acuucaaac caacuucgac uuggccgagg uucugggcuc agcuccaaacu gcaacugagc aaagauaccu	1020
	acgaugauga cuuuggacaau cuucuggcuc agaucggcga ccaghuacgcc gaccugnucc	1080
	uugcggcuua gaaccugucg gacggcaucc ugcuguccga cauccugcg gucaauaccg	1140
	aaauuacuaa agcaccacuc ucggcaucca ugaucgaagc auacgaugaa caccaccagg	1200
	auucacaccu ccugaaagca cuggugcggc agcagcuccc ugagaaaauac aggaaaauu	1260
	ucuuuugauc guccaagaac ggaucgccc ggaucacga cggcgcgcg agccaagg	1320
	aaucuacaa guucaucaag ccgauccugg aaaagaugga uggcacggaa gaacucugg	1380
	ucaaacugaa uagagaggau cugcuccga acaacggac cuucgaaac ggaucgacc	1440
	cgcaccagau ccaccucggc gaacugcag gaaaagauc ccauccugcg gcggcaggag	1500
	cguuccucaa agacaacaga gaaaagauc gaaagauuu gaccuuucg aucccguauc	1560
	acguggggcc gcucgagaga gguaacucc gcuuugcuug gaaagacuga agucagagg	1620
	aaacgaucac cccauggaac uucgaaagg ucgugacuu uggugacaa agggcgagc gcccaucgu	1680
	ucaucgaacg gaugacuac uucgaaaga aucugccgaa ugagaaaggc cugccuaagc	1740
	acucacuuc guaugaauc uuacugugc uuacccugac cccaaaguc aaauacguga	1800
	cugagggaau gcgcaagccu gcguuuuugu ccggcgagca gaaaaggcc aucguggacu	1860
	ugcuguucaa acccaaccgc aggugacug uuaagcaacu caaagaggac uacuuuaaga	1920
	agaucgaaug cuuugacuc gucgagauu ccgggguuu gaaluaagau acgcgucgc	1980
	ugggaaccua ccaugauc cucaagaua ucaaggacaa agacuuccug gaaacgagg	2040
	agaaugagga cauccucgaa gauuugugc uuacccugac ccuuuucgaa gaucgcaaa	2100
	ugaucgaa gaagggaaa accuacguc accuugucga cgaucggug augaaacagu	2160
	ugaaacgccc gcgguacac gguuggggc accauccugc ggcugucgag caagcugauc aacggaauc	2220
	gggacaaa gaagcggaaag accauccugc auuuucugaa guccgauggu uuugccaacc	2280
	gcaacuuc auagcucauc caugacgahu cgcuuaccuu uaaggaggau uaccgaaagg	2340
	cacaagugc gggacaagg gaaucguc accgaacacu cgcacaucug gcggggucgc	2400
	cggcaauuaa gaagggaauc caccagacug uuaagggugu cgacgagcug gugaagguga	2460
	uggggagaca uaagccugaa acauuguga ucgagaucc gagagaaaa caaacuacuc	2520
	agaagggaca gaagaaucc cgggagcggg ugaagcgcau cgaggaggga aucaaggaa	2580
	ugggcuccca auuccugaaa gagcaucccg uggaaaauac ucagcugcag aacgagaagc	2640
	uuuaccugua cuauuucua auuggcagg acauguaucg cgaacaagaa cuggauaua	2700
	aucggcuc cgaauacgac gucgacaca ucgucuccca aucauuucug aggaaugaa	2760
	gcaucgaa caagguucuc acuagaucag acaaaaaccg gggaaaagaa gaaacguc	2820
	ccagcgaaga agucugaa agaugaaaga auuacuggag gcaacuucug aacgcaaac	2880
	ucaucacua cgcgaaguu gacaaccuga ccaagcaga agggggagga cucagcagc	2940
	uggacaaggc ugguuucauc aaacggcagc uggugagac ucgcaaauc acgaagcaug	3000
	uggcccagau ucucgacuc cgaugaaua cuagucga cgaaaaacgaa agcugauc	3060
	gggaggugaa ggugaucacc cucaagagca agcucgugc cgauuuuccgg aaagacuucc	3120

Description	Sequence	SEQ ID NO
	<p> aguuuacaa ggugcgggag auuaacaacu accaucaacgc ucacgaagcgu uaccucaaag 3180 cugugggugg gacggcguug auuaagaagu acccaaaaacu ggaguccgaa uucgucuaag 3240 gagauuacaa ggucuaacgac gucgcgaaga ugaauGCCaa gucgggacag gaaauuggga 3300 aagcgaacgc uaaguuacuuc uucuaucuga auaucaugaa cuucuucaag accgaaauca 3360 ccucggcuua cggcgaagau aggaacacgc cgcuaucuga aacuaauggu gagacuggug 3420 aaucugugug ggauaaaggga cgggacuucg ccacggucucc caagtuuccc agaucgccc 3480 aagugaaau uguuaagaaa accgaaaguc agaccggugg guucucgaag gaaucucc 3540 ugccaaagc caaucggau aagcuuauug cccgcaagaa ggauugggac ccgaaaaagu 3600 acgguggguu cgacuccccu accgugcggu acucgguguu ggugugggcc aaaguggaaa 3660 agggcaaac aaagaagcuc aagagcguca aggagcugcu gggaucacc aucaungaga 3720 gguccagcuu ugagaaaaac ccgaucgacu ucuuugaagc caaggauac aaagagguga 3780 agaaagaccu gaucauacaa cuuccaaaagu acucccuguu cgaacucgaa acggggagg 3840 agcgaugcu cgcucagcc ggggaacugc aaaagggcaa cgaacuggcc cucccguc 3900 aaucgucuaa cuucugtuac uggcgucac acuaacaaaa gcugaaagg gacccagagg 3960 acaacgaaca gaacagcug uucgucgagc agcaacaa gcaucucgag gagaucauc 4020 aacagauc ggaauucagc agagaguga ucuuugcaga cgcuaaccuu gacaaagucc 4080 ucucggcau cauaaagcau cgcgacaagc cgaucagaga acagggcgag acaucaucc 4140 accuguaac ucucaccaac cugggcgcgc cagcgcuuu uaaguacuuu gauaccacca 4200 uugaccgcaa gagauacacc ucaacuaaag aagucugga cgcaccuccg auccaucaaa 4260 gcauaccgg acuuuaugaa acucggauc aucucucaca gcucggaggga gacaaaagac 4320 cggcugccc caagaaggcc ggacaggcaa agaagaagaa augaauaag gcuggagccu 4380 cgguggccau gcuucugcc cuuugggcu cccccagcc cuccucccc uuccugcacc 4440 cguaccccc uggucuuuga auuaagucug aguggcgcc ucuaga 4486 </p>	
dCAS9 as described in Qi et al	<p> MDKKYSIGLAIGTNSVGWAVITDEYKVPKFKVLGNTDRHSIKKNLIGALLFDSGETAETRLKRTARRRYTRKRNRCYLQELIF SNEMAKVDDSFHRLSESLVEEDKKHERHPIFGNIVDEVAYHEKYPTIYHLRKKLVDSTDKADLRLIYLALAHMIKFRGHFLIEG DLNPDNSDVKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRRENLIQAOLPGEKKNLGNLIALSLGLTPNFKSNFDL AEDAKLQSKDTYDDDLNLLAQIGDQYADFLAAKNLSDAILLSDILRYNTEITKAPLSASMIKRYDEHHQDLTLLKALVRQQLP EKYKEIFFDQSKNGYAGYIDGASQEEFYKFIKPILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHGELHAILRRQEDFY PFLKDNREKIEKILTRIPYVVGPLARGNSRFAMTRKSEETITPWNFEVVVDKASAQSAQSFIERMTNFDKNLPNEKVLPKHSLLYE YFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNKVTVKQKEDYFKKIECFDVSVEISGVEDRFRNASLGTYHDLKLIK DKDFLDNEENEDILEDIVLTLTFEDREMIEERLKYAHLFDDKVMKQLKRRRYTGWRLSRKLLINGIRDKQSGKTIIDFLKSDGF ANRNFMLIHDDSLTFKEDIQKAQVSGQDSLHEHIANLAGSPAIKKGIQTVKVVDELVKVMGRHKPENIVEMARENQTTQKQ KNSRMRKRIEEGIKELGSOILKEHPVENTQLQEKLYLYLQNGRDMYDQELDINRLSDYDVAIVPQSFLKDDSIDNKLITRS DKNRGKSDNVPSEEVVKMKNYWRQLLNAKLITQRKFDNLTKAERGGSLDKAGFIKRLQVETRQITKHVAQIILDSRMNTKYDEN DKLIREVKVITLKSILVSDFRKDFQFYKVIKREINNYHHAHDAYLNAVVGVALIKKYPKLESEFVYGDYKVDVVRKMIKSEQEI GKA </p>	61

Description	Sequence	SEQ ID NO
<p>As describe in Gilbert et al 2013. Amino acid sequence of dCas9-KRAB (dCas9- HA tag- 2xSV40NLS- KRAB)</p>	<p>TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFAVRKVL SMPQVNI VVKTEVQTGGF SKESILPKRNSDK LIARKKDWDPKKYGGFSDPTVAYSVLVAVAKVEGSKKLSVKELLGITIMERSSEFEKNPIDFLEAKGYKEVKKDLI IKLPKYSLF ELENGRKRMLASAGELQKGNELALPSKYVNFYLASHYKELKSGPEDNEQKQFLVEQHKHYLDEIIEQISEFSKRVILADANLKV LSAYNKHRRDKPIREQAENI IHLFTLTNLGAPAAFKYFDTTIDRKRYTSKTEKVEDATLIHQSI TGLYETRIDLSQLGGD</p> <p>MDKKYSIGLAIGTNSVGVAVITDEYKVPSSKFKVLTGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIF SNEMAKVDDSFHRLSEFLVEEDKKHERHPIFGNI VDEVAYHEKYPTIYHLRKKLVSDTKADLRLIYLALAHMIKFRGHFLIEG DLNPDNSDVDKLFIQLVQTYNQLFEENPINASGYDAKAL SARLSKSRRLLENLIAQLPGEKKNGLFNGLIALSGLT PNFKSNFDL AEDAKLQLSKDTYDDDLNLLAQIGDQYADLF LAAKNLSDAI LLSDI LRVNTETKAPLSASMIKRYDEHHQDLTLLKALVRQQLP EKYKEIFFDQSKNGYAGIDGGASQEEFYKFKP ILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFY PFLKDNREKIEKILTFRIPYVGPLARGNSRFAMTRKSEETITPWNFEVVYDVGASQAQSFIERMTNFDKNLPNEKVLPKHSLLYE YFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQKEDYFKKIECFDSVEISGVEDRFNASLGTYHDDLKIIK DKDFLDNEENEDILEDIVLTLTFEDREMIEERLKYAHLFDDKVMKQKRRRYTGWGRLSRKLINGIRDKQSGKTI LDFLKSDGF ANRNFMLIHDDSLTFKEDIQKAQVSGQDLSHEHIANLAGSPA I KKGILQTVKVVDELVKVMGRHKPENIVIEMARENQTTQKQK QNSRERMKRIEEGIKELGSOI LKEHPVENTQONEKLYLYLQNGRDMYDQELDINRLSDYDVAIVPQSF LKDDSIDNKNVITRS DKNRGKSDNVPSEEVVKKMKNYWRQLLNAKLI TQPKFDNLTKAERGGLSELDKAGFIKQQLVETROI TKHVAQI LDSRMNTKYDEN DKLIREVKVI TLKSKLVSDFRKFDFQYKVVREINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGYDVKVYDVRKMIKSEQEIGKA TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFAVRKVL SMPQVNI VVKTEVQTGGF SKESILPKRNSDK LIARKKDWDPKKYGGFSDPTVAYSVLVAVAKVEGSKKLSVKELLGITIMERSSEFEKNPIDFLEAKGYKEVKKDLI IKLPKYSLF ELENGRKRMLASAGELQKGNELALPSKYVNFYLASHYKELKSGPEDNEQKQFLVEQHKHYLDEIIEQISEFSKRVILADANLKV LSAYNKHRRDKPIREQAENI IHLFTLTNLGAPAAFKYFDTTIDRKRYTSKTEKVEDATLIHQSI TGLYETRIDLSQLGGDAYPYDVPD YASLGSGPSKKRKRKVEDPKKRKVDGIGSGSNGSGGGGMDAKSLTAWSR TLVTFKDV FVDFTR EEWKLLD TAQQIVYRNVMLE NYKNLVS LGYQLTKPVDVILRLEKGEEP</p>	<p>62</p>
<p>Amino acid sequence of dCas9-BFP- KRAB ((dCas9- HA tag- 2xSV40NLS- BFP- KRAB)</p>	<p>MDKKYSIGLAIGTNSVGVAVITDEYKVPSSKFKVLTGNTDRHSIKKNLIGALLFDSGETAEATRLKRTARRRYTRRKNRICYLQEIF SNEMAKVDDSFHRLSEFLVEEDKKHERHPIFGNI VDEVAYHEKYPTIYHLRKKLVSDTKADLRLIYLALAHMIKFRGHFLIEG DLNPDNSDVDKLFIQLVQTYNQLFEENPINASGYDAKAL SARLSKSRRLLENLIAQLPGEKKNGLFNGLIALSGLT PNFKSNFDL AEDAKLQLSKDTYDDDLNLLAQIGDQYADLF LAAKNLSDAI LLSDI LRVNTETKAPLSASMIKRYDEHHQDLTLLKALVRQQLP EKYKEIFFDQSKNGYAGIDGGASQEEFYKFKP ILEKMDGTEELLVKNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFY PFLKDNREKIEKILTFRIPYVGPLARGNSRFAMTRKSEETITPWNFEVVYDVGASQAQSFIERMTNFDKNLPNEKVLPKHSLLYE YFTVYNELTKVKYVTEGMRKPAFLSGEQKKAIVDLLFKTNRKVTVKQKEDYFKKIECFDSVEISGVEDRFNASLGTYHDDLKIIK DKDFLDNEENEDILEDIVLTLTFEDREMIEERLKYAHLFDDKVMKQKRRRYTGWGRLSRKLINGIRDKQSGKTI LDFLKSDGF ANRNFMLIHDDSLTFKEDIQKAQVSGQDLSHEHIANLAGSPA I KKGILQTVKVVDELVKVMGRHKPENIVIEMARENQTTQKQK QNSRERMKRIEEGIKELGSOI LKEHPVENTQONEKLYLYLQNGRDMYDQELDINRLSDYDVAIVPQSF LKDDSIDNKNVITRS DKNRGKSDNVPSEEVVKKMKNYWRQLLNAKLI TQPKFDNLTKAERGGLSELDKAGFIKQQLVETROI TKHVAQI LDSRMNTKYDEN DKLIREVKVI TLKSKLVSDFRKFDFQYKVVREINNYHHAHDAYLNAVVGTAIIKKYPKLESEFVYGYDVKVYDVRKMIKSEQEIGKA TAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFAVRKVL SMPQVNI VVKTEVQTGGF SKESILPKRNSDK LIARKKDWDPKKYGGFSDPTVAYSVLVAVAKVEGSKKLSVKELLGITIMERSSEFEKNPIDFLEAKGYKEVKKDLI IKLPKYSLF</p>	<p>63</p>

Description	Sequence	SEQ ID NO
	<p>ELENGRKRMLASAGELQKGNELALPSKYVNFLLASHYEKLGKSPEDNEQQLFVEQHKHYLDEIIFQISEFSKRVILADANLQKVI LSAYNKHHRDKPIREQAENI IHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKVEVDATLIHQSI TGLYETRIDLSQLGGDAYPYDVPD YASLGSQSPKKRKRVEDPKKKRVDGIGSGSNGSGSESELIKENMHMKLYMGTVDNHHFKCTSEGGKPYEGTQTMRIKVVEGGP LPFAFDI LATSFLYGSKTF INHTQGI PDFFKQFPPEGTWERVTTTYEDGGVLTATQDTSLQDGLIYVVKIRGVNFTSNGPVMQKK TLGWEAFTE TLYPADGGLEGRNDMALKLVGGSHLIANI KTYRSRKKPAKMLKMPGVYVYDYLRLERIKANNETIYVEQHEVAARYC DLPSKLGHLNGGGGGMDAKSLTAWSR TLTVFKDVFVDF TREEWKLLDTAQQIVYRNVMLNENYKLVSLGYQLTKPDVILRLKKEGE EP</p>	64
<p>Amino acid sequence of dCas9-NLS-FLAG-VP64 (as described in Maeder et al)</p>	<p>MDKKYSIGLAIGTNSVGWAVITDEYKVPKFKFVLTGNTDRHSIKKNLIGALLFDSGETAEATRLKRFARRRYTRRKNRNICYLQEIF SNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFGNI VDEVAYHEKYPTIYHLRKKLVSDTDKADLRLLIYLALAHMIKFRGHFLIEG DLNPDNSDVDKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRLENLIIAQLPGEKKNGLFENLIALSLGLTFNFKSNFDL AEDAKLQLSKDTYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVNTEITKAPLSASMIKRYDEHHQDLLLLKALVRQQLP EKYKEIFFDQSKNGYAGIDGGASQEEFYKFIKPILEKMDGTEELVKNREDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFY PFLKDNREKIEKILTFRIPYVGLARGNSRFAMTRKSEETITPWNFEVYVDKASQAQSFIERMTNFDKNLPNEKVLPKHSLLYE YFTVYNELTKVYVTEGMRKPAFLSGEQKKAIVDLLFKTRKVTYKQKEDYFKKIECFDSEVIEISGVEDRNASLGTYYHDLKLIK DKDFLDNEENEDILEDIVLTLTFEDREMIERLKYAHLFDDKVMKQLKRRRYTGWRLSRKLLINGIRDKQSKTILDFLKSDFG ANRNFMLIHDDSLTFREDIQKAQVSGQSDSLHEHIANLAGSPA I KKGILQTVKVVDELVKVMGRHKPENIVIEMARENQTTQKQ KNSRERMKRIEEGIKELGSOILKEHPVENTQONEKLYLYLQNGRDMYDQELDINRLSDYDVAIVPQSFLKDDSIDNKKVLTTRS DKNRGKSDNVPSEEVVKKMKNYWRQLLNAKLITQRKFDNLTKAERGGSELKAGFIKRQLVETROI TKHVAQIILDSRMNTKYDEN DKLIREVKVITLKSLLVDFRKKDFQYKRVREINNYHHAHDAYLNAVGTALIKKYPKLESEFVYGDYKVDYRKMIAKSEQEI GKA TAKYFFYSIMNFFKTEITLANGEIRKRPLIETNGETGEI VWDKGRDFATVRKVL SMPQVNI VVKTEVQTTGGFSKESILPKRNSDK LIARKKDWDPKKGDFDPTVAYSVLVAVKVERGSKKLSVKELLGITIMERSSEFKENPIDFLEAKGYEVKDKLIIKLPKYSILF ELENGRKRMLASAGELQKGNELALPSKYVNFLLASHYEKLGKSPEDNEQQLFVEQHKHYLDEIIFQISEFSKRVILADANLQKVI LSAYNKHHRDKPIREQAENI IHLFTLTNLGAPAAFKYFDTTIDRKRYTSTKVEVDATLIHQSI TGLYETRIDLSQLGGDGSPPKKRRK VSSDYKDHGDYKDHDIYKDDDDKMAPKKKRKGVMGDKKYSIGLAIGTNSVGWAVITDEYKVPKFKFVLTGNTDRHSIKKNLIGAL MDYKDHGDYKDHDIYKDDDDKMAPKKKRKGVMGDKKYSIGLAIGTNSVGWAVITDEYKVPKFKFVLTGNTDRHSIKKNLIGAL LFDSGETAEATRLKRFARRRYTRRKNRNICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFGNI VDEVAYHEKYPTIYH LRKKLVSDTDKADLRLLIYLALAHMIKFRGHFLIEGDLNPDNSDVDKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRLE NLIAQLPGEKKNGLFENLIALSLGLTFNFKSNFDLAEDAKLQLSKDTYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVN TEITKAPLSASMIKRYDEHHQDLLLLKALVRQQLPEKYKEIFFDQSKNGYAGIDGGASQEEFYKFIKPILEKMDGTEELLVKLN EDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFLKDNREKIEKILTFRIPYVGLARGNSRFAMTRKSEETITPWNFEV VDKGASAQSFIERMTNFDKNLPNEKVLPKHSLLYEFTVYNELTKVYVTEGMRKPAFLSGEQKKAIVDLLFKTRKVTYKQKED YFKKIECFDSEVIEISGVEDRNASLGTYYHDLKLIKDKDFLDNEENEDILEDIVLTLTFEDREMIERLKYAHLFDDKVMKQLKR RRYTGWRLSRKLLINGIRDKQSGKTI LDFLKSDFGANRNFMLIHDDSLTFREDIQKAQVSGQSDSLHEHIANLAGSPA I KKGILQ TVKVDLVLKVMGRHKPENIVIEMARENQTTQKQKNSRERMKRIEEGIKELGSOILKEHPVENTQONEKLYLYLQNGRDMYD QELDINRLSDYDVAIVPQSFLKDDSIDNKKVLTGNTDRHSIKKNLIGALMDYKDHGDYKDHDIYKDDDDKMAPKKKRKGVMGDKKYSIGLAIGTNSVGWAVITDEYKVPKFKFVLTGNTDRHSIKKNLIGAL DKAGFIKRQLVETROI TKHVAQIILDSRMNTKYDENKLIREVKVIITLKSRLVDFRKKDFQYKRVREINNYHHAHDAYLNAVGTAL</p>	64
<p>Amino acid sequence of FLAG-NLS dCas9-NLS-VP64-HA (as described in Perez-Pinera et al)</p>	<p>MDYKDHGDYKDHDIYKDDDDKMAPKKKRKGVMGDKKYSIGLAIGTNSVGWAVITDEYKVPKFKFVLTGNTDRHSIKKNLIGAL LFDSGETAEATRLKRFARRRYTRRKNRNICYLQEIFSNEMAKVDDSFHRLSEESFLVEEDKKHERHPIFGNI VDEVAYHEKYPTIYH LRKKLVSDTDKADLRLLIYLALAHMIKFRGHFLIEGDLNPDNSDVDKLFIQLVQTYNQLFEENPINASGVDAKAILSARLSKSRLE NLIAQLPGEKKNGLFENLIALSLGLTFNFKSNFDLAEDAKLQLSKDTYDDDLNLLAQIGDQYADLFLAAKNLSDAILLSDILRVN TEITKAPLSASMIKRYDEHHQDLLLLKALVRQQLPEKYKEIFFDQSKNGYAGIDGGASQEEFYKFIKPILEKMDGTEELLVKLN EDLLRKQRTFDNGSIPHQIHLGELHAILRRQEDFYFLKDNREKIEKILTFRIPYVGLARGNSRFAMTRKSEETITPWNFEV VDKGASAQSFIERMTNFDKNLPNEKVLPKHSLLYEFTVYNELTKVYVTEGMRKPAFLSGEQKKAIVDLLFKTRKVTYKQKED YFKKIECFDSEVIEISGVEDRNASLGTYYHDLKLIKDKDFLDNEENEDILEDIVLTLTFEDREMIERLKYAHLFDDKVMKQLKR RRYTGWRLSRKLLINGIRDKQSGKTI LDFLKSDFGANRNFMLIHDDSLTFREDIQKAQVSGQSDSLHEHIANLAGSPA I KKGILQ TVKVDLVLKVMGRHKPENIVIEMARENQTTQKQKNSRERMKRIEEGIKELGSOILKEHPVENTQONEKLYLYLQNGRDMYD QELDINRLSDYDVAIVPQSFLKDDSIDNKKVLTGNTDRHSIKKNLIGALMDYKDHGDYKDHDIYKDDDDKMAPKKKRKGVMGDKKYSIGLAIGTNSVGWAVITDEYKVPKFKFVLTGNTDRHSIKKNLIGAL DKAGFIKRQLVETROI TKHVAQIILDSRMNTKYDENKLIREVKVIITLKSRLVDFRKKDFQYKRVREINNYHHAHDAYLNAVGTAL</p>	65

Description	Sequence	SEQ ID NO
<p>dCas9 (bacteria) S. pyogenes >bac teria dCas9</p>	<p>IKKYPKLESEFVYGDYKVDVRKMIKSEQEIGKATAKYFFYSNIMNFFKTEITLANGEIRKRPLIETNGETGEIVWDKGRDFATV RKVLSMPQVNIKKTEVQTGGFSKESILPKRNSDKLIARKKDWDPKKYGGFSDSPTVAYSVLVVAKYKGSKKLKSVKELLGITIM ERSSFENKPIDFLEAKGYKEVKKDLIIKLPKYSLEFLENGRKRMLASAGLQKGNELALPSKYVNFYLASHYKELKGSPEQK QLFVEQHKHYLDEIIIEQISEFSCRVI LADANLKVLSAYNKHRDKPIREQAENI IHLFTLNLGAPAAFKYFDFTTIDRKRYTSTKE VLDATLIHQSI TGLYETRIDLSQLGGDPIAGSRASPKKRVKVRADALDDFDLMLGSDALDDFDLMLGSDALDDFDLMLGSDA LDDFDLMLINYPYDVPDYAS gacgtcttaagaccactttcacatcttaagttgtttttctaatccgcataatgatcaattcaagggccgaat aagaaggtggctctgacacttgatcaataatcgatagcttgcgtaataatggcgccatactat cagtagtggtttcccttcttttagcacttgatgctcttgatcttccaaatcgcaacctaaagt aaaaagccacagcgtgagtgatataatgacatctctagtgaaaaccctgttgccataaaaaggct aattgatttccagagttcactgtttttctgtagccgtgacctaactgacttttgcctccatcgc gatgacttagtaaaagcactaaaacttttagggttattacgtaaaaaactttgccagctttcccttc taaagggcaaaagtgagtgatggtgcctatctaacatctcaatggctaaggctcgagcaaaagcccgctta tttttacatgccaatcaaatgtaggctcttaacccctagcttggggagtttacgggtgtttaaac cttcgattccgacctcaatgagcagctcaatggcgttaaatcactttactttatctaatctagacat cattaaattcctaaattttgtgacactctatcgttgatagattatttaccactccctatcagtgatag agaaaagaattcaaaagatcaaaagagagagagatctatggataagaaatactcaataggcttagcta TCGGCACAAATAGCGTCGGATGGCGGTGATCACTGATGAATATAAGGTTCCGCTAAAAAGTTCAAGGT TCTGGAAAATACAGCCGCCACAGTATCAAAAAAAATCTTATAGGGGCTCTTTTATTTGACAGTGGAGAG ACAGCGGAACCGACTCGTCTCAAACGGACAGCTCGTAGAAGGTATACACGTCGGAAGAAATCGTATTTGTT ATCTACAGGAGATTTTTCAAATGAGATGGCGAAAGTAGATGATAGTTTCTTCACTCGACTTGAAGAGTC TTTTTGGTGGAAAGACAAAGAACGATGAACGTCATCTCTTTTTGGAAATATAGTAGATGAAGTTGCT TATCATGAGAAATATCCAACTATCTATCATCTGCGAAAAAAAATGGTAGAFTCTACTGATAAAGCGGAT TGGCTTAACTATTTGGCTTAGCCATATGATTAAGTTTCGTGGTCAATTTTTGATTTGAGGGGAGATTT AAATCTGATAATAGTGATGTGGACAAACTATTTATCCAGTTGGTACAAACCTACAATCAATTAATTTGAA GAAAAACCTATTAACGCAAGTGGAGTAGATGCTAAAAGCGATTTCTTCTGCACGATGAGTAAATCAAGAC GATTAGAAAAATCTCATTTGCTCAGTCCCGGTGAGAAAGAAAAATGGCTTATTTGGGAATCTCATTTGCTTT GTCATTTGGTTTGACCCCTAAATTTAAATCAAAATTTGATTTGGCAGAGATGCTAAAATACAGCTTCA AAAGATACTTACGATGATGATTTAGATAATTTATGGCGCAAATGGGATCAATATGCTGATTTGTTTT TGGCAGCTAAGAAATTTACAGATGCTATTTTACTTTCAGATACTCTAAGAGTAAATACTGAAAATAACTAA GGTCCCTATCAGCTCAATGATTAACCGCTACGATGAACATCATCAAGACTTGACTCTTTTTAAAAGCT TTAGTTGACAAACACTCCAGAAAAGTATAAAGAAATCTTTTTGATCAATCAAAAAACGGATATGCGAG GTTATTTGATGGGAGCTAGCCAAAGAAATTTATAAATTTATCAAACTAATTTAGAAAAATGGA TGGTACTGAGGAAATTTAGTGAATTAATCTGTAAGATTTGCTGGCGAAGCAACGGACCTTTTGCAAC GGCTCTATTTCCCAATCAAAATCACTTGGGTGAGTGCATGCTATTTTGAGAAAGACAAAGAACTTTTATC CATTTTAAAAGACAAATCGTGAGAAAGATTGAAAAAATCTTGACTTTTTTCGAATTCCTTATTAATGTTGGTCC</p>	<p>110</p>

Description	Sequence	SEQ ID NO
	<p>ATTGGCGGTGGCAATAGTCTGTTTTGCATGGATGGATGACTCGGAAGTCTGAAGAAACAATTAACCCATGGAAAT TTTTGAAGAAAGTTGTCGATAAAGGTGCTTCAGCTCAATCATTTTATGAACGCATGACAAAATTTTGAATAAAA ATCTTCCAAAATGAAAAAGTACTACAAAACATAGTTTGTCTTTATGAGTATTTTACGGTTTATAACGAAAT GACAAAAGGTCAAATACTGTTACTGAAGGAAATGCGAAAACACAGCATTTCTTTCCAGGTGAACAGAGAAAAGCC ATTGTTGATTTACTCTTCAAAAACAATCGAAAAGTAAACCCGTTAAGCAATTAAGAAAGATTATTTCAAAA AAAATAGAAATGTTTGTAGTGTGAAAATTTCCAGGAGTGAAGATAGATTTAAATGCTTCATAGGTACTTA CCATGATTTGCTAAAAATTAATAAAGATAAAGATTTTGGATAATGAAGAAAATGAAGATATCTTAGAG GATAATGTTTTAACATGACCTTATTTGAAGATAGGAGATGATGAGGAAAAGACTTAAAACATATGCTC ACCTCTTTGATGATAAGGTGATGAAACAGCTTAAACGTCGCCGTTATACCTGGTTGGGACGTTTTGCTCG AAAAATGATTAATGGTATAGGGATAAGCAATCTGGCAAAAACAATATTAGATTTTTTGAATCAGATGGT TTTGCCAAATCGCAATTTATGACGCTGATCCATGATGATAGTTTGACATTTAAAGAAAGACATTTCAAAAAG CACAAAGTCTGGACAAGGCGATAGTTTACATGAACATATTTGCAAAATTTAGCTGGTAGCCCTGCTATTA AAAAGGTAATTTACAGACTGTAAAAGTTGTTGATFGAATTTGTTCAAAAGTAAATGGGGCGGCATAAGCCAGAA AATACTGTTATGAAAATGGCACGTAAGAAAATCGACAACTCAAAAAGGCCAGAAAATAATCCGGAGAGCGTA TGAAAACGAAATCGAAGAAAGTAAAGAAATTTAGGAAGTCAAGTTCTTAAAGAGCATCTCTGTTGAAAATAC TCAATTTGCAAAATGAAAAGCTCTATCTCTATTTATCTCCAAAATGGAAGAGACATGATGTGGACCCAGAA TTAGATATTAATCGTTAAAGTATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGAT CAATAGACAAATAAGGCTTAAACGCTTCTGATAAAAATCGTGTAAATCGGATAAACGTTCCAAAGTGAAGA AGTAGTCAAAAAGATGAAAACATTTGGAGACAACTTCAAAACCCCAAGTAAATCACTCAACCGTAAAGTTT GATAATTTAACGAAAAGTGAACGTTGGAGGTTTGGAGTGAACCTGATAAAGCTGGTTTTATCAAAACGCCAAAT TGGTTGAAAATCGCCAAATCACTAAGCATGTTGGCACAATAATTTGGATAGTCGCATGAATACTAAAATACGA TGAAAATGATAAACTTAATCGAGAGGTTAAAGTGAATCACTTAAAATCTAAAATTAAGTTTCTGACTTCCGA AAAAGATTTCCAAATCTATAAAGTACGTGAGATAAACAATTAACAATGATGATGATGATGATGATGATGATGATG CCGTCGTTGGAACTGCTTGTAAAGAAATCCAAAACCTTGAATCGGAGTTTGTCTATGGTGAATATAA AGTTTATGATGTTGTAATAATGATTTGCTAAGTCTGAGCAAGAAATAGGCAAAAAGCAACCGCAAAAATATTC TTTTACTTAATCATGAACCTTCTTCAAAAACAGAAAATACACTTGCAAAATGGAGAGATTCGCAAAACGCC CTCTAAATCGAAAATTAATGGGAAAACGAGAAAATGTTGTTGGATAAAGGGCGAGATTTTCCACAGTGGC CAAAGTATGTCCTATGCCCAAAGTCAAATAATGTTCAAAGAAAACAGAAATACAGACAGGCGGATTTCTCCAAG GAGTCAAATTTACCAAAAAGAAAATTCGGACAAGCTTATGCTCGTAAAAGAAAGACTGGGATCCAAAATAAT ATGGTGGTTTTGATAGTCCAAACGGTAGCTTATTCAGTCTTAGTGGTTGCTTAAAGTTGAAAAGGGAAATC GAAAGAAAGTAAAATCCGTTAAAAGAGTTACTAGGATCACAATTAATGAAAAGAAAGTTCCTTTGAAAATAAAT CCGATTTGACTTTTTTAGAACTAAAAGGATAAAGAAAGTTAAAAGAAAGTAAAATCAATTAACACTACCTAAAT ATAGTCTTTTTGAGTATGAAAACGGTCTGTAACGGATGCTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGTGGT TGAGCTGGCTCTGCCAAGCAAAATATGTGAAATTTTTATATTTAGCTAGTCATATGAAAAGTTGAAAGGTT AGTCCAGAAATAACGAAACAAAACAAATTTGTTTTGTGGAGCAGCATAAGCATTAATAGATGAGATTAATG AGCAAAATCAGTGAATTTCTAAAGCGTGTATTTTAGCAGATGCCAAATTTAGATAAAAGTTCTTTAGTGCATA</p>	

Description	Sequence	SEQ ID NO
<p>Cas9 " > Cas 9 Gene</p>	<p>TAACAAACATAGAGACAAACCAATAACGGTGAACAAACAGAAAAATATTATTTCATTTATTTACGTTGACGAAT CTTGGAGTCCCGCTGCTTTAAATAATTTTGATACAAACAATTTGATCGTAAACGATATACGTCACAAAAG AAGTTTAGATGCCACTTTATCCATCAATCCATCACTGGTCTTTATGAAACACGCATTGATTTGAGTCA GCTAGGAGGTGACTAACtcgagtaaggatctccaggcatcaataaaaaagaaagctcagtcgaaagact gggcccttcgcttttatacgttggttggtgaaagcctctactagagtcacactggctcaaccttcggg ttggcctttctcgcttatacctaggatataatccgcttctcgcctcaactgactcgtcagctcggctcg ttcgactcggcgagcgaaatggcttacgaaagggcggagatctccctggaagatgccaggaagatact taacagggaaagtgagagggccggcaagcgtttccataggtccgccccctgacaagcatcacg aaatcgacgtcaaatcagtggtgggaaacccgacagactataaagataccaggcgtttccccctgg cggctccctcgtgctcctcgtttcctgcttccggtttaccggtgtcaatccgctgttatggccgctg ttgctcatccacgctgacactcagttccgggtaggcagttcgcctcaagctggactgtatgcaagaa cccccgttcagtcgaccgctgcttaccggttaactatcgtcttgagtcacaacccggaaagacatg caaagcaccactggcagcaccctggttaattgattagagggttagcttgaagtcagtcgcccgggtt aaggctaaactgaaagacaagttttggtgactgctcctccaaagcagttacctcggttcacaagagtt ggtagctcagagaaacctcgaaaaaccgctcgaaggggttttttcggtttcagagcaagagattacg cgcagacaaaacgatctcaagaagatcatcttataatcagataaaatattctagatttcagtgcaat ttatctctcaaatgtagcactgaaagtcagccacatacagataaagttagttagtctgaggtcattctc accataaaaaacgcccggcaaccgagcttctgaacaaatccagatggagtctgaggtcattact ggtatctacaacaggtccaagcagctcgaatacaaatcagccccctgcccactcatcgcagctac tgttgtaattcattagcatctcgcgacatggaagccatcacaacagcagatgataacctgaaatccgca gcgcatcagcacccttgcctcgtataatctgccaatggtgaaaaacggggcgaagagttgtc cataatggccacgtttaaatacaaaactggtgaaactcaccagggttggtgagacgaaaaacataatc tcaataaacctttagggaataaggccaggttttcaaccgtaacacgcccacatcttgcgaataatgtgta gaaactcgggaaatcgtcgtggtattcaactcagagcagatgaaaaacgtttcagtttgctcatggaaaaa ggtgtaacaaggtgaaactatccatatacaccagctcaccctcttcaatggccatcgaataatccgga tgagcatcatcaggcgggcaagaaatgtgaataaaagccggataaaaacttgcttatttttcttaacg tctttaaagggccttaataatccagctgaaacggtctggttataggtacattgagcaactgactgaaatgc ctcaaaaatgttcttaacagatgcatgggataatcaacggtggtataccagtgatttttttctccat tttagcttcttagctccbgaaaaatctcgtataactcaaaaaatcagccccggtagtgatcttatttcat ggtgaaagtggaaaccttaccgtgcccgtcaagctctcaatcttccgcaagatctc"</p> <p>gccaccATGGACAAGAAGTACTCCATTGGGCTCGATATGGCACAAACAGCGTCGGCTGGGCCGTCAATTA CGGACGAGTACAAGGTGCCGAGCAAAAAATTCAAAGTTCTGGGCAATACCGATCGCCACAGCATAAAGAA GAACCTCATTTGGCGCCCTCTGTTCGACTCCGGGAGACGGCCGAAAGCCACCGGGCTCAAAAGAACAGCA CGGGCAGATAACCCGAGAAAGAAATCGGATCTGCTACCTGCAGGAGATCTTAGTAATGAGTGGCTA AGGTGGATGACTCTTTCTCCATAGGCTGGAGGAGTCCCTTTTGGTGGAGAGGATAAAAAGCACGAGCG CCACCCAAATCTTTGGCAATATCGTGGACGAGGTGGCGGTACCATGAAAAGTACCCCAACCATATATCATCTG</p>	<p>111</p>

Description	Sequence	SEQ ID NO
	AGGAAGAAGCTGTAGACAGTACTGATAAAGGCTGACTTGGGTTGATCTATCTCGGGTGGCCGCAATGA TCAAAATTCGGGGACACTTCTCATCGAGGGGACCTGAACCCAGACACACAGCGATGTCGACAAAACCTTT TATCCAATCGTTTCCAGACTTACAATCAGCTTTTGAAGAGAACCCGATCAACGCATCCGGAGTTGACGCC AAAGCAAATCTGAGCGCTAGGCTGTCCAAATCCCGGGGCTCGAAAACCTCAFCGCACAGCTCCCTGGGG AGAAGAACAACGGCTGTGGTAACTTATCGCCCTGTCACTCGGCTGACCCCAACTTTAAATCTAA CTTGACCTGGCCGAAGATGCCAAGCTTCAACTGAGCAAAAGACACACTACGATGATGATCGACAACTG CTGGCCAGATCGGCGACAGTACGCAGACCTTTTGGGGCAAGAACCTGTACAGCGCCATTCTGC TGAGTGATAATCTGCGAGTGAACACGGAGATACCAAAGCTCCGCTGAGCGCTAGTATGATCAAGCGCTA TGATGAGCACCCACAGACTTGACTTTGCTGAGGCCCTTGTACAGACGAACTGCCGTGAGAAATCAAG GAAAATTTCTTCGATCAGTCTAAAATGGCTACCCCGGATACATTCAGCGGGAGCAAGCCAGGAGGAA TTTACAAAATTTAAGCCCACTTTGGAAAAAATGGACGGCACCGAGGAGCTGTGTAAGCTTTAACAG AGAAGATCTGTTGGCAAAACAGCGCACTTTCGAAATGGAAGCATCCCCACAGATTCACCTGGCGAA CTGCACGCTATCCTCAGCGGCAAGAGGATTTTACCCCTTTTGAAGAATAACAGGGAAAAAGATTGAGA AAATCCTCACATTTCCGATACCTACTATGAGGCCCTTCGCCCGGGGAAAATCCAGATTCGCGTGGAT GACTCGAAATCAGAGAGACCATCACTCCCTGGAATTCGAGGAAGCTGTGATAAGGGGCCCTCTGCC CAGTCTTCATCGAAAAGGATGACTAATTTGATAAAAACTGCCTAACGAAAAGGCTGTCCATAACACT CTCTGTGTACGAGTACTCACAGTTTATAACGAGCTCACCAAGGTCAAATACGTACAGAAAGGATGAG AAAGCCAGCATTCCTGTGTGAGAGCAGAAGAAAGCTATCTGTGACCTCTCTCAAGACGAAACCCGAAA GTTACCCTGAAAACAGCTCAAAAGAGACTATTTAAAAGAAATGAAATGTTTCGACTCTGTTGAAATCAGCG GAGTGGAGGATCGCTTCAACGCATCCCTGGGACGTATCAGCATCTCCCTGAAAAATCAATAAAGACAGGA CTTCTGGACAATGAGGAGAACGAGGACATTTCTGAGGACATTTGCTCTCACCTTACGTTGTTTGAAGAT AGGGAGATGATTTGAAGACGCTTGAAAACTTACGCTCATCTCTCGACGACAAAAGTCAATGAAAACAGCTCA AGAGCGCCGATAACAGGATGGGGCGGCTGTCAAGAAAACCTGATCAATGGGATCCGAGACAAAGCAGAG TGGAAAGACAAATCCTGGATTTCTTAACTCCGATGGATTTGCCAACCGGAACTTCAATGCAGTTGATCCAT GATGACTCTCTCACCTTAAAGGAGGACATCCAGAAAACAAAGTTTCTGSCCAGGGGACAGCTTTCAG AGCACATCGCTAATCTGCAGGTAGCCAGCTATCAAAAAGGGAATACTGCAGACCGTTAAGGTCGTGGA TGAACCTCGTCAAAGTAAATGGGAAGGCAATAGCCGAGAAATATCGTTATCGAGATGGCCCGAGAGAACAA ACTACCAGAGGACAGAAGAACAGTAGGGAAGGATGAAGAGGATGAAGAGGTTGAAAGAGGTTAAAAGAACTGG GGTCCCAAATCCTTAAAGAACACCCAGTTGAAAACACCCAGCTTCAGAAATGAGAAAGCTCTACCTGTACTA CCTGCAGAACCGCAGGACATGTACGTGGATCAGGAATGGAATCAATCGGCTCCCGACTACGACCTG GATCATATCGTCCCGAGCTTTCTCAAAGATGATCTATTGATAATAAAGTGTGACAAAGATCCGATA AAAATAGAGGGAAGAGTAAACGTCCTCCCTCAGAAAGGTTGTCAAGAAAATGAAAAATTAATGGCGGCA GCTGTGAACGCCAACTGATCACACAACGGAGTTCGATAAATCTGACTAAGCTGAACGAGGTGGCTG TCTGAGTTGGATAAAGCCGGCTTCAATAAAGGAGGAGTGTGAGACAAAGCTGATCCACCAAGCAGTTGG CCCAAATTCGATTCAGCATGAACCAAGTACGATGAAAATGACAAAACCTGATTCGAGAGGTGAAAGT TATTACTCTGAAGTCTAAGCTGGTCTCAGATTTTCAGAAAAGGACTTTTTCAGTTTTATAAAGGTGAGAGATC	

Description	Sequence	SEQ ID NO
Human codon optimized Cas9 ">Cas 9 ORF	AACAAATTACCACCATGGCATGATGCCCTACCTGAATGCAGTGGTAGGCACCTGCACATTATCAAAAAATATC CCAAGCTTTGAAATCGAAATTTGTTTACGGAGACATAAAAGTGTACGATGTAGGAAAATGATCGCAAAGTC TGAGCAGGAAAATAGCAAGCCACCCTAAGTACTTCTTTTACAGCAATAFATGAATTTTTTCAAGACC GAGATTACACTGGCCAAATGGAGAGATTCCGAAAGCCACCTTATCGAAAACAAACGGAGAAAACAGGAGAAA TCGTGTGGACAAGGTAGGGATTTCCGGACATCCCGAAGGCTCTCTCCGAAAGGAAAGTATCTCCGAAAAGCAACAGCGACAAG TAAAAAGCCGAAAGTACAGACCGGAGGCTTCTCAAAGGAAAGTATCTCCGAAAAGGAAAGCAACAGCGACAAG CTGATCGCACGCAAAAAGATTTGGACCCCAAGAAATACGGCGGATTCGATTCCTACAGTCGCTTACA GTGTACTGGTTGTGGCCAAAGTGGAAAAGGAAAGTCTAAAAAATCAAAAAGCGTCAAGGAACTGCTGGG CATCAAAATCATGGAGCGATCAAGCTTCGAAAAAAAACCCCATCGACTTTCGAGGCGAAAAGGATAAAA GAGGTCAAAAAGACCTCATCATTAAGCTTCCCAAGTACTCTCTCTTTGAGCTTGAAAACGGCCCGGAAAAC GAATGCTCGTAGTGGGGGAGCTGCAGAAAAGGTAACGAGCTGGCACTGCCCTCTAAAATACGTTTAAATTT CTTGTATCTGGCCAGCCACTATGAAAAGCTCAAAGGTTCTCCGAAAGATAATGAGCAGAAAGCAGCTGTTTC GTGGAACAAACAAACACTACTCTTGTATGATGAGATCAFCGAGCAAAATAAGCGAAATTCCTCAAAAAGAGTGAATCC TCGCCGACGGTAACTCGATAAGGTGCTTTCTGCTTACAAATAAGCACAGGATAAGCCCATCAGGGAGCA GGCAGAAAACATTAATCCACTTGTTTACTCTGACCAACTGGGCGCCCTGCAGCCTTCAAGTACTTCGAC ACCACCATAGACAGAAAAGCGGTACACCTCTACAAAAGGAGGCTCTGGACGCCACACTGATTCATCATAGTCAA TTACGGGGCTCTATGAAAACAAAGAAATCGACCTCTCTCAGTCTGGTGGAGACAGCAGGGGCTGACCCCAAGAA GAAGAGGAAAGGTGTGA	112

Description	Sequence	SEQ ID NO
	actgaagagctgctggtgaagctgaacaggggagacctgctgaggaagcagagggacctttgataaatggctccattccacatcagatacacctgggagagctgatgcaatcctccgagcagaggagattctctatcctttcctgaaggataaccgggagaagatagagaagatcctgacctcaggatccctattacgtcggccctctggttagaggcaactccgcttgccttgatgacaggaatactgaggagacaattactccttggaaactctgaagaggtcgtggaataagggcgcaagcccagctcatcctcgaacagtgaccacaatttcgataaagaacctgccccaacgagaaggtcctgccccaaacattcaactctgacagatttcaacgctataacgagctgactaaagtgaagtacgtgacgagggcatgaggaagctgctcctcctgcccggagagcagaagaagggctatcgtgatccttcaagactaatagaaggtgacagtgaacagctcaaggagattactttaaagaagatcgaatgctttgactcagtggaattctctggctgagggaccgctttaaagccagcctgggcacttaccatgatcgtgagaatacaagacaagaattcctcgtataatgaggagaacagaggacatcctggaagatacgtgctgacccctgactcgtcgaggatagagagatgatcgaagagcctgaagaccctatgccccatctgttgacgataaaagtcataaacagctcaagcggcggctacactgggtgggtgagactctccaggaaactcaataaacggcattccggacaaacagagcgaagaccatcctggattcctgaaatccgacggattcctaacaggaaacttcaagctgactcctgacgttcaagatttaaaagggagacatccagaaagggcacaaggtgagcgtcaagcagcctgacagacatcgccaaacctcgtggatcaccggcctaaagaaaggaatactgcagacagctcaagctcgtggacgaaactcgtcaagtgatgggtcggcacaaagccaggaatactggtatcgaaatggcaagggagaaaccaaaccacagaaagggccagaaagaaactcctggaaacggatgaaagaatcgaaggggaatgaaggagctggaactcagatactgaaggagaccctgtggagaaatcacagctccagaacgagaaactctacctgactacccagaaacgggggacatgtaagttgaccaggaactcgcacatacaccggctgaccgatgacgtggaccaatggtccacagctcctcctcaagatgactccatgacaacaaggtgctgaccagatccgataaagaatcggcgtgaagctgacaatgtccatcagaagaggtggtcaagaagatgaagaattactggcggcagctcctcaacgcaactgacccagcggaaagtggacaatctgactaaggcagaagagggctgagcgaactcgaacagccggcttataagaggcaactgggcgaaacaccagatccaaacacgtggcaaaatcctcgaactcgaactaggaagaaactcgtgagaaactcgaactcgtgagaaagtgaaagtgaataactcgaagcaagctggtcctgactccggaaggacttcaatctacaaagtccgaaataaacaattaccatcgtcagatgacctcaatgctgctggcaccctgatcaagaataaccctaaactggagctgagttcgtgacggtgactataaagcttacgatgaggaagatgataagctgagaagagattggcaaacccgcaagctactctctactcaataatctttaaagactgataaccctggctaacctggctaacgggaaatccggaaagcggccactgatcgaacaaacgggaaacaggaatacgtgtgggataaaggcagactcgcactgtgcggaaggtgctccatgccacaagtcaatcgtgaaagaaccgaaagtcagaccggcggatctcaaaaggagagatcctgccaaagcgaactcgaagctgacccaggaagaagattgggacccaagaagatggcgggttcgattcctccacagtggcttaccgttctggtcgtggcaaaagtggaaagcaggtcccaagaaactcaagctgtaagagagctcggaaataactataaggagataccagcttcgagaagaatcccaatcgattcctggaaagctaaagggctataaaagaagtgaagaagatctcatcatcaactgccccaaagttactcttggagctggagaatgtaggaagcggctccggcggagctgcagaaaggaaacggag	

Description	Sequence	SEQ ID NO
	ctggctctgccctccaaatacgtgaacttccctgtatctggcctcccactacgagaaaactcaaaggtagcc ctgaagacaatgagcagaagcaactctttgttgagcaacataaacactacctggacgaaatcattgaaca gattagcgagttcagcaagcgggttattctggccgatgcaaacctcgataaagtgtgagcgcataaat aagcacagggaagccaattcgcgaacaagcagagaaattatccacctcttactctgactaatctgg gcgctcctgctgccttcaagtatctgatatacaactattgacaggaagcgtacacctctacccaaaagt tctcgatgccaccctgatcacaccagtcattaccggactgtacgagactcgcatcgacctgtctcagctc ggcggcgactag"	

Table 6: 5' UTR

Description	Sequence	SEQ ID NO.
Synthetic polynucleotide 5' UTR	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAATAACGACTCACTATAGGGAAAATAAGAGAGAAAAAGAGCTAAAGAAATATAAGA GCCACC	71
sgRNA 5' UTR	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAATAACGACTCACTATAGGG	72
5UTR-001; Synthetic UTR	GGGAAAATAAGAGAGAAAAAGAGAGTAAGAGAAAATAAAGAGCCACC	15
5UTR-002; Upstream UTR	GGGAGATCAGAGAGAAAAAGAGAGTAAGAGAAAATAAAGAGCCACC	16
5UTR-003; Upstream UTR	GGAAATAAAAGTCTCAACACACATATACAAAACAAACGAAATCTCAAGCAATCAAGCATTTCTACTTCTATTCAGCAATTTAAATCA TTCTTTTAAAGCAAAAGCAATTTCTGAAAAATTTTACACCATTTACGAAACGATAGCAAC	17
5UTR-004; Upstream UTR	GGGAGACAAGCUUGGCAUUCGGUACUGUUGGUAAGCCACC	18

Table 7: 3' UTR

Description	Sequence	SEQ ID NO.
Ex. 9 3' UTR	TGATAAATAGctg gagectogtgccatgett cttgccccctt gggcctcccc ccagccccctc ctcccccttcc tgcacccgtacccccgtgt ctttgaataaa agtctgagtg ggcggc tCTAGA	81
miR122 3' UTR	TGATAAATAGCTGGAGCCTGGTGGCCATGCTTCTTGCCCTTGGCCCTCCCCAGCCCTCCTCCCTCCCTGCACCCCGTACCC CCAAAACACCAATTGTCACACTCCAGTGGTCTTTGAAATAAAGTCTGAGTGGGGGCTCTAGA	82
3UTR-001; Creatine Kinase	GGCCCTGCCACCTGCCACCGACTGCTGGAACCCAGCAGTGGAGGGCCCTGGCCACCACAGAGTCTGCTCCCTCACTCCCTCGCCC CGCCCCCTGTCCAGAGTCCACCTGGGGCTCTCCACCTTCTCAGAGTCCAGTTCACACCAGAGTTCACACCAATGGGCTC CATCCCTGGATTCTGGCCAAATGAAAATACTCCCTGGCAGGGTCTCTTCTTTCCAGAGCTCCACCCAAACCAGGAGCTTAGT TAA TGGAGAGCTCCAGCACACTCGGAGCTTGTGCTTCTCCACGCAAGCGGATAAATAAAAGCAATTGGTGGCCCTTTGGTCTTT GAAATAAGCCTGAGTAGGAAGTCTAGA	19
3UTR-002; Myoglobin	GCCCCCTGCCCTCCCAACCCCACTCTGGCCCCGGGTTCAAGAGAGAGCGGGGCTGTGATCTCGTGTAGCCATATAGAGTTTGC TTCTGAGTGTCTGTCTTGTAGTAGAGGTTGGCAGAGAGCTGAGGGCTGGGGTGTGAAAGTGGCTTTGCATGCC AGCGATGCGCCTCCCTGTGGATGTCACTACCCCTGGAAACCGGAGTGGCCCTGGCTCACTGTGTTCTGCATGGTGTGGATCTGA ATTAA TTGTCTTCTTAAATCCCAACCGAATCTTCCAACTCCAACTGGCTGTAACCCCAATCCAAAGCCATTAACCTACA CCTGACAGTAGCAATTTGCTGATTAATCACTGGCCCCCTTGAAGACACAGAGATGTCCCTTTGCAATGAGGAGGAGATCTGGGCTGG GGGGCCAGCTGGGAAGCAATTGACTATCTGGAACCTGTGTGCTCCCTCAGGATGCGAGTACCTCAGTGGTTTTAATAAAAA CAACTGCAACATCTATGGTCTTTGAAATAAAGCCTGAGTAGGAAGTCTAGA	20
3UTR-003; α - actin	ACACACTCACCTCCAGCAGCGACTTCTCAGGACGACGAACTTCTCAATGGGGGGCGGTGAGCTCCAGCCACCCCGCAGTCA CTTCTTTGTAACAACITTCGGTGTCTGCCATCGTAAACTGCACACAGTGTATAACGTGTACATACATTAACCTTATACCTCAITTT TGTTATTTTTCGAAACAAAAGCCCTGTGGAAGAAAATGGAACACTTGAAGAAGCATTAAGTCAATCTCTGTTAAGCTGCGTAAATGGT CTTTGAAATAAGCCTGAGTAGGAAGTCTAGA	21
3UTR-004; Albumin	CATCACATTTAAAAGCATCTCAGCCTACCATGAGAAATAAGAGAAAATAAAGATCAAAAAGCTTATTCATCTGTTTTCTTTTT CGTTGGTGTAAAAGCCACCCCTGTCTAAAAAACAATAAATTTCTTTAATCATTTTGCCTCTTTCTCTGTGCTTCAATTAATAAAA AATGGAAGAATACTAATAGAGTGGTACAGCACTGTATA TTTTCAAAGATGHTGTCTATCCTGAAAATCTGTAGGTTCTGTGGAA GTCCAGTGTCTCTCTTATCCACTTCGGTAGAGGATTTCTAGTTTTCTTGTGGGCTAATAAATAAATCAATAATACTCTTTCTTAA TGGTCTTTGAAATAAGCCTGAGTAGGAAGTCTAGA	22
3UTR-005; α - globin	GCTGCCCTCTGCGGGCTTGGCTTCTGGCCATGCCCTTCTCTCCCTTGCACTGTACCTCTTGGTCTTTGAATAAAGCCTGAG TAGGAAGCGGCCGCTCGAGCATGCACTAGA	23
3UTR-006; G- CSF	GCCAAGCCCTCCCCATCCCATGTAATTTATCTCTAATTAATAATTTATGCTAATTAAGCCTCATATTTAAAAGACAGGGAAGAGCAGA ACGGAGCCCCAGGCCCTCTGTCTCTCCCTGCATTTCTGAGTTTCATCTCCCTGCCTGTAGCAGTGAAGAAAAGCTCTGTCTCTCC CATCCCCGGACTGGGAGGTAGATAAGTAAATACCAGATTTATTAATGACTGCTCCCGCCCTGGCTGTGCAATGGGCACT GGGATGAGCCGCTGTGAGCCCTGGTCTGAGGGTCCCACTGGAGATTCAGGTTCCACGTGGGAGACAAGAA ATCCCTGTTTAAATA TTTAAAACAGCAGTGTTCCTCCCACTCTGGGTCCTTGACCCCTCACTCTGGCCCTGCACAGCGGCC	24

<p>3UTR-007; Coll1a2; collagen, type I, alpha 2</p>	<p>CTGCATCCCTTGGCTGTGAGGCCCTTGGACAAGCAGAGGTGGCCAGAGCTGGGAGGCAATGGCCCTGGGTCCCACGAAATTTGGCTG GGGAAATCTCGTTTTTTCTTAAGACTTTTGGGACATGGTTTGGACTCCCGAACATCACCAGCCGGTCTCTGTGTTTTTTCTGGGTGGC CTGGGACACCTGCCCTGCCCCACGAGGTCAGGACTGTGACTCTTTTATGGGCCAGGAGGCTCCCTGGACATTTGCCCTTGTCTGG ACGGGACTGGGATGTGGAGGGAGCAGACAGAGGAAATCATGTGAGCCCTGTGTGAAAGGAAAGTCCACTGTCAACCCTCCAC CTCTTCACCCCCACCAGTGTCCCTCCACCTGCACATTTGTAACCTGAACTTCAGGATAATAAAGTGTGTCCTCCATGGTCT TTGAAATAAAGCCTTGTAGGAAGCCGGCTCGAGCATGCATCTAGA ACTCAATCTAAAATAAAGAAAGAAAATTTGAAATAAACTTTCTCTTCCCATTTTCTTCTTTTAACTGAAAGTGAAT CCTTCCATTTCTTCTGCACATCTACTTGTAAATTTGGCCAAAGAAAAGAAAGGATTTGATCAGAGCATTTGTGCAATACAGT TTGATTAACCTCTTCCCGCTCCCCAAAATTTGAAATTTTTTTTTCAACACTTACACCTGTATAGAAAATGTCAACCCTTG TAAGAAAACCAAAAATAAAAATAAAAATAAAAATAAAAATAAAAATAAAAATAAAAATAAAAATAAAAATAAAAATAAAAATAAAA TTTAAAACCCAACTCCAAAGGTTTAAACTACCTAAAACACTTTCCCATGAGTGTGATCCACATTTGAGGTGCTGACCTAGAC AGAGATGAACCTGAGGCTCTGTGTGTGTGTTTGTTCATAATAACAAGGTGCTAATTAATAGTATTTTCAGATACTTGAAGAATGTTGAT GGTGTAGAAGAAATTTGAGAAGAAATACTCCTGTATGAGTGTATCGTGTGGTGTATTTTTTAAAATAATTTGATTTAGCATTCAT AATTTCCATCTTATTTCCAAATAAAGTATGCAGATTTATTTGCCAAAATCTTCTTCAGATTCAGCATTTGTCTTTGCCAGTCTCA TTTTTCATCTTTCTTCCATGTTCCACAGAAAGCTTTGTTCTTGGCAAGCAGAAAATAAATAATGTTACCTATTTGTATATGTGAGA TGTTAATAATAAATTTGTGAAAATAATAAATAAAGCATGTTTGGTTTTCAAAAGAACATAAT</p>	25
<p>3UTR-008; Col1a2; collagen, type VI, alpha 2</p>	<p>CGCCGCGCCCGGGCCCGAGTCGAGGGTCTGAGCCCAACCCCTGTCATGTTGCTAAGCGGGCCCGGTCCCACACGCGCCAGCAC CGTGTCTACTCGGACGAGCCCTGGCCCTGCACCTCTCCAGCTCTCCACGGGGTCCCGTAGCCCGGCCCGCCCGCCAGCCCC AGGCTCCCCAGGCCCTCCGAGGCTGCCCGCCCTCCCTCCCTGCAAGCCATCCCAAGGCTCCTGACCTACCTGGCCCCCTGAGCT CTGGACAAGCCCTGACCAATAAAGGCTTTGAACCCAT</p>	26
<p>3UTR-009; RPN1; ribophorin I</p>	<p>GGGGTAGAGCCCTCTCCGACAGCTGGAGAGCGGGCAAGAGGGGGTTATAGGATTTGGTGGTTTTGTGTTTTGTTTTTAA GCGGTGGGAAAATGGCACAATTTACTCTGTGGGAGATGCAACACTGAGAGCCAAAGGGTGGGATTTGGGATAATTTTTTATATAA AAGAAGTTTTTCCACTTTGAAATGCTAAAAGTGGCAATTTTCTATAGTGCAGTCACTCTCTCATTTTAAAATAAGGACGTGGCC AGGCACGGTGGCTCATGCCTGTAATCCAGCACTTTGGAGGCCGAGGCAAGGCTCACGAGGTGAGGATCGAGACTATCCCTG GCTAACACGGTAAAACCTGTCTACTAAAAGTACAAAATAATAAGGATGAGTGGGCTGTTGGGAGGCTGAGTCCAGCTACTCGG GAGGTGAGGACGAGGAAAGGATGAATCCAAGAGGAGAGCTTGCAGTGCAGTGCAGCTGAGATCACGCTATGACCTCCAGCTGGGCAA CAGTGTAAAGACTCTGTCTCAATA GGCCCTGCCCGTCCGACTGCCCCAGAAAACCTCTCTGCCCTGCAAGTGAAGTCTTCAGTGAGCCCTCCCCAGCCAGCCCTT CCCTGGCCCCGGATGATA TATTTTCCATCCCTCCCTGCTGCTCTTGGCACCCCATGCTGCCCTCAGGAGACAGGAGGAGGCTTGGGCTGCACCT CCTACCTCCACAGAACGCAACCCACTGGGAGGCTGGTGTGAGCCCTCCCTGTATAAGACACTTTGCCAAGGCTCT CCCCCTCGCCCCATCCCTGCTTGGCCGCTCCCAAGCTTCTGAGGGCTAAATCTGGGAAAGGAGATCTTTGTGCCCTGTC TGGAAGACGTGGCTCTGGGTGAGTAGGGGGAAAGGATGGAGTGTTTAGTCTTTGGGGAGGCCCCCAACCCAGCCCCCAA CTCCAGGGGACCTATGAGATGGCCATGCTCAACCCCTCCAGACAGGCTCCCTGCTCTCCAGGGCCCCCAGGTTTCCCA GGGCTGGAGACTTCCCTGTTGTAACAATTTCTCCAGCTCCCTCCCTGGGACCCCAAGGAGTGGGCGCACACCCAGGAAAGGAA AGCGGGCAGCCCGTTTTGGGGACGTGAACGTTTTTAAATAATTTTGTGTAATTTCTTAACTAAAATAACACAGATAATTTGTTATA AATAATAATTGT</p>	27
<p>3UTR-010; LRP1; low density lipoprotein receptor- related protein 1</p>	<p>GGCCCTGCCCGTCCGACTGCCCCAGAAAACCTCTCTGCCCTGCAAGTGAAGTCTTCAGTGAGCCCTCCCCAGCCAGCCCTT CCCTGGCCCCGGATGATA TATTTTCCATCCCTCCCTGCTGCTCTTGGCACCCCATGCTGCCCTCAGGAGACAGGAGGAGGCTTGGGCTGCACCT CCTACCTCCACAGAACGCAACCCACTGGGAGGCTGGTGTGAGCCCTCCCTGTATAAGACACTTTGCCAAGGCTCT CCCCCTCGCCCCATCCCTGCTTGGCCGCTCCCAAGCTTCTGAGGGCTAAATCTGGGAAAGGAGATCTTTGTGCCCTGTC TGGAAGACGTGGCTCTGGGTGAGTAGGGGGAAAGGATGGAGTGTTTAGTCTTTGGGGAGGCCCCCAACCCAGCCCCCAA CTCCAGGGGACCTATGAGATGGCCATGCTCAACCCCTCCAGACAGGCTCCCTGCTCTCCAGGGCCCCCAGGTTTCCCA GGGCTGGAGACTTCCCTGTTGTAACAATTTCTCCAGCTCCCTCCCTGGGACCCCAAGGAGTGGGCGCACACCCAGGAAAGGAA AGCGGGCAGCCCGTTTTGGGGACGTGAACGTTTTTAAATAATTTTGTGTAATTTCTTAACTAAAATAACACAGATAATTTGTTATA AATAATAATTGT</p>	28
<p>3UTR-011;</p>	<p>ATATTAAGGATCAAGCTGTAGCTAATAATGCCACCTCTGCAGTTTTTGGGAACAGGCAATAAAGTATCAGTATACATGTTGATGT</p>	29

<p>Ntl1; cardiotrophin -like cytokine factor 1</p>	<p>ACATCTGTAGCAAAAGCTTTGGAGAAAAATGAAGACTGAAGAAAAAAGCAAAAACCTGTATAGAGAGATTTTCAAAAAGCAGTAAT CCCTCAATTTTAAAAAAGGATTGAAAAATCTAAAATCTTAAATGCTTTCTGTGCATATTTTTGTTGTTAGGAATCAAAAAGTATTTTATAAAG GAGAAAAACAGCCTCATTTAGATGTAGTCTCTGTGGATTTTTTATAGCCTCCAGTAACCCAGAAAATGTTTAAAAACATAAGTG TTTAGGATTTCAAGACAAATATACATGGCTCGTGAATATCTGACACAATGTAAAACATTCAGGCGCACCTGCATTTATGTTTTTT TTTTCAACAAATGTGACTAAATGAACTTTTTAAGCTCTGAGCTGCTCCCTGCAATTCACCCGAGTTTGAATTAATCAAT CAAACTAGTTTTTAAATTTGACTCTCAGAGTCTATATTCAAGGGCACAATTTCTCAGTACTATTTTAATACATTAAGG ACTAAAATTTCTTCAGAGATGCTGAAAACAAATCAATTGTTTTTATAGTTTCAATTAGAATACCAATGAAAACATAACAACCTTGA TTAGTAAATAGTATTTTTGAAAGTCCCATTTCTAATTTGAGATCTCTTTAAATTCGATCAACTATAAATGTGTAGTACATATAAG TGCACTTGAGTGGAAATTCACAATTTGACTAATAAATGAGTTCAATCAATTTGGCAAGTGATGTGGCAATTAATCTCTGGTGACAAAA GATFAAAAATCAAAATTTCTGCTGTTTACAAATATCAAGGAAGACCCTGCTACTATGAAAATAGATGACATTAATCTGCTTTCACCTGT TTATAATACGGATGGATTTTCAAAATCAGTGTGTTTTTGAGGCTTATGTAATTGATGACATTTGAGAGAAAATGGTGGCTTT TTTTAGCTACCTCTTTTGTTCATTTAAGCACCAGTAAAAGATCATGCTTTTTTATAGAAAGTAGATTTTCTTGTGACTTTGCTATC GTGCTAAAGCTCTAAATATAGGTGAATGTGTGATGAATACATCAGATTTTGTCTCTCTATAATAATTAGTTTGGTACTAAGTTTC TCAAAAAAATTAACACATGAAGACAATCTCTAAAACAGAAAAAAGAAAGTACTGTAACAAAATTTTGGTTACTGTAATGCTCGCGTTTTAG TGAGTTTAAAACACACAGTATCTTTGGTTTTATAATCAGTTTTCTATTTTGGTGTGCTGAGATTAAGATCTGTGATGTGTGTG GTGTGTGTGCTGTTTGTGTAAAGCAGAAAAAGATTTTTTAAAAATTTAAAGTATAAGTATAAAATGCAAATTTGTAATGACTTTAGA TCACTAGTAAACTCAGGCTGAAATATACCATGATATACTAATAGAAAGAAATAAACACCATCTTTATTCCTGCCCCTTTCTTC TCTCAAAAGTGTGTGTTATATCTAGAAAGAGCAATTTGATTTCTTGAAAAAGGTAGTCTCCAGCAGTTTAAACTAAAAAT AAATCATCTTGGATTTTATTTATTTTGGTCTATAGTAAAAAATTTTTAAATTTATAATATTTTTATTTTAGTATATCTTATTTCTTGGCT ATTTGCCAACTTGTGCATCAATTTGTTAATGAAATTTAAATTTTCTGCTGTCATGCTTTTATTTTACCTTTATTTGTTAGGATA TTTTAAAGGATTTTGTATATAAAATTTCTTAATAATAATTTCCAAAAGTTAGTGGACTTAGATTAATAATGACAAAAATCTA AAAAACAAAAATGATTTTATACATTTCTATTTCTCTTTTCCAAATAAGTCATAAATTTGGTACATAAATTTGGTAGATAATGACTTATTT TATTTTGTATATTACATATACTTTATAGATTTAAAGTATAATAATAATAAAAAATTTATTTGTACCTTAGCTGTCACCCAAA AAAAAAAAATTTCTGTAGGTAGTGAATGCTAATTTAGTATTTGTCTTTAAGGCTTGTAACTATCTTTTATTTCTCATTTTGTCT TTAAATTAGGATTTTGTTTTAAATTTACTCATTAAGCAAAAAATGATATAAATCCCATTTACTGGGTAATACCCAAAAGGATTA AAATCATGCTGTCTATAAGACACATGCACACGTATGTTTATGACACCTAATTCACAAATAGCAAAAGACTTGAACCAACCCAAAATG TCCATCAATGATAGCTTGAATAAGAAAATGTGCACAATAACCCATGGAATACTATGACCCATAAAAAGGATGAGTTCATGTC CTTTGTAGGGACATGGATAAAGCTGAAAACCATCTTTGAGCAAACATAATGGCAAGGACAAAACCCAAAACATGCATGTTCTCAC TCATAGGTGGAAATGAAACAATGAAACACCTGGACAAGGTTGGGAAACACACACACCCAGGCTTCATGGGGTGGGGGAGT GGGAGGGATAGCATTAGGAGATACCTAATGTAATGATGAGTTAATGGGTGACGACACCAACATGGCACATGTATAACATATG TAGCAAACCTGCACGTGTGCACATGACCCTAGAACTTAAAGTATAATAAAAAAAAAGAAAACAGAAGCTATTTATAAAGAA GTTATTTGCTGAAAATAATGATGATCTTTCCCATTTAAAAAATAAAGAAAATTTGGGGTAAAAAACACAAATATAATTGTATTTCTTGA AAAATTCAAAGAGGTGGATGAGTGAAGTGTCTCACCAAAAAGTATAAATAATGAGGTAAATGCACATATAAATAGAAAGATTT TGTCATTTCCAAATGTATATATACTTTAAAATATGTATACAAATAAATATAACATTAATAAATAAAGTAAATAATGTA</p>
<p>3UTR-012; Col6al; collagen, type VI, alpha 1</p>	<p>CCCACCTGACGGCGCAAAAACCTGTCTCCCAACCTTCCCCTCATACATAAAGCAAGTAAATGTTAGTCGAAATTTTCC CGACCAACTGATTCGGTAGATTTTTTAAAGAAAAGCTTGGAAAGCCAGGACACAACCGTGTGCTGCTTGTGACAGGGTCTT CCGGGCTCAGCCCTGAGTTGGCATCACCTCGGCAAGGCCCTCTGGGGCTCAGCCCTGAGTACTGACAGGGCCCTCT GAGGCTCAGCCCTGAGCTGGGCTACCTGTGCGAGGCCCCCTGTGGGGCTCAGGCTGAGGCTCAGCCCTGCACACCCCGG GCTCTCCCTGCCCCTGCTGCCCCGCCCCTGCTGCTGGCCAGCTCCTCCCTTAGGCACCTCTCTGCTGCATCCCAAGGCC</p>

	<p>CGTGCTCTTTGGGAAGGAGACCTGGCCCTGCTACCCACGGACTGGGTGGCTCGACCTCCTGAAATCCCCAGAACAACACCCCCC TGGCTGGGGTGGTCTGGGAAACCATCGTGGCCCGCTCCCGCTACTCCTTTTAAGCTT TTGGCCAGGCTGACCCCTTTGGACCTTTCTTTTGGCCGACAACCACTGCCACGAGCCTCTGGGACCTCGGGGTCCCAGGGAAC CCAGTCCAGCTCCTGGCTGTTGACTTCCATTGGCTTTGGAGCCACCAATCAAAGAGATTCAAAGAGATTCTGCAGGCCAGAGG CGGAACACACCTTTATGGCTGGGCTCTCCGTGGTCTTGACCCAGCCCTGGAGACACCAATTCACATTTTACTGCTTTGTAGTG ACTCGTCTCTCAACCTGCTTCTGAAAAAACAAGCCCTTCCCACTTCCATGGGTGAGACTTGAGCAGAACAAGG GCTTCCCAAGTTGCCAGAAAGACTGTCTGGGTGAGAAAGCCATGGCCAGAGTTCTCCAGGCACAGGTGTTCACCCAGGGACTT CTGCTCAAAGTTTGGGTAAAGACACCTGGATCAGACTCCAAGGCTGCCCTGAGTCTGGACTTCTGGCTCCATGGCTGGTCA GAGACAAAACCGTAGTCCCTGGAGACAGGACTCCAGAGAACCTCTTGGGAGACAGAAGGCAATCTGTGCACAGCTCGATCTTC TACTTGCTGTGGGAGGGAGTGACAGGTCCACACACACTGGGTACCCCTGTCTGGATGCCCTGAAAGAGAGGGGACAGACC GTCAGAAACTGGAGAGTTCTATAAAGGTCAITTAACCA</p>	<p>33</p>
<p>3UTR-016; P1od1; procollagen- lysine, 2- oxoglutarate 5-di oxygenase 1</p>	<p>TCCTCCGGACCCAGCCTCAGGATTCCTGATGCTCCAAGGGACTGATGGGGCTGGATGAAAGTGGCACAGTCAGCTTCCCCTGG GGGCTGGTGTCAATGTTGGCTCCTGGGGGGGGGACGGCTGGCATTTACGCAATGCTGCCACCCAGGTCCACCTGTCTCCAC TTTACAGCCTCCAAGTCTGTGGCTTCCCTTCTGTCTCCGAGGGCTTGGCTTCTCGTGTCCAGTGAAGTGTCAAGTATC GGCTAACTTAGAGAAAGCCCGCCCTCCCTTCTCCGTCTGTCCCAAGAGGGTCTGCTCTGAGCCTGGCTTCCTAGGTGGCTCGG CCTCAGCTGCCCTGGTTGTGGCCGCCCTAGCATCCTGTATGCCACAGCTACTGGAATCCCCTGCTGCTCCGGGCCAAGCTTCT GGTGATTAATGAGGGCATGGGTGGTCCCTCAAGACTTCCCTACCTTTTGTGGAACCAAGTGTGCTCAAAGACAGTGTCCCC TCCACAGCTGGTGGCCAGGGCAGGGATCCTCAGTATAGCCCGTGAACCTGTACCCACACAGCCCAAGAGGCTGGCCCTGAAACCCCTGG CTTCCAGCCATCTCATCGCCAGCCTCCTCCTGGACTTGGCCCCCAGCCCTTCCCCACACAGCCCAAGAGGGTCCCAGAGCT GACCCCACTCCAGGACCTAGGCCAGCCCTCAGCCTCATCTGGAGCCCTGAAGACCAGTCCCAACCCACTTCTGGCCCTCATCT GACACTGCTCCGCAATCCTGCTGTGTCTGTTCCATGTTCCGTTCCATCCAAATACACTTTCTGGAAACAAA</p>	<p>34</p>
<p>3UTR-017; α- globin</p>	<p>GCTGGAGCCTCGGTGGCCATGCTTCTTGGCCCTTGGGGCTCCCCCAGCCCTCCTCCCTTCTCTGACCCCGTACCCCCCGTGGTCT TTGAATAAAGTCTGAGTGGGCGGG</p>	<p>35</p>

Table 8: DNA constructs encoding synthetic polynucleotides (mRNA) coding for CRISPR related proteins

Description	Sequence	SEQ ID NO:
DNA sequence coding for mRNA transcript of Cas9 (Mali)	TCAAGCTTTGGACCTCGTACAGAAGCTAATACGACTCACTATAGGAAATAAGAGAGAAAAAGAGTAAGAAATAAAGA GCCACCATGGACAGAAATACAGCATCGGCTGGATATGGAACTAACAGCGTTGGATGGCAGTGCATACCGACGAGTACAAGGT GCCGAGCAAGAAGTTCAAGGTGCTGGGAACTGACCGCCATTCAAATTAAGAAAAAACCTCATTGGAGCACTGCTTTTGGACTCGG GTGAGACTGCCGAAGCTACAGGCTCAAACGCCACGACGACGAGCGGTACACCCGCCGCAAGAAATCGCATCTGCTATCTGCAAGAG ATCTTTTCCAAACGGAGATGGGAAGTTGACGACAGCTTTTTTCCACCGGCTGGAAGAGAGCTTCCCTGTTGGAAGAGGACAAAAAGCA CGAAAGGCATCCAAATCTTCGGTAAACATCGTGGACGAAGTGGCGTATACGAAAAAGTACCCCTACCATCTACCATCTGCGGAAAGGAGC TGGTCGATTCCACGGATAAGGCAGACCTGAGACTGATCTACCTGGCTTTGGCCCATATGATCAAAATCCCGGGCCATTTCCCTGATC GAGGGGACCTTAACCCGGATAACTCGGATGTCGACAAGCTGTTTCATCCAGCTGGTCCAAAACGTAACCAACTGTTTGGAGAAA TCCCATCAACCGCTTCGGGGTGGACGCCAAAGCAATCTCTCCCGCGCTGAGCAAGTACCGGCTGACTCCAAACTTTAAGTCAAATTC AGTCCCGGGAGAAAAAGAAAATGGACTGTTGGAACTGATCGCGCTGTCGCTCGGCTGACTCCAAACTTTAAGTCAAATTC GATTTGGCCGAAGATGCCAAGCTGCAGCTGCAAAAGGACACTTACGACGACGACCTGGCAATCTGCTGGCCAGATTTGGGACCA ATACGCAGACCTGTTTGGCCCGGAAGAACCTGAGCGACCCCTTCTTGTCCGATATTTCTGAGAGTCAATACCGAAATCACTA AGGTCCTGCTCCGCTTCAATGATCAAGCGCTACGATGAACACCCACAGGATCTCACTCTGCTCAAAGCCCTCGTGAGACAACAA TTGCCTGAAAAGTACAAGGAGATCTTCTTCGACCAGAGCAAAAACGGCTACGACGCTACATCGATGGAGGAGCGTCAACAAGAA GTTCTCAAAGTTCAATCAAGCCAACTTTGGAGAAGATGGACGGTACTGAAAGAACTCCTTGTGAAGCTGAAATAGGGAGGATTTGCTCA GAAAGCAGCGGACTTTTGACAAACGGCTCGATCCCTCATCAGATTCACCTCGGTGAGCTGCAATGCCATCCTTCGGCCCAAGAGGAT TTTTACCCCTTCCCTGAAAGGATAATCGCGAGAAAATCGAAAAGATCCTGACGTTTCAAGATTCCTTACCTGCTGAGCCGCTGGCCG CCGTAACTCGCGGTTTGCATGGATGACTCGCAAGTCAAGAGAAAATCACTTCCGAAACGAGAAAGTTCCTTCAAAAGCACTCCCTGTTG CCTCCGCCAGTCAATTCATCGAACGCATGACCAACTTCGACAAGAACTTCGACAAGAACTTCGCAAGAACTTCGCAAGCACTCCCTGTTG TACGAATACTTCCCGGTACAAATGAGCTGACCAAAAGTTAAGTATGTCAACGAGCCAAACCGGAAAGGTGACCGTCAAGCAACTCAAGGAGGACTACTTCAAGAA ACAAAAGAGGCCATCGTCGACCTCCTTCAAGACCAACCGGAAAGGTGACCGTCAAGCAACTCAAGGAGGACTACTTCAAGAA TCGAATGCTTTGACTCGGTGAAAATCAGCGGAGTGGAGGACCGGTTTAAACCGCTCACTGGGTACCTACCATGATCTCCTGAAAAATC ATCAAAGACAAGGACTTCTGGACAACGAAAGAAAACGAGGACATCCTGGAAAGATATTTGTGTGACCTGACCGTGTTCGAGGACCG GGAAATGATCGAGGAAAGGCTTAAGACCTACGCACACCTCTTCGATGACAAAAGTATGAAAGCAACTGAAAGCGGCGGAGATATACTG GCTGGGGGAGGCTCTCCCGGAAGCTCAATTAATGGAAATCAGAGACAAAAGTCCGGTAAAACATACTCTCGACTTCTCAAGTCCGAT GGGTTCGCCAACCGGAACTTCAATGCAGCTGATCCACGATGATTCCTTGACTTCAAGGAAGATATCCAGAAAGGCGCAAGTGAAGCGG ACAGGGAGATTCGTTGCACGAACATAATCGCTAAATCTCGCCGGATCCCAAGCCATCAAGAAAAGGAAATCTTCAGACCCGTGAAGGTGG TGGATGAACCTGGTGAAGTATGGGGCCCAAAAACAGAGAACTATGATGGATGGCCCGGAGAAATCAGACCCACTCAGAAAG GGACAAAAGAACTCCAGAGAGCGGATGAAACCGCATCGAGGAAAGGCTTGGTAGCCAAAATCTTGAAGGAAACACCCCGGT	41

Description	Sequence	SEQ ID NO:
	<p>CGAGAACACCCAGCTCCAGAACGAAAAGCTTTACCTGTACTACCTCCAAAATGGACGGGACATGTACGTCGACCAGGAATTTGGACA TCAACAGACTCAGCGACTACGATGTGGACCATAATGTGCCACAGTCCITTTCTTAAGGACGACAGCATCGATAACAAAAGTGTCACT AGATCAGACAAAATCGCGGAAAATCAGACRAATGTGCCATCGGAAGAGTGTCAAGAAGATGAAAAAATACTGGAGACAGCTGCT CAATGCCAAAATTCACCCAGCGGAAGTTCGACAACTTACCAAGCGGAGCGGAGGATTTGCCGAATCGACAAGCCGGCT TCATCAAAAGGCCTGGTGGAAAACCCGGCAGATCACTAAACACGTCGAAAACCTGGTGTGGACTTTAGAAAAGGATTTCCAGTTCTA GAGAA TGACAAAGCTGATTAGGGAAGTCAAGGTCACTACTGAAAGTCAAACTGGTGTGGACTTTAGAAAAGGATTTCCAGTTCTA CAAAGTCCGGAGATTAACAAC TACCACCGTCA TGACGCTACCTGAA TGCAGTGTGGGACCCGCGTGTATCAAGAAATATC CGAAGCTGGAA TCCGAGTTCGTGTACGGAGATTAACAAGTGTACGACGTGCGCAAGATGATCGCCAAAGTCGGAACAGGAAA TCGGA AAGGCTACTGCAAAGTACTTCTTCTACTCAAACATCATGAAC TTTCTTCAAACCGGAGATCACGCTCGGAAACGGGAAA TCCGAA AAGGCCGCTCATTTGAAACCAAACGGGAAACCGGGGAGATCGTGTGGGCAAGGGAAGGATTTTGGCACTGTGAGGAAGGTGTGT CCATGCCGCAAGTCAATATTTGAAAAGACGGAAGTGCAAAACCGGAGATTCAGCAAAGAAATCCATCTTCCAAAAGCGCAACTCG GACAAAATCATCGCGCAAGAAAGGATTTGGACCCCAAGAAAATACGGTGGCTTTGACAGCCCTACTGTGGCTTACTCCGCTCCTCGT CGTGGCTAAAAGTGGAAAAGGTTAAATCCAAAAAGTCAAA TCGGTGAAGAGCTCCTGGAAATCACGATCATGGAGCGGTGAGCT TCGAAAAGAAATCC TATTGATTTCTTGGAGGCAAGGCTACAAGGAAATCAAGAAAGACCTGATCATCAAGCTCCCGAAAGTACAGC CTC TCCGAGCTCGAAAACGGCAGAAAAGAGGATGCTGGCATCAGCGGGAGAA TGCAGAAAGGAAAACGAACTGGCACTGCCGTCCAA GTACGTGAA TTTTCTCTATCTGGTAGCCACTACGAAAAGTGAAGGATCGCCGAGGACAAACGAGCAAAAA CAAC TGTTCGTGG AGCAGCAAGCACTACTGATGAGATCATCGAGCAGATCTCCGAA TTTTCGAAAACGCGTGTCTTCCGCAATGCCAAATCTGGAT AAAGTGTGTCCGCTTACAACAAGCATCGGATAAACCGATCCGCGAACAGGCAGAAAACATCATCTGTTCAC TTTTGAACCAA TCTGGGAGCGCTGCCGCTTAAAGTACTTCGACACCACTAT TGATAGAAAAGCGCTACACCTCGACCAAGGAA GTGCTGGACCGTA CCC TGATCCACCAGTCCATCACCGGACTCTACGAAACTCGCAT TGACCTGTCCCAGCTTGGAGGAGATTCACGGGCCGATCCAAAG AAAAGCGCAAGGCTGATAA TAGGCTGGAGCCTCGGTGGCCATGCTTCTTGGCCCTTGGGCCCTCCCCCAGCCCCCTCCTCCCTT CCTGCACCCCTACCCCGTGGTCTTTGAAATAAAGTCTGAGTGGGGGCTCTAGA</p>	42
DNA sequence coding for mRNA transcript of Cas9 (Cong)	<p>TCAAGCTTTTGGACCTCGTACAGAAAGCTAATACGACTCACTATAGGAAAATAAGAGAGAAAAGAGTAAGAAGAAAATAAAGA GCCACCATGGACTACAAGGACCACGACGGGACTACAAGACCATGACATCGATTAACAAGGATGACGATGACAAAATGGCACCGAA GAAGAAGAGAAAAGTCCGAAATTCACGGGTGCCGCGGACAAAGATPACTCAATCGGACTGGATTCGGACGAAACAGCGTGG GTTGGCAGTGTACCCGAGAAATACAAGTCCGAGCAAGAAAGTTCAAAGTGTGGAAAATCCGATCGCCATTCGATCAAGAAA AATCTGATTTGGCGCTCCTGTTTCGACTCGGAGAGACTGCCGAGCCACTAGACTGAAGAGGACCGCTAGGCCCGCTACACGAG GCGAAAACCCGCTGCTATCTTCAAGAAAATCTTCTCAAACGAGATGCCAAAGTGGACGACTCTTTTCCATCCGCTGGAAAG AATCATTTCTGGTGGAGGAGCAAGAAGACGAAACGCCATCCCATTTTCGCAACATTTGTCGACGAAAGTGGCTTATCATGAGAAAG TATCCGACTATCCACTTGAGAAAAGGCTGGTGGACTCCACTGACAAGGCAGATCTGCGGTTGATCTACCTCGCACTGGCCCA TATGATCAAATTCGGGGACATTTCTCTCATCGAGGGCACCTTAATCCGCAAAATTCGATGTGGAATAAGCTTTTCAATCCAGCTGG TCCAGACCTACAACCAACTGTTTGAAGAAAATCCAA TCAA TGGAGCGGTGTCTGATGCAAAAAGGCCATCCTGAGCGCCCTCTCG</p>	42

Description	Sequence	SEQ ID NO:
	AAAAGCAGAAGGCTCGAAAACCTGATCGCCACAGCTTGCCCTGGAGAGAAAGAAAGCGGCCTCTTCGGCAATCTCATCGCATTTGCCCT GGGACTGACTCCAAAATCAAAATCCAACCTCGACTTGCCGAGGACGCGCAAACTGCAACTGAGCAAAAGATACCTACGATGATGACT TGGACAAATCTTCTGGCTCAGATCGCGCAGCTAGCCCGGACTGTTCTTTCGGCTTAAGAACCTTCGGACGCCATCTCGCTGTCTCC GACTCTCGCGCTCAATACCGAAATTAATAAGACCACCTCTCGGATCCATGATCAAGAGATACGATGAACACACCCAGGATCT CACCTCTGAAAGCAGCTGGTGGCCAGCAGCTCCCTGAGAAATCAAGGAAATCTTCTTTGATCAGTCCAAAGAACCGGATACGCCG GATACATCGACGGCGCGAGCCAAAGAGAAATTTACAAGTTTCACTAACCGGATCTTGAAAATGATGGCACCGGAAAGAACTC CTGGTCAAACTGAATFAGAGAGGATCTGCTCCGCAAAACAACGGACTTCGATAACGGATCGATCCCGCACCCAGATCCACCTCGCGA ACTGATGCCATCCTCGCGCGGCAAGGACTTTTACCCGTTCTCAAGACAACAGAAAAGAFCGAGAAGATCTTGACTTTTC GCATCCCGTACTACGTGGGCCCCGCTCGCGAGAGGTAACCTCCCGCTTTGCTGGATGACTAGAAAAGTCAGAGGAAACGATCACCCCA TGGAACTTCGAAGAGGTGGTTGACAAAGGAGCGGCCCAATCGTTTCACTGAACGGATGACTTAACCTCGATAAGAACTCTGCCGAA TGAGAAGTCTCCCTAAGCACTCACTTCTGTATGAAATACCTTACTGTGTATAACGAACTCACCAAAAGTCAAAATCCGTGACTGAGG GAAATGCGCAAGCTCGCTTTTGTCCGGGAGCAAAAAGGCCATCTGGACTTGCTGTTTCAAAACCAACCGCAAGGTGACTGTT AAGCAACTCAAAGAGACTACTTTAAGAAAGTCGAATGCTTTGACTGGTCTGAGATTTCCGGGTTGAAAGATAGATTCACACCGCTC GCTGGAACTTACCAATGATCTCTCAAGATTAACAAGGACAAAGACTTCTTGATAAACGAGGAGAAATGAGGACATCTCGAAAGATA TTGTGTTACCTGACCTTTTCGAAGATCGGAAATGATCGAAAGACCGCTGAAAACCTACGCTCACCTGTTCCAGCATAGGTG ATGAAAACAGTTGAAAACCGCGGCTACACCGGTTGGGCGGCTGTCCGCAAGCTGATCAACGGAAATCGGGACAAACAGAGCGG AAAGACCATCTCGATTTTGTGAAGTCCGATGTTTTCGCAACCGCAACTTCATGCACTCATCCATGACGATTCGCTTACCTTA AGGAGGATA TCCAGAAAGCACAAGTGTCCGGACAAGGGGATTCGCTCCACAAACATCGCCAAATCTGGCGGGTTCGCCGGCAAT AAGAAAGGAAATCTCCAGACTGTTAAGGTGGTCCAGACTGTTGAGGAGTGTGGGAGACATAAGCTGAAAACATTTGTGATCGA GATGGCGAGAGAAAATCAAACACTACTCAGAAAGGACAGAAAGAAATTCGCGGAGCGGATGAAGCGCATCGAGGAGGAAATCAAGGAAC TGGGCTCCCAATCCTGAAAAGAGCATCCCGTGGAAAATACTCAGCTCGAAGACGAGAAAGCTTTTACCTGTACTATCTTCAAAAATGGC AGGGACATGATCGACCAAGAACTGGATATCAATCGGCTCTCCGATFACGACGTCGATCAGTACATCGTCCCCCAATCATCTCGAA GGATGATAGCATCGATAACAAGGTGCTACTAGATCAGACAAAACCGGGGAAAGTCAGATAAGTCCCGCAGCAAGAAAGTCTGTGA AGAAGATGAAGAAATACGGAGGCACTTCTGAAACGGCAAACTCATCTCAGCGCAAGTTCCGAAACCTGACCAAACTGACCAAAAGG GGAGGACTCAGCGAGTGGACAAGGCTGGTTTCACTAAACGGCAGCTGGTGGAGACTCGCCAAATCAAGAAAGTGTGGCCAGAT TCTCGACTCGCGCATGAAATACTAAGTACGACGAAAACGATAAGCTGATCCGGGAGGTGAAGGTGATCACCCCTCAAGAGCAAGCTCG TGTCCGATTTCCGGAAGACTTCCAGTTTCAAGGTTCCGGGAGATTAACAACCTACCATCACGCTCACGACGCTTACCTCAATGCT GTGGTGGGACGGGTTGATTAAGAAATCCCAAACTGGAGTCCGAAATCTGCTACGGAGATTAACAAGTCTACGACGTCGCGCAA GATGATTTGCCAAGTCGGAGCAGGAAATTTGGGAAAGCGACTGCTAAGTACTTCTTCTACTCGAATACTATGAACCTTCTTCAAGCCG AAACTACCTTGGCTAACGGGAGATCAGGAAACCGCCGCTGATCGAAACATAATGGTGGAGACTGGTGAATTCGTGTGGGATAAAGGA CGGGACTTCGCGACGGTCCGAAGTCTCAGCATCCGCAAGTGAATGTTTAAGAAAACCGAAGTGCAGACCCGGTGGGTCTC GAAAGGAAATCCATCTGCCAAAAGCGCAACTCGGATTAAGCTTATTTGCCCGCAAGAAAGGATTTGGGACCCGAAAAGATCAGGTGGTTCG ACTCCCTACCGTGGGCTACTCCGTTGTTGGTGGCCAAAAGTGGAAAGAGGCAAAATCAAAGAGTCAAGAGCTCAAGGAGCTG CTGGAAATCACCATCATGGAGAGGTCCAGCTTTGAGAAAACCCGATTCGACTTCTTGGAAACCCGAGGAAAGCGCATGCTGCCCTCAGCCCGGAACTGC AGACTGATCATCAAACCTCCAAAGTACTCCCTGTTCGAACTCGAAAACGGGAGGAAAGCGCATGCTGCCCTCAGCCCGGAACTGC AAAAGGGCAACGAACTGGCCCTCCCGTCAAAAATACGTCAACTTCCCTGTACTTGGCGTCACTACGAAAAGCTGAAAAGGATCCCCA	

Description	Sequence	SEQ ID NO:
	<p>GAGGACAACGAAACAGAAACAGTGTTCGTCGACGACACAAGCACTCCTGGACGAGATCATCGAAACAGATCTCGGAATTCAGCAA GAGAGTGATCTTGGCAGACGCTAACCTTGCACAAGTCTCTCGGCATACAATAAGCATCGCGACAAGCCGATCAGAGAAACAGCGGG AGAACATATCCACCTGTTCACCTCACCACTGGCGCCAGCGGCTTTAAAGTACTTTGATFACCACCATTTGACCGCAAGAGA TACACCTCAACTAAAGAGTGTGGACGCAACCTTGATCCATAAAGCATCACCGGACTTTATGAAACTCGGATCGATCTCTACA GCTCGGAGGACAAAAGACCGGCTGCCACCAAGAAGCCGGACAGGCAAGAAGAAATGATAA TAGGCTGGAGCCCTCGGTGG CCAATGCTTCTTGGCCCTTGGCCCTCCCCCAGCCCTCCTCCCTTCTGCACCCGTAACCCCTGGTCTTTGAAATAAAGTCTGAG TGGGGGCTCTAGA</p>	43
<p>DNA sequence coding for mRNA transcript of dCas9-KRAB</p>	<p>TCAAGCTTTTGGACCTCGTACAGAAGCTAATACGACTCACTATAGGGAATAAAGAGAGAAAAGAAGAGTAAGAAGAAAATAAAGA GCCACCATGGACAAGAAAATCAATCGGACTTGCTATCGGAACATAACGCTGGATGGCCCTGCTCACTACTGACGAATACAAGGT GCCCTCAAAGAAGTCAAGGTCCTGGAAAATAACGATAGACACTCCATCAAAAAGAACTGTATCGGGCACATGTTGTTCCGACTCGG GAGAGACTGCAGAAACCAAGGCTCAAGCGACTGGAGGGCCGGTACACCCGGAGAAACCCGATCTGCTACCTCCAAAGAG ATTTTCAGCAACGAGATGGCAAAGGTCGATGATTCGTTCTTTACCCGCTTGAGGAGTCGTTCTTGTGAGGAGGACAAAAAGCA TGAAGACATCCGATCTTCGAAAACATCGTGGACGAAGTGCATACCATGAAAAGTACCCCTACCATCTACCATCTCAGAAAAGAAAC TCGTTCGATTCAACTGATAAGGCCGACTTGGGGTGATCTACTGGCTTGGCGCACATGATCAAGTTTCGGGGTCACTTCTCATC GAGGTGATCTCAACCCGGACAATTCGGACGTGACAAAACCTTCTCATCCAACTGGTCCAGACGTAACAACAGCTGTTCCGAAAGAAA TCCGATCAACGCCAAGCGGAGTGAAGCCAAAGCCATCTGTGCGCCCGCTCTGAAAGTCGCTCGCTGGAAAATCTGATTCGCTC AGCTCCCGGGGAAAAGAAGAAATGGCCCTGTTTGGAAAACCTCATCGCACTGCTCCCTCGGGTACTCCCAACTTCAAACTCGAACTTT GACTTGGCTGAGGATGCAAAAGTCAACTTCCAAAAGACACTTACGATGATGACTTGGACAATCTCTGGCGCAGATCGGGGATCA GTATGCTGACCTGTCTCGCGGCAAGAACCTGTCTGATGCCATCTCTGCTCCGATACTCTGAGAGTGAACACTGAGATCACCA AGGGCCTCTGAGCGCTCGATGATCAAAACGTTACGATGAACACCATCAGGACCTCACTCTTCTGAAAGGCTTGGTCCGGCAGCAG CTTCGGGAAAAGTACAAAAGAGATCTTCTTCGACCAGTCGAAAACCGCTACCGCGGATACATTTGATGCGCGGCAAGCCAGGAGGA ATCTATAAGTTTTACAAAACCGATCCTGGAGAAGATGGACGGCACTGAAAGAACTTCTGGTCAAGCTGAAATCGAGAGGATCTGCTCC GGAAGCAGCGGACCTTCGACAAATGGGTCTATCCCTCACAAAATCCATCTCGGCGAGCTGCAATGCGATTTGAGGCGCCAGGAGGAC TTCTACCCATTCCTGAAAAGCAAATCGGGAGAAAATCGAAAAGATTTCTGACGTTCCGCAATCCATACACTACGTCGGGCCACTTGGCGG GGGTAAATTCGAGATTCGCTGGATGACCGGGAAGTCCGAAAGAAACCATCACGCCGTGAAATTTCCGAAAGAGTGGTCGACAAAGGGAG CCAGCGCACAGTCTTCATGAGCGCATGACCAAATTCGACAAAATTCGCAAAAATTCGCGAACGAGAAAGTCTCTCCGAAAGCATTCACTGCTG TACGAATACCTTACCGGTACAAACGAACTGACCAAGGTGAAGTACGTCAAGGAAATGAGAAAAGTCTGCTTCTGAGCGGAGAGA ACAGAAAGAGGCCATTTGTTGACTCTCTTCAAGACTAATCGAAAAGTACCGTGAAGCAGTAAAGAGGATTAATTCAAAAGAAAGA TCGAAATGTTTCGACTCCGTGAAAATCAGCGGCTGGAGGATAGATTCACCGGCTCCCTTGGGACTTACCACGACCTCCCTTAAGATC ATCAAGGATAAGGATTTCTTCGACAAATGAGGAAAACGAAGATATCTCGGAGGACATCGTCTGACTTGACCCCTTTTGAGGACCG GGAGATGATCGAGGAGAGACTCAAAGCCTACGCGCACTGTTTTCGACGACAAAAGTATGAAGCAACTTAAACGACGCGCTACACCG GCTGGGCGAGACTGTCACGCAAGTTGATCAACGGAATAGAGATAACAGTCCGAAAAGACCTCTGGACTTCTGAAAGTCCGAT GGAATCGCCAAACCGGAAATTTCAATGACGCTATCCATGACACTCAATGACTTTCAAGGAGGATTCGAAAAGGCCAAAGTGAAGCGG CCAAGGGACTCCCTTCAGAACACATCGGAAAATTTGGCCGATCACCGGATTAAGAGGGAATCTCGACACCGTGAAGGTGG TGGACGAGCTGGTGAAGTGTGATGGGACGGCAAGCCGGAACATCTGATCGAGATGGCCAGAGAGAACCCAGACGACTCAAAAG GGCCAGAAAGAACTCGCGCGAACCGCATGAAGAGAAATGAAGAGGGAATTAAGGAACTGGGATCGCAGATCTTGAAGGAGCACCTGT</p>	43

Description	Sequence	SEQ ID NO:
	<p>CGAAAAATACCTCAGAACGAGAAGCTGTACCTGTACTATACTTCAAACGGCAGGGACATGATGTGACCAAGAGCTCGACA TTAACCGCCTGTCCGATATGACGTGGACGCCATCGTCCGAGAGCTTCTCAAGGACGATCCATCGACAAACAAAAGTGTCAACC CGCAGCACAAGATAGAGGGAAGTCGGATAACGTCCTTTCGAAAGGAGTGGTGAAGAAATGAAAGATTAAGTGGCGCAGCTCCT GAAATGCAAGCTCAACCCAAACGGAAGTTTGACAACCTCACCAAGAGGAAAGAGGAGGACTGTCCGAAATGGATAAGCCCGGTT TCATCAAGCCGCAATGGTGGAACTCGGCAAAATCAACCAAGATGTGGACATGTGGACATGAAACACCAACCAAGTATCGAC GAGAACGATAAGCTGATCCGCGAGGTCAAGGTGATCACCTCAAAGCAAACTTGTGTGACAGCTTCCGAAAGGACTTCCAAATCTTA CAAGGTCGCGGAAATCAAACAACATACCACCGTCAATGACGCATACCTGAACCGTGTGGTCCGACTGCCCTCATCAAGAGTAC CTAAACTCGAAAGCGAAATTTGTGTACGGGACTACAAGGTGTACGATGTCCGGAAGATGATCGCGAAATCCGAGCAGGAGATCGGA AAGGCACTGCTAAGTACTTTTCTACTCGAACAATCATGAACCTTCTCAAACCCGAAATCACCTGGCTAATGGCGAGATCAGAAA GGCCCGCTGATCGAAAACCAACCGCGAAACCGGTGAAATCGTGTGGGACAAAGGCGCGGATTTCCGTACTGTGAGAAAAGGTCCTTT CCAATGCCGCAAGTGAATATCGTCAAAGAACTGAGGTGACAGCTGGCGGATTTCCAAGGAATCGATCCTCCCAAAGAGGAACTCA GATAAGCTCATCGCGGAAAAGGATTTGGACCTTAAGAAAGTACGGAGATTTGATAGCCCAACTGTGGCTACTCTGTGCTCGT GGTGGCCAAAGTCGAGAAAGGAAAGTCGAAAGAAAGTTGAAATCCGTGAAAGAACTTTGGGAATCACTATCATGGAGCGGTCTCAT TTGAAAAGAACCCCAATCGACTTCCGTGAAAGCCAAAGGATACAAGAAAGTCAAGAAAGCTGATCACTAAGCTCCCTAAGTACAGC CTGTTCGAACTGGAGAACGGAAGGAAACGGATGCTGGCTTCCGCCGGGAACTGCAAAAGGGCAATGAGCTGGCCCTCCCATCGAA ATACGTGAACCTTCTACTACTTGCCTCCCATTAACGAAAGTTGAAGGGTCAACCGGACCAATGAGCAGAGAAACAGCTCTTTTGTG AACAAACAACTACTGACGAAATCAAGAAATCAAGAAATCAAGAAATCAAGAAATCAAGAAATCAAGAAATCAAGAAATCAAGAAAT AAAGTGTGTCCGGTACAACAAGCACCGCAATAAGCCGATACGGGAAAGGCTGAGAACTCAATCACTCTTCACTCTCACTAA TCTGGAGCCCCCGCCTTCAAGTACTTGTACTACTACCTCGAACCGAAAGATACAGGACCAAGAAAGTCTCGATCGATGC CCCTGATCCACCACTCATACTTGTGTGTAAGAAACCGGAAATCGATGTCACAGCTCGGAGGAGATGCGTACCCCTACGATGTC CCCGACTACGCTCACTCGGTAGCGGCAGCCCGAAGAAAGAAAGGTTGAGGAGCCCGAAGAAAGAAAGGAAAGGTTGACGGGAT CGGAAGCGGATCGAATGGATCGTACGAGGAGGTGGCGGAGGTGAGACCGCAAAATCACTTACGGCTGGTCAACGGACCTTGGTGA CCTTTAAAGACGTGTTCGTGGATTTCAACAGGAAAGAAATGGAACTGTGGACACCCCGCAGCAGATCGTGTACCCGAAATGTGATG CTGGAAACTACAAAATTTGGTGTCCCTGGGTATCAACTCACTAAGCCAGATGTCATCTTTAGACTGGAAAAGGGAGAAAGAACCC GTGATAATAGGCTGGAGCCTCGGTGGCCATGCTTTTGGCCCTTGGCCCTCCCGCAGCCCTCTCCCTTCTTCCACCCCGTACC CCCGTGGTCTTTTGAATAAAGTCTGAGTGGGCGGCTCTAGA</p>	44
DNA plasmid sequence of dCas9-NLS- FLAG-VP64	<p>TCAAGCTTTTGGACCTCGTACAGAAAGTATAACGACTCACTATAGGAAAATAAGAGAGAAAAGAAAGAGTAAGAAAATAAAGA GCCACATGGACAAGAAATACCTAACTCGGACTTGCTATCGGAATCAAGCTGAGGCTGGATGGGCGCTCATTAATGACGAATACAAGT GCCCTCAAAGAAGTCAAGGTCCTGGGAAATACCGATAGACACTCCATCAAAAAGAACTGATCGGGCAGCTGTGTTCGACTCGG GAGAGACTGCAGAGCAACCAAGCTCAAGCCGACTGCGAGGCGCCGTAACCCCGGAGAAAGAACCCGATCTGCTACCTCCAAAGAG ATTTTCAGCAACCGAGATGGCAAGGTCGATTCGTTCTTTCACCGCTTGAAGGAGTCTGTTCTGTCGAGGAGGACAAAAAGCA TGAAGACATCCGATCTTCGAAACATCTGTGGACGAAGTGCATACCTGAAAGTACCCATCCATCTACCATCTCAGAAAAGAAAC TCGTGATTTCAACTGATAAGGCCGACTTGGCGTGTACTACTGGCTGTGGCGCACATGATCAAGTTTCGGGGTCACTTTCTCATC GAGGTGATCTCAACCCGGACAATCCGACGTTGACAAAATCTTCACTCAACTGGTCCAGACGTAACCAAGCTGTTCGAAAGAAA TCCGATCAACCGCAAGCGGAGTGAACCCCAAGCCATCTGTTCGGCCCGCTCTCGAAGTCCGCGTCCCTGAAATACTGTGATGCTC AGCTCCCGGGGAAAAAGAAAGAAATGGCCCTGTGTTGAAACCTCACTCGCACTGTCCCTCGGGCTGACTCCCAACTTCAAATCGAACTTT</p>	44

Description	Sequence	SEQ ID NO:
	<p>GACTTGGCTGAGGATGCAAAAGCTGCAAACTCTCAAAAGACACTTACGATGATGACCTGGACAAATCTCTGGCGCAGATCGGGGATCA GTATGCTGACCTGTCTCTGGCGGCAAGAACTGTCTGATGCCATCCTGTCTCCGATATCCTGAGAGTGAACACTGAGATCACCA AGGCGCTCTGAGCGCTCGATGATCAAAAGCTGACGATGAAACACCTCAGGACCTCACTCTTCTGAAAGCTTTGGTGGCGCAGCAG CTTCCGGAAAAGTACAAAGAGATCTTCTTCCGACAGTCAAAAGAAACCGGTGACCCGGATACATTTGATGGCGGCGCAAGCCAGGAGGA ATTCTATAAGTTTATCAAAACCGATCCTGGAGAAAGTGGACGGCACTGAAACACTTCTGTCAAGTGAATCGAGAGGATCTGCTCC GGAAGCAGCGGACCTCGACAAATGGTCTATCCCTCACCAAATCCATCTCGCGGAGTGCATGCGATCTGAGGCGCCACAGGAGAC TTCACCCATTCGAAAAGACAAATCGGGAGAAAATCGAAAAGATCTGACGTTCCGATTCATACCTACGTTGGGCGCCACTTGGCGG GGGTAATTCGAGATTCGCTTGGATGACGCGGAACTCCGAAAGAAACCAATCACGCCCTGGAATTCGAAAGAAAGTGGTCGACAAAGGAG CCAGCGCACAGTCTTCAATGAGCGCATGACCAAATTCGACAAAATTCGCCGAAACGAGAAAGTCTGCCGAAAGCATTCACCTGCTG TACGAAATCTTACCGTGTACAACGAACTGACCAAGGTGAAGTACGTCACCGAGGAAATGAGAAAGCCTGCTTCTCTGAGCGGAGAG ACAGAAAGAGGCCATTTGACCTCTCTCAAGACTAATCGCAAAGTACCGTGAAGCAGCTTAAAGAGGATTAATTCAAAAGA TCGAAATGTTTCGACTCCGTGGAAATCAGCGGCTGGAGGATAGATTCACCGCTCCCTTGGGACTTACCACGACCTCCTTAAAGATC ATCAAGGATAAGGATTTCTTCGACAAATGAGGAAAACGAAAGATATCTGGAGGACATCGTTCTGACTCTGACCCCTTTTGGAGACCG GGAGATGTCGAGGAGACTCAAGACCTACCGGCACCTGTTGACGACAAAAGTGAAGGCACTTAAACGACGCGCTACACCG GCTGGGCAGACTGTACGCAAGTTGATCAACGGAAATAGAGATAACAGTCCGAAAGACCTCTGGACTTCTTGAAGTCCGAT GGATTCGCCAACCCGGAATTTCAATGACGCTCATCCATGACGACTCAATGACTTCAAGGAGGATTCGAAAGGCGCCAAAGTGAAGCGG CCAAGGGACTCCTTCACGAAACACATCGCAAATTTGGCCGGATCACAGCAGATTAAGAAAGGAAATCTGCAGACCGTGAAGTGG TGGACGAGCTGTTGAAAGTATGGACGGCACAAGCCGAAAACATCGTGTGAGATCGAGATGGCCAGAGAAACACGACGACTCAAAAG GGCCAGAAAGAACTCGCGGAAACGATGAAAGAAATAGAAAGGAAATTAAGAACTGGGATCGGAGATCTTGAAGGAGCACCTGT CGAAAATACTCAACTCCAGAACGAGAACTGTACTGTACTTCTCAAAAACGGCAGGACATGTATGTCGACCAAGAGCTCGACA TTAACCGCTGTCCGATTAAGCGTGGACGCCATCGTCCGACAGCTTCTCAAGGACGATTCATCGACAAACAAAAGTGTCTCAC CGAGCGACAAGATPAGAGGGAAGTCCGATAACGTCCTTCCGAAAGGTTGGTGAAGAAAGTGAAGAAATTAATGCGCGCAGCTCT GAAATGCAAGCTTACCCCAACGGAAAGTTGACAACTCACCAAGGCAAGAAAGAGGAGGACTGTCCGAAATGGATAAGGCGGTT TCATCAAGCGACAATGGTGGAAAATCGGCAAAATACCAAGCATGTGGCACAGATTCGACTCCCGTATGAACACCAAGTACGAC GAGAACGATAAGCTGATCCGCGAGGTCAAGTGTACCTCAACCAAGCAAACTGTGTGACAGACTCCGAAAGGACTTCCAAATCTTA CAAGGTCGCGGAAAATCAACAACACTACCACCGCTCATGACGCATACCTGAAACGCTGTGGTCCGGAATGCCCCATCAAGAAATACC CTAAAACCTCGAAAGCGAAATTTGTGTACGGCGACTACAAAAGTGTACGATGTCGGAAGATGATCGGAAAATCCGAGCAGGAGATCGGA AAGCGACTGTAGTACTTTTCTACTCGAAATCATGAACCTTCTCAAAACCGAAAATCACTTGGTAAATGGCGAGATCAGAAA GGCCCGCTGATCGAAAACCAACCGGAAACCGGTGAAATCGTGTGGGCAAGGGCCCGGATTTCCGTACTGTGAGAAAAGGCTCTTT CCAATGCCGCAAGTGAATATCGTCAAAAAGACTGAGGTGACAGCTGGCGGATTTCCAAGAAATCGATCTCCCAAGAGGAACTCA GATAAGCTCATCGCGGGAAAAGGATTTGGACCTTAAGAAAGTACGGAGATTTGATAGCCCAACTGTGGCCTACTCTGTGTGCTGCT GGTGGCCAAAAGTCGAAAAGGAAAAGTCGAAAGGTTGAAAATCCGTGAAAGAACTCTTTGGAAATCAATCATGGAGCGGTCTGTCAT TTGAAAAGAACCCCAATCGACTTCTTGGAGCCAAAGGATACAAAAGTCAAGAAAGACTGATCAATCAAGCTCCCTAAGTACAGC CTGTTCGAACTGGAGAACCGGAAACCGGATGCTGGCTTCCGCGCGGCTGCTGCTTCCGAAAGGCAATGAGCTGGCCCTCCCAATCGAA ATACGTGAACTTCTCTACCTTGCCTCCCAATACGAAAAGTTGAAGGGTCAACCGAGGACAAATGAGCAGAAAACAGCTCTTTGTG AACAAACACAAACTACTGACGAAATCATCGAAACAAAATCAGCGAGTTTCAAGCAAGCGCTCAATCTTGGCGGACCGCAACCTGGAT</p>	

Description	Sequence	SEQ ID NO:
	AAAAGTGTGTCGGCGTACAACAAGCACCAGGTAAGCCGATACGGGGAACAGGCTGAGAACAATCAATTCACCTCTTCACTCTCACTAA TCTGGGAGCCCCCGCCCTTCAAGTACTTTGATACCTACCATCGACCAGGAAAGAGATACACGAGACCAAGGAAAGTGTCTCGATGCCA CCCGTGATCCACAGTCCATTAAGTGTGTACGAAACGCGAAATCGATCTGTACAGCTCGGAGGAGATGGGTACCGGAAAGAA CGAAAGTCAAGTCCGATACAAGGATACGACGACGAGACTACAAGGACCATGACATCGACTTAAGAGACGACGACGAAAGCCCGC TGGAGCCGTGGATCCGGACCGCCGACCGCTTGGATGACTTCGACCTTGACATGCTGGATCCGACGCACTTGTATTTGATG TCGATATGCTTGGCAGCGACGACTGGACGATTTCCAGCTTCGACTCGGATCGGATCGGCTCGACGACTTCGATCTGGATATG CTGTGATAATAGGCTGGAGCCTCGGTGGCCATGCTTCTTGGCCCTTGGGCTTCCCGCCAGCCCTCTCCCTTCTCCGACCCGTA CCCCCGTGGTCTTTGAATAAAGTCTGAGTGGCGGCTCTAGA	45
Plasmid sequence of dCas9-KRAB- miR122	TCAAGCTTTTGGACCTCGTACAGAAGCTAAATACGACTCACTATAGGGAATAAGAGAGAAAAGAAGAGTAAGAAGAAAATAAAGA GCCACCATGGACAAGAACTCAATCGGACTCGCAATCGGAACCAATTCAGTCGGTGGGCACTCATACCCGATGAATACAAGGT GCCGTGAAAGAAGTTCAAAGTCTTGGTAACTGACAGACATTCGATCAAAAAGAAACCTGATCGGAGCCTTGTCTTGTGATTCAG GCGAAACCGCCGGAAGCTACCCGGTTGAAACGAACTGCTAGACCGCTACACGCGCCGCAAGAACCGGATCTGCTACCTCCAAGAA ATCTTCTCGAACGAAATGGTAAAGTGGACGACTCGTCTTTCACCGGCTCGAGGAGTCAATTCCTTGTGGAGGAAATAAGAAGCA CGAAAGACACCCGATCTTCGGCAACA TCGTGGACGAAGTCCGCTACACGAAAGATACCCGACTA TCTACCATCTCCGGAAAGAAC TCGTGGATAGCACCCGATAAGCCGATCTGCGACTGATCTACCTCGCGCTGGCCCATATGATTAAGTTCGCGGGCACCTTCTCATC GAAAGGACCTGATCCAGAACCTCGGACCTGGATAAGCTGTTTTATCCAGCTGGTGACAGCTTACAATCAATGTTTTGAAAGAAA CCTATCAACCGCTCGGGTGGACGCAAGGCCATCTGAGCGCGGCTCAAAAATCCAGACGCTGGAATACTGATAGCC AACTGCCGGGAGAAAACCGCCCTGTTTGGAAATCTTA TCGCCCTGTCCCTGGACTGACCCCAACTTCAAGTCGAACTTC GACTTGGCCGAGGATCGAAGCTCCAGCTCAGCAAGAACACTCAGACGATGACCTCGATAACCTGTGGCCAGATCGGTGACCA GTA TGTGATCTCTTGGCGGCCAAGAACCTGTCAGACGCAATTCCTGCTCCGACATCCCTGCGGGTGAATACTGAGATCACTA AAGCCCCATTGAGCGGCTCGATGATCAAAAAGATACGACGACACACAGGATCTGACTCTCC TCAAGGCACCTGGTCCGCCAACAG CTCCCGGAAAAGTACAAAAGAGATCTTCTTTGACCAATCCAAAACGGATACGCTGGTTACATAGACGCGGAGCGTCAAGAAAGA GTTCTACAAAGTTCAAGCCCTATCCTGGAAAAGATGGACGGGACCGAGAACTCTCTGTTAAGCTCAATAGGGAGGATCTGTGC GCAAGCAACGCAGTTCGACAA TGGAAAGCATCCCCCATCAGATCCACTGGGGGAGCTCCACGGGATCCTGAGGCCCCAGGAAGAT TTCTA CCCCATTTCTGAAAGGACAA TAGAGAGAAA TCGAAAAGATCCTGACTTTCCGAAATCCCGTACTACGTGGCCCCCTCCACG GGGAACTCACGGTTTCC TGGATGACTCGCAAA TCCGAA GAAACCAATACCCCTGGAATTTTCGAGGAGGTGGTCGATAAAGCGG CCTCAGCCCCAGTCTCATCGAAAAGATGACCAACTTTGCAAGAACCTCCCAATGAGAAGGTGCTGCCAAAACATAGCCTGTG TACGAGTACTTTACTGTGTATAACGAACTCAAGAGTGAATACTGTAACCGAGGAAATCGCAAGCCCGGCAATTTCTGTCTGGCGGA ACAGAAGAGGCAATGTGGACTTGTCTTTCAAACCAACCGGAAAGGTGACCGTGAACAGCTCAAGGAAGATTACTTTAAGAAAGA TCGAGTGTTCGATAGCTCGAAA TTTCCGGGGTGGAAAGATCGCTTCAA TCGAAGCTTTGGGACGTACCACGATCTGCTTAAGATC ATTAAGGACAAAGGATTTCTTGACAAACGAAAGAACGAGGATATTTCTCGAGGATATCGTCTTACCCCTGACTCTGTTTGGAGTAG AGAAAATGATCGAGGAGAGATGAAAACCTTACGCAACCTTCTCGACGATTAAGGTGATGAAACAGTGA AAAAGCGGTAGATACACTG GTTGGGAAAGGCTGTGAGAAAAGTGA TCAACGGAATAGGCAAGCTCCGAAAAACCACTCTCTGGATTTCTCAAAGTCCGAC GGTTCGCCAACCCGCAACTTCA TCGACTGATCCACGATGATTCCTGACTTCAAAGAGGATTC AAGAGGACCAAGTGTCCGG ACAAGGAGACTCACTCCACGAGCATATCGCTAATCTCGCCGATCGCCAGCTATCAAGAAAGGGAATCTTGCAGACTGTCAAGGTGG TGGACGGAAC TGGTGAAGTGTG TGGAAAGGCATTAAGCCGGGAATATCTGTGATCGAGATGGCGGAGGGA AAAACCCAGACCCCAAGAAA	45

Description	Sequence	SEQ ID NO:
	<p>GGACAGAAAAACAGCCGGGAACGCATGAAGCGCATCGAAGAGGGAATCAAAAGAGCTTGGGAGCCAAAATCCTCAAAGAACACCCCTGT GGAAAAATACCCAACTGCAGAAATGAGAAGCTTTACCTGTATTACCTCCAAAACGGCGCGACATGTACGTTGACCAGGAAATGGACA TTAACCGGCTTTCCGACTACGATGTGGAGCTATCGTCCCGAGTCCCTTCCGAAAGACGATTCGATCGACAAATAAGGTCCTGACT AGATCAGACAAGAAATCCGGGAAGTACAGCAACCGTGCCTAGCGAAGAGGTCGTTAAGAAGATGAAGAAATACTGGCCCGCAGCTGT GAACGGAAAGCTTATCACTCAGCCAAAGTTCCAGCAACTCACCAGGCAAGAAAGCGGATTTGCGAGCTCGACAAAATGGCT TCAATCAAGCCGCTCGTGGAACTCGCCAGATTAATGATGTGGCCAGATCTGGACAGCCGATGAACTACTAAGTATGAT GAGAAATGACAAGCTCATCCGGGAGGTGAAGTTCATACCTGAACTGAGTCCGACTTCCGGAAGGACTTCCAAATCTTA CAAAGTCAGAGAAATCAACAAATTAACATACCGCATGACCCCTACTTGAATGCAGTGTGGTACTGCCCTCATCAAGAAAATACC CAAAGCTTGAAGCGAGTTGTCTACGGAGCTACAAGGTGTACGACGTCGGAAGATGATCGCCAAAATCGGAACAGGAAATGGG AAGCGCACCGCTAAGTACTTCTTACTCGAATATCATGAAATTTCTTCAAGACCCGAGATCACGCTTGCAAAATGGCGAAAATCCGGAA GGGCCCTCATCGAAAACCAACGGGAAACCGGAGAAATCGTGTGGACAAAGGTTCGGATTTGGGACCGTCCGAAAGGTTCTTA GCATGCCTCAAGTGAACATCGTCAAGAAAACGGAAGTGCAGACTGGAGGCTCAGCAAGGAGTCCATCTCCCGAAAACGCAACTCC GACAACTGATCGCACGCAAGAAAGACTGGGACCCGAAAGAAATACGGAGCTTCGATTCGCCGACTGTGGCTTACTCGGTCTGGT TGTGGCCAAAGGTGGAAAAGGAAAGTCCAGAAAGTGAAGTCCGTCAAGAGCTCCTCGAATCACCATCAAGGACGGTCAAGCT TCGAGAAAACCCCAATTGACTTCTTGGAGGCAAGGGGTACAAGAGGTGAAGAAGGATCTGATCATCAAACTGCCGAAGTACAGC CTCTTTGAGCTCGAAAACGGACGCAAAAGGATGCTGGCTCCGCCGGAGACTGCAAAAGGAAAGAGCTTGCCTTGCCTTCCAA GTACGTGAACCTTCTGTACTGGCATCCACTACGAAAACCTGAAGGATCGCGGAGGACAAACGAAACAGAAAGCAGCTGTTGTGG AACAAACAAGCATATCTGGATGAAATCACTGAAACAAATCAGCGAATCTCAAAAAGGTTGATTTGGCCGACGCCAACCTGGAT AAAGTCTTTCCGCCACAACAACAATCGCGACAAGCCGATCCGGGAGCAGCCGAAACAATCACTACCTGTTTACCCCTGACTAA TCTGGTGGCCCCGGCTTTCAAATACTTCGATACCACGATCGACCGGAAAGAGATACACCAGCAACAAAGAGGTTGTGGACCGGA CCCTCATCCACCAATCTATACCAGGCTCTATGAAACTAGGATCGACTCAGCCAGCTGGGAGGCGATGCCCTTACGATGTC CCGGACTACGCTCGTGGGATCCGGATCCGAAAGAAAGGAGCGGAGGTCGAGGACCCAAAAGAAAAGCGCAAAAGTGGATGGGAT CGGTAGCGGTTCCAAACGGTTCTCGGTGGCGGGAGGCGGCAATGGAATGCTAAAGTCACTTACCGCTGGTTCGGGACGCTGGTGA CTTTCAAAGATGTGTGGATTTCACTCGTGAAGAAATGGAAATGCTGGACACTGCCCAACAGATCGTCTACCCCAACGTCATG CTTGAACACTACAAAACCTCGTGTGCTGGATATCAGCTGACCAAGCCGACGCTGATCTTGAGACTGGAGAAGGGCGGAAAGAAC TTGATAATAGGCTGGAGCCTCGGTGGCCATGCTTCTTGCCCTTGGCCCTCCCGACGCCCTCCTCCCTTCCCTGCACCCCGTACC CCCCAAAACCCATTGTACACTCCAGTGGCTTTTGAATAAAGTCTGAGTGGCGGCTCTAGA</p>	
DNA Plasmid sequence of dCas9-VP64- miR122	<p>TCAAGCTTTGGACCTCGTACAGAAGCTAAATACGACTCACTATAGGAAAATAAGAGAGAAAAGAAAGATAAGAAAATAAAGA GCCACCATGGACAAGAAATCAATCGGACTCGCAATCGGAACCAATTCAGTCGGTGGCGACTATTACCCGATGAATACAAGGT GCCGTGAAAGAAGTTCAAAGTCTGGGTAACTGACAGACATTCGATCAAAAAGAACTTGTGCGAGCCCTGCTGTTTGTGATTCAG GCGAAACCGCCGAAGCTACCCGGTTGAAACGAATCTAGACCGCCGCTACACGCGCCGAAGAACCGGATCTGCTACCTCCAAAGAA ATCTTCTCGAACGAAATGGTAAGGTGGACGACTCGTTCTTACCGGCTGAGGAGTCACTTCTGTGGAGGAAAGATAAGAAAGCA CGAAAGACACCCGATCTCGGCAACAATCGTGGACGAAGTCCGCTACACGAAAAGTACCCGACTACTACCAATCTCCGGAAAGAAC TCGTGGATAGCACCGATTAAGCCGATCTGCGATGATCTACCTCGGCTGGCCCAATATGATTAAGTTCGGGGGACCTTCCCTCATC GAAGGGACCTGATCCAGACAACCTCGGACGTGGATAAGCTGTTTATCCAGCTGGTGCAGACTTACAATCAATTTGTTGAAGAAA CCCTATCAACCGCTTGGGGTGGACGCAAAAGGCCATCCTGAGCGCGGCTGTCAAAAATCCAGACGGCTGGAATACTGTAGGCC</p>	46

Description	Sequence	SEQ ID NO:
	<p>AACTGCCGGGGGAGAAAAAAGGGCCCTGTTTGGAAAATCTTATCGCCCTGTCCCTGGGACTGACCCCAACTTCAAGTCGAATTC GACTTTGGCCGAGGATCGAAGCTCCAGCTCAGCAAGACACCTACGACGATGACCTCGATAACCTGTGTGGCCAGATCGGTGACCA GTATGCTGATCTTCTTGGCGGCCAAGAACCTGTAGAGCAATTCGTCTCCGACATCTCCGAGTCCGCGGTGAATCTGAGATCACTA AAGCCCAATTGAGCGGTCGATGATCAAAAATACGACGAGCACCCAGAGATCTGACTCTCCTAAGGCATGTCGCGCCCAACAG CTCCCGAAAAAGTACAAAAGATCTTCTTGGACCAATCCAAAAACCGATACCTGTTTACATAGACGCGGAGCGTCCACAGAAGA GTTCTCAAAGTTCAAGCCCTATCTTGGAAAATGAGCGGACCGGAACTCTCTGGTTAAGTCAA TAGGGAGGATCTGCTGC GCAAGCAACGCATCGACAAATGGAAGCTCCCATCAGATCCACCTGGGGAGCTCCACGCGATCTGAGGCGCCAGGAAGAT TTCTACCCATTTCTGAAGGACAAATAGAGAGAAAATCGAAAAGATCTGACTTTCCGAATCCCGTACTACGTGGCCCCGCTCGCACG GGAAACTCACGGTTTGCCTGGATGACTCGCAAAATCCGAAGAAAACCAATACCCCTGGAAATTTCCGAGGAGGTGGTCGATAAAGGG CCTCAGCCAGTCTGATCGAAAAGAAATGACCAACTTTGACAAGAACCCTCCAAAATGAGAAGGTGCTGCCAAAACATAGCCTGCTG TACGAGTACTTTACTGTATAACGAACCTACCAAGGTGAAAATACGTGACCCGAGGAAATCGCRAAGCCGGCATTTCTGTCCGGCGA ACAGAAGAGGCAATGTGGACTTGTCTTCAAAAACCAACCGAAGGTGACCTGAAAACAGCTCAAGGAAGATTAATTTAAGAAAGA TCGAGTGTTCGATAGCTTCGAAAATTTCCGGGGTGAAGAATCGCTTCAAATGCAAGCTTTGGACGTACCACGATCTGCTTAAAGATC ATTAAGGACAAAGGATTTCTTGACAAACGAGAAACGAGGATATTTCCGAGGATACTGCTGACCTGACTCTGTTGAGGATAG AGAAATGATCGAGGAGATTTGAAAATTTACGACACCTCTTCGACGATAAGGTGATGAAAACGCTGAAAAGGCGTAGATACACTG GTTGGGAAAGGCTCGAGAAAGTGTCAACGGAAATAGGGACAAGCAGTCCGGAAAACCATCTCTGGATTTCCCTCAAAGTCCGAC GGTTTCGCCAACCGCACTTCATGCAGTGTACACGATGATTCCTGACCTCAAAGAGGATATCCAGAAAGGCACAAGTTCCTGG ACAAGGACTCACTCCACGAGCATACTCGTAAATCTCGCCGATCGCCAGCTATCAAGAAGGATACTCAAGAGGAAATCTGACACTGTCAAAGTGG TGGACGAACTGGTGAAGTATGGGAAGGATAAGCCGGAGAAATCTGTGATCGAGATGGCGAGGAAAACCAAGCACCCAGAAA GGACAGAAAACAGCCGGAAACGATGAAGCGCATGAAGCGCATCAAGAGGAAATCAAAAGACTTTGGAGCCAAAATCTCAAAGAACACCTGT GGAAAATACCCAACTGCAGAAATGAAAGCTTTACTGTATTAACCTCCAAAACCGGCGGACATGTACGTTGACCCAGGAAATGGACA TTAAACCGGCTTTCCGACTACGATGTGGAGGTTTACTGCTCCCGAGTCTTCTGAAAAGACGATTCGATCGACAAATAAGGTCCTGACT AGATCAGACAAAGATCGGGGAAAGTCAACAACGTCCTTAGCAAGAGTCTTAAGAAGAATGAAGAAATTAAGTGGCGCCAGCTGCT GAAAGGAAAGCTTACTCAGCGCAAGTTCCGACAACTCACCAAGGCAAGAAAGGCGGATTTGTCGGAGCTCGACAAAAGCTGGCT TCACTAAGCCGCTGCTCGAAACTCGCCAGATTAAGCATGTGGCGAGATCTTGGACAGCCGATGAA TACTAAGTATGAT GAGAAATGACAAGCTCATCCGGAGGTGAAGTCACTACCTGAAAGTCCAGCTGGTGTCCGACTTCCGGAAGGACTTCCCAATCTTA CAAAAGTCAGAGAAATCAACAAATTAACCATCAGCGCATGACCCCTACTTGAATGCAAGTGGTGGTACTGCCCTCATCAAGAAATACC CAAAGCTTGAAAGCGAGTTGTCTACGGAGACTCAAGGTGTACGAGTCCGGAAGATGATCGCCAAAATCGGAACAGGAAATGGGG AAGCGCAAGCTTACTTCTACTCGAATATCATGAAATTTCTCAAGACCAGATCACGCTTCAAAATGGCGAAAATCCGGAA CGGCCCCCTCATCGAAAACCAACGGGAAACCGGAGAAATCGTGTGGACAAAGGTCGGATTTTGGCAGCCGTCGAAAAGGTTTA GCATGCTCAAGTGAACATCGTCAAGAAAACCGAAGTGCAGACTGGAGGCTTCAGCAAGGAGTCCATCTCCCGAAAACCGCAACTCC GACAAAATGATCGACGCAAGAAAAGTGGGACCCGAAAGAAAATACGGAGGCTTCGATTCGCCGACTGTGGCTTACTCGGTCCTGGT TGTGGCCAAAGGTGGAAGGAAAAGTCCAGAAAGTGAAGTCCGTCAAGGACTCTCCGAAATCAACATCATGGAACGGTCAAGCT TCGAGAAAACCCCAATGACTTCTTGGAGGCAAGGGGTACAAGAGGTGAAGAGGATCTGATCACTCAAACTGCCGAAAGTACAGC CTCTTTGAGCTCGAAAACCGACGCAAAAGGATGCTGGGCTCCGCGGAGAGCTGCAAAAGGGAAAACGAGCTTGGCTTGGCTTCCAA GTACGTTAACTTCTGTACTTGGCATCCCACTACGAAAAAATGAAAGGATCGCCGGAGGACAAACGAAACAGAAAGCAGCTGTTTGTGG</p>	

Description	Sequence	SEQ ID NO:
	AACAAACAAGCATTATCTGGATGAAATCATCGAACAAAATCAGCGAATTCTCAAAAAGGGTGAATCTTGGCCCGACGCCAACCTGGAT AAAGTGCTTTCGGCTACAACAACAACATCGCGACAAGCCGATCCGGGAGCAGGCCGAAAACATCATTCACCTGTTTACCCCTGACTAA TCTGGGTGCGCCCGCGCTTCAAAATACCTCGATACCACGATCGACCGGAAAGAGATACACCAGCACCAAAGAGGTGTTGGACCCGA CCTCATCCACCAATCTATTACCGGCTCTATGAAACTAGGATCGACCTCAGCCAGCTGGGAGGCGATGGATCCCCAAAGAAAG AGAAAAGTGTCTCCGACTACAAGGATCATGATGGGACTATAAAGATCATGATATTGATTACAAAGGACGACGACCAAGGCCGC TGGAGGAGGAGGTTCGGCCCGCCGATGCTCTCGACGACTTCGACCTCGACATGCTGGGATCCGACGCCCTGGACGACTTTGATC TGGATATGCTGGGCTCGGACGCCCTTGATGACTTCGATCTGGACATGCTGGGTCGGATGCACGACGACCTTCGACCTTGATATG CTGTGATAATAGGCTGGAGCCTCGGTGGCCATGCTTCTGGCCCTTGGGCTCCCGCCAGCCCTTCCCTTCCCTGACCCGTA CCCCCAACACCAATGTCACACTCCAGTGGTCTTGAATAAAGTCTGAGTGGGGCTCTAGA	

Table 9: sgRNA constructs

Description	Sequence	SEQ ID NO:
VEGF V1 sgRNA construct sequence	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAAATACGACTCACTATAGGGTGTGCAGACGGCAGTCACTAGGGTTTTAGAGCTAGAAA TAGCAAGTTAAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC	101
VEGF V2 sgRNA construct sequence	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAAATACGACTCACTATAGGGAGCAGCGTCTTCGAGAGTGAGGGTTTTAGAGCTAGAAA TAGCAAGTTAAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC	102
VEGF V3 sgRNA construct sequence	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAAATACGACTCACTATAGGGTGAAGTGTGGTTTTAGAGCTAGAAA TAGCAAGTTAAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC	103
VEGF V4 sgRNA construct sequence	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAAATACGACTCACTATAGGGTGGAGTGTGGTTTTAGAGCTAGAAA TAGCAAGTTAAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC	104
sgRNA plasmid template	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAAATACGACTCACTATAGGGTTTTAGAGCTAGAAA TAGCAAGTTAAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC AAATAGCAAGTTAAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGTGC TTTTAGAGCTAG	105
Moderna sgRNA 5' UTR	TCAAGCTTTTGGACCCCTCGTACAGAAAGCTAAATACGACTCACTATAGGG	106
Synthetic sgRNA guide scaffold sequence	GTTTTAGAGCTAGAAAATAGCAAGTTAAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCCG GTGGTGC	100

Table 10: Synthetic polynucleotides encoding a CRISPR related protein and synthetic sgRNA

Description	Sequence	SEQ ID NO
<p>mRNA sequence of dCas9- HA tag- 2xSV40NLS- KRAB</p>	<p>GGGAAUUAAGAGAGAAAAAGAGUUAAGAAAGAAAUUUAAGAGCCACCAUGGACAAGAAAUAUCAAUUGGACUUUGCUAUCGG AACUAAACAGGUGGUAUGGGCCGUCAUUAACUGACGAAUACAAGGUGCCUCAAGAAAGUUAAGGUCCUGGGAAAUAUACCGAUAGAC ACUCCAUCAAAAAGAACCCUGAUCGGGCACUGUUGUUCGACUCGGGAGAGACUCGACAGAAACAAACAGGUCUACGCCACUCCGAGG CGCGGUAACCCCGGAGGAAGAACCGCAUCUGCAUCCUCCAGAAUUUUCAGCAACGAGAUUGGCAAGAGUUCGUAUUCGUUCUU UCACCGCUUGAGGAGUCGUUCUUGCAGGAGGACAAAAGCAUGAAAAGACAUCCGAUUCUGGAAACAUUGGACGAAAGUCG</p> <p>CAUACCAUGAAAAGUAACCCUACCAUCCAUUCAGAAAAGAAACUCGUCGAUUCAAUGAUAAAGCCGACUUUGCGGCUGAUCUAC CUGGCUUGGGCCACAUGAUCAAGUUUCGGGUCACUUUCUACUGAGGGUGAUUCAACCCGGACAUAUCCGACGUUGACAAAACU CUUCAUCCAAUCUGGUCCAGACGUAACACAGCUGUUCGAAAGAAAUCGCAUACCGCAAGCGGAGUGGACCCCAAAACCCAUUCUGU CGGCCGCCUCUGAAGUCGCCUGGAAAUCUGAUUGUCAGUCUCCCGGGGCAAAAAGAAUUGGCCUGUUUGGAAAACCCUC</p> <p>AUCGCACUGUCCUCGGGUCGACUCCCAACUUCAAAUUCGAAUUCUUGGCUUGGAGGAGUACAAGCUGCAACUCUCCAAAGACAC UUA CGAUGAUGACCCUGGACAAUCUCCUGGCGCAGAUCCGGGAAUCAGUAUGUCUGACUUGUUCUUGGGCCGCAAGAACCCUGUCUGAUG CCAUCCUGUCUCCGAUUAUCCUGAGUGAACAACUGAGAUCAACAAAGGCCUCUGAAGGCCUCUGAUAUCAAACCCUACGAUGAA CACCAUCAGGACCCUACUCUUCUGAAAGCUUUGGUGGGCAGCAGUUCCGGAAAAGUACAAAGAGAUUCUUCGACCCAGUCGAA</p> <p>AAAACGGUACCGCCGGAUACAUAUGGCGCGCAAGCCAGGAGAAUUCUAUAAUUUAUCAAACCCGAUUCUGGAGAAAGAUUGGACG GCACUGAAGAAUCUUCUGGUCAAGCUGAAUCGAGAGAUUCUGUCCGGAAGCAGGACCUUCGACAAUUGGUCUAUCCUACCCAA AUCCAUCUCGGGAGCUGCAUGCGAUUUCGAGGCCAGGAGGAUUCUACCCAUCUUCGAAAGACAUCGCGGAGAAAUCGGA AAA GAUUCUGACGUUCCGCAUCCCAUAUACGUCGGGCCAUUGCGCGGGUAAAUCGAGAUUCGCCUGGAGCGGGAAGUCCGAAAG</p> <p>AAACCAUCA CGCCGUGGAAUUUCGAAAAGAGUGGUCGACAAAGGAGCCAGCCAGCAGUCCUUAUUGAGCGCAUUCGAAUUUCGAC</p> <p>AAAAAUCUGCCGAACGAGAGGUCUGCCGGAAGCAUUCACUGCUGUACGAAUACUUUACCGUGUACAACGAAACUGACCAAGGUGAA GUACGUCACCCGAGGGAAUGAGAAAACCCUGUUUCUGAGCGGAGAAACAGAAAGAGCCAUUUGUUGACCCUUCUUAAGACUAAAUC GCAAUGUACCCGUGAAGCAUCUUAAGAGGAUUAUCUCAAAGAAUCGAAUUGUUCGACUCCGUGGAAAUCAGCGCGUGGAGGAU</p> <p>AGAUUCAACCGGUCUCCUGGACUUAACACGACCCUUAAGAUCAUCAAGGAUAGGAUUUUCUGACAUAAGGAAACGAAAGA UAUCCUGGAGGACAUCCGUUCGACUCUACCCUUCUUGAGGACCGGGAGAUUCGAGGAGAGACUCAAGACCUUACGGCACCUUGU</p> <p>UUGACGACAAAAGUGAUGAAGCAACUUAACCGCAGGGCCUACACCGGUGGCGAGACUGUACGCAAGUUUGAUAACGGAUUUAGA GAUAAACAGUCCGGAAAACCAUCCUGGACUUCCUGAAGUCCGAUUGCCCAACCGCAAUUUUCAUGCAGCUUCCAUUGACGA</p> <p>CUCAUUGACUUUCAAGGAGAUUCCAAAAGGCCCAAGUGAGCGGCCAAGGGGACUCCUUAACGAAACAUCGCAAAAUUUGGCCG GAUCACCGCAUUAAGAAAGGAAUCCUGCAGACUUGAAGGUGGAGCAGCGUUGGAAAGAGUUGGACGGCACAAAGCCGGAA</p> <p>AACAUCUGAUCGAGAUUGGCCAGAGAGAAACAGACGACUCAAAGGGCCAGAAACUCGCGGCAACGCAUCGUAAGAAUAGAAGA GGGAAUUAAGGAAACUGGGACUGCGAUCUUGAAGGAGCACCCUGUGAAAUAUCUACAUCAGAACGAGAAAGCUGUACUUGUAUC</p> <p>AUCUUAACAAACGGCAGGGACAUGUAUUGUCGACCAAGAGCAGUACAUUAAACCGCCUGUCCGAUUAUGACGUGGACGCCAUUCGUGCCG CAGAGCUUUUCUCAAAGGACGUUCCAUCCGACAAACAAAGUGUUCACCCGACGCAAGAAUAGAGGGAAGUCGGAAUAAACGUCCCUUC</p> <p>GGAAAGGUGGUGAAAAGAUAGAAUUAUCUGGCGGCAGUCCUGAAUGCAAAAGCUACCCCAACGGAAAGUUUGACAAACCUCA CCAAGGCAGAAAAGAGGAGGACUUGCCGAAUUGGAUAAGGCCGCUUUAUCAAGCGGACAAUUGGUGGAAAUCUGGGCAAAUUAACCAAG CAUUGGGCACAGAUUCUGGACUCCCGUAUGAAACCAAGUACGAGAAACGAUAAAGCUGAUCGCGGAGGUAACCGGUAUCACCCU</p>	<p>51</p>

	<p>CAAAAGCAAACUUUGUGUCAGACUUCCGGAAAGGACUUCCAAUUUUAACAAGGUCGCGGAAAUACAACAUCACCACCGCUCAUGACG CAUACUGAAACGCUGUGGUCGGGACUGCCCUCAUCAAGAAAGUAACCCUAAAUCGAAAGCGAAUUGUUAUUGUGUACGGCGACUACAAAGUG UACGAUUCGGAAAGAUUCGCGAAAUCGAGCAGGAGAUUCGAAAGGGCAGUCGUAAAGUACUUUUUUAUCUCCGAAACAUCAGAA CUUCUAAAACCGAAAUACACCCUGGCUAAUGGCGAGAUCAAGAAAGCGCCCGUAGUAGAAACCAACGGCGAAAACCGGUGAAAUUCG UGUGGACAAGGGCCGGAUUUCGUACUGUGAGAAAGGUUUUUAUGCCGCAAGUAAUAUCGUCAAAAAGACUGAGGUGCAG ACUGGGGAUUUCCAAAGAAUCGAUCCUCCAAAAGAGACUCAGAUAAAGUCUACUCGCGCGAAAAGAAAGGAUUGGGAUCCUAAAGAA GUACGGAGAUUUUAUAGCCCAACUGUGGCCUACUCUGUGUCUGUGGCGCCAAAAGUCGAGAAAGAAAGAAAGUUCGAAAGUUGAAAU CCGUGAAAGAAACUCUUGGGAUUCACUAUCAUGGAGCGGUCUCAUUUGAAAGAAACCCAAUCAGACUUCUCCUGGAAAGCCAAAGGUAUC AAAGAGUCAAGAAAGACUGUAUCAAGUCCUAAAGUACAGCCUUAAGUUCGAAACUGGAGAAACGGAAAGGAAACGGUAGCUGGCUUC CGCCGGCAACUGCAAAAGGGCAUAGAGCUGGCCUCCUCCUAGAAUACGUGAAACUCCUACUCCUUGCCUCCAUUACGAAAAGU UGAAGGCUACCCGAGGACAAUGAGCAAAAAGCUCUUUUUUGAAACAACAACAACUACUCCUGGACGAAAUCAUCGAAACAAAUC AGCGAUUCAGCAAGCGGUCUUCUGGGGACCGGAAACUUGGAUAAAGUCUGUCGCGGUACAACAAGCACCGCGGAUAGCCGGAU ACGGAAACAGGUCGAGAACUACUUCACU UCGACCGCAAGAGAUACACGAGCAACAAGGAAUGUCGUAUCCACCCUGAUCCACAGUCUACUACUACUACUACUACUACUACUACUACU AUCGAUCUGACAGCUCGGAGGAGUCCUACU AAGAAAGGUGGAGGACCCGAAAGAAAGAGGAAAGUUCGCGGACGGAUCGAAAGCGGAUCGAAUUGGAUCGUAAGGGGAGGUGGCGGAG GUAUGGACGCAAAAUCAUUCGGCCUGGUCACGGACCUUGGUGACUUUAAAAGCUGUUUCGUAUUUACCAAGGAAAGAAUAGCCG AAACUUGUGGACACCGCCAGCAGUUCGUAUCGGAUUGGUAUCGGAUUCGGAUUCGGAUUCGGAUUCGGAUUCGGAUUCGGAUUCGGA CACUAAAGCAGAGUUCUUAAGACUGGAAAGGAGAAACCGUGAAUAAAGGUGGAGCCUCCUGGUGGCAUGUUCUUGCC CUUGGCCUCCCCCAGCCCCU t) tail</p>	<p>52</p>
<p>mRNA sequence of dCas9-NLS- FLAG-VP64</p>	<p>GGAAUUAAGAGAGAAAAGAGUAAGAAGAAAUAUAAAGCCACCAUGGACAAGAAAUAUCAAAUCGGAUUGGCUUUGCUAUCGGAACU AACAGCUGGGAUGGGCCGUAUUCUGACGAAUACAAGGUGCCUCAAAGAAUUAAGGUCUGGAAAUACCGAUAGACACUC CAUCAAAAGAAACCCUGAUCGGGGCACUGUUGUUCGACUCGGGAGAGACUCGAGAAAGCAACAGGCUCAAGCGCACUCGCGAGGCGCC GGUACCCGGAGGAAACCGCAUCUGCUACCCUCCAAAGAUUUUCAGCAACGAGAUUGGCAAAAGGUCGUAUUCGUUUUUCAC CGCUUGAGGAGUCGUUCUUGCAGGAGGACAAAAGCAUAAAGACAUCGGAUCUUCGGAACAUCGUGGACGAAAGUCGCAUA CCAUAAAAGUAACCUACU CUCUGGCGACAUCAAGUUUUGGGUCACUUUCUACUGAGGUGAUCUACCCCGGACAAUUCGACGUUGACAAACUCUUC AUCCAUCUGGUCAGACGUACACAGCUGUUCGAAAGAAAUCGUAUCCGCAAGCGGAGUGGACGCCAAAGCAUUCUGUCGGC CCGCCUUCGAAAGUCGCGUCUGGAAAUCUGAUUGCUCAGCUCGCGGCGAAAAGAAUUGGCCUUGUUUGGAAAACCUCAUCG CACUUCUCCUGGGUCACUCCAAUCAAUCGAAUUCUUGACUUGGUGGAGGAGUCAAAGCUGCAUCUCCAAAGACACUUAC GAUGUACUGGACAAUCUCCUGGCGAGAUCCGGGAUCAGUAUGCUGACCUUUCUGGCGGCAAGAAACCUUGUCUGAUGCCAU CCUGUCUCCGUAUCCUGAGUAGACACUGAGUACCAAGGCGCUCUGAGCGCCUAGUAUCAACCGCUACGUAACACCC AUCAGGACUACUUCUGAAGGUUUUGGCGGACAGUUCGGAAGAAUACAAGAGAUUUUUCGACAGUCGAGGAAAGAAAC GGUACGCCGGAUACAUAUGGCGCGCAAGCCAGGAAUUCUAUAAAGUUUAUCAAAACCGAUUCUCCUGGAGAAUGGACGGCAC UGAAGAUCUUCUGUCAAGUGAUUCGAGGAAUCUGUCUCCGGAAGCAGCGGAUUCUUCGACAAUUGGUCUAUCCUACCAAUUC AUCUCGGGAGCUGCAUUCGGAUUCGAGGCGCCAGGAGCAUUCACCAUCCUGAAAGACAAUCGGGAGAAAUCGAAAGAAU CUGACGUUCGCAUUCUACUACGUCGGGCCACUUGCGCGGGUAAUUCGAGAUUCGCUUGGAGUUCGCGGGAAGUCGAAAGAA CAUCACGCCGUGGAAUUUCGAAAGAGUGGUCGACAAAGGGAGCCAGCGCACAGUCCUUAUUGAGCGCAUGACCAAUUUCGACAAA</p>	

<p>AUCUGCCGAAACGAGAAAGGUCUCCUGCCGAAAGCAUUCACUGUCGUUACGAAUAUUUACCGUGUAACAACGAAACUGACCAAGGUGAAAGUAC GUCACCGAGGAAUGAGAAAGCCUGUUCUGAGCGGGAAACAGAAAGAGCCCAUUGUUGACUCCUUCUUAAGACUAUUAUCGCAA AGUACCCGUGAAGCAGCUUAAAAGAGAUUAUUUAAAAGAUUUAUUUUCGACUCCUGUGAAAUCACCGCGGUGGAGGAUAGAU UCAACCGUCCUUGGACUUACACGACCCUUAAGAUCAUAAAGGAUAAGGAUUUCUCGACAAUUGAGGAAAACGAAGAUAUC CUGGAGACAUUGUUCUGACUCGACCCUUCUUGAGGACCGGAGAUAGUACGAGGAGAGACUCAAGACCUCACCGCAUUGUUUGA CGACAAAGUGAUGAAACAUAUAAAACGACGGCCUAACCGGUGGGGACAGUCUACGCAAGUUGAUCACGGAAUUAAGAGAUA AACAGCCGAAAGACCAUCCUGGACUUCUGAAGUCCGAGUGGAUUGCCAACCGGAAUUUAUGCAUCUCCAUAGCAGCAGCUCA UUGACUUUCAAAGGAGUAUCCAAAAGCCCAAGUGAGCGGCCAAAGGGACUCCUUCACGAAACAUCGCAAAUUGGCCGGAUC ACCAGCGUAUAAAGGAAUUCUGCAGACCGUGAAGGUGGACGAGCGUGGAAAGUAGUGGACGGCACAAAGCCGAAACA UCGUGAUCGAGAUUGCCAGAGAAACAGACGACUCAAAAAGGCGCAGAAAGAACUCGCGAAACGCAUGAAGAGAAUAGAAGAGGGA AUAAAGGAAUCUGGGAUCGAGAUUUUAAAGGAGCACCCUUGUGAAAUAUCUACAUCAGAACGAGAAAGCUGUACUUAUCU UCAAAACGGCAGGACAUGUAUUGCAGCAAGAGCUCGACAUUAACCGCCUGUCCGAUUUAGCUGGACGCCAUCUGGCCGAGA GCUUUCAAAGGACGAUUCACAAACAAGUGCUCACCCGACGACAAGAAUAGAGGGAAGUCGGAUAACGUCUCCUUCGGAA GAGGUGGAAAAAGAUAGAAUUAUCUGGCGGACGUCUGAAUGCAAAAGCUCUACCCAAAGGAAUUGACAAACCCUACCCAA GGCAGAAAGAGGAGGACUGUGGAAUUGGAUAAAGGCGGUUAUCAAGCGACAAUUGGUGGAAACUCGGCAAAUUAACAAGCAUG UGGCACAGAUUCUGGACUCCCGUAUGAACACCAAGUACGACAGAAACGUAAGCUGUCCGAGGUAAGGUGAACCCUUCAAA AGCAAACUUUGUCAGAUUCCCGAAAGGACUUCUUAUUUACAGGUCGCGAAAUCACAAACUACCCACCGCUCAGCAGCAUA CCUGAACGUCUGUGCGGACUCCUUAACAAGAAUACGAAUUCGAAACGGAUUUUGUACGGCGACUACAAGUGUACG AUGCCGGAAGAUAGUCGGAUUCGAGAGGAGUCGGAAGGCGACUGCUAAGUACUUUUUUAUCUCGCAACAUCAGAAAUUC UUCAAAACGAAUACCCUGGUAUUGGCGAGAUACAGAAAGCCCGCUGAUCGAAACCAACGCGAAACCGGUAUUCGUGUG GGACAGGCGCGAUUUCGUACUGUGAGAAAGUCCUUAUGCCGCAAGUUAUCUGUCAAAGAAUUAUGUCAAAGAAUUGAGGUCAGACUG CGGUAUUUCCAAAGAAUCGACUCCCAAGAGAAUCAGAUAAUCUACCGCGGAAAAGGAUUGGGAUCCUUAAGAAUUCG GGAGAUUUGAUGCCAAUCUGUGCCUACUCUGUGUCUGGUGGCGCAAAGUCGAGAAAGGAAAUCGCAAGAAUUGAAUUCG GAAAGAACUCUUGGGAUACAUUAGUAGGAGGUGUCUAUUUAAAAGAAACCAUUCGACUUCUGGAAAGCAAGGAAUACAAAG AAGUCAAGAAAGACUUGAUCUACAAGCUCCUAAAGUACAGCCUGUUCGAAACUGGAGAACGGAAGGAAACGGAUUGGCUUCGCGC GGGAAUCUGCAAAGGGCAUAGAGCUGGCGCCUCCCAUCGAAUAUACGUGAACUUCUUAUCCUUGCCUUAUACGAAAAGUUGAA GGGCUACCCGAGGACAAUGAGCAAAACAGCUCUUUUGAACAACAACAACUACUGGACGAAUUCAGAAACAACAAACAGCG AGUUCAGCAAGCGGUAUUCUGGCGGACGGAACCCUGGUAUAAAGUGCUGUCCCGUAACAAGCAACCGCGUAAGCCGAUACGG GAACAGGUGAGAAACAUCUACUUCUACUCUACUUAUUGGAGCCCGCGCGCUUAAGUAUUAAGUACUUAAGCAGAAUACCGAUCG CCGCAAGAUACAGGACCAAGGAAUGUCUGAUGCCACCCUGAUCACAGUUCUAUACUGGUCUAUACGAAACGCGAAUCG AUCUGUCACAGCUCGGAGGAGUUGGUCACCGAAAAGAAACGGAUAGUCAGCUCGGAUUAACAAGAUACGACGGAGACUAACAAG GACCAUGACUACGUAUAAAGGACGACGACAAAGCCCGUGGAGCGGUGGAUCGGGACGCGCGGACCCUUGGAUGACUUCGA CCUUGACUUCUGGACCGCAGCUUUGAUAUUUUGAUCUGAUAUGCUUGGACGACGACUGGACGAAUUCGACCCUUCGAC UGUCGGAUCGGAUGCGUCGACGACUUCGUAUCUGGAAUUGUGUAUAAUAGGUCUGGAGCCUCGUGGCCUUCUUGCCCU UGGGCUUCCCGACCCUUCUUCGACCCCGGUCUUAUUAUAAAGUUCUGAGUGGGCGC++A (140nt) tail</p>	<p>53</p>
<p>mRNA sequence of dCas9- KRAB-miR122</p>	<p>53</p>

<p>GCUACGCGCCGCAAGAACCGGATUCUGCUAACCCUCCAAAGAAAUCUUCUCGAAACGAAAUUGGCUAAGGUGGACGACUCUGUUCUUCAC CGGUCGAGGAGUCAUUCUUGGAGGAGGAAUAAGAGGAGCAAGAAAGACACCCGUAUCUUCGGCAACAUCUGGACGAAGUCGGUA CCACGAAAAGUACCCGACUACUACCUCCGGAAAGGUCUGGUAAGCAACCGAUAAGGCCGAUCUCGGCAUCUACUACCUCCG CGCUGGCCAUUAUAAGUUUCGGCCACUUCUCUAGAAAGGACCGUAAUCAGACAACUCGGACUGGUAAGUCUGUUA AUCCAGCUGGUCAGACUUAACAUAUUGUUAAGAAAACCUUAACAACCGUCUGGGUGGACGCAAGGCAUCCUGAGCGC GGGCUUUCAAAAUCGACGGCUGGAAAUCUGAUAAGCCCAUCGCGGGGAGAAAAGAAAAGGAAAUCUUAUCG CCCUCUCCUGGAGUAGACCCCACUUAAGUAGAAUCUCAGUAUCUGCCGAGGAGGAAAGUCGCAAGGAAAGACACCUAC GACGUAACCUAGUAACCUUUGCCCAAGAUCCGUGACAGUAUCUUCUUGGCGCCCAAGAACCCUGUCAGACGCAAU UCUGUCUCCGACAUCCUGCGGUGAAUAUCUGAUAACAAAGCCCAUUGAGCGCGCUAUAAGAAAGUACGACGAGCAC ACCAGAUUCGACUCCUCAAGGCAUCUGUCCGCAACAGCUCGCGAAAAGUAAAGAGAAUUCUUCUUCGACCAAUCCAAAAC GGAUACGUGUUAUAUAGACGGGAGCGUACAAAGAAGUUUAACAAGUUAACAAGCUCUAACAAACCUUCGAAAAGAGACGGAC CGAGGAUCUCCUGUUAAGCUAAUAGGAGGAUCUGUCGGCCAAAGCAACGCCAUUCGACAAUGGAAACUCCCAUCAGAUCC ACCUGGGAGUCUCCGCAUCUUGAGGCGCCAGGAAGAUUUAUCCCAUUCUGAAGGACAAUAGAGAGAAAUCGAAAAGAU CUGACUUCCGAAUCCCGUACUACUGGGCCCGUCGACGGGAAACUCAACCGUUCGCGGAGUACUCGCAAAAUCGGAAGAAC CAUUAACCGUGGAAUUCGAGGAGGUGUCUAAGAGCGCCACAGCUCGUAUCGAAAGAAUACCAACUUUGACAAAG ACUCCAAAUAAGAGAGGUGUCGCAAAAUAAGCUCGUGUACGUAUCUUAUCUGUUAUAACGAAUCACAAAGGUAUUAC GUGACCGAGGAAUGCCGAAGCCGCAUUCUGUCGGGCAAGGAAAGGAAAUUGGACUUCGUAUAUACCAACCCGAA GGUACCGUAAAAGCUAAGGAAAGAUUAUCUUAAGAAGUUCGUAUUGGUAUUCGUAUGGAAAUUCGGGGGGAAGAUCCGU UCAUAGCAAGCCUUGGAGUACCACGACUCUGCUUAAGAUAUAAGGCAAGGAUUCUUCGACAAACGAAAGAAACGAGGAUUA CUCGAGUAUCGUCGACCUCGACUCUGUUUGAGGUAAGAUAUAGAAAUAUCGAGAGAAUAUAGAAAUCUUCGACACCUUCU CGUAAGGUAUAGAACAGUUAAGGCUAGUAUAUCUGGUUGGAAAGGCUAGAAAGCUGAAACGAGAAUAGGGACA AGCAGUCGGAUAAACUUCUGGUAUUCCUAAGUCCGCGGUAUCCCAAGCGAAUAUCGUAUGCAGCUGUACGAGUAUCC CUGACGUAAGAGGUAUCCAGAAAGCAAGUUCGCGCAAGAGAGACUCCAGCAGCAUAUCGCUAAUCUGCCCGAUC GCCAGUAUAAGGAAUUCUGCAGACUCUUAAGGUGGACGAAACUGGUAAGGUAUAGGAAAGGCAUAGCCGAGAAUA UCGUAUCGAGUUGGAGGAAACCAAGCAGCCCAGAAAGGACAAAGAAACAGCCGGGAACGUAAGGCAUCGAAAGGGA AUCAAAGACUUGGAGCCAAUCCUCAAGAACCCUUGGAAAUAACCAAUCUGCAAAUAGAAAGCUUAUCUGUUAUACCU CCAAACGGGCGCAGUAUGACGUAGCCAGGAAUUGGCAUUAACCGGUUCGAAUAGGUAUCGAGGACGUUCCGCGAGU CUUCUGAAAGACGUAUCGACAAUAAGGUCCUGACUAAGACAAAGAAUCGAGAAAGGUAAGGAAAGGAAAGGUAAGGAA GAGGUCUUAAGAAAGAAUUAUCUGGCGCCAGCUGAAACGGAAGGCUUAUCACUCAGCGCAAGUUCGAAACCUACCCAA GGCAGAAAGGCGGUAUUCGGAGCUGACAAAGCUGGCUUAUAAGCGCCAGCUCGUCGAAAUAUCGAAUUAUAAGCAUG UGGCGAGAUCCUGGACAGCCGUAUUAUAAGUAUGAAGAAUGACAAGCUUACCGGGAGGUAAGGUAUCACCCUGAAG UCCAGCUGGUCGACUUCGGAAGGACUUCCAAUUCACAAAGUCAGAAAUCAAUAUAUCUACCGCGCAUACGCGCAUCG CUUGAUUGCAGUGGUGGUACUGCCUACAAGAAUUAACAAAGGUAAGGUAUUGGUAUUCGAGGUAUAAAGGUAACG ACGUCGGAAAGUAUCGCAUUCGGAAACAGGAAUUGGAAAGGCGACCUGAAAGUACUUCUUAUCGAAUUAUAGAAUUC UCAAAGCCGAGUACGCUUAAGGAAUUGCGAAUCCGGAGCCCUUAUCGAAACCAACCGGAGAAACCGGAGAAUCGUGUG GGACAAGGUCGCAUUUGCGAACCGUUAAGGUAUUCGUAUUAUGAUUAUCGUAUUAAGGAAUUAAGGAAUUAAGGAAUUA GAGGCUUCAGCAUAGGUAUUCGCGCAUUCGCGUCCGAAAGGUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAA GAGGCUUCGAAUUCGCGCACUGUGCUAUCGCGUCCUUGGUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAAUUA CAAAGGCUCCUGGAAUACCAUAUGGAACGGUACAGCUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAA CAAGGAGCUCUGGAAUACCAUAUGGAACGGUACAGCUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAAUUAAGGAA</p>
--

mRNA sequence of dCas9- VP64-miR122	<p>AGGUGAAAGGGAUCUGAUCAUCAAAAUCGCGGAAAGUACAGCCUUCUUGAGUCUCGAAACCGGACGCAAAAAGGUAUGCUGCCUCCGCC GGAGGUGCAAAAAGGAAACGGAGCUUGCCUUGCCUCCAAAGUAAGUGAAACUUCUUGUAUCUGGCAUCCCAUCAGAAACACUGAA GGAUCCGCGGAGGACAACGAAACAGAGCAGCUGUUCUGGAAACAACAAGCAUAUCUGGAUGAAUUAUCGAAACAACAAUCAGCG AAUUCACAAAAGGUGAUUUGCGCGACGCCAACCCUGGAUAAAGUUCUUCGCGCAACAACAACUUCGCGCAAGCCGGAUCCCGG GAGCAGCCGAAACAUAUACCUUUAACCCUGACUAUCUGGUGCGCCCGGCUUUCAAAUAUCUGAUACACGUAUCACGUAUCGA CCGGAAGAUACACCAACCAAGAGGUGUUGGACGCGCACCCUCAUCCACCAUUAUUAACCGGCCUUAUGAAACUAAGGAUCG ACCUAGCCAGCUGGGAGGGGAGUCCUUAAGAUUCCGCGGACUAAGCCUCGUGGUAUCUGGUAUCUCCGAAAGAAAGCGG AAGUCAGGACCCAAAAGAAAGCGCAAGUGGAUGGUAUCGGUACCGGUUCCACCGGUUCCGUGGUGCGGCGGAGGCGGCCAU GGAUGCUAAGUCACUUAACCGCCUGGUCGCGGACGCGUGGUAUUAAGAUUGUUCGUGGAUUUACUGUGAGGAUUGGAAU UGCUGACACUGCCCAACAGAUUCGCAACGUAUGUUGAAACUAACAAACCCUGUGCGUGGGAUACAGCUGAGCC AAGCCGACGUGAUUCUGAGACUGGAAAGGCGGAAAGAACCUUGAUAAUAGGUCGGAACCCUGGUGGCCAUUCUUCUUGCCCUUG GGCUCCCCAGCCCUUCUUCUGCACCCGUACCCCCCAACACCAUUGUACACACUCCAGUGGUCUUCUUGAAUAAAAGUCU GAGUGGCGGC+A (140nt) tail</p>	54
	<p>GGGAAUAAAGAGAGAAAGAAAGUAAGAAAGAAUUAAGAGCCACCAUGGACAAGAAAGUACUCAAUUCGGACUCGCAAUUCGGAACC AAUUCAGUCGCGGUCGAGUCUAUACCGAUGAAUACAAGGUGCCUGGAAAGAUUCAAAAGUCCUGGGUAAACACUGACAGACAUUC GAUCAAAAAGAACCCUGACUGGAGCCUUCGUGUUAUGAUUACAGCGAAACCGCCGAAAGUACCCGGUUGAAACGAAACUGCUAGACGCC GCUACCGCGCCGCAAGAACCGGAUCUGCUAACCCCAAGAAAUUUCGAAACGAAAUUGGCUAAGGUGACGGAUCUGUUUUUAC CGGUCAGGAGUCAUUCUUCUGGAGGAAGAUAAAGAAAGCAACCCGCAUUCUGGCCAAUUCGUGGACGAAUGCGGUA CCACGAAAAGUACCCGCAUUAUCAUUCGCGGAAAGAGCUCGUGGAUAGCACCAGAAAGCCGUAUCUGCGACUGAUUCACUCUG CGUGGCCAAUUGAUUAGUUCGCGGCACUUCUUCAGAAAGGACCGUAUCCAGACAACACUCGGACGUGGAUAAAGCUGUUU AUCCAGCUGGUCAGACUUAACAUAUUGUUUAGAAACCUUAUACACGUCUGGGUGGACGCAAAAGCCAUUCUGAGCGC GGCGUGUAAAUCCAGACGGCUGGAAAUCUGUAAGCCCAACUGCCGGGAGAAAGAAAACGGCCUGUUGGAAAUCUUUACG CCUUGCCUGGGACUGACCCCAACUUAAGUCAAUUCGACUUGCCGAGGAGUGGAAAGCUCAGCUCAGCAAAAGACACCUAC GACGAUACCUUGUUGCCAGAUCCGUGGACAGUAUGCUGAUUCUUCUUGCGGCCAAGAACCCUGUCAGACGCAU UCUGCUCCGACAUCCUGGGGUAUACUGAGAUACUAAAAGCCCACUUGAGCGCGGUAUCAAAAAGUAUCAAAAAGUAACGAGCACC ACCAGAUUCUGACUCUCUCAAAGGCAUCUGGUCGCCCAACAGCUCGCCGAAAGUAACAAGAGAUUCUUUUUCCAAUCCAAAAC GGAUACGUGUUAUAUAGCGCGGAGCUCACAAAGAGUUAACAAGUUAACAAGUUAUCCUUGAUCUUGAUCUUGAUCUUGAUCUUGA CGAGGAUUCUGUUAAGCUCUAUAGGAGGAUCUGUCGCGCAAGCAACGCAUGUUGCAUUGGAAGCAUCCCAUCAGAUCC ACCUGGGAGCUCACCGCUUCUGAGGCCAGGAAUUAUCCCAUUGAAGCAUUAAGAGAAAGAAUAGAGAGAAAUUGAAGAAUUC CUGACUUUCCGAAUCCCGUACUACUGGGCCCGCUCGCAAGGAAACUCACCGGUUGCUGGAUACUCGAAAUCCGAAAGAAAC CAUUAACCCUGGAAUUCGAGGAGGUGUCGAUAAAGGGCCUCAGCCCAGUCGUAUCGAAAGAAUGAACAAUUCUUGAACAAGA ACCUCCAAAUGAGAGGUCGCAAAACAUAAGCCUGUACGUAUUAUCUGUUAUACGAAACUCCAGGUAUAAACCAACCGGAA GUGACCGAGGGAAUGCGCAAGCCGUAUUCUGUCGGCGGAAACAGAAAGAGGCAUUGUGACUUGUUAUAAACCAACCGGAA GGUGACGUGAAACAGCUCAAGGAAGAUUAUUAUAAAGAUUCGAGUUAUUAUUGGUAUUAUUCGGGGGUGGAAGAUUCGCU UCAUUGCAAAGCCUUGGGACUACACGAUCUGCUUAAAGUAUUAAGCAAGGAUUUCUUGGAAACGAAAGAACGAGGAUUAU CUCGAGGAUAUCUGUCCUGACCCUGACUCUGUUAUGGGAUAGAAAUUGAUGAGGAGGAUUAUUAUCGACACCCUCUUCGA CGUAAGGUGAUGAAAACGUCGUAAGGCGUAUAACUGGUAUAGGCGGUCGUAUAGCUACACGAAUUAAGGACA AGCAGUCCGGAAAAAACAUUCUUGGAUUCUUAAGUCCGCAAGUUCGCAACCGCAACUUAUCGAGCUGAUCCAGUUAUUC CUGAGUUCAAAAGAGGAUUAUCCAGAAAGCACAAAGUGUCCCGGACAAGGAGACUCACUCCACGAGCAUUAUCGCUAUAUCGCGCGGGAUC</p>	

<p>GCCAGCUAUAAGAAAGGAAUUCUGCAGACUGUCAAGGUGUGGACGAAUCUGGUGAAAGUGAUGGGAAGGCAUAAGCCGGAGAAUA UCGUGAUCGAGAUUGGAGGGGAAAACAGACGACCCAGAAAGGACAGAAAACACCGCGGAAACGCAUGAAGCGCAUCGAAGAGGGA AUCAAAAGACUUGGAGCCAAAUCUCUAAAAGAACACCCUUGGAAAUAUCCCAAUCUGCAGAAUGAAGAAUCUUACUGUAUUACCU CCAAAACGGCCGGAUGUACGUGACAGGAAUUGGACAUAAACCGGCUUCCGACUACGUGGACGCUUUCGCGCAGU CCUUCUGAAAAGACGAUUCGACAAUAAAGGUCUGAGUAUAGCAAGAAUUCGGGAAAGUCAGACAAACGUCUAGCGAA GAGGUCUUAAGAAGUAAGAAUUAUCUGGCGCCAGCUGUCUGAAACGCGAAGCUUAUCACUCAGCGCAAGUUCGACAAACCUACCAA GGCAGAAAGAGCGGGAUUGUCGGAGCUCGACAAAAGUCUGGUUUAUCAAAGCCAGCUCUCGAAAACUGCCAGAUUAUAAGCAUG UGGCGCAGAUUCUGGACAGCCGCAUGAAUAUAAGUAUGAUGAAUGCAAGCUCUACCCGGGAGGUAAGGUCACUACCCGCUAAG UCCAAAGCUGGUCGCAUUCGGAAGGACUUCCAAUUCUACAAAGUCAGAGAAUCAAACAUAUCCAAUCCAGCAGCAGUAGCCGCUA CUUGAAUGCAGUGGUGGUAUCGCCUCAUCAAAGAAUACCAAAGCUUAGAAAGCGUUUGUCUACGGAGACUACAAGGUGUACG ACGUCCGGAAGUAUCGCCAAUUCGGAACAGGAAUUGGAAAGCGCAGCUCUUAAGUAUCUUCUAUCGAAUAUAUGAAUUUC UUCAAGACCGAGAUACGCUUGCAAAAUGGCGAAAUCGCGAAGCGGCCCUCAUCGAAAACCAACGGAGAAAACCGGAGAAAUCGUGUG GGACAAGGUCGGAUUUUGCGACCGUCCGAAAGGUUCUAGCAUGCCUAAAGUAAUCUGUCAAGAAAACGGAAAGUGCAGACUG GAGGCUUCAGCAAGGAGUCCAUUCUCCGAAAACGCAACUCCGACAAAACUGAUCGACGCAAGAAAGACUGGGACCCGGAAGAAUAC GGAGGCUUCGAUUCGCCGACUGGCUUACUCGGUUCUGGCUAAGGUGGCAAGGUGGAAAAGGAAAGUCCAAAGAGCUGAAGUCCCGU CAAGGACUUCUGGAAUCAUCAUGGAACGGUAAGUUCGAGAAAACCCAAUUGACUUCUGGAGGCAAAAGGGGUAACAAGG AGGUGAAGAAAGUAUCAGUAUCAAAUCUGCGAAGUAAGCUCGAAAACGGACGAAAGGUAAGGUAAGGUCUCCGCG GGAGGUCGCAAAAGGAAACGAGCUUCCUUAAGUAUGGAAACUUCUGUAUCUGGCAUCCCAUACGAAAACUGAA GGGUUCGCGGAGGACAAACGAAAGCAGCUGUUUGGAAACAAACAAAGCAUUAUCUGGAUGAAUUAUCGAAACAAUACAGCG AAUUCUAAAAGGUGUAUCUUGGCCGACGCCAACUUGGAAAGUUCUUCGCUUCCGCUACAAACAAACUCCGCAAGCCGAUCCGG GAGAGCCGAAAACAUAUACUCUUAUACCCUGAUAUUCUGGUGCGCCGCGGCUUUAUUAUUCGAAUACUUCGAAUACCAAGC CCGGAAGAUACACAGCAACCAAGAGGUGUUGGACCGCACCUCAUCCAAUUAUUAUACCGGCCUUAUGAAACUAGGAUCG ACCUCAGCAGCUGGAGGGAUGAUUCCCAAGAAAGAGAAAGUUCUCCGACUACAGGAUCAAGGAUCAUGAUGGGGACUAUAAA GAUCAUGAUUUAUUAACAAGGACGACGACAAAGGCGGUGGAGGAGGUAUUCGCGCCGCGCGGUAUCGACGACUUCG CCUCGACUUCUGGUAUCCGACGCCUUCGACUUAUGAUCUGGAUAUUGCUGGGUUCGGACGCCUUAUGAUGACUUCGUAUCUGGACA UGCGGGGUCGGAUGCACUGGACGACUUCGACCUUGAUAUGCUGGUAUUAUAGGUCUGGAGCCUUGGUGGCAUUCUUGCCCU UGGGCCUCCCCCAGCCUCCUCCUCCUCCGACCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCUCCU CUGAGUGGGCGC+A (140nt) tail</p>	<p>55</p>
<p>mRNA sequence of Cas9 (Mali)</p>	<p>GCGAAUAAGAGAGAAAAGAAAGAGUAAGAAAGAAAUAUAAAGAGCCACCAUGGACAAAGAAAUAACAGCAUCGCGCCUGGUAUUGGAAACU AACAGCUUGGAGGAGGAGUGAUCACCGACGAGUAACAAGGUGCCGAGCAAGAAAGUUCAAAGGUGCUGGGGAAACUGACCCGCAUUC AAUUAAGAAAACUCAUUGGAGCUCUUAUUGAUCUGGUGAGACUGCCGAAAGUAUACAGGUCUCAAACGCAACCGCACGACGAC GGUACACCCCGCAAGAAUCGCAUCUGCAAGAGAUUUUCCAAACGAGAUUGGCAAGGUAAGCAGCUUUUUUCCAC CGGCUUGAAAGAGAUUCCUUGGAAAGAGGACAAAAGCAAGAAAGCAUCCAAUUCUGGUAACAUCGUGGACGAAAGUGGCGUA UCACGAAAAGUACCCUACCAUCUUGCGGAAAGGUGGUAUUCACGGAUAAAGCAGACCUUGAGAGUAUCUACCUUG CUUUGGCCAUUAUGAUCAAAUCGCGGCCAUUUCUGAUCGAGGGGGAUUAUCCCGGUAUACUCGGAUACUCGGAUUCGAAAGCUGUUC AUCCAGCUGGUCCAAACGUAUAACCAACUGUUUAGGAAAUAUCCCAUCAACGCUUCGGGGGUGGACGCCAAAGCAAUCCUUCUCCGC CGCCUAGCAAGUCACGGCGGCUUGAAAACUUGAUCGCGCGGAGAAAAGAAAUAUGGACUUGUUUGGAAUUCUGAUCG CGCUGCUCGGCCUGACUCCAAAACUUUAAGUCAAAUUCGUAUUGGCCGAAAGUCCAAAGGAGCAGCUGCAAAAGGACACUUC GACGACGACCUUGGACAAUCUGCUGGCCAGAUUGGGGACCAAUAACGACAGACCCUUGUUCUUGGCCGGAAGAACCCUGAGCGACGCCAU</p>

<p> UUUCUGUCCGAAUUAUCUGAGUCAAUACCGAAAUCAUAAAGGCUCCGUGUCCGCUUCAAUGAUAAGCGCUACGAUGAACACC ACCAGAUUCACUCUGCAAAGCCUCUGAGACAAUUAUCCUGAAAAGUAAGAGGAUUAUUCUGGACCGAGACAAAAC GGUACCGCAGCUACUCGAUGGAGGAGGUCACAAGAGAUUCACAAAGUUAUCAAGUUAUCAAGGCCAAUUCUUGGAGAAGAGGACGGUAC UGAAGAACUCCUUGUGAAGCUGAAUAGGAGGAUUGUCUCAGAAAAGCAGCGGACUUAUGAACAGGCUUCCUCACUGAUUC ACCUCGGUGAGCUGCAUGCCUCCUUGGCGCAAGAGAUUUUAUCCUUCGAAAGAAUAUUCGCGAGAAAUCGAAAAGAU CUGACGUUCAGAAAUCUACUGUGGAGCCUGGCGCGGUAUUCGCGGUGUUGCAUGGAUGACUCGCAAGUCAGAGGAAAC UAUCAUCCUUGGAAUUUGAGGAGGUCGUAUAGGGGACUCCUCCAGUUAUCAUGAACGCAUGAACCAUUCUGACAAGA AUUCUCCGAAACGAAAGGUUCCUAAAGCACUCCUUCUUGUAUAGAAUUAUCAUCCUGUAUACAGAGCCAAAGUUUAUUA GUCACCGAGGCAUGAGAAAGCCGCUUCUACAGCGGCGAAUCAAAGAGGCAUUCUGCACUCCUUCUUAAGACCAACCGGAA GGUGACCGUAAGCAACUCAAAGAGACUAUUAAGAGAAUGAAUUCUUAUCCUGGUAUUCGUAUUCGAAUUCAGCGGAGGAGGACCGGU UUAACGGUCAUUGGUAUCCUACAUGAUUCCUGAAAUAUCAUAAAGACAAGGAUUCUCUGGACAAACGAAAGAAAACGAGGACAUC CUGGAAGAUAUUGUUGAUCUUGACCCUGACCGUUCGAGGACCGGAAUUAUGACGAGGAAAGGCUUAAGACCUACCGCACACCUUCUCGA UGACAAAGUGAUGAAUCAAUGAAGCGGCGGAGUAUUAUCUGGCGGGGAGGCUUCUCCGGAAGCUUAUUAUGGAAUCAGAGACA AACAGUCGGUAAAUAUCUUCGACUUCUCAAAGUCCGAAUGGGUUCGCAACCGGAACUUAUGCAGUCGACGUAUCCACGAUUAUCC UUGACUUAAGGAAGAUUCCAGAAAGCGCAAUGAGGAGGAGGAGAUUCUGUACGAAACAUAUCGUAUUCGCGCGGUAUC CCCAGCAUAAGAAAGAAUCUUGCAGACCGUGAAGGUGGUGAUGAAUCUGGUGAAAGUGAUGGCGGCGCCAAAACAGAGAAACA UCGUCAUUGAGUUGCCCGGAGAAUCAGACCAUCAGAAAGGCAAAAAGAACUCCAGAGAGCGGAUAAAACCGCAUCGAGGAAGGC AUCAAAGACUUGUAGCCAAUCCUGAAGAACACCCCGUUCGAGAACACCCAGCUCAGAACGAAAGAAAGCUUAUUCUUAUCCU CCAAAUUGGACGGGACAUUGUACGUCGACAGGAAUUGGACUAACAAGACUCAGCAGCAUCAGUGGAGCCAUUAUUGUCCACAGU CCUUUUAAGGACGACAGCAUCGAUAAUCAAAGUGUCUACVAGAUACAGACAAAUAUCGCGGAAAUACAGACAAUUGUCCUACGGA GAGUUGUCAAAGAAUAAAUAUCUGGAGACAGUCUCAAUGCCAAUUAUACCCAGCGGAAGUUCGACAAUCCUUAACCA GGCCGCGGAGGAUUGUCCGAAUCUGACAAAGCCCGUUAUCAAAGGCGUGGUGAAACCCGGCAGUACUACUAAAACAG UGGCCAGAUCCUCGAAUUCGCGAUAAUACGAUGAAUGAACAGCUGAUAUAGGAAUUAAGGUAUAGGUAUAGGUAUACUCUGAAG UCGAAACUGGUGCGGACUUUAGAAAGAUUUCAGUUUAUCAAAGUCCGCGAGAUAAACAUAUCCACCGCUACUACGACGCUA CCUGAAUGCAUUGGGCACCGCGCUGAUCAAAGAAUUCGGAAGUCUGAAUCCGAGUUCGUGUACGGAGAUUAUCAAAGUGUA ACGUGCGAAAGAUUCGCCAAGUCGGAACAGGAAUUCGGAAGGCUACUGCAAAGUAUUCUUAUCUCAAAUCAUAGAAUUC UUCAAAACGGAGAUACCGCUCGGAAACGGCGAAUUCGGAAGAGCCUUAUUAUUAUUGAAACCAACGGAGAAACCGGGAGAU GGACAAAGGAAAGGAAUUGGCGACUGUGAGGAAAGGUGUUGUUAUGCCGCAAGUAAUUGUAAAAGAGCGAAAGUGCAAACCG GAGGAUUCAGCAAGAAUCCAUCCUCCAAAGCGCAUCUGGACAAAACUCAUCGCGCGAAAGAAAGGUAUUGGACCCAAAGAAUUA GGUGGUUUGACAGCCUUAUGUGGUUACUCCUGUCCUGUUGGCUAAAUGGUAAGGUAUUCGAAAGCUCAAUUCGAAUCG GAAAGGCUCCUGGAAUACAGAUACUGGAGCGGUCGAGUUCGAAAGAAUCCUAUUGAUUUCUGGAGGCGAAAGGCUA AAGUCAAGAAAGAACUUGAUCAUACAGUCCCGAAGUACAGCUUCGAGCUCGAAACGGCAGAAAGAGGUAUGGCUACAGCG GGAAUUGCAGAAAGGAAACGAAUCUGGCAUGCCGUAAGUAUUUUUUAUCUGGUAAGCCUACUACGAAAGGUAU GGAUCCGCGGAGGACACGCAAAAACAACUUGUUGGAGCAGCACAAGCACUACUGGUAUGAGAUUACGAGCAGAUUCUCCG AAUUCUGAAAACGCGUGAUCCUUGCGGAUGCCAAUCUGGUAUAAAGUUGUGGCUUAUCAAAGCAUCGGGAUAAACCGAUUC GAAACAGGAGAAACAUAUUAUCUUGUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUA UAGAAAGGCUACACCGCAAGGAAAGGUCUGGACGCCUCCUGAUCCACCGAUUUAUUAUUAUUAUUAUUAUUAUUAUUA ACCUUGCCAGCUUGGAGGAGAUUACCGGGCGGUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUAUUA CUUCUUGCCUUGGGCCUCCCGACGCCCCUCCUGCACCCCGUACCCCGGUGGUCUUUGAAUAAAGUUCUGAGUGGGC </p>

mRNA sequence of Cas9 (Cong)	<p>GGGAAUUAAGAGAGAAAAGAGUAAGAAAGAAAUAUAAGAGCCACCAUGGACUACAAGGACCAGCAGCGGAGACUACAAGAACCAU GACUUCGAUUAACAAGGAGUACGACGACAAUUAUGGCAACCGAAGAAAGAGAAAAGGUCGGAAUUAACGGGGUCCGGCCGGGACAA GAAGUACUAAUCGGACUGGAUUCGGCACGAACAGCGUGGUGGCGAGUGUACACCGACGAUAACAAGGUGCCGAGCAAGAAAGU UCAAGUGCUGGGAAAUAACCGAUCCGAUCCGAUUAAGAAAUAUUGAUUGGCGCUCUGUUCGACUCUGGGAGAGACUGCCGGAG GCCACUAGACUGAAGAGGACCGUAGGCGCCGCUACAGAGCGCAAAAACCGCAUCUGCUAUUUAUUAAGAAUUCUUCAAAACGA GAUGGCCAAGGUGGACGACUCCUUUUUCCAUCCGGUCGGAAGAAUUAUUUCUGUGGAGGAGGACAAGAAAGCAGAAACGCCAUUCCCA UUUUCGGCAACAUUGUCGACGGAUGGGCCUAUCAUGAGAAUCCCGACUAUCAACUACGAAAGAAAGGUGGAGACUCCACU GACAAAGGACAUUCUGGGUUAUCAUCUGCACUGGCCAUUAUGAUAUAAUUCGGGGACACUUAUUCUUCGAGGCGGACCUUAA UCCCGACAAUCCGAUGUGGAUAGCUUUUUAUCCAGCUGGUCAGACCUACAACCAACUGUUUGAAGAAAUCUAAUCAAUUGCGA GCGGUGCAUGCAAGGCCAUCCUGAGCGCCCGCCUCUGAAAAGCAGAAAGGUCGAAAACCCUGAUCGCACAGUUGCCUGGAGAG AAGAAGAACGGCCUCUUCGGCAUUCUACUGCAUUGUCCUUGGACUGACUCCAAACUUCAAUCCAAUUCGACUUGGCCGAGGA CGCCAAACUGCAAUCUGACCAAGAUACCUACGAUGAUGAUUGGACAAUUCUGGUCUAGAUCCGGCACCAUUAUCGCCGACCUUG UCCUUGCGCUAAAGAACUUGUCGGACGCCAUCCUGUCUCCGACAUCCUGCGGUCUAAUACCGAAUUAUCAAAGCACCAUCUCUCG GCAUCCAUAGAUCAAGAGAUACGAAACACCAACAGGACUCCUCCUGAAAAGCAGUGGCGGCAGCAGCUCUUGAGAAAUA CAAAGAAUUCUUGAUUCAGUCCAAAGAACGGAUACGCCGGAUACAUCGACGGCGCGGAGCCAAAGAAUUCUACAAGUUA UCAAGCCGAUCCUGGAAAAGUUGGAUUGGACGGAAGAAUUCUCUGUUAACUUGAAUGAGAGGAUUCUGUCUCCGCAAAACAACGGACC UUCGUAACGGAUCCUACCAAAGUCAAUAUCGUGACUGAGGAAUUGCCAAAGCUCUCCUGGCGGACGAGGAGGACUUUUAACCGGUUCCU CAAAGACAAACAGAGAAAAGAUCCGAAAGAUUCUUGACUCCUGUAUCCUGGCGCUCUCCGCGGAGGAAUCCUCCCGU UUGCUUGGAGUACUAGAAAAGUCAGAGAAAACGAUACCCCAUUGAAACUUCGAAAGGUGGUGACAAAAGGAGCGCCCAUUCG UUAUCGAAACGGAUACUUAUCGUAUAAGAAUUCUGCCGAUGAAGAGUCCUGCUUAAGCAUCUCAAAGAGGACUACUUUAAAGAAUCCGAAUUGAC UGUGUAUAACGAAUCUACCAAAGUCAAUAUCGUGACUGAGGAAUUGCCAAAGCUCUCAAAGAGGACUACUUUAAAGAAUCCGAAUUGAC UCGUGGACUUGCUGUUAACCAACCGCAAGGUGACUUAAGCAUCUCAAAGAGGACUACUUUAAAGAAUCCGAAUUGAC UCGGUCGAGAUUCCGGGUGGAAUAGAUUAUCAACCGCUCGCGUGGGAAACCUACCAUGAUCUCCUCAAAGAAUUAUCAAAGGACAAAAG CUUCCUGGAUAAACGAGGAGAAUAGAGCAUCCUCGAAAGAUUUGUGUUAUCCUCCUGACCCUUUUUCGAAAGAUCCGAAAUAUUAUCGAAAG AACGCCUAGAAACCUACGCUCACUUGUUCGACGUAAGGUGAUGAAACAGUUGAAACGCGCGGCUAACAAGGUGGCGGCGGUG UCGGCAAGCUGAUCAACGGAAUUCGGGACAAACAGAGCGGAAAGACCAUCCUCGUAUUUCUGAAUCCCGAUGGUUUUUGCCAAACCG CAACUUCUAGCAGCUACUCCAUAGCAGUUCGUUACCUUUAAGGAGAUUCCAGAGGACCAAGUGUCGGGACAAAGGGGAUUCGCG UCCACGAAACAUCCGCAAUCUGGCGGGUCCCGGCAAUUAAGAAAGGAAUCCUCCAGACUGUUAAAGGUGGUCGACGAGCUGGUG AAGGUGAUGGGGAGACAUAAGCCUGAAAACAUUUGAUCGAGAUUGGCGAGAGAAAUAUCAAAUCAAAUCAAGAGGACAGAAAGAAUUC CCGGAGCGGAGUAGCGCAUCGAGAGGAAUCAAAGAAUCUGGCUCCCAAUCCUGAAAGAGCAUCCCGUGGAAAUAUCUCAGC UGCAGAACGAGAAACUUUACCUUAUCUUAUCAAUUGGAGGACAUUGACGUCGCAAGAAUCUGGAUUAUCAUCCGCUUCU GAUUAACGACGUCGAUACAUUCUCCCAAUCAUUCUUGAAGGAUUAUCAUCGUAUAAACAAGGUGUCACUAUAUCAGACAAAA CCGGGAAAGUCAGAUAAACGUCUCCAGCGAAGAAAGUCUGAAGAAAGAAUUAUCUGGAGGCAACUUCUGAACGCCAAACUCA UCACUAGCGCAAGUUCGACAAUCGACAAAAGCAGAAAGGAGGACUACGCGAGCUGGCAAGGCGUGGUUUCAUCAAACGGCAG CUGGAGGACUCGCCAAUACGAAAGCAUGUGGCCAGAUUCUGCAUCUGCGCAUGAAUAUAUAAAGCAGCAAGAAACGUAUAGCU GAUCCGGAGGUGAGUACCCUCAAAGCAACCUUGUCCGAAUUCUCCGAAAGACUUCAGGUUUAUCAUAAUCCGCGGAG UUAACAAUACCAUACGCUACGACGCUUACCUCAAUUGCUGUGGUGGGGACCGGCUUGAUAUAAAGAAUCCCAAAAACUUGGAGUCC GAAUUCGUCUAACGGAGAUUAACAAGGUCUACGACGUGGCGCAAGAUUAUUGCCAAAGUCGAGCAGGAAAUUGGGAAAAGCGACUGCUAA</p>
------------------------------	---

<p>GUA CUUCUUCUCGAAUAUCAUGAACUUCUUAAGACCGAAAUAACCCUGGCUAAACGGCGAGAUCAGGAAAACGGCCGCGUAUCG AAACUAAUGGUGAGACUGGUGAAAUCGUGUGGGAUAAGGACGGGAUUUCGCCACGGUCGCAAGGUCCUCAGCAUGCCGCAAGUG AAUUAUUAAGAAAACCGAAGUGCAGACCGGUGGUUCGAAAGAAUCCAUCCUGGCAAGCGCAACUCGGAAUAGCUUAUUGC CCGCAAGAAAGAUUGGACCCGAAAAGUA CGGUGGUUCGACUCCCUACCCUGGCGUACUCGGUGUUGGUGGCGCCAAAAGUGG AAAAGGCAAAUCAAGAAAGCUCAAGAGCGUCAAGGAGCUGUGGAAUACCAUGGAGAGGUCCAGCUUUGAGAAAAACCCCG AUCGACUUCUUGGAA GCCAAGGGAUACAAAAGAGGUAAGAAAGACCUGAUCAUCAAACUUCCAAAGUAUCUCCUGUUCGAAUCUCGA AAACGGAGGAAAGCGCAUGCUCGCCUCAGCCGGGGAACUGGCAAAAGGGCAACGAAUCUGGCCUCCCGUCAAAAUAACGUCAA CUUCC UGUACUUGGCGUCACACUACGAAAAGCUGAAAAGAUCCCCAGAGGACAAACGAAACAGAAAACAGCUUUCGUCGAGCAGCAAGCAC UACCCUGGACGAGAUCAUCGAAACAGAUUCUCGGAAUUCAGCAAGAGAGUGAUUCUUGGCAGACGCUAAACCUUGACAAAAGUCCUCUGGC AUACAAUAGCAUCGCGACAAGCCGAUCAGAGAACAGGCGGAGAAACAUCAUCCACUGUUCACUCUCACCAACCCUGGCGGCGCCAG CGCUUUUAAGUA CUUUGAUACCAUUGACCGCAAGAGAUACCUCAAUAAGAAAGUGCUGGACGCAACCCUGAUCCAUCAA AGCAUACCCGGACUUUAUGAAAUCUGGAUCGAUCUCUCACAGCUCGGAGGAGACAAAAGACCCGGCUGCCACCAAGAAAGGCCGGACA GGCAAAGAAAGAAAUGAUAUAAGGCUUGGAGCCUCGGUGGCAUGCUUUCUUGCCCCUUGGGCCUCCCCCCAGCCUCCUCCU UCCUGCACCCCGUACCCCGUGGUCUUUGAAUAAAAGUCUGAGUGGGGGG</p>	
--	--

Table 11: Synthetic sgRNA

Description	Sequence	SEQ ID NO:
VEGF V1 sgRNA	GGGUGGCAGACGGCAGUCACUAGGGUUUAGAGCUAGAAAUAAGCAAGUUAAAAUAAGGCUAGUCCGUUUAUCAACUUUGAAAAAGUG GCACCGAGUCGGGUGC	91
VEGF V2 sgRNA	GGGUGAGAGAGUGUGCGGUGGGUUUAGAGCUAGAAAUAAGCAAGUUAAAAUAAGGCUAGUCCGUUUAUCAACUUUGAAAAAGUGG CACCGAGUCGGUGC	92
VEGF V3 sgRNA	GGGUGAGAGAGUGUGCGGUGGGUUUAGAGCUAGAAAUAAGCAAGUUAAAAUAAGGCUAGUCCGUUUAUCAACUUUGAAAAAGUGG CACCGAGUCGGUGC	93
VEGF V4 sgRNA	GGGUUGGAGCGGGGAGAGGCCAGGGUUUAGAGCUAGAAAUAAGCAAGUUAAAAUAAGGCUAGUCCGUUUAUCAACUUUGAAAAAGUG GCACCGAGUCGGUGC	94
basic sgRNA template is based on Gilbert, Qi, Maeder,	GGNN GUGGCACCGAGUCGGGUGGUGC	90

Table 12: Additional Sequences as deposited in Addgene database (plasmid repository, Cambridge, MA)

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
dCas9 (bacteria)	<i>S. pyogenes</i>	Plasmid 44249: pdCas9-bacteria	D10A H840A (catalytically inactive)	aTc-inducible expression of a catalytically inactive bacterial Cas9 (<i>S. pyogenes</i>) for bacterial gene knockdown
Cas9	N/A	Plasmid 41815: hCas9	human codon-optimized	Expresses human codon optimized Cas9 nuclease for genome engineering
Human codon optimized Cas9	N/A	Plasmid 44758: pST1374-NLS-flag-linker-Cas9	codon-optimized	Expresses human codon optimized Cas9 nuclease for genome engineering
Cas9_D10A	N/A	Plasmid 41816: hCas9_D10A	human codon-optimized, D10A nickase	Expresses human codon optimized Cas9 D10A mutant which functions as a nickase for genome engineering

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
dead Cas9 with 3X NLS	<i>H. sapiens</i> (human)	Plasmid 44246: pdCas9-humanized	D10A H840A (catalytically inactive)	Expression of a catalytically inactive, human codon-optimized Cas9 under the control of Murine Stem Cell retrovirus LTR promoter for mammalian gene knockdown
guide RNA (bacteria)	N/A	Plasmid 44251: pgRNA-bacteria	N/A	Expression of customizable guide RNA (gRNA) for bacterial gene knockdown
dCas9 fused to BFP	<i>H. sapiens</i> (human)	Plasmid 44247: pdCas9::BFP-humanized	D10A H840A	Expression of a catalytically inactive, human codon-optimized Cas9-BFP fusion under the control of Murine Stem Cell retrovirus LTR promoter for mammalian gene knockdown

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
wild-type Cas9	<i>S. pyogenes</i>	Plasmid 44250: pwtCas9-bacteria	N/A	aTc-inducible expression of wild-type Cas9 (<i>S. pyogenes</i>) for bacterial gene knockdown
guide RNA	<i>H. sapiens</i> (human)	Plasmid 44248: pgRNA-humanized	N/A	Expression of customizable guide RNA (gRNA) under control of murine U6 promoter, also contains a CMV-puro-t2A-mCherry expression cassette for mammalian gene knockdown
Codon optimized Cas9	N/A	Plasmid 45945: pHsp70-Cas9	Silent change from C to A at bp 834 of insert	The codon-optimized Cas9 nuclease under the control of the <i>Drosophila</i> hsp70 promoter used in Gratz, et al. (2013). Plasmid is low copy number.
Csy4	<i>Pseudomonas aeruginosa</i>	Plasmid 41091: pHMGWA-Pa14Csy4	Wild type	

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
Csy4	<i>Pseudomonas aeruginosa</i>	Plasmid 41092: pHMGWA- Pa14Csy4H29A	His29Ala mmutation to abolish ribonuclease activity	
mammalian codon- optimized <i>Streptococcus</i> <i>pyogenes</i> Cas9 - 3X Flag	<i>Streptococcus</i> <i>pyogenes</i>	Plasmid 43861: JDS246		Expresses mammalian codon optimized Cas9 nuclease with C- term 3X FLAG from CMV and T7 promoters
<i>Streptococcus</i> <i>pyogenes</i> Cas9	<i>Streptococcus</i> <i>pyogenes</i>	Plasmid 42251: MLM3613		Expresses Cas9 nuclease (<i>Streptococcus</i> <i>pyogenes</i>) from CMV and T7 promoters
None		Plasmid 43860: MLM3636		guide RNA (gRNA) expression vector used to create a gRNA to a specific sequence, uses U6 promoter
<i>Streptococcus</i> <i>pyogenes</i> Cas9-3X Flag		Plasmid 42252: MLM3639		
Cas9	<i>Streptococcus</i> <i>thermophilus</i>	Plasmid 39314: pMJ824		

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
Cas9	<i>Listeria innocua</i>	Plasmid 39313: pMJ823		
Cas9	<i>Streptococcus pyogenes</i>	Plasmid 39312: pMJ806		
Cas9	<i>Streptococcus pyogenes</i>	Plasmid 39315: pMJ825		
Cas9	<i>Streptococcus pyogenes</i>	Plasmid 39316: pMJ826		
Cas9	<i>Neisseria meningitidis</i>	Plasmid 39317: pMJ839		
Cas9	<i>Streptococcus pyogenes</i>	Plasmid 39318: pMJ841	Aspartate 10 to Alanine (D10A) and Histidine 840 to Alanine (H840A)	
Cas9	<i>S. pyogenes</i>	Plasmid 43945: p3s-Cas9HC		
Human Optimized <i>S. pyogenes</i> Cas9	<i>H. sapiens</i> (human), <i>S. cerevisiae</i> (budding yeast); <i>S.pyogenes</i>	Plasmid 43802: p414-TEF1p-Cas9-CYC1t		
Human Optimized <i>S. pyogenes</i> Cas9	<i>H. sapiens</i> (human), <i>S. cerevisiae</i> (budding yeast); <i>S.pyogenes</i>	Plasmid 43804: p415-Gall-Cas9-CYC1t		
CAN1.y gRNA	<i>S. cerevisiae</i> (budding yeast)	Plasmid 43803: p426-SNR52p-gRNA.CAN1.Y-SUP4t		

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
codon optimized Cas9		Plasmid 46294: pBS-Hsp70-Cas9	Silent change from C to A at bp 834 of insert	A codon-optimized Cas9 nuclease under the control of the Drosophila hsp70 promoter.
tracr/Cas9		Plasmid 42876: pCas9		Bacterial expression of Cas9 nuclease, tracrRNA and crRNA guide
Cas9-2A-GFP	Synthetic	Plasmid 44719: pCas9_GFP		Co-expression of human codon-optimized Cas9 nuclease and GFP, plasmid optimized for expression in human pluripotent stem cells

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
Cas9D10A-2A-GFP	Synthetic	Plasmid 44720: pCas9D10A_GFP		Co-expression of human codon-optimized Cas9 (D10A) mutant nickase and GFP, plasmid optimized for expression in human pluripotent stem cells
dCas9	<i>S. Pyogenes</i>	Plasmid 47106: pcDNA-dCas9	D10A, H840A	Expresses inactivated <i>S. pyogenes</i> dCas9 (D10A, H840A) in mammalian cells
dCas9-VP64	<i>S. Pyogenes</i>	Plasmid 47107: pcDNA-dCas9-VP64	D10A, H840A	Expresses inactivated <i>S. pyogenes</i> dCas9 (D10A, H840A) fused to VP64 transactivator domain in mammalian cells
CRISPR-BsaI		Plasmid 42875: pCRISPR		A crRNA expression plasmid for targeting a specific sequence.

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
CRISPR::rpsL		Plasmid 44505: pCRISPR::rpsL		A crRNA expression plasmid specific to the rpsL allele.
nls-zcas9-nls	Synthetic	Plasmid 47929: pCS2-nCas9n		expression of an optimized Cas9 for genome-editing in zebrafish
codon optimized Cas9_SV40 NLS with intron	Synthetic; <i>S. pyogenes</i>	Plasmid 46168: Peft-3::cas9-SV40_NLS::tbb-2 3'UTR	intron inserted at position 3113-3163, and SV40 NLS inserted from position 5192-5227	
dCas9-BFP fusion	<i>H. sapiens</i> (human)	Plasmid 46910: pHR-SFFV-dCas9-BFP		Human expression vector containing SFFV promoter, dCas9 that is fused to 2x NLS and tagBFP

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
dCas9-BFP-KRAB fusion	<i>H. sapiens</i> (human)	Plasmid 46911: pHR-SFFV-dCas9-BFP-KRAB		Human expression vector containing SFFV promoter, dCas9 that is fused to 2x NLS, tagBFP and a KRAB domain
Cas9	Synthetic; <i>Streptococcus pyogenes</i>	Plasmid 42234: pMJ920	codon-optimized synthetic DNA sequence	
dCas9-p65AD-BFP fusion	<i>H. sapiens</i> (human)	Plasmid 46913: pMSCV-LTR-dCas9-p65AD-BFP		Human expression vector containing MSCV LTR promoter, dCas9 that is fused to 2x NLS, p65 activation domain and tagBFP
dCas9-VP64-BFP fusion	<i>H. sapiens</i> (human)	Plasmid 46912: pMSCV-LTR-dCas9-VP64-BFP		Human expression vector containing MSCV LTR promoter, dCas9 that is fused to 2x NLS, VP64 and tagBFP

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
SPgRNA	Synthetic	Plasmid 47108: pSPgRNA		Expresses a <i>S. pyogenes</i> Cas9/dCas9 guide RNA in mammalian cells
Cas9	Synthetic	Plasmid 46757: pT3TS-nCas9n		
tyr Target		Plasmid 46761: pT7tyrgRNA		
dCas9		Plasmid 46920: pTDH3-dCas9		Yeast CEN/ARS vector (Leu2) that contains dCas9 fused to NLS controlled by TDH3 promoter
dCas9-Mxi1		Plasmid 46921: pTDH3-dCas9-Mxi1		Yeast CEN/ARS vector (Leu2) that contains dCas9 fused to NLS and Mxi1 domain controlled by TDH3 promoter
sgCD71-2		Plasmid 46918: pU6-sgCD71-2		Human pSico-based U6 vector containing murine U6 promoter and sgRNA targeting endogenous CD71 gene

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
sgCXCR4-2		Plasmid 46917: pU6-sgCXCR4-2		Human pSico-based U6 vector containing murine U6 promoter and sgRNA targeting endogenous CXCR4 gene
U6-SpTracrRNA-EF1a-hSpRNaseIII-mCherry	<i>S. pyogenes</i>	Plasmid 41863: pX097-U6-SpTracrRNA-EF1a-hSpRNaseIII-mCherry		This plasmid is used to reconstitute the complete Type II CRISPR system from <i>S. pyogenes</i> and contains the host factor RNaseIII and tracrRNA.
Humanized <i>S. pyogenes</i> Cas9	<i>S. pyogenes</i>	Plasmid 42229: pX260-U6-DR-BB-DR-Cbh-NLS-hSpCas9-NLS-H1-shorttracr-PGK-puro		This plasmid separately encodes a human codon-optimized SpCas9, a tracrRNA and customizable crRNA.

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
humanized <i>S. pyogenes</i> Cas9	<i>S. pyogenes</i>	Plasmid 42337: pX261-U6-DR-hEmx1-DR-Cbh-NLS-hSpCas9-NLS-H1-shorttracr-PGK-puro	Humanized	Dual expression plasmid of human codon-optimized SpCas9 and a gRNA to the human Emx1 locus, can be used to test SpCas9 cleavage in cell lines of choice.
humanized <i>S. pyogenes</i> Cas9	<i>S. pyogenes</i>	Plasmid 42230: pX330-U6-Chimeric_BB-CBh-hSpCas9	Humanized	A human codon-optimized SpCas9 and chimeric guide RNA expression plasmid.
humanized <i>S. pyogenes</i> Cas9 (D10A) nickase	<i>S. pyogenes</i>	Plasmid 42333: pX334-U6-DR-BB-DR-Cbh-NLS-hSpCas9n (D10A) - NLS-H1 - shorttracr-PGK-puro	D10A nickase-converting mutation	This plasmid separately encodes a human codon-optimized SpCas9 nickase, a tracrRNA and a customizable crRNA.
humanized <i>S. pyogenes</i> Cas9 (D10A) nickase	<i>S. pyogenes</i>	Plasmid 42335: pX335-U6-Chimeric_BB-CBh-hSpCas9n (D10A)	D10A nickase-converting mutation	A human codon-optimized SpCas9 nickase and chimeric guide RNA expression plasmid.

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
None		Plasmid 41824: gRNA_Cloning Vector		An empty gRNA expression vector, used to create a gRNA to a specific sequence.
gRNA_AAVS1-T1		Plasmid 41817: gRNA_AAVS1-T1		Expresses a guide RNA (gRNA) to target human AAVS1 (T1 target sequence) for genome engineering
gRNA_AAVS1-T2		Plasmid 41818: gRNA_AAVS1-T2		Expresses a guide RNA (gRNA) to target human AAVS1 (T2 target sequence) for genome engineering
gRNA_DNMT3a-T1		Plasmid 41821: gRNA_DNMT3a-T1		Expresses a guide RNA (gRNA) to target DNMT3a (T1 target sequence) for genome engineering

Gene / Insert	Species	Plasmid Name	Mutation	Purpose
gRNA_DNMT3a-T2		Plasmid 41822: gRNA_DNMT3a-T2		Expresses a guide RNA (gRNA) to target DNMT3a (T2 target sequence) for genome engineering
gRNA_DNMT3b		Plasmid 41823: gRNA_DNMT3b		Expresses a guide RNA (gRNA) to target DNMT3b for genome engineering

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word “comprise” or variations such as “comprises” or “comprising” is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

Claims

1. A composition comprising:
 - (a) an mRNA comprising:
 - (i) at least one 5'-cap structure;
 - (ii) a 5'-UTR;
 - (iii) an open reading frame encoding a polypeptide,
wherein the polypeptide has an amino acid sequence that is at least 80% identical to SEQ ID NO: 7, 8, 9, 61, 62, 63, 64, or 65, and
wherein the open reading frame consists of nucleotides selected from 1-methylpseudouridine, cytidine, adenosine, and guanosine; and
 - (iv) a 3'-UTR;
 - (b) an sgRNA comprising:
 - (i) a first region of 20-25 linked nucleosides complementary to either strand of a 5'-UTR of the gene of interest;
 - (ii) a second region located at the 3'-terminus of the first region comprising a guide RNA scaffold sequence;
 - (iii) at least one phosphorothioate linkage; and
 - (iv) at least one 2'-methoxy-nucleotide;and
 - (c) a plurality of lipid nanoparticles comprising:
 - (i) a biodegradable cationic lipid;
 - (ii) a non-cationic lipid;
 - (iii) a sterol; and
 - (iv) a PEGylated lipid,wherein the lipid nanoparticles encapsulate the mRNA and/or the sgRNA.
2. The composition of claim 1, wherein the mRNA further comprises a poly-A tail.
3. The composition of claim 2, wherein the poly-A tail is from 80 to 140 nucleotides in length.
4. The composition of claim 3, wherein the poly-A tail is 100 nucleotides in length.

5. The composition of any one of claims 1 to 4, wherein the biodegradable cationic lipid comprises at least one ester linkage.

6. The composition of claim 5, wherein the biodegradable cationic lipid comprises an internal ester.

7. The composition of claim 5, wherein the biodegradable cationic lipid comprises a terminal ester.

8. The composition of any one of claims 1 to 7, wherein the polypeptide comprises a sequence having at least 80% identity to SEQ ID NO: 61.

9. The composition of claim 8, wherein the polypeptide comprises a sequence that is 100% identical to SEQ ID NO: 61.

10. The composition of any one of claims 1 to 9, wherein the open reading frame further encodes an effector domain.

11. The composition of claim 10, wherein the effector domain is selected from the group consisting of KRAB, VP64, p65AD, and Mxi.

12. The composition of claim 11, wherein the effector domain is KRAB or VP64.

13. The composition of any one of claims 1 to 12, wherein the 3'-UTR comprises at least one miR binding site or one miR binding site seed.

14. The composition of claim 13, wherein the 3'-UTR comprises a miR-122 or miR-142 or miR-146 binding site.

15. The composition of any one of claims 1 to 14, wherein the guide RNA scaffold sequence comprises SEQ ID NO: 82.


16. The composition of any one of claims 1 to 15, wherein the mRNA is produced using an in vitro transcription method or using a chemical synthesis method.

17. The composition of any one of claims 1 to 16, wherein the mRNA is substantially purified.

18. The composition of any one of claims 1 to 17, further comprising a pharmaceutically acceptable excipient.

19. The composition of claim 18, wherein the pharmaceutically acceptable excipient is selected from the group consisting of a solvent, aqueous solvent, non-aqueous solvent, dispersion media, diluent, dispersion, suspension aid, surface active agent, isotonic agent, thickening or emulsifying agent, preservative, polymer, peptide, protein, cell, hyaluronidase, and mixtures thereof.

Figure 1






Figure 2






Figure 3

A)



B)



C)




Figure 4




Figure 5