

(12) STANDARD PATENT APPLICATION (11) Application No. AU 2017202349 A1
(19) AUSTRALIAN PATENT OFFICE

(54) Title
CNS DELIVERY OF THERAPEUTIC AGENTS

(51) International Patent Classification(s)
A61K 38/46 (2006.01)

(21) Application No: **2017202349** (22) Date of Filing: **2017.04.10**

(43) Publication Date: **2017.04.27**

(43) Publication Journal Date: **2017.04.27**

(62) Divisional of:
2011270668

(71) Applicant(s)
Shire Human Genetic Therapies, Inc.

(72) Inventor(s)
Calias, Pericles;Pan, Jing;Powell, Jan;Charnas, Lawrence;McCauley, Thomas;Wright, Teresa Leah;Pfeifer, Richard;Shahrokh, Zahra

(74) Agent / Attorney
Pizzseys Patent and Trade Mark Attorneys Pty Ltd, PO Box 291, WODEN, ACT, 2606, AU

ABSTRACT

The present invention provides an effective and less invasive approach for direct delivery of therapeutic agents to the central nervous system (CNS). In some embodiments, the present invention provides methods including a step of administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme.

2017202349 10 Apr 2017

CNS DELIVERY OF THERAPEUTIC AGENTS

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to United States Provisional Patent Applications serial numbers 61/358,857 filed June 25, 2010; 61/360,786, filed July 1, 2010; 61/387,862, filed September 29, 2010; 61/435,710, filed January 24, 2011; 61/442,115, filed February 11, 2011; 61/476,210, filed April 15, 2011; and 61/495,268 filed on June 9, 2011; the entirety of each of which is hereby incorporated by reference. This application relates to US applications entitled “Methods and Compositions for CNS Delivery of Heparan N-Sulfatase,” filed on even date; “Methods and Compositions for CNS Delivery of Iduronate-2-Sulfatase,” filed on even date; “Methods and Compositions for CNS Delivery of β -Galactocerebrosidase,” filed on even date; “Methods and Compositions for CNS Delivery of Arylsulfatase A,” filed on even date; “Treatment of Sanfilippo Syndrome Type B,” filed on even date; the entirety of each of which is hereby incorporated by reference.

BACKGROUND

[0002] Enzyme replacement therapy (ERT) involves the systemic administration of natural or recombinantly-derived proteins and/or enzymes to a subject. Approved therapies are typically administered to subjects intravenously and are generally effective in treating the somatic symptoms of the underlying enzyme deficiency. As a result of the limited distribution of the intravenously administered protein and/or enzyme into the cells and tissues of the central nervous system (CNS), the treatment of diseases having a CNS etiology has been especially challenging because the intravenously administered proteins and/or enzymes do not adequately cross the blood-brain barrier (BBB).

[0003] The blood-brain barrier (BBB) is a structural system comprised of endothelial cells that functions to protect the central nervous system (CNS) from deleterious substances in the blood stream, such as bacteria, macromolecules (e.g., proteins) and other hydrophilic molecules, by limiting the diffusion of such substances across the BBB and into the underlying cerebrospinal fluid (CSF) and CNS.

2017202349 10 Apr 2017

[0004] There are several ways of circumventing the BBB to enhance brain delivery of a therapeutic agent including direct intra-cranial injection, transient permeabilization of the BBB, and modification of the active agent to alter tissue distribution. Direct injection of a therapeutic agent into brain tissue bypasses the vasculature completely, but suffers primarily from the risk of complications (infection, tissue damage, immune responsive) incurred by intra-cranial injections and poor diffusion of the active agent from the site of administration. To date, direct administration of proteins into the brain substance has not achieved significant therapeutic effect due to diffusion barriers and the limited volume of therapeutic that can be administered. Convection-assisted diffusion has been studied via catheters placed in the brain parenchyma using slow, long-term infusions (Bobo, et al., Proc. Natl. Acad. Sci. U.S.A 91, 2076-2080 (1994); Nguyen, et al. J. Neurosurg. 98, 584-590 (2003)), but no approved therapies currently use this approach for long-term therapy. In addition, the placement of intracerebral catheters is very invasive and less desirable as a clinical alternative.

[0005] Intrathecal (IT) injection, or the administration of proteins to the cerebrospinal fluid (CSF), has also been attempted but has not yet yielded therapeutic success. A major challenge in this treatment has been the tendency of the active agent to bind the ependymal lining of the ventricle very tightly which prevented subsequent diffusion. Currently, there are no approved products for the treatment of brain genetic disease by administration directly to the CSF.

[0006] In fact, many have believed that the barrier to diffusion at the brain's surface, as well as the lack of effective and convenient delivery methods, were too great an obstacle to achieve adequate therapeutic effect in the brain for any disease.

[0007] Many lysosomal storage disorders affect the nervous system and thus demonstrate unique challenges in treating these diseases with traditional therapies. There is often a large build-up of glycosaminoglycans (GAGs) in neurons and meninges of affected individuals, leading to various forms of CNS symptoms. To date, no CNS symptoms resulting from a lysosomal disorder has successfully been treated by any means available.

2017202349 10 Apr 2017

[0008] Thus, there remains a great need to effectively deliver therapeutic agents to the brain. More particularly, there is a great need for more effective delivery of active agents to the central nervous system for the treatment of lysosomal storage disorders.

SUMMARY

[0009] The present invention provides an effective and less invasive approach for direct delivery of therapeutic agents to the central nervous system (CNS). The present invention is, in part, based on the unexpected discovery that a replacement enzyme for a lysosomal storage disease can be directly introduced into the cerebrospinal fluid (CSF) of a subject in need of treatment at a high concentration (e.g., greater than about 3 mg/ml, 4 mg/ml, 5 mg/ml, 10 mg/ml or more) such that the enzyme effectively and extensively diffuses across various surfaces and penetrates various regions across the brain, including deep brain regions. More surprisingly, the present inventors have demonstrated that such high protein concentration delivery can be achieved using simple saline or buffer-based formulations and without inducing substantial adverse effects, such as severe immune response, in the subject. Therefore, the present invention provides a highly efficient, clinically desirable and patient-friendly approach for direct CNS delivery for the treatment of various diseases and disorders that have CNS components, in particular, lysosomal storage diseases. The present invention represents a significant advancement in the field of CNS targeting and enzyme replacement therapy.

[0010] Among other things, the present invention provides methods of intrathecal (IT) administration of a therapeutic agent (e.g., a replacement enzyme) to a subject in need of treatment. In some embodiments, a replacement enzyme can be a recombinant, gene-activated or natural enzyme. As used herein, the terms “intrathecal administration,” “intrathecal injection,” “intrathecal delivery,” or grammatic equivalents, refer to an injection into the spinal canal (intrathecal space surrounding the spinal cord). In some embodiments, “intrathecal administration” or “intrathecal delivery” according to the present invention refers to IT administration or delivery via the lumbar area or region, i.e., lumbar IT administration or delivery. As used herein, the term “lumbar region” or “lumbar area” refers to the area between the third and fourth lumbar (lower back) vertebrae and, more inclusively, the L2-S1 region of the spine. It is contemplated that lumbar IT administration or delivery distinguishes over cisterna

2017202349 10 Apr 2017

magna delivery (i.e., injection via the space around and below the cerebellum via the opening between the skull and the top of the spine) in that lumbar IT administration or delivery according to our invention provides better and more effective delivery to the distal spinal canal, while cisterna magna delivery, among other things, typically does not deliver well to the distal spinal canal.

[0011] In one aspect, the present invention provides methods including a step of administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme at a concentration of greater than about 5 mg/ml (e.g., greater than 6 mg/ml, 7 mg/ml, 8 mg/ml, 9 mg/ml, 10 mg/ml, 15 mg/ml, 20 mg/ml, 25 mg/ml, 30 mg/ml, 40 mg/ml, 50 mg/ml, 75 mg/ml, or 100 mg/ml).

[0012] In some embodiments, the composition further comprises one or more of (i) a buffering agent, (ii) a surfactant, or (iii) a tonicifier. In some embodiments, the composition has a pH of approximately 3.0-8.0 (e.g., 4.0-7.5, 5.0-7.5, 5.5-7.7, 5.5-7.0, 6.0-7.0, 6.5-7.5, 6.5-7.0, or 5.5-6.5). In some embodiments, the composition comprises a replacement enzyme in a formulation that is not synthetic CSF.

[0013] In some embodiments, the composition is administered at a single dose volume of less than about 15 mL (e.g., less than about 10 ml, 9 ml, 8 ml, 7 ml, 6 ml, 5 ml, 4 ml, 3 ml, 2 ml, 1.5 ml, 1.0 ml, or 0.5 ml).

[0014] In some embodiments, the intrathecal administration of the composition does not result in substantial adverse effect in the subject. In certain embodiments, the intrathecal administration of the composition does not result in an adaptive T-cell mediated immune response.

[0015] In yet another aspect, the present invention provides methods including a step of administering to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme, which administering involves intrathecal administration of the composition in absence of concurrent immunosuppressant therapy. In some embodiments,

2017202349 10 Apr 2017

the method does not involve an immune tolerance induction in the subject being treated. In certain embodiments, the method does not involve a pre-treatment or preconditioning of the subject using T-cell immunosuppressive agent.

[0016] In a further aspect, the present invention provides methods including a step of administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme at a therapeutically effective dose and an administration interval such that at least about 10% (e.g., at least about 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) of normal levels or activities of the lysosomal enzyme in one or more tissues of brain, spinal cord and peripheral organs is achieved.

[0017] In some embodiments, the one or more tissues of brain to which the enzyme is delivered comprise a meningeal tissue. In some embodiments, the meningeal tissue is selected from the group consisting of pia mater, dura mater, and arachnoid tissue.

[0018] In some embodiments, the one or more tissues of brain to which the enzyme is delivered comprise a tissue of the cerebrum. In certain embodiments, the tissue of the cerebrum is a surface or shallow tissue of the cerebrum. In certain embodiments, the surface or shallow tissue of the cerebrum is selected from the group consisting of pia mater tissues, cerebral cortical ribbon tissues, hippocampus, tissues within 4 mm from the surface of the surface of the cerebrum, Virchow Robin space, blood vessels within the VR space, the hippocampus, portions of the hypothalamus on the inferior surface of the brain, the optic nerves and tracts, the olfactory bulb and projections, and combinations thereof.

[0019] In some embodiments, the tissue of the cerebrum to which the enzyme is delivered is a deep tissue of the cerebrum. In certain embodiments, the deep tissue of the cerebrum is selected from the group consisting of tissues internal to the cerebral cortical ribbon, tissues below 4 mm from the surface of the surface of the cerebrum, tissues below 6 mm from the surface of the surface of the cerebrum, tissues below 10 mm from the surface of the surface of the cerebrum, the diencephalon, the hypothalamus, thalamus, prethalamus, and subthalamus, the metencephalon, the cerebral peduncles, the red nucleus, the cranial nerve III nucleus, deep

2017202349 10 Apr 2017

grey matter, the lentiform nuclei, the basal ganglia, caudate, putamen, amygdala, globus pallidus, and combinations thereof.

[0020] In some embodiments, the one or more tissues of brain to which the enzyme is delivered comprise a tissue of the cerebellum. In certain embodiments, the tissue of the cerebellum is selected from the group consisting of tissues of the molecular layer, tissues of the Purkinje cell layer, tissues of the Granular cell layer, cerebellar peduncles, and combination thereof. In some embodiments, the tissue of the cerebellum is a deep tissue of the cerebellum. In certain embodiments, the deep tissue of the cerebellum is selected from the group consisting of tissues of the Purkinje cell layer, tissues of the Granular cell layer, deep cerebellar white matter tissue, and deep cerebellar nuclei tissue.

[0021] In some embodiments, the one or more tissues of brain to which the enzyme is delivered comprise a tissue of the brainstem. In certain embodiments, the tissue of the brainstem is selected from the group consisting of brain stem white matter tissue and/or brain stem nuclei tissue.

[0022] In some embodiments, the one or more tissues of the spinal cord to which the enzyme is delivered is a surface or shallow tissue of the spinal cord. In certain embodiments, the surface or shallow tissue of the spinal cord is selected from the group consisting of pia matter, the tracts of white matter, and tissue within 4 mm from the surface of the surface of the spinal cord. In some embodiments, the one or more tissues of the spinal cord is a deep tissue of the spinal cord. In certain embodiments, the deep tissue of the spinal cord is selected from the group consisting of spinal cord grey matter and ependymal cells, and tissue below 4 mm from the surface of the surface of the spinal cord.

[0023] In some embodiments, the one or more tissues of brain to which the enzyme is delivered comprise surface or shallow tissues. In certain embodiments, the surface or shallow tissues are selected from the group consisting of pia mater, dura mater, and arachnoid tissues of meningeal, pia mater tissues, cerebral cortical ribbon tissues, tissues within 4 mm from the surface of the surface of the cerebrum, and combination thereof.

2017202349 10 Apr 2017

[0024] In some embodiments, the tissues to which the enzyme is delivered comprise deep tissues. In certain embodiments, the deep brain tissues are selected from deep white matter, of the cerebrum, deep gray matter of the spinal cord, corpus callosum, periventricular tissue, thalamus, basal ganglia, diencephalon, fimbria, tissues below the cerebral cortical ribbon, tissues below 4 mm from the surface of the surface of the cerebrum, tissues below 6 mm from the surface of the surface of the cerebrum, tissues below 10 mm from the surface of the surface of the cerebrum, Purkinje cell layer, tissues of the Granular cell layer, deep cerebellar white matter tissue, and deep cerebellar nuclei tissue, and combination thereof.

[0025] In some embodiments, the therapeutically effective dose ranges from 0.005 mg/kg brain weight to 100 mg/kg brain weight. In certain embodiments, the therapeutically effective dose is greater than 1 mg/kg brain weight (e.g., greater than 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 mg/kg brain weight). In certain embodiments, the therapeutically effective dose is greater than 10 mg/kg brain weight. In certain embodiments, the therapeutically effective dose is greater than 30 mg/kg brain weight.

[0026] In some embodiments, the administration interval is once every two weeks. In some embodiments, the administration interval is once every month. In some embodiments, the administration interval is once every two months. In some embodiments, the administration interval is twice per month. In some embodiments, the administration interval is once every week. In some embodiments, the administration interval is twice or several times per week. In some embodiments, the administration is continuous, such as through a continuous perfusion pump.

[0027] In another aspect, the present invention provides methods including a step of administering to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme, which administering involves intrathecal administration of the composition so that the replacement enzyme is delivered to a deep brain tissue at least 5 mm below the external surface (e.g., at least 6 mm, 7 mm, 8 mm, 9 mm, 10 mm, 11 mm, 12 mm, or deeper below the external surface). In some embodiments, the replacement enzyme is delivered

2017202349 10 Apr 2017

to a deep brain tissue at least 10 mm below the external surface. In certain embodiments, the replacement enzyme is specifically delivered to cellular lysosomes of the deep brain tissue.

[0028] In some embodiments, the deep brain tissues to which the enzyme is delivered are selected from deep white matter, of the cerebrum, deep gray matter of the spinal cord, corpus collosum, periventricular tissue, thalamus, fimbria, tissues below the cerebral cortical ribbon, tissues below 4 mm from the surface of the surface of the cerebrum, tissues below 6 mm from the surface of the surface of the cerebrum, tissues below 10 mm from the surface of the surface of the cerebrum, Purkinje cell layer, tissues of the Granular cell layer, deep cerebellar white matter tissue, and deep cerebellar nuclei tissue, and combination thereof.

[0029] In yet another aspect, the present invention provides methods including a step of administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme that is produced from human cells.

[0030] In some embodiments, the lysosomal storage disease is selected from the group consisting of aspartylglucosaminuria, cholesterol ester storage disease, Wolman disease, cystinosis, Danon disease, Fabry disease, Farber lipogranulomatosis, Farber disease, fucosidosis, galactosialidosis types I/II, Gaucher disease types I/II/III, globoid cell leukodystrophy, Krabbe disease, glycogen storage disease II, Pompe disease, GM1- gangliosidosis types I/II/III, GM2- gangliosidosis type I, Tay Sachs disease, GM2- gangliosidosis type II, Sandhoff disease, GM2- gangliosidosis, α -mannosidosis types I/II, β -mannosidosis, metachromatic leukodystrophy, mucopolipidosis type I, sialidosis types I/II, mucopolipidosis types II /III, mucopolipidosis type IV, I-cell disease, mucopolipidosis type IIIC pseudo-Hurler polydystrophy, mucopolysaccharidosis type I, mucopolysaccharidosis type II, Hunter syndrome, mucopolysaccharidosis type IIIA, Sanfilippo syndrome type A, B, or D (mucopolysaccharidosis type IIIB, mucopolysaccharidosis type IIIC, mucopolysaccharidosis type IIID), mucopolysaccharidosis type IVA, Morquio syndrome, mucopolysaccharidosis type IVB, mucopolysaccharidosis type VI, mucopolysaccharidosis type VII, Sly syndrome, mucopolysaccharidosis type IX, multiple sulfatase deficiency, neuronal ceroid lipofuscinosis, CLN1 Batten disease, CLN2 Batten disease, Niemann-Pick disease types

2017202349 10 Apr 2017

A/B, Niemann-Pick disease type C1, Niemann-Pick disease type C2, pycnodysostosis, Schindler disease types I/II, Gaucher disease and sialic acid storage disease

[0031] In some embodiments, the lysosomal storage disease is selected from the group consisting of Hunters Syndrome, metachromatic leukodystrophy (MLD) disease, Sanfilippo syndrome type A, Sanfilippo syndrome type B, and globoid cell leukodystrophy (GLD) disease. In certain embodiments, the replacement enzyme is selected from the group consisting of recombinant iduronate-2-sulfatase (I2S), arylsulfatase A (ASA), heparan N-sulfatase (HNS), alpha-N-acetylglucosaminidase (Naglu) and β -galactosidase (GLC). In some embodiments, the replacement enzyme contains mannose-6-phosphate (M6P) residues. In some embodiments, the replacement enzyme is a fusion protein comprising a lysosomal targeting moiety.

[0032] In some embodiments, the replacement enzyme is delivered to neurons, glial cells, perivascular cells and/or meningeal cells. In certain embodiments, the replacement enzyme is further delivered to the neurons in the spinal cord.

[0033] In some embodiments, the intrathecal administration further results in systemic delivery of the replacement enzyme in peripheral target tissues. In certain embodiments, the peripheral target tissues are selected from liver, kidney, and/or heart, endothelium, bone marrow and bone marrow derived cells, spleen, lung, lymph node, bone and cartilage, ovary and testis.

[0034] In some embodiments, the intrathecal administration results in lysosomal localization of the replacement enzyme in brain target tissues, spinal cord neurons and/or peripheral target tissues. In some embodiments, the intrathecal administration results in reduction of GAG storage in the brain target tissues, spinal cord neurons and/or peripheral target tissues. In certain embodiments, the GAG storage is reduced by at least 20%, 40%, 50%, 60%, 80%, 90%, 1-fold, 1.5-fold, or 2-fold as compared to a control.

[0035] In some embodiments, the intrathecal administration results in reduced vacuolization in neurons. In some embodiments, the neurons comprise Purkinje cells.

[0036] In some embodiments, the intrathecal administration results in increased enzymatic activity of the replacement enzyme in the brain target tissues, spinal cord neurons

2017202349 10 Apr 2017

and/or peripheral target tissues. In certain embodiments, the enzymatic activity is increased by at least 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold or 10-fold as compared to a control. In certain embodiments, the increased enzymatic activity is at least approximately 10 nmol/hr/mg, 20 nmol/hr/mg, 40 nmol/hr/mg, 50 nmol/hr/mg, 60 nmol/hr/mg, 70 nmol/hr/mg, 80 nmol/hr/mg, 90 nmol/hr/mg, 100 nmol/hr/mg, 150 nmol/hr/mg, 200 nmol/hr/mg, 250 nmol/hr/mg, 300 nmol/hr/mg, 350 nmol/hr/mg, 400 nmol/hr/mg, 450 nmol/hr/mg, 500 nmol/hr/mg, 550 nmol/hr/mg or 600 nmol/hr/mg.

[0037] In some embodiments, the enzymatic activity is increased in the lumbar region. In certain embodiments, the increased enzymatic activity in the lumbar region is at least approximately 500 nmol/hr/mg, 600 nmol/hr/mg, 700 nmol/hr/mg, 800 nmol/hr/mg, 900 nmol/hr/mg, 1000 nmol/hr/mg, 1500 nmol/hr/mg, 2000 nmol/hr/mg, 3000 nmol/hr/mg, 4000 nmol/hr/mg, 5000 nmol/hr/mg, 6000 nmol/hr/mg, 7000 nmol/hr/mg, 8000 nmol/hr/mg, 9000 nmol/hr/mg, or 10,000 nmol/hr/mg.

[0038] In some embodiments, the lysosomal storage disease is associated with peripheral symptoms and the method further comprises administering the replacement enzyme intravenously to the subject. In certain embodiments, the intravenous administration is no more frequent than weekly administration (e.g., no more frequent than biweekly, monthly, once every two months, once every three months, once every four months, once every five months, or once every six months). In certain embodiments, the intravenous administration is more frequent than monthly administration, such as twice weekly, weekly, every other week, or twice monthly. In some embodiments, intravenous and intrathecal administrations are performed on the same day. In some embodiments, the intravenous and intrathecal administrations are not performed within a certain amount of time of each other, such as not within at least 2 days, within at least 3 days, within at least 4 days, within at least 5 days, within at least 6 days, within at least 7 days, or within at least one week. In some embodiments, intravenous and intrathecal administrations are performed on an alternating schedule, such as alternating administrations weekly, every other week, twice monthly, or monthly. In some embodiments, an intrathecal administration replaces an intravenous administration in an administration schedule, such as in a schedule of intravenous administration weekly, every other week, twice monthly, or monthly, every third or fourth or fifth administration in that schedule can be replaced with an intrathecal administration

2017202349 10 Apr 2017

in place of an intravenous administration. In some embodiments, an intravenous administration replaces an intrathecal administration in an administration schedule, such as in a schedule of intrathecal administration weekly, every other week, twice monthly, or monthly, every third or fourth or fifth administration in that schedule can be replaced with an intravenous administration in place of an intravenous administration. In some embodiments, intravenous and intrathecal administrations are performed sequentially, such as performing intravenous administrations first (e.g., weekly, every other week, twice monthly, or monthly dosing for two weeks, a month, two months, three months, four months, five months, six months, or a year or more) followed by intrathecal administrations (e.g., weekly, every other week, twice monthly, or monthly dosing for more than two weeks, a month, two months, three months, four months, five months, six months, or a year or more). In some embodiments, intrathecal administrations are performed first (e.g., weekly, every other week, twice monthly, monthly, once every two months, once every three months dosing for two weeks, a month, two months, three months, four months, five months, six months, or a year or more) followed by intravenous administrations (e.g., weekly, every other week, twice monthly, or monthly dosing for more than two weeks, a month, two months, three months, four months, five months, six months, or a year or more).

[0039] In some embodiments, the lysosomal storage disease is associated with peripheral symptoms and the method includes administering the replacement enzyme intrathecally but does not involve administering the replacement enzyme intravenously to the subject. In certain embodiments, the intrathecal administration of the replacement enzymes ameliorates or reduces one or more of the peripheral symptoms of the enzyme replacement deficiency of the subject.

[0040] In another aspect, the present invention provides methods of treating Hunters Syndrome including a step of administering intrathecally to a subject in need of treatment a recombinant iduronate-2-sulfatase (I2S) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the Hunters Syndrome is reduced in intensity, severity, or frequency, or has delayed onset. In some embodiments, the at least one symptom or feature of the Hunters Syndrome is cognitive impairment; white matter lesions; dilated perivascular spaces in the brain parenchyma, ganglia, corpus callosum, and/or brainstem; atrophy; and/or ventriculomegaly.

2017202349 10 Apr 2017

[0041] In yet another aspect, the present invention provides methods of treating metachromatic leukodystrophy (MLD) disease including a step of administering intrathecally to a subject in need of treatment a recombinant arylsulfatase A (ASA) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the MLD disease is reduced in intensity, severity, or frequency, or has delayed onset. In some embodiments, the at least one symptom or feature of the MLD disease is increased intracranial pressure, hydrocephalus ex vacuo, accumulated sulfated glycolipids in the myelin sheaths in the central and peripheral nervous system and in visceral organs, progressive demyelination, axonal loss within the CNS and PNS, and/or motor and cognitive dysfunction.

[0042] In still another aspect, the present invention provides methods of treating Sanfilippo syndrome type A (Sanfilippo A) disease including a step of administering intrathecally to a subject in need of treatment a recombinant heparan N-sulfatase (HNS) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the Sanfilippo A disease is reduced in intensity, severity, or frequency, or has delayed onset.

[0043] In another aspect, the present invention provides methods of treating Sanfilippo syndrome type B (Sanfilippo B) disease including a step of administering intrathecally to a subject in need of treatment a recombinant alpha-N-acetylglucosaminidase (Naglu) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the Sanfilippo B disease is reduced in intensity, severity, or frequency, or has delayed onset.

[0044] In some embodiments, the at least one symptom or feature of the Sanfilippo A or Sanfilippo B disease is hearing loss, impaired speech development, deficits in motor skills, motoric hyperactivity, progressive cognitive impairment, , aggressiveness and/or sleep disturbances.

[0045] In some embodiments, the recombinant Naglu enzyme is a fusion protein comprising Naglu and a lysosomal targeting moiety. In certain embodiments, the lysosomal targeting moiety is IGF-II.

2017202349 10 Apr 2017

[0046] In another aspect, the present invention provides methods of treating globoid cell leukodystrophy (GLD) disease including a step of administering intrathecally to a subject in need of treatment a recombinant β -galactosidase (GLC) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the GLD disease is reduced in intensity, severity, or frequency, or has delayed onset. In some embodiments, the at least one symptom or feature of the GLD disease is irritability, convulsion, mental deterioration, deafness, blindness, myoclonic seizures, excessive muscle tone, developmental delay, regression of developmental skills, hypersensitivity, tremor, ataxia, spasticity, episodic severe vomiting, leukodystrophy, cerebral atrophy, impaired development of globoid cells and/or demyelination.

[0047] In yet another aspect, the present invention provides devices for intrathecal administration, including a fluid access port; a hollow body having a first flow orifice in fluid communication with the fluid access port and a second flow orifice configured for insertion into spinal cord; and a securing mechanism for securing the insertion of the hollow body in the spinal cord. In some embodiments, the securing mechanism comprises one or more nobs mounted on the surface of the hollow body and a sutured ring adjustable over the one or more nobs. In some embodiments, the fluid access port comprises a reservoir. In certain embodiments, the fluid access port is implantable. In certain embodiments, the fluid access port is an injectable port. In some embodiments, the fluid access port is a mechanical pump.

[0048] As used in this application, the terms “about” and “approximately” are used as equivalents. Any numerals used in this application with or without about/approximately are meant to cover any normal fluctuations appreciated by one of ordinary skill in the relevant art.

[0049] Other features, objects, and advantages of the present invention are apparent in the detailed description that follows. It should be understood, however, that the detailed description, while indicating embodiments of the present invention, is given by way of illustration only, not limitation. Various changes and modifications within the scope of the invention will become apparent to those skilled in the art from the detailed description.

2017202349 10 Apr 2017

BRIEF DESCRIPTION OF THE DRAWINGS

- [0050] The drawings are for illustration purposes only, not for limitation.
- [0051] Figure 1 illustrates an exemplary diagram of an intrathecal drug delivery device (IDDD) with a securing mechanism.
- [0052] Figure 2A depicts exemplary locations within a patient's body where an IDDD may be placed; Figure 2B depicts various components of an intrathecal drug delivery device (IDDD); and Figure 2C depicts an exemplary insertion location within a patient's body for IT-lumbar injection.
- [0053] Figure 3 depicts exemplary results summarizing vehicles tested in adult monkeys.
- [0054] Figure 4 depicts exemplary results illustrating the stability of hGalC in a thermal screen of hGalC as a function of pH.
- [0055] Figure 5 depicts exemplary results illustrating the specific activity of hGalC as a function of pH.
- [0056] Figure 6 depicts exemplary results illustrating a thermal screen of hGalC as a function of salt concentration.
- [0057] Figure 7 depicts exemplary results illustrating sedimentation velocity runs of GalC comparing different ionic strengths in 5mM Na phosphate, pH 6.0 buffer. Figure 7A depicts exemplary results using 50mM NaCl and hGalC. Figure 7B depicts exemplary results illustrating 150mM NaCl and hGalC. Figure 7C depicts exemplary results illustrating 500mM NaCl and hGalC. Figure 7D depicts exemplary results illustrating 150mM NaCl and mouse GalC.
- [0058] Figure 8 depicts exemplary results illustrating GalC AUC profile as a function of salt concentration (1mg/mL GalC, 5mM Na phosphate, pH 6.0)(Y axis = $s^*g(s^*)$; X axis = s^*).
- [0059] Figure 9 depicts exemplary results illustrating a dilution series of hGalC in universal buffer, pH 6.0 (Y-axis = $\langle g(s^*)/C_0 \rangle (1/\text{svedberg})$; X-axis = $s^* (\text{svedbergs})$).

2017202349 10 Apr 2017

- [0060] Figure 10 depicts exemplary results illustrating a GalC AUC profile as a function of pH (1mg/mL, 3mM citrate, phosphate and borate buffer with 50mM NaCl).
- [0061] Figure 11 depicts exemplary results illustrating the baseline reading from a WDA analysis at the highest concentration at pH 6.0, in 5mM Na phosphate and 150mM NaCl.
- [0062] Figure 12 depicts exemplary results illustrating the stressed reading from a WDA analysis at the highest concentration at pH 6.0, in 5mM Na phosphate and 150mM NaCl.
- [0063] Figure 13 graphically compares and overlays baseline and stressed GalC samples.
- [0064] Figure 14 depicts exemplary results illustrating a dilution series of hGalC in the presence of 1% NaTC.
- [0065] Figure 15 depicts exemplary results illustrating a dilution series of hGalC in the presence of 1% NaTC (1.0mg/mL and 0.3mg/mL).
- [0066] Figure 16 depicts exemplary results illustrating the intrinsic fluorescence of hGalC (1mg/mL) in different buffers and pHs.
- [0067] Figure 17 depicts exemplary results illustrating the circular dichroism of hGalC as a function of pH.
- [0068] Figure 18 depicts exemplary results illustrating the group mean concentration of radioactivity in serum, blood and red blood cells of male Sprague-Dawley rats following a single intrathecal dose of ^{125}I -hGalC.
- [0069] Figure 19 depicts exemplary results illustrating the group mean concentrations of radioactivity in serum, heart, kidneys, liver, lungs, spleen of male Sprague-Dawley rats following a single intrathecal dose of ^{125}I -hGalC.
- [0070] Figure 20 depicts exemplary results illustrating the group mean concentrations of radioactivity in serum, heart, kidneys, liver, lungs, spleen of male Sprague-Dawley rats following a single intravenous bolus injection of ^{125}I -hGalC.

2017202349 10 Apr 2017

[0071] Figure 21 depicts exemplary results illustrating the group mean concentrations of radioactivity in serum, heart, kidneys, liver, lungs, spleen of male Sprague-Dawley rats following a single intrathecal dose and intravenous bolus injection of ^{125}I -hGalC.

[0072] Figure 22 depicts exemplary results illustrating the mean concentrations of radioactivity in serum and various tissues (adipose tissues (kidney fast), adrenal glands, bone (femur), muscle (skeletal), sciatic nerve)) of male Sprague-Dawley rats following a single intrathecal dose of ^{125}I -hGalC.

[0073] Figure 23 depicts exemplary results illustrating the mean concentrations of radioactivity in serum and various tissues (adipose tissues (kidney fast), adrenal glands, bone (femur), muscle (skeletal), sciatic nerve)) of male Sprague-Dawley rats following a single intravenous bolus injection of ^{125}I -hGalC.

[0074] Figure 24 depicts exemplary results illustrating the mean concentrations of radioactivity in serum and tissues (adipose tissues (kidney fast), adrenal glands, bone (femur), muscle (skeletal), sciatic nerve)) of male Sprague-Dawley rats following a single intrathecal dose and intravenous bolus injection of ^{125}I -hGalC.

[0075] Figure 25 depicts exemplary results illustrating the mean concentrations of radioactivity in serum, cerebrospinal fluid and various other tissues of male Sprague-Dawley rats following a single intrathecal dose of ^{125}I -hGalC.

[0076] Figure 26 depicts exemplary results illustrating the mean concentrations of radioactivity in serum, cerebrospinal fluid and various other tissues of male Sprague-Dawley rats following a single intravenous bolus injection of ^{125}I -hGalC.

[0077] Figure 27 depicts exemplary results illustrating the mean concentrations of radioactivity in serum, cerebrospinal fluid and tissues of male Sprague-Dawley rats following a single intrathecal dose and intravenous bolus injection of ^{125}I -hGalC.

[0078] Figure 28 depicts exemplary results illustrating the mean concentrations of radioactivity in serum and tissues of male Sprague-Dawley rats following a single intrathecal dose of ^{125}I -hGalC.

2017202349 10 Apr 2017

- [0079] Figure 29 depicts exemplary results illustrating the mean concentrations of radioactivity in serum and tissues of male Sprague-Dawley rats following a single intravenous bolus injection of ¹²⁵I-hGalC.
- [0080] Figure 30 depicts exemplary results illustrating the mean concentrations of radioactivity in serum and tissues of male Sprague-Dawley rats following a single intrathecal dose and intravenous bolus injection of ¹²⁵I-hGalC.
- [0081] Figure 31 depicts exemplary results illustrating that IP administration of rmGalC reduces brain psychosine levels in twitcher mice. Data represents mean ± SEM for n=4 mice per treatment group.
- [0082] Figure 32 depicts exemplary results illustrating increased survival with ICV only and ICV/IP rmGalC therapy.
- [0083] Figure 33 depicts exemplary results illustrating that brain psychosine is significantly reduced after ICV and ICV/IP injections of rmGalC in twitcher mice.
- [0084] Figure 34 depicts exemplary results illustrating improvement in histological markers observed in twitcher mice treated with 40 ug of rmGalC. Glial fibrillary acidic protein (GFAP) was used as an astrocytes marker. Iba 1 was used as a microglia/macrophage marker. Lysosomal associated membrane protein-1 (LAMP-1) was used as a lysosomal marker.
- [0085] Figure 35 depicts exemplary results illustrating psychosine re-accumulation following a single ICV injection of rmGalC or vehicle.
- [0086] Figure 36 depicts exemplary results illustrating percent survival in twitcher mice treated with a single ICV injection of rmGalC at PND19/20. Data represents n= 8 per group.
- [0087] Figure 37 depicts exemplary results illustrating percent survival in mice treated ICV/IP with rmGalC and rhGalC.
- [0088] Figure 38 depicts exemplary results illustrating gait analysis of mice treated with a single ICV injection of rmGalC and rhGalC.

2017202349 10 Apr 2017

- [0089] Figure 39 depicts exemplary results illustrating an antigenic response to rmGalC or rhGalC in twitcher mice.
- [0090] Figure 40 depicts exemplary results illustrating psychosine levels in the CSF of naïve and rhGalC-treated GLD dogs.
- [0091] Figure 41 depicts exemplary results illustrating IHC staining of IT injected GalC in the cerebrum with Group 1 polyclonal antibody.
- [0092] Figure 42 depicts exemplary results illustrating IHC staining of IT injected GalC in the cerebrum with Group 2 antibody.
- [0093] Figure 43 depicts exemplary results illustrating IHC staining of IT injected GalC in the cerebrum with Mouse monoclonal antibody.
- [0094] Figure 44 depicts exemplary results illustrating IHC staining of IT injected GalC in the cerebrum with Mouse monoclonal antibody.
- [0095] Figure 45 depicts exemplary results illustrating IHC staining of IT injected GalC in the liver with Mouse monoclonal antibody.
- [0096] Figure 46 depicts exemplary results illustrating IHC staining of IT injected GalC in the liver with Group 2 polyclonal antibody.
- [0097] Figure 47 depicts exemplary results illustrating mean GalC activity in the brain.
- [0098] Figure 48 depicts exemplary results illustrating mean GalC activity in the liver.
- [0099] Figure 49 depicts exemplary results illustrating GalC immunostaining in the brain at 10X.
- [0100] Figure 50 depicts exemplary results illustrating GalC immunostaining in the brain at 40X.
- [0101] Figure 51 depicts exemplary results illustrating Iba staining of activated microglia at 40X.

2017202349 10 Apr 2017

[0102] Figure 52 depicts exemplary results illustrating LFB/PAS staining in the brain at 10X.

[0103] Figure 53 illustrates exemplary I2S IHC demonstrated I2S detected in the neurons (arrows) in the cerebral and cerebellar cortex including a layer of meningeal cells covering the surface of the brain (arrow heads) following intrathecal injections of 3 doses of I2S. Staining of I2S IHC in 2 dose injected brains was weaker (photo not shown). There was no observed positive I2S staining for any type of cells in the brain of vehicle control animals. 40X.

[0104] Figure 54 depicts exemplary reversal of pathology in the brain of IKO mice after intrathecal-lumbar I2S injection. H&E stained brain tissues showed numerous cellular storage vacuoles (arrows) in the vehicle control animals. Cellular vacuolation was reduced throughout the brain in both 2 dose (photo not shown) and 3 dose injected mice. Marked reduction was found in the 3 dose injected ones. 40X.

[0105] Figure 55 depicts exemplary immunohistochemical staining of LAMP-1, there was a marked reduction of lysosomal activity in the brains after 2 doses (photo not shown) and 3 doses of I2S treatment compared with vehicle controlled mice. The reduction was characterized by the decrease in the number of LAMP-1 positive cells and lighter staining intensity in the regions throughout the brain. 40X.

[0106] Figure 56 illustrates exemplary morphometry results from a comparison of the mean LAMP-1 positive area among wild-type (WT), vehicle untreated and I2S (2 and 3 doses) mice in the cerebral cortex (Cortex), caudate nucleus (CP), thalamus (TH), white matter (WM) and cerebellum (CBL) confirmed that there were significant reductions in the LAMP-1 positive staining in all areas of the brain evaluated. Data are represented as the mean \pm s.d. # = $P < 0.05$; * = $P < 0.01$; ** = $P < 0.001$.

[0107] Figure 57 depicts exemplary electron micrographs of brain cells showed pathological improvements at the ultrastructural level. Neurons of vehicle treated mice had lamellated inclusions, zebra body-like structures, and vacuoles containing granular storage material (insert), which was reduced in I2S injected mice. Oligodendrocytes of vehicle treated mice showed large electron-lucent storage vacuoles (arrow) while oligodendrocytes of I2S-

2017202349 10 Apr 2017

injected mice had minimal vacuolation. Scale bar: in neurons, 2 μ m; in oligodendrocytes, 500 nm.

[0108] Figure 58 depicts exemplary immunohistochemistry results demonstrating I2S detected in sinusoidal cells of the liver following intrathecal injections of 3 doses of I2S. 2S IHC staining in 2 dose injected livers was weaker (photo not shown). There was no observed positive I2S staining in the liver of vehicle controlled animals. 40X.

[0109] Figure 59 depicts exemplary tissue from the liver. Severe cellular vacuolation and abnormally high lysosomal activity is revealed by H&E staining and strong LAMP-1 immunostaining were found in vehicle controlled animals compared to WT ones. Marked reduction of cellular vacuolation and LAMP-1 immunostaining was found after intertheal treatment with 3 and 2 (photo not shown) doses of I2S treatment. H&E staining revealed intracytoplasmic vacuolization was almost completely disappear with a nearly normal liver cell structure. H&E, 40X; LAMP-1, 20X.

[0110] Figure 60 depicts exemplary tissues showing cerebrum of a 3mg treatment group animal. Positive I2S staining was observed in meningeal cells. 4X.

[0111] Figure 61 depicts exemplary tissues showing cerebrum of a 30 mg treatment group animal. Positive I2S staining was observed in neurons and meningeal cells. 4X.

[0112] Figure 62 depicts exemplary tissues showing cerebrum of 100 mg treatment group animal. Positive I2S staining in neurons and meningeal cells was stronger than in 3 and 30 mg treated animals. 4X.

[0113] Figure 63 depicts exemplary tissues showing cerebrum of a 150 mg treatment group animal. A large population of neurons was observed as being I2S positive along with strongly positive meningeal cells.

[0114] Figure 64 depicts exemplary tissues showing I2S positive neurons and glial cells, along with meningeal cells, within layer I of the cerebrum in a 30 mg treatment group animal. 40X.

2017202349 10 Apr 2017

- [0115] Figure 65 depicts exemplary tissues showing I2S positive neurons, glial cells, along with perivascular cells, within layer III of the cerebrum in a 30 mg treatment group animal. 40X.
- [0116] Figure 66 depicts exemplary tissues showing I2S positive neurons and glial cells within the layer VI of cerebrum adjacent to the white matter in a 30 mg treatment group animal. 40X.
- [0117] Figure 67 depicts exemplary tissues showing strongly positive I2S staining in the neurons (cerebrum) of a 150 mg treatment group animal. 100X.
- [0118] Figure 68 depicts exemplary tissue showing I2S immunostaining of the cervical spinal cord in a 150 mg treatment group. 4X.
- [0119] Figure 69 depicts exemplary tissue showing strong I2S immunostaining in the lumbar spinal cord of a 150 mg treatment group animal. 4X.
- [0120] Figure 70 depicts exemplary tissue showing strongly positive I2S immunostaining of meningeal cells, glial cells, and epi/peri/endoneurium (connective cells) in the lumbar section of a 150 mg treatment group animal. 40X.
- [0121] Figure 71 depicts an image showing that neurons in the lumbar spinal cord of a 150 mg treatment group animal were strongly I2S positive. 40X .
- [0122] Figure 72 depicts exemplary results from a liver from a 3 mg treatment group animal. Only sinusoidal cells were I2S positive. 40X.
- [0123] Figure 73 depicts exemplary results from a liver from a 30 mg treatment group animal. Sinusoidal cells and hepatocytes were I2S positive. 40X.
- [0124] Figure 74 depicts exemplary results from a liver from a 100 mg treatment group animal. I2S immunostaining was strong in the sinusoidal cells and the hepatocytes. 40X.
- [0125] Figure 75 depicts exemplary results from a liver from a 150 mg treatment group animal. Strongly positive I2S staining was identified in sinusoidal cells and hepatocytes. 40X.

2017202349 10 Apr 2017

- [0126] Figure 76 depicts exemplary results from a heart from a 3 mg treatment group animal. I2S immunostaining was negative. 40X.
- [0127] Figure 77 depicts exemplary results from a heart from a 30 mg treatment group animal. Interstitial cells were I2S positive. 40X.
- [0128] Figure 78 depicts exemplary results from a heart from a 100 mg treatment group animal. Positive interstitial cell staining for I2S was observed. 40X.
- [0129] Figure 79 depicts exemplary results from a heart from a 150 mg treatment group animal. Strongly positive interstitial cell staining for I2S was observed. 40X.
- [0130] Figure 80 depicts exemplary results from a kidney from a 3 mg treatment group animal. I2S immunostaining was negative. 40X.
- [0131] Figure 81 depicts exemplary results from a kidney from a 30 mg treatment group animal. Glomerular and interstitial cells were I2S positive.
- [0132] Figure 82 depicts exemplary results from a kidney from a 100 mg treatment group animal. Increased glomerular and interstitial cell staining for I2S was observed. 40X.
- [0133] Figure 83 depicts exemplary results from a kidney from a 150 mg treatment group animal. Positive I2S staining of proximal tubular, glomerular and interstitial cells was observed. 40X.
- [0134] Figure 84 illustrates results of immunohistochemistry (IHC) studies evaluating the CNS tissues of cynomolgus monkeys administered weekly doses of iduronate-2-sulfatase (I2S). As determined by (IHC), there was cellular deposition of I2S throughout the CNS. In the gray matter I2S was detected in the neurons of the cerebrum, cerebellum, brain stem, and spinal cord of all groups in a dose-dependent manner. In the surface gray matter of the higher dose groups, large numbers of cerebral neurons were positive for I2S staining in the surface cortex (Figure 84A). I2S was also detected in neurons in the thalamus (Figure 84B), hippocampus (Figure 84C), caudate nucleus (Figure 84D) and spinal cord (Figure 84E). Meningial and perivascular cells were also I2S staining positive (Figure 84F). The identified scale bars correspond to 25um.

2017202349 10 Apr 2017

[0135] Figure 85 graphically compares the clearance of iduronate-2-sulfatase (I2S) in the cranial and spinal pools by plotting the amount of I2S in such pools relative to the time following administration.

[0136] Figure 86 illustrates the dose dependant gray matter deposition of intrathecally-administered iduronate-2-sulfatase (I2S) to non-human primates over six months. The illustrated staining intensity corresponds with accumulation of iduronate-2-sulfatase in the thalamus. In the present Figure 86, the nuclei are counterstained by DAPI and appear as blue and protein (I2S) appears as green.

[0137] Figure 87 illustrates the dose dependant accumulation of intrathecally-administered iduronate-2-sulfatase (I2S) to non-human primates following a single injection and following multiple injections over a six month period. The illustrated staining intensity corresponds with accumulation of I2S protein in the cerebral cortex.

[0138] Figure 88 demonstrates the cellular localization of iduronate-2-sulfatase (I2S) throughout the cerebrum of a primate. Figure 88A illustrates the cross-sectional view of brain tissue extracted from the cerebrum of the primate, while Figure 88B illustrates that particular areas of the region corresponding to three areas of white matter tissue (designated W1, W2 and W3), the white matter near the ventricle (VW) and the surface gray matter (SG) tissues of the section identified in Figure 88A.

[0139] Figure 89A – D illustrate neuronal and oligodendrocyte uptake and axonal association of intrathecally-administered iduronate-2-sulfatase (I2S) to primates following monthly injections over a six month period. In particular, Figure 89A, Figure 89B, Figure 89C and Figure 89D are illustrative of a filament staining of the cerebrum tissues of the primate intrathecally administered iduronate-2-sulfatase (I2S) and respectively correspond to the three areas of the white matter (W1, W2 and W3) and the surface gray matter (SG) regions identified in Figure 87B. Figure 89A illustrates oligodendrocyte uptake of intrathecally-administered I2S in the white matter (W1) tissues. Figure 89B and Figure 89C illustrate oligodendrocyte uptake and axonal association of the intrathecally-administered I2S in the W2 and W3 white matter tissues respectively. Figure 89D illustrates neuronal uptake of the intrathecally-administered I2S in the surface gray matter (SG) tissues.

2017202349 10 Apr 2017

[0140] Figure 90 illustrates the cellular identification of iduronate-2-sulfatase in the white matter near the ventricle (VW) of a non-human primate (arrows in top left). As depicted in the superimposed image (arrows in bottom right), the iduronate-2-sulfatase is not associated with myelin (top right). In the present Figure 90, the nuclei are counterstained by DAPI (bottom left) Protein (I2S) appears in the top left box.

[0141] Figure 91 illustrates staining in the tissues of healthy Beagle dogs that were intracerebroventricularly (ICV) or intrathecally (IT) administered a single injection of iduronate-2-sulfatase (I2S). As depicted in images a-h, I2S was widely distributed throughout the gray matter of both the IT and ICV groups as determined by immunohistochemistry (IHC). Images a and b illustrate that in the cerebral cortex, neurons were positive for I2S in all six neuronal layers, from the surface molecular layer to the deep internal layer in both IT and ICV groups. Images c and d illustrate that in the cerebellar cortex of the IT and ICV groups I2S was detected in neurons, including Purkinje cells. Similarly, images e and f illustrate that in both IT and ICV groups a large population of neurons in the hippocampus were positive for I2S. Finally, images g and h demonstrate that I2S-positive neurons were also found in the thalamus and caudate nucleus in the both the IT and ICV groups. In the present Figure 91, I2S staining is indicated with arrows.

[0142] Figure 92 comparatively illustrates corpus callosum tissues of iduronate-2-sulfatase knock-out (IKO) mice that were either untreated or were administered I2S intrathecally. As depicted, the treated IKO mice exhibited a reduction of cellular vacuolation characteristic of certain lysosomal storage disorders in the corpus callosum and fornix tissues of the I2S-treated IKO mouse.

[0143] Figure 93A illustrates a marked reduction in the presence of lysosomal associated membrane protein 1 (LAMP1), a lysosomal disease pathological biomarker, in the surface cerebral cortex tissues of the treated IKO mouse (Figure 93A) relative to the untreated IKO control mouse (Figure 93B) under both 20X and 40X magnification.

[0144] Figure 94 depicts an exemplary intrathecal drug delivery device (IDDD).

2017202349 10 Apr 2017

- [0145] Figure 95 depicts an exemplary PORT-A-CATH[®] low profile intrathecal implantable access system.
- [0146] Figure 96 depicts an exemplary intrathecal drug delivery device (IDDD).
- [0147] Figure 97 depicts an exemplary intrathecal drug delivery device (IDDD), which allows for in-home administration for CNS enzyme replacement therapy (ERT).
- [0148] Figure 98 is an exemplary illustration showing the effect of vacuolization after a single intra-cerebral injection of idursulfase in neurons (Purkinje cells).
- [0149] Figure 99 is an exemplary illustration showing I2S activity in the brain by dose and region.
- [0150] Figure 100 is an exemplary illustration showing data of immunohistochemical localization of Idursulfase at different depths of the cerebral cortex.
- [0151] Figure 101 is an exemplary illustration showing I2S activity in the spinal cord of monkey following intrathecal dosing with idursulfase.
- [0152] Figure 102 is an exemplary illustration showing I2S activity in monkey liver, heart and kidney after intrathecal dosing with Idursulfase.
- [0153] Figure 103 depicts an exemplary schematic for an escalation Hunter-IT trial program.
- [0154] Figure 104 is an exemplary illustration showing the measurements of I2S concentrations in various sections of brain tissue after 30 mg dose. Different plots correspond to different times of measurement.
- [0155] Figure 105 is an exemplary illustration showing the measurements of I2S concentration after administration over time via various routes of administration for various product concentrations.
- [0156] Figure 106 depicts PET Imaging of ¹²⁴I-labeled Idursulfase-IT in Cynomolgus Monkeys at t=5 hours Following IV, IT-L, or ICV dosing.

2017202349 10 Apr 2017

[0157] Figure 107 is an exemplary illustration showing ASA concentration data in serum after intravenous administration.

[0158] Figure 108 is an exemplary illustration showing ASA concentration data in serum after IT-lumbar administration.

[0159] Figure 109 is an exemplary illustration showing ASA concentration in CSF after IV administration.

[0160] Figure 110 is an exemplary illustration showing the ASA concentration in CSF after IT-lumbar administration.

[0161] Figure 111 depicts exemplary photo-micrographs of brain tissue, meninges, infiltrates (mid and high dose groups, both sexes) after treatment.

[0162] Figure 112 depicts another exemplary photo-micrographs of brain tissue, meninges, infiltrates (mid and high dose groups, both sexes) after treatment.

[0163] Figure 113 depicts exemplary photo-micrographs of brain tissue, perivascular, infiltrates (mid dose males; high dose females) after treatment.

[0164] Figure 114 depicts exemplary Alcian blue staining of spinal cord of immunotolerant MLD Mice treated with rhASA1 and results illustrating sulfatide reduction as determined by Alcian blue staining of the cervical spinal cord in animals that received intrathecal injections of rhASA at days 1, 8, 15 and 22 at doses of 520 mg/kg brain weight or vehicle control. As demonstrated, treatment with intrathecally injected rhASA resulted in reduction of sulfatide accumulation in the spinal cord, including in the cervical region of the spinal cord.

[0165] Figure 115 illustrates exemplary morphometry analysis of Alcian blue stained spinal cord sections from immunotolerant MLD Mice treated with rhASA1 and results illustrating optical density of Alcian blue in total spinal cord (T-Spinal Cord), total gray matter (T-GM), lumbar gray matter (L-GM), cervical gray matter (C-GM), total white matter (T-WM), lumbar white matter (L-WM), and cervical white matter (C-WM) as determined by morphometry

2017202349 10 Apr 2017

analysis. As demonstrated, a statistically significant reduction in Alcian blue staining was observed in animals treated with rhASA as compared to a vehicle control.

[0166] Figure 116 depicts exemplary reduction of LAMP staining in white matter (Fimbria) of immunotolerant MLD mice treated with rhASA1 depicts exemplary results illustrating LAMP-1 levels in fimbria as determined by immunohistochemistry. Magnification = 20X. As demonstrated, treatment with intrathecally injected rhASA resulted in reduction of LAMP-1 in the cerebral white matter.

[0167] Figure 117 illustrates exemplary morphometry analysis of LAMP staining of brain from immunotolerant MLD mice treated with rhASA1 and results illustrating LAMP-1 staining intensity in corpus collosum (CC), fimbria (F), cerebellar white matter (CB-WM) and brain stem (BS) of animals treated with 20 mg/kg intravenous rhASA, 300 mg/kg brain weight intrathecal rhASA, 520 mg/kg brain weight intravenous rhASA, or vehicle control.

[0168] Figure 118 is an exemplary illustration showing the concentration of ASA in brain punches of vehicle-dosed juvenile Cynomolgus monkeys following every-other-week (EOW) IT dosing for 6-months (main necropsy).

[0169] Figure 119 is an exemplary illustration showing the concentration of ASA in brain punches of juvenile Cynomolgus monkeys following EOW IT dosing of rhASA1 at 1.8mg/dose for 6-months (main necropsy).

[0170] Figure 120 is an exemplary illustration showing the concentration of ASA in brain punches of juvenile Cynomolgus monkeys following EOW IT dosing of rhASA1 at 6.0 mg/dose for 6- months (main necropsy).

[0171] Figure 121 is an exemplary illustration showing the concentration of rhASA in brain punches of juvenile cynomolgus monkeys following EOW IT dosing of rhASA1 at 18.6 mg/dose for 6- months (main necropsy).

[0172] Figure 122 is an exemplary illustration showing the concentration of rhASA in brain punches of juvenile cynomolgus monkeys following EOW IT dosing (PBS-control) for 6-months (main necropsy).

2017202349 10 Apr 2017

[0173] Figure 123 is an exemplary illustration showing the concentration of rhASA in brain punches of juvenile cynomolgus monkeys following EOW IT dosing of vehicle for 6-months (main necropsy).

[0174] Figure 124 is an exemplary illustration showing the concentration of rhASA in brain punches of juvenile cynomolgus 1 monkeys following EOW IT Dosing of rhASA1 at 1.8 mg/dose for 6- months (main necropsy).

[0175] Figure 125 is an exemplary illustration showing the concentration of rhASA in brain punches of juvenile cynomolgus monkeys following EOW IT dosing of rhASA1 at 6.0 mg/dose for 6 months (main necropsy).

[0176] Figure 126 is an exemplary illustration showing the concentration of rhASA in brain punches of juvenile cynomolgus following EOW IT dosing of rhASA1 at 18.6 mg/dose for 6-months (main necropsy).

[0177] Figure 127 is an exemplary illustration showing the concentration of rhASA in selected punches from surface of brain for device control, vehicle, 1.8 mg, 6.0 mg and 18.6 mg treated animals. (male and female separate, device control data is from recovery necropsy, all other data from main necropsy)

[0178] Figure 128 is an exemplary illustration showing the concentration of rhASA in selected punches from deep white area of brain for device control, vehicle, 1.8 mg, 6.0 mg and 18.6 mg treated animals. (male and female separate, device control data is from recovery necropsy, all other data from main necropsy)

[0179] Figure 129 is an exemplary illustration showing the concentration of rhASA in selected punches from deep grey area of brain for device control, vehicle, 1.8 mg, 6.0 mg and 18.6 mg treated animals. (male and female separate, device control data is from recovery necropsy, all other data from main necropsy)

[0180] Figure 130 is an exemplary illustration showing the concentration of rhASA in selected punches from various regions in device control, vehicle, 1.8.mg, 6.0 mg and 18.6 mg

2017202349 10 Apr 2017

treated animals. (male and female combined, device control data is from recovery necropsy, all other data from main necropsy)

[0181] Figure 131 is an exemplary illustration showing the concentration of rhASA in spinal cord sections of juvenile cynomolgus monkeys following EOW IT dosing for 6-months (recovery necropsy).

[0182] Figure 132 is an exemplary illustration showing the concentration of rhASA in liver of juvenile cynomolgus monkeys following EOW IT dosing for 6-Months (047-021) (recovery necropsy).

[0183] Figure 133 is an exemplary illustration showing the anatomical locations of brain punches in the subcortical WM, periventricular WM (and deep white matter) and subcortical WM.

[0184] Figure 134 is an exemplary illustration showing the anatomical locations of brain punches in the corpus callosum and pericallosal subcortical WM, internal capsule – GPi, internal capsule – caudate nucleus, deep white matter, subcortical WM and cortex, putamen, and temporal subcortical WM and cortex.

[0185] Figure 135 is an exemplary illustration showing the anatomical locations of brain punches in the deep grey matter, deep WM (frontal periventricular and subcortical), and subcortical white and cortex – superficial sagittal.

[0186] Figure 136 is an exemplary illustration showing the anatomical locations of brain punches in the corpus callosum and pericallosal subcortical WM, deep subcortical WM, deep grey, periventricular WM, subcortical WM and hippocampus.

[0187] Figure 137 is an exemplary illustration showing the anatomical locations of brain punches in the corpus callosum and deep WM.

[0188] Figure 138 is an exemplary illustration showing the anatomical locations of brain punches in the subcortical WM – occipital lobe and cerebellar white matter, including dentate nucleus (WM).

2017202349 10 Apr 2017

[0189] Figure 139A-G illustrate the concentration of recombinant human arylsulfatase A (ASA) in extracted tissue punches from the brain tissues of adult and juvenile cynomolgus monkeys administered either a vehicle, 1.8mg rhASA or 18.6mg rhASA. Each of Figure 139A-G corresponds to a region of the brain tissue depicted in Figure 134.

[0190] Figure 140A and Figure 140B is an exemplary illustration showing the comparison of the concentrations of recombinant human arylsulfatase A (ASA) detected in the deep white matter (Figure 140A) or in the deep grey matter (Figure 140B) brain tissues of adult and juvenile cynomolgus monkeys which were intrathecally (IT) or intracerebroventricularly (ICV) administered rhASA.

[0191] Figure 141A is an exemplary illustration showing the concentrations of ASA detected in several tissue punches obtained from juvenile (<12 months of age) cynomolgus monkeys IT-administered an 18.6 or a 1.8mg dose of recombinant human arylsulfatase A (rhASA). As illustrated in both Figure 140A-B, the concentration of ASA delivered to the tissues were within, or otherwise exceeded the target therapeutic concentration of 2.5ng/mg rhASA. The anatomical regions of brain tissue which correspond to each of the punch numbers depicted in Figure 140A and Figure 140B are the: subcortical white matter (1); periventricular white matter and deep white matter (2); subcortical white matter (3); subcortical white matter (4); internal capsule (5); internal capsule caudate nucleus (6); deep white matter (7); subcortical white matter and cortex (8); putamen (9); temporal subcortical white matter and cortex (10), deep grey matter (11), deep grey matter (12), frontal periventricular & subcortical (13); subcortical white matter, cortex superficial perifalxian (14); corpus callosum and pericallosal subcortical white matter (15); deep subcortical white matter (16); deep grey matter (17); deep grey matter (18); periventricular white matter (19); deep subcortical white matter (20); hippocampus (21); corpus callosum (22); deep white matter (23); subcortical white matter, occipital lobe (24); and cerebellar white matter (25).

[0192] Figure 142A illustrates the area of deep white matter tissue extracted from a cynomolgus monkey IT-administered 1.8mg of ASA. Figure 142B illustrates immunostaining of the deep white matter tissue and distribution of ASA in relevant cells. In Figure 142B, the protein (ASA) is illustrated in the right bottom box. Figure 142C illustrates that the IT-

2017202349 10 Apr 2017

administered ASA showed organelle co-localization in the deep white matter tissues of the cynomolgus monkey and in particular in the lysosomes. In Figure 142C, the ASA immunostaining is illustrated in the top left box..

[0193] Figure 143 compares the distribution of ¹²⁴I-labeled arylsulfatase A (ASA) using PET scanning 24 hours following either IT- or ICV-administration of such labeled ASA to a cynomolgus monkey.

[0194] Figure 144 illustrates the distribution of ¹²⁴I-labeled ASA immediately following ICV administration to a cynomolgus monkey, and compares the distribution of IT-administered ¹²⁴I-labeled ASA within 2-5 hours. As demonstrated, IT administration delivered the ¹²⁴I-labeled ASA to the same initial compartments (cisternae and proximal spine) as that shown for the ICV administration.

[0195] Figure 145 depicts exemplary ICV and IT administration in a mouse model.

[0196] Figure 146A depicts an exemplary result illustrating CSF concentrations of HNS as a function of time at 1.5, 4.5 and 8.3 mg doses following 6 months of dosing. Figure 146B details an exemplary result illustrating Anti-HNS antibody concentrations in the CSF after 6 months of IT administration of 1.5, 4.5 and 8.3 mg doses in monkeys. Data are shown for male and females combined. Figure 146C details an exemplary result illustrating Anti-HNS antibody concentrations in the CSF after 6 months of IT administration of 1.5, 4.5 and 8.3 mg doses in monkeys following 6 months of dosing. Data are shown for male and females combined. The two highest concentrations (32,205 ng/mL and 15,467 ng/mL) post IT dose 6 at 8.3 mg of HNS were excluded from the plot because no CSF samples were taken predose 6.

[0197] Figure 147 depicts an exemplary result illustrating representative images of tissue sections from the meninges and parenchyma of the brain stained with hematoxylin and eosin. Figure 147A depicts an exemplary result illustrating a low-power view of neutrophilic infiltrates local to the IT catheter in a DC monkey. Figure 147B depicts an exemplary result illustrating a high-power view of eosinophilic infiltrates in the meninges of a high-dose (8.3 mg/dose) monkey; the overall severity of infiltrates was similar to the mid-dose (4.5 mg/dose) group (not shown). Figure 147C depicts an exemplary result illustrating a high-power view of a low-dose

2017202349 10 Apr 2017

(1.5 mg/dose) monkey showing eosinophils in the perivascular space (brain parenchyma). Figure 147D depicts an exemplary result illustrating a low-dose monkey (1.5 mg/dose) showing eosinophils in the perivascular space and adjoining parenchyma. Figure 147E depicts an exemplary result illustrating eosinophils in the spinal cord parenchyma (indicated by arrows) of a low-dose group animal; neurons in the area are normal. Figure 147F depicts an exemplary result illustrating eosinophils and an area of microgliosis (arrows indicate eosinophils; the box indicates an area of microgliosis) in a low-dose (1.5 mg/dose) monkey. There are several large neurons in the area, all of which are normal. Scale bars: 200 μ m.

[0198] Figure 148 depicts an exemplary result illustrating HNS enzyme activity in monkey spinal cords and brains. Figure 148A/B depicts an exemplary result illustrating activity in the spinal cords of (A) male and (B) female monkeys. Slice -3 = lumbar, slices 3, 6 = thoracic, and slice 9 = cervical; 0 = catheter tip. Figure 148C/D depicts an exemplary result illustrating HNS activity in the brains of (C) male and (D) female monkeys. Slices are numbered rostral to caudal (3 to 15). All tissue samples were collected approximately 24 hours after the last dose or 4 weeks after the last dose for the recovery animals. DC, device control. The data represent mean \pm SEM for n = 4 monkeys per treatment group.

[0199] Figure 149 depicts an exemplary result illustrating enzyme activity in monkey brain and liver. Figure 149A depicts an exemplary result illustrating HNS activity distribution in the high-dose (8.3 mg/dose) group monkey brain. The fold-change in activity for surface, deep, and very deep (periventricular) areas of the brain compared with endogenous levels (DC group) is shown. All tissue samples were collected approximately 24 hours after the last dose or 4 weeks after the last dose for the recovery animals. The data represent mean \pm SEM for n = 6 monkeys (both sexes), brain slices 6 and 9. Data for two monkeys were not included; at necropsy the catheters were not found to be patent. Figure 149B shows HNS activity in monkey liver. All tissue samples were collected approximately 24 hours after the last dose or 4 weeks after the last dose for the recovery animals. DC, device control. Rec, recovery. The data represent mean \pm SEM for n = 4 monkeys per treatment group except for the low-dose (4.5 mg/dose) female group (n = 3).

2017202349 10 Apr 2017

[0200] Figure 150 depicts an exemplary result illustrating HNS localization in juvenile cynomolgus monkey cerebellum: 3-month interim cohort. Figure 150A depicts an exemplary result illustrating cerebellum of a vehicle control animal (0 mg/dose) negative for HNS immunostaining; 20x magnification. Figure 150B depicts an exemplary result illustrating cerebellum of a low-dose (1.5 mg/dose) animal showing minimal positive staining limited to the molecular layer; 20x magnification. Figure 150C depicts an exemplary result illustrating cerebellum of a mid-dose (4.5 mg/dose) animal showing minimal staining in the outer granular layer; 20x magnification. Figure 150D depicts an exemplary result illustrating moderate staining in the cerebellum of a high-dose (8.3 mg/dose) animal including molecular, outer granular layer, and Purkinje cells; 20x magnification.

[0201] Figure 151 depicts an exemplary study of the concentration of HNS in the head region plotted with time in the first 20 minutes after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

[0202] Figure 152 depicts an exemplary study of the concentration of HNS in the brain plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

[0203] Figure 153 depicts an exemplary study of the concentration of HNS in the brain region plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

[0204] Figure 154 depicts an exemplary study of the concentration of HNS in the head region plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

[0205] Figure 155 depicts an exemplary study of the concentration of HNS in the proximal spine plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

[0206] Figure 156 depicts an exemplary study of the concentration of HNS in the mid-spine plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

[0207] Figure 157 depicts an exemplary study of the concentration of HNS in the distal spine plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

[0208] Figure 158 depicts an exemplary study of the concentration of HNS in the liver plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg.

2017202349 10 Apr 2017

[0209] Figure 159 depicts an exemplary study of the concentration of HNS in the brain plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg, individual (top) and mean \pm SD (bottom).

[0210] Figure 160 depicts an exemplary study of the hepatic concentration of HNS plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg, individual (top) and mean \pm SD (bottom).

[0211] Figure 161 depicts an exemplary study of the renal concentration of HNS plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg, individual (top) and mean \pm SD (bottom).

[0212] Figure 162 depicts an exemplary study of the heart concentration of HNS plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg, individual (top) and mean \pm SD (bottom).

[0213] Figure 163 depicts an exemplary study of the skin concentration of HNS plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg, individual (top) and mean \pm SD (bottom).

[0214] Figure 164 depicts an exemplary study of the brain concentration of HNS plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg (top), and a comparison of the non-compartmental PK parameters in the brain (bottom).

[0215] Figure 165 depicts an exemplary study of the liver concentration of HNS plotted with time after IT dosing of ^{124}I -HNS at 1 and 10 mg/kg (top), and a comparison of the non-compartmental PK parameters in the liver (bottom).

[0216] Figure 166 illustrates exemplary primary fibroblast cells from normal human were used for cellular internalization study of rhNaglu and Naglu-IGFII. Cellular uptake of rhNaglu was minimum, while the cellular uptake of Naglu-IGFII was much pronounced. The saturating curve of Naglu-IGFII internalization indicated a receptor mediated uptake. This uptake was inhibited by IGFII, but not by mannose-6-phosphate.

2017202349 10 Apr 2017

[0217] Figure 167 depicts results of an exemplary confocal microscopy study using a Sanfilippo B subject's fibroblast cells (GM01426). Extensive internalization of Naglu-IGFII, and co-localization of Naglu-IGFII with Lamp-1 was observed.

[0218] Figure 168 is an exemplary illustration showing Naglu activity in wild type (WT), Naglu-/- (KO) and heterozygote Naglu+/- (Het) mouse. Total deficiency of Naglu in Sanfilippo B mouse was observed in brain, liver, kidney and spleen.

[0219] Figure 169 depicts superior and lateral view of the mouse brain to indicate the site of intracisternal injection (IC) and the sectioning plane for histology analyses. Middle micrograph, a transversal section of mouse brain viewed at 1x magnitude. Boxed area indicates the field for 4x microscopy image in the bottom micrograph. Bottom micrograph, 4x image of histology slide. Box A and B indicates the field of 40x microscopy image in Figure 170 and Figure 171.

[0220] Figure 170 depicts exemplary immunohistochemistry of the cerebral cortex in Sanfilippo B mice 7 days after intracisternal injection (IC) 40x. Both rhNaglu and Naglu-IGFII exhibited extensive cellular uptake in neurons as well as in glial cells, and the distribution and cellular uptake patterns were very similar between the two proteins. (anti-human Naglu monoclonal antibody).

[0221] Figure 171 depicts exemplary LAMP-1 immunostaining of the cerebral cortex at 40x magnification. Comparing to the brain of wild type mouse, increased lysosomal storage was observed in the brain of vehicle treated Sanfilippo B mouse, as demonstrated by the increased LAMP-1 immunostaining positive spots. The brain of both rhNaglu and Naglu-IGFII treated Sanfilippo B mouse exhibited reduction of lysosomal storage that was very similar to wt mouse.

[0222] Figure 172A illustrates widespread reduction of cellular vacuolation in the white matter tissues of Naglu-deficient mice IT-administered Naglu relative to the same Naglu-deficient mice that were administered the vehicle. Figure 172B illustrates a marked reduction in lysosomal associated membrane protein 1 (LAMP1) immunostaining in the white matter tissues of Naglu-deficient mice intrathecally-administered Naglu relative to the same Naglu-deficient mice that were administered a vehicle.

2017202349 10 Apr 2017

[0223] Figure 173 quantitatively illustrates and compares the concentration of LAMP measured in the cerebral cortex, caudate nucleus and putamen (CP), thalamus (TH), cerebellum (CBL) and white matter (WM) of the Naglu-deficient mice which were administered Naglu relative to both the wild-type and Naglu-deficient mice that were administered a vehicle. The LAMP-positive areas in each area of brain tissue analyzed were further reduced following the intrathecal administration of three doses of Naglu over the course of seven days (Figure 173A) relative to two doses of Naglu over the course of two weeks (Figure 173B).

[0224] Figure 174 illustrates an exemplary midsagittal anatomical diagram of human CNS is used as a reference in this figure, to demonstrate the site of IT injection in wt cannulated Rat. Arrows indicate the approximate anatomic location of IT injection in the spinal cord the cerebral cortex region where tissues were taken for immunohistochemistry study.

[0225] Figure 175 illustrates exemplary Naglu activity in the brain after IT injection. Naglu activity was significantly higher in the brain of Naglu-TAT and Naglu-IGFII injected wt rat.

[0226] Figure 176 depicts exemplary Naglu immunostaining of the cerebral cortex of rhNaglu, Naglu-TAT, Naglu-IGFII, Naglu-kif and PerT-Naglu treated wt cannulated rat 24hr after IT injection 20x. Naglu-IGFII was the only protein exhibited extensive distribution well into the parenchyma of the brain. Cellular uptake into neurons and glial cells was also evident in Naglu-IGFII treated rat. On the other hand, in rhNaglu, Naglu-TAT, Naglu kif and PerT-Naglu treated groups, the protein remained in the meninges (M).

[0227] Figure 177 depicts exemplary high power magnification of the selected slides from Figure 176. Upper panel, in the rhNaglu treated wt cannulated rat, rhNaglu remained at the meninges (M), no positive staining found in the parenchyma of the brain. Lower panel, in Naglu-IGFII treated wt cannulated rat, extensive distribution was observed well into the parenchyma of the brain, and cellular uptake was observed in neurons and glial cells.

[0228] Figure 178 illustrates exemplary Naglu activity in brain and liver 24hr after last IT injection. Among the three treated groups, Naglu activity in the brain did not show significant differences, the same was true for the Naglu activity in the liver. This result indicated that the

2017202349 10 Apr 2017

Naglu activity detected in the brain and liver was largely due to the last injection, which occurred 24hr prior to sacrifice.

[0229] Figure 179 illustrates exemplary total GAG level in the brain and liver after IT injection of Naglu-IGFII. Total GAG in the brain of vehicle treated Sanfilippo B mice exhibited progressive increases, a reflection of accumulative effect as the Sanfilippo B mice aging. A statistically significant reduction of GAG in the brain was observed in 3x injection group ($p < 0.05$). Statistically significant reductions of GAG in liver were also observed in 2x and 3x injection groups ($p < 0.05$). The quicker and more drastic change of GAG level in liver than in the brain is a phenomenon that also has been observed in IT delivery of I2S for Hunter Syndrome.

[0230] Figure 180 depicts exemplary biodistribution of Naglu in the brain of Sanfilippo B mice after IT injection. Naglu immunofluorescent staining revealed the Naglu-IGFII protein on the meninges (M) and parenchyma of the brain. Cellular uptake was observed in the 2x and 3x injection groups. G: glial cells.

[0231] Figure 181 is an exemplary illustration showing a coronal section of the mouse brain. Boxes indicate where the pictures for LAMP-1 immunostaining were taken. To demonstrate the extent of protein distribution and efficacy, cerebral cortex and subcortical tissues such as caudate nucleus, thalamus and white matter were selected for LAMP-1 immunostaining.

[0232] Figure 182 is an exemplary illustration showing the LAMP-1 immunostaining of cerebral cortex at 40x magnification. Comparing to the brain of wild type mouse, increased lysosomal storage was observed in the brain of vehicle treated Sanfilippo B mouse, as seen by the increased LAMP-1 immunostaining positive spots. Reduction of lysosomal storage after Naglu-IGFII IT injection was evident by the reduced size of positive spots of 2x injection treated Sanfilippo B mouse brain, and the reduced size and number of positive spots of the 3x injection treated Sanfilippo B mouse brain.

[0233] Figure 183 is an exemplary illustration showing LAMP-1 immunostaining of the caudate nucleus, a subcortical nucleus (40x). Similar to what was seen in the cerebral cortex, increased lysosomal storage was observed in the brain of vehicle treated Sanfilippo B mouse, as seen by the increased LAMP-1 immunostaining positive spots. Reduction of lysosomal storage

2017202349 10 Apr 2017

after Naglu-IGFII IT injection was evident by the reduced size of positive spots of 2x injection treated Sanfilippo B mouse brain and by the reduced size and number of positive spots of the 3x injection treated Sanfilippo B mouse brain.

[0234] Figure 184 is an exemplary illustration showing LAMP-1 immunostaining of the thalamus, a diencephalic nuclei (40x). Reduction of lysosomal storage after Naglu-IGFII IT injection was evident by the reduced size of positive spots of 2x injection treated Sanfilippo B mouse brain and by the reduced size and number of positive spots of the 3x injection treated Sanfilippo B mouse brain.

[0235] Figure 185 is an exemplary illustration showing LAMP-1 immunostaining of white matter (40x). The longitudinal track of neuron axon fibers distinguishes the white matter from grey matters presented in Figures 181-184. Nonetheless, the same pattern of increases of lysosomal storage could be seen in vehicle treated Sanfilippo B mouse's brain when compared to the wild type mouse. Reduction of lysosomal storage after Naglu-IGFII IT injection was evident by the reduced size and reduced number of positive spots in the 2x and 3x injection treated Sanfilippo B mouse brain.

[0236] Figure 186 is an exemplary illustration showing LAMP-1 immunostaining of the cerebellar cortex. The morphology of cerebellar cortex was evident by the densely populated granular neurons, the hypocellular Molecular layer, and the single layer of Purkinje neurons between the granular neurons and the molecular layer. Purkinje neurons were identified by the large cytoplasm and occasional dendrites protruding into the Molecular layer.

[0237] Figure 187 is an exemplary illustration showing Naglu staining in the brain, spinal cord and liver. In the brain and spinal cord, injected Naglu was detected in meninges (M) only by IHC and no Naglu positive staining was detected in any other regions. In the liver, sinusoidal cells (S) were Naglu positive and no Naglu uptake was found in hepatocytes (H).

[0238] Figure 188 is an exemplary illustration showing LAMP immunostaining and H & E staining of the liver and spinal cord. Compared with the vehicle animals, LAMP staining was decreased throughout in both livers and spinal cords treated with Naglu. H & E staining showed

2017202349 10 Apr 2017

that cellular vacuolation in hepatocytes was reduced in the treated group compared with vehicle treated animals.

[0239] Figure 189 A and Figure 189 B is an exemplary illustration showing H & E staining of the brain and morphology improvement of the brain after 6 every other week (EOW) IT injections of Naglu for 3 months. In the treated brain, the cellular vacuolation (arrows) in all examined regions decreased compared with the vehicle group.

[0240] Figure 190 A and Figure 190 B are exemplary illustrations showing LAMP immunostaining in various brain regions after 6 IT Naglu injections for 3 months. Compared with the vehicle treated group, Naglu IT administration to Sanfilippo B mice resulted in a reduction of lysosomal activity in all examined regions revealed by LAMP immunostaining. This reduction was characterized by the decrease in the number of LAMP positive cells, smaller cell size and lighter staining. A marked reduction was found in the cerebellum and brainstem, which are located in the caudate part of the brain close to the spinal cord, compared with other brain regions. A clear reduction was also found in the deep brain regions, including the white matter, hippocampus, and thalamus.

[0241] Figure 191 A and Figure 190 B are exemplary illustrations showing Iba IHC in various brain regions after 6 IT Naglu injections for 3 months, which revealed activation of microglial cells. Compared with vehicle treated group, no decrease in the number of positive cells and staining intensity was observed in the Naglu treated group. However, the cellular morphology of positive microglial cells changed with reduced cell size in all examined brain regions compared to large and vacuolated ones in the vehicle group (inserts).

[0242] Figure 192 A and Figure 192 B are exemplary illustrations showing GFAP IHC in various brain regions after 6 IT Naglu injections for 3 months, which revealed astrocytic activation. Compared with the vehicle treated group, GFAP positive staining was decreased in the cerebellum and brainstem, and slightly decreased in other examined regions.

2017202349 10 Apr 2017

DEFINITIONS

[0243] In order for the present invention to be more readily understood, certain terms are first defined below. Additional definitions for the following terms and other terms are set forth throughout the specification.

[0244] *Approximately or about:* As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[0245] *Amelioration:* As used herein, the term “amelioration” is meant the prevention, reduction or palliation of a state, or improvement of the state of a subject. Amelioration includes, but does not require complete recovery or complete prevention of a disease condition. In some embodiments, amelioration includes increasing levels of relevant protein or its activity that is deficient in relevant disease tissues.

[0246] *Biologically active:* As used herein, the phrase “biologically active” refers to a characteristic of any agent that has activity in a biological system, and particularly in an organism. For instance, an agent that, when administered to an organism, has a biological effect on that organism, is considered to be biologically active. In particular embodiments, where a protein or polypeptide is biologically active, a portion of that protein or polypeptide that shares at least one biological activity of the protein or polypeptide is typically referred to as a “biologically active” portion.

[0247] *Bulking agent:* As used herein, the term “bulking agent” refers to a compound which adds mass to the lyophilized mixture and contributes to the physical structure of the lyophilized cake (e.g., facilitates the production of an essentially uniform lyophilized cake which maintains an open pore structure). Exemplary bulking agents include mannitol, glycine, sodium chloride, hydroxyethyl starch, lactose, sucrose, trehalose, polyethylene glycol and dextran.

2017202349 10 Apr 2017

[0248] *Cation-independent mannose-6-phosphate receptor (CI-MPR)*: As used herein, the term “cation-independent mannose-6-phosphate receptor (CI-MPR)” refers to a cellular receptor that binds mannose-6-phosphate (M6P) tags on acid hydrolase precursors in the Golgi apparatus that are destined for transport to the lysosome. In addition to mannose-6-phosphates, the CI-MPR also binds other proteins including IGF-II. The CI-MPR is also known as “M6P/IGF-II receptor,” “CI-MPR/IGF-II receptor,” “IGF-II receptor” or “IGF2 Receptor.” These terms and abbreviations thereof are used interchangeably herein.

[0249] *Concurrent immunosuppressant therapy*: As used herein, the term “concurrent immunosuppressant therapy” includes any immunosuppressant therapy used as pre-treatment, preconditioning or in parallel to a treatment method.

[0250] *Diluent*: As used herein, the term “diluent” refers to a pharmaceutically acceptable (e.g., safe and non-toxic for administration to a human) diluting substance useful for the preparation of a reconstituted formulation. Exemplary diluents include sterile water, bacteriostatic water for injection (BWHI), a pH buffered solution (e.g. phosphate-buffered saline), sterile saline solution, Ringer's solution or dextrose solution.

[0251] *Dosage form*: As used herein, the terms “dosage form” and “unit dosage form” refer to a physically discrete unit of a therapeutic protein for the patient to be treated. Each unit contains a predetermined quantity of active material calculated to produce the desired therapeutic effect. It will be understood, however, that the total dosage of the composition will be decided by the attending physician within the scope of sound medical judgment.

[0252] *Enzyme replacement therapy (ERT)*: As used herein, the term “enzyme replacement therapy (ERT)” refers to any therapeutic strategy that corrects an enzyme deficiency by providing the missing enzyme. In some embodiments, the missing enzyme is provided by intrathecal administration. In some embodiments, the missing enzyme is provided by infusing into bloodstream. Once administered, enzyme is taken up by cells and transported to the lysosome, where the enzyme acts to eliminate material that has accumulated in the lysosomes due to the enzyme deficiency. Typically, for lysosomal enzyme replacement therapy to be effective, the therapeutic enzyme is delivered to lysosomes in the appropriate cells in target tissues where the storage defect is manifest.

2017202349 10 Apr 2017

[0253] *Improve, increase, or reduce:* As used herein, the terms “improve,” “increase” or “reduce,” or grammatical equivalents, indicate values that are relative to a baseline measurement, such as a measurement in the same individual prior to initiation of the treatment described herein, or a measurement in a control individual (or multiple control individuals) in the absence of the treatment described herein. A “control individual” is an individual afflicted with the same form of lysosomal storage disease as the individual being treated, who is about the same age as the individual being treated (to ensure that the stages of the disease in the treated individual and the control individual(s) are comparable).

[0254] *Individual, subject, patient:* As used herein, the terms “subject,” “individual” or “patient” refer to a human or a non-human mammalian subject. The individual (also referred to as “patient” or “subject”) being treated is an individual (fetus, infant, child, adolescent, or adult human) suffering from a disease.

[0255] *Intrathecal administration:* As used herein, the term “intrathecal administration” or “intrathecal injection” refers to an injection into the spinal canal (intrathecal space surrounding the spinal cord). Various techniques may be used including, without limitation, lateral cerebroventricular injection through a burrhole or cisternal or lumbar puncture or the like. In some embodiments, “intrathecal administration” or “intrathecal delivery” according to the present invention refers to IT administration or delivery via the lumbar area or region, i.e., lumbar IT administration or delivery. As used herein, the term “lumbar region” or “lumbar area” refers to the area between the third and fourth lumbar (lower back) vertebrae and, more inclusively, the L2-S1 region of the spine.

[0256] *Linker:* As used herein, the term “linker” refers to, in a fusion protein, an amino acid sequence other than that appearing at a particular position in the natural protein and is generally designed to be flexible or to interpose a structure, such as an α -helix, between two protein moieties. A linker is also referred to as a spacer.

[0257] *Lyoprotectant:* As used herein, the term “lyoprotectant” refers to a molecule that prevents or reduces chemical and/or physical instability of a protein or other substance upon lyophilization and subsequent storage. Exemplary lyoprotectants include sugars such as sucrose or trehalose; an amino acid such as monosodium glutamate or histidine; a methylamine such as

2017202349 10 Apr 2017

betaine; a lyotropic salt such as magnesium sulfate; a polyol such as trihydric or higher sugar alcohols, e.g. glycerin, erythritol, glycerol, arabitol, xylitol, sorbitol, and mannitol; propylene glycol; polyethylene glycol; Pluronic; and combinations thereof. In some embodiments, a lyoprotectant is a non-reducing sugar, such as trehalose or sucrose.

[0258] *Lysosomal enzyme:* As used herein, the term “lysosomal enzyme” refers to any enzyme that is capable of reducing accumulated materials in mammalian lysosomes or that can rescue or ameliorate one or more lysosomal storage disease symptoms. Lysosomal enzymes suitable for the invention include both wild-type or modified lysosomal enzymes and can be produced using recombinant and synthetic methods or purified from nature sources. Exemplary lysosomal enzymes are listed in Table 1.

[0259] *Lysosomal enzyme deficiency:* As used herein, “lysosomal enzyme deficiency” refers to a group of genetic disorders that result from deficiency in at least one of the enzymes that are required to break macromolecules (e.g., enzyme substrates) down to peptides, amino acids, monosaccharides, nucleic acids and fatty acids in lysosomes. As a result, individuals suffering from lysosomal enzyme deficiencies have accumulated materials in various tissues (e.g., CNS, liver, spleen, gut, blood vessel walls and other organs).

[0260] *Lysosomal Storage Disease:* As used herein, the term “lysosomal storage disease” refers to any disease resulting from the deficiency of one or more lysosomal enzymes necessary for metabolizing natural macromolecules. These diseases typically result in the accumulation of un-degraded molecules in the lysosomes, resulting in increased numbers of storage granules (also termed storage vesicles). These diseases and various examples are described in more detail below.

[0261] *Polypeptide:* As used herein, a “polypeptide”, generally speaking, is a string of at least two amino acids attached to one another by a peptide bond. In some embodiments, a polypeptide may include at least 3-5 amino acids, each of which is attached to others by way of at least one peptide bond. Those of ordinary skill in the art will appreciate that polypeptides sometimes include “non-natural” amino acids or other entities that nonetheless are capable of integrating into a polypeptide chain, optionally.

2017202349 10 Apr 2017

[0262] *Replacement enzyme:* As used herein, the term “replacement enzyme” refers to any enzyme that can act to replace at least in part the deficient or missing enzyme in a disease to be treated. In some embodiments, the term “replacement enzyme” refers to any enzyme that can act to replace at least in part the deficient or missing lysosomal enzyme in a lysosomal storage disease to be treated. In some embodiments, a replacement enzyme is capable of reducing accumulated materials in mammalian lysosomes or that can rescue or ameliorate one or more lysosomal storage disease symptoms. Replacement enzymes suitable for the invention include both wild-type or modified lysosomal enzymes and can be produced using recombinant and synthetic methods or purified from nature sources. A replacement enzyme can be a recombinant, synthetic, gene-activated or natural enzyme.

[0263] *Soluble:* As used herein, the term “soluble” refers to the ability of a therapeutic agent to form a homogenous solution. In some embodiments, the solubility of the therapeutic agent in the solution into which it is administered and by which it is transported to the target site of action (e.g., the cells and tissues of the brain) is sufficient to permit the delivery of a therapeutically effective amount of the therapeutic agent to the targeted site of action. Several factors can impact the solubility of the therapeutic agents. For example, relevant factors which may impact protein solubility include ionic strength, amino acid sequence and the presence of other co-solubilizing agents or salts (e.g., calcium salts). In some embodiments, the pharmaceutical compositions are formulated such that calcium salts are excluded from such compositions. In some embodiments, therapeutic agents in accordance with the present invention are soluble in its corresponding pharmaceutical composition. It will be appreciated that, while isotonic solutions are generally preferred for parenterally administered drugs, the use of isotonic solutions may limit adequate solubility for some therapeutic agents and, in particular some proteins and/or enzymes. Slightly hypertonic solutions (e.g., up to 175mM sodium chloride in 5mM sodium phosphate at pH 7.0) and sugar-containing solutions (e.g., up to 2% sucrose in 5mM sodium phosphate at pH 7.0) have been demonstrated to be well tolerated in monkeys. For example, the most common approved CNS bolus formulation composition is saline (150mM NaCl in water).

[0264] *Stability:* As used herein, the term “stable” refers to the ability of the therapeutic agent (e.g., a recombinant enzyme) to maintain its therapeutic efficacy (e.g., all or the majority

2017202349 10 Apr 2017

of its intended biological activity and/or physiochemical integrity) over extended periods of time. The stability of a therapeutic agent, and the capability of the pharmaceutical composition to maintain stability of such therapeutic agent, may be assessed over extended periods of time (e.g., for at least 1, 3, 6, 12, 18, 24, 30, 36 months or more). In general, pharmaceutical compositions described herein have been formulated such that they are capable of stabilizing, or alternatively slowing or preventing the degradation, of one or more therapeutic agents formulated therewith (e.g., recombinant proteins). In the context of a formulation a stable formulation is one in which the therapeutic agent therein essentially retains its physical and/or chemical integrity and biological activity upon storage and during processes (such as freeze/thaw, mechanical mixing and lyophilization). For protein stability, it can be measure by formation of high molecular weight (HMW) aggregates, loss of enzyme activity, generation of peptide fragments and shift of charge profiles.

[0265] *Subject:* As used herein, the term “subject” means any mammal, including humans. In certain embodiments of the present invention the subject is an adult, an adolescent or an infant. Also contemplated by the present invention are the administration of the pharmaceutical compositions and/or performance of the methods of treatment in-utero.

[0266] *Substantial homology:* The phrase “substantial homology” is used herein to refer to a comparison between amino acid or nucleic acid sequences. As will be appreciated by those of ordinary skill in the art, two sequences are generally considered to be “substantially homologous” if they contain homologous residues in corresponding positions. Homologous residues may be identical residues. Alternatively, homologous residues may be non-identical residues will appropriately similar structural and/or functional characteristics. For example, as is well known by those of ordinary skill in the art, certain amino acids are typically classified as “hydrophobic” or “hydrophilic” amino acids., and/or as having “polar” or “non-polar” side chains. Substitution of one amino acid for another of the same type may often be considered a “homologous” substitution.

[0267] As is well known in this art, amino acid or nucleic acid sequences may be compared using any of a variety of algorithms, including those available in commercial computer programs such as BLASTN for nucleotide sequences and BLASTP, gapped BLAST, and PSI-

2017202349 10 Apr 2017

BLAST for amino acid sequences. Exemplary such programs are described in Altschul, et al., Basic local alignment search tool, *J. Mol. Biol.*, 215(3): 403-410, 1990; Altschul, et al., *Methods in Enzymology*; Altschul, et al., "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", *Nucleic Acids Res.* 25:3389-3402, 1997; Baxevanis, et al., *Bioinformatics : A Practical Guide to the Analysis of Genes and Proteins*, Wiley, 1998; and Misener, et al., (eds.), *Bioinformatics Methods and Protocols* (Methods in Molecular Biology, Vol. 132), Humana Press, 1999. In addition to identifying homologous sequences, the programs mentioned above typically provide an indication of the degree of homology. In some embodiments, two sequences are considered to be substantially homologous if at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more of their corresponding residues are homologous over a relevant stretch of residues. In some embodiments, the relevant stretch is a complete sequence. In some embodiments, the relevant stretch is at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500 or more residues.

[0268] *Substantial identity:* The phrase "substantial identity" is used herein to refer to a comparison between amino acid or nucleic acid sequences. As will be appreciated by those of ordinary skill in the art, two sequences are generally considered to be "substantially identical" if they contain identical residues in corresponding positions. As is well known in this art, amino acid or nucleic acid sequences may be compared using any of a variety of algorithms, including those available in commercial computer programs such as BLASTN for nucleotide sequences and BLASTP, gapped BLAST, and PSI-BLAST for amino acid sequences. Exemplary such programs are described in Altschul, et al., Basic local alignment search tool, *J. Mol. Biol.*, 215(3): 403-410, 1990; Altschul, et al., *Methods in Enzymology*; Altschul et al., *Nucleic Acids Res.* 25:3389-3402, 1997; Baxevanis et al., *Bioinformatics : A Practical Guide to the Analysis of Genes and Proteins*, Wiley, 1998; and Misener, et al., (eds.), *Bioinformatics Methods and Protocols* (Methods in Molecular Biology, Vol. 132), Humana Press, 1999. In addition to identifying identical sequences, the programs mentioned above typically provide an indication of the degree of identity. In some embodiments, two sequences are considered to be substantially identical if at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%,

2017202349 10 Apr 2017

95%, 96%, 97%, 98%, 99% or more of their corresponding residues are identical over a relevant stretch of residues. In some embodiments, the relevant stretch is a complete sequence. In some embodiments, the relevant stretch is at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500 or more residues.

[0269] *Synthetic CSF*: As used herein, the term “synthetic CSF” refers to a solution that has pH, electrolyte composition, glucose content and osmolarity consistent with the cerebrospinal fluid. Synthetic CSF is also referred to as artificial CSF. In some embodiments, synthetic CSF is an Elliott’s B solution.

[0270] *Suitable for CNS delivery*: As used herein, the phrase “suitable for CNS delivery” or “suitable for intrathecal delivery” as it relates to the pharmaceutical compositions of the present invention generally refers to the stability, tolerability, and solubility properties of such compositions, as well as the ability of such compositions to deliver an effective amount of the therapeutic agent contained therein to the targeted site of delivery (e.g., the CSF or the brain).

[0271] *Target tissues*: As used herein, the term “target tissues” refers to any tissue that is affected by the lysosomal storage disease to be treated or any tissue in which the deficient lysosomal enzyme is normally expressed. In some embodiments, target tissues include those tissues in which there is a detectable or abnormally high amount of enzyme substrate, for example stored in the cellular lysosomes of the tissue, in patients suffering from or susceptible to the lysosomal storage disease. In some embodiments, target tissues include those tissues that display disease-associated pathology, symptom, or feature. In some embodiments, target tissues include those tissues in which the deficient lysosomal enzyme is normally expressed at an elevated level. As used herein, a target tissue may be a brain target tissue, a spinal cord target tissue and/or a peripheral target tissue. Exemplary target tissues are described in detail below.

[0272] *Therapeutic moiety*: As used herein, the term “therapeutic moiety” refers to a portion of a molecule that renders the therapeutic effect of the molecule. In some embodiments, a therapeutic moiety is a polypeptide having therapeutic activity.

2017202349 10 Apr 2017

[0273] *Therapeutically effective amount:* As used herein, the term “therapeutically effective amount” refers to an amount of a therapeutic protein (e.g., replacement enzyme) which confers a therapeutic effect on the treated subject, at a reasonable benefit/risk ratio applicable to any medical treatment. The therapeutic effect may be objective (i.e., measurable by some test or marker) or subjective (i.e., subject gives an indication of or feels an effect). In particular, the “therapeutically effective amount” refers to an amount of a therapeutic protein or composition effective to treat, ameliorate, or prevent a desired disease or condition, or to exhibit a detectable therapeutic or preventative effect, such as by ameliorating symptoms associated with the disease, preventing or delaying the onset of the disease, and/or also lessening the severity or frequency of symptoms of the disease. A therapeutically effective amount is commonly administered in a dosing regimen that may comprise multiple unit doses. For any particular therapeutic protein, a therapeutically effective amount (and/or an appropriate unit dose within an effective dosing regimen) may vary, for example, depending on route of administration, on combination with other pharmaceutical agents. Also, the specific therapeutically effective amount (and/or unit dose) for any particular patient may depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific pharmaceutical agent employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and/or rate of excretion or metabolism of the specific fusion protein employed; the duration of the treatment; and like factors as is well known in the medical arts.

[0274] *Tolerable:* As used herein, the terms “tolerable” and “tolerability” refer to the ability of the pharmaceutical compositions of the present invention to not elicit an adverse reaction in the subject to whom such composition is administered, or alternatively not to elicit a serious adverse reaction in the subject to whom such composition is administered. In some embodiments, the pharmaceutical compositions of the present invention are well tolerated by the subject to whom such compositions is administered.

[0275] *Treatment:* As used herein, the term “treatment” (also “treat” or “treating”) refers to any administration of a therapeutic protein (e.g., lysosomal enzyme) that partially or completely alleviates, ameliorates, relieves, inhibits, delays onset of, reduces severity of and/or reduces incidence of one or more symptoms or features of a particular disease, disorder, and/or

2017202349 10 Apr 2017

condition (e.g., Hunters syndrome, Sanfilippo syndrome type B). Such treatment may be of a subject who does not exhibit signs of the relevant disease, disorder and/or condition and/or of a subject who exhibits only early signs of the disease, disorder, and/or condition. Alternatively or additionally, such treatment may be of a subject who exhibits one or more established signs of the relevant disease, disorder and/or condition.

DETAILED DESCRIPTION

[0276] The present invention provides, among other things, improved methods for effective direct delivery of a therapeutic agent to the central nervous system (CNS). As discussed above, the present invention is based on unexpected discovery that a replacement enzyme for a lysosomal storage disease can be directly introduced into the cerebrospinal fluid (CSF) of a subject in need of treatment at a high concentration without inducing substantial adverse effects in the subject. More surprisingly, the present inventors found that the replacement enzyme may be delivered in a simple saline or buffer-based formulation, without using synthetic CSF. Even more unexpectedly, intrathecal delivery according to the present invention does not result in substantial adverse effects, such as severe immune response, in the subject. Therefore, in some embodiments, intrathecal delivery according to the present invention may be used in absence of concurrent immunosuppressant therapy (e.g. without induction of immune tolerance by pre-treatment or pre-conditioning).

[0277] In some embodiments, intrathecal delivery according to the present invention permits efficient diffusion across various brain tissues resulting in effective delivery of the replacement enzyme in various target brain tissues in surface, shallow and/or deep brain regions. In some embodiments, intrathecal delivery according to the present invention resulted in sufficient amount of replacement enzymes entering the peripheral circulation. As a result, in some cases, intrathecal delivery according to the present invention resulted in delivery of the replacement enzyme in peripheral tissues, such as liver, heart, and kidney. This discovery is unexpected and can be particularly useful for the treatment of lysosomal storage diseases that have both CNS and peripheral components, which would typically require both regular intrathecal administration and intravenous administration. It is contemplated that intrathecal delivery

2017202349 10 Apr 2017

according to the present invention may allow reduced dosing and/or frequency of *iv* injection without compromising therapeutic effects in treating peripheral symptoms.

[0278] The present invention provides various unexpected and beneficial features that allow efficient and convenient delivery of replacement enzymes to various brain target tissues, resulting in effective treatment of lysosomal storage diseases that have CNS indications.

[0279] Various aspects of the invention are described in detail in the following sections. The use of sections is not meant to limit the invention. Each section can apply to any aspect of the invention. In this application, the use of “or” means “and/or” unless stated otherwise.

Lysosomal Storage Diseases and Replacement Enzymes

[0280] Inventive methods according to the present may be used to treat any lysosomal storage diseases, in particular those lysosomal storage diseases having CNS etiology and/or symptoms, including, but are not limited to, aspartylglucosaminuria, cholesterol ester storage disease, Wolman disease, cystinosis, Danon disease, Fabry disease, Farber lipogranulomatosis, Farber disease, fucosidosis, galactosialidosis types I/II, Gaucher disease types I/II/III, globoid cell leukodystrophy, Krabbe disease, glycogen storage disease II, Pompe disease, GM1-gangliosidosis types I/II/III, GM2-gangliosidosis type I, Tay Sachs disease, GM2-gangliosidosis type II, Sandhoff disease, GM2-gangliosidosis, α -mannosidosis types I/II, β -mannosidosis, metachromatic leukodystrophy, mucopolipidosis type I, sialidosis types I/II, mucopolipidosis types II/III, I-cell disease, mucopolipidosis type IIIC pseudo-Hurler polydystrophy, mucopolysaccharidosis type I, mucopolysaccharidosis type II, Hunter syndrome, mucopolysaccharidosis type IIIA, Sanfilippo syndrome (type A, B, C or D), mucopolysaccharidosis type IIIB, mucopolysaccharidosis type IIIC, mucopolysaccharidosis type IIID, mucopolysaccharidosis type IVA, Morquio syndrome, mucopolysaccharidosis type IVB, mucopolysaccharidosis type VI, mucopolysaccharidosis type VII, Sly syndrome, mucopolysaccharidosis type IX, multiple sulfatase deficiency, neuronal ceroid lipofuscinosis, CLN1 Batten disease, CLN2 Batten disease, Niemann-Pick disease types A/B, Niemann-Pick disease type C1, Niemann-Pick disease type

2017202349 10 Apr 2017

C2, pycnodysostosis, Schindler disease types I/II, Gaucher disease and sialic acid storage disease.

[0281] In some embodiments, lysosomal storage diseases to be treated using inventive methods of the present invention include Hunters Syndrome, metachromatic leukodystrophy (MLD) disease, Sanfilippo syndrome type A, Sanfilippo syndrome type B, and globoid cell leukodystrophy (GLD) disease.

[0282] A detailed review of the genetic etiology, clinical manifestations, and molecular biology of the lysosomal storage diseases are detailed in Scriver et al., eds., *The Metabolic and Molecular Basis of Inherited Disease*, 7th Ed., Vol. II, McGraw Hill, (1995). Thus, the enzymes deficient in the above diseases are known to those of skill in the art, some of these are exemplified in the Table below:

Table 1

Disease Name	Enzyme Deficiency	Substance Stored
Pompe Disease	Acid- α 1, 4-Glucosidase	Glycogen α 1-4 linked Oligosaccharides
GM1 Gangliosidosis	β -Galactosidase	GM ₁ Gangliosides
Tay-Sachs Disease	β -Hexosaminidase A	GM ₂ Ganglioside
GM2 Gangliosidosis: AB Variant	GM ₂ Activator Protein	GM ₂ Ganglioside
Sandhoff Disease	β -Hexosaminidase A&B	GM ₂ Ganglioside
Fabry Disease	α -Galactosidase A	Globosides
Gaucher Disease	Glucocerebrosidase	Glucosylceramide
Metachromatic Leukodystrophy	Arylsulfatase A	Sulphatides
Krabbe Disease	Galactosylceramidase	Galactocerebroside
Niemann Pick, Types A & B	Acid Sphingomyelinase	Sphingomyelin
Niemann-Pick, Type C	Cholesterol Esterification Defect	Sphingomyelin
Niemann-Pick, Type D	Unknown	Sphingomyelin
Farber Disease	Acid Ceramidase	Ceramide
Wolman Disease	Acid Lipase	Cholesteryl Esters
Hurler Syndrome	α -L-Iduronidase	Heparan &

2017202349 10 Apr 2017

(MPS IH)		Dermatan Sulfates
Scheie Syndrome (MPS IS)	α -L-Iduronidase	Heparan & Dermatan, Sulfates
Hurler-Scheie (MPS IH/S)	α -L-Iduronidase	Heparan & Dermatan Sulfates
Hunter Syndrome (MPS II)	Iduronate Sulfatase	Heparan & Dermatan Sulfates
Sanfilippo syndrome type A (MPS IIIA)	Heparan N-Sulfatase	Heparan Sulfate
Sanfilippo syndrome type B (MPS IIIB)	α -N-Acetylglucosaminidase	Heparan Sulfate
Sanfilippo syndrome type C (MPS IIIC)	Acetyl-CoA-Glucosaminide Acetyltransferase	Heparan Sulfate
Sanfilippo syndrome type D (MPS IIID)	N-Acetylglucosamine-6-Sulfatase	Heparan Sulfate
Morquio B (MPS IVB)	β -Galactosidase	Keratan Sulfate
Maroteaux-Lamy (MPS VI)	Arylsulfatase B	Dermatan Sulfate
Sly Syndrome (MPS VII)	β -Glucuronidase	
α -Mannosidosis	α -Mannosidase	Mannose/Oligosaccharides
β -Mannosidosis	β -Mannosidase	Mannose/Oligosaccharides
Fucosidosis	α -L-Fucosidase	Fucosyl Oligosaccharides
Aspartylglucosaminuria	N-Aspartyl- β -Glucosaminidase	Aspartylglucosamine Asparagines
Sialidosis (Mucopolipidosis I)	α -Neuraminidase	Sialyloligosaccharides
Galactosialidosis (Goldberg Syndrome)	Lysosomal Protective Protein Deficiency	Sialyloligosaccharides
Schindler Disease	α -N-Acetyl-Galactosaminidase	
Mucopolipidosis II (I-	N-Acetylglucosamine-1-Phosphotransferase	Heparan Sulfate

2017202349 10 Apr 2017

Cell Disease)		
Mucopolidosis III (Pseudo-Hurler Polydystrophy)	Same as ML II	
Cystinosis	Cystine Transport Protein	Free Cystine
Salla Disease	Sialic Acid Transport Protein	Free Sialic Acid and Glucuronic Acid
Infantile Sialic Acid Storage Disease	Sialic Acid Transport Protein	Free Sialic Acid and Glucuronic Acid
Infantile Neuronal Ceroid Lipofuscinosis	Palmitoyl-Protein Thioesterase	Lipofuscins
Mucopolidosis IV	Unknown	Gangliosides & Hyaluronic Acid
Prosaposin	Saposins A, B, C or D	

Replacement Enzymes

[0283] Inventive methods according to the present invention may be used to deliver any replacement enzymes. As used herein, replacement enzymes suitable for the present invention may include any enzyme that can act to replace at least partial activity of the deficient or missing lysosomal enzyme in a lysosomal storage disease to be treated. In some embodiments, a replacement enzyme is capable of reducing accumulated substance in lysosomes or that can rescue or ameliorate one or more lysosomal storage disease symptoms.

[0284] In some embodiments, a suitable replacement enzyme may be any lysosomal enzyme known to be associated with the lysosomal storage disease to be treated. In some embodiments, a suitable replacement enzyme is an enzyme selected from the enzyme listed in Table 1 above. In some embodiments, a replacement enzyme suitable for the present invention is iduronate-2-sulfatase (I2S), arylsulfatase A (ASA), heparan N-sulfatase (HNS), alpha-N-acetylglucosaminidase (Naglu) or β -galactosidase (GLC).

[0285] In some embodiments, a replacement enzyme suitable for the invention may have a wild-type or naturally occurring sequence. In some embodiments, a replacement enzyme suitable for the invention may have a modified sequence having substantial homology or identify

2017202349 10 Apr 2017

to the wild-type or naturally-occurring sequence (e.g., having at least 70%, 75%, 80%, 85%, 90%, 95%, 98% sequence identity to the wild-type or naturally-occurring sequence).

[0286] A replacement enzyme suitable for the present invention may be produced by any available means. For example, replacement enzymes may be recombinantly produced by utilizing a host cell system engineered to express a replacement enzyme-encoding nucleic acid. Alternatively or additionally, replacement enzymes may be produced by activating endogenous genes. Alternatively or additionally, replacement enzymes may be partially or fully prepared by chemical synthesis. Alternatively or additionally, replacement enzymes may also be purified from natural sources.

[0287] Where enzymes are recombinantly produced, any expression system can be used. To give but a few examples, known expression systems include, for example, egg, baculovirus, plant, yeast, or mammalian cells.

[0288] In some embodiments, enzymes suitable for the present invention are produced in mammalian cells. Non-limiting examples of mammalian cells that may be used in accordance with the present invention include BALB/c mouse myeloma line (NSO/I, ECACC No: 85110503); human retinoblasts (PER.C6, CruCell, Leiden, The Netherlands); monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol., 36:59,1977); human fibrosarcoma cell line (e.g., HT1080); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells +/-DHFR (CHO, Urlaub and Chasin, Proc. Natl. Acad. Sci. USA, 77:4216, 1980); mouse sertoli cells (TM4, Mather, Biol. Reprod., 23:243-251, 1980); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1 587); human cervical carcinoma cells (HeLa, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad. Sci., 383:44-68, 1982); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[0289] In some embodiments, inventive methods according to the present invention are used to deliver replacement enzymes produced from human cells. In some embodiments,

2017202349 10 Apr 2017

inventive methods according to the present invention are used to deliver replacement enzymes produced from CHO cells.

[0290] In some embodiments, replacement enzymes delivered using a method of the invention contains a moiety that binds to a receptor on the surface of brain cells to facilitate cellular uptake and/or lysosomal targeting. For example, such a receptor may be the cation-independent mannose-6-phosphate receptor (CI-MPR) which binds the mannose-6-phosphate (M6P) residues. In addition, the CI-MPR also binds other proteins including IGF-II. In some embodiments, a replacement enzyme suitable for the present invention contains M6P residues on the surface of the protein. In some embodiments, a replacement enzyme suitable for the present invention may contain bis-phosphorylated oligosaccharides which have higher binding affinity to the CI-MPR. In some embodiments, a suitable enzyme contains up to about an average of about at least 20% bis-phosphorylated oligosaccharides per enzyme. In other embodiments, a suitable enzyme may contain about 10%, 15%, 18%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60% bis-phosphorylated oligosaccharides per enzyme. While such bis-phosphorylated oligosaccharides may be naturally present on the enzyme, it should be noted that the enzymes may be modified to possess such oligosaccharides. For example, suitable replacement enzymes may be modified by certain enzymes which are capable of catalyzing the transfer of N-acetylglucosamine-L-phosphate from UDP-GlcNAc to the 6' position of α -1,2-linked mannoses on lysosomal enzymes. Methods and compositions for producing and using such enzymes are described by, for example, Canfield et al. in U.S. Pat. No. 6,537,785, and U.S. Pat. No. 6,534,300, each incorporated herein by reference.

[0291] In some embodiments, replacement enzymes for use in the present invention may be conjugated or fused to a lysosomal targeting moiety that is capable of binding to a receptor on the surface of brain cells. A suitable lysosomal targeting moiety can be IGF-I, IGF-II, RAP, p97, and variants, homologues or fragments thereof (e.g., including those peptide having a sequence at least 70%, 75%, 80%, 85%, 90%, or 95% identical to a wild-type mature human IGF-I, IGF-II, RAP, p97 peptide sequence).

2017202349 10 Apr 2017

[0292] In some embodiments, replacement enzymes suitable for the present invention have not been modified to enhance delivery or transport of such agents across the BBB and into the CNS.

Intrathecal Delivery

[0293] According to the present invention, a replacement enzyme is delivered to the CNS. In some embodiments, a replacement enzyme is delivered to the CNS by administering into the cerebrospinal fluid (CSF) of a subject in need of treatment. In some embodiments, intrathecal administration is used to deliver a desired replacement enzyme into the CSF. As used herein, intrathecal administration (also referred to as intrathecal injection) refers to an injection into the spinal canal (intrathecal space surrounding the spinal cord). Various techniques may be used including, without limitation, lateral cerebroventricular injection through a burrhole or cisternal or lumbar puncture or the like. Exemplary methods are described in Lazorthes et al. *Advances in Drug Delivery Systems and Applications in Neurosurgery*, 143-192 and Omayya et al., *Cancer Drug Delivery*, 1: 169-179, the contents of which are incorporated herein by reference.

[0294] According to the present invention, an enzyme may be injected at any region surrounding the spinal canal. In some embodiments, an enzyme is injected into the lumbar area or the cisterna magna or intraventricularly into a cerebral ventricle space. As used herein, the term “lumbar region” or “lumbar area” refers to the area between the third and fourth lumbar (lower back) vertebrae and, more inclusively, the L2-S1 region of the spine. Typically, intrathecal injection via the lumbar region or lumbar area is also referred to as “lumbar IT delivery” or “lumbar IT administration.” The term “cisterna magna” refers to the space around and below the cerebellum via the opening between the skull and the top of the spine. Typically, intrathecal injection via cisterna magna is also referred to as “cisterna magna delivery.” The term “cerebral ventricle” refers to the cavities in the brain that are continuous with the central canal of the spinal cord. Typically, injections via the cerebral ventricle cavities are referred to as intraventricular Cerebral (ICV) delivery.

2017202349 10 Apr 2017

[0295] In some embodiments, “intrathecal administration” or “intrathecal delivery” according to the present invention refers to lumbar IT administration or delivery, for example, delivered between the third and fourth lumbar (lower back) vertebrae and, more inclusively, the L2-S1 region of the spine. It is contemplated that lumbar IT administration or delivery distinguishes over cisterna magna delivery in that lumbar IT administration or delivery according to our invention provides better and more effective delivery to the distal spinal canal, while cisterna magna delivery, among other things, typically does not deliver well to the distal spinal canal.

Stable Formulations for IT Delivery

[0296] In some embodiments, desired enzymes are delivered in stable formulations for intrathecal delivery. Certain embodiments of the invention are based, at least in part, on the discovery that various formulations disclosed herein facilitate the effective delivery and distribution of one or more therapeutic agents (e.g., enzymes) to targeted tissues, cells and/or organelles of the CNS. Among other things, formulations described herein are capable of solubilizing high concentrations of therapeutic agents (e.g., proteins or enzymes) and are suitable for the delivery of such therapeutic agents to the CNS of subjects for the treatment of diseases having a CNS component and/or etiology. The compositions described herein are further characterized by improved stability and improved tolerability when administered to the CNS of a subject (e.g., intrathecally) in need thereof.

[0297] Before the present invention, traditional unbuffered isotonic saline and Elliott’s B solution, which is artificial CSF, were typically used for intrathecal delivery. A comparison depicting the compositions of CSF relative to Elliott’s B solution is included in Table 2 below. As shown in Table 2, the concentration of Elliott’s B Solution closely parallels that of the CSF. Elliott’s B Solution, however contains a very low buffer concentration and accordingly may not provide the adequate buffering capacity needed to stabilize therapeutic agents (e.g., proteins), especially over extended periods of time (e.g., during storage conditions). Furthermore, Elliott’s B Solution contains certain salts which may be incompatible with the formulations intended to deliver some therapeutic agents, and in particular proteins or enzymes. For example, the calcium

2017202349 10 Apr 2017

salts present in Elliott's B Solution are capable of mediating protein precipitation and thereby reducing the stability of the formulation.

Table 2

Solution	Na ⁺ mEq/L	K ⁺ mEq/L	Ca ⁺⁺ mEq/L	Mg ⁺⁺ mEq/L	HCO ₃ ⁻ mEq/L	Cl ⁻ mEq/L	pH	Phosphorous mg/L	Glucose mg/L
CSF	117- 137	2.3	2.2	2.2	22.9	113- 127	7.31	1.2-2.1	45-80
Elliott's B Sol'n	149	2.6	2.7	2.4	22.6	132	6.0- 7.5	2.3	80

Thus, in some embodiments, formulations suitable for intrathecal delivery according to the present invention are not synthetic or artificial CSF.

[0298] In some embodiments, formulations for intrathecal delivery have been formulated such that they are capable of stabilizing, or alternatively slowing or preventing the degradation, of one or more therapeutic agents formulated therewith (e.g., recombinant proteins). As used herein, the term "stable" refers to the ability of the therapeutic agent (e.g., a recombinant enzyme) to maintain its therapeutic efficacy (e.g., all or the majority of its intended biological activity and/or physiochemical integrity) over extended periods of time. The stability of a therapeutic agent, and the capability of the pharmaceutical composition to maintain stability of such therapeutic agent, may be assessed over extended periods of time (e.g., preferably for at least 1, 3, 6, 12, 18, 24, 30, 36 months or more). In the context of a formulation a stable formulation is one in which the therapeutic agent therein essentially retains its physical and/or chemical integrity and biological activity upon storage and during processes (such as freeze/thaw, mechanical mixing and lyophilization). For protein stability, it can be measure by formation of high molecular weight (HMW) aggregates, loss of enzyme activity, generation of peptide fragments and shift of charge profiles.

[0299] Stability of the therapeutic agent is of particular importance. Stability of the therapeutic agent may be further assessed relative to the biological activity or physiochemical integrity of the therapeutic agent over extended periods of time. For example, stability at a given time point may be compared against stability at an earlier time point (e.g., upon formulation day

2017202349 10 Apr 2017

0) or against unformulated therapeutic agent and the results of this comparison expressed as a percentage. Preferably, the pharmaceutical compositions of the present invention maintain at least 100%, at least 99%, at least 98%, at least 97%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55% or at least 50% of the therapeutic agent's biological activity or physiochemical integrity over an extended period of time (e.g., as measured over at least about 6-12 months, at room temperature or under accelerated storage conditions).

[0300] In some embodiments, therapeutic agents (e.g., desired enzymes) are soluble in formulations of the present invention. The term "soluble" as it relates to the therapeutic agents of the present invention refer to the ability of such therapeutic agents to form a homogenous solution. Preferably the solubility of the therapeutic agent in the solution into which it is administered and by which it is transported to the target site of action (e.g., the cells and tissues of the brain) is sufficient to permit the delivery of a therapeutically effective amount of the therapeutic agent to the targeted site of action. Several factors can impact the solubility of the therapeutic agents. For example, relevant factors which may impact protein solubility include ionic strength, amino acid sequence and the presence of other co-solubilizing agents or salts (e.g., calcium salts.) In some embodiments, the pharmaceutical compositions are formulated such that calcium salts are excluded from such compositions.

[0301] Thus, suitable formulations for intrathecal administration may contain a therapeutic agent (e.g., enzyme) of interest at various concentrations. In some embodiments, suitable formulations may contain a protein or enzyme of interest at a concentration up to about 300 mg/ml (e.g., up to about 250 mg/ml, up to 200 mg/ml, up to 150 mg/ml, up to 100 mg/ml, up to 90 mg/ml, up to 80 mg/ml, up to 70 mg/ml, up to 60 mg/ml, up to 50 mg/ml, up to 40 mg/ml, up to 30 mg/ml, up to 25 mg/ml, up to 20 mg/ml, up to 10 mg/ml). In some embodiments, suitable formulations may contain a protein or enzyme of interest at a concentration ranging between about 0-300 mg/ml (e.g., about 1-250 mg/ml, about 1-200 mg/ml, about 1-150 mg/ml, about 1-100 mg/ml, about 10-100 mg/ml, about 10-80 mg/ml, about 10-70 mg/ml, about 1-60 mg/ml, about 1-50 mg/ml, about 10-150 mg/ml, about 1-30 mg/ml). In some embodiments, formulations suitable for intrathecal delivery may contain a protein of interest at a concentration

2017202349 10 Apr 2017

of approximately 1 mg/ml, 3 mg/ml, 5 mg/ml, 10 mg/ml, 15 mg/ml, 20 mg/ml, 25 mg/ml, 50 mg/ml, 75 mg/ml, 100 mg/ml, 150 mg/ml, 200 mg/ml, 250 mg/ml or 300 mg/ml.

[0302] In some embodiments, isotonic solutions are used. In some embodiments, slightly hypertonic solutions (e.g., up to 300 mM (e.g., up to 250 mM, 200 mM, 175mM, 150 mM, 125 mM) sodium chloride in 5mM sodium phosphate at pH 7.0) and sugar-containing solutions (e.g., up to 3% (e.g., up to 2.4%, 2.0%, 1.5%, 1.0%) sucrose in 5mM sodium phosphate at pH 7.0) have been demonstrated to be well tolerated in monkeys. In some embodiments, a suitable CNS bolus formulation composition is saline (e.g., 150mM NaCl in water).

[0303] Many therapeutic agents, and in particular the proteins and enzymes of the present invention, require controlled pH and specific excipients to maintain their solubility and stability in the pharmaceutical compositions of the present invention. Table 3 below identifies certain exemplary aspects of protein formulations considered to be important for maintaining the solubility and stability of the protein therapeutic agents of the present invention.

Table 3

Parameter	Typical Range/Type	Rationale
pH	5 to 7.5	For stability Sometimes also for solubility
Buffer type	acetate, succinate, citrate, histidine, phosphate or Tris	To maintain optimal pH May also affect stability
Buffer concentration	5-50 mM	To maintain pH May also stabilize or add ionic strength
Tonicifier	NaCl, sugars, mannitol	To render iso-osmotic or isotonic solutions
Surfactant	Polysorbate 20, polysorbate 80	To stabilize against interfaces and shear
Other	Amino acids (e.g. arginine) at tens to hundreds of mM	For enhanced solubility or stability

[0304] The pH of the pharmaceutical composition is an additional factor which is capable of altering the solubility of a therapeutic agent (e.g., an enzyme or protein) in an aqueous pharmaceutical composition,. In some embodiments, pharmaceutical compositions of the present invention conytain one or more buffers. In some embodiments, compositions according

2017202349 10 Apr 2017

to the invention contain an amount of buffer sufficient to maintain the optimal pH of said composition between about 4.0-8.0, between about 5.0-7.5, between about 5.5-7.0, between about 6.0-7.0 and between about 6.0-7.5. In other embodiments, the buffer comprises up to about 50 mM (e.g., up to about 45 mM, 40 mM, 35 mM, 30 mM, 25 mM, 20 mM, 15 mM, 10 mM, 5 mM) of sodium phosphate. Suitable buffers include, for example acetate, succinate, citrate, phosphate, other organic acids and tris(hydroxymethyl)aminomethane ("Tris"). Suitable buffer concentrations can be from about 1 mM to about 100 mM, or from about 3 mM to about 20 mM, depending, for example, on the buffer and the desired isotonicity of the formulation. In some embodiments, a suitable buffering agent is present at a concentration of approximately 1 mM, 5 mM, 10 mM, 15 mM, 20 mM, 25 mM, 30 mM, 35 mM, 40 mM, 45 mM, 50 mM, 55 mM, 60 mM, 65 mM, 70 mM, 75 mM, 80 mM, 85 mM, 90 mM, 95 mM, or 100 mM.

[0305] In some embodiments, formulations contain an isotonicity agent to keep the formulations isotonic. As used in connection with IT delivery, by "isotonic" is meant that the formulation of interest has essentially the same osmolarity as human CSF. Isotonic formulations will generally have an osmolarity from about 240 mOsm/kg to about 350 mOsm/kg. Isotonicity can be measured using, for example, a vapor pressure or freezing point type osmometers. Exemplary isotonicity agents include, but are not limited to, glycine, sorbitol, mannitol, sodium chloride and arginine. In some embodiments, suitable isotonic agents may be present in formulations at a concentration from about 0.01 – 5 % (e.g., 0.05, 0.1, 0.15, 0.2, 0.3, 0.4, 0.5, 0.75, 1.0, 1.25, 1.5, 2.0, 2.5, 3.0, 4.0 or 5.0%) by weight.

[0306] In some embodiments, formulations may contain a stabilizing agent to protect the protein. Typically, a suitable stabilizing agent is a non-reducing sugar such as sucrose, raffinose, trehalose, or amino acids such as glycine, arginine and methionine. The amount of stabilizing agent in a formulation is generally such that the formulation will be isotonic. However, hypertonic formulations may also be suitable. In addition, the amount of stabilizing agent must not be too low such that an unacceptable amount of degradation/aggregation of the therapeutic agent occurs. Exemplary stabilizing agent concentrations in the formulation may range from about 1 mM to about 400 mM (e.g., from about 30 mM to about 300 mM, and from about 50 mM to about 100 mM), or alternatively, from 0.1% to 15% (e.g., from 1% to 10%, from 5% to 15%, from 5% to 10%) by weight. In some embodiments, the ratio of the mass amount of the

2017202349 10 Apr 2017

stabilizing agent and the therapeutic agent is about 1:1. In other embodiments, the ratio of the mass amount of the stabilizing agent and the therapeutic agent can be about 0.1:1, 0.2:1, 0.25:1, 0.4:1, 0.5:1, 1:1, 2:1, 2.6:1, 3:1, 4:1, 5:1, 10:1, or 20:1. In some embodiments, suitable for lyophilization, the stabilizing agent is also a lyoprotectants.

[0307] The pharmaceutical compositions, formulations and related methods of the invention are useful for delivering a variety of therapeutic agents to the CNS of a subject (e.g., intrathecally, intraventricularly or intracisternally) and for the treatment of the associated diseases. The pharmaceutical compositions of the present invention are particularly useful for delivering proteins and enzymes to subjects suffering from lysosomal storage disorders.

[0308] In some embodiments, it is desirable to add a surfactant to formulations. Exemplary surfactants include nonionic surfactants such as Polysorbates (e.g., Polysorbates 20 or 80); poloxamers (e.g., poloxamer 188); Triton; sodium dodecyl sulfate (SDS); sodium laurel sulfate; sodium octyl glycoside; lauryl-, myristyl-, linoleyl-, or stearyl-sulfobetaine; lauryl-, myristyl-, linoleyl- or stearyl-sarcosine; linoleyl-, myristyl-, or cetyl-betaine; lauroamidopropyl-, cocamidopropyl-, linoleamidopropyl-, myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-betaine (e.g., lauroamidopropyl); myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-dimethylamine; sodium methyl cocoyl-, or disodium methyl ofeyle-aurate; and the MONAQUATTM series (Mona Industries, Inc., Paterson, N.J.), polyethyl glycol, polypropyl glycol, and copolymers of ethylene and propylene glycol (e.g., Pluronic, PF68, etc). Typically, the amount of surfactant added is such that it reduces aggregation of the protein and minimizes the formation of particulates or effervescences. For example, a surfactant may be present in a formulation at a concentration from about 0.001 – 0.5% (e.g., about 0.005 – 0.05%, or 0.005 – 0.01%). In particular, a surfactant may be present in a formulation at a concentration of approximately 0.005%, 0.01%, 0.02%, 0.1%, 0.2%, 0.3%, 0.4%, or 0.5%, etc.

[0309] In some embodiments, suitable formulations may further include one or more bulking agents, in particular, for lyophilized formulations. A “bulking agent” is a compound which adds mass to the lyophilized mixture and contributes to the physical structure of the lyophilized cake. For example, a bulking agent may improve the appearance of lyophilized cake (e.g., essentially uniform lyophilized cake). Suitable bulking agents include, but are not limited

2017202349 10 Apr 2017

to, sodium chloride, lactose, mannitol, glycine, sucrose, trehalose, hydroxyethyl starch.

Exemplary concentrations of bulking agents are from about 1% to about 10% (e.g., 1.0%, 1.5%, 2.0%, 2.5%, 3.0%, 3.5%, 4.0%, 4.5%, 5.0%, 5.5%, 6.0%, 6.5%, 7.0%, 7.5%, 8.0%, 8.5%, 9.0%, 9.5%, and 10.0%).

[0310] Formulations in accordance with the present invention can be assessed based on product quality analysis, reconstitution time (if lyophilized), quality of reconstitution (if lyophilized), high molecular weight, moisture, and glass transition temperature. Typically, protein quality and product analysis include product degradation rate analysis using methods including, but not limited to, size exclusion HPLC (SE-HPLC), cation exchange-HPLC (CEX-HPLC), X-ray diffraction (XRD), modulated differential scanning calorimetry (mDSC), reversed phase HPLC (RP-HPLC), multi-angle light scattering (MALS), fluorescence, ultraviolet absorption, nephelometry, capillary electrophoresis (CE), SDS-PAGE, and combinations thereof. In some embodiments, evaluation of product in accordance with the present invention may include a step of evaluating appearance (either liquid or cake appearance).

[0311] Generally, formulations (lyophilized or aqueous) can be stored for extended periods of time at room temperature. Storage temperature may typically range from 0 °C to 45 °C (e.g., 4°C, 20 °C, 25 °C, 45 °C etc.). Formulations may be stored for a period of months to a period of years. Storage time generally will be 24 months, 12 months, 6 months, 4.5 months, 3 months, 2 months or 1 month. Formulations can be stored directly in the container used for administration, eliminating transfer steps.

[0312] Formulations can be stored directly in the lyophilization container (if lyophilized), which may also function as the reconstitution vessel, eliminating transfer steps. Alternatively, lyophilized product formulations may be measured into smaller increments for storage. Storage should generally avoid circumstances that lead to degradation of the proteins, including but not limited to exposure to sunlight, UV radiation, other forms of electromagnetic radiation, excessive heat or cold, rapid thermal shock, and mechanical shock.

[0313] In some embodiments, formulations according to the present invention are in a liquid or aqueous form. In some embodiments, formulations of the present invention are

2017202349 10 Apr 2017

lyophilized. Such lyophilized formulations may be reconstituted by adding one or more diluents thereto prior to administration to a subject. Suitable diluents include, but are not limited to, sterile water, bacteriostatic water for injection and sterile saline solution. Preferably, upon reconstitution, the therapeutic agent contained therein is stable, soluble and demonstrates tolerability upon administration to a subject

[0314] The pharmaceutical compositions of the present invention are characterized by their tolerability. As used herein, the terms “tolerable” and “tolerability” refer to the ability of the pharmaceutical compositions of the present invention to not elicit an adverse reaction in the subject to whom such composition is administered, or alternatively not to elicit a serious adverse reaction in the subject to whom such composition is administered. In some embodiments, the pharmaceutical compositions of the present invention are well tolerated by the subject to whom such compositions is administered.

Device for Intrathecal Delivery

[0315] Various devices may be used for intrathecal delivery according to the present invention. In some embodiments, a device for intrathecal administration contains a fluid access port (e.g., injectable port); a hollow body (e.g., catheter) having a first flow orifice in fluid communication with the fluid access port and a second flow orifice configured for insertion into spinal cord; and a securing mechanism for securing the insertion of the hollow body in the spinal cord. As a non-limiting example shown in Figure 1, a suitable securing mechanism contains one or more nobs mounted on the surface of the hollow body and a sutured ring adjustable over the one or more nobs to prevent the hollow body (e.g., catheter) from slipping out of the spinal cord. In various embodiments, the fluid access port comprises a reservoir. In some embodiments, the fluid access port comprises a mechanical pump (e.g., an infusion pump). In some embodiments, an implanted catheter is connected to either a reservoir (e.g., for bolus delivery), or an infusion pump. The fluid access port may be implanted or external

[0316] In some embodiments, intrathecal administration may be performed by either lumbar puncture (i.e., slow bolus) or via a port-catheter delivery system (i.e., infusion or bolus).

2017202349 10 Apr 2017

In some embodiments, the catheter is inserted between the laminae of the lumbar vertebrae and the tip is threaded up the thecal space to the desired level (generally L3-L4) (Figure 2).

[0317] Relative to intravenous administration, a single dose volume suitable for intrathecal administration is typically small. Typically, intrathecal delivery according to the present invention maintains the balance of the composition of the CSF as well as the intracranial pressure of the subject. In some embodiments, intrathecal delivery is performed absent the corresponding removal of CSF from a subject. In some embodiments, a suitable single dose volume may be e.g., less than about 10 ml, 8 ml, 6 ml, 5 ml, 4 ml, 3 ml, 2 ml, 1.5 ml, 1 ml, or 0.5 ml. In some embodiments, a suitable single dose volume may be about 0.5-5 ml, 0.5-4 ml, 0.5-3 ml, 0.5-2 ml, 0.5-1 ml, 1-3 ml, 1-5 ml, 1.5-3 ml, 1-4 ml, or 0.5-1.5 ml. In some embodiments, intrathecal delivery according to the present invention involves a step of removing a desired amount of CSF first. In some embodiments, less than about 10 ml (e.g., less than about 9 ml, 8 ml, 7 ml, 6 ml, 5 ml, 4 ml, 3 ml, 2 ml, 1 ml) of CSF is first removed before IT administration. In those cases, a suitable single dose volume may be e.g., more than about 3 ml, 4 ml, 5 ml, 6 ml, 7 ml, 8 ml, 9 ml, 10 ml, 15 ml, or 20 ml.

[0318] Various other devices may be used to effect intrathecal administration of a therapeutic composition. For example, formulations containing desired enzymes may be given using an Ommaya reservoir which is in common use for intrathecally administering drugs for meningeal carcinomatosis (Lancet 2: 983-84, 1963). More specifically, in this method, a ventricular tube is inserted through a hole formed in the anterior horn and is connected to an Ommaya reservoir installed under the scalp, and the reservoir is subcutaneously punctured to intrathecally deliver the particular enzyme being replaced, which is injected into the reservoir. Other devices for intrathecal administration of therapeutic compositions or formulations to an individual are described in U.S. Pat. No. 6,217,552, incorporated herein by reference. Alternatively, the drug may be intrathecally given, for example, by a single injection, or continuous infusion. It should be understood that the dosage treatment may be in the form of a single dose administration or multiple doses.

[0319] For injection, formulations of the invention can be formulated in liquid solutions. In addition, the enzyme may be formulated in solid form and re-dissolved or suspended

2017202349 10 Apr 2017

immediately prior to use. Lyophilized forms are also included. The injection can be, for example, in the form of a bolus injection or continuous infusion (e.g., using infusion pumps) of the enzyme.

[0320] In one embodiment of the invention, the enzyme is administered by lateral cerebro ventricular injection into the brain of a subject. The injection can be made, for example, through a burr hole made in the subject's skull. In another embodiment, the enzyme and/or other pharmaceutical formulation is administered through a surgically inserted shunt into the cerebral ventricle of a subject. For example, the injection can be made into the lateral ventricles, which are larger. In some embodiments, injection into the third and fourth smaller ventricles can also be made.

[0321] In yet another embodiment, the pharmaceutical compositions used in the present invention are administered by injection into the cisterna magna, or lumbar area of a subject.

[0322] In another embodiment of the method of the invention, the pharmaceutically acceptable formulation provides sustained delivery, e.g., "slow release" of the enzyme or other pharmaceutical composition used in the present invention, to a subject for at least one, two, three, four weeks or longer periods of time after the pharmaceutically acceptable formulation is administered to the subject.

[0323] As used herein, the term "sustained delivery" refers to continual delivery of a pharmaceutical formulation of the invention *in vivo* over a period of time following administration, preferably at least several days, a week or several weeks. Sustained delivery of the composition can be demonstrated by, for example, the continued therapeutic effect of the enzyme over time (e.g., sustained delivery of the enzyme can be demonstrated by continued reduced amount of storage granules in the subject). Alternatively, sustained delivery of the enzyme may be demonstrated by detecting the presence of the enzyme *in vivo* over time.

2017202349 10 Apr 2017

Delivery to Target Tissues

[0324] As discussed above, one of the surprising and important features of the present invention is that therapeutic agents, in particular, replacement enzymes administered using inventive methods and compositions of the present invention are able to effectively and extensively diffuse across the brain surface and penetrate various layers or regions of the brain, including deep brain regions. In addition, inventive methods and compositions of the present invention effectively deliver therapeutic agents (e.g., replacement enzymes) to various tissues, neurons or cells of spinal cord, including the lumbar region, which is hard to target by existing CNS delivery methods such as ICV injection. Furthermore, inventive methods and compositions of the present invention deliver sufficient amount of therapeutic agents (e.g., replacement enzymes) to blood stream and various peripheral organs and tissues.

[0325] Thus, in some embodiments, a therapeutic protein (e.g., a replacement enzyme) is delivered to the central nervous system of a subject. In some embodiments, a therapeutic protein (e.g., a replacement enzyme) is delivered to one or more of target tissues of brain, spinal cord, and/or peripheral organs. As used herein, the term “target tissues” refers to any tissue that is affected by the lysosomal storage disease to be treated or any tissue in which the deficient lysosomal enzyme is normally expressed. In some embodiments, target tissues include those tissues in which there is a detectable or abnormally high amount of enzyme substrate, for example stored in the cellular lysosomes of the tissue, in patients suffering from or susceptible to the lysosomal storage disease. In some embodiments, target tissues include those tissues that display disease-associated pathology, symptom, or feature. In some embodiments, target tissues include those tissues in which the deficient lysosomal enzyme is normally expressed at an elevated level. As used herein, a target tissue may be a brain target tissue, a spinal cord target tissue and/or a peripheral target tissue. Exemplary target tissues are described in detail below.

Brain Target Tissues

[0326] In general, the brain can be divided into different regions, layers and tissues. For example, meningeal tissue is a system of membranes which envelops the central nervous system,

2017202349 10 Apr 2017

including the brain. The meninges contain three layers, including dura matter, arachnoid matter, and pia matter. In general, the primary function of the meninges and of the cerebrospinal fluid is to protect the central nervous system. In some embodiments, a therapeutic protein in accordance with the present invention is delivered to one or more layers of the meninges.

[0327] The brain has three primary subdivisions, including the cerebrum, cerebellum, and brain stem. The cerebral hemispheres, which are situated above most other brain structures and are covered with a cortical layer. Underneath the cerebrum lies the brainstem, which resembles a stalk on which the cerebrum is attached. At the rear of the brain, beneath the cerebrum and behind the brainstem, is the cerebellum.

[0328] The diencephalon, which is located near the midline of the brain and above the mesencephalon, contains the thalamus, metathalamus, hypothalamus, epithalamus, prethalamus, and pretectum. The mesencephalon, also called the midbrain, contains the tectum, tegumentum, ventricular mesocoelia, and cerebral peduncels, the red nucleus, and the cranial nerve III nucleus. The mesencephalon is associated with vision, hearing, motor control, sleep/wake, alertness, and temperature regulation.

[0329] Regions of tissues of the central nervous system, including the brain, can be characterized based on the depth of the tissues. For example, CNS (e.g., brain) tissues can be characterized as surface or shallow tissues, mid-depth tissues, and/or deep tissues.

[0330] According to the present invention, a therapeutic protein (e.g., a replacement enzyme) may be delivered to any appropriate brain target tissue(s) associated with a particular disease to be treated in a subject. In some embodiments, a therapeutic protein (e.g., a replacement enzyme) in accordance with the present invention is delivered to surface or shallow brain target tissue. In some embodiments, a therapeutic protein in accordance with the present invention is delivered to mid-depth brain target tissue. In some embodiments, a therapeutic protein in accordance with the present invention is delivered to deep brain target tissue. In some embodiments, a therapeutic protein in accordance with the present invention is delivered to a combination of surface or shallow brain target tissue, mid-depth brain target tissue, and/or deep brain target tissue. In some embodiments, a therapeutic protein in accordance with the present

2017202349 10 Apr 2017

invention is delivered to a deep brain tissue at least 4 mm, 5 mm, 6 mm, 7 mm, 8 mm, 9 mm, 10 mm or more below (or internal to) the external surface of the brain.

[0331] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to one or more surface or shallow tissues of cerebrum. In some embodiments, the targeted surface or shallow tissues of the cerebrum are located within 4 mm from the surface of the cerebrum. In some embodiments, the targeted surface or shallow tissues of the cerebrum are selected from pia mater tissues, cerebral cortical ribbon tissues, hippocampus, Virchow Robin space, blood vessels within the VR space, the hippocampus, portions of the hypothalamus on the inferior surface of the brain, the optic nerves and tracts, the olfactory bulb and projections, and combinations thereof.

[0332] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to one or more deep tissues of the cerebrum. In some embodiments, the targeted surface or shallow tissues of the cerebrum are located 4 mm (e.g., 5 mm, 6 mm, 7 mm, 8 mm, 9 mm, or 10 mm) below (or internal to) the surface of the cerebrum. In some embodiments, targeted deep tissues of the cerebrum include the cerebral cortical ribbon. In some embodiments, targeted deep tissues of the cerebrum include one or more of the diencephalon (e.g., the hypothalamus, thalamus, prethalamus, subthalamus, etc.), metencephalon, lentiform nuclei, the basal ganglia, caudate, putamen, amygdala, globus pallidus, and combinations thereof.

[0333] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to one or more tissues of the cerebellum. In certain embodiments, the targeted one or more tissues of the cerebellum are selected from the group consisting of tissues of the molecular layer, tissues of the Purkinje cell layer, tissues of the Granular cell layer, cerebellar peduncles, and combination thereof. In some embodiments, therapeutic agents (e.g., enzymes) are delivered to one or more deep tissues of the cerebellum including, but not limited to, tissues of the Purkinje cell layer, tissues of the Granular cell layer, deep cerebellar white matter tissue (e.g., deep relative to the Granular cell layer), and deep cerebellar nuclei tissue.

[0334] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to one or more tissues of the brainstem. In some embodiments, the targeted one or more tissues of the brainstem include brain stem white matter tissue and/or brain stem nuclei tissue.

2017202349 10 Apr 2017

[0335] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to various brain tissues including, but not limited to, gray matter, white matter, periventricular areas, pia-arachnoid, meninges, neocortex, cerebellum, deep tissues in cerebral cortex, molecular layer, caudate/putamen region, midbrain, deep regions of the pons or medulla, and combinations thereof.

[0336] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to various cells in the brain including, but not limited to, neurons, glial cells, perivascular cells and/or meningeal cells. In some embodiments, a therapeutic protein is delivered to oligodendrocytes of deep white matter.

Spinal Cord

[0337] In general, regions or tissues of the spinal cord can be characterized based on the depth of the tissues. For example, spinal cord tissues can be characterized as surface or shallow tissues, mid-depth tissues, and/or deep tissues.

[0338] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to one or more surface or shallow tissues of the spinal cord. In some embodiments, a targeted surface or shallow tissue of the spinal cord is located within 4 mm from the surface of the spinal cord. In some embodiments, a targeted surface or shallow tissue of the spinal cord contains pia matter and/or the tracts of white matter.

[0339] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to one or more deep tissues of the spinal cord. In some embodiments, a targeted deep tissue of the spinal cord is located internal to 4 mm from the surface of the spinal cord. In some embodiments, a targeted deep tissue of the spinal cord contains spinal cord grey matter and/or ependymal cells.

[0340] In some embodiments, therapeutic agents (e.g., enzymes) are delivered to neurons of the spinal cord.

2017202349 10 Apr 2017

Peripheral Target Tissues

[0341] As used herein, peripheral organs or tissues refer to any organs or tissues that are not part of the central nervous system (CNS). Peripheral target tissues may include, but are not limited to, blood system, liver, kidney, heart, endothelium, bone marrow and bone marrow derived cells, spleen, lung, lymph node, bone, cartilage, ovary and testis. In some embodiments, a therapeutic protein (e.g., a replacement enzyme) in accordance with the present invention is delivered to one or more of the peripheral target tissues.

Biodistribution and bioavailability

[0342] In various embodiments, once delivered to the target tissue, a therapeutic agent (e.g., a replacement enzyme) is localized intracellularly. For example, a therapeutic agent (e.g., enzyme) may be localized to exons, axons, lysosomes, mitochondria or vacuoles of a target cell (e.g., neurons such as Purkinje cells). For example, in some embodiments intrathecally-administered enzymes demonstrate translocation dynamics such that the enzyme moves within the perivascular space (e.g., by pulsation-assisted convective mechanisms). In addition, active axonal transport mechanisms relating to the association of the administered protein or enzyme with neurofilaments may also contribute to or otherwise facilitate the distribution of intrathecally-administered proteins or enzymes into the deeper tissues of the central nervous system.

[0343] In some embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention may achieve therapeutically or clinically effective levels or activities in various target tissues described herein. As used herein, a therapeutically or clinically effective level or activity is a level or activity sufficient to confer a therapeutic effect in a target tissue. The therapeutic effect may be objective (i.e., measurable by some test or marker) or subjective (i.e., subject gives an indication of or feels an effect). For example, a therapeutically or clinically effective level or activity may be an enzymatic level or activity that is sufficient to ameliorate symptoms associated with the disease in the target tissue (e.g., GAG storage).

2017202349 10 Apr 2017

[0344] In some embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention may achieve an enzymatic level or activity that is at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% of the normal level or activity of the corresponding lysosomal enzyme in the target tissue. In some embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention may achieve an enzymatic level or activity that is increased by at least 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold or 10-fold as compared to a control (e.g., endogenous levels or activities without the treatment). In some embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention may achieve an increased enzymatic level or activity at least approximately 10 nmol/hr/mg, 20 nmol/hr/mg, 40 nmol/hr/mg, 50 nmol/hr/mg, 60 nmol/hr/mg, 70 nmol/hr/mg, 80 nmol/hr/mg, 90 nmol/hr/mg, 100 nmol/hr/mg, 150 nmol/hr/mg, 200 nmol/hr/mg, 250 nmol/hr/mg, 300 nmol/hr/mg, 350 nmol/hr/mg, 400 nmol/hr/mg, 450 nmol/hr/mg, 500 nmol/hr/mg, 550 nmol/hr/mg or 600 nmol/hr/mg in a target tissue.

[0345] In some embodiments, inventive methods according to the present invention are particularly useful for targeting the lumbar region. In some embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention may achieve an increased enzymatic level or activity in the lumbar region of at least approximately 500 nmol/hr/mg, 600 nmol/hr/mg, 700 nmol/hr/mg, 800 nmol/hr/mg, 900 nmol/hr/mg, 1000 nmol/hr/mg, 1500 nmol/hr/mg, 2000 nmol/hr/mg, 3000 nmol/hr/mg, 4000 nmol/hr/mg, 5000 nmol/hr/mg, 6000 nmol/hr/mg, 7000 nmol/hr/mg, 8000 nmol/hr/mg, 9000 nmol/hr/mg, or 10,000 nmol/hr/mg.

[0346] In general, therapeutic agents (e.g., replacement enzymes) delivered according to the present invention have sufficiently long half time in CSF and target tissues of the brain, spinal cord, and peripheral organs. In some embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention may have a half-life of at least approximately 30 minutes, 45 minutes, 60 minutes, 90 minutes, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 12 hours, 16 hours, 18 hours, 20 hours, 25 hours, 30 hours, 35 hours, 40 hours, up to 3 days, up to 7 days, up to 14 days, up to 21 days or up to a month. In some embodiments, In some embodiments, a therapeutic agent (e.g., a

2017202349 10 Apr 2017

replacement enzyme) delivered according to the present invention may retain detectable level or activity in CSF or bloodstream after 12 hours, 24 hours, 30 hours, 36 hours, 42 hours, 48 hours, 54 hours, 60 hours, 66 hours, 72 hours, 78 hours, 84 hours, 90 hours, 96 hours, 102 hours, or a week following administration. Detectable level or activity may be determined using various methods known in the art.

[0347] In certain embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention achieves a concentration of at least 30 μ g/ml in the CNS tissues and cells of the subject following administration (e.g., one week, 3 days, 48 hours, 36 hours, 24 hours, 18 hours, 12 hours, 8 hours, 6 hours, 4 hours, 3 hours, 2 hours, 1 hour, 30 minutes, or less, following intrathecal administration of the pharmaceutical composition to the subject). In certain embodiments, a therapeutic agent (e.g., a replacement enzyme) delivered according to the present invention achieves a concentration of at least 20 μ g/ml, at least 15 μ g/ml, at least 10 μ g/ml, at least 7.5 μ g/ml, at least 5 μ g/ml, at least 2.5 μ g/ml, at least 1.0 μ g/ml or at least 0.5 μ g/ml in the targeted tissues or cells of the subject (e.g., brain tissues or neurons) following administration to such subject (e.g., one week, 3 days, 48 hours, 36 hours, 24 hours, 18 hours, 12 hours, 8 hours, 6 hours, 4 hours, 3 hours, 2 hours, 1 hour, 30 minutes, or less following intrathecal administration of such pharmaceutical compositions to the subject).

Treatment of Lysosomal Storage Diseases By Intrathecal Administration

[0348] The lysosomal storage diseases represent a group of relatively rare inherited metabolic disorders that result from defects in lysosomal function. The lysosomal diseases are characterized by the accumulation of undigested macromolecules, including those enzyme substrates, within the lysosomes (see Table 1), which results in an increase in the size and number of such lysosomes and ultimately in cellular dysfunction and clinical abnormalities.

[0349] Inventive methods described herein can advantageously facilitate the delivery of one or more therapeutic agents (e.g., one or more replacement enzymes) to targeted organelles. For example, because lysosomal storage disorders such as Hunter syndrome are characterized by

2017202349 10 Apr 2017

an accumulation of glycosaminoglycans (GAG) in the lysosomes of affected cells, the lysosomes represent an desired target organelle for the treatment of the lysosomal storage disorders.

[0350] Inventive methods and compositions of the present invention are particularly useful for treating those diseases having a CNS etiology or component. Lysosomal storage diseases having a CNS etiology or component, include for example and without limitation Sanfilippo syndrome Type A, Sanfilippo syndrome type B, Hunter syndrome, metachromatic leukodystrophy and globoid cell leukodystrophy. Prior to the present invention, traditional therapies are limited in that they are administered to subjects intravenously, and are generally only effective in treating the somatic symptoms of the underlying enzyme deficiency. The compositions and methods of the present invention may advantageously be administered directly into the CNS of a subject suffering from a disease having such a CNS etiology thereby achieving a therapeutic concentration within the affected cells and tissues of the CNS (e.g., the brain), thus overcoming the limitations associated with traditional systemic administration of such therapeutic agents.

[0351] In some embodiments, inventive methods and compositions of the invention are useful for treating both the neurologic and the somatic sequelae or symptoms of lysosomal storage disorders. For example, some embodiments of the invention relate to compositions and methods of delivering one or more therapeutic agents to the CNS of a subject (e.g., intrathecally, intraventricularly or intracisternally) for the treatment of the CNS or neurologic sequelae and manifestations of a lysosomal storage disease, while also treating the systemic or somatic manifestations of that lysosomal storage disease. For example, some compositions of the present invention may be administered to a subject intrathecally, thereby delivering one or more therapeutic agents to the CNS of the subject and treating the neurological sequelae, coupled with the intravenous administration of one or more therapeutic agents to deliver such therapeutic agents to both the cells and tissues of the systemic circulation (e.g., cells and tissues of heart, lungs, liver, kidney or lymph nodes) to thereby treat the somatic sequelae. For example, a subject having or otherwise affected by a lysosomal storage disease (e.g., Hunter syndrome) may be administered a pharmaceutical composition comprising one or more therapeutic agents (e.g., iduronate-2-sulfatase) intrathecally at least once per week, biweekly, monthly, bimonthly or more to treat the neurologic sequelae, while a different therapeutic agent is administered to the

2017202349 10 Apr 2017

subject intravenously on a more frequent basis (e.g., once per day, every other day, three times a week or weekly) to treat the systemic or somatic manifestations of the disease.

[0352] For example, patients suffering from Hunter syndrome exhibit histological changes in the brains which may include atrophy, cortical neuronal swelling, cerebral white matter reduction, dilated perivascular spaces and Purkinje cell dendrite swelling. Magnetic resonance imaging/spectroscopy studies have shown that severe diffuse lesions involving the white matter, brain atrophy, and hydrocephalus were more common in patients with cognitive impairment compared to those without impairment. (Vedolin, L., et al., AJNR Am J Neuroradiol (2007) 28, 1029-1033). Even patients without extreme neurologic sequelae such as mental retardation or developmental delays were shown to have brain abnormalities that included atrophy, ventriculomegaly, and enlarged perivascular spaces. (Matheus, MG, et al., Neuroradiology (2004) 46, 666-672.)

[0353] As a non-limiting example, mucopolysaccharidosis type IIIA (MPS IIIA; Sanfilippo syndrome type A) is the most severe form of Sanfilippo syndrome type A and affects approximately 1 in 100,000 people worldwide. Sanfilippo syndrome type A (Sanfilippo A) is characterized by a deficiency of the enzyme heparan *N*-sulfatase (HNS), an exosulfatase involved in the lysosomal catabolism of glycosaminoglycan (GAG) heparan sulfate (Neufeld EF, et al. The Metabolic and Molecular Bases of Inherited Disease (2001) pp. 3421-3452). In the absence of this enzyme, GAG heparan sulfate accumulates in lysosomes of neurons and glial cells, with lesser accumulation outside the brain.

[0354] As a non-limiting example, mucopolysaccharidosis type IIIB (MPS IIIB; Sanfilippo syndrome type B disease) is an autosomal recessive disorder that is characterized by a deficiency of the enzyme alpha-N-acetyl-glucosaminidase (Naglu). In the absence of this enzyme, GAG heparan sulfate accumulates in lysosomes of neurons and glial cells, with lesser accumulation outside the brain.

[0355] As a non-limiting example, globoid cell leukodystrophy (GLD) is a rare autosomal recessive lysosomal storage disorder caused by defective function of galactocerebrosidase (GALC). GALC is a soluble lysosomal acid hydrolase enzyme which degrades galactosylceramide, a normal component of myelin, into galactose and ceramide, and

2017202349 10 Apr 2017

psychosine (galactosylsphingosine), a toxic byproduct of galactosylceramide synthesis, into galactose and sphingosine. GALC deficiency leads to neurologic injury of the central and peripheral nervous systems (CNS and PNS respectively) in two related, but distinct pathways. The first pathway leads to excessive psychosine accumulation with resultant apoptosis of myelinating cells. In the second pathway, galactosylceramide accumulates and is phagocytosed in activated microglia, producing the characteristic globoid cell for which the disease is named. In contrast to other lysosomal storage diseases which accumulate undegraded substrate, there is generally no increase in total galactosylceramide in neural tissue.

[0356] A defining clinical feature of this disorder is central nervous system (CNS) degeneration, which results in loss of, or failure to attain, major developmental milestones. The progressive cognitive decline culminates in dementia and premature mortality. The disease can manifest itself in young children (Early-onset GLD), or in individuals of any age (Late-onset GLD). The lifespan of an individual affected with Early-onset GLD typically does not extend beyond the age of two years. Late-onset GLD can appear in individuals of any age and the progression of the disease can vary greatly.

[0357] Metachromatic Leukodystrophy Disease (MLD), is an autosomal recessive disorder resulting from a deficiency of the enzyme Arylsulfatase A (ASA). ASA, which is encoded by the ARSA gene in humans, is an enzyme that breaks down cerebroside 3-sulfate or sphingolipid 3-O-sulfogalactosylceramide (sulfatide) into cerebroside and sulfate. In the absence of the enzyme, sulfatides accumulate in the nervous system (e.g., myelin sheaths, neurons and glial cells) and to a lesser extent in visceral organs. The consequence of these molecular and cellular events is progressive demyelination and axonal loss within the CNS and PNS, which is accompanied clinically by severe motor and cognitive dysfunction.

[0358] A defining clinical feature of this disorder is central nervous system (CNS) degeneration, which results in cognitive impairment (e.g., mental retardation, nervous disorders, and blindness, among others).

[0359] As a non-limiting example, MLD can manifest itself in young children (Late-infantile form), where affected children typically begin showing symptoms just after the first year of life (e.g., at about 15-24 months), and generally do not survive past the age of 5 years.

2017202349 10 Apr 2017

MLD can manifest itself in children (Juvenile form), where affected children typically show cognitive impairment by about the age of 3-10 years, and life-span can vary (e.g., in the range of 10-15 years after onset of symptoms). MLD can manifest itself in adults (Adult-onset form) and can appear in individuals of any age (e.g., typically at age 16 and later) and the progression of the disease can vary greatly.

[0360] Thus, in some embodiments, inventive methods and compositions deliver one or more therapeutic agents (e.g., one or more replacement enzymes) to one or more organelles (e.g., the lysosomes) of target tissues and cells of the brain, spinal cord and/or peripheral organs to effect treatment various lysosomal storage diseases. As used herein, the terms, “treat” or “treatment,” as used herein, refers to amelioration of one or more symptoms associated with the disease, prevention or delay of the onset of one or more symptoms of the disease, and/or lessening of the severity or frequency of one or more symptoms of the disease.

[0361] In some embodiments, treatment refers to partially or complete alleviation, amelioration, relief, inhibition, delaying onset, reducing severity and/or incidence of neurological impairment in a patient suffering from or susceptible to a lysosomal disease. As used herein, the term “neurological impairment” includes various symptoms associated with impairment of the central nervous system (e.g., the brain and spinal cord). Symptoms of neurological impairment may include, for example, developmental delay, progressive cognitive impairment, hearing loss, impaired speech development, deficits in motor skills, hyperactivity, aggressiveness and/or sleep disturbances, among others.

[0362] In some embodiments, treatment refers to decreased lysosomal storage (e.g., macromolecules stored such as GAG) in various tissues. In some embodiments, treatment refers to decreased lysosomal storage in brain target tissues, spinal cord neurons, and/or peripheral target tissues. In certain embodiments, lysosomal storage is decreased by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% or more as compared to a control. In some embodiments, lysosomal storage is decreased by at least 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold or 10-fold as compared to a control. In some embodiments, lysosomal storage is measured by the presence of lysosomal storage granules (e.g., zebra-striped morphology).

2017202349 10 Apr 2017

[0363] In some embodiments, treatment refers to reduced vacuolization in neurons (e.g., neurons containing Purkinje cells). In certain embodiments, vacuolization in neurons is decreased by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% or more as compared to a control. In some embodiments, vacuolization is decreased by at least 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold or 10-fold as compared to a control.

[0364] In certain embodiments, treatment according to the present invention results in a reduction (e.g., about a 5%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 90%, 95%, 97.5%, 99% or more reduction) or a complete elimination of the presence, or alternatively the accumulation, of one or more pathological or biological markers which are associated with the lysosomal storage diseases. Such reduction or elimination may be particularly evident in the cells and tissues of the CNS (e.g., neurons and oligodendrocytes). For example, in some embodiments, upon administration to a subject the pharmaceutical compositions of the present invention demonstrate or achieve a reduction in the accumulation of the biomarker lysosomal associated membrane protein 1 (LAMP1) in the CNS cells and tissues of the subject (e.g., in the cerebral cortex, cerebellum, caudate nucleus and putamen, white matter and/or thalamus). LAMP1 is a glycoprotein highly expressed in lysosomal membranes and its presence is elevated many patients with a lysosomal storage disorder. (Meikle, et al. Clin Chem. (1997)43:1325-1335.) The presence or absence of LAMP1 in patients (e.g., as determined by LAMP staining) with a lysosomal storage disease therefore may provide a useful indicator of lysosomal activity and a marker for both the diagnosis and monitoring of lysosomal storage diseases.

[0365] Accordingly, some embodiments of the present invention relate to methods of reducing or otherwise eliminating the presence or accumulation of one or more pathological or biological markers associated with a disease (e.g., a lysosomal storage disease). Similarly, some embodiments of the invention relate to methods of increasing the degradation (or the rate of degradation) of one or more pathological or biological markers (e.g., LAMP1) associated with lysosomal storage diseases.

2017202349 10 Apr 2017

[0366] In some embodiments, treatment refers to decreased progression of loss of cognitive ability. In certain embodiments, progression of loss of cognitive ability is decreased by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% or more as compared to a control. In some embodiments, treatment refers to decreased developmental delay. In certain embodiments, developmental delay is decreased by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 100% or more as compared to a control.

[0367] In some embodiments, treatment refers to increased survival (e.g. survival time). For example, treatment can result in an increased life expectancy of a patient. In some embodiments, treatment according to the present invention results in an increased life expectancy of a patient by more than about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 100%, about 105%, about 110%, about 115%, about 120%, about 125%, about 130%, about 135%, about 140%, about 145%, about 150%, about 155%, about 160%, about 165%, about 170%, about 175%, about 180%, about 185%, about 190%, about 195%, about 200% or more, as compared to the average life expectancy of one or more control individuals with similar disease without treatment. In some embodiments, treatment according to the present invention results in an increased life expectancy of a patient by more than about 6 month, about 7 months, about 8 months, about 9 months, about 10 months, about 11 months, about 12 months, about 2 years, about 3 years, about 4 years, about 5 years, about 6 years, about 7 years, about 8 years, about 9 years, about 10 years or more, as compared to the average life expectancy of one or more control individuals with similar disease without treatment. In some embodiments, treatment according to the present invention results in long term survival of a patient. As used herein, the term “long term survival” refers to a survival time or life expectancy longer than about 40 years, 45 years, 50 years, 55 years, 60 years, or longer.

[0368] The terms, “improve,” “increase” or “reduce,” as used herein, indicate values that are relative to a control. In some embodiments, a suitable control is a baseline measurement, such as a measurement in the same individual prior to initiation of the treatment described herein, or a measurement in a control individual (or multiple control individuals) in the absence

2017202349 10 Apr 2017

of the treatment described herein. A “control individual” is an individual afflicted with the same disease, who is about the same age and/or gender as the individual being treated (to ensure that the stages of the disease in the treated individual and the control individual(s) are comparable).

[0369] The individual (also referred to as “patient” or “subject”) being treated is an individual (fetus, infant, child, adolescent, or adult human) having the disease or having the potential to develop the disease. The individual can have residual endogenous lysosomal enzyme expression and/or activity, or no measurable activity. For example, the individual having Sanfilippo syndrome type A may have HNS expression levels that are less than about 30-50%, less than about 25-30%, less than about 20-25%, less than about 15-20%, less than about 10-15%, less than about 5-10%, less than about 0.1-5% of normal HNS expression levels.

Immune Tolerance

[0370] Generally, intrathecal administration of a therapeutic agent (e.g., a replacement enzyme) according to the present invention does not result in severe adverse effects in the subject. As used herein, severe adverse effects induce, but are not limited to, substantial immune response, toxicity, or death. As used herein, the term “substantial immune response” refers to severe or serious immune responses, such as adaptive T-cell immune responses.

[0371] Thus, in many embodiments, inventive methods according to the present invention do not involve concurrent immunosuppressant therapy (i.e., any immunosuppressant therapy used as pre-treatment/pre-conditioning or in parallel to the method). In some embodiments, inventive methods according to the present invention do not involve an immune tolerance induction in the subject being treated. In some embodiments, inventive methods according to the present invention do not involve a pre-treatment or preconditioning of the subject using T-cell immunosuppressive agent.

[0372] In some embodiments, intrathecal administration of therapeutic agents can mount an immune response against these agents. Thus, in some embodiments, it may be useful to render the subject receiving the replacement enzyme tolerant to the enzyme replacement therapy.

2017202349 10 Apr 2017

Immune tolerance may be induced using various methods known in the art. For example, an initial 30-60 day regimen of a T-cell immunosuppressive agent such as cyclosporin A (CsA) and an antiproliferative agent, such as, azathioprine (Aza), combined with weekly intrathecal infusions of low doses of a desired replacement enzyme may be used.

[0373] Any immunosuppressant agent known to the skilled artisan may be employed together with a combination therapy of the invention. Such immunosuppressant agents include but are not limited to cyclosporine, FK506, rapamycin, CTLA4-Ig, and anti-TNF agents such as etanercept (see e.g. Moder, 2000, *Ann. Allergy Asthma Immunol.* 84, 280-284; Nevins, 2000, *Curr. Opin. Pediatr.* 12, 146-150; Kurlberg et al., 2000, *Scand. J. Immunol.* 51, 224-230; Ideguchi et al., 2000, *Neuroscience* 95, 217-226; Potteret et al., 1999, *Ann. N.Y. Acad. Sci.* 875, 159-174; Slavik et al., 1999, *Immunol. Res.* 19, 1-24; Gaziev et al., 1999, *Bone Marrow Transplant.* 25, 689-696; Henry, 1999, *Clin. Transplant.* 13, 209-220; Gummert et al., 1999, *J. Am. Soc. Nephrol.* 10, 1366-1380; Qi et al., 2000, *Transplantation* 69, 1275-1283). The anti-IL2 receptor (.alpha.-subunit) antibody daclizumab (e.g. Zenapax.TM.), which has been demonstrated effective in transplant patients, can also be used as an immunosuppressant agent (see e.g. Wiseman et al., 1999, *Drugs* 58, 1029-1042; Beniaminovitz et al., 2000, *N. Engl. J. Med.* 342, 613-619; Ponticelli et al., 1999, *Drugs R. D.* 1, 55-60; Berard et al., 1999, *Pharmacotherapy* 19, 1127-1137; Eckhoff et al., 2000, *Transplantation* 69, 1867-1872; Ekberg et al., 2000, *Transpl. Int.* 13, 151-159). Additional immunosuppressant agents include but are not limited to anti-CD2 (Branco et al., 1999, *Transplantation* 68, 1588-1596; Przepiorcka et al., 1998, *Blood* 92, 4066-4071), anti-CD4 (Marinova-Mutafchieva et al., 2000, *Arthritis Rheum.* 43, 638-644; Fishwild et al., 1999, *Clin. Immunol.* 92, 138-152), and anti-CD40 ligand (Hong et al., 2000, *Semin. Nephrol.* 20, 108-125; Chirmule et al., 2000, *J. Virol.* 74, 3345-3352; Ito et al., 2000, *J. Immunol.* 164, 1230-1235).

Administration

[0374] Inventive methods of the present invention contemplate single as well as multiple administrations of a therapeutically effective amount of the therapeutic agents (e.g., replacement enzymes) described herein. Therapeutic agents (e.g., replacement enzymes) can be administered

2017202349 10 Apr 2017

at regular intervals, depending on the nature, severity and extent of the subject's condition (e.g., a lysosomal storage disease). In some embodiments, a therapeutically effective amount of the therapeutic agents (e.g., replacement enzymes) of the present invention may be administered intrathecally periodically at regular intervals (e.g., once every year, once every six months, once every five months, once every three months, bimonthly (once every two months), monthly (once every month), biweekly (once every two weeks), weekly).

[0375] In some embodiments, intrathecal administration may be used in conjunction with other routes of administration (e.g., intravenous, subcutaneously, intramuscularly, parenterally, transdermally, or transmucosally (e.g., orally or nasally)). In some embodiments, those other routes of administration (e.g., intravenous administration) may be performed no more frequent than biweekly, monthly, once every two months, once every three months, once every four months, once every five months, once every six months, annually administration.

[0376] As used herein, the term "therapeutically effective amount" is largely determined base on the total amount of the therapeutic agent contained in the pharmaceutical compositions of the present invention. Generally, a therapeutically effective amount is sufficient to achieve a meaningful benefit to the subject (e.g., treating, modulating, curing, preventing and/or ameliorating the underlying disease or condition). For example, a therapeutically effective amount may be an amount sufficient to achieve a desired therapeutic and/or prophylactic effect, such as an amount sufficient to modulate lysosomal enzyme receptors or their activity to thereby treat such lysosomal storage disease or the symptoms thereof (e.g., a reduction in or elimination of the presence or incidence of "zebra bodies" or cellular vacuolization following the administration of the compositions of the present invention to a subject). Generally, the amount of a therapeutic agent (e.g., a recombinant lysosomal enzyme) administered to a subject in need thereof will depend upon the characteristics of the subject. Such characteristics include the condition, disease severity, general health, age, sex and body weight of the subject. One of ordinary skill in the art will be readily able to determine appropriate dosages depending on these and other related factors. In addition, both objective and subjective assays may optionally be employed to identify optimal dosage ranges.

2017202349 10 Apr 2017

[0377] A therapeutically effective amount is commonly administered in a dosing regimen that may comprise multiple unit doses. For any particular therapeutic protein, a therapeutically effective amount (and/or an appropriate unit dose within an effective dosing regimen) may vary, for example, depending on route of administration, on combination with other pharmaceutical agents. Also, the specific therapeutically effective amount (and/or unit dose) for any particular patient may depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific pharmaceutical agent employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and/or rate of excretion or metabolism of the specific fusion protein employed; the duration of the treatment; and like factors as is well known in the medical arts.

[0378] In some embodiments, the therapeutically effective dose ranges from about 0.005 mg/kg brain weight to 500 mg/kg brain weight, e.g., from about 0.005 mg/kg brain weight to 400 mg/kg brain weight, from about 0.005 mg/kg brain weight to 300 mg/kg brain weight, from about 0.005 mg/kg brain weight to 200 mg/kg brain weight, from about 0.005 mg/kg brain weight to 100 mg/kg brain weight, from about 0.005 mg/kg brain weight to 90 mg/kg brain weight, from about 0.005 mg/kg brain weight to 80 mg/kg brain weight, from about 0.005 mg/kg brain weight to 70 mg/kg brain weight, from about 0.005 mg/kg brain weight to 60 mg/kg brain weight, from about 0.005 mg/kg brain weight to 50 mg/kg brain weight, from about 0.005 mg/kg brain weight to 40 mg/kg brain weight, from about 0.005 mg/kg brain weight to 30 mg/kg brain weight, from about 0.005 mg/kg brain weight to 25 mg/kg brain weight, from about 0.005 mg/kg brain weight to 20 mg/kg brain weight, from about 0.005 mg/kg brain weight to 15 mg/kg brain weight, from about 0.005 mg/kg brain weight to 10 mg/kg brain weight.

[0379] In some embodiments, the therapeutically effective dose is greater than about 0.1 mg/kg brain weight, greater than about 0.5 mg/kg brain weight, greater than about 1.0 mg/kg brain weight, greater than about 3 mg/kg brain weight, greater than about 5 mg/kg brain weight, greater than about 10 mg/kg brain weight, greater than about 15 mg/kg brain weight, greater than about 20 mg/kg brain weight, greater than about 30 mg/kg brain weight, greater than about 40 mg/kg brain weight, greater than about 50 mg/kg brain weight, greater than about 60 mg/kg brain weight, greater than about 70 mg/kg brain weight, greater than about 80 mg/kg brain weight,

2017202349 10 Apr 2017

greater than about 90 mg/kg brain weight, greater than about 100 mg/kg brain weight, greater than about 150 mg/kg brain weight, greater than about 200 mg/kg brain weight, greater than about 250 mg/kg brain weight, greater than about 300 mg/kg brain weight, greater than about 350 mg/kg brain weight, greater than about 400 mg/kg brain weight, greater than about 450 mg/kg brain weight, greater than about 500 mg/kg brain weight.

[0380] In some embodiments, the therapeutically effective dose may also be defined by mg/kg body weight. As one skilled in the art would appreciate, the brain weights and body weights can be correlated. Dekaban AS. "Changes in brain weights during the span of human life: relation of brain weights to body heights and body weights," *Ann Neurol* 1978; 4:345-56. Thus, in some embodiments, the dosages can be converted as shown in Table 4.

Table 4 Dosage conversion

Correlation between Brain Weights, body weights and ages of males

Age (year)	Brain weight (kg)	Body weight (kg)
3 (31-43 months)	1.27	15.55
4-5	1.30	19.46

[0381] In some embodiments, the therapeutically effective dose may also be defined by mg/15 cc of CSF. As one skilled in the art would appreciate, therapeutically effective doses based on brain weights and body weights can be converted to mg/15 cc of CSF. For example, the volume of CSF in adult humans is approximately 150 mL (Johanson CE, et al. "Multiplicity of cerebrospinal fluid functions: New challenges in health and disease," *Cerebrospinal Fluid Res.* 2008 May 14;5:10). Therefore, single dose injections of 0.1 mg to 50 mg protein to adults would be approximately 0.01 mg/15 cc of CSF (0.1 mg) to 5.0 mg/15 cc of CSF (50 mg) doses in adults.

[0382] It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the enzyme replacement therapy and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed invention.

2017202349 10 Apr 2017

Kits

[0383] The present invention further provides kits or other articles of manufacture which contains the formulation of the present invention and provides instructions for its reconstitution (if lyophilized) and/or use. Kits or other articles of manufacture may include a container, an IDDD, a catheter and any other articles, devices or equipment useful in interthecal administration and associated surgery. Suitable containers include, for example, bottles, vials, syringes (e.g., pre-filled syringes), ampules, cartridges, reservoirs, or lyo-jects. The container may be formed from a variety of materials such as glass or plastic. In some embodiments, a container is a pre-filled syringe. Suitable pre-filled syringes include, but are not limited to, borosilicate glass syringes with baked silicone coating, borosilicate glass syringes with sprayed silicone, or plastic resin syringes without silicone.

[0384] Typically, the container may hold formulations and a label on, or associated with, the container that may indicate directions for reconstitution and/or use. For example, the label may indicate that the formulation is reconstituted to protein concentrations as described above. The label may further indicate that the formulation is useful or intended for, for example, IT administration. In some embodiments, a container may contain a single dose of a stable formulation containing a therapeutic agent (e.g., a replacement enzyme). In various embodiments, a single dose of the stable formulation is present in a volume of less than about 15 ml, 10 ml, 5.0 ml, 4.0 ml, 3.5 ml, 3.0 ml, 2.5 ml, 2.0 ml, 1.5 ml, 1.0 ml, or 0.5 ml. Alternatively, a container holding the formulation may be a multi-use vial, which allows for repeat administrations (e.g., from 2-6 administrations) of the formulation. Kits or other articles of manufacture may further include a second container comprising a suitable diluent (e.g., BWFI, saline, buffered saline). Upon mixing of the diluent and the formulation, the final protein concentration in the reconstituted formulation will generally be at least 1 mg/ml (e.g., at least 5 mg/ml, at least 10 mg/ml, at least 25 mg/ml, at least 50 mg/ml, at least 75 mg/ml, at least 100 mg/ml). Kits or other articles of manufacture may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, IDDDs, catheters, syringes, and package inserts with instructions for use.

2017202349 10 Apr 2017

[0385] The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the invention. All literature citations are incorporated by reference.

EXAMPLES

Examples of IT Delivery of GalC Protein

EXAMPLE 1: PHYSIOCHEMICAL CHARACTERIZATION OF GALC FORMULATION FOR INTRATHECAL DELIVERY

[0386] The present Example describes physiochemical characterization of GalC in order to understand its behavior and stability under different solution conditions during intrathecal (IT) delivery of the protein.

[0387] Among other things, the present Example describes a GalC formulation which is important for successful IT delivery of GalC. In some embodiments, this formulation includes 5 mM Na phosphate + 150 mM NaCl, pH 6.0 + 0.005% polysorbate 20. In some embodiments, this formulation includes <5 mM, <10 mM, <15 mM and <20 mM Na phosphate. In some embodiments, this formulation includes a pH \geq 5.5 and \leq pH 7.0, with 150 mM NaCl.

[0388] PBS delivery vehicles of varying phosphate molarity and pH were tested in adult cynomolgous monkeys (Figure 3). 5 mM phosphate in a pH range of 5.5-7.0 showed no adverse effect whereas 20 mM phosphate between pH 7.0-7.5 and 10-20 mM phosphate between pH 7.5-8.0 showed an adverse effect in the monkeys (Figure 3). Thermal stability of hGalC (1mg/ml) in 3 mM citrate, phosphate and borate buffer with 50 mM NaCl, was investigated as a function of pH within the range of pH 5.0-8.0 (Figure 4). hGalC specific activity was measured at baseline (20-25°C) and at 2 weeks at 40°C with the highest specific activity retained between pH 6.0-6.5 (Figure 4). hGalC specific activity was additionally measured at 3 months at 5°C with the highest specific activity retained between pH 6.0-6.5 (Figure 5). The melting temperature of hGalC was measured as a function of pH (Table 5) and also measured independently in different formulations (Table 6).

2017202349 10 Apr 2017

Table 5: Melting Temperature of hGalC (1mg/mL) as a Function of pH

<u>pH of</u>	<u>Tm (°C)</u>
<u>Universal</u>	
<u>Buffer</u>	
4.5*	61.6
5.0*	63.0
6.0	60.8
6.5	58.9
7.0	57.3
7.5	56.5

*[GalC]<1 mg/mL due to precipitation

Table 6: Melting Temperature of hGalC (1mg/mL) in Different Formulations

<u>Formulation (pH 6.0)</u>	<u>Tm (°C)</u>
5 mM phosphate, 50 mM NaCl	61.6
5 mM phosphate, 150 mM NaCl	60.2
5 mM phosphate, 500 mM NaCl	59.5
5 mM phosphate, 5% Dextrose	63.8
5 mM phosphate, 150 mM NaCl, 1%NaTC	56.8

[0389] Thermal stability of hGalC, as determined by retention of hGalC specific activity at ~3 weeks at 5°C and 2 weeks at 40°C, was also evaluated as a function of salt concentration (Figure 6). Results showed that hGalC retained high specific activity after 3 weeks at 5°C in a variety of salt concentrations ranging from 5 mM phosphate + 50 mM NaCl (abbreviated herein as 5+50) to 50 mM phosphate + 150 mM NaCl (abbreviated herein as 50+150), at pH 6.5 (Figure 6).

2017202349 10 Apr 2017

Sedimentation Analysis of hGalC

[0390] Sedimentation velocity is an analytical ultracentrifugation (AUC) method that measures the rate at which molecules move in response to centrifugal forces generated in a centrifuge and is a useful technique for determine protein association state in solution. The first sedimentation velocity run was a dilution series of human GalC in 5 mM Na phosphate, pH 6.0 with 150 mM NaCl (Figure 7) to assess the sample for self-association and/or nonideality. The dilution series was plotted as normalized $g(s^*)$ curves ($g(s^*)$ vs s^*) at each concentration. The general shift in the curves to lower s values upon dilution indicates dissociation, and this is a rapidly reversible self-associating system. Comparing different ionic strengths (Figure 7A, B & C), it is apparent that the sets of curves shift to lower s values upon raising the ionic strength indicating that ionic interactions are also involved in the association process and that the self association is decreased at higher salt concentrations.

[0391] The mouse GalC was also run at the same time at 150 mM NaCl to compare with hGalC. Comparing corresponding ionic strengths (150 mM NaCl), it is apparent that the free energy of self-association of mGalC is less than that of hGalC. The curves in Figure 7 were cut off at about 20S to show the dissociation more clearly; however, when these runs are analyzed using the wide distribution analysis (WDA) and the results are plotted on a log scale, higher aggregates ($s^*>20S$) can clearly be seen. The aggregation to high oligomers (Figure 8) is especially visible at 50 mM NaCl, somewhat decreased in 10 mM NaCl and significantly reduced, but present, in 500 mM NaCl at pH 6.0. The WDA curve from the highest concentration from each of the ionic strengths is plotted in Figure 8.

Self-association in universal buffer at pH 6.0

[0392] Under these conditions in the universal buffer, the self association appears to be of about the same magnitude as in the phosphate buffer, pH 6.0, as seen in Figure 9. The effect of pH on the energetics of hGalC self-association in universal buffer was also investigated. Dilution series were performed at pH 4.5, 5.0, 6.0, 6.5, 7.0 and 7.5. The samples at pH 4.5 and 5.0 were insoluble with essentially 100% of the hGalC having precipitated leaving nothing to measure in the supernatant.

2017202349 10 Apr 2017

[0393] The effect of pH is clearly shown in Figure 10 where the least amount of self-association is observed at pH 7.5 and considerable self-association is observed at pH 6.0. The trend is similar to that seen with variations in ionic strength with higher pH. Increasing both ionic strength and pH shifts the equilibrium to favor the smaller oligomers at the highest concentration (all about 1.0 mg/mL). Decrease in concentration by 1/3 serial dilutions (see Figure 7) shifts the equilibrium toward the smallest species which appears to have a sedimentation coefficient of about 5.2S. The peak that occurs at about 10-13S likely represents a tetramer of the 5S species. Efforts to fit these data to a self-association model have so far been unsuccessful and is likely due to the inherent micro-heterogeneity arising from variable degrees of glycosylation.

Self-association in universal buffer at pH 6.0

[0394] The stressed and baseline samples of GalC in 5mM Na phosphate, pH 6.0, with 150 mM NaCl were compared in a dilution series experiment (red→blue→green→black). The results for the lowest concentration (black) ~0.03mg/mL have been smoothed which is why the curve seems to have less noise. In the stressed sample there is an aggregate around $\ln(s^*) = 3.0$ (~20S) that is present in much higher concentration than in the baseline sample. It represents a nearly constant fraction of the sample as evidenced by its persistence upon dilution in the normalized plots (Figure 11, Figure 12, Figure 13). It is therefore an irreversible aggregate with a molar mass of at least 500 kg/mol.

hGalC with sodium taurocholate in solution

[0395] In sodium taurocholate (NaTC)(1%), the self association is significantly reduced. The main boundary is shifted to lower s values and the higher oligomerization is suppressed (Figure 14).

hGalC with 5% dextrose

[0396] The addition of 5% dextrose to GalC in 5mM Na phosphate, pH 6.0 resulted in the formation of large aggregates (Figure 15). The peak at 18S corresponds to a minimum molar mass of about 440 kDa and the peak at 56S corresponds to a minimum molar mass of 2.4 MDa

2017202349 10 Apr 2017

with a tail extending beyond 150S, corresponding to molar masses greater than 10.0 MDa. There is very little change in this pattern upon dilution from 1.0 to 0.3 mg/mL indicating that these oligomers are mostly irreversible on the time scale of the sedimentation experiment, a period of 5-6 hours

hGalC intrinsic fluorescence

[0397] Intrinsic fluorescence studies of hGalC (using 23 Trp) were performed to evaluate the role of pH and salt concentration on molecular interactions (Figure 16 and Figure 17). Molecular interactions were the least (highest relative fluorescence between 330nm-350nm) in either 500mM NaCl or 1% NaTC (Figure 16). A small change in the secondary structure was observed as a function of pH. Precipitation was observed at pH 4.5 and 5.0 (Figure 17).

Summary

[0398] To evaluate the relative solubility of hGalC and mGalC, a polyethylene glycol (PEG)-induced solid phase approach was used (Middaugh et al., J. Biol. Chem. 1979, 254, 367-370). This approach allows for the relative solubility of proteins to be measured in a quantifiable manner. Solubility measurements were performed by introducing buffered solutions (5 mM sodium phosphate with 150 mM NaCl, pH 6.0) of each GalC to the different concentrations of PEG (10kDa). Plots of log protein solubility vs. PEG concentrations produced a linear trend. Extrapolation of the apparent solubility to zero PEG concentration was made to obtain the relative solubility of each protein. Relative solubility of the mGalC vs. hGalC did not show any difference. In solubility experiments of hGalC, no precipitation or loss of activity was observed after 3 weeks at 2-8°C (in 5mM sodium phosphate with different salt concentrations, pH 6.0-6.5). Solubility at ~30 mg/mL was achieved with the formulation 5 mM Na phosphate + 150 mM NaCl, pH 6.0, and no precipitation was observed after 50 days at 2-8°C.

[0399] The AUC data suggest that the “native” state of GalC is a concentration dependent reversible association to higher order oligomers. The biophysical data suggest that there may be a functional and structural importance to the higher order oligomers. At higher pH values, there is less retention of activity, lower T_m values and a more homogenous system as determined by AUC. In 5 mM sodium phosphate with 150 mM NaCl, pH 6.0, there is likely an

2017202349 10 Apr 2017

equilibrium between monomer, tetramer and other higher order species. Furthermore, pH does not dramatically affect the AUC profiles in the pH range of 6.5-7.5. Overall, the GalC system is a rapidly reversible, highly self-associating system in the tested buffers.

EXAMPLE 2: PHARMACOKINETICS AND TISSUE DISTRIBUTION OF RADIOACTIVITY IN SPRAGUE-DAWLEY RATS FOLLOWING A SINGLE INTRATHECAL DOSE OR A SINGLE INTRAVENOUS BOLUS INJECTION OF ¹²⁵I-HGALC

[0400] The present Example depicts an exemplary result illustrating pharmacokinetics and tissue distribution of ¹²⁵I-hGALC in male Sprague-Dawley rats following a single intrathecal dose or a single intravenous bolus injection. The concentration and content of radioactivity in whole blood, serum, red blood cells, cerebrospinal fluid (CSF) and tissues were measured and non-compartmental pharmacokinetic analyses were performed on the resulting data. The intrathecal and intravenous routes were selected as they are the intended routes of administration in humans. The dose levels were selected based on potential human exposure, existing toxicity and pharmacokinetic data and any limitations imposed by the test article. The rat was selected for the study because it is an accepted species for use in pharmacokinetic and tissue distribution studies. The number of animals used in this study was the minimum needed to adequately assess the expected variability at each time point and meet the experimental objectives.

Materials and Methods

Test System

[0401] 82 male Sprague-Dawley rats (*Rattus norvegicus*) were received from Charles River Canada Inc. (St. Constant, Quebec, Canada) on 15 April 2009. At the onset of treatment, the animals were approximately 10-11 weeks old. A further 9 male rats were received from Charles River Canada on 28 April 2009; these animals were approximately 9 weeks old on arrival and were required to ensure that sufficient cannulated animals were available in order to complete dosing of the study.

2017202349 10 Apr 2017

[0402] The bodyweights of the male rats ranged from 342 to 453 g at the onset of treatment. The body weights of all but one of the male rats on dosing were higher than the range stated in the protocol (250-350 g), however this minor deviation was not considered to have affected the study or the data obtained since the animals were healthy and the actual body weight was used for dose administration.

Animal Management

[0403] Following arrival at PCS-MTL, all animals were subjected to a general physical examination by a qualified member of the veterinary staff. No significant abnormalities were detected in the animals received. Animals were housed individually in stainless steel cages with a wire-mesh bottomed floor and an automatic watering valve. The environmental enrichment program was in accordance with the appropriate SOP. Each cage was clearly labelled with a colour-coded cage card indicating study, group, animal numbers and sex. Each animal was uniquely identified using the AIMS[®] tattoo system. Environmental conditions during the study conduct were controlled at a target temperature and relative humidity of 19 to 25°C and 30 to 70%, respectively. The photoperiod was 12 hours light and 12 hours dark except when interrupted due to scheduled activities.

Diet

[0404] All animals had free access to a standard certified pelleted commercial laboratory diet (PMI Certified Rodent Diet 5002: PMI Nutrition International Inc.) except during designated procedures. Maximum allowable concentrations of contaminants in the diet (e.g., heavy metals, aflatoxin, organophosphate, chlorinated hydrocarbons, PCBs) are controlled and routinely analyzed by the manufacturers. Municipal tap water, suitable for human consumption (filtered through a 0.5 µm bacteriostatic polycarbonate filter) was available to the animals ad libitum except during designated procedures. It was considered that there were no known contaminants in the dietary materials that could interfere with the objectives of the study.

Acclimation and Randomization

2017202349 10 Apr 2017

[0405] At least 6 days (for animals received on 15 April 2009) or 3 days (for the 9 additional animals received on 28 April 2009) were allowed between the receipt of the animals and surgery to place the intrathecal cannula, to allow the animals to become acclimated to the physical and environmental conditions. During the acclimation period, all animals were weighed and randomized, using a computer-based randomization procedure. Randomization was performed following stratification using body weight as the parameter. Animals at the extremes of the body weight range were not assigned to groups.

[0406] The animals were assigned to the study groups as follows:

Table 7

Group	Route of Administration and Dose		Projected Dose Volume		Animal Numbers
	Intravenous	Intrathecal	Intravenous (mL/kg)	Intrathecal (mL)	
1	-	60 µg	-	0.02	1001-1024
2	1 mg/kg	-	3.33	-	2001-2024
3 ^a	1 mg/kg	60 µg	3.33	0.02	3001-3024

a: The IV dose was administered within 5 minutes after the intrathecal dose.

Each rat in Groups 1 and 2 received a nominal radiochemical dose of approximately 3 µCi/animal. Each rat in Group 3 received a nominal radiochemical dose of approximately 6 µCi/animal.

Intrathecal dose formulation

[0407] The intrathecal dose formulation was prepared on the day of first administration of the intrathecal dose. Sufficient ¹²⁵I-hGALC solution was measured and added to sufficient measured unlabelled hGALC solution. A measured volume of vehicle was added and the whole mixed gently. A solution of concentration 3 mg/mL at a target radioactivity level of

2017202349 10 Apr 2017

approximately 150 $\mu\text{Ci}/\text{mL}$ was prepared. The resulting formulation was filtered through a low protein binding filter (0.22 μm GV PVDF filter unit) into a sterile vessel and kept refrigerated (2-8°C), protected from light, pending use for dosing.

Intravenous dose formulation

[0408] The intravenous dose formulation was prepared on the day of first administration of the intravenous dose. Sufficient ^{125}I -hGALC solution was measured and added to sufficient measured unlabelled hGALC solution. A measured volume of vehicle was added and the whole mixed gently. A solution of concentration 0.3 mg/mL at a target radioactivity level of approximately 3 $\mu\text{Ci}/\text{mL}$ was prepared. The resulting formulation was filtered through a low protein binding filter (0.22 μm GV PVDF filter unit) into a sterile vessel and kept refrigerated (2-8°C), protected from light, pending use for dosing.

Analysis of the Dose Formulations

[0409] Each radiolabelled dose formulation was analyzed at PCS-MTL on each day of dosing by liquid scintillation spectroscopy to determine the radioactivity concentration before and after treatment. The radioactivity concentration was determined by preparing appropriate dilutions of the dose formulation in vehicle and duplicate aliquots of each dilution were analyzed. The remaining dose formulations were discarded following completion of analysis (including repeat analysis).

Calculation of Specific Activity of Test Article

[0410] The specific activity of the test article in the dose formulations was calculated from the mean (pre and post dose) measured levels of radioactivity and the total mass of test article (based on the concentrations provided) in the dose formulations.

Clinical Observations

[0411] All animals were examined twice daily for mortality and signs of ill health and reaction to treatment throughout the acclimation and study periods, except on the days of arrival

2017202349 10 Apr 2017

and termination of the study, on which days the animals were only examined once. A detailed examination was performed weekly.

Body Weight

[0412] Individual body weights were measured once during acclimation, before surgery and on the day prior to dose administration. Only the body weights recorded on the day prior to dose administration were reported.

Surgery

[0413] A minimum of 6 days (or 3 days for the 9 additional animals) was allowed between the receipt of the animals and the surgery to allow the animals to become accustomed to the laboratory environmental conditions. All animals, including the spares, received a single intramuscular injection of Benzathine Penicillin G + Procaine Penicillin G antibiotic on the day of surgery and again 2 days following surgery. In general, Buprenorphine 0.05 mg/kg was administered subcutaneously prior to surgery and approximately 8 hours post first administration, and as deemed necessary thereafter. For some animals, Buprenorphine was administered approximately 6 hours post first administration instead of 8-12 hours. Considering the half-life of Buprenorphine in rats, this deviation from the protocol did not affect the health of these animals, and thus had not impact on the validity or data obtained in the study.

[0414] The animals were prepared for surgery by shaving from the cranium to the dorso-thoracic region of the neck. The animals were anesthetized with isoflurane/oxygen gas prior to surgery and maintained under isoflurane gas anesthesia throughout the surgical procedure. Prior to surgery, and at the end of the surgical procedure, while under anesthesia, a bland lubricating ophthalmic agent was administered to each eye. Prior to the surgery, and on 2 other occasions at approximately 24-hour intervals following the first administration, each animal received an anti-inflammatory (Carprofen at 5 mg/kg) by subcutaneous injection.

[0415] The animal was positioned within the stereotaxic table. A skin incision, of approximately 2 cm, was made from the caudal edge of the cranium to the neck. The dorsal neck muscles were separated in order to expose the atlanto-occipital membrane. A retractor was used

2017202349 10 Apr 2017

to facilitate access to the membrane. The atlanto-occipital membrane was incised and the intrathecal catheter was slowly inserted caudally until the catheter was located in the lumbar region. Excess fluid was removed using cotton-tipped swabs and the atlanto-occipital membrane was dried. Immediately thereafter, adhesive was used to anchor the catheter bulb to the membrane. Once the glue had dried and the catheter was solidly anchored, the retractors were removed. A small loop was made with the catheter on the cranium and the bulb was attached using a suture of non-absorbable material. Once the catheter was secured, it was passed to the dorsal thoracic region where an incision was made to place an access port. This was sutured in place using non-absorbable material.

[0416] Prior to closing the neck muscles, a 2 mL flush of warm saline (i.e.: approximately 37.5°C) was made in the wound. The muscles were closed using simple interrupted sutures of absorbable material. The access port site was flushed with 2 mL of warm saline and the skin was closed using a continuous subcuticular suture of absorbable suture material. A topical antibiotic ointment was administered to surgical sites post-surgery and once daily thereafter until considered unnecessary.

[0417] The dead volume of the catheter and access port was determined at the time of surgery. A patency check was performed once during the pre-treatment period between the surgery day and the treatment day.

Treatment

[0418] A period of at least 7 days was allowed between the surgical implantation of the catheter/access port and treatment initiation to allow for adequate recovery. Prior to intrathecal dosing, the access port area was shaved, if necessary. The puncture site was cleaned using chlorhexidine gluconate and water, and the site wiped with soaked gauze of sterile water followed by 3 passages of povidone iodine 10%. The access port was punctured with a needle connected to the dosing syringe and the test article was administered slowly. After dosing, the site was wiped with iodine in order to limit contamination.

[0419] On Day 1 of the study, Group 1 animals were administered the formulated ¹²⁵I-hGALC by slow bolus intrathecal injection into the subcutaneous lumbar access port

2017202349 10 Apr 2017

followed by a saline flush of 0.04 mL to deliver a target dose level of 60 µg/animal and a radioactivity dose of approximately 3 µCi/animal.

[0420] On Day 2 of the study, Group 3 animals were administered formulated ¹²⁵I-hGALC by slow bolus intrathecal injection into the subcutaneous lumbar access port followed by a saline flush of 0.04 mL to deliver a target dose level of 60 µg/animal and a radioactivity dose of approximately 3 µCi/animal. Within 5 minutes of the slow bolus intrathecal injection, Group 3 animals also received an intravenous injection via an intravenous catheter into the tail vein (3.33 mL/kg) followed by a 0.6 mL saline flush to deliver a target dose level of 1 mg/kg, with an approximate radioactivity level of 3 µCi/animal.

[0421] On Day 3 of the study, Group 2 animals were administered formulated ¹²⁵I-hGALC by intravenous injection via an intravenous catheter into the tail vein (3.33 mL/kg) followed by a 0.6 mL saline flush to deliver a target dose level of 1 mg/kg animal and a radioactivity dose of approximately 3 µCi/animal.

[0422] The volume administered was based on the most recent practical body weight of each animal. The weights of the syringes filled with formulated ¹²⁵I-hGALC and empty after delivery to the animals were recorded. The dose delivered to each animal was calculated on the basis of the net weight of dosage formulation expelled from the syringe and the measured radioactivity concentration in the formulated dose.

[0423] During dosing, gauzes were available to absorb any small amounts of reflux of dose formulation and the test article loss was accounted for by liquid scintillation counting according to a project specific procedure. The syringes and intravenous catheters used for administration of formulated test article were retained. The intravenous catheters and selected intrathecal access port/catheters were analyzed for the level of radioactivity according to a project specific procedure.

2017202349 10 Apr 2017

*Sample Collection**Blood/Serum and Tissues*

[0424] A terminal blood sample (maximum possible volume) was collected at 10 minutes, 30 minutes and 1, 3, 6, 24, 48 and 96 h post dose from 3 animals/time point for Groups 1 to 3. The intrathecal administration preceded the intravenous administration in Group 3, and the timing for the terminal blood sample was based on the time of the intravenous administration. Terminal blood samples were collected from the abdominal aorta of rats (Groups 1, 2 and 3, and 3 spare animals) euthanized under isoflurane anesthesia by exsanguination from the abdominal aorta. Approximately 3 mL of blood (Groups 1, 2 and 3) was transferred to a suitable tube containing K₃-EDTA, to furnish whole blood samples and was kept on wet ice pending processing. For Groups 2 and 3, and the spare animals, an additional 1.5 mL of blood was transferred into tubes containing sodium citrate for analysis of prothrombin time (PTT), activated partial thromboplastin time (APTT) and fibrinogen. Blood samples were stored on wet ice, pending centrifugation at 2700 RPM and 4°C for 15 minutes. Plasma samples were stored frozen at approximately -80°C, before shipment and analysis at a laboratory designated by the Applicant. Plasma from the spare animals was to serve as blank samples for the analysis of PTT, APTT and fibrinogen. Where insufficient blood volume was obtained to perform all analyses (Groups 1, 2 and 3), then blood for radioactivity analysis had the priority.

[0425] The remaining blood (Groups 1, 2 and 3, and 3 spare animals) was transferred into tubes containing clotting activator for serum production and was allowed to clot, at room temperature, over a period of approximately 30 minutes before centrifugation. The samples collected from the spare animals were used to assess the clotting of blood samples from non-treated animals.

[0426] Following exsanguination, the following tissues were collected from 3 animals/time point from Groups 1 to 3, as indicated: Adipose tissue (kidney fat), Adrenal glands, Bone (femur), Brain, Eyes, Heart, Kidneys, Large intestine, Large intestine content, Liver, Lungs, Muscle (skeletal), Sciatic nerve, Small intestine, Small intestine content, Spinal cord

2017202349 10 Apr 2017

(lumbar, thoracic, cervical), Spleen, Stomach, Stomach content, Thyroid/parathyroid gland, Urinary bladder content.

[0427] Upon collection, tissues were weighed and then processed and analyzed for total radioactivity. All tissues mentioned above, as well as terminal blood and serum, were also collected from a spare animal and were used to determine background levels of radioactivity. The remaining carcasses were kept frozen (-10°C to -20°C) in the designated freezer in order to allow for radioactive decay before being disposed as biological waste. The carcass of the first animal at each time point from Groups 1 and 3 were retrieved from the freezer, thawed and the access port and catheter removed, flushed with water and verified for residual radioactivity.

Cerebrospinal Fluid

[0428] Cerebrospinal fluid (CSF) samples were collected from all animals at necropsy immediately before euthanasia. Three animals/time-point from Groups 1 to 3 were euthanized at 10 minutes, 30 minutes and 1, 3, 6, 24, 48 and 96 h post dose. A sample (maximum possible volume) of CSF was removed via the cisterna magna, using a stereotaxic table were necessary to hold the head in alignment. CSF was transferred into a plain tube and placed on wet ice. A portion (approximately 20 µL) was processed and analyzed for total radioactivity content. CSF was also collected from a spare animal and was used to determine background levels of radioactivity.

Determination of Background Radioactivity Levels

[0429] The blood, serum and tissues collected from the spare animal, were used for the determination of background radioactivity levels for blood, serum and tissues of animals in Groups 1, 2 and 3. The CSF collected from the spare animal, was used for the determination of background radioactivity levels for CSF.

Sample Processing for Radioactivity Measurements

[0430] All samples were weighed following collection, except for blood, plasma, serum and CSF. For all groups, duplicate 100 µL weighed aliquots of whole blood collected on K₃-EDTA, were taken for analysis of radioactivity. Protein precipitation using trichloroacetic

2017202349 10 Apr 2017

acid (TCA) of whole blood was performed as follows: an equivalent volume of a 15% aqueous solution of TCA was added to duplicate 100 μ L weighed aliquots of whole blood. Samples (100 μ L whole blood + 100 μ L TCA) were mixed by vortexing and then centrifuged at 4°C for approximately 15 minutes at 10000 rpm, and the supernatant decanted into a separate tube. Both the supernatant and the pellet were analyzed for radioactivity content.

[0431] The blood for serum collection was kept at room temperature for approximately 30 minutes, to allow for clotting, before being centrifuged at 4°C at 2700 rpm (1250 rcf) for approximately 10 minutes to separate serum. Serum samples were then kept on wet ice pending aliquotting for radioactivity analysis (2 x 100 μ L weighed aliquots). The packed red blood cells (obtained after serum separation) were kept on wet ice pending processing for radioactivity analysis. Remaining serum was stored frozen (-10°C to -20°C). Duplicate 100 μ L weighed aliquots of whole blood and red blood cells (obtained after serum separation, mixed with an equal volume of deionized water (w/v) and homogenized with a Polytron emulsifier) were solubilized in Soluene-350, decolorized with hydrogen peroxide (30% w/v), and mixed with liquid scintillation fluid for analysis of radioactivity.

[0432] The TCA blood precipitate pellet was solubilized in 35% tetraethylammonium hydroxide (TEAH), decolorized with hydrogen peroxide (30% w/v), and mixed with liquid scintillation fluid for radioactivity measurement. Urinary bladder contents, TCA blood supernatant, duplicate weighed aliquots of dose formulations (diluted) and serum were mixed directly with liquid scintillation fluid for radioactivity measurement. Duplicate weighed aliquots of CSF (approximately 10 μ L/aliquot) were solubilized in 35% TEAH prior to mixing with liquid scintillation fluid for radioactivity measurement.

[0433] Tissue samples were solubilized in toto in 35% TEAH. Duplicate aliquots were then mixed with liquid scintillation fluid prior to radioactivity measurement. Large intestine contents were homogenized in a known volume of water. Duplicate weighed aliquots of large intestine content (LINC) homogenates, stomach contents (STC) and small intestine contents (SINC) were solubilized in 35% TEAH and mixed with liquid scintillation fluid for radioactivity measurement.

2017202349 10 Apr 2017

Radioactivity Measurements

[0434] Radioactivity measurements were conducted by liquid scintillation spectroscopy according to Standard Operating Procedures (SOP). Each sample was counted for 5 minutes or to a two-sigma error of 0.1%, whichever occurred first. All counts were converted to absolute radioactivity (DPM) by automatic quench correction based on the shift of the spectrum for the external standard. The appropriate background DPM values were subtracted from all sample DPM values. Following background subtraction, samples that exhibited radioactivity less than or equal to the background values were considered as zero for all subsequent manipulations.

*Data Analysis**Radioactivity Concentration*

[0435] All radioactivity measurements were entered into a standard computer database program (Debra Version 5.2) for the calculation of concentrations of radioactivity (dpm/g and mass eq/g) and percentage-administered radioactivity in sample. Blood, serum, tissues and CSF concentrations of radioactivity in dpm/g and mass eq/g were calculated on the basis of the measured specific activity (dpm/mg or appropriate mass unit) of radiolabelled test article in the dose solutions. The radioactivity concentration in blood samples was converted to mass eq/mL on the basis of the density of rat blood. Total tissue content was calculated for the total organ weights.

Pharmacokinetics

[0436] The pharmacokinetic (PK) profile of total radioactivity in blood, serum, CSF and tissues was characterized by non-compartmental analysis of the concentration versus time data using validated computer software (WinNonlin, version 3.2, Pharsight Corp., Mountain View, California, USA). Models were selected based on the intravenous and extravascular routes of administration. Concentration values reported as not detectable or quantifiable were not estimated; they were treated as absent samples. Concentration data were obtained from different animals at each time point, and mean values were used to generate a composite pharmacokinetic profile. The 10-minute sampling for Group 1 (Animal Nos. 1001, 1002, 1003) and Group 2

2017202349 10 Apr 2017

(Animal Nos. 2001, 2002, 2003), and the 48-hour for Group 1 (Animal Nos. 1019, 1020) deviated by more than 10% or 6 minutes of the nominal timepoint. This deviation from the protocol did not affect the validity of the study or the data obtained, since the mean time was calculated and used in the pharmacokinetic analyses.

[0437] The area under the radioactivity concentration vs. time curve (AUC) was calculated using the linear trapezoidal method (linear interpolation). When practical, the terminal elimination phase of the PK profile was identified based on the line of best fit (R^2) using at least the final three observed concentration values. The slope of the terminal elimination phase was calculated using log-linear regression using the unweighted concentration data. Parameters relying on the determination of k_{el} were not reported if the coefficient of determination (R^2) was less than 0.8, or if the extrapolation of the AUC to infinity represented more than 20% of the total area.

Results

Analysis of the Dosing Formulations (Table 8)

[0438] On each day of dosing, aliquots of each formulation were analyzed by liquid scintillation spectroscopy prior to and following dose administration to all groups, and the specific activity of the test article calculated from these analyses. The overall mean radioactivity concentration (\pm S.D.) in the formulation for intrathecal administration was $345.4 \times 10^6 \pm 4.92 \times 10^6$ dpm/g ($155.60 \mu\text{Ci/g}$) for Group 1 and $334.4 \times 10^6 \pm 5.87 \times 10^6$ dpm/g ($150.62 \mu\text{Ci/g}$) for Group 3. The overall mean radioactivity concentration in the formulation for intravenous administration was $4.4 \times 10^6 \pm 4.22 \times 10^5$ dpm/g ($1.97 \mu\text{Ci/g}$) for Group 2 and $4.7 \times 10^6 \pm 2.31 \times 10^5$ dpm/g ($2.11 \mu\text{Ci/g}$) for Group 3. The specific activity of the test article in the intrathecal formulation was calculated as $51.16 \mu\text{Ci/mg}$ for the Group 1 dose and $49.53 \mu\text{Ci/mg}$ for the Group 3 dose. The specific activity of the test article in the intravenous formulation was calculated as $6.53 \mu\text{Ci/mg}$ for the Group 2 dose and $6.99 \mu\text{Ci/mg}$ for the Group 3 dose.

Table 8: Summary Results of the Concentration of Radioactivity in the Dosing Formulations by Liquid Scintillation Spectroscopy

Group No.	Route of Administration	Occasion	Mean Concentration of Radioactivity		
			(dpm/g)		(μ Ci/g)
			Mean \pm SD	CV	Mean
1	Intrathecal	Pre-dose	348445137 \pm 3391878	0.97%	156.96
		Post-dose	342426851 \pm 4484476	1.31%	154.25
		Overall	345435994 \pm 4924300	1.43%	155.60
2	Intravenous Bolus Injection	Pre-dose	4091887 \pm 61669	1.51%	1.84
		Post-dose	4672629 \pm 430335	9.21%	2.10
		Overall	4382258 \pm 421765	9.62%	1.97
3	Intrathecal	Pre-dose	332418463 \pm 3013337	0.91%	149.74
		Post-dose	336332353 \pm 7582128	2.25%	151.50
		Overall	334375408 \pm 5868250	1.75%	150.62
3	Intravenous Bolus Injection	Pre-dose	4827255 \pm 92783	1.92%	2.17
		Post-dose	4545578 \pm 247903	5.45%	2.05
		Overall	4686417 \pm 231271	4.93%	2.11

Animal Body Weights and Doses Administered (Table 9)

[0439] The mean body weights of the rats in Groups 1, 2 and 3 on the day prior to dosing were 405 g (range 373 g to 452 g), 410 g (range 367 g to 453 g), and 395 g (range 342 g to 444 g), respectively. The calculated mean dose of 125 I-hGALC administered intrathecally to Group 1 animals was $41 \pm 0.014 \mu\text{g}/\text{animal}$, this was equivalent to a radiochemical dose of $2.12 \pm 0.72 \mu\text{Ci}/\text{animal}$. The mean dose of 125 I-hGALC administered by the intravenous route to Group 2 animals was $1.00 \pm 0.02 \text{ mg}/\text{kg}$ ($2.69 \pm 0.14 \mu\text{Ci}/\text{animal}$). For Group 3, the calculated mean dose

2017202349 10 Apr 2017

of ¹²⁵I-hGALC administered intrathecally and intravenously was 1.08 ± 0.04 mg/kg (5.72 ± 0.31 μCi/animal).

Table 9: Group Mean Body Weights and Specifications of ¹²⁵I-hGALC Dose Administered to Male Sprague-Dawley Rats

Group	Body Weight (kg)	Route of Administration	Radioactivity ^a					
			DPM/animal	μCi/animal	μCi/kg	ng/animal	mg/kg	μg/animal
1	0.405 ± 0.022	IT	4,715,057 ± 1,000,366	3.12 ± 0.72	5.26 ± 1.86	0.041 ± 0.014	0.102 ± 0.037	43
2	0.410 ± 0.021	IV	5,961,365 ± 306,654	2.69 ± 0.14	6.55 ± 0.15	0.411 ± 0.022	1.00 ± 0.023	-
3 ^b	0.395 ± 0.027	IT and IV	13,698,351 ± 886,160	5.72 ± 0.31	14.5 ± 0.62	0.425 ± 0.034	1.08 ± 0.042	-

[0440] The mean chemical dose and the radiochemical dose administered to rats in Group 1 were lower (approximately 32% and 29%, respectively) than the target dose levels and this constituted a deviation from the protocol. However, since the actual doses administered to the animals were used throughout the calculations, these lower values were considered not to affect the validity of the study or the data obtained.

Clinical Observations

[0441] No treatment related clinical signs were observed in any of the rats following administration of ¹²⁵I-hGALC intrathecally at 60 μg/animal and/or intravenously at 1 mg/kg.

Clotting Assessment

[0442] At the earlier time points (10 minutes to 6 hours post dose) it was noted that blood collected from treated animals did not fully clot within the 30 minutes allowed. However the blood collected from 3 untreated spare rats clotted readily, suggesting some interference of the test article with the clotting process. Clotting times of less than or greater than 30 minutes constituted a deviation from the protocol. However, the longer clotting times were required for some samples in order to provide some serum for analysis. A review of the results obtained revealed no correlation between concentration values obtained in serum and the length of time

2017202349 10 Apr 2017

the blood took to clot. Therefore, this extended or shortened clotting time did not affect the validity of the study or the data obtained.

Pharmacokinetics of Total Radioactivity in Blood, Serum, Red Blood Cells, CSF and Tissues

Total Radioactivity Concentrations in Blood, Serum and Red Blood Cells (Table 10, Table 11, Table 12, Figures 18-21)

[0443] Mean concentrations of radiolabelled material in serum of male rats following intrathecal and/or intravenous doses of ^{125}I -hGALC are given in Table 10. Mean concentrations of radiolabelled material in whole blood and in red blood cells are presented in Table 11. Mean data are presented graphically in Figure 18. Mean percentage of radioactivity recovered in supernatant and pellet of blood following TCA precipitation are presented in Table 12.

Group 1 (Intrathecal Mean Dose of 41 $\mu\text{g}/\text{animal}$)

[0444] Following intrathecal dosing, the highest mean concentration (C_{max}) of radiolabelled material in serum and blood were observed at 3 hours following dosing ($0.108 \pm 0.026 \mu\text{g eq/g}$ and $0.093 \pm 0.023 \mu\text{g eq/g}$ respectively). Radioactivity levels in blood remained relatively constant between 3 and 6 hours post dose whereas radioactivity levels in serum declined slightly. Thereafter, radioactivity concentrations in serum and blood declined and were below the limit of quantitation (LOQ) by 48 hours post dose. For red blood cells, C_{max} was observed at 6 hours post dose and was $0.089 \pm 0.024 \mu\text{g eq/g}$. Thereafter, red blood cells radioactivity concentrations declined and were below LOQ by 48 hour post dose. Mean blood to serum ratios following the intrathecal dose were less than 1 throughout the study period (range from 0.7 to 0.9), indicating that the radiolabelled material was not particularly associated with the blood cells. The values of the red blood cell to serum ratios (ranging from 0.8 to 0.9) also supported that radioactivity was not substantially associated with blood cells. The percentage of the dose found in the blood was estimated, using a standard blood volume/body weight (*i.e.* 64.0 mL/kg). At t_{max} (the time at which the highest radioactivity concentration occurred), approximately 6% of the administered dose was associated with blood.

Group 2 (Intravenous Mean Dose of 1.00 mg/kg)

2017202349 10 Apr 2017

[0445] Following intravenous administration, the highest mean concentration (C_{\max}) of radiolabelled material in serum ($14.864 \pm 0.853 \mu\text{g eq/g}$) and blood ($10.228 \pm 0.447 \mu\text{g eq/g}$) were observed at 10 minutes following dosing (*i.e.* the first time point analyzed). Thereafter, radioactivity concentrations in serum and blood declined slowly but were still detectable at 96 hours post dose (serum: $0.088 \pm 0.006 \mu\text{g eq/g}$, 0.59% of C_{\max} ; blood: $0.051 \pm 0.002 \mu\text{g eq/g}$, 0.50% of C_{\max}), with the estimated percent of dose in blood decreasing from 68.4% to 0.3%. For red blood cells, a C_{\max} of $5.136 \pm 1.529 \mu\text{g eq/g}$ was observed at 10 minutes post dose. Thereafter, red blood cells radioactivity concentrations declined and were below LOQ by 96 hours post dose. Mean blood to serum ratios following the intravenous dose were less than 1 throughout the study period (range from 0.6 to 0.8), indicating that the radiolabelled material was not particularly associated with the blood cells. The values of the red blood cell to serum ratios (ranging from 0.4 to 0.6) also supported that radioactivity was not substantially associated with blood cells.

Group 3 (Intrathecal Followed by Intravenous Dose: 1.08 mg/kg (Combined Dose))

[0446] Following the intrathecal dose (target $60 \mu\text{g/animal}$) and the intravenous dose (1 mg/kg), the highest mean concentration (C_{\max}) of radiolabelled material in serum ($14.675 \pm 0.810 \mu\text{g eq/g}$) and blood ($9.974 \pm 0.558 \mu\text{g eq/g}$) were observed at 10 minutes following dosing (*i.e.* the first time point analyzed). Thereafter, radioactivity concentrations in serum and blood declined slowly but were still detectable at 96 hours post dose (serum: $0.077 \pm 0.010 \mu\text{g eq/g}$, 0.52% of C_{\max} ; blood: $0.037 \pm 0.033 \mu\text{g eq/g}$, 0.37% of C_{\max}), with the extrapolated percent of dose in blood decreasing from 32.6% to 0.1%. For red blood cells, a C_{\max} of $6.113 \pm 1.748 \mu\text{g eq/g}$ was observed at 10 minutes post dose. Thereafter, red blood cells radioactivity concentrations declined and were below the limit of quantification by 96 hours post dose. Radiolabelled material was not particularly associated with the blood cells as shown by the mean blood to serum and red blood cell to serum ratios of less than 1 (ranging from 0.7 to 0.8 and 0.4 to 0.7, respectively).

2017202349 10 Apr 2017

Table 10a – Group Mean Concentration of Radioactivity in Serum of Male Sprague-Dawley Rats following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 41 µg/animal

Time Point	Radioactivity Concentration ⁵	
	DFM/g	µg eq/g
10 min	504 ± 462	0.004 ± 0.004
30 min	4125 ± 2327	0.036 ± 0.020
1 h	5705 ± 1535	0.050 ± 0.014
3 h	12311 ± 2960	0.108 ± 0.026
6 h	11473 ± 2596	0.101 ± 0.023
24 h	884 ± 122	0.008 ± 0.001
48 h	0 ± 0	0.000 ± 0.000
96 h	0 ± 0	0.000 ± 0.000

2017202349 10 Apr 2017

Table 10b – Group Mean Concentration of Radioactivity in Serum of Male Sprague-Dawley Rats following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 1.00 mg/kg

Time Point	Radioactivity Concentration ^a	
	DFM/g	µg eq/g
10 min	215632 ± 12377	14.864 ± 0.853
30 min	157259 ± 14339	10.840 ± 0.988
1 h	106804 ± 6790	7.362 ± 0.468
3 h	47009 ± 3754	3.240 ± 0.259
6 h	31898 ± 2417	2.199 ± 0.167
24 h	6584 ± 194	0.454 ± 0.013
48 h	3523 ± 503	0.243 ± 0.035
96 h	1278 ± 86	0.088 ± 0.006

2017202349 10 Apr 2017

Table 10c – Group Mean Concentration of Radioactivity in Serum of Male Sprague-Dawley Rats following a Single Intrathecal and Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 3: At a Mean Dose of 1.08 mg/kg

Time Point	Radioactivity Concentration ^a	
	DFM/g	µg eq/g
10 min	227675 ± 12574	14.675 ± 0.810
30 min	171721 ± 10165	11.069 ± 0.655
1 h	127621 ± 7783	8.226 ± 0.502
3 h	66561 ± 1164	4.290 ± 0.073
6 h	54374 ± 4044	3.505 ± 0.261
24 h	8894 ± 686	0.573 ± 0.044
48 h	3623 ± 458	0.233 ± 0.030
96 h	1199 ± 157	0.077 ± 0.010

2017202349 10 Apr 2017

Table 11a – Group Mean Concentration and Content of Radioactivity in Blood and Blood to Serum Ratios of Male Sprague-Dawley Rats following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 41 µg/animal

Time Point	Radioactivity Concentration *			Blood to Serum Ratio	Percent of Dose
	DPM/g	µg eq/g	µg eq/mL		
10 min	110 ± 364	0.002 ± 0.003	0.002 ± 0.003	0.696 ^b	0.074 ± 0.128
30 min	3579 ± 1918	0.032 ± 0.017	0.033 ± 0.018	0.878 ± 0.029	1.821 ± 0.351
1 h	4933 ± 1446	0.043 ± 0.013	0.046 ± 0.013	0.860 ± 0.027	3.890 ± 0.253
3 h	10617 ± 2586	0.093 ± 0.023	0.098 ± 0.024	0.862 ± 0.006	5.582 ± 0.554
6 h	10530 ± 2507	0.093 ± 0.022	0.097 ± 0.023	0.917 ± 0.033	4.664 ± 0.576
24 h	677 ± 118	0.006 ± 0.001	0.006 ± 0.001	0.764 ± 0.032	0.600 ± 0.114
48 h	0 ± 0	0.000 ± 0.000	0.000 ± 0.000	n/a	0.000 ± 0.000
96 h	0 ± 0	0.000 ± 0.000	0.000 ± 0.000	n/a	0.000 ± 0.000

2017202349 10 Apr 2017

Table 11b – Group Mean Concentration and Content of Radioactivity in Blood and Blood to Serum Ratios of Male Sprague-Dawley Rats following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 2: At a Mean Dose of 1.00 mg/kg

Time Point	Radioactivity Concentration ^a			Blood to Serum Ratio	Percent of Dose
	DPM/g	µg eq/g	µg eq/mL		
10 min	148373 ± 6480	10.228 ± 0.447	10.739 ± 0.469	0.683 ± 0.012	68.393 ± 3.453
30 min	107195 ± 5739	7.389 ± 0.396	7.759 ± 0.415	0.683 ± 0.036	49.317 ± 1.788
1 h	77163 ± 694	5.319 ± 0.048	5.585 ± 0.051	0.724 ± 0.040	36.460 ± 0.174
3 h	35469 ± 3124	2.445 ± 0.215	2.567 ± 0.226	0.754 ± 0.007	16.355 ± 1.166
6 h	24364 ± 1639	1.679 ± 0.113	1.763 ± 0.119	0.764 ± 0.007	11.184 ± 0.612
24 h	4794 ± 160	0.330 ± 0.011	0.347 ± 0.011	0.729 ± 0.030	2.218 ± 0.076
48 h	2259 ± 233	0.156 ± 0.016	0.163 ± 0.017	0.644 ± 0.028	1.042 ± 0.141
96 h	738 ± 29	0.051 ± 0.002	0.053 ± 0.003	0.579 ± 0.052	0.341 ± 0.011

2017202349 10 Apr 2017

Table 11c – Group Mean Concentration and Content of Radioactivity in Blood and Blood to Serum Ratios of Male Sprague-Dawley Rats following a Single Intrathecal Dose and Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 3: At a Mean Dose of 1.98 mg/kg

Time Point	Radioactivity Concentration *			Blood to Serum Ratio	Percent of Dose
	DPM/g	µg eq/g	µg eq/mL		
10 min	154742 ± 8651	9.974 ± 0.558	10.473 ± 0.586	0.680 ± 0.009	32.599 ± 1.331
30 min	117563 ± 4922	7.578 ± 0.317	7.957 ± 0.333	0.685 ± 0.018	24.596 ± 1.523
1 h	92086 ± 2612	5.936 ± 0.181	6.233 ± 0.191	0.723 ± 0.022	19.132 ± 1.432
3 h	52419 ± 244	3.379 ± 0.016	3.548 ± 0.016	0.788 ± 0.017	11.283 ± 0.344
6 h	43097 ± 4071	2.778 ± 0.262	2.917 ± 0.276	0.792 ± 0.019	9.263 ± 0.836
24 h	6561 ± 78	0.423 ± 0.005	0.444 ± 0.006	0.740 ± 0.054	1.345 ± 0.080
48 h	2362 ± 398	0.152 ± 0.026	0.160 ± 0.027	0.650 ± 0.029	0.465 ± 0.083
96 h	581 ± 513	0.037 ± 0.033	0.039 ± 0.035	0.684 ± c	0.124 ± 0.109

2017202349 10 Apr 2017

Table 11d – Group Mean Concentration and Content of Radioactivity in Red Blood Cells and Red Blood Cells to Serum Ratios of Male Sprague-Dawley Rats following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 41µg/animal

Time Point	Radioactivity Concentration ^a		RB Cells to Serum Ratio	Percent of Dose
	DPM/g	µg eq/g		
10 min	0 ± 0	0.000 ± 0.000	n/a	0.000 ± 0.000
30 min	3044 ± 1261	0.027 ± 0.011	0.793 ± 0.148	0.213 ± 0.067
1 h	4434 ± 1396	0.039 ± 0.012	0.773 ± 0.039	0.357 ± 0.336
3 h	9768 ± 2664	0.086 ± 0.023	0.789 ± 0.031	0.734 ± 0.300
6 h	10086 ± 2682	0.089 ± 0.024	0.876 ± 0.083	0.616 ± 0.200
24 h	287 ± 497	0.003 ± 0.004	0.841 ^b	0.044 ± 0.075
48 h	0 ± 0	0.000 ± 0.000	n/a	0.000 ± 0.000
96 h	0 ± 0	0.000 ± 0.000	n/a	0.000 ± 0.000

Table 11e – Group Mean Concentration and Content of Radioactivity in Red Blood Cells and Red Blood Cells to Serum Ratios of Male Sprague-Dawley Rats Following a Single Intravenous Bolus Injection of ^{125}I -hGALC

Group 2: At a Mean Dose of 1.00 mg/kg

Time Point	Radioactivity Concentration ^a		RB Cells to Serum Ratio	Percent of Dose
	DPM/g	$\mu\text{g eq/g}$		
10 min	74506 ± 22185	5.136 ± 1.529	0.350 ± 0.119	4.110 ± 2.794
30 min	59201 ± 14694	4.081 ± 1.013	0.377 ± 0.086	2.600 ± 1.087
1 h	52799 ± 23155	3.639 ± 1.596	0.487 ± 0.196	3.229 ± 2.403
3 h	28039 ± 3432	1.933 ± 0.237	0.599 ± 0.083	1.709 ± 0.734
6 h	19662 ± 2540	1.355 ± 0.175	0.616 ± 0.057	1.143 ± 0.315
24 h	3714 ± 292	0.256 ± 0.020	0.564 ± 0.040	0.164 ± 0.111
48 h	1619 ± 482	0.112 ± 0.033	0.453 ± 0.082	0.076 ± 0.064
96 h	0 ± 0	0.000 ± 0.000	n/s	0.000 ± 0.000

2017202349 10 Apr 2017

Table 11f – Group Mean Concentration and Content of Radioactivity in Red Blood Cells and Red Blood Cells to Serum Ratios of Male Sprague-Dawley Rats Following a Single Intrathecal Dose and Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 3: At a Mean Dose of 1.98 µg/kg

Time Point	Radioactivity Concentration ^a		RB Cells to Serum Ratio	Percent of Dose
	DPM/g	µg eq/g		
10 min	94643 ± 27122	6.113 ± 1.748	0.414 ± 0.104	3.640 ± 1.162
30 min	65477 ± 23687	4.220 ± 1.527	0.378 ± 0.117	2.266 ± 1.583
1 h	61906 ± 14623	3.990 ± 0.943	0.489 ± 0.130	2.253 ± 1.300
3 h	38985 ± 8524	2.513 ± 0.549	0.586 ± 0.128	0.992 ± 0.458
6 h	37327 ± 4497	2.406 ± 0.280	0.685 ± 0.038	1.479 ± 0.417
24 h	5250 ± 334	0.338 ± 0.022	0.591 ± 0.032	0.139 ± 0.070
48 h	2109 ± 319	0.136 ± 0.021	0.581 ± 0.022	0.080 ± 0.017
96 h	0 ± 0	0.000 ± 0.000	n/a	0.000 ± 0.000

2017202349 10 Apr 2017

Table 12a – Mean Percent Radioactivity Recovered in Supernatant and Pellet of Blood from Male Sprague-Dawley Rats Following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 0.10 mg/kg

Time Point	Percent Recovery of Radioactivity ^a	
	Pellet	Supernatant
10 min	100 ± 0	0 ± 0
30 min	75.1 ± 10.7	24.9 ± 10.7
1 h	71.8 ± 11.7	28.2 ± 11.7
3 h	81.2 ± 13.8	18.8 ± 13.8
6 h	67.3 ± 13.5	32.7 ± 13.5
24 h	100 ± 0	0 ± 0
48 h	100 ± 0	0 ± 0
96 h	100 ± 0	0 ± 0

2017202349 10 Apr 2017

Table 12b – Mean Percent Radioactivity Recovered in Supernatant and Pellet of Blood from Male Sprague-Dawley Rats Following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 2: At a Mean Dose of 1.00 mg/kg

Time Point	Percent Recovery of Radioactivity ^a	
	Pellet	Supernatant
10 min	99.2 ± 0.03	0.85 ± 0.03
30 min	97.5 ± 0.32	2.48 ± 0.32
1 h	95.8 ± 0.56	4.23 ± 0.56
3 h	92.5 ± 0.17	7.49 ± 0.17
6 h	90.7 ± 0.45	9.26 ± 0.45
24 h	100 ± 0	0 ± 0
48 h	100 ± 0	0 ± 0
96 h	100 ± 0	0 ± 0

Table 12c – Mean Percent Radioactivity Recovered in Supernatant and Pellet of Blood from Male Sprague-Dawley Rats Following a Single Intrathecal Dose and Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 3: At a Mean Dose of 1.08 mg/kg

Time Point	Percent Recovery of Radioactivity ^a	
	Pellet	Supernatant
10 min	99.0 ± 0.11	1.02 ± 0.11
30 min	95.9 ± 0.49	4.07 ± 0.49
1 h	94.5 ± 0.56	5.55 ± 0.56
3 h	88.1 ± 5.34	11.9 ± 5.34
6 h	88.9 ± 1.03	11.1 ± 1.03
24 h	90.7 ± 3.48	9.33 ± 3.48
48 h	100 ± 0	0 ± 0
96 h	100 ± 0	0 ± 0

2017202349 10 Apr 2017

¹²⁵I-Precipitable in Whole Blood

(Table 12)

[0447] The mean values for recovery of radioactivity in pellet and supernatant following precipitation in whole blood by trichloroacetic acid (TCA) for Groups 1, 2 and 3 are summarized in Table 12. When using a 15% aqueous solution of TCA to precipitate the proteins in whole blood, the radioactivity was mainly recovered in the pellet of the blood (ranging from 100% to 67% in Group 1; 100% to 91% in Group 2; 100% to 88% in Group 3) suggesting that the majority of circulating radioactivity was associated with protein and therefore not reflective of free ¹²⁵iodine.

Radioactivity Concentration in Tissues and Cerebrospinal Fluid (CSF)

(Table 13, Table 14, Table 15, Figures 19 – 30)

[0448] Mean concentrations of radioactivity in tissues and CSF of rats following a single intrathecal and/or intravenous dose of ¹²⁵I-hGALC are given in Table 13. Mean data are presented graphically in Figures 19 – 30. Mean tissue to serum ratios are presented in Table 14 and the recovery of the administered dose in the tissues, CSF and gastrointestinal and urinary bladder contents are given in Table 15.

2017202349 10 Apr 2017

Table 13a – Group Mean Concentration of Radioactivity in Tissues, Cerebrospinal Fluid of Male Sprague-Dawley Rats Following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 41 µg/animal

Sample	Concentration of Radioactivity, µg eq/g ^a			
	10 min	30 min	1 h	3 h
Adipose Tissue (Kidney Fat)	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Adrenal Glands	0.000 ± 0.000	0.014 ± 0.006	0.017 ± 0.006	0.021 ± 0.005
Bone Femur	0.000 ± 0.000	0.011 ± 0.006	0.016 ± 0.005	0.040 ± 0.012
Brain	0.000 ± 0.000	0.003 ± 0.003	0.004 ± 0.004	0.005 ± 0.001
Cerebrospinal Fluid (CSF)	0.000 ^b	0.000 ^b	0.000 ^b	0.000 ± 0.000
Eyes	0.000 ± 0.000	0.006 ± 0.004	0.013 ± 0.003	0.027 ± 0.008
Heart	0.001 ± 0.002	0.014 ± 0.006	0.017 ± 0.005	0.028 ± 0.006
Kidneys	0.004 ± 0.004	0.042 ± 0.023	0.052 ± 0.014	0.096 ± 0.018
Large Intestine	0.000 ± 0.000	0.009 ± 0.004	0.013 ± 0.003	0.024 ± 0.010
Liver	0.000 ± 0.000	0.012 ± 0.007	0.015 ± 0.006	0.029 ± 0.008
Lungs	0.002 ± 0.003	0.020 ± 0.010	0.027 ± 0.008	0.050 ± 0.014
Muscle (Skeletal)	0.000 ± 0.000	0.007 ± 0.003	0.010 ± 0.002	0.014 ± 0.003
Sciatic Nerve	0.000 ± 0.000	0.008 ± 0.003	0.012 ± 0.011	0.043 ± 0.017
Small Intestine	0.000 ± 0.000	0.011 ± 0.005	0.015 ± 0.005	0.046 ± 0.013
Spinal Cord (Lumbar, Thoracic, Cervical)	0.000 ± 0.000	0.004 ± 0.004	0.006 ± 0.002	0.009 ± 0.001
Spleen	0.000 ± 0.000	0.014 ± 0.003	0.019 ± 0.005	0.040 ± 0.010
Stomach	0.003 ± 0.002	0.022 ± 0.010	0.037 ± 0.017	0.203 ± 0.101
Thyroid/Parathyroid Gland	0.020 ± 0.010	0.149 ± 0.083	0.278 ± 0.147	2.031 ± 1.228

Sample	Concentration of Radioactivity, µg eq/g ^a			
	8 h	24 h	48 h	96 h
Adipose Tissue (Kidney Fat)	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Adrenal Glands	0.020 ± 0.002	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Bone Femur	0.041 ± 0.007	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Brain	0.004 ± 0.001	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Cerebrospinal Fluid (CSF)	0.000 ^b	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Eyes	0.024 ± 0.003	0.001 ± 0.001	0.000 ± 0.000	0.000 ± 0.000
Heart	0.026 ± 0.004	0.001 ± 0.002	0.000 ± 0.000	0.000 ± 0.000
Kidneys	0.002 ± 0.012	0.012 ± 0.001	0.008 ± 0.002	0.005 ± 0.001
Large Intestine	0.024 ± 0.003	0.002 ± 0.002	0.000 ± 0.000	0.000 ± 0.000
Liver	0.030 ± 0.002	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Lungs	0.055 ± 0.012	0.004 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Muscle (Skeletal)	0.012 ± 0.002	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Sciatic Nerve	0.030 ± 0.013	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Small Intestine	0.041 ± 0.015	0.004 ± 0.001	0.000 ± 0.000	0.000 ± 0.000
Spinal Cord (Lumbar, Thoracic, Cervical)	0.008 ± 0.003	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Spleen	0.036 ± 0.007	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Stomach	0.163 ± 0.060	0.008 ± 0.001	0.003 ± 0.000	0.002 ± 0.001
Thyroid/Parathyroid Gland	2.453 ± 0.554	4.126 ± 1.073	4.127 ± 1.635	1.927 ± 1.585

2017202349 10 Apr 2017

Table 13b – Group Mean Concentration of Radioactivity in Tissues, Cerebrospinal Fluid of Male Sprague-Dawley Rats Following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 2: At a Mean Dose of 1.00 mg/kg

Sample	Concentration of Radioactivity, $\mu\text{g eq/g}^{\text{a}}$			
	10 min	30 min	1 h	3 h
Adipose Tissue (Kidney Fat)	0.138 ± 0.054	0.158 ± 0.019	0.128 ± 0.007	0.092 ± 0.000
Adrenal Glands	2.327 ± 2.435	7.090 ± 0.547	4.360 ± 0.374	1.873 ± 0.070
Bone Femur	1.568 ± 0.013	1.584 ± 0.223	1.286 ± 0.166	0.887 ± 0.090
Brain	0.252 ± 0.041	0.236 ± 0.017	0.195 ± 0.018	0.083 ± 0.002
Cerebrospinal Fluid (CSF)	0.137 ± 0.238	0.000 ± 0.000	0.000 ^b	0.210 ± 0.363
Eyes	0.110 ± 0.010	0.207 ± 0.016	0.406 ± 0.027	0.344 ± 0.049
Heart	1.215 ± 0.122	1.108 ± 0.039	0.999 ± 0.052	0.558 ± 0.093
Kidneys	3.027 ± 0.330	2.872 ± 0.139	2.288 ± 0.149	1.657 ± 0.190
Large Intestine	0.328 ± 0.072	0.467 ± 0.110	0.492 ± 0.103	0.397 ± 0.031
Liver	11.335 ± 1.436	8.688 ± 0.788	5.904 ± 0.367	3.590 ± 0.192
Lungs	11.584 ± 0.906	29.629 ± 2.125	18.456 ± 3.906	8.526 ± 0.815
Muscle (Skeletal)	0.128 ± 0.011	0.261 ± 0.039	0.175 ± 0.025	0.189 ± 0.007
Sciatic Nerve	0.173 ± 0.023	0.338 ± 0.108	0.594 ± 0.059	0.689 ± 0.036
Small Intestine	0.424 ± 0.004	0.691 ± 0.031	0.786 ± 0.125	0.932 ± 0.166
Spinal Cord (Lumbar, Thoracic, Cervical)	0.293 ± 0.028	0.272 ± 0.009	0.277 ± 0.008	0.142 ± 0.010
Spleen	6.595 ± 0.625	5.952 ± 1.316	4.187 ± 0.311	2.010 ± 0.333
Stomach	0.433 ± 0.028	0.939 ± 0.204	1.430 ± 0.076	2.484 ± 0.139
Thyroid/Parathyroid Gland	4.483 ± 1.194	23.335 ± 2.598	37.990 ± 11.900	147.644 ± 58.596

Sample	Concentration of Radioactivity, $\mu\text{g eq/g}^{\text{a}}$			
	6 h	24 h	48 h	96 h
Adipose Tissue (Kidney Fat)	0.077 ± 0.007	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Adrenal Glands	1.313 ± 0.031	0.339 ± 0.033	0.142 ± 0.013	0.074 ± 0.010
Bone Femur	0.726 ± 0.033	0.108 ± 0.016	0.034 ± 0.030	0.000 ± 0.000
Brain	0.066 ± 0.009	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Cerebrospinal Fluid (CSF)	0.185 ± 0.321	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Eyes	0.336 ± 0.080	0.033 ± 0.006	0.000 ± 0.000	0.000 ± 0.000
Heart	0.440 ± 0.032	0.075 ± 0.011	0.040 ± 0.002	0.000 ± 0.000
Kidneys	1.418 ± 0.108	0.337 ± 0.021	0.199 ± 0.009	0.099 ± 0.010
Large Intestine	0.376 ± 0.077	0.054 ± 0.009	0.026 ± 0.003	0.000 ± 0.000
Liver	3.179 ± 0.188	1.020 ± 0.091	0.506 ± 0.046	0.126 ± 0.014
Lungs	3.187 ± 0.079	2.958 ± 1.012	0.325 ± 0.114	0.069 ± 0.003
Muscle (Skeletal)	0.153 ± 0.018	0.008 ± 0.014	0.000 ± 0.000	0.000 ± 0.000
Sciatic Nerve	0.643 ± 0.063	0.025 ± 0.043	0.000 ± 0.000	0.000 ± 0.000
Small Intestine	0.691 ± 0.131	0.094 ± 0.025	0.041 ± 0.012	0.000 ± 0.000
Spinal Cord (Lumbar, Thoracic, Cervical)	0.128 ± 0.017	0.014 ± 0.013	0.000 ± 0.000	0.000 ± 0.000
Spleen	1.667 ± 0.091	0.565 ± 0.046	0.250 ± 0.028	0.111 ± 0.009
Stomach	1.628 ± 0.310	0.129 ± 0.057	0.047 ± 0.005	0.015 ± 0.013
Thyroid/Parathyroid Gland	267.423 ± 177.568	380.829 ± 84.988	294.321 ± 52.953	218.917 ± 43.086

2017202349 10 Apr 2017

Table 13c – Group Mean Concentration of Radioactivity in Tissues, Cerebrospinal Fluid of Male Sprague-Dawley Rats Following a Single Intrathecal Dose and Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 3: At a Mean Dose of 1.88 mg/kg

Sample	Concentration of Radioactivity, $\mu\text{g eq/g}^a$			
	10 min	30 min	1 h	3 h
Adipose Tissue (Kidney Fat)	0.140 ± 0.028	0.178 ± 0.051	0.188 ± 0.020	0.161 ± 0.008
Adrenal Glands	5.767 ± 1.678	5.487 ± 1.129	4.868 ± 0.930	2.010 ± 0.331
Bone Femur	1.227 ± 0.137	1.707 ± 0.160	1.571 ± 0.071	1.261 ± 0.030
Brain	0.283 ± 0.062	0.276 ± 0.010	0.230 ± 0.008	0.133 ± 0.023
Cerebrospinal Fluid (CSF)	2.087 ± 2.912	0.380 ± 0.371	0.598 ± 1.035	0.105 ± 0.182
Eyes	0.110 ± 0.018	0.372 ± 0.042	0.539 ± 0.019	0.611 ± 0.079
Heart	1.034 ± 0.049	1.315 ± 0.156	1.188 ± 0.028	0.845 ± 0.039
Kidneys	2.864 ± 0.353	3.324 ± 0.265	3.390 ± 0.183	2.822 ± 0.020
Large Intestine	0.251 ± 0.026	0.567 ± 0.051	0.716 ± 0.098	0.681 ± 0.102
Liver	10.181 ± 0.606	8.475 ± 0.204	6.237 ± 0.341	3.740 ± 0.655
Lungs	3.133 ± 0.356	5.162 ± 0.564	5.305 ± 0.194	2.727 ± 0.198
Muscle (Skeletal)	0.119 ± 0.006	0.297 ± 0.011	0.411 ± 0.009	0.298 ± 0.015
Sciatic Nerve	0.244 ± 0.037	0.558 ± 0.023	0.994 ± 0.096	1.043 ± 0.057
Small Intestine	0.304 ± 0.093	0.778 ± 0.037	1.149 ± 0.110	1.481 ± 0.152
Spinal Cord (Lumbar, Thoracic, Cervical)	0.327 ± 0.062	0.319 ± 0.025	0.283 ± 0.044	0.227 ± 0.019
Spleen	5.042 ± 0.902	4.721 ± 0.302	3.740 ± 0.406	2.186 ± 0.218
Stomach	0.465 ± 0.068	1.028 ± 0.175	2.450 ± 0.569	4.434 ± 1.455
Thyroid/Parathyroid Gland	3.191 ± 1.542	21.727 ± 8.873	30.411 ± 18.766	139.771 ± 37.999

Sample	Concentration of Radioactivity, $\mu\text{g eq/g}^a$			
	6 h	24 h	48 h	96 h
Adipose Tissue (Kidney Fat)	0.131 ± 0.003	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Adrenal Glands	1.412 ± 0.137	0.301 ± 0.014	0.118 ± 0.013	0.069 ± 0.016
Bone Femur	1.163 ± 0.066	0.148 ± 0.012	0.029 ± 0.026	0.000 ± 0.000
Brain	0.098 ± 0.012	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Cerebrospinal Fluid (CSF)	0.000 ^b	0.000 ^b	0.000 ^b	0.000 ± 0.000
Eyes	0.574 ± 0.083	0.064 ± 0.006	0.010 ± 0.009	0.000 ± 0.000
Heart	0.723 ± 0.057	0.101 ± 0.008	0.038 ± 0.007	0.006 ± 0.011
Kidneys	2.046 ± 0.228	0.515 ± 0.019	0.248 ± 0.029	0.124 ± 0.005
Large Intestine	0.726 ± 0.173	0.074 ± 0.014	0.027 ± 0.004	0.000 ± 0.000
Liver	3.156 ± 0.143	0.996 ± 0.055	0.418 ± 0.036	0.137 ± 0.018
Lungs	1.830 ± 0.133	0.223 ± 0.007	0.076 ± 0.020	0.033 ± 0.008
Muscle (Skeletal)	0.253 ± 0.029	0.032 ± 0.002	0.000 ± 0.000	0.000 ± 0.000
Sciatic Nerve	1.039 ± 0.133	0.056 ± 0.008	0.000 ± 0.000	0.000 ± 0.000
Small Intestine	1.102 ± 0.101	0.136 ± 0.027	0.033 ± 0.008	0.000 ± 0.000
Spinal Cord (Lumbar, Thoracic, Cervical)	0.202 ± 0.032	0.026 ± 0.003	0.000 ± 0.000	0.000 ± 0.000
Spleen	1.648 ± 0.109	0.395 ± 0.017	0.152 ± 0.009	0.083 ± 0.009
Stomach	4.242 ± 1.361	0.463 ± 0.257	0.064 ± 0.014	0.031 ± 0.005
Thyroid/Parathyroid Gland	182.099 ± 38.422	296.957 ± 57.793	199.316 ± 26.285	45.962 ± 23.164

2017202349 10 Apr 2017

Table 14a – Group Mean Tissue, Cerebrospinal Fluid to Serum Radioactivity Ratios of Male Sprague-Dawley Rats Following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 41 µg/animal

Sample	Tissue, CSF to Serum Ratio ^a			
	30 min ^b	30 min	1 h	7 h
Adipose Tissue (Kidney Fat)	n/a	n/a	n/a	0.071 ^b
Adrenal Glands	n/a	0.421 ± 0.118	0.324 ± 0.033	0.196 ± 0.012
Bone Femur	n/a	0.368 ± 0.019	0.319 ± 0.038	0.369 ± 0.028
Brain	n/a	0.097 ^b	0.119 ± 0.018	0.045 ± 0.006
Cerebrospinal Fluid (CSF)	n/a	n/a	n/a	n/a
Eyes	n/a	0.177 ± 0.032	0.216 ± 0.020	0.253 ± 0.007
Heart	0.333	0.412 ± 0.078	0.329 ± 0.008	0.265 ± 0.012
Kidneys	0.976	1.157 ± 0.040	1.036 ± 0.062	0.695 ± 0.058
Large Intestine	n/a	0.249 ± 0.027	0.220 ± 0.023	0.220 ± 0.048
Liver	n/a	0.351 ± 0.028	0.303 ± 0.036	0.265 ± 0.015
Lungs	0.576	0.565 ± 0.059	0.533 ± 0.034	0.532 ± 0.003
Muscle (Skeletal)	n/a	0.197 ± 0.032	0.192 ± 0.031	0.134 ± 0.021
Sciatic Nerve	n/a	0.349 ^b	0.317 ^b	0.382 ± 0.074
Small Intestine	n/a	0.318 ± 0.033	0.312 ± 0.033	0.426 ± 0.018
Spinal Cord (Lumbar, Thoracic, Cervical)	n/a	0.144 ^b	0.117 ± 0.017	0.082 ± 0.010
Spleen	n/a	0.395 ± 0.036	0.382 ± 0.013	0.368 ± 0.007
Stomach	0.377	0.627 ± 0.072	0.723 ± 0.252	1.801 ± 0.619
Thyroid/Parathyroid Gland	4.443	4.035 ± 0.750	5.680 ± 2.612	17.423 ± 8.215

Sample	Tissue, CSF to Serum Ratio ^a			
	6 h	24 h	48 h	96 h
Adipose Tissue (Kidney Fat)	0.057 ± 0.012	n/a	n/a	n/a
Adrenal Glands	0.197 ± 0.026	n/a	n/a	n/a
Bone Femur	0.407 ± 0.022	n/a	n/a	n/a
Brain	0.040 ± 0.005	n/a	n/a	n/a
Cerebrospinal Fluid (CSF)	n/a	n/a	n/a	n/a
Eyes	0.245 ± 0.023	0.284 ^b	n/a	n/a
Heart	0.258 ± 0.022	0.343 ^b	n/a	n/a
Kidneys	0.821 ± 0.066	1.491 ± 0.128	n/a	n/a
Large Intestine	0.250 ± 0.074	0.395 ^b	n/a	n/a
Liver	0.293 ± 0.029	n/a	n/a	n/a
Lungs	0.347 ± 0.009	0.489 ± 0.105	n/a	n/a
Muscle (Skeletal)	0.115 ± 0.013	n/a	n/a	n/a
Sciatic Nerve	0.406 ± 0.030	n/a	n/a	n/a
Small Intestine	0.400 ± 0.059	0.520 ± 0.021	n/a	n/a
Spinal Cord (Lumbar, Thoracic, Cervical)	0.079 ± 0.009	n/a	n/a	n/a
Spleen	0.337 ± 0.009	n/a	n/a	n/a
Stomach	1.604 ± 0.478	1.083 ± 0.155	n/a	n/a
Thyroid/Parathyroid Gland	24.297 ± 0.831	327.002 ± 160.186	n/a	n/a

2017202349 10 Apr 2017

Table 14b – Group Mean Tissue, Cerebrospinal Fluid to Serum Radioactivity Ratios of Male Sprague-Dawley Rats Following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 1.66 mg/kg

Sample	Tissue, CSF to Serum Ratio ^a			
	10 min	30 min	1 h	3 h
Adipose Tissue (Kidney Fat)	0.009 ± 0.003	0.015 ± 0.003	0.017 ± 0.001	0.028 ± 0.003
Adrenal Glands	0.589 ± 0.133	0.654 ± 0.010	0.594 ± 0.089	0.580 ± 0.039
Bone Femur	0.106 ± 0.006	0.146 ± 0.019	0.174 ± 0.012	0.273 ± 0.008
Brain	0.017 ± 0.002	0.022 ± 0.003	0.027 ± 0.002	0.026 ± 0.002
Cerebrospinal Fluid (CSF)	0.028 ^b	n/a	n/a	0.188 ^b
Eyes	0.007 ± 0.001	0.028 ± 0.002	0.053 ± 0.008	0.106 ± 0.009
Heart	0.082 ± 0.004	0.103 ± 0.006	0.136 ± 0.007	0.171 ± 0.016
Kidneys	0.203 ± 0.011	0.266 ± 0.019	0.311 ± 0.015	0.512 ± 0.043
Large Intestine	0.022 ± 0.004	0.043 ± 0.009	0.067 ± 0.012	0.123 ± 0.019
Liver	0.766 ± 0.119	0.802 ± 0.026	0.805 ± 0.085	1.110 ± 0.055
Lungs	0.781 ± 0.076	1.903 ± 0.100	2.486 ± 0.472	2.642 ± 0.316
Muscle (Skeletal)	0.008 ± 0.001	0.024 ± 0.004	0.037 ± 0.002	0.059 ± 0.004
Sciatic Nerve	0.012 ± 0.002	0.052 ± 0.012	0.080 ± 0.009	0.213 ± 0.007
Small Intestine	0.029 ± 0.002	0.064 ± 0.007	0.107 ± 0.019	0.255 ± 0.032
Spinal Cord (Lumbar, Thoracic, Cervical)	0.019 ± 0.002	0.025 ± 0.002	0.038 ± 0.001	0.044 ± 0.007
Spleen	0.443 ± 0.016	0.547 ± 0.090	0.571 ± 0.075	0.618 ± 0.065
Stomach	0.029 ± 0.004	0.088 ± 0.013	0.194 ± 0.012	0.743 ± 0.028
Thyroid/Parathyroid Gland	0.305 ± 0.092	2.074 ± 0.319	5.106 ± 1.355	46.707 ± 21.839

Sample	Tissue, CSF to Serum Ratio ^a			
	6 h	24 h	48 h	96 h
Adipose Tissue (Kidney Fat)	0.035 ± 0.003	n/a	n/a	n/a
Adrenal Glands	0.554 ± 0.045	0.747 ± 0.061	0.593 ± 0.104	0.845 ± 0.120
Bone Femur	0.330 ± 0.004	0.234 ± 0.030	0.225 ^b	n/a
Brain	0.030 ± 0.002	n/a	n/a	n/a
Cerebrospinal Fluid (CSF)	0.232 ^b	n/a	n/a	n/a
Eyes	0.132 ± 0.024	0.073 ± 0.012	n/a	n/a
Heart	0.200 ± 0.010	0.165 ± 0.024	0.167 ± 0.025	n/a
Kidneys	0.649 ± 0.086	0.741 ± 0.025	0.833 ± 0.169	1.118 ± 0.086
Large Intestine	0.171 ± 0.035	0.119 ± 0.020	0.109 ± 0.050	n/a
Liver	1.449 ± 0.096	2.245 ± 0.142	2.109 ± 0.302	1.440 ± 0.229
Lungs	1.453 ± 0.071	6.505 ± 2.210	1.343 ± 0.431	0.780 ± 0.033
Muscle (Skeletal)	0.069 ± 0.003	0.052 ^b	n/a	n/a
Sciatic Nerve	0.292 ± 0.011	0.189 ^b	n/a	n/a
Small Intestine	0.316 ± 0.065	0.207 ± 0.057	0.175 ± 0.070	n/a
Spinal Cord (Lumbar, Thoracic, Cervical)	0.050 ± 0.003	0.047 ^b	n/a	n/a
Spleen	0.760 ± 0.084	1.247 ± 0.134	1.030 ± 0.098	1.263 ± 0.068
Stomach	0.774 ± 0.176	0.397 ± 0.129	0.197 ± 0.047	0.263 ^b
Thyroid/Parathyroid Gland	124.616 ± 86.507	613.613 ± 169.527	1231.604 ± 285.895	2484.660 ± 471.997

2017202349 10 Apr 2017

Table 14c – Group Mean Tissue, Cerebrospinal Fluid to Serum Radioactivity Ratios of Male Sprague-Dawley Rats Following a Single Intrathecal Dose and Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 3: At a Mean Dose of 1.88 mg/kg

Sample	Tissue, CSF to Serum Ratio ^a			
	10 min	30 min	1 h	3 h
Adipose Tissue (Kidney Fat)	0.010 ± 0.002	0.016 ± 0.005	0.023 ± 0.003	0.037 ± 0.002
Adrenal Glands	0.649 ± 0.076	0.493 ± 0.072	0.388 ± 0.077	0.468 ± 0.060
Bone Femur	0.083 ± 0.008	0.155 ± 0.012	0.191 ± 0.013	0.294 ± 0.005
Brain	0.019 ± 0.003	0.025 ± 0.001	0.028 ± 0.003	0.035 ± 0.005
Cerebrospinal Fluid (CSF)	0.204 ^b	0.053 ^b	0.221 ^b	0.072 ^b
Eyes	0.007 ± 0.001	0.034 ± 0.005	0.066 ± 0.007	0.143 ± 0.021
Heart	0.071 ± 0.001	0.139 ± 0.015	0.145 ± 0.011	0.197 ± 0.012
Kidneys	0.195 ± 0.013	0.302 ± 0.041	0.413 ± 0.022	0.638 ± 0.016
Large Intestine	0.018 ± 0.002	0.051 ± 0.005	0.087 ± 0.011	0.159 ± 0.023
Liver	0.694 ± 0.007	0.768 ± 0.061	0.759 ± 0.023	0.872 ± 0.024
Lungs	0.214 ± 0.030	0.465 ± 0.025	0.646 ± 0.043	0.635 ± 0.036
Muscle (Skeletal)	0.008 ± 0.001	0.027 ± 0.002	0.050 ± 0.003	0.070 ± 0.005
Sciatic Nerve	0.017 ± 0.003	0.050 ± 0.004	0.122 ± 0.018	0.243 ± 0.012
Small Intestine	0.021 ± 0.007	0.071 ± 0.007	0.146 ± 0.016	0.326 ± 0.029
Spinal Cord (Lumbar, Thoracic, Cervical)	0.022 ± 0.005	0.029 ± 0.001	0.035 ± 0.007	0.059 ± 0.004
Spleen	0.342 ± 0.044	0.427 ± 0.036	0.455 ± 0.044	0.530 ± 0.049
Stomach	0.032 ± 0.005	0.094 ± 0.020	0.300 ± 0.078	1.030 ± 0.348
Thyroid/Parathyroid Gland	0.217 ± 0.108	1.960 ± 0.776	3.781 ± 2.521	32.561 ± 8.787

Sample	Tissue, CSF to Serum Ratio ^a			
	6 h	24 h	48 h	96 h
Adipose Tissue (Kidney Fat)	0.038 ± 0.003	n/a	n/a	n/a
Adrenal Glands	0.405 ± 0.059	0.527 ± 0.029	0.514 ± 0.108	0.918 ± 0.317
Bone Femur	0.333 ± 0.006	0.258 ± 0.026	0.183 ^b	n/a
Brain	0.028 ± 0.001	n/a	n/a	n/a
Cerebrospinal Fluid (CSF)	n/a	n/a	n/a	n/a
Eyes	0.163 ± 0.012	0.113 ± 0.019	0.068 ^b	n/a
Heart	0.206 ± 0.001	0.177 ± 0.018	0.164 ± 0.008	0.246 ^b
Kidneys	0.583 ± 0.022	0.900 ± 0.057	1.071 ± 0.104	1.623 ± 0.270
Large Intestine	0.207 ± 0.044	0.131 ± 0.031	0.116 ± 0.018	n/a
Liver	0.902 ± 0.028	1.744 ± 0.121	1.815 ± 0.316	1.770 ± 0.023
Lungs	0.522 ± 0.009	0.391 ± 0.040	0.321 ± 0.044	0.428 ± 0.084
Muscle (Skeletal)	0.072 ± 0.011	0.056 ± 0.008	n/a	n/a
Sciatic Nerve	0.296 ± 0.016	0.293 ^b	n/a	n/a
Small Intestine	0.314 ± 0.008	0.239 ± 0.063	0.148 ± 0.019	n/a
Spinal Cord (Lumbar, Thoracic, Cervical)	0.057 ± 0.007	0.048 ± 0.009	n/a	n/a
Spleen	0.471 ± 0.033	0.692 ± 0.067	0.661 ± 0.104	1.087 ± 0.230
Stomach	1.206 ± 0.373	0.807 ± 0.616	0.274 ± 0.092	0.405 ± 0.112
Thyroid/Parathyroid Gland	52.475 ± 13.382	525.935 ± 143.883	854.144 ± 52.674	571.941 ± 305.367

2017202349 10 Apr 2017

Table 15a – Group Mean Radioactivity Content in Tissues, Cerebrospinal Fluid, Gastrointestinal Tract and Urinary Bladder Contents of Male Sprague-Dawley Rats Following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 41 µg/inimal

Sample	Percent of Dose ^a			
	10 min	30 min	1 h	3 h
Adrenal Glands	0.000 ± 0.000	0.003 ± 0.000	0.004 ± 0.000	0.003 ± 0.001
Brain	0.000 ± 0.000	0.010 ± 0.009	0.027 ± 0.024	0.023 ± 0.003
Cerebrospinal Fluid (CSF)	0.000 ^b	0.000 ^b	0.000 ^b	0.000 ± 0.000
Eyes	0.000 ± 0.000	0.003 ± 0.001	0.011 ± 0.002	0.017 ± 0.001
Heart	0.002 ± 0.003	0.037 ± 0.003	0.066 ± 0.009	0.077 ± 0.004
Kidneys	0.027 ± 0.025	0.253 ± 0.086	0.553 ± 0.057	0.632 ± 0.104
Liver	0.000 ± 0.000	0.417 ± 0.057	0.748 ± 0.108	0.907 ± 0.101
Lungs	0.003 ± 0.005	0.055 ± 0.009	0.123 ± 0.021	0.166 ± 0.013
Sciatic Nerve	0.000 ± 0.000	0.001 ± 0.001	0.002 ± 0.002	0.003 ± 0.001
Spinal Cord (Lumbar, Thoracic, Cervical)	0.000 ± 0.000	0.004 ± 0.004	0.013 ± 0.001	0.012 ± 0.000
Spleen	0.000 ± 0.000	0.024 ± 0.009	0.054 ± 0.010	0.066 ± 0.011
Thyroid/Parathyroid Gland	0.001 ± 0.001	0.007 ± 0.002	0.024 ± 0.012	0.108 ± 0.021
<u>Gastrointestinal Tract:</u>				
Small Intestine	0.000 ± 0.000	0.193 ± 0.054	0.364 ± 0.069	0.763 ± 0.107
Small Intestine Contents	0.000 ± 0.000	0.309 ± 0.062	0.778 ± 0.084	2.611 ± 0.291
Large Intestine	0.000 ± 0.000	0.090 ± 0.018	0.165 ± 0.037	0.230 ± 0.070
Large Intestine Contents	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000	0.598 ± 0.114
Stomach	0.008 ± 0.007	0.086 ± 0.009	0.216 ± 0.094	0.236 ± 0.336
Stomach Contents	0.000 ± 0.000	0.489 ± 0.052	1.774 ± 0.326	3.094 ± 0.346
Urinary Bladder Contents	0.003 ± 0.003	0.110 ± 0.030	0.156 ± 0.077	1.207 ± 1.029
Sample	Percent of Dose ^a			
	6 h	24 h	48 h	96 h
Adrenal Glands	0.002 ± 0.000	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Brain	0.015 ± 0.001	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Cerebrospinal Fluid (CSF)	0.000 ^b	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Eyes	0.013 ± 0.003	0.001 ± 0.001	0.000 ± 0.000	0.000 ± 0.000
Heart	0.056 ± 0.006	0.005 ± 0.008	0.000 ± 0.000	0.000 ± 0.000
Kidneys	0.452 ± 0.115	0.130 ± 0.034	0.059 ± 0.007	0.029 ± 0.006
Liver	0.775 ± 0.078	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Lungs	0.132 ± 0.019	0.019 ± 0.005	0.000 ± 0.000	0.000 ± 0.000
Sciatic Nerve	0.003 ± 0.001	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Spinal Cord (Lumbar, Thoracic, Cervical)	0.000 ± 0.001	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Spleen	0.048 ± 0.007	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Thyroid/Parathyroid Gland	0.125 ± 0.037	0.348 ± 0.013	0.195 ± 0.075	0.086 ± 0.023
<u>Gastrointestinal Tract:</u>				
Small Intestine	0.571 ± 0.165	0.117 ± 0.027	0.000 ± 0.000	0.000 ± 0.000
Small Intestine Contents	1.740 ± 0.925	0.385 ± 0.047	0.000 ± 0.000	0.000 ± 0.000
Large Intestine	0.199 ± 0.022	0.029 ± 0.025	0.000 ± 0.000	0.000 ± 0.000
Large Intestine Contents	0.864 ± 0.100	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Stomach	0.537 ± 0.117	0.059 ± 0.023	0.015 ± 0.002	0.009 ± 0.007
Stomach Contents	3.996 ± 1.013	0.758 ± 0.167	0.122 ± 0.107	0.000 ± 0.000
Urinary Bladder Contents	0.525 ± 0.284	0.178 ± 0.130	0.014 ^b	0.021 ± 0.028

2017202349 10 Apr 2017

Table 15b – Group Mean Radioactivity Content in Tissues, Cerebrospinal Fluid, Gastrointestinal Tract and Urinary Bladder Contents of Male Sprague-Dawley Rats Following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 2: At a Mean Dose of 1.00 mg/kg

Sample	Percent of Dose ^a			
	10 min	30 min	1 h	3 h
Adrenal Glands	0.113 ± 0.023	0.117 ± 0.022	0.059 ± 0.009	0.026 ± 0.002
Brain	0.129 ± 0.022	0.120 ± 0.011	0.098 ± 0.005	0.044 ± 0.001
Cerebrospinal Fluid (CSF)	0.001 ± 0.002	0.000 ± 0.000	0.000 ^b	0.001 ± 0.001
Eyes	0.007 ± 0.001	0.021 ± 0.001	0.028 ± 0.007	0.024 ± 0.005
Heart	0.582 ± 0.055	0.319 ± 0.012	0.292 ± 0.025	0.161 ± 0.024
Kidneys	2.168 ± 0.172	1.966 ± 0.081	1.658 ± 0.014	1.168 ± 0.068
Liver	41.711 ± 3.901	31.161 ± 1.954	20.702 ± 1.140	15.039 ± 0.875
Lungs	4.024 ± 0.305	7.047 ± 0.512	6.456 ± 1.094	2.842 ± 0.248
Sciatic Nerve	0.001 ± 0.001	0.003 ± 0.003	0.004 ± 0.001	0.006 ± 0.003
Spinal Cord (Lumbar, Thoracic, Cervical)	0.045 ± 0.005	0.042 ± 0.008	0.039 ± 0.003	0.023 ± 0.002
Spleen	1.234 ± 0.045	1.014 ± 0.093	0.784 ± 0.123	0.393 ± 0.013
Thyroid/Parathyroid Gland	0.024 ± 0.001	0.100 ± 0.009	0.186 ± 0.050	0.947 ± 0.340
<u>Gastrointestinal Tract:</u>				
Small Intestine	0.749 ± 0.121	1.477 ± 0.237	1.548 ± 0.188	1.335 ± 0.191
Small Intestine Contents	0.590 ± 0.056	1.921 ± 0.346	3.737 ± 1.427	5.446 ± 2.102
Large Intestine	0.327 ± 0.072	0.478 ± 0.076	0.529 ± 0.065	0.412 ± 0.008
Large Intestine Contents	0.000 ± 0.000	0.345 ± 0.029	0.517 ± 0.133	0.782 ± 0.083
Stomach	0.176 ± 0.032	0.437 ± 0.050	0.632 ± 0.047	0.992 ± 0.059
Stomach Contents	0.343 ± 0.127	1.537 ± 0.287	5.330 ± 0.937	10.263 ± 1.971
Urinary Bladder Contents	0.100 ± 0.041	0.409 ± 0.179	0.675 ± 0.060	0.945 ± 0.571

Sample	Percent of Dose ^a			
	6 h	24 h	48 h	96 h
Adrenal Glands	0.019 ± 0.002	0.005 ± 0.001	0.002 ± 0.000	0.001 ± 0.000
Brain	0.034 ± 0.004	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Cerebrospinal Fluid (CSF)	0.003 ± 0.005	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Eyes	0.023 ± 0.007	0.002 ± 0.001	0.000 ± 0.000	0.000 ± 0.000
Heart	0.121 ± 0.007	0.022 ± 0.003	0.012 ± 0.001	0.000 ± 0.000
Kidneys	0.909 ± 0.075	0.251 ± 0.006	0.140 ± 0.003	0.072 ± 0.002
Liver	10.311 ± 0.361	3.891 ± 0.203	1.980 ± 0.065	0.498 ± 0.016
Lungs	1.027 ± 0.037	0.991 ± 0.289	0.112 ± 0.037	0.033 ± 0.001
Sciatic Nerve	0.004 ± 0.000	0.000 ± 0.001	0.000 ± 0.000	0.000 ± 0.000
Spinal Cord (Lumbar, Thoracic, Cervical)	0.017 ± 0.002	0.002 ± 0.002	0.000 ± 0.000	0.000 ± 0.000
Spleen	0.362 ± 0.028	0.100 ± 0.003	0.046 ± 0.010	0.022 ± 0.001
Thyroid/Parathyroid Gland	1.405 ± 0.030	1.333 ± 0.115	1.440 ± 0.604	0.997 ± 0.329
<u>Gastrointestinal Tract:</u>				
Small Intestine	1.537 ± 0.438	0.173 ± 0.033	0.074 ± 0.031	0.000 ± 0.000
Small Intestine Contents	3.051 ± 0.706	0.500 ± 0.184	0.254 ± 0.101	0.000 ± 0.000
Large Intestine	0.380 ± 0.063	0.054 ± 0.007	0.030 ± 0.003	0.000 ± 0.000
Large Intestine Contents	1.053 ± 0.118	0.396 ± 0.058	0.155 ± 0.134	0.000 ± 0.000
Stomach	0.744 ± 0.196	0.078 ± 0.023	0.021 ± 0.003	0.006 ± 0.003
Stomach Contents	0.294 ± 0.070	1.053 ± 0.057	0.296 ± 0.159	0.000 ± 0.000
Urinary Bladder Contents	1.531 ± 1.303	0.075 ^b	0.019 ± 0.021	0.007 ± 0.002

2017202349 10 Apr 2017

Table 15c – Group Mean Radioactivity Content in Tissues, Cerebrospinal Fluid, Gastrointestinal Tract and Urinary Bladder Contents of Male Sprague-Dawley Rats Following a Single Intrathecal Dose and Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 1.83 mg/kg

Sample	Percent of Dose ^a			
	10 min	30 min	1 h	3 h
Adrenal Glands	0.066 ± 0.013	0.037 ± 0.010	0.052 ± 0.009	0.015 ± 0.002
Brain	0.071 ± 0.009	0.071 ± 0.007	0.058 ± 0.002	0.039 ± 0.008
Cerebrospinal Fluid (CSF)	0.046 ± 0.071	0.002 ± 0.002	0.010 ± 0.017	0.002 ± 0.003
Eyes	0.004 ± 0.001	0.012 ± 0.002	0.018 ± 0.001	0.021 ± 0.003
Heart	0.147 ± 0.038	0.188 ± 0.026	0.165 ± 0.008	0.136 ± 0.014
Kidneys	1.003 ± 0.107	1.157 ± 0.073	1.179 ± 0.061	0.985 ± 0.006
Liver	18.955 ± 0.713	14.647 ± 0.420	10.032 ± 1.037	6.754 ± 0.213
Lungs	0.506 ± 0.053	0.811 ± 0.104	0.871 ± 0.037	0.457 ± 0.031
Sciatic Nerve	0.001 ± 0.000	0.002 ± 0.000	0.003 ± 0.001	0.003 ± 0.002
Spinal Cord (Lumbar, Thoracic, Cervical)	0.028 ± 0.007	0.025 ± 0.005	0.023 ± 0.005	0.018 ± 0.004
Spleen	0.468 ± 0.027	0.435 ± 0.037	0.329 ± 0.032	0.217 ± 0.007
Thyroid/Parathyroid Gland	0.008 ± 0.004	0.053 ± 0.019	0.073 ± 0.041	0.392 ± 0.071
<u>Gastrointestinal Tract:</u>				
Small Intestine	0.206 ± 0.046	0.641 ± 0.033	1.118 ± 0.264	1.176 ± 0.044
Small Intestine Contents	0.288 ± 0.037	1.150 ± 0.013	2.414 ± 0.038	4.314 ± 1.755
Large Intestine	0.131 ± 0.013	0.272 ± 0.015	0.360 ± 0.032	0.335 ± 0.051
Large Intestine Contents	0.000 ± 0.000	0.212 ± 0.070	0.351 ± 0.089	0.606 ± 0.101
Stomach	0.100 ± 0.017	0.208 ± 0.034	0.493 ± 0.124	0.931 ± 0.293
Stomach Contents	0.161 ± 0.029	0.866 ± 0.191	2.870 ± 1.090	8.789 ± 1.443
Urinary Bladder Contents	0.029 ± 0.021	0.102 ± 0.031	0.834 ± 0.663	0.273 ± 0.087

Sample	Percent of Dose ^a			
	6 h	24 h	48 h	96 h
Adrenal Glands	0.010 ± 0.002	0.002 ± 0.000	0.001 ± 0.000	0.001 ± 0.000
Brain	0.024 ± 0.003	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Cerebrospinal Fluid (CSF)	0.000 ^b	0.000 ^b	0.000 ^b	0.000 ± 0.000
Eyes	0.019 ± 0.004	0.002 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Heart	0.106 ± 0.030	0.015 ± 0.001	0.005 ± 0.001	0.001 ± 0.002
Kidneys	0.609 ± 0.063	0.103 ± 0.013	0.081 ± 0.007	0.048 ± 0.002
Liver	5.085 ± 0.292	1.653 ± 0.097	0.757 ± 0.043	0.281 ± 0.022
Lungs	0.297 ± 0.011	0.038 ± 0.001	0.012 ± 0.003	0.006 ± 0.002
Sciatic Nerve	0.004 ± 0.001	0.000 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Spinal Cord (Lumbar, Thoracic, Cervical)	0.016 ± 0.003	0.002 ± 0.000	0.000 ± 0.000	0.000 ± 0.000
Spleen	0.146 ± 0.020	0.039 ± 0.003	0.015 ± 0.003	0.009 ± 0.000
Thyroid/Parathyroid Gland	0.496 ± 0.064	0.973 ± 0.162	0.567 ± 0.080	0.124 ± 0.008
<u>Gastrointestinal Tract:</u>				
Small Intestine	0.954 ± 0.153	0.115 ± 0.046	0.032 ± 0.007	0.000 ± 0.000
Small Intestine Contents	2.054 ± 0.707	0.289 ± 0.019	0.113 ± 0.029	0.000 ± 0.000
Large Intestine	0.339 ± 0.097	0.036 ± 0.009	0.013 ± 0.002	0.000 ± 0.000
Large Intestine Contents	0.971 ± 0.085	0.290 ± 0.115	0.041 ± 0.071	0.000 ± 0.000
Stomach	0.943 ± 0.323	0.102 ± 0.084	0.014 ± 0.004	0.008 ± 0.002
Stomach Contents	3.501 ± 3.698	0.610 ± 0.365	0.175 ± 0.053	0.000 ± 0.000
Urinary Bladder Contents	0.114 ± 0.034	0.199 ± 0.266	0.003 ± 0.002	0.005 ± 0.000

Group 1 (Intrathecal Mean Dose of 41 µg/animal)

[0449] Following the intrathecal dose, there was a general distribution of ¹²⁵I-labelled material into all of the tissues examined, however, radioactivity levels in the CSF were below the LOQ. The highest mean concentrations of ¹²⁵I-labelled material in tissues of male rats were

2017202349 10 Apr 2017

observed at 48 hours post dose in thyroid/parathyroid gland ($4.127 \pm 1.635 \mu\text{g eq/g}$) and at 3 hours post dose in stomach ($0.203 \pm 0.101 \mu\text{g eq/g}$), kidneys ($0.096 \pm 0.014 \mu\text{g eq/g}$) and lungs ($0.058 \pm 0.014 \mu\text{g eq/g}$). Levels were lower in the other tissues with t_{max} values generally observed between 3 and 6 hours post dose. The lowest C_{max} values were observed in brain ($0.005 \pm 0.001 \mu\text{g eq/g}$) and kidney fat ($0.006 \pm 0.000 \mu\text{g eq/g}$). By 48 and 96 hours post dose the radioactivity levels in the majority of the tissues were below the limit of detection, the exceptions being thyroid/parathyroid gland, kidneys and stomach. At 96 hours post dose, the highest mean concentration was observed in thyroid/parathyroid gland ($1.927 \pm 1.585 \mu\text{g eq/g}$, 46.7% of C_{max}) followed by the kidneys ($0.005 \pm 0.001 \mu\text{g eq/g}$, 5.2% of C_{max}) and the stomach ($0.002 \pm 0.001 \mu\text{g eq/g}$, 1% of C_{max}).

[0450] Tissue to serum ratios were generally less than 1 for the tissues up to 24 hours post-intrathecal dose. The exceptions were the thyroid/parathyroid gland, kidneys and stomach. The highest ratios were, by far, observed for the thyroid/parathyroid gland. By 48 and 96 hours post dose, tissue to serum ratios could not be calculated since serum concentrations were below the LOQ.

[0451] The levels of radioactivity recovered in all tissues were less than 1% of the administered dose with the highest proportions observed in liver (0.91%) at 3 hours post dose. At 1 hour post dose, proportions greater than 1% of the administered dose were only found in stomach contents (1.8%). By 3 hours post-dosing, proportions of greater than 1% of the administered dose were detected in small intestine contents (2.6%), stomach contents (5.0%) and urinary bladder contents (1.2%). At 6 hours post-dosing, proportions of greater than 1% of the administered dose were found in small intestine contents (1.7%) and stomach contents (4.0%). By 96 hours post dose, small amounts of ^{125}I -hGALC-derived radioactivity (less than 0.1%) was still recovered in kidneys, thyroid/parathyroid gland, stomach and urinary bladder contents, with the highest recoveries observed in the thyroid/parathyroid gland (0.09%).

Group 2 (Intravenous Mean Dose of 1.00 mg/kg)

[0452] Following intravenous administration, the highest mean concentration (C_{max}) of radiolabelled material in tissues of Group 2 rats were observed in thyroid/parathyroid glands ($294.521 \pm 52.953 \mu\text{g eq/g}$; at 48 hours post dose), followed by lungs ($20.629 \pm 2.125 \mu\text{g eq/g}$;

2017202349 10 Apr 2017

30 minutes post dose), liver ($11.335 \pm 1.436 \mu\text{g eq/g}$; 10 minutes post dose), adrenal glands ($8.827 \pm 2.435 \mu\text{g eq/g}$; 10 minutes post dose), spleen ($6.595 \pm 0.625 \mu\text{g eq/g}$; 10 minutes post dose) and kidneys ($3.027 \pm 0.330 \mu\text{g eq/g}$; 10 minutes). The t_{max} values for the tissues occurred between 10 minutes and 3 hours post dose except for the thyroid/parathyroid glands (48 hours post dose). The lowest mean radioactivity C_{max} values were observed in kidney fat ($0.158 \pm 0.019 \mu\text{g eq/g}$), CSF ($0.210 \pm 0.363 \mu\text{g eq/g}$), brain ($0.252 \pm 0.041 \mu\text{g eq/g}$), skeletal muscle ($0.275 \pm 0.025 \mu\text{g eq/g}$) and spinal cord ($0.293 \pm 0.028 \mu\text{g eq/g}$). By 96 hours post-dosing, radioactivity was still detected, in 7 of the 18 tissues analyzed, with the highest mean concentrations being detected in the thyroid/parathyroid glands ($218.917 \pm 45.098 \mu\text{g eq/g}$, 74.3% of C_{max}), followed by liver ($0.126 \pm 0.014 \mu\text{g eq/g}$, 1.1% of C_{max}), spleen ($0.111 \pm 0.009 \mu\text{g eq/g}$, 1.7% of C_{max}) and kidneys ($0.099 \pm 0.010 \mu\text{g eq/g}$, 3.3% of C_{max}).

[0453] At 10 minutes post dose, mean tissue-to-serum ratios were less than 1 for all tissues analyzed. By 30 minutes and 1 hour post dose, mean tissue-to-serum ratios were greater than 1 for lungs and thyroid/parathyroid gland. At 3 and 6 hours post dose, mean tissue-to-serum ratios were greater than 1 for liver, lungs and thyroid/parathyroid gland. At 24 and 48 hours post dose liver, lungs, spleen and thyroid/parathyroid gland had mean tissue-to-serum ratios above 1. At 96 hours post dose, mean tissue-to-serum ratios were greater than 1 for kidneys, liver, spleen and thyroid/parathyroid gland. The highest tissue-to-serum ratios were observed in thyroid/parathyroid glands (2485 at 96 hours), lungs (6.5 at 24 hours) and liver (2.2 at 24 hours).

[0454] In terms of proportion of the radioactivity administered, the highest mean values in tissues were observed in the liver (41.7% at 10 minutes post dose), lungs (7.0% at 30 minutes), kidneys (2.2% at 10 minutes), small intestine (1.5% at 1 hour) and thyroid/parathyroid glands (1.4% at 48 hours). In gastro-intestinal tract contents, the highest mean values were 10.3% of the dose in stomach contents (at 3 hours post dose), 5.4% in small intestine contents (at 3 hours post dose) and 1.1% in large intestine contents (6 hours). By 96 hours post dosing, the highest proportions of the administered dose were detected in thyroid/parathyroid glands (1.0%), liver (0.5%), and kidneys (0.1%). At this time point post dose, less than 0.01% of the administered dose remained in the stomach and urinary bladder contents.

2017202349 10 Apr 2017

Group 3 (Intrathecal Followed by Intravenous Dose: 1.08 mg/kg (Combined Dose))

[0455] Following the intrathecal and the intravenous dose, the highest mean concentration (C_{\max}) of radiolabelled material in tissues of Group 3 rats were observed in thyroid/parathyroid glands ($296.957 \pm 57.793 \mu\text{g eq/g}$; at 24 hours post dose), followed by liver ($10.181 \pm 0.600 \mu\text{g eq/g}$; 10 minutes post dose), adrenal glands ($9.567 \pm 1.678 \mu\text{g eq/g}$; 10 minutes post dose), lungs ($5.305 \pm 0.194 \mu\text{g eq/g}$; 1 hour post dose), spleen ($5.042 \pm 0.902 \mu\text{g eq/g}$; 10 minutes post dose), stomach ($4.454 \pm 1.455 \mu\text{g eq/g}$; 3 hour, post dose), kidneys ($3.390 \pm 0.183 \mu\text{g eq/g}$; 1 hour) and CSF ($2.087 \pm 2.912 \mu\text{g eq/g}$; 10 minutes). The t_{\max} values for the tissues occurred between 10 minutes and 3 hours post dose except for the large intestine (6 hours post dose) and thyroid/parathyroid glands (24 hours post dose). The lowest mean radioactivity C_{\max} values were observed in kidney fat ($0.188 \pm 0.020 \mu\text{g eq/g}$), brain ($0.283 \pm 0.062 \mu\text{g eq/g}$), spinal cord ($0.327 \pm 0.062 \mu\text{g eq/g}$) and skeletal muscle ($0.411 \pm 0.009 \mu\text{g eq/g}$). By 96 hours post-dosing, radioactivity was still detected, in 8 of the 18 tissues analyzed, the highest mean concentrations being detected in the thyroid/parathyroid glands ($43.962 \pm 23.164 \mu\text{g eq/g}$, 14.8% of C_{\max}), followed by liver ($0.137 \pm 0.018 \mu\text{g eq/g}$, 1.3% of C_{\max}), kidneys ($0.124 \pm 0.005 \mu\text{g eq/g}$, 3.7% of C_{\max}), spleen ($0.083 \pm 0.009 \mu\text{g eq/g}$, 1.6% of C_{\max}) and adrenal glands ($0.069 \pm 0.016 \mu\text{g eq/g}$, 0.7% of C_{\max}).

[0456] At 10 minutes post dose, mean tissue-to-serum ratios were less than 1 for all tissues analyzed. By 30 minutes and 1 hour post dose, mean tissue-to-serum ratios were greater than 1 for thyroid/parathyroid gland. At 3 and 6 hours post dose, mean tissue-to-serum ratios were greater than 1 for stomach and thyroid/parathyroid gland. At 24 hours post dose liver and thyroid/parathyroid gland had mean tissue-to-serum ratios above 1. At 48 and 96 hours post dose, mean tissue-to-serum ratios were greater than 1 for kidneys, liver and thyroid/parathyroid gland and for the spleen (96 hours). The highest tissue-to-serum ratios were observed in thyroid/parathyroid glands (854 at 48 hours), liver (1.8 at 48 hours) and kidneys (1.6 at 96 hours).

[0457] In terms of proportion of the radioactivity administered, the highest mean values in tissues were observed in the liver (19.0% at 10 minutes post dose), kidneys (1.2% at 1 hour) and small intestine (1.2 at 3 hours). In gastro-intestinal tract contents, the highest mean values

2017202349 10 Apr 2017

were 8.8% of the dose in stomach contents (at 3 hours post dose), 4.3% in small intestine contents (at 3 hours post dose) and 1.0% in large intestine contents (6 hours). By 96 hours post dosing, the highest proportions of the administered dose were detected liver (0.3%), in thyroid/parathyroid glands (0.1%), and kidneys (0.05%). At this time point post dose, less than 0.01% of the administered dose remained in the adrenal glands, heart, lungs, spleen, stomach and urinary bladder contents.

Pharmacokinetics of Radioactivity in Blood, Serum, Red Blood Cells, CSF and Tissues

(Table 16 and Table 17)

[0458] Mean pharmacokinetic parameters for radioactivity in blood, serum, red blood cells, CSF and tissues of rats following a single intrathecal and/or intravenous dose of ¹²⁵I-hGALC are given in Table 16 and Table 17.

2017202349 10 Apr 2017

Table 16: Disposition Kinetics of the Total Radioactivity in Serum, Blood and Red Blood Cells of Male Sprague-Dawley Rats Following a Single Intrathecal Dose of ¹²⁵I-hGALC

Serum								
Group I: At a Mean Dose of 41 µg/animal								
t _{max} (h)	C _{max} (µg eq/g)	t _{1/2} (h)	AUC _{0-24h} (µg eq·h/g)	k (h ⁻¹)	R ²	t _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	%Extrapolation AUC _{0-∞}
3	0.108	24	1.48	0.130	0.938	3.34	1.34	4.09
Blood								
t _{max} (h)	C _{max} (µg eq/g)	t _{1/2} (h)	AUC _{0-24h} (µg eq·h/g)	k (h ⁻¹)	R ²	t _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	%Extrapolation AUC _{0-∞}
3	0.0930	24	1.33	0.138	0.923	5.62	1.37	3.16
Red Blood Cells								
t _{max} (h)	C _{max} (µg eq/g)	t _{1/2} (h)	AUC _{0-24h} (µg eq·h/g)	k (h ⁻¹)	R ²	t _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	%Extrapolation AUC _{0-∞}
6	0.0250	24	1.24	0.170	0.920	4.08	1.25	1.41

2017202349 10 Apr 2017

Table 16b – Disposition Kinetics of the Total Radioactivity in Serum, Blood and Red Blood Cells of Male Sprague-Dawley Rats Following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Serum										
Group 2: At a Mean Dose of 1.80 mg/kg										
t_{max} (h)	C_{max} ($\mu\text{g eq/g}$)	$t_{1/2}$ (h)	AUC_{0-24h} ($\mu\text{g eq}\cdot\text{h/g}$)	k (h^{-1})	R^2	$t_{1/2}$ (h)	AUC_{0-24h} ($\mu\text{g eq}\cdot\text{h/g}$)	%Extrapolation AUC_{0-24h}	V_z (mL/kg)	Cl (mL/h/kg)
0	20.1	96	71.1	0.0226	0.997	30.7	73.0	5.21	591	13.3
Blood										
t_{max} (h)	C_{max} ($\mu\text{g eq/g}$)	$t_{1/2}$ (h)	AUC_{0-24h} ($\mu\text{g eq}\cdot\text{h/g}$)	k (h^{-1})	R^2	$t_{1/2}$ (h)	AUC_{0-24h} ($\mu\text{g eq}\cdot\text{h/g}$)	%Extrapolation AUC_{0-24h}	V_z (mL/kg)	Cl (mL/h/kg)
0	14.0	96	51.2	0.0256	0.994	27.1	52.2	3.75	735	18.2
Red Blood Cells										
t_{max} (h)	C_{max} ($\mu\text{g eq/g}$)	$t_{1/2}$ (h)	AUC_{0-24h} ($\mu\text{g eq}\cdot\text{h/g}$)	k (h^{-1})	R^2	$t_{1/2}$ (h)	AUC_{0-24h} ($\mu\text{g eq}\cdot\text{h/g}$)	%Extrapolation AUC_{0-24h}	V_z (mL/kg)	Cl (mL/h/kg)
0	6.40	48	33.9	0.0655	0.941	10.9	35.7	4.94	441	28.0

2017202349 10 Apr 2017

Table 16c – Disposition Kinetics of the Total Radioactivity in Serum, Blood and Red Blood Cells of Male Sprague-Dawley Rats Following a Single Intrathecal Dose and Intravenous Bolus Injection of ¹²⁵I-hGALC

Serum										
Group 3: At a Mean Dose of 1.88 mg/kg										
<i>t</i> _{max} (h)	<i>C</i> _{max} (µg eq/g)	<i>t</i> _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	<i>k</i> (h ⁻¹)	<i>R</i> ²	<i>t</i> _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	%Extrapolation AUC _{0-∞}	<i>V</i> _z (mL/kg)	Cl (mL/h/kg)
0	16.9	96	898	0.0272	0.983	25.5	92.6	3.06	429	11.7
Blood										
<i>t</i> _{max} (h)	<i>C</i> _{max} (µg eq/g)	<i>t</i> _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	<i>k</i> (h ⁻¹)	<i>R</i> ²	<i>t</i> _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	%Extrapolation AUC _{0-∞}	<i>V</i> _z (mL/kg)	Cl (mL/h/kg)
0	11.4	96	669	0.0332	0.990	20.9	68.0	1.64	478	15.9
Red Blood Cells										
<i>t</i> _{max} (h)	<i>C</i> _{max} (µg eq/g)	<i>t</i> _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	<i>k</i> (h ⁻¹)	<i>R</i> ²	<i>t</i> _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	%Extrapolation AUC _{0-∞}	<i>V</i> _z (mL/kg)	Cl (mL/h/kg)
0	7.56	48	482	0.0721	0.947	9.61	51.8	3.69	293	21.2

2017202349 10 Apr 2017

Table 17a – Disposition Kinetics of the Total Radioactivity in Tissues and Cerebrospinal Fluid of Male Sprague-Dawley Rats Following a Single Intrathecal Dose of ¹²⁵I-hGALC

Group 1: At a Mean Dose of 41 µg/animal

Samples	t _{max} (h)	C _{max} (µg eq/g)	t _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	k (h ⁻¹)	R ²	t _{1/2} (h)	AUC _{0-∞} (µg eq·h/g)	%Extrapolation (AUC _{0-∞})
Adipose Tissue (Kidney Fat)	6	0.0060	6	0.0215	a	a	a	a	a
Adrenal Glands	3	0.0210	6	0.109	a	a	a	a	a
Bone (Femur)	6	0.0410	6	0.166	a	a	a	a	a
Brain	3	0.0050	6	0.0247	a	a	a	a	a
Cerebrospinal Fluid (CSF)	6	0.000	b	0.000	b	b	b	b	b
Eyes	3	0.0270	24	0.345	0.164	0.990	4.23	0.351	1.74
Heart	3	0.0280	24	0.379	0.167	0.987	4.16	0.385	1.56
Kidneys	3	0.0960	96	1.84	0.0118	0.979	58.6	2.37	13.6
Large Intestine	3	0.0240	24	0.347	0.125	0.993	5.54	0.363	4.46
Liver	6	0.0360	6	0.141	a	a	a	a	a
Lungs	3	0.0580	24	0.891	0.134	0.987	5.18	0.821	3.60
Muscle (Skeletal)	3	0.0140	6	0.0623	a	a	a	a	a
Sciatic Nerve	6	0.0500	6	0.201	a	a	a	a	a
Small Intestine	3	0.0460	24	0.606	0.121	0.992	5.74	0.639	3.18
Spinal Cord (lumbar, thoracic, cervical)	3	0.0090	6	0.0436	a	a	a	a	a
Spleen	3	0.0400	6	0.183	a	a	a	a	a
Stomach	3	0.203	96	2.60	0.0177	0.831	39.1	1.71	4.16
Thyroid/Parathyroid Gland	48.1	4.13	96	313	c	0.882	c	c	35.2

a No reportable results as the terminal phase could not be identified

b PK parameters not estimated due to samples being < LLOQ

c Values are not reported because the AUC_{0-∞} was extrapolated by more than 20% or R² is < 0.8.

2017202349 10 Apr 2017

Table 17b – Disposition Kinetics of the Total Radioactivity in Tissues and Cerebrospinal Fluid of Male Sprague-Dawley Rats Following a Single Intravenous Bolus Injection of ¹²⁵I-hGALC

Group 2: At a Mean Dose of 1.89 mg/kg

Sample:	t _{max} (h)	C _{max} (µg eq/g)	t _{1/2} (h)	AUC _{0-t_{last}} (µg eq·h/g)	k (h ⁻¹)	R ²	t _{1/2} (h)	AUC _{0-inf} (µg eq·h/g)	%Extrapolation (AUC _{0-inf})
Adipose Tissue (Kidney Fat)	0.5	0.158	8	0.617	a	0.920	a	a	56.0
Adrenal Glands	0	10.9	96	43.1	0.0291	0.927	34.6	46.8	7.89
Bone (Femur)	0.5	1.58	48	15.3	0.0777	0.965	8.92	15.7	2.79
Brain	0	0.268	6	0.735	a	0.897	a	a	26.8
Cerebrospinal Fluid (CSF)	3	0.216	6	0.854	b	b	b	b	b
Eyes	1	0.406	24	5.35	0.113	0.981	6.15	5.64	5.19
Heart	0	1.35	48	18.2	0.0726	0.939	8.54	16.8	5.12
Kidneys	0	3.18	96	48.7	0.0167	0.988	41.6	46.7	12.7
Large Intestine	1	0.492	48	7.30	0.0658	0.938	10.5	7.70	5.14
Liver	0	14.6	96	109	0.0290	1.00	23.9	105	4.15
Lungs	0.5	20.6	96	165	0.0497	0.939	13.9	167	0.832
Muscle (Skeletal)	1	0.275	24	2.64	0.154	0.996	4.50	2.69	1.93
Sciatic Nerve	3	0.689	24	9.62	0.166	0.987	4.18	9.77	1.54
Small Intestine	3	0.832	48	15.2	0.0693	0.952	10.0	13.8	4.29
Spinal Cord (lumbar, thoracic, cervical)	0	0.315	24	2.39	0.115	0.991	6.54	2.51	4.87
Spleen	0	7.27	96	56.1	0.0218	0.994	31.8	61.2	8.33
Stomach	3	2.40	96	31.9	0.0330	0.945	21.0	32.3	1.41
Thyroid/Parathyroid Gland	48	295	96	24989	b	b	b	b	b

a Values are not reported because the AUC_{0-inf} was extrapolated by more than 20% or R² is <0.8.

b No reportable results as the terminal phase could not be identified.

2017202349 10 Apr 2017

Table 17c – Disposition Kinetics of the Total Radioactivity in Tissues and Cerebrospinal Fluid of Male Sprague-Dawley Rats Following a Single Intrathecal Dose and Intravenous Bolus Injection of ^{125}I -hGALC

Group 3: At a Mean Dose of 1.05 mg/kg

Sample:	$t_{1/2}$ (h)	C_{max} ($\mu\text{g eq/g}$)	$t_{1/2}$ (h)	$\text{AUC}_{0-t_{1/2}}$ ($\mu\text{g eq}\cdot\text{h/g}$)	k (h^{-1})	R^2	$\%$ (h)	$\text{AUC}_{0-\infty}$ ($\mu\text{g eq}\cdot\text{h/g}$)	%Extrapolation ($\text{AUC}_{0-t_{1/2}}$)
Adipose Tissue (Kidney Fat)	1	0.188	6	0.954	a	0.999	a	a	65.6
Adrenal Glands	0	12.6	96	43.0	0.0354	0.835	19.6	45.8	4.25
Bone (Femur)	0.5	1.71	48	21.9	0.0889	0.985	7.98	22.3	1.50
Brain	0	0.287	6	1.03	a	0.992	a	a	36.1
Cerebrospinal Fluid (CSF)	0	4.89	3	1.94	a	0.775	a	a	5.55
Eyes	3	0.611	48	9.88	0.0947	0.928	7.32	9.99	1.06
Heart	0.5	1.32	96	15.7	0.0391	1.00	17.7	15.9	0.967
Kidneys	1	3.38	96	57.9	0.0190	0.960	36.2	64.4	19.1
Large Intestine	6	0.726	48	12.4	0.0764	0.911	9.07	12.8	2.77
Liver	0	11.2	96	96.5	0.0269	0.926	25.7	102	5.00
Lungs	1	5.31	96	44.1	0.0252	0.932	27.5	45.4	2.88
Muscle (Skeletal)	1	0.411	24	4.37	0.110	0.997	6.31	4.86	6.25
Sciatic Nerve	3	1.94	24	15.6	0.147	0.983	4.71	16.0	2.38
Small Intestine	3	1.40	48	20.2	0.0851	0.974	8.14	20.6	1.88
Spinal Cord (lumbar, thoracic, cervical)	0	0.331	24	3.52	0.105	0.994	6.58	3.77	6.53
Spleen	0	5.21	96	46.9	0.0347	0.860	20.0	49.3	4.85
Stomach	3	4.45	96	72.1	0.0557	0.878	12.4	72.6	0.766
Thyroid/Parathyroid Gland	24	297	96	16776	0.0272	0.923	25.4	12390	8.72

a Values are not reported because the $\text{AUC}_{0-\infty}$ was extrapolated by more than 20% or R^2 is <0.8 .*Blood, Serum and Red Blood Cells*

[0459] Following the intrathecal dose (Group 1: 41 $\mu\text{g}/\text{animal}$), the mean calculated areas under the radioactivity concentration vs. time curves from time zero to the last measurable time point ($\text{AUC}_{0-\text{last}}$) for serum, whole blood and red blood cells were 1.48 $\mu\text{g eq}\cdot\text{h/g}$, 1.33 $\mu\text{g eq}\cdot\text{h/g}$ and 1.24 $\mu\text{g eq}\cdot\text{h/g}$, respectively. The apparent terminal $t_{1/2}$ values reported for radioactivity in serum, whole blood and red blood cells were 5.34, 5.02 and 4.08 hours, respectively. The elimination rate constant, k , was calculated as 0.130 h^{-1} , 0.138 h^{-1} and 0.170 h^{-1} in serum, whole blood and red blood cells, respectively. $\text{AUC}_{0-\text{inf}}$ was calculated as 1.54 $\mu\text{g eq}\cdot\text{h/g}$, 1.37 $\mu\text{g eq}\cdot\text{h/g}$ and 1.25 $\mu\text{g eq}\cdot\text{h/g}$ in serum, whole blood and red blood cells, respectively. The elimination phases for radioactivity from serum, whole blood and red blood cells were well-defined, as evidenced by the very low percentage extrapolation values (4.0, 3.2 and 1.4%, respectively) required for calculation of $\text{AUC}_{0-\text{inf}}$.

2017202349 10 Apr 2017

[0460] Following the intravenous dose (Group 2: 1.00 mg/kg), the mean $AUC_{0-t_{last}}$ values for serum, whole blood and red blood cells were 71.1 $\mu\text{g eq}\cdot\text{h/g}$, 51.2 $\mu\text{g eq}\cdot\text{h/g}$ and 33.9 $\mu\text{g eq}\cdot\text{h/g}$, and the apparent terminal $t_{1/2}$ values were 30.7, 27.1 and 10.9 hours, respectively. The value of k was calculated as 0.0226 h^{-1} , 0.0256 h^{-1} and 0.0635 h^{-1} in serum, whole blood and red blood cells, respectively. The elimination phases for radioactivity from serum, whole blood and red blood cells were well-defined and $AUC_{0-\infty}$ was calculated as 75.0 $\mu\text{g eq}\cdot\text{h/g}$ (extrapolation 5.21%), 53.2 $\mu\text{g eq}\cdot\text{h/g}$ (extrapolation 3.75%) and 35.7 $\mu\text{g eq}\cdot\text{h/g}$ (extrapolation 4.94%) in serum, whole blood and red blood cells, respectively. The apparent volume of distribution (V_z) was greatest in whole blood (735 mL/kg) followed by serum (591 mL/kg) and red blood cells (441 mL/kg). Clearance of the test article was estimated at 13.3 mL/h/kg from serum and 18.8 mL/h/kg for whole blood.

[0461] Following the intrathecal dose and intravenous dose (combined 1.08 mg/kg) to Group 3 animals, the mean $AUC_{0-t_{last}}$ values for serum, whole blood and red blood cells were 89.8 $\mu\text{g eq}\cdot\text{h/g}$, 66.9 $\mu\text{g eq}\cdot\text{h/g}$ and 49.2 $\mu\text{g eq}\cdot\text{h/g}$, respectively. The apparent terminal $t_{1/2}$ values reported for radioactivity in serum, whole blood and red blood cells were 25.5, 20.9 and 9.61 hours, respectively, with k as 0.0272 h^{-1} , 0.0332 h^{-1} and 0.0721 h^{-1} . Again, the elimination phases for all three matrices were well-defined, with $AUC_{0-\infty}$ calculated as 92.6 $\mu\text{g eq}\cdot\text{h/g}$, 68.0 $\mu\text{g eq}\cdot\text{h/g}$ and 51.0 $\mu\text{g eq}\cdot\text{h/g}$ (extrapolation of 3.06%, 1.64% and 3.69%) in serum, whole blood and red blood cells, respectively. The V_z was greater in whole blood (478 mL/kg) followed by serum (429 mL/kg) and red blood cells (293 mL/kg). Clearance values were 15.9 mL/h/kg for whole blood and 11.7 mL/h/kg for serum.

Tissues

[0462] The highest $AUC_{0-t_{last}}$ value in tissues from rats, following an intrathecal dose of ^{125}I -hGALC (Group 1: 41 $\mu\text{g/animal}$), was observed in thyroid/parathyroid gland (313 $\mu\text{g eq}\cdot\text{h/g}$), followed by stomach (2.60 $\mu\text{g eq}\cdot\text{h/g}$) and kidneys (1.84 $\mu\text{g eq}\cdot\text{h/g}$). For several tissues, it was not possible to estimate k or any parameters derived from k (i.e. $t_{1/2}$ and $AUC_{0-\infty}$) since the % extrapolation of the AUC to infinity was greater than 20% or due to lack of data in the terminal phase. For those tissues where it could be estimated (eyes, heart, kidneys, large

2017202349 10 Apr 2017

intestine, lungs, small intestine and stomach), k ranged from 0.01 to 0.17 h^{-1} and the $t_{1/2}$ generally ranged from 4 to 6 h, the exceptions being 58.6 h for kidneys and 39.1 h for stomach.

[0463] Following the intravenous dose (Group 2; 1.00 mg/kg), the highest values for $\text{AUC}_{0-\text{last}}$ were observed in thyroid/parathyroid gland (24989 $\mu\text{g eq}\cdot\text{h/g}$), followed by lungs (165 $\mu\text{g eq}\cdot\text{h/g}$), liver (100 $\mu\text{g eq}\cdot\text{h/g}$), spleen (56.1 $\mu\text{g eq}\cdot\text{h/g}$), adrenal glands (43.1 $\mu\text{g eq}\cdot\text{h/g}$) and kidneys (40.7 $\mu\text{g eq}\cdot\text{h/g}$). The lowest $\text{AUC}_{0-\text{last}}$ values were observed for kidney fat (0.617 $\mu\text{g eq}\cdot\text{h/g}$) and brain (0.735 $\mu\text{g eq}\cdot\text{h/g}$). Parameters derived from k were not reported for tissues where the elimination phase was poorly defined (thyroid/parathyroid gland and CSF), or where the extrapolation to $\text{AUC}_{0-\text{inf}}$ was greater than 20% (kidney fat and brain). Only the $\text{AUC}_{0-\text{inf}}$ values for liver and lungs were greater than that of serum (75 $\mu\text{g eq}\cdot\text{h/g}$). The highest reported $\text{AUC}_{0-\text{inf}}$ value was for lungs (167 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 0.832%), followed by liver (105 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 4.15%), spleen (61.2 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 8.33%), adrenal glands (46.8 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 7.89%) and kidneys (46.7 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 12.7%).

[0464] The lowest reported value for $\text{AUC}_{0-\text{inf}}$ value was calculated for spinal cord (2.51 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 4.87%) followed by muscle (2.69 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 1.93%) and eyes (5.64 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 5.19%). The longest calculable $t_{1/2}$ in tissues was 41.6 hours for kidneys, followed by 34.6 hours for the adrenal glands and 31.8 hours for the spleen. The shortest reported $t_{1/2}$ was 4.18 hours for sciatic nerve.

[0465] For Group 3, after an intrathecal and an intravenous dose (1.08 mg/kg, combined dose), the highest values for $\text{AUC}_{0-\text{last}}$ was observed in thyroid/parathyroid gland (16776 $\mu\text{g eq}\cdot\text{h/g}$) followed by liver (96.5 $\mu\text{g eq}\cdot\text{h/g}$), stomach (72.1 $\mu\text{g eq}\cdot\text{h/g}$), kidneys (57.9 $\mu\text{g eq}\cdot\text{h/g}$), spleen (46.9 $\mu\text{g eq}\cdot\text{h/g}$), lungs (44.1 $\mu\text{g eq}\cdot\text{h/g}$) and adrenal glands (43.9 $\mu\text{g eq}\cdot\text{h/g}$). The lowest $\text{AUC}_{0-\text{last}}$ values were observed for kidney fat (0.954 $\mu\text{g eq}\cdot\text{h/g}$) and brain (1.03 $\mu\text{g eq}\cdot\text{h/g}$). Parameters derived from k were not reported for tissues where the extrapolation to $\text{AUC}_{0-\text{inf}}$ was greater than 20% (kidney fat and brain) or R^2 lower than 0.8 (CSF). Only the $\text{AUC}_{0-\text{inf}}$ values for thyroid/parathyroid gland and liver were greater than that of serum (92.6 $\mu\text{g eq}\cdot\text{h/g}$). The highest reported $\text{AUC}_{0-\text{inf}}$ value was for thyroid/parathyroid gland (18390 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 8.78%), followed by liver (102 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 5.0%), stomach (72.6 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 0.766%), kidneys (64.4 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 10.1%), spleen (49.3 $\mu\text{g eq}\cdot\text{h/g}$;

2017202349 10 Apr 2017

extrapolation 4.85%), adrenal glands (45.8 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 4.25%) and lungs (45.4 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 2.88%). The lowest reported value for $\text{AUC}_{0-\text{inf}}$ value was calculated for spinal cord (3.77 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 6.55%) followed by muscle (4.66 $\mu\text{g eq}\cdot\text{h/g}$; extrapolation 6.25%). The longest calculable $t_{1/2}$ in tissues was 36.4 hours for kidneys, followed by 27.5 hours for lungs, 25.7 hours for liver and 25.4 hours thyroid/parathyroid gland. The shortest reported $t_{1/2}$ was 4.71 hours for sciatic nerve.

Discussion

[0466] Following intrathecal administration, the highest mean concentrations of radioactivity in serum and whole blood were observed at 3 hours post dose suggesting relatively rapid distribution of dose-related material to the systemic circulation. Following intravenous administration, the highest mean concentrations of radioactivity in serum and whole blood were observed at the first time point measured. Concentrations in serum were always higher than those in whole blood, as reflected by blood-to-serum ratios of less than 1. This indicated that dose-related material was not particularly associated with the blood cells of any groups at any time post dose. Following TCA precipitate of blood proteins, the radioactivity was mainly recovered in the pellet suggesting that the majority of circulating radioactivity was protein associated, indicating that radioactivity distribution observed was not largely reflective of the disposition of free ^{125}I iodine.

[0467] When comparing Group 2 (intravenous dose 1.00 mg/kg) to Group 3 (intrathecal and intravenous combined dose 1.08 mg/kg), concentrations in Group 3 serum and whole blood appeared to be generally similar to those of Group 2. The decline of radioactivity in both matrices for both groups was also very similar, as assessed by blood-to-serum ratios. Comparing $\text{AUC}_{0-\text{last}}$ and $\text{AUC}_{0-\text{inf}}$ for Group 2 and Group 3 serum and blood, indicated that exposure to dose-related material was slightly higher for Group 3 animals.

[0468] In Group 1, levels of radioactivity in CSF were very low, a finding which does not appear to be in accordance with the administration of the test article directly to the intrathecal space, although very low levels were observed in brain. However, radioactivity was observed in the systemic circulation, and in systemic tissues, shortly following dosing, suggesting that dose-related material was fairly rapidly distributed from the intrathecal space following

2017202349 10 Apr 2017

administration. Higher levels in the stomach and intestinal contents suggested that dose-related material was excreted via feces, although direct measurement in the excreta was not performed in this study. In addition, high levels in the urinary bladder contents also suggest excretion via urine. Other than high levels in the thyroid/parathyroid glands, considered to reflect loss of the iodine label and persistence of the label in this tissue rather than distribution/persistence of the test article itself, high levels of radioactivity were observed in liver, lungs, adrenal glands spleen and kidneys; tissues which were likely to be involved in the metabolism and/or excretion of the test article.

[0469] Distribution of radioactivity was general and widespread by the first time point post dose in Groups 2 and 3. The highest concentrations were generally associated with the liver, lungs, kidneys, spleen, adrenal gland, and in particular, the thyroid/parathyroid glands. Thus the pattern of distribution of radioactivity in tissues of all three groups was largely similar. Again, high levels of radioactivity observed in the thyroid/parathyroid glands of all animals, particularly considering the marked concentration increase with increasing time post dose, probably indicated loss of the iodine label and persistence of the label in this tissue rather than distribution/persistence of the test article itself. CSF levels were higher in these groups, as compared to Group 1, at early timepoints post dose, suggesting that radiolabelled material was able to cross the blood-brain barrier. Slightly higher levels were observed in this matrix in Group 3, as compared to Group 2, again at early timepoints post dose, suggesting that this concentration was accounted for by test article-related material distributing from the intravenous dose and material directly injected into the intrathecal space. The below LOQ values observed for Group 1 may therefore be a consequence of very low concentrations in very small sample volumes, being below the quantitation possible by this analytical method.

[0470] Tissue-to-serum ratios were generally less than 1 in the majority of tissues of all groups by 96 hours post dose, indicating that dose-related material was distributed into the tissues and was generally cleared more readily from the tissues than from the serum. For all groups, exposure of the majority of the tissues to dose-related material (as assessed by $AUC_{0-t_{last}}$) was less than that of serum.

Conclusion

2017202349 10 Apr 2017

[0471] Following administration of a single intrathecal (nominal 60 ug/animal) and/or intravenous bolus dose of ^{125}I -hGALC to male rats (nominal concentrations of 1 mg/kg), concentrations of radioactivity in blood, serum, red blood cells, CSF and tissues were determined.

[0472] The highest observed concentrations of radioactivity in both serum and whole blood occurred at 3 hours post dose following intrathecal administration, indicating relatively rapid distribution to the systemic circulation, or at the first time point post dose (10 minutes) following intravenous dosing. Concentrations in serum were higher than in blood, indicating that test article-related material was not particularly associated with the blood cells. Distribution of radioactivity into tissues was general and widespread by early time points post dose and, in general, the pattern of distribution to tissues was similar between all three groups. For all groups, exposure of the majority of the tissues to dose-related material (as assessed by $\text{AUC}_{0-\text{last}}$) was less than that of serum. High concentrations in thyroid/parathyroid glands for all three groups were considered to indicate loss of the iodine label rather than distribution and persistence of dose-related material in this tissue. By 96 hours post intravenous dose, radioactivity was still detected in a few of the tissues examined.

EXAMPLE 3: PRE-CLINICAL STUDY OF ICV AND ICV/IP RMGALC INJECTION AND EXTENDED SURVIVAL IN TWITCHER MICE

[0473] The present Example demonstrates one embodiment of a preclinical study illustrating extended survival in twitcher mice provided with weekly IP injections of rmGALC. In the present embodiment, improved myelination was observed in the sciatic nerve, along with reduced psychosine levels and gross motor function (i.e., gait) improvement. In some embodiments, twitcher mice treated with a single ICV or ICV/IP rmGALC injection exhibited increased survival and up to a 63% reduction in the levels of brain psychosine. The positive results in important endpoints (i.e., survival, brain psychosine levels) following a single ICV administration of rmGALC along with the very minimal improvement in these endpoints following the addition of systemic administration (ICV/IP) suggest that a CNS only regimen is a viable clinical option for the treatment of GLD.

2017202349 10 Apr 2017

Introduction

[0474] Globoid Cell Leukodystrophy (GLD) is an autosomal recessive lysosomal storage disorder that occurs at an incidence of approximately 1:100,000 births (1.9:100,000 births in Scandinavian countries). A progressive peripheral (PNS) and central (CNS) nervous system disorder, GLD is the result of genetic mutations causing a deficiency in the enzyme activity of galactocerebrosidase (GALC) to degrade substrate lipids [i.e., galactosylceramide to galactose and ceramide; galactosylsphingosine (psychosine) to galactose and sphingosine]. This disorder is characterized by a complete loss of oligodendrocytes and myelin as well as the presence of galactosylceramide-engorged macrophages (“globoid” cells).

[0475] The clinical features of this disease present in two forms: infantile and late-onset. The infantile form of GLD (also known as Krabbe disease) occurs in 90% of all patients diagnosed with GALC deficiency, and symptoms are usually observed within 3-6 months after birth; there are reports of symptoms manifesting as early as 2-3 weeks of age (Wenger, D.A. *et al.*, 2001, *Galactosylceramide Lipidosis: Globoid Cell Leukodystrophy (Krabbe Disease)*, in *The Metabolic and Molecular Bases of Inherited Disease*, C.R. Scriver, Beaudet, A. L., Sly, W.S., and Valle, D, Editor. 2001, McGraw-Hill. p. 3669-3687; incorporated herein as reference). The late-onset variant of this disease usually presents clinically by 10 years of age, however, patients diagnosed at 40 years of age have been reported (Wenger, D.A. *et al.*, 2001, *Galactosylceramide Lipidosis: Globoid Cell Leukodystrophy (Krabbe Disease)*, in *The Metabolic and Molecular Bases of Inherited Disease*, C.R. Scriver, Beaudet, A. L., Sly, W.S., and Valle, D, Editor. 2001, McGraw-Hill. p. 3669-3687; incorporated herein as reference). The decline of function in late-onset patients proceeds gradually over a period of several years.

[0476] Systemic enzyme replacement therapy (ERT) has provided benefit for patients suffering from lysosomal storage disorders (LSDs) such as Gaucher disease, Fabry disease, and Hunter syndrome (Wenger, D.A. *et al.*, 2001, *Galactosylceramide Lipidosis: Globoid Cell Leukodystrophy (Krabbe Disease)*, in *The Metabolic and Molecular Bases of Inherited Disease*, C.R. Scriver, Beaudet, A. L., Sly, W.S., and Valle, D, Editor. 2001, McGraw-Hill. p. 3669-3687; Neufeld, E.F., 2004, *Enzyme Replacement therapy*. Lysosomal disorders of the Brain, ed. F.M.a.W. Platt, S.V. 2004: Oxford University Press. 327-338; Desnick, R.J., 2004. *J. Inherit.*

2017202349 10 Apr 2017

Metab. Dis., **27**(3): p. 385-410; all of which are incorporated herein as reference). ERT for GLD has not been pursued with rigor, perhaps because the disease affects both the PNS and CNS. Current treatments for patients with GLD include hematopoietic cell transplant (HCT), however this procedure has its limitations due to significant adverse events (i.e., 30% treatment-related mortality, lifelong immunosuppressive therapy) and efficacy only in presymptomatic patients.

[0477] The twitcher mouse is the most common experimental animal model used to study GLD, and constitutes the bulk of experimental work on this disease (Wenger, D.A., 2000, *Mol. Med. Today*, 6(11): p. 449-451; incorporated herein as reference), but other naturally occurring animal models of GLD exist with variable degrees of characterization. Spontaneous mutation exists in West Highland White/Cairn Terriers (Kobayashi T., *et al.*, 1980, *Brain Res.*, 202:479-83; incorporated herein as reference), polled Dorset Sheep (Pritchard D., *et al.*, 1980, *Vet. Pathol.*, 17:399-405), the domestic cat (Johnson K., 1970, *J. Am. Vet. Med. Assoc.*, 157:2057-64; incorporated herein as reference) and non-human primate Rhesus macaque (Baskin G., *et al.*, 1998, *Lab Anim. Sci.*, 48:476-82; incorporated herein as reference).

[0478] The initial nerve allograft studies demonstrated that the ability to improve peripheral nerve function of twitcher mouse Schwann cells was mediated by enzyme replacement into allograft twitcher cells in situ and that long term recovery of injured twitcher peripheral myelinating cells was possible. This technology, however, could not be generalized as an overall therapy of the twitcher mouse (Baskin G., *et al.*, 1998, *Lab Anim. Sci.*, 48:476-82; incorporated herein as reference). In affected mice, HCT demonstrated significant improvement in the life span and weight gain of affected animals, however variable efficacy is observed with viability documented between 44 days to more than 100 days (in mice receiving myeloreductive conditioning) (Lin, D., *et al.*, 2007, *Mol. Ther.*, **15**(1): p. 44-52; Hoogerbrugge, P.M., *et al.*, 1998, *J. Clin. Invest.*, **81**(6): p. 1790-4; both of which are herein incorporated as reference). The typical life span of untreated mice in these investigations was approximately 40 days.

[0479] In these and other studies, neither the rate of remyelination nor existing brain pathology was improved in treated mice versus untreated controls (Yeager A., *et al.*, 1984, *Science*, 225:1052-4; Toyoshima, E., *et al.*, 1986, *J. Neurol. Sci.*, **74**(2-3), p. 307-18; both of which are herein incorporated as reference). Substrate inhibition targeting sphingosine synthesis

2017202349 10 Apr 2017

using L-cycloserine, either alone or in combination with HCT, increases twitcher mouse life span (LeVine S., *et al.*, 2000, *J. Neurosci. Res.*, 60:231-6; Biswas S., *et al.*, 2003, *Neurosci. Lett.*, p347:33-6; both of which are herein incorporated as reference). L-cycloserine is too toxic for human use, unlike its enantiomer D-cycloserine, which is indicated for treatment of anxiety. Gene therapy experiments have shown the ability to generate enzyme in transfected cells and to improve lifespan in twitcher mice, either in monotherapy or combination with HCT (Lin, D., *et al.*, 2007, *Mol. Ther.*, **15**(1): p. 44-52; incorporated herein as reference). Substrate reduction, HCT, and gene therapy all provide the most significant efficacy when used in presymptomatic animals, with either no or limited impact on disease in symptomatic animals. Therefore, ERT may provide a viable option in the treatment of GLD, especially when given to pre-symptomatic patients.

Results

[0480] Systemically administered enzyme replacement therapy using a HEK 293 derived murine GALC (rmGALC; 5 mg/kg), given peripherally as multiple intraperitoneal (IP) injections, improved the life span of twitcher mice and decreased psychosine accumulation by 15% when compared against vehicle-treated animals (Table 18, Figure 31).

2017202349 10 Apr 2017

Table 18: IP administration of rmGALC improves survival in twitcher mice

Dose Group	Survival (days)			Mann-Whitney Analysis (vs. vehicle)
	Mean	Min	Max	
Untreated	42.6	39	45	0.49
Vehicle	43.2	37	48	n/a
1 mg/kg	43.0	40	46	0.61
5 mg/kg	48.9	46	54	0.0003
10 mg/kg	49.2	47	54	0.0003

[0481] Mice treated IP with rmGALC performed better in gait testing, and sciatic nerve histopathology was improved compared to untreated or vehicle-treated animals. Peripherally (IP) administered rmGALC was minimally delivered to the brain resulting in a slight decrease in brain psychosine. However, there did not appear to be any change in brain histopathology. Therefore, the results observed in twitcher mice treated with repeated weekly systemic administration (IP) of rmGALC (5 mg/kg) resulted in a survival benefit, a slight decrease in brain psychosine levels, and an improvement in gross motor function.

Single ICV and combined ICV/IP rmGALC in twitcher mice

[0482] Results indicate that the high dose ICV/IP treatment group survived on average 50 days (120 µg/5 mpk) with the vehicle treated animals surviving only 36 days (Figure 32). Mice treated with ICV rmGALC only showed a dose-responsive mean survival time of 42 days (40 µg) and 48 days (120 µg). A single 120 µg ICV injection reduced the brain level of psychosine (63%) whereas a single ICV injection of 40 µg rmGALC resulted in a 39% decrease in psychosine (Figure 33). Although ICV/IP administration did not provide any additional benefit in psychosine reduction compared to ICV alone, the 48% observed reduction in psychosine

2017202349 10 Apr 2017

levels observed with the combined regimen was significantly lower than that observed with weekly IP treatments alone (15%). In addition, an improvement in brain histology at sites distal to the injection site was observed with ICV treatments at the 40 μ g level (Figure 34). These results confirmed the activity and biodistribution in the brain of rmGALC following direct ICV injection. However, mice treated with ICV rmGALC only failed to demonstrate restoration of sciatic nerve fiber morphology or myelination and only slight improvements in gross motor function (e.g., gait analysis). The significant improvement in key endpoints (i.e., survival, brain psychosine levels) following a single ICV administration of rmGALC suggests a lack of sufficient enzyme concentration in the systemic circulation.

Clinical Dosing Parameters: Psychosine Reaccumulation rate in twitcher Mice

[0483] The following studies were performed in the twitcher mouse model in an effort to define an appropriate clinical dose range:

- Brain psychosine re-accumulation rate in twitcher mice following a single ICV injection at PND19.
- Dose finding studies using rmGALC combined intraperitoneal (IP) + intracerebroventricular (ICV) injections in twitcher mice

[0484] In order to assess the rate of psychosine reaccumulation in the central nervous system, twitcher mice were treated with a single ICV injection of 12 μ g or 40 μ g of rmGALC at PND19. Groups of mice (n = 3) were sacrificed 24hr after the injection (PND20) and then every three days subsequently. Brain tissue was removed and submitted for psychosine analysis, histopathology, and enzyme activity analysis. A subset of animals was monitored for survival (n = 8), and motor function (gait analysis) was analyzed at PND 40.

[0485] Psychosine levels in brain homogenate following a single ICV injection was analyzed via mass spectrometry (LCMS Ltd., North Carolina), and suggests a rapid decrease in psychosine within 24hr of rmGALC administration (Figure 35). The trend of psychosine reduction was maintained for 24 day period post enzyme administration. In addition, the decrease in psychosine concentration appeared dose dependent over this period as compared with

2017202349 10 Apr 2017

vehicle-treated animals: Vehicle treated (average: 4.5 ng/ml psychosine) vs. 12 µg rmGALC (average: 2.5 mg/ml psychosine) vs. 40 µg/ml rmGALC (average: 1.6 ng/ml psychosine). Of interest, the increasing psychosine levels observed in both dose groups at the end of the study (days 28-32 post-treatment) suggests that ERT may not be successful if administered on a monthly basis. A more frequent dosing schedule may be required. Due to the small number of animals at each sampling time point, variability in the results was evident. However, based on these results, it is evident that psychosine reaccumulation occurs approximately on a 4 week (28 day) schedule.

[0486] When the survival time was analyzed, the results indicated that both the 12 µg/mL and 40 µg/mL rmGALC treatment groups had a median survival of 48 days (12 µg/mL) and 50.5 days (40 µg/mL) with the vehicle treated animals surviving 40 days (Figure 36). Unexpectedly, mice treated with 40 µg human GALC (rhGALC) showed a survival benefit only to 42 days as compared with the vehicle treated animals surviving 40 days. The reason(s) for this reduced efficacy with rhGALC is not known, but will be discussed in a later section. However, from the results of this study, it is apparent that even at lower doses of rmGALC are effective at showing a survival benefit in the twitcher mouse model.

Clinical dosing parameters: rmGALC and rhGALC dose ranging study in twitcher mice

[0487] Previous results indicated that twitcher mice treated with ICV/IP rmGALC (120 µg and 5 mpk) lived 14 days longer than vehicle-treated animals. However, twitcher mice treated only with direct CNS injections showed a dose-responsive improvement in mean survival of 12 days (120 µg ICV) and 6 days (40 µg ICV). A dose of 120 µg in the murine brain translates to a dose of 300 mg/kg brain in patients; it was therefore important to investigate the efficacy of lower doses of rmGALC. In addition, an early lot of rhGALC was examined for efficacy in the twitcher mouse. Groups of mice were treated with weekly IP injections (5 mg/kg) of rmGALC starting at PND 10 plus a single ICV injection of either 12 µg (30 mg/kg brain weight) or 26 µg (60 mg/kg brain weight) of rmGALC or rhGALC at PND19. At PND39, a subset of mice (n=3/group) were sacrificed for tissue harvest (brain, sciatic nerve, liver, sera). Brain tissue was submitted for psychosine analysis, histopathology, and enzyme activity quantification. The remaining animals survival (n = 8) were monitored for survival and gait analysis.

2017202349 10 Apr 2017

Discussion

[0488] The results of this dose finding study show a survival benefit for rmGALC administration with a trend towards dose dependence (Figure 37). The 12 $\mu\text{g}/5$ mpk and 26 $\mu\text{g}/5$ mpk combination doses of rmGALC extended the mean life span of the twitcher mouse to 44.5 and 45.5 days respectively as compared with 40.5 days for vehicle-treated animals. There was no survival benefit for the 12 $\mu\text{g}/5$ mpk (38 days) and 26 $\mu\text{g}/5$ mpk (39.5 days) doses of rhGALC. The 26 $\mu\text{g}/5$ mpk rhGALC dose extended the lifespan of the affected twitcher mice by 1.5 days, however neither dose of rhGALC reached the days of survival for the vehicle-treated animals (Figure 37). As observed previously with animals systemically-treated (IP) with rmGALC, an improvement in gait analysis was observed for all animals receiving the combined ICV/IP administration of rmGALC, while animals treated with a single ICV injection showed less benefit in motor function (Figure 38). As observed for the benefit in lifespan, no benefit in gait analysis was observed in mice treated with rhGALC. However, the specific activity of rhGALC was found to be approximately 33% of the rmGALC in vitro activity (Table 19). Therefore, these current results suggest that even at lower doses of rmGALC, there is a benefit in both survival and motor function and reinforces the opportunity for ERT for the treatment of GLD. It is evident that psychosine reaccumulation occurs approximately on a 4 week (28 day) schedule.

Table 19: rmGALC and rhGALC activity

Lot	Mean Activity ($\mu\text{mol}/\text{hr}/\text{mg}$)	% rmGALC
rmGALC R5 (3.44 mg/mL)	154.48 \pm 87.5	n/a
rhGALC Lot 73 (8.74 mg/mL)	51.35 \pm 16.2	33

2017202349 10 Apr 2017

The antigenicity of rmGALC and rhGALC is to be expected as the twitcher mouse is a null model [i.e., they are cross-reacting immunologic material (CRIM)-negative]. Overall, the maximum serum antibody titer in rhGALC-treated mice (ICV/IP regimen) was significantly higher than mice treated with a comparable ICV/IP rmGALC regimen (Figure 39). Although antibodies were also present in mice treated with direct CNS injections, the maximum titer was several fold lower than animals receiving ICV/IP treatment. The possibility exists that neutralizing antibodies may have been generated.

[0489] A study with GALC-deficient canines was initiated to characterize the antigenicity of rhGALC. In this study, affected animals (6 weeks after birth) were treated with 2 mg/kg weekly IV and/or 2.25 mg (30 mg/kg brain weight) IT administration of Human GALC or vehicle alone. Additional treatments were administered at 8 weeks and monthly for the remainder of the study (until ~16 weeks after birth). CSF was removed prior to euthanasia and analyzed for antibody formation and psychosine levels (Figure 40).

[0490] Previous studies with recombinant human heparin N-sulfatase in the Huntaway dog model of MPSIIIA demonstrated a marked antibody response to the exogenous enzyme, resulting in the need for tolerization of the animals in the study. However, preliminary results examining CSF from naïve and rhGALC-treated dogs showed an apparent reduction in psychosine levels as compared with untreated controls (Figure 40).

EXAMPLE 4: BRAIN AND LIVER HISTOLOGY/LABELING OF IT-INJECTED GALC IN MICE

[0491] The present Example describes one embodiment of IT-injected hGalC and mGalC in mice and the corresponding detection and localization of GalC antibody in various tissues.

Experimental Design

2017202349 10 Apr 2017

Experimental Design:

Group	N	Treatment	Dose (µg)	Injection volume (µL)	Route	Frequency	Sacrifice
A	6	Vehicle control	0	10 µl	IT	Three weekly injections	24hr post final injection
B	6	hGalC (Research)	100				
C	6	hGalC (PD)					
D	6	mGalC					

Tissue collection and histology staining

[0492] There were only three animals available for histological analysis from Group B and C, respectively. Samples from the brains and livers were fixed in 10% neutral buffered formalin for subsequent paraffin embedding. Five µm paraffin sections were prepared for immunohistochemistry (IHC) of I2S to detect injected proteins. Three anti-GalC antibodies were used for IHC staining of GalCA.

1. Mouse monoclonal antibody (generated by Dr. Eckman's lab)
2. Rabbit polyclonal antibody (generated by Group 1)
3. Rabbit polyclonal antibody (generated by Group 2)

A highly sensitive ABC + Tyramide fluorescence amplification method was used to label the targeted protein. The staining results showed GalC positive cells as green, with nuclei as DAPI blue counterstain, and background areas as black.

Results

2017202349 10 Apr 2017

[0493] Group 1 polyclonal antibody had a strong cross-reaction with endogenous proteins in mouse brains. Even in vehicle control brains, all CNS cells were stained strongly positive. The injected proteins were not identified with such strong background (Figure 41). Group 2 polyclonal antibody had weaker cross-reaction with endogenous proteins in mouse brains, but CNS cells in vehicle control brains were still positive. The injected proteins were not detected above the background (Figure 42). Mouse monoclonal antibody had acceptable specificity, with much lower signals in vehicle control brains (data not shown). After IT injection, all proteins were detected in the meninges on the surface of the brain. Both hGalC of Group 1 and Group 2 were detected in the CNS cells (neurons and glial cells) in the regions below the meninges, with relatively stronger signals in hGalC of Group 1 treated animals. No positive neurons and glial cells were detected in mGalC treated brains (Figure 43). In the cerebellum, hGalC produced staining in the meninges and on the surface of the granular zone, whereas mGalC did not (Figure 44). Mouse monoclonal antibody worked in the mouse brain but showed strong cross-reactivity with sinusoidal cells in the liver and could not be used to assess cellular uptake of IT injected proteins in the liver (Figure 45). Group 2 polyclonal antibody showed specificity in liver tissues with much lower signals in vehicle control brains. All IT injected proteins were detected in both sinusoidal cells and hepatocytes in the livers after treatment, with fewer positive cells and weaker signals in the hGalC of Group 1 treated animals (Figure 46). Although no higher GalC activity was found in any treated groups, positive staining was found in the meninges and the CNS cells in surrounding regions, indicating IHC is sensitive in detecting injected protein which has been taken up at the cellular level (Figure 47). In the liver, mGalC showed higher activity however IHC via Group 2 Ab detected very little difference between mGalC and hGalC (Figure 48). Low detectable activity with Group 1 Ab in hGalC was consistent with the low observed IHC levels.

Summary

[0494] After IT injection, all injected proteins were detected in the meninges of the cerebrum via IHC. Cellular uptake of injected hGalC of both Group 1 and Group 2 was detected in CNS cells (neurons and glial cells), with relatively stronger signals in hGalC of Group 1 treated brains. No positive neurons and glial cells were detected in mGalC treated brains. In the cerebellum, in addition to positive signal in the meninges, injected hGalC of both Group 1 and

2017202349 10 Apr 2017

Group 2 were found in a layer of cells on the surface of the granular zone. In the livers of all treated groups, injected proteins were detected in the sinusoidal cells and hepatocytes suggesting eventual uptake of intrathecal I2S into the circulatory system. mGalC and hGalC of Group 2 had similar strong staining signals versus hGalC of Group 1.

EXAMPLE 5: BRAIN HISTOLOGY/LABELING OF IT-INJECTED GALC IN DOGS

[0495] The present Example describes one embodiment of IT-injected GalC in dogs and the corresponding detection and localization of GalC antibody in the brain. In this embodiment, IT injected protein was detected in the meninges and in the regions of surface cortex below the meninges. ICV injected protein was found in periventricle regions (Figure 49). GalC IHC showed diffused extracellular staining pattern in the cortex after IT injection, with negative signal in neurons (circled)(Figure 50). A limited decrease of activated microglial cells with positive Iba staining was observed in ICV injected periventricle regions and IT injected cortex (Figure 51). No morphological change (Globoid cells) was found in the cortex with LFB/PAS in vehicle group and no difference was observed across the groups. Globoid cells (arrow) marked by Iba staining were decreased after ICV treatment in 4 limited areas of periventricle regions (Figure 52).

Examples of IT Delivery of I2S Protein

EXAMPLE 6: BIODISTRIBUTION OF IT DELIVERED I2S

[0496] The major objective of this study was to determine whether recombinant human I2S could be delivered to the brain of adult MPS II mice by the intrathecal-lumbar route

2017202349 10 Apr 2017

Table 20: **SIX GROUPS OF 8-12 WEEK OLD MALE MICE WERE TREATED AS FOLLOWS:**

Group	N	Strain	Treatment	Volume	Dose	Dose/Brain weight	Route
A	3	IKO	I2S	10 μ L	260 μ g	520 mg/kg	IT-lumbar
B	3	IKO	I2S	10 μ L	260 μ g	520 mg/kg	IT-lumbar
C	3	IKO	Untreated	N/A	N/A	N/A	N/A
D	1	IKO	I2S	10 μ L	260 μ g	520 mg/kg	IT-lumbar
E	3	IKO	Untreated	N/A	N/A	N/A	N/A
F	3	C57Bl/6	Untreated	N/A	N/A	N/A	N/A

Injection schedule: Animals received up to 3 injections of idursulfase (10 μ L) via the intrathecal-lumbar route:

- Groups A & D: Administered 3 doses of I2S on days 1, 8, and 15
- Group B: Administered 2 doses of I2S on days 1 and 8
- Groups C & E: Untreated control (IKO) mice
- Group F: Untreated wild-type control mice

MATERIALS AND METHODS

Animals:

[0497] Mice were housed in groups of up to 4 per cage in a colony room under a 12-hour light-dark cycle. Rodent diet (LabDiet-5001, St Louis, MO) and water (Lexington, MA municipal water purified by reverse osmosis) was available *ad libitum* for the duration of the experiment. Care of animals was conducted in accordance with the guidelines described in the Guide for the Care and Use of Laboratory Animals (National Academy Press, Washington D.C., 1996). The current IKO breeding colony was established from four carrier female mice heterozygous for the IKO mutation that were obtained from Dr. Joseph Muenzer (University of North Carolina). Carrier females were bred with male mice of the C57BL/6 background strain (C57BL/6NTac, Taconic, Hudson, NY), producing heterozygous females and hemizygous male knockout mice, as well as wild-type male and female littermates. All offspring were genotypes by PCR analysis of tissue DNA. All mice used in this experiment were males identified as either hemizygous IKO (-/0) or wild-type (WT) littermate (+/0) mice between 8 and 12 weeks of age.

2017202349 10 Apr 2017

Idursulfase:

[0498] Twenty-two mL I2S [Recombinant human idursulfase] was dialyzed against four changes of 2L phosphate buffered saline (PBS). The I2S was then concentrated by Vivaspin column and resuspended in a final volume of 1 mL PBS, followed by filter sterilization using a 0.2 µm filter. The final concentration was 51 mg/mL.

Intrathecal-lumbar Injections:

[0499] Adult mice were anesthetized using 1.25% 2,2,2 tribromoethanol (Avertin) at 200-300 µL/ 10 grams body weight (250-350 mg/kg) by intraperitoneal injection. Dorsal hair was removed between the base of the tail and the shoulder blades and the shaved area was swabbed with povidine/betadine scrub followed by isopropyl alcohol. A small midline skin incision (1-2 cm) was made over the lumbosacral spine and the intersection of the dorsal midline and the cranial aspect of the wings of the ilea (singular ileum) identified. The muscle in the iliac fossa (gluteus medius) is a heart shaped muscle and the two sides of the top of the “heart” approximate the location of the wings of the ilea. A 32 gauge needle attached to a gas tight 10-20 µL glass Hamilton syringe was inserted until resistance was felt from the underlying bone. Injection of 10 µL of test article at an approximate rate of 2 µL/20 seconds (10 µL/2 minutes) was performed. The skin incision was closed using wound clips as appropriate and the animal was allowed to recover in a recovery chamber before being returned to the appropriate cage.

Histology Procedures:

[0500] Animals were sacrificed at one hour after the final injection.

[0501] Brain and liver tissues were collected and fixed in 10% neutral buffered formalin, then processed and embedded in paraffin. Five µm sections were prepared for hematoxylin/eosin (H&E) and immunohistochemistry (IHC) staining.

Hematoxylin and Eosin Staining:

2017202349 10 Apr 2017

[0502] Brain and liver sections were stained with H&E. The staining results showed nuclei as purple and cytoplasm as pink to red. H&E stained slides were used for histopathological morphology evaluation.

Immunohistochemistry:

[0503] For I2S biodistribution evaluation, deparaffinized and rehydrated brain and liver sections were incubated overnight with mouse monoclonal antibody 2C4-2B2 (Maine Biotechnology Services, Portland, ME) against recombinant human I2S to detect injected I2S (or an irrelevant mouse IgG as a negative control antibody; Vector Laboratories, Burlingame, CA). Following an overnight incubation at 2-8°C, a secondary goat anti-mouse IgG conjugated with horseradish peroxidase was added. After additional 30 minutes of incubation at 37°C, Tyramide-Alexa Fluor 488 labeling solution (Invitrogen Corp., Carlsbad, CA) was added for an additional 10 minutes. Sections were coverslipped using an antifading mounting medium (VectaShield; Vector Laboratories) containing 1.5 µg/ml 4'-6-diamidino-2-phenylindole (DAPI) as a nuclear counterstain and observed with a multiple channel Nikon fluorescent microscope. The staining results showed I2S positive cells as green, with nuclei as blue, and background areas as black.

[0504] For efficacy analysis, brain and liver sections were stained with a rat anti-LAMP-1 (lysosomal associated membrane protein as a lysosomal marker) IgG (Santa Cruz Biotechnology, Santa Cruz, California) as the primary antibody. A rat IgG as an irrelevant antibody was used as negative control. The ABC (avidin biotin complex kits from Vector Labs, Burlingame, California) method was used to amplify the targeted marker.

[0505] Briefly, deparaffinized sections were rehydrated and incubated with the primary antibody. Following overnight incubation at 2-8°C, a secondary biotinylated rabbit anti-rat IgG (Vector Labs, Burlingame, California) was added and incubated 30 minutes at 37°C, then samples were washed and treated with avidin-biotin-peroxidase complex (Vector Laboratories) for 30 minutes. For color development, 3,3'-diaminobenzidine tetrahydrochloride (DAB) was used as the chromagen. Sections were then counterstained with hematoxylin and coverslipped. The staining results showed LAMP-1 positive cells as brown and nuclei as blue.

2017202349 10 Apr 2017

[0506] The representative photos were taken and the area of LAMP-1 positive cells was analyzed with Image-Pro Plus software (Media Cybernetics, Inc., Bethesda, MD) and comparative statistics were performed using student's t-test.

Electron Microscope Method:

[0507] Brain tissues from 3 doses of I2S treated animals were fixed in 2.5% PFA/2.5% glutaraldehyde in 0.1M sodium cacodylate buffer pH 7.4 at 4 degrees for over night. Then the samples were washed in cacodylate buffer (0.1M, pH7.4) and post-fixed in osmium tetroxide, dehydrated in alcohols and propylene oxide and embedded in Epon resin. Ultrathin sections were cut at 100nm, stained with lead citrate and examined in a Tecnai™ G² Spirit BioTWIN transmission electron microscope.

Results

[0508] In the brain as determined by immunohistochemistry (IHC), no I2S was found in vehicle control animals. In contrast, meningeal cells, neurons of the cerebrum and cerebellum were positively stained for I2S in I2S injected animals. The staining signal was stronger in animals administered 3 doses (Figure 53).

[0509] In brain tissues of vehicle-treated IKO mice, cellular vacuolation, a histopathological hallmark of lysosomal storage diseases, was found throughout brains compared to wild type animals. In I2S treated IKO mice, there was widespread reduction of cellular vacuolation from the surface cerebral cortex, caudate nucleus, thalamus, cerebellum, to the white matter compared to untreated ones (Figure 54). Abnormally high lysosomal activity was found by lysosomal-associated membrane protein-1 (LAMP-1) staining, an indicator of lysosomal activity and disease state, in microglial, meningeal and perivascular cells of vehicle-treated IKO mice when compared to wild type animals (Figure 55). The I2S intrathecal-treated mice had marked reductions in LAMP-1 immunostaining. This reduction was characterized by the decrease in the number of LAMP-1 positive cells and lighter staining. The reduction was found throughout whole brain from the surface cerebral cortex, caudate nucleus, thalamus, cerebellum to white matter (Figure 56) in both 2 and 3 dose of I2S treated animals. Morphometrical analysis of LAMP-1 immunostaining of various brain regions confirmed that there were significant reductions in the LAMP-1 positive staining in all areas of the brain evaluated (Figure 56).

2017202349 10 Apr 2017

[0510] Electron microscopy examination of brain cells in vehicle-treated IKO mice revealed the enlarged vacuoles containing amorphous granular storage material and inclusions with lamellated and zebra body-like structures. These typical pathological features of lysosomal storage at the ultrastructural level were reduced in I2S intrathecal-lumbar injected mice (Figure 57).

[0511] In the liver, there was no positive staining of I2S in the vehicle treated animals. In the I2S intrathecal injected mice, a large amount of injected I2S was clearly found in sinusoidal cells (Figure 58), which indicated the injected I2S within the intrathecal space circulated with CSF and was then absorbed through the arachnoid granulations into the circulatory system.

[0512] In liver tissues of vehicle-treated IKO mice, severe cellular vacuolation and abnormally high lysosomal activity demonstrated by H&E staining and strong LAMP-1 immunostaining were found compared to WT mice. Marked reduction of cellular vacuolation and LAMP-1 immunostaining in livers was found after intrathecal treatment with I2S. H&E staining revealed intracytoplasmic vacuolization that almost completely disappeared with a nearly normal liver cell structure (Figure 59).

[0513] In IKO mice, recombinant human I2S was delivered to the brain by the intrathecal-lumbar route and injected I2S cause widespread histopathological improvement in a variety of regions in the brain.

- Injected I2S was detected in meningeal cells and neurons in the brain.
- Reduced cellular vacuolation throughout the brain at both light and electron microscopy levels.
- Reduced LAMP-1 lysosomal marker throughout the brain.
- Intrathecal injected I2S entered the peripheral circulation and improved liver morphology and histological marker.

EXAMPLE7: TOXICOLOGY OF IT DELIVERY OF I2S

2017202349 10 Apr 2017

[0514] This example illustrates the clinical signs associated with idursulfase via monthly bolus intrathecal lumbar doses in cynomolgus monkeys. To achieve this, 14 male, cynomolgus monkeys were randomly assigned to five treatment groups as shown in the following Table 21.

Table 21: **Experimental Design**

Group	Number of Animals	Nominal Dose (mg)	Dose Volume (ml)
1	3	0	1
2	3	3	1
3	3	30	1
4	3	150	1
5	2	100	1

[0515] Animals in all groups were dosed three times at monthly intervals IT at the level of the lumbar spine. The 1 ml dose volume was flushed from the catheter system with 0.3 ml of PBS. One to two days prior to each dosing, approximately 2 ml of CSF was collected from an IT spinal tap at the level of the cisterna magna. Blood samples (2 ml) were also collected at this time. Blood (2 ml) and CSF (0.1 ml) were collected from Group 5 animals predose, 0.5, 1, 2, 4, 8, 24, and 48 hours post dose after the first dose. Clinical signs were recorded at least twice daily. A necropsy was performed approximately 24 hours after the third dose and selected tissues were harvested and saved.

[0516] On Day 1, all three Group 4 (150 mg) animals exhibited minimal tending to hind quarters within 3-12 minutes post dose, lasting 5-15 minutes; this sign was deemed related to the test article. There were no changes in body weight, food consumption and neurological/physical examination parameters that were considered related to the test article.

[0517] The analysis of the serum and CSF samples and the dosing solution analyses are presented. Variations in endogenous idursulfase activity were observed in different tissues from the cynomolgus monkey; brain and spinal cord had greater endogenous activity than other peripheral organs examined, including liver, heart, and kidney. Idursulfase administration was associated with dose-dependent increases in idursulfase activity in various brain regions, as well as in the brainstem and spinal cord. IT delivery did not result in an observable difference in distribution between the right and left cerebral hemispheres. There was a clear dose-dependent increase in idursulfase activity in the following organs: brain, liver, heart, and kidney.

Immunostaining for idursulfase in the brain demonstrated a dose-dependent increase in staining

2017202349 10 Apr 2017

intensity. In the 3 mg group, meningeal cell and limited glial cell staining beneath the meninges was observed; neuronal staining was not evident in animals from the 3 mg treatment group. Idursulfase staining was positive and dose dependent in the spinal cord, with the highest staining intensity in the lumbar region, where IT administration of idursulfase occurred. Idursulfase staining intensity in liver, kidney, and heart was dose-dependent and consistent with increased idursulfase activity in these organs.

[0518] In conclusion, IT administration of idursulfase at doses up to 150 mg delivered at monthly intervals had no adverse effects. Thus, the no observed adverse effect level (NOAEL) was interpreted to be 150 mg, the highest dose tested in this study. Idursulfase administration was associated with dose-dependent increases in idursulfase activity in the CNS and resulted in systemic levels in the liver, kidney, and heart.

[0519] The test article, idursulfase, was supplied as dosing solutions in 154 mM NaCl, 0.005% Polysorbate 20, pH 5.3 – 6.1. The nominal concentrations of the supplied dosing solutions were 0, 3, 30 or 150 mg/ml. The test article was stored in a freezer at -82° to -79 °C. Phosphate buffered saline (PBS), pH 7.2, was used as a flushing agent after the doses were administered and after serial CSF collections. The PBS was obtained from Gibco, Invitrogen Corporation.

Test Article Dosing Preparation

[0520] On the first day of dosing for each time interval, one vial of each concentration was removed from the -80°C chest freezer and allowed to thaw on the countertop to room temperature. Once thawed, the vials for Groups 1, 2, and 3 were labeled, weighed and 1 ml was withdrawn through a 0.22 µm filter for each animal scheduled for dosing. After all of the doses were administered, the vials were reweighed and placed in the refrigerator.

[0521] The following day (day of dosing for Animal 003, Group 4, and Group 5) dosing solutions for Groups 1 and 4 were removed from the refrigerator and placed on the countertop to reach room temperature. Once room temperature was obtained, the vials for Groups 1 and 4 were weighed, Group 4 vial was labeled, and 1 ml was withdrawn through the filter for each animal scheduled for dosing in Groups 1 and 4. The dosing solution for Group 5 was then prepared by injecting the appropriate amount of Group 4 dosing solution and Group 1 (vehicle) into a sterile

2017202349 10 Apr 2017

polypropylene vial. The amount added from Groups 1 and 4 were recorded. The solution was mixed by gently inverting the vial and 2-1 ml doses were withdrawn through the filter for the animals in Group 5. The vials for Groups 1 and 4 were reweighed upon completion of dosing and all the vials (Groups 1-5) were placed in a freezer.

[0522] Fourteen animals were randomly assigned to treatment groups as described in Table 21.

[0523] The IT route of administration was selected because this is an intended route for human administration. The doses of idursulfase that were selected for this study (3, 30, 100, and 150 mg/ml) were chosen to assess the biodistribution of varying doses levels of enzyme within the non-human primate central nervous system (CNS) after three consecutive monthly bolus IT lumbar injections.

Clinical Observations

[0524] The overall incidence of clinical signs was minimal. None of the animals in Group 1 (control), Group 2 (3 mg), Group 3 (30 mg), or Group 5 (100 mg) had clinical signs that were considered related to the test article at any time during the study.

[0525] On Day 1, all three Group 4 (150 mg) animals (012 -014) exhibited minimal tending to hind quarters within 3-12 minutes post dose, lasting 5-15 minutes. This sign was considered related to the test article and was not observed in any of the lower dose groups. There were no other clinical signs immediately after the first dose or on the days immediately following test article administration. The only other sign observed for the Group 4 animals was a single episode of emesis for Animal 013 on Day 35.

[0526] Administration of the test article as a single, monthly intrathecal bolus was not associated with any adverse gross or microscopic change when taking into consideration the changes inherent with an implanted drug delivery device. All groups, including the control group, had microscopic changes in the meninges indicating inflammatory reactions to the drug delivery system. In the animals that received doses of the test article of 30 mg and greater, there was a tendency for the inflammatory reaction in the meninges to have a more pronounced eosinophilic component but this difference was not considered to be biologically significant.

2017202349 10 Apr 2017

[0527] Because the differences between the control and test article treated animals were so slight, the no observed adverse effect level (NOAEL) was interpreted to be 150 mg, the highest dose tested in this study.

[0528] The overall inflammatory reaction in the meninges in all groups (including controls) was slightly more pronounced than generally encountered in an intrathecal study of this duration in monkeys. However, this was considered to possibly be related to some characteristic of the vehicle or to the act of dosing 24 hours prior to necropsy.

[0529] Brain idursulfase staining was positive in all treated animals except one animal in the 3 mg group, with the highest staining intensity found in the 150 mg group (Figures 60, 61, 62 and 63). In the 3 mg group, only meningeal cells and a few glial cells beneath the meninges were positive; no injected idursulfase was detected in neurons. In the higher dose groups (30, 100 and 150 mg), large populations of cerebral neurons were strongly positive for idursulfase staining, along with meningeal cells, glial cells and perivascular cells. Idursulfase immunostaining revealed a wide distribution of injected idursulfase in cerebral neurons from the neurons within layer I at the surface near the meninges, to the ones within the deeper layer VI adjacent to the white matter (Figures 64, 65 and 66). Marked staining of neurons was also observed for the 150 mg dose group (Figure 67). In all animals (dose group from 30 -150 mg), no marked difference in the neuronal idursulfase staining was found between frontal, middle, and rear sections of the brain.

[0530] Idursulfase staining was positive in the spinal cords of all animals, with the highest staining intensity in the lumbar region (Figures 68 and 69). Idursulfase immunostaining was also dose dependent. Neurons, meningeal cells, glial cells, perivascular cells and epi/peri/endoneurium (connective cells) surrounding nerve fibers were strongly positive for idursulfase staining in the 150 mg group (Figures 70 and 71).

[0531] In the liver, positive staining for idursulfase was found in sinusoidal cells (Kupffer cells and endothelial cells) of all animals. Idursulfase, however, was not detected in hepatocytes for the 3 mg treatment group (Figure 72), while positive idursulfase staining in the hepatocytes was found in the higher dose groups, with the greatest staining intensity in the 150 mg treatment group (Figures 73, 74 and 75).

2017202349 10 Apr 2017

[0532] There was no positive staining for idursulfase in animals from the 3 mg treatment group (Figure 76). In contrast, interstitial cells were positively stained for idursulfase in the 30, 100 and 150 mg groups, with marked staining being observed in the 150 mg group - in terms of positive cell number and staining intensity (Figures 77, 78 and 79).

Kidney

[0533] Little or no injected idursulfase was detected in animals from the 3 mg dose group (Figure 80). Positive idursulfase staining, however, was found in the glomerular cells and interstitial cells in the 30 and 100 mg groups (Figures 81 and 82). In the 150 mg group, idursulfase immunostaining additionally revealed idursulfase staining of proximal tubular cells, along with marked staining of glomerular and interstitial cells (Figure 83).

DISCUSSION

[0534] There were no test article-related clinical signs or effects on body weight, food consumption, physical examination findings and neurological examination findings. On Day 1, the Group 4 (150 mg) animals exhibited minimal tending to hind quarters within 3-12 minutes post dosing, lasting 5 to 15 minutes; this sign was judged to be related to the test article.

[0535] Idursulfase administration was associated with dose-dependent increases in idursulfase activity in various brain regions, as well as the brain stem and spinal cord. The highest level of staining intensity in the spinal cord was in the lumbar region, where IT administration of idursulfase occurred. IT administration of idursulfase also resulted in systemic exposure with dose-dependent staining intensity in the liver, kidney, and heart. Animals that received doses of the test article at 30 mg and greater had a tendency for the inflammatory reaction in the meninges to have a more pronounced eosinophilic component

[0536] IT administration of idursulfase at doses up to 150 mg delivered at monthly intervals had no adverse effects. Thus, the no observed adverse effect level (NOAEL) was interpreted to be 150 mg, the highest dose tested in this Example. Idursulfase administration was associated with dose-dependent increases in idursulfase activity in the CNS and resulted in systemic levels in the liver, kidney, and heart.

2017202349 10 Apr 2017

EXAMPLE 8: PK (Serum and CSF) of IT delivered I2S

[0537] This example provides serum and cerebrospinal fluid (CSF) analysis associated with a 6-Month Toxicity Study of Idursulfase Administered Via Monthly Bolus Intrathecal Lumbar Injections and Weekly Bolus Intravenous Injections in Cynomolgus Monkeys for test article (TA) concentration.

EXPERIMENTAL DESIGN

[0538] The objective of t was to evaluate repeat dose intrathecal (IT) administration of idursulfase (12s) from a toxicology and safety pharmacology perspective over a six month period. The study design is shown in Table 22.

Table 22: Study Design

Group No.	Number of Animals	IV Dose (mg/kg)	No. of IV Doses	IT Dose (mg)	No. of IT Doses
1	6	DC (saline)	23	DC (PBS)	6
2	12	0 (IV vehicle)	23	0 (IT vehicle)	6
3	12	0.5	23	0	6
4	6	0.5	23	30	6
5	12	0.5	23	100	6

DC = Device Control; Animals in group 1 not dosed with vehicle or test article.

Test Article

Identification: Idursulfase IV Dosing - Lot No. FDC06-001 (2.0 mg/mL)

IT Dosing - idursulfase (0 mg/mL)

idursulfase (3 mg/mL)

idursulfase (30 mg/mL)

idursulfase (100mg/mL)

Assay Methods:

[0539] Analyses were conducted using an ELISA (Enzyme Linked Immunosorbent Assay) for determining idursulfase concentration. The limit of detection (LOD) = 1.25 ng/mL prior to multiplying by dilution factor. Samples were screened at a 1:50 dilution, therefore the assay sensitivity is 62.5 ng/mL. Samples falling beyond the high end of the calibration curve

2017202349 10 Apr 2017

were further diluted and retested at an appropriate dilution that resulted in a value within the range of the curve. Selected samples were additionally analyzed using an enzyme activity assay. The LOD for this assay is 0.18 mU/mL at a minimal sample dilution of 1:150.

[0540] Animals in groups 1 and 2 that were dosed with saline or vehicle, respectively, all had serum idursulfase levels ranging between 138 ng/mL and <62.5 ng/mL (or <LOD) throughout the period of IV and IT dosing. Of 200 CSF samples tested from group 1 and 2 animals, 62 demonstrated levels of I2S above the assay LOD. Of these, 7 values were high (>1,000 ng/mL). One other CSF sample collected pre IT dose 3 tested above 1,000 ng/mL of I2S. The samples were then tested for idursulfase activity. In each case the activity results indicated the presence of I2S and when the approximate concentration of I2S was calculated based on the activity levels, the results were within 20% of those obtained by the antigen ELISA. (See Table 23) Additional randomly chosen CSF samples with antigen ELISA results <LOD were also tested using the enzyme activity assay to rule out any non-specific activity.

Table 23: Investigation Results from CSF samples

Animal Number	Group	Dose	Dose Number	Dose Mode	Time Point	ELISA Result (mg/mL)	Activity Result (mU/mL)	Calculated ng/mL Based on Activity	Calculated as % of Measured
003	1	Saline	5	IT	Predose	1392	4.7	1173	119%
003	1	Saline	6	IT	Predose	7322	29.9	7469	96%
004	1	Saline	2	IT	2 hr post	17045	62.1	15527	110%
006	1	Saline	6	IT	4 hr post	16435	70.7	17682	93%
006	1	Saline	1	IT	Predose	1320	5.3	1319	100%
0016	2	Vehicle	1	IT	2 hr post	3070	11	2743	112%
017A	2	Vehicle	mo. 3	IV	4 hr post	2236	8.8	2194	102%
046	5	100 mg/kg	3	IT	Predose	2086	7	1750	119%

[0541] In this study, serum and CSF samples were analyzed for idursulfase concentration. Serum samples were collected according to the following schedule:

IV Doses: predose and 2 hours post doses 1 through 10, predose and 4 hours post doses 11 through 23, and at necropsy.

2017202349 10 Apr 2017

IT Doses: predose and 2 hours post doses 1 and 2, predose and 4 hours post doses 3 through 6, and at necropsy.

CSF samples were collected according to the following schedule:

IV Doses: predose and 2 hours post dose 1, and 4 hours post doses 3 and 6.

IT Doses: predose and 2 hours post doses 1 and 2, predose and 4 hours post doses 3 through 6, and at necropsy.

[0542] Generally, serum idursulfase seemed to clear faster than CSF idursulfase. Serum idursulfase levels in groups 1 and 2 animals that were dosed with saline or vehicle, respectively, were less than or equal to 138 ng/mL at all time points tested. Some animals had levels below the assay limit of detection (LOD).

[0543] Fewer CSF samples from groups 1 and 2 were above the assay LOD, with 7 notable exceptions that resulted in high (>1,000 ng/mL) levels. One CSF sample collected from an animal pre IT dose 3, also tested above 1,000 ng/mL idursulfase.

[0544] The samples giving these out-of-trend results were retested and confirmed. In addition, these samples were tested for idursulfase enzyme activity. These activity results also confirmed high idursulfase levels within 20% of those obtained by the idursulfase mass assay (Table 23).

[0545] The specificity of the activity assay was validated within this sample cohort by randomly testing CSF samples with idursulfase mass units below LOD and confirmed that idursulfase levels in these samples were indeed LOD (data not shown).

EXAMPLE 9. BIODISTRIBUTION OF IT DELIVERED I2S

[0546] Having successfully demonstrated that intrathecal administration is an efficacious way of delivering I2S to the tissues of the CNS, additional studies were conducted to determine whether IT-administered I2S is capable of distributing into the deep tissues of the brain and whether there is cellular localization of IT-administered I2S. A recombinant human iduronate-2-sulfatase (I2S) formulation was prepared and formulated in a vehicle of 154 mM NaCl, 0.005% polysorbate 20 at a pH of 6.0.

2017202349 10 Apr 2017

[0547] Non-human primates were administered either 3mg, 30mg, or 100mg of I2S on a monthly basis by way of an implanted intrathecal port for six consecutive months. The design of the study is summarized in Table 24 below.

Table 24

Group	n	IV Dose (mg/kg) ^a	IT Dose (mg) ^a	Last Day on Study (number of animals)	
				6 Months	Recovery
1	6	DC (NS)	DC (PBS)	6	-
2	12	0 (vehicle)	0 (IT vehicle)	6	6
3	12	0.5	3	6	6
4	6	0.5	30	6	-
5	12	0.5	100	6	6

^aIdursulfase unless otherwise specified. DC (device control); IT (intrathecal); IV (intravenous); NS (normal saline); PBS (phosphate-buffered saline, pH 7.2).

[0548] Repeat monthly administration of I2S to the non-human primates for six months was well tolerated at the highest dose tested and not associated with any significant adverse toxicologic events. Twenty-four hours following the administration of the sixth and final dose of I2S, the subject non-human primates were sacrificed and CNS tissues of such non-human primates were examined.

[0549] As determined by immunohistochemistry (IHC), there was widespread cellular deposition of I2S throughout the cells and tissues of the CNS. I2S protein was detected in all tissues of the brain by IHC, with a deposition gradient from the cerebral cortex to the ventricular white matter. In the gray matter I2S was detected in the neurons of the cerebrum, cerebellum, brain stem, and spinal cord of all groups in a dose-dependent manner. In the surface gray matter of the higher dose groups, large numbers of cerebral neurons were positive for I2S staining in the surface cortex (Figure 84A). I2S was also detected in neurons in the thalamus (Figure 84B),

2017202349 10 Apr 2017

hippocampus (Figure 84C), caudate nucleus Figure 84D) and spinal cord (Figure 84E). Meningial and perivascular cells were also positive for I2S staining (Figure 84F).

[0550] As depicted in Figure 85 and Figure 86, distribution of IT-administered I2S into the tissues of the CNS and in particular deposition in the gray matter, thalamus and cerebral cortex of the subject non-human primates is evident. Furthermore, Figure 86 and Figure 87 illustrate that the IT-administered I2S accumulates in the depicted CNS tissues of the subject non-human primates in a dose dependant manner. Co-localization staining also revealed that IT administration of I2S associates with both neurons and oligodendrocytes. The IT-administered I2S also distributes and localizes throughout the cerebrum of the subject non-human primates as evidenced by Figure 88. In particular, Figures 89A-D illustrate neuronal uptake and axonal association of the I2S following IT-administration to the non-human primates, as demonstrated by filament staining. Also of particular interest, the present studies illustrate that I2S is selective for neuronal cells and such neuronal cells facilitate the distribution of intrathecally-administered I2S into the deep tissues of the brain and appears to be associated with axonal structures, indicating an anterograde axonal transport of I2S.

[0551] Table 25 below present the pharmacokinetic data of various administration routes and doses for a separate animal study.

Table 25

Dose	AUClast	Body weight	Brain weight	Dose	
<i>unit</i>	<i>hr*ng/mL</i>	<i>kg</i>	<i>kg</i>	<i>mg/kg BW</i>	<i>mg/kg Br wt</i>
0.5 mg/kg	8331	2.7	0.1	0.5	5
1 mg, IT	1933	3.1	0.1	0.32	10
10 mg, IT	31316	2.7	0.1	3.66	100
30 mg, IT	140345	2.9	0.1	10.34	300

2017202349 10 Apr 2017

[0552] ^{124}I -labeled I2S was administered to test animals as shown in Table 26 below and PET scan results are shown in Figure 106

Table 26

Group	Animals/ Group	Route	Test Article	Dose
1	1	ICV	[124I]- idursulfase	3 mg
2	4	IT-L	[124I]- idursulfase	3 mg
3	4	IV	[124I]- idursulfase	0.1 mg/kg
4	4	IV	[124I]- idursulfase	1 mg/kg

[0553] The present studies also demonstrated the cellular identification of IT-administered I2S in white matter brain tissue near the ventricles of the subject non-human primates following IT-administration. While the I2S staining density in the white matter was generally lower than the gray matter, I2S was detected within oligodendrocytes (Figure 90). In particular, Figure 90 illustrates the cellular identification of I2S in white matter brain tissues and further demonstrates that I2S does not appear to associate with myelin.

[0554] In addition to demonstrating the distribution of IT-administered I2S deep into the tissues of the brain, the present studies also confirmed localization of I2S into the target organelles, and importantly localization of I2S into the lysosomes which are affected organelles in the lysosomal storage disorders, such as Hunter's syndrome. In particular, I2S was located within the lysosomes and also detected within axons. Figure 90 illustrates the localization of IT-administered I2S within the lysosomes of oligodendrocytes of the subject non-human primate, thereby confirming that IT-administered I2S is capable of distributing into the deep tissues of the brain and is capable of cellular localization.

[0555] In order to discern whether the delivered I2S retained biological activity, levels of I2S in the brain were measured utilizing a specific activity assay. The activity in the brain of the 3 mg IT group 24 hours after the last dose was not apparently different from the basal levels in

2017202349 10 Apr 2017

the device control and vehicle control animals. Enzyme activity in the brain of 30 mg and 100 mg IT dosed animals was above baseline at necropsy (24 hours post-dose).

Further animal tests to discern the location of I2S delivery to the brain is shown in Figure 104 and in Table 27 below.

Table 27: Location of samples

Sample Number	Structure	Slice Number	Sample Number	Structure	Slice Number
1	Cerebral cortex-superficial (L)	4	14	Thalamus (L)	8
2	Cerebral cortex-superficial (R)	4	15	Thalamus (R)	8
3	Caudate nucleus (R)	6	16	Hypothalamus (L)	8
4	Caudate nucleus (L)	6	17	Hypothalamus (R)	8
5	Corpus callosum	6	18	Hippocampus (L)	8
6	Cerebral cortex(frontal)-superficial (L)	8	19	Hippocampus (R)	8
7	Cerebral cortex(frontal)-superficial (R)	8	20	White matter- deep (L)	10
8	White matter-superficial (L)	8	21	White matter- superficial (R)	10
9	White matter-superficial (R)	8	22	Corpus callosum	10
10	White matter- deep (L)	8	23	White matter- deep (L)	12
11	White matter- deep (R)	8	24	White matter- deep (R)	12
12	Cerebal cortex (temporal)-superficial (L)	8	25	Cerebellum (R)	14
13	Cerebal cortex (temporal)-superficial (R)	8			

2017202349 10 Apr 2017

EXAMPLE 10. BIODISTRIBUTION OF IT DELIVERY IN BEAGLE DOGS

[0556] The I2S distribution patterns observed in the foregoing example was also recapitulated in healthy Beagle dogs given a single IT or ICV dose. Male Beagle dogs were randomized using computer-generated numbers into two groups (Group 1 (ICV), N=3; Group 2 (IT); N=4). All had catheters implanted in the subarachnoid space at the lumbar spine or in the left lateral cerebral ventricle (for dosing) and in the cisterna magna (for sampling). All catheters terminated in a subcutaneous titanium access port. An additional dog was used as an un-dosed surgical control.

[0557] A single bolus 1 ml injection of I2S (30 mg/ml in 20 mM sodium phosphate, pH 6.0; 137 mM sodium chloride; 0.02% polysorbate-20), was administered IT or ICV, followed by a 0.3 ml flush with phosphate buffered saline (PBS; pH 7.2). Clinical signs were monitored and sacrifice occurred 24 hours following the dose. Brain and spinal cord tissue samples were collected for quantitative I2S analyses as determined by ELISA, I2S enzyme activity and IHC, and compared between the study groups.

[0558] I2S was widely distributed throughout the gray matter of both IT and ICV groups as determined by IHC. In the cerebral cortex, neurons were positive for I2S in all six neuronal layers, from the surface molecular layer to the deep internal layer in both IT and ICV groups, as illustrated by Figure 91 (images a and c). In the cerebellar cortex of the IT and ICV groups, I2S was detected in neurons, including Purkinje cells, as illustrated by Figure 91 (images c and d). In both IT and ICV groups a large population of neurons in the hippocampus were positive for I2S, as demonstrated by Figure 91 (images e and f). I2S positive neurons were also found in the thalamus and caudate nucleus in both of the groups, as illustrated in Figure 91 (images g and h).

[0559] The present studies therefore confirm the ability of IT-administered enzymes to distribute into the deep cells and tissues of the brain and support the utility of IT-administered enzymes such as I2S for the treatment of the CNS manifestations associated with lysosomal storage diseases, such as Hunter's syndrome.

2017202349 10 Apr 2017

EXAMPLE 11. IN VIVO EFFICACY OF IT DELIVERED I2S*Iduronate- 2-sulfatase Deficient Mouse Model*

[0560] Having demonstrated that IT-administered I2S is capable of distributing into the deep tissues of the brain and cellular localization of I2S, further studies were conducted to determine the therapeutic efficacy of IT-administered I2S. A genetically-engineered iduronate-2-sulfatase knock-out (IKO) mouse model of Hunter syndrome was developed to study the ability of the IT-administered I2S to alter disease progression. The I2S knock-out mouse model was developed using a targeted disruption of the I2S locus which results in an accumulation of glycosaminoglycans (GAG) in tissues and organs. The IKO mouse model exhibits many of the physical characteristics of Hunter syndrome seen in humans, including the characteristic coarse features and skeletal defects. In addition, the IKO mouse model demonstrates elevated glycosaminoglycan (GAG) levels in urine and in tissues throughout the body, as well as widespread cellular vacuolization which was observed histopathologically.

[0561] In the present study, commercially-available I2S (Elaprase®) was concentrated and re-suspended in phosphate buffered saline (PBS). Six groups of male IKO mice, 8-12 weeks old, were treated with I2S (10µl; 26 mg/ml). Groups A and B (N=3) were intrathecally administered three 260µg doses (at days 1, 8, and 15) and two 260µg doses (at days 1 and 8) of I2S, respectively. Group D was also treated with three intrathecally administered 260µg doses at days 1, 8, and 15. Group C and E (N=3) were untreated control groups and group F (N=3) was an untreated wild-type control. Control mice were administered a vehicle without I2S. Mice were sacrificed after 1 hour following the last injection, followed by tissue preparation for immunohistochemistry (IHC) and histopathological analysis.

[0562] Following the third injection, there was widespread reduction of cellular vacuolation in the surface cerebral cortex, caudate nucleus, thalamus and the cerebellum in I2S-treated mice compared to vehicle-treated mice. Reductions in cellular vacuolation were also found in the white matter after IT treatment. Distribution of I2S to the brain tissues of the IKO mouse was evident following IT-administration.

[0563] Three weekly IT administrations of I2S in the IKO mice also demonstrated a marked reduction in CNS cellular vacuolization at both light and electronic microscopic levels.

2017202349 10 Apr 2017

Following IT administration of I2S, a reduction of cellular vacuolation was evident relative to untreated IKO mice, suggesting that IT-administered I2S is capable of altering disease progression. As illustrated in Figure 92, a reduction of cellular vacuolation was evident in the corpus callosum and fornix of the IKO mice following IT-administration of I2S.

[0564] Additionally, electron microscopy demonstrated a reduction in the presence of storage inclusions in neurons in the gray matter and vacuolation in oligodendrocytes in the white matter. In particular, the IKO mice IT-administered I2S also demonstrated a reduction in palisaded lamellar bodies (“zebra bodies”) which are characteristic of certain lysosomal storage diseases. In particular, Figure 57 represents an electron microscope scan illustrating a reduction of the characteristic zebra bodies in the neurons of the IKO mouse that was administered I2S, relative to the untreated IKO mouse. Similarly, Figure 57 illustrates an electron microscope scan of oligodendrocytes in the corpus callosum.

[0565] In addition, the IT administrations of I2S to the IKO mice also demonstrated a marked reduction in the lysosomal disease pathological biomarker lysosomal associated membrane protein 1 (LAMP1) immunostaining, an indicator of lysosomal activity and disease state, in the surface cerebral cortex, caudate nucleus, thalamus, cerebellum and white matter. As illustrated in Figure 93A, a marked reduction in LAMP1 immunostaining is evident in the treated IKO mouse surface cerebral cortex tissue relative to the untreated IKO control mouse surface cerebral cortex tissue illustrated in Figure 93B, reflecting an improvement in disease pathology.

[0566] Figure 56 quantitatively illustrates and compares the concentration of LAMP1 measured in μm^2 areas of brain tissue. Morphometrical analysis of LAMP-1 immunostaining of various brain regions confirmed that there were significant reductions in the LAMP-1 positive staining in all areas of the brain evaluated. As shown in Figure 56, in each area of brain tissue evaluated (the cortex, caudate nucleus and putamen (CP), thalamus (TH), cerebellum (CBL) and white matter (WM)) the LAMP-positive area was reduced in the treated IKO mice relative to the untreated IKO control mice, and approached the LAMP-positive area of the wild-type mice. Particularly notable is that the LAMP-positive areas in each area of brain tissue analyzed were further reduced with continued treatment duration.

2017202349 10 Apr 2017

[0567] Reduction of abnormally high lysosomal activity correlated with dramatic morphological improvements in all areas of the brain. These results confirm that IT-administered I2S is capable of altering progression of lysosomal storage diseases, in a genetically-engineered IKO mouse model, further confirming the ability of IT-administered enzymes such as I2S to treat the CNS manifestations associated with lysosomal storage diseases, such as Hunter's syndrome.

EXAMPLE 12 - TREATMENT OF HUNTER'S DISEASE PATIENTS

[0568] Direct CNS administration through, e.g., IT delivery can be used to effectively treat Hunter's Disease patients. This example illustrates a multicenter dose escalation study designed to evaluate the safety of up to 3 dose levels every other week (EOW) for a total of 40 weeks of I2S administered via an intrathecal drug delivery device (IDDD) to patients with late infantile Hunter's Disease. Various exemplary intrathecal drug delivery devices suitable for human treatment are depicted in Figure 89 - Figure 92.

[0569] Up to 20 patients will be enrolled:

Cohort 1: 5 patients (Lowest Dose)

Cohort 2: 5 patients (Intermediate Dose)

Cohort 3: 5 patients (Highest Dose)

5 patients will be randomized to no treatment.

[0570] Patients are selected for the study based on inclusion of the following criteria: (1) appearance of first symptoms prior to 30 months of age; (2) ambulatory at the time of screening (defined as the ability to stand up alone and walk forward 10 steps with one hand held); (3) presence of neurological signs at time of screening. Typically, patients history of hematopoietic stem cell transplantation are excluded.

[0571] Safety of ascending doses of I2S administered by IT injection for 40 weeks in children with late infantile Hunter's Disease is determined. In addition, the clinical activity of

I2S on gross motor function, and single and repeated-dose pharmacokinetics in serum and concentrations in cerebrospinal fluid (CSF) are assessed.

IT Delivery of rhASA Protein

EXAMPLE 13: TOXICOLOGY STUDY OF IT-DELIVERED RECOMBINANT ASA

[0572] To assess the ability of other intrathecally-administered recombinant enzymes to distribute into the cells and tissues of the CNS, GLP study was conducted to evaluate repeat dose intrathecal (IT) administration of recombinantly-prepared human arylsulfatase A (rhASA) from a toxicology and safety pharmacology perspective over a one-month period in juvenile (less than 12 months of age) cynomolgus monkeys. The formulation of rhASA was prepared and formulated in a vehicle of 154 mM NaCl, 0.005% polysorbate 20 at a pH of 6.0.

[0573] To achieve this, nine male and nine female juvenile cynomolgus monkeys were randomly assigned by body weight to one of three treatment groups as shown in the following Table 28. The animals (with the exception of 1 male animal for Dose 1) received 0.6 mL short-term IT infusion of 0, 3 or 31 mg/mL of rhASA (total dose of 0, 1.8 or 18.6 mg) every other week for a total of three doses per animal. Body weights, clinical observations, neurological and physical examinations, clinical pathology, ophthalmologic examinations, and toxicokinetic sampling were monitored. All of the animals were necropsied on Day 29, 30 or 31 (~24 hours after the last IT dose). Selected tissues were harvested, saved and examined microscopically.

Table 28

Group	Number of Animals	Nominal Dose Concentration (mg/mL)	Dose Volume (mL)	Administered Dose (mg)
1	3M, 3F	0	0.6	0
2	3M, 3F	3	0.6	1.8
3	3M, 3F	31	0.6	18.6

2017202349 10 Apr 2017

[0574] The concentrations of rhASA detected in the CNS tissues of the cynomolgus monkeys were analyzed by ELISA and compared to a therapeutic target of 10% of normal human rhASA concentrations, corresponding to approximately 2.5ng/mg of tissue. Tissue samples or punches were extracted from different areas of the brains of the cynomolgus monkeys and further analyzed for the presence of rhASA. Figure 133 – Figure 138 illustrates the tissues from which the punches were extracted. The punched tissue samples reflected an increase in the concentrations of rhASA, as reflected in Figure 139A-G, with a deposition gradient from the cerebral cortex to the deep white matter and deep gray matter.

[0575] Concentrations of rhASA detected using the same punch from both the IT and ICV routes of administration for six monkeys administered the 18.6mg dose of rhASA, are illustrated in Figure 140A-B. The concentrations of rhASA detected in the deep white matter (Figure 140A) and in the deep grey matter (Figure 140B) brain tissues of adult and juvenile cynomolgus monkeys intrathecally- (IT) or intracerebroventricularly- (ICV) administered rhASA were comparable.

[0576] The punched tissue samples extracted from the brains of adult and juvenile cynomolgus monkeys were then analyzed to determine the concentrations of rhASA deposited in the extracted tissue sample, and to compare such concentrations to the therapeutic target concentration of 2.5ng rhASA per mg protein (corresponding to 10% of the normal concentration of rhASA in a healthy subject). As illustrated in Figure 141A, in each tissue sample punch analyzed the 18.6mg dose of IT-administered rhASA resulted in an rhASA concentration which exceeded the target therapeutic concentration of 2.5ng/mg of protein. Similarly, when a 1.8mg dose of rhASA was IT-administered to juvenile cynomolgus monkeys, each tissue sample punch analyzed demonstrated a concentration of rhASA either within or exceeding the therapeutic concentration of 2.5ng/mg of protein and the median rhASA concentrations were above the therapeutic target for all tissue punches tested (Figure 141B).

[0577] To determine whether IT-administered rhASA was distributing to the relevant cells, tissue was analyzed from the deep white matter of a cynomolgus monkey IT-administered 1.8mg of rhASA, from the area illustrated in Figure 142A. Immunostaining of the deep white matter tissue revealed distribution of rhASA in the cynomolgus monkey in oligodendrocyte cells,

as illustrated by Figure 142B. Similarly, Figure 142C illustrates that the IT-administered rhASA demonstrated co-localization in the deep white matter tissues of the cynomolgus monkey. In particular, under staining co-localization in target organelles, such as the lysosome, is evident (Figure 142C), supporting the conclusion that IT-administered rhASA is capable of distributing to the relevant cells, tissues and organelles of the CNS, including the lysosomes of oligodendrocytes.

EXAMPLE 14. ICV- VS. IT- ADMINISTRATION

[0578] rhASA labeled with the positron emitter ^{124}I was prepared and formulated in a vehicle of 154 mM NaCl, 0.005% polysorbate 20 at a pH of 6.0. A volume of the formulation equivalent to 3mg of rhASA (corresponding to approximately 38mg/kg of brain) was administered to adult cynomolgus monkeys via intracerebroventricular (ICV) and intrathecal (IT) routes of administration. The cynomolgus monkeys were subject to high-resolution PET scan imaging studies (microPET P4) to determine distribution of the administered ^{124}I -labeled rhASA.

[0579] PET imaging data (Figure 143) illustrates that both the ICV- and IT-administered ^{124}I -labeled rhASA effectively distributed to the tissues of the CNS, and in particular the ^{124}I -labeled rhASA administered through the IT-lumbar catheter immediately and uniformly spread in the cerebrospinal fluid (CSF) over the length of the spine. In particular, as depicted in Figure 143, following ICV- and IT-administration, therapeutic concentrations of ^{124}I -labeled rhASA were detected in the CNS tissues of the subject cynomolgus monkey, including the brain, spinal cord and CSF. The concentrations of rhASA detected in such CNS tissues, and in particular in the tissues of the brain, exceeded the therapeutic target concentration of 2.5ng/mg of protein.

[0580] While the distribution of rhASA protein was comparable for both IT and ICV routes of administration, ICV resulted in notably less deposition within the spinal column, as evidence by Figure 143.

[0581] Twenty four hours following administration of the formulation, both the ICV- and IT-administered ^{124}I -labeled rhASA effectively distributed to the tissues of the CNS. In particular, twenty four hours following IT-administration 12.4% of the administered dose was in the cranial region, compared to 16.7% of the ICV-administered dose. Accordingly, the

2017202349 10 Apr 2017

concentrations of rhASA detected in such CNS tissues, and in particular in the tissues of the brain, when rhASA was administered IT approached those concentrations detected following ICV-administration of the same dose.

[0582] ICV injection of the ^{124}I -labeled rhASA results ICV injection results in the immediate transfer of the injected volume to the cisterna magna, cisterna pontis, cisterna interpeduncularis and proximal spine, as illustrated in Figure 144. As also illustrated in Figure 144, within 2-5 hr IT administration delivered the ^{124}I -labeled rhASA to the same initial compartments (cisternae and proximal spine) as shown for the ICV administration. Twenty four hours following both ICV- and IT-administration distribution of the ^{124}I -labeled rhASA was comparable in the cisternal and proximal spine area, as illustrated in Figure 145.

[0583] These results confirm that rhASA can be delivered to a subject using the less invasive IT route of administration and thereby achieve therapeutic concentrations in target cells and tissues.

[0584] The lysosomal storage diseases represent a family of genetic disorders caused by missing or defective enzymes which result in abnormal substrate accumulation. While the peripheral symptoms associated with several of these diseases can be effectively mitigated by intravenous administration of recombinant enzymes, intravenous administration of such recombinant enzymes are not expected to significantly impact the CNS manifestations associated with a majority of the lysosomal storage disease. For example, recombinant human iduronate-2-sulfatase (Idursulfase, Elaprase®; Shire Human Genetic Therapies, Inc. Lexington, MA) is approved for treatment of the somatic symptoms of Hunter syndrome but there is no pharmacologic therapy for the treatment of the neurologic manifestations of Hunter Syndrome, which can include delayed development and progressive mental impairment. This is in part due to the nature of I2S, which is a large, highly-glycosylated enzyme with a molecular weight of approximately 76kD and that does not traverse the blood brain barrier following intravenous administration.

[0585] The present inventors have therefore undertaken a program to investigate the intrathecal (IT) delivery of intrathecal formulations of recombinant human enzymes, such as, for example, iduronate-2-sulfatase (I2S), arylsulfatase A (rhASA) and alpha-N-

2017202349 10 Apr 2017

acetylglucosaminidase (Naglu). The results presented herein represent the first to demonstrate that IT-lumbar administration of a recombinant lysosomal proteins result in the delivery of a significant fraction of the administered protein to the brain and in particular result in the widespread deposition of such proteins in neurons of the brain and spinal cord in both cynomolgus monkeys and dogs. Immunohistochemical analyses of the CNS tissues demonstrated that the protein is targeted to the lysosome, the site of pathologic glycosaminoglycan accumulation in the lysosomal storage disorders. Furthermore, the morphologic improvements demonstrated in the IKO mouse model of Hunter syndrome, the Naglu-deficient mouse model of Sanfilippo syndrome type B, and the ASA knockout mouse model of metachromatic leukodystrophy (MLD) reinforces the observation that IT-administered enzyme is distributed to the appropriate tissues and transported to the appropriate cellular compartments and organelles.

[0586] The similarities observed in brain distribution patterns detected after IT-lumbar and ICV administration of I2S is suggestive of bulk flow and active remixing of the CSF. Thus in a clinical setting, both the IT and the ICV administration routes are potentially feasible, however, the observed deposition of I2S in the spinal cord following IT administration provides a clear advantage in addressing spinal sequelae and components of lysosomal storage diseases such as Hunter syndrome. Moreover, spinal injection ports are less invasive and expected to be more suitable for chronic use, especially in pediatric subjects.

[0587] Evidence from perivascular cell staining and protein translocation dynamics observed by the foregoing PET imaging studies indicate that enzyme moves within the perivascular space, presumably by pulsation-assisted convective mechanisms. An additional mechanism of transport is suggested by the observed association of I2S with neurofilaments, indicative of active axonal transport. The latter presumably begins with protein interaction with neuronal mannose-6-phosphate (M6P) receptors, which are widely expressed on cells of the spinal cord and brain and which upon direct administration to the brain parenchyma may cause I2S enzyme to be readily absorbed by target cells. (Begley, et al., *Curr Pharm Des* (2008) 14: 1566-1580).

[0588] While axonal transport of lysosomal enzymes have previously been implied by indirect methods *in vivo* and by imaging *in vitro*, the current studies provide the first direct

2017202349 10 Apr 2017

evidence of axonal transport of non-virally or expressed enzymes delivered via the CSF. Thus, protein delivery from the CSF to the brain surface and deeper into the brain tissues seems to depend on active transfer processes, none of which have been previously described or elucidate for protein or enzyme delivery to the cells, tissues and organelles of the brain.

[0589] Contrary to the prevailing viewpoint that the flow dynamics of the parenchyma interstitium and CSF would prevent the distribution of IT-lumbar administered proteins to the white matter of the brain, the instant studies clearly demonstrate that IT delivery of a lysosomal enzyme results in protein distribution and accumulation in all brain tissues and deposition in the lysosomal compartment of target cells which are the site of pathologic glycosaminoglycan accumulation. (See, e.g., Fenstermacher et al., *Ann N Y Acad Sci* (1988) 531:29-39 and DiChiro et al., *Neurology* (1976) 26:1-8.) Together with the less invasive nature of IT-lumbar delivery, this route offers a clinically relevant means of delivering biologic therapeutics to the brain, particularly in children.

EXAMPLE 15- TOXICOLOGY

[0590] This example illustrate repeat dose intrathecal (IT) administration of rhASA from a toxicology and safety pharmacology perspective over a six-month period. The IT test article for this study was rhASA. Thirty-six male and 36 female cynomolgus monkeys were randomly assigned to five treatment groups. The animals in Group 1 were untreated implant device control (port & catheter) and were not dosed with the vehicle or test article; however, these animals were dosed with 0.6 mL of PBS on a schedule matching the test article dosing schedule. The animals in Groups 2-5 received 0.6 mL IT infusion of 0, 3, 10 or 31 mg/mL of rhASA (total dose of 0, 1.8, 6.0, or 18.6 mg) every other week (i.e. a total of 12 doses). Animals were necropsied at 6 months (24 hours post last IT dose), and the remaining 4 animals/sex/group were necropsied at the end of a 4-week recovery period. Selected tissues were harvested, saved and examined microscopically.

[0591] In general, the test article related changes could be categorized into two major types and were present at all dose levels (1.8, 6.0 and 18.6 mg/dose). Increase of infiltrates (of

2017202349 10 Apr 2017

white blood cells, usually with a prominent eosinophilic component) in the meninges, the brain parenchyma, the spinal cord parenchyma, trigeminal ganglion, and occasionally the spinal nerve roots/ganglia (or the epineurium surrounding those structures). This increase was interpreted to be due to the presence of the test article (a protein) in the intrathecal space and in the nervous system tissues. Slight, focal increase of microglial cells in the spinal cord and brain in occasional animals (microgliosis was not observed in any high dose animals). Both categories of morphologic changes were interpreted to be a response to the presence of the test article. There was no evidence of neuronal necrosis in any animal. None of the test article related changes were related to any biologically adverse reactions in the brain, spinal cord, spinal nerve roots or ganglia. Specifically, there was no evidence of neuronal necrosis or a biologically important glial response. There were no test article related lesions in the non-nervous system tissues.

[0592] Following a one-month recovery period (a dosing free period), the test article related changes had either entirely resolved or were limited to remnants of the prior increase in the inflammatory response associated with the presence of the test article. There were no adverse morphologic effects in the recovery animals. As based on a blinded microscopic examination assigning a semi-quantitative staining score, immunohistochemical staining for Arylsulfatase A (rhASA; the test article) was increased in the brain and spinal cord in various cell types, except neurons, for all test article treated groups at the terminal sacrifice. This increase was also apparent in the Kupffer cells of the liver. Following the 1-month recovery period, rhASA staining in the test article treated animals (all dose groups) had returned to control (device and/or vehicle control) levels. In one low dose recovery male, there were multiple foci of astrocytosis and neuronal loss, indicating multiple areas of prior ischemia, in the cerebral cortex. Although the exact pathogenesis of these lesions in this animal was not apparent, the lack of similar lesions in any other test article treated animals, including the high dose animals that received 10X the dose, indicated these lesions were not related to the test article.

[0593] The IT test article for this study was rhASA. Thirty-six male and 36 female cynomolgus monkeys were randomly assigned to five treatment groups. The animals in Group 1 were untreated implant device control (port & catheter) and were not dosed with the vehicle or test article; however, these animals were dosed with 0.6 mL of PBS on a schedule matching the test article dosing schedule. The animals in Groups 2-5 received 0.6 mL IT infusion of 0, 3, 10 or

2017202349 10 Apr 2017

31 mg/mL of rhASA (total dose of 0, 1.8, 6.0, or 18.6 mg) every other week (i.e. a total of 12 doses). Animals were necropsied at 6 months (24 hours post last IT dose), and the remaining 4 animals/sex/group were necropsied at the end of a 4-week recovery period. Selected tissues were harvested, saved and examined microscopically. The table below reflects the study design as it pertained to the pathology aspect of this study.

[0594] At the time of sacrifice, the brain was cut in a brain matrix at approximately 3 mm coronal slice thickness. The first slice and every second slice thereafter were fixed in formalin for histopathological evaluation and immunohistochemical analysis. The brain was processed as full coronal sections. These sections included at a minimum the following brain regions.

- Neocortex (including frontal, parietal, temporal and occipital cortex): brain sections 1 to 8 (and slice 9 when present)
- Paleocortex (olfactory bulbs and/or piriform lobe): brain sections 1 to 3
- Basal ganglia (including caudate and putamen): brain sections 3 and 4
- Limbic system (including hippocampus and cingulate gyri): brain sections 4 and 5
- Thalamus/hypothalamus and midbrain regions including substantia nigra: brain sections 4 and 5
- Cerebellum, pons and medulla oblongata: brain sections 6 to 8 (and slice 9 when present) .

[0595] The brain sections are listed in the data tables as sections 1 to 8/9 (a section 9 was provided by the testing facility for some animals). Sectioning varied slightly between animals. The brain sections (1 through 8/9) provided above were the approximate location of the various anatomic areas. The brain sections are listed in the data tables as individual sections, with diagnoses pertinent to that section, to facilitate potential, future additional slide review (if any). During data interpretation, individual brain anatomic sites (as listed above) were compared in order to identify any unique test article effects (i.e. unique to a particular brain region). At TPS, all brain sections from all animals were embedded in paraffin, sectioned at 5 microns, stained with hematoxylin and eosin (H&E) and examined microscopically. In addition, brains from the control and high dose animals were stained with Fluoro-Jade B (a stain increasing the sensitivity of evaluating the brain for neuronal degeneration) and a Bielschowsky's silver stain (a procedure that allows for direct visualization of axons, dendrites and neuronal filaments) and examined.

[0596] The spinal cord (cervical, thoracic and lumber) was cut into one centimeter sections. The first slice and every other slice thereafter were fixed in formalin for

2017202349 10 Apr 2017

histopathological evaluation and immunohistochemical analysis. The spinal cord sections (cervical, thoracic (including the catheter tip) and lumbar) from all animals were sectioned at approximately 5 microns, stained with H&E and examined with transverse and oblique sections taken at each level. Serial spinal cord sections from the control and high dose groups were additionally stained with Bielschowsky's silver stain and anti-GFAP (an immunohistochemical stain that allows for the direct visualization of astrocytes and their processes).

[0597] Dorsal spinal nerve roots and ganglion (taken at mid-cervical, mid-thoracic, and mid-lumbar) were embedded in paraffin, with serial sections stained with H&E. In addition, serial sections from the control and high dose groups were stained with Bielschowsky's silver stain.

[0598] For the sciatic, tibial and sural nerve sections from all animals: A longitudinal section of each nerve was embedded in paraffin, sectioned at approximately 5 microns and stained with H&E. A cross section of each nerve was post-fixed in osmium, embedded in Spurr's resin, sectioned at approximately 1 to 2 microns and stained with toluidine blue. Osmium post-fixation and resin embedding provides for superior preservation of the myelin in peripheral nerves and thus a more detailed examination of the nerve.

[0599] All tissues collected and gross lesions harvested at necropsy from all animals were also embedded in paraffin, stained with H&E, and examined microscopically. Histopathological processing and evaluations and immunohistochemical analyses were performed by TPS.

Methods: Arylsulfatase A (rhASA) Staining

[0600] Positive control slides were supplied by the study sponsor. The slides were liver sections from mice injected with rhASA. The positive control slides all showed ample evidence of rhASA in Kupffer cells (sinusoidal macrophages) in the liver. The positive control slides are stored with the other slides from this study. All evaluations of the rhASA stained sections were initially conducted blinded to the treatment group of the animal. This was accomplished by having the pathologist initially read the rhASA stained slides with the animal number on the label obscured (by an assistant with knowledge of the actual animal being evaluated), dictating the score (severity grade) during evaluation, and having the same assistant immediately record

2017202349 10 Apr 2017

the staining score (severity grade) into the data tables. The animal ID was then verified by both the study neuropathologist and the assistant to guarantee accurate data entry. This procedure was conducted so as to not introduce any bias into the judging of the overall intensity of staining with the immunohistochemical stain for the detection of intracellular rhASA. The relative degree of staining of neurons, meningeal macrophages, perivascular macrophages and glial cells (astrocytes and microglial cells but likely predominantly microglial cells) was graded in all the brain and spinal cord sections. The average severity scores at each brain and spinal cord level for each group was totaled (by group) and recorded as a total under the tissue heading brain, general, rhASA staining and spinal cord, general, rhASA staining.

[0601] In general, rhASA staining in neurons of the brain was a measure of the neurons in the cerebral cortex and other nuclear areas in the brain. rhASA staining in meningeal macrophages was evidence of uptake of the test article by meningeal macrophages and/or endogenous rhASA in meningeal macrophages. rhASA staining of perivascular macrophages was a measure of uptake of rhASA by macrophages in the brain/spinal cord (or endogenous rhASA), although it should be noted that the perivascular space in the brain and spinal cord (the Virchow-Robins space) is continuous with the meninges. In general, the grading of rhASA staining in the glial cells was predominantly a measure of uptake of the test article/penetration of the test article into the gray and/or white matter, especially of the cerebral cortex (the corona radiata is the white matter beneath the cerebral cortex). The rhASA staining in the white matter appeared to be in astrocytes and microglial cells.

[0602] The following grading scheme was used to score the degree of rhASA staining the various cell types (neurons, glial cells, macrophages).

Grade Explanation (% of the possible cells stained)

- 1 Less than 10%
- 2 Greater than 10 to 25%
- 3 Greater than 25 to 50%
- 4 Greater than 50 to 75%
- 5 Greater than 75%

[0603] Note this scheme is not strictly quantitative. It was used as an efficient, semi-quantitative method to assess the brain and spinal cord for the degree of staining with rhASA. It was noted by the Study Neuropathologist that not all neuronal areas had equal rhASA staining. It

2017202349 10 Apr 2017

was also noted that there was endogenous neuronal staining in some control animals and that cells of the choroid plexus and neurons of the dorsal root ganglia tended to stain strongly for rhASA even in control animals. Staining of the choroid plexus and dorsal root ganglia was not graded but was noted by the study neuropathologist to be prominent, even in control animals.

[0604] Note: All dose groups: Low Dose = 1.8 mg/dose; Mid dose = 6.0 mg/dose; High dose = 18.6 mg/dose. There were no test article related lesions in the non-nervous system tissues except for increased rhASA staining in the liver of all dose groups (male and female; see below).

TERMINAL SACRIFICE ANIMALS (6 MONTHS OF EVERY OTHER WEEK DOSING): rhASA STAINED SECTIONS

[0605] There was an increase of rhASA staining in the following tissues/cell types. When considering a test article effect on the degree of rhASA staining in a particular cell type in a particular dose group, the staining levels in the concurrent vehicle control and the device control (sacrificed with the recovery sacrifice animals) were considered for comparison.

[0606] Brain, Meninges, Macrophages (all dose groups, males and females)

- Brain, Perivascular, Macrophages (all dose groups, males and females)
- Brain, Glial Cells (all dose groups, males and females)
- Spinal Cord, Meninges, Macrophages (all dose groups, males and females)
- Spinal Cord, Perivascular, Macrophages (all dose groups, males and females)
- Spinal Cord, Glial Cells (mid and high dose males and females)
- Liver, Kupffer Cells (all dose groups, males and females)

[0607] Because of endogenous staining, ARSA staining levels in the neurons of the brain and spinal cord were the most difficult to specifically define. The rhASA staining demonstrated consistently increased levels of rhASA in the meningeal and brain/spinal cord perivascular macrophages and also within glial cells. There were no detectable differences of rhASA staining in neurons between the control and test article treated animals.

RECOVERY SACRIFICE ANIMALS (6 MONTHS OF EVERY OTHER WEEK DOSING FOLLOWED BY A ONE-MONTH PERIOD WITHOUT DOSING)

2017202349 10 Apr 2017

[0608] In general, test article related changes were either totally resolved or were notably diminished in those animals allowed a one-month period without dosing prior to necropsy. The following microscopic changes were present at an incidence and/or severity that indicated a possible relationship to the test article.

[0609] Test Article Related Microscopic Changes (Recovery Animals)

- Brain, Meninges, Infiltrates (mid and high dose groups, both sexes) (Figure 111 and Figure 112)
- Brain, Meninges, Infiltrates, % Eosinophils (mid dose males; high dose females)
- Brain, Perivascular, Infiltrates (mid dose males; high dose females) (Figure 113)
- Brain, Perivascular, Infiltrates, % Eosinophils (mid dose males; high dose females)
- Brain, Gray Matter, Infiltrates (all dose groups, both sexes)
- Brain, Gray Matter Infiltrates, % Eosinophils (low dose males)
- Brain, Gray Matter, Eosinophils, Necrosis (low dose males)
- Spinal Cord, Meninges, Infiltrates (mid and high dose males; low and high dose females)
- Spinal Cord, Meninges, Infiltrates, % Eosinophils (mid dose males; low dose females)
- Spinal Cord, Gray Matter, Infiltrates (low dose females)
- Spinal Cord, Gray Matter, Infiltrates, % Eosinophils (low dose females)
- Dorsal Root Ganglion and Roots, Epineurium, Infiltrates (mid dose females)
- Spinal Nerve Roots and Ganglia, Infiltrates, Eosinophils (mid and high dose males; all doses, females)
- Trigeminal Ganglion, Infiltrates, Eosinophils (mid dose males and females)

[0610] All these changes were interpreted to represent remnants of the increased inflammatory changes noted in the terminal sacrifice animals. As in the terminal sacrifice animals, there was no evidence the increase of inflammatory cell infiltrates still present in some recovery animals represented morphologic changes that were causing any adverse effects. There were no test article related lesions in the non-nervous system tissues.

RECOVERY SACRIFICE ANIMALS (6 MONTHS OF EVERY OTHER WEEK DOSING FOLLOWED BY A ONE-MONTH PERIOD WITHOUT DOSING): ARSA STAINING

[0611] There was no indication of increased rhASA staining in the recovery males or females as compared to the device and/or vehicle controls. In the brain of the low, mid and high dose recovery males, there was actually an indication of decreased rhASA staining in some cell types (this varied among the treatment groups) as compared to the device and/or vehicle controls. The reason for this, including whether or not this was an actual effect, was not apparent. One

2017202349 10 Apr 2017

possible explanation would be that administration of exogenous rhASA may cause some decrease in endogenous rhASA production. A similar finding was not present in the spinal cord of the males. In the recovery males and females, staining in the liver was similar to that noted in controls.

[0612] In general, the test article related changes could be categorized into two major types and were present at all dose levels (1.8, 6.0 and 18.6 mg/dose).

[0613] Increase of infiltrates (of white blood cells, usually with a prominent eosinophilic component) in the meninges, the brain parenchyma, the spinal cord parenchyma, trigeminal ganglion, and occasionally the spinal nerve roots/ganglia (or the epineurium surrounding those structures). This increase was interpreted to be due to the presence of the test article (a protein) in the intrathecal space and in the nervous system tissues.

[0614] Slight, focal increase of microglial cells in the spinal cord and brain in occasional animals (microgliosis was not observed in any high dose animals). Both categories of morphologic changes were interpreted to be a response to the presence of the test article. There was no evidence of neuronal necrosis in any animal. Evaluation of the rhASA stained sections is pending as of the writing of this interim report. None of the test article related changes were related to any biologically adverse reactions in the brain, spinal cord, spinal nerve roots or ganglia. Specifically, there was no evidence of neuronal necrosis or a biologically important glial response. There were no test article related lesions in the non-nervous system tissues. Following a one-month recovery period (a dosing free period), the test article related changes had either entirely resolved or were limited to remnants of the prior increase in the inflammatory response associated with the presence of the test article. There were no adverse morphologic effects in the recovery animals.

[0615] As based on a blinded microscopic examination assigning a semi-quantitative staining score, immunohistochemical staining for Arylsulfatase A (rhASA; the test article) was increased in the brain and spinal cord in various cell types, except neurons, for all test article treated groups. This increase was also apparent in the Kupffer cells of the liver. Following the 1-month recovery period, rhASA staining in the test article treated animals (all dose groups) had returned to control (device and/or vehicle control) levels. In one low dose recovery male, there

2017202349 10 Apr 2017

were multiple foci of astrogliosis and neuronal loss, indicating multiple areas of prior ischemia, in the cerebral cortex. Although the exact pathogenesis of these lesions in this animal was not apparent, the lack of similar lesions in any other test article treated animals, including the high dose animals that received 10X the dose, indicated these lesions were not related to the test article. At the time of the issuance of this preliminary report, and based strictly on the gross and microscopic findings (on the paraffin embedded, hematoxylin and eosin stained sections) in this study, the no observed adverse effect level (NOAEL) was 18.6 mg.

EXAMPLE 16 - PHARMAKINETIC DATA

6 Month Animal Data

[0616] This example provides interpretive analysis for serum and CSF concentrations of rhASA and anti- rhASA serum antibodies from Northern Biomedical Research, Inc.

[0617] The objective of the example was to evaluate repeat dose intrathecal (IT) administration of rhASA from a toxicology and safety pharmacology perspective in juvenile (<12 months of age) cynomolgus monkeys. A total of 12 doses were given in a six month period. Animals were necropsied 24 hours or one-month after the last dose. The study design is shown in Table 29.

Table 29: Study Design of 047-021 (rhASA 1-09-015)

Group	No. of Animals	Study Design			
		Nominal Dose Concentration (mg/mL)	Administered Dose (mg)	No. of Animals, 6 Month Sacrifice	No. of Animals, 1 Month Recovery Sacrifice
1	4M, 4F	DC	0	-	4M, 4F
2	8M, 8F	0	0	4 M, 3 F ^a	4M, 4F
3	8M, 8F	3	1.8	4 M, 4 F	4M, 4F
4	8M, 8F	10	6.0	4 M, 4 F	4M, 4F
5	8M, 8F	31	18.6	4 M, 4 F	4M, 4F

DC = Device Control; Animals in Group 1 were not dosed with vehicle or test article.

^a Vehicle Control Animal No. 044 was sacrificed early on Day 50 due to a leaking catheter

2017202349 10 Apr 2017

Assay Methods - Antibody Analysis

[0618] Quantitation of anti- rhASA antibodies in the serum and CSF from cynomolgus monkeys was conducted using a validated method. Briefly, the assay begins by blocking a MSD streptavidin coated plate, followed by incubation with biotin-labeled rhASA. After a washing step, diluted samples, calibrators, and QCs are added to the plate and incubated. After an additional wash step, SULFO TAG-labelled drug is added and incubated. A final wash step is performed and MSD read buffer is added. Plates are read immediately. The signal data in relative luminescence units (RLU) are analyzed using SOFTMax Pro templates.

Serum and CSF Concentration

[0619] Quantitation of rhASA in the serum and CSF from cynomolgus monkeys was conducted using a validated method. The method is based on Enzyme-Linked Immunosorbent Assay (ELISA) technology. Briefly, a microtiter plate is coated with a rabbit polyclonal antibody (SH040) raised against recombinant human Arylsulfatase A (rhASA). After incubation with rhASA reference standards and test samples, bound rhASA protein is detected by horseradish peroxidase (HRP)-conjugated anti-ASA monoclonal antibody (clone 19-16-3). The plate is then incubated with a substrate for HRP, TMB peroxidase. This enzyme-substrate reaction is stopped by the addition of 2N sulfuric acid (H₂SO₄) and the absorbance of each well is measured at the absorbance wavelength 450 nm with a reference wavelength 655 nm. The concentrations of rhASA in samples are calculated using the rhASA calibration curve in the same plate.

[0620] The summary of serum concentrations of rhASA are presented in Table 30.

[0621] The summary CSF concentrations of rhASA are presented in Table 31.

[0622] The summary anti- rhASA serum antibody concentrations are presented in Table 32.

[0623] The summary anti- rhASA CSF antibody concentrations are presented in Table 33.

[0624] Incidence of antibodies by group and sex is presented in Table 36.

2017202349 10 Apr 2017

Table 30: Summary of Serum Concentration of rhASA in Cynomolgus Monkeys

Group 1: Vehicle control	Male			Female		
	Mean	SD	n	Mean	SD	n
	ng/m L	ng/m L		ng/m L	ng/m L	
Prior to Dose 2	0	0	4	0	0	4
Post Dose 2	0	0	4	0	0	4
Prior to Dose 4	0	0	4	0	0	4
Post Dose 4	0	0	4	0	0	4
Prior to Dose 6	0	0	4	0	0	4
Post Dose 6	0	0	4	0	0	4
Prior to Dose 8	0	0	4	0	0	4
Post Dose 8	0	0	4	0	0	4
Prior to Dose 10	0	0	4	0	0	4
Post Dose 10	0	0	4	0	0	4
Prior to Dose 12	0	0	4	0	0	4
Post Dose 12	0	0	4	0	0	4
Mid Recovery	0	0	4	0	0	4
Recovery Necropsy	0	0	4	0	0	4

Group 2: 0 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
	ng/m L	ng/m L		ng/m L	ng/m L	
Prior to Dose 2	0	0	8	0	0	7
Post Dose 2	0	0	8	0	0	7
Prior to Dose 4	0	0	8	0	0	7
Post Dose 4	0	0	8	0	0	7
Prior to Dose 6	0	0	8	0	0	8
Post Dose 6	0	0	8	0	0	8
Prior to Dose 8	0	0	8	0	0	8
Post Dose 8	0	0	8	0	0	8
Prior to Dose 10	0	0	8	0	0	7
Post Dose 10	0	0	8	0	0	7
Prior to Dose 12	0	0	8	0	0	7
Post Dose 12 (Prior to 6-month Necropsy)	0	0	8	0	0	8
Mid Recovery	0	0	4	0	0	4
Recovery Necropsy	0	0	4	0	0	4

2017202349 10 Apr 2017

Table 30 (con't): Summary of Serum Concentration of rhASA in Cynomolgus Monkeys

Group 3: 1.8 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
	ng/m L	ng/m L		ng/m L	ng/m L	
Time point						
Prior to Dose 2	0	0	8	0	0	8
Post Dose 2	49.2	46.8	8	40.3	27.3	8
Prior to Dose 4	0	0	8	0	0	8
Post Dose 4	0	0	8	0	0	8
Prior to Dose 6	0	0	8	0	0	8
Post Dose 6	0	0	8	0	0	8
Prior to Dose 8	0	0	8	0	0	8
Post Dose 8	0	0	8	0	0	8
Prior to Dose 10	0	0	8	0	0	8
Post Dose 10	0	0	8	0	0	8
Prior to Dose 12	0	0	8	0	0	8
Post Dose 12 (Prior to 6-month Necropsy)	0	0	8	0	0	8
Mid Recovery	0	0	4	0	0	4
Recovery Necropsy	0	0	4	0	0	4

Group 4: 6.0 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
	ng/m L	ng/m L		ng/m L	ng/m L	
Time point						
Prior to Dose 2	0	0	8	0	0	8
Post Dose 2	173.6	69.5	8	143.2	89.0	8
Prior to Dose 4	0	0	8	0	0	8
Post Dose 4	17	49	8	63.8	119.9	8
Prior to Dose 6	0	0	8	0	0	8
Post Dose 6	0	0	8	0	0	8
Prior to Dose 8	0	0	8	0	0	8
Post Dose 8	0	0	8	0	0	8
Prior to Dose 10	0	0	8	0	0	8
Post Dose 10	0	0	8	0	0	8
Prior to Dose 12	0	0	8	0	0	8
Post Dose 12 (Prior to 6-month Necropsy)	0	0	8	0	0	8
Mid Recovery	0	0	4	0	0	4
Recovery Necropsy	0	0	4	0	0	4

2017202349 10 Apr 2017

Table 30 (con't): Summary of Serum Concentration of rhASA in Cynomolgus Monkeys

Group 5: 18.6 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
	L	L		L	L	
Prior to Dose 2	0	0	8	0	0	8
Post Dose 2	348.0	272.9	8	562.3	204.3	8
Prior to Dose 4	0	0	8	0	0	8
Post Dose 4	105.7	274.6	8	172.0	141.3	8
Prior to Dose 6	0	0	8	0	0	8
Post Dose 6	20.4	38.4	8	88.6	121.4	8
Prior to Dose 8	0	0	8	0	0	8
Post Dose 8	0	0	8	54.0	89.4	8
Prior to Dose 10	0	0	8	0	0	8
Post Dose 10	0	0	8	6	18	8
Prior to Dose 12	0	0	8	0	0	8
Post Dose 12 (Prior to 6-month Necropsy)	0	0	8	0	0	8
Mid Recovery	0	0	4	0	0	4
Recovery Necropsy	0	0	4	0	0	4

Table 31: Summary of CSF Concentrations in Cynomolgus Monkeys

Group 1: Vehicle Control	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	0	0	4	0	0	4
Post Dose 2	0	0	4	0	0	4
Prior to Dose 4	0	0	4	0	0	4
Post Dose 4	0	0	4	0	0	4
Prior to Dose 6	0	0	4	0	0	4
Post Dose 6	0	0	4	0	0	4
Prior to Dose 8	0	0	4	0	0	4
Post Dose 8	0	0	4	0	0	4
Prior to Dose 10	0	0	4	0	0	4
Post Dose 10	0	0	3	0	0	4
Prior to Dose 12	0	0	3	0	0	4
Post Dose 12	0	0	3	0	0	4
Mid Recovery	0	0	3	0	0	4
Recovery Necropsy	0	0	4	0	0	4

Group 2: 0 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	0	0	6	0	0	7
Post Dose 2	0	0	5	0	0	7

2017202349 10 Apr 2017

Prior to Dose 4	0	0	5	0	0	6
Post Dose 4	0	0	5	0	0	5
Prior to Dose 6	0	0	5	0	0	5
Post Dose 6	0	0	5	0	0	5
Prior to Dose 8	0	0	5	0	0	5
Post Dose 8	0	0	5	0	0	5
Prior to Dose 10	0	0	4	0	0	5
Post Dose 10	0	0	4	0	0	5
Prior to Dose 12	0	0	4	0	0	5
Post Dose 12 (Prior to 6-month Necropsy)	0	0	5	0	0	5
Mid Recovery	0	0	2	0	0	3
Recovery Necropsy	0	0	4	0	0	4

Table 31 (con't): Summary of CSF Concentrations in Cynomolgus Monkeys

Group 3: 1.8 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	42491	59255	7	42217	47300	6
Post Dose 2	95886	22626	7	125717	61723	6
Prior to Dose 4	17664	24372	6	50829	41891	6
Post Dose 4	106783	42823	6	138400	49908	6
Prior to Dose 6	39400	50105	4	45817	38404	6
Post Dose 6	95275	12836	4	104080	37423	5
Prior to Dose 8	25799	31589	4	58086	43821	5
Post Dose 8	148750	34664	4	119200	66556	5
Prior to Dose 10	25927	31380	4	30380	30328	5
Post Dose 10	89975	29494	4	105200	44603	5
Prior to Dose 12	29746	34267	4	82780	65906	5
Post Dose 12 (Prior to 6-month Necropsy)	32030	39155	7	47331	49015	6
Mid Recovery	0	0	3	0	0	2
Recovery Necropsy	0	0	4	0	0	4
Group 4: 6.0 mg						
Group 4: 6.0 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	75203	67002	8	146979	233673	6
Post Dose 2	360000	179276	8	267667	103369	6
Prior to Dose 4	58064	77210	8	53285	73340	5
Post Dose 4	369250	241251	8	305517	152232	6
Prior to Dose 6	77253	91407	8	97987	146762	6
Post Dose 6	418600	200098	5	369000	232238	5
Prior to Dose 8	66342	80374	5	11592	23072	4

2017202349 10 Apr 2017

Post Dose 8	329400	209841	5	340500	135128	4
Prior to Dose 10	119420	148408	5	74031	104609	2
Post Dose 10	412000	149278	5	245500	161927	2
Prior to Dose 12	68651	92902	5	74577	105251	2
Post Dose 12 (Prior to 6-month Necropsy)	141833	173933	7	58986	99016	4
Mid Recovery	0	0	3	0	NA	1
Recovery Necropsy	0	0	4	0	0	4

Table 31 (con't): Summary of CSF Concentrations in Cynomolgus Monkeys

Group 5: 18.6 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	289917	291188	7	201339	250774	8
Post Dose 2	734429	298352	7	920143	448409	7
Prior to Dose 4	150238	210302	7	169895	185675	6
Post Dose 4	984857	570039	7	965167	425924	6
Prior to Dose 6	265479	252067	7	288879	226889	6
Post Dose 6	758143	102009	7	1270000	558533	6
Prior to Dose 8	190529	240081	7	196021	199396	6
Post Dose 8	1003429	538271	7	989800	585072	5
Prior to Dose 10	176297	272500	7	168864	191087	6
Post Dose 10	1013000	390673	7	773400	103717	5
Prior to Dose 12	142334	196793	5	430542	436534	6
Post Dose 12 (Prior to 6-month Necropsy)	291525	350251	7	252142	381200	6
Mid Recovery	0	0	3	0	0	2
Recovery Necropsy	0	0	4	0	0	4

Table 32: Summary of Anti- rhASA Antibody Concentration in Serum

Group 1: Vehicle control	Male			Female		
	Mean	SD	n	Mean	SD	n
Time Point	ng/mL	ng/ml		ng/mL	ng/mL	
Predose 2	0	0	4	0	0	4
Predose 4	0	0	4	0	0	4
Predose 6	0	0	4	0	0	4
Predose 8	0	0	4	0	0	4
Predose 10	0	0	4	0	0	4
Predose 12	0	0	4	0	0	4
Mid Recovery	0	0	4	0	0	4
Recovery Necropsy	0	0	4	0	0	4

Group 2: 0 mg	Male			Female		
	Mean	SD	n	Mean	SD	n

2017202349 10 Apr 2017

Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Predose 2	0	0	8	0	0	8
Predose 4	0	0	8	0	0	8
Predose 6	0	0	8	0	0	7
Predose 8	0	0	8	0	0	7
Predose 10	0	0	8	0	0	7
Predose 12	0	0	8	0	0	7
Necropsy (24 hr after last dose)	0	0	4	0	0	4
Mid Recovery	0	0	4	0	0	4
Recovery Necropsy	0	0	4	0	0	4

Group 3: 1.8 mg	Male			Female		
	Mean	SD	n	Mean	SD	N
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Predose 2	0	0	8	0	0	8
Predose 4	18409	21371	8	27648	37504	8
Predose 6	75913	64863	8	85625	79871	8
Predose 8	132163	95576	8	151900	97818	8
Predose 10	392338	606626	8	290675	186213	8
Predose 12	499438	735028	8	524438	569523	8
Necropsy (24 hr after last dose)	261625	157865	4	733550	928411	4
Mid Recovery	339250	265888	4	377175	218955	4
Recovery Necropsy	712500	110712 9	4	295525	174718	4

Table 32 (con't): Summary of Anti- rhASA Antibody Concentration in Serum

Group 4: 6.0 mg	Male			Female		
	Mean	SD	n	Mean	SD	N
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Predose 2	0	0	8	0	0	8
Predose 4	30419	30561	8	64000	89510	8
Predose 6	143693	128094	8	191750	150511	8
Predose 8	325750	190651	8	305850	224707	8
Predose 10	669125	515458	8	832188	846241	8
Predose 12	946125	651530	8	106077 5	108888 9	8
Necropsy (24 hr after last dose)	713500	598812	4	104756 8	113204 8	4
Mid Recovery	156600 0	708132	4	975500	114973 4	4
Recovery Necropsy	111325 0	554510	4	793000	991450	4

2017202349 10 Apr 2017

Group 5: 18.6 mg	Male			Female		
	Mean	SD	n	Mean	SD	N
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Predose 2	0	0	8	0	0	8
Predose 4	56873	39107	8	39994	53411	8
Predose 6	311638	237796	8	193263	208952	8
Predose 8	482875	270130	8	399363	360425	8
Predose 10	100675 0	857916	8	866875	894776	8
Predose 12	141900 0	138227 6	8	134150 0	137377 1	8
Necropsy (24 hr after last dose)	165000	147463	4	407300	268570	4
Mid Recovery	288425 0	136312 8	4	210150 0	209042 0	4
Recovery Necropsy	250425 0	111804 2	4	150600 0	152468 2	4

Table 33: Summary of Anti- rhASA Antibody Concentration in CSF

Group 1: Vehicle control	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery	0	0	4	0	0	4
Predose 2	0	0	4	0	0	4
Predose 4	0	0	4	0	0	4
Predose 6	0	0	4	0	0	4
Predose 8	0	0	4	0	0	4
Predose 10	0	0	4	0	0	4
Predose 12	0	0	3	0	0	4
Mid Recovery	0	0	3	0	0	4
Recovery Necropsy	0	0	4	0	0	4

Group 2: 0 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery	0	0	7	0	0	6
Predose 2	0	0	6	0	0	7
Predose 4	0	0	5	0	0	6
Predose 6	0	0	5	0	0	5
Predose 8	0	0	5	0	0	5
Predose 10	0	0	4	0	0	5
Predose 12	0	0	4	0	0	5
Necropsy (24 hr after last dose)	0	0	3	0	0	2

2017202349 10 Apr 2017

Mid Recovery	0	NA	1	0	0	3
Recovery Necropsy	0	0	4	0	0	4

Group 3: 1.8 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery	0	0	7	0	0	8
Predose 2	0	0	7	0	0	6
Predose 4	0	0	6	41	101	6
Predose 6	685	1317	4	632	1413	5
Predose 8	2238	2596	4	2180	4875	5
Predose 10	3393	5038	4	5560	12433	5
Predose 12	6436	8266	4	12700	28398	5
Necropsy (24 hr after last dose)	14848	12401	4	21442	32382	4
Mid Recovery	29307	40617	3	18700	283	2
Recovery Necropsy	21060	30010	3	13078	7181	4

Table 33 (con't): Summary of Anti- rhASA Antibody Concentration in CSF

Group 4: 6.0 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery	0	0	7	0	0	8
Predose 2	0	0	7	0	0	6
Predose 4	99	172	7	84	187	5
Predose 6	1117	1862	8	1473	2775	6
Predose 8	3987	5580	5	20824	27320	4
Predose 10	6600	9679	5	2715	1237	2
Predose 12	5285	7279	5	955	1237	2
Necropsy (24 hr after last dose)	16870	16350	4	63000	63000	3
Mid Recovery	66233	42238	3	16800	NA	1
Recovery Necropsy	53600	14388	3	28880	29890	4

Group 5: 18.6 mg	Male			Female		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery	0	0	7	0	0	6
Predose 2	0	0	7	0	0	8
Predose 4	102	192	7	0	0	6
Predose 6	233	351	7	1506	3234	6
Predose 8	3378	5931	7	6367	9865	6
Predose 10	16327	24035	7	19567	27542	6
Predose 12	11596	16406	5	15143	24351	6
Necropsy (24 hr after last dose)	5168	7427	4	12135	10341	4
Mid Recovery	54700	26439	3	46315	62770	2

2017202349 10 Apr 2017

Recovery Necropsy	50725	29217	4	37790	35967	4

Table 34: Serum and CSF Concentrations of rhASA, Male and Female Combined (ng/mL)

Group 1: Vehicle control	Serum rhASA (ng/mL)			CSF rhASA (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	0	0	8	0	0	8
Post Dose 2	0	0	8	0	0	8
Prior to Dose 4	0	0	8	0	0	8
Post Dose 4	0	0	8	0	0	8
Prior to Dose 6	0	0	8	0	0	8
Post Dose 6	0	0	8	0	0	8
Prior to Dose 8	0	0	8	0	0	8
Post Dose 8	0	0	8	0	0	8
Prior to Dose 10	0	0	8	0	0	8
Post Dose 10	0	0	8	0	0	7
Prior to Dose 12	0	0	8	0	0	7
Post Dose 12	0	0	8	0	0	7
Mid Recovery	0	0	8	0	0	7
Recovery Necropsy	0	0	8	0	0	8

Group 2: 0 mg	Serum rhASA (ng/mL)			CSF rhASA (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	0	0	16	0	0	13
Post Dose 2	0	0	16	0	0	12
Prior to Dose 4	0	0	16	0	0	11
Post Dose 4	0	0	16	0	0	10
Prior to Dose 6	0	0	15	0	0	10
Post Dose 6	0	0	15	0	0	10
Prior to Dose 8	0	0	15	0	0	10
Post Dose 8	0	0	15	0	0	10
Prior to Dose 10	0	0	15	0	0	9
Post Dose 10	0	0	15	0	0	9
Prior to Dose 12	0	0	15	0	0	9
Post Dose 12 (Prior to 6-month Necropsy)	0	0	15	0	0	10
Mid Recovery	0	0	8	0	0	5
Recovery Necropsy	0	0	8	0	0	8

2017202349 10 Apr 2017

Table 34 (con't): Serum and CSF Concentrations of rhASA, Male and Female Combined (ng/mL)

Group 3: 1.8 mg	Serum rhASA (ng/mL)			CSF rhASA (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	0	0	16	42365	51844	13
Post Dose 2	44.7	37.3	16	109654	45639	13
Prior to Dose 4	0	0	16	34247	36982	12
Post Dose 4	0	0	16	122592	47311	12
Prior to Dose 6	0	0	16	43250	40831	10
Post Dose 6	0	0	16	100167	27992	9
Prior to Dose 8	0	0	16	43736	40298	9
Post Dose 8	0	0	16	132333	53926	9
Prior to Dose 10	0	0	16	28401	28890	9
Post Dose 10	0	0	16	98433	37220	9
Prior to Dose 12	0	0	16	59209	58253	9
Post Dose 12 (Prior to 6-month Necropsy)	0	0	16	39092	42786	13
Mid Recovery	0	0	8	0	0	5
Recovery Necropsy	0	0	8	0	0	8

Group 4: 6.0 mg	Serum rhASA (ng/mL)			CSF rhASA (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	0	0	16	105964	157408	14
Post Dose 2	158.4	78.7	16	320429	153832	14
Prior to Dose 4	0	0	16	56226	72638	13
Post Dose 4	40.6	91.7	16	341936	203284	14
Prior to Dose 6	0	0	16	86139	113563	14
Post Dose 6	0	0	16	393800	206033	10
Prior to Dose 8	0	0	16	42009	65286	9
Post Dose 8	0	0	16	334333	169995	9
Prior to Dose 10	0	0	16	106452	130375	7
Post Dose 10	0	0	16	364429	160707	7
Prior to Dose 12	0	0	16	70344	87227	7
Post Dose 12 (Prior to 6-month Necropsy)	0	0	16	111707	151129	11
Mid Recovery	0	0	8	0	0	4
Recovery Necropsy	0	0	8	0	0	8

2017202349 10 Apr 2017

Table 34 (con't): Serum and CSF Concentrations of rhASA, Male and Female Combined (ng/mL)

Group 5: 18.6 mg	Serum rhASA (ng/mL)			CSF rhASA (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time point	ng/mL	ng/mL		ng/mL	ng/mL	
Prior to Dose 2	0	0	16	242676	264338	15
Post Dose 2	455.1	257.8	16	827286	378379	14
Prior to Dose 4	0	0	16	159311	191264	13
Post Dose 4	138.8	213.7	16	975769	488021	13
Prior to Dose 6	0	0	16	276279	231010	13
Post Dose 6	54.5	93.8	16	994385	453568	13
Prior to Dose 8	0	0	16	193064	213058	13
Post Dose 8	27.0	67.1	16	997750	531567	12
Prior to Dose 10	0	0	16	172866	228817	13
Post Dose 10	3.2	13	16	913167	319975	12
Prior to Dose 12	0	0	16	299538	365275	11
Post Dose 12 (Prior to 6-month Necropsy)	0	0	16	273348	349718	13
Mid Recovery	0	0	8	0	0	5
Recovery Necropsy	0	0	8	0	0	8

Table 35: Serum and CSF Anti- rhASA Antibody, Male and Female Combined (ng/mL)

Group 1: Vehicle control	Serum Anti- rhASA Antibody (ng/mL)			CSF Anti- rhASA Antibody (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery	0	0	8	0	0	8
Predose 2	0	0	8	0	0	8
Predose 4	0	0	8	0	0	8
Predose 6	0	0	8	0	0	8
Predose 8	0	0	8	0	0	8
Predose 10	0	0	8	0	0	8
Predose 12	0	0	8	0	0	7
Mid Recovery	0	0	8	0	0	7
Recovery Necropsy	0	0	8	0	0	8

Group 2: 0 mg	Serum Anti- rhASA Antibody (ng/mL)			CSF Anti- rhASA Antibody (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	

2017202349 10 Apr 2017

Surgery				0	0	13
Predose 2	0	0	16	0	0	13
Predose 4	0	0	16	0	0	11
Predose 6	0	0	15	0	0	10
Predose 8	0	0	15	0	0	10
Predose 10	0	0	15	0	0	9
Predose 12	0	0	15	0	0	9
Necropsy (24 hr after last dose)	0	0	8	0	0	5
Mid Recovery	0	0	8	0	0	4
Recovery Necropsy	0	0	8	0	0	8

Table 35 (con't): Serum and CSF Anti- rhASA Antibody, Male and Female Combined (ng/mL)

Group 3: 1.8 mg	Serum Anti- rhASA Antibody (ng/mL)			CSF Anti- rhASA Antibody (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery				0	0	15
Predose 2	0	0	16	0	0	13
Predose 4	23028	29871	16	21	72	12
Predose 6	80769	70467	16	656	1284	9
Predose 8	142031	93979	16	2206	3796	9
Predose 10	341506	436656	16	4597	9386	9
Predose 12	511938	635340	16	9916	20970	9
Necropsy (24 hr after last dose)	497588	666122	8	18145	22972	8
Mid Recovery	358213	226397	8	25064	29302	5
Recovery Necropsy	504013	766860	8	16499	18552	7

Group 4: 6.0 mg	Serum Anti- rhASA Antibody (ng/mL)			CSF Anti- rhASA Antibody (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery				0	0	15
Predose 2	0	0	16	0	0	13
Predose 4	47209	66899	16	93	170	12
Predose 6	167721	137276	16	1269	2205	14
Predose 8	315800	201572	16	11470	19344	9
Predose 10	750656	682110	16	5490	8143	7
Predose 12	1003450	868860	16	4048	6328	7
Necropsy (24 hr after last dose)	880534	857199	8	36640	45439	7

Mid Recovery	1270750	938646	8	53875	42430	4
Recovery Necropsy	953125	763122	8	39474	26274	7

Table 35 (con't): Serum and CSF Anti- rhASA Antibody, Male and Female Combined (ng/mL)

Group 5: 18.6 mg	Serum Anti- rhASA Antibody (ng/mL)			CSF Anti- rhASA Antibody (ng/mL)		
	Group in total			Group in total		
	Mean	SD	n	Mean	SD	n
Time Point	ng/mL	ng/mL		ng/mL	ng/mL	
Surgery				0	0	13
Predose 2	0	0	16	0	0	15
Predose 4	48433	46054	16	55	146	13
Predose 6	252450	224723	16	821	2204	13
Predose 8	441119	310702	16	4757	7781	13
Predose 10	936813	849893	16	17822	24652	13
Predose 12	1380250	1331905	16	13531	20189	11
Necropsy (24 hr after last dose)	286150	238760	8	8652	9129	8
Mid Recovery	2492875	1686472	8	51346	36819	5
Recovery Necropsy	2005125	1347857	8	44258	31114	8

Table 36: Incidence of Anti- rhASA Antibodies at Necropsy

Group	Serum Antibody-Positive Animals (positive/total tested)				CSF Antibody-Positive Animals (positive/total tested)			
	M		F		M		F	
	6-month Necropsy	Recovery Necropsy	6-month Necropsy	Recovery Necropsy	6-month Necropsy	Recovery Necropsy	6-month Necropsy	Recovery Necropsy
1 (DC)	NA	0/4	NA	0/4	NA	0/4	NA	0/4
2 (vehicle)	0/4	0/4	0/4	0/4	0/3	0/4	0/2	0/4
3 (1.8 mg IT)	4/4	4/4	4/4	4/4	4/4	3/3	3/4	4/4
4 (6.0 mg IT)	4/4	4/4	4/4	4/4	4/4	3/3	2/3	4/4
5 (18.6 mg IT)	4/4	4/4	4/4	4/4	3/4	4/4	4/4	4/4

[0625] The quantitation limit for rhASA in cynomolgus monkey serum is 39.1 ng/mL, and all serum samples from Groups 1 and 2 were below quantitation limit (BQL), see Table 30.

2017202349 10 Apr 2017

Serum levels of rhASA were tested prior to and at 24 hours after Doses 2, 4, 6, 8, 10, and 12 (6-month necropsy), midway through the recovery period, and prior to the recovery necropsy. rhASA levels were undetectable in Group 3 (1.8 mg/dose), Group 4 (6.0 mg/dose), and Group 5 (18.6 mg/dose) prior to Doses 2, 4, 6, 8, 10, and 12, After Dose 12, midway through the recovery period, and prior to the recovery necropsy. After Dose 2, the levels of rhASA in serum were dose-related. After Dose 4 (Group 3), Dose 6 (Groups 3 and 4), and Doses 8 and 10 (Groups 3 and 4 and Group 5 males), rhASA levels were undetectable. Serum levels of rhASA declined in Group 4 (6.0 mg/dose) after Dose 4 and in Group 5 (18.6 mg/dose) after Doses 4 and 6 for males and Doses 4, 6, 8, and 10 for females. This apparent decline in serum rhASA levels may be related to the increasing concentration of anti-rhASA antibodies. There were no apparent sex differences in serum levels of rhASA, given the sample variability and small group numbers in this study.

[0626] The quantitation limit for rhASA in cynomolgus monkey CSF is 19.5 ng/mL, and all CSF samples from Groups 1 and 2 were BQL, see Table 31. rhASA was detectable in CSF prior to and after Doses 2, 4, 6, 8, 10, and 12 (6-month necropsy) in all dosed groups. The levels were higher postdose (approximately 24 hours postdose) and were dose related. The levels in CSF were much greater than those in serum. There were no apparent sex differences in CSF levels of rhASA, given the sample variability and small group numbers in this study. rhASA was not detectable midway through the recovery period and prior to the recovery necropsy in all dosed groups. CSF levels at the Dose 12 (necropsy) collections for rhASA treated groups were lower than levels postdose 8 and 11. Potential reasons for lower rhASA levels at necropsy include the larger volume taken (~2.25 mL total for cell counts, chemistry, rhASA and anti-RHASA antibody) at necropsy vs. those taken at in-life dosing interval (up to 0.5 mL pre- or postdose for rhASA concentration). Additionally, some animals did not have patent catheters at necropsy, and samples were taken via a CM tap rather than via the catheter. This route consistently yielded lower rhASA concentrations as compared with sampling via the catheter. This is likely due to the limited rostrocaudal direction of CSF bulk flow that is acknowledged to occur in vertically-oriented animals like monkeys and man (e.g., it is well known that constituents of CSF exhibit marked rostrocaudal gradients throughout an individuals lifetime).

2017202349 10 Apr 2017

[0627] Anti-rhASA antibodies in serum were detected in every animal treated with RHASA at some time point, see Table 32. Animals are defined as positive for anti-rhASA antibodies if the level of anti-rhASA antibody was above the quantitation limit (78.1 ng/mL). Animals remained positive for anti-rhASA antibodies once they seroconverted. No animals were positive for anti-rhASA antibodies at the predose 2 timepoint. All rhASA animals except Male No. 026 (Group 4; 6.0 mg/dose) were positive for serum anti-rhASA antibodies at the predose 4 timepoint. Male No. 026 was positive for serum antibody at the predose 6 timepoint. In Group 5 (18.6 mg/kg), the necropsy antibody samples had lower antibody levels. This apparent decrease may be due to the presence of rhASA interfering with the assay. The titer was generally higher in the mid- and high-dose groups (6.0 and 18.6 mg/dose) than the low dose animals (1.8 mg/dose). The presence of anti-rhASA antibodies is an expected result from treating cynomolgus monkeys with a recombinant human protein¹. Given the variability in the results, there was no apparent sex differences.

[0628] All animals with detectable anti-rhASA antibodies in CSF had detectable rhASA antibodies in serum as well, with the exception of Female Nos. 049 (Group 3; 1.8 mg/dose) and 057 (Group 4; 6.0 mg/dose). The variability in the antibody concentration and incidence precludes determination of a dose response. Animals are defined as positive for anti-RHASA antibodies if the level of anti-rhASA antibody was above the quantitation limit (78.1 ng/mL)

[0629] Combined values for males and females for serum and CSF RHASA levels and for anti-RHASA antibodies are shown in Table 34 and Table 35. Combined male and female results are similar to the individual sexes, discussed above.

2017202349 10 Apr 2017

EXAMPLE 17 - EFFICACY

[0630] In this example, 11 Wild-type control (mASA +/- hASA -/-) mice were assigned to Group A and received no treatment. thirty-four (34) hASAC69S/ASA -/- mice were assigned to each of 5 dose groups and received vehicle (Group B) or rhASA at doses of 20 mg/kg (intravenous [IV]; Group C) or 0.04, 0.12, and 0.21 mg (Groups D, E, and F, respectively) on Days 1, 9, 15/16, and 22. All IV doses were administered via a tail vein. All intrathecal (IT) doses were administered as an infusion in a volume of 12 μ L at an approximate range of 2 μ L/20 seconds (Table 37).

Table 37: Study Design

Group	No. of Animals	Animal Type	Treatment	Dose	Route	Total No. of Injections	Sacrifice	Dose in mg/kg brain weight ^a
A	11	Wild-type control (mASA +/- hASA -/-)	None	NA	NA	NA	NA	NA
B	9	hASAC69S/ASA -/-	Vehicle Control	Vehicle	IT lumbar	4 (Days 1, 9, 15/16 ^b , and 22)	24 hours after the fourth dose	0
C	5		rhASA	20 mg/kg	IV (tail vein)			NA
D	5		rhASA	0.04 mg	IT lumbar			100
E	5		rhASA	0.12 mg	IT lumbar			300
F	10		rhASA	0.21 mg	IT lumbar			520

NA = not applicable; IT = intrathecal; IV = intravenous.

^a Brain weight for mice is approximately 0.0004 kg.

^b Groups C, D, and E were dosed on Day 15; Groups B and E were dosed on Day 16.

2017202349 10 Apr 2017

[0631] The ASA knockout mouse hASAC69S/ASA(-/-) is an accepted model of MLD, and has been used to test potential treatments for this disease. The intrathecal route is the intended route of administration in humans. The intravenous route of administration has been tested for this compound and a similar compound in MLD mice. An intravenous control group has been added as a positive control for histological changes expected in peripheral organs. Animals received 100, 300, or 520 mg/kg of brain weight (0.04, 0.12, 0.21 mg, respectively) of rhASA. The dose levels normalized to brain weight selected for this study correspond to doses that are planned for use in humans or have been used in toxicology studies or in previous efficacy models of lysosomal storage diseases. These doses were not expected to have any toxicity.

Receipt

Species	Mice (<i>Mus musculus</i>)
Strain	hASAC69S/ASA (-/-) mice and wild type controls
Age	Approximately 14-17 months at arrival
No. of Groups	6
No. of Animals	34 ASA knockout mice + 11 wild type controls

Following arrival, each animal was examined to assess health status.

Housing

Animals were group housed in high-temp polycarbonate filter-top cages, with CareFresh paper bedding and water bottles. Each cage was clearly labeled with a cage card indicating project, group and animal numbers, and sex. Each animal was uniquely identified using an ear punch system.

The targeted conditions for animal room environment and photoperiod were as follows:

Temperature	22°C ± 3°C
Humidity	50% ± 20%
Light cycle	12 hours light and 12 hours dark

[0632] All available wild type animals (11) were assigned to Group A and were numbered 35 through 45. ASA (-/-) hASA (+/-) animals were assigned consecutive numbers (1 through 34) as they were removed from their cages, weighed, and ear punched during acclimation. Animals were then assigned to the treatment groups using Research Randomizer

2017202349 10 Apr 2017

(www.randomizer.org) on January 3, 2011. the first 9 numbers were assigned to Group B, the next 5 to Group C, the next 5 to Group D, the next 5 to Group E, and the final 10 to Group F. Animals were assigned as follows:

Table 38: Animal Assignment

Group	N	Animal Numbers
A	11	35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45
B	9	7, 13, 17, 22, 23, 24, 28, 29, 30
C	5	6, 16, 19 ^a , 21, 32
D	5	5, 9, 14, 18, 27
E	5	1, 2, 4, 8, 11
F	10	3 ^b , 10, 12, 15, 20, 25, 26, 31, 33, 34

^a Animal No. 19 could not be located at the time of dosing.

^b Animal No. 3 died before dosing began.

Test Article and Vehicle

Test Article

Identity rhASA
Description human recombinant Arylsulfatase A (ARSA)
Storage Conditions Approximately 4°C

Vehicle

Identity rhASA Vehicle (154 mM NaCl, 0.005% polysorbate 20, pH ~6.0)
Storage Condition Approximately 4°C

Preparation of Vehicle

[0633] The vehicle was used as provided. The vehicle was warmed on the bench top (ambient). Once the vehicle was warmed, the material was mixed by gently swirling and inverting. The bottles were not vortexed or shaken. The bottle was dried before accessing the material. Any remaining vehicle was returned to the refrigerator (1°C-8°C).

Dose Formulation Preparation

[0634] rhASA was diluted with vehicle to achieve the necessary concentrations. The test article was warmed on the bench top (ambient). Once the test article was warmed, the material was mixed by gently swirling and inverting. The bottles were not vortexed or shaken.

Dyes to track injections:

2017202349 10 Apr 2017

[0635] An infrared dye (such as IRDye®, LI-COR Biosciences, Lincoln, NE) was utilized for tracking the injections. Dyes such as this have been used in intrathecal injections as a survival procedure after intrathecal administration. The dye was mixed with the test article before administration; 1 nmole of dye in 1 μ L was added to the test article. In addition to the infrared dye, 1 μ L of FD&C blue #1 (0.25%) was used for tracking injections. This blue dye is a common food additive and is generally considered safe and non-toxic.

Lumbosacral IT Injection of rhASA or vehicle

[0636] Animals in Groups B, D, E, and F received intrathecal injections on Days 1, 9, 15 or 16, and 22.

[0637] Adult mice were anesthetized using 1.25% 2,2,2 tribromoethanol (Avertin) at 200-300 μ L/ 10 grams body weight (250-350 mg/kg) by intraperitoneal injection. Dorsal hair was removed between the base of the tail and the shoulder blades using a clippers. The shaved area was cleaned with povidine/betadine scrub followed by isopropyl alcohol. A small midline skin incision (1-2 cm) was made over the lumbosacral spine, and the intersection of the dorsal midline and the cranial aspect of the wings of the ilea (singular ileum) was identified. The muscle in the iliac fossa (gluteus medius) is a heart shaped muscle. The two sides of the top of the “heart” approximate the location of the wings of the ilea. A 32-gauge needle attached to a gas tight 10-20 μ L glass Hamilton syringe was inserted until resistance was felt from the underlying bone. Injection of 10 μ L of test article, 1 μ L of infrared dye, and 1 μ L of FD&C blue #1 (total injection volume of 12 μ L) was performed at an approximate rate of 2 μ L/20 seconds (12 μ L/2 minutes). The skin incision was closed using wound clips. The success of the injection was judged by imaging to determine if the infrared dye had distributed throughout the CNS, as well as the visible blue dye. After imaging, the animal was allowed to recover in a recovery chamber.

Intravenous Injection of rhASA

[0638] Animals in Group C received intravenous injections on Days 1, 9, 15, and 22.

[0639] For IV injections, animals were anesthetized using isoflurane, if required, and were placed in a restrainer. The tail vein was dilated by warming by flicking the tail gently with the finger. The injection site was then wiped with 70% ethanol. Alternatively, the animal was

2017202349 10 Apr 2017

placed in a warm chamber (40°C) for 1-1.5 minutes. A 28- to 30-gauge needle was used to inject test material. The volume of injection was 5-10 mL/kg.

[0640] Approximately 24 hours after the fourth dose, animals in Groups B-F were euthanized. Animals were subjected to different tissue collection procedures, as detailed below. Animals in Group A were not treated; however, they were euthanized on January 27 or 28, 2011 and subjected to tissue collection procedures, as detailed below.

Serum (all animals)

[0641] A terminal blood sample (approximately 0.5 mL) was collected from all animals (Groups A-F) via retroorbital puncture under isoflurane anesthesia. A glass tube was placed in the orbit, gently penetrating the area behind the eye and thus disrupting the venous drainage located behind the eye. Blood was collected by capillary action and/or gravity flow. Following blood collection, pressure was applied to the orbit to stop the bleeding.

[0642] The whole blood samples were processed to serum and frozen at < -80° C. The serum was stored at -80° C and analyzed for antibodies.

Tissues for Light Microscopy Investigations (Groups A-F; 5 mice per group)

[0643] After blood collection, animals were euthanized via CO₂ asphyxiation. A tail snip was collected prior to perfusion and frozen for possible genotyping. The pericardial cavity was exposed. Three (3) mice per group were transcidentally perfused with heparinized saline solution (1 U/mL sodium heparin in 0.9% NaCl, sterile-filtered) chilled ice-cold and then with 4% paraformaldehyde at approximately 4°C. The brain was removed, and the abdomen was cut to expose the internal organs further. The brain and carcass were placed in paraformaldehyde, except for the tail snip which was frozen.

Tissues for Lipid Analysis (Groups A, B, and F; 6, 4, and 5 animals, respectively)

[0644] After blood collection, animals were euthanized via CO₂ asphyxiation. A tail snip was collected prior to perfusion and frozen for possible genotyping. The pericardial cavity was exposed. For lipid analyses, 4-6 mice per group were transcidentally perfused with heparinized saline solution (1 U/mL sodium heparin in 0.9% NaCl, sterile-filtered) chilled ice-cold.

2017202349 10 Apr 2017

Table 39: Tissues Collected for Lipid Analysis

Tissues Collected for Lipid Analysis	
Brain (separated into left and right hemispheres and weighed)	Kidney (2)
Spinal cord (removed from spinal column)	
Sciatic nerve (2) (dissected free from muscle)	Tail snip (prior to perfusion)

Upon collection, tissues were weighed and then frozen, either on dry ice or by placing in a -80°C freezer. The brain was separated into left and right hemispheres. The right will be utilized for lipid analysis by MS. The left will be analyzed for possible N-acetyl-L-aspartate (NAA) analysis. Tissues were stored at -80° C until analysis.

Table 40: Sample Storage Conditions

Type of Sample	Storage Temperature
Serum	frozen at <i>circa</i> -80°C
tissues for lipid analysis	frozen at <i>circa</i> -80°C
Tail snips	frozen at <i>circa</i> -80°C
Tissues for light microscopy	Approximately 4°C

[0645] rhASA reduced sulfatide storage in the spinal cord of MLD mice, particularly in the white matter, Figure 114. Morphometry analysis of the spinal cord demonstrated that the optical density of alcian blue staining was statistically significantly reduced after rhASA dosing, Figure 115. rhASA treated MLD mice also exhibited reduced lysosomal activity in the brain, Figure 116. This reduction was statistically significant in the high-dose group (0.21 mg- 520 mg/kg brain weight) compared with vehicle treated animals, Figure 117.

[0646] Immunotolerant MLD mice (hASAC69S/ASA(-/-)) over 1 year in age received intrathecal-lumbar administration of rhASA one time each week for 4 weeks (a total of 4 doses). Doses were vehicle (154 mM NaCl, 0.005% polysorbate 20, pH ~6.0), 0.04, 0.12, 0.21 mg/dose (normalized doses were 100, 300 and 520 mg/kg of brain weight, respectively). At terminal timepoints efficacy was evaluated by immunohistochemistry assessment of sulfatide clearance and lysosome activity within the brain and spinal cord. Spinal cord and brain sections were stained using alcian blue stain targeting sulfatides in tissues. Brain sections were also stained for the presence of lysosomal-associated membrane protein (LAMP), an indicator of lysosomal

processes. Additionally, morphometry analysis was performed on alcian blue and LAMP stained sections of the spinal cord (cervical, thoracic and lumbar) and brain.

[0647] These preliminary results demonstrate efficacy of intrathecal lumbar administration of rhASA. Compared to vehicle control mice, rhASA treated MLD mice exhibit evidence of improvement within the histological markers of disease, such as reduced sulfatide storage (noted by alcian blue staining) and lysosomal activity in the brain. These histopathological changes were observed near the site of administration (lumbar region of the spinal cord), in the distal spinal cord, as well as in the distal portions of the brain.

EXAMPLE 18 – BIODISTRIBUTION 2

Overview

[0648] In this study, 36 male and 36 female juvenile cynomolgus monkeys (< 12 months at initiation) were assigned to each of 5 dose groups and received rhASA at doses of 0 (device control; animals were dosed with 0.6 mL of PBS), 0 (vehicle control), 1.8, 6.0, or 18.6 mg (Groups 1, 2, 3, 4, and 5, respectively) every other week for 6 months for a total of 12 doses. All doses were administered as an infusion in a volume of 0.6 mL, followed by a flush of 0.5 mL PBS given over approximately 10 minutes (Table 41).

Table 41 : Study Design

Study Design					
Group	No. of Animals	Nominal Dose Concentration (mg/mL)	Administered Dose (mg)	No. of Animals, 6 Month Sacrifice	No. of Animals, 1 Month Recovery Sacrifice
1	4M, 4F	DC	0	-	4M, 4F
2	8M, 8F	0	0	4 M, 3 F ^a	4M, 4F
3	8M, 8F	3	1.8	4 M, 4 F	4M, 4F
4	8M, 8F	10	6.0	4 M, 4 F	4M, 4F
5	8M, 8F	31	18.6	4 M, 4 F	4M, 4F

DC = Device Control; Animals in Group 1 were not dosed with vehicle or test article.

^a Vehicle Control Animal No. 044 was sacrificed early on Day 50 due to a leaking catheter

MATERIAL AND METHODS

Tissue Collection

2017202349 10 Apr 2017

[0649] The brains were cut in a brain matrix at 3 mm thick coronal slice thickness. Each brain was sectioned into full coronal slices including: neocortex (including frontal, parietal, temporal, and occipital cortex), paleocortex (olfactory bulbs and/or piriform lobe), basal ganglia (including caudate and putamen), limbic system (including hippocampus and cingulate gyri), thalamus/hypothalamus, midbrain regions (including substantia nigra), cerebellum, pons, and medulla oblongata. The locations from which individual tissue samples were obtained (via 4-mm biopsy punch) are shown in Figures 133-138. The images in Figures 133-138 are from the University of Wisconsin and Michigan State Comparative Mammalian Brain Collections, (also the National Museum of Health and Medicine). Punch number 22 was not collected, as this structure was not present during necropsy. All brain samples were frozen and stored at -60°C or below prior to analysis for rhASA using an enzyme-linked immunosorbent assay.

[0650] The first brain slice and every second slice thereafter were fixed in formalin for histopathological evaluation and immunohistochemical. The second brain slice and every second slice thereafter were frozen for test article concentration analysis. Prior to freezing, samples of brain were taken from the right portion of the even-numbered, test article analysis brain slices for biodistribution analysis. The location of the brain samples were photographed at necropsy and the brain slice number was recorded. The samples were obtained using either a 4-mm circular punch or cut with a scalpel to optimize the amount of white matter collected. All punches were frozen and stored at -60°C or below for test article analysis. The remainder of the brain slice was frozen and stored at -60°C or below for possible test article analysis.

[0651] The spinal cord (cervical, thoracic and lumbar) was cut into one-centimeter sections. The first slice and every second slice thereafter was fixed in formalin for histopathological and immunohistochemical analysis. The second slice of spinal cord and every second slice thereafter was frozen and stored at -60°C or lower for test article analysis. The distribution of slices was adjusted so that the slice with the tip of the intrathecal catheter (Slice 0) was fixed in formalin and analyzed for histopathology.

Preparation of Brain, Liver, and Spinal Extracts and Determination of rhASA Concentration

2017202349 10 Apr 2017

[0652] Brain punches, spinal cord, and liver samples were analyzed using a validated method in compliance with the United States Food and Drug Administration (FDA) Good Laboratory Practice (GLP) regulations 21 CFR, Part 58 and with applicable Midwest BioResearch standard operating procedures. Tissue samples were homogenized in lysis buffer, centrifuged to remove any tissue debris, and stored at -80°C until assayed. rhASA concentration in the soluble fractions of the homogenates was determined by an ELISA using polyclonal rabbit antibody SH040 as the capture antibody and HRP (horseradish peroxidase)-conjugated anti-ASA monoclonal antibody 19-16-3 as the detection antibody. After a wash step to remove unbound materials, tetramethylbenzidine (TMB) substrate solution reacted with the peroxide in the presence of HRP-conjugated antibody to produce a colorimetric signal that was proportional to the amount of rhASA bound by the anti ASA antibody in the initial step. The resulting amount of rhASA in each tissue homogenate was interpolated from a standard curve.

[0653] Samples were also analyzed by a bicinchoninic acid (BCA) protein determination assay to obtain the concentration of protein in the sample. The protein concentration for each sample was determined by interpolation of an albumin standard curve. rhASA concentration results were then normalized to total protein in tissue extracts, as determined by bicinchoninic acid assay.

[0654] The rhASA levels of all punches for the vehicle, 1.8 mg/dose, 6.0 mg/dose, and 18.6 mg/dose groups are shown in Figure 118, Figure 119, Figure 120, and Figure 121, respectively. The rhASA levels of all punches for the recovery animals for the device control, vehicle, 1.8 mg/dose, 6.0 mg/dose, and 18.6 mg/dose groups are shown in Figure 122, Figure 123, Figure 124, Figure 125, and Figure 126, respectively

[0655] The rhASA levels for selected punches that were taken near the surface (meninges) of the brain are shown in Figure 127. rhASA levels for selected punches that are considered to contain mostly deep white brain matter are shown in Figure 128. White matter is composed of bundles of myelinated nerve cell processes (or axons). Selected punches which contain mostly material from the deep grey brain matter are shown in Figure 129. Grey matter contains neural cell bodies, in contrast to white matter. The values of rhASA in selected punches from the surface, deep white and deep grey are shown for each dose group in Figure 130.

2017202349 10 Apr 2017

[0656] Spinal cord concentration data is shown in Figure 131.

[0657] Liver concentration data is shown in Figure 132.

[0658] rhASA concentration levels in the liver, spinal cord, and brain of the device and vehicle-dosed control groups were in some cases measurable. The levels in liver and spinal cord were lower than any of the rhASA -treated groups. The level of rhASA measured in the device control and vehicle-dosed animals represents a cross-reactivity between the anti-rhASA antibody used in the ELISA with the native cynomolgus monkey protein. The reported values in the device control and vehicle tissues do not represent quantitative values for cynomolgus monkey rhASA in the tissues, because the degree of cross-reactivity between the antibody and cynomolgus ASA is not known, and the fact that the assay standards use rhASA. However, the variation in the levels of rhASA detected between device control and vehicle-dosed tissues is may be interpreted as demonstrated variability in the relative amounts of cynomolgus rhASA in different tissues and anatomical regions.

[0659] The rhASA levels in spinal cord slices ranged from 160–2352, 1081–6607, and 1893–9252 ng/mg protein in males and 0–3151, 669–6637, and 1404–16424 ng/mg protein in females for the 1.8, 6.0, and 18.6 mg/dose groups, respectively (Figure 127). Levels of rhASA were higher in the lumbar region of the spine than in the cervical region. Levels of rhASA protein detected in the liver were dose responsive in the rhASA treated groups and were very low in the vehicle group. Mean rhASA levels were 88, 674, and 2424 in males and 140, 462, and 1996 ng/mg protein in females for the 1.8, 6.0, and 18.6 mg/dose groups, respectively (Figure 128).

[0660] Overall, the level of rhASA appeared to be dose-related in samples prepared from the spinal cord slices and liver of the rhASA -dosed groups. Many of the brain regions tested demonstrated a clear dose relationship between rhASA levels and rhASA administration, while others were more equivocal. In general, rhASA levels in the brain increased with rhASA dose.

2017202349 10 Apr 2017

EXAMPLE19: PHARMACOKINETIC (PK) AND BIODISTRIBUTION OF IT VS. IV ADMINISTERED rhASA

[0661] The objective this study is to evaluate the pharmacokinetic (PK) and biodistribution of various therapeutic replacement enzymes after intrathecal (IT) and intravenous (IV) administration to cynomolgus monkeys.

[0662] In this study, a total of twelve male and twelve female cynomolgus monkeys with intrathecal-lumbar (IT-L) catheters were randomly assigned by body weight into four treatment groups for Phase 1a (I2S administration) and Phase 1b (rhASA administration).

[0663] Blood and CSF (from IT-CM catheter) were collected at specified intervals post dosing for both phases. After the last samples were collected from Phase 1a, the animals were allowed a 7-day washout period before initiation of Phase 1b.

[0664] After the last samples were collected from Phase 1b, the animals will be allowed a 7-day washout period between initiation of Phase 2. A total of 12 male and female cynomolgus monkeys from Phase 1b were randomly assigned by body weight into 12 treatment groups of I2S (Groups 1a-6a) and rhASA (Groups 1b-6b).

[0665] The absolute bioavailability of rhASA in serum following IT-L administration is ~30 to 40%. In contrast, only 0.5% of the IV dose is bioavailable in CSF.

[0666] Exposure to rhASA in serum increases in a more than proportional manner following IT-L administration.

[0667] Following IT-L administration, exposure to rhASA in CSF increases in a less than proportional manner as dose increases.

2017202349 10 Apr 2017

Table 42 - Summary PK Parameters of rhASA in Serum of Cynomolgus Monkeys

Mean (CV%)	Serum Arylsulfatase A			
	Arylsulfatase A (Phase 1b: IV 1 mg/kg)	Arylsulfatase A (Phase 1b: IT-L 1.8 mg)	Arylsulfatase A (Phase 1b: IT-L 6 mg)	Arylsulfatase A (Phase 1b: IT-L 18.6 mg)
N	8	6	8	8
AUC _{0-t} (ng•h/mL)	10505 (16.9)	2219 (41.9)	10352 (31.9)	17583 (28.2)
AUC _{0-∞} (ng•h/mL)	11069 (17.2)	NC (NC)b	9634 (28.9)c	20789 (27.8)d
C _{max} (ng/mL)	11911 (20.0)	363 (40.4)	1160 (29.9)	1621 (25.1)
T _{max} (h)	0.08 (0.08, 0.08)	4.00 (2.00, 4.00)	4.00 (1.00, 4.00)	3.00 (1.00, 4.00)
t _{1/2} (h)	6.55 (31.8)	NC (NC)b	6.77 (21.4)c	7.40 (32.8)d
CL or CL/F (mL/h)	261 (17.0)	NC (NC)b	654 (25.0)c	944 (25.4)d
V _z or V _z /F (mL)	2418 (32.4)	NC (NC)b	6523 (41.3)c	9686 (25.8)d

Table 43 - Summary PK Parameters of rhASA in CSF of Cynomolgus Monkeys

Mean (CV%)	CSF Arylsulfatase A			
	Arylsulfatase A (Phase 1b: IV 1 mg/kg)	Arylsulfatase A (Phase 1b: IT-L 1.8 mg)	Arylsulfatase A (Phase 1b: IT-L 6 mg)	Arylsulfatase A (Phase 1b: IT-L 18.6 mg)
N	4	6	8	8
AUC _{0-t} (ng•h/mL)	1629 (179.8)	1267266 (86.6)	5334329 (68.8)	8028775 (71.2)
AUC _{0-∞} (ng•h/mL)	8221 (NC)b	1595942 (79.1)c	4291829 (84.2)d	9406664 (64.5)e
C _{max} (ng/mL)	69.3 (94.2)	345167 (48.7)	1039079 (73.6)	1841125 (62.8)
T _{max} (h)	6.00 (1.00, 8.00)	0.08 (0.08, 4.00)	0.29 (0.08, 4.00)	2.04 (0.08, 4.00)
t _{1/2} (h)	37.6 (NC)b	23.6 (68.3)c	17.1 (31.3)d	13.4 (29.3)e
CL or CL/F (mL/h)	392 (NC)b	1.95 (74.1)c	38.1 (214.8)d *	3.04 (66.1)e
V _z or V _z /F (mL)	21237 (NC)b	80.6 (110.4)c	1090 (215.1)d	67.6 (81.2)e

2017202349 10 Apr 2017

Table 44 – Bioavailability of rhASA in Serum and CSF

Absolute Bioavailability Comparison			
	Arylsulfatase A (Phase 1b: IT-L 1.8 mg)	Arylsulfatase A (Phase 1b: IT-L 6 mg)	Arylsulfatase A (Phase 1b: IT-L 18.6 mg)
Fabs (%)	NC	39.9	27.3

[0668] The bioavailability of rhASA in serum following IT-L administration is ~30-40%. In contrast, only 0.5% of the dose administered by IV route is bioavailable in CSF.

Table 45 - CSF:Serum Partition

CSF:Plasma Partition			
Arylsulfatase A (Phase 1b: IV 1 mg/kg)	Arylsulfatase A (Phase 1b:IT-L 1.8 mg)	Arylsulfatase A (Phase 1b: IT-L 6 mg)	Arylsulfatase A (Phase 1b:IT-L 18.6 mg)
0.74	NC	445	452

EXAMPLE 20 - TREATMENT OF MLD PATIENTS

[0669] Direct CNS administration through, e.g., IT delivery can be used to effectively treat MLD patients. This example illustrates a multicenter dose escalation study designed to evaluate the safety of up to 3 dose levels every other week (EOW) for a total of 40 weeks of rhASA administered via an intrathecal drug delivery device (IDDD) to patients with late infantile MLD. Various exemplary intrathecal drug delivery devices suitable for human treatment are depicted in Figure 94, Figure 95, Figure 96 and Figure 97.

[0670] Up to 20 patients will be enrolled:

Cohort 1: 5 patients (Lowest Dose)

Cohort 2: 5 patients (Intermediate Dose)

Cohort 3: 5 patients (Highest Dose)

2017202349 10 Apr 2017

5 patients will be randomized to no treatment.

[0671] Patients are selected for the study based on inclusion of the following criteria: (1) appearance of first symptoms prior to 30 months of age; (2) ambulatory at the time of screening (defined as the ability to stand up alone and walk forward 10 steps with one hand held); (3) presence of neurological signs at time of screening. Typically, patients history of hematopoietic stem cell transplantation are excluded.

[0672] Safety of ascending doses of rhASA administered by IT injection for 40 weeks in children with late infantile MLD is determined. In addition, the clinical activity of rhASA on gross motor function, and single and repeated-dose pharmacokinetics in serum and concentrations in cerebrospinal fluid (CSF) are assessed.

Examples of IT Delivery of HNS

EXAMPLE 21: CHRONIC INTRATHECAL ADMINISTRATION OF HEPARAN N-SULFATASE

[0673] This example demonstrates that intrathecal administration can be used to effectively deliver a lysosomal enzyme, such as recombinant human heparan N-sulfatase (rhHNS), into brain tissues for the treatment of the neurologic symptoms of mucopolysaccharidosis IIIA (MPS IIIA; Sanfilippo syndrome type A), the defining clinical feature of this disorder. Experiments described in this example demonstrate that chronic IT administration of rhHNS was well tolerated with dose-related enzyme activity detected in the brain, spinal cord and liver.

[0674] In summary, an intrathecal (IT) formulation of recombinant human heparan N-sulfatase (HNS) has been developed for the treatment of the neurologic symptoms of mucopolysaccharidosis IIIA (MPS IIIA; Sanfilippo syndrome type A), the defining clinical feature of this disorder. Since the average age of MPS IIIA patients is 4.5 years, the pivotal toxicology studies for HNS were conducted in juvenile cynomolgus monkeys to evaluate the effects on the developing brain. Monkeys were implanted with an intrathecal (IT)-lumbar drug

2017202349 10 Apr 2017

delivery device and dosed every other week by short-term infusion (1.5, 4.5, or 8.3 mg/dose HNS for 6 months; 12 doses), with device and vehicle controls receiving phosphate-buffered saline or vehicle, respectively. Eight animals per group (4/sex) were necropsied at 3 and 6 months (device-control group necropsied at 3 months), and 8 animals from the vehicle group and the 3 HNS dose groups were necropsied 1 month after the final IT dose. No HNS-related clinical signs or gross central nervous system lesions were observed. Compared to controls, there were cellular infiltrates of slight-to-minimal mean severity in the meninges/perineurium surrounding the brain/spinal cord correlating with transient increases in cerebrospinal fluid (CSF) leukocytes, predominantly eosinophils, which largely resolved 1-month post-final dose. These changes were not associated with any adverse morphologic changes in the brain or spinal cord. There appeared to be a dose related trend toward higher mean CSF HNS levels and in tissue HNS activity levels in the brain, spinal cord, and liver. The no-observed-adverse-effect-level was 8.3 mg/dose given every other week, the highest dose administered, indicating that HNS may be safely administered intrathecally at various concentration including concentrations higher than 8.3 mg/dose.

Sanfilippo Syndrome type A

[0675] Mucopolysaccharidosis type IIIA (MPS IIIA; Sanfilippo Syndrome type A), a rare lysosomal storage disorder affecting approximately 1 in 100,000 people worldwide, results from the absence or defective function of heparan N-sulfatase (HNS) (Neufeld EF, et al. *The Metabolic and Molecular Bases of Inherited Disease* (2001) pp. 3421-3452), an exosulfatase involved in the lysosomal catabolism of glycosaminoglycan (GAG) heparan sulfate. In the absence of this enzyme, GAG heparan sulfate accumulates in lysosomes of neurons and glial cells, with lesser accumulation outside the brain. The defining clinical feature of this disorder is central nervous system (CNS) degeneration, which results in loss of, or failure to attain, major developmental milestones. The progressive cognitive decline culminates in dementia and premature mortality.

IT Delivery of rhHNS

[0676] Since the average age of MPS IIIA patients is 4.5 years, the pivotal toxicology studies for HNS were conducted in juvenile cynomolgus monkeys (species selection based upon genetic and anatomic similarity to humans) to evaluate the effects on the developing brain. The

2017202349 10 Apr 2017

age equivalence of monkeys to humans as cited in the literature ranges from 7.6 months to 12.1 months for children 30 to 40 months old (Hood RD, Developmental and Reproductive Toxicology: A practical approach (2006) p. 276). As part of this effort, a 6-month toxicology study was conducted in juvenile cynomolgus monkeys to evaluate IT lumbar administration of HNS. The data obtained from a prior 1-month juvenile cynomolgus monkey toxicity study guided the dose level selection and design of the 6-month repeated-dose juvenile monkey study. To our knowledge, this is the first study involving the chronic IT administration of ERT in juvenile nonhuman primates.

[0677] Fifty-six male and 56 female juvenile cynomolgus monkeys (*Macaca fascicularis*) approximately 6 to 9 months old and weighing 0.82 to 1.81 kg were used in this study. Monkeys were fed 15 biscuits of PMI-Certified Primate Diet 5048 (Richmond, IN) daily. Water was provided *ad libitum* via a filtered automatic water system and was withheld during urine collection periods. Monkeys were group-housed (two per cage) for 2 to 4 weeks in stainless steel cages upon arrival with the exception of the 3-month monkeys; these were individually housed in stainless steel cages. For the duration of the study, all monkeys were housed in individual stainless steel cages in rooms with controlled temperature and humidity with a cycle of 12 hours of light and 12 hours of darkness.

[0678] Prior to study initiation, all monkeys were implanted surgically with SC ports and IT catheters. Prednisolone sodium succinate (IV, 30 mg/kg) and flunixin meglumine (intramuscular [IM], 2 mg/kg) were administered prior to surgery. The monkeys were pretreated with SC atropine sulfate (0.04 mg/kg), sedated with IM ketamine HCl; 8 mg/kg), intubated, and maintained on approximately 1 L/min of oxygen and 2.0% isoflurane. An incision was made over the dorsal processes of the lumbar spine (L₄, L₅, or L₆), and a hemilaminectomy was made for the insertion of a tapered polyurethane catheter (25 cm in length, 0.9 mm outer diameter x 0.5 mm inner diameter, with six side holes of 0.33 mm diameter) at L₃, L₄, or L₅. The catheter was inserted through a small dural incision and was advanced approximately 10 cm anterograde to the area of the thoracolumbar junction. A titanium SC port was attached to the IT catheter and implanted in the SC tissue. Proper catheter placement was confirmed by myelogram using Isovue-300 (0.8 ml; Bracco Diagnostics, Inc., Princeton, NJ). After recovering from surgery,

2017202349 10 Apr 2017

monkeys received butorphanol tartrate (IM, 0.05 mg/kg) and ceftiofur sodium (IM, 5.0 mg/kg twice daily for 2 days).

[0679] In this example, HNS was provided in an IT formulation vehicle including 5 mM sodium phosphate, 145 mM sodium chloride, and 0.005% polysorbate 20 (pH 7.0). EOW doses of HNS were administered as a short-term infusion over approximately eleven minutes: 0.6 mL (4 minutes) followed with a flush of 0.5 mL phosphate-buffered saline (PBS) (7 minutes). Monkeys in the vehicle-control group received the IT formulation alone; DC monkeys received PBS (pH 7.2) IT.

Morbidity and mortality

[0680] There were no HNS-related deaths or early sacrifices. There were no HNS-related clinical signs noted at dosing or during the daily observations. Misplacement, pruritis, tremors, and ataxia observed during and after dosing resolved within a few minutes to approximately 4 hours of administration, and were considered a volume-related response rather than a reaction to HNS or the vehicle. Clinical signs observed during and immediately after dosing were seen at a comparable incidence in control groups (DC and/or vehicle-dosed group); there was no evidence of a dose response. In general, the incidence of clinical signs at dosing decreased with each subsequent dose. There were no HNS-related changes in body weight, food consumption, and physical and neurologic findings, or alterations in ECG or ophthalmology examinations.

Clinical pathology

[0681] There were no changes considered related to HNS in hematology, serum chemistry, coagulation, or urinalysis parameters at any interval.

CSF cell counts and chemistry

[0682] There were dose-related increases in mean CSF leukocyte counts for all groups, including DC and 0 mg/dose groups, 24 hours postdose. There was a general increase in leukocyte counts with each dose administered. Collection of CSF from approximately one half of the monkeys prior to dosing showed that these effects had abated in the 2 weeks since the

2017202349 10 Apr 2017

previous dose. After dose 5, in addition to an increase in leukocytes, higher group mean CSF total protein and albumin were observed for the HNS-dosed males in the 4.5 and 8.3 mg/dose groups (up to 4- to 5-fold) compared with the predose mean ($P \leq 0.05$ versus the DC and the 0 mg/dose group); less of a trend was evident in the female HNS-dosed groups.

HNS concentrations and antibody analysis

[0683] Typically, the mean HNS levels in serum were < limit of detection (LOD) for all test groups for all time points. The HNS concentration in CSF from monkeys in the DC- and vehicle-dosed control group was generally below the limit of quantification (LOQ). Although no statistical analyses were performed, there appeared to be a dose-related trend towards higher mean HNS levels in CSF in the 1.5, 4.5, and 8.3 mg/dose groups. The predose CSF mean HNS levels were significantly lower than the postdose CSF levels. The mean HNS concentrations for the 6-month cohort (both sexes) at study termination (main and recovery necropsy) are summarized in Table 46. At a given dose level, mean concentrations of HNS in the CSF appeared to be maintained in the same range (Figure 146A) despite the anti-HNS antibody levels in the serum and CSF, which continued to rise throughout the study.

Table 46: CSF HNS concentrations at study termination (main and recovery necropsies).

Group	Main Necropsy		Recovery Necropsy	
	n	Mean ± SD ^a (ng/mL)	n	Mean ± SD (ng/mL)
Vehicle	8	-	8	NA
1.5 mg IT	8	516,366 ± 1,024,084	8	NA
4.5 mg IT	7	377,460 ± 304,996	7	NA
8.3 mg IT	8	419,492 ± 345,975	8	NA

CSF, cerebrospinal fluid; HNS, human heparan N-sulfatase; n = number of samples above the LOQ; IT, intrathecal; SD, standard deviation.

^a = samples collected approx. 24 hours postdose.

NA = no samples available for analysis or samples below the LOQ.

[0684] In the 6-month/recovery cohort, none of the monkeys in the device control group (PBS only) or those dosed with vehicle developed anti-HNS antibodies in serum or CSF at any

2017202349 10 Apr 2017

time point tested. All monkeys in the 1.5, 4.5, and 8.3 mg/dose groups tested negative (<LOD) for anti-HNS antibodies in serum and CSF samples collected prestudy (for CSF) and at predose 2. By the end of the study, all monkeys tested positive for anti-HNS antibodies in serum.

[0685] All monkeys in the 1.5 mg/dose and 8.3 mg/dose groups and six of eight monkeys in the 4.5 mg/dose group tested positive for anti-HNS antibodies in the CSF at one or more time points. Since two monkeys in the 4.5 mg group had no sample collected at any time point including necropsy, these results would appear to indicate that all monkeys dosed with HNS produced an antibody response.

[0686] At all three dose levels, anti-HNS antibody concentrations in serum were detected after dose 2, and levels increased markedly after dose 4. Although no statistical analyses were performed, there appeared to be a dose-related trend towards higher serum antibody concentration; by the end of the study, levels were comparable across the 3 HNS dose groups (Figure 146B). Anti-HNS antibody levels in the serum were always higher than in the CSF over the time course of this study (from 9 to 236-fold serum/CSF antibody concentrations); the highest ratios of serum to CSF concentrations (98 and 236-fold) were seen at 8.3 mg dose level in the earlier course of dosing (6 and 10 weeks).

[0687] Anti-HNS antibody concentrations in the serum increased 9-, 16-, and 16-fold at 1.5 mg, 4.5 mg, and 8.3 mg/dose levels, respectively, in the early time of dosing (from week 6 to week 14). During the same time period, CSF antibody concentrations increased 30-, 41-, and 52-fold at 1.5 mg, 4.5 mg, and 8.3 mg/dose levels, respectively (Figure 146B); substantial levels remained after the 1-month dose-free recovery phase (

Table 47).

2017202349 10 Apr 2017

Table 47: CSF anti-HNS antibody concentrations at study termination (main and recovery necropsies).

Group	Main Necropsy ^a		Recovery Necropsy	
	n	Mean ± SD (ng/mL)	n	Mean ± SD (ng/mL)
Vehicle	8	-	8	-
1.5 mg IT	8	351,456 ± 244,171	8	299,512 ± 226,654
4.5 mg IT	7	147,187 ± 213,095	7	193,045 ± 157,896
8.3 mg IT	8	185,227 ± 315,858	8	238,727 ± 185,785

CSF, cerebrospinal fluid; HNS, human heparan N-sulfatase; IT, intrathecal; n, number of sample above the limit of quantification; SD, standard deviation.

^aSamples collected approximately 1 week prior to dosing.

[0688] Anti-HNS antibodies appeared later in the CSF than in serum (Figure 146C). No apparent dose-related differences of antibody concentrations in the serum or CSF were observed (statistical analysis was not done due to small sample sizes); there was no observable difference between males and females in antibody responses.

[0689] In the presence of anti-HNS antibody in the CSF, the mean concentrations of HNS in the CSF appeared to be maintained, suggesting that the presence of anti-HNS antibodies in the serum and CSF did not alter the concentration level of the IT-dosed HNS. The 6-month/recovery cohort analyses of the 6-month repeat-dose administration of HNS indicated that the anti-HNS antibody concentrations for the 3-month interim and 6-month cohort sacrifice monkeys were comparable (Figure 146C).

Gross and histopathologic findings

[0690] At all dose levels (although not at all sacrifice intervals, gender-specific, nor in a dose-related manner), eosinophilic infiltrates (Figure 147A) were present in the parenchyma of the brain (predominantly gray matter), spinal cord (gray and white matter), dorsal spinal nerve roots/ganglia and the trigeminal ganglia (mid-dose males only) (Figure 147B-D). The infiltrates

2017202349 10 Apr 2017

were interpreted to be secondary to the meningeal/perineurium infiltrates and/or to the presence of (penetration by) HNS within the parenchyma of the tissue. Although there were numerous inflammatory type changes, the monkeys appeared to tolerate administration of HNS and none of the infiltrates were considered related to or causing adverse morphologic changes in the nervous system parenchyma. Specifically, there was no evidence of neuronal necrosis/degeneration and no glial response related to HNS administration.

[0691] Microgliosis in the gray matter of the brain and spinal cord, in association with cellular infiltrates, predominantly eosinophilic, was relatively common in a previously performed 1-month juvenile monkey toxicity study; these changes were relatively uncommon by the 3-month interim sacrifice in the 6-month study, but residual evidence of such a response could still be seen in the 6-month cohort (Figure 147F). Microglial reactions tend to be a relatively early event in the reaction to some (typically protein-based) centrally administered (or centrally-reactive) test articles. The eosinophilic infiltrates did correlate with increased number of eosinophils in the CSF of HNS-dosed monkeys, although the cells were not present in sufficient numbers to elicit an adverse reaction.

[0692] At all dose levels, eosinophilic infiltrates were observed in the dorsal spinal nerve roots/ganglia for most HNS-dosed groups, regardless of gender. The infiltrates in the various nervous system tissues were interpreted to be secondary to the meningeal/perineurium infiltrates and/or to the presence of (penetration by) HNS within the parenchyma of the tissue. In the recovery sacrifice monkeys, HNS-related effects were generally either absent or reduced to control levels. Some changes, such as microgliosis in the spinal cord, were completely resolved after the recovery period. None of the HNS-related changes appeared to be associated with any adverse structural microscopic changes in the brain or spinal cord. There was no neuronal necrosis noted in the brain, spinal cord, or ganglia.

HNS enzyme activity

[0693] In the 6-month/recovery cohorts, HNS enzyme activity in the spinal cord and brain of the vehicle-dosed group (0.0-0.154 nmol/hr/mg protein) were similar to levels shown in tissues from the 3-month interim cohort (0.0-0.0.154 nmol/hr/mg protein). Enzyme activity levels in the spine were higher (approximately an order of magnitude higher in the lumbar spine)

2017202349 10 Apr 2017

than levels measured in brain or liver, the 4.5 mg and 8.3 mg/dose groups having similar levels. The HNS enzyme activity in spinal cord slices ranged from 3.9-18.6, 13.1-67.1, and 3.6-69.2 nmol/hr/mg protein in males (Figure 148A) and 1.8-16.2, 4.5-61.2, and 21.1-66.0 nmol/hr/mg protein in females (Figure 148B) for the 1.5, 4.5, and 8.3 mg/dose groups, respectively. In spinal tissue after a 1-month recovery period, enzyme activity levels returned to levels consistent with vehicle control values.

[0694] The HNS enzyme activity in brain slices ranged from 0.03-16.0, 0.30-55.7, and 0.15-21.2 nmol/hr/mg protein in males (Figure 148C), and 0.04-5.1, 0.0-14.4 and 0.9-33.2 nmol/hr/mg protein in females (Figure 148D) for the 1.5, 4.5, and 8.3 mg/dose groups, respectively. In brain tissue after recovery, enzyme activity levels returned to levels consistent with control values.

[0695] The fold-change in activity for different areas of the brain compared with endogenous levels (DC group) is shown in Figure 149A. Although a trend toward increased distribution was noted in surface samples, lumbar-IT administered HNS could be shown to penetrate to periventricular areas of the brain.

[0696] In the 6-month cohort/recovery cohorts, mean activity levels in liver were 0.50, 2.41, and 6.65 nmol/hr/mg protein in males and 1.04, 4.15, and 7.62 nmol/hr/mg protein in females for the 1.5, 4.5, and 8.3 mg/dose groups, respectively (Figure 149B). Levels in vehicle control monkeys were 0.089 nmol/hr/mg protein for males and 0.083 nmol/hr/mg protein for females. Following the recovery period, HNS activity levels in liver were comparable to baseline control levels for all dose groups.

Immunohistochemistry

[0697] HNS delivery to the CNS via bolus IT injection in the 3-month interim and 6-month/recovery cohorts resulted in delivery of immunoreactive test article to the pia-arachnoid tissues of the spinal cord and brain. In the monkeys that received IT HNS, the immunoreactive material was consistently present in meningeal and perivascular macrophages (brain/spinal cord) and variably present in the adjacent glial and neuronal cell populations. The lack of staining in vehicle-dosed control monkeys (Figure 150A) demonstrated the specificity of the antibody to

2017202349 10 Apr 2017

human HNS. Generally, the immunoreactivity was dose related (i.e., using a semi-quantitative grading scale, increased immunohistochemical staining was noted in a generally dose-dependent manner). HNS delivery to the CNS via bolus IT resulted in positive immunostaining in the cerebral cortex and cerebellum (Figure 150B-D); however, immunoreactivity was not consistently evident in the caudate/putamen region, midbrain, or deeper regions of the pons or medulla. Immunoreactivity was evident in the livers (in sinusoidal lining cells including Kupffer cells, but not in hepatocytes) of all monkeys administered HNS. Immunoreactivity was not evident in the one female sacrificed early (4.5 mg/dose group) because of a leaking catheter that could not be repaired.

[0698] In the 1.5 mg/dose group, essentially full recovery was evident with the exception of liver and the meninges of the brain and spinal cord where some residual immunoreactivity was evident. At higher doses (4.5 and 8.3 mg/dose), the intensity and incidences of immunoreactivity were lower than at the end of dosing. At all dose levels, the levels of HNS in spinal cord, brain, and liver approximated those seen in vehicle-dosed controls after the 1-month recovery period.

[0699] In this study, EOW delivery of HNS administered IT for 6 months was generally well tolerated. No remarkable changes were observed in body weight, clinical status, ophthalmologic/neurologic/physical examinations, ECGs, organ weights, or gross organ appearance. Findings were limited to transient changes in CSF clinical pathology accompanied by slight to mild meningeal infiltrates and epidural inflammation, with nearly complete reversal in all but the highest dose group following the recovery period. Widespread distribution of HNS throughout the brain and spinal cord was observed.

[0700] IT administration of HNS EOW elicited an inflammatory response characterized by residual leukocyte infiltration and effusion of albumin noted at 24 hours postdose and at necropsy. Without wishing to be bound by any particular theory, this presumably reflects a transient, localized, and incomplete opening of the BBB related to changes in the tight junctions near the catheter tip, resulting in entry of leukocytes and plasma proteins into the CSF (Simard JM, et al. *Lancet Neurol.* (2007) 6, 258–268; Stamatovic SM, et al. *Curr. Neuropharmacol.*

2017202349 10 Apr 2017

(2008) 6, 179–192). This may be the result of two components: one related to the dose administration procedures or volume and another related to IT administration of a protein.

[0701] The transient changes in BBB permeability (no significant differences between dose groups and controls 24 hours postdose at the main necropsy), were not accompanied by any clinical signs.

[0702] There appeared to be a dose-related trend for higher mean CSF HNS levels; at a given dose level, mean concentrations of HNS in the CSF appeared to be maintained in the same range despite the increasing anti-HNS antibody levels in the serum and CSF.

[0703] Meningeal cellular infiltration of slight-to-minimal mean severity was observed in the brains and spinal cords of HNS-dosed juvenile monkeys. This microscopic change was also noted in vehicle-dosed controls, indicating some of the response was related to IT catheter placement, as well as a nonspecific inflammatory response to foreign protein. The introduction of a biologic/protein into the IT space, especially one that penetrates the CNS, nearly always elicits some degree of an inflammatory response (Hovland DN, et al. *Toxicol. Pathol.* (2007) 35, 1013–1029; Butt MT, *Toxicol. Pathol.* (2011) 39, 213–219), which, if present in numbers that damage adjacent tissue, would represent an adverse effect. In the current study, however, these cells (predominantly eosinophils) appeared to represent a marker of tissue reaction/penetration and were not found in sufficient quantities to qualify as an adverse effect. None of the HNS-related changes appeared to be associated with any adverse structural microscopic changes in the brain or spinal cord. There was no neuronal necrosis noted in the brain, spinal cord, or ganglia.

[0704] Evaluation of anti-test article antibodies is an important aspect of the toxicity studies because of the potential impact of neutralizing or binding antibodies on the clearance or biodistribution of test article (Ponce RP, et al. *Regul. Toxicol. Pharmacol.* (2009) 54, 164–182). In this study, since dose-related and quantitatively similar levels of HNS enzyme activity were noted in the brain and spinal cord of the 3-month interim and 6-month cohorts, and mean concentrations of HNS in the CSF appeared to be maintained in the same range despite the increasing anti-HNS antibody levels in the serum and CSF, suggesting no neutralizing activity.

2017202349 10 Apr 2017

[0705] There appeared to be a dose-related trend toward higher levels of HNS enzyme activity in spinal cord, brain, and liver, that was highest near the injection site in the lumbar region of the spinal cord and uniform in the brain, with no significant differences rostral to caudal and between right and left hemispheres. No evidence for HNS accumulation was noted in the brain and spinal cord tissue of the 6-month cohort as compared with the 3-month interim cohort. Although a trend toward increased distribution was noted in surface samples, lumbar-IT administered HNS penetrated to deep, periventricular areas of the brain. The HNS enzyme activity in the liver suggested the HNS redistributed systemically after IT delivery; no HNS-related adverse effects were observed in the liver after evaluation of clinical and anatomic pathology parameters in the pivotal toxicity studies.

[0706] In general, the immunohistochemistry results corroborated the tissue enzyme activity in that dose-related immunoreactivity was observed in the spinal cord and brain pia-arachnoid meninges and in the nervous tissues (neurons, glial cells) in the immediate proximity of the meninges. There was good gray matter penetration of the cerebrum and cerebellum after bolus IT injection or short-term IT infusion. Although immunoreactivity was not evident in deeper structures such as the basal ganglia or the central regions of the thalamus/hypothalamus, midbrain or the pons/medulla, enzyme activity results indicate that lumbar-IT administered HNS penetrated to deep, periventricular areas of the brain. Thus, immunohistochemistry may be less sensitive technique for detecting biodistribution of a test article. Immunoreactivity was evident in Kupffer cells and the endothelial cells (cells capable of phagocytosis) of the liver, but not parenchymal cells (hepatocytes).

[0707] The 6-month/recovery cohort analyses of the 6-month repeated-dose IT toxicity study in juvenile monkeys indicated that HNS-related changes in the 3-month interim and 6-month sacrifice monkeys were comparable, including in-life parameters, clinical and anatomic pathology, concentrations of HNS and anti-HNS antibodies in CSF and serum, and distribution/subcellular location of HNS in spinal cord, brain, and liver. In the recovery sacrifice monkeys, HNS effects were either absent or significantly reduced. Thus, the no-observed-adverse-effect-level for the 6-month juvenile monkey study was 8.3 mg/dose, the highest dose administered.

2017202349 10 Apr 2017

[0708] Monitoring changes in CSF cellularity and protein concentrations appears to be a reliable correlate of the morphological changes noted on histopathologic evaluation and may be useful in patients treated IT with HNS; these changes were considered to be an expected reaction to an IT-administered protein and were largely resolved after the recovery period. These data from animal models provide confidence for pursuing IT therapy as a treatment strategy for the neurological manifestations of lysosomal storage diseases. This juvenile nonhuman primate toxicology study demonstrates the feasibility and tolerability of administering HNS via an IT lumbar drug delivery device to pediatric patients. The nonadverse CNS pathology and lack of adverse clinical signs have supported the recent investigational medical product dossier approval and indicated that IT-administered HNS can safely and effectively treat CNS symptoms of Sanfillippo A syndrome.

MATERIALS AND METHODS

[0709] Exemplary materials and methods used in various experiments described in this examples are provided below.

Study design and HNS dosing

[0710] The monkeys were randomized into five treatment groups; group 1 was untreated (implant device control [DC], port and catheter) and was not dosed with the vehicle or test article. Groups 2 through 5 received 0.6 mL of 0, 2.5, 7.5 or 13.8 mg/mL HNS IT, (i.e., a total dose of 0, 1.5, 4.5, or 8.3 mg) EOW. Four monkeys/sex/group were necropsied at 3 months (interim necropsy; 24 hours after the 6th dose), four monkeys/sex/group (except the DC group, which were necropsied at 3 months) were necropsied at 6 months of dosing (main necropsy; 24 hours after the 12th dose), and the remaining four monkeys/sex/group were necropsied at the end of a 1-month recovery period. At necropsy, selected tissues were harvested, processed, and examined microscopically.

[0711] HNS was provided in an IT formulation vehicle consisting of 5 mM sodium phosphate, 145 mM sodium chloride, and 0.005% polysorbate 20 (pH 7.0). Every other week doses of HNS were administered as a short-term infusion over approximately eleven minutes: 0.6

2017202349 10 Apr 2017

mL (4 minutes) followed with a flush of 0.5 mL phosphate-buffered saline (PBS) (7 minutes). Monkeys in the vehicle-control group received the IT formulation alone; DC monkeys received PBS (pH 7.2) IT.

Clinical evaluation

[0712] Clinical signs and morbidity and mortality observations were recorded at least twice daily starting at the first dose. Body weights were measured prior to surgery, on the day of surgery, weekly during the study, and at necropsy. Food consumption was monitored daily starting before surgery. Physical (heart rate, respiration, body temperature, auscultation, gait, disposition, abdominal palpation, lymph nodes, and general appearance) and neurologic (level of consciousness, tracking) examinations were performed before the study was initiated, each month during the study, and before necropsy. Motor functions, cerebral reflexes (pupillary, blink, and corneal reflex), and spinal reflexes (sensory foot, knee jerk, cutaneous, proprioceptive, and tail reflex) were also assessed. Electrocardiographic (ECG; leads I, II, and III) and ophthalmologic examinations were completed prior to the first dose of HNS and in the week before the interim (3-month) or the main (6-month) necropsy. Ophthalmic examinations were performed by indirect ophthalmoscope, the monkeys were sedated with ketamine HCl (IM, 8 mg/kg), and eyes were dilated with 1% tropicamide.

Clinical pathology

[0713] Blood samples were collected from fasted monkeys for hematology and serum chemistry prior to the study start, after IT doses 1, 3, 5, 7, 9 and 11, mid-recovery, and at necropsy. Urine samples were collected via pan catch predose, once monthly during the dosing and recovery period, and prior to necropsy. CSF samples were collected via the lumbar catheter for total cell count and chemistry analysis at the time of surgery, and 24 hours following IT doses 1, 3, 5, 7, 9, 11, mid-recovery, and at necropsy; on occasion, samples were not collected due to partial catheter obstruction. Because higher than expected CSF leukocyte counts were noted, the 3-month dose 5 CSF samples were collected from half the monkeys in each group before dosing and from the remaining monkeys 24 hours after dosing. The predose sample collection occurred at least 1 day prior to dosing so as not to significantly alter the CSF volume just prior to dosing. For the 6-month and recovery monkeys, CSF for total cell count and chemistry was collected

2017202349 10 Apr 2017

from half the monkeys in each group before dosing and from the remaining monkeys 24 hours after dosing. If a monkey had a nonsampling catheter due to an obstruction, a spinal tap (cisterna magna) was performed at the necropsy.

HNS analysis

[0714] Blood samples for HNS analysis were collected from a peripheral vein prior to and 24 hours post IT doses 2, 4, 6, 8, 10, 12; mid-recovery, and at necropsy. CSF samples were collected via the lumbar catheter prior to and 24 hours post IT doses 2, 4, 6, 8, 10, 12, mid-recovery, and at necropsy. HNS concentrations were determined by enzyme-linked immunosorbent assay. The capture antibody was a polyclonal rabbit anti-HNS IgG and the detection antibody was a horseradish peroxidase-conjugate of the same rabbit anti-HNS IgG. The LOD was 0.22 ng/mL; thus, the LOQ was calculated to be 0.66 ng/mL. Serum and CSF samples were screened in duplicate at 1:100 and 1:5 dilutions; samples exceeding the high end of the calibration curve were further diluted and retested.

Anti-HNS antibody analysis

[0715] Blood for antibody analysis was collected from a peripheral vein approximately 1 week prior to IT doses 2, 4, 6, 8, 10, 12; mid-recovery, and at necropsy. CSF samples for antibody analysis were collected at surgery, and via the lumbar catheter approximately 1 week prior to IT doses 2, 4, 6, 8, 10, 12; mid-recovery; and at necropsy. A Meso Scale Discovery (MSD[®]) technology electrochemiluminescent bridge test was used for detection of anti-HNS antibodies. The assay is a general, but sensitive, screening method for anti-HNS antibodies from any species and all immunoglobulin isotypes. The LOD was 5 ng/mL, and the samples were screened in duplicate at a 1:20 dilution, resulting in an effective assay sensitivity of 100 ng/mL. Samples exceeding the high end of the calibration curve were further diluted and retested.

Necropsy and preparation of tissues

[0716] Monkeys underwent a full necropsy either 24 hours after the final IT dose (main necropsy) or at the end of the 1-month recovery period (recovery necropsy). All monkeys were sedated with ketamine HCl (IM, 8 mg/kg), were maintained on an isoflurane/oxygen mixture,

2017202349 10 Apr 2017

and received an IV bolus of heparin sodium (200 IU/kg). Monkeys were perfused via the left cardiac ventricle with room temperature 0.001% sodium nitrite in saline at a rate of 200 ml/min for 12 min (~2400 ml). After collection, tissue samples were then fixed in 10% neutral buffered formalin for histopathologic examination/immunohistochemical analysis or were frozen on dry ice and stored at -60°C or lower for analysis of HNS activity.

[0717] The brain was cut in a brain matrix (MBM-2000C, ASI Instruments, Inc., Warren, MI) at 3-mm coronal slice thickness. The slices were numbered, with the most rostral slice designated as slice 1. Slices 1, 4, 7, 10, 13, and 16 were processed for histopathology and slices 2, 5, 8, 11, 14, and 17 (if available) were processed for immunohistochemistry. Slices 3, 6, 9, 12, and 15 were frozen for analysis of HNS activity. The spinal cords (cervical, thoracic, and lumbar portions) were cut into 1-cm sections. The first slice and every third slice thereafter were processed for histopathologic evaluation and the second slice and every third slice thereafter were processed for immunohistochemical analysis. The third slice and every third slice thereafter were frozen for HNS analysis. The distribution of slices was adjusted so that the slice containing the tip of the intrathecal catheter (slice 0) was fixed in formalin and analyzed for histopathology. Duplicate samples of ~5 g of the liver were taken from two separate lobes and frozen for HNS analysis and an additional sample of ~5 g was fixed for immunohistochemical analysis.

Histopathology

[0718] The brains, spinal cords, dorsal spinal nerve roots/ganglion, sciatic, tibial and sural nerves, a complete tissue list (typical for preclinical drug safety studies of this duration in this species), and any gross lesions were harvested at necropsy from all monkeys. Tissue sections were embedded in paraffin and stained with hematoxylin and eosin (in addition to any special staining/embedding procedures noted below) for comprehensive microscopic evaluation.

[0719] Brain sections from the prepared paraffin blocks from the device and vehicle-control groups, and the high-dose monkeys were stained with Fluoro-Jade B (a stain increasing the sensitivity of evaluating neuronal degeneration) and Bielschowsky's silver (a procedure that allows direct visualization of axons, dendrites, and neuronal filaments). The

2017202349 10 Apr 2017

Fluoro-Jade B stained slides were examined under fluorescent lighting using a fluorescein isothiocyanate filter cube.

[0720] Spinal cords were sectioned serially, with a transverse and oblique sections taken at the cervical, thoracic, and lumbar regions (one slice examined at each level) including sections at the catheter tip; an additional transverse section was taken from the cauda equina region. Dorsal spinal roots and ganglia (midcervical, midthoracic, and midlumbar) were processed and examined. Peripheral nerves (sciatic, tibial, and sural) were sectioned longitudinally, embedded in paraffin and stained with hematoxylin and eosin (H&E). Cross sections were postfixed in osmium, embedded in Spurr's resin, sectioned (2 μ m) and stained with toluidine blue. Serial spinal cord sections, as well as dorsal spinal nerve roots and ganglia, from the device and vehicle control groups and the high-dose group were stained with Bielschowsky's silver. Spinal cord sections from these groups also were stained with anti-glial fibrillary acidic protein, an immunohistochemical stain that allows for direct visualization of astrocytes and their processes.

Preparation of tissue extracts for quantitative analysis

[0721] Frozen brain slices 3, 6, 9, 12, and 15 were dissected by separating the left and right hemispheres. Surface tissue was taken by measuring 4 mm from the surface, and the remaining tissue in each hemisphere was considered deep tissue. If present (e.g., slices 6 and 9), an additional periventricular sample was cut from the coronal slices. Since only one-half of the brain (the right side) was processed (the left side was retained frozen), the sectioning resulted in two to three samples per slice: right surface, right deep, and, if present, right periventricular (i.e., Ventricle deep; Vdeep). Cerebellar and brain stem tissues, when present, were isolated prior to separating the hemispheres and were processed independently. Spinal cord sections were prepared similarly, weighed, and homogenized.

[0722] Tissue samples were homogenized in lysis buffer (1 ml/0.25 g tissue) formulated with 10 mM Tris, 5 mM ethylenediaminetetracetic acid, 0.1% Igepal supplemented with Alpha Complete protease inhibitor minitables (Roche Diagnostics, Indianapolis, IN) using TeenA Lysing Matrix A tubes or conical polypropylene tubes. Samples were processed for 40 seconds in the Fastprep-24 automated homogenizer (MP Biomedicals, Solon, OH) or PowerGen Model 125 powered homogenizer (Omni International, Kennesaw, GA). Once homogenized, samples

2017202349 10 Apr 2017

were subjected to five freeze-thaw cycles using an ethanol/dry ice bath and a 37°C water bath and then centrifuged at 4°C to pellet tissue debris; supernatants were stored at -80°C until assayed. HNS activity was determined using a specific substrate (4-methylumbelliferyl- α -D-N-sulphoglucosaminide) with a 2-step fluorometric assay.

Tissue processing and staining for immunohistochemistry

[0723] Six formalin-fixed coronal brain slices (slice numbers 2, 5, 8, 11, 14, and 17) of 3-mm thickness from each monkey were numbered 1 to 6 rostral to caudal. Generally, slices 1 to 4 contained basal nuclei/thalamus/midbrain and cerebrum, and the caudal two slices contained cerebellum and brain stem (medulla oblongata) tissue. Brain, spinal cord and liver sections (from the same paraffin blocks as those used for H&E and the various special stains) were immunohistochemically stained for HNS. A specific mouse monoclonal antibody (clone 2C7; Maine Biotech, Portland, ME) was used to detect intracellular uptake of IT-administered HNS; this reagent demonstrated no cross-reactivity with endogenous cynomolgus monkey HNS. Negative controls were performed using an irrelevant mouse IgG. Deparaffinized slides were incubated with primary mouse anti-HNS antibody overnight at 2 to 8°C. A secondary goat anti-mouse biotinylated immunoglobulin G was added and incubated for 30 minutes at 37°C. Avidin/biotinylated horseradish peroxidase complex was added and incubated for 30 minutes. Slides were incubated in peroxidase substrate diaminobenzidine solution until the desired stain intensity developed. Nuclei were counterstained with hematoxylin.

Statistical analyses

[0724] Body weights, body weight changes, food consumption, respiratory rate, body temperature, heart rate, CSF cell count, CSF chemistry, clinical pathology data, urine data, and absolute and relative organ weights were analyzed by a one-way analysis of variance and a comparison of the device and vehicle control groups to each HNS-dosed group by Dunnett's test. In addition, the statistical analysis compared the two control groups to each other. Analysis was two-tailed for significance levels of 5% and 1%. All data are presented as mean \pm standard deviation.

2017202349 10 Apr 2017

EXAMPLE 22: HEPARAN N-SULFATASE BIODISTRIBUTION AND PHARMACOKINETIC STUDIES

[0725] The experiments in this example were designed to determine tissue distribution of HNS in rats after a single intravenous or intrathecal dose (1 or 10 mg/kg) of HNS. For example, among other things, the purpose of these experiments were to characterize the biodistribution (BD) properties of HNS in rats using positron emission tomography (PET); to compare distribution patterns of HNS when given in different routes (IV or IT) and at different doses (1 or 10 mg/kg); and to determine pharmacokinetic properties of HNS in each of the interest organs in these dosing regimens.

[0726] Pharmacokinetic (PK) and biodistribution (BD) profiles of ¹²⁴I-sulfamidase (HNS) were studied by tissue PET imaging in rats after single intravenous (IV) or intrathecal (IT) administration of 1 or 10 mg/kg of ¹²⁴I-HNS. Radioactivity-time data in the region of interest were obtained from dynamic images in the first 20 min and from static images at 0.05 (only for IT administration), 1, 2, 4, 8, 24, 48, 96 and 192 hours post IV or IT dosing.

[0727] Four rats in each of four groups (1 mg/kg IV, 1 mg/kg IT, 10 mg/kg IV and 10 mg/kg IT) were used in this study. Radioactivity-time data were measured in the head, brain (including cerebrospinal fluid, CSF), spine and liver regions after IT administration; and in the blood, brain (including CSF), liver, kidney, heart (including lungs) and skin after IV administration. The data were corrected by the decay half-life of ¹²⁴iodine (100.2 hours), expressed as percentage of injected dose (%ID) of a region of interest or %ID per gram (%ID/g) of the imaged tissues, and then normalized for the body weight of 200 grams. The total amounts (ug) or concentrations (ug/g) of the dosed protein in the region of interest were calculated from the corresponding %ID or %ID/g data.

[0728] In the first 20 min after IT dosing, total amount of HNS in the head region was reduced at a constant rate of 0.002/min – 0.011/min (λ_z) at 1 and 10 mg/kg. Clearance rates and distribution volumes were not used for pharmacokinetic comparisons between the two doses and the two administration routes in this report (see Results section for more information). The constant rates of elimination from the brain were essentially the same at two test doses (λ_z : 0.016/hr versus 0.014/hr for 1 and 10 mg/kg, respectively) with a similar half-life of about two

2017202349 10 Apr 2017

days as determined by static imaging up to 192 hours after IT dosing. The values of C_{max} and AUC (0-last or 0-infinite) were proportional to the administered doses. A linear PK behavior was indicated in the dose range of 1 to 10 mg/kg given in these IT single-dosing regimens.

Concentration gradients were observed from the proximal to distal sections of the spine at both dose levels.

[0729] After IT dosing, HNS protein was measurable in the liver up to 96 hours at 1 mg/kg and up to 192 hours at 10 mg/kg of HNS. The concentrations in the liver reached the peak 2 hours at 1 mg/kg, and 7 hours at 10 mg/kg. The elimination was $0.030 \pm 0.011/\text{hr}$ (mean λ_z) at 1 mg/kg, which was not significantly different from that at 10 mg/kg (λ_z $0.017 \pm 0/\text{hr}$) ($p=0.10$), with a corresponding $t_{1/2}$ (28 versus 42 hours at the doses of 1 and 10 mg/kg, respectively).

[0730] After IV dosing, the elimination half-lives in the liver, kidney, heart and skin were 47 ± 10 and 38 ± 13 hours for the liver, 54 ± 25 and 29 ± 16 hours for the kidney, 36 ± 15 and 42 ± 19 hours for the heart, and 40 ± 21 and 31 ± 13 hours for the skin at 1 and 10 mg/kg, respectively; while the half-lives in the brain were 71 ± 23 and 60 ± 53 hours. The mean values of C_{max} for the liver, skin, kidney, heart and brain were 9.6, 0.30, 0.25, 0.22, and 0.08 $\mu\text{g/g}$ at 1 mg/kg and 132, 7.9, 3.9, 3.7 and 1.8 $\mu\text{g/g}$ at 10 mg/kg. After the C_{max} values from individual animal were normalized for dose, the C_{max}/dose values at 10 mg/kg were significantly higher than that at 1 mg/kg in all these organs (most p values <0.05 , $p=0.06$ for the liver). The values of AUC_{last} for the liver, skin, kidney, heart and brain were 525, 16, 14, 9 and 7 $\text{hr}\cdot\mu\text{g/g}$ at 1 mg/kg; and 6747, 276, 183, 201 and 86 $\text{hr}\cdot\mu\text{g/g}$ at 10 mg/kg. After normalization, the AUC_{last}/dose values at 10 mg/kg were significantly higher than that at 1 mg/kg in the skin ($p<0.01$), marginally different in the heart ($p=0.06$), and not significantly different in the liver, brain and kidney (all p values >0.34).

[0731] When the same dose of HNS was injected, intrathecal administration resulted in a three-log greater brain exposure than that with intravenous administration. The elimination half-life in the brain was 2 days by IT and 3 days by IV administration. However, hepatic exposures after IT dosing were similar to that after IV dosing at the same dose of HNS. The exposure (C_{max} and AUC_{last}) for the liver by IT/IV at 1 mg/kg and 10 mg/kg were in a range of 0.4 - 1.2.

2017202349 10 Apr 2017

Experimental Design

[0732] The central nervous system (CNS) is vulnerable in most lysosome storage diseases and is seriously damaged in some types of these diseases, such as Sanfilippo syndrome type A or B (mucopolysaccharidosis III), Metachromatic Leukodystrophy (MLD) and Hunter Syndrome. As described herein, it is contemplated that, due to poor penetration through blood-brain barrier when administered peripherally, direct administration of enzymatic proteins into the CNS may increase their concentrations in the central nervous tissues and further enhance their therapeutic effects. Intrathecal (IT, or cisterna magna) administration was investigated and compared with IV administration at different dose levels in this study.

[0733] PET is a non-invasive, repeatable and quantitative technology to provide dynamic change of drug concentration over time in the organ of interest. The dynamic concentration-time data from target organs (active sites, rather than in blood circulation) are valuable, and are directly related to the biological activity of the dosed drug. Furthermore, the information on tissue exposures from PET study in animals can be used to guide the selection of the first-dose in human.

*Materials and Methods***Test Articles**

[0734] Heparin N-Sulfatase (HNS) was formulated at a concentration of 20 mg/mL of HNS in 5mM sodium phosphates buffer with 145 mM sodium chloride at pH 7.0. The material was purified by RP-HPLC and contained 98.7% of Heparin N-Sulfatase with 99.9% of dimer. HNS was labeled with ¹²⁴Iodine.

Sample Source

[0735] Radioactivity images were from rats after IV and IT dosing ¹²⁴I-H-N-sulfatase at 1 and 10 mg/kg.

Animals

2017202349 10 Apr 2017

[0736] Sixteen male Sprague-Dawley rats were purchased from Charles River Laboratories (190 ± 60 g, $n = 16$), and were separated into four groups ($n = 4$). Single IV or IT injection at two different doses (1 mg/kg and 10 mg/kg) was given to each group of these rats (total 4 groups). The dose and injected volume were individualized based on the body weight of each animal. In two IV-treated groups, sedation was induced by IV injection of sodium pentobarbital at a dose of 35 mg/kg. Intravenous doses were injected in a bolus through a tail vein. In two IT-treated groups, animals were anesthetized by intra-peritoneal administration of sodium pentobarbital at a dose of 50 mg/kg. Intrathecal doses were administered over 1 min at cisterna magna level through the atlanto-occipital membrane. The actual administered radioactivity was measured by PET, and served as the injected dose.

Experimental and/or Assay Method(s)

[0737] Dynamic images (every 2 min) were obtained in the first 20 minutes in the regions of the heart (including the lungs), liver and kidneys post IV injection; and in the head region post IT administration of both doses. Static imaging was acquired in the regions including the brain (including cerebrospinal fluid, CSF), liver, kidney, heart (including the lungs), muscle, skin and bone in IV-treated group; and in the region of head, brain (including CSF) and liver of IT-treated animals at 0.05 (only available for IT groups), 1, 2, 4, 8, 24, 48, 96 and 192 hours post-dosing. The images were reconstructed and the three body sections were fused into one image.

Data Analyses

[0738] PET data were expressed in nanocurie (nCi) per mL (for fluid) or per gram (for tissue). Relative activity was obtained for the brain, liver, kidneys, skeletal muscle, stomach, heart (with lungs) and skin regions in static images. Absolute activity in the whole head or brain regions was obtained for the animals that received IT injections. Radioactivity per millimeter of spinal column was determined in the IT injected animals at three selected sections: the proximal (neck), mid (against upper edge of the liver), and distal (1 cm from the distal end of the protein containing compartment) spine.

2017202349 10 Apr 2017

[0739] All data were corrected by the decay half-life of ^{124}I (100.2 hours) and normalized for registration efficacy based on calibration with a ^{124}I source with externally measured activity. The data were then expressed as percentage of injected dose (%ID) of a whole region (the head and brain) or %ID per gram (%ID/g) of a tissue, and then normalized for a body weight of 200 grams [data normalization: (%ID or %ID/g) / body weight of the animal x 200]. The normalization was adopted to reduce the variability of the data, as only four animals were used in each group.

[0740] In this example, HNS protein concentrations or amount were calculated using the injected protein dose to each animal: protein concentration (ug/g) = (%ID/g) x (mg/kg of injected dose x 1000 x 0.2); total amount of the dosed protein (ug) in a region of interest = %ID x (mg/kg of injected dose x 1000 x 0.2), here the injected dose was 1 mg/kg or 10 mg/kg and 0.2 is the normalizing factor for body weight. Group mean and standard deviation of each PK parameter were calculated based on the individual non-compartmental data in each of the four groups. A Student *t-test* was performed to compare the values of λ_z , $t_{1/2}$, C_{\max} and AUC between the two test doses and the two administration routes. Statistical significance was defined as a *p*-values less than 0.05 ($p < 0.05$).

Results

[0741] The amounts (ug) or concentrations (ug/g) of HNS in the following tables, figures and PK analyses were calculated by multiplying the injected protein dose (1 mg/kg or 10 mg/kg) with the corresponding values of %ID or %ID/g.

Intrathecal treatment with ^{124}I -HNS at doses of 1 and 10 mg/kg

[0742] The amount of the dosed protein (ug) in the head region from dynamic images was plotted as a function of time in Figure 151. The concentration (ug/g) in the brain regions from static images was plotted as a function of time in Figure 152. The total amount of injected protein (ug) in the brain and head regions from static images were plotted with time in Figure 153 and Figure 154, respectively. Concentration-time curves (ug/mm) at the proximal, mid and distal spine are shown in Figure 155 to Figure 157. Figure 158 shows the changes of HNS concentration (ug/g) in the liver with time after IT administration of ^{124}I -HNS at 1 and 10 mg/kg.

2017202349 10 Apr 2017

[0743] The total amount-time (ug) or concentration-time (ug/g) data were analyzed by non-compartmental models (WinNonlin 5.2, Pharsight, Mountain View, CA). The PK parameters, such as the constant rate of elimination (λ_z), peak concentration (C_{max}), terminal half-life ($t_{1/2}$), area under curve (AUC_{last} and AUC_{0-inf}) and others were estimated from the data of each individual animal.

[0744] Clearance rates and distribution volumes were estimated however, they were not used for PK comparisons between the two doses and the two administration routes in this report for two reasons (1) this study focused on biodistribution of HNS in solid tissues, rather than on blood PK; and (2) the radioactivity in the brain region was the sum of those from the brain tissue (solid) and CSF (liquid), which could not be separated from each other in the study. The λ_z was evaluated, and used for comparison, because it indicated a percentage of the injected dose eliminated per unit of time.

[0745] The group means and standard deviations (SD) were calculated and compared between two test doses. These PK parameters are tabulated in

Table 48 below

Table 48

1 mg/kg IT	Brain (ug/g)*		Liver		Brain (ug) [#]		Head (ug) [#]		Proximal		Mid		Distal	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
λ_z	0.016	0.003	0.030	0.011	0.017	0.002	0.016	0.002	0.025	0.012	0.020	0.008	0.028	0.016
$t_{1/2}$	45	7	28	16	42	5	45	7	32	13	39	16	30	12
T_{max}	0.1	0.0	2.3	1.3	2.0	4.0	0.1	0.0	0.3	0.5	1.8	1.5	1.0	0.0
C_{max}	257.0	89.9	4.9	1.3	68.6	8.0	200.1	0.0	0.5	0.1	0.2	0.0	0.1	0.0
AUC_{last}	8393	2457	204	50	3809	622	8216	782	9	3	7	3	2	1
$AUC_{int.}$	8942	2416	216	57	4030	643	8904	1069	11	3	8	3	3	2
MRT_{last}	46	6	32	13	44	5	46	5	31	17	34	20	16	5

10 mg/kg IT

λ_z	0.014	0.001	0.017	0.000	0.014	0.001	0.010	0.001	0.018	0.008	0.014	-	0.006	-
$t_{1/2}$	49	4	42	1	51	5	70	9	45	18	50	-	123	-
T_{max}	0.1	0.0	7.0	2.0	0.1	0.0	0.1	0.0	0.3	0.5	8.7	13.3	8.0	-
C_{max}	2628	265	105	41	836	117	1844	314	6	4	1	0	1	-
AUC_{last}	83962	10083	7987	3276	59115	8624	128751	15723	83	67	35	20	38	-
$AUC_{int.}$	89460	12098	8345	3424	63836	9466	151405	15123	98	66	60	-	73	-
MRT_{last}	56	1	51	1	58	2	65	3	31	2	32	7	61	-

1 mg/kg IV	Brain (ug/g)*		Liver		Kidney		Heart		Skin	
λ_z	0.011	0.005	0.015	0.003	0.016	0.009	0.021	0.006	0.021	0.010
$t_{1/2}$	71	23	47	10	54	25	36	15	40	21

2017202349 10 Apr 2017

T _{max}	7	12	5	4	10	12	2	1	5	4
C _{max}	0.1	0.0	9.6	1.5	0.2	0.1	0.2	0.0	0.3	0.1
AUC _{last}	7	2	525	104	14	5	9	3	16	4
AUC _{inf.}	9	3	576	138	16	6	10	3	18	5
MRT _{last}	61	16	47	5	47	18	36	13	41	16

10 mg/kg IV

λ _z	0.102	0.180	0.021	0.012	0.035	0.024	0.020	0.010	0.026	0.012
t _{1/2}	60.5	53.1	37.8	13.4	28.4	16.4	41.6	18.6	31.0	12.7
T _{max}	13	12	2	1	12	11	16	9	3	1
C _{max}	1.8	0.2	131.6	26.8	3.9	0.7	3.7	0.7	7.9	2.3
AUC _{last}	86	66	6747	2837	183	123	201	89	276	40
AUC _{inf.}	118	98	7171	3029	198	131	230	110	292	43
MRT _{last}	43	32	40	14	33	21	41	18	33	13

[0746] In the first 20 min after dosing, total amount (ug) of HNS in the head region was reduced at a constant rate of 0.002 – 0.011 per min (λ_z , 0.005±0.004/min) at 1 mg/kg and 0.003 – 0.010 per min (0.007±0.003/min) at 10 mg/kg. These constant rates of elimination were not significantly different at these two dose levels ($p=0.57$, Figure 151).

[0747] The concentration-time curve (ug/g from 0.05 to 192 hours) for the brain indicated a bi-phasic profile (Figure 152). The early phase lasts for about two hours. The terminal phase follows first-order kinetics. The constant rates of elimination from the brain were very similar at two tested doses (0.0016 ± 0.003 and 0.014 ± 0.001 per hour) with a similar half-life of about two days (45 ± 7 and 49 ± 4 hours at 1 and 10 mg/kg, respectively). The values of peak concentrations (257 ± 90 and 2628 ± 265 ug/g) and AUC_{last} (8393 ± 2457 and 83962 ± 10083 hr.ug/g at 1 and 10 mg/kg, respectively) increase approximately ten-fold when the dose was increased from 1 to 10 mg/kg. These observations indicated a linear PK behavior in the dose range of 1 to 10 mg/kg given in these IT single dosing regimens. The peak concentration appeared in the brain 3 min (T_{max}) after IT dosing.

[0748] The total amount-time curve (ug from 0.05 to 192 hours) in the brain and head regions followed the same bi-phasic pattern as seen with concentration-time curves (ug/g) in the brain (Figure 153 and Figure 154). The values of C_{max} in the brain region were significantly lower than that in the head region (69 ± 8 versus 200 ± 0 at 1 mg/kg, $p<0.01$; and 836 ± 117 versus 1844 ± 314 ug, $p<0.01$ at 10 mg/kg, respectively). The constant rates of elimination were 0.017 ± 0.002/hr and 0.014 ± 0.001/hr for the brain, and 0.016 ± 0.002 and 0.010 ± 0.001/hr for the head region at 1 and 10 mg/kg, respectively. The values of mean residual time were 42 ± 5

2017202349 10 Apr 2017

versus 51 ± 5 hours for the brain ($p=0.048$), and 45 ± 7 versus 70 ± 9 hours for the head ($p<0.01$) at 1 and 10 mg/kg, respectively. These observations suggested that the dosed protein was eliminated from both regions more rapidly at lower dose than at higher doses. The mean half-lives were in a range of 42 to 70 hours in these regions after IT dosing 1 mg/kg and 10 mg/kg of HNS.

[0749] A concentration gradient was observed from the proximal, to the mid and to the distal sections of the spine at both dose levels (data not shown). After IT dosing, the peak concentration (ug/mm of spine column) was seen around 30 min (0 to 1 hour) at the proximal, 1 to 4 hours at the mid (except of one rat being 24 hours) and 1 to 8 hours at the distal section. The half-lives in these sections were variable (mean $t_{1/2}$: 32 ± 13 and 45 ± 18 hours for the proximal, 39 ± 16 and about 50 hours for the mid, and 30 ± 12 and about 123 hours for the distal sections of spine at 1 mg/kg and 10 mg/kg, respectively). The mean values of peak concentrations were roughly proportional to the doses at each of these three sections at 1 and 10 mg/kg of ^{124}I -HNS (0.5 versus 6.0, 0.2 versus 0.9 and 0.1 versus 0.5 ug/mm at the proximal, mid and distal sections of the spine, respectively). The mean values of AUClast followed the same proportional pattern as seen in the peak concentration (9.5 versus 83, 6.8 versus 35, and 2 versus 38 hr.ug/mm at the proximal, mid and distal sections, respectively).

[0750] Even though HNS was not detectable in most peripheral organs, it was measurable in the liver from as early as 1 hour (the first imaging time point after dosing) to 96 hours (three of four animals) at 1 mg/kg and to 192 hours (all four rats) at 10 mg/kg after IT dosing (Figure 158). The concentrations in the liver reached the peak 2 hours after IT dosing of 1 mg/kg, and 7 hours after IT dosing of 10 mg/kg, which was followed by an elimination phase with first-order kinetics. The constant rate of elimination was faster at 1 mg/kg ($\lambda_z 0.030 \pm 0.011/\text{hr}$) than that at 10 mg/kg ($\lambda_z 0.017 \pm 0/\text{hr}$) ($p=0.10$), with a corresponding shorter $t_{1/2}$ (28 ± 16 versus 42 ± 1 hours at the doses of 1 and 10 mg/kg, respectively, $p=0.76$). The value of AUClast at 1 mg/kg reduced about 40-fold in comparison with that at 10 mg/kg (204 ± 50 versus 7987 ± 3276 ug/g, respectively).

Intravenous treatment with ^{124}I -HNS at doses of 1 and 10 mg/kg

2017202349 10 Apr 2017

[0751] The concentration in the brain, liver, kidney, heart (including lung tissue) and skin were plotted as a function of time after IV dosing 1 and 10 mg/kg of HNS as shown in Figure 159 through Figure 163, respectively. Since the first static imaging time point for these organs was one hour after dosing, the initial phase of these concentration-time curves cannot be observed in this study. The concentration-time curves for the liver, kidney, heart and skin showed a flat phase from 1 to 8 hours after IV dosing. This flat phase lasted for 24 hours in the brain post-dosing, suggesting that the brain took up the IV dosed protein slower than that by the peripheral organs. The remaining data indicated a terminal elimination phase with approximately first-order kinetics.

[0752] The elimination half-lives in the liver, kidney, heart and skin 47 ± 10 and 38 ± 13 hours for the liver, 54 ± 25 and 29 ± 16 hours for the kidney, 36 ± 15 and 42 ± 19 hours for the heart and 40 ± 21 and 31 ± 13 hours for the skin at 1 and 10 mg/kg, respectively; while the half-lives in the brain were 71 ± 23 and 60 ± 53 hours (Rat 3 in 10 mg/kg group was excluded for insufficient data to determine $t_{1/2}$) at 1 and 10 mg/kg, respectively. No statistical differences were seen between the half-lives at 1 and 10 mg/kg in these organs, with an exception of p value < 0.03 for kidney.

[0753] The mean values of C_{max} for the liver, skin, kidney, heart and brain were 9.6, 0.3, 0.25, 0.22, and 0.08 ug/g at 1 mg/kg and 132, 7.9, 3.9, 3.7 and 1.8 ug/g at 10 mg/kg. The ratios of C_{max} at 10 mg/kg to the corresponding values at 1 mg/kg were 14, 26, 16, 17 and 23 for these organs. After the C_{max} values from individual animal were normalized for dose, the $C_{max}/dose$ values at 10 mg/kg were significantly higher than that at 1 mg/kg in all these organs (most p values < 0.05 , $p=0.06$ for the liver). The values of AUC_{last} for the liver, skin, kidney, heart and brain were 525, 16, 14, 9.3 and 7 hr.ug/g at 1 mg/kg; and 6747, 276, 183, 201 and 86 hr.ug/g at 10 mg/kg. The ratios of AUC_{last} at 10 mg/kg to the corresponding values of AUC_{last} at 1 mg/kg were 13, 17, 13, 22 and 12 for these organs, respectively. After normalization, the $AUC_{last}/dose$ values at 10 mg/kg were significantly higher than that at 1 mg/kg in the skin ($p < 0.01$), marginally different in the heart ($p=0.06$), and not significantly different in the liver, brain and kidney (all p values > 0.34).

2017202349 10 Apr 2017

[0754] These observations suggested (1) the half-lives in most organs were about 2 days, with the exception of the brain (about 3 days); (2) the exposure per gram in the liver was larger than that of the skin, heart and kidney, which are larger than that of the brain; (3) with a ten-fold increase in dose (10 / 1 mg/kg), the values of C_{max} at 10 mg/kg from all tested organs increased more than 10 times than that at 1 mg/kg.

[0755] The peak concentration in the brain was reached 1- 24 hours (T_{max}) after IV dosing.

Comparison of IV versus IT treatments

[0756] The concentration-time curves in the brain and liver after IV and IT administration at 1 and 10 mg/kg are compared in Figure 164 and Figure 165, respectively. The ratios of C_{max} in the brain by IT/IV at 1 and 10 mg/kg were 3212 and 1501, respectively. These ratios of AUC_{0-192hr} were 1136 and 978. These observations indicated that, when the same dose of HNS was injected, intrathecal administration resulted in an approximately three-log greater exposure of the brain than that with intravenous administration. The elimination half-life in the brain was 2 days (45 and 49 hours at 1 and 10 mg/kg) by IT and 3 days (71 and 60 hours at 1 and 10 mg/kg) by IV administration at both dose levels.

However, hepatic exposures after IT dosing were similar to that after IV dosing at the same dose of HNS. The ratios of C_{max} in the liver by IT/IV at 1 mg/kg and 10 mg/kg were 0.5 and 0.8, and the ratios of AUC_{last} were 0.4 and 1.2, respectively.

Conclusions

[0757] Pharmacokinetic and biodistribution profiles of ¹²⁴I-sulfamidase (HNS) were studied by tissue PET images in rats after single intravenous or intrathecal administration of 1 or 10 mg/kg of ¹²⁴I-sulfamidase. Concentration-time data were obtained both dynamically (the first 20 min) and statically in the regions of interest at 0.05, 1, 2, 4, 8, 24, 48, 96 and 192 hours post dosing. By dynamic imaging after IT dosing, total amount of HNS in the head region was reduced at a similar constant rate of 0.005/min – 0.007/min (mean λ_z) in the first 20 min. By static imaging, the rates of elimination from the brain were essentially the same at two tested

2017202349 10 Apr 2017

doses (λ_z : 0.016/hr versus 0.014/hr for 1 and 10 mg/kg, respectively) with a similar half-life about two days.

[0758] The values of C_{max} and AUC_{last} were proportional to the administered doses, and a linear PK behavior was indicated in the dose range of 1 to 10 mg/kg given in these IT single dosing regimens.

[0759] Concentration gradients were observed from the proximal to distal spine at both dose levels.

[0760] After IT dosing, the peak concentration was seen around 20 min at the proximal, 1 to 4 hours at the mid and 1 to 8 hours at the distal sections. Linear PK behavior was also indicated in the different sections of the spine.

[0761] After IT dosing, HNS protein was measurable in the liver from very early time up to 96 hours at 1 mg/kg and 192 hours at 10 mg/kg. The rate of elimination was faster at 1 mg/kg (λ_z 0.030/hr) than that at 10 mg/kg (λ_z 0.017/hr), with a corresponding shorter $t_{1/2}$ at the lower dose (28 ± 16 versus 42 ± 1 hours at the doses of 1 and 10 mg/kg, respectively).

[0762] After IV dosing, the elimination half-lives in the liver, kidney, heart and skin 47 ± 10 and 38 ± 13 hours for the liver, 54 ± 25 and 29 ± 16 hours for the kidney, 36 ± 15 and 42 ± 19 hours for the heart and 40 ± 21 and 31 ± 13 hours for the skin at 1 and 10 mg/kg, respectively; while the half-lives in the brain were 71 ± 23 and 60 ± 53 hours. The mean values of C_{max} for the liver, skin, kidney, heart and brain were 9.6, 0.30, 0.25, 0.22, and 0.08 ug/g at 1 mg/kg and 132, 7.9, 3.9, 3.7 and 1.8 ug/g at 10 mg/kg. After the C_{max} values from individual animal were normalized for dose, the $C_{max}/dose$ values at 10 mg/kg were significantly higher than that at 1 mg/kg in all these organs (most p values <0.05 , $p=0.06$ for the liver). The values of AUC_{last} for the liver, skin, kidney, heart and brain were 525, 16, 14, 9.3 and 7 hr.ug/g at 1 mg/kg; and 6747, 276, 183, 201 and 86 hr.ug/g at 10 mg/kg. After normalization, the $AUC_{last}/dose$ values at 10 mg/kg were significantly higher than that at 1 mg/kg in the skin ($p<0.01$), marginally different in the heart ($p=0.06$), and not significantly different in the liver, brain and kidney (all p values >0.34).

2017202349 10 Apr 2017

EXAMPLE 23: Treatment of Sanfilippo syndrome type A (Sanfilippo A) patients with HNS

[0763] Direct CNS administration through, e.g., IT delivery can be used to effectively treat Sanfilippo A patients. This example illustrates a multicenter dose escalation study designed to evaluate the safety of up to 3 dose levels every other week (EOW) for a total of 40 weeks of rhHNS administered via an intrathecal drug delivery device (IDDD) to patients with Sanfilippo A. Various exemplary intrathecal drug delivery devices suitable for human treatment are depicted in Figures 94-97.

[0764] Up to 20 patients will be enrolled:

Cohort 1: 5 patients (Lowest Dose)

Cohort 2: 5 patients (Intermediate Dose)

Cohort 3: 5 patients (Highest Dose)

5 patients will be randomized to no treatment.

[0765] Patients are selected for the study based on inclusion of the following criteria:

[0766] Safety of ascending doses of HNS administered by IT injection for 40 weeks in patients with Sanfilippo A is determined. In addition, the clinical activity of HNS on cognitive function, and single and repeated-dose pharmacokinetics in serum and concentrations in cerebrospinal fluid (CSF) are assessed.

EXAMPLES OF IT DELIVERY OF NAGLU-IGFII**EXAMPLE 24: In vitro study of rhNaglu and Naglu-IGFII**

[0767] The mechanism of cellular uptake by each of the Naglu variants was studied using two strains of Sanfilippo B patient fibroblast cells, GM02391 (P359L) and GM 01426 (E153K), and a normal human fibroblast cell line. Attributed to M6P receptor expression on the cell line,

2017202349 10 Apr 2017

fibroblast cells are traditionally used by researchers for the study of lysosomal enzymes cellular uptake.

[0768] Cellular uptake studies were done by incubation of fibroblast cells with rhNaglu or Naglu-IGFII for four hours at 37°C. Cells were washed and lysed after incubation, and Naglu enzymatic activity in cell lysates was measured. Incubation of rhNaglu with fibroblast cells resulted in barely detectable amount of enzyme intracellularly. In contrast, incubation of Naglu-IGFII with fibroblast cells resulted in pronounced level of enzyme intracellularly (Figure 166). The amount of internalized Naglu-IGFII reached saturation as the amount of enzyme used for incubation increased. The dose dependant saturating uptake is a typical finding for receptor mediated cellular uptake. Furthermore, the internalization of Naglu-IGFII was not inhibited by exogenous M6P, but was inhibited by exogenous IGFII completely (Figure 166). This result indicated that Naglu-IGFII internalization into fibroblast cells is dependant on M6P/IGFII receptor in a glycosylation independent manner.

[0769] An experiment was also conducted to study the trafficking of rhNaglu and Naglu-IGFII to lysosomes. Sanfilippo B patient fibroblast cells (GM01426) were used for this study. Detection of rhNaglu and Naglu-IGFII was examined by staining the cells with anti-human Naglu polyclonal antibody after initial incubation of the proteins with the cells. Immunofluorescent staining of LAMP-1 (lysosomal associated membrane protein 1) was used for the detection of lysosomes. Co-localization of rhNaglu and Naglu-IGFII with lysosomes was visualized by confocal microscopy (Figure 167).

[0770] Extensive internalization of Naglu-IGFII was observed after 4 hours of incubation of the protein with the cells, co-localization of Naglu-IGFII with lysosomes was demonstrated. Contrarily, rhNaglu failed to show internalization in the same time frame, and no co-localization with the lysosomes was observed. This result further provided the evidence that Naglu-IGFII was internalized into cells and transported to the correct cellular compartment, the lysosomes. The half life of internalized Naglu-IGFII in Sanfilippo B patient fibroblast cells was determined to be 1.5 days (data not shown).

2017202349 10 Apr 2017

EXAMPLE 25: In vivo studies in mouse models*Wild type (wt) cannulated rat*

[0771] In addition to the Sanfilippo B mouse model, the wt cannulated rat, a non-deficient animal model, was also used for molecule screening *in vivo*. The wt cannulated rats had surgically implanted cannula at the upper lumbar and lower thoracic region of the spinal cord, and a single injection of 35ul to the CSF was done through the cannula. The criteria assessed for molecule screening using this animal model were Naglu activity assay and immunohistochemistry of the brain and spinal cord.

Sanfilippo syndrome type B mouse model

[0772] The mouse model of Sanfilippo syndrome type B (Naglu^{-/-} mouse, Sanfilippo B mouse) was generated by E. Neufeld and colleague (Li HH, et al., PNAS 96(25):14505–14510; 1999). The exon 6 of the mouse's Naglu gene is disrupted by insertion of a selection marker, neomycin resistant gene. The resulting homozygote Naglu^{-/-} mouse are completely Naglu deficient (Figure 168), and have total GAG accumulation in liver and kidney. Despite the total deficiency of Naglu, these mice are generally healthy and have life span of 8-12 month. Changes of other lysosomal enzymes' expression happen at age around 5 months, these changes include compensatory increase of β -galactosidase, α -glucosidase, β -glucuronidase and β -hexosaminidase in liver and brain, elevation of α -L-iduronidase in liver but not in brain, and the reduction of neuraminidase in liver and brain. Death usually occurs as a result of urinary retention and urinary infection. The Sanfilippo B mouse model has been studied extensively in the literature to depict Sanfilippo B pathological changes. The phenotype related to CNS pathology of Naglu^{-/-} mouse is reported to be hypo-activity at the age 4.5 month, but hyperactivity at other ages has also been observed.

[0773] The neuro-pathological changes in Naglu^{-/-} mouse are described as vacuoles and inclusion bodies in neurons, macrophages and epithelial cells as observed by EM (electron-microscopy). These pathological changes typically start at 33 days of age, and progressively worsen as animals get older. Activated astrocyte and microglial cells are also demonstrated by histo-pathological analysis. Biochemical analysis of two gangliosides, GM2 and GM3, showed

2017202349 10 Apr 2017

5 fold and 9 fold increase the brain. (Since GM2 and GM3 are not direct substrates of Naglu, and it could be challenging to demonstrate significant reduction after ERT for short period of time, they were not used as end biomarkers for POC).

[0774] Biochemical analysis was done by measurement of Naglu enzyme activities and GAG levels, histological analysis was done by anti-human Naglu antibody, anti-LAMP-1 antibody, anti-Iba-1 antibody and anti-GFAP antibody immunohistochemistry. The anti-human Naglu antibody used for this study was a mouse monoclonal antibody that doesn't bind endogenous murine Naglu in wt mouse or the mutated Naglu in Sanfilippo B mouse. LAMP-1 immunostaining used an antibody binds to lysosomal membrane protein, lysosomal associated membrane protein-1. Iba-1 staining used an antibody binds to ionized calcium-binding adaptor protein that is specific for microglial and macrophage cells. GFAP staining used an antibody that binds to glial fibrillary acidic protein which is specific for astrocytes.

In vivo biological activity screening by intracranial (IC) injection into Sanfilippo B mouse

[0775] The objective of this study was to evaluate the biological activity of Naglu enzymes *in vivo*. In this study, proteins were administered through IC injection into the brain of the Sanfilippo B mouse. The age of Sanfilippo B mice for the study was closely matched to be at 8 weeks of age. The IC injection route offered the best case scenario to evaluate the efficacy of the molecules. Naglu proteins were assessed by the ability to be taken up into neuronal cells and to reduce lysosomal storage. Immunohistochemistry was used to assess biodistribution. And lysosomal storage was characterized by the number and the size of positive staining using LAMP-1 immunostaining.

[0776] IC injection was done by direct injection through the skull of the Sanfilippo B mouse into the right cerebrum cortex. Two microliters, or 35µg of Naglu protein was injected into each animal. Sacrifices of the animals took place 7-days after injection. The time of sacrifice was pre-determined in a pilot study where sacrifices of the animal took place 3, 7, and 14 day after injection. From the pilot study, it was determined that 7 days post injection is the optimum time for immunohistochemical study. Brain sections were cut transversally (Figure 169), and Naglu and Lamp-1 immunostaining were performed. Cellular uptake into both the neurons and the glial cells in rhNaglu and Naglu-IGFII treated Sanfilippo B mouse was demonstrated by

2017202349 10 Apr 2017

immunohistochemistry using an anti-human Naglu antibody (Figure 170 and Figure 171). There was no significant difference between rhNaglu and Naglu-IGFII treated Sanfilippo B mouse in regards to the cellular uptake was observed. Additionally, LAMP-1 immunostaining of the brain tissue of both the rhNaglu and the Naglu-IGFII treated mouse indicates significant level of reduction of lysosomal storage. The level of lysosomal storage reduction in both rhNaglu and Naglu-IGFII treated groups was almost at the same level of normal wt mouse.

[0777] Reduction of lysosomal storage was also observed in Naglu-TAT, Naglu-Kif and PerT-Naglu tested Sanfilippo B mice after IC injection (data not shown). This study demonstrated the *in vivo* biological activity of all of the variants of Naglu.

[0778] In a separate study, Naglu-deficient mice were IT-administered a vehicle or alternatively one, two or three weekly doses of a recombinant Naglu-IgF-II fusion protein construct (Naglu) in PBS. An untreated wild-type group of mice served as an untreated wild-type control and were administered a vehicle without Naglu. Mice were sacrificed after 24 hours following the final injection, followed by tissue preparation for immunohistochemistry (IHC) and histopathological analysis.

[0779] Distribution of Naglu to the brain tissues of the Naglu-deficient mice was evident following IT-administration of the recombinant Naglu. As illustrated in Figure 172A, IT-administration of the recombinant Naglu to the Naglu-deficient mice resulted in the widespread reduction of cellular vacuolation in the white matter tissues compared to Naglu-deficient mice which were IT-administered the vehicle. Similarly, and as illustrated in Figure 172B, morphometrical analysis revealed a marked reduction in LAMP1 immunostaining in the white matter tissues of the treated mice relative to the untreated Naglu-deficient mice, thereby reflecting an improvement in disease pathology.

[0780] As shown in Figures 173A-B, in each area of brain tissue evaluated (the cortex, caudate nucleus and putamen (CP), thalamus (TH), cerebellum (CBL) and white matter (WM)) the LAMP-positive area was reduced in the Naglu-treated mice relative to the untreated Naglu-deficient control mice, and approached the LAMP-positive area of the wild-type mice. Particularly notable is that the LAMP-positive areas in each area of brain tissue analyzed were

2017202349 10 Apr 2017

further reduced following the IT-administration of two or three doses (Figure 173B) relative to a single dose (Figure 173A) of Naglu.

[0781] These results confirm that IT-administered Naglu is capable of altering progression of lysosomal storage diseases such as Sanfilippo syndrome type B in the Naglu-deficient mouse model, further confirming the ability of IT-administered enzymes such as Naglu to treat the CNS manifestations associated with lysosomal storage diseases, such as Sanfilippo syndrome type B.

Molecule screening by intrathecal (IT) injection into wt cannulated Rat

[0782] This study directly mimics a port-mediated approach for drug administration. Naglu protein was administered via IT injections into wt cannulated rats to determine biodistribution into the parenchyma of the brain.

[0783] The cannula in these animals was placed in the upper lumbar and lower thoracic portion of the spinal cord (Figure 174). Animals were injected with 35 μ l, or 385 μ g of rhNaglu, Naglu-TAT, Naglu-IGFII and PerT-Naglu, through the cannula (due to the solubility limitation, Naglu Kif was injected with only 38.5 μ g, which is 10 fold less than the rest of the Naglu). Sacrifices happened 4hr and 24hr after injections.

[0784] Brain and spinal cord tissues were collected and measured by the Naglu activity assay. In the brain of treated animals, Naglu-TAT and Naglu-IGFII treated animals exhibited higher activity than the rhNaglu and all other Naglu variants treated animals (Figure 175). As a general trend, the Naglu activity was significantly higher in the spinal cord than in the brain for all treated animals (data not shown). This phenomenon may indicate that proteins were taken up more at the site closer to the IT injection.

[0785] Immunohistochemistry analysis indicated that the biodistribution of the Naglu-IGFII treated group was more extensive in the brain than all other Naglu variants treated group 24 hr after IT injections (Figure 176 and Figure 177). In the rhNaglu treated animals the protein was observed in the meninges of the brain only. In the spinal cord section, IHC indicated some

2017202349 10 Apr 2017

cellular uptake of rhNaglu in the neurons of the grey matter, but to a much lesser extent than Naglu-IGFII uptake in the neurons of spinal cord (data not shown).

[0786] In Naglu-TAT IT injected group, even though highest Naglu activity was observed in brain tissue by biochemical analysis, but IHC failed to indicate any Naglu-TAT penetration into the parenchyma of the brain, other than remaining on the meninges. Besides from Naglu-IGFII, all of the other Naglu variants failed to show biodistribution beyond the meninges, a strong testimony of the dependency on M6P/IGFII receptors for the cellular uptake of Naglu in the brain after IT injection. This study pointed to Naglu-IGFII as the lead molecule for drug development for Sanfilippo B.

EXAMPLE 26: PROOF OF CONCEPT STUDY USING NAGLU-IGFII

Experimental design

[0787] The proof of concept study was designed to show both biodistribution and the reversal of lysosomal storage after IT injection of Naglu-IGFII in Sanfilippo B mouse. For this study, three groups of Sanfilippo B mice at 8 weeks of age were treated with an IT injection of Naglu-IGFII. Each IT injection constituted a 10ul volume or 260 ug of Naglu-IGFII. There were three treated groups, 1x injection, 2x injection and 3x injections group. For the 1x injection group, a single dose of protein was administrated at day 0. Animals were sacrificed 24 hr after injection. For the 2x injection group, two IT injections were administrated at day 0 and day 7, and animals were sacrificed 24hr after the last injection. For the 3x injection group, IT injections were administrated at day 0, day 7 and day 14, and animals were sacrificed 24hr after the last injection. Three groups of vehicle treated mouse were also included. For the vehicle control groups, Sanfilippo B mice were injected with vehicle at the same time interval as the treated groups and sacrificed the same way as the treated groups.

[0788] Both biochemical and histological analyses were applied to evaluate the outcome of the study. The biochemical analyses include a Naglu activity assay to measure the amount of enzymes in the tissue and a total GAG assay to evaluate the reduction of lysosomal storage. Liver and brain were the two subjected tissue for biochemical analyses (Figure 178 and Figure

2017202349 10 Apr 2017

179). The histological analyses include H&E staining of the tissues for morphological evaluation (data not shown), and immunohistochemical staining with anti-human Naglu antibody, LAMP, Iba and GFAP (data for Iba and GFAP staining not shown).

[0789] The anti-human Naglu antibody used for this study was a mouse monoclonal antibody that doesn't bind endogenous murine Naglu in wt mouse or the mutated Naglu in Sanfilippo B mouse. LAMP-1 immunostaining used an antibody binds to lysosomal associated membrane protein. Iba-1 staining used an antibody binds to ionized calcium-binding adaptor protein that is specific for microglial and macrophage cells. GFAP staining used an antibody that binds to glial fibrillary acidic protein which is specific for astrocytes.

[0790] Representative microscopic pictures of Naglu immunofluorescence are shown in Figure 180. Figure 181 shows a representative section schematic of the brain. Even though Naglu-IGFII was detected into the cerebral cortex which is closer to the meninges, it was not found in the subcortical region such as the caudate nucleus, the thalamus and the white matter (data not shown). Since the immunostaining of LAMP-1, Iba-1 and GFAP of the same subcortical areas did demonstrate reversal of lysosomal storage, it was believed that the negative immunostaining of Naglu in the deep brain areas was probably due to the sensitivity of the Naglu immunofluorescence.

[0791] Representative microscopic pictures of Lamp-1 immunostaining are shown in Figures 182 through Figures186. To demonstrate the extent of protein distribution and efficacy, cerebral cortex and subcortical regions, such as caudate nucleus, thalamus and white matter, and cerebellar cortex were selected for immunohistological analysis. The result from Iba-1 and GFAP immunostaining (data not shown) indicated that what was seen in the LAMP-1 immunostaining was the combined effect of the changes of microglial cells and astrocytes, the two cell types that were reported to be affected in Sanfilippo B mouse model (Li 2002, Ohmi 2002) in addition to neurons. Due to technical limitations, LAMP-1 immunostaining was not able to reveal lysosomal storage in neurons. To best observe the lysosomal accumulation in neurons, such vacuoles and inclusions, electron microscopy is usually utilized (EM was not included in current study).

2017202349 10 Apr 2017

[0792] It will be appreciated that the identification of cell types was limited to neurons and glial cells. The neurons were typically identified by the relatively large and pale nucleus that contains one or more densely stained nucleoli, and the frequently detectable cytoplasm. The glial cells were generally identified by the small dense nucleus and the inconspicuous cytoplasm. The distinction between the different types of glial cells, such as astrocytes, microglial cells, ependymal cells and oligodendrocytes, is typically best done by staining with cell type specific markers.

[0793] In addition to the reduction of lysosomal storage exhibited by the LAMP-1 immunostaining, the Iba-1 immunostaining indicated the reduction of cell size and number of processes in microglial cells, and GFAP immunostaining indicated the reduction of cell size and length/number of processes in astrocytes, in the cerebral cortex, caudate nucleate, thalamus, white matter and cerebellum after IT injections of Naglu-IGFII (data not shown). Furthermore, histopathological analysis by H&E staining (hematoxylin and eosin) of the brain tissues from the same areas as examined for immunohistochemistry, demonstrated the reduction of vacuoles in glial cell after 3x IT injection of Naglu-IGFII. All of the result mentioned above also suggested the dose-related effect of Naglu-IGFII IT injections.

[0794] The biochemical analyses of Sanfilippo B mice after IT injection of Naglu-IGFII detected Naglu activity in the brain and liver. Efficacy of the Naglu-IGFII was demonstrated by total GAG reduction in the brain and liver. Immunohistochemistry demonstrated the biodistribution of Naglu-IGFII in the parenchyma of the brain. Immunostaining of LAMP-1, Iba-1, GFAP and histopathological analysis by H&E staining exhibited reduction of lysosomal storage, the reduction of size and process by microglial and astrocytes in not only the cerebral cortical area of the brain, but also in the subcortical areas, white matter and cerebellar cortex of the brain.

Conclusions

[0795] Among other things, it has been demonstrated that the fusion protein, Naglu-IGFII, exhibited enzymatic activity *in vitro* toward a substrate that has similar structure to the native substrate of Naglu. *In vitro* cellular uptake study demonstrated that the molecule was taken up to cells by the M6P/IGFII receptor in a manner that was independent of M6P

2017202349 10 Apr 2017

glycosylation. Internalized Naglu-IGFII was shown to co-localize with lysosomes. Naglu-IGFII was shown to reduce lysosomal storage *in vivo* after IC injection into the Sanfilippo B mouse. In comparison to rhNaglu and other Naglu fusions and modifications, Naglu-IGFII surpassed them all in penetrating into the parenchyma of the brain of wt cannulated rat after IT injection. Finally, IT injection of Naglu-IGFII into Sanfilippo B mice demonstrated extensive distribution well beyond the meninges, and observed reversal of lysosomal storage in the cerebral cortex as well as in the subcortical regions. Taken together, these data indicate that Naglu-IGFII is a candidate drug for treatment of Sanfilippo B disease.

EXAMPLE 27: TOXICITY, PHARMACOKINETICS (PK) AND TISSUE BIODISTRIBUTION STUDIES OF IT DELIVERED NAGLU-IGFII

Proof of Concept Studies in Mouse

[0796] Three groups (n=3) of Naglu (-/-) mice were injected with 10 uL containing 260 ug of Naglu-IGFII given as a single bolus IT lumbar injection. The 260 ug dose translates into a 520 mg/kg brain weight dose (mouse brain = 0.0005 kg). One group was injected at Day 0 and sacrificed 24 hr post injection. A second group was injected on Days 0 and 7, and sacrificed 24 hr after the last injection. The third group was injected on Days 0, 7, and 14, and sacrificed 24 hr after the last injection. Each Naglu-IGFII -dosed group was paired with a vehicle control group in order to control for age/disease severity.

[0797] Naglu enzyme activity in the brain and the liver was similar for the three Naglu-IGFII-dosed groups. Comparing rhNaglu enzyme activity in the liver to brain, more than 10-fold rhNaglu enzyme activity was found in the liver. It was contemplated that since levels of rhHNS enzyme activity were comparable in the brain and liver after 1-, 3-, and 6-months of dosing in the pivotal toxicity studies in rats and juvenile monkeys, some portion of rhNaglu dose given to the Naglu (-/-) mice may not have been delivered IT, but rather systemically. Nevertheless, the total GAG level in the brain showed a statistically-significant reduction ($p < 0.05$) after 3 IT injections. A dose-related trend for total GAG level reduction was seen in the livers, which was statistically-significant ($p < 0.05$) in the groups receiving 2 or 3 doses.

2017202349 10 Apr 2017

[0798] The biodistribution of Naglu-IGFII after IT injection was observed well beyond meninges into the parenchyma of the brain, but deep subcortical regions were negative for anti-Naglu antibody immunostaining. A reduction of lysosomal activity by lysosomal-associated membrane protein (LAMP) immunostaining was observed in the groups given 2 or 3 doses only. Areas of lysosomal activity reduction included cerebral cortex and deep subcortical regions of caudate nucleus, thalamus, and white matter. Thus, the reduction of various immunostaining parameters in Naglu-IGFII -dosed animals suggested that therapeutic levels of NAGLU might be present despite the absence of anti-NAGLU immunostaining. An attenuated inflammatory response was evidenced by reduction of glial fibrillary acidic protein (GFAP) immunostaining of astrocytes and reduction of ionized calcium-binding adaptor molecule (Iba) staining of microglia/macrophages in groups given 2 or 3 doses only. Areas of analysis included cerebral cortex and deep subcortical regions of caudate nucleus, thalamus, and white matter.

Studies in Rat

[0799] The S-D rat was selected as the rodent species for toxicological evaluation of IT-administered Naglu-IGFII . As a result, sixteen rats (eight per sex) are dosed with recombinant Naglu-IGFII at the maximal feasible dose (MFD), and at approximately ¼ and ½ the MFD (low- and mid-dose levels, respectively) every 4 days for a total of 8 doses.

[0800] Single-dose PK/biodistribution study in S-D rats is performed to determine CSF and serum concentration, or tissue distribution, respectively, following IT-L administration to male and female animals.

[0801] Toxicology studies are designed to evaluate IT-L administration of Naglu-IGFII from a toxicology and safety pharmacology (neurologic, respiratory, and cardiovascular safety) perspective in both male and female animals. Toxicological evaluation in these studies includes clinical observations, body weights, food consumption, clinical pathology, appropriate safety pharmacology assessments (by physical examination or electrocardiography), gross tissue and microscopic evaluation. A limited number of CSF and serum samples are collected and analyzed for Naglu-IGFII, and for antibodies to the test article. Naglu-IGFII tissue distribution and subcellular localization are quantified by enzyme activity assay and immunohistochemistry,

2017202349 10 Apr 2017

respectively. Additionally, selected studies include a recovery period to assess the reversibility, or potential delayed appearance, of any noted significant toxicological findings.

Studies in Monkeys

[0802] The cynomolgus monkey was been selected as the nonrodent species for toxicological evaluations of IT-administered Naglu-IGFII due to their genetic and anatomical similarity to humans and hence is thought to be the more relevant species. Given that the planned patient population for the Sanfilippo syndrome type B clinical trials is pediatric, a chronic 6-month toxicology study in juvenile cynomolgus monkeys featuring intrathecal drug deliver device (IDDD) administration of Naglu-IGFII is performed. Juvenile cynomolgus monkeys are generally less than 1 year of age at initiation of study (approximately 7-9 months of age) and weigh between 900 g to 1,500 g at study initiation. The data obtained from a 1-month repeated-dose juvenile cynomolgus monkey toxicity study guide the dose level selection and design of the 6-month juvenile monkey study. The repeated-dose toxicology studies are designed to mimic the expected clinical route (IT-L bolus) and frequency of administration (every other week; EOW) over a period of 1 through 6 months.

[0803] As described above, toxicology studies are designed to evaluate IT-L administration of Naglu-IGFII from a toxicology and safety pharmacology (neurologic, respiratory, and cardiovascular safety) perspective in both male and female animals. Toxicological evaluation in these studies includes clinical observations, body weights, food consumption, clinical pathology, appropriate safety pharmacology assessments (by physical examination or electrocardiography), gross tissue and microscopic evaluation. A limited number of CSF and serum samples are collected and analyzed for Naglu-IGFII, and for antibodies to the test article. Naglu-IGFII tissue distribution and subcellular localization are quantified by enzyme activity assay and immunohistochemistry, respectively. Additionally, selected studies include a recovery period to assess the reversibility, or potential delayed appearance, of any noted significant toxicological findings.

EXAMPLE 28: EOW INTRATHECAL ADMINISTRATION OF NAGLU-IGFII

2017202349 10 Apr 2017

[0804] This example was designed to determine the feasibility of IT-lumbar dosing EOW for 6 injections (3 month study) in the Naglu -/- mouse model. This dosing regimen may be more clinically relevant as compared to weekly dosing.

[0805] Eight week old Naglu -/- male and female mice were studied according to the following experimental design:

Table 49: Experimental Design for EOW IT Delivery of Naglu-IGFII

Group	N	Treatment	Dose	Frequency	Sacrifice
A	3	Vehicle	N/A	IT injection EOW for 3 months (total of 6 injections)	24 h after last injection
B	6	Naglu-IGFII	60 mg/kg brain weight (30 ug)	IT injection EOW for 3 months (total of 6 injections)	24 h after last injection

[0806] Physiological studies, including Naglu activity assay on liver, brain and serum, anti-Naglu antibody assay on serum, and BCA assay on liver and brain, were performed. Histological studies, including Naglu IHC on brain, spinal cord and liver, and Lamp staining on brain and spinal cord, were performed.

[0807] Brain, spinal cord and liver were collected and fixed in 10% NBF. Five µm paraffin sections were prepared for histological staining. Immunohistochemical (IHC) staining of Naglu was used to detect cellular uptake of the injected protein. H&E staining was used to observe morphological changes. LAMP, an indicator of lysosomal activity and disease state, GFAP and Iba-1, two CNS pathological markers for activated astrocytes and microglial cells, were used for histopathological improvement evaluation.

[0808] Naglu immunostaining of brain, spinal cord and liver of vehicle and Naglu-IGFII treated mice demonstrated that, in the brain and spinal cord, injected Naglu was detected in meninges (M) only by IHC and no Naglu positive staining was detected in any other regions (Figure 187). In the liver, sinusoidal cells (S) were Naglu positive and no Naglu uptake was found in hepatocytes (H).

2017202349 10 Apr 2017

[0809] LAMP immunostaining and H & E staining of the liver and spinal cord of vehicle and Naglu-IGFII treated mice demonstrated that, compared with the vehicle animals, LAMP staining was decreased throughout in both livers and spinal cords treated with Naglu. H&E staining showed cellular vacuolation in hepatocytes was evidently reduced in the treated group compared with vehicle treated animals(Figure 188 and Figure 189).

[0810] H & E staining of the brain of vehicle and Naglu-IGFII treated mice demonstrated a morphology improvement in the brain after 6 every other week IT injection of Naglu-IGFII for 3 months. In the treated brain, the cellular vacuolation (arrows) in all examined regions decreased compared with the vehicle group (Figure 190)

[0811] LAMP IHC in various brain regions after 6 IT Naglu injections for 3 months demonstrated that, compared with the vehicle treated group, Naglu IT administration to SFB mice resulted in a reduction of lysosomal activity in all examined regions revealed by LAMP immunostaining (Figure 190). This reduction was characterized by the decrease in the number of LAMP positive cells, smaller cell size and lighter staining. A marked reduction was found in the cerebellum and brainstem, which are located in the caudate part of the brain close to the spinal cord, compared with other brain regions. A clear reduction was also found in the deep brain regions, including the white matter, hippocampus and thalamus.

[0812] Iba IHC in various brain regions after 6 IT Naglu injections for 3 months revealed activation of microglial cells (Figure 191). Compared with vehicle treated group, no decrease in the number of positive cells and staining intensity was observed in Naglu treated group. However, the cellular morphology of positive microglial cells changed with reduced cell size in all examined brain regions compared to large and vacuolated one in the vehicle group (inserts).

[0813] GFAP IHC in various brain regions after 6 IT Naglu injections for 3 months revealed astrocytic activation (Figure 192). Compared with the vehicle treated group, GFAP positive staining was decreased in the cerebellum and brainstem, and slightly decreased in other examined regions.

[0814] With respect to cellular uptake, these data demonstrate that in the brain and spinal cord, Naglu was detected in meningeal cells only after 6 time every other week Naglu IGFII IT

2017202349 10 Apr 2017

injection for 3 month. Naglu was undetectable by IHC in any other regions of the brain and spinal cord. In the liver, Naglu positive staining was found in sinusoidal cells.

[0815] In the brain and spinal cord, after 6 every other week IT injection of Naglu-IGFII for 3 months, histopathological improvement was seen throughout the brain and spinal cord even though injected Naglu was undetectable by IHC. H&E staining demonstrated cellular vacuolation reduction in all examined brain regions. LAMP staining decreased throughout treated spinal cords and in all evaluated brain regions including the white matter, hippocampus and thalamus which are deep brain areas, with marked decrease in the cerebellum and brainstem in the Naglu-IGFII treated group. The decreased staining pattern of GFAP staining for astrocytes was consistent with LAMP staining while not dramatically decreased as LAMP. Iba-1 staining showed reduction of the cell size of microglial cells in all examines brain regions. In the liver, H&E staining demonstrated cellular vacuolation reduction with marked reduction in LAMP staining in the Naglu treated group.

EXAMPLE 29: TREATMENT OF SANFILIPPO B PATIENTS

[0816] Direct CNS administration through, e.g., IT delivery can be used to effectively treat Sanfilippo syndrome type B (Sanfilippo B) patients. This example illustrates a multicenter dose escalation study designed to evaluate the safety of up to 3 dose levels every other week (EOW) for a total of 40 weeks of Naglu-IGFII and/or rhNaglu administered via an intrathecal drug delivery device (IDDD) to patients with Sanfilippo B Syndrome. Various exemplary intrathecal drug delivery devices suitable for human treatment are depicted in Figures 94-97.

[0817] Up to 20 patients will be enrolled:

Cohort 1: 5 patients (Lowest Dose)

Cohort 2: 5 patients (Intermediate Dose)

Cohort 3: 5 patients (Highest Dose)

5 patients will be randomized to no treatment.

[0818] Patients are selected for the study based on inclusion of the following criteria:

2017202349 10 Apr 2017

[0819] Safety of ascending doses of Naglu-IGFII administered by IT injection for 40 weeks in patients with Sanfilippo B is determined. In addition, the clinical activity of Naglu-IGFII and/or rhNaglu on cognitive function and single and repeated-dose pharmacokinetics in serum and concentrations in cerebrospinal fluid (CSF) are assessed.

[0820] Typically, a therapeutically effective amount of Naglu-IGFII and/or rhNaglu is administered intrathecally at regular intervals, depending on the nature and extent of the disease's effects, and on an ongoing basis. Administration at an "interval," as used herein, indicates that the therapeutically effective amount is administered periodically (as distinguished from a one-time dose). The interval can be determined by standard clinical techniques. In some embodiments, Naglu-IGFII and/or rhNaglu is administered intrathecally bimonthly, monthly, twice monthly, triweekly, biweekly, weekly, twice weekly, thrice weekly, or daily. The administration interval for a single individual need not be a fixed interval, but can be varied over time, depending on the needs of the individual. For example, in times of physical illness or stress, if anti-Naglu antibodies become present or increase, or if disease symptoms worsen, the interval between doses can be decreased.

[0821] While certain compounds, compositions and methods described herein have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the compounds of the invention and are not intended to limit the same.

[0822] The articles "a" and "an" as used herein in the specification and in the claims, unless clearly indicated to the contrary, should be understood to include the plural referents. Claims or descriptions that include "or" between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention also includes embodiments in which more than one, or the entire group members are present in, employed in, or otherwise relevant to a given product or process. Furthermore, it is to be understood that the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, descriptive terms, etc., from one or more of the

2017202349 10 Apr 2017

listed claims is introduced into another claim dependent on the same base claim (or, as relevant, any other claim) unless otherwise indicated or unless it would be evident to one of ordinary skill in the art that a contradiction or inconsistency would arise. Where elements are presented as lists, (e.g., in Markush group or similar format) it is to be understood that each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements, features, etc., certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements, features, etc. For purposes of simplicity those embodiments have not in every case been specifically set forth in so many words herein. It should also be understood that any embodiment or aspect of the invention can be explicitly excluded from the claims, regardless of whether the specific exclusion is recited in the specification. The publications, websites and other reference materials referenced herein to describe the background of the invention and to provide additional detail regarding its practice are hereby incorporated by reference.

We claim:

1. A method comprising a step of:
administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme at a concentration of greater than about 5 mg/ml.
2. The method of claim 1, wherein the replacement enzyme is at a concentration of greater than about 10 mg/ml.
3. The method of claim 1 or 2, wherein the composition further comprises one or more of (i) a buffering agent, (ii) a surfactant, or (iii) a tonicifier.
4. The method of any one of the preceding claims, wherein the composition has a pH of approximately 3.0-8.0.
5. The method of any one of preceding claims, wherein the composition is administered at a single dose volume of less than 5 mL.
6. The method of any one of preceding claims, wherein the composition is administered at a single dose volume of less than 3 mL.
7. The method of any one of preceding claims, wherein the intrathecal administration of the composition does not result in substantial adverse effect in the subject.
8. The method of any one of the preceding claims, wherein the intrathecal administration of the composition does not result in adaptive T-cell mediated immune response.
9. A method comprising a step of:
administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme in a formulation that is not synthetic CSF.
10. The method of claim 9, wherein the replacement enzyme is at a concentration of greater than about 5 mg/ml.
11. A method comprising a step of:
administering to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a

- replacement enzyme for the lysosomal enzyme, which administering involves intrathecal administration of the composition in absence of concurrent immunosuppressant therapy.
12. The method of claim 11, wherein the method does not involve an immune tolerance induction in the subject being treated.
 13. The method of claim 12, wherein the method does not involve a pre-treatment or preconditioning of the subject using T-cell immunosuppressive agent.
 14. A method comprising a step of:
administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme at a therapeutically effective dose and an administration interval such that at least 10% of normal levels or activities of the lysosomal enzyme in one or more tissues of brain, spinal cord and peripheral organs is achieved.
 15. The method of claim 14, wherein the one or more tissues of brain comprise a menengal tissue.
 16. The method of claim 15, wherein the menengal tissue is selected from the group consisting of pia mater, dura mater, and arachnoid tissue.
 17. The method of claim 16, wherein the one or more tissues of brain comprise a tissue of the cerebrum.
 18. The method of claim 17, wherein the tissue of the cerebrum is a surface or shallow tissue of the cerebrum.
 19. The method of claim 18, wherein the surface or shallow tissue of the cerebrum is selected from the group consisting of pia mater tissues, cerebral cortical ribbon tissues, tissues within 4 mm from the surface of the surface of the cerebrum, and combination thereof.
 20. The method of claim 17, wherein the tissue of the cerebrum is a deep tissue of the cerebrum.
 21. The method of claim 20, wherein the deep tissue of the cerebrum is selected from the group consisting of tissues below the cerebral cortical ribbon, tissues below 4 mm from the surface of the surface of the cerebrum, tissues below 6 mm from the surface of the surface of the cerebrum, tissues below 10 mm from the surface of the surface of the cerebrum, and combination thereof.

22. The method of claim 14, wherein the one or more tissues of brain comprise a tissue of the cerebellum.
23. The method of claim 22, wherein the tissue of the cerebellum is selected from the group consisting of tissues of the molecular layer, tissues of the Purkinje cell layer, tissues of the Granular cell layer, and combination thereof.
24. The method of claim 22, wherein the tissue of the cerebellum is a deep tissue of the cerebellum.
25. The method of claim 24, wherein the deep tissue of the cerebellum is selected from the group consisting of tissues of the Purkinje cell layer, tissues of the Granular cell layer, deep cerebellar white matter tissue, and deep cerebellar nuclei tissue.
26. The method of claim 14, wherein the one or more tissues of brain comprise a tissue of the brainstem.
27. The method of claim 26, wherein the tissue of the brainstem is selected from the group consisting of brain stem white matter tissue and/or brain stem nuclei tissue.
28. The method of claim 14, wherein the one or more tissues of the spinal cord is a surface or shallow tissue of the spinal cord.
29. The method of claim 28, wherein the surface or shallow tissue of the spinal cord is selected from the group consisting of pia matter, the tracts of white matter, and tissue within 4 mm from the surface of the surface of the spinal cord.
30. The method of claim 14, wherein the one or more tissues of the spinal cord is a deep tissue of the spinal cord.
31. The method of claim 30, wherein the deep tissue of the spinal cord is selected from the group consisting of spinal cord grey matter and ependymal cells, and tissue below 4 mm from the surface of the surface of the spinal cord.
32. The method of claim 14, wherein the one or more tissues of brain comprise surface or shallow tissues.
33. The method of claim 32, wherein the surface or shallow tissues are selected from the group consisting of pia mater, dura mater, and arachnoid tissues of menengal, pia mater tissues, cerebral cortical ribbon tissues, tissues within 4 mm from the surface of the surface of the cerebrum, and combination thereof.

34. The method of claim 14, wherein the tissues comprise deep tissues.
35. The method of claim 34, wherein the deep brain tissues are selected from deep white matter, of the cerebrum, deep gray matter of the spinal cord, corpus collosum, periventricular tissue, thalamus, fimbria, tissues below the cerebral cortical ribbon, tissues below 4 mm from the surface of the surface of the cerebrum, tissues below 6 mm from the surface of the surface of the cerebrum, tissues below 10 mm from the surface of the surface of the cerebrum, Purkinje cell layer, tissues of the Granular cell layer, deep cerebellar white matter tissue, and deep cerebellar nuclei tissue, and combination thereof.
36. The method of any one of claims 14-35, wherein the therapeutically effective dose ranges from 0.005 mg/kg brain weight to 100 mg/kg brain weight.
37. The method of any one of claims 14-35, wherein the therapeutically effective dose is greater than 1 mg/kg brain weight.
38. The method of any one of claims 14-35, wherein the therapeutically effective dose is greater than 10 mg/kg brain weight.
39. The method of any one of claims 14-35, wherein the therapeutically effective dose is greater than 30 mg/kg brain weight.
40. The method of any one of claims 14-39, wherein the administration interval is once every two weeks.
41. The method of any one of claims 14-39, wherein the administration interval is once every month.
42. The method of any one of claims 14-39, wherein the administration interval is once every two months.
43. A method comprising a step of:
administering to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme, which administering involves intrathecal administration of the composition so that the replacement enzyme is delivered to a deep brain tissue at least 5 mm below the external surface.
44. The method of claim 43, wherein the replacement enzyme is delivered to a deep brain tissue at least 10 mm below the external surface.

45. The method of claim 43, wherein the replacement enzyme is specifically delivered to lysosomes of the deep brain tissue.
46. The method of any one of claims 43, 44, and 45, wherein the deep brain tissues are selected from deep white matter, of the cerebrum, deep gray matter of the spinal cord, corpus callosum, periventricular tissue, thalamus, fimbria, tissues below the cerebral cortical ribbon, tissues below 4 mm from the surface of the surface of the cerebrum, tissues below 6 mm from the surface of the surface of the cerebrum, tissues below 10 mm from the surface of the surface of the cerebrum, Purkinje cell layer, tissues of the Granular cell layer, deep cerebellar white matter tissue, and deep cerebellar nuclei tissue, and combination thereof.
47. A method comprising a step of:
administering intrathecally to a subject suffering from or susceptible to a lysosomal storage disease associated with reduced level or activity of a lysosomal enzyme, a composition comprising a replacement enzyme for the lysosomal enzyme that is produced from human cells.
48. The method of any one of the preceding claims, wherein the lysosomal storage disease is selected from the group consisting of aspartylglucosaminuria, cholesterol ester storage disease, Wolman disease, cystinosis, Danon disease, Fabry disease, Farber lipogranulomatosis, Farber disease, fucosidosis, galactosialidosis types I/II, Gaucher disease types I/II/III, globoid cell leukodystrophy, Krabbe disease, glycogen storage disease II, Pompe disease, GM1-gangliosidosis types I/II/III, GM2-gangliosidosis type I, Tay Sachs disease, GM2-gangliosidosis type II, Sandhoff disease, GM2-gangliosidosis, α -mannosidosis types I/II, β -mannosidosis, metachromatic leukodystrophy, mucopolipidosis type I, sialidosis types I/II, mucopolipidosis types II /III, I-cell disease, mucopolipidosis type IIIC pseudo-Hurler polydystrophy, mucopolysaccharidosis type I, mucopolysaccharidosis type II, Hunter syndrome, mucopolysaccharidosis type IIIA, Sanfilippo syndrome types A, B or C, mucopolysaccharidosis type IIIB, mucopolysaccharidosis type IIIC, mucopolysaccharidosis type IIID, mucopolysaccharidosis type IVA, Morquio syndrome, mucopolysaccharidosis type IVB, mucopolysaccharidosis type VI, mucopolysaccharidosis type VII, Sly syndrome, mucopolysaccharidosis type IX, multiple sulfatase deficiency, neuronal ceroid lipofuscinosis, CLN1 Batten disease, CLN2 Batten disease, Niemann-Pick disease types A/B, Niemann-Pick disease type C1, Niemann-Pick disease type C2, pycnodysostosis, Schindler disease types I/II, Gaucher disease and sialic acid storage disease.

49. The method of any one of the preceding claims, wherein the lysosomal storage disease is selected from the group consisting of Hunters Syndrome, metachromatic leukodystrophy (MLD) disease, Sanfilippo syndrome type A, Sanfilippo syndrome type B, and globoid cell leukodystrophy (GLD) disease.
50. The method of claim 49, wherein the replacement enzyme is selected from the group consisting of recombinant iduronate-2-sulfatase (I2S), arylsulfatase A (ASA), heparan N-sulfatase (HNS), alpha-N-acetylglucosaminidase (Naglu) and β -galactosidase (GLC).
51. The method of any one of the preceding claims, wherein the replacement enzyme contains mannose-6-phosphate (M6P) residues.
52. The method of any one of the preceding claims, wherein the replacement enzyme is a fusion protein comprising a lysosomal targeting moiety.
53. The method of any one of the preceding claims, wherein the replacement enzyme is delivered to neurons, glial cells, perivascular cells and/or meningeal cells.
54. The method of any one of the preceding claims, wherein the replacement enzyme is further delivered to the neurons in the spinal cord.
55. The method of any one of the preceding claims, wherein the intrathecal administration further results in systemic delivery of the replacement enzyme in peripheral target tissues.
56. The method of claim 55, wherein the peripheral target tissues are selected from liver, kidney, spleen and/or heart.
57. The method of any one of the preceding claims, wherein the intrathecal administration results in lysosomal localization of the replacement enzyme in brain target tissues, spinal cord neurons and/or peripheral target tissues.
58. The method of any one of the preceding claims, wherein the intrathecal administration results in reduction of GAG storage in the brain target tissues, spinal cord neurons and/or peripheral target tissues.
59. The method of claim 58, wherein the GAG storage is reduced by at least 20%, 40%, 50%, 60%, 80%, 90%, 1-fold, 1.5-fold, or 2-fold as compared to a control.
60. The method of any one of the preceding claims, wherein the intrathecal administration results in reduced vacuolization in neurons.
61. The method of claim 60, wherein the neurons comprises Purkinje cells.

62. The method of any one of the preceding claims, wherein the intrathecal administration results in increased enzymatic activity of the replacement enzyme in the brain target tissues, spinal cord neurons and/or peripheral target tissues.
63. The method of claim 62, wherein the enzymatic activity is increased by at least 1-fold, 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold or 10-fold as compared to a control.
64. The method of claim 62 or 63, wherein the increased enzymatic activity is at least approximately 10 nmol/hr/mg, 20 nmol/hr/mg, 40 nmol/hr/mg, 50 nmol/hr/mg, 60 nmol/hr/mg, 70 nmol/hr/mg, 80 nmol/hr/mg, 90 nmol/hr/mg, 100 nmol/hr/mg, 150 nmol/hr/mg, 200 nmol/hr/mg, 250 nmol/hr/mg, 300 nmol/hr/mg, 350 nmol/hr/mg, 400 nmol/hr/mg, 450 nmol/hr/mg, 500 nmol/hr/mg, 550 nmol/hr/mg or 600 nmol/hr/mg.
65. The method of claim 64, wherein the enzymatic activity is increased in the lumbar region.
66. The method of claim 65, wherein the increased enzymatic activity in the lumbar region is at least approximately 500 nmol/hr/mg, 600 nmol/hr/mg, 700 nmol/hr/mg, 800 nmol/hr/mg, 900 nmol/hr/mg, 1000 nmol/hr/mg, 1500 nmol/hr/mg, 2000 nmol/hr/mg, 3000 nmol/hr/mg, 4000 nmol/hr/mg, 5000 nmol/hr/mg, 6000 nmol/hr/mg, 7000 nmol/hr/mg, 8000 nmol/hr/mg, 9000 nmol/hr/mg, or 10,000 nmol/hr/mg.
67. The method of any one of the preceding claims, wherein the lysosomal storage disease is associated with peripheral symptoms and the method further comprises administering the replacement enzyme intravenously to the subject.
68. The method of claim 67, wherein the intravenous administration is no more frequent than monthly administration.
69. The method of any one of the preceding claims, wherein the lysosomal storage disease is associated with peripheral symptoms and wherein the method does not involve administering the replacement enzyme intravenously to the subject.
70. A method of treating Hunters Syndrome comprising a step of administering intrathecally to a subject in need of treatment a recombinant iduronate-2-sulfatase (I2S) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the Hunters Syndrome is reduced in intensity, severity, or frequency, or has delayed onset.

71. The method of claim 70, wherein the at least one symptom or feature of the Hunters Syndrome is cognitive impairment; white matter lesions; dilated perivascular spaces in the brain parenchyma, ganglia, corpus callosum, and/or brainstem; atrophy; and/or ventriculomegaly.
72. A method of treating metachromatic leukodystrophy (MLD) disease comprising a step of administering intrathecally to a subject in need of treatment a recombinant arylsulfatase A (ASA) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the MLD disease is reduced in intensity, severity, or frequency, or has delayed onset.
73. The method of claim 73, wherein the at least one symptom or feature of the MLD disease is increased intracranial pressure, hydrocephalus ex vacuo, accumulated sulfated glycolipids in the myelin sheaths in the central and peripheral nervous system and in visceral organs, progressive demyelination and axonal loss within the CNS and PNS, and/or motor and cognitive dysfunction.
74. A method of treating Sanfilippo syndrome type A (Sanfilippo A) disease comprising a step of
administering intrathecally to a subject in need of treatment a recombinant heparan N-sulfatase (HNS) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the Sanfilippo A disease is reduced in intensity, severity, or frequency, or has delayed onset.
75. A method of treating Sanfilippo syndrome type B (Sanfilippo B) disease comprising a step of
administering intrathecally to a subject in need of treatment a recombinant alpha-N-acetylglucosaminidase (Naglu) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the Sanfilippo B disease is reduced in intensity, severity, or frequency, or has delayed onset.
76. The method of claim 75 or 76, wherein the at least one symptom or feature of the Sanfilippo A or Sanfilippo B disease is hearing loss, delayed speech development, deficits in motor skills, hyperactivity, mental retardation, aggressiveness and/or sleep disturbances.
77. The method of claim 77, wherein the recombinant Naglu enzyme is a fusion protein comprising Naglu and a lysosomal targeting moiety.
78. The method of claim 78, wherein the lysosomal targeting moiety is IGF-II.

79. A method of treating globoid cell leukodystrophy (GLD) disease comprising a step of administering intrathecally to a subject in need of treatment a recombinant β -galactosidase (GLC) enzyme at a therapeutically effective dose and an administration interval such that at least one symptom or feature of the GLD disease is reduced in intensity, severity, or frequency, or has delayed onset.
80. The method of claim 80, wherein the at least one symptom or feature of the GLD disease is irritability, convulsion, mental deterioration, deafness, blindness, myoclonic seizures, excessive muscle tone, developmental delay, regression of developmental skills, hypersensitivity, tremor, ataxia, spasticity, episodic severe vomiting, leukodystrophy, cerebral atrophy, development of globoid cells and/or demyelination.
81. A device for intrathecal administration, comprising
a fluid access port;
a hollow body having a first flow orifice in fluid communication with the fluid access port and a second flow orifice configured for insertion into spinal cord; and
a securing mechanism for securing the insertion of the hollow body in the spinal cord.
82. The device of claim 82, wherein the securing mechanism comprises one or more nobs mounted on the surface of the hollow body and a sutured ring adjustable over the one or more nobs.
83. The device of claim 82 or 83, wherein the fluid access port is comprises a reservoir.
84. The device of any one of claims 82-84, wherein the fluid access port is implantable.
85. The device of any one of claims 82-85, wherein the fluid access port is an injectable port.
86. The device of any one of claims 82-85, wherein the fluid access port is a mechanical pump.

2017202349 10 Apr 2017




Figure 1

2017202349 10 Apr 2017




Figure 2A




Figure 2B

2017202349 10 Apr 2017




Figure 2C

2017202349 10 Apr 2017

GalC Drawings

Summary of Vehicles Tested in Adult Monkeys




Figure 3

10 Apr 2017

2017202349

Thermal Screen of hGalC as A Function of pH

1 mg/mL GalC in 3 mM citrate, phosphate, and borate buffer with 50 mM NaCl




Figure 4

2017202349 10 Apr 2017

Specific Activity of hGalC as A Function of pH

1 mg/mL, GalC in 3 mM citrate, phosphate, and borate buffer with 50 mM NaCl




Figure 5

2017202349 10 Apr 2017




Figure 6




Figure 7

2017202349 10 Apr 2017

AUC Profile as the Function of Salt Concentration 1 mg/mL, 5 mM Na phosphate, pH 6.0




Figure 8

2017202349 10 Apr 2017




Figure 9

2017202349 10 Apr 2017

AUC Profile as the Function of pH 1 mg/mL, 3 mM citrate, phosphate, and borate buffer with 50 mM NaCl




Figure 10

2017202349 10 Apr 2017




Figure 11

2017202349 10 Apr 2017




Figure 12

2017202349 10 Apr 2017




Figure 13

2017202349 10 Apr 2017




Figure 14

2017202349 10 Apr 2017




Figure 15

2017202349 10 Apr 2017




Figure 16

Figure 16

2017202349 10 Apr 2017

Circular Dichroism of hGalC as a Function of pH

1 mg/mL GalC in 3 mM citrate, phosphate, and borate buffer with 50 mM NaCl




Figure 17

2017202349 10 Apr 2017




Figure 18

2017202349 10 Apr 2017




Figure 19




Figure 20

2017202349 10 Apr 2017




Figure 21

2017202349 10 Apr 2017




Figure 22




Figure 23

2017202349 10 Apr 2017




Figure 24

2017202349 10 Apr 2017




Figure 25




Figure 26

2017202349 10 Apr 2017




Figure 27

2017202349 10 Apr 2017




Figure 28




Figure 29

2017202349 10 Apr 2017




Figure 30

2017202349 10 Apr 2017




Figure 31

2017202349 10 Apr 2017




Figure 32

2017202349 10 Apr 2017




Figure 33

2017202349 10 Apr 2017




Figure 34

2017202349 10 Apr 2017




Figure 35

2017202349 10 Apr 2017




Figure 36

2017202349 10 Apr 2017




Figure 37

2017202349 10 Apr 2017




Figure 38

2017202349 10 Apr 2017




Figure 39

2017202349 10 Apr 2017




Figure 40

2017202349 10 Apr 2017




Figure 41

2017202349 10 Apr 2017




Figure 42

2017202349 10 Apr 2017




Figure 43

2017202349 10 Apr 2017




Figure 44

2017202349 10 Apr 2017




Figure 45

2017202349 10 Apr 2017




Figure 46

2017202349 10 Apr 2017

Brain




Figure 47

Liver




Figure 48

2017202349 10 Apr 2017





Figure 49

2017202349 10 Apr 2017

Cortex region, IT Injection



Periventricle region, ICV Injection




Figure 50

2017202349 10 Apr 2017




Figure 51

2017202349 10 Apr 2017





Figure 52


2017202349 10 Apr 2017

I2S Drawings

**Vehicle
cerebral cortex**



**3 dose I2S
cerebral cortex**



**3 dose I2S
cerebellar cortex**




Figure 53

2017202349 10 Apr 2017




Figure 54

2017202349 10 Apr 2017




Figure 55

2017202349 10 Apr 2017




Figure 56




Figure 57

2017202349 10 Apr 2017




Figure 58




Figure 59

2017202349 10 Apr 2017




Figure 60




Figure 61

2017202349 10 Apr 2017




Figure 62




Figure 63

2017202349 10 Apr 2017




Figure 64




Figure 65

2017202349 10 Apr 2017




Figure 66




Figure 67

2017202349 10 Apr 2017




Figure 68




Figure 69

2017202349 10 Apr 2017




Figure 70




Figure 71

2017202349 10 Apr 2017




Figure 72




Figure 73

2017202349 10 Apr 2017




Figure 74




Figure 75

2017202349 10 Apr 2017




Figure 76




Figure 77

2017202349 10 Apr 2017




Figure 78




Figure 79

2017202349 10 Apr 2017




Figure 80




Figure 81

2017202349 10 Apr 2017




Figure 82




Figure 83

2017202349 10 Apr 2017




Figure 84

2017202349 10 Apr 2017




Figure 85

2017202349 10 Apr 2017




Figure 86




Figure 87

2017202349 10 Apr 2017





Figure 88

2017202349 10 Apr 2017




Figure 89A




Figure 89B

2017202349 10 Apr 2017




Figure 89C




Figure 89D

2017202349 10 Apr 2017




Figure 90

2017202349 10 Apr 2017




Figure 91

2017202349 10 Apr 2017




Figure 92

2017202349 10 Apr 2017





Figure 93A

2017202349 10 Apr 2017





Figure 93B

2017202349 10 Apr 2017




Figure 94

Port-A-Cath Low Profile Intrathecal Implantable Access System




Figure 95

2017202349 10 Apr 2017




Figure 96






Figure 97

2017202349 10 Apr 2017


I₁ Purkinje cell



Wild Type



I25-KO-vehicle



I25-KO-idursulfase

Figure 98

2017202349 10 Apr 2017




Figure 99

2017202349 10 Apr 2017




Figure 100

2017202349 10 Apr 2017




Figure 101.

2017202349 10 Apr 2017





Figure 102.

2017202349 10 Apr 2017



10 mg, 6 months
30 mg, 6 months
100 mg, 6 months
No-treatment

Figure 103

2017202349 10 Apr 2017




Figure 104

2017202349 10 Apr 2017






Figure 105


2017202349 10 Apr 2017



**IV (1
mg/kg)**



**IT-L (3
mg)**



**ICV (3
mg)**

Figure 106

2017202349 10 Apr 2017

ASA/MLD Drawings




Figure 107

2017202349 10 Apr 2017




Figure 108

2017202349 10 Apr 2017





Figure 109

2017202349 10 Apr 2017



Small sample size rule applied to calculation of summary variable

Figure 110

2017202349 10 Apr 2017




Figure 111:




Figure 112:

2017202349 10 Apr 2017





Figure 113:


2017202349 10 Apr 2017



WT= Wild type; IT treated mice received 4 weekly doses of 520 mg/kg brain weight of rhASA (0.21 mg), 4X magnification.

Figure 114


2017202349 10 Apr 2017



** P<0.001; * P<0.05; T-Spinal cord = total spinal cord; T-GM = total gray matter; L-GM = lumbar gray matter; C-GM = cervical gray matter; T-WM = total white matter; L-WM = lumbar white matter; C-WM = cervical white matter

Figure 115

2017202349 10 Apr 2017



WT= Wild type; IT treated mice received 4 weekly doses of 520 mg/kg brain weight of HGT-1110 (0.21 mg), 20X magnification.

Figure 116

2017202349 10 Apr 2017




Figure 117

2017202349 10 Apr 2017




Figure 118

2017202349 10 Apr 2017




Figure 119

2017202349 10 Apr 2017




Figure 120

2017202349 10 Apr 2017




Figure 121

2017202349 10 Apr 2017




Figure 122

2017202349 10 Apr 2017




Figure 123

2017202349 10 Apr 2017




Figure 124

2017202349 10 Apr 2017




Figure 125

2017202349 10 Apr 2017




Figure 126

2017202349 10 Apr 2017

107/127




Figure 127:

2017202349 10 Apr 2017

108/127




Figure 128

2017202349 10 Apr 2017

109/127




Figure 129

2017202349 10 Apr 2017




Figure 130

2017202349 10 Apr 2017

111/127





Figure 131

2017202349 10 Apr 2017

112/127





Figure 132

2017202349 10 Apr 2017

62-133

Slice 4

#458



1 cm


University of Wisconsin-Madison Brain Collection

Figure 133: Anatomical Locations of Brain Punches

2017202349 10 Apr 2017

62-133

#780



1 cm

University of Wisconsin-Madison Brain Collection

Figure 134: Anatomical Locations of Brain Punches

2017202349 10 Apr 2017

62-133

#962





Figure 135: Anatomical Locations of Brain Punches

2017202349 10 Apr 2017

62-133

Slice 10

#1102



- Anatomical label
- 15. Corpus callosum and pericallosal subcortical WM
 - 16. Deep Subcortical WM
 - 17. Deep grey
 - 18. Deep grey
 - 19. Periventricular WM
 - 20. Subcortical WM
 - 21. hippocampus

1 cm

University of Wisconsin-Madison Brain Collection


Figure 136:

2017202349 10 Apr 2017

62-133

Slice 12

#1240



Anatomical label
22. Corpus callosum
23. Deep WM

1 cm

University of Wisconsin-Madison Brain Collection


Figure 137

2017202349 10 Apr 2017

62-133

Slice 14

#1486



Anatomical label

24. Subcortical WM, occipital lobe

25. Cerebellar White Matte, including dentate nucleus (WM)

1 cm

University of Wisconsin-Madison Brain Collection

Figure 138

2017202349 10 Apr 2017




Figure 139A




Figure 139B




Figure 139C

2017202349 10 Apr 2017




Figure 139D




Figure 139E




Figure 139F

2017202349 10 Apr 2017




Figure 139G

2017202349 10 Apr 2017




Figure 140A




Figure 140B

2017202349 10 Apr 2017




Figure 141A




Figure 141B

2017202349 10 Apr 2017




Figure 142A




Figure 142B

2017202349 10 Apr 2017




Figure 142C

2017202349 10 Apr 2017




Figure 143

2017202349 10 Apr 2017




Figure 144




Figure 145





Fig. 146A and 146B

Antibody Concentrations in the Serum and CSF after Repeatedly IT Dosing at All Three Dose Levels




Fig. 146C

Representative Images From Study

Figure	1A
Animal	078
Group	1F
Slide	1
Mag.	40X
Stain	HE
Tissue	BRAIN/MENINGES
Note	NEUTROPHILIC INFILTRATES IN THE MENINGES OF A CONTROL ANIMAL. EPITHELIAL CELLS OCCUR VARIABLY IN ANIMALS WITH INTRA-THORACIC CATHETERS.
Figure	1C
Animal	046
Group	5M
Slide	1
Mag.	40X
Stain	HE
Tissue	BRAIN/MENINGES
Note	MENINGEAL EPITHELIAL CELLS IN A HIGH DOSE MALE MENTY OF THE CELLS ARE EOSINOPHILS




Fig. 147A/B

Representative Images From Study

Figure	2A								
Animal	083								
Group	IF								
Slide	6								
Mag.	40X	Stain	HE						
Tissue	BRAIN								
Note	THE ARROW INDICATES AN EOSINOPHIL IN THE PERIVASCULAR SPACE OF A LOW EDGE FEMALE								
Figure	2C								
Animal	013								
Group	IM								
Slide	3								
Mag.	40X	Stain	HE						
Tissue	BRAIN								
Note	EOSINOPHILS IN THE PERIVASCULAR SPACE AND THE ADJACENT PARENCHYMA OF THE BRAIN IN A LOW EDGE MALE								

Fig. 147C/D

Representative Images From Study

Figure	3A
Animal	023
Group	EM
Slide	7
Mag.	40X
Stain	HE
Tissue	SPINAL CORD, CERVICAL
Note	THE ARROWS INDICATE ECSTROPHILS IN THE SPINAL CORD PARENCHYMA. NOTE THE SURROUNDING NEURONS ARE NORMAL.
Figure	3C
Animal	024
Group	EM
Slide	8
Mag.	40X
Stain	HE
Tissue	SPINAL CORD, THORACIC
Note	ARROWS INDICATE ECSTROPHILS. THE BOX INDICATES AN AREA OF POSSIBLE NEUROGLIOSIS. THERE ARE SEVERAL LARGE NEURONS. ALL NORMAL.

Fig. 147E/F




Fig. 148A/B






Fig. 148C/D



8.3 mg dose group




Fig. 149A




Fig. 149B

2017202349 10 Apr 2017




Fig. 150A/B/C/D




Fig. 151




Fig. 152




Fig. 153




Fig. 154




Fig. 155




Fig. 156




Fig. 157




Fig. 158





Fig. 159





Fig. 160




Fig. 161

2017202349 10 Apr 2017




Fig. 162






Fig. 163



Comparison of non-compartmental PK parameters in the brain

Parameter	Unit	IT		IV		Ratio IT/IV
		Mean	SD	Mean	SD	
A_z	1/Nr	0.016	0.003	0.011	0.665	
$t_{1/2}$	hr	45	7	71	23	
C_{max}	ug/g	257	86	0.1	0.6	3212
AUC_{0-192}	hr*ug/g	6386	2457	7	2	1136
A_z	1/Nr	0.014	0.001	0.102	0.160	
$t_{1/2}$	hr	49	4	60	50	
C_{max}	ug/g	2628	265	1.8	0.2	1501
AUC_{0-192}	hr*ug/g	83962	16693	86	99	978

Fig. 164



Comparison of non-compartmental PK parameters in the liver

Parameter	Unit	IT		IV		Ratio
		Mean	SD	Mean	SD	
Az	1/hr	0.030	0.011	0.015	0.003	IT/IV
t _{1/2}	hr	26	16	47	10	
C _{max}	ug/g	4.3	1.3	9.6	1.5	0.5
AUC _{0-192h}	hr*ug/g	204	50	525	104	0.4
Az	1/hr	0.017	0.000	0.021	0.012	
t _{1/2}	hr	42	1	38	13	
C _{max}	ug/g	8.05	4.1	13.15	2.7	0.8
AUC _{0-192h}	hr*ug/g	758.7	32.9	874.7	283.7	1.2

Fig. 165




Figure 2-13 Primary fibroblast cells from normal human (Shire HF1156 cells) were used for cellular internalization study of rhNaglu and Naglu-IGFII. Cellular uptake of rhNaglu was minimum, while the cellular uptake of Naglu-IGFII was much pronounced. The saturating curve of Naglu-IGFII internalization indicated a receptor mediated uptake. This uptake was inhibited by IGFII, but not by mannose-6-phosphate

Figure 166




Figure 2-14. Confocal microscopy study using SanB patient's fibroblast cells (GM01426). Extensive internalization of Naglu-IGFII, and co-localization of Naglu-IGFII with Lamp-1 was observed.

Figure 167





Figure 2-15. Naglu activity in wild type (WT), Naglu-/- (KO) and heterozygote Naglu+/- (Het) mouse. Total deficiency of Naglu in SanB mouse was observed in brain, liver, kidney and spleen.

Figure 168



Sectioning Plan







Figure 169

Vehicle Control



rhNaglu



Naglu-IGFII




Figure 170




Figure 171






Figure 172

1 W Naglu-IGF-II IT Treatment



A

2 W Naglu-IGF II IT Treatments



B

Figure 173

2017202349 10 Apr 2017




Figure 174




Figure 175




Figure 176

2017202349 10 Apr 2017





Figure 177



*Note: split y-axis

Figure 178




Figure 179

2017202349 10 Apr 2017




Figure 180

2017202349 10 Apr 2017




Figure 181




Figure 182

2017202349 10 Apr 2017




Figure 183

2017202349 10 Apr 2017




Figure 184

2017202349 10 Apr 2017




Figure 185

2017202349 10 Apr 2017




Figure 186




Fig. 187




Fig. 188

2017202349 10 Apr 2017




Fig. 189A

2017202349 10 Apr 2017




Fig. 189B

2017202349 10 Apr 2017




Fig. 190A

2017202349 10 Apr 2017




Fig. 190B

2017202349 10 Apr 2017




Fig. 191A

2017202349 10 Apr 2017




Fig. 191B

2017202349 10 Apr 2017




Fig. 192A

2017202349 10 Apr 2017




Fig. 192B