

NACIONES " "  
UNIDAS" "

EP"



Programa de las  
Naciones Unidas  
para el Medio  
Ambiente"



PNUMA

F luvt0NKO K/CFC"

"

WP GR\*F GRKJECT"Y I 6: IK H9""

"42"f g"ugr vko dtg"f g"4238"

Qtki kpcnGURC § QN"

Ugr vko c"Twplp"p" f grEgo k2 "Cuguqt"ElgpVHeq"{"V2 epleq"UVCE+" "  
f grRtqvqeqm" T grvxxq'c'ru'f tgcu"{'c'rc' Hqt c"{'Hwpc'Ukx gust gu"  
Gur gclm gpv'Rtqvgi kf cu"URCY +gp"rc" T gi k p" f grl tcp'Ectkdg"  
"

O lco k" Hqt kf c. '4'6'6'f g'paxlgo dtg'f g'4238"

"  
"  
"  
"  
"  
"  
"  
"

### Propuestas de especies presentadas por las Partes para inclusión en los Anexos del Protocolo SPAW

*Por razones de economía y de medio ambiente, se solicita amablemente a los Delegados descargar de Internet y traer sus copias de los Documentos de Trabajo y Documentos de Información de la Reunión y no solicitar copias adicionales.*

"

, Guv'f qewo gpvq"j c'ukf q'tgr tqf wekf q'ukp"gf lek p'hqto crf'



Estados Unidos de América

Nominación del mero estriado (*Epinephelus striatus*)

para su inclusión en el anexo III

Estados Unidos de América

Nominación del mero estriado (*Epinephelus striatus*) para su inclusión en el anexo III del Protocolo relativo a las Áreas y Flora y Fauna Silvestres Especialmente Protegidas en la Región del Gran Caribe del Convenio para la Protección y el Desarrollo del Medio Marino en la Región del Gran Caribe (Protocolo SPAW, por sus siglas en inglés)

I. Requisitos de la nominación

Gp"mqu"ctvfwqu"33."3; "f gn'Rtqvqeqm"URCY "{ "gp"rcu"fkgevtlegu"{"etkgtkqu"cf qr vcf qu"r qt" rcu"Rctvqu"fg"eqphqto kf cf "eqp"gn'ctvfwq"43."ug"guvcdngegp"mqu"tgs wkukqu"eqp"tgur geq" c"rc" ppo kpek»p"fg"gur geku"0" Nqu"rtqegf ko kgpvqu"r ctc" gpo gpf ct" mqu"cpgzqu." eqpvpgk qu" gp" gn' ctvfwq"33\*6+." guvkr wrcp"s wg" ðewcns wktc"fg"rcu"Rctvqu"r wgf g"ppo kpc" wpc"gur gekg"fg"hnqtc"q" hcwpc"gp"r grki tq"fg"gz vpek»p"q"co gpc| cf c"r ctc"uw"lpenwuk»p"q"uwr tguk»p"gp"guvqu"cpgzqu"{"s wg." f gur w"u"fg"rc"tgxkuk»p"{"rc"gxcmcek»p"ghgewcf cu"r qt"gn'Eqo k² "Cuguqt"Elkpvfwheq"{"V² epkq."rcu" Rctvqu"tgxkuctª p"rcu"ppo kpekqpgu."rcu"rtwgdcu"fqewo gpvrgu"{"mqu"lphqto gu"fg"gn'Eqo k² "Cuguqt" Elkpvfwheq"{"V² epkq"{"eqpukf gtctª p"rc"lpenwuk»p"fg"rc"gur gekg"gp"gn'kucf q0" F lej c"ppo kpek»p" f gdtª "ghgewctug"fg"eqphqto kf cf "eqp"rcu"fkgevtlegu" {"etkgtkqu"cf qr vcf qu"r qt" rcu"Rctvqu"cn' co r ctq"fg"gn'ctvfwq"430" F g"r qt"u"rc"rtgugpv"ppo kpek»p"cdqtf c"mqu"ðEtkgtkqu"o qf kkecf qu" r ctc"rc"lpenwuk»p"fg"gur geku"gp"mqu"cpgzqu"fg"gn'Rtqvqeqm"tgnvwxq" c"rcu"ftgeu"{"Hnqtc"{"Hcwp" Ukkgutgu" Gur gekm gpvg" Rtqvgi kf cu" \*URCY +f g"4236" {" gn'rtqegf ko kgpv" r ctc" r t g u g p v c t " {" cr tqdct"ppo kpekqpgu"fg"gur geku"r ctc"uw"lpenwuk»p"q"uwr tguk»p"gp"mqu"cpgzqu" K" K" {" K60" Hkpcm gpvg."gn'ctvfwq"3; \*5+"dtkpf c"gn'vkr q"fg"lphqto cek»p"s wg"ug"fgdg"lpenwk."gp"rc"o gf kf c"fg"m" r qukdrg."gp"mqu"lphqto gu"r gtvpkpgvqu"rcu"gur geku"r tqvgi kf cu"

Gn'ctvfwq"3"fg"gn'Rtqvqeqm"URCY "fg"hlpg"s wg"gn'cpgzq"KKK"gn'ðcpgzq"cn'Rtqvqeqm"s wg" eqpvkpgp"rc"rkuc"ceqtf cf c"fg"gur geku"fg"hnqtc"{"hcwpc"o ctkpc"{"equvgtc"s wg"ug"r wgf gp"wkkt ct"fg" hqto c'tcekpncn'{"uquvpgkdr"{"s wg"tgs wktgp"fg"rcu"o gf kf cu"fg"r tqveek»p"lpf kecf cu"gp"gn'ctvfwq" 33\*3+e-í0" " Cf goª u." gn' ctvfwq" 33" fg" gn' Rtqvqeqm" gur gekhlec" s wg" ðecf c" Rctv" f gdtª ." gp" eqqr gtcek»p"eqp"rcu"fgª u"Rctvqu." hqto wrct." cf qr vct" {" cr nect" r rcpgu"r ctc"rc"i guk»p" {" gn' cr tqxgej co kgpv"fg"vrgu"gur geku" ð0"

II. Declaración de la nominación y descripción de los apéndices

F g"eqphqto kf cf "eqp" guvqu"tgs wkukqu." mqu" Gucf qu" Wpkf qu"ppo kpcp"cn' o gtq" guvklcf q" \*Epinephelus striatus+r ctc"uw"lpenwuk»p"gp"gn'cpgzq"KKK"fg"gn'Rtqvqeqm"URCY 0"Etggo qu"s wg"gn' ekem"dkqn"i leq"{"mqu"r cvtqpgu"o ki tcvqtq"fg"guvc"gur gekg"tgs wktgp"fg"wp"gp"phqs wg"tgi kqpcn'fg" eqqr gtcek»p"r ctc"uw"eqpugt»xcek»p."cn'eqo q"m"gzki g"gn'ctvfwq"33\*3+0"

Gn' Ugtxleq" P cekqpcn'fg" Rgus wgt"cu" O ctkpcu" \*P O HU." r qt" uwu"uki rcu"gp"lpi rª u"fg"mqu" Gucf qu" Wpkf qu"cpwpek»gn'4"fg"ugr vgo dtg"fg"4236" wpc"tguqmwek»p."cn'ecdq"fg"34"o gugu."cegtec" fg"rc"r gvlek»p"r t g u g p v c f c" r q t " Y k f G c t y " I w c t f k c p u " r c t c " r k u c t " g n ' o g t q " g u v k l c f q " \*Epinephelus striatus+eqo q"co gpc| cf q"q"gp"r grki tq"fg"gz vpek»p."fg"eqphqto kf cf "eqp"rc"rg" {"fg" Gur geku"gp" Rgrki tq"fg"gz vpek»p"fg"mqu"Gucf qu" Wpkf qu" \*GUC." r qt" uwu"uki rcu"gp"lpi rª u"o" Guc"tguqmwek»p"ug" r vdrk»"nvgi q"fg"hkprk ct"wp"gzco gp"fg"rc"ukwcek»p"gp"s wg"ug"gpewgpvtc"gn' o gtq" guvklcf q0"

F gur w<sup>2</sup> u'f g"gzco kpcr "mqu" f cvqu" ekp v h e qu" { "eqo gtekr gu" f kur qpkr gu" o<sup>a</sup> u' h k f g f ki pqu. "gn' P O HU" f g vto k p » "s w g" gn' o gtq" gu t k f q" ewo r i f c" eqp" r c" f g h k p k e p" f g" gur g e k g" co g p c l c f c" { "r t q r w u q" u w" k p e n u k e p" g p" gn' r k u n c f q" f g" r c" GUC 0" Gn' c p w p e k q" e q p v g p" c" w p" k p h q t o g" d k q n i k e q. "gn' ewc n' u g" c p g z c" eqo q" cr<sup>2</sup> p f k e g" C" f g" r c" r t g u g p v g" p q o k p c e k e p 0" Gn' U g t x l e k q" P c e k q p c n' f g" R g u s w g t f c u" O c t k p c u" f g" G u x f q u" W p k f q u" t g e k d k e" eqo g p v t k q u" r A d r i e q u. " k p e n u q" eqo g p v t k q u" k p v g t p c e k q p c r g u. " u q d t g" r c" k p e n u k e p" r t q r w g u c" g p" gn' r k u n c f q" f g" r c" GUC" { " f g u r w<sup>2</sup> u' f g" v q o c t" g p" e q p u k f g t c e k e p" m u" eqo g p v t k q u" r A d r i e q u. "gn' P O HU" tomó la decisión f g h k p k x c de listar gn' o gtq" gu t k f q" eqo q" co g p c l c f q" \* e q p" r t q d c d k f c f" f g" g p v t c t" g p" r g n i t q" f g" g z v k p e k e p" g p" gn' h w w t q" r t g x k u d n g" + c n' c o r c t q" f g" r c" GUC 0" Gn' f l e x c o g p" h k p c n' u q d t g" r c" k p e n u k e p" g p" gn' r k u n c f q" f g" r c" GUC" g u" gn' cr<sup>2</sup> p f k e g" D" f g" r c" r t g u g p v g" p q o k p c e k e p 0

Gn' cr<sup>2</sup> p f k e g" E" f g" r c" r t g u g p v g" p q o k p c e k e p" g u" gn' k p h q t o g" f g" r c" r t k o g t c" t g w p k e p" f g n' I t w r q" f g" V t c d c l q" u q d t g" C i t g i c e k q p g u" f g" F g u x g. " e q p h q t o c f q" r q t" gn' E q p u g l q" f g" C f o k p k u t c e k e p" R g u s w g t c" f g n' E c t k d g" \* E H O E. " r q t" u w u" u k i r c u" g p" k p i r f u+. " r c" E q o k u k e p" f g" R g u e c" r c t c" gn' C v i p v k e q" E g p v t q/ Q e e k f g p v c n' \* E Q R C E Q + " r c" Q t i c p k c e k e p" f g n' U g e v q t" R g u s w g t q" { " C e w f e q r c" f g" E g p v t q" C o<sup>2</sup> t l e c" \* Q U R G U E C + " { " gn' O g e c p k u o q" T g i k a p c n' f g" R g u e c" f g n' E c t k d g" \* E T H O. " r q t" u w u" u k i r c u" g p" k p i r f u+. " O k c o k" 4; / 5 3" f g" q e w d t g" f g" 4 2 3 5 0" " N c" F g e r t c e k e p" f g" O k c o k' t g f c e v c f c" r q t" m u" r c t v k e r c p y g u" f g n' v c m g t" t g e q o g p f ». " i n t e r a l i a, " g u v c d r e g e t" v g o r q t c f c u" t g i k a p c r g u" c t o q p k c f c u" f g" x g f c" r c t c" g u r g e k g u" g u r g e h e c u" s w g" u g" u c d g" u g" e q p i t g i c p" r c t c" f g u x c t" \* e q o g p l c p f q" e q p" gn' o g t q" g u t k f q" { " c i t g i c p f q" q v t c u. " u g i A p" e q t t g u r q p f c +. " t g e q r g e v c t" { " e q o r c t v k" f c v q u" d k q n i k e q u" { " e q o g t e k r g u" u q d t g" r c u" g u r g e k g u" " N c" F g e r t c e k e p" v c o d k e p" j k q" w p" n c o c f q" r c t c" r c" i g u k e p" { " e q p u g t x c e k e p" t g i k a p c n' f g" g u r g e k g u" f g" r g e g u" s w g" u g" e q p i t g i c p" r c t c" f g u x c t" F g" u w o q" k p v g t<sup>2</sup> u' r c t c" r c u" R c t v g u" f g n' R t q v e q m" U R C Y. " r c" F g e r t c e k e p" f g" O k c o k' t g e q o g p f » " ö s w g" m u" o k g o d t q u" f g" r c" E Q R C E Q" r t q r w u k t c p" r k u n c t" g p" gn' c p g z q" K K f g n' R t q v e q m" U R C Y " r c u" g u r g e k g u" s w g" u g" e q p i t g i c p" r c t c" f g u x c t" \* g p" r c t v k e w r t. " gn' o g t q" g u t k f q" { " gn' o g t q" i k i c p v g + i ö 0

### III. Requisitos corroborados de nominación para apoyar la inclusión en el anexo III

C" e q p k p w e k e p. " u g" r t g u g p v c" w p c" t g u g ° c" f g" r c" k p h q t o c e k e p" u q d t g" gn' o g t q" g u t k f q" \* E p i n e p h e l u s s t r i a t u s + r c t c" e q t t q d t c t" m u" t g s w k u k q u" f g" p q o k p c e k e p" r t g u g p v c f q u" g p" r c" u g e e k e p" I. **Requisitos de la nominación** f g n' r t g u g p v g" f q e w o g p v q 0" G u x t g u g ° c" c r q { c" r c" k p e n u k e p" f g n' o g t q" g u t k f q" g p" gn' c p g z q" K K f g n' R t q v e q m" U R C Y 0" U g" r w f g" g p e q p v t c t" k p h q t o c e k e p" o<sup>a</sup> u' f g v c m f c" g p" gn' k p h q t o g" d k q n i k e q" { " g p" gn' f l e x c o g p" h k p c n' u q d t g" r c" k p e n u k e p" g p" gn' r k u n c f q" f g" r c" GUC" \* c r<sup>2</sup> p f k e g u" C" { " D + 0

#### A. Artículo 19(3) – Información que debe incluirse en los informes pertinentes a especies protegidas, en la medida de lo posible

##### 1. Artículo 19(3)(a) – Nombres científicos y comunes de la especie

P q o d t g" e k p v h e q < " E p i n e p h e l u s s t r i a t u s " \* D m q e m" 3 9; 4 +

P q o d t g u" e q o w p g u < " O g t q" g u t k f q. " e j g t p c. " e j g t p c" e t k q m. " o g t q" f g" P c u u c w'

##### 2. Artículo 19(3)(2) – Poblaciones estimadas de la especie y sus zonas de distribución geográfica

Gn'o gtq" gutkcf q" guv<sup>a</sup> " eqo r wguvq" f g" wpc" uqr" r qdrcek»p" gp" vqf c" uw" | qpc" f g" f kvtkdwek»p" i gqi t<sup>a</sup> hlec" { "pq"ug"j cp"kf gpwkecf q"erctco gpvg"uwdgutwewtcu" f ghpkf cu" f g" r qdrcek»p" \*J kpgi ctf pgt" { "Tqugp"3; 94. "Ugf dggt { "et al"03; ; 8. "J cvgrg { "4227-0" "Cwps wg"wp" guwf kq" tgekgpv" \*Lcemuqp" et al. "4236+" lphqto » "wpc" f kgt gpekcek»p" i gp<sup>2</sup> vlec. "pq" kpf lec" wp" cnq" i tcf q" f g" uwd/ gutwewtccek»p" r qdrcekqpcn" gp" vqf c" r" | qpc" f g" f kvtkdwek»p"0" Ewcpf q" ug" eqpukf gtc" gn" guwf kq" f g" " Lcemuqp" et al. " gp" gn" eqpvz vq" f gn" eÀo wq" o<sup>a</sup> u" i tpcf g" f g" r wdkecekqpgu. "s wgf c" ekgtv" kpegt vkf wo dtg" gp" ewcpvq" c" r" uwd/ gutwewtccek»p" r qdrcekqpcn" f gn'o gtq" gutkcf q"0

Gn'o gtq" gutkcf q" ug" gpewgptc" gp" vqf q" gn'o ct" Ectkdg" f guf g" r u" Dgto wf cu" j cuvc" gn" uw" f g" Dtcukf0" " Gu" pcvxq" f g" mu" uki wkgpvu" r c" ugu" Cpi wkr" = Cpxi wc" { "Dctdwc" = Ctwc" = Dcj co cu" = Dctdcf qu" = Dgrleg" = r u" Dgto wf cu" = Kuru" Ecko<sup>a</sup> p" = Eqm qo dlc" = Equc" Tlec" = Ewdc" = Ewtc±cq" = F qo kplec" = Tgr Adrlec" F qo kplecpc" = I w{cpc" Hcpeguc" = I tpcf c" = I wcf cnw g" = I wvgo crc" = I w{cpc" = J ck" = J qpf wtcu" = Lco clec" = O<sup>2</sup> zleq" = O qpwtctc" = Cpxkru" J qrcpf gucu" = Plectci wc" = Rcpco<sup>a</sup> = Rwtg vq" Tleq" = Ucp" Etku»dcn" { "P lxxgu" = Ucpw" Nwe" = Ucp" Xlekgpv" { "ru" I tpcf kpcu" = Uwtlpcu" = Vtkpf cf" { " Vqdc i q" = Kuru" Vwtequ" { " Eclequ" = Guvcf qu" Wpkf qu" \*Hqtkf c" = Kuru" Rgtkf<sup>2</sup> tlecu" O gpqtgu" f g" Guvcf qu" Wpkf qu" = Tgr Àdrlec" Dqrxctkpc" f g" Xgpgj wgrc" = Kuru" X" i gpgu" Dtk<sup>2</sup> plectu" = Kuru" X" i gpgu" f g" GGWWO

**3. Artículo 19(3)(c) – Estado de la protección legal, con referencia a la legislación o regulación nacional pertinente "**

Eqo q" ug" o cphgu»" cttkdc. "gn" P O HU" j c" vqo cf q" r" f gekul»p" f ghpkkxc" f g" r kuct" gn" o gtq" gutkcf q" eqo q" co gpc| cf q" \*eqp" r tqdcdkrf cf" f g" gptct" gp" r gni tq" f g" gz vpek»p" gp" gn" hwwtq" r tgxkldrg" = cn" co r ctq" f g" r" GUC0" Gn" f lewco gp" hpcn" uqdtg" r" kpenmuk»p" gp" gn" r kucf q" f g" r" GUC" gu" gn" r<sup>2</sup> pf leg" D" f g" r" r tguvgpv" p qo kpecek»p0

**4. Artículo 19(3)(d) – Interacciones ecológicas con otras especies y necesidades específicas de hábitat "**

K wcn" s wg" r" o c { qt" c" f g" mu" r gegu" f g" ctt gekhg. "gn" j<sup>a</sup> dkcv" f gn'o gtq" gutkcf q" eco dlc" c" o gf kf c" s wg" gn" r gl" "etgeg0" Nqu" o gtqu" gutkcf qu" o w{ r gs wg" qu" ug" gpewgptcp" gp" o cek qu" f g" o cetqcn cu. "ngej qu" f g" r cuqu" o ctkpqu" { "eqtcrgu" \*Gi i rguvq"3; ; 7. "F cj ni tgp"3; ; : + " ukwcf qu" gp" r tgcu" egtepcu" c" r" equnc" c" r tqhwf kf cf gu" gptg"3" { "6" o O" Ug" j c" f guetkq" s wg" gn" o letq" j<sup>a</sup> dkcv" f gn'o gtq" gutkcf q" tgeke<sup>2</sup> p" cugpvf q" gu" gn" kpvgtkqt" f g" mu" o cek qu" f g" eqtcn" \*Porites" ur r 0" ewdkgvqu" r qt" o cucu" f g" o cetqcn cu" \*r tko qtf kcm gpvg" Laurencia" ur r 0: " cwps wg" c" o gpwf q" gn" j<sup>a</sup> dkcv" ug" j c" ukf q" ekcf q" uko r rgo gpvg" eqo q" Laurencia0" " Nc" gutwewtc" tgvewrct" cdkgtv" f g" guvqu" o cek qu" f g" eqtcn" ewdkgvqu" f g" cni cu" r tqr qtekqpedc" wp" tghwi kq" { "hcekkcdc" gn" o qxko kpvq" f g" mu" kpf kxf vqu" f gptq" f g" mu" kpvgtulekqu" f g" mu" o cek qu" \*Gi i rguvq"3; ; 7+0" Tgekgpvgo gpvg. "co dk<sup>2</sup> p" ug" j cp" tgeqrgvxf qu" o gtqu" gutkcf qu" c" 3: " o " gp" o qp" vewqu" f g" " gueqo dtqu" f g" drcps wkmqu. "Malacanthus plumieri." gpeqpv<sup>a</sup> pf qug" pq" o gpqu" f g" 5" r gegu" lwpvqu" \*Eqrkp" et al"03; ; 9+0" Cu" o kuo q. "ug" j c" kphqto cf q" s wg" ug" cuqekp" eqp" r u" eqpej cu" f gugej cf cu" f g" ectceqn" tqucf q" q" eco dwg. "Strombus gigas." { "qvtqu" f gugej qu" cttgf gf qt" f g" mu" rgej qu" f g" Thalassia" \*Erc { f qp" et al. "4232. "Y kmmwpf ". "eqo gpvctkq" r gtuqpcn"0"

Nqu" o gtqu" lwxgpkrgu" gutkcf qu" r gs wg" qu" uq" eqo wpgu" gp" mu" rgej qu" uqo gtqu" f g" r cuqu" o ctkpqu. "gp" r u" o cetqcn cu" { "crtgf gf qt" f g" mu" o cek qu" f g" eqtcn" f g" Porites" ur r 0c"

o g f k f c 's w g ' g o r k g l c p ' c ' c r g l c t u g ' f g ' u w u ' j ' a d k c w ' f g ' c u g p w c o k g p v q ' q ' o l e t q ' j ' a d k c w ' \* T c p f c m ' 3 ; : 5 . " G i i n g u x p p ' 3 ; ; 7 + 0 " E q p h q t o g ' e t g e g p " r n u " l w x g p k r g u . " u g " t c u r f c p " o c t " c f g p v t q " f g u f g " r n u " r c t e j g u " f g " c t t g e k h g u " r k q t c r g u " j c u v c " r c u " | q p c u " r t q i t g u l x c o g p v g " o a u " r t q h w p f c u " f g " r n u " c t t g e k h g u " h t q p w c r g u 0 " E q o q " c f w n q u . " r n u " o g t q u " g u t k c f q u " u g " e r c u k h e c p " e q o q " r g e g u " f g " c t t g e k h g " { " u g " g p e w g p v t c p " g p " r n u " c t t g e k h g u " f g " v q f q " g n ' E c t k d g 0 " W k k k c p " g n ' c t t g e k h g " e q o q " t g h w i k q " { . " r q t " e q p u k i w k g p v g . " p q " p g e g u k c p " f g " w p " j a d k c v ' x l x q " p k ' f g " e q t c n ' x l x q . " u k p q " A p l e c o g p v g " f g " e k g t c " g u t w e w t c . " w n ' e q o q " j g p f k f w t c u " f g " r n u " c t t g e k h g u " q " g u t w e w t c u " c t w k h e k r g u 0 " "

Gp " e w c p v q " c " r c " k p v t c e e k p " g e q n i k e c . " u g " e c t g e g " g p " i t c p " r c t v g " f g " k p h q t o c e k p " c e g t e c " f g " r c " f g r t g f c e k p " u q d t g " r n u " o g t q u . " c w p s w g " u g " j c " k p h q t o c f q " f g " v d w t q p g u " s w g " c v c e c p " c " r n u " b o g t q u " g u t k c f q u " g p " r n u " c i t g i c e k q p g u " f g " f g u x g " g p " r n u " K u r c u " X i i g p g u " \* Q n g p " { " N c R r e g " 3 ; 9 ; + { " j c { " w p " k p h q t o g ' f g " e c p k d c r k u o q " g p " g u v c " g u r g e k g " \* U k x c " N g g " 3 ; 9 6 + 0 " P q " u g " q d u g t x » " p k p i w p c " f g r t g f c e k p " u q d t g " r n u " r g e g u " g p " f g u x g " g p " N c u " D c j c o c u . " c " r g u c t " f g " r c " r t g u g p e k " f g " v d w t q p g u " g p " g n ' a t g c " \* E q n k p " 3 ; ; 4 + 0 " U g " t g e w g t » " w p " r g l " o w k r c f q . " r q u l d r g o g p v g " c v c e c f q " r q t " w p c " d c t t c e w f c " q " w p " v d w t » p . " f g u r w u " f g " r k d g t c t " k p f k x f w q u " o c t e c f q u . " e t k c f q u " g p " r c d q t c v q t k q " { " u k p " g z r g t k p e k " g p " w p " c t t g e k h g " f g " r n u " K u r c u " X i i g p g u " \* T q d g t w " e t a l . " 3 ; ; 7 + 0 " N c " r t g h g t g p e k " f g " r n u " l w x g p k r g u " r q t " r n u " o c e t q c r i c u " { " p q " r q t " r e j q u " f g " r c u q u " b o c t k p q u " r q e q " f g u r w u " f g " u w " c u g p w c o k g p v q " r t q d c d r g o g p v g " u g " t g r e k q p g . " g p " r c t v g . " c " r n u " o c { q t g u " p k x g r g u " f g " f g r t g f c e k p " g p " r n u " r e j q u " f g " r c u q u " o c t k p q u " \* P c f g e w " c p f " G i i n g u x p p " 3 ; ; 8 + 0 " N q u " k p h q t o g u " u q d t g " r c " f g r t g f c e k p " g h g e w c f c " r q t " g n ' r g l " g u e q t r k p " q " r g l " n g » p " g p v g " r n u " r g s w g o q u " r g e g u " f g " c t t g e k h g " { " g p " r n u " r t k o g t c u " g v c r c u " f g n ' e l e m q " d k q n i l e q " e q p u k w f " g p " w p c " k p s w k g w f " r q t " v q f q " g n ' E c t k d g . " c " o g f k f c " s w g " g u v c " g u r g e k g " k p x c u q t c " u g " g z w k g p f g " \* C n l k p u " c p f " J k z q p " 4 2 2 : - 0 " "

Rqeq " u g " j c " r w d n e c f q " u q d t g " r c " e q o r g v g p e k " k p v t c g u r g e h e c " " q " k p v t g u r g e h e c " g p " g n ' o g t q " g u t k c f q 0 " N q u " l w x g p k r g u " o w g u t c p " c i t g u k p " j c e k " r n u " e q p g u r g e h e c u " f g n ' o k u o q " w o c o q " { " g z j k d g p " c i t g u k p " k p v t g u r g e h e c " \* I 0 F w p j c o . " E c t k d d g c p " O c t k p g " T g u g c t e j " E g p v g t . " e k " H i q t k f c " U c v g " O c t k p g " N c d q t c v q t { . " k p h q t o g " u k p " r w d n e c t " c n ' E c t k d d g c p " O c t k p g " T g u g c t e j " E g p v g t . " 4 ; " f g " o c t l q " f g " 3 ; ; : + 0 " E w c p f q " u g " g p e w g p v t c p " f q u " c f w n q u " p q " t g r t q f w e v q t g u " q " w p " c f w n q " { " w p " l w x g p k i " i t c p f g . " g n ' r g l " o a u " r g s w g o q " c f s w k e t g " g n ' r c v t » p " d l e q m t " f g u e t k q " r c t c " r n u " r g e g u " s w g " u g " e q p i t g i c p . " g p " w p c " u g o c n ' c r c t g p v g " f g " u w o k u k p . " r c t c " n w g i q " i k c t " r e v g t c m g p v g " { " r q t " r n " i g p g t c n ' c r g l c t u g " \* E q n k p " 3 ; ; 4 . " R 0 " E q n k p . " E q t c n " T g g h " T g u g c t e j " H q w p f c v k p " o " R c r w . " e q o g p v t k q " r g t u a p c n i c " I 0 U c f q x { . " P O H U . " 3 ; ; 2 + 0 " "

Gn ' o g t q " g u t k c f q " g u " w p " f g r t g f c f q t " f g " c n q " p l x g n ' g p " r n u " c t t g e k h g u " e q t c r k p q u 0 " N q u " o g t q u " g u t k c f q u " u q p " h q t t e l g c f q t g u " p q " g u r g e k c r k c f q u " f g " u w e e k p " e q p " g o d q u e c f c " \* T c p f c m ' 3 ; 8 7 . " V j q o r u q p " c p f " O w p t q " 3 ; 9 : + s w g " u g " t c i c p " r n u " r t g u c u " g p v t c u " \* Y g t p g t " 3 ; 9 6 . " 3 ; 9 9 + 0 " P w o g t q u q u " g u w f k q u " f g u e t k d g p " s w g " r n u " o g t q u " g u t k c f q u " u q p " r k u e " x q t q u " e w c p f q " u q p " c f w n q u " \* T c p f c m ' c p f " D t q e m ' 3 ; 8 2 . " T c p f c m ' 3 ; 8 7 . " T c p f c m ' 3 ; 8 9 . " R e t t k u j " 3 ; ; 9 . " E c t v g t " e t a l . " 3 ; ; 6 . " G i i n g u x p p " e t a l . " 3 ; ; : + 0 " G u v c " g u r g e k g " e q p u w o g " o w e j q u " v k r q u " { " w o c o q u " f g " c r k o g p v q u " { " u g " v t c u r f c " g p v g " f k h g t g p v g u " j a d k c w . " v c r g u " e q o q " r e j q u " f g " r c u q u " o c t k p q u " { " c t t g e k h g u " e q t c r k p q u . " g p " f k h g t g p v g u " g v c r c u " f g " u w " e l e m q " d k q n i l e q " q " h c u g u " t g r t q f w e v x c u " q " o k g p v t c u " g u v p " e c | c p f q 0 " "

5. " **Artículo 19(3)(e) / "Planes de gestión y recuperación para especies en peligro de extinción y amenazadas "** "

P q"ucdgo qu" f g" plpi Àp" r rcp" r gt vkgpv" f g" i guk»p" q" tgewr gtcekl»p" gp" mqu" Guvcf qu" Wplk qu" q" gp" plpi wpc" qtc" r ctvg" cwps wg" j c { " wpc" co r rlc" i co c" f g" o gecpkuo qu" f g" tgi wrcekl»p" r qt" glgo r mq" gp" Ncu" Dcj co cu. "GGOWW0" Dgrleg. "rcu" Dgto wf cu. "Kircu" Ecko<sup>3</sup> p. " Ewdc. " O<sup>2</sup> zleq. " Vvtequ" { " Eclequ. " Eqmqo dlc" { " qvtqu" s wg" gz kurgp" gp" vqf c" r" | qpc" f g" f kvtkdwek»p" i gqi t<sup>3</sup> hcc" f gr'o gtq" gutkcf q. "eqp" gn' hkp" f g" rko kct" r" ecr wtc" { " cu" o cpvpgt" uw'cdwpc" cpekl0

**6. Artículo 19(3)(f) - Programas de investigación y publicaciones disponibles científicas y técnicas pertinentes a la especie**

Rqt" hcxqt. " tgh<sup>2</sup>tcug" c" mqu" cr<sup>2</sup>pfkegu" rctc" gpeqvtct" wpc" rhuvc" f g" rcu" r wdrlcekl»p" gultghgt gpeku" r tgr ctcf cu" r qt" kpxgukf cf qtgu" r ctvewrtgu" { " r tqi tco cu" f g" kpxgukf cekl»p0"

**7. Artículo 19(3)(g) - Amenazas a las especies protegidas, sus hábitats y sus ecosistemas conexos, especialmente las amenazas que se originan fuera de la jurisdicción de la Parte**

Ncu" r tkpekr crgu" co gpc| cu" cn' o gtq" gutkcf q" uqp" r" ecr wtc" gzegukxc" { " r" qdugtxcpekl»p" kpcf gewcf c" f g" rcu" r g { gu0" Nc" gz vceek»p" hqecrk| cf c" { " hwtvgo" gf kcpvg" r" r gucc" gp" mqu" ukkqu" f g" ci tgi cek»p" f g" f guqyg" gu" wpc" i tcp" kps wgwf "gp" tgrcekl»p" eqp" r" co gpc| c" f g" ecr wtc" gzegukxc0" Nc" gur gekg" j c" f gucr ctgekf q" o c { qto gpvg" eqo q" r gl" f g" ko r qtcpekl»p" eqo gtekn' gp" r" o c { qt" c" f g" mqu" r c" ugu" f g" r" tgi k»p" f gdlf q" c" guvcu" co gpc| cu" r tkpekr crgu0" Nc" ecpvkf cf " eqpqekf c" f g" ci tgi cekppgu" f g" f guqyg" f g" r" gur gekg" ug" j c" xgpkf q" cdclq=" cni wpcu" { c" pq" ug" hqto cp. " vcrgu" eqo q" gp" Ecv' Ec { " Dko kpk" r" r tko gtc" s wg" ug" f guetkdk»p" r ctc" r" ekpkek" \*Gt kico cp" et al. "4235+" { " gp" O cj cj wcn" O<sup>2</sup> zleq" \*Ci wkrct/ Rgtgc. "C04236+" { " vqf cu" rcu" tguvcvgu" ci tgi cekppgu" eqpqekf cu" guv' p" eqpukwkwf cu" r qt" 32" xgegu" o gpqu" r" ecpvkf cf " f g" r gegu" s wg" gp" wpc" qecul»p" wxlgtqp0 Gnl' hwtvgo" cvtcevxq" f g" rcu" ci tgi cekppgu" f g" f guqyg" eqo q" f gukppqu" f g" r gucc. "uw" ko r qtcpekl»p" gp" o wej cu" r gus wgt' cu" f g" vgo r qtcf c" { " r" cr ctgpvg" cdwpc" cpekl»p" f gr' r gl" gp" rcu" ci tgi cekppgu" j cegp" s wg" rcu" ci tgi cekppgu" f g" f guqyg" ugcp" r ctvewrtco gpvg' uwuegr vdrngu" c" r" uqdtgr gucc0

Nc" xwpgtcdkrf cf " f g" guvc" gur gekg" c" r" ecr wtc" gzegukxc" gzi g" gxcvct" uk' mqu" o gecpkuo qu" gz kurgpvu" f g" tgi wrcekl»p" guv' p" eqpvtqrpf q" q" o kki cpf q" r" co gpc| c" f g" uqdtggzr mqcekl»p0 Nqu" o gecpkuo qu" f g" tgi wrcekl»p" r ctc" guvc" gur gekg" uqp" kpcf gewcf qu. "mq" s wg" kpenw' g" eqpvgo r rct" uk' r" qdugtxcpekl»p" f g" guqu" o gecpkuo qu" gu" r" cf gewcf c0 Gnl' i tcf q" j cux" gn' ewcn' mqu" o gecpkuo qu" f g" tgi wrcekl»p" eqpvtqrpf q" r qf t' cp" eqpvtqrct" rcu" co gpc| cu" s wg" eqpvtkw' gp" c" r" f kuo k»pwek»p" f gr' o gtq" gutkcf q" eqpukw' g" wv" hcevqt" emxg0' Gnl' f levc" gp" hpcn' uqdtg' uw' kpenwul»p" gp" gn' hkrvcf q" f g" r" GUC" f g" Guvcf qu" Wplk qu" \*cr<sup>2</sup> pf keg" C+ " gzco k»p" mqu" o gecpkuo qu" f g" tgi wrcekl»p" f g" Ncu" Dcj co cu. " Dgrleg. " rcu" Dgto wf cu. " Kircu" Ecko<sup>3</sup> p. " Ewdc. " O<sup>2</sup> zleq. " Vvtequ" { " Eclequ. " GGOWW0" Eqmqo dlc. " Tgr Àdrlec" F qo kpekcpc" { " rcu" Kircu" X' i gpgu" Dtk<sup>3</sup> pkecu0"

**B. Artículo 21 – Asentamiento de directrices o criterios comunes**

õNqu" etkgtkqu" o qf hccf qu" r ctc" rhuvc" gur gekgu" gp" mqu" cpgz qu" f gn' Rtqvceqm" tgrvwxq" c" rcu" f tgcu" { " Hqtc" { " Hcwp" Ukkxgutgu" Gur gekcm gpvg" Rtqvgi kf cu" \*URCY + { " gn' r tqegf ko kpvq" r ctc"

r t g u p v c t " { " c r t q d c t " p q o k p c e k q p p g u " f g " g u r g e l g u " r c t c " u w " k p e n m u k > p " q " u w r t g u k > p " g p " n q u " c p g z q u " K " K K  
c p f " K K K f g " 4 2 3 6 0 " g p w o g t c d c p " n q u " h c e v t g u " g u r g e " h e q u " s w g " u g " f g d " c p " k p e n m k " g p " g n " c p " a r k u k u " f g " r e u "  
c o g p c | c u " f g " w p c " g x c n w e k > p " e k g p v " h e c " f g n " g u v c f q " c o g p c | c f q " q " g p " r g r i t q " f g " g z v k p e k > p " f g " r e u "  
g u r g e l g u " p q o k p c f c u " 0 R t c " h k p g u " f g " r e " g x c n w e k > p " e k g p v " h e c " f g " r e " e q p f l e k > p " f g " c o g p c | c f c " q " g p "  
r g r i t q " f g " g z v k p e k > p . " c s w " u g " g z c o k p c p " n q u " h c e v t g u " g u r g e " h e c o g p v g " f g u e t k q u " g p " n q u " e t k g t k q u " { " u g "  
c d q t f c p " e q p " o " a u " f g v c m g " g p " n q u " c r " 2 p f l e g u " C " { " D " f g " r e " r t g u g p v g " p q o k p c e k > p 0 " "  
"

## 1. Tamaño de las poblaciones

N c u " r k u v " t q l c " f g " g u r g e l g u " c o g p c | c f c u " f g " r e " W p k > p " k p v t p c e k q p c n " r c t c " r e " E q p u g t x c e k > p " f g "  
r e " P c w t c r g | c " \* W E P + " l p h q t o c " s w g " u q p " t e t q u " n q u " g u k o c f q u " f g n " v o c " o q " f g " r e " r q d r e k > p " f g n " o g t q "  
g u t k c f q " r q t " r c " u = " r g t q " v o d l e p " g u k o c " s w g " g n " v o c " o q " f g " r e " r q d r e k > p " c e w c n " g u " f g " @ 2 . 2 2 2 "  
k p f k x k f v q u " o c f w t q u " e q p " w p c " f k u o k p w e k > p " g u k o c f c " f g " r e " r q d r e k > p " f g " c n " o g p q u " " 8 2 " . " g p "  
e q o r c t c e k > p " e q p " r e u " t g u " A n k o c u " i g p g t c e k q p p u " \* 4 9 / 5 2 " c " o q u " 0 E q o q " r e " f k u o k p w e k > p " g u k o c f c " f g n "  
8 2 " " e w o r n g " e q p " w p q " f g " n q u " e t k g t k q u " f g " r e " W E P " r c t c " w p c " g u r g e l g u " g p " c n q " t k g u i q " f g " g z v k p e k > p " g p "  
r e " p c w t c r g | c . " r e " W E P " e r u k h e c " g n " o g t q " g u t k c f q " e q o q " o g p " r g r i t q " f g " g z v k p e k > p 0 " N c " W E P "  
r t k o g t q " e r u k h e c " g n " o g t q " g u t k c f q " e q o q " g p " r g r i t q " f g " g z v k p e k > p " g p " 3 ; ; 8 = " r e " g x c n w e k > p " o " a u "  
t g e k g p v g " q e w t t k > " g p " 4 2 2 5 " { " g n " o g t q " g u t k c f q " o c p w x q " r e " e r u k h e c e k > p " f g " o g p " r g r i t q " f g " g z v k p e k > p 0 "  
f g " r e " W E P 0 " " "

## 2. Evidencia de disminución

C r c t v g " f g n " g u k o c f q " f g " r e " W E P . " g z k u g " w p c " h e n e " f g " g x c n w e k q p p u " r q d r e k q p c r g u " q "  
g u k o c f q u " f g " r q d r e k > p " u w h e k p v g u " r c t c " g n " o g t q " g u t k c f q . " f g " o c p g t c " s w g " r e u " v p f g p e k u " g p " r e u "  
c i t g i c e k q p p u " f g " f g u x x g " r w g f g p " w k r k c t u g " e q o q " r c t " a o g t q u " f g " r e u " v p f g p e k u " f g " r q d r e k > p 0 " G u "  
r t q d c d r g " s w g " g n " g u v c f q " f g " r e u " c i t g i c e k q p p u " f g " f g u x x g " t g h r g l g " r e " r q d r e k > p " i n d e n " r q t s w g " n q u "  
c f w n q u " o k i t e p " c " r e u " c i t g i c e k q p p u " f g " f g u x x g " r c t c " g u v c t " r t g u g p v g u " g p " n q u " A p l e q u " g x g p v q u "  
t g r t q f w e v x q u " e q p q e k f q u 0 J k r > t l e c o g p v g . " u g " j c d " c p " k f g p v k h e c f q " 7 2 " u k k q u " f g " c i t g i c e k > p " f g "  
f g u x x g " g p " v q f q " g n " E c t k d g " \* U c f q x { " f g " O k e j g u q p " e t a l . " 4 2 2 : + 0 F g " g u v q u " 7 2 . " r t q d c d r g o g p v g " c A p "  
r g t o c p g e g p " o g p q u " f g " 4 2 " \* U c f q x { " f g " O k e j g u q p " e t a l . " 4 2 2 : + 0 G u " o " a u . " c w p s w g " r e u " e c p v k f c f g u " f g "  
r g e g u " g p " n q u " u k k q u " f g " c i t g i c e k > p " u g " e q p v c d c p " ] g p " w p c " q e c u k > p \_ " g p " f g e g p c u " f g " o k r e u " \* 5 2 . 2 2 2 6  
3 2 2 . 2 2 2 " r g e g u = U b k j " 3 ; 9 4 + . " c j q t c " u g " j c p " t g f w e k f q " c " o g p q u " f g " 5 . 2 2 2 " g p " n q u " u k k q u " f q p f g " u g " j c p "  
g h g e w c f q " e q p v g u " \* U c f q x { " f g " O k e j g u q p " e t a l . " 4 2 2 : + 0 G p " i g p g t c n " r e u " g u r g e l g u " f g " r e t i c " x l k c " { "  
n g p v q " e t g e k o k g p v q " \* c r g u " e q o q " n q u " r c t i q u " { " n q u " o g t q u " e q p " r g t " q f q u " r k o k c f q u " f g " f g u x x g . " { "  
r q u k d r g o g p v g " e q p " u q m " w p c " g u t g e j c " x g p v c p c " f g " t g e n w c o k g p v q . " u q p " u w e g r v d r g u " f g "  
u q d t g g z r n v c e k > p " \* D c p p g t q v " e t a l . " 3 ; : 9 . " R q n x k p c " { " T c n r q p " 3 ; : 9 + 0 " " "

G p " v q f q " g n " E c t k d g " u g " l p h q t o c " s w g " g p " t g r g w k c u " q e c u k q p p u " u g " j c p " f g u e w d k g t v q " { " e c r w t c f q "  
c i t g i c e k q p p u " f g " f g u x x g " f g " o g t q u " g u t k c f q u . " r e u " e w c r g u " n w g i q " j c p " f g l c f q " f g " g z k u k k " q " g z k u g p " g p "  
f g p u k f c f g u " v p " d c l c u " s w g " g n " f g u x x g " h c e c u c 0 N q u " o g t q u " g u t k c f q u " h w g t q p " r g u e c f q u " g z e n w u k x c o g p v g "  
f w t c p v g " r " h q t o c e k > p " f g " r e u " c i t g i c e k q p p u " g p " r e " f 2 e c f c " f g " 3 ; 9 2 " g p " r e u " D g t o w f c u " 0 N c u " e c r w t c u "  
e q o g t e k r g u " g p " 3 ; 9 7 " h w g t q p " f g " 9 7 0 2 2 2 " v q p g r c f c u = r c t c " 3 ; : 3 . " r e u " e c r w t c u " j c d " c p " f g u e g p f k f q " c "  
3 2 0 2 2 " v q p g r c f c u " \* U c f q x { " f g " O k e j g u q p " c p f " G t k u o c p " 4 2 3 4 + 0 N q u " e w c v t q " u k k q u " f g " c i t g i c e k > p " f g "  
f g u x x g " e q p q e k f q u " f g l c t q p " f g " h q t o c t u g " r q e q " f g u r w 2 u " { " c A p " p q " u g " j c p " t g e w r g t c f q " f g u f g " g p v q p e g u "  
\* U c f q x { " f g " O k e j g u q p " c p f " G t k u o c p " 4 2 3 4 + 0 G p " O c j c j w e n " S w k v c p c " T q q . " O 2 z l e q . " u g " h q t o c d c p "  
e c f c " c " o q " c i t g i c e k q p p u " f g " j c u w " 3 7 0 2 2 2 " r g e g u " g p " g n " o k u o q " u k k q = r g t q " f g d k f q " c " r e " o c { q t " r t g u k > p "  
f g " r e " r g u e c " g p " n q u " c " o q u " p q x g p v c u . " p q " u g " j c p " h q t o c f q " c i t g i c e k q p p u " f g u f g " 3 ; ; 8 " { " p q " u g " j c p " r w g u v q "  
g p " r t a e v e c " r e u " o g f k c u " f g " i g u k > p " f g u k p c f c u " c " r t q v g i g t " r e u " c i t g i c e k q p p u " f g " f g u x x g " \* C i w k r t /

Rgtgc"4229+0'Nqu"o gtqu" gutkcf qu" gtcp" dwuecf qu" gzenwukxco gpvf" f wtcpvg" r" hqto cekp" f g" rnu" ci tgi cekppgu" gp" Ewdc=" f gdlf q" c" gmj. "42" f g" rnu" 43" ci tgi cekppgu" eqpkef cu" { c" pq" ug" hqto cp" o" a" u" \*Erctq" et al. "422; +0'Gp" Dgrleg. "gn'wco c° q" f g" r" ci tgi cekp" gp" gn'I nqxtg" u" Tggh" j c" f kuo kpwk q" qej gpc" r qt" elgpv" gp" rnu" Anko qu" 47" c° qu" \*f g" 37" 022" r gegu" c" 5022+0' Cf go" a" u. "cr gpcu" 4" f g" rnu"; " ci tgi cekppgu" eqpkef cu" vqf cx" f" c" ug" hqto cdc" p" r ctc" 4223" { " gucu" ug" j cd" f" cp" tgf wekf q" f g" 52" 022" r gegu" c" 302267" 022" r gegu" 0' Vtcdclq" tgekpgvu" j cp" kf gp" v" k" ecf q" 37" ukkqu" f g" ci tgi cekp" f g" f guqyg" gp" Dgrleg" 0' Ugvg" f g" guvqu" ukkqu" ug" o" qpkqtgctq" p" f wtcpvg" wp" r gt" f" qf q" f g" 32" c° qu" \*422564234+0' Nc" ecpv" k" cf" f g" r gegu" eqpvcf qu" gp" rnu" ukvg" ukkqu" j c" r gto cpgek q" o" w" f" dclc" \*ekpeq" ukkqu" v" k" p" gp" o" gpqu" f g" 392" r gegu. " rnu" qvtqu" f qu" v" k" p" gp" 3072" { " 3072+." ukp" p" kpi wpc" ug" o" cn' f g" tgewr gtcekp" \*i twr q" f g" v" tcdclq" URCI " f g" Dgrleg. "4234+0' Ug" ucdg" s" wg" j cp" qewttk" q" ukwcekppgu" uko krtgu" gp" Ncu" Dcj co cu. " rnu" Kircu" X" f" i" gpgu" f g" GGWWO" Rwtv" q" Tleq" { " J" qpf wtcu" \*Ucf qx { " f g" O" ke" j" guq" p" cpf" Gt" kuo cp" 4234. " xgt" v" co dk" p" J" kn' c" p" f" Ucf qx { " f g" O" ke" j" guq" p" 4235+0' }

Qtqu" kpf kecf qtgu" o" a" u" f g" r" f kuo kpwk" p" f g" r" r" qdrcek" p" uq" p" gn' wco c° q" tgf wekf q" { lq" r" o" gpqt" gf cf" f g" rnu" r gegu" gp" o" wej cu" f g" rnu" ci tgi cekppgu" f g" f guqyg" s" wg" uwdukr" gp" 0' Gu' tctq" qd" v" p" gt" kpf kxk" vqu" f g" o" a" u" f g" 34" c° qu" f g" gf cf" gp" rnu" r gus wgt" f" cu" g" zr" m" v" cf" cu. " o" k" p" v" t" cu" s" wg" gp" rnu" | qpcu" f g" r g" u" ec" "gzegukx" c" ug" gpew" p" v" t" cp" r gegu" o" wej q" o" a" u" l" x" gp" gu" gp" r" tqo" gf" kq" 0' Nc" gf cf" o" a" z" k" c" g" u" k" o" cf" c" r ctc" r" r" qdrcek" p" o" w" f" g" zr" m" v" cf" c" f g" rnu" Kircu" X" f" i" gpgu" f g" GGWWO" gu" f g"; " c° qu" \*Qn" gp" cpf" Nc" Rrcg" 3; 9; + "34" c° qu" gp" gn' p" q" t" v" g" f g" Ewdc. "39" c° qu" gp" gn' uw" f g" Ewdc" \*Erctq" et al. "3; ; 2+ { "43" c° qu" gp" Ncu" Dcj co cu. \*Ucf qx { " cpf" Eq" k" p" 3; ; 7+0' Nc" o" c { qt" f" c" f g" rnu" kpf kxk" vqu" ecr w" t" cf" qu" gp" wpc" ci tgi cekp" f g" f guqyg" f g" rnu" Kircu" X" f" i" gpgu" f g" GGWWO" v" p" f" c" wpc" NV" f g" cr" tqz" k" o" cf" co" gpv" g" p" v" t" g" 722" { "822" o" o" \*Qn" gp" cpf" Nc" Rrcg" 3; 9; +0' P" go" gj" et al. "4228+ f" g" u" e" w" dt" k" gt" q" p" s" wg" rnu" o" gtqu" gutkcf qu" cf" w" n" qu" gp" wp" ukkqu" f" k" h" gt" gp" v" g" f g" ci tgi cekp" f g" f guqyg" \*i" t" co" o" cpkni" Dcpni" gp" rnu" Kircu" X" f" i" gpgu" f g" GGWWO" o" gf" f" c" p" gp" v" t" g" 6: 2" { " : 22" o" o" . " uk" gp" f" q" u" go" gl" cp" v" g" r" n" q" pi" kwf" " v" q" w" ni" \*NV+ r" tqo" gf" kq" r" ctc" o" ce" j" qu" \*825" o" o" . " p" ? " 3: + { " j" go" dtcu" \*7; 3" o" o" . " p" ? " 66+0' }

**3. Restricciones en su zona de distribución geográfica**

Nc" f kurtkdwel" p" eqphkto cf c" f gn' o" gtq" gutkcf q" kpenw" g" ce" w" cm" g" p" v" g" ð" Ncu" Dgto w" f" cu" { " Hqtk" c" \*GGWW+." r qt" v" qf" cu" Ncu" Dcj co cu" { " gn' o" ct" Ectk" d" g" o" r" 0" gl" 0" J" g" go" utc" cpf" T" cpf" cm" 3; ; 5+0' Nc" gur" gek" g" u" v" " co" r" n" co" gpv" g" f kurtkdwel" c" { " uw" a" tgc" ce" w" cn' f g" f kurtkdwel" p" gu" r" ct" gek" c" c" uw" a" tgc" j" k" v" t" k" ec" 0' Gu" c" eqpenw" ul" p" ug" d" cuc" gp" r" u" ge" ek" p" u" q" dt" g" \ qpc" E" qo" r" r" g" v" c" f g" F" kurtkdwel" p" f" gn' l" p" h" q" to" g" dkq" n" i" leq" \*cr" 2" p" f" leq" C+." r" e" w" cn' eqpenw" { »" s" wg" r" k" p" h" q" to" cekp" f" k" u" r" q" p" k" d" r" g" u" w" i" k" g" t" g" s" wg" uw" a" tgc" ce" w" cn' f g" f kurtkdwel" p" gu" gs" v" k" x" c" r" gp" v" g" c" uw" f kurtkdwel" p" j" k" v" t" k" ec. " c" w" ps" wg" r" c" d" w" p" f" c" pek" ug" j" c" ci" q" v" cf" q" 0' }

**4. Grado de fragmentación de la población**

P" q" j" c { " p" k" pi" À" p" k" p" f" lek" q" f g" s" wg" r" h" ci" o" gpv" cek" p" f g" r" r" qdrcek" p" ugc" wpc" co" gpc" |" c" q" r" g" t" c" v" k" c" r" ctc" gn' o" gtq" gutkcf q" 0' Vcn' eqo" q" ug" g" zr" t" g" u" »" c" tt" k" dc" gp" KKK" 0. " gn' o" gtq" gutkcf q" gu" v" " eqo" r" w" gu" q" f g" wpc" u" q" r" r" qdrcek" p" gp" v" qf" c" uw" | qpc" f g" f kurtkdwel" p" i" g" qi" t" a" h" ec" uk" p" g" x" k" f" g" pek" f g" wpc" eqpvtcek" p" f gn' a" tgc" f g" f kurtkdwel" p" 0' }

**5. Biología y comportamiento de la especie, así como otros aspectos de la dinámica poblacional**

Eqo q"ug"o gpekq»"cpvgtkqto gpvg."gn'eqo r qtvcō kgpvq" f g" f guqyg" f g" guv" gur gekg" r" xwngk" g" uwegr vdrng" c" r" uqdtggzr mvcel»p0' Gr' o gtq" gvtkcf q" o ki tc" c" ukkqu" r txxkukdrgu" gp" o qo gpvqu" r txxkukdrgu" r ctc" f guqxt" f wcpvg" uqm" wpcu" ewcpvcu" ugo cpcu" ecf c" c° q0' Ug" ucdg" swg" r" tgr tqf weel»p" uqm" qewttg" f wcpvg" r" u" ci tgi cekqpgu" cpwergu. "gp" r" u" ewergu" f guqxcp" eqrgevxco gpvg" i tcpf gu" ecpvk cf gu" f g" o gtqu" gvtkcf qu. "s wg" xcp" f guf g" r" u" f gegpcu" j cuvc" r" u" f gegpcu" f g" o krgu" \*Uo kj "3; 94." Qnrgp" cpf "NcRreg"3; 9; . "Eqrkp" et al. "3; ; 9." Hkp"3; ; 2." Hkp"3; ; 4." Eqrkp"3; ; 4+0' O wej qu" r gegu" xkclcp" i tcpf gu" f kucpeku" r ctc" mgi ct" c" mqu" mwi ctgu" r txxkukdrgu" f wcpvg" r" u" r qecu" ugo cpcu" ecf c" c° q. "s wg" ug" gz kppf gp" f wcpvg" o gugu. "ewcpf q" qewttg" gn' f guqyg" { "mgi q" tgi tguv" c" uwa" cttgekhu" f g" qtki gp" \*Ucf qx { "cpf "Gmwpf "3; ; ; +0"

Gr' o gtq" gvtkcf q" ug" eqpi tgi c" gp" i tcpf gu" ecpvk cf gu" r ctc" f guqxt" ecf c" c° q" r" ci tgi cek»p" o a u" i tcpf g" guwf kcf c" vgp" c" wp" guvko cf q" f g" 52.222/322.222" r gegu" gp" f guqyg" \*Uo kj "3; 94+" gp" Dko kpk" Dc j co cu0' J cuvc" f qpf g" ug" ucdg. " vqf c" r" cevxkcf cf " tgr tqf wevxk" qewttg" gp" guvcu" ci tgi cekqpgu" s wg" ug" hqto cp" eqpukvpgvo gpvg" gp" ukkqu" \*ōj wgequ" f g" o gtqu" { " o qo gpvqu" gur ge" h" equ0' Ncu" ci tgi cekqpgu" j cp" eqpukvq" gp" egpvpcu" gu. "o krgu" q. "j k»tkeco gpvg. " f gegpcu" f g" o krgu" f g" kpf kxf wqu" { " j cp" r gtukvq" gp" wlecekqpgu" eqpkef cu" f wcpvg" r gt" qf qu" f g"; 2" c° qu" q" o a u" \*Uo kj "3; 94." Qnrgp" cpf "NcRreg"3; 9; . "Eqrkp" et al. "3; ; 9." Hkp"3; ; 2." 3; ; 4." Eqrkp"3; ; 4." Ectvgt" et al. "3; ; 6." Ucf qx { "3; ; 9." T0Emtq. "Ncdqtcvqt { "qh' Hkuj " Geqrmi { "ō' Ewdc. "eqo gpvctkq" r gtuqpcn" c" [ 0' Ucf qx { . "P O HU."3; ; 3+0'

**6. Otras condiciones que aumenten claramente la vulnerabilidad de la especie**

Vcn'eqo q"ug" f guetkdk»" gp" r" ugeek»p" f g" co gpc| cu" KKC0" cttkdc. "pwgustq" f lewco gp" r ctc" uw" kpenuk»p" gp" gn' r kncf q" f g" r" GUC" f g" GGOWW0' f guewdtk»" s wg" r" ecr wtc" gzegukc" { " r" cr rkecel»p" kpcf gewcf c" f g" r" r" { " uqp" r" u" f qu" r tpekr crgu" co gpc| cu" r ctc" gn' o gtq" gvtkcf q0' Nqu" cr 2 pf legu" C" { " D" qht gegp" o a u" f gvcng. "cf go a u" f g" r" ugeek»p" KKC00'

**7. Importancia de la especie para mantener ecosistemas y hábitats frágiles o vulnerables**

Eqo q" f grtgf cf qt" uwr tgo q" gp" gequkugo cu" h<sup>a</sup> i krgu" f g" cttgekhu. " gn' o gtq" gvtkcf q" f gugo r g° c" hmpkqpgu" geqn»i kecu" s wg" c" Ap" ug" guv" p" cerctcpf q" \*O wo d { " et al. "4228+0' Uw" r tguvpek" o cpvkpg" c" mqu" j gtd" xqtqu" { " r" r" tguk»p" f g" j" gtdkxqt" c" uqdtg" r" u" cni cu" f g" cttgekhu. " m" s wg" r tguv" wp" dgpghkq" ko r qtvcpg" c" mqu" eqtcrgu" \*O wo d { " et al. "4228+0' Gr' o gtq" gvtkcf q" f grtgf cf qt" r wgf g" c { wf ct" c" rko kct" gn' ko r cevq" f gn' kpxcukxq" r g| " gueqtr k»p" r gtq" gp" guv" o qo gpvq" r" gxf gpek" guv" rglqu" f g" ugt" eqpenw { gpvg" \*O wo d { " et al. "4233+0' Ug" j c" gur gewxf q" s wg" uw" cwvpek" chgevc" r" f kvgpuk»p" geqn»i kec" f g" mqu" f grtgf cf qtgu" o a u" r gs wgo qu. " kpenwk qu" mqu" o gtqu" r gs wgo qu. " eqp" eco dkqu" tguwncpvgu" gp" r" u" tgrcekqpgu" t» hkecu" f g" mqu" gequkugo cu" f g" cttgekhu" \*Ucnkpi u" 422: . " O wo d { " et al. "4234+0'

**C. Artículo 11(1) – Utilidad de los esfuerzos regionales de cooperación ""**

Ug" ucdg" s wg" mqu" o gtqu" gvtkcf qu" o ki tcp" ekgpvqu" f g" nk»o gvtqu" cvcxgucpf q" htpvgtcu" lwtkuf keekpvcrgu" r ctc" mgi ct" c" mqu" ukkqu" gur ge" h" equ" f g" f guqyg0' Vcn'eqo q" ug" j c" r tguvpcf q" gp" pwo gtqucu" qecukqpgu. " guvqu" ukkqu" f g" ci tgi cekqpgu" f g" f guqyg" uqp" t cpukqt kqu. " gur ge" h" equ" f g" wp" mwi ct" { " uwrgp" ugt" eqpkef qu" r qt" mqu" r guecf qt gu" mcecrgu" s wg" mqu" r guecp" kpvpguco gpvg" f wcpvg" gn' r gt" qf q" f g" f guqyg" \*Dqrf gp. "4222+0' Nc" ecr wtc" gzegukc" gu" gur gekm gpvg" r tggewr cpvg" f gdkf q" c" r" u" i tcpf gu" eqpegv tcekqpgu" f g" o gtqu" gvtkcf qu" gp" wlecekqpgu" egtccpcu" c" r" equvc0""

"

Vcn' eqo q"ug" r t g u g p v c " g p " g n " c r <sup>2</sup> p f k e g " E " f g " n " r t g u g p v g " p q o k p c e k p . " g n " k p h q t o g " f g " n " r t k o g t c " t g w p k p " f g n " I t w r q " f g " V t c d c l q " u q d t g " C i t g i c e k q p p u " f g " F g u q x g " e q p h q t o c f q " r q t " E H O E I " E Q R C E Q I Q U R G U E C I E T H O " t g e q o g p f » . " *inter alia* , " g u c d n g e g t " v g o r q t c f c u " t g i k p p c r g u " c t o q p k c f c u " f g " x g f c " r c t c " g u r g e k g u " g u r g e k e c u " s w g " u g " u c d g " u g " e q p i t g i c p " r c t c " f g u q x c t " \* e q o g p | c p f q " e q p " g n ' o g t q " g u t k c f q " { " c i t g i c p f q " q v t c u . " u g i A p " e q t t g u r q p f c + " t g e q r g e v c t " { " e q o r c t v k " f c v q u " d k q n i k e q u " { " e q o g t e k c r g u " r c t c " n u " g u r g e k g u " { " n " i g u k p " { " e q p u g t x c e k p " t g i k p p c n " f g " n u " g u r g e k g u " f g " r g e g u " s w g " u g " e q p i t g i c p " r c t c " f g u q x c t 0 " " "

**Nassau Grouper, *Epinephelus striatus* (Bloch 1792)**

**Biological Report**

**PURPOSE**

Vj ku'tgr qtv'owo o ctk gu'cpf "u{pvj guk gu'dkqmi kecnkphqto cvkqp"eqxgtkpi "P cuucw'i tqwr gt." *Epinephelus striatus*. "vj tqwi j qw'ku'pcwtcnf kutkdwkqp0"K'uggm'vq'r t gugpv'vj g'dgu'cxckrdng" kphqto cvkqp'htqo 'r wdkuj gf "cpf 'wpr wdkuj gf 'uqwtegu."\*gd 0'rkgtcwtg'ugctej gu.'kpvgtxkgy u#0"Vj ku' f qewo gpv'f qgu'pqvtgr t gugpv'c'f gekukap'd{ "P O HU'qp'y j gvj gt'vj ku'czqp'uj qwf "dg'r tqr qugf 'hqt" rkuvki "cu'vj tgcvgpgf "qt"gpf cpi gtgf "wvf gt'vj g'Gpf cpi gtgf "Ur geku'Cev0"

## Table of Contents

<b>INTRODUCTION .....</b>	<b>3</b>
<b>1. NATURAL HISTORY.....</b>	<b>3</b>
<b>1.a. Description of species.....</b>	<b>3</b>
<b>1.b. Taxonomy and distinctive characteristics .....</b>	<b>20</b>
<b>1.c. Range wide distribution .....</b>	<b>22</b>
1.c.i. Historical Distribution .....	22
1.c.ii. Influences on Distribution .....	23
<b>1.d. Biological characteristics .....</b>	<b>23</b>
1.d.i. Age, growth and mortality.....	23
1.d.ii. Ecological Roles.....	27
1.d.iii. Population connectivity/population genetics.....	30
<b>2. THREATS OR STRESSORS .....</b>	<b>31</b>
<b>2.a. Anthropogenic Effects .....</b>	<b>31</b>
<b>2.b. Habitat loss or degradation .....</b>	<b>34</b>
<b>2.c. Climate change implications .....</b>	<b>35</b>
<b>2.d. Limits to recruitment/depensation .....</b>	<b>35</b>
<b>2.e. Disease, parasites, and abnormalities .....</b>	<b>37</b>
<b>2.f. Aquaculture – successes, failures, potential threats .....</b>	<b>37</b>
<b>3. DESCRIPTION OF FISHERIES AND FISHERIES MANAGEMENT AND CONSERVATION .....</b>	<b>38</b>
<b>3.a. Abundance indices and trends over time.....</b>	<b>39</b>
<b>3.b. COUNTRY ACCOUNTS.....</b>	<b>44</b>
ANGUILLA.....	45
BAHAMAS .....	46
BELIZE.....	54
BERMUDA .....	59
BRITISH VIRGIN ISLANDS.....	63
CAYMAN ISLANDS.....	64
COLOMBIA .....	68
CUBA .....	69
DOMINICAN REPUBLIC .....	72
HONDURAS .....	73
JAMAICA .....	75
LESSER ANTILLES, CENTRAL AMERICA, AND SOUTH AMERICA .....	77
MEXICO .....	79
TURKS AND CAICOS ISLANDS.....	83
UNITED STATES (FLORIDA).....	86
UNITED STATES (PUERTO RICO).....	90
UNITED STATES (VIRGIN ISLANDS).....	93
<b>LITERATURE CITED.....</b>	<b>97</b>

"

"

"

"

"



tgrkgh'eqtcrnt'gghu'qt'tqemf'uwduwcv'lp'erget'y cvgtu'Ucf qx { 'cpf 'Gmwpf '3; ; ; + 'cnj qwi j 'y' g { 'ecp' dg'hqwpf 'htqo 'y' g'uj qtgrkpg'v'cdqw'322/352'o 0"Ncti gt'cf wmu'v'gpf 'v'q'qeew { 'f ggr gt.'o qtg' twi qug.'tggh'ctgcu'Ugo o gpu'gv'crf04229c+0'Dqj 'cf wmu'cpf 'lwxgpkgu'y kn'wug'gkj gt'pcwtcrf'qt' ct'k'k'k'crnt'gghu'Uo kj '3; 93.'Dggw'cpf 'J kzqp'3; ; 6.'Eqrkp'gv'crf03; ; 9+0

Cu'c'v'qr 'r tgf cvqt'lp'tggh'gequ{ ugo u.'y' g'P cuucw'i tqwr gt'ugt'xgu'geqmi kecn'hwpev'kpu'y' cv' ctg'w'k'm'd'g'kpi 'emct'k'k'g'f '\*O wo d { 'gv'crf04228+0'Ku'r t'gugpeg'o c'k'p'v'k'p'u'i t'c| gtu'cpf 'i t'c| kpi 'r t'guuwt'g' qp'tggh'cni c'r tqxkf kpi 'cp'ko r qt'v'p'v'd'g'p'g'h'k'v'q'uv'p { 'eqt'cni '\*O wo d { 'gv'crf04228+0'Vj g'r t'gf cvqt { ' P cuucw'i tqwr gt'o c { 'j gr 'iko k'v'y' g'ko r cev'q'h'y' g'k'p'x'c'k'x'g'k'q'p'h'k'j . 'dw'v'y' g'g'x'k'f' g'peg'ku'ht'ht'qo " eqpenwuk'g'cv'y' ku'ko g '\*O wo d { 'gv'crf04233+0'Ku'cdugpeg'j' cu'dggp'ur gew'v'g'f 'v'q'c'h'g'ev'geqmi kecn' t'g'g'c'ug'ht'uo cmgt'r t'gf cvqtu.'k'pen'f' kpi 'uo c'm'i tqwr gtu.'y' k'j 't'guw'nc'p'v'ej' cpi gu'lp'y' g't'q'r'j' k'e" t'g'c'v'k'p'uj' k'r u'lp't'ggh'gequ{ ugo u'Uc'cr'k'p'i u'422: . 'O wo d { 'gv'crf04234+0

Cu'y' k'j 'o qu'v'rti' g'o c't'k'p'g't'ggh'k'k'j' gu.'P cuucw'i tqwr gt'f go q'p'ut'cv'g'c'd'k'r c't'k'g'k'k'g'e { 'erg' y' k'j 'f go g'tu'c'n'l'w'x'g'p'k'g'u'cpf 'cf wmu'dw'r' g'nci' k'e'gi' i' u'cpf 'h'c't'x'c'g'0'T'gr' t'q'f' w'v'k'p'ku'q'p'n' 'h'p'q'y' p'v'q' " qeew'f' w'k'p'i 'c'p'p'w'c'n'c'i' i' t'g'i' c'v'k'p'u.'k'p'y' j' k'ej 'r'c't'i' g'p'w'o d'g't'u'q'h'P cuucw'i tqwr gt.'t'c'p'i' k'p'i 'ht'qo " f'q'l' g'p'u'v'q'v'g'p'u'q'h'y' q'w'c'p'f' u.'eq'm'g'ev'k'x'g'n'f' 'ur' c'y' p '\*Uo' k'j '3; 94.'Q'nu'g'p'cpf 'N'c'R'rc'g'3; 9; . 'Eqrkp'g'v' crf03; ; 9.'H'p'g'3; ; 2.'H'p'g'3; ; 4.'Eqrkp'3; ; 4+0'O' c'p { 'h'k'j 'v't'c'x'g'n'k'p'i 'f'k'uc'p'egu'v'q'c't't'k'x'g'cv' r' t'g'f' l'ev'c'd'ng'r' n'c'eg'u'f' w'k'p'i 'y' g'h'g'y' 'y' g'g'm'u.'ur' t'g'c'f' 'q'x'g't' 'u'g'x'g't'c'n'o' q'p'v'j' u.'g'c'ej' "{ g'c't'y' j' g'p'ur' c'y' p'k'p'i " qeew'tu'cpf 'y' g'p't'g'w't'p'v'q'y' g'k't'j' q'o g't'g'g'h'u'Ucf' qx { 'cpf 'Gmwpf '3; ; ; +0'H'g't'v'k'k'f' c'v'k'p'ku'g'z'v'g't'p'c'f'0' H'g't'v'k'k'f' g'f' 'g'i' i' u'j' c'v'ej' 'c'h'g't'45'v'q'62'j' q'w't'u'f' g'r' g'p'f' k'p'i 'q'p'g'p'x'k'q'p'o' g'p'v'c'n'v'g'o' r' g't'c'w't'g'u'0' 'C'h'g't' j' c'v'ej' k'p'i . 'r' g'nci' k'e' r'c't'x'c'n'f' w't'c'v'k'p'o' c { 't'c'p'i' g'ht'qo '64/92'f' c { 'u'y' k'j 'v't'c'p'u'h'q't'o' c'v'k'p'ht'qo 'r' g'nci' k'e'v'q' " f'go' g'tu'c'n'l'q't'o' 'q'ee'w't'k'p'i 'k'p'ig'u'u'v'y' c'p'q'p'g'y' g'g'm'i'R'q'y' g'm'c'p'f' 'V'w'eng't'3; ; 4.'V'w'eng't'c'p'f' 'Y' q'q'f' y' c't'f' " 3; ; 6+0'P' g'y' n'f' 'u'g'w'r'g'f' 'h'k'j ' \*o' g'c'p'?' '530' 'o' o' 'V'q'v'c'n'N'g'p'i' v'y' '\*VN+. 'u'c'p'f' c't'f' 'f' g'x'k'c'v'k'p' '\*UF' +?' '40' . 'P' ? '53+'p'g'c't' 'G'z'w'o' c' 'E'c { u.'D'c'j' c'o' c'u.'y' g't'g'h'q'w'p'f' 'y' k'j' k'p' 'e'q't'c'n'c'n'w'o' r' u' \*P'or'it'es' 'ur' r' 0' 'e'q'x'g't'g'f' 'd { " o' c'u'g'u'q'h'o' c'e't'q'c'n'i' c'g' \*r' t'k'o' c't'k'k'f' 'y' g't'g'f' 'c'n'i' c' 'L'au're'n'c'i'a' 'ur' r' 00" "

### **Nassau Grouper Juvenile Stages**

**Newly settled juveniles (~2.5 – 5 cm TL).** "H'q'm'y' k'p'i 'u'g'w'r'g'o' g'p'v.'P cuucw'i tqwr gt'l'w'x'g'p'k'g'u'c't'g' t'g'r' q't'v'g'f' 'v'q'k'p'j' c'd'k'o' c'e't'q'c'n'i' c'n'c'n'w'o' r' u.'u'g'c'i' t'c'u'u'd'g'f' u.'c'p'f' 'e'q't'c'n' \*G'i' i' n'g'u'q'p'3; ; 7.'F'c'j' n'i' t'g'p' " 3; ; ; +0'O' qu'v'q'h'y' j' c'v'k'u'p'q'y' p'c'd'q'w'v'y' g'g'c't'r'k'g'u'v'k'k'g' 'u'c'i' g'u'eq'o' g'u'ht'qo' 'c' 'u'g't'k'g'u'q'h' 'u'w'f' k'g'u' eq'p'f' w'ev'g'f' 'ht'qo' '3; ; 9/3; ; 6'p'g'c't' 'N'g'g' 'U'q'c'n'k'p'i 'K'uc'p'f' 'k'p'y' g'G'z'w'o' c' 'E'c { u'k'p'y' g'D'c'j' c'o' c'u'0' 'V'j' g' " u'w'x'g'f' u'c'p'f' 'g'z'r' g't'k'o' g'p'u'k'p'o' c'p'i' t'q'x'g'f'k'p'g'f' 'r'c'i' q'q'p'u'c'p'f' 'v'k'f' c'n'c'et'g'g'm'i' \*3/6' 'o' 'f' g'g'r' +.'u'g'c'i' t'c'u'u'd'g'f' u' " c'p'f' 'u'c'p'f' l'r' c'v'ej' 't'g'g'h'j' c'd'k'c'v'j' g'r' g'f' 'k'f' g'p'v'k'h' { 'y' g'P cuucw'i tqwr gt'v' 'u'g't'k'g'u'q'h' 'q'p'v'q'i' g'p'g'v'k'j' c'd'k'c'v' " e'j' c'p'i' g'u'0' 'U'q'o' g'x'c't'k'c'v'k'p'g'z'k'u'u'k'p'y' g'g'z'c'ev'd'q'f' { 'u'k'f' g'c'v'y' j' k'ej' 'j' c'd'k'c'v'v'j' k'h'u'q'ee'w't' 'd'w'v'uj' k'h'u'c't'g' " eq'o' o' q'p' 'c'e't'q'u'u' 'u'w'f' k'g'u'0' 'O' k'e't'q'j' c'd'k'c'v'q'h'p'g'y' n'f' 'u'g'w'r'g'f' 'P' cuucw'i tqwr gt'y' c'u'f' g'u'et'k'd'g'f' 'c'u'y' k'j' k'p' " e'q't'c'n'c'n'w'o' r' u' \*P'or'it'es' 'ur' r' 0' 'e'q'x'g't'g'f' 'd { 'o' c'u'g'u'q'h'o' c'e't'q'c'n'i' c'g' \*r' t'k'o' c't'k'k'f' 'L'au're'n'c'i'a' 'ur' r' 0: " c'n'j' q'w'i' j' 'q'h'w'p' 'y' g'j' c'd'k'c'v'j' c'u'k'o' r' n'f' 'd'g'g'p' 'e'k'x'g'f' 'c'u' 'L'au're'n'c'i'a'0' 'V'j' g'q'r' g'p' 'r'c'w'k'g'q'h'y' g'c'n'i' c'n' " e'q'x'g't'g'f' 'e'q't'c'n'c'n'w'o' r' u'r' t'q'x'k'f' g'f' 'e'q'x'g't'c'p'f' 'h'c'c'k'k'c'v'g'f' 'y' g'o' q'x'g'o' g'p'v'q'h'k'p'f' k'k'f' w'c'n'u'y' k'j' k'p'y' g' " k'p'v't'u'k'g'u'q'h'y' g' 'e'n'w'o' r' u' \*G'i' i' n'g'u'q'p'3; ; 7+0' 'U'g'x'g't'c'n'p'g'y' n'f' / 'u'g'w'r'g'f' 'P' cuucw'i tqwr gt' \*w' 'v'q' : '+y' g't'g' " h'q'w'p'f' 'e'm'q'g'v'q'i' g'y' g't' 'k'p'p'g'k'i' j' d'q't'k'p'i' 'c'n'i' c'n'c'n'w'o' r' u'0' 'C'd'w'p'f' c'p'eg'q'h' 'r'c'v'g' / r'c't'x'c'n'v'q' 'g'c't'n'f' / l'w'x'g'p'k'g' "

"  
"  
"  
"

P cuucw'i tqwr gt'y cu'uwducpvkcm' j ki j gt'kp"Laurencia"ur r 0j cdkcw'v'j cp'kp'ugci tcuu0"Y kj kp'v'j g" Dcttcvgtt"De{ "o cetqeni cn'lu{ ugo . 'r gtegpv'cni cn'leqxt' y cu'eqtg'grcvf' y kj 'r quv'ugwrgo gpv' i tqwr gt'f'gpukv{ =qj gt'j' cdkcv'ej ctcevgt'kuk'eu'we'j 'cu'cni cn'f'kur'cego gpv'xq'no g.'cpf' 'v'j g' pwo dgtu'qh'j' qrgu.'rgf' i gu.'cpf' 'eqtcn' y gtg'pqv' \*Gi i ngu'qp'3; ; 7+0"Vj g'hwpevk'qpcn't'gr'v'k'q'puj' kr " dgwy'ggp'r' gtegpv'cni cn'leqxt' "cpf' 'r quv'ugwrgo gpv'f' gpukv{ 'y cu'k'p'gct' "cpf' 'r qu'k'k'x'g' \*Gi i ngu'qp' 3; ; 7+0" T gegp'w' /ugw'rgf' 'P cuucw'i tqwr gt'j' cxg'cnu'q' dggp'eqm'ge'v'f' 'ht'qo 'v'k'gh'kuj' . "Malacanthus plumieri.'twddng'o qwpf' u'cv'3: 'o . 'y kj 'cu'o cp{ 'cu'5' 'h'kuj' 'v'qi' g'v'j' gt' \*Eq'rk'p' 'gv'cni'03; ; 9+0"Vj g{ 'j' cxg' dggp' t'gr' q'v'f' 'cu'cuu'q'ek'cv'f' 'y kj 'f'k'uect'f' g'f' 's'w'ggp'eq'pej' . "Strombus gigas.'uj' g'm' 'cpf' 'q'v'j' g'f' gdt'ku' ct'q'wpf' "Thalassia' d'g'f' u' \*E'rc { f' q'p' 'gv'cni'04232. "D0Y' kem'wpf' . "Ect'k'dd'gcp' "O' ct'k'p'g' "T' g'ug'cte'j' "E'gp'v'gt' . " r'gtu'0'eqo' o' 0'v'q' [ 0'U'cf' q'x' { . "P' O' H'U.'3; ; 2+'kp' 'v'j' g'V'w'm' 'cpf' "E'ck'equ' "K'ur'p'f' u' . "c'n'j' q'w'j' 'v'j' g'g'z'cev' 'h'kuj' " uk' gu'q'd'ug't'x'g'f' "ct'g'p'q'v'erg'ct'0"R'qu'v'ugwrgo' gpv'uw't'x'k'cni'kp' "o' cetqeni' cn'j' cdkcw'ku'j' ki j gt' 'v'j' cp'kp' " u'gci' tcuu' d'g'f' u' . "uj' q'y' k'pi' "c' 'r'k'ng'f' "c'f' cr' v'k'x'g' "c'f' x'cp'w'ci' g' 'h'q't' 'v'j' g'f' go' q'p'ut'cv'f' 'j' cdkcv'ug'ge'v'k'p' " \*F' c'j' ni' t'gp' "cpf' "Gi i ngu'qp'4222+0" "

**Early juveniles (~4.5 – 15cm TL).** "Uo' cni'lw'x'g'p'k'rg'P' cuucw'i tqwr gt'ct'g'eqo' o' q'p'kp'uj' c'm'q'y' " u'gci' tcuu' d'g'f' u' . "o' cetqeni' c'g' . "cpf' "ct'q'wpf' "em'no' r' u'q'h' "Porites"ur' r' 0'eq't'cni'cu'v'j' g{ 'd'gi' kp'v'q'uj' k'w'ht'qo' " ugwrgo' gpv'j' cdkcw'qt' "o' k'et'q'j' cdkcw' \*T' c'p'f' c'm'3; ; 5. "Gi i ngu'qp'3; ; 7+0"Vj g't'gr'v'k'q'puj' kr' "dgwy'ggp' " Laurencia' c'p'f' 'p'gy' "ugwrg't' "c'p'f' 'g'c't'n'f' "lw'x'g'p'k'rg'f' g'p'uk'k'g'u' y' cu' 'o' c'k'p'v'k'p'g'f' "w'p'k'i'c'd'q'w'7" "o' q'p'v'j' u'c'h'ng't' " ugwrgo' gpv'0' "Ch'ng't' 'v'j' c'v'v'ko' g' . "o' q't'v'erk'f' "cu' y' g'm' 'cu' 'o' q'x'go' gpv'v'q' 'r' c'v'ej' "t'g'gh'j' cdkcv't'g'h'ng'e'v'ej' c'p'i' gu' " kp'f' k'ut'k'd'w'k'q'p' "c'p'f' "c'd'w'p'f' c'p'eg' \*Gi i ngu'qp'3; ; 7+0" D'c'p'f' "t'c'p'ug'ew' 'r' g't'h'q'to' g'f' "p'g'c't' "N'gg' "U'q'en'k'pi' " K'ur'p'f' . "D'c'j' c'o' cu' "6/7" "o' q'p'v'j' u'c'h'ng't' 'v'j' g'ugwrgo' gpv'r' g't'k'q'f' \*L'w'p'g'3; ; 3/; 5+ 'h'q'w'p'f' 'v'j' c'v'g'c't'n'f' " lw'x'g'p'k'rg'u'f' go' q'p'ut'cv'f' "c'uw'd'v'g'ej' c'p'i' g'kp' "o' k'et'q'j' cdkcv'=: ' 'y' g't'g' "u'q'rk'ct' { 'y' k'j' kp'qt' "c'f' l'ce'gp'v'v'q' " cni' cn'eq'x'g't'g'f' "eq't'cni'emo' r' u'0" T'g'gh'j' cdkcw' . "k'p'en'w'f' k'pi' "u'q'rw'k'q'p'j' q'rgu' "c'p'f' "rg'f' i' gu' . "v'q'q'm'q'p' " eqo' r' c't'c'v'k'x'g'n'f' "i' t'g'c'v'gt' "k'o' r' q't'v'c'p'eg' "cu'j' cdkcw' "h'q't' "g'c't'n'f' "lw'x'g'p'k'rg'u'cu'v'j' g{ 'i' t'gy' 0" T'g'r' g'c'v'f' "o' q'p'v'j' n'f' " eg'p'u'w'gu'q'h'c' "r' t'g'u'wo' g'f' "eq'j' q't'v'k'p'f' k'ec'v'f' 'v'j' c'v'lw'x'g'p'k'rg'f' g'p'uk'v'f' "f' g'et'g'c'ug'f' "uj' c't'r' n'f' "c'h'ng't' "ugwrgo' gpv' " w'p'k'i' 'h'kuj' "go' g't'i' g'f' "ht'qo' "c'ni' cn'j' cdkcv'c'v'ug'x'g't'cni' b' q'p'v'j' u'q'h'c'i' g' . "c'p'f' 'v'j' g't'g'c'h'ng't' "t'go' c'k'p'g'f' "t'gr'v'k'x'g'n'f' " eq'p'ux'c'p'v' \*F' c'j' ni' t'gp'3; ; : +0" Q'p'uj' c'm'q'y' "eq'p'ut'w'ew'g'f' "d'm'em'it'g'gh'u'kp'v'j' g' "X'k'i' kp' "K'ur'p'f' u' "52/ : 2" "o' o' " VN'p'gy' n'f' "ugwrg'f' l'g'c't'n'f' "lw'x'g'p'k'rg'u'q'ee'w' k'g'f' "u'o' c'm'i'ug'r' c't'c'v'g' "d'w't't'q'y' u' d'g'p'g'c'v'j' 'v'j' g't'g'gh'j' y' k'rg' "r'c't'i' g't' " lw'x'g'p'k'rg'u'q'ee'w' k'g'f' 'j' q'rgu'kp'v'j' g't'g'gh'u' \*D'gg'w' "c'p'f' "J' k'z'q'p'3; ; 6+0' "

F'w'k'p'i' 'v'j' g' 'u'c'o' r' n'k'p'i' 'r' g't'k'q'f' "kp'3; ; 5" c't'q'wp'f' "N'gg' "U'q'en'k'pi' "K'ur'p'f' . "o' g'c'p' "uk' g'k'p'et'g'c'ug'f' " ht'qo' "530"v'q' : 70" "o' o' "VN" \*z' "32" "o' o' l'o' q'p'v'j' +0" I' t'q'y' v'j' "t'c'v'gu' y' g't'g' "eq'p'uk'v'g'p'v' y' k'j' "v'j' q'ug' "t'gr' q't'v'f' " h'q't' "g'c't'n'f' "lw'x'g'p'k'rg'u'k'p'j' c'd'k'k'p'i' "c't'v'h'k'ek'c'n'r' c'v'ej' "t'g'gh'u'kp'v'j' g' "W'U'X'K'00" \*D'gg'w' "c'p'f' "J' k'z'q'p'3; ; 6+0" " J' cdkcv'w'uci' g'q'h'p'gy' n'f' "ugwrg'f' "lw'x'g'p'k'rg'u' t'gr' q't'v'f' n'f' "o' c'z'k'o' k' gu'uw't'x'k'cni' y' j' k'rg'j' cdkcv'uj' k'm'u' "h'q't' " g'c't'n'f' "lw'x'g'p'k'rg'u' "c'p'f' "lw'x'g'p'k'rg'u' "h'c'ek'k'c'v'g' "k'p'et'g'c'ug'f' "i' t'q'y' v'j' "t'c'v'gu' \*F' c'j' ni' t'gp' "c'p'f' "Gi i ngu'qp'4222+0' "

**Juveniles (~15 – 35; 30 – 50cm TL).** "Lw'x'g'p'k'rg'P' cuucw'i tqwr gt'ct'g't'gr'v'k'x'g'n'f' "u'q'rk'ct' { "c'p'f' . " y' j' k'rg'v'j' g{ 't'go' c'k'p' "ur' g'ek'h'le' "c't'g'c'u' "h'q't' "g'z'v'p'f' g'f' "r' g't'k'q'f' u' \*D'c't'f' c'ej' "3; 7: +. "v'j' g{ "o' c' { "g'z'j' k'd'k' " f' k'uk'p'ev'q'p'v'q'i' g'p'g'v'k'e' "uj' k'm'u'kp'j' cdkcv'c'p'f' "f' k'g'v'cu'uk' gu'k'p'et'g'c'ug'0" Lw'x'g'p'k'rg'u'kp'v'j' g' "D'c'j' c'o' cu'uj' k'hw'g'f' " ht'qo' "o' cetqeni' cn'j' cdkcw'v'q' "p'ew't'cni' "c'p'f' "c't'v'h'k'ek'c'n'r' c'v'ej' "t'g'gh'u' "q'x'g't' "c'5/ "o' q'p'v'j' "r' g't'k'q'f' "c'v'342/372" "

"  
"  
"  
"



**Nassau Grouper Adult Stage**

*Size and age at maturity.* "O crg"cpf 'hgo crg"P cuucw'i tqwr gt "v{r kccm{ 'o cwtg'dgy ggp'622" cpf '672"o o "UN"\*662"cpf '726"o o "VN+:'y kj 'o quv'kpf kxf wcu'cwckpki 'ugz wcu'lo cwtk{ 'd{ 'cdqw' 722"o o "UN"\*779"o o "VN+cpf 'cdqw'6/7" { gctu'qh'ci g"\*Vcdrg'3+ 'cmj qwi j 'y j g'uo cmguv'o cwtg'hkuj " tgeqtf gf 'kp'Ewdc'y cu'c"o crg'kp'y j g'582/5; 2o o "VN'uk{ g'emuu"\*Erctq'gv'cr03; ; 2+0"Qnugp"cpf " NcRrreg"\*3; 9; +tgr qtvgf 'C; 7"\*y j g'ci g'cv'y j kej "; 7' "qh'y j g'cu{ o r vqke'hgpi y j 'ku'tgej gf +ku'370 " { gctu0"kp'uco r rpi 'hkuj gt { 'ecvej gu'cv'O cj cj wen'uqwj gtp'S wkpvcpc'Tqq. 'O gzleq. 'f wtkpi '3; ; 3/ 3; ; 5"cpf '3; ; 9. 'Ci wkrt/Rgtgtc"\*4226+tgr qtvgf 'y j g'uo cmguv'o crg'cu'5; 2"o o "VN'cpf 'y j g'uo cmguv' hgo crg'cu'682"o o "VN'O quv'kpf kxf wcu'ecwi j v'ltqo "c'WUOX00ur cy plpi 'ci i tgi cvkqp'y gtg" dgy ggp'cdqw'722"cpf '822"o o "VN"\*Qnugp"cpf 'NcRrreg'3; 9; +0"P go gy 'gv'cr0"\*4228+hqwpf 'y j cv' cf wv'P cuucw'i tqwr gt "cv'c'f khtgtpv'ur cy plpi 'ci i tgi cvkqp'ukg"\*1 tco o cpkn'Dcpm'kp'y j g'WUOX00 teci gf 'dgy ggp'6: 2"cpf ": 22"o o 'y kj 'cxgtci g'vqcn'hgpi y j 'hqt"o crgu"\*825"o o . 'p'? '3: +cpf " hgo crgu"\*7; 3"o o . 'p'? '66+dgkpi 'uko kct0'Htqo "qvqkj 'ci lpi 'y qtm'y j g'o lpo wo 'ci g'cv'ugz wcu' o cwtk{ 'ku'dgy ggp'6"cpf ": " { gctu"\*Dwij 'gv'cr03; ; 8. '4228+ " y kj 'o quv'hkuj 'ur cy plpi 'd{ 'ci g' 9- "{ gctu"\*Dwij 'gv'cr0'4228+0" P cuucw'i tqwr gt 'tckugf 'htqo 'y j g' gi i 'kp'ecr vxxk{ 'o cwtg' 'cv'49/ 4: 'o qp'y u"\*622/672"o o " UN'662/726"o o "VN+\*"Vwengt" cpf 'Y qqf y ctf '3; ; 6+0"Uk{ g." tc'y gt 'y j cp'ci g. 'o c{ 'dg'y j g' o clqt 'f gy'to kpcpv'qh'ugz wcu' o cwtcvkqp"\*Ucf qx { "cpf 'Gmw'pf " 3; ; ; +0"

**Table 1. Summary of Age and Length Parameters for Nassau grouper, *Epinephelus striatus* (from Table 3, Sadovy and Eklund 1999; "Bush et al., in press" refers to Bush et al. 2006)**

	<i>Epinephelus striatus</i>
Age and length at maturity	5 yrs, 580 mm SL (Virgin Islands) (Olsen and LaPlace, 1979)  420-450 mm SL females 400-450 mm SL males\4+ yrs (Cayman Islands) (Colin et al., 1987; Bush et al., in press) 500 mm TL (minimum size ripe males) (Cayman Islands) (Tucker et al., 1993) 425 mm SL females; 402 mm SL males, immatures are 3-6 yrs (otolith growth zones not validated) (Bahamas) (Sadovy and Colin, 1995) 483 mm TL (North Carolina-Florida) (SAFMC, text footnote 24) 480 mm TL (Jamaica) (Thompson and Munro, 1978)
Age and length at first capture	< 300 mm TL & 4-5 yrs (Virgin Islands) (Olsen and LaPlace, 1979; CFMC, text footnote 26) 6-7 yrs (Cayman Islands) (Bush et al., in press) 275-625 mm TL (mean = 570) (Jamaica) (Thompson and Munro, 1978) 300-500 mm TL depending on size limits (North Carolina-Florida) (SAFMC, text footnote 46) 450 mm TL (South Florida) (Bohnsack, 1990)
Maximum age and length	1200 mm TL (CFMC, text footnote 26) 9 yrs, 910-960 mm SL (Olsen and LaPlace, 1979) (Virgin Islands) 17 yrs, 710 mm TL, 6700 g (Cuba) (Claro et al., 1990) 755 mm SL (Bermuda) (Bardach et al., 1958) 840 mm TL (Jamaica) (Thompson and Munro, 1978) 640 mm TL (Netherlands Antilles) (Nagelkerken, 1981) 29 yrs, 850mm FL (Cayman Islands) (Bush et al., in press)

o cwtk{ 'ku'dgy ggp'6"cpf ": " { gctu"\*Dwij 'gv'cr03; ; 8. '4228+ " y kj 'o quv'hkuj 'ur cy plpi 'd{ 'ci g' 9- "{ gctu"\*Dwij 'gv'cr0'4228+0" P cuucw'i tqwr gt 'tckugf 'htqo 'y j g' gi i 'kp'ecr vxxk{ 'o cwtg' 'cv'49/ 4: 'o qp'y u"\*622/672"o o " UN'662/726"o o "VN+\*"Vwengt" cpf 'Y qqf y ctf '3; ; 6+0"Uk{ g." tc'y gt 'y j cp'ci g. 'o c{ 'dg'y j g' o clqt 'f gy'to kpcpv'qh'ugz wcu' o cwtcvkqp"\*Ucf qx { "cpf 'Gmw'pf " 3; ; ; +0"

*Habitat and Home*

Range0""Cnj qwi j 'y j gtg'ecp'dg" qxgtrr 'dgy ggp'lwxgpkng'cpf " cf wv'j cdkcw'y j g'ku'pqto cm{ " c'r qukxg'eqtt grv'kqp'dgy ggp" uk{ g'cpf 'f gr y j 0"P cuucw'i tqwr gt " ctg'f kwtpcn'qt "etgr wuewrt 'kp" y j gk' o qxgo gpw"\*Eqmgwg"cpf " Vcrdq'3; 94+cpf 'f q'pqv'wuwcm{ " o qxg'ht'htqo 'eqxgt"\*Uctem' cpf 'F cxku'3; 88+0"Vj tgg" uqplecm{ 'vci i gf 'hkuj 'y gtg'o quv' cexkg'kp'y j g'j qwtu'r tkqt'vq'cpf "

"  
"  
"  
"

hqmjy kpi "uwptkug"cpf "uwpuv"ECTvgt "gv'cr03; ; 6+0"Vy q"qh'vj g'hkuj "o qxgf "tcpf qo n{ "y kj kp"e"382" o "z": 2'o "tgevcp i ng"fwtkpi "vj g'f c{ .t'gwtpkpi "kp"vj g'gxgpkpi "v'y j gtg'vj g{ "j cf "kpklcvgf "f ckn{ "cevxxkku"ECTvgt "gv'cr03; ; 6+0"Uwrxcp"cpf "f g"l ctkpg/Y lej cvksun{ "3; ; 6+"guko cvgf "vj cv kpf kxf wcu"o qxgf "cv'ngcu/622"o If c{ "cpf "42"o "qt"o qtg'htqo "vj gk"j qo g'tgghu"O gcp"j qo g/ t'cpi g'ctgc"y cu"ecrewcvgf "cv'3: .527o <sup>4u</sup>- 1/7.: 28\*"UG+Dqrf gp"4223+0"P cuucw'i tqwr gt"j cf "rcti gt" j qo g'tcpi gu'cv'nguu'utwewtcm{ "eqo r ngz "t'gghu"cpf "t'guqwtg"cxckcdkxv{ "j cdkcv"cpf "r tg{ "+" kphwpgegu"j qo g'tcpi g'uk' g'o qtg'vj cp"dqf { "uk' g\*"Dqrf gp"4223+0"Dqrf gp"4223+"kpxguk' cvgf "f kgn{ "cevxxk{ "r cwgtpu"xk"eqpvkpwqu"ceqwuk"vgrgo gvt { "cpf "hqwpf "P cuucw'i tqwr gtu'ctg"o qtg"cevxxg" f kxtpcm{ "cpf "nguu"cevxxg"pqewt'pcm{ "y kj "cevxxk{ "r gcm'cv'3222"cpf "4222"j qwtu"O"p go gj "cpf " eqy qtngtu\*"Wpkxgtuk{ "qh'vj g"Xkti kp"Kucpf u."o cpwuetkr v'kp"r tgr +j cxg'hqwpf "c"uki pkl'ecpv" r qukkxg't'grv'kpuj kr "dgy ggp"dqf { "uk' g'cpf "j qo g'tcpi g."hqt'hkuj "uci i gf "kp"Nco guj wt"Dc{ ."U0' Lqj p."y kj "o gcp"o kpo wo "eqpxgz"r qn{ i qp\*"O ER+xctk'v'kpu'htqo " : ; 07/ ; ; 350 "o <sup>4</sup>O"t'gegpv" uwf kgu'kp"e"o ctkpg't'gugt'xg'kp"Ewdc"uwi i gu'vj cv't'grv'kxg'f gpuk'ku"o c{ "eqpvt'qn'o qxgo gpw." ej cpi gu'kp"mqecv'kp."cpf ."r quukdn{ ."j qo g'tcpi g'uk' g\*"Co cti »u"gv'cr04232+0"

*Depth ranges*0"Cf wv/P cuucw'i tqwr gt"ctg'i gpgtcm{ "cuuqekcvgf "y kj "uj cmjy "t'ggh"j cdkcw" vq"fr gj u"qh'322"o 0"t'gr qt'u'htqo "hkuj kpi "cevxxkku"kp"vj g'Nggy ctf "Kucpf u"uj qy "vj cv'cnj qwi j " P cuucw'i tqwr gt"y cu'hkuj gf "vq"352"o ."vj g'i tgc'v'v'v'tcr "ecvej gu'y gtg'htqo "74/82"o "Dtqy pgn'cpf " T'clpg{ "3; 93+0"kp"Xgpgl wgr. "P cuucw'i tqwr gt"y gtg'ekgf "cu'eqo o qp"vq"62"o "kp"vj g'Ctej kr gnci q" Nqu"t'qs wgu."dw't'ctg'kp"pqtvj gcuvtp"ku'cpf u"Et'gtxki »p"3; 88+0"t'gegpv"uci i kpi "uwf kgu'kp"Dgrk' g" j cxg'uj qy p"vj cv'kpf kxf wcu't'gi wrctn{ "f guegpf "vq"fr gj u"qh'cv'ngcu/477o "Ucct"gv'cr04229+0"Vj g" uj kv'kp"fr gj "hqmjy gf "ur cy pl'pi "cpf "y cu'u'pej tqpwu"vq"cp"cxgtci g"qh'930 "o "020"UG+ "y kj " c"o czko wo "fr gj "qh'477"o ."cpf "r gtukwgf "cdqw'5"o qp'v' u."vj tqwi j qw'vj g'y k'vgt "ur cy pl'pi " ugcup'kp"Dgrk' g0"Ucct"cpf "eq/cwj qtu"4229+ "j { r qv' guk' gf "vj cv'vj gug'f ggr "o ki t'v'kpu"o ki j " h'ek'k'cvgf "r j { uk'qmi k'ecnt'geq'xgt { "cpf lqt"vj cv'ur cy pl'pi "o ki j v'eqpv'kpwg"cv'fr gj ."dw'vj g't'wg" r wtr qug't'gs vkt'gu'hw'w'g't'gugctej 0'

*Sizes and size distribution*0"O gcp"o crg"cpf "hgo crg"uk' gu'ctg'uko kct"y kj kp"e"i kxgp"ctgc." qt"cv'c"ur gek'le"ci i t'gi cv'k'p'uk'g."y kj "uqo g'k'p'f'k'ec'v'k'p'vj cv'uk' gu'qh'dqj "ugz gu'f gen'k'p'g'kp"ctgcu" y kj kp"e"ur gek'le"t'gi k'p"y kj "j ki j gt"gzr m'k'v'k'p'p"t'g'x'k'g' gf "kp"Ucf qx { "cpf "Gm'w'p'f"3; ; ; +0"Hqt" gzco r ng."kp"Dgrk' g."vj g'cxgtci g'ngpi vj "qh'dqj "ugz gu'y cu'322"o o "uo cmgt"kp"ecvej gu'htqo " g'zr m'k'w'gf "eqo r ctgf "vq"v'p'gzr m'k'w'gf "ci i t'gi cv'k'p'u"ECTvgt "gv'cr03; ; 6+0"kp' kxf wcu"p cuucw'i tqwr gt" ecp'h'x'g'hqt"cmo qu'v'vj tgg'f gecf gu."dw'o qu'v'hkuj "eqn'ge'v'gf "ctg"u'w'du'cv'k'cm{ "uo cmgt"cpf " r t'guwo cdn{ "{ qwpi gt0"Dwuj "gv'cr04228+ "t'gr qt'v'gf "vj cv'vj g'q'rf gu'v'P cuucw'i tqwr gt"kp"vj gk"u'wf { "kp" vj g'Ec{o cp"Kucpf u"y cu"4; "{ gctu."dcugf "qp"cp"ci gkpi "u'wf { "vuk'pi "uci k'w'cn'q'v'rk'j u0'

*Reproductive mode.*"Vj g'P cuucw'i tqwr gt"y cu'qtki k'pcm{ "eqpuk' gtgf "vq"dg"e"o q'p'c'p'f' t'le" r tqv'j { p'qwu"j gto crj tqf k'g."h'ng"o qu'v'vj gt"i tqwr gtu."y kj "cm'o crgu'f g'k'k'k'pi "htqo "vj g'ugz" ej cpi g'qh'cf wv'hgo crgu"Uo kj "3; 93."Erctq"gv'cr03; ; 2."Ectvgt"gv'cr03; ; 6+0"Gxk'f g'peg"qh'vj g" ej cpi g'htqo "cf wv'hgo crg"vq"cf wv'o crg"kp"vj g'P cuucw'i tqwr gt"K0'hkuj "w'p'f'gti q'k'pi "ugz wcu" v'c'p'uh'qto cv'k'p"y j gtgd { "vj g'i q'p'c'f'u'uj qy "f'gi g'p'g't'c'v'k'p"qh'o cwt'g"v'ku'w'g"qh'q'p'g'ugz"cpf " r tqv'j g'k'v'k'p"qh'ht'gr tqf w'v'k'x'g"v'ku'w'g"qh'vj g"q'v'j gt+"j qy gxgt."y cu'y gcm"Ucf qx { "cpf "Uj cr k'q"3; ; 9."

:"

"  
"  
"  
"

Uj cr ktq"3; : 9-0"Qvj gt'ej ctcevgtkueu'y gtg'hqwpf "vq'dg'kpeqpukv'p'v'y kj "c'f'kci pquku'qh"  
 o qpcpf tke'r tqvqi {p{ 'uwej "cu'vj g'utqpi "o crg'hgo crg'uk' g'qxgtcr. "vj g'r tgugpeg'qh'o crgu'vj cv'  
 f gxgnr 'f ktgev' "htqo "vj g'lwxgpkng'r j cug."cpf "vj g'o cv'pi "u{wgo "Eqrkp"3; ; 4."Ucf qx {"cpf "Eqrkp"  
 3; ; 7-0"

P cuucw'i tqwr gt'r cuu'vj tqwi j "c'lwxgpkng'dkugz wcn'r j cug"\*vj g'i qpcf u'eqpukv'qh'dqj "  
 ko o cwtg'ur gto cvqi gple'cpf "ko o cwtg'qxctkcp "kuuwg+\*Vcdng"4+."cpf "o cwtg'f ktgev' "cu'o crg'qt "  
 hgo crg"\*Ucf qx {"cpf "Eqrkp"3; ; 7-0"Cnj qwi j "vj g'P cuucw'i tqwr gt'ku'ecr cdng'qh'ej cpi kpi "ugz "  
 hmqy kpi "j qto qpg'kplgevkp/qpg'P cuucw'i tqwr gt'tgr tqf wegf "cu'c'hgo crg'cpf "uwdugs wgpv' "cu'c"  
 o crg'er r tqzko cvgn' "8"o qpj u'rcvt. "hmqy kpi "cp'NJ TJ /c"ko r rcpv'kp'ecr v'k'k' "Y OY cvpcdg"  
 cpf "Y OJ gcf."Ectkddgcp'O ctkpg'Tgugctej "Egpygt."r gtu0eqo o 0'vq' [ 0'Ucf qx {"P O HU."3; ; 4."  
 Y cvpcdg'gv'cr03; ; 7d+/pcwten'ugz "ej cpi g'j cu'pqv'dggp"eqphko gf 0'Vj g'emug'ch'k'k' "qh'vj ku"  
 ur gekgu'y kj "qj gt'j gto cr j tqf kke'ugttcpkf u'ceeqpw'ht'vj g'i qpcf "utwewt'g'qh'vj ku'ur gekgu'cpf "  
 cnj qwi j "k'o c' "t'g'v'c' "ecr cek' "hqt'pcwten'ugz "ej cpi g'cxckrdng'g'x'k' g'peg'k'p'k'ecv'gu'vj cv'vj ku'ku"  
 pqv'v' r kcn'cpf "vj cv'vj g'P cuucw'i tqwr gt'ku'r tko ctk' "i qpqej qtk'k' "ugr ctcv'ugz gu+\*Ucf qx {"cpf "  
 Eqrkp"3; ; 7-0"

**Table 2. Gonadal maturity according to size for Nassau grouper (from Sadovy and Eklund 1999)**

Stages of gonadal maturation for 230 *Epinephelus striatus* collected in the Bahamas between May 1988 and October 1990 (from Sadovy and Colin, 1995- Fig. 3). Bisexual fish are those in which the gonads contain both ovarian and testicular tissue and include both immature (both male and female tissue immature) and mature (in parentheses) bisexuals.

Size class (mm SL)	Female		Male		
	Bisexual	Immature	Mature	Immature	
151-200	1	1			
201-250	2				
251-300	8	3			1
301-350	11	3			1
351-400	15	2			
401-450	4	1	2	1	1
451-500	9		10		23
501-550	4(1)		36		15
551-600	(3)		33		9
601-650			13		4
651-700			5		6
701-750			1		1
Total	58	10	100	1	61

hgcw'gu."uwej "cu'vj g'gpf u'qh'kurcpf u'qt'r tq'gevkpu'r tqo qpvtkgu'qh'vj g'tggh'ugcy ctf "htqo "vj g'  
 i gpgtcnt'ggh'eqpvqwt "Eqrkp"gv'cr03; : 9."J g{o cp'cpf "Mlghxg"422: -0"Vq'mqecv'c'ukg."i tqwr gt "  
 eqwf "uy ko "wr /'qt'f'qy p/ewttgpv'cmqi "vj g'uj gnh'dt'gcn'v'q'tgcej "vj g'o quv'ugcy ctf "wr /ewttgpv'  
 gzv'pukp'qh'vj g'tggh'y j gtg'ci i tgi cv'k'p'ukgu'ctg'i gpgtcn' "mqecv' "Ectvgt"3; : 8."Eqrkp"gv'cr0

Spawning migrations. "Vj g'P cuucw'i tqwr gt "  
 ci i tgi cv'gu'k'p'rci g'pwo dgtu'v'ur cy p'gcej "{ gct=" "  
 vj g'rci gu'ci i tgi cv'k'p'uwf k'f "j cf "cp'guko cv'f "  
 52.222/322.222"ur cy p'kpi "h'uj "Uo kj "3; 94+k'p "  
 Dko k'p'Dcj co cu'0'Cu'ht'cu'ku'hpqy p."cm "  
 tgr tqf we'k'g'cev'k'k' "qeewtu'k'p'vj gug "  
 ci i tgi cv'k'p'u'vj cv'htqo "eqpukv'p'v' "cv'ur gekke "  
 ukgu'oi tqwr gt "j qrgu'0'cpf "ko gu'0'Ci i tgi cv'k'p'u "  
 j cxg'eqpukv'g'qh'j w'p'f tgf u."vj qwucpf u."qt. "  
 j kv'qt'k'cm' ".v'pu'qh'vj qwucpf u'qh'k'p'k'k' w'cn'cpf "  
 j cxg'r g'ukv'g' "cv'hpqy p'rqec'k'p'u'ht'r g'k'q'f u'qh "  
 ; 2{" gct'qt"o qtg'Uo kj "3; 94."Qnug'p'cpf "  
 NrRreg"3; 9; . "Eqrkp"gv'cr03; : 9."H'k'p"3; ; 2. "  
 3; ; 4."Eqrkp"3; ; 4."Ectvgt'gv'cr03; ; 6."Ucf qx {" "  
 3; ; 9."T0Ertq."Ncdqtcv'qt {"qh'H'uj "Geqmi {"o "  
 Ewdc."r gtu'0'eqo o 0'vq' [ 0'Ucf qx {"P O HU."3; ; 3-0 "  
 K'ku'p'qv'hpqy p'j qy "P cuucw'i tqwr gt "  
 ugr'ev'cpf "mqecv'ci i tgi cv'k'p'ukgu'qt'y j {"vj g {" "  
 ci i tgi cv'g'v'ur cy p'0'Ci i tgi cv'k'p'u'ctg'v' r k'cm' "  
 mqecv' "pgct'uki p'k'ecp'v'i gqo qtr j qmi k'cn' "

;"

"  
 "  
 "  
 "

3; : 9+0"Vj g'vko lpi "cpf "u{pej tqpk cvkqp"qh'ur cy plpi "o c{"dg"f gvgto kpgf "d{"vj g'pgeguukv{"hqt" y kf gn{ "f kur gtugf "cf wnu"v"eqqtf kpcvg"vj gkt "tgr tqf wevkxg"cevkkkku."o c{"hcekrkcvg"gi i "f kur gtucn" o c{"o kpo k g"gi i "f kur gtucn"qt"o kpo k g'r tgf cvkqp"qp"cf wnu"qt"gi i u"Eqrk"3; ; 4+0

Rtkqt"vq"ur cy plpi . "huj "o ki tcvg"vqy ctf "ci i tgi cvkqp"ukgu"lp"i tqwr u'pwo dgtkpi "dgwy ggp"47" cpf "722."o qxlpi "r ctcnri"v"vj g"eqcu"vt"cmppi "vj g'uj grh'gf i g"Eqrk"3; ; 4."Ectvgt"gv'crl03; ; 6." Ci wkrt/Rgtgtc"cpf "Ci wkrt/F ctkc"3; ; 8+0"Qxgt"7"{"gctu"qh"qdugtxcvkqpu"\*4224/4228+"lp"vj g" Ec{o cp"Krcpf u."o ki tcvkpi "P cuucw'i tqwr gt'y gtg'tgr qtvgf "cv'vj g'uj grh'gf i g."v{r kcmf "cv'f gr vj u" tcpu lpi "htqo "42"v"55"o 0"Ceeqtf lpi "v"Y j c{"rgp"gv'crl0\*4229+<öO ki tcvkpi "i tqwr gt'y gtg'o cknf " lp"vj g'f ctnleiqnt"r j cug."cnj qwi j "vj g'y j kg'dgmf "r j cug'y cu'pqv'wpeqo o qp0"Rgcni'pwo dgtu"qh" o ki tcvkpi "i tqwr gt'u'y gtg'qdugt'xgf "4"v"5"fc{"u'chgt"hwmb' qqp'y kj "enwvgtu"qh'wr "v"322"i tqwr gt'u" v'cxgrkpi "vqi gj gt"cmppi "vj g'y cni"vqy ctf u"vj g'ci i tgi cvkqp"ukg0"P cuucw'i tqwr gt"o ki tcvkpi "v"vj g" I tco o cplni'Dcpmlur cy plpi "ukg"qh'i"U0Vj qo cu."WUOX00"o qxgf "cmppi "c"rkpget"tgg"522/722"o " kpij qtg'tcvj gt"vj cp"uy ko o lpi "cmppi "vj g'cewcn'uj grh'gf i g"P go gj "gv'crl0422; +0

"öCorridas de desoveö"ur cy plpi "twpu+.y j kej "tghgtu"dqj "v"vj g"o ki tcvkqp"qh'huj "vqy ctf " c"ur cy plpi "ukg"cpf "v"vj g"ci i tgi cvkqp"kuh'gh"y gtg'htu'f guetkdgf "lp"P cuucw'i tqwr gt"htqo "Ewdc" lp"3: : 6"d{"Xkrctq"Fc| . "cpf "rcvt"d{"T wkctv/O cpf c{"cpf "Lxctgl /Hgtpcpf gl "3; ; 88+0"Cm'vj tgg" y qtngtu"pqvgf "vj cv'huj gt'u'tgr qtvgf "ur cy plpi "twpu"qeewtkpi "o cknf "dgwy ggp"P qxgo dgt"cpf " Hgdtwet {"cpf "cv'f khgtgpv'o qqp"r j cugu0"Kku'pqv'hpqy p"y j gj gt"corridas"ctg"gzenukxgnf " cuuqekcvf "y kj "ur cy plpi "qt"qeew"cv'vqj gt"vko gu."wpcuuqekcvf "y kj "tgr tqf wevkxg"cevkkk{0

"F wtkpi "vj g'ugxgten'o qpj "ur cy plpi "ugcuqp"geej "gct."P cuucw'i tqwr gt"o qxg'htqo "vj gkt" tgul'gpv'cnj cdkew"v"ur cy plpi "ci i tgi cvkqp"j cdkew0"Ur cy pgtu'er r gct"v"uj qy "uqo g'ukg" hkf grk{"v"vj g'uco g"ci i tgi cvkqp"ukgu"{"gct"chgt"{"gct0"O qxgo gpv'cy c{"htqo "tgul'gpv'tgghu"qeewtu" cu"ur cy plpi "vko g'er r tqcej gu"cpf "f kucpegu"v'cxgrgf "xct{"f gr gpf lpi "qp"f kucpeg"v"ci i tgi cvkqp" ukgo"F kucpeg"v'cxgrgf "ku"j ki j n{"xctkcdng0"Uqo g'huj "o qxg"qpnf "c'hgy "hkrqo gvgtu."dw'uqo g" kpf kxf wcu"ctg"npqy p"v"v'cxgrn'wr "v"ugxgtenj wpf tgf "hkrqo gvgtu"v"tgr tqf weg0Qdugtxcvkqpu"qh" o ki tcvkpi "i tqwr u'qh'huj . "qp"qt"dghtg"vj g'hwmb' qqp"qh'ur cy plpi . "kpf kcvg"vj cv'cv'rgcu"uqo g'huj " v'cxgrn'v"ci i tgi cvkqp"ukgu"lp"i tqwr u'tcpu lpi "htqo "c'hgy "huj "wr "v"cdqw"722"kpf kxf wcu"Eqrk" 3; ; 4+0"Ugxgtenf q| gp'huj "y gtg'qdugt'xgf "r cuukpi "unqy n{"cmppi "vj g"52/62"o "uj grh'dtgcni'eqpv'w" cv'ugxgten'ncrkkku"cmppi "c'tggh'lp"Dgrk g"lp"rcvg"Qevqdg"cpf "gctn{"P qxgo dgt"Ectvgt gv'crl0 3; ; 6+lp"qvj gt"y qtf u."c"o qpj "qt"vy q"dghtg"ur cy plpi "y cu'hkngn{0"Kp"J qpf wcu."i tqwr gtu" pqto cmf "mqecvf "6: "no "htqo "cp"ci i tgi cvkqp"ukg"fkucr r gctgf "htqo "tgul'gpv'tgghu"cv'ur cy plpi " vko g"Hpg"3; ; 4+0"Qpg"vci i gf "huj "lp"vj g'Dcj co cu'eqxgtgf "c"fkucpeg"qh'cv'rgcu"v"332"no "lp"vy q" o qpj u"v"cp"ci i tgi cvkqp"ukg"Eqrk"3; ; 4+0"Cpqj gt'huj . "vci i gf "qp"cp"ci i tgi cvkqp"ukg"lp" Dgrk g."y cu'tgecr wtgf "4"{"gctu"rcvt"462"no "pqtvj "qh'vj g'vci i lpi "ukg"Ectvgt"gv'crl03; ; 6+0"C" P cuucw'i tqwr gt"\*7: "eo "VN+vci i gf "y kj "cp"gzvgtpcn'vci "hqt"c"j qo g'tcpu g"uwf {"lp"vj g'egp'tcn' Dcj co cu'y cu'tgrgcugf "lp"Lwn{"3; ; 9"cpf "tgecr wtgf "3: 7"fc{"u'rcvt"d{"c'huj gto cp"cv'vj g"Nqpi " Krcpf "ur cy plpi "ci i tgi cvkqp"cr r tqzko cvgn{"442"no "htqo "vj g'tgrgcug"r qkpv"Dqrf gp"4222+0" Qpi qkpi "tgugetej "lp"vj g"Gzwo c"Uqwpf . "Dcj co cu'j cu'vcengf "huj "wr "v"422"no "347"o k+y kj " rkngn "guko cvgu"qh'wr "v"552"no "427"o k'cu"vj g{"o qxg"v"ur cy plpi "ukgu"E0F cj ni tgp."Rgtt {"

"  
"  
"  
"

Kpukwng'htq'O ctkpg'Uekppeg'Ectkddgcp'O ctkpg'Tgugctej 'Egpgvt.'r gtu0eqo o 0\q'T0J km'P O HU' UGHUE.'4235+0"Ur cy pgtu'o ki tcvki 'cmipi 'rcti gt'eqvki wqwu'tggh'tcceu'uggo 'v'q'o qxg'i tgcvtg' f kucpegu'v'ci i tgi cvg'vj cp'vj qug'qp'uo cmkucpuf u'qt'cvqm=vj g'eqpvtckp'v'ku'hkng' 'vj gkt' tgmecpeg'qt'kpcdkk'v'v'pcxki cvg'gzv'tgo g'y cvgt'f gr vj u'v'q'tgcej 'uwkcdng'j cdkcv'\*Uctt'g'v'cn' 4229+0"

Htqo 'ceqwuke'vci i kpi 'uwf lgu'ctqwpf 'I nqxtai'Tggh 'Dgrk' g.'Uctt'g'v'cn'0\*4229+0' gcuwtgf " cxgtci g'uy ko o kpi 'ur ggf'qh'P cuucw'i tqwr gt'o ki tcvki 'v'q'cpf 'Htqo 'vj g'ur cy plpi 'ukg'cu'3Q 2'0' 2027'\*UG+no lj qwt0"Vj g'ur ggf'qh'o qxgo gpv'v'q'vj g'ur cy plpi 'ukg'y cu'kf gpv'kcn'v'vj g'ur ggf'qh' v'cxng'icy c{'Htqo 'vj g'ur cy plpi 'ukg'0"Vj g{'pqv'f'vj cv'ugxgtcn'vci i gf 'i tqwr gtu'y gtg'tgeqtf gf 'cv' tgegkxgtu'52'no 'cy c{'Htqo 'vj g'ur cy plpi 'ukg'cpf 'cv'vj g'ur cy plpi 'ukg'rguu'vj cp'46'j qwtu'rcvt0" Vj g{'hqwpf 'ugz'dcugf 'f khgt'pegu'kp'uy ko o kpi 'ur ggf'y kj 'o gcp'ur ggf'qh'o crgu.'4Q'0'204'\*UG+." dgkpi 'uki pkkcepv'f'hcugt'vj cp'hgo crg'i tqwr gtu.'30 '0'204'\*UG+no lj t0"Vj g{'cuq'wugf 'cm' uy ko o kpi 'ugi o gpw'vj cv'y gtg'@'no 'v'q'gxcwcv'v'ko g'qh'f c{'qh'i tqwr gt'o qxgo gpw'v'q'vj g' ur cy plpi 'ukg'd{.'cpf'hqwpf'vj cv'38'vci i gf'hkuj 'o qxgf'qpn'f'v'v'kpi 'vj g'f c{'\*f'ghkpgf'cu'3'j'dghqtg' uwptkug'vj tqw j 'v'q'wpu'gv'+cpf'."hkuj 'o qxgf'dqj 'f'v'v'kpi 'vj g'f c{'cpf'cv'pki j v0I tqwr gt'uy ko " ur ggf'u'f'v'v'kpi 'vj g'f c{'cxgtci gf'3Q 8'0'2025'\*UG+no lj t'cpf'y gtg'uki pkkcepv'f'hcugt'vj cp'o gcp' i tqwr gt'uy ko 'ur ggf'u'cv'pki j v'306'0'208'no lj t+0'

Qdugt'x'cv'kpu'ui i gu'vj cv'kpf kxf wcu'ecp't'gwtp'v'q'vj gkt'qtki kpcn'j qo g'tggh'hqmy kpi " ur cy plpi 0'Ugxgtcn'rcti g'cf wv'P cuucw'i tqwr gt'kp'vj g'Dcj co cu.'engctn'f'uy qnpg'y kj 'i co gvu." f kucr r gctgf 'Htqo 'tgukf gpv'kcn'ct'gcu'htq'r g'kqf u'tcpi kpi 'Htqo '32'f c{'u'dghqtg.'v'c'hgy 'f c{'u'chgt." vj g'hwn'o qpp'qh'F gego dgt'3; ; 0"Vj g{'tgo ckpgf'kp'j qo g'ctgeu'htq'vj g'Lcpwct{'3; ; 2'hwn'o qpp' cpf'y gtg'uggp'pgkj gt'v'q'uy gn'y kj 'i co gvu'pqt'v'q'gzj kdk'eqw'vuj kr'dgj c'xkqt.'ui i gu'v'kpi 'vj cv' p'q'v'cn'o cwtg'hkuj 'ci i tgi cvg'qt'ur cy p'kp'gxgt{'ci i tgi cvkqp'o qpvj 'kp'gcej 'tgr tqf wv'kxg'ugcu'p' \*ROEqrkp.'Eqten'Tggh'Tgugctej 'Hqwpf'cv'kqp'o'Rcrw.'r gtu0eqo o 0\q'[ 0Ucf qx{'P O HU.'3; ; 2+0" Dqrf gp'\*4223+0'tgr qt'v'f'vci i gf'hkuj 't'gwtp'kpi 'v'q'j qo g'tgghu'kp'uwdu'gs wgpv'f'gctu0'Uqple'v'cenc'pi " uwf lgu'ctqwpf 'Nkvr'Ec{o cp'Kucp'f'j'cxg'f'go q'p'v'cv'f'vj cv'ur cy pgtu'o c{'t'gwtp'v'q'vj g' ci i tgi cvkqp'ukg'kp'uweegu'kxg'o qpvj u'y kj 't'gwtpu'v'q'qt'v'qy ctf'u'vj gkt'gukf gpv'kcn't'gghu'kp'dgy ggp' \*Ugo o gpu'g'v'cn'04229+0'Ncti gt'hkuj 'ctg'o qtg'rkng'f'v'q't'gwtp'cpf'ur cy p'kp'uweegu'kxg'o qpvj u' vj cp'uo cmgt'hkuj '\*Ugo o gpu'g'v'cn'04229+0'"

Spawning habitat."Ur cy plpi 'ci i tgi cvkqp'ukgu'v'f'r kcm'f'q'ee'w't'p'gct'vj g'gf i g'qh'kpuwrt" r'ncv'htqo u.'cu'k'wng'cu'72'o 'Htqo 'vj g'uj qtg.'cpf'emug'v'q'c'f'tqr/qh'h'kpv'f'ggr'y cvgt'qxgt'c'y kf g'8/ 82'o +f'gr vj 't'cpi g'cpf'f'kxgtuk'f'qh'u'wdu'tcv'v'f'r gu'\*Etcki '3; 88.'Uo kj '3; 94.'Dwtpgw'J gtngu' 3; ; 97.'Qnugp'cpf'NcRr'eg'3; 9; .Eqrkp'g'v'cn'03; ; 9.'Ectvgt'3; ; ; .Hkp'3; ; 2.'Dggw'cpf'Hk'kf'ncpf'gt' 3; ; 4.'Eqrkp'3; ; 4.'Ci w'krt/Rgt'g'tc'3; ; 6+0'Uk'gu'ctg'ej ctcevgt'k'v'kcm'f'uo cm'j ki j n'f'ektewo uetkdgf " ctgeu.'o gcuwtkpi 'ugxgtcn'j w'p'f'tgf' "o g'v'gtu'kp'f'kco g'v'gt.'y kj 'u'q'h'eqtcn.'ur q'pi gu.'uw'p'f'eqtcn' qwetqr u.'cpf'uc'p'f'f'gr t'gu'k'p'u'\*Etcki '3; 88.'Uo kj '3; 94.'Dwtpgw'J gtngu'3; 97.'Qnugp'cpf' NcRr'eg'3; 9; .Eqrkp'g'v'cn'03; ; 9.'Ectvgt'3; ; ; .Hkp'3; ; 2.'Dggw'cpf'Hk'kf'ncpf'gt'3; ; 4.'Eqrkp' 3; ; 4.'Ci w'krt/Rgt'g'tc'3; ; 6+0'Cd'q'w'82/ : 2'ci i tgi cvkqp'ukgu'j'cxg'dggp't'geqtf gf.'o qu'w'f' 'Htqo " kpuwrt'ctgeu.'cnj qwi j 'o cp{'pq'rupi gt'htqo 0'T'gegp'v'y qtn'j cu'kf gp'v'k'kf'f' i g'qo qtr j qm'j kcn'

"  
"  
"  
"

ulo kretkkgu'kp'ur cy plpi 'uksu'vj cv'o c{ 'dg'wughw'kp'cr r n{ kpi 'tgo qvg'ugpukpi 'v'ej pls wgu'vq' f kueqxtg'r t'gxlqwun{ 'w'pnpqy p'ur cy plpi 'uksu'\*Mqdctc'cpf 'J g{ o cp'4232+0'C'v'ur cy plpi " ci i tgi cvkqp'uksu.'P cuucw'i tqwr gt'v'gpf 'vq'o g'c'p'gt'ctqwpf 'kp'c'ou'uci kpi 'ctgcö'cf l'cegpv'v'j g'eqt'g' ctgc'y j gtg'ur cy plpi 'ce'v'x'k'v' 'ce'w'cm' 'v'cngu'r m'eg'\*Mcf kuqp'g'v'c'f'04232.'P go gj '4234+0'Vj gug' ci i tgi cvkqp'uci kpi 'ctgcu'j cxg'dggp'tgr qtvgf 'cv'f gr vj u'qh'8/72'o 0'Cu'w'pugv'cr r tqcej gu.'v'j g' ur cy pgtu'v'f r k'ecm' 'o q'x'g'ugcy ctf.'kp'v'urki j v'f 'f ggr gt'y cvgt '\*52/82o +0'Ur cy plpi 'twuj gu'j cxg' dggp'f guetkdgf 'g'k'j gt'cu'c'eqnwo p'qt'eqpg'q'h'h'kuj 'q'h'f k'htg'gpv'eqnqt'r j cugu'tkukpi 'vq'y k'j kp'42/47' o 'q'h'v'j g'y cvgt'uw'h'ceg'qt'cu'c'ugt'k'gu'q'h't'wuj gu'd{ 'uo c'm'i tqwr u'q'h'o c'ngu'h'qm'y kpi 'c'ukpi ng'hgo c'ng' \*Qnugp'cpf 'NcR'ceg'3; 9; . 'Ectvgt'3; ; 8.'Ci w'k'ct/Rgtg'tc'cpf 'Ci w'k'ct/F c'x'k'c'3; ; 8+0'

Cm'ur cy plpi . 'cu'ht'cu'ku'npqy p. 'qeewtu'kp'f k'uk'p'ev'ci i tgi cvkqpu'cv'uksu'vj cv'tgo c'kp' eqpukuv'p'v'x'g't'qpi 'v'ko g'r g'tk'qf u'0'Vj g't'g'ct'g'p'q't'gr qt'v'q'h'r c'k'ur cy plpi 0'Ur cy plpi " ci i tgi cvkqpu'j cxg'dggp'tgr qtvgf 'ht'qo 'v'j g'D'c'j co cu.'D'g'k' g.'D'g'to w'f c.'D't'k'kuj 'X'k'i k'p'K'ur'c'p'f u.' Ec{ o cp'K'ur'c'p'f u.'E'w'dc.'J q'p'f w'cu.'L'co c'k'ec.'O g'z'k'eq.'R'w'g't'v'q'T'k'eq.'V'w't'm'i'c'p'f 'E'c'k'eq'u'c'p'f 'v'j g'W'U'U' X'k'i k'p'K'ur'c'p'f u.\*Q'nu'q'p'c'p'f 'N'c'R'ceg'3; 9; . 'E'q'r'k'p'g'v'c'f'03; ; 9.'E'c't'v'g't'3; ; . . 'E'q'r'k'p'3; ; 4.'C'i w'k'ct/ R'g't'g't'c'c'p'f 'C'i w'k'ct/F c'x'k'c'3; ; 8.'R'c'l' 'c'p'f 'I' t'k'o u'j cy '4223+0'U'w'ur g'ev'f 'q't'c'p'g'ef q'v'c'n'g'x'k'f g'p'eg' c'nu'q'k'f g'p'v'h'k'gu'ur cy plpi 'ci i tgi cvkqpu'kp'N'qu'T'q's w'gu.'X'g'p'g' w'g'r '\*D'q'q'o j q'y g't'g'v'c'f'04232+'c'p'f " Q'f 'R't'q'x'k'f g'p'eg'\*R't'c'f'c'g'v'c'f'04226+'k'p'E'q'm'o d'k'æ'U'c'p'c'p'f t'2 u'C't'ej k'r g'nci q'0'P g'k'j g't'ci i tgi cvkqp' p'q't'ur cy plpi 'j cu'dggp'tgr qtvgf 'ht'qo 'U'q'w'j 'C'o g't'k'ec'c'nj q'w'i j 't'k'r g'P cuucw'i tqwr gtu'ct'g' ht'g's w'p'w'v' 'v'c'ng'p'kp'eg't'v'c'k'p'ct'g'cu'\*H'0'E'g't'x'k'i »p.'H'w'p'f c'ek'q'p'E'k'g'p'v'h'k'ec'N'qu'T'q's w'gu'o'X'g'p'g' w'g'r." r gtu'0'eqo o 0'v'q' 'I' 0'U'c'f q'x{ . 'P' O' H'U.'3; ; 3+0'Ci i tgi cvkqp'ur cy plpi 'ku'h'k'ng'y k'ug'w'p'npqy p'ht'qo 'v'j g' N'gu'ug't'c'p'v'k'ng'u.'ht'qo 'E'g'p'v't'c'n'c'o g't'k'ec'u'q'w'j 'q'h'J q'p'f w'cu.'q't'ht'qo 'H'q't'k'f c'0'V'j g'g'p'x'k'q'p'o g'p'v'c'n' c'p'f 'u'q'ek'n'v't'k'i i gtu'v'j cv'ec'w'ug'P cuucw'i tqwr gt'v'q'ci i tgi cv'g'ct'g'p'q'v'y g'n'w'p'f gtu'v'q'q'f . 'c'nj q'w'i j " ej c'p'i k'p'i 'h'w'p'c't'f'k'i j v'eq'p'f k'k'q'pu.'y cvgt'v'go r g't'c'w't'g.'e'w't'g'p'v.'h'g'c't'p'g'f 'd'g'j c'x'k'q't.'q't'c'eqo d'k'p'cv'k'q'p' q'h'v'j g'ug'q't'q'v'j g't'h'c'ev'q'tu'ct'g'v'j g'r q'u'w'v'v'g'f 'd'c'u'k'uh'q't'ci i tgi cvkqp'h'q'to cvkqp'\*E'q'r'k'p'g'v'c'f'03; ; 9." E'c't'v'g't'3; ; . . 'V'w'eng't'g'v'c'f'03; ; 5.'F'q'o g'k'g't'c'p'f 'E'q'r'k'p'3; ; 9.'U'c'f q'x{ 'c'p'f 'G'm'w'p'f'3; ; . . 'R'c'l' 'c'p'f " I' t'k'o u'j cy '4223+0'

Spawning timing0'Vj g'P cuucw'i tqwr gtu'v'j g'm'npqy p'tgr tqf w'v'x'g'o q'f g'q'h'q'to k'p'i " v'c'p'uk'ep'v'ur cy plpi 'ci i tgi cvkqpu'ku'i g'p'g't'c'm' 'r t'g'f l'ev'c'd'ng'y k'j k'p'c'r t'g'u'et'k'd'g'f 'ctgc'0'Ci i tgi cvkqpu' qeew'cv'r t'g'f l'ev'c'd'ng'v'ko g'u'c'p'f 'r m'egu'g'cej "{ g'c't'c't'q'w'p'f 'v'j g'v'ko g'q'h'v'j g'h'm'o q'q'p.'w'w'c'm'f " d'g'y g'g'p'F g'ego d'g't'c'p'f 'O'c't'ej '\*t'g'x'k'g'y g'f 'k'p'U'c'f q'x{ 'c'p'f 'G'm'w'p'f'3; ; . + 'c'nj q'w'i j 'k'p'D'g'to w'f c" ci i tgi cvkqp'ur cy plpi 'qeew't'g'f 'k'p'v'j g'p'q't'y g't'p'w'o o g't'r g't'k'q'f 'ht'qo 'O'c{ 'v'q'L'w'v'f '\*D'c't'f'c'ej 'g'v'c'f'0 3; 7: +0'Q'nu'g'p'c'p'f 'N'c'R'ceg'3; 9; +t'gr qtvgf 'ur cy plpi 'qeew't'k'p'i 'q'p'v'j g'h'k'u'v'h'm'o q'q'p'c'h'g't'v'j g' y k'p'v'g't'u'q'n'k'eg'0'Y q't'n'k'p'i 'ht'qo 'i q'p'c'f 'g'z'c'o k'p'c'v'k'q'pu.'O'w'p't'q'c'p'f 'e'q'm'g'c'i w'gu'\*3; 95+t'gr qtvgf " P cuucw'i tqwr gt'ht'qo 'L'co c'k'ec'æ'q'h'h'uj q't'g'q'eg'c'p'k'e'd'c'p'm'i'v'q'd'g'k'p'ur cy plpi 'eq'p'f k'k'q'p' r t'g'f q'o k'p'c'p'v'f 'k'p'H'g'd't'w'c't' . 'd'w'c'nu'q'v'q'c'ng'u'g't'f'g'i t'g'g'k'p'c'r t'k'i'c'p'f 'O'c{ 0'T'g'eg'p'v'g'x'k'f g'p'eg' u'w'i i g'u'u'v'j cv'ur cy plpi 'ku'c'nu'q'qeew't'k'p'i 'cv'y j cv'c'r r g'c't'v'q'd'g't'g'eq'p'uk'w'w'g'f 'q't'p'q'x'g'n'ur cy plpi " u'k'g'u'kp'd'q'v'j 'R'w'g't'v'q'T'k'eq'c'p'f 'v'j g'W'U'U'K'0'f' w'k'p'i 'L'x'p'g'\*T'0'C'r r g'f'q'q't'p.'W'p'k'x'g't'uk'v'f 'q'h'R'w'g't'v'q' T'k'eq/O'c{ c'i Å'g'l' . 'F'g'r c't'vo g'p'v'q'h'O'c't'k'p'g'U'ek'g'p'eg'0'r gtu'0'eqo o 0'v'q'T'0'J k'm'P O' H'U'U'G'H'U'E.'4234=T'0' P go gj . 'W'p'k'x'g't'uk'v'f 'q'h'v'j g'X'k'i k'p'K'ur'c'p'f u'/'E'g'p'v'g't'h'q't'O'c't'k'p'g'c'p'f 'G'p'x'k'q'p'o g'p'v'c'n'U'w'f'k'gu.'r gtu'0'

"  
"  
"  
"

eqo o 0\q"T0J km"P O HU"UGHUE."4234=F 0Qmgp."Ej lgh"Uelgpkuv"/"U0Vj qo cu"Hkuj gto gpau"  
 Cuuqekvqp"tgr qt\kpi "vj g'hkpf kpi u'qh'T0I qo gl "XKF HY ."r gtu0eqo o 0\q"T0J km"P O HU"UGHUE."  
 4234+tcvj gt "vj cp"fwtkpi "vj g'y kpvgt"o qpvy u."cmj qwi j "hwtj gt"y qtniku"pggf gf "vq'hwm" f qewo gpv"  
 vj gug"qdugtxcvku0"Ur cy plki "qeewtu'hqt"wr "vq"307"j qwtu'ctqwpf "vj g'vko g"qh'lwpuv'g'ht"ugxgtcn"  
 f c{u'kp"geej "qh'ugxgtcn'o qpvy u"\*Y j c{ngp"gv'cn04229+0"Vj g'i qpcf quqo cvk'kpf gz"\*I UK"qh"  
 hgo crgu"\*Q0"vj g'tgrvkg"qxct{/vq/dqf {"y gli j w'ku'c"i qpf "kpf kecvt"qh'ur cy plki "ugcuqpcrk{"\*Hi 0  
 3+0"

Vj g'tgr tqf wevkg"

ugcuqp"kp"vj g'P cuucwi tqwr gt"ku"  
 dtkgh'cpf "gxkf gpn{"cuuqekvqf "  
 y kj "vgo r gtcwtg"cpf "o qqp"  
 r j cug."ceeqtf kpi "vq"i UK"  
 i qpcf cnj kuvmji {."  
 o cetqueqr ke."cpf "qqe{vg"  
 f lco vgt"cpn{ugu0"Cv'ny gt"  
 rkvwfu."tgr tqf wevkg"cevkvk"  
 ruw'ht"cdqw'qp'g'y ggm'r gt"  
 o qpvy ."ht"qpg"v"vj tgg"o qpvy u"  
 gcej {"gct."dgw ggp"F gego dgt"  
 cpf "Hgdwtct{"\*Hi 03+."gkj gt"  
 r gcnkpi "kp"Lcpwct{"\*Uo kj "  
 3; 94."Qmgp"cpf "NcRrcg"3; 9; ."  
 Erctq"gv'cn03; ; 2."Eqrkp"3; ; 4."  
 Rqy gni'cpf "Vwengt"3; ; 4."




Figure 1. Percent frequency of different gonad development stages for female Nassau grouper by month collected from Belize from 1984-86 (n=1,232) [redrafted from Carter et al. 1994]

Ci vkrct/Rgtgtc"3; ; 6."O kmg<sup>3</sup>+qt"dgw ggp"Lcpwct {"cpf "Cr tkn"Vj qo ruqp"cpf "O wptq"3; 9; -0"K"  
 o qtg"pqtj gtn{ "rkvwfu"\*Q0"Dgto wf c+."vj g'tgr tqf wevkg"ugcuqp"hcni"dgw ggp"O c {"cpf "Cwi wuv."  
 r gcnkpi "kp"Lwn{"\*Nc"i qteg"3; 5; . "Uo kj "3; 93."DwtpgwJ gtngu"3; 97+0"Gzegr vkuu"v"vj g'r quikng"  
 rkvwfu kpcnr cwgt'p'y gtg'vj g'ecr wtg"qh't gegpn{/ur cy pgf "hgo crgu"kp"Ugr vgo dgt"kp"Ewdc"eqwr rnf "  
 y kj "vj g"qdugtxcvkuq."qh'c"i tqwr "qh'P cuucwi tqwr gt"cv'4; "o "f gr vj "kp"vj g'uco g'qecvkuq"\*Erctq"gv'  
 cn03; ; 2+0"

Ur cy plki "ku"j ki j n{ "u{pej tqpk gf "cpf "qeewtu'dtkghn{ "y kj kp"cdqw'c'y ggm'qh'hwm'o qqp."qt"  
 dgw ggp'hwm'cpf "pgy "o qqp"\*Uo kj "3; 93."Eqrkp"3; ; 4."Vwengt"gv'cn03; ; 5."Ci vkrct/Rgtgtc"3; ; 6."  
 Ectvt"gv'cn03; ; 6."Vwengt"cpf "Y qqf y ctf "3; ; 6+."y kj kp"vj g'pcttqy "vgo r gtcwtg'tcpi g'qh'47/  
 48Å"cpf "qxgt"cy kf g'tcpi g'qh'f c{/ngpi vj u"\*Eqrkp"3; ; 4."Vwengt"gv'cn03; ; 5."Ectvt"gv'cn03; ; 6+0"  
 Y j c{ngp"gv'cn04229+j cxg'r tqr qugf "c'r tgf kevkg'i wkf g'ht"vj g'Ec{o cp "Kmpf u'vj cv'kh'vj g'ur cp"qh"  
 vko g'htqo "vj g'y kpvgt"uqnvleg"v"Lcpwct {au'hwm'o qqp"ku'rguu'vj cp"52"f c{u."vj gp"Hgdwtct {"y cu'vj g"  
 o clqt"ur cy plki "o qpvy 0Eqpxgtugn{ ."h'ky cu'i tgcvt"vj cp"52"f c{u."Lcpwct {"y cu'vj g'o clqt"

<sup>3</sup>O kmg. "Y 03; ; 60Ur cy plki "ci i tgi cvkpu'qh'vj g'P cuucwi tqwr gt."Epinephelus striatus."cpf "cuuqekvqf "Hkuj gt {"kp"  
 Dgik g0Cf xcpegu"kp"tggh'Uelgpegu."Qevqdg"48/4: .3; ; 6."Wpkgtuks{"qh'O kco k'Hqtkf c0Wpr wdr0f cvc."r 03; 0  
 35"


"  
 "  
 "  
 "

ur cy pl̄pi 'o qpvj 0"Qvj gt'tgugctej gtu'j cxg'tgeqi pl̄ gf 'vj cv'vj g'v̄ko l̄pi 'qh'vj g'hwm'o qpvj .gctn'qt" r̄v'g'lp'vj g'o qpvj .ecp'i kxg'cp'lpf l̄c̄v̄k̄q̄p'qh'y j gp'vj g'r gen'ur cy pl̄pi 'y km'q̄eewt" \*T0Cr r grf qqtp." Wpl̄xgtukv' 'qh'Rwgtvq'Tleq/O c{ci Ågl .F gr ctvo gpv'qh'O ctk̄pg'Uel̄gpeg.'r gtu'leqo o 0'vq'T0J km" P O HU."4233=O 0Uej @gt."Wpl̄xgtukv' 'qh'Rwgtvq'Tleq/O c{ci Ågl .F gr ctvo gpv'qh'O ctk̄pg'Uel̄gpeg." r gtu'leqo o 0'vq'T0J km" P O HU."4233+0"

Ugc'uwthceg'vgo r gtcwtg."cu'k'hcm'dg{qpf "48ÅE"vq'ugcuqpcn'ny u."j cu'cnuq'dggp'r tqr qugf " cu'c'ng{ 'eqpvtqn'qp'ur cy pl̄pi 'v̄ko l̄pi " \*Eqr̄kp"3; ; 4+0"Uko k̄rct'cuuqek̄v̄k̄pu'dgwy ggp'tgr tqf w̄v̄k̄q̄p." v̄go r gtcwtg."cpf "n̄w̄pct'r j cug'y gtg'cnuq'pqvgf "lp'ecr v̄kxg'cpko cn0"P cuucw'i tqwr gt'tckugf "Itqo " gi i "vq'o cwtk'v'lp'Hqtkf c"cpf "Dgto wf c'w̄pf gt'eqpf k̄k̄q̄pu'qh'co d̄k̄gpv̄iki j v."v̄go r gtcwtg."cpf " ucr̄k̄p̄k̄v'."gzj k̄dk̄gf "qxctk̄cp"o cwt̄v̄k̄q̄p."qxw̄v̄k̄q̄p."dgj cxk̄qt."cpf "eqm̄t'ej cpi gu'ej ct̄cev̄gtk̄v̄ke"qh' ur cy pl̄pi ."cv'48ÅE."cnj qwi j "pq'ur cy pl̄pi 'y cu'q̄dugt̄xgf " \*Vwengt"cpf "Y qqf y ctf "3; ; 6+0"

Vgo r gtcwtg'ku'gk̄f gpv'v'c"o qtg'lo r qt̄v̄cpv̄uko w̄w̄u'hqt'ur cy pl̄pi 'vj cp'f c{ 'rgpi vj . " ceeqtf l̄pi "vq'r c̄w̄gt̄pu'qh'x̄q̄n̄w̄pct { 'ur cy pl̄pi 'lp'ecr v̄kxg'hkuj 0"Y j k̄ng'ur cy pl̄pi 'q̄eewt̄gf "cv" v̄go r gtcwtg'ucpi l̄pi "Itqo "450"/"490 ÅE."46/49ÅE"y cu'vj g'o quv'uw̄k̄cd̄ng'dcugf "qp'ur cy pl̄pi " Itgs w̄p̄e { "cpf "x̄q̄n̄w̄o g."cpf "gi i "cpf "h̄et̄x̄cn̄f ḡx̄gn̄r o gpv' \*Vwengt"3; ; 6."Y c̄v̄cp̄cd̄g'gv'c̄r03; ; 7c." Vwengt'gv'c̄r03; ; 8+0P cuucw'i tqwr gt'ur cy pgf "ur qp̄v̄cp̄ḡq̄w̄un' "qp̄g'f c{ 'r tk̄qt"vq'vj g'pgy "o q̄q̄p'lp" Cr tk̄i3; 85"lp'cp'cs w̄ctk̄w̄o "lp'Ew̄dc'w̄pf gt'ct̄v̄h̄ek̄cn̄iki j v'cpf "y cv̄gt"v̄go r gtcwtg'qh'460 ÅE" \*I wk̄ctv̄O c̄pf c{ 'c̄pf "Lw̄³ tgl /H̄gt̄p̄cp̄f gl "3; 88+0'

Spawning behavior. "Hkuj "i gpgtcm' "i cvj gt'pgct"vj g'ur cy pl̄pi "uk̄g"e'f c{ "qt"vy q'r tk̄qt"vq" l̄p̄k̄k̄v̄k̄q̄p'qh'ur cy pl̄pi 0"Uw̄txg{ u'ecp'k̄f gpv̄kh' "w̄p̄w̄w̄cm' "j ki j "pwo d̄gtu'qh'lp̄f k̄k̄f w̄cnu'gk̄j gt" l̄p̄v̄gt̄cev̄k̄pi "qt't̄ḡun̄k̄pi "qp̄lp̄gct"vj g'd̄q̄w̄qo 0"Rtk̄qt"vq'ur cy pl̄pi .lp̄f k̄k̄f w̄cnu'o km̄ct̄q̄w̄pf "qx̄gt"vj g" uwd̄ut̄cv̄g'gzj k̄dk̄k̄pi "q̄pg'qh'h̄q̄w'f k̄v̄k̄p̄v̄k̄g"eqm̄t'r j cugu" \*3+"d̄ct̄tgf " \*p̄q̄to cn=" \*4+"d̄leq̄m̄t=" \*5+" y j k̄g'd̄gm' =qt" \*6+"f c̄tn̄i j cug" \*Hki 04+0"Vj gtg'ctg'lp̄v̄gti t̄cf c̄v̄k̄q̄pu'qh'vj gug'r c̄w̄gt̄pu.'y kj "t̄cr k̄f "



(Diagrams adapted from Sadovy & Eklund, 1999. Photograph by Andy Stockbridge)

Figure 2. Color phases of Nassau grouper. From Paz and Grimshaw 2001b.

ej cpi gu'co qpi 'r c̄w̄gt̄pu'r qūk̄d̄ng" \*Eqr̄kp"3; ; 4+0"Vj g'd̄ct̄tgf " \*v' l̄rc̄cn" eqm̄t'r j cug'ku'h̄q̄w̄pf "co qpi "hkuj "lp'vj g" ci i t̄gi c̄v̄k̄q̄p'lp'vj g'o q̄tp̄k̄pi 0"Vj g" d̄leq̄m̄t'r j cug."h̄t̄u'f ḡūet̄k̄d̄gf "d{ "Uo kj " \*3; 94+.q̄eewt̄u'lp'd̄q̄vj "o c̄rgu'c̄pf " h̄go c̄rgu'c̄pf "ku'f qo l̄p̄cpv'f v̄tk̄pi "vj g" r̄v̄g'c̄h̄gt̄p̄q̄q̄p'y kj "o quv'h̄kuj "d̄geqo l̄pi " d̄leq̄m̄t̄gf "d{ "f w̄um'y j gp'ur cy pl̄pi " q̄eewt̄u" \*Eqr̄kp"3; ; 4+0"lp'vj ku'r j cug."vj g" w̄r r gt'd̄qf { "cpf "j gcf "d̄geqo g'f c̄tn̄i" y j k̄g'vj g'd̄gm' .ny gt'uk̄f gu.'h̄r u."c̄pf " c̄m̄h̄k̄pu'dw'vj g'f q̄t̄uc̄n̄ct̄g'y j k̄g0"C" y j k̄g'g{ ḡd̄ct'ku'r tqo l̄p̄gpv'qp'vj g'j gcf " \*Eqr̄kp"3; ; 4+0"lp'vj g'y j k̄g'd̄gm' "r j cug." uggp"co qpi "r t̄guwo gf "h̄go c̄rgu'y kj "

"  
"  
"

dwi kpi 'cdf qo gpu' r tqdcndf 'hwn'qh'qxc+'vj g'pqto cn'eqmt'r cwgt'ku'o qf kkgf 'uwej 'vj cv'vj g' cdf qo kpcn'ctgc'ku'f kwpewf 'y j kg'Eqrkp'3; ; 4+0"Vj g'rcuv'r cwgt'p.'vj g'Sf ctn\$'r j cug.'ku'hqwpf 'kp' eqwt'kpi 'cpf'ur cy plpi 'huj =vj g'dqf {'cpf 'hpu'dgeqo g'f ctni tc {'vq'drcenly kj 'vj g'dcttgf 'r cwgt'p' xkukng'dgpgev 'vj g'f ctni ki o gpvc'kp0"Vj g'ug'huj 'ctg'r tqdcndf 'hgo cngu'tgcf {'vq'ur cy p'ukpeg' vj g{'cr r gct'vq'rgcf 'i tqwr /ur cy plpi 'gxgpw'Eqrkp'3; ; 4+0"

Eqwt'vuj kr 'ku'kpf kcvgf 'd{'y q'dgj cxlqtu'y j lej " qeewt'rcvg'kp'vj g'chgt'pqqp<ohqmy kpi o'cpf "oektkpi o' "Eqrkp'3; ; 4+0"ohqmy kpi o'qeewtu'cu'qpg'qt'o qtg'huj 'kp' vj g'dleqmt'r j cug'uy ko 'enqunf 'dgj kpf 'cp'cr r ctgpv'hgo cng' y j kg'oektkpi o'qeewtu'cu'c'dleqmt'r j cug'huj 'ektngu'c' dcttgf'qt'f ctni j cug'huj 0"Rtqi tguukp'htqo 'eqwt'vuj kr 'vq' ur cy plpi 'o c{'f gr gpf 'qp'ci i tgi cvkq'uk' g'dw'i gpgtcmf " qeewtu'cu'hqmy u0"Vqy ctf u'vj g'rcvg'chgt'pqqp'huj 'o qxg' r tqi tguukgnf 'j ki j gt'kp'vj g'y cvgt'eqno p.'y kj 'cp' kpetgcukpi 'pwo dgt'gzj kdkkpi 'vj g'dleqmt'r j cug'Eqrkp' 3; ; 4.'Ectvgt'gv'cn03; ; 6+0"Vj g'ci i tgi cvkq'vj gp'o qxgu' kp'v'f ggr gt'y cvgt'uj qt'v'f 'dghgt'ur cy plpi "Eqrkp'3; ; 4." Vwengt'gv'cn03; ; 5.'Ectvgt'gv'cn03; ; 6+'d {'y j lej 'vko g'cm' kpf kxf wcu'ctg'gkj gt'o'f ctni j cug'o'qt'odleqmt0"Dleqmt' huj 'vj gp'hqmy 'f ctni j cug'huj 'enqunf 'cpf 'i tqwr / ur cy plpi 'qeewtu'kp'uw'd/i tqwr u'qh'5/47'huj "Hki 05+0"Uko krc't'ceeqwpu'qh'ur cy plpi 'dgj cxlqt' htqo 'vj g'WUUK00'f guetkdgf 'vj g'ci i tgi cvgf 'huj 'cu'c'eqpg'Hki '60'kp'vj g'y cvgt'eqno p'tcvj gt'vj cp' dgkpi 'f kur gtugf 'cetqu'vj g'dqwqo "Qnugp'cpf 'NcRrcg'3; 9; +0"

Uo cmgt'ci i tgi cvkqu'v'gpf 'vq'kpenf g'hgy gt'dleqmt'r j cug'huj 'cpf 'i gpgt'cn'cvkxk' 'cpf " eqmt'ej cpi gu'ctg'nguu'kwpug'Eqrkp'3; ; 4.'Ci wkrct/Rgtgc'cpf 'Ci wkrct/F cxkrc'3; ; 8+0"Ur cy plpi " kpxqkxg'c'tcr kf 'j qtk'qpvcn'uy ko 'hqmy gf 'd{'c'ektkpi 'cuegpv'qh'uo cm'uw'd/i tqwr u'kp'v'vj g'y cvgt' eqno p.'y kj 'tgrgcug'qh'ur gto 'cpf 'gi i u'cpf 'c'tcr kf " tgwtp'qh'vj g'htci o gpv'f 'uw'd/i tqwr 'vq'vj g'uwduxcvg0" I co gv'g'tgrgcug'ku'y gm'cdq'xg'vj g'dqwqo 0Qp'vj g'dcuku' qh'qdugt'xcv'kpu'qh'qxgt'72"ur cy plpi 'gxgpw.'vj g'gct'rkuv' cpf 'rcv'ur cy plpi 'qeewt'gf 'y kj kp'42'o kpwgu'qh' uwpugv'cpf 'o qu'v'y kj kp'32'o kpwgu'qh'uwpugv'Eqrkp' 3; ; 4+0"J {f'cvkq'qh'xkgm'qi gple'gi i u'qeewtu'kp'vj g' chgt'pqqp'uj qt'v'f 'dghgt'ur cy plpi 0"





Figure 4. Depiction of spawning rush. From Olsen and LaPlace 1979.



Spawning Behavior of Nassau Grouper (Adapted from Sadovy & Eklund, 1999)

Figure 3. Depiction of spawning rush. From Sadovy and Eklund 1999.


Cnj qwi j 'ci i tgi cvkqu'htqo 'o qtg'vj cp'qpeg'c'v'c' r ct'kewrct'ukg'f wtkpi 'c'tgr tqf wv'xg'ugcuq'p.'k'ku'wpergct' y j gj gt'vj g'uco g'kpf kxf wcu'r ct'v'ekr cvg'gcej 'vko g0" J qy gxgt.'ugxgct'nhgo cngu'htqo 'qpg'ci i tgi cvkq'p' eqpvc'kpgf 'tkr g'cpf 'uw'd/tkr g'qqe {'v'v'v'q'v' gj gt'y kj 'r quv'

"  
"  
"  
"

qxwrcvt { "hmlergu" \*y j lej 'tgo clp' chvgt "o cwtg" qqe { vgu'j cxg' dggp' tgrgcugf + "uwi i guv'kpi "vj cv" kpf kxkf wcn' hgo crgu' ur cy p' tgr gcvgf n' { "qp' f khtgtpv' f c { u' f wtkpi "qpg" ci i tgi cvkqp " \*Uo kj "3; 94." Ucf qx { . "P O HU. "r gtu0' qdu0' "O qtgqxgt. "gzco kpcvkqp' qh' ur cy plki "qp' xkf gqvr g' kpf kecvgf "vj cv" f wtkpi "5/6" uweeguukxg' i co gvg' tgrgcugu' d { "c' uwd/ i tqwr "y kj lp' c" 37/42" ugeqpf "r gtkqf. "vj g' uco "g" hgo crg' r' gf "cm' ur cy plki "gxgpw. "ci clp' kpf kecvkpi "o wvkr ng' "gi i "tgrgcugu' k' "qpg' gxgpki " \*Eqrkp " 3; ; 4+0" P q' f cv' ctg' cxkrcdng. "j qy gxgt. "cf f tguakpi "y j gvj gt "gcej "o cwtg' hgo crg' ur cy pu' k' "gxgt { " ci i tgi cvkqp "o qpj . "qt' kpf ggf. "gcej " { gct0

K' rcti gt "ci i tgi cvkqpu. "c' ercgt "kpet gcug' k' "vj g' r' tqr qt vkap' qh' vj g' dleqmt "r j cug' v' "qvj gt " eqmt "r j cugu' htqo "2027" gctn' { "k' vj g' ci i tgi cvkqp "v' "2062" q' vj g' f c { "qh' ur cy plki "uwi i guv' g' "vj g' eqmt "r j cug' kpf kecvgf "d' gj cxkqtcn' c' pf "r j { ukqmi kecn' r' tgr ctgf pguu' v' "ur cy p' " \*Ctej gt "gv' cr04234+0" Y j kg' P cuucw' i tqwr gt "k' "i tqwr u' qh' cu' hgy "cu' 42" huj "y gt g' uggp' v' "ur cy p. "Eqrkp " \*3; ; 4+ "tgr qt vgf " uwej "uo cm' i tqwr u' r r gctgf "v' u' j qy "uwduxpvcem' { "hgy gt "huj "k' vj g' dleqmt "r j cug' vj cv' { r kecm' " r tgegf gu' ur cy plki 0k' vj g' Ec { o cp "Kurcpf u. "huj "k' uo cm' ci i tgi cvkqpu' i cvj gtgf "qp' uksg' hqt "h' ppi gt " vj cp' vj qug' k' "rcti g' i tqwr u' \*D0Ugo o gpu. "Uetkr r u' k' p' ukxwv' qh' Qegcpqi tcr j { "Wpkxgtukv' { "qh' Ecn' hqt p' k' "o "Ucp' F k' gi q. "r gtu0' eqo o 0v' [ 0Ucf qx { . "Wpkxgtukv' { "qh' J qpi "Mqpi . "4234+ "r tguwo cdn' " gzvgpf kpi "qt' f g' r { kpi "ur cy plki 0"

*Fecundity.* "Hgewpf k' { "guvko cvgu' htqo "y kf /ecwi j v' P cuucw' i tqwr gt "ctg' hgy "cpf "xctkgf . "dw' uwi i guv' c' o gcp' tgrcvxg' hgewpf k' { "qh' dgvy ggp "5" c' pf "7" gi i ulo i "qh' tkr g' qxct { . "f gr gpf kpi "qp' vj g' o gvj qf "wugf . "k' vj g' y qtf u. "y j lej "uwi gu' qh' qqe { vgu' ctg' k' p' emf gf "k' "gi i "eqwpu0" Guvko cvgu' htqo " Dgrk' g' \*Hi 07+ { k' r' gf "c' o gcp' tgrcvxg' hgewpf k' { "qh' 60 "gi i ulo i "qxct { "y gki j v' c' pf "c' o gcp' v' wcn' pwo dgt "qh' qqe { vgu' \*uwi g' w' pur gek' hgf + "qh' 6.422.222 " \*t' cpi g' ? "572.222/8.722.222 " hqt "hgo crgu' htqo "522" v' "922" o o "UN+ " \*Ectvgt "gv' cr03; ; 6+0" Guvko cvg' pwo dgt "qh' gi i u' k' vj g' tkr g' qxct { " \*; 20" i + "qh' c' 667" o o "UN" k' k' wcn' htqo "Dgto wf c' y cu' 9: 7.323 " \*Dctf cej "gv' cr03; 7: +0" k' vj g' Xki k' " Kurcpf u. "hgewpf k' { "guvko cvgu' o cf g' htqo "64" o cwtg' hgo crgu' i cxg' c' o gcp' xcnwv' qh' 60 9 "gi i ulo i "qh' qxct { " \*uf 0? "4054+ y kj "o gcp' gi i "r tqf wvkap' qh' 6.: 22.222 "gi i u' \*Qugp' c' pf "NcRrcg' 3; 9; +0" J qy gxgt. "ukpeg' vj ku' r' wgt "guvko cvg' k' p' emf gu' r tg/ xkgm' i gple "qqe { vgu. "y j lej "o c { "p' v' t' getv' k' v' k' v' vj g' xkgm' i gple "uqen' r' tkqt "v' ur cy plki . "k' ku' eqpukf gtgf "v' dg' cp' qxgt guvko cvg0" Hgewpf k' { "guvko cvgu' y gt g' cnq' o cf g. "dcugf " qp' xkgm' i gple "qqe { vgu' qpn' . "htqo " Dcj co cu' huj "r tqf wvkap' "c' o gcp' tgrcvxg' hgewpf k' { "qh' 40 "gi i ulo i "tkr g' qxct { " \*uf 0? " 30; =p? "86+ c' pf "c' o gcp' hgewpf k' { "qh' 938.886 " \*t' cpi g' ? "33.946" / "6.549.662 " hqt " hgo crgu. "697/8: 8" o o "UN+0" Guvko cvgu' qh' qqe { v' r tqf wvkap' htqo "cpko cni' k' pf wvgt "v' ur cy p' k' "ecr v' k' v' ctg' emugt "v' vj qug" dcugf "uqen' { "qp' xkgm' i gple "qqe { v' eqwpu0"



1984-1996  
**Figure 5. Fecundity of female Nassau groupers as a function of size (from Carter et al 1994)**



\*Tlej ctf u'gv'cr04227-0"Vj g'ngpi 'f qtucn'cpf 'r grkle'ur kpgu'ctg'htci krg'cpf 'hgy 'ur geko gpu'j cxg" ur kpgu'kpcev.'y j lej 'o cmgu'kf gpv'k'ecv'kp'r tqdrgo cvk'kh'o g'k'ueu'cpf 'ur kpggv'o qtr j qmji { " ecppqv'dg'cuuguugf 0" Tlej ctf u'gv'cr0\*4227+r tqxkf g'c'r tqxkukqpcn'hg{ 'v'f kuetko kpcv'g'rtxcg'qh' uqo g'v'ur geku'qt'ur geku'i tqwr u0

Nctxcn'cpf "gctn{ 'lwxgpkrg'r j cugu'ctg'y gmf guetkdgf 'hqt 'y g'P cuucw'i tqwr gt "dgecwug'qh" uweeguuhw'lecr vxg'htg'v'k'k'cv'kp'cpf 'ur cy plpi 0P gy n{ 'j cvej gf 'rtxcg'eqmgevgf 'htqo 'kpf wegf " ur cy plpi 'o gcwv'gf "30/30 'o o 'pqvej qtf 'ngpi vj '\*P N+\*Rqy gm'cpf "Vwengt"3; ; 4+0"Nctxcg'j cf " r ki o gpv'gf "g{ gu'6: 'j qwtu'r quv'j cvej kpi 'cpf "dgi cp'hggf kpi 'y kj kp'82'j qwtu'\*Vwengt'gv'cr03; ; 3+0 F gxgnr o gpv'j cu'dggp'f guetkdgf 'hqt'rdqtcvqt{/tgctgf 'ur geko gpu'htqo 'y g'gi i 'v'c'350'o o 'UN' rctxc'er r tqzko cvgn{ '62'f c{ u'r quv'j cvej kpi '\*Rqy gm'cpf "Vwengt"3; ; 4+\*Hki u035C/'35I 'kp'Ucf qx{ " cpf "Gmwpf"3; ; ; +0'Hkpu'f gxgnr 'kp'vj g'qtf gt'qh'r grkle.'htuv'f qtucn'ecw'cn'r gevqtcn'cpcn'cpf " ugeqpf "f qtucn'0"Vj g'cf wv'eqo r ngo gpv'qhr' tpekr cn'ecw'cn'hp'tc{ u'y cu'c'w'k'p'gf 'cv'80'o o 'UN'cpf " qh'f qtucn'ur kpgu'cv'vj g'r quv'ngz'k'p'uci g'cv'cr r tqzko cvgn{ '80'o o 'UN'y kj 'eqo r ngv'k'p'qh'htuv'cpf " ugeqpf "f qtucn'cpf "cpcn'k'p'cv'90'o o 'UN'0"Rt ghngz'k'p'rtxcg'dgeqo g'ngz'k'p'rtxcg'q'x'gt'vj g'tcpi g' qh'70/'70'o o 'P N'cpf 'ngz'k'p'v'q'r quv'ngz'k'p'qeewtu'dgy ggp'80'cpf '80'o o 'P N'\*Rqy gm'cpf " Vwengt"3; ; 4+0"Nctxcg'y gtg'r rcpmqple'wpv'k'64/'92'f c{ u'r quv'j cvej kpi 'y kj 't'c'puh'qto cv'kp' qeewt'k'pi 'kp'nguu'vj cp'q'p'g'y ggm'\*Rqy gm'cpf "Vwengt"3; ; 4.'Vwengt'cpf "Y q'q'f y ctf "3; ; 6+0"

P gy n{/j cvej gf 'rtxcg'ctg'k'p'epur'k'w'q'w'ur{ 'r ki o gpv'gf 'cpf 'urki j v{ 'ewtxgf 'ctqwpf 'vj g' { qm' uce'y j gp'ct'v'k'ekm{ 't'gctgf '\*Rqy gm'cpf "Vwengt"3; ; 4+0"Y krf /ecwi j v'rtxcg'gzj kdk'ugxg'tcn'uo cm" f gpf tkle'o g'rcpqr j qtgu'qp'vj g'upqw'\*Uo kj "3; 93.'Nctqej g<sup>4</sup>+0"l qm'uce'rtxcg'y kj 'c'f gxgnr kpi " o qwj 'j cxg'c'ej ctcevg'tk'ue'r ki o gpv'r cw'gtp'kp'vj g'htqo 'qh'c'f k'k'p'ev'ok'p'x'gt'v'gf 'ucf f ng'o'qp'vj g" xgp't'cn'o k'f r'k'p'g'cpf 'r'v'g't'cn'uw't'c'eg'qh'vj g'ecw'cn'r gf w'p'eng'\*Rqy gm'cpf "Vwengt"3; ; 4+cpf " ur geko gpu">43'o o 'UN'cnuq'rcen'vj g'ecw'cn'r gf w'p'eng'd'rq'vej 'y j lej 'ku'h'q'wpf 'kp'cm'h'k'uj "@7'o o " \*Uo kj "3; 93+0Rki o gpv'r cw'gtpu'ej cpi g'o ctngf n{ 'f w'k'pi 'vj g'ngz'k'p'uci g.'cpf " { q'w'pi 'r quv'ngz'k'p' rctxcg'\*>80 'o o 'UN'+ctg'uko k'rc't'v'q'rc'v'ngz'k'p'rtxcg'0'k'uo cm'l'w'x'g'p'k'gu'vj gtg'ku'c'ej ctcevg'tk'ue" r'k'p'g'qh'd'rcen'ur qw'c'ru'ppi 'vj g'dcugu'qh'vj g'f qtucn'tc{ u'r quv'g't'k'q't'v'q'vj g'h'h'j 'ur k'p'g'\*Uo kj "3; 83+0' Vj g'r cw'gtp'qh'x'gt'v'k'ec'n'd'ct'u'uggo u'v'q'f gxgnr 'cv'cd'q'w'62'o o 'kp'ur geko gpu'htqo 'y g'D'c'j co cu" \*Uo kj "3; 83+0'

Rt ghngz'k'p'cpf 'ngz'k'p'gr k'p'gr j gr'k'p'cg'rtxcg'ctg'f k'h'k'ew'v'q'k'f gpv'k'h{ 'r qu'k'k'gn{ 'cu" *Epinephelus striatus*.'cnj qwi j 'egt'v'k'p'eqo d'k'p'cv'k'p'u'qh'r ki o gpv.'h'p'ur k'p'g'gw.'cpf 'ur k'p'g'ngpi vj u" p'ctt'qy 'f qy p'r qu'k'k'k'k'k'g'u'\*M'g'p'f cm'3; 9; .L'qj pu'q'p'cpf 'M'g'g'p'g't"3; ; 6.'Rqy gm'cpf "Vwengt"3; ; 4+0" Y kj 'r quv'ngz'k'p'rtxcg'i t'g'cv'gt'vj cp'90'o o 'UN'k'ku'r qu'k'k'k'g'v'q'ugr c't'c'v'g'P cuucw'i tqwr gt 'htqo " q'v'j gt'i tqwr gtu."g'z'egr v'ht' *E. adscensionis*."qp'vj g'd'c'uku'qh'f qtucn'cpf "cpcn'k'p'tc{ 'eqw'p'w.'ur k'p'g'gv' eq'p'h'i w'c'v'k'p'.ugeqpf 'htuv'f qtucn'hp'ur k'p'g'ngpi vj 't'g'r'v'x'g'v'q'UN.'cpf 'ecr w'g'ng'ecv'k'p'\*Rqy gm' cpf "Vwengt"3; ; 4+0"

Nctxcg'cw'k'p'c'o czko wo 'uk'g'qh'52'o o 'UN'\*cxg'tci g'4506'o o +d{ '58'f c{ u'ch'v'gt" r t'guwo r v'x'g'ur cy plpi '\*Uj gp'ngt'gv'cr03; ; 5+0"Nctxcg'eqmgevgf "32'f c{ u'ch'v'gt'r tqd'cd'ng'ur cy plpi "

.....  
4"Nctqej g.'Y c{ pg0U'q'p'g'h'uj 'G'p'x'k'q'p'o gp'v'cn'cpf "V'c'z'q'p'q'o l'e'U'g't'x'legu.'D'qz'438.'G'p'q'ud'w'i 'H'c'm.'XV'276720W'p'r w'd'r0 f'c'c'0'

"  
"  
"  
"

o gcuwtgf '8/32'o o "UN0"Qxgt"37/fc{ 'r gtlkf .: /44'fc{ u'chvgt 'y g'hwml'o qqp."rtxcn'lk' gu"  
kpetgcugf 'ltqo '70'vq'32'o o "UN"i tggpy qqf '3; ; 3."Uj gpmgt'gv'cr03; ; 5+0"Rgrci le 'lwxgpkrgu'y gtg"  
eqmgvxf 'wr 'vq'68'fc{ u'hqmny kpi 'c'r tguwo r vxg'ur cy plpi 'o qqp."cpf 'dgpvj le 'lwxgpkrgu'y gtg'htuv"  
hqwpf "qp'ct'v'k'ekn'cpf 'pcwnt'gghu'cv'69'fc{ u0'Rgrci le 'lwxgpkrgu'cngp'lp'ej cpgn'pgu'lwuv'r tkt"  
vq'ugwgo gpv'o gcuwtgf '44/49'o o "UN"Eqrkp'3; ; 4."Eqrkp'gv'cr03; ; 9+0"Vtcpu'k'kp'ltqo 'rtxcn'vq"  
lwxgpkrg'r j cugu'qeev'u'cv'8/9'y ggmu'ht' y kf 'hkuj' c'pf '8/32'y ggmu'ht' 'hkuj' 't'kugf 'wpf gt 'ct'v'k'ekn'  
eqpf k'k'qpu'ltqo 'k'pf wegf 'ur cy pu0'Vj g'y kf /ecwi j v'rtxcg'i tgy 'o qtg'umy n' 'y' cp'rtxcg'ltqo "  
k'pf wegf 'ur cy pu"Uj gpmgt'gv'cr03; ; 5."Vwengt'cpf "Y qqf y ctf '3; ; 6."Eqrkp'gv'cr03; ; 9+0"ltqo "  
j cvej gt { 'uwf lgu. 'rtxcn'f w'c'v'k'p'ku'guko cvgf 'vq'tcpi g'dgy ggp'47'cpf '97'fc{ u"Ngku'3; ; 9."  
Vwengt'cpf "Y qqf y ctf '3; ; 6+0"Qv'rkj 'cpcn'uku'q'h'p'gy n' 'ugwrgf 'lwxgpkrgu'lp'y g'Dcj co cu'  
guko cvgf 'r grci le 'rtxcn'f w'c'v'k'p'cu'tcpi kpi 'ltqo '59'vq'67'fc{ u"Eqrkp'gv'cr03; ; 9+0'

Rtguwo r vxg'f ckn' 'kpetgo gpv'lp' r'r k'n'q'h'y kf /ecwi j v'rtxcg'k'pf k'ecv'g'c'rtxcn'r gtlkf 'qh'  
57/'62'fc{ u'cpf 'uwr r q't'v'ht'v'k'k' cv'k'p'cv'y' g'hwml'o qqp0'C'o gcp'rtxcn'r gtlkf 'qh'6308'fc{ u'y' cu'  
k'pf k'ecv'g'f 'ltqo 'p'g'v'ecwi j v'uc'o r r'gu"Eqrkp'3; ; 4."Eqrkp'gv'cr03; ; 9+0"Rt'gugwgo gpv'q'v'rkj "  
kpetgo gpv'y gtg'f k'k'p'ev'cpf 'g'cukn' 'eq'w'p'v'g'f "y qy g'xgt. 'ugwgo gpv'o ct'm'y gtg'p'q'v'cu'cr r ctgp'0'k'  
y cu'cuwo gf 'y' cv'y' g'htuv'q'v'rkj 'kpetgo gpv'ht'o u'chvgt' { qm'icduqtr v'k'p.'cv'rg'cu'v'6'fc{ u'r quv'  
h'g't'v'k'k' cv'k'p'cpf 'y' tgg'f c{ u'r quv'j cvej . 'ukpeg'rtxcg't'g'ctgf 'k'p'cs w'ct'k'w' 'vq'y' g'uv'ci g'qh' { qm'luce"  
cduqtr v'k'p'uj qy gf 'p'q'g'x'k'f g'peg'q'h'kpetgo gpv'ht'o cv'k'p' "Eqrkp'gv'cr03; ; 9+0'

Nctxcg'qh'Epinephelus striatus ecppq'v'dg'f k'k'p'i v'kuj gf 'ltqo "E. adscensionis" \*t'qen'ij k'pf +"  
o g't'k'v'k'ecm' { cu'eq'w'p'u'cpf 'r ki o g'p'v'k'p'ct'g'p'g'ctn' { k'f g'p'v'ecr0'Dq'v' "Epinephelus striatus" c'pf "E."  
adscensionis" j c'x'g'uo cm' 'uko r ng. 'c'p'f 'u't'ck'i j v'ur k'p'g'rg'w. 'c'p'f 'ecppq'v'dg'ugr c't'c'v'g'f 'ltqo "E. morio"  
\*t'g'f 'i t'q'w' g't+, E. guttatus \*t'g'f 'j k'pf +c'p'f E. drummondhayi \*ur g'em'g'f 'j k'pf +v'p'v'k'if g'x'g'nr o gpv'qh'  
y' g'c'p'c'n'k'p'ku'eq'o r ng'v' "T'k'ej ctf u'gv'cr04227+0"

"Larval distribution and recruitment0" P cuucw'i t'q'w' g't'rtxcg'ct'g't'ct'gn' { t'g'r q't'v'g'f 'ltqo "  
q'h'uj q't'g'y cv'gtu"Ngku'3; ; 9+c'p'f 'r'k'w'g'ku'np'qy p'q'h'y g'k' "o q'x'g'o gpv'qt'f k'v'k'k'w'k'p.'q'y' g't' 'y' cp'  
r'ko k'g'f 'f'c'v'q'p'ugwgo gpv'r cv'gt'pu0'Chvgt'c'o gcp'57/62'fc{ 'r grci le 'rtxcn'r gtlkf . 'rtxcg't'get'v'k'  
ltqo 'cp'q'eg'c'p'le'g'p'x'k't'q'p'o gpv'k'p'v'q'f go g'tu'cn' 'd'c'p'n'ij c'd'k'c'w'v'y' t'q'w' j 'v'k'f c'n'ej c'p'p'g'u'Eqrkp'3; ; 4+0'  
Vj k'u't'get'w'k'o gpv'r t'q'eg'u'ec'p'd'g'd't'k'g'h'c'p'f 'k'p'v'p'ug. 'c'p'f 'ku'cr r ct'g'p'v' { f't'k'g'p'd' { r t'g'x'c'k'k'p'i 'y' k'p'f u."  
ewt'g'p'w. 'c'p'f 'h'w'p'ct' r j c'ug' "Uj gpmgt'gv'cr03; ; 5+0"Rgrci le 'rtxcg'y' gtg'eqmgvxf '20 /38'no "q'h'i'Ngg"  
U'q'en'k'p'i "K'ur'c'p'f . 'D'c'j co cu. 'c'v'p'k'i j v. 'cv'4/72o 'f' g'r y' u'c'p'f 'ltqo 'v'k'f c'n'ej c'p'p'g'u'g'c'f k'p'i 'q'p'v' 'y' g'  
G'z'w'o c' 'D'c'p'n'if w'k'p'i 'y' g'f c { "i tggpy qqf '3; ; 3+0"J qy g'xgt. 'y' g'k'p'n'id'g'y ggp'ur cy plpi 'uk'g'u'c'p'f "  
ugwgo gpv'uk'g'u'ku'p'q'v'y g'm'w'p'f g'tu'v'q'q'f 0'Nctxcn'luco r r'k'p'i 'c'f l'c'eg'p'v'q'c'ur cy plpi 'ci i t'gi cv'k'p'cv'  
O'c'j c'j w'cn' "O'g'z'k'eq' "X" us w'g' / [ g'q'o c'p'u'gv'cr03; ; ; +h'c'k'g'f 'vq'ecr w'g'g'x'g'p'q'p'g'P cuucw'i t'q'w' g't'  
rtxcg0'D { 'y' c { 'q'h'g'z'r n'c'p'v'k'p.'y' g'c'w'j q'tu's w'g'u'k'q'p'g'f 'd'q'y' 'y' g'k' "o g'y' q'f q'm'i { "c'p'f 'y' g'  
t'q'd'w'p'g'u'q'h'y' g'h'q'ec'n'ur cy plpi 'cu'c'f f'k'k'q'p'c'n'g'z'r n'c'p'v'k'p'u0'

Vj g'i g'q'o q't'r j q'm'i { 'q'h'ur cy plpi 'uk'g'u'j cu'rg'f 't'g'ug'c't'ej g'tu'v'q'cu'wo g'y' cv'q'h'uj q't'g'  
v'c'p'ur q't'v'y cu'c'f g'uk't'cd'ng'r' t'q'r g't'v' { 'q'h'ug'ng'v'g'f 'uk'g'u0'J qy g'xgt. 'ewt'g'p'w'k'p'y' g'x'k'k'p'k'v' { 'q'h'  
ci i t'gi cv'k'p'uk'g'u'f q'p'q'v'p'g'eg'u'c't'k'n' { h'c'x'q't'q'h'uj q't'g'gi i 'v'c'p'ur q't'v'ng'c'x'k'p'i 'q'r g'p'y' g'r qu'k'd'k'k'v' { 'y' cv'  
u'q'o g'uv'q'em'i'ct'g'cv'rg'cu'v'r c't'v'k'm' { u'g'r'h't'get'w'k'k'p'i 0'H'q't'g'z'co r ng. 'f' t'q'i w'g'u' "h'q'c'w'u'y' j k'ej 'f' t'k'h'y' k'y' "

"  
"  
"  
"

y cvgt "ewttgpw+f gr nq { gf "pgct"vj g'r qkp'qh'i co gvg'tgngcug"cv'gcuvgtp"Nqpi "Kucpf."Dcj co cu." o qxgf "hkwg"htqo "vj g'uj gnh'gf i g'ht"ugxgtcnf c { u'ko o gf kvgn { hqmqy kpi "ur cy plpi "cpf "qpg"gpf gf " wr "kpuj qtg"Eqnk"3; ; 4+0"K'uko kct"uwf kgu"ctqwpf "c"ur cy plpi "ci i tgi cvkq"ukg"cv"Nkwg"Ec { o cp." uwthceg"xgnqkx { 'r tqhkg"ftkhgtu'tgngcugf "qp"vj g'pki j v'qhr gem'ur cy plpi "uj qy gf "uki pkkcepv'gff { " hqto cvkq"uq"vj cv'f tkhgtu"vgpf gf "v"tgo clp"pgct"qt"tgwtp"v"vj g'ur cy plpi "tgg'h'dw'f tkhgtu" tngcugf "qp"vj g'f c { u'r tgegf kpi "vgpf gf "v"b qxg"cy c { "k"o qtg'qh'c"utcki j v'rkpg'y kj "vj g'f qo kpcpv' ewttgpw"J gr r gm'gv'cr04233+0"Cf f kkpncit'gugctej "ku'pggf gf "v"wpf gtucpf "vj gug'ur cvkn' f { pco leu0'

F cw"qp'tgetwko gpv'qh'rtxcg"qp"v"tgghu'umi i guv'vj cv'vj gk"qp"uj qtg'tcpur qt'v'ecp'tgn { j gcxkn { "qp"etquu/uj gnh'y kpf u'cpf "ewttgpw"cpf "qeewtu"kp"uj qt'v'r wngu'f wtkpi "j ki j n { 'rko ksgf " r gtlkf u'gcej " { gct"Uj gpngt'gv'cr03; ; 5+0"tgetwko gpv'qh'P cuucw'i tqwr gt'rtxcg"qeewtu"cv'cp" cxgtci g'qh'54"o o "VN"Gi i nguq"3; ; 7+cpf "y cu'o qpkq"tgf "ht"c'97/f c { 'r gtlkf "htqo "o kf / F gego dgt"vj tqwi j "Hgdwtct { "wukpi "ej cpgn'p'gwu'uwur gpf gf "k"v'kf cn'r cuugu'dgy ggp"ku'c'p'f u'qp"vj g" gf i g'qh'vj g"Gzwo c"Uqwpf."Dcj co cu0"Cuuwo kpi "vj cv'vj g'hwn'tgetwko gpv'y kpf qy "y cu'uco r ngf." : 8' "qh'vj g'v'cn'ppwcn'tgetwko gpv'qh'P cuucw'i tqwr gt'qeewt'gf "k"vj ku'ctgc'f wtkpi "c"ukpi ng'6/f c { " uqto ."y j kg"cpqj gt'32' "tgetwkgf "f wtkpi "c"ugeqpf "uqto "gxgpv0"F wtkpi "vj g'uco r rki "r gtlkf." 35' "qh'cm'rtxcg"uco r ngf "y gtg"p cuucw'i tqwr gt."y j lej "tgetwkgf "f wtkpi "r ct'kwctn { "uj qtv." f kuetg'v'r wngu'y j gp'eqo r ctgf "v"q'qj gt'vcz'c'ngp"vj tqwi j qw'vj g'uwf { 0"Y j kg"gtcn { "tgetwko gpv' qeewtu"kp"v"dqj "eqtcn'o cetqcn' cg"cpf "ugci tcuu'dgf u."uwdugs wgpw { "j ki j gt"cdw'pf cpegu"kp"eqtcn' o cetqcn' cg"ctg'r tqdcdn { "f wg'v"q"c"eqo dlp'cvkq"qh'c'v'kxg"ugng'v'kq"ht"eqtcn'o cetqcn' cg"cpf "j ki j " r quv'ugwgo gpv'r tgf cvkq"kp"ugci tcuu"p cf gw'cpf "Gi i nguq"3; ; 8+0"

**1.b. Taxonomy and distinctive characteristics**

- Phylum: Ejqtf cv""
- Class: Cevkqr vgt { i k""
- Order: Rgtekhqto gu""
- Family: Ugttcpkf cg""
- Subfamily: Gr kpg'j grkpcg""
- Genus: *Epinephelus* "
- Species: *striatus* "




Figure 6. Nassau grouper adult

Tgegpvi' ppg'v'c'z'q'p'q'o { 'umi i guv'vj cv'hc'o kn { 'i tqwr kpi u'o c { "dg"ej cmgpi gf "Ectki "cpf " J cvkpi u'4229."Ectki "gv'cr04223+cpqpgj grguu."rtgxkqu'f guetkr v'kpu'ctg'r t'gug'p'vgf "wp'v'ki" o qf h'k'ec'v'kpu'ctg'y kf gn { "ceegr v'gf 0'

Tgcej kpi "c"o czko wo "uk { g'qh'344eo "6: kp+v'v'cn'ngpi vj "VN+"J wo cpp'cpf "F gmqcej " 4224."H'q'gug'cpf "R'cwn { "4232+cpf "o czko wo "y gki j v'qh'47"mi "J ggo utc'cpf "T'cpf cm'3; ; 5+ "vj g" P cuucw'i tqwr gt "ku'qpg'qh'vj g'rti gt "ugt'c'pkf u'qh'vj g'v'qr k'cn'Y guv'gtp "C'v'c'p'k'c'cpf "E'ct'k'd'g'c'p'f " ecp'h'x'g'ht'p'g'c'tn { "5" f ge'cf gu0"Uko kct "v"b cp { "qj gt "i tqwr gt "ur geku."P cuucw'i tqwr gt "l'w'g'p'k'gu"

"  
"  
"  
"

cpf "cf wnu"ctg"mpqy p"htq"vj gk"rti g"i cr gu"cpf "r tqv"wf kpi "lcy u."y j kej . "y j gp"qr gpgf "ter kf n{ ." r tqf weg"uwv"vqp"vj cv"lcek"kcvgu"hgdf kpi 0"Vj g"P cuucw"i tqwr gt"ecp"dg"f kpkpi wkuj gf "Itqo "qyj gt" i tqwr gt"u"cv"cm"l"hg"j kxqt { "r j cugu"d { "vj g"ej ctcevgt"kuke"xgt"vecn"ldct"r cwgtp"cpf "f ctm"i"ucf f ngö" eqm"tcv"vqp"cm"pi "vj g"f qtucnr ctv"qh"vj g"ecwf cnr gf wperg0""

Uo kj "3; 93+"kf gp"v"khgf "cp"ö"Epinephelus striatus"Ur gekgu/I tqwr ö"eqo r tkugf "qh"E. striatus."E. guttatus"\*t"gf "j kpf +."E. morio"\*t"gf "i tqwr gt-<"ö"E. striatus"cpf "E. guttatus"ctg"uq"uko krc"t "vj cv"uwp/drgcej gf "f kur m { "ur geko gpu"ctg"f khl"ewv"vq"kf gp"v"kh { "cmj qwi j "vj gtg"ctg"ugxgtcno g"tkuke" ej ctcevgtu"vj cv"ecp"dg"uwgf "vq"ugr ctcvg"vj go 0"T gf "i tqwr gt"\*E. morio+"f khtgtu"lp"l"kp"q"wnkpgu."dw" qyj gty kug"utqpi n{ "tgugo drgu"vj g"qyj gt "vy q"ur gekgu0"Vj g { "ctg"cm"i"o qf gtcvgn { "rti g"l"kuj gu"y kj " vr gt"lpi "cpf "uqo gy j cv"eqo r tguugf "dqf { "q"wnkpgu0"T gf "i tqwr gt"j cu"ur qwgf "cpf "dctt"gf "t"cpul"gpv" eqm"t"r j cugu"cpf "kpf kxk" wcu"l"p"vj gug"eqm"t"r j cugu"dgct"c"tgo ctn"æ"dr"tgugo drpe"p"vq"P cuucw" i tqwr gt0"Vj ku"l"u"tgh"gevgf "lp"vj g"Dgto wf cp"eqo o qp"pco g"f ggt"j co ngv"ht"E. morio, eqp"t"cu"l"pi " y kj "j co ngv"\*y kj qw"v"o qf kht"gt+"htq"E. striatus0"Vj gug"vj tgg"ur gekgu"ctg"egt"v"l"p"n { "emug"vq"ge"cej " qyj gt"cpf "y gm"l"ugr ctcvgf "Itqo "qyj gt"Co g"t"kecp"i tqwr gt"u"ö"

Vj g"l"qm"y kpi "f guetk"v"qpu"ctg"dcugf "r tgf qo kpcpv"n { "qp"Uo kj "3; 93+"Cegt"q"gv"cr"0"3; ; 3+" cpf "J ggo utc"cpf "T"cpf cm"3; ; 5+"cu"r tgu"p"vgf "lp"Ucf qx { "cpf "Gm"wpf "3; ; ; <"

Vj g"P cuucw"i tqwr gt."Epinephelus striatus"\*Dm"ej "39; 4+"ku"v"o qf gtcv"v"uk" gf "Epinephelus" y kj "rti g"g { gu"cpf "c"tqdw"v"dqf { 0"Dqf { "f gr vj "ku"l" kpkpe"v"n { "nguu"vj cp"j" gcf "ngpi vj . "f gr vj " eqpv"l"p"gf "408"vq"40 "vko gu"l"p"UN"\*ht"l"kuj "382"vq"552o o "UN+0"J gcf "ngpi vj "ku"eqpv"l"p"gf "406"vq"408" vko gu"l"p"UN="lpvgt"qtdkcn"l"eqpxgz="r tgqr gteng"gxgpn { "ugttcv"v."y kj qw"l"ucn"l"gpv"v"cp"i ng="r quvgt"kt" pqu"tku"uqo gy j cv"p"rti gf "cpf "gm"pi cvgf "qt"eqo o c/uj cr gf "lp"rti g"cf wnu0"I tqwpf "eqm"t"ku" i gpgtcm { "dwh"y kj "7"f ctm"i"dtqy p"xgt"vecn"ldct"u"cpf "c"rti g"dr"cn"l"ucf f ng"dm"vej "qp"vq"r"qh"ecwf cnr" r gf wperg="c"tqy "qh"dr"cn"l"ur qu"dg"ny "cpf "dgi kpf "g { g0"F kpkpe"v"v"kg" f ctm"l"wp"l"pi /ht"m"i"o ctm"i" dgi kpp"l"pi "cv"ht"qp"v"qh"vr r gt"lcy . "gz"v"p"l"pi "f qtucm { "cm"pi "lpvgt"qtdkcn"l"gi kpp."cpf "dkh"t"ecv"l"pi "qp" vq"r"qh"j" gcf "dgi kpf "vj g" { gu="cp"qyj gt" f ctm"i"dc"p"l" Itqo "vr"qh"up"q"v"vj tqw"j "g { g"cpf "vj gp"ewt"xl"pi " wr y ctf "vq"o gg"v"ku"l"gm"y "l"uw"v"dg"htg" f qtucn"l"p"q"tki kp0"Uqo g"l"kuj "j cxg"l"t"gi wct"r cng"ur qu"v"cpf " dm"vej gu"cm"l"qxgt"vj g"j gcf "cpf "dqf { "y j kng"ur geko gpu"ht"qo "f ggr"y cvgt"ctg"uqo gy j cv"r kpn"kuj "qt" tgf f kuj "xgp"t"cm { 0"Vj g"l"puk" f g"qh"vj g"o qwj "ku"t"gf . "vj g"v"gg"y "ctg"ecp"l"p"kh"qto "cpf "xkn"kh"qto "cpf "ctg" l"p"v"y q"ugt"l"gu"l"p"ge"cej "lcy "Uo kj "3; 9: +0"Vj g"t"cp"i g"qh"l"eqm"t"ku"y kf g0E"qm"t"r cwgtp"ecp"ej cpi g" y kj l"p"o l"p"wg"u"ht"qo "cm" qu"v"y j kg"vq"d"l"eqm"t"gf "vq"v"p"kh"qto n { "f ctm"i"dtqy p."cee"q"t" f l"pi "vq"vj g" dgj cxk"q"tcn"l"ucv"v"qh"vj g"l"kuj "N"qpi ng { "3; 39."E"q"l"p"3; ; 4."J ggo utc"cpf "T"cpf cm"3; ; 5."Ect"vgt"gv"cr"0' 3; ; 6+0"C" f kpkpe"v"v"kg"d"l"eqm"t"gf "r cwgtp"ku"l"uggp"y j gp"v"y q"cf wnu"qt"cp"cf wv"v"cpf "rti g"l"wxg"p"kg" o gg"v"cpf "ku"l"t"gs wgp"v"n { "qdug"t"xgf "lp"ur cy pl"pi "ci i tgi cv"l"qpu"\*J ggo utc"cpf "T"cpf cm"3; ; 5+0"Q"pn { " f qw"l"ct"q"wpf "vj g" { gu"cpf "vj g"dm"vej "qp"vj g"ecwf cnr gf wperg" f q"p"q"v"ej cpi g"Uo kj "3; 93+0" Lwxg"p"kgu"gz j kdk"v"o"eqm"t"r cwgtp"u"ko krc"t"vq"cf wnu"\*g"0 0"Uk"kc"Ngg"3; 99+0"

"  
"  
"  
"  
"

**1.c. Range wide distribution**

Vj g'P cuucw'i tqwr gtai'eqphko gf 'f kvtkdwkqp'ewttgpvn' 'kpenmf gu'odgto wf c'cpf 'Hqtktf c' \*WUC+'y tqwi j qw'vj g'Dej co cu'cpf 'Ectkddgcp'Ugc'\*Hi 09+'Uo kj '3; 93.'Cegtq'cpf 'I ct| qp/Hgttgkt'3; ; 3.'J ggo utc'cpf 'Tcpf cm'3; ; 5.'Egtxki qp'3; ; 6+0'Vj g'r t gxlqwu'tgr qt v'qh'E. striatus' Itqo 'y g'Dtc| klcp'eqcu'uwj 'qh'y g'gs wcvqt '\*Hi 0636'\*f kvtkdwkqp'o cr +lp'J ggo utc'cpf " Tcpf cm'3; ; 5.'r 0459+'ku'wpuwducpvkvvgf o'\*Etcki 'gv'cr04233+0'Vj g'P cuucw'i tqwr gt'j cu'dggp" f qewo gpv'f 'lp'y g'y guvtp'I wh'qh'O gzleq.'v'y g'y guv'qh'y g'l wecvp'Rgpkpuwr.'O gzleq.'cv' Cttgk'g'Crcetcpgu'\*pqt vj 'qh'Rtqi tguq+'J krf gdtcpf 'gv'cr03; 86+0'K'y cu'ekgf 'cu'c'tctg'qt" vcpukgp'ur gelgu'lp'y g'pqt vj y guvtp'I wh'qh'O gzleq.'qhh'Vgzcu'\*I wvgt'cpf 'Mpcr r '3; 73'lp" J qgug'cpf 'O qqtg'3; 99+0'Hqrg{ 'gv'cr0\*4229+'tgr qt v'f 'y g'htuv'r j qvqi ter j gf "cpf 'eqphko gf " uki j vpi 'lp'y g'Hqy gt'I ctf gp'Dcpm'P cvkqpcn'O ctkpg'Ucpewct { .y j lej 'ku'hqecv'f 'lp'y g' pqt vj y guv'I wh'qh'O gzleq.'cr r tqzko cvgn{ '3: 2'no 'uqwj gcuv'qh'I crk guvq. 'Vgzcu'P cuucw' i tqwr gt'ku'i gpgtcm{ 'tgr megf 'geqni kcm{ 'lp'y g'gcvgt'P wh'd{ 'Epinephelus morio'\*Uo kj " 3; 93+lp'ctgcu'pqt vj 'qh'Mg{ 'Y guv'qt 'y g'Vqtwi cu'O cp{ 'qh'y g'gctrkt'f guetk vqpu'gzv'f 'y g' tpci g'wr 'y g'Cvcpv'e'eqcu'v'P qt vj 'Ectqkpc.'dw'eqphko cvkq'ku'ewttgpvn' 'rcenkpi 0"

Vj g'P cuucw'i tqwr gt'ku'rkv'f 'cu'op cvkxgo'v'y g'hqmy kpi 'eqwv'kgulucv'g'\*Eqtpkuj 'cpf " Gmwpf "42250+ Cpi wkr=Cpki we'cpf 'Detdwc=Ctwc=Dcj co cu=Detdcf qu=Dgrk g=Dgto wf c=" Ec{o cp'Krcpf u=Eqmo dlc=Eque'Tlec=Ewdc=Ewtc±cq=F qo kpec=F qo kpecp'Tgr wdre=Ht gpej " I wkpc=I tgpfc=I wcf gnqr g=I wcvgo cr=I w'cpc=J ckk=J qpf wcu=Lco clec=O gzleq=" O qpvttecv=P gj gtrpf u'Cpvrku'\*Ewtc±cq=P lectci we=Rcpco c=Rwgtv'Tleq=Uckp'v'Mku'w'cpf " P gxku=Uckp'v'Nwec=Uckp'v'Xkpegp'v'cpf 'y g'I tgpfc kpu=Uwtlpc g=Vtkp'cf 'cpf 'Vqdc q=Vwmu' cpf 'Eclequ'Krcpf u=Wpkgf 'Ucv'g'\*Hqtktf c=Wpkgf 'Ucv'g'O kqt 'Qwn{ kpi 'Krcpf u'Ectkddgcp<




Figure 7. Range of Nassau grouper (*Epinephelus striatus*). Habitat zones include shoreline to insular or continental shelf throughout the indicated range.

KQ0'P cxcuuc'Krcpf +=  
Xgpgj wgr=Dtkskj 'Xkti kp"  
Krcpf u=WUOXkti kp"  
Krcpf u0"  
"

**1.c.i. Historical Distribution**

P cuucw'i tqwr gt" qv'rkj u'j c'xg'dggp" t'gt'kxgf 'Itqo 'c'xctk'v{ " qh'ukgu'no kf f gpu+'lp" r tgj kv'qtle'hkj kpi " eqo o w'pkkgu'qh'y g" Ectkddgcp.'cpf 'y g" ur gelgu'tgr t'gug'v'f 'cp"

ko r qt'wv'eqo r qp'gp'v'qh"

"  
"  
"  
"

vj gug'eqo o wplkgu0"Qvqrkj u'y gtg'tgrvkggnl "cdwpcf cpv'cv'ukgu"qp'gcuvgtp'CPvki wc"\*CF '722/3372+."kp'I tgpfc c"\*CF '2/722.'CF '3222/3722+."Ucp"Ucrkcf qt.'Dej co cu"\*CF ': 72/3322+."U0' Lqj p.'Xki kp"Kucpf u"\*CF '922/3422+."Hrktkf c'y guv'eqcu"\*4222/3222"DE.'CF '372/522.'CF '622/3222+."U0Nvekc.'Y guv'Kpf lgu"\*CF '2/3722+cpf "qp"vj g'pqt vj "eqcu'qh'Lco ckec"\*pq'f cvg+\*Y kpi "gv' cr03; 8: "cpf "Y kpi "cpf "T gkl "3; : 4."cu'ekvgf "kp"Ucf qx {"cpf "Gmwpf "3; ; ; +0'

**1.c.ii. Influences on Distribution**

Rtlo ct { 'f gvgto kpcpw'qh'f kvtkdwkqp'kp'P cuucw'i tqwr gt'ctg'pqv'hpqy p'cmj qwi j 'y cvgt' emtkv\.'j cdkcv.'cpf "uwdutcvg'v{r g'er r gct "v"dg'ko r qt wcpv"\*Uo kj "3; 93."Gi i nguqp"3; ; 7+0Vj ku' ur gekgu'ku'o quv'cdwpcf cpv'kp'engct'y cvgtu'qp"j ki j /tgrkgh'eqtcn'qt'tqem\ 'tgghu0Uo cm'lwxgpkngu'ctg' cuuqekcvf 'y kj "o cetqcn' cg.'ugci tcuu'dgf u.'qt"Porites'emo r u0Vj g'o gcp'f gr vj 'tapi g'qh'vj g" P cuucw'i tqwr gt"\*2/352'o +0' c { "dg'kphwvpegf "o qtg'd { "vj g'cxckrdkrk\ "qh'uwxkdrj'j cdkcv'vj cp'd { " hqf "tguqwtegu.'ukpeg'f kgv'ku'j ki j n\ "xctkgf "cpf "o qtg'c'hwpev'kp'qh'dqf { "uk' g'vj cp'qh'y cvgt'f gr vj 0' F gur kg'cf wmu'o ki tvkpi "mpj 'f kucpegu'v'g'cej "ur cy plpi "ukgu"\*Uctt'gv'cr04229+." r tqzko kv\ "v"vj gug'ukgu'f wtkpi "pqp/tgr tqf wvkg'g'r g'kqf u'ku'cr r ctg'pvn\ "pqv'etk'ecn'cmj qwi j "vj g" ci i tgi cvkqp'ukgu'vj go ugrkgu'o c { "dg'guugpv'kcn'ht'g'r tqf wv'kqp'g'k'j gt'dgecwug'qh'r j { ukecn' ej ctcev'gt'k'ku'qh'vj g'uwdutcvg'qt'dgecwug'qh'vj g'qegcpqi tcr j ke'eqpf k'k'qpu'cv'vj g'ukg0'Vj g'muu'qh' m'ecn'uqem'kp'c'pwo dgt'qh'k'puwrt'ctgcu"\*g'f 0'Dgto wf c'cpf "Rwgtvq'Tleq'+uwi i guv'vj cv'uo g" r qr wv'k'qpu'ctg'r ct'k'cn\ "ugr/tget w'k'k'pi ."cmj qwi j "hwt vj gt' i g'p'g'ke'uwf lgu'ctg'p'geguuct { "v'v'gu'vj ku' j { r qvj guku"\*Ucf qx { "3; ; 5+0'

**1.d. Biological characteristics**

**1.d.i. Age, growth and mortality**

" I tqy vj "kp'P cuucw'i tqwr gt'j cu'dggp"gzco kpgf "d { "uk' g'htgs w'g'pe { "cpcn\ ugu."ci i kpi " uwf lgu.'h'grf "qdugt'x'cv'k'pu."cpf 't'gcf kpi "cppwrt' t'kpi u'kp'uci kvcn'qvqrkj u"\*Vcdrg'5.'Hki 0: +0'O quv' uwf lgu'k'p'f kecv'g'ter kf "i tqy vj ."cdqw'32o o lo qp'vj "h'qt'uo cm'lwxgpkngu0'O gcp'o qp'vj n\ "i tqy vj "qh' P cuucw'lwxgpkngu"52/492"o o "VN"qp'ct'v'k'ecn'cpf "pcwrt'cn'tgg'hu'kp'vj g'Xki kp"Kucpf u'y cu": 06"vq" 330"o o lo qp'vj ."f gvgto kpgf "f wtkpi "uk'x'kuwcn'le'g'puwugu'qxgt"33"o qp'vj u."\*Dggv'cpf "J kzqp"3; ; 6+0' Uko k'ctn\ ."lwxgpkngu'uco r ngf "cv'Ngg'Uqen'kpi "Kucpf "kp'vj g'Dej co cu'i tgy "cv'cdqw'32"o o lo qp'vj " dgw'ggp"54"cpf ": 7"o o "VN"\*Gi i nguqp"3; ; 7+0P gct "ugz wcn'o cwtkv\ "cv'cdqw'6/9" { gctu."P cuucw' i tqwr gt' i tqy vj "umy u'v'q'cdqw'4o o lo qp'vj ."y kj "m'y gt'tcv'gu'kp'rti gt'qt'ugz wcn\ "o cwtg'h'kuj " \*Dwuj "gv'cr04228+0"

O cti kpcn'k'petgo gpv'cpcn\ uku'qh'uci kvcn'qvqrkj u'wui i guvgf "vj cv'i tqy vj "l' qpgu'y gtg'hqto gf " c'ppwcm\ "cpf "vj cv'c'ppwcn'k'petgo gpv'f gr qukv'kp'q'ee'wt'gf "htqo 'Cr tkn'v'q'O c { "kp'Ewdc"\*Emtq'gv'cr0' 3; ; 2+0'Vj g'i tqy vj "l' qpgu'f gr qukv'gf "kp'qvqrkj u'y gtg'x'cn'f cvgf "cu'c'ppwcn'w'k'k'pi "qz { v'g't'ce { er'k'p'g" \*QVE+0' ctn'kpi "v'gej pl'k' w'gu=q'v'qrkj "ngi k'k'k'k'v\ "y cu'cr r tqzko cvgn\ ": 2/; 7' " \*Dwuj "gv'cr03; ; 8+0'

F cw'htqo "uecngu'cpf "qvqrkj u'k'p'f kecv'g'vj cv'h'kuj 't'g'cej "622/672"o o "UN"\*k'g'0'ugz wcn' o cwtkv\ +kp'cr r tqzko cvgn\ "6/9" { gctu0"J qy g'x'gt."gu'ko cv'gu'qh'uk' g'cv'ci g'f g'k'x'gf "htqo "ngpi vj /"

"  
"  
"  
"

It gs wpe { "f cvc " \*Vcdng " 5 + uwi i guv'o qtg'tcr kf "i tqy vj " \*Qmrgp " cpf " NcRrceg " 3 ; 9 ; + 0 " Vj ku'cr r ctgpv " f kuetgr cpe { " dgvy ggp " qvqkij / " cpf " rgi vj / dcugf " o gj qf u'qh'ci g " f gvgto kpcvkqp " eqwrf " tguwn / Itqo "

**Table 3. Size at age data for Nassau grouper (from Sadovy and Eklund 1999; "Bush et al., in press" should refer to Bush et al. 2006)**


*Epinephelus striatus* size-at-age data for ages 1-13 years. All lengths are in mm (standard/SL or total/TL lengths as indicated). Ageing method is given.

Source	Age (years)												
	1	2	3	4	5	6	7	8	9	10	11	12	13
a	293	354	390	464	537	561	634	659	-	-	-	-	-
b	160	270	-	410	480	540	570	600	640	650	660	700	710
c	175	253	309	358	401	436	468	497	519	542	563	580	591
d	174	254	315	366	414	451	483	518	559	583	594	617	-
e	235	370	435	500	543	605	660	720	760	800	-	-	-

a: Buesa, 1987; Cuba, aged by scales/TL.  
 b: Bush et al., in press; Cayman Islands, aged by otoliths/TL lengths estimated from observed growth curve.  
 c & d: Claro et al., 1990; SW and NE Cuba, respectively, fish from both areas were aged by otoliths and backcalculation/TL.  
 e: Olsen and LaPlace, 1979; Jamaica, aged by length-frequency data/SL.

vj g "wpcxckrdkx { "qh'ci g "ercuu/3 " kpf kxf wcu. "t guwnkpi " kq'qrf gt " \*kqo " ci g " 4 - " { gct u + " kpf kxf wcu " f guki pcvgf " cu'ci g " 3 " { gct "ercuu " \*Ucf qx { " cpf " Gmwpf " 3 ; ; ; + 0 " O qtgqxt. " rgi vj " It gs wpe { " cpcn { uku " ecp " dg " rguu " t grkcdng " hqt " rgi vj / rkgf " ur geku " vj cp " qvqkij / dcugf " uwf kgu " cu " qrf gt " eqj qt wu " uqap " dgi kq " vq " o gti g " kvq " gcej " qy gt " qdewtkpi " kpf kxf wcu " ci g " ercuugu 0 "

"Xqp " Dgt wcrph { " i tqy vj " r cto gvgtu " f gkxgf " hqt " vj g " P cuuw " i tqw gt " y kj " vj g " Dtqf { " i tqy vj " eqghkelpv " \*M " t cpi g " Itqo " 2085 / " 208 : 7 " \*Vcdng " 6 + 0 "



**Figure 8. Growth curve for Nassau grouper sampled from aggregations between 1987 and 1992 in the Cayman Islands (from Bush et al, 2006)**

"  
 "  
 "  
 "

**Table 4. Von Bertalanffy growth equation parameters for Nassau grouper**

Von Bertalanffy growth equation parameters for Nassau grouper, *Epinephelus striatus*. Standard equation for length-at-age is:  $L_t = L_\infty (1 - e^{-K(t-t_0)})$ . Lengths are in cm (length type indicated).

Source & method	Locality	Length type	Parameter		
			$L_\infty$	$t_0$	K
Olsen & LaPlace, 1979	Virgin Islands	SL	97.4	0.488 <sup>1</sup>	0.185
Claro et al., 1990	Cuba (SW)	TL	94.0	-3.27	0.063
	Cuba (NE)		76.0	-1.12 <sup>2</sup>	0.127
Thompson & Munro, 1978	Jamaica	TL	90.0 <sup>3</sup>	-	0.090
Baisre & Pérez, 1981	Cuba	-	92.8	-	0.100

<sup>1</sup> Appears also as  $t_0 = -0.488$  in some places.

<sup>2</sup> Appears also as -4.13 in Abstract.

<sup>3</sup>  $L_\infty$  assumed, based on tagging data from Randall, 1962, 1963.

I tqy vj 'tcvgu'y gtg'cnuq'f gvgto kpgf 'kp'hkgrf'qdugtxcv'kqpu'cpf'vci i kpi 'uwx'kgu'0'kp'yj g' Xkti kp'Krcpf u.'cplk'cm'vci i gf'hqt'nguv'y cp'522'f c{u'{'kgrf gf'yj g'hqmy kpi 'i tqy vj 'tcvgu'2397/' 472'o o 'VN'i tgy 'cdqw'60/7'o o lo qpjv '=473/'547'o o 'VN'cdqw'50'o o lo qpjv '=548/673'o o " VN'cdqw'30 4'o o lo qpjv '\*Tcpf cm'3; 84."

**Table 5. Age and size parameters for Nassau grouper. [excerpt from Sadovy and Eklund 1999: Bush et al, in press refers to Bush et al, 2006, CFMC footnote 26 refers to CFMC 1985, SAFMC footnote 24 refers to SAFMC 1983]**

<i>Epinephelus striatus</i>	
Age and length at maturity	5 yrs, 580 mm SL (Virgin Islands) (Olsen and LaPlace, 1979)  420-450 mm SL females 400-450 mm SL males 4+ yrs (Cayman Islands) (Colin et al., 1987; Bush et al., in press) 500 mm TL (minimum size ripe males) (Cayman Islands) (Tucker et al., 1993) 425 mm SL females; 402 mm SL males, immatures are 3-6 yrs (otolith growth zones not validated) (Bahamas) (Sadovy and Colin, 1995) 483 mm TL (North Carolina-Florida) (SAFMC, text footnote 24) 480 mm TL (Jamaica) (Thompson and Munro, 1978)
Age and length at first capture	< 300 mm TL & 4-5 yrs (Virgin Islands) (Olsen and LaPlace, 1979; CFMC, text footnote 26) 6-7 yrs (Cayman Islands) (Bush et al., in press) 275-625 mm TL (mean = 570) (Jamaica) (Thompson and Munro, 1978) 300-500 mm TL depending on size limits (North Carolina-Florida) (SAFMC, text footnote 46) 450 mm TL (South Florida) (Bohnsack, 1990)
Maximum age and length	1200 mm TL (CFMC, text footnote 26) 9 yrs, 910-960 mm SL (Olsen and LaPlace, 1979) (Virgin Islands) 17 yrs, 710 mm TL, 6700 g (Cuba) (Claro et al., 1990) 755 mm SL (Bermuda) (Bardach et al., 1958) 840 mm TL (Jamaica) (Thompson and Munro, 1978) 640 mm TL (Netherlands Antilles) (Nagelkerken, 1981) 29 yrs, 850mm FL (Cayman Islands) (Bush et al., in press)

3; 85."Vcdrg": 'kp'Ucf qx{'cpf'Gmwpf'3; ; ; +0" J qy gxgt.'i tqy vj 'tcvgu'y gtg'gxkf gpw'f wpf gtgunko cvgf 'dgecwug'qh'i tqy vj 'uwr r tguukqp' f wg'vq'vci i kpi '\*Vj qo r uqp'cpf'O wptq'3; 9: +0" Hkuj 'yj cv'tgo ckpgf 'kp'yj g'hkgrf'hqt'535'vq'959" f c{u'j cf'j ki j gt'i tqy vj 'tcvgu.'xct{ kpi 'htqo '6" vq'808'o o lo qpjv 'hqt'hkuj 'kp'yj g'478/'5: 2'o o " VN'uk'g0'I tqy vj 'kp'P cuucw'i tqwr gt'y cu'cnuq" o gcuwtgf 'd{'ecrewr'vki 'y gli j v'kpetgo gpw'qh" o ctngf'hkuj 'kp'yj g'hkgrf '<y gli j v'kpetgcug'hqt'9" kpf kxf wcu'kp'yj g'922'i 'uk'g'eruu'y cu'42/" 72' 'r gt'{'gct'y kj 'cp'cxgtci g'qh'5: ' " \*Dctfcej 'cpf'O gpl'gn'3; 79+'j qy gxgt.'yj g' cwj qtu'awi i guvgf 'c'f genkp'kp'i tqy vj 'tcv' chngt'lcy 'vci u'y gtg'cr r rkgf'y j gp'f cw'y gtg" eqo r ctgf 'vq'f ct/vci i gf'hkuj 0'Ci g'uk'g" r ctco gvgtu'ctg'r tguvgf 'kp'Vcdrg'7'cpf " ngpi vj /y gli j vtgr'vqpuj kr u'hqt'wcpf ctf.'vqven" cpf'hqtn'ngpi vj u.'cpf'VN/UN'tgr'vqpuj kr u'ctg" uj qy p'kp'Vcdrg'80"

"  
"  
"  
"

**Table 6. Length-weight and length-length conversion parameters for Nassau grouper (excerpt from Sadovy and Eklund 1999)**

Length-weight and length-length parameters for *Epinephelus striatus* and *E. itajara*. The length-weight relationship is defined as:  $W=aL^b$ , where L is length (standard/SL, fork/FL or total/TL, as indicated) in mm and W is body weight in grams (guttated, G, or unguttated, UG, as indicated- where known). The standard length-total length relationship is defined as:  $TL=a+bSL$ .

Locality	Restrictions <sup>1</sup>	Parameter		Source
		a	b	
<b>Length-weight</b>				
<i>Epinephelus striatus</i>				
Virgin Islands	180-760 (SL) n=241	0.0097	3.23	Olsen and LaPlace, 1979
(St. Thomas/St. John)	330-770 (SL) n=73	$1.43 \times 10^{-6}$	3.38	Bohnsack and Harper, 1988 (UG)
Puerto Rico	210-645 (FL) n=60	$1.26 \times 10^{-5}$	3.04	Bohnsack and Harper, 1988 (UG)
Cuba (NE)	n=75 (TL)	0.1980	2.98	Ciaro et al., 1990
Cuba (SW)	n=270 (TL)	0.0052	3.30	Ciaro et al., 1990
Jamaica	325-825 (TL) n=112	0.0107	3.11	Thompson and Munro, 1978 (UG)
Belize	180-802 (SL) n=930	0.0107	3.08	Carter et al., 1994 (UG implied)
Florida	203-516 (TL) n=9	$3.8 \times 10^{-6}$	3.23	Bohnsack and Harper, 1988 (UG)
Bahamas	174-724 (SL)	$2.14 \times 10^{-5}$	3.03	Sadovy and Colin, 1995 (UG)
<b>Total length-standard length</b>				
<i>Epinephelus striatus</i>				
Cuba	n=330	2.24	1.11	Ciaro et al., 1990
Jamaica	430-750 n=26	3.00	1.09	Thompson and Munro, 1978
Bahamas	174-695 n=33	2.81	1.13	Sadovy and Colin, 1995

<sup>1</sup> Restrictions are upper and lower sizes in samples analyzed.

*Longevity* "Vj g' b czko wo 'ci g'tgeqtf gf 'hqt' P cuucw'i tqwr gt 'ku'4; '{ gctu.'wukpi 'uci kwcn' qvqrkj u'htqo 'y j g'Ec {o cp'Kurcpf u'\*Dwuj 'gv'cn03; ; 8.'4228+'\*Hki 0: +0'Wukpi 'rgpi yj /htgs wgpe { " cpcn {uku.'y j kej 'vgpf u'vq'gzemf g' {qwp i gt'cpko cnu.'c'yj gqtgkccn'o czko wo 'ci g'cv'; 7' "cu {o r vqke" uk' g'ku'38" { gctu0'Qvj gt'o czko wo 'ci g'guko cvgu'kpenf g'kpf kxf wcu'qh'wr 'vq"; '{ gctu'kp'yj g' j gcxkn { "gzr mkgf 'Xkti kp'Kurcpf u'hkuj gt { '\*Qnugp'cpf 'NcRrcg'n 9; +.'34" { gctu'kp'pqtvj gtp'Ewdc.'39" { gctu'kp'uqwj gtp'Ewdc '\*Erctq'gv'cn03; ; 2+'cpf '43" { gctu'htqo 'y j g'Dcj co cu.'cuuwo kpi .'cu" f go qpucvqf 'kp'uqo g'hqecvqpu.'yj cvtkpi u'ctg'htqo gf 'cppwcm { '\*Ucf qx { 'cpf 'Eqrkp'3; ; 7+0'Vj gug" f khtgpegu'kp'o czko wo 'ci g'guko cvgu'ctg'f wg'vq'yj g'uco r ngu'cxckrdng'ht'ci kpi 'cpf " o gvj qf qm i kecni f khtgpegu0'kpf kxf wcu'qh'o qtg'yj cp'34" { gctu'qh'ci g'ctg'pqveqo o qp'kp" hkuj g'kgu.'y j kb'o qtg'j gcxkn { 'hkuj gf 'ctgcu' { kgr kpi 'o wej " { qwp i gt' hkuj "qp'cxgtci g0'I gpgtcvqkq" vko g'\*yj g'cxgtci g'ci g'qh'r ctgpw'kp'yj g'r qr wrcvqkq+'ku'guko cvgf 'cu'; /32" { gctu'dcugf 'qp'cxgtci g' hkuj 'uk' g'htqo 'cp'wpgzr mkgf 'ci i tgi cvkq'kp'Dgnk' g.'y j g'i tqy yj 'ewtxg'htqo 'y j g'hkg'Ec {o cp" Kurcpf 'ur cy kpi 'ci i tgi cvkqpu.'cpf 'y j g'UN/VN'eqpxgtukq'ewtxg'htqo 'Ucf qx { 'cpf 'Eqrkp'3; ; 7+0'

*Mortality rates.* "Guko cvgu'qh'pcwcn'o qtvrk { "O +.'dcugf 'qp'rgpi yj /htgs wgpe { 'f cv'htqo " P cuucw'i tqwr gt 'cngp'qp'wpgzr mkgf 'dcpm'kp'Leo clec. 'tapi gf 'htqo '2089'vq'2052'\*Vj qo r uqp'cpf "

"  
"  
"  
"

O wptq'3; 9: +0"Vqven'ò qtvrk\{\* + 'wukpi 'hgi y 'htgs wpe{ 'f cvc. 'y cu'guko cvgf 'cv'207'lp'Ewdc0  
Y kj 'c'mqy 'pcwten'ò qtvrk\{\*O +f gyto kpgf 'vq'dg'208: . 'y ku'lpf lecvu'c'hkuj kpi 'o qtvrk\{\*H"qh"  
209"\*Dckutg'cpf 'Rcgl '3; : 3+0'  
"

### 1.d.ii. Ecological Roles

*As Prey.* "Kphqto cvkqp'qp'r tgf cvkqp'wr qp'i tqwr gtu'ku'rcti gn\ 'rcenkpi . 'cnj qwi j 'uj ctmi"  
y gtg'tgr qtvgf 'vq'cwceniP cuucw'i tqwr gtu'cv'ur cy plpi 'ci i tgi cvkqu'lp'y g'Xkti kp'Kucpf u"\*Qungp"  
cpf 'NcRrcg'3; 9; +cpf 'y gtg'ku'qpg'tgr qt'v'qh'ecppkdcruo 'lp'y ku'ur gekgu"\*Ukxc'Ngg'3; 96+0"P q"  
r tgf cvkqp'y cu'qdugtxgf "qp'ur cy plpi 'hkuj 'lp'y g'Dcj co cu. 'f gur ksg'y g'r tgupeg'qh'uj ctmi'lp'y g"  
ctgc"\*Eqrkp'3; ; 4+0"Qpg'o wkrvgf 'hkuj 'y cu'tgeqxtgf . 'r quukdn\ 'cwcengf 'd{ 'c'dcttcwfc'qt'uj ctmi'  
hqmqy kpi 'tgrcug'qh'vci i gf . 'rcdqtcvt{/tgetgf . 'pckxg'lpf kxf wcu'qpva'c'tgg'h'lp'y g'Xkti kp'Kucpf u"  
\*Tqdgwu'gv'c'03; ; 7+0"Gctn\ 'r quv'ugwgo gpv'lwgpkrg'r tghgtgpegu'hqt'o cetqci cg'tcy gt'y cp"  
ugci tcuu'dgf u'ctg'r tqdcdn\ 'tgrvgf . 'lp'r ctv.'vq'j ki j gt'rgxgn'qh'r tgf cvkqp'lp'ugci tcuu'dgf u"\*P cf gcw"  
cpf 'Gi i nguqp'3; ; 8+0"Tr gtwa'qh'hkaphkuj 'r tgf cvkqp'qp'uo cm'tggh'hkuj 'cpf 'uo cm'rhg'uci gu'ctg'c"  
eqpegt'y tqwi j qw'y g'Ectkddgcp'cu'y g'lp'xcukxg'ur tgcf 'j cu'y kf gpgf "\*"Cndkpu'cpf 'J kzqp'422: +0'

*As Competitors.* "Nkwg'ku'r wdrkuj gf "qp'gkj gt'kvtc/"qt'kvtg/ur gekhe"eqo r gvkqp'lp"  
P cuucw'i tqwr gt0"Lwgpkrgu"gzj kdk/ci i tguukqp'vqy ctf u'uko kct/uk\ gf "eqpur gekheu'cpf 'f kur r {"  
kvtur gekhe'ci i tguukqp'"L0F wj co . 'Ectkddgcp'O ctkpg'Tgugctej 'Egpgvt. 'e lq'Hqtkf c'Ucvg"  
O ctkpg'Ncdqtcvt{ . 'wpr wdr0tgr qt'v'vq'y g'Ectkddgcp'O ctkpg'Tgugctej 'Egpgvt. 'O ctej '4; . '3; : ; +0"  
Y j gp'y q'pqp/tgr tqf wv'xg'cf wnu. 'qt'cp'cf wv'cpf 'rci g'lwgpkrg. "gpeqwpvt'qpg'cpqy gt. 'y g"  
uo cmgt'hkuj 'ces wktgu'y g'dleqmt'r cwgt'f guetkdgf 'hqt'ci i tgi cvkpi 'hkuj 'lp'cr r ctgpv'uwo kuukqp. "  
y gp'wtpu'rcvgtcm\ "cpf 'wuwcm\ 'uy ko u'cy c{ "\*"Eqrkp'3; ; 4. 'R0Eqrkp. 'Eqtcn'Tggh'Tgugctej "  
Hqwpf cvkqp'o 'Rcruw. 'r gtu0'eqo o 0'vq' 0'Ucf qx{ . 'P O HU.'3; ; 2+0'

*As Predators.* "Vj g'P cuucw'i tqwr gt'ku'c'vqr /rgxgn'r tgf cvqt'qp'eqtcn'tgghu0'P cuucw'i tqwr gt"  
ctg'wpur geknk\ gf /co dwuj /uwekqp'hqtcu'gtu"\*Tcpf cm'3; 87. "Vj qo r uqp'cpf 'O wptq'3; 9: +v'cv'  
uy cmqy 'r tg{ 'y j qrg"\*Y gtpgt'3; 96.'3; 99+0'P wo gtqwu'uwf kgu'f guetkdg'P cuucw'i tqwr gt'cu"  
r kckxqtqwu'cu'cf wnu"\*Tcpf cm'cpf 'Dtqeni'3; 82. 'Tcpf cm'3; 87. 'Tcpf cm'3; 89. 'Rcttkuj '3; : 9. 'Ectygt"  
gv'c'03; ; 6. 'Gi i nguqp'gv'c'03; ; : +0"Vj ku'ur gekgu'vcngu'o cp{ 'v\ r gu'cpf 'uk\ gu'qh'hqf "cpf 'o qxgu"  
co qpi 'f khgtgpv'j cdkcw. 'uwej 'cu'ugci tcuu'dgf u'cpf 'eqtcn'tgghu. 'cv'f khgtgpv'rhg/j kuxqt{ 'uci gu'qt"  
tgr tqf wv'xg'r j cugu. 'qt'y j krg'j wv'kpi 0'"

"  
"  
"  
"

I tqwr gtu'ctg'wpuv geknġ gf. "dqwqo /f y gmkpi . "uqkxct { 'r tgf cvqtu' \*Tcpf cm'cpf "Dtqem' 3; 82. "Tcpf cm'3; 87. "3; 89-0" Hggf kpi "cngu'r mæg' yj tqwi j qw' yj g' f kgn'e { erg' cmj qwi j "o quv' h' guj " hqf 'ku' h' w' p' k' p' u' xqo cej u' eqngewf 'k' p' yj g' g' ctn' { "o qtpkpi "cpf 'cv' f wum' \*Tcpf cm'3; 89-0" Go r v' { " u' xqo cej u' y' g' t' g' cnuq' p' q' v' f' yj tqwi j qw' f' c { rki j v' j' qwtu' \*Ukkc' "Ngg'3; 96-0" k' p' k' k' f' wcn' h' g' g' f' d' { "

**Table 7. Food items recorded in the stomachs of Nassau grouper**

Food items recorded in the stomachs of the Nassau grouper, *Epinephelus striatus* (from Randall, 1965, 1967; Silva Lee, 1974; Claro et al., 1990; Carter et al., 1994).

Nekton	Mullidae	Benthic animals
Fishes	<i>Pseudupeneus maculatus</i>	Molluscs
Acanthuridae	Muraenidae	Gastropods
<i>Acanthurus</i> sp.	<i>Gymnothorax moringa</i>	<i>Strombus gigas</i>
<i>Acanthurus coeruleus</i>	<i>Gymnothorax</i> sp.	<i>Strombus</i> sp.
Apogonidae	<i>Enchelycore nigricans</i>	<i>Fasciolaria tulipa</i>
Atherinidae	<i>Lycodontis moringa</i>	Bivalves
Balistidae	<i>Muraena miliaris</i>	<i>Barbatia cancellaria</i>
<i>Balistes vetula</i>	<i>Muraena</i> sp.	Pelecypods
Bothidae	Ostraciidae	Crustaceans
Carangidae	<i>Lactophrys</i> sp.	Isopods
<i>Caranx ruber</i>	Pomacentridae	Stomatopods
Clupeidae	<i>Chromis cyanea</i>	<i>Gonodactylus perstedii</i>
<i>Harengula humeralis</i>	<i>Chromis multilineata</i>	<i>Pseudosquilla ciliata</i>
<i>Harengula clupeola</i>	<i>Pomacentrus fuscus</i>	<i>Squilla</i> sp.
<i>Jenkinsia lamprotaenia</i>	<i>Pomacentrus</i> sp.	Shrimps/prawns
Gerreidae	<i>Abudefduf saxatilis</i>	Alpheids
<i>Gerres cinereus</i>	<i>Microspathodon chrysurus</i>	Carideans
Haemulidae	Priacanthidae	Penaeids
<i>Haemulon aurolineatum</i>	<i>Priacanthus cruentatus</i>	Lobsters
<i>Haemulon flavolineatum</i>	Scaridae	<i>Panulirus argus</i>
<i>Haemulon album</i>	<i>Sparisoma aurofrenatum</i>	<i>Panulirus guttatus</i>
<i>Haemulon sciurus</i>	<i>Sparisoma rubripinne</i>	<i>Justitia longimana</i>
<i>Haemulon plumieri</i>	<i>Sparisoma chrysopterum</i>	<i>Palinurellus gundlachi</i>
<i>Haemulon</i> sp.	<i>Sparisoma</i> sp.	Hermit crabs
Holocentridae	<i>Scarus vetula</i>	<i>Paguristes depressus</i>
<i>Sargocentron vexillarium</i>	<i>Scarus croicensis</i>	<i>Petrochirus diogenes</i>
<i>Myripristis jacobus</i>	<i>Scarus</i> sp.	Crabs
<i>Holocentrus rufus</i>	Serranidae	<i>Calappa flammea</i>
<i>Holocentrus</i> sp.	<i>Hypoplectrus puella</i>	<i>Calappa</i> sp.
Labridae	<i>Cephalopholis fulva</i>	<i>Stenorhynchus seticornis</i>
<i>Halichoeres garnoti</i>	<i>Epinephelus striatus</i>	<i>Mithrax verrucosus</i>
<i>Halichoeres bivittatus</i>	Synodontidae	<i>Mithrax cinctimanus</i>
<i>Halichoeres</i> sp.	<i>Synodus intermedius</i>	<i>Mithrax</i> sp.
<i>Hemipteronotus</i> sp.	<i>Synodus</i> sp.	<i>Macrocoelema</i> sp.
<i>Clepticus parrae</i>	Urolophidae	<i>Petrolisthes galathinus</i>
Lutjanidae	<i>Urolophus jamaicensis</i>	<i>Chronus ruber</i>
<i>Lutjanus synagris</i>	Molluscs	<i>Portunus sebae</i>
<i>Lutjanus</i> sp.	Squids	<i>Portunus</i> sp.
<i>Ocyurus chrysurus</i>	<i>Loligo</i> sp.	Xanthids
Monacanthidae	Cuttlefish/octopi	Grapsids
<i>Monacanthus ciliatus</i>		
<i>Monacanthus</i> sp.		
<i>Cantherines pullus</i>		

tr k' n' f' k' v' k' p' i' 'y' g' i' k' m' l' e' q' x' g' t' u' 'v' q' g' p' i' w' h' r' t' g' { 'd' { 'u' w' e' k' q' p' \*Vj qo r uq' p' c' p' f' 'O wptq3; 9: . 'E' c' t' v' g' t' " 3; : 8+ 'c' p' f' 'c' m' g' 'c' y' k' f' g' 'x' c' t' k' e' v' { 'c' p' f' 'u' k' g' 't' c' p' i' g' 'q' h' 'h' u' j' g' u' 'c' p' f' 'k' p' x' g' t' v' g' d' t' c' v' g' u' . 'd' q' y' 'd' g' p' y' k' e' 'c' p' f' 'r' g' r' i' k' e' " \*V' c' d' r' u' g' u' '9' 'c' p' f' ' : -0' Y' k' j' 'k' p' e' t' g' c' u' k' p' i' 'c' i' g' . 'y' j' g' t' g' 'k' u' 'c' 'u' j' k' m' h' t' q' o' "e' q' p' u' w' o' k' p' i' "e' t' w' u' c' e' g' c' p' u' 'v' q' 'c' m' k' p' i' " h' k' u' j' g' u' . 'r' e' c' t' i' g' t' 'd' k' x' c' m' x' g' u' . 'h' q' d' u' v' g' t' . 'c' p' f' 'i' c' u' t' q' r' q' f' u' \*g' d' 0' 'G' i' i' n' g' u' v' q' p' 'g' v' 'c' i' n' 0' 3; ; : -0' J' q' y' g' x' g' t' . 'y' j' g' " t' g' r' v' k' q' p' u' j' k' r' 'd' g' y' g' g' p' 'h' u' j' 'u' k' g' 'c' p' f' 'r' t' g' { 'u' k' g' 'u' j' q' y' u' o' w' e' j' "x' c' t' k' v' k' q' p' . 'y' k' j' 'r' e' c' t' i' g' 'h' u' j' 'g' e' v' k' p' i' 'u' o' c' m' i' " r' t' g' { 'c' p' f' 'x' l' e' g' 'x' g' t' u' c' 0' 'Q' p' g' 't' g' r' q' t' v' f' q' e' w' o' g' p' v' g' f' 'c' '7: 2' " o' 'H' N' P' c' u' u' c' w' i' t' q' w' r' g' t' 'u' y' c' m' q' y' k' p' i' 'c' '842" 4: "

"  
"  
"  
"

o o "Gymnothorax=dw'qxgtem"o gcp'r tgf { 'uk' g'y cu'cdqw'37' "qh'yj g'P cuucw'i tqwr gt'hqtnlhpi yj "

\*Ukxc'Ngg'3; 96+0"Hqwt'uwf'kgu'r tqxkf g'c'hggf kpi 'r tqhkg'qh'yj g'P cuucw'i tqwr gt' "Vcdng'9+0"Hkuj "

r tgf qo kpcvfg . 'y kj 'uectkf u'cpf 'rctdkf u'o quv'eqo o qpnf 'kf gp'vkhgf . 'r quukdn { 'dgecwug'yj g'hqto gt "

ecp'dg'tgcf kf { 'tgeqi pk' gf 'htqo 'uqo cej 'eqpvgpw'd { 'y gk' 'wpls wg'f gp'vkkqp "T'cpf cm'3; 87+0 "

Etcdy'g'g'g'o quv'eqo o qp'kpxgt'vgtc'vgo' "Cm'j qwi j 'j gto k'etcdy'cpf 'y g'qr gtew'g'qh "

Strombus'cpf 'Fasciolaria'y g'g'hqwpf . 'uqo cej u'f kf 'pqv'eqp'v'k'p'uj gm'0'k'p'qpg'Ewdcp'uwf { . 'y g' "

o quv'cdwpf cpv'kgo u'\*d { 'y gki j v'y g'g'i twpw' . 'r cttq'v'kuj gu' . 'cpf 'qevqr wu'y kj 'c' 'uwi i gukqp' 'y cv' "

o qtg'i twpw'y g'g'vcn'gp'k'p'y k'p'vgt'o qp'vj u'\*E'ctq'g'v'cr'03; ; 2+0 "

**Table 8. Comparison of 4 studies of stomach contents of Nassau grouper**

Principal categories of food items encountered in the stomachs of Nassau grouper, *Epinephelus striatus*, in four studies: (A) Virgin Islands (Randall, 1965, percent by volume); (B) Cuba (Silva Lee, 1974, percent frequency occurrence); (C) Cuba (Claro et al., 1990, percent by weight); (D) Belize (Carter et al., 1994, percent frequency occurrence).

Food category	A	B	C	D
Nekton, fish	53	39	71	58
Benthic, crustaceans				
crabs	23	29	8	16
stomatopods	6	3	<1	1
shrimp/prawn	5	8	<1	3
spiny lobster	4	6	4	6
hermit crab	1	<1	-	1
isopod	<1	<1	-	-
unidentified	1	4	-	3
Nekton, molluscs				
cephalopods	5	9	15	6
Benthic, molluscs				
gastropods	2	2	2	1
Unidentified	-	-	-	5

mecn'r tgf cvqtu'cpf 'ctg'f g'g'p'f gf 'd { 'm'pi 'xgpqo qwu'hk'ur k'p'gu' . 'uwej 'y cv' 'g'xgp'y j gp'uj ctmu'qt "

recti g'i tqwr gt'f q'c'w'cem'yj g' { 'j' cxg'dggp'qdugt'xgf 'ko o gf k'vgn' { 't'g'v'g'v'k'pi 'y kj qw'qdxk'wu'k'plwt { "

v'q' 'y g' 'rkqphkuj "Ucf qx { . 'r gtu'0'qdu'0'0'P qpg'y g'guu' . 'y g'g' 'ku'c' 'r wdrkuj gf 't'gr qt'v'qh'hkuj gto gp'k'p' 'y g' "

D'ej co cu'ecr wtk'pi "qpg'v'ki g't'i tqwr gt' "Mycteroperca tigris" + 'cpf 'y q'P cuucw'i tqwr gt' "E. striatus" + "

g'cej 'y kj "c'rkqphkuj 'k'p'ku'u'qo cej "O'crlmq'xk "g'v'cr'0422: + 'd'oi h'x'g'P cuucw'i tqwr gtu' "E. striatus." "

ecwi j v'q'hh'G'rgwj gtc'K'ur'p'f "cv'cp'cr r tqzko cv'f gr yj 'qh'36'o "qp'7'O ctej "422: . 'y g'g'f ku'ge'v'f'0 "

Vy q'qh'yj g'uqo cej u'eqp'v'k'p'gf 't'gf "rkqphkuj 0"Vj g'htu'v'i tqwr gt' "699/o o "UN" + 'eqp'v'k'p'gf "c' 'r' ct'v'k'cm' { "

f ki gu'gf 'rkqphkuj . 'kf gp'v'kh'cdng'q'pn' { 'd { 'y g'o qtr j qm'j { 'cpf 'o w'k'r'k'ek'v' { 'qh'yj g'tgo c'k'p'k'pi 'h'k'p' 't'c { u'0 "

Vj g'uge'q'p'f 'urki j v'k' 'r'cti g't'i tqwr gt' "6: 4/o o "UN" + 'eqp'v'k'p'gf "c' 't'gf "rkqphkuj "qh'359/o o "UN." y j k'ej "

y cu'k'p' 'cm' qu'v'r tk'k'p'g'eq'p'f k'k'q'p'0' "Uqo g'qh'yj ku'hggf k'pi 'o c { 't'gu'w'v'ht'qo "c'w'go r w'u'v'q'eq'p'f k'k'q'p' "

mecn'r tgf cvqtu'v'q'hggf "qp' 'y g'p'q'p'p'v'k'x'g'ur g'elgu'0'F k'x'g'tu'k'p' 'y g'Ec { o cp'K'ur'p'f u'j cxg'v'k'p'gf "

y k'f "P cuucw'i tqwr gt'v'q'eq'puwo g'rkqphkuj . 'y kj qw'v'y g'i tqwr gt'uj qy k'pi "k'm'gh'g'ew' "Y 0J g { o cp. "

"  
"  
"  
"

k'p'D'grk' g' 'y g'r tgf qo k'p'cp'v'h'q'q'f . 'd { "

r gte'gp'v'ci g'ht'gs w'p'e { 'qh'q'ee'wt'g'p'eg . 'y cu'hkuj . "

y kj "c'j k'j j 'r gte'gp'v'ci g'qh'et'w'x'ce'g'c'p'u . "

g'ur g'ek'cm' { 'et'cdy' . 'cpf "c' 'uo c'm'p'wo d'g't'qh' "

i c'ust'qr q'f u' . 'egr j c'm'r q'f u' . 'cpf 'r g'rg'e { r q'f u'0' "Vj g' "

r t'k'p'ek'r c'n'r t'g { 'hkuj 'h'co k'k'g'u'y g'g'i t'wpw' 'cpf "

u'p'c'r r g'tu' "E'ct'v'g't'g'v'cr'03; ; 6+0' "N'kng'q'yj g't "

i t'qwr g'tu' . 'P cuucw' 'h'q'm'q'y "cpf 'hggf 'y kj 'q'yj g't "

r t'gf c'v'qtu' . 'uwej "cu'v't'ki i g't'hkuj . 'q'ev'qr wu' . 'qt' "g'gn' "

"E'ct'v'g't'g'v'cr'03; ; 6. "U'w'n'k'x'c'p' 'cpf 'f g'I c't'k'p'g/ "

Y k'ej c'v'k'um' { '3; ; 6. "T'q'd'g't'w'g'v'cr'03; ; 7. "U'c'f q'x { "

r g'tu'0'q'du'0' 'r t'gu'wo c'd'n' { 'd'g'p'g'h'k'k'p'i 'ht'qo 'ur q'ku' "

o c'f g'c'x'c'k'c'd'ng' 'f k't'g'e'w'v' . 'qt' 'ht'qo 'f k'w'w't'd'c'p'eg'qh' "

r t'g { 'ur g'elgu'0' "

Uqo g'c'p'g'ef q'v'cr' 'cpf 'r j q'v'qi t'c'r j k'e "

g'x'k'f g'p'eg' 'r tqx'k'f g'f 'd { 'hkuj g'tu' 'cpf 'f k'x'g'tu' "

u'wi i gu'u' 'y c'v'p'c'v'k'x'g' 'i tqwr gt' 'ur g'elgu' 'c't'g'r t'g { k'pi "

qp' 'y j g'k'p'x'c'uk'x'g' 't'gf 'rk'q'p'h'kuj 'y kj 'u'qo g' "

t'gi w'r'ct'k'v' { 0' "N'k'q'p'h'kuj "c't'g' 'i g'p'g't'cm' { 'w'p'h'co k'k'ct' "v'q "



i tqwr gt. "wukpi "RET" \*Rqn( o gtcug'Ej clp'Tgcevqap+/co r rkhgf "o vFPC"i gpgu'cpf "pwerget"  
o letqucvgrkngu."tgxgcrnf "pq"ergetn( "f ghkpgf "r qr wrvqap"uudutwewt( "hqt" gkj gt "ur gekgu"cv'v'j g"  
i gqi tcr j le"mcevkqpu"uco r rnf. "k0Dgrk g. "Ewdc. "Dcj co cu. "Hrktk c' hqt "P cuucw'i tqwr gt "Ugf dgtt { "  
gv'cr03; ; 8-0"Vj gug'f cv'kpf kecvg'y cv'ur cy plpi "ci i tgi cvkqpu"ctg'pqv'gzenvukxgn( "ugrh/tgetwvki "  
cpf "vj cv'v'j g'rctxcn'luci gu'ecp" f kur gtug"qxgt "i tgcvf'kucpegu. "j qy gxgt "vj g'tgrv'xg'ko r qtvcpeg'qh"  
ugrh/tgetwko gpv'cpf "rctxcnko o ki tcvkqap"vq"mecn( qr wrvqapu'y cu'pqv'erget "Ugf dgtt { "gv'cr03; ; 8-0""  
Tgegpv'cf xcegu'o ki j v'dg'cr r rkhgf "vq"gzco kpg'uqwtg"qt'pwtugt { "ctgcu'cpf "uj klu'kp" hkuj "dgvy ggp"  
j cdkcuv'y kj "eqpv'cvkpi "o letqej go kecnluki pcwtgu0'

Tguvnu'qh'dqv'j "J cvrg { "4227+'cpf "Ugf dgtt { "gv'cr0\*3; ; 8+'kpf kecvg'c "ukpi ng'r cpo keve"  
r qr wrvqap'qh'P cuucw'i tqwr gt "kp"vj g'pqt'v'j gtp'Ectkddgcp"dcukp'y kj "j ki j "i gpg'hny "dgvy ggp"  
Hrktk c. "Ewdc. "Dgrk g'cpf "vj g'Dcj co cu0"J qy gxgt. "vj g' "f q'pqv's wcpvkh( "vj g'eqppgevqap0" Tguvnu'qh'  
cp'qpi qkpi "Rj F "uwf { "wukpi "o qtg'hkpg/uecrg"i gpgv'ke'v'gej pls wgu'o c { "r tqxkf g'c"o qtg'f gckrgf "  
wpf gtucpf kpi "qh'r qr wrvqap"utwewt0" \*Crgzku'Lcemqap. "Rj F 0tugctej "kp"r tqi tguu. "F gr ctvo gpv"  
qh'Geqrni { ( "Gxqnvkqapct { "Dkqrni { . "Wpkxgtukv( "qh'Ecnkhtpk. "Ucpvc'Et w( +0'  
"

## 2. THREATS OR STRESSORS

Mg { "vj tgcw'ctg'r tguv'v'gf "cnj qwi j "vj g' "ctg'wprkngn( "vq"dg'cm'v'j cv'ctg'r quukdrg0'

### 2.a. Anthropogenic Effects

*Fishing effects.* "Vy q'f khtgtpv'cur gew'qh'hkuj kpi "ghhev'P cuucw'i tqwr gt "uqemu. "hkuj kpi "  
ghhtv'v'j tqwi j qw'v'j g'pqp/ur cy plpi "o qp'v'j u'cpf "hkuj kpi "ghhtv'v'j kgevgf "cv'ur cy plpi "ci i tgi cvkqpu"  
qt"o ki tcvqt { "ceegu'v'q'ur cy plpi "ci i tgi cvkqpu0""

P cuucw'i tqwr gt "ctg'hkuj gf "eqo o gtekm( "cpf "tgetgcvkqpcm( "vj tqwi j qw'v'j g" { gct'd { "  
j cpf rkp. "mipi rkp. "hkuj "tcr u. "ur gct"i wpu. "cpf "i kmpgw" \*P O HU I gpgtcn'Ecpxcu'Ncpf kpi "U{ uvg0 +0"  
Ci i tgi cvkqpu"ctg"o clpn( "gzn'rkkgf "d { "j cpf rkp"u'qt" d { "hkuj "tcr u. "cnj qwi j "i kmpgw'y gtg'dgkpi "  
wugf "kp"O gzkeq'kp"vj g'gctn( "vq"o kf/3; ; 2u" \*Ci vkrct/Rgtgtc"4226+0"Ucf qx { "cpf "Gmwpf "3; ; ; +"  
uj qy "f gerkp'gu'kp"rcpf kpi u. "ecvej "r gt "wkv'ghhtv'v' \*ERWG+"cpf . "d { "ko r necvkqap. "cdwvf cpeg'kp"vj g"  
rv'g'3; ; 2u'cpf "gctn( "3; ; 2u'v'j tqwi j qw'ku'tcpi g. "y j kej "j cu'rgf "P cuucw'i tqwr gt "vq"pqy "dg"  
eqpukf gtgf "eqo o gtekm( "gzvkp'ev'kp"c"pwo dgt'qh'ctgcu" \*Ucf qx { "cpf "Gmwpf "3; ; ; +0" Tgegpv'  
tgr qt w'htqo "vj tqwi j qw'v'j g'P cuucw'i tqwr gt w'htqo g'f qewo gpv'eqpv'kwgf "r qr wrvqap" f gerkp'gu'cpf "  
mqu'qh'ci i tgi cvkqpu" \*Ucf qx { "f g'O kej guqp"4234+0'

Vj g'ci i tgi cvkxg'tgr tqf wvqap'uv'rg"i cvj g'kpi "cv'r tgf kevcdng"ukgu'kp"rcti g'eqpegpv'cvkqpu"vq"  
ur cy p'f wtkpi "c'hgy "y ggmu" \*qxgt "c'hgy "o qp'v'j u+"gej " { gct'/"o cngu'v'j g'P cuucw'i tqwr gt "xwpgtcdng"  
cu'c'vcti gv'qh'hkuj kpi "hkn"o cp { "qj gt "tggh'ur geku'v'j cv'htqo "rcti g'ci i tgi cvkqpu"vq"ur cy p0"kp"o cp { "  
r rnegu. "ci i tgi cvkqap/hkuj kpi "qpeg"r tqf wegf "o quv'qh'v'j g'cppwcn'rcpf kpi u'qh'v'j g'ur geku" \*gd 0'Erctq"  
gv'cr03; ; 2. "Dwuj "3; ; 4-0" Dgecwug'P cuucw'i tqwr gt "ctg'qpn( "mpqy p'vq'tgr tqf weg'kp"ur cy plpi "  
ci i tgi cvkqpu. "tgo qxkpi "tkr g'kpf kxkf wcu'f wtkpi "ur cy plpi "j cu'v'j g'r qv'pvcn'vq"i tgcw( "kphwv'peg"  
r qr wrvqap" f { pco leu'cpf "hwwt g'hkuj gt { " { kgrf u" \*Uj cr ktq"3; ; 9-0"Vj g'hcev'v'j cv'o wej "qh'v'j g'ecvej "kp"

"  
"  
"  
"

o cp{ "eqwptkgu"j kxqtkecm{ "eco g'ltqo 'ur cy plpi 'ci i tgi cvkqpu"\*Qmgp'cpf "NgRmeg"3; 9: ." Ci wkrt/Rgtgtc"3; ; 6."Ucf qx{ "cpf "Gmwpf "3; ; ; +hkngn{ 'o ci plhkgf 'y g'ghgewu'v'j g'gzvqv'v'j cv vti gvgf 'ci i tgi cvkqpu'j cxg'eqmcr ugf 'kp'o cp{ "eqwptkgu"\*Ucf qx{ 'f g'O kej guqp"4234+0"Ku" f genpogu'j cxg'eqo r tqo kugf 'y g'geqmi kecnhwpevqp'qh'c'o clqt 'vqr 'r tgf cvqt 'kp'y g'tggh'gequ{ uvg " \*Tcpf cm3; ; 9.'O wo d{ "gv'cr04228.'O wo d{ "gv'cr04234+0"

"Rtkqt 'v' tgi wrcvqpu'r tqj kdkkpi 'y g'j' ctxguv'cpf 'r quuguukqp.'y g'WUOXki kp'Krcpf u'cpf " Rwtgvtq"Tequ't'ggh'hkuj g'kgu'eqo o qpn{ 'vqmiP cuucw'i tqwr gtu'cv'ci i tgi cvkqpu'ukgu"\*UCHO E"3; ; 2." EHO E"3; ; 5+0"P cuucw'i tqwr g'j' cxg'cnuq'dggp'ecwi j v'ltqo 'ugxgtcn'ukgu'qh'v'j g'Lco clecp'eqcu' cpf 'qh'v'j g'pqt'v'j gtp'eqcu'qh'v'j g'F qo kplecp'T gr wdke"\*Vj qo r uqp'cpf 'O wptq"3; ; 5."Ucf qx{ " 3; ; 9+0"Kp'O gzleq.'cv'ngcu'ugxgp'ci i tgi cvkqpu'ukgu'j cxg'dggp'hkuj gf 'cmipi 'y g' l' wecvp'Rgpkuwrc" ukpeg'v'j g'dgi kppkpi 'qh'v'j g'42<sup>v</sup>'egpwt{ "3; 32/3; 42+\*Ci wkrt/Rgtgtc"3; ; 6+0""Vj qo r uqp"\*3; 67+ " f guetkdgf 'qpg'rcti g'ci i tgi cvkqpu'ukg'qh'Ec{ 'I mt{ .Dgrk g.'y cv'j cf 'dggp'hkuj gf 'hqt'o cp{ " f gecf gu"cpf 'r quwrcv'v'j cv'q'v'j g'eqpi tgi cvkqpu'qeevt.'dw'j cf 'guecr gf 'f gvevqp'dgecvug'qh' v'j g'kt'gr j go gtcn'pcwtg=q'v'j g'ukgu'j cxg'dggp'kf gpv'khgf 'ukpeg'Vj qo r uqp'au'y qtm'Rcl'cpf " I tko uj cy "4223+0"Kp'Ewdc.'43'ur cy plpi 'ci i tgi cvkqpu'ukgu'y g'g'kf gpv'khgf =qpn{ "32'qh'v'j gug" ci i tgi cvkqpu'ukgu'y g'g'P cuucw'i tqwr g'ur cy plpi 'ci i tgi cvkqpu'ukgu'0"Qh'v'j g'32'P cuucw'i tqwr g' " ci i tgi cvkqpu'ukgu.'y q'y g'g'oc'i i tgi cvkqpu'ucwuguo'y g'g'kf gpv'khgf 'cu'of genkpgf o'cpf 'gki j v'y g'g' kf gpv'khgf 'cu'ouj ctr n{ 'f genkpgf o"\*Erctq'cpf 'Nkpf go cp"4225+0"Kp'Cvrcpke'y cvgtu.'P cuucw'i tqwr g' " j cxg'dggp'ecwi j v'kp'y g'Hqtkf c'Mg{u'cpf 'y g'Dcj co cu"\*Daj puceni4225+0"Vj g'Dcj co kcp" F gr ctwo gpv'qh'Hkuj g'kgu'tgr q'v'v'j cv'lp"3; ; 4."qxgt"42'ur cy plpi 'mqecvqpu'y g'g'hkuj gf "T'O Vj qo r uqp.'Dcj co cu'F gr ctwo gpv'qh'Hkuj g'kgu.'r gtu0'eqo o 0'v'q' l' 0Ucf qx{ .P O HU."3; ; 4+ " cmj qwi j "y g'ewtgpv'ucwu'qh'o cp{ 'ku'wmpqy p0"C'tgugctej 'v'kr 'f wtkpi 'y g'ur cy plpi 'ugcuq'qh' 4235"\*DOGtuo cu.'UETHC'P gy urgwgt"39.'Lxpg"4235+'hckrgf 'v' h'kp' cp{ 'uki p'qh'P cuucw'i tqwr gtu' cv'v'j g'ur cy plpi 'ci i tgi cvkqpu'ukg'v'j cv'y cu'v'j g'qtki kpcn'ukg'f guetkdgf 'd{ "Uo kj "3; 94+0"Vj g'g'ctg" pq'npqy p'ur cy plpi 'ci i tgi cvkqpu'ukgu'kp'Hqtkf c'y cvgtu0"Kp'y g'I wh'qh'O gzleq.'P cuucw'i tqwr g' " y g'g'ecwi j v'r tko ctkn{ 'qh'h'uwj y guv'Hqtkf c.'y kj 'eqo o gtekn'cpf 'tgetgcvqpcn'ecvej 'tgr q'v'v'j " ltqo 'y g'uwj y guv'Hqtkf c'Mg{u0"Daj 'tgetgcvqpcn'cpf 'eqo o gtekn'ecvej gu'qh'P cuucw'i tqwr g' " y g'g'j ki j g'ltqo 'y g'Hqtkf c/I wh'qh'O gzleq'v'j cp'ltqo 'y g'Hqtkf c/Cvrcpke'eqcu'ltqo "3; ; 8/ 3; ; 5"\*P O HUI gpgtcn'Ecpxcu'Ncpf kpi u'U{ uvg "0"Chgt"3; ; 3.'y gug'f khgtgpegu'y g'g'r tqdcdn{ " r ctvcm{ 'f wg'v'q'hkuj g'g' { 'tgi wrcvqpu'dcplkpi 'cm'ecr wtg'qh'P cuucw'i tqwr gtu'ltqo 'y g'WUOCvrcpke" y cvgtu.'y qwi j 'pqv'ltqo 'y g'I wh'qh'O gzleq'=j ctxguv'cpf 'r quuguukqp'ctg'pqy 'dcp'p'kf 'kp'cm'WUO' y cvgtu"\*EHO E"3; ; 2."UCHO E"3; ; 3.'I O HO E"3; ; 8.'eqo r krgf 'kp'Ucf qx{ "cpf "Gmwpf "3; ; ; +0"

*Age composition of fishery catches*0"P cuucw'i tqwr g'uco r rnf 'ltqo 'ecvej gu'cv'7'ur cy plpi " ci i tgi cvkqpu'kp'y g'Ec{ o cp'Krcpf au'ltqo "3; ; 9/3; ; 4'i gpgtcm{ 'hgm'y kj kp'ci g'ercuugu'4/; "{ gctu" cpf 'kpenf gf 'o cp{ 'ko o cwtg'kpf kxkf wenu"\*Dwij 'gv'cr04228+0"P q'uk' g'cv'ci g'f khgtgpegu'dgy ggp" o cngu'cpf 'hgo cngu'j cxg'i gpgtcm{ 'dggp'pqvgf "Dwij 'gv'cr04228+0"Qxgt": 2' "qh'v'j g'uco r ngu'cngp" \*p"?": 38+ltqo 'c'npqy p'ci i tgi cvkqpu'kp'y g'Xki kp'Krcpf u'dgy ggp"3; 96'cpf "3; 9: 'y g'g'ci gf "6/8" { gctu"\*cu'guko cvgf 'd{ 'r tqdk'cpcn{ uku+\*Qmgp'cpf "NcRmeg"3; 9; +y j krg'bo quv'hkuj 'rcpf gf 'ltqo " ci i tgi cvkqpu'kp'y g'Ec{ o cp'Krcpf u'ltqo "3; ; 9/3; ; 4'y g'g'ci gf "9/; "{ gctu"\*Dwij 'gv'cr04228+0"

"  
"  
"  
"

Ci g'emcuugu8/; "f qo kpcvzf "cm'ncpf kpi u'kp'uqwj y guvgtp'Ewdc"cpf "5/: "{ gctu'kp'pqtjy gcuvgtp'Ewdc" dgvy ggp'yj g'gctn{"3; 82u'cpf "rcvg"3; : 2u'y kj "72" "qh'ncpf kpi u'eqo kpi 'htqo "ci i tgi cvkqup"Erctq" gv'cr03; ; 2+0'

*Size composition of fishery catches.* C'o czko wo 'hgpi yj 'qh'3.442'o o "VN"cpf 'y gli j v'qh' 45/49"ni "ctg'tgeqtf gf "hqt"yj g'P cuucw'i tqwr gt "Gxgto cpp"3; 22."Tcpf cm'3; 85."Uo kj "3; 93." Dwguc"3; : 9+0"O quv'hkuj 'kp'o ctngvu."j qy gxgt."ctg'eqpukf gtcn{"uo cmgt "100"4/33"mi +\*Uo kj " 3; 93+0"Y gli j w'qh'ci i tgi cvkpi 'hkuj 'tapi gf 'htqo "7/34"mi . 'y kj "c'o czko wo "qh'36"ni \*Uo kj "3; 93." 3; 9: ."Ci w'krt/Rgt gtc"3; ; 6+0"i tqwr gt "w'vq"; 82'o o "UN"y gtg'vcnrgp'kp'yj g'Xkti kp'Kncpf u" cmj qwi j 'hkuj 'rci gt 'yj cp'cdqw'922'o o 'y gtg'wpeqo o qp"92"qh': 38'hkuj 'uco r rnf +\*Qnrgp"cpf " NcRrcg"3; 9; +0"O czko wo 'yj g'atg'v'ecno gcp'hgpi yj "N'o'htqo 'yj g'xqp'Dgtvncph{"i tqy yj 'hwpev'kqp" /'xqp'Dgtvncph{"3; 79+j cu'dggp'guko cvgf "cv'dgvy ggp"982/3.34; 'o o "VN"\*Vj qo ruqp"cpf "O wptq" 3; 9: ."Qnrgp"cpf "NcRrcg"3; 9; ."Erctq"gv'cr03; ; 2+0'

Cu'ucvzf 'r t'gxlqunf . 'o gcp'o crg'cpf 'hgo crg'uk' gu'ctg'uko krt 'y kj kp'c'i kxgp'ctgc."qt'cv'c" ur gekhe'ci i tgi cvkqp'ukg0"Vj gtg'ku'uqo g'kpf kec'kqp'yj cv'uk' gu'qh'dqy "ugz gu'f gen'kp'kp'ctgcu'qh' j ki j gt'g'zr m'kcv'kqp'xgtuwu'wpgzr m'kgf 'r qr w'v'kqup'y kj kp'c'ur gekhe'tgi kqp "Ectvgt'gv'cr03; ; 6+ "Hki 0; +0"

Y j gp'g'zr m'kcv'kqp'ku" j ki j ."ecvej gu'ctg'rci gn{" eqo r tkugf "qh'lwxgpkrgu" \*i tqy yj "q'xgthkuj kpi +0Hqt" gzco r ng."kp'Dgnk' g.'yj g" cxgtci g'hgpi yj "qh'dqy "ugz gu" y cu'322'o o "uo cmgt 'kp" ecvej gu'htqo "g'zr m'kgf" eqo r ctgf "v'wpgzr m'kgf" ci i tgi cvkqup"\*Hki 0; +0"kp" qpn{"y q'ecugu'y gtg'hgo crgu" uki pk'kecpv{"r'api gt 'yj cp" o crgu."y j krg'o crgu'y gtg" pgxgt'rci gt 'yj cp'hgo crgu" \*Vj qo ruqp"cpf "O wptq" 3; 9: ."Ucf qx{"cpf "Eqnkp" 3; ; 7+0"O quv'ecvej gu" eqpukv'gf "qh'lwxgpkrgu'kp" j g'cxkn{"g'zr m'kgf "ctgcu'qh' Rwtvq'Tleq."Hqtkf c"\*Hki u0' 44"cpf "45"kp'Ucf qx {"cpf " Gmwpf "3; ; + "cpf "Ewdc" \*Gur k'p'quc"3; ; 2+0'




Figure 21

Length-frequency distributions of male and female Nassau grouper, *Epinephelus striatus*, taken from aggregations [males = white bars, females = black bars] in Belize: (A) unexploited site (n=694); (B) exploited site (n=485) (Carter et al., 1994).

Figure 8. Length-frequency distributions by sex for exploited and unexploited sites in Belize

**Table 9. Mean sizes and sex ratio across a gradient of fishing pressure (excerpt from Sadovy and Eklund 1999)**

Mean sizes and sex ratios of aggregating and non-aggregating Nassau grouper, *Epinephelus striatus*, in the western Atlantic, from lightly (top of table) to heavily (bottom of table) exploited areas. Fishing intensity implied by descriptions of current and historic fishing activity (from Sadovy and Colin, 1995). Number in parentheses refers to references.

Sex ratio F:M	Mean SL in mm		Max SL in mm	Gear used <sup>1</sup>	Location and source <sup>2</sup>
	F	M			
0.57:1 (n=750)	526	529	750	T,H	Bermuda, offshore banks > 60 m deep (1)
0.72:1 (n=163)		554 <sup>3</sup>	718	H,T	Jamaica, offshore (3)
1.5:1 (n=694)	517	521	802	H,S	Belize, aggregation (4)
2.0:1 (n=42)	502	487	568	G	Mexico, aggregation (2)
0.67:1 (n=70)	514	503 <sup>4</sup>	657	G	Mexico, aggregation (2)
1.0:1 (n=940)	589	585	940	T,H	U.S.V.I., aggregation (5)
1.9:1 (n=95)	516	512	640	H	Caymans, aggregation (6)
2.0:1 (n=140)	506	538	772	H	Caymans, aggregation (7)
2.2:1 (n=717)	418	420	760	H,S	Belize, nonaggregated (4)
2.4:1 (n=485)	418	420	690	H,S	Belize, aggregation (4)
2.5:1 (n=216)	549	517 <sup>4</sup>	700	T	Bahamas, aggregation (8)
4.0:1 (n=319)	>500 <sup>3</sup>		710	S,T	Cuba, (9) (only adults assessed)

<sup>1</sup> Gear used: T=trap; H=handline; S=spear; G=gillnet.  
<sup>2</sup> Sources: (1) Bardach et al., 1958; (2) Sosa-Cordero & Cárdenas-Vidal, 1997; Aguilar-Perera, 1994; (3) Thompson & Munro, 1978; (4) Carter et al., 1994; (5) Olsen and LaPlace, 1979; (6) Colin et al., 1987; (7) Bush (text footnote 42); (8) Colin, 1992; (9) Claro et al., 1990.  
<sup>3</sup> Males and females combined.  
<sup>4</sup> Females significantly larger than males at p<0.05, otherwise no sex difference in size.

**2.b. Habitat loss or degradation**

F wtkpi 'ku'xctkqwa'rhg'j kuxqt { 'uuci gu. 'yj g'P cuucw'i tqwr gt 'wugu'o cp { 'f khgt gpv' eqo o wpkkgu'qt'j cdkcv'v' r gu'y kj kp'vj g'eqtcr't ggh'gequ { uvg0 0''Vj g'kpet gcug'kp'wt dcp. 'kpf wutken'' cpf 'qwt ku'f gxgnr o gpw'vj tqwi j qw'vj g'ur geku' t'cpi g'ko r ceu'equcen'o cpi tqxgu. 'ugci tcuu' dgf u. 'guwctkgu. 'cpf 'rhxg'eqtcr' O cj qp'3; ; 2-#0'Nqu'qh'lwxgpkrg'j cdkcv. 'uwej 'cu'o cetqcri cg. ' ugcitcui'dgf u. 'cpf 'o cpi tqxg'ej cpgnu'ku'rhgn' { 'v'pgi cvkxgn' { 'chgevt'getwko gpvt'cvgu0''Cu' uj qy p'kp'vj g'Dcj co cu'F cj ni tgp'cpf 'Gi i nguqp'4223+. 'j cdkcv'r tghgt gpegu'qt'ugrgevkqp'o c { 'dg' ng { 'v'gctn' { 'uwxkxcr'cpf 'uwdugs wgpvr qr wcvkqp'uk' g'cpf 'hqu'qh'vj qug'r tghgtg' { 'eqtcr'cni cni' ugwgo gpvj cdkcv'o c { 'r qug'c'vj tgecv'v' i tqwr gt 'r qr wcvkqp' M'w'ho cp'cpf 'Tqo gtq'4233-#0'Rqqt' y cvgt's wcrk' { 'ku'c'vj tgecv'v' dqy 'eqtcr'cpf 'o cetqcri cg'kp'pgctuj qtg'ctgcu0''kpetgcugf'' ugf ko gpvcvkp'tguwtkpi 'htqo 'r qqt'rcpf 'f gxgnr o gpvr tcevkgeu'cf f u'wtdkf kv' { 'cpf 'r qmwcpvu'kp'v' pgctuj qtg'j cdkcv'cpf 'ecp'ej cpi g'y cvgt'huqy 'r cwgtpu'kp'etggm. 'y j gtg'pgy n' { 'ugwrg' 'lwxgpkrgu' o c { 'dg'hqwpf 0''F tgf i kpi 'qr gtcvkqp'ctg'cnuq'ecr cdrg'qh'f gwtq { kpi 'o cetqcri cni'dgf u'vj cv'o c { 'dg' wugf 'cu'i tqwr gt 'pwtugt { 'ctgcu0''Chgevu'v'P cuucw'i tqwr gt 'yj tqwi j 'j cdkcv'rhqu'qt'f gi tcf cvkqp'ctg' uwo o ctkt gf 'dguv'd { 'Ugo o gpu'gv'cni'422: c+<

oY j kg'P cuucw'i tqwr gt 'ctg'v' r kcm' { 'vj qwi j v'qh'cu'utkew' { 'c'tggh'cuuqekcv'f 'ur geku. 'vj g' { ' v'cpukkkp'vj tqwi j 'c'ugt'ku'qh'qpqi gpgv'e'uj kku. 'htqo 'r rcpmqple'rtxcg. 'v'pgctuj qtg'ugc/ i tcuu'cpf 'cni cg'j cdkcv'v' r tgf qo kpcv'n' { 'tgg'h'j cdkcv'g'0 0'hqtg'tggh'cpf 'tgg'h'etguv'0''Gxgp'' y kj kp'tgg'h'j cdkcv'v' j gtg'cr r gctv'v'q'dg'qpqi gpgv'e' uqt kpi. 'uwej 'vj cv'vj g'rti gt'' kpf kxkf wcu'v'gpf 'v'q'qeer { 'vj g'f ggr gt. 'o qtg'twi qug'tggh'ctgcu0''Gcej 'qh'vj gug'i gpgtcr' j cdkcv'j cu'v'pfti qpg'cpf 'eqpv'kwgu'v'w'pfti q'ej cpi g0''Qr gp/qegcp'rtxcn'j cdkcv'ku' dgkpi 'kphwpegf 'd { 'vj g'qpi qkpi 'kpetgcug'kp'qegcp'ugc/ uwxhceg'vgo r gtcwv'gu0''Vj gug' ej cpi gu'kp'vgo r gtcwv'g'o c { 'kphwpeg'j cdkcv's wcrk' { 'f ktgevn' { 'vj tqwi j 'r j { 'ukqmi kcr' utguu. 'qt'kpf ktgevn' { 'vj tqwi j 'ko r ceu'v' r tg { 'cpf 'r tgf cvgt'f gpukkgu' C'pf gtuqp'3; ; : -#0''

''  
''  
''  
''

Ugci tcuugu'ctg'lp'f gerkg'i mdcn\ "Nqv g'gv'cr04228+=v'j g'f gerkg'qh'wtvg'i tcuu'lp'v'j g'  
Ectkddgcp'o c { 'tgf weg'v'j g'co qwpv'qh'uwxcdrg'j cdkcv'ht'pgy n' ugwgf 'P cuucw'i tqwr gt."  
cpf "o c { 'kphwpeg'v'j g'cdwfp cpeg'qh'r tg { 'kgo u'ht'pgy 'tgetwku0"Eqtcn'tggh'dkqi gple"  
utwewt'g'ku'lp'f gerkg."qy kpi 'k'rti g'r ctv'v'v'j g'f tco cve'f gerkg'lp'cetqr qtk'eqtcn0"  
Hwtv'j gto qtg.'v'j g'qpi qkpi 'f gegcug'lp'qegcp'cekf kv' 'ku'rkng\ 'v'j cxg'c'f tco cve'kphwpeg"  
qp'v'j g'ceetgkqp'tcv'qh'eqtcn'ur geku'lp'v'j g'hwatg"\*J qgi j /I wfdgti 'gv'cr04229+0"K'ku"  
r quukdr'v'j cv'v'j g'qpi qkpi 'cpf 'r tqlgev'f'f gerkg'lp'dkqi gple'utwewt'g'qp'Ectkddgcp'eqtcn'  
tgg'hu'y km'j cxg'c'f tco cve'ko r cev'qp'v'j g'cxckrdkiv' 'cpf 's wcrk\ 'qh'tggh'j cdkcv'ht'o cwtg'  
P cuucw'i tqwr gt0"

Uwxcdrg'j cdkcv'ht'v'j g'P cuucw'i tqwr gt'ku'cnuq'rkng\ 'v'dg'lp'f gerkg"\*Ugo o g'gv'cr0'  
422: c. 'Nqv g'gv'cr04228+0'Qh'v'j g'42.222'no 'qh'eqtcn'tggh'guko cvgf 'ht'v'j g'Ectkddgcp'lp'v'j g'  
o kf/3; ; 2u.'4; ' 'y cu'guko cvgf 'v'dg'wfp gt'j ki j 'tkun'qh'f gi tcf cvkqp'htqo 'j wo cp'cev'kkgu.'54' "  
ku'cv'o gf kwo 'tkun'cpf '5; ' 'ku'cv'ny 'tkun'Dt { cpv'gv'cr03; ; : +0'C'f gecf g'ci q.'I ctf pgt'cpf "  
eqy qtngtu"\*4225+ff qewo gpv'f 'dcukp/y kf g'muugu'qh'j ctf 'eqtcn'eqxgt'htqo 'cdqw/72' 'v'cdqw'  
32' 0"Y kj 'pq'lpf kec'v'pu'qh'tgeqxt { 'qh'uergtcev'kpcp'eqtcn'eqxgt.'k'ku'rkng\ 'v'cv'o cp { "  
Ectkddgcp'tggh'hu'km'eqpv'kw'v'v'j 'mqug'v'j tgg/f ko gpukpcn'utwewt'g'v'j tqwi j 'wpeqo r gpucv'f "  
dkqgtqkqp'cpf 'kpetgcug'lp'o cetqcn'cn'eqxgt"\*O eEmcej cp'gv'cr04224+0"

Wpf gt'pcwt'cn'eqpf k'k'pu'v'j g'P cuucw'i tqwr gt'cr r gtu'v'v'j r tghgt'ergct'y cvgtu"\*Cnk'pu'gv'cr0'  
422; + 'dw'ku'hckn\ 'v'ngtcpv'qh'c'tcpi g'qh'y cvgt's wcrkkgu'qpg'cf wv'utxkg'f'ht'o qtg'v'j cp'ugxgp'  
{ gtu'lp'v'j g'qr'f'P gy 'I qtn'Cs wctkwo 'lp'y j kej 'v'j g'y cvgt'cv'ko gu'dgeco g'pgctn\ 'ht'g'j 'cpf 'y cu'  
htgs wgpw\ 's wkg'r qmwgf "\*Vqy pugpf '3; 27+0"

**2.c. Climate change implications**

P cuucw'i tqwr gt'j cxg'dggp'hqwpf 'cetqu'c'tcpi g'qh'vgo r gtcwt'gu'y kj 'v'j g'qpn\ 'ko r r'ckv'kqp'  
dgkpi 'v'j cv'ur cy plpi 'qeewtu'y j gp'ugc'utw'ceg'vgo r gtcwt'gu'ctg'cr r tqzko cvgn\ "47AE0"K'ugc"  
utw'ceg'vgo r gtcwt'gu'tkug.'v'j g'i gqi tcr j k'tcpi g'qh'v'j g'ur geku'o c { 'uj k'lp'v'j g'ur qpug'v'j cp { "  
ej cpi gu0'Qpg'qh'v'j g'q'v'j g'r qv'p'v'cn'gh'g'ew'qh'enko cvg'ej cpi g'eqwf 'tgr'v'v'v'j g'v'j g'mu'qh'utwewt'cn'  
j cdkcv'lp'v'j g'eqtcn'tggh'gequ'f'vgo u"\*O wfp c { 'gv'cr0422: +0"Qegcp'cekf k'k'v'kqp'ku'cv'kcr cvgf 'v'  
ch'gev'v'j g'lp'v'j tk\ 'qh'eqtcn'tggh'hu'cpf 'ej cpi kpi 'ugc'ng'gn'eqwf 'o qf kh\ 'v'j g'f gr v'j 'tgi ko g'y kj 'uwej "  
tcr kf kv\ 'v'j cv'eqtcn'cpf 'eqtcn'tggh'hu'km'dg'ch'gev'f "\*O wfp c { 'gv'cr0422: +0"K'petgcug'f'ugc'utw'ceg"  
vgo r gtcwt'gu'j cxg'dggp'tgur qpukdr'ht'eqtcn'mu'v'j tqwi j 'dngcej kpi 'cpf 'f'kugcug'cpf 'dkqgtqkqp'  
o c { 'tgf weg'5/f ko gpukpcn'utwewt'g'lp'ch'gev'f'ctgcu"\*C'rkctg' /Hkr 'gv'cr0422; + 'tgf w'kpi 'cf wv'  
j cdkcv'ht'P cuucw'i tqwr gt"\*Eqrgo cp'cpf 'M'qg'pki "4232."Tqi gtu'cpf 'Dggw'4223+0'Ej cpi gu'lp'  
tgr tqf w'v'kg'qwr w'qt'ugcu'pcn'v'ko kpi 'ctg'cnuq'r quukdr'y kj 'w'pnpqy p'eqpugs w'p'egu'ht"  
r qr w'v'kqp'cdwfp cpeg0"K'petgcug'f'i mdcn'vgo r gtcwt'gu'ctg'cnuq'r tgf kev'f 'v'j 'ej cpi g'r ctcukg/j quv'  
tgr'v'k'puj kr u'cpf 'o c { 'r t'gugpv'w'pnpqy p'eqpegtpu"\*J c'xg'm'gv'cr04224.'O cteqi rkug'4223+0"

**2.d. Limits to recruitment/depensation**

F gr gpuc'kqp.'cnuq'tghgt'gf 'v'cu'v'j g'C'mgg'gh'gev.'qeewtu'y j gp'v'j g'cdwfp cpeg'qt'f'gpukv\ "  
qh'lpf k'kf w'nu'f'q'r u'dgnqy 'c'et'k'v'cn'v'j t'g'j qf 'cpf 'tgr tqf w'v'kqp'dgeqo gu'lp'gh'gev'v'j g'lp'

"  
"  
"  
"



cu'Dgto wf c."qt'Rwgtvq'Tleq.'y j gtg'ci i tgi cvkqpu'pq'npqi gt'hqto . 'P cuucw'i tqwr gt'ctg'pqy 'tctgn' " vngp'qt'qdugtxgf 'cpf 'y g'qpn' 'tgr qt'u'qh'P cuucw'ctg'htqo 'y gug'tctg'hkuj gt { 'kpvgtcevkqpu' "

**2.e. Disease, parasites, and abnormalities**

Rctcukgu'qeewt'kp'dqvj 'y kf/ecwi j v'cpf 'ewnkxcvfg 'P cuucw'i tqwr gt.'r tgf qo kpcpwn' 'kp'yj g' xkuegtc'cpf 'i qpcf u0'Gpe { uvfg 'rctxci'vcr gy qto u'ctg'eqo o qp'kp'yj g'xkuegtc'cpf 'c'tgf f kuj 'dtqy p" pgo cvqf g'qeewtu'kp'yj g'i qpcf u'\*Vj qo ruqp'cpf 'O wptq'3; 9: +0'Rctcukle'kuqr qf u'ctg'hqwpf 'kp" pquutku'\*Vj qo ruqp'cpf 'O wptq'3; 9: +0'Vj g'f ki gpgvle'tgo cvqf g'*Helicometra tortc*'\*r { nqtke" ecgec+.'*Lecithochirum parvum*'cpf '*L. microstomum*'\*uqo cej +.'cpf '*Sterrhurus musculus*' \*uqo cej +y gtg'kf gpvkhgf 'kp'Hqtkf c/ecwi j v'hkuj '\*O cpvgt'3; 69.'Qxgtutggv'3; 8; +0'

F luggcu'cpf 'cdpqto crikgu'ctg'pqv'f guetkdgf 0'Cnj qwi j 'ugxgtcn'ur gekgu'qh'y guvtp" Cwvple'i tqwr gtu'ctg'npqy p'vq'dg'eki wcvzle '\*gur gekm' { 'y j gp'rcti g+.'P cuucw'i tqwr gtu'j cxg" dggp'yj qwi j v'vq'dg'wplhqto n' 'pqp/vzle'yj tqwi j qw'yj gk'tcpi g'\*J cnvgtf'3; 89.'Lqt { 'cpf 'Kgtuqp" 3; ; +y kj 'y g'kpvgtgukpi 'gzegr vqp'qh'qpg'uo cni'vzle'P cuucw'i tqwr gt'kp'yj g'Xkti kp'Kurcpf u" \*Dtqy pgm'cpf 'Tckpg' { '3; 93+0'Gzetguegpegu'y gtg'pqvfg 'qp'qvqkij u'cpf 'qpg'hkuj 'j cf 'c'eqo r ngvgn' o crhqto gf 'uci kwcnr ckt'y kj 'y g'y j qng'qh'yj g'eqpecxg'uwthceg'qxgti tqy p'y kj 'c'rcti g" gzetguegpeg'\*Vj qo ruqp'cpf 'O wptq'3; 9: +0'

**2.f. Aquaculture – successes, failures, potential threats**

Vj g'P cuucw'i tqwr gt'ku'eqpukf gtgf 'c'r tko g'ur gekgu'ht'cs wcewntg'\*Vwengt'3; ; 4c.'3; ; 4d+0" Kp'yj g'rcv'3; ; 2u'cpf 'kpqv'yj g'3; ; 2u.'eqpukf gtedrg'r tqi tgu'y cu'o cf g'kp'j cvej gt { 'ur cy plpi 'cpf " tgetkpi 'qh'i tqwr gtu'wpf gt'cs wetkwo 'eqpf kkpqu'\*Vwengt'3; ; 4c.'Y cvpcdg'gv'cn03; ; 7c.'3; ; 7d." Vwengt'gv'cn03; ; 8+0'

Hgo crg'P cuucw'i tqwr gtu'y gtg'lpf wegf 'v'q'qxwrcv'wukpi 'j wo cp'ej qtkpke'i qpcf qtqr kp" \*J EI +lplgevkqu.'nwgkpk kpi 'j qto qpg/tgrgculpi 'j qto qpg'cpcmqi '\*NJ TJ c+'cpf 'ectr 'r kwkct { " j qo qi gpcvg'\*ERJ +.'qt'eqo dkpcvkpu'yj gtgqh'\*Vwengt'3; ; 4d.'Mng { 'gv'cn03; ; 6.'Y cvpcdg'gv'cn0 3; ; 7d+0'Hgo crgu'y kj 'o gcp'qqe { 'v'f lco gvgtu'tcpi kpi 'htqo '6: 4/783'o letqo gvgtu'\* o +y gtg" uwkcdrg'ht'j qto qpg/lpf wegf 'ur cy plpi '\*Y cvpcdg'gv'cn03; ; 7d+0'Vwengt'gv'cn0\*3; ; 8+f guetkdgf " hqwt'o gyj qf u'ht'cej kexkpi 'hgtvkk gf 'gi i u.'kpenf kpi 'eqo dkpcvkpu'qh'lpf wegf 'qt'pcwtcn' qxwrcvqp'cpf 'ctv'kckcn'htvkk cvqp'y kj 'htguj 'o kn'qt'pcwtcn'ur cy plpi 'kp'vcpm0'

Hgtvkk cvqp'tcvgu'kp'ctv'kckcm' 'lpf wegf 'ur cy pu'tcpi gf 'htqo '3: /322' 'cpf 'j cvej kpi " uweegu'tcpi gf 'htqo '8: /322' '\*J gcf 'gv'cn03; ; 8.'Vwengt'gv'cn03; ; 8+0'O wnr ng'ur cy pu'qeewt'gf " qp'eqpugew'xg'f c { u'cpf 'j cvej gt { 'tgetgf 'lwxgpkng'P cuucw'i tqwr gtu'i tgy 'v'307/'40'mi 'kp'4' { gctu" \*Vwengt'cpf 'Y qqf y ctf '3; ; 5+0'

Hqmjy kpi 'j qto qpg'lplgevkqu.'P cuucw'i tqwr gt'hgo crgu'r tqf wegf 'enwej gu'qh'dgy ggp" 45.222'cpf '822.222'o cwtg'gi i u'r gt'mi 'qh'dqf { 'y gli j v'y kj 'rcti g'hgo crgu'ecr cdrg'qh' { kgrf kpi " cm quv'7.222.222'gi i u0'Mng { 'gv'cn0\*3; ; 6+'tgr qt'vfg 'qpg'v'q'y q'enwej gu'r tqf wegf 'f wtkpi 'yj g" pcwtcn'tgr tqf we'xg'ugcuqp.'y kj 'gcej 'enwej 'v'qvrkpi '72.222/822.222'gi i u'r gt'nrqi tco 'dqf { "

"  
"  
"  
"

y gki j x0J gcf "gv'cr0\*3; ; 8+hqwpf "vj cv'hgo crgu'eqwf "ur cy p"vy q"vq"vj tgg"vko gu'cv'lpvgtxcnu'qh'4: " vq'97"fc {u."r tqf wekpi "422.222/"4.222.222"gi i u'r gt'hgo crg"\*76.222"cpf "562.222"gi i ulni "dqf { " y gki j v'y kj "hgo crgu'tcpi kpi "kp"uk g'ltqo "507/80 "mi 0"Vwengt"gv'cr0\*3; ; 3+pqvfg "enwej gu'qh" 722.222"vq"922.222"ht "hgo crgu'tcpi kpi "ltqo "5/7"mi "388.888"vq"362.222"gi i ulni + "y j krg" Y cvpcdg"gv'cr0\*3; ; 7d+tg r qtvgf "utkr r gf "hgo crgu'qh"604/34"mi "tgrgculpi "dgw ggp"; 7.222"cpf " 6.972.222"gi i u"\*44.83; /5; 7.: 55"gi i ulni + "y kj "c"uki pkhcepv'tgrv'kpuj kr "dgw ggp"dqf { "y gki j v" cpf "gi i u'utkr r gf " \*{ "?" 205: 7z/2077: ; =t4"? 2062."p"? "63."r >2023={ "ku"gi i u'utkr r gf "cpf "z "ku"dqf { " y gki j v'kp"mi +0"

Nctxcn'utwxkcn'vq'htuv'hggf kpi "y cu'i gpgtcm { "j ki j . "y kj "f gerkpgu"vj gtgchgt "f gr gpf kpi "qp" hggf kpi "tgi ko g0"Uwtxkcn'qh'rtxcg"vq'htuv'hggf kpi "kp"qpg'ugv'qh'gZR gtko gpw'y cu'87' "Vwengt" 3; ; 4d+dw'y cu'hqwpf "vq" f gerkp"vq"cdqw'3' "d { "f c { "84"r quv'j vej kpi "kp"cpqy gt "Y cvpcdg"gv'cr0 3; ; 6."3; ; 8+rtxcn'utwxkcn'f gerkp"vq" qpeg"vj g" { qmluce "y cu'cduqtdgf 0"Hggf kpi "y kj "q { uvgt" vqej qr j qtgu"cpf "ukxgf "tq'vhtu."eqo dlpf . "cej kxgf "j ki j gt'rtxcn'utwxkcn'cvgu"vj cp'hggf kpi " y kj "vpukxgf "tq'vhtu"cmppg"Y cvpcdg"gv'cr03; ; 6+cpf "uo cni'r tg { "uk g'y cu'ko r qtvcv"Y cvpcdg" gv'cr03; ; 8+0"Tuwmu'qh'hggf kpi "gZR gtko gpw'lpf kecvgf "vj cv'ewwtgf "lwxgpkrgu'tgs vkt g'c" f kvct { " r tqvkl'rgxgn'cdqxg'77' "cpf "cp"gpgti { /vq/r tqvkl'tcvk'qh'dgny "4: Q "nLi "hqt"qr vko wo "i tqy vj " \*Gmku"gv'cr03; ; 8+0"Eqpv'tn'qh'wtdwrgpeg."ucrkp { . "cpf "iki j v'kpvuk { "ko r tqxgu'utwxkcn'vq"vj g" htuv'hggf kpi "uci g" \*Gmku"gv'cr03; ; 9d+0"Kpetgcugf "i tqy vj "cpf "hggf kpi "cvgu'qeevttgf "y kj " kpetgcugf "y cvgt "vgo r gtcwtgu" \*Gmku"gv'cr03; ; 9c+0"

GZR gtko gpw'vq" f gvgt o kpg"vj g'uweegu'tcvg'qh'rtxcn'P cuucw'i tqwr gt "ewwtg" \*Y cvpcdg"gv' cr03; ; 7c."3; ; 7d+cpf "utwxkcn'qh'tgrcugf "j vej gt { /tgctgf "lwxgpkrgu" \*T qdgtu"gv'cr03; ; 6+ "j cxg" dggp"eqpf wevgf 0" Cnj qwi j "vgo r gtcwtg" o cplr wrcvqp" o ki j v'dg" wugf "vq" eqpf kkp "P cuucw'i tqwr gt" vq"ur cy p"cp { "o qpj "qh"vj g" { gct " \*Vwengt"gv'cr03; ; 8+ "j vej kpi "uweegu'y cu"j ki j gt "dgw ggp"48/" 4: Å "eqo r ctgf "vq"j vej kpi "cv52Å " \*Y cvpcdg"gv'cr03; ; 7d+0" P cuucw'i tqwr gt "lwxgpkrgu" \*52; /" 589" o o "VN+tgctgf "ltqo "gi i u" \*p"? "49+cv"J ctdqt "Dtcpej "y gtg" wugf "vq" vgu'vj g' hgcuklkr { "qh" tguqen kpi "tgg" \*T qdgtu"gv'cr03; ; 6+lp" U0Vj qo cu0" F gur kg"uqo g" o qtcrk { "cpf "f kur gtucn "c" hgy " vci i gf "huj "y gtg" qdugt xgf "vr "vq" plpg" o qpj u'chgt "tgrcug0" Vj g'r qv'p'v'cn'qh'P cuucw'i tqwr gt" uqem'gpj cpego gpv."cu'y kj "cp { "qy gt" i tqwr gt "ur geku."j cu" { gv'vq" dg" f gvgt o kpgf " \*T qdgtu"gv'cr0 3; ; 6+0" Ugtkqu'eqpegtpu'cdqw'vj g'i gpgv "eqpugs vqpegu'qh'kpv'qf wv'kpu'cpf "cdqw'r quukrg" r tqdrgo u'qh'lwxgpkrg"j cdkc'v'cxck'cdkrk { . "kpv'qf wv'kpu"qh'o crcf er vgf "kp' kxk' wcu."qt "kpcdkr { "vq" mqecv'v'cf kkp'cn'ur cy p'kpi "ci i tgi cv'kpu."eqv'kpwg"vq" dg'tckugf 0 "

### 3. DESCRIPTION OF FISHERIES AND FISHERIES MANAGEMENT AND CONSERVATION

F cv'qp'tgetvko gpv'lpvq"vj g'huj gt { "lpf kecvg"vj cv'ci g"cpf "uk g'htuv'uuegr vldrg"vq" ecr wtg" ctg"6/9" { gctu"cpf "497- "o o "VN."tgr ge'v'xgn { 0" k' uqo g'ctgcu."o quv'qh'vj g'ecvej "ku."qt"j cu'dggp." eqo r qugf "qh'lwxgpkrgu" \*g { 0Rwgt vq "T leq"cpf "Ewdc+ "Rwgt vq "T leq "Huj gt kgu" T gugctej "Ncdqtcvqt { " 3; ; 3."Emtq"gv'cr03; ; 2+0" Qnugp"cpf "NcRwreg" \*3; 9; +ecr ewrcvgf "ci g'qh'htuv'ecr wtg"cv'6/7" { gctu."

"  
"  
"  
"

cnj qwi j 'ko o cwtg'hkuj 'qh'4"{gctu">522"o o "VN+y gtg'cnuq'tgetwkgf 0"O gcp'uk' g'qh'tgetwko gpv'  
kpv'v'j g'hkuj gt { 'kp'Lco ckec'y cu'guko cvgf "cv792"o o "VN"\*cdqw7" { gctu'qrf '+qp'qegcple'dcpmi'hqt "  
j cpf rkgp'cpf 'hkuj 'vcr 'hkuj g'kgu=v'j g'o kpk wo 'hpi v' 'ecr wtgf 'y cu'497"o o "VN'cpf 'v' g'hwni  
tgvppkqp'hpi v' 'y cu'847"o o "VN"\*Vj qo ruqp'cpf "O wptq"3; 9: +0"O qf cnci gu'tgr qtvgf 'hqt"c"  
Ec{o cp"Kucpf'u'ci i tgi cvkqp'cpf "c'uvqeni'k'Ewdc'y gtg'8/: "{gctu"\*Erctq'gv'cr03; ; 2."Dwuj "gv'cr0'  
4228+.'uwi i gukpi 'v' cv'kpf kxf wcu'y gtg'pqv'hwni { 'tgetwkgf 'wv'k'v'j ku'ci g'tcpi g0"

### 3.a. Abundance indices and trends over time

*Stock assessments.* Hgy 'hqtto cni'uvqeni'cuuguo gpw'j cxg'dggp'eqpf wevgf 'hqt'v'j g'P cuucw'i tqwr gt." rkngn' 'dgecwug'qh'iko kxf 'f cvc0"Vj g'o quv'tgegpv'r wdrkuj gf "cuuguo gpv."eqpf wevgf 'kp'v'j g' Dcj co cu. 'kpf kecvgu'hkuj kpi 'ghqtv'k'v'j g'Dcj co cu'pggf u'v'q'dg'tgf wegf 'htqo 'v'j g'3; ; : 'v'q'4223" r'xgn'qv'j gty kug'v'j g'uvqeni'ctg'rkngn' { 'v'q'dg'qxgtg'zr r'k'kgf 't'gnc'v'x'g'v'q'dk'q'ni kecn't'ghgt'gpeg'r q'k'p'w0" \*Ej gwpi "gv'cr04235+0"Vj g'r qr wcvkqp'f { pco ke'o qf g'kpi 'd { 'Ej gwpi "gv'cr04235+'h'q'wpf < o'cuwo kpi 'v'j cv'v'j g'em'uw'g'qh'v'j g'ur cy p'kpi 'ci i tgi cvkqp'ugcuqp'ku'r gthgevn' 'ko r ngo gpv'gf 'cpf " gphqtegf . 'v'j g'o gf kcp'x'c'w'g'qh'FURT"? '57' 'qp'pqp/ur cy p'kpi 'hkuj 'y qwf 'dg'72' 'qh'v'j g'hkuj kpi " o qt'v'k'v' { 'qh'v'j g'3; ; : 'v'q'4223' r'xgn'0"Vj g'7' 'cpf "; 7' 'eqph'k'f gpeg'iko ku'ct'g'guko cvgf 'v'q'dg'rguu' v'j cp'42' 'cpf "o qt'g'v'j cp'322' 'qh'v'j g'hkuj kpi "o qt'v'k'v' { 'cv'v'j g'3; ; : 'v'q'4223' r'xgn't'gur ge'v'x'gn' { 0" 'k'p'qv'j gt'y qtf u.'k'h'3+'hkuj kpi "o qt'v'k'v' { 't'cv'g'qh'pqp/ur cy p'kpi 'hkuj 'ct'g'o c'k'p'v'k'p'gf 'cv'v'j g'3; ; : 'v'q' 4223' r'xgn'cpf "4+'hkuj kpi "qp'ur cy p'kpi 'ci i tgi cvkqp'u'k'p'gi r'ki kdr. 'v'j g'o gf kcp'ur cy p'kpi " r qv'p'kcn'ur cy pgt'dkqo cu't'gnc'v'x'g'v'j g'v'p'g'zr r'k'kgf 'r'xgn'ku'g'zr ge'vgf 'v'q'dg'ct'q'wpf '47' '\*7" cpf "; 7' 'EKqh'42'cpf '52' . 't'gur ge'v'x'gn' { +0"Vj ku'r'xgn'ku'ki p'k'k'ec'p'v' { 'd'g'm'y 'v'j g't'ghgt'gpeg'iko k'v'q'h' 57' 'qh'ur cy p'kpi 'r qv'p'kcn'o gcp'kpi 'v'j cv'v'j gt'g'ku'c'j ki j 'ej c'peg'qh'tgetwko gpv'qxgthkuj kpi " dgecwug'qh'v'j g'hqy 'ur cy p'kpi 'uvqeni'dkqo cuu0"

F wtkpi 'v'j g'htu'WU0uw'x'g' { 'qh'v'j g'hkuj gt { 't'gu'wtegu'qh'Rwgt'v'q'Tleq.'v'j g'P cuucw'i tqwr gt" y cu'p'q'vgf 'cu'c'eqo o qp'cpf 'xgt { 'ko r qt'v'p'v'h'q'f 'hkuj . 't'ge'j kpi 'c'y g'ki j v'qh'72'ndu0\*440'mi '+qt" o qt'g'Gx'gto c'pp'3; 22+0'D { '3; 92."P cuucw'i tqwr gt'y cu'uv'k'v'j g'h'q'w'v'j "o quv'eqo o qp'uj cmqy / y cvgt'ur ge'k'g'u'rc'p'f gf 'k'p'Rwgt'v'q'Tleq'\*Vj qo ruqp'3; 9: +.'cpf 'k'v'y cu'eqo o qp'k'p'v'j g't'ggh'hkuj " hkuj gt { 'qh'v'j g'X'k'i k'p'Kucpf'u.'y j gt'g'cp'ci i tgi cvkqp'k'p'v'j g'3; 92u'eq'p'v'k'p'gf "cp'guko cvgf '4.222/ 5.222'k'p'f k'k'f wcu'\*Qm'gp'cpf "Nc'R'rc'eg'3; 9; +0"F wtkpi 'v'j g'3; ; 2u.'r qt'v'uco r r'kpi 'k'p'v'j g'WU0X00' u'j qy gf 'v'j cv'P cuucw'i tqwr gt'cee'q'w'p'vgf 'hqt'44'r gte'gp'v'q'h'i tqwr gt'rc'p'f kpi u'y k'j ": 7'r gte'gp'v'q'h'v'j g' P cuucw'i tqwr gt'ec'v'j 'eqo kpi 'htqo 'ur cy p'kpi 'ci i tgi cvkqp'u'F0Qm'gp.'Ej k'gh'Ue'k'p'v'k'u'o'U0' Vj qo cu'Hkuj gto gp'u'Cu'q'ek'v'k'p.'r gtu0'eqo o 0'v'q'L0T'v'g'v'g't.'P O HU.'Qev'q'dgt.'4235+0'D { '3; ; 3." o'v'j g'P cuucw'i tqwr gt'j c'f +r t'ce'v'k'ec'm' { 'f'k'uc'r r gct'gf 'htqo 'v'j g'h'q'ec'v'j gu'c'p'f 'v'j g'q'p'g'u'v'j cv'f 'k'f + cr r gct '\*y gt'g'+uo cm'le'qo r ct'gf 'y k'j 'r t'g'x'k'q'w' { gctu0'\*EHO E'3; ; 7+'cpf "d { '3; ; 8.'v'j g'P cuucw' i tqwr gt'y cu'eq'p'k'f gt'gf 'eqo o gte'k'cm' { 'g'z'v'p'ev'k'p'v'j g'WU0X'k'i k'p'Kucpf'u'r'w'g't'v'q'Tleq't'gi k'q'p' " \*D'q'j p'uc'eni'g'v'cr03; ; 8+0"C'd'q'w'3.222'mi 'y gt'g'rc'p'f gf 'htqo 'v'j g'T'ggh'Hkuj 'hkuj gt { 'f wtkpi 'v'j g'r'c'w'g't' j c'h'q'h'v'j g'3; ; 2u'k'p'Rwgt'v'q'Tleq.'o quv'q'h'v'j go 'y gt'g'rguu'v'j cp'722"o o . 'k'p'f k'ec'v'kpi 'v'j g' { 'y gt'g' rkngn' 'ugz'w'cm' { 'ko o cwt'g'\*U'c'f q'x { '3; ; 9+0"

Nk'w'g'ku'np'qy p'cd'q'w'v'j g'f { pco keu'qh'v'p'g'zr r'k'kgf 'uvqeni'qh'P cuucw'i tqwr gt'cm'j qwi j " uqo g'qh'v'j g'f cv'htqo 'v'j g'3; ; 2u'i k'x'g'wu'uqo g'k'p'uki j v'\*E'ct'v'g't'gv'cr03; ; 6+0"Ur cy p'kpi 'uvqeni'

"  
"  
"  
"

dlqo cuu'r gt'tgetwk'j cu'pqv'dggp's wcpv'k'gf 'hqt'v'j g'ur gekgu'dw'h'cpf kpi u'f c'c'engctn' 'uj qy "c" ej tqpqm'j kecn'v'gpf 'htqo "cdw'p'f cpeg'v'q'tctk'v' 'kp'o cp{' "ctgcu"\*g'f 0"Ucf qx { "3; ; 9+0"Qh'r ct'v'ewrct" eqpegtp'j cu'dggp'v'j g'tcr kf "cpf "gzv'tgo g'f genk'p'g'lp'pwo dgtu'vcngp'htqo "tcf k'k'qpcn'ci i tgi cv'k'qp" uk'gu"\*Ucr'g'v'cr'04223+0"K'p'i gpgten'urqy /i tqy kpi . 'n'p'i /r'k'x'gf 'ur gekgu"\*uwej "cu'upcr r gtu'cpf " i tqwr gtu'+y kj "rko k'gf 'ur cy pl'pi "r g'k'qf u'cpf . 'r quukdn' . 'y kj "qpn' { "c"pcttqy "tgetwko gp'v'y k'p'f qy " ctg'uwuegr v'drg'v'q'x'g'tzr n'k'c'v'k'qp"\*Dcpgtq'v'g'v'cr'03; ; 9. 'R'q'n'x'k'p'c'cpf "T'c'n'v'q'p'3; ; 9+0"J qf i uqp" cpf "N'k'gd'g'ngt"\*4224+'p'q'v'f 'v'j cv'P cuucw'i tqwr gt'y g'tg'cdugp'v'htqo " : 4 "qh'uj cmqy "E'ct'k'dd'g'cp't'g'ghu" \*5632o +f w'k'p'i "c'7/ { gct'r g'k'qf "\*"3; ; 9/4223+"q'h'w'p'f gty cvgt'uw'x'g' { u'h'qt'v'j g'T'g'g'h'E'j gen'r t'q'l'g'ev'0" Vj ku'ku'f g'k'x'gf 'htqo "w'p'f gty cvgt'uw'x'g' { u'k'p'o quv'e'q'w'v'k'gu'k'p'v'j g't'c'p'i g'q'h'v'j g'ur gekgu'0"

M'p'qy p'ur cy pl'pi "ci i tgi cv'k'qpu'q'h'P cuucw'i tqwr gt'ctg'f'kur m { gf 'k'p'H'k'i v'tg' ; "cu'cx'k'x'd'rg" k'p'r w'd'r'k'uj gf "cpf "i t'c { "r'k'g't'c'w't'g'cpf "k'p'v't'x'k'gy u"\*Ucf qx { "f'g'O ke'j guqp'g'v'cr'0422: +0"F'c'v'c'j' c'x'g" dggp'ct'ej k'x'gf "k'p'v'j g'ue't'h'c'q'iti "y g'duk'g'f'c'v'c'd'c'ug'0"Vj g'o cr "uj qy u'cm'h'p'qy p'ci i tgi cv'k'qpu" t'gr q't'v'f "v'q'g'z'k'v'uk'peg'3: : 6\*c+0"K'p'v'j g'h'gy "ec'ug'u'y j g'tg'ci i tgi cv'k'qp'p'wo dgtu'y g'tg'g'u'k'o cv'g'f . " cdw'p'f cpegu't'c'p'i gf "htqo "cr r tqzko cv'gn' "32.222"v'q'u'qo gy j g'tg'd'g'y ggp'52.222"cpf "322.222"h'k'uj " \*Uo k'j "3; 94. "Q'u'ng'p'cpf "N'c'R'nc'g'3; 9; . "E'q'n'k'p'g'v'cr'03; ; 9. "H'k'p'g'3; ; 2.'3; ; 4. "E'ct'v'g't'g'v'cr'03; ; 6." Ucf qx { "3; ; 9+0"ht'eqo r ct'k'q'p. "k'v'c'n'q'uj qy u'v'j q'ug'ci i tgi cv'k'qp'uk'gu't'gr q't'v'f "v'q'g'z'k'v'cu'q'h" cdq'w'4229"\*d+0"Vj g'em'q'ug'f "ekt'eng'u't'gr t'g'ug'p'v'uk'gu'd'g'n'k'x'g'f "v'q'g'z'k'v.'y kj "h'k'uj "p'wo dgtu'g'u'k'o cv'g'f " cv'd'g'y ggp'322"cpf "5222"\*g'u'k'o cv'gu'htqo "h'k'uj k'p'i "cpf "f'k't'g'ev'q'd'ug't'x'c'v'k'qpu+0"Vj g'q'r gp'ekt'eng'u" t'gr t'g'ug'p'v'uk'gu'k'p'E'w'd'c'v'k'n'i'd'g'n'k'x'g'f "v'q'r tqf w'eg'uo cm'ec'v'ej gu'q'h'P cuucw'i tqwr gt'dw'uk'gu'j' c'x'g" p'q'v'd'ggp'cu'gu'ug'f "f'k't'g'ev'0"

Y j k'g'j' g'c'x { "h'k'uj k'p'i "q'p'ur cy pl'pi "ci i tgi cv'k'qpu'o c { "j' c'x'g'd'ggp'c'r t'ko ct { "f't'k'x'g't'q'h" r'qr w'v'k'q'p'f' gen'k'p'gu"\*Ucf qx { "f'g'O ke'j guqp'cpf "G't'k'uo cp"4234+."q'y gt'h'c'ev'qt'u'o c { "ch'g'ev" r'qr w'v'k'q'pu'c'v'c'p'c'v'k'q'p'c'n'g'x'g'r'0"J g'c'x { "h'k'uj k'p'i "q'h'c'f'w'w'u'cy c { "htqo "qt'f'w'k'p'i "ur cy pl'pi "t'w'p'u.'v'j g' k'p'v'g'p'uk'x'g'ec'r w't'g'q'h'l'w'x'g'p'k'gu."g'k'j gt'v'j tqw'j "f'k't'g'ev'v'ct'i g'v'k'p'i "\*"g'f 0"ur g'ct'h'k'uj k'p'i +qt'w'uk'p'i "uo cm' o guj "t'cr u'qt'p'g'w.'y k'n'le'qo r tqo k'ug'r'qr w'v'k'q'p'w'c'd'k'k'v' { "cpf "ur cy pl'pi "r'q'v'g'p'v'c'n'cpf "n'q'u'qt" f'gi t'cf cv'k'qp'q'h'j' c'd'k'c'v'eq'w'f "ch'g'ev'r'qr w'v'k'q'pu'd'g'ec'w'ug't'g'g'h'cu'q'ek'cv'g'f "j' c'd'k'c'w'ct'g'w'ug'f "cu" u'j g'ng't'c'v'c'm'i'h'k'g'j' k'v'q't { "r'j' c'ug'u'o c { "cm'j' c'x'g'f'g'v't'ko g'p'v'c'n'g'h'g'ev'u"\*g'f 0"U'go o g'p'u'g'v'cr'04229c+." v'j q'w'j "k'v'ku'p'q'v'eng't'k'h'q'p'g'h'c'ev'q't'ku'o q't'g'f'g'v't'ko g'p'v'c'n'v'j cp'v'j g'q'v'j gtu."qt'k'h'v'j g'ug'f'g'ng'v't'k'q'w'u" g'h'g'ev'u'y q't'm'k'p'eqo d'k'p'c'v'k'q'p'0""

"

"  
"  
"  
"




Figure 10. Maps showing locations of known Nassau grouper spawning aggregations both historically (a) and as of about 2007 (b) according to available information-not all sites have been validated. Inset shows full geographic range, main concentrations (shaded) and extended areas (dashed lines). Each closed circle represents 1, or occasionally 2, reported site(s). Open circles are “probable” sites. (Sources: Smith 1972; Sadovy and Eklund 1999; Sala et al. 2001; Whaylen et al. 2004; Belize Spawning Aggregation Working Group, unpublished data; R. Claro, unpublished data; E. Sala, unpublished data, as presented in Sadovy de Mitcheson et al. 2008)

*Ecological assessments.* "Vj g'P cuucw'i tqwr gt'y cu'hqto gtn{ "qpg"qh'y g'o quv'eqo o qp"cpf " ko r qtvcv'eqo o gtekrn' tqwr gtu'kp'y g'kpuwrc'tqwr kcrn'y guvgtp"Cvrcpvk"cpf "Ectkddgcp"\*Uo kj " 3; 9: .Tcpf cm'3; : 5.'Cr r gnf qqtp'gv'cn'03; : 9.'Ucf qx{ '3; ; 9:0"F genkpgu'kp"ncpf kpi u."ecvej 'r gt'wplv' 63"

"  
"  
"  
"

qh'ghqtv\*ERWG+. 'cpf . 'd { 'lo r nkecvkp. 'cdwfp cpeg. 'j cxg'dggp'tgr qtvgf 'y tqwi j qw'ku'tcpi g. 'cpf 'k' ku'pqy 'eqpukf gtgf 'vq'dg'eqo o gtekmf 'gz vkev\* 'y g'ur gekgu'ku'gz vkev'ht 'hkuj gt { 'r vtr qugu'f wg'vq' nqy 'ecvej 'r gt 'wplv'ghqtv+'lp'c'pwo dgt'qh'tgcu0'kphqto cvkp'qp'r cuv'cpf 'r t gupv'cdwfp cpeg'cpf " f gpukv { . 'lp'dqv 'ci i tgi cvkp'cpf 'pqp/ci i tgi cvkp'j edkcv.'ku'dcugf "qp'c'eqo dlpvcv'qh'cpgef qvni' ceeqwpv. 'xkuwcn'egpuwu'uwtxg { u. 'cpf 'hkuj g'kgu'f cvc0"Vj g'rceni'qh'ur gekgu'ngxgr'hkuj g'kgu'f cvc" ugxgtgnf 'rko ku'hkuj gt { 'f gr gpf gpv'cpcnf uku'qh'v' g'ur gekgu'v' tqwi j qw'ku'tcpi g0'Hkuj gt { " kpf gr gpf gpv'uwtxg { u'r tqxkf g'y g'qpnf 'dtqcf 'uecrg'f cvc'y kj 'y j lej 'vq'ceegu'ewtgpv'r qr wrcv'kp' eqpf kkp0"Uwej 'uwf kgu'ctg'tghgt'ppegf 'lp'y g'hqmy kpi 'ugev'kpu. 'cu'cxckrdrg0'Wphqtwpvcgnf . " vko g'ugt'kgu'f cvc'ctg'i gpgtcmf 'rcenkpi 'cpf 'eqo r ct'ku'pu'dgy ggp'tgghu'qt'dgy ggp'eqwv'kgu'ctg' vj g'qpnf 'r quukdr'g'y c { u'vq'eqo r ctg'cu'c'o gcuwtg'qh'tgr'v'xg'cdwfp cpeg0

C'pwo dgt'qh'qti cpl cvkp'qt'ci gpek'j cxg'w'p'f gt vcn'p'uwtxg { u'vq'gr'w'ekf cvg'y g'uw'wu'qh' eqtcn'tgghu'cpf 'tgg'h'hkuj 'r qr wrcv'kpu'v' tqwi j qw'v'j g'y guv'gtp'c'v'p'v'k. 'cu'y gni'cu'v'j gt 'r ct'w'qh'v'j g' y qtrf 0'T guwu'htqo 'v'j gug'o qpkqt'kpi 'uwf kgu'\*Mgnku'p'g'v'c'0422; +q'ht'g'uo g'kpf kcv'k'p'qh' tgr'v'xg'cdwfp cpeg'lp'xct'k'wu'ncv'k'p'u'ht'P cuucw'i tqwr gt '\*Vcdrg'32+. 'cnj qwi j 'i gpgtcmf " f k'ht'g'p'v'o g'j qf u'ctg'go r nq { gf 'cpf 't'gu'wu'ecppq'v'dg'f'kt'g'w'f 'eqo r ct'gf 0'Uki j v'kpi 'ht'gs w'p'e { 'cpf " f gpukv { 'o c { 'q'ht'g'k'p'ht'o cvkp'0'T guwu'htqo 'c'v'p'v'k'cpf 'I wh'T'cr'k'f 'T'ggh'Cu'gu'uo gpv'Rt'qi t'co " \*CI TTC+'uj qy 'hgy 'P cuucw'i tqwr gtu'v'j tqwi j qw'v'j g'k'uwtxg { u0"Vj g'uki j v'kpi 'ht'gs w'p'e { " \*r tq'rt'v'k'p'qh'c'm'uwtxg { u'y kj 'cv'ng'cu'v'p'g'P cuucw'r t'g'p'v'v'v'c'p'i gf 'ht'qo 'h'gu'v'j cp'3' 'vq'ng'uu'v'j cp' 32' 0'F gpuk'k'gu'y q'w'f 'uecrg'w'r 'vq't'c'p'i g'ht'qo '3'vq'37'hkuj lj gev't'g'y kj 'c'o gcp'qh'708'hkuj lj gev't'g' cet'qu'u'c'm'ct'g'cu'uwtxg { gf 0'P QCC'au'E'qtcn'T'ggh'Ge'qu { u'vgo 'O qpkqt'kpi 'Rt'qi t'co \*ETGO R+'j cu' eqpf v'ev'g'f 'uwf kgu'lp'R'w'gt'v'q'T'leq'cpf 'v'j g'WUOX'k'i k'p'K'ur'p'f u'uk'p'eg'4222'cpf 'uki j v'kpi 'ht'gs w'p'e { " j cu't'c'p'i gf 'ht'qo '2'vq'20' 'cpf 'f'gpukv { 'j cu't'c'p'i gf 'ht'qo '2'vq'20'hkuj lj gev't'g'0'F cvc'ht'qo " W'p'k'g't'ukv { 'qh'v'j g'X'k'i k'p'K'ur'p'f u' \*WX'KX'ku'0'Uw'0'uc'o r'k'p'i 'cu'r'ct'v'qh'v'j g'k'lw'k'uf'k'v'k'p'c'n'f'eqtcn't'ggh' o qpkqt'kpi '\*h'w'p'f'gf'd { 'v'j g'P QCC'E'qtcn'T'ggh'E'q'p'ug't'x'cv'k'p'Rt'qi t'co +. 'y cu'p'q'v't'g'cf'k'f 'cxckrdrg' vq'eqo r w'g'uki j v'kpi 'ht'gs w'p'e { 'dw'f'gpuk'k'gu'y g't'g'6'hkuj lj gev't'g'0'P QCC'au' \*P O HU'HT'XE+'cpf " H'q't'k'f'c'au'Hkuj 'cpf 'Y'k'f'r'h'g'E'q'p'ug't'x'cv'k'p'E'q'o o k'uk'p' \*HHY EE 'X'ku'0'Uw'0'uw'f'k'gu'v'j cv'h'q'ew'u'q'p' v'j g'H'q't'k'f'c'M'g { u'k'p'f'k'ev'g'uki j v'kpi 'ht'gs w'p'e'k'gu't'c'p'i gf 'd'gy ggp'4/32' =f'gpuk'k'gu'ht'qo 'd'qv'j " uw'f'k'gu'y g't'g'3'hkuj lj gev't'g' \*Vcdrg'32-0'Dg { q'p'f 'v'j gug'o qpkqt'kpi 'uwtxg { u.'v'j g'T'ggh' G'p'x'k'q'p'o g'p'v'cn'G'f'w'cv'k'p'H'q'w'p'f'cv'k'p' \*TGGH+'ur'q'p'u'q't'u'cpf 'u'w'r'q't'v'u'x'q'w'p'v'g't'f'k'x'g/d'c'ug'f " uwtxg { u'cet'qu'u'v'j g't'g'i k'p'0'Q'd'ug't'x'g't'u't'c'p'ni'ur' gekgu'cdwfp cpeg'cu'U'k'p'i ng'?'3. 'Hgy' '?'4/32. 'O cp { '?' " 33/322. 'cpf 'C'dw'p'f'cpv'?'q'x'g't'322't'c'v'j g't'v'j cp't'g'eq't'f'k'p'i 'r't'g'ek'ug'p'w'o d'g't'u'0"Vj g'f'c'v'c't'g'v'j gp' ec'r'w'v'g'f'cu'c'F'gpukv { 'k'p'f'gz \*F'gp+'y j lej 'ku'c'o gcuwtg'qh't'gr'v'xg'cdwfp cpeg'y j gp'v'j g'ur gekgu' ku'uggp'dw'f'q'gu'p'q'v'i k'x'g'cp'k'p'f'k'ev'k'p'qh'rceni'qh'q'ee'w't'g'p'eg'cpf 'cu'c'U'ki j v'kpi 'H'ht'gs w'p'e { '\* UH+: " y j lej 'ku'c'o gcuwtg'qh'j'qy 'q'h'ng'p'v'j g'ur gekgu'y cu'q'd'ug't'x'g'f'0"Vj g'F'gp'cpf " UH'ue'q't'g'u'eq'w'f'd'g' o w'nr'k'g'f'vq'r't'q'x'k'f'g'c'o gcuwtg'qh'ur gekgu'cdwfp cpeg. 'y j lej 'ceeqw'p'u'ht'q'j' g't'q'q'd'ug't'x'cv'k'p'u'0" Y j g't'g'TGGH'uwtxg { 'k'p'ht'o cvkp'ku'cxckrdrg. 'k'ku'k'p'ev'f'gf'lp'y'j'g'h'q'm'y k'p'i 'E'q'w'p't { 'C'ee'q'w'p'u'0" Vj g'f'c'v'c't'g'p'q'v'p'g'eg'u'c't'k'f'cu'g'cu { 'v'q'k'p'v'g't'r'g'v'cu'f'g'uk't'gf'y kj qw'cf'f'k'k'q'p'c'n'ur'cv'k'c'n'f'eq'p'v'g'v' \*g'f'0'o c'p'c'i go g'p'v't'g'i k'o g+0

"  
"  
"  
"

**Table10. Fishery Independent Surveys from various sources. Sighting Frequency is the number of surveys in which at least one Nassau grouper was encountered; Density is the total number counted per unit area, standardized by area of each survey type. AGRRR info: Kramer 2003.**

Survey	Location/Extent	Year	Num. E. Stri. Observed	Num. Surveys w/ E. Stri.	Total Num. of Surveys	Sighting Frequency	Density (Num/m <sup>2</sup> )
AGRRR	Andros Island, Bahamas	1998	23	23	295	0.078	0.0013
AGRRR	Abaco Islands, Bahamas	1999	4	4	130	0.031	0.0005
AGRRR	Lighthouse Atoll, Belize	1999	1	1	110	0.009	0.0002
AGRRR	Glovers, Turneffe, Barrier Reefs, Belize	2000	6	6	349	0.017	0.0003
AGRRR	Little and Grand Cayman	1999	23	20	341	0.059	0.0011
AGRRR	Batabano, Cuba	2001	29	27	686	0.039	0.0007
AGRRR	Sabana and Camaguey, Cuba	2001	6	6	368	0.016	0.0003
AGRRR	Jardines de la Reina, Cuba	2001	7	7	535	0.013	0.0002
AGRRR	Boca del Toro and Comarca de Kuna, Panama	2002	4	4	451	0.009	0.0001
AGRRR	Caicos, Turks, and Mouchair Banks, Turks and Caicos	1999	25	25	279	0.09	0.0015
AGRRR	Culebra, Vieques, and Cayos de la Cordillera, PR	2003	2	2	174	0.011	0.0002
CREMP	La Parguera, PR	Average 2000-2007	2	2	1010	0.002	0.000025
CREMP**	Vieques, PR	2007	0	0	75	0	0
AGRRR	Biscayne National Park and Keys NMS, Florida	2003	8	7	381	0.018	0.0003
FFWCC Vis. Sur.	Keys NMS (Key Largo to Key West)	Average 1999-2007	79	76	7396	0.01	0.0001
NMFS FRVC	Keys NMS (Key Largo to Dry Tortugas)	Average 2000-2007	210	198	8563	0.0208	0.0001
AGRRR	St Croix, St Thomas, USVI and Guana, BVI	1999	1	1	144	0.007	0.0001
AGRRR	St Thomas, St John, USVI and Aneгада, Virgin Gorda, BVI	2000	6	6	100	0.06	0.001
UVI Vis. Sur.	St. Thomas, USVI	Average 2003-2007	8	N/A*	290	N/A*	0.0004
CREMP USVI	St. John and St. Croix, USVI	Average 2001-2008	14	13	2638	0.005	0.00005

\* Lack of raw dataset prevented computation of surveys in which Nassau grouper were sighted, and hence, sighting frequency as well  
 \*\*This data not included in computation of density and sighting frequency trends for CREMP visual surveys in Puerto Rico

"  
 "  
 "  
 "  
 "  
 "  
 "  
 "  
 "  
 "  
 "  
 "  
 "

### 3.b. COUNTRY ACCOUNTS

" Hgy 'r qr wrcvqp'cpcn{ ugu'qt 'uqemlcuuguo gpw'j cxg'dggp'eqpf wevgf 'qp'v'j g'P cuucw' i tqwr gt0'Vj gtghqtg'y g'uwo o ctk g'dgmjy 'huj gt { 'tgpf u'ecvej gu'qt'ecvej 'r gt'wplk'qh'ghqtv+'qxtg' vko g.'huj gt { /kpf gr gpf gpv'wvf gty cvgt'tggh'huj 'uwxg{ u.'ukl gu'rcpf gf . 'cpf 'c'pcttqy 'tapi g'qh' dlqmj kecn'uwf lgu'v'q'kphqto 'ewttgpvr qr wrcvqp'ucwu0'I gpgvle'y qtnlv'f cvg'wui i guu'c'ukpi ng' r cpo leve'r qr wrcvqp'eqppgevgf 'v'j tqwi j qw'ku'tcpi g0'Uwf lgu'qh'ektewrcvqp'r cvgtpu'cv'ur cy plpi " ci i tgi cvkqp'ukgu'i gpgtcm{ 'j cxg'kpf kecvgf 'v'j g'r tgupeg'qh'gf f lgu'cpf 'mqecn'gvpv'kqp'o gej cpkuo u' vj cv'tguwn'lp'ugrh'tgetwko gpv'lp'o quv'ctgcu'cmj qwi j 'c'o kz 'qh'mqecn'cpf 'mqpi /f kucpeg'gi i 'cpf " mctxcn'tepur qt'vr r gctu'o quv'hkngn{ 'cpf 'uqo gy j cv'wpr tgf kecvdrg0'

J qfi i uqp'cpf 'Nkgdgrt '\*4224+'pqvgf 'v'j cv'P cuucw'i tqwr gt'y gtg'cdugpv'htqo ": 4' "qh' uj cmjy 'Ectkddgcp'tgghu'f wtkpi 'c'7/{ gct'r gtkqf 'qh'wvf gty cvgt'uwxg{ u'ht'v'j g'TgghEj genlr tqlgex0' Qh'384'tgghu'uwxg{ gf 'hqt'P cuucw'i tqwr gt.'qpn{ 'gki j v'tgghu'j cf 'o qtg'v'j cp'qpg'huj 0'Qh'v'j g'328" vqcn'huj 'eqwpgf 'f wtkpi 'hxg' { gctu'qh'o qpkqtkpi . '98'y gtg'hqwpf 'qp'v'j q'tgghu'lp'v'j g'Y qtrf " J gtkci g'Ukg'lp'Ucp' Cpft'2 u' Ctej kr gmi q'lp'Eqmqo dlc. 'y j gtg'ur gct'huj kpi 'ku'r tqj kdkgf 'qp'dqj " tgghu0'k'v'j g'Cvrcvle'tgi kqp.'i tqwr gt'cdwvf cpeg'\*kpenf kpi 'P cuucw'i tqwr gt+'f genkpgf 'htqo "3; ; ; " \*305'i tqwr gt'0504'r gt'322'o 4+'v'4222 '\*2047'02076'r gt'322'o 4+'cmj qwi j 'v'j ku'ku'pqv' uvc'v'kcm{ 'uki p'k'lecp0'Vj ku'tgpf 'ku'eqpukv'gpv'y kj 'wvf gty cvgt'uwxg{ u'lp'o quv'eqwv'k'cetquu' vj g'tcpi g'qh'v'j g'ur geku0'

O cp { 'qh'v'j g'eqwv'k'g'u'y j gtg'P cuucw'i tqwr gt'j cxg'dggp'tgr qtvgf 'j cxg'o gej cpkuo u'v'q' tgr qt'v'huj gt { 'rcpf kpi u.'gkj gt'cu'c'o gcpu'qh'wvf gtucpf kpi 'mqecn'o cpci go gpv'ucwu'cpf 'pggf u'qt' cu'c'r ctv'ekr cpv'lp'dtqcf gt'tgi kpcn'qt'lpv'gt'pcv'kpcn'o cpci go gpv'qt'eqpugtxcvkqp'ghqtv'u'\*kq0' Hqaf 'cpf 'Ci tlewwtg'Qti cpk cvkqp'qh'v'j g'Wpkgf 'P cvkpu0'Wphqt wpcvgn{ . 'o cp { 'f q'pqv'eqmgev' f cv'cv'v'j g'ur geku'ngxgn'dw'tcv'j gt'eqmgev'f cvc.'rcpf kpi u'qt'ghqtv'ht'k'pucpeg.'qpn{ 'cv'uqo g' hwpev'kpcn'i tqwr 'qt'hc'o kn{ 'ngxgn0'Y j kng'v'j ku'o c { 'dg'o kf n{ 'kphqto cvxg.'k'ku'tctgn{ 'wughw'lp' wvf gtucpf kpi 'huj gt { 'ko r cevu'v'q'kpf kxf wcn'ur geku0'O quv'qh'y j cv'ku'npqy p'qh'v'j g'ewttgpv'ucwu' qh'P cuucw'i tqwr gt'uqemu'o wu'dg'f gtxgf 'htqo 'tgugctej 'qt'o qpkqtkpi 'ghqtv'qt'cu'k'v'gtr'tgcvkqp' qh'v'j g'ucteg'f cvc0''

K'cf f k'kqp'v'q'v'j g'eqwv' { 'ceeqwpu'v'j cv'hqmqy . 'KWE'P 'hku'v'j g'hqmqy kpi 'cu' kucpf uleqwpv'k'g'u'y j gtg'P cuucw'i tqwr gt'ku'eqpukf gtgf 'v'q'dg'pcv'xg'\*J ggo uvc'cpf 'Tcpf cm'3; ; 5+0" Vq'f cvg.'hkwg'geqmqi kecn'qt'huj gt'k'g'u'f cvc.'kphqto cvkqp'qt'cpgef qcn'g'xkf gpeg'ku'cxckrdng'v'q' r tqxkf g'k'puki j v'k'p'v'j g'ucwu'qh'P cuucw'i tqwr gt'lp'v'j g'hqmqy kpi 'lwt'kuf ke'v'k'p'u'<C'p'ki wc'cpf " Dctdwf c=Ctwdc=Eqvc=Tlec=Ewte+cq=F qo k'p'lec=H'g'pej 'I w'k'pc=I t'g'pcf c=I wcf grqwr g= I wcvgo cr=I w'c'pc=J c'k'k=O qp'v'gt'tcv=P gj gtr'rcpf u'C'p'v'k'ngu'\*E wte+cq=P lectci wc=Rcpco c= Uclpv'M'ku'cpf 'P gxku=Uclpv'Nwlc=Uclpv'X'k'pegv'cpf 'v'j g'I t'g'pcf k'p'gu=Uwt'k'pco g=Vt'k'p'kf cf 'cpf " Vqdc'i q=W'p'k'gf 'U'cv'gu'O k'p'qt'Q'wn'f kpi 'K'uc'pf u'\*k'q0'P c'x'cuuc=cpf 'X'g'pgl w'gr0'U'qo g'qh'v'j g'ug' mqecv'k'pu'ctg'eqo d'k'p'gf 'k'p'v'j g'uge'v'k'p'g'p'v'k'ngf '0'N'gu'gt'C'p'v'k'ngu.'E'g'p'tcn'cpf 'U'q'w'j 'C'o g't'lec0''

"  
"  
"  
"

**ANGUILLA**

"

Vj g'hqmjy kpi "kphqto cvkqp'y cu'qdvckp'gf 'xk'Lco gu'E0I wo du.'F ktgevqt"qh'Hkuj g'kgu'cpf 'O ctkpg'  
Tguqwtegu.'O kpkut { 'qh'J qo g'Chcktu.'I qxgtpo gpv'qh'Cpi wkm0'r gtu0'eqo o 0'vq'T0J km'P O HU'  
UGHUE.'42350"

"

**Anguilla – Populations**

"

Nkwg'kphqto cvkqp'ku'cxckrdng'htqo 'r wdrkuj gf 'uqwtegu'qp'yj g'ucwuw'qh'P cuucw'i tqwr gt'kp"  
Cpi wkm0'y cvgtu0'Ceeqtf kpi 'vq'yj g'Hkuj g'kgu'F gr ctwo gpv'kp"4234<0Y kj 'tgi ctf u'vq'yj g'P cuucw'  
i tqwr gt'k'ku'pqv'xgt { "cdw'pf cpv'kp'Cpi wkm0'Qh'kgu'cv'yj g'F gr ctwo gpv'j cxg'tgr qt'v'gf "qpn { "  
uggkpi "qpg'qt'vy q'lwxgpkgu'qp'yj gk'f kxgu'cpf "qvj gt'kp/y cvgt'y qtn0"Y g'f q'pqvj cxg'yj g'  
j k'w'q'k'c'n'f'c'v'k'p'Cpi wkm'vq'f'gv'to k'p'g'yj gk'htqo gt'cdw'pf c'peg.'j qy gxgt'k'ku'dgn'x'g'f'yj cv'yj g' { "  
y gtg'o qtg'cdw'pf cpv'yj cp'yj g' { "ctg'pqy . 'lwf i kpi 'htqo 'r cuv'hkuj "ecvej "qdugt'xc'v'k'pu0"

"

**Anguilla – Fisheries**

"

P q'f'c'v'ctg'cxckrdng'htqo 'r wdrkuj gf 'uqwtegu'qp'yj g'hkuj g'kgu'yj cv'v'cng'qt'j cxg'v'cng'p'P cuucw'  
i tqwr gt'0'Ceeqtf kpi 'vq'yj g'Hkuj g'kgu'F gr ctwo gpv'kp"4234<0Vj g'P cuucw'i tqwr gt'ku'c'ur gekgu'yj cv'  
y cu'qdugt'x'g'f'kp'hkuj "ecvej gu'kp'yj g': 2u'cpf 'r tkqt'vq'yj cv'pqv'cp { 'i t'ge'v'co q'w'p'u+'d'w'p'qy 'yj g' { "ctg"  
pqv'c'r ctv'qh'yj g'ewtt'gpv'hkuj "ecvej gu'hkuj 'v'cr u'cpf 'h'p'gu+0'C'hkuj "ecvej 'f'c'v'e'q'm'g'v'k'p'r tqi t'co "  
lj cu'q'pn { 'd'ggp\_'ko r r'go gp'v'gf 'cv'yj g'f'gr ctwo gpv'kp'yj g'r'cu'v'h'q'w' { g'ct'u'cpf 'u'q'cp'c'p'c'n { u'ku'qh'  
j k'w'q'k'c'n'f'g'p'f' u'ku'p'q'v'r'qu'k'd'ng'0"J qy gxgt'í vj ku'ur gekgu'ku'p'q'v'r' t'g'ug'p'v'kp'ewtt'gpv'hkuj "ecvej gu0"

"

**Anguilla – Conservation and Management**

"

0Vj gtg'ctg'pq'hpqy p'ur cy p'kpi "ci i tgi cvkqp'uk'gu'cpf "yj gtg'ctg'pq'ur gekn'eq'pugt'xc'v'k'p'qt"  
o cpci go gpv'tgi w'v'k'p'u'kp'r' r'eg0"

"

"

"

"


"

**BAHAMAS**

**Bahamas – Abundance and Distribution**

"

" Vj g'Dcj co cu'y kj 'ku'o cp { 'kurpf u'cpf "gz vgpukxg'uj cmqy 'tgg'h'ctgcu'xgt { 'r quakn { 'j qrf u' qt'j grf "qpg'qh'vj g'rcti guv'r qr wrcvqpu'qh'P cuucw'i tqwr gt 'vj tqwi j qw'ku'tcpi g0"Vj g'ur geku'j cu' mpi "dggp'vj g'o clqt'rcpf gf 'hphkuj 'hqt'vj g'eqwpx { 'cpf 'vj g'htuv'cpf 'rcti guv'gxgt'tgr qtvgf " ur cy plpi "ci i tgi cvkqp\*y kj "cp'guko cvgf "52.222"vq'322.222'hkuj +y cu'f qewo gpvgf 'Itqo 'vj g"



**Figure 11. Approximate locations of Nassau grouper spawning aggregation sites in the Bahamas.**

Dcj co cu'lp'Dko lpk\*Uo kj " 3; 94-0"D { 'vj g'rcvg" 3; ; 2ulgctn { "4222u.'vj g" P cuucw'i tqwr gt" r qr wrcvqpu'u+lp'vj g" Dcj co cu'y cu'hkngn { 'hwm { " gZR mkgf "v'qxgt/gZR mkgf " \*Gj tj ctf v'cpf "F grxgcwz" 4229.'Ej gwpi "gv'cr04235-0" Dqyj 'hkuj g'ltgu'rcpf lpi u'cpf" o gcp'dqf { 'uk'g'lp'ecvej gu" j cxg'f genkpgf 'ukpeg'vj g" 3; ; 2u.'f gur kg'c'o loko wo " uk'g'tgi wrcvqpu.'r tqvgevkqp" f wtkpi 'vj g'ur cy plpi " ci i tgi cvkqp'ugcuqp." guvcdhkuj o gpv'qh'ugxgtcn' r tqvgevgf "ci i tgi cvkqp'ukgu."

cpf "o ctkpg'r tqvgevgf "ctgcu\*Ej gwpi "gv'cr04235-0"Vj gtg'ku'pq'lpf kcvkqp'vj cv'vj gug'f genkpgu'ctg" f vg'v'q'tgf wegf 'hkuj lpi "ghhqtv'qt'v'ej cpi gu'lp'hkuj lpi 'r tcevegu=qxgthkuj lpi 'ku'o quv'rkngn { 'vj g" ecwug0'T gf wrcvqpu'lp'pwo dgtu'qh'hkuj "qdugtxgf 'lp'tgg'h'uwtxg { u.'cnuq'umi i guv'vj cv'r qr wrcvqpu'ctg" f genkplpi 0"Qpg'o clqt'eqpegt'ku'y kj 'r qcej lpi . 'gur gekm { 'd { 'pqp/Dcj co kcpu'Dcj co kcp" hkuj gto gp'rcti gn { 'cdkf g'd { 'vj g'ugcuqpcn'eqwuxgu'hqt'P cuucw'i tqwr gt \*O 0Dtc { pgp.'Dcj co cu' F gr ctvo gpv'qh'O ctkpg'T guqwtegu.'r gtu0eqo o 0vq [ 0Ucf qx { . 'Wpkxgtuk { 'qh'J qpi 'Mqpi .'4234-0"

O qtg'vj cp'42"ci i tgi cvkqpu'j cxg'dggp'tgr qtvgf 'Itqo 'vj g'Dcj co cu.'dw'xgt { 'hgy 'j cxg" dggp'uwf kgf 'lp'cp { 'f gvckl'cpf 'vj g'ewtgpv'ucwu'qh'vj g'i tgcv'o clqtkv { 'ku'wmpqy p0'Ewo wrcvkg" f cv'htqo "TGGH\*4225/4235+uj qy 'j ki j 'pwo dgtu'qh'uki j vki u'qh'624'P cuucw'i tqwr gt 'lp'3693" uwtxg { u'f gpuk { 'lpf gz'307.'uki j vki 'Itgs wgpe { '490' +lp'vj g'pqt'vj 'Dcj co cu.'594; 'P cuucw' i tqwr gt 'lp'8749'uwtxg { u'f gpuk { 'lpf gz'308.'uki j vki 'Itgs wgpe { '790' +lp'vj g'egp'tcn'Dcj co cu." cpf "6; 'P cuucw'i tqwr gt 'lp'97'uwtxg { u'f gpuk { 'lpf gz'308.'uki j vki 'Itgs wgpe { '870' +lp'vj g'uqwj " Dcj co cu'cetqu'vj g'32/ { gct'r g'kqf 0"Gzco kpcvqpu'qh'vko g'r g'kqf u'qh'3; ; 2/; 7'xu0422: /35'f q'pqv"

"  
"  
"  
"

uj qy 'i t gcv'f khhgtgpegu'lp'uki j v'kpi 'Itgs wgpe { 'qt 'f g'pukv { 'lpf gz. 'dw'ur cvkcnlo cpci go gpv' | qpg'f cvc" ctg'rcenkpi 'uq'eqo r ctdckkv { 'qh'uksgu'ku'pqv'npqy p"

\*j wr <ly y y 0gghQti If d l tgr qt vulf kuvur gekguIVY C l22; 94225/23/234235/26/29-0"C v'c'p'ke'cpf " I wh'T cr kf 'Tggh'Cuuguuo gpv'\*CI TTC+'uwxg { u'lp' Cpf tq'K'rcpf '\*3; ; : +'cpf 'C'dceq'K'rcpf u" \*3; ; ; +'hqwpf 'tgrcvkxgn { 'rjy 'pwo dgtu'qh'gpeqwpvgtu'\*uki j v'kpi 'Itgs wgpeku'qh'90 ' 'cpf '508' . " tgr ge'v'x'gn { =f g'puk'ku'qh'35'huj lj gevctg'cpf '7'huj lj gevctg. 'tgr ge'v'x'gn { +'\*CI TTC'f cvc. 'Itqo 'V0' M'gnkuqp. 'P O HU'UGHUE 'o'Dgcwhqtv'Ncdqtcvqt { . 'P E+0"Vj gug'ctg'dgy ggp'590' 'cpf '3606' 'qh' vj g'f g'puk'ku'ekvgf 'd { 'Dctfcej 'lp'c'tgrcvkxgn { 'iki j v' { 'ko r cev'g' 'Dgto wf c'lp'vj g'3; 72u0""

Gz v'g'puk'x'g'cpf 'tgr gev'g'f 'uwxg { u'qh'ur cy pl'pi 'ci i tgi cvk'pu'o c { 'r tqxkf g'uqo g'g'xkf g'peg'qh" vtgpf u'lp'cdwpf c'peg'kh'ghqtv'ku'eqpukv'gpv'cpf 'v'ko l'pi 'qh'uwxg { u'tgrcvkx'g'v'q'ur cy pl'pi 'cev'x'kv { 'ecp" dg'cuuwtg'f'0"Cm'j qwi j 'u { v'go cvk'uwxg { u'j cxg'pqv'dggp'eqo o qp. 'uqo g'uwf k'gu'ecp'j ki j rki j v' o clqt'ej cpi gu'0"Cv'Ec'v'Ec { . 'Uo kj '\*3; 94+.'lp'vj g'htuv'uek'p'v'k'he'tgr qt v'qh'c'P cuucw'i tqwr gt" ur cy pl'pi 'ci i tgi cvk'qp. 'f qewo gpv'g'f 'v'g'pu'qh'vj qwucpf u'\*52.222/322.222+'qh'ur cy pl'pi 'P cuucw' i tqwr gtu'0"C' uwxg { 'lp'Lcpwct { '4235'f v'k'pi 'vj g'hw'no qqp'r g'k'qf 't'g'x'k'v'g'f 'vj g'ci i tgi cvk'qp'uks'g" tgr qt v'g'f 'd { 'Uo kj 'lp'3; 940"Vj g'uks'g'cpf 'gz v'g'puk'x'g'uwtqwpf l'pi 'ctgcu'\*60' h'p'gct' o k'gu'+cm'pi 'vj g" tgg'h'y g'tg'uwxg { g'f 'o v'w'k' r'g'v'ko gu'0"P q'g'x'k'f g'peg'qh'ur cy pl'pi 'h'uj 'qt'c'x'k'cd'g'ur cy pl'pi " ci i tgi cvk'qp'y cu'h'qwpf 0'Y j gp's v'g't'k'g'f . 'h'ec'h'uj gtu'uck'f 'vj g'ci i tgi cvk'qp'j cf 'f'k'uc'r g'ct'g'f 'd { 'vj g" g'ctn { '3; : 2u'\*j wr <ly y y 0uehcQti l'ko ci guluvt'k'g'ul'f h'lp'gy u'ng'w'g' l'p'gy u39ah'k'p'c'r'f h'"

**Bahamas - Fisheries"**

P cuucw'i tqwr gt'ctg'v'cti g'v'g'f 'd { 'ct'v'uc'p'cn'l'w'd'uk'v'g'peg. 't'get'g'cv'k'p'c'n'c'p'f 'eqo o g'tek'n' \*k'p'c'n'f l'pi 'h'qt'g'z'r q'tv'h'uj g't'k'gu'0'D'we'j c'p'\*4222+'l'p'f k'ev'g'f 'vj cv'vj g'uj c'm'qy 'd'c'p'm'u'v'j t'q'w' j q'w" I t'g'cv'c'p'f 'N'k'w'g'D'c'j co cu'D'ep'm'u. 'vj g'Ec { 'U'c'n'D'ep'm'c'p'f 'vj g'E't'q'q'ng'f 'K'rc'p'f 'c'p'f 'C'c'm'k'p'u'K'rc'p'f " D'ep'm'u'y g't'g'v'j g'o clqt'h'uj l'pi 'i t'q'wp'f u'h'qt'P cuucw'i tqwr gt'0'K'p'v'j g'D'c'j co cu. 'h'uj g'to gp'w'ug" j c'p'f h'p'g'u. 't'c'r u. 'c'p'f 'ur g'c'tu'\*k'p'c'n'f l'pi 'eqo r t'g'u'q't'ule'qo r t'g'u'g'f 'c'k't+'v'q'v'c'ng'P cuucw'i tqwr gt" \*U'c'f q'x { '3; ; 9+0"Vj g'v'ug'q'h'c'ur g'c'ti v'p'k'u'k'ng'i c'n'd'w'c'ur g'c't'y k'j 'c'ur'k'pi '\*g'f 0'J cy c'k'c'p'ur'k'pi +'k'u" ng'i c'r'0'U'r g'c't'h'uj l'pi 'c'p'f 'h'uj /v'c'r r l'pi . 'lp'r c't'v'k'w'r't. 't'g'u'w'v'lp'uki p'k'h'c'ep'v'f 'j ki j g't'ERWG'v'j c'p" q'v'j g't'h'uj l'pi 'o g'v'j q'f u'\*E'j g'w'p'i 'g'v'c'r'0'4235+0'T'g'i v'w'v'k'p'u'd'gi c'p'v'q'v'k'o k'u'q'o g'ci i tgi cvk'qp'h'uj l'pi " l'p'3; ; : 'c'p'f 'y g't'g'k'o r ng'o gpv'g'f . 'cu'c'5/o q'p'v'j 'e'm'q'u'w't'g. 'p'c'v'k'p'c'm'f 'lp'42270'H'uj l'pi 'h'qt'P cuucw' i tqwr gt'lp'q'v'j g't'o q'p'v'j u'eq'p'v'k'p'w'g'u'0""

K'p'v'g'to u'qh'y g'ki j v'c'p'f 'x'c'n'w'g. 'P cuucw'i tqwr gt'j cu'd'ggp'v'j g'h'q'w'v'j 'o q'u'v'k'o r q't'v'c'p'v' eqo o g'tek'n'h'uj g't { 't'g'u'q'w't'eg'lp'v'j g'D'c'j co cu'G'z'e'n'w'k'x'g'G'eq'p'q'o k'e\ q'p'g'd'g'j l'p'f 'ur l'p' { 'm'd'v'g't. " u'p'c'r r g'tu'c'p'f 's'w'g'g'p'eq'p'ej '\*D'we'j c'p'4222+0'K'p'4229. 'vj g'o q'u'v't'g'eg'p'v'v'w'o o c't { 'c'x'c'k'c'd'g'g. 'P cuucw' i tqwr gt'eqo r t'k'ug'f '4' 'd { 'd'q'v'j 'y g'ki j v'c'p'f 'x'c'n'w'g'q'h'v'j g't'g'eq't'f g'f 'eqo o g'tek'n'h'c'p'f l'pi u'qh'c'm' eqo o g'tek'm'f { 'g'z'r m'k'g'f 'ur g'ek'gu'lp'v'j g'D'c'j co cu'\*H'c'2'422; +=ur l'p' { 'm'd'v'g't'o c'ng'u'v'r 'vj g'i t'g'c'v'g'u'v' o clqt'k'f { 'qh'v'j g'eq'w'p't { 'u'eqo o g'tek'n'h'c'p'f l'pi u'0'P cuucw'i tqwr gt'lp'4229'c'ee'q'w'p'v'g'f 'h'qt'95' 'qh'c'm' eqo o g'tek'n'i tqwr gt'rc'p'f l'pi u'lp'v'j g'eq'w'p't { =t'get'g'cv'k'p'c'n'c'p'f 'u'w'd'uk'v'g'peg'v'ug'f'c'v'c'q'h'v'j g'ur g'ek'gu' ctg'p'q'v'c'x'c'k'c'd'g'g'\*H'c'Q'422; +0"C'ug'c'h'q'q'f 'eq'p'uw'o r v'k'q'p' u'w't'x'g'f 'lp'4225/4226'd { 'V'c'r'w'g/

"  
"  
"  
"

O eO cpwu'cpf "J c| gm'gunko cvgf "vj cv'vj g'hkuj g'kgo' qpkqtkpi "u{vgo "kp'vj g'Dcj co cu'f kf "pqv" f qewo gpv'; 6' "qh'vqcn'i tqwr gt'ecvej "dcugf "qp'equpuo r vkqp'cpf "tcf g'ucvknku" \*NOO eO cpwu." Vj g'I mdcn'Gpxkqpo gpv'Hckkx\ "Vtcpuqwpf ct { "Y cvgtu'Cuuguuo gpv'Rtqi tco o g.'wpr wd0f cvc." r gtu0eqo o 0'vq'T0J km"P O HU."4236+0"Vj gtghqg.'tgr qtvgf "ncpf kpi u'ctg'r tqlgvegf "vq'tgr tgugpv' qpn\ '8' "qh'vj g'vqcn'r tqf wevkqp'pggf gf "vq'o ggv'gZR qtv'cpf "equpuo r vkqp'rgxgn0"

O vej "qh'vj g'cppwcn'ncpf kpi u'j kvqtkcm\ "eco g'ltqo "ur cy plpi "ci i tgi cvkqpu" \*Eqrkp"3; ; 4=" cu'o cp { "cu'53'f khtgpv'ukgu" \*Vcdng"33+ j cxg'dggp'tgr qtvgf " \*Ucf qx { "f g'O ke j guqp"4234+0"Y j kng" DTGGH"3; ; : + "tgr qtvgf "dgvy ggp"35"cpf "53"ci i tgi cvkqpu."45"j cxg'dggp"eqphko gf "d { "f kgev" qdugtxcvkqp'qt'ecvej "o qpkqtkpi " \*Ucf qx { "cpf "Gmwpf"3; ; ; +0"ncpf kpi u'f cvc'ltqo "3; ; 7/4228" uj qy gf "vj cv'o quv'P cuucw'i tqwr gt'y gtg'ncpf gf "dgvy ggp" F gego dgt "vq'Hgdtwct { ".cmj qwi j "ewtgpv' tgi wrcvkqpu'tgustkv'hkuj kpi "hqt" P cuucw'i tqwr gt'f wtkpi "o quv'qh'vj cv'r g'kqf 0"

Gctn\ 'tgugctej "d { "Uo kj "3; 94+ "cpf "Eqrkp"3; ; 4+ "kf gpv'kkgf "ur cy plpi "ci i tgi cvkqpu'ukgu'kp" vj g'3; 92/3; : 2u'y kj "pwo dgtu'qh'ur cy pgtu'tcpi kpi "ltqo "j wpf tgf u'vq'v'g'pu'qh'vj qwucpf u0" Uwdugs wgpv'tgugctej "j cu'tctgn\ "hqw'f "cdwpf cpegu'pgctn\ "uq"j ki j 0"kp'cp'cwgo r v'vq'tguxg { "vj g" ukgu'f qewo gpv'gf "d { "Eqrkp"3; ; 4+ "tgugctej gtu'ltqo "P qtj 'Ectqrkpc"Ucv'g'Wpkxgtukv\ "kp'lcwct { " 4224"eqpf wevgf "f kxgt"cpf "j { "ftqceqwukc"uwxg { u'ctqwpf "Nqpi "Kucpf."Dcj co cu' \*I cueqki pg" \*4224."F 0Gi i gnv'q. "P qtj 'Ectqrkpc"Ucv'g'Wpkxgtukv\ ." r gtu0eqo o 0'vq'T0J km"P O HU"UGHUE." 4236+0" P qpg'qh'vj g'ukgu'xkukgf "j cf "o qtg'vj cp"4: "hkuj "cpf "pqpg'qh'vj g'hkuj "qdugt'xgf "gzj kdkgf " ur cy plpi "dg'cxkqt0"K'ku'r quukdng'vj cv'ur cy plpi "j cf "qee'wtgf "vj g'r t'gxlqwu'o qpj u'cmj qwi j " f kweuukqpu'y kj "ctgc'hkuj gtu'cpf "hkuj "o ctngv'gtu'ngf "vj g'tgugctej gtu'vq'dng'kxg'vj cv'ur cy plpi " ci i tgi cvkqpu'pq'mpi gt'qee'wtgf "cv'vj gug'ukgu" \*F 0Gi i gnv'q. "P qtj 'Ectqrkpc"Ucv'g'Wpkxgtukv\ ." wpr wd0f cvc.' r gtu0eqo o 0'vq'T0J km"P O HU"UGHUE."4236+0"

Vj g'ci i tgi cvkqpu'ukg'cv'J ki j "Ec { "y cu'cmj'tgr qtvgf "vq'eqpukv'qh'cp'qtf gt'qh'o ci plkwf g" hgy gt'ur cy pgtu'vj cp'ku'j kvqtkcn'uk\ g0'F kxgt'gunko cvgu'tcpi gf "ltqo "322"/"3.222"hkuj "hqt"3; ; ; /" 4222" \*Gj tj ctf v'cpf "F grxgcwz"4229."I cueqki pg"4224."Tc { "gv'cn04222+0"kp"3; ; ; ."4222."cpf " 4223."j { "ftqceqwukc"uwxg { u'y gtg'w'pf gt'wngp'cv'J ki j "Ec { "cu'c'pqxgn'cuuguuo gpv'qh'vj g'pwo dgt" qh'hkuj "kp'ukpi ng'ur cy plpi "ci i tgi cvkqpu" \*Gj tj ctf v'cpf "F grxgcwz"4229+0"Vj ku'uwf { "tgr qtvgf " gunko cvgu'qh'32.745"3; ; ; +; ".522" \*4222+ "cpf "34.: 79" \*4223+ "hkuj "dcugf "qp'ceqwukc"uki pcn' utgpi vj "dw'vj gtg'ku'rkwng'f gvckn'qp'vj g'uco r r'kpi "o gj qf "qt'vj g'lp/y cvgt" qdugt'xcvkqpu'pgegu'ct { "vq" xcnf cvg'vj g'f cvc0"kp"4222"/"4223."f kxgtu'tgwtpgf "vq'vj g'ncv'kqp'cpf "f kf "pqv'ncv'g'cp" ci i tgi cvkqpu."eqpenw' kpi "hkuj "o c { "pqv'j cxg'ci i tgi cvgf "cv'vj g'ukg" \*I 0Ectrgv'p" Tc { ".Wpkxgtukv\ "qh" Xkti kpkc." r gtu0eqo o 0'cu'tgr qtvgf "kp'I cueqki pg"4224+0"Vj gtg'y cu'pq'uw'xg { "f cvc'cxckrdng'hqt" vj g'J ki j "Ec { "ukg'kp"4224."dw'ecvej "y cu'mjy "hqmjy kpi "c'vj tgg'o" { gct'o qtcv'qtkwo " \*L0Dktej ." Uo cm'J qr g'Dc { ".Cpf tq. " r gtu0eqo o 0'vq'L0I cueqki pg."Wpkxgtukv\ "qh'Xkti kpkc."4224+0"Vj g' pwo dgt'qh'ur cy plpi "P cuucw'i tqwr gt'cv'vj g'J ki j "Ec { "ci i tgi cvkqpu'y cu'g'xkf gpv' "f getg'culpi " tgr'v'xg'vq"j kvqtkc" gunko cvgu" \*Tc { "gv'cn 4222+0"

"  
"  
"  
"

**Table 4. Aggregation sites in the Bahamas in 1998**  
*Bahamas aggregations - modified from BREEF (1998): DoF = Department of Fisheries*

Location "	Number "	Discussion/Source "
Ec{vEc{ "	3"	322.222'lpf kxf wcn'qdugt xgf "3; 920 <sup>33</sup> "4235"uwxg{ "cv'Lcpwct { 'hwn'o qqp'cpf " nqecn'lpvgt xly u'lpf kcv'g'v'v'j g'ci i tgi cvkqp'j cu'npqi 'egcugf 'v'q'hto "UETHC" P gy urgwgt '39-0'
I tgecv'kuce'Nki j v'"	3"	O c{ 'j cxg'f kcr r gctgf 'f wg'v'q' huj kpi 'r tguwtg'htqo 'Hqtkf c' "4"
Cpf tqu"	4"	Nqecv'kpu'J ki j 'Ec{ 'cpf 'V'kpngt 'Tqem' "5"
Cpf tqu"	5'o qtg'"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'npqy p0 <sup>6</sup> "P qv'cr r ctgpnw{ 'npqy p'd{ 'nqecn' huj gto gp'uj'r tqdcdn{ 'pqv'twg0"
Nqpi "Krcpf'"	3"	F ger'kpg'kp'ecvej gu'htqo 'ugxg'tcn'v'j qwucpf 'v'q'nguu'v'j cp'322'huj 0 <sup>7</sup> "J qy gxgt. " 3; ; 9'ecvej gu'qp'v'j g'ci i tgi cvkqp'tgr qt'v'gf 'v'q'dg'i qqf 'd{ 'Nqpi "Krcpf 'huj gto gp0" Qpg'ci i tgi cvkqp'usk'lp'Nqpi "Krcpf 'tgr qt'v'gf 'd{ 'huj gto gp'lp'Ucp'Ucixcf qt'cu' v'j gkt'pgct gu'v'ci i tgi cvkqp'"
Long Island "	2 more "	Nqecv'kpu'cpf 'uqwtg pqv'i kxgp0 <sup>8</sup> "F qH'dgnkxg'v'j cv'v'j g'g'ctg'wy q'ci i tgi cvkqp' usku'lp'Nqpi "Krcpf 0"
Gzwo c'Ec{ u'"	3"	Htqo 'y qtnid{ 'Rcv'Eqk' "9"
Ec{v'Krcpf'"	3"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'i kxgp <sup>10</sup> "
Dgtt{ "Krcpf u'"	6"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'i kxgp'"
P gy 'Rtqxf gpeg'"	3"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'i kxgp0P qv'tgr qt'v'gf 'd{ 'P gy 'Rtqxf gpeg' huj gto gp'uj'wp'kngn{ 'v'q'dg'twg0"
Tci i gf "Krcpf'"	3"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'i kxgp0K'gzkuw'o c{ 'dg'v'j tgc'v'p'gf 'd{ 'huj kpi " r tguwtg'htqo 'q'v'j g't'eqw'p't'k'u'"
Ec{ 'Ucn'"	3"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'i kxgp0K'gzkuw'o c{ 'dg'v'j tgc'v'p'gf 'd{ 'huj kpi " r tguwtg'htqo 'q'v'j g't'eqw'p't'k'u'"
Grgwj gtc'"	6"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'i kxgp'"
Cem'kpu'"	3"	Nqecv'kpu'cpf 'uqwtg'qh'tgr qt'v'pqv'i kxgp0K'gzkuw.'o c{ 'dg'v'j tgc'v'p'gf 'd{ " huj kpi 'r tguwtg'htqo 'q'v'j g't'eqw'p't'k'u0"
Cdceq'"	5"	F k'ewu'k'p'u'y kj "J qr g'v'qy p'cpf 'O ctuj "J ctdqwt'huj gto gp'"
I t'cpf 'Dej co cu'"	6"	Tgr qt'v'gf 'v'q'dg'npqy p'cpf 'huj gf 'd{ 'c'I t'cpf 'Dej co c'dcugf 'huj kpi 'eqo r cp{ " * "4"
Minimum total "	6"	Ci i tgi cvkqp'eqph'kto gf 'lp't'gegpv'uekpv'k'le'k'k'g'c'w'g'"
Approximation "	35"	Ci i tgi cvkqp'eqph'kto gf 'd{ 'nqecn'tgr qt'u'cpf 'k'g'c'w'g'"
Maximum total "	53"	Cm'tgr qt'u'cdq'xg.'u'qo g'q'h'y j kej 'ctg'k'c'k'n{ 'wp'k'ngn{ '"

"Sources: (1) Smith, 1972; (2) Reported by CL Smith in the early 1970s; (3) From discussions with fishermen- Dr. Tim Turnbull (4) 5 spawning aggregations in Andros reported in Sadovy (1997) (5) Colin 1992; (6) Sadovy (1997); (7) Dr. Tim Turnbull, Sadovy (1997); (8) Sadovy (1997)-also source for Berry Islands, New Providence, Ragged Island, Cal Say, Eleuthera and Acklins; (9) Vallierre Deleveaux, Bahamas Dept. of Fisheries "

kp'c't'gegpv'f g'v'k'g'f 'c'p'c'n{ u'ku.'ecvej gu'htqo "3; ; 6'v'q'422; 'y g'g'c'cu'gu'ugf 'w'k'p'i 'huj g't{/ o qf g'k'p'i 'cr r tqcej gu'Ej g'w'p'i 'g'v'c'i'04235-0'Vj g'u'w'f { 'uj qy gf 'v'j cv'v'q'cn'rc'p'f kpi u'q'h'P cu'cw' i tqwr g't'lp'v'j g'Dej co cu'f ger'k'p'gf 'i tcf w'c'm'f 'htqo "3; ; 6'v'q'422; "H'k'i 035-0'E'qo r gp'uc'v'p'i 'h'q't" w'p'tgr qt'v'gf 'ecvej "eq'p'x'g't'v'gf 'htqo 'Ej g'w'p'i 'g'v'c'i'04235+'v'j g'Dej co cu'P cu'cw'i tqwr g't'ecvej gu' u'j q'w'f 'j cxg'dggp'g'uk'o cv'gf 'cv'ct'q'w'p'f "32.: 22'v'lp'3; ; 6'v'q'ct'q'w'p'f "4822'v'lp'422; .c'f get'g'c'ug'v'q" q'p'n' "46' 'q'h'v'j g'ecvej 'lp'3; ; 60'O qt'g'q'x'g't.'v'j g'r'q'r qt'v'k'p'q'h'P cu'cw'i tqwr g't'lp'v'j g'v'q'cn'huj g't{ " rc'p'f kpi u'c'm'ur g'el'gu+'lp'v'j g'Dej co cu'c'nu'q'f ger'k'p'gf 'htqo "32' 'v'q'6' 'f w'k'p'i 'v'j k'u'r g't'k'q'f. " u'w'i i gu'k'p'i 'v'j cv'v'j g'f ger'k'p'g'lp'rc'p'f kpi u'y cu'p'q'v'o kt'q't'g'f 'lp'q'v'j g't'g'z'r n'q'k'g'f 'c'z'c'y j kej 'y q'w'f " j cxg'lp'f k'c'v'gf "c'ej c'p'i g'lp'huj kpi 'g'h'q't'v'q't'o c'tng'v'eq'p'f k'k'q'p'u0'Vj k'u'w't'q'p'i n' 'u'w'i i gu'u'c' "

"  
 "  
 "  
 "  
 "



ecwi j v'f wtkpi 'ur cy plpi 'o ki tcvkpu'qt'y gtg'ugzwcm{ 'ko o cwtg0'Hkuj gtu'tgr qtvf 'y cv'  
-ei i tgi cvkpu'qh'o ki tcvkpi 'hkuj . 'y j kej 'y gtg'pqv'tkr g. 'y gtg's wkg'eqo o qp. 'dw'vj ku'dgij cxlqt 'ku'  
pqv'y kf gn' 'tgr qtvf 0'

Kphqto cvkqp'qp'tcf g'ku'rti gn' 'iko kkgf 'v'q'y kj kp/eqwpt { 'ucngu'gzeqr v'htq'i tqwr gt'ko r qtw'  
htqo 'y g'Dej co cu'lpv'vj g'Wpkgf 'Ucvgu0'O ctngv'uwtxg{ u'tgxgnf kthgtgpv'cur gew'qh'vj g'hkuj gt { "  
vj cp'vj qug'i ngco gf 'lwv'htqo 'rpf kpi u'f cv0'Ceeqtf kpi 'v'q'b qpj n' 'hkuj gto gp'lpvgtxky u'cpf "  
rpf kpi 'cdw'pf cpeg'uwtxg{ u'eqpf wevgf 'cv'O qpwi w'tco r '\*P cuucw+: 'c'ng{ 'o ctngv'qwwgv'htqo 'O c { "  
4229'v'Qevdgt'4229. 'y g'equv'htq'c'6/607'ni 'P cuucw'i tqwr gt'cxgtci gf 'WU&570220'Qh'c'v'v'cn'qh'  
76.222'hkuj 'rpf gf 'f wtkpi 'y g'8/o qpj 'uwtxg{ 'r g'kqf . 'P cuucw'i tqwr gtu'o cf g'w' 'cp'cxgtci g'qh'  
32' '\*d { 'pwo dgt+'o qpj n' '\*Q0cdqw'7.622'lpf kxkf wcn'hkuj +'y kj 'Lwpg'dgkpi 'y g'nyy guv'\*6' + 'cpf "  
Qevdgt'dgkpi 'y g'j k j guv'\*35' + '\*E wuj kqp'cpf 'Uwrkcp/Ugcrg { '4229+0'Vj ku'uwf { 'cnuq'pqvgf 'y cv'  
c'uk' gcdrg'r tqwr qv'kqp'qh'P cuucw'i tqwr gt 'y gtg'o ctngv'f 'd { 'ugngtu'y j q'r wtej cugf 'y go 'htqo "  
rti g'uecrg'eqo o gtekn'hkuj g'kgu'lp'P gy 'Rtqxkf gpeg0'Vj wu. 'y g'v'v'cn'cdw'pf cpeg'pqvgf 'lp'vj g'  
uwf { 'f k' 'pqv'uqng' 'tgr t'gugp'vj g'gh'qt'v'qh'O qpwi w'dcugf 'hkuj gto gp0'

" Vj gtg'ku'c'j kuvt { 'qh'g'zr q'v'cvkqp'qh'i tqwr gt'htqo 'y g'Dej co cu'v'q'vj g'Wpkgf 'Ucvgu0'  
Vj gtg'ku'ur gew'cvkqp'vj cv'eqpv'kwgf 'ko r q'v'cvkqp'qh'vj g'i g'p'g'k' i tqwr gt'ercu'k'hec'v'k'p'o c { 'k'p'v'f g'  
P cuucw'i tqwr gt. 'uk'peg'k'j cu'tcf k'k'q'p'cm { 'tgr t'gugp'vgf '92' 'qt'o q'tg'qh'vj g'Dej co k'p'i tqwr gt "  
rpf kpi u' '\*Ucf qx { 'f g'O kej guqp'4234+0Cv'vj ku'v'ko g'vj g'f cv'ctg'pqv'cxck'cdrg'v'q'eqph'ko 'y g'  
o ci pkw'f g'qh'vj gug'ko r q'v'v'qt'vj g'gh'ev'qh'eqpv'kwgf "g'zr q'v'v'qp'vj g'ucw'u'qh'P cuucw'i tqwr gtu'lp'  
vj g'Dej co cu0'E'q'ng'ev'k'p'cpf 'c'p'cn { uku'qh'vj gug'f cv'uj q'w'f 'dg'c'r t'k'q'k'v { 0

## The Bahamas – Conservation and Management

Kp'vj g'Dej co cu.'dqj 'ur cv'cn'cpf 'ugcu'q'p'cn'r tqv'ev'xg'o gcu'v'gu'ctg'lp'r m'eg'htq'vj g'  
o c'pci go gp'v'qh'vj g'P cuucw'i tqwr gt'lp'vj g'Dej co cu0'Kp'vj g'3; : 2u'c'o k'p'ko wo 'uk' g'qh'5'ndu0\*3058'  
ni +'y cu'lp'v'q'f wegf . 'ugcu'q'p'cn'r'eq'uw'v'gu'qh'ug'x'g't'cn'ur cy plpi 'ci i tgi cvkqp'uk'gu'y gtg'htu'v'  
ko r ngo gp'vgf 'lp'3; ; . 'cpf 'cp'c'p'p'w'cn'ow q/o qpj o '\*x'ct'k'cdrg'cee'q'f kpi 'v'q'h'wn'o q'p'+h'kuj gt { "  
en'uw'v'g'v'q'eq'k'p'k'f g'y kj 'y g'ur cy plpi 'r g'k'q'f 'y cu'htu'v'ko r ngo gp'vgf 'lp'F g'ego dgt'42250'Vj ku'  
en'uw'v'g'y cu'gz'v'g'f gf 'v'q'vj t'gg'o qpj u'lp'4227'v'g'p'eqo r cu'vj g'ur cy plpi 'r g'k'q'f 'htqo 'F g'ego dgt'  
vj tqw'j 'H'g'd't'w'c { 0'Vj g'en'uw'v'g'ku'c'r r r'k'gf "qp'c" { g'c't'n { 'd'c'uk'u'c'p'f 'k'o c { 'd'g'uj q't'v'g'p'gf "q't'q'y g'ty k'ug"  
k'p'h'w'g'p'eg'f 'd { 'u'w'ej 'h'c'v'q'tu'cu'vj g'g'eq'p'qo { '\*Ucf qx { 'c'p'f 'G'm'w'p'f "3; ; ; +0'H'q't'g'z'co r ng. 'h'q'm'y kpi "  
vj g'g'eq'p'qo k'e'f'q'y p'w'p'q'h'422: 'y g'en'uw'v'g'y cu'rh'ng'f 'v'q'ng'u'g'p'vj g'g'eq'p'qo k'e'd'w'f g'p'q'h'c'eq'ug'f "  
hkuj gt { 'v'q'h'kuj gto gp0'F wtkpi 'y g'ci i tgi cvkqp'r g'k'q'f . 'f wtkpi 'y j kej 'y gtg'ku'c'p'v'k'p'cn'd'cp'qp'  
P cuucw'i tqwr gt'ec'v'ej gu. 'rti g'p'wo dgtu'qh'hkuj 'y gtg'd'g'k'p'i 'v'c'ng'p'cee'q'f kpi 'v'q'h'kuj gt'cee'q'w'p'u'  
y kj 'r'j q'v'q'f'q'ewo gp'v'cv'k'p'cpf 'eq'p'h'ko kpi 't'gr q'v'v'q'h'r q'cej kpi 'qh'vj g'ur g'ek'gu'f wtkpi 'y g'  
ci i tgi cvkqp'ug'cu'q'p' '\*D'ej co cu'T'g'g'h'G'f w'ev'k'p'cn'r'H'q'w'p'f'cv'k'p'v' ]DTGGH\_ 'w'p'r w'd'0'f'cv'c+0'

Vj g'G'z'wo c'Ec { u'N'c'p'f 'c'p'f 'U'g'c'R'c't'm'h't'u'v'g'u'v'c'd'r'kuj gf 'lp'3; 7; . 'j cu'd'gg'p'eq'ug'f 'v'q'h'kuj kpi "  
uk'peg'3; : 8. 'y w'ur' t'q'v'ev'k'p'i 'dqj 'p'w't'ug't { 'c'p'f 'c'f w'n'j c'd'k'c'v'htq'P cuucw'i tqwr gt'c'p'f 'q'y g't'f'g'ng'v'f "  
o c't'k'p'g'ur g'ek'gu. 'u'w'ej 'cu's'w'gg'p'eq'ej . 'ur k'p { 'n'q'd'v'g't'c'p'f 'o c't'k'p'g'w't'v'g'u'0'G'x'k'f g'peg'htqo 'y g'G'z'wo c'

"  
"  
"  
"

Ec{u'Ncpf "cpf "Ugc'Rctnluj qy u'c'engct'f khtgpgpeg'kp'vj g'pwo dgt'cpf 'uk' g'qh'cm'rti g'i tqwr gt " ur gekgu'dgwy ggp'hkuj gf "cpf 'pqp/hkuj gf "ctgcu0"Vj g'dkqo cuu'qh'P cuucw'i tqwr gt'y cu'uj qy p'v'q'dg" ucw'kucem' 'i tgcvg't'kpukf g'cpf 'y kj kp'7"no 'qh'vj g'Rctn'dqwpf ctkgu'cpf 'tgr tqf wexg'qwr w'gi i " r tqf wexkp+y cu'ecw'w'v'f 'cu'ukz 'ko gu'j ki j gt'vj cp'qwukf g'vj g'r ctm'Uw'nc'gv'cr03; ; 9+0T gegpv' uwf kgu'd{ 'F cj ni tgp'gv'cr0\*wpr wd0f cwc+j' cxg'uggp'cf f kkp'cn'k'petgcugu'kp'dkqo cuu'htqo 'rguu'vj cp" 522'i B22'o 4'kp'4222/4226'v'pgct'3322'i B22'o 4'kp'vj g'4227/422; 'r g'ktqf "cpf 'o qtg'vj cp'3322" i lo 4'htqo "4232"/42350"Vj g'ewt'gpv'rgx'gn'ku'cdqw'vy keg'vj cv'uggp'd{ 'Uw'nc'gv'cr0\*3; ; 9+lp'vj g" o kf/3; ; 2u0'Qvj gt'ukgu.'kpenw'kpi 'vj g'Uqwj 'Dgtt{ 'K'rcpf u'O ctkpg'T gugt'xg'f gerctgf "qp" F gego dgt'4; .422: + 'Uqwj y guv'P gy 'Rtqxf gpeg'P cvk'p'cn'Rctn'ukg.'cpf 'pqt'vj 'Gz wo cu'uwf { 'ukg" cnu'j' cxg'uj qy p'uqo g'k'petgcugu'kp'dkqo cuu'kp't'gegpv'uw'xg{u.'dw'vj g't'gur qpug'ku'o wej 'rguu'vj cp" vj cv'uggp'kp'vj g'Gz wo c'Ec{u'Ncpf "cpf "Ugc'Rctn0

Hkuj kpi 'y cu'emugf 'hqt'c'32'f c{ 'r g'ktqf 'ctqwpf 'vj g'hwm'o qppu'qh'F ge'vj tqwi j 'Hgdwtct { " 3; ; : /4228'v'r tqv'ev'ur cy plpi 'P cuucw'i tqwr gt'cv'vj g'J ki j 'Ec{ 'ci i tgi cvk'p'ukg'cpf 'vj g'gcuw'gt'p" eqcuu'qh'Nqpi 'K'rcpf 0'Qp'F gego dgt'38.'4225.'vj g'Dcj co cu'F kt gev't'qh'Hkuj g'k'gu'cpp'q'w'pegf 'vj g' hktuv'gxgt'emugf 'ugcuq'p'hqt'vj g'ur gekgu.'vj wu'r tqj kdkkpi 'vj tqwi j qw'vj g'eqw'w' { 'vj g'ö'cnkpi . " r'cp'f kpi . 'r tqegu'kpi . 'ugn'kpi 'cpf 'q'htg'kpi 'hqt'uc'ng'qh'ht'guj 'P cuucw'i tqwr gt'ö'f wtkpi 'ur cy plpi " r g'ktqf u0'Uwdugs wgp'w' { 'vj g'em'uw't'g'y cu't'gr n'egf 'd{ 'cp'cpp'w'cm' { 't'gpgy cdng'p'cv'k'py kf g'em'uw't'g'qh' hkuj kpi 'hqt'vj g'P cuucw'i tqwr gt'f wtkpi 'vj g'y k'p'v't'o qpvj u'F gego dgt'v'q'Hgdwtct { +\*F gr ctvo gpv'qh' O ctkpg'T guq'w'tegu'4229.'Ej gwpi 'gv'cr04235+0'Nqec'n'pqp/i qxg'tpo gpv'qti cpk' cvk'p'u'P I Qu+'ctg" y qtn'kpi 'v'j' cxg'vj ku'ej cpi gf 'v'c'r gto cp'gpv't'cvj gt'vj cp'cp'cpp'w'cm' { 't'gpgy cdng'o gcuw't'g" \*DTGGHqti +0""

Vj gtg'ctg'cnu'ugx'g'cni' gct'eqp't'qni'kp'vj g'Dcj co cu't'gr'x'cp'v'htq.'dw'p'qv'ur gek'ke'v'q.'vj g' P cuucw'i tqwr gt'0'Hkuj kpi 'y kj 'UEWDC'cpf 'vj g'wug'qh'g'zr'q'uk'x'gu.'r q'ku'q'pu.'cpf 'ur g'cti w'pu'ku" r tqj kdkgf . 'cnj qwi j 'ur'kpi 'ur g'ct'u'ctg'cm'qy gf 0"Vj g'wug'qh'd'ng'cej 'qt'q'y gt'p'qz'k'q'wu'qt'r q'ku'q'p'q'wu" uwd'uc'p'egu'hqt'hkuj kpi . 'qt'r qu'gu'ku'q'p'qh'u'wej 'uwd'uc'p'egu'q'p'd'q'ctf 'c'hkuj kpi 'x'gu'gn'y kj qw" y tkw'gp'cr r tq'x'cn'q'h'vj g'O k'p'k'v't'g' . 'ku'r tqj kdkgf 0'I qxg'tpo gpv'r q'rk' { 't'g'ut'k'ew'eqo o g't'ek'ni'hkuj kpi " v'q'vj g'p'cv'x'g'r q'r w'cv'k'p'cpf . 'cu'c'eq'p'ugs w'g'peg . 'cm'x'gu'gn'hkuj kpi 'y kj kp'vj g'Dcj co cu'Gz'en'w'k'x'g" Hkuj gt { '\ qpg'o w'w'd'g'hwm' { 'qy p'gf 'd{ 'c'Dcj co k'p'ek'k' gp't'g'uk'f kpi 'kp'vj g'Dcj co cu0

Ur gct'hkuj kpi 'y kj kp'q'pg'o k'g'qh'vj g'eq'cu'v'qh'P gy 'Rtqxf gpeg'cpf 'H'ggr q't'v'cpf "422" { ctf u' qh'vj g'eq'cu'v'qh'cm'q'vj gt'H'co k'k' 'K'rcpf u'ku'r tqj kdkgf . 'cu'ku'vj g'wug'qh'h'k'g'cto u'qt'g'zr'q'uk'x'gu'0'Hqt" p'g'v.'c'o k'p'ko wo 'o guj 'uk' g'qh'4'k'p'0'ku'p'gegu'ct { . 'gz'egr v'y j gp'hkuj kpi 'i qi i ng/g { g'\*dk / g { g'uecf + " qt'r k'raj ctf 0'Hkuj 'v't'r u'ctg't'gs w'k'gf 'v'q'j cxg'ug'h'f g'ut'w'ev'r cp'gn'cpf 'o k'p'ko wo 'o guj 'uk' gu'qh'3'd { " 4'k'p'0'ht'g'ev'cpi w'ct'y k'g'o guj 'v't'r u'c'p'f "307'k'p'0'i t'g'cv'gu'v'rgpi vj 'qh'o guj +hqt'j g'z'ci q'p'cn'y k'g" o guj 'v't'r u'0'C'r gto k'ku't'gs w'k'gf 'v'q'ug'm'ec'v'ej 0'C'r gto k'ku't'gs w'k'gf 'v'q'w'ug'c'k'eqo r t'guu'qt'u'hqt" hkuj kpi 'r w't'r q'ugu'cpf 'vj g'wug'qh'eqo r t'guu'qt'u'ku't'g'ut'k'ev'gf 'v'q'vj g'r g'ktqf "3'C'wi w'w'53'O c'tej 'cpf 'v'q" f gr vj u'qh'32/42'o 0"Vj g'ecr w't'g'qh'i tqwr gt'cpf 't'q'en'hkuj 'y g'ki j kpi 'rguu'vj cp"5'ndu'0'ku'r tqj kdkgf 0" F cj ni tgp'\*r g'tu'0'eqo o 0'j' cu'p'q'v'gf 'c'5'nd'0'P cuucw'i tqwr gt'ku'q'pn' { 'cd'q'w'67'eo 'n'q'pi . 't'q'wi j n' { "5'eo " u'j q't'v't'vj cp'vj g'o k'p'ko wo 'uk' g'qh'o c'w't'k'v' { 'hqt'h'go c'rgu'0" 'F cj ni tgp'uw'i i gu'gf 'vj cv'cp'k'pet'g'cu'g'kp" vj g'ec'v'ej 'r'ko k'v'q'79'eo 'y q'w'f 'g'p'uw't'g'vj cv'cv'rg'cu'v'97' 'qh'hkuj 'eq'w'f 'ur cy p'd'gh'q't'g'rgi c'ni'hkuj gt { "

"  
"  
"  
"

tgo qxcn0'

Ej gpi 'gv'cn0\*4235+'uwi i guvgf "c'tgf wevkp'kp'hkuj kpi "ghqtv'f wtkpi 'vj g'pqp/ur cy plpi " r gtkqf u'htqo 'vj g'3; ; : /4223'ngxgn'vq'ko r tqxg'P cuucw'i tqwr gt 'uwuxkpcdkrkv{0"Vj g{ "cnuq'uxt guugf " vj g'poggf 'vq'gpuwtg'vj cv'r qcej kpi 'ku'eqpvtqmgf 'f wtkpi 'vj g'ur cy plpi 'ugcuqpu0"Vj gug'o gcuwtgu" y qwf "cf f tguu'vj g'o ckp"eqpegtpu'gzr tguugf "d{ 'hkuj gt u'cdqw'vj g'P cuucw'i tqwr gt 'hkuj gt { 'kp" kpvgtxlgy u0"Cmj qwi j "c'tgf wevkp'kp'hkuj kpi "o qtvrk{ 'vj tqwi j 'tgf wevkp"qh'hkuj kpi "ghqtv."o c {" chgev'vj g'uj qt v'vgt "geqpqo le'dgpghku.'vj g'hkuj gt { 'y qwf 'r gthqto "dgwgt'geqpqo lecm{ "cpf 'cu'c" hqf "uqwtg"qxgt 'vj g'qpi /vgt "Ej gwpi 'gv'cn0'4235+0"

Vj gtg'ku'pq'o gej cpkuo 'kp'vj g'Dcj co cu'hqt'f gerctkpi "c'ur gelgu"ögpfcpi gtgf .ö" övj tgcvgpgf .ö"qt"ör tqvgevgf Ö"Vq'cf xkug'vj g'r vdrke'cpf 'f gxgnr 'uwr r qt v'htq. "cpf 'wpf gtucpf kpi " qh'vj g'poggf 'hqt'r tqvgevxg'o gcuwtgu."qwtgcej "eco r cki pu'y gtg"eqpf wevgf "qp'vj g'wkrk{ 'qh'vj g" ugcupcn'hkuj kpi "emquwtgu'cpf 'vq'f kueqwtci g'vj g'r wtej cug'qh'P cuucw'i tqwr gt 'd{ "equwo gt u'f wtkpi " vj g'r tqvgevgf "ugcuqp0"Vj g'kpxcukxg'rkqphkuj 'y cu'uwi i guvgf . 'y kj 'uqo g'uweegu. "cu'cp'cngt pcvkxg" hkuj kpi 'cti gv'cpf 'hqf 'ej qlg0"

"

""

"

"  
"  
"  
"

# BELIZE

## Belize -- Populations

"

Dgrk g.'y kj 'ku'gz vgpukg'tggh'u{uwo "cpf "eqcuvkpg.'y cu'qpeg'c"o clqt'j cdkcv'ht'v'j g" P cuucw'i tqwr gt'cpf "d {"cm'ceeqpwu.'y g'ur gekgu'y cu'j kvqtkecm {"gz v'tgo gn' "cdwpf cpv\*"Etcki " 3; 88+0"i tqwr gt'ci i tgi cvkqpu'j cxg'dggp'ugxgtgn {"tgf wegf "cv'o cp {"m'ecv'kpu'kp"Dgrk g.'cv'ukgu" uvej "cu'O gz'leq'Tqemu.'Tkug'cpf "Hcm'Dcpm'cpf "Ec {g'I mqt {"cu'kpf'kecv'gf "d {"I tggp'Tggh'u" pcv'kqpen'I tqwr gt'Ur cy plpi "Ci i tgi cvkqp'Cuuguo gpv\*"4223+0"Guvko cvgf "pwo dgtu'qh'hkuj "vcn'gp" Itqo "ur cy plpi "ci i tgi cvkqpu'y cu'v'j g'o clp'kpf'kecv'qt'cxck'rdng'qh'r qr w'cv'kqpu'uk' g'y kj "v'j g'ug'qpeg" gzeggf'kpi "cp'guvko cvgf "52'v'j qwucpf "hkuj "f w'kpi "l'wuv'qpg'ur cy plpi "ugcu'qp'cv'l'wuv'qpg"

ci i tgi cvkqp'uksg\*"Ec {g'I mqt {"+cpf " tgr'qtwa'qh'v'g'pu'qh'v'j qwucpf u'qh'hkuj " y gtg'qpeg'v'j g'p'qto "Etcki "3; 88+0" Cv'Ec {g'I mqt {".'y j gtg'i tqwr gt" ecvej gu'tgcej gf "4'v'qpu'r gt'f c {"k'p" v'j g'nc'v'g'3; 82au.'c'lc'p'w'et {"4223" uwt'xg {"m'ec'v'gf "43'hkuj O'hkuj gto gp'cv' v'j g'uksg'ecw' j v'qpn {"; "hkuj "f w'kpi " hqwt'f c {"u'qh'k'p'v'g'pug'hkuj kpi " \*J g{o cp'cpf "Y cf g'4227+0

K'p'Dgrk g.'v'j gtg'ctg'cv'rg'cu'v'37" npqy p'ur cy plpi "ci i tgi cvkqp'uksgu" \*Hki 037+ "v'j cv'q'ee'w'cm'pi "v'j g'dctt'k'gt" tgg'h'cpf "qp'q'w'gt'cv'qm'O'cm'uksgu" q'ee'w'y kj k'p'342'o "qh'v'j g'uj g'h'gf i g." y kj "v'j g'cxg'tci g'f k'uc'p'eg'v'q'v'j g'uj g'h'

gf i g'dgkpi "cdq'w': 2'o 0'O quv'uksgu'ctg'p'gct'k'p'h'g'ev'k'p'r'q'k'p'w'q'h'eq'p'x'gz'uj'cr'gf "ugcy'ctf / gz'v'p'f'kpi "t'g'gh'u\*"y kj k'p'582'o "qh't'g'gh'r'tqo'q'p'v'q't'k'gu+\*"M'q'd'ct'c'cpf "J g{o cp'4229.'M'q'd'ct'c'422; +0" V'j g'ug'h'g'c'w't'gu'j'cxg'd'ggp'w'ugf "v'q'v't {"v'q'k'f'g'p'v'h {"w'p'np'qy p'ur cy plpi "ci i tgi cvkqpu'kp'Dgrk g'cpf " q'v'j gt'r'ctw'q'h'v'j g'P cuucw'i tqwr gtu't'cpi g'dw'j'cxg'd'ggp'u'w'ee'g'u'w'v'q'f'cv'g'O

Ur gekgu'ur gek'he'c'p'p'w'cn'ic'p'f'kpi u'f'c'v'cv'v'j g'p'cv'k'q'p'cn'ix'g'gn'ct'g'p'q'v'cx'ck'rdng.'cn'j'q'w'i'j " u'c't'v'k'pi "k'p'4225.'cp'g'h'q't'v'y cu'w'p'f'g't'c'n'gp'v'q'o'q'p'k'q't'p'w'o' d'g't'q'h'P cuucw'i tqwr gtu'cv'r't'k'q't'k'v {" ur cy plpi "uksgu"O'T'g'eg'p'v'o'q'p'k'q't'k'pi {"k'gr'f'g'f'eq'w'p'w'q'h'c'h'gy "j'w'p'f't'g'f' "hkuj "k'p'o'qu'v't'go'cl'p'k'pi " ci i tgi cvkqpu'uwt'xg'gf {"\*V'cd'ng'32+.'cpf "c'h'gy "v'j q'w'uc'p'f' "hkuj "cv'q'v'j gtu\*"Dgrk g'Ur cy plpi " Ci i tgi cvkqp'y q't'n'k'pi "i tqwr <

\*j w'v'<leq'm'd'q't'cv'k'q'pu'v' eu'q't'i IF g'h'c'w'n'c'ur z'Ac'rc'ku?eq'm'd'q't'cv'k'q'pu'v' eu'q't'i l'ur'ci ( +0'F'qi "H'gc" Ec {g'y cu'j'k'j' r'k'i'j'v'gf "cu'c'uksg'y'j'gt'g'k'm'gi'c'n'h'kuj'kpi "j'cu'eq'p'v'k'p'w'gf "cpf "v'j g'p'w'o' d'gtu'q'h'ur cy p'g'tu" j'cxg'i't'g'c'w' {"f'g'et'g'c'ug'f' "eq'o' r'c't'g'f' "v'q'uksgu'y'kj "i'q'q'f' "g'p'h'q't'ego' gpv\*"g'f' 0'P'G'R'q'k'p'v'k'p'I' m'x'g'tu'T'g'gh"




Figure 13. Known Nassau grouper spawning sites (noted by an orange circle) on the east coast of Belize.

"  
"  
"  
"

cpf "Ucpf dqtg'Ec { 'kp'Nki j vj qwug+<sup>5</sup>Dgk g'Ur cy plpi 'Ci i tgi cvkqp'y qtnkpi 'i tqwr -0'

**Table 5. Number of Nassau grouper at priority spawning aggregation sites in Belize. (Belize Spawning Aggregation Working Group Information Circular 10, November 2012.) Effort is variable as noted in footnotes.**

Maximum Nassau Grouper Counts for 2003 – 2012										
Site	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
Rocky Point	0	200	200	0 <sup>1</sup>	N/A	0	N/A	N/A	0	2
Dog Flea Caye, Turneffe	1,500	100 <sup>3</sup>	-	2 <sup>2</sup>	N/A	N/A	N/A	N/A	4	4
Sandbore, Lighthouse	1,800	2,500	1,800	1,205 <sup>4</sup>	1,495	1,250	2,050	2,000	1,300	1,350
Caye Glory	1,000	1,000	350	7 <sup>5</sup>	69	405	3,000	N/A	400	120
NE Pt., Glover's	2,400	1,700	2,240	3,000	800	1,190	1,100	3,328	1,800	1,050
Gladden Spit <sup>5</sup>	250	450	360	700	500	1,106	260	238	375	164
Nicholas Caye	52	~50	80	48	80	100	25	30	45	85

<sup>1</sup> Only one dive; <sup>2</sup> Site only monitored in February; <sup>3</sup> Site only monitored in January; <sup>4</sup> Probably missed peak spawning due to bad weather, <sup>5</sup> Numbers for Gladden Spit revised by SEA in 2011)

F gerkpgu'lp'vj g'qxgtcm'cdwfp cpeg'qh'P cuucw'i tqwr gt'ecp'dg'kphgttgf 'Htqo 'ur cy plpi " ci i tgi cvkqp'eqwpu'0'O quv'qh'vj g'f gerkpgu'qeewtg'f'r tkqt'vq'vj g'kpkkcvkqp'qh'ur cy plpi 'ci i tgi cvkqp' o qpkqtkpi '\*Vcdrg'32+0'CvI mxgtai'Tggh'vj g'ur cy plpi 'ci i tgi cvkqp'y j kej 'j ctdqg'f'37.222" P cuucw'i tqwr gtu'lp'3; 97'j cf 'f gerkp'g'f'd { ': 2' 'vq'rguu'vj cp'5.222'i tqwr gtu'lp'3; ; ; '\*Ucr'gv'crf' 4223+'cpf 'vq'cdqw'c'vj qwucpf 'kp'4233'cpf '4234'\*q' 0Ucf qx { . 'Wpkxgtukv' 'qh'J qpi 'Mqpi . 'r gtu' qdu'4234+0'Qpn' '4'qh'vj g' ; 'ci i tgi cvkqp'ukgu'kf gp'v'k'kf 'kp'3; ; 6'\*Ectvgt'gv'crf'3; ; 6+'j cf 'o qtg' vj cp'372'P cuucw'i tqwr gtu'=vj g'tguv'qh'vj g'ukgu'j cf 'dggp'hkuj gf 'qw'\*J g { o cp'4223. 'Rc| 'cpf " I tko uj cy '4223+0'Ec { g'I mqt { . 'cnuq'hpqy p'cu'Go kn { . 'y cu'gzr m'k'kf 'hqt'qxgt' : 2' { gtu'y kj " f gerkp'gu'cwt'kdwgf 'vq'rcem'qh'o cpci go gp'v'cpf 'c'hwet'cvkxg'hkuj kpi 'kpf wut { 'vj cv'cwt'cev'f' o cp { " hkuj gtu'\*Rc| 'cpf 'Vtw' '4229+0'Cm'hpqy p'ci i tgi cvkqp'ukgu'j cxg'w'p'f gti qpg'f tco cvk'f gerkp'gu'lp' vj g'cdw'p'f cpeg'qh'ur cy plpi 'hkuj 'qxgt'vj g'rcu'v'y q'f gecf gu'0'Ewt'gp'v'ci i tgi cvkqp'r tqv'gcvkqp'f qgu' p'q'v'cr r gct'vq'dg'tguv'qt'kpi 'vj ku'ur geku'c'mj qwi j 'cm qu'v'egt'cvkpn' { 'vj g'gh'qt'w'j' cxg'uv'go o gf " f gerkp'g'0'"

Ci i tgi cvkqp'pwo dgtu'cuugu'gf 'f wt'kpi 'vj g'r g'k'kf '4225/4229'tcpi gf 'Htqo 'c'j ki j 'qh'5.222" hkuj 'cv'I mxgtai'Tggh'vq'my u'qh'rguu'vj cp'32'hkuj 'cv'vj tgg'q'vj gt'ukgu.'cmj qwi j 'k'y cu'p'q'v'f 'vj cv' uwt'xg { u'y g'tg'p'q'v'cny c { u'cu'eqo r r'gv'cu'f gult'gf 0'Vj g'4234'f cv'uj qy gf 'y q'ukgu'y kj 'hgy gt' vj cp'7'hkuj . 'vj tgg'ukgu'y kj 'rguu'vj cp'422'hkuj 'cpf 'y q'ukgu'y kj 'dgy ggp'3222/3722'hkuj 0" Cr r ct'gp'v'f gerkp'gu't'guwo gf 'ch'gt 'vj g'422; /4232'uw'xg { u'0'"

Ugxgt'cn'uw'f kgu'j cxg'gzco k'p'gf 'o q'xgo gpw'qh'v'ci i gf 'hkuj 0'Vj g'bo q'xgo gpw'qh'P cuucw' i tqwr gt'cm'pi 'vj g'dctt'k'gt'eq'cu'cn't'ggh'j' cxg'dggp't'geq't'f gf 'kp'gz'egu'qh'422'mo '\*Ectvgt'gv'crf' 3; ; 6+0'CvI mxgtai'Tggh'\*cp'cv'm'±'P cuucw'i tqwr gt'uj qy gf 'ut'qpi 'h'f g'k'v' 'vq'dq'vj 'p'q'p/ tgr tqf w'v'k'x'g'cpf 'ur cy plpi 'ct'geu'qp'vj g'cv'qm'cpf 'o c { 'p'q'v'bo ki t'cv'g'cv'cm'\*Uctt'gv'crf'4229+0" Dcu'gf "qp'vj g'h'p'f kpi u'qh'ce'q'w'k'e'v'rgo g't { . 'P cuucw'i tqwr gt'gz'j k'k'k'gf 'i tgc'v' { 'u'pej t'q'p'q'w' u' o ki t'cv'k'p'v'q'ur cy plpi 'ukgu'f wt'kpi 'h'm'no q'p'p'u'ht'qo 'F gego dgt'vj tqwi j 'O cte'j 'f gur k'g'vj g'k' " q'vj gty kug'v'ur'k'ct { 'j cdku'0'I tqwr u'qh'72/322'hkuj 'j' cxg'dggp'q'dugt'x'gf 'o q'x'k'pi 'vq'ci i tgi cvkqp' ukgu'0'T gr tqf w'v'k'x'g'cf w'nu'o q'x'g'ht'qo 'vj g'k'uj cm'j 'y cv'gt'j cdk'cv'f wt'kpi 'vj g'y k'p'v'gt' h'm'no q'p'p'u' cpf 'o ki t'cv'g'v'q'vj g'uco g'ur cy plpi 'uk'g'w'r 'vq'h'q'w'v'ko gu'r g't { gct'uv' { kpi 'cp'cx'gt'ci g'qh'3308'f c { u' cv'vj g'uk'g'f wt'kpi 'vj g'y k'p'v'gt' h'm'no q'p'p'u'\*Uctt'gv'crf'4229+0'Wuk'pi 'v'ci i kpi 'y kj 'XGO EQ'X38"

"  
"  
"  
"



Hkuj kpi 'hqt'P cuucw'i tqwr gt'qwukf g'qh'v'j g'ci i tgi cvkqp'uksgu'eqpvkpwgu'v'q'dg'ko r qtvcpv0"  
Dgi kppkpi 'kp'v'j g'3; 92u.'P cuucw'i tqwr gt'y gtg'vcnmp'v'j tqwi j qw'v'j g'gct'RC| "cpf "Vtwn{ "4229+ "  
y kj "Ucrn"gv'cr0\*4223+pqv'kpi 'v'j cv'36' "qh'v'j g'cf wv'r qr wv'v'kqp'ku'tgo qxgf "cppwcm{ "d{ "{gct/  
tqwpf "ur gct'hkuj kpi 0"khqtto cvkqp"qp'ngpi v'j "qh'P cuucw'i tqwr gt'ecwi j v'qwukf g'qh'v'j g'ur cy pl'pi "  
ugcuqp"uwi i guu'v'j g'uactv'qh'c'tgeqxtg {0'Cv'I mxgt'au'tggh'rkngn{ "c'rcti gn{ "ugrht'getv'k'kpi "ctgc."  
uwxg{u'qh'hkuj gt'ecvej gu'ht'qo "4226'v'q'4232'uwi i guv'cp'kpetgcug'kp'cxgtci g'ngpi v'j "qh'P cuucw'  
i tqwr gt'ht'qo "c'b' gcp'qh'593'o o "6'6; 5'b' o "VN'kp'4229'v'q'785'o o "VN'kp'4232"LOI kduqp."  
Y kf rkhg'Eqpugt'xcv'kqp'Uqekgv{ "/Dgrk' g'Ekv{ .Dgrk' g.'r'gtu0'eqo o 0'v'q' [ 0Ucf qx{ .Wpkxgtukv{ "J qpi "  
Mqpi . "4232+0""  
"

**Belize – Conservation and Management**

"

O cpci go gpv'qh'v'j g'P cuucw'i tqwr gt'j cu'c'ngpi 'j kuvqt { 'kp'Dgrk' g'gxgp'v'j qwi j 'cppwcn'  
ncpf kpi u'ht'v'j g'ur geku'ct'g'pqv'cxck'cdng0""kpuvcf . 'v'j g'ucw'u'ku'f gv'to kpgf "d{ "pwo dgtu'qh'hkuj "cv'  
ur cy pl'pi "ci i tgi cvkqp'cpf "cnuq'd{ "hkuj gto gp'g'zr g'k'p'eguc'cpf "ur q'ctf k'e'tgr qt'v'0"Vj g'Hkuj g'tkgu"  
F gr ct'vo gpv'ku't'gur qpukdng'ht'v'j g'o qpkqt'kpi . 'eqpv'qn'cpf "uwxg'k'nc'peg'qh'v'j g'hkuj kpi 'kpf wut { "  
\*Ecteco q'422: +0"Vj g'ht'uv'o gcuwt'g'v'q'r tqv'ev'P cuucw'i tqwr gt'y cu'c'ugcuqpcn'eqmwt'g'y kj kp'v'j g'  
I mxgt'au'Tggh'O ct'kpg'Tgugt'xg'kp'3; ; 5=v'j g'ct'gc'y cu'emugf 'ht'qo 'F gego dgt'3'v'q'O ctej "3'qh'v'j g'  
hqmqy kpi "{gct0'k'3; ; 8.'v'j g'pgy "o ct'kpg'tgugt'xg."Dcecrnt'Ej k'eq."cnuq'k'p'nmf gf "c'ugcuqpcn'eqmwt'g'  
| qpg'ht'v'j g'r tqv'ev'kqp'qh'v'j g'P cuucw'i tqwr gt'ur cy pl'pi "ci i tgi cvkqp"RC| "cpf "Vtwn{ "4229+0"  
O k'p'ko wo "cpf "b' cz'ko wo "ecr wt'g'uk' gu'y gt'g'k'p'v'qf weg'f "c'f gecf g'ci q'Ucrn"gv'cr0'4223+Ect'v'gt'gv'  
cr0'3; ; 6=J g{o cp'cpf "Tgs w'gpc'4224=LOI kduqp."Y kf rkhg'Eqpugt'xcv'kqp'Uqekgv{ "/Dgrk' g'Ekv{ . "  
Dgrk' g.'r'gtu0'eqo o 0'v'q' [ 0Ucf qx{ .Wpkxgtukv{ "J qpi "Mqpi . "4232=Ucf qx{ 'f'g'O kej guqp'gv'cr0'  
422: +0"

I kxgp'i tqy kpi 'k'p'v'gt'gu'v'cpf "eqpegtp'ht'v'j g'ur geku.'kp'4223'v'j g'Dgrk' g'P cvkqpcn'  
Ur cy pl'pi "Ci i tgi cvkqp"Y qtnkpi "I tqwr 'y cu'guv'cd'kuj gf 0'F wtkpi "4224."c'eqcrk'kqp'qh'ugxgp"  
P I Qu.'i qxgtpo gpv.'hkuj gtu."cpf "q'v'j gt'ucng'j qrf gtu'y qtn'gf "u'weeg'uh'wm{ "v'j guv'cd'kuj "r tqv'ev'xg"  
ngi k'ur'v'kqp'ht'33'qh'v'j g'npqy p'P cuucw'i tqwr gt'ur cy pl'pi "ukgu."cpf "v'q'k'p'v'qf weg'c'ht'wt/o qpyj "  
enugf 'tgr tqf w'v'xg'ugcuqp'kp'4225"Q'EQppqt'4224."I kduqp'422: +0"Ugxgp'qh'v'j qug'33'ukgu"  
\*Vcdng'32+ct'g'o qpkqt'gf "cu'tgi w'ctn{ "cu'r quukdng'cpf "k'p'nmf g<'T'qem{ "R'0'Dcecrnt'Ej k'eq'O ct'kpg'  
Tgugt'xg+."F qi h'rc'Ec { g'V'wtpghg'K'ur'p'f u+."Uc'p'f d'q'g'N'ki j v'j q'wug'Tggh+."Go k'nf IEc { g'I nqt { . "  
I ncf f'gp'Ur k'v'f ncf f'gp'Ur k'v'f'cpf "Uk'm'Ec { gu'O ct'kpg'Tgugt'xg+."P q'v'j g'cu'v'R'q'k'p'v'f I mxgt'au'Tggh'  
O ct'kpg'Tgugt'xg+cpf "P k'ej q'ru'Ec { g'Ucr qf k'nc'Ec { gu'O ct'kpg'Tgugt'xg+0"Vj g'Y qtnkpi "I tqwr "  
o g'gw't'gi w'ctn{ "v'j 'uj ct'g'f'c'v'c'cpf "f'g'x'gn'r "o cpci go gpv'ut'cv'gi k'gu'y y y **Ur ci dgrk' g'Q'iti** =t'g'v'k'xg'f "  
qp'37'Cr tk'4234+cpf "o qpkqt'kpi "eqpv'kpwgu'cv'ugxg'tcn'uksgu0'

k'p'4225.'v'j q'Uc'w'w'qt { "k'p'ut'wo gp'u'y gt'g'g'p'cev'f 0"Vj g'ht'uv'f ger'ct'gf "33'ukgu.'k'p'nmf kpi "  
oGo k'nf o'Ec { g'I nqt { +."cu'o ct'kpg'tgugt'xg'u'enugf "v'j 'hkuj kpi "cm{gct'tqwpf 0"Vj qug'uksgu'v'j cv'y gt'g"  
y j qm{ "qt'r ct'v'cm{ "h'ec'v'gf "kp'o ct'kpg'tgugt'xg."dw'pq'v'k'p'nmf gf "kp'cp{ "ugcuqpcn'eqmwt'g'qt"  
eqpugt'xcv'kqp" | qpg."eqwf "dg'w'ugf "qpn{ "d{ "t'cf k'k'qpcn'hkuj gto gp't'geqo o g'p'f gf "d{ "v'j g't'gur gev'xg'g"

"  
"  
"  
"

eq/o cpci gtu'qh'vj g'tgugtxg'cpf 'y kj 'ur gekn'hegpug'i tcpvgf 'd{ 'vj g'Hkuj g'kgu'cf o kpkmtcvqt0'Vj g" ugeqpf "Ucwwqt{ "kputwo gpv'gucdrkuj gf "c'hqwt/o qpvj "emugf "ugcuqp"vq'r tqgev'ur cy plpi "P cuucw" i tqwr gt."gzvpgf lpi 'htgo "F gego dgt'vq'O ctej 0'Hkuj gto gp'cv'O cwi tg'Ec{g'cpf "P qt'vj gtp"Vy q" Ec{gu."j qy gxgt."y gtg'cmqy gf "vq'hkuj 'f wtkpi 'vj g'ur cy plpi "ugcuqp."dw'qpn{ "vpf gt'ur gekn'hegpug" i tcpvgf "d{ 'vj g'Hkuj g'kgu'cf o kpkmtcvqt."c'eqpf kkpq"qh'y j lej "y cu'vj cv'cm'ecvej "y qwf "dg'xgt'kkgf " d{ "c'Hkuj g'kgu'Qh'hegt \*Rc| "cpf "Vtwn{ "4229+lp"qtf gt 'vq'o qpkqt "uqenlwtwewt g0'Vj gug" gzeqr vqpu'o cf g'vj g'pvcvqpcnr tqgev'vq'f khlwv'vq'gphqteg'cpf "j gpeg'uvct vki "lp'vj g"4232/4233" ugcuaq."ur gekn'hegpugu'vq'hkuj 'hqt"P cuucw'i tqwr gt'cv'vj gug'vy q'ukgu'f wtkpi "vj g'emugf "ugcuqp" y gtg'pq'hqpi gt'kuwgf 0'Vj gug'hkpcn'vy q'ukgu."j qy gxgt."ctg'pqv{gvf guki pcvgf "cu'hwm{ 'r tqvevgf " ctgcu'emugf "vq'hkuj lpi 0'Vj gtghqtg."35'qh'vj g"37'npqy p'ci i tgi cvkqp'ukgu'ctg'hwm{ "emugf "vq" hkuj lpi "f wtkpi "vj g'ur cy plpi "ugcuqp0'Qh'vj g'tgo clkpi "vy q'npqy p'ci i tgi cvkqp'ukgu."O cwi tg" Ec{g'uj qwf "dg'r tqvevgf "y j gp'vj g'Vwtpghg"Krcpf u'o ctkpg'tgugtxg'ku'f gerctgf 0'Dgrk g'ku'ukm' uggmkpi "P qt'vj "Vy q'Ec{g}u'r tqgev'vq'p0'

Kp"gtcn{ "Cr tkn422; ."vj g'O kpkugt'qh'Hkuj g'kgu'uki pgf "kvq'rcy "cf f kkpccn'o gcuwtgu'vq'j gr " o cpci g'cpf 'r tqgev'vj g'P cuucw'i tqwr gt0'Vj gug'kpenmf g'o kpk wo "cpf "o czko wo "uk g'uko ku'qh" 732"o o \*42'kpej gu+cpf "982"o o \*52'kpej gu+ "tgr gev'xgn{."cpf "c'r rppgf "dcp'qp'ur gct'hkuj lpi " y kj kp'cm'o ctkpg'tgugtxg{\*gv'vq'dg'ko r ngo gpvgf +0'Hwtj gto qtg."cu'c'rci g'r tqr qt'vq'qh'hkphkuj " ctg'rcpf gf "cu'hkngw."vj g'pgy "tgi wv'vqpu'tgs wkt'vj cv'cm'P cuucw'i tqwr gt'dg'rcpf gf "y j qrg."cpf "kh' hknvgf "o wv'vj cxg'c'3/4'kpej \*47/72"o o +unkp'r cvej \*Vj g'Dgrk g'Ur cy plpi "Ci i tgi cvkqp" Y qtnkpi "I tqwr "422; +0'Qvj gt "i gct'tgukt'vqpu'ctg'lp'r rneg'hqt'tggh'hkuj gu'i gpgtcm{ "vq'ckf "lp'vj gkt" o cpci go gpv.'uwej "cu'pq'ur gct'hkuj lpi "qp'eqo r tguugf "ckt0"

I kduqp"gv'cr0\*4229+lpf kecvf "vj cv'vj g'r tqxkukp'qh'cuukcpeg'hqt"o cpci go gpv'cpf " gphqtego gpv."cpf "uwuclpki "vj g'r qnk'ecn'ly kn'cv'vj g'j ki j guv'rgxgn."y qwf "dg'pgeguuct { "vq'gphqteg" vj g'rcy u'vq'gpj cpeg'vj g'r tqgev'vq'qh'P cuucw'i tqwr gt'ur cy plpi "ci i tgi cvkqp'ukgu'ctg'hwm{ g0'Vj gtg" j cu'dggp"gzv'pukg'r wdrke"qwtgcej "lp'vj g'eqv'vq' { "vq'kphqto "vj g'r wdrke"qh'vj g'o cpci go gpv" o gcuwtgu'cpf "vj g'pggf "vq'r tqgev'vj g'P cuucw'i tqwr gt."kpenmf lpi "hko ."VX."tcf kq."gve0'Crnj qwi j " o ctngf "tgeqxt'kgu'j cxg'pqv{gv'dggp'pqvgf "hmqy lpi "ko r ngo gpvcvq'qh'o cpci go gpv.'k'ku" cm qu'vegt'vq'vj cv'vj ku'j cu'r tgxgpvgf "hwtj gt'f gen'pku'cpf "o qtg'vko g'y kn'dg'pggf gf "hqt'tgeqxt { " vq'dg'gxf gp0'Vj g'o wnk'ugevqt'pvcvqpcn'ly qtnkpi "i tqwr "o qf gn'lp'Dgrk g'cr r gctu'vq'j cxg'dggp" xgt { "gh'ge'v'xg'lp'i cvj gtlpi "uwr r qtv'hqt"o cpci go gpv'o gcuwtgu'cpf "o c { "ugt'xg'cu'c'wughw'o qf gr0"

"

"  
"  
"  
"

**BERMUDA**

**Bermuda – Populations**

Vj g'w'pf gtuc'pf kpi "qh'r qr w'v'k'p'ej cpi g'c'p'f "u'c'w'u'q'h'P cuuc'w'i t'q'w' g't'k'p'D'g't'o w'f'c'o w'w'v' dg'f' g't'k'x'g'f "h't'q'o "c'e'q'o d'k'p'c'k'p'q'h'g'e'q'm'i l'ec'n'u'w'f' l'g'u'c'p'f "h'k'u'j' g't' { 'f' g'r' g'p'f' g'p'v'f' c'v'e't'g'r' q't'w'c'u' ur' g'e'k'g'u/ur' g'e'k'h'e' l'p'h'q't'o c'v'k'p'k'u'p'q'v'c'x'c'k'r'd'g'0'k'p'c'j' k'u'q't'l'ec'n'e'q'p'v'g'z'v'i' t'q'w' g't'u'j' c'x'g'f' q'o' k'p'c'v'g'f' " D'g't'o w'f'c'u' h'k'u'j' g't'k'g'u'0'D'c't'f' c'ej' "g'v'c'r'0\*3; 7: +f' k'u'e'w'u'g'f' "y'j' g'c'd'w'p'f' c'p'eg'c'p'f' "k'o' r' q't'v'c'p'eg'q'h'i' t'q'w' g't'u' v'q' "y'j' g'k'u'r'p'f' "y'j' k'g'r' t'q'x'k'f' k'p'i' "l'p'h'q't'o c'v'k'p'q'p'c'u'r' g'e'u'q'h'v'j' g'k'd'k'q'm'i' { 0'F' g'p'u'k'v' { "q'h'P' c'u'c'w'i' t'q'w' g't' q'p' "u'j' c'm'q'y' "t'g'g'h'u'k'p'D'g't'o w'f'c'k'p' "y'j' g'3; 72u'y' c'u'g'u'k'o' c'v'g'f' "c'v'34'h'k'u'j' "r' g't'c'e't'g'\*568' l'j' g'e'v'c't'g'+ 'y' k'j' " y'j' g'h'k'u'j' "y' g'k'i' j' k'p'i' "c'p'c'x'g't'c'i' g'q'h'308'm'i' \*4064' "r'd'u'0'D'c't'f' c'ej' "c'p'f' "O' g'p'l' g'r'i'3; 79+0'D'c't'f' c'ej' "g'v'c'r'0 \*3; 7: +g'u'k'o' c'v'g'f' "y'j' c'v'i' t'q'w' g't'u'e'q'o' r' t'k'u'g'f' "c'r' r' t'q'z'k'o' c'v'g'n' { '92' "q'h'v'q'v'c'n'h'q'q'f' /h'k'u'j' "r'c'p'f' k'p'i' u'f' w't'k'p'i' " y'j' g'r' g't'k'q'f' "q'h'v'j' g'k' "u'w'f' { \*o' k'f' "3; 72u+ 'y' k'j' "u'p'c'r' r' g't'u'e'q'p'v'k'd'w'k'p'i' "42' "v'q' "y'j' g'v'q'v'c'r'0" E'w'o' w'v'c'x'g'f' f' c'v'e' "h't'q'o' "TGGH\*4225/4235+ "t'g'r' q't'v'g'f' "p'l'k'p'g'P' c'u'c'w'i' t'q'w' g't'k'p' "37; 6' "u'w't'x'g' { u'f' "f' g'p'u'k'v' { "k'p'f' g'z' "308. " u'k'i' j' v'k'p'i' "h't'g's' w'g'p'e' { "208' + "c'e't'q'u'i' "y'j' g'32/ { g'e't' "r' g't'k'q'f' "

\*j' w'r' <l'y' y' 0'g'g'h'q't'i' l'f' d' l'g'r' q't'w'v'f' k'u'v'l'ur' g'e'k'g'u' "V'Y' C' 122; 9'4225/23/23'4235/26/29+0" "V'j' g'u'g'f' c'v'e' " k'p'f' l'ec'v'g'c' "u'k'p'i' n'g' "P' c'u'c'w'i' t'q'w' g't' "y' c'u' "u'w't'x'g' { g'f' "q'p'c' "f' k'x'g' "q'p' "q'p'n' { "208' "q'h'v'j' g'f' k'x'g'u'0" "Y' j' k'g' "y'j' g' " u'w't'x'g' { u'f' "q' "p'q'v'j' c'x'g'c' "y' c' { "v'q' "e'q'p'x'g't'v' "v'q' "c't'g'e'n' "e'q'o' r' c't'k'u'q'p'u' "y'j' g' "h't'g's' w'g'p'e' { "q'h' "q'ee'w't'g'p'eg' "k'u's' w'k'g' "




Figure 14. Proportions of Groupers and Snappers in Commercial Landings from Bermuda 1975-1992.

n'y' "e'q'o' r' c't'g'f' "v'q' "g'e't'r'k'g't' "e'q'p'f' k'k'q'p'u' \*g'f' 0'D'c't'f' c'ej' " c'p'f' "O' g'p'l' g'r'i'3; 79+0'

k'p' "3; 97. "c' "h'k'u'j' g't'k'g'u' "u'c'v'k'u'k'e'u'r' t'q'i' t'c'o' " d'g'e'c'o' g' "h'w'm' { "q'r' g't'c'v'k'p'c'n'r' t'q'x'k'f' k'p'i' "e'c'v'ej' "c'p'f' "g'h'q't'v' f' c'v'e' "h't'q'o' "y'j' g'k'p'f' w'u't' { "q'p'c' "e'q'o' r' w'u'q't' { "d'c'u'k'u'0' "k'p' " y'j' g' "h'k'u'v' { g'e't' "q'h'v'j' g'r' t'q'i' t'c'o' " "H'k'i' 038+ "i' t'q'w' g't'u' " e'q'o' r' t'k'u'g'f' "6908' "q'h'v'j' g'v'q'v'c'n' "r'c'p'f' g'f' "y' g'k'i' j' v'q'h' " h'q'q'f' /h'k'u'j' "v'q'v'c'n' "653" o' w'v'y' j' k'g' "u'p'c'r' r' g't'u'e'q'p'v'k'd'w'g'f' " ; 0' " "N'w'e'n'j' w't'u'v'c'p'f' "Y' c't'f' "3; ; 8+0" "N'c'p'f' k'p'i' u' " f' g'e'n'k'p'g'f' "f' t'c'u'v'k'c'm' { "d'g'y' g'g'p' "3; 97" "c'p'f' "3; : 3" "N'w'e'n'j' w't'u'v' "3; ; 8+0" "V'j' g'i' t'q'w' g't' "r'c'p'f' k'p'i' u'c'v'v'j' k'u' " v'k'o' g'y' g't'g'f' q'o' k'p'c'v'g'f' "d' { "t'g'f' "j' k'p'f' " "H'k'i' 039+0" "D' { " 3; ; ; . "u'r' g'e'k'g'u' "e'q'o' r' q'u'k'k'q'p' "j' c'f' "d'g'g'p' "t'g'f' w'eg'f' "

u'k'i' p'h'k'ec'p'v' { "y' k'j' "y'j' g'i' t'q'w' g't' "r'c'p'f' k'p'i' u' "d'g'k'p'i' "t'g'f' w'eg'f' "v'q' "3; 0' "q'h'v'j' g'v'q'v'c'n' "y'j' k'g' "u'p'c'r' r' g't'u' "y' g't'g' " r'e't'i' g'n' { "w'p'ej' c'p'i' g'f' "c'v'3208" " "H'k'i' 038+0"

V'j' g'q'x'g't'c'n'r' c'w'g't'p' "k'p' "r'c'p'f' k'p'i' u' "q'h'i' t'q'w' g't'u'f' g'e'n'k'p'g'f' "u'j' c't'r' n' { "h't'q'o' "c'd'q'w' "453" o' v'k'p' "3; 97" "v'q' " c'r' r' t'q'z'k'o' c'v'g'n' { "7: "o' v'k'p' "3; ; 3" "H'k'i' 039+ "h'q'm'q'y' g'f' "d' { "c'p' "k'p'et'g'c'u'k'p'i' "t'g'p'f' "w'p'v'k'i' "3; ; ; 0'F' w't'k'p'i' "y'j' c'v' " v'k'o' g' "y'j' g' "u'r' g'e'k'g'u' "e'q'o' r' q'u'k'k'q'p' "q'h'v'j' g'i' t'q'w' g't' "e'c'v'ej' "e'j' c'p'i' g'f' "o' c't'n'g'f' n' { "f' w't'k'p'i' "y'j' g'3; ; 2u' "h't'q'o' "t'g'f' " j' k'p'f' "v'q' "y'j' q' "u'o' c'm'g't' "u'r' g'e'k'g'u' "e'q'p'g' { "c'p'f' "e't'g'q'n'g' /h'k'u'j' "+ "e'q'o' r' t'k'u'k'p'i' "c'm' q'u'v' "72' "q'h'v'q'v'c'n' "r'c'p'f' k'p'i' u' "k'p' " 3; ; ; "N'w'e'n'j' w't'u'v'c'p'f' "Y' c't'f' "3; ; 8+0" "C' "h'k'u'j' "r' q'v'd'ep' "y' c'u'r' w' "k'p'v'q' "g'h'g'ev' "k'p' "C'r' t'k'i' "3; ; 2' "k'p' "c'p' "g'h'q't'v' "v'q' " c'm'q'y' "y'j' g' "t'g'e'q'x'g't' { "q'h' "t'g'g'h' "h'k'u'j' "u'v'q'e'm'u' "y'j' k'ej' "j' c'f' "d'g'g'p' "u'w'd'l'g'ev'g'f' "v'q' "j' g'c'x' { "h'k'u'j' k'p'i' "r' t'g'u'w't'g' "y' k'j' " h'k'u'j' "r' q'u' "t'c'r' u+0'

"  
"  
"  
"

Hqmqy kpi 'vj g'hkuj 'r qv'dcp'kp'3; ; 2.'vj g' vqcnf tqr gt'rcpf kpi u'rgxgnf genkpgf 'd{'7: ' " cpf'tgo ckpgf 'uxdng'y tqwi j '3; ; 4\*'Hki wtg'38+0" Cp'cpcn{uku'qh'y g'tgpf u'kp'kp'kxf wcnf tqr gt' ur gekgu'kp'kcvgu'y g'tgrvkg'eqpvtkdwkqp'qh' gcej 'ur gekgu'v'y ku'i gpgtcnr cwgtp0"P cuucw' i tqr gt'rcpf kpi u'uj qy 'c'uvgr 'f genkpg'htqo 'qxgt' 55'o v'kp'3; 97'vq'rgu'y cp'4'o v'kp'3; ; 3.'c'f tqr " qh'; 7Q' 'kp'rcpf kpi u0'F gur kg'qxgt'32/{gctu'qh' pq/vcng'r tqvkvqp'qh'y g'P cuucw'i tqr gt'kp' Dgto wf c.'vj gt'g'j cu'pqv'dggp'cp'cr r tgekdrng' tgeqxt { 'cpf 'pwo dgtu'tgo ckpgf 'gz vgo gn{ 'mqy " cu'qh'3; ; ; 'cpf 'kp'v'y g'gctn{ '4222u\*'Ucf qx { 'cpf " Gmwpf '3; ; ; . 'Ugo o gpu'gv'cr0422: c+0'Vj g'ur gekgu'j cf 'pqv'uj qy p'cp { 'gxf gpeg'qh'c'uwdugs wgpv' tgeqxt { 'd{ '4227'Nwenj wtu'4227+0'cmj qwi j 'vj g{ 'ctg'vkn'eqpukf gtg' tctg.'vj gt'g'ctg'uqo g' cpgef qcntr qt w'd { 'f kxgtu'qh'o qtg'P cuucw'i tqr gt'kp'vj g'r cu'32/37" {gctu\*'D0Nwenj wtu." Dgto wf c'F gr ctvo gpv'qh'Ci tlewwtg.'Hkuj g'kju.'cpf 'Rctm.'F kxkukqp'qh'Hkuj g'kju.'r gtu0eqo o 0vq' [ 0Ucf qx { . 'Wpkxgtuk{ 'qh'J qpi 'Mqpi . '4234+0'Cu'ht'cu'ku'hpqy p. 'P cuucw'i tqr gt'ur cy plpi " ci i tgi cvkpu'pq'iqpi gt'hqto 'kp'Dgto wf c\*'D0Nwenj wtu.'Dgto wf c'F gr ctvo gpv'qh'Ci tlewwtg.' Hkuj g'kju.'cpf 'Rctm.'F kxkukqp'qh'Hkuj g'kju.'r gtu0eqo o 0vq' [ 0Ucf qx { . 'Wpkxgtuk{ 'qh'J qpi 'Mqpi . ' 4234+0'




Figure 15. Proportion of grouper species in landing from Bermuda commercial catch

### Bermuda – Fishing

"

Kp'vj g'3; 72u.'vj g'cppwcn'hqf/hkuj 'j ctvgn'vqcnf 'cr r tqzko cvgn{ '672.222'mi '\*672'o w' cpf'eqpukv'qh'cp'gunko cvgf '92' 'i tqr gt' 'i tqr gt'cpf'tqen'kuj '+42' 'upcr r gt='; ' 'lcm' o cengtgn'cpf 'wpc'cpf '3' 'qj gt'ur gekgu'g0'j qi hkuj '+\*Dctfcej 'gv'cr03; 7: +0'Ceeqtf kpi 'vq' hkuj gt { 'tgeqtf u'cxckrdng'ukpeg'3; 97.'eqo o gteknf tqr gt'rcpf kpi u'f genkpgf 'kp'Dgto wf c'f gur kg' cp'kpetgcug'kp'ghqtv'qxgt'vj g'r g'kqf '\*Dcpgtqv'gv'cr03; : 9+0'D{ '3; ; ; . 'vj g'vqcn'ecvej 'qh'hqf/hkuj " j cf 'kpetgcugf 'vq'cdqw'843.222'mi '\*843'o v+r gt' { gct0'Vj g'eqo r qukkqp'qh'y g'ecvej 'kp'3; ; ; " uj qy gf 'uki p'k'ecp'ej cpi gu.'3: 0' 'qh'y g'ecvej 'eqpukv'qh'i tqr gt'=3208' 'upcr r gt'=37' " lcmu'=47' 'wpcu'cpf 'tgrv'ur gekgu'cpf '53' 'y cu'eqo r tkugf 'qh'o kuegmcpqwu'tggh'hkuj . 'uwej 'cu' r cttq'vkuj . 'r qti { . 'i twpv.'vki i g'kuj . 'j qi hkuj 'cpf 'Dgto wf c'ej wd0'Vj g'uj kv'htqo 'c'ecvej " f qo kpcv'gf 'd{ 'i tqr gt'cpf 'upcr r gt'vq'qp'f qo kpcv'gf 'd{ 'j gtdkxqtqwu'tggh'hkuj . 'uwej 'cu'r cttq'vkuj " cpf 'uwti gqphkuj . 'tguwngf 'htqo 'vj g'ugxgt'f genkpg'kp'vj g'r tghgtgf 'vcti gv'ur gekgu' 'i tqr gtu' '\*DwtpgwJ gtngu'cpf 'Dcpgu'3; ; 8+0'Y j kg'cmf tqr gtu'y gt'g'chgevgf . 'co qpi 'vj qug'o quv' ugxgtgnf 'tgf wegf 'y cu'y g'P cuucw'i tqr gt0'Ncpf kpi u'qh'P cuucw'i tqr gt'f genkpgf 'htqo '38' 'qh' vqcnf tqr gt'\*cm'ur gekgu'+ecvej . 'd{ 'y gli j v'kp'3; 97'vq'>3' 'kp'3; ; ; '\*Dcpgtqv'gv'cr03; : 9." Tgr qt'v'qh'y g'Eqo o kukqp'qh'kps vkt { . 'Dgto wf c'3; ; 3+0'

"  
"  
"  
"

P cuucw'i tqwr gt'y gtg'hkuj gf 'r tko ctkn'f wtkpi 'ci i tgi cvkqp'r gtkqf u'wukpi 'j cpf rkpqu.'tcr u." cpf 'ur gcti wpu=eqo o gtekn'hkuj gto gp'gZR mkgf 'eqpepvtcvkqpu'hqt'i gpgtcvkqpu'Dctfcej 'gv'crf' 3; 7: .DwtpgwJ gtngu'3; 97+0"Ci i tgi cvkqpu'y gtg'npqy p'htqo 'y g'Ej cmgpi gt'cpf 'Cti wu" \*Rrcpci gpgv+'dcpm0"Vj tgg'ukgu'y gtg'hkuj gf 'wvkn'y g'o kf/3; 92u'DwtpgwJ gtngu'3; 97+0"D{ " 3; : 3."cm'hqwt'npqy p'j kxqtkecn'ci i tgi cvkqp'ukgu'pq'npqi gt'htqo gf 'cpf 'j cf 'r tqdcdn'etcuj gf " ceeqtf kpi 'vq'hkuj gt'ceeqwpw'Dcppgtqv'gv'crf'3; : 9."Nwenj wtuv'3; ; 8+0"F gur kg'uwdugs wgpv' r tqvevqkq."y g'hkuj gt { 'hqt'y ku'ur geku'ku'eqpukf gtgf 'eqo o gtekm'gzvkev'Dcppgtqv'gv'crf'3; : 9=Nwenj wtuv'3; ; 8=D0Nwenj wtuv.'Dgto wf c'F gr ctvo gpv'qh'Ci tlewwtg.'Hkuj gtkgu."cpf 'Rctmu." F kxkukq'qh'Hkuj gtkgu.'r gtu0eqo o 0'vq' l 0Ucf qx { .Wpkxgtuk{ 'qh'J qpi 'Mqpi . 'Ugr 04234+0"

K'cr r gctu'y cv'y g'ur cy plpi 'uqem'dkqo cuu'y cu'tgf wegf 'dgnqy "c'etk'kecn'dw'wmpqy p" r xgn'uq'y cv'y g'r qr wcvkqp'j cu'cr r ctgpv' dggp'wpcdn'vq'tgeqxgt 'ugg'Ucf qx { '3; ; 8+0"O gcp'uk' g" cpf 'htgs wpe { 'qh'uki j vki 'j cu'tghrgevgf 'y gug'ej cpi gu0"O gcp'uk' g'uco r nrf 'cv'qhuj qtg'dcpm'lp" y g'o kf/3; 72u'y cu'cr r tqzko cvgn'842'o o 'HN'Dctfcej 'gv'crf'3; 7: +y kj 'eqpukf gtcn' 'uo cmgt" kpf kxf wcu'kpj qtg0'Hqmny kpi 'y g'eqmcr ug'qh'y g'ci i tgi cvkqpu'lp'3; : 3."qpn' 'lwxgpkp'P cuucw' i tqwr gt'y gtg'uggp.'dw'qpn' 'tctgn' 'kpj qtg'LOY ctf . 'Dgto wf c'F gr ctvo gpv'qh'Ci tlewwtg." Hkuj gtkgu."cpf 'Rctmu.'F kxkukq'qh'Hkuj gtkgu.'r gtu0eqo o 0'vq' l 0Ucf qx { .P O HU.'3; ; 4+0"Vj gtg" j cxg'dggp'cpgef qv'n'ceeqwpw'qh'ecvej gu'qh'P cuucw'i tqwr gt.'kpxqk'kpi 'i qqf /uk' gf 'hkuj . 'j qy gxgt." ukpeg'P cuucw'i tqwr gtu'ctg'r tqvevgf 'hkuj gto gp'ctg'tgnwcvp'vq'tgr qt'v'ecvej kpi 'qt'r quugukpi 'y go 0" Hkuj 'ctg'qh'ngp'hkngvf 'vq'cxqkf 'f gvevqk'vq'y g'gz wgpv'qh'cp { 'r gtegxgf 'kpetgcug'ku'wmpqy p'D0' Nwenj wtuv.'Dgto wf c'F gr ctvo gpv'qh'Ci tlewwtg.'Hkuj gtkgu."cpf 'Rctmu.'F kxkukq'qh'Hkuj gtkgu.'r gtu0 eqo o 0'vq' l 0Ucf qx { .Wpkxgtuk{ 'qh'J qpi 'Mqpi . 'Ugr 04234+0" "

## Bermuda – Conservation and Management

Vj g'gctn'ku'v'hkuj gtkgu'o cpci go gpv'o gcuwtg'vq'eqpugt'xg'ur cy plpi 'ci i tgi cvkqpu'lp" Dgto wf c'qeewtgf 'lp'3; 960"Vj ku'kpxqk'gf 'y g'ugcuqpcn'eqmwtg'6"o qpvy u+qh'vy q'tgf 'j kpf " ci i tgi cvkqp'ukgu0"Vj ku'o cpci go gpv'cevqk'y cu'ecmgf 'hqt'd { 'eqo o gtekn'hkuj gto gp'cpf 'y g' tgi wcvkqp'y cu'gpcev' d { 'y g'Hkuj gtkgu'F gr ctvo gp0"Vj g'ugcuqpcn'eqmwtg'qh'y g'tgf 'j kpf " ci i tgi cvkqp'ukgu'ku'wkn'lp'ghge'v'53" { gctu'rcvgt'cmj qwi j 'y gtg'j cxg'dggp'uqo g'o qf k'kecvkqpu'qh" dqwpf ctkgu'cpf 'y g'uk' g'qh'y g'r tqvevgf 'ctgcu0"Hqmny kpi 'y ku'o gcuwtg.'ecvej gu'eqpv'kwgf 'vq" f gerkg'dw'y gp'uxcdk'k' gf 'lp'y g'npqi gt'vgo 0"Eqo r r'kepeg'qt'gphqtego gpv'ku'pqv'y gm' f qewo gpvgf 0"P cuucw'i tqwr gt'ci i tgi cvkqpu'ugcy ctf 'qh'y gug'tgf 'j kpf 'ukgu'y gtg'pqv'r tqvevgf " wpf gt'y g'tgi wcvkqpu'cpf 'y gtg'j gcxkn' 'hkuj gf 0"Cu'c'tguwn.'P cuucw'i tqwr gt'rcpf kpi u'f gerkgf " ; 7' 'htqo '3; 97/3; : 3'cpf 'cm'npqy p'ci i tgi cvkqpu'f kuc r getgf 0"Dci 'hko ku'\*4'hkuj '+cpf " o kplko wo 'uk' g'tgut'levkqpu'\*578"o o 'HN'+y gtg'lp'ghge'v'ht'y g'P cuucw'i tqwr gt'r tkqt'vq'3; ; 2" \*Nwenj wtuv'3; ; 2+0'

P cuucw'i tqwr gt'lp'Dgto wf c'j cxg'dggp'o cpci gf 'ukpeg'3; ; 8'y kj 'pq/vcn'cpf 'pq/ r quugukq'tgi wcvkqpu'dw'lp'ur kg'qh'y qug'eqpugt'xv'kqpu'o gcuwtgu.'P cuucw'i tqwr gt'j cu'o cf g'pq" cr r tgekdng'tgeqxgt { 0"Vj g'ur geku'ku'eqo r ngv'n' 'r tqvevgf 'y tqwi j 'r tqj kdkkq'qp'vcn'cpf "

"  
"  
"  
"

r quuguukqp"cpf "r quukdn{ "dgpghku"htqo "pwo gtqwu'pq/vcng"o ctkg'tgugtxgu"\*D0Nwenj wtuv."  
Dgto wf c'F gr ctvo gpv'qh'Ci tlewwwtg."Hkuj gtlgu."cpf "Rctm."F kxkukqp"qh'Hkuj gtlgu."r gtu0eqo o 0\q"  
[ 0Ucf qx{."Wpkxgtuk{ "qh'J qpi "Mqpi ."Ugr v04234-0 "

"  
"  
"  
"

## BRITISH VIRGIN ISLANDS

### British Virgin Islands – Abundance and Distribution

Nkwg'kphqto cvkqp'ku'cxckrdrg'qp'P cuucw'i tqwr gt'kp'yj g'Dtkkuj 'Xkti kp'Kucpf u'DXK' cni qwi j 'cpgef qvri'ceeqpwu'ui i guv'yj cv'eqpukf gtcdrng'rcpf kpi u'ukm'qeewt'cni qwi j 'pqv'htqo " ci i tgi cvkqpu'0'Ewo wrcvkg'f cv'htqo 'TGGH'\*4225/4235+'uj qy 'uki j vki u'qh'329'P cuucw'i tqwr gt' kp'4225'uwtxg{'u'f gpukv{'kpf gz'304.'uki j vki 'htgs wqpe {'76' +cetquu'yj g'32/{gct'r gtkqf " \*j wr <ly y y 0ggf'0ti lf d ltr qt vulf kvlur geku'IVY C 122; 94225/23/23 4235/26/29-0'Tgs wguu'htqo " wrf cvgf 'kphqto cvkqp'yj tqwi j 'yj g'kuj g'kqu'f gr ctwo gpv'htqo 'yj ku'ucwu'tgr qt v'j cxg'tgegkxgf "pq" tgur qpug'0

kp'yj g'o kf/3; ; 2u.'rci g'P cuucw'i tqwr gt'yj g'g'ukm'dgkpi 'ecwi j v'gcu'qh'Relctqu'Rqkpv." Xkti kp'I qtf c.'dw'yj gug'yj g'g'kpekf gpvri'ecvej gu'cpf 'pqv'cti gvgf 'ecvej gu'O wptq'cpf 'Dnm'4227-0" O qtg'tgegpm'. 'kuj gtu'tgr qt v'yj cv'o gf kwo /uk' gf 'P cuucw'i tqwr gt'ctg'ukm's wkg'eqo o qp'dw'yj cv' ci i tgi cvkqpu'ctg'pq'htqo gt'cevkg'nf 'cti gvgf 0'Qpn' 'c'hy 'P cuucw'i tqwr gt'yj g'g'rcpf gf 'cv'yj g' DXKHkuj g'kqu'Ego r ngz'f wtkpi 'yj g'yj k'vgt'o qpyj u'qh'4225'\*O wptq'cpf 'Dnm'4227-0'Dcugf 'qp'yj g' hpf kpi u'qh'c'uwtxg{'eqpf wevgf 'kp'Lcpwet {'v'Hgdtwet {'4225.'O wptq'cpf 'Dnm'\*4227+'htqwpf "pq" gxkf gpeg'qh'cp{'ur cy plpi "ci i tgi cvkqp'htqo "c'r t'gkq'wun' 'tgr qt v'gf 'ukg'qp'yj g'Ucdc'uj g'0'Hkuj gtu' k'vgt'xky gf 'erko gf 'yj cv'yj g{'eqwrf 'ecvej '42/62'P cuucw'i tqwr gtu'r gt'f c{'cv'yj g'ukg'37/42'}{gctu' ci q'0 " "

### British Virgin Islands – Conservation and Management

P cuucw'i tqwr gt'ecp'dg'uggp'htq'ucrg'kp'yj g'DXKHkuj g'kqu'Ego r ngz'cpf 'kp'uw' gto ctngvu' Vj g'g'ku'c'emugf 'ugcuq'htq'rcpf kpi 'P cuucw'i tqwr gt'dgy ggp'O ctej '3'cpf 'O c{'53'\*O wptq'cpf " Dnm'4227-0'

**CAYMAN ISLANDS**

**Cayman Islands – Populations**

Vj g'P cuucw'i tqwr gt'o c { 'ukm'dg'tgrvkg' "cdw'p'cpv'kp'vj g'Ec {o cp'Kuc'p'f'u'eqo r'ctgf "vq" o cp { "qvj gt'mecv'kpu'\*Rcv'p'i km'Ugo o gpu'cpf "Ugo o gpu'4225+'cee'qtf kpi "vq'xku'w'ru'w'xg {u'cpf " vj g'uc'wu'qh'ug'x'g'te'n'ur cy p'kpi "ci i tgi cv'kpu'0"Ewo w'v'k'g'f'c'v' "It'qo "TGGH\*4225/4235+'uj qy " uki j kpi u'qh'3: 79'P cuucw'i tqwr gt'kp'5968'w'v'x'g {u'f'g'p'uk' { "k'p'f'g'z'30. "uki j kpi "It'gs'w'g'p'e { "6; 8' -+ cet'qu'u'vj g'32/ {gct'r'g'tk'f' \*j'w'r <ly y y 0gg'f'q'ti l'f'd'l't'g'r'q't'v'lf'k'v'l'ur'g'ek'gu'V'Y' C'122; 9'4225/23/ 23'4235/26/29+0"kp'vj g'Ec {o cp'Kuc'p'f'u' "vj g'P cuucw'i tqwr gt' "huj gt { 'y'cu'q'peg'eq'p'uk'f'gt'gf "vq'dg" qp'vj g'd't'k'p'm'q'h'eq'm'r'ug'g'x'g'p'vj q'w'i j "huj kpi 'y'cu'o' c'p'c'i'gf'0"Vj g'P cuucw'i tqwr gt' "u'q'em'i'kp'vj g' Ec {o cp'Kuc'p'f'u'er'r'g'ct'vq' "j'c'x'g'uj'q'y'p' "u'q'o'g'f'g'i't'g'g'q'h't'g'uk'k'p'eg'w'p'f'gt' "huj kpi 'r't'g'u'w't'g'f'w'g'vq" vj g'ewo w'v'k'g'g'h'g'ew'q'h'k'p'ergo'g'p'v'y'g'c'v'j'gt' "f'v't'k'p'i 'vj g'ci i tgi cv'k'p'ug'cu'q'pu' \*100' "h'o'k'k'p'i 'huj kpi " q'r'r'q't'w'p'k'k'g'u' "u'q'o'g' " r't'q'v'g'v'k'p' "It'qo " r'q'cej'kpi 'y'k'j' "vj g" t'g'i'w'r'c't'r't'g'ug'p'eg'q'h" t'g'ug'c't'ej'g'tu'cv'vj g'uk'g" f'v't'k'p'i 'vj g'ur'cy'p'k'p'i " u'g'cu'q'p.'r'qu'k'd'ng" t'g'et'w'ko'g'p'v' "It'qo " p'g'c't'd { "q'h'uj'q't'g" d'c'p'm.'c'p'f'c'r'qu'k'd'ng" u'j'k'h'k'p'i'q'h" ci i tgi cv'k'p'uk'g'u'vj'cv' t'g'o'c'k'p'w'p'h'uj'g'f' "q't" w'p'p'q'y'p' \*Y'j'c {ng'p" g'v'c'r'0'4229+0" T'g'ug'c't'ej'g'tu'q'd'ug't'x'g'f" u'j'k'h'k'p'i'q'h'vj'g" ci i tgi cv'g'f' "ur'cy'p'g'tu" qp'vj g'ue'c'ng'q'h'ug'x'g't'e'n' j'w'p'f't'g'f'o'g'v'g'tu' \*Y'j'c {ng'p'g'v'c'r'0'4229+'c'p'f' "vj'g't'g'c't'g' "u'q'o'g't'g'r'q't'u'q'h' "u'ko'k'r'c't' "uj'k'h'u'cv'q'y'gt' "uk'g'u' \*C'i'w'k'r'c't'4228+'v'j'c'v'o'c'ng' "v'j'k'u'c'r'qu'k'd'k'k'v' {0' "o'K'k'u'r'qu'k'd'ng' "vj'g't'g'c't'g' "q'y'gt' "o'k'p'q't' "uc'v'ng'k'g" ci i tgi cv'k'p'uk'g'u'vj'cv't'g'o'c'k'p'w'p'h'uj'g'f' . "d'w'k'k'u'w'p'k'ng'ng' "c'p'f' "q'x'g't' "v'j'g' "u'v'38" {g'c'tu' "e'c'v'ej'g'u'j'c'x'g" u'g'c'f'k' "f'g'ek'p'g'f' "kp'vj g'P cuucw'i tqwr gt' "huj gt { \*R'0'D'w'uj' . "Ec {o cp'Kuc'p'f'u'f'g'r'c't'w'o'g'p'v'q'h" G'p'x'k'q'p'o'g'p'v' "r'g'tu'0'q'd'ug't'x'0'c'u't'g'r'q't'v'g'f' "kp'Y'j'c {ng'p'g'v'c'r'0'4226+0' "Huj kpi "q'p'vj g'uk'g'u'r't'q'f'w'eg'f" vj q'w'uc'p'f'u'q'h' "huj "c'p'p'w'c'm' { "c'p'f' "kp'vj g'3; 92'u'g'x'g'p'k'p'e'n'w'f'g'f' "vj g'uc'ng'q'h' "e'c'v'ej' "vq' "L'co'c'k'ec'p' "x'g'u'g'n'u' \*Y'j'c {ng'p'g'v'c'r'0'4226+0'




Figure 16. Cayman Islands (islands west to east Grand Cayman, Little Cayman and Cayman Brac)

j'w'p'f't'g'f'o'g'v'g'tu' \*Y'j'c {ng'p'g'v'c'r'0'4229+'c'p'f' "vj'g't'g'c't'g' "u'q'o'g't'g'r'q't'u'q'h' "u'ko'k'r'c't' "uj'k'h'u'cv'q'y'gt' "uk'g'u' \*C'i'w'k'r'c't'4228+'v'j'c'v'o'c'ng' "v'j'k'u'c'r'qu'k'd'k'k'v' {0' "o'K'k'u'r'qu'k'd'ng' "vj'g't'g'c't'g' "q'y'gt' "o'k'p'q't' "uc'v'ng'k'g" ci i tgi cv'k'p'uk'g'u'vj'cv't'g'o'c'k'p'w'p'h'uj'g'f' . "d'w'k'k'u'w'p'k'ng'ng' "c'p'f' "q'x'g't' "v'j'g' "u'v'38" {g'c'tu' "e'c'v'ej'g'u'j'c'x'g" u'g'c'f'k' "f'g'ek'p'g'f' "kp'vj g'P cuucw'i tqwr gt' "huj gt { \*R'0'D'w'uj' . "Ec {o cp'Kuc'p'f'u'f'g'r'c't'w'o'g'p'v'q'h" G'p'x'k'q'p'o'g'p'v' "r'g'tu'0'q'd'ug't'x'0'c'u't'g'r'q't'v'g'f' "kp'Y'j'c {ng'p'g'v'c'r'0'4226+0' "Huj kpi "q'p'vj g'uk'g'u'r't'q'f'w'eg'f" vj q'w'uc'p'f'u'q'h' "huj "c'p'p'w'c'm' { "c'p'f' "kp'vj g'3; 92'u'g'x'g'p'k'p'e'n'w'f'g'f' "vj g'uc'ng'q'h' "e'c'v'ej' "vq' "L'co'c'k'ec'p' "x'g'u'g'n'u' \*Y'j'c {ng'p'g'v'c'r'0'4226+0'

"  
"  
"  
"

Vj g'tg'7'tcf k'kqpen'ci i tgi cvkqp'ukgu'eqphkto gf 'kp'yj g'Ec {o cp'K'rcpf u.'qpg'qh'y j kej ."  
 qh'Nkwg'Ec {o cp'v'y g'v'gpf . 'ku'hkn' 'y g'rci g'v'ci i tgi cvkqp' "kp'v'gto u'qh'hkuj "pwo dgtu+'npqy p"  
 kp'tgegpv'ko gu'cp {y j g'tg'y kj kp'yj g'i gqi tcr j le'tcpi g'qh'yj g'ur gekgu0"Vj g'Nkwg'Ec {o cp'ukg'ku"  
 m'ecv'gf "qp'c'tggh'r tqo qpvt { 'qp'yj g'y g'v'gtp'gf i g'qh'Nkwg'Ec {o cp'K'rcpf "T'cpf "g'v'cr04227+0"  
 Y j c { r'gp'g'v'cr0\*4226+'tgeqtf gf "tqo 'w'p'f gty cvgt "qdugtxcv'kpu+'y cv'yj g'cxgtci g'g'v'ko cvgf "  
 pwo dgt'qh'P cuucw'i tqwr gt'r t'gugpv'cv'yj g'ci i tgi cvkqp'ukg'qh'Nkwg'Ec {o cp'kp'4224'y cu'7.422"  
 kp'f k'kf w'cu'y q'f c { u'chgt 'hwm'o qqp0"Vj g'o g'ep'uk' g'qh'ci i tgi cvkpi "i tqwr gt'y cu'842"o o "VN'cpf "  
 yj g'q'xgtcm'hgo c'ng'v'q'o c'ng'ugz'tcv'k'y cu'3-3(80"Y j c { r'gp'g'v'cr0\*4226+'tgr qt'v'yj cv'hgo c'ng'g'zj kdk'  
 f'ctnr j cug'cpf "o c'ng'g'zj kdk'dleq'rt'r j cug'cv'yj g'r q'kp'v'qh'i co g'v'g'tgr'cug'cnj qwi j "kp'yj g'ngcf/wr "  
 v'ur cy pl'pi "dqy "ugz'gu'o ki j v'f'kur m' { "dqy "eq'rtu'cv'qy gt'v'ko gu'"Ctej gt'g'v'cr04234+0"C "  
 j { f'tq'ceq'w'ke'u'w'f { "qh'yj g'ci i tgi cvkqp'uw'i i g'v'g'f "y g'r t'gug'peg'qh'o qt'g'hkuj "y cp'eq'w'p'v'g'f "d { "  
 f'kxgtu'f w'g'v'q'yj g'h'cv'yj cv'yj g'ci i tgi cvkqp'cr'r g'ctgf "v'dg'ur t'gcf "r'cvej k'f "q'xgt'c'y kf gt'ctgc'yj cp"  
 yj cv'eq'xgtgf "d { "f'kxgtu'qp'yj g'q'yj gt'j cpf . 'hkuj "em'q'ug'v'q'yj g'u'w'v'g'v'yj g'tg'p'q'v'g'f "d { "f'kxgtu'dw'p'q'v"  
 j { f'tq'ceq'w'ke'cm { =c'eqo d'kp'cv'k'p'qh'f'kxgtu'cp'f "j { f'tq'ceq'w'ke'u'ku'uw'i i g'v'g'f "h'q't'uwej "uw'f'kgu"  
 \*Vc { r'qt'g'v'cr04228+0'

Kp'yj g'Ec {o cp'K'rcpf u.'cm'ur cy pl'pi "ci i tgi cvkqp'ukgu'ctg'm'ecv'gf "y kj kp'72"o "qh'yj g'uj g'h"  
 gf i g'\*52"qt'62"o "f'gr yj '+'cpf "cf l'cegpv'v'q'f ggr "y cvgt "@422"o +0"J gr r gm'g'v'cr0\*422: +r'q'q'ugf "  
 yj cv'ur cy pl'pi "o ki j v'dg'v'ko gf "v'cm'qy "r'tx'cg'v'q't'g'w'p'qp'm'ec'n'i { t'gu'v'q'Ec {o cp'K'rcpf "y cvgtu"  
 uw'i i g'v'k'pi "y cv'yj g'eq'p'f k'k'qp'qh'm'ec'n'r q'r w'v'k'p'u'o c { "dg'et'k'ke'cn'v'q'yj g'k't'm'pi /v'gto "uw'v'k'p'cd'k'v'0"  
 M'q'd'ctc'\*422; +t'g'x'g'c'ng'f "y cv'cm'7'd'gu'v'npqy p'Ec {o cp'K'rcpf u'ur cy pl'pi "ci i tgi cvkqp'ukgu'ctg"  
 m'ecv'gf "cv'eq'p'x'g'z/uj cr gf "ug'cy c'tf "g'z'v'g'p'f'k'pi "t'g'ghu't'g'gh'r tqo qp'v'q't'kgu+l'w'w'k'pi "kp'v'q'f ggr "y cvgt."  
 y kj kp'3"n'o "qh't'g'gh'r tqo qp'v'q't { "v'kr u'0'

**Cayman Islands – Fishing**

Vj g'Ec {o cp'K'rcpf u'q'peg'j cf "c'uo cm'h'q'ec'n'tcf k'k'q'p'cn'hkuj gt { 'h'q't'P cuucw'i tqwr gt'y kj "  
 ; 2' "qt'o qt'g'qh'yj g'rc'p'f'k'pi u'eqo k'pi "t'qo "y j g'7"yj gp'npqy p'c'pp'w'cn'ur cy pl'pi "ci i tgi cvk'p'u"  
 \*Y j c { r'gp'g'v'cr04226d+0"Vj g'tcf k'k'q'p'cn'hkuj k'pi "ew'w'w'g'g'x'q'k'g'f "kp'v'q'q'p'g'ge'q'p'q'o k'ecm { "f'gr g'p'f g'p'v"  
 qp'o c't'k'p'g'v'q'w't'kuo "cp'f "h'k'p'c'p'eg'q'x'g't'yj g'r cu'v'52" { g'ct'u' \*D'wuj "g'v'cr04228+0"V'w'eng't "g'v'cr0\*3; ; 5+"  
 t'gr qt'v'g'f "h'k'x'g'P cuucw'i tqwr gt'ur cy pl'pi "ci i tgi cvkqp'ukgu'j k'v'q't'k'ecm { "kp'yj g'eq'w'p't { < "q'p'g'cv'yj g"  
 u'q'w'j g'cu'v'eq't'p'g'tu'qh'g'cej "qh'yj g'yj t'g'g'k'ur'c'p'f u.'q'p'g'cv'yj g'u'q'w'j y g'v'g't'p'eq't'p'g't'qh'I t'c'p'f "Ec {o cp."  
 cp'f "cp'q'yj gt'cv'yj g'u'q'w'j g'cu'v'eq't'p'g't'qh'yj g'Vy g'x'k'g'O k'g'D'c'p'm'u'y g'v'q'h'I t'c'p'f "Ec {o cp0"Vj g"  
 ci i tgi cvk'p'u'cv'yj g'g'cu'v'g't'p'g'p'f u'q'h'yj g'k'ur'c'p'f u'y g't'g'yj g'o q'u'v'y g'm'npqy p.'cp'f "t'cf k'k'q'p'c'm { "  
 g'z'r m'k'g'f "uk'p'eg'yj g'g'ct'n { "3; 22u'y kj "y j g'w'ug'qh'uo cm'l'q'r g'p'd'q'c'w'cp'f "j cp'f "h'k'p'g'u' \*D'wuj "g'v'cr0'  
 4228+0" M'0'R'0'V'k'd'd'g'u'q'h'Ec {o cp'D't'ce' \*r' g'tu'0'eqo o 0'kp'E'q'r'k'p'3; ; 9+t'gr qt'v'g'f "j c'x'k'pi "hkuj gf "y j g'g"  
 ci i tgi cvk'pi "m'ec'v'k'p'u'uk'p'eg'3; 47/3; 48."cp'f "j k'u'h'v'yj gt'j cf "hkuj gf "y j go "uk'p'eg'cd'q'w'3; 250'

Kp'4223. 'hkuj gto gp'h'q'w'p'f "ci i tgi cv'gf "P cuucw'i tqwr gt'qp'yj g'y g'v'g'p'f "qh'Nkwg'Ec {o cp'  
 K'rcpf \*Y j c { r'gp'g'v'cr04226. 'D'wuj "g'v'cr04228+; "cnj qwi j "dc'ug'f "qp'o qt'g't'gegp'v'f k'ue'w'uk'p'u'y kj "  
 g'f g'tu'kp'yj g'hkuj k'pi "eqo o w'p'k'f. "k'c'r r g'ct'u'yj cv'yj g'y g'v'g'p'f "ur cy pl'pi "uk'g'y cu'hkuj gf "g'ct'r'k'gt "kp'

"  
 "  
 "  
 "



qh'P cuucw'i tqwr gt'vcngp'htqo 'y g'hkuj gt {'r tqo r vgf 'y g'ko r ngo gpvcvqp'qh'c'o qpkqtkpi 'r tqi tco " d{'y g'F gr ctwo gpv'qh'y g'Gpxktqpo gpv'Dwuj 'gv'cr04228+0'

Kp'y g'3; ; 2u.'ugxgtcrlo cpci go gpv'o gcuwtgu'y gtg'v'kgf 0'Kp'3; ; 7.'cp'ocngt'pcw'l gct" Hkuj kpi o'twcvgi {'y cu'tgeqo o gpf gf 'dw'y cu'pqv'ko r ngo gpv'gf 'f wg'v'q'rcem'qh'r qrk'kecn'uw'r qt v' \*Dwuj 'gv'cr04228+0'Kp'3; ; . 'y g'y tgg'o ckp'ur cy plpi 'ctgcu'cv'y g'gcuwtp'gpf u'qh'y g'kurcpf u'y gtg" hqto cm{'f guki pcvgf 'cu'ot'gultevgf 'O ctkpg'ctgcu'o'htq' y j lej 'ceegu't'gs wktgf 'r'egpukpi "d{'y g' O ctkpg'E'qpu'gt'xcv'kqp'Dqctf '\*y g'ucwwqt {'cwj qtkv{'t gur qpukdrg'htq'y g'cf o kpkurcv'kqp'qh'y g' O ctkpg'E'qpu'gt'xcv'kqp'Ncy +\*Dwuj 'gv'cr04228+0'Kp'y g'3; ; 2u.'rgi kurcv'kqp'r tqj kdkgf 'ur gct'hkuj kpi " cv'ur cy plpi 'ci i tgi cvkqp'ukgu'0'Kp'H'gdwtct {'4224.'r tqv'ev'xg'rgi kurcv'kqp'f ghkpgf 'c'ur cy plpi " ugcuqp'cu'P qxgo dgt'3'v'q'O ctej '53.'cpf 'y g'ocngt'pcw'l gct'Hkuj kpi o'twv'g'y cu'r cuugf 0'Vj ku'rcy " cmqy gf 'hkuj kpi 'g'xgt {'qy gt'{'gct'y kj 'y g'htuv'pqp/hkuj kpi "{'gct'uvct'v'ki 'y kj '4225.'cpf 'cnuq'ugv'c" ecvej 'rko k'v'qh'34'P cuucw'i tqwr gt'r gt'dqcv'r gt'f c{'f wtkpi 'hkuj kpi "{'gctu'0'Vj g'rcy 'f ghkpgf 'y g'qpg" pcw'kecn'o kng'\*po +\*opq'v'cr r kpi o' qpgu'ctqwpf 'gcej 'ur cy plpi 'ukg.'cpf 'ugv'c'o k'ko wo 'uk' g'rko k'v' qh'34'k'ej gu'ht'P cuucw'i tqwr gt'kp'4224'kp't gur qpug'v'q'lwxgpk'gu'dgkpi 'vcngp'd{'hkuj 'v'cr u'k'pukf g' y g'uqwpf u'\*Y j c{'ngp'gv'cr04226.'Dwuj 'gv'cr04228+0'Kp'4225.'ur gcti wpu'y gtg't'gultevgf 'htqo 'wug" y kj kp'3'pcw'kecn'o kng'qh'cp{'f guki pcvgf 'i tqwr gt'ur cy plpi 'ctgc'\*F I UC+'htqo 'P qxgo dgt" y j tqw j 'O ctej 0"

Gh'ge'v'xg'F gego dgt'4; .4225.'hkuj kpi 'y cu'emugf 'cv'cm'f guki pcvgf 'P cuucw'i tqwr gt" ur cy plpi 'ukgu'htq'c'r g'kqf 'qh': "{'gctu'0'Kp'cf qr v'ki 'y ku'f gekukap.'y g'O ctkpg'E'qpu'gt'xcv'kqp'Dqctf " pqv'gf 'y cv'y q'qh'y g'ukz'ctgcu'y gtg'ohkuj gf 'qww'cpf 'y tgg'kp'ugt'k'wu'f ger'kpg'0'"C'eeqtf kpi 'v'q" t'gugctej 't'guw'u'htqo 'uwxg'u'qp'y g'Nkw'g'Ec{o cp'y guv'gpf 'ur cy plpi 'ukg.'y g'pwo dgt'qh" ur cy pgtu'k'pet'gcu'f 'htqo 'cr r tqz'ko cvgn{'4.722'hkuj 'v'q'6.222'hkuj 'qxgt'y g'gki j v'{'gct'r tqv'ev'kqp" r g'kqf '\*Ugo o gpu'gv'cr04229c+0'Vj g'e'qpu'gt'xcv'kqp'o gcuwt'g'y cu't'gpgy gf 'htq'c'hw'v'j gt': "{'gctu'kp" 4233'cpf . 'kpf g'gf . 'pwo dgtu'qh'hkuj 'ctg'uj qy kpi 'r tqo k'kpi 'uki pu'qh'k'pet'gcu'g'kp'cv'rgcu'v'qpg" ci i tgi cvkqp'ukg'\*F gr ctwo gpv'qh'Gpxktqpo gpv'4233.'J gr r gm'gv'cr04234+0'Kp'422: . 'k'y cu" r tqj kdkgf 'v'q'vcng'cp{'P cuucw'i tqwr gt'd{'ur gcti w'p'cp{'y j gtg'kp'Ec{o cp'y cvgtu'y kj 'pq'v'cr r kpi " y kj kp'3'po 'qh'c'r tqv'ev'gf 'ci i tgi cvkqp'f wtkpi 'y g'ur cy plpi 'ugcuqp'\*P qx03'6'O ct053+0'Ugcup'cn' cpf 'ur cv'kecn'o gcuwt'gu'ucv'g'y cv'pq'P cuucw'i tqwr gt'ku'v'q'dg'vcngp'htqo 'cp{'F I UC'htqo 'P qxgo dgt" v'q'O ctej 'wp'v'ki'423; 0'V'q'v'cn'ctgc'qh'y g'ewt'gpv': 'F I UC'au'ku'39078'no 40'H'qo 'y g't'guw'u'qh'c" o ctm't'gecr w'g'u'wf {'qp'Ec{o cp'Dtce.'Ec{o cp'Kur'cpf 'hkuj gto gp'cr r gct'v'q'ecvej 'u'w'h'k'ek'gpv'cf w'u' i tqwr gt'q'wukf g'y g'ur cy plpi 'ugcuqp'v'q'ugt'k'wun{'ko r cv'r qr w'cv'k'pu'\*Ugo o gpu'gv'cr04234+0'

Vj g'kpf k'ec'v'k'pu'qh't'ge'q'xgt {'\*cu'f g'v'gto k'pgf 'd{'k'pet'gcu'f 'cdw'pf c'peg'qh'hkuj +'ctg" g'peq'wt'ci kpi 'kp'Nkw'g'Ec{o cp'cpf 'qp'Ec{o cp'Dtce=} qy g'xgt.'y g't'g'j cu'd'ggp'pq't'ge'gp'v'uw'xg{'qh' y g'ur cy plpi 'ci i tgi cvkqp'0'Vj gtg'ctg'hgy 'i tqwr gt'cv'I t'cpf 'Ec{o cp.'j qy g'xgt.'cpf 'y g'j k'j " hkuj kpi 'r t'guw't'g'uw't'q'wpf kpi 'y g'uo cm'pq'vcng'ctgc'ci i tgi cvkqp'ukg.'cu'y gm'cu'r q'cej kpi . 'cr r gct" v'q'ng'gr 'y g'r qr w'cv'k'qp'f gr t'gu'gf '\*Ugo o gpu'gv'cr04234+0'Vj gtg'ku'pq'g'x'k'f g'peg'r'ct'x'cg'htqo 'y g' Ec{o cp'Kur'cpf u'eq'v'k'd'w'g'v'q'y j gt'k'pf k'k'f w'cn'0'

"

"  
"  
"  
"

**COLOMBIA**

"

**Colombia – Populations**

Vj gtg'ku'ikw'f'c'c'cxckrd'q'p'yj g'ucwu'qh'P cuucw'i tqwr gt'kp'Eqmo dlc0"

Ewo wrc'k'g'f'c'c'ht'qo "TGGH"4225/4235+'tgr qt'v'33'P cuucw'i tqwr gt'kp'623'uwtxg{u' \*f'gpukv{ 'kpf gz '3.'uki j v'kpi 'ht'gs w'p'e{ '40' +cet'quu'v'j g'32/{gct'r'gt'k'q'f'ht'qo 'v'j g'r'qr wrc'v'g'f' " k'ur'c'p'f'u'q'h'v'j g'Ucp'C'p'f't'2'u'C'tej k'r'g'nci q'Ucp'C'p'f't'2'u'k'ur'c'p'f'. 'Rt'q'x'k'f'g'p'ekc.'c'p'f'Ucp'w'c' " Ec'v'c'k'p'c' \*j wr <1y y y 0ggf'q'ti lf d lt gr qt wulf kuv'ur geku'IVY C 122; 914225/23/2314235/26/ 29+0"kp'c't'gr qt'v'd{ 'Rt'c'f'c'g'v'c'f'0\*4226+'c't'v'k'uc'p'c'k'ur'g'k'uj g'to gp'k'p'f'k'ec'v'g'f'v'j c'v'k'p'v'j g'Ucp' " C'p'f't'2'u.'Rt'q'x'k'f'g'p'ekc.'c'p'f'Ucp'w'c'Ec'v'c'k'p'c'c'tej k'r'g'nci q'Q'f' "Rt'q'x'k'f'g'p'eg+'q'p'v'j g'p'q't'v'j g'c'u'v' " c'p'f' "u'q'w'j 'd'c'p'm.'i'q'ec'v'c'k'p'c'g'q'r'g'q'p'eg'k'uj g'f' "P cuucw'i tqwr gt'f'w'k'p'i 'ur'c'y'p'k'p'i 'c'i i t'g'i c'v'k'p'u' " ht'qo 'c'r'r't'q'z'k'o c'v'g'n{ 'h'k'x'g'f'k'ht'g'p'v'uk'v'g'u'0Q'ee'c'uk'q'p'c'm{.'c'h'g'y' "P cuucw'i tqwr gt'c't'g'uk'm' " ec'w'i j v.'d'w'r'c'u'v'c'd'w'p'f'c'p'eg'u'j c'f' "p'q'v'd'g'g'p' "u'g'g'p'k'p'c'f'g'ec'f'g' "Rt'c'f'c'g'v'c'f'0\*4226+0"kp'v'j g' " u'w'f' {. 'v'g'p'uk'v'g'u'y' g't'g'k'f'g'p'k'k'g'f' "c'u'r'q'v'g'p'v'c'k'ur'c'y'p'k'p'i 'c'i i t'g'i c'v'k'p'uk'v'g'u.'k'p'en'f'k'p'i 'h'k'x'g'ht' " P cuucw'i tqwr gt.'k'uj g'f' "h'q't'o c'p'f' "{gct'u.'c'n'j'q'w'i j 'p'q'y' "q'p'n{ 'c'h'g'y' 'k'p'f'k'k'f'w'c'u'c't'g'g'x'g't' "u'g'g'p' " \*Rt'c'f'c'g'v'c'f'0\*4226+0' "

"

**Colombia – Fishing**

Eqmo dlc't'gr qt'v'g'f' "v'q'HCQ'c'o c'z'k'o w'o "q'h'342'o v'q'h'P cuucw'i tqwr gt'k'p'f'g'f' "k'p'v'j g'g'c't'n{ " 3; ; 2u'U'c'f'q'x' { 'f'g'O'k'ej'g'u'q'p' "4234+0"J'q'y'g'x'g't.'d' { 'v'j g'g'c't'n{ "4222u'v'j g'h'k'uj g't' { 'o'c' { 'j'c'x'g' " eq'm'r'ug'f' "y'k'j' "p'q'k'p'f'k'p'i u't'gr qt'v'g'f' "v'q'v'j g'HCQ'uk'p'eg'0"p'q'k'c't'i g'ur'c'y'p'k'p'i 'c'i i t'g'i c'v'k'p'u'j'c'x'g' " d'g'g'p't'gr qt'v'g'f' "h'q't'v'j k'u'ur'gek'u'ht'qo 'Eqmo dlc0'Eqo o g't'ek'n'h'k'uj k'p'i 'eqo r'c'p'k'u't'gr qt'v'g'f' "P cuucw' " i tqwr gt't'gr t'g'ug'p'v'g'f' "34' "q'h'k'p'i k'p'g'ec'v'ej'g'u'q'h'k'c't'i g'ug't'c'p'k'f'u'k'p'Ucp'C'p'f't'2'u'd'g'y'g'g'p' "4228'/" 4229="c'i i t'g'i c'v'k'p'u'q'h'72'qt' "u'q'P cuucw'i tqwr gt'j'c'x'g'd'g'g'p't'gr qt'v'g'f' \*J (E)OJ q'q'ng't.'W'p'k'x'g't'uk'f'c'f' " P c'ek'q'p'c'n'f'g'Eqmo dlc.'U'g'f'g'E'c't'k'd'g.'r'g't'u'0'eqo o 0'v'q' [ 0U'c'f'q'x' {. 'W'p'k'x'g't'uk'f' "q'h'J'q'p'i 'M'q'p'i . " 4234+0' "

"

**Colombia – Conservation and Management**

"

Kp'v'j g'Ucp'C'p'f't'2'u'C'tej k'r'g'nci q'q'h'Eqmo dlc.'v'j g't'g'c't'g'c'p'w'o d'g't'q'h'c't'g'c'u'v'j c'v'c't'g' " f'g'uk'i p'c'v'g'f' "c'u'p'q'v'c'ng'h'k'uj k'p'i | q'p'g'u="k'p'4222.'v'j g'c'tej k'r'g'nci q'y'c'u'f'g'er'c't'g'f' "d' { 'W'P'G'U'EQ'c'u'v'j g' " U'g'c'h'm'y'g't' "D'k'ur'j'g't'g' "T'g'ug't'x'g'0"kp'4226.'k'c't'i g'r'q't'v'k'p'u'q'h'v'j g'c'tej k'r'g'nci q'y'g't'g'f'g'er'c't'g'f' "c'u'c' " u'f'u'g'o "q'h'o c't'k'p'g'r't'q'v'g'v'g'f' "c't'g'c'u'y'k'j "x'c't' {k'p'i | q'p'g'u'q'h'h'k'uj g't'k'g'u'o c'p'c'i g'o g'p'v'j'q'y'g'x'g't' " g'p'h'q't'ego g'p'v'k'u'k'c't'i g'n{ "k'c'k'p'i 0'T'k'i j'v'q'v'k'uj "k'c'y'u'c'n'q' "t'g's'w'k'g'v'j c'v'h'k'uj g'to g'p.'r'c't'v'k'w'r'c't'n{ "g'r'f'g't' " h'k'uj g'to g'p.'d'g'c'm'y'g'f' "v'q'h'k'uj "c'v'c' "u'w'd'uk'v'g'p'eg' "g'x'g'n'g'x'g'p'y'k'j k'p'v'j g'p'q'v'c'ng' | q'p'g'u' "O 0Rt'c'f'c' " Eq't'c'k'p'c.'Ucp'C'p'f't'g'u.'Eqmo dlc.'r'g't'u'0'eqo o 0T'0J k'n'P O H U."4232+0"p'q'q'v'j g't'g'i w'c'v'k'p'u'eq'w'f' " dg'k'f'g'p'k'k'g'f' "v'j c'v'o k'i j'v'd'g'p'g'h'k'P cuucw'i tqwr gt'y'k'j k'p'Eqmo dlc'p'y'c'v'g't'u'0' " "

"

"

"

"

"

# CUBA

"

## Cuba – Populations

"

" Dkqrqi kecn'uwf lgu"qp"vj g'P cuucw'i tqwr gt"j cxg'pqv'dggp"wpf gtvcngp'lp'tgegpv" { gctu'dw' dlqrqi kecn'cpf 'hkuj gt { 'f gvcku'o c { 'dg'hqwpf 'lp'Ercdq'gv'crf0\*3; ; 2.'422; +0"Ercdq'gv'crf0\*4223+'cpf "



Figure 17. Confirmed Nassau grouper spawning aggregation sites of Cuba.

Ercdq'cpf "Nkpf go cp"4225" f qewo gpvqf "npqy p"ur cy plpi " ci i tgi cvkqp"ukgu"qh'upcr r gt"cpf " i tqwr gt."o quv'qh'vj go "o wnk ur gekgu="lphqto cvkqp'y cu" r tko ctkl' 'hkuj gt { /f gr gpf gpv' tcvj gt"vj cp'ltqo "wpf gty cvgt" uwxg { u0"Vj g'gctrkguv" f qewo gpvcvqp"qh'P cuucw'i tqwr gt" ci i tgi cvkqp'cpf "ugcuqpcn" o ki tvkqp'u'y cu'ltqo "Ewdc'lp"vj g" 3: 22u'lpf kecvpi "c"uwxuqpcn' hkuj gt { 'cv'vj cv'ko g"\*Xkrctq'F kcl " 3: : 6+0"Nkwng'lp'hqto cvkqp"qp"vj g" ewtgpv'ucwu"qh'vj g"ur gekgu'ku" cxckrdng"\*Hcdkcp'Rkpc. "Egptq'f g" kpxgunki cekqpgu'f g'Gequkugo cu"

Eqvgtqu."Ec { q'Ereq. "Ewdc."r gtu0'eqo o 0'vq' [ 0Ucf qx { . "Wpkxgtuky { 'qh'J qpi "Mqpi ."4233+0" Ewo wrcvkg'f cv'ltqo "TGGH"\*4225/4235+'uj qy "uki j vki u'qh'5: "P cuucw'i tqwr gt'lp'342'uwxg { u" \*f gpukv { 'lpf gz'308."uki j vki 'ltgs wgpe { "530' + "cetquu'vj g'32/ { gct'r gtlqf 0"Vj g'dwmi'qh'vj gug" uco r ngu"\*p? 327."55"P cuucw'i tqwr gt'l'f gpukv { 'lpf gz'<308."uki j vki 'ltgs wgpe { <5306+'y gtg'ltqo " vj g'y guv'ukf g'qh'Ewdc"\*j [vr <ly y y 0ggf0ti lf d ltr qt vlf kvhr gekuIVY C 022; 94225/23/ 234235/26/29+0'](http://www.fishbase.org/species/luvarus/022994225/23/234235/26/29+0)

## Cuba – Fishing

"

Vtcr 'hkuj lpi "j cu'dggp"vj g'r tko ct { "o gvj qf "hqt'ecvej lpi "i tqwr gt"\*O wptq'cpf "Vj qo r uqp" 3; : 5+0"Dqcu'ctg'v'r kecm { 'pqp/o gej cpk { gf'cpf 'hguu'vj cp'8'o "hupi "\*Ercdq'gv'crf03; ; 2."Dckutg" 3; : 5+0"Vj g' Cpvmgcp"\*cttqy j gcf +hkuj "tcr u'ctg'y qqf gp/htco gf'y kj "i cmxcpk { gf'y ktg'o guj "cpf " qpg'qt'y q'gpw'cpeg'hwppgnu"\*O wptq'3; : 5c+0"Vj g'ukpi ng'hwppgn'0ej gxtqp'tcr u0'ctg'eqo o qpn { " wugf 'lp'vj g'gcuvgtp'Ectkddgcp."cpf "vj g'0U0'qt'0\ 0'uj cr gf'tcr u.'y kj "f wcn'gpw'cpeg'hwppgnu.'ctg" hqwpf 'lp'Ewdc"cpf "Lco clec0"O quv'tcr u'j cf "o guj "uk { gu'dgwy ggp"47/72"o o "\*O wptq'3; : 5c+0"

"

"

"

"

"

J kuvtkecm{.'y g'P cuucw'i tqwr gt'y cu" co qpi 'y g'o quv'ko r qtvcv'v'kphkuj 'ur gekgu" rcpf gf 'kp'Ewdcp'hkuj gtkgu."{kgrf kpi 'uqo g'qh' y j g'j ki j guv'ecvej gu'hqt'y j g'ur gekgu'cp{y j gtg' y kj kp'ku'i gqi tcr j ke'tcpi g0'I kxgp'y g'xgt { " j ki j 's wcrk{\ 'qh'rcpf kpi u'f cv'hqt'ng{ " eqo o gtekn'ur gekgu'kp'y j g'eqwpt {.'y j lej " gzv'gpf u'ltqo 'y j g'3; 82u'cpf.'hqt'uqo g" ur gekgu'y cu'tgeqtf gf 'o qp'y n{.'y j g'ku'cp" gzegmgp'v'cpf "wpo cvej gf 'tgeqtf "qh'rcpf kpi u' hqt'y j ku'ur gekgu'qxgt'cm quv'7'f gecf gu" \*Erctq'gv'cr04224.'422; +0'Hkuj kpi 'r tguuwt g" qp'y j g'P cuucw'i tqwr gt'kpetgcugf 'pqvcdn{ " chvgt'3; 7; .t'gecej kpi '3.922'o v'kp'3; 85.'chvgt" y j lej 'ko g'rcpf kpi u'f genkpgf "Hki 042+0'Vj g'f gvckrgf 'f cvcugv'ltqo 'y j g'3; 82u'uj qy u'y cv'y g'i tgcv' o clqtkv{\ 'qh'rcpf kpi u'y cu'vncgp'ltqo 'ur cy plpi 'ci i tgi cvkqp'ukgu'cpf 'ko gu.'72' 'qh'y j g'cppwcn' ecvej 'ltqo 'F gego dgt'v'q'Hgdtwct { \*Hki 043+0'O quv'ecvej gu'qh'P cuucw'i tqwr gt'\*57/72' 'qh'y j g' pcvkqpcn'ecr wtg'qh'y j g'ur gekgu'+y j g'j kuvtkecm{\ 'vncgp'kp'y j g'Ctej kr gnci q'Ucdepc/Eco ci Ag{ " \*pqt'y j /egpvt'cn'ctcg+.'cnj qwi j 'vr 'wv'kn'3; 8; "cp'ko r qtvcv'r tqwr gt'v'kp'qh'y j ku'ecvej 'y cu'qdv'kpgf " ltqo 'y j g'Dcj co cu'uj grh'0'C'uqo gy j cv'uwf f gp'eqmcr ug.'uwi i guv'kpi "c'hyperstability eqpf kkp'k'p" y j lej 'eqpegpvt'v'kqpu'qh'hkuj .g0 0'ci i tgi cvkpi 'hqt'ur cy plpi .o cumlc'i gpgtcn'r qr wrcv'kqp'f genkpg+ " qeewttgf 'kp'y j g'rcv'3; 92u.'f gur kg'uqo g'r tqv'ek'kg'o cpci go gp'0'Vj g'f cv'cnu'uj qy 'y j cv." f gur kg'c'i tcf wcn'kpetgcug'kp'hkphkuj 'rcpf kpi u'Hki 042+dgvy ggp'3; 84'cpf '3; ; : \*Erctq'gv'cr0 4223+.'r tqdcdn{\ 'f w'v'q'kpetgcukpi 'hkuj kpi 'ghhqt'v'P cuucw'i tqwr gt'uj qy gf 'c'r tgekr kqwu'f genkpg." utqpi n{\ 'uwi i guv'kpi 'y j cv'k'ku'o qtg'xwpgtcdrg'v'q'hkuj kpi .qt'o qtg'j gcxkn{\ 'cti gv'g'f .y j cp'qy j gt'tgg'h hkuj 'ur gekgu'\*Erctq'gv'cr0422; .Ucf qx { 'f g" O ke j guqp'gv'cr0422: +0'

O quv'rcpf kpi u'qh'P cuucw'i tqwr gt'kp" Ewdc'y j g'g'tgr qtvgf n{\ 'vncgp'd { 'hkuj 'tcr u'cpf ." qh'y j g'42'qt'uj'j kuvtkecm{\ 'tgr qtvgf " ci i tgi cvkqp'ukgu.'pqp'g'j cxg'dggp'eqphkto gf "v'q" ukn'ihqto 'kp'uki p'k'k'ecp'v'pwo dgtu'kp't'gegpv' { gct'u'cnj qwi j 'cdqw'; 'j cxg'dggp'tgr qtvgf 'kp" y j g'o quv't'gegpv'n{\ 'cxck'cdrg'hkuj gt'ceeqwpu'0" Erctq'gv'cr0422; <'o'F w'v'q'f genk'kpi 'hkuj " { kgrf u'qxgt'ko g'cpf 'y j g'tguw'kpi 't'gf w'v'kqp" kp'r tqh'kcdk'k'v{\ 'qh'hkuj kpi "qp'ci i tgi cvkqpu." hkuj kpi 'ghhqt'v'qp'y j g'ur cy plpi 'ci i tgi cvkqpu" f genkpgf 0"Vj g'r gcm'ecvej gu'pqv'g'f "chvgt'3; : 2




Figure 190. Fishery landings in Cuba (1962-1998)




Figure 18. Seasonal landings from Cuba, noting decrease of catches of Nassau grouper during spawning season

"  
"  
"  
"

qeewtqf "o cknf "f wtkpi "ur cy plpi "o ki tcvkpu'y j gp"vj g'xwpgtcdkklv "qh'hkuj gu'vq'hkuj kpi " i gctu'uwej "cu'ugv'pgvu'y cu"j ki j 0"P gxgtvj gnguu."vj gtg'r gtukvqf "cp"ko r qtwcpv'tgetgcvkqpcn' hkuj gt { ".wukpi "dqj "j qqm'cpf "hkg"cpf "ur gct/i wp."qp"vj g'ur cy plpi "ci i tgi cvkqp"ukgu'lp"vj g" pqtj gtp"Ewdcp"ktej kr gnci q0"Vj g'ukl g'qh'vj ku'hkuj gt { "ku'wmpqy p"f wg"vq"rcem'qh'ucvkukecn' kphqto cvkqp"cpf "f kxgtu"j cxg'pqv'uwtxg{ gf "ur cy plpi "ci i tgi cvkqp"ukgu'vq"cuuguu"vj g'pwo dgtu" qh'hkuj "cuugo drkpi "vq"ur cy p0"Vj g'o ckn"ci i tgi cvkqp"ukgu'lp"uqwj gtp"Ewdc"\*Rwpcn>p"f g'E0' I wcpq"cpf "Dpeq"f g'Lci wc+"ctg"pq"mpj gt"tgi wrctn' "hkuj gf "f wg"vq"vj g"f khkeww"ceeguikdkklv " qh'vj gug'ukgu0Qxgtcm"t gncvkgnl "hgy "xkcdng"ur cy plpi "ci i tgi cvkpu"ctg"vj qwi j v'vq"r gtukv'lp" Ewdc"vqf c { 0""

## Cuba – Conservation and Management

Ewdc"j cu"l'mpi "cpf "y gm'f qewo gpvqf "j kvqt { "qh'gzr nkcvkqp"cpf "o cpci go gpv'qh'vj g" P cuucw'i tqwr gt."y j kej "y cu'qpeg"cp"ko r qtwcpv'eqo o gtekn'ur geku'ncpf gf "lp"vj g'eqwpt { 0"Ewdcp" hrgvu'cnuq'hkuj gf "gzvpuvkgnl "hqt"vj g'ur geku'qwu'f g'qh'Ewdcp"y cvgtu."r ctvkwrcn' "lp"vj g" Dcj co cu"\*Erctq"gv'cn0422; +0"Vj g'hkuj gt { "y cu'rci gn' "dcugf "qp"ecvej gu'vcngp"f wtkpi "vj g" ur cy plpi "ci i tgi cvkqp"ugcuqp"\*Hki 043+0"lp"vj g'3; 92u."ci i tgi cvkqp"ecvej gu'uwf f gpn' "f tqr r gf ." kpf kecvpi "c"ugxgtg'tgf wevq'lp"vj g'hkuj gt { "y j kej "y cu'pqv'cwkdwcdng"vq"ej cpi g'lp"ghhqtv'qt"qvj gt" hcevtu"cu'ht"cu'eqwf "dg"v'vgt o kpgf "Erctq"gv'cn0422; +0"F cv"qp"ewtgpv'ucwu'qh'vj g'hkuj gt { "ctg" wpcxckcdng0"Vj gtg"ctg'tgr qtvqf "vq"r gtukv'c'r quikdng"; "qww'qh'4243'r tgxkqwn' "hpqy p" ci i tgi cvkqp"ukgu'cnj qwi j "vj gug"j cxg'pqv'dggp"xcrkf cvqf "tgegpv' 0"

Uqpeg"vj g'3; : 2u."o cp { "tgi wcvkpu"j cxg"dggp"kvqf wegf "vq"cf f tguu'r ctvkwrcn'ur geku." kuuwu."uwej "cu'f gerkgu'lp"ecvej gu."qt"tgi kqpu."g0 0ugcuqpcn'ur cy plpi "enqwt gu."i gct"dcpu."hkuj kpi " ghhqtv'eqptqn"gv0"Vj gug"y gtg'qh'gp"kvqf wegf "hqt"uj qtv'r gkqf u'qh'vko g"cpf "d { "r ctvkwrcn" Hkuj kpi "Cuqekcvkpu0"Hqt"P cuucw'i tqwr gt."vj gtg"y cu'cp"cm quv'eqo r rvg"cdugpeg'qh'ur geku/ ur gekh'e'r tqvgevxg'o cpci go gpv."y kj "vj g'gzegr vq"qh'c'o kpk wo "ngi cn'ukl g"\*54eo "VN? 792i +" vj cv'ku'vq"uo cm'ht"vj g'ur geku'dcugf "qp"ukl g'cv'o cwtk{ 0"Qh'uqo g'dgpghk'vq"vj g'P cuucw'i tqwr gt" y gtg'dci "rko ku'ht"tgetgcvkqpcn'hkuj kpi ."tgi wcvkpu'vq"ketgcug'ugrgevxk{ "qh'ugxgtcn'hkuj kpi "i gctu" \*o guj "ukl g+vq"cxqkf "vj g'ecvej "qh'lwxgpkgu."eqptqn'qh'ugv'pgv'wug."cpf "rko ku'f wtkpi "ur cy plpi " ci i tgi cvkqp"vko g."cpf "eqptqn'qh'ur gcti wp'wug."dqj "eqo o gtekm' "cpf "tgetgcvkqpcn' 0"O ctkg" r tqvgeqf "ctgcu"j cxg"dggp"kvqf wegf 0"lp"4224."vj g'vqcn'pwo dgt"qh'tgetgcvkqpcn'kegpugu'y cu" rko kqf "vq"5.722"ht"vj g'y j qrg"eqwpt { "j qr kpi "vq"tgf wegf "f kgeqf "hkuj kpi "r tguuwtg0"Gphqtego gpv" qh'vj gug"tgi wcvkpu"j cu'dggp"xctkqwn' "ghhgevxg"\*Erctq"gv'cn0422; +dwt'geqxt { "qh'vj g'ur geku'ku" pqvtgeqtf gf 0"

"  
"  
"  
"

**DOMINICAN REPUBLIC**

**Dominican Republic - Populations**

"

Vj g'ewttgpv'ucwu'qh'P cuucw'i tqwr gt'ku'rti gn{ 'wvnpqy p'cmj qwi j 'kpf kec'kpu'ctg'yj cv'yj g" ur geku'j cu'dggp'rti gn{ 'f gr ngvf 'htqo 'mecn'tgghu'\*100 cvgq. 'Eqpuglq'F qo kplecpq'f g'Rguec'{" 'Cewkewwmtc.'Gf h0Ugetgvt'f'g'Ci tlewwmtc.'r gtu0eqo o 0\q'T0J km" P O HU."4234+0'T gr qtwa" uwi i guv'yj cv'rti g'hkuj 'ecp'ukm'dg'uggp'kp'yj g'hkuj 'o ctngvu'qp'yj g'pqtvj 'eqcu'v'\*100 cvgq. 'Eqpuglq' F qo kplecpq'f g'Rguec'{" 'Cewkewwmtc.'Gf h0Ugetgvt'f'g'Ci tlewwmtc.'r gtu0eqo o 0\q'T0J km" P O HU."4234+'cmj qwi j 'yj g'mecv'kpu'htqo 'y j lej 'yj qug'ecvej gu'f gtxg'ctg'wvnpqy p'\*dw'ugg" Dcj co cu."cdqxg+0'Ewo wr'v'xg'f'c'v'htqo 'TGGH"\*4225/4235+'uj qy 'uki j v'kpi u'qh'qpn{ '6'P cuucw" i tqwr gt'kp'338'uwtxg{u'f'gpukv{ 'kpf gz'306.'uki j v'kpi 'htgs wpe{ '506' '+cetquu'yj g'32/{ gct'r g'kqf 0" Cm'uki j v'kpi 'kp'yj gug'uco r ngu'\*p?: 6.'6'P cuucw'i tqwr gt'l'f'gpukv{ 'kpf gz'<306.'uki j v'kpi 'htgs wpe{ '< 60 ' '+y gtg'htqo 'O cp{ cpk'q'Dc{ 'v'q'Ecdq'Gpi cpq'qp'yj g'pqtvj 'eqcu'v' \* [vr <ly y y 0ggfQti lf d lt gr qt wlf kvhr gekuIVY C 122; 94225/23/234235/26/29](#)+0'F c'v'htqo " Ucf qx{ '\*3; ; 9+'kpf kec'v'f'qpg'npqy p'ur cy pl'kpi 'ci i tgi cv'k'p'htqo 'Rwp'v'T wuk'cmj qwi j 'ucwu' y cu'kuv'g'f'cv'yj g'v'ko g."cu'or tqdcdn{ 'f'kuc' r gctgf 0""Wp'f gty cvgt'eqtcn'tggh'x'kuwcn'egpuwugu'kp'yj g" F qo kplecp'T gr wdrle'r tqf wegf 'pq'tgeqtf u'qh'P cuucw'i tqwr gt'\*Uej o kv'c'p'f'Uwn'k'c'p'3; ; 6+0" "

**Dominican Republic – Fishing "**

"

Vtcr 'hkuj kpi 'j cu'dggp'yj g'r tko ct{ 'o gyj qf 'hqt'ecvej kpi 'i tqwr gt'kp'yj g'F qo kplecp" Tgr wdrle'\*O wptq'c'p'f'Vj qo r uq'p'3; ; 5+0'P q'rc'p'f'kpi u'j cv'xg'dggp'tgr qt'v'f'htqo 'yj g'F qo kplecp" Tgr wdrle'hqt'o cp{ '{ gct'u'c'p'f'yj g'ur geku'cr r gct'u'v'j'j' cv'xg'dggp'ugxgtgn{ 'f gr ngvf 'kp'mecn'y cvgtu0" Rqcej kpi 'd{ 'F qo kplecp'x'gu'gn'k'p'Dcj co kcp'y cvgtu'hqt'yj ku'ur geku'j cu'dggp'tgr qt'v'f 0" "

**Dominican Republic – Conservation and Management**

"

Nkw'g'k'p'htqo cv'k'p'ku'cx'ck'rdng'f'guet'k'k'p'i 'ur gek'k'le'hkuj kpi 'tgi wr'v'k'p'u'j qy gxgt'k'ku'tgr qt'v'f " yj cv'uk'peg'yj g'o kf/3; ; 2u.'pq'ecvej 'qt'uc'rg'qh't'kr g'hgo c'rgu'k'p'ur cy pl'kpi 'ugcu'q'p'ku'cmjy gf " \*Dqj pucem3; ; ; .Ucf qx{ 'c'p'f'Gm'w'p'f'3; ; ; .Dqz'c'p'f'Dq'p'k'rc'O glk'422: +0'C'v'rg'cu'v'q'p'g" o ct'k'p'g'r ct'mj cu'dggp'gu'cd'rkuj gf 'y kj 'hkuj kpi 'tgi wr'v'k'p'u'cmj qwi j 'pq'k'p'htqo cv'k'p'ku'cx'ck'rdng" qp'P cuucw'i tqwr gt'r t'gug'peg'kp'yj g'r ctn0" "

"

"

"

"

"

"

**HONDURAS**

**Honduras – Populations**

F gur kg'vj g'geqpqo le'ko r qtvcepg'qh'vj g'P cuucw'i tqwr gt'kp'J qpf wcu'vj gtg'ctg'hgy 'f cvc'qp'vj g' ur gekgu'qt'ku'hkuj gt {.'gkj gt'ctvucpcn'qt'eqo o gtekn'0'O wej 'qh'vj g'geqmi kecn'uwf lgu'j cxg" cr r gctgf 'kp'tgr qtwa'vj cv'ctg'pqv'tgcf kn' "cxckrdrg"\*ugg'ekcvkpu'kp'Hqpugec'gv'cn'4226+0" Ewo wrcvkg'f cvc'htqo 'TGGH'\*4225/4235+tr qt v': 2; 'P cuucw'i tqwr gt'kp'5269'uwtxg{u'\*f gpukv' " kpf gz'306.'uki j vki 'htgs wgpe{ '4808' +cetquu'vj g'32/{gct'r gtlqf 00 quv'qh'vj g'uki j vki 'kp'vj gug" uco r ngu'eco g'htqo 'Tqcvp'\*p"? '3: : 6.'7: 7'P cuucw'i tqwr gt/f gpukv' 'kpf gz'<306.'uki j vki " htgs wgpe{ <5308' +cpf 'Wkr '\*p'? '3293.'424'P cuucw'i tqwr gt/f gpukv' 'kpf gz'<304.'uki j vki " htgs wgpe{ <3: 0' +\*j wr <ly y 0ggf0ti lf d ltr qt wlf kuvlr geku'VY C 122; 9 14225/23/23 14235/ 26/29+0'P q'i qxgtpo gpv'wkv'qt'kpukwkwq'eqngew'f cvc'qp'vj g'ur geku'0'Vq'r tqxkf g'cp'qxgxkgy " qh'vj g'ur geku.'c'tgxkgy 'y cu'eqo o kuukppgf '\*Dqz'cpf 'Dqpkrc'O glc'422: +0'Vj g'qpn' 'qvj gt" r wdkuj gf 'uwf lgu'mecv'f 'ctg'vj qug'd { 'Hkg'\*3; ; 2.'3; ; 4+.'y j kej 'f qewo gpv'vj g'tcr kf 'f go kug'qh' qpg'ci i tgi cvkqp'ukg'0"

Vj g'Dqz'cpf 'Dqpkrc'O glc'\*422: +tr qt v'hqwpf 'vj cv'P cuucw'i tqwr gt'rcpf kpi u'kpetgcugf " wr 'wvkv'vj g'gp'qh'vj g'3; : 2u'cpf 'gctn' { '3; ; 2u'cpf 'vj gp'f gerkp'f . 'ku'kpi "eqo o gtekn'ko r qtvcepg" kp'42250'kp'vj g'gctn' { '3; ; 2u.'vj gtg'y cu'gxkf gpeg'qh'wpeqpv'tqngf 'hkuj kpi 'qh'P cuucw'i tqwr gt" ur cy plki 'ci i tgi cvkqp'0'Hq'gzco r ng.'cv'qpg'ukg'emug'vq'I wpcclc.'mecn'cpf 'hqtgki p'xguugu" tgf wegf 'vj g'ci i tgi cvkqp'htqo 'cr r tqzko cvn' { '32.222'hkuj 'v'ngu'vj cp'722'kp'4" { gctu='hkuj gtu" tgo qxgf '35086'v'\*52.222'rdu'0'r gt'ugcuq'p '\*Hkg'3; ; 2.'3; ; 4+0'Qvj gt'ci i tgi cvkqp'r tqdcdn' "




Figure 20. Confirmed and suspected (yellow circle) spawning sites in Honduras

qee'wtgf 'kp'vj g'ctgc'j kvqtkecm' " dw'ukpeg'f gerkp'f . 'cee'qf kpi 'vq" cpgef qvnl'kuj gt'cee'qpw'\*Dqz " cpf 'Dqpkrc'O glc'422: +0'

Hw'vj gt'gxkf gpeg'qh" f gerkp'gu'qh'vj ku'ur geku'ku" tgh'gevgf 'kp'tgf wegf 'gZR qt w'qh" P cuucw'cpf 'tgf 'i tqwr gtu'kp'vj g" ruv'hgy 'f gecf gu'\*Dqz'cpf " Dqpkrc'O glc'422: +0'Rgcml' gZR qt w'qee'wtgf 'f wtkpi 'vj g" P cuucw'i tqwr gt'ur cy plki " ugcuq'p'dw'f gerkp'f 'ugxgtgn' { qxgtcm'dgy ggp'3; ; 7'cpf '42260" Cpgef qvnl'gr qt w'htqo 'hkuj kpi " eqo o wpk'ku'wi i gu'vj cv'vj g" -I tqwr gt'w'Lq { 0'ukg'cpf "

"  
"  
"  
"

o ki tevkqp'tqwgukp'v'ur cy plpi 'ctgcu'j cxg'dggp'kpvpgukxgn{ 'huj gf 'ukpeg'vj g'ncv'3; ; 2u'cpf 'vj cv' vj g'huj 'ku'pqy 'vpeqo o qp0'Vj g'422: 'tgr qtveqpenmf gu'vj cv'vj g'ur gekgu'ku'pqy 'c'o wej 'uo cmgt' r tqr qtvkp'qh'tgg'huj 'vcnpg'kp'vj g'eqwpt {.'tgr tgugpvkpi '>7' 'qh'kpeqo g'htqo 'vj g'huj gt {0" P cuucw'i tqwr gt'f genkpgf 'htqo '9' 'd{ 'y gli j v'qh'gzr qtuv'v'vj g'WUC'kp'3; ; 8'v'20' 'kp'42290" Huj kpi "eqo o wpkkgu'tgr qt'vj cv'P cuucw'i tqwr gt'ctg'dgkpi 'tgr rcegf 'kp'vj g'rcpf kpi u'd{ " *Mycteroperca venenosa.*"{ gmy hkp'i tqwr gt0'Ecvej "qh'P cuucw'i tqwr gt'v'gpf u'v'gd'kpek g'pvcn'v'q' vj cv'qh'upcr r gtu'cpf 'mqduvt' huj g'kgu0"

**Honduras – Fishing "**

Nqecnhuj gto gp'cpf 'eqo o gtekn'dqcu'kp'vj g'Dc{ "Krcpf u'j cxg'gzr mqkpf 'P cuucw' i tqwr gt=Tqcvp.'Nc'Egkdc.'cpf 'I wpcclc'ctg'vj g'o c'kp'eqo o gtekn'huj gt { 'egpvgtu'ht'vj g' eqwpt {.'kpenf kpi 'vj g'rcpf kpi u'qh'P cuucw'i tqwr gt0'Ur cy plpi 'ci i tgi cvkpu'y g'g'huj gf 'y kj " vcr u'cpf 'ur gtu'\*Dqz'cpf 'Dqpknc'O glk'422: +0'O quv'P cuucw'i tqwr gt'rcpf gf 'y g'g'gzr qtvgf 'v'q' vj g'WUC'\*cdqw'; 7' +v'j g'g'j cu'p'g'x'g't'dggp'cp'ko r qt'v'p'v'o ctngv'ht'vj g'ur gekgu'y kj kpi " J qpf wcu'\*Dqz'cpf 'Dqpknc'O glk'422: +0'Vj g'qpg'f qewo g'p'v'g'f 'ur cy plpi 'kp'J qpf wcu." Ecif g'c'f g'ri'F kcdm.'q'wuk'f g'I wpcclc'er r gtu'v'q'j cxg'dggp'gtcf k'ecv'g'f 'kp'vj g'g'ctn{ '3; ; 2u'\*Hkg" 3; ; 2.'3; ; 4+'cnj qwi j 'vj g'g'ctg'pq'ur r qt'v'kpi 'dkqmi k'cnf'c'v'q'p'ku'ewt'gpv'eqpf k'k'q'p0'Huj gtu' j cxg'tgr qtvgf 'o cp{ 'q'v'j g't'q'ecv'k'pu'vj cv'ctg'h'kn'gn{ 'v'q'dg'ur cy plpi 'ukgu.'cnj qwi j 'vj g'k'ewt'gpv' eqpf k'k'q'p'ku'w'p'p'qy p0'K'ku'vj qwi j v'vj cv'q'pn{ 'vj g'o qtg'k'p'ceegu'k'd'ng'ukgu.'uwej 'cu'Dcpeq" Eco r lej g.'ctg'ukn'h'kn'gn{ 'v'q'j cxg'ci i tgi cvkpu'\*Dqz'cpf 'Dqpknc'O glk'422: +0'

Qpg'k'pucpeg'qh'r qcej kpi l'gphqtego g'p'v'y cu'f qewo g'p'v'g'f 'kp'Hgdwtct { '422; 0'Hqwt " J qpf wcp'huj gto gp'htqo 'Rwgt'v'Eq'v'gl 'y g'g'ctt'g'ug'f 'y j k'g'c'v'k'x'gn{ 'pki j v'huj kpi 'kp'Dgrk' g' y cvgtu'cv'vj g'emugf 'P cuucw'i tqwr gt'ukg'kp'I rcf f gp'Ur k'cpf 'Ukmi'Ec { gu'O ct'k'p'g'T'gugtxg" \*I UEO T+0'Vj g'k'ecvej . 'kpenf kpi '3; 'P cuucw'i tqwr gtu'y cu'k'p'x'g'p'v'q't'k'g'f 'cpf 'vj g{ 'y g'g'h'k'p'g'f " cr r tqzko cvgn{ '&3; .4220'Vy q'huj gto gp.'wpcdng'v'q'r c{ 'vj g'k'h'k'p'g'u.'y g'g't'go cpf gf 'v'l'ckn' \*Dgrk' g'Ur cy plpi 'Ci i tgi cvkpu'Y qtnkpi 'I tqwr 'k'p'htqo cvkpu'Ektewct 'P q09.'Lwpg'422; +0' "

**Honduras – Conservation and Management "**

Vj g'g'ku'pq'rgi k'rcv'k'p'vj cv'eqpvt'qnu'huj kpi 'kp'vj g'upcr r gt li tqwr gt 'huj gt { 'kp'vj g'eqwpt { " cnj qwi j 'vcr u'cpf 'ur g'ct'ctg'kn'gi cn'kp'vj g'Dc{ "Krcpf u'0'C'drcenib ctngv'g'x'k'f g'p'v'f { 'eqpvk'p'w'gu" r ct'v'k'ewrctn{ 'kp'vj g'kn'gi cn'uc'ng'qh'huj 'd{ 'mqduvt' huj gto gp.'dw'ku'gz'v'g'p'v'cpf 'ko r cv'ctg' " w'p'p'qy p'\*Dqz'cpf 'Dqpknc'O glk'422: +0'Uqo g'huj 'h'c'x'g'vj g'eqwpt { 'kn'gi cm{ 'q'p'x'gu'gn'cpf " uqo g'ctg'vcnpg'kn'gi cm{ 'q'p'h'qecn'd'q'cu'p'q'v'h'eg'p'ug'f 'v'q'vcn'g'huj '\*Dqz'cpf 'Dqpknc'O glk'422: +0' Eqphk'f g'p'v'cn'k'p'v'g't'x'k'g'y u'k'p'f k'ecv'g'f 'vj cv'f v'k'pi 'vj g'ur cy plpi 'ugcu'p'q'h'vr 'v'q'3.222'rd'0'qh" i tqwr gt'r gt'd'q'cv'y g'g'q'peg'rcpf gf 'ecwukpi 'h'qecn'uc'w't'cv'k'p'v'cpf 't'g'f w'ekpi 'uc'ng'r t'legu'\*Dqz'cpf " Dqpknc'O glk'422: +0' "

"  
"  
"  
"



Vj g'Hkuj gtlgu'F kxkukp'eqmgeu'ecvej 'cpf 'ghqtv'f cve'w'pf gt 'vj g'NTU\*Nlegpukpi 'cpf 'Tgi kurtv'kqp"  
U{uygo +0"Lco clec'cnuq'gpvgtu'vj g'f cve'eqmgevgf 'w'pf gt 'c'ucv'kuecni'uco r r'kpi 'htco g'kpv'vj g'VKR"  
\*Vtkr 'kpvgtx'ky 'Rtqi tco 'f c'vdcug'f g'xgnr gf 'd{ 'EHTCOR\*ECTEQO 'Hkuj gtlgu'T guqwtg"  
Cuuguu gpv'cpf 'O cpci go gpv'Rtqi tco o g+0"C's wgt { 'v'vj g'ECTEQOR'f cve'o cpci gt 'h'kngf 'v"  
wpeq'xgt'cp{ 'f cve'r gt'kpgpv'v'q'P cuucw'i tqwr gt '\*O 0Etgct{ . 'Gpxk'qpo gpv'cn'F cve'O cpci gt."  
Ectkddgcp'E'qcu'cn'F cve'E'gpvt'g. 'E'gpvt'g'ht' 'O ct'kpg'U'el'gpegu. 'W'pkx'gtukv{ 'qh'vj g'Y guv'k'pf k'gu."  
O qpc. 'M'kpi u'vqp. 'Lco clec'Y K'r gtu'0'eqo o 0'v'q'T'0'J km'P O HU. 'F ge04234+0'

Vtcr 'h'kuj kpi 'j cu'dggp'vj g'r tko ct { 'o g'vj qf 'ht'ecvej kpi 'i tqwr gt 'k'P Lco clec '\*O wptq'cpf "  
Vj qo r uqp"3; : 5+0"Vj g'C'p'v'kngcp '\*cttqy j gcf +h'kuj 'v'cr u'ct'g'y q'qf gp/h'co gf 'y k'j 'i c'ix'cpk' gf "  
y k'g'o guj 'cpf 'q'pg'qt'v'y q'g'p't'c'peg'h'wpp'gn'u '\*O wptq"3; : 5c+0"Vj g'u'k'pi ng'h'wpp'gn'i'0'ej g'xt'q'p'v'cr u'o"  
ct'g'eqo o q'pn{ 'w'ugf 'k'p'vj g'g'cu'v'gt'p'E'ct'k'dd'g'cp.'cpf 'vj g'\$'U'\$'qt'\$\ '\$'uj cr gf 'v'cr u.'y k'j 'f w'cn'  
g'p't'c'peg'h'wpp'gn'u.'ct'g'h'q'w'pf 'k'p'E'w'dc'cpf 'Lco clec'0'0 quv'v'cr u'j c'x'g'o guj 'u'k' gu'd'g'y g'gp'47/72"  
o o '\*O wptq"3; : 5c+0'

K'P Lco clec. 'h'kuj kpi 'u'w'x'g{ u'eq'pf w'ev'gf 'k'p'vj g'g'ct'n{ "3; 92u't'gu'w'ngf 'k'p'P cuucw'i tqwr gt "  
ERWG'qh'306'mi 'r gt'h'k'pg'j qw'k'p'42/52'o 'q'h'y cvgt'cpf '30'mi 'r gt'h'k'pg'j qw'k'p'52/'67'o '\*O wptq."  
3; : 5d+0"Y k'j 'vj g'c'f x'gpv'q'h'o q'v'qt'k' gf 'd'q'cw'cpf 'o g'ej c'p'k' gf 'i g'ctu.'k'p'v'g'p'ug'g'z'r m'k'c'v'k'p'ng'f 'v'q"  
m'y gt'ecvej 't'c'v'gu'q'h'c'm'i't'g'gh'h'kuj 'cpf 'vj g'f'k'uc'r r g'ct'c'peg'q'h'u'q'o g'ur g'el'gu'ht'q'o 'o w'nk'ur g'el'gu"  
ecvej gu'\*U'g'x'g'p'u'q'p"3; : 3+0"C'p'w'pf g'ty cvgt'u'w'x'g{ 'q'h'i't'g'gh'h'kuj gu'k'p'Lco clec'k'p"3; : 8't'g'x'g'c'ng'f 'p'q"  
i tqwr gtu'\*M'q'u'ny 'g'v'c'i'0'3; : : +cpf 'd{ "3; : ; 'P cuucw'i tqwr gt'y g't'g't'ct'gn{ 'ecwi j v'\*U'c'f'q'x{ "3; : 9+0'

## Jamaica – Conservation and Management

"

P q'ur g'el'cn'i't'gi w'c'v'k'p'u'g'z'k'v'h'qt'P cuucw'i tqwr gt. 'ur g'el'h'ec'm{ 0'Lco clec'j cu'k'f g'p'v'h'k'gf "  
ct'g'cu'cu'O RCu.'d'w'vj g'f'g'uki p'c'v'k'p'y cu'g'p'c'ev'gf 'q'p'n{ "4" { g'ctu'ci q. 'u'q'p'q'v'c'h'q'v'q'h'ej c'pi gu'ct'g"  
g'z'r g'ev'gf "{ g'v'\*MOC'k'ng'p'W'pkx'gtukv{ 'qh'Y guv'k'pf k'gu.'r gtu'0'eqo o 0'v'q'T'0'J km'P O HU.'4234+0'

"  
"  
"  
"

## LESSER ANTILLES, CENTRAL AMERICA, AND SOUTH AMERICA

### Lesser Antilles, Central America, and South America – Populations

"

P cuucw'i tqrw gt 'ctg'hpqy p 'v'q'qeewt 'qp'v'j g'pqt v'j gtp 'eqcu'qh'Uqwj 'Co gtkec.'dw' ci i tgi cvkqu'j cxg'pgxgt 'dggp'tgeqtf gf 'htqo 'v'j g'eqp'v'k'p'w'cn'luj grh'gxgp'y j gtg'uwduv'p'k'cn' hku'j gt'kgu'j cxg'gz'kuvf 'uwej 'cu'lp'Eqrqo dlc '\*Ucf qx { 'cpf 'Gmwpf '3; ; ; +0'Nkwg'cdwpf cpeg' kphqto cvkqp'eqwf 'dg'hqwpf 'htqo 'Xgpg' wgr. 'cmj qwi j 'v'j g { 'y gtg'tgr qtvgf 'cv'rgcu'htqo 'Nqu' Tqs wgu '\*Egtxki »p'3; ; 6. 'Dqgo j qy gt 'g'v'cn'04232+0'Ewo wr'v'xg'f'c'v'htqo 'TGGH'\*4225/4235+' tgr qt'v'4'P cuucw'i tqrw gt 'lp'54'uw'xg' { u'f'gpukv { 'lpf'gz'3. 'uki j v'kpi 'htgs wpe { '80' ' +cetqu'v'j g' 32/ { gct'r'gtkqf'0' 'C'f'f'k'k'q'p'cn'uw'xg' { u'ct'g'rkuvf 'hqt'Xgpg' wgr '\*p'?' '36: +.'dw'rqecv'k'qu'ctg'pqv' i kxgp'cpf 'P cuucw'i tqrw gt 'y gtg'pqv'tgeqtf gf "

[\\*j wr <ly y y 0ggHQti lf d ltr qt vulf kuvur gekuIVY C 122; 94225/23/234235/26/29+0'''](#)

kp'v'j g'Nguugt' Cp'v'k'ngu. 'P cuucw'i tqrw gt 'y gtg'tgr qtvgf 'lp'4227'v'q'dg'xgt { 'uecteg'lp'U0' Gwuc'v'ku '\*O wptq'cpf 'Drqm'4227+0'Qp'v'j g' Cp'v'ki wc/Dctd'w'c'dcpm'O wptq'cpf 'Drqm'4227+' tgr qtvgf 'c'ur'cy'p'kpi 'ci i tgi cvkqp'ukg'lp'Lcpwct { 'cpf 'Hgdwtct { '4225'cv'Mp'qmi'lp'v'j g'egp't'cn'ctgc' qh'v'j g'uj grh'qh' Cp'v'ki wc/Dctd'w'c'Dcpn'0'Ewo wr'v'xg'f'c'v'htqo 'TGGH'\*4225/4235+'tgr qt'v'345' P cuucw'i tqrw gt 'lp'v'j g'Nggy ctf 'Krcpf u' \*k'0' 'Cpi v'k'nc. 'U0'0 ct'v'p'U0'0 cct'v'p. 'U0' Dct'v'j qm'qo { . 'Udc. 'U0'Gwuc'v'ku. 'U0'M'ku. 'P g'x'ku. 'Cp'v'ki wc. 'I wcf'gr'w'g. 'cpf 'F'qo'k'p'ec+'lp' 3: 37'uw'xg' { u'f'gpukv { 'lpf'gz'30. 'uki j v'kpi 'htgs wpe { '80' ' +cetqu'v'j g'32/ { gct'r'gtkqf'0' Uki j v'kpi u'lp'v'j g'Y'k'p'f'y'ctf 'Krcpf u' \*k'0' 'O ct'v'p'k'w'g. 'U0'N'w'ec. 'U0'X'k'p'eg'p'v. 'Dctd'c'f'qu. 'V'j g' I t'g'p'c'f'k'p'gu. 'I t'g'p'c'f'c+'hqt'v'j g'uc'g'r'gtkqf' \*p'?' '5226. '34'P cuucw'i tqrw gt 'l'f'gpukv { 'lpf'gz'<30. ' uki j v'kpi 'htgs wpe { <20' ' +uwi i guv'v'j cv'P cuucw'i tqrw gt 'ctg'o'wej 'o'qtg'uecteg"

[\\*j wr <ly y y 0ggHQti lf d ltr qt vulf kuvur gekuIVY C 122; 94225/23/234235/26/29+0'J'0'](#)

Qz'g'p'h'q'tf '\*Qz'g'p'h'q'tf. 'Egp'v't'g'hqt' 'T'gu'q'w'teg' 'O'c'p'ci'go'gp'v'cpf 'G'p'x'k'q'p'o'gp'v'cn'U'w'f'k'gu. 'W'p'k'g't'ukv { qh'v'j g'Y'gu'v'k'p'f'k'gu. 'E'cx'g'J'k'm'E'co'r'w'u. 'D'ctd'c'f'qu. 'r'g'tu'0'e'q'o'0'v'q' 'T'0'J'k'm' 'P' 'O' 'H'U. '4234+'u'v'c'v'f' " v'j cv'uj g'j' cu'pq'v'uggp'c' 'P cuucw'i tqrw gt 'lp'52' { gct'u'qh'f'k'k'p'ki 'hqt' 't'g'g'h't'g'ug'c'tej 'lp' 'D'ctd'c'f'qu'0'

kp'v'k'p'k'f'c'f' 'cpf 'V'q'd'c'i'q. 'P cuucw'i tqrw gt 'ctg'eq'p'uk'f'gt'gf 'v'q'dg'm'q'ecm' { 'gz'v'p'ev' \*D'q'w'ej'qp' g'v'cn'0422: +0'E'q'p'v'c'ew'v'q'v'j g'h'ku'j'gt'kgu'f'gr'ct'vo'gp'v'g'r'k'ek'g'f'v'j g't'g'ur'q'p'ug'v'j cv'P cuucw'i tqrw gt 'ctg' s'v'k'g't'ctg' 'cpf 'p'g'x'gt' 'uj'qy 'w' 'lp'v'j g'h'ku'j 'o'ct'ng'v' \*L'0'c'rg'o'w. 'F'gr'ct'vo'gp'v'q'h' 'H'ku'j'gt'kgu. 'r'g'tu'0' eq'o'0'v'q' 'T'0'J'k'm' 'P' 'O' 'H'U. '4234+0'

"

### Lesser Antilles, Central America, and South America – Fishing

"

kp'v'j g'Nguugt' Cp'v'k'ngu. 'r'cti'gt'i' tqrw'gtu'ctg'h'ku'j'gf'y'k'j'j' 'j' 'cpf'rk'p'gu'cpf'y'k'j' 't'cr'u'htqo' '6/: ' o' 'n'q'pi' 'd'q'c'w'gs'w'k'r'gf'y'k'j' ': /'v'q'6: /j'qt'ugr'qy'gt'q'w'd'q'c'tf' 'g'p'i'k'p'gu' \*O'c'j'q'p'3; ; 2= 'd'g'ec'w'ug'v'j'g' 'uj'grh'ku'v'q'p'ett'qy' 'q'h'h'v'j g'Nguugt' Cp'v'k'ng'cp'Krcpf'u'v'j'gt'g'j'cu'd'ggp'p'q'i't'g'cv'p'ggf' 'hqt'r'cti'gt' d'q'c'w'0'I' tqrw'gtu'ctg'u'q'o'g'v'o'gu'ec'w'i'v'q'h'h'v'j'g'f'g'gr'gt' 'u'nr'gu'w'uk'p'i' 'g'r'g'ev't'k'e' 't'g'gn'q't' 'o'ge'j'c'p'k'gf' " y'k'p'ej'gu'hqt'j'c'w'k'p'i' 't'cr'u' \*O'c'j'q'p'3; ; 2+0'N'kw'g'k'p'h'q'to'cvkqp'ku'c'x'c'k'c'd'ng't'gi'ctf'k'p'i' 'q'v'j'gt' "

"

"

"

"

huj g'ktgu'ltqo 'vj g'ctgc'vj cv'cti gv'P cuucw'i tqwr gt0'

**Lesser Antilles, Central America, and South America – Conservation and Management**

kp'c'pki wc/Dctdwf c'vj g'Hkuj g'ktgu'Cev.'P q0B6"qh'3; : 5 cpf 'vj g'Hkuj g'ktgu'T gi wr'v'kpu." P q0B2"qh'3; ; 2.'ctg'vj g'r tko ct { 'hgi kur'v'kxg'dcuku'hqt'hkuj g'ktgu'o cpci go gpv'cpf 'f gxgnqr o gpv'qh'cm' hkuj g'ktgu'kpenw'f kpi 'vj g'\*P cuucw+'i tqwr gt'hkuj gt {0'Vj g'Cev'cpf 'T gi wr'v'kpu'o cmg'r tqxkukqp'hqt<3+" hkuj g'ktgu'o cpci go gpv'grgo gpva.'uwej "cu'hkuj kpi 'h'egpukpi ."gpj cpegf 'hkuj g'ktgu'tgugctej "cpf " gphqtego gpv.'vj g'tgi kum'v'kqp"qh'hkuj kpi 'xguugnu'cpf 'vj g'gucdrkuj o gpv'qh'c'hkuj g'ktgu'cf xluqt { " eqo o k'v'gg'cpf =4+"eqpugt'v'kqp'o gcuw'tgu.'uwej "cu'r tqj kdkkpi 'vj g'wug'qh'egt'v'k'p'hkuj kpi "o gvj qf u" cpf "i gct.'ugv'kpi 'ur gekgu'uk' g'tgum'v'k'p'u."gucdrkuj kpi "emugf "ugcuqpu.'cpf "etg'v'kpi "o ctk'pg" tgugt'xgu0'Y kj 'vj g'cuuk'v'peg'ltqo "HCQ."kp'k'v'gf "kp"4225.'vj g'Hkuj g'ktgu'Cev.'P q044"qh'4228'y cu" r cuugf "cpf "gpcev'gf "\*"J qtuhqtf "422; +'v'q'dgwgt'crki p'h'qecni'tgi wr'v'k'p'u'y kj 'evtt'gpv'k'p'v'gt'p'v'k'p'cni' hkuj g'ktgu'r'cy u'kpenw'f kpi 'vj g'Gwtqr gcp'ugchqf 'r tqxkuk'p'u.'r tko ctk' { "dgp'gh'kpi "g'zr qtu0'K'cnuq'i cxg" vj g'O k'p'k'ngt'lo r tqxgf "o cpci go gpv'ecr cdk'k'k'gu.'uwej "cu'o qxkpi "o quv'hkuj g'ktgu'ltqo "qr gp'ceeguu'v'q" h'egpugf "qt'r gto k'v'gf 'hkuj kpi 0'Y j k'g'P cuucw'i tqwr gt'ku'p'qv'ur gek'k'ecm' { "o cpci gf "qt'r tqv'gev'gf ." emugf "ugcuqpu'y gtg'eqpukf gtgf "kp"422: "hqt'P cuucw'i tqwr gt'cpf "tgf "j k'p'f ."vj g'o qtg'f qo k'p'cpv' ur gekgu'kp'vj g'h'qecni' tqwr gt'hkuj gt { 0'

kp'I wcf gnqr g'cpf 'O ctk'v'k'p's wg.'vj g'tg'ctg'r r'p'u'v'q'r tqv'gev'vj g'ur gekgu'\*HI qwf k'p'." T gi k'p'cni'Cev'k'k'v' { 'Egpv'gt'hqt'Ur gekcm' { 'Rtqv'gev'gf 'Ct'gcu'cpf "Y k'f r'k'g'URCY II CE "o'WP GR.'r gtu0' eqo o 0'v'q' [ 0'Ucf qx { ."W'p'k'xgtukv' { "qh'J qpi 'M'qpi ."4233+"cmj qwi j "p'q'f g'v'k'ku'ctg'cx'ck'rdng'cv'vj ku" v'ko g0'

Qvj g't'qecv'k'p'u'uggo "v'q'j cxg'c'hgy "ur cv'k'ni'enquw'tgu'\*g0' 0'U0'Nwek+'vj cv'y qwf "dgp'gh'k'v' P cuucw'i tqwr gt'dw'v'j g' { 'y g'tg'p'qv'f g'uki pgf "hqt'vj g'ur gekgu'qt'vj g'kt'ci i tgi cv'k'p'u0' "

"  
"  
"  
"

MEXICO

Mexico – Populations

C"qvcn'qh'4: 'ci i tgi cvkqp'ukgu'j cxg'dggp'tgr qtvgf 'lp'O gzleq '\*Ci wkrt'gv'cr0422; +dww." qpnf 'hqw '\*Hi 045+j cxg'dggp'xgt'kkgf '\*Ci wkrt/Rgtgtc'gv'cr0422; +0"Vj g'vy q'dguv'uwf kfg 'ukgu' '\*Ci wkrt/Rgtgtc'4228+lpenwf g'O cj cj wcn'y j lej 'cr r ctgpnf 'pq'iqpi gt'hqto u't'gegpv'ej gemi'htqo " Ft0C0Ci wkrt/Rgtgtc'hqwpf 'pq'hkuj 'ur cy plpi 'lp'4235"]C0Ci wkrt/Rgtgtc.'F gr ctvco gpvq'f g" Dkqni 'p'O ct'kpc.'Hcewncf 'f g'O gf lekpc'Xgvt'kpc'k'{" \ qqvgepk.'Wpkxgtukf cf 'Cw»pqo c'f g" [ wecv'p.'O 2 zleq.'r gtu'eqo o 0'vq'T0J km'P O HU.'42340+.'cpf 'Zecrm'y g'rcti guv'hpqy p" ci i tgi cvkqp'lp'O gzleq0"J kvqtkcmf.'ci i tgi cvkpu'qh'wr 'vq'37.222'hkuj 'hqto gf 'gcej " { gct'cv




Figure 21. Nassau grouper spawning aggregation sites confirmed off Quintana Roo, Mexico.

O cj vj wcn'dw'f wg'vq'kpetgcu'f " hkuj lpi 'r tguwtg'lp'yj g'3; ; 2u" ci i tgi cvkpu'j cxg'pqv'htqo gf 'ukpeg" 3; ; 8"\*Ci wkrt/Rgtgtc'gv'cr0422; +0" F gur kg'eqpugt'cvkqp'eqpegtpu.'yj g" P cuucw'i tqwr gt'tge'kx'gu'rkwr" o cpci go gpv'\*gzegr v'y j gtg'pqvgf " dgm'y +.'y kj 'yj g'rko kfg 'gzegr v'kqp" qh'y g'Zecrm'ukg.'y j lej 'ku'lp'c" pcvkqpcn'r ct'no'lp'cf f k'kqp'vq'yj g'ug" vy q'qecv'kpu.'y q'q'yj gt" ci i tgi cvkqp'ukgu'j cxg'dggp" eqp'ht'o gf 'd { 'f k'kpi '\*P lej gj cdlp" cpf 'Ucp'Lwcp'Ej gpej qo ce-0"Qpg" qecv'kqp'j cf ': 22'i tqwr gtu'cpf 'y cu' hktuv'kf gpv'kkgf 'd { 'hkuj gt'cee'qwp'u'lp' 4227"\*Ci wkrt/Rgtgtc'gv'cr0422; +0"

Cp'cf f k'kqpcn'46'ci i tgi cvkqp'ukgu'j cxg'dggp'tgr qtvgf 'htqo 'hkuj gt'lpv'gt'x'ky u'cn'pi 'yj g'eqcu'v'cpf " qp'Ej k'pej qttq'Dcpm'dw'j cxg'pqv' { gv'dggp'xgt'kkgf 'vq'j cxg'P cuucw'i tqwr gt0"Vj g'ug'cf f k'kqpcn' qecv'kpu'o c { 'dg'cn'pi 'o ki tcvqt { 'tqwg'u't'cyj gt'yj cp'cewcn'ci i tgi cvkqp'ukgu'\*Uqc/Eqt'f gtq'gv'cr0' 4224.'Ci wkrt/Rgtgtc'gv'cr0422; =C0Ci wkrt/Rgtgtc.'F gr ctvco gpvq'f g'Dkqni 'p'O ct'kpc.'Hcewncf " f g'O gf lekpc'Xgvt'kpc'k' {" \ qqvgepk.'Wpkxgtukf cf 'Cw»pqo c'f g' [ wecv'p.'O 2 zleq.'r gtu'eqo o 0' vq'T0J km'P O HU.'4234+0'

F g'v'k'kf 'tgr qt'v'ctg'cx'k'k'rd'ng'ht'vy q'ukgu.'Zecrm'icpf 'O cj cj wcn'yj g'r'w'gt.'j ki j n'f " ceegu'k'd'ng'vq'hkuj lpi 0"Vj g'ci i tgi cvkqp'ukg'cv'Zecrm'y cu'yj g'rcti guv'hpqy p'lp'O gzleq'y kj "6.322" hkuj 'tgr qtvgf 'lp'4226/7.'wr 'htqo '425'hkuj 'lp'4223/4"O gf l'pc/S wgl'gv'cr04226.'Ci wkrt/Rgtgtc" 4228.'Dq'k'q/O qi wgn'4229+0"Wp'f gty cvgt'uw'xg { u'cv'O cj cj wcn'f wtkpi 'yj g'tgr tqf wv'kxg'ugcu'pu'qh' F gego dgt'cpf 'Lcpwct { 'htqo '3; ; 3'vq'3; ; 9.'tgr qtvgf 'i tqwr u'qh'd'gy ggp'72"cpf ": 22'P cuucw' i tqwr gtu'o q'x'kpi 'cn'pi 'yj g'ht'gt'ggh'd'qtf gt'3'mo 'uq'wj 'qh'yj g'v'cf k'kqpcn'ci i tgi cvkqp'ukg0"lp' 9; "

"  
"  
"  
"

F gego dgt'3; ; 5.'37'i tqwr gt'u'y gtg'qdugt'xgf "cv'yj g'uksg.'y j krg'pq'ci i tgi cvkqp'y cu'hqwpf 'f wtkpi " vj g'3; ; 8'cpf'3; ; 9'ugcuqpu'\*Ci vkrct/Rgtgtc'4228+'.uwi i guvki 'vj cv'yj g'ci i tgi cvkqp'j cf "pqv" hqto gf "qt'hqto gf "gnugy j gtg'0"Gz vgpukxg'ugctej gu'qh'yj g'ctgc'd { 'f kxgtu'hckrgf "vq'mqecv'cp { 'P cuucw' i tqwr gt'y kj lp'hkqo gvgtu'qh'yj g'ur cy plpi 'uksg'\*C0Ci vkrct/Rgtgtc.'F gr ctwco gpvq'f g'Dkqmi 'c" O ctkpc.'Hcewncf 'f g'O gf kelpc'Xgvtk'pctk'{" \ qqvgepk.'Wpkxgtukf cf 'Cw»pqo c'f g'l wecv' p." O<sup>2</sup>zleq.'r gtu'eqo o 0'vq'T0J km'P O HU.'4234+0'Ci vkrct/Rgtgtc'\*4228+'.uwi i guvgf 'vj cv'f gerkp'g'cpf " cr r ctgpv'f kucr r gctcepg'qh'lpf kxkf wcu'htqo 'vj g'v'cf kxkpcn'ci i tgi cvkqp'uksg'qh'0 cj cj wcn'y cu'f wg' vq'qxgthkuj kpi "qxgt'vj g'rcuv'72" { gctu0""

Vj g'qvj gt'uwf kcf "ci i tgi cvkqpu'qeeew'cv'õGn'Drcps wkl crõ"qp'vj g'uqwj "eqcu'qh'S wkpvcpc" Tqq'cpf 'Rwpc'I c'krcp'\*O gf kpc/S wgl'gv'cn'04226+0'Hkuj gt'lpvgt'xkgy u'uwi i guvgf 'vj g'r'tgugpeg'qh' ugxg'tcn'gz vcp'ur cy plpi "ci i tgi cvkqpu'qp'vj g'qh'uj qtg'Ej kpej qttq'Dcpm'dw'vj gug'j cxg'pqv'dggp" xcnf cvgf '\*Ci vkrct/Rgtgtc'gv'cn'0422; +0'Vj gtg'ku'rkvw'lpf kcvkqp'qh'qxg'tcm'r qr wrcv'kp'ucwu'qh'yj g' P cuucw'i tqwr gt'lp'O gzleq'dw'eqpegtp'gzkuu'cdqw'vj g'qxgthkuj kpi "qh'cp { 'tgo c'k'kpi 'ur cy plpi " ci i tgi cvkqpu0"

Ewo wrcv'xg'f cv'htqo 'TGGH'\*4225/4235+'tgr qtv'536'P cuucw'i tqwr gt'lp'7; 38'uwtxg { u'lp' vj g'O gzlecp'Ectkddgcp'\*f gpuk { 'lpf gz'304.'uki j vki 'htgs wge { '705' +cetqu'vj g'32/ { gct'r gt'kqf 0' Vj g'rcu' gv'pwo dgt'qh'yj gug'uwtxg { u'y gtg'eqpf wevgf "cv'Kur'Eql wo gn'\*p"? '743: +y kj 'uki j vki u' qh'49; 'P cuucw'i tqwr gt'uo'0? '705' . 'f 00? '304+0'Vj g'eqcu'v'kp'g'kpenf kpi "Xgtcetw' 'cnuq'rkuv'33" P cuucw'i tqwr gt'htqo '847'uwtxg { u'y kj 'uki j vki 'htgs wge { 'qh'304' "

[\\*j wr <ly y 0 ggh0ti lf d lt gr qt vulf kuvur geku IVY C 122; 9 4225/23/23 4235/26/29+0'](#)

"  
**Mexico – Fishing "**  
"

Vj g'P cuucw'i tqwr gt'j cu'mpi "dggp'cp'ko r qtvcv'hqaf "cpf "eqo o gtekn'hkuj 'lp'O gzleq." g'zr m'k'kf "hqt'qxgt'92" { gctu0"K'vj g'O gzlecp'Ectkddgcp.'y j krg'ugeqpf ct { "cu'c'hkuj kpi 'vcti gv'vq" vj g'Ectkddgcp'ur kp { 'm'duvgt '\*Panulirus argus'+cpf 'vj g's wggp'eqpej '\*Strombus gigas'+ 'vj g' P cuucw'i tqwr gt'j cu'dggp'ugcuqpcmf 'ko r qtvcv'cpf 'i gp'g'tcm { 'cngp'cv'ku'ur cy plpi " ci i tgi cvkqpu'lp'F gego dgt.'Lcpwct { . 'cpf 'Hgdwtct { '\*Ci vkrct/Rgtgtc'gv'cn'0422; +0'Vj gtg'j cu'dggp" eqpegtp'd { 'hkuj gtu.'dkqmi kuv.'cpf 'hkuj gtlgu'cwj qtkk'gu'qxgt'f gerkp'gu'lp'ecvej gu'cmj qwi j 'vj gtg' ctg'pq'ur geku/ur gekle'rcpf kpi u'f cv'eqm'gevgf '\*C0Ci vkrct/Rgtgtc.'F gr ctwco gpvq'f g'Dkqmi 'c" O ctkpc.'Hcewncf 'f g'O gf kelpc'Xgvtk'pctk'{" \ qqvgepk.'Wpkxgtukf cf 'Cw»pqo c'f g'l wecv' p." O<sup>2</sup>zleq.'r gtu'eqo o 0'vq'T0J km'P O HU.'4234+0'

Cv'O cj cj wcn'O gzleq.'hkuj gto gp'wugf '5'v'r gu'qh'hkuj kpi 'i gctu'k'0'j qqm'cpf /rkpg." ur gcti wp.'cpf 'i kmpgw'+hqt'g'zr m'k'kpi 'vj g'ci i tgi cvkqp'\*Ci vkrct/Rgtgtc'3; ; 6+0'htqo 'vj g'gctn { " 3; 72u'vq'vj g'3; 92u.'j qqm'cpf /rkpg'y cu'wugf =ur get'i wpu'y gtg'wugf 'lp'vj g'rcv'g'3; 82u'vj tqwi j 'vj g' gctn { 3; ; 2u'0'Vj g'gh'ek'gpe { 'qh'ur gcti wpu'rgf "vq'c'f gerkp'g'lp'cppwcn'rcpf kpi u'\*Ci vkrct/Rgtgtc" 3; ; 6+0'I kmpgw'y gtg'wugf 'htqo '3; ; : 'cpf 'c'htg'ur gcti wpu'y gtg'dc'ppgf '\*3; ; 5+cv'ur cy plpi " ci i tgi cvkqpu.'i kmpgw'\*37/420'eo 'o guj '+wug'k'pet'g'cugf 'cu'dctt'k'g't'pgu'ctqwpf "ci i tgi cvkqp'uk'gu" cpf "dm'enkpi 'o ki cvkqp'tq'w'gu'0'O gcp'uk' g'ht' 'i kmpgw'f 'hkuj 'ecwi j v'cv'y q'ci i tgi cvkqpu'uk'gu'y cu'

"  
"  
"  
"

cdqW'822'o o "VN"Uquc/Eqtfgtq'cpf'Ectf gpcu/Xkf cn3; ; 9+0"K'S wkpvcpc'Tqq.'O gzlecp"  
huj gto gp'ctg'npqy p"vq'ecr wtg'i tqwr gt'd { "v'kpi 'c'kxg'hgo cng'i tqwr gt'v'c'kpg.'r wkpki 'j gt'wr "  
tr kfn'.'cpf'pgwki 'v'g'o cngu'v'v'hmny 'j gt'v'v'g'uwthceg"COci wkrct/Rgtgtc.'F gr ctwco gpvq'f g"  
Dkqmi 'c'O ctkpc.'Hcewncf 'f g'O gf lekpc'Xgvtkptk{" \ qqvgepk.'Wpkxgtukf cf 'Cw»pqo c'f g"  
[ wecv'p.'O<sup>2</sup>zleq.'r gtu'eqo o 0'v'T0J km'P O HU.'4234+0'

Cv'O cj cj wcn'k'S wkpvcpc'Tqq.'huj gto gpau'ceeqwpu\*cu'gctn' 'cu'v'g'3; 72u+'kpf lecvg"  
ecvej gu'qh'wr 'v'46'v'qh'P cuucw'i tqwr gt'r gt'tgr tqf wev'xg'ugcuqp'f k'gew' 'htqo 'v'g'ur cy plki "  
ci i tgi cvkqp'qhi'O cj cj wcn'Vj ku'ecvej 'tgr tgu'g'v'v'qpn' '6'v'7'f c { u'qh'huj kpi 'f wtkpi 'F gego dgt "  
cpf 'Lcpwct { 'wukpi 'qpn' 'j qm/cpf /hpg'i gct '\*Ci wkrct'4228+0'Vj gug'rcpf kpi u'eqpvcuv'uj ctr n' "  
y kj 'f cv'i cvj gtgf 'htqo 'v'g'eqo o gtekn'ecvej '\*wukpi 'i kmpgw'y kj '37/eo 'o guj 'uk' g+'f wtkpi 'v'g "  
tgr tqf wev'xg'ugcuqp'u'gcej 'F gego dgt'cpf 'Lcpwct { 'htqo '3; ; 3/3; ; 90'D { 'v'g'gctn' 'v'q'o k/3; ; 2u "  
rcpf kpi u'htqo 'v'g'O cj cj wcn'ci i tgi cvkqp'k'v'g'o qpv' 'qh'F gego dgt 'j cf 'f tqr r gf 'v'q'5'o v'cpf "  
rcpf kpi u'htqo 'Lcpwct { 'ci i tgi cvkqp'u'f tqr r gf 'v'q'3'o v'\*Ci wkrct/Rgtgtc'4228+0'

## Mexico – Conservation and Management

K'v'g'O gzlecp'Ectkddgpc'Ugc.'v'g'tg'y'gtg'pq'tcf kkpnc'huj gt { 'tgi wrcv'kpu'\*gd 0uk'g."  
s wqcu.'cpf 'huj kpi 'i gct'tgutlev'kpu+'htqo 'huj gt { 'cwj qtk'ku'i qxgtpkpi 'v'g'zr m'kcvkqp'qh'  
P cuucw'i tqwr gt'ci i tgi cvkqp'u'J qy gxgt.'tgi wrcv'kpu'y'gtg'gucdr'kuj gf 'hmqy kpi 'uekpv'khe "  
f qewo gpvcv'kqp'qh'f ger'kpu'cv'O cj cj wcn'\*Ci wkrct/Rgtgtc'3; ; 6+0'Vy q'r tqj kdkk'kpu'v'cv'ch'qtf "  
r tqv'ev'kqp'v'P cuucw'i tqwr gt'y'gtg'gpcev'f <3+'ur gct/huj kpi 'y'cu'dc'ppgf 'cv'cp { 'ur cy plki "  
ci i tgi cvkqp'uk'gu'k'p'u'qwj gtp'S wkpvcpc'Tqq'k'3; ; 5=cpf '4+'rcvt'k'3; ; 9'v'g'huj kpi 'qh'cp { "  
i tqwr gt'ur geku'y'cu'dc'ppgf 'f wtkpi 'F gego dgt'cpf 'Lcpwct { '\*Ci wkrct/Rgtgtc'4228+0'J qy gxgt."  
v'gug'o gcuwt'gu'y'gtg'vgo r qtct { . 'pq'h'qpi gt'k'p'gh'gev.'cpf 'y'gtg'g'x'k'f'gpw' 'pqv't'gur gev'f 'd { "  
huj gto gp'0'Cu'ku'eqo o qp'k'p'u'q'o cp { 'ctgcu.'rcn'qh'gph'qtego gp'v'j'cu'd'ggp'c'r'gtuku'gpv'r'q'drgo "  
\*Ci wkrct/Rgtgtc'gv'cr'0'422; +0'

K'4225.'c'em'ugf'ugcuqp'hqt'cm'i tqwr gt'y'cu'ko r ngo gpv'f'htqo 'Hgd'twct { '37'v'q'O ctej '37"  
cpf 'cr r'k'gu'v'cm'y cvgtu'qh'v'g'O gzlecp'GG\ 'htqo 'Eco r'gej'g'cpf 'l wecv'p'\*I wh'qh'O gzleq+'cpf "  
S wkpvcpc'Tqq'\*Ectkddgpc'+ucv'gu.'cu'y'gm'cu'htqo 'T'k'Ucp'Rgf tq.'dgw'ggp'Ved'cueq'cpf "  
Eco r'gej'g'ucv'gu'v'v'g'Dgr'k'g'dqtf'gt'0'Y'j'k'g'o'cl'pn' 'qh'gt'kpi 'r tqv'ev'kqp'hqt'tgf 'i tqwr gt.'E.  
morio, P cuucw'i tqwr gt'ku'cnu'k'pen'f'gf 'cu'c'r'qj'kdk'f'ur geku'\*Ci wkrct/Rgtgtc'gv'cr'0'422: 0'Vj'ku'  
rcy 'r tqj'kdku'v'g'tgo qxcn'qh'qj'gt'i tqwr gt'ur geku'\*k'pen'f'kpi '{ gm'y'k'p'i tqwr gt.'Mycteroperca  
venenosa+'f wtkpi 'v'g'tgr tqf wev'xg'ugcuqp'0'K'ku'f'k'h'ew'n'v'q'cu'guu'v'g'gh'gewu'qh'v'ku'r tqj'kdk'k'p'  
i kxgp'v'g'cdugpeg'qh'eqp'v'kpw'wu'r'qr wrcv'kqp'o'qpk'qt'kpi 'r'ktq'v'v'g'dcp'0'

D { 'v'g'g'p'f'qh'4234.'c'o'cpci go gpv'r'rcp'y'cu'v'j'cxg'i'qpg'k'p'v'g'gh'gev'k'p'v'g'u'qwj'gtp'I wh'  
qh'O gzleq'cpf'Ectkddgpc'hqt'cm'eqo o gtekm' 'gzr'k'k'gf 'i tqwr gtu'\*cdqW'39'ur geku'='v'g'r'rcp'  
j'cu'pqv'd'ggp'ko r ngo gpv'f.'dw'v'g'tg'ku'zr'gevcv'k'p'v'cv'k'y'kn'd'g'r'w'k'p'v'r'rc'eg'k'p'4236'\*C'O'  
Ci wkrct/Rgtgtc.'F gr ctwco gpvq'f g'Dkqmi 'c'O ctkpc.'Hcewncf 'f g'O gf lekpc'Xgvtkptk{" "  
\ qqvgepk.'Wpkxgtukf cf 'Cw»pqo c'f g'l wecv'p.'O<sup>2</sup>zleq.'r gtu'eqo o 0'v'T0J km'P O HU.'4234+0'

"  
"  
"  
"

Vj ku'o cpci go gpv'r rcp"ku"cp"lpkkcwxg"qh'yj g'hgf gtcn'i qxgtpo gpv'yj tqwi j "ku"qhheg."P CRGUEC."  
uwr r qtvgf "d{ "uekpkuu"dcugf "qp'yj g'dguv'uekpkkle"npqy ngf i g"cxckrdrg0"Hqt'yj g'htuv'ko g."  
ur cy plpi "P cuucwi tqwr gt'y kn'dg'ur gekkccm{ 'r tqvgevgf "dgw ggp'F gego dgt"3"cpf "Lcpwct{ '53"  
cppwcm{."o clpn{ 'hqt'yj g'O gzkcp"Ectkddgcp'y j gtg'yj ku'ur gekgu'ku'o qtg"cdwfp cp0"Cp"cf f kkkpcn'  
dcp'hqt'ecvej lpi "cm'i tqwr gtu'kp'yj g'I wh'qh'O gzkcp'y kn'gzvgpf "htqo "Lcpwct{ "37"vq'O ctej "36"  
cppwcm{0"Y kj kp'yj g'lwtkuf kkkp"qh'O gzkcp'kp'yj g'Ectkddgcp"Ugc."cm'ci i tgi cvlpi "i tqwr gtu."uwej "  
cu'drcen'i tqwr gt."*Myceteroperca bonaci*."y kn'dg'r tqvgevgf 0"Ucwwu"qh'yj g'dcp"cv'yj ku'ko g'ku"  
wpegtvckp0

"  
"  
"  
"

**TURKS AND CAICOS ISLANDS**

"

**Turks and Caicos Islands – Populations**

"

P cuucw'i tqwr gt 'kp'v'j g'Vwtm'cpf 'Eclequ'Kucpf u'VEK'ctg'pqv'uwdlgev'f 'v'uki p'k'ecp'v' eqo o gtecl'nc'kuj kpi 'r' tguuvtg='v'j g' 'ctg'eqpukf gt gf 'v'q'dg'kp'j' gcmj { 'eqpf k'k'qp'y' kj 't'gr'v'x'gn' { 'j' ki j ' f gpuk'k'gu'kp' 'u'qo g'ctgcu'Vwr r gt '4224+0'Vwr r gt '\*4224+'cpf 'Vwr r gt'cpf 'T'wf'f '\*4224+'t'gr'qt'v'gf " f gpuk'k'gu'kp' 'v'j g't'cpi g'qh'2067'v'q'20 'kp'f'k'k'f'w'c'u'r' gt '322'us'w'ct'g'o' g'v'gt'u'\*67/; 2lj'gev't'g+'y' kj " j' ki j' gt 'f' gpuk'k'gu'qp'f' ggr' gt 't'gg'h'ucpf' 'pq'f'k'h'gt'g'peg'kp' 'h'kuj 'ngpi' v'j' 'd' { 'f' gr' v'j' '\*Vwr r gt'4224.'Vwr r gt' cpf' 'T'wf'f'4224.'T'wf'f'4225c.'T'wf'f'.'4226+0'E'j' k'r'r' q'p'g'g'v'c'r'0\*4222+'t'gr'qt'v'gf "c'f' gpuk'v' { 'qh'2067/ 2084'P cuucw'i tqwr gt 'r' gt '322'us'w'ct'g'o' g'v'gt'u'cv'U'q'w'j' 'E'clequ'uk'gu'0'V'j' g'ug' 'h'ki' w't'gu'eqo' r' ctg' " h'x'q't'cd'n' { 'y' kj '2023'r' gt '322'us'w'ct'g'o' g'v'gt'u'kp' 'v'j' g'f' gr'ng'v'f' 'H'q't'k'f'c' 'c't'g'c'cpf' '2088/2042'r' gt '322' us'w'ct'g'o' g'v'gt'u'kp' 'v'j' g'D'c'j' co' cu'kp'p'q'p'ur' cy' p'k'pi' 'k'o' gu'0'E'wo' w'r'v'x'g'f'c'v' 'h't'q'o' 'T'GGH'\*4225/ 4235+'t'gr'qt'v': : 7'P cuucw'i tqwr gt 'kp'3567'uw'x'g' { 'u'f' gpuk'v' { 'kp'f'g'z'30.'uki' j' v'k'pi' 'h't'g's'w'p'e' { " 870 ' -'c'et'qu'v'j' g'32/{g'c't'r'gt'k'q'f'0'Y' kj 'v'j' g'g'z'egr'v'k'p'q'h'U'c'n'Ec' { '\*u'00'? '3: 0' + 'c'm'v'j' gt' uw'x'g' { 'uk'gu'j' c'x'g'uki' j' v'k'pi' 'h't'g's'w'p'e'k'u't'c'pi' k'pi' 'h't'q'o' '7; 0' 'v'q'322' "

[\\*j wr <ly y 0ggHQtI lf d ltgr qt vulf kvlur gekulVY C 02; 94225/23/234235/26/29+0'](#)

Dgecw'ug'h'kuj kpi 'r' tguuvtg'ku'm'qy . 'h'k'gr'f' 'u'w'f' l'gu'ecp'r' t'q'x'k'f' g'x'c'n'w'c'd'ng'k'p'uki' j' w'u'k'p'v'q' 'v'j' g' geq'ni' { 'cpf' 'd'k'q'ni' { 'q'h'P cuucw'i tqwr gt'0'P cuucw'i tqwr gt 'h't'q'o' 'U'q'w'j' 'E'clequ'j' c'x'g'd'gg'p't'gr'qt'v'gf " v'q'v't'c'x'g'n'r'r' t'q'z'k'o' c'v'gn' { '62'h'o' 'v'q'c' 'r'c'ti' g'ur' cy' p'k'pi' 'ci' i' t'gi' c'v'k'q'p' 'c'v'R'j' k'k'r' u'T'g'g'h' 'q'h'v'j' g'k'uc'p'f' 'q'h' G'cu'v'E'clequ'c't'q'w'p'f' 'v'j' g'h'w'm'i'o' q'q'p' 'k'p' 'l'c'p'w'c't' { '\*T'wf'f' '4225c+0'V'j' k'u'c'i' i' t'gi' c'v'k'q'p' 'k'u't'c't'g'n' { 'h'kuj' g'f' 'f'w'g' k'u't'go' q'v'g' 'h'q'ec'v'k'p' 'c'p'f' 't'q'w'j' 'u'g'cu'\*T'wf'f' '4225c+'c'f'f' k'k'q'p'c'n'l'p'h'q't'o' c'v'k'q'p' 'c'd'q'w'v'j' k'u'c'i' i' t'gi' c'v'k'q'p' 'k'u' u'c't'eg'0'U'w'f' l'gu'j' c'x'g' 'u'j' q'y' p' 'l'w'x'g'p'k'gu' 'u'g'w'g' 'k'p'uj' q't'g' '\*E'nc' { 'f' q'p' 'c'p'f' 'M'q'g'v' '4229+0'k'p' 'c'p' " w'p'f' g't'y' c'v'g't' uw'x'g' { 'eq'p'f' w'ev'g'f' 'h't'q'o' '42' O'c' { 'v'q'45' C'w'i' w'u'v'4229' u'q'w'j' 'q'h'U'q'w'j' 'E'clequ.'42; " P cuucw'i tqwr gt 'l'w'x'g'p'k'gu' '\*>34'eo' 'V'N'+y' g't'g'q'd'ug't'x'g'f' 'y' kj' k'p' 'q't' 'e'm'ug'v'q' '\*42'o' +'u'g'c'i' t'c'u'u'd'g'f' u'0' " U'q'r'k'c't' { 'eq'p'ej' 'u'j' g'm'u'y' g't'g'q'ee'w' k'g'f' 'd' { 'g'c't'n' { 'l'w'x'g'p'k'g'P cuucw'i tqwr gt 'd'w'v'j' g'ug'y' g't'g' 'r'c'ti' g'n' { " c'd'ug'p'v' 'h't'q'o' 'u'g'c'i' t'c'u'u'c't'g'c'u'v'q' 'v'j' g'p'q't'v'j' 'q'h'F'q'x'g' 'E'c' { 'r' q'u'uk'd'n' { 'd'gecw'ug'v'j' g'ug'j' c'd'k'c'v'u'c't'g' 'k'p' 'e'm'ug' " r' t'q'z'k'o' k'v' { 'v'q' 'r'c'p'f' 'c'p'f' 'v'j' g'c'v'k'x'k'k'gu' 'q'h' 'r'c'ti' g'x'g'u'g'n'u'c'u'v'j' g' { 'c't'g'j' g'c'f' k'pi' 'v'q' 'c'p'f' 'h'g'c'x'k'pi' 'v'j' g'p'g'c't'd' { " f' q'en'0' "

"

**Turks and Caicos Islands – Fishing**

"

V'j' g'P cuucw'i tqwr gt 'ku'j' ki j' n' { 'x'c'n'w'g'f' 'h'q't'v'j' g'm'q'ec'n'v'q'w't'k'u'o' 'c'p'f' 't'g'u'c'w't'c'p'v'o' c't'ng'u'c'p'f' 'k'u' c'n'q' 'k'o' r'q't'c'p'v'k'p' 'v'j' g'f' k'x'g't' 'v'q'w't'k'u'o' 'u'g'ev'q't'0'N'q'ec'n'r' q'r' w'r'v'k'q'p'u'c't'g'c'u'u'w'o' g'f' 'v'q'd'g' 'k'p' 'i' q'q'f' 'j' g'c'm'j' 0' " O'c'k'p' 'v'c'ti' g'v'ur' g'ek'gu'k'p' 'v'j' g'h'kuj' g't' { 'c't'g's' w'g'g'p' 'eq'p'ej' 'c'p'f' 'h'q'd'ug't' 'c'n'j' q'w'i' j' 'ec'v'ej' 'q'h' 'u'ec'ng' 'h'kuj' . " k'p'en'f' k'pi' 'P cuucw'i tqwr gt. 'h'q't'v'j' g'm'q'ec'n'o' c't'ng'u'j' c'u't'g'eg'p'v' { 'k'p'et'g'c'ug'f' '\*w'p'r' w'd'r'k'uj' g'f' 'F' gr' v'0'q'h' G'p'x'k't'q'p'o' g'p'v'c'p'f' 'E'q'c'u'c'n' 'T'g'u'q'w't'eg'u. 'V'wt'm'i'c'p'f' 'E'clequ'K'uc'p'f' u'P' c'v'k'q'p'c'n' 'T'g'r'q't'v.'422: +0'V'j' g'VEK' c't'g'o' q'x'k'p'i' 'v'q'f' k'x'g't'uk'h' { 'h'kuj' g't'k'gu'k'p'en'f' k'pi' 'h'q't' 'u'ec'ng' 'h'kuj' . 'y' j' k'ej' 'c't'g'g'x'k'f' g'p'v' { 'w'p'f' g't'w'k'k'k' g'f' " \*w'p'r' w'd'r'k'uj' g'f' 'F' gr' v'0'q'h'G'p'x'k't'q'p'o' g'p'v'c'p'f' 'E'q'c'u'c'n' 'T'g'u'q'w't'eg'u. 'V'wt'm'i'c'p'f' 'E'clequ'K'uc'p'f' u'P' c'v'k'q'p'c'n' "

"  
"  
"  
"

Tgr qtv."422: +0

Ncti gt "dqcvu'y kj "grgextle'tggm'j cxg'pqy "gucdrkuj gf 'c'rqecr'no ctngv'hqt'vj gkt'ecvej "cpf " kpetgcugf "j ctxgux0"Vj g'i tqy vj 'lp'vqwtkuo "j cu'kpetgcugf "vj g'f go cpf "kp'vj g'rqecr'no ctngv'hqt " P cuucw'i tqwr gt0'O cp { 'rqecr'hkuj gtu'i clp'cf f kxqpcr'kpeqo g'htqo "vcti gv'pi 'tgg'hkuj 0"Dgecwug'pqv' cml'uecr'g'hkuj "ctg'vcngp'vq'rkpgugf "r tqeguqtu.'k'ku'f khhkwn'vq'npqy "vqcr'ecvej 0"Twf f "cpf "Vwr r gt " \*4224+'tgr qtvgf "vj cv'vj g'ncpf kpi "r tkegu'hqt "vj g'P cuucw'i tqwr gt "cv'f qemuf g'tgej gf "WU&502'r gt " ni "y j krg'hkuj gto gp'o ki j v'ugm'P cuucw'i tqwr gt "f k'gevn' "vq'tgucwcpw'hqt'wr "vq"WU&8702'r gt "ni 0" Uqo g'Uqwj "Eckequ'hkuj gtu'j cxg'dgi wp'vq'vcti gv'i tqwr gt "o qtg'tgegpvn' "cu'vj g'xcnw'g'qh'vj g'ecvej " ku'qh'gp'y qt'vj "vj g'g'zr gpug'qh'v'cxgnkpi "82'no "vq'Rtqxkf gpekrgu'vq'ugm'vj gkt'ecvej "y j gp'k' gzeeggf u'cdqw'322'ni "Twf f ."4225d+0"

P cuucw'i tqwr gt "ctg'cp'ko r qvcv'eqo r qpgpv'qh'vj g'o gpw'qh'tgucwcpw'hqt'rqecr' eqpuwo r vkp0"Vj g'P cuucw'i tqwr gt "ku'c'r qr wrt'i tqwr gt "dgecwug'uqo g'qvj gt "i tqwr gt "ur geku'\*g0 vki gt "i tqwr gt."Mycteroperca tigris'+0 c { 'eqpvc'k'eki wcvz'k'p'y j lej "rko ku'vj gkt'ucrgu'kp" t'gucwcpw0"Twf f "4226+'hqwpf "vj cv'vj g'k'p'v'qf wv'k'p'qh'cp'ko r qt'v'v'ct'k'h'qp'hkuj "uki p'k'k'ecpvn' " kpetgcugf "f go cpf "hqt'rqecr'P cuucw'i tqwr gt0'O cp { "P cuucw'i tqwr gt "ecw'j v'qp'vj g'Uqwj "Eckequ" hkuj kpi "i tqwpf u'ctg'vcngp'd { 'mqduv'g'f kxgtu'y j q'qr r qt'w'p'k'v'ecm' "ur gct'hkuj "Twf f "4225d+0"

Hgy "f cv'ctg'cxck'rdng'qp'vqcr'ecvej "qh'P cuucw'i tqwr gt "dw'fko k'gf "ERWG'f cv'uwi i guv' t'g'v'k'x'gn' "mqy "ecvej "tcv'u'eqo r ctgf "vq'qvj gt "tgg'hkuj gu0"Vwr r gt "cpf "Twf f "4224+'hqwpf "ERWG" hqt'P cuucw'i tqwr gt "vq'dg'20'mi "r gt'j qwt'eqo r ctgf "y kj "50'mi "r gt'j qwt'hqt "cml'tgg'hkuj 0"Hkuj " cdw'p'cpeg."cu'k'p'f'ecv'g'f "d { "ERWG."ku'mqy gt "d { '72' "qt'o qtg'k'p'hkuj gf "tcv'j gt "vj cp'rki j vn' "hkuj gf "qt" w'p'hkuj gf "k'g'0'r tqv'ge'v'g'f +ct'gcu'dw'f k'h'gt'g'f "k'w'g'dgy ggp'vj g'ncw'g't'y q' | q'p'gu'\*Vwr r gt "cpf "Twf f " 4224+0"Vwr r gt "cpf "Twf f "4224+'hqwpf "pq'f k'h'gt'g'p'gu'k'p'uk' g."cdw'p'cpeg'qt "dkqo cuu'dgy ggp" | q'p'gu'qh'f k'h'gt'g'p'v'hkuj kpi "k'p'v'p'uk' { "cpf "uwi i guv'g'f "vj cv'hkuj kpi "k'p'v'p'uk' { "y cu'w'p'k'ngn' { "vq'g'zr r'k'p" vj g'i t'g'cv'g't'cdw'p'cpeg'cpf "dkqo cuu'qp'f ggr gt "tgg'u0"

K'p'cf f k'k'q'p'vq'h'q'f . "P cuucw'i tqwr gtu'r tqxkf g'p'q'p/gz'v'c'v'k'x'g'geq'p'qo k'e'xcnw'g'\*g0 0'p'q'p/ r'g'v' cn'ec'v'j /cpf /t'g'g'cug'hkuj kpi "cpf "y k'f r'k'g'x'k'y kpi "+vq'f k'x'gtu'hqt "vqwtkuo 0'c'p'k'p'et'g'cug'k'p" P cuucw'i tqwr gt "cdw'p'cpeg'cpf lqt'o gcp'uk' g'cf f u'xcnw'g'vq'vj g'f k'x'g'g'zr g't'k'p'eg'dgecwug'o quv' f k'x'gtu'j cxg'r t'g'h'gt'g'p'gu'hqt'x'k'y kpi "o qtg'hkuj "cpf "o cp { "f k'x'gtu'g'zr t'guu'r t'g'h'gt'g'p'gu'hqt'x'k'y kpi " r'cti gt "hkuj "Twf f "4225c+0"Twf f "cpf "Vwr r gt "4224+'cnuq'tgr qtvgf "vj cv'up'q't'ng'rgtu'cu'y gm'cu'f k'x'gtu' r t'g'h'gt'x'k'y kpi "r'cti gt "cpf lqt'o qtg'cdw'p'cp'v'P cuucw'i tqwr gt0"

**Turks and Caicos Islands – Conservation and Management**

Qpg'ur cy pl'pi "ci i t'gi cv'k'p'uk'g'ku'r tqv'ge'v'g'f "htqo "hkuj kpi "k'p'P qt'v'j y guv'Rq'k'p'v'O c't'k'p'g" P cv'k'p'c'ri'Retm'Rtqxkf gpekrgu'\*F GE T "4226="P cv'k'p'c'ri'Retm'Q't'f k'p'c'p'eg'cpf "Uwdu'k'f k'ct { " Ngi kur'v'k'p'ECRO: 2"qh'3; : : +0"K'p'vj g'Vw't'm'i'cpf "Eckequ"K'ur'p'f u.'vj g'o cl'p'ci i t'gi cv'k'p'uk'g'ku" t'go q'g'cpf "t'q'w'j "y g'v'j gt "f w'k'pi "vj g'ur cy pl'pi "ugcu'p'j cu'i g'p'g't'cm' { "t'g'v't'k'v'g'f "hkuj kpi "cv'k'k'v' { 0" U'g'cu'p'c'ri'eq'w't'gu'o c { "r'nc { "c't'q'ng'k'p'hkuj g't'k'gu'o c'p'ci go gp'v'r r'p'p'k'pi "k'p'vj g'h'w'w'g'dw'k'p'vj g" u'j qt'v'g'to "ctg'p'q'v'uki p'h'k'ec'p'v'h'c'v'q'tu'hqt'P cuucw'i tqwr gt "eq'p'ug't'x'cv'k'p'k'p'vj g'Vw't'm'i'cpf "Eckequ"

"  
"  
"  
"

Kirpuf u\*Twf f .4225d-0"Hwml'r tqvgevkqp"qh'guugpvkcnP cuucw'i tqwr gt'j cdkxcv'cpf "ur cy plpi "  
o ki tcvkqp"eqttkf qtu"qp'yj g'xgt { 'pcttqy "htkpi g'qh'Eckequ'Dcpniy qwf 'ko r qug"geppqo ke"j ctf uj kr "  
qp"mecn'hkuj gtu'y j q'f gr gpf "qp'yj qug'ctgcu'ht'eqo o gtekn'ur geku\*ur kp { 'mqduvgtu+'cpf "  
uwdukv'peg'hkuj kpi \*Twf f "4226-0"Vwr r gt'cpf "Twf f "\*4224+'uwi i guvgf 'yj cv'ugcuqpcn'ur cy plpi "  
emquwtgu'kp'yj g'Vwtm'cpf 'Eckequ"Kirpuf u'o ki j v'j cxg'vq'dg'ugxgtcn'o qpj u'kp'ngpi yj \*gd 0'  
P qxgo dgt'yj tqwi j "O ctej +'vq'dg'ghgevkxg0"F gur kvg'tgrcvkxgn' hkwng'hkuj gt { 'hewu'qp'yj g'P cuucw'  
i tqwr gt.'yj gtg'ku'eqpuwo gt'kpvgtgu'kp'yj g'ur geku\*c'utqpi 'mecn'vqwtkuo 'ugevqt+'cpf 'c'uki pkkcpv'  
r tqwr qt vkqp"qh'hkuj 'kp'qpg'tgegpv'uwf { 'y cu'cngp'dgmjy 'yj g'uk g/cv'o cwtcvkqp'uq'r tguwtg'ku"  
gzr gevgf 'vq'i tqy 'kp'yj g'cdugpeg'qh'o cpci go gpv\*Ncpf uo cp'gv'cn0422; +0'  
"

"  
"  
"  
"

**UNITED STATES (FLORIDA)**

**United States (Florida) – Populations**

Cm j qwi j 'y gtg'ctg'hgy 'f cxc'qp'j kvqtke'cdwfp cpeg'qh'P cuucw'i tqwr gt'qhh'y j g'WUO' o ckrp'f . 'k'cr r gctu'yj cv'cdwfp cpeg'y cu'qpeg'j ki j 'kp'uqwj gtp'Hqtkf c' \*Ur tkpi gt'cpf 'O eGtrgcp.' 3; 84+0" Cpgef qvcr'tgr qtwa'htqo 'ur gcthkuj gtu'pqvgf 'rti g'f ckn' { 'ecvej gu'kp'y j g'3; 72u' \*Dqj pucem' 3; ; 2+0" Kpvtxky u'qh'Hqtkf c'Mg{ u' t'gukf gpvu' uwi i guvgf 'y cv'P cuucw'i tqwr gt'y gtg'qpeg'ecwi j v'kp' o vej 'i tgcvt'pwo dgtu'htqo 'y j g'wr r gt'Hqtkf c'Mg{ u'cpf 'y j g'Dej co cu' \*Ucf qx { 'cpf 'Gmwpf '3; ; ; +0' Uctem' \*3; 8: +tgr qtvgf 'P cuucw'i tqwr gt'htgs wgpv' { 'cv'Crki cvqt 'Tggh'kp'y j g'Hqtkf c'Mg{ u' }

J kvqtkecm' . 'P cuucw'i tqwr gt'y cu'c'eqo r qpgpv'qh'y j g'i tqwr gt'hkuj gt { 'kp'Hqtkf c.' uwi i gukpi 'qpeg'j genj { '\*uwr qr wcvkq'p'u+'kp'uqwj gcuvgp'WUO' o ckrp'f 'y cvgtu' \*Ucf qx { 'cpf ' Gmwpf '3; ; ; +0' K'eqpvcuv.'pqy 'y j g'ur geku'ku'tctgn' { 'gpeqwpvgf ' \*Ucf qx { 'cpf 'Gmwpf '3; ; ; +0' K' y j g'F t { 'Vqtwi cu.'y j gtg'P cuucw'i tqwr gt'y gtg'qpeg'cdwfp cpv.'qpn' { 'qpg'kp'kxf wcn'y cu'tgeqtf gf " kp'3; ; 6'qw'qh'3: 5'r qkv'egpuwugu'cpf 'pqp'kp'59'r tgf cvqt 'egpuwugu' \*Uwnc'gv'cr03; ; ; +0' Qp' Gndqy 'Tggh'Hqtkf c'Mg{ u.'o gcp'P cuucw'i tqwr gt'f gpukku'y gtg'2023/'2026' hkuj 'r gt'322'o "4'kp" 3; ; 5/; 6' \*Uwnc'gv'cr03; ; ; + 'y kj 'hgy 'uggp'qp'egpuwuf'kxgu'y tqwi j 'y j g'Hqtkf c'Mg{ u'0' Egpwugu' eqo r ctkpi 'ctgcu'r tqvgevgf 'cpf 'vpr tqvgevgf 'htqo 'hkuj kpi 'kpf kecvgf 'y cv'P cuucw'i tqwr gt.'y j gtg' r tqvgevgf . 'j cf 'c'j ki j gt'f gpuk' { 'cpf 'y j g'qpg'qh'y j g'f qo kpcpv'i tqwr gt'ur geku'qdugtxgf ' \*Uwnc'gv' cr03; ; 6+0F gur kg'32/42" { gctu'qh'pq/vcng'r tqvgevgf'qh'y j g'P cuucw'i tqwr gt'kp'y j g'Hqtkf c'Mg{ u.' P cuucw'i tqwr gt'j cu'o cf g'pq'cr r tgekdng'tgeqxt { 'cpf 'pwo dgtu'tgo ckp'gz'vgo gn' 'mqy " \*Ugo o gpu'gv'cr0'4229c.' F qp'F gO ctk'r gtu0'eqo o 04234+0'"

Tggh'hkuj 'uwxg { u'd { 'y j g'P O HU'Uqwj gcu'vkuj g'kgu'Uekppeg'Egpvgt'u' \*UGHUE+'Tggh' Vgco 'tgxgcnf'mqy 'f gpukku'htqo '3; ; 2/3; ; 6'kp'uqwj gtp'Hqtkf c' \*Hki 046+'qh'5.73: 'xkuwcn'r qkv'




Figure 22. Counts of Nassau grouper observed in SEFSC reef fish visual census in the Florida Keys from 1980-2011

"  
"  
"  
"

eqwpu'P cuucw'i tqwr gt'y gtg'tgeqtf gf "4; "lko gu.'yj g'pwo dgt'f genkpi 'vq' | gtq'lp'3; ; 50"Dqj 'yj g' pwo dgt'qh'P cuucw'i tqwr gt'cpf 'yj g'pwo dgt'qh'uwxg{ u'kpetgcugf 'Htqo '3; ; 7'wv 'vq'4227"\*Hki 047+0"




Figure 23. Mean Percent Frequency of Occurrence of Nassau grouper in 3-year Intervals (except for 2010-2011), data from SEFSC

Htqo '3; ; 2'vq'3; ; 8.'Nqqg'Mg{ 'cpf 'O qrcuugu'y gtg'yj g'qpnf 'r tqvewgf '\*o ctkpg'tgugtxg+'ukvgo'0"kp' 3; ; 9.'yj g'o ctkpg'tgugtxg' | qpgu'hqt 'yj g'Hqtkf c'Mg{ u'P cvkqpcn'O ctkpg'Ucpewct { 'y gtg' guwdrkj gf .lpenmf kpi 'URCu'\*Ucpewct { 'Rtgugtxcvkq'P'ctgcu+'cpf 'yj g'Mg{ u'y kf g' uco r nkpi 'f guki p'y cu'f gxgnr gf 'vq'o qpkqt' tgg'h'gequ{ u'go 'eqpf kkpqu'0Vj tqwi j qw'yj g' tcpi g'qh'uwxg{ u'\*Hki 047+'Htgs wpe { 'qh' qeewt'ppeg'hqt 'P cuucw'i tqwr gt'y cu'hqy 'cpf' eqo r ctdng'dqj 'kpukf g'cpf 'qwu'kf g'qh'o ctkpg' tgugtxgu'2'vq'3Q ' 'qh'uco r ngu'lpenmf gf " P cuucw'i tqwr gt'\*P O HU'UGHUE'f'c'v'." uwr r nkf 'd{ 'L'0Dmpf gcw.'4234+0"C'o cr 'qh' yj g'f kwtkdwkq'qh'r quk'xg'gpeqwpvgtu' uwi i guu'yj g{ 'ctg'f kwtkdwgf 'yj tqwi j qw' O qptqg'Eqwv{ 'cpf 'f qgu'pqv'uwi i guv'cp{ " engct'r cwgt'p'\*Hki 048+0"Ugr ctcvg'f'c'v'hqt"




Figure 24. Distribution of samples with positive counts of Nassau grouper (SEFSC data, map by J. Blondeau)

"  
"  
"  
"

uwxg{u'dgwy ggp"4222/4229"\*Mg{ "Ncti q"vq" Ft { "Vqtwi cu+ 'y kj ": 785'uwxg{u'qdugt xgf "432"  
P cuucw'i tqwr gt"qp"3; : "qh'yj g'uwxg{u"\*f gpukv{ <20223 lo "4"cpf "408' "uki j vki "Htgs wgpe{ "+"  
\*UGHUE"fcv. 'uwr r rkgf "d{ "V0Mgmkup"4234+0"

C'cti g'pwo dgt"cpf "f kxgtukv{ "qh'cf f kxqpcn'hkuj gt { /kpf gr gpf gpv'uwxg{u'd{ "uvcg'r qtv"  
uco r rgtu'qxgt 'y g'rcuv'f gecf g'j cxg'tguwngf "kp'tgeqt f u'qh'c'hgy "j wpt gf "P cuucw'i tqwr gt"rcpf gf "  
\*Cnglcpf tq'Cequc. "HY EE. 'r gtu0eqo o 00"Cf f kxqpcn'wpt gty cvgt "uwxg{u'd{ "y g'Hrqt kf c'Hkuj "  
cpf "Y kf rkg'Ego o kuukap"j cxg'dggp"eqpf wevgf "Htqo "3; ; ; /4229"wkpi "y q'uco r rki "  
cr r tqcej gu"3+"hpgct "t cpugeu"\*c "vqcn'qh"32"kp f kxk wcn"P cuucw'i tqwr gt"y gt g'tgeqt f gf "Htqo "  
3349"t cpugeu"\*52o "d{ "32o "y kf g="cpf "4+r qkpveqpwu"\*c "vqcn'qh"8; "kp f kxk wcn"P cuucw'  
i tqwr gt"y gt g'tgeqt f gf "Htqo "95; ; \*7"o "tcf kxu+"uwxg{u+0" F wtkpi "gki j v' { gctu'qh'uwxg{u'9; "  
P cuucw'i tqwr gtu"\*q w'qh"5; 49"vqcn'i tqwr gtu+"y gt g"qdugt xgf "y kj "; 4' "qh'yj g"P cuucw'i tqwr gt"  
dgwy ggp"57"cpf "92"eo "kp"rgpi yj \*100 eEcy ng{. "F kgevt. "F kx0qh'O ct kpg"Hkuj gt ku"  
O cpci go gpv'rgwt "v"UGT Q+0'

Ewo wrcvxg'f cvc"Htqo "TGGH"\*4225/4235+"tgr qt v'3544"P cuucw'i tqwr gt"kp"; 928'uwxg{u'  
\*f gpukv{ "kpf gz "304. "uki j vki "Htgs wgpe{ "3508' "+qxgt "y g"32/ { gct 'r gt kqf 0"Uwxg{u'wr "y g'gcu'  
eqcu'qh'Hrqt kf c'vq" Lwr kgt "krgv'tgr qtv": 5"P cuucw'i tqwr gt"kp"8985"uwxg{u"\*f gpukv{ "kpf gz "304."  
uki j vki "Htgs wgpe{ "304' "+cpf "qp"y g'y guv'eqcu'qh'Hrqt kf c'Htqo "Ecr g'Ucdrg"vq"Vco r c'Dc{ "34"  
P cuucw'i tqwr gt"kp"7; 2"uwxg{u"\*f gpukv{ "kpf gz "4. "uki j vki "Htgs wgpe{ "4' "+  
[\\*j wr <ly y 0ggqti lf d ltr qt vulf kuvur geku IVY C 122; 94225/23/234235/26/29+0'](#)

P q"P cuucw'i tqwr gt"ur cy plpi "ci i tgi cvkqp"ukgu"j cxg'dggp'tgr qt vgf "kp"Hrqt kf c'y cvgtu0'  
"

### United States (Florida) – Fishing

Ego o gtekn'rcpf lpi u'qh'P cuucw'i tqwr gt"qh'Hrqt kf c'u' C'rcpve"eqcu'y gt g'ecw j v'  
r tko ctkf "d{ "j cpf rkgu. "cnj qwi j "ecvej gu"Htqo "ur gct hkuj lpi "vqcn'lo qt g'yj cp"qpq's wct vt "qh'yj g"  
eqo o gtekn'rcpf lpi u'kp"3; ; ; .3; ; 3. "cpf "3; ; 4" \*ekgf "kp"Ucf qx{ "cpf "Gmwpf "3; ; ; +0"Ego o gtekn'  
hkuj gt ku"fcv' r tkt "vq"3; ; 8" f kf "pqv'f kxkpi wkuj "rcpf lpi u'vq"ur geku" \*gd 0"i tqwr gt+"u" f gvc krgf "fcv"  
hqt "P cuucw'i tqwr gt"rcpf lpi u'ctg'pqv'cxk'cdrg' r tkt "vq"yj cv"3; ; 8"cu'yj g{ "y gt g'i tqwr gf "y kj "qy j gt"  
i tqwr gt"ur geku"0"O quv'tget gcvkqpcn'ecvej "kp"y g" WUOC'rcpve"eco g"Htqo "r tkxcv'ltgpvcn'dqcu0'

Ego o gtekn'rcpf lpi u'qh'P cuucw'i tqwr gt"Htqo "y g'gcuvgtp" I wh'qh'O gz leq. "y gt g'd{ "  
j cpf rkgu"cpf "rpi rkgu"cpf "ceeqwpgf "hqt": 2/322' "qh'P cuucw'i tqwr gt"eqo o gtekn{ "rcpf gf . "d{ "  
y gki j v' "Htqo "3; ; 8/3; ; 4" \*Ucf qx{ "cpf "Gmwpf "3; ; ; +0" kpekf gpvcn'ecvej "qh'P cuucw'i tqwr gt"cnj"  
qeevt gf "kp" hkuj "tcr u. "y kj "y g'pwo dgt "qh'tcr /ecw j v'i tqwr gtu'kpet gculpi "ulpeg"3; ; 6" \*I O HO E"  
3; ; ; +0" k' yj g"3; ; 2u. "o quv'ecvej "Htqo "y g'tget gcvkqpcn'hkuj gt { "y cu' "Htqo "r tkxcv'ltgpvcn'dqcu"  
\*f gvc krgf "kp"Ucf qx{ "cpf "Gmwpf "3; ; ; +0" Cp"cpn{ "uki"qh'yj g'j gcf dqcv'ugevt "qh'yj g'hkuj gt { "uj qy gf "  
c' r gcn'lp"j gcf dqcv'ecvej gu'kp"3; ; 3/3; ; 4" ctqwpf "306"o v'y kj "c"uvgr "f ger kpg"vq"cdqw"2057"o v'd{ "  
3; ; ; \*Dqj pucni4225+0"D{ "o cvej lpi "t gpf u'y kj "Ewdcp" hkuj gt ku. "k'ku"pqv'wpt gcuqpcdrng"vq"  
eqpenmf g'yj cv'j gcf dqcv'ecvej gu'kp"y g'Hrqt kf c'Mg{ u'lp"y g"3; 82u'y qwf "j cxg'dggp"5"vq"6"vko gu"  
j ki j gt "yj cp"gz kxkpi "cpf "vgo r qtcn{ "tko krgf "rcpf lpi u'fcv" \*Dqj pucni4225+0'

:: "

"  
"  
"  
"



UNITED STATES (PUERTO RICO)

United States (Puerto Rico) – Populations

Rwgtvq'Tleq'qpeg'j cf 'uki pkhecpv'ncpf kpi u'cpf . 'j gpeg. '\*uwr+r qr wrcvkap\*u+'qh'P cuucw' i tqwr gt'cpf 'cv'hcuv'qpg'uwduxcpv'nci i tgi cvkqp'kp'ku'uqwj y guv'eatpgt.'ceeqt f kpi 'vq'cpgcf qvci' tgr qt u' \*Ucf qx { '3; ; 5+0'Vj ku'ci i tgi cvkqp'cr r gct u'vq'j cxg'ncpi 'ukpeg'f kucr r gctgf 'cpf 'ncpf kpi u'qh' vj g'ur gekgu.'ceeqt f kpi 'vq'tgi wrct'r qt v'uwtxg{u'eqpf wevgf 'd{ 'vj g'i qxgtpo gpvwi':Ncdqtcvqt'kq'f g' kpxguki cekqpgu'Rgus wgtcuo' \*Hkuj gt { 'Tgugctej 'Ncdqtcvqt { '+qxtg'ugxgcnf' gecf gu.'f tqr r gf 'vq' pgi rki kdng'ngxgn'dghqtg'vj g'ur gekgu'y cu'hwnc{ 'r tqvgevgf '\*kp'eqo o qpy gcnj 'cpf 'hgf gtcn'y cvgtu+'kp' 42260' Cnj qwi j 'qpn{ 'c'ukpi ng' \*r gtj cr u'tghqto kpi +'ur cy plpi 'ci i tgi cvkqp'j cu'tgegpv' dggp'hqwpf " \*Uej @gt'gv'cr04234+'vj gtg'y gtg'qeeucqpcn'tgr qt u'qh'lwxgpkng'ugwngo gpv'kp'ncn'y cvgtu' uwi i guv'kpi 'gkj gt'ur cy plpi 'ci i tgi cvkqp'u'cv'wmpqy p'ukgu'kp'vj g'tgi kqp.'cpf lqt'vj cv'o cvkpi 'kp' uo cmgt 'i tqwr u'\*g0 0'r cktgf 'kpf kxf wcu+'qeewu'0'K'ku'cnq'r quukng'vj cv'ncxgc'ctg'eqo kpi 'qp' ewtgpv'htqo 'f kucpv'kucpf u'kp'vj g'tgi kqp '\*Ci wrct/Rgtgtc'gv'cr04228+0

Uxggtcn'uwf kgu'j cxg'dggp'eqpf wevgf 'ctqwpf 'vj g'kucpf u'qh'Rwgtvq'Tleq'kp'tgegpv' { gct u'd { " xkuwcn'uwtxg{u0'Cv'O qpc'cpf 'O qpkq.'uo cm'kucpf u'vq'vj g'y guv'qh'Rwgtvq'Tleq.'kp'4222'cpf '4227.' 9'P cuucw'i tqwr gt'lwxgpkngu'y gtg'hqwpf 'kp'uj cmqy 'ugci tcuu'cpf 'twdng'j cdkcv'y kj kp'vj g'tggh' nci qap0'K'p'y kpvgt'4226.'4'cf wv'P cuucw'i tqwr gt'y gtg'hqwpf 'kp'eqtcn'tggh'qh'uqwj gtp'O qpc" f wtkpi 'uwtxg{u'ht' i tqwr gt'ur cy plpi 'ci i tgi cvkqp'u' \*Ci wrct/Rgtgtc'gv'cr04228+0' Ceeqt f kpi 'vq' wpf gty cvgt'xkuwcn'uwtxg{u'htqo '4226'vq'4229'cv'O qpc'Kucpf . 'Rwgtvq'Tleq.'vj g'cdwpcpeg'qh' P cuucw'i tqwr gt'ku'gz vtgo gn{ 'ncy 'cpf 'ku'f kntkdwkqp'ku'iko kgf 'vq'ur gekke'f gr vj u'cpf 'j cdkcv' v'r gu'ceeqt f kpi 'vq' hku' 'uk' g'ercu' \*Uej @gt'gv'cr04229+0' P q'ur cy plpi 'ci i tgi cvkqp'u'qh'P cuucw' i tqwr gt'y gtg'gpeqwpvgtgf '\*Cr r grf qatp'r gtu0'qdu0'gxgp'vj qwi j 'tgr qt u'htqo 'hku' gto gp'f guetkdgf " cdwpcp'v'ci i tgi cvkqp'u'f cvkpi 'dcen'f gecf gu' \*Uej @gt'gv'cr04229+0' Gctn{ 'lwxgpkngu' \* > '32'eo 'VN+" ctg'qeeucqpcn' { 'qdugtxgf . 'uwi i guv'kpi 'uweegu'hwnt'gr tqf wev'kqp'uqo gy j gtg'cv'qt'pgct'O qpc" \*Uej @gt'gv'cr04229+'cnj qwi j 'vj g'ncxnc'nc'qh'cdqw'62'f c { u' \*Eqkq'gv'cr03; ; 9+'y qwf 'r tqxkf g' v'o g'ht'ncxgc'vq'tgcej 'O qpc'htqo 'o qtg'f kucpv'ncv'kqp'u'y j gtg'ci i tgi cvkqp'u'ctg'ukn'r tgu'p0

Ewtgpv'nc' t'gugctej 'ku'w'pf gty c { 'cv'vj tgg'i tqwr gt'ur cy plpi 'ukgu'qh'vj g'y guv'gp'eqcu'qh' Rwgtvq'Tleq0'Vj ku'y qn'ku'wukpi 'r cuuk'g'ceqwu'ke'o qpkqt'kpi 'cpf 'f kxgtu'vq's wcpv'kh' 'ur cv'cn' gzv'p'ur cy pgt'cdwpcpeg.'cpf 'ur cy plpi 'v'o kpi 0'Cv'qpg'qh'vj g'vj tgg'ukgu.'vj g'tgugctej gtu'j cxg" kf gpv'k'kf "c'uo cm'pwo dgt'qh'P cuucw'i tqwr gt'cuuqekcv'f'y kj 'ur cy pgtu'qh'q'vj gt'ur gekgu' \*Uej @gt' gv'cr04234+0' C f f k'kqpcn'y qtm'ku'dgkpi 'w'pf g'v'ncng'vq'o gcuwt'g'cpf 'ej ctcevt'k' g'vj g'ur cy plpi 'qh' P cuucw'i tqwr gt'cv'vj ku'ukg' \*T0Cr r grf qatp'cpf 'O 0'Uej @gt'r gtu0'eqo o 0'0'Qpg'qh'vj g'r gew'ct'k'k'gu' qh'vj g'r quukn' { 'ot'geqpu'kw'w'gf o'ur cy plpi 'ci i tgi cvkqp'ku'vj cv'vj g'v'o kpi 'uggo u'vq'f'k'ht'htqo 'vj g' v'ctf k'kqpcn'y kpvgt'o qpvj u'cpf 'g'xkf gpeg' uwi i guv'k'o c { 'dg'qeewt'kpi 'o qpvj u'ncv'gt'vj cp'g'zr gev'f 0'

Ewo wrcv'k'g'f cv'htqo 'TGGH' \*4225/4235+'tgr qt v'54'P cuucw'i tqwr gt'kp'345; 'uwtxg{u' \*f gpuk' { 'kpf gz'30. 'uki j v'kpi 'htgs w'p'e { '408' '+'cetqu'vj g'32/ { gct'r g'kqf 0'Qh'vj g'P cuucw'i tqwr gt' kpen'f gf 'kp'vj gug'uwtxg{u'cm quv'qpg/vj kf 'qh'vj go 'ctg'htqo 'vj g'kucpf 'qh'E wgd'tc'y j gtg'vj g'

"  
"  
"  
"

Nwku'Rg° c'P q/Vcng'O ctłpg'T gugt xg'ku'hqecvfg 0"

[\\*j wr <ly y y 0ggħti lf d ltr qt vulf kuvlur gekuIVY C 122; 9 4225/23/23 4235/26/29+0'](#)

**United States (Puerto Rico) – Fishing**

kp'vj g'WUOXkti kp'Kırcpf u'cpf 'Rwgtvq'Tleq.'tggh'hkuj 'ctg'ecwi j v'd { 'hkuj 'vtr 'y kj 'uqo g' ur gcthkuj kpi 'cpf 'j cpf rkpki 0Vj g'dqcw'uugf 'ctg'uo em'tcpi kpi 'htqo '36'vq'62'h0kp'vj g'vtr " hkuj gt {ô nguu'vj cp'9Q "o "ımpı \*Cr r grf qqtp'cpf 'O {gtu'3; ; 5.'Ci ct'gv'cı0'4227.'EHO E+0'Hkuj gt u' j cxg'vcti gvfg 'P cuucw'i tqwr gt'ur cy plıpi 'ci i tgi cvkqu'ułpeg'vj g'3; 72u0'Ceeqtł kpi 'vq'hkuj gt" kpvgtxıgy u'P cuucw'i tqwr gt'ırcpf kpi u'htqo 'O qpc'Kırcpf 'tcpi gf 'htqo '449'ni '\*722'r qwpf u+vq'8: 3" ni '\*3.722'r qwpf u+r gt'7/9'f c { 'vkr 'dghqtg'vj g'3; ; 2u.'dw'ıwdugs wgpvı { 'f gerkpıgf 'uq'vj cv'hkuj kpi " vkr u'vq'O qpc'Kırcpf 'y gtg'pq'ımpı gt'hgculdrg'\*Uej @gt'gv'cı0'4229+0'

Rwgtvq'Tleq'j cu'ımpı 'eqmıgevgf 'uqo g'ırcpf kpi u'f cv'cv'vj g'ur geku'ıgxgrıhtqo 'ku'hkuj kpi " eqo o wpkıku0'K'ku'vj wu'y gm'f qewo gpvgf 'vj cv'vj g'P cuucw'i tqwr gt.'f qo kpcpv'kp'vj g'3; 72u'vq" 3; 92u.'j cu'ıłpeg'xcpkuj gf 'htqo 'vj g'eqo o gteknıhkuj gt { \*RTFP T'4234+0'Vj g'ur geku'y cu" gxıf gpvı { 'j gcxkı { 'hkuj gf . 'ıpenıf kpi 'f vıtkıpi 'ku'ur cy plıpi 'r gıtkıf u.'y kj 'uo cmgt '\*ko o cwtg'ıkkı gf + " hkuj 'vcngp'kp'hkuj 'vtr u'\*Ucf qx { '3; ; 5.'Ucf qx { 'cpf 'Gmıwpf '3; ; ; . 'Ucf qx { 'r gıtu0'ıdu0'0'F vıtkıpi 'vj g' gctıf '3; ; 2u.'ırcpf kpi u'f gerkpıgf 'cpf . 'd { '3; ; : /3; ; ; . 'P cuucw'i tqwr gt. 'vj g'f qo kpcpv'eqo o gteknı' i tqwr gt'ıłpeg'vj g'3; 72u.'y cu'tctg'cpf 'tgr tıgıgpvgf 'qpnı '4' "qh'cmıfı tqwr gt'ırcpf kpi u'cpf '20ı' "qh" cmıf go gtucııhkuj 'ur geku'\*RTFP T'4234+0'K'y cu'eqpıkf gtgf 'gz vıpev'eqo o gteknı { 'dghqtg'3; ; 2" \*O cvqu/Ectcdcmı'422: =cmı qıı j 'vj g'ur geku'ıvıcmı'cr r gctı'kp'ırcpf kpi u'tgr qtıu'y j gtg'k'j cu" cxgtı gf 'cr r tqzko cvıgı { '33.222'r qwpf u'c" { gct'htqo '3; ; 6/42280"

Uko krt'ımpı /vgto 'f gerkpıgy gtg'ıggp'kp'eqo o gteknıırcpf kpi u'htqo 'Rwgtvq'Tleq'cpf 'vj g' WUOXkti kp'Kırcpf u'Eqo o gteknıırcpf kpi u'qh'P cuucw'i tqwr gt'kp'Rwgtvq'Tleq'tgr tıgıgpvgf 'c'o clqt" eqo r qpgpv'qh'vj g'hkuj gt { 'kp'vj g'ırvıg'3: 22u'\*Y kreız'3; ; ; . 'P lej qıı'3; 4; +dw'f gerkpıgf 'vq'cp" kpi kpi pıhıecpv'eqo r qpgpv'd { 'vj g'3; ; 2u0'Cr r grf qqtp'gv'cı0'3; ; ; 4+ıtr qtıvgf 'vj cv'P cuucw'i tqwr gt" ceeqıwpvgf 'ıht'363'qıw'qh'48.4; 6'vıvıcnıhkuj gu'ıco r ıgf 'kp'3; ; 7'cpf 'qpnı '5: 'qıw'qh'48.276'hkuj " ıco r ıgf 'kp'3; ; 2'\*Dıj pıcem'4225+0'

**United States (Puerto Rico) – Conservation and Management**

Vj g'o cpci go gpv'qh'hkuj gt { 'tguqıtegu.'ıpenıf kpi 'P cuucw'i tqwr gt. 'ku'ıj ctgf 'dgy ggp'vj g'ııecıı' lıvıkf leıvıpcııhkuj gt { 'o cpci gıu'qh'Rwgtvq'Tleq'cpf 'vj g'Ectkddıgcp'Hkuj gtıku'O cpci go gpv'Eqıpeki' y kj 'uqo g'cvıj qtıku'ur rıv'dgy ggp'eqo o qpy gıcnı hıgttkıqtıcnıy cvıtu'cpf 'ıgf gıcnıy cvıtu0C" o kpi wo 'ıkkı g'ıht'P cuucw'i tqwr gt'y cu'kpııqıf vıegf 'kp'3; ; 7'cpf . 'ghıgeıvıxg'P qıxgo dıgt'3; ; 2.'vcıng" cpf 'r quıguıkuı'qh'vj g'ur geku'y gtg'r tıj kdkıgf 'kp'WUOııgf gıcnıy cvıtu'EHO E'3; ; 8+0'kp'Rwgtvq" Tleq.'vj g'ur geku'y cu'ıımı { 'r tqıgevgf 'kp'dıj 'ıvcıv'cpf 'ıgf gıcnıy cvıtu'd { '42260'Dgecvıwg'o quıv'qh" vj g'ecr wtg'qh'P cuucw'i tqwr gt'kp'vj g'WUOEctkddıgcp'qıewıu'kp'vıgttkıqtıcnıy cvıtu'\*Rwgtvq'Tleq" cpf 'vj g'WUOXkti kp'Kırcpf u+'y j gtg'ıgf gıcnıhkuj gtıku'tıgıvıvııı'ıı'pıv'cr r nı . 'vj g'kpııqıf vıvıkuı'qh"

"  
"  
"  
"

r tqgevkkp'lp'Rwgtvq'Tleq'lwtkuf kkkpncly cvgtu'lp'4226'y cu'r ctvkwrcnq' 'ko r qtvcpv'Vcdrg'33+0'

Table 6. Summary of Nassau grouper regulations in the U.S. Caribbean (García-Moliner and Sadovy 2008); PR = Puerto Rico, St. Thomas and St. Croix = U.S.V.I.

Year	Reef Fish FMP Regulations
3; ; 7"	O kp"Uk g"34ö"q"46ö"petgculpi "3'lp1{t="Ugcuqpcn'enuwtg"*r tqj kdkkqp"qp'vcng+Htqo 'Lcpwct { "3"vq" O ctej "53"gej "{gct'lp'Hgf gtcn'y cvgtu"
3; ; 2"	P q"j ctXgUV'qt'r quuguukqp'lp"WU'hgf gtcn'y cvgtu"; /422"po ="Ugcuqpcn'enuwtg"cv'Tgf "J kpf "Dcpni'U0' Vj qo cu"F ge/Hgd+"]3; ; ; "pq/vcng_"
3; ; 5"	Ugcuqpcn'enuwtg'hqt'tgf "j kpf "cv'Vqwtö cirkp"*RT+"cpf "Nepi "Dcpni'U0'Etqkz+"
3; ; 8"	Ugcuqpcn'enuwtg'hqt'tgf "j kpf "Dclq"f g"Uleq."Cdtk"Nc"Ugctc"*RT+"
4226"	P q"j ctXgUV'qt'r quuguukqp'lp'Rwgtvq'Tleq'uvvg'y cvgtu"*vq"; "po ="pq'hkngvki "cv'ugc"
4227"	Cm'ugcuqpcn'ctgc'enuwt gu<r tqj kdk'dqwqo "vgpf kpi "i gct="pq'hkngvki "hkuj "cv'ugc"
4228"	P q"j ctXgUV'qt'r quuguukqp'lp"WUXX00="pq'hkngvki "cv'ugc"

"

"  
"  
"  
"



vj tgg'P cuucw'i tqwr gt'y gtg'qdugt'xgf 'kp'yj g'uwwf { 'tgi kqp'qxgt'yj g'eqwtug'qh'yj g'ukz '{gctu'qh' o qpkqtkpi . 'i kxkpi 'o gcp'cdw'pf cpeg. 'htgs w'p'e { . 'cpf 'o gcp'dkqo cuu'qh'201. '>2023. 'cpf '8; 0 'i tco . ' t'gur gev'kxgn { '\*Rkwo cp'gv'cn'422: +0




Figure 26. Mean biomass of Nassau grouper from St. John surveys (A. Friedlander, unpub. data).

Hqmqy kpi 'yj g'eqm'r ug'qh'yj g'P cuucw'i tqwr gt'hu'j gt { 'kp'yj g'WUXKkp'yj g'rcv'g'3; 92u'Qm'gp' cpf 'NcR'rc'g'3; 9; + 'yj gtg'y cu'pq'uki p'k'hecp'v'ur cy pl'kpi 'ci i tgi cv'kqp'ht'yj ku'ur gekgu'qp'yj g'uj grh' uqwj 'qh'U0Vj qo cu'qt'U0Lqj p0'J qy gxgt. 'hu'j gto gp'ctg'tgr qt'v'kpi 'c'r quukdr'g'tgewtt'gpeg'uqwj 'qh' U0Lqj p'F'0'Qm'gp'r gtu0'eqo o 0'4233+'cpf 'Mcf kuqp'gv'cr0'4232+'cpf 'P go gj 'gv'cr0'4228+'uwi i guv' yj cv'yj gtg'o ki j v'dg'cp'ci i tgi cv'kqp'tg/ht'o kpi 'cv'qpg'qh'yj g'ukgu'uqwj 'qh'U0Vj qo cu' 'Ceeqtf kpi " vq'f kxgt'w'xt'xg { u'eqpf w'ev'gf 'kp'4223/4226. 'c'uo cmi'P cuucw'i tqwr gt'ci i tgi cv'kqp'j cu'dggp'qdugt'xgf " cv'yj g'I tco o cp'kmi'Dcpm'c'f ggr 't'ggh'52/62'o + 'm'ecv'gf "qp'yj g'uj grh'gf i g'uqwj 'qh'U0Vj qo cu' \*Mcf kuqp'gv'cr0'4232. 'P go gj 'gv'cr0'4228+0'kp'4224. 'uo cmi'env'ugtu'qh'P cuucw'i tqwr gt. 'r quukdr' " tgr t'gugp'v'kpi 'yj g'gct'r'kgu'v'uci gu'kp'yj g't'geq'xgt { 'qh'c'ur cy pl'kpi 'ci i tgi cv'kqp. 'y gtg'p'q'v'gf 'cv' I tco o cp'kmi'Dcpm'y j k'g'kp'O cte'j '4225. 'c'ukpi ng'env'ugt'qh'P cuucw'i tqwr gt. 'p'q'v'r t'g'x'k'w'ur' " t'geqtf gf 'kp'g'k'j gt'F gego dgt'qt' 'Lcpwct { . 'y cu'qdugt'xgf 'cv'yj g'uco g'ukg' \*P go gj 'gv'cr0'4228+0' Vj gtg'y cu. 'j qy gxgt. 'p'q'ergct'g'x'k'f gpeg' \*e. g. 'dgj cxk'qt. 'eqm'tcv'kqp+'yj cv'P cuucw'i tqwr gt " uweegu'uh'wm { 'ur cy pgf 'kp'4224'qt'4225'cv'I tco o cp'kmi'Dcpm'0'kp'Cr t'ki'4226. 'cdq'w'82'P cuucw' i tqwr gtu'ci i tgi cv'gf 'qp'yj g'I tco o cp'kmi'Dcpm'6'q'w'qh'82'hu'j 'y gtg'uggp'kp'd'leq'rt' 'r j cug'dw'p'q' " eqw'uj kr'qt'ur cy pl'kpi 'y cu'qdugt'xgf ' \*P go gj 'gv'cr0'4228+0' T'gegp'v'y qtm'd { 'P go gj 'cpf " eqy qtn'gtu' \*r gtu0'eqo o 0'j cu'f q'ewo gp'v'gf 'u'qo g'lp'et'g'cugf 'ug'w'go gp'v'it'get'w'ko gp'v' \*4226/4228+'kp' pgctuj qtg'j cdk'cw'kp'd'q'j 'U0Vj qo cu'cpf 'U0Lqj p. 'cpf 'yj g { 'j cxg'f go qp'ut'cv'gf 'uweegu'v't'cen'kpi " P cuucw'i tqwr gt'vq'yj g'I tco o cp'kmi'Dcpm'ur cy pl'kpi 'ukg0'K'ku'r quukdr'g'yj cv'c' { gct'qt'y q'qh'w'x'q'pi " t'get'w'ko gp'v'q'ee'w't'gf 'y k'j 't'gu'w'kpi 'uo cmi'lp'et'g'cugu'kp'm'ec'ni'cdw'pf cpeg' \*P go gj 'gv'cr0'kp' 'r tgr +0' Ceeqtf kpi 'vq' 'Mcf kuqp'gv'cr0'4232+'o'Qp'U0'Et'q'kz. 'y j gtg'p'q'P cuucw'i tqwr gt'ci i tgi cv'kqp' ku'd'g'r'k'x'g'f 'vq'gz'k'w. 'hu'j gto gp'cpf 'f k'x'g'qr g't'cv'qtu'ci tgg'yj cv'i tqwr gt'ctg'cm' qu'v'eqo r ng'v'gn' " cdugp'v'ht'qo 'yj g'k'ku'q'rc'v'gf 'uj grh' 'I gtu'qp'O ct'v'k'p'g'j . 'hu'j gto cp'r gtu0'eqo o 0' 'O k'ej gr'g' R'w'j j . 'f k'x'g

"  
"  
"  
"

dwukpguu"qy pgt"cpf"qr gtcvqt"r gtu0eqo o 00Qpn{ "qpg"j cu'dggp"qdugtxgf "kp"ukz "{ gctu"qh'hkuj " uwtxg{ u'eqpf wexgf "cppwcm{ "qp"36"ukgu"ctqwpf "U0Etqkz "P go gj "Wpr wd0f cxc+06"

**United States (Virgin Islands) – Fishing**

Kp"vj g"WUOXki kp"Krcpf u'cpf "Rwgtvq"tleq. 'tggh'hkuj "ctg"ecwi j v'd{ 'hkuj "tcr "y kj "uqo g" ur gcthkuj kpi "cpf "j cpf nkpki 0"Vj g"U0Etqkz "hkuj gt { "vgpf u"vq"dg"c" f kxgt/f qo kpcvzf "hkuj gt { ." y j gtgcu"kp"U0Vj qo cu"vj g'hkuj gt { "vgpf u"vq"dg"tcr /f qo kpcvzf "Qnugp."r gtu0eqo o 00"Vtcr u'ctg" f guki pgf "y kj "dkqf gi tcf cdng"r cpnu"cpf "o guj "uk gu"j cxg'dggp"cf lwuvf "tgr gcvgf n{ "vq"tgf weg"vj g" d{ ecvej "qh'uo cm'hkuj 0"i kxgp"vj g"o guj "uk g. 'lwxgpkng"p cuucw'i tqwr gt"y qwr "dg'tgcf k{ 'tgcvkpgf "kp" tcr u0"

Cu'tgr qtvgf "d{ "O wptq"cpf "Drcm"4227+0i tqwr gt"ci i tgi cvkpu"kp"WUOX00y cvgtu"y gtg" j gcxk{ "gzr mqkgf "htqo "vj g"3; 82u"vj tqwi j "vj g"3; : 2u"y kj "vj g'i tgcvgu"ghhqt"vj cxkpi "uvtvgf "pqt"vj "qh"vj g"Rwgtvq"tlecp"krpcpf "qh"ewgdtc0"Ci i tgi cvkpu"qp"vj g"Dcttceqwc"Dcpm"pqt"vj "qh"U0 Vj qo cu. "y gtg'hkuj gf "vq"gz vkevqp"d{ "vj g"rcvg"3; 92u."r tqf wekpi "cu"o wej "cu"40"o v"o gtle" vqu+7222"rdu0"qh'i tqwr gt"r gt"fc { "cv"ku"r gcnö"MOVvtdg."r gtu"eqo o +0"

**United States (Virgin Islands) – Conservation and Management**

Kp"vj g"3; 92u."vj g"eqo o gteknj ctxguv"qh"vj g"p cuucw'i tqwr gt"kp"vj g"WUOX00tgej gf "ku" j ki j guvtgeqtf gf "r qkpv"cpf "k'y cu"cnq"kp"vj ku'f gecf "vj cv'y gm/f qewo gpvgf "f gerkpgu"qeewtgf "cv" qpg"ko r qtcvp"ci i tgi cvkqp"ukg"Qnugp"cpf "NcRceg"3; 9; +0"Nqecn'hkuj gto gp"y gtg"uq"eqpegtpgf " y kj "ecvej "ngxgn"vj cv"kp"3; 98."U0Vj qo cu'hkuj gto gp"tgs wguvf "vq"vj g"mecn'i qxgtpo gpv"vj cv" vj g'i tqwr gt"dcpm"dg"emugf "hqt"7" { gctu0"Vj gk"qpn{ "eqpf kxqp"y cu"vj cv"vj g"emuwg"y cu"vq"dg" ceeqo r cplgf "d{ "gphqtego gp0"Vj ku'tgs wguv"y cu'ki pqtgf "gpvktgn{ "Qnugp."UVHC."r gtu0eqo o 0vq" L0T wvgt. "P O HU."4235+0"kp"3; ; 2."vj g"Ectkddgcp"Hkuj gt { "O cpci go gpvEqwpeki"EHO E+"gpcevgf " c"r tqj kdkxqp"qp"0hkuj kpi "hqt"qt"r quugukqp"qh"p cuucw'i tqwr gt"kp"qt"htqo "vj g"WU'Ectkddgcp" Gzenukxg"Geppqo k\ apgö"vj tqwi j "ku"Uj cmqy /y cvgt"tggh'hkuj "O cpci go gpvRrcp0"kp"cf f kxqp." vj g"EHO E."y kj "uwr r qtv"qh"mecn'hkuj gto gp."guvdrkuj gf "c"pq/vcng"o ctkgp"r tqvgevgf "ctgc"qh"vj g" uqwj y guv"eqcu"qh"U0Vj qo cu."J kpf "Dcpm"O ctkgp"Eqpugtxcvqp"F kntlev"Dtqy p"4229+lpvgpf gf " vq"r tqvevtgf "j kpf "cpf "tgf "j kpf "ur cy plki "ci i tgi cvkpu0"Vj g"J kpf "Dcpm"O ctkgp"Eqpugtxcvqp" F kntlev"y cu"htu"uwdlgev"vq"cu"ugcuqpen'emuwg."dgi kppki "kp"3; ; 2"Dggw"cpf "Hkfg rcpf gt"3; ; ; ." P go gj "4227."P go gj "gv"cn04228+0"r tqvev"ur cy plki "ci i tgi cvkpu"qh"tgf "j kpf ."hmqy gf "d{ " { gct/tqwpf "emuwg"vq"hkuj kpi "kp"3; ; : "F R P T"4227+0"Vj g'emugf "ctgc"j cu'dggp"ghgexg"cv" tguqtkpi "tgf "j kpf "gxgp"vj qwi j "eqo r nkepg"j cu."cv"ko gu."dggp"s wguvqpcdn"LOTKgtc."UGTQ." r gtu0eqo o 0wpr wd0f cxc+."cnj qwi j "c"U0Vj qo cu'hkuj gto cp"y cu"cttgugf "cpf "r tqgewgf "kp" 422: "d{ "P QCC"Ncy "Gphqtego gpv"FOQnugp."UVHC."r gtu0eqo o 0vq"L0T wvgt. "P O HU."4235+0

Kp"WUOXki kp"Krcpf u'vgttkqtkcn'y cvgtu."vj g"ur geku."r tkqt"vq"4228."dgpghkgf "htqo "i gpgtcn' hkuj gtkgu"tgutlevkpu."wej "cu"i gct"tgutlevkpu"cpf "twgu"qp"vj g"o ctngvki "qh'hkuj ."cpf "vj qug"

"  
"  
"  
"

cr r n( kpi 'v'ur gekklem( 'r tqvev( 'ukgu.'uwej 'cu'v'j g'Xkti kP "Krcpf u'EqtentTggh'P cvkqpcn'  
O qpwo gpv\*pq/vcng+'Xkti kP "Krcpf u'P cvkqpcn'Rctn)\*pq'eqo o gtekn'hkuj kpi +'Dweni'Krcpf 'Tggh'  
P cvkqpcn'O qpwo gpv\*pq/vcng+'cpf 'ugxgtcn'WUOXO'o ctkpg'tgugtxgu" kP "4228.'v'j g'WUOXO'  
kpukwg( 'tgi wrvkvpu'v'q'r tqj kdk'j' ctxgu'qt'r quuguukqp'qh'P cuucw'i tqwr gt'kP "WUOXO'y cvgt'cpf "  
v'q'r tqj kdk'hkgv'kpi 'cv'ugc"\*I cte'c/O qnkpgt'cpf "Ucf qx{ "4229-0" kP "4232.'cu'r ctv'qh'c'UgcI tcv'  
r tqlgev'v'dtkpi 'cwgpvkqp'v'v'j g'P cuucw'i tqwr gt.'v'j g'U0Vj qo cu'Hkuj gto cpau'Cuqekcvkqp "  
\*UVHC +'f kwtkwg( 'pggf rgu'v'q'xgpv'uy ko "dnf fgtu'cpf 'tgeqtf 'cm'P cuucw'i tqwr gt'ecwi j v'htqo "  
Cr tki'v'q'Lwn( '0'P cuucw'i tqwr gt'y gtg'tgi wrctn( 'ecwi j v.'cmj qwi j "pqv'cu'cdw( cpw( 'cu'kP'v'j g'r cuv'  
\*F 0Qnpg.'UVHC.'r gtu'eqo o 0'v'q'LO'v'gvt.'P O HU.'4235-0'

Hqmjy kpi 'tgugctej 'hqt'o cp{ '{gctuf'qewo gpv'kpi 'i tqwr gt'ur cy pl'kpi 'cpf'o ki tcvkqp."  
P go gvj 'gv'cn0\*4228+'lwi i guvgf 'v'v'v'j g'ugcuqpcn'emqwtg'qh'v'j g'I tco o cpkn'Dcpn'htqo 'Hgdwtct { "  
3'v'q'Cr tki'52'eqwf 'r tqxkf g'r tqvev'kqp '\*xlc'o cpci go gpv'o gcwtgu'kP'c'o wnk'ur geku'ur cy pl'kpi "  
ci i tgi cvkqp'ukg+'hqt'v'j g'r qv'p'v'kcm( 'tghqto kpi 'P cuucw'i tqwr gt'ur cy pl'kpi 'ci i tgi cvkqp'0'Vj g"  
I tco o cpkn'Dcpn'ur cy pl'kpi 'ci i tgi cvkqp'ukg'j' cu'dggp'ugcuqpcn( 'r tqvev'g'htqo 'Hgdwtct { "  
v'j tqwi j 'Cr tki'ukpeg'4228'dw'tgegpv'gxkf gpeg'htqo 'ceqwukc'v'ci i kpi 'cpf'j { f tqj qpg"  
xqecrk( cvkqpu'lwi i guv'v'j cv'P cuucw'ci i tgi cvg'v'q'ur cy p'cv'v'j g'I tco o cpkn'Dcpn'htqo 'Lcpwct { "  
v'j tqwi j 'O c{ 'y j lej 'o c{ 'y cttepv'cp'gz v'pukqp'qh'v'j g'I tco o cpkn'Dcpn'emqvgf 'ugcuq'v'q'hkxg"  
o qpj u'0'Vj g'J kP 'Dcpn'O ctkpg'Eqpugt'cvkqp'F kwtkev.'U0'Vj qo cu.'tgo ckpu'emqvgf 'v'q'hkuj kpi "  
{gct/tqwpf.'r tqvev'kpi 'c'tgf'j' kP 'ur cy pl'kpi 'ci i tgi cvkqp'cpf'c'hqto gt'P cuucw'i tqwr gt'ur cy pl'kpi "  
ukg0'

"  
"  
"  
"

## LITERATURE CITED

- Cegtg. 'C0R0'cpf 'L0I ct| qp/Hggtgk03; ; 300 gtqu."ej gtpcu{"ecdtkmru'f gn'Ectkdg"Eqnqo dlcpcq \*Rkuegu'Ugttcplf cg-'  
gr kprj gkpcg<gr kprj gkpk0Ecnf culc'38\*9: +577/5980"
- Ci ct. 'L00'0 0Uj kxpk 'L0T0Y cvgtu.'O 0Xcfr<sup>2</sup>u/Rk| kpk'V00 wtc{.'L0Mkmg{.'cpf 'F 0Uwo cp042270U.S. Caribbean  
Fish Trap Fishery Costs and Earnings Study0P QCC'Vgej plecnO go qtcpcf wo 'P O HU/UGHUE/'756.'349'r 0"
- Ci wkrct/Rgtgc.'C03; ; 60Rtgrko kpc{ "qdugtxcvqpu'qh'hj g'ur cy plpi "ci i tgi cvkqp"qh'P cuucw'i tqwr gt."Epinephelus  
striatus."cv'O clcj wcn'S wkpvcpc'Tqq.'O gzleq0Proceedings of the Gulf and Caribbean Fisheries Institute,"  
43:334/3440"
- Ci wkrct/Rgtgc.'C042260F gvgevkqp'qh'hkuj kpi "ghgeu'qp'c'P cuucw'i tqwr gt'ur cy plpi "ci i tgi cvkqp"tqo "uqwj gtp"  
S wkpvcpc'Tqq.'O gzleq0Proceedings of the Gulf and Caribbean Fisheries Institute."55:766/7780"
- Ci wkrct/Rgtgc.'C042280F kuc r gctcpeg'qh'c'P cuucw'i tqwr gt'ur cy plpi "ci i tgi cvkqp"qh'h'j g'uqwj gtp'O gzleq0"  
Ectkddgcp'eqcu0Marine Ecology Progress Series."327:4: ; /4; 80"
- Ci wkrct/Rgtgc.'L0C042220Uqenl'cpcn'uku'hqt'cp'ci i tgi cvkpi "tgg'hkuj . 'j g'P cuucw'i tqwr gt"Epinephelus striatus  
\*Rkuegu'Ugttcplf cg+'htqo "j g'O gzleq0'Ectkddgcp0O U'Vj guku."WRT/O c{ci Agl .": 7'r r 0"
- Ci wkrct/Rgtgc.'C0'cpf 'Ci wkrct/F cxk'Y 03; ; 80C'ur cy plpi "ci i tgi cvkqp"qh'P cuucw'i tqwr gt"Epinephelus striatus"  
\*Rkuegu'Ugttcplf cg+'lp"j g'O gzleq0'Ectkddgcp0Environmental Biology of Fishes,"45:573/5830"
- Ci wkrct/Rgtgc.'C0'0 0Uej @gt."cpf 'O 0P go gvj 042280Qeewtgpeg'qh'lwxgpkp'P cuucw'i tqwr gt."Epinephelus striatus  
\*Vrgquvkg'Ugttcplf cg+'qh'0 qpc'Kurpf . 'Rwgtvq'Tleq-'eqpu'f gtcv'qpu'qh'tgetwko gpv'r qv'p'v'cn0Caribbean  
Journal of Science."42(2):483/4870"
- Ci wkrct/Rgtgc.'C0'E0I qpl<sup>a</sup> rgl /Ucru."cpf 'J 0Xkmgj cu/J gtp<sup>a</sup> pf gl 0422; 0Hkuj kpi . 'o cpci go gpv'cpf "eqpugtxcvqpu'qh'  
j g'P cuucw'i tqwr gt."Epinephelus striatus."lp'j g'O gzleq0'Ectkddgcp0Proceedings of the Gulf and  
Caribbean Fisheries Institute."61:535/53; 0"
- Ckngp.'M0C0'cpf 'V0Utggv03; ; 50Lco clec0Rctv'K0K0'<Marine Fishery Resources of the Antilles: Lesser Antilles,  
Puerto Rico and Hispaniola, Jamaica, Cuba.'r 037; /3: 20HCQ'Hkuj gtlgu'Vgej plecn'Rcr gt0548.'457'r r 0"
- Cndkpu'O (C0'cpf 'O (C0J kzqp0422: 0Kpxcukg'lpf q/Rcekhe'kqphkuj "Pterois volitans'tgf weg'tgetwko gpv'qh'Cv'p'v'e"  
eqtcn'tggh'hkuj gu0Marine Ecology Progress Series."367:455/45: 0"
- Cndkpu'cpf 'J kzqp042330Y qtuv'ecug'uegpctkq<r qv'p'v'cn'ip'i /vgt0 "ghgeu'qh'kpxcukg'r tgf cvt{ 'kqphkuj "\*Pterois  
volitans+'qp'Cv'p'v'e"cpf 'Ectkddgcp'eqtcn'tggh'eqo o wpk'gu0Environmental Biology of Fishes."Qprkpg"  
Htu0'F QK"320229lu32863/233; 9; 7/30"
- Cndkpu.'O (C0'0 (C0J kzqp.'cpf 'l 0Ucf qx{0422; 0Vj tgcvgpgf 'hkuj gu'qh'h'j g'y qtrf <Epinephelus striatus \*Dnjej ."  
39; 4+\*Ugttcplf cg+0Environmental Biology of Fishes."86:52; /5320"
- Ckrctgl /Hkr 'N0'P M0F wkr{.'L0C0I km'K0 0Eqvq."cpf 'C0T0Y cndkpuqp0422; 0Hrcwgp'pi "qh'Ectkddgcp'eqtcn'tgghu'  
tgi kq/y kf g'f gerlpgu'lp'cte'j kgewtcr'leqo r r'gzk'0Proceedings of the Royal Society B0276:523; 652470"
- Co cti >u.'HR0'I 0 0Uc'p'p.'C00f gn'Ecu'kmq.'C0 0Htgp' pf gl . 'HO 0Dncpeq."cpf 'Y (C0f g'rc'Tgf 042320Cp"  
g'zr g'klo gpv'qh'hkuj "ur kmq'xgt'htqo "c"o ctkpg'tgugtxg'lp'Ew'c0Environmental Biology of Fishes,"87:58565940"
- Crr r g'f qatp.'T0'cpf 'U00 g{gtu03; ; 50Rwgtvq'Tleq'cpf 'J kur cpl'qm0Rctv'K0K0'<Marine Fishery Resources of the  
Antilles: Lesser Antilles, Puerto Rico and Hispaniola, Jamaica, Cuba.'r 0; ; /37: 0HCQ'Hkuj gtlgu'Vgej plecn'  
Rcr gt0548.'457'r r 0"

"  
Crr grf qqtp. "T0U0I 0F0F gppka."cpf "Q00 qpvgttquc"Nqr gl 03; : 90Tgxkgy "qh'uj ctgf "f go gtucn't guqwtegu"qh'Rwgtvq"  
Tleq"cpf "y g'Nguugt"Cpvknngu't gi kqp."r 058/328."kn:"Tgr qtv'cpf "r tqeggf kpi u'qh'y g"gzr gtveqpuwncvqp"qp"  
uj ctgf "Hkuj gt { 't guqwtegu"qh'y g'Nguugt"Cpvknngu't gi kqp."O c { ci wgl . "Rwgtvq" Tleq.": /34"Ugr vgo dgt"3; : 80HCQ"  
Hkuj gt lgu" Tgr qtv"5: 5: 49: "r r 0"  
"  
Crr grf qqtp. "T0U. L0Dggw. L0Dqj pucem"U0Dqif gp. "F 00 cvqu. "U00 g { gtu. "C0Tquctkq. "I 0Ucf qx { . "cpf "Y 0Vqdkcu0"  
3; ; 40Shallow water reef fish stock assessment for the U.S. Caribbean0P QCC"Vgej plecn"O go qtcpf wo "  
P O HU/UGHUE/526."92"r r 0"  
"  
Ctej gt. "U0M0"U0C0J gr r gm "D0Z0Ugo o gpu. "E0K0Rcwppi gm Ugo o gpu. "R0I 0Dwuj . "E0O eEq { . "cpf "D0E0Lqj puqp0"  
42340Rcwgtpu"qh'eqnqt"r j cug"lpf lecvg"ur cy p'vko kpi "cv'c" P cuucw'i tqwr gt "Epinephelus striatus ur cy plpi "  
ci i tgi cvkqp0"Current Zoology 7: \*3+<95-: 50"  
"  
Dckutg. "L0C03; ; 50Ewdc0Rctv"K0Kp. "Marine Fishery Resources of the Antilles: Lesser Antilles, Puerto Rico and  
Hispaniola, Jamaica, Cuba."r 03: 3/"4570HCQ" Hkuj gt lgu"Vgej plecn"Rcr gt"548."457"r 0"  
"  
Dckutg. "L0C0"cpf "L0Rcgl 03; : 30Nqu't gewtuqu'r gus wgtqu'f grictej kr grci q "Ewdcpq0Y GECH"Uwf 0: . "9; "r r 0"  
"  
Dcrf y kp. "E0E0I 0F0Lqj puqp. "cpf "R0N0Eqrkp03; ; 30Nctxcg"qh"Diplorion bifasciatum."Belonoperca chabanaudi"cpf "  
Grammistes sellineatus"\*Ugttcplf cg-<Gr kpr j grkpcg+y kj "c"eqo r ct kuqp"qh'npqy p"rtxcg"qh'qj gt"  
gr kpr j grkpgu0"Bulletin of Marine Science."48(1):89/; 50"  
"  
Dcpgtqv"U0R0"Y 0Y 0Hqz. "I. "cpf "L0G0Rqy gtu03; : 90Tgr tqf wevkg"utcvgi kgu"cpf "y g"o cpci go gpv'qh'upcr r gtu"cpf "  
i tqwr gtu"lp"y j "I wh'qh'O gzleq"cpf "Ectkddgcp."r 0783/825."kn:"L0L0Rqrxkpc"cpf "U0Tcnwqp"\*gf u0Vtqr lecn"  
Upcr r gtu"cpf "I tqwr gtu-<Dkqmi { "cpf "Hkuj gt lgu"O cpci go gpv0Y guxkgy "Rtguu."Dqwf gt. "EQ0"  
"  
Dctfcej . "L0G03; 7: 0Qp"y j"o qxgo gpw"qh'egtvcpl"Dgto wf c'tgg'hkuj gu0"Ecology"5; \*3+<35; /3680"  
"  
Dctfcej . "L0G0"cpf "F 0Y 0O gpl gr03; 790Hkgrf "cpf "rdqtcvqt { "qdugtxcvqp"qp"y g'i tqy yj "qh'uqo g"Dgto wf c'tgg'h"  
kuj gu0"Proceedings of the Gulf and Caribbean Fisheries Institute."9:328/3340"  
"  
Dctfcej . "L0G0"E0N0Uo kj . "cpf "F 0Y 0O gpl gr03; 7: 0Dgto wf c'hkuj gt lgu'tgugctej "r tqi tco "hkpcn'tgr qt v0Dgto wf c"  
Vtcf g" F gxgnr o gpv"Dqctf. "J co knqp."7; "r 0"  
"  
Dctvj qmqo gy . "C0"cpf "L0C0Dqj pucen042270C"tgxkgy "qh'ecvej /cpf /tgrgcug" cpi r kpi "o qtvrk { "y kj "ko r rlecvcqp"htq"  
pq/vcng'tgugtxgu0"Reviews in Fish Biology and Fisheries."15:34; 6376"  
"  
Dgcwo ctkei g. "F 0U"cpf "N0I 0Dwmqen03; 980Dkqmi lecn'tgugctej "qp"upcr r gtu"cpf "i tqwr gtu"cu'tgrcvgf "vq"hkuj gt { "  
o cpci go gpv'tgs wktgo gpw."r 0: 8/"; 6."in:"J 0F0Dwruku."I r 0cpf "C0E0Lqpgu"\*gf u0t: "Rtqeggf kpi u'qh'y g"  
Eqmqv wkw "qp"upcr r gt/i tqwr gt "hkuj gt { "t guqwtegu"qh'y g"y guvtp"egpvcn" C vrpvle"Qegcp.0Hrc0Ugc" I tcpv"  
Eqn0Tgr 0390555"r r 0"  
"  
Dggwu."L03; ; 80Vj g"ghgevu"qh'hkuj kpi "cpf "hkuj "vcr u'qp"hkuj "cuugo drci gu'y kj kp"Xkti kp"Krcpf u"P cvkpcn"Rctm'icpf "  
Dwem"Krcpf "Tgg'h" P cvkpcn"O qpwo gpv0C"vgej plecn'tgr qtv'htq"y j g" WU0F gr ctvo gpv'qh'k'p'vgtkqt. "P cvkpcn"  
Rctm'Ugtxleg0"Vgej plecn" Tgr qtv"XKR"R"7I; 8043"r 0"  
"  
Dggwu" L0"cpf "C0Hkgrf rcpf gt03; ; ; 0Gxcnvcvqp"qh'c"eqpugtxcvqp"utcvgi { <c"ur cy plpi "ci i tgi cvkqp"enquwtg"htq"tgf "  
j kpf . "Epinephelus guttatus."lp"y j g" WU0Xkti kp"Krcpf u0"Environmental Biology of Fishes,"55:; 3/; : 0"  
"  
Dggwu."L0"cpf "O 0C0J kzqp03; ; 60F kvtkdvwkqp."r gtu kvvpeg."cpf "i tqy yj "qh'i tqwr gtu"\*Rkuegu-<Ugttcplf cg+qp"ct v'k'lecn"  
cpf "pcwctn'cvej "tgg'h"lp"y j g"Xkti kp"Krcpf u0"Bulletin of Marine Science."55:692/"6: 50"  
"  
"  
"; : "  
"  
"  
"  
"

Dggv. 'L0'cpf 'E0Tqi gtu042220F gerlpg'qh'hkuj gt { 'tguqwtegu'lp'o ctkpg'r tqvgevf 'ctgcu'lp'yj g'WUOXkxi kp'Krcpf u'Vj g' pggf 'hqt' 'o ctkpg'tgugt xgu' *Proceedings of the International Coral Reef Symposium* 09(1):66; /6760

Vj g'Dgrk g'Ur cy plpi 'Ci i tgi cvkqp'Y qtnkpi 'I tqwr 0422; 0kphqto cvkqp'Ektewrt 'P q'90Lxpg'422; . '4r r 0htqo " j wr <ly y y ur ci dgrk g0ti INpnEren0ur zAkrgvengv?9f qDqgO 3RTm 5f( vcdk?4: 29( npi wci g?gp/WU' tgvlgxgf "qp'37'F gego dgt'42330

Vj g'Dgrk g'Ur cy plpi 'Ci i tgi cvkqp'Y qtnkpi 'I tqwr 042340kphqto cvkqp'Ektewrt 'P q'320P qxgo dgt'4234. '4r r 0htqo " j wr <ly y y ur ci dgrk g0ti IRqtvcu15 IRF HuUr Ci aP gy urgwta32ahkpcrf f h'tgvlgxgf "qp'37'Cr tki'42350

D3/4 mg. 'L0G0'cpf 'E0' 0Ej cr nk03; 8: 0*Fishes of the Bahamas and Adjacent Tropical Waters* 0Nkxkpi uqpp'RwdroEq0" Y {ppgy qqf . 'RC.'993'r r 0'

Dqj pucem'L0C03; ; 0Rtqvgevkqp'qh'i tqwr gt'ur cy plpi 'ci i tgi cvkqp'0Eqcucn'Tguqwtegu'F kxkukqp'Eqpvt0P q0: : / : / 28.' : 'r 0'

Dqj pucem'L0C03; ; 20Drcen'cpf 'P cuucw'i tqwr gt' hkuj gt { 'vtgpf u0Crr gpf lz 'lp'Uqwj 'Cvcpvle'Tggthkuj 'Rrcp" F gxnqr o gpv'Vgco 'Tgr qt'v'qh'yj g'Upcrr r g/I tqwr gt 'Cuuguuo gpv'qh'yj g'Uqwj 'Cvcpvle'Hkuj gt { 'O cpci go gpv' Eqwpekn'3: 'r 0'

Dqj pucem'L0C042250Uj knkpi 'dcugnkpgu.'o ctkpg'tgugt xgu.'cpf 'Ngqr qrf a'dk'vle"gvj le0*Gulf and Caribbean Research* 14(2):3/90

Dqj pucem'L0C0'cpf 'UR0Dppgtq03; ; 80C'uv'vqpet { 'xkucn'legpuwa'vgej pls wg'hqt's wcpvkc'v'xgn' 'cuuguulpi " eqo o wpkv' 'utvewtg'qh'eqtcn'tggh'hkuj gu0P QCC'Vgej 0Tgr 0P O HU'63037'r 0'

Dqj pucem'L0C0'cpf 'F 0G0J ctr gt03; ; 0*Length-weight relationships of selected marine reef fishes from the southeastern United States and the Caribbean.* 'P QCC'Vgej 0O go q0P O HU'UGHE/437053'r r 0'

Dqj pucem'L0C0'F 0N0Uwj gtr'cpf . 'C0Dty p. 'F 0G0J ctr gt. 'cpf 'F 0D00 eEngmcp03; ; 80Cp'cpcn' uku'qh'yj g'Ectkddgcp" dkqucv'k'v'k'ecnf c'v'cd'cug'hqt'3; ; 70E'qcu0T'gu0F gx0Tgr 0EHO E'Eqpvt0P q0ETF/: 8t: 9/32.'57'r 0'

Dqrf gp. 'UM03; ; 60C'uvo o ct { 'qh'dkqmi k'ecn'cpf 'hkuj gt { 'f'cv'qp'tgf 'j' lpf '(*Epinephelus guttatus*) cpf "eqpg{ " \**Cephalopholis fulva* uqemu'lp'yj g'WUOXkxi kp'Krcpf u0P QCC IP O HU'O kco k'Ncdqtcvqt { 'Eqpvtkd0P q0' ; 5t: 6/54.'55'r 0'

Dqrf gp. 'UM042220Nqpi /f k'v'peg'o qxgo gpv'qh'c'P cuucw'i tqwr gt '\**Epinephelus striatus*+v'ur cy plpi 'ci i tgi cvkqp'lp" vj g'egpvt'cn'Dcj co cu0*Fishery Bulletin*, 98(3):864/8670

Dqrf gp. 'UM042230Wukpi 'Wintcuuple'Vrgo gvt { 'v'F gvgto kpg'J qo g'Tcpi g'qh'c'Eqt'cn'Tgg'h'hkuj 0in: 'L0F0Ukdt'v'cpf " L0N0P k'ng'p'v'gf u0: 'Rtq'eg'gf lpi u'qh'yj g'U' o r qukw' "qp'Vci i lpi "cpf "Vt'cen'kpi 'O ctkpg'Hkuj 'y kj 'Grg'v'v'q'ple" F g'x'legu. 'Hgd't'wct { '4222. 'J cy c'k'v'Ur t'kpi gt 'Rwd'rkuj lpi -0T g'x'lg'y u'<O gy qf u'cpf "Vgej p'q'q'ni l'gu'lp'Hkuj " Dk'q'q'ni { 'cpf 'Hkuj gt'ku'3-389/3: : 0'

Dqrf q/O qi vgn' M00 042290Ect'cev'g'tk' cel'p" { 'f'guet'kr'el'p'r' gus wgtc'f'gr'lu'kk'q'f'g'ci t'wr'cel'p'f'g'r'gegu'0'Gn'd'nc'ps'v'k' c'nl'6 U'c'p'v'c'lw'k'c'ö. 'S'w'p'v'c'p'c' 'T'q'q. "O 2 z'leq'<'k'p'x'k'g't'p'q'4226/Rt'k'lo' c'x'g't'c'42270'V'g'u'k'u' 'O' c'g'u't'v'c'0'G'E'Q'U'W'T.'; 2'r r 0'

Dqgo j qy gt. 'L0'0 0Tqo gtq. 'L0R'qu'c'f'c. 'U0M'q'd'c'tc. 'cpf 'Y 0J g { o cp0422: 0k'gp'v'k'ec'v'k'qp'qh't'ggh'hkuj 'ur cy plpi " ci i tgi cvkqp'uk'gu'lp'Nqu'T'qs wgu'Ctej k'rg'nci q'P cvk'q'p'cn'R'ct'm'X'g'p'g'l w'gr'0*Proceedings of the Gulf and Caribbean Fisheries Institute.* 60:77; /7870

Dqgo j qy gt. 'L0'0 0Tqo gtq. 'L0R'qu'c'f'c. 'U0M'q'd'c'tc. 'cpf 'Y 0J g { o cp042320Rt'gf'k'v'k'qp'cpf 'x'g't'k'k'ec'v'k'qp'qh'r'qu'k'd'rg" t'ggh'hkuj 'ur cy plpi 'ci i tgi cvkqp'uk'gu'lp'Nqu'T'qs wgu'Ctej k'rg'nci q'P cvk'q'p'cn'R'ct'm'X'g'p'g'l w'gr'0*Journal of Fish Biology* 77: 44/: 620

; ; "

"  
"  
"  
"

"

Dqvej qp.'E0'R0Rqt vkmq.'I 0Dqvej qp/P cxctq.'O 0Nqkw.'R0J qgvlgv.'M0F g'O g{gt.'F 0O cetcg.'J 0Cto utqpi .'X0' F cvcf kp.'U0J ctf kpi .'L0O cmgrc.'T0Rctnkpuqp.'L'Y 0Xcp'Dqej qxg.'F 0Nkt o cp.'L'J gtrcp.'C0Dcngt.'N0' Eqmf q'cpf'UE0'Kcce0'422: 0Ucwu'qh'EqtcrnTggh'T guqwtegu'qh'v'j g'Nguugt' Cpvkngu<Vj g'Hi gpej 'Y guv' kpf lgu.'Vj g'P gvj gtrcpf u' Cpvkngu.'Cpi vkmr.'Cpvi wc.'I tgpfc c.'Vtkpkf cf'cpf'Vqdc q'0'k<Y kmkpuqp.'E0' \*gf 00Ucwu'qh'EqtcrnTggh'qh'v'j g'Y qtrf <422: 0I mqdcn'EqtcrnTggh'O qpkqtkpi 'P gvj qtrncpf 'Tggh'cpf " Tckphqtgv'Tgugctej 'Egpvgt.'Vqy puxkng.'C wwtcrkc0r 487/4: 20'

"

Dqz.'U0'cpf'Dqpkrc'O gl'c0'422: 0Gn'gucf q'f g'rc'eqpugt xcelp'{"gZR nqvelep'f gr'O gtq'P cuucw'gp'rc'Equc' Cvf pvek' f g'J qpf wcu'Lvpg'422: . 'VP E'Tgr qtv.'r r 073"

"

DTGGH'Dej co cu'Tggh'Gpxtkppo gpv'Gf wecvkqpcn'Hqwpf cvkqp+03; ; 0Hkuj gtkgu'O cpci go gpv'Cevkqp'Rrcp'ht'v'j g' Dej co cu'0Tgr qtv'v'q'j g'Dej co cu'F gr ctvo gpv'qh'Hkuj gtkgu'Vj g'Dej co cu'Tggh'Gpxtkppo gpv'Gf wecvkqpcn' Hqwpf cvkqp'cpf'O ceCkrngt'Gnkqw'cpf'Rctvgtu'Nvf 0"

"

Dtqy p.'X042290Vj g'r qrte{o cnkpi 'r tqeguu'cpf'v'j g'r qrkku'qh'o cpci kpi 'v'j g'P cuucw'i tqwr gt'cpf'qvj gt'o ctkgp' tguqwtegu'0Proceedings of the Gulf and Caribbean Fisheries Institute.'60:827/8280'

"

Dtqy pgm'Y 0'0'cpf'Y 0G0Tckpg{03; 930Tgugctej'cpf'f gxnrc o gpv'qh'f ggr'y cvgt'eqo o gtekn'cpf'ur qtv'Hkuj gtkgu' ctqwpf'v'j g'Xki k'Krcpf u'r rcvgcw0Xki k'Krcpf u'Gcaqni lecn'Tgugctej'Ucvkqp'Eqpvtkd0P q05.'; : 'r r 0'

"

Dt{cpv.'F 0'N0Dwtng.'L0Y 0O eO cpwu.'cpf'O 0Ur crf kpi 03; ; 0Tggh'cv'Tkum'c'O cr/dcugf'kpf lecvt'qh'Rqvgpvcn' Vj tgcvu'v'j g'Y qtrf au'EqtcrnTgghu'0Y qtrf 'Tguqwtegu'kpuvkwg.'Y cuj kpi vq.'F 0E078'r r 0'

"

Dvej cp.'M042220Vj g'Dej co cu'Marine Pollution Bulletin.'41(1-6);: 6/3330'

"

Dwguc.'T0L03; : 90I tqy v'j 'tcvg'qh'tqr lecn'f go gtucn'Hkuj gu'Marine Ecology Progress Series'36:3; 3/3; ; 0'

"

Dwtpgw/J gtngv.'L03; 970Eqpvtkdwkqp'v'j g'dkqni {"qh'v'j g'tgf'j kpf.'Epinephelus guttatus.'c'eqo o gtekn'f " ko r qtvcv'v'gtcpkf'Hkuj 'htqo 'v'j g'tqr lecn'y gungtp'Cwv'p'0Rj F 'T kuugt'cvkqp.'W00 kco k'Eqtcrn'I cdrgu.'376'r 0'

"

Dwtpgw/J gtngv.'Lco gu.'cpf'Lqj p'C0Dctpgu03; ; 80Dcpkpi 'v'j g'wug'qh'r qvu'cpf'qvj gt'o cpci go gpv'kvtqf weg'f'kp" Dgto wf c'v'q'r tqgev'f gerkpi 't'ggh'Hkuj 'urqemu'0Proceedings of the Gulf and Caribbean Fisheries Institute' 44:45; /4780'

"

Dwuj .'R0 0'Gdcpnu.'I 0E0'cpf'Ncpg.'GF 03; ; 80Xcnf cvkqp'qh'v'j g'ci gkpi 'v'gej pls vg'ht'v'j g'P cuucw'i tqwr gt' \*Epinephelus striatus+kp'v'j g'Ec{o cp'Krcpf u.'r 0372/37: . 'in:'H0Cttgi v'kp/Ucpej gl . 'L0N00 wptq.'O 0E0' Dcn qu.'cpf'F 0Rcw'f' \*gf u0'Biology, fisheries and culture of tropical groupers and snappers'0'ENCTO " Eqph0Rtqe06: . '66; r r 0'

"

Dwuj .'R0 0'F 0G0Ncpg.'I 0E0'Gdcpnu/Rgtkg.'M0Nwng.'D0Lqj puqp.'E00 eEq{.'L0Dqj y gm'cpf.'G0Rctv'p'042280' Vj g'P cuucw'i tqwr gt'ur cy plpi 'ci i tgi cvkqp'Hkuj gt{'qh'v'j g'Ec{o cp'Krcpf u'o'cp'j kvtlecn'cpf'o cpci go gpv' r gtr gev'xg'0Proceedings of the Gulf and Caribbean Fisheries Institute.'57:737/7460'

"

Ecteco q.'T0'lt0422: 0P cvkqpcn'Tgr qtv'qh'v'j g'P cuucw'I tqwr gt'\*Epinephelus striatus+'Hkuj gt{'qh'Dgrk g0Dgrk g' Hkuj gtkgu'F gr ctvo gpv.'O kpkut {'qh'Ci tlewnwtg'(' Hkuj gtkgu.'Qevqdg'422: 0'

"

Ectngv'p'Tc{'I 0'0'0 0O eEqto lemTc{'E0C0Nc{o cp.'cpf'D0T0Uknko cp042220Investigations of Nassau grouper breeding aggregations at High Cay, Andros: implication for a conservation strategy'0'Hkpcn'tgr qtv'0Vj g' F gr ctvo gpv'qh'Hkuj gtkgu.'O kpkut {'qh'Ci tlewnwtg'cpf'Hkuj gtkgu.'P cuucw.'Vj g'Dej co cu'

"

Ectvgt.'L03; : 800 qppri j v'o cvkpi 'qh'v'j g'o wnkwf gu'0Animal Kingdom Magazine'89(6):85/930'

"

Ectvgt.'L03; : : 0I tqwr gt'o cvkpi 'tkwcn'q'p'c'Ectkldgcp't'ggh'0Underwater Naturalist'17:: /330'

"  
"  
"  
"  
"

"

Ectvgt.'103; : ; 01 tqwr gt'ugz'lp'Dgrk g0Natural History.'Qev'82/'8; 0

"

Ectvgt.'10'I 000 cttqy .'cpf'X0Rt {qt03; ; 60Cur gewu'qh'yj g'geqmi { "cpf 'tgr tqf wevqp'qh'P cuucw'i tqwr gt."  
*Epinephelus striatus*.'qh'yj g'eqcu'qh'Dgrk g.'Egvt'cn'Co gteco0Proceedings of the Gulf and Caribbean  
 Fisheries Institute, 43:8763330

"

Egtxki »p.'H03; 880Nqu'Rgegu'O ct'kpcu'f g'Xgpgl wgrc0Xqu0Kc'p'f'K0Hwpf 0Nc'Ucng'Elgpe0P c'v0'; 73'r 0'

"

Egtxki »p.'H03; ; 60Los peces marinas de Venezuela, Fundacion Cientifica Los Roques.'E0Hwpf cel8p'Elgpe'Whec'rcu'  
 Tqs wgu'gf 0e.'Ectcecu'Xgpgl wgrc.'4pf 'gf k'k'p'Xq'5.'4; 7'r 0'

"

EHO E' \*Ectkddgcp'Hkuj gt { 'O cpci go gpv'Eqwpekn03; ; 70Hkuj gt { 'O cpci go gpv'Rrcp.'Hkpcn'Gp'xk'qpo gpv'cn'K6 r cev'  
 Ucvgo gpv.'cpf 'F tch'Tgi wrcvt { 'K6 r cev'Tg'x'g'y . 'hqt' 'y' g'Uj cmqy /y cvgt'Tggh'Hkuj 'Hkuj gt { 'qh'Rwgtvq'Tleq'  
 cpf 'y' g'WUOX'kti lp'Krc'p'f' u.'326'r 0'

"

EHO E' \*Ectkddgcp'Hkuj gt { 'O cpci go gpv'Eqwpekn03; ; 50Co gpf o gpv'4'vq'yj g'Hkuj gt { 'O cpci go gpv'Rrcp' hqt' 'y' g'  
 Uj cmqy /y cvgt'Tggh'Hkuj 'Hkuj gt { 'qh'Rwgtvq'Tleq'cpf 'y' g'WUOX'kti lp'Krc'p'f' u04; 'r 0'

"

EHO E' \*Ectkddgcp'Hkuj gt { 'O cpci go gpv'Eqwpekn03; ; 80Tgi wrcvt { 'Co gpf o gpv'vq'yj g'Hkuj gt { 'O cpci go gpv'Rrcp'  
 hqt' 'y' g'Tggh'Hkuj 'Hkuj gt { 'qh'Rwgtvq'Tleq'cpf 'y' g'Wp'k'g'f' 'Ucv'gu'X'kti lp'Krc'p'f' u'Eq'p'eg't'p'k'p' 'T'g'f' 'J' k'p'f' "  
 Ur cy p'k'p' 'Ci i tgi cv'k'p' 'En'qu'w'gu'k'p'cn'f' k'p' 'c' 'Tgi wrcvt { 'K6 r cev'Tg'x'g'y 'cpf 'cp' 'Gp'x'k'qpo gpv'cn'Cu'gu'uo gpv0'  
 Cwi w'w'0j [wr <ly y <ectkddgcp'ho e'eqo IUECP P GF' 42HO RUITGGH/HKJ ' 42HO RQ vo "](#)

"

Ej gwpi . 'Y 0Y 0N0' [ 0Ucf qx { . 'O 0V0Dtc {pgp.'cpf 'N0 0I kv'g'pu042350Ct'g'yj g'rcu'v'tgo cl'k'p'k'p' 'P cuucw'i tqwr gt'  
 \*Epinephelus striatus'+hkuj g't'k'gu'w'w'c'k'p'c'd'g'v'j g'ecug'lp' 'y' g'Dcj co cu0Endangered Species Research."  
 20:49/5; 0

"

Ej kcr r qpg.'O 0'T0U'w'nc.'cpf 'M0U'w'k'x'cp' 'Ugcrg { 042220I tqwr gtu' \*R'lu'egu' <Ugt'c'p'k'f' cg+'lp' 'hkuj gf 'cpf 'r tq'v'g'v'g'f' 'ct'g'cu'  
 qh'yj g'Hq'k'f' c'Mg' {u.'Dcj co cu'cpf 'p'q't'yj g'tp' 'Ectkddgcp0Marine Ecology Progress Series. "198:483/4940"

"

Erntm' O 0T0' Q'0C'p'f' gtu'q'p.'T'0K'0E'j t'k' 'H'c'p'ek'u.'cpf 'F 0O 0V'iceg { 042220Vj g'g'h'g'ewu'qh'eqo o gte'k'cn'g'zr m'k'c'v'k'p'  
 qp'q't'c'p' g't'q'w'j { \*Hoplostethus atlanticus+'H'q'o 'y' g'eq'p'k'p'gp'v'cn'ur' g'qh'yj g'Ej cvj co 'T'k'ug.'P gy '\ g'nc'p'f' ."  
 H'q'o '3; 9; 'v'q'3; ; 90Fisheries Research. "45:439645: 0"

"

Erntq.'T0'c'p'f' 'M'0E'0N'p'f' go cp'042250Ur cy p'k'p' 'ci i tgi cv'k'p' 'u'k'gu'q'h'up'c'r r gt' 'cpf' 'i' tqwr gt' 'ur' g'el'gu' \*N'w'c'p'k'f' cg' 'cpf' "  
 Ugt'c'p'k'f' cg+'q'p' 'y' g'k'p'uw'rc't' 'uj' g'h'q'h' 'E'w'd'c'0Gulf and Caribbean Research"14(2); ; 3/3280

"

Erntq.'T0'I cte'f'c' /Eci k'f' g.'C0'U'g't'tc.'N00'I cte'f'c' /C't'g'c'i c.'10R03; ; 20Ect'c'v'g't' 'j'w'le'cu'd'k'q'n'i leq'r gus wgt'cu'f' g'rc'  
 ej g't'p'c' 'e't'k'q'nc' \*Epinephelus striatus \*D'rq'ej +' \*R'lu'egu' <Ugt'c'p'k'f' cg+'gp' 'rc' 'r' rc'w'ch'q'to c' 'e'w'd'c'p'c'0Cienc. Biol045-<  
 456640

"

Erntq.'T0'M'0E'0N'p'f' go cp.'cpf 'N0I'0R'ct'g'p'v'042240Ecology of the Marine Fishes of Cuba0Y cuj k'p' i v'q'p.'F E-<  
 Uo k'j u'q'p'k'p' 'k'p'w'k'w'k'p' 'R't'g'u'0'

"

Erntq.'T0'L'0C'0D'c'k'ut'g.'M'0E'0N'p'f' go cp.'10R0I cte'f'c' /C't'g'c'i c'042230Ew'd'c'p' 'hkuj g't'k'gu' <j' k'v'q't'k'ec'n'v't'g'p'f' u' 'cpf' 'e'w't't'g'p'v'  
 u'c'w'u.'r 03; 6/43; . 'in:'T0'Erntq.'M'0E'0N'p'f' go cp.'N0I'0R'ct'g'p'v' \*g'f' u0'Ecology of the Marine Fishes of Cuba0'  
 Uo k'j u'q'p'k'p' 'k'p'w'k'w'k'p' 'R't'g'u.'Y cuj k'p' i v'q'p.'F 0E0475'r 0'

"

Erntq.'T0' [ 0Ucf qx { 'f'g'O ke'j g'gu'p.'M'0E'0N'p'f' go cp.'C0I'0I cte'f'c' /Eci k'f' g'0422; 0J k'v'q't'k'ec'n'c'p'c'n' 'u'k' 'q'h' 'E'w'd'c'p'  
 eqo o gte'k'cn' 'hkuj k'p' 'g'h'q't'v' 'cpf' 'y' g'g'h'g'ewu'qh' 'o cpci go gpv' 'k'p'v'g't'x'g'p'v'k'p'u'q'p' 'k'o r q't'c'p'v't'g'g'h' 'hkuj g'u' 'H'q'o '3; 826  
 42270Fisheries Research 99(1):9/38"

"

Ernt {f' q'p.'1042260Ur cy p'k'p' 'ci i tgi cv'k'p'u'q'h'eq't'c'n't'g'g'h' 'hkuj g'u' <ej c't'c'v'g't'k'w'leu.'j { r q'y' g'uku.'y' t'g'c'u' 'cpf' 'o cpci go gpv0'



Ewuj kqp. 'P 00' cpf 'M0Uwnkxcp/Ugcrg{042290Ncpf kpi u. 'ghqt v' cpf 'uqekq/geqppo leu'qh'c'uo cml'uecrg'eqo o gtekn'  
Hkuj gt { 'kp'vj g'Dcj co cu0Proceedings of the Gulf and Caribbean Fisheries Institute **60**:384/3880'  
"

Fcj ni tgp'E0R03; ; 0Population dynamics of early juvenile Nassau grouper: an integrated modeling and field study0  
Rj (F 0f kuugtwkqp0P qt vj 'Ectqrkpc'Ucvg'Wpkxgtuks{0Tcrgk j . 'P E'WUC0367'r r 0'  
"

'Fcj ni tgp'E0R0' cpf 'F 0D0Gi i nguq042220Geqmi lecnRtqeguugu'Wpf gtn{ kpi 'Qpvqi gpgvle'J cdkcv'Uj kku'lp'c'Eqtcn'  
Tggh'Hkuj 0Ecology'**81(8)**:4449/44620'  
"

Fcj ni tgp'E0R0' cpf 'F 0D0Gi i nguq042230Ur cvkq/vgo r qtcn'xctkcdkks{ 'lp'cdwpcpeg.'uk'g' cpf 'o letqj cdkcv'  
cuuqekcvkpu'qh'gctn{ 'lwxgpkrg'P cuucw'i tqwr gt 'Epinephelus striatus'lp'cp'qh'tggh'pwugt { 'u'vgo 0Marine  
Ecology Progress Series. '**217**:367/3780'  
"

Fcj ni tgp'gv'cn' ]kp'r tgr \_"  
"

Fgr ctvo gpv'qh'O ctkgT'Guqwtegu042290Vj g'Dcj co cu0Cr tkr'42290Dtkghkpi 'Fqewo gpvu.'77r r 0'  
"

F P GT '\*Rwgtvq'Tleq'F gr ctvo gpv'qh'P cwten' cpf 'Gpxkqpo gpcnT'Guqwtegu-042260Rwgtvq'Tleq'Hkuj kpi 'Tgi wrcvqpu0'  
P a898: 0Uwr rigo gpcwt { 'Higt0j wr <ly y y f tpc0 qdkgtpqt t l''  
"

F R P T '\*F kxkukq'qh'Rncppkpi ' cpf 'P cwtenT'Guqwtegu.'WU'Xki lp'Kncpf u-042270Ego o gtekn( 'Tgetcvkqpcn'Hkuj gt a'  
kphqto cvkq' 'Dqqmgv0F gr ctvo gpv'qh'Rncppkpi ' cpf 'P cwtenT'Guqwtegu.'F kxkukq'qh'Hkuj '( 'Y kf rktg' cpf "  
F kxkukq'qh'Gpxkqpo gpcn'Gphqtego gpv0Lwn{ '42270j [wr <ly y y 0kHkuj cpf y kf rktg@qo 0'](#)  
"

Fqo gkg. 'O 0N0' cpf 'R0N0Eqk03; ; 90Vtqr lecn'tggh'Hkuj 'ur cy plki 'ci i tgi cvkqpu'f ghkgf' cpf 'tgxky gf 0Bulletin of  
Marine Science. '**60(3)**:8; ; /9480'  
"

Gi i nguq'F 0D03; ; 70Tgetwko gpv'lp'P cuucw'i tqwr gt 'Epinephelus striatus' r quv'ugwgo gpv'cdwpcpeg.'o letqj cdkcv'  
hgcwtgu' cpf 'qpvqi gpgvle'j cdkcv'uj kku0Marine Ecology Progress Series. '**124**:: /440'  
"

Gi i nguq. 'F 0D0' L0L0I tqxgt. ' cpf 'T0P 0Nk' ekw03; ; 0Qpvqi gpgvle'f lgv'uj kku'lp'P cuucw'i tqwr gt <tqr j le' hknai gu' cpf "  
r tgf cvqt { 'lo r cev0Bulletin of Marine Science. '**63(1)**:333/3480'  
"

Gj tj ctf v. 'P 00' cpf 'X0M0Y 0F grxgcwz042290Vj g'Dcj co cu'P cuucw'i tqwr gt '\*Epinephelus striatus'+Hkuj gt { ' /'y q'  
cuuguo gpv'o gj qf u'cr r rkgf 'v'c'f'cvc'/'f ghkgpv'eqcucn'r qr wrcvqpu0Fisheries Research, **87**:39/490'  
"

Gnku.'U0'I 0Xkrc.' cpf 'Y 0Q0Y cvpcdg03; ; 80I tqy vj ' cpf 'hggf 'wkrk' cvkq'qh'j cvj gt { /tgetgf 'lwxgpkrg'P cuucw'  
i tqwr gt 'hgf 'hqt'r tcevecn'f lgu0Progressive Fish-Culturist. '**58(3)**:89/3940'  
"

Gnku.'U0E0'Y 0Q0Y cvpcdg. ' cpf 'G0R0' Gnku03; ; 9c0Vgo r gtcwt'g'ghgeu'qp' hggf 'wkrk' cvkq' cpf 'i tqy vj 'qh'r quv'  
ugwgo gpv'uci g'P cuucw'i tqwr gt 0Transactions of the American Fisheries Society. '**126(2)**:52; /'5370'  
"

Gnku.'G0R0'Y 0Q0Y cvpcdg. 'UE0' Gnku.'L0I lpq l c. ' cpf 'C00 qtly cng03; ; 9d0Ghgeu'qh'wtdwngpeg.'ucrkpkv{ . ' cpf 'hki j v'  
kpv'gpkv{ 'qp'j cvj kpi 'tcvg' cpf 'uwxkcn'qh'rtxcn'P cuucw'i tqwr gt. 'Epinephelus striatus0Journal of Applied  
Aquaculture, '**7(3)**:55/650'  
"

Gtf o cp. 'F 0U03; 980Ur cy plki 'r cvgtpu'qh'Hkuj gu'htqo 'vj g'pqt vj gcvgtp'Ectkddgcp0Eqpvtkd0Ci tle0Rgus 0'F gr v0'  
Ci tle0Ego o qpy gcnj 'qh'Rwgtvq'Tleq. ': 3/590'  
"

Gxgto cpp. 'D0Y 03; 220Fishes and Fisheries of Porto Rico0WU0Ego o kukq'qp'Hkuj ' cpf 'Hkuj gt lgu.'572'r 0'  
"

HCQ0422; 0P cvkqpcn'Hkuj gt { 'Ugevqt'Qxgtxky 0Vj g'Ego o qpy gcnj 'qh'vj g'Dcj co cu0Hkuj gt { ' cpf 'Cs wcewmtg'  
Eqwpt { 'Rtqkrgu.'HCQ.'HkF IERIDJ U0c { '422; =: 'r r 0'  
"

Hlpg.'L0E03; ; 20I tqwr gtu'lp'Nqyg<Ur cy plpi 'ci i tgi cvkpu'qh'P cuucw'i tqwr gtu'lp'J qpf wcu0'Sea Frontiers.'r '64/670'  
 "
 Hlpg'L0E03; ; 40I tggf { 'hqt'I tqwr gtu'0'Wildlife Conservation'0'0 c{ 'Lwpg'3; ; 4-3/70'  
 "
 Hlpg{.'M0C0'E0E0c0f cy . 'G0J l0ngtuq0'42290"Gulf of Mexico Science'0'Hktuv'eqphko gf 'tgeqt'f'qh'P cuucw'i tqwr gt'  
*Epinephelus striatus* \*Ruegu<Ugttcplf cg+'lp'y g'Hujy gt'I ctf gp'Dcpmi'P cvkpcni'0 ctkpg'Ucpewct { 0'Rr 0384/  
 3870  
 Hlpgug.'T0c0f 'F 0'Rcwn{ 0'42320'Hkuj dcug0'y y 0'Hkuj dcug0'qti 0'  
 "
 I ctekc/O qrlpgt.'I 0'c0f 'l 0'Ucf qx { 0'422: 0'Vj g'ecug'hqt'tgi kqpcni'0 cpci go gpv'qh'y g'P cuucw'i tqwr gt.'*Epinephelus  
 striatus*0'Proceedings of the Gulf and Caribbean Fisheries Institute.'60:798/8240"  
 "
 I ctf pgt.'V0C0'K0 0'E½² . 'L0C0I km'C0I t0cpv'c0f 'C0F0Y cvkpuq0'42250'Nqpi /vgt0 'tgi kq/y kf g'f'genkpu'lp'  
 Ectkddgcp'eqtcnu'0'Science.'301:; 7: /; 820'  
 "
 I cueqki pg.'L042240'Nassau Grouper and Queen Conch in the Bahamas. Status and Management Options'  
 O ceCkngt'Gnkq'w'c0f 'Rctvpgtu'N0'hqt'Vj g'Dej co cu'Tggh'Gpxk'qpo gpv'Gf wecvkpcni'Hqwpf cvkq'0'DTGGH0'  
 66'r r 0'  
 "
 I kduqp.'L0'T'0'Rq'w'I 0'Rc| . 'K0 cln'P 0'Tgs wpc0'42290'Gzr gtlqpegu'qh'y g'Dgrk' g'ur cy plpi 'ci i tgi cvkpu'y qtnkpi "  
 i tqwr 0'Proceedings of the Gulf and Caribbean Fisheries Institute.'59:677/6840"  
 "
 I kduqp.'L0422: 0'0 cpci lpi 'c'P cuucw'I tqwr gt'Hkuj gt { '0'C'Ecug'Uwf { 'Itqo 'Dgrk'g0'Proceedings of the Gulf and  
 Caribbean Fisheries Institute.'60:825/8260'  
 "
 I O HO E'¶I wh'qh'O gzleq'Hkuj gt { 'O cpci go gpv'Eqwpekn'03; ; 0'Co gpf o gpv'P wo dgt'3'vq'y g'Hkuj gt { 'O cpci go gpv'  
 Rrcp'hqt'y g'Tggh'Hkuj 'Hkuj gt { 'qh'y g'I wh'qh'O gzleq'0'Vco r c.'Hqtkf c0578'r 0'  
 "
 I qqf { gct.'E0R03; ; 50'Ur cy plpi 'uvqen'ldkqo cuu'r gt'tgetwk'lp'Hkuj gt'ku'o cpci go gpv'hqwpf cvkq'c0f 'ewttgpv'wug.'r 0'  
 89/' 3.'in:'U0U0 kj . 'L0L0J wpx'c0f 'F 0'Tkctf '¶gf u0t.'Tkn'gxcn'cvkq'c0f 'dkmqi k0cni'tghgt'peg'r qlpw'hqt'  
 Hkuj gtlgu'o cpci go gpv.'Ecp0'Ur ge0'Rwdri'Hkuj 0'Cs wcv0'Uek0'3420'  
 "
 I qtgcw.'V0L03; ; 40'Dngej lpi 'c0f 'tgghe'qo o wplv { 'ej cpi g'lp'Lco clec<3: 73/3; ; 30'American Zoologist,'32:8: 5/8; 70'  
 "
 I tggpy qqf . 'E0D03; ; 30'F kntkdwkq'c0f 'hggf lpi 'j cdku'qh'ictxeni'Gr kgr j grkpg'i tqwr gtu'lp'Gzwo c'Uqwpf . 'Dej co cu'  
 O U'y guku.'Hqtkf c'Kpukwq'qh'Vgej pqmqi { . 'O gndqwt'pg.'HN.'83'r r 0'  
 "
 I tqxgt.'L0L03; ; 50'Vtqr j le'geqmqi { 'qh'r grci le'gctn{/lw'gpkrg'P cuucw'i tqwr gt.'*Epinephelus striatus*.'f wtkpi 'cp'gctn'  
 r j cug'qh'tgetvko gpv'lpw'f go gtucl'j cdkc'w0'Bulletin of Marine Science.'53:3339/33470'  
 "
 I tqxgt.'L0L03; ; 60'Hggf lpi 'j cdku'qh'gctn{/lw'gpkrg'P cuucw'i tqwr gt'0'Bahamas Journal of Science'2(1):44/480'  
 "
 I tqxgt.'L0L0'F 0'D0Gi i nguq'p.'c0f 'L0 0'Uj gpngt'03; ; : 0'Vt0pukq'p'Itqo 'r grci le'vq'f go gtucl'r j cug'lp'gctn{/lw'gpkrg'  
 P cuucw'i tqwr gt.'*Epinephelus striatus*<Rki o gpvc'kq'p.'us wco cvkq'p.'c0f 'qpqi gp { 'qh'f kg'0'Bulletin of Marine  
 Science.'62(1); ; 9/3350'  
 "
 I wketv'O c0f c { . 'F 0'c0f 'H0Lw' tgl /Hgt'p0f gl 03; 880'F gucttqmq'go dtkq'pctkq' { 'r tko gtqu'guwf kqu'ictxergu'f g'ic'  
 ej gtpc'etkqmc.'*Epinephelus striatus*'\*Dmgej +\*Rgtelkqto gu<Ugttcplf cg+0'Cecf go le'Ekgpeku'f g'Ewdc.'  
 Kpukwq'f g'Qegcpmqi k0c0'Nc'J cdcpc'1:57/670'  
 "
 I wpygt.'I 0'c0f 'N0M'p0crr 03; 730'Hkuj gu.'pgy . 'tctg'qt'ugrf qo 'tgeqt'f gf 'Itqo 'y g'Vgzcu'eqcu'0'Texas Journal of  
 Science,'3(1):356/35: 0'  
 "

"  
 "  
 "  
 "  
 "



"

J qtubqtf. "K0422; 0Vj g'I tqwr gt'Hkuj gt { 'qh' Cpki wc' cpf 'Dctdwf c0Tgr qtv'qp' Tgi kqpcn' Y qtmij qr 'qh' P cuucw' I tqwr gt. " Ectwci gpc. 'Eqmqo dlc. '42/43' Qevqdg' 422: . 't' gxlugf 'Lcpwct { '422; . '34' r' r' 0

"

J wi j gu. "VR03; ; 60Ecvustqr j gu. 'r' j cug/uj kku. " cpf 'rcti g/uecrg' f gi tcf cvkqp' qh' c' Ectkddgcp' eqtcnt' ggh' Science " 265:3769/37730

"

J wo cpp. 'R0' cpf 'P 0F gNqcej 042240 Reef Fish Identification -- Florida-Caribbean-Bahamas \*5<sup>th</sup> 'Gf kxqp' 0P gy " Y qtrf 'Rvdrkcvkqpu. 'Kpe0Lcemuqpxkng. 'Hqt' k' c' 0"

"

KWEP 0422504225 " KWEP 'Tgf 'Nku' qh' Vj tgcvgpgf 'Ur gelgu' 0y y y (kweptgf rku' Qti 0F qy pnycf gf 'qp' 3: 'P qxgo dgt' 42250

"

Lqj puqp. 'F 0' cpf 'Mggpgt. 'R03; ; 60Ck' 'v' k' g' p' w' h' e' c' v' k' p' qh' 'C' o' g' t' k' e' c' p' i' tqwr gt 'rctxcg' 0 Bulletin of Marine Science. " 34(1):328/3560

"

Lqpgu. 'I 0R0' UORcpgu. 'U0' 0Vj qttqrf 042270 Eqtent' ggh' Hkuj 'rctxcg' ugwrg' emug' 'v' j' qo g' 0 " Current Biology " 15(14):3536/353: 0

"

Lqt { . 'F 0G0' cpf 'G00' k' g' t' u' g' 03; ; 0' Ur gelgu' r' tqh' k' g' u' < 'h' g' j' k' u' q' t' k' u' c' p' f' 'g' p' x' k' q' p' o' g' p' w' c' i' t' s' w' k' t' g' o' g' p' w' 'q' h' 'e' q' c' u' w' c' i' h' k' u' j' g' u' " c' p' f' 'p' x' g' t' v' d' t' e' v' u' \*' u' q' w' j' 'H' q' t' k' c' + // 'd' r' e' m' t' g' f' . 'c' p' f' 'P' c' u' u' c' w' i' t' q' w' r' g' t' u' 0' W' L' 0' H' k' u' j' 'Y' k' f' r' 0' U' g' t' x' 0' D' k' i' r' 0' T' g' r' 0' : 4\*330B32+0WU0Cto { 'Eqtr u' qh' Gpi kpggtu. 'VT' GN: 4/6043' r' r' 0

"

Mcf kuqp. 'G0' T' 0' 0' P' go g' j' . 'L0' D' r' p' f' g' c' w' 'V0' U' o' k' j' . 'c' p' f' 'L0' E' c' r' p' c' p' 042320 P cuucw' i' tqwr gt \* Epinephelus striatus + 'lp' U' 0' Vj qo cu. 'WU' X' k' i' k' p' 'K' u' r' p' f' u' . 'y' k' j' 'g' x' k' f' g' p' e' g' 'h' q' t' 'c' 'u' r' c' y' p' k' p' i' 'c' i' i' t' g' i' c' v' k' q' p' 'u' k' s' g' t' g' e' q' x' g' t' { 0 Proceedings of the Gulf and Caribbean Fisheries Institute. " 62:495/49; 0

"

Mcwo cp. 'N0' cpf 'G0T' qo g' t' q' 042330 Hkuj g' t' k' u' 'd' c' u' g' f' 'q' p' 'D' g' r' k' g' c' p' 'd' k' q' f' k' x' g' t' u' k' v' { < 'y' j' { 'v' j' g' { } g' u' q' 'x' w' p' g' t' c' d' r' g' 'v' q' 'q' h' u' j' q' t' g' " q' k' n' r' 0357/363. " in: 'O' N' F' 0' R' c' m' q' o' c' t' g' u' 'c' p' f' 'F' 0' R' c' w' n' f' \*' g' f' u' 0: 'Too Precious to Drill: the Marine Biodiversity of Belize' 0 Hkuj g' t' k' u' 'E' g' p' t' g' 'T' g' u' g' t' e' j' 'T' g' r' q' t' u' 3; \*8+0 Hkuj g' t' k' u' 'E' g' p' t' g' . 'W' p' k' x' g' t' u' k' v' { 'q' h' 'D' t' k' u' j' 'E' q' n' w' o' d' l' c' ] KUP " 33; : /8949\_0

"

Mmgg { . 'E' 0' F' 0' 'C' 0' 0' q' t' k' y' c' n' g' . 'I' 0' 0' k' f' c' o' q' v' . 'X' 0' P' l' e' q' n' 'c' p' f' 'Y' 0' Y' c' v' e' p' c' d' g' 03; ; 60Vj g' w' u' g' 'q' h' 'N' J' T' J' /c' h' q' t' 'l' p' f' w' e' g' f' " u' r' c' y' p' k' p' i' 'q' h' 'h' x' g' 'f' k' h' g' t' g' p' v' u' r' g' e' l' g' u' 'q' h' 'o' c' t' k' p' g' 'v' r' g' q' u' w' 'h' u' j' g' u' 0' C' d' u' s' t' c' e' v' h' q' t' 'Y' q' t' r' f' 'C' s' w' c' e' w' n' w' t' g' ; 60Y' q' t' r' f' " C' s' w' c' e' w' n' w' t' 'U' q' e' k' v' { 0P gy " Q' t' r' g' e' p' u' 0' N' C' 0' L' c' p' w' c' t' { '36/3: . '3; ; 60369' r' 0

"

Mgmkuqp. 'V0' U' 0' J' g' r' r' g' m' 'D0' U' g' o' g' p' u' . 'P' 0' G' j' t' j' c' t' f' v' . 'K0' \ k' p' n' 0422; 0T' g' x' l' g' y' 'q' h' 'F' c' v' e' d' c' u' g' u' 'C' x' c' k' e' d' r' g' 'c' p' f' 'V' t' g' p' f' u' 'l' p' " C' d' w' p' f' c' p' e' g' 'T' g' r' c' v' k' p' i' 'v' q' 'P' c' u' u' c' w' i' t' q' w' r' g' t' 'Epinephelus striatus' l' p' 'W' p' k' s' g' f' 'U' c' v' g' u' 'c' p' f' 'W' L' 0' V' g' t' t' k' q' t' k' e' n' 'Y' c' v' g' t' u' 'k' p' e' n' f' k' p' i' 'T' g' e' q' o' o' g' p' f' c' v' k' p' u' 'h' q' t' 'c' 'H' q' t' o' c' n' 'U' c' w' u' 'T' g' x' l' g' y' 0' 'Y' q' t' m' i' t' q' w' r' 'T' g' r' q' t' 036' r' r' 0

"

Mgpf cm. 'C0Y' 0' L' t' 03; 9; 00' q' t' r' j' q' m' i' k' e' c' n' l' e' q' o' r' c' t' k' u' q' p' u' 'q' h' 'P' q' t' v' j' 'C' o' g' t' k' e' c' p' 'u' g' c' 'd' c' u' u' 'r' c' t' x' c' g' \*' R' k' u' e' g' u' < 'U' g' t' c' p' k' f' c' g' +P' QCC " V' g' e' j' 0T' g' r' 0P' O' H' U' E' k' e' 064: . '72' r' 0

"

Mgpf cm. 'C0Y' 0' L' t' . 'c' p' f' 'O' (R0' H' c' j' c' { 03; 9; 0N' c' t' x' c' 'q' h' 'v' j' g' 'u' g' t' t' c' p' k' f' 'h' u' j' 'Gonioplectrus hispanus' y' k' j' 'e' q' o' o' g' p' u' 'q' p' 'k' u' " t' g' r' c' v' k' p' u' j' k' r' u' 0 Bulletin of Marine Science. " 29:339/3430

"

Mqdetc. 'U0422; 0T' g' i' k' q' p' c' n' 'C' p' c' n' f' u' k' u' 'q' h' 'U' g' c' h' q' q' t' 'E' j' c' t' e' v' g' t' k' u' k' e' u' 'c' v' 'T' g' g' h' 'H' k' u' j' " U' r' c' y' p' k' p' i' 'C' i' i' t' g' i' c' v' k' q' p' 'U' k' s' g' u' 'l' p' 'v' j' g' " E' c' t' k' d' d' g' c' p' 0Rj (F 0F k' u' g' t' c' v' k' p' . 'V' g' z' c' u' C( O' 'W' p' k' x' g' t' u' k' v' \*' I' g' q' i' t' e' r' j' { +0

"

Mqdetc. 'U0' c' p' f' 'J' g' { o' c' p' . 'Y' (F 042290 E' q' o' r' c' t' c' v' x' g' i' g' q' o' q' t' r' j' k' e' 'c' p' c' n' f' u' k' u' 'q' h' 'P' c' u' u' c' w' i' t' q' w' r' g' t' 'u' r' c' y' p' k' p' i' 'c' i' i' t' g' i' c' v' k' q' p' " u' k' s' g' u' 'l' p' 'D' g' r' k' g' 'c' p' f' 'v' j' g' 'E' c' { o' c' p' 'K' u' r' c' p' f' u' 0 Proceedings of the Gulf and Caribbean Fisheries Institute. " 60:795/ 7980

"

Mqdetc. 'U0' c' p' f' 'J' g' { o' c' p' . 'Y' (F 042320 U' g' c' 'd' q' w' q' o' 'i' g' q' o' q' t' r' j' q' m' i' { 'q' h' 'o' w' n' k' u' r' g' e' l' g' u' 'u' r' c' y' p' k' p' i' 'c' i' i' t' g' i' c' v' k' q' p' 'u' k' s' g' u' 'l' p' " D' g' r' k' g' 0 Marine Ecology Progress Series. " 405:465/4760

"

Mqunqy . 'L0C0'H0J cprg{ . 'cpf 'T0Y k0mwpf03; ; 0Ghgevu'qh'hkuj kpi 'qp'tgg'hkuj 'eqo o wpkkgu'cv'Rgf tq'Dcpn'icpf 'Rqtv' Tq{cni'Ec{u. 'Lco clec0Marine Ecology Progress Series'43:423/4340

"

Mqunqy . 'L0C0'M0Ckngp. 'U0Cwkn'cpf 'C0Ergo gpwup03; ; 60Ecevej 'cpf 'ghqtv'cpcn{uku'qh'yj g'tgg'hkuj gtlgu'qhi'Lco clec' cpf 'Dgrk g0Fishery Bulletin'92:959/9690

"

Mqunqy . 'L0C0'I 0Y 0Dqgj rgtv. 'L0F 00 0I qtf qp. 'T0N0J cgf tlej . 'R0Nqtcepeg. 'cpf 'P 0Rctk042220Eqpvkpgpwn'urq g'cpf " f ggr /ugc'hkuj gtlgu'ko r nlec'vkpu'ht'c'ht ci krg'gequ{ uvg0 ICES Journal of Marine Science. '57:76: 67790"

"

Mico gt. 'R042250U{pyj guku'qh'eqtcn'tgg'hj gcnj 'kpf lecvqtu'ht'vj g'y guvgp' Cwv'pke'lt guwmu'qh'yj g'CI TTC" Rtqi tco o g'3; ; 964222-0Atoll Research Bulletin, '496:367: 0

"

Nc'I qteg. 'L0Qrxkt' \*gf 003; 5; 0The Book of Fishes0P cvkqpcn'I gqi tcr j le'Uqekgv. 'Y cuj kpi vqp. 'F 0E0'589'r 0

"

Ncpf uo cp. 'U0L0'E0Lcf qv. 'O 0Cuj rg{ . 'cpf 'L0C0D0Erc{ f qp0422; 0Kpxgukl cvkqp'qh'yj g'P cuucw'i tqwr gt' \*Epinephelus striatus+hkuj gt { 'kp'yj g'Vwmu'cpf 'Eclequ'Kuc'p'f u'ko r nlec'vkpu'ht'eqpugt'xcv'kqp'cpf 'o cpci go gpv0" Proceedings of the Gulf and Caribbean Fisheries Institute. '61:: 4/; ; 0

"

Ngku. 'L00 0; ; 80Nctxcn'f gxgrqr o gpv'kp'iqw'ur gekgu'qh'kpf q/Rcelhe'eqtcn'tqww'Plectropomus' \*Rluegu'Ugttc'pkf cg< Gr kpg'j g'kpcg'ly kj 'cp'cpcn{uku'qh'yj g'tgr'cvkpuj k' u'qh'yj g'i gpw0Bulletin Marine Science. '38(3):747/7740

"

Ngku. 'L00 0; ; 90Tgxkgy 'qh'yj g'gctn{ 'hkg'j kvqt { 'qh'vtqr lecn'i tqwr gtu' \*Ugttc'pkf cg+'cpf 'upcr r gtu' \*Nwlc'pkf cg+'r 03: ; /" 45: . 'in: 'L0L0Rqrxkpc'cpf 'U0Tcnvqp' \*gf u0: 'Tropical Snappers and Groupers: Biology and Fisheries Management'0Y guxkgy 'Rtguu0Dqwf gt0

"

Nqpi rg{ . 'Y 0 03; 390Uwf lgu'wr qp'vj g'dkqmi lecn'uki p'htec'peg'qh'c'pko c'neqmt'cvkqp0K0Vj g'eqmtu'cpf 'eqmt'ej cpi gu' qh'Y gu'kpf kcp'tgg'hkuj gu0Journal of Experimental Zoology. '23(3):755/8230

"

Nwenj wtu. 'D0G03; ; 80Vtgp'f u'kp'eqo o gtekn'hkuj gt { 'rcpf kpi u'qhi' tqwr gtu'cpf 'upcr r gtu'kp'Dgto wf c'htqo "3; 97"vq" 3; ; 4'cpf 'cuuqek'cv'f 'hkuj gt { 'o cpci go gpv'kuwgu. 'r "4: 8/4; 9. 'in: 'H0Cttgi wp/Ucpej gl . 'L0N00 wptq. 'O 0E0' Dcni qu. 'cpf 'F 0Rcwn{ \*gf u00Biology, Fisheries and Culture of Tropical Groupers and Snappers. 'KENCTO " Pq06: . '66; 'rr 0"

"

Nwenj wtu. 'D0G042250F gxgrqr o gpv'qh'c' Ectkddgcp'tgi kqpcn'eqpugt'xcv'kqp'utcvgi { 'ht'tgg'hkuj 'ur cy plpi " ci i tgi cvkpu. Proceedings of the Gulf and Caribbean Fisheries Institute. '54:88: /89: 0"

"

Nwenj wtu. 'D0G042260E wttgp'v'ucwmu'qh'eqpugt'xcv'kqp'cpf 'o cpci go gpv'qht'gg'hkuj 'ur cy plpi 'ci i tgi cvkpu'kp'yj g' I tgevt'Ectkddgcp0Proceedings of the Gulf and Caribbean Fisheries Institute. '55:752/7640"

"

Nwenj wtu. 'D0G042270Gxcn'cvkqp'qh'hkuj gtlgu'o cpci go gpv'cpf 'eqpugt'xcv'kqp'o gcuwt'gu'cnngp'v'q'r tqgcv'i tqwr gt' ur cy plpi 'ci i tgi cvkpu'kp'yj g'y kf gt'Ectkddgcp<Ecug'uwf lgu'qh'Dgto wf c. 'Dgrk g'cpf 'Ec{o cp'Kuc'p'f u0 Proceedings of the Gulf and Caribbean Fisheries Institute. '58:48: /48; 0

"

Nwenj wtu. 'D0G0'cpf 'L0C0Y ctf 03; ; 80Cpcn{uku'qh'vtgp'f u'kp'Dgto wf c)'Hkuj gt { 'Uc'v'w'lecn'F cvdcug'htqo "3; 97"vq" 3; ; 2. 'y kj 'tghgt'peg'v'q'hkuj gt { 'o cpci go gpv'o gcuwt'gu'ko r ngo gpv'gf 'f wtkpi 'vj ku'r gtlqf 0Proceedings of the Gulf and Caribbean Fisheries Institute. '44:528/5460

"

O cj qp. 'T03; ; 20Fishery Management Options for Lesser Antilles Countries (Antigua and Barbuda, Barbados, Dominica, Granada, St. Kitts and Nevis, St. Lucia, St. Vincent and the Grenadines). 'HCQ'Vgej 0Rcr 05350 HCQ. 'Tqo g0

"

O cj qp. 'T03; ; 50Nguugt' C'p'v'kgu0Rctv'K'r 03; ; . 'in: 'Marine Fishery Resources of the Antilles: Lesser Antilles, Puerto Rico and Hispaniola, Jamaica, Cuba0HCQ'Vgej 0Rcr 0548. '457'r r 0

"  
"  
"  
"

"

O crlnqkxk . 'C0'V0G0'xcp'Nggwy gp.'cpf 'UP 0'Eq'xg'422: 0'Tggh'Uk'gu<'Rt'gf'cv'kqp'qp'vj'g'lp'xcuk'g'tgf'rk'p'hkuj . 'Pterois volitans'\*Rkuegu<'Ueqtr'cgpk'cg+.'d{ 'pc'v'xg'i' tqwr'gtu'lp'vj'g'Dcj'co'cu'0'Coral Reefs, '27:723"

"

O cpvgt. 'J 0'Y 03; 690'Vj'g'f'ki'gpg'vle'v'tgo'cv'qf'gu'qh'o'ct'k'p'g'hkuj'gu'qh'V'qt'wi'cu.'H'rt'kf'c'0'American Midland Naturalist.' '38(2):479/6380'

"

O cteqi'rl'gug.'F'10'42230'K' r'rl'ec'v'k'pu'qh'er'ko'cv'g'ej'cpi'g'hqt'r'ct'cuk'kuo'qh'c'p'ko'cnu'lp'vj'g'cs'w'v'le'"  
g'p'x'k'q'po'g'p'0'E'c'p'f'k'p' 'L'q'w't'p'c'n'q'h' \ 'q'q'm'i' { '9; \* : +<3553/35740'

"

O cvqu'E'ct'c'd'cm'q.'F'0422: 0'N'gu'u'q'pu'h'g'c't'p'g'f' 'h't'q'o' 'R'w'g't'v'q' 'T'le'q'u' 'e'q'o' o' g't'ek'n'h'kuj'g't' { '3; : : /422: 0'P'ro'c'ee'd'ings'of'the' Gulf and Caribbean Fisheries Institute, '61:345/34; 0'

"

O eE'rc'p'c'j'cp.'V'0'P'0'R'q'n'w'p'k'p.'c'p'f' 'V'0'F'q'p'g'0'42240'G'eq'm'i' l'ec'n'i'v'c'u'g'u'c'p'f' 'v'j'g't'g'u'k'g'p'eg'q'h'e'q't'c'n't'g'gh'0' 'C'onservation Ecology'6(2):3: 0'

"

O g'f'k'p'c/S'w'g'l.'C'0'C'0'F'0'J' g't't'g'c/R'c'x'q'p.'T' 'R'q'q'v'N'q'r'g'l' . 'G'0'U'q'c/E'q't'f'g't'q.'M'0'D'q'r'k'q/O'q'i'w'g'n'c'p'f' 'Y'0'J' c'f'c'f'0'42260'C' " r't'g'k'o'k'p'c't' { 'u'w't'x'g' { 'q'h'v'j'g'p'c'u'c'w'i' tqwr'gt' 'E'p'ine'p'h'e'l'u's' s't'r'i'a't'u's' u'r'c'y'p'l'k'p'i' 'c'i' i' t'g'i'c'v'k'q'p'c'v'0'G'i'D'r'c'p's'w'k'c'r'o'b'lp'"  
v'j'g' 'u'q'w'j' 'e'q'c'u'v'q'h'i'S'w'k'p'w'c'p'c' 'T'q'q.'O'g'z'k'eq'0'P'ro'c'ee'd'ings'of'the' Gulf and Caribbean Fisheries Institute.' " '55:779/78; 0'

"

O q't'c.'E'0'c'p'f' 'R'H'0'U'c'g'0'42240'C't'g'r'q'r'w'c'v'k'p'pu'q'h'e'q't'c'n't'g'gh'h'kuj' "q'r'g'p'q't' 'e'm'q'ug'f' A'T'r'e'n'd's' i'n' E'c'o'l'o'g'y' & E'v'o'l'u't'i'o'n'" '17(9):644/64: 0'

"

O q't't'k'u.'L'0'0'c'p'f' 'L'0'N'0'c'n'k'p'u'0'422; 0'H'g'g'f'k'p'i' "g'eq'm'i' { 'q'h'lp'xcuk'g'rk'p'hkuj' '\*Pterois volitans'+lp'vj'g'Dcj'co'k'p'"  
c't'ej'k'r'g'r'c'i'q'0' 'E'n'v'ir'o'n'm'e'n't'a'l' B'io'l'o'g'y'of'F'is'h'es, '86:5: ; /5; : 0'

"

O q'w't'c.'T'0'N'0'42230'U'g't't'c'p'k'f'c'g'0'In.'R'0'c'0'D'w'e'n'w'r' "c'p'f' 'P'0'c'0'0'g'p'g'l'g'u' \*g'f' u'0'E'c'v' m'i'q' 'f'c'u'G'ur'g'el'g'u'f'g'R'g'k'z'g'u'0'c't'k'p'j'q'u' "  
f'q' 'D't'c'uk'0'0'w'ug'w'f'g' \ 'q'q'm'i'k'c' 'W'p'k'x'g't'ul'f'c'f'g'f'g'U'q' 'R'c'w'q'"

"

O w'o'd' { . 'R'0'0' 'C'0'F'0'J' c't'd'q't'p'g.'c'p'f' 'F'0'F'0'D't'w'o' d'c'w'i'j' 042330I' tqwr'gt' 'cu'c' 'p'c'w't'c'n'd'k'q'eq'p't'q'n'i'q'h'lp'xcuk'g'rk'p'hkuj' 0'P'L'0'S' 'O'N'E' 8\*8+g'437320f'q'k'3208593' l'q'w't'p'c'f'r'q'p'g'0'2437320"

"

O w'o'd' { . 'R'0'0' 'E'0'R'0'F'c'j'ni't'g'p.'C'0'F'0'J' c't'd'q't'p'g.'E'0'K'0'M'c'r'r'g'n' 'H'0'0' l'ej'g'r'k' 'F'0'F'0'D't'w'o' d'c'w'i'j' . 'M'0'G'0'J' q'm'g'u.'L'0'0' 'O'g'p'f'g'u.'M'0'D't'q'c'f' . 'L'P'0'U'c'p'ej'k'l'eq.'M'0'D'w'ej' . 'U'0'D'q'z.' 'T'0'Y'0'U'q'h'g' "c'p'f' 'C'0'D'0'I' k'n'0'42280'H'kuj'k'p'i' . 'v't'q'r'j'k'e'"  
e'c'ue'c'f'g'u.'c'p'f' 'v'j'g'r' 't'q'eg'u'u'q'h'i' t'c'l'k'p'i' "q'p' 'e'q't'c'n't'g'gh'0'S'c'i'e'n'c'e' "311:; : 323"

"

O w'o'd' { . 'R'0'0' 'T'0'U'0'U'g'p'g'em' 'C'0'0'G'f'y'c't'f'u.'T'0'F'g't't'c't'k' 'T'0'E'q'r'g'o'c'p.'C'0'F'0'J' c't'd'q't'p'g.'c'p'f' 'L'0'R'0'I' k'd'u'q'p'0'42340'H'kuj'k'p'i' "  
f'q'y'p'c' 'E'c't'k'd'd'g'c'p' 'h'q'q'f' 'y'g'd' 't'g'm'z'g'u'v't'q'r'j'k'e' 'e'c'ue'c'f'g'u'0'M'a'r'i'ne' E'c'o'l'o'g'y' P'r'o'g'r'e's's' S'e'r'i'e's' "445:35/460"

"

O w'p'f'c' { . 'R'0'N'0' 'I' 'R'L'q'p'g'u.'O' 'L'0'R't'c'v'ej'g'w.'C'0'0'Y' k'r'k'c'o' u'0422: 'E'r'ko'c'v'g'ej'cpi'g'c'p'f' 'v'j'g'h'w'w't'g'h'q't' 'e'q't'c'n't'g'gh'h'kuj'g'u'0' "  
F'is'h'and'F'is'h'e'r'i'e's.' "9(3):483/4: 70"

"

O w'p't'q.'L'0'N'03; : 5e0Ej'cr'vgt'3<'E'q't'c'n't'g'gh'h'kuj' "c'p'f' 'h'kuj'g't'k'g'u'q'h'v'j'g' 'E'c't'k'd'd'g'c'p' 'U'g'c.'r'03/; . 'in,'L'0'N'0' w'p't'q' \*g'f'0:' "  
E'c't'k'd'd'g'c'p' 'E'q't'c'n't'g'gh'h'kuj'g't' { "T'g'u'q'w't'eg'u'<'K'EN'c'T'O' 'U'w'f'k'g'u'c'p'f' 'T'g'x'k'g'y' u.'X'q'r'0'90'k'p'v'g't'p'c'v'k'p'c'n'E'g'p'v'g't' 'h'q't' "  
N'k'k'p'i' 'c'p'f' 'C's'w'v'le' 'T'g'u'q'w't'eg'u' 'O'c'p'c'i'g'o'g'p'v' 'O'c'p'k'r.'R'j'k'r'k'r'k'p'g'u'0'E'q'p't'k'd'034704p'f' 'g'f'0'498'r'0'

"

O w'p't'q.'L'0'N'03; : 5d0Ej'cr'vgt'6<'V'j'g' 'e'q'o' r'q'u'k'k'q'p'c'p'f' "o'c'i'p'k'w'f'g'q'h'h'p'g' 'e'c'v'ej'g'u'lp' 'L'c'o'c'l'ec'p' 'y'c'v'g't'u.'r'048/54'in:'L'0'N'0' "  
O'w'p't'q' \*g'f'0:' 'E'c't'k'd'd'g'c'p' 'E'q't'c'n't'g'gh'h'kuj'g't' { "T'g'u'q'w't'eg'u'<'K'EN'c'T'O' 'U'w'f'k'g'u'c'p'f' 'T'g'x'k'g'y' u.'X'q'r'0'90' "  
k'p'v'g't'p'c'v'k'p'c'n'E'g'p'v'g't' 'h'q't' 'N'k'k'p'i' 'c'p'f' 'C's'w'v'le' 'T'g'u'q'w't'eg'u' 'O'c'p'c'i'g'o'g'p'v' 'O'c'p'k'r.'R'j'k'r'k'r'k'p'g'u'0'E'q'p't'k'd'034704p'f' "  
g'f'0'498'r'0'

"

O w'p't'q.'L'0'N'03; : 5e0Ej'cr'vgt'7<'V'j'g' 'e'q'o' r'q'u'k'k'q'p'c'p'f' "o'c'i'p'k'w'f'g'q'h'v't'c'r' 'e'c'v'ej'g'u'lp' 'L'c'o'c'l'ec'p' 'y'c'v'g't'u.'r'055/6; . 'in:'L'0'N'0' "  
O'w'p't'q' \*g'f'0:' 'E'c't'k'd'd'g'c'p' 'E'q't'c'n't'g'gh'h'kuj'g't' { "T'g'u'q'w't'eg'u'<'K'EN'c'T'O' 'U'w'f'k'g'u'c'p'f' 'T'g'x'k'g'y' u.'X'q'r'0'90'



"

Qxgtutggv."T00 03; 8; 0F ki gpgvle'tgo cvqf gu'qh'o ctkpg'vrgquv'hkuj gu'htqo 'Dkuec{pg'Dc{.'Hqtkf c0Tulane Studies in Zoology and Botany."15(4):33; /3970

"

Rcvpi km'Ugo o gpu.'E0K0cpf "D0Z0Ugo o gpu042250Vj g'ucwuw'qh'tggh'hkuj gu'lp'y g'Ec{o cp'Kucpf u'\*DY K0Ucwu'qh' eqtcn'tgghu'lp'y g'Y guvtp'Cvcpvle<T guwuu'qh'lp'kkn'uwtxg{.'Cvcpvle'I wrh'Tcr kf "Tggh'Cuuguo gpv' \*CI TTC+Rtqi tco 0Atoll Research Bulletin."498:448/469."

"

Rcwgtuq."J 00'0'U0'0Vj qttqrf.'cpf 'L00'0'Uj gpnng03; ; ; 0Cpcn{uku'qh'q'v'rkj 'ej go kv{ 'lp'P cuucw'i tqwr gt " \*Epinephelus striatus+htqo 'y g'Dcj co cu'cpf 'Dgrk g'wukpi 'uqnwkp/ducgf "KER/O U0Coral Reefs."18:393/39: 0

"

Rcl.'I 0'cpf 'I tko uj cy . "V04223c0Ucwu'tgr qt'v'qp'P cuucw'i tqwr gt'htq'Dgrk g.'egp'tcn'Co gtlec0Uelgpv'kle'tgr qt'v'qh' y g'I tggp'Tggh'Gpxkqpo gpvcn'k'pukswg0Ucp'Rgf tq'Vqy p.'Co dgti tku'Ec{g.'Dgrk g"

"

Rcl.'I 0'cpf 'I tko uj cy . "V04223d0Ucwu'Tgr qt'v'qp'P cuucw'i tqwr gt'Ci i tgi cvkpu'lp'Dgrk g.'Egpt'cn'Co gtlec0' Proceedings of the First National Workshop on the Status of Nassau Groupers in Belize: Working Towards Sustainable Management.'cv'Dgrk g'Ek{.'52'Lwn'4223.'I tggp'Tggh'Gpxkqpo gpvcn'k'pukswg'r 049/580"

"

Rcl.'I 00'cpf 'V0I tko uj cy 04223e0Nkg'J kvqt { 'Ej ctcevgt'kukc'('O cpci go gpv'Tgeqo o gp'cvkpu'htq'y g'P cuucw' I tqwr gt'Rqr wcvkp'lp'Dgrk g'0'Tgr qt'v'd{ 'I tggp'Tggh'Gpxkqpo gpvcn'k'pukswg0\*WP GRII GHI tcpv'P q0D\ G' ; : 'I 74/RTQ'25+ "

"

Rcl.'I 0'cpf 'G0Vtwn'042290Vj g'P cuucw'i tqwr gt'ur cy plpi 'ci i tgi cvkpp'cv'Ec{g'I rqt {.'Dgrk g<C"dtlgh'j kvqt {0C" ecug'uwf { 'd{ 'Vj g'P cwtg'Eqpugt'xpe{.'O guqco gtlecp'Tggh'Rtqi tco 086r r 0"

"

RFV'\*Rcp'F g'xgnr o gpv'Vgco +03; ; 2c0Uqwj 'Cvcpvle'tggh'hkuj <Rcp'F g'xgnr o gpv'Vgco 'tgr qt'v'q'y g'Uqwj " Cvcpvle'Huj gt{ 'O cpci go gpv'Eqwpekn'749'r 0

"

RFV'\*Rcp'F g'xgnr o gpv'Vgco +03; ; 2d0Vj g'r qv'p'kn'qh'o ctkpg'hkuj gt{ 't'gugt'xgu'htq'tggh'hkuj 'o cpci go gpv'lp'y g' WLUUqwj gtp'Cvcpvle0P QCC0P O HU.'Eqcucn'Tguq'wtegu'F k'kukp.'Eqpv'0P q0ETF l ; ; / 226063'r r 0

"

Rgnkpu0L003; : 50Vj g'Dgrk g'Dcttkt'Tggh'Gequ'f ugo <Cp'cuuguo gpv'qh'ku'tguq'wtegu.'eqpugt'xcv'kp'ucwuw'cpf " o cpci go gp'0P gy 'l qtnl'qqm'i kcn'Uqelgv' 'Tgr qt'v.'36: 'r 0

"

Rkwo cp.'U00'U0'0J kg.'E0HI 0Lgh'g{.'E0Ecnf qy . 'O 00M'gpf cm 'O 0G0O qpceq.'cpf '\ 0J knku/Ucctt0422: 0Hkuj " Cuugo dnci gu'cpf 'Dgpj le'J cdkcu'qh'Dweni'Kucpf 'Tggh'P cvkqpen'0 qpwo gpv'\*U0Etqkz.'WLUXki lp" Kucpf u'+cpf 'y g'Uwt'qwpf lpi 'Ugcuec g<C'Ej ctcevgt'k'cvkpp'qh'Ur cvkn'cpf 'Vgo r qtcn'Rcwgt'pu'0P QCC" Vgej plcn'0 go qtcpf wo 'P QUP'EEQU'930Ukxgt'Ur tkpi . 'O F 0; 8r r 0"

"

Rqixkpc.'L00'cpf "U0Tcnv'p'\*gf u+03; ; 90Tropical Snappers and Groupers: Biology and Fisheries Management0'r r 0 783/8250Y gux'ky 'Rtguu.'Dqwf gt.'EQ0

"

Rqnxkp.'P 0X'E0'cpf 'E00'0Tqdg'w03; ; 50I tgcvg't'dkqo cuu'cpf 'xcnw'g'qh'v'cti g'v'eqtcn'tggh'hkuj gu'lp'y q'uo cml' Ectkddgcp'o ctkpg'tgugt'xgu'0Marine Ecology Progress Series."100:389/3980

"

Rqy gm'CD0'cpf 'L0Y 0Vwengt'lt03; ; 40Gi i 'cpf 'h'etx'cnf g'xgnr o gpv'qh'h'edq'tc'v'qt{/tgctgf 'P cuucw'i tqwr gt." Epinephelus striatus"\*Ruegu<Ugtt'cp'k'cg+0Bulletin of Marine Science,"50(1):393/3: 70

"

Rtcf c.'O 0E0'I 0Rgpcm'j c.'U0Rq'uf c.'P 0J qy ctf.'R0J gttq.'N0Ucn'p'cu.'G0Ec'utq.'H0Ec'dg' cu.'cpf 'J 0Tq'd'lp'p'p'0' 42260Hkuj 'ur cy plpi 'ci i tgi cvkpu'lp'y g'Ucp'P'f tgu'C'tej k' g'nci q.'c'htu'v'cr r tqzko cvkpp'0Hkpcn'Tgr qt'v' Eqtcn'k'p'cpf 'Vj g'Qegcp'Eqpugt'xpe{.'72'r r 0

"

Tcf enq. 'F 0X0'CF 00 qvej gm' [ P 0Udknp. 'T0Ertq' O cf twi c. 'cpf 'C0Ukxc' Ngg03; 970Cegtec' f'rc' rpi kwf 'f'g' rqu'r gegu'eqo gtekrnu'gp'ecr wtcu'f'g'rc' ] qpc'pqtqeekf gpvni'f'g'Ewdc0Ugtkg'Qegcpqmi kec0P q04: 0Cecf go kc' f'g'Ekpcku'f'g'Ewdc0Kpukswq'f'g'Qegcpqmi kc0J cdcpc0Ewdc.'; 'r r 0'

Tcpf . 'R0'E0Vc { m. 'cpf 'F 0Gi i rnuvq042270C' xkf gq' o gj qf 'hqt' s wcpvkh' kpi 'uk' g' f' kvt kd wkqp. 'f' gpukv' { 'cpf 'y' tgg/ f' lo gpukqpcn'ur cvkn'ust wewt'g'qh'tgg'hkuj 'ur cy plpi 'ci i tgi cvkpu0Go gti kpi 'Vgej pqmi lgu' Cduwcew0' *Proceedings of the Gulf and Caribbean Fisheries Institute.* "56:64; /6520"

Tcpf cm 'LOG03; 840Vci i kpi 'tgg'hkuj gu'lp' y' g' Xkti lp' Krcpf u0' *Proceedings of the Gulf and Caribbean Fisheries Institute* "14:42364630"

Tcpf cm 'LOG03; 850Cff kkpncn'tgeqxtlgu'qh'vci i gf 'tgg'hkuj gu'ltqo 'y' g' Xkti lp' Krcpf u0' *Proceedings of the Gulf and Caribbean Fisheries Institute* "15:377/3790"

Tcpf cm 'LOG03; 870Hqf' j' cdku'qh' y' g' P cuucw'i tqwr gt \*'Epinephelus striatus'-0Cuucw0' Krcpf 'O ct0Ncdu' Ectkd08' y' " O gvkpi 0Lcp03; 87-35/ "380"

Tcpf cm 'LOG03; 890Hqf' j' cdku'qh'tgg'hkuj gu'qh' y' g' Y gu'kpf lgu'Uwv lgu'lp' Vtqr kecn'Qegcpqi tcr j { . 'O lco k5:8876 : 690"

Tcpf cm 'LOG03; : 50Ectkddgcp' T ggh' Hkuj gu'0Ugeqpf 'gf kkp0' VHD 0Rvdrlec' vkpu. 'P gr wpg' Ek' { 0P L0572' r 0'

Tcpf cm 'LOG0' cpf 'XG0Dtqen03; 820Qdugt'xc'vkpu'qp' y' g' geqmi { 'qh'gr kpg' j' g' nkg' 'cpf nwlcpk' 'hkuj gu'qh' y' g' Uqekv' " Krcpf u' y' kj 'go r j' cuku'qp' 'hqqf' j' cdku'0' *Transactions of the American Fisheries Society.* "89(1); /380"

Tc { 'I E0' O 0 eEqto kem' Tc { . 'E0C0Nc { o cp. 'cpf 'D0'0Ukno cp042220' kpxguki cvkpu'qh' P cuucw' I tqwr gt " Dtggf kpi 'Ci i tgi cvkpu'cv' J ki j' 'Ec { . 'Cpf tq' < K r rlec' vkpu' hqt' c' Eqpugt'xc' vkp' Utcvgi { 0T gr qt' v' y' g' " F gr ctvo gpv'qh' Hkuj gtlgu. 'P cuucw' 'Dcj co cu' "

Tggf . 'E0W03; 6300 ct' kpg' 'hkg' lp' "Vgzcu' y' cvgtu'0' *Texas Academy Sci. Publ. Nat. Hist.* "Xqr040: : 'r r 0'

TGGH042340' T ggh' Gpxktqpo gpvni' Gf wecvkq' Hqwpf cvkq' Xqnpv' ggt' Uwt' xg { 'Rtql' gev' F cvdcug' 0Y qtrf 'Y kf g' Y gd' " grgevt' qple' r' wdrlec' vkp' 0' y y 0' ggh' qti 0' f' cvg' qh' f' qy pncf < F gego dgt' 42340' "

Tgr qtv'qh' y' g' Eqo o kuuq' qh' kps wkt { 03; ; 30' Tgr qtv'qh' y' g' Eqo o kuuq' qh' kps wkt { "v' g' zco kpg' 'cpf' o' cng' " tgeqo o gp' f' cvkpu' hqt' y' g' hwwt' g' qh' y' g' 'hkuj kpi 'lp' wwt { 'cpf' hqt' y' g' hwwt' g' r' tqv' gev' qh' y' g' o' ct' kpg' " gpxktqpo gpv' lp' Dgto wf c0I qxgtpo gpv' qh' Dgto wf c0' Hgdtwct { . '3; ; 30' "

Tlej ctf u' Y 0' E0E0Dcrf y' lp. 'cpf 'C0T 3/4 ng042280Ej cr vgt' 33; 0Ugt' cpl' cg' < Ugc' d'cuugu. 'r 03447/3554' in: "Y 0' Tlej ctf u' \*gf 0: " *Early Stages of Atlantic Fishes: An Identification Guide for the Western Central North Atlantic.* "Xqr0' K0ETE 'Rtguu. 'Dqec' Tcvq. 'HN. "3557' r 0' "

Tqdg' tu. 'E00 0' 'cpf' P 0' XE0R' qnwpk03; ; 30' Ct' g' o' ct' kpg' t' gugt' xgu' g' h' g' evk' g' lp' o' cpci go gpv' qh' t' ggh' hkuj gtlgu' *Reviews in Fish Biology and Fisheries* 1:87; /30' "

Tqdg' tu. 'E00 0' P 0S wpp. 'L0Y 0' Vwengt' 'I' 0' 'cpf' R0P 0' Y qqf y' ctf 03; ; 70' k' v' tqf' we' vkq' qh' j' v' e' j' gt { /t' g' t' gf' 'P cuucw' " i tqwr gt' v' q' c' eqtcn' t' ggh' gpxktqpo gpv' 0' *North American Journal of Fisheries Management,* "15(1):37; /3860' "

Tqi gtu. 'E00' 'cpf' 'L0Dggw042230F gi tcf' cvkq' qh' o' ct' kpg' gequ' u' wgo u' 'cpf' 'f' gen' kpg' qh' 'hkuj gt { 't' gu' wtegu' lp' o' ct' kpg' " r tqv' gev' f' ct' gcu' lp' y' g' WU' Xkti lp' Krcpf u0' *Environmental Conservation,* "28(4):534/5440' "

Twf f . 'O 0C04225c0' k' p' ukw' k' qpcn' cpcn' uku' qh' o' ct' kpg' t' gugt' xgu' 'cpf' 'hkuj gtlgu' i' qxgt' p' cpeg' r' qile { "g' zr' g' t' lo' gpw' < c' ecug' " uwf { "qh' P cuucw'i tqwr gt' eqpugt' xc' vkp' lp' y' g' Vwt' mi' 'cpf' 'Eclequ' Krcpf u0' Rj 0' f' 0' y' guku' 0' Y ci gpki gp' " Wpkxgtuks { . 'Vj g' P g' j' gtr' p' f' u. "498r r 0' "

"  
"  
"  
"



Ucf qx { 'f g' O ke j gu pp. 'I 0' C0Eqt pluj . 'O 0F qo gkgt. 'R0Eqrkp. 'O 0T wuugm 'cpf 'M0Nkp f go cp0422: 0C' I m dcn'  
Dcu grkp g' hqt 'Ur cy plpi 'Ci i tgi cvkpu' qh' Tggh' Hkuj gu *Conservation Biology* **22(5):3455/34660**

UCHO E \*Uqwj 'Cvcpvle' Hkuj gt { 'O cpci go gpv'Eqwpekn03; ; 50Hkuj gt { 'O cpci go gpv'Rrcp. 'Tgi wrcvt { 'K6 r cev'  
Tgx lgy . 'cpf 'Hkpcn' Gpxktqpo gpvcn' K6 r cev' Ucvgo gpv' hqt 'y j g' 'Upcrr gt 'I tqwr gt 'Hkuj gt { 'qh' y j g' Uqwj 'Cvcpvle'  
Tgi kqp. '395' r 0

UCHO E \*Uqwj 'Cvcpvle' Hkuj gt { 'O cpci go gpv'Eqwpekn03; ; 20Co gpf o gpv'P wo dgt'4. 'Tgi wrcvt { 'K6 r cev' Tgx lgy . "  
Tgi wrcvt { 'Hgzldkrk' 'Cpcn' uku' 'cpf 'Gpxktqpo gpvcn' Cuuguo gpv' hqt 'Hkuj gt { 'O cpci go gpv' Rrcp' hqt 'y j g'  
Upcrr gt 'I tqwr gt 'Hkuj gt { 'qh' y j g' Uqwj 'Cvcpvle' Tgi kqp. '69' r r 0

Ucr. 'G0' cpf 'G0Dcmgugtqu042220' *Conservation status and dynamics of the Glover's Reef, Belize, spawning  
aggregation 0December 1999 - January 20000* Uelgpvle 'tgr qt v' y j g' 'Y kf rkg' Eqpugtxcvqp 'Uqelgv' u'  
I ruxgt u' Tggh' O ctkpg' Tgugctej 'Ucvkqp. 'Dgrk' g. 'Egpcn' Co gtlec0

Ucr. 'G0' G0Dcmgugtqu. 'cpf 'T00' Uctt042230' Tcr kf 'f gerkp' qh' P cuucw' i tqwr gt 'ur cy plpi 'ci i tgi cvkpu' k' Dgrk' g <  
Hkuj gt { 'o cpci go gpv' cpf 'eqpugtxcvqp' pggf u0' *Fisheries*. **26(10):45/520**

Uej @gt. 'O 0' O 0P go gj . 'cpf 'T0C' r r grf qqtp042290' Rcu' v' g' z' r' m' k' c' v' k' p' 'cpf 'r t' g' u' g' p' v' f' k' u' t' k' d' w' k' p' 'qh' P cuucw' i tqwr gt 'cv'  
O qpc' 'Krcpf . 'Rwgt' v' 'Tleq' 'r quvg' 'cpf 'cdut' cev' 0' *Proceedings of the Gulf and Caribbean Fisheries Institute*. "  
**60:8970**

Uej @gt. 'O 0' 0' V00' Tqy gm 'O 0' 0' P go gj . 'T00' C' r r grf qqtp042340' Uqwpf 'r tqf v' w' k' p' 'cuucw' i tqwr gt 'y kj 'tgr tqf v' w' x' g'  
dgi c' x' l' q' t' 'qh' P cuucw' i tqwr gt 'Epinephelus striatus' 'cv' ur cy plpi 'ci i tgi cvkpu' 0' *Endangered Species Research*. "  
**19:4; /5: 0**

Uej o kw. 'G0H' 'cpf 'M00' Uwnkcp03; ; 60Tgugctej 'crr' r' k' c' v' k' p' u' qh' 'x' q' n' w' p' v' g' g' t' i' g' p' g' t' c' v' g' f' 'e' q' t' c' n' t' g' g' h' 'h' k' u' j' 'u' w' x' g' u' 0' V' j' g'  
P cwt' g' Eqpugt' x' c' p' e' { 'cpf 'y j g' 'Wp' k' x' g' t' u' k' v' { 'qh' O' k' c' o' k' F' g' r' c' t' v' o' gpv' qh' D' k' q' m' i' { 'T' g' r' q' t' v' 0' E' q' t' c' n' I' c' d' r' g' u' 'H' q' t' k' f' c' 0'  
5: 'r r 0

UETHC \*4225- 'UETHC' P gy urgwgt 'P wo dgt '60F gego dgt '42250' Uqelgv' 'hqt' y j g' Eqpugt' xc' v' k' p' 'qh' T' g' g' h' 'Hkuj "  
Ci i tgi cvkpu' 0' y y 0' *ethc0qti* "

Ugf dgtt { 'I 0' 0' F 0' 0' U' g' x' g' p' u' q' p' . 'cpf 'T0Y' 0' E' j' c' r' o' c' p' 03; ; 80' U' q' e' n' i' k' f' g' p' v' h' e' c' v' k' p' 'k' p' 'r' q' v' p' v' c' m' { 'y j' t' g' c' v' p' g' f' 'ur' g' e' l' g' u' 'qh'  
i' t' q' w' r' g' t' 'V' g' r' q' u' v' g' k' < 'U' g' t' t' e' p' k' f' c' g' < 'G' r' k' p' g' r' j' g' r' k' p' c' g' + 'k' p' 'C' v' c' p' v' e' 'c' p' f' 'E' c' t' k' d' d' g' c' p' 'Y' c' v' g' t' u' 0' 'H' k' p' c' n' 'T' g' r' 0' 0' C' T' H' R' "  
I' t' e' p' v' p' q' 0' P' C' 69' H' H' 2' 3' 4' 0' U' q' w' j' 'E' c' t' q' i' k' p' c' F' g' r' v' 0' qh' P' c' w' t' c' n' T' g' u' q' w' t' e' g' u' . 'O' c' t' k' p' g' T' g' u' q' w' t' e' g' u' T' g' u' g' c' t' e' j' 'K' p' u' k' w' w' g' 0'  
73' r r 0

Ugo o gpu. 'D0Z0' R0D' wuj . 'U0J' g' r' g' m' 'D0L' qj' p' u' q' p' . 'E00' e' E' q' { . 'E0R' c' w' g' p' i' k' m' Ugo o gpu' 042340' C' p' "in situ' x' k' u' c' n' i' o' c' t' m' i'  
t' g' e' c' r' w' t' g' o' g' y' q' f' "v' q' 'c' u' u' g' u' 'y j g' 'c' d' w' p' f' c' p' e' g' 'q' h' 'ur' c' y' p' g' t' u' 'c' v' 'c' p' "ci i tgi cvkpu' 'u' k' g' 0' *Proceedings of the Gulf and  
Caribbean Fisheries Institute*. "**64:446/4480**

Ugo o gpu. 'D0Z0' M0G0N' wng. 'R0I' 0' D' wuj . 'E0R' c' w' g' p' i' k' m' Ugo o gpu. 'D0L' qj' p' u' q' p' . 'E00' e' E' q' { . 'cpf 'U0J' g' r' g' m' 042290'  
k' p' x' g' u' k' i' c' v' k' p' i' 'y j g' t' g' r' t' q' f' v' e' w' x' g' 'o' k' i' t' c' v' k' p' 'c' p' f' 'ur' c' v' c' n' i' g' e' q' r' m' i' { 'qh' P' cuucw' i tqwr gt 'Epinephelus striatus' + 'q' p'  
N' k' w' g' 'E' c' { o' c' p' 'K' r' c' p' f' 'w' u' k' p' i' 'c' e' q' w' u' k' e' 'c' i' u' 0' 'c' p' 'q' x' g' t' x' l' g' y' 0' *Proceedings of the Gulf and Caribbean Fisheries  
Institute*. "**58:3; 3/3; : 0**

Ugo o gpu. 'D0Z0' R0D' wuj . 'U0J' g' r' g' m' 'D0L' qj' p' u' q' p' . 'E00' e' E' q' { . 'E0R' c' w' g' p' i' k' m' Ugo o gpu. 'cpf 'N0Y' j' c' { r' g' p' 0422: c' 0'  
E' j' c' t' v' k' p' i' "c' e' q' w' t' u' g' 'h' q' t' P' cuucw' i tqwr gt 't' g' e' q' x' g' t' { 'k' p' 'y j g' 'E' c' t' k' d' d' g' c' p' < 'y j' c' v' y' g' o' x' g' 'h' g' c' t' p' g' f' 'c' p' f' 'y j' c' v' y' g' 'u' k' n' i' p' g' g' f'  
v' q' 'h' p' q' y' 0' *Proceedings of the Gulf and Caribbean Fisheries Institute*, **60:829/82; 0**"

Ugo o gpu. 'D0Z0' U0J' g' r' g' m' 'R0D' wuj . 'D0L' qj' p' u' q' p' . 'E00' e' E' q' { . 'E0R' c' w' g' p' i' k' m' Ugo o gpu. 'cpf 'N0Y' j' c' { r' g' p' 0422: d0C' p'  
k' p' x' t' c' / 'c' p' f' 'k' p' v' g' t' / 'c' p' p' v' c' n' i' c' p' c' n' i' u' k' u' 'q' h' P' cuucw' i tqwr gt 'u' k' g' t' k' u' t' k' d' w' k' p' u' 'h' t' q' o' 'c' 't' g' e' g' p' v' i' 'r' t' q' v' e' v' g' f' 'ur' c' y' p' l' p' i' "  
ci i tgi cvkpu' k' p' y j g' 'E' c' { o' c' p' 'K' r' c' p' f' u' 0' *Proceedings of the Gulf and Caribbean Fisheries Institute*. "**60:7: 7/7: 80**

"

Uj cr ktq. 'F Q 03; : 90Tgr tqf vevkqp'lp'i tqwr gtu.'r 04; 7/549. 'in. 'LLO'Rqrxkpc'cpf 'UOT'cnvqp '\*gf u0: 'Tropical Snappers and Groupers: Biology and Fisheries Management'0Y guxkgy 'Rt guo'Dqwr gt. 'EQ0'

"

Uj gpngt. 'LLO 0'GOF 00 cf f qz. 'G0Y kuj kpunk 'C0Rgctn 'UOT'0Vj qttqrf. 'cpf 'P 0Uo kj 03; ; 50Qpuj qtg'tcpur qt v'qh' ugwrqo gpvuci g'P cuucw'i tqwr gt '\*Epinephelus striatus+cpf 'qyj gt 'Hkuj gu'lp'Gzwo c'Uqwpf. 'Dcj co cu0 Marine Ecology Progress Series. '98:53/650'

"

Ukxc 'Ngg. 'C'0H03; 960J 'a dksqu'cnko gpvctkqu'f g'nc'ej gtpc'etkqmc 'Epinephelus striatus' Dnjej '{ 'cri wpu'f cvqu'uqdtg'uw' dlqmqi kc0Serie Oceanologica Academia de Ciencias de Cuba'25:5/360'

"

Ukxc 'Ngg. 'C'0H03; 990P qvc'uqdtg'nc'eqm'celx'p' { 'nc'eqpf vev'f g'nc'ej gtpc'etkqmc '\*Epinephelus striatus Dnjej +0' Cecf go kc'f g'Ekpkelcu'f g'Ewdc. 'Kpukw'g'f g'Qegcpmqi kc. 'Kphqto g'Ekgp'w'heq 'Vgepleq'P q036. 'Ewdc. ': 'r r 0'

"

Uwnc. 'T0'O 0Ej kr r qpg. 'MO 0Uwnkxcp. 'cpf 'T0'Y tki j x03; ; 80J cdkcv'cpf 'Nhg'lp'vj g'Gzwo c'Ec {u. 'vj g'Dcj co cu' vj g'ucwu'qh'i tqwr gtu'cpf 'eqtcn'tggh'lp'vj g'P qt vj gtp'Ec {u'0Vj g'P cwtg'Eqpugt xcp {0"

"

Uwnc. 'T0'O 0Ej kr r qpg. 'MO 0Uwnkxcp. 'VORqwu. 'LLO 0Ngx { . 'GHO'Uej o kv'cpf 'I 0O ggugt03; ; : 0F gpukv { . 'ur gelgu' cpf 'uk g'f kmtkdwkqp'qh'i tqwr gtu '\*Ugttcpkf cg+'lp'vj tgg'j cdkcv'cv'Gndqy 'Tggg. 'Hqtkf c'Mg {u'0Bulletin Marine Science. '62:43; /44: 0'

"

Uo kj . 'E0N03; 830U { pqr uku'qh'dkqmqi lecn'f cvc"qp'i tqwr gtu '\*Gr lpg'j gnuw cpf 'cmkgf 'i gpgtc+'qh'vj g'y guvgt'P qt vj " Cvc'p'v'0HCQ 'Hkuj 0Dkq'0U { pqr 0P q045. '83'r r 0'

"

Uo kj . 'E0N03; 930C 'tgxkukqp'qh'vj g'Co gtlecp'i tqwr gtu '<Epinephelus"cpf 'cmkgf 'i gpgtc'0Bulletin of the American Museum of Natural History. '146:8; /4630'

"

Uo kj . 'E0N03; 940C 'ur cy plpi 'ci i tgi cvkqp'qh'P cuucw'i tqwr gt. 'Epinephelus striatus \*Dnjej +0Transactions of the American Fisheries Society, 101:479/483"

"

Uo kj . 'E0N03; 9: 0Ugttcpkf cg0in. 'Y 0Huej gt '\*gf 0: 'HCQ'Ur gelgu'K gpv'hecvkqp'Uj ggv'u'qt 'Hkuj gt { 'Rwr qugu. " Y guvgt'P'egpvc'cn'0Cvc'p'v'k. 'Hkuj lpi 'Ctgc'530Xq'0KX. 'X0HCQ. 'Tqo g0]Wpr ci kpcv'f 0'

"

Uo kj /Xcpk . 'Y 0H'DD0E'qmgw'g. 'cpf 'D0G0N'wenj wtu03; ; : 0Fishes Of Bermuda: History, Zoogeography, Annotated Checklist, and Identification Keys0Co gtlecp'Uqelgv' 'qh'Kj vj { qmqi kuw'cpf 'J gtr g'v'qmqi ku'Ur gelcn' Rv'd'hecvkqp'P q060Reviews in Fish Biology and Fisheries0'

"

Uquc/Eqtf gtq 'G0'C0O gf kpc/S wgl. 'T0J gttgtc. 'cpf 'Y 0Ci w'act 'F' a xkx042240Ci tgi celqpgu'tgr tqf vevkcu'f g'r gegu' gp'gn'U'ungo c'Ctt'gelk'cn'0 guqco gtlecpq '<Eqpuw'nat'f'P celqpcn'0 2 zleq0'Kphqto g'r tgr ctcf q'r ctc'gn'leqpuw'nat'f' k'p'v'gt'p'celqpcn'T gugctej 'Rvc'p'kpi 'Kpe0' { 'Rtq { gev'uco 0Gequw. 'Ej gwo cn'S w'k'v'cpc'Tqq047'r r 0"

"

Ur tkpi gt. 'XQ 0'cpf 'C'LOO eGtr'gcp03; 840C 'uwf { 'qh'vj g'dgj cxkqt'qh'uqo g'vci i gf 'Uqwj gtp'Hqtkf c'eqtcn'tggh'kuj gu0 American Midland Naturalist. '67:5: 8/5; 90'

"

Ucnc'kpi u. 'E0F 0422: 0'Kpf k'gev'gh'g'ew'qh'cp'g'zr m'k'g'f 'r t'gf cvqt'qp't'getwko gpv'qh'eqtcn'tggh'kuj gu0'Ecology0: ; \*: +0'

"

Uctem 'Y 0C0'K03; 8: 0C 'kuw'qh'kuj gu'qh'0Cnki cvqt 'Tggg. 'Hqtkf c'y kj 'eqo o gpw'qp'vj g'pcwtg'qh'vj g'Hqtkf c'tggg' Hkuj 'h'wpc'0Undersea Biology. '1:7/580'

"

Uctem 'Y 0C0'K'cpf 'Y 0R0F cxk03; 880P ki j v'j cdku'qh'kuj gu'cv'0Cnki cvqt 'Tggg. 'Hqtkf c'0Ichthyologica'38(4):535/ 5780'

"

Uctt. 'T00 0'G0U'nc. 'G0D'cngugtqu. 'cpf 'O 0\ cdcn:042290Ur cvkcn'f { pco leu'qh'vj g'P cuucw'i tqwr gt 'Epinephelus striatus lp'c'Ec'tk'dd'gcp'cv'q'0Marine Ecology Progress Series. '343:45; /46; 0'

"

Ugxgquqp. 'F 0M03; : 30A review of the marine resources of the Western Central Atlantic Fisheries Commission (WECAFC) Region 0HCQ 'Hkuj 0Vgej 0Rcr 04330HCQ. 'Tqo g. '354' r 0

"

Uwrxcp. 'MO 0' cpf 'O 0f g' I ctkpg/Y kej cksun{ 03; ; 60Gpgti g'leu'qh'lwxgpkrg" *Epinephelus* 'i tqwr gtu'k' r cev'qh' uwo o gt 'ygo r gtcwtgu' cpf "cevkxk{ 'r cvgtpu'qp' i tqy vj 'tcvgtu' 0Proceedings of the Gulf and Caribbean Fisheries Institute, **43**:36: /3890

"

Uwrxcp/Ugcrg{ . 'MO' V0Tcj o kpi . 'cpf 'O 0Tqmg042240Uk g. 'ugz' 'tcvqk. 'cpf 'hgewpf k{ "qh'P cuucw'i tqwr gt "Epinephelus striatus'+h' cpf gf 'f wtkpi 'ur cy plpi 'ugcuqp 'kp' vj g' EgpwtcnDcj co cu' 0Proceedings of the Gulf and Caribbean Fisheries Institute. **53**:694/6: 30"

"

Vc{ nqt. 'L0E0' F 0D0Gi i nguqp. 'cpf 'R0U0T cpf 042280P cuucw'i tqwr gt "Epinephelus striatus'+ur cy plpi "ci i tgi cvkqpu' j { f tqceqw'wle' uwtxg{ u' cpf 'i gquc'w'wle' cnc' cpcn{ uku' 0Kp' <Go gti kpi "Vgej pqmji kgu' Hqt "Tgg'h' Hkuj g' tkgu' Tgugctej " Cpf "O cpci go gpw' 0P QCC "Rtqhgukqpcn' Rcr gtu' P O HU' \*7+0P QCC. "Ugcwrg. "Y C. 'r 03: /47"

"

Vj qo r uqp. 'GH03; 670The Fisheries of British Honduras 0F gxgrq o gpv' cpf "Y grtctg' kp' vj g' Y guv' kpf kgu. 'Cf xqecv'g' Eq0' Dtkf i gqy p. 'Dctdcf qu' 0Dwn' 021:3/540

"

Vj qo r uqp. 'T0Y 03; 9: 0Tguw'u' qh' vj g' WPFR' IHCQ' Dcj co cul' ggr 'y cvgt' h'kuj gt { 'uwtxg{ '3; 94/3; 970Proceedings of the Gulf and Caribbean Fisheries Institute. **30**:66/920

"

Vj qo r uqp. 'T0' cpf 'L0N0O wptq' 03; 9: 0Cur gew' qh' vj g' d'kqmi { "cpf "geqmi { "qh' Ectkddgcp' tgg'h' Hkuj gu' Ugttcplf cg" \*j kpf u' cpf 'i tqwr gtu' 0Journal of Fish Biology. **12**:337/3680

"

Vj qo r uqp. 'T0' cpf 'L0N0O wptq' 03; ; 50Ej cr vgt '9' <Vj g' d'kqmi { . "geqmi { "cpf "dkqpqo leu' qh' vj g' j kpf u' cpf 'i tqwr gtu. " Ugttcplf cg. 'r 07; /: 3. "in: 'L0N0O wptq' \*gf 0: "Caribbean Coral Reef Fishery Resources. "KENC TO 'Uwf kgu' cpf ' Tgxlg' y u. "Xqr' 090' k' vgt' pc' w' k' p' c' n' Egpvgt 'hqt "Nlxkpi "cpf 'Cs w' wle' Tguq' wtegu' O cpci go gpv. 'O cplrc. 'Rj kkr r kpgu' Eqp' w' kd' 0347. '4pf "gf 0' 498' r 0

"

Vqy pugpf . 'E0' 03; 270Tgr qt v' qh' vj g' f' k' gevqt 'qh' vj g' cs w' wtkwo 0P k' vj "Cp' w' c' n' Tgr qt v' qh' vj g' P gy 'I q' t' n' l' q' q' m' i' k' e' c' n' Uqekv{ . 'D' k' q' m' i' { 09: : ; /3250

"

Vwengt. 'L0Y 0' L' t' 03; ; 4c0Ur cy plpi "ugttcplf 'Hkuj gu' lp' ecr w' k' k' { 0Kp. "Aquaculture '92 -Growing towards the 21st Century. 'r 043; 0

"

Vwengt. 'L0Y 0' L' t' 03; ; 4d0I tqwr gt 'ew' w' w' g' h' q' t' vj g' Ectkddgcp' 0Rtqi t' gu' t' gr qt v' 0Proceedings of the Gulf and Caribbean Fisheries Institute. **41**:7: 90

"

Vwengt. 'L0Y 0' L' t' 03; ; 60Ur cy plpi 'd{ 'ecr w' k' g' ugttcplf 'Hkuj gu' c' c' t' g' x' l' g' y 0Journal of the World Aquaculture Society. **25**:567/57; 0

"

Vwengt. 'L0Y 0' L' t' 0' cpf 'RP 0Y qqf y ctf 03; ; 50P cuucw'i tqwr gt 'cs w' c' ew' w' w' g. 'r 0585/599. "in: "H0C' t' t' g' i' w' k' p' /Ucpej gl . 'L0N0' O wptq. 'O (E0D' c' n' i' qu. 'cpf 'F 0R' c' w' n' { "gf u' 0: "Biology, Fisheries, and Culture of Tropical Groupers and Snappers. "KENC TO 'Eqph' 0Rtqe' 06: . '66; 'r 0

"

Vwengt. 'L0Y 0' L' t' 0' cpf 'RP 0Y qqf y ctf 03; ; 60I tqy vj 'cpf 'f gxgrq o gpv' qh' f' qo gu' wle' l' w' x' g' p' k' r' g' P cuucw'i tqwr gtu' 0Proceedings of the Gulf and Caribbean Fisheries Institute. **43**:5; ; /5; 30

"

Vwengt. 'L0Y 0' R' I' 0D' w' j' . 'cpf 'U' 0V' 0U' r' { d' c' w' i' j' 03; ; 50Tgr tqf w' e' w' k' g' r' cvgtpu' qh' Ec{ o cp' "K' r' c' p' f' u' P cuucw'i tqwr gt" \*Epinephelus striatus'+r qr w' r' w' k' p' u' 0Bulletin of Marine Science, **52**:; 836; 8; 0"

"

Vwengt. 'L0Y 0' L' t' 0' R' P' 0Y qqf y ctf . 'cpf 'F' 0' U' g' p' p' g' y' 03; ; 80X' q' n' w' p' v' c' t' { 'ur cy plpi "qh' e' c' r' w' k' g' P cuucw'i tqwr gtu' Epinephelus striatus' kp' c' eqpetg' g' t' c' e' g' y' c' { 0Journal of the World Aquaculture Society. **27**(4):595/5: 50

"  
"  
"  
"

"

Vwengt. 'L0Y 0'Lt0'LOGRctuqpu.'I (E0Gdcpmu.'cpf 'R0 0'Dwuj 03; ; 30'kpf wegf 'ur cy plpi 'qh'P cuucw'i tqwr gt"  
*Epinephelus striatus*Journal of the World Aquaculture Society. '22:3: 9/3; 30"

"

Vwr r gt. 'O 042240Guugpvkci'hkuj 'j cdkcv'cpf "o ctkpg'tgugt xgu'htq'i tqwr gtu'lp'yj g'Vwmu'cpf "Eclequ'Krcpf u0"  
*Proceedings of the Gulf Caribbean Fisheries Institute.*'53:828/8440"

"

Vwr r gt. 'O 0 0'cpf 'Twf f. 'O (C042240Ur gelgu'ur gekhe 'ko r cew'qh'c'uo cml'o ctkpg'tgugt xg'qp'tgg'hkuj 'r tqf wevkp'cpf "  
 hkuj kpi 'r tqf wevkv' 'lp'yj g'Vwmu'cpf "Eclequ'Krcpf u0*Environmental Conservation.* '29:6: 6/6; 40"

"

N0X<sup>a</sup> us wgl / [ gqo cpu. 'W0Qtf q° gl /N>r gl 'cpf "G0Uquc/Eqtfgtq03; ; 0Hkuj 'rtxcg'cf lcegpv'v'c'eqtcn'tgg'h'lp'yj g"  
 y guvtp'Ectkddgcp'Ugc'qhi'O cj wcn'O gzleq0*Bulletin of Marine Science.* 62(1):44; /4670"

"

Xkrtq'F kl. 'F 003: : 60Eqttkf c' { 'cttkdc| qp'f'g'cni wpu'r gegu'ewdcpqu00 cpwgnI qo gl 'f'g'r'O c| c. 'Nc'J cdcpc. 'Ewdc'"

"

Y cxcpcdg. 'Y 0Q0'E0U0Ngg. 'UE0Gnku. 'GROGnku. 'cpf 'GROGnku03; ; 7c0J cvej gt { 'uwf { "qh'yj g'ghgevu'qh'vgo r gtcwtg'qp'gi i u"  
 cpf "{ qmuce'rtxcg'qh'yj g'P cuucw'i tqwr gt "*Epinephelus striatus*Journal of the World Aquaculture, '136:363/3690"

"

Y cxcpcdg. 'Y 0Q0'UE0Gnku. 'GROGnku. 'Y 0F0J gcf. 'E0F0Mngg { 'C00 qtly cng. 'E/U'Ngg. 'cpf 'R0M0Dlghrcpi 03; ; 7d0'  
 Rtqi tgu'lp'eqpvtqmgf "dtggf kpi "qh'P cuucw'i tqwr gt "*Epinephelus striatus*+dtqqf uqemid { "j qto qpg"  
 kpf wevkp0*Aquaculture,* '138:427/43; 0"

"

Y cxcpcdg. 'Y 0Q0'UE0Gnku. 'GROGnku. 'I 0Nqr gl. 'R0Dcuu. 'L0I kpgl c. 'cpf 'C00 qtly cng03; ; 80Gxcnwv'kp'qh'htuv'  
 hggf kpi 'tgi ko gpu'ht'rtxcn'P cuucw'i tqwr gt "*Epinephelus striatus*"cpf 'r tgrko kpc { 'r kv'uecrg'ewmwg"  
 y tqw j 'o gco qtr j quku0*Journal of the World Aquaculture Society,* '27(3):545/5530"

"

Y cxcpcdg. 'Y 0Q0'E0U0Ngg. 'UE0Gnku. 'GROGnku. 'Y 0F0J gcf. 'E0F0Mngg { 'I 00 k'co qv. 'M0Nkw. 'cpf 'L0I kpgl c0'  
 3; ; 60Gzr gtko gpwn'ewmwg'qh'rtxcn'P cuucw'i tqwr gt "*E. striatus*+<yj g'ghgevu'qh'vgo r gtcwtg'qp'gi i 'cpf "  
 { qmuce'uci gu'cpf 'qh'r tg { 's wckv' 'qp'uw'xkcn'c'v'ht'uv'hggf kpi 0Cduwcev'ht'Y qtrf 'Cs wcewmtg'; 6. 'r 04: 90'  
 Y qtrf "Cs wcewmtg"Uqelgv'0P gy 'Qtrgcpu. 'NC. 'Lcp036/'3: .3; ; 60"

"

Y gmu. 'L0Y 0'cpf 'L0E0Ncpi 03; 950U { ugo cve 'ku'qh'Lco clecp'uj cmjy /y cvt 'uergtcev'kpc0*Bulletin of Marine Science.*"  
 23:77/7: 0"

"

Y j c { rpg. 'N0'E0K0Rcvgpi km'Ugo o gpu. 'D0Z0Ugo o gpu. 'R0 0'Dwuj. 'cpf 'O 0F0Dqctf o cp042260Qdugt xcv'kpu'qh'c"  
 P cuucw'i tqwr gt, *Epinephelus striatus.* 'ur cy plpi 'ci i tgi cvkqp'ukg'lp'Nkwg'Ec { o cp. 'Ec { o cp 'Krcpf u."  
 kpenmf kpi 'o wv'ur gelgu'ur cy plpi 'kphqto cvkqp0*Environmental Biology of Fishes.* '70:527/5350"

"

Y j c { rpg. 'N0'R0Dwuj. 'D0Lqj puqp. 'M0Nwng. 'E00 eEtq { . 'U0J gr r gm 'D0Ugo o gpu. 'O 0F0Dqctf o cp042290'  
 Ci i tgi cvkqp'f { pco leu'cpf 'rguqupu'hgctpgf 'htqo 'h'xg' { gctu'qh'o qpkqtkpi "cv'c'P cuucw'i tqwr gt "*Epinephelus*  
*striatus*+ur cy plpi 'ci i tgi cvkqp'lp'Nkwg'Ec { o cp. 'Ec { o cp 'Krcpf u 'DY K0*Proceedings of the Gulf and*  
*Caribbean Fisheries Institute,* '59:6356643"

"

Y kreqz. 'Y (C03: ; ; 0Vj g'kuj gtkgu'cpf 'kuj 'v'cf g'qh'Rqtv'q'Tleq. 'r 04966: . 'in. 'kpxguki cvkqp'qh'yj g'cs wv'le 'tguqwtegu'  
 cpf 'kuj gtkgu'qh'Rqtv'q'Tleq0'WU0Ego o kuukqp'qh'Hkuj 'cpf 'Hkuj gtkgu0"

"

Y kpi. 'G0U0'cpf 'G0U0Tgk' 03; ; 40Rtg kvqtle 'kuj kpi 'geqpqo lgu'qh'yj g'Ectkddgcp0*Journal of New World*  
*Archaeology,* 5(2):35/540"

"

Y kpi. 'G0U0'E0C0J qhho cp. 'Lt. 'cpf 'E0G0Tc { 03; 8: 0Xgtvgdtevg'tgo clpu'htqo 'kpf kcp'ukgu'qp'c'pki wc. 'Y guv'kpf lgu0'  
*Caribbean Journal of Science.* '8(3&4):345/35; 0"

"

Y qqf rg { . 'LF 03; ; 70*Tropical Americas Regional Report on the Issues and Activities Associated with Coral Reefs*  
*and Related Ecosystems*0Rtgr ctgf 'ht'v'j g'3; ; 7'k'v'gt'cv'kpcn'Eqtcn'Tgg'h'k'k'c'v'xg'Y qtmuj qr. 'F wo ci wv'g"

Ek{ . 'Rj kkr r kpgu0'

"

Y qqf rg{ . 'LF0'MOF g'O g{ gt. 'R0Dwuj . 'I 0Gdcpmu/Rgvtkg. 'L0I ct| qp/Hgttgktc. 'G0Mrgkp. 'N0Rqtu. 'cpf 'E0Y kuzp03; ; : 0'  
Ucwu'qh'eqtcrit'gghu'kp'y g'uqwj /egpwtci'Ectkddgcp0'k<Status of Coral Reefs of the World: 19980E0'  
Y kmpuqp "gf 0'Cwutcrkcp'kpukwg'qh'O ct|pg'Uelgpeg0'

"

"

"

"

United States of America

Nomination of the Nassau grouper (*Epinephelus striatus*) for inclusion in Annex III  
Appendix B\_FINAL Nassau grouper ESA rule

also reduce, eliminate, or prevent unnecessary differences in regulatory requirements.

Similarly, the Trade Agreements Act of 1979 (Pub. L. 96-39), as amended by the Uruguay Round Agreements Act (Pub. L. 103-465), prohibits Federal agencies from establishing any standards or engaging in related activities that create unnecessary obstacles to the foreign commerce of the United States. For purposes of these requirements, Federal agencies may participate in the establishment of international standards, so long as the standards have a legitimate domestic objective, such as providing for safety, and do not operate to exclude imports that meet this objective. The statute also requires consideration of international standards and, where appropriate, that they be the basis for U.S. standards.

PHMSA participates in the establishment of international standards in order to protect the safety of the American public, and we have assessed the effects of the interim final rule to ensure that it does not cause unnecessary obstacles to foreign trade. Accordingly, this rulemaking is consistent with Executive Order 13609 and PHMSA's obligations.

List of Subjects

49 CFR Part 107

Administrative practices and procedure, Hazardous materials transportation, Packaging and containers, Penalties, Reporting and recordkeeping requirements.

49 CFR Part 171

General information, Regulations, and Definitions.

In consideration of the foregoing, 49 CFR Chapter I is amended as follows:

PART 107—HAZARDOUS MATERIALS PROGRAM PROCEDURES

1. The authority citation for part 107 is revised to read as follows:

Authority: 49 U.S.C. 5101-5128, 44701; Pub. L. 101-410 section 4; Pub. L. 104-121, sections 212-213; Pub. L. 104-134, section 31001; Pub. L. 114-74 section 4 (28 U.S.C. 2461 note); 49 CFR 1.81 and 1.97.

2. Revise § 107.329 to read as follows:

§ 107.329 Maximum penalties.

(a) A person who knowingly violates a requirement of the Federal hazardous material transportation law, an order issued thereunder, this subchapter, subchapter C of the chapter, or a special permit or approval issued under this subchapter applicable to the transportation of hazardous materials or

the causing of them to be transported or shipped is liable for a civil penalty of not more than \$77,114 for each violation, except the maximum civil penalty is \$179,933 if the violation results in death, serious illness or severe injury to any person or substantial destruction of property. There is no minimum civil penalty, except for a minimum civil penalty of \$463 for violations relating to training. When the violation is a continuing one, each day of the violation constitutes a separate offense.

(b) A person who knowingly violates a requirement of the Federal hazardous material transportation law, an order issued thereunder, this subchapter, subchapter C of the chapter, or a special permit or approval issued under this subchapter applicable to the design, manufacture, fabrication, inspection, marking, maintenance, reconditioning, repair or testing of a package, container, or packaging component which is represented, marked, certified, or sold by that person as qualified for use in the transportation of hazardous materials in commerce is liable for a civil penalty of not more than \$77,114 for each violation, except the maximum civil penalty is \$179,933 if the violation results in death, serious illness or severe injury to any person or substantial destruction of property. There is no minimum civil penalty, except for a minimum civil penalty of \$463 for violations relating to training.

3. In Appendix A to subpart D of part 107, Section II.B. ("Penalty Increases for Multiple Counts"), the first sentence of the second paragraph is revised to read as follows:

Appendix A to Subpart D of Part 107—Guidelines for Civil Penalties

Under the Federal hazmat law, 49 U.S.C. 5123(a), each violation of the HMR and each day of a continuing violation (except for violations relating to packaging manufacture or qualification) is subject to a civil penalty of up to \$77,114 or \$179,933 for a violation occurring on or after August 1, 2016.

PART 171—GENERAL INFORMATION, REGULATIONS, AND DEFINITIONS

4. The authority citation for part 171 is revised to read as follows:

Authority: 49 U.S.C. 5101-5128, 44701; Pub. L. 101-410 section 4; Pub. L. 104-134, section 31001; Pub. L. 114-74 section 4 (28 U.S.C. 2461 note); 49 CFR 1.81 and 1.97.

5. In § 171.1, paragraph (g) is revised to read as follows:

§ 171.1 Applicability of Hazardous Materials Regulations (HMR) to persons and functions.

\* \* \* \* \*

(g) Penalties for noncompliance. Each person who knowingly violates a requirement of the Federal hazardous material transportation law, an order issued under Federal hazardous material transportation law, subchapter A of this chapter, or a special permit or approval issued under subchapter A or C of this chapter is liable for a civil penalty of not more than \$77,114 for each violation, except the maximum civil penalty is \$179,933 if the violation results in death, serious illness or severe injury to any person or substantial destruction of property. There is no minimum civil penalty, except for a minimum civil penalty of \$463 for a violation relating to training.

Issued in Washington, DC, on June 14, 2016 under authority delegated in 49 CFR part 1.97.

Marie Therese Dominguez,

Administrator, Pipeline and Hazardous Materials Safety Administration.

[FR Doc. 2016-15404 Filed 6-28-16; 8:45 am]

BILLING CODE 4910-60-P

DEPARTMENT OF COMMERCE

National Oceanic and Atmospheric Administration

50 CFR Part 223

[Docket No. 1206013326-6497-03]

RIN 0648-XA984

Endangered and Threatened Wildlife and Plants: Final Listing Determination on the Proposal To List the Nassau Grouper as Threatened Under the Endangered Species Act

AGENCY: National Marine Fisheries Service (NMFS), National Oceanic and Atmospheric Administration (NOAA), Commerce.

ACTION: Final rule; request for information.

SUMMARY: We, NMFS, are publishing this final rule to implement our determination to list the Nassau grouper (Epinephelus striatus) as threatened under the Endangered Species Act of 1973, as amended (ESA). We have completed a status review of the Nassau grouper in response to a petition submitted by WildEarth Guardians. After reviewing the best scientific and commercial data available, including the status review and comments received on the proposed rule, we have determined that the Nassau grouper

meets the definition of a threatened species. While the species still occupies its historical range, overutilization through historical harvest has reduced the number of individuals which in turn has reduced the number and size of spawning aggregations. Although harvest of Nassau grouper has diminished due to management measures, the reduced number and size of spawning aggregations and the inadequacy of law enforcement continue to present extinction risk to Nassau grouper. Based on these considerations, described in more detail within this action, we conclude that the Nassau grouper is not currently in danger of extinction throughout all or a significant portion of its range, but is likely to become so within the foreseeable future. We also solicit information that may be relevant to the designation of critical habitat for Nassau grouper, including information on physical or biological features essential to the species' conservation, areas containing these features, and potential impacts of a designation.

**DATES:** The effective date of this final rule is July 29, 2016. Information on features, areas, and potential impacts, that may support designation of critical habitat for Nassau grouper must be received by August 29, 2016.

**ADDRESSES:** Information regarding this final rule may be obtained by contacting NMFS, Southeast Regional Office, 263 13th Avenue South, Saint Petersburg, FL 33701. Supporting information, including the Biological Report, is available electronically on the NMFS Web site at: [http://sero.nmfs.noaa.gov/protected\\_resources/listing\\_petitions/species\\_esa\\_consideration/index.html](http://sero.nmfs.noaa.gov/protected_resources/listing_petitions/species_esa_consideration/index.html).

You may submit information regarding potential critical habitat designation to the Protected Resources Division by either of the following methods:

- **Electronic Submissions:** Submit all electronic comments via the Federal eRulemaking Portal. Go to [www.regulations.gov/](http://www.regulations.gov/)#!/docketDetail;D=NOAA-NMFS-2015-0130, click the "Comment Now!" icon, complete the required fields, and enter or attach your comments.

- **Mail:** Submit written information to the Protected Resources Division, NMFS Southeast Regional Office, 263 13th Avenue South, Saint Petersburg, FL 33701.

**FOR FURTHER INFORMATION CONTACT:** Adam Brame, NMFS, Southeast Regional Office (727) 209-5958; or Lisa Manning, NMFS, Office of Protected Resources (301) 427-8466.

**SUPPLEMENTARY INFORMATION:**

## Background

On September 3, 2010, we received a petition from the WildEarth Guardians to list speckled hind (*Epinephelus drummondhayi*), goliath grouper (*E. itajara*), and Nassau grouper (*E. striatus*) as threatened or endangered under the ESA. The petition asserted that (1) the present or threatened destruction, modification, or curtailment of habitat or range; (2) overutilization for commercial, recreational, scientific, or educational purposes; (3) inadequacy of existing regulatory mechanisms; and (4) other natural or manmade factors are affecting the continued existence of and contributing to the imperiled statuses of these species. The petitioner also requested that critical habitat be designated for these species concurrent with listing under the ESA. Due to the scope of the WildEarth Guardians' petition, as well as the breadth and extent of the required evaluation and response, we provided species-specific 90-day findings (76 FR 31592, June 1, 2011; 77 FR 25687, May 1, 2012; 77 FR 61559, October 10, 2012).

On October 10, 2012, we published a 90-day finding for Nassau grouper with our determination that the petition presented substantial scientific and commercial information indicating that the petitioned action may be warranted (77 FR 61559). At that time, we announced the initiation of a formal status review and requested scientific and commercial information from the public on: (1) The status of historical and current spawning aggregation sites; (2) historical and current distribution, abundance, and population trends; (3) biological information (life history, genetics, population connectivity, etc.); (4) management measures, regulatory mechanisms designed to protect spawning aggregations, and enforcement information; (5) any current or planned activities that may adversely impact the species; and (6) ongoing or planned efforts to protect and restore the species and its habitat.

As part of the status review process to determine whether the Nassau grouper warrants listing under the ESA, we completed a Biological Report and an extinction risk analysis (ERA). The Biological Report summarizes the taxonomy, distribution, abundance, life history, and biology of the species. The Biological Report also identifies threats or stressors affecting the status of the species as well as a description of the fisheries, fisheries management, and conservation efforts. The Biological Report incorporates information received in response to our request for information (77 FR 61559, October 10,

2012) and comments from three independent peer reviewers. We used the Biological Report to complete a threats evaluation and an ERA to determine the status of the species.

After completing the Biological Report and considering the information received on the 90-day finding, we published a proposed rule to list Nassau grouper as a threatened species on September 2, 2014 (79 FR 51929). During a 90-day comment period, we solicited comments on our proposal from the public and any other interested parties.

## Listing Determinations Under the ESA

We are responsible for determining whether the Nassau grouper is threatened or endangered under the ESA (16 U.S.C. 1531 *et seq.*). Section 4(b)(1)(A) of the ESA requires us to make listing determinations based solely on the best scientific and commercial data available after conducting a review of the status of the species and after taking into account efforts being made by any state or foreign nation to protect the species. To be considered for listing under the ESA, a group of organisms must constitute a "species," which is defined in section 3 of the ESA to include taxonomic species and "any subspecies of fish, or wildlife, or plants, and any distinct population segment of any species of vertebrate fish or wildlife which interbreeds when mature."

Section 3 of the ESA defines an endangered species as "any species which is in danger of extinction throughout all or a significant portion of its range" and a threatened species as one "which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range." Thus, we interpret an "endangered species" to be one that is presently in danger of extinction. A "threatened species," on the other hand, is not currently in danger of extinction but is likely to become so in the foreseeable future. In other words, a key statutory difference between a threatened and endangered species is the timing of when a species may be in danger of extinction, either presently (endangered) or in the foreseeable future (threatened).

Under section 4(a) of the ESA, we must determine whether any species is endangered or threatened due to any of the following five factors: (A) The present or threatened destruction, modification, or curtailment of its habitat or range; (B) overutilization for commercial, recreational, scientific, or educational purposes; (C) disease or predation; (D) the inadequacy of

existing regulatory mechanisms; or (E) other natural or manmade factors affecting its continued existence (sections 4(a)(1)(A) through (E)). We are required to make listing determinations based solely on the best scientific and commercial data available after conducting a review of the status of the species and after taking into account efforts being made by any state or foreign nation to protect the species.

In determining whether the Nassau grouper meets the standard of endangered or threatened, we followed a stepwise approach. First we considered the specific life history, ecology, and status of the species as documented in the Biological Report. We then considered information on factors adversely affecting and posing extinction risk to the species in a threats evaluation. In this evaluation we assessed the threats affecting the status of the species using the factors identified in ESA section 4(a)(1). We considered the nature of the threats and the species response to those threats. We also considered each threat identified, both individually and cumulatively. Once we evaluated the threats, we assessed the efforts being made to protect the species to determine if these conservation efforts were adequate to mitigate the existing threats and alter extinction risk. Finally, we considered the public comments received in response to the proposed rule. In making this finding, we have relied on the best available scientific and commercial data.

### Summary of Comments Received

Below we address the comments received on the proposed listing for Nassau grouper. In response to our request for public comments, we received 17 written responses. The overall feedback was supportive of the rule with the exception of three commenters, who believe current regulations within the United States are sufficient in protecting this species. No comments addressed threats to Nassau grouper throughout the rest of their range. We did not receive any information on additional conservation efforts being taken.

*Comment 1:* Multiple commenters supported the proposed rule to list Nassau grouper as a threatened species and further encouraged regional collaboration to develop adequate management measures.

*Response:* We agree that regional collaboration will strengthen efforts to consistently manage and conserve the species, and we hope this listing will encourage collaborative efforts. In some cases, adding a species to the

endangered species list leads to increased funding opportunities and potential for collaboration between state and federal partners, as well as stakeholders. We will seek regional collaborative conservation efforts within the Caribbean region to further the conservation of the species.

*Comment 2:* We received comments that the existing management measures implemented by Fishery Management Councils are already effective at protecting Nassau grouper within U.S. waters, (including U.S. territorial waters of Puerto Rico and the U.S. Virgin Islands) and that the listing may add unnecessary burdens on our domestic fisheries.

*Response:* We agree that the South Atlantic Fishery Management Council and the Caribbean Fishery Management Council have taken significant steps to protect and rebuild the Nassau grouper population in U.S. waters. Unfortunately, a large part of the species' range and population is outside of U.S. jurisdiction and is therefore not directly aided by Council protections. We must make our determination based on the best scientific and commercial data available, independent of the potential burdens to our other domestic fisheries. This standard has been applied when making the Nassau grouper final listing determination.

*Comment 3:* Some comments expressed concern over the economic consequences of listing Nassau grouper, including possible effects on commercial fishermen.

*Response:* We are unable to consider economic impacts in a listing determination. The ESA requires us to make listing determinations by evaluating the standards and factors in section 4 of the ESA, and based solely on the best scientific and commercial data available. Listing Nassau grouper as a threatened species would not create any immediate additional regulatory requirements directly affecting commercial fishermen. Potential future regulations affecting conservation of Nassau grouper, including take and import regulations may be proposed via a separate rulemaking process which would include consideration of certain economic impacts (e.g., impacts on small businesses) and opportunities for public input. Individuals that require federal permits or funding for actions that might affect Nassau grouper might need to make adjustments to their activities to avoid jeopardizing Nassau grouper, and to avoid or minimize take of the species, but that would be a determination for a specific section 7 consultation in the future.

*Comment 4:* Several comments indicated that spawning aggregation sites need to be protected and that proper enforcement of both existing and future rules is paramount in protecting the species.

*Response:* We agree that the lack of adequate protections for Nassau grouper spawning aggregations and the inadequacy of law enforcement are major contributors to the species' decline throughout its range. These threats were rated 'high' during the ERA as explained in the proposed rule and, as such, were taken into consideration when making our final listing determination.

*Comment 5:* One commenter supported the rule stating, "We agree that the best available science demonstrates that Nassau grouper is likely to be at risk of extinction in the foreseeable future, and may in fact be in danger of extinction now." They further encouraged swift designation of critical habitat to protect spawning aggregation sites, nursery and juvenile habitat, and feeding habitat.

*Response:* We acknowledge the concern raised by the commenter that the species may be in danger of extinction now and provide further detail below as to how we reached our listing determination in this final rule. With regard to critical habitat, section 4(a)(3)(A) of the ESA (16 U.S.C. 1533(a)(3)(A)) requires that, if prudent and determinable, critical habitat be designated concurrently with the listing of a species. We do not currently have sufficient information to determine what physical and biological features within Nassau grouper habitats facilitate the species' life history strategy and thus are essential to the species' conservation. Therefore, we cannot yet determine what areas meet the definition of critical habitat under the ESA. Because critical habitat is not currently determinable, we will not designate critical habitat concurrently with this final rule. Designation of critical habitat may occur via a subsequent rule-making process if we can identify critical habitat and designation is prudent. We are soliciting information on features, areas, and impacts of designation, that may support designation of critical habitat for Nassau grouper.

*Comment 6:* One commenter suggested the use of size restrictions, monitoring, closed fishing seasons for the protection of spawning aggregations, and the use of marine protected areas as measures to protect the species.

*Response:* We summarize in this rule the existing regulations currently in place throughout the Caribbean Sea that

include many of these suggested practices. Within U.S. waters, measures to protect Nassau grouper are already in place under the Magnuson-Stevens Act and State and Territorial fishery management authorities. As a species listed as threatened under the ESA, any federal action implemented, authorized or funded that “may affect” Nassau grouper will require consultation to ensure the action is not likely to jeopardize the species’ continued existence. We may also implement additional protective regulations for Nassau grouper under section 4(d) of the ESA if we determine such regulations are necessary and advisable for the conservation of this threatened species. Issuance of a 4(d) rule would be a separate rule-making process that would include specific opportunities for public input.

*Comment 7:* The U.S. Navy identified three Navy installations or properties that are within the geographic range of Nassau grouper. They expressed concern over their ability to utilize and maintain those areas with a listing and designation of critical habitat. In particular, the Navy expressed concern over their ability to conduct maintenance dredging and requested we consult with them prior to proposing critical habitat.

*Response:* A rule to list Nassau grouper will require federal agencies to assess whether any actions implemented, authorized, or funded within the range of the species “may affect” Nassau grouper, and consult with NMFS to ensure their actions are not likely to jeopardize the species’ continued existence. The rule-making process for identifying critical habitat is separate from this final listing rule and would include opportunities for public participation and input, as well as coordination with all military branches. Unlike ESA listing decisions, the designation of critical habitat requires us to consider economic, national security, and other impacts of the designation.

*Comment 8:* One commenter opposed the proposed rule to list Nassau grouper as a threatened species stating this is “merely a precursor to an attempt to form a basis for a push for Marine Protection Areas.”

*Response:* The proposed rule to list Nassau grouper was the result of the petition we received from WildEarth Guardians, our 90-day finding that the petition presented substantial information that listing may be warranted, and our 12-month finding that listing as a threatened species was warranted. Section 4(b)(1)(A) of the ESA requires us to make listing

determinations based solely on the best scientific and commercial data available after conducting a review of the status of the species and after taking into account efforts being made by any state or foreign nation to protect the species. We have not proposed any additional regulations affecting management of Nassau grouper as a result of the proposed listing rule. However, we will need to determine whether we can identify critical habitat for this species, and if so, make an appropriate designation of critical habitat. A critical habitat designation could have implications for fishing activities. Any designation of critical habitat would include opportunities for public input. As previously mentioned, we could also implement additional protective regulations for Nassau grouper under section 4(d) of the ESA, if we determine they are necessary and advisable for the conservation of this threatened species. Issuance of a 4(d) rule would be a separate rule-making process that would include specific opportunities for public input.

#### Changes From the Proposed Rule

In addition to responding to the comments, we made a number of changes in this final rule. These included making revisions to the Biological Review section (most notably in the Population Structure and Genetics, and the Fishing Impacts on Spawning Aggregations subsections), including a more detailed description of our role in the Threats Evaluation, providing more detail in the Extinction Risk Analysis section, and clarifying the role of foreign conservation measures as they relate to making our final listing determination. We made several of these changes to provide clarity on how we reached our listing determination in response to the comment that, “. . . Nassau grouper is likely to be at risk of extinction in the foreseeable future, and may in fact be in danger of extinction now.”

#### Biological Review

This section provides a summary of key biological information presented in the Biological Report (Hill and Sadovy de Mitcheson 2013), which provides the baseline context and foundation for our listing determination.

#### Species Description

The Nassau grouper, *E. striatus* (Bloch 1792), is a long-lived, moderate sized serranid fish with large eyes and a robust body. Coloration is variable, but adult fish are generally buff, with five dark brown vertical bars, a large black saddle blotch on top of the base of the

tail, and a row of black spots below and behind each eye. Color pattern can also change within minutes from almost white to bicolored to uniformly dark brown, according to the behavioral state of the fish (Longley 1917, Colin 1992, Heemstra and Randall 1993, Carter *et al.* 1994). A distinctive bicolor pattern is seen when two adults or an adult and large juvenile meet and is frequently observed at spawning aggregations (Heemstra and Randall 1993). There is also a distinctive dark tuning-fork mark that begins at the front of the upper jaw, extends back between the eyes, and then divides into two branches on top of the head behind the eyes. Another dark band runs from the tip of the snout through the eye and then curves upward to meet its corresponding band from the opposite side just in front of the dorsal fin. Juveniles exhibit a color pattern similar to adults (*e.g.*, Silva Lee 1977).

Maximum age has been estimated as 29 years, based on an ageing study using sagittal otoliths (Bush *et al.* 2006). Most studies indicate a rapid growth rate for juveniles, which has been estimated to be about 10 mm/month total length (TL) for small juveniles, and 8.4 to 11.7 mm/month TL for larger juveniles (Beets and Hixon 1994, Eggleston 1995). Maximum size is about 122 cm TL and maximum weight is about 25 kg (Heemstra and Randall 1993, Humann and Deloach 2002, Froese and Pauly 2010). Generation time (the interval between the birth of an individual and the subsequent birth of its first offspring) is estimated as 9–10 years (Sadovy and Eklund 1999).

#### Distribution

The Nassau grouper’s confirmed distribution currently includes “Bermuda and Florida (USA), throughout the Bahamas and Caribbean Sea” (*e.g.*, Heemstra and Randall 1993). The occurrence of Nassau grouper from the Brazilian coast south of the equator as reported in Heemstra and Randall (1993) is “unsubstantiated” (Craig *et al.* 2011). The Nassau grouper has been documented in the Gulf of Mexico, at Arrecife Alacranes (north of Progreso) to the west off the Yucatan Peninsula, Mexico, (Hildebrand *et al.* 1964). Nassau grouper is generally replaced ecologically in the eastern Gulf by red grouper (*E. morio*) in areas north of Key West or the Tortugas (Smith 1971). They are considered a rare or transient species off Texas in the northwestern Gulf of Mexico (Gunter and Knapp 1951 in Hoese and Moore 1998). The first confirmed sighting of Nassau grouper in the Flower Garden Banks National Marine Sanctuary, which is located in the northwest Gulf of Mexico

approximately 180 km southeast of Galveston, Texas, was reported by Foley *et al.* (2007). Many earlier reports of Nassau grouper up the Atlantic coast to North Carolina have not been confirmed. The Biological Report (Hill and Sadovy de Mitcheson, 2013) provides a detailed description of their distribution.

#### *Habitat and Depth*

The Nassau grouper is primarily a shallow-water, insular fish species that has long been valued as a major fishery resource throughout the wider Caribbean, South Florida, Bermuda, and the Bahamas (Carter *et al.* 1994). The Nassau grouper is considered a reef fish, but it transitions through a series of developmental shifts in habitat. As larvae, they are planktonic. After an average of 35–40 days and at an average size of 32 mm TL, larvae recruit from an oceanic environment into demersal habitats (Colin 1992, Eggleston 1995). Following settlement, juvenile Nassau grouper inhabit macroalgae (primarily *Laurencia* spp.), coral clumps (*Porites* spp.), and seagrass beds (Eggleston 1995, Dahlgren 1998). Recently-settled Nassau grouper have also been collected from rubble mounds, some from tilefish (*Malacanthus plumieri*), at 18 m depth (Colin *et al.* 1997). Post-settlement, small Nassau grouper have been reported with discarded queen conch shells (*Strombus gigas*) and other debris around *Thalassia* beds (Randall 1983, Eggleston 1995).

Juvenile Nassau grouper (12–15 cm TL) are relatively solitary and remain in specific areas for months (Bardach 1958). Juveniles of this size class are associated with macroalgae, and both natural and artificial reef structure. As juveniles grow, they move progressively to deeper areas and offshore reefs (Tucker *et al.* 1993, Colin *et al.* 1997). Schools of 30–40 juveniles (25–35 cm TL) were observed at 8–10 m depths in the Cayman Islands (Tucker *et al.* 1993). No clear distinction can be made between types of adult and juvenile habitats, although a general size segregation with depth occurs—with smaller Nassau grouper in shallower inshore waters (3.7–16.5 m) and larger individuals more common on deeper (18.3–54.9 m) offshore banks (Bardach *et al.* 1958, Cervigón 1966, Silva Lee 1974, Radakov *et al.* 1975, Thompson and Munro 1978).

Recent work by Nemeth and coworkers in the U. S. Virgin Islands (U.S.V.I.; manuscript, in prep) found more overlap in home ranges of smaller juveniles compared to larger juveniles and adults have larger home ranges with less overlap. Mean home range of adult

Nassau grouper in the Bahamas was  $18,305 \text{ m}^2 \pm 5,806$  (SD) with larger ranges at less structurally-complex reefs (Bolden 2001). The availability of habitat and prey was found to significantly influence home range of adults (Bolden 2001).

Adult Nassau grouper tend to be relatively sedentary and are generally associated with high-relief coral reefs or rocky substrate in clear waters to depths of 130 m. Generally, adults are most common at depths less than 100 m (Hill and Sadovy de Mitcheson, 2013) except when at spawning aggregations where they are known to descend to depths of 255 m (Starr *et al.* 2007).

#### *Diet and Feeding*

Adult Nassau grouper are unspecialized, bottom-dwelling, ambush-suction predators (Randall 1965, Thompson and Munro 1978). Numerous studies describe adult Nassau grouper as piscivorous (Randall and Brock 1960, Randall 1965, Randall 1967, Carter *et al.* 1994, Eggleston *et al.* 1998). Feeding can take place around the clock although most fresh food is found in stomachs collected in the early morning and at dusk (Randall 1967). Young Nassau grouper (20.2–27.2 mm standard length; SL) feed on a variety of plankton, including pteropods, amphipods, and copepods (Greenwood 1991, Grover *et al.* 1998).

#### *Population Structure and Genetics*

Early genetic analyses indicated high gene flow throughout the geographic range of Nassau grouper but were unable to determine the relative contributions of populations (Hinegardner and Rosen 1972, Hateley 2005). A study of Nassau grouper genetic population structure, using mitochondrial DNA (mtDNA) and nuclear microsatellite DNA, revealed no clearly defined population substructuring based on samples from Belize, Cuba, Bahamas, and Florida. These data indicated that spawning aggregations are not exclusively self-recruiting and that larvae can disperse over great distances, but the relative importance of self-recruitment and larval immigration to local populations was unclear (Sedberry *et al.* 1996). Similarly, a study by Hateley (2005) that analyzed samples from Belize, Bahamas, Turks and Caicos, and Cayman Islands using enzyme electrophoresis indicated low to intermediate levels of genetic variability. Results from this study provided no evidence for population substructuring by sex or small-scale spatial distribution, or for macrogeographic stock separation. These results are consistent with a

single panmictic population within the northern Caribbean basin with high gene flow through the region.

A recent study, published subsequent to the Biological Report, analyzed genetic variation in mtDNA, microsatellites, and single nucleotide polymorphisms for Nassau grouper (Jackson *et al.* 2014). The study identified three potential “permeable” barriers to dispersal and concluded that large-scale oceanographic patterns likely influence larval dispersal and population structuring (regional genetic differentiation). However, the evidence of population structuring was limited. In pairwise analyses of genetic distance between the sample populations (using *F*<sub>st</sub> for microsatellites and  $\Phi$ <sub>st</sub> for mtDNA), zero (of 171) comparisons based on microsatellite DNA were statistically significant, only 47 (of 153) comparisons based on mtDNA were statistically significant (*p* < 0.00029), and there was no indication of isolation by distance in any of the genetic datasets. Overall, while this study indicated some instances of genetic differentiation, the results do not indicate a high degree of population structuring across the range. When the Jackson *et al.* study is considered in the context of the larger body of literature, there remains some uncertainty as to population substructuring for Nassau grouper.

#### *Reproductive Biology*

The Nassau grouper was originally considered to be a monandric protogynous hermaphrodite, meaning males derive from adult females that undergo a change in sex (Smith 1971, Claro *et al.* 1990, Carter *et al.* 1994). While it is taxonomically similar to other hermaphroditic groupers, the Nassau grouper is now primarily considered a gonochore with separate sexes (Sadovy and Colin 1995). Juveniles were found to possess both male and female tissue, indicating they can mature directly into either sex (Sadovy and Colin 1995). Other characteristics such as the strong size overlap between males and females, the presence of males that develop directly from the juvenile phase, the reproductive behavior of forming spawning aggregations, and the mating system were found to be inconsistent with the protogynous reproductive strategy (Colin 1992, Sadovy and Colin 1995).

Both male and female Nassau grouper typically mature at 4–5 years of age and at lengths between 40 and 45 cm SL (44 and 50 cm TL). Size, rather than age, may be the major determinant of sexual maturation (Sadovy and Eklund 1999).

Nassau grouper raised from eggs in captivity matured at 40–45 cm SL (44–50 cm TL) in just over 2 years (Tucker and Woodward 1994). Yet, the minimum age at sexual maturity based on otoliths is between 4 and 8 years (Bush *et al.* 1996, 2006). Most fish have spawned by age 7+ years (Bush *et al.* 2006).

Fecundity estimates vary by location throughout the Caribbean. Mean fecundity estimates are generally between 3 and 5 eggs/mg of ripe ovary. For example, Carter *et al.* (1994) found female Nassau grouper between 30–70 cm SL from Belize yielded a mean relative fecundity of 4.1 eggs/mg ovary weight and a mean total number of 4,200,000 oocytes (range = 350,000 – 6,500,000). Estimated number of eggs in the ripe ovary (90.7 g) of a 44.5 cm SL Nassau grouper from Bermuda was 785,101 (Bardach *et al.* 1958). In the U.S.V.I., mean fecundity was 4.97 eggs/mg of ovary (s.d. = 2.32) with mean egg production of 4,800,000 eggs (Olsen and LaPlace 1979); however, this may be an overestimate as it included premature eggs that may not develop. Fecundity estimates based only on vitellogenic oocytes, from fish captured in the Bahamas indicated a mean relative fecundity of 2.9 eggs/mg ripe ovary (s.d. = 1.09; n = 64) and a mean egg production of 716,664 (range = 11,724 – 4,327,440 for females between 47.5–68.6 cm SL). Estimates of oocyte production from Nassau grouper induced to spawn in captivity are closer to the lower estimates based solely on vitellogenic oocyte counts.

#### *Spawning Behavior and Habitat*

Nassau grouper form spawning aggregations at predictable locations around the winter full moons, or between full and new moons (Smith 1971, Colin 1992, Tucker *et al.* 1993, Aguilar-Perera 1994, Carter *et al.* 1994, Tucker and Woodward 1994). Aggregations consist of hundreds, thousands, or, historically, tens of thousands of individuals. Some aggregations have persisted at known locations for periods of 90 years or more (see references in Hill and Sadovy de Mitcheson 2013). Pair spawning has not been observed.

About 50 individual spawning aggregation sites have been recorded, mostly from insular areas in the Bahamas, Belize, Bermuda, British Virgin Islands, Cayman Islands, Cuba, Honduras, Jamaica, Mexico, Puerto Rico, Turks and Caicos, and the U.S.V.I.; however, many of these may no longer form (Figure 10 in Hill and Sadovy de Mitcheson 2013). Recent evidence suggests that spawning is occurring at

what may be reconstituted or novel spawning sites in both Puerto Rico and the U.S.V.I. (Hill and Sadovy de Mitcheson 2013). Suspected or anecdotal evidence also identifies spawning aggregations in Los Roques, Venezuela (Boomhower *et al.* 2010) and Old Providence in Colombia's San Andrés Archipelago (Prada *et al.* 2004). Neither aggregation nor spawning has been reported from South America, despite the fact ripe Nassau grouper are frequently caught in certain areas (F. Cervigón, Fundacion Cientifica Los Roques-Venezuela, pers. comm. to Y. Sadovy, NMFS, 1991). Spawning aggregation sites have not been reported in the Lesser Antilles, Central America south of Honduras, or Florida.

“Spawning runs,” or movements of adult Nassau grouper from coral reefs to spawning aggregation sites, were first described in Cuba in 1884 by Vilaro Diaz, and later by Guitart-Manday and Juarez-Fernandez (1966). Nassau grouper migrate to aggregation sites in groups numbering between 25 and 500, moving parallel to the coast or along shelf edges or even inshore reefs (Colin 1992, Carter *et al.* 1994, Aguilar-Perera and Aguilar-Davila 1996, Nemeth *et al.* 2009). Distance traveled by Nassau grouper to aggregation sites is highly variable; some fish move only a few kilometers (km), while others move up to several hundred km (Colin 1992, Carter *et al.* 1994, Bolden 2000). Ongoing research in the Exuma Sound, Bahamas has tracked migrating Nassau grouper up to 200 km, with likely estimates of up to 330 km, as they move to aggregation sites (Hill and Sadovy de Mitcheson 2013).

Observations suggest that individuals can return to their original home reef following spawning. Bolden (2001) reported 2 out of 22 tagged fish returning to home reefs in the Bahamas one year after spawning. Sonic tracking studies around Little Cayman Island have demonstrated that spawners may return to the aggregation site in successive months with returns to their residential reefs in between (Semmens *et al.* 2007). Sixty percent of fish tagged at the west end spawning aggregation site in Little Cayman in January 2005 returned to the same aggregation site in February 2005 (Semmens *et al.* 2007). Larger fish are more likely to return to aggregation sites and spawn in successive months than smaller fish (Semmens *et al.* 2007).

It is not known how Nassau grouper select and locate aggregation sites or why they aggregate to spawn. Spawning aggregation sites are typically located near significant geomorphological features, such as projections

(promontories) of the reef as little as 50 m from the shore, and close to a drop-off into deep water over a wide (6–60 m) depth range (Craig 1966, Smith 1972, Burnett-Herkes 1975, Olsen and LaPlace 1979, Colin *et al.* 1987, Carter 1989, Fine 1990, Beets and Friedlander 1998, Colin 1992, Aguilar-Perera 1994). Sites are characteristically small, highly circumscribed areas, measuring several hundred meters in diameter, with soft corals, sponges, stony coral outcrops, and sandy depressions (Craig 1966, Smith 1972, Burnett-Herkes 1975, Olsen and LaPlace 1979, Colin *et al.* 1987, Carter 1989, Fine 1990, Beets and Friedlander 1999, Colin 1992, Aguilar-Perera 1994). Recent work has identified geomorphological similarities in spawning sites that may be useful in applying remote sensing techniques to discover previously unknown spawning sites (Kobara and Heyman 2010).

The link between spawning sites and settlement sites is also not well understood. Researchers speculate the location of spawning sites assists offshore transport of fertilized eggs. However, currents nearby aggregation sites do not necessarily favor offshore egg transport, indicating some locations may be at least partially self-recruiting (*e.g.*, Colin 1992). In a study around a spawning aggregation site at Little Cayman, surface velocity profile drifters released on the night of peak spawning tended to remain near or returned to the spawning reef due to eddy formation, while drifters released on the days preceding the peak spawn tended to move away from the reef in line with the dominant currents (Heppell *et al.* 2011).

Spawning aggregations form around the full moon between December and March (reviewed in Sadovy and Eklund 1999), though this may occur later (May–August) in more northerly latitudes (La Gorce 1939, Bardach *et al.* 1958, Smith 1971, Burnett-Herkes 1975). The formation of spawning aggregations is triggered by a very narrow range of water temperatures between 25°–26 °C. While day length has also been considered as a trigger for aggregation formation (Colin 1992, Tucker *et al.* 1993, Carter *et al.* 1994), temperature is evidently a more important stimulus (Hill and Sadovy de Mitcheson 2013). The narrow range of water temperature is likely responsible for the later reproductive season in more northerly latitudes like Bermuda.

Spawning occurs for up to 1.5 hours around sunset for several days (Whaylen *et al.* 2007). At spawning aggregation sites, Nassau grouper tend to mill around for a day or two in a “staging area” adjacent to the core area where

spawning activity later occurs (Colin 1992, Kadison *et al.* 2010, Nemeth 2012). Courtship is indicated by two behaviors that occur late in the afternoon: “following” and “circling” (Colin 1992). The aggregation then moves into deeper water shortly before spawning (Colin 1992, Tucker *et al.* 1993, Carter *et al.* 1994). Progression from courtship to spawning may depend on aggregation size, but generally fish move up into the water column, with an increasing number exhibiting the bicolor phase (Colin 1992, Carter *et al.* 1994).

Spawning involves a rapid horizontal swim or a “rush” of bicolor fish following dark fish closely in either a column or cone rising to within 20–25 m of the water surface where group-spawning occurs in sub-groups of 3–25 fish (Olsen and LaPlace 1979, Carter 1986, Aguilar-Perera and Aguilar-Davila 1996). Following the release of sperm and eggs, there is a rapid return of the fragmented sub-group to the bottom. All spawning events have been recorded within 20 minutes of sunset, with most within 10 minutes of sunset (Colin 1992).

Repeated spawning occurs at the same site for up to three consecutive months generally around the full moon or between the full and new moons (Smith 1971, Colin 1992, Tucker *et al.* 1993, Aguilar-Perera 1994, Carter *et al.* 1994, Tucker and Woodward 1994). Participation by individual fish across the months is unknown. Examination of female reproductive tissue suggests multiple spawning events across several days at a single aggregation (Smith 1972, Sadovy, NMFS, pers. obs.). A video recording shows a single female in repeated spawning rushes during a single night, repeatedly releasing eggs (Colin 1992). It is unknown whether a single, mature female will spawn continuously throughout the spawning season or just once per year.

#### Status Assessments

Few formal stock assessments have been conducted for the Nassau grouper. The most recent published assessment, conducted in the Bahamas, indicates fishing effort, and hence fishing mortality (F), in the Bahamas needs to be reduced from the 1998–2001 levels, otherwise the stocks are likely to be overexploited relative to biological reference points (Cheung *et al.* 2013). The population dynamic modeling by Cheung *et al.* (2013) found: “assuming that the closure of the spawning aggregation season is perfectly implemented and enforced, the median value of  $F_{SPR}$  (the fishing mortality rate that produces a certain spawning

potential ratio) = 35 percent on non-spawning fish would be 50 percent of the fishing mortality of the 1998 to 2001 level. The 5 percent and 95 percent confidence limits are estimated to be less than 20 percent and more than 100 percent of the fishing mortality at the 1998 to 2001 level, respectively. In other words, if (1) fishing mortality (F) rates of non-spawning fish are maintained at the 1998 to 2001 level, and (2) fishing on spawning aggregations is negligible, the median spawning potential (spawner biomass relative to the unexploited level) is expected to be around 25 percent (5 and 95 percent confidence interval (CI) of 20 and 30 percent, respectively). This level is significantly below the reference limit of 35 percent of spawning potential, meaning that there is a high chance of recruitment overfishing because of the low spawning stock biomass.”

The Nassau grouper was formerly one of the most common and important commercial groupers in the insular tropical western Atlantic and Caribbean (Smith 1978, Randall 1983, Appeldoorn *et al.* 1987, Sadovy 1997). Declines in landings and catch per unit of effort (CPUE) have been reported throughout its range, and it is now considered to be commercially extinct (*i.e.*, the species is extinct for fishery purposes due to low catch per unit effort) in a number of areas, including Jamaica, Dominican Republic, U.S.V.I., and Puerto Rico (Sadovy and Eklund 1999). Information on past and present abundance and density, at coral reefs and aggregation sites, is based on a combination of anecdotal accounts, visual census surveys, and fisheries data. Because grouper species are reported collectively in landings data, there are limited species-specific data to determine catch of Nassau grouper throughout its range.

While fisheries dependent data are generally limited for the species throughout its range, there are some 1970s and 1980s port-sampling data from the U.S.V.I. and Puerto Rico. In the U.S.V.I., Nassau grouper accounted for 22 percent of total grouper landings, and 85 percent of the Nassau grouper catch came from spawning aggregations (D. Olsen, Chief Scientist—St. Thomas Fishermen’s Association, pers. comm. to J. Rueter, NMFS, October 2013). The first U.S. survey of the fishery resources of Puerto Rico noted the Nassau grouper was common and a very important food fish, reaching a weight of 22.7 kg or more (Evermann 1900). The Nassau grouper was still the fourth-most common shallow-water species landed in Puerto Rico in the 1970s (Thompson 1978), and it was common in the reef fish fishery of the U.S.V.I. (Olsen and

LaPlace 1979). By 1981, “the Nassau grouper ha[d] practically disappeared from the local catches and the ones that d[id] appear [were] small compared with previous years” (CFMC 1985). By 1986, the Nassau grouper was considered commercially extinct in the U.S. Caribbean (Bohnsack *et al.* 1986). About 1,000 kg of Nassau grouper landings were reported in the Puerto Rico Reef Fish Fishery during the latter half of the 1980s, and most of them were less than 50 cm indicating they were likely sexually immature (Sadovy 1997).

A number of organizations and agencies have conducted surveys to examine the status of coral reefs and reef-fish populations throughout the western Atlantic. Results from these monitoring studies offer some indication of relative abundance of Nassau grouper in various locations, although different methods are often employed and thus results of different studies cannot be directly compared (Kellison *et al.* 2009). The Atlantic and Gulf Rapid Reef Assessment Program (AGRRA), which samples a broad spectrum of western Atlantic reefs, includes few reports of Nassau grouper, as sighting frequency (proportion of all surveys with at least one Nassau grouper present) ranged from less than 1 percent to less than 10 percent per survey from 1997–2000. Density of Nassau grouper ranged from 1 to 15 fish/hectare with a mean of 5.6 fish/hectare across all areas surveyed (AGRRA). NOAA’s Coral Reef Ecosystem Monitoring Program (CREMP) has conducted studies on coral reefs in Puerto Rico and the U.S.V.I. since 2000, and sighting frequency of Nassau grouper has ranged from 0 to 0.5 percent with density between 0 to 0.5 fish/hectare. Data from SCUBA surveys conducted by the University of the Virgin Islands report a density of 4 Nassau grouper/hectare per survey across reef habitat types in the U.S.V.I. SCUBA surveys by NOAA in the Florida Keys across reef habitat types have sighting frequencies of 2–10 percent per survey, with a density of 1 Nassau grouper/hectare (NOAA’s NMFS FRVC). In addition to these surveys, Hodgson and Liebler (2002) noted that Nassau grouper were absent from 82 percent of shallow Caribbean reefs surveyed (3–10 m) during a 5-year period (1997–2001) for the ReefCheck project.

#### Fishing Impacts on Spawning Aggregations

Because we lack sufficient stock assessments or population estimates, we considered the changes in spawning aggregations as a proxy for the status of the current population. We believe the

status of spawning aggregations is likely to be reflective of the overall population because adults migrate to spawning aggregations for the only known reproductive events. Historically, 50 spawning aggregation sites had been identified throughout the Caribbean (Sadovy de Mitcheson *et al.* 2008). Of these 50, less than 20 probably still remain (Sadovy de Mitcheson *et al.* 2008). Furthermore, while numbers of fish at aggregation sites once numbered in the tens of thousands (30,000–100,000 fish; Smith 1972), they have now been reduced to less than 3,000 at those sites where counts have been made (Sadovy de Mitcheson *et al.* 2008). Based on the size and number of current spawning aggregations the Nassau grouper population appears to be just a fraction of its historical size.

In general, slow-growing, long-lived species (such as snappers and groupers) with limited spawning periods, and possibly with narrow recruitment windows, are susceptible to overexploitation (Bannerot *et al.* 1987, Polovina and Ralston 1987). The strong appeal of spawning aggregations as targets for fishing, their importance in many seasonal fisheries, and the apparent abundance of fish at aggregations make spawning aggregations particularly susceptible to over-exploitation. There are repeated reports from across the Caribbean where Nassau grouper spawning aggregations have been discovered and fished to the point that the aggregation ceased to form, or formed at such low densities that spawning was no longer viable. For example, the commercial fishing of Nassau grouper aggregations in Bermuda resulted in decreased landings from 75,000 tons in 1975 to 10,000 tons by 1981 (Luckhurst 1996, Sadovy de Mitcheson and Erisman 2012). The four known spawning aggregation sites in Bermuda ceased to form shortly thereafter and have yet to recover (Sadovy de Mitcheson and Erisman 2012). However, Nassau grouper are still present in Bermuda and reported observations have slightly increased over the last 10–15 years (B. Luckhurst, Bermuda Department of Agriculture, Fisheries, and Parks, Division of Fisheries, pers. comm. to Y. Sadovy, University of Hong Kong, 2012). In Puerto Rico, historical spawning aggregations no longer form, though a small aggregation has recently been found, and may be a reconstitution of one of the former aggregations (Schärer *et al.* 2012). In Mahahual, Quintana Roo, Mexico, aggregations of up to 15,000 fish formed each year, but due to increased fishing pressure in the 1990's,

aggregations have not formed in Mahahual since 1996 (Aguilar-Perera 2006). Inadequate enforcement of management measures designed to protect spawning aggregations in Mexico has further affected aggregations (Aguilar-Perera 2006), though at least three aggregation sites remain viable. In Cuba, Nassau grouper were almost exclusively targeted during aggregation formation; because of this, there have been severe declines in the number of Nassau grouper at 8 of the 10 aggregations and moderate declines in the other 2 (Claro *et al.* 2009). Similar situations are known to have occurred in the Bahamas, U.S.V.I., Puerto Rico, and Honduras (Sadovy de Mitcheson and Erisman 2012, see also Hill and Sadovy de Mitcheson 2013).

Overexploitation has also occurred in Belize. Between 1975 and 2001 there was an 80 percent decline in the number of Nassau grouper (15,000 fish to 3,000) at the Glover's Reef aggregation (Sala *et al.* 2001). Additionally, a 2001 assessment concluded that only 2 of the 9 aggregation sites identified in 1994 remained viable, and those had been reduced from 30,000 fish to 3,000–5,000 fish (Heyman 2002). More recent monitoring (2003–2012) at the two sites at Glover's Reef indicates further declines in the sizes of these aggregations. A maximum of 800–3,000 Nassau grouper were counted per year at these sites over the ten years of monitoring (Belize SPAG Working Group 2012).

Further indicators of population decline through over-exploitation include reduced size and/or age of fish harvested compared to maximum sizes and ages. Nassau grouper can attain sizes of greater than 120 cm (Heemstra and Randall 1993, Humann and Deloach 2002, Froese and Pauly 2010) and live as long as 29 years (Bush *et al.* 2006). However, it is unusual to obtain individuals of more than 12 years of age in exploited fisheries, and more heavily fished areas yield much younger fish on average. The maximum age estimates in heavily exploited areas are depressed—9 years in the U.S.V.I. (Olsen and LaPlace 1979), 12 years in northern Cuba, 17 years in southern Cuba (Claro *et al.* 1990), and 21 years in the Bahamas (Sadovy and Colin 1995). Similarly, there is some indication that size at capture of both sexes declined in areas of higher exploitation versus unexploited populations within a specific region (Carter *et al.* 1994). When exploitation is high, catches are largely comprised of juveniles. For example, most catches of Nassau grouper in heavily exploited areas of Puerto Rico, Florida (Sadovy and Eklund 1999), and

Cuba (Espinosa 1980) consisted of juveniles. In exploited U.S.V.I. aggregations, harvest of Nassau grouper larger than 70 cm TL was uncommon (Olsen and LaPlace 1979).

While direct fishing of spawning aggregations was a primary driver of Nassau grouper population declines as indicated by the observed declines in spawning aggregations (Sadovy de Mitcheson and Erisman 2012), other factors also affect abundance. For example, removal of adults from spawning runs and intensive capture of juveniles, either through direct targeting (*e.g.*, spearfishing) or using small mesh traps or nets, also occur (Hill and Sadovy de Mitcheson 2013). In addition to the high fishing pressure in some areas, poaching also appears to be affecting some populations (*e.g.*, in the Cayman Islands; Semmens *et al.* 2012).

#### *NMFS's Conclusions From the Biological Report*

*The species is made up of a single population over its entire geographic range.* As summarized above, multiple genetic analyses indicate that there is high gene flow throughout the geographic range of the Nassau grouper, and no clearly defined population substructuring has been identified (Hinegardner and Rosen 1972, Sedberry *et al.* 1996, Hateley 2005). Although a recent study (Jackson *et al.* 2014) reported genetic differentiation, it does not provide evidence to support biological differences between populations. We believe further studies are needed to verify and expand upon the work presented by Jackson *et al.* (2014). Based on the best available information, we conclude there is a single population of Nassau grouper throughout the Caribbean.

*The species has patchy abundance, with declines identified in many areas.* The Biological Report describes the reduction in both size and number of spawning aggregations throughout the range. Patchy abundance throughout the range of a species is common due to differences in habitat quality/quantity or exploitation levels at different locations. However, dramatic, consistent declines of Nassau grouper have been noted throughout its range. In many areas throughout the Caribbean, the species is now considered commercially extinct and numerous spawning aggregations have been extirpated with no signs of recovery.

*The species possesses life history characteristics that increase vulnerability to harvest, including slow growth to a large size, late maturation, formation of large spawning aggregations, and occurrence in shallow*

*habitat*. This conclusion is based on the Description of the Species in the Biological Report (Hill and Sadovy de Mitcheson 2013). Slow growth and late maturation expose sub-adults to harvest prior to reproduction. Sub-adult and adult Nassau grouper form large conspicuous spawning aggregations. These aggregations are often in shallow habitat areas that are easily accessible to fishermen and thus heavily exploited. Despite these life-history vulnerabilities, there are remaining spawning aggregations that, while reduced in size and number, still function and provide recruits into the population.

*The species is broadly distributed, and its current range is similar to its historical range.* The Range-wide Distribution section of the Biological Report (Hill and Sadovy de Mitcheson 2013) concluded that the current range is equivalent to the historical range, though abundance has been severely depleted.

### Threats Evaluation

The threats evaluation was the second step in the process of making an ESA listing determination for Nassau grouper as described above in "Listing Determinations under the ESA". The Extinction Risk Analysis Group (ERAG), which consisted of 12 NOAA Fisheries Science Center and Regional Office personnel, was asked to independently review the Biological Report and assess 4 demographic factors (abundance, growth rate/productivity, spatial structure/connectivity, and diversity) and 13 specific threats (see ERA Threat Table under supporting documents). The group members were asked to provide qualitative scores based on their perceived severity of each factor and threat.

Members of the ERAG were asked to independently evaluate the severity, scope, and certainty for these threats currently and in the foreseeable future (30 years from now). The foreseeable future was based on the upper estimate of generation time for Nassau grouper (9–10 years) as described by Sadovy and Eklund (1999) and an age at maturity of 8 years (Bush *et al.* 1996, 2006). We chose 30 years, which would potentially allow recruitment of 2–3 generations of mature individuals to appear in spawning aggregations as a result of fishery management actions. Given the limited information we have to predict the impacts of threats, we felt the 30 year timeframe was the most appropriate to assess threats in the foreseeable future.

Members of the ERAG were asked to rank each of four demographic factors and 13 identified threats as "very low

risk," "low risk," "moderate risk," "increasing risk," "high risk," or "unknown." "Very low risk" meant that it is unlikely that the demographic factor or threat affects the species' overall status. "Low risk" meant that the demographic factor may affect species' status, but only to a degree that it is unlikely that this factor significantly elevates risk of extinction now or in the future. "Moderate risk" meant that the demographic factor or threat contributes significantly to long term risk of extinction, but does not constitute a danger of extinction in the near future. "Increasing risk" meant that the present demographic risk or threat is low or moderate, but is likely to increase to high risk in the foreseeable future if present conditions continue. Finally, "high risk" meant that the demographic factor or threat indicates danger of extinction in the near future. Each member of the ERAG evaluated risk on this scale, and we then interpreted these rankings against the statutory language for threatened or endangered to determine the status of Nassau grouper. We did not directly relate the risk levels with particular listing outcomes, because the risk levels alone are not very informative. Acknowledging the differences in terminology between the ERAG risk scale and the ESA statutory definitions of threatened and endangered, we relied upon our own judgment and expertise in reviewing the ERA to determine the status of Nassau grouper and form our final listing determination.

ERAG members were also asked to consider the potential interactions between demographic factors and threats. If the demographic factor or threat was ranked higher due to interactions with other demographic factors or threats, each member was asked to then identify those factors or threats that caused them to score the risk higher or lower than it would have been if it were considered independently. We then examined the independent responses from each ERAG member for each demographic factor and threat and used the modal response to determine the level of threat to Nassau grouper.

Climate change and international trade regulations (*e.g.*, the Convention on International Trade in Endangered Species (CITES), as described in the Biological Report) were categorized by the ERAG as "unknown." Habitat alteration, U.S. federal regulations, disease/parasites/abnormalities, and aquaculture were ranked as "very low risk" to "low risk." State/territorial regulations, growth rate/productivity, abundance, spatial structure/

connectivity, commercial harvest, foreign regulations, artificial selection, and diversity were ranked as "moderate risk" to "increasing risk." Historical harvest (the effect of prior harvest on current population status), fishing at spawning aggregations, and inadequate law enforcement were classified as "high risk." The demographic factors and threats are described below by the five ESA factors with the corresponding ERAG ranking and our analysis.

#### A. The Present or Threatened Destruction, Modification, or Curtailment of Its Habitat or Range

Spatial structure/connectivity and habitat alteration were considered under ESA Factor A; this included habitat loss or degradation, and the loss of habitat patches, critical source populations, subpopulations, or dispersal among populations.

Nassau grouper use many different habitat types within the coral reef ecosystem. The increase in urban, industrial, and tourist developments throughout the species range impacts coastal mangroves, seagrass beds, estuaries, and live coral (Mahon 1990). Loss of juvenile habitat, such as macroalgae, seagrass beds, and mangrove channels is likely to negatively affect recruitment rates. Habitat alteration was ranked by the ERAG as a "low risk" threat to Nassau grouper. We agree with the ERAG that habitat alteration presents a low risk to the species and is unlikely to contribute to the threat of extinction presently or over the foreseeable future. The use of many different habitat types by Nassau grouper may spread the risk of impacts associated with habitat loss to a point that reduces overall extinction risk to the species.

The range of Nassau grouper is influenced by spatial structure and connectivity of the population. As described in Hill and Sadovy de Mitcheson (2013), a study of genetic population structure in Nassau grouper revealed no clearly defined population substructuring at the geographic locations sampled, *i.e.*, Belize, Cuba, Bahamas, and Florida (Sedberry *et al.* 1996). Based on ERAG scores, spatial structure/connectivity was characterized as an "increasing" risk for Nassau grouper. We agree with the ERAG ranking and believe this increasing risk is due, in part, to the declining number and size of spawning aggregations, which affects population structure. Given the increasing risk associated with this demographic factor we believe it could lead the species to become endangered over the foreseeable future.

### *B. Overutilization for Commercial, Recreational, Scientific, or Educational Purposes*

Based on ERAG rankings, historical harvest and fishing at spawning aggregations are two of the three most severe threats (the third being inadequate law enforcement) to Nassau grouper. Historical harvest and fishing at spawning aggregations were both classified as “high” risk threats to Nassau grouper. Curiously, the ERAG rankings for commercial harvest, which often includes the fishing on spawning aggregations, were lower and indicated current commercial harvest was a “moderate” threat for Nassau grouper. We believe this lower ranking may be related to the fact that the species has declined to the point that commercial harvest is not as large a threat as in decades past. This is also related to abundance which was similarly classified as a “moderate” risk for Nassau grouper.

Two different aspects of fishing affect Nassau grouper abundance: Fishing effort throughout the non-spawning months and directed fishing at spawning aggregations or on migrating adults. In some countries Nassau grouper are fished commercially and recreationally throughout the year by handline, longline, fish traps, spear guns, and gillnets (NMFS General Canvas Landing System). Fishing at spawning aggregations is mainly conducted by handlines or by fish traps, although gillnets were being used in Mexico in the early to mid-1990s (Aguilar-Perera 2004). Declines in landings, catch per unit effort (CPUE) and, by implication, abundance in the late 1980s and early 1990s occurred throughout its range, which has led Nassau grouper to now be considered commercially extinct in a number of areas (Sadovy and Eklund 1999). Population declines and loss of spawning aggregations continue throughout the Nassau grouper’s range (Sadovy de Mitcheson 2012).

We agree with the ERAG’s assessment for the threat of abundance. It is clear that the abundance of Nassau grouper has diminished dramatically over the past several decades. This decline is a direct impact of historical harvest and the overfishing of spawning aggregations. The current abundance of Nassau grouper is not causing or contributing to the species currently being in danger of extinction but does raise concern for the status of the species over the foreseeable future if abundance continues to decline.

We disagree with the ERAG’s “high risk” rating for historical harvest. We

believe that while historical harvest has reduced the population size of Nassau grouper, which has in turn affected the ability of the population to recover, we don’t agree that this threat continues to be a “high risk”. It seems more appropriate to consider the ERAG’s risk assessment for the abundance of the current population in making our listing determination.

Predictable spawning aggregations make Nassau grouper a vulnerable fishing target. In many places, annual landings for Nassau grouper were mostly from aggregation-fishing (e.g., Claro *et al.* 1990, Bush *et al.* 2006). Because Nassau grouper are only known to reproduce in spawning aggregations, removing ripe individuals from the spawning aggregations greatly influences population dynamics and future fishery yields (Shapiro 1987). Harvesting a species during its reproductive period increases adult mortality and diminishes juvenile recruitment rates. The loss of adults and the lack of recruitment greatly increase a species’ extinction risk. The collapse of aggregations in many countries (Sadovy de Mitcheson 2012) was likely a result of overharvesting fish from spawning aggregations (Olsen and LaPlace 1979, Aguilar-Perera 1994, Sadovy and Eklund 1999). As Semmens *et al.* (2012) noted from the results of a mark-recapture study on Cayman Brac, Cayman Island fishermen appear to catch sufficient adult grouper outside the spawning season to seriously impact population size. It appears that fishing at spawning aggregations has depressed population size such that fishing operations away from the aggregations are also impacting population status.

We agree that fishing at spawning aggregations has reduced the population of Nassau grouper and has affected its current status. While the ERAG determined this is a “high risk” threat, we are less certain about our determination. We believe that this threat is in large part exacerbated by the inadequacy of regulatory mechanisms as discussed further below under Factor D. If existing regulatory mechanisms and corresponding law enforcement were adequate, this threat would be less of a concern. In the absence of adequate law enforcement, we believe that fishing at spawning aggregations is increasing the extinction risk of Nassau grouper.

The final threat analyzed for Factor B was artificial selection. The ERAG scores indicated artificial selection was a “moderate” threat; however, ranking of this threat was widely distributed amongst ERAG members, indicating a high level of uncertainty about the effects of artificial selection on Nassau

grouper. We recognize the uncertainty associated with this threat and believe more information is needed. That said, we do not believe available information indicates artificial selection is currently impacting the species’ risk of extinction.

### *C. Disease*

There is very little information on the impacts of disease, parasites, and abnormalities on Nassau grouper, yet the species is not known to be affected by any specific disease or parasite. Given this, NMFS agrees with the ERAG ranking indicating a “very low risk” threat from disease, parasites, and abnormalities. We do not believe any of these threats will rise to the level of impacting the species’ status over the foreseeable future.

### *D. Inadequacy of Existing Regulatory Mechanisms*

Consideration of the inadequacy of existing regulatory mechanisms, includes whether enforcement of those mechanisms is adequate. The relevance of existing regulatory mechanisms to extinction risk for an individual species depends on the vulnerability of that species to each of the threats identified under the other factors of ESA section 4, and the extent to which regulatory mechanisms could or do control the threats that are contributing to the species’ extinction risk. If a species is not currently, and not expected within the foreseeable future to become, vulnerable to a particular threat, it is not necessary to evaluate the adequacy of existing regulatory mechanisms for addressing that threat. Conversely, if a species is vulnerable to a particular threat (now or in the foreseeable future), we do evaluate the adequacy of existing measures, if any, in controlling or mitigating that threat. In the following paragraphs, we will discuss existing regulatory mechanisms for addressing the threats to Nassau grouper generally, and assess their adequacy for controlling those threats. In the Extinction Risk Analysis section, we determine if the inadequacy of regulatory mechanisms is a contributing factor to the species’ status as threatened or endangered because the existing regulatory mechanisms fail to adequately control or mitigate the underlying threats.

#### *Summary of Existing Regulatory Mechanisms*

As discussed in detail in the Biological Report (Hill and Sadovy de Mitcheson 2013), a wide array of regulatory mechanisms exists throughout the range of Nassau grouper that are intended to limit harvest and

thus maintain abundance. Existing regulatory mechanisms include minimum size restrictions, seasonal closures, spatial closures, and gear and access restrictions. We summarize some of these regulatory mechanisms below by country.

The Bahamas has implemented a number of regulatory mechanisms to limit harvest. In the 1980s, the Bahamas introduced a minimum size of 3 lbs. (1.36 kg) for Nassau grouper. This was followed in 1998 with a 10-day seasonal closure at several spawning aggregations. An annual “two-month” fishery closure was added in December 2003 to coincide with the spawning period and was extended to three months in 2005 to encompass the December through February spawning period. Up until 2015, the implementation of the 3-month closure was determined annually and could be shortened or otherwise influenced by such factors as the economy (Sadovy and Eklund 1999). In 2015, the annual assessment of the closure was removed ensuring a fixed 3-month closure each year moving forward (Fisheries Resources [Jurisdiction and Conservation] [Amendment] Regulations 2015). During the 3-month closure there is a national ban on Nassau grouper catches; however, the Bahamas Reef Educational Foundation (BREEF; unpub. data), has reported large numbers of fish being taken according to fisher accounts with photo-documentation and confirming reports of poaching of the species during the aggregation season.

The Bahamas has implemented several other actions that aid the conservation of Nassau grouper. There are marine parks in the Bahamas that are closed to fishing year round and therefore protect Nassau grouper. The Exuma Cays Land and Sea Park, first established in 1959, has been closed to fishing since 1986, thus protecting both nursery and adult habitat for Nassau grouper and other depleted marine species. Other sites, including the South Berry Islands Marine Reserve (established on December 29, 2008), Southwest New Providence National Park, and North Exumas Study Site have also been established and closed to fishing. Several gear restrictions in the Bahamas are also protective of Nassau grouper. Fishing with SCUBA and the use of explosives, poisons, and spearguns is prohibited in the Bahamas, although snorkeling with sling spears is allowed. The use of bleach or other noxious or poisonous substances for fishing, or possession of such substances on board a fishing vessel, without written approval of the

Minister, is prohibited. Commercial fishing in the Bahamas is restricted to only the native population and, as a consequence, all vessels fishing within the Bahamas Exclusive Fishery Zone must be fully owned by a Bahamian citizen residing in the Bahamas.

In Belize, the first measure to protect Nassau grouper was a seasonal closure within the Glover’s Reef Marine Reserve in 1993; the area was closed from December 1 to March 1 to protect spawning aggregations. A seasonal closure zone to protect Nassau grouper spawning aggregations was included when the Bacalar Chico marine reserve was established in 1996 (Paz and Truly 2007). Minimum and maximum capture sizes were later introduced (Hill and Sadovy de Mitcheson 2013 and citations therein).

In 2001 the Belize National Spawning Aggregation Working Group established protective legislation for 11 of the known Nassau grouper spawning sites within Belize. Seven of those 11 sites are monitored as regularly as possible. The Working Group meets regularly to share data and develop management strategies ([www.spagbelize.org](http://www.spagbelize.org); retrieved on 15 April 2012). In 2003, Belize introduced a four-month closed season to protect spawning fish (O’Connor 2002, Gibson 2008). However, the 2003 legislation also allowed for exemptions to the closures by special license granted by the Fisheries Administrator, provided data be taken on any Nassau grouper removed. These special licenses made it difficult to enforce the national prohibition and in 2010 Belize stopped issuing permits to fish for Nassau grouper during the 4-month spawning period, except at Maugre Caye and Northern Two Caye.

In 2009, Belize issued additional protective measures to help manage and protect the Nassau grouper. These include minimum and maximum size limits of 20 inches and 30 inches, respectively. Belize has also introduced a plan to ban spear fishing within all marine reserves (yet to be implemented). Furthermore, as a large proportion of finfish are landed as fillets, the new regulations require that all Nassau grouper be landed whole, and if filleted must have a 1-inch by 2-inch skin patch (The Belize Spawning Aggregation Working Group 2009). Other gear restrictions are in place to generally aid in the management of reef fish, such as no spearfishing on compressed air.

Although Bermuda closed red hind aggregation sites in 1974, Nassau grouper aggregation sites located seaward of these sites were not included and continued to be fished. In 1990, a

two-fish bag limit and minimum size restriction (35.6 cm FL) were enacted in Bermuda (Luckhurst 1996). Since 1996, Nassau grouper has been completely protected through a prohibition on take and possession and likely benefits from numerous no-take marine reserves (Hill and Sadovy de Mitcheson 2013).

In the Cayman Islands, the three main (“traditional”) grouper “holes” were officially protected in the late 1970’s and only residents were allowed to fish by lines during the spawning season (Hill and Sadovy de Mitcheson 2013). In 1986, increasing complaints from fishermen of a decline in both numbers and size of Nassau grouper taken from the fishery prompted the implementation of a monitoring program by the Department of the Environment (Bush *et al.* 2006).

Following the development of the monitoring program, the Cayman Islands implemented a number of management measures. In the early 1990s, legislation prohibited spearfishing at spawning aggregation sites. In 1998, the three main grouper holes at the eastern end of the islands were formally designated as “Restricted Marine Areas” where access requires licensing by the Marine Conservation Board (Bush *et al.* 2006). In February 2002, protective legislation defined a spawning season as November 1 to March 31, and an “Alternate Year Fishing” rule was passed. This law allowed fishing of the spawning aggregations to occur every other year with the first non-fishing year starting in 2003. A catch limit of 12 Nassau grouper per boat, per day during fishing years was also set. The 2002 law defined a one nautical mile (nm) “no trapping” zone around each spawning site, and set a minimum size limit of 12 inches for Nassau grouper in response to juveniles being taken by fish traps inside the sounds (Whaylen *et al.* 2004, Bush *et al.* 2006). In 2003, spearguns were restricted from use within 1 nm of any designated grouper spawning area from November through March. In 2008, it was prohibited to take any Nassau grouper by speargun anywhere in Cayman waters. Effective December 29, 2003, the Marine Conservation Board, closed fishing at all designated Nassau grouper spawning sites for a period of 8 years. The conservation measure was renewed for a further 8 years in 2011.

In Cuba, there is a minimum size limit for Nassau grouper though this regulation is largely unprotective. The minimum size of 32 cm TL (or 570g) for Nassau grouper is less than the reported average size at maturity of 50 cm TL, indicating that Nassau grouper can be harvested before having the opportunity

to reproduce. Of some benefit to Nassau grouper are more general fishing regulations such as bag limits for recreational fishing, regulations to increase selectivity of fishing gears to avoid the catch of juveniles, limits of net use during spawning aggregation time, and controls of speargun use, both commercially and recreationally. Marine protected areas have also been introduced throughout the country. In 2002, the total number of recreational licenses was limited to 3,500 for the whole country hoping to reduce directed fishing pressure nationally.

In Mexico, following scientific documentation of declines of Nassau grouper at Mahahual (Aguilar-Perera 1994), two regulations were enacted: (1) In 1993 spear-fishing was banned at any spawning aggregation site in southern Quintana Roo; and (2) in 1997 the fishing of any grouper species was banned during December and January (Aguilar-Perera 2006). Then, in 2003, a closed season for all grouper was implemented from February 15 to March 15 in all waters of the Mexican Exclusive Economic Zone. Although aimed at protecting red grouper this closure also protects Nassau grouper during a part of its spawning season (Aguilar-Perera *et al.* 2008). A management plan was to have gone into effect in 2012 to protect all commercially exploited groupers in Mexico's southern Gulf of Mexico and Caribbean Sea; yet at this time the plan has not been implemented.

In the Turks and Caicos Islands, the only documented Nassau grouper spawning aggregation site is protected from fishing in Northwest Point Marine National Park, Providenciales (DECR 2004; National Parks Ordinance and Subsidiary Legislation CAP. 80 of 1988). Similar to situations in other countries, protection of Nassau grouper habitat and spawning migration corridors on the narrow ledge of Caicos Bank is problematic as it would impose economic hardship on local fishers who depend on those areas for commercial species (e.g., spiny lobsters) and subsistence fishing (Rudd 2001).

In U.S. federal waters, including those federal waters around Puerto Rico and the U.S.V.I., take and possession of Nassau grouper have been prohibited since 1990. Since 1993, a ban on fishing/possessing Nassau grouper was implemented for the state of Florida and has since been enacted in all U.S. state waters. The species was fully protected in both state and federal waters of Puerto Rico by 2004. The Caribbean Fishery Management Council, with support of local fishermen, established a no-take marine protected area off the

southwest coast of St. Thomas, U.S.V.I. in 1990. This area, known as the Hind Bank Marine Conservation District (HBMCD), was intended to protect red hind and their spawning aggregations, as well as a former Nassau grouper spawning site (Brown 2007). The HBMCD was first subject to a seasonal closure beginning in 1990 (Beets and Friedlander 1999, Nemeth 2005, Nemeth *et al.* 2006) to protect spawning aggregations of red hind, and was later closed to fishing year-round in 1998 (DPNR 2005). Additional fishing restrictions in the U.S.V.I. such as gear restrictions, rules on the sale of fish, and protected areas such as the Virgin Islands Coral Reef National Monument and Buck Island Reef National Monument where all take is prohibited, Virgin Islands National Park (commercial fishing prohibited), and several U.S.V.I. marine reserves offer additional protection to Nassau grouper. In 2006, the U.S.V.I. instituted regulations to prohibit harvest and possession of Nassau grouper in territorial waters and filleting at sea was prohibited (García-Moliner and Sadovy 2008).

In Colombia, the San Andrés Archipelago has a number of areas that are designated as no-take fishing zones, and in 2000 the entire archipelago was declared by the United Nations Educational, Scientific and Cultural Organization (UNESCO) as the Seaflower Biosphere Reserve. In 2004, large portions of the archipelago were declared as a system of marine protected areas with varying zones of fisheries management; however, enforcement is largely lacking (M. Prada, Coralina, San Andres, Colombia, pers. comm. R. Hill, NMFS, 2010). Right-to-fish laws in Colombia also require that fishermen be allowed to fish at a subsistence level even within the no-take zones (M. Prada, Coralina, San Andres, Colombia, pers. comm. R. Hill, NMFS, 2010).

There are other Caribbean countries that have either few management measures in place or have yet to implement any conservation measures for Nassau grouper. We are not aware of special conservation or management regulations for Nassau grouper in Anguilla. In Antigua-Barbuda, while Nassau grouper is not specifically managed or protected, closed seasons were considered in 2008 for Nassau grouper and red hind, though the status of these closed seasons is not known. In the British Virgin Islands, there is a closed season for landing Nassau grouper between March 1 and May 31 (Munro and Blok 2005). In the Dominican Republic the catch and sale of ripe female Nassau grouper during

the spawning season is not allowed (Bohnsack 1989, Sadovy and Eklund 1999, Box and Bonilla Mejia 2008) and at least one marine park has been established with fishing regulations. In Guadeloupe and Martinique, there are plans to protect the species (F. Gourdin, Regional Activity Center for Specially Protected Areas and Wildlife—UNEP, pers. comm. to Y. Sadovy, University of Hong Kong, 2011) although no details are available at this time. In Honduras, there is no legislation that controls fishing in the snapper/grouper fishery; however, traps and spears are illegal in the Bay Islands. There are no Nassau grouper special regulations in Jamaica; yet, some marine protected areas were designated in 2011.

#### Analysis of Existing Regulatory Mechanisms

The ERAG considered several threats under Factor D including law enforcement, international trade regulations, foreign regulations in their jurisdictional waters, U.S. federal laws, and U.S. state and territorial laws. The ERAG determined that these threats substantially contribute to the overall risk to the species. Inadequate law enforcement was noted by several ERAG members as influencing their scoring for abundance, fishing of spawning aggregations, commercial harvest, and historical harvest. Inadequate law enforcement led to higher risk scores for each of these threats. The ERAG scored law enforcement as a “high risk” threat for Nassau grouper. ERAG rankings for the other threats were widely distributed. The inadequacy of foreign regulations in jurisdictional waters was considered an “increasing” risk while the risk of international trade regulations was “unknown.” The remaining two categories of regulations (U.S. Federal and State of Florida/U.S. territory regulations) were considered “low risk” and “moderate risk” respectively. While the ERAG rankings for threats impacting the adequacy of regulatory mechanisms were generally moderate, we believe the concern about fishing at spawning aggregations (“high risk” according to the ERAG) is due in part to the inadequacy of existing regulatory mechanisms.

Overall, we believe existing regulatory mechanisms throughout the species' range (international trade, foreign, U.S. federal, and U.S. state and territorial regulations) vary in their effectiveness, especially in addressing the most serious threat to Nassau grouper—fishing of spawning aggregations. In some countries, an array of national regulatory mechanisms, increases in marine protected areas, and customary

management may be effective at addressing fishing of spawning aggregations. For example, the Exuma Cays Land and Sea Park (Bahamas), has been closed to fishing for over 25 years and protects both nursery and adult habitat for Nassau grouper and other marine species. In that park, there is a clear difference in the number, biomass, and size of Nassau grouper in comparison to adjacent areas where fishing is permitted (Sluka *et al.* 1997).

We note, however, that many countries have few, if any, specific Nassau grouper regulations. Instead they rely on general fisheries regulations (*e.g.*, Anguilla, Antigua-Barbuda, Colombia, and Cuba all rely only on size limits, while Guadeloupe and Martinique, Honduras, Jamaica, Mexico, St. Lucia, and the Turks and Caicos rely on a variety of general fishing regulations). Additionally, where Nassau grouper-specific regulations do exist, the ERAG scores indicated that law enforcement still presents a high risk threat to the species. We agree with the ERAG's risk assessment and believe that law enforcement in many foreign countries is less than adequate, thus rendering the regulations ineffective.

Some foreign regulations may be ephemeral, unprotective of migrating adults, or inadequate to conserve the viability of a species. In some cases, regulations do not completely protect all known spawning aggregations (*e.g.*, Belize, where 2 spawning aggregations are fished by license). In another instance, we found no protections for Nassau grouper in any foreign country during the period they move to and from spawning aggregation sites. Foreign regulations in some countries specify exemptions for "historical," "local," or artisanal fishermen (*e.g.*, Colombia). Finally, some particular types of regulations are insufficient to protect the species (*e.g.*, minimum size limits in both the Bahamas and Cuba are less than size-at-maturity).

In some places, such as Bermuda, no recovery has been documented after years of regulations (B. Luckhurst, Bermuda Department of Agriculture, Fisheries, and Parks, pers. comm. to Y. Sadovy, University of Hong Kong, September, 2012). In other places (*e.g.*, Cayman Islands) there are indications of potential recovery at spawning aggregation sites, but fishing continues to keep the population depressed (Semmens *et al.* 2012) and inconsistent surveys do not provide data adequate to realize impacts. Additionally, larval recruitment is highly variable due to currents in the Caribbean basin. Some populations may receive larval input from neighboring spawning

aggregations, while other local circulation patterns may entrain larvae (Colin *et al.* 1987) making the population entirely self-recruiting.

In conclusion, although many countries have taken regulatory measures to conserve Nassau grouper, the species faces an ongoing threat due to the inadequacy of regulatory mechanisms to prevent or remediate the impacts of other threats that are elevating the species' extinction risk, particularly fishing of spawning aggregations.

#### *E. Other Natural or Manmade Factors Affecting Its Continued Existence*

The ERAG considered climate change as a threat to Nassau grouper including global warming, sea level rise, and ocean acidification for Factor E. Although Nassau grouper occur across a range of temperatures, spawning occurs when sea surface temperatures range between 25 °C–26 °C (Colin 1992, Tucker and Woodward 1996). Because Nassau grouper spawn in a narrow window of temperatures, a rise in sea surface temperature outside that range could impact spawning or shift the geographic range of it to overlap with waters within the required temperature parameters. Increased sea surface temperatures have also been linked to coral loss through bleaching and disease. Further, increased global temperatures are also predicted to change parasite-host relationships and may present additional unknown concerns (Harvell *et al.* 2002, Marcogliese 2001). Rising sea surface temperatures are also associated with sea level rise. If sea level changed rapidly, water depth at reef sites may be modified with such rapidity that coral and coral reefs could be affected (Munday *et al.* 2008).

Another potential effect of climate change could be the loss of structural habitat in coral reef ecosystems as ocean acidification is anticipated to affect the integrity of coral reefs (Munday *et al.* 2008). Bioerosion may reduce the 3-dimensional structure of coral reefs (Alvarez-Filip *et al.* 2009), reducing adult habitat for Nassau grouper (Coleman and Koenig 2010, Rogers and Beets 2001). Results of the ERAG scores indicated that climate change was an "unknown risk" to Nassau grouper. We agree with the assessment of the ERAG and believe there is not enough information at this time to determine how climate change is affecting the extinction risk of Nassau grouper now or in the foreseeable future.

The ERAG also considered threats from aquaculture to Nassau grouper under Factor E and determined that

aquaculture was a "very low" risk threat to Nassau grouper. Experiments to determine the success rate of larval Nassau grouper culture (Watanabe *et al.* 1995a, 1995b) and survival of released hatchery-reared juveniles have been conducted and feasibility of restocking reefs has been tested (Roberts *et al.* 1995) in St. Thomas, U.S.V.I. However, the potential of Nassau grouper stock enhancement, as with any other grouper species, has yet to be determined (Roberts *et al.* 1995). Serious concerns about the genetic consequences of introducing Nassau grouper raised in facilities, possible problems of juvenile habitat availability, introduction of maladapted individuals, and the inability of stocked individuals to locate traditional spawning locations, continue to be raised. Given the number of concerns with aquaculture and the fact that some spawning aggregations remain, we believe that it is unlikely that Nassau grouper aquaculture will develop further. Therefore we agree with the ERAG that aquaculture presents a very low extinction risk to Nassau grouper and is not contributing to the species' current status.

Demographic factors of abundance, population growth rate/productivity and diversity were also considered by the ERAG under Factor E. Each ERAG member considered whether the species is likely to be able to maintain a sustainable population size and adequate genetic diversity. They also considered whether the species is at risk due to a loss in the breeding population, which leads to a reduction in survival and production of eggs and offspring. Trends or shifts in demographic or reproductive traits were considered when assessing the ranking of threats by each ERAG member to identify a decline in population growth rate. The ERAG scores indicated that abundance of Nassau grouper was a "moderate risk," growth rate/productivity was an "increasing risk," and that diversity was a "moderate risk." We agree with these rankings and believe they are supported by the declining number and size of spawning aggregations, which affects growth rate/productivity and diversity.

#### NMFS's Conclusions From Threats Evaluation

The most serious threats to Nassau grouper are fishing at spawning aggregations and inadequate law enforcement. These threats, considered under Factors B and D, were rated by the ERAG as "high risk" threats to the species. We agree with the ERAG's assessment that these threats are currently affecting the status of Nassau grouper, putting it at a heightened risk

of extinction. A variety of other threats were identified by the ERAG as also impacting the status of this species. Growth rate/productivity (Factor E), spatial structure/connectivity (Factors A and E), and effectiveness of foreign regulations (Factor D) were identified by the ERAG as “increasing risks.” Artificial selection (Factor B), abundance (Factors B and E), diversity (Factor E), commercial harvest (Factors B and D), and effectiveness of state and territory regulations (Factor D) were determined to be “moderate risks.” NMFS concurs that these threats have the potential to adversely affect the status of Nassau grouper over the foreseeable future.

### Extinction Risk Analysis

We must assess the ERA results and make a determination as to whether the Nassau grouper is currently in danger of extinction, or likely to become so within the foreseeable future. We first evaluated the current status of the Nassau grouper in light of the four demographic factors. Based on our assessment of the ERA in regards to these demographic factors (abundance, growth rate/productivity, spatial structure and connectivity, and diversity) we do not believe the Nassau grouper is currently in danger of extinction. Each of these demographic factors was ranked by the ERAG as a moderate or increasing risk to the species' current status.

We acknowledge that the abundance of Nassau grouper has been dramatically reduced in relation to historical records, but we do not believe abundance is currently so low that the species is at risk of extinction from stochastic events, environmental variation, anthropogenic perturbations, lack of genetic diversity, or compensatory processes. Although the reduced abundance of Nassau grouper has diminished the size and number of spawning aggregations, spawning is still occurring and abundance is increasing in some locations (*e.g.* Cayman Islands and Bermuda) where adequate protections are effectively being implemented. The abundance of Nassau grouper is now patchily distributed throughout the Caribbean with areas of higher abundance correlated with those areas with effective regulations. We believe the abundance of Nassau grouper in these protected areas is large enough to sustain the overall population and limit extinction risk. However, we also believe that further regulations will be necessary in other countries to counteract past population declines and ultimately recover the population of Nassau grouper throughout the Caribbean.

Abundance is closely related with the other three demographic factors. Growth rate/productivity, spatial structure and connectivity, and diversity are all negatively affected by decreased abundance associated with overexploitation. Historical overfishing has led to a decreased average length and earlier age at maturity in exploited populations, which affects the species' ability to maintain the population growth rate above replacement level. Reductions in the number and distribution of spawning aggregations has the potential to affect larval and juvenile dispersal. This can further affect genetic diversity within the population. However, we don't believe that any of these demographic factors have been adversely affected to the point that Nassau grouper is currently in danger of extinction. As described previously, the species continues to occupy its current range, spawning is still occurring in several locations thus continuing to deliver new recruits to the population, and recovery of spawning aggregations has been documented in locations with adequate regulatory mechanisms and enforcement. The size of Nassau grouper is also increasing in areas where protections are in place (*e.g.*, Belize and U.S.V.I.), indicating that current abundance is not adversely affecting growth rate and productivity at these locations.

After considering the current status of Nassau grouper based on the four demographic factors, we next assessed how the identified threats are expected to affect the status of the species, including its demographic factors, over the foreseeable future. The ERAG identified a variety of threats that have the potential to impact Nassau grouper. The ERAG ranked and we agreed that several threats (habitat alteration, disease, aquaculture, and U.S. federal regulations) ranked as “very low” or “low” risk, will have little to no effect on the extinction risk of Nassau grouper within the foreseeable future. Several other threats (commercial harvest, artificial selection, foreign regulations within jurisdictional waters, and regulations of the U.S. and its territories), were ranked as moderate or increasing risks to the status of Nassau grouper. We agree that collectively these threats could cause Nassau grouper to become in danger of extinction within the foreseeable future.

Finally, the ERAG identified three threats that present a “high” risk to the status of Nassau grouper over the foreseeable future. We agree with the ERAG's assessment that fishing of spawning aggregations combined with inadequate law enforcement is currently

adversely affecting the status of Nassau grouper as discussed above, but disagree with the ERAG's ranking of historic harvest as a high risk. These high risk threats will continue to elevate the extinction risk of Nassau grouper over the foreseeable future. Both threats directly affect the current abundance of the species, its ability to maintain population growth rate, the population structure of the species, and its diversity in terms of genetics and overall ecology.

As previously described, the ERAG analyzed inadequate law enforcement as a standalone threat under Factor D, inadequacy of existing regulatory mechanisms, and ranked it as a “high risk” threat. We agree that existing regulations, and enforcement of existing regulations, are inadequate to control the threat posed by fishing on spawning aggregations, and thus this threat under Factor D is contributing to the extinction risk and status of Nassau grouper.

Based on the information in the Biological Report and the results from the ERA, we conclude that ESA Factors B (overutilization for commercial, recreational, scientific, or educational purposes), D (inadequacy of regulatory mechanisms), and E (other natural or manmade factors) are contributing to a threatened status for Nassau grouper. Overutilization in the form of historical harvest has reduced population size and led to the collapse of spawning aggregations in many locations. While some countries have made efforts to curb harvest, fishing at spawning aggregation sites remains a “high risk” threat. Further contributing to the risk of Nassau grouper extinction is the inadequacy of regulatory control and law enforcement, which leads to continued overutilization (low abundance), reduced reproductive output, and reduced recruitment. If growth and sexual recruitment rates cannot balance the loss from these threats, populations will become more vulnerable to extinction over the future (Primack 1993).

### Protective Efforts

Section 4(b)(1)(A) of the ESA requires the Secretary, when making a listing determination for a species, to take into consideration those efforts, if any, being made by any State or foreign nation to protect the species. To evaluate the efficacy of domestic efforts that have not yet implemented or that have been demonstrated to be effective, the Services developed a joint “Policy for Evaluation of Conservation Efforts When Making Listing Decisions” (“PECE”; 68 FR 15100; March 28, 2003).

The PECE is designed to ensure consistent and adequate evaluation on whether domestic conservation efforts that have been recently adopted or implemented, but not yet proven to be successful, will result in recovering the species to the point at which listing is not warranted or contribute to forming the basis for listing a species as threatened rather than endangered. The PECE is expected to facilitate the development of conservation efforts by states and other entities that sufficiently improve a species' status so as to make listing the species as threatened or endangered unnecessary.

The PECE establishes two overarching criteria to use in evaluating efforts identified in conservations plans, conservation agreements, management plans or similar documents: (1) The certainty that the conservation efforts will be implemented; and (2) the certainty that the efforts will be effective. While section 4(b)(1)(A) requires that we evaluate both domestic and foreign conservation efforts, it does not set out particular criteria for doing so. While the particular framework of the PECE policy only directly applies to consideration of domestic efforts, we have discretion to evaluate foreign efforts using a similar approach and find that it is reasonable to do so here. In our discretion, we evaluated foreign conservation efforts to protect and recover Nassau grouper that are either underway, but not yet fully implemented, or are only planned, using these overarching criteria.

Conservation efforts with the potential to address identified threats to Nassau grouper include, but are not limited to, fisheries management plans, education about overfishing and fishing of spawning aggregations, and projects addressing the health of coral reef ecosystems. These conservation efforts may be conducted by countries, states, local governments, individuals, NGOs, academic institutions, private companies, individuals, or other entities. They also include global conservation organizations that conduct coral reef and/or marine environment conservation projects, global coral reef monitoring networks and research projects, regional or global conventions, and education and outreach projects throughout the range of Nassau grouper.

The Biological Report summarizes known conservation efforts, including those that have yet to be fully implemented or have yet to demonstrate effectiveness. Conservation efforts that we considered that are yet to be fully implemented include Mexico's 2012 proposed management plan, Antigua-Barbuda's 2008 closed season proposal,

and Guadeloupe and Martinique's plans to protect the species. Because these proposed plans are several years old with no updates or known implementation, we find that there is not a sufficient basis to conclude that there is a reasonable certainty of implementation or effectiveness. We also considered the marine protected areas implemented by Jamaica in 2011, though based on Jamaica's historic overfishing and difficulty in enforcing existing regulations, we find that there is not a sufficient basis to conclude that these marine protected areas present a reasonable certainty of effectiveness in reducing threats that contribute to Nassau grouper's extinction risk. We carefully considered the other conservation efforts summarized in the Biological Report and acknowledge that time is required to see the benefit of mature adults in the spawning aggregations; however, the continued decline in number and size of Nassau grouper spawning aggregations indicates the effectiveness of those conservation efforts is currently unknown and thus there is insufficient basis to conclude there is a reasonable certainty of effectiveness. While some conservation efforts have been partially successful on localized scales, Nassau grouper appear to still be overutilized and at heightened risk of extinction based on the ERA. After taking into account these conservation efforts, our evaluation of the section 4(a)(1) factors is that the conservation efforts do not reduce the risk of extinction of Nassau grouper to the point at which listing is not warranted.

#### *Significant Portion of Range*

There are two situations under which a species is eligible for listing under ESA: A species may be endangered or threatened throughout all of its range or a species may be endangered or threatened throughout only a "significant portion of its range" (SPOIR). Although the ESA does not define "SPOIR," NMFS and the U.S. Fish and Wildlife Service (USFWS) published a final policy clarifying their interpretation of this phrase (79 FR 37577; July 7, 2014). Under the policy, if a species is found to be endangered or threatened throughout only a significant portion of its range, the entire species is subject to listing and must be protected everywhere. A portion of a species' range is "significant" if ". . . the species is not currently endangered or threatened throughout its range, but the portion's contribution to the viability of the species is so important that, without the members in that portion, the species

would be in danger of extinction, or likely to become so in the foreseeable future, throughout all of its range." Thus, if the species is found to be threatened or endangered throughout its range, we do not separately evaluate portions of the species' range.

Although the SPOIR Policy had yet to go into effect during our status review of Nassau grouper, we considered the interpretations and principles contained in the 2014 Draft Policy with regards to the Nassau grouper and completed an assessment of potential "SPOIR," which is documented in the ERA. However, throughout the status review process NMFS determined threats and risks to the status of Nassau grouper are affecting the species over the entirety of its range. Because the threats and risks are widespread throughout the entire range of this species, there is no portion of the range that can be considered "significant."

#### **Listing Determination**

Based on the Biological Report, the Threats Evaluation, the Extinction Risk Analysis, and Protective Efforts we determined that the Nassau grouper warrants a threatened status under the ESA. We summarize the results of our comprehensive status review as follows: (1) The species is made up of a single population over a broad geographic range, and its current range is indistinguishable from its historical range; (2) the species possesses life history characteristics that increase vulnerability to unregulated harvest; (3) historical harvest greatly diminished the population of Nassau grouper and the species has yet to recover from this overexploitation; (4) spawning aggregations have drastically declined in size and number across the species' range; (5) there are two threats the ERAG rated as "high risk," that we agree are affecting the current status of the species and will continue to do so over the foreseeable future—fishing at spawning aggregations and inadequate law enforcement; and (6) historical harvest has abated, though existing regulatory mechanisms and law enforcement have not been effective in preventing fishing at many spawning aggregation sites. Conservation efforts in some nations (U.S., Puerto Rico, U.S.V.I., and Belize) have almost certainly prevented further declines. Given the life history characteristics of Nassau grouper, more time will be needed to determine if these protective measures are successful in recovering the population. Collectively, the information obtained during the status review indicates the species is not currently in danger of extinction

(though reduced in number, the species maintains its historical range and still forms spawning aggregations at some sites), but it is likely to become endangered within the foreseeable future (based on continued risk of harvest, especially at spawning aggregation sites inadequately controlled by regulations and law enforcement). Accordingly, we have determined that the Nassau grouper warrants listing as a threatened species under the ESA.

### Effects of Listing

Conservation measures provided for species listed as endangered or threatened under the ESA include recovery plans (16 U.S.C. 1533(f)), critical habitat designations (16 U.S.C. 1533(a)(3)(A)), Federal agency consultation requirements (16 U.S.C. 1536), and protective regulations (16 U.S.C. 1533(d)). Recognition of the species' status through listing promotes conservation actions by Federal and state agencies, private groups, and individuals, as well as the international community. Both a recovery program and designation of critical habitat could result from this final listing. Given its broad range across the Caribbean Sea, a regional cooperative effort to protect and restore Nassau grouper is necessary. We anticipate that protective regulations for Nassau grouper will also be necessary for the conservation of the species. Federal, state, and the private sectors will need to cooperate to conserve listed Nassau grouper and the ecosystems upon which they depend.

### Identifying ESA Section 7 Consultation Requirements

Section 7(a)(2) of the ESA and NMFS/FWS regulations require Federal agencies to consult with us on any actions they authorize, fund, or carry out if those actions may affect the listed species or designated critical habitat. Based on currently available information, we can conclude that examples of Federal actions that may affect Nassau grouper include, but are not limited to, artificial reef creation, dredging, pile-driving, military activities, and fisheries management practices.

### Critical Habitat

Critical habitat is defined in section 3 of the ESA (16 U.S.C. 1532(5)) as: (1) The specific areas within the geographical area occupied by a species, at the time it is listed in accordance with the ESA, on which are found those physical or biological features (a) essential to the conservation of the species and (b) that may require special

management considerations or protection; and (2) specific areas outside the geographical area occupied by a species at the time it is listed upon a determination that such areas are essential for the conservation of the species. "Conservation" means the use of all methods and procedures needed to bring the species to the point at which listing under the ESA is no longer necessary. Critical habitat may also include areas unoccupied by Nassau grouper if those areas are essential to the conservation of the species.

Section 4(a)(3)(A) of the ESA (16 U.S.C. 1533(a)(3)(A)) requires that, to the maximum extent prudent and determinable, critical habitat be designated concurrently with the listing of a species. Pursuant to 50 CFR 424.12(a), designation of critical habitat is not determinable when one or both of the following situations exist: Data sufficient to perform required analyses are lacking; or the biological needs of the species are not sufficiently well known to identify any area that meets the definition of "critical habitat." Although we have gathered information through the status review and public comment periods on the habitats occupied by this species, we currently do not have enough information to determine what physical and biological features within those habitats facilitate the species' life history strategy and are thus essential to the conservation of Nassau grouper, and may require special management considerations or protection. To the maximum extent prudent and determinable, we will publish a proposed designation of critical habitat for Nassau grouper in a separate rule. Designations of critical habitat must be based on the best scientific data available and must take into consideration the economic, national security, and other relevant impacts of specifying any particular area as critical habitat. Once critical habitat is designated, section 7 of the ESA requires Federal agencies to ensure that they do not fund, authorize, or carry out any actions that are likely to destroy or adversely modify that habitat. This requirement is in addition to the section 7 requirement that Federal agencies ensure that their actions do not jeopardize the continued existence of listed species.

### Identification of Those Activities That Would Constitute a Violation of Section 9 of the ESA

Because we are proposing to list Nassau grouper as threatened, the ESA section 9 prohibitions do not automatically apply. Therefore,

pursuant to ESA section 4(d), we will evaluate whether there are protective regulations we deem necessary and advisable for the conservation of Nassau grouper, including application of some or all of the take prohibitions. If protective regulations are deemed necessary, a proposed 4(d) rule would be subject to public comment.

### Policies on Peer Review

In December 2004, the Office of Management and Budget (OMB) issued a Final Information Quality Bulletin for Peer Review establishing minimum peer review standards, a transparent process for public disclosure of peer review planning, and opportunities for public participation. The OMB Bulletin, implemented under the Information Quality Act (Pub. L. 106-554) is intended to enhance the quality and credibility of the Federal government's scientific information, and applies to influential or highly influential scientific information disseminated on or after June 16, 2005. To satisfy our requirements under the OMB Bulletin, we obtained independent peer review of the Biological Report. Five independent specialists were selected from the academic and scientific community, Federal and state agencies, and the private sector for this review (with three respondents). All peer reviewer comments were addressed prior to dissemination of the final Biological Report and publication of this final rule.

### Solicitation of Information

We are soliciting information on features and areas that may support designation of critical habitat for Nassau grouper. Information provided should identify the physical and biological features essential to the conservation of the species and areas that contain these features. Areas outside the occupied geographical area should also be identified if such areas themselves are essential to the conservation of the species. Essential features may include, but are not limited to, features specific to the species' range, habitats, and life history characteristics within the following general categories of habitat features: (1) Space for individual growth and for normal behavior; (2) food, water, air, light, minerals, or other nutritional or physiological requirements; (3) cover or shelter; (4) sites for reproduction and development of offspring; and (5) habitats that are protected from disturbance or are representative of the historical, geographical, and ecological distributions of the species (50 CFR 424.12(b)). ESA implementing regulations at 50 CFR 424.12(h) specify that critical habitat shall not be

designated within foreign countries or in other areas outside of U.S. jurisdiction. Therefore, we request information only on potential areas of critical habitat within waters in U.S. jurisdiction.

For features and areas potentially qualifying as critical habitat, we also request information describing: (1) Activities or other threats to the essential features or activities that could be affected by designating them as critical habitat, and (2) the positive and negative economic, national security and other relevant impacts, including benefits to the recovery of the species, likely to result if these areas are designated as critical habitat.

**References**

A complete list of the references used in this final rule is available at: ([http://sero.nmfs.noaa.gov/protected\\_resources/listing\\_petitions/species\\_esa\\_consideration/index.html](http://sero.nmfs.noaa.gov/protected_resources/listing_petitions/species_esa_consideration/index.html)).

**Classifications**

*National Environmental Policy Act*

The 1982 amendments to the ESA, in section 4(b)(1)(A), restrict the information that may be considered when assessing species for listing. Based on this limitation of criteria for a listing decision and the opinion in *Pacific Legal Foundation v. Andrus*, 675 F. 2d 825 (6th Cir. 1981), NMFS has concluded that ESA listing actions are not subject to the environmental assessment requirements of the National

Environmental Policy Act (See NOAA Administrative Order 216–6).

*Executive Order 12866, Regulatory Flexibility Act and Paperwork Reduction Act*

As noted in the Conference Report on the 1982 amendments to the ESA, economic impacts cannot be considered when assessing the status of a species. Therefore, the economic analysis requirements of the Regulatory Flexibility Act are not applicable to the listing process. In addition, this final rule is exempt from review under Executive Order 12866. This final rule does not contain a collection-of-information requirement for the purposes of the Paperwork Reduction Act.

*Executive Order 13132, Federalism*

In keeping with the intent of the Administration and Congress to provide continuing and meaningful dialogue on issues of mutual state and Federal interest, the proposed rule was provided to the relevant agencies in each state in which the subject species occurs, and these agencies were invited to comment. We did not receive comments from any state agencies.

*Executive Order 12898, Environmental Justice*

Executive Order 12898 requires that Federal actions address environmental justice in the decision-making process. In particular, the environmental effects of the actions should not have a

disproportionate effect on minority and low-income communities. This final rule is not expected to have a disproportionately high effect on minority populations or low-income populations.

**List of Subjects in 50 CFR Part 223**

Endangered and threatened species, Exports, Transportation.

Dated: June 21, 2016.

**Samuel D Rauch, III,**  
*Deputy Assistant Administrator for Regulatory Programs, National Marine Fisheries Service.*

For the reasons set out in the preamble, we amend 50 CFR part 223 as follows:

**PART 223—THREATENED MARINE AND ANADROMOUS SPECIES**

■ 1. The authority citation for part 223 continues to read as follows:

**Authority:** 16 U.S.C. 1531–1543; subpart B, § 223.201–202 also issued under 16 U.S.C. 1361 *et seq.*; 16 U.S.C. 5503(d) for § 223.206(d)(9).

■ 2. In § 223.102, amend the table in paragraph (e) by adding an entry under the “Fishes” subheading for “Grouper, Nassau” in alphabetical order to read as follows:

**§ 223.102 Enumeration of threatened marine and anadromous species.**

\* \* \* \* \*  
(e) \* \* \*

Species <sup>1</sup>		Description of listed entity	Citation(s) for listing determination(s)	Critical habitat	ESA rules
Common name	Scientific name				
*	*	*	*	*	*
FISHES					
*	*	*	*	*	*
Grouper, Nassau .....	<i>Epinephelus striatus</i> ..	Entire species .....	[Insert <b>Federal Register</b> citation], June 29, 2016.	NA	NA
*	*	*	*	*	*

<sup>1</sup>Species includes taxonomic species, subspecies, distinct population segments (DPSs) (for a policy statement, see 61 FR 4722, February 7, 1996), and evolutionarily significant units (ESUs) (for a policy statement, see 56 FR 58612, November 20, 1991).

\* \* \* \* \*

[FR Doc. 2016-15101 Filed 6-28-16; 8:45 am]

BILLING CODE 3510-22-P

**DEPARTMENT OF COMMERCE****National Oceanic and Atmospheric Administration****50 CFR Part 600**

[Docket No. 111014628-6513-02]

RIN 0648-BB54

**Magnuson-Stevens Fishery Conservation and Management Act Provisions; Implementation of the Shark Conservation Act of 2010**

**AGENCY:** National Marine Fisheries Service (NMFS), National Oceanic and Atmospheric Administration (NOAA), Commerce.

**ACTION:** Final rule.

**SUMMARY:** This final action updates agency regulations consistent with provisions of the Shark Conservation Act of 2010 (SCA) and prohibits any person from removing any of the fins of a shark at sea, possessing shark fins on board a fishing vessel unless they are naturally attached to the corresponding carcass, transferring or receiving fins from one vessel to another at sea unless the fins are naturally attached to the corresponding carcass, landing shark fins unless they are naturally attached to the corresponding carcass, or landing shark carcasses without their fins naturally attached. This action amends existing regulations and makes them consistent with the SCA.

**DATES:** Effective July 29, 2016.

**ADDRESSES:** Copies of the Environmental Assessment (EA)/Regulatory Impact Review (RIR)/Final Regulatory Flexibility Analysis (FRFA) prepared for this action can be obtained from: Erin Wilkinson, National Marine Fisheries Service, 1315 East-West Highway, Room 13437, Silver Spring MD 20910. An electronic copy of the EA/RIR/FRFA document as well as copies of public comments received can be viewed at the Federal e-rulemaking portal: <http://www.regulations.gov/> (Docket ID: NOAA-NMFS-2012-0092).

**FOR FURTHER INFORMATION CONTACT:** Erin Wilkinson by phone at 301-427-8561, or by email: [erin.wilkinson@noaa.gov](mailto:erin.wilkinson@noaa.gov) or [sca.rulemaking@noaa.gov](mailto:sca.rulemaking@noaa.gov).

**SUPPLEMENTARY INFORMATION:****I. Overview of the Shark Conservation Act**

Background information and an overview of the Shark Conservation Act

can be found in the preamble of the proposed rule published on May 2, 2013 (78 FR 25685). Copies are available from NMFS (see **ADDRESSES**), or can be viewed electronically at the Federal E-Rulemaking portal for this action: <http://www.regulations.gov>.

**II. Major Components of the Final Action**

Retaining a shark fin while discarding the shark carcass (shark finning) has been prohibited in the United States since the 2000 Shark Finning Prohibition Act. The 2010 SCA included provisions that amended the Magnuson-Stevens Fishery Conservation and Management Act (MSA) to prohibit any person from: (1) Removing any of the fins of a shark (including the tail) at sea; (2) having custody, control, or possession of a fin aboard a fishing vessel unless it is naturally attached to the corresponding carcass; (3) transferring a fin from one vessel to another vessel at sea, or receiving a fin in such transfer, unless the fin is naturally attached to the corresponding carcass; or (4) landing a fin that is not naturally attached to the corresponding carcass, or landing a shark carcass without its fins naturally attached. For the purpose of the SCA and these regulations, “naturally attached,” with respect to a shark fin, means to be attached to the corresponding shark carcass through some portion of uncut skin.

This action amends NMFS’ regulations consistent with these provisions of the SCA. Specifically, the rule amends regulations at 50 CFR part 600, subpart N, to prohibit the removal of shark fins at sea, namely, the possession, transfer and landing of shark fins that are not naturally attached to the corresponding carcass, and the landing of shark carcasses without the corresponding fins naturally attached. In the preamble to the proposed rule, NMFS noted that it interprets the prohibitions in subpart N as applying to sharks, not skates and rays, and solicited public comment on whether clarification was needed in the regulatory text on this issue. See 78 FR 25685, 25686 (May 2, 2013). NMFS received only one public comment on this point, which was supportive of this interpretation, and NMFS thus affirms in this final rule that the prohibitions do not apply to skates and rays.

This final rule also updates subpart N to be consistent with section 103(b) of the SCA regarding an exception for individuals engaged in commercial fishing for smooth dogfish. Interpretation of that exception was addressed in a rule finalized in

November 2015, for Amendment 9 to the 2006 Consolidated Atlantic Highly Migratory Species Fishery Management Plan (November 24, 2015; 80 FR 73128). That final rule, among other things, allows for the at-sea removal of smooth dogfish fins provided that fishing occurs within 50 nautical miles of shore along the Atlantic Coast from Maine through the east coast of Florida; smooth dogfish fin weight does not exceed 12 percent of the carcass weight on board; smooth dogfish make up at least 25 percent of the total retained catch, by weight; and the fisherman/vessel holds both federal and state permits appropriate for the retention of smooth dogfish.

This final rule also combines the existing §§ 600.1203 and 600.1204 into one section. The text throughout 50 CFR part 600, subpart N, is amended to make it consistent with the provisions of the SCA.

The MSA authorizes the Secretary to regulate fisheries seaward of the inner boundary of the U.S. exclusive economic zone (EEZ), which is defined as a line coterminous with the seaward boundary of each U.S. coastal state. 16 U.S.C. 1802(11). Thus, as noted in the proposed rule, the SCA provisions apply to any person subject to the jurisdiction of the United States, including persons on board U.S. and foreign vessels, engaging in activities prohibited under the statute with respect to sharks harvested seaward of the inner boundary of the EEZ. See 78 FR 25685, 25686 (May 2, 2013). Federal regulations pertaining to the conservation and management of specific shark fisheries are set forth in parts 635, 648, and 660 of title 50 of the Code of Federal Regulations. For Atlantic highly migratory species fisheries, as a condition of its Federal permit, a vessel’s fishing, catch, and gear are subject to federal requirements even when fishing in state waters. See 50 CFR 635.4(a)(10) (noting also that, when fishing within the waters of a state with more restrictive regulations, persons aboard the vessel must comply with those requirements). This rule amends 50 CFR part 600, subpart N, and does not supersede or amend any other federal regulation or requirement related to the conservation and management of sharks.

The SCA also amended the High Seas Driftnet Fishing Moratorium Protection Act, which provides for identification and certification of nations to address illegal, unreported, or unregulated fishing; bycatch of protected living marine resources; and, as amended by the SCA, shark catches. 16 U.S.C. 1826h-1826k. With regard to sharks, the High Seas Driftnet Fishing Moratorium

Spawning Aggregations Workshop Report  
FIRST MEETING OF THE CFMC/WECAFC/OSPESCA/CRFM WORKING  
GROUP ON SPAWNING AGGREGATIONS  
Miami, United States of America, 29–31 October 2013


## WESTERN CENTRAL ATLANTIC FISHERY COMMISSION

Report of the

---

### FIRST MEETING OF THE CFMC/WECAFC/OSPESCA/CRFM WORKING GROUP ON SPAWNING AGGREGATIONS

Miami, United States of America, 29–31 October 2013



Copies of FAO publications can be requested from:

Sales and Marketing Group  
Food and Agriculture Organization  
of the United Nations

E-mail: [publications-sales@fao.org](mailto:publications-sales@fao.org)

Fax: +39 06 57053360

Web site: [www.fao.org/icatalog/inter-e.htm](http://www.fao.org/icatalog/inter-e.htm)

WESTERN CENTRAL ATLANTIC FISHERY COMMISSION

Report of the

FIRST MEETING OF THE CFMC/WECAFC/OSPESCA/CRFM WORKING GROUP  
ON SPAWNING AGGREGATIONS

Miami, United States of America, 29–31 October 2013

FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS  
Sub-regional Office for the Caribbean  
Bridgetown, Barbados, 2014

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

ISBN 978-92-5-108139-6 (print)  
E-ISBN 978-92-5-108140-2 (PDF)

© FAO, 2014

FAO encourages the use, reproduction and dissemination of material in this information product. Except where otherwise indicated, material may be copied, downloaded and printed for private study, research and teaching purposes, or for use in non-commercial products or services, provided that appropriate acknowledgement of FAO as the source and copyright holder is given and that FAO's endorsement of users' views, products or services is not implied in any way.

All requests for translation and adaptation rights, and for resale and other commercial use rights should be made via [www.fao.org/contact-us/licence-request](http://www.fao.org/contact-us/licence-request) or addressed to [copyright@fao.org](mailto:copyright@fao.org).

FAO information products are available on the FAO website ([www.fao.org/publications](http://www.fao.org/publications)) and can be purchased through [publications-sales@fao.org](mailto:publications-sales@fao.org).

## **PREPARATION OF THIS DOCUMENT**

This is the report of the first meeting of the Caribbean Fisheries Management Council (CFMC), the Western Central Atlantic Fishery Commission (WECAFC), the Central American Organization of the Fisheries and Aquaculture Sector (OSPESCA), and the Caribbean Regional Fishery Mechanism (CRFM) Working Group on Spawning Aggregations, held in Miami, from 29 to 31 October 2013.

The joint Working Group was established by the fourteenth session of WECAFC in February 2012 and this first meeting was co-organized and sponsored by the CFMC of the United States Department of Commerce, WECAFC and FAO.

The FAO Secretariat to the meeting consisted of Dr Raymon van Anrooy, WECAFC Secretary. Administrative and logistical support was provided by CFMC, and coordinated by Mr Miguel Rolon, Executive Director of CFMC and convener of this Working Group, with assistance from Ms Diana Martino and Ms Maria de los Angeles Irizarry. Dr Yvonne Sadovy of the University of Hong Kong, technically coordinated and facilitated the meeting.

This report contains a summary of the presentations, discussions, conclusions and recommendations of the meeting. The conclusions adopted and recommendations made are presented in the form of a “Declaration of Miami” and a Recommendation to the fifteenth session of WECAFC on the establishment of a regional closed season for fisheries in the WECAFC area to protect spawning aggregations of groupers and snappers. The national summary reports presented at the meeting will be published separately with support from CFMC.

**FAO Western Central Atlantic Fishery Commission. 2014.**

*Report of the first meeting of the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations, Miami, United States of America, 29–31 October 2013.*

FAO Fisheries and Aquaculture Report. No. 1059. Bridgetown, Barbados, FAO. 29 pp.

#### **ABSTRACT**

The first meeting of the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations, was held in Miami, United States of America from 29 to 31 October 2013. The meeting brought together 23 experts working on spawning aggregations of fishes from all over the Western Central Atlantic region. The Working Group noted with concern the ongoing declines in stocks of many aggregating species and particularly groupers and snappers in the Wider Caribbean Region, the reduced numbers of their aggregations and the relatively smaller size of remaining aggregations. The Working Group also verified that the status of Nassau Grouper, Goliath Grouper (and several other species) stocks in the Wider Caribbean Region should be considered “overexploited”, and that some stocks can even be regarded as “depleted”. The Working Group further emphasized the high ecological and biological value of reef fishes that aggregate to spawn (including groupers and snappers) for the ecosystem and aquatic biodiversity in the region, as well as for achieving regional food security and livelihood objectives. The Working Group compiled information on the spawning fish aggregation management and conservation measures in place and examined their effectiveness. The meeting issued a “Declaration of Miami”, which included a recommendation to the fifteenth session of WECAFC on the establishment of a regional closed season for Nassau Grouper fisheries in the WECAFC area to protect spawning aggregations of this species.

## CONTENTS

	<b>Page</b>
INTRODUCTION	1
OPENING OF THE MEETING	1
ATTENDANCE	2
ELECTION OF CHAIRPERSONS AND RAPPORTEURS	2
ADOPTION OF THE AGENDA	2
GLOBAL PERSPECTIVE OF AGGREGATING SNAPPERS AND GROUPERS	2
HISTORICAL BACKGROUND OF WECAFC'S WORK ON SPAWNING AGGREGATIONS OF KEY SPECIES	3
PRESENTATIONS OF NATIONAL STATUS REPORTS	3
WORKING GROUP DISCUSSIONS	13
GENERAL DISCUSSIONS	17
PREPARATION AND ADOPTION OF THE REPORT TO WECAFC	18
CLOSURE OF THE MEETING	18
APPENDIX A     Agenda	19
APPENDIX B     List of participants	20
APPENDIX C     Declaration of Miami	23
APPENDIX D     Draft Recommendation to the fifteenth session of WECAFC	26
APPENDIX E     Terms of Reference of the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations	29

## ABBREVIATIONS AND ACRONYMS

CFMC	Caribbean Fishery Management Council
CITES	Convention on International Trade in Endangered Species of Wild Fauna and Flora
CRFM	Caribbean Regional Fisheries Mechanism
EEZ	exclusive economic zone
ESA	Endangered Species Act (USA)
FSA	fish spawning aggregation
GCFI	Gulf and Caribbean Fisheries Institute
INPESCA	Institute for Fisheries and Aquaculture (Nicaragua)
IUCN	International Union for Conservation of Nature
MPAs	marine protected areas
MSC	Marine Stewardship Council
NGO	non-governmental organization
NMFS	National Marine Fisheries Service
NOAA	National Oceanic and Atmospheric Administration (USA)
OSPESCA	Central American Organization of the Fisheries and Aquaculture Sector
SAG	Scientific Advisory Group
SCRFA	Society for the Conservation of Reef Fish Aggregations
SIDS	Small Island Developing States
SPAW Protocol	Protocol Concerning Specially Protected Areas and Wildlife
STAC	Scientific Meeting and Technical Advisory Committee
Code	Code of Conduct for Responsible Fisheries (FAO)
USVI	United States Virgin Islands
WECAFC	Western Central Atlantic Fishery Commission

## **INTRODUCTION**

1. Spawning Aggregations of reef fishes, particularly groupers, have been the focus of various regional meetings in the Caribbean. A Regional Workshop on Nassau grouper, which was coordinated by the Caribbean Fishery Management Council (CFMC) and the Western Central Atlantic Fishery Commission (WECAFC), was held on 20 and 21 October 2008, prior to the thirteenth session of WECAFC. The National Oceanographic and Atmospheric Administration (NOAA) and the National Marine Fisheries Service (NMFS) of the United States of America sponsored that workshop. Representatives from 17 countries attended the workshop. The two main items on the agenda were: a regional summary of the status of the Nassau grouper fishery in the region and the compilation of country status reports. The Regional Workshop made various recommendations which were presented to WECAFC. These recommendations included:

- a) a proposal for establishment of a WECAFC/CFMC ad hoc Working Group on Nassau Grouper;
- b) that management of Nassau Grouper be more effective at the national level;
- c) closed seasons are one of the most effective ways to protect spawning aggregations, when the species is more vulnerable to fishing; and
- d) countries that do not have a closed season from December to February should establish one.

2. The thirteenth session of WECAFC (Colombia, October 2008) endorsed the recommendations and added that the main purpose of the Working Group is to foster regional cooperation in the management and conservation and restoration of Nassau grouper stocks in the WECAFC region; and to include coordination and harmonization of efforts for the management and conservation of the Nassau grouper. The thirteenth session recommended a regional coherent management approach, supported by national level implementation efforts.

3. Various Gulf and Caribbean Fisheries Institute (GCFI) annual conferences in recent years incorporated sessions or presentations on Spawning Aggregations or Nassau Grouper management and conservation. Moreover, the Society for Conservation of Reef Fish Aggregations (SCRFA) (later revised to Science and Conservation of Fish Aggregations) has been very active in raising awareness and building capacity on aggregations over the last decade. Numerous scientists, researchers, fishers and projects have been working on spawning aggregations and related issues lately.

4. At the fourteenth session of WECAFC, held in Panama City in February 2012, the Commission noted the limited activities of the Working Group on Nassau Grouper and Mr Miguel Rolon (CFMC) kindly offered to revive the working group as CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations. The respective terms of reference were developed and endorsed by WECAFC (available in Appendix E) and funding was sought in support of Working Group activities. Mr Rolon called, as convener, the Working Group together to meet in Miami, United States of America. Moreover, Dr Sadovy compiled a status report on Nassau grouper which was delivered to the CFMC following the thirteenth session of WECAFC.

5. The principal objective of this first Working Group meeting was to bring together key experts to examine the available biological and socio-economic information from Caribbean countries involved in the fisheries of groupers and snappers and other species that aggregate to spawn. It was aimed to use the information to provide (as Working Group) advice on the management and implementation of regional strategies and regulations to protect spawning aggregations.

## **OPENING OF THE MEETING**

6. The first meeting of the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations was held in Miami from 29 to 31 October 2013. The meeting was kindly hosted by the

Caribbean Fisheries Management Council (CFMC). Welcoming remarks were delivered by Mr Miguel Rolon on behalf of CFMC and as convener of the Working Group, and by Dr Raymon van Anrooy on behalf of FAO/WECAFC.

## **ATTENDANCE**

7. The following countries and territories attended the meeting: Bahamas, Belize, Brazil, Caribbean Netherlands, Cayman Islands, Cuba, China (Hong Kong SAR), Mexico, Nicaragua, Puerto Rico, United States Virgin Islands and the United States of America. CMFC, CRFM, PEW and WECAFC/FAO, as well as various Spawning Aggregations experts were also in attendance. The list of 23 participants, including Working Group members and other participants, can be found in Appendix B.

## **ELECTION OF CHAIRPERSONS AND RAPPORTEURS**

8. Dr Yvonne Sadovy was elected Chairperson of the Meeting. She was assisted by Dr Raymon Van Anrooy who also agreed to act as Rapporteur.

## **ADOPTION OF THE AGENDA**

9. The Meeting adopted the agenda as shown in Appendix A.

## **GLOBAL PERSPECTIVE OF AGGREGATING SNAPPERS AND GROUPERS**

10. Dr Yvonne Sadovy presented a “Global Perspective of Aggregating Snappers and Groupers”. The presentation covered what is known of the fish taxa that aggregate to spawn (mainly groupers and snappers according to the SCRFA database), the main habitats that spawning occurs globally, multispecies spawning, timing, relative to lunar phase, of spawning, and spawning behaviour. The talk covered the high importance of aggregating species within reef fisheries and hence their significance for food security and for earnings, with particular focus on groupers and snappers used both domestically and for exports. It was noted that export trade can drive particularly heavy focus on aggregating and migrating fish to fulfill the need to complete large shipments and catch large numbers of fish quickly (the cases of Honduras and Fiji were presented) which can contribute to overfishing of the species. Aggregating species are typically fished both during the aggregation season as well as outside of it and the case was presented to protect aggregations, the source of the next generation, and only fish outside of aggregations to ensure continuation of the fishery (of aggregating species) in the long term.

11. The status of aggregations globally was presented with most of known status found to be declining and little effective management in place. The case of the Nassau grouper throughout its geographic range was presented in detail for the lessons learned, as was the role of overfishing of aggregations in producing the threatened and near-threatened listings (International Union for Conservation of Nature [IUCN] Red List criteria) of several grouper species. Finally, the challenges and opportunities of management of aggregating species were presented highlighting particularly the issues of the value of aggregating species, the challenges of assessing their status (hyper-stability) from aggregation catches and the illusion of plenty that large number of fish gathered at one time and place can give. Management options were presented and statements of concern from various forums summarized.

12. The presentation triggered a lot of discussion on a wide range of issues. The importance of getting disaggregated trade data on grouper imports and exports, the potential impact of climate change and variability on reef fish species that aggregate to spawn, hyperstability issues of stocks of fishes that aggregate to spawn, the need for local fisherfolk participation in development and implementation of spawning aggregation conservation measures and the availability of and access to fisheries manuals and other awareness raising materials on spawning aggregations, were among the issues discussed. It was argued that there is an imbalance in terms of regulatory and management

measures in place for aquatic species, as lobster, queen conch and turtle are often covered by these measures, but groupers and other fishes that aggregate to spawn frequently (or even typically) are not. The Working Group agreed that the SCRFA website ([www.scrfa.org](http://www.scrfa.org)), with its database and visibility and training materials could be used as repository for researchers/experts in the region who would like to share information on spawning aggregations.

## **HISTORICAL BACKGROUND OF WECAFC'S WORK ON SPAWNING AGGREGATIONS OF KEY SPECIES**

13. Dr Van Anrooy made a presentation which covered the history, objectives and core activities of WECAFC, membership issues, and the work of the seven current working groups. Most information presented, as well as reports and publications of WECAFC working groups are available at the WECAFC website in three languages, accessible at: [www.fao.org/fishery/rfb/wecafc/en](http://www.fao.org/fishery/rfb/wecafc/en). Dr Van Anrooy provided further some historical background to the Working Group on Spawning Aggregations and detailed (on behalf of the convener) the Terms of Reference of the Working Group. He also presented the data and information that FAO has on grouper landings in the WECAFC mandate area. In summary, the total landings of groupers in the WECAFC region were estimated in 2010 to be 16 400 tonnes and in 2011 some 14 400 tonnes. This is equivalent to 1.3 percent and 1 percent of total capture fisheries production in the region in these years. Mexico (51 percent), USA (28 percent), Venezuela (9 percent) and the Dominican Republic (5 percent) are the largest producers in terms of volumes of grouper harvested. The FAO data showed that since the 1980s there is a clear downward trend in landings of groupers. It was noted that the USA is the largest importer of grouper and grouper products from the region.

14. The discussion succeeding the presentation related to the role of non-coastal member states of WECAFC and how they could be incorporated better in regional fisheries management and to a perceived need to be able to make binding fisheries management recommendations in the region.

## **PRESENTATIONS OF NATIONAL STATUS REPORTS**

15. National Summary Reports were prepared by expert participants from most countries attending the Working Group meeting. These summary reports and other research outcomes provided are made available in full in a separate report, along with an updated regional status overview. Also the representative of the CRFM provided an overview of the work of the Mechanism on aggregating species.

16. The presentations of the overviews were received with interest by the Working Group.

17. Summarized below are the presentations made by the experts and issues raised by participants during the discussions following the presentations.

18. **The United States Virgin Islands (USVI).** Dr Richard S. Nemeth of the Center for Marine and Environmental Studies of the University of the Virgin Islands presented the status of spawning aggregations in the USVI.

19. In the United States Virgin Islands at least 20 species from five families (*Lutjanidae*, *Epinephelidae*, *Carangidae*, *Balistidae*, *Kyphosidae*) are known or suspected to form transient fish spawning aggregations (FSA). FSA's are important life history events characterized by very predictable locations and timing where the spawning adults are the primary source of annual reproductive effort. These characteristics make spawning aggregations very vulnerable to fishing which may severely deplete local populations: a scenario that has occurred repeatedly in the USVI and elsewhere in the Caribbean, especially the collapse of Nassau grouper (*Epinephalus striatus*) spawning aggregations.

20. Understanding the status of spawning aggregations is critical to their management. In the USVI, nearly all of the species that form transient spawning aggregations are either declining or have

insufficient information to evaluate their status, even though management regulations have been in place for five to ten years. These regulations include three US federal marine protected areas, three federal and local seasonal area closures and three areas with limited protection. Additional regulations include no-take for Nassau and Goliath grouper (*E. itajara*) and three endangered parrotfish (*Scarus guacamaia*, *S. coelestinus*, *S. coeruleus*) and seasonal catch restrictions on groupers (February to April) and snappers (April to June). In only one case has a species (red hind *E. guttatus* on St. Thomas) shown recovery due to protection of its spawning aggregation site. This is in stark contrast to the St. Croix red hind spawning population which has shown continuous decline for the past ten years in terms of size of males and females, sex ratios, population abundance and biomass even though it has received similar protection. A lack of basic biological information is hindering our understanding of these differences in response to management actions.

21. A minimum level of research is needed to provide Caribbean countries a baseline on which to establish FSA monitoring protocols (i.e. port surveys, underwater fish counts, bathymetric and habitat mapping). This basic information as well as more sophisticated studies can provide guidance for implementing precautionary management regulations. For example, a study in the USVI using acoustic telemetry to track grouper movements found that area requirements around spawning sites showed a strong positive relationship based on fish size. The largest species (yellowfin grouper, *Mycteroperca venenosa*) required 10–12 km<sup>2</sup>, Nassau grouper required 5–6 km<sup>2</sup> and tiger grouper (*M. tigris*) 3–4 km<sup>2</sup>. This information is broadly applicable to other countries and can be used to guide managers to define spatial and temporal closed areas and justify boundaries to stakeholders through a variety of outreach and informal education efforts.

22. The discussion which followed Dr Nemeth's presentation focused on the recommendations from the study, expressed a need to investigate the differences in effects of implementation of various management measures, and noted that the status of the stocks continued to decline.

23. **Puerto Rico.** Dr Michelle T. Schärer-Umpierre of the Department of Marine Sciences of the University of Puerto Rico presented the status of spawning aggregations in Puerto Rico.

24. Spawning aggregations of groupers and snappers have been confirmed for a handful of sites in the Puerto Rican archipelago. Various vulnerable, threatened and endangered species of grouper have been documented at some of these multi-species spawning sites with the aid of passive acoustic monitoring studies. Many of these species are extremely rare and hence they are not detected in fishery-independent studies; therefore the study of aggregations provides an efficient method to monitor their populations.

25. Of the spawning aggregation sites highlighted in Puerto Rico one is permanently protected from fishing year-round, three have seasonal protections and a three remain unprotected despite research documenting them. Current seasonal bans for some of the species that aggregate to spawn are applied island-wide, but differ in compatibility between local and federal regulations in the exclusive economic zone (EEZ). Compliance and enforcement efforts at sea are very limited and the effectiveness of seasonal bans is not perceived in local restaurants and markets and there is no export of these products to other locations.

26. Current fishery-dependent data available for these species is unsuitable for trends analyses. Difficulties associated with inconsistent data collection methods, lack of species-specific landings, misreporting from commercial fisheries and little or no information from the recreational sector make population evaluations problematic. The fishery-dependent recreational fishery data available is limited and contains high uncertainty due to the rarity of many of these species.

27. Questions asked after this presentation related to the effectiveness of management measures applied, occurrence of illegal fisheries, confiscation of illegally caught red hind, and why a buffer area around sites where fish aggregate to spawn is needed.

28. **Cayman Islands.** Mr Phillippe Bush, of the Marine Conservation Board of the Department of Environment of the Cayman Islands presented the “Historical and proposed future management of the Nassau grouper spawning aggregations of the Cayman Islands”.

29. Monitoring of the Nassau grouper spawning aggregation-based fishery of the Cayman Islands began in 1987, due to earlier complaints and reports from fishermen of decreasing catches and fish size. Fifteen years of data (1987–2001) from three main historical spawning aggregations showed declining trends in catch, size, and CPUE. In 2001 and 2002, approximately 4 000 fish were taken from a newly discovered spawning aggregation at the west end of Little Cayman, essentially halving a pre-fishing aggregation estimated at 7 000–8 000.

30. As a result of public outcry, 2003 saw the first (and only) “no-take” year based on “alternate year fishing” regulations, and a defined spawning season of November through March. Based on ageing and validation work done earlier, an 8 year (2004–2011) fishing ban prohibited the taking of Nassau grouper from any of the eight designated grouper spawning areas. In 2011, a second consecutive eight year ban (2012–2019) was implemented. In 2012, a conspicuous recruitment pulse of 1–2 year old juvenile Nassau grouper (total length ranging from 12–26 cm) occurred. This was the first time a recruitment of this magnitude was detected in ten years since the cessation of the fishery in 2003. Current frequent sightings of larger sub adults (30–40 cm) suggest much lower level recruitment events in prior years. This underpins importance of long term protection in maximizing chances of meaningful recruitment events. Thus, realistically, protection for depressed Caribbean stocks should therefore be in perpetuity.

31. The currently proposed legislation includes:

- Placing the species on protected status lists (i.e. prohibit the taking of the species anywhere in Cayman waters). Once recovered, the productivity of its population can provide a healthy non-spawning season fishery. (This is the most desirable option).
- Implement an annual closed season throughout Cayman waters for Nassau grouper from November through March.
- Impose a daily catch limit of two fish/person/day in open season.
- Impose a slot size limit of 45–60 cm.
- Ban the taking of Nassau grouper from all designated grouper spawning areas indefinitely.
- Change the current boundaries of designated grouper spawning areas to more realistically accommodate the potential shifting of spawning aggregations.

32. The discussion that succeeded the presentation revolved around the success of the eight year bans and why a permanent ban may be necessary, the proposed legislation and the question whether fishing should be allowed on recovered populations.

33. Dr Brice Semmens of the Scripps Institution of Oceanography of the University of California presented some work of the Grouper Moon research program in the Cayman Islands.

34. The Grouper Moon research program, a collaborative effort between the Reef Environmental Education Foundation (REEF) and the Cayman Islands Department of the Environment, uses a diverse array of field techniques in order to study the population and spatial biology of Nassau grouper (*Epinephalus striatus*). The Cayman Islands maintains a uniquely large (healthy) spawning aggregation of Nassau grouper (~4 000 fish), in addition to several heavily depleted spawning aggregations of the species. Acoustic tagging studies on both the healthy and depleted spawning aggregations indicate that all or nearly all reproductively mature individuals aggregate each year, and do not make abyssal migrations between islands. The acoustic data also suggest that individual grouper may visit multiple aggregation sites before ultimately coalescing at a single site. Finally, acoustic data revealed that larger (more fecund) fish aggregate longer than smaller fish, and that regardless of size, all fish appear to aggregate over a longer period of time at depleted spawning sites.

35. Taken together, these findings suggest a set of behavioral characteristics that present a mechanistic underpinning to the apparent hyper-stability in aggregating species; hyper-stability refers to the fact that catch per unit effort remains relatively constant despite steep declines in catch. The fact that hyper-stability is mediated by spawning behaviors suggests that efforts to harvest aggregating species during their spawning season will likely stymy traditional fisheries management and assessment approaches.

36. Questions after the presentation related to the size of the spawning aggregation and the time spent by fish on the aggregation site, the reason why catchability is higher in smaller populations, and the movement of fish to and from spawning aggregations.

37. **The Bahamas.** Mr Lester Gittens of the Department of Marine Resources of the Ministry of Agriculture, Marine Resources and Local Government presented the status of spawning aggregations in The Bahamas.

38. Though many fish species aggregate in The Bahamas, the Nassau grouper, Yellowfin grouper, Mutton snapper and Lane snapper have been targeted at spawning aggregations. Other than the iconic Nassau grouper, not much is known about the status of these resources either by species or by individual spawning aggregation. While quantitative evidence of management success is largely limited to a study that showed a greater diversity of groupers in the Exuma Cays Land and Sea Park, fishers also make anecdotal reports that there are greater numbers of small Nassau grouper. Nevertheless, despite the use of closed seasons in most years since the late 1990s (along with other older management measures) and abounding educational efforts led by non-governmental organizations (NGOs), the overall Nassau grouper fishery is estimated to range from fully exploited to overexploited. In addition, studies of a few individual Nassau grouper aggregations showed the disappearance of some aggregations and greatly reduced numbers in others.

39. Like regional counterparts, The Bahamas is challenged with finding the right combination of enforceable management measures that simultaneously facilitate food security, sustainability and the ability of fishers to right now earn a living. This can only be achieved by excising the current scourge of poaching (foreign and local) in addition to further embracing informed management decisions. Likewise, informed management implies that there is information to base decisions on. More resources must be contributed towards enhancing these sources of information including surveillance for enforcement purposes, accurate monitoring of landings, scientific research and stock assessments.

40. In the discussion that took place after the presentation the success of a closed season combined with a sales ban compared with a closed season without sales ban was an issue. The impact of the closed season on the stocks of Nassau grouper was discussed also and the need for a regional recommendation on the use of fish traps with biodegradable panels.

41. **Mexico.** Dr Alfonso Aguilar-Perera of the Universidad Autónoma de Yucatán presented the status of spawning aggregations in the Southern Gulf of Mexico and Mexican Caribbean.

42. In the Southern Gulf of Mexico and Mexican Caribbean, a lack of detailed knowledge prevails on the current conditions and fishery status of fish spawning aggregations (FSAs). Limited scientific documentation has revealed that grouper species (*Epinephelus striatus*, *E. itajara*, *E. guttatus*, *Mycteroperca bonaci*, *M. tigris*, *M. venenosa*) and snappers species (*Lutjanus analis*, *L. cyanopterus*, *L. synagris*, *L. jocu*, *Ocyurus chrysurus*) are opportunistically exploited during spawning aggregations. The practice of exploiting groupers, such as the Nassau and the Goliath, has been progressively fading because of population declines of these groupers. In fact, no fishermen community now strongly depends economically on fishing these aggregations.

43. There are no legal provisions by the Mexican Government for management of FSAs in the region. Most attention is paid to management of the red grouper, *E. morio* and the red snapper,

*Lutjanus campechanus*. The only regulation for grouper fishing is a one-month ban (February 15 to March 15 every year) established in 2005 for all grouper species (about 17). Also, a normative regulation (NOM-065-PESC-2007) established in 2010 provides complementary criteria to regulate the grouper fishing. None of these latter legal instruments consider the existence of FSAs. The equivalence of a Species Red List in Mexico is the NOM-059-SEMARNAT-2010, which only includes one commercially, marine, exploited teleost: *Totoaba macdonaldi* endemic to the Gulf of California.

44. The presentation contained a range of proposals to improve management, including introduction of co-management for FSAs. The subsequent discussion stressed the need for a regional coherent approach and that stock assessments should be conducted for more species.

45. **Cuba.** Mr Servando Valle of the Centre for Fisheries Research of Cuba presented, on behalf of Mr Rodolfo Claro, the status of spawning aggregations in Cuba.

46. Traditionally, the catches of reef-associated fin fishes in Cuba have shown strong seasonal trends, mainly associated with the reproductive periods of the most economically important species, among them snappers (*Lutjanidae*) and groupers (*Serranidae*). This seasonality in catch trends is determined by the increased vulnerability of aggregating species to fishing during the reproductive period and the resulting focus of fishing activity on spawning aggregations which yields a large proportion of annual catches of such species. Since the responses of aggregating species to fishing vary according to the biology of different targeted species, an understanding of the impacts of fishing and the consequences of management can only be understood by species-specific analyses in the context of the coastal fishery and its management history as a whole. The history of the coastal, reef-associated, fishery of Cuba is one of increasing and decreasing fishing pressure and variable management effectiveness that ultimately led to substantial declines in most key commercial species.

47. Snappers and the Nassau grouper are traditionally considered to be the major fin fish resources in Cuba, but many of these species have declined over the last four decades. The fishery of snappers and groupers typically concentrated on seasonal spawning aggregations. Twenty-two spawning aggregation sites were identified around the Cuban Shelf. Most of these sites are sequentially used by several species in different times. Some other sites may be found, but probably most important sites for massive spawning of targeted species are included.

48. Direct observations of spawning events have been rare in Cuba and more information on the population size of past and current spawning aggregations is needed. Active spawning aggregations due to their discrete nature and high productivity are clearly important resources. This emphasizes the need to validate aggregation information when available. Use of these spawning aggregation sites may vary temporally under natural conditions or be fully eliminated due to fishing pressure; therefore, efforts to confirm the existence of nominal aggregation sites and monitor their production through time will be essential to optimal reserve design and management.

49. Following the presentation the effects of a closed season were discussed as well as the identification of historical sites of Nassau grouper spawning aggregations. It was argued that the spawning aggregations in Cuba are often far at sea and that rough weather plus the distance to the site are not permitting the fishers to fish at many of these spawning aggregations.

50. **Belize.** Mr Mauro Gongora of the Fisheries Department presented the status of spawning aggregations in Belize.

51. Since 2003, the Government of Belize passed legislation to protect several commercially important fish species at 11 spawning aggregation sites distributed along the coast and in the three atolls of Belize. The major declines in the number of fish species, and in particular, the Nassau grouper (*Epinephelus striatus*) in the spawning aggregation sites, as demonstrated by studies done,

prompted the passing of Statutory Instrument numbers 161 and 162 of 2003 to protect the Nassau grouper and several other fish species.

52. Currently, seven spawning aggregation sites are monitored regularly. The monitoring teams are guided by the Reef Fish Spawning Aggregation Monitoring Protocol for the Mesoamerican Reef and Wider Caribbean. The inconsistency in spawning aggregation data collection as a result of the lack of resources has not helped the spawning aggregation working group to determine whether a particular fish species or more that aggregate to spawn in Belize have either recovered or have declined even further. It is clear that more resources are urgently needed to conduct additional field research and fisheries law enforcement activities at these sites to deter illegal fishing. This is a major challenge and needs to be addressed through a coordinated national and regional approach.

53. The presentation was followed by a discussion on the monitoring data, timing of monitoring, manpower available for monitoring and the concern about Nassau grouper aggregations moving between Belize and Mexico's EEZs, which requires subregional collaborative research.

54. **Nicaragua.** Mr Renaldy Barnuty Navarro of the Nicaraguan Institute for Fisheries and Aquaculture (INPESCA) made a presentation on the status of finfish fisheries in the Caribbean Sea of Nicaragua.

55. Finfish fisheries are the most important in Nicaragua in terms of volume landed and from the social point of view, because it is carried out mainly by artisanal fishermen. In Nicaragua finfish fisheries usually operate in environments dominated by multispecies landings comprising mainly snappers and groupers (*Lutjanidae* and *Serranidae*), snook (*Centropomus spp*), sharks (*Carcharhinidae*, *Triakidae*) and croakers (*Sciaenidae*). The highest landings of finfish originate from the Pacific and are clearly dominated by snappers. The boats used for fishing are mostly fiberglass boats with lengths between 5–10 m and outboard motors up to 75 HP. The crew of two to three people uses a variety of gears, such as gill nets, trammel nets, cast nets and lines with hooks.

56. In the case of finfish landings in the Caribbean Sea, the snappers (*Lutjanus spp.*) show a clear predominance followed by the snooks (*Centropomus spp*), and the group of other fish, followed by groupers (*Epinephelus spp.*) and sharks (*Carcharhinidae and Triakide*). For all species, there are growing trends in landings over the last five years.

57. The landings of snapper species in the Caribbean of Nicaragua are increasing and this is mainly due to the improvement of national and international market prices and an increased fishing effort mainly by the industrial fleet which is using traps. In the case of groupers, stability is observed in the landings over the last five years. The snapper and grouper landings in 2012 were equivalent to respectively USD2.4 million and 270 000 pounds round weight.

58. The trends of landings for the Caribbean groupers show that they were decreasing until 2009. After this period, there is a stabilization in the order of the thirty thousand pounds harvested per month. Major grouper species that are landed are the black grouper (*Mycteroperca bonaci*) Warsaw grouper (*Hyporthodus nigrurus*) and yellow grouper (*Epinephelus flavolimbatus*). The monthly snappers landings showed an increasing trend, the main species that are landed is yellowtail snapper (*Ocyurus chrysurus*) with 71 percent, followed by 7 percent yellow eye snapper (*Lutjanus vivanus*) and the black end snapper (*Lutjanus buccanella*).

59. In Nicaragua there have been a few studies on the biology and dynamics population of finfish. INPESCA, responsible for the management and wise use of fishery resources of the country and as the competent authority for the application of Law 489, Law on Fisheries and Aquaculture and Regulations, established minimum sizes for fish species catches from the Caribbean Sea and the Pacific Ocean in Nicaragua. (Executive Resolution 003-2012). The minimum size is established based on studies and regulations established and conducted in other countries, such as Mexico, Jamaica and

the United States of America and by applying the precautionary principle and the Code of Conduct for Responsible Fisheries (the Code) of FAO.

60. Measures established to promote the use of the minimum size:

- Mesh size regulation for gill nets, traps and the size of hooks used in fishing - target fish by a Technical Standard Fishing Gear and Methods.
- Releasing live fish caught that are below the minimum size.
- Prohibition of fishing in breeding and nursery areas.
- Implementation of the Code.
- From 2012 onwards, monthly biological sampling of snappers in the Pacific Ocean.

61. In Nicaragua, closed seasons or quotas for finfish fisheries have not been established and today these fisheries are considered open-access fisheries. In the case of sharks, an indefinite closed season for species that penetrate inland waters exists.

62. The presentation was followed by some discussion on how minimum fish size regulations are enforced in practice. Examples of collaboration between the fishing authorities, navy, coast guard and police were given. The limited monitoring and few stock assessment studies being done were issues of concern raised.

63. **Caribbean Netherlands.** Mr Pieter Van Baren of the Rijksdienst Caribisch Nederland presented the status of spawning aggregations of commercially exploited aggregating species of the Caribbean Netherlands.

64. Various grouper and snapper species are exploited commercially. This is being done in an artisanal manner with hook and line and fish traps being used as gear. The status of FSA's in the Caribbean Netherlands is largely unknown. Currently, there is one known targeted multispecies (red hind and queen triggerfish) FSA off the coast of Saba. The red hind (*Epinephelus gattatus*) is being targeted commercially whereas the queen triggerfish (*Balistes vetula*) for recreational use. Red hind is being exported from Saba, mainly to St. Maarten.

65. In 2005, a study on the FSA was carried out of which the outcome was that the FSA was moderately exploited. Since then fishing pressure has increased tremendously and it is being presumed that the spawning aggregation is heavily overfished. As of 1<sup>st</sup> December 2013, the FSA will be closed during the months of December, January and February for the next five years. During this time the FSA will be monitored and after five years, the measure will be evaluated to see if prolonging is required and if additional measures are necessary.

66. On Bonaire and St. Eustatius there have been reporting's of FSA's in the past. Currently, there are no known FSA's near these islands.

67. The discussion that succeeded the presentation referred to a recommendation on Nassau grouper, which came out of the fourth Scientific Meeting and Technical Advisory Committee (STAC) of the SPAW protocol that discussed whether there was a need to protect marbled grouper and the threat of increased fishing pressure on FSAs following their identification through research efforts.

68. **Brazil.** Dr Athila Bertocini, of the Federal University of the State of Rio de Janeiro made a presentation on reef fish aggregations in Southern Brazil: Pró-Arribada and Meros do Brasil Initiatives.

69. The presentation was followed by some questions that related to the incentives for fishers and other stakeholders to monitor goliath grouper sticks and FSAs, the type of environmental education and the focus of the research projects on dusky grouper and goliath grouper.

70. **USA – Atlantic Coast.** Dr Ken Lindeman of the Florida Institute of Technology presented an overview of “Snapper and Grouper Spawning Aggregation Information for the United States Atlantic Coast”. His presentation summarized joint research work with G. Sedberry, M. Meadows, M. Burton, T. Kellison, N. Farmer, M. Reichert, D. DeMaria, C. Koenig, D. Morley, A. Acosta, C. Taylor, W. Heyman, S. Harter, and A. David.

71. They surveyed literature, unpublished data, and interviewed fishers to identify known and potential spawning aggregation sites for the snapper and grouper reef fish faunal complex of the United States Atlantic coast. Focal species included the 14 *lutjanid* and 18 *serranid* species (five and four genera, respectively) managed under the Snapper-Grouper Fishery Management Plan of the United States South Atlantic Fishery Management Council on diverse reef systems from the lower Florida Keys through North Carolina. Criteria were based on Domeier and Colin (1997) and use of local fisher information to supplement research data.

72. Nine snapper species have confirmed or potential spawning aggregation sites identified on the United States Atlantic coast. Mutton and cubera snapper had the most known spawning sites (13–15). There is considerable evidence of simple migratory spawning and some evidence of spawning aggregations for *L griseus*, *L campechanus* and five other species. In total, >40 confirmed or potential *lutjanid* spawning sites were identified. Confirmed and potential spawning aggregation sites were identified for seven grouper species with 20–30 total sites. Of these, at least nine goliath grouper aggregations have been confirmed by Koenig and Coleman (2013) in the Jupiter Inlet area of East Florida (27°N).

73. The majority of known or potential aggregation sites for the southeast United States snapper-grouper reef fish complex are subject to few specific management measures to ensure aggregation sustainability; however where no-take areas are enforced, in situ data are positive for some aggregation sites. Monitoring and research have often been constrained by funding and few data to fully characterize potentially important spawning aggregations are available for the majority of sites.

74. The presentation was followed by discussion on the comparative effectiveness of spawning season closures. It was also noted that the Society for the Conservation of Reef Fish Aggregations had many outreach materials on its website and that fisher outreach should be scaled up through outreach campaigns as part of management measures.

75. The second day of the Working Group meeting started with summarizing the main findings and conclusions from the first day. A number of important additional observations were made related to the following issues:

- The multi-species, multi gear reef fisheries and aspects of fishing down the food web, given that many of the more vulnerable aggregating species are at the top of the food web.
- The public-value of aggregating species (food, tourism, earnings), along with ecological and biological values.
- The need to acknowledge as part of the ecosystem value that top predators contribute to the Caribbean marine ecosystem and the ecosystem role of groupers and other species that aggregate to spawn.
- The need to have a minimum standard regional closed-season for aggregating species, like there exists for lobster at the sub-regional level, given the dispersive larval phase of aggregating species and challenges for enforcing regulations, especially when there is international trade.
- The possibility to develop eco-tourism around spawning aggregations.
- The involvement of fishers in the management of spawning aggregations, as well as in spawning aggregation research and verification of spawning aggregations.

- The outreach and fishers exchange programmes that could contribute to increasing awareness and involvement of fishers in the management of spawning aggregations.
- The need to put in place threatened species legislation, as many countries in the region lack such legislation.
- The need for enforcement of existing regulations and monitoring of aggregating species.

76. Some additional questions were posed to the presenters of the national status reports. These questions related to the enforcement of fish size, catch and gear regulations, the ways to reduce fishing pressure and fleet capacity, the functioning of spawning aggregation working groups at national level (e.g. in Belize), alternative employment options for fishers during closed seasons, social development programmes that fishers can tap into, and the involvement of fishers in spawning aggregation monitoring programmes.

### **Biogeography of transient reef-fish spawning aggregations in the Wider Caribbean**

77. Dr Shinichi Kobara of Texas A&M University presented a brief summary of the recently published review paper, “Biogeography of transient reef-fish spawning aggregations in the Caribbean: a synthesis for future research and management.” The review evaluates all currently known and documented transient reef fish species and their spawning aggregation sites in the Wider Caribbean. In this region, 37 species of fish from ten families form transient FSAs and there are at least 108 geographically discrete transient FSA sites. Nassau grouper aggregations were the most commonly documented spawning aggregations (55 sites) and 32 sites had multispecies aggregations.

78. Dr Kobara emphasized the importance of bathymetric data collection in characterizing spawning aggregations. Even relatively crude bathymetric information can support site characterization and help design of appropriately sized marine protected areas (MPAs). Bathymetric information can also help in the understanding and modeling of hydrodynamics – water mass movement around the spawning site – and thus the influences on larval transport from the site. Finally, and perhaps most importantly, bathymetric data can be used to predict previously unknown spawning sites.

79. There are 18 multispecies sites that have bathymetric data available in this region. For every site, the spawning aggregation occurred at a shelf edge, adjacent to relatively deep water, and a reef promontory. Although it might not be applicable for every single-species spawning aggregation site (e.g. red hind spawning aggregation sites in Puerto Rico), the geomorphological approach has been used to predict and find a previously unknown multispecies spawning aggregation site in Belize. The approach might prove feasible in other locations as well.

80. Dr William D. Heyman of LGL Ecological Research Associates, Inc., continued the presentation on the research undertaken.

81. Many large groupers and snapper species can be considered as components of a snapper-grouper complex – a suite of species that share similar life history characteristics that are harvested as part of multi-species fisheries throughout the wider Caribbean. Many of these species are over-fished; some are threatened or endangered. Though many are considered data-poor species in that their status has not been successfully evaluated. These fishes are generally long-lived, late to reproductive maturity, and spawn in massive transient aggregations – all contributing to their vulnerability to over-exploitation.

82. Dr Heyman categorized research on aggregations into eight levels with increasing cost and sophistication. He identified the minimum data needed for management action: a site map and characterization using fisher interviews, fishery dependent surveys, and underwater visual counts and documentation with photos or video. He further documented that research can be conducted and sites

protected more efficiently by involving local aggregation fishermen in all aspects of the research and management process.

83. Dr Heyman offered support for the hypothesis that multi-species spawning aggregations occur predictably at the tips of reef promontories, at shelf edges in 15–60 m water depth, adjacent to deepwater (>200 m). This search image has been used to predict the location of multi-species spawning aggregations in Belize and Mexico, and may prove useful throughout the wider Caribbean and the Gulf of Mexico. He offered a vision of the future whereby a network of multi-species spawning aggregations are protected and monitored with a standard protocol, promoting recovery of the Wider Caribbean snapper grouper complex.

84. The discussion that succeeded the presentations focused on the interconnection of the grouper/snapper complex and the need to update a spawning aggregation monitoring manual.

### **ESA, CITES, SPAW PROTOCOL**

85. Ms Stephania Bolden of the National Oceanic and Atmospheric Administration, National Marine Fisheries Service and Southeast Regional Office made a presentation on the regulatory tools: the Endangered Species Act (ESA), the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and the Protocol Concerning Specially Protected Areas and Wildlife (SPAW).

86. The presentation outlined what CITES involves and does and discussed the CITES appendices. The benefits of CITES listing in an appendix were noted as well. An overview was given also of the United States ESA and the various ESA sections of relevance to the process of listing endangered marine aquatic species under ESA.

87. The Cartagena Convention, the only legally binding environmental treaty for the region, was discussed as well, with emphasis on the SPAW Protocol. Ms Bolden detailed that the SPAW Protocol assists governments in the Wider Caribbean region to:

- protect and recover certain species;
- protect areas and ecosystems;
- develop technical and scientific research on these areas and species, and exchange and coordinate information concerning research or monitoring programmes.

88. The SPAW Protocol establishes the principle of coordination of measures, criteria and guidelines corresponding to these different objectives. The Protocol includes three species lists to protect listed flora (Annex I), fauna (Annex II), and species of flora and fauna to be maintained at a sustainable level (Annex III). The Protocol became international law in 2000 and 16 countries have ratified; however SPAW ratification is not necessary for collaborative activities.

89. Ms Bolden added that at the fifth Meeting of the STAC to the Protocol Concerning Specially Protected Areas and Wildlife (SPAW) in the Wider Caribbean region, a report was submitted by the Government of Cuba with recommendations for listing of species under the SPAW Annexes, which given time limitations, could not be previously considered. That list included, amongst others, the Nassau grouper.

90. The Working Group took note of the three regulatory tools and recognized that CITES and SPAW listings are only useful if they receive follow-up from the countries. It was noted that CITES is not a management body as such and that all CITES decisions are to be carried out by the national governments. There was some resistance among various of the country delegates to have any of the grouper species listed under CITES. It was argued further that most Nassau grouper currently caught

ends up at domestic markets and that thus the CITES tool would not add much to the management of spawning aggregations.

91. In terms of ESA the ongoing process following the proposal for the listing of queen conch was explained. It was noted that ESA listing will have far reaching consequences and that any ESA related follow-up recovery plans are limited to the USA EEZ only; there is unlikely to be management support for other countries.

92. The listing of Nassau grouper under an appendix of the SPAW protocol was an option favoured by many experts in the Working Group. It was recognized that the distribution of Nassau grouper has dwindled at the regional level. While at the national level, in some cases, the stocks are not under threat it is a different situation when looking at the regional trends of landings of Nassau grouper by fisheries and occurrence of spawning aggregations. It was noted however that there are large gaps in terms of ratification of the SPAW protocol in the region and that the recommendations of it are non-binding, i.e. voluntary.

## WORKING GROUP DISCUSSIONS

93. The meeting decided to split into three break-out working groups:

- Group 1: Importance of aggregating species for food and income and the need to manage/conservate them.
- Group 2: Research and monitoring of aggregating species
- Group 3: Educational and outreach – experiences and challenges

94. **Group 1** (Importance of aggregating species for food and income and the need to manage/conservate them) summary of the group discussions:

- It is a “no-brainer” that the aggregations that are commercially fished need to be protected or managed in order to have populations of fishes that form aggregations in the long term (includes most reef fishes such as parrotfish, surgeonfishes, snappers and groupers as well as pelagic sp. i.e. flying fish).
- Fisheries regulations regarding traps should be revised to make sure they do not impact the species that form spawning aggregations (timing, mesh size, biodegradable panels, etc.).
- It is important to highlight the cultural, economic and ecological value of the fish populations for the livelihood of many Caribbean communities.
- There was much discussion regarding the need for highlighting the importance of fish spawning aggregations for “food security” in some of these countries since many coastal communities depend on many different coral reef species which aggregate.
- The urgency of this matter was highlighted since we have been discussing among us for many years the need to protect FSA.
- It would be useful to develop a report card for the Caribbean countries, similar to “Reefs at Risk” to present which countries have FSAs and how they are doing.
- Fisheries management bodies haven’t all incorporated the protection of FSA as a priority and this is worrisome, but perhaps due to short-term alternatives such as aquaculture that don’t really shift effort but add to it.
- An effort has to be made to highlight the urgent need for protecting FSAs to fisheries managers and include the list of solutions that they can use to act, not leave it at “here is the problem and deal with it”.

- The cascading benefits of protecting FSA due to ecological integrity i.e. improved coral reefs.
- The plight of FSAs needs to be encapsulated into a marketing campaign for which the information already exists (SCRFA) and the audience should be two-fold one bottom-up (public and constituency) and another top-down (government levels).
- We need to develop a marketing plan focused on the importance of FSAs for the fisheries species, then seek the support of NGOs and other partners to be able to implement across Caribbean and the Gulf of Mexico.
- Mexico could be a good pilot project for the implementation of this marketing plan.
- Some type of eco-label could be applied to products from sustainable “non” FSA fisheries.
- Perhaps some “earned media” coverage could be used to highlight the importance of protecting FSA.
- The message of the importance of FSAs should be transmitted through the voice of fishers, i.e. the ‘Ambassadors’ that can take the message to their government agencies and stakeholders.

95. The discussion that followed Group 1’s presentation added that, for coastal communities’ food security, and for long-term general food security, the management of FSAs is essential. It was noted that no new eco-label may be needed as there are so many around already and that some linkage with the Marine Stewardship Council (MSC) certification scheme may be an option.

96. **Group 2** (Research and monitoring of aggregating species) summarized its discussion in the following table:

	<b>Realities/ current situation</b>	<b>Needs/GAPS desired situation</b>	<b>Approaches to bridge gaps/ Action required and who to act</b>
<b>Identification</b>			
Issues: ongoing work	<ul style="list-style-type: none"> <li>• Discovery based monitoring (detailed maps – occurrence)</li> <li>• Acoustic monitoring</li> <li>• Cuba has lengthy historical landing information</li> </ul>	Know what is/are most important spawning sites regionally Know the migration patterns of each species – to inform management and conservation	In Turks and Caicos islands, no fishing for Nassau grouper, thus it would be opportunity to identify spawning sites.
<b>Research</b>			
Issues:	Countries fearful of other countries getting hands on data  Researchers that collect data and information in other countries need to provide info/findings to these countries	<ul style="list-style-type: none"> <li>• Effective data and info exchange at regional level</li> <li>• Raw level data can be kept internal; only aggregated data be shared</li> <li>• All Small Island Developing States (SIDS) should identify spawning aggregation sites</li> <li>• Socio-economic dependence on aggregations and perception related to conservation of aggregations</li> </ul>	Website – database – digital library established.  NGOs conservation community is needed to collect data.  Transfer of technology and knowledge from south-east Asia may be useful

		<ul style="list-style-type: none"> <li>• Ecological indicators</li> <li>• Reference points for management</li> <li>• Stock assessment methods for spawning aggregation species</li> <li>• Restocking of wild stocks might provide options in some cases</li> <li>• Threshold density for aggregating species?</li> <li>• Artificial fertilization – aquaculture – survival rate of larvae is low</li> </ul>	
<b>Monitoring</b>			
Issues:	<p>Shifts in aggregations are occurring (MPAs were established but now fish aggregate elsewhere – albeit nearby – in some cases)</p> <p>Difficult to find aggregations without fishers knowledge/information</p> <p>Allow a few cooperating fishers to fish on specific days (note that fish won't bite on actual spawning day) only – to ensure data/ information requirements are met, and only a small part of the fish is caught (use of single hook and line – one day per month -17 vessels – two fishers per vessel in Belize; estimated that 18–20% is caught). Limited access, rights based fisheries.</p> <p>Visual surveys are done additionally as well as cross-checking of info with exporters.</p> <p>NGO community has created public awareness and conservation efforts promotion.</p>	<p>Legislation + voluntary collaboration with fishers</p> <p>Monitoring manual to be updated and endorsed by WECAFC, CRFM, etc.</p>	<p>Fisherfolk cooperation is required to identify and monitor spawning aggregation sites</p> <p>May be needed to issue special permit to allow few fishers to fish site for limited time – in return for full collaboration/info from fishers</p> <p>Acoustic monitoring may include also vessels monitoring during aggregation events</p>

97. The presentation of the above group discussion was followed by a plenary discussion in which the issue of identification of spawning sites got most attention. Some experts regarded it important not to reveal the spawning sites in cases where there are no management measures in place to protect the spawning aggregation. Moreover, it was argued that there are often no immediate benefits for fishers to inform the government or researchers of existing spawning sites. Local institutional capacity is often too limited to monitor and manage spawning aggregation sites effectively and more remote aggregation sites are often difficult to protect/monitor due to the high costs related to doing so.

98. **Group 3** (Educational and outreach – experiences and challenges) summary of discussions:

99. What is the goal? We need to have people broadly understand, but also to ACT.  
WHO DOES the OUTREACH? Fishermen

“Low Hanging Fruit”

1. Capture all of the existing outreach information in a single website that links to all existing education and outreach information.
  - SCRFA website
  - Project Goliath
  - TAMU Geography UTube
  - Reef resilience Website (TNC)
  - Exchange videos
  - Spawning aggregation working group site
  - REEF – kids programme in Cayman Islands, live uplink from aggregation
2. Support local fishermen leaders as spokespersons and Ambassadors
  - Awards for student paper
  - Award for best fishermen
  - Gladding Award Winner
  - Training fishermen on policies and public speaking
  - CFMC to pay for travel expenses for fishers to attend
  - Travel expenses for fishermen to attend GCFI
3. Messaging might be targeted for various audiences
  - Fishermen = value in supporting sustainable source of seafood
  - Divers – a great dive experience
  - Broad general public – anyone who knows about the sea should know
  - Decision makers/managers – protect multi-species aggregations year round
4. Tools to use:
  - Live Traveling educational shows
    - ✓ Mero-movil Grouper-mobile
    - ✓ Sailboat – educational boat
    - ✓ Carnival or travelling festival
    - ✓ Children’s play
  - Fisher exchanges
  - Social media
    - ✓ YouTUBE
    - ✓ Facebook
    - ✓ iTunes University
  - Get companies who want to advertise to contribute and use the platforms for their own purpose of advertising.
5. Existing Television or Video outlets
  - Wild Krat’s Episode
  - National Geographic Film
6. Certification of species – Work with MSC to have them consider whether a species aggregates to spawn and if a fish was caught outside time and location of spawning aggregations – as part of the certification process

7. Fishermen Ambassador Programme
  - Through GCFI
  - Gladding Award Winners
8. MESSAGE SHOULD COME FROM FISHERMEN
  - Videos of fishermen talking to fishermen
  - Fisher exchange videos
  - “At Sea Level”
9. Possible Donors: Ballard Foundation, watch leaders – Ocean Exploration Trust
10. Donors must also remember that regional bodies need support
11. Teacher training programme
  - Packages for interested schools – grouper day, curricula

100. In the discussion that followed Group 3’s presentation the CFMC Secretariat referred to the importance to have a teacher’s manual for educating school children on spawning fish aggregations.

## GENERAL DISCUSSIONS

101. Following an example provided by the CFMC/WECAFC Working Group Secretariat, the participants were requested to complete overview tables of fisheries management measures for aggregating groupers and snappers in each of the participating countries. The overview table included input controls (e.g. closed areas, closed seasons, gear restrictions, method restrictions, effort restrictions, and licenses) as well as output controls (e.g. harvest restrictions, length limits, bag/catch limits, fish holding restrictions, sale/market restrictions, trade restrictions and landing requirements). The completed overview tables will be published in the updated regional status overview in a separate report.

102. Ms Elizabeth Mohammed of the Caribbean Regional Fisheries Mechanism (CRFM) Secretariat made a presentation on the work of CRFM in relation to fish spawning aggregations.

103. The CRFM is an inter-governmental organization which seeks to promote and facilitate the responsible utilization of the region's fisheries and other aquatic resources for the economic and social benefits of the population of the region. Currently the Mechanism comprises seventeen member States, of which spawning aggregations are documented thus far to be of importance to Jamaica, Belize, the Bahamas and Antigua and Barbuda.

104. Through the CRFM Reef and Slope Fish Resource Working Group, which meets at the CRFM Annual Scientific Meetings, data analyses have been conducted for several fisheries targeting snappers (*Lutjanus purpureus*, *L. synagris*), groupers (*Epinephelus guttatus*, *E. striatus*) and other reef and slope species. Management objectives do not focus specifically on fish spawning aggregations, except perhaps in the case of Belize and Jamaica, but address inter alia the need for long-term sustainability of the resource, application of the ecosystem approach, rebuilding depleted fish stocks in nearshore areas, protection of essential fish habitat, regulation of fishing effort, fishing areas and size of fish in the catch and control of the alien invasive species, *Pterois* spp. (Indo-Pacific lionfish).

105. Current management measures include, to varying degrees among member States, effort regulation through licensing systems, mesh size regulation, closed seasons, reduction in ghost fishing and establishment of marine protected areas. Weak monitoring, control and surveillance capability continues to impede effective management. Generally, stock assessment results have been inconclusive due to uncertainties regarding stock identification, distribution and level of sharing

among countries and inadequate catch, effort and biological data. Consequently, to improve the quality of stock assessments and management advice provided, future data collection and research efforts should focus in these areas as well as collection of industry socio-economic data; identification of spawning locations; consideration of environmental data in assessment modelling; biomass, ecological and economic evaluation of fish spawning aggregations, assessment of socio-economic impacts of management measures on fishing communities and examination of alternative livelihood options. Public awareness and education on the need to identify and protect spawning aggregations for long term sustainability of the resource should target decision-making bodies such as the CRFM Ministerial Council as well as direct stakeholders such as the Caribbean Network of Fisherfolk Organizations. The latter could be instrumental in acquiring local ecological knowledge on fish spawning aggregations to inform management.

106. In the discussion that followed the presentation questions were asked about the sub-regional flying fish management plan and when it would come into effect, the effects of fish aggregating devices (FADs) in fisheries, how the working group of CRFM relates to the joint working group with CFMC, OSPECA and WECAFC, and about the need to work jointly on public outreach to increase understanding on spawning aggregations.

#### **PREPARATION AND ADOPTION OF THE REPORT TO WECAFC**

107. The Working Group was presented by the meeting Secretariat with a draft declaration in which the main discussions, conclusions and recommendations from the meeting were combined. The draft declaration was discussed and modified by the Working Group. The final version of the Declaration of Miami, as approved by the Working Group, can be found in Appendix C. The Declaration contains an annex with the recommendation to the sixth session of the Scientific Advisory Group (SAG) of WECAFC and the fifteenth session of WECAFC on the “Establishment of a regional closed season for fisheries in the WECAFC area to protect spawning aggregations of groupers and snappers”. The sixth session of the SAG reviewed and endorsed the Recommendation on 3 November 2013. The Recommendation to the fifteenth session of WECAFC can be found in Appendix D.

#### **CLOSURE OF THE MEETING**

108. Mr Miguel Rolon, on behalf of CFMC, thanked the Working Group members and other meeting participants, the co-organizers, the members of the CFMC/WECAFC meeting Secretariat, chairperson and interpreters for their active participation and their contributions to the success of the meeting.

109. The meeting was declared closed by Mr Rolon, on Thursday 31 October 2013, at 13:00 hours.

**Agenda**

1. Opening of the meeting
2. Election of the Chairpersons and rapporteurs
3. Adoption of the agenda and arrangements for the meeting
4. Global perspective of aggregating snappers and groupers
5. Historical background of WECAFC's Work on Spawning Aggregations of key species
6. Presentations of national status reports by each of the participants
7. Biogeography of Transient Reef Fish Spawning Aggregations in the Caribbean
8. ESA, CITES, SPAW Protocol
9. Working Group Discussions
10. General discussions
11. Preparation and adoption of the report to WECAFC
12. Closure of the meeting

## List of participants

**BAHAMAS****Gittens, Lester**

Dept. of Marine Resources  
 Ministry of Agriculture Marine Resources  
 and Local Government  
 PO Box N3028  
 Nassau, Bahamas  
 Tel: (242) 393-1777  
 Email: [lestergittens@bahamas.gov.bs](mailto:lestergittens@bahamas.gov.bs)

**BELIZE****Gongora, Mauro**

Fisheries Department  
 Princess Margaret Drive  
 PO Box 148, Belize City  
 Tel: (501) 224-4552  
 Fax: (501) 223-2187/2983  
 E-mail: [species@btl.net](mailto:species@btl.net)  
[megongora@hotmail.com](mailto:megongora@hotmail.com)

**BRAZIL****Bertoncini, Athila**

Universidade Federal do Estado do Rio de Janeiro – UNIRIO  
 Laboratorio de Ictiologia Teorica e Aplicada – LICTA  
 Ave. Pasteur 458 CCET/IBIO  
 Rio de Janeiro RJ Brazil 24290-240  
 Tel: 5521 980101440  
 Email: [athilapeixe@gmail.com](mailto:athilapeixe@gmail.com)  
[www.merosdobrazil.org](http://www.merosdobrazil.org)

**CAYMAN ISLANDS****Bush, Phillippe**

Secretary, Marine Conservation Board  
 Dept. of Environment, Cayman Islands  
 Government  
 CI Environmental Center, 580 North Sound  
 Road GT  
 PO Box 10202GT (KY1-1002)  
 Tel: (345) 949-8469  
 Email: [phillippe.bush@gov.ky](mailto:phillippe.bush@gov.ky)

**CUBA****Valle Gomez, Servando V.**

Centro de Investigaciones Pesqueras  
 Calle 246 No. 503  
 e/5ta. Avenida y Mar  
 Reparto Barlovento  
 Municipio Playa, La Habana  
 Tel: 209 8055  
 Email: [servando@cip.alinet.cu](mailto:servando@cip.alinet.cu)

**HONG KONG****Yvonne, Sadovy**

School of Biological Sciences  
 Room 3S-01  
 The Kadoorie Biological Sciences Building  
 The University of Hong Kong  
 Pokfulam Road, Hong Kong, China  
 Tel: 852 2299 0603  
 Email: [yjsadovy@hku.hk](mailto:yjsadovy@hku.hk)

**MEXICO****Aguilar- Perera, Alfonso**

Universidad Autónoma de Yucatán  
 Facultad Medicina Veterinaria y Zootecnia  
 Carr. Merida –Xtmakuil, Km. 15.5 A.P 116  
 C.P. 97135  
 Merida, Yucatán, México  
 Tel: +52 (999) 942-3200 Ext. 17  
 Email: [alfaguilar@gmail.com](mailto:alfaguilar@gmail.com)

**NETHERLANDS****Van Baren, Pieter**

Policy Advisor Agriculture & Fisheries  
 Ministry of Economic Affairs  
 Rijksdienst Caribisch Nederland  
 Kaya Internashonal z/n | Kralendijk | Bonaire  
 PO Box 357 | Kralendijk | Bonaire  
 Tel: (work): (+599) 715 83 21  
 Mobile: (+599) 795 90 79  
 Fax: (+599) 717 83 30  
 Email: [pieter.vanbaren@rijksdienstcn.com](mailto:pieter.vanbaren@rijksdienstcn.com)  
<http://www.rijksdienstcn.com>

**NICARAGUA****Barnuty, Renaldy A.**

Instituto Nicaragüense de la Pesca  
y la Acuicultura (INPESCA)  
Km 3½ Carretera Norte, frente a Banpro  
San Luis  
Managua  
Tel: (505) 248 7149/248 7151; Ext 131  
Fax: (505) 248 7151  
E-mail: [rbarnutti@inpesca.gob.ni](mailto:rbarnutti@inpesca.gob.ni)

**PUERTO RICO****Schärer, Michelle**

Department of Marine Sciences  
University of Puerto Rico  
Mayaguez, PR 00681-9000  
Tel: (787) 319-4172  
Email: [michelle.scharer@upr.edu](mailto:michelle.scharer@upr.edu)

**UNITED STATES OF AMERICA****Kobara, Shinichi**

Assistant Research Scientist GCOOS  
GISManager and Application Developer  
Department of Oceanography  
Eller O&M Building  
MS 3146  
Texas A&M University College Station  
Texas 77843-3146  
Tel: (979) 845-4089  
Email: [shinichi@tamu.edu](mailto:shinichi@tamu.edu)  
[Shin.kobara@gmail.com](mailto:Shin.kobara@gmail.com)

**Heyman, William D.**

Senior Marine Scientist  
LGL Ecological Research Associates, Inc.  
4103 Texas Avenue #211  
Bryan, Texas 77807  
Tel: (979) 775-2008  
Mobile: (301) 335-3230  
Email: [wheyman@lgl.com](mailto:wheyman@lgl.com)

**Semmens, Brice**

Scripps Institution of Oceanography  
University of California, San Diego  
9500 Gilman Drive  
La Jolla, CA, 92039-0202  
Tel: (206) 529-7858  
Email: [Brice.semmens@gmail.com](mailto:Brice.semmens@gmail.com)

**Lindeman, Ken**

Professor  
Florida Institute of Technology  
D.E.I.S.  
150 W. University Boulevard  
Melbourne, Florida 32901  
Tel: (321) 271-7547  
Email: [lindeman@fit.edu](mailto:lindeman@fit.edu)

**US VIRGIN ISLANDS****Nemeth, Richard**

Center for Marine Environmental Studies  
University of the Virgin Islands  
2 John Brewers Bay  
St. Thomas, USVI 00802-9990  
Tel: (340) 693-1393  
Email: [rnemeth@uvi.edu](mailto:rnemeth@uvi.edu)

**FOOD AND AGRICULTURE  
ORGANIZATION  
OF THE UNITED NATIONS (FAO)****Van Anrooy, Raymon**

Fishery and Aquaculture Officer/  
Secretary of WECAFC  
Sub-Regional Office for the Caribbean  
2<sup>nd</sup> Floor, United Nations House  
Marine Gardens, Hastings  
Christ Church BB11000, Barbados  
PO Box 631-C  
Bridgetown, Barbados  
Tel: 1 (246) 426-7110/11 Ext. 249  
Fax: 1 (246) 427-6075  
Email: [Raymon.vanAnrooy@fao.org](mailto:Raymon.vanAnrooy@fao.org)

**CARIBBEAN REGIONAL FISHERIES  
MECHANISM (CRFM)****Mohammed, Elizabeth**

Programme Manager  
Research and Resource Assessment  
Programme  
CRFM Secretariat  
Halifax Street  
Kingstown  
St Vincent and the Grenadines  
Tel: +1 784 457 3474  
Fax: +1 784 457 3475  
Email: [elizabeth.mohammed@crfm.int](mailto:elizabeth.mohammed@crfm.int)

**NATIONAL OCEANIC AND  
ATMOSPHERIC  
ADMINISTRATION/NATIONAL  
MARINE FISHERIES  
SERVICE/SOUTHEAST REGIONAL  
OFFICE**

**Bolden, Stephania**

Chief, Species Conservation Branch  
Protected Resources Division  
NOAA/NMFS/SERO  
263 13th Avenue South  
St. Petersburg, FL 33701  
Tel: (727) 824-5312  
Fax: (727) 824-5309  
Email: [stephania.bolden@noaa.gov](mailto:stephania.bolden@noaa.gov)

**CARIBBEAN FISHERY MANAGEMENT  
COUNCIL**

**Rolon, Miguel A.**

Executive Director  
270 Munoz Rivera Avenue  
Suite 401  
San Juan, Puerto Rico 00918  
Tel: (787) 766-5926  
Fax: (787) 766-6239  
Email: [Miguel\\_rolon\\_cfmc@yahoo.com](mailto:Miguel_rolon_cfmc@yahoo.com)

**Martino, Diana**

Assistant to the Executive Director  
And Executive Secretary  
270 Munoz Rivera Avenue  
Suite 401  
San Juan, Puerto Rico 00918  
Tel: (787) 766-5926  
Fax: (787) 766-6239  
Email: [Diana\\_martino\\_cfmc@yahoo.com](mailto:Diana_martino_cfmc@yahoo.com)

**Irizarry, Maria de los Angeles**

Fiscal Officer  
270 Munoz Rivera Avenue  
Suite 401  
San Juan, Puerto Rico 00918  
Tel: (787) 766-5926  
Fax: (787) 766-6239  
Email: [mairizarry\\_cfmc@yahoo.com](mailto:mairizarry_cfmc@yahoo.com)

**OBSERVERS**

**Dunmire, Leda**

Manager, US Oceans SE  
The Pew Charitable Trusts  
114 Bee Street  
Tavernier, Florida 33070  
Tel: (305) 393-0934  
Email: [LDunmire@pewtrusts.org](mailto:LDunmire@pewtrusts.org)

**Helmick, Emily**

Senior Associate, US Oceans SE  
The Pew Charitable Trusts  
1095 Military Trail #8522  
Jupiter, Florida 33458  
Tel: (561) 670-3397  
Email: [ehelmick@pewtrust.org](mailto:ehelmick@pewtrust.org)

### Declaration of Miami

The Members of the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations:

*Recalling* the Terms of Reference of the joint Working Group, as established by the 14<sup>th</sup> session of the Western Central Atlantic Fishery Commission (Panama City, 6–9 February 2012);

*Noting* with concern the ongoing declines in stocks of many aggregating species and particularly groupers and snappers in the Wider Caribbean Region, the reduced numbers of spawning aggregations, the relatively smaller size of remaining aggregations and the resulting reduced economic and food opportunities;

*Having verified* with scientific methods and based on the information available that the status of Nassau Grouper, Goliath Grouper (and several other species) stocks in the Wider Caribbean Region should be considered “overexploited”, and that some stocks can even be regarded as “depleted”;

*Stressing* the high ecological and biological value of fishes that aggregate to spawn (including groupers and snappers) for the ecosystem and aquatic biodiversity in the region, and that fishing down the food web needs to be avoided;

*Noting* that the biological connectivity of both adults and larvae of some species of snapper and grouper are geographically extensive and hence cross national boundaries;

*Mindful* of the importance of groupers and snapper fisheries for local food security and of the social and economic value of these fisheries for coastal communities in the region;

*Noting* that the actual number of fishers targeting spawning aggregations (as opposed to species that have the aggregating habit) is low. Consequently, while management aimed to conserve spawning aggregations may reduce short-term profits for few fishers, it should enhance long-term sustainable fisheries for many other fishers that fish outside of aggregations. As such spawning aggregations are best considered as capital in a savings account that is guarded to allow provision of annual interest (more fish) to the fishery sector when conducted outside of the aggregation period;

*Concerned* about the increasing demand for grouper and snapper in the international market, which will almost certainly further increase fishing pressure on aggregating species in the region and is strongly implicated in illegal, unregulated and unreported trade;

*Reiterating* the recommendations from the CFMC/WECAFC Regional Workshop on Nassau Grouper (Cartagena, Colombia, October 2008), which called for a regional closed season and establishment of regional collaboration on grouper research and management;

*Recognizing* that in recent years, national level management and conservation efforts targeting spawning aggregations and aggregating species have shown mixed results in the Caribbean, and that introductions of closed seasons and/or site closures in some of the countries in the Wider Caribbean region and in other regions have proven successful in protecting aggregations, particularly when networks of such reserves are implemented by neighbouring countries or regionally. Simultaneous sales controls active fisherfolk involvement can also increase effectiveness;

*Recognizing* that fishers and their organizations have a key role in fisheries management and that there is a need for their active involvement in the research, conservation, and adaptive management of spawning aggregations of fishes;

*Recognizing* that many groupers and snappers spawn in multi-species spawning aggregations and that these aggregations are both extremely valuable and extremely vulnerable to overfishing in the absence of management;

*Further recognizing* the efforts at local, national and regional level to conserve aggregating fish species fisheries in line with the FAO Code of Conduct for Responsible Fisheries, the 1995 UN Fish Stocks Agreement, the precautionary approach and the Ecosystem Approach to Fisheries (EAF), the 2009 FAO Agreement on Port State Measures to Prevent, Deter and Eliminate Illegal, Unreported and Unregulated Fishing, the IUCN Red list, a Recommendation to better protect and manage fish spawning aggregations (adopted by the 4th IUCN World Fisheries Congress, 2004), the work of the Science and Conservation of Fish Aggregations (SCRFA), the Statement of Concern adopted by the second Inter-Tropical Marine Ecosystem Management Symposium in March 2003 on aggregations, and the recommendations of the 4th Scientific Meeting and Technical Advisory Committee (STAC) of the SPAW protocol;

*Convinced* that scientific research on aggregating species and spawning aggregations (e.g. local traditional knowledge, specific stock assessment methods, biology, ecology and life cycle, social and economic value, and reference points for conservation and management of aggregating fish species) should continue to inform fisheries decision makers on *inter alia* suitable input and output measures for fisheries management, appropriate harvesting strategies, consistent with the Ecosystem Approach to Fisheries as well as trade controls and measures to enhance capacities for enforcement and compliance;

*Committed* to individually and collectively taking measures and actions to further improve the management and conservation of fish aggregations and aggregating species in the Wider Caribbean Region;

1. RECOMMEND the endorsement and implementation of the enclosed draft Recommendation to the 6th WECAFC Scientific Advisory Group and 15th session of WECAFC on the establishment of a regional suit of harmonized closed seasons for specific species (starting with Nassau Grouper and adding others as appropriate) in the WECAFC area to protect spawning of overexploited aggregating species (see Annex A);
2. RECOMMEND that the range countries collect and share species specific national and international trade data for Nassau Grouper and other fish species that aggregate to spawn;
3. RECOMMEND that WECAFC members propose the listing of species that aggregate to spawn (in particular Nassau Grouper and Goliath Grouper) under Annex III<sup>1</sup> of the SPAW Protocol , to the Scientific and Technical Advisory Committee (STAC);
4. RECOMMEND that WECAFC, CFMC, CRFM and OSPESCA support the development of a regional plan for the management and conservation of fish species that aggregate to spawn (targeting groupers and snappers), in accordance with the best available scientific evidence to be presented to the 16<sup>th</sup> session of WECAFC in 2016 for review, consideration and regional adoption;
5. RECOMMEND that member countries assess the timing, location and status, of all known transient multi-species spawning aggregations. A list of sites should be prioritized for monitoring, conservation and management based on status and institutional capacity for management at each site;

---

<sup>1</sup> Containing threatened and endangered species of marine and coastal fauna that may be utilized on a sustainable basis, but for which management measures are necessary in collaboration with other range States.

6. FURTHER RECOMMEND that these assessments be conducted along with local fishers who are presently fishing those aggregations, in part to gather their support and in part to offer economic alternatives to fishing those aggregations;
7. SOLICIT the support for, and the direct and immediate implementation by the countries in the Wider Caribbean Region of the above listed recommendations; and
8. REQUESTS THE RESPECTIVE SECRETARIATS to present this declaration and its annexes for discussion and endorsement to the 15th Session of WECAFC, which is scheduled to be held in Trinidad and Tobago in March 2014, as well as to the next session of the Caribbean Fisheries Forum of CRFM and the next ministerial meeting of OSPESCA and communicate with the SPAW Secretariat for appropriate follow-up.

**Annex A: Recommendation to the sixth WECAFC Scientific Advisory Group  
and fifteenth session of WECAFC**

**ON THE ESTABLISHMENT OF A REGIONAL CLOSED SEASON FOR FISHERIES IN  
THE WECAFC AREA TO PROTECT SPAWNING AGGREGATIONS OF GROUPERS AND  
SNAPPERS**

The Western Central Atlantic Fishery Commission (WECAFC),

*RECALLING* that the objective of the Commission is to promote the effective conservation, management and development of the living marine resources within the area of competence of the Commission, in accordance with the FAO Code of Conduct for Responsible Fisheries, and address common problems of fisheries management and development faced by members of the Commission;

*RECALLING* the recommendations of the Regional Workshop on the Management of Nassau Grouper and the agreement of the 13th session of WECAFC (both held in Colombia, October 2008) with these recommendations on the management of Nassau Grouper;

*REAFFIRMING* its commitments, made at the 14th session, through establishing the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations with an aim to provide advice on the management and implementation of regional strategies and regulations to protect spawning aggregations and aggregating species;

*RECOGNIZING* the conclusions of the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations, which convened in Miami, USA, 29-31 October 2013, reviewed the status of some species that aggregate to spawn in the WECAFC Area and discussed a large variety of management and conservation options;

*CONSIDERING* that the Scientific Advisory Group (SAG) at its 6th Session assessed that several fish species that aggregate to spawn (in particular Nassau Grouper and Goliath Grouper) are overexploited, some with a high risk of collapse, and that sustainable management requires that measures aimed at limiting the fishing of spawning aggregations and aggregating species are implemented;

*NOTING* that both the Working Group and the SAG advise of the need to establish a harmonized regional closed season for commercial and recreational fisheries of fish species that aggregate to spawn;

*NOTING* that both the Working Group and the SAG advise the WECAFC members to establish year round no-take marine protected areas at known transient multi-species spawning aggregation sites;

*RECOGNIZING* that various WECAFC members have already established closed seasons for commercial grouper fishing and/or closed areas to protect spawning aggregations;

*NOTING* that many of the spawning aggregations of grouper and snapper in the Caribbean have seriously declined or disappeared in the last two decades and that immediate action is required to stop further reduction in spawning areas and depletion of the stocks;

*CONSIDERING* that current management and conservation efforts targeting spawning aggregations and addressing aggregating species have shown mixed results in the Caribbean, and that application of closed seasons in other regions has proven to be successful when implemented regionally, it is fundamental to limit the fishing effort in areas where adults of important species aggregate to spawn to allow these stocks to reproduce, and, in many cases to recover, thereby allowing for their sustainable exploitation and ongoing contribution to long-term food security and social and economic objectives of the governments in the WECAFC region;

*CONSIDERING* that more scientific information and research is needed with a view to better understanding the relevance of areas on the continental shelf and slope for the protection of spawners in known aggregations and sensitive habitats, as well as to better know the level and spatial distribution of the fishing effort exerted on aggregating species in general;

*PENDING* the delivery of this additional information by the Working Group and the SAG;

*ADOPTS* in conformity with the provision of Article 6 (h) of the Revised Statutes of the WECAFC the RECOMMENDATION that:

1. Members of WECAFC [shall] identify and monitor all known and exploited spawning aggregation areas of groupers and snappers and inform the SAG of any changes in these areas.
2. Members of WECAFC [shall] issue a regional seasonal closure for all commercial and recreational fishing activities of Nassau Grouper (*Epinephelus striatus*) in the identified areas for the period 1 December – 31 March.
3. For the fisheries restricted area referred to in paragraph 1 above, Members [shall] call the attention of the appropriate national and international authorities in order to protect spawning aggregations from the impact of any other human activity jeopardizing the spawning aggregation areas, and ensure enforcement of closed seasons during the reproductive period, if necessary by also implementing sales bans during the closed season.
4. Members of WECAFC [shall] not permit any export of Nassau grouper and Nassau grouper products (e.g. roe, fillets) for the duration of the regional seasonal closure.
5. Member shall conduct research to ascertain the ecological, social and economic impacts of the proposed management measures to inform future management decision-making.
6. Members [shall] prepare national grouper and snapper fisheries management and conservation plans.
7. Members [shall] communicate to the WECAFC Secretariat the measures taken to adhere to the above paragraphs.
8. The WECAFC Secretariat [shall], together with the Members, establish an outreach and communication campaign on the closed areas and regionally agreed closed season.
9. The WECAFC Secretariat [shall], together with the Members, seek to mobilize resources to assist the Members in the implementation of research, monitoring and management measures.
10. Boundaries of the identified spawning areas, spawning seasons and conditions to fish therein, as referred to in previous paragraphs may change on the basis of Working Group and SAG advice coming from additional knowledge.

11. WECAFC, CFMC, and as appropriate CRFM and OSPESCA, [shall] support the development of a regional plan for the management and conservation of fish species that aggregate to spawn (targeting groupers and snappers), in accordance with the best available scientific evidence to be presented to the 16th session of WECAFC in 2016 for review, consideration and regional adoption.
12. Members [shall] assess the timing, location and status, of all known transient multi-species spawning aggregations. A list of aggregations should be prioritized for monitoring, conservation and management based on status and institutional capacity for management at each site.
13. Members [shall] conduct assessments along with local fishers who are presently fishing those aggregations, in part to gather their support and in part to offer economic alternatives to fishing those aggregations.
14. Members [shall] solicit support for direct and immediate implementation by the countries in the Wider Caribbean Region of the above listed recommendations.

**Terms of Reference of the CFMC/WECAFC/OSPESCA/CRFM Working Group  
on Spawning Aggregations**

Convener: Miguel Rolón (CFMC)

The working group will carry out the following tasks:

- Compile and analyze data on spawning aggregations in the member countries and monitor any changes.
- Seek partnerships with other institutions that could provide assistance in the monitoring, evaluation, and recommendations for management for protection and conservation of spawning aggregations.
- Provide advice on the management and implementation of regional strategies and regulations to protect spawning aggregations.
- Report to the appropriate institutions at each session.

The first meeting of the CFMC/WECAFC/OSPESCA/CRFM Working Group on Spawning Aggregations, was held in Miami, United States of America, from 29 to 31 October 2013. The meeting brought together experts working on spawning aggregations of fish from all over Western Central Atlantic region. The Working Group noted with concern the ongoing declines in stocks of many aggregating species and particularly groupers and snappers in the Wider Caribbean Region, the reduced numbers of aggregations and the relatively smaller size of remaining aggregations. The Working Group also verified that the status of Nassau grouper, Goliath grouper (and several other species) stocks in the Wider Caribbean region should be considered “overexploited”, and that some stocks can even be regarded as “depleted”. The meeting issued a “Declaration of Miami”, which included a recommendation to the fifteenth session of WECAFC on the establishment of a regional closed season for Nassau Grouper fisheries in the WECAFC area to protect spawning aggregations of this species.

ISBN 978-92-5-108139-6 ISSN 2070-6987



9 7 8 9 2 5 1 0 8 1 3 9 6

I3585ETri/1/11.13

Propuesta del Reino de los Países Bajos  
para la Lista de Especies de Tiburones y Mantarrayas  
en los Anexos del Protocolo SPAW

# Propuesta del Reino de los Países Bajos para la Lista de Especies de Tiburones y Mantarrayas en los Anexos del Protocolo SPAW

## Resumen Ejecutivo

Gn'T glkq'f g'mu'Rc'f'gu'Dclqu'r tqr qpg'wp'p'Ào gtq'f g'gur gekgu'f g'v'kdw'qpgu'{' 'o cpv'ctc' { cu'r ctc' kpenwuk'p'gp'rc'rkuc'f g'mu'c'pgzqu'KK' { "KKf gn'Rtqv'eqm'URCY 0'"

### Para el Anexo II:

#### 1. "Pez Sierra Peine (*Pristis pectinata*)"

Ecn'k'ecf q'r ctc'kpenwuk'p'gp'rc'rkuc'f gn'c'pgzq'4'f gn'Rtqv'eqm'URCY . 'ugi Àp<

- / Et'kgtkq'3. 'f g'gdk'q'c'g'xkf g'p'ek'f g't'gf we'ek'p'{' 'Itci o g'p'v'ek'p'f g'rc'rkuc'q'dr'ek'p'0'
- / Et'kgtkq'6. 'f cf q's w'g'rc'gur gek'g'gu'v' rkuc'f c'eqo q'et'f'keco g'p'v'g'gp'rc'rkuc'q'gr'ki tq'f g'g'z'v'ek'p' r qt'rc'WEP '\*Y kg' { "et al., 4235+0'
- / Et'kgtkq'7. 'f g'gdk'q'cn'eqo g'tek'q'gp' uk'ttcu' t'qu't'crgu' { "uw'kpenwuk'p'gp'rc'rkuc'f gn' Cr<sup>2</sup> pf leg'Kf g'EKGUO'
- / Et'kgtkq'8. 'f g'gdk'q's w'g'rc'eqqr g't'ek'p'gp'v'g'rc'f'gu' gu'p'ge'guct'k' r ctc' r tq'v'gi gt'rc'u' gur gek'gu'f w'cp'v'g'uw'u'o ki t'ek'q'p'gu'g'uc'ek'q'p'crgu'0'

Gn'r gl' "uk'ttc' r g'k'p'g' \*Pristis pectinata+ j c'uk'f q'ecuk'v'q'v'm g'p'v'g'z'v'k' r cf q'f g'i t'cp'f gu'g' t'gcu' f g'uw'cp'v'ki w'c'g'c'f g'f k'ut'k'w'ek'p'gp'gn'Qe<sup>2</sup> cpq' C'v'f p'v'ek'q'q'ek'f g'p'v'cn'f g'gdk'q'c'rc'rkuc'g'uec' \*t'gf gu'f g' c'tt'c'ut'g' { "t'gf gu'f g' d'cl'w'c'+ { "rc'o q'f k'k'ek'ek'p'f g'uw'j<sup>a</sup> d'k'c'v'0'Nqu' t'gi k'ut'qu' p'gi c'v'x'qu'f g'rc'u'g'uw'f k'qu'ek'p'v'f'kequ'rc'u'q'd'ug't'x'ek'q'p'gu'c'p'g'ef »v'ek'cu'f g'rc'u'r g'uec'f q't'gu' { "rc'u'f c'v'qu' u'q'd't'g' f g'ugo d'c't's w'gu' r t'q'eg'f g'p'v'gu' f gn' u'g'eq't' r g'us w'g't'q' u'q'd't'g' uw'g'c'f g' f k'ut'k'w'ek'p'j k'w'v't'k'ec'k'p'h'g't'g'p'w'c't'gf we'ek'p'f g'x'; 7' "f g'rc'rkuc'q'dr'ek'p'gp'wp'rc'rkuc'q'f g'v'g'u'i g'p'g't'ek'q'p'gu'f g'g'f g'3; 84'j c'v'c'rc'h'g'ej c-0'Ncu'gur gek'gu'f g'rc'rkuc'k'k'c'Rt'k'w'f c'g'rc'g'gu' uk'ttc'+u'q'p'rc'u'g'rc'uo q'd't'c'p's v'k'qu'o<sup>a</sup> u'co g'p'c'f q'u'c'p'k'x'gn'o w'p'f k'rc'0'

#### 2. "Tiburón ballena (*Rhincodon typus*)"

Ecn'k'ecf q'r ctc'kpenwuk'p'gp'rc'rkuc'f gn'c'pgzq'4'f gn'Rtqv'eqm'URCY . 'ugi Àp<

- / Et'kgtkq'3. 'f g'gdk'q'c't'gf we'ek'p'f g'rc'rkuc'q'dr'ek'p'0'
- / Et'kgtkq'6. 'f cf q's w'g'rc'WEP 'rc'u'j c'rkuc'f q'eqo q'co g'p'c'f q'u'c'p'k'x'gn'o w'p'f k'rc' { "uw' r q'dr'ek'p'gp'gn' C'v'f p'v'ek'q'eqo q'x'w'p'g't'cd'rg'0'
- / Et'kgtkq'7. 'f cf q's w'g'rc'gur gek'g'gu'v' rkuc'f c'gp'gn' Cr<sup>2</sup> pf leg'Kf g'EKGUO'
- / Et'kgtkq'8. 'f cf q's w'g'rc'ek'k'ec' r ctc'eqqr g't'ek'p't'gi k'q'p'cn'f g'gdk'q'c'rc'rkuc'f g'o ki t'ek'p' p'q't'v'g'uw' s'w'g'ug'z'v'ep'f g'c'v'c'x<sup>2</sup> u'f g'rc'u'l'w'k'uf l'ek'q'p'gu'f g'o w'ej q'u'r c'f'gu'0'

Gn'v'kdw'»p' d'cm'g'p'c' gu'w'p'c' gur gek'g'c'nc'o g'p'v'g'o ki t'c'v'q't'k' { "j c'd'k'c'gp'rc'u'o c't'gu'v't'qr k'c'rgu' { "v'go r n'ef qu'0'Ncu'ec'r w't'cu'j' cp'f k'uo k'p'w'f q' { "cn'r c't'ge'g't'ug'j' cp'ci q'v'cf q'rc'u'r q'dr'ek'q'p'gu'gp'x'ct'k'qu' r c'f'gu'f g'gdk'q'c'rc'rkuc'g'uec'eq'p'ctr »p'gp'i t'cp'f gu'eq'p'eg'p'v'ek'q'p'gu'f g'g'uw'gur gek'g'gp'q'to g'f g' o q'x'lo k'ep'v'qu' r'ep'v'qu' { "eq'p'f w'ec'x'w'p'g't'cd'rg'0'Gn'v'kdw'»p' d'cm'g'p'c'j c'o q'v'c'f q'w'p'f g'ue'g'p'qu' i g'p'g't'cn'f gn'85' "gp'gn'R'ce'f'keq'k'p'f k'q'gp'rc'u'À'ko qu'97' c'o qu' \*t'gu'i g'p'g't'ek'q'p'gu'0'Gp'gn' C'v'f p'v'ek'q'rc'rkuc'f g'g'g'g'rc'rkuc'q'dr'ek'p'ug'eq'p'k'f g't'c'o g'p'q't'c'x'52' . { "ug'k'p'h'g't'g's w'g'rc' t'gf we'ek'p'i m'd'cn'gu'x'72' 0'

#### 3. "Tiburón oceánico punta blanca (*Carcharhinus longimanus*)"

Ecriñkef q'r ctc'kpenwuk»p"gp"rc"rkuc" f gn'Crpgzq"4" f gn'Rtqvceqmq"URCY . "ugi Àp<

- / Etkgtkq"3." {c"s wg"uw"r qdrcek»p"j c" f kuo kpwkf q"pqvdrgo gpvg"gp"rc"tgi k»p"pqtqguv" { "egpvtq"qeekf gpvcn'f gn'Cr vñ pveq0
- / Etkgtkq"6." {c"s wg"rc"KWE P"m"j c"rkucf q"eqo q"xwpgtcdrg" { "et"keco gpvg"gp"r grki tq" f g" gz vpeke»p"gp"rc"tgi k»p"pqtqguv" { "egpvtq"qeekf gpvcn'f gn'Cr vñ pveq" f gdlf q" c"uw"gpqto g" tgf week»p0
- / Etkgtkq"7." {c"s wg"hi wtc"gp"gn'Cr<sup>2</sup> pf leg"KKf g'E K/GU0
- / Etkgtkq"8." {c"s wg"guv" "enukñkef q'r qt"rc"E qpxgpeke»p" f g"rcu" P cekqpgu" Wpkf cu"uqdtg" gn' F gtgej q" f gn'O ct"eqo q" wpc" gur gekg" cnco gpvg" o ki tcvqtk0"

Gn'kdw»p"qeg<sup>a</sup> pleq"r wpc"drpecc"gu" wpc" gur gekg" cnco gpvg" o ki tcvqtk. "ew" c" r qdrcek»p" ug" j c" tgf wekf q" o<sup>a</sup> u" f gn'; 2' " f wcpvg" w" r gt" qf q" s wg" cdctec" tgu" i gpgtcekqpgu0 f gdlf q" c" uw" pcwrcgl c" o ki tcvqtk. " rc" eqpugt xcek»p" gñkec| " f g" guvc" gur gekg" tgs wgtk<sup>a</sup> " eqqr gtcek»p" kpvgtcekqpcn0'

**Para el Anexo III**

**1." Mantarrayas (*Manta birostris*, *Manta alfredi* y *Manta cf. birostris*)"**

Ecriñkef c'r ctc'kpenwuk»p"gp"rc"rkuc" f gn'Crpgzq"KKf gn'Rtqvceqmq"URCY . "ugi Àp<

- / Etkgtkq"3." f gdlf q" c" tgf week»p" { "Itci o gpvcek»p" f g"rcu" r qdrcekqpgu0
- / Etkgtkq"6." r qts wg" vqf cu" guvcu" gur gekgu" guv" p" rkucf cu" eqo q" xwpgtcdrgu" r qt"rc" WEP 0
- / Etkgtkq"7." f cf q" s wg" guv" "rkucf c" gp" gn'Cr<sup>2</sup> pf leg"KKf g'E K/GU0
- / Etkgtkq"8." f gdlf q" s wg" ug" j c" kpenwuk q" gp" gn'Cr<sup>2</sup> pf leg"K { "KKf g"rc" EO U" { "gp" gn'Crpgzq" K f gn'O go qtc pf q" f g" Gpvpgf ko kpvq" uqdtg" kdw"tqpgu" { "tgs wgtgp" eqqr gtcek»p" r ctc" uw" eqpugt xcek»p0'

Nc"o cpvcttc { c" i ki cpvg" \*o cpvc" dktquwku" + { "rc" o cpvcttc { c" f g" cttgek" \*o cpvc" crhtgf k: "eqp" wpc" ur wguvc" vtegtc" gur gekg" gpf<sup>2</sup> o lec" gp"rc" tgi k»p" f gn'Ectkdg. "O cpvc" eh0 dktquwku. "uqp" gn' i<sup>2</sup> pgtq" f g" o cpvcttc { cu" o<sup>a</sup> u" i tcpf g. "m" ewcn' j ceg" s wg" uw" xlf c" ugc" gur gekm gpvg" eqpugt xcf qtc" { " xwpgtcdrg" cn' ci qvco kpvq0 Rqt" qtc" r ctvg. "c" r guct" f g" gxlk gpekc" f g" rti cu" o ki tcekqpgu. "rcu" r qdrcekqpgu" tgi kpcrcgu" r ctgegp" ugt" r gs wgo cu. "gucuco gpvg" f kwtkdwkf cu" { "Itci o gpvcf cu. "m" ewcn' uki pñkec" s wg" gu" r req" r tqdcdrg" s wg" rcu" tgf weekqpgu" kf gpñkef cu" ugcp" o kki cf cu" r qt"rc" kpo ki tcek»p0'

**2." Tiburón martillo (*Sphyrna lewini*, *Sphyrna mokarran* y *Sphyrna zygaena*)"**

Ecriñkef q'r ctc'kpenwuk»p"gp"gn'rkucf q" f gn'Crpgzq"KKf g"URCY . "ugi Àp<

- / Etkgtkq"3." {c"s wg" j c { "cdwvf cpvg" gxlk gpekc" f g" rc" tgf week»p" f g" r qdrcekqpgu" gp" gn' Cr vñ pveq" qeekf gpvcn" { c"s wg" uqp" xwpgtcdrgu" c" rc" uqdtggzr mqveke»p" { "vkgpgp" w" dclq" r qvpekn' f g" tgewr gtcek»p" f gdlf q" c" wpc" dclc" vcuc" f g" etgelo kpvq" kvt" pugec" { " tgr tqf week»p" ngpv. "cu" eqo q" wpc" etgekpvvg" ecr wtc" f kki kf c" g" lpekf gpvcn' gp"rc" tgi k»p" pqtqguv" { "egpvtq"qeekf gpvcn'f gn'Cr vñ pveq0
- Etkgtkq"40" Ug" co gtkec" w" gphqs wg" ecwgrnuq" { c"s wg" rc" ecpkcf "gzcevc" f g" rc" r tguke»p" r gus wgtc" { "rc" eqttgur qpf kpvvg" vcuc" f g" o qtcvckf cf "gu" quewtc0' P q" gu" h' eki" j cegt" kphgtgpekc" gur geñkecu" c" gur gekgu" f gdlf q" c" rcu" f kñewncf gu" cuqekf cu" eqp" rc" kpeccelcf cf "r ctc" f kwpk vkt" gpv" g" U. *zygaena*, *S. lewini*, y *S. mokarran*. { "gn' f kñekf guvcf q" f g" eqpugt xcek»p" f gn' kdw»p" o ctvkm0'

- Etkgtkq "6. "f cf q's wg"rc" WĒEP "rkuc"gn'gucf q'f g'eqpugt xcek»p'f gn'vdtw»p"o ct vkmq "eqo q" gp"r grki tq'r ctc" S. mokarran. "S. lewini" \*co dqu" c" guecnc" o wpf kni { "rc" uwdr qdrcck»p" gp" rc' tgi k»p' pqtqguw" { "egpvtq" qeekf gpvcnf gn' C vñ pñeq-+ { "xwpgtcdrg'r ctc" S. zygaena' }
- Etkgtkq "7. " { c' s wg" rc" hco krc "f gn' vdtw»p" o ct vkmq "ug" gpewgpvtc" gp" rc" rkuc "f gn' Cr<sup>2</sup> pf leg" Kk' g' EKVGU' }
- Etkgtkq "8. " { c' s wg" rqu" vdtwqpgu" o ct vkmq " hki wtcp" rkucf qu" gp" gn' Cpgzq" K f g" rc" Eqpxgpek»p' f g" rcu" P cekqpgu" Wpk cu" uqdtg" gn' F gtgej q' f gn' O ct " \*EP WF O + " { . " r qt" m" wpvq. ' tgs wkg' gp" eqqr gtcek»p' k»vtpcekqpcnr ctc" uw' eqpugt xcek»p' }

Gn' vdtw»p" o ct vkmq " gu" wpc" gur gekg" f g" vdtw»p" ekewo i nqdcn' qtkwvf q' f g" rcu" ci wcu" equvgtcu" e<sup>a</sup> rkf cu. " wgo r rcf cu" { " vqr kcrngu' S. lewini" vkgpg" gn' r qvpekn' f g" tgewr gtcek»p" o<sup>a</sup> u' dclq" gp" eqo r ctcek»p" eqp" qvtcu" gur gekgu" f g" vdtwqpgu' Ncu" vcucu" f g" etgelo kgpvq' f g" rc" r qdrcck»p" f g'vgo kpcf cu" r ctc" r qdrcckqpgu" gp" gn' qe<sup>2</sup> cpq" Rceñkq { " C vñ pñeq" uqp" dclcu' Cp<sup>a</sup> rkuku" f g" wpf gpek' f g" cdwvf cpek' f g" rqu" f cvqu" f g" vcuc' f g" ecr wtc" j cp" tgr qtvcf q' i tcpf gu" o gto cu" gp" cdwvf cpek' gp" w" tpci q' f g" 82/ ; ; ' " gp" c<sup>o</sup> qu' tgekgpvu' Wpc" gxcncck»p' f g" r qdrcckqpgu" eqp" kphqto cek»p' f g" ecr wtc. " wpf gpekcu" f g" cdwvf cpek' { " dkqmi " c" gur geñkq " c" S. lewini" gp" gn' C vñ pñeq" pqtqekf gpvcn' kpf kcp" wpc" tgf week»p' f gn' : 5' " gpvtg" 3 ; : 3' { " 42270' Ncu" vcucu" f g" ecr wtc" guvcf ctk' cf cu' f g" rc" r gus wgtñc' f g' r crpi tgr gn' i keq' f g' Gucf qu' Wpk qu" o wgvcp' wpc" tgf week»p' f gn' : ; ' " gp" Sphyrna spp' 0gpvtg" 3 ; : 8' { " 4222. " { " wpc" tgf week»p' f gn' 98' " gpvtg" 3 ; ; 4' { " 42270' Ncu" crvcu" f g' vdtw»p" o ct vkmq " uqp" o w' " xcmqcf cu" { " uqp" w" etgekgpv" drcpeq" gp" cri wpcu" <sup>a</sup> tgcu" gp" tgr wgvu" c" rc" etgekgpv" f go cpf c' f g' crvcu" f g' vdtw»p' 0Ug" gpeqvt»" s wg" rcu" gur gekgu" f g' vdtw»p" o ct vkmq " S. zygaena" { " S. lewini" tgr tguvcp' r qt" m" o gpqu" gpvtg" gn' 6' " { " 7' " f g" rcu" crvcu" uwdcucf cu" gp" J qpi " Mqpi . " gn' egpvtq" eqo gtekn' o<sup>a</sup> u' i tcpf g' f g' crvcu" f g' vdtw»p' 0Ncu" crvcu" gp" gn' o gtecf q" UCT" f g' J qpi " Mqpi " ug" r wgf gp" gxcnwt" i gp<sup>2</sup> vkeco gpv" { " ug" j c" f go qvtcf q' s wgtk' kpcp' f g' rcu' ewgpecu" gp" gn' C vñ pñeq" qeekf gpvcn' }

**Propuesta**

**Teniendo en cuenta:"**

Nc' pgegukf cf " f g' r tqvgi gt" c" rqu" vdtwqpgu" { " o cpvttc { cu. " ugi Æp" m" gxf gpek' uw' xwpgtcdkrf cf " gzegeklqpcn' c" rc" uqdtgr guc" { " ci qco kgpvq" c" rti q" r n| q. " f gdlq" c" uw' etgelo kgpvq" ngpvq. " o cf wtgl " vtfñc" { " r gs wgo cu" eco cf cu. " f cf q' s wg" rcu" j go dtcu" pq" ug" tgr tqf wegp" j cuvc" s wg" ewo r ngp" f lgl " c<sup>o</sup> qu" { " f cp" c" nñ " c" w" p' p' Ao gtq" r gs wgo q' f g' etñcu" f gur w<sup>2</sup> u" f g" w" rti q" go dtcl q' 0Nqu" vgo r qu" f g' i gpgtcek»p' f g' rcu" gur gekgu" f g' vdtwqpgu" r wgf gp" ugt" f g' j cuvc" 72' c<sup>o</sup> qu. " m' ewcn' m' eqmec" gp" rc' ecvgi qtñc' f g' tgr tqf week»p" o<sup>a</sup> u' dclc' f g' rc' HCQ=

S wg" rcu" r qdrcckqpgu" f g' vdtw»p" ug" j cp" xkrq" uqo gvk' cu" c" wpc" r guc" gzegekxc" { " j cp" ulf q" ugxgtco gpv" ci qvcf cu" gp" vqf q" gn' qe<sup>2</sup> cpq" C vñ pñeq. " { " f cf q' s wg" rcu" j go dtcu" tgr tqf wevqtcu" uqp" o<sup>a</sup> u' i tcpf gu" { " o<sup>a</sup> u' r tgekc' cu. " uwrgp" ugt" gn' r tkpek' cn' drcpeq' f g' rcu' r gus wgtñcu="

S wg" rqu" vdtwqpgu" ug" j cp" eqpxgtvk' q" gp" wpc" o gtecpñc" ko r qtvcpv" { " o w' " cr tgekc' c" r qt" uw' ectpg. " r tkpek' cm' gpv" wu" crvcu. " gn' lpi tgf kgpv" en' xg' r ctc" rc" uqr c' f g' crvc' f g' vdtw»p. " s wg" gu" w" o cplct" gp" Cuk' 0Nc' i tcp' f go cpf c' f g' r tqf wevqu" f g' vdtw»p" { " rc" kpcvc' xwpgtcdkrf cf " f g' rqu" vdtwqpgu" c' rc' r tgu' p' f g' rc' r guc' j c' m' xcf q' c' wpc' tgf week»p' ci w' c' gp" gn' p' Ao gtq' f g' vdtwqpgu" gp" vqf q' gn' o wpf q. " tgf wekpf q' cn' wpcu' r qdrcckqpgu" gp" o<sup>a</sup> u' f gn' ; 2' 0"

"

S w g " r " o c { q t " p " f g " r u " g u r g e k u " f g " v d w t q p g u " { " o c p v t t c { c u " u q p " o k i t e v t k u " q " v g p g p " f g u r r | c o k g p v q u " t c p u h t q p v g t k q u " { . " r q t " n j " v c p v q . " r " r t q v g e e k p " { " i g u v k p " e q q r g t c v k c " g p v t g " n j u " r c " p u g u " u q p " p g e g u c t k u " r c t c " o c p g l c t " g u w c u " g u r g e k u " =

S w g " g n " T g k p q " f g " n j u " R c " p u g u " D c l q u " g u v c d n g e k " g n " U c p w c t k q " } c t c t k " r c t c " v d w t q p g u " { " o c o " h g t q u " o c t k p q u " g p " r u " c i w c u " f g " D q p c k t g " { " U c d c " =

**Recordando:"**

G n ' c t v p e w n j ' 3 3 ' 6 0 \* c + f g n ' R t q v e q m j =

**Tomando nota de:"**

N c " l p h t o c e k p " t g s w g t k f c " g p " g n ' c t v p e w n j " 3 ; 6 =

N q u " e t k v g t k q u " { " g n " r t q e g f k o k g p v q " f g " r k u c f q " c f q r v c f q u " r q t " n " Q e v c x c " E q p h g t g p e k " f g " r u " R c t v g u " \* E Q R " : + g p " E c t v i g p c . " E q m o d k . " g p " f l e k g o d t g " f g " 4 2 3 6 0

The Kingdom of the Netherlands proposes the following species of Sharks and Rays for listing on the SPAW Annexes as indicated below:

## For Annex II

### 1. Smalltooth Sawfish (*Pristis pectinata*)

#### Overview

The Smalltooth Sawfish qualifies for listing on the SPAW Protocol Annex 2, especially according to: criteria 1 due to evidence of decline and population fragmentation, criterion 4 as the species is listed as critically endangered by IUCN (Wiley *et al.*, 2013), criterion 5 due to the trade in the rostral saws and CITES listing on Appendix I and criterion 6 as cooperation between countries is needed to protect the species during their seasonal migrations

The Smalltooth Sawfish (*Pristis pectinata*) has been wholly or nearly extirpated from large areas of its former range in the Western Atlantic Ocean by fishing (trawl and inshore netting) and habitat modification. Negative records from scientific surveys, anecdotal fisher observations, and fish landings data over its historic range infer a population reduction  $\geq 95\%$  over a period of three generations (i.e., 1962 to present). While the population found in the United States appears to have stabilized with some evidence of increase, information from other areas is lacking. The remaining populations are inferred to be small and fragmented based on the lack of records. The species can only be reliably encountered in the Bahamas (where suitable habitat is available) and the United States (Georgia south to Louisiana). It is rare but present in Honduras, Belize, and Cuba. While historic threats to Smalltooth Sawfish have been reduced in places like the United States, threats still exist today from areas where Sawfish are unprotected and habitat modification and inshore netting still occurs. All species of the family Pristidae (the Sawfishes) are the most threatened elasmobranchs globally (Dulvy *et al.*, 2014).

#### Species information

##### a. Scientific and common names of the species

- 1.1. Class: Chondrichthyes, subclass Elasmobranchii
- 1.2. Order: Rajiformes
- 1.3. Family: Pristidae
- 1.4. Species: *Pristis pectinata* (Latham, 1974)
- 1.5. Scientific synonyms: *Pristis serra* (Bloch and Schneider 1801), *Pristis granulosa* (Bloch and Schneider 1801), *Pristis acutirostris* (Duméril 1865), *Pristis leptodon* (Duméril 1865), *Pristis megalodon* (Duméril 1865), *Pristis occa* (Duméril 1865), *Pristis woermanni* (Fischer 1884), *Pristis evermanni* (Fischer 1884), and *Pristis anandalei* (Chaudhuri 1908).
- 1.6. Common names:
  - English: Smalltooth Sawfish, Wide Sawfish
  - Spanish: Pejepeine, Pez Sierra, Espadachin, Espadon, Pejes sierra, Pez espada, Pez rastrillo
  - French: Poisson-scie, Requin-scie

## **b. Estimated population of species and its geographic ranges**

Smalltooth Sawfish were widely distributed throughout the tropical and subtropical marine and estuarine waters of the Western Atlantic Ocean. They were found from Uruguay through the Caribbean and Central America, the Gulf of Mexico, and the Atlantic coast of the United States (Faria *et al.*, 2013). However, this range has contracted and the Smalltooth Sawfish has been likely extirpated from large areas of its former range. The species is currently known to occur in the southeastern United States, Bahamas, Cuba, Honduras, and Belize. Reports of Smalltooth Sawfish outside of the Atlantic Ocean are likely misidentifications of other Sawfish species (Faria *et al.*, 2013). In the United States, the Smalltooth Sawfish population appears to have declined dramatically during the middle and later parts of the 20th century (Simpfendorfer, 2002). Based on the contraction of the range and declines in landings, it is likely that the population in the United States at the end of the 20th century was less than 5% of its size at the time of European settlement (Simpfendorfer, 2002). Based on genetic sampling, estimates of the current effective population size range of the United States population of Smalltooth Sawfish were from 269.6–504.9 individuals (95% Confidence Interval 139.3–1,515; Chapman *et al.* 2011). Outside United States waters, no data on population size or trends in abundance exist and the only information on trends in the population can be inferred from capture records. While early records of this species include most countries throughout Central and South America, records and reports indicate the Smalltooth Sawfish can only be reliably encountered in the Bahamas where suitable habitat is available, and in Honduras, Belize, and Cuba (R. Graham pers. comm. 2012).

Using data from reported encounters from 1998 to 2008, Wiley and Simpfendorfer (2010) evaluated Smalltooth Sawfish habitat use patterns in the US. There was an inverse relationship between Sawfish size and extent of northern distribution, with animals less than 200 cm having a wider latitudinal distribution and occurring farthest north, and animals greater than 200 cm reported mostly in southern Florida (Wiley and Simpfendorfer, 2010). Most encounters occurred in estuarine and nearshore habitats, and their locations were not randomly distributed, having a positive association with inshore mangrove and seagrass habitats. While Sawfish were reported in depths to 73 m, there was a significant relationship between size and depth, with smaller animals occurring in shallower waters (Wiley and Simpfendorfer, 2010).

Data from acoustic telemetry and tag-recapture information indicates Smalltooth Sawfish (less than 100 cm) had the smallest home ranges, a low linearity of movement, and a preference for very shallow mud banks (Simpfendorfer *et al.*, 2010). Juveniles greater than 100 cm demonstrated larger home ranges, preference for shallow mud/sand banks, and remained close to mangrove shorelines. Tide was found to be the main factor influencing movement on short time scales. Sawfish <150 cm. STL spend the majority of their time in water <0.5 m deep, while larger juveniles spend most of their time in water 0.5–1.0 m. deep. Juveniles >130 cm had high levels of site fidelity for specific nursery areas for periods up to almost 3 months, but the smaller juveniles had relatively short site fidelity to specific locations (Simpfendorfer *et al.*, 2010). For adult Sawfish, unpublished data from pop-off archival satellite transmitting (PAT) tags indicate Smalltooth Sawfish spend the majority of their time in shallow waters (<10 m deep) and prefer temperatures between 22°C and 28°C (J.K. Carlson, unpublished data). The maximum-recorded depth for Smalltooth Sawfish is 88 m.

The population of Smalltooth Sawfish may have stabilized in the United States. Carlson and Osborne (2012) reported the relative abundance of Sawfish increased at an average rate of about 3–5% per year since 1989 based on of voluntary dockside interviews of sports fishers. Despite a low population size in the United States, the Smalltooth Sawfish population will probably retain >90% of its current genetic diversity over the next century (Chapman et al. 2011).

Faria *et al.* (2013) state that both morphology and genetics support the current specific status of the Smalltooth Sawfish (*Pristis pectinata*) and proposed a modification of the distribution of the species to an Atlantic only range. No geographical structure of Smalltooth Sawfish populations has been detected, but the Western and Eastern Atlantic populations of the Smalltooth Sawfish represent separate units for conservation purposes.

A recent paper by Dulvy *et al.* (2014) shows that all seven species of the family Pristidae are the most threatened elasmobranchs in the world, as a result of their high exposure to coastal shallow-water fisheries and their large body size.

### **c. Status of legal protection, with reference to relevant national legislation or regulation**

#### **International**

##### ***Convention on the International Trade of endangered Species (CITES)***

All Sawfish species are listed under Appendix I of CITES. This means that CITES recognizes that the species is threatened with extinction and that all international commercial trade in wild specimens is prohibited. See [www.cites.org](http://www.cites.org)

##### **Convention for the protection of Migratory Species (CMS) – Memorandum of Understanding (MOU) on the Conservation of Migratory Sharks**

All sawfish species are listed on Annex 1 of the Memorandum of Understanding (MOU) on the Conservation of Migratory Species. The Shark MOU is the first global instrument for the conservation of migratory species of sharks. Signatories to the MOU commit to the objective of achieving and maintaining a favorable conservation status for migratory sharks based on the best available scientific information, in particular the sharks listed on Annex 1 of the MOU, recognizing that successful shark conservation and management require the fullest possible cooperation among governments, intergovernmental organizations, nongovernmental organizations, and all stakeholders

##### ***IPOA Sharks***

Since the 1990s there are several shark protection plans, both internationally at intergovernmental and non-governmental level, as well as at national level by several nations in the Wider Caribbean region. Within the framework of the Code of Conduct for Responsible Fisheries, the FAO (Food and Agriculture Organization) developed the International Plan of Action for the Conservation and Management of Sharks (IPOA Sharks) in 1999. The objective of IPOA Sharks is to ensure the conservation and management of sharks and their long-term sustainable use. IPOA Sharks is voluntary and intends to give

states guidelines on how to establish a National Plan of Action (NPOA) through guiding principles and procedures for implementation.

## **National Protection**

### **USA**

The United States listed *Pristis pectinata* on the US Endangered Species Act in 2003, following earlier protection in the State waters of Florida and Louisiana and protection under the USA Atlantic & Gulf Coasts Fishery Management Plan since 1997. This remnant population in the Gulf of Mexico is considered to have survived because of the benefits of large marine and coastal protected areas, including the establishment of the Everglades National Park in 1947, and as a result of a number of conservation measures during the 1990s, including species protection in Florida and Louisiana and a ban on all forms of entangling fishing nets in Florida State waters (Simpfendorfer 2002). A Recovery Plan has been adopted for this species (NMFS, 2006). The decline in this population may have ceased as a result of these measures.

Outside United States waters, Nicaragua imposed a permanent ban on targeted Sawfish fishing in Lake Nicaragua. In Brazil, the Smalltooth Sawfish is protected by the Ministry of Environment and in Mexico, the take of all Sawfishes is banned.

### **Honduras**

In June 2011 Honduras created the first shark sanctuary in America and declared all its marine waters in both the Pacific and Caribbean as a permanent shark sanctuary. This had been preceded in 2010 by a shark fishing moratorium and created the first shark sanctuary of the Americas amounting to about 240,000 km<sup>2</sup> of national waters, most of which lie along the 700 km-long Caribbean coast of the nation.

### **Bahamas**

The Bahamas have had a longline fishing ban since 1993 and consequently there has been no commercial shark fishing activity. This longline ban has effectively made the whole archipelago of the Bahamas a shark “no-take” zone. The last export of shark from the Bahamas was a lot of 2 metric tons in 2004. In July 2011 the Bahamas went a step further and legally banned all shark fishing. That law firmly turns all 630,000 sq km of Bahamian waters into a shark sanctuary<sup>17</sup>. The fines for shark fishing were raised from 3000 to 5000 USD per incident.

### **Venezuela**

Towards implementing its Plan de Acción Nacional (PAN) de conservación for sharks, in June 2012 Venezuela joined the rest of the Americas in outlawing the finning of sharks in its waters and established a 3,730 km<sup>2</sup> shark sanctuary surrounding the touristic archipelago of Los Roques. Recent research (e.g. Tavares 2005, 2008 2009) had demonstrated the importance of the shallow waters of Los Roques as a shark nursery area.

The Dominican Republic has, together with Belize and six other Central American countries, united under the name SICA (Central American Integration System), signed an agreement to prohibit shark finning. This ban is also applicable to fishing vessels in international waters under the flag of SICA member states. This arrangement OSP-05-11 entered into force in 1 January 2012.

### **Kingdom of the Netherlands**

— **St. Maarten**

On the 12th of October 2011 the government of St. Maarten issued a temporary moratorium on shark fishing. The shark fishing moratorium prohibits the take and landing of sharks and requires immediate release of incidentally caught sharks, under penalty of a maximum of 500,000 Antillean Guilders or 3 months in prison.

— **Caribbean Netherlands**

In 2015, the Dutch government designated the Yarari sanctuary for sharks and marine mammals in the Economic Exclusive Zones of Saba and Bonaire, declaring that provisions will be considered and implemented as necessary to regulate activities that may have a negative impact on sharks.

— **Bonaire**

In 2008 the island of Bonaire passed a nature ordinance providing full protection for a list of species of plants and animals. This list includes all sharks and rays

#### **d. Ecological interactions with other species and specific habitat requirements**

Little is published about the ecological role and trophic ecology of Sawfish. It is known that the sawfish is a high order predator in riverine environments, and while consuming a wide range of prey types, it predominantly feeds on bony fishes (Thorburn, 2006). Adults are likely to be important predators of teleost fish and peneaid prawns in coastal marine ecosystems. Bigelow and Schroeder (1953) reported that Sawfish in general predominantly prey on small schooling fish, such as mullets and clupeids. Bigelow and Schroeder also reported that they feed to some extent on crustaceans and other bottom dwelling inhabitants. They use their rostrum to stun schooling fishes with sideswipes of the snout.

#### **e. Management and recovery plans for the species**

Since the U.S. Smalltooth Sawfish population was listed as endangered in 2003, the commercial bycatch and recreational fisheries, as well as habitat loss have greatly decreased (some of the actions already existed before 2003). There has been a ban on inshore fishing nets in Florida waters for more than a decade and there are prohibitions and fines against intentionally capturing, harming or harassing Sawfish)<sup>1</sup>.

#### **f. Research programs and available scientific and technical publications relevant to the species**

Currently the major aim of Sawfish research in the U.S. (Florida) is monitoring the Sawfish population to determine if the population is rebounding or at the very least stabilizing, in order to evaluate the effectiveness of protective measures. This monitoring information will provide important data about the ecology, reproduction and life history of the species, which will enable more effective conservation efforts to protect the Smalltooth Sawfish. It is important that this monitoring program continues well into the future as the recuperation of this species will take some time due to its life history characteristics (<https://www.flmnh.ufl.edu/fish/Sawfish/conservation/about/>).

---

<sup>1</sup> <https://www.flmnh.ufl.edu/fish/Sawfish/conservation/about/>

In July of 2016, the annual Joint Meeting of Herpetologists and Ichthyologists will be about the biology and ecology of sawfishes, possibly facilitating new research opportunities and improving coordination of current research efforts.

**g. Threats to the species, its habitats and associated ecosystems, especially threats which originate outside the jurisdiction of the Party**

The principal threat to the Sawfishes is from target and utilized bycatch (or byproduct) fisheries. Their long tooth-studded saw makes them extraordinarily vulnerable to entanglement in any sort of net gear. Bycatch mortality in net fisheries was the major reason for the decline of *Pristis pectinata* in the United States (Seitz and Poulakis, 2006). There have been some large-scale target Sawfish fisheries: in Lake Nicaragua and possibly in Brazil from 1960s to 1980s (bycatch is still landed in this range State). Populations are now so depleted, however, that commercial targeting of Sawfish stocks is no longer economic. Most Sawfishes have been and still are killed in broad-spectrum commercial and artisanal fisheries, particularly set net and trawl fisheries that target a very wide range of fishes and invertebrates. Sawfishes are retained in these fisheries, just as they were in former target fisheries, because of the very high value of their products (particularly meat, fins and rostral saws, also liver oil and skin). They are also targeted or bycatch and retained opportunistically for the same reasons. Sawfish fins occur but are now extremely rare in the Asian dried shark fin trade and may have once had their own trade name given their value (D. Chapman pers obs). Trophy angling for very large specimens has been reported (Simpfendorfer, 2005; McClenachan, 2009). The Nicaraguan government imposed a temporary moratorium on targeted fishing for Sawfishes in Lake Nicaragua in the early 1980s (Thorson, 1982), after the population collapsed following intensive fishing in the 1970s. The aim was to allow the population to recover, but no such recovery has occurred (McDavitt, 2002). It appears that even bycatch mortality is sufficient to prevent population growth.

Sawfish are regularly used for their meat; however, most of the consumption is local and so they appear to be only occasionally traded beyond local markets (NMFS, 2009). The meat is white and tender, particularly in juveniles, and is one of the most valuable and preferred of all elasmobranchs (sharks and rays) sold in the city of Belém, Pará State, Brazil (Charvet-Almeida, 2002) and caught by Guinéan fishers (Doubouya, 2004). A large individual can yield several hundred kg of valuable meat (Last and Stevens 1994). The rostral saws can be very valuable as curios (particularly those from the largest specimens). In North Brazil (Pará State) Charvet-Almeida (2002) reports that large saws (>1.5 m) are ordered by buyers before fishing starts and may be worth up to US\$ 300 to the fisherman, depending upon size. There is a significant market in Chinese Taipei for Sawfish saws that are part of the ceremonial equipment/weapons of spirit mediums (there are an estimated 23,000 of these mediums in Taiwan). The small saws, from newborn and juvenile Sawfish, are sold as curios, or ground up as a local treatment for asthma (in Brazil), or exported for use in traditional Chinese medicine.

Habitat degradation and loss also threaten Sawfishes throughout their range (CITES, 2007). The Smalltooth Sawfish relies on a variety of specific habitat types including estuaries and mangroves; these are all affected by human development (CITES, 2007). Agricultural and urban development, commercial activities, dredge-and-fill operations, boating, erosion, and diversions of freshwater runoff as a result of continued coastal and catchment development has caused substantial loss or modification of these habitats (CITES, 2007).

The other significant problem is that the species are only protected by a very few range States. Any national conservation initiative intended to prevent these Critically Endangered species from being driven further towards extinction is unlikely to be successful if Sawfishes are not protected during their seasonal migrations through other range States' waters. This is a particular problem when the population is distributed along a coastline that is divided into a large number of small countries, as is the case in the Central Caribbean.

Sawfish rostra are often traded as curios, ceremonial weapons, or for use in traditional medicines, and artificial spurs for cock fights (NMFS, 2009). Rostra have long been a favorite marine curiosity (Migdalski, 1981), with large rostra commanding impressive prices (McDavitt 1996). These rostral teeth are mostly obtained from Brazil, Ecuador, Panama and various Caribbean countries (CITES, 2007). Sawfish skin has been used to produce leather, which, like shark leather, is considered of very high quality (NMFS, 2009). The leather is used to make belts, boots, purses, and even to cover books (NMFS, 2009).

## 2. Whale Shark (*Rhincodon typus*)

### Overview

Little is known about the life history of the Whale Shark but it attains a maximum size of 15-20 m and is likely to live to up to 60-100 years (Van Beek *et al.*, 2014). It is a highly migratory species and is widespread in tropical-temperate seas (Debrot *et al.*, 2013). Catches have declined and populations have apparently been depleted in several countries by harpoon fisheries targeting localized concentrations of this huge, slow-moving and behaviorally vulnerable species. There is also incidental capture in other fisheries. Directed fisheries, high value in international trade, a K-selected life history, highly migratory nature, and low abundance make this species vulnerable to exploitation. The Whale Shark is endangered worldwide and vulnerable in the Atlantic, according to IUCN. The species qualifies for listing on SPAW Annex 2 according to: criterion 1 due to the population decline, criterion 4 due to the vulnerable IUCN status, criterion 5 as the species is on Appendix II of CITES, and it qualifies for regional cooperative efforts under criterion 6 due to the north-south seasonal migration path which spans several countries jurisdictions.

### Species information

#### a. Scientific and common names of the species

1.1 Class: Chondrichthyes (subclass Elasmobranchii)

1.2 Order: Orectolobiformes

1.3 Family: Rhincodontidae

1.4 Species: *Rhincodon typus* (Smith 1828)

1.5 Scientific synonyms: Primarily variant spellings: *Rhiniodon typus*, *Rhineodon typus* Smith, 1828; Genus Rhinchodon Smith; Genus Rineodon Müller and Henle, 1838; Genus Rhineodon Müller and Henle, 1838; Genus Rhinodon and Rhineodon typicus Müller and Henle, 1839; Genus Rhiniodon Swainson, 1839; Genus Rhinecodon Agassiz, 1845; Genus Rhinodon Smith, 1849.

Other synonyms: *Micristodus punctatus* Gill, 1865. *Rhinodon pentalineatus* Kishinouye, 1901.

1.6 Common names:

English: Whale shark

French: Requin-baleine

Spanish: Tiburón ballena, pez dama (chequer-board fish)

Papiamentu: Tintorero

#### b. Estimated population of species and its geographic ranges;

Whale Sharks are found in all tropical and warm temperate seas except for the Mediterranean. They are occasionally recorded in oceanic waters but are most commonly reported in feeding aggregations close to the coast. Although widely distributed, they are generally infrequently recorded except in a few apparently favored coastal areas, where they are usually seen in relatively large numbers (tens to low hundreds) for only a few months of the year. The distribution records are characterized by highly seasonal appearances, with aggregations of Whale Sharks appearing for a few months in locations where their zooplankton food is abundant as a result of regular fish or invertebrate spawning events

(Fowler, 2000; Norman, in press; Heyman *et al.*, 2001). The species is certainly highly migratory, with satellite tracking of individuals demonstrating some very long-distance and long-term migrations, including a journey of over 2000 kilometers. Whether these migrations are solely driven by feeding events or linked to other aspects of their life history is yet to be determined. Genetic analysis showed little genetic differentiation on a global scale, although there is some genetic variance between the Atlantic, and the Indo-Pacific region.

Satellite tracking by Hueter *et al.* (2013) revealed movements of Whale Sharks into parts of the Caribbean Sea and the sharks' use of this tropical environment for up to several months. Accounts of Whale Sharks off Trinidad, Haiti, and the Bahamas are mentioned in the early literature but substantive, contemporary reports of *R. typus* in the eastern Caribbean Sea are lacking. However, the ECOCEAN database reports encounters from several islands in this area including Aruba, Dominica, Grenada, Puerto Rico, and the US Virgin Islands (as reviewed by Hueter *et al.* 2013). Compagno (2001) reported Whale Sharks off central Brazil, Colombia, Panama, and Venezuela. In a compilation of Whale Shark sightings over a 51-year period, Romero *et al.* (2000) reported 20 specimens of *R. typus* off Venezuela between the months of August and February with most sightings from a region of highly productive upwelled water. Debrot *et al.* (2013) documented 24 records of Whale Sharks for the Dutch Caribbean. Their results suggest a higher abundance of Whale Sharks in the southern, leeward part of the Dutch Caribbean, likely associated with seasonal upwelling-driven productivity known for the southeastern Caribbean area. A bimodal seasonal pattern as documented elsewhere for Venezuela was not pronounced in the Leeward Dutch islands and Whale Sharks were recorded in 9 months of the year. In the Windward Dutch islands all records so far were for the winter months of December-February.

There appears to be spatial and seasonal population segregation, with animals of similar size and largely the same sex often reported in the same area (Norman, 1999), while other age classes and a predominance of the other sex are found elsewhere (Eckert and Stewart, 2001; Graham, 2007). By analogy with other large migratory sharks, different age classes and sexes may undertake different migrations. Thus, juveniles may have different migration patterns from mature fish, and mature males and females may also have migration patterns of different lengths over different distances.

The global status of the Whale Shark is assessed as Endangered by IUCN (Pierce and Norman, 2016). They infer that approximately 75% of the global Whale Shark population occurs in the Indo-Pacific, and 25% in the Atlantic. In the Indo-Pacific, a population reduction of 63% is inferred over the last three generations (75 years), and in the Atlantic a population reduction of more than 30% is inferred. Combining data from both regions, it is likely that the global Whale Shark population has declined by >50% over the last 75 years.

Pierce and Norman (2016) base their inferred decline of  $\geq 30\%$  in the Atlantic subpopulation on data from tuna fleet observers off a likely center of abundance for this subpopulation. Between 1980 and 2010 there was a decline in sightings per unit effort (SPUE) off western Africa, with SPUE peaking in 1995 and declining thereafter (Sequeira *et al.* 2014; Table 1 in the supplementary material). In absolute terms, sightings decreased from about 500 during the 1990s to around 150 during the 2000s. Peak-month sightings also declined by approximately 50% over this time (Sequeira *et al.* 2014). At Gladden Spit in Belize, Whale Shark sightings declined from a mean of 4 to 6 sharks per day between 1998 and 2001 to less

than 2 per day in 2003 (Graham and Roberts 2007), with reports from diving guides indicating that numbers have remained low until 2016 (Pierce and Norman, 2016).

Pierce and Norman (2016) note that for the Atlantic subpopulation size regional counts of identified sharks or modelled abundance estimates are available from many of the larger known aggregation or feeding areas. Ramírez-Macías et al. (2012) photo-identified 350 individual Whale Sharks from Holbox Island in Mexico between 2005 and 2008, and estimated that 521–809 sharks participate in this aggregation. Aerial surveys from this area and the adjacent Caribbean coast have counted up to 420 sharks in a single aerial survey (de la Parra Venegas et al. 2011). The largest-known aggregation as of February 2016 occurs seasonally off the Yucatan coast of Mexico, with over 1,100 identified sharks (Norman et al. submitted). Satellite-tagged sharks from this aggregation have been tracked to the northern Gulf of Mexico (Hueter et al. 2013), where aggregations of up to 100 sharks have been reported (Hoffmayer et al. 2005), south to Belize where 106 individual sharks were identified between 1998 and 2003 (Graham and Roberts 2007), and off the island of Utila, Honduras, where 95 sharks were identified between 1999 and 2011 (Fox et al. 2013).

There is no detailed study of Whale Shark life history; estimates of age at maturity range from 9 to over 20 or 30 years, generation time from 24 to over 60 years, and longevity from 60 to over 100 years (e.g. Wintner, 2000). Even if the most conservative (lowest) estimates are taken, this is a very low-productivity, low-resilience species. Calculating life history parameters using Fishbase ([www.fishbase.org](http://www.fishbase.org)) and the 20 meter long shark reported by Chen *et al.* (1997) yields an estimate of 0.08/year intrinsic rate of population increase.

Gestation period and the interval between births are both unknown; only one litter of about 300 small near-term pups of 48-58 cm TL that grew rapidly in captivity has been reported (Joung *et al.*, 1996; Leu *et al.*, 1997). By analogy with the Nurse Shark *Ginglymostoma cirratum* (Castro, 2000), the only other Orectolobid shark for which detailed reproductive data are available, pregnancy may last for less than a year, but birth is likely followed by a long resting period and litters born only every two years. This strategy might explain the small number of pregnant females observed. The initial rapid growth of pups (Leu *et al.*, 1997) would explain the scarcity of records of very small Whale Sharks. Growth would slow rapidly at maturity (Pauly, 2002). A Whale Shark about 20m long and 34t in weight (as reported landed in Taiwan by Chen *et al.* 1997) could be over 100 years old.

There are several documented declines in seasonal catches by directed fisheries for the Whale Shark, with these declines having occurred in some areas over only a few years in relatively recent and short-lived intensive fisheries. Local populations have apparently declined drastically in some places, while fishing effort and price have greatly increased. Most of these fisheries are too recent and/or populations too poorly monitored to determine whether these declines would result in long-term (many decades) reductions in local populations even if closed. This may well be the case, by analogy with other large sharks, as a result of low productivity and rebound potential and a lack of migration into the area of unfished stocks from other sources. It is not known to what degree fishing in one area affects population(s) in other areas, although the fact that at least some of the sharks migrate long distances within ocean basins suggests that the effects may not be purely local. Thus, a fishery in one may affect numbers sighted in another area or even in a different region. There is increasing concern that unexplained declines in numbers sighted seasonally in apparently unfished areas such as Thailand and South Africa could be the result of fisheries impacting

these populations elsewhere. The rapid collapse of localized fisheries for this widely distributed and apparently seasonally migratory species could be explained by the tendency for Whale Sharks to be philopatric and to return regularly to the same seasonal feeding locations. Despite their very wide-ranging nature, they are, therefore, effectively part of local stocks that are particularly vulnerable to depletion by fisheries activity.

### **c. Status of legal protection, with reference to relevant national legislation or regulation**

#### **International**

##### ***Convention on the International Trade of endangered Species (CITES)***

The Whale Shark is listed in Appendix II of the Convention on Trade in Endangered Species of Flora and Fauna (CITES). This means that all transboundary trade has to be licensed, based on an analysis of the effects of the removal from the wild, or culture of the species – a Non-Detriment Finding ([www.cites.org](http://www.cites.org)).

##### **Convention for the protection of Migratory Species (CMS) – Memorandum of Understanding (MOU) on the Conservation of Migratory Sharks**

The whale shark is listed on Appendix II of the Convention on Migratory Species (CMS), and on Annex I of the Shark MoU, with the objective of achieving and maintaining a favorable conservation status for migratory sharks based on the best available scientific information, in particular the sharks listed on Annex 1 of the MOU, recognizing that successful shark conservation and management require the fullest possible cooperation among governments, intergovernmental organizations, nongovernmental organizations, and all stakeholders

##### ***United Nations Convention on the Law of the Sea (UNCLOS).***

The Whale Shark is also listed on Annex I (Highly Migratory Species) of UNCLOS, requiring cooperation, directly or through appropriate international organizations, to ensure the conservation and sustainable use of such species.

##### ***IPOA Sharks:***

Since the 1990s there are several shark protection plans, both internationally at intergovernmental and non-governmental level, as well as at national level by several nations in the Wider Caribbean region. Within the framework of the Code of Conduct for Responsible Fisheries the FAO (Food and Agriculture Organization) developed the International Plan of Action for the Conservation and Management of Sharks (IPOA Sharks) in 1999. The objective of IPOA Sharks is to ensure the conservation and management of sharks and their long-term sustainable use. IPOA Sharks is voluntary and intends to give states guidelines on how to establish a National Plan of Action (NPOA) through guiding principles and procedures for implementation.

#### **National Protection**

National legislations in the Caribbean region applying to sharks (as reviewed by Van Beek *et al.*, 2014) is as follows:

**US Caribbean Region:**

NOAA fisheries service presented the amendment 4 to the 2006 Consolidated Atlantic Highly Migratory Species (HMS) Fishery Management Plan (FMP). The PowerPoint states that “in 2010, Puerto Rico reported approximately 11.8 mt of commercial shark landings and less than one megaton was reported by St. Thomas and St. John combined. These landings were not species specific and it is unknown if they were harvested from Federal or Territorial waters”. Proposed management measures for small-scale HMS commercial fisheries include specific authorized gears and retention limits for sharks.

**US Gulf of Mexico and (Caribbean) Florida:**

Following years of declines in catches, and concern about the protection status of many shark species, in 1993 the USA established a Federal Management Plan for Shark Fisheries in the Atlantic Ocean, particularly directed at the coastal bottom long-line fishery. Since 1993 several amendments of the original plan have been implemented and local state governments have tied in by implementing complementary legislation. Measures included successively restrictive catch quotas, finning limitations, area closures, seasonal closures, adjustments of size limits, limits to retention in recreational fisheries, establishment of protected species lists, establish a shark research fishery and the use of regional and species specific quotas.

**Honduras:**

In June 2011 Honduras created the first shark sanctuary in America and declared all its marine waters in both the Pacific and Caribbean as a permanent shark sanctuary. This had been preceded in 2010 by a shark fishing moratorium and created the first shark sanctuary of the Americas amounting to about 240,000 km<sup>2</sup> of national waters, most of which lie along the 700 km-long Caribbean coast of the nation.

**Bahamas:**

The Bahamas have had a longline fishing ban since 1993 and consequently there has been no commercial shark fishing activity. This longline ban has effectively made the whole archipelago of the Bahamas a shark “no-take” zone. The last export of shark from the Bahamas was a lot of 2 metric tons in 2004. In July 2011 the Bahamas went a step further and legally banned all shark fishing. That law firmly turns all 630,000 sq km of Bahamian waters into a shark sanctuary<sup>17</sup>. The fines for shark fishing were raised from 3000 to 5000 USD per incident.

**Venezuela:**

Towards implementing its Plan de Acción Nacional (PAN) de conservación for sharks, in June 2012 Venezuela joined the rest of the Americas in outlawing the finning of sharks in its waters and established a 3,730 km<sup>2</sup> shark sanctuary surrounding the touristic archipelago of Los Roques. Recent research (e.g. Tavares 2005, 2008 2009) had demonstrated the importance of the shallow waters of Los Roques as a shark nursery area.

The Dominican Republic has, together with Belize and six other Central American countries, united under the name SICA (Central American Integration System), signed an agreement to prohibit shark finning. This ban is also applicable to fishing vessels in international waters under the flag of SICA member states. This arrangement OSP-05-11 entered into force in 1 January 2012.

**Kingdom of the Netherlands**

— **St. Maarten**

On the 12th of October 2011 the government of St. Maarten issued a temporary moratorium on shark fishing. The shark fishing moratorium prohibits the take and landing of sharks and requires immediate release of incidentally caught sharks, under penalty of a maximum of 500,000 Antillean Guilders or 3 months in prison.

— **Caribbean Netherlands**

In 2015, the Dutch government designated the Yarari sanctuary for sharks and marine mammals in the Economic Exclusive Zones of Saba and Bonaire, declaring that provisions will be considered and implemented as necessary to regulate activities that may have a negative impact on sharks.

— **Bonaire**

In 2008 the island of Bonaire passed a nature ordinance providing full protection for a list of species of plants and animals. This list includes all sharks and rays

#### **d. Ecological interactions with other species and specific habitat requirements**

The role of the Whale Shark in its ecosystem is unknown but, as a large plankton feeder, it may be similar to that of the smaller baleen whales. The Whale Shark is one of only three species of shark that filter feeds, the other two being the Megamouth (*Megachasma pelagios*) and Basking Shark (*Cetorhinus maximus*; Compagno, 1984). Unlike these two, the Whale Shark does not rely on forward motion for filtration, but is able to hang vertically in the water and suction feed by closing its gill slits and opening its mouth (Compagno, 1984). *R. typus* is believed to be able to sieve zooplankton as small as 1 mm in diameter through the fine mesh of their gill-rakers, and typically feeds on a variety of planktonic and nektonic prey, small crustaceans and schooling fishes and even occasionally ingesting small tuna and squid. Although the species occasionally feeds on eggs released by spawning aggregations of reef fish, this localized predatory activity is not considered likely to have a significant effect upon populations of the prey species (only a minute proportion of fertilized teleost eggs result in recruitment of adults to the population). Whale Sharks are known by traditional tuna fishermen to be associated with schools of tuna and have been used as natural 'fish aggregation devices' by tuna purse seiners in the Caribbean. Predators include killer whale, *Orcinus orca* and, for juveniles, blue marlin and blue shark.

#### **e. Management and recovery plans for the species**

National regulations are providing the strongest form of protection for Whale Sharks with total ban on fishing in Honduras in 1999, Belize and most recently in Mexico (as reviewed in Graham, 2007).

## **f. Research programs and available scientific and technical publications relevant to the species**

Observations of the Whale Shark in the Leeward Dutch Caribbean have shown a Whale Shark feeding in tuna schools positioned in a stationary vertical stance by opening and closing its mouth at the water surface (Debrot *et al.*, 2013). It was surrounded by schools of 2-ft-long yellowfin tunas (*Thunnus albacares*) that were preying on smaller baitfish which in turn sought protection in compact schools around the Whale Shark. This implies potential feeding benefit to the Whale Shark derived from the predatory activities of the tunas. Hoffmeyer *et al.* (2005) has remarked that the highest diversity of pelagic fish aggregations in the Gulf of Mexico are associated with Whale Sharks. It is proposed that the causal mechanism for this may be based on pelagic schools of baitfish seeking protection from (a diversity of) predators by schooling tightly around Whale Sharks. Colman (1997) and Hoffmeyer *et al.* (2005) have similarly described Whale Sharks feeding from a stationary vertical stance, also referred to as “suction-feeding” (Hoffmeyer *et al.*, 2005). However, the bulk of local observers indicate that most feeding behavior witnessed in the Dutch Caribbean concerned ram surface feeding (gill pumping movement) directed towards surface shoals of baitfish (Taylor, 2007). Other research suggests that Whale Sharks are gregarious and form seasonal aggregations in some coastal waters (De la Parra Venegas *et al.*, 2011). The authors describe an aggregation that occurs annually north of Cabo Catoche, off Isla Holbox on the Yucatán Peninsula of Mexico. Another, much denser aggregation of Whale Sharks (dubbed “the Afuera”) is described as appearing to occur off to the east of the tip of the Yucatán Peninsula in the Caribbean Sea, with 420 Whale Sharks aggregating in an area 18 km<sup>2</sup> in 2009 (De la Parra Venegas *et al.*, 2011). The authors note that plankton studies indicated that the sharks were feeding on dense homogenous patches of fish eggs, identified as belonging to little tunny, *Euthynnus alletteratus*. This contrasts with the annual Cabo Catoche aggregation nearby, where prey consists mostly of copepods and scombrid shrimp. Increased sightings at the Afuera coincide with decreased sightings at Cabo Catoche, and both groups have the same sex ratio, implying that the same animals are likely involved in both aggregations; tagging data support this idea. With two Whale Shark aggregation areas, high coastal productivity and a previously unknown scombrid spawning ground, the northeastern Yucatán marine region is a critical habitat that deserves more concerted conservation efforts (De la Parra Venegas *et al.*, 2011).

## **g. Threats to the species, its habitats and associated ecosystems, especially threats which originate outside the jurisdiction of the Party**

Small-scale harpoon and entanglement fisheries have taken place in various regions of the world, including India, Pakistan, Taiwan, the Philippines, and the Maldives. These took Whale Sharks primarily for their meat, liver oil, and/or fins. Liver oil was traditionally used for water-proofing boat hulls. The huge fins are low quality but of high value as restaurant “signboards” in East Asia, and the soft meat (known as “tofu shark”) are in great demand in Taiwan (Province of China).

In the Caribbean, since there are no targeted fisheries, threats to Whale Sharks stem primarily from unregulated tourism, aquaria collections and boat collisions. Research on

Whale Shark behavior indicates that patterns of movement exist, most notably following the bathymetric contours of the Mesoamerican Barrier Reef . This north-south path coincides with an important shipping lane that links the United States with the Mesoamerican reef countries. The volume of shipping and more recently cruise boat traffic and its potential impact to the regional Whale Shark population is undetermined. Coastal Development, cruise ship tourism, rising oil and gas exploration and land-based sources of pollution may pose additional yet site-variable direct and indirect threats to the region's Whale Shark population (Graham, 2007).

### 3. Oceanic Whitetip Shark - *Carcharhinus longimanus*

#### Overview

The Oceanic Whitetip Shark qualifies for inclusion under Annex 2 of SPAW under Criterion 1 and 4. The global status of the species is assessed by IUCN as *Vulnerable*, but in the Northwest and Western Central Atlantic as *Critically Endangered* because of the enormous declines that have been reported (Baum *et al.*, 2015). Two estimates of trends in abundance from standardized catch rate indices were made from independent datasets. An analysis of the US pelagic longline logbook data between 1992 and 2000, which covers the Northwest and Western Central Atlantic regions, estimated declines of 70%. An analysis of the Gulf of Mexico, which used data from US pelagic longline surveys in the mid-1950s and US pelagic longline observer data in the late-1990s, estimated a decline of 99.3% over this forty year time period or 98% over three generations (30 years), although this may be an overestimation. Fishing pressure on this species must be considerably decreased through reduction in fishing effort, catch limits, measures to enhance chances of survival after capture and possibly also through the implementation of large-scale oceanic non-fishing areas. Because of its migratory nature, effective conservation of this species will require international cooperation. The Whitetip is listed under CITES Appendix II and listed as highly migratory under UNCLOS. This also makes it eligible for the SPAW listing under criterion 5.

#### Species information

##### a. Scientific and common names of the species;

1.1 Class: Chondrichthyes

1.2 Order: Carcharhiniformes

1.3 Family: Carcharhinidae

1.4 Species: *Carcharhinus longimanus* (Poey, 1861)

1.5 Scientific synonyms: *Pterolamiops longimanus* (Poey, 1861), *Carcharius obtusus* (Garman, 1881), *Carcharius insularum* (Zinder, 1904), *Pterolamiops magnipinnis* (Smith, 1958), and *Pterolamiops budkeri* (Fourmanoir, 1961).

1.6 Common names:

English: Oceanic Whitetip Shark, Brown Milbert's sand bar shark, brown shark, nigarno shark, whitetip, whitetip shark, white-tip shark, and whitetip whaler

French: Requin océanique

Spanish: Tiburón punta blanca oceánico, aletiblanco oceánico, cazón, galano

##### b. Estimated population of species and its geographic ranges

The Oceanic Whitetip Shark is a globally widespread shark, ranging across entire oceans in tropical and subtropical waters. It is an oceanic-epipelagic shark, usually found far offshore in the open sea in waters 200 m deep, between about 30°N and 35°S in all oceans; it is normally found in surface waters, although it has been recorded to 152 m. It has occasionally been recorded inshore but is more typically found offshore or around oceanic islands and areas with narrow continental shelves (Fourmanoir, 1961, Compagno, 2005, Last and Stevens, 1994). Temperatures of waters in which it regularly occurs are 18 to 28°C, with water above 20°C preferred. Although one whitetip was caught in water of 15°C it tends to withdraw from waters that are cooling below this, as in the Gulf of Mexico in winter

(Compagno, 2005.). The location of nurseries has not been reported, but very young Oceanic Whitetip Sharks have been found well offshore along the southeastern US, suggesting offshore nurseries over the continental shelves (Compagno, 2005).

Smith *et al.* (1998) investigated the intrinsic rebound potential of Pacific sharks and found that Oceanic Whitetips have a moderate rebound potential, because of their relatively fast growth and early maturation. The population dynamics and structure of this species are unknown. Distribution appears to depend on the size and sex and the nursery areas appear to be oceanic (Seki *et al.*, 1998). Larger individuals are caught deeper than smaller ones and there is geographic and sexual segregation (Anderson and Ahmed, 1993).

Despite being initially described as the most common pelagic shark throughout the warm-temperate and tropical waters of the Atlantic (Mather and Day, 1954) and beyond the continental shelf in the Gulf of Mexico (Bullis, 1961), enormous declines are estimated to have occurred in the Northwest and Western Central Atlantic. Two estimates of trends in abundance from standardized catch rate indices have been made from independent datasets. An analysis of the US pelagic longline logbook data between 1992 and 2000, which covers the Northwest and Western Central Atlantic regions, estimated declines of 70% (Baum *et al.* 2003) and 57% from 1992 to 2005 (Cortés *et al.*, 2007). An analysis of the Gulf of Mexico, which used data from US pelagic longline surveys in the mid-1950s and US pelagic longline observer data in the late-1990s, estimated a decline of 99.3% over this forty year time period (Baum and Myers, 2004). When trends in abundance from the former analysis are extrapolated back to the mid-1950s, they match the latter analysis almost exactly (99.8%). Over a period of three generations (30 years), the estimated decline is 98%. However, the latter study has recently been criticized because temporal changes in fishing gear and practices over the time period were not taken fully into account and the study may, therefore, have exaggerated or underestimated the magnitude of the declines (Burgess *et al.*, 2005; Baum *et al.*, 2005).

### **c. Status of legal protection, with reference to relevant national legislation or regulation**

#### **International**

##### ***Convention on the International Trade of endangered Species (CITES)***

The Oceanic Whitetip is listed under Appendix II of CITES in 2013. This means that although the species is not necessarily currently threatened with extinction, it may become so unless trade is strictly regulated to avoid utilization incompatible with their survival. International (commercial) trade is permitted but regulated through a licensing system ([www.cites.org](http://www.cites.org)).

##### **Convention for the protection of Migratory Species (CMS) – Memorandum of Understanding (MOU) on the Conservation of Migratory Sharks**

*Carcharinus longimanus* is listed on Annex 1 of the Memorandum of Understanding (MOU) on the Conservation of Migratory Species. The Shark MOU is the first global instrument for the conservation of migratory species of sharks. Signatories to the MOU commit to the

objective of achieving and maintaining a favorable conservation status for migratory sharks based on the best available scientific information, in particular the sharks listed on Annex 1 of the MOU, recognizing that successful shark conservation and management require the fullest possible cooperation among governments, intergovernmental organizations, nongovernmental organizations, and all stakeholders

### ***IPOA Sharks***

There are since the 1990s several shark protection plans, both internationally at intergovernmental and non-governmental level, as well as at national level by several nations in the Wider Caribbean region. Within the framework of the Code of Conduct for Responsible Fisheries the FAO (Food and Agriculture Organization) developed the International Plan of Action for the Conservation and Management of Sharks (IPOA Sharks) in 1999. The objective of IPOA Sharks is to ensure the conservation and management of sharks and their long-term sustainable use. IPOA Sharks is voluntary and intends to give states guidelines on how to establish a National Plan of Action (NPOA) through guiding principles and procedures for implementation.

### **National Protection**

National legislations in the Caribbean region applying to sharks (as reviewed by Van Beek *et al.*, 2014) are as follows:

#### ***US Caribbean Region***

NOAA fisheries service presented the amendment 4 to the 2006 Consolidated Atlantic Highly Migratory Species (HMS) Fishery Management Plan (FMP). The powerpoint states that “in 2010, Puerto Rico reported approximately 11.8 mt of commercial shark landings and less than one megaton was reported by St. Thomas and St. John combined. These landings were not species specific and it is unknown if they were harvested from Federal or Territorial waters”. Proposed management measures for small-scale HMS commercial fisheries include specific authorized gears and retention limits for sharks.

#### ***US Gulf of Mexico and (Caribbean) Florida***

Following years of declines in catches, and concern about the protection status of many shark species, in 1993 the USA established a Federal Management Plan for Shark Fisheries in the Atlantic Ocean, particularly directed at the coastal bottom long-line fishery. Since 1993 several amendments of the original plan have been implemented and local state governments have tied in by implementing complementary legislation. Measures included successively restrictive catch quotas, finning limitations, area closures, seasonal closures, adjustments of size limits, limits to retention in recreational fisheries, establishment of protected species lists, establish a shark research fishery and the use of regional and species specific quotas.

#### ***Honduras***

In June 2011 Honduras created the first shark sanctuary in America and declared all its marine waters in both the Pacific and Caribbean as a permanent shark sanctuary. This had been preceded in 2010 by a shark fishing moratorium and created the first shark sanctuary of the Americas amounting to about 240,000 km<sup>2</sup> of national waters, most of which lie along the 700 km-long Caribbean coast of the nation.

### ***Bahamas***

The Bahamas have had a longline fishing ban since 1993 and consequently there has been no commercial shark fishing activity. This longline ban has effectively made the whole archipelago of the Bahamas a shark “no-take” zone. The last export of shark from the Bahamas was a lot of 2 metric tons in 2004. In July 2011 the Bahamas went a step further and legally banned all shark fishing. That law firmly turns all 630,000 sq km of Bahamian waters into a shark sanctuary<sup>17</sup>. The fines for shark fishing were raised from 3000 to 5000 USD per incident.

### ***Venezuela***

Towards implementing its Plan de Acción Nacional (PAN) de conservación for sharks, in June 2012 Venezuela joined the rest of the Americas in outlawing the finning of sharks in its waters and established a 3,730 km<sup>2</sup> shark sanctuary surrounding the touristic archipelago of Los Roques. Recent research (e.g. Tavares 2005, 2008 2009) had demonstrated the importance of the shallow waters of Los Roques as a shark nursery area.

The Dominican Republic has, together with Belize and six other Central American countries, united under the name SICA (Central American Integration System), signed an agreement to prohibit shark finning. This ban is also applicable to fishing vessels in international waters under the flag of SICA member states. This arrangement OSP-05-11 entered into force in 1 January 2012.

### ***Kingdom of the Netherlands***

#### **— *St. Maarten***

On the 12th of October 2011 the government of St. Maarten issued a temporary moratorium on shark fishing. The shark fishing moratorium prohibits the take and landing of sharks and requires immediate release of incidentally caught sharks, under penalty of a maximum of 500,000 Antillean Guilders or 3 months in prison.

#### **— *Caribbean Netherlands***

In 2015, the Dutch government designated the Yarari sanctuary for sharks and marine mammals in the Economic Exclusive Zones of Saba and Bonaire, declaring that provisions will be considered and implemented as necessary to regulate activities that may have a negative impact on sharks.

#### **— *Bonaire***

In 2008 the island of Bonaire passed a nature ordinance providing full protection for a list of species of plants and animals. This list includes all sharks and rays.

## **d. Ecological interactions with other species and specific habitat requirements**

Oceanic Whitetip Sharks are high trophic-level predators in the open ocean, feeding mainly on teleosts and cephalopods (Backus, 1956), but some studies have also reported that they prey on sea birds and marine mammals, among others (Compagno, 1984). Based on the diet of the oceanic white shark, Cortés (1999) determined that its trophic level was 4.2 (maximum=5.0).

### **e. Management and recovery plans for the species**

Conservation and management action are urgently required for this species; the only known conservation measure at present is a broad, multi-species pelagic shark quota for U.S. Atlantic waters. Specifically, fishing pressure on this species must be considerably decreased through reduction in fishing effort, catch limits, measures to enhance chances of survival after capture and possibly also through the implementation of large-scale oceanic non-fishing areas. Effective conservation of this species will require international cooperation. The Oceanic Whitetip is listed as a highly migratory species under the 1995 UN Agreement on the Conservation and Management of Straddling Fish Stocks and Highly Migratory Fish Stocks (UNFSA). The Agreement specifically requires coastal States and fishing States to cooperate and adopt measures to ensure the conservation of these listed species. To date, there is little progress in this regard. See United Nations Convention on the Law of the Sea for further details. Also of relevance is the FAO International Plan of Action for the Conservation and Management of Sharks (IPOA-Sharks) which specifically recommends that Regional Fisheries Organisations (RFO) carry out regular shark population assessments and that member States cooperate on joint and regional shark management plans. This is of particular importance for pelagic sharks such as *C. longimanus* whose stocks are exploited by more than one State on the high seas. Although steps are being taken by some RFOs to collect species-specific data on pelagic sharks, and to ban the practice of shark finning, to date no RFO has limited shark catches or drafted a "Shark Plan" as suggested in the IPOA-Shark guidelines (R. Cavanagh, pers. comm).

### **f. Research programs and available scientific and technical publications relevant to the species**

Research is being carried out on the Oceanic Whitetip Sharks (*Carcharhinus longimanus*) in the western North Atlantic following severe declines in abundance and the identification of the need for conservation measures (Howey-Jordan *et al.*, 2013). The research brings to light the spatial and temporal distribution of the individuals and the potential interaction with fishing gear during their migrations. Individuals have been tagged with pop-up satellite archival tags near Cat Island in the central Bahamas 1–8 May 2011 to provide information about the horizontal and vertical movements of this species. The individuals remained within 500 km of the tagging area for about 30 days and then dispersed across 16,422 km<sup>2</sup> of the western North Atlantic. Maximum individual displacement from the tagging site ranged from 290–1940 km after times at liberty from 30–245 days, with individuals moving to several different destinations (the northern Lesser Antilles, the northern Bahamas, and north of the Windward Passage). Many sharks returned to The Bahamas after ~150 days. Sharks spent 99.7% of their time shallower than 200 m and did not exhibit differences in day and night mean depths. All individuals made short duration (mean = 13.06 minutes) dives into the mesopelagic zone (down to 1082 m and 7.75°C), which occurred significantly more often at night. Ascent rates during these dives were significantly slower than descent rates, suggesting that these dives are for foraging. The sharks tracked appear to be most vulnerable to pelagic fishing gear deployed from 0–125 m depths, which they may encounter from June to October after leaving the protected waters of The Bahamas EEZ.

**g. Threats to the species, its habitats and associated ecosystems, especially threats which originate outside the jurisdiction of the Party**

Oceanic Whitetip Sharks have been caught in large numbers virtually everywhere they occur, particularly in pelagic longline and driftnet fisheries. This species was initially described as the most common pelagic shark beyond the continental shelf in the Gulf of Mexico (Wathne, 1959; Bullis, 1961), and throughout the warm-temperate and tropical waters of the Atlantic and Pacific (Mather and Day 1954, Strasburg 1957). In the Gulf of Mexico, for example, between 2 and 25 of these sharks were usually observed following the vessel during longline retrieval on the exploratory surveys in the 1950s and their abundance was considered as a serious problem because of the high proportion of tuna they damaged (Bullis and Captiva, 1955; Backus *et al.*, 1956; Wathne, 1959). Recent shark papers on the Gulf of Mexico have either not mentioned this species or have dismissed it as rare, not recognizing its former prevalence in the area (Baum and Myers 2004).

Few data are available on the catch rate of these sharks, and this is a serious hindrance to assessing the status of this species in regions other than the Northwest Atlantic and Eastern Central Pacific. According to Berkeley and Campos (1988), Oceanic Whitetip Sharks constituted 2.1% of the shark bycatch in the swordfish fishery along the east coast of Florida in 1981 to 1983. Information collected by at-sea scientific observers on U.S.-flagged longline vessels in the western North Atlantic Ocean indicates that Oceanic Whitetip is the 8th most abundant pelagic species caught. However, the low abundance of this species likely reflects the distribution of the fishery, as most U.S.-flagged vessels fish at the northernmost part of the range of the Oceanic Whitetip Shark (Beerkircher *et al.*, 2002). The United States reports that commercial fisheries land very few Oceanic Whitetip Sharks. Except for two peaks of about 1,250 and 1,800 sharks landed in 1983 and 1998, respectively, total catches never exceeded 450 individuals per year. However, the proportion of the catch of Oceanic Whitetip Shark increases in areas of the Atlantic Ocean that are more tropical than temperate. For example, Oceanic Whitetip Sharks were present in 4.72% of eastern tropical Atlantic French and Spanish tuna purse-seine sets (Santana *et al.*, 1997). Domingo (2004) reported that the Uruguayan longline fleet observer program in 1998-2003 recorded catch rates of 0.006 sharks/1,000 hooks in Uruguayan and adjacent high seas South Atlantic waters (latitude 260-370, 16-230C) but catch rates increased to 0.09 sharks/1,000 hooks in international waters off western equatorial Africa. Only Brazil, Mexico, Spain, St. Lucia and the United States have reported catches to ICCAT and, as indicated by Clarke (2008), these data are likely inaccurate and therefore may under-represent the magnitude of catches in the Atlantic Ocean. This species has been recorded as part of the catch of oceanic longline industrial fisheries in the Colombian Caribbean, with mean catch sizes of 128 +/- 62.35 cm TL, which corresponds to juveniles and may be impacting likely development areas (Caldas and Correa, 2010).



## 1. Manta rays (*Manta birostris*, *Manta alfredi*, *Manta cf. birostris*)

### Overview

The giant Manta Ray *Manta birostris* and the reef Manta Ray *Manta alfredi*, with a third putative species endemic to the Caribbean region, *Manta cf. birostris*, are the largest genus of rays, making their life history especially conservative, and rendering them vulnerable to depletion. Moreover, despite evidence for long migrations, regional populations appear to be small, sparsely distributed, and fragmented, meaning localized declines are unlikely mitigate by immigration. Both species of Manta Ray have recently been reassessed for the IUCN Red List, which looks at different species against a range of criteria to see what issues are of concern to the species survival. Both species of manta are considered to be ‘Vulnerable’ on this listing. Giant mantas have also recently been listed on Appendix I and II under the Convention on Migratory Species (CMS), and both species are listed in Appendix II of the Convention on International Trade of Endangered Species (CITES). Listing of the Manta Ray in Annex 3 of SPAW would thus be consistent with international agreements and would be compliant with criteria 4 (IUCN), 5 (CITES) and 6 (regional cooperation). Criterion 1 is met due to the decline and fragmentation of the populations.

### Species information

#### a. Scientific and common names of the species

1.1 Class: Chondrichthyes (Subclass: Elasmobranchii)

1.2 Order: Rajiformes

1.3 Family: Mobulidae

1.4 Genus and species:

All species of Genus *Manta birostris* (Donndorff 1798), *Manta alfredi* (Kreffft, 1868), *Manta cf. birostris* (putative) and any other putative *Manta* species.

1.5 1.5a. Scientific synonyms:

*M. birostris*: *Manta hamiltoni* (Hamilton & Newman 1849); *Raja birostris* (Donndorff, 1798)

*M. alfredi*: *Deratoptera alfredi* (Kreffft, 1868); *Manta fowleri* (Whitney, 1936)

1.6 Common names:

*M. birostris*: English: Oceanic Manta Ray, Giant Manta Ray, Chevron Manta Ray, Pacific Manta Ray, Pelagic Manta Ray

Spanish: Manta Comuda, Manta Diablo, Manta Gigante, Manta Raya, Manta Voladora.

*M. alfredi*: English: Reef Manta Ray, Coastal Manta Ray, Inshore Manta Ray, Prince Alfred’s Ray, Resident Manta Ray.

#### b. Estimated population of species and its geographic ranges

The Giant Manta Ray *M. birostris* occurs in tropical, sub-tropical and temperate waters of the Atlantic, Pacific and Indian Ocean. The reef Manta *M. alfredi* is found in tropical and subtropical waters (Marshall *et al.*, 2009; Kashiwagi *et al.*, 2011; Couturier *et al.*, 2012). A

possible subspecies *Manta cf birostris* appears to be a regional endemic with a reported distribution throughout the Gulf of Mexico, the Caribbean, and along the eastern coast of the United States. *Manta birostris* are thought to be seasonal visitors along productive coastlines with regular upwelling, in oceanic island groups, and near offshore pinnacles and seamounts. They visit cleaning stations on shallow reefs, are sighted feeding at the surface inshore and offshore, and are also occasionally observed in sandy bottom areas and seagrass beds (Marshall *et al.*, 2011). *M. alfredi* is commonly sighted inshore but is also observed around offshore coral reefs, rocky reefs, and seamounts. This species is often resident in or along productive near-shore environments, such as island groups, atolls, or continental coastlines, and may also be associated with areas or events of high primary productivity (e.g., upwelling; Homma *et al.*, 1999; Dewar *et al.*, 2008; Kitchen-Wheeler, 2010; Anderson *et al.*, 2011; Deakos *et al.*, 2011; Marshall *et al.*, 2011a). *Manta cf birostris* exhibits similar habitat preferences to *M. alfredi*.

The Manta Ray is a migratory species, A global investigation of major aggregation sites revealed that the Giant Manta Ray may be a more oceanic and a more migratory species than the Reef Manta Ray (A. Marshall *et al.*, unpubl. data). Rare or seasonal sightings of the Giant Manta Ray at locations such as northern New Zealand (Duffy and Abbott, 2003), southern Brazil (Luiz *et al.*, 2009) and Uruguay (Milessi and Oddone, 2003), the Azores Islands, the Similan Islands, Thailand (A. Marshall, unpubl. data) and the eastern coast of the United States (Bigelow and Schroeder, 1953), suggests that this species undergoes significant seasonal migrations.

Despite these data, preliminary satellite tracking studies and international photo-identification matching projects have suggested a high degree of fragmentation between regional populations of this species, suggesting that movements across ocean basins may be rare. Satellite tracking results have been able to reveal that the Giant Manta Ray is capable of large migrations (over 1,100 km straight line distance) and have monitored individual movements across international borders, across large bodies of water, and into international waters (A. Marshall *et al.*, unpubl. data; R. Rubin, pers. comm.). Satellite tracking studies using archival PAT tags have registered movements of the Giant Manta Ray from Mozambique to South Africa (a distance of 1,100 km), from Ecuador to Peru (190 km), from the Yucatan, Mexico into the Gulf of Mexico (448 km). This species is capable of deep dives and has been both seen at depth and tracked down to depths exceeding 1,000 meters (A. Marshall *et al.*, unpubl. data), as reviewed by Marshall *et al.* (2011).

Despite the long distance migrations, it is believed that regional populations are rather small. Individuals demonstrated a degree of site fidelity to specific regions, as well as critical habitats within them. Because of the global nature of their individual distributions, absolute population sizes will always be difficult to assess. Currently, the overall total global population sizes of both Manta species are unknown, but subpopulations appear, in most cases, to be less than 1,000 individuals, as well as sparsely distributed, and highly fragmented. Also, limited interchanging between populations is suggested, likely due to their resource and habitat needs, meaning declines are not likely to be mitigated by immigration.. The degree of interchange of individuals between subpopulations is assumed to be low because there are currently no data that support such interchange, despite active efforts to do so (A. Marshall *et al.*, unpubl. data). The giant Manta Ray, unlike the reef Manta Ray, is not often encountered in schools of more than 30 fish when feeding. In general, they are less

frequently observed than reef Manta Rays, despite having a larger distribution across the world.

Depletion has been documented in some monitored subpopulations in the Philippines, Indonesia, and Mexico. Fishermen and divers in Mozambique, Madagascar, Sri Lanka, Thailand, and Australia have offered much anecdotal evidence of population declines over the last decade as a result of increased fishing (TRAFFIC, 2013). Overall, the rate of population reduction appears to be high in several regions, up to as much as 80% over the last three generations (approximately 75 years), and globally a decline of >30% is strongly suspected.

### **c. Status of legal protection, with reference to relevant national legislation or regulation**

#### **International**

##### ***Convention on the International Trade of endangered Species (CITES )***

The genus of Manta Rays is listed in Appendix II of the Convention on International Trade in Endangered Species (CITES). This means that all transboundary trade has to be licensed, based on an analysis of the effects of the removal from the wild, or culture of the species – a Non-Detriment Finding ([www.cites.org](http://www.cites.org)).

##### **Convention for the protection of Migratory Species (CMS) – Memorandum of Understanding (MOU) on the Conservation of Migratory Sharks**

Manta rays are listed under the Convention on Migratory Species (CMS) and in Annex 1 of the Shark MoU with the objective of international cooperation for their conservation. Signatories to the MOU commit to the objective of achieving and maintaining a favorable conservation status for migratory sharks based on the best available scientific information, in particular the sharks listed on Annex 1 of the MOU, recognizing that successful shark conservation and management require the fullest possible cooperation among governments, intergovernmental organizations, nongovernmental organizations, and all stakeholders

##### ***IPOA Sharks:***

There are since the 1990s several shark protection plans, both internationally at intergovernmental and non- governmental level, as well as at national level by several nations in the Wider Caribbean region. Within the framework of the Code of Conduct for Responsible Fisheries the FAO (Food and Agriculture Organization) developed the International Plan of Action for the Conservation and Management of Sharks (IPOA Sharks) in 1999. The objective of IPOA Sharks is to ensure the conservation and management of sharks and their long-term sustainable use. IPOA Sharks is voluntary and intends to give states guidelines on how to establish a National Plan of Action (NPOA) through guiding principles and procedures for implementation.

#### **National Protection**

National legislations in the Caribbean region applying to sharks (as reviewed by Van Beek *et al.*, 2014) are as follows:

**US Caribbean Region:**

NOAA fisheries service presented the amendment 4 to the 2006 Consolidated Atlantic Highly Migratory Species (HMS) Fishery Management Plan (FMP). The PowerPoint states that “in 2010, Puerto Rico reported approximately 11.8 mt of commercial shark landings and less than one megaton was reported by St. Thomas and St. John combined. These landings were not species specific and it is unknown if they were harvested from Federal or Territorial waters”. Proposed management measures for small-scale HMS commercial fisheries include specific authorized gears and retention limits for sharks.

**US Gulf of Mexico and (Caribbean) Florida:**

Following years of declines in catches, and concern about the protection status of many shark species, in 1993 the USA established a Federal Management Plan for Shark Fisheries in the Atlantic Ocean, particularly directed at the coastal bottom long-line fishery. Since 1993 several amendments of the original plan have been implemented and local state governments have tied in by implementing complementary legislation. Measures included successively restrictive catch quotas, finning limitations, area closures, seasonal closures, adjustments of size limits, limits to retention in recreational fisheries, establishment of protected species lists, establish a shark research fishery and the use of regional and species specific quotas.

**Honduras:**

In June 2011 Honduras created the first shark sanctuary in America and declared all its marine waters in both the Pacific and Caribbean as a permanent shark sanctuary. This had been preceded in 2010 by a shark fishing moratorium and created the first shark sanctuary of the Americas amounting to about 240,000 km<sup>2</sup> of national waters, most of which lie along the 700 km-long Caribbean coast of the nation.

**Bahamas:**

The Bahamas have had a longline fishing ban since 1993 and consequently there has been no commercial shark fishing activity. This longline ban has effectively made the whole archipelago of the Bahamas a shark “no-take” zone. The last export of shark from the Bahamas was a lot of 2 metric tons in 2004. In July 2011 the Bahamas went a step further and legally banned all shark fishing. That law firmly turns all 630,000 sq km of Bahamian waters into a shark sanctuary<sup>17</sup>. The fines for shark fishing were raised from 3000 to 5000 USD per incident.

**Venezuela:**

Towards implementing its Plan de Acción Nacional (PAN) de conservación for sharks, in June 2012 Venezuela joined the rest of the Americas in outlawing the finning of sharks in its waters and established a 3,730 km<sup>2</sup> shark sanctuary surrounding the touristic archipelago of Los Roques. Recent research (e.g. Tavares 2005, 2008 2009) had demonstrated the importance of the shallow waters of Los Roques as a shark nursery area.

The Dominican Republic has, together with Belize and six other Central American countries, united under the name SICA (Central American Integration System), signed an agreement to prohibit shark finning. This ban is also applicable to fishing vessels in international waters

under the flag of SICA member states. This arrangement OSP-05-11 entered into force in 1 January 2012.

## **Kingdom of the Netherlands**

### ***Kingdom of the Netherlands***

#### — ***St. Maarten***

On the 12th of October 2011 the government of St. Maarten issued a temporary moratorium on shark fishing. The shark fishing moratorium prohibits the take and landing of sharks and requires immediate release of incidentally caught sharks, under penalty of a maximum of 500,000 Antillean Guilders or 3 months in prison.

#### — ***Caribbean Netherlands***

In 2015, the Dutch government designated the Yarari sanctuary for sharks and marine mammals in the Economic Exclusive Zones of Saba and Bonaire, declaring that provisions will be considered and implemented as necessary to regulate activities that may have a negative impact on sharks.

#### — **Bonaire**

In 2008 the island of Bonaire passed a nature ordinance providing full protection for a list of species of plants and animals. This list includes all sharks and rays.

## **d. Ecological interactions with other species and specific habitat requirements**

The role of the *Manta spp.* in their ecosystem is not fully known but, as large plankton feeders, it may be similar to that of the smaller baleen whales. As large species which feed low in the food chain, *Manta spp.* can be viewed as indicator species for the overall health of the ecosystem. Studies have suggested that removing large, filter-feeding organisms from marine environments can result in significant, cascading species composition changes (Springer *et al.*, 2003).

## **e. Management and recovery plans for the species**

Manta rays are protected under CITES Appendix II, meaning that the species is not necessarily currently threatened with extinction, but may become so unless trade is strictly regulated to avoid utilization incompatible with their survival. International (commercial) trade is permitted but regulated. See: [www.cites.org](http://www.cites.org)

The reef manta (*Manta alfredi*) and giant manta (*Manta birostris*) are protected in all waters where EU fleets are allowed to fish according to the EU TAC and quatum regulation (EU 2016/71) which will be reformed during 2016.

## **f. Research programs and available scientific and technical publications relevant to the species**

The spatio-temporal distribution of devil ray (*Manta birostri*) was studied using satellite tracking off the Yucatan peninsula in Mexico (Graham *et al.*, 2012). The authors describe that the Manta Rays were associated with seasonal upwelling events and thermal fronts off the peninsula, and made short-range shuttling movements, foraging along and between them. The majority of locations were received from waters shallower than 50 m deep, representing thermally dynamic and productive waters (Graham *et al.*, 2012).

The biology of elasmobranchs is among the most poorly known and least understood of all the marine vertebrate groups (Fowler, 2005). This is particularly so for the Caribbean region in specific. Within the Caribbean Sea, research efforts are made to assess distribution, habitat use, population structure, and trophic ecology of sharks using acoustic telemetry, satellite tagging methods, genetic analysis and stable isotope research. The occurrence and relative abundance of sharks are investigated using Baited Remote Underwater Videos (BRUVs). The skillful use of modern techniques such as genetic analyses, telemetry, and Baited Remote Video monitoring can help circumvent the often-low abundance (and low sampling) of many species, and should help develop powerful new insights and introduce new techniques to the region where capacity and technology have lagged behind.

## **g. Threats to the species, its habitats and associated ecosystems, especially threats which originate outside the jurisdiction of the Party**

The main threat to both Manta species is fishing, both targeted and incidental. Manta rays are currently killed or captured by a variety of methods including harpooning, netting and trawling. These rays are easy to target because of their large size, slow swimming speed, aggregative behavior, predictable habitat use, and lack of human avoidance. Specifically for the Caribbean, exploitation rates are unknown because of lacking landings data from fisheries.

Manta ray products have a high value in international trade markets. Their gill rakers are particularly sought after and are used in Asian medicinal products. This market has resulted in directed fisheries for Manta Rays, which are currently targeting these rays in unsustainable numbers. Over 1,000 Manta Rays are caught per year in some areas (Alava *et al.*, 2002; Dewar, 2002; White *et al.*, 2006; Anderson *et al.*, 2010). Artisanal fisheries also target both species for food and local products (White *et al.*, 2006; Marshall *et al.*, 2011).

Aside from directed fisheries, Manta Rays are also incidentally caught as bycatch in both large-scale fisheries and small netting programs such as shark control bather protection nets (Young 2001, C. Rose, pers. comm.).

As a result of sustained pressure from targeted fisheries and bycatch certain monitored subpopulations appear to have been rapidly depleted (e.g., Indonesia and the Philippines; Anon, 1997; Alava *et al.*, 2002). Targeting either species of *Manta* at critical habitats or aggregation sites, where individuals can be caught in large numbers in a short time frame, is a particular threat. Regional populations of both species appear to be small, and localized declines are unlikely to be mitigated by immigration. This situation is exacerbated by the

conservative life history of these rays, which constrain their ability to recover from a depleted state.

Cryptic threats such as mooring line entanglement and boat strikes can also wound Manta Rays, decrease fitness or contribute to unnatural mortality (Marshall *et al.*, 2011; Deakos *et al.*, 2011; F. McGregor pers. obs.). In Maui, Hawaii, 10% of the population has amputated or non-functioning cephalic fins, most likely caused from entanglement in monofilament fishing line (Deakos *et al.*, 2011). Many other threats have been postulated and identified such as habitat degradation, climate change, pollution (from oil spills), ingestion of micro plastics and irresponsible tourism practices.

Dive tourism involving this species is a growing industry and it has been demonstrated that sustainable tourism significantly enhances the economic value of such species in comparison to short-term returns from fishing (Anderson *et al.*, 2010). However, rapidly growing tourism (including in-water interactions and recreational boating traffic) if unmanaged, is likely to affect localized use of and visitation rates to critical cleaning and feeding habitats (Osada 2010; Deakos *et al.*, 2011). Their natural behavior can also be affected by excessive ecotourism (F. McGregor unpubl. data, A. Marshall unpubl. data).

## 2. Hammerhead sharks – *Sphyrna lewini*, *Sphyrna mokarran*, *Sphyrna zygaena*

### Overview

*Sphyrna lewini*, *Sphyrna mokarran* and *Sphyrna zygaena* are circumglobal shark species residing in coastal warm temperate and tropical coastal seas. *S. lewini* have among the lowest recovery potential when compared to other species of sharks. Population growth rates determined for populations in the Pacific and Atlantic Ocean are low ( $r=0.08-0.10$  yr<sup>-1</sup>) and fall under the low productivity category ( $r<0.14$ ) as defined by Food and Agriculture Organization of the United Nations (FAO). Abundance trend analyses of catch-rate data specific to *S. lewini* and to a hammerhead complex of *S. lewini*, including *Sphyrna mokarran* and *Sphyrna zygaena*, have reported large declines in abundance ranging from 60-99% over recent years. A stock assessment using information on catch, abundance trends and biology specific to *S. lewini* from the northwest Atlantic Ocean indicate a decline of 83% from 1981-2005. Standardized catch rates from the U.S. pelagic longline fishery show declines in *Sphyrna* spp. of 89% between 1986 and 2000 and declines of 76% between 1992 and 2005. Hammerhead fins are highly valued and they are being increasingly targeted in some areas in response to increasing demand for shark fins. Hammerhead shark species *S. zygaena* and *S. lewini* were found to represent at least 4-5% of the fins auctioned in Hong Kong, the world's largest shark fin trading center. Fins from the Hong Kong SAR market can be genetically assessed and have been shown to originate western Atlantic Ocean basins.

The listing of the Sphyrnidae family to Annex III of SPAW is warranted by the proliferating evidence for declining populations in the West-Atlantic Ocean, their vulnerability to overexploitation and low recovery potential due to a low intrinsic growth rate and slow reproduction, and sustaining targeted catch and bycatch in the Northwest and Western Central Atlantic ocean. The exact amount of fishing pressure and the corresponding mortality rate is obscure, and especially species-specific inferences cannot easily be made, because of the difficulties associated with the inability to distinguish between *S. zygaena*, *S. lewini*, and *S. mokarran*. The precautionary approach should be taken because of these constraints, and the dire situation of the conservation status of hammerhead sharks, which is assessed by IUCN as Endangered for *S. mokarran* and *S. lewini* (both on a global scale and the Northwest and Western Central Atlantic subpopulation) and Vulnerable for *S. zygaena*. The family of hammerheads is listed under Appendix II of CITES and in Annex I of the United Nations Convention on the Law of the Sea (UNCLOS) and should therefore be subject to its provisions concerning fisheries management in international waters.

In summary, the three hammerhead species are eligible for listing under SPAW Annex 3 (III) according to the criteria 1 (decline in population), 4 (IUCN listing), 5 (CITES) and 6 (the importance of regional cooperation to protect the species).

### Species information

#### a. Scientific and common names of the species

The family of Sphyrnidae, or hammerhead sharks, with primarily the following three species:

- Smooth hammerhead - *Sphyrna zygaena*
- Great hammerhead - *Sphyrna mokarran*

– Scalloped hammerhead - *Sphyrna lewini*

1. Class: Chondrichthyes (Subclass: Elasmobranchii)
2. Order: Carcharhiniformes
3. Family: Sphyrnidae
4. a Genus, species: *Sphyrna lewini* (Griffith and Smith, 1834)
5. a Scientific synonyms: *Cestracion leeuwenii* (Day 1865), *Zygaena erythraea* (Klunzinger 1871), *Cestracion oceanica* (Garman 1913), *Sphyrna diplana* (Springer 1941), *Sphyrna couardi* (Cadenat, 1951), *Zygaena lewini* (Griffith & Smith, 1834)
6. a Common names: English: scalloped hammerhead, bronze hammerhead shark, hammerhead, hammerhead shark, kidney-headed shark, scalloped hammerhead shark, and southern hammerhead shark,  
French: requin marteau halicorne  
Spanish: tiburón-martillo, cachona, cornuda común  
Portuguese: tubarão martelo, tubarão-martelo-entalhado, cambeva, cambeva-branca, cambevota, vaca, vacota, panã  
Papiamentu: tribon martin, krus
4. b Genus, species: *Sphyrna mokarran* (Rüppell, 1837)
5. b Scientific synonyms: *Zygaena mokarran* (Rüppell, 1837)
6. b Common names:  
Great Hammerhead, Squat-headed Hammerhead Shark, Hammerhead Shark  
French: Sorosena, Grand Requin-marteau, Marieau Millet, Poisson Pantouflier  
Spanish: Cornuda, El Tiburon, Guardia Civil, Pez Martillo, Tiburon  
Papiamentu: tribon martin, krus
4. c Genus, species: *Sphyrna zygaena* (Linnaeus 1758)
5. c Scientific synonyms:
6. c Common names: Smooth Hammerhead  
French: Requin-marteau commun, Requin marteau lisse  
Papiamentu: tribon martin, krus

## **b. Estimated population of species and its geographic ranges**

### ***Sphyrna lewini***

*S. lewini* is a coastal and semi-oceanic hammerhead shark that is circumglobal in coastal warm temperate and tropical seas, from the surface and intertidal to at least 275 m depth. Although it is wide ranging, there is genetic evidence for multiple subpopulations, with a separate subpopulations in the Northwest and Western Central Atlantic. Where catch data are available, significant declines have been documented: both species-specific estimates for *S. lewini* and grouped estimates for *Sphyrna* spp. combined suggest declines in abundance of 50-90% over periods of up to 32 years in several areas of its range, including the northwest Atlantic. Interviews with fishermen also suggest declining trends. Similar declines are also inferred in areas of the species' range from which specific data are not available, but fishing pressure is known to be high. Estimates of trends in abundance are available from two long-term research surveys conducted on the U.S. east coast, both of which indicate this species has undergone substantial declines in this region (98% between 1972 and 2003, and an

order of magnitude between 1975 and 2005). A third survey comparing catch rates between 1983/84 with those in 1993-95 showed a decline of two-thirds, while a survey beginning more recently showed increases in catch rates of juveniles. Standardized catch rates from the U.S. pelagic longline fishery show declines in *Sphyrna* spp. of 89% between 1986 and 2000 (according to the logbook data) and declines of 76% between 1992 and 2005 (according to observer data). The other information for this species from this region comes from Belize, where it has been heavily fished since the 1980s and fishermen have reported dramatic declines, which led to the end of the fishery. Guatemalan fishermen sustain fishing pressure in Belize (Baum *et al.*, 2007).

Recent studies indicate that the Northwest Atlantic, Caribbean Sea and Southwest Atlantic populations of this species are each genetically distinct from each other, and from Eastern Central Atlantic and Indo-Pacific populations (D. Chapman and M. Shivji, Nova, unpublished data). The boundaries between each population are not yet completely defined due to sampling constraints, but the "Caribbean Sea" population includes Belize and Panama and the "U.S. Gulf Of Mexico" sample covers from Texas to southwestern Florida, the boundary or transition zone will be in between Texas and Northern Belize (D. Chapman and M. Shivji, pers. comm.). Given the major declines reported in many areas of this species' range, increased targeting for its high value fins, low resilience to exploitation and largely unregulated, continuing fishing pressure from both inshore and offshore fisheries, this species is assessed by IUCN as Endangered globally, as well as in the Northwest and Western Central Atlantic (Baum *et al.*, 2007). Hayes *et al.* (2009) conducted an assessment in the Northwest Atlantic using two surplus production models. Population size in 1981 was estimated to be between 142,000 and 169,000 sharks, but decreased to about 24,000 sharks in 2005 (an 83-85% reduction). A new stock assessment by the NMFS for the northwestern Atlantic was released April 2011 Under the Magnuson Stevens Act. The stock assessment estimated that a total allowable catch (TAC) of 2,853 scalloped hammerhead sharks per year (or 69 percent of the 2005 catch) would allow a 70 percent probability of rebuilding to MSY in 10 years. Great hammerhead (*S. mokarran*) and smooth hammerhead (*S. zygaena*) are also part of the Atlantic Large Coastal Shark Complex, but are assessed at the complex level. The overfished and overfishing status of this complex is unknown as of the 4th quarter of 2011 (NMFS 4th Quarter 2011 stock status).

### ***Sphyrna mokarran***

*S. mokarran* ranges widely throughout the tropical waters of the world, from latitudes 40°N to 35°S. It is apparently nomadic and migratory, with some populations moving polewards in the summer, as off Florida and in the South China Sea. There is a pupping and nursery ground in a coastal mangrove estuarine area of southern Belize (R.T. Graham, pers. obs). The large, widely distributed, tropical hammerhead shark is largely restricted to continental shelves.

Although there is very little species-specific data available, the absence of recent records give cause to suspect a decline of at least 80% in the past 25 years. Fishing proceeds unmanaged and unmonitored, resulting in an assessment of Critically Endangered in the Eastern Atlantic. Although not targeted in the Northwest Atlantic and Gulf of Mexico it is taken as by-catch in several fisheries and suffers greater than 90% vessel mortality. Two time series data sets (pelagic logbook, large pelagic survey) have shown a decline in the catch of *Sphyrna* spp. since 1986. Difficulties in species identification and accurate recording make an assessment of this species very difficult, however low survival at capture makes it highly

vulnerable to fishing pressure, whether directed or incidental. It is therefore assessed by IUCN as Endangered in the Northwest Atlantic and Gulf of Mexico, based on a suspected decline of at least >50% over the past 10 years. The decline is poorly documented and has not been curtailed (Denham *et al.*, 2007).

### ***Sphyrna zygaena***

Specific data on populations of this species are generally unavailable in many areas because hammerhead shark catches are often grouped to include several *Sphyrna* species. Furthermore, this species has sometimes been confused with the *S. lewini* in the Caribbean and these two species are probably misidentified with each other. *Sphyrna zygaena* is one of the larger hammerhead sharks, found worldwide in temperate and tropical seas, with a wider range than other members of its family. It is semi-pelagic and occurs on the continental shelf. Although few data are available on the hammerhead's life-history characteristics, it is a large hammerhead shark and presumably at least as biologically vulnerable as *S. lewini*. Few species-specific data are available to assess population trends because catches of hammerhead sharks are often grouped together under a single category. Very often these sharks are finned and the carcasses discarded. This species has sometimes been confused with *S. lewini* in the tropics and these two species are probably misidentified with each other in some areas. Time series data on population trends in hammerhead sharks, including *S. zygaena*, are available from the Northwest and Western Central Atlantic and the Mediterranean Sea. In the Northwest and Western Central Atlantic, where *S. zygaena* is outnumbered by *S. lewini* by about ten to one, analysis of U.S. pelagic longline logbook data estimated that Sphyrnidae (including *S. lewini*, *S. mokarran* and *S. zygaena*) declined in abundance by 89% since 1986. In the Mediterranean Sea, where *S. zygaena* outnumbers *S. lewini*, compilation and meta-analysis of time series abundance indices estimated that Sphyrnidae (including *S. lewini*, *S. mokarran* and *S. zygaena*) declined by >99% in abundance and biomass since the early 19th century. The species is currently assessed by IUCN as Vulnerable (Casper *et al.*, 2005) and further investigation into threats, population trends, catches and life-history parameters throughout its range are required to determine whether it may warrant a higher category in the future.

## **c. Status of legal protection, with reference to relevant national legislation or regulation**

### **International**

#### ***Convention on the International Trade of endangered Species (CITES)***

The hammerhead species *S. lewini*, *S. mokarran* and *S. zygaena* are all listed in Appendix II of the Convention on International Trade in Endangered Species (CITES). This means that all transboundary trade has to be licensed, based on an analysis of the effects of the removal from the wild, or culture of the species – a Non-Detriment Finding ([www.cites.org](http://www.cites.org))

#### ***United Nations Convention on the Law of the Seas (UNCLOS)***

The family Sphyrnidae is listed on Annex I, Highly Migratory Species, of the UN Convention on the Law of the Sea. States are urged to cooperate over the management of these species. No such management yet exists.

## ***Convention for the protection of Migratory Species (CMS) – Memorandum of Understanding (MOU) on the Conservation of Migratory Sharks***

The Memorandum of Understanding on the conservation of migratory sharks (Sharks MoU) of the Convention on the Conservation of Migratory Species of Wild Animals (CMS) is a legally non-binding instrument of the CMS and the first global instrument for the conservation of migratory shark species. Signatories to the MOU commit to the objective of achieving and maintaining a favorable conservation status for migratory sharks based on the best available scientific information, in particular the sharks listed on Annex 1 of the MOU, recognizing that successful shark conservation and management require the fullest possible cooperation among governments, intergovernmental organizations, nongovernmental organizations, and all stakeholders

### **National Protection**

National legislations in the Caribbean region applying to sharks (as reviewed by Van Beek *et al.*, 2014) are as follows:

#### ***IPOA Sharks***

There are since the 1990s several shark protection plans, both internationally at intergovernmental and non- governmental level, as well as at national level by several nations in the Wider Caribbean region. Within the framework of the Code of Conduct for Responsible Fisheries the FAO (Food and Agriculture Organization) developed the International Plan of Action for the Conservation and Management of Sharks (IPOA Sharks) in 1999. The objective of IPOA Sharks is to ensure the conservation and management of sharks and their long-term sustainable use. IPOA Sharks is voluntary and intends to give states guidelines on how to establish a National Plan of Action (NPOA) through guiding principles and procedures for implementation. Caribbean countries with an IPOA are: Antigua and Barbuda (in draft), Costa Rica (2010 – not official), Bolivarian Republic of Venezuela (2013 – not official; FAO, 2016).

### **National Legislation**

#### **USA**

In the U.S., *S. lewini*, *S. mokarran* and *S. zygaena* are included in the Large Coastal Shark complex management unit, on U.S. Highly Migratory Species Fishery Management Plan (National Marine Fisheries Service: Federal Fisheries Management Plan for Atlantic Tuna, Swordfish and Sharks). There are, however, no management measures specific to this species, and no stock assessments.

#### ***US Caribbean Region***

NOAA fisheries service presented the amendment 4 to the 2006 Consolidated Atlantic Highly Migratory Species (HMS) Fishery Management Plan (FMP). The powerpoint states that “in 2010, Puerto Rico reported approximately 11.8 mt of commercial shark landings and less than one megaton was reported by St. Thomas and St. John combined. These landings were not species specific and it is unknown if they were harvested from Federal or Territorial waters”. Proposed management measures for small-scale HMS commercial fisheries include specific authorized gears and retention limits for sharks.

#### ***Kingdom of the Netherlands***

— ***St. Maarten***

On the 12th of October 2011 the government of St. Maarten issued a temporary moratorium on shark fishing. The shark fishing moratorium prohibits the take and landing of sharks and requires immediate release of incidentally caught sharks, under penalty of a maximum of 500,000 Antillean Guilders or 3 months in prison.

— ***Caribbean Netherlands***

In 2015, the Dutch government designated the Yarari sanctuary for sharks and marine mammals in the Economic Exclusive Zones of Saba and Bonaire, declaring that provisions will be considered and implemented as necessary to regulate activities that may have a negative impact on sharks.

— ***Bonaire***

In 2008 the island of Bonaire passed a nature ordinance providing full protection for a list of species of plants and animals. This list includes all sharks and rays.

***US Gulf of Mexico and (Caribbean) Florida***

Following years of declines in catches, and concern about the protection status of many shark species, in 1993 the USA established a Federal Management Plan for Shark Fisheries in the Atlantic Ocean, particularly directed at the coastal bottom long-line fishery. Since 1993 several amendments of the original plan have been implemented and local state governments have tied in by implementing complementary legislation. Measures included successively restrictive catch quotas, finning limitations, area closures, seasonal closures, adjustments of size limits, limits to retention in recreational fisheries, establishment of protected species lists, establish a shark research fishery and the use of regional and species specific quotas.

***Honduras***

In June 2011 Honduras created the first shark sanctuary in America and declared all its marine waters in both the Pacific and Caribbean as a permanent shark sanctuary. This had been preceded in 2010 by a shark fishing moratorium and created the first shark sanctuary of the Americas amounting to about 240,000 km<sup>2</sup> of national waters, most of which lie along the 700 km-long Caribbean coast of the nation.

***Bahamas***

The Bahamas have had a longline fishing ban since 1993 and consequently there has been no commercial shark fishing activity. This longline ban has effectively made the whole archipelago of the Bahamas a shark “no-take” zone. The last export of shark from the Bahamas was a lot of 2 metric tons in 2004. In July 2011 the Bahamas went a step further and legally banned all shark fishing. That law firmly turns all 630,000 sq km of Bahamian waters into a shark sanctuary<sup>17</sup>. The fines for shark fishing were raised from 3000 to 5000 USD per incident.

***Venezuela***

Towards implementing its Plan de Acción Nacional (PAN) de conservación for sharks, in June 2012 Venezuela joined the rest of the Americas in outlawing the finning of sharks in its waters and established a 3,730 km<sup>2</sup> shark sanctuary surrounding the touristic archipelago of Los Roques. Recent research (e.g. Tavares 2005, 2008 2009) had demonstrated the importance of the shallow waters of Los Roques as a shark nursery area.

The Dominican Republic has, together with Belize and six other Central American countries, united under the name SICA (Central American Integration System), signed an agreement to

prohibit shark finning. This ban is also applicable to fishing vessels in international waters under the flag of SICA member states. This arrangement OSP-05-11 entered into force in 1 January 2012.

#### **d. Ecological interactions with other species and specific habitat requirements**

The diet of *Sphyrna mokarran* includes fish (mainly demersal species), other elasmobranchs, crustacea and cephalopods (Compagno in prep. b). Strong *et al.* (1990) observed a large (*ca* 4 m) Great Hammerhead feeding on a southern stingray *Dasyatis americana* (disc width 1.5 m). Adult *S. lewini* feed on mesopelagic fish and squids. In certain areas stingrays of the *Dasyatis* family are the preferred food. Pups and juveniles feed mainly on benthic reef fishes (e.g., scarids and gobiids), demersal fish and crustaceans. (Baum *et al.*, 2007). For *S. zygaena* less than 2 m in length from the waters off South Africa, Smale (1991) reported that the diet was dominated by inshore squid (mostly *Loligo v. reynaudii*), with teleosts such as hake, horse mackerel and ribbonfish also being important. Crustaceans and elasmobranchs have also been reported from stomach analyses (Bass *et al.*, 1975; Compagno, 1984; Smale 1991; Last and Stevens, 1994).

Sharks and rays are often predators feeding at a high trophic level and are therefore thought to exert a significant top-down control over the ecosystem. Both empirical studies and ecosystem modeling studies demonstrated that the decline of large coastal elasmobranch species could induce a trophic cascade, as well as decreased ecosystem functioning and resilience. Because of their large size they occupy ecological niches first occupied by large predatory reptilians and have likely played a critical role in the evolution of marine mammals as well as other predators and prey species (Ferretti *et al.*, 2010). Sharks are largely seen as feeding generalists and typically take a wide range of prey and therefore likely have limited effect on mortality rates in individual species (Ellis and Musick, 2007). They are typically wide ranging and interconnect food webs across wide geographic ranges (Musick *et al.*, 2000). The ecological role each species can play in this is likely influenced by their distribution across habitats. Most shark species (90%) are restricted to near-shore waters of the continental shelves whereas some species (e.g., hammerhead, tiger shark) migrate between the pelagic and near-shore habitats and only few are fully pelagic in habits.

We know very little about the specific roles of sharks in Caribbean coral reef ecosystems, and hammerheads are no exception, but current models and theories suggest that their loss causes multiple effects throughout local food webs and could lead to reef collapse. A study by Rezende *et al.* (2009) highlighted the importance of sharks for the organization, and potentially also for the stability and biodiversity of the Caribbean food webs. Modelling suggests that sharks are important regulators of grouper biomass on Caribbean reefs (Bascompte *et al.*, 2005) and potentially important for the biological control of the invasive lionfish *Pterois volitans* (Albins and Hixon, 2008; Arias-Gonzalez *et al.*, 2011). Other work suggests the role of sharks in regulating grouper biomass has an indirect positive effect on parrotfish biomass and grazing capacity (Chapman *et al.*, 2006). The model of Arias-Gonzalez *et al.* (2011) predicts that lionfish will replace sharks as apex predators as a result of a decrease in sharks due to overfishing throughout the region. The ecological effects of loss of sharks as top predators is difficult to understand and generally obscured by the fact that ecosystems have simultaneously been undergoing many other major changes. *S. lewini* is a high trophic level predator in coastal and open ocean ecosystems. It has a diverse diet,

feeding on crustaceans, teleosts, cephalopods and rays (Compagno, 1984). An analysis of its stomach contents revealed that the males feed on 42% of *Ancistrocheirus lesueurii* (Orbigny 1842), a species of mesopelagic cephalopod (Klimley, 1987). On the other hand, females consumed 63% mesopelagic squid species, *Mastigoteuthis* sp and *Moroteuthis robusta* (Verrill, 1876). Cortés (1999) determined the trophic level to be 4.1 (maximum=5.0) for *S. lewini*, based on diet information. Navia *et al.* (2010) propose that this is the second most topologically important species for the maintenance of the structure of the community in the central fishing zone in the Colombia Pacific.

### **e. Management and recovery plans for the species**

See Van Beek *et al.* (2014) for a complete overview of Dutch national management and recovery plans.

There is a management plan in place in US waters, regulating catches from fishing and the scalloped hammerhead (*Sphyrna lewini*) population has stabilized since the plan was put into place in 1994 (Hayes *et al.*, 2009). Scalloped hammerheads, which are among the faster growing species in the complex, have a relatively high probability of recovering quickly. Despite its slow life history characteristics, this scalloped hammerhead population appears to have a 58% or greater probability of recovery within a decade if the 2005 catch is maintained or decreased (Hayes *et al.*, 2009).

### **f. Research programs and available scientific and technical publications relevant to the species**

The biology of elasmobranchs is among the most poorly known and least understood of all the marine vertebrate groups (Fowler, 2005). This is particularly so for the Caribbean region. Within the Caribbean Sea, research efforts are underway to assess distribution, habitat use, population structure, and trophic ecology of sharks using acoustic telemetry, satellite tagging methods, genetic analysis and stable isotope research. The occurrence and relative abundance of sharks are investigated using Baited Remote Underwater Videos (BRUVs). The skillful use of modern techniques such as genetic analyses, telemetry, and Baited Remote Video monitoring can help circumvent the often-low abundance (and low sampling) of many species, and should help develop powerful new insights and introduce new techniques to the region where capacity and technology have lagged behind.

### **g. Threats to the species, its habitats and associated ecosystems, especially threats which originate outside the jurisdiction of the Party.**

Baum *et al.* (2003) have shown a decline of 89% of hammerheads (primarily scalloped hammerheads (*Sphyrna lewini*)) in the northwestern Atlantic, including the Caribbean between 1986 and 2000.

For pelagic species, fishing is identified as the main threat, which is corroborated by studies that have demonstrated the extent of overfishing of large predators in the Caribbean (e.g. Bonfil, 1997; Stallings, 2009; Pandolfi *et al.*, 2003). Pelagic sharks are all found to be declining, albeit at different rates (Cortés *et al.*, 2007; Baum and Blanchard, 2010). A decadal dataset (1994–2003) of the Venezuelan longline fisheries recorded (by order of

importance) landings of the Blue Shark (*Prionace glauca*), Night shark (*C. signatus*), Silky Shark, Great Hammerhead (*Sphyrna mokarran*), and the Shortfin Mako (*Isurus oxyrinchus*; Tavares and Arocha, 2008). A study after bycatch rates of the Venezuelan longline fleet showed a major bycatch of great, and smooth hammerhead (Arocha *et al.*, 2002).

Due to the distinctive head shape of this genus, it is typical for catches to be reported at the genus level, *Sphyrna* spp. Therefore, it is rare to find fisheries statistics that are specific to one species of hammerhead shark. Species identification (*S. mokarran* vs. *S. lewini*) is a large obstacle in the proper assessment of this species. Catches of Sphyrnidae have been reported only from the Atlantic Ocean since 1991 and these landings are undoubtedly under-reported. The catch was near 2,200 tons in 2004 (Maguire *et al.*, 2006). Only *S. zygaena* and *S. lewini* are reported as individual species in the Food and Agriculture Organisation (FAO) fisheries statistics, but hammerhead catches are often grouped in one category as, *Sphyrna* species, which makes identification of actual catches of *S. zygaena* difficult. The high at-vessel fishing mortality for hammerheads makes the threat of fishing even greater for these species. This species' fins are highly valued and they are being increasingly targeted in some areas in response to increasing demand for shark fins. Hammerhead shark species *S. zygaena* and *S. lewini* were found to represent at least 4-5% of the fins auctioned in Hong Kong, the world's largest shark fin trading center (Clarke *et al.*, 2006). Fins from the Hong Kong SAR market can be genetically assessed and have been shown to originate western Atlantic Ocean basins. In a study by Chapman *et al.* (2009) approximately 21% of the samples were sourced from the western Atlantic.

Hammerhead shark fins are generally high value compared to other species because of their high fin ray count (*S. Clarke unpubl. data*). It is estimated that between 1.3 and 2.7 million *S. zygaena* or *S. lewini* are represented in the shark fin trade each year or, in biomass, 49,000 to 90,000 mt (Clarke *et al.*, 2006). Longline fleets exert intense fishing pressure throughout the Northwest Atlantic (Baum *et al.*, 2003). Baum *et al.* (2003) estimated that hammerhead sharks (grouped data for *S. lewini*, *S. mokarran* and *S. zygaena*) have declined in abundance by 89% since 1986 (95% confidence interval (CI): 86 to 91%) in their analysis of U.S. pelagic longline logbook data. This group is primarily composed of *S. lewini*; in Virginia Institute of Marine Science sampling programs since 1973, *S. lewini* outnumbered *S. zygaena* by more than ten to one (Ha, 2006).

Recent research shows that large, oceanic sharks may actually depend on shallow coastal areas during part of their life cycle (e.g. Carrier and Pratt, 1998; Tavares, 2008; Clarke *et al.*, 2011; Daly-Engel *et al.*, 2012; Hammerschlag *et al.*, 2012). This makes many sharks vulnerable to habitat destruction in coastal areas, as caused by man (Jennings *et al.*, 2008) and possibly, on the long-term by climate change (Field *et al.*, 2009). The dependence of sharks on habitat quality has hardly been studied so far (Field *et al.*, 2009). One important dimension of habitat quality is that of food availability. Sharks are potentially affected by shortage of prey due to competition for the same resources by their largest piscivorous competitor, namely man, but this has received even less attention.

### ***S. lewini***

*Sphyrna lewini* is taken as both a target and bycatch by trawls, purse seines, gillnets, fixed bottom longlines, pelagic longlines and inshore artisanal fisheries. The latter catch large numbers of pups and juveniles in some regions. The species' aggregating habit makes them

vulnerable to capture in large schools. This also means that they may appear more abundant in landings, where they are caught in high, localized concentrations. Intense fishing pressure can deplete regional stocks rapidly, and re-colonization of depleted areas from neighboring regions is expected to be a slow and complex process. This species is expected to have a low resilience to exploitation because of its life-history characteristics. Also, the aggregating habit of *S. lewini* makes it very vulnerable to capture. In the nursery zones (<10 m) south and southeast of Brazil the newborn are intensively fished through coast gillnets, prawn trawls and pair trawls, as well as recreational capture (Haimovici & Mendonça, 1996; Kotas 2004; Kotas *et al.*, 2005; Vooren *et al.*, 2005).

In the USA, this species is caught in both commercial coastal shark bottom longline and gillnet fisheries and the pelagic longline fishery, where it suffers high mortality (Piercy *et al.*, 2007). It is also taken in recreational shark fisheries. The USA pelagic longline fishery has operated since the 1960s and encompasses the entire range of this species in the Northwest and Western Central Atlantic, from the equator to about 50°N. Although this is quite a fecund shark, its late age at maturity in this region (15 years) will render it quite vulnerable to overexploitation, and limit its recovery potential.

Estimates of trends in abundance of *Sphyrna* spp. are available from standardized catch rate indices of the U.S.A. pelagic longline fishery, from logbook data between 1986 and 2000 and from observer data between 1992 and 2005. The area covered by this fishery, ranging from the equator to about 50°N, encompasses the range of this species in these two regions. Although this fishery will not sample individuals closest to the coast, the sample size of hammerheads recorded in the logbook data (the majority of which are thought to be *S. lewini*) is substantial, with over 60,000 recorded during this period. This subpopulation of Scalloped Hammerhead sharks is estimated from the logbook data to have declined by 89% over the 15 year time period, from 1986-2000 (Baum *et al.*, 2003), which is less than one generation. A more recent analysis of the pelagic longline observer data indicates that *Sphyrna* spp. declined by 76% between 1992 and 2005 (Baum *et al.*, in prep.). The pelagic longline fishery has operated in these regions since the 1960s, thus declines from 1986 were certainly not from virgin population abundance.

Off the Atlantic coast of Belize hammerheads were fished heavily by longline in the 1980s and early 1990s (R.T. Graham, pers. obs.). Hammerheads are a favored target species for their large fins. Interviews with fishermen indicate that the abundance and size of Sphyrnids has declined dramatically in the past 10 years as a result of over exploitation, leading to a halt in the Belize based shark fishery (R.T. Graham, pers. obs.). However, the pressure is still sustained by fishers driving into Belizean waters from Guatemala (R.T. Graham, pers. obs.). *Sphyrna lewini* is also taken in various fisheries along the Caribbean coast of South America. It is taken in artisanal gillnet fisheries targeting mackerel off Guyana, Trinidad and Tobago and in pelagic tuna fisheries of the eastern Caribbean (Chan A Shing, 1999).

### ***S. mokarran***

*Sphyrna mokarran* is taken by target and bycatch, fisheries (Dudley and Simpfendorfer, 2006; Zeeberg *et al.*, 2006) and is regularly caught in the Caribbean, with longlines, fixed bottom nets, hook-and-line, and possibly with pelagic and bottom trawls (Compagno, *in prep*). Hammerhead sharks, with *S. mokarran* in particular, have been noted as a favored target species due to the size of their fins (R.T. Graham, pers. comm). Fin prices are rising

above US\$50/lb in the neighboring countries of Guatemala, driven by Asian buyers, according to interviews (R.T. Graham, pers. obs). Bonfil (1994) gives an overview of global shark fisheries. This species is mentioned specifically with reference to fisheries in Brazil, East USA and Mexico, however *Sphyrna* spp. are mentioned in the majority of tropical fisheries cited.

This species is caught primarily as a bycatch in the pelagic longline, bottom longline and net fisheries along the northwest Atlantic and Gulf of Mexico. It is also caught in the recreational fishery. The species represents 0.7% of the species catch and suffers from greater than 90% at-vessel fishing mortality in the U.S. bottom longline fishery (Commercial Shark Fishery Observer Program unpubl. data). The U.S. pelagic fishery logbook data has shown a decline close to 90%, however this dataset is known for inaccurate data reporting (Beerkircher *et al.*, 2002). There is probably a lack of reporting of the catch of Great Hammerheads because this species is routinely finned and discarded, which is illegal in the US Atlantic Federal Waters (Commercial Shark Fishery Observer Program, unpub. data). Both the pelagic and bottom longline observer programs have recorded a 2 to 3:1 ratio for *S. Lewini* to *S. mokarran*. The meat is not valuable but the fins are high grade and bring in a good price, thus finning still occurs in the U.S. fishery.

There appear to be little data for landings and catch effort for this species in Central America and the Caribbean. Off the coast of Belize hammerheads were fished heavily by longline in the 1980s and early 1990s. Interviews with fishermen indicate that the abundance and size of Sphyrnids has declined dramatically in the past 10 years as a result of over exploitation, leading to a halt in the Belize based shark fishery (R.T. Graham, pers. obs). However, the pressure is still sustained by fishers driving into Belizean waters from Guatemala (R.T. Graham pers. obs). The Cuban directed shark fishery (longline) recorded between 1983 and 1991 *S. mokarran* (subadults and juveniles) as one of 23 species caught. Since 1992 small increases in mean sizes were noted, indicating partial recovery of the species. In Mexico between November 1993 and December 1994 (Tamaulipas, Veracruz, Tabasco, Campeche and Yucatan) 901 vessels were monitored every day. *Sphyrna mokarran* represented 86% of the total catch.

### ***S. zygaena***

*Sphyrna zygaena* is caught with a variety of gears, including with pelagic longlines, handlines, gillnets, purse seines and pelagic and bottom trawls (Bonfil, 1994; Compagno in prep; Maguire *et al.*, 2006). This shark is undoubtedly caught in shark fisheries in most parts of its range, but it is not always reported separately from other hammerhead species. Bonfil (1994) reported that this species is caught as bycatch in a number of non-shark fisheries, particularly pelagic longline and gillnet fisheries that operate close to temperate and subtropical continental shelves. The capture of *S. zygaena* in many of these fisheries is infrequent (Bonfil, 1994). Although size data are limited, catches in pelagic fisheries appear to be dominated by larger individuals, while juveniles are common in inshore shelf fisheries.

## **References**

- Alava, M.N.R., Dolumbaló, E.R.Z., Yaptinchay, A.A. and Trono, R.B. 2002. Fishery and trade of Whale Sharks and Manta Rays in the Bohol Sea, Philippines. Pp. 132-148. In: S.L. Fowler, T.M. Reed and F.A. Dipper (eds). *Elasmobranch Biodiversity, Conservation and Management: Proceedings of the International Seminar and Workshop*. Sabah, Malaysia, July 1997. Occasional paper of the IUCN Species Survival Commission No. 25.
- Anderson, R., Adam, M., Kitchen-Wheeler, A. and Stevens, G. 2010. Extent and economic value of Manta Ray watching in Maldives. *Tourism in Marine Environments* 7: 15-27.
- Anderson, R.C. and Ahmed, H. 1993. *The shark fisheries in the Maldives*. FAO, Rome, and Ministry of Fisheries, Male, Maldives.
- Anderson, R.C., Adam, M.S., and Goes, J.I. 2011. From monsoons to mantas: seasonal distribution of Manta alfredi in the Maldives. *Fisheries Oceanography*, 20(2): 104-113.
- Anon. 1997. Fisheries conservation crisis in Indonesia: Massive destruction of marine mammals, sea turtles and fish reported from trap nets in pelagic migratory channel. <http://darwin.bio.uci.edu/~sustain/bio65/indonesia/indon97e.htm>.
- Backus, R.H., Springer, S. and Arnold Jr., E.L. 1956. A contribution to the natural history of the white-tip shark, *Pterolamiops longimanus* (Poey). *Deep-Sea Research* 3: 176–188.
- Baum, J., Medina, E., Musick, J.A. & Smale, M. 2015. *Carcharhinus longimanus*. The IUCN Red List of Threatened Species 2015: e.T39374A85699641. <http://dx.doi.org/10.2305/IUCN.UK.2015.RLTS.T39374A85699641.en>.
- Baum, J.K. and Myers, R.A. 2004. Shifting baselines and the decline of pelagic sharks in the Gulf of Mexico. *Ecology Letters*. 7(3): 135–145.
- Baum, J.K., Kehler, D. and Myers, R.A. 2005. Robust estimates of decline for pelagic shark populations in the northwest Atlantic and Gulf of Mexico. *Fisheries* 30: 27–30.
- Beerkircher, L.R., Cortés, E., and Shivji, M. 2002. Characteristics of Shark Bycatch Observed on Pelagic Longlines off the Southeastern United States, 1992–2000. *Marine Fisheries Review*, 64(4) 40–49.
- Berkeley, S.A. and Campos, W.L. 1988. Relative abundance and fishery potential of pelagic sharks along Florida's east coast. *Mar. Fish. Rev.* 50(1): 9–16.
- Bigelow, H.B. and Schroeder, W.C. 1953. Sawfish, guitarfish, skates and rays. In: Tee-Van (ed.), *Fishes of the Western North Atlantic, Part 2*. Sears Foundation for Marine Research, Yale University, New Haven, pp 508-514.
- Bullis H.R.J. and Captiva, F.J. 1955. Preliminary report on exploratory long-line fishing for tuna in the Gulf of Mexico and the Caribbean sea. *Commercial Fisheries Review* 17: 1–20.
- Bullis, H.R.J. 1961. Observations on the feeding behavior of white-tip sharks on schooling fishes. *Ecology* 42: 194–195.
- Burgess, G.H., Beerkircher, L.W., Cailliet, G.M., Carlson, J.K., Cortes, E., Goldman, K.J., Grubbs, R.D., Musick, J.A., Musyl, M.K. and Simpfendorfer, C.A. 2005. Is the collapse of shark populations in the northwest Atlantic Ocean and Gulf of Mexico real? *Fisheries* 30: 10.

- Caldas, J.P. and Correa, J.L. 2010. Captura de tiburones asociada a la pesca industrial com palangre oceânico en el mar Caribe colombiano. Libro de Resúmenes II Encuentro de Colombiano sobre Condrictios. Cali, Colombia. P 35.
- Carlson, J.K. and Osborne, J. 2012. Relative abundance of smalltooth Sawfish (*Pristis pectinata*) based on the Everglades National Park Creel Survey. NOAA Technical Memorandum NMFS-SEFSC-626, 15 p.
- Castro, J.I. 2000. The biology of the nurse shark, *Ginglymostoma cirratum*, off the Florida east coast and the Bahama Islands. *Environmental Biology of Fishes* 58(1): 1-22.
- Chapman, D.D., Simpfendorfer, C.A., Wiley, T.R., Poulakis, G.R., Tringali, M., Carlson, J.K., and Feldheim, K.A. 2011. Genetic diversity despite population collapse in a critically endangered marine fish: the Smalltooth Sawfish (*Pristis pectinata*). *Journal of Heredity* 102(6): 643-652.
- Charvet-Almeida, P. 2002. Sawfish trade in the North of Brazil. *Shark News* 14: 9. Newsletter of the IUCN Shark Specialist Group. Naturebureau, Newbury, UK.
- Chen, C.T., Liu, K.M. and Joung, S.J. 1997. Preliminary report on Taiwan's Whale Shark fishery. *TRAFFIC Bulletin* 17(1): 53-57.
- Clarke, S. 2008. Estimating historic shark removals in the Atlantic using shark fin trade data and Atlantic specific area, tuna catch and effort scaling factors. *ICCAT SCRS* 2008/139.
- Colman, J.G. 1997. A review of the biology and ecology of the Whale Shark. *Journal of Fish Biology* 51:1219-1234.
- Compagno L.J.V. (2001) *Sharks of the World: An Annotated and Illustrated Catalogue of Shark Species Known to Date. Volume 2 Bullhead, Mackerel and Carpet Sharks (Heterodontiformes, Lamniformes and Orectolobiformes)*. Food and Agriculture Organization of the United Nations, species catalogue for fishery purposes, Rome, Italy.
- Compagno, L.J.V. 1984. *FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. FAO Fisheries Synopsis No. 125, Volume 4, Part 1.*
- Compagno, L.J.V. 1984. *Sharks of the World. An annotated and illustrated catalogue of shark species to date. Part I (Hexanchiformes to Lamniformes)*. FAO Fisheries Synopsis, FAO, Rome.
- Compagno, L.J.V. 2005. Checklist of living Chondrichthyes. In: Hamlett, W.C. (ed.) *Reproductive biology and phylogeny of chondrichthyes: sharks, rays and chimaeras*, Vol. 3. Endfield, USA: Science Publishers: 503-548.
- Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES). 2007. Proposal 17 Inclusion of all species of the family Pristidae in Appendix I of CITES. Fourteenth meeting of the Conference of the Parties The Hague (Netherlands), 3-15 June 2007. <http://www.cites.org/eng/cop/14/prop/e14-p17.pdf>.
- Cortés, E. 1999. Standardized diet compositions and trophic levels of sharks. *ICES Journal of Marine Science* 56:707-17

- Cortés, E., C. Brown, and L. R. Beerkircher. 2007. Relative abundance of pelagic sharks in the western North Atlantic Ocean, including the Gulf of Mexico and Caribbean Sea. *Gulf Caribbean Research Report* 19: 37-52.
- Couturier, L.I.E., Marshall, A.D., Jaine, F.R.A., Kashiwagi, T., Pierce, S.J., Townsend, K.A., Weeks, S.J., Bennet, M.B., and Richardson, A.J. 2012. Biology, ecology and conservation of the Mobulidae. *Journal of Fish Biology*, 80: 1075-1119.
- De la Parra Venegas, R., R. Hueter, J. González Cano, J. Tyminski, J. Gregorio Remolina, M. Maslanka, A. Ormos, L. Weigt, B. Carlson and A. Dove, 2011. An unprecedented aggregation of Whale Sharks, *Rhincodon typus*, in Mexican coastal waters of the Caribbean Sea. *PLoS ONE* 6(4):e18994.
- Deakos, M., Baker, J., and Bejder, L. 2011. Characteristics of a Manta Ray (*Manta alfredi*) population off Maui, Hawaii, and implications for management. *Marine Ecology Progress Series*, 429: 245-260.
- Debrot, A.O., Leon de, R., Esteban, N., Meesters, H.W.G. (2013) Observations on the Whale Shark (*Rhincodon typus*) in the Dutch Caribbean. *Caribbean Journal of Science*, Vol. 47 [2-3]: 344-349.
- Dewar, H. 2002. Preliminary report: Manta harvest in Lamakera. Report from the Pflieger Institute of Environmental Research and the Nature Conservancy.
- Dewar, H., Mous, P., Domeier, M., Muljadi, A., Pet, J., Whitty, J. 2008. Movements and site fidelity of the giant Manta Ray, *Manta birostris*, in the Komodo Marine Park, Indonesia. *Marine Biology*, 155, (2); 121-133.
- Domingo, A. 2004. ¿Adónde fue el Longimanus?. *Elasmovisor*, Bol. SBEEL, July, Brazil, 6pp.
- Doumbouya, F. 2004. Rapport sur la preparation du PAN-Requin Guinéen et le dossier de la raie *Pristis*. Report to the Sub-Regional Fisheries Commission from the Département Pêche Artisanale, Ministère de la Pêche et de l'Aquaculture, Conakry, Republic de Guinée.
- Duffy, C.A.J. and Abbott, D. 2003. Sightings of mobulid rays from northern New Zealand, with confirmation of the occurrence of *Manta birostris* in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* 37: 715-721.
- Dulvy, N.K, Fowler, S.L., Musick, J.A., Cavanagh, R.D., Kyne, P.M., Harrison, L.R., et al. (2014) Extinction risk and conservation of the world's sharks and rays. *eLife* 2014;3:e00590
- Eckert, S.A. and B.S. Stewart. 2001. Telemetry and satellite tracking of Whale Sharks, *Rhincodon typus*, in the Sea of Cortez, Mexico, and the north Pacific Ocean. *Environmental Biology of Fishes* 60: 299-308.
- Faria, V.V., McDavitt, M.T., Charvet, P., Wiley, T.R., Simpfendorfer, C.A. and Naylor, G.J.P. 2013. Species delineation and global population structure of Critically Endangered Sawfishes (*Pristidae*). *Zoological Journal of the Linnean Society* 167: 136-164.
- Fourmanoir, P. 1961. Requins de la Côte Ouest de Madagascar. *Memoires de L'Institut Scientifique de Madagascar. Série F. Oceanographie. ORSTOM. Tome IV.*

Fowler, S.L. 2000. Whale shark *Rhincodon typus*. Policy and research scoping study. Nature Conservation Bureau, Newbury, UK.

Fowler, S.L., Cavanagh, R.D., Camhi, M., Burgess, G.H., Cailliet, G.M., Fordham, S.V., Simpfendorfer, C.A. and Musick, J.A. (comp. and ed.). 2005. Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes. Status Survey. IUCN/ SSC Shark Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK. x + 461 pp.

Graham, R.T. 2007. Behaviour and conservation of Whale Sharks on the Belize Barrier Reef

Graham, R.T. and Roberts, C.M. 2007. Assessing the size, growth rate and structure of a seasonal population of whale sharks (*Rhincodon typus* Smith 1828) using conventional tagging and photo identification. *Fisheries Research* 84: 71-80.

Graham, R.T., Hickerson, E., Castellanos, D.W., Remolina, F., Maxwell, S. 2012. Satellite Tracking of Manta Rays Highlights Challenges to Their Conservation. *PLoS ONE* 7(5): e36834. Doi:10.1371/journal.pone.0036834

Heyman, W., Graham, R., Kjerfve, B. and Johannes, R.E. 2001. Whale sharks *Rhincodon typus* aggregate to feed on fish spawn in Belize. *Marine Ecology Progress Series* 215: 275–282.

Hoffmayer, E. R., J. S. Franks, and J. P. Shelley. 2005. Recent Observations of the Whale Shark (*Rhincodon typus*) in the northcentral Gulf of Mexico. *Gulf and Caribbean Research* 17: 1-4.

Homma, K., Maruyama, T., Itoh, T., Ishihara, H., and Uchida, S. 1999. Biology of the Manta Ray, *Manta birostris* Walbaum, in the Indo-Pacific. In: Seret, B. and Sire, J.Y. (eds) *Indo-Pacific fish biology: Proc 5th Int Conf Indo-Pacific Fishes*, Noumea, 1997. Ichthyological Society of France, Paris, p 209–216.

Howey-Jordan LA, Brooks EJ, Abercrombie DL, Jordan LKB, Brooks A, Williams S et al (2013) Complex movements, philopatry and expanded depth range of a severely threatened pelagic shark, the oceanic whitetip (*Carcharhinus longimanus*) in the Western North Atlantic. *PLoS One* 8(2):e56588.

Hueter RE, Tyminski JP, de la Parra R (2013) Correction: Horizontal Movements, Migration Patterns, and Population Structure of Whale Sharks in the Gulf of Mexico and Northwestern Caribbean Sea. doi: info:doi/10.1371/annotation/491b9b6c-7f77-4fbo-b336-572078aec830

Joung, S.J., C.T. Chen, C. Eugenie, S. Uchida and W.Y.P. Huang. 1996. The Whale Shark *Rhincodon typus* is a livebearer: 300 embryos found in one 'megamma' supreme. *Environmental Biology of Fishes* 46: 219–223.

Kashiwagi, T., Marshall, A.D., Bennett, M.B. and Ovenden, J.R. 2011. Habitat segregation and mosaic sympatry of the two species of Manta Ray in the Indian and Pacific Oceans, *Manta alfredi* and *M. birostris*. *Marine Biodiversity Records* 4: e53.

Kitchen-Wheeler, A. 2010. Visual identification of individual Manta Ray (*Manta alfredi*) in the Maldives Islands, Western Indian Ocean. *Marine Biology Research*, 6(4):351-363

- Last, P.R. and Stevens, J.D. 2009. *Sharks and Rays of Australia*. Second Edition. CSIRO Publishing, Collingwood.
- Leu, M.Y., Chang, W.B. and Fang, L.S. 1997. The success of keeping a baby Whale Shark from its fetal stage in Taiwan. Proceedings of the Fourth International Aquarium Congress, Tokyo, Japan, June 1996.
- Luiz, O.J. Jr, Balboni, A.P., Kodja, G., Andrade, M. and Marum, H. 2009. Seasonal occurrences of *Manta birostris* (Chondrichthyes: Mobulidae) in southeastern Brasil. *Ichthyological Research* 56: 96-99.
- Marshall, A., Bennett, M.B., Kodja, G., Hinojosa-Alvarez, S., Galvan-Magana, F., Harding, M., Stevens, G. & Kashiwagi, T. 2011a. *Manta birostris*. The IUCN Red List of Threatened Species 2011: e.T198921A9108067.
- Marshall, A.D., Compagno, L.J.V. and Bennett, M.B. 2009. Redescription of the genus *Manta* with resurrection of *Manta alfredi* (Krefft, 1868) (Chondrichthyes; Myliobatoidei; Mobulidae). *Zootaxa* 2301: 1-28.
- Marshall, A.D., Dudgeon, C. and Bennett, M.B. 2011. Size and structure of a photographically identified population of *Manta Rays Manta alfredi* in southern Mozambique. *Marine Biology* 158(5): 1111-1124.
- Mather, F.J.I. and Day, C.G. 1954. Observations of pelagic fishes of the tropical Atlantic. *Copeia* 1954: 179-188.
- McClenachan, L. 2009. Documenting loss of large trophy fish from the Florida Keys with historical photographs. *Conservation Biology* 23 (3), 636-643.
- McDavitt, M. 1996. The cultural and economic importance of Sawfishes (family Pristidae). *Shark News* 8: 10-11.
- McDavitt, M. T. 2002. Lake Nicaragua revisited: conversations with a former Sawfish fisherman. *Shark News* 14: 5. Newsletter of the Shark Specialist Group. Naturebureau, Newbury, UK.
- Migdalski, E.C. 1981. *Fish Mounts and Other Fish Trophies*. John Wiley & Sons, New York.
- Millessi, A.C. and Oddone, M.C. 2003. Primer registro de *Manta birostris* (Donndurff 1798) (Batouidea: Mobulidae) En el Rio de la Plata, Uruguay. *Gayana* 67: 127-130.
- National Marine Fisheries Service (NMFS). 2009. Recovery Plan for Smalltooth Sawfish (*Pristis pectinata*). Prepared by the Smalltooth Sawfish Recovery Team for the National Marine Fisheries Service. Silver Spring.
- Norman, B.M. 1999. Aspects of the biology and ecotourism industry of the Whale Shark *Rhincodon typus* in north-western Australia. MPhil. Thesis (Murdoch University, Western Australia).
- Norman, B.M., Holmberg, J.A., Arzoumanian, Z., Reynolds, S., Wilson, R.P., Gleiss, A.C., Rob, D., Pierce, S.J., de la Parra, R., Galvan, B., Ramirez-Macias, D., Robinson, D., Fox, S., Graham, R., Rowat, D., Potenski, M., Levine, M., Mckinney, J.A., Hoffmayer, E., Dove, A.,

Hueter, R., Ponzo, A., Araujo, G., Aca, E., David, D., Rees, R., Duncan, A., Rohner, C.A., Hearn, A., Acuna, D., Berumen, M.L., Vazquez, A., Green, J., Bach, S.S., Schmidt, J.V. and Morgan, D.L. Submitted. Understanding constellations: 'citizen scientists' elucidate the global biology of a threatened marine mega-vertebrate. *Bioscience*.

Osada, K. 2010. Relationship of zooplankton emergence, Manta Ray abundance and SCUBA diver usage Kona, Hawaii. Masters Thesis, University of Hawaii.

Pauly, D. 2002. Growth and mortality of the basking shark *Cetorhinus maximus* and their implications for management of Whale Sharks *Rhincodon typus*. In: S.L. Fowler, T.M. Reed and F.A. Dipper (eds), *Elasmobranch Biodiversity, Conservation and Management*. Proceedings of the International Seminar and Workshop, Sabah, Malaysia, July 1997, pp. 199–208. IUCN SSC Shark Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.

Pierce, S.J. & Norman, B. 2016. *Rhincodon typus*. The IUCN Red List of Threatened Species 2016: e.T19488A2365291. Downloaded on 04 July 2016.

Ramírez-Macías, D., Meekan, M., de la Parra-Venegas, R., Remolina-Suárez, F., Trigo-Mendoza, M. and Vázquez-Juárez, R. 2012. Patterns in composition, abundance and scarring of whale sharks *Rhincodon typus* near Holbox Island, Mexico. *Journal of Fish Biology* 80: 1401-1416.

Romero, A., A. I. Agudo, S. M. Green, and G. Notarbartolo di Sciara. 2001. Cetaceans of Venezuela: Their distribution and conservation status. *NOAA Technical Reports, NMFS* 151:1-60.

Santana, J.C., A., Molina de Delgado, R., Molinda de Delgado, J. Ariz, J.M. Stretta, and G. Domalain. 1997. Lista faunística de las especies asociadas a las capturas de atún de las flotas de cerco comunitarias que faenan en las zonas tropicales de los océanos Atlántico e Índico. *ICCAT Collective Volume of Scientific Papers* 48:129-137.

Seitz, J.C. & G.R. Poulakis. 2006. Anthropogenic effects on the Smalltooth Sawfish (*Pristis pectinata*) in the United States. *Marine Pollution Bulletin* 52:1533-1540.

Seki, T., Taniuchi, T., Nakano, H. and Shimizu, M. 1998. Age, growth, and reproduction of the Oceanic Whitetip Shark from the Pacific Ocean. *Fisheries Science Tokyo* 64: 14–20.

Sequeira, A.M.M., Mellin, C., Fordham, D.A., Meekan, M.G. and Bradshaw, C.J.A. 2014. Predicting current and future global distributions of whale sharks. *Global Change Biology* 20: 778-789

Simpfendorfer, C. A. 2005. Threatened fishes of the world: *Pristis pectinata* Latham, 1794 (Pristidae). *Environmental Biology of Fishes* 73: 20.

Simpfendorfer, C.A. 2002. Smalltooth Sawfish: The USA's First Endangered Elasmobranch? *Endangered Species UPDATE* 19: 45-49.

Simpfendorfer, C.A., Wiley, T.R., and Yeiser, B.G. 2010. Improving conservation planning for an endangered Sawfish using data from acoustic telemetry. *Biological Conservation* 143(6): 1460-1469.

- Smith, S.E., Au, D.W., Show, C. 1998. Intrinsic rebound potentials of 26 species of Pacific sharks. *Marine and Freshwater Research* 49: 663–678.
- Springer, A.M., Estes, J.A., van Vliet, G.B., Williams, T.M., Doak, D.F., Danner, E.M., Forney, K.A., and Pfister, B. 2003. Sequential megafaunal collapse in the North Pacific Ocean: An ongoing legacy of industrial whaling? *PNAS*, 100(21): 12223-12228.
- Taylor, J. G. 2007. Ram filter-feeding and nocturnal feeding of Whale Sharks (*Rhincodon typus*) at Ningaloo Reef, Western Australia. *Fish. Res.* 84: 65-70.
- Thorburn, D.C. 2006. Biology, ecology and trophic interactions of elasmobranchs and other fishes in riverine waters of Northern Australia. PhD thesis, Murdoch University.
- Thorson, T.B. 1982. Life history implications of a tagging study of the large-tooth Sawfish, *Pristis perotteti*, in the Lake Nicaragua-Rio San Juan system. *Environmental Biology of Fishes*, 7(3): 207-228, figures 1-5.
- TRAFFIC (2013) Fact sheet for the 16th Meeting of the Conference of the Parties (CoP16) to the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES). [http://www.sharkadvocates.org/cites\\_4sharks\\_manta\\_fact\\_sheet.pdf](http://www.sharkadvocates.org/cites_4sharks_manta_fact_sheet.pdf)
- Van Beek, I.J.M., Debrot, A.O., Walker, P.A., Kingma, I. (2014) Shark protection plan for the Dutch Caribbean EEZ. Imares, Report number C209/13.
- Van Beek, I.J.M., Debrot, A.O., Walker, P.A., Kingma, I. (2014) Shark protection plan for the Dutch Caribbean EEZ. Imares, Report number C209/13.
- Van Beek, I.J.M., Debrot, A.O., Walker, P.A., Kingma, I. (2014) Shark protection plan for the Dutch Caribbean EEZ. Imares, Report number C209/13.
- Wathne F. 1959. Summary report of exploratory long-line fishing for tuna in Gulf of Mexico and Caribbean sea, 1954–1957. *Commercial Fisheries Review* 21: 1–26.
- White, W.T., Giles, J., Dharmadi and Potter, I.C. 2006. Data on the bycatch fishery and reproductive biology of mobulid rays (*Myliobatiformes*) in Indonesia. *Fisheries Research* 82: 65-73.
- Wiley, T., Carlson, J. & Smith, K. 2013. *Pristis pectinata* (Western Atlantic subpopulation). The IUCN Red List of Threatened Species 2013: e.T43398250A43398253. <http://dx.doi.org/10.2305/IUCN.UK.2013-1.RLTS.T43398250A43398253.en>.
- Wiley, T.R. and Simpfendorfer, C.A. 2010. Using public encounter data to direct recovery efforts for the endangered Smalltooth Sawfish (*Pristis pectinata*). *Endangered Species Research* 12: 179-191.
- Wintner, S.P. 2000. Preliminary study of vertebral growth rings in the Whale Shark, *Rhincodon typus*, from the east coast of South Africa. *Environmental Biology of Fishes* 59: 441–451.
- Young, N. 2001. An analysis of the trends in by-catch of turtle species, angelsharks and batoid species in the protective gillnets off KwaZulu-Natal, South Africa. Thesis, University of Reading.

## **Criteria for SPAW listing**

Criterion 1. Is the listing of the species warranted by the size of the population, evidence of decline, restrictions on its range of distribution, degree of population fragmentation, biology and behavior of the species, as well as other aspects of population dynamics, or other conditions clearly increasing the vulnerability of the species?

[If applicable] Criterion 2. Why is a precautionary approach necessary i.e., the lack of full scientific certainty about the exact status of the species is not to prevent the listing of the species on the appropriate annex?

Criterion 3. [In particular with respect to species proposed for Annex III], what are the levels and patterns of use and how successful are national management programs?

Criterion 4. Does the evaluation according to IUCN criteria, applied in a Caribbean context, i.e., the status of the population at the regional level, warrant listing of the species?

Criterion 5. Is the species subject to local or international trade, and is the international trade of the species regulated under CITES or other instruments?

Criterion 6. How important and useful are regional cooperative efforts for the protection and recovery of the species? [Include strengthening of existing cooperative efforts through global MEAs such as CMS]

Criterion 7. The species is not an endemic species [or there are specific reasons why cooperative action is important for its recovery].

Criterion 8. The species is not a sub-species.

Criterion 9. The status of the population at the regional level warrants listing, not only of a sub-population.

Criterion 10. Is the species essential to the maintenance of such fragile and vulnerable ecosystems/habitats, as mangrove ecosystems, seagrass beds and coral reefs and is the listing of the species felt to be an "appropriate measure to ensure the protection and recovery"?

Propuesta especies  
Cuba



**Taller Nacional para la Revisión de los Anexos al Protocolo SPAW.  
8 y 9 de Diciembre de 2011  
Jardín Botánico Nacional. La Habana. Cuba**

Se propone cambiar la Cotorra (*Amazona leucocephala*) para el Anexo III.

**Amazona leucocephala (Cotorra)**

La especie se encuentra amenazada con categoría de Vulnerable, sin embargo, una de las medidas de manejo para su conservación es la cría en cautiverio, razón por la cual se propone **pasar del Anexo II al III**. Existe un Plan de Manejo y Desarrollo por parte de la Empresa Nacional para la Protección de la Flora y la Fauna (ENPFF), del Ministerio de la Agricultura, para la cría en cautiverio y posible comercialización en el futuro, sin afectar las poblaciones silvestres y teniendo en cuenta los planes de manejo de las áreas protegidas donde habitan, ya que una de las principales amenazas es la captura de pichones para el comercio ilegal de mascotas.

**Características:**

La Cotorra Cubana, *Amazona leucocephala*, se distribuye en Cuba, Bahamas e Islas Caimán. Actualmente se reconocen cuatro subespecies: *A. l. leucocephala*; *A. l. bahamensis*; *A. l. hesterna* y *A. l. caymanensis* (Collar, 1997).


En Cuba, esta especie se distribuye por todo el país. Antiguamente se consideraba abundante y bien distribuida, pero en la actualidad sus bandos han sido muy reducidos en número. Los hábitats ocupados por esta especie incluyen bosques pluviales, bosques siempreverde mesófilo, bosques siempreverde de ciénaga, bosques aciculifolio de pinos, manglares y complejo de vegetación de costa rocosa. Aunque se localiza preferiblemente en hábitats naturales y bien conservados. *A. leucocephala* ha establecido una población en ambientes urbanos de La Habana, posiblemente de aves de jaula que se han escapado (Berovides y Cañizares, 2004).

Muchas poblaciones se encuentran hoy en áreas protegidas por toda Cuba, pero solo la población de la Reserva Ecológica "Los Indios" en la Isla de la Juventud ha sido manejada con éxito para su recuperación (Berovides *et al.*, 1995, 1996; Gálvez *et al.*, 1998).

La Cotorra Cubana anida en huecos abandonados de carpinteros y cavidades naturales de, prácticamente, cualquier especie de árbol, en dependencia de su disponibilidad, aunque son mucho más frecuentes los nidos en palmas de los géneros *Colpothrinax*, *Roystonea* y *Sabal*, y en troncos con huecos de mangle prieto (*Avicennia germinans*). A pesar de que hay poblaciones como la de Los Indios, que se reproducen en hábitat de sabanas abiertas, generalmente prefiere bosques intrincados y bien conservados con árboles maduros. Los nidos de Cotorras se localizan generalmente aislados unos de otros y los adultos se muestran muy cautelosos y silenciosos en las cercanías del nido. La nidificación se extiende desde marzo hasta finales de mayo. La puesta comprende usualmente entre 2 y 4 huevos, el periodo de incubación dura de 26 a 28 días y los pichones permanecen en el nido entre 56 y 60 días (Collar, 1997).

Se alimenta de una amplia variedad de flores, frutos y semillas de varias especies de plantas. González *et al.* (1987) registraron 18 especies de plantas consumidas por esta especie en la Ciénaga de Zapata y Gálvez *et al.* (1998) encontraron 39 especies vegetales consumidas en Los Indios, Isla de la Juventud. En un estudio reciente acerca de las preferencias en la dieta de esta especie en Alturas de Banao, se encontró que *A. leucocephala* utiliza los recursos tróficos proporcionalmente a sus disponibilidad en el ambiente (García, 2009).

Tiene una distribución nacional, el tamaño de sus poblaciones ha declinado y el hábitat está fragmentado, el cual ha disminuido 20% en los últimos 50 años (Berovides y Cañizares, 2004; Cañizares y Berovides, 2008).



Área de ocupación donde se ha registrado la especie

**Grado de protección:**

Nombre del área protegida donde se localiza la especie	Provincia
PN Guanahacabibes	Pinar del Río
APRM Mil Cumbres	Pinar del Río
RE Los Indios	Isla de la Juventud
PN Ciénaga de Zapata	Matanzas
RE Lomas de Banao	Sancti Spíritus
RF Tunas de Zaza	Sancti Spíritus
RF Delta del Agabama	Sancti Spíritus
PNP Topes de Collantes	Sancti Spíritus
PNP Hanabanilla	Villa Clara
RE Pico San Juan	Cienfuegos
APRM Humedales del norte de Ciego de Ávila	Ciego de Ávila
PNP Sierra de Najasa	Camagüey
APRM Sierra del Chorrillo	Camagüey
APRM Sierra de Cubitas	Camagüey
PN Pico Cristal	Holguín
PN La Mensura, Pilotoss	Holguín
PN Desembarco del Granma	Granma
RF Delta del Cauto	Las Tunas-Granma
PN Alejandro de Humboldt	Guantánamo-Holguín
RE Quibiján-Duaba	Guantánamo

Protegida en la Resolución 160/2011 en el Apéndice I como especie de especial significado de la República de Cuba

**Categoría de amenaza:**

Cuba: VU A2(a,c,d); B1b(i,ii,iii).

UICN: NT.

**Justificación de los criterios:**

Actualmente el comercio internacional de *A. leucocephala* está legalmente prohibido; sin embargo, a escala regional, el tráfico ilegal de esta especie es uno de los más preocupantes en Cuba. En un estudio desarrollado desde 1998 hasta 2008 en la región Central de Cuba, donde la Cotorra anida tanto en árboles en el bosque, como en oquedades de los farallones de piedra caliza (Pico San Juan), más de 90% de los nidos de Cotorra que se encontraron fueron

saqueados. El bajo porcentaje que sobrevive se debe, básicamente, a la inaccesibilidad de los nidos. En las condiciones actuales las medidas de protección sobre esta especie son totalmente ineficientes y el elevado precio de estas aves en el mercado negro constituye un fuerte incentivo para la actividad ilícita. Datos de un inventario realizado en varios municipios de la Habana, acerca de la tenencia de cotorras como mascotas, reflejan que, el porcentaje de viviendas donde se tienen estas aves es de cerca de 10%, aunque hay zonas residenciales como el Vedado, donde más de 30% de las viviendas poseen cotorras como mascotas (Patricia Rodríguez 2010, *Com. Pers.*)

A pesar de que no existen datos precisos de la magnitud del decline de las cotorras en Cuba en los últimos años, en conteos simultáneos realizados en áreas naturales y antrópicas de las montañas de Cuba central en marzo de 2009 se estima entre 90 y 100 Cotorras, para un área de más de 200 km<sup>2</sup>, lo que representa una densidad muy baja (0,5 ind/km<sup>2</sup>).

La principal amenaza para *A. leucocephala* es la captura de pichones para el comercio ilegal de mascotas, lo que en muchas ocasiones provoca la destrucción de los sitios de nidificación. Aunque también es afectada por la fragmentación y pérdida del hábitat, la deforestación y los huracanes.

Todas las poblaciones de la especie se encuentran amenazadas. Con excepción de las poblaciones del norte de la región oriental de Cuba, la mayoría se encuentran muy reducidas y con efectivos poblacionales muy bajos, por lo que las amenazas de extinciones locales son elevadas.

#### **Acciones que se deben acometer para su conservación:**

- Se debe trabajar en el manejo de hábitat y poblaciones silvestres, realizar monitoreos y trabajos de educación ambiental con el pueblo en general y en particular con la población humana que vive en los alrededores del área donde habita la especie. Además, se puede implementar la reproducción en cautiverio.
- Los datos anteriormente aportados se basan en censos, estudios de campos, observaciones informales de campo y la literatura disponible.

#### **Referencias:**

- Berovides, V. y M. Cañizares. 2004. Diagnóstico del decline de los psitácidos cubanos y su posible solución. *Rev. Biol.* 18(2):109-112.
- Cañizares, M. y V. Berovides. 2008. Distribución y abundancia de los bandos de psitácidos de Cuba Central. *Mesoamericana* 11(4):36-44.
- Collar, N. J. 1997. Family Psittacidae (Parrots). En del Hoyo, J., A. Elliott y J. Sargatal (Eds.) *Handbook of the Birds of the World*. Vol. 4. Sandgrouse to Cuckoos. Lynx Ediciones. Barcelona.
- Forshaw, J. M. 1989. *Parrots of the World*. Melbourne, Australia: Lansdowne Editions (3<sup>rd</sup> Ed.).
- Gálvez, X., V. Berovides y J. Fernández..1996. *Aratinga euops*. Taller para la conservación, análisis planificado de una selección de especies cubanas. La Habana, Cuba: Zoológico de la Habana / Facultad de Biología, Universidad de la Habana / CITMA / Parque Zoológico Nacional / Flora y Fauna / UICN, 47-55.
- García, L. 2009. Ecología trófica de los psitácidos en la reserva ecológica "Alturas de Banao", Sancti Spíritus. Tesis de Diploma. Facultad de Biología. Universidad de la Habana. 51 pp.
- Garrido, O. H. y A. Kirkconnell. 2000. *Field guide to the birds of Cuba*. Cornell University press, Ithaca: 207-208.
- González, H., D. Rodríguez y M. E.

#### **SE PROPONE AÑADIR AL ANEXO II LA ESPECIE:**

- *Passerina ciris* (Mariposa)

***Passerina ciris* (Mariposa)**

Es una especie migratoria considerada como Vulnerable dentro de la Lista Roja de especies amenazadas de Cuba. Ha habido una disminución de sus poblaciones debido a la captura indiscriminada con fines de lucro, para ser utilizada como ave ornamental.

**Características:**

La Mariposa presenta dos poblaciones reproductivas disyuntas. La del oeste de Estados Unidos, que habita en Kansas, Oklahoma, Texas, Arkansas, Louisiana, y al sur de los estados de Chihuahua, Coahuila, Nuevo León y Tamaulipas en México, que pasa el invierno en Centro América. La otra población, la del este de Estados Unidos, se reproduce en la porción costera de Carolina del Norte y del Sur, Georgia y el noreste de Florida. Esta pasa el invierno en el sureste de Louisiana y Alabama, en la Península de Florida, en Las Bahamas y Cuba (Íñigo-Elías *et al.*, 2002; Rich *et al.*, 2004). En Cuba la especie se distribuye en todo el territorio, aunque es más común en la parte centro oriental. Está catalogada como un común transeúnte y un raro residente invernal. Su alimentación es basada en semillas, fundamentalmente, aunque puede ingerir frutas. Es muy común verlas en formaciones vegetales secundarias o en áreas que colinden con zonas donde existan gramíneas. Esta especie no se reproduce en Cuba (González *et al.*, 1999; Garrido y Kirkconnell, 2000; Rodríguez, 2000).

Tiene una distribución nacional, el tamaño de sus poblaciones es desconocido en Cuba y el hábitat está fragmentado, el cual ha disminuido en los últimos 30 años.



Área de ocupación donde se ha registrado la especie

Nombre del área protegida donde se localiza la especie	Provincia	Grado de protección:
PN Guanahacabibes	Pinar del Río	
APRM Sierra del Rosario	Pinar del Río	
PN Ciénaga de Zapata	Matanzas	
RF Las Picúas-Cayo Cristo	Villa Clara	
RF Lanzasillo-Pajonal-Fragoso	Villa Clara	
APRM Humedales del norte de Ciego de Ávila	Ciego de Ávila	
APRM Humedales de Cayo Romano	Camagüey	
RF Río Máximo	Camagüey	
RE Caletones	Holguín	Categoría de

**amenaza:**

**Cuba: Vu A1 (a, b, c d, e); B2 (iii).**

**UICN: NT.**

**Protegida en la Resolución 160/2011 en el Apéndice I como especie de especial significado de la República de Cuba**

**Justificación de los criterios:**

Las poblaciones de Mariposa han sufrido una disminución de 55% en sus efectivos poblacionales en los últimos años 30 años. Una de las causas, es la pérdida de hábitat en las

zonas de cría en Norteamérica, la pérdida de hábitat en las zonas de invernada y el comercio ilegal a que está sometida la especie. La población que habita en el sureste de Estados Unidos y el Caribe presenta la mayor tasa de disminución poblacional con 3,9% anual (Íñigo-Elías *et al.*, 2002; Rich *et al.*, 2004).

En Cuba particularmente, el desarrollo turístico que se está llevando a cabo en la cayería norte de nuestro archipiélago, está ocasionando que se pierdan hábitats importantes para la alimentación y descanso de la Mariposa, durante la migración y la residencia invernal. Además, el auge que ha tenido el comercio de aves silvestres, en la última década, ha repercutido negativamente en sus poblaciones. En un estudio realizado en el año 2001, en tres localidades de Cuba, se demostró que anualmente se comercializan ilegalmente más de 400 individuos de la especie (Ayón, 2001). Este valor, sesgado por el número de localidades trabajadas, e intercambios recientes con pajareros, sugiere que el número real debe ser, al menos, el doble de lo estimado en dicho estudio.

**Acciones que se deben acometer para su conservación:**

Se debe trabajar en el manejo de hábitat, realizar monitoreos y trabajos de educación ambiental con el pueblo en general y en particular con los cazadores ilegales y los que comercializan la especie.

Los datos anteriormente aportados se basan en censos, encuestas, estudios de campos, observaciones informales de campo y la literatura disponible.

**Referencias:**

- Ayón, X. 2001. Uso y comercio de las aves silvestres cubanas. Tesis para la obtención del Título de Master Sobre Gestión, Conservación y Control de Especies Sometidas a Comercio Internacional. Universidad Internacional de Andalucía, Sede Antonio Machado, Baeza, Jaén, España. 109 pp.
- Garrido, O. H. y A. Kirkconnell. 2000. *Field guide to the birds of Cuba*. Cornell University press, Ithaca: 207-208.
- González Alonso, H., A. Llanes Sosa, B. Sánchez Oria, D. Rodríguez Batista, E. Pérez Mena, P. Blanco Rodríguez, R. Oviedo Prieto y A. Pérez Hernández. 1999. Estado de las comunidades de aves residentes y migratorias en ecosistemas cubanos en relación con el impacto provocado por los cambios globales. 1989-1999. *Informe Final* depositado en el Inst. de Ecología y Sistemática. 111 pp.
- Íñigo-Elías, E. E., K. V. Rosenberg y J. V. Wells. 2002. The Danger of Beauty. Birdscope, newsletter Summer 2002 Vol 16, no 3, of the Cornell Lab of Ornithology, Summer 2002. <http://birds.cornell.edu/publications/birdscope>.
- Llanes A., H. González, B. Sánchez y E. Pérez 2002. Lista de las aves registradas para Cuba. Aves de Cuba, Ed. H. González. UPC Print, Vaasa, Finland. 147-155.
- Llanes A., E. Pérez, H. González, A. Pérez. en prensa. *Wilsonia pussilla pileolata*, nueva subespecie para Cuba y adiciones a la ornitofauna a la Península de Guanahacabibes. *J. Carib. Ornithol.*
- Raffaele, H., J. Wiley; O. H. Garrido; A. Keith y J. Raffaele 1998. *A guide to the birds of the West Indies*. Princeton University press. Princeton: 424-425.
- Rich, T. D., C. J. Beardmore, H. Berlanga, P. J. Blancher, M. S. W. Bradstreet, G. S. Butcher, D. Demarest, E. H. Dunn, W. C. Hunter, E. Íñigo-Elías, J. A. Kennedy, A. Martell, A. Panjabi, D. N. Pashley, K. V. Rosenberg, C. Rustay, S. Wendt y T. Will. 2004. Partners in Flight North American Landbird Conservation Plan. Cornell Lab of Ornithology. Ithaca, NY.
- Rodríguez, D. 2000. Composición y estructura de las comunidades de aves en tres formaciones vegetales de Cayo Coco, Archipiélago de Sabana-Camagüey, Cuba. Tesis Doctorado, 99 Pp.

**Autor:** Eneider Ernesto Pérez Mena.

**ANEXO III.** Lista de Especies de Flora Marina y Costera Protegidas en Virtud del Artículo 11(1)(c)

Se propone pasar *Dendrocygna arborea* al Anexo II.

## **GRUPO DE TRABAJO REPTILES, ANFIBIOS Y MAMÍFEROS**

### **REPTILES**

Se propone cambiar para el Anexo III a la especie:

*Crocodylus acutus*

Es una especie de amplia distribución y existen programas de aprovechamiento aprobados por CITES para poblaciones silvestres. No se encuentra en Peligro Crítico y en la región del Caribe, las poblaciones están aparentemente estables. Se propone que se traslade del Anexo II al III.

## **GRUPO MOLUSCOS**

Se propone **INCLUIR** en el ANEXO III a *Liguus fasciatus* (Müller, 1774)

Presente en Florida, Cuba.

Muchos taxa considerados hasta ahora subespecies son costeros, sufren destrucción y fragmentación de hábitats (tala, carbón, cultivo, urbanización, captura directa con fines artesanales)

Ej. Cuba *L. f. sanctamariae*, *caroli*, *caribbaeus*, *leonorae*, *goodrichi s.s.*, etc

Ej. U.S. Subespecie del grupo *L. f. pictus* de los Cayos de la Florida, *L. f. septentrionalis* y varias de las variedades de *L. f. solidus*. Es susceptible al efecto directo del cambio climático.

a) *Liguus fasciatus* (Müller, 1774) Anexo III

Sinónimos:

Ubicación taxonómica: Familia Orthalicidae s.s., Superfamilia Orthalicoidea, Sigmurethra, Styломmatophora, Pulmonata, Clase Gastropoda, Phylum Mollusca.

b) Distribución: Es la única especie del género con un área de distribución tan extensa como toda Cuba, Isla de Pinos y la Florida, *L. virgineus* Linné solo en la Española y las otras solo en el occidente de Cuba. Esta especie puede producir variaciones discretas de patrón de coloración reconocibles para pequeñas zonas geográficas, por lo que se han descrito erróneamente en total más de 100 subespecies en todo su rango de habitación, y dentro de estas generar varios morfos de color.

Pudiera ser la de mejor capacidad adaptativa a juzgar por su distribución, pero varias de estas categorías taxonómicas están sufriendo decline y procesos de extinción.

Se necesitan primeramente estudios de taxonomía para definir sus especies, subespecies y grupos subordinados pues distintos taxa no suelen tener comportamiento ecológico idéntico y por tanto su programa de conservación debe tomar esto en cuenta. Además cuantificar la diversidad que encierra y tomar correctamente los datos de ubicación de poblaciones y sus variables ecológicas. Aunque existen vacíos de información de grandes extensiones de

territorio, alguna información de ecología básica se ha reunido en la mayoría de las publicaciones señaladas mas abajo.

c) Está incluido como especie protegida en el decreto ley de medio ambiente 160 en el apéndice 1 de protección total pero a pesar de esto se continúa su extracción. A pesar de haber estado sometida a la sobrecolecta y se han perdido algunas razas microgeográficas y otras están amenazadas, **no está en CITES, no está en lista roja de IUCN, ni en Catalog of Life.** En el Taller para la Conservación Análisis y Manejo Planificado de especies cubanas (CAMPII) Fernández *et al.*, 1997 proponen *L. fasciatus* como especie en peligro a incluirse en CITES y había sido considerada amenazada por Kay en 1995.

**Nota: ninguna de las otras 3 sp de Liguus cubanos, que si tienen distribución reducida estan en ninguna de estas.**

d) Requerimientos, interacciones ecológicas y amenazas: Es una especie arborícola que se alimenta raspando los complejos de hongos algas y líquenes de la superficie de las hojas y troncos de las especies de árboles y arbustos que prefiere, solo baja al suelo, y eso si hay buena cobertura de hojarasca, para poner los huevos en una cavidad que abre. Se distribuye en parches o agrupaciones en el area que ocupa la población, a veces muy reducida (Fernandez,2000). Prefiere bosque semideciduo (Fernandez y Berovides, 2000 y otros) sobre carso es capaz de explotar otras formaciones vegetales Utiliza arboles de corteza mas bien lisa y preferentemente de tronco grueso y se situa hacia el interior de la vegetacion. Por ejemplo el sabicú o soplillo *Lysiloma latisiliquum*, el almácigo *Bursera simarouba*, guao de monte *Metopium toxiferum*, y de costa *M.brownii*, *Mastichodendrum foetidissimum*, *Piscidia piscipula*, uvilla, *Coccoloba diversifolia*, *Krugiodendron ferreum*, guairaje *Eugenia axilaris*, dágame, anacahuita *Sterculia apetala*, granadillo *Brya ebenus* entre otros. En invierno se cementan a los troncos y ramas para hibernar a alturas mas bien bajas de unos centímetros a unos 4m pero en verano ascienden a veces hasta el dosel a 20m , puede explorar tanto hojas como ramas y su mayor densidad tambien esta asociada a una buena cobertura en la canopia del bosque , debido a todo lo anterior su presencia en buen numero es un indicador de salud del mismo A esto puede sumarse una distribucion cercana a gaussiana por clases tallas, sin estar afectado o ausente algun grupo de edad y una presencia de varios morfos de color, que es el comportamiento tipico de la especie, una reduccion en estos tambien es indice de disturbio.Es depredado por pájaros, guareao (*Aramus guarauna* ), arriero (*Saurothera merlini*) y otros, hormigas (una picada de *W.auropunctata* mata a un *L.fasciatus* adulto en menos de 12hrs y se han observado *S.geminata* depredandolo,también arañas,sin descartar otros artrópodos, grandes chipojos comedores de moluscos pudieran ingerir juveniles, al igual que algunas pequeñas serpientes. En Florida tambien los comen los mapaches y *Euglandina rosea* un gran gastropodo oleacinido voraz. Son suceptibles a enfermedades y parasitos pero no esta estudiado por el momento.Los afectan los largos periodos de sequía, inundaciones, huracanes que los dañan directamente y que tumban muchos de los grandes árboles hospedero (a veces más del 50 % de ellos como el Andrew en Florida), fuegos naturales, etc. Pero los que más impacto causan son los derivados de la acción humana, fragmentación de los bosques por tala con diversos usos, por incendios con o sin carácter intencional, minería, obras ingenieriles

desde gran a pequeña envergadura (carreteras, trasvases, pedraplenes, urbanización, etc), vale incluir aparte las acciones de desmonte, quema y modificación del terreno para cultivos, la invasión de plantas herbáceas que constituyen barrera de dispersión y otras que compiten con su hospedero y que no toleran como tal. Animales introducidos salvajes y tanto los domésticos como sus variantes salvajes producen grandes estragos: ratas, puercos, etc. Cualquier sustancia tóxica contaminante producto de la actividad humana puede ser asimilada por la piel, incluso el humo los daña.

La maduración sexual ocurre entre los 3 a 4 años de edad, entre julio y agosto ocurren los apareamientos y después de tres a seis semanas ponen los huevos (Davidson, 1965) sobre Noviembre empiezan a entrar en el letargo invernal por frío y falta de precipitaciones. Con las lluvias de abril-Mayo emergen los juveniles y salen los adultos de la hibernación. Las fluctuaciones de densidad no son grandes si no está perturbado pero obedecen a estos procesos mencionados. La capacidad de autofertilización parcial fue reconocida para la especie Hillis *et al.*, 1987; Hillis, 1989; Hillis *et al.*, 1991). *L. fasciatus* de la Florida tuvo un periodo de incubación de seis meses y el tamaño de nidada varía de 8 a 14 huevos (Blackwell, 1940) y hasta dos docenas aunque Fernández y Berovides (2001) registraron valores superiores (18 – 41), con 28.6 huevos promedio y un periodo de incubación más prolongado (6.6 a 8 meses) y las alteraciones del microhábitat afectaron la viabilidad de los huevos. La sequía retarda las eclosiones en moluscos (Pollard 1975). El tamaño de nidada en otros moluscos depende de la edad de los individuos y de las condiciones ambientales.

#### **f) programas de investigación y publicaciones científicas y técnicas disponibles acerca de las especies.**

Como parte de la descripción de lo que en la actualidad se consideran subespecies, Jaume en 1952 y 54 brinda datos sobre la distribución de estas y de otras especies y subespecies conocidas, así como literatura asociada. También fue uno de los primeros en poner por escrito en 1943 la preocupación por su conservación, aunque en el proceso de descripción colectaba miles de ejemplares para verificar la supuesta estabilidad de los morfos de color.

Alvarez y Berovides 1989 en Cayo Romano estiman la densidad de *L.fasciatus* entre 0.76 y 8 ind/100m<sup>2</sup>, la hayan similar a otras poblaciones observadas, al parecer normal y evalúan el efecto de 6 variables ecológicas en la distribución de los patrones de color. Observan preferentemente 1 ind./árbol, selección de árboles gruesos (+ de 9cm diámetro), buena cobertura de follaje, reposo a más de 2m.

Berovides y Alfonso 1995 en S.Chorrillo Camagüey estiman la influencia de la incidencia de depredación en robustez de la concha y distribución de 3 morfos de color de *L.fasciatus*. Alertan de lo fácil que resultaría para el hombre alterar la estructura genética y el tamaño de la población.

Fernández, I., L. Bidart, A. Fernández y V. Berovides, 1997 en el Informe del 2do Taller de Conservación, Análisis y Manejo Planificado (CAMP II), La Habana presentan la Hoja de datos del taxón *Liguus fasciatus*.

Referidos a ecología de *L.f.achatinus* en Holguín:

Fernandez, A. y Berovides 91, Fernandez, A. y Berovides,2000 en el Yayal Holguín, hallaron densidades de 0.12-0.17 animales/m<sup>2</sup> sin afectación por tala y la casi extincion al talarse el bosque. Los mismos autores en 1995 determinaron densidades de 0.09-0.31 con promedio 0.20 ind./m<sup>2</sup> para la población de Pedernales, hoy extirpada.

Bidart,L. *et al.*, 1992, González, et al,1997.

La tesis e maestría de A.Fernandez detrmno aspectos de la composición de la vegetación y su estructura y otras variables predictivas de los cambios espaciotemporales de densidad de *L.f.achatinus* en el Yayal, Holguín y su utilidad como indicador de salud del bosque.

Para ecología de *Lf.sanctamariae* se cita a Fernández I. et al.95 Fernández y Perera, 1997 que relacionan algunas variables ambientales que pudieran determinar la distribucion de los morfos de color en el cayo de igual nombre.

Espinosa y Ortea en 1999 relacionan la mayoría e las especies y subespecies del genero con algunas de sus localidades, entre otros moluscos terrestres.

Tambien se relacionan abajo algunas publicaciones referentes a descripciones de taxa infraespecificos y una breve busqueda bibliografica que muestra que en los ultimos años esincluido en estudios de conservacion y ecologia, algunos en Cuba.

Young 1951 y 1958 y Blackwell, 1940 explican las causas que amenazan a esta especie en la Florida y Voss, 1976; Brown, 1978 y Bennetts *et al.*, 2000 comentan sobre relaciones molusco-hospedero, en especial *L.latisiliquum*.

Mas relacionados con aspectos geneticos y ecologicos los de Roth y Bogan ,1984 para alelos de color de la concha, los de Young 1960, ambos en Florida.

H.A.Pilsbry describe varias subespecies en 1912 y 1946 y menciona algunas preferencias de arboles en distintas zonas de Florida.

e) No existe ningún plan de manejo **implementado** en Cuba para contribuir con su protección aunque si hay propuestas en documentos no publicados como las mencionadas para *L.f.achatinus* por A.Fernández en Holguín y Maceira et al.2011 en Granma, en las que se alerta de su decline fundamentalmente por accion humana. Dichos planes incluirían monitoreo de las poblaciones, cría ex situ, reforestar con vegetación primaria, intentar eliminar las plantas y animales invasores, garantizar las condiciones de trabajo de los conservadores y la educacion ambiental de los comunitarios. En algunas de las áreas protegidas visitadas se tiene conocimiento de la presencia de la especie de la necesidad de conservarla por los especialistas, pero los medios de transporte, comunicación y la actitud de los locales no garantizan una vigilancia efectiva.

Se ha criado en cautiverio en Florida y se han hecho acciones de traslado e implantación de pequeñas colonias con cierto éxito, aunque sin estudiar a fondo las posibles consecuencias. Si hay planes de conservación y manejo en activo en las zonas protegidas.