

A biochemical mechanism for time-encoding memory formation within individual synapses of Purkinje cells

Ayush Mandwal^{1,*}, Javier G. Orlandi¹, Christoph Simon², Jörn Davidsen^{1,3,*}

1 Complexity Science Group, Department of Physics and Astronomy, University of Calgary, Calgary, Alberta, Canada

2 Institute for Quantum Science and Technology, Department of Physics and Astronomy, University of Calgary, Calgary, Alberta, Canada

3 Hotchkiss Brain Institute, University of Calgary, Calgary, Alberta, Canada

* ayush.mandwal@ucalgary.ca

* davidsen@phas.ucalgary.ca

Abstract

Within the classical eye-blink conditioning, Purkinje cells within the cerebellum are known to suppress their tonic firing rates for a well defined time period in response to the conditional stimulus after training. The temporal profile of the drop in tonic firing rate, i.e., the onset and the duration, depend upon the time interval between the onsets of the training conditional and unconditional stimulus. Direct stimulation of parallel fibers and climbing fiber by electrodes was found to be sufficient to reproduce the same characteristic drop in the firing rate of the Purkinje cell. In addition, the specific metabotropic glutamate-based receptor type 7 (mGluR₇), which resides on the Purkinje cell synapses, was found responsible for the initiation of the response, suggesting an intrinsic mechanism within the Purkinje cell for the temporal learning. In an attempt to look for a mechanism for time-encoding memory formation within individual Purkinje cells, we propose a biochemical mechanism based on recent experimental findings. The proposed model tries to answer key aspects of the “Coding problem” of Neuroscience by focussing on the Purkinje cell’s ability to encode time intervals through training. According to the proposed mechanism, the time memory is encoded within the dynamics of a set of proteins — mGluR₇, G-protein, G-protein coupled Inward Rectifier Potassium ion channel, Protein Kinase A and Protein Phosphatase 1 — which self-organize themselves into a protein complex. The intrinsic dynamics of these protein complexes can differ and thus can encode different time durations. Based on their amount and their collective dynamics within individual synapses, the Purkinje cell is able to suppress its own tonic firing rate for a specific time interval. The time memory is encoded within the effective rate constants of the biochemical reactions and altering these rates constants means storing a different time memory. The proposed mechanism is verified by a simplified mathematical model and corresponding dynamical simulations of the involved biomolecules, yielding testable experimental predictions.

Author summary

Hebbian plasticity is a widely accepted form of learning that can encode memories in our brain. Spike-timing dependent plasticity resulting in Long-term Potentiation or Depression of synapses has become the default mechanistic explanation behind memory

formation within a neuronal population. However, recent experiments of conditional eyeblink response in Purkinje cells have challenged this point of view by showing that these mechanisms alone cannot account for time memory formation in the Purkinje cell. To explain the underlying mechanism behind this novel synaptic plasticity, we introduce a biochemical mechanism based on protein interactions occurring within a single synapse. These protein interactions and the associated effective rate constants are sufficient to encode time delays by auto-induced inhibition on a single excitatory synapse, suggesting that synapses are capable of storing more information than previously thought.

Introduction

How do we store memories in our brain? How do we retrieve and edit them when required? Recent experimental findings have shed some light onto these fundamental questions. Experiments have shown that memories are held within specific neuronal populations [1–3]. Such populations, referred as *memory engram cells* [4, 5] store memory either by forming or eliminating synapses [6, 7] or by altering synaptic strengths between neurons [8, 9] within the population. These forms of learning and memory formation fall under the widely accepted Hebbian learning paradigm [10]. However, the individual contribution of each synapse to the engrams, and how changes in synaptic strength affects memories, remain poorly understood. The problem of information encoding was raised by C.R. Gallistel [11] and termed as the “Coding Question”, one of the fundamental open questions in Neuroscience today. Recent experiments on Purkinje cells, one of the major neuronal populations in the Cerebellum and essential for motor coordination, have shed some light on the Coding Problem. Those experimental results have illustrated that the memory of time interval duration can be encoded within individual Purkinje cells, and does not require a whole neuronal population [12, 13]. In addition, the stored time memory can be accessed and changed anytime. This result has also challenged the prevailing doctrine of Hebbian learning by showing that traditional changes of synaptic strength alone cannot explain the Purkinje cell response after learning [14].

Purkinje cells can learn to encode a specific time memory through Classical or Pavlovian conditioning. This kind of Associative learning can occur when a biologically potent stimulus, such as food, is paired with a neutral stimulus, such as a bell, that precedes it. Depending upon the response the potent stimulus elicits, e.g., saliva flow, and the exact protocol followed, Classical Conditioning can be categorized into various kinds. One of them being classical motor conditioning, such as the eye blink conditioning, where a neutral conditional stimulus (CS) in the form of a light or a sound can trigger an eye blink response before the onset of an unconditional stimulus (US) that elicits a blink reflex response [15, 16]. In other words, CS triggers a response that predicts the time of arrival of the US. Such a conditional response appears after successive training sessions, where a CS is followed by an US after a fixed time interval “T”, called the interstimulus time interval (ISI) [17]. At the cellular level, the eye blink response is causally related to a suppression of the tonic firing of individual Purkinje cells, which regulate the activity of ocular muscles [17, 18]. Because of such causal connection, the suppression of the firing rate of the Purkinje cell is termed as the conditional response of the Purkinje cell.

Previous mechanistic explanations considered Long-term Depression (LTD) of selective synapses between parallel fibers and Purkinje cells (pf-PC) as the main mechanism behind the conditional response in the Purkinje cell [19]. Based on the widely accepted Marr-Albus model of the cerebellum [20, 21], this suggests that the time memory of the response is encoded within the network dynamics of Granule cell neurons and inhibitory interneurons, found within the molecular layer of the Cerebellum

between Mossy fibers and Purkinje cells. However, recent experiments on ferrets were able to identify the source of the conditional response at the level of *individual* Purkinje cells by showing that the direct stimulation of parallel fibers and climbing fibers using electrodes was sufficient for Purkinje cells to learn the specific time interval duration [12]. These experiments also showed that a glutamate-based metabotropic receptor type 7 (mGluR7) which resides on Purkinje cells synapses, initiates the conditional response [13] by opening G-protein coupled Inward Rectifier Potassium (GIRK) ion channels [22]. This implies that there exists a specific biochemical mechanism within the Purkinje cell that can encode and store temporal information.

Unlike other memory formation mechanisms requiring neuronal assemblies, temporal signatures can be encoded within a single Purkinje cell, but the specific mechanism remains poorly understood. Here, we propose a biochemical description, based on past experimental findings, that is able to explain time memory formation, consolidation and access.

Biochemical description

As mentioned above, the conditional response at the level of an individual Purkinje cell appears after several repetitions of two stimuli: A CS from the parallel fibers followed by an US from the climbing fiber after a fixed ISI. A sufficient condition for the learning process to be called completed is that a CS without an applied US can initiate the conditional response — the suppression of the tonic firing rate — with the given ISI. Below we discuss the conditional response of the Purkinje cell and the associated detailed biochemistry we propose. We separate it into three parts: during training, after training and training with different ISIs. The first part focuses on two questions: What makes a Purkinje cell learn a conditional response, and how does the cell learn a conditional response of a specific duration? The later two parts describe the most crucial aspects of the conditional response, i.e., its formation after training, along with other features of the conditional response, which were experimentally observed.

During training

What makes a Purkinje cell learn a conditional response? The activation of the conditional response was found to be initiated by the activation of mGluR₇ receptors [13]. Purkinje cells express mGluR₇ receptors on their entire cell body and dendritic branches [23], yet no conditional response was observed before training [12]. The fact that in the presence of CS Purkinje cells behave similarly before training and after training if mGluR₇ antagonists such as 6-(4-Methoxyphenyl)-5-methyl-3-(4-pyridinyl)-isoxazolo[4,5-c]pyridin-4(5H)-one hydrochloride (MMPiP) or LY341495 are present implies that the learning of the conditional response is associated with the expression of mGluR₇ receptors on the synapse. In other words, during training mGluR₇ receptors are being transported from the perisynaptic zone to the postsynaptic zone of the synapse via some biochemical mechanism that is activated during training. Once placed on the synapse, mGluR₇ receptors activate $G_{i/o}$ type G-proteins in the presence of glutamate from the parallel fibers. In turn, the $G_{\beta\gamma}$ subunits of the $G_{i/o}$ type G-proteins activate the GIRK ion channels [22, 24]. However, as alternative hypotheses to the absence of mGluR₇ from the postsynaptic zone before training two other options are conceivable: 1) GIRK ion channels are absent at the synapse; or 2) the expression of $G_{i/o}$ type G-proteins was low at the synapse. However, we can rule these out. Immunohistochemistry analysis showed the presence of GIRK subtypes GIRK2/3 ion channels on PC synapses — which are innervated by parallel fibers — in the absence of any kind of prior conditional training [25]. This rules out 1). If 2) were

true and the G-protein expression would change during training, this would affect not only the conditional response profile but also various other physiological properties of the Purkinje cell. This is because different types of G-proteins play crucial roles in signal transductions and provide various physiological properties to the cell [26]. Since no change in the tonic firing rate has been observed before and after conditional training [12], we believe that other physiological properties of the cell may also remain unaltered. Thus, the translocation of mGluR₇ receptors to the synapse is the most likely result of the training and we assume that the amount of other proteins such as GIRK ion channels and G-protein is constant for all different durations of conditional training.

How does the Purkinje cell learn a conditional response of a specific duration? To train a Purkinje cell to memorize a specific duration “T”, two stimuli separated by the same duration, called the Interstimulus Interval (ISI), are required. The first stimulus must come from the parallel fibers and the second stimulus comes from the climbing fiber [17]. Generally, translocation of mGluR type receptors to and from the synapse happens by Clathrin-mediated Endocytosis (CME) [27]. Also, G-protein coupled receptor kinases (GRKs) and in some cases protein kinases such as Protein kinase C (PKC) can phosphorylate receptors and initiate their endocytosis and translocation via CME [27]. Therefore, we propose that the PKC initiates trafficking of mGluR₇ receptor via CME. This is supported by the fact that the presence of two stimuli, one from the parallel fiber and the other one from the climbing fiber, also makes PKC activation most favorable [28]. In particular, the presence of either one of the two stimuli is not sufficient for the Purkinje cell to learn the conditional response [17]. While dendritic spines of Purkinje cells — which parallel fibers are innervating — express mGluR₁ receptors [29] and those receptors can potentially activate PKC [30], evidently PKC activation is not the downstream effect of mGluR₁ receptor activation alone [31, 32]. Similarly, climbing fiber stimuli alone cannot activate PKC. While Long-Term Depression (LTD) of pf-PC synapses occurs via PKC activation [33], it only happens when both parallel fibers and climbing fiber are active [28]. Hence, PKC can become active during training and helps in translocation of mGluR₇ receptors to the synapse. However, PKC alone is not responsible for the translocation of mGluR₇ receptors as Purkinje cells cannot be trained for ISI durations shorter than 100ms [34] but PKC can become active even when both CS and US occur at the same time. Currently, we cannot make any suggestion for proteins, which might be involved in addition to PKC during conditional learning.

To ensure storage of a specific time duration memory, such translocation process must stop after some time. This can happen by inhibiting PKC activation, which can be achieved by either removal of mGluR₁ receptors from the synapse or by preventing the rise of the intracellular Ca⁺² ion concentration within the synapse. The latter option is the most suitable one because translocated mGluR₇ receptors at the synapse can open GIRK ion channels, which will drop the membrane potential and thus prevent opening of Voltage-gated Ca⁺² ion channels. Activation of GIRK ion channels causing a drop in tonic firing rate during training has been observed in experiments [35, 36]. Moreover, if mGluR₁ receptors were to be removed from the synapse, then the retraining process with a different ISI would not happen as there would be none or very few mGluR₁ receptors left on the synapse to produce Diacylglycerol (DAG), a necessary membrane-bound biomolecule for PKC activation [30]. As Purkinje cells can be retrained [36], the amount of mGluR₁ receptors cannot decrease on the synapse. Therefore, we conclude that as training progresses the intracellular Ca⁺² ion concentration decreases to a level that is no longer sufficient to activate PKC, which prevents further translocation of mGluR₇ receptors to the synapse and so a steady state will be reached. When a steady state has been reached, then we can say that the Purkinje cell has learned the conditional response of duration “T” as shown in fig.(1). This mechanism of learning also suggests that the training period will be longer for a

long duration conditional response as observed in the experiment [12]. As the net amount of the receptor translocated during training depends upon its net duration, long duration training means more transportation of the receptor to the synapse, which produces a long duration conditional response. We will explain how a higher amount of receptors can produce a long duration conditional response below.




Fig 1. mGluR₇ receptor distribution before and after conditional training in the Purkinje cell. Before training, mGluR₇ receptors are localised at perisynaptic areas of the synapses. After training, as pointed out by the blue arrows, these receptors localised themselves at the postsynaptic area of the synapse via CME.

After training

The conditional response with a duration of hundreds of milliseconds can be initiated by a CS of as little as 20 milliseconds duration [12]. This means that just the activation of the mGluR₇ receptors by CS is enough to initiate the conditional response. In addition, the dynamics of the G-protein activation by mGluR₇ receptors and the binding of G-protein subunits to the GIRK ion channels are usually too slow to explain the fast dynamics of the conditional response initiation observed in the experiment [12]. Such fast activation of GIRK ion channels correlates with some past studies. It was proposed and later verified experimentally that the mGluR₇ receptor forms a protein complex with the G-protein of G_{i/o} type, which is located in direct vicinity to a GIRK ion channel as facilitated by a Regulator of G-protein signaling protein 8 (RGS8) [37, 38]. RGS8 proteins are expressed in dendritic spines of the Purkinje cell [39] and they have a special property to accelerate both activation and deactivation of the G-protein causing fast opening and closing of GIRK ion channels [38]. In particular, it has two domains.

RGS8's core domain is responsible for rapid deactivation of the G-protein. An additional N-terminal domain helps in the rapid G-protein activation, possibly by enhancing the coupling between G-protein and the receptor [40]. Thus, the fast dynamics of the conditional response is possible by forming a protein complex of mGluR₇ receptor, G-protein and GIRK ion channel facilitated by a RGS8 protein.

There are two additional important properties of the conditional response, which we must consider: 1) The conditional response is lost after repetitive CS [12], and 2) the conditional response is independent of CS duration. These two properties are in fact related to each other. The former implies the removal of mGluR₇ receptors from the synapse and the latter implies that some protein blocks the receptor's active site to prevent reactivation of the conditional response. Dephosphorylation of the mGluR₇ receptor by Protein Phosphatase 1 (PP1), which causes their rapid internalization [41], can explain both these properties of the conditional response. Rapid internalization of any receptor is initiated by the binding of a protein called Arrestin protein, which prevents the receptor to transmit any signal further [30]. Since CS activates the conditional response, it implies that PP1 is inactive before the conditional response. Furthermore, the conditional response is mediated via activation of the G_{i/o} type G-protein whose G_α subunit blocks the production of Cyclic adenosine monophosphate (cAMP) by Acetyl Cyclase (AC) and, thus, results in reduced Protein Kinase A (PKA) activity. Also, due to the tonic firing of the Purkinje cell, Calmodulin, which regulates intracellular Ca⁺² ion concentration [42], can stimulate Acetyl cyclase (AC) [43] to produce cAMP molecules and increase PKA activity. It is also known that PKA can phosphorylate mGluR₇ receptors [44] as well as PP1 regulatory proteins such as Dopamine- and cAMP-Regulated neuronal Phosphoprotein (DARPP-32) or Inhibitor-1 (I-1) [45], which prevents dephosphorylation of receptors. Thus, phosphorylation of the receptor and inhibition of PP1 activity by PKA help in the retention of the memory for a long time. As PKA is essential for the conditional response, this protein could bind to the receptor via a special PKA anchoring protein called A-kinase anchoring proteins (AKAP) [46]. Similarly, PP1 can also bind close to the receptor via another scaffold protein namely Spinophilin [47], which is expressed in dendritic spines of neurons across various regions of the brain including Cerebellum [48]. Such a close agglomeration of various proteins together forms a protein complex, which is self-sufficient in its own regulation and dynamics.

In short, the underlying biochemical mechanisms of the conditional response can be described as follows. Before CS, the PP1 protein is inactive because of PKA activity. The release of glutamate during CS activates mGluR₇ receptors on the Purkinje cell synapses [step 1 of Fig. (2)], which in turn activates G-proteins [step 2 of Fig. (2)]. Each unit of G-protein splits into a G_α subunit and a G_{βγ} subunit. One unit of G_α subunit binds to an AC enzyme to block the production of cAMP molecules. This in turn deactivates PKA as Phosphodiesterase enzyme (PDE) hydrolyses the remaining cAMP molecules [30] [step 3 of Fig. (2)]. At the same time the G_{βγ} subunit binds to the GIRK ion channel, which becomes fully active upon binding of four G_{βγ} subunits [24]. As PKA activity decreases, PP1 activity rises due to dephosphorylation of DARPP-32 or I-1 by Protein Phosphatases such as PP2A [45,49] [step 4 of Fig. (2)]. The rise of PP1 activity causes dephosphorylation of mGluR₇ receptors [step 5 of Fig. (2)] and initiates their rapid internalization. However, rapid internalization of a receptor is still a slow process compared to the conditional response as it involves many protein interactions and, hence, the receptor is not immediately displaced from the synapse after dephosphorylation. However, after dephosphorylation, Arrestin protein blocks the active site of the mGluR₇ receptor to prevent reactivation of the G-protein [50]. After receptor dephosphorylation, the active G-protein is deactivated by the RGS8 protein [step 6 of Fig. (2)]. As G-protein activity reduces, GIRK ion channels also shut down. In the

absence of active G-protein, PKA activity begins to rise again [step 7 of Fig. (2)] due to rise in activity of AC enzymes in the presence of Calmodulin. Active PKA deactivates PP1 [step 8 of Fig. (2)] by phosphorylating DARPP-32 or I-1 and finally, PKA also phosphorylates mGluR₇ receptors [step 9 of Fig. (2)] to prevent their internalization and prepare the Purkinje cell for another conditional response. It is likely that the reactivation of PKA takes some time, which might explain why CS cannot initiate another conditional response while CS is still on. 214
215
216
217
218
219
220




Fig 2. Interactions between different biochemicals involved in our proposed mechanism. The numbers on the top of the arrows highlight the order in which the different reactions occur during the conditional response. Conditional response initiates with the release of glutamate from parallel fibers denoted by I as input in (1), which activates mGluR₇ receptors. In (2), active receptors activate G-proteins, which deactivate PKA through (3). As PKA activity reduces, PP1 activity rises through (4) causing dephosphorylation of the receptor (5). As receptor activity reduces, RGS8 reduces G-protein activity (6), which allows PKA activity to rise again (7). Active PKA will deactivate PP1 (8) and lastly phosphorylate dephosphorylated receptors to prevent their rapid internalization (9). 221
222
223
224
225
226

In fig.(2), the rate at which GIRK ion channels open and close depends upon the rate at which intermediate reactions occur. In other words, the time memory of the training is stored within the effective rate constants arising from these reactions. In a complete cycle of GIRK ion channel activation and deactivation, altering only effective rate constants for both activation and deactivation processes is sufficient to store a different time memory of the conditional response. 227
228
229
230
231
232
233
234
235

Training with different ISI duration and time-encoding protein complexes

Training with a different ISI duration means storage of a different time memory. As discussed above, the time memory is encoded within effective rate constants of the biochemical reactions, which regulate the gating dynamics of the GIRK ion channel and altering these rate constants means storing a different time memory. However, there are two additional questions we need to answer in order to get a complete understanding of time memory storage in biochemical reactions: 1) How do these biochemical reactions get tuned so finely to store a specific time duration memory? 2) Among all possible 227
228
229
230
231
232
233
234
235

effective rate constants of the proposed biochemical mechanism, which rate constants are most likely to get affected by choosing a different ISI for the training?

The reason behind 1) is that there are several GIRK ion channels present at the synapse. Each GIRK ion channel requires four units of $G_{\beta\gamma}$ subunits to open completely [24]. This means that each GIRK ion channel forms a protein complex with four units of G-protein, receptor and RGS8 protein along with PKA and PP1 proteins with their anchoring proteins together. As each of these protein complexes has their own intrinsic dynamics, which regulate how fast the GIRK ion channel opens and closes upon stimulation, we can call each of these protein complexes “Time-Encoding protein Complexes” (TEC). Within each TEC, the rate of G-protein activation by the receptor and the rate of binding of G-protein subunits to the GIRK ion channel decide the overall rate of opening of GIRK ion channels i.e., the onset of conditional response. After the onset of the conditional response the rates of PKA deactivation, PP1 activation, dephosphorylation of the receptor and the deactivation of G-protein by RGS8 decide the overall duration of the conditional response since at the end of these biochemical reactions the GIRK ion channel begins to close. Thus, each TEC encodes time information of the conditional response completely in terms of effective rate constants of different biochemical interactions and stores this time memory by forming a protein complex. Formation of a protein complex as TEC ensures strong consolidation of memory with less chances of errors in the information storage. If the rates were to be changed so would the memory as well. The rates can be affected by the translocation of extra mGluR₇ receptors to the synapse during conditional training. These extra mGluR₇ receptors can form clusters with receptors — which are part of TEC — with the help of a scaffold protein, Protein Interacting with C Kinase - 1 (PICK1) [51]. Such cluster formation can affect TEC’s intrinsic dynamical properties by influencing the protein interaction of the mGluR₇ receptor with the G-protein facilitated by RGS8. As a result, RGS8’s ability to accelerate the dynamics of the conditional response might be affected, which results in a delayed onset of the conditional response. Such clustering of receptors can also affect the dynamics of the PKA protein anchored close to the receptor via the AKAP protein, thus affecting also the time duration of the conditional response. To summarize, we propose that at individual synapses the interaction of extra mGluR₇ receptors with TECs can affect the dynamics of TECs and collectively these varied TEC units help to produce the conditional response of any specific time duration in the Purkinje cell.

Interestingly, time memories stored at synapses in the form of TECs are not permanent but can be altered or edited through retraining. Retraining can happen via two ways, (i) erase the memory first and then store another memory by training with a different ISI interval, or (ii) retrain the Purkinje cell with a different ISI without erasing the previous memory. Experimentally, time memory can be erased by repeating unpaired representation of CS and US several times [12], which can also be explained by our proposed mechanism. Due to the action of PP1 on the receptor, every time CS initiates a conditional response, some of the receptors at the synapse might undergo rapid internalization from which phosphorylation of the receptor by PKA cannot bring them back and, hence, these receptors will be removed from the synapse. Because of rapid internalization, retraining with the same or a different ISI will be faster as many receptors are close to the synapse. This rapid relearning phenomenon is called “Saving” and it takes only a few minutes to recall the old memory [17]. If retraining with a different ISI is performed without erasing the old memory then the conditional response profile differs in terms of the onset of the conditional response while the duration remains intact when compared with the case where retraining does involve erasing the old memory prior to retraining [17,52]. This difference implies that there might be some membrane-bound proteins or a scaffold protein such as PICK1 that are interacting with

TECs. If the two ISI durations differ significantly then surprisingly, the Purkinje cell pauses twice in the presence of a sufficiently long duration CS [12]. Depending upon the time difference between the two different ISIs, different conditional responses were observed implying that there might be additional interactions among TECs, which give rise to a wide variety of conditional responses [53].

From the above description, it follows that in principle a single synapse can completely contain a time duration memory, which can be altered through retraining. However, a single synapse probably will not be sufficient to suppress the tonic firing rate of the whole Purkinje cell. This is because the spontaneous tonic firing rate of the Purkinje cell [54] appears to be due to voltage-dependent resurgent Na^+ ion channels, which are distributed over the *entire* somata and dendritic regions of the cell [55, 56]. The resurgent Na^+ channels have the property to become active and inactive during depolarization, as well as to reactivate during repolarization due to the presence of a positive membrane potential. The latter results in a spontaneous rapid sequence of action potentials [57]. In order to suppress the spontaneous firing rate of the cell, these resurgent Na^+ ion channels need to be deactivated by the hyperpolarizing membrane potential of the entire cell. Activation of GIRK ion channels by CS can hyperpolarize the membrane at a synaptic regions and deactivate resurgent Na^+ ion channels near those synaptic region. Their deactivation will lower the overall membrane potential and the activation of multiple pf-PC synapses distributed over the dendrites can overcome the resurgent Na^+ current and suppresses tonic firing rate of the cell. Thus, a finite fraction of the total pf-PC synapses can produce a suppression in tonic firing rate of the Purkinje cell for a specific duration and the corresponding memory is encoded at the respective synapses.

In the next section, we will introduce and discuss a mathematical model of our proposed biochemical mechanism and subsequently make few predictions that can be tested experimentally.

Mathematical model

To model the conditional response behavior of the Purkinje cell, we start with an established dynamical model of the Purkinje cell which incorporates many properties of the cell within a realistic biophysical framework [58], see Materials and Methods for details. Before training, GIRK ion channels cannot be opened because mGluR_7 receptors are not present at the synapse. However, after training, mGluR_7 receptors are present at the synapse to open GIRK ion channels. Therefore, adding an additional term for the gating of the GIRK ion channel in the Purkinje cell model will allow it to exhibit the conditional response. As GIRK ion channels reside at synapses, the additional term of GIRK ion channel gating must be added in the dendritic equation of the Purkinje cell model. Eq.(1) defines the gating of the GIRK ion channels, in which g_{GIRK} is the net conductance of GIRK ion channels per unit area, h_{GIRK} is the gating parameter and V_{GIRK} is the voltage dependence of the GIRK ion channel obtained from the I-V characteristics curve [59]. Since no experimentally measured value exist for g_{GIRK} to the best of our knowledge, we choose its value to match the conditional response observed experimentally.

$$\begin{aligned} I_{GIRK} &= -g_{GIRK}h_{GIRK}V_{GIRK}(V_d) \\ V_{GIRK}(V_d) &= -0.02(1.3V_d + 50.0)/(1.0 + \exp((V_d + 40)/10.0)) \end{aligned} \quad (1)$$

Gating of the GIRK ion channel depends upon the availability of the Phosphatidylinositol 4,5-bisphosphate (PIP_2) molecules [24]. This molecule has low affinity for the GIRK ion channel but binds efficiently after binding of $\text{G}_{\beta\gamma}$ subunit to

the GIRK ion channel. The amount of PIP₂ on the synaptic membrane is low but its quantity is often replenished by various biochemical processes to maintain its concentration fairly constant upon consumption or degradation [60]. Therefore, the amount of active G_{βγ} subunits can determine the gating dynamics of the GIRK ion channel. As G-protein is closely associated with the GIRK ion channel, we can assume fast binding of the G_{βγ} subunit to the GIRK ion channel. With these assumptions, we can equate the normalized G-protein activity with the GIRK ion channel gating parameter h_{GIRK} as shown in eq.(6) below.

G-protein activity depends upon the activity of the mGluR₇ receptor along with other proteins as shown in fig.(2), which self-organize to form discrete units of TECs. Since we don't know the number of TECs and their detailed intrinsic dynamics, we choose to model the collective dynamics of TECs and different biochemical interactions within them in an effective way. Hence, instead of using discrete variables for the activity of different biochemicals, we use continuous variables to capture the “average” dynamics of different biochemicals by considering all TECs together. In addition, AKAP proteins — which anchor PKA close to cAMP production machineries — do accelerate the activation of PKA, but not sufficiently fast to generate a conditional response [61,62]. It is possible that other proteins such as Homer Proteins might also be involved in the TEC, facilitating cross-talk between various target proteins [63]. Thus, due to lack of knowledge of various protein interactions and their strengths within individual TECs, we only attempt to create a conceptual minimal model whose aim is to reproduce features of the conditional response, namely 1) the conditional response should be independent of CS duration, and 2) changing the dynamics of PKA and G-protein should be sufficient to produce conditional responses of different durations. Our conceptual minimal model consists of four main biochemicals - mGluR₇, G-protein, PKA and PP1. In order to simplify and minimize the number of parameters, we used nonlinear terms to model their overall behaviour as observed *in vivo*.

The nondimensional dynamical equations for the proposed biochemical mechanism within individual TECs are as follows:

$$\tau_1 \frac{du}{dt} = \frac{1}{\alpha + x} u(u_0 - u) - \beta u, \quad (2)$$

$$\tau_2 \frac{dv}{dt} = (v - v_2)(v_1 - v)(v - v_0) - \gamma wv + I, \quad (3)$$

$$\tau_3 \frac{dw}{dt} = -\delta uw + \eta(w_0 - w) + w(w_0 - w), \quad (4)$$

$$\tau_4 \frac{dx}{dt} = (v - x) \quad (5)$$

$$h_{GIRK} = x/v_0 \quad (6)$$

where u , v , w and x are the activities of PKA, mGluR₇ receptor, PP1 and G-protein respectively. In the above model, all the parameters and variables are positive and dimensionless quantities except for τ_i for $i = 1, 2, 3, 4$, which have dimension of time.

Eqs.(2 – 5) match the pictorial diagram shown in Fig. 3, which depicts various variables and their dependencies. In order to understand various terms within each equation, let us focus on each equation individually. Before CS, when G-protein is still inactive, i.e., $x \sim 0$, AC enzyme produces cAMP molecules facilitated by the Calmodulin protein. As the activity of AC enzyme increases, cAMP production also increases, which increases PKA activity. This behaviour of PKA activity is modeled in eq.(2) by the first term $u(u_0 - u)/\alpha$ for $x = 0$, where u_0 is the maximum PKA activity possible. Active PKA phosphorylates phosphodiesterase enzyme (PDE), which hydrolyses cAMP molecules to Adenosine monophosphate (AMP) [64]. Activity of PDE

depends upon the activity of PKA since it can be dephosphorylated by Protein Phosphatases such as PP2A. Thus, PDE activity depends on the PKA activity to reduce the net PKA activity. This behaviour of PDE is modeled by the second term $-\beta u$ in eq.(2), where the parameter β signifies the strength of the PDE action on PKA deactivation. Upon parallel fiber stimulation, glutamate activates the mGluR₇ receptor, which activates G-protein to produce a G_α subunit to block the cAMP molecule production. This behaviour is modeled by the prefactor $1/(\alpha + x)$ of $u(u_0 - u)$ in eq.(2), where x denotes G-protein activity. The prefactor corresponds to blocking of AC enzyme, which is obtained from Hill's equation with Hill's coefficient equal to 1 as only one unit of G_α protein binds to AC. For more details on Hill's equation, see Materials and Methods. The constant α denotes the disassociation constant K_d of AC and G_α subunit and it has a small value due to their strong bonding. τ_1 signifies the overall time scale of the PKA dynamics. Since according to the proposed mechanism different conditional responses are the result of changes in the dynamics of PKA activation and deactivation, the value of τ_1 will increase or decrease for a conditional response of longer or shorter duration, respectively.




Fig 3. A conceptual minimal model of conditional response in the Purkinje cell.

Eq.(3) models the activity of the mGluR₇ receptor. The first term is a cubic polynomial, which captures the switching property of the mGluR₇ receptor corresponding to the unaltered conditional response with changing the CS durations. In the cubic polynomial $v_0 \gg v_1 \gtrsim v_2$, where v_0 signifies the amount of receptors, which are associated with the G-protein, v_1 is the threshold activity that needs to be crossed to initiate the conditional response and v_2 is the net finite intrinsic activity of the receptor [65]. As each mGluR₇ receptor can form a protein complex with only one unit of G-protein, the value of v_0 remains constant as we assumed that the amount of G-protein is constant for all different conditional response trainings. The values of v_1 and v_2 depend upon intrinsic properties of the receptor itself, which we also assumed to be constant. After deactivation of PKA by G-protein activity, activity of PP1 rises, which dephosphorylates the receptor resulting in the blockade of the receptor's active site to activate G-proteins further. This is modeled as $-\gamma w v$ in eq.(3), which denotes the lowering of net receptor activity due to dephosphorylation by PP1. It is a product

because PP1 interacts with the mGluR₇ receptor directly during dephosphorylation. The factor γ denotes the strength of the influence of PP1 on mGluR₇ receptors. As Spinophilin binds PP1 close to the mGluR₇ receptor by binding to RGS8 [66] and not the receptor, the strength of influence of PP1 on mGluR₇, i.e., the value of γ , can be assumed constant for different conditional responses. Finally, I denotes the strength of the CS in the form of glutamate release from parallel fibers to activate mGluR₇ receptors and τ_2 signifies the overall time scale of receptor activation and deactivation. The value of τ_2 is taken to be smaller than the fastest onset of the conditional response observed in the experiment [12].

Eq.(4) models the activity of the PP1 protein. Its activity is regulated by PKA as a suppressor by phosphorylating DARPP-32 or I-1 protein, which is modeled as $-\delta uw$. It is a product because PKA interacts directly with the PP1 regulatory protein, which binds to PP1 to block its activity. The factor δ signifies the strength of the PKA influence on PP1 activity. As the activity of PKA decreases, PP1 activity rises by the action of other Phosphatase proteins such as PP2A/B [45] and by itself [67]. The rise due to other Phosphatase proteins is given by $\eta(w_0 - w)$, while the rise of PP1 by itself is given by $w(w_0 - w)$, where w_0 is the maximum activity of PP1 and the factor η controls the strength of the influence of other Phosphatase proteins on the rise of PP1 activity. τ_3 signifies the overall time scale of PP1 activation and deactivation. For simplicity, we assume all the variables in this equation to be constant.

Eq.(5) models G-protein activity. Since we assumed that the amount of G-protein on the synapse is constant, net G-protein activity will be the same for all different conditional responses. As a result, G-protein activity will be limited to ' v_0 ', which is the total amount of G-protein present at the synapse and modeled as $(v - x)$. τ_4 signifies the time scale of activation and deactivation of the G-protein. As different conditional response trainings involve a different amount of mGluR₇ receptors, the net dynamics of G-protein activation and deactivation by RGS8 is affected by extra mGluR₇ receptors interacting with TECs. Therefore, depending upon training, the value of τ_4 can be small or big. This will result in a short delay or a long delay in the onset of the conditional response, respectively. When v reduces due to PP1 activity, $(v - x) < 0$. This signifies the deactivation of the G-protein due to the action of the RGS8 protein.

In eqs. (3) and (4), the terms $-\gamma wv$ and $-\delta uw$ signify the interaction of PP1 with mGluR₇ and PKA with PP1, respectively. Yet, there are no corresponding terms in eq.(2) of PKA and eq.(4) of PP1 because those interactions are enzymatic in nature and have very short time scales compared to the response, which we are trying to model. Hence, the activity of PKA and PP1 does not change when they interact with other proteins. Note that the τ_i factors on the LHS of each equation make them dimensionless.

Results

Properties of the model

Since experimental results have shown that the conditional response is independent of CS durations, the activation of the G-protein must also satisfy this property as it regulates GIRK ion channels. This behavior is indeed captured by our mathematical model. It also successfully captures the dynamics of other biochemicals - PKA, mGluR₇, PP1 and G-protein as proposed in the mechanism that is shown in Fig. (4). As per our proposed mechanism, before CS, PKA activity is high while activity of mGluR₇ receptor, G-protein and PP1 is low. Upon CS, mGluR₇ receptors become active, which in turn activate G-protein. Due to activation of G-protein, PKA activity drops down, which causes a rise in the PP1 activity. When PP1 activity is sufficiently high, it causes deactivation of mGluR₇ receptors, which then causes deactivation of the G-protein by

RGS8. When the stimulus is turned off, activities of various proteins return back to their original states as shown in the top panel of Fig. (4). However, if the stimulus remains on for a long time, then even very low G-protein activity can prevent the rise of PKA activity to a value that would be high enough to block PP1. As a result, PP1 activity will be crucial to cause dephosphorylation of mGluR₇ receptors and to block their active sites to prevent the initiation of another conditional response as shown in the middle panel of Fig. (4). Very faint G-protein activity in case of long duration CS can be observed in the bottom panel of Fig. (4). This is enough to prevent reactivation of the conditional response.

454
455
456
457
458
459
460
461
462




Fig 4. Temporal behaviour of PKA, mGluR₇, PP1 and G-protein. Time varying dimensionless quantities of PKA, mGluR₇, PP1 and G-protein upon short (top panel) and long (middle panel) stimulus durations represented by black horizontal bar at the bottom of each panel. In the bottom panel, activity of the G-protein is shown as in the upper two panels but for two different stimulus durations together, indicated by the red and black bars at the bottom. Both responses are almost identical implying that the G-protein activity is indeed independent of stimulus duration. Parameters for our dynamical model of PKA, mGluR₇, G-protein and PP1 are: $\alpha = 0.1$, $u_0 = 11.0$, $\beta = 48$, $v_0 = 4.0$, $v_1 = 1.01$, $v_2 = 1.0$, $\gamma = 2.0$, $\delta = 5.0$, $\eta = 0.2$, $w_0 = 6.0$, $I = 0.1$, $\tau_1 = 2100ms$, $\tau_2 = 6ms$, $\tau_3 = 60ms$, $\tau_4 = 7.9ms$.

Specifically, the chosen parameter values will determine the specific extreme values of the various variables and their temporal profiles. While these values vary with the chosen parameters, the overall properties of the model will not be affected as long as two features are preserved: i) The G-protein activation remains largely independent of the CS duration, and ii) no oscillatory response emerges. These two features are essential in order to reproduce experimental results. We have verified that these features are preserved over a large range of parameter values for our model. Specifically,

changes in size of up to 100% of the values we use in the different figures will render above mentioned two features unchanged. Thus, our conceptual model can robustly capture the main experimental results.

By combining the dynamical equations of mGluR₇, G-protein, PKA and PP1 with the Purkinje cell model, we can generate the conditional response dynamics of the Purkinje cells as shown in Figs. (5,6). In Fig.(5) we show that the suppression of firing rates during the conditional response of ISI = 150ms is independent of CS durations as observed in the experiments [12]. Result shown in Fig.(5) can be considered as an average response of the firing rate during the conditional response given the deterministic nature of our model.




Fig 5. Conditional response in the model is independent of conditional stimulus duration. The width of the light green vertical bar corresponds to the duration of the ISI = 150 msec and the black bar at the bottom signifies the conditional stimulus duration, which is short in the top panel and long in the bottom panel. Parameters for the dynamical model of PKA, mGluR₇, G-protein and PP1 are the same as in Fig. 4. For all other parameters of the Purkinje cell model, see Materials and Methods.

In order to obtain a conditional response of longer duration, more mGluR₇ receptors need to be inserted into the synapse. These extra receptors cause a rise in the value of τ_1 and τ_4 as discussed earlier. Different τ_1 and τ_4 values, which we have used for reproducing different conditional responses, are summarized in Table 1. Fig.(6) shows different long duration conditional responses, which match with the experimental results [12]. Fig.(6) shows the drop in firing rate for ISI = 200ms, while Fig.(6) shows the drop in firing rate for ISI = 400ms as an additional case. Indeed, the three conditional response firing patterns obtained from our model shown in the left panel of

Fig. (7) match with the experimental results [13]. In addition, our proposed mechanism also explains why the time-memory remains unaffected in the presence of mGluR₇ antagonist 6-(4-Methoxyphenyl)-5-methyl-3-(4-pyridinyl)isoxazolo[4,5-c]pyridin-4(5H)-one hydrochloride (MMPIP) as observed in the experiment [13]. Specifically, because of the presence of MMPIP, fewer mGluR₇ receptors are left to activate GIRK ion channels, which leads to a smaller drop in firing rate. However, reducing the net amount of active mGluR₇ does not inhibit internal interactions between receptor and other proteins involved in our proposed mechanism. Hence, the time-memory, which is encoded within effective rate constants of biochemical reactions, is unaffected by MMPIP as shown in Fig. (7). Note that in Fig. (7) the action of an increasing dose of MMPIP is simulated by decreasing the value of the parameter g_{GIRK} . As the corresponding values of g_{GIRK} have not been measured experimentally as mentioned above, we choose suitable values of g_{GIRK} for the given parameter value of the Purkinje cell model Materials and Methods.




Fig 6. Different conditional responses of the Purkinje cell obtained from the mathematical model. For $ISI = 200ms$ (top panel), the firing rate drops and then rises slowly, which is consistent with the experimental results. The values of τ_1 and τ_4 are 7.2s and 33.3ms, respectively. For higher $ISI = 400ms$ (bottom panel), the drop and rise of the firing rate is observed to be even slower compared to $ISI = 200ms$ at the top panel. The values of τ_1 and τ_4 for $ISI = 400ms$ are 18.0s and 120.0ms, respectively. The width of the light green vertical bar corresponds to the duration of the ISI interval. The black horizontal bar at the bottom represents the conditional stimulus duration.

Table 1. Model parameters for different conditional responses of the Purkinje cell.

ISI (ms)	τ_1 (s)	τ_4 (ms)
150.0	2.1	7.9
200.0	7.2	33.3
300.0	15.0	79.0
400.0	18.0	120.0




Fig 7. Conditional response profiles for different ISIs and different amounts of MMPiP. Conditional response profiles obtained from the model for different values of τ_1 and τ_4 (see Table 1, all other parameters as in Fig. 5) (left panel), and in the presence of mGluR₇ receptor's antagonist MMPiP (right panel). The latter leads to a decrease in the net amount of active mGluR₇ and, hence, the amount of active GIRK ion channels, which corresponds to smaller values of g_{GIRK} (see Eq. 1). Here, $\tau_1 = 18.0s$, $\tau_4 = 120.0ms$ and all other parameters as in Fig. 5. Note that the normalized firing activity is calculated here by taking the inverse of the time interval between two successive spikes and dividing it by the firing frequency before the onset of the conditional response.

Changing values of both τ_1 and τ_4 simultaneously is one possibility to model different conditional responses within the framework of our model. We would like to point out that changing either one of the two alone does not reproduce the experimental behavior. Based on existing studies, we have no clear evidence for changes in the value of any other model parameter given our proposed mechanism. Therefore as a first approximation, we have assumed them to not change at all.

Model predictions

Based on our proposed model, we can make two predictions that can be tested easily through experiments. 1) If PP1 is knocked out then active mGluR₇ receptors will never deactivate after they become active from CS and hence the G-protein will remain active. This implies that the Purkinje cell will not fire again after receiving CS as shown in Fig. (8). 2) On the other hand, as PKA inhibits PP1 activity, knocking out PKA activation will activate PP1, which will dephosphorylate mGluR₇ receptors and hence the G-protein cannot be activated. This implies that the Purkinje cell will not exhibit a

conditional response as shown in Fig. (8).

515




Fig 8. Model predictions for knockout experiments. In our mathematical model, PP1 can be knocked out by setting $w_0 = 0.1$ (top left panel), which prevents the Purkinje cell to fire again after the initiation of the conditional response (bottom left panel). PKA can be knocked out by setting $u_0 = 0.1$ in our model (top right panel), which prevents the Purkinje cell to initiate a conditional response (bottom right panel).

However, in reality biological cells are very robust and have redundancy mechanisms to overcome such behaviours. As a result, there might be still a weak conditional response observed after knocking out PKA or a slow deactivation of G-protein after knocking out PP1, but in both cases significant effects on the conditional response should be observed.

516
517
518
519
520

Specific experimental options to test the proposed model

521

There are various experimental options to check whether our proposed mechanism for the conditional response is valid or not, including the two model predictions mentioned above.

522
523
524

1. As PP1 desensitizes the mGluR₇ receptor during conditional response, blocking of PP1, using Okadaic acid, for example, must affect the deactivation rate of GIRK ion channels during the conditional response. This would test the first model prediction.
2. As PKA is an essential biochemical for the resensitization of the receptor and maintaining low PP1 activity, reducing PKA activity in the cell will prevent the Purkinje cell from suppressing its firing rate as PP1 will desensitize the receptor and therefore GIRK ion channels will not be activated. This can be verified by using, for example, cAMPS-Rp or triethylammonium salt, which will block the cAMP production and, hence, PKA. This would test the second model prediction.

525
526
527
528
529
530
531
532
533
534

3. If mGluR₁ receptors are activating PKC then blocking of mGluR₁ receptors using CPCCOEt during training will not initiate trafficking of mGluR₇ receptors and thus no conditional response should be observed even after extensive training. 535
536
4. Use of RGS8 knockout specimen should allow only long duration conditional response: Without RGS8 protein, the activation and deactivation of G-protein will be much slower and will produce only long conditional response durations. In addition, only sufficient long CS will be able to initiate the conditional response as mGluR₇ receptors will take longer time to activate G-protein in the absence of RGS8 protein, which accelerates G-protein activation. 537
538
539
540
541
542
543

Discussion 544

In this article, we introduced a potential biochemical mechanism to explain time-encoding memory formation within a single synapse of a Purkinje cell. This time-encoding memory is stored in an excitatory synapse, but it is associated with an inhibitory response, i.e., the suppression of the Purkinje cell's tonic firing rate in the presence of an excitatory stimulus, namely glutamate discharge from the parallel fiber. During conditional training, Purkinje cells imprint the time information by expressing an appropriate amount of mGluR₇ receptors on the synapse, while encoding time information in the form of effective rate constants. The memory is stored by forming a protein complex, i.e., a Time-Encoding protein Complex (TEC). Alterations of effective rate constants within TECs will change its temporal signature, while the removal of receptors from the synapse will cause memory loss. However, during retraining, the previous memory can quickly be reacquired and it becomes accessible again. Our idea of TEC is similar to the "Timer Proteins" previously proposed by Ref. [12], but in contrast, it does not require an active selection of feedforward protein activations to produce a specific conditional response. Recently, a different biochemical mechanism was proposed for time memory learning, which uses Ca⁺² ion dynamics for storing different time information [68]. That model does not incorporate the documented role of the GIRK ion channel and it also predicts faster learning for long duration conditional responses, which is not compatible with previous experimental findings [12]. 545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563

As previously mentioned, in our model the time information of the conditional response is stored in the TECs found on individual synapses, implying that the substrate or the Engram of a time memory can reside at individual synapses, not in a cell or a cell assembly. This result is in line with the synaptogenic point of view of memory substrates [10], where single synapses play a large role in memory formation. In contrast, another point of view puts more emphasis on the intrinsic plasticity of a whole neuronal cell compared to the synaptic plasticity of individual synapses [69]. Intrinsic plasticity considers changes in the electrophysiological properties of the cell by changing the expression of Voltage-dependent Ca/K ion channels and many other kinds of ion channels, which are expressed by neurons and which decide neural firing rate as well as the sensitivity of the cell upon stimulation. However, neither points of view can fully account for the development of the conditional response in the Purkinje cell, since it neither involves the formation or elimination of pf-PC synapses [12,13], nor LTD of pf-PC synapses [14] nor any change in the electrophysiological properties of the cell [12]. Thus, Purkinje cells show a novel form of synaptic plasticity and provide an example of monosynaptic memory encoding. In addition, considering this fact and that each Purkinje cell makes at least one synapse with up to 200,000 parallel fibers passing through the dendritic tree of the cell [70], the storage capacity of a Purkinje cell might be much higher than previously thought and the Purkinje cell might be considered as a multi-information storage device. Specifically, one might be able to encode a specific 564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583

time interval by stimulating only a subset of parallel fibers and encode another time interval by stimulating a separate subset of fibers. In this case, a specific time memory out of the whole set can be selectively retrieved when the respective set of parallel fibers becomes active upon stimulation, producing the conditional response for the previously encoded time interval.

One can get a rough estimate of the total number of unique time memories that can be stored by an individual Purkinje cell by taking the ratio of the collective hyperpolarization current produced by GIRK ion channels from all synapses and the minimum required hyperpolarization current in order to noticeably suppress the tonic firing rate of the cell. To determine the minimum hyperpolarization current, one can assume that its value is approximately equal to the net resurgent Na^+ current produced by the Purkinje cell as measured in [55]. As mentioned in the “Training with different ISI duration” subsection, the resurgent Na^+ ion channel has the capability to spontaneously generate rapid sequences of action potentials [55,57]. However, such resurgent currents have been observed in other neuronal classes as well, which have distinct firing activity pattern compared to the Purkinje cell [71,72] implying that resurgent Na^+ ion channels alone do not contribute to spontaneous high firing rates of the Purkinje cell as shown in [73]. As a result, our assumption for the minimum hyperpolarization current is just a first order approximation. In order to determine the collective hyperpolarization current, one needs to know the conductances of the GIRK ion channels and their respective densities on the synapses. Although there are experimentally measured values for single GIRK ion channel conductances [74,75], no absolute density quantification of GIRK ion channels for Purkinje cells has been done as far as we know. Only relative abundances of GIRK ion channels at Purkinje cell’s dendritic spines are known [25]. Hence, it is currently not possible to determine the collective hyperpolarization current and, thus, the total number of unique time memories that can be stored by an individual Purkinje cell. This remains an interesting challenge for the future.

As an alternative approach, one could aim to establish experimentally that an individual Purkinje cell can indeed store at least two different time memories at separate sets of pf-PC synapses. As discussed above, stimulating separate sets of parallel fibers can in principle initiate different conditional responses. While this can be achieved by electrodes [12], it is challenging less so in terms of potential experimental protocols for conditional training [35] but rather due to difficulties in selecting specific fibers. An alternative could be to stimulate granule cells in the Granule layer of the Cerebellum [76] since parallel fibers are axonal branches of the granule cells. By stimulating a selected sub-population of granule cells and a specific branch of the climbing fiber, a subset of pf-PC synapses of an individual Purkinje cells can be trained for a specific ISI. Stimulating granule cells may appear as a drawback as they also excite other GABAergic interneurons, namely Golgi, stellate, and basket cells, which directly or indirectly can influence Purkinje cell firing activity [76]. However, their excitation did not appear to influence the conditional response profile of the Purkinje cell as shown experimentally [12]. Hence, stimulating subsets of granule cells experimentally — potentially using optogenetics [77] — might be a good way to test the capability of a Purkinje cell as a multi-information storage device in the future.

Note added: New support for our proposed biochemical mechanism for time-encoding memory formation comes from the observation that the mGluR_1 receptor is necessary for the learning process, while it is not for the activation of the conditional response [78]. Despite using a different experimental approach, it basically verifies point 3 listed in the section “Specific experimental options to test the proposed model”. Specifically, the observation matches with our proposed mechanism since the latter assumes that the mGluR_1 receptor is responsible for the learning via facilitating trafficking of the mGluR_7

receptor to the synapses. Further experimental verification of our proposed mechanisms potentially along the lines outlined above remains an exciting challenge for the future.

Materials and Methods

Purkinje cell model

To model the conditional response behavior of the Purkinje cell, we start with an established dynamical model of the Purkinje cell [58] as summarized by eqs.(7) to (11). Specifically, it aims to model the dynamics of the Purkinje cell by incorporating many properties of the Purkinje cell within a realistic biophysical framework. In contrast to the original formulation [58], eqs.(7) to (11) already incorporate the features specific to our situation: In eq.(7), the input current term I_i , which originally signified an external electrical stimulus, now signifies the intrinsic current causing the tonic firing of the Purkinje cell [79,80]. Moreover, we added the influence of the GIRK ion channel in eq.(8), which only becomes relevant after training — see also eq.(1). Here, g_{GIRK} is the net conductance of GIRK ion channels per unit area, h_{GIRK} is the gating parameter and V_{GIRK} is the voltage dependence of the GIRK ion channel obtained from the I-V characteristics curve of the ion channel [59].

Except for g_{GIRK} , all values of the model are taken from [58] and listed below. As far as we know, there is no literature on the specific g_{GIRK} values. As a result, we chose a value of g_{GIRK} that matches the experimentally observed conditional response profiles.

Somatic voltage equation:

$$C_s \frac{dV_s}{dt} = \frac{(V_d - V_s)}{R} - g_{Na} m_\infty h (V_s - E_{Na}) - g_K (1 - h) (V_s - E_K) - g_{leak} (V_s - E_{leak}) - g_{I_H} i h (V_s - E_{ih}) + I_i \quad (7)$$

Dendritic voltage equation:

$$C_d \frac{dV_d}{dt} = \frac{(V_s - V_d)}{R} - g_{leak} (V_d - E_{leak}) - g_{Kd(slow)} n_d (V_d - E_K) - g_{GIRK} h_{GIRK} V_{GIRK} (V_d) \quad (8)$$

$$V_{GIRK} (V_d) = -0.02(1.3V_d + 50.0)/(1.0 + \exp((V_d + 40)/10.0))$$

Na⁺ activation equation:

$$m_\infty = \frac{1}{1 + \exp[-(V - V_{1/2})/k]}, \quad V_{1/2} = -40.0mV, \quad k = 3.0mV$$

$$\frac{dh}{dt} = \frac{h_\infty - h}{\tau_h} = \frac{1}{1 + \exp[-(V - V_{1/2})/k]}, \quad V_{1/2} = -40.0mV, \quad k = -3.0mV \quad (9)$$

$$\tau_h(V) = \frac{295.4}{4(V + 50)^2 + 400} + 0.012$$

Hyperpolarizing activated cation current (I_h):

$$\frac{dI_h}{dt} = \frac{I_{h_\infty} - I_h}{\tau_{I_h}} = \frac{1}{1 + \exp[-(V - V_{1/2})/k]}, \quad V_{1/2} = -80.0mV, \quad k = -3.0mV, \quad (10)$$

$$\tau_{I_h} = 100ms$$

Slow K⁺ activation equation:

$$\frac{dn_d}{dt} = \frac{n_{d_\infty} - n_d}{\tau_{nd}} = \frac{1}{1 + \exp[-(V - V_{1/2})/k]}, \quad V_{1/2} = -35.0mV, \quad k = 3.0mV \quad (11)$$

$$\tau_{ih} = 15ms$$

Parameters value of the Purkinje cell model

$$\begin{aligned}C_s &= 1.5\mu F/cm^2, C_d = 1.5\mu F/cm^2, R = 0.75, g_{Na} = 40mS/cm^2, \\g_{Ks} &= 8.75mS/cm^2, g_{IH} = 0.03mS/cm^2, g_{Kd(slow)} = 12mS/cm^2 \\g_{leak} &= 0.032mS/cm^2, E_{Na^+} = 45mV, E_{K^+} = -95mV, E_{IH} = -20.0, \\E_{leak} &= -77mV, g_{GIRK} = 0.38mS/cm^2, I_i = 0.198\mu A\end{aligned}$$

Hill's equation

Hill's equation gives the fraction of protein saturated by a ligand at a given concentration of ligand in a solution [81]. Since, in the case of Acetyl cyclase (AC), only one unit of G_α subunit binds to it, Hill's coefficient will be 1. The fraction of AC bound by a G_α subunit to the total available amount of AC at a concentration $[G_\alpha]$ is given by

$$\frac{[Blocked\ AC]}{[AC]} = \frac{[G_\alpha]}{K_d + [G_\alpha]}, \quad (12)$$

where, K_d is the disassociation constant of AC and G_α subunit. However, we are interested in free AC. So, the fraction of free AC will be given by

$$\frac{[Active\ AC]}{[AC]} = 1 - \frac{[Blocked\ AC]}{[AC]} = \frac{K_d}{K_d + [G_\alpha]}. \quad (13)$$

Since the concentration of $[G_\alpha]$ is proportional to its activity, eq.(13) can be written as

$$\frac{[Active\ AC]}{[AC]} = \frac{K_d}{K_d + x}, \quad (14)$$

where x denotes the G-protein activity.

Acknowledgments

Authors thanks Dr. Ray W. Turner at the University of Calgary for helpful discussion.

References

1. Liu X, Ramirez S, Pang PT, Puryear CB, Govindarajan A, Deisseroth K, et al. Optogenetic stimulation of a hippocampal engram activates fear memory recall. *Nature*. 2012;484:381. doi:10.1038/nature11028.
2. Yiu A, Mercaldo V, Yan C, Richards B, Rashid A, Hsiang HL, et al. Neurons Are Recruited to a Memory Trace Based on Relative Neuronal Excitability Immediately before Training. *Neuron*. 2014;83:722–735. doi:10.1016/j.neuron.2014.07.017.
3. Kim J, Kwon JT, Kim HS, Josselyn SA, Han JH. Memory recall and modifications by activating neurons with elevated CREB. *Nature Neuroscience*. 2014;17:65. doi:10.1038/nn.3592.
4. Tonegawa S, Liu X, Ramirez S, Redondo R. Memory Engram Cells Have Come of Age. *Neuron*. 2015;87:918–931. doi:10.1016/j.neuron.2015.08.002.
5. Tonegawa S, Morrissey MD, Kitamura T. The role of engram cells in the systems consolidation of memory. *Nature Reviews Neuroscience*. 2018;19:485–498. doi:10.1038/s41583-018-0031-2.

6. Yang G, Pan F, Gan WB. Stably maintained dendritic spines are associated with lifelong memories. *Nature*. 2009;462:920. doi:10.1038/nature08577.
7. Ryan TJ, Roy DS, Pignatelli M, Arons A, Tonegawa S. Engram cells retain memory under retrograde amnesia. *Science*. 2015;348:1007–1013. doi:10.1126/science.aaa5542.
8. Bannerman DM, Bus T, Taylor A, Sanderson DJ, Schwarz I, Jensen V, et al. Dissecting spatial knowledge from spatial choice by hippocampal NMDA receptor deletion. *Nature Neuroscience*. 2012;15:1153. doi:10.1038/nn.3166.
9. Nabavi S, Fox R, Proulx CD, Lin JY, Tsien RY, Malinow R. Engineering a memory with LTD and LTP. *Nature*. 2014;511:348. doi:10.1038/nature13294.
10. Poo Mm, Pignatelli M, Ryan TJ, Tonegawa S, Bonhoeffer T, Martin KC, et al. What is memory? The present state of the engram. *BMC Biology*. 2016;14:40. doi:10.1186/s12915-016-0261-6.
11. Gallistel CR. The Coding Question. *Trends in Cognitive Sciences*. 2017;21:498–508. doi:10.1016/j.tics.2017.04.012.
12. Johansson F, Jirenhed DA, Rasmussen A, Zucca R, Hesslow G. Memory trace and timing mechanism localized to cerebellar Purkinje cells. *Proceedings of the National Academy of Sciences*. 2014;111:14930–14934. doi:10.1073/pnas.1415371111.
13. Johansson F, Carlsson HE, Rasmussen A, Yeo C, Hesslow G. Activation of a Temporal Memory in Purkinje Cells by the mGluR7 Receptor. *Cell Reports*. 2015;13:1741–1746. doi:10.1016/j.celrep.2015.10.047.
14. Johansson F, Jirenhed DA, Rasmussen A, Zucca R, Hesslow G. Absence of Parallel Fibre to Purkinje Cell LTD During Eyeblink Conditioning. *Scientific Reports*. 2018;8:14777. doi:10.1038/s41598-018-32791-7.
15. Hesslow G, Jirenhed DA, Rasmussen A, Johansson F. Classical conditioning of motor responses: What is the learning mechanism? *Neural Networks*. 2013;47:81–87. doi:10.1016/j.neunet.2013.03.013.
16. McCormick DA, Clark GA, Lavond DG, Thompson RF. Initial localization of the memory trace for a basic form of learning. *Proceedings of the National Academy of Sciences*. 1982;79:2731–2735. doi:10.1073/pnas.79.8.2731.
17. Jirenhed DA, Bengtsson F, Hesslow G. Acquisition, Extinction, and Reacquisition of a Cerebellar Cortical Memory Trace. *The Journal of Neuroscience*. 2007;27:2493–2502. doi:10.1523/jneurosci.4202-06.2007.
18. Hesslow G, Yeo CH. A Neuroscientist's Guide to Classical Conditioning. 2002; p. 86–146. doi:10.1007/978-1-4419-8558-3_4.
19. Marr D. A Theory of Cerebellar Cortex. *The Journal of Physiology*. 1969;202:437–470. doi:10.1113/jphysiol.1969.sp008820.
20. Medina JF, Garcia KS, Nores WL, Taylor NM, Mauk MD. Timing Mechanisms in the Cerebellum: Testing Predictions of a Large-Scale Computer Simulation. *Journal of Neuroscience*. 2000;20:5516–5525. doi:10.1523/jneurosci.20-14-05516.2000.

21. Lepora NF, Porrill J, Yeo CH, Dean P. Sensory Prediction or Motor Control? Application of Marr–Albus Type Models of Cerebellar Function to Classical Conditioning. *Frontiers in Computational Neuroscience*. 2010;4:140. doi:10.3389/fncom.2010.00140.
22. Johansson F. A Purkinje cell Timing Mechanism. On the Physical Basis of a Temporal Duration Memory. Lund University; 2015.
23. Phillips T, Makoff A, Murrison E, Mimmack M, Waldvogel H, Faull R, et al. Immunohistochemical localisation of mGluR7 protein in the rodent and human cerebellar cortex using subtype specific antibodies. *Molecular Brain Research*. 1998;57:132–141. doi:10.1016/s0169-328x(98)00081-3.
24. Sadjia R, Alagem N, Reuveny E. Gating of GIRK Channels Details of an Intricate, Membrane-Delimited Signaling Complex. *Neuron*. 2003;39:9–12. doi:10.1016/s0896-6273(03)00402-1.
25. Fernández-Alacid L, Aguado C, Ciruela F, Martín R, Colí J, Cabañero MJ, et al. Subcellular compartment-specific molecular diversity of pre- and post-synaptic GABAB-activated GIRK channels in Purkinje cells. *Journal of Neurochemistry*. 2009;110:1363–1376. doi:10.1111/j.1471-4159.2009.06229.x.
26. Syrovatkina V, Alegre KO, Dey R, Huang XY. Regulation, Signaling, and Physiological Functions of G-Proteins. *Journal of Molecular Biology*. 2016;428:3850–3868. doi:10.1016/j.jmb.2016.08.002.
27. Suh YH, Chang K, Roche KW. Metabotropic glutamate receptor trafficking. *Molecular and Cellular Neuroscience*. 2018;doi:10.1016/j.mcn.2018.03.014.
28. Ito M, Sakurai M, Tongroach P. Climbing fibre induced depression of both mossy fibre responsiveness and glutamate sensitivity of cerebellar Purkinje cells. *The Journal of Physiology*. 1982;324:113–134. doi:10.1113/jphysiol.1982.sp014103.
29. Nusser Z, Mulvihill E, Streit P, Somogyi P. Subsynaptic segregation of metabotropic and ionotropic glutamate receptors as revealed by immunogold localization. *Neuroscience*. 1994;61:421–427. doi:10.1016/0306-4522(94)90421-9.
30. Telser A. *Molecular Biology of the Cell*, 4th Edition. Shock. 2002;18:289. doi:10.1097/00024382-200209000-00015.
31. Tempia F, Miniaci MC, Anchisi D, Strata P. Postsynaptic Current Mediated by Metabotropic Glutamate Receptors in Cerebellar Purkinje Cells. *Journal of Neurophysiology*. 1998;80:520–528. doi:10.1152/jn.1998.80.2.520.
32. Canepari M, Papageorgiou G, Corrie JET, Watkins C, Ogden D. The conductance underlying the parallel fibre slow EPSP in rat cerebellar Purkinje neurones studied with photolytic release of L-glutamate. *The Journal of Physiology*. 2001;533:765–772. doi:10.1111/j.1469-7793.2001.00765.x.
33. Leitges M, Kovac J, Plomann M, Linden DJ. A Unique PDZ Ligand in PKC α Confers Induction of Cerebellar Long-Term Synaptic Depression. *Neuron*. 2004;44:585–594. doi:10.1016/j.neuron.2004.10.024.
34. Wetmore DZ, Jirenhed DA, Rasmussen A, Johansson F, Schnitzer MJ, Hesslow G. Bidirectional Plasticity of Purkinje Cells Matches Temporal Features of Learning. *The Journal of Neuroscience*. 2014;34:1731–1737. doi:10.1523/jneurosci.2883-13.2014.



35. Boele HJ, Peter S, Brinke MMT, Verdonschot L, IJpelaar ACH, Rizopoulos D, et al. Impact of parallel fiber to Purkinje cell long-term depression is unmasked in absence of inhibitory input. *Science Advances*. 2018;4:eaas9426. doi:10.1126/sciadv.aas9426.
36. Jirenhed DA, Hesslow G. Learning Stimulus Intervals-Adaptive Timing of Conditioned Purkinje Cell Responses. *The Cerebellum*. 2011;10:523–535. doi:10.1007/s12311-011-0264-3.
37. Zhang Q, Pacheco MA, Douppnik CA. Gating properties of girk channels activated by $\alpha\omega$ - and $G\alpha i$ -Coupled Muscarinic m2 Receptors in *Xenopus* Oocytes: The Role of Receptor Precoupling in RGS Modulation. *The Journal of Physiology*. 2002;545:355–373. doi:10.1113/jphysiol.2002.032151.
38. Saitoh O, Kubo Y, Miyatani Y, Asano T, Nakata H. RGS8 accelerates G-protein-mediated modulation of K⁺ currents. *Nature*. 1997;390:525–529. doi:10.1038/37385.
39. Saitoh O, Odagiri M. RGS8 expression in developing cerebellar Purkinje cells. *Biochemical and Biophysical Research Communications*. 2003;309:836–842. doi:10.1016/j.bbrc.2003.08.083.
40. Jeong SW, Ikeda SR. Differential regulation of G protein-gated inwardly rectifying K⁺ channel kinetics by distinct domains of RGS8. *The Journal of Physiology*. 2001;535:335–347. doi:10.1111/j.1469-7793.2001.00335.x.
41. Suh YH, Park JY, Park S, Jou I, Roche PA, Roche KW. Regulation of Metabotropic Glutamate Receptor 7 (mGluR7) Internalization and Surface Expression by Ser/Thr Protein Phosphatase 1. *Journal of Biological Chemistry*. 2013;288:17544–17551. doi:10.1074/jbc.m112.439513.
42. Fierro L, DiPolo R, Llano I. Intracellular calcium clearance in Purkinje cell somata from rat cerebellar slices. *The Journal of Physiology*. 1998;510:499–512. doi:10.1111/j.1469-7793.1998.499bk.x.
43. Neil SM, Lakey T, Tomlinson S. Calmodulin regulation of adenylate cyclase activity. *Cell Calcium*. 1985;6:213–226. doi:10.1016/0143-4160(85)90007-7.
44. Sorensen SD, Macek TA, Cai Z, Saugstad JA, Conn PJ. Dissociation of Protein Kinase-Mediated Regulation of Metabotropic Glutamate Receptor 7 (mGluR7) Interactions with Calmodulin and Regulation of mGluR7 Function. *Molecular Pharmacology*. 2002;61:1303–1312. doi:10.1124/mol.61.6.1303.
45. Aggen JB, Nairn AC, Chamberlin R. Regulation of protein phosphatase-1. *Chemistry & Biology*. 2000;7:R13–R23. doi:10.1016/s1074-5521(00)00069-7.
46. Greenwald EC, Saucerman JJ. Bigger, Better, Faster: Principles and Models of AKAP Anchoring Protein Signaling. *Journal of Cardiovascular Pharmacology*. 2011;58:462. doi:10.1097/fjc.0b013e31822001e3.
47. Allen PB, Ouimet CC, Greengard P. Spinophilin, a novel protein phosphatase 1 binding protein localized to dendritic spines. *Proceedings of the National Academy of Sciences*. 1997;94:9956–9961. doi:10.1073/pnas.94.18.9956.
48. Ouimet CC, Katona I, Allen P, Freund TF, Greengard P. Cellular and subcellular distribution of spinophilin, a PP1 regulatory protein that bundles F-actin in dendritic spines. *Journal of Comparative Neurology*. 2004;479:374–388. doi:10.1002/cne.20313.


49. Jr HCH, Greengard P, Tung HYL, Cohen P. DARPP-32, a dopamine-regulated neuronal phosphoprotein, is a potent inhibitor of protein phosphatase-1. *Nature*. 1984;310:310503a0. doi:10.1038/310503a0.
50. Hausdorff WP, Caron MG, Lefkowitz RJ. Turning off the signal: desensitization of beta-adrenergic receptor function. *The FASEB Journal*. 1990;4:2881–2889. doi:10.1096/fasebj.4.11.2165947.
51. Boudin H, Doan A, Xia J, Shigemoto R, Haganir RL, Worley P, et al. Presynaptic Clustering of mGluR7a Requires the PICK1 PDZ Domain Binding Site. *Neuron*. 2000;28:485–497. doi:10.1016/s0896-6273(00)00127-6.
52. Johansson F, Hesslow G, Medina JF. Mechanisms for motor timing in the cerebellar cortex. *Current Opinion in Behavioral Sciences*. 2016;8:53–59. doi:10.1016/j.cobeha.2016.01.013.
53. Jirenhed DA, Rasmussen A, Johansson F, Hesslow G. Learned response sequences in cerebellar Purkinje cells. *Proceedings of the National Academy of Sciences*. 2017;114:6127–6132. doi:10.1073/pnas.1621132114.
54. Häusser M, Clark BA. Tonic Synaptic Inhibition Modulates Neuronal Output Pattern and Spatiotemporal Synaptic Integration. *Neuron*. 1997;19(3):665–678. doi:10.1016/s0896-6273(00)80379-7.
55. Raman IM, Bean BP. Resurgent Sodium Current and Action Potential Formation in Dissociated Cerebellar Purkinje Neurons. *Journal of Neuroscience*. 1997;17:4517–4526. doi:10.1523/jneurosci.17-12-04517.1997.
56. Schaller KL, Caldwell JH. Expression and distribution of voltage-gated sodium channels in the cerebellum. *The Cerebellum*. 2003;2:2–9. doi:10.1080/14734220309424.
57. Khaliq ZM, Gouwens NW, Raman IM. The Contribution of Resurgent Sodium Current to High-Frequency Firing in Purkinje Neurons: An Experimental and Modeling Study. *Journal of Neuroscience*. 2003;23:4899–4912. doi:10.1523/jneurosci.23-12-04899.2003.
58. Fernandez FR, Engbers JDT, Turner RW. Firing Dynamics of Cerebellar Purkinje Cells. *Journal of Neurophysiology*. 2007;98:278–294. doi:10.1152/jn.00306.2007.
59. Bichet D, Haass FA, Jan LY. Merging functional studies with structures of inward-rectifier K⁺ channels. *Nature Reviews Neuroscience*. 2003;4:957–967. doi:10.1038/nrn1244.
60. Loew LM. Where does all the PIP2 come from? *The Journal of Physiology*. 2007;582:945–951. doi:10.1113/jphysiol.2007.132860.
61. Feliciello A, Gottesman ME, Avvedimento EV. The biological functions of A-kinase anchor proteins1 1Edited by P. E. Wright. *Journal of Molecular Biology*. 2001;308:99–114. doi:10.1006/jmbi.2001.4585.
62. Dodge-Kafka KL, Soughayer J, Pare GC, Michel JJC, Langeberg LK, Kapiloff MS, et al. The protein kinase A anchoring protein mAKAP coordinates two integrated cAMP effector pathways. *Nature*. 2005;437:574–578. doi:10.1038/nature03966.


63. Shiraishi-Yamaguchi Y, Furuichi T. The Homer family proteins. *Genome Biology*. 2007;8:206. doi:10.1186/gb-2007-8-2-206.
64. Fimia GM, Sassone-Corsi P. Cyclic AMP signalling. *Journal of Cell Science*. 2001;114:1971–2.
65. Kammermeier PJ. Constitutive activity of metabotropic glutamate receptor 7. *BMC Neuroscience*. 2015;16:17. doi:10.1186/s12868-015-0154-6.
66. Fujii S, Yamazoe G, Itoh M, Kubo Y, Saitoh O. Spinophilin inhibits the binding of RGS8 to M1-mAChR but enhances the regulatory function of RGS8. *Biochemical and Biophysical Research Communications*. 2008;377:200–204. doi:10.1016/j.bbrc.2008.09.096.
67. Ingebritsen TS, Cohen P. The Protein Phosphatases Involved in Cellular Regulation. *European Journal of Biochemistry*. 1983;132:255–261. doi:10.1111/j.1432-1033.1983.tb07357.x.
68. Majoral D, Zemmar A, Vicente R. A Model for Time Interval Learning in the Purkinje Cell. *bioRxiv*. 2018; p. 448027. doi:10.1101/448027.
69. Titley HK, Brunel N, Hansel C. Toward a Neurocentric View of Learning. *Neuron*. 2017;95:19–32. doi:10.1016/j.neuron.2017.05.021.
70. Tyrrell T, Willshaw D. Cerebellar cortex: its simulation and the relevance of Marr's theory. *Philosophical Transactions of the Royal Society of London Series B: Biological Sciences*. 1992;336:239–257. doi:10.1098/rstb.1992.0059.
71. Mossadeghi B, Slater N. Persistent and resurgent sodium currents in cerebellar unipolar brush cells. In: *Soc Neurosci Abstr*. vol. 24; 1998. p. 1078.
72. Raman IM, Gustafson AE, Padgett D. Ionic Currents and Spontaneous Firing in Neurons Isolated from the Cerebellar Nuclei. *Journal of Neuroscience*. 2000;20:9004–9016. doi:10.1523/jneurosci.20-24-09004.2000.
73. Akemann W, Knöpfel T. Interaction of Kv3 Potassium Channels and Resurgent Sodium Current Influences the Rate of Spontaneous Firing of Purkinje Neurons. *The Journal of Neuroscience*. 2006;26:4602–4612. doi:10.1523/jneurosci.5204-05.2006.
74. Kofuji P, Davidson N, Lester HA. Evidence that neuronal G-protein-gated inwardly rectifying K⁺ channels are activated by G beta gamma subunits and function as heteromultimers. *Proceedings of the National Academy of Sciences*. 1995;92:6542–6546. doi:10.1073/pnas.92.14.6542.
75. Kubo Y, Adelman JP, Clapham DE, Jan LY, Karschin A, Kurachi Y, et al. International Union of Pharmacology. LIV. Nomenclature and Molecular Relationships of Inwardly Rectifying Potassium Channels. *Pharmacological Reviews*. 2005;57:509–526. doi:10.1124/pr.57.4.11.
76. Purves D, Augustine G, Fitzpatrick D, et al. *Circuits within the Cerebellum*. U.S. National Library of Medicine; 1970. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK10865/>.
77. Yamanaka A. Optogenetics, Light-Sensing Proteins and Their Applications. 2015; p. 241–248. doi:10.1007/978-4-431-55516-2_16.


78. Nakao H, Kishimoto Y, Hashimoto K, Kitamura K, Yamasaki M, Nakao K, et al. mGluR1 in cerebellar Purkinje cells is essential for the formation but not expression of associative eyeblink memory. *Scientific Reports*. 2019;9:7353. doi:10.1038/s41598-019-43744-z.
79. Häusser M, Clark BA. Tonic Synaptic Inhibition Modulates Neuronal Output Pattern and Spatiotemporal Synaptic Integration. *Neuron*. 1997;19:665–678. doi:10.1016/s0896-6273(00)80379-7.
80. Shin SL, Hoebeek FE, Schonewille M, Zeeuw CID, Aertsen A, Schutter ED. Regular Patterns in Cerebellar Purkinje Cell Simple Spike Trains. *PLoS ONE*. 2007;2:e485. doi:10.1371/journal.pone.0000485.
81. Gesztelyi R, Zsuga J, Kemeny-Beke A, Varga B, Juhasz B, Tosaki A. The Hill equation and the origin of quantitative pharmacology. *Archive for History of Exact Sciences*. 2012;66:427–438. doi:10.1007/s00407-012-0098-5.


Normalized firing activity %















→ Activation

—| Inhibition

★ Active state

dep. Dephosphorylated state

pho. Phosphorylated state




Before Training

Parallel fiber

-  mGluR7
-  AMPA
-  GIRK
-  G-protein
-  glutamate

Climbing fiber

Purkinje cell's dendritic branch



After Training

Parallel fiber

Climbing fiber

Purkinje cell's dendritic branch

