


Assessment of Pacific mackerel (*Scomber japonicus*) for U.S. management in the 2023-24 and 2024-25 fishing years

Peter T. Kuriyama¹
Juan P. Zwolinski²
Caitlin Allen Akselrud¹
Kevin T. Hill¹

¹ Fisheries Resources Division, Southwest Fisheries Science Center, NOAA National Marine Fisheries Service, 8901 La Jolla Shores Drive, La Jolla, CA 92037, USA

²Institute of Marine Sciences University of California Santa Cruz, Earth and Marine Sciences Building, Santa Cruz, CA 95064, USA (affiliated with SWFSC)



Assessment of Pacific mackerel (*Scomber japonicus*) for U.S. management in the 2023-24 and 2024-25 fishing years

Contents

Executive Summary	i
Stock	i
Catches	i
Data and Assessment	ii
Spawning Stock Biomass and Recruitment	iii
Stock Biomass for PFMC Management	v
Exploitation Status	vi
Ecosystem Considerations	viii
Harvest Control Rules	viii
Management Performance	ix
Research and Data Needs	x
1 Introduction	1
1.1 Stock Structure and Management Units	1
1.2 Distribution and Movement	1
1.3 Life History	2
1.4 Fishery Descriptions	4
1.5 Ecosystem Considerations	5
1.6 Management History	6
1.7 Management Performance	7
2 Data	7
2.1 Fishery-dependent data	7
2.1.1 Landings	8
2.1.2 Age compositions	9
2.1.3 Ageing error	10

2.1.4	Empirical weight-at-age	11
2.2	Fishery-independent data: Acoustic-trawl survey	11
2.2.1	Overview	11
2.2.2	Index of Abundance	11
2.2.3	Age compositions	13
2.2.4	Ageing error	13
2.2.5	Empirical weight-at-age	14
2.3	Nearshore sampling	14
2.4	Biological Parameters	15
2.4.1	Stock Structure	15
2.4.2	Growth	15
2.4.3	Maturity	15
2.4.4	Natural mortality	15
2.5	Available data sets not used in assessment	16
3	Stock Assessment Model	17
3.1	History of modeling approaches	17
3.2	2019 STAR Panel Recommendations	17
3.3	Base model description	21
3.3.1	Time period and time step	21
3.3.2	Forecast	21
3.3.3	Stock-recruit relationship	22
3.3.4	Catchability	22
3.3.5	Selectivity	23
3.3.6	Likelihoods components and model parameters	23
3.3.7	Bridging analysis	23
3.4	Base model results	23
3.4.1	Likelihoods and quantities of interest	23
3.4.2	Selectivity estimates and fits to fishery and survey age-compositions	24
3.4.3	Fit to survey index of abundance	24
3.4.4	Stock-recruitment relationship	24
3.4.5	Population numbers- and biomass-at-age estimates	24

3.4.6	Biomass and recruitment	24
3.4.7	Fishing mortality	25
3.5	Modeling Diagnostics	25
3.5.1	Convergence	25
3.5.2	Historical analysis	25
3.5.3	Likelihood profiles	25
3.5.4	Sensitivity to alternative data weighting	26
3.5.5	Sensitivity to alternative catch values	26
3.5.6	Sensitivity to 2022 and 2023 AT survey biomass values	26
3.5.7	Retrospective analysis	26
4	Harvest Control Rules	27
5	Research and Data Needs	28
6	Acknowledgements	28
7	Tables	29
8	Figures	51
9	References	96
10	Appendix A: Age and Maturity Assessment of Pacific mackerel (<i>Scomber japonicus</i>)	102
11	Appendix B: Supplementary details following STAR Panel	118
11.1	Justification for time-varying catchability (Q)	118
11.2	Historical catch time series	120
11.3	Density-dependence in Pacific mackerel	
11.4	Difference in ageing methods between the fishery and survey data	

Executive Summary

The following Pacific mackerel (*Scomber japonicus*) stock assessment was reviewed at a STAR Panel in April 2023.

Stock

This assessment focuses on the Pacific mackerel sub-stock that is found along the Pacific coast north of Punta Abreojos (Baja California) that extends north to areas off southern California, and even further during favorable oceanographic periods to waters off the U.S. Pacific Northwest. This sub-stock is harvested by fishermen in the U.S. and Mexico, and is the population considered in this assessment. Stock structure of the species off the Pacific coast of North America is not known definitively.

Catches

The assessment includes commercial and recreational landings from calendar years 2008 to 2022 and from Mexico and the US. Mexico landings reflect catches off Baja California from commercial purse seine fleets operating off Ensenada and in Magdalena Bay. US landings include values from commercial fishing in California, Oregon, Washington, bycatch from the Pacific whiting at-sea fishery, and recreational catches from California. Landings from each region by model year are shown in Table [ES-1](#).

Table ES-1: Landings (mt) of Pacific mackerel by region and fishing year from 2008-2022. Mexican landings were from Magdalena Bay, BCS (MAG) and Ensenada, BC (ENS). US landings are from California (CA), Oregon (OR), Washington (WA). Additionally, California recreational landings are included (CA-REC). The total (TOT) landings are summed across all regions and used as input to the stock assessment.

Model Year	MAG	ENS	MEX-TOT	CA	OR	WA	CA-REC	USA-TOT	TOT
2008	689	114	803	4,198	58	9	279	4,543	5,346
2009	49	0	49	3,279	54	5	269	3,607	3,656
2010	312	1,605	1,917	2,047	48	2	216	2,313	4,229
2011	1,081	1,151	2,232	1,665	202	83	124	2,074	4,306
2012	7,219	171	7,390	3,202	1,588	719	99	5,608	12,998
2013	2,071	482	2,553	11,165	438	173	133	11,909	14,462
2014	2,757	1,342	4,099	3,651	1,215	502	225	5,593	9,692
2015	3,663	5,515	9,179	4,435	7	1	243	4,686	13,865
2016	5,730	5,977	11,707	2,523	4	22	209	2,757	14,464
2017	2,224	585	2,810	1,513	45	4	245	1,808	4,617
2018	3,422	12,330	15,752	2,199	112	10	180	2,501	18,252
2019	16,777	2,297	19,074	3,783	50	5	78	3,916	22,990
2020	26,136	5,232	31,368	500	101	3	87	691	32,060
2021	7,649	1,760	9,409	847	86	0	73	1,007	10,416
2022	7,649	7,361	15,010	543	366	26	56	990	16,000

Data and Assessment

The integrated assessment model was developed using Stock Synthesis 3 (SS3; version 3.30.20), and includes fishery and survey data collected from 2008 to 2021. The 2022 AT survey value is unavailable, and as a result the model spans 2008-2021 with model forecasts for 2022-2025. An empirical catch value for 2022 is used in the forecast file. The model is based on a July-June biological year (aka ‘model year’). Catches and biological samples for the fisheries were pooled into a single fishing fleet, for which selectivity was modeled annually. A single AT survey index of abundance from SWFSC surveys (2008-2021) was included in the model.

The base model incorporates the following specifications:

- Model spans 2008-2021 with forecasts for 2022-2025;
- Sexes were combined; ages 0-8+;
- One fishery (MexCal), with annual selectivity patterns;
- The fishing fleet had age-based selectivity (time-varying and 2dAR option in SS3);
- AT survey age-based selectivity is assumed to be uniform (fully-selected) above age-1 with age-0 selectivity estimated;
- Length-based selectivity fixed at 1 for all lengths and for the AT survey and fishing fleet;
- AT survey age compositions with effective sample sizes set to 1 per cluster (externally);
- Fishery age compositions with effective sample sizes calculated by dividing the number of fish sampled by 25 (externally) and lambda weighting=1 (internally);

- Beverton-Holt stock-recruitment relationship with steepness set to 0.75;
- Natural mortality (M) estimated to be age-specific (Lorenzen);
- Recruitment deviations estimated for 2008-2021;
- Virgin recruitment estimated, and total recruitment variability (σ_R) fixed at 0.75; and
- AT survey with catchability (Q) estimated annually with deviations for 2008-2015 and one block for 2016-2021 with a prior calculated from the 2021 AT survey estimates from the US and Mexico.

Spawning Stock Biomass and Recruitment

Time series of estimated spawning stock biomass (SSB, shown as million mt) from the base model and associated 95% confidence intervals are displayed in Figure ES-1 and Table ES-2. The initial level of SSB was estimated to be 136,664 mt. SSB for 2022 is projected to be 43,864 mt and 46,167 mt in 2023 and 50,372 mt in 2024.




Figure ES-1: Estimated spawning stock biomass time series (mt) with 95% confidence intervals (dashed lines).




Figure ES-2: Time series of estimated recruitment (age-0, billions of fish) with 95% confidence intervals.

Time series of estimated recruitment (age-0, billions of fish) abundance is presented in Figure ES-2 and Table ES-2 for the base model. The initial level of recruitment was estimated to be 1,157,070 age-0 thousands of fish.

Table ES-2: Spawning stock biomass (SSB) and recruitment (1000s of fish) estimates and asymptotic standard errors for the base model.

Year	SSB	SSB SD	Recruits	Recruits SD
Virgin	108,420	20,365	917,938	314,313
Initial	136,664	84,074	1,157,070	833,104
2008	129,863	71,817	1,038,020	619,869
2009	102,687	44,607	475,290	316,083
2010	82,111	31,628	1,089,720	621,740
2011	85,120	29,231	2,180,780	994,322
2012	122,554	36,514	686,338	318,134
2013	118,786	34,710	442,058	193,156
2014	89,487	26,120	488,121	208,439
2015	60,413	17,585	642,887	250,514
2016	50,421	14,176	1,070,520	403,597
2017	63,936	16,709	249,293	113,096
2018	56,224	14,628	806,313	308,286
2019	48,433	13,255	984,076	394,732
2020	46,687	14,062	569,096	250,638
2021	36,646	13,880	473,902	225,892

Table ES-3: Total (age-0+) and summary (age-1+) biomass values (mt) estimated on July 1 of each year.

Yr	Age0+	Age1+	Age1+ SD
2008	267,376	145,702	80,025
2009	199,239	143,527	59,558
2010	231,941	104,207	39,935
2011	382,198	126,574	46,377
2012	329,161	201,137	63,538
2013	214,028	157,958	46,576
2014	173,411	111,498	33,172
2015	205,618	78,630	22,464
2016	138,890	74,037	20,323
2017	161,050	106,420	28,281
2018	93,438	71,610	18,243
2019	82,303	74,160	20,001
2020	141,663	69,479	20,093
2021	93,822	49,977	18,251
2022	122,333	46,682	18,176
2023	132,029	55,680	31,631
2024	138,272	60,785	37,941
2025	143,969	65,477	41,681

Stock Biomass for PFMC Management

Stock biomass, used for calculating annual harvest specifications, is defined as the sum of the biomass for Pacific mackerel ages one and older (age-1+, mt) at the start of the management year. Time series of estimated stock biomass from the base model are presented in Figure [ES-3](#) and Table [ES-3](#). The base model stock biomass was estimated to be 49,977 mt in 2021 and is projected to be 46,682 mt in 2022.




Figure ES-3: Estimated stock biomass (age-1+ fish; mt) time series for the base model with 95% confidence intervals.

Exploitation Status

Exploitation rate is defined as the calendar year Pacific mackerel catch divided by the total mid-year biomass (July-1, ages-0+). Based on the base model estimates, the U.S. exploitation rate has been below 30% in the model period and peaked at 27% in 2020. Exploitation rates for Pacific mackerel, calculated from the base model, are presented in Figure ES-4 and Table ES-4.




Figure ES-4: Annual exploitation rates (calendar year landings divided by July total biomass) for the base model.

Table ES-4: Annual exploitation rate (calendar year landings / June total biomass) by country and calendar year.

Calendar Year	Total biomass	Catch	Exp. Rate
2008	267,376	4,500	0.02
2009	199,239	6,002	0.03
2010	231,941	2,480	0.01
2011	382,198	4,745	0.01
2012	329,161	9,682	0.03
2013	214,028	15,100	0.07
2014	173,411	9,531	0.05
2015	205,618	11,379	0.06
2016	138,890	19,257	0.14
2017	161,050	6,112	0.04
2018	93,438	16,924	0.18
2019	82,303	17,285	0.21
2020	141,663	38,173	0.27
2021	93,822	9,444	0.10
2022	122,333	15,999	0.13
2023	132,029	3,061	0.02
2024	138,272	NA	
2025	143,969	NA	

Ecosystem Considerations

Pacific mackerel are part of the CPS assemblage of the northeastern Pacific Ocean, which represents an important forage base in the California Current Ecosystem (CCE). Pacific mackerel do not typically represent a dominant species of this assemblage in any given year, with abundances likely less than more productive CPS, such as northern anchovy and Pacific sardine. However, mackerel population biomass can increase to relatively high levels during periods of favorable oceanographic conditions, which likely occur less regularly than observed for anchovy and sardine stocks. Relatedly, periods of low recruitment success driven by prevailing oceanic phenomena can lead to low population abundance over extended periods of time. Readers should consult Field et al. (2001), PFMC (1998, 2021), and NMFS (2022) for comprehensive information regarding environmental processes generally hypothesized to influence small pelagic species that inhabit the CCE.

Harvest Control Rules

A federal fishery management plan (FMP) for CPS, including Pacific mackerel, was implemented by the PFMC in January 2000 (PFMC 1998). The FMP’s harvest policy for Pacific mackerel, originally implemented by the State of California, was based on simulation analysis

conducted during the mid-1980s (MacCall et al. 1985), with the addition of a proration to account nominally for the portion of the assessed stock assumed to inhabit U.S. waters (PFMC 1998). The following maximum sustainable yield (MSY) control rule for Pacific mackerel has been generally used for management from the early 2000s to the present:

$$Harvest = (Biomass - Cutoff)E_{MSY} * Distribution$$

where Harvest is the harvest guideline (HG), Cutoff (18,200 mt) is the lowest level of estimated biomass above which harvest is allowed, E_{MSY} (30%, also referred to as exploitation fraction in earlier PFMC documents) is the proportion of biomass above the Cutoff that can be harvested by fisheries, and Distribution (70%) is the average proportion of total Biomass (ages 1+) assumed to reside in U.S. waters. The HGs under the federal FMP are applied to a July to June fishing year. Detailed description of the current management actions applicable to Pacific mackerel, including quotas and related fishing quantities (e.g., allowable biological catch-ABC, annual catch limit-ACL, overfishing limit-OFL, etc.), can be found in the most recent CPS SAFE document (PFMC 2021). Also, see Harvest Control Rules for U.S. Management (2019-20 and 2020-21).

Total annual harvest of Pacific mackerel by the Mexico fishery is not regulated by quotas, but there has been minimum legal size limits (e.g., 25.5 cm) imposed in the past. International management agreements between the U.S. and Mexico regarding transboundary stocks, such as Pacific mackerel, have not been developed to date (see Research and data needs below).

Management Performance

From 1985 to 1991, the catch exceeded 136,000 mt and no state quota restrictions were in effect. State quotas for 1992-00 fishing years averaged roughly 24,000 mt. The HGs averaged roughly 15,000 mt from 2001-06. In 2007, the HG was increased substantially to 40,000 mt and remained at this level until 2009, when the calculated HG (55,408 mt) was reduced by management to 10,000 mt based on limited landings in recent years, with the quota applicable through the 2010-11 fishing year that included an additional 1,000 mt incidental landing allowance (11,000 mt). Following the full stock assessment conducted in 2011, a harvest guideline of roughly 31,000 mt was implemented for two consecutive fishing years. Catch-based projection assessments were used to set HGs for 2013-14 (~39,000 mt) and 2014-15 (~29,000 mt). HGs have remained at roughly 20,000-25,000 mt since 2015. Note that from a management context, the CPS fishery has not fully utilized HGs since the late 1990s, with total landings far below recommended catches (Table ES-5).

Table ES-5: Pacific mackerel US overfishing limits (OFL), allowable biological catches (ABC), annual catch limits (ACL), harvest guidelines (HG) since 2008. Total US landings (USA-TOT) and the percentage of ACL are also shown. Model year 2008, for example includes landings from July 1, 2008 to June 30, 2009 to align with fishery management timeframes.

Model Year	OFL	ABC	ACL	HG	USA-TOT	PercHG
2008	NA	NA	40,000	NA	4,543	11%
2009	NA	NA	10,000	NA	3,607	36%
2010	NA	NA	11,000	NA	2,313	21%
2011	44,336	42,375	40,514	30,386	2,074	7%
2012	44,336	42,375	40,514	30,386	5,608	18%
2013	57,316	52,358	52,358	39,268	11,909	30%
2014	32,992	30,138	29,170	24,170	5,593	23%
2015	25,291	23,104	21,469	20,469	4,686	23%
2016	24,983	22,822	21,161	20,161	2,757	14%
2017	30,115	27,510	26,293	25,293	1,808	7%
2018	27,662	25,269	23,840	22,840	2,501	11%
2019	14,931	13,169	11,109	10,109	3,916	39%
2020	11,772	10,289	7,950	6,950	691	10%
2021	12,145	9,446	8,323	7,323	1,007	14%
2022	9,644	7,501	5,822	4,822	990	21%

Research and Data Needs

Extending the AT survey into Mexican waters should continue to be a top priority. The data collected on these surveys are valuable for the stock assessment (see prior based on 2021 Q value) and will enable future research into the movement and distribution of Pacific mackerel (and other CPS like Pacific sardine).

Thanks to the full time staff at the SWFSC, the AT survey age data are no longer a major data need. Efforts to coordinate with state agencies and, perhaps in the future, Mexican agencies should continue as age-composition data are crucial for stock assessment.

The harvest control rule utilized in the Pacific mackerel federal CPS-FMP was developed in the mid-1980s based on estimated abundance and spawner-recruit data available at that time. Harvest strategies should be re-examined using updated data and simulation methods.

1 Introduction

1.1 Stock Structure and Management Units

The full range of Pacific mackerel (*Scomber japonicus*, also referred to as chub or blue mackerel) in the northeastern Pacific Ocean is from southeastern Alaska to Banderas Bay (Puerto Vallarta), Mexico, including the Gulf of California (Hart 1971). Although stock structure of this species off the Pacific coast of North America is not known definitively, it is generally hypothesized that three spawning aggregations exist currently: one in the Gulf of California; one in the vicinity of Cabo San Lucas (Baja California, Mexico); and one along the Pacific coast north of Punta Abreojos (Baja California) that extends north to areas off southern California, and even further during favorable oceanographic periods to waters off the U.S. Pacific Northwest. The latter sub-stock is harvested by fishermen in the U.S. and Mexico, and is the population considered in this assessment.

The Pacific Fishery Management Council (PFMC) manages the northeastern Pacific Ocean stock along the Pacific coast of North America as a single unit, with no area- or sector-specific allocations. However, the formal Fishery Management Plan (FMP) harvest control rule does include a stock distribution adjustment, based on a long-term assumption that on average, roughly 70% of this transboundary population resides in U.S. waters in any given year (PFMC 1998).

1.2 Distribution and Movement

Although the northeastern Pacific Ocean stock ranges from southeastern Alaska to southern Baja California, the species is more common from Monterey Bay, CA to Cabo San Lucas, Mexico (Figure 1). Over the last few decades, the stock has been observed to more fully occupy the northernmost portions of its range in response to warmer oceanographic conditions that have persisted in the northeastern Pacific Ocean, being found at times as far north as British Columbia, Canada (Ware and Hargreaves 1993, Hargreaves and Hungar 1995). To date, there exists only a general understanding of the seasonal movement patterns exhibited by this species along the coast of North America (Fry Jr and Roedel 1949, Roedel 1949, Parrish and MacCall 1978, Hill et al. 1999), with northward movement from waters off Baja and southern California beginning in the late spring/summer to feed in productive areas of upwelling off Oregon and Washington (potentially, more extensive geographical range during El Niño events, MBC (1987)); and southerly movement in the late fall/winter back to spawning grounds off southern and Baja California. Pacific mackerel sampled from Pacific Northwest incidental fisheries (e.g., Pacific hake and salmon spp.) during the mid-1990s indicated the fish were generally older and larger than those captured in the southern California fishery (Hill et al. 1999). In recent years, the stock has been observed to be relatively abundant in waters off the Pacific Northwest as documented in cruise reports for the acoustic-trawl (AT) survey, conducted annually since the mid-2000s by the Southwest Fisheries Science

Center (SWFSC), e.g., Stierhoff et al. (2019a) and Zwolinski et al. (2019). Thus, the stock is assumed to be most abundant in U.S. waters during the summer and fall months of each year; however, determination of the exact portion of the population that occupies U.S. waters each summer/fall is necessarily problematic and subject to some level of uncertainty.

It is further hypothesized that the stock exhibits east-west (inshore-offshore) movement along the U.S. Pacific coast, with increased inshore abundance during July to November and increased offshore abundance during March to May (Cannon 1967, Sciences 1987). Pacific mackerel usually occur within 30 km of shore, but have been captured as far as 400 km offshore (Fitch 1969, Frey 1971, Sciences 1987, Allen et al. 1990). Pacific mackerel adults are found in water ranging from 10 to 22.2°C (Sciences 1987) and larvae are found in water around 14°C (Allen et al. 1990). Adult fish are commonly found near shallow banks. Juveniles are found off sandy beaches, around kelp beds, and in open bays. Adults are found from the surface to 300 m depth (Allen et al. 1990). Pacific mackerel often school with other small pelagic species, particularly jack mackerel and Pacific sardine, likely based on size/age attributes as well (Parrish and MacCall 1978).

1.3 Life History

Pacific mackerel found off the Pacific coast of North America are the same species found elsewhere in the Pacific and Indian Oceans (Collette and Nauen 1983). Synopses regarding the biology of Pacific mackerel are presented in Kramer (1969) and Schaefer (1980). Spawning occurs from Point Conception, California to Cabo San Lucas from 3 to over 300 km offshore (Moser et al. 1993). Off California, spawning occurs from March to October (primarily, late April through August) at depths to 100 meters (Knaggs and Parrish 1973). Off central Baja California, spawning can occur year round at some level. Around Cabo San Lucas, spawning occurs primarily from late fall to early spring. Pacific mackerel are believed to seldomly spawn north of Point Conception (Fritzsche 1978, Sciences 1987).

As exhibited by similar CPS, Pacific mackerel have indeterminate fecundity and appear to spawn whenever sufficient food is available and favorable oceanographic conditions prevail. Individual fish may spawn eight times or more each year and can release batches of at least 68,000 eggs per spawning. Actively spawning fish appear capable of spawning daily or every other day (Dickerson et al. 1992). New research on Mediterranean Sea *S. japonicus* reproduction showed a wide range of relative fecundity by length and weight [420 to 2,553 eggs per cm for total length, and 76 to 379 eggs per gram for total weight; Farrag et al. (2022)]. The length at 50% maturity for this population was 19.7 cm for females and 19.5 cm for males. Farrag et al. (2022) also include a literature summary of length at first maturity for global *S. japonicus* populations, which ranges from about 18-30 cm. Research from the western North Pacific Ocean shows an effect of maternal age on egg and larval success (Yoneda et al. 2022). Yoneda et al. (2022) found significantly larger and more nutrient-rich eggs, higher starvation tolerance, larger body size, and faster growth rates of larvae from 3 year-old females compared to 1 year-old females. Currently, Stock Synthesis does not

provide an option for directly increasing egg or larval survival based on female age, therefore any model explorations would have to indirectly address this relationship by increasing the number of eggs by length or weight.

Pacific mackerel larvae eat copepods and other zooplankton, including fish larvae (Collette and Nauen 1983, Sciences 1987). Juvenile and adult mackerel feed on small fish (e.g., northern anchovy), fish larvae, squid, and pelagic crustaceans, such as euphausiids (Clemmens and Wilby 1961, Turner and Sexsmith 1967, Fitch 1969, Fitch and Lavenberg 1971, Frey 1971, Hart 1971, Collette and Nauen 1983). Pacific mackerel larvae are subject to predation from a number of invertebrate and vertebrate planktivores. Juveniles and adults are eaten by larger fishes, marine mammals, and seabirds. Principal predators include porpoises, California sea lions, pelicans, and large piscivorous fish, such as sharks and tunas. Pacific mackerel likely school as a defense against predation, often with other CPS, such as jack mackerel and Pacific sardine.

Population dynamics of the Pacific mackerel stock off U.S. Pacific coast, particularly California, have been extensively studied in the past and of particular importance was pioneering research conducted during the 1970s and 1980s, e.g., Parrish (1974), Parrish and MacCall (1978), Mallicoate and Parrish (1981), MacCall et al. (1985), and Prager and MacCall (1988). Since the mid-1990s, various age-structured population dynamics models have been used to assess the Pacific mackerel stock for providing management advice (e.g., Jacobson et al. 1994, Hill and Crone 2005, Crone et al. 2009, Crone and Hill 2015), see History of modeling approaches below.

Pacific mackerel experience cyclical periods of notable abundance, a phenomenon exhibited by CPS in general, which are characterized by relatively short life spans and highly variable productivity/abundance driven primarily by large-scale environmental factors (e.g., Pacific Decadal Oscillation (PDO), North Pacific Gyre Oscillation (NPGO), and related oceanographic drivers, such as sea-surface temperature, sea-surface height, upwelling, chlorophyll, etc.). Analysis of mackerel scale-deposition data (Soutar and Isaacs 1974) indicates that periods of high biomass, such as during the 1930s and 1980s, are relatively rare events that might be expected to occur, on average, about once every 60 years (MacCall et al. 1985). Results from the ongoing assessment of this stock generally support past research, with periods of high recruitment success observed no more frequently than every few decades. As presented above, recruitment is generally variable both spatially and temporally in the northeastern Pacific Ocean, and unlikely to be related strongly to spawning stock size (Parrish 1974, Parrish and MacCall 1978).

One of the largest recorded Pacific mackerel was 63.0 cm in length (FL) and weighed 2.9 kg (Roedel 1938, Hart 1971), but the largest Pacific mackerels taken by commercial fishing (CA) were a 47.8 cm FL fish and a 1.72 kg fish. The oldest recorded age for a Pacific mackerel was 14 years, but most commercially caught Pacific mackerel recorded by CDFW are less than 4 years old, with few living beyond age 8 and larger than 45 cm. Historical data of generally older and larger Pacific mackerel sampled from Pacific Northwest incidental fisheries in the 1990s exists, and have been reported on previously (Hill et al. 1999). The oldest Pacific

mackerel from Washington state was 16 years old and measured 69 cm.

As addressed in earlier assessments/reviews, size-at-age relationships by sex and sex ratio data indicated no notable sexual dimorphism in growth or mortality rate is exhibited by this species. Combined sex models have been used in all Pacific mackerel assessments used to advise management.

1.4 Fishery Descriptions

Pacific mackerel are currently harvested by three fisheries (Table 1 and Figure 2): the USA commercial fishery that primarily operates out of southern California, as well as Oregon and Washington; a sport fishery based largely in southern California; and the Mexico commercial fishery that is based in Ensenada, Baja California and Magdalena Bay, Baja California Sur. In the commercial fisheries, Pacific mackerel are landed by the same boats that catch Pacific sardine, northern anchovy, jack mackerel, and market squid (commonly referred to as the west coast ‘wetfish’ fleet). In recent years, Oregon and Washington have landed limited amounts of Pacific mackerel, with a combined annual average catch of roughly 500 mt over the last decade. Pacific mackerel are also (incidentally) harvested in small volumes by whiting trawlers and salmon trollers. Available information concerning bycatch and discard mortality of Pacific mackerel, as well as other members of the small pelagic fish assemblage of the California Current, is presented in PFMC (2021). Limited information from observer programs implemented in the past indicated little bycatch of other species and/or discard of Pacific mackerel in the commercial purse seine fishery off the U.S. Pacific coast.

The history of California’s Pacific mackerel fishery has been reviewed by Croker (1933), Croker (1938), Roedel (1952), and Klingbeil (1983). Historically, Pacific mackerel have been landed in moderate amounts, supporting a viable fishery off California during the 1930s and 1940s and more recently, during the 1980s and early 1990s. During the early years of the fishery, Pacific mackerel were taken by lampara and pole-and-line boats, which were replaced in the 1930s by the same purse seine fleet that fished for Pacific sardine. Before 1929, Pacific mackerel were taken incidentally, in relatively small volumes with sardine and sold as a fresh product (Frey 1971). Canning of Pacific mackerel began in the late 1920s and increased as greater processing capacities and more marketable ‘packs’ were developed. Landings decreased in the early 1930s due to the economic depression and subsequent decline in demand, but increased significantly by the mid-1930s (66,400 mt in 1935-36). During this period, Pacific mackerel were second only to Pacific sardine in total (annual) landings. Subsequently, harvests underwent a long-term decline and for many years, a continued demand for canned mackerel exceeded supply. Supply reached record low levels in the early 1970s, at which time the State of California implemented a ‘moratorium’ on the directed fishery, allowing only limited amounts of incidental landings.

The moratorium was lifted following a period of ‘recovery’ that spanned from the mid to late 1970s. During the 1980s through mid-1990s, catches of Pacific mackerel by California

fishermen supported an economically viable fishery. The market for canned mackerel during the 1980s through early 1990s fluctuated substantially due largely to economic factors. Domestic demand for canned Pacific mackerel eventually waned and the last mackerel cannery in California closed in 1992. Presently, the limited landings of Pacific mackerel caught by U.S. fishermen are used for human consumption (e.g., canned, frozen, fresh) or pet food.

Pacific mackerel are caught by recreational anglers in southern California using commercial passenger fishing vessels (CPFV), private boats, piers, beaches, etc., but are not typically considered a target species (Young 1969), with comparatively minimal catches to landings from commercial operations (Table 1). Pacific mackerel are also harvested in California's recreational fishery as bait for directed fishing on larger pelagic species, such as tunas, sharks, and billfishes. Additionally, Pacific mackerel are caught by anglers in central California, Oregon, and Washington, but typically, in only limited amounts. The sport harvest of Pacific mackerel in California comprises a very small fraction of the total landings of Pacific mackerel, e.g., over the last decade, recreational catch is less than 5% of the total weight landed (Table 1). Although some mackerel are likely discarded in some recreational fishing sectors for this non-targeted species, accurate determination of discard magnitude from available creel survey data is not straightforward, potentially subject to problematic sampling biases in the field.

In summary, Pacific mackerel landings in the U.S. have remained low over the last two decades, with total annual landings averaging roughly 7,000 mt since the late 1990s (Table 1). Relatedly, mackerel catches from fisheries have not realized allowable yields via stipulated harvest guidelines imposed since the late 1990s (see Table 2 and “Management performance” below).

The Mexico fishery for Pacific mackerel is primarily based in Ensenada and to a lesser extent, Magdalena Bay, Baja California Sur. The Mexico purse seine fleet has slightly larger vessels, but is similar to southern California's fleet with respect to gear (mesh size) and fishing practices. The fleet operates in the vicinity of the nearby ports and also targets other CPS. Demand for Pacific mackerel in Baja California increased after World War II. Mexico landings remained stable for several years, increased to over 10,000 mt in the mid-1950s, declined to under 500 mt during the mid-1970s, and remained relatively low through the late 1980s. Landings of Pacific mackerel in Ensenada peaked during the 1990s, but have remained relatively low over the last two decades. For the most part, the Ensenada fishery has been generally comparable in volume to the southern California fishery since 1990 (averaging about 10,000 mt/yr), with some differences for particular years (Table 1). In Mexico, harvested Pacific mackerel have been canned for human consumption or reduced to fish meal.

1.5 Ecosystem Considerations

Pacific mackerel are part of the CPS assemblage of the northeastern Pacific Ocean, which represents an important forage base in the California Current Ecosystem (CCE). Pacific mackerel do not typically represent a dominant species of this assemblage in any given year,

with abundances likely less than more productive CPS, such as northern anchovy and Pacific sardine. However, mackerel population biomass can increase to relatively high levels during periods of favorable oceanographic conditions, which likely occur less regularly than observed for anchovy and sardine stocks. Relatedly, periods of low recruitment success driven by prevailing oceanic phenomena can lead to low population abundance over extended periods of time. Readers should consult Field et al. (2001), PFMC (1998, 2021), and NMFS (2022) for comprehensive information regarding environmental processes generally hypothesized to influence small pelagic species that inhabit the CCE.

1.6 Management History

The state of California first implemented formal management associated with the Pacific mackerel stock in 1970, after the stock was thought to have declined substantially during the mid-1960s. A moratorium was placed on the fishery at this time, with a small allowance for incidental catch in mixed-fish landings. In 1972, legislation was enacted that imposed a quota based on the estimate of age-1+ biomass (>1-yr old fish) generated from formal stock assessments. Some very strong year classes in the late 1970s led to a brief period of moderately high stock abundance, which was followed by the fishery being reopened under a quota system in 1977. From 1977 to 1985, various adjustments were made to quotas for the directed harvest of Pacific mackerel and related incidental catch limits. It is important to note that even during the moratorium, substantial allowances were made for incidental catches associated with this species (Parrish and MacCall 1978).

State regulations enacted in 1985 imposed a moratorium on directed fishing when the total biomass was less than 18,200 mt, and limited incidental landings of Pacific mackerel to 18% (about 3,000 mt) during such periods. At this time, the ‘fishing year’ was set to extend from July 1st to June 30th of the following year. In summary, seasonal quotas, equal to 30% of the total biomass in excess of 18,200 mt, were allowed when the biomass was between 18,200 and 136,000 mt, with no quota limitations in effect when the total biomass was estimated to be 136,000 mt or higher.

A federal fishery management plan (FMP) for CPS, including Pacific mackerel, was implemented by the PFMC in January 2000 (PFMC 1998). The FMP’s harvest policy for Pacific mackerel, originally implemented by the State of California, was based on simulation analysis conducted during the mid-1980s (MacCall et al. 1985), with the addition of a proration to account nominally for the portion of the assessed stock assumed to inhabit U.S. waters (PFMC 1998). The following maximum sustainable yield (MSY) control rule for Pacific mackerel has been generally used for management from the early 2000s to the present:

$$Harvest = (Biomass - Cutoff)E_{MSY} * Distribution$$

where Harvest is the harvest guideline (HG), Cutoff (18,200 mt) is the lowest level of estimated biomass above which harvest is allowed, E_{MSY} (30%, also referred to as exploitation fraction

in earlier PFMC documents) is the proportion of biomass above the Cutoff that can be harvested by fisheries, and Distribution (70%) is the average proportion of total Biomass (ages 1+) assumed to reside in U.S. waters. The HGs under the federal FMP are applied to a July to June fishing year. Detailed description of the current management actions applicable to Pacific mackerel, including quotas and related fishing quantities (e.g., allowable biological catch-ABC, annual catch limit-ACL, overfishing limit-OFL, etc.), can be found in the most recent CPS SAFE document (PFMC 2021). Also, see Harvest Control Rules for U.S. Management (2019-20 and 2020-21).

Total annual harvest of Pacific mackerel by the Mexico fishery is not regulated by quotas, but there has been minimum legal size limits (e.g., 25.5 cm) imposed in the past. International management agreements between the U.S. and Mexico regarding transboundary stocks, such as Pacific mackerel, have not been developed to date (see Research and data needs below).

1.7 Management Performance

From 1985 to 1991, the catch biomass exceeded 136,000 mt and no state quota restrictions were in effect. State quotas for 1992-00 fishing years averaged roughly 24,000 mt. The HGs averaged roughly 15,000 mt from 2001-06. In 2007, the HG was increased substantially to 40,000 mt and remained at this level until 2009, when the calculated HG (55,408 mt) was reduced by management to 10,000 mt based on limited landings in recent years, with the quota applicable through the 2010-11 fishing year that included an additional 1,000 mt incidental landing allowance (11,000 mt). Following the full stock assessment conducted in 2011, a harvest guideline of roughly 31,000 mt was implemented for two consecutive fishing years. Catch-based projection assessments were used to set quotas for 2013-14 (~39,000 mt) and 2014-15 (~29,000 mt). Quotas have remained at roughly 20,000-25,000 mt since 2015. Note that from a management context, the CPS fishery has not fully utilized HGs since the late 1990s, with total landings far below recommended catches (see Table 2 for harvest regulations from 2008-18).

2 Data

The available data between 2008 and 2021 are shown in Figure 3. Data for model year 2022 were available but not finalized nor included in this base model.

2.1 Fishery-dependent data

Fishery data for assessing Pacific mackerel included landings from California, Oregon, and Washington commercial fisheries, the California recreational fishery, and the Mexico commer-

cial fishery from Ensenada, BC and Magdalena Bay, BCS. Additionally, port sample data (ages, lengths, and weights) from from California’s commercial fishery were included.

Since 1929, CDFW has collected biological data for Pacific mackerel landed in the southern California fishery (primarily, San Pedro). Limited samples have also been collected from the Monterey fishery when available. Sample data collected from 2008 through 2022 were incorporated in this assessment (Table 4). There was one fishery sample from San Pedro from August 2022 (model year 2022) which was not included. Biological samples from the commercial fishery generally include whole body weight, fork length, sex, maturity (visual), and otoliths for age determination. Currently, CDFW strives to collect 12 ‘random’ (port) samples per month (typically, 25 fish per sample) to determine length/age compositions, as well as catch-at-age, weight-at-age, etc. for the directed fishery.

Additionally, port sampling data for the commercial fishery in Mexico have been collected by the National Fisheries Institute (INAPESCA) since 1989; however, this information has not been made formally available to date and thus, commercial fishery data from the California purse seine fleet were assumed to be representative of the combined fisheries. Lack of data from the Pacific Northwest and Baja California may not be a serious problem for some years when catches were low. However, in some recent years, Baja California catches have equaled or exceed California catches by volume (Table 1), which necessarily increases the likelihood that potential biases associated with the omission of (and subsequent assumptions concerning) sample data from the Mexico and Pacific Northwest fisheries.

Pacific mackerel are aged by CDFW biologists based on identification of annuli in whole sagittae. Historically, a birth date of May 1st was used to assign year class (Fitch 1951). In 1976, ageing protocols changed to a July 1st birth date, which coincided with an increasing population, resumed fishery sampling, and a change in the management season from a May 1st opening to a July 1st start date. Fishery inputs for this assessment were compiled by ‘biological year,’ based on the birth dates used to assign age. The biological year used in this assessment is synonymous with the ‘fishing year’ defined previously, as well as with ‘fishing season’ as reported in the historical literature (from 1976 onwards). All landings and biological compositions included in this assessment were developed on a fishing year (July – June) basis. Sample sizes associated with biological data used in this assessment are presented in (Table 4).

2.1.1 Landings

The assessment includes commercial and recreational landings from calendar years 2008 to 2022. Catch estimates are based on model years and presented by region in Table 1 and Figure 2. Commercial catch statistics compiled in the CPS assessment data base are from the state fishery agencies CDFW (T. Nguyen, pers. comm.), Oregon Department of Fish and Wildlife (ODFW, C. Schmitt, pers. comm.), and Washington Department of Fish and Wildlife (WDFW, L. Wargo, pers. comm.). California recreational catch (mt) time series

from 2008 to the present are based on all sport fishery modes (man-made, beach/bank, party/charter, and private/rental) and obtained from CDFW (K. Lynn, pers. comm.).

As in the last assessment (Crone et al. 2019), commercial and recreational catch have been combined into one fishery, given similar selectivity properties between the two fisheries and the limited sport-related catches. To date, the sport fishery has contributed only limited catches to the overall landings of this species. Discards were assumed to be negligible, as in previous assessments, in both the commercial and recreational fisheries associated with this species. The total values summed across region are shown in Table 5 and Figure 4

Mexico landings reflect catches in Baja California from commercial purse seine fleets operating off Ensenada and in Magdalena Bay. Commercial landings from 2008 to 2022 were taken from the National Commission of Aquaculture and Fishing (CONAPESCA) website that archives Mexico’s fishery yearbook statistics e.g. CONAPESCA (2020).

Landings values were updated at the STAR panel to reflect removals from the Pacific whiting at-sea fishery for 2008-2022. Additionally, Washington state landings in 2012 were corrected to exclude 126 mt of ‘unspecified’ mackerel previously ascribed to Pacific mackerel in the PacFIN database. The differences between the catch time series are shown in Table 3, and the base model and associated sensitivities used this updated catch time series.

2.1.2 Age compositions

Presently, age data are only available from the California commercial fishery, which typically contributes the majority of fish landed at U.S. Pacific coast ports (Table 1). Biological sampling directed towards Pacific mackerel has recently begun in the states of Oregon and Washington, but only limited information is available at this time. Sample sizes (number of fishing trips) and number of measured individuals (specimens) associated with biological compositions included or considered in this assessment are presented in Table 4.

The nominal age compositions were weighted by the total monthly landings (L_m). Port samplers biologically sample 25 individual fish per landed haul. The following steps were used to develop the weighted age-composition time series (Figure 5):

- identified an ‘age-plus’ group (8+) for combining older fish into a single group and enumerate the number of individual fish (n) sampled in each month (m), age (a), and calendar year (y)

$$n_{m,a,y}$$

- Sum total biological sample weight (B) by m and y and calculate mean weight (w) of sampled fish by m, a, y :

$$B_{m,y}$$

$$\bar{w}_{m,a,y}$$

- Calculate proportions (A) in the biological samples by m, a, y

$$A_{m,a,y} = (\bar{w}_{m,a,y} * n_{m,a,y}) / B_{m,y}$$

- Calculate the total landings L by m, a, y

$$L_{m,a,y} = A_{m,a,y} * L_{m,y}$$

- The number of fish (F) in the catch were then calculated m, a, y

$$F_{m,a,y} = L_{m,a,y} / \bar{w}_{m,a,y}$$

and summed by a and model year (MY). Model years span July of year y to June of $y + 1$.

$$F_{a,MY} = \sum_{z=July,y}^{June,y+1} F_{a,z}$$

- The final proportion P at a and MY is

$$P_{a,MY} = F_{a,MY} / \sum_{a=0}^8 F_{MY}$$

Total numbers of ages measured were divided by 25, which is the typical number of fish collected per sampled fishing load. This calculation was used to set the sample sizes for age composition data included in the assessment model. Age compositions were input as proportions.

2.1.3 Ageing error

Pacific mackerel are routinely aged by fishery biologists at CDFW and the SWFSC based on the number of annuli, defined to be the interface between an inner translucent growth increment (Fitch 1951). Ageing error vectors were based on double-read methods and calculated based on the methodology described in Punt et al. (2008). The two ageing error vectors for calendar years 2015-2016 and 2017-2018 for the fishery-dependent data are shown in Table 6 and Figure 6. Additional details on CDFW ageing methodology can be found in Fitch (1951) and past stock assesment reports.

2.1.4 Empirical weight-at-age

A matrix of empirically derived weight-at-age (WAA) data were used in the model to convert estimated numbers-at-age in the model to biomass-at-age. Additionally, the WAA data were a substitute for directly estimating growth in the base model from available age and length composition data (Figure 7). WAA values for each age and model year were calculated with unweighted averages. A specific WAA value had to be calculated from a minimum of three measured fish. Within a cohort, ages without observations were linearly interpolated. A cohort without observations greater than a specific age were assumed to have constant weight-at-age values. For example, the 2013 cohort (Figure 7) did not have any age 6-8 fish measured, and the WAA value for age 5 was assumed to be applied to ages 6-8. The 2020 cohort did not have an age-0 WAA value, and this value was assumed to be the pooled age-0 WAA value across all cohorts.

2.2 Fishery-independent data: Acoustic-trawl survey

2.2.1 Overview

This assessment uses a single time series of biomass from the SWFSC's acoustic-trawl (AT) survey. Acoustic sampling of marine environments for determining abundance of fish populations is a standard practice worldwide that continues to receive more focused research in fisheries science, e.g., see Simmonds and MacLennan (2005) for general theory and application of fisheries acoustics, and ICES (2018) for an example of a long-term program for surveying trans-national, wide-ranging small pelagic fish communities. In February 2018, a second review was held for purposes of critically evaluating the AT survey methods in general, as well as determining the utility of these survey data for informing abundance of CPS in both ongoing and future assessments of the small pelagic fish assemblage of the California Current (PFMC 2018). The panel concluded that AT data represent the best scientific information available on an annual basis for assessing abundance of all members of the CPS assemblage (except Pacific herring), and approved the use of these data for directly (survey-based) or indirectly (model-based) assessing the status of the stock, depending on the species of interest (PFMC 2018).

2.2.2 Index of Abundance

Data from the summer SWFSC AT survey from 2008 and 2012-2021 were used in this assessment. There is no 2022 AT survey observation for Pacific mackerel. The time series used here is slightly different than that used in Crone et al. (2019). The previous time series borrowed a target strength (TS) value and length-weight relationship from South African Jack mackerel to translate abundance at length to biomass. The TS to length relationship

for Jack Mackerel (*Trachurus symmetricus*) was derived from echosounder measurements of backscattering from in situ Horse Mackerel (*Trachurus trachurus*) of South Africa (Barange et al. 1996). Because European Horse Mackerel (*Trachurus mediterraneus*) have similar TS to those of the Atlantic Mackerel [Palermino et al. (2021); *Scomber colias*, previously *Scomber japonicus*], the same TS to length relationship was used for Pacific Mackerel (*Scomber japonicus*) and Jack Mackerel.

The borrowed length-weight relationship resulted in AT survey empirical weight-at-age values that were lower than those from the fishery data. As a result, the STAT used a recently published Pacific mackerel length-weight relationship (Palance et al. 2019), which was calculated based on AT survey trawl samples. This Pacific mackerel length-weight relationship was used to convert abundance-at-length data to biomass, and the difference between the two biomass time series was about 9% on average. The one exception was the 2015 observation which had a previously published estimate of 7,146 mt but is now 1,353 mt with the updated length-weight relationship (Figure 8). The CVs associated with each estimate were assumed to be unchanged. The values of abundance by fork length and abundance by age are shown in Tables 7 and 8.

The summer 2008 survey biomass was estimated to be 58,511 mt with a CV of 0.38. The previous estimate was 55,000 mt (Demer et al. 2012).

The summer 2012 survey biomass estimate was 119,038 mt with a CV of 0.34. The summer 2013 estimate was 9,168 mt with a CV of 0.61. The previous estimates were 109,951 mt and 8,245 mt, respectively (Zwolinski et al. 2014).

The summer 2014 survey biomass was 9,159 with a CV of 0.56. The previous estimate was 10,423 mt. There is no report associated with this survey but the values were calculated with the same methods as other cruises (Zwolinski, personal communication). The values for this survey were calculated specifically for the 2019 benchmark (Crone et al. 2019).

The summer 2015 survey spanned roughly Haida Gwaii, British Columbia, Canada to San Diego, CA, USA with 79 east-west transects covering 3,150 nmi and 158 Nordic trawls (Stierhoff et al. 2018). The biomass estimate is 1,353 with a CV of 0.52. The previous published biomass estimate is 7,146 mt (Stierhoff et al. 2021). This difference is due to the reanalysis of the echograms and is not related to the update of the length-weight relationship.

The summer 2016 survey spanned roughly Cape Scott, British Columbia, Canada to San Diego, CA, USA with 103 east-west transects covering 4,627 nmi and 118 Nordic trawls (Stierhoff et al. 2018b). The biomass estimate was 35,401 with a CV of 0.52. The previous published biomass estimate was 32,782 mt (Stierhoff et al. 2021b).

The summer 2017 survey spanned roughly Cape Scott, British Columbia, Canada to Point Conception, CA, USA with 105 east-west transects covering 3,540 nmi and 83 Nordic trawls (Stierhoff et al. 2018c). The biomass estimate was 45,319 with a CV of 0.26. The previous published biomass estimate was 41,139 mt (Zwolinski et al. 2019).

The summer 2018 survey spanned Cape Scott, British Columbia, Canada to San Diego, CA with 127 east-west transects covering 6,104 nmi and 169 Nordic trawls (Stierhoff et al. 2019a). The biomass estimate was 31,739 mt with a CV of 0.22. The previous published biomass estimate was 33,351 mt (Stierhoff et al. 2019b).

The summer 2019 survey spanned Cape Scott, British Columbia, Canada to San Diego, CA with 140 east-west transects covering 6,691 nmi and 163 Nordic trawls (Stierhoff et al. 2020). The biomass estimate was 27,750 with a CV of 0.24. The previously published biomass estimate was 26,577, with 24,643 found in the core area and 1,934 nearshore (Stierhoff et al. 2020b).

The summer 2021 survey survey spanned Cape Flattery, WA to Punta Abreojos, Mexico with 141 east-west transects covering 6,749 nmi (Renfree et al. 2022). The biomass estimate was 23,830 with a CV of 0.24. The previously published biomass estimate was 21,998 mt (Stierhoff et al. 2023). There were an estimated 14,202 mt (65%) in Mexican and 7,796 mt (35%) in US waters [see Figure 9; Stierhoff et al. (2023)]

The full time series is shown in Figure 10.

2.2.3 Age compositions

Age composition data are shown in Figure 11. Estimates of abundance-at-length were converted to abundance-at-age using survey-specific age-length keys for the summer surveys (Figure 12). Age-length keys were constructed using ordinal generalized additive regression models from the R package *mgcv* (Wood 2017). A generalized additive model with an ordinal categorical distribution fits an ordered logistic regression model in which the linear predictor provides the expected value of a latent variable following sequentially ordered logistic distributions. Unlike previous iterations in which the conditional age-at-length was modeled as a multinomial response function ‘multinom’ from the R package ‘nnet’, and hence, disregarding the order of the age classes, the order logistical framework provides a more strict structure for the conditional age-at-length, which might, arguably, be beneficial with small sample sizes. The survey age compositions were weighted (i.e input sample sizes in Stock Synthesis) by the number of positive clusters in each cruise. This is in contrast to the calculation for the fishery age compositions, which considered a sample to be the number of total aged fish / 25.

2.2.4 Ageing error

Ageing error vectors were calculated based on the methodology described in Punt et al. (2008) and Thorson et al. (2012). The ageing error vectors are shown in Figure 6. There was one ageing error vector for the AT survey data from 2019-2021 (Table 6 and Figure 6), which included ages read by SWFSC staff. Ageing error vectors calculated from SWFSC

and CDFW staff had higher CVs, particularly ages 0-1. A model sensitivity that used this SWFSC/CDFW ageing error vector did not have an impact on age-1+ biomass. Further details on the ageing methodology are available in Appendix A.

2.2.5 Empirical weight-at-age

AT survey weight-at-age time series (Figure 7) were calculated for every survey using the following process: 1) the AT-derived abundance-at-length was converted to biomass-at-length using a time-invariant length-to-weight relationship (Palance et al. 2019); 2) the biomass- and numbers-at-length were converted to biomass-at-age and numbers-at-age, respectively, using the above-mentioned age-length keys; and 3) mean weights-at-age were calculated by dividing biomass-at-age by the respective numbers-at-age. The protocols for filling and interpolating missing values were the same as those described in the empirical weight-at-age section for the fishery data.

In the previous assessment, the AT survey and fishery weight-at-age values were assumed to be the same. This assessment utilizes updated age compositions, produced by the Life History Group at the SWFSC.

2.3 Nearshore sampling

The acoustic-trawl survey has had three methods for extrapolating or observing nearshore biomass: model extrapolation, unmanned surface vehicles, and fishing vessel acoustic-trawl methods (Stierhoff et al. 2020b).

With model extrapolation, the easternmost portions of transects are extrapolated to the 5-m isobath in the unsampled nearshore areas. Thus, the length and species compositions associated with the end of the transects are extrapolated to the 5-m isobath.

Unmanned surface vehicles (USVs) generally cover portions of the coast rather than the entire coast. The ability to collect USV observations has depended on the number of USVs available for use and on local wind conditions. The USVs collect acoustic data but do not collect associated biological samples. As a result, the nearest trawl compositions are assumed to be representative of the nearshore acoustic observations when calculating species-specific biomass values.

Fishing vessel acoustic-trawl methods involve equipping vessels with acoustic echosounders and conducting a maximum of one purse seine set during daylight hours. In the case of abundant coastal pelagic species or an unsuccessful daytime set, a set is conducted at night.

Nearshore biomass estimates for Pacific mackerel are: 5.97 mt in 2015 from model extrapolation (Stierhoff et al. 2021), 3,102 mt in 2016 from model extrapolation (Stierhoff et al.

2021b), 1,105 mt in 2017 from model extrapolation (Zwolinski et al. 2019), 1,320 mt in 2018 from model extrapolation (Stierhoff et al. 2019b), 1,934 mt in 2019 from acoustic-trawl fishing vessels (Stierhoff et al. 2020b), and 1,507 mt in 2021 from acoustic-trawl fishing vessels (Stierhoff et al. 2023).

2.4 Biological Parameters

2.4.1 Stock Structure

Fishery and survey observations from the west coast of the US (California, Oregon, and Washington) and catch values from Mexico (Baja California and Baja California Sur) were assumed to be part of the same stock. Pacific mackerel are found throughout the Northeast Pacific Ocean as described in the introduction.

2.4.2 Growth

Growth was assumed to not be sexually dimorphic, consistent with the assumptions in previous stock assessments (e.g. Crone et al. 2019). The assessment model used empirical weight-at-age values to account for Pacific mackerel growth. This approach is also consistent with the assessments of other US coastal pelagic species. Estimating growth internally in the stock assessment may be difficult due to variation in time and space and potential confounding between length-based selectivity, age-based availability to fishing/survey gear, and variable growth parameters.

2.4.3 Maturity

Maturity was modeled with a fixed vector of fecundity multiplied by maturity at age. The equation: $Maturity = \frac{1}{1 + \exp(\text{slope} * \text{age} - \text{age}_{inflection})}$ was used to estimate maturity at age from 494 female mackerel collected during spring and summer AT surveys from 2010-2021. The fixed maturity-at-age vector used as input for the population is shown in Table 9 and Figure 13

2.4.4 Natural mortality

In past assessments, natural mortality rate (M) was assumed to be $0.5yr^{-1}$ and constant over time for all ages. Parrish and MacCall (1978) estimated natural mortality for Pacific mackerel using early catch curves ($M = 0.3-0.5yr^{-1}$), regression of Z on f ($M = 0.5yr^{-1}$), and

comparative studies of maximum age [$M = 0.3-0.7yr^{-1}$; Beverton (1963)] and growth rate [$M = 0.4-0.6yr^{-1}$; Beverton and Holt (1959)]. The above research and overall conclusions considered the regression of Z on f to be the most reliable method, with the estimate $M = 0.5yr^{-1}$ falling within the range of the plausible estimates.

Given past uncertainty associated with assumed rates of M to consider for Pacific mackerel, as well as other members of the small pelagic species assemblage of the CCE, M was estimated in this assessment with a longevity-based prior described in Hamel and Cope (2022). The maximum age assumed for the prior was age-8, which is also the beginning of the plus group assumed in this assessment. The prior on M was lognormal with a mean of -0.393 (0.675 in linear space; $5.40 / 8$ the assumed age max) and SD of 0.31 (Hamel and Cope 2022).

2.5 Available data sets not used in assessment

The STAT investigated three fishery-independent data sets, that were ultimately not incorporated to this assessment: Investigaciones Mexicanas de la Corriente de California (IMECOCAL), California Cooperative Oceanic Fisheries Investigations (CalCOFI), and the Rockfish Recruitment and Ecosystem Assessment Survey (RREAS). IMECOCAL and CalCOFI seasonally sample eggs and ichthyoplankton in fixed grids in Mexican and US waters, respectively. The challenge with these data sets is that there is not a straightforward method of directly incorporating data from these early life stages, directly into the assessment framework. The RREAS data set has sparse observations for Pacific mackerel (134 individuals observed from 1990-2018).

Previous assessments have used a number of alternative indices of abundance (e.g. Hill et al. 1999). Aerial spotter data have been standardized with delta log-normal models to generate an index from 1963 to 1997. Commercial Passenger Fishing Vessel skippers provided catch and effort data to CDFW beginning in 1936. This index has calculated an index by calculating the number of mackerel caught per 1000 angler-hours for 10nmi blocks in California. Power Plant Impingement values have been provided from Southern California Edison. Samples of fish become entrained in cooling water at ten power plants along southern California coast. The Alaska Fisheries Science Center has a triennial bottom trawl survey that targets groundfish and has had bycatch of Pacific mackerel.

The 2022 AT survey biomass estimate is available for Pacific mackerel. However, due to logistical constraints, the survey area off Washington, Oregon, and part of northern California had to be conducted with fishing vessel acoustic-purse seine sampling. Acoustic-trawl sampling aboard the SWFSC's R/V *Reuben Lasker* began off northern California and proceeded south. Due to these differences with the preceding AT survey protocols, the 2022 biomass estimate and associated age compositions were not included in this benchmark.

Catch data and biological compositions are available prior to 2008, but the potential benefits of extending the modeling timeframe were not clear given the relatively short lifespan of

Pacific mackerel (and similar CPS). The model begins in 2008 to align with the beginning of estimates of Pacific mackerel biomass from the AT survey.

3 Stock Assessment Model

3.1 History of modeling approaches

Parrish and MacCall (Parrish and MacCall 1978) were the first to provide stock status determinations for Pacific mackerel using an age-structured population model (virtual population analysis, VPA). Beginning in the mid-1990s, the ADEPT model, which was based on the ADAPT VPA and modified for Pacific mackerel (Jacobson 1993, Jacobson et al. 1994), was used to evaluate stock status and establish management quotas for approximately 10 years. The assessment conducted in 2004 (for 2004-05 management) represented the final ADEPT-based analysis for this stock (see Hill and Crone 2004). The forward-simulation model ASAP (Legault and Restrepo 1998) was reviewed and adopted for Pacific mackerel at the STAR Panel conducted in 2004 (Hill and Crone 2004). The ASAP model was used for assessments and management advice from 2005 through 2008. The STAR Panel conducted in 2009 supported decisions to begin using the Stock Synthesis (SS) model for conducting formal stock assessments of Pacific mackerel (Crone et al. 2009, PFMC 2009); the SS model has been used for all assessments since 2009. A full (benchmark) stock assessment and review for this species were conducted in 2011 (Crone et al. 2011), with a harvest guideline (HG) serving for two fishing years. In 2013 and 2014, catch-based projections were conducted and used to set the HGs (Crone 2013, Crone and Hill 2014). In 2015, a benchmark assessment was conducted for purposes of providing management advice that served for two (fishing) years, 2015-16 and 2016-17 (Crone and Hill 2015). A catch-only projection was conducted in May 2017 that provided HGs for managing the Pacific mackerel resource for fishing years 2017-18 and 2018-19 (Crone and Hill 2017). The most recent benchmark assessment was conducted in 2019 (Crone et al. 2019).

3.2 2019 STAR Panel Recommendations

High priority

1. Improve collaboration with fishery researchers from Mexico. As noted in previous assessment reviews, a large fraction of the catch is taken off Mexico, and efforts should be made to obtain length, age and related biological data from the Mexican fisheries. Inclusion of the AT surveys in the assessment has increased the need for comparable surveys within Mexican waters because such information could be used to develop a nearly comprehensive index of the abundance of the transboundary stock of Pacific

mackerel. Alternatively, collaborative research extending the AT survey into Mexican waters would also achieve the goal of encompassing the full range of Pacific Mackerel.

- The AT survey began surveying Mexican waters in 2021. This was the result of extensive work by members of the Advanced Survey Technologies and Life History Group at the SWFSC.
2. Continue to refine the indices of abundance. The Panel considers an AT survey to be an appropriate way to index the abundance of CPS such as Pacific mackerel. The PFMC conducted reviews of the AT survey in 2011 (PFMC 2011) and in 2018 (PFMC 2018). Some of the recommendations from those reviews have been implemented (e.g. Zwolinski and Demer, 2014). However, most of the recommendations, even those from the 2011 review, have yet to be addressed. The following are a subset of tasks to better realize the potential of the AT survey for Pacific mackerel:
 - a. Trawl sampling during the day to address the potential for differences in fish represented by the signal from the acoustic sampling during the day versus trawl sampling at night to capture the species, length and age composition of the sampled fish.
 - This will be one component of experimental trawling scheduled for summer 2023.
 - b. Refine the target strength estimates for Pacific mackerel.
 - This may be evaluated in the future.
 - c. Provide separate estimates of age-0 and age-1+ Pacific mackerel biomass from the AT survey. There appears to be more uncertainty in the enumeration of age-0 mackerel than of other age classes due to the spatial distribution and age-specific selectivity patterns.
 - This calculation is possible but has not been provided.
 3. Standard data processing procedures should be developed for CPS, similar to those developed for groundfish species, and a ‘data document’ developed that provides, in considerable detail, how the basic data sources (e.g., catches, CPFV indices, etc.) are constructed. Much of this information has been published in the past, but a single (and ‘living’) document describing the basic data will assist assessment authors and future review panels.

- See this document and Appendix A for documentation
4. Investigate the spatial distribution, especially the range, of the Pacific mackerel population over time and whether this changes with population size and/or environmental conditions. In particular, an environmentally based index of spatial distribution might prove useful for developing priors for AT survey catchability for use in future assessments.
 - See response to recommendation number 1
 5. Improve collection of age data, coordination of ageing laboratories and cross validation efforts to standardize reads between laboratories and develop bias adjustments.
 - a. Increase support for current port sampling and laboratory analysis programs for CPS, particularly in the Pacific Northwest. Biological (e.g. length, age, sex) data on mackerel caught in the Pacific Northwest should be collected. These data could further assist in understanding whether and to what extent selectivity for the commercial fishery is domeshaped. The aging of Pacific sardine in the Pacific Northwest should be coordinated with laboratories conducting ageing in California.
 - b. Analysis of data from the multistage approach to age/length composition sampling has indicated that most of the variability occurs between commercial trips as opposed to replicate sampling of a landing within a landing. The number of trips sampled is relatively low due to the infrequent fishing and need to coordinate sampling with industry to increase the effective sample size. Many samples from the Pacific Northwest have not been processed and should be aged with methods consistent with those currently employed by the CDFW from the commercial fishery.
 - c. Ageing of survey collections for the survey age production laboratory at SWFSC needs increased collaboration to increase precision in reads. Reading of otoliths from the AT survey should be prioritized to alleviate the need for using age length keys to convert lengths to ages with greater potential for bias and imprecision. Production ageing of otoliths from the AT survey needs validation and verification of age reads between observers or laboratories should be conducted to provide reads consistent with those currently provided by CDFW for commercial landings, relying on experienced age readers as the basis for comparison between laboratories.
 - d. Cross reads should be conducted between laboratories or, preferably, reads simply done by CDFW staff to provide greater consistency and precision. Ageing bias can be identified using cross-reads of the same otoliths among laboratories.
 - The SWFSC hired full time staff in the Life History Program to improve the collection and processing of age data, standardize ageing protocols, cross-validate reads, improve ageing precision, and develop bias adjustments. Three SWFSC readers aged 1,762 Pacific mackerel collected from the 2012-2022 AT surveys for this assessment, including

samples collected from the Pacific Northwest. The SWFSC readers trained with the best CDFW reader and generated a standardized protocol, and 317 Pacific mackerel were cross-read by all four readers. This collaborative effort significantly improved the quality of age data, as bias among readers was low and precision was high (See Appendix A). A forthcoming Tech Memo will summarize ageing efforts by the SWFSC Life History Program for Pacific mackerel in greater detail. Additionally, there are plans to reach out to Pacific mackerel age reading labs in the Pacific Northwest to examine interagency comparisons.

6. Revisit the harvest control rules and reference points for Pacific mackerel. The basis for the current harvest cutoff are derived from analyses performed by MacCall et al. (1985) over 30 years ago using data, biological assumptions (e.g. about selectivity and natural mortality), and methods (virtual population analysis) that are not reflected in the current stock assessment. If the underlying data and assumptions used by MacCall et al. (1985) are no longer considered relevant to the current population as reflected in the ALT_19 assessment model, it is likely time to revise the scientific basis for these reference points.
 - The harvest control rules have not been revisited. Catches have been below harvest guidelines in the time frame of this model (2008-2021).

Medium priority

1. Examine whether parameters such as growth rate and asymptotic size have changed over time.
 - Growth was not modeled internal to the assessment.
2. Conduct a study to update the information used to determine maturity-at-length (and maturity-at-age).
 - See Appendix A

Low Priority

1. Explore the feasibility of modeling non-landed mortalities of sublegal-sized fish in the Mexican fishery
 - This has not been explored yet.

3.3 Base model description

A number of features have been modified for the 2023 benchmark assessment (Figure 14):

- Use of SS3.30.20, which was the most recent version (v3.30.21 has since been released)
- Extension of main recruitment deviation period to 2021
- Equal weight ($\lambda=1$) for fishery and AT survey age compositions. The previous model downweighted ($\lambda=0.5$) the AT survey age compositions, which were derived from an ALK developed from fishery-dependent data.
- Addition of SR_regime block parameter. Previously the model, which begins in 2008, was assumed to be starting from equilibrium conditions. Estimation of this additional parameter accounts for the model period beginning in a fished state which more closely matches the reality of the stock's history.
- The 2021 AT survey had observations from both US and Mexican waters. These observations informed the prior for values of Q
- Time-varying Q estimated with deviations for 2008-2015 and one block for 2016-2021. The prior was centered at 0.308 with SD of 0.28. The rationale for this decision is outlined in Appendix B.
- Time-varying fishery selectivity, modeled to have the random-walk (one selectivity parameter per age; option 17 in SS3) with parameter deviations estimated with the two-dimensional auto-regressive smoother. This treatment was also used in the 2021 anchovy benchmark assessment (Kuriyama et al. 2020).
- Age-specific, time-invariant natural mortality across ages 0-8. An average value of M is estimated in SS3, with a longevity-based prior assuming a maximum age of 8 per Hamel and Cope (2022).

3.3.1 Time period and time step

The modeled timeframe begins in 2008 and extends through 2021, to match the availability of the AT survey data (Figure 3). Annual timesteps are used in this assessment and the model year is aligned with the fishing year which spans July of one calendar year to June of the following calendar year. For example model year 2021 represents July, 2021 to June, 2022.

The goal of this assessment is to estimate terminal year stock biomass and forecast biomass levels for the following two fishing years. Extension of this model prior to 2008 may result in different estimates of scaling parameters but may not result in significantly different biomass estimates for recent years.

3.3.2 Forecast

Stock biomass was forecasted for model years 2022, 2023, and 2024. There are 2022 AT survey biomass data available but these were not included due to logistical challenges that

limited the survey protocols. The catch values used in the forecast file were data for 2022 and catch values averaged from 2019-2022 for the years 2023 and 2024. The fishery selectivity pattern in the forecast file was assumed to be the selectivity curve estimated for 2021.

3.3.3 Stock-recruit relationship

Equilibrium recruitment (R_0) and initial recruitment equilibrium offset (SR_{regime}) were estimated in the base model. Steepness (h) and average recruitment variability (σ_R) were fixed at 0.75 and 0.75, respectively. These were the values used in the previous stock assessment (Crone et al. 2019). Recruitment deviations were estimated as separate vectors for the early and main data periods in the model. A recruitment bias adjustment ramp (Methot and Taylor 2011) was applied to the early period and adjusted recruitment in the main period of the model.

3.3.4 Catchability

There is a high degree of variability in the index of abundance that is unlikely to be due to recruitment and natural mortality. For example, in 2012 the AT survey estimate was about 120,000 mt and the biomass estimates from 2013-2015 ranged from 1,353 to 9,168 mt. The STAT assumed that this decrease in biomass was due to a change in catchability (Q) rather than a large mortality event coupled with low recruitment. Pacific mackerel catchability could vary through time due to time-varying availability (i.e. migrations and movement) or due to gear avoidance.

The STAT modeled Q to be time-varying with annual deviations for 2008-2015 and one block for 2016-2021. The prior was centered at 0.308 with a SD of 0.28. The prior on Q was calculated based on the data from the 2021 AT survey which included observations from the US core survey grid (6,840mt; CV=0.279), US nearshore (1,680mt; CV=0.390), and Mexican core survey grid (15,310mt; CV=0.275). Values for Mexico nearshore biomass were calculated by sampling 100,000 values from the 2021 biomass mean and CV values (US core, US nearshore, and Mexico core) assuming normal distributions. The Mexico nearshore values were calculated assuming the US core to nearshore ratios were the same in Mexico. The proportion of biomass in the US was calculated as: $(US_{core} + US_{nearshore}) / (US_{core} + US_{nearshore} + Mex_{core} + Mex_{nearshore})$.

The mean proportion was 0.308 and the SD of the sampled values was 0.28.

The spatial observations for the AT survey vary through time, and the subsequent variability in the index of abundance was modeled with time-varying Q rather than a time-varying M for example. More details regarding sensitivity analyses run at the STAR panel are outlined in Appendix B.

3.3.5 Selectivity

Fishery selectivity was estimated to be time-varying with the 2dAR feature in SS3 (Xu et al. 2019). The base selectivity form was estimated as a “random walk” using SS3 terminology. In practice, the “random walk” form estimates a selectivity parameter for each age, and deviations around this base curve are estimated to be temporally independent. Parameters for ages 0-3 were time-varying where ages 4-8+ were time-invariant. The SE value for the deviations was 1.0, and values of 0.2, 0.5, and 0.7 were explored at the STAR panel. Decreasing the SE values resulted in smoother curves but poorer fits to the age composition data. The goal of this configuration was to capture the year-to-year variability in the fishery age composition data.

AT survey age-0 selectivity was estimated to be time-invariant. Other CPS assessments (e.g. Kuriyama et al. 2020) estimated age-0 selectivity to be time-varying. However, estimating time-varying selectivity for the AT survey resulted in a high estimate of M (roughly 1 for the average value across all ages). Biologically it does not seem possible that M for Pacific mackerel is greater than that for Pacific sardine and northern anchovy, and the STAT decided to estimate age-0 selectivity to be time-invariant.

3.3.6 Likelihoods components and model parameters

A complete list of model parameters estimated in the base model is shown in Table 10. The total objective function was based on the likelihood components from fits to the AT survey abundance index and fishery and AT survey age compositions (Table 11).

3.3.7 Bridging analysis

Figure 14 shows the addition of each major feature to the 2019 benchmark model. The additions of the Q blocks and time-varying fishery selectivity resulted in the largest changes in summary biomass estimates.

3.4 Base model results

3.4.1 Likelihoods and quantities of interest

The total likelihood value was 115.003 and the gradient was 4.580e-05. Likelihood values from the age-compositions and parameter deviations constituted a majority of the total likelihood. The forecast summary biomass values for model years 2022, 2023, and 2024 are 46,682, 55,680, and 60,785 mt, respectively.

3.4.2 Selectivity estimates and fits to fishery and survey age-compositions

Time-varying age-based selectivities were estimated for the fishery (Figure 15). Fits to the fishery age-composition data were relatively good, as the flexible 2dAR selectivity captured year-to-year variability (Figures 16 and 17). The fits to the survey age compositions are shown in Figure 18 and 19.

3.4.3 Fit to survey index of abundance

The base model, with time-varying Q values, fit all the AT survey indices of abundance (Figures 20 and 21). The values of Q are shown in Figure 22 and the values of age-specific M in Figure 23.

3.4.4 Stock-recruitment relationship

Recruitment was modeled using a Beverton-Holt stock-recruitment relationship (Figure 24). The recruitment deviations are presented in Figure 25. Asymptotic standard errors for recruitment deviations are shown in Figure 26 and the recruitment bias adjustment plot is shown in Figure 27. Note steepness and σ_R were both fixed at 0.75.

3.4.5 Population numbers- and biomass-at-age estimates

The population age distributions (by numbers of fish) are shown in Figure 28 and Table 12. Corresponding estimates of population biomass-at-age, total biomass (age-0+, mt) and summary biomass (age-1+, mt) are shown in Table 13.

3.4.6 Biomass and recruitment

Time series of estimated spawning stock biomass (SSB; mt) and associated 95% confidence intervals are presented in Table 14 and Figure 29. The estimated recruitment time series is shown in Table 14 and Figure 30.

Total and summary biomass values are shown in Table 15 and Figure 31. Summary biomass values are 49,977 mt in 2021, 46,682 mt in 2022, 55,680 mt in 2023 and 60,785 mt in 2024.

3.4.7 Fishing mortality

Estimated fishing mortality (apical F) time series are presented in Figure 32. Exploitation rates are shown in Table 16 and Figure 33.

3.5 Modeling Diagnostics

3.5.1 Convergence

Convergence was evaluated by starting model parameters from values jittered from the maximum likelihood estimates. Starting parameters were jittered by 5% for 50 replicates and 10% for 20 replicates. A lower likelihood was not found, and nearly all the replicates for both scenarios converged to the maximum likelihood value from the base model. The hessian was invertible in the base model.

3.5.2 Historical analysis

The historical analysis for summary biomass is shown in Figure 34. The assessments shown are from 2005, 2011, 2015, and 2019.

3.5.3 Likelihood profiles

There was not much information in the age compositions nor the AT index of abundance to estimate steepness (Table 17 and Figure 35). Steepness was fixed at 0.75 in the base model. There is a relatively weak data conflict between the survey and age compositions as steepness decreases below 0.75.

For the profile on Q , the value was fixed for the 2016-2021 block with time-varying estimates for the deviations. Neither the age compositions nor survey data seemed to have any information on catchability (Table 18 and Figure 36). Specifically the survey data contained little information to estimate catchability (Figure 36).

The AT survey age compositions seemed to contain the most information to estimate M and all the data sets were in relative agreement (Table 19 and Figure 37).

3.5.4 Sensitivity to alternative data weighting

The base model was run with age compositions reweighted according to the Francis method (Francis 2011) to evaluate model sensitivity to data weighting. The variance adjustment values were 4.161 for the fishery age comps and 0.508 for the AT survey age comps (Table 20). Parameter estimates, biomass estimates, and likelihood values are shown in Table 20 and Figure 38. With Francis reweighting, the 2021 summary biomass value increase from 40,024 in the base model to 43,962.

The base model was also run with downweighted age compositions ($\lambda = 0.5$ rather than 1 in the base model) to evaluate model sensitivity to data weighting. Parameter estimates, biomass estimates, and likelihood values are shown in Table 21 and Figure 39.

3.5.5 Sensitivity to alternative catch values

The base model was run with with two alternative scenarios based on the 2008-2021 catch values. One assumed forecast catch values were 5,699mt (the 25th percentile of 2008-2021 catch values) and the other assumed 20,366mt (the 75th percentile). The difference in values for 2024 and 2025 were 69,010 and 79,866 mt for the 25th percentile scenario and 62,523 and 68,524mt for the 75th percentile scenario (Table 22 and Figure 40).

3.5.6 Sensitivity to 2022 and 2023 AT survey biomass values

Models were run with values of 2022 and 2023 AT survey biomass ranging from 10,000 to 100,000 mt in increments of 10,000. The block on Q was extended to cover 2016-2022 or 2016-2023 in these model sensitivity runs. Parameter estimates, biomass estimates and likelihood values are shown for 2022 in Table 23 and 2023 in Table 24. The likelihood profiles show that a wide range of survey biomass values are plausible in both 2022 (Figure 41) and 2023 (Figure 42). Estimates of the summary biomass are shown in Figures ?? and ??.

3.5.7 Retrospective analysis

There was a retrospective pattern when re-running the model with one year of data dropped at a time (Figure 45). Pacific mackerel and CPS more generally have recruitment variability which partly explains the retrospective pattern. The base model has a time-varying Q value which is likely another source of the retrospective pattern.

4 Harvest Control Rules

Since 2000, the Pacific mackerel stock has been managed under a Federal Management Plan (FMP) harvest policy, stipulating that an optimum yield for this species be set according to the following harvest control rule [(HCR; PFMC (1998))]:

$$Harvest = (Biomass - Cutoff) * E_{MSY} * Distribution$$

where Harvest is the harvest guideline (HG), Biomass is age 1+ stock biomass (mt) in the respective fishing year (under the assumption that 55,681 mt in July 2023 and 60,785 mt in July 2024), Cutoff (18,200 mt) is the lowest level of estimated biomass above which harvest is allowed, E_{MSY} (30%, also referred to as Fraction) is the proportion of biomass above the Cutoff that can be harvested by fisheries, and Distribution (70%) is the average proportion of stock biomass (ages 1+) assumed in U.S. waters (PFMC 1998). Harvest stipulations under the federal FMP are applied to a July-June fishing year. The base model HG estimate for July 2023 was 7,871 mt (Figure 46) and 8,943 for July 2024 (Figure 47).

Overfishing limits (OFL) and Acceptable Biological Catch (ABC) statistics have been included in the management process since the adoption of Amendment 13 (PFMC 2011). OFL and ABC are defined as:

$$OFL = Biomass * E_{MSY} * Distribution$$

and

$$ABC_{Pstar} = Biomass * Buffer_{Pstar} * E_{MSY} * Distribution$$

where $Buffer_{Pstar}$ is calculated based on technical guidance from the SSC as documented in Wetzel and Hamel (2023). ABC buffers are based on assumed level of assessment uncertainty (σ) in the terminal model year, combined with additional uncertainty to account for natural mortality rate and time elapsed since the assessment. Stock biomass in the terminal model year (2022) had a CV=0.389, so the assessment σ (0.376) is lower than the default values for Category 1 ($\sigma = 0.5$) and Category 2 ($\sigma = 1.0$) assessments. The annual linear increase (r) in σ for the 2023 and 2024 projection years was calculated following Wetzel and Hamel (2023), where natural mortality rate ($M=0.851$) and time elapsed are considered. The annual linear increase was calculated to be:

$$r = 0.52 * M * \sigma_{baseline} = 0.4426 * \sigma_{baseline}$$

such that:

$$\sigma_y = \sigma_{baseline} * (1 + 0.4426(y_{management} - y_{assessment}))$$

Calculated values for a Category 1 assessment were $\sigma = 0.6106$ and $\sigma = 0.7213$ for 2023 and 2024, respectively. Calculated values for a Category 2 assessment were $\sigma = 0.7213$ and $\sigma = 0.9426$ for 2023 and 2024, respectively. Based on these methods, a range of Pstar buffers, along with corresponding ABCs for the 2023 and 2024 management years are presented in Figures 46 and 47.

5 Research and Data Needs

Extending the AT survey into Mexican waters should continue to be a top priority. The data collected on these surveys are valuable for the stock assessment (see fixed 2021 Q value) and will enable future research into the movement and distribution of Pacific mackerel (and other CPS like Pacific sardine).

Thanks to the full time staff at the SWFSC, the AT survey age data are no longer a major data need. Efforts to coordinate with state agencies and, perhaps in the future, Mexican agencies should continue as age-composition data are crucial for stock assessment.

The harvest control rule utilized in the Pacific mackerel federal CPS-FMP was developed in the mid-1980s based on estimated abundance and spawner-recruit data available at that time. Harvest strategies should be re-examined using updated data and simulation methods.

6 Acknowledgements

This assessment is possible with a number of contributions and analysis from multiple groups at the SWFSC and state agencies. Thanks to Emmanis Dorval, Brad Erisman, Owyn Snodgrass, Kelsey James, Brittany Schwartzkopf, Jon Walker, and Bev Macewicz of the Life History Group at the SWFSC; Kevin Stierhoff, Dave Demer and others from the Advanced Survey Technologies Group; Andrew Thompson from the CalCOFI Group; Steve Teo and Huihua Lee of the Population Dynamics Group; and Yuhong Gu of the Database Management Group. Thanks to Dianna Porzio, Kirk Lynn, and Trung Nguyen of CDFW and staff at WDFW and ODFW. Thanks as well to the STAR panel members: Andre Punt, Theresa Tsou, Chris Free and the CIE reviewer Joe Powers.

7 Tables

Table 1: Landings (mt) of Pacific mackerel by region and fishing year (1999-2022). Landings values from 2008-2022 were included in the assessment (see horizontal line). Mexican landings were from Magdalena Bay, BCS (MAG) and Ensenada, BC (ENS). US landings are from California (CA), Oregon (OR), Washington (WA). Additionally, California recreational landings are included (CA-REC). The total (TOT) landings are summed across all regions and used as input to the stock assessment. Note that model years include data from two calendar years. For example, model year 1999 includes landings from July 1, 1999 to June 30, 2000 to align with the fishery management timeframes.

Model Year	MAG	ENS	MEX-TOT	CA	OR	WA	CA-REC	USA-TOT	TOT
1999	97	2,524	2,621	3,634	0	0	26	3,660	6,281
2000	0	6,530	6,530	20,936	139	48	325	21,449	27,979
2001	372	3,631	4,003	8,436	303	271	571	9,580	13,584
2002	3,050	7,278	10,328	3,541	128	249	254	4,171	14,499
2003	222	2,396	2,618	5,972	159	53	323	6,508	9,125
2004	83	1,628	1,711	5,012	111	24	544	5,690	7,402
2005	7	3,078	3,085	4,572	314	22	411	5,320	8,405
2006	19	1,967	1,986	7,870	669	42	372	8,953	10,939
2007	28	2,190	2,218	6,208	698	38	310	7,254	9,472
2008	689	114	803	4,198	58	9	279	4,543	5,346
2009	49	0	49	3,279	54	5	269	3,607	3,656
2010	312	1,605	1,917	2,047	48	2	216	2,313	4,229
2011	1,081	1,151	2,232	1,665	202	83	124	2,074	4,306
2012	7,219	171	7,390	3,202	1,588	719	99	5,608	12,998
2013	2,071	482	2,553	11,165	438	173	133	11,909	14,462
2014	2,757	1,342	4,099	3,651	1,215	502	225	5,593	9,692
2015	3,663	5,515	9,179	4,435	7	1	243	4,686	13,865
2016	5,730	5,977	11,707	2,523	4	22	209	2,757	14,464
2017	2,224	585	2,810	1,513	45	4	245	1,808	4,617
2018	3,422	12,330	15,752	2,199	112	10	180	2,501	18,252
2019	16,777	2,297	19,074	3,783	50	5	78	3,916	22,990
2020	26,136	5,232	31,368	500	101	3	87	691	32,060
2021	7,649	1,760	9,409	847	86	0	73	1,007	10,416
2022	7,649	7,361	15,010	543	366	26	56	990	16,000

Table 2: Pacific mackerel US overfishing limits (OFL), allowable biological catches (ABC), annual catch limits (ACL), harvest guidelines (HG) since 2008. Total US landings (USA-TOT) and the percentage of ACL are also shown. Model year 2008, for example includes landings from July 1, 2008 to June 30, 2009 to align with fishery management timeframes.

Model Year	OFL	ABC	ACL	HG	USA-TOT	PercHG
2008	NA	NA	40,000	NA	4,543	11%
2009	NA	NA	10,000	NA	3,607	36%
2010	NA	NA	11,000	NA	2,313	21%
2011	44,336	42,375	40,514	30,386	2,074	7%
2012	44,336	42,375	40,514	30,386	5,608	18%
2013	57,316	52,358	52,358	39,268	11,909	30%
2014	32,992	30,138	29,170	24,170	5,593	23%
2015	25,291	23,104	21,469	20,469	4,686	23%
2016	24,983	22,822	21,161	20,161	2,757	14%
2017	30,115	27,510	26,293	25,293	1,808	7%
2018	27,662	25,269	23,840	22,840	2,501	11%
2019	14,931	13,169	11,109	10,109	3,916	39%
2020	11,772	10,289	7,950	6,950	691	10%
2021	12,145	9,446	8,323	7,323	1,007	14%
2022	9,644	7,501	5,822	4,822	990	21%

Table 3: Landings (mt) of Pacific mackerel without Pacific hake fishery bycatch (Old catch) and with (New Catch) and the difference between the two values arranged by model year.

Model Year	Old	New	Difference
2008	5,346	5,346	0
2009	3,656	3,656	0
2010	4,229	4,229	0
2011	4,305	4,305	0
2012	12,997	12,874	-123
2013	14,461	14,461	0
2014	9,691	9,707	15
2015	13,865	13,891	26
2016	14,464	14,473	9
2017	4,617	4,703	86
2018	18,252	18,352	100
2019	22,989	22,989	0
2020	32,059	32,062	2
2021	10,415	10,528	113
2022	16,000	16,128	128

Table 4: Pacific mackerel samples from the California commercial fishery and AT survey. The numbers of samples, ages, and age 8+ fish are shown for the fishery. For the AT survey, there were no age 8+ fish and the number of aged fish are shown. The numbers of lengths and weights are the same as the number of ages.

Model year	N samples	Fishery		Survey
		N fish	N 8+	N fish
2008	29	725	2	0
2009	17	440	18	0
2010	18	512	15	0
2011	26	775	4	0
2012	48	1,198	3	449
2013	72	1,800	7	9
2014	56	1,396	1	45
2015	18	447	0	26
2016	20	494	0	82
2017	9	222	0	110
2018	6	148	0	371
2019	10	250	0	289
2021	8	200	0	183
2022	1	25	0	198

Table 5: Pacific mackerel catch (mt) by landing year input to the base model. The model year for 2008, for example, includes landings from July 1, 2008 to June 30, 2009. Catch data for 2022 were used in the base model forecast file as the last model year in the assessment was 2021.

Model Year	Catch (mt)
2008	5,346
2009	3,656
2010	4,229
2011	4,305
2012	12,874
2013	14,461
2014	9,707
2015	13,891
2016	14,473
2017	4,703
2018	18,352
2019	22,989
2020	32,062
2021	10,528
2022	16,128

Table 6: Standard deviations of ageing error, arranged by age, for Pacific mackerel. Ageing error from the AT survey and fishery.

Age	Fishery	AT survey
0	0.32	0.00
1	0.32	0.00
2	0.55	0.15
3	0.79	0.23
4	1.04	0.27
5	1.31	0.30
6	1.59	0.31
7	1.88	0.32
8	2.19	0.32

Table 7: Abundance by fork length (cm) for AT summer surveys from 2012 to 2022.

FL (cm)	2012	2013	2014	2015	2016	2017	2018	2019	2021	2022
8	0	0	0	0	4,135,821	0	0	0	0	41,814,427
9	0	0	0	0	0	0	6,743,924	6,743,924	428,113	36,269,442
10	0	0	0	0	4,098,922	0	60,695,315	60,695,315	776,215	25,256,609
11	0	0	0	0	495,151	0	135,203,988	135,203,988	93,942	78,306,355
12	0	0	589,930	0	10,534	0	83,032,095	83,032,095	760,693	36,190,083
13	0	0	0	0	513,877	0	45,019,544	45,019,544	2,200,508	4,934,019
14	0	0	0	0	3,400,322	0	28,271,563	28,271,563	6,707,487	9,970,176
15	0	0	589,930	0	140,120,589	0	102,859,438	102,859,438	6,924,410	17,995,740
16	0	0	2,359,721	0	140,445,041	0	85,131,501	85,131,501	3,858,857	14,103,694
17	0	0	589,930	0	564,583	0	18,780,235	18,780,235	8,936,143	11,092,929
18	0	0	1,179,860	0	222,670	0	17,884,006	17,884,006	11,165,214	14,111,066
19	0	0	1,769,790	0	2,221,024	0	17,589,955	17,589,955	21,076,531	4,113,360
20	0	0	589,930	0	144,282,995	0	1,207,190	1,207,190	19,608,695	1,842,523
21	26,264,946	0	0	0	12,701,738	0	1,235,522	1,235,522	30,395,251	2,727,661
22	4,420,079	4,965	0	67,679	11,239,310	0	16,150,698	16,150,698	26,348,708	1,317,896
23	2,698,532	0	0	184,835	11,193,303	63,950	0	0	23,062,284	976,320
24	43,651,664	0	0	248,469	12,680,136	4,307,611	238,131	238,131	16,299,526	412,623
25	76,410,284	0	0	744,452	4,932,854	15,681,142	1,366,016	1,366,016	5,622,562	501,368
26	162,917,641	4,965	707,811	1,418,233	1,262,309	38,091,584	2,736,261	2,736,261	1,931,577	575,014
27	161,713,912	558,272	0	905,898	792,413	47,794,765	1,954,689	1,954,689	371,503	1,659,187
28	40,953,968	7,264,697	0	1,041,195	557,164	36,028,892	4,451,299	4,451,299	0	693,934
29	20,881,761	8,694,120	1,225,926	462,819	1,034,677	13,328,999	7,394,546	7,394,546	24,672	1,009,929
30	6,088,585	6,907,247	1,663,349	31,089	1,312,437	5,232,239	10,182,669	10,182,669	123,358	30,972
31	1,212,517	1,776,998	5,111,446	4,432	1,617,476	3,708,441	10,542,879	10,542,879	409,107	483,707
32	145,477	2,153,637	6,561,372	0	1,796,604	5,918,203	1,402,458	1,402,458	49,343	46,458
33	246,982	1,233,623	3,435,199	361,579	1,306,108	3,140,715	619,747	619,747	471,483	15,486
34	855,801	0	709,506	8,864	0	1,457,915	76,341	76,341	5,274,991	15,486
35	855,801	156,805	1,375,500	26,657	89,120	860,964	0	0	335,092	46,458
36	0	0	687,750	22,224	178,240	575,634	0	0	496,155	0
37	648,328	0	0	0	0	150,781	0	0	24,672	0
38	0	0	0	0	0	89,099	0	0	1,476,761	152,226
39	0	0	0	0	0	0	0	0	0	0
40	29,529	0	0	0	0	0	0	0	0	0

Table 8: Abundance by age for AT summer surveys from 2012 to 2022.

Age	2012	2013	2014	2015	2016	2017	2018	2019	2021	2022
0	194,517,355	194,517,355	8,114,309	1,796,645	466,835,981	32,409,605	622,895,074	846,230,237	136,466,340	299,590,444
1	311,577,301	311,577,301	3,934,681	2,783,473	28,436,595	93,456,933	10,787,379	35,066,954	48,554,378	5,677,958
2	39,268,492	39,268,492	5,311,950	470,493	4,269,920	44,594,499	24,239,648	12,298,455	2,521,360	786,103
3	3,401,988	3,401,988	6,318,553	265,103	3,263,356	3,231,279	2,577,133	18,454,929	4,454,624	414,419
4	11	11	4,487,425	178,617	6	938,142	134,235	2,198,805	1,649,663	76,865
5	931,954	931,954	980,029	34,098	399,567	1,696,157	136,529	977,360	1,286,213	91,540
6	20	20	2	0	0	104,313	4	10	321,283	27,816
7	NA	298,687	2	0	0	2	4	10	0	0

Table 9: Proportion of mature mackerel by age. The number of mature fish, number of total fish, and predicted proportion of mature fish by age from a binomial GLM are shown.

Age	N mature	Total fish	Predicted
0	16	106	0.12
1	88	189	0.49
2	105	120	0.87
3	66	66	0.98
4	8	8	1.00
5	5	5	1.00
6	-	-	1.00
7	1	1	1.00

Table 10: Parameter estimates in the base model. Estimated values, standard deviations (SDs), bounds (minimum and maximum), estimation phase (negative values indicate that a parameter was not estimated), status (indicates if parameters are near bounds), and prior type information (mean, SD) are shown.

Parameter	Value	Phase	Bounds	Status	SD
NatM.Lorenzen_averageFem_GP_1	0.8512	3	(0.3, 1.1)	OK	0.1074
SR.LN(R0)	13.7299	1	(5, 20)	OK	0.3424
SR_regime_BLK3repl_2007	0.2315	1	(-15, 15)	OK	0.6005
Early_InitAge_6	0.0046	3	(-6, 6)	act	0.7517
Early_InitAge_5	0.0072	3	(-6, 6)	act	0.7518
Early_InitAge_4	0.0386	3	(-6, 6)	act	0.7475
Early_InitAge_3	0.4717	3	(-6, 6)	act	0.6802
Early_InitAge_2	-0.0410	3	(-6, 6)	act	0.6734
Early_InitAge_1	-0.3502	3	(-6, 6)	act	0.6271
Main_RecrDev_2008	0.2485	1	(-6, 6)	act	0.4392
Main_RecrDev_2009	-0.4801	1	(-6, 6)	act	0.5022
Main_RecrDev_2010	0.4053	1	(-6, 6)	act	0.3876
Main_RecrDev_2011	1.1293	1	(-6, 6)	act	0.2770
Main_RecrDev_2012	-0.0590	1	(-6, 6)	act	0.2845
Main_RecrDev_2013	-0.4966	1	(-6, 6)	act	0.3082
Main_RecrDev_2014	-0.3727	1	(-6, 6)	act	0.2869
Main_RecrDev_2015	-0.0506	1	(-6, 6)	act	0.2642
Main_RecrDev_2016	0.4867	1	(-6, 6)	act	0.2339
Main_RecrDev_2017	-1.0057	1	(-6, 6)	act	0.3140
Main_RecrDev_2018	0.1863	1	(-6, 6)	act	0.2465
Main_RecrDev_2019	0.4092	1	(-6, 6)	act	0.2527
Main_RecrDev_2020	-0.1321	1	(-6, 6)	act	0.3374
Main_RecrDev_2021	-0.2685	1	(-6, 6)	act	0.3100
ForeRecr_2022	0.0000	4	(-6, 6)	act	0.7500
ForeRecr_2023	0.0000	4	(-6, 6)	act	0.7500
ForeRecr_2024	0.0000	4	(-6, 6)	act	0.7500
ForeRecr_2025	0.0000	4	(-6, 6)	act	0.7500
LnQ_base_AT(2)	-1.8024	1	(-5, 5)	OK	0.2090
LnQ_base_AT(2)_BLK4repl_2016	-1.1159	1	(-4.59, 5.41)	OK	0.2359
AgeSel_P2_FISHERY(1)	1.2516	2	(-5, 9)	OK	0.4641
AgeSel_P3_FISHERY(1)	0.0263	2	(-5, 9)	OK	0.5318
AgeSel_P4_FISHERY(1)	0.0010	2	(-5, 9)	OK	0.7657
AgeSel_P5_FISHERY(1)	-0.5394	2	(-5, 9)	OK	1.2003
AgeSel_P2_AT(2)	0.1308	2	(0, 9)	OK	0.2996
LnQ_base_AT(2)_DEVmult_2008	-0.4994	1	(-10, 10)	act	0.8324
LnQ_base_AT(2)_DEVmult_2009	0.0000	1	(-10, 10)	NO_MOVE	1.0000
LnQ_base_AT(2)_DEVmult_2010	0.0000	1	(-10, 10)	NO_MOVE	1.0000
LnQ_base_AT(2)_DEVmult_2011	0.0000	1	(-10, 10)	NO_MOVE	1.0000

LnQ_base_AT(2)_DEVmult_2012	-1.2086	1	(-10, 10)	act	0.7539
LnQ_base_AT(2)_DEVmult_2013	1.1007	1	(-10, 10)	act	0.8118
LnQ_base_AT(2)_DEVmult_2014	0.9860	1	(-10, 10)	act	0.7895
LnQ_base_AT(2)_DEVmult_2015	3.2101	1	(-10, 10)	act	0.6126
FISHERY_ARDEV_y2008_A0	0.2424	3	(-10, 10)	act	0.7511
FISHERY_ARDEV_y2008_A1	-0.5635	3	(-10, 10)	act	0.7829
FISHERY_ARDEV_y2008_A2	-0.3345	3	(-10, 10)	act	0.8461
FISHERY_ARDEV_y2008_A3	0.6389	3	(-10, 10)	act	0.8313
FISHERY_ARDEV_y2008_A4	0.0232	3	(-10, 10)	act	0.9767
FISHERY_ARDEV_y2008_A5	-0.0056	3	(-10, 10)	act	0.9958
FISHERY_ARDEV_y2009_A0	-0.5934	3	(-10, 10)	act	0.8058
FISHERY_ARDEV_y2009_A1	0.4889	3	(-10, 10)	act	0.7589
FISHERY_ARDEV_y2009_A2	-0.1044	3	(-10, 10)	act	0.8836
FISHERY_ARDEV_y2009_A3	0.1394	3	(-10, 10)	act	0.9253
FISHERY_ARDEV_y2009_A4	0.0617	3	(-10, 10)	act	0.9675
FISHERY_ARDEV_y2009_A5	0.0070	3	(-10, 10)	act	1.0016
FISHERY_ARDEV_y2010_A0	-0.2210	3	(-10, 10)	act	0.7125
FISHERY_ARDEV_y2010_A1	0.1281	3	(-10, 10)	act	0.7612
FISHERY_ARDEV_y2010_A2	0.0552	3	(-10, 10)	act	0.8035
FISHERY_ARDEV_y2010_A3	0.0219	3	(-10, 10)	act	0.9257
FISHERY_ARDEV_y2010_A4	0.0214	3	(-10, 10)	act	0.9641
FISHERY_ARDEV_y2010_A5	-0.0048	3	(-10, 10)	act	0.9943
FISHERY_ARDEV_y2011_A0	0.7787	3	(-10, 10)	act	0.7052
FISHERY_ARDEV_y2011_A1	0.2370	3	(-10, 10)	act	0.7405
FISHERY_ARDEV_y2011_A2	-0.4047	3	(-10, 10)	act	0.8801
FISHERY_ARDEV_y2011_A3	-0.4615	3	(-10, 10)	act	0.8821
FISHERY_ARDEV_y2011_A4	-0.1278	3	(-10, 10)	act	0.9521
FISHERY_ARDEV_y2011_A5	-0.0194	3	(-10, 10)	act	0.9907
FISHERY_ARDEV_y2012_A0	0.7409	3	(-10, 10)	act	0.6649
FISHERY_ARDEV_y2012_A1	-0.0879	3	(-10, 10)	act	0.6697
FISHERY_ARDEV_y2012_A2	-0.2956	3	(-10, 10)	act	0.7773
FISHERY_ARDEV_y2012_A3	-0.2296	3	(-10, 10)	act	0.9081
FISHERY_ARDEV_y2012_A4	-0.1259	3	(-10, 10)	act	0.9277
FISHERY_ARDEV_y2012_A5	-0.0024	3	(-10, 10)	act	0.9969
FISHERY_ARDEV_y2013_A0	2.0499	3	(-10, 10)	act	0.6589
FISHERY_ARDEV_y2013_A1	-0.8610	3	(-10, 10)	act	0.7223
FISHERY_ARDEV_y2013_A2	-0.7114	3	(-10, 10)	act	0.7257
FISHERY_ARDEV_y2013_A3	-0.4969	3	(-10, 10)	act	0.8362
FISHERY_ARDEV_y2013_A4	-0.0429	3	(-10, 10)	act	0.9619
FISHERY_ARDEV_y2013_A5	0.0544	3	(-10, 10)	act	1.0083
FISHERY_ARDEV_y2014_A0	1.0635	3	(-10, 10)	act	0.6532
FISHERY_ARDEV_y2014_A1	-0.5969	3	(-10, 10)	act	0.7478
FISHERY_ARDEV_y2014_A2	-0.4107	3	(-10, 10)	act	0.7928
FISHERY_ARDEV_y2014_A3	-0.2878	3	(-10, 10)	act	0.7990
FISHERY_ARDEV_y2014_A4	0.1746	3	(-10, 10)	act	0.9508
FISHERY_ARDEV_y2014_A5	0.0414	3	(-10, 10)	act	1.0173
FISHERY_ARDEV_y2015_A0	0.0477	3	(-10, 10)	act	0.6777
FISHERY_ARDEV_y2015_A1	-0.3176	3	(-10, 10)	act	0.7321
FISHERY_ARDEV_y2015_A2	0.3692	3	(-10, 10)	act	0.8641
FISHERY_ARDEV_y2015_A3	0.0366	3	(-10, 10)	act	0.9360
FISHERY_ARDEV_y2015_A4	-0.1217	3	(-10, 10)	act	0.9157
FISHERY_ARDEV_y2015_A5	-0.0135	3	(-10, 10)	act	0.9932
FISHERY_ARDEV_y2016_A0	-1.4574	3	(-10, 10)	act	0.7438
FISHERY_ARDEV_y2016_A1	0.5794	3	(-10, 10)	act	0.7129
FISHERY_ARDEV_y2016_A2	1.0106	3	(-10, 10)	act	0.7988
FISHERY_ARDEV_y2016_A3	-0.0405	3	(-10, 10)	act	0.9740
FISHERY_ARDEV_y2016_A4	-0.0677	3	(-10, 10)	act	0.9642
FISHERY_ARDEV_y2016_A5	-0.0234	3	(-10, 10)	act	0.9870
FISHERY_ARDEV_y2017_A0	0.1543	3	(-10, 10)	act	0.7895
FISHERY_ARDEV_y2017_A1	-0.4137	3	(-10, 10)	act	0.7692
FISHERY_ARDEV_y2017_A2	0.2454	3	(-10, 10)	act	0.8603
FISHERY_ARDEV_y2017_A3	0.0184	3	(-10, 10)	act	0.9816
FISHERY_ARDEV_y2017_A4	-0.0088	3	(-10, 10)	act	0.9929
FISHERY_ARDEV_y2017_A5	0.0023	3	(-10, 10)	act	0.9993

FISHERY_ARDEV_y2018_A0	-0.5851	3	(-10, 10)	act	0.8178
FISHERY_ARDEV_y2018_A1	0.9818	3	(-10, 10)	act	0.8146
FISHERY_ARDEV_y2018_A2	-0.3503	3	(-10, 10)	act	0.8660
FISHERY_ARDEV_y2018_A3	-0.0509	3	(-10, 10)	act	0.9595
FISHERY_ARDEV_y2018_A4	0.0044	3	(-10, 10)	act	0.9990
FISHERY_ARDEV_y2018_A5	0.0001	3	(-10, 10)	act	1.0000
FISHERY_ARDEV_y2019_A0	-0.7501	3	(-10, 10)	act	0.7617
FISHERY_ARDEV_y2019_A1	0.0909	3	(-10, 10)	act	0.7749
FISHERY_ARDEV_y2019_A2	0.1224	3	(-10, 10)	act	1.0154
FISHERY_ARDEV_y2019_A3	0.4972	3	(-10, 10)	act	0.8856
FISHERY_ARDEV_y2019_A4	0.0362	3	(-10, 10)	act	1.0071
FISHERY_ARDEV_y2019_A5	0.0031	3	(-10, 10)	act	1.0014
FISHERY_ARDEV_y2020_A0	-0.1656	3	(-10, 10)	act	0.9346
FISHERY_ARDEV_y2020_A1	0.4628	3	(-10, 10)	act	0.8811
FISHERY_ARDEV_y2020_A2	-0.1527	3	(-10, 10)	act	0.9218
FISHERY_ARDEV_y2020_A3	-0.0877	3	(-10, 10)	act	0.9675
FISHERY_ARDEV_y2020_A4	-0.0550	3	(-10, 10)	act	0.9767
FISHERY_ARDEV_y2020_A5	-0.0018	3	(-10, 10)	act	0.9991
FISHERY_ARDEV_y2021_A0	-1.3047	3	(-10, 10)	act	0.7206
FISHERY_ARDEV_y2021_A1	-0.1281	3	(-10, 10)	act	0.7003
FISHERY_ARDEV_y2021_A2	0.9614	3	(-10, 10)	act	0.7542
FISHERY_ARDEV_y2021_A3	0.3025	3	(-10, 10)	act	0.8615
FISHERY_ARDEV_y2021_A4	0.0687	3	(-10, 10)	act	1.0305
FISHERY_ARDEV_y2021_A5	0.0953	3	(-10, 10)	act	1.0375

Table 11: Likelihood components, parameters, and biomass estimates.

	Description	Value	
Likelihood	TOTAL	115.003	
	Catch	0	
	Equil_catch	0	
	Survey	-6.303	
	Length_comp	0	
	Age_comp	29.552	
	Recruitment	0.095	
	InitEQ_Regime	0.028	
	Forecast_Recruitment	0	
	Parm_priors	2.791	
	Parm_softbounds	0.003	
	Parm_devs	88.837	
	Crash_Pen	0	
	Parameter	NatM_Lorenzen_averageFem_GP_1	0.851
SR_LN(R0)		13.73	
SR_BH_steep		0.75	
SR_sigmaR		0.75	
SR_regime_BLK3repl_2007		0.232	
LnQ_base_AT(2)		-1.802	
LnQ_base_AT(2)_BLK4repl_2016		-1.116	
LnQ_base_AT(2)_dev_se		0.25	
LnQ_base_AT(2)_dev_autocorr		0	
LnQ_base_AT(2)_DEVmult_2008		-0.499	
LnQ_base_AT(2)_DEVmult_2009		0	
LnQ_base_AT(2)_DEVmult_2010		0	
LnQ_base_AT(2)_DEVmult_2011		0	
LnQ_base_AT(2)_DEVmult_2012		-1.209	
LnQ_base_AT(2)_DEVmult_2013		1.101	
LnQ_base_AT(2)_DEVmult_2014		0.986	
LnQ_base_AT(2)_DEVmult_2015		3.21	
Biomass (mt)		2020 Age1+	69,479
		2021 Age1+	49,977

Table 12: Pacific mackerel numbers-at-age (thousands of fish) estimated in the base model

Model Year	Age 0	Age 1	Age 2	Age 3	Age 4	Age 5	Age 6	Age 7	Age 8+
VIRG	917,938	276,684	103,273	42,810	18,820	8,571	3,990	1,884	1,732
INIT	1,157,070	348,764	130,176	53,963	23,723	10,803	5,030	2,375	2,183
2008	1,038,020	221,136	116,338	83,311	24,573	10,881	5,053	2,375	2,183
2009	475,290	307,718	80,422	46,636	33,510	10,882	5,038	2,384	2,183
2010	1,089,720	142,634	109,772	32,493	19,840	14,993	5,049	2,378	2,188
2011	2,180,780	324,100	49,825	42,716	13,436	8,718	6,931	2,382	2,186
2012	686,338	646,682	117,019	20,287	18,462	6,035	4,046	3,271	2,188
2013	442,058	195,633	221,643	45,178	8,265	8,004	2,778	1,908	2,612
2014	488,121	101,094	69,282	86,300	18,376	3,505	3,667	1,309	2,167
2015	642,887	132,723	35,236	26,388	34,472	7,658	1,606	1,728	1,667
2016	1,070,520	179,320	41,043	9,951	8,807	13,687	3,456	755	1,625
2017	249,293	319,349	50,698	10,970	3,752	3,676	6,254	1,628	1,142
2018	806,313	73,290	113,448	19,053	4,460	1,634	1,696	2,950	1,326
2019	984,076	231,090	11,757	37,410	6,149	1,679	732	797	2,043
2020	569,096	284,372	61,284	3,393	9,706	2,307	752	344	1,362
2021	473,902	156,784	58,879	18,321	1,052	3,582	1,026	353	819

Table 13: Pacific mackerel biomass-at-age estimated for the base model

Model year	0	1	2	3	4	5	6	7	8+	Total Age0+	Total Age1+
VIRG	107,598	65,480	34,141	17,310	8,507	4,044	2,035	996	909	241,021	133,423
INIT	135,629	82,539	43,036	21,819	10,724	5,098	2,565	1,255	1,146	303,811	168,182
2008	121,674	52,334	38,461	33,686	11,108	5,135	2,577	1,255	1,146	267,376	145,702
2009	55,712	72,825	26,587	18,857	15,148	5,135	2,570	1,260	1,146	199,239	143,527
2010	127,734	33,756	36,290	13,138	8,969	7,075	2,575	1,256	1,148	231,941	104,207
2011	255,624	76,702	16,472	17,272	6,074	4,114	3,535	1,259	1,147	382,198	126,574
2012	128,024	146,604	30,604	7,941	8,111	2,916	2,094	1,718	1,149	329,161	201,137
2013	56,071	58,224	72,206	16,218	3,631	3,868	1,437	1,002	1,372	214,028	157,958
2014	61,913	35,042	26,825	35,756	8,459	1,694	1,897	687	1,138	173,411	111,498
2015	126,988	31,685	14,452	10,683	15,397	3,798	831	907	875	205,618	78,630
2016	64,853	42,996	13,106	4,081	3,721	6,932	1,951	397	854	138,890	74,037
2017	54,630	79,867	14,298	3,647	1,539	1,638	3,801	1,031	600	161,050	106,420
2018	21,828	20,319	38,351	7,154	1,728	670	756	1,792	839	93,438	71,610
2019	8,143	48,899	4,310	15,594	2,714	748	300	355	1,241	82,304	74,161
2020	72,184	39,424	21,224	1,553	5,089	1,018	422	141	607	141,663	69,479
2021	43,846	21,884	15,838	8,813	578	1,878	453	198	336	93,823	49,977

Table 14: Spawning stock biomass (SSB) and recruitment (1000s of fish) estimates and asymptotic standard errors for the base model.

Year	SSB	SSB SD	Recruits	Recruits SD
Virgin	108,420	20,365	917,938	314,313
Initial	136,664	84,074	1,157,070	833,104
2008	129,863	71,817	1,038,020	619,869
2009	102,687	44,607	475,290	316,083
2010	82,111	31,628	1,089,720	621,740
2011	85,120	29,231	2,180,780	994,322
2012	122,554	36,514	686,338	318,134
2013	118,786	34,710	442,058	193,156
2014	89,487	26,120	488,121	208,439
2015	60,413	17,585	642,887	250,514
2016	50,421	14,176	1,070,520	403,597
2017	63,936	16,709	249,293	113,096
2018	56,224	14,628	806,313	308,286
2019	48,433	13,255	984,076	394,732
2020	46,687	14,062	569,096	250,638
2021	36,646	13,880	473,902	225,892

Table 15: Total (age-0+) and summary (age-1+) biomass values (mt) estimated on June 1 of each year.

Year	Age-0+	Age-1+
2008	267,376	145,702
2009	199,239	143,527
2010	231,941	104,207
2011	382,198	126,574
2012	329,161	201,137
2013	214,028	157,958
2014	173,411	111,498
2015	205,618	78,630
2016	138,890	74,037
2017	161,050	106,420
2018	93,438	71,610
2019	82,303	74,160
2020	141,663	69,479
2021	93,822	49,977

Table 16: Annual exploitation rate (calendar year landings / total age-0+ biomass values).

Year	Exploitation rate
2008	0.02
2009	0.03
2010	0.01
2011	0.01
2012	0.03
2013	0.07
2014	0.05
2015	0.06
2016	0.14
2017	0.04
2018	0.18
2019	0.21
2020	0.27
2021	0.10

Table 17: Parameter estimates, summary biomass (age 1+; mt), and total likelihood values associated with fixed values of steepness ranging from 0.25 to 1. The base model steepness value was 0.75.

	Steepness									
	0.25	0.3	0.4	0.5	0.6	0.7	0.75	0.8	0.9	1
NatM.Lorenzen.averageFem.GP_1	0.869	0.866	0.861	0.857	0.854	0.852	0.851	0.850	0.849	0.848
SR.LN(R0)	15.108	14.481	14.035	13.880	13.799	13.748	13.730	13.714	13.690	13.672
SR.regime.BLK3repl.2007	-0.366	-0.132	0.115	0.188	0.215	0.228	0.232	0.234	0.237	0.239
LnQ_base_AT(2)	-1.922	-1.864	-1.835	-1.821	-1.812	-1.805	-1.802	-1.800	-1.796	-1.794
LnQ_base_AT(2).BLK4repl.2016	-1.109	-1.093	-1.089	-1.095	-1.104	-1.112	-1.116	-1.119	-1.126	-1.131
LnQ_base_AT(2).DEVmult.2008	0.114	-0.194	-0.342	-0.410	-0.455	-0.487	-0.499	-0.510	-0.529	-0.543
LnQ_base_AT(2).DEVmult.2012	-1.024	-1.117	-1.157	-1.178	-1.193	-1.204	-1.209	-1.212	-1.219	-1.224
LnQ_base_AT(2).DEVmult.2013	1.217	1.162	1.135	1.121	1.111	1.104	1.101	1.098	1.094	1.091
LnQ_base_AT(2).DEVmult.2014	1.089	1.045	1.017	1.003	0.995	0.988	0.986	0.984	0.981	0.978
LnQ_base_AT(2).DEVmult.2015	3.164	3.183	3.190	3.197	3.203	3.208	3.210	3.212	3.216	3.219
2020 Age-1+ bio	68,452	67,459	66,989	67,509	68,300	69,102	69,479	69,834	70,475	71,030
2021 Age-1+ bio	44,700	45,509	46,332	47,422	48,531	49,529	49,977	50,391	51,124	51,746
2022 Age-1+ bio	37,816	40,101	41,957	43,552	44,964	46,161	46,683	47,158	47,988	48,683
2023 Age-1+ bio	30,362	36,246	43,296	48,340	51,982	54,629	55,680	56,592	58,086	59,252
2024 Age-1+ bio	20,355	29,377	41,730	50,163	55,706	59,398	60,785	61,950	63,786	65,154
Total likelihood	116.766	115.797	115.251	115.087	115.029	115.008	115.003	115.001	115.001	115.003

Table 18: Parameter estimates, summary biomass (age 1+; mt), and total likelihood values associated with fixed values of 2016-2021 log catchability (Q) values. The deviations for Q values for 2008-2015 were estimated. Column headers show the Q values in normal space.

	Fixed 2021 catchability (Q)										
	0.1	0.2	0.3	0.33	0.4	0.5	0.6	0.7	0.8	0.9	1
NatM.Lorenzen.averageFem.GP.1	0.863	0.860	0.853	0.851	0.846	0.839	0.834	0.829	0.825	0.822	0.820
SR.LN(R0)	14.567	14.063	13.787	13.730	13.604	13.472	13.372	13.293	13.229	13.176	13.131
SR.BH.steep	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750
SR.regime.BLK3repl.2007	-0.079	0.116	0.213	0.232	0.270	0.306	0.332	0.352	0.369	0.383	0.396
LnQ.base.AT(2)	-2.046	-1.904	-1.821	-1.802	-1.761	-1.715	-1.678	-1.646	-1.619	-1.595	-1.574
LnQ.base.AT(2).BLK4repl.2016	-2.303	-1.609	-1.204	-1.116	-0.916	-0.693	-0.511	-0.357	-0.223	-0.105	0.000
LnQ.base.AT(2).DEVmult.2008	-0.160	-0.367	-0.477	-0.499	-0.550	-0.603	-0.644	-0.676	-0.701	-0.722	-0.738
LnQ.base.AT(2).DEVmult.2012	-0.676	-1.006	-1.174	-1.209	-1.284	-1.364	-1.426	-1.476	-1.517	-1.551	-1.579
LnQ.base.AT(2).DEVmult.2013	1.495	1.256	1.127	1.101	1.042	0.978	0.928	0.886	0.852	0.822	0.797
LnQ.base.AT(2).DEVmult.2014	1.463	1.174	1.018	0.986	0.915	0.840	0.781	0.733	0.692	0.658	0.628
LnQ.base.AT(2).DEVmult.2015	3.544	3.350	3.235	3.210	3.153	3.089	3.037	2.992	2.952	2.916	2.883
2020 Age-1+ bio	222,890	111,583	75,398	69,479	58,158	48,467	42,458	38,460	35,647	33,572	31,971
2021 Age-1+ bio	189,304	88,279	55,371	49,977	39,637	30,740	25,176	21,434	18,774	16,797	15,273
2022 Age-1+ bio	177,338	82,774	51,795	46,683	36,824	28,239	22,784	19,057	16,367	14,339	12,756
2023 Age-1+ bio	185,634	92,761	61,056	55,680	45,125	35,651	29,437	25,076	21,851	19,360	17,368
2024 Age-1+ bio	189,369	98,197	66,282	60,785	49,875	39,894	33,198	28,387	24,743	21,867	19,525
Total likelihood	128.629	119.077	117.052	115.003	117.350	118.677	120.545	122.725	125.086	127.553	130.075

Table 19: Parameter estimates, summary biomass (age 1+; mt), and total likelihood values associated with fixed values of average age-specific natural mortality ($j_i M_j / i_i$). Note that for this configuration steepness was fixed at 0.75.

	Average age-specific natural mortality (M)								
	0.3	0.4	0.5	0.6	0.7	0.8	0.851	0.9	1
SR_LN(R0)	12.014	12.339	12.681	13.009	13.313	13.592	13.730	13.859	14.119
SR_BH_steep	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750
SR_regime_BLK3repl.2007	-0.367	-0.255	-0.135	-0.020	0.085	0.179	0.232	0.279	0.368
LnQ_base_AT(2)	-1.353	-1.441	-1.545	-1.640	-1.723	-1.790	-1.802	-1.814	-1.835
LnQ_base_AT(2)_BLK4repl.2016	-0.687	-0.810	-0.926	-1.010	-1.070	-1.110	-1.116	-1.122	-1.138
LnQ_base_AT(2)_DEVmult.2008	-1.133	-1.093	-0.977	-0.833	-0.684	-0.545	-0.499	-0.458	-0.379
LnQ_base_AT(2)_DEVmult.2012	-2.068	-1.966	-1.812	-1.631	-1.439	-1.259	-1.209	-1.162	-1.071
LnQ_base_AT(2)_DEVmult.2013	0.533	0.642	0.775	0.897	1.002	1.084	1.101	1.115	1.143
LnQ_base_AT(2)_DEVmult.2014	0.503	0.611	0.734	0.840	0.925	0.986	0.986	0.985	0.983
LnQ_base_AT(2)_DEVmult.2015	2.920	3.012	3.088	3.144	3.191	3.226	3.210	3.196	3.170
2020 Age-1+ bio	50,902	56,674	62,249	65,666	67,302	68,238	69,479	70,897	74,465
2021 Age-1+ bio	38,528	43,854	48,307	50,320	50,552	50,033	49,977	50,024	50,375
2022 Age-1+ bio	48,413	52,802	55,025	53,917	50,705	47,062	46,683	46,454	46,302
2023 Age-1+ bio	52,651	56,399	58,268	57,736	56,149	54,888	55,680	56,648	59,191
2024 Age-1+ bio	51,916	55,377	57,641	58,380	58,606	59,252	60,785	62,424	66,249
Total likelihood	155.673	139.571	129.178	122.410	118.544	117.102	115.003	117.083	117.842

Table 20: Variance adjustment, parameter estimates, summary biomass (age-1+; mt) and total NLL from the base model and a model with Francis reweighting of age compositions.

	Base model	Francis
Fishery age comps	-	4.161
AT Survey age comps	-	0.508
NatM_Lorenzen_averageFem_GP_1	0.851	0.826
SR_LN(R0)	13.730	13.766
SR_BH_steep	0.750	0.750
SR_regime_BLK3repl_2007	0.232	0.116
LnQ_base_AT(2)	-1.802	-1.793
LnQ_base_AT(2)_BLK4repl_2016	-1.116	-1.278
LnQ_base_AT(2)_DEVmult_2008	-0.499	-0.549
LnQ_base_AT(2)_DEVmult_2012	-1.209	-1.325
LnQ_base_AT(2)_DEVmult_2013	1.101	1.073
LnQ_base_AT(2)_DEVmult_2014	0.986	1.041
LnQ_base_AT(2)_DEVmult_2015	3.210	3.281
2020 Age-1+ bio	69,479	81,480
2021 Age-1+ bio	49,977	58,478
2022 Age-1+ bio	46,682	54,189
2023 Age-1+ bio	55,680	63,634
2024 Age-1+ bio	60,785	69,408
Total likelihood	115.003	117.227

Table 21: Parameter estimates, summary biomass (age-1+; mt) and total NLL from the base model and a model with fishery and AT survey age compositions downweighted. Fishery age compositions had lambda of 0.5 and AT survey age compositions had a lambda of 0.5 for each of the respective runs.

	Base model	Fishery down	AT survey down
NatM_Lorenzen_averageFem_GP_1	0.851	0.848	0.832
SR_LN(R0)	13.730	13.673	13.702
SR_BH_steep	0.750	0.750	0.750
SR_regime_BLK3repl_2007	0.232	0.268	0.232
LnQ_base_AT(2)	-1.802	-1.789	-1.779
LnQ_base_AT(2)_BLK4repl_2016	-1.116	-1.052	-1.151
LnQ_base_AT(2)_DEVmult_2008	-0.499	-0.484	-0.491
LnQ_base_AT(2)_DEVmult_2012	-1.209	-1.221	-1.342
LnQ_base_AT(2)_DEVmult_2013	1.101	1.081	1.049
LnQ_base_AT(2)_DEVmult_2014	0.986	0.945	0.989
LnQ_base_AT(2)_DEVmult_2015	3.210	3.167	3.208
2020 Age-1+ bio	69,479	65,216	72,666
2021 Age-1+ bio	49,977	46,366	51,732
2022 Age-1+ bio	46,682	43,377	48,010
2023 Age-1+ bio	55,680	51,838	56,779
2024 Age-1+ bio	60,785	56,533	61,932
Total likelihood	115.003	109.775	103.493

Table 22: Summary biomass forecast values with assumed future catch values equal to the 25th percentile catch value (perc25; 5,699mt) and 75th percentile catch value (perc75; 17,177mt). The benchmark model uses an assumed average catch value from 2019-2022 of 20,366mt.

Year	benchmark	perc25	perc75
2022	46,683	46,683	46,683
2023	55,680	55,680	55,680
2024	60,785	69,010	62,523
2025	65,477	79,866	68,524

Table 23: Parameter estimates, summary biomass (age-1+; mt) and total NLL with values of 2022 AT survey biomass ranging from 10,000 to 100,000mt.

	10,000	20,000	30,000	40,000	50,000	60,000	70,000	80,000	90,000	100,000
NatM.Lorenzen_averageFem_GP_1	0.857	0.852	0.851	0.852	0.853	0.854	0.854	0.855	0.856	0.857
SR_LN(R0)	13.611	13.692	13.723	13.738	13.748	13.754	13.759	13.763	13.767	13.770
SR_BH_steep	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750
SR_regime_BLK3repl_2007	0.300	0.252	0.235	0.228	0.226	0.224	0.224	0.224	0.224	0.224
LnQ_base_AT(2)	-1.776	-1.791	-1.800	-1.806	-1.811	-1.815	-1.819	-1.821	-1.824	-1.826
LnQ_base_AT(2)_BLK4repl_2016	-1.058	-1.107	-1.116	-1.115	-1.113	-1.110	-1.108	-1.106	-1.104	-1.102
LnQ_base_AT(2)_DEVmult_2008	-0.546	-0.520	-0.504	-0.493	-0.484	-0.476	-0.470	-0.465	-0.460	-0.456
LnQ_base_AT(2)_DEVmult_2012	-1.256	-1.232	-1.215	-1.200	-1.189	-1.179	-1.171	-1.164	-1.158	-1.153
LnQ_base_AT(2)_DEVmult_2013	1.055	1.082	1.096	1.106	1.114	1.120	1.124	1.128	1.132	1.135
LnQ_base_AT(2)_DEVmult_2014	0.936	0.966	0.981	0.992	1.000	1.006	1.010	1.015	1.018	1.021
LnQ_base_AT(2)_DEVmult_2015	3.200	3.205	3.209	3.212	3.215	3.218	3.220	3.222	3.224	3.226
2020 Age-1+ bio	56,998	65,846	68,930	69,979	70,318	70,400	70,384	70,328	70,258	70,185
2021 Age-1+ bio	34,296	44,997	49,159	50,790	51,472	51,781	51,929	52,002	52,037	52,052
2022 Age-1+ bio	29,641	41,336	45,838	47,480	48,066	48,256	48,288	48,252	48,189	48,115
2023 Age-1+ bio	18,464	35,399	49,541	62,734	75,444	87,814	99,910	111,771	123,431	134,911
2024 Age-1+ bio	29,685	46,053	58,146	68,801	78,743	88,224	97,368	106,244	114,901	123,373
Total likelihood	117.215	114.168	113.605	113.623	113.818	114.072	114.344	114.617	114.886	115.147

Table 24: Parameter estimates, summary biomass (age-1+; mt) and total NLL with values of 2023 AT survey biomass ranging from 10,000 to 100,000mt.

	10,000	20,000	30,000	40,000	50,000	60,000	70,000	80,000	90,000	100,000
NatM.Lorenzen_averageFem_GP_1	0.857	0.850	0.850	0.852	0.854	0.856	0.858	0.859	0.860	0.861
SR_LN(R0)	13.617	13.688	13.717	13.735	13.748	13.758	13.765	13.771	13.775	13.779
SR_BH_steep	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750	0.750
SR_regime_BLK3repl_2007	0.292	0.248	0.235	0.231	0.230	0.230	0.231	0.231	0.232	0.233
LnQ_base_AT(2)	-1.768	-1.787	-1.797	-1.804	-1.810	-1.814	-1.818	-1.821	-1.824	-1.827
LnQ_base_AT(2)_BLK4repl_2016	-1.050	-1.110	-1.117	-1.115	-1.111	-1.107	-1.103	-1.100	-1.097	-1.095
LnQ_base_AT(2)_DEVmult_2008	-0.558	-0.527	-0.509	-0.495	-0.484	-0.475	-0.468	-0.461	-0.456	-0.451
LnQ_base_AT(2)_DEVmult_2012	-1.274	-1.241	-1.220	-1.204	-1.190	-1.179	-1.170	-1.162	-1.155	-1.148
LnQ_base_AT(2)_DEVmult_2013	1.042	1.076	1.093	1.104	1.112	1.119	1.125	1.129	1.133	1.137
LnQ_base_AT(2)_DEVmult_2014	0.921	0.960	0.978	0.989	0.997	1.004	1.009	1.014	1.017	1.021
LnQ_base_AT(2)_DEVmult_2015	3.191	3.205	3.209	3.211	3.212	3.214	3.215	3.217	3.218	3.220
2020 Age-1+ bio	59,505	66,832	68,928	69,624	69,819	69,808	69,711	69,580	69,438	69,297
2021 Age-1+ bio	37,103	46,046	49,053	50,258	50,768	50,972	51,033	51,024	50,982	50,926
2022 Age-1+ bio	32,558	42,396	45,706	46,966	47,434	47,558	47,527	47,425	47,294	47,154
2023 Age-1+ bio	23,852	39,811	49,788	56,762	61,686	65,080	67,347	68,822	69,765	70,361
2024 Age-1+ bio	15,157	34,714	50,978	65,648	79,308	92,253	104,674	116,703	128,432	139,924
Total likelihood	116.613	114.216	113.645	113.587	113.720	113.931	114.175	114.430	114.686	114.939

8 Figures




Figure 1: Map of Pacific mackerel stock distribution, spawning range, and fisheries. Created by Paul Crone.




Figure 2: Pacific mackerel landings (mt) by major fishing region in Mexico (a) and USA (b). Landings from Ensenada (BC) and Magdalena Bay (BCS) are shown in the top panel. Landings from California (CA), California recreational sector (CA-REC), Oregon (OR), and Washington (WA) are shown in the bottom panel. Landings were grouped by model year which spans July 1 to June 30 of the following calendar year.




Figure 3: Summary of data sources used in the base model. Note, length compositions were available for the years shown and 2019 and 2021, but the base model was not fit to any length-composition data.




Figure 4: Catch time series input to the stock assessment. Catches from all fishing regions were summed by model year.




Figure 5: Age composition data for the fishery arranged by model year. The input sample sizes (numbers of measured fish/25) are shown in the top right of each panel. One sample (25 measured fish) was available for model year 2022 but not included in the assessment.




Figure 6: Ageing error estimated for the fishery and AT survey.




Figure 7: Weight-at-age data for Pacific mackerel arranged by fleet (columns) and cohort model year (rows). Numbers shown in the bottom right are the number of individual fish measured for each cohort. Panels are arranged by cohort because missing weight-at-age values were interpolated as necessary by cohort.




Figure 8: Acoustic-trawl survey biomass time series used in the 2019 benchmark assessment (red) and 2023 benchmark (blue). The differences are due to an updated length-weight relationship for Pacific mackerel, and for the 2015 estimate a reanalysis of the echogram. The 95% CIs are shown as the vertical bars.




Figure 9: Biomass densities (colored points) of Pacific mackerel, per stratum in the core survey regions from the summer 2021 AT survey. Thick gray lines represent acoustic transects. A majority of the biomass density was observed in Mexican waters (65%).




Figure 10: AT survey index of abundance values in untransformed space.




Figure 11: Age composition data for the AT survey arranged by model year. The input sample sizes are the numbers of clusters per model year.




Figure 12: Annual age-length keys derived from summer AT survey samples from 2012 to 2022. There were pooled age-length keys for 2013-2015 and 2021-2022 due to low sample sizes.




Figure 13: Pacific mackerel maturity-at-age. The observed proportion mature are shown (points; point size represents number of fish) and the predicted proportion mature based on a binomial GLM.




Figure 14: Model bridging between the 2019 (dashed lines) and 2023 benchmark assessments. Features were modified one at a time and added cumulatively. The panels show the 2019 benchmark model (Benchmark2019), the 2019 benchmark model with SS3.30.20 (SS3.30.20), updated data through 2021 (New data), modified fishery and AT selectivity patterns (Modify selex patterns), recruitment deviation period to 2021 (Extend rec dev period), age compositions equally weighted at 1 (Even data weighting), SR_regime parameter (Add SR_regime block), deviations for 2008-2015 and block for 2016-2021 Q values (dev and block Q s), age-specific M (Age-specific M), and the current base model (Benchmark2023).




Figure 15: Age-based selectivity patterns for the fishery (time-varying) and AT survey.




Figure 16: Fits to the age-composition time series for the fishery in the base model. Values in the top right are input sample sizes (N_{adj}) and effective sample sizes given the fit of the model (N_{eff}).




Figure 17: Residuals from the fits to the age-composition time series for the fishery data in the base model.




Figure 18: Fits to the age-composition time series for the AT survey in the base model. Values in the top right are input sample sizes (N_{adj}) and effective sample sizes given the fit of the model (N_{eff}).




Figure 19: Residuals from the fits to the age-composition time series for the AT survey data in the base model.




Figure 20: Fit to the index data (blue line) for the AT survey in normal space. Vertical lines indicate 95% uncertainty intervals around index values based on the model assumption of lognormal error.




Figure 21: Fit to the index data (blue line) for the AT survey in log space. Vertical lines indicate 95% uncertainty intervals around index values based on the model assumption of lognormal error.




Figure 22: Catchability (Q) values through time. Blocks span the years 2008-2012, 2013-2015, 2016-2019 and 2021. The 2021 Q value was fixed and the other blocks were estimated with a prior centered at the 2021 Q value.




Figure 23: Time-invariant, age-specific natural mortality values.




Figure 24: Estimated stock-recruit (Beverton-Holt) relationship with steepness fixed at 0.75. Year numbers indicate the first, last, and years with (log) deviations > 0.5.




Figure 25: Recruitment deviations with 95% intervals for the base model $\sigma_R = 0.75$.




Figure 26: Asymptotic standard errors for the estimated recruitment deviations.




Figure 27: Recruitment bias adjustment plot for the early, main, and forecast periods.




Figure 28: Population numbers at age from the base model.




Figure 29: Estimated spawning stock biomass time series (mt) with 95% confidence intervals (dashed lines).




Figure 30: Estimated recruitment time series (billions fish) with 95% confidence intervals (dashed lines).




Figure 31: Estimated summary (age-1+) biomass (mt) with 95% confidence intervals.




Figure 32: Continuous fishing mortality (F) estimates.




Figure 33: Annual exploitation rates (calendar year landings/ July total biomass).




Figure 34: Historical analysis comparing the estimates of summary biomass for the base model to the 2019 benchmark, and a selection of models dating back to 2005. The top panel shows the longer time series of biomass, and the bottom panel shows time series dating back to 2000.




Figure 35: Likelihood profile for values of steepness (h) ranging from 0.25 to 1. Values within 1.92 units of the MLE (dashed horizontal line) are within the 95% confidence interval and the vertical dashed line is the fixed steepness value of 0.75 assumed in the base model.




Figure 36: Likelihood profile for values of catchability (Q) ranging from 0.1 to 1. Values within 1.92 units of the MLE (dashed horizontal line) are within the 95% confidence interval. Note that the $\text{Ln}Q$ values were implemented in SS although the values are shown in normal space here.




Figure 37: Likelihood profile for values of natural mortality (M) ranging from 0.3 to 1 and steepness fixed at 0.75. Values within 1.92 units of the MLE (dashed horizontal line) are within the 95% confidence interval.




Figure 38: Age-1+ summary biomass (mt) values estimated from the base model (solid line) and the model with Francis reweighting (dashed line) for the age composition from the fishing and AT survey fleets.




Figure 39: Age-1+ summary biomass (mt) values estimated from the base model and models with AT survey age compositions and fishery age compositions downweighted individually ($\lambda = 0.5$) in the total likelihood calculation.




Figure 40: Forecasted summary biomass values assuming future catch values equal to the 25th percentile catch value (perc25; 5,699mt) and 75th percentile catch value (perc75; 17,177mt). The benchmark model uses an assumed average catch value from 2019-2022 of 20,366mt.




Figure 41: Likelihood profile for values of 2022 AT survey biomass ranging from 10,000 mt to 100,000 mt in increments of 10,000. Values within 1.92 units of the MLE (dashed horizontal line) are within the 95% confidence interval. The block on Q was extended to be 2016-2022 in these runs.




Figure 42: Likelihood profile for values of 2023 AT survey biomass ranging from 10,000 mt to 100,000 mt in increments of 10,000. Values within 1.92 units of the MLE (dashed horizontal line) are within the 95% confidence interval. The block on Q was extended to be 2016-2023 in these runs.




Figure 43: Summary biomass estimates for models with 2022 AT survey biomass values ranging from 10,000 mt to 100,000 mt in increments of 10,000.




Figure 44: Summary biomass estimates for models with 2023 AT survey biomass values ranging from 10,000 mt to 100,000 mt in increments of 10,000.




Figure 45: Retrospective analysis of summary biomass estimates. One year of data is removed at a time for each model run.

Harvest Control Rule Formulas					
OFL = BIOMASS * E_{MSY} * DISTRIBUTION					
$ABC_{P\text{-star}} = BIOMASS * BUFFER_{P\text{-star}} * E_{MSY} * DISTRIBUTION$					
HG = (BIOMASS - CUTOFF) * FRACTION * DISTRIBUTION					
Harvest Formula Parameters					
BIOMASS (ages 1+, mt)	55,681				
P-star	0.45	0.40	0.35	0.30	0.25
ABC Buffer _{Cat 1}	0.9261	0.8567	0.7903	0.7260	0.6624
ABC Buffer _{Cat 2}	0.9133	0.8330	0.7573	0.6851	0.6148
$E_{MSY} \equiv$ FRACTION	0.30				
CUTOFF (mt)	18,200				
DISTRIBUTION (U.S.)	0.70				
Harvest Control Rule Values (MT)					
OFL =	11,693				
$ABC_{Cat 1} =$	10,829	10,017	9,241	8,489	7,745
$ABC_{Cat 2} =$	10,680	9,740	8,856	8,010	7,188
HG =	7,871				

Figure 46: Pacific mackerel harvest control rules for fishing year 2023.

Harvest Control Rule Formulas					
OFL = BIOMASS * E_{MSY} * DISTRIBUTION					
ABC _{P-star} = BIOMASS * BUFFER _{P-star} * E_{MSY} * DISTRIBUTION					
HG = (BIOMASS - CUTOFF) * FRACTION * DISTRIBUTION					
Harvest Formula Parameters					
BIOMASS (ages 1+, mt)	60,785				
P-star	0.45	0.40	0.35	0.30	0.25
ABC Buffer _{Cat 1}	0.9133	0.8330	0.7573	0.6851	0.6148
ABC Buffer _{Cat 2}	0.8883	0.7876	0.6954	0.6100	0.5295
E_{MSY} ≡FRACTION	0.30				
CUTOFF (mt)	18,200				
DISTRIBUTION (U.S.)	0.70				
Harvest Control Rule Values (MT)					
OFL =	12,765				
ABC _{Cat 1} =	11,659	10,633	9,667	8,745	7,847
ABC _{Cat 2} =	11,339	10,053	8,877	7,787	6,759
HG =	8,943				

Figure 47: Pacific mackerel harvest control rules for fishing year 2024.

9 References

- Allen, M.J., Wolotira Jr., R.J., Sample, T.M., Noel, S.F., and Iten, C.R. 1990. West coast of North America coastal and oceanic zones strategic assessment: Data Atlas. NOAA Seattle, WA. *Invertebrate and Fish* **145**.
- Barange, M., Hampton, I., and Soule, M. 1996. Empirical determination of in situ target strengths of three loosely aggregated pelagic fish species. *ICES Journal of Marine Science* **53**(2): 225–232.
- Beverton, R.J.H. 1963. Maturation, growth and mortality of clupeid and engraulid stocks in relation to fishing. app. P.-V. Reun. Cons. Perm. Int. Explor. Mer. **154**: 44–67.
- Beverton, R.J.H., and Holt, S.J. 1959. A review of the lifespans and mortality rates of fish in nature, and their relation to growth and other physiological characteristics. *In* CIBA foundation symposium-the lifespan of animals (colloquia on ageing). John Wiley & Sons Ltd, Chichester, UK. pp. 142–180.
- Cannon, R. 1967. How to fish the Pacific Coast. Lane Books, Menlo Park, CA.
- Clemmens, W.A., and Wilby, G.V. 1961. Fishes of the Pacific coast of Canada. *In* Bulletin No. 68. Fisheries Research Board of Canada.
- Collette, B.B., and Nauen, C.E. 1983. Scombrids of the world. *FAO Fisheries Synopsis* **2**(125).
- CONAPESCA. 2020. Anuario Estadístico de Acuicultura y Pesca. Available from <https://www.gob.mx/conapesca/documentos/anuario-estadistico-de-acuicultura-y-pesca>.
- Crocker, R.S. 1933. The California mackerel fishery. California Division of Fish and Game Fish Bulletin **40**: 149.
- Crocker, R.S. 1938. Historical account of the Los Angeles mackerel fishery. California Division of Fish and Game Fish Bulletin **52**: 62.
- Crone, P.R. 2013. Pacific mackerel (*Scomber japonicus*) biomass projection estimate for USA management. Pacific Fishery Management Council, Portland, OR.
- Crone, P.R., Hill, J.P., K. T. Zwolinski, and Kinney, M.J. 2019. Pacific mackerel (*Scomber japonicus*) stock assessment for U.S. management in the 2019-20 and 2020-21 fishing years. Pacific Fishery Management Council, Portland, OR.
- Crone, P.R., and Hill, K.T. 2015. Pacific mackerel (*Scomber japonicus*) stock assessment for USA management in the 2015-16 fishing year. Pacific Fishery Management Council, Portland, OR.
- Crone, P.R., and Hill, K.T. 2017. Pacific mackerel biomass projection estimate for USA management in 2017-18 and 2018-19. Pacific Fishery Management Council, Portland, OR.
- Crone, P.R., and Hill, K.T.H. 2014. Pacific mackerel biomass projection estimate for USA management (2014-15). Pacific Fishery Management Council, Portland, OR.
- Crone, P.R., Hill, K.T., McDaniel, J.D., and Lo, N.C.H. 2009. Pacific mackerel (*Scomber japonicus*) stock assessment for USA Management in the 2009-10 fishing year. Pacific Fishery Management Council, Portland, OR.
- Crone, P.R., Hill, K.T., McDaniel, J.D., and Lynn, K. 2011. Pacific mackerel (*Scomber japonicus*) stock assessment for USA Management in the 2011-12 fishing year. Pacific Fishery Management Council, Portland, OR.
- Demer, D.A., Zwolinski, J.P., Byers, K.A., Cutter, G.R., Renfree, J.S., Sessions, T.S., and

- Macewicz, B.J. 2012. Prediction and confirmation of seasonal migration of pacific sardine (*sardinops sagax*) in the california current ecosystem. *Fishery Bulletin* **110**(1): 52–70.
- Dickerson, T.L., Macewicz, B.J., and Hunter, J.R. 1992. Spawning frequency and batch fecundity of chub mackerel, *Scomber japonicus*, during 1985. *CalCOFI Reports* **33**: 130–140.
- Farrag, M.M., Ismail, R.F., El-Haweet, A.A., Fattah, M.F.A., and Osman, A.G. 2022. Reproductive parameters of the chub mackerel, *scomber japonicus* (houuttuyn, 1782) from the east mediterranean sea, egypt. *Iranian Journal of Ichthyology* **9**(2): 96–110.
- Field, J.C., Francis, R.C., and Strom, A. 2001. Toward a fisheries ecosystem plan for the northern California Current. *CalCOFI Reports* **42**: 74–87.
- Fitch, J.E. 1951. Age composition of the southern California catch of Pacific mackerel 1939-40 through 1950-51. *California Division of Fish and Game Fish Bulletin* **83**: 1–73.
- Fitch, J.E. 1969. *Offshore fishes of California*. California Department of Fish and Game, Sacramento, CA.
- Fitch, J.E., and Lavenberg, R.J. 1971. *Marine food and game fishes of California*. University of California Press, Berkeley, CA.
- Francis, R.C. 2011. Data weighting in statistical fisheries stock assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(6): 1124–1138.
- Frey, H.W. (*Editor*). 1971. *California's living marine resources and their utilization*. California Department of Fish; Game.
- Fritzsche, R.A. 1978. *Development of fishes in the Mid-Atlantic Bight, Vol. 5 Chaetodontidae through Ophidiidae*. U.S. Fish; Wildlife Service.
- Fry Jr, D.H., and Roedel, P.M. 1949. Tagging experiments on the Pacific mackerel (*Pneumatophorus diego*). *California Division of Fish and Game Fish Bulletin* **73**: 64.
- Hamel, O.S., and Cope, J.M. 2022. Development and considerations for application of a longevity-based prior for the natural mortality rate. *Fisheries Research* **256**: 106477.
- Hargreaves, N.B., and Hungar, R.M. 1995. Robertson creek chinook assessment and forecast for 1994 and 1995. Part B: early marine mortality. *PSARC Report* **S95-03**: 55.
- Hart, J.L. 1971. *Pacific fishes of Canada*. Fisheries Research Board of Canada, Bulletin 180.
- Hill, K.T., and Crone, P.R. 2004. Stock assessment of Pacific mackerel (*Scomber japonicus*) with recommendations for the 2004-2005 management season. Pacific Fishery Management Council, Portland, OR.
- Hill, K.T., and Crone, P.R. 2005. Assessment of the Pacific Mackerel (*Scomber japonicus*) stock for U.S. management in the 2005-2006 season. Pacific Fishery Management Council, Portland, OR.
- Hill, K.T., Yaremko, M., and Jacobson, L.D. 1999. Status of the Pacific Mackerel Resource and Fishery in 1998. CDFG Marine Region Administrative Report **99-3**.
- ICES. 2018. Report of the Working Group on Acoustic and Egg Surveys for Sardine and Anchovy in ICES Areas 7, 8 and 9 (WGACEGG). 19-23 November, Nantes, France. ICES CM 2018 **EOSG:03**: 355p.
- Jacobson, L.D. 1993. ADEPT: Software for VPA analysis using Gavaris's procedure. National Marine Fisheries Service, Southwest Fisheries Science Center Admin. Rep. **LJ-93-02**.
- Jacobson, L.D., Konno, E.S., and Pertierra, J.P. 1994. Status of Pacific mackerel and trends in abundance during 1978-1993 (with data tables). National Marine Fisheries Service, SWFSC Admin. Rep. **LJ-94-08**: 33.

- Kamimura, Y., Taga, M., Yukami, R., Watanabe, C., and Furuichi, S. 2021. Intra- and inter-specific density dependence of body condition, growth, and habitat temperature in chub mackerel (*scomber japonicus*). *ICES Journal of Marine Science* **78**(9): 3254–3264.
- Klingbeil, R.A. 1983. Pacific mackerel: a resurgent resource and fishery of the California Current. *CalCOFI Reports* **24**: 35–45.
- Knaggs, E.H., and Parrish, R.H. 1973. Maturation and growth of Pacific mackerel population, *Scomber japonicus houuttuyn*. *California Department of Fish and Game* **59**: 114–120.
- Kramer, D. 1969. Synopsis of the biological data on the Pacific mackerel, *Scomber japonicus* Houuttuyn (northeast Pacific). *FAO UN, Fisheries Synopsis* **40**: 1–18.
- Kuriyama, P.T., Zwolinski, J.P., Teo, S.L.H., and Hill, K.T. 2020. Assessment of the Northern anchovy (*Engraulis mordax*) central subpopulation in 2021 for U.S. management. Pacific Fishery Management Council, Portland, OR.
- Legault, C.M., and Restrepo, V.R. 1998. A flexible forward age-structured assessment program. *ICCAT Working Document SCRS/98/58*: 15.
- MacCall, A.D., Klingbeil, R.A., and Methot, R.D. 1985. Recent increased abundance and potential productivity of Pacific mackerel (*Scomber japonicus*). *CalCOFI Reports* **26**: 119–129.
- Mallicoate, D.L., and Parrish, R.H. 1981. Seasonal growth patterns of California stocks of northern anchovy, *Engraulis mordax*, Pacific mackerel, *Scomber japonicus*, and jack mackerel, *emphTrachurus symmetricus*. *CalCOFI Reports* **22**: 69–81.
- Methot, R.D., and Taylor, I.G. 2011. Adjusting for bias due to variability of estimated recruitments in fishery assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **68**(10): 1744–1760.
- Moser, H.G., Charter, R.L., Smith, P.E., Ambrose, D.A., Charter, S.R., Meyer, C.A., Sandknop, E.M., and Watson, W. 1993. Distributional atlas of fish larvae and eggs in the California Current region: taxa with 1000 or more total larvae, 1951 through 1984. *CalCOFI Atlas* **31**: 233.
- (NMFS), N.M.F.S. 2022. 2021-2022 California Current Ecosystem Status Report. Pacific Fishery Management Council, Portland, OR.
- Palance, D.G., Macewicz, B.J., Stierhoff, K., Demer, D.A., and Zwolinski, J.P. 2019. Length conversions and mass–length relationships of five forage-fish species in the California current ecosystem. *Journal of Fish Biology* **95**(4): 1116–1124.
- Palermino, A., De Felice, A., Canduci, G., Biagiotti, I., Costantini, I., Malavolti, S., and Leonori, I. 2021. First target strength measurement of trachurus mediterraneus and scomber colias in the mediterranean sea. *Fisheries Research* **240**: 105973.
- Parrish, R.H. 1974. Exploitation and recruitment of Pacific Mackerel, *Scomber japonicas*, in the northeastern Pacific. *CalCOFI Reports* **17**: 136–140.
- Parrish, R.H., and MacCall, A.D. 1978. Climatic variation and exploitation in the Pacific mackerel fishery. *California Division of Fish and Game Fish Bulletin* **167**: 110.
- PFMC. 1998. Coastal pelagic species fishery management plan: Amendment 8 to the northern anchovy fishery management plan (Appendix B). Pacific Fishery Management Council, Portland, OR.
- PFMC. 2009. Pacific mackerel STAR panel meeting report. Pacific Fishery Management Council. Portland, OR.
- PFMC. 2011. Acoustic-trawl survey method for coastal pelagic species: Report of methodology

- review panel meeting. Pacific Fishery Management Council. Portland, OR.
- PFMC. 2018. Methodology review panel report: acoustic trawl methodology review for use in coastal pelagic species stock assessments. Pacific Fishery Management Council, Portland, OR.
- PFMC. 2021. Status of the Pacific Coast Coastal Pelagic Species Fishery and Recommended Acceptable Biological Catches: Stock Assessment and Fishery Evaluation 2021 including information through June 2021. Pacific Fishery Management Council. Portland, OR.
- Prager, M.H., and MacCall, A.D. 1988. Revised estimates of historical spawning biomass of the Pacific mackerel, *Scomber japonicus*, Calif. CalCOFI Reports **29**: 91–101.
- Punt, A.E., Smith, K., D. Krusisc Golub, and Robertson, S. 2008. Quantifying age-reading error for use in fisheries stock assessments, with ap eastern scalefish and shark Fishery. Canadian Journal of Fisheries and Aquatic Sciences **65**: 1991–2005.
- Renfree, J.S., Bowlin, N.M., Erisman, B.E., Rojas-González, R.I., Johnson, G.E., Mau, S.A., Murfin, D.W., Schwartzkopf, B.D., Sessions, T.S., Stierhoff, K.L., and others. 2022. Report on the summer 2021 california current ecosystem survey (CCES)(2107RL), 6 july to 15 october 2021, conducted aboard NOAA ship reuben lasker, mexican research vessel dr. Jorge carranza fraser, fishing vessels lisa marie and long beach carnage, and uncrewed surface vehicles. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-669**.
- Roedel, P. 1938. Record-size mackerel in Santa Monica Bay. California Division of Fish and Game Fish Bulletin **24**: 423.
- Roedel, P.M. 1949. Movements of Pacific mackerel as demonstrated by tag recoveries. California Division of Fish and Game Fish Bulletin **35**: 281–291.
- Roedel, P.M. 1952. A review of the Pacific mackerel (*Pneumatophorus diego*) fishery of the Los Angeles region with special reference to the years 1939-1951. California Division of Fish and Game Fish Bulletin **38**: 253–273.
- Schaefer, K.M. 1980. Synopsis of biological data on the chub mackerel, *Scomber japonicus* Houtuyn, 1782, in the Pacific Ocean. Inter-Amer. Trop. Tuna Comm., Spec. Rep. **2**: 395–446.
- Sciences, M.A.E. 1987. Ecology of important fisheries species offshore California. MBC Applied Environmental Sciences.
- Simmonds, E.J., and MacLennan, D.N. 2005. Fisheries acoustics: Theory and practice. Wiley-Blackwell.
- Soutar, A., and Isaacs, J.D. 1974. Abundance of pelagic fish during the 19th and 20th centuries as recorded in anaerobic sediment off the Californias. Fishery Bulletin **72**(2): 257–273.
- Stierhoff, K.L., Renfree, J.S., Rojas-González, R.I., Vallarta-Zárate, J.R., Zwolinski, J.P., and Demer, D.A. 2023. Distribution, biomass, and demographics of coastal pelagic fishes in the california current ecosystem during summer 2021 based on acoustic-trawl sampling. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-676**.
- Stierhoff, K.L., Zwolinski, J.P., and Demer, D.A. 2019b. Distribution, biomass, and demography of coastal pelagic fishes in the California Current Ecosystem during summer 2018 based on acoustic-trawl sampling. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-613**.
- Stierhoff, K.L., Zwolinski, J.P., and Demer, D.A. 2021b. Distribution, biomass, and demogra-

- phy of coastal pelagic fishes in the California Current Ecosystem during summer 2016 based on acoustic-trawl sampling. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-649**.
- Stierhoff, K.L., Zwolinski, J.P., and Demer, D.A. 2020b. Distribution, biomass, and demography of coastal pelagic fishes in the California Current Ecosystem during summer 2018 based on acoustic-trawl sampling. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-626**.
- Stierhoff, K.L., Zwolinski, J.P., and Demer, D.A. 2021. Distribution, biomass, and demography of coastal pelagic fishes in the California Current Ecosystem during summer 2015 based on acoustic-trawl sampling. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-648**.
- Stierhoff, K.L., Zwolinski, J.P., Palance, D.G., Renfree, J.S., Mau, S.A., Murfin, D.W., and Sessions, D., T.S. 2019a. Report on the 2018 California Current Ecosystem (CCE) Survey (1807RL), 26 June to 23 September 2018, conducted aboard NOAA Ship Reuben Lasker. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-609**.
- Stierhoff, K.L., Zwolinski, J.P., Renfree, J.S., and Demer, D.A. 2018c. Report on the collection of data during the summer 2017 California Current Ecosystem Survey (1706RL), 19 June to 11 August 2017, conducted aboard fisheries survey vessel *Reuben Lasker*. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-593**.
- Stierhoff, K.L., Zwolinski, J.P., Renfree, J.S., Johnson, G.E., Mau, S.A., Murfin, D.W., Sessions, T.S., and Demer, D.A. 2020. Report on the Summer 2019 California Current Ecosystem Survey (1907RL), 13 June to 9 September 2019, conducted aboard NOAA Ship Reuben Lasker, Fishing Vessels Lisa Marie and Long Beach Carnage, and three unmanned sailboats. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-625**.
- Stierhoff, K.L., Zwolinski, J.P., Renfree, J.S., Mau, S.A., Murfin, D.W., and Demer, D.A. 2018. Report on the SWFSC's collection of data during the 2015 Joint U.S.-Canada Integrated Acoustic and Trawl Survey of Pacific Hake and Coastal Pelagic Species (SaKe 2015; 1507SH) within the California Current Ecosystem, 15 June to 10 September 2015, conducted aboard Fisheries Survey Vessel Bell M. Shimada. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-603**.
- Stierhoff, K.L., Zwolinski, J.P., Renfree, J.S., Mau, S.A., Murfin, D.W., Palance, D.G., Sessions, T.S., and Demer, D.A. 2018b. Report on the collection of data during the summer 2016 California Current Ecosystem Survey (1606RL), 28 June to 22 September 2016, conducted aboard fisheries survey vessel *Reuben Lasker*. NOAA Technical Memorandum, NMFS **NOAA-TM-NMFS-SWFSC-604**.
- Takasuka, A., Yoneda, M., and Oozeki, Y. 2021. Density-dependent egg production in chub mackerel in the kuroshio current system. *Fisheries Oceanography* **30**(1): 38–50.
- Thorson, J.T., Stewart, I.J., and Punt, A.E. 2012. `nwfscAgeingError`: a user interface in R for the Punt et al. (2008) method for calculating ageing error and imprecision. Available from <https://github.com/pfmc-assessments/nwfscAgeingError>.
- Turner, C.H., and Sexsmith, J.C. 1967. Marine baits of California. California Department of Fish; Game.
- Ware, D.M., and Hargreaves, N.B. 1993. Occurrence of Pacific (chub) mackerel off the B.C. coast in 1993. *PICES Press* **2**: 12–13.
- Wetzel, C.R., and Hamel, O.S. 2023. Applying a probability harvest control rule to account for

- increased uncertainty in setting precautionary harvest limits from past stock assessments. *Fisheries Research* **262**: 106658.
- Wood, S. 2017. *Generalized Additive Models: An Introduction with R*, 2nd edition. Chapman; Hall/CRC.
- Xu, H., Thorson, J.T., Methot, R.D., and Taylor, I.G. 2019. A new semi-parametric method for autocorrelated age-and time-varying selectivity in age-structured assessment models. *Canadian Journal of Fisheries and Aquatic Sciences* **76**(2): 268–285.
- Yoneda, M., Kitano, H., Nyuji, M., Nakamura, M., Takahashi, M., Kawabata, A., Matsuyama, M., Shimizu, A., Tsuzaki, T., Togashi, H., and others. 2022. Maternal spawning experience and thermal effects on offspring viability of chub mackerel and their influence on reproductive success. *Frontiers in Marine Science* **9**: 2701.
- Young, P.H. 1969. The California partyboat fishery 1947-1967. California Department of Fish and Game, *Fishery Bulletin* **145**: 91.
- Zwolinski, J.P., Demer, D.A., Cutter Jr, G.R., Stierhoff, K., and Macewicz, B.J. 2014. Building on fisheries acoustics for marine ecosystem surveys. *Oceanography* **27**(4): 68–79. JSTOR.
- Zwolinski, J.P., Stierhoff, K.L., and Demer, D.A. 2019. Distribution, biomass, and demography of coastal pelagic fishes in the California Current Ecosystem during summer 2017 based on acoustic-trawl sampling. NOAA Technical Memorandum **NMFS-SWFSC-610**.

10 Appendix A: Age and Maturity Assessment of Pacific mackerel (*Scomber japonicus*)

Owyn E. Snodgrass¹, Emmanis Dorval^{1,2}, Kelsey C. James¹, Brittany D. Schwartzkopf¹, Jonathan M. Walker^{1,3}, Dianna L. Porzio⁴, Beverly J. Macewicz¹, and Brad E. Erisman¹

¹ NOAA Fisheries, SWFSC Fisheries Resources Division, 8901 La Jolla Shores Dr., La Jolla, CA 92037

² Lynker under contract with Southwest Fisheries Science Center, 202 Church St., SE / #536, Leesburg, VA 20175

³ University of California Santa Cruz, The Cooperative Institute for Marine, Earth, and Atmospheric Systems (CIMEAS) under partnership with NOAA Fisheries, 1156 High St., Santa Cruz, CA 95064

⁴ California Department of Fish and Wildlife, Marine Region, 4665 Lampson Ave., Suite C, Los Alamitos, CA 90720

Summary

The goal of this report is to provide updated information on age and maturity of Pacific Mackerel (*Scomber japonicus*) for consideration in the 2023 benchmark stock assessment. In section 1, we provide an ageing dataset and estimates of ageing errors for Pacific Mackerel otoliths collected from 2012 to 2022 during fishery-independent surveys. In section 2, we provide an updated estimate of length and age at sexual maturity for Pacific Mackerel based on samples collected from 2010 to 2021 during fishery-independent surveys.

1. Ageing of Pacific Mackerel

Background

Historically, biological samples of Pacific Mackerel were collected solely from commercial fishery landings by the California Department of Fish and Wildlife (CDFW). Consequently, all age data incorporated into assessments were fishery-dependent. The Southwest Fisheries Science Center (SWFSC) began archiving Pacific Mackerel otoliths in 2007 to provide fishery-independent biological samples for consideration in assessments, although this species was not a primary target species. To provide a more robust sample archive to generate length and age compositions for acoustic biomass estimates, Pacific Mackerel became a primary target species in 2012 and were sampled following the same protocol as Pacific Sardine (*Sardinops sagax*) and Northern Anchovy (*Engraulis mordax*) (Dorval et al. 2022).

SWFSC staff produced Pacific Mackerel ages from whole, unpolished otoliths collected during SWFSC surveys. The procedure described by Fitch (1951) was used to estimate ages with the assumption that observable growth increments were deposited during the progression of seasons. An annulus was assigned when “the interface between an inner translucent growth increment and the successive outer opaque growth increment” (Fitch 1951, Yaremko 1996) was observed. The

application of this method was to immerse the otolith in distilled water, view using a stereo microscope, and count the number of annuli observed on the distal side of the otolith in less than three minutes. Although Pacific Mackerel has an extended spawning season, a July 1 birthdate was assigned for all individual Pacific Mackerel collected in U.S. waters, albeit an unknown number of these individual fish could have been born prior to or following this date. After annuli were counted without knowledge of size, sex, or capture date, the birthdate, capture date, and analysis of the most distal pair of growth increments were used to assign final ages by readers (see Yaremko 1996).

Sample Collection

Pacific Mackerel otoliths were collected during SWFSC summer acoustic trawl method (ATM) surveys conducted from July through October (Dorval et al. 2022). Collections spanned from the Canadian-US border to the US-Mexican border (2012-2022) (Figure 1). Pacific Mackerel were randomly subsampled ($n = 50$) from the larger catch and measured for fork length (FL; mm) and weighed (g). If fewer than 50 were caught, all Pacific Mackerel were measured and weighed. Sagittal otoliths were then extracted from up to 25 Pacific Mackerel and stored dry.




Figure 1. Catch locations for Pacific Mackerel (*Scomber japonicus*) during SWFSC spring and summer trawl surveys (2010-2022).

Age-reading

Whole otoliths were immersed in distilled water with the distal side facing up and then read from the posterior region, using a stereo microscope at 25 X magnification. Three SWFSC age

readers, identified as readers 15, 17, and 18, participated in the age determination process, using the conventional technique of otolith age-reading described in Yaremko (1996). All agers used in this study were certified agers. Further, the SWFSC ATM survey age dataset is consistent with fishery ages produced by CDFW for the 2019 and 2023 stock assessments, as the best CDFW age reader (reader 2) was involved in the training process of the three SWFSC readers above. Furthermore, a random set of survey otolith samples (n=317, summer 2012-2017) was assigned to readers 2, 15, 17, and 18 for ageing in order to evaluate bias and precision among these four readers.

A total of 1,762 ages from 2012 to 2022 were produced by readers 17 and 18. From each summer survey, otolith samples were randomly selected by haul and by length bin (50 mm FL), and approximately 50% of the selected samples were randomly allocated to each of these two readers. This selection scheme maintained the spatial and temporal integrity of the trawl sampling and the distribution of length-at-age in space and time. Due to time constraints, a subset of total otoliths collected were aged from 2013 to 2019 that accounted for length bin, year, and geographic location. Each individual fish was assigned a final age based on the capture date and an assumed July 1 birthdate (see Yaremko 1996) and the analysis of the most distal pair of growth increments.

Further, 36% of the total number of otolith samples aged by readers 17 and 18 were randomly selected and double-read by these two readers and reader 15 to produce a consensus age reading vector identified as reader CA. The CA ageing vector included ages that all three readers agreed upon and additional ages determined from simultaneous onsite readings under the same stereo microscope until they reached 100% agreement. As such, the CA ageing vector was assumed to be the best ages, and accordingly was considered unbiased in the computation of ageing errors. This method was previously reviewed and approved by Pacific Sardine STAR panels in 2011 for ages produced by the Department of Fisheries and Oceans (DFO) laboratory (Hill et al. 2011, Dorval et al. 2013) and in 2020 for ages produced by SWFSC (Kuriyama et al. 2020).

The computation of age-reading errors was based on the method described by Punt et al. (2008), using the `nwfscAgeingError` R package (Thorson et al. 2012). We computed ageing error matrices based on otoliths that were aged by reader CA, 17, and 18 while assuming that: (1) ageing bias depends on reader and the true age of a fish; (2) the age-reading error standard deviation depends on reader and the true age; and (3) age-reading error is normally distributed around the expected age (see Punt et al. 2008). For the purpose of this report, we were mostly interested in estimating the *SDs-at-age* for age data collected during the 2012-2022 trawl surveys, following similar methods used in the past for Pacific Sardine and Pacific Mackerel assessments (Hill et al. 2011; Dorval et al. 2013; Crone et al. 2019; Kuriyama et al. 2020). We defined various model scenarios, comparing models that assumed equal or unequal *SDs* among readers. As in previous assessments, Model C (Dorval et al. 2013) was selected as the best model, using Akaike Information Criterion with a correction for finite sample sizes. This model assumed that all three readers (CA, 17, and 18) were unbiased and had equal *SDs*. One dataset set, including age data from 2012 to 2022, was used to compute ageing errors for the trawl

surveys. The functional form of random ageing-error precisions was assumed to follow a curvilinear *SD* and a curvilinear *CV* based on a three-parameters, Hollings-form relationship of *SD* or *CV* with true age (see Punt et al. 2008; Thorson et al. 2012, Dorval et al. 2013). Further, the maximum *SD* allowed in model runs was 40.

Results and Discussion

The length distribution of Pacific Mackerel subsampled and measured during summer trawl surveys from 2012 to 2022 ranged from 53 mm FL to 402 mm FL (Figure 2a). A total of 1,762 fish were aged, with ages ranging from 0 to 7 years (Figure 2b). Aged samples were comprised mostly of young fish, with individuals aged at 0, 1, 2, and 3 years representing 46%, 29%, 16%, and 6% of the total number of otoliths aged, respectively. Older fish (4-7 years in age) made up only 2.3% of the samples aged, and thus these age classes might not have been well represented in the summer trawl surveys. There were large overlaps in length distributions among age classes (Figure 3). Although ages through 2022 were produced, the final assessment model only used ages through 2021 due to uncertainties in the 2022 biomass estimate from the SWFSC ATM survey (PFMC 2023).

Age-Reading Errors

Age-reading errors for the survey data were computed using 643 otoliths collected from 2012 to 2022. Ages were estimated with a high level of precision. Ageing agreement for these 643 otoliths between readers 17 and 18 was 100% from age 0 to age 2, 94% at age 3, 75% at age 4, and 70% at age 5 (Figure 4). Only 2 fish were aged greater than 5 years, but these readers disagreed on the age of these fish. In the consensus ageing vector, one of these fish was assigned an age 5 and the other an age 6. As a result, *SDs-at-age* estimated from Model C were very low, varying from 0.001 to 0.319 (Table 1).

Pacific Mackerel of ages 4 years and older (Figure 4) were the only ages where readers agreed 75% of the time or less. This age group is more frequent in the Pacific Northwest and/or in offshore waters that are not well covered by current trawl surveys. Only 26 Pacific Mackerel out of 1,762 were in the 4⁺ age group. Older age classes generally have lower agreement. Interpreting increments at the edge of older fish otoliths is more challenging, because annuli are much closer together and it is more difficult to differentiate a check from an annulus (Yaremko 1996).

Bias plots showed a high level of accuracy among readers 2, 15, 17, and 18 (Figure 5), indicating that on average all of these readers can produce the same age compositions given a set of survey or fishery otolith samples. The CDFW ageing error vector used in this stock assessment had higher standard deviations from age 3 to 7 than survey-based estimates (Figure 6). However, these differences did not result from the application of ageing criteria between the two laboratories; rather they most likely reflect differences in the process of sample selection and in the level of agreement targeted within each lab, when conducting multiple readings for

estimating ageing errors. Primary differences between CDFW and SWFSC are summarized in Table 2.

A current drawback is that no age validation has been published for Pacific Mackerel in the eastern North Pacific. The absence of validation of the periodicity of increment formation in each and every age group can lead to systematic bias in age determination (Campana 2001). Shiraishi et al. (2008) confirmed annual periodicity of annuli in Pacific Mackerel from southwest Japan through captive growth of known-age fish up to 2 years old and edge analysis in wild Pacific Mackerel up to 6 years old. The SWFSC conducted a captive growth experiment of Pacific Mackerel and preliminary results suggest annual periodicity of annuli in fish up to approximately 2 years old (K.C. James et al. unpublished data). While this research is not for every age class, and there still is a possibility of bias from unvalidated ages, it lends confidence to the accuracy of ages provided to the stock assessment.

While all otolith samples were collected during SWFSC ATM surveys, it is important to note that the entire length range of Pacific Mackerel was not sampled for this study. The ATM survey is designed to produce abundance estimates for multiple coastal pelagic species based on their acoustic signatures. Additionally, trawl net avoidance and rates of capture likely varies by species and fish length.




Figure 2. a) Length and b) age distribution of aged Pacific Mackerel (*Scomber japonicus*) collected from summer SWFSC acoustic trawl surveys (2012-2022).




Figure 3. Age-at-length for Pacific Mackerel (*Scomber japonicus*) collected from summer SWFSC acoustic trawl surveys (2012-2022).

Table 1. Coefficient of variation (CV) and standard deviation (SD) at age estimated for Pacific Mackerel (*Scomber japonicus*) collected from summer SWFSC acoustic trawl surveys (2012-2022). All estimates were calculated using the latest version of the nwfscAgeingError R package (Thorson et al. 2012) based on the assumptions that, within the SWFSC laboratory, there was no bias in ageing among readers, and readers had similar SD.

Survey	Collection Year	Data set ID	Sample size	Number of readers	Agemat model		
					Age	CV	SD
SFWSC Trawl Survey	2012-2022	1	643	3	0	0.001	0.001
					1	0.001	0.001
					2	0.074	0.148
					3	0.076	0.229
					4	0.068	0.273
					5	0.060	0.298
					6	0.052	0.311
7	0.046	0.319					




Figure 4. Age bias plots from the Agemat model for readers CA, 17, and 18 for Pacific Mackerel (*Scomber japonicus*) collected from summer SWFSC acoustic trawl surveys (2012-2022).




Figure 5. Age bias plots (mean age \pm SE and bubble plots) of readers 2, 15, 17,18 for Pacific Mackerel (*Scomber japonicus*) collected from summer SWFSC acoustic trawl surveys (2012-2022).




Figure 6. Survey and fishery ageing error vectors used in the final assessment model (CDFW and SWFC vectors) and sensitivity tests (CDFW-SWFSC vector). Blue dotted line shows ageing errors estimated from 2012-2022 summer survey samples; orange dotted line shows ageing errors estimated from summer 2012-2017 survey samples; and black dotted line shows ageing error estimated from 2008-2017 fishery samples. Numbers in parentheses indicate SWFSC and CDFW readers that produced age readings for estimating ageing errors.

Table 2. Comparison of CDFW and SWFSC methodologies for computing ageing errors.

Component	Fishery ageing data	Survey ageing data
Ageing criteria	Same as acoustic trawl survey	Same as fishery port sampling
Sample selection for Double readings	From every third - fourth port sample	From every acoustic trawl sample
Target of ageing precision	75% agreement among readers	100% agreement among readers
Data used to compute ageing errors	Double readings before 75% agreement	Double readings at 100% agreement
Rationale	Not possible to adjust or drop assessment ages after double readings	Possible to adjust or drop assessment ages after double readings
	Avoid bias in CV and SD because assessment data cannot be adjusted after double readings	Minimize CV and SD as assessment data can be adjusted after double readings

2. Length and Age at Maturity of Pacific Mackerel

Background

The Pacific Mackerel is a multiple batch spawner with indeterminate fecundity, asynchronous oocyte development, and a relatively high spawning frequency (Knaggs and Parrish 1973; Peña et al. 1986; Asano and Tanaka 1989; Dickerson et al. 1992). In the northeast Pacific, spawning of Pacific Mackerel typically occurs from Point Conception to Cabo San Lucas from 3 to over 300 km offshore (Moser et al. 1993), although small juveniles have been reported off Oregon and Washington in recent years (Stierhoff et al. 2019). Pacific Mackerel have a protracted spawning season throughout their range, with peak spawning off California and central Baja California, Mexico, occurring during the spring through summer months and some spawning occurring during all months of the year (Ahlstrom 1959; Kramer 1969; Knaggs and Parrish 1973; Schaefer 1980; Gluyas-Millán 1994). Similar to other broadcast-spawning marine fishes, both spawning frequency and spawning season duration are believed to increase with female size and age (Knaggs and Parrish 1973; Dickerson et al. 1992).

Recent stock assessments for Pacific Mackerel used maturity schedules from Dickerson et al. (1992), in which the fraction of mature females was estimated by fitting a logistic regression model to maturity data (Crone and Hill 2015; Crone et al. 2019). A more recent study was conducted from 2009 to 2012 for purposes of re-evaluating maturity-at-age for Pacific Mackerel, which used simple logistic regression to estimate 50% maturity at 27 cm FL and 2.2 years of age (Crone and Hill 2015). The results of the more recent study were similar to those based on Dickerson et al. (1992), and consequently, the maturity schedules used in past assessments were again applied in both 2015 and 2020 (Crone and Hill 2015; Crone et al. 2019). Estimated maturity schedules for Pacific Mackerel off California are similar to those reported in Mexico. For example, Gluyas-Millán (1994) concluded that 50% of female Pacific Mackerel off Vizcaino Bay, Mexico, are mature by 293 mm standard length (SL).

Material and Methods

Samples of ovarian tissues were collected from female Pacific Mackerel during SWFSC spring and summer surveys conducted from 2010 through 2021 to generate updated estimates of length- and age-at-maturity. Males were not included in this study, because previous studies have concluded there to be no notable differences in growth, maturity, or mortality rate in Pacific Mackerel by sex (see Crone et al. 2019). Consequently, combined sex models have been used in all stock assessments used to advise management in U.S. Pacific waters (Crone et al. 2019). Each gonad sample was placed in a tissue-tek cassette and preserved in 10% neutral buffered formalin in preparation for histological processing and examination. Samples were later embedded in paraffin, sectioned at 6 μ m, mounted on slides, stained with Mayer's haematoxylin-eosin, and observed under a compound microscope (Humason 1972). Past studies on reproductive development in Pacific Mackerel emphasized the importance of using histological criteria for maturity assessments, as all stages of ovarian development cannot be discerned with the unaided eye (Asano and Tanaka 1989; Dickerson et al. 1992).

Standardized terminology for describing reproductive development in marine fishes (Brown-Peterson et al. 2011) was used to classify each sampled female Pacific Mackerel as either immature (never spawned) or mature (previously spawned or first spawning) (Figure 7). Females with ovaries containing no oocytes undergoing vitellogenesis but numerous oocytes in the cortical alveolar stage of development were classified as mature, because fish sampled at this phase of development usually spawn at some point during the season (Murua and Saborido-Rey 2003; Wright 2007; Lowerre-Barbieri et al. 2011a,b). Additional histological features used to distinguish between immature females and mature, regenerating females included the thickness of the ovarian wall, the presence of muscle bundles or atretic follicles, and the level of organization within the lamellar structure (Lowerre-Barbieri et al. 2011a,b).

Following common practice, the length and age at sexual maturity for Pacific Mackerel was estimated using an analytical method based on logistic, non-linear regression (Hunter et al. 1992; Macewicz et al. 1996; Roa et al. 1999; Lo et al. 2005; Basilone et al. 2006). Specifically, we followed the methods described by McBride (2016), which used a binomial model in R (R Core Team 2022) to estimate the length and age at 25, 50, and 95% maturity and the uncertainty around the predicted relationship between length or age and percent maturity (Formula: Maturity \sim FL). Maturity data were pooled across all survey years to generate sample sizes across all length and age classes that were sufficient to produce a realistic ogive estimate without sample distribution bias. The use of a pooled maturity data set was consistent with recent stock assessments for Pacific Mackerel, in which age-length keys used to estimate age compositions were comprised of pooled age and length data (see Crone and Hill 2015 and Crone et al. 2019).




Figure 7. Histological sections of gonads of female Pacific Mackerel (*Scomber japonicus*) collected from SWFSC spring and summer trawl surveys (2010-2021): (a) Immature female with only previtellogenic oocytes; (b) Mature, developing female with numerous oocytes in early cortical alveoli stage; (c) Mature, spawning capable female with numerous vitellogenic oocytes; (d) Mature, actively spawning female with hydrated oocytes.

Results and Discussion

A total of 911 gonad samples of female Pacific Mackerel were examined histologically, classified as either immature (juvenile) or mature (adult), and then used to generate an estimate of length at maturity. Age data were available for 494 of these sampled females to generate an estimate of age at maturity. Females ranged in length from 174 to 402 mm FL and in age from 0 to 7 years (Figure 8a,b). Immature females ranged in length from 174 to 329 mm FL and in age from 0 to 2 years. Mature females were 207-402 mm FL and 0-7 years of age.

The estimated length at maturity (L_{50}) for all sampled females ($n = 911$) was 274 ± 1.26 mm FL with all females (L_{95}) larger than 309 ± 2.60 mm FL predicted to be mature (Figure 9a; Table 3). The estimated age at maturity (A_{50}) for all sampled females ($n = 494$) was 1.01 ± 0.06 years with all females older than 2.52 ± 0.15 years predicted to be mature (Figure 9b; Table 4).




Figure 8. Histograms showing a) length ($n = 911$) and b) age distribution ($n = 494$) by maturity state for female Pacific Mackerel (*Scomber japonicus*) collected from SWFSC spring and summer trawl surveys (2010-2021) and analyzed histologically for reproductive condition.

The estimated length at maturity reported here (274 mm SL) is nearly identical to the value used in recent stock assessments for Pacific Mackerel, whereas the estimated age at maturity (1.01

years) is lower than the previous estimate (2.2 years; see Crone et al. 2015, Crone et al. 2019). We attribute the difference in estimated age at maturity to increased precision and accuracy in ageing methods combined with an increase in availability of smaller age classes to the trawl survey. Collectively, the results of this and past studies indicate that maturity schedules in Pacific Mackerel off the U.S. Pacific coast have remained constant over the past several decades.




Figure 9. a) Length-based ($n = 911$) and b) age-based ($n = 494$) maturity ogives of female Pacific Mackerel (*Scomber japonicus*) based on samples collected from SWFSC spring and summer trawl surveys (2010-2021). Data are shown as jittered tick marks along the lower (immature fish) and upper (mature fish) x-axis. The solid line represents the predicted curve, and the dashed lines depict the 95% confidence intervals.

Table 3. Mean predicted probability of being mature and standard deviation for Pacific Mackerel (*Scomber japonicus*) in 50 mm fork-length bins from the length-based ogive for samples collected from SWFSC spring and summer trawl surveys (2010-2021).

Fork-length bin	Mean predicted probability	Standard deviation
151-200 mm FL	0.00083	5.34e-04
201-250 mm FL	0.03	3.22e-02
251-300 mm FL	0.52	2.54e-01
301-350 mm FL	0.97	2.52e-02
351-400 mm FL	0.99	4.11e-04
401-450 mm FL	0.99	6.32e-06

Table 4. Predicted probability of being mature for each age with 95% confidence intervals for Pacific Mackerel (*Scomber japonicus*) from the age-based ogive for samples collected from SWFSC spring and summer trawl surveys (2010-2021).

Age (years)	Predicted probability	95% confidence interval
0	0.12	0.08-0.17
1	0.49	0.43-0.55
2	0.87	0.82-0.91
3	0.98	0.95-0.99
4	0.99	0.99-0.99
5	0.99	0.99-0.99

Acknowledgements

We thank all SWFSC staff and volunteers who collected Pacific Mackerel from the SWFSC ATM survey and previous staff from the Life History Program at SWFSC who processed and prepared the otoliths and gonads to be read and analyzed.

References

- Ahlstrom, E.H. 1959. Vertical distribution of pelagic fish eggs and larvae off California and Baja California. *Fishery Bulletin* 60:107-146.
- Asano, K., and Tanaka, S. 1989. Ovarian maturation and spawning of the Japanese Common Mackerel *Scomber japonicus*. *Nippon Suisan Gakkaishi* 55:1715–1726.
- Basilone, G., Guisande, C., Patti, B., Mazzola, S., Cuttitta, A., Bonanno, A., Vergara, A.R., and Maneiro, I. 2006. Effect of habitat conditions on reproduction of the European anchovy (*Engraulis encrasicolus*) in the Strait of Sicily. *Fisheries Oceanography* 15:271–280.
- Brown-Peterson, N.J., Wyanski, D.M., Saborido-Rey, F., Macewicz, B.J., and Lowerre-Barbieri, S.K. 2011. A standardized terminology for describing reproductive development in fishes. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 3:52–70.

Campana, S. E. 2001. Accuracy, precision, and quality control in age determination, including a review and abuse of age validation methods. *Journal of Fish Biology* 59:197–242.

Crone, P.R., and Hill, K.T. 2015. Pacific Mackerel (*Scomber japonicus*) stock assessment for USA management in the 2015-16 fishing year. Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 101, Portland, OR, 97220. 131 p.

Crone, P.R., Hill, K.T., Zwolinski, J.P., and Kinney, M.J. 2019. Pacific Mackerel (*Scomber japonicus*) stock assessment for U.S. management in the 2019-20 and 2020-21 fishing years. Pacific Fishery Management Council, Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 101, Portland, OR 97220. 112 p.

Dickerson, T.L., Macewicz, B.J. and Hunter, J.R. 1992. Spawning frequency and batch fecundity of chub mackerel, *Scomber japonicus*, during 1985. *California Cooperative Oceanic Fisheries Investigative Reports* 33:130–140.

Dorval, E., J.D. McDaniel, D.L. Porzio, R. Felix-Uraga, V. Hodes, S. Rosenfield. 2013. Computing and selecting ageing errors to include in stock assessment models of Pacific sardine (*Sardinops sagax*). *California Cooperative Oceanic Fisheries Investigative Reports* 54:1–13.

Dorval, E., Porzio, D., Schwartzkopf, B.D, James, K.C. Vasquez, L., and Erisman, B. 2022. Sampling methodology for estimating life history parameters of coastal pelagic species along the U.S. Pacific Coast. U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-660. 46 p.

Fitch, J. E. 1951. Age composition of the southern California catch of Pacific Mackerel 1939–40 through 1950–51. *California Department of Fish and Game* 83:1–73.

Gluyas-Millán, M.G. 1994. Reproduction, age and length of first maturity of Pacific Mackerel *Scomber japonicus* (Houttuyn, 1872) in Vizcaíno Bay. *Ciencias Marinas* 20:409–419.

Humason, G.L. 1972. *Animal Tissue Techniques*. 3rd Edition, W.H. Freeman and Company, San Francisco.

Hunter J.R., Macewicz, B.J., Lo, N.C.H., and Kimbrell, C.A. 1992. Fecundity, spawning, and maturity of female Dover sole, *Microstomus pacificus*, with an evaluation of assumptions and precision. *Fishery Bulletin* 90:101–128.

Hill, K. T., Crone, P.R., Lo, N.C.H., Macewicz, B.J., Dorval, E., McDaniel, J.D., and Gu, Y. 2011. Assessment of the Pacific sardine resource in 2011 for U.S. management in 2012. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-SWFSC-487. 265 pp.

Knaggs, E.H., and Parrish, R.H. 1973. Maturation and growth of Pacific Mackerel, *Scomber japonicus* Houttuyn. *California Department of Fish and Game* 59:114-120.

Kramer, D. 1969. Synopsis of the biological data on the Pacific Mackerel, *Scomber japonicus* Houttuyn (northeast Pacific). FAO (Food and Agri. Org.), U.N., Fish. Synopsis 40:1–18.

Kuriyama, P.T., Zwolinski, J.P., Hill, K.T., and Crone, P.R. 2020. Assessment of the Pacific Sardine resource in 2020 for U.S. management in 2020-2021. U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-625. 189 pp.

Lo, N.C.H., Macewicz, B.J., and Griffith, D.A. 2005. Spawning biomass of Pacific sardine (*Sardinops sagax*), from 1994-2004, off California. California Cooperative Oceanic Fisheries Investigative Reports 46:93–112.

Lowerre-Barbieri, S.K., Brown-Peterson, N.J., Murua, H., Tomkiewicz, J., Wyanski, D., and Saborido-Rey, F. 2011a. Emerging issues and methodological advances in fisheries reproductive biology. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 3:32–51.

Lowerre-Barbieri, S.K., Ganas, K., Saborido-Rey, F., Murua, H., and Hunter, J.R. 2011b. Reproductive timing in marine fishes: variability, temporal scales, and methods. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 3:71–91.

Macewicz, B.J., Castro-Gonzalez, J.J., Coterio-Altamrano, C.E., and Hunter, J.R. 1996. Adult reproductive parameters of Pacific sardine (*Sardinops sagax*) during 1994. California Cooperative Oceanic Fisheries Investigative Reports 37:140–151.

McBride R. 2016. Maturity Schedules: matching data with models. Available at: <https://rpubs.com/fishmcbride/222042>. (September 2022).

Moser, H.G., Charter, R.L., Smith, P.E., Ambrose, D.A., Charter, S.R., Meyer, C.A., Sandknop, E.M., and Watson, W. 1993. Distributional atlas of fish larvae and eggs in the California Current region: taxa with 1000 or more total larvae, 1951 through 1984. *CalCOFI Atlas* 31. 233 p.

Murua, H., and Saborido-Rey, F. 2003. Female reproductive strategies of marine fish species of the North Atlantic. *Journal of Northwest Atlantic Fishery Science* 33:23–31.

Peña, N., Alheit, J., and Nakama, M.E., 1986. Fecundidad parcial de Caballa del Perú (*Scomber japonicus peruanus*). *Boletín IMARPE* 10:93-104.

Pacific Fisheries Management Council (PFMC). 2023. Coastal Pelagic Species Subcommittee of the Scientific and Statistical Committee Review of aspects related to assessment of Pacific Sardine and other CPS Species. Available at: <https://www.pcouncil.org/documents/2023/03/coastal-pelagic-species-subcommittee-of-the-scientific-and-statistical-committee-review-of-aspects-related-to-assessment-of-pacific-sardine-and-other-cps-species-march-20-21-2023-meeting.pdf/>

Punt, A. E., Smith, D.C., Krusisc Golub, K., and Robertson, S. 2008. Quantifying age-reading error for use in fisheries stock assessments, with application to species in Australia's southern and eastern scalefish and shark Fishery. *Canadian Journal Fisheries and Aquatic Sciences* 65: 1991–2005.

R Core Team. 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Roa, R., Ernst, B., and Tapia, F. 1999. Estimation of size at sexual maturity: an evaluation of analytical and resampling procedures. *Fishery Bulletin* 97:570–580.

Schaefer, K.M. 1980. Synopsis of biological data on the chub mackerel, *Scomber japonicus* Houtuyn, 1782, in the Pacific Ocean. Inter-Amer. Trop. Tuna Comm., Spec. Rep. 2:395-446.

Shiraishi, T., Okamoto, K., Yoneda, M., Sakai, T., Ohshimo, S., Onoe, S., Yamaguichi, A., and Matsuyama, M. 2008. Age validation, growth and annual reproductive cycle of chub mackerel *Scomber japonicus* off the waters of northern Kyushu and in the East China Sea. *Fisheries Science* 74: 947-954.

Stierhoff, K.L., Zwolinski, J.P., and Demer, D.A. 2019. Distribution, biomass, and demography of coastal pelagic fishes in the California Current Ecosystem during summer 2018 based on acoustic-trawl sampling. U.S. Dep. Commer., NOAA Tech. Memo., NMFS-SWFSC-613. 83 pp.

Thorson, J.T., Stewart, J.J., and Punt, A.E. 2012. nwfscAgeingError: a user interface in R for the Punt et al. (2008) method for calculating ageing error and imprecision. Available from: <http://github.com/nwfsc-assess/nwfscAgeingError>.

Wright, P.J. 2007. Understanding the maturation process for field investigations of fisheries-induced evolution. *Marine Ecology Progress Series* 335:279–283.

Yaremko, M. L. 1996. Age determination in Pacific sardine, *Sardinops sagax*. U.S. Dep. Commer., NOAA Tech. Memo., NMFS SWFSC-223. 33 pp.

11 Appendix B: Supplementary details following STAR Panel

11.1 Justification for time-varying catchability (Q)

The AT survey index of abundance data have relatively large fluctuations that might be explained by a number of time-varying processes. Discussions at the STAR panel focused on accounting for this variability with two model configurations: time-varying Q values and time-varying natural mortality M . The base model accounts for variability in the AT survey data by estimating annual deviations on Q for 2008-2015 and an estimated Q for years 2016-2021, with a prior based on the proportion of biomass observed in the 2021 AT survey in US and Mexican waters. This decision was the result of a number of sensitivity runs.

Values of Q might be changing due to variability in Pacific mackerel distributions and movements. There does not seem to be any apparent spatial pattern in Pacific mackerel distributions from the AT survey. In 2008, the biomass was observed off northern CA and central OR (Demer et al. 2012). In 2012, biomass was observed continuously from Vancouver Island to roughly San Francisco, CA (Zwolinski et al. 2014). Pacific mackerel were found only off Oregon in 2015 (Stierhoff et al. 2021), and distributed continuously from Vancouver Island to the US-Mexico border in 2016 (Stierhoff et al. 2021b). In 2017, mackerel were observed throughout WA, OR, and northern CA [Zwolinski et al. (2019); note this survey ended in Morro Bay, CA]. In 2018, mackerel were found throughout the US west coast, although there were no observations between Mendocino and Monterey, CA (Stierhoff et al. 2019b). A similar pattern was seen in 2019 (Stierhoff et al. 2020b) and 2021 (Stierhoff et al. 2023). The 2021 survey continued into Mexican waters, and mackerel were observed from Ensenada to Punta Abreojos (Stierhoff et al. 2023). Maps of the mackerel distributions are shown in Figures B-1 and B-2. These distributions might seem to be patchy because the stock seems to be at a relatively low biomass state.

The AT surveys had year-to-year changes in survey protocols from 2008-2015. Examples include longer distances between trawl transect lines and changes in target species for survey design. Protocols were standardized from 2016-2021, despite the variability in survey footprint.




Figure B-1: Maps of Pacific mackerel abundances from the cited AT survey reports. Maps for summers 2008, 2012, 2015, and 2016 are shown from left to right




Figure B-2: Maps of Pacific mackerel abundances from the cited AT survey reports. Maps for summers 2017, 2018, 2019, and 2021 are shown left to right

Estimating a time-invariant Q resulted in model fits that miss three of the ten index values. The STAT evaluated a number of alternative models that explored configurations that might improve the fits of the indices.

The STAT considered models that had different time blocks on Q . One sensitivity had two blocks, one for 2008-2015 and another for 2016-2021. These blocks were informed by the degree of consistency in survey protocols with input from Juan Zwolinski. The configuration with two blocks did not improve fits to the index data. Another model run at the other end of the spectrum estimated annual deviations in Q . This run led to higher M values but Q values that were lower than those estimated in other models. The period from 2016-2021 has comparatively similar biomass values as might be expected given the consistency in survey protocols for this period. Allowing deviations in this period may not be justified given input from the AT survey group and the biomass estimates.

One consideration was dropping 2013-2015 AT index data, which were about 10,000 mt and less than the estimates before and after. In these sensitivities, the model fit all seven index values but time-invariant M increased to roughly 1, which was higher than estimates from other runs. Additionally, the decision to drop three of the ten AT survey indices of abundance did not seem to be appropriate given the already short time period of the base model.

Time-varying M was evaluated but not adopted by the STAT. One sensitivity heavily weighted the AT index of abundance and estimated annual deviations around M (and estimated a single time-invariant Q). If all the variability in the AT index was modeled with time-varying M , estimates of M ranged from about 0.5 to 3.5. The high values did not seem biologically plausible. Another run did not upweight the index data and estimated time-varying M . This run still missed the same three index values and increased the scale of model biomasses.

Ultimately the STAT settled on a base model that estimates annual deviations for Q values in 2008-2015, rather than dropping data values, and one Q block for 2016-2021 when survey protocols were fixed. This Q block was centered with a prior based on the biomass estimates in US and Mexican waters. This approach allows flexibility for the early years of the survey while utilizing available data with some structure in 2016-2021.

11.2 Historical catch time series

Long-term landings dating back to 1926 are presented in Table B-1. Note, that the landings time series in the base model includes bycatch in the At-Sea Pacific Whiting fishery, but these values are not included in Table B-1.

Table B-1: Landings (mt) of Pacific mackerel by region and fishing year (1926-2022). Mexican landings were from Magdalena Bay, BCS (MAG) and Ensenada, BC (ENS). US landings are from California (CA), Oregon (OR), Washington (WA). Additionally, California recreational landings are included (CA-REC).

Model Year	MAG	ENS	MEX-TOT	CA	OR	WA	CA-REC	USA-TOT	TOT
1926	—	—	—	1,630	—	—	—	1,630	1,630
1927	—	—	—	2,928	—	—	—	2,928	2,928
1928	—	—	—	17,874	—	—	—	17,874	17,874
1929	—	—	—	25,716	—	—	—	25,716	25,716
1930	—	—	—	5,809	—	—	—	5,809	5,809
1931	—	—	—	6,873	—	—	—	6,873	6,873
1932	—	—	—	4,922	—	—	—	4,922	4,922
1933	—	—	—	33,055	—	—	—	33,055	33,055
1934	—	—	—	51,467	—	—	—	51,467	51,467
1935	—	—	—	66,400	—	—	—	66,400	66,400
1936	—	—	—	45,697	—	—	—	45,697	45,697
1937	—	—	—	31,954	—	—	—	31,954	31,954
1938	—	—	—	34,502	—	—	—	34,502	34,502
1939	—	—	—	45,341	—	—	—	45,341	45,341
1940	—	—	—	48,786	—	—	—	48,786	48,786
1941	—	—	—	32,547	—	—	—	32,547	32,547
1942	—	—	—	21,872	—	—	—	21,872	21,872
1943	—	—	—	35,291	—	—	—	35,291	35,291
1944	—	—	—	36,644	—	—	—	36,644	36,644
1945	—	—	—	23,588	—	—	—	23,588	23,588
1946	—	851	851	26,715	—	—	—	26,715	27,566
1947	—	1,262	1,262	17,975	—	—	200	18,175	19,437
1948	—	515	515	17,329	—	—	281	17,610	18,125
1949	—	1,352	1,352	22,708	—	—	130	22,837	24,189
1950	—	2,029	2,029	15,372	—	—	92	15,464	17,493
1951	—	1,320	1,320	14,472	—	—	65	14,537	15,857
1952	—	1,052	1,052	9,171	—	—	103	9,273	10,326
1953	—	1,177	1,177	4,005	—	—	84	4,089	5,266
1954	—	5,681	5,681	12,342	—	—	442	12,784	18,465
1955	—	9,798	9,798	12,200	—	—	203	12,403	22,201
1956	—	10,725	10,725	25,938	—	—	172	26,110	36,835
1957	—	2,034	2,034	25,509	—	—	210	25,719	27,753
1958	—	449	449	11,238	—	—	188	11,426	11,875
1959	—	495	495	18,725	—	—	112	18,837	19,332
1960	—	2,981	2,981	17,724	—	—	117	17,841	20,823
1961	—	5,964	5,964	20,094	—	—	141	20,235	26,199
1962	—	3,231	3,231	20,527	—	—	143	20,670	23,901
1963	—	7,966	7,966	15,517	—	—	220	15,737	23,703
1964	—	8,618	8,618	11,283	—	—	87	11,370	19,988
1965	—	7,615	7,615	3,442	—	—	222	3,665	11,279
1966	—	5,290	5,290	1,848	—	—	267	2,115	7,405
1967	—	948	948	619	—	—	146	765	1,713
1968	—	107	107	1,492	—	—	96	1,588	1,695
1969	—	201	201	809	—	—	158	967	1,168
1970	—	400	400	277	—	—	158	435	835
1971	—	500	500	90	—	—	321	411	911

Table B-1: Landings (mt) of Pacific mackerel by region and fishing year (1926-2022). Mexican landings were from Magdalena Bay, BCS (MAG) and Ensenada, BC (ENS). US landings are from California (CA), Oregon (OR), Washington (WA). Additionally, California recreational landings are included (CA-REC). (*continued*)

Model Year	MAG	ENS	MEX-TOT	CA	OR	WA	CA-REC	USA-TOT	TOT
1972	—	200	200	28	—	—	304	332	532
1973	—	100	100	52	—	—	249	301	401
1974	—	471	471	43	—	—	120	163	634
1975	—	1,809	1,809	141	—	—	199	340	2,149
1976	—	1,271	1,271	2,654	—	—	167	2,821	4,092
1977	—	5,165	5,165	7,748	—	—	837	8,586	13,751
1978	—	7,372	7,372	18,446	—	—	1,355	19,801	27,173
1979	—	5,150	5,150	28,755	—	—	1,953	30,708	35,858
1980	304	4,242	4,546	27,972	—	—	2,685	30,657	35,203
1981	942	6,213	7,155	38,407	—	—	1,423	39,830	46,985
1982	144	4,185	4,329	30,626	—	—	1,416	32,042	36,371
1983	77	4,188	4,264	36,309	—	—	1,544	37,853	42,118
1984	310	5,451	5,761	39,240	—	—	1,467	40,707	46,468
1985	238	7,959	8,197	37,615	—	—	1,016	38,631	46,828
1986	25	8,941	8,965	44,298	—	—	859	45,157	54,123
1987	94	2,027	2,120	44,838	—	—	1,264	46,102	48,223
1988	592	6,017	6,608	41,968	—	—	689	42,656	49,265
1989	332	23,392	23,724	25,063	—	—	618	25,681	49,406
1990	0	30,961	30,961	39,974	—	—	616	40,590	71,551
1991	153	34,403	34,557	30,268	—	—	680	30,948	65,505
1992	17	6,153	6,170	25,584	—	—	464	26,047	32,217
1993	10	9,514	9,524	10,787	—	—	609	11,396	20,920
1994	159	13,143	13,302	9,372	—	—	1,063	10,435	23,737
1995	552	2,816	3,368	7,615	—	—	1,013	8,628	11,996
1996	3,906	10,183	14,089	9,788	—	—	686	10,473	24,563
1997	4,249	22,611	26,860	23,413	—	—	804	24,217	51,076
1998	2,229	40,586	42,815	19,578	—	—	430	20,008	62,823
1999	1,028	7,559	8,587	7,170	—	—	153	7,323	15,910
2000	0	6,530	6,530	20,936	139	48	325	21,449	27,979
2001	372	3,631	4,003	8,436	303	271	571	9,580	13,584
2002	3,050	7,278	10,328	3,541	128	249	254	4,171	14,499
2003	222	2,396	2,618	5,972	159	53	323	6,508	9,125
2004	83	1,628	1,711	5,012	111	24	478	5,624	7,336
2005	7	3,078	3,085	4,572	314	22	411	5,320	8,405
2006	19	1,967	1,986	7,870	669	42	372	8,953	10,939
2007	28	2,190	2,218	6,208	698	38	310	7,254	9,472
2008	689	114	803	4,198	58	9	279	4,543	5,346
2009	49	0	49	3,279	54	5	269	3,607	3,656
2010	312	1,605	1,917	2,047	48	2	216	2,313	4,229
2011	1,081	1,151	2,232	1,665	202	83	124	2,074	4,306
2012	7,219	171	7,390	3,202	1,588	719	99	5,608	12,998
2013	2,071	482	2,553	11,165	438	173	133	11,909	14,462
2014	2,757	1,342	4,099	3,651	1,215	502	225	5,593	9,692
2015	3,663	5,515	9,179	4,435	7	1	243	4,686	13,865
2016	5,730	5,977	11,707	2,523	4	22	209	2,757	14,464
2017	2,224	585	2,810	1,513	45	4	245	1,808	4,617
2018	3,422	12,330	15,752	2,199	112	10	180	2,501	18,252

Table B-1: Landings (mt) of Pacific mackerel by region and fishing year (1926-2022). Mexican landings were from Magdalena Bay, BCS (MAG) and Ensenada, BC (ENS). US landings are from California (CA), Oregon (OR), Washington (WA). Additionally, California recreational landings are included (CA-REC). (*continued*)

Model Year	MAG	ENS	MEX-TOT	CA	OR	WA	CA-REC	USA-TOT	TOT
2019	16,777	2,297	19,074	3,783	50	5	78	3,916	22,990
2020	26,136	5,232	31,368	500	101	3	87	691	32,060
2021	7,649	1,760	9,409	847	86	0	73	1,007	10,416
2022	7,649	7,361	15,010	543	366	26	56	990	16,000

11.3 Density-dependence in Pacific mackerel

Pacific mackerel have been at relatively low biomass levels from 2008-2022 (see historical analysis from Figure 34). Thus, while there may potentially be density-dependence in a number of biological processes, the recent stock biomass values may preclude observation of such effects.

Recently, there have been studies of Pacific mackerel in the northwestern Pacific Ocean documenting density-dependence in egg production (Takasuka et al. 2021) and body condition and growth (Kamimura et al. 2021). However, studies of density-dependence in US Pacific mackerel are less common. Parrish and MacCall (1978) found density-dependent spawner-recruit models explain 24% of the recruitment variability.

The base model does not have any explicitly density-dependent processes. The model has no growth relationship and uses empirical weight-at-age data. Density-dependence might be observed in the empirical weight-at-age values, but because the biomass has been relatively low in recent years, the effect may not be present in these data. The base model also has a Beverton-Holt spawner-recruit curve and natural mortality is time-invariant. The base model configuration may be revisited in the future

11.4 Difference in ageing methods between the fishery and survey data

The fishery and AT survey age data have slightly different protocols. Otoliths for double readings are selected from every third to fourth port sample in the fishery data and from every trawl sample in the AT survey. The target of ageing precision is 75% among readers in the fishery data and 100% for the AT survey. Double readings before 75% agreement are used to compute the fishery ageing errors and agreement is 100% when computing ageing error for the AT survey. See Appendix A for more details on the different methodologies.