

PR 10 499-310 p. 186

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460  <b>NOTICE OF PESTICIDE:</b> <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION NO. 499-310	DATE OF ISSUANCE APR 29 1991
	TERM OF ISSUANCE Conditional	
	NAME OF PESTICIDE PRODUCT Whitmire PT 566WB Pyrethrum Insect Fogger	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Whitmire Research Laboratories, Inc.  
 3568 Tree Court Industrial Boulevard  
 St. Louis, MO 63122

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
  - a. Add the phrase "EPA Registration No. 499-310."
  - b. On the front panel, delete the phrase "... but not limited to ..." in reference to sites for application.
  - c. Delete "Allyl homolog of Cinerin 1" from the ingredient statement. It no longer accurately describes the allethrin in this formulation.
  - d. Under FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, change the sentence to read "When using this product apply only when plant is not in operation."

**BEST AVAILABLE COPY**

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE APR 29 1991
---------------------------------	---------------------

EPA Form 8570-4 (Rev. 8-76) THIS COPY MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

3. Delete the last sentence in that paragraph "Careful crack and crevice application for inspection purposes is permissible at all times."

4. Submit a 1-year storage stability study on this formulation. This study is due 15 months from the date of registration.

5. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

*R. F. Mountfort*  
Richard F. Mountfort  
Product Manager (10)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosures

**BEST AVAILABLE COPY**

FRONT PANEL:

7/23/90

Whitmire  
PT® 566WB  
Pyrethrum  
Insect Fogger

KILLS: Anguomois Grain Moths, Ants, Bed Bugs, Booklice, Carpet Beetles, Centipedes, Chocolate Moths, Cigarette Beetles, Clover Mites, Cluster Flies, Cockroaches, Confused Flour Beetles, Crickets, Drug Store Beetles, Fleas, Flies, Fruit Flies, Gnats, Grain Mites, Granary Weevils, Horn Flies, House Flies, Indian Meal Moths, Mediterranean Flour Moths, Millipedes, Mosquitoes, Mud Daubers, Red Flour Beetles, Rice Weevils, Sawtoothed Grain Beetles, Silverfish, Small Flying Moths, Sow Bugs, Spiders, Stable Flies, and Wasps.

For Use in and Around, but not limited to: Campgrounds, Food Storage Areas, Homes, Hospitals, Hotels, Meat Packing and Food Processing Plants, Motels, Resorts, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Trucks, Planes), Utilities, Warehouses, and other Commercial and Industrial Buildings.

Not for use in Federally inspected ~~meat~~ plants.

ACTIVE INGREDIENTS:

Pyrethrum . . . . .	0.25Z
d-trans Allethrin (Allyl homolog of Cinerin I) . . . . .	0.25Z
Piperonyl Butoxide, Technical* . . . . .	1.00Z
n-Octyl Bicycloheptene Dicarboximide . . . . .	1.00Z

INERT INGREDIENTS: . . . . . 97.50Z

\*Equivalent to 0.8Z (Butylcarbityl)(6-Propylpiperonyl) Ether and 0.2Z Related Compounds

EPA Reg. No. 499-GRN

EPA Est. No. 499-MO-1

KEEP OUT OF REACH OF CHILDREN  
CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. Do not induce vomiting. Vomiting may cause aspiration pneumonia.

If inhaled: Remove patient to fresh air. Apply artificial respiration if indicated.

If on skin: Remove contaminated clothing and wash affected skin areas with soap and water. Get medical attention if irritation persists.

If in eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

See side panel for additional precautionary statements.

NET WEIGHT:

**BEST AVAILABLE COPY**

ACCEPTED  
with COMMENTS  
in EPA Letter dated  
APR 29 1991  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
499-310

Side Panel:

PT® 566WB  
Pyrethrum

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or absorbed through skin. Do not remain in treated area and ventilate the area after treatment is completed. Avoid breathing of vapor. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using and before smoking or eating. Do not contaminate water, food, or foodstuffs. Remove all pets and birds. Cover fish tanks or remove before spraying.

**FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS:** Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product ~~as a space spray~~, apply only when plant is not in operation. Careful crack and crevice application for inspection purposes is permissible at all times.

**NOTICE TO APPLICATOR:** During any general volumetric treatment, wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute of Occupational Safety and Health under the provisions of 30 CFR Part 11.

**ENVIRONMENTAL HAZARDS-** This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Apply this product only as specified on this label.

**PHYSICAL OR CHEMICAL HAZARDS -** Extremely Flammable. Contents under pressure. Keep away from heat, sparks, and open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted, or varnished surfaces or directly into any electronic equipment such as radios, televisions, computers. etc.

**DIRECTIONS FOR USE-** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**HOW TO USE WHITMIRE CRACK AND CREVICE INJECTION SYSTEM:** Whitmire PT® 566WB is for use only with the supplied actuator and injection tubes or other Whitmire equipment. Inject PT® 566WB into cracks and crevices or void space where insects may be harboring, living and breeding. Place injector tip 1/2" into cracks and crevices, holes and other small openings. Release approximately one-second spray per spot. For a serious infestation, place spots approximately 12" apart. For light infestation or for preventative treatment, place spots approximately three feet apart. In long running cracks, move injector tip along the crack while spraying at the rate of three linear foot per second for a light infestation. For heavy infestations move injector tip along at one linear foot per second. For inaccessible closed voids calculate the voids cubic area and treat at rate of five to ten seconds per three cubic feet. Several holes may be required in long-running voids. Repeat as necessary.

**BEST AVAILABLE COPY**

Side Panel (cont.)

Flying Insects: Cluster Fly, Flies, Fruit Flies, Gnats, Mosquitoes, Small Flying Moths. GENERAL VOLUMETRIC TREATMENT: Close all windows and doors and direct fog upward at a rate of 1 to 3 seconds per 1000 cu. ft. (1/2 to 1 1/2 seconds per 1000 cu. ft. for 8 & 15 lb. size due to delivery rate). Disperse in all locations. Keep area closed for 15 min. Open and ventilate before reoccupying. Repeat treatment as necessary.

Ants, Booklice, Centipedes, Clover Mites, Crickets, Millipedes, Roaches, Silverfish, Sow Bugs, and Spiders. Inject into cracks and crevices in all hiding places behind baseboards, sinks, cabinets, meter boxes, door frames, windows. General Volumetric treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, fans and close doors and windows before spraying. Spray at rate of 20 seconds per 1000 cu. ft. of space (for 8 & 15 lb. size: 10 seconds per 1000 cu. ft.) Direct spray toward area suspected of harboring the greatest insect infestations. Dispense in all locations contacting as many insects as possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

Bed Bugs: Take bed apart and treat cracks, joints and interior of framework. Mattresses and box springs should be treated, especially tufts, folds, and edges. Picture frames, moldings and all cracks and crevices in the room should also be treated. Do not use in hospital patient quarters.

Mud Dauber and Wasp Nests: Spray mist directly on insects and their nests from approximately 2 ft. distance. Contacted insects will fly away from fog. Applications should be made in late evening when insects are at rest. Spray into hiding and breeding places contacting as many insects as possible.

Carpet Beetles, Cigarette Beetles, Confused Flour Beetles, Drug Store Beetles, Grain Mites in exposed adult and larvae stages, Granary Weevils, Red Flour Beetles, Rice Weevils, and Sawtoothed Grain Beetles. Contact as many insects as possible. Inject into cracks and crevices, including processing and handling equipment. Drift around cartons, and containers, and other areas where these insects tend to congregate. General Volumetric Treatment: Spray 20 seconds per 1000 cu. ft. of space (for 8 & 15 lb. size: Spray 10 seconds per 1000 cu. ft.) Repeat treatment weekly or as insects appear. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Anguomois Grain Moth, Chocolate Moth, Indian Meal Moth, Mediterranean Flour Moth: Close all doors and windows. Direct spray into all parts of room, especially around stored product containers, pallets, and darkened areas. Spray 5 to 10 seconds for each 1000 cu. ft. of space (for 8 & 15 lb. size: spray 2 1/2 to 5 seconds per 1000 cu. ft.) Keep area closed for 15 minutes following treatment. Open and ventilate before reoccupying. Repeat treatment weekly or as necessary. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

6/16

Side Panel (cont.)

Fleas in buildings-General Volumetric Treatment: Hold aerosol 36" above floor and direct spray toward floor and lower walls at a rate of 10 seconds per 100 sq. ft. (5 seconds per 100 sq. ft. for 15 lb. unit) making sure that all floor, sofa and chair surfaces are contacted. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment after 7 days or as necessary.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS: (Attics, false ceilings, wall and equipment voids.) For spray application with Whitmire Crack & Crevice injection equipment.

For exposed stages of ants, beetles, centipedes, cockroaches, crickets, millipedes, silverfish, sowbugs and spiders harboring in void, calculate void volume and treat at a rate of 20 seconds per 1000 cu. ft. (for 15 lb. size: 10 seconds per 1000 cu. ft.) Repeat treatment as necessary.

OUTDOOR GROUND APPLICATION: Flies, Gnats and Mosquitoes in open areas near building and in campgrounds. Best results are obtained when wind speed is five (5) MPH or less. Apply at a rate of 60-80 seconds per acre. (For 8 & 15 lb. size: Spray 30-40 seconds per acre). Spray in wide swaths across area to be treated. Allow spray drift to penetrate dense foliage. Repeat treatment as necessary.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool area away from heat or open flame.

Pesticide Disposal: Waste resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Replace cap and discard empty container in trash, sanitary landfill, or by other approved state and local procedures. Do not puncture or incinerate.

Copyright, 1990  
Whitmire Research Laboratories, Inc.  
St. Louis, MO 63122

**BEST AVAILABLE COPY**