

LESCO

ACCEPTED

APR 11 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10404-14

RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY

FOR RETAIL SALE TO AND USE BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

Diazinon 3.33% + Fertilizer



NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

To Control Both Soil and Surface-Feeding Insects
Ready-to-Use - Contains 3.33% Diazinon
This product must not be used on golf courses or sod farms.

Overall Lawn Application: LESCO Diazinon 3.33% + XX-X-X Fertilizer controls the following lawn insects on both grass and dichondra lawns:

Ants	Chiggers	Cutworms	Leafhoppers
Armyworms	Chinch bugs	Digger Wasps	Millipedes
Bermudagrass mites	Clover mites	Earwigs	Springtails(Collembola)
Billbugs	Crickets	Lawn moths	White grubs of: -Japanese beetle -European chafer -Southern chafer
	Sowbugs	(Sod Webworms)	

Active Ingredient:
0,0-diethyl 0-(2-isopropyl-6-methyl-4 pyrimidinyl) phosphorothioate 3.33%

Inert Ingredients: 96.67%

Total 100.00%

GUARANTEED ANALYSIS

Total Nitrogen (N) XX.00%

X.XX% Ammoniacal Nitrogen

X.XX% Urea Nitrogen*

Available Phosphate (P₂O₅) X.XX%

Soluble Potash (K₂O) X.XX%

Sulfur (S) Total X.XX%

X.XX% Free Sulfur (S)

DERIVED FROM: Fertilizer ingredients will vary and will be determined by analysis of finished product.

Chlorine (Cl) Max. X.XX%

* X.XX% Slowly Available Nitrogen from LESCO™ Poly Plus® Sulfur Coated Urea.

Brush off walks and painted surfaces following application to avoid discoloration.

F335 Made in U.S.A.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes or clothing. May cause contact sensitization following repeated contact with skin of susceptible individuals. If sensitization reactions result, contact a physician. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Do not contaminate ornamental fish ponds. Do not allow children or pets onto treated areas until granules have been watered into the soil and the grass is dry.

STATEMENT OF PRACTICAL TREATMENT
If Swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.
If Inhaled: Remove victim to fresh air and apply artificial respiration if indicated.
If On Skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.
If In Eyes: Rinse eyes with water for at least 15 minutes and call a physician immediately.
NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS
This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks and geese), feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Follow all watering instructions in the Directions For Use, however, stop watering before puddling occurs. For lawns, do not apply this product to sites occupied by waterfowl, or within 75 feet of any body of water which may attract waterfowl, such as ponds or lakes. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to fish and other aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS
Birds, especially waterfowl (ducks or geese) feeding on treated areas may be killed. To reduce risks to birds:

- Do not treat areas larger than 15,000 sq. ft.
- Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20.
- Do not exceed maximum permitted label rates, since application rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
- Avoid overlapping treated areas.
- Follow all watering instruction in the Directions for Use, however, stop watering before puddling occurs.
- Do not apply this product to sites occupied by waterfowl, or within 75 feet of any body of water which may attract waterfowl, such as lakes or ponds.
- If it is necessary to repeat applications, wait a minimum of three weeks between applications. Make no more than 4 applications per year.

STORAGE AND DISPOSAL

Pesticide Storage: Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place and avoid excess heat. Carefully open container. After partial use, fold and roll back bag, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In Case of Spill: Avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills. If dry material: Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Apply LESCO Diazinon 3.33% + XX-X-X Fertilizer to lawn with fertilizer spreader at the rate shown. Application should be made when grass is dry. Water lawn lightly after application. Repeat as necessary. Apply at a rate of 3 pounds per 1,000 sq. ft. (an area of 20 ft. x 50 ft.)

Billbugs, White Grubs of Japanese beetle, European chafer, and Southern chafer: Apply 3 lbs./1,000 sq. ft. of lawn. Water lawn thoroughly after application. For grub control, make application any time between late July and early October. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

Hyperodes Weevil (New York Only): Apply 3 lbs./1,000 sq. ft. of lawn. Water lawn thoroughly after application. Treat problem areas in mid-April and again in mid-May.

APPROXIMATE SPREADER SETTINGS FOR LESCO ROTARY SPREADER:*

#Product/ 1,000 sq. ft.	#N/ 1,000 sq. ft.	#A/A Spreading 4.36	Effective 9 feet	Third 1	Approx.* G
3	0.72	4.36	9 feet	1	G
Cyclone* 3 lbs./1,000 sq. ft.	Once Over 4.75	Twice Over 4.00	Spread Width 7 feet		

*IMPORTANT: These settings are only approximate. Age, condition of spreader, and operator speed can cause wide variation. Be sure to calibrate your spreader with each application.

WARRANTY

LESCO warrants that this product conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this product is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

LESCO is a trademark of LESCO, Inc.
Poly Plus is a registered trademark of LESCO, Inc.
C:\WPWIN6\WPDOCS\LEPA\FERT.DZN Rev. 03/87/WPK
Net Weight: 50 Lb (22.72 kg)

REC'D APR -2

EPA Reg. No. 10404-14
EPA Est. No. 10404-OH-04
#000000

F335
Manufactured by: LESCO, Inc.
20005 Lake Road, P.O. Box 16915
Rocky River, Ohio 44116

